

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.1-Sine/179-4.1.1.1

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 11:49pm

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24
2	detailed summary tables of results	25
2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	55
3	Listing of integrals	59
3.1	$\int (a + a \sin(c + dx))^3 dx$	62
3.2	$\int (a + a \sin(c + dx))^2 dx$	68
3.3	$\int (a + a \sin(c + dx)) dx$	73
3.4	$\int \frac{1}{a+a \sin(c+dx)} dx$	78
3.5	$\int \frac{1}{(a+a \sin(c+dx))^2} dx$	83
3.6	$\int \frac{1}{(a+a \sin(c+dx))^3} dx$	89
3.7	$\int (a - a \sin(c + dx))^3 dx$	96

3.8	$\int (a - a \sin(c + dx))^2 dx$	102
3.9	$\int (a - a \sin(c + dx)) dx$	107
3.10	$\int \frac{1}{a - a \sin(c + dx)} dx$	112
3.11	$\int \frac{1}{(a - a \sin(c + dx))^2} dx$	117
3.12	$\int \frac{1}{(a - a \sin(c + dx))^3} dx$	123
3.13	$\int (a + a \sin(c + dx))^{7/2} dx$	130
3.14	$\int (a + a \sin(c + dx))^{5/2} dx$	137
3.15	$\int (a + a \sin(c + dx))^{3/2} dx$	143
3.16	$\int \sqrt{a + a \sin(c + dx)} dx$	149
3.17	$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx$	154
3.18	$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx$	160
3.19	$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx$	166
3.20	$\int (a - a \sin(c + dx))^{7/2} dx$	173
3.21	$\int (a - a \sin(c + dx))^{5/2} dx$	180
3.22	$\int (a - a \sin(c + dx))^{3/2} dx$	186
3.23	$\int \sqrt{a - a \sin(c + dx)} dx$	192
3.24	$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx$	197
3.25	$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx$	203
3.26	$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx$	209
3.27	$\int (-a + a \sin(c + dx))^{7/2} dx$	216
3.28	$\int (-a + a \sin(c + dx))^{5/2} dx$	223
3.29	$\int (-a + a \sin(c + dx))^{3/2} dx$	229
3.30	$\int \sqrt{-a + a \sin(c + dx)} dx$	235
3.31	$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx$	240
3.32	$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx$	246
3.33	$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx$	252
3.34	$\int (a + a \sin(c + dx))^{4/3} dx$	259
3.35	$\int (a + a \sin(c + dx))^{2/3} dx$	264
3.36	$\int \sqrt[3]{a + a \sin(c + dx)} dx$	269
3.37	$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx$	274
3.38	$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx$	279
3.39	$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx$	284
3.40	$\int (a + a \sin(c + dx))^n dx$	289
3.41	$\int (a - a \sin(c + dx))^n dx$	294
3.42	$\int (2 + 2 \sin(c + dx))^n dx$	299
3.43	$\int (2 - 2 \sin(c + dx))^n dx$	304
3.44	$\int \frac{1}{5 + 3 \sin(c + dx)} dx$	309

3.45	$\int \frac{1}{(5+3\sin(c+dx))^2} dx$	314
3.46	$\int \frac{1}{(5+3\sin(c+dx))^3} dx$	321
3.47	$\int \frac{1}{(5+3\sin(c+dx))^4} dx$	329
3.48	$\int \frac{1}{5-3\sin(c+dx)} dx$	338
3.49	$\int \frac{1}{(5-3\sin(c+dx))^2} dx$	343
3.50	$\int \frac{1}{(5-3\sin(c+dx))^3} dx$	350
3.51	$\int \frac{1}{(5-3\sin(c+dx))^4} dx$	358
3.52	$\int \frac{1}{-5+3\sin(c+dx)} dx$	367
3.53	$\int \frac{1}{(-5+3\sin(c+dx))^2} dx$	372
3.54	$\int \frac{1}{(-5+3\sin(c+dx))^3} dx$	379
3.55	$\int \frac{1}{(-5+3\sin(c+dx))^4} dx$	387
3.56	$\int \frac{1}{-5-3\sin(c+dx)} dx$	396
3.57	$\int \frac{1}{(-5-3\sin(c+dx))^2} dx$	401
3.58	$\int \frac{1}{(-5-3\sin(c+dx))^3} dx$	408
3.59	$\int \frac{1}{(-5-3\sin(c+dx))^4} dx$	416
3.60	$\int \frac{1}{3+5\sin(c+dx)} dx$	425
3.61	$\int \frac{1}{(3+5\sin(c+dx))^2} dx$	431
3.62	$\int \frac{1}{(3+5\sin(c+dx))^3} dx$	438
3.63	$\int \frac{1}{(3+5\sin(c+dx))^4} dx$	447
3.64	$\int \frac{1}{3-5\sin(c+dx)} dx$	457
3.65	$\int \frac{1}{(3-5\sin(c+dx))^2} dx$	463
3.66	$\int \frac{1}{(3-5\sin(c+dx))^3} dx$	470
3.67	$\int \frac{1}{(3-5\sin(c+dx))^4} dx$	479
3.68	$\int \frac{1}{-3+5\sin(c+dx)} dx$	489
3.69	$\int \frac{1}{(-3+5\sin(c+dx))^2} dx$	495
3.70	$\int \frac{1}{(-3+5\sin(c+dx))^3} dx$	502
3.71	$\int \frac{1}{(-3+5\sin(c+dx))^4} dx$	511
3.72	$\int \frac{1}{-3-5\sin(c+dx)} dx$	521
3.73	$\int \frac{1}{(-3-5\sin(c+dx))^2} dx$	527
3.74	$\int \frac{1}{(-3-5\sin(c+dx))^3} dx$	534
3.75	$\int \frac{1}{(-3-5\sin(c+dx))^4} dx$	543
3.76	$\int (a + b \sin(c + dx))^{7/2} dx$	553
3.77	$\int (a + b \sin(c + dx))^{5/2} dx$	564
3.78	$\int (a + b \sin(c + dx))^{3/2} dx$	574
3.79	$\int \sqrt{a + b \sin(c + dx)} dx$	583
3.80	$\int \frac{1}{\sqrt{a+b\sin(c+dx)}} dx$	589

3.81	$\int \frac{1}{(a+b \sin(c+dx))^{3/2}} dx$	595
3.82	$\int \frac{1}{(a+b \sin(c+dx))^{5/2}} dx$	602
3.83	$\int \frac{1}{(a+b \sin(c+dx))^{7/2}} dx$	612
3.84	$\int (a + b \sin(c + dx))^{4/3} dx$	623
3.85	$\int (a + b \sin(c + dx))^{2/3} dx$	629
3.86	$\int \sqrt[3]{a + b \sin(c + dx)} dx$	635
3.87	$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx$	641
3.88	$\int \frac{1}{(a+b \sin(c+dx))^{2/3}} dx$	647
3.89	$\int \frac{1}{(a+b \sin(c+dx))^{4/3}} dx$	653
3.90	$\int (a + b \sin(c + dx))^n dx$	659
3.91	$\int (3 + 4 \sin(c + dx))^n dx$	665
3.92	$\int (3 - 4 \sin(c + dx))^n dx$	670
3.93	$\int (4 + 3 \sin(c + dx))^n dx$	675
3.94	$\int (4 - 3 \sin(c + dx))^n dx$	680
3.95	$\int (-3 + 4 \sin(c + dx))^n dx$	685
3.96	$\int (-3 - 4 \sin(c + dx))^n dx$	690
3.97	$\int (-4 + 3 \sin(c + dx))^n dx$	695
3.98	$\int (-4 - 3 \sin(c + dx))^n dx$	701
4	Appendix	707
4.1	Listing of Grading functions	707
4.2	Links to plain text integration problems used in this report for each CAS725	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [98]. This is test number [179].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (98)	0.00 (0)
Mathematica	100.00 (98)	0.00 (0)
Maple	74.49 (73)	25.51 (25)
Fricas	74.49 (73)	25.51 (25)
Giac	66.33 (65)	33.67 (33)
Mupad	51.02 (50)	48.98 (48)
Maxima	44.90 (44)	55.10 (54)
Reduce	44.90 (44)	55.10 (54)
Sympy	44.90 (44)	55.10 (54)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	76.531	10.204	13.265	0.000
Fricas	59.184	7.143	8.163	25.510
Giac	59.184	5.102	2.041	33.673
Maple	58.163	12.245	4.082	25.510
Maxima	28.571	16.327	0.000	55.102
Sympy	10.204	22.449	12.245	55.102
Mupad	0.000	51.020	0.000	48.980
Reduce	0.000	44.898	0.000	55.102

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	25	100.00	0.00	0.00
Maple	25	100.00	0.00	0.00
Giac	33	100.00	0.00	0.00
Mupad	48	0.00	100.00	0.00
Maxima	54	100.00	0.00	0.00
Reduce	54	100.00	0.00	0.00
Sympy	54	92.59	7.41	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.06
Fricas	0.10
Giac	0.14
Reduce	0.17
Rubi	0.33
Maple	0.43
Mathematica	0.69
Sympy	1.36
Mupad	20.51

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	85.02	0.99	72.00	1.00
Mupad	89.66	1.28	83.00	1.26
Giac	93.38	1.22	81.00	1.09
Reduce	104.23	1.23	81.00	1.13
Mathematica	113.15	1.42	105.50	1.37
Maple	127.58	1.24	68.00	0.96
Maxima	129.14	1.56	115.00	1.63
Fricas	153.52	1.67	115.00	1.23
Sympy	694.70	7.01	425.50	5.89

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {98}

Mathematica {34, 36, 38, 84, 85, 86, 87, 88, 89, 90, 93, 98}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	55

2.1 List of integrals sorted by grade for each CAS

Rubi	26
Mma	26
Maple	27
Fricas	27
Maxima	28
Giac	28
Mupad	28
Sympy	29
Reduce	29

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 20, 21, 22, 27, 28, 29, 35, 37, 39, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 88, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

B grade { 4, 10, 16, 23, 30, 34, 36, 38, 84, 89 }

C grade { 17, 18, 19, 24, 25, 26, 31, 32, 33, 40, 41, 42, 43 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 7, 8, 9, 10, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 27, 28, 29, 30, 31, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75 }

B grade { 19, 26, 32, 33, 76, 77, 78, 79, 80, 81, 82, 83 }

C grade { 5, 6, 11, 12 }

F normal fail { 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 20, 21, 22, 23, 24, 27, 28, 29, 30, 31, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75 }

B grade { 16, 18, 19, 25, 26, 32, 33 }

C grade { 76, 77, 78, 79, 80, 81, 82, 83 }

F normal fail { 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 7, 8, 9, 10, 44, 45, 48, 49, 52, 53, 56, 57, 60, 61, 62, 64, 65, 66, 68, 69, 70, 72, 73, 74 }

B grade { 5, 6, 11, 12, 46, 47, 50, 51, 54, 55, 58, 59, 63, 67, 71, 75 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 27, 28, 29, 30, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75 }

B grade { 18, 25, 26, 32, 33 }

C grade { 24, 31 }

F normal fail { 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 16, 17, 23, 30, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 79, 80 }

C grade { }

F normal fail { }

F(-1) timedout fail { 13, 14, 15, 18, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 76, 77, 78, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-2) exception fail { }

Sympy

A grade { 2, 3, 4, 8, 9, 10, 60, 64, 68, 72 }

B grade { 1, 5, 6, 7, 11, 12, 44, 48, 52, 56, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75 }

C grade { 45, 46, 47, 49, 50, 51, 53, 54, 55, 57, 58, 59 }

F normal fail { 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { 13, 20, 27, 76 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75 }

C grade { }

F normal fail { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	44	43	72	54	121	55	52	156
N.S.	1	1.00	0.70	0.68	1.14	0.86	1.92	0.87	0.83	2.48
time (sec)	N/A	0.238	3.851	7.242	0.034	0.076	0.124	0.125	0.173	28.280

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	34	32	47	41	78	38	36	123
N.S.	1	1.00	0.76	0.71	1.04	0.91	1.73	0.84	0.80	2.73
time (sec)	N/A	0.182	0.502	0.422	0.034	0.102	0.095	0.125	0.161	26.097

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	27	17	16	18	19	16	17	25
N.S.	1	1.00	1.69	1.06	1.00	1.12	1.19	1.00	1.06	1.56
time (sec)	N/A	0.139	0.037	0.082	0.029	0.083	0.056	0.126	0.262	25.714

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	48	22	27	42	27	21	30	21
N.S.	1	1.00	2.09	0.96	1.17	1.83	1.17	0.91	1.30	0.91
time (sec)	N/A	0.173	0.265	0.079	0.036	0.080	0.407	0.130	0.218	25.750

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	54	54	38	117	95	221	47	60	76
N.S.	1	0.98	0.98	0.69	2.13	1.73	4.02	0.85	1.09	1.38
time (sec)	N/A	0.248	0.463	0.158	0.045	0.071	0.710	0.112	0.160	25.737

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	87	76	48	203	147	558	73	112	133
N.S.	1	1.05	0.92	0.58	2.45	1.77	6.72	0.88	1.35	1.60
time (sec)	N/A	0.330	0.424	0.211	0.038	0.084	1.368	0.121	0.162	25.902

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	46	45	72	54	121	55	52	156
N.S.	1	1.00	0.73	0.71	1.14	0.86	1.92	0.87	0.83	2.48
time (sec)	N/A	0.241	5.378	7.089	0.029	0.083	0.124	0.125	0.169	28.209

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	36	34	47	41	78	38	36	123
N.S.	1	1.00	0.80	0.76	1.04	0.91	1.73	0.84	0.80	2.73
time (sec)	N/A	0.185	0.641	0.404	0.030	0.087	0.091	0.119	0.170	25.933

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	27	16	15	17	19	15	15	25
N.S.	1	1.00	1.80	1.07	1.00	1.13	1.27	1.00	1.00	1.67
time (sec)	N/A	0.142	0.039	0.071	0.027	0.092	0.060	0.114	0.150	25.641

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	51	22	28	40	27	21	30	21
N.S.	1	1.00	2.22	0.96	1.22	1.74	1.17	0.91	1.30	0.91
time (sec)	N/A	0.175	0.261	0.077	0.027	0.076	0.384	0.119	0.157	25.771

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	56	56	38	118	95	219	47	62	78
N.S.	1	0.98	0.98	0.67	2.07	1.67	3.84	0.82	1.09	1.37
time (sec)	N/A	0.251	0.390	0.149	0.037	0.073	0.693	0.121	0.161	25.951

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	90	74	48	204	147	556	73	112	135
N.S.	1	1.05	0.86	0.56	2.37	1.71	6.47	0.85	1.30	1.57
time (sec)	N/A	0.335	0.458	0.209	0.039	0.066	1.318	0.127	0.160	25.908

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F(-1)	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	119	127	154	75	0	140	0	132	80	0
N.S.	1	1.07	1.29	0.63	0.00	1.18	0.00	1.11	0.67	0.00
time (sec)	N/A	0.433	3.145	0.273	0.000	0.088	0.000	0.143	0.175	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	93	117	65	0	115	0	102	58	0
N.S.	1	1.04	1.31	0.73	0.00	1.29	0.00	1.15	0.65	0.00
time (sec)	N/A	0.336	1.943	0.162	0.000	0.096	0.000	0.138	0.180	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	89	53	0	76	0	67	34	0
N.S.	1	1.00	1.51	0.90	0.00	1.29	0.00	1.14	0.58	0.00
time (sec)	N/A	0.247	0.757	0.115	0.000	0.091	0.000	0.136	0.152	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	F	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	65	43	0	50	0	36	14	33
N.S.	1	1.00	2.50	1.65	0.00	1.92	0.00	1.38	0.54	1.27
time (sec)	N/A	0.173	0.083	0.115	0.000	0.076	0.000	0.125	0.159	0.222

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	73	75	0	193	0	61	28	49
N.S.	1	1.00	1.55	1.60	0.00	4.11	0.00	1.30	0.60	1.04
time (sec)	N/A	0.190	0.224	0.328	0.000	0.098	0.000	0.140	0.158	0.405

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	108	125	0	252	0	133	38	0
N.S.	1	1.00	1.40	1.62	0.00	3.27	0.00	1.73	0.49	0.00
time (sec)	N/A	0.261	0.344	0.117	0.000	0.114	0.000	0.135	0.154	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	112	196	199	0	320	0	153	48	0
N.S.	1	1.05	1.83	1.86	0.00	2.99	0.00	1.43	0.45	0.00
time (sec)	N/A	0.356	0.345	0.122	0.000	0.106	0.000	0.145	0.169	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F(-1)	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	131	157	76	0	143	0	132	90	0
N.S.	1	1.07	1.28	0.62	0.00	1.16	0.00	1.07	0.73	0.00
time (sec)	N/A	0.448	3.503	0.220	0.000	0.078	0.000	0.202	0.176	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	96	133	66	0	116	0	102	64	0
N.S.	1	1.04	1.45	0.72	0.00	1.26	0.00	1.11	0.70	0.00
time (sec)	N/A	0.354	2.251	0.149	0.000	0.105	0.000	0.180	0.166	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	103	54	0	79	0	68	40	0
N.S.	1	1.00	1.69	0.89	0.00	1.30	0.00	1.11	0.66	0.00
time (sec)	N/A	0.258	0.926	0.102	0.000	0.095	0.000	0.158	0.169	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	F	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	66	44	0	50	0	36	16	34
N.S.	1	1.00	2.44	1.63	0.00	1.85	0.00	1.33	0.59	1.26
time (sec)	N/A	0.177	0.049	0.113	0.000	0.074	0.000	0.133	0.179	25.976

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	76	74	0	190	0	42	31	0
N.S.	1	1.00	1.62	1.57	0.00	4.04	0.00	0.89	0.66	0.00
time (sec)	N/A	0.194	0.311	0.311	0.000	0.089	0.000	0.141	0.166	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	122	123	0	246	0	206	40	0
N.S.	1	1.00	1.54	1.56	0.00	3.11	0.00	2.61	0.51	0.00
time (sec)	N/A	0.277	0.421	0.100	0.000	0.109	0.000	0.185	0.180	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	115	160	194	0	321	0	303	51	0
N.S.	1	1.05	1.45	1.76	0.00	2.92	0.00	2.75	0.46	0.00
time (sec)	N/A	0.372	0.586	0.097	0.000	0.100	0.000	0.198	0.165	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F(-1)	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	135	156	77	0	144	0	134	82	0
N.S.	1	1.06	1.23	0.61	0.00	1.13	0.00	1.06	0.65	0.00
time (sec)	N/A	0.456	2.858	0.167	0.000	0.084	0.000	0.200	0.179	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	99	119	67	0	117	0	104	58	0
N.S.	1	1.04	1.25	0.71	0.00	1.23	0.00	1.09	0.61	0.00
time (sec)	N/A	0.349	1.828	0.165	0.000	0.097	0.000	0.180	0.169	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	93	55	0	80	0	70	36	0
N.S.	1	1.00	1.48	0.87	0.00	1.27	0.00	1.11	0.57	0.00
time (sec)	N/A	0.252	0.691	0.116	0.000	0.105	0.000	0.156	0.174	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	F	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	65	45	0	51	0	38	14	33
N.S.	1	1.00	2.32	1.61	0.00	1.82	0.00	1.36	0.50	1.18
time (sec)	N/A	0.178	0.059	0.116	0.000	0.103	0.000	0.126	0.183	0.250

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	75	76	0	192	0	40	28	0
N.S.	1	1.00	1.56	1.58	0.00	4.00	0.00	0.83	0.58	0.00
time (sec)	N/A	0.193	0.171	0.253	0.000	0.104	0.000	0.146	0.166	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	108	134	0	252	0	215	38	0
N.S.	1	1.00	1.33	1.65	0.00	3.11	0.00	2.65	0.47	0.00
time (sec)	N/A	0.269	0.283	0.118	0.000	0.109	0.000	0.174	0.168	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	118	146	211	0	327	0	313	48	0
N.S.	1	1.04	1.29	1.87	0.00	2.89	0.00	2.77	0.42	0.00
time (sec)	N/A	0.371	0.427	0.119	0.000	0.106	0.000	0.204	0.170	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	223	0	0	0	0	0	36	0
N.S.	1	1.00	3.33	0.00	0.00	0.00	0.00	0.00	0.54	0.00
time (sec)	N/A	0.253	1.843	0.000	0.000	0.000	0.000	0.000	0.174	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	124	0	0	0	0	0	16	0
N.S.	1	1.00	1.88	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.259	0.470	0.000	0.000	0.000	0.000	0.000	0.176	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	169	0	0	0	0	0	16	0
N.S.	1	1.00	2.56	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.258	0.695	0.000	0.000	0.000	0.000	0.000	0.164	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	70	0	0	0	0	0	16	0
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.260	0.332	0.000	0.000	0.000	0.000	0.000	0.177	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	215	0	0	0	0	0	16	0
N.S.	1	1.00	3.26	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.256	1.130	0.000	0.000	0.000	0.000	0.000	0.235	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	130	0	0	0	0	0	36	0
N.S.	1	1.00	1.88	0.00	0.00	0.00	0.00	0.00	0.52	0.00
time (sec)	N/A	0.259	0.535	0.000	0.000	0.000	0.000	0.000	0.225	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	70	0	0	0	0	0	14	0
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.253	0.185	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	80	0	0	0	0	0	15	0
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.267	0.294	0.000	0.000	0.000	0.000	0.000	0.162	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	46	0	0	0	0	0	14	0
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.195	0.217	0.000	0.000	0.000	0.000	0.000	0.159	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	49	0	0	0	0	0	14	0
N.S.	1	1.00	0.83	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	0.197	0.223	0.000	0.000	0.000	0.000	0.000	0.157	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	56	20	26	26	46	49	19	40
N.S.	1	1.00	1.81	0.65	0.84	0.84	1.48	1.58	0.61	1.29
time (sec)	N/A	0.184	0.060	0.142	0.104	0.092	0.367	0.122	0.166	26.027

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	61	91	63	93	59	388	95	64	83
N.S.	1	1.09	1.62	1.12	1.66	1.05	6.93	1.70	1.14	1.48
time (sec)	N/A	0.265	0.212	0.230	0.110	0.132	0.911	0.122	0.173	25.950

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	91	113	91	173	94	918	121	131	111
N.S.	1	1.12	1.40	1.12	2.14	1.16	11.33	1.49	1.62	1.37
time (sec)	N/A	0.368	0.427	0.352	0.127	0.081	1.966	0.122	0.155	26.004

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	121	133	117	253	130	1693	147	191	187
N.S.	1	1.14	1.25	1.10	2.39	1.23	15.97	1.39	1.80	1.76
time (sec)	N/A	0.503	0.563	0.535	0.121	0.116	4.553	0.125	0.170	26.516

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	56	20	26	26	46	50	19	40
N.S.	1	1.00	1.70	0.61	0.79	0.79	1.39	1.52	0.58	1.21
time (sec)	N/A	0.183	0.054	0.138	0.108	0.110	0.387	0.120	0.156	25.853

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	63	91	63	93	59	384	96	64	83
N.S.	1	1.09	1.57	1.09	1.60	1.02	6.62	1.66	1.10	1.43
time (sec)	N/A	0.267	0.298	0.237	0.112	0.121	0.916	0.123	0.155	26.006

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	93	113	91	173	94	915	122	131	111
N.S.	1	1.12	1.36	1.10	2.08	1.13	11.02	1.47	1.58	1.34
time (sec)	N/A	0.387	0.746	0.353	0.117	0.118	2.043	0.122	0.168	25.976

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	123	133	117	253	130	1690	148	191	187
N.S.	1	1.14	1.23	1.08	2.34	1.20	15.65	1.37	1.77	1.73
time (sec)	N/A	0.506	0.674	0.535	0.112	0.103	4.661	0.124	0.163	26.423

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	56	20	26	26	48	50	19	40
N.S.	1	1.00	1.70	0.61	0.79	0.79	1.45	1.52	0.58	1.21
time (sec)	N/A	0.183	0.044	0.070	0.109	0.088	0.361	0.126	0.160	25.859

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	63	91	63	93	59	384	96	64	83
N.S.	1	1.09	1.57	1.09	1.60	1.02	6.62	1.66	1.10	1.43
time (sec)	N/A	0.261	0.092	0.099	0.112	0.101	0.913	0.119	0.167	0.002

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	93	113	91	173	94	915	122	131	112
N.S.	1	1.12	1.36	1.10	2.08	1.13	11.02	1.47	1.58	1.35
time (sec)	N/A	0.365	0.411	0.243	0.111	0.090	1.951	0.123	0.172	25.999

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	123	133	117	253	130	1690	148	191	187
N.S.	1	1.14	1.23	1.08	2.34	1.20	15.65	1.37	1.77	1.73
time (sec)	N/A	0.486	0.123	0.212	0.111	0.107	4.479	0.123	0.170	0.002

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	56	20	26	26	49	49	19	40
N.S.	1	1.00	1.81	0.65	0.84	0.84	1.58	1.58	0.61	1.29
time (sec)	N/A	0.183	0.046	0.086	0.115	0.121	0.381	0.128	0.178	25.855

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	61	91	63	93	59	389	95	64	83
N.S.	1	1.09	1.62	1.12	1.66	1.05	6.95	1.70	1.14	1.48
time (sec)	N/A	0.254	0.114	0.141	0.110	0.097	0.906	0.124	0.168	0.002

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	91	114	91	173	94	921	121	131	112
N.S.	1	1.12	1.41	1.12	2.14	1.16	11.37	1.49	1.62	1.38
time (sec)	N/A	0.359	0.660	0.331	0.109	0.107	1.960	0.129	0.177	26.054

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	121	133	117	253	130	1695	147	191	187
N.S.	1	1.14	1.25	1.10	2.39	1.23	15.99	1.39	1.80	1.76
time (sec)	N/A	0.485	0.424	0.361	0.128	0.110	4.497	0.122	0.171	0.002

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	42	63	36	49	46	42	36	34	19
N.S.	1	0.67	1.00	0.57	0.78	0.73	0.67	0.57	0.54	0.30
time (sec)	N/A	0.216	0.053	0.115	0.026	0.082	0.321	0.121	0.166	26.021

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	70	126	68	115	88	466	81	98	64
N.S.	1	0.80	1.43	0.77	1.31	1.00	5.30	0.92	1.11	0.73
time (sec)	N/A	0.296	0.288	0.263	0.028	0.112	0.719	0.128	0.180	25.972

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	100	180	100	195	133	1227	107	187	115
N.S.	1	0.88	1.59	0.88	1.73	1.18	10.86	0.95	1.65	1.02
time (sec)	N/A	0.417	0.576	0.302	0.034	0.084	1.444	0.137	0.166	27.202

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	130	235	132	275	181	2356	133	269	168
N.S.	1	0.94	1.70	0.96	1.99	1.31	17.07	0.96	1.95	1.22
time (sec)	N/A	0.556	1.016	0.441	0.036	0.094	3.219	0.132	0.168	28.666

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	44	65	36	49	46	42	36	34	19
N.S.	1	0.68	1.00	0.55	0.75	0.71	0.65	0.55	0.52	0.29
time (sec)	N/A	0.222	0.053	0.123	0.029	0.142	0.325	0.132	0.172	26.030

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	72	130	68	115	88	462	81	98	64
N.S.	1	0.80	1.44	0.76	1.28	0.98	5.13	0.90	1.09	0.71
time (sec)	N/A	0.293	0.349	0.269	0.030	0.080	0.761	0.126	0.192	25.983

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	102	184	100	195	133	1224	107	187	116
N.S.	1	0.89	1.60	0.87	1.70	1.16	10.64	0.93	1.63	1.01
time (sec)	N/A	0.424	0.826	0.312	0.034	0.087	1.414	0.126	0.178	27.197

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	132	241	132	275	181	2353	133	269	168
N.S.	1	0.94	1.72	0.94	1.96	1.29	16.81	0.95	1.92	1.20
time (sec)	N/A	0.549	1.110	0.451	0.036	0.092	3.248	0.126	0.169	28.212

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	44	65	36	49	46	42	36	34	19
N.S.	1	0.68	1.00	0.55	0.75	0.71	0.65	0.55	0.52	0.29
time (sec)	N/A	0.217	0.044	0.066	0.032	0.085	0.323	0.127	0.171	25.426

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	72	130	68	115	88	462	81	98	64
N.S.	1	0.80	1.44	0.76	1.28	0.98	5.13	0.90	1.09	0.71
time (sec)	N/A	0.296	0.132	0.069	0.030	0.082	0.755	0.135	0.168	0.003

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	102	183	100	195	133	1224	107	187	115
N.S.	1	0.89	1.59	0.87	1.70	1.16	10.64	0.93	1.63	1.00
time (sec)	N/A	0.407	0.618	0.269	0.036	0.144	1.481	0.132	0.162	26.711

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	132	241	132	275	181	2353	133	269	168
N.S.	1	0.94	1.72	0.94	1.96	1.29	16.81	0.95	1.92	1.20
time (sec)	N/A	0.536	0.014	0.124	0.039	0.103	3.240	0.125	0.171	0.002

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	42	63	36	49	46	44	36	34	19
N.S.	1	0.67	1.00	0.57	0.78	0.73	0.70	0.57	0.54	0.30
time (sec)	N/A	0.214	0.045	0.083	0.028	0.102	0.324	0.127	0.234	25.469

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	70	126	68	115	88	468	81	98	64
N.S.	1	0.80	1.43	0.77	1.31	1.00	5.32	0.92	1.11	0.73
time (sec)	N/A	0.291	0.189	0.125	0.034	0.092	0.731	0.129	0.276	0.002

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	100	179	100	195	133	1229	107	187	116
N.S.	1	0.88	1.58	0.88	1.73	1.18	10.88	0.95	1.65	1.03
time (sec)	N/A	0.413	0.769	0.296	0.037	0.094	1.434	0.130	0.170	26.870

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	130	235	132	275	181	2358	133	269	168
N.S.	1	0.94	1.70	0.96	1.99	1.31	17.09	0.96	1.95	1.22
time (sec)	N/A	0.524	0.898	0.146	0.037	0.116	3.236	0.129	0.169	0.002

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	256	267	220	1040	0	479	0	0	98	0
N.S.	1	1.04	0.86	4.06	0.00	1.87	0.00	0.00	0.38	0.00
time (sec)	N/A	1.409	1.240	1.506	0.000	0.115	0.000	0.000	0.204	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	207	213	185	890	0	435	0	0	68	0
N.S.	1	1.03	0.89	4.30	0.00	2.10	0.00	0.00	0.33	0.00
time (sec)	N/A	0.993	1.155	0.587	0.000	0.121	0.000	0.000	0.168	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	168	142	656	0	394	0	0	38	0
N.S.	1	1.01	0.85	3.93	0.00	2.36	0.00	0.00	0.23	0.00
time (sec)	N/A	0.734	0.953	0.418	0.000	0.104	0.000	0.000	0.182	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	61	239	0	351	0	0	13	55
N.S.	1	1.00	0.98	3.85	0.00	5.66	0.00	0.00	0.21	0.89
time (sec)	N/A	0.280	2.580	0.804	0.000	0.095	0.000	0.000	0.169	25.708

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	61	126	0	143	0	0	26	55
N.S.	1	1.00	0.98	2.03	0.00	2.31	0.00	0.00	0.42	0.89
time (sec)	N/A	0.286	0.133	0.122	0.000	0.095	0.000	0.000	0.163	25.849

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	111	87	443	0	460	0	0	42	0
N.S.	1	1.00	0.78	3.99	0.00	4.14	0.00	0.00	0.38	0.00
time (sec)	N/A	0.422	0.379	0.181	0.000	0.139	0.000	0.000	0.188	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	231	241	166	495	0	687	0	0	58	0
N.S.	1	1.04	0.72	2.14	0.00	2.97	0.00	0.00	0.25	0.00
time (sec)	N/A	1.071	1.149	0.237	0.000	0.163	0.000	0.000	0.172	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	318	198	582	0	985	0	0	74	0
N.S.	1	1.09	0.68	1.99	0.00	3.37	0.00	0.00	0.25	0.00
time (sec)	N/A	1.471	1.555	0.313	0.000	0.123	0.000	0.000	0.172	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	109	244	0	0	0	0	0	40	0
N.S.	1	1.03	2.30	0.00	0.00	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	0.260	1.881	0.000	0.000	0.000	0.000	0.000	0.180	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	118	0	0	0	0	0	14	0
N.S.	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	0.13	0.00
time (sec)	N/A	0.254	0.324	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	118	0	0	0	0	0	14	0
N.S.	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	0.13	0.00
time (sec)	N/A	0.261	0.312	0.000	0.000	0.000	0.000	0.000	0.178	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	118	0	0	0	0	0	14	0
N.S.	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	0.13	0.00
time (sec)	N/A	0.256	0.296	0.000	0.000	0.000	0.000	0.000	0.163	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	116	0	0	0	0	0	14	0
N.S.	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	0.13	0.00
time (sec)	N/A	0.255	0.264	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	106	111	262	0	0	0	0	0	39	0
N.S.	1	1.05	2.47	0.00	0.00	0.00	0.00	0.00	0.37	0.00
time (sec)	N/A	0.255	2.097	0.000	0.000	0.000	0.000	0.000	0.166	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	104	104	120	0	0	0	0	0	14	0
N.S.	1	1.00	1.15	0.00	0.00	0.00	0.00	0.00	0.13	0.00
time (sec)	N/A	0.253	0.360	0.000	0.000	0.000	0.000	0.000	0.170	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	83	0	0	0	0	0	14	0
N.S.	1	1.00	1.15	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.233	0.195	0.000	0.000	0.000	0.000	0.000	0.175	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	84	0	0	0	0	0	14	0
N.S.	1	1.00	1.22	0.00	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.229	0.192	0.000	0.000	0.000	0.000	0.000	0.221	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	72	64	99	0	0	0	0	0	14	0
N.S.	1	0.89	1.38	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.224	0.157	0.000	0.000	0.000	0.000	0.000	0.258	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	69	96	0	0	0	0	0	14	0
N.S.	1	1.03	1.43	0.00	0.00	0.00	0.00	0.00	0.21	0.00
time (sec)	N/A	0.226	0.179	0.000	0.000	0.000	0.000	0.000	0.164	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	83	0	0	0	0	0	14	0
N.S.	1	1.00	1.24	0.00	0.00	0.00	0.00	0.00	0.21	0.00
time (sec)	N/A	0.229	0.185	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	84	0	0	0	0	0	14	0
N.S.	1	1.00	1.31	0.00	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.227	0.200	0.000	0.000	0.000	0.000	0.000	0.189	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	93	95	95	0	0	0	0	0	14	0
N.S.	1	1.02	1.02	0.00	0.00	0.00	0.00	0.00	0.15	0.00
time (sec)	N/A	0.249	0.170	0.000	0.000	0.000	0.000	0.000	0.165	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	98	110	100	0	0	0	0	0	14	0
N.S.	1	1.12	1.02	0.00	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.260	0.181	0.000	0.000	0.000	0.000	0.000	0.174	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [76] had the largest ratio of [1.2857099999999991]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.00	12	0.250
2	A	2	2	1.00	12	0.167
3	A	1	1	1.00	10	0.100
4	A	2	2	1.00	12	0.167
5	A	4	4	0.98	12	0.333
6	A	6	6	1.05	12	0.500
7	A	3	3	1.00	13	0.231
8	A	2	2	1.00	13	0.154
9	A	1	1	1.00	11	0.091
10	A	2	2	1.00	13	0.154
11	A	4	4	0.98	13	0.308
12	A	6	6	1.05	13	0.462
13	A	8	8	1.07	14	0.571
14	A	6	6	1.04	14	0.429
15	A	4	4	1.00	14	0.286
16	A	2	2	1.00	14	0.143
17	A	4	3	1.00	14	0.214
18	A	6	5	1.00	14	0.357
19	A	8	7	1.05	14	0.500
20	A	8	8	1.07	15	0.533
21	A	6	6	1.04	15	0.400

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	4	4	1.00	15	0.267
23	A	2	2	1.00	15	0.133
24	A	4	3	1.00	15	0.200
25	A	6	5	1.00	15	0.333
26	A	8	7	1.05	15	0.467
27	A	8	8	1.06	16	0.500
28	A	6	6	1.04	16	0.375
29	A	4	4	1.00	16	0.250
30	A	2	2	1.00	16	0.125
31	A	4	3	1.00	16	0.188
32	A	6	5	1.00	16	0.312
33	A	8	7	1.04	16	0.438
34	A	4	4	1.00	14	0.286
35	A	4	4	1.00	14	0.286
36	A	4	4	1.00	14	0.286
37	A	4	4	1.00	14	0.286
38	A	4	4	1.00	14	0.286
39	A	4	4	1.00	14	0.286
40	A	4	4	1.00	12	0.333
41	A	4	4	1.00	13	0.308
42	A	2	2	1.00	12	0.167
43	A	2	2	1.00	12	0.167
44	A	2	2	1.00	12	0.167
45	A	5	5	1.09	12	0.417
46	A	8	8	1.12	12	0.667
47	A	11	11	1.14	12	0.917
48	A	2	2	1.00	12	0.167
49	A	5	5	1.09	12	0.417
50	A	8	8	1.12	12	0.667
51	A	11	11	1.14	12	0.917
52	A	2	2	1.00	12	0.167
53	A	5	5	1.09	12	0.417

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	7	7	1.12	12	0.583
55	A	11	11	1.14	12	0.917
56	A	2	2	1.00	12	0.167
57	A	5	5	1.09	12	0.417
58	A	7	7	1.12	12	0.583
59	A	11	11	1.14	12	0.917
60	A	5	4	0.67	12	0.333
61	A	8	7	0.80	12	0.583
62	A	11	10	0.88	12	0.833
63	A	14	13	0.94	12	1.083
64	A	5	4	0.68	12	0.333
65	A	8	7	0.80	12	0.583
66	A	11	10	0.89	12	0.833
67	A	14	13	0.94	12	1.083
68	A	5	4	0.68	12	0.333
69	A	8	7	0.80	12	0.583
70	A	10	9	0.89	12	0.750
71	A	14	13	0.94	12	1.083
72	A	5	4	0.67	12	0.333
73	A	8	7	0.80	12	0.583
74	A	10	9	0.88	12	0.750
75	A	14	13	0.94	12	1.083
76	A	18	18	1.04	14	1.286
77	A	15	15	1.03	14	1.071
78	A	12	12	1.01	14	0.857
79	A	4	4	1.00	14	0.286
80	A	4	4	1.00	14	0.286
81	A	7	7	1.00	14	0.500
82	A	15	15	1.04	14	1.071
83	A	18	18	1.09	14	1.286
84	A	5	4	1.03	14	0.286
85	A	5	4	1.00	14	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	5	4	1.00	14	0.286
87	A	5	4	1.00	14	0.286
88	A	5	4	1.00	14	0.286
89	A	5	4	1.05	14	0.286
90	A	5	4	1.00	12	0.333
91	A	4	3	1.00	12	0.250
92	A	4	3	1.00	12	0.250
93	A	4	3	0.89	12	0.250
94	A	4	3	1.03	12	0.250
95	A	4	3	1.00	12	0.250
96	A	4	3	1.00	12	0.250
97	A	5	4	1.02	12	0.333
98	A	6	5	1.12	12	0.417

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + a \sin(c + dx))^3 dx$	62
3.2	$\int (a + a \sin(c + dx))^2 dx$	68
3.3	$\int (a + a \sin(c + dx)) dx$	73
3.4	$\int \frac{1}{a + a \sin(c + dx)} dx$	78
3.5	$\int \frac{1}{(a + a \sin(c + dx))^2} dx$	83
3.6	$\int \frac{1}{(a + a \sin(c + dx))^3} dx$	89
3.7	$\int (a - a \sin(c + dx))^3 dx$	96
3.8	$\int (a - a \sin(c + dx))^2 dx$	102
3.9	$\int (a - a \sin(c + dx)) dx$	107
3.10	$\int \frac{1}{a - a \sin(c + dx)} dx$	112
3.11	$\int \frac{1}{(a - a \sin(c + dx))^2} dx$	117
3.12	$\int \frac{1}{(a - a \sin(c + dx))^3} dx$	123
3.13	$\int (a + a \sin(c + dx))^{7/2} dx$	130
3.14	$\int (a + a \sin(c + dx))^{5/2} dx$	137
3.15	$\int (a + a \sin(c + dx))^{3/2} dx$	143
3.16	$\int \sqrt{a + a \sin(c + dx)} dx$	149
3.17	$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx$	154
3.18	$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx$	160
3.19	$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx$	166
3.20	$\int (a - a \sin(c + dx))^{7/2} dx$	173
3.21	$\int (a - a \sin(c + dx))^{5/2} dx$	180
3.22	$\int (a - a \sin(c + dx))^{3/2} dx$	186
3.23	$\int \sqrt{a - a \sin(c + dx)} dx$	192
3.24	$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx$	197
3.25	$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx$	203
3.26	$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx$	209

3.27	$\int (-a + a \sin(c + dx))^{7/2} dx$	216
3.28	$\int (-a + a \sin(c + dx))^{5/2} dx$	223
3.29	$\int (-a + a \sin(c + dx))^{3/2} dx$	229
3.30	$\int \sqrt{-a + a \sin(c + dx)} dx$	235
3.31	$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx$	240
3.32	$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx$	246
3.33	$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx$	252
3.34	$\int (a + a \sin(c + dx))^{4/3} dx$	259
3.35	$\int (a + a \sin(c + dx))^{2/3} dx$	264
3.36	$\int \sqrt[3]{a + a \sin(c + dx)} dx$	269
3.37	$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx$	274
3.38	$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx$	279
3.39	$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx$	284
3.40	$\int (a + a \sin(c + dx))^n dx$	289
3.41	$\int (a - a \sin(c + dx))^n dx$	294
3.42	$\int (2 + 2 \sin(c + dx))^n dx$	299
3.43	$\int (2 - 2 \sin(c + dx))^n dx$	304
3.44	$\int \frac{1}{5 + 3 \sin(c + dx)} dx$	309
3.45	$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx$	314
3.46	$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx$	321
3.47	$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx$	329
3.48	$\int \frac{1}{5 - 3 \sin(c + dx)} dx$	338
3.49	$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx$	343
3.50	$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx$	350
3.51	$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx$	358
3.52	$\int \frac{1}{-5 + 3 \sin(c + dx)} dx$	367
3.53	$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx$	372
3.54	$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx$	379
3.55	$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx$	387
3.56	$\int \frac{1}{-5 - 3 \sin(c + dx)} dx$	396
3.57	$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx$	401
3.58	$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx$	408
3.59	$\int \frac{1}{(-5 - 3 \sin(c + dx))^4} dx$	416
3.60	$\int \frac{1}{3 + 5 \sin(c + dx)} dx$	425
3.61	$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx$	431
3.62	$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx$	438

3.63	$\int \frac{1}{(3+5 \sin(c+dx))^4} dx$	447
3.64	$\int \frac{1}{3-5 \sin(c+dx)} dx$	457
3.65	$\int \frac{1}{(3-5 \sin(c+dx))^2} dx$	463
3.66	$\int \frac{1}{(3-5 \sin(c+dx))^3} dx$	470
3.67	$\int \frac{1}{(3-5 \sin(c+dx))^4} dx$	479
3.68	$\int \frac{1}{-3+5 \sin(c+dx)} dx$	489
3.69	$\int \frac{1}{(-3+5 \sin(c+dx))^2} dx$	495
3.70	$\int \frac{1}{(-3+5 \sin(c+dx))^3} dx$	502
3.71	$\int \frac{1}{(-3+5 \sin(c+dx))^4} dx$	511
3.72	$\int \frac{1}{-3-5 \sin(c+dx)} dx$	521
3.73	$\int \frac{1}{(-3-5 \sin(c+dx))^2} dx$	527
3.74	$\int \frac{1}{(-3-5 \sin(c+dx))^3} dx$	534
3.75	$\int \frac{1}{(-3-5 \sin(c+dx))^4} dx$	543
3.76	$\int (a + b \sin(c + dx))^{7/2} dx$	553
3.77	$\int (a + b \sin(c + dx))^{5/2} dx$	564
3.78	$\int (a + b \sin(c + dx))^{3/2} dx$	574
3.79	$\int \sqrt{a + b \sin(c + dx)} dx$	583
3.80	$\int \frac{1}{\sqrt{a+b \sin(c+dx)}} dx$	589
3.81	$\int \frac{1}{(a+b \sin(c+dx))^{3/2}} dx$	595
3.82	$\int \frac{1}{(a+b \sin(c+dx))^{5/2}} dx$	602
3.83	$\int \frac{1}{(a+b \sin(c+dx))^{7/2}} dx$	612
3.84	$\int (a + b \sin(c + dx))^{4/3} dx$	623
3.85	$\int (a + b \sin(c + dx))^{2/3} dx$	629
3.86	$\int \sqrt[3]{a + b \sin(c + dx)} dx$	635
3.87	$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx$	641
3.88	$\int \frac{1}{(a+b \sin(c+dx))^{2/3}} dx$	647
3.89	$\int \frac{1}{(a+b \sin(c+dx))^{4/3}} dx$	653
3.90	$\int (a + b \sin(c + dx))^n dx$	659
3.91	$\int (3 + 4 \sin(c + dx))^n dx$	665
3.92	$\int (3 - 4 \sin(c + dx))^n dx$	670
3.93	$\int (4 + 3 \sin(c + dx))^n dx$	675
3.94	$\int (4 - 3 \sin(c + dx))^n dx$	680
3.95	$\int (-3 + 4 \sin(c + dx))^n dx$	685
3.96	$\int (-3 - 4 \sin(c + dx))^n dx$	690
3.97	$\int (-4 + 3 \sin(c + dx))^n dx$	695
3.98	$\int (-4 - 3 \sin(c + dx))^n dx$	701

3.1 $\int (a + a \sin(c + dx))^3 dx$

Optimal result	62
Mathematica [A] (verified)	62
Rubi [A] (verified)	63
Maple [A] (verified)	64
Fricas [A] (verification not implemented)	65
Sympy [B] (verification not implemented)	65
Maxima [A] (verification not implemented)	66
Giac [A] (verification not implemented)	66
Mupad [B] (verification not implemented)	67
Reduce [B] (verification not implemented)	67

Optimal result

Integrand size = 12, antiderivative size = 63

$$\int (a + a \sin(c + dx))^3 dx = \frac{5a^3x}{2} - \frac{4a^3 \cos(c + dx)}{d} + \frac{a^3 \cos^3(c + dx)}{3d} - \frac{3a^3 \cos(c + dx) \sin(c + dx)}{2d}$$

output `5/2*a^3*x-4*a^3*cos(d*x+c)/d+1/3*a^3*cos(d*x+c)^3/d-3/2*a^3*cos(d*x+c)*sin(d*x+c)/d`

Mathematica [A] (verified)

Time = 3.85 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int (a + a \sin(c + dx))^3 dx = \frac{a^3(30c + 30dx - 45 \cos(c + dx) + \cos(3(c + dx)) - 9 \sin(2(c + dx)))}{12d}$$

input `Integrate[(a + a*Sin[c + d*x])^3,x]`

output

$$(a^3(30c + 30dx - 45\cos[c + dx] + \cos[3(c + dx)] - 9\sin[2(c + dx)]))/(12d)$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3124, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sin(c + dx) + a)^3 dx \\ & \quad \downarrow \text{3042} \\ & \int (a \sin(c + dx) + a)^3 dx \\ & \quad \downarrow \text{3124} \\ & \int (a^3 \sin^3(c + dx) + 3a^3 \sin^2(c + dx) + 3a^3 \sin(c + dx) + a^3) dx \\ & \quad \downarrow \text{2009} \\ & \frac{a^3 \cos^3(c + dx)}{3d} - \frac{4a^3 \cos(c + dx)}{d} - \frac{3a^3 \sin(c + dx) \cos(c + dx)}{2d} + \frac{5a^3 x}{2} \end{aligned}$$

input

$$\text{Int}[(a + a*\text{Sin}[c + d*x])^3, x]$$

output

$$(5a^3x)/2 - (4a^3\cos[c + dx])/d + (a^3\cos[c + dx]^3)/(3d) - (3a^3\cos[c + dx]*\sin[c + dx])/(2d)$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3124 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrig[(a + b*sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 7.24 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.68

method	result
parallelsch	$\frac{a^3(30dx-45\cos(dx+c)+\cos(3dx+3c)-9\sin(2dx+2c)-44)}{12d}$
risch	$\frac{5a^3x}{2} - \frac{15a^3\cos(dx+c)}{4d} + \frac{a^3\cos(3dx+3c)}{12d} - \frac{3a^3\sin(2dx+2c)}{4d}$
derivativdivides	$-\frac{a^3(2+\sin(dx+c)^2)\cos(dx+c)}{3} + 3a^3\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) - 3a^3\cos(dx+c) + a^3(dx+c)$
default	$-\frac{a^3(2+\sin(dx+c)^2)\cos(dx+c)}{3} + 3a^3\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) - 3a^3\cos(dx+c) + a^3(dx+c)$
parts	$a^3x - \frac{a^3(2+\sin(dx+c)^2)\cos(dx+c)}{3d} - \frac{3a^3\cos(dx+c)}{d} + \frac{3a^3\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d}$
norman	$\frac{5a^3x}{2} - \frac{22a^3}{3d} - \frac{3a^3\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{3a^3\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} + \frac{15a^3x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{2} + \frac{15a^3x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{2} + \frac{5a^3x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6}{2} - \frac{6a^3\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{\left(1+\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)^3}$

input `int((a+a*sin(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/12*a^3*(30*d*x-45*cos(d*x+c)+cos(3*d*x+3*c)-9*sin(2*d*x+2*c)-44)/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.86

$$\int (a + a \sin(c + dx))^3 dx$$

$$= \frac{2 a^3 \cos(dx + c)^3 + 15 a^3 dx - 9 a^3 \cos(dx + c) \sin(dx + c) - 24 a^3 \cos(dx + c)}{6 d}$$

input `integrate((a+a*sin(d*x+c))^3,x, algorithm="fricas")`

output `1/6*(2*a^3*cos(d*x + c)^3 + 15*a^3*d*x - 9*a^3*cos(d*x + c)*sin(d*x + c) - 24*a^3*cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. 2(58) = 116.

Time = 0.12 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.92

$$\int (a + a \sin(c + dx))^3 dx$$

$$= \left\{ \begin{array}{l} \frac{3a^3 x \sin^2(c+dx)}{2} + \frac{3a^3 x \cos^2(c+dx)}{2} + a^3 x - \frac{a^3 \sin^2(c+dx) \cos(c+dx)}{d} - \frac{3a^3 \sin(c+dx) \cos(c+dx)}{2d} - \frac{2a^3 \cos^3(c+dx)}{3d} - \frac{3a^3 \cos^3(c+dx)}{3d} \\ x(a \sin(c) + a)^3 \end{array} \right.$$

input `integrate((a+a*sin(d*x+c))**3,x)`

output `Piecewise((3*a**3*x*sin(c + d*x)**2/2 + 3*a**3*x*cos(c + d*x)**2/2 + a**3*x - a**3*sin(c + d*x)**2*cos(c + d*x)/d - 3*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) - 2*a**3*cos(c + d*x)**3/(3*d) - 3*a**3*cos(c + d*x)/d, Ne(d, 0)), (x*(a*sin(c) + a)**3, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.14

$$\int (a + a \sin(c + dx))^3 dx = a^3 x + \frac{(\cos(dx + c)^3 - 3 \cos(dx + c))a^3}{3d} + \frac{3(2dx + 2c - \sin(2dx + 2c))a^3}{4d} - \frac{3a^3 \cos(dx + c)}{d}$$

input `integrate((a+a*sin(d*x+c))^3,x, algorithm="maxima")`output `a^3*x + 1/3*(cos(d*x + c)^3 - 3*cos(d*x + c))*a^3/d + 3/4*(2*d*x + 2*c - sin(2*d*x + 2*c))*a^3/d - 3*a^3*cos(d*x + c)/d`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.87

$$\int (a + a \sin(c + dx))^3 dx = \frac{5}{2} a^3 x + \frac{a^3 \cos(3dx + 3c)}{12d} - \frac{15 a^3 \cos(dx + c)}{4d} - \frac{3 a^3 \sin(2dx + 2c)}{4d}$$

input `integrate((a+a*sin(d*x+c))^3,x, algorithm="giac")`output `5/2*a^3*x + 1/12*a^3*cos(3*d*x + 3*c)/d - 15/4*a^3*cos(d*x + c)/d - 3/4*a^3*sin(2*d*x + 2*c)/d`

Mupad [B] (verification not implemented)

Time = 28.28 (sec) , antiderivative size = 156, normalized size of antiderivative = 2.48

$$\int (a + a \sin(c + dx))^3 dx = \frac{5a^3 x}{2} - \frac{\frac{5a^3(c+dx)}{2} - 3a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 - \frac{a^3(15c+15dx-44)}{6} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 \left(\frac{15a^3(c+dx)}{2} - \frac{a^3(45c+45dx-36)}{6}\right) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{15a^3(c+dx)}{2} - \frac{a^3(45c+45dx-36)}{6}\right) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \left(\frac{15a^3(c+dx)}{2} - \frac{a^3(45c+45dx-36)}{6}\right) + 3a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)^3}$$

input `int((a + a*sin(c + d*x))^3,x)`output `(5*a^3*x)/2 - ((5*a^3*(c + d*x))/2 - 3*a^3*tan(c/2 + (d*x)/2)^5 - (a^3*(15*c + 15*d*x - 44))/6 + tan(c/2 + (d*x)/2)^4*((15*a^3*(c + d*x))/2 - (a^3*(45*c + 45*d*x - 36))/6) + tan(c/2 + (d*x)/2)^2*((15*a^3*(c + d*x))/2 - (a^3*(45*c + 45*d*x - 96))/6) + 3*a^3*tan(c/2 + (d*x)/2))/(d*(tan(c/2 + (d*x)/2)^2 + 1)^3)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

$$\int (a + a \sin(c + dx))^3 dx = \frac{a^3(-2 \cos(dx + c) \sin(dx + c)^2 - 9 \cos(dx + c) \sin(dx + c) - 22 \cos(dx + c) + 15dx + 22)}{6d}$$

input `int((a+a*sin(d*x+c))^3,x)`output `(a**3*(- 2*cos(c + d*x)*sin(c + d*x)**2 - 9*cos(c + d*x)*sin(c + d*x) - 2*2*cos(c + d*x) + 15*d*x + 22))/(6*d)`

3.2 $\int (a + a \sin(c + dx))^2 dx$

Optimal result	68
Mathematica [A] (verified)	68
Rubi [A] (verified)	69
Maple [A] (verified)	70
Fricas [A] (verification not implemented)	70
Sympy [A] (verification not implemented)	71
Maxima [A] (verification not implemented)	71
Giac [A] (verification not implemented)	71
Mupad [B] (verification not implemented)	72
Reduce [B] (verification not implemented)	72

Optimal result

Integrand size = 12, antiderivative size = 45

$$\int (a + a \sin(c + dx))^2 dx = \frac{3a^2x}{2} - \frac{2a^2 \cos(c + dx)}{d} - \frac{a^2 \cos(c + dx) \sin(c + dx)}{2d}$$

output

```
3/2*a^2*x-2*a^2*cos(d*x+c)/d-1/2*a^2*cos(d*x+c)*sin(d*x+c)/d
```

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.76

$$\int (a + a \sin(c + dx))^2 dx = -\frac{a^2(-6(c + dx) + 8 \cos(c + dx) + \sin(2(c + dx)))}{4d}$$

input

```
Integrate[(a + a*Sin[c + d*x])^2,x]
```

output

```
-1/4*(a^2*(-6*(c + d*x) + 8*Cos[c + d*x] + Sin[2*(c + d*x)]))/d
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3123}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(c + dx) + a)^2 dx$$

↓ 3042

$$\int (a \sin(c + dx) + a)^2 dx$$

↓ 3123

$$-\frac{2a^2 \cos(c + dx)}{d} - \frac{a^2 \sin(c + dx) \cos(c + dx)}{2d} + \frac{3a^2 x}{2}$$

input `Int[(a + a*Sin[c + d*x])^2,x]`

output `(3*a^2*x)/2 - (2*a^2*Cos[c + d*x])/d - (a^2*Cos[c + d*x]*Sin[c + d*x])/(2*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3123 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^2, x_Symbol] := Simp[(2*a^2 + b^2)*(x/2), x] + (-Simp[2*a*b*(Cos[c + d*x]/d), x] - Simp[b^2*Cos[c + d*x]*(Sin[c + d*x]/(2*d)), x]) /; FreeQ[{a, b, c, d}, x]`

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.71

method	result
parallelrisch	$-\frac{a^2(-6dx+8\cos(dx+c)+\sin(2dx+2c)-8)}{4d}$
risch	$\frac{3a^2x}{2} - \frac{2a^2\cos(dx+c)}{d} - \frac{a^2\sin(2dx+2c)}{4d}$
parts	$a^2x + \frac{a^2\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} - \frac{2a^2\cos(dx+c)}{d}$
derivativedivides	$\frac{a^2\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) - 2\cos(dx+c)a^2 + a^2(dx+c)}{d}$
default	$\frac{a^2\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) - 2\cos(dx+c)a^2 + a^2(dx+c)}{d}$
norman	$\frac{\frac{a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{d} + \frac{4a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{d} + \frac{3a^2x}{2} - \frac{a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + 3a^2x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{3a^2x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{2} + \frac{4a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$
orering	$x(a + a\sin(dx+c))^2 - \frac{5(a+a\sin(dx+c))a\cos(dx+c)}{2d} + \frac{5x(2a^2d^2\cos(dx+c)^2 - 2(a+a\sin(dx+c))ad^2\sin(dx+c))}{4d^2}$

input `int((a+a*sin(d*x+c))^2,x,method=_RETURNVERBOSE)`output `-1/4*a^2*(-6*d*x+8*cos(d*x+c)+sin(2*d*x+2*c)-8)/d`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.91

$$\int (a + a\sin(c + dx))^2 dx = \frac{3a^2dx - a^2\cos(dx+c)\sin(dx+c) - 4a^2\cos(dx+c)}{2d}$$

input `integrate((a+a*sin(d*x+c))^2,x, algorithm="fricas")`output `1/2*(3*a^2*d*x - a^2*cos(d*x + c)*sin(d*x + c) - 4*a^2*cos(d*x + c))/d`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.73

$$\int (a + a \sin(c + dx))^2 dx = \begin{cases} \frac{a^2 x \sin^2(c+dx)}{2} + \frac{a^2 x \cos^2(c+dx)}{2} + a^2 x - \frac{a^2 \sin(c+dx) \cos(c+dx)}{2d} - \frac{2a^2 \cos(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \sin(c) + a)^2 & \text{otherwise} \end{cases}$$

input `integrate((a+a*sin(d*x+c))**2,x)`output `Piecewise((a**2*x*sin(c + d*x)**2/2 + a**2*x*cos(c + d*x)**2/2 + a**2*x - a**2*sin(c + d*x)*cos(c + d*x)/(2*d) - 2*a**2*cos(c + d*x)/d, Ne(d, 0)), (x*(a*sin(c) + a)**2, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int (a + a \sin(c + dx))^2 dx = a^2 x + \frac{(2 dx + 2 c - \sin(2 dx + 2 c))a^2}{4 d} - \frac{2 a^2 \cos(dx + c)}{d}$$

input `integrate((a+a*sin(d*x+c))^2,x, algorithm="maxima")`output `a^2*x + 1/4*(2*d*x + 2*c - sin(2*d*x + 2*c))*a^2/d - 2*a^2*cos(d*x + c)/d`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.84

$$\int (a + a \sin(c + dx))^2 dx = \frac{3}{2} a^2 x - \frac{2 a^2 \cos(dx + c)}{d} - \frac{a^2 \sin(2 dx + 2 c)}{4 d}$$

input `integrate((a+a*sin(d*x+c))^2,x, algorithm="giac")`

output $3/2*a^2*x - 2*a^2*\cos(d*x + c)/d - 1/4*a^2*\sin(2*d*x + 2*c)/d$

Mupad [B] (verification not implemented)

Time = 26.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.73

$$\int (a + a \sin(c + dx))^2 dx = \frac{3a^2 x}{2} - \frac{a^2 \left(\frac{3c}{2} + \frac{3dx}{2} \right) - a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 - a^2 \left(\frac{3c}{2} + \frac{3dx}{2} - 4 \right) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \left(2a^2 \left(\frac{3c}{2} + \frac{3dx}{2} \right) - a^2 (3c + 3dx) \right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)^2}$$

input $\text{int}((a + a*\sin(c + d*x))^2,x)$

output $(3*a^2*x)/2 - (a^2*((3*c)/2 + (3*d*x)/2) - a^2*\tan(c/2 + (d*x)/2)^3 - a^2*((3*c)/2 + (3*d*x)/2 - 4) + \tan(c/2 + (d*x)/2)^2*(2*a^2*((3*c)/2 + (3*d*x)/2) - a^2*(3*c + 3*d*x - 4)) + a^2*\tan(c/2 + (d*x)/2))/(d*(\tan(c/2 + (d*x)/2)^2 + 1)^2)$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int (a + a \sin(c + dx))^2 dx = \frac{a^2(-\cos(dx + c)\sin(dx + c) - 4\cos(dx + c) + 3dx + 4)}{2d}$$

input $\text{int}((a+a*\sin(d*x+c))^2,x)$

output $(a**2*(-\cos(c + d*x)*\sin(c + d*x) - 4*\cos(c + d*x) + 3*d*x + 4))/(2*d)$

3.3 $\int (a + a \sin(c + dx)) dx$

Optimal result	73
Mathematica [A] (verified)	73
Rubi [A] (verified)	74
Maple [A] (verified)	75
Fricas [A] (verification not implemented)	75
Sympy [A] (verification not implemented)	76
Maxima [A] (verification not implemented)	76
Giac [A] (verification not implemented)	76
Mupad [B] (verification not implemented)	77
Reduce [B] (verification not implemented)	77

Optimal result

Integrand size = 10, antiderivative size = 16

$$\int (a + a \sin(c + dx)) dx = ax - \frac{a \cos(c + dx)}{d}$$

output

```
a*x-a*cos(d*x+c)/d
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int (a + a \sin(c + dx)) dx = ax - \frac{a \cos(c) \cos(dx)}{d} + \frac{a \sin(c) \sin(dx)}{d}$$

input

```
Integrate[a + a*Sin[c + d*x],x]
```

output

```
a*x - (a*Cos[c]*Cos[d*x])/d + (a*Sin[c]*Sin[d*x])/d
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(c + dx) + a) dx$$

$$\downarrow \text{2009}$$

$$ax - \frac{a \cos(c + dx)}{d}$$

input `Int[a + a*Sin[c + d*x],x]`

output `a*x - (a*Cos[c + d*x])/d`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
default	$ax - \frac{a \cos(dx+c)}{d}$	17
risch	$ax - \frac{a \cos(dx+c)}{d}$	17
parts	$ax - \frac{a \cos(dx+c)}{d}$	17
parallelrisc	$\frac{a(-1-\cos(dx+c))}{d} + ax$	20
derivativdivides	$\frac{(dx+c)a - a \cos(dx+c)}{d}$	22
orering	$x(a + a \sin(dx + c)) - \frac{a \cos(dx+c)}{d} - xa \sin(dx + c)$	36
norman	$\frac{ax + ax \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{2a \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d}}{1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}$	52

input `int(a+a*sin(d*x+c),x,method=_RETURNVERBOSE)`output `a*x-a*cos(d*x+c)/d`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (a + a \sin(c + dx)) dx = \frac{adx - a \cos(dx + c)}{d}$$

input `integrate(a+a*sin(d*x+c),x, algorithm="fricas")`output `(a*d*x - a*cos(d*x + c))/d`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int (a + a \sin(c + dx)) dx = ax + a \begin{cases} -\frac{\cos(c+dx)}{d} & \text{for } d \neq 0 \\ x \sin(c) & \text{otherwise} \end{cases}$$

input `integrate(a+a*sin(d*x+c),x)`output `a*x + a*Piecewise((-cos(c + d*x)/d, Ne(d, 0)), (x*sin(c), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (a + a \sin(c + dx)) dx = ax - \frac{a \cos(dx + c)}{d}$$

input `integrate(a+a*sin(d*x+c),x, algorithm="maxima")`output `a*x - a*cos(d*x + c)/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (a + a \sin(c + dx)) dx = ax - \frac{a \cos(dx + c)}{d}$$

input `integrate(a+a*sin(d*x+c),x, algorithm="giac")`output `a*x - a*cos(d*x + c)/d`

Mupad [B] (verification not implemented)

Time = 25.71 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.56

$$\int (a + a \sin(c + dx)) dx = ax - \frac{2a}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)}$$

input `int(a + a*sin(c + d*x),x)`

output `a*x - (2*a)/(d*(tan(c/2 + (d*x)/2)^2 + 1))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int (a + a \sin(c + dx)) dx = \frac{a(-\cos(dx + c) + dx)}{d}$$

input `int(a+a*sin(d*x+c),x)`

output `(a*(- cos(c + d*x) + d*x))/d`

3.4 $\int \frac{1}{a+a \sin(c+dx)} dx$

Optimal result	78
Mathematica [B] (verified)	78
Rubi [A] (verified)	79
Maple [A] (verified)	80
Fricas [A] (verification not implemented)	80
Sympy [A] (verification not implemented)	81
Maxima [A] (verification not implemented)	81
Giac [A] (verification not implemented)	81
Mupad [B] (verification not implemented)	82
Reduce [B] (verification not implemented)	82

Optimal result

Integrand size = 12, antiderivative size = 23

$$\int \frac{1}{a + a \sin(c + dx)} dx = -\frac{\cos(c + dx)}{d(a + a \sin(c + dx))}$$

output `-cos(d*x+c)/d/(a+a*sin(d*x+c))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 48 vs. 2(23) = 46.

Time = 0.27 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.09

$$\int \frac{1}{a + a \sin(c + dx)} dx = \frac{2 \sin\left(\frac{1}{2}(c + dx)\right) \left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)}{d(a + a \sin(c + dx))}$$

input `Integrate[(a + a*Sin[c + d*x])^(-1),x]`

output `(2*Sin[(c + d*x)/2]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(d*(a + a*Sin[c + d*x]))`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a \sin(c + dx) + a} dx$$

↓ 3042

$$\int \frac{1}{a \sin(c + dx) + a} dx$$

↓ 3127

$$\frac{\cos(c + dx)}{d(a \sin(c + dx) + a)}$$

input `Int[(a + a*Sin[c + d*x])^(-1),x]`

output `-(Cos[c + d*x]/(d*(a + a*Sin[c + d*x])))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$-\frac{2}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}$	22
default	$-\frac{2}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}$	22
risch	$-\frac{2}{da\left(e^{i(dx+c)}+i\right)}$	23
norman	$\frac{2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}$	31
parallelrisch	$\frac{2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)}$	31

input `int(1/(a+a*sin(d*x+c)),x,method=_RETURNVERBOSE)`output `-2/d/a/(tan(1/2*d*x+1/2*c)+1)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \frac{1}{a + a \sin(c + dx)} dx = -\frac{\cos(dx + c) - \sin(dx + c) + 1}{ad \cos(dx + c) + ad \sin(dx + c) + ad}$$

input `integrate(1/(a+a*sin(d*x+c)),x, algorithm="fricas")`output `-(cos(d*x + c) - sin(d*x + c) + 1)/(a*d*cos(d*x + c) + a*d*sin(d*x + c) + a*d)`

Sympy [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{1}{a + a \sin(c + dx)} dx = \begin{cases} -\frac{2}{ad \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + ad} & \text{for } d \neq 0 \\ \frac{x}{a \sin(c) + a} & \text{otherwise} \end{cases}$$

input `integrate(1/(a+a*sin(d*x+c)),x)`output `Piecewise((-2/(a*d*tan(c/2 + d*x/2) + a*d), Ne(d, 0)), (x/(a*sin(c) + a), True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{1}{a + a \sin(c + dx)} dx = -\frac{2}{\left(a + \frac{a \sin(dx+c)}{\cos(dx+c)+1}\right) d}$$

input `integrate(1/(a+a*sin(d*x+c)),x, algorithm="maxima")`output `-2/((a + a*sin(d*x + c)/(cos(d*x + c) + 1))*d)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{1}{a + a \sin(c + dx)} dx = -\frac{2}{ad \left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right)}$$

input `integrate(1/(a+a*sin(d*x+c)),x, algorithm="giac")`output `-2/(a*d*(tan(1/2*d*x + 1/2*c) + 1))`

Mupad [B] (verification not implemented)

Time = 25.75 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{1}{a + a \sin(c + dx)} dx = -\frac{2}{ad \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1 \right)}$$

input `int(1/(a + a*sin(c + d*x)),x)`output `-2/(a*d*(tan(c/2 + (d*x)/2) + 1))`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \frac{1}{a + a \sin(c + dx)} dx = \frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1 \right)}$$

input `int(1/(a+a*sin(d*x+c)),x)`output `(2*tan((c + d*x)/2))/(a*d*(tan((c + d*x)/2) + 1))`

3.5 $\int \frac{1}{(a+a \sin(c+dx))^2} dx$

Optimal result	83
Mathematica [A] (verified)	83
Rubi [A] (verified)	84
Maple [C] (verified)	85
Fricas [A] (verification not implemented)	86
Sympy [B] (verification not implemented)	86
Maxima [B] (verification not implemented)	87
Giac [A] (verification not implemented)	87
Mupad [B] (verification not implemented)	88
Reduce [B] (verification not implemented)	88

Optimal result

Integrand size = 12, antiderivative size = 55

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx = -\frac{\cos(c + dx)}{3d(a + a \sin(c + dx))^2} - \frac{\cos(c + dx)}{3d(a^2 + a^2 \sin(c + dx))}$$

output `-1/3*cos(d*x+c)/d/(a+a*sin(d*x+c))^2-1/3*cos(d*x+c)/d/(a^2+a^2*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.98

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx = -\frac{-3 + 4 \cos(c + dx) + \cos(2(c + dx)) - 4 \sin(c + dx) + \sin(2(c + dx))}{6a^2d(1 + \sin(c + dx))^2}$$

input `Integrate[(a + a*Sin[c + d*x])^(-2),x]`

output `-1/6*(-3 + 4*Cos[c + d*x] + Cos[2*(c + d*x)] - 4*Sin[c + d*x] + Sin[2*(c + d*x)])/(a^2*d*(1 + Sin[c + d*x])^2)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sin(c + dx) + a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx) + a)^2} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{\int \frac{1}{\sin(c+dx)a+a} dx}{3a} - \frac{\cos(c + dx)}{3d(a \sin(c + dx) + a)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{\sin(c+dx)a+a} dx}{3a} - \frac{\cos(c + dx)}{3d(a \sin(c + dx) + a)^2} \\
 & \quad \downarrow \text{3127} \\
 & -\frac{\cos(c + dx)}{3ad(a \sin(c + dx) + a)} - \frac{\cos(c + dx)}{3d(a \sin(c + dx) + a)^2}
 \end{aligned}$$

input `Int[(a + a*Sin[c + d*x])^(-2),x]`

output `-1/3*Cos[c + d*x]/(d*(a + a*Sin[c + d*x])^2) - Cos[c + d*x]/(3*a*d*(a + a*Sin[c + d*x]))`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3127 $\text{Int}[(a_ + (b_)\sin[(c_ + (d_)(x_)])^{-1}, x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/(d*(b + a*\text{Sin}[c + d*x])), x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

rule 3129 $\text{Int}[(a_ + (b_)\sin[(c_ + (d_)(x_)])^{n_}, x_Symbol] \rightarrow \text{Simp}[b*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^n/(a*d*(2*n + 1))), x] + \text{Simp}[(n + 1)/(a*(2*n + 1)) \ \text{Int}[(a + b*\text{Sin}[c + d*x])^{n + 1}, x], x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.69

method	result	size
risch	$-\frac{2i(i+3e^{i(dx+c)})}{3da^2(e^{i(dx+c)}+i)^3}$	38
parallelrisc	$\frac{-4-6\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2-6\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{3da^2\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}$	48
derivativdivides	$-\frac{\frac{2}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1}+\frac{2}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^2}-\frac{4}{3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}}{da^2}$	53
default	$-\frac{\frac{2}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1}+\frac{2}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^2}-\frac{4}{3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}}{da^2}$	53
norman	$\frac{-\frac{2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{ad}-\frac{2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2}{ad}-\frac{4}{3ad}}{a\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^3}$	63

input $\text{int}(1/(a+a*\sin(d*x+c))^2,x,\text{method}=_RETURNVERBOSE)$

output $-2/3*I*(I+3*\exp(I*(d*x+c)))/d/a^2/(\exp(I*(d*x+c))+I)^3$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.73

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx$$

$$= \frac{\cos(dx + c)^2 + (\cos(dx + c) - 1) \sin(dx + c) + 2 \cos(dx + c) + 1}{3(a^2 d \cos(dx + c)^2 - a^2 d \cos(dx + c) - 2a^2 d - (a^2 d \cos(dx + c) + 2a^2 d) \sin(dx + c))}$$

input `integrate(1/(a+a*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/3*(cos(d*x + c)^2 + (cos(d*x + c) - 1)*sin(d*x + c) + 2*cos(d*x + c) + 1)/(a^2*d*cos(d*x + c)^2 - a^2*d*cos(d*x + c) - 2*a^2*d - (a^2*d*cos(d*x + c) + 2*a^2*d)*sin(d*x + c))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. 2(46) = 92.

Time = 0.71 (sec) , antiderivative size = 221, normalized size of antiderivative = 4.02

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx$$

$$= \begin{cases} -\frac{6 \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{3a^2 d \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right) + 9a^2 d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 9a^2 d \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 3a^2 d} - \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3a^2 d \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right) + 9a^2 d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 9a^2 d \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 3a^2 d} \\ \frac{x}{(a \sin(c) + a)^2} \end{cases}$$

input `integrate(1/(a+a*sin(d*x+c))**2,x)`

output `Piecewise((-6*tan(c/2 + d*x/2)**2/(3*a**2*d*tan(c/2 + d*x/2)**3 + 9*a**2*d*tan(c/2 + d*x/2)**2 + 9*a**2*d*tan(c/2 + d*x/2) + 3*a**2*d) - 6*tan(c/2 + d*x/2)/(3*a**2*d*tan(c/2 + d*x/2)**3 + 9*a**2*d*tan(c/2 + d*x/2)**2 + 9*a**2*d*tan(c/2 + d*x/2) + 3*a**2*d) - 4/(3*a**2*d*tan(c/2 + d*x/2)**3 + 9*a**2*d*tan(c/2 + d*x/2)**2 + 9*a**2*d*tan(c/2 + d*x/2) + 3*a**2*d), Ne(d, 0)), (x/(a*sin(c) + a)**2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 117 vs. $2(51) = 102$.

Time = 0.05 (sec) , antiderivative size = 117, normalized size of antiderivative = 2.13

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx = -\frac{2 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 2 \right)}{3 \left(a^2 + \frac{3 a^2 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3 a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{a^2 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right) d}$$

input `integrate(1/(a+a*sin(d*x+c))^2,x, algorithm="maxima")`

output `-2/3*(3*sin(d*x + c)/(cos(d*x + c) + 1) + 3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 2)/((a^2 + 3*a^2*sin(d*x + c)/(cos(d*x + c) + 1) + 3*a^2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + a^2*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)*d`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.85

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx = -\frac{2 \left(3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 + 3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 2 \right)}{3 a^2 d \left(\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1 \right)^3}$$

input `integrate(1/(a+a*sin(d*x+c))^2,x, algorithm="giac")`

output `-2/3*(3*tan(1/2*d*x + 1/2*c)^2 + 3*tan(1/2*d*x + 1/2*c) + 2)/(a^2*d*(tan(1/2*d*x + 1/2*c) + 1)^3)`

Mupad [B] (verification not implemented)

Time = 25.74 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.38

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx = -\frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 3\right)}{3}{a^2 d \left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right) + \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)^3}$$

input `int(1/(a + a*sin(c + d*x))^2,x)`output `-(2*cos(c/2 + (d*x)/2)^2*sin(c/2 + (d*x)/2) - (2*cos(c/2 + (d*x)/2)*(cos(c/2 + (d*x)/2)^2 - 3))/3/(a^2*d*(cos(c/2 + (d*x)/2) + sin(c/2 + (d*x)/2))^3)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.09

$$\int \frac{1}{(a + a \sin(c + dx))^2} dx = \frac{\frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{3} - \frac{2}{3}}{a^2 d \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3 + 3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}$$

input `int(1/(a+a*sin(d*x+c))^2,x)`output `(2*(tan((c + d*x)/2)**3 - 1))/(3*a**2*d*(tan((c + d*x)/2)**3 + 3*tan((c + d*x)/2)**2 + 3*tan((c + d*x)/2) + 1))`

3.6 $\int \frac{1}{(a+a \sin(c+dx))^3} dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [C] (verified)	92
Fricas [A] (verification not implemented)	92
Sympy [B] (verification not implemented)	93
Maxima [B] (verification not implemented)	94
Giac [A] (verification not implemented)	94
Mupad [B] (verification not implemented)	95
Reduce [B] (verification not implemented)	95

Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = -\frac{\cos(c + dx)}{5d(a + a \sin(c + dx))^3} - \frac{2 \cos(c + dx)}{15ad(a + a \sin(c + dx))^2} - \frac{2 \cos(c + dx)}{15d(a^3 + a^3 \sin(c + dx))}$$

output

```
-1/5*cos(d*x+c)/d/(a+a*sin(d*x+c))^3-2/15*cos(d*x+c)/a/d/(a+a*sin(d*x+c))^2-2/15*cos(d*x+c)/d/(a^3+a^3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.92

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \frac{10 - 15 \cos(c + dx) - 6 \cos(2(c + dx)) + \cos(3(c + dx)) + 15 \sin(c + dx) - 6 \sin(2(c + dx)) - \sin(3(c + dx))}{30a^3d(1 + \sin(c + dx))^3}$$

input

```
Integrate[(a + a*Sin[c + d*x])^(-3),x]
```

output

```
(10 - 15*Cos[c + d*x] - 6*Cos[2*(c + d*x)] + Cos[3*(c + d*x)] + 15*Sin[c +
d*x] - 6*Sin[2*(c + d*x)] - Sin[3*(c + d*x)])/(30*a^3*d*(1 + Sin[c + d*x]
)^3)
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3129, 3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sin(c + dx) + a)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx) + a)^3} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{2 \int \frac{1}{(\sin(c+dx)a+a)^2} dx}{5a} - \frac{\cos(c + dx)}{5d(a \sin(c + dx) + a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \int \frac{1}{(\sin(c+dx)a+a)^2} dx}{5a} - \frac{\cos(c + dx)}{5d(a \sin(c + dx) + a)^3} \\
 & \quad \downarrow \text{3129} \\
 & \frac{2 \left(\frac{\int \frac{1}{\sin(c+dx)a+a} dx}{3a} - \frac{\cos(c+dx)}{3d(a \sin(c+dx)+a)^2} \right)}{5a} - \frac{\cos(c + dx)}{5d(a \sin(c + dx) + a)^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \left(\frac{\int \frac{1}{\sin(c+dx)a+a} dx}{3a} - \frac{\cos(c+dx)}{3d(a \sin(c+dx)+a)^2} \right)}{5a} - \frac{\cos(c + dx)}{5d(a \sin(c + dx) + a)^3} \\
 & \quad \downarrow \text{3127}
 \end{aligned}$$

$$2\left(\frac{\cos(c+dx)}{3ad(a\sin(c+dx)+a)} - \frac{\cos(c+dx)}{3d(a\sin(c+dx)+a)^2}\right) - \frac{\cos(c+dx)}{5d(a\sin(c+dx)+a)^3}$$

input `Int[(a + a*Sin[c + d*x])^(-3),x]`

output `-1/5*Cos[c + d*x]/(d*(a + a*Sin[c + d*x])^3) + (2*(-1/3*Cos[c + d*x]/(d*(a + a*Sin[c + d*x])^2) - Cos[c + d*x]/(3*a*d*(a + a*Sin[c + d*x]))) / (5*a)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] :> Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.21 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.58

method	result	size
risch	$\frac{-\frac{4}{15} + \frac{8e^{2i(dx+c)}}{3} + \frac{4ie^{i(dx+c)}}{3}}{da^3(e^{i(dx+c)}+i)^5}$	48
paralelrisch	$\frac{-\frac{14}{15} - 2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 - 4 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3 - \frac{16 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{3} - \frac{8 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{3}}{da^3\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^5}$	74
derivativedivides	$\frac{\frac{4}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^4} - \frac{8}{5\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^5} - \frac{2}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1} + \frac{4}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{16}{3\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^3}}{da^3}$	85
default	$\frac{\frac{4}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^4} - \frac{8}{5\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^5} - \frac{2}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1} + \frac{4}{\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^2} - \frac{16}{3\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^3}}{da^3}$	85
norman	$\frac{-\frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{da} - \frac{14}{15ad} - \frac{4 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{da} - \frac{8 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{3ad} - \frac{16 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{3ad}}{a^2\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)^5}$	101

input

```
int(1/(a+a*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

$$\frac{4}{15} * (-1 + 10 * \exp(2 * I * (d * x + c)) + 5 * I * \exp(I * (d * x + c))) / d / a^3 / (\exp(I * (d * x + c)) + I)^5$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.77

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \frac{2 \cos(dx + c)^3 - 4 \cos(dx + c)^2 - (2 \cos(dx + c)^2 + 6 \cos(dx + c) - 3) \sin(dx + c) - 15(a^3 d \cos(dx + c))^3 + 3a^3 d \cos(dx + c)^2 - 2a^3 d \cos(dx + c) - 4a^3 d + (a^3 d \cos(dx + c))^2 - 2a^3 d \cos(dx + c)}{15(a^3 d \cos(dx + c))^3 + 3a^3 d \cos(dx + c)^2 - 2a^3 d \cos(dx + c) - 4a^3 d + (a^3 d \cos(dx + c))^2 - 2a^3 d \cos(dx + c)}$$

input

```
integrate(1/(a+a*sin(d*x+c))^3,x, algorithm="fricas")
```

output

```
-1/15*(2*cos(d*x + c)^3 - 4*cos(d*x + c)^2 - (2*cos(d*x + c)^2 + 6*cos(d*x + c) - 3)*sin(d*x + c) - 9*cos(d*x + c) - 3)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d + (a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d)*sin(d*x + c))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 558 vs. $2(73) = 146$.

Time = 1.37 (sec) , antiderivative size = 558, normalized size of antiderivative = 6.72

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(a+a*sin(d*x+c))**3,x)
```

output

```
Piecewise((-30*tan(c/2 + d*x/2)**4/(15*a**3*d*tan(c/2 + d*x/2)**5 + 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 + 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) + 15*a**3*d) - 60*tan(c/2 + d*x/2)**3/(15*a**3*d*tan(c/2 + d*x/2)**5 + 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 + 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) + 15*a**3*d) - 80*tan(c/2 + d*x/2)**2/(15*a**3*d*tan(c/2 + d*x/2)**5 + 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 + 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) + 15*a**3*d) - 40*tan(c/2 + d*x/2)/(15*a**3*d*tan(c/2 + d*x/2)**5 + 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 + 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) + 15*a**3*d) - 14/(15*a**3*d*tan(c/2 + d*x/2)**5 + 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 + 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) + 15*a**3*d), Ne(d, 0)), (x/(a*sin(c) + a)**3, True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 203 vs. $2(77) = 154$.

Time = 0.04 (sec) , antiderivative size = 203, normalized size of antiderivative = 2.45

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \frac{2 \left(\frac{20 \sin(dx+c)}{\cos(dx+c)+1} + \frac{40 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{30 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{15 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 7 \right)}{15 \left(a^3 + \frac{5 a^3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{10 a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{10 a^3 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{5 a^3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{a^3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)} d$$

input `integrate(1/(a+a*sin(d*x+c))^3,x, algorithm="maxima")`

output `-2/15*(20*sin(d*x + c)/(cos(d*x + c) + 1) + 40*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 30*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 15*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 7)/((a^3 + 5*a^3*sin(d*x + c)/(cos(d*x + c) + 1) + 10*a^3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 10*a^3*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 5*a^3*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + a^3*sin(d*x + c)^5/(cos(d*x + c) + 1)^5)*d)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \frac{2 \left(15 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^4 + 30 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 + 40 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 + 20 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 7 \right)}{15 a^3 d (\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 1)^5}$$

input `integrate(1/(a+a*sin(d*x+c))^3,x, algorithm="giac")`

output `-2/15*(15*tan(1/2*d*x + 1/2*c)^4 + 30*tan(1/2*d*x + 1/2*c)^3 + 40*tan(1/2*d*x + 1/2*c)^2 + 20*tan(1/2*d*x + 1/2*c) + 7)/(a^3*d*(tan(1/2*d*x + 1/2*c) + 1)^5)`

Mupad [B] (verification not implemented)

Time = 25.90 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.60

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \left(7 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + 20 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) + 40 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 30 \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^3 + 20 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^4\right)}{15 a^3 d \left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right) + \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)^5}$$

input `int(1/(a + a*sin(c + d*x))^3,x)`output `-(2*cos(c/2 + (d*x)/2)*(7*cos(c/2 + (d*x)/2)^4 + 15*sin(c/2 + (d*x)/2)^4 + 30*cos(c/2 + (d*x)/2)*sin(c/2 + (d*x)/2)^3 + 20*cos(c/2 + (d*x)/2)^3*sin(c/2 + (d*x)/2) + 40*cos(c/2 + (d*x)/2)^2*sin(c/2 + (d*x)/2)^2))/(15*a^3*d*(cos(c/2 + (d*x)/2) + sin(c/2 + (d*x)/2))^5)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.35

$$\int \frac{1}{(a + a \sin(c + dx))^3} dx = \frac{\frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{5} - \frac{4 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{3} - \frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{3} - \frac{8}{15}}{a^3 d \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5 + 5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + 10 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3 + 10 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}$$

input `int(1/(a+a*sin(d*x+c))^3,x)`output `(2*(3*tan((c + d*x)/2)**5 - 10*tan((c + d*x)/2)**2 - 5*tan((c + d*x)/2) - 4))/(15*a**3*d*(tan((c + d*x)/2)**5 + 5*tan((c + d*x)/2)**4 + 10*tan((c + d*x)/2)**3 + 10*tan((c + d*x)/2)**2 + 5*tan((c + d*x)/2) + 1))`

3.7 $\int (a - a \sin(c + dx))^3 dx$

Optimal result	96
Mathematica [A] (verified)	96
Rubi [A] (verified)	97
Maple [A] (verified)	98
Fricas [A] (verification not implemented)	99
Sympy [B] (verification not implemented)	99
Maxima [A] (verification not implemented)	100
Giac [A] (verification not implemented)	100
Mupad [B] (verification not implemented)	101
Reduce [B] (verification not implemented)	101

Optimal result

Integrand size = 13, antiderivative size = 63

$$\int (a - a \sin(c + dx))^3 dx = \frac{5a^3x}{2} + \frac{4a^3 \cos(c + dx)}{d} - \frac{a^3 \cos^3(c + dx)}{3d} - \frac{3a^3 \cos(c + dx) \sin(c + dx)}{2d}$$

output

$5/2*a^3*x+4*a^3*\cos(d*x+c)/d-1/3*a^3*\cos(d*x+c)^3/d-3/2*a^3*\cos(d*x+c)*\sin(d*x+c)/d$

Mathematica [A] (verified)

Time = 5.38 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int (a - a \sin(c + dx))^3 dx = \frac{a^3(30c + 30dx + 45 \cos(c + dx) - \cos(3(c + dx)) - 9 \sin(2(c + dx)))}{12d}$$

input

`Integrate[(a - a*Sin[c + d*x])^3,x]`

output

$$(a^3(30c + 30dx + 45\cos[c + dx] - \cos[3(c + dx)] - 9\sin[2(c + dx)]))/(12d)$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 3124, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a - a \sin(c + dx))^3 dx \\ & \quad \downarrow \text{3042} \\ & \int (a - a \sin(c + dx))^3 dx \\ & \quad \downarrow \text{3124} \\ & \int (a^3(-\sin^3(c + dx)) + 3a^3 \sin^2(c + dx) - 3a^3 \sin(c + dx) + a^3) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{a^3 \cos^3(c + dx)}{3d} + \frac{4a^3 \cos(c + dx)}{d} - \frac{3a^3 \sin(c + dx) \cos(c + dx)}{2d} + \frac{5a^3 x}{2} \end{aligned}$$

input

$$\text{Int}[(a - a\text{Sin}[c + d*x])^3, x]$$

output

$$(5a^3x)/2 + (4a^3\cos[c + d*x])/d - (a^3\cos[c + d*x]^3)/(3d) - (3a^3\cos[c + d*x]*\sin[c + d*x])/(2d)$$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3124 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrig[(a + b*sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 7.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

method	result
parallelrisch	$\frac{a^3(30dx+45\cos(dx+c)-\cos(3dx+3c)-9\sin(2dx+2c)+44)}{12d}$
risch	$\frac{5a^3x}{2} + \frac{15a^3\cos(dx+c)}{4d} - \frac{a^3\cos(3dx+3c)}{12d} - \frac{3a^3\sin(2dx+2c)}{4d}$
derivativedivides	$\frac{a^3(2+\sin(dx+c)^2)\cos(dx+c)}{3} + 3a^3\left(\frac{-\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + 3a^3\cos(dx+c) + a^3(dx+c)$
default	$\frac{a^3(2+\sin(dx+c)^2)\cos(dx+c)}{3} + 3a^3\left(\frac{-\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + 3a^3\cos(dx+c) + a^3(dx+c)$
parts	$a^3x + \frac{3a^3\cos(dx+c)}{d} + \frac{3a^3\left(\frac{-\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{a^3(2+\sin(dx+c)^2)\cos(dx+c)}{3d}$
norman	$\frac{16a^3\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{d} + \frac{5a^3x}{2} + \frac{22a^3}{3d} - \frac{3a^3\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{3a^3\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{d} + \frac{15a^3x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{2} + \frac{15a^3x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{2} + \frac{5a^3x}{2}$ $\left(1+\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2\right)^3$

input `int((a-a*sin(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/12*a^3*(30*d*x+45*cos(d*x+c)-cos(3*d*x+3*c)-9*sin(2*d*x+2*c)+44)/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.86

$$\int (a - a \sin(c + dx))^3 dx$$

$$= \frac{2a^3 \cos(dx + c)^3 - 15a^3 dx + 9a^3 \cos(dx + c) \sin(dx + c) - 24a^3 \cos(dx + c)}{6d}$$

input `integrate((a-a*sin(d*x+c))^3,x, algorithm="fricas")`

output `-1/6*(2*a^3*cos(d*x + c)^3 - 15*a^3*d*x + 9*a^3*cos(d*x + c)*sin(d*x + c) - 24*a^3*cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. 2(58) = 116.

Time = 0.12 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.92

$$\int (a - a \sin(c + dx))^3 dx$$

$$= \begin{cases} \frac{3a^3 x \sin^2(c+dx)}{2} + \frac{3a^3 x \cos^2(c+dx)}{2} + a^3 x + \frac{a^3 \sin^2(c+dx) \cos(c+dx)}{d} - \frac{3a^3 \sin(c+dx) \cos(c+dx)}{2d} + \frac{2a^3 \cos^3(c+dx)}{3d} + \frac{3a^3 \cos^3(c+dx)}{3d} \\ x(-a \sin(c) + a)^3 \end{cases}$$

input `integrate((a-a*sin(d*x+c))**3,x)`

output `Piecewise((3*a**3*x*sin(c + d*x)**2/2 + 3*a**3*x*cos(c + d*x)**2/2 + a**3*x + a**3*sin(c + d*x)**2*cos(c + d*x)/d - 3*a**3*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*a**3*cos(c + d*x)**3/(3*d) + 3*a**3*cos(c + d*x)/d, Ne(d, 0)), (x*(-a*sin(c) + a)**3, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.14

$$\int (a - a \sin(c + dx))^3 dx = a^3 x - \frac{(\cos(dx + c))^3 - 3 \cos(dx + c)}{3d} a^3 + \frac{3(2dx + 2c - \sin(2dx + 2c))a^3}{4d} + \frac{3a^3 \cos(dx + c)}{d}$$

input `integrate((a-a*sin(d*x+c))^3,x, algorithm="maxima")`

output `a^3*x - 1/3*(cos(d*x + c)^3 - 3*cos(d*x + c))*a^3/d + 3/4*(2*d*x + 2*c - sin(2*d*x + 2*c))*a^3/d + 3*a^3*cos(d*x + c)/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.87

$$\int (a - a \sin(c + dx))^3 dx = \frac{5}{2} a^3 x - \frac{a^3 \cos(3dx + 3c)}{12d} + \frac{15 a^3 \cos(dx + c)}{4d} - \frac{3 a^3 \sin(2dx + 2c)}{4d}$$

input `integrate((a-a*sin(d*x+c))^3,x, algorithm="giac")`

output `5/2*a^3*x - 1/12*a^3*cos(3*d*x + 3*c)/d + 15/4*a^3*cos(d*x + c)/d - 3/4*a^3*sin(2*d*x + 2*c)/d`

Mupad [B] (verification not implemented)

Time = 28.21 (sec) , antiderivative size = 156, normalized size of antiderivative = 2.48

$$\int (a - a \sin(c + dx))^3 dx = \frac{5a^3 x}{2} - \frac{\frac{5a^3(c+dx)}{2} - 3a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 - \frac{a^3(15c+15dx+44)}{6} + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 \left(\frac{15a^3(c+dx)}{2} - \frac{a^3(45c+45dx+36)}{6}\right) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \left(\frac{15a^3(c+dx)}{2} - \frac{a^3(45c+45dx+36)}{6}\right) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \left(\frac{15a^3(c+dx)}{2} - \frac{a^3(45c+45dx+36)}{6}\right) + 3a^3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)^3}$$

input `int((a - a*sin(c + d*x))^3,x)`output `(5*a^3*x)/2 - ((5*a^3*(c + d*x))/2 - 3*a^3*tan(c/2 + (d*x)/2)^5 - (a^3*(15*c + 15*d*x + 44))/6 + tan(c/2 + (d*x)/2)^4*((15*a^3*(c + d*x))/2 - (a^3*(45*c + 45*d*x + 36))/6) + tan(c/2 + (d*x)/2)^2*((15*a^3*(c + d*x))/2 - (a^3*(45*c + 45*d*x + 96))/6) + 3*a^3*tan(c/2 + (d*x)/2))/(d*(tan(c/2 + (d*x)/2)^2 + 1)^3)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

$$\int (a - a \sin(c + dx))^3 dx = \frac{a^3(2 \cos(dx + c) \sin(dx + c)^2 - 9 \cos(dx + c) \sin(dx + c) + 22 \cos(dx + c) + 15dx - 22)}{6d}$$

input `int((a-a*sin(d*x+c))^3,x)`output `(a**3*(2*cos(c + d*x)*sin(c + d*x)**2 - 9*cos(c + d*x)*sin(c + d*x) + 22*cos(c + d*x) + 15*d*x - 22))/(6*d)`

3.8 $\int (a - a \sin(c + dx))^2 dx$

Optimal result	102
Mathematica [A] (verified)	102
Rubi [A] (verified)	103
Maple [A] (verified)	104
Fricas [A] (verification not implemented)	104
Sympy [A] (verification not implemented)	105
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	105
Mupad [B] (verification not implemented)	106
Reduce [B] (verification not implemented)	106

Optimal result

Integrand size = 13, antiderivative size = 45

$$\int (a - a \sin(c + dx))^2 dx = \frac{3a^2x}{2} + \frac{2a^2 \cos(c + dx)}{d} - \frac{a^2 \cos(c + dx) \sin(c + dx)}{2d}$$

output

```
3/2*a^2*x+2*a^2*cos(d*x+c)/d-1/2*a^2*cos(d*x+c)*sin(d*x+c)/d
```

Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int (a - a \sin(c + dx))^2 dx = \frac{a^2(6(c + dx) + 8 \cos(c + dx) - \sin(2(c + dx)))}{4d}$$

input

```
Integrate[(a - a*Sin[c + d*x])^2,x]
```

output

```
(a^2*(6*(c + d*x) + 8*Cos[c + d*x] - Sin[2*(c + d*x)]))/(4*d)
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3042, 3123}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a - a \sin(c + dx))^2 dx$$

↓ 3042

$$\int (a - a \sin(c + dx))^2 dx$$

↓ 3123

$$\frac{2a^2 \cos(c + dx)}{d} - \frac{a^2 \sin(c + dx) \cos(c + dx)}{2d} + \frac{3a^2 x}{2}$$

input `Int[(a - a*Sin[c + d*x])^2,x]`

output `(3*a^2*x)/2 + (2*a^2*Cos[c + d*x])/d - (a^2*Cos[c + d*x]*Sin[c + d*x])/(2*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3123 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^2, x_Symbol] := Simp[(2*a^2 + b^2)*(x/2), x] + (-Simp[2*a*b*(Cos[c + d*x]/d), x] - Simp[b^2*Cos[c + d*x]*(Sin[c + d*x]/(2*d)), x]) /; FreeQ[{a, b, c, d}, x]`

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.76

method	result
parallelrisch	$\frac{a^2(6dx+8\cos(dx+c)-\sin(2dx+2c))-8}{4d}$
risch	$\frac{3a^2x}{2} + \frac{2a^2\cos(dx+c)}{d} - \frac{a^2\sin(2dx+2c)}{4d}$
parts	$a^2x + \frac{a^2\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right)}{d} + \frac{2a^2\cos(dx+c)}{d}$
derivativedivides	$\frac{a^2\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + 2\cos(dx+c)a^2 + a^2(dx+c)}{d}$
default	$\frac{a^2\left(-\frac{\cos(dx+c)\sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2}\right) + 2\cos(dx+c)a^2 + a^2(dx+c)}{d}$
norman	$\frac{\frac{a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{d} + \frac{3a^2x}{2} - \frac{a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{d} + 3a^2x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{3a^2x\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{2} - \frac{4a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{d} - \frac{4a^2\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{d}}{\left(1 + \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2}$
orering	$x(a - a\sin(dx+c))^2 + \frac{5(a-a\sin(dx+c))a\cos(dx+c)}{2d} + \frac{5x(2a^2d^2\cos(dx+c)^2 + 2(a-a\sin(dx+c))ad^2\sin(dx+c))}{4d^2}$

input `int((a-a*sin(d*x+c))^2,x,method=_RETURNVERBOSE)`output `1/4*a^2*(6*d*x+8*cos(d*x+c)-sin(2*d*x+2*c))-8)/d`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.91

$$\int (a - a\sin(c + dx))^2 dx = \frac{3a^2dx - a^2\cos(dx+c)\sin(dx+c) + 4a^2\cos(dx+c)}{2d}$$

input `integrate((a-a*sin(d*x+c))^2,x, algorithm="fricas")`output `1/2*(3*a^2*d*x - a^2*cos(d*x + c)*sin(d*x + c) + 4*a^2*cos(d*x + c))/d`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.73

$$\int (a - a \sin(c + dx))^2 dx = \begin{cases} \frac{a^2 x \sin^2(c+dx)}{2} + \frac{a^2 x \cos^2(c+dx)}{2} + a^2 x - \frac{a^2 \sin(c+dx) \cos(c+dx)}{2d} + \frac{2a^2 \cos(c+dx)}{d} & \text{for } d \neq 0 \\ x(-a \sin(c) + a)^2 & \text{otherwise} \end{cases}$$

input `integrate((a-a*sin(d*x+c))**2,x)`output `Piecewise((a**2*x*sin(c + d*x)**2/2 + a**2*x*cos(c + d*x)**2/2 + a**2*x - a**2*sin(c + d*x)*cos(c + d*x)/(2*d) + 2*a**2*cos(c + d*x)/d, Ne(d, 0)), (x*(-a*sin(c) + a)**2, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int (a - a \sin(c + dx))^2 dx = a^2 x + \frac{(2 dx + 2 c - \sin(2 dx + 2 c))a^2}{4 d} + \frac{2 a^2 \cos(dx + c)}{d}$$

input `integrate((a-a*sin(d*x+c))^2,x, algorithm="maxima")`output `a^2*x + 1/4*(2*d*x + 2*c - sin(2*d*x + 2*c))*a^2/d + 2*a^2*cos(d*x + c)/d`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.84

$$\int (a - a \sin(c + dx))^2 dx = \frac{3}{2} a^2 x + \frac{2 a^2 \cos(dx + c)}{d} - \frac{a^2 \sin(2 dx + 2 c)}{4 d}$$

input `integrate((a-a*sin(d*x+c))^2,x, algorithm="giac")`

output $3/2*a^2*x + 2*a^2*\cos(d*x + c)/d - 1/4*a^2*\sin(2*d*x + 2*c)/d$

Mupad [B] (verification not implemented)

Time = 25.93 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.73

$$\int (a - a \sin(c + dx))^2 dx = \frac{3a^2 x}{2} - \frac{a^2 \left(\frac{3c}{2} + \frac{3dx}{2}\right) - a^2 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3 - a^2 \left(\frac{3c}{2} + \frac{3dx}{2} + 4\right) + \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \left(2a^2 \left(\frac{3c}{2} + \frac{3dx}{2}\right) - a^2 (3c + 3a)\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1\right)^2}$$

input $\text{int}((a - a*\sin(c + d*x))^2,x)$

output $(3*a^2*x)/2 - (a^2*((3*c)/2 + (3*d*x)/2) - a^2*\tan(c/2 + (d*x)/2)^3 - a^2*((3*c)/2 + (3*d*x)/2 + 4) + \tan(c/2 + (d*x)/2)^2*(2*a^2*((3*c)/2 + (3*d*x)/2) - a^2*(3*c + 3*d*x + 4)) + a^2*\tan(c/2 + (d*x)/2))/(d*(\tan(c/2 + (d*x)/2)^2 + 1)^2)$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int (a - a \sin(c + dx))^2 dx = \frac{a^2(-\cos(dx + c)\sin(dx + c) + 4\cos(dx + c) + 3dx - 4)}{2d}$$

input $\text{int}((a-a*\sin(d*x+c))^2,x)$

output $(a**2*(-\cos(c + d*x)*\sin(c + d*x) + 4*\cos(c + d*x) + 3*d*x - 4))/(2*d)$

3.9 $\int (a - a \sin(c + dx)) dx$

Optimal result	107
Mathematica [A] (verified)	107
Rubi [A] (verified)	108
Maple [A] (verified)	109
Fricas [A] (verification not implemented)	109
Sympy [A] (verification not implemented)	110
Maxima [A] (verification not implemented)	110
Giac [A] (verification not implemented)	110
Mupad [B] (verification not implemented)	111
Reduce [B] (verification not implemented)	111

Optimal result

Integrand size = 11, antiderivative size = 15

$$\int (a - a \sin(c + dx)) dx = ax + \frac{a \cos(c + dx)}{d}$$

output

```
a*x+a*cos(d*x+c)/d
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int (a - a \sin(c + dx)) dx = ax + \frac{a \cos(c) \cos(dx)}{d} - \frac{a \sin(c) \sin(dx)}{d}$$

input

```
Integrate[a - a*Sin[c + d*x],x]
```

output

```
a*x + (a*Cos[c]*Cos[d*x])/d - (a*Sin[c]*Sin[d*x])/d
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a - a \sin(c + dx)) dx$$

$$\downarrow \text{2009}$$

$$\frac{a \cos(c + dx)}{d} + ax$$

input `Int[a - a*Sin[c + d*x],x]`

output `a*x + (a*Cos[c + d*x])/d`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$ax + \frac{a \cos(dx+c)}{d}$	16
risch	$ax + \frac{a \cos(dx+c)}{d}$	16
parts	$ax + \frac{a \cos(dx+c)}{d}$	16
derivativedivides	$\frac{(dx+c)a+a \cos(dx+c)}{d}$	21
parallelrisc	$-\frac{a(-1-\cos(dx+c))}{d} + ax$	21
orering	$x(a - a \sin(dx + c)) + \frac{a \cos(dx+c)}{d} + xa \sin(dx + c)$	35
norman	$\frac{ax+ax \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2 - \frac{2a \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{d}}{1+\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2}$	52

input `int(a-a*sin(d*x+c),x,method=_RETURNVERBOSE)`output `a*x+a*cos(d*x+c)/d`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int (a - a \sin(c + dx)) dx = \frac{adx + a \cos(dx + c)}{d}$$

input `integrate(a-a*sin(d*x+c),x, algorithm="fricas")`output `(a*d*x + a*cos(d*x + c))/d`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int (a - a \sin(c + dx)) dx = ax - a \begin{cases} -\frac{\cos(c+dx)}{d} & \text{for } d \neq 0 \\ x \sin(c) & \text{otherwise} \end{cases}$$

input `integrate(a-a*sin(d*x+c),x)`output `a*x - a*Piecewise((-cos(c + d*x)/d, Ne(d, 0)), (x*sin(c), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (a - a \sin(c + dx)) dx = ax + \frac{a \cos(dx + c)}{d}$$

input `integrate(a-a*sin(d*x+c),x, algorithm="maxima")`output `a*x + a*cos(d*x + c)/d`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (a - a \sin(c + dx)) dx = ax + \frac{a \cos(dx + c)}{d}$$

input `integrate(a-a*sin(d*x+c),x, algorithm="giac")`output `a*x + a*cos(d*x + c)/d`

Mupad [B] (verification not implemented)

Time = 25.64 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.67

$$\int (a - a \sin(c + dx)) dx = ax + \frac{2a}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 1 \right)}$$

input `int(a - a*sin(c + d*x),x)`

output `a*x + (2*a)/(d*(tan(c/2 + (d*x)/2)^2 + 1))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int (a - a \sin(c + dx)) dx = \frac{a(\cos(dx + c) + dx)}{d}$$

input `int(a-a*sin(d*x+c),x)`

output `(a*(cos(c + d*x) + d*x))/d`

3.10 $\int \frac{1}{a - a \sin(c + dx)} dx$

Optimal result	112
Mathematica [B] (verified)	112
Rubi [A] (verified)	113
Maple [A] (verified)	114
Fricas [A] (verification not implemented)	114
Sympy [A] (verification not implemented)	115
Maxima [A] (verification not implemented)	115
Giac [A] (verification not implemented)	115
Mupad [B] (verification not implemented)	116
Reduce [B] (verification not implemented)	116

Optimal result

Integrand size = 13, antiderivative size = 23

$$\int \frac{1}{a - a \sin(c + dx)} dx = \frac{\cos(c + dx)}{d(a - a \sin(c + dx))}$$

output `cos(d*x+c)/d/(a-a*sin(d*x+c))`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. 2(23) = 46.

Time = 0.26 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.22

$$\int \frac{1}{a - a \sin(c + dx)} dx = \frac{2(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) \sin(\frac{1}{2}(c + dx))}{d(a - a \sin(c + dx))}$$

input `Integrate[(a - a*Sin[c + d*x])^(-1),x]`

output `(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*Sin[(c + d*x)/2])/(d*(a - a*Sin[c + d*x]))`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a - a \sin(c + dx)} dx$$

↓ 3042

$$\int \frac{1}{a - a \sin(c + dx)} dx$$

↓ 3127

$$\frac{\cos(c + dx)}{d(a - a \sin(c + dx))}$$

input `Int[(a - a*Sin[c + d*x])^(-1),x]`

output `Cos[c + d*x]/(d*(a - a*Sin[c + d*x]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] :> Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$-\frac{2}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}$	22
default	$-\frac{2}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}$	22
norman	$-\frac{2}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}$	22
parallelrisc	$-\frac{2}{da\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}$	22
risc	$\frac{2}{da\left(e^{i(dx+c)}-i\right)}$	23

input `int(1/(a-a*sin(d*x+c)),x,method=_RETURNVERBOSE)`output `-2/d/a/(tan(1/2*d*x+1/2*c)-1)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.74

$$\int \frac{1}{a - a \sin(c + dx)} dx = \frac{\cos(dx + c) + \sin(dx + c) + 1}{ad \cos(dx + c) - ad \sin(dx + c) + ad}$$

input `integrate(1/(a-a*sin(d*x+c)),x, algorithm="fricas")`output `(cos(d*x + c) + sin(d*x + c) + 1)/(a*d*cos(d*x + c) - a*d*sin(d*x + c) + a*d)`

Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \frac{1}{a - a \sin(c + dx)} dx = \begin{cases} -\frac{2}{ad \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - ad} & \text{for } d \neq 0 \\ \frac{x}{-a \sin(c) + a} & \text{otherwise} \end{cases}$$

input `integrate(1/(a-a*sin(d*x+c)),x)`output `Piecewise((-2/(a*d*tan(c/2 + d*x/2) - a*d), Ne(d, 0)), (x/(-a*sin(c) + a), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.22

$$\int \frac{1}{a - a \sin(c + dx)} dx = \frac{2}{\left(a - \frac{a \sin(dx+c)}{\cos(dx+c)+1}\right)d}$$

input `integrate(1/(a-a*sin(d*x+c)),x, algorithm="maxima")`output `2/((a - a*sin(d*x + c)/(cos(d*x + c) + 1))*d)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{1}{a - a \sin(c + dx)} dx = -\frac{2}{ad \left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right)}$$

input `integrate(1/(a-a*sin(d*x+c)),x, algorithm="giac")`output `-2/(a*d*(tan(1/2*d*x + 1/2*c) - 1))`

Mupad [B] (verification not implemented)

Time = 25.77 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{1}{a - a \sin(c + dx)} dx = -\frac{2}{ad \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1 \right)}$$

input `int(1/(a - a*sin(c + d*x)),x)`

output `-2/(a*d*(tan(c/2 + (d*x)/2) - 1))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \frac{1}{a - a \sin(c + dx)} dx = -\frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{ad \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1 \right)}$$

input `int(1/(a-a*sin(d*x+c)),x)`

output `(- 2*tan((c + d*x)/2))/(a*d*(tan((c + d*x)/2) - 1))`

3.11 $\int \frac{1}{(a - a \sin(c + dx))^2} dx$

Optimal result	117
Mathematica [A] (verified)	117
Rubi [A] (verified)	118
Maple [C] (verified)	119
Fricas [A] (verification not implemented)	120
Sympy [B] (verification not implemented)	120
Maxima [B] (verification not implemented)	121
Giac [A] (verification not implemented)	121
Mupad [B] (verification not implemented)	122
Reduce [B] (verification not implemented)	122

Optimal result

Integrand size = 13, antiderivative size = 57

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = \frac{\cos(c + dx)}{3d(a - a \sin(c + dx))^2} + \frac{\cos(c + dx)}{3d(a^2 - a^2 \sin(c + dx))}$$

output `1/3*cos(d*x+c)/d/(a-a*sin(d*x+c))^2+1/3*cos(d*x+c)/d/(a^2-a^2*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.98

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = \frac{-3 + 4 \cos(c + dx) + \cos(2(c + dx)) + 4 \sin(c + dx) - \sin(2(c + dx))}{6a^2d(-1 + \sin(c + dx))^2}$$

input `Integrate[(a - a*Sin[c + d*x])^(-2),x]`

output `(-3 + 4*Cos[c + d*x] + Cos[2*(c + d*x)] + 4*Sin[c + d*x] - Sin[2*(c + d*x)])/ (6*a^2*d*(-1 + Sin[c + d*x])^2)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - a \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a - a \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{\int \frac{1}{a - a \sin(c + dx)} dx}{3a} + \frac{\cos(c + dx)}{3d(a - a \sin(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{a - a \sin(c + dx)} dx}{3a} + \frac{\cos(c + dx)}{3d(a - a \sin(c + dx))^2} \\
 & \quad \downarrow \text{3127} \\
 & \frac{\cos(c + dx)}{3ad(a - a \sin(c + dx))} + \frac{\cos(c + dx)}{3d(a - a \sin(c + dx))^2}
 \end{aligned}$$

input `Int[(a - a*Sin[c + d*x])^(-2),x]`

output `Cos[c + d*x]/(3*d*(a - a*Sin[c + d*x])^2) + Cos[c + d*x]/(3*a*d*(a - a*Sin[c + d*x]))`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3127 $\text{Int}[\{(a_) + (b_) \cdot \sin[(c_) + (d_) \cdot (x)]\}^{-1}, x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d \cdot x] / (d \cdot (b + a \cdot \text{Sin}[c + d \cdot x])), x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

rule 3129 $\text{Int}[\{(a_) + (b_) \cdot \sin[(c_) + (d_) \cdot (x)]\}^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b \cdot \text{Cos}[c + d \cdot x] \cdot \{(a + b \cdot \text{Sin}[c + d \cdot x])^n / (a \cdot d \cdot (2 \cdot n + 1))\}, x] + \text{Simp}[(n + 1) / (a \cdot (2 \cdot n + 1)) \text{ Int}[(a + b \cdot \text{Sin}[c + d \cdot x])^{(n + 1)}, x], x] \text{ ; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2 \cdot n]$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.67

method	result	size
risch	$-\frac{2i(-i+3e^{i(dx+c)})}{3da^2(e^{i(dx+c)}-i)^3}$	38
parallelrisc	$\frac{-4-6 \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2+6 \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{3da^2\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}$	48
derivativedivides	$\frac{-\frac{4}{3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}-\frac{2}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2}-\frac{2}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1}}{da^2}$	55
default	$\frac{-\frac{4}{3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}-\frac{2}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2}-\frac{2}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1}}{da^2}$	55
norman	$\frac{\frac{2 \tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{ad}-\frac{4}{3ad}-\frac{2 \tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2}{ad}}{a\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}$	63

input $\text{int}(1/(a-a \cdot \sin(dx+c))^2, x, \text{method}=_RETURNVERBOSE)$

output $-2/3 \cdot I \cdot (-I+3 \cdot \exp(I \cdot (dx+c))) / d / a^2 / (\exp(I \cdot (dx+c))-I)^3$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.67

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = \frac{\cos(dx + c)^2 - (\cos(dx + c) - 1) \sin(dx + c) + 2 \cos(dx + c) + 1}{3(a^2 d \cos(dx + c))^2 - a^2 d \cos(dx + c) - 2a^2 d + (a^2 d \cos(dx + c) + 2a^2 d) \sin(dx + c)}$$

input `integrate(1/(a-a*sin(d*x+c))^2,x, algorithm="fricas")`

output `-1/3*(cos(d*x + c)^2 - (cos(d*x + c) - 1)*sin(d*x + c) + 2*cos(d*x + c) + 1)/(a^2*d*cos(d*x + c)^2 - a^2*d*cos(d*x + c) - 2*a^2*d + (a^2*d*cos(d*x + c) + 2*a^2*d)*sin(d*x + c))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(44) = 88.

Time = 0.69 (sec) , antiderivative size = 219, normalized size of antiderivative = 3.84

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = \begin{cases} -\frac{6 \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right)}{3a^2 d \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right) - 9a^2 d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 9a^2 d \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 3a^2 d} + \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3a^2 d \tan^3\left(\frac{c}{2} + \frac{dx}{2}\right) - 9a^2 d \tan^2\left(\frac{c}{2} + \frac{dx}{2}\right) + 9a^2 d \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 3a^2 d} \\ \frac{x}{(-a \sin(c) + a)^2} \end{cases}$$

input `integrate(1/(a-a*sin(d*x+c))**2,x)`

output `Piecewise((-6*tan(c/2 + d*x/2)**2/(3*a**2*d*tan(c/2 + d*x/2)**3 - 9*a**2*d*tan(c/2 + d*x/2)**2 + 9*a**2*d*tan(c/2 + d*x/2) - 3*a**2*d) + 6*tan(c/2 + d*x/2)/(3*a**2*d*tan(c/2 + d*x/2)**3 - 9*a**2*d*tan(c/2 + d*x/2)**2 + 9*a**2*d*tan(c/2 + d*x/2) - 3*a**2*d) - 4/(3*a**2*d*tan(c/2 + d*x/2)**3 - 9*a**2*d*tan(c/2 + d*x/2)**2 + 9*a**2*d*tan(c/2 + d*x/2) - 3*a**2*d), Ne(d, 0)), (x/(-a*sin(c) + a)**2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. $2(55) = 110$.

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 2.07

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = -\frac{2 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 2 \right)}{3 \left(a^2 - \frac{3 a^2 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3 a^2 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{a^2 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} \right) d}$$

input `integrate(1/(a-a*sin(d*x+c))^2,x, algorithm="maxima")`

output `-2/3*(3*sin(d*x + c)/(cos(d*x + c) + 1) - 3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 2)/((a^2 - 3*a^2*sin(d*x + c)/(cos(d*x + c) + 1) + 3*a^2*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - a^2*sin(d*x + c)^3/(cos(d*x + c) + 1)^3)*d`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = -\frac{2 \left(3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 - 3 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 2 \right)}{3 a^2 d \left(\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1 \right)^3}$$

input `integrate(1/(a-a*sin(d*x+c))^2,x, algorithm="giac")`

output `-2/3*(3*tan(1/2*d*x + 1/2*c)^2 - 3*tan(1/2*d*x + 1/2*c) + 2)/(a^2*d*(tan(1/2*d*x + 1/2*c) - 1)^3)`

Mupad [B] (verification not implemented)

Time = 25.95 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.37

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = -\frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) + \frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 3\right)}{3 a^2 d \left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right) - \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)^3}$$

input `int(1/(a - a*sin(c + d*x))^2,x)`output `-(2*cos(c/2 + (d*x)/2)^2*sin(c/2 + (d*x)/2) + (2*cos(c/2 + (d*x)/2)*(cos(c/2 + (d*x)/2)^2 - 3))/3/(a^2*d*(cos(c/2 + (d*x)/2) - sin(c/2 + (d*x)/2))^3)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.09

$$\int \frac{1}{(a - a \sin(c + dx))^2} dx = \frac{-2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3 - 2}{3a^2 d \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3 - 3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$$

input `int(1/(a-a*sin(d*x+c))^2,x)`output `(- 2*(tan((c + d*x)/2)**3 + 1))/(3*a**2*d*(tan((c + d*x)/2)**3 - 3*tan((c + d*x)/2)**2 + 3*tan((c + d*x)/2) - 1))`

3.12 $\int \frac{1}{(a - a \sin(c + dx))^3} dx$

Optimal result	123
Mathematica [A] (verified)	123
Rubi [A] (verified)	124
Maple [C] (verified)	126
Fricas [A] (verification not implemented)	126
Sympy [B] (verification not implemented)	127
Maxima [B] (verification not implemented)	128
Giac [A] (verification not implemented)	128
Mupad [B] (verification not implemented)	129
Reduce [B] (verification not implemented)	129

Optimal result

Integrand size = 13, antiderivative size = 86

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx = \frac{\cos(c + dx)}{5d(a - a \sin(c + dx))^3} + \frac{2 \cos(c + dx)}{15ad(a - a \sin(c + dx))^2} + \frac{2 \cos(c + dx)}{15d(a^3 - a^3 \sin(c + dx))}$$

output `1/5*cos(d*x+c)/d/(a-a*sin(d*x+c))^3+2/15*cos(d*x+c)/a/d/(a-a*sin(d*x+c))^2+2/15*cos(d*x+c)/d/(a^3-a^3*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx = \frac{10 - 15 \cos(c + dx) - 6 \cos(2(c + dx)) + \cos(3(c + dx)) - 15 \sin(c + dx) + 6 \sin(2(c + dx)) + \sin(3(c + dx))}{30a^3d(-1 + \sin(c + dx))^3}$$

input `Integrate[(a - a*Sin[c + d*x])^(-3),x]`

output

```
(10 - 15*Cos[c + d*x] - 6*Cos[2*(c + d*x)] + Cos[3*(c + d*x)] - 15*Sin[c +
d*x] + 6*Sin[2*(c + d*x)] + Sin[3*(c + d*x)])/(30*a^3*d*(-1 + Sin[c + d*x
])^3)
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3042, 3129, 3042, 3129, 3042, 3127}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - a \sin(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a - a \sin(c + dx))^3} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{2 \int \frac{1}{(a - a \sin(c + dx))^2} dx}{5a} + \frac{\cos(c + dx)}{5d(a - a \sin(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \int \frac{1}{(a - a \sin(c + dx))^2} dx}{5a} + \frac{\cos(c + dx)}{5d(a - a \sin(c + dx))^3} \\
 & \quad \downarrow \text{3129} \\
 & \frac{2 \left(\frac{\int \frac{1}{a - a \sin(c + dx)} dx}{3a} + \frac{\cos(c + dx)}{3d(a - a \sin(c + dx))^2} \right)}{5a} + \frac{\cos(c + dx)}{5d(a - a \sin(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \left(\frac{\int \frac{1}{a - a \sin(c + dx)} dx}{3a} + \frac{\cos(c + dx)}{3d(a - a \sin(c + dx))^2} \right)}{5a} + \frac{\cos(c + dx)}{5d(a - a \sin(c + dx))^3} \\
 & \quad \downarrow \text{3127}
 \end{aligned}$$

$$\frac{\cos(c+dx)}{5d(a-a\sin(c+dx))^3} + \frac{2\left(\frac{\cos(c+dx)}{3ad(a-a\sin(c+dx))} + \frac{\cos(c+dx)}{3d(a-a\sin(c+dx))^2}\right)}{5a}$$

input `Int[(a - a*Sin[c + d*x])^(-3),x]`

output `Cos[c + d*x]/(5*d*(a - a*Sin[c + d*x])^3) + (2*(Cos[c + d*x]/(3*d*(a - a*Sin[c + d*x])^2) + Cos[c + d*x]/(3*a*d*(a - a*Sin[c + d*x])))/(5*a)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3127 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] :> Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.21 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.56

method	result	size
risch	$-\frac{4(-1+10e^{2i(dx+c)}-5ie^{i(dx+c)})}{15(e^{i(dx+c)}-i)^5 da^3}$	48
parallelrisch	$\frac{-\frac{14}{15}-2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^4+4\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3-\frac{16\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2}{3}+\frac{8\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{3}}{da^3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^5}$	74
derivativedivides	$\frac{-\frac{4}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^4}-\frac{16}{3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}-\frac{4}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2}-\frac{8}{5\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^5}-\frac{2}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1}}{da^3}}$	85
default	$\frac{-\frac{4}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^4}-\frac{16}{3\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^3}-\frac{4}{\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^2}-\frac{8}{5\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^5}-\frac{2}{\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1}}{da^3}}$	85
norman	$\frac{-\frac{14}{15ad}+\frac{4\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^3}{da}-\frac{2\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^4}{da}+\frac{8\tan\left(\frac{dx}{2}+\frac{c}{2}\right)}{3ad}-\frac{16\tan\left(\frac{dx}{2}+\frac{c}{2}\right)^2}{3ad}}{a^2\left(\tan\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)^5}$	101

input `int(1/(a-a*sin(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{-4/15*(-1+10*\exp(2*I*(d*x+c))-5*I*\exp(I*(d*x+c)))/(\exp(I*(d*x+c))-I)^5/d/a}{^3}$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.71

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx$$

$$= \frac{2 \cos(dx + c)^3 - 4 \cos(dx + c)^2 + (2 \cos(dx + c)^2 + 6 \cos(dx + c) - 3) \sin(dx + c) - 9}{15 (a^3 d \cos(dx + c))^3 + 3 a^3 d \cos(dx + c)^2 - 2 a^3 d \cos(dx + c) - 4 a^3 d - (a^3 d \cos(dx + c))^2 - 2 a^3 d \cos(dx + c)}$$

input `integrate(1/(a-a*sin(d*x+c))^3,x, algorithm="fricas")`

output

```
1/15*(2*cos(d*x + c)^3 - 4*cos(d*x + c)^2 + (2*cos(d*x + c)^2 + 6*cos(d*x
+ c) - 3)*sin(d*x + c) - 9*cos(d*x + c) - 3)/(a^3*d*cos(d*x + c)^3 + 3*a^3
*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d - (a^3*d*cos(d*x + c)^2
- 2*a^3*d*cos(d*x + c) - 4*a^3*d)*sin(d*x + c))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 556 vs. $2(71) = 142$.

Time = 1.32 (sec) , antiderivative size = 556, normalized size of antiderivative = 6.47

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(a-a*sin(d*x+c))**3,x)
```

output

```
Piecewise((-30*tan(c/2 + d*x/2)**4/(15*a**3*d*tan(c/2 + d*x/2)**5 - 75*a**
3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 - 150*a**3*d*tan(
c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) - 15*a**3*d) + 60*tan(c/2 + d
*x/2)**3/(15*a**3*d*tan(c/2 + d*x/2)**5 - 75*a**3*d*tan(c/2 + d*x/2)**4 +
150*a**3*d*tan(c/2 + d*x/2)**3 - 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*
d*tan(c/2 + d*x/2) - 15*a**3*d) - 80*tan(c/2 + d*x/2)**2/(15*a**3*d*tan(c/
2 + d*x/2)**5 - 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2
)**3 - 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) - 15*a*
**3*d) + 40*tan(c/2 + d*x/2)/(15*a**3*d*tan(c/2 + d*x/2)**5 - 75*a**3*d*tan
(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)**3 - 150*a**3*d*tan(c/2 + d
*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) - 15*a**3*d) - 14/(15*a**3*d*tan(c/2
+ d*x/2)**5 - 75*a**3*d*tan(c/2 + d*x/2)**4 + 150*a**3*d*tan(c/2 + d*x/2)
**3 - 150*a**3*d*tan(c/2 + d*x/2)**2 + 75*a**3*d*tan(c/2 + d*x/2) - 15*a**
3*d), Ne(d, 0)), (x/(-a*sin(c) + a)**3, True))
```


Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. $2(83) = 166$.

Time = 0.04 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.37

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx = \frac{2 \left(\frac{20 \sin(dx+c)}{\cos(dx+c)+1} - \frac{40 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{30 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{15 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 7 \right)}{15 \left(a^3 - \frac{5 a^3 \sin(dx+c)}{\cos(dx+c)+1} + \frac{10 a^3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{10 a^3 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{5 a^3 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - \frac{a^3 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} \right)} d$$

input `integrate(1/(a-a*sin(d*x+c))^3,x, algorithm="maxima")`

output
$$-2/15*(20*\sin(d*x + c)/(\cos(d*x + c) + 1) - 40*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 30*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 - 15*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 - 7)/((a^3 - 5*a^3*\sin(d*x + c)/(\cos(d*x + c) + 1) + 10*a^3*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 - 10*a^3*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 + 5*a^3*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 - a^3*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5)*d)$$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.85

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx = \frac{2 \left(15 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^4 - 30 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^3 + 40 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right)^2 - 20 \tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) + 7 \right)}{15 a^3 d (\tan \left(\frac{1}{2} dx + \frac{1}{2} c \right) - 1)^5}$$

input `integrate(1/(a-a*sin(d*x+c))^3,x, algorithm="giac")`

output
$$-2/15*(15*\tan(1/2*d*x + 1/2*c)^4 - 30*\tan(1/2*d*x + 1/2*c)^3 + 40*\tan(1/2*d*x + 1/2*c)^2 - 20*\tan(1/2*d*x + 1/2*c) + 7)/(a^3*d*(\tan(1/2*d*x + 1/2*c) - 1)^5)$$

Mupad [B] (verification not implemented)

Time = 25.91 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.57

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx$$

$$= \frac{2 \cos\left(\frac{c}{2} + \frac{dx}{2}\right) \left(7 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - 20 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^3 \sin\left(\frac{c}{2} + \frac{dx}{2}\right) + 40 \cos\left(\frac{c}{2} + \frac{dx}{2}\right)^2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 30 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^3\right)}{15 a^3 d \left(\cos\left(\frac{c}{2} + \frac{dx}{2}\right) - \sin\left(\frac{c}{2} + \frac{dx}{2}\right)\right)^5}$$

input `int(1/(a - a*sin(c + d*x))^3,x)`output `(2*cos(c/2 + (d*x)/2)*(7*cos(c/2 + (d*x)/2)^4 + 15*sin(c/2 + (d*x)/2)^4 - 30*cos(c/2 + (d*x)/2)*sin(c/2 + (d*x)/2)^3 - 20*cos(c/2 + (d*x)/2)^3*sin(c/2 + (d*x)/2) + 40*cos(c/2 + (d*x)/2)^2*sin(c/2 + (d*x)/2)^2))/(15*a^3*d*(cos(c/2 + (d*x)/2) - sin(c/2 + (d*x)/2))^5)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.30

$$\int \frac{1}{(a - a \sin(c + dx))^3} dx$$

$$= \frac{-\frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{5} - \frac{4 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{3} + \frac{2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{3} - \frac{8}{15}}{a^3 d \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5 - 5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4 + 10 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3 - 10 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}$$

input `int(1/(a-a*sin(d*x+c))^3,x)`output `(2*(- 3*tan((c + d*x)/2)**5 - 10*tan((c + d*x)/2)**2 + 5*tan((c + d*x)/2) - 4))/(15*a**3*d*(tan((c + d*x)/2)**5 - 5*tan((c + d*x)/2)**4 + 10*tan((c + d*x)/2)**3 - 10*tan((c + d*x)/2)**2 + 5*tan((c + d*x)/2) - 1))`

3.13 $\int (a + a \sin(c + dx))^{7/2} dx$

Optimal result	130
Mathematica [A] (verified)	130
Rubi [A] (verified)	131
Maple [A] (verified)	133
Fricas [A] (verification not implemented)	134
Sympy [F(-1)]	134
Maxima [F]	134
Giac [A] (verification not implemented)	135
Mupad [F(-1)]	135
Reduce [F]	136

Optimal result

Integrand size = 14, antiderivative size = 119

$$\int (a + a \sin(c + dx))^{7/2} dx = -\frac{256a^4 \cos(c + dx)}{35d\sqrt{a + a \sin(c + dx)}} - \frac{64a^3 \cos(c + dx)\sqrt{a + a \sin(c + dx)}}{35d} - \frac{24a^2 \cos(c + dx)(a + a \sin(c + dx))^{3/2}}{35d} - \frac{2a \cos(c + dx)(a + a \sin(c + dx))^{5/2}}{7d}$$

output

```
-256/35*a^4*cos(d*x+c)/d/(a+a*sin(d*x+c))^(1/2)-64/35*a^3*cos(d*x+c)*(a+a*
sin(d*x+c))^(1/2)/d-24/35*a^2*cos(d*x+c)*(a+a*sin(d*x+c))^(3/2)/d-2/7*a*co
s(d*x+c)*(a+a*sin(d*x+c))^(5/2)/d
```

Mathematica [A] (verified)

Time = 3.15 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.29

$$\int (a + a \sin(c + dx))^{7/2} dx = \frac{a^3(1 + \sin(c + dx))^3 \sqrt{a(1 + \sin(c + dx))} (1225 \cos(\frac{1}{2}(c + dx)) + 245 \cos(\frac{3}{2}(c + dx)) - 49 \cos(\frac{5}{2}(c + dx)))}{140d (\cos(\frac{1}{2}(c + dx)))}$$

input `Integrate[(a + a*Sin[c + d*x])^(7/2),x]`

output
$$-1/140*(a^3*(1 + \text{Sin}[c + d*x])^3*\text{Sqrt}[a*(1 + \text{Sin}[c + d*x])]*(1225*\text{Cos}[(c + d*x)/2] + 245*\text{Cos}[(3*(c + d*x))/2] - 49*\text{Cos}[(5*(c + d*x))/2] - 5*\text{Cos}[(7*(c + d*x))/2] - 1225*\text{Sin}[(c + d*x)/2] + 245*\text{Sin}[(3*(c + d*x))/2] + 49*\text{Sin}[(5*(c + d*x))/2] - 5*\text{Sin}[(7*(c + d*x))/2]))/(d*(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]))^7)$$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 3126, 3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sin(c + dx) + a)^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (a \sin(c + dx) + a)^{7/2} dx \\ & \quad \downarrow \text{3126} \\ & \frac{12}{7} a \int (\sin(c + dx)a + a)^{5/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{5/2}}{7d} \\ & \quad \downarrow \text{3042} \\ & \frac{12}{7} a \int (\sin(c + dx)a + a)^{5/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{5/2}}{7d} \\ & \quad \downarrow \text{3126} \\ & \frac{12}{7} a \left(\frac{8}{5} a \int (\sin(c + dx)a + a)^{3/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{3/2}}{5d} \right) - \\ & \quad \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{5/2}}{7d} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{12}{7}a \left(\frac{8}{5}a \int (\sin(c+dx)a+a)^{3/2} dx - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{3/2}}{5d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{5/2}}{7d}$$

↓ 3126

$$\frac{12}{7}a \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{\sin(c+dx)a+adx} - \frac{2a \cos(c+dx)\sqrt{a \sin(c+dx)+a}}{3d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{3/2}}{5d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{5/2}}{7d}$$

↓ 3042

$$\frac{12}{7}a \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{\sin(c+dx)a+adx} - \frac{2a \cos(c+dx)\sqrt{a \sin(c+dx)+a}}{3d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{3/2}}{5d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{5/2}}{7d}$$

↓ 3125

$$\frac{12}{7}a \left(\frac{8}{5}a \left(-\frac{8a^2 \cos(c+dx)}{3d\sqrt{a \sin(c+dx)+a}} - \frac{2a \cos(c+dx)\sqrt{a \sin(c+dx)+a}}{3d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{3/2}}{5d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx)+a)^{5/2}}{7d}$$

input `Int[(a + a*Sin[c + d*x])^(7/2),x]`

output `(-2*a*Cos[c + d*x]*(a + a*Sin[c + d*x])^(5/2))/(7*d) + (12*a*((-2*a*Cos[c + d*x]*(a + a*Sin[c + d*x])^(3/2))/(5*d) + (8*a*((-8*a^2*Cos[c + d*x])/(3*d*Sqrt[a + a*Sin[c + d*x]]) - (2*a*Cos[c + d*x]*Sqrt[a + a*Sin[c + d*x]))/(3*d)))/5)/7`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.63

method	result	size
default	$\frac{2(1+\sin(dx+c))a^4(\sin(dx+c)-1)(5\sin(dx+c)^3+27\sin(dx+c)^2+71\sin(dx+c)+177)}{35\cos(dx+c)\sqrt{a+a\sin(dx+c)}d}$	75

input `int((a+a*sin(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output `2/35*(1+sin(d*x+c))*a^4*(sin(d*x+c)-1)*(5*sin(d*x+c)^3+27*sin(d*x+c)^2+71*sin(d*x+c)+177)/cos(d*x+c)/(a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.18

$$\int (a + a \sin(c + dx))^{7/2} dx = \frac{2(5a^3 \cos(dx + c)^4 + 27a^3 \cos(dx + c)^3 - 54a^3 \cos(dx + c)^2 - 204a^3 \cos(dx + c) - 128a^3 + (5a^3 \cos(dx + c)^3 - 22a^3 \cos(dx + c)^2 - 76a^3 \cos(dx + c) + 128a^3) \sin(dx + c) \sqrt{a \sin(dx + c) + a})}{35(d \cos(dx + c) + d \sin(dx + c) + d)}$$

input `integrate((a+a*sin(d*x+c))^(7/2),x, algorithm="fricas")`

output `2/35*(5*a^3*cos(d*x + c)^4 + 27*a^3*cos(d*x + c)^3 - 54*a^3*cos(d*x + c)^2 - 204*a^3*cos(d*x + c) - 128*a^3 + (5*a^3*cos(d*x + c)^3 - 22*a^3*cos(d*x + c)^2 - 76*a^3*cos(d*x + c) + 128*a^3)*sin(d*x + c))*sqrt(a*sin(d*x + c) + a)/(d*cos(d*x + c) + d*sin(d*x + c) + d)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^{7/2} dx = \text{Timed out}$$

input `integrate((a+a*sin(d*x+c))**(7/2),x)`

output `Timed out`

Maxima [F]

$$\int (a + a \sin(c + dx))^{7/2} dx = \int (a \sin(dx + c) + a)^{7/2} dx$$

input `integrate((a+a*sin(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(7/2), x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.11

$$\int (a + a \sin(c + dx))^{7/2} dx = \frac{\sqrt{2}(1225 a^3 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) + 245 a^3 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{3}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) + 49 a^3 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{5}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) + 5 a^3 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{7}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sqrt{a}}{d}$$

input `integrate((a+a*sin(d*x+c))^(7/2),x, algorithm="giac")`

output `1/140*sqrt(2)*(1225*a^3*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-1/4*pi + 1/2*d*x + 1/2*c) + 245*a^3*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-3/4*pi + 3/2*d*x + 3/2*c) + 49*a^3*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-5/4*pi + 5/2*d*x + 5/2*c) + 5*a^3*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-7/4*pi + 7/2*d*x + 7/2*c))*sqrt(a)/d`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^{7/2} dx = \int (a + a \sin(c + dx))^{7/2} dx$$

input `int((a + a*sin(c + d*x))^(7/2),x)`

output `int((a + a*sin(c + d*x))^(7/2), x)`

Reduce [F]

$$\begin{aligned} \int (a + a \sin(c + dx))^{7/2} dx &= \sqrt{a} a^3 \left(\int \sqrt{\sin(dx + c) + 1} dx \right. \\ &+ \int \sqrt{\sin(dx + c) + 1} \sin(dx + c)^3 dx + 3 \left(\int \sqrt{\sin(dx + c) + 1} \sin(dx + c)^2 dx \right) \\ &\left. + 3 \left(\int \sqrt{\sin(dx + c) + 1} \sin(dx + c) dx \right) \right) \end{aligned}$$

input `int((a+a*sin(d*x+c))^(7/2),x)`

output `sqrt(a)*a**3*(int(sqrt(sin(c + d*x) + 1),x) + int(sqrt(sin(c + d*x) + 1)*sin(c + d*x)**3,x) + 3*int(sqrt(sin(c + d*x) + 1)*sin(c + d*x)**2,x) + 3*int(sqrt(sin(c + d*x) + 1)*sin(c + d*x),x))`

3.14 $\int (a + a \sin(c + dx))^{5/2} dx$

Optimal result	137
Mathematica [A] (verified)	137
Rubi [A] (verified)	138
Maple [A] (verified)	140
Fricas [A] (verification not implemented)	140
Sympy [F]	141
Maxima [F]	141
Giac [A] (verification not implemented)	141
Mupad [F(-1)]	142
Reduce [F]	142

Optimal result

Integrand size = 14, antiderivative size = 89

$$\int (a + a \sin(c + dx))^{5/2} dx = -\frac{64a^3 \cos(c + dx)}{15d\sqrt{a + a \sin(c + dx)}} - \frac{16a^2 \cos(c + dx)\sqrt{a + a \sin(c + dx)}}{15d} - \frac{2a \cos(c + dx)(a + a \sin(c + dx))^{3/2}}{5d}$$

output

$$-64/15*a^3*\cos(d*x+c)/d/(a+a*\sin(d*x+c))^(1/2)-16/15*a^2*\cos(d*x+c)*(a+a*\sin(d*x+c))^(1/2)/d-2/5*a*\cos(d*x+c)*(a+a*\sin(d*x+c))^(3/2)/d$$

Mathematica [A] (verified)

Time = 1.94 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.31

$$\int (a + a \sin(c + dx))^{5/2} dx = \frac{(a(1 + \sin(c + dx)))^{5/2} (150 \cos(\frac{1}{2}(c + dx)) + 25 \cos(\frac{3}{2}(c + dx)) - 3 \cos(\frac{5}{2}(c + dx)) - 150 \sin(\frac{1}{2}(c + dx)))}{30d (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^5}$$

input

`Integrate[(a + a*Sin[c + d*x])^(5/2), x]`

output

```
-1/30*((a*(1 + Sin[c + d*x]))^(5/2)*(150*Cos[(c + d*x)/2] + 25*Cos[(3*(c +
d*x))/2] - 3*Cos[(5*(c + d*x))/2] - 150*Sin[(c + d*x)/2] + 25*Sin[(3*(c +
d*x))/2] + 3*Sin[(5*(c + d*x))/2]))/(d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/
2]))^5)
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sin(c + dx) + a)^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sin(c + dx) + a)^{5/2} dx \\
 & \quad \downarrow \text{3126} \\
 & \frac{8}{5} a \int (\sin(c + dx)a + a)^{3/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{5} a \int (\sin(c + dx)a + a)^{3/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3126} \\
 & \frac{8}{5} a \left(\frac{4}{3} a \int \sqrt{\sin(c + dx)a + a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) + a}}{3d} \right) - \\
 & \quad \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8}{5} a \left(\frac{4}{3} a \int \sqrt{\sin(c + dx)a + a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) + a}}{3d} \right) - \\
 & \quad \frac{2a \cos(c + dx)(a \sin(c + dx) + a)^{3/2}}{5d}
 \end{aligned}$$

$$\frac{8}{5}a \left(-\frac{8a^2 \cos(c+dx)}{3d\sqrt{a\sin(c+dx)+a}} - \frac{2a \cos(c+dx)\sqrt{a\sin(c+dx)+a}}{3d} \right) - \frac{2a \cos(c+dx)(a\sin(c+dx)+a)^{3/2}}{5d}$$

input `Int[(a + a*Sin[c + d*x])^(5/2),x]`

output `(-2*a*Cos[c + d*x]*(a + a*Sin[c + d*x])^(3/2))/(5*d) + (8*a*((-8*a^2*Cos[c + d*x])/(3*d*Sqrt[a + a*Sin[c + d*x]]) - (2*a*Cos[c + d*x]*Sqrt[a + a*Sin[c + d*x]])/(3*d)))/5`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{2(1+\sin(dx+c))a^3(\sin(dx+c)-1)(3\sin(dx+c)^2+14\sin(dx+c)+43)}{15\cos(dx+c)\sqrt{a+a\sin(dx+c)}d}$	65

input `int((a+a*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`output `2/15*(1+sin(d*x+c))*a^3*(sin(d*x+c)-1)*(3*sin(d*x+c)^2+14*sin(d*x+c)+43)/cos(d*x+c)/(a+a*sin(d*x+c))^(1/2)/d`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.29

$$\int (a + a \sin(dx + c))^{5/2} dx = \frac{2(3a^2 \cos(dx + c)^3 - 11a^2 \cos(dx + c)^2 - 46a^2 \cos(dx + c) - 32a^2 - (3a^2 \cos(dx + c)^2 + dx))^{5/2}}{15(d \cos(dx + c) + d \sin(dx + c) + d)}$$

input `integrate((a+a*sin(d*x+c))^(5/2),x, algorithm="fricas")`output `2/15*(3*a^2*cos(d*x + c)^3 - 11*a^2*cos(d*x + c)^2 - 46*a^2*cos(d*x + c) - 32*a^2 - (3*a^2*cos(d*x + c)^2 + 14*a^2*cos(d*x + c) - 32*a^2)*sin(d*x + c))*sqrt(a*sin(d*x + c) + a)/(d*cos(d*x + c) + d*sin(d*x + c) + d)`

Sympy [F]

$$\int (a + a \sin(c + dx))^{5/2} dx = \int (a \sin(c + dx) + a)^{\frac{5}{2}} dx$$

input `integrate((a+a*sin(d*x+c))**(5/2),x)`

output `Integral((a*sin(c + d*x) + a)**(5/2), x)`

Maxima [F]

$$\int (a + a \sin(c + dx))^{5/2} dx = \int (a \sin(dx + c) + a)^{\frac{5}{2}} dx$$

input `integrate((a+a*sin(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(5/2), x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.15

$$\int (a + a \sin(c + dx))^{5/2} dx = \frac{\sqrt{2}(150 a^2 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) + 25 a^2 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{3}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) + 3 a^2 \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{5}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sqrt{a}}{d}$$

input `integrate((a+a*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `1/30*sqrt(2)*(150*a^2*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-1/4*pi + 1/2*d*x + 1/2*c) + 25*a^2*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-3/4*pi + 3/2*d*x + 3/2*c) + 3*a^2*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-5/4*pi + 5/2*d*x + 5/2*c))*sqrt(a)/d`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^{5/2} dx = \int (a + a \sin(c + dx))^{5/2} dx$$

input `int((a + a*sin(c + d*x))^(5/2),x)`output `int((a + a*sin(c + d*x))^(5/2), x)`**Reduce [F]**

$$\int (a + a \sin(c + dx))^{5/2} dx = \sqrt{a} a^2 \left(\int \sqrt{\sin(dx + c) + 1} dx \right. \\ \left. + \int \sqrt{\sin(dx + c) + 1} \sin(dx + c)^2 dx + 2 \left(\int \sqrt{\sin(dx + c) + 1} \sin(dx + c) dx \right) \right)$$

input `int((a+a*sin(d*x+c))^(5/2),x)`output `sqrt(a)*a**2*(int(sqrt(sin(c + d*x) + 1),x) + int(sqrt(sin(c + d*x) + 1)*sin(c + d*x)**2,x) + 2*int(sqrt(sin(c + d*x) + 1)*sin(c + d*x),x))`

3.15 $\int (a + a \sin(c + dx))^{3/2} dx$

Optimal result	143
Mathematica [A] (verified)	143
Rubi [A] (verified)	144
Maple [A] (verified)	145
Fricas [A] (verification not implemented)	146
Sympy [F]	146
Maxima [F]	146
Giac [A] (verification not implemented)	147
Mupad [F(-1)]	147
Reduce [F]	147

Optimal result

Integrand size = 14, antiderivative size = 59

$$\int (a + a \sin(c + dx))^{3/2} dx = -\frac{8a^2 \cos(c + dx)}{3d\sqrt{a + a \sin(c + dx)}} - \frac{2a \cos(c + dx)\sqrt{a + a \sin(c + dx)}}{3d}$$

output `-8/3*a^2*cos(d*x+c)/d/(a+a*sin(d*x+c))^(1/2)-2/3*a*cos(d*x+c)*(a+a*sin(d*x+c))^(1/2)/d`

Mathematica [A] (verified)

Time = 0.76 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.51

$$\int (a + a \sin(c + dx))^{3/2} dx = \frac{(a(1 + \sin(c + dx)))^{3/2} (9 \cos(\frac{1}{2}(c + dx)) + \cos(\frac{3}{2}(c + dx)) - 9 \sin(\frac{1}{2}(c + dx)) + \sin(\frac{3}{2}(c + dx)))}{3d (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^3}$$

input `Integrate[(a + a*Sin[c + d*x])^(3/2),x]`

output

$$-1/3*((a*(1 + \sin[c + d*x]))^{3/2}*(9*\cos[(c + d*x)/2] + \cos[(3*(c + d*x))/2] - 9*\sin[(c + d*x)/2] + \sin[(3*(c + d*x))/2]))/(d*(\cos[(c + d*x)/2] + \sin[(c + d*x)/2])^3)$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sin(c + dx) + a)^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (a \sin(c + dx) + a)^{3/2} dx \\ & \quad \downarrow \text{3126} \\ & \frac{4}{3}a \int \sqrt{\sin(c + dx)a + a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) + a}}{3d} \\ & \quad \downarrow \text{3042} \\ & \frac{4}{3}a \int \sqrt{\sin(c + dx)a + a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) + a}}{3d} \\ & \quad \downarrow \text{3125} \\ & -\frac{8a^2 \cos(c + dx)}{3d\sqrt{a \sin(c + dx) + a}} - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) + a}}{3d} \end{aligned}$$

input

$$\text{Int}[(a + a*\sin[c + d*x])^{3/2},x]$$

output

$$(-8*a^2*\cos[c + d*x])/(3*d*\sqrt{a + a*\sin[c + d*x]}) - (2*a*\cos[c + d*x]*\sqrt{a + a*\sin[c + d*x]})/(3*d)$$

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :=> Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

method	result	size
default	$\frac{2(1+\sin(dx+c))a^2(\sin(dx+c)-1)(\sin(dx+c)+5)}{3\cos(dx+c)\sqrt{a+a\sin(dx+c)}d}$	53

input `int((a+a*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/3*(1+sin(d*x+c))*a^2*(sin(d*x+c)-1)*(sin(d*x+c)+5)/cos(d*x+c)/(a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.29

$$\int (a + a \sin(c + dx))^{3/2} dx = \frac{2(a \cos(dx + c))^2 + 5a \cos(dx + c) + (a \cos(dx + c) - 4a) \sin(dx + c) + 4a \sqrt{a \sin(dx + c) + a}}{3(d \cos(dx + c) + d \sin(dx + c) + d)}$$

input `integrate((a+a*sin(d*x+c))^(3/2),x, algorithm="fricas")`

output `-2/3*(a*cos(d*x + c)^2 + 5*a*cos(d*x + c) + (a*cos(d*x + c) - 4*a)*sin(d*x + c) + 4*a)*sqrt(a*sin(d*x + c) + a)/(d*cos(d*x + c) + d*sin(d*x + c) + d)`

Sympy [F]

$$\int (a + a \sin(c + dx))^{3/2} dx = \int (a \sin(c + dx) + a)^{\frac{3}{2}} dx$$

input `integrate((a+a*sin(d*x+c))**(3/2),x)`

output `Integral((a*sin(c + d*x) + a)**(3/2), x)`

Maxima [F]

$$\int (a + a \sin(c + dx))^{3/2} dx = \int (a \sin(dx + c) + a)^{\frac{3}{2}} dx$$

input `integrate((a+a*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.14

$$\int (a + a \sin(c + dx))^{3/2} dx = \frac{\sqrt{2}(9 a \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) + a \operatorname{sgn}(\cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) \sin(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c)) \sqrt{a}}{3 d}$$

input `integrate((a+a*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `1/3*sqrt(2)*(9*a*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-1/4*pi + 1/2*d*x + 1/2*c) + a*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-3/4*pi + 3/2*d*x + 3/2*c))*sqrt(a)/d`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^{3/2} dx = \int (a + a \sin(c + dx))^{3/2} dx$$

input `int((a + a*sin(c + d*x))^(3/2),x)`

output `int((a + a*sin(c + d*x))^(3/2), x)`

Reduce [F]

$$\int (a + a \sin(c + dx))^{3/2} dx = \sqrt{a} a \left(\int \sqrt{\sin(dx + c) + 1} dx + \int \sqrt{\sin(dx + c) + 1} \sin(dx + c) dx \right)$$

input `int((a+a*sin(d*x+c))^(3/2),x)`

output `sqrt(a)*a*(int(sqrt(sin(c + d*x) + 1),x) + int(sqrt(sin(c + d*x) + 1)*sin(c + d*x),x))`

3.16 $\int \sqrt{a + a \sin(c + dx)} dx$

Optimal result	149
Mathematica [B] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	151
Fricas [B] (verification not implemented)	151
Sympy [F]	151
Maxima [F]	152
Giac [A] (verification not implemented)	152
Mupad [B] (verification not implemented)	152
Reduce [F]	153

Optimal result

Integrand size = 14, antiderivative size = 26

$$\int \sqrt{a + a \sin(c + dx)} dx = -\frac{2a \cos(c + dx)}{d\sqrt{a + a \sin(c + dx)}}$$

output `-2*a*cos(d*x+c)/d/(a+a*sin(d*x+c))^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 65 vs. 2(26) = 52.

Time = 0.08 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.50

$$\int \sqrt{a + a \sin(c + dx)} dx = \frac{2(-\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) \sqrt{a(1 + \sin(c + dx))}}{d(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}$$

input `Integrate[Sqrt[a + a*Sin[c + d*x]],x]`

output `(2*(-Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*Sqrt[a*(1 + Sin[c + d*x])])/(d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a \sin(c + dx) + a} dx$$

↓ 3042

$$\int \sqrt{a \sin(c + dx) + a} dx$$

↓ 3125

$$\frac{2a \cos(c + dx)}{d \sqrt{a \sin(c + dx) + a}}$$

input `Int[Sqrt[a + a*Sin[c + d*x]],x]`

output `(-2*a*Cos[c + d*x])/(d*Sqrt[a + a*Sin[c + d*x]])`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.65

method	result	size
default	$\frac{2(1+\sin(dx+c))(\sin(dx+c)-1)a}{\cos(dx+c)\sqrt{a+a\sin(dx+c)}d}$	43
risch	$-\frac{i\sqrt{2}\sqrt{-a(-2-2\sin(dx+c))}(e^{i(dx+c)}-i)(e^{i(dx+c)}+i)}{(e^{2i(dx+c)}-1+2ie^{i(dx+c)})d}$	74

input `int((a+a*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(1+sin(d*x+c))*(sin(d*x+c)-1)*a/cos(d*x+c)/(a+a*sin(d*x+c))^(1/2)/d`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(24) = 48.

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.92

$$\int \sqrt{a + a \sin(c + dx)} dx = -\frac{2\sqrt{a \sin(dx + c) + a}(\cos(dx + c) - \sin(dx + c) + 1)}{d \cos(dx + c) + d \sin(dx + c) + d}$$

input `integrate((a+a*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output `-2*sqrt(a*sin(d*x + c) + a)*(cos(d*x + c) - sin(d*x + c) + 1)/(d*cos(d*x + c) + d*sin(d*x + c) + d)`

Sympy [F]

$$\int \sqrt{a + a \sin(c + dx)} dx = \int \sqrt{a \sin(c + dx) + a} dx$$

input `integrate((a+a*sin(d*x+c))**(1/2),x)`

output `Integral(sqrt(a*sin(c + d*x) + a), x)`

Maxima [F]

$$\int \sqrt{a + a \sin(c + dx)} dx = \int \sqrt{a \sin(dx + c) + a} dx$$

input `integrate((a+a*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(a*sin(d*x + c) + a), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.38

$$\int \sqrt{a + a \sin(c + dx)} dx = \frac{2\sqrt{2}\sqrt{a}\operatorname{sgn}\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)\right)\sin\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)}{d}$$

input `integrate((a+a*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `2*sqrt(2)*sqrt(a)*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))*sin(-1/4*pi + 1/2*d*x + 1/2*c)/d`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.27

$$\int \sqrt{a + a \sin(c + dx)} dx = -\frac{2 \cos(c + dx) \sqrt{a (\sin(c + dx) + 1)}}{d (\sin(c + dx) + 1)}$$

input `int((a + a*sin(c + d*x))^(1/2),x)`

output `-(2*cos(c + d*x)*(a*(sin(c + d*x) + 1))^(1/2))/(d*(sin(c + d*x) + 1))`

Reduce [F]

$$\int \sqrt{a + a \sin(c + dx)} dx = \sqrt{a} \left(\int \sqrt{\sin(dx + c) + 1} dx \right)$$

input `int((a+a*sin(d*x+c))^(1/2),x)`

output `sqrt(a)*int(sqrt(sin(c + d*x) + 1),x)`

3.17 $\int \frac{1}{\sqrt{a+a \sin(c+dx)}} dx$

Optimal result	154
Mathematica [C] (verified)	154
Rubi [A] (verified)	155
Maple [A] (verified)	156
Fricas [A] (verification not implemented)	157
Sympy [F]	157
Maxima [F]	158
Giac [A] (verification not implemented)	158
Mupad [B] (verification not implemented)	158
Reduce [F]	159

Optimal result

Integrand size = 14, antiderivative size = 47

$$\int \frac{1}{\sqrt{a+a \sin(c+dx)}} dx = -\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2} \sqrt{a+a \sin(c+dx)}}\right)}{\sqrt{ad}}$$

output

$-2^{(1/2)} * \operatorname{arctanh}(1/2 * a^{(1/2)} * \cos(dx+c) * 2^{(1/2)} / (a+a * \sin(dx+c))^{(1/2)}) / a^{(1/2)} / d$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.55

$$\int \frac{1}{\sqrt{a+a \sin(c+dx)}} dx = \frac{(2+2i)(-1)^{3/4} \operatorname{arctanh}\left(\left(\frac{1}{2} + \frac{i}{2}\right) (-1)^{3/4} (-1 + \tan\left(\frac{1}{4}(c+dx)\right))\right) (\cos\left(\frac{1}{2}(c+dx)\right) + \sin\left(\frac{1}{2}(c+dx)\right))}{d \sqrt{a(1 + \sin(c+dx))}}$$

input

`Integrate[1/Sqrt[a + a*Sin[c + d*x]],x]`

output

$$\left((2 + 2I)^{-3/4} \operatorname{ArcTanh}\left[\frac{1}{2} + \frac{I}{2}\right] (-1)^{-3/4} (-1 + \operatorname{Tan}\left[\frac{c + dx}{4}\right]) \right) \cdot (\operatorname{Cos}\left[\frac{c + dx}{2}\right] + \operatorname{Sin}\left[\frac{c + dx}{2}\right]) / (d \operatorname{Sqrt}[a(1 + \operatorname{Sin}[c + dx])])$$
Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{a \sin(c + dx) + a}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sqrt{a \sin(c + dx) + a}} dx \\ & \quad \downarrow \text{3128} \\ & \frac{2 \int \frac{1}{2a - \frac{a^2 \cos^2(c+dx)}{\sin(c+dx)a+a}} d \frac{a \cos(c+dx)}{\sqrt{\sin(c+dx)a+a}}}{d} \\ & \quad \downarrow \text{219} \\ & \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2} \sqrt{a \sin(c+dx) + a}}\right)}{\sqrt{ad}} \end{aligned}$$

input

$$\operatorname{Int}[1/\operatorname{Sqrt}[a + a \operatorname{Sin}[c + d*x]], x]$$

output

$$-\left(\left(\operatorname{Sqrt}[2] \operatorname{ArcTanh}\left[\frac{\operatorname{Sqrt}[a] \operatorname{Cos}[c + d*x]}{\operatorname{Sqrt}[2] \operatorname{Sqrt}[a + a \operatorname{Sin}[c + d*x]]}\right]\right)\right) / (\operatorname{Sqrt}[a] * d)$$

Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3128 Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.60

method	result	s
default	$-\frac{(1+\sin(dx+c))\sqrt{-a(\sin(dx+c)-1)}\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{-a(\sin(dx+c)-1)}\sqrt{2}}{2\sqrt{a}}\right)}{\sqrt{a}\cos(dx+c)\sqrt{a+a\sin(dx+c)}}d$	7
risch	$\frac{2i(e^{i(dx+c)}+i)\sqrt{2}e^{-i(dx+c)}}{d\sqrt{-a(ie^{2i(dx+c)}-i-2e^{i(dx+c)})}e^{-i(dx+c)}} - \frac{2i(e^{i(dx+c)}+i)\left(a^{\frac{3}{2}}+\operatorname{arctan}\left(\frac{\sqrt{-ie^{i(dx+c)}}a}{\sqrt{a}}\right)a\sqrt{-ie^{i(dx+c)}}\right)\sqrt{2}e^{-i(dx+c)}}{da^{\frac{3}{2}}\sqrt{-a(ie^{2i(dx+c)}-i-2e^{i(dx+c)})}e^{-i(dx+c)}}$	1

```
input int(1/(a+a*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -(1+sin(d*x+c))*(-a*(sin(d*x+c)-1))^(1/2)*2^(1/2)/a^(1/2)*arctanh(1/2*(-a*
(sin(d*x+c)-1))^(1/2)*2^(1/2)/a^(1/2))/cos(d*x+c)/(a+a*sin(d*x+c))^(1/2)/d
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 193, normalized size of antiderivative = 4.11

$$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx$$

$$= \left[\frac{\sqrt{2} \log \left(-\frac{\cos(dx+c)^2 - (\cos(dx+c)-2) \sin(dx+c) - 2\sqrt{2}\sqrt{a \sin(dx+c)+a}(\cos(dx+c)-\sin(dx+c)+1) + 3 \cos(dx+c)+2}{\cos(dx+c)^2 - (\cos(dx+c)+2) \sin(dx+c) - \cos(dx+c)-2} \right)}{2\sqrt{ad}}, \right.$$

$$\left. - \frac{\sqrt{2}\sqrt{-\frac{1}{a}} \arctan \left(-\frac{\sqrt{2}\sqrt{a \sin(dx+c)+a}\sqrt{-\frac{1}{a}}(\cos(dx+c)-\sin(dx+c)+1)}{2(\cos(dx+c)+\sin(dx+c)+1)} \right)}{d} \right]$$

input `integrate(1/(a+a*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output `[1/2*sqrt(2)*log(-(cos(d*x + c))^2 - (cos(d*x + c) - 2)*sin(d*x + c) - 2*sqrt(2)*sqrt(a*sin(d*x + c) + a)*(cos(d*x + c) - sin(d*x + c) + 1)/sqrt(a) + 3*cos(d*x + c) + 2)/(cos(d*x + c)^2 - (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2))/(sqrt(a)*d), -sqrt(2)*sqrt(-1/a)*arctan(-1/2*sqrt(2)*sqrt(a*sin(d*x + c) + a)*sqrt(-1/a)*(cos(d*x + c) - sin(d*x + c) + 1)/(cos(d*x + c) + sin(d*x + c) + 1))/d]`

Sympy [F]

$$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a \sin(c + dx) + a}} dx$$

input `integrate(1/(a+a*sin(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(a*sin(c + d*x) + a), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a \sin(dx + c) + a}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(a*sin(d*x + c) + a), x)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.30

$$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx$$

$$= \frac{\sqrt{2}(\log(|\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1|) - \log(|\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1|))}{2\sqrt{a}d\operatorname{sgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))}$$

input `integrate(1/(a+a*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*(log(abs(sin(-1/4*pi + 1/2*d*x + 1/2*c) + 1)) - log(abs(sin(-1/4*pi + 1/2*d*x + 1/2*c) - 1)))/(sqrt(a)*d*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c)))`

Mupad [B] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx = -\frac{F\left(\frac{\pi}{4} - \frac{c}{2} - \frac{dx}{2} \mid 1\right) \sqrt{\frac{2(a+a \sin(c+dx))}{a}}}{d \sqrt{a + a \sin(c + dx)}}$$

input `int(1/(a + a*sin(c + d*x))^(1/2),x)`

output `-(ellipticF(pi/4 - c/2 - (d*x)/2, 1)*((2*(a + a*sin(c + d*x)))/a)^(1/2))/(d*(a + a*sin(c + d*x))^(1/2))`

Reduce [F]

$$\int \frac{1}{\sqrt{a + a \sin(c + dx)}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{\sin(dx+c)+1}}{\sin(dx+c)+1} dx \right)}{a}$$

input `int(1/(a+a*sin(d*x+c))^(1/2),x)`

output `(sqrt(a)*int(sqrt(sin(c + d*x) + 1)/(sin(c + d*x) + 1),x))/a`

3.18 $\int \frac{1}{(a+a \sin(c+dx))^{3/2}} dx$

Optimal result	160
Mathematica [C] (verified)	160
Rubi [A] (verified)	161
Maple [A] (verified)	162
Fricas [B] (verification not implemented)	163
Sympy [F]	163
Maxima [F]	164
Giac [B] (verification not implemented)	164
Mupad [F(-1)]	165
Reduce [F]	165

Optimal result

Integrand size = 14, antiderivative size = 77

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2}\sqrt{a+a \sin(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{\cos(c + dx)}{2d(a + a \sin(c + dx))^{3/2}}$$

```
output -1/4*arctanh(1/2*a^(1/2)*cos(d*x+c)*2^(1/2)/(a+a*sin(d*x+c))^(1/2))*2^(1/2)
/a^(3/2)/d-1/2*cos(d*x+c)/d/(a+a*sin(d*x+c))^(3/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.40

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \frac{(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) (-\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) + 2d(a(1 + \sin(\frac{1}{2}(c + dx))))}{2d(a(1 + \sin(\frac{1}{2}(c + dx))))^2}$$

```
input Integrate[(a + a*Sin[c + d*x])^(-3/2),x]
```

output

```
((Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*(-Cos[(c + d*x)/2] + Sin[(c + d*x)/2] + (1 + I)*(-1)^(3/4)*ArcTanh[(1/2 + I/2)*(-1)^(3/4)*(-1 + Tan[(c + d*x)/4])]*(1 + Sin[c + d*x]))/(2*d*(a*(1 + Sin[c + d*x]))^(3/2))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \sin(c + dx) + a)^{3/2}} dx$$

↓ 3042

$$\int \frac{1}{(a \sin(c + dx) + a)^{3/2}} dx$$

↓ 3129

$$\frac{\int \frac{1}{\sqrt{\sin(c+dx)a+a}} dx}{4a} - \frac{\cos(c + dx)}{2d(a \sin(c + dx) + a)^{3/2}}$$

↓ 3042

$$\frac{\int \frac{1}{\sqrt{\sin(c+dx)a+a}} dx}{4a} - \frac{\cos(c + dx)}{2d(a \sin(c + dx) + a)^{3/2}}$$

↓ 3128

$$-\frac{\int \frac{1}{2a - \frac{a^2 \cos^2(c+dx)}{\sin(c+dx)a+a}} d \frac{a \cos(c+dx)}{\sqrt{\sin(c+dx)a+a}}}{2ad} - \frac{\cos(c + dx)}{2d(a \sin(c + dx) + a)^{3/2}}$$

↓ 219

$$-\frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2}\sqrt{a \sin(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{\cos(c + dx)}{2d(a \sin(c + dx) + a)^{3/2}}$$

input `Int[(a + a*Sin[c + d*x])^(-3/2),x]`

output `-1/2*ArcTanh[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[a + a*Sin[c + d*x]])]/(Sqrt[2]*a^(3/2)*d) - Cos[c + d*x]/(2*d*(a + a*Sin[c + d*x])^(3/2))`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.62

method	result
default	$-\frac{\left(\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a-a \sin(dx+c)} \sqrt{2}}{2\sqrt{a}}\right) a^2 \sin(dx+c) + \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a-a \sin(dx+c)} \sqrt{2}}{2\sqrt{a}}\right) a^2 + 2\sqrt{a-a \sin(dx+c)} a^{\frac{3}{2}}\right) \sqrt{-a(\sin(dx+c))}}{4a^{\frac{7}{2}} \cos(dx+c) \sqrt{a+a \sin(dx+c)} d}$

input `int(1/(a+a*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output

```
-1/4/a^(7/2)*(2^(1/2)*arctanh(1/2*(a-a*sin(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*
a^2*sin(d*x+c)+2^(1/2)*arctanh(1/2*(a-a*sin(d*x+c))^(1/2)*2^(1/2)/a^(1/2))
*a^2+2*(a-a*sin(d*x+c))^(1/2)*a^(3/2))*(-a*(sin(d*x+c)-1))^(1/2)/cos(d*x+c
)/(a+a*sin(d*x+c))^(1/2)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. $2(62) = 124$.

Time = 0.11 (sec) , antiderivative size = 252, normalized size of antiderivative = 3.27

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{2}(\cos(dx + c)^2 - (\cos(dx + c) + 2) \sin(dx + c) - \cos(dx + c) - 2) \sqrt{a} \log\left(\frac{\cos(dx + c) - \sin(dx + c) + 1}{\cos(dx + c) + 2}\right) + 4\sqrt{a} \sin(dx + c) - (a \cos(dx + c) - 2a) \sin(dx + c) + 2a}{8(a^2 d \cos(dx + c)^2 - a^2 d \cos(dx + c) - 2a^2 d \sin(dx + c) + a^2 d \cos(dx + c) + 2a^2 d) \sin(dx + c)}$$

input

```
integrate(1/(a+a*sin(d*x+c))^(3/2),x, algorithm="fricas")
```

output

```
1/8*(sqrt(2)*(cos(d*x + c)^2 - (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x +
c) - 2)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*sin(d*x + c) +
a)*sqrt(a)*(cos(d*x + c) - sin(d*x + c) + 1) + 3*a*cos(d*x + c) - (a*cos(d
*x + c) - 2*a)*sin(d*x + c) + 2*a)/(cos(d*x + c)^2 - (cos(d*x + c) + 2)*si
n(d*x + c) - cos(d*x + c) - 2)) + 4*sqrt(a*sin(d*x + c) + a)*(cos(d*x + c)
- sin(d*x + c) + 1))/(a^2*d*cos(d*x + c)^2 - a^2*d*cos(d*x + c) - 2*a^2*d
- (a^2*d*cos(d*x + c) + 2*a^2*d)*sin(d*x + c))
```

Sympy [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a \sin(c + dx) + a)^{3/2}} dx$$

input

```
integrate(1/(a+a*sin(d*x+c))**(3/2),x)
```

output

```
Integral((a*sin(c + d*x) + a)**(-3/2), x)
```

Maxima [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{3/2}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(-3/2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. $2(62) = 124$.

Time = 0.13 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.73

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{2} \left(\frac{\log(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1)}{\operatorname{asgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} - \frac{\log(-\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1)}{\operatorname{asgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} - \frac{2 \sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c)}{(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))^2} \right)}{8\sqrt{ad}}$$

input `integrate(1/(a+a*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `1/8*sqrt(2)*(log(sin(-1/4*pi + 1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))) - log(-sin(-1/4*pi + 1/2*d*x + 1/2*c) + 1)/(a*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))) - 2*sin(-1/4*pi + 1/2*d*x + 1/2*c)/((sin(-1/4*pi + 1/2*d*x + 1/2*c))^2 - 1)*a*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c)))/(sqrt(a)*d)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx$$

input `int(1/(a + a*sin(c + d*x))^(3/2),x)`output `int(1/(a + a*sin(c + d*x))^(3/2), x)`**Reduce [F]**

$$\int \frac{1}{(a + a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{\sin(dx+c)+1}}{\sin(dx+c)^2 + 2\sin(dx+c)+1} dx \right)}{a^2}$$

input `int(1/(a+a*sin(d*x+c))^(3/2),x)`output `(sqrt(a)*int(sqrt(sin(c + d*x) + 1)/(sin(c + d*x)**2 + 2*sin(c + d*x) + 1),x))/a**2`

3.19 $\int \frac{1}{(a+a \sin(c+dx))^{5/2}} dx$

Optimal result	166
Mathematica [C] (verified)	166
Rubi [A] (verified)	167
Maple [B] (verified)	169
Fricas [B] (verification not implemented)	169
Sympy [F]	170
Maxima [F]	170
Giac [A] (verification not implemented)	171
Mupad [F(-1)]	171
Reduce [F]	171

Optimal result

Integrand size = 14, antiderivative size = 107

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = -\frac{3 \arctanh\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2} \sqrt{a+a \sin(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} - \frac{\cos(c + dx)}{4d(a + a \sin(c + dx))^{5/2}} - \frac{3 \cos(c + dx)}{16ad(a + a \sin(c + dx))^{3/2}}$$

output `-3/32*arctanh(1/2*a^(1/2)*cos(d*x+c)*2^(1/2)/(a+a*sin(d*x+c))^(1/2))*2^(1/2)/a^(5/2)/d-1/4*cos(d*x+c)/d/(a+a*sin(d*x+c))^(5/2)-3/16*cos(d*x+c)/a/d/(a+a*sin(d*x+c))^(3/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.83

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \frac{(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) (8 \sin(\frac{1}{2}(c + dx)) - 4(\cos(\frac{1}{2}(c + dx))))}{(a + a \sin(c + dx))^{5/2}}$$

input `Integrate[(a + a*Sin[c + d*x])^(-5/2),x]`

output

```
((Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*(8*Sin[(c + d*x)/2] - 4*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) + 6*Sin[(c + d*x)/2]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 - 3*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^3 + (3 + 3*I)*(-1)^(3/4)*ArcTanh[(1/2 + I/2)*(-1)^(3/4)*(-1 + Tan[(c + d*x)/4])]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^4)/(16*d*(a*(1 + Sin[c + d*x]))^(5/2))
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3129, 3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \sin(c + dx) + a)^{5/2}} dx$$

↓ 3042

$$\int \frac{1}{(a \sin(c + dx) + a)^{5/2}} dx$$

↓ 3129

$$\frac{3 \int \frac{1}{(\sin(c+dx)a+a)^{3/2}} dx}{8a} - \frac{\cos(c + dx)}{4d(a \sin(c + dx) + a)^{5/2}}$$

↓ 3042

$$\frac{3 \int \frac{1}{(\sin(c+dx)a+a)^{3/2}} dx}{8a} - \frac{\cos(c + dx)}{4d(a \sin(c + dx) + a)^{5/2}}$$

↓ 3129

$$\frac{3 \left(\frac{\int \frac{1}{\sqrt{\sin(c+dx)a+a}} dx}{4a} - \frac{\cos(c+dx)}{2d(a \sin(c+dx)+a)^{3/2}} \right)}{8a} - \frac{\cos(c + dx)}{4d(a \sin(c + dx) + a)^{5/2}}$$

↓ 3042

$$\frac{3 \left(\frac{\int \frac{1}{\sqrt{\sin(c+dx)a+a}} dx}{4a} - \frac{\cos(c+dx)}{2d(a \sin(c+dx)+a)^{3/2}} \right)}{8a} - \frac{\cos(c + dx)}{4d(a \sin(c + dx) + a)^{5/2}}$$

$$\begin{array}{c}
 \downarrow \text{3128} \\
 3 \left(\frac{\int \frac{1}{2a - \frac{a^2 \cos^2(c+dx)}{\sin(c+dx)a+a}} d \frac{a \cos(c+dx)}{\sqrt{\sin(c+dx)a+a}}}{2ad} - \frac{\cos(c+dx)}{2d(a \sin(c+dx)+a)^{3/2}} \right) \\
 \hline
 8a - \frac{\cos(c+dx)}{4d(a \sin(c+dx)+a)^{5/2}} \\
 \downarrow \text{219} \\
 3 \left(-\frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2}\sqrt{a \sin(c+dx)+a}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{\cos(c+dx)}{2d(a \sin(c+dx)+a)^{3/2}} \right) \\
 \hline
 8a - \frac{\cos(c+dx)}{4d(a \sin(c+dx)+a)^{5/2}}
 \end{array}$$

input `Int[(a + a*Sin[c + d*x])^(-5/2),x]`

output `-1/4*Cos[c + d*x]/(d*(a + a*Sin[c + d*x])^(5/2)) + (3*(-1/2*ArcTanh[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[a + a*Sin[c + d*x]])]/(Sqrt[2]*a^(3/2)*d) - Cos[c + d*x]/(2*d*(a + a*Sin[c + d*x])^(3/2)))/(8*a)`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c
+ d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n
+ 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x]
&& EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. $2(88) = 176$.

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.86

method	result
default	$-\frac{\left(-3\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a-a\sin(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)\right)a^2 \cos(dx+c)^2 + 6\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a-a\sin(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)a^2 \sin(dx+c) + 6\sqrt{a-a\sin(dx+c)}a^{\frac{3}{2}}}{32a^{\frac{9}{2}}(1+\sin(dx+c))\cos(dx+c)\sqrt{a+a\sin(dx+c)}}$

input

```
int(1/(a+a*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
-1/32/a^(9/2)*(-3*2^(1/2)*arctanh(1/2*(a-a*sin(d*x+c))^(1/2)*2^(1/2)/a^(1/2))
)*a^2*cos(d*x+c)^2+6*2^(1/2)*arctanh(1/2*(a-a*sin(d*x+c))^(1/2)*2^(1/2)/
a^(1/2))*a^2*sin(d*x+c)+6*(a-a*sin(d*x+c))^(1/2)*a^(3/2)*sin(d*x+c)+6*2^(1/2)
*arctanh(1/2*(a-a*sin(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*a^2+14*(a-a*sin(d*
x+c))^(1/2)*a^(3/2)*(-a*(sin(d*x+c)-1))^(1/2)/(1+sin(d*x+c))/cos(d*x+c)/(
a+a*sin(d*x+c))^(1/2)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. $2(88) = 176$.

Time = 0.11 (sec) , antiderivative size = 320, normalized size of antiderivative = 2.99

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \frac{3\sqrt{2}(\cos(dx + c)^3 + 3 \cos(dx + c)^2 + (\cos(dx + c))^2 - 2 \cos(dx + c) - 4)}{\dots}$$

input

```
integrate(1/(a+a*sin(d*x+c))^(5/2),x, algorithm="fricas")
```

output

```
1/64*(3*sqrt(2)*(cos(d*x + c)^3 + 3*cos(d*x + c)^2 + (cos(d*x + c)^2 - 2*cos(d*x + c) - 4)*sin(d*x + c) - 2*cos(d*x + c) - 4)*sqrt(a)*log(-(a*cos(d*x + c)^2 - 2*sqrt(2)*sqrt(a*sin(d*x + c) + a)*sqrt(a)*(cos(d*x + c) - sin(d*x + c) + 1) + 3*a*cos(d*x + c) - (a*cos(d*x + c) - 2*a)*sin(d*x + c) + 2*a)/(cos(d*x + c)^2 - (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2)) + 4*(3*cos(d*x + c)^2 + (3*cos(d*x + c) - 4)*sin(d*x + c) + 7*cos(d*x + c) + 4)*sqrt(a*sin(d*x + c) + a)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d + (a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d)*sin(d*x + c))
```

Sympy [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a \sin(c + dx) + a)^{5/2}} dx$$

input

```
integrate(1/(a+a*sin(d*x+c))**(5/2),x)
```

output

```
Integral((a*sin(c + d*x) + a)**(-5/2), x)
```

Maxima [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{5/2}} dx$$

input

```
integrate(1/(a+a*sin(d*x+c))^(5/2),x, algorithm="maxima")
```

output

```
integrate((a*sin(d*x + c) + a)^(-5/2), x)
```

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.43

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \frac{\sqrt{2} \left(\frac{3 \log(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1)}{a^2 \operatorname{sgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} - \frac{3 \log(-\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1)}{a^2 \operatorname{sgn}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} - \frac{2 \left(3 \sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) \right)^3 - 5 \sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c)}{\left(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) \right)^2 - 1} \right)}{64 \sqrt{ad}}$$

input `integrate(1/(a+a*sin(d*x+c))^(5/2),x, algorithm="giac")`output `1/64*sqrt(2)*(3*log(sin(-1/4*pi + 1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))) - 3*log(-sin(-1/4*pi + 1/2*d*x + 1/2*c) + 1)/(a^2*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c))) - 2*(3*sin(-1/4*pi + 1/2*d*x + 1/2*c))^3 - 5*sin(-1/4*pi + 1/2*d*x + 1/2*c))/((sin(-1/4*pi + 1/2*d*x + 1/2*c))^2 - 1)^2*a^2*sgn(cos(-1/4*pi + 1/2*d*x + 1/2*c)))/(sqrt(a)*d)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx$$

input `int(1/(a + a*sin(c + d*x))^(5/2),x)`output `int(1/(a + a*sin(c + d*x))^(5/2), x)`**Reduce [F]**

$$\int \frac{1}{(a + a \sin(c + dx))^{5/2}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{\sin(dx+c)+1}}{\sin(dx+c)^3 + 3 \sin(dx+c)^2 + 3 \sin(dx+c) + 1} dx \right)}{a^3}$$

input `int(1/(a+a*sin(d*x+c))^(5/2),x)`

output $(\sqrt{a} \cdot \text{int}(\sqrt{\sin(c + d \cdot x) + 1} / (\sin(c + d \cdot x)^3 + 3 \sin(c + d \cdot x)^2 + 3 \sin(c + d \cdot x) + 1), x)) / a^3$

3.20 $\int (a - a \sin(c + dx))^{7/2} dx$

Optimal result	173
Mathematica [A] (verified)	173
Rubi [A] (verified)	174
Maple [A] (verified)	176
Fricas [A] (verification not implemented)	177
Sympy [F(-1)]	177
Maxima [F]	177
Giac [A] (verification not implemented)	178
Mupad [F(-1)]	178
Reduce [F]	179

Optimal result

Integrand size = 15, antiderivative size = 123

$$\int (a - a \sin(c + dx))^{7/2} dx = \frac{256a^4 \cos(c + dx)}{35d\sqrt{a - a \sin(c + dx)}} + \frac{64a^3 \cos(c + dx)\sqrt{a - a \sin(c + dx)}}{35d} + \frac{24a^2 \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{35d} + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d}$$

output

```
256/35*a^4*cos(d*x+c)/d/(a-a*sin(d*x+c))^(1/2)+64/35*a^3*cos(d*x+c)*(a-a*
sin(d*x+c))^(1/2)/d+24/35*a^2*cos(d*x+c)*(a-a*sin(d*x+c))^(3/2)/d+2/7*a*cos
(d*x+c)*(a-a*sin(d*x+c))^(5/2)/d
```

Mathematica [A] (verified)

Time = 3.50 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.28

$$\int (a - a \sin(c + dx))^{7/2} dx = \frac{a^3(-1 + \sin(c + dx))^3 \sqrt{a - a \sin(c + dx)} (1225 \cos(\frac{1}{2}(c + dx)) + 245 \cos(\frac{3}{2}(c + dx)) - 49 \cos(\frac{5}{2}(c + dx)))}{140d (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}$$

input `Integrate[(a - a*Sin[c + d*x])^(7/2),x]`

output
$$-1/140*(a^3*(-1 + \sin[c + d*x])^3*\sqrt{a - a*\sin[c + d*x]}*(1225*\cos[(c + d*x)/2] + 245*\cos[(3*(c + d*x))/2] - 49*\cos[(5*(c + d*x))/2] - 5*\cos[(7*(c + d*x))/2] + 1225*\sin[(c + d*x)/2] - 245*\sin[(3*(c + d*x))/2] - 49*\sin[(5*(c + d*x))/2] + 5*\sin[(7*(c + d*x))/2]))/(d*(\cos[(c + d*x)/2] - \sin[(c + d*x)/2]))^7$$

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3042, 3126, 3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a - a \sin(c + dx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (a - a \sin(c + dx))^{7/2} dx \\ & \quad \downarrow \text{3126} \\ & \frac{12}{7}a \int (a - a \sin(c + dx))^{5/2} dx + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d} \\ & \quad \downarrow \text{3042} \\ & \frac{12}{7}a \int (a - a \sin(c + dx))^{5/2} dx + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d} \\ & \quad \downarrow \text{3126} \\ & \frac{12}{7}a \left(\frac{8}{5}a \int (a - a \sin(c + dx))^{3/2} dx + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \right) + \\ & \quad \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{12}{7}a \left(\frac{8}{5}a \int (a - a \sin(c + dx))^{3/2} dx + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d}$$

↓ 3126

$$\frac{12}{7}a \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{a - a \sin(c + dx)} dx + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d}$$

↓ 3042

$$\frac{12}{7}a \left(\frac{8}{5}a \left(\frac{4}{3}a \int \sqrt{a - a \sin(c + dx)} dx + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d}$$

↓ 3125

$$\frac{12}{7}a \left(\frac{8}{5}a \left(\frac{8a^2 \cos(c + dx)}{3d \sqrt{a - a \sin(c + dx)}} + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \right) + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{5/2}}{7d}$$

input `Int[(a - a*Sin[c + d*x])^(7/2),x]`

output `(2*a*Cos[c + d*x]*(a - a*Sin[c + d*x])^(5/2))/(7*d) + (12*a*((2*a*Cos[c + d*x]*(a - a*Sin[c + d*x])^(3/2))/(5*d) + (8*a*((8*a^2*Cos[c + d*x])/(3*d*Sqrt[a - a*Sin[c + d*x]]) + (2*a*Cos[c + d*x]*Sqrt[a - a*Sin[c + d*x]])/(3*d))))/5)/7`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.62

method	result	size
default	$\frac{2(\sin(dx+c)-1)a^4(1+\sin(dx+c))(5\sin(dx+c)^3-27\sin(dx+c)^2+71\sin(dx+c)-177)}{35\cos(dx+c)\sqrt{a-a\sin(dx+c)}d}$	76

input `int((a-a*sin(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output `2/35*(sin(d*x+c)-1)*a^4*(1+sin(d*x+c))*(5*sin(d*x+c)^3-27*sin(d*x+c)^2+71*sin(d*x+c)-177)/cos(d*x+c)/(a-a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.16

$$\int (a - a \sin(c + dx))^{7/2} dx = \frac{2(5a^3 \cos(dx + c)^4 + 27a^3 \cos(dx + c)^3 - 54a^3 \cos(dx + c)^2 - 204a^3 \cos(dx + c) - 128a^3 - (5a^3 \cos(dx + c) - d \sin(dx + c)) \sqrt{a - a \sin(dx + c)})}{35(d \cos(dx + c) - d \sin(dx + c))}$$

input `integrate((a-a*sin(d*x+c))^(7/2),x, algorithm="fricas")`

output `-2/35*(5*a^3*cos(d*x + c)^4 + 27*a^3*cos(d*x + c)^3 - 54*a^3*cos(d*x + c)^2 - 204*a^3*cos(d*x + c) - 128*a^3 - (5*a^3*cos(d*x + c) - 22*a^3*cos(d*x + c)^2 - 76*a^3*cos(d*x + c) + 128*a^3)*sin(d*x + c))*sqrt(-a*sin(d*x + c) + a)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`

Sympy [F(-1)]

Timed out.

$$\int (a - a \sin(c + dx))^{7/2} dx = \text{Timed out}$$

input `integrate((a-a*sin(d*x+c))**(7/2),x)`

output `Timed out`

Maxima [F]

$$\int (a - a \sin(c + dx))^{7/2} dx = \int (-a \sin(dx + c) + a)^{7/2} dx$$

input `integrate((a-a*sin(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((-a*sin(d*x + c) + a)^(7/2), x)`

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.07

$$\int (a - a \sin(c + dx))^{7/2} dx = \frac{\sqrt{2}(1225 a^3 \cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) \operatorname{sgn}(\sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) - 245 a^3 \cos(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c) \operatorname{sgn}(\sin(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c)) + 49 a^3 \cos(-\frac{5}{4} \pi + \frac{5}{2} dx + \frac{5}{2} c) \operatorname{sgn}(\sin(-\frac{5}{4} \pi + \frac{5}{2} dx + \frac{5}{2} c)) - 5 a^3 \cos(-\frac{7}{4} \pi + \frac{7}{2} dx + \frac{7}{2} c) \operatorname{sgn}(\sin(-\frac{7}{4} \pi + \frac{7}{2} dx + \frac{7}{2} c))) \sqrt{a}}{d}$$

input `integrate((a-a*sin(d*x+c))^(7/2),x, algorithm="giac")`

output `-1/140*sqrt(2)*(1225*a^3*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - 245*a^3*cos(-3/4*pi + 3/2*d*x + 3/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) + 49*a^3*cos(-5/4*pi + 5/2*d*x + 5/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - 5*a^3*cos(-7/4*pi + 7/2*d*x + 7/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))*sqrt(a)/d`

Mupad [F(-1)]

Timed out.

$$\int (a - a \sin(c + dx))^{7/2} dx = \int (a - a \sin(c + dx))^{7/2} dx$$

input `int((a - a*sin(c + d*x))^(7/2),x)`

output `int((a - a*sin(c + d*x))^(7/2), x)`

Reduce [F]

$$\int (a - a \sin(c + dx))^{7/2} dx = \sqrt{a} a^3 \left(\int \sqrt{-\sin(dx + c) + 1} dx \right. \\ \left. - \left(\int \sqrt{-\sin(dx + c) + 1} \sin(dx + c)^3 dx \right) \right. \\ \left. + 3 \left(\int \sqrt{-\sin(dx + c) + 1} \sin(dx + c)^2 dx \right) \right. \\ \left. - 3 \left(\int \sqrt{-\sin(dx + c) + 1} \sin(dx + c) dx \right) \right)$$

input `int((a-a*sin(d*x+c))^(7/2),x)`

output `sqrt(a)*a**3*(int(sqrt(-sin(c+d*x)+1),x) - int(sqrt(-sin(c+d*x)+1)*sin(c+d*x)**3,x) + 3*int(sqrt(-sin(c+d*x)+1)*sin(c+d*x)**2,x) - 3*int(sqrt(-sin(c+d*x)+1)*sin(c+d*x),x))`

3.21 $\int (a - a \sin(c + dx))^{5/2} dx$

Optimal result	180
Mathematica [A] (verified)	180
Rubi [A] (verified)	181
Maple [A] (verified)	183
Fricas [A] (verification not implemented)	183
Sympy [F]	184
Maxima [F]	184
Giac [A] (verification not implemented)	184
Mupad [F(-1)]	185
Reduce [F]	185

Optimal result

Integrand size = 15, antiderivative size = 92

$$\int (a - a \sin(c + dx))^{5/2} dx = \frac{64a^3 \cos(c + dx)}{15d\sqrt{a - a \sin(c + dx)}} + \frac{16a^2 \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{15d} + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d}$$

output `64/15*a^3*cos(d*x+c)/d/(a-a*sin(d*x+c))^(1/2)+16/15*a^2*cos(d*x+c)*(a-a*sin(d*x+c))^(1/2)/d+2/5*a*cos(d*x+c)*(a-a*sin(d*x+c))^(3/2)/d`

Mathematica [A] (verified)

Time = 2.25 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.45

$$\int (a - a \sin(c + dx))^{5/2} dx = \frac{a^2(-1 + \sin(c + dx))^2 \sqrt{a - a \sin(c + dx)} (150 \cos(\frac{1}{2}(c + dx)) + 25 \cos(\frac{3}{2}(c + dx)) - 3 \cos(\frac{5}{2}(c + dx)))}{30d (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}$$

input `Integrate[(a - a*Sin[c + d*x])^(5/2),x]`

output

```
(a^2*(-1 + Sin[c + d*x])^2*sqrt[a - a*Sin[c + d*x]]*(150*cos[(c + d*x)/2]
+ 25*cos[(3*(c + d*x))/2] - 3*cos[(5*(c + d*x))/2] + 150*sin[(c + d*x)/2]
- 25*sin[(3*(c + d*x))/2] - 3*sin[(5*(c + d*x))/2]))/(30*d*(cos[(c + d*x)/
2] - sin[(c + d*x)/2])^5)
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (a - a \sin(c + dx))^{5/2} dx \\
& \quad \downarrow \text{3042} \\
& \int (a - a \sin(c + dx))^{5/2} dx \\
& \quad \downarrow \text{3126} \\
& \frac{8}{5} a \int (a - a \sin(c + dx))^{3/2} dx + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{8}{5} a \int (a - a \sin(c + dx))^{3/2} dx + \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \\
& \quad \downarrow \text{3126} \\
& \frac{8}{5} a \left(\frac{4}{3} a \int \sqrt{a - a \sin(c + dx)} dx + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \right) + \\
& \quad \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d} \\
& \quad \downarrow \text{3042} \\
& \frac{8}{5} a \left(\frac{4}{3} a \int \sqrt{a - a \sin(c + dx)} dx + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \right) + \\
& \quad \frac{2a \cos(c + dx)(a - a \sin(c + dx))^{3/2}}{5d}
\end{aligned}$$

$$\frac{8}{5}a \left(\frac{8a^2 \cos(c+dx)}{3d\sqrt{a-a\sin(c+dx)}} + \frac{2a \cos(c+dx)\sqrt{a-a\sin(c+dx)}}{3d} \right) + \frac{2a \cos(c+dx)(a-a\sin(c+dx))^{3/2}}{5d}$$

input `Int[(a - a*Sin[c + d*x])^(5/2),x]`

output `(2*a*Cos[c + d*x]*(a - a*Sin[c + d*x])^(3/2))/(5*d) + (8*a*((8*a^2*Cos[c + d*x])/(3*d*Sqrt[a - a*Sin[c + d*x]]) + (2*a*Cos[c + d*x]*Sqrt[a - a*Sin[c + d*x]])/(3*d)))/5`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.72

method	result	size
default	$-\frac{2(\sin(dx+c)-1)a^3(1+\sin(dx+c))(3\sin(dx+c)^2-14\sin(dx+c)+43)}{15\cos(dx+c)\sqrt{a-a\sin(dx+c)}d}$	66

input `int((a-a*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `-2/15*(sin(d*x+c)-1)*a^3*(1+sin(d*x+c))*(3*sin(d*x+c)^2-14*sin(d*x+c)+43)/cos(d*x+c)/(a-a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.26

$$\int (a - a \sin(c + dx))^{5/2} dx = \frac{2(3a^2 \cos(dx + c)^3 - 11a^2 \cos(dx + c)^2 - 46a^2 \cos(dx + c) - 32a^2 + (3a^2 \cos(dx + c)^2 + 14a^2 \cos(dx + c)) \sqrt{-a \sin(dx + c) + a}}{15(d \cos(dx + c) - d \sin(dx + c) + d)}$$

input `integrate((a-a*sin(d*x+c))^(5/2),x, algorithm="fricas")`

output `-2/15*(3*a^2*cos(d*x + c)^3 - 11*a^2*cos(d*x + c)^2 - 46*a^2*cos(d*x + c) - 32*a^2 + (3*a^2*cos(d*x + c)^2 + 14*a^2*cos(d*x + c) - 32*a^2)*sin(d*x + c))*sqrt(-a*sin(d*x + c) + a)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`

Sympy [F]

$$\int (a - a \sin(c + dx))^{5/2} dx = \int (-a \sin(c + dx) + a)^{\frac{5}{2}} dx$$

input `integrate((a-a*sin(d*x+c))**(5/2),x)`

output `Integral((-a*sin(c + d*x) + a)**(5/2), x)`

Maxima [F]

$$\int (a - a \sin(c + dx))^{5/2} dx = \int (-a \sin(dx + c) + a)^{\frac{5}{2}} dx$$

input `integrate((a-a*sin(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((-a*sin(d*x + c) + a)^(5/2), x)`

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.11

$$\int (a - a \sin(c + dx))^{5/2} dx = \frac{\sqrt{2}(150 a^2 \cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) \operatorname{sgn}(\sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) - 25 a^2 \cos(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c) \operatorname{sgn}(\sin(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c)))}{30 d}$$

input `integrate((a-a*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `-1/30*sqrt(2)*(150*a^2*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - 25*a^2*cos(-3/4*pi + 3/2*d*x + 3/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) + 3*a^2*cos(-5/4*pi + 5/2*d*x + 5/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))*sqrt(a)/d`

Mupad [F(-1)]

Timed out.

$$\int (a - a \sin(c + dx))^{5/2} dx = \int (a - a \sin(c + dx))^{5/2} dx$$

input `int((a - a*sin(c + d*x))^(5/2),x)`output `int((a - a*sin(c + d*x))^(5/2), x)`**Reduce [F]**

$$\begin{aligned} \int (a - a \sin(c + dx))^{5/2} dx &= \sqrt{a} a^2 \left(\int \sqrt{-\sin(dx + c) + 1} dx \right. \\ &+ \int \sqrt{-\sin(dx + c) + 1} \sin(dx + c)^2 dx \\ &\left. - 2 \left(\int \sqrt{-\sin(dx + c) + 1} \sin(dx + c) dx \right) \right) \end{aligned}$$

input `int((a-a*sin(d*x+c))^(5/2),x)`output `sqrt(a)*a**2*(int(sqrt(-sin(c + d*x) + 1),x) + int(sqrt(-sin(c + d*x) + 1)*sin(c + d*x)**2,x) - 2*int(sqrt(-sin(c + d*x) + 1)*sin(c + d*x),x))`

3.22 $\int (a - a \sin(c + dx))^{3/2} dx$

Optimal result	186
Mathematica [A] (verified)	186
Rubi [A] (verified)	187
Maple [A] (verified)	188
Fricas [A] (verification not implemented)	189
Sympy [F]	189
Maxima [F]	189
Giac [A] (verification not implemented)	190
Mupad [F(-1)]	190
Reduce [F]	190

Optimal result

Integrand size = 15, antiderivative size = 61

$$\int (a - a \sin(c + dx))^{3/2} dx = \frac{8a^2 \cos(c + dx)}{3d\sqrt{a - a \sin(c + dx)}} + \frac{2a \cos(c + dx)\sqrt{a - a \sin(c + dx)}}{3d}$$

output

```
8/3*a^2*cos(d*x+c)/d/(a-a*sin(d*x+c))^(1/2)+2/3*a*cos(d*x+c)*(a-a*sin(d*x+c))^(1/2)/d
```

Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.69

$$\int (a - a \sin(c + dx))^{3/2} dx = \frac{a(-1 + \sin(c + dx))\sqrt{a - a \sin(c + dx)}(9 \cos(\frac{1}{2}(c + dx)) + \cos(\frac{3}{2}(c + dx)) + 9 \sin(\frac{1}{2}(c + dx)) - \sin(\frac{3}{2}(c + dx)))}{3d(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^3}$$

input

```
Integrate[(a - a*Sin[c + d*x])^(3/2),x]
```

output

$$-1/3*(a*(-1 + \sin[c + dx])*sqrt[a - a*\sin[c + dx]]*(9*\cos[(c + dx)/2] + \cos[(3*(c + dx))/2] + 9*\sin[(c + dx)/2] - \sin[(3*(c + dx))/2]))/(d*(\cos[(c + dx)/2] - \sin[(c + dx)/2])^3)$$
Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a - a \sin(c + dx))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (a - a \sin(c + dx))^{3/2} dx \\ & \quad \downarrow \text{3126} \\ & \frac{4}{3}a \int \sqrt{a - a \sin(c + dx)} dx + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \\ & \quad \downarrow \text{3042} \\ & \frac{4}{3}a \int \sqrt{a - a \sin(c + dx)} dx + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \\ & \quad \downarrow \text{3125} \\ & \frac{8a^2 \cos(c + dx)}{3d \sqrt{a - a \sin(c + dx)}} + \frac{2a \cos(c + dx) \sqrt{a - a \sin(c + dx)}}{3d} \end{aligned}$$

input

$$\text{Int}[(a - a*\sin[c + dx])^(3/2),x]$$

output

$$(8*a^2*\cos[c + dx])/(3*d*sqrt[a - a*\sin[c + dx]]) + (2*a*\cos[c + dx]*sqrt[a - a*\sin[c + dx]])/(3*d)$$

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{2(\sin(dx+c)-1)a^2(1+\sin(dx+c))(\sin(dx+c)-5)}{3\cos(dx+c)\sqrt{a-a\sin(dx+c)}d}$	54

input `int((a-a*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/3*(sin(d*x+c)-1)*a^2*(1+sin(d*x+c))*(sin(d*x+c)-5)/cos(d*x+c)/(a-a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.30

$$\int (a - a \sin(c + dx))^{3/2} dx = \frac{2(a \cos(dx + c)^2 + 5a \cos(dx + c) - (a \cos(dx + c) - 4a) \sin(dx + c) + 4a) \sqrt{-a \sin(dx + c)}}{3(d \cos(dx + c) - d \sin(dx + c) + d)}$$

input `integrate((a-a*sin(d*x+c))^(3/2),x, algorithm="fricas")`

output `2/3*(a*cos(d*x + c)^2 + 5*a*cos(d*x + c) - (a*cos(d*x + c) - 4*a)*sin(d*x + c) + 4*a)*sqrt(-a*sin(d*x + c) + a)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`

Sympy [F]

$$\int (a - a \sin(c + dx))^{3/2} dx = \int (-a \sin(c + dx) + a)^{\frac{3}{2}} dx$$

input `integrate((a-a*sin(d*x+c))**(3/2),x)`

output `Integral((-a*sin(c + d*x) + a)**(3/2), x)`

Maxima [F]

$$\int (a - a \sin(c + dx))^{3/2} dx = \int (-a \sin(dx + c) + a)^{\frac{3}{2}} dx$$

input `integrate((a-a*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((-a*sin(d*x + c) + a)^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.11

$$\int (a - a \sin(c + dx))^{3/2} dx = \frac{\sqrt{2} \left(9 a \cos \left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c \right) \operatorname{sgn} \left(\sin \left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c \right) \right) - a \cos \left(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c \right) \operatorname{sgn} \left(\sin \left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c \right) \right) \right)}{3 d}$$

input `integrate((a-a*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `-1/3*sqrt(2)*(9*a*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - a*cos(-3/4*pi + 3/2*d*x + 3/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))*sqrt(a)/d`

Mupad [F(-1)]

Timed out.

$$\int (a - a \sin(c + dx))^{3/2} dx = \int (a - a \sin(c + dx))^{3/2} dx$$

input `int((a - a*sin(c + d*x))^(3/2),x)`

output `int((a - a*sin(c + d*x))^(3/2), x)`

Reduce [F]

$$\int (a - a \sin(c + dx))^{3/2} dx = \sqrt{a} a \left(\int \sqrt{-\sin(dx + c) + 1} dx - \left(\int \sqrt{-\sin(dx + c) + 1} \sin(dx + c) dx \right) \right)$$

input `int((a-a*sin(d*x+c))^(3/2),x)`

output `sqrt(a)*a*(int(sqrt(-sin(c+d*x)+1),x) - int(sqrt(-sin(c+d*x)+1)*sin(c+d*x),x))`

3.23 $\int \sqrt{a - a \sin(c + dx)} dx$

Optimal result	192
Mathematica [B] (verified)	192
Rubi [A] (verified)	193
Maple [A] (verified)	194
Fricas [A] (verification not implemented)	194
Sympy [F]	194
Maxima [F]	195
Giac [A] (verification not implemented)	195
Mupad [B] (verification not implemented)	195
Reduce [F]	196

Optimal result

Integrand size = 15, antiderivative size = 27

$$\int \sqrt{a - a \sin(c + dx)} dx = \frac{2a \cos(c + dx)}{d \sqrt{a - a \sin(c + dx)}}$$

output `2*a*cos(d*x+c)/d/(a-a*sin(d*x+c))^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 66 vs. $2(27) = 54$.

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.44

$$\int \sqrt{a - a \sin(c + dx)} dx = \frac{2(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) \sqrt{a - a \sin(c + dx)}}{d(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}$$

input `Integrate[Sqrt[a - a*Sin[c + d*x]],x]`

output `(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*Sqrt[a - a*Sin[c + d*x]])/(d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a - a \sin(c + dx)} dx$$

↓ 3042

$$\int \sqrt{a - a \sin(c + dx)} dx$$

↓ 3125

$$\frac{2a \cos(c + dx)}{d\sqrt{a - a \sin(c + dx)}}$$

input `Int[Sqrt[a - a*Sin[c + d*x]],x]`

output `(2*a*Cos[c + d*x])/(d*Sqrt[a - a*Sin[c + d*x]])`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.63

method	result	size
default	$-\frac{2(\sin(dx+c)-1)(1+\sin(dx+c))a}{\cos(dx+c)\sqrt{a-a\sin(dx+c)}d}$	44
risch	$-\frac{i\sqrt{2}\sqrt{a(2-2\sin(dx+c))}(e^{i(dx+c)}+i)(e^{i(dx+c)}-i)}{(e^{2i(dx+c)}-2ie^{i(dx+c)}-1)d}$	73

input `int((a-a*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`output `-2*(sin(d*x+c)-1)*(1+sin(d*x+c))*a/cos(d*x+c)/(a-a*sin(d*x+c))^(1/2)/d`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.85

$$\int \sqrt{a - a \sin(c + dx)} dx = \frac{2 \sqrt{-a \sin(dx + c) + a} (\cos(dx + c) + \sin(dx + c) + 1)}{d \cos(dx + c) - d \sin(dx + c) + d}$$

input `integrate((a-a*sin(d*x+c))^(1/2),x, algorithm="fricas")`output `2*sqrt(-a*sin(d*x + c) + a)*(cos(d*x + c) + sin(d*x + c) + 1)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`**Sympy [F]**

$$\int \sqrt{a - a \sin(c + dx)} dx = \int \sqrt{-a \sin(c + dx) + a} dx$$

input `integrate((a-a*sin(d*x+c))**(1/2),x)`output `Integral(sqrt(-a*sin(c + d*x) + a), x)`

Maxima [F]

$$\int \sqrt{a - a \sin(c + dx)} dx = \int \sqrt{-a \sin(dx + c) + a} dx$$

input `integrate((a-a*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-a*sin(d*x + c) + a), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.33

$$\begin{aligned} \int \sqrt{a - a \sin(c + dx)} dx \\ = -\frac{2\sqrt{2}\sqrt{a} \cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) \operatorname{sgn}\left(\sin\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)\right)}{d} \end{aligned}$$

input `integrate((a-a*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `-2*sqrt(2)*sqrt(a)*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))/d`

Mupad [B] (verification not implemented)

Time = 25.98 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int \sqrt{a - a \sin(c + dx)} dx = -\frac{2 \cos(c + dx) \sqrt{-a (\sin(c + dx) - 1)}}{d (\sin(c + dx) - 1)}$$

input `int((a - a*sin(c + d*x))^(1/2),x)`

output `-(2*cos(c + d*x)*(-a*(sin(c + d*x) - 1))^(1/2))/(d*(sin(c + d*x) - 1))`

Reduce [F]

$$\int \sqrt{a - a \sin(c + dx)} dx = \sqrt{a} \left(\int \sqrt{-\sin(dx + c) + 1} dx \right)$$

input `int((a-a*sin(d*x+c))^(1/2),x)`

output `sqrt(a)*int(sqrt(-sin(c+d*x)+1),x)`

3.24 $\int \frac{1}{\sqrt{a-a \sin(c+dx)}} dx$

Optimal result	197
Mathematica [C] (verified)	197
Rubi [A] (verified)	198
Maple [A] (verified)	199
Fricas [A] (verification not implemented)	200
Sympy [F]	200
Maxima [F]	201
Giac [C] (verification not implemented)	201
Mupad [F(-1)]	201
Reduce [F]	202

Optimal result

Integrand size = 15, antiderivative size = 47

$$\int \frac{1}{\sqrt{a-a \sin(c+dx)}} dx = \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2} \sqrt{a-a \sin(c+dx)}}\right)}{\sqrt{ad}}$$

output $2^{(1/2)} * \operatorname{arctanh}(1/2 * a^{(1/2)} * \cos(dx+c) * 2^{(1/2)} / (a - a * \sin(dx+c))^{(1/2)}) / a^{(1/2)} / d$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.62

$$\int \frac{1}{\sqrt{a-a \sin(c+dx)}} dx = \frac{(2+2i)\sqrt[4]{-1} \arctan\left(\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt[4]{-1} (1 + \tan\left(\frac{1}{4}(c+dx)\right))\right) (\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right))}{d \sqrt{a-a \sin(c+dx)}}$$

input `Integrate[1/Sqrt[a - a*Sin[c + d*x]],x]`

output

```
((-2 - 2*I)*(-1)^(1/4)*ArcTan[(1/2 + I/2)*(-1)^(1/4)*(1 + Tan[(c + d*x)/4])*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])]/(d*Sqrt[a - a*Sin[c + d*x]]))
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx$$

↓ 3042

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx$$

↓ 3128

$$\frac{2 \int \frac{1}{2a - \frac{a^2 \cos^2(c+dx)}{a - a \sin(c+dx)}} d\left(-\frac{a \cos(c+dx)}{\sqrt{a - a \sin(c+dx)}}\right)}{d}$$

↓ 219

$$\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2} \sqrt{a - a \sin(c+dx)}}\right)}{\sqrt{ad}}$$

input

```
Int[1/Sqrt[a - a*Sin[c + d*x]],x]
```

output

```
(Sqrt[2]*ArcTanh[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[a - a*Sin[c + d*x]])]/(Sqrt[a]*d))
```

Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3128 Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[-2/d
Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]],
x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.57

method	result
default	$-\frac{(\sin(dx+c)-1)\sqrt{a(1+\sin(dx+c))}\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{a(1+\sin(dx+c))}\sqrt{2}}{2\sqrt{a}}\right)}{\sqrt{a}\cos(dx+c)\sqrt{a-a\sin(dx+c)}}d$
risch	$\frac{2i(e^{i(dx+c)}-i)\sqrt{2}e^{-i(dx+c)}}{d\sqrt{a(ie^{2i(dx+c)}+2e^{i(dx+c)}-i)e^{-i(dx+c)}}} + \frac{2i(-e^{i(dx+c)}+i)\left(\operatorname{arctan}\left(\frac{\sqrt{ie^{i(dx+c)}a}}{\sqrt{a}}\right)a\sqrt{ie^{i(dx+c)}a+a^{\frac{3}{2}}}\right)\sqrt{2}\sqrt{ia(e^{2i(dx+c)}-2i)}}{d\sqrt{-ia(-e^{2i(dx+c)}+2ie^{i(dx+c)}+1)e^{i(dx+c)}}a^{\frac{3}{2}}\sqrt{a(ie^{2i(dx+c)}+2e^{i(dx+c)}})}$

```
input int(1/(a-a*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -(sin(d*x+c)-1)*(a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1/2)*arctanh(1/2*(a*(1
+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1/2))/cos(d*x+c)/(a-a*sin(d*x+c))^(1/2)/d
```


Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 190, normalized size of antiderivative = 4.04

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx$$

$$= \left[\frac{\sqrt{2} \log \left(-\frac{\cos(dx+c)^2 + (\cos(dx+c)-2) \sin(dx+c) + \frac{2\sqrt{2}\sqrt{-a \sin(dx+c)+a}(\cos(dx+c)+\sin(dx+c)+1)}{\sqrt{a}} + 3 \cos(dx+c)+2}{\cos(dx+c)^2 + (\cos(dx+c)+2) \sin(dx+c) - \cos(dx+c) - 2} \right)}{2\sqrt{ad}}, \frac{\sqrt{2}\sqrt{-\frac{1}{a}} \arctan\left(\frac{\cos(dx+c)+\sin(dx+c)+1}{\cos(dx+c)-\sin(dx+c)+1}\right)}{d} \right]$$

input `integrate(1/(a-a*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output `[1/2*sqrt(2)*log(-(cos(d*x + c))^2 + (cos(d*x + c) - 2)*sin(d*x + c) + 2*sqrt(2)*sqrt(-a*sin(d*x + c) + a)*(cos(d*x + c) + sin(d*x + c) + 1)/sqrt(a) + 3*cos(d*x + c) + 2)/(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2))/(sqrt(a)*d), sqrt(2)*sqrt(-1/a)*arctan(-1/2*sqrt(2)*sqrt(-a*sin(d*x + c) + a)*sqrt(-1/a)*(cos(d*x + c) + sin(d*x + c) + 1)/(cos(d*x + c) - sin(d*x + c) + 1))/d]`

Sympy [F]

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{-a \sin(c + dx) + a}} dx$$

input `integrate(1/(a-a*sin(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(-a*sin(c + d*x) + a), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{-a \sin(dx + c) + a}} dx$$

input `integrate(1/(a-a*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(-a*sin(d*x + c) + a), x)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx = \frac{\sqrt{2} \arctan \left(i \cos \left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c \right) \right)}{\sqrt{-a} \operatorname{sgn} \left(\sin \left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c \right) \right)}$$

input `integrate(1/(a-a*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `sqrt(2)*arctan(I*cos(-1/4*pi + 1/2*d*x + 1/2*c))/(sqrt(-a)*d*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx$$

input `int(1/(a - a*sin(c + d*x))^(1/2),x)`

output `int(1/(a - a*sin(c + d*x))^(1/2), x)`

Reduce [F]

$$\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx = -\frac{\sqrt{a} \left(\int \frac{\sqrt{-\sin(dx+c)+1}}{\sin(dx+c)-1} dx \right)}{a}$$

input `int(1/(a-a*sin(d*x+c))^(1/2),x)`

output `(- sqrt(a)*int(sqrt(- sin(c + d*x) + 1)/(sin(c + d*x) - 1),x))/a`

3.25 $\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx$

Optimal result	203
Mathematica [C] (verified)	203
Rubi [A] (verified)	204
Maple [A] (verified)	205
Fricas [B] (verification not implemented)	206
Sympy [F]	206
Maxima [F]	207
Giac [B] (verification not implemented)	207
Mupad [F(-1)]	208
Reduce [F]	208

Optimal result

Integrand size = 15, antiderivative size = 79

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c + dx)}{\sqrt{2} \sqrt{a - a \sin(c + dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}}$$

```
output 1/4*arctanh(1/2*a^(1/2)*cos(d*x+c)*2^(1/2)/(a-a*sin(d*x+c))^(1/2))*2^(1/2)
/a^(3/2)/d+1/2*cos(d*x+c)/d/(a-a*sin(d*x+c))^(3/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.54

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \frac{(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)) + (1 + i)\sqrt[4]{-1} \arctan((\frac{1}{2} + \frac{i}{2})\sqrt[4]{-1}))}{2ad(-1 + \sin(c + dx))\sqrt{a - a \sin(c + dx)}}$$

```
input Integrate[(a - a*Sin[c + d*x])^(-3/2), x]
```

output

```
-1/2*((Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2] + (1 + I)*(-1)^(1/4)*ArcTan[(1/2 + I/2)*(-1)^(1/4)*(1 + Tan[(c + d*x)/4])]*(-1 + Sin[c + d*x])))/(a*d*(-1 + Sin[c + d*x])*Sqrt[a - a*Sin[c + d*x]])
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx}{4a} + \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx}{4a} + \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3128} \\
 & \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}} - \frac{\int \frac{1}{2a - \frac{a^2 \cos^2(c + dx)}{a - a \sin(c + dx)}} d\left(-\frac{a \cos(c + dx)}{\sqrt{a - a \sin(c + dx)}}\right)}{2ad} \\
 & \quad \downarrow \text{219} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \cos(c + dx)}{\sqrt{2} \sqrt{a - a \sin(c + dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}}
 \end{aligned}$$

input `Int[(a - a*Sin[c + d*x])^(-3/2),x]`

output `ArcTanh[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[a - a*Sin[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) + Cos[c + d*x]/(2*d*(a - a*Sin[c + d*x])^(3/2))`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.56

method	result
default	$-\frac{\left(\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a(1+\sin(dx+c))}\sqrt{2}}{2\sqrt{a}}\right)a^2 \sin(dx+c) - \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a(1+\sin(dx+c))}\sqrt{2}}{2\sqrt{a}}\right)a^2 - 2\sqrt{a(1+\sin(dx+c))}a^{\frac{3}{2}}\right)\sqrt{a(1+\sin(dx+c))}}{4a^{\frac{7}{2}} \cos(dx+c)\sqrt{a-a \sin(dx+c)}d}$

input `int(1/(a-a*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output

```
-1/4/a^(7/2)*(2^(1/2)*arctanh(1/2*(a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1/2)
)*a^2*sin(d*x+c)-2^(1/2)*arctanh(1/2*(a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1
/2))*a^2-2*(a*(1+sin(d*x+c)))^(1/2)*a^(3/2))*(a*(1+sin(d*x+c)))^(1/2)/cos(
d*x+c)/(a-a*sin(d*x+c))^(1/2)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. $2(64) = 128$.

Time = 0.11 (sec) , antiderivative size = 246, normalized size of antiderivative = 3.11

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{2}(\cos(dx + c)^2 + (\cos(dx + c) + 2) \sin(dx + c) - \cos(dx + c) - 2) \sqrt{a} \log\left(\frac{\cos(dx + c) + \sin(dx + c) + 1}{\cos(dx + c) - \sin(dx + c) - 1}\right) + 3a \cos(dx + c) + (a \cos(dx + c) - 2a) \sin(dx + c) + 2a}{8(a^2 d \cos(dx + c)^2 - a^2 d \cos(dx + c) - 2a^2 d + (a^2 d \cos(dx + c) + 2a^2 d) \sin(dx + c))} + C$$

input

```
integrate(1/(a-a*sin(d*x+c))^(3/2),x, algorithm="fricas")
```

output

```
1/8*(sqrt(2)*(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x +
c) - 2)*sqrt(a)*log((-a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(-a*sin(d*x + c) +
a)*sqrt(a)*(cos(d*x + c) + sin(d*x + c) + 1) + 3*a*cos(d*x + c) + (a*cos(
d*x + c) - 2*a)*sin(d*x + c) + 2*a)/(cos(d*x + c)^2 + (cos(d*x + c) + 2)*s
in(d*x + c) - cos(d*x + c) - 2)) - 4*sqrt(-a*sin(d*x + c) + a)*(cos(d*x +
c) + sin(d*x + c) + 1))/(a^2*d*cos(d*x + c)^2 - a^2*d*cos(d*x + c) - 2*a^2
*d + (a^2*d*cos(d*x + c) + 2*a^2*d)*sin(d*x + c))
```

Sympy [F]

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(-a \sin(c + dx) + a)^{3/2}} dx$$

input

```
integrate(1/(a-a*sin(d*x+c))**(3/2),x)
```

output

```
Integral((-a*sin(c + d*x) + a)**(-3/2), x)
```

Maxima [F]

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(-a \sin(dx + c) + a)^{3/2}} dx$$

input `integrate(1/(a-a*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((-a*sin(d*x + c) + a)^(-3/2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 206 vs. 2(64) = 128.

Time = 0.18 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.61

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \frac{2\sqrt{2} \log\left(-\frac{\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1}{\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1}\right)}{a^{3/2} \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} + \frac{\sqrt{2}\left(\sqrt{a} - \frac{2\sqrt{a}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1)}{\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1}\right)(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))}{a^2 (\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1) \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} \quad 16d$$

input `integrate(1/(a-a*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `1/16*(2*sqrt(2)*log(-(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)))/(a^(3/2)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) + sqrt(2)*(sqrt(a) - 2*sqrt(a)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1))*(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)/(a^2*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) - sqrt(2)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(a^(3/2)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx$$

input `int(1/(a - a*sin(c + d*x))^(3/2),x)`output `int(1/(a - a*sin(c + d*x))^(3/2), x)`**Reduce [F]**

$$\int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{-\sin(dx+c)+1}}{\sin(dx+c)^2 - 2\sin(dx+c)+1} dx \right)}{a^2}$$

input `int(1/(a-a*sin(d*x+c))^(3/2),x)`output `(sqrt(a)*int(sqrt(-sin(c + d*x) + 1)/(sin(c + d*x)**2 - 2*sin(c + d*x) + 1),x))/a**2`

3.26 $\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx$

Optimal result	209
Mathematica [C] (verified)	209
Rubi [A] (verified)	210
Maple [B] (verified)	212
Fricas [B] (verification not implemented)	212
Sympy [F]	213
Maxima [F]	213
Giac [B] (verification not implemented)	214
Mupad [F(-1)]	214
Reduce [F]	215

Optimal result

Integrand size = 15, antiderivative size = 110

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \frac{3 \arctanh\left(\frac{\sqrt{a} \cos(c + dx)}{\sqrt{2} \sqrt{a - a \sin(c + dx)}}\right)}{16 \sqrt{2} a^{5/2} d} + \frac{\cos(c + dx)}{4d(a - a \sin(c + dx))^{5/2}} + \frac{3 \cos(c + dx)}{16ad(a - a \sin(c + dx))^{3/2}}$$

output

```
3/32*arctanh(1/2*a^(1/2)*cos(d*x+c)*2^(1/2)/(a-a*sin(d*x+c))^(1/2))*2^(1/2)/a^(5/2)/d+1/4*cos(d*x+c)/d/(a-a*sin(d*x+c))^(5/2)+3/16*cos(d*x+c)/a/d/(a-a*sin(d*x+c))^(3/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.45

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \frac{(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) (11 \cos(\frac{1}{2}(c + dx)) + 3 \cos(\frac{3}{2}(c + dx)))}{(a - a \sin(c + dx))^{5/2}}$$

input

```
Integrate[(a - a*Sin[c + d*x])^(-5/2), x]
```

output

```
((Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(11*Cos[(c + d*x)/2] + 3*Cos[(3*(c + d*x))/2] + 11*Sin[(c + d*x)/2] + (3 + 3*I)*(-1)^(1/4)*ArcTan[(1/2 + I/2)*(-1)^(1/4)*(1 + Tan[(c + d*x)/4])]*(-3 + Cos[2*(c + d*x)] + 4*Sin[c + d*x]) - 3*Sin[(3*(c + d*x))/2]))/(32*a^2*d*(-1 + Sin[c + d*x])^2*sqrt[a - a*Sin[c + d*x]])
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 3129, 3042, 3129, 3042, 3128, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3129} \\
 & \frac{3 \int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx}{8a} + \frac{\cos(c + dx)}{4d(a - a \sin(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{1}{(a - a \sin(c + dx))^{3/2}} dx}{8a} + \frac{\cos(c + dx)}{4d(a - a \sin(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3129} \\
 & \frac{3 \left(\frac{\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx}{4a} + \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}} \right)}{8a} + \frac{\cos(c + dx)}{4d(a - a \sin(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \left(\frac{\int \frac{1}{\sqrt{a - a \sin(c + dx)}} dx}{4a} + \frac{\cos(c + dx)}{2d(a - a \sin(c + dx))^{3/2}} \right)}{8a} + \frac{\cos(c + dx)}{4d(a - a \sin(c + dx))^{5/2}}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3128} \\
 \frac{3 \left(\frac{\cos(c+dx)}{2d(a-a\sin(c+dx))^{3/2}} - \frac{\int \frac{1}{2a - \frac{a^2 \cos^2(c+dx)}{a-a\sin(c+dx)}} d \left(-\frac{a \cos(c+dx)}{\sqrt{a-a\sin(c+dx)}} \right)}{2ad} \right)}{8a} + \frac{\cos(c+dx)}{4d(a-a\sin(c+dx))^{5/2}} \\
 \downarrow \text{219} \\
 \frac{3 \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{a}\cos(c+dx)}{\sqrt{2}\sqrt{a-a\sin(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\cos(c+dx)}{2d(a-a\sin(c+dx))^{3/2}} \right)}{8a} + \frac{\cos(c+dx)}{4d(a-a\sin(c+dx))^{5/2}}
 \end{array}$$

input `Int[(a - a*Sin[c + d*x])^(-5/2), x]`

output `Cos[c + d*x]/(4*d*(a - a*Sin[c + d*x])^(5/2)) + (3*(ArcTanh[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[a - a*Sin[c + d*x]])]/(2*Sqrt[2]*a^(3/2)*d) + Cos[c + d*x]/(2*d*(a - a*Sin[c + d*x])^(3/2)))/(8*a)`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c
+ d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n
+ 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x]
&& EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 193 vs. $2(91) = 182$.

Time = 0.10 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.76

method	result
default	$-\frac{\left(3\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a(1+\sin(dx+c))\sqrt{2}}}{2\sqrt{a}}\right) \sin(dx+c)^2 a^2 - 6\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a(1+\sin(dx+c))\sqrt{2}}}{2\sqrt{a}}\right) a^2 \sin(dx+c) - 6\sqrt{a(1+\sin(dx+c))} a^{\frac{3}{2}}\right)}{32a^{\frac{9}{2}}(\sin(dx+c)-1) \cos(dx+c) \sqrt{a-a \sin(dx+c)}}$

input

```
int(1/(a-a*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
-1/32/a^(9/2)*(3*2^(1/2)*arctanh(1/2*(a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1
/2))*sin(d*x+c)^2*a^2-6*2^(1/2)*arctanh(1/2*(a*(1+sin(d*x+c)))^(1/2)*2^(1
/2)/a^(1/2))*a^2*sin(d*x+c)-6*(a*(1+sin(d*x+c)))^(1/2)*a^(3/2)*sin(d*x+c)+3
*2^(1/2)*arctanh(1/2*(a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1/2))*a^2+14*(a*(
1+sin(d*x+c)))^(1/2)*a^(3/2)*(a*(1+sin(d*x+c)))^(1/2)/(sin(d*x+c)-1)/cos(
d*x+c)/(a-a*sin(d*x+c))^(1/2)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 321 vs. $2(91) = 182$.

Time = 0.10 (sec) , antiderivative size = 321, normalized size of antiderivative = 2.92

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \frac{3\sqrt{2}(\cos(dx+c)^3 + 3\cos(dx+c)^2 - (\cos(dx+c)^2 - 2\cos(dx+c) - 4))}{32a^{9/2}(\sin(dx+c)-1)\cos(dx+c)\sqrt{a-a\sin(dx+c)}}$$

input

```
integrate(1/(a-a*sin(d*x+c))^(5/2),x, algorithm="fricas")
```

output

```
1/64*(3*sqrt(2)*(cos(d*x + c)^3 + 3*cos(d*x + c)^2 - (cos(d*x + c)^2 - 2*cos(d*x + c) - 4)*sin(d*x + c) - 2*cos(d*x + c) - 4)*sqrt(a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(-a*sin(d*x + c) + a)*sqrt(a)*(cos(d*x + c) + sin(d*x + c) + 1) + 3*a*cos(d*x + c) + (a*cos(d*x + c) - 2*a)*sin(d*x + c) + 2*a)/(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2)) - 4*(3*cos(d*x + c)^2 - (3*cos(d*x + c) - 4)*sin(d*x + c) + 7*cos(d*x + c) + 4)*sqrt(-a*sin(d*x + c) + a)/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d - (a^3*d*cos(d*x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d)*sin(d*x + c))
```

Sympy [F]

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(-a \sin(c + dx) + a)^{5/2}} dx$$

input

```
integrate(1/(a-a*sin(d*x+c))**(5/2),x)
```

output

```
Integral((-a*sin(c + d*x) + a)**(-5/2), x)
```

Maxima [F]

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(-a \sin(dx + c) + a)^{5/2}} dx$$

input

```
integrate(1/(a-a*sin(d*x+c))^(5/2),x, algorithm="maxima")
```

output

```
integrate((-a*sin(d*x + c) + a)^(-5/2), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 303 vs. $2(91) = 182$.

Time = 0.20 (sec) , antiderivative size = 303, normalized size of antiderivative = 2.75

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \frac{12\sqrt{2} \log\left(\frac{\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1}{\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1}\right)}{a^{\frac{5}{2}} \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} - \frac{\sqrt{2} \left(\sqrt{a} - \frac{8\sqrt{a}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1)}{\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) + 1} + \frac{18\sqrt{a}(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1)^2}{(\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1)^2 \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))} \right)}{a^3 (\cos(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c) - 1)^2 \operatorname{sgn}(\sin(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c))}$$

input `integrate(1/(a-a*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `1/256*(12*sqrt(2)*log(-(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1))/(a^(5/2)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) - sqrt(2)*(sqrt(a) - 8*sqrt(a)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1) + 18*sqrt(a)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)^2/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)^2*(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)^2/(a^3*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)^2*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))) - (8*sqrt(2)*a^(7/2)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1) - sqrt(2)*a^(7/2)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)^2*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)^2)/a^6)/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx$$

input `int(1/(a - a*sin(c + d*x))^(5/2),x)`

output `int(1/(a - a*sin(c + d*x))^(5/2), x)`

Reduce [F]

$$\int \frac{1}{(a - a \sin(c + dx))^{5/2}} dx = -\frac{\sqrt{a} \left(\int \frac{\sqrt{-\sin(dx+c)+1}}{\sin(dx+c)^3 - 3\sin(dx+c)^2 + 3\sin(dx+c) - 1} dx \right)}{a^3}$$

input `int(1/(a-a*sin(d*x+c))^(5/2),x)`

output `(- sqrt(a)*int(sqrt(- sin(c + d*x) + 1)/(sin(c + d*x)**3 - 3*sin(c + d*x)**2 + 3*sin(c + d*x) - 1),x))/a**3`

3.27 $\int (-a + a \sin(c + dx))^{7/2} dx$

Optimal result	216
Mathematica [A] (verified)	216
Rubi [A] (verified)	217
Maple [A] (verified)	219
Fricas [A] (verification not implemented)	220
Sympy [F(-1)]	220
Maxima [F]	220
Giac [A] (verification not implemented)	221
Mupad [F(-1)]	221
Reduce [F]	222

Optimal result

Integrand size = 16, antiderivative size = 127

$$\int (-a + a \sin(c + dx))^{7/2} dx = \frac{256a^4 \cos(c + dx)}{35d\sqrt{-a + a \sin(c + dx)}} - \frac{64a^3 \cos(c + dx)\sqrt{-a + a \sin(c + dx)}}{35d} + \frac{24a^2 \cos(c + dx)(-a + a \sin(c + dx))^{3/2}}{35d} - \frac{2a \cos(c + dx)(-a + a \sin(c + dx))^{5/2}}{7d}$$

output

```
256/35*a^4*cos(d*x+c)/d/(-a+a*sin(d*x+c))^(1/2)-64/35*a^3*cos(d*x+c)*(-a+a*
*sin(d*x+c))^(1/2)/d+24/35*a^2*cos(d*x+c)*(-a+a*sin(d*x+c))^(3/2)/d-2/7*a*
cos(d*x+c)*(-a+a*sin(d*x+c))^(5/2)/d
```

Mathematica [A] (verified)

Time = 2.86 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.23

$$\int (-a + a \sin(c + dx))^{7/2} dx = \frac{a^3(-1 + \sin(c + dx))^3 \sqrt{a(-1 + \sin(c + dx))} (1225 \cos(\frac{1}{2}(c + dx)) + 245 \cos(\frac{3}{2}(c + dx)) - 140 \cos(\frac{5}{2}(c + dx)))}{140 \sqrt{a(-1 + \sin(c + dx))}}$$

input `Integrate[(-a + a*Sin[c + d*x])^(7/2),x]`

output $(a^3(-1 + \sin[c + dx])^3 \sqrt{a(-1 + \sin[c + dx])} (1225 \cos[(c + dx)/2] + 245 \cos[(3(c + dx))/2] - 49 \cos[(5(c + dx))/2] - 5 \cos[(7(c + dx))/2] + 1225 \sin[(c + dx)/2] - 245 \sin[(3(c + dx))/2] - 49 \sin[(5(c + dx))/2] + 5 \sin[(7(c + dx))/2])) / (140 d (\cos[(c + dx)/2] - \sin[(c + dx)/2]))^7$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3126, 3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sin(c + dx) - a)^{7/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sin(c + dx) - a)^{7/2} dx \\
 & \quad \downarrow \text{3126} \\
 & -\frac{12}{7} a \int (a \sin(c + dx) - a)^{5/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{12}{7} a \int (a \sin(c + dx) - a)^{5/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3126} \\
 & -\frac{12}{7} a \left(-\frac{8}{5} a \int (a \sin(c + dx) - a)^{3/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{3/2}}{5d} \right) - \\
 & \quad \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{5/2}}{7d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{12}{7}a \left(-\frac{8}{5}a \int (a \sin(c+dx) - a)^{3/2} dx - \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{3/2}}{5d} \right) - \\
& \quad \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{7d} \\
& \quad \downarrow \text{3126} \\
& -\frac{12}{7}a \left(-\frac{8}{5}a \left(-\frac{4}{3}a \int \sqrt{a \sin(c+dx) - a} dx - \frac{2a \cos(c+dx)\sqrt{a \sin(c+dx) - a}}{3d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{5d} \right) \\
& \quad \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{7d} \\
& \quad \downarrow \text{3042} \\
& -\frac{12}{7}a \left(-\frac{8}{5}a \left(-\frac{4}{3}a \int \sqrt{a \sin(c+dx) - a} dx - \frac{2a \cos(c+dx)\sqrt{a \sin(c+dx) - a}}{3d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{5d} \right) \\
& \quad \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{7d} \\
& \quad \downarrow \text{3125} \\
& -\frac{12}{7}a \left(-\frac{8}{5}a \left(\frac{8a^2 \cos(c+dx)}{3d\sqrt{a \sin(c+dx) - a}} - \frac{2a \cos(c+dx)\sqrt{a \sin(c+dx) - a}}{3d} \right) - \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{5d} \right) \\
& \quad \frac{2a \cos(c+dx)(a \sin(c+dx) - a)^{5/2}}{7d}
\end{aligned}$$

input `Int[(-a + a*Sin[c + d*x])^(7/2),x]`

output `(-2*a*Cos[c + d*x]*(-a + a*Sin[c + d*x])^(5/2))/(7*d) - (12*a*((-2*a*Cos[c + d*x]*(-a + a*Sin[c + d*x])^(3/2))/(5*d) - (8*a*((8*a^2*Cos[c + d*x])/(3*d*Sqrt[-a + a*Sin[c + d*x]]) - (2*a*Cos[c + d*x]*Sqrt[-a + a*Sin[c + d*x])]/(3*d))))/5)/7`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :=> Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.61

method	result	size
default	$\frac{2(\sin(dx+c)-1)a^4(1+\sin(dx+c))(5\sin(dx+c)^3-27\sin(dx+c)^2+71\sin(dx+c)-177)}{35\cos(dx+c)\sqrt{-a+a\sin(dx+c)}d}$	77

input `int((-a+a*sin(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output `2/35*(sin(d*x+c)-1)*a^4*(1+sin(d*x+c))*(5*sin(d*x+c)^3-27*sin(d*x+c)^2+71*sin(d*x+c)-177)/cos(d*x+c)/(-a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.13

$$\int (-a + a \sin(c + dx))^{7/2} dx = \frac{2(5a^3 \cos(dx + c)^4 + 27a^3 \cos(dx + c)^3 - 54a^3 \cos(dx + c)^2 - 204a^3 \cos(dx + c) - 128a^3 + dx)}{35(d \cos(dx + c) - d \sin(dx + c) + d)}$$

input `integrate((-a+a*sin(d*x+c))^(7/2),x, algorithm="fricas")`output `2/35*(5*a^3*cos(d*x + c)^4 + 27*a^3*cos(d*x + c)^3 - 54*a^3*cos(d*x + c)^2 - 204*a^3*cos(d*x + c) - 128*a^3 - (5*a^3*cos(d*x + c)^3 - 22*a^3*cos(d*x + c)^2 - 76*a^3*cos(d*x + c) + 128*a^3)*sin(d*x + c))*sqrt(a*sin(d*x + c) - a)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`**Sympy [F(-1)]**

Timed out.

$$\int (-a + a \sin(c + dx))^{7/2} dx = \text{Timed out}$$

input `integrate((-a+a*sin(d*x+c))**(7/2),x)`output `Timed out`**Maxima [F]**

$$\int (-a + a \sin(c + dx))^{7/2} dx = \int (a \sin(dx + c) - a)^{7/2} dx$$

input `integrate((-a+a*sin(d*x+c))^(7/2),x, algorithm="maxima")`output `integrate((a*sin(d*x + c) - a)^(7/2), x)`

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.06

$$\int (-a + a \sin(c + dx))^{7/2} dx = \frac{\sqrt{2}(1225 a^3 \cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) \operatorname{sgn}(\sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) - 245 a^3 \cos(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c) \operatorname{sgn}(\sin(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c)) + 49 a^3 \cos(-\frac{5}{4} \pi + \frac{5}{2} dx + \frac{5}{2} c) \operatorname{sgn}(\sin(-\frac{5}{4} \pi + \frac{5}{2} dx + \frac{5}{2} c)) - 5 a^3 \cos(-\frac{7}{4} \pi + \frac{7}{2} dx + \frac{7}{2} c) \operatorname{sgn}(\sin(-\frac{7}{4} \pi + \frac{7}{2} dx + \frac{7}{2} c))) \sqrt{-a}}{d}$$

input `integrate((-a+a*sin(d*x+c))^(7/2),x, algorithm="giac")`

output `1/140*sqrt(2)*(1225*a^3*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - 245*a^3*cos(-3/4*pi + 3/2*d*x + 3/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) + 49*a^3*cos(-5/4*pi + 5/2*d*x + 5/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - 5*a^3*cos(-7/4*pi + 7/2*d*x + 7/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))*sqrt(-a)/d`

Mupad [F(-1)]

Timed out.

$$\int (-a + a \sin(c + dx))^{7/2} dx = \int (a \sin(c + dx) - a)^{7/2} dx$$

input `int((a*sin(c + d*x) - a)^(7/2),x)`

output `int((a*sin(c + d*x) - a)^(7/2), x)`

Reduce [F]

$$\int (-a + a \sin(c + dx))^{7/2} dx = \sqrt{a} a^3 \left(- \left(\int \sqrt{\sin(dx + c) - 1} dx \right) \right. \\ \left. + \int \sqrt{\sin(dx + c) - 1} \sin(dx + c)^3 dx - 3 \left(\int \sqrt{\sin(dx + c) - 1} \sin(dx + c)^2 dx \right) \right. \\ \left. + 3 \left(\int \sqrt{\sin(dx + c) - 1} \sin(dx + c) dx \right) \right)$$

input `int((-a+a*sin(d*x+c))^(7/2),x)`

output `sqrt(a)*a**3*(- int(sqrt(sin(c + d*x) - 1),x) + int(sqrt(sin(c + d*x) - 1)*sin(c + d*x)**3,x) - 3*int(sqrt(sin(c + d*x) - 1)*sin(c + d*x)**2,x) + 3*int(sqrt(sin(c + d*x) - 1)*sin(c + d*x),x))`

3.28 $\int (-a + a \sin(c + dx))^{5/2} dx$

Optimal result	223
Mathematica [A] (verified)	223
Rubi [A] (verified)	224
Maple [A] (verified)	226
Fricas [A] (verification not implemented)	226
Sympy [F]	227
Maxima [F]	227
Giac [A] (verification not implemented)	227
Mupad [F(-1)]	228
Reduce [F]	228

Optimal result

Integrand size = 16, antiderivative size = 95

$$\int (-a + a \sin(c + dx))^{5/2} dx = -\frac{64a^3 \cos(c + dx)}{15d \sqrt{-a + a \sin(c + dx)}} + \frac{16a^2 \cos(c + dx) \sqrt{-a + a \sin(c + dx)}}{15d} - \frac{2a \cos(c + dx) (-a + a \sin(c + dx))^{3/2}}{5d}$$

output

```
-64/15*a^3*cos(d*x+c)/d/(-a+a*sin(d*x+c))^(1/2)+16/15*a^2*cos(d*x+c)*(-a+a*
*sin(d*x+c))^(1/2)/d-2/5*a*cos(d*x+c)*(-a+a*sin(d*x+c))^(3/2)/d
```

Mathematica [A] (verified)

Time = 1.83 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.25

$$\int (-a + a \sin(c + dx))^{5/2} dx = \frac{(a(-1 + \sin(c + dx)))^{5/2} (150 \cos(\frac{1}{2}(c + dx)) + 25 \cos(\frac{3}{2}(c + dx)) - 3 \cos(\frac{5}{2}(c + dx)) + 15 \cos(\frac{7}{2}(c + dx)))}{30d (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}$$

input

```
Integrate[(-a + a*Sin[c + d*x])^(5/2), x]
```


output

```
((a*(-1 + Sin[c + d*x]))^(5/2)*(150*Cos[(c + d*x)/2] + 25*Cos[(3*(c + d*x)
)/2] - 3*Cos[(5*(c + d*x))/2] + 150*Sin[(c + d*x)/2] - 25*Sin[(3*(c + d*x)
)/2] - 3*Sin[(5*(c + d*x))/2]))/(30*d*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]
)^5)
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 3126, 3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sin(c + dx) - a)^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sin(c + dx) - a)^{5/2} dx \\
 & \quad \downarrow \text{3126} \\
 & -\frac{8}{5} a \int (a \sin(c + dx) - a)^{3/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{8}{5} a \int (a \sin(c + dx) - a)^{3/2} dx - \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3126} \\
 & -\frac{8}{5} a \left(-\frac{4}{3} a \int \sqrt{a \sin(c + dx) - a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) - a}}{3d} \right) - \\
 & \quad \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{3/2}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{8}{5} a \left(-\frac{4}{3} a \int \sqrt{a \sin(c + dx) - a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) - a}}{3d} \right) - \\
 & \quad \frac{2a \cos(c + dx)(a \sin(c + dx) - a)^{3/2}}{5d}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3125} \\
 -\frac{8}{5}a \left(\frac{8a^2 \cos(c+dx)}{3d\sqrt{a\sin(c+dx)-a}} - \frac{2a \cos(c+dx)\sqrt{a\sin(c+dx)-a}}{3d} \right) - \\
 \frac{2a \cos(c+dx)(a\sin(c+dx)-a)^{3/2}}{5d}
 \end{array}$$

input `Int[(-a + a*Sin[c + d*x])^(5/2),x]`

output `(-2*a*Cos[c + d*x]*(-a + a*Sin[c + d*x])^(3/2))/(5*d) - (8*a*((8*a^2*Cos[c + d*x])/(3*d*Sqrt[-a + a*Sin[c + d*x]]) - (2*a*Cos[c + d*x]*Sqrt[-a + a*Sin[c + d*x]])/(3*d)))/5`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.71

method	result	size
default	$\frac{2(\sin(dx+c)-1)a^3(1+\sin(dx+c))(3\sin(dx+c)^2-14\sin(dx+c)+43)}{15\cos(dx+c)\sqrt{-a+a\sin(dx+c)}d}$	67

input `int((-a+a*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2/15*(sin(d*x+c)-1)*a^3*(1+sin(d*x+c))*(3*sin(d*x+c)^2-14*sin(d*x+c)+43)/cos(d*x+c)/(-a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.23

$$\int (-a + a \sin(c + dx))^{5/2} dx = \frac{2(3a^2 \cos(dx+c)^3 - 11a^2 \cos(dx+c)^2 - 46a^2 \cos(dx+c) - 32a^2 + (3a^2 \cos(dx+c)^2 + 14a^2 \cos(dx+c) - 32a^2) \sin(dx+c)) \sqrt{a \sin(dx+c) - a}}{15(d \cos(dx+c) - d \sin(dx+c) + d)}$$

input `integrate((-a+a*sin(d*x+c))^(5/2),x, algorithm="fricas")`

output `-2/15*(3*a^2*cos(d*x + c)^3 - 11*a^2*cos(d*x + c)^2 - 46*a^2*cos(d*x + c) - 32*a^2 + (3*a^2*cos(d*x + c)^2 + 14*a^2*cos(d*x + c) - 32*a^2)*sin(d*x + c))*sqrt(a*sin(d*x + c) - a)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`

Sympy [F]

$$\int (-a + a \sin(c + dx))^{5/2} dx = \int (a \sin(c + dx) - a)^{5/2} dx$$

input `integrate((-a+a*sin(d*x+c))**(5/2),x)`

output `Integral((a*sin(c + d*x) - a)**(5/2), x)`

Maxima [F]

$$\int (-a + a \sin(c + dx))^{5/2} dx = \int (a \sin(dx + c) - a)^{5/2} dx$$

input `integrate((-a+a*sin(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) - a)^(5/2), x)`

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.09

$$\int (-a + a \sin(c + dx))^{5/2} dx = \frac{\sqrt{2}(150 a^2 \cos(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c) \operatorname{sgn}(\sin(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c)) - 25 a^2 \cos(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c) \operatorname{sgn}(\sin(-\frac{3}{4} \pi + \frac{3}{2} dx + \frac{3}{2} c)))}{30 d}$$

input `integrate((-a+a*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `-1/30*sqrt(2)*(150*a^2*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - 25*a^2*cos(-3/4*pi + 3/2*d*x + 3/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) + 3*a^2*cos(-5/4*pi + 5/2*d*x + 5/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))*sqrt(-a)/d`

Mupad [F(-1)]

Timed out.

$$\int (-a + a \sin(c + dx))^{5/2} dx = \int (a \sin(c + dx) - a)^{5/2} dx$$

input `int((a*sin(c + d*x) - a)^(5/2),x)`output `int((a*sin(c + d*x) - a)^(5/2), x)`**Reduce [F]**

$$\int (-a + a \sin(c + dx))^{5/2} dx = \sqrt{a} a^2 \left(\int \sqrt{\sin(dx + c) - 1} dx \right. \\ \left. + \int \sqrt{\sin(dx + c) - 1} \sin(dx + c)^2 dx - 2 \left(\int \sqrt{\sin(dx + c) - 1} \sin(dx + c) dx \right) \right)$$

input `int((-a+a*sin(d*x+c))^(5/2),x)`output `sqrt(a)*a**2*(int(sqrt(sin(c + d*x) - 1),x) + int(sqrt(sin(c + d*x) - 1)*sin(c + d*x)**2,x) - 2*int(sqrt(sin(c + d*x) - 1)*sin(c + d*x),x))`

3.29 $\int (-a + a \sin(c + dx))^{3/2} dx$

Optimal result	229
Mathematica [A] (verified)	229
Rubi [A] (verified)	230
Maple [A] (verified)	231
Fricas [A] (verification not implemented)	232
Sympy [F]	232
Maxima [F]	232
Giac [A] (verification not implemented)	233
Mupad [F(-1)]	233
Reduce [F]	233

Optimal result

Integrand size = 16, antiderivative size = 63

$$\int (-a + a \sin(c + dx))^{3/2} dx = \frac{8a^2 \cos(c + dx)}{3d\sqrt{-a + a \sin(c + dx)}} - \frac{2a \cos(c + dx)\sqrt{-a + a \sin(c + dx)}}{3d}$$

output

$8/3*a^2*\cos(d*x+c)/d/(-a+a*\sin(d*x+c))^(1/2)-2/3*a*\cos(d*x+c)*(-a+a*\sin(d*x+c))^(1/2)/d$

Mathematica [A] (verified)

Time = 0.69 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.48

$$\int (-a + a \sin(c + dx))^{3/2} dx = \frac{(a(-1 + \sin(c + dx)))^{3/2} (9 \cos(\frac{1}{2}(c + dx)) + \cos(\frac{3}{2}(c + dx)) + 9 \sin(\frac{1}{2}(c + dx)) - \sin(\frac{3}{2}(c + dx)))}{3d (\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))^3}$$

input

`Integrate[(-a + a*Sin[c + d*x])^(3/2), x]`

output

$$\left((a(-1 + \sin(c + dx)))^{3/2} (9 \cos((c + dx)/2) + \cos(3(c + dx)/2) + 9 \sin((c + dx)/2) - \sin(3(c + dx)/2)) / (3d(\cos((c + dx)/2) - \sin((c + dx)/2)) \right)^3$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3126, 3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sin(c + dx) - a)^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (a \sin(c + dx) - a)^{3/2} dx \\ & \quad \downarrow \text{3126} \\ & -\frac{4}{3}a \int \sqrt{a \sin(c + dx) - a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) - a}}{3d} \\ & \quad \downarrow \text{3042} \\ & -\frac{4}{3}a \int \sqrt{a \sin(c + dx) - a} dx - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) - a}}{3d} \\ & \quad \downarrow \text{3125} \\ & \frac{8a^2 \cos(c + dx)}{3d \sqrt{a \sin(c + dx) - a}} - \frac{2a \cos(c + dx) \sqrt{a \sin(c + dx) - a}}{3d} \end{aligned}$$

input

$$\text{Int}[(-a + a \sin[c + d*x])^{3/2}, x]$$

output

$$\frac{(8a^2 \cos[c + d*x]) / (3d \sqrt{-a + a \sin[c + d*x]}) - (2a \cos[c + d*x] * \sqrt{-a + a \sin[c + d*x]}) / (3d)}$$

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3125 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3126 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[a*((2*n - 1)/n) Int[(a + b*Sin[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && IGtQ[n - 1/2, 0]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{2(\sin(dx+c)-1)a^2(1+\sin(dx+c))(\sin(dx+c)-5)}{3 \cos(dx+c) \sqrt{-a+a \sin(dx+c)} d}$	55

input `int((-a+a*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/3*(sin(d*x+c)-1)*a^2*(1+sin(d*x+c))*(sin(d*x+c)-5)/cos(d*x+c)/(-a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.27

$$\int (-a + a \sin(c + dx))^{3/2} dx = \frac{2(a \cos(dx + c)^2 + 5a \cos(dx + c) - (a \cos(dx + c) - 4a) \sin(dx + c) + 4a) \sqrt{a \sin(dx + c) - a}}{3(d \cos(dx + c) - d \sin(dx + c) + d)}$$

input `integrate((-a+a*sin(d*x+c))^(3/2),x, algorithm="fricas")`

output `-2/3*(a*cos(d*x + c)^2 + 5*a*cos(d*x + c) - (a*cos(d*x + c) - 4*a)*sin(d*x + c) + 4*a)*sqrt(a*sin(d*x + c) - a)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`

Sympy [F]

$$\int (-a + a \sin(c + dx))^{3/2} dx = \int (a \sin(c + dx) - a)^{\frac{3}{2}} dx$$

input `integrate((-a+a*sin(d*x+c))**(3/2),x)`

output `Integral((a*sin(c + d*x) - a)**(3/2), x)`

Maxima [F]

$$\int (-a + a \sin(c + dx))^{3/2} dx = \int (a \sin(dx + c) - a)^{\frac{3}{2}} dx$$

input `integrate((-a+a*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) - a)^(3/2), x)`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11

$$\int (-a + a \sin(c + dx))^{3/2} dx = \frac{\sqrt{2} \left(9a \cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) \operatorname{sgn}\left(\sin\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)\right) - a \cos\left(-\frac{3}{4}\pi + \frac{3}{2}dx + \frac{3}{2}c\right) \operatorname{sgn}\left(\sin\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)\right) \right)}{3d}$$

input `integrate((-a+a*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `1/3*sqrt(2)*(9*a*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)) - a*cos(-3/4*pi + 3/2*d*x + 3/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))*sqrt(-a)/d`

Mupad [F(-1)]

Timed out.

$$\int (-a + a \sin(c + dx))^{3/2} dx = \int (a \sin(c + dx) - a)^{3/2} dx$$

input `int((a*sin(c + d*x) - a)^(3/2),x)`

output `int((a*sin(c + d*x) - a)^(3/2), x)`

Reduce [F]

$$\int (-a + a \sin(c + dx))^{3/2} dx = \sqrt{a} a \left(- \left(\int \sqrt{\sin(dx + c) - 1} dx \right) + \int \sqrt{\sin(dx + c) - 1} \sin(dx + c) dx \right)$$

input `int((-a+a*sin(d*x+c))^(3/2),x)`

output

```
sqrt(a)*a*( - int(sqrt(sin(c + d*x) - 1),x) + int(sqrt(sin(c + d*x) - 1)*s  
in(c + d*x),x))
```

3.30 $\int \sqrt{-a + a \sin(c + dx)} dx$

Optimal result	235
Mathematica [B] (verified)	235
Rubi [A] (verified)	236
Maple [A] (verified)	237
Fricas [A] (verification not implemented)	237
Sympy [F]	237
Maxima [F]	238
Giac [A] (verification not implemented)	238
Mupad [B] (verification not implemented)	238
Reduce [F]	239

Optimal result

Integrand size = 16, antiderivative size = 28

$$\int \sqrt{-a + a \sin(c + dx)} dx = -\frac{2a \cos(c + dx)}{d\sqrt{-a + a \sin(c + dx)}}$$

output `-2*a*cos(d*x+c)/d/(-a+a*sin(d*x+c))^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 65 vs. 2(28) = 56.

Time = 0.06 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.32

$$\begin{aligned} & \int \sqrt{-a + a \sin(c + dx)} dx \\ &= \frac{2(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) \sqrt{a(-1 + \sin(c + dx))}}{d(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))} \end{aligned}$$

input `Integrate[Sqrt[-a + a*Sin[c + d*x]],x]`

output

```
(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*Sqrt[a*(-1 + Sin[c + d*x]))/(d*(
Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 3125}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a \sin(c + dx) - a} dx$$

↓ 3042

$$\int \sqrt{a \sin(c + dx) - a} dx$$

↓ 3125

$$-\frac{2a \cos(c + dx)}{d \sqrt{a \sin(c + dx) - a}}$$

input

```
Int[Sqrt[-a + a*Sin[c + d*x]],x]
```

output

```
(-2*a*Cos[c + d*x])/(d*Sqrt[-a + a*Sin[c + d*x]])
```

Defintions of rubi rules used

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3125

```
Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*b*(Cos
[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq
Q[a^2 - b^2, 0]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.61

method	result	size
default	$\frac{2(\sin(dx+c)-1)(1+\sin(dx+c))a}{\cos(dx+c)\sqrt{-a+a\sin(dx+c)}}d$	45
risch	$-\frac{i\sqrt{2}\sqrt{-a(2-2\sin(dx+c))}(e^{i(dx+c)}+i)(e^{i(dx+c)}-i)}{(e^{2i(dx+c)}-2ie^{i(dx+c)}-1)d}$	74

input `int((-a+a*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(sin(d*x+c)-1)*(1+sin(d*x+c))*a/cos(d*x+c)/(-a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.82

$$\int \sqrt{-a + a \sin(c + dx)} dx = \frac{2 \sqrt{a \sin(dx + c) - a} (\cos(dx + c) + \sin(dx + c) + 1)}{d \cos(dx + c) - d \sin(dx + c) + d}$$

input `integrate((-a+a*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output `2*sqrt(a*sin(d*x + c) - a)*(cos(d*x + c) + sin(d*x + c) + 1)/(d*cos(d*x + c) - d*sin(d*x + c) + d)`

Sympy [F]

$$\int \sqrt{-a + a \sin(c + dx)} dx = \int \sqrt{a \sin(c + dx) - a} dx$$

input `integrate((-a+a*sin(d*x+c))**(1/2),x)`

output `Integral(sqrt(a*sin(c + d*x) - a), x)`

Maxima [F]

$$\int \sqrt{-a + a \sin(c + dx)} dx = \int \sqrt{a \sin(dx + c) - a} dx$$

input `integrate((-a+a*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(a*sin(d*x + c) - a), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.36

$$\begin{aligned} \int \sqrt{-a + a \sin(c + dx)} dx \\ = -\frac{2\sqrt{2}\sqrt{-a} \cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) \operatorname{sgn}\left(\sin\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)\right)}{d} \end{aligned}$$

input `integrate((-a+a*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `-2*sqrt(2)*sqrt(-a)*cos(-1/4*pi + 1/2*d*x + 1/2*c)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))/d`

Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.18

$$\int \sqrt{-a + a \sin(c + dx)} dx = -\frac{2 \cos(c + dx) \sqrt{a (\sin(c + dx) - 1)}}{d (\sin(c + dx) - 1)}$$

input `int((a*sin(c + d*x) - a)^(1/2),x)`

output `-(2*cos(c + d*x)*(a*(sin(c + d*x) - 1))^(1/2))/(d*(sin(c + d*x) - 1))`

Reduce [F]

$$\int \sqrt{-a + a \sin(c + dx)} dx = \sqrt{a} \left(\int \sqrt{\sin(dx + c) - 1} dx \right)$$

input `int((-a+a*sin(d*x+c))^(1/2),x)`

output `sqrt(a)*int(sqrt(sin(c + d*x) - 1),x)`

3.31 $\int \frac{1}{\sqrt{-a+a \sin(c+dx)}} dx$

Optimal result	240
Mathematica [C] (verified)	240
Rubi [A] (verified)	241
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	243
Sympy [F]	243
Maxima [F]	244
Giac [C] (verification not implemented)	244
Mupad [F(-1)]	244
Reduce [F]	245

Optimal result

Integrand size = 16, antiderivative size = 48

$$\int \frac{1}{\sqrt{-a+a \sin(c+dx)}} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2}\sqrt{-a+a \sin(c+dx)}}\right)}{\sqrt{ad}}$$

output

$2^{(1/2)}*\arctan(1/2*a^{(1/2)}*\cos(d*x+c)*2^{(1/2)/(-a+a*\sin(d*x+c))^{(1/2)})/a^{(1/2)}/d$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.56

$$\int \frac{1}{\sqrt{-a+a \sin(c+dx)}} dx = \frac{(2+2i)\sqrt[4]{-1} \arctan\left(\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt[4]{-1}(1 + \tan\left(\frac{1}{4}(c+dx)\right))\right) (\cos\left(\frac{1}{2}(c+dx)\right) - \sin\left(\frac{1}{2}(c+dx)\right))}{d\sqrt{a(-1+\sin(c+dx))}}$$

input

`Integrate[1/Sqrt[-a + a*Sin[c + d*x]],x]`

output

$$\left((-2 - 2i)^{-1/4} \operatorname{ArcTan}\left[\frac{1}{2} + \frac{i}{2}\right] (-1)^{-1/4} (1 + \operatorname{Tan}[(c + dx)/4]) \right) \cdot (\operatorname{Cos}[(c + dx)/2] - \operatorname{Sin}[(c + dx)/2]) / (d \sqrt{a(-1 + \operatorname{Sin}[c + dx])})$$
Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 3128, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx \\ & \quad \downarrow \text{3128} \\ & \frac{2 \int \frac{1}{\frac{-a^2 \cos^2(c+dx)}{a \sin(c+dx)-a} - 2a} d \frac{a \cos(c+dx)}{\sqrt{a \sin(c+dx)-a}}}{d} \\ & \quad \downarrow \text{217} \\ & \frac{\sqrt{2} \arctan\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2} \sqrt{a \sin(c+dx)-a}}\right)}{\sqrt{ad}} \end{aligned}$$

input

$$\text{Int}[1/\text{Sqrt}[-a + a*\text{Sin}[c + d*x]],x]$$

output

$$\left(\sqrt{2} \operatorname{ArcTan}\left[\frac{\sqrt{a} \operatorname{Cos}[c + dx]}{\sqrt{2} \sqrt{-a + a \operatorname{Sin}[c + dx]}}\right] \right) / (\sqrt{a} * d)$$

Definitions of rubi rules used

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.58

method	result
default	$\frac{(\sin(dx+c)-1)\sqrt{-a(1+\sin(dx+c))}\sqrt{2}\arctan\left(\frac{\sqrt{-a(1+\sin(dx+c))}\sqrt{2}}{2\sqrt{a}}\right)}{\sqrt{a}\cos(dx+c)\sqrt{-a+a\sin(dx+c)}d}$
risch	$\frac{2i(e^{i(dx+c)}-i)\sqrt{2}e^{-i(dx+c)}}{d\sqrt{-a(ie^{2i(dx+c)}+2e^{i(dx+c)}-i)e^{-i(dx+c)}}} + \frac{2i(-e^{i(dx+c)}+i)\left(a^{\frac{3}{2}}-\operatorname{arctanh}\left(\frac{\sqrt{-ie^{i(dx+c)}}a}{\sqrt{a}}\right)a\sqrt{-ie^{i(dx+c)}}a\right)\sqrt{2}\sqrt{-ia(e^{2i(dx+c)}+2e^{i(dx+c)}-i)e^{-i(dx+c)}}}{d\sqrt{ia(-e^{2i(dx+c)}+2ie^{i(dx+c)}+1)e^{i(dx+c)}}a^{\frac{3}{2}}\sqrt{-a(ie^{2i(dx+c)}+2e^{i(dx+c)}-i)e^{-i(dx+c)}}}$

input `int(1/(-a+a*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `(sin(d*x+c)-1)*(-a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1/2)*arctan(1/2*(-a*(1+sin(d*x+c)))^(1/2)*2^(1/2)/a^(1/2))/cos(d*x+c)/(-a+a*sin(d*x+c))^(1/2)/d`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 192, normalized size of antiderivative = 4.00

$$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx$$

$$= \left[\frac{\sqrt{2} \sqrt{-\frac{1}{a}} \log \left(-\frac{2\sqrt{2}\sqrt{a \sin(dx+c)} - a \sqrt{-\frac{1}{a}} (\cos(dx+c) + \sin(dx+c) + 1) + \cos(dx+c)^2 + (\cos(dx+c) - 2) \sin(dx+c) + 3 \cos(dx+c) + 2}{\cos(dx+c)^2 + (\cos(dx+c) + 2) \sin(dx+c) - \cos(dx+c) - 2} \right)}{2d} \right],$$

input `integrate(1/(-a+a*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output `[1/2*sqrt(2)*sqrt(-1/a)*log(-(2*sqrt(2)*sqrt(a*sin(d*x + c) - a)*sqrt(-1/a)*(cos(d*x + c) + sin(d*x + c) + 1) + cos(d*x + c)^2 + (cos(d*x + c) - 2)*sin(d*x + c) + 3*cos(d*x + c) + 2)/(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2))/d, sqrt(2)*arctan(-1/2*sqrt(2)*sqrt(a*sin(d*x + c) - a)*(cos(d*x + c) + sin(d*x + c) + 1)/(sqrt(a)*(cos(d*x + c) - sin(d*x + c) + 1)))/(sqrt(a)*d)]`

Sympy [F]

$$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx$$

input `integrate(1/(-a+a*sin(d*x+c))**(1/2),x)`

output `Integral(1/sqrt(a*sin(c + d*x) - a), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a \sin(dx + c) - a}} dx$$

input `integrate(1/(-a+a*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(a*sin(d*x + c) - a), x)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx = \frac{\sqrt{2} \arctan\left(-i \cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right)\right)}{\sqrt{a} d \operatorname{sgn}\left(\sin\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right)\right)}$$

input `integrate(1/(-a+a*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `sqrt(2)*arctan(-I*cos(-1/4*pi + 1/2*d*x + 1/2*c))/(sqrt(a)*d*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx$$

input `int(1/(a*sin(c + d*x) - a)^(1/2),x)`

output `int(1/(a*sin(c + d*x) - a)^(1/2), x)`

Reduce [F]

$$\int \frac{1}{\sqrt{-a + a \sin(c + dx)}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{\sin(dx+c)-1}}{\sin(dx+c)-1} dx \right)}{a}$$

input `int(1/(-a+a*sin(d*x+c))^(1/2),x)`

output `(sqrt(a)*int(sqrt(sin(c + d*x) - 1)/(sin(c + d*x) - 1),x))/a`

3.32 $\int \frac{1}{(-a+a \sin(c+dx))^{3/2}} dx$

Optimal result	246
Mathematica [C] (verified)	246
Rubi [A] (verified)	247
Maple [B] (verified)	248
Fricas [B] (verification not implemented)	249
Sympy [F]	250
Maxima [F]	250
Giac [B] (verification not implemented)	250
Mupad [F(-1)]	251
Reduce [F]	251

Optimal result

Integrand size = 16, antiderivative size = 81

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = -\frac{\arctan\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2}\sqrt{-a+a \sin(c+dx)}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{\cos(c + dx)}{2d(-a + a \sin(c + dx))^{3/2}}$$

```
output -1/4*arctan(1/2*a^(1/2)*cos(d*x+c)*2^(1/2)/(-a+a*sin(d*x+c))^(1/2))*2^(1/2)
)/a^(3/2)/d+1/2*cos(d*x+c)/d/(-a+a*sin(d*x+c))^(3/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.33

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \frac{(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{2d(a(-1 + \sin(c + dx)))^{3/2}}$$

```
input Integrate[(-a + a*Sin[c + d*x])^(-3/2),x]
```

output

```
((Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]
] + (1 + I)*(-1)^(1/4)*ArcTan[(1/2 + I/2)*(-1)^(1/4)*(1 + Tan[(c + d*x)/4]
)]*(-1 + Sin[c + d*x]))/(2*d*(a*(-1 + Sin[c + d*x]))^(3/2))
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 3129, 3042, 3128, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \sin(c + dx) - a)^{3/2}} dx$$

↓ 3042

$$\int \frac{1}{(a \sin(c + dx) - a)^{3/2}} dx$$

↓ 3129

$$\frac{\cos(c + dx)}{2d(a \sin(c + dx) - a)^{3/2}} - \frac{\int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx}{4a}$$

↓ 3042

$$\frac{\cos(c + dx)}{2d(a \sin(c + dx) - a)^{3/2}} - \frac{\int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx}{4a}$$

↓ 3128

$$\frac{\int \frac{1}{-\frac{a^2 \cos^2(c + dx)}{a \sin(c + dx) - a} - 2a} d \frac{a \cos(c + dx)}{\sqrt{a \sin(c + dx) - a}}}{2ad} + \frac{\cos(c + dx)}{2d(a \sin(c + dx) - a)^{3/2}}$$

↓ 217

$$\frac{\cos(c + dx)}{2d(a \sin(c + dx) - a)^{3/2}} - \frac{\arctan\left(\frac{\sqrt{a} \cos(c + dx)}{\sqrt{2} \sqrt{a \sin(c + dx) - a}}\right)}{2\sqrt{2}a^{3/2}d}$$

input `Int[(-a + a*Sin[c + d*x])^(-3/2),x]`

output `-1/2*ArcTan[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[-a + a*Sin[c + d*x]])]/(Sqrt[2]*a^(3/2)*d) + Cos[c + d*x]/(2*d*(-a + a*Sin[c + d*x])^(3/2))`

Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c + d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n + 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. 2(66) = 132.

Time = 0.12 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.65

method	result
default	$\frac{\left(-\sqrt{2} \arctan\left(\frac{\sqrt{-a \sin(dx+c)-a\sqrt{2}}}{2\sqrt{a}}\right) a^2 \sin(dx+c) + 2\sqrt{-a \sin(dx+c)-a} a^{\frac{3}{2}} + \sqrt{2} \arctan\left(\frac{\sqrt{-a \sin(dx+c)-a\sqrt{2}}}{2\sqrt{a}}\right) a^2\right) \sqrt{-a(1+\sin(dx+c))}}{4a^{\frac{7}{2}} \cos(dx+c) \sqrt{-a+a \sin(dx+c)} d}$

input `int(1/(-a+a*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `1/4*(-2^(1/2)*arctan(1/2*(-a*sin(d*x+c)-a)^(1/2)*2^(1/2)/a^(1/2))*a^2*sin(d*x+c)+2*(-a*sin(d*x+c)-a)^(1/2)*a^(3/2)+2^(1/2)*arctan(1/2*(-a*sin(d*x+c)-a)^(1/2)*2^(1/2)/a^(1/2))*a^2*(-a*(1+sin(d*x+c)))^(1/2)/a^(7/2)/cos(d*x+c)/(-a+a*sin(d*x+c))^(1/2)/d`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. $2(66) = 132$.

Time = 0.11 (sec) , antiderivative size = 252, normalized size of antiderivative = 3.11

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{2}(\cos(dx + c)^2 + (\cos(dx + c) + 2)\sin(dx + c) - \cos(dx + c) - 2)\sqrt{-a} \log\left(-\frac{a \cos(dx+c)^2 + 2\sqrt{2}\sqrt{a \sin(dx+c)}}{8(a^2 d \cos(dx + c)^2 - a^2 d \cos(dx + c) - 2a^2 d + (a^2 d \cos(dx + c) + 2a^2 d)\sin(dx + c))}\right)}{8(a^2 d \cos(dx + c)^2 - a^2 d \cos(dx + c) - 2a^2 d + (a^2 d \cos(dx + c) + 2a^2 d)\sin(dx + c))}$$

input `integrate(1/(-a+a*sin(d*x+c))^(3/2),x, algorithm="fricas")`

output `-1/8*(sqrt(2)*(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2)*sqrt(-a)*log(-(a*cos(d*x + c)^2 + 2*sqrt(2)*sqrt(a*sin(d*x + c) - a)*sqrt(-a)*(cos(d*x + c) + sin(d*x + c) + 1) + 3*a*cos(d*x + c) + (a*cos(d*x + c) - 2*a)*sin(d*x + c) + 2*a)/(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) - 2)) + 4*sqrt(a*sin(d*x + c) - a)*(cos(d*x + c) + sin(d*x + c) + 1))/(a^2*d*cos(d*x + c)^2 - a^2*d*cos(d*x + c) - 2*a^2*d + (a^2*d*cos(d*x + c) + 2*a^2*d)*sin(d*x + c))`

Sympy [F]

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a \sin(c + dx) - a)^{\frac{3}{2}}} dx$$

input `integrate(1/(-a+a*sin(d*x+c))**(3/2),x)`

output `Integral((a*sin(c + d*x) - a)**(-3/2), x)`

Maxima [F]

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a \sin(dx + c) - a)^{\frac{3}{2}}} dx$$

input `integrate(1/(-a+a*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) - a)^(-3/2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 215 vs. 2(66) = 132.

Time = 0.17 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.65

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{2} \left(\frac{2 \left(\cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right) - 1\right)}{\cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right) + 1} - 1 \right) \left(\cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right)}{\sqrt{-aa} \left(\cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right) \operatorname{sgn}\left(\sin\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right)\right)} - \frac{2 \sqrt{2} \log\left(-\frac{\cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right)}{\cos\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right)}\right)}{\sqrt{-a} \operatorname{sgn}\left(\sin\left(-\frac{1}{4} \pi + \frac{1}{2} dx + \frac{1}{2} c\right)\right)} \frac{1}{16 d}$$

input `integrate(1/(-a+a*sin(d*x+c))^(3/2),x, algorithm="giac")`

output

```
1/16*(sqrt(2)*(2*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1) - 1)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)/(sqrt(-a)*a*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) - 2*sqrt(2)*log(-(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1))/(sqrt(-a)*a*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) + sqrt(2)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(sqrt(-a)*a*(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))))/d
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a \sin(c + dx) - a)^{3/2}} dx$$

input

```
int(1/(a*sin(c + d*x) - a)^(3/2), x)
```

output

```
int(1/(a*sin(c + d*x) - a)^(3/2), x)
```

Reduce [F]

$$\int \frac{1}{(-a + a \sin(c + dx))^{3/2}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{\sin(dx+c)-1}}{\sin(dx+c)^2 - 2\sin(dx+c) + 1} dx \right)}{a^2}$$

input

```
int(1/(-a+a*sin(d*x+c))^(3/2), x)
```

output

```
(sqrt(a)*int(sqrt(sin(c + d*x) - 1)/(sin(c + d*x)**2 - 2*sin(c + d*x) + 1), x))/a**2
```

3.33 $\int \frac{1}{(-a+a \sin(c+dx))^{5/2}} dx$

Optimal result	252
Mathematica [C] (verified)	252
Rubi [A] (verified)	253
Maple [B] (verified)	255
Fricas [B] (verification not implemented)	255
Sympy [F]	256
Maxima [F]	256
Giac [B] (verification not implemented)	257
Mupad [F(-1)]	257
Reduce [F]	258

Optimal result

Integrand size = 16, antiderivative size = 113

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \frac{3 \arctan\left(\frac{\sqrt{a} \cos(c+dx)}{\sqrt{2}\sqrt{-a+a \sin(c+dx)}}\right)}{16\sqrt{2}a^{5/2}d} + \frac{\cos(c + dx)}{4d(-a + a \sin(c + dx))^{5/2}} - \frac{3 \cos(c + dx)}{16ad(-a + a \sin(c + dx))^{3/2}}$$

output

```
3/32*arctan(1/2*a^(1/2)*cos(d*x+c)*2^(1/2)/(-a+a*sin(d*x+c))^(1/2))*2^(1/2)/a^(5/2)/d+1/4*cos(d*x+c)/d/(-a+a*sin(d*x+c))^(5/2)-3/16*cos(d*x+c)/a/d/(-a+a*sin(d*x+c))^(3/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.29

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \frac{(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) (11 \cos(\frac{1}{2}(c + dx)) + 3 \cos(\frac{3}{2}(c + dx)))}{(-a + a \sin(c + dx))^{5/2}}$$

input

```
Integrate[(-a + a*Sin[c + d*x])^(-5/2), x]
```

output

```
((Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(11*Cos[(c + d*x)/2] + 3*Cos[(3*(c + d*x))/2] + 11*Sin[(c + d*x)/2] + (3 + 3*I)*(-1)^(1/4)*ArcTan[(1/2 + I/2)*(-1)^(1/4)*(1 + Tan[(c + d*x)/4])]*(-3 + Cos[2*(c + d*x)] + 4*Sin[c + d*x]) - 3*Sin[(3*(c + d*x))/2]))/(32*d*(a*(-1 + Sin[c + d*x]))^(5/2))
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {3042, 3129, 3042, 3129, 3042, 3128, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a \sin(c + dx) - a)^{5/2}} dx$$

↓ 3042

$$\int \frac{1}{(a \sin(c + dx) - a)^{5/2}} dx$$

↓ 3129

$$\frac{\cos(c + dx)}{4d(a \sin(c + dx) - a)^{5/2}} - \frac{3 \int \frac{1}{(a \sin(c + dx) - a)^{3/2}} dx}{8a}$$

↓ 3042

$$\frac{\cos(c + dx)}{4d(a \sin(c + dx) - a)^{5/2}} - \frac{3 \int \frac{1}{(a \sin(c + dx) - a)^{3/2}} dx}{8a}$$

↓ 3129

$$\frac{\cos(c + dx)}{4d(a \sin(c + dx) - a)^{5/2}} - \frac{3 \left(\frac{\cos(c + dx)}{2d(a \sin(c + dx) - a)^{3/2}} - \frac{\int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx}{4a} \right)}{8a}$$

↓ 3042

$$\frac{\cos(c + dx)}{4d(a \sin(c + dx) - a)^{5/2}} - \frac{3 \left(\frac{\cos(c + dx)}{2d(a \sin(c + dx) - a)^{3/2}} - \frac{\int \frac{1}{\sqrt{a \sin(c + dx) - a}} dx}{4a} \right)}{8a}$$

$$\frac{\cos(c+dx)}{4d(a\sin(c+dx)-a)^{5/2}} - \frac{3 \left(\frac{\int \frac{1}{-\frac{a^2 \cos^2(c+dx)}{a\sin(c+dx)-a} - 2a} d - \frac{a \cos(c+dx)}{\sqrt{a\sin(c+dx)-a}}}{2ad} + \frac{\cos(c+dx)}{2d(a\sin(c+dx)-a)^{3/2}} \right)}{8a}$$

↓ 3128

$$\frac{\cos(c+dx)}{4d(a\sin(c+dx)-a)^{5/2}} - \frac{3 \left(\frac{\cos(c+dx)}{2d(a\sin(c+dx)-a)^{3/2}} - \frac{\arctan\left(\frac{\sqrt{a}\cos(c+dx)}{\sqrt{2}\sqrt{a\sin(c+dx)-a}}\right)}{2\sqrt{2}a^{3/2}d} \right)}{8a}$$

↓ 217

input `Int[(-a + a*Sin[c + d*x])^(-5/2),x]`

output `Cos[c + d*x]/(4*d*(-a + a*Sin[c + d*x])^(5/2)) - (3*(-1/2*ArcTan[(Sqrt[a]*Cos[c + d*x])/(Sqrt[2]*Sqrt[-a + a*Sin[c + d*x]])]/(Sqrt[2]*a^(3/2)*d) + Cos[c + d*x]/(2*d*(-a + a*Sin[c + d*x])^(3/2)))/(8*a)`

Defintions of rubi rules used

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3128 `Int[1/Sqrt[(a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2/d Subst[Int[1/(2*a - x^2), x], x, b*(Cos[c + d*x]/Sqrt[a + b*Sin[c + d*x])]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]`

rule 3129

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*Cos[c
+ d*x]*((a + b*Sin[c + d*x])^n/(a*d*(2*n + 1))), x] + Simp[(n + 1)/(a*(2*n
+ 1)) Int[(a + b*Sin[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x]
&& EqQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(94) = 188.

Time = 0.12 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.87

method	result
default	$-\frac{\left(3\sqrt{2} \arctan\left(\frac{\sqrt{-a \sin(dx+c)-a}\sqrt{2}}{2\sqrt{a}}\right)a^2 \cos(dx+c)^2 + 6\sqrt{2} \arctan\left(\frac{\sqrt{-a \sin(dx+c)-a}\sqrt{2}}{2\sqrt{a}}\right)a^2 \sin(dx+c) + 6\sqrt{-a \sin(dx+c)-a} a^{\frac{3}{2}}\right)}{32a^{\frac{9}{2}}(\sin(dx+c)-1)\cos(dx+c)\sqrt{-a+a \sin(dx+c)}}$

input

```
int(1/(-a+a*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
-1/32*(3*2^(1/2)*arctan(1/2*(-a*sin(d*x+c)-a)^(1/2)*2^(1/2)/a^(1/2))*a^2*cos
(d*x+c)^2+6*2^(1/2)*arctan(1/2*(-a*sin(d*x+c)-a)^(1/2)*2^(1/2)/a^(1/2))*
a^2*sin(d*x+c)+6*(-a*sin(d*x+c)-a)^(1/2)*a^(3/2)*sin(d*x+c)-6*2^(1/2)*arct
an(1/2*(-a*sin(d*x+c)-a)^(1/2)*2^(1/2)/a^(1/2))*a^2-14*(-a*sin(d*x+c)-a)^(
1/2)*a^(3/2)*(-a*(1+sin(d*x+c)))^(1/2)/a^(9/2)/(sin(d*x+c)-1)/cos(d*x+c)/
(-a+a*sin(d*x+c))^(1/2)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. 2(94) = 188.

Time = 0.11 (sec) , antiderivative size = 327, normalized size of antiderivative = 2.89

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \frac{3\sqrt{2}(\cos(dx+c)^3 + 3\cos(dx+c)^2 - (\cos(dx+c))^2 - 2\cos(dx+c) - 4)\sin(dx+c) - 2\cos(dx+c)}{64(a^3 d \cos(dx+c) + \dots)}$$

input

```
integrate(1/(-a+a*sin(d*x+c))^(5/2),x, algorithm="fricas")
```


output

```
-1/64*(3*sqrt(2)*(cos(d*x + c)^3 + 3*cos(d*x + c)^2 - (cos(d*x + c)^2 - 2*
cos(d*x + c) - 4)*sin(d*x + c) - 2*cos(d*x + c) - 4)*sqrt(-a)*log(-(a*cos(
d*x + c)^2 - 2*sqrt(2)*sqrt(a*sin(d*x + c) - a)*sqrt(-a)*(cos(d*x + c) + s
in(d*x + c) + 1) + 3*a*cos(d*x + c) + (a*cos(d*x + c) - 2*a)*sin(d*x + c)
+ 2*a)/(cos(d*x + c)^2 + (cos(d*x + c) + 2)*sin(d*x + c) - cos(d*x + c) -
2)) - 4*(3*cos(d*x + c)^2 - (3*cos(d*x + c) - 4)*sin(d*x + c) + 7*cos(d*x
+ c) + 4)*sqrt(a*sin(d*x + c) - a))/(a^3*d*cos(d*x + c)^3 + 3*a^3*d*cos(d*
x + c)^2 - 2*a^3*d*cos(d*x + c) - 4*a^3*d - (a^3*d*cos(d*x + c)^2 - 2*a^3*
d*cos(d*x + c) - 4*a^3*d)*sin(d*x + c))
```

Sympy [F]

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a \sin(c + dx) - a)^{\frac{5}{2}}} dx$$

input

```
integrate(1/(-a+a*sin(d*x+c))**(5/2),x)
```

output

```
Integral((a*sin(c + d*x) - a)**(-5/2), x)
```

Maxima [F]

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a \sin(dx + c) - a)^{\frac{5}{2}}} dx$$

input

```
integrate(1/(-a+a*sin(d*x+c))^(5/2),x, algorithm="maxima")
```

output

```
integrate((a*sin(d*x + c) - a)^(-5/2), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 313 vs. $2(94) = 188$.

Time = 0.20 (sec) , antiderivative size = 313, normalized size of antiderivative = 2.77

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \frac{\sqrt{2} \left(\frac{8 \left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) - 1\right)}{\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) + 1} - \frac{18 \left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) - 1\right)^2}{\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) + 1\right)^2} - 1 \right) \left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) + 1\right)^2}{\sqrt{-aa^2} \left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right) - 1\right)^2 \operatorname{sgn}\left(\sin\left(-\frac{1}{4}\pi + \frac{1}{2}dx + \frac{1}{2}c\right)\right)} +$$

input `integrate(1/(-a+a*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `1/256*(sqrt(2)*(8*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1) - 18*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)^2/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)^2 - 1)*(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)^2/(sqrt(-a)*a^2*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)^2*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) + 12*sqrt(2)*log(-(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1))/(sqrt(-a)*a^2*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))) + (8*sqrt(2)*sqrt(-a)*a^2*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c)))/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1) - sqrt(2)*sqrt(-a)*a^2*(cos(-1/4*pi + 1/2*d*x + 1/2*c) - 1)^2*sgn(sin(-1/4*pi + 1/2*d*x + 1/2*c))/(cos(-1/4*pi + 1/2*d*x + 1/2*c) + 1)^2)/a^5/d`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a \sin(c + dx) - a)^{5/2}} dx$$

input `int(1/(a*sin(c + d*x) - a)^(5/2),x)`

output `int(1/(a*sin(c + d*x) - a)^(5/2), x)`

Reduce [F]

$$\int \frac{1}{(-a + a \sin(c + dx))^{5/2}} dx = \frac{\sqrt{a} \left(\int \frac{\sqrt{\sin(dx+c)-1}}{\sin(dx+c)^3 - 3\sin(dx+c)^2 + 3\sin(dx+c) - 1} dx \right)}{a^3}$$

input `int(1/(-a+a*sin(d*x+c))^(5/2),x)`

output `(sqrt(a)*int(sqrt(sin(c + d*x) - 1)/(sin(c + d*x)**3 - 3*sin(c + d*x)**2 + 3*sin(c + d*x) - 1),x))/a**3`

3.34 $\int (a + a \sin(c + dx))^{4/3} dx$

Optimal result	259
Mathematica [B] (warning: unable to verify)	259
Rubi [A] (verified)	260
Maple [F]	261
Fricas [F]	262
Sympy [F]	262
Maxima [F]	262
Giac [F]	263
Mupad [F(-1)]	263
Reduce [F]	263

Optimal result

Integrand size = 14, antiderivative size = 67

$$\int (a + a \sin(c + dx))^{4/3} dx = \frac{2 \cdot 2^{5/6} a \cos(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right) \sqrt[3]{a + a \sin(c + dx)}}{d(1 + \sin(c + dx))^{5/6}}$$

output

```
-2*2^(5/6)*a*cos(d*x+c)*hypergeom([-5/6, 1/2], [3/2], 1/2-1/2*sin(d*x+c))*(a+a*sin(d*x+c))^(1/3)/d/(1+sin(d*x+c))^(5/6)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 223 vs. 2(67) = 134.

Time = 1.84 (sec) , antiderivative size = 223, normalized size of antiderivative = 3.33

$$\int (a + a \sin(c + dx))^{4/3} dx = \frac{(a(1 + \sin(c + dx)))^{4/3} \left(20\sqrt[3]{2} \cos\left(\frac{1}{4}(2c + \pi + 2dx)\right) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{6}; \frac{5}{6}; \sin^2\left(\frac{1}{4}(2c + \pi + 2dx)\right)\right) \right)}{8d\sqrt{\cos^2\left(\frac{1}{4}(2c + \pi + 2dx)\right)}}$$

input `Integrate[(a + a*Sin[c + d*x])^(4/3),x]`

output `((a*(1 + Sin[c + d*x]))^(4/3)*(20*2^(1/3)*Cos[(2*c + Pi + 2*d*x)/4]*HypergeometricPFQ[{-1/2, -1/6}, {5/6}, Sin[(2*c + Pi + 2*d*x)/4]^2] - Sqrt[2 - 2*Sin[c + d*x]]*(10*2^(1/3)*Cos[(2*c + Pi + 2*d*x)/4] + 3*Cos[c + d*x]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^(2/3)*Sin[(2*c + Pi + 2*d*x)/4]^(1/3)))/(8*d*Sqrt[Cos[(2*c + Pi + 2*d*x)/4]^2]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^(8/3)*Sin[(2*c + Pi + 2*d*x)/4]^(1/3))`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sin(c + dx) + a)^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a \sin(c + dx) + a)^{4/3} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{a \sqrt[3]{a \sin(c + dx) + a} \int (\sin(c + dx) + 1)^{4/3} dx}{\sqrt[3]{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a \sqrt[3]{a \sin(c + dx) + a} \int (\sin(c + dx) + 1)^{4/3} dx}{\sqrt[3]{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{3130} \\
 & - \frac{2 \cdot 2^{5/6} a \cos(c + dx) \sqrt[3]{a \sin(c + dx) + a} \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d(\sin(c + dx) + 1)^{5/6}}
 \end{aligned}$$

input `Int[(a + a*Sin[c + d*x])^(4/3),x]`

output `(-2*2^(5/6)*a*Cos[c + d*x]*Hypergeometric2F1[-5/6, 1/2, 3/2, (1 - Sin[c + d*x])/2]*(a + a*Sin[c + d*x])^(1/3))/(d*(1 + Sin[c + d*x])^(5/6))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

Maple [F]

$$\int (a + a \sin(dx + c))^{\frac{4}{3}} dx$$

input `int((a+a*sin(d*x+c))^(4/3),x)`

output `int((a+a*sin(d*x+c))^(4/3),x)`

Fricas [F]

$$\int (a + a \sin(c + dx))^{4/3} dx = \int (a \sin(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral((a*sin(d*x + c) + a)^(4/3), x)`

Sympy [F]

$$\int (a + a \sin(c + dx))^{4/3} dx = \int (a \sin(c + dx) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*sin(d*x+c))**(4/3),x)`

output `Integral((a*sin(c + d*x) + a)**(4/3), x)`

Maxima [F]

$$\int (a + a \sin(c + dx))^{4/3} dx = \int (a \sin(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(4/3), x)`

Giac [F]

$$\int (a + a \sin(c + dx))^{4/3} dx = \int (a \sin(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^(4/3), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^{4/3} dx = \int (a + a \sin(c + dx))^{4/3} dx$$

input `int((a + a*sin(c + d*x))^(4/3),x)`

output `int((a + a*sin(c + d*x))^(4/3), x)`

Reduce [F]

$$\int (a + a \sin(c + dx))^{4/3} dx = a^{\frac{4}{3}} \left(\int (\sin(dx + c) + 1)^{\frac{1}{3}} dx + \int (\sin(dx + c) + 1)^{\frac{1}{3}} \sin(dx + c) dx \right)$$

input `int((a+a*sin(d*x+c))^(4/3),x)`

output `a**(1/3)*a*(int((sin(c + d*x) + 1)**(1/3),x) + int((sin(c + d*x) + 1)**(1/3)*sin(c + d*x),x))`

3.35 $\int (a + a \sin(c + dx))^{2/3} dx$

Optimal result	264
Mathematica [A] (verified)	264
Rubi [A] (verified)	265
Maple [F]	266
Fricas [F]	266
Sympy [F]	267
Maxima [F]	267
Giac [F]	267
Mupad [F(-1)]	268
Reduce [F]	268

Optimal result

Integrand size = 14, antiderivative size = 66

$$\int (a + a \sin(c + dx))^{2/3} dx = \frac{2\sqrt[6]{2} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right) (a + a \sin(c + dx))^{2/3}}{d(1 + \sin(c + dx))^{7/6}}$$

output

```
-2*2^(1/6)*cos(d*x+c)*hypergeom([-1/6, 1/2], [3/2], 1/2-1/2*sin(d*x+c))*(a+a
*sin(d*x+c))^(2/3)/d/(1+sin(d*x+c))^(7/6)
```

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.88

$$\int (a + a \sin(c + dx))^{2/3} dx = \frac{3(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) \left(-2 \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \sin^2\left(\frac{1}{4}(2c + \pi + 2dx)\right)\right) + \sqrt{2 - 2\sin(c + dx)}\right)}{2d(\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) \sqrt{2 - 2\sin(c + dx)}}$$

input

```
Integrate[(a + a*Sin[c + d*x])^(2/3), x]
```

output

```
(-3*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(-2*Hypergeometric2F1[1/6, 1/2,
7/6, Sin[(2*c + Pi + 2*d*x)/4]^2] + Sqrt[2 - 2*Sin[c + d*x]])*(a*(1 + Sin[
c + d*x]))^(2/3))/(2*d*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*Sqrt[2 - 2*Si
n[c + d*x]])
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(c + dx) + a)^{2/3} dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(c + dx) + a)^{2/3} dx$$

$$\downarrow \text{3131}$$

$$\frac{(a \sin(c + dx) + a)^{2/3} \int (\sin(c + dx) + 1)^{2/3} dx}{(\sin(c + dx) + 1)^{2/3}}$$

$$\downarrow \text{3042}$$

$$\frac{(a \sin(c + dx) + a)^{2/3} \int (\sin(c + dx) + 1)^{2/3} dx}{(\sin(c + dx) + 1)^{2/3}}$$

$$\downarrow \text{3130}$$

$$-\frac{2\sqrt[6]{2} \cos(c + dx) (a \sin(c + dx) + a)^{2/3} \text{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d(\sin(c + dx) + 1)^{7/6}}$$

input

```
Int[(a + a*Sin[c + d*x])^(2/3),x]
```

output

```
(-2*2^(1/6)*Cos[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 3/2, (1 - Sin[c + d*
x])/2]*(a + a*Sin[c + d*x])^(2/3))/(d*(1 + Sin[c + d*x])^(7/6))
```

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

Maple [F]

$$\int (a + a \sin(dx + c))^{\frac{2}{3}} dx$$

input `int((a+a*sin(d*x+c))^(2/3),x)`

output `int((a+a*sin(d*x+c))^(2/3),x)`

Fricas [F]

$$\int (a + a \sin(c + dx))^{2/3} dx = \int (a \sin(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((a*sin(d*x + c) + a)^(2/3), x)`

Sympy [F]

$$\int (a + a \sin(c + dx))^{2/3} dx = \int (a \sin(c + dx) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*sin(d*x+c))**(2/3),x)`

output `Integral((a*sin(c + d*x) + a)**(2/3), x)`

Maxima [F]

$$\int (a + a \sin(c + dx))^{2/3} dx = \int (a \sin(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(2/3), x)`

Giac [F]

$$\int (a + a \sin(c + dx))^{2/3} dx = \int (a \sin(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^(2/3), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^{2/3} dx = \int (a + a \sin(c + dx))^{2/3} dx$$

input `int((a + a*sin(c + d*x))^(2/3),x)`output `int((a + a*sin(c + d*x))^(2/3), x)`**Reduce [F]**

$$\int (a + a \sin(c + dx))^{2/3} dx = a^{2/3} \left(\int (\sin(dx + c) + 1)^{2/3} dx \right)$$

input `int((a+a*sin(d*x+c))^(2/3),x)`output `a**(2/3)*int((sin(c + d*x) + 1)**(2/3),x)`

3.36 $\int \sqrt[3]{a + a \sin(c + dx)} dx$

Optimal result	269
Mathematica [B] (warning: unable to verify)	269
Rubi [A] (verified)	270
Maple [F]	271
Fricas [F]	272
Sympy [F]	272
Maxima [F]	272
Giac [F]	273
Mupad [F(-1)]	273
Reduce [F]	273

Optimal result

Integrand size = 14, antiderivative size = 66

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \frac{2^{5/6} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right) \sqrt[3]{a + a \sin(c + dx)}}{d(1 + \sin(c + dx))^{5/6}}$$

output

```
-2^(5/6)*cos(d*x+c)*hypergeom([1/6, 1/2], [3/2], 1/2-1/2*sin(d*x+c))*(a+a*sin(d*x+c))^(1/3)/d/(1+sin(d*x+c))^(5/6)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 169 vs. 2(66) = 132.

Time = 0.70 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.56

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \frac{\sqrt[3]{2} \left(2 \cos\left(\frac{1}{4}(2c + \pi + 2dx)\right) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{6}; \frac{5}{6}; \sin^2\left(\frac{1}{4}(2c + \pi + 2dx)\right)\right) + \left(-\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right) \right)}{d \sqrt{\cos^2\left(\frac{1}{4}(2c + \pi + 2dx)\right)} \left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right)^{2/3} \sqrt[3]{\sin\left(\frac{1}{4}(2c + \pi + 2dx)\right)}}$$

input `Integrate[(a + a*Sin[c + d*x])^(1/3),x]`

output `(2^(1/3)*(2*Cos[(2*c + Pi + 2*d*x)/4]*HypergeometricPFQ[{-1/2, -1/6}, {5/6}, Sin[(2*c + Pi + 2*d*x)/4]^2] + (-Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*Sqrt[1 - Sin[c + d*x]])*(a*(1 + Sin[c + d*x]))^(1/3)/(d*Sqrt[Cos[(2*c + Pi + 2*d*x)/4]^2]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^(2/3)*Sin[(2*c + Pi + 2*d*x)/4]^(1/3))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{a \sin(c + dx) + a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{a \sin(c + dx) + a} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{\sqrt[3]{a \sin(c + dx) + a} \int \sqrt[3]{\sin(c + dx) + 1} dx}{\sqrt[3]{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{a \sin(c + dx) + a} \int \sqrt[3]{\sin(c + dx) + 1} dx}{\sqrt[3]{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{2^{5/6} \cos(c + dx) \sqrt[3]{a \sin(c + dx) + a} \text{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d(\sin(c + dx) + 1)^{5/6}}
 \end{aligned}$$

input `Int[(a + a*Sin[c + d*x])^(1/3),x]`

output $-\left(\frac{2^{5/6} \cos[c + dx] \operatorname{Hypergeometric2F1}\left[\frac{1}{6}, \frac{1}{2}, \frac{3}{2}, (1 - \sin[c + dx])\right]}{2} (a + a \sin[c + dx])^{1/3} / (d(1 + \sin[c + dx])^{5/6})\right)$

Defintions of rubi rules used

rule 3042 $\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3130 $\operatorname{Int}[(a_ + (b_)\sin[(c_ + (d_)(x_)])^{n_}, x_Symbol] \rightarrow \operatorname{Simp}[(-2^{(n + 1/2)})a^{(n - 1/2)}b(\cos[c + dx]/(d\sqrt{a + b\sin[c + dx]}))\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - b(\sin[c + dx]/a))\right], x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \operatorname{EqQ}[a^2 - b^2, 0] \ \&\& \ !\operatorname{IntegerQ}[2n] \ \&\& \operatorname{GtQ}[a, 0]$

rule 3131 $\operatorname{Int}[(a_ + (b_)\sin[(c_ + (d_)(x_)])^{n_}, x_Symbol] \rightarrow \operatorname{Simp}[a^{\operatorname{IntPart}[n]}((a + b\sin[c + dx])^{\operatorname{FracPart}[n]} / (1 + (b/a)\sin[c + dx])^{\operatorname{FracPart}[n]}) \operatorname{Int}[(1 + (b/a)\sin[c + dx])^n, x], x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \operatorname{EqQ}[a^2 - b^2, 0] \ \&\& \ !\operatorname{IntegerQ}[2n] \ \&\& \ !\operatorname{GtQ}[a, 0]$

Maple [F]

$$\int (a + a \sin(dx + c))^{\frac{1}{3}} dx$$

input $\operatorname{int}((a+a\sin(dx+c))^{1/3},x)$

output $\operatorname{int}((a+a\sin(dx+c))^{1/3},x)$

Fricas [F]

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \int (a \sin(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((a*sin(d*x + c) + a)^(1/3), x)`

Sympy [F]

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \int \sqrt[3]{a \sin(c + dx) + a} dx$$

input `integrate((a+a*sin(d*x+c))**(1/3),x)`

output `Integral((a*sin(c + d*x) + a)**(1/3), x)`

Maxima [F]

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \int (a \sin(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(1/3), x)`

Giac [F]

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \int (a \sin(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+a*sin(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^(1/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = \int (a + a \sin(c + dx))^{1/3} dx$$

input `int((a + a*sin(c + d*x))^(1/3),x)`

output `int((a + a*sin(c + d*x))^(1/3), x)`

Reduce [F]

$$\int \sqrt[3]{a + a \sin(c + dx)} dx = a^{\frac{1}{3}} \left(\int (\sin(dx + c) + 1)^{\frac{1}{3}} dx \right)$$

input `int((a+a*sin(d*x+c))^(1/3),x)`

output `a**(1/3)*int((sin(c + d*x) + 1)**(1/3),x)`

3.37
$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx$$

Optimal result	274
Mathematica [A] (verified)	274
Rubi [A] (verified)	275
Maple [F]	276
Fricas [F]	276
Sympy [F]	277
Maxima [F]	277
Giac [F]	277
Mupad [F(-1)]	278
Reduce [F]	278

Optimal result

Integrand size = 14, antiderivative size = 66

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = -\frac{\sqrt[6]{2} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d \sqrt[6]{1 + \sin(c + dx)} \sqrt[3]{a + a \sin(c + dx)}}$$

output `-2^(1/6)*cos(d*x+c)*hypergeom([1/2, 5/6], [3/2], 1/2-1/2*sin(d*x+c))/d/(1+sin(d*x+c))^(1/6)/(a+a*sin(d*x+c))^(1/3)`

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \frac{3\sqrt{2} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \sin^2\left(\frac{1}{4}(2c + \pi + 2dx)\right)\right)}{d \sqrt{1 - \sin(c + dx)} \sqrt[3]{a(1 + \sin(c + dx))}}$$

input `Integrate[(a + a*Sin[c + d*x])^(-1/3), x]`

output

```
(3*Sqrt[2]*Cos[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sin[(2*c + Pi + 2
*d*x)/4]^2])/(d*Sqrt[1 - Sin[c + d*x]]*(a*(1 + Sin[c + d*x]))^(1/3))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{a \sin(c + dx) + a}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{a \sin(c + dx) + a}} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{\sqrt[3]{\sin(c + dx) + 1} \int \frac{1}{\sqrt[3]{\sin(c + dx) + 1}} dx}{\sqrt[3]{a \sin(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\sin(c + dx) + 1} \int \frac{1}{\sqrt[3]{\sin(c + dx) + 1}} dx}{\sqrt[3]{a \sin(c + dx) + a}} \\
 & \quad \downarrow \text{3130} \\
 & -\frac{\sqrt[6]{2} \cos(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d \sqrt[6]{\sin(c + dx) + 1} \sqrt[3]{a \sin(c + dx) + a}}
 \end{aligned}$$

input

```
Int[(a + a*Sin[c + d*x])^(-1/3),x]
```

output

```
-((2^(1/6)*Cos[c + d*x]*Hypergeometric2F1[1/2, 5/6, 3/2, (1 - Sin[c + d*x]
)/2])/(d*(1 + Sin[c + d*x])^(1/6)*(a + a*Sin[c + d*x])^(1/3)))
```

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

Maple [F]

$$\int \frac{1}{(a + a \sin(dx + c))^{\frac{1}{3}}} dx$$

input `int(1/(a+a*sin(d*x+c))^(1/3),x)`

output `int(1/(a+a*sin(d*x+c))^(1/3),x)`

Fricas [F]

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((a*sin(d*x + c) + a)^(-1/3), x)`

Sympy [F]

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \int \frac{1}{\sqrt[3]{a \sin(c + dx) + a}} dx$$

input `integrate(1/(a+a*sin(d*x+c))**(1/3),x)`

output `Integral((a*sin(c + d*x) + a)**(-1/3), x)`

Maxima [F]

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(-1/3), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^(-1/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \int \frac{1}{(a + a \sin(c + dx))^{1/3}} dx$$

input `int(1/(a + a*sin(c + d*x))^(1/3),x)`output `int(1/(a + a*sin(c + d*x))^(1/3), x)`**Reduce [F]**

$$\int \frac{1}{\sqrt[3]{a + a \sin(c + dx)}} dx = \frac{\int \frac{1}{(\sin(dx+c)+1)^{\frac{1}{3}}} dx}{a^{\frac{1}{3}}}$$

input `int(1/(a+a*sin(d*x+c))^(1/3),x)`output `int(1/(sin(c + d*x) + 1)**(1/3),x)/a**(1/3)`

3.38 $\int \frac{1}{(a+a \sin(c+dx))^{2/3}} dx$

Optimal result	279
Mathematica [B] (warning: unable to verify)	279
Rubi [A] (verified)	280
Maple [F]	281
Fricas [F]	282
Sympy [F]	282
Maxima [F]	282
Giac [F]	283
Mupad [F(-1)]	283
Reduce [F]	283

Optimal result

Integrand size = 14, antiderivative size = 66

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \frac{\cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right) \sqrt[6]{1 + \sin(c + dx)}}{\sqrt[6]{2d(a + a \sin(c + dx))^{2/3}}}$$

output

```
-1/2*cos(d*x+c)*hypergeom([1/2, 7/6],[3/2],1/2-1/2*sin(d*x+c))*(1+sin(d*x+c))^(1/6)*2^(5/6)/d/(a+a*sin(d*x+c))^(2/3)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 215 vs. 2(66) = 132.

Time = 1.13 (sec) , antiderivative size = 215, normalized size of antiderivative = 3.26

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \frac{2 \left(-3 \cos\left(\frac{1}{2}(c + dx)\right) \left(\cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right) \right) + \frac{(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\dots} \right)}{\dots}$$

input `Integrate[(a + a*Sin[c + d*x])^(-2/3),x]`

output `(2*(-3*Cos[(c + d*x)/2]*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) + ((Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^(4/3)*(-2*Cos[(2*c + Pi + 2*d*x)/4]*HypergeometricPFQ[{-1/2, -1/6}, {5/6}, Sin[(2*c + Pi + 2*d*x)/4]^2] + Sqrt[Cos[(2*c + Pi + 2*d*x)/4]^2]*(2*Cos[(2*c + Pi + 2*d*x)/4] + 3*Sin[(2*c + Pi + 2*d*x)/4])))^(1/3))/(2^(1/6)*Sqrt[1 - Sin[c + d*x]]*Sin[(2*c + Pi + 2*d*x)/4]^(1/3)))/(d*(a*(1 + Sin[c + d*x]))^(2/3))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sin(c + dx) + a)^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx) + a)^{2/3}} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{(\sin(c + dx) + 1)^{2/3} \int \frac{1}{(\sin(c + dx) + 1)^{2/3}} dx}{(a \sin(c + dx) + a)^{2/3}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(\sin(c + dx) + 1)^{2/3} \int \frac{1}{(\sin(c + dx) + 1)^{2/3}} dx}{(a \sin(c + dx) + a)^{2/3}} \\
 & \quad \downarrow \text{3130} \\
 & - \frac{\sqrt[6]{\sin(c + dx) + 1} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{\sqrt[6]{2d(a \sin(c + dx) + a)^{2/3}}}
 \end{aligned}$$

input `Int[(a + a*Sin[c + d*x])^(-2/3),x]`

output `-((Cos[c + d*x]*Hypergeometric2F1[1/2, 7/6, 3/2, (1 - Sin[c + d*x])/2]*(1 + Sin[c + d*x])^(1/6))/(2^(1/6)*d*(a + a*Sin[c + d*x])^(2/3))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

Maple [F]

$$\int \frac{1}{(a + a \sin(dx + c))^{\frac{2}{3}}} dx$$

input `int(1/(a+a*sin(d*x+c))^(2/3),x)`

output `int(1/(a+a*sin(d*x+c))^(2/3),x)`

Fricas [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((a*sin(d*x + c) + a)^(-2/3), x)`

Sympy [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a \sin(c + dx) + a)^{2/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))**(2/3),x)`

output `Integral((a*sin(c + d*x) + a)**(-2/3), x)`

Maxima [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(-2/3), x)`

Giac [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^(-2/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx$$

input `int(1/(a + a*sin(c + d*x))^(2/3),x)`

output `int(1/(a + a*sin(c + d*x))^(2/3), x)`

Reduce [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{2/3}} dx = \frac{\int \frac{1}{(\sin(dx+c)+1)^{2/3}} dx}{a^{2/3}}$$

input `int(1/(a+a*sin(d*x+c))^(2/3),x)`

output `int(1/(sin(c + d*x) + 1)**(2/3),x)/a**(2/3)`

3.39 $\int \frac{1}{(a+a \sin(c+dx))^{4/3}} dx$

Optimal result	284
Mathematica [A] (verified)	284
Rubi [A] (verified)	285
Maple [F]	286
Fricas [F]	287
Sympy [F]	287
Maxima [F]	287
Giac [F]	288
Mupad [F(-1)]	288
Reduce [F]	288

Optimal result

Integrand size = 14, antiderivative size = 69

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \frac{\cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{2^{5/6} a d \sqrt[6]{1 + \sin(c + dx)} \sqrt[3]{a + a \sin(c + dx)}}$$

output

```
-1/2*cos(d*x+c)*hypergeom([1/2, 11/6],[3/2],1/2-1/2*sin(d*x+c))*2^(1/6)/a/
d/(1+sin(d*x+c))^(1/6)/(a+a*sin(d*x+c))^(1/3)
```

Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.88

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \frac{3(\cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) (\cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) (\sqrt{2 - 2 \sin(c + dx)} - 2 \operatorname{Hypergeometric2F1}(\frac{1}{2}, \frac{11}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))))}{5d\sqrt{2 - 2 \sin(c + dx)}(a(1 + \sin(c + dx)))}$$

input

```
Integrate[(a + a*Sin[c + d*x])^(-4/3),x]
```

output

```
(-3*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])*(Sqrt[2 - 2*Sin[c + d*x]] - 2*Hypergeometric2F1[1/6, 1/2, 7/6, Sin[(2*c + Pi + 2*d*x)/4]^2]*(1 + Sin[c + d*x]))/(5*d*Sqrt[2 - 2*Sin[c + d*x]]*(a*(1 + Sin[c + d*x]))^(4/3))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \sin(c + dx) + a)^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a \sin(c + dx) + a)^{4/3}} dx \\
 & \quad \downarrow \text{3131} \\
 & \frac{\sqrt[3]{\sin(c + dx) + 1} \int \frac{1}{(\sin(c + dx) + 1)^{4/3}} dx}{a \sqrt[3]{a \sin(c + dx) + a}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt[3]{\sin(c + dx) + 1} \int \frac{1}{(\sin(c + dx) + 1)^{4/3}} dx}{a \sqrt[3]{a \sin(c + dx) + a}} \\
 & \quad \downarrow \text{3130} \\
 & \frac{\cos(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{2^{5/6} a d \sqrt[6]{\sin(c + dx) + 1} \sqrt[3]{a \sin(c + dx) + a}}
 \end{aligned}$$

input

```
Int[(a + a*Sin[c + d*x])^(-4/3), x]
```

output $-\left(\frac{\cos[c + dx] \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{11}{6}, \frac{3}{2}, \frac{1 - \sin[c + dx]}{2}\right]}{2^{5/6} a d (1 + \sin[c + dx])^{1/6} (a + a \sin[c + dx])^{1/3}}\right)$

Definitions of rubi rules used

rule 3042 $\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear } Q[u, x]$

rule 3130 $\operatorname{Int}[(a_ + (b_)\sin[(c_ + (d_)(x_)])^{n_}, x_Symbol] \rightarrow \operatorname{Simp}[(-2^{(n + 1/2)})a^{(n - 1/2)}b(\cos[c + dx]/(d\sqrt{a + b\sin[c + dx]}))\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - b(\sin[c + dx]/a))\right], x] \text{ ; FreeQ}\{a, b, c, d, n\}, x] \ \&\& \operatorname{EqQ}[a^2 - b^2, 0] \ \&\& \operatorname{!IntegerQ}[2n] \ \&\& \operatorname{GtQ}[a, 0]$

rule 3131 $\operatorname{Int}[(a_ + (b_)\sin[(c_ + (d_)(x_)])^{n_}, x_Symbol] \rightarrow \operatorname{Simp}[a^{\operatorname{IntPart}[n]}((a + b\sin[c + dx])^{\operatorname{FracPart}[n]} / (1 + (b/a)\sin[c + dx])^{\operatorname{FracPart}[n]}) \operatorname{Int}[(1 + (b/a)\sin[c + dx])^n, x], x] \text{ ; FreeQ}\{a, b, c, d, n\}, x] \ \&\& \operatorname{EqQ}[a^2 - b^2, 0] \ \&\& \operatorname{!IntegerQ}[2n] \ \&\& \operatorname{!GtQ}[a, 0]$

Maple [F]

$$\int \frac{1}{(a + a \sin(dx + c))^{4/3}} dx$$

input $\operatorname{int}(1/(a+a\sin(dx+c))^{4/3},x)$

output $\operatorname{int}(1/(a+a\sin(dx+c))^{4/3},x)$

Fricas [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(4/3),x, algorithm="fricas")`

output `integral(-(a*sin(d*x + c) + a)^(2/3)/(a^2*cos(d*x + c)^2 - 2*a^2*sin(d*x + c) - 2*a^2), x)`

Sympy [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a \sin(c + dx) + a)^{4/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))**(4/3),x)`

output `Integral((a*sin(c + d*x) + a)**(-4/3), x)`

Maxima [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^(-4/3), x)`

Giac [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a \sin(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+a*sin(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^(-4/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx$$

input `int(1/(a + a*sin(c + d*x))^(4/3),x)`

output `int(1/(a + a*sin(c + d*x))^(4/3), x)`

Reduce [F]

$$\int \frac{1}{(a + a \sin(c + dx))^{4/3}} dx = \frac{\int \frac{1}{(\sin(dx+c)+1)^{1/3} \sin(dx+c) + (\sin(dx+c)+1)^{1/3}} dx}{a^{4/3}}$$

input `int(1/(a+a*sin(d*x+c))^(4/3),x)`

output `int(1/((sin(c + d*x) + 1)**(1/3)*sin(c + d*x) + (sin(c + d*x) + 1)**(1/3)),x)/(a**(1/3)*a)`

3.40 $\int (a + a \sin(c + dx))^n dx$

Optimal result	289
Mathematica [C] (verified)	289
Rubi [A] (verified)	290
Maple [F]	291
Fricas [F]	291
Sympy [F]	292
Maxima [F]	292
Giac [F]	292
Mupad [F(-1)]	293
Reduce [F]	293

Optimal result

Integrand size = 12, antiderivative size = 74

$$\int (a + a \sin(c + dx))^n dx = \frac{2^{\frac{1}{2}+n} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right) (1 + \sin(c + dx))^{-\frac{1}{2}-n} (a + a \sin(c + dx))^n}{d}$$

output

```
-2^(1/2+n)*cos(d*x+c)*hypergeom([1/2, 1/2-n], [3/2], 1/2-1/2*sin(d*x+c))*(1+sin(d*x+c))^(1/2-n)*(a+a*sin(d*x+c))^n/d
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 5 in optimal.

Time = 0.19 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.95

$$\int (a + a \sin(c + dx))^n dx = \frac{2^n B_{\frac{1}{2}(1+\sin(c+dx))}\left(\frac{1}{2} + n, \frac{1}{2}\right) \sqrt{\cos^2(c + dx)} \sec(c + dx) (1 + \sin(c + dx))^{-n} (a(1 + \sin(c + dx)))^n}{d}$$

input

```
Integrate[(a + a*Sin[c + d*x])^n,x]
```

output

$$(2^n \text{Beta}[(1 + \text{Sin}[c + d*x])/2, 1/2 + n, 1/2] * \text{Sqrt}[\text{Cos}[c + d*x]^2] * \text{Sec}[c + d*x] * (a * (1 + \text{Sin}[c + d*x]))^n) / (d * (1 + \text{Sin}[c + d*x])^n)$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \sin(c + dx) + a)^n dx \\ & \quad \downarrow \text{3042} \\ & \int (a \sin(c + dx) + a)^n dx \\ & \quad \downarrow \text{3131} \\ & (\sin(c + dx) + 1)^{-n} (a \sin(c + dx) + a)^n \int (\sin(c + dx) + 1)^n dx \\ & \quad \downarrow \text{3042} \\ & (\sin(c + dx) + 1)^{-n} (a \sin(c + dx) + a)^n \int (\sin(c + dx) + 1)^n dx \\ & \quad \downarrow \text{3130} \\ & \frac{2^{n+\frac{1}{2}} \cos(c + dx) (\sin(c + dx) + 1)^{-n-\frac{1}{2}} (a \sin(c + dx) + a)^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d} \end{aligned}$$

input

$$\text{Int}[(a + a*\text{Sin}[c + d*x])^n, x]$$

output

$$-((2^{(1/2 + n)} * \text{Cos}[c + d*x] * \text{Hypergeometric2F1}[1/2, 1/2 - n, 3/2, (1 - \text{Sin}[c + d*x])/2]) * (1 + \text{Sin}[c + d*x])^{(-1/2 - n)} * (a + a*\text{Sin}[c + d*x])^n) / d$$

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :=> Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

Maple [F]

$$\int (a + a \sin(dx + c))^n dx$$

input `int((a+a*sin(d*x+c))^n,x)`

output `int((a+a*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (a + a \sin(c + dx))^n dx = \int (a \sin(dx + c) + a)^n dx$$

input `integrate((a+a*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((a*sin(d*x + c) + a)^n, x)`

Sympy [F]

$$\int (a + a \sin(c + dx))^n dx = \int (a \sin(c + dx) + a)^n dx$$

input `integrate((a+a*sin(d*x+c))**n,x)`

output `Integral((a*sin(c + d*x) + a)**n, x)`

Maxima [F]

$$\int (a + a \sin(c + dx))^n dx = \int (a \sin(dx + c) + a)^n dx$$

input `integrate((a+a*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((a*sin(d*x + c) + a)^n, x)`

Giac [F]

$$\int (a + a \sin(c + dx))^n dx = \int (a \sin(dx + c) + a)^n dx$$

input `integrate((a+a*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((a*sin(d*x + c) + a)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(c + dx))^n dx = \int (a + a \sin(c + dx))^n dx$$

input `int((a + a*sin(c + d*x))^n,x)`output `int((a + a*sin(c + d*x))^n, x)`**Reduce [F]**

$$\int (a + a \sin(c + dx))^n dx = \int (\sin(dx + c) a + a)^n dx$$

input `int((a+a*sin(d*x+c))^n,x)`output `int((sin(c + d*x)*a + a)**n,x)`

3.41 $\int (a - a \sin(c + dx))^n dx$

Optimal result	294
Mathematica [C] (verified)	294
Rubi [A] (verified)	295
Maple [F]	296
Fricas [F]	296
Sympy [F]	297
Maxima [F]	297
Giac [F]	297
Mupad [F(-1)]	298
Reduce [F]	298

Optimal result

Integrand size = 13, antiderivative size = 74

$$\int (a - a \sin(c + dx))^n dx = \frac{2^{\frac{1}{2}+n} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 + \sin(c + dx))\right) (1 - \sin(c + dx))^{-\frac{1}{2}-n} (a - a \sin(c + dx))^n}{d}$$

output

$2^{1/2+n} \cos(d*x+c) \operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}-n\right], \left[\frac{3}{2}\right], \frac{1}{2} + \frac{1}{2} \sin(d*x+c)\right) (1 - \sin(d*x+c))^{-1/2-n} (a - a \sin(d*x+c))^n / d$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 5 in optimal.

Time = 0.29 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.08

$$\int (a - a \sin(c + dx))^n dx = \frac{2^n B_{\frac{1}{2}(1-\sin(c+dx))}\left(\frac{1}{2} + n, \frac{1}{2}\right) \sec(c + dx) (1 - \sin(c + dx))^{\frac{1}{2}-n} \sqrt{1 + \sin(c + dx)} (a - a \sin(c + dx))^n}{d}$$

input

`Integrate[(a - a*Sin[c + d*x])^n, x]`

output

$$-\left(2^n \text{Beta}\left[\frac{1 - \sin[c + dx]}{2}, \frac{1}{2} + n, \frac{1}{2}\right] \text{Sec}[c + dx] (1 - \sin[c + dx])^{1/2 - n} \sqrt{1 + \sin[c + dx]} (a - a \sin[c + dx])^n\right) / d$$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 3131, 3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a - a \sin(c + dx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int (a - a \sin(c + dx))^n dx \\ & \quad \downarrow \text{3131} \\ & (1 - \sin(c + dx))^{-n} (a - a \sin(c + dx))^n \int (1 - \sin(c + dx))^n dx \\ & \quad \downarrow \text{3042} \\ & (1 - \sin(c + dx))^{-n} (a - a \sin(c + dx))^n \int (1 - \sin(c + dx))^n dx \\ & \quad \downarrow \text{3130} \\ & \frac{2^{n+\frac{1}{2}} \cos(c + dx) (1 - \sin(c + dx))^{-n-\frac{1}{2}} (a - a \sin(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(\sin(c + dx) + 1)\right)}{d} \end{aligned}$$

input

$$\text{Int}[(a - a \sin[c + dx])^n, x]$$

output

$$\left(2^{1/2 + n} \text{Cos}[c + dx] \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1 + \sin[c + dx]}{2}\right] (1 - \sin[c + dx])^{-1/2 - n} (a - a \sin[c + dx])^n\right) / d$$

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]]))*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

rule 3131 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[a^IntPart[n]*((a + b*Sin[c + d*x])^FracPart[n]/(1 + (b/a)*Sin[c + d*x])^FracPart[n]) Int[(1 + (b/a)*Sin[c + d*x])^n, x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && !GtQ[a, 0]`

Maple [F]

$$\int (a - a \sin(dx + c))^n dx$$

input `int((a-a*sin(d*x+c))^n,x)`

output `int((a-a*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (a - a \sin(c + dx))^n dx = \int (-a \sin(dx + c) + a)^n dx$$

input `integrate((a-a*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((-a*sin(d*x + c) + a)^n, x)`

Sympy [F]

$$\int (a - a \sin(c + dx))^n dx = \int (-a \sin(c + dx) + a)^n dx$$

input `integrate((a-a*sin(d*x+c))**n,x)`

output `Integral((-a*sin(c + d*x) + a)**n, x)`

Maxima [F]

$$\int (a - a \sin(c + dx))^n dx = \int (-a \sin(dx + c) + a)^n dx$$

input `integrate((a-a*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-a*sin(d*x + c) + a)^n, x)`

Giac [F]

$$\int (a - a \sin(c + dx))^n dx = \int (-a \sin(dx + c) + a)^n dx$$

input `integrate((a-a*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((-a*sin(d*x + c) + a)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a - a \sin(c + dx))^n dx = \int (a - a \sin(c + dx))^n dx$$

input `int((a - a*sin(c + d*x))^n,x)`output `int((a - a*sin(c + d*x))^n, x)`**Reduce [F]**

$$\int (a - a \sin(c + dx))^n dx = \int (-\sin(dx + c) a + a)^n dx$$

input `int((a-a*sin(d*x+c))^n,x)`output `int((-sin(c + d*x)*a + a)**n,x)`

3.42 $\int (2 + 2 \sin(c + dx))^n dx$

Optimal result	299
Mathematica [C] (verified)	299
Rubi [A] (verified)	300
Maple [F]	301
Fricas [F]	301
Sympy [F]	302
Maxima [F]	302
Giac [F]	302
Mupad [F(-1)]	303
Reduce [F]	303

Optimal result

Integrand size = 12, antiderivative size = 60

$$\int (2 + 2 \sin(c + dx))^n dx = -\frac{2^{\frac{1}{2}+2n} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d \sqrt{1 + \sin(c + dx)}}$$

output

```
-2^(1/2+2*n)*cos(d*x+c)*hypergeom([1/2, 1/2-n], [3/2], 1/2-1/2*sin(d*x+c))/d
/(1+sin(d*x+c))^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 5 in optimal.

Time = 0.22 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.77

$$\int (2 + 2 \sin(c + dx))^n dx = \frac{4^n B_{\frac{1}{2}(1+\sin(c+dx))}\left(\frac{1}{2} + n, \frac{1}{2}\right) \sqrt{\cos^2(c + dx)} \sec(c + dx)}{d}$$

input

```
Integrate[(2 + 2*Sin[c + d*x])^n,x]
```

output $(4^n \text{Beta}[(1 + \text{Sin}[c + d*x])/2, 1/2 + n, 1/2] \text{Sqrt}[\text{Cos}[c + d*x]^2] \text{Sec}[c + d*x])/d$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2 \sin(c + dx) + 2)^n dx$$

$$\downarrow \text{3042}$$

$$\int (2 \sin(c + dx) + 2)^n dx$$

$$\downarrow \text{3130}$$

$$-\frac{2^{2n+\frac{1}{2}} \cos(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx))\right)}{d \sqrt{\sin(c + dx) + 1}}$$

input $\text{Int}[(2 + 2*\text{Sin}[c + d*x])^n, x]$

output $-((2^{(1/2 + 2*n)} \text{Cos}[c + d*x] \text{Hypergeometric2F1}[1/2, 1/2 - n, 3/2, (1 - \text{Sin}[c + d*x])/2]) / (d \text{Sqrt}[1 + \text{Sin}[c + d*x]]))$

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

Maple [F]

$$\int (2 + 2 \sin(dx + c))^n dx$$

input `int((2+2*sin(d*x+c))^n,x)`

output `int((2+2*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (2 + 2 \sin(c + dx))^n dx = \int (2 \sin(dx + c) + 2)^n dx$$

input `integrate((2+2*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((2*sin(d*x + c) + 2)^n, x)`

Sympy [F]

$$\int (2 + 2 \sin(c + dx))^n dx = 2^n \int (\sin(c + dx) + 1)^n dx$$

input `integrate((2+2*sin(d*x+c))**n,x)`

output `2**n*Integral((sin(c + d*x) + 1)**n, x)`

Maxima [F]

$$\int (2 + 2 \sin(c + dx))^n dx = \int (2 \sin(dx + c) + 2)^n dx$$

input `integrate((2+2*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((2*sin(d*x + c) + 2)^n, x)`

Giac [F]

$$\int (2 + 2 \sin(c + dx))^n dx = \int (2 \sin(dx + c) + 2)^n dx$$

input `integrate((2+2*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((2*sin(d*x + c) + 2)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (2 + 2 \sin(c + dx))^n dx = \int (2 \sin(c + dx) + 2)^n dx$$

input `int((2*sin(c + d*x) + 2)^n,x)`output `int((2*sin(c + d*x) + 2)^n, x)`**Reduce [F]**

$$\int (2 + 2 \sin(c + dx))^n dx = \int (2 \sin(dx + c) + 2)^n dx$$

input `int((2+2*sin(d*x+c))^n,x)`output `int((2*sin(c + d*x) + 2)**n,x)`

3.43 $\int (2 - 2 \sin(c + dx))^n dx$

Optimal result	304
Mathematica [C] (verified)	304
Rubi [A] (verified)	305
Maple [F]	306
Fricas [F]	306
Sympy [F]	307
Maxima [F]	307
Giac [F]	307
Mupad [F(-1)]	308
Reduce [F]	308

Optimal result

Integrand size = 12, antiderivative size = 59

$$\int (2 - 2 \sin(c + dx))^n dx = \frac{2^{\frac{1}{2}+2n} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(1 + \sin(c + dx))\right)}{d \sqrt{1 - \sin(c + dx)}}$$

output

$2^{(1/2+2*n)}*\cos(d*x+c)*\operatorname{hypergeom}([1/2, 1/2-n], [3/2], 1/2+1/2*\sin(d*x+c))/d/(1-\sin(d*x+c))^{(1/2)}$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 5 in optimal.

Time = 0.22 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int (2 - 2 \sin(c + dx))^n dx = -\frac{4^n B_{\frac{1}{2}(1-\sin(c+dx))}\left(\frac{1}{2} + n, \frac{1}{2}\right) \sqrt{\cos^2(c + dx)} \sec(c + dx)}{d}$$

input

`Integrate[(2 - 2*Sin[c + d*x])^n,x]`

output

```

-((4^n*Beta[(1 - Sin[c + d*x])/2, 1/2 + n, 1/2]*Sqrt[Cos[c + d*x]^2]*Sec[c
+ d*x])/d)

```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3130}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int (2 - 2 \sin(c + dx))^n dx \\
 \downarrow \text{3042} \\
 \int (2 - 2 \sin(c + dx))^n dx \\
 \downarrow \text{3130} \\
 \frac{2^{2n+\frac{1}{2}} \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}, \frac{1}{2}(\sin(c + dx) + 1)\right)}{d \sqrt{1 - \sin(c + dx)}}
 \end{array}$$

input

```

Int[(2 - 2*Sin[c + d*x])^n,x]

```

output

```

(2^(1/2 + 2*n)*Cos[c + d*x]*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1 + Sin[
c + d*x])/2])/(d*Sqrt[1 - Sin[c + d*x]])

```

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3130 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[(-2^(n + 1/2))*a^(n - 1/2)*b*(Cos[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])*Hypergeometric2F1[1/2, 1/2 - n, 3/2, (1/2)*(1 - b*(Sin[c + d*x]/a))], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[2*n] && GtQ[a, 0]`

Maple [F]

$$\int (2 - 2 \sin(dx + c))^n dx$$

input `int((2-2*sin(d*x+c))^n,x)`

output `int((2-2*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (2 - 2 \sin(c + dx))^n dx = \int (-2 \sin(dx + c) + 2)^n dx$$

input `integrate((2-2*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((-2*sin(d*x + c) + 2)^n, x)`

Sympy [F]

$$\int (2 - 2 \sin(c + dx))^n dx = \int (2 - 2 \sin(c + dx))^n dx$$

input `integrate((2-2*sin(d*x+c))**n,x)`

output `Integral((2 - 2*sin(c + d*x))**n, x)`

Maxima [F]

$$\int (2 - 2 \sin(c + dx))^n dx = \int (-2 \sin(dx + c) + 2)^n dx$$

input `integrate((2-2*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-2*sin(d*x + c) + 2)^n, x)`

Giac [F]

$$\int (2 - 2 \sin(c + dx))^n dx = \int (-2 \sin(dx + c) + 2)^n dx$$

input `integrate((2-2*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((-2*sin(d*x + c) + 2)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (2 - 2 \sin(c + dx))^n dx = \int (2 - 2 \sin(c + dx))^n dx$$

input `int((2 - 2*sin(c + d*x))^n,x)`output `int((2 - 2*sin(c + d*x))^n, x)`**Reduce [F]**

$$\int (2 - 2 \sin(c + dx))^n dx = \int (-2 \sin(dx + c) + 2)^n dx$$

input `int((2-2*sin(d*x+c))^n,x)`output `int((- 2*sin(c + d*x) + 2)**n,x)`

3.44 $\int \frac{1}{5+3\sin(c+dx)} dx$

Optimal result	309
Mathematica [A] (verified)	309
Rubi [A] (verified)	310
Maple [A] (verified)	311
Fricas [A] (verification not implemented)	311
Sympy [B] (verification not implemented)	312
Maxima [A] (verification not implemented)	312
Giac [A] (verification not implemented)	312
Mupad [B] (verification not implemented)	313
Reduce [B] (verification not implemented)	313

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{5+3\sin(c+dx)} dx = \frac{x}{4} + \frac{\arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{2d}$$

output `1/4*x+1/2*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.81

$$\int \frac{1}{5+3\sin(c+dx)} dx = \frac{\arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right)}{2d}$$

input `Integrate[(5 + 3*Sin[c + d*x])^(-1),x]`

output `ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])]/(2*d)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{3 \sin(c + dx) + 5} dx$$

↓ 3042

$$\int \frac{1}{3 \sin(c + dx) + 5} dx$$

↓ 3136

$$\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4}$$

input `Int[(5 + 3*Sin[c + d*x])^(-1),x]`

output `x/4 + ArcTan[Cos[c + d*x]/(3 + Sin[c + d*x])]/(2*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.65

method	result	size
derivativedivides	$\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}}{4}\right)}{2d}$	20
default	$\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}}{4}\right)}{2d}$	20
risch	$\frac{i \ln(e^{i(dx+c)} + 3i)}{4d} - \frac{i \ln(e^{i(dx+c)} + \frac{i}{3})}{4d}$	40
parallelrisch	$-\frac{i\left(\ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) - \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 + 4i\right)\right)}{4d}$	42

input `int(1/(5+3*sin(d*x+c)),x,method=_RETURNVERBOSE)`output `1/2/d*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{5 + 3 \sin(c + dx)} dx = \frac{\arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right)}{4d}$$

input `integrate(1/(5+3*sin(d*x+c)),x, algorithm="fricas")`output `1/4*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(22) = 44$.

Time = 0.37 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48

$$\int \frac{1}{5 + 3 \sin(c + dx)} dx = \begin{cases} \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \frac{3}{4}}{4}\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{3 \sin(c) + 5} & \text{otherwise} \end{cases}$$

input `integrate(1/(5+3*sin(d*x+c)),x)`

output `Piecewise(((atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(2*d), Ne(d, 0)), (x/(3*sin(c) + 5), True))`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{5 + 3 \sin(c + dx)} dx = \frac{\arctan\left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4}\right)}{2d}$$

input `integrate(1/(5+3*sin(d*x+c)),x, algorithm="maxima")`

output `1/2*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4)/d`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{1}{5 + 3 \sin(c + dx)} dx = \frac{dx + c + 2 \arctan\left(\frac{-3 \cos(dx+c) + \sin(dx+c) + 3}{\cos(dx+c) - 3 \sin(dx+c) - 9}\right)}{4d}$$

input `integrate(1/(5+3*sin(d*x+c)),x, algorithm="giac")`

output $1/4*(d*x + c + 2*\arctan(-(3*\cos(d*x + c) + \sin(d*x + c) + 3)/(\cos(d*x + c) - 3*\sin(d*x + c) - 9)))/d$

Mupad [B] (verification not implemented)

Time = 26.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int \frac{1}{5 + 3 \sin(c + dx)} dx = \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \frac{3}{4}}{4}\right)}{2d} - \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d}$$

input `int(1/(3*sin(c + d*x) + 5),x)`

output `atan((5*tan(c/2 + (d*x)/2))/4 + 3/4)/(2*d) - (atan(tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{1}{5 + 3 \sin(c + dx)} dx = \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}}{4}\right)}{2d}$$

input `int(1/(5+3*sin(d*x+c)),x)`

output `atan((5*tan((c + d*x)/2) + 3)/4)/(2*d)`

3.45 $\int \frac{1}{(5+3 \sin(c+dx))^2} dx$

Optimal result	314
Mathematica [A] (verified)	314
Rubi [A] (verified)	315
Maple [A] (verified)	316
Fricas [A] (verification not implemented)	317
Sympy [C] (verification not implemented)	318
Maxima [A] (verification not implemented)	318
Giac [A] (verification not implemented)	319
Mupad [B] (verification not implemented)	319
Reduce [B] (verification not implemented)	320

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx = \frac{5x}{64} + \frac{5 \arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{32d} + \frac{3 \cos(c + dx)}{16d(5 + 3 \sin(c + dx))}$$

output

```
5/64*x+5/32*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d+3/16*cos(d*x+c)/d/(5+3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.62

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx = \frac{25 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right) + \frac{6(-5+5 \cos(c+dx)-3 \sin(c+dx))}{5+3 \sin(c+dx)}}{160d}$$

input

```
Integrate[(5 + 3*Sin[c + d*x])^(-2),x]
```

output

```
(25*ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])] + (6*(-5 + 5*Cos[c + d*x] - 3*Sin[c + d*x]))/(5 + 3*Sin[c + d*x]))/(160*d)
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3 \sin(c + dx) + 5)^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(3 \sin(c + dx) + 5)^2} dx$$

$$\downarrow \text{3143}$$

$$\frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} - \frac{1}{16} \int -\frac{5}{3 \sin(c + dx) + 5} dx$$

$$\downarrow \text{27}$$

$$\frac{5}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)}$$

$$\downarrow \text{3042}$$

$$\frac{5}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)}$$

$$\downarrow \text{3136}$$

$$\frac{5}{16} \left(\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4} \right) + \frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)}$$

input

```
Int[(5 + 3*Sin[c + d*x])^(-2),x]
```

output $(5*(x/4 + \text{ArcTan}[\text{Cos}[c + d*x]/(3 + \text{Sin}[c + d*x])]/(2*d)))/16 + (3*\text{Cos}[c + d*x])/(16*d*(5 + 3*\text{Sin}[c + d*x]))$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_) \text{ ; FreeQ}[b, x]]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3136 $\text{Int}[(a_) + (b_)*\text{sin}[(c_) + (d_)*(x_)]^{(-1)}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2 - b^2, 2]\}, \text{Simp}[x/q, x] + \text{Simp}[(2/(d*q))*\text{ArcTan}[b*(\text{Cos}[c + d*x]/(a + q + b*\text{Sin}[c + d*x]))], x]] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{GtQ}[a^2 - b^2, 0] \ \&\& \ \text{PosQ}[a]$

rule 3143 $\text{Int}[(a_) + (b_)*\text{sin}[(c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^{(n + 1)}/(d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\text{Sin}[c + d*x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
default	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
risch	$\frac{5 e^{i(dx+c)} + 3i}{8d(3 e^{2i(dx+c)} - 3 + 10i e^{i(dx+c)})} - \frac{5i \ln(e^{i(dx+c)} + \frac{i}{3})}{64d} + \frac{5i \ln(e^{i(dx+c)} + 3i)}{64d}$
parallelrisch	$\frac{(-125i - 75i \sin(dx+c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) + (75i \sin(dx+c) + 125i) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 + 4i\right) + 60 \cos(dx+c) + 3}{960 \sin(dx+c)d + 1600d}$

input `int(1/(5+3*sin(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(2*(9/400*tan(1/2*d*x+1/2*c)+3/80)/(tan(1/2*d*x+1/2*c)^2+6/5*tan(1/2*d*x+1/2*c)+1)+5/32*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4))`

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx$$

$$= \frac{5(3 \sin(dx + c) + 5) \arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right) + 12 \cos(dx + c)}{64(3d \sin(dx + c) + 5d)}$$

input `integrate(1/(5+3*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/64*(5*(3*sin(d*x + c) + 5)*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c)) + 12*cos(d*x + c))/(3*d*sin(d*x + c) + 5*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.91 (sec) , antiderivative size = 388, normalized size of antiderivative = 6.93

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(5+3*sin(d*x+c))**2,x)`

output

```
Piecewise((x/(5 - 3*sin(2*atan(3/5 - 4*I/5)))**2, Eq(c, -d*x - 2*atan(3/5 - 4*I/5))), (x/(5 - 3*sin(2*atan(3/5 + 4*I/5)))**2, Eq(c, -d*x - 2*atan(3/5 + 4*I/5))), (x/(3*sin(c) + 5)**2, Eq(d, 0)), (125*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 150*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 125*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 36*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 60/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.66

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx = \frac{12 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 5 \right)}{\frac{6 \sin(dx+c)}{\cos(dx+c)+1} + \frac{5 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 5} + 25 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4} \right) \frac{1}{160 d}$$

input `integrate(1/(5+3*sin(d*x+c))^2,x, algorithm="maxima")`

output

```
1/160*(12*(3*sin(d*x + c)/(cos(d*x + c) + 1) + 5)/(6*sin(d*x + c)/(cos(d*x + c) + 1) + 5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 5) + 25*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.70

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx$$

$$= \frac{25 dx + 25 c + \frac{24 (3 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5)}{5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 6 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5} + 50 \arctan\left(-\frac{3 \cos(dx+c) + \sin(dx+c) + 3}{\cos(dx+c) - 3 \sin(dx+c) - 9}\right)}{320 d}$$

input `integrate(1/(5+3*sin(d*x+c))^2,x, algorithm="giac")`

output

```
1/320*(25*d*x + 25*c + 24*(3*tan(1/2*d*x + 1/2*c) + 5)/(5*tan(1/2*d*x + 1/2*c)^2 + 6*tan(1/2*d*x + 1/2*c) + 5) + 50*arctan(-(3*cos(d*x + c) + sin(d*x + c) + 3)/(cos(d*x + c) - 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 25.95 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.48

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d}$$

$$+ \frac{\frac{9 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{200} + \frac{3}{40}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1\right)}$$

input `int(1/(3*sin(c + d*x) + 5)^2,x)`

output

```
(5*atan((5*tan(c/2 + (d*x)/2))/4 + 3/4))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) + ((9*tan(c/2 + (d*x)/2))/200 + 3/40)/(d*((6*tan(c/2 + (d*x)/2))/5 + tan(c/2 + (d*x)/2)^2 + 1))
```


Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.14

$$\int \frac{1}{(5 + 3 \sin(c + dx))^2} dx$$

$$= \frac{15 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) + 3}{4}\right) \sin(dx+c) + 25 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) + 3}{4}\right) + 6 \cos(dx+c)}{32d(3 \sin(dx+c) + 5)}$$

input

```
int(1/(5+3*sin(d*x+c))^2,x)
```

output

```
(15*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x) + 25*atan((5*tan((c + d*x)/2) + 3)/4) + 6*cos(c + d*x))/(32*d*(3*sin(c + d*x) + 5))
```

3.46 $\int \frac{1}{(5+3 \sin(c+dx))^3} dx$

Optimal result	321
Mathematica [A] (verified)	321
Rubi [A] (verified)	322
Maple [A] (verified)	324
Fricas [A] (verification not implemented)	325
Sympy [C] (verification not implemented)	325
Maxima [B] (verification not implemented)	326
Giac [A] (verification not implemented)	327
Mupad [B] (verification not implemented)	327
Reduce [B] (verification not implemented)	328

Optimal result

Integrand size = 12, antiderivative size = 81

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx = \frac{59x}{2048} + \frac{59 \arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{1024d} + \frac{3 \cos(c + dx)}{32d(5 + 3 \sin(c + dx))^2} + \frac{45 \cos(c + dx)}{512d(5 + 3 \sin(c + dx))}$$

output

```
59/2048*x+59/1024*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d+3/32*cos(d*x+c)/d/(5+3*sin(d*x+c))^2+45/512*cos(d*x+c)/d/(5+3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.40

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx = \frac{59 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right) + \frac{546 \cos(c+dx)+9(-59+9 \cos(2(c+dx))-60 \sin(c+dx)+15 \sin(2(c+dx)))}{(5+3 \sin(c+dx))^2}}{1024d}$$

input

```
Integrate[(5 + 3*Sin[c + d*x])^(-3), x]
```

output

$$(59*\text{ArcTan}[(2*(\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]))/(\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2])] + (546*\text{Cos}[c + d*x] + 9*(-59 + 9*\text{Cos}[2*(c + d*x)] - 60*\text{Sin}[c + d*x] + 15*\text{Sin}[2*(c + d*x)]))/((5 + 3*\text{Sin}[c + d*x])^2)/(1024*d)$$
Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(3 \sin(c + dx) + 5)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(3 \sin(c + dx) + 5)^3} dx \\ & \quad \downarrow \text{3143} \\ & \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} - \frac{1}{32} \int -\frac{10 - 3 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx \\ & \quad \downarrow \text{25} \\ & \frac{1}{32} \int \frac{10 - 3 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx + \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{32} \int \frac{10 - 3 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx + \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{3233} \\ & \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} - \frac{1}{16} \int -\frac{59}{3 \sin(c + dx) + 5} dx \right) + \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{27} \\ & \frac{1}{32} \left(\frac{59}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{45 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \end{aligned}$$

$$\begin{array}{c}
 \downarrow 3042 \\
 \frac{1}{32} \left(\frac{59}{16} \int \frac{1}{3 \sin(c+dx)+5} dx + \frac{45 \cos(c+dx)}{16d(3 \sin(c+dx)+5)} \right) + \frac{3 \cos(c+dx)}{32d(3 \sin(c+dx)+5)^2} \\
 \downarrow 3136 \\
 \frac{1}{32} \left(\frac{59}{16} \left(\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4} \right) + \frac{45 \cos(c+dx)}{16d(3 \sin(c+dx)+5)} \right) + \frac{3 \cos(c+dx)}{32d(3 \sin(c+dx)+5)^2}
 \end{array}$$

input `Int[(5 + 3*Sin[c + d*x])^(-3),x]`

output `(3*Cos[c + d*x])/(32*d*(5 + 3*Sin[c + d*x])^2) + ((59*(x/4 + ArcTan[Cos[c + d*x]/(3 + Sin[c + d*x])]/(2*d)))/16 + (45*Cos[c + d*x])/(16*d*(5 + 3*Sin[c + d*x]))) / 32`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

rule 3233

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12

method	result
derivativdivides	$\frac{\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{1280} + \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1280} + \frac{273}{256} + \frac{59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right)}{1024}}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5} d$
default	$\frac{\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{1280} + \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1280} + \frac{273}{256} + \frac{59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right)}{1024}}{d}$
risch	$\frac{\frac{885ie^{2i(dx+c)}}{256} + \frac{177e^{3i(dx+c)}}{256} - \frac{723e^{i(dx+c)}}{256} - \frac{135i}{256}}{\left(3e^{2i(dx+c)} - 3 + 10ie^{i(dx+c)}\right)^2} d - \frac{59i \ln\left(e^{i(dx+c)} + \frac{i}{3}\right)}{2048d} + \frac{59i \ln\left(e^{i(dx+c)} + 3i\right)}{2048d}$
parallelrisc	$\frac{1062 - 59i(59 - 9 \cos(2dx + 2c) + 60 \sin(dx + c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) - 59i(-59 + 9 \cos(2dx + 2c) - 60 \sin(dx + c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 + 4i\right)}{2048(59 - 9 \cos(2dx + 2c) + 60 \sin(dx + c))}$

input

```
int(1/(5+3*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(50*(963/64000*tan(1/2*d*x+1/2*c)^3+11739/320000*tan(1/2*d*x+1/2*c)^2+
2313/64000*tan(1/2*d*x+1/2*c)+273/12800)/(5*tan(1/2*d*x+1/2*c)^2+6*tan(1/2
*d*x+1/2*c)+5)^2+59/1024*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.16

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx$$

$$= \frac{59 (9 \cos(dx + c)^2 - 30 \sin(dx + c) - 34) \arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right) - 540 \cos(dx + c) \sin(dx + c) - 1092}{2048 (9 d \cos(dx + c)^2 - 30 d \sin(dx + c) - 34 d)}$$

input `integrate(1/(5+3*sin(d*x+c))^3,x, algorithm="fricas")`

output `1/2048*(59*(9*cos(d*x + c)^2 - 30*sin(d*x + c) - 34)*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c)) - 540*cos(d*x + c)*sin(d*x + c) - 1092*cos(d*x + c))/(9*d*cos(d*x + c)^2 - 30*d*sin(d*x + c) - 34*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.97 (sec) , antiderivative size = 918, normalized size of antiderivative = 11.33

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx = \text{Too large to display}$$

input `integrate(1/(5+3*sin(d*x+c))**3,x)`

output

```
Piecewise((x/(5 - 3*sin(2*atan(3/5 - 4*I/5)))**3, Eq(c, -d*x - 2*atan(3/5 - 4*I/5))), (x/(5 - 3*sin(2*atan(3/5 + 4*I/5)))**3, Eq(c, -d*x - 2*atan(3/5 + 4*I/5))), (x/(3*sin(c) + 5)**3, Eq(d, 0)), (36875*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 88500*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 126850*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 88500*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 36875*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 19260*tan(c/2 + d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*t...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(73) = 146$.

Time = 0.13 (sec) , antiderivative size = 173, normalized size of antiderivative = 2.14

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx$$

$$= \frac{12 \left(\frac{3855 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3913 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1605 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + 2275 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} + \frac{86 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{25 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 25} + 1475 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4} \right)$$

$$25600 d$$

input

```
integrate(1/(5+3*sin(d*x+c))^3,x, algorithm="maxima")
```

output

```
1/25600*(12*(3855*sin(d*x + c)/(cos(d*x + c) + 1) + 3913*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1605*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 2275)/(60*sin(d*x + c)/(cos(d*x + c) + 1) + 86*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 25*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 25) + 1475*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.49

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx$$

$$= \frac{1475 dx + 1475 c + \frac{24 \left(1605 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 3913 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 3855 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2275 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^2} + 2950 \arctan\left(-\frac{3 \cos(dx + c) + \sin(dx + c) + 3}{\cos(dx + c) - 3 \sin(dx + c) - 9}\right)}{51200 d}$$

input

```
integrate(1/(5+3*sin(d*x+c))^3,x, algorithm="giac")
```

output

```
1/51200*(1475*d*x + 1475*c + 24*(1605*tan(1/2*d*x + 1/2*c)^3 + 3913*tan(1/2*d*x + 1/2*c)^2 + 3855*tan(1/2*d*x + 1/2*c) + 2275)/(5*tan(1/2*d*x + 1/2*c)^2 + 6*tan(1/2*d*x + 1/2*c) + 5)^2 + 2950*arctan(-(3*cos(d*x + c) + sin(d*x + c) + 3)/(cos(d*x + c) - 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 26.00 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.37

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx = \frac{59 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{1024 d} - \frac{59 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{1024 d}$$

$$+ \frac{\frac{963 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{1280} + \frac{11739 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{1280} + \frac{273}{256}}{d \left(5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 5 \right)^2}$$

input

```
int(1/(3*sin(c + d*x) + 5)^3,x)
```


output

```
(59*atan((5*tan(c/2 + (d*x)/2))/4 + 3/4))/(1024*d) - (59*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) + ((2313*tan(c/2 + (d*x)/2))/1280 + (11739*atan(c/2 + (d*x)/2)^2)/6400 + (963*tan(c/2 + (d*x)/2)^3)/1280 + 273/256)/(d*(6*tan(c/2 + (d*x)/2) + 5*tan(c/2 + (d*x)/2)^2 + 5)^2)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.62

$$\int \frac{1}{(5 + 3 \sin(c + dx))^3} dx$$

$$= \frac{531 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c)^2 + 1770 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c) + 1475 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4}\right)}{1024d (9 \sin(dx + c)^2 - 1)}$$

input

```
int(1/(5+3*sin(d*x+c))^3,x)
```

output

```
(531*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x)**2 + 1770*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x) + 1475*atan((5*tan((c + d*x)/2) + 3)/4) + 270*cos(c + d*x)*sin(c + d*x) + 546*cos(c + d*x) + 81*sin(c + d*x)**2 + 270*sin(c + d*x) + 225)/(1024*d*(9*sin(c + d*x)**2 + 30*sin(c + d*x) + 25))
```

3.47 $\int \frac{1}{(5+3 \sin(c+dx))^4} dx$

Optimal result	329
Mathematica [A] (verified)	329
Rubi [A] (verified)	330
Maple [A] (verified)	333
Fricas [A] (verification not implemented)	333
Sympy [C] (verification not implemented)	334
Maxima [B] (verification not implemented)	335
Giac [A] (verification not implemented)	335
Mupad [B] (verification not implemented)	336
Reduce [B] (verification not implemented)	337

Optimal result

Integrand size = 12, antiderivative size = 106

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx = \frac{385x}{32768} + \frac{385 \arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{16384d} + \frac{\cos(c + dx)}{16d(5 + 3 \sin(c + dx))^3} + \frac{25 \cos(c + dx)}{512d(5 + 3 \sin(c + dx))^2} + \frac{311 \cos(c + dx)}{8192d(5 + 3 \sin(c + dx))}$$

output `385/32768*x+385/16384*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d+1/16*cos(d*x+c)/d/(5+3*sin(d*x+c))^3+25/512*cos(d*x+c)/d/(5+3*sin(d*x+c))^2+311/8192*cos(d*x+c)/d/(5+3*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.25

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx = \frac{1925 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right) + \frac{-239470+219735 \cos(c+dx)+83970 \cos(2(c+dx))-13995 \cos(3(c+dx))-305091 \sin(c+dx)}{2(5+3 \sin(c+dx))^3}}{81920d}$$

input `Integrate[(5 + 3*Sin[c + d*x])^(-4), x]`

output

```
(1925*ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] -
Sin[(c + d*x)/2])] + (-239470 + 219735*Cos[c + d*x] + 83970*Cos[2*(c + d*
x)] - 13995*Cos[3*(c + d*x)] - 305091*Sin[c + d*x] + 105300*Sin[2*(c + d*x
)] + 8397*Sin[3*(c + d*x)])/(2*(5 + 3*Sin[c + d*x])^3)/(81920*d)
```

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3 \sin(c + dx) + 5)^4} dx$$

↓ 3042

$$\int \frac{1}{(3 \sin(c + dx) + 5)^4} dx$$

↓ 3143

$$\frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} - \frac{1}{48} \int -\frac{3(5 - 2 \sin(c + dx))}{(3 \sin(c + dx) + 5)^3} dx$$

↓ 27

$$\frac{1}{16} \int \frac{5 - 2 \sin(c + dx)}{(3 \sin(c + dx) + 5)^3} dx + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

↓ 3042

$$\frac{1}{16} \int \frac{5 - 2 \sin(c + dx)}{(3 \sin(c + dx) + 5)^3} dx + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

↓ 3233

$$\frac{1}{16} \left(\frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} - \frac{1}{32} \int -\frac{62 - 25 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx \right) + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

↓ 25

$$\frac{1}{16} \left(\frac{1}{32} \int \frac{62 - 25 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \int \frac{62 - 25 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 3233 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} - \frac{1}{16} \int -\frac{385}{3 \sin(c + dx) + 5} dx \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 27 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 3136 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \left(\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4} \right) + \frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}
\end{aligned}$$

input `Int[(5 + 3*Sin[c + d*x])^(-4),x]`

output `Cos[c + d*x]/(16*d*(5 + 3*Sin[c + d*x])^3) + ((25*Cos[c + d*x])/(32*d*(5 + 3*Sin[c + d*x])^2) + ((385*(x/4 + ArcTan[Cos[c + d*x]/(3 + Sin[c + d*x]))/(2*d)))/16 + (311*Cos[c + d*x])/(16*d*(5 + 3*Sin[c + d*x]))/32)/16`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \text{ Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3136 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(-1)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = \text{Rt}[\text{a}^2 - \text{b}^2, 2]\}, \text{Simp}[\text{x}/\text{q}, \text{x}] + \text{Simp}[(2/(d*q))*\text{ArcTan}[\text{b}*(\text{Cos}[\text{c} + \text{d}*x]/(\text{a} + \text{q} + \text{b}*\text{Sin}[\text{c} + \text{d}*x]))], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{PosQ}[\text{a}]$
- rule 3143 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(n_)}, \text{x_Symbol}] \rightarrow \text{Simp}[-\text{b})*\text{Cos}[\text{c} + \text{d}*x]*((\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(n+1)}/(\text{d}*(n+1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((n+1)*(a^2 - b^2)) \text{ Int}[(\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(n+1)}*\text{Simp}[\text{a}*(n+1) - \text{b}*(n+2)*\text{Sin}[\text{c} + \text{d}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{n}, -1] \ \&\& \ \text{IntegerQ}[2*\text{n}]$
- rule 3233 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]^{(m_)}*((\text{c}_.) + (\text{d}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[-(\text{b}*c - \text{a}*d)*\text{Cos}[\text{e} + \text{f}*x]*((\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(m+1)}/(\text{f}*(m+1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((m+1)*(a^2 - b^2)) \text{ Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(m+1)}*\text{Simp}[(\text{a}*c - \text{b}*d)*(m+1) - (\text{b}*c - \text{a}*d)*(m+2)*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntegerQ}[2*\text{m}]$

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} + \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} + \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} + \frac{10287}{4096} + \frac{385}{16384} \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4}\right)}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3} dx$
default	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} + \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} + \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} + \frac{10287}{4096} + \frac{385}{16384} \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4}\right)}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3} dx$
risch	$\frac{-239470 e^{3i(dx+c)} + 86625 i e^{4i(dx+c)} - 218466 i e^{2i(dx+c)} + 10395 e^{5i(dx+c)} + 73575 e^{i(dx+c)} + 8397 i}{12288 (3 e^{2i(dx+c)} - 3 + 10 i e^{i(dx+c)})^3} dx - \frac{385 i \ln(e^{i(dx+c)})}{32768 d}$
parallelrisch	$-31683960 + 48125i(-27 \sin(3dx+3c) + 981 \sin(dx+c) - 270 \cos(2dx+2c) + 770) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) + 48125i(-7$

input `int(1/(5+3*sin(d*x+c))^4,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{250 \cdot (39933/5120000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c))^5 + 672723/25600000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^4 + 2870073/64000000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 + 604899/12800000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 145233/5120000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + 10287/1024000}{(5 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 6 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + 5)^3} + \frac{385}{16384} \cdot \arctan\left(\frac{5}{4} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + \frac{3}{4}\right) \right)$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.23

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{11196 \cos(dx + c)^3 + 385 (135 \cos(dx + c)^2 + 9 (3 \cos(dx + c)^2 - 28) \sin(dx + c) - 260) \arctan\left(\frac{5 \sin(dx + c)}{4 \cos(dx + c)}\right)}{32768 (135 d \cos(dx + c)^2 + 9 (3 d \cos(dx + c)^2 - 28 d) \sin(dx + c) - 260 d)}$$

input `integrate(1/(5+3*sin(d*x+c))^4,x, algorithm="fricas")`

output

```
1/32768*(11196*cos(d*x + c)^3 + 385*(135*cos(d*x + c)^2 + 9*(3*cos(d*x + c)
)^2 - 28)*sin(d*x + c) - 260)*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c)
) - 42120*cos(d*x + c)*sin(d*x + c) - 52344*cos(d*x + c))/(135*d*cos(d*x +
c)^2 + 9*(3*d*cos(d*x + c)^2 - 28*d)*sin(d*x + c) - 260*d)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.55 (sec) , antiderivative size = 1693, normalized size of antiderivative = 15.97

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(5+3*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(5 - 3*sin(2*atan(3/5 - 4*I/5)))**4, Eq(c, -d*x - 2*atan(3/5
- 4*I/5))), (x/(5 - 3*sin(2*atan(3/5 + 4*I/5)))**4, Eq(c, -d*x - 2*atan(3/
5 + 4*I/5))), (x/(3*sin(c) + 5)**4, Eq(d, 0)), (6015625*(atan(5*tan(c/2 +
d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**6/(
256000000*d*tan(c/2 + d*x/2)**6 + 921600000*d*tan(c/2 + d*x/2)**5 + 187392
0000*d*tan(c/2 + d*x/2)**4 + 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000
*d*tan(c/2 + d*x/2)**2 + 921600000*d*tan(c/2 + d*x/2) + 256000000*d) + 216
56250*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi
))*tan(c/2 + d*x/2)**5/(256000000*d*tan(c/2 + d*x/2)**6 + 921600000*d*tan(
c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)**4 + 2285568000*d*tan(c/2
+ d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 + 921600000*d*tan(c/2 + d*x
/2) + 256000000*d) + 44034375*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor
((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(256000000*d*tan(c/2 + d*x/
2)**6 + 921600000*d*tan(c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)**4
+ 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 + 9
21600000*d*tan(c/2 + d*x/2) + 256000000*d) + 53707500*(atan(5*tan(c/2 + d*
x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**3/(25
6000000*d*tan(c/2 + d*x/2)**6 + 921600000*d*tan(c/2 + d*x/2)**5 + 18739200
00*d*tan(c/2 + d*x/2)**4 + 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000*d
*tan(c/2 + d*x/2)**2 + 921600000*d*tan(c/2 + d*x/2) + 256000000*d) + 44...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. $2(96) = 192$.

Time = 0.12 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.39

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{36 \left(\frac{403425 \sin(dx+c)}{\cos(dx+c)+1} + \frac{672110 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{637794 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{373735 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{110925 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + 142875 \right) + 48125 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} \right)}{2048000 d}$$

input `integrate(1/(5+3*sin(d*x+c))^4,x, algorithm="maxima")`

output

$$\frac{1}{2048000} \cdot \left(36 \cdot \left(\frac{403425 \sin(dx+c)}{\cos(dx+c)+1} + \frac{672110 \sin^2(dx+c)}{(\cos(dx+c)+1)^2} + \frac{637794 \sin^3(dx+c)}{(\cos(dx+c)+1)^3} + \frac{373735 \sin^4(dx+c)}{(\cos(dx+c)+1)^4} + \frac{110925 \sin^5(dx+c)}{(\cos(dx+c)+1)^5} + 142875 \right) + 48125 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} \right) \right) / d$$
Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.39

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{48125 dx + 48125 c + \frac{72 \left(110925 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 373735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 637794 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 672110 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 403425 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 142875 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^3}{4096000 d}$$

input `integrate(1/(5+3*sin(d*x+c))^4,x, algorithm="giac")`

output

```
1/4096000*(48125*d*x + 48125*c + 72*(110925*tan(1/2*d*x + 1/2*c)^5 + 37373
5*tan(1/2*d*x + 1/2*c)^4 + 637794*tan(1/2*d*x + 1/2*c)^3 + 672110*tan(1/2*
d*x + 1/2*c)^2 + 403425*tan(1/2*d*x + 1/2*c) + 142875)/(5*tan(1/2*d*x + 1/
2*c)^2 + 6*tan(1/2*d*x + 1/2*c) + 5)^3 + 96250*arctan(-(3*cos(d*x + c) + s
in(d*x + c) + 3)/(cos(d*x + c) - 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 26.52 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.76

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{385 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{16384 d} - \frac{385 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{16384 d}$$

$$+ \frac{\frac{39933 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{2560000} + \frac{672723 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{12800000} + \frac{2870073 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{32000000} + \frac{604899 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400000} + \frac{145233 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2560000} + \frac{10287}{512000}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{5} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{25} + \frac{1116 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{125} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{25} + \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1 \right)}$$

input

```
int(1/(3*sin(c + d*x) + 5)^4,x)
```

output

```
(385*atan((5*tan(c/2 + (d*x)/2))/4 + 3/4))/(16384*d) - (385*(atan(tan(c/2
+ (d*x)/2)) - (d*x)/2))/(16384*d) + ((145233*tan(c/2 + (d*x)/2))/2560000 +
(604899*tan(c/2 + (d*x)/2)^2)/6400000 + (2870073*tan(c/2 + (d*x)/2)^3)/32
000000 + (672723*tan(c/2 + (d*x)/2)^4)/12800000 + (39933*tan(c/2 + (d*x)/2
)^5)/2560000 + 10287/512000)/(d*((18*tan(c/2 + (d*x)/2))/5 + (183*tan(c/2
+ (d*x)/2)^2)/25 + (1116*tan(c/2 + (d*x)/2)^3)/125 + (183*tan(c/2 + (d*x)/
2)^4)/25 + (18*tan(c/2 + (d*x)/2)^5)/5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.80

$$\int \frac{1}{(5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{51975 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c)^3 + 259875 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c)^2 + 433125 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c) + 240625 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) + 27990 \cos(c + dx) \sin(c + dx)^2 + 105300 \cos(c + dx) \sin(c + dx) + 102870 \cos(c + dx) + 12636 \sin(c + dx)^3 + 63180 \sin(c + dx)^2 + 105300 \sin(c + dx) + 58500}{(81920 dx (27 \sin(c + dx)^3 + 135 \sin(c + dx)^2 + 225 \sin(c + dx) + 125))}$$

input `int(1/(5+3*sin(d*x+c))^4,x)`output `(51975*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x)**3 + 259875*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x)**2 + 433125*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x) + 240625*atan((5*tan((c + d*x)/2) + 3)/4) + 27990*cos(c + d*x)*sin(c + d*x)**2 + 105300*cos(c + d*x)*sin(c + d*x) + 102870*cos(c + d*x) + 12636*sin(c + d*x)**3 + 63180*sin(c + d*x)**2 + 105300*sin(c + d*x) + 58500)/(81920*d*(27*sin(c + d*x)**3 + 135*sin(c + d*x)**2 + 225*sin(c + d*x) + 125))`

3.48 $\int \frac{1}{5-3\sin(c+dx)} dx$

Optimal result	338
Mathematica [A] (verified)	338
Rubi [A] (verified)	339
Maple [A] (verified)	340
Fricas [A] (verification not implemented)	340
Sympy [B] (verification not implemented)	341
Maxima [A] (verification not implemented)	341
Giac [A] (verification not implemented)	341
Mupad [B] (verification not implemented)	342
Reduce [B] (verification not implemented)	342

Optimal result

Integrand size = 12, antiderivative size = 33

$$\int \frac{1}{5-3\sin(c+dx)} dx = \frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d}$$

output `1/4*x-1/2*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.70

$$\int \frac{1}{5-3\sin(c+dx)} dx = -\frac{\arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right)}{2d}$$

input `Integrate[(5 - 3*Sin[c + d*x])^(-1),x]`

output `-1/2*ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])]/d`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx$$

↓ 3042

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx$$

↓ 3136

$$\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d}$$

input `Int[(5 - 3*Sin[c + d*x])^(-1),x]`

output `x/4 - ArcTan[Cos[c + d*x]/(3 - Sin[c + d*x])]/(2*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.61

method	result	size
derivativedivides	$\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}}{4}\right)}{2d}$	20
default	$\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}}{4}\right)}{2d}$	20
risch	$\frac{i \ln(e^{i(dx+c)} - 3i)}{4d} - \frac{i \ln(e^{i(dx+c)} - \frac{i}{3})}{4d}$	40
parallelrisch	$-\frac{i \left(\ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) - \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 + 4i\right) \right)}{4d}$	42

input `int(1/(5-3*sin(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2/d*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx = \frac{\arctan\left(\frac{5 \sin(dx+c) - 3}{4 \cos(dx+c)}\right)}{4d}$$

input `integrate(1/(5-3*sin(d*x+c)),x, algorithm="fricas")`

output `1/4*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(22) = 44$.

Time = 0.39 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx = \begin{cases} \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c+dx}{2}\right) - \frac{3}{4}}{4}\right) + \pi \left\lfloor \frac{\frac{c+dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{5 - 3 \sin(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/(5-3*sin(d*x+c)),x)`

output `Piecewise(((atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(2*d), Ne(d, 0)), (x/(5 - 3*sin(c)), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx = \frac{\arctan\left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} - \frac{3}{4}\right)}{2d}$$

input `integrate(1/(5-3*sin(d*x+c)),x, algorithm="maxima")`

output `1/2*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4)/d`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.52

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx = \frac{dx + c + 2 \arctan\left(\frac{3 \cos(dx+c) - \sin(dx+c) + 3}{\cos(dx+c) + 3 \sin(dx+c) - 9}\right)}{4d}$$

input `integrate(1/(5-3*sin(d*x+c)),x, algorithm="giac")`

output

$$\frac{1}{4} \cdot (d \cdot x + c + 2 \cdot \arctan((3 \cdot \cos(d \cdot x + c) - \sin(d \cdot x + c) + 3) / (\cos(d \cdot x + c) + 3 \cdot \sin(d \cdot x + c) - 9))) / d$$
Mupad [B] (verification not implemented)

Time = 25.85 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx = \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{3}{4}}{4}\right)}{2d} - \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d}$$

input

$$\operatorname{int}(-1/(3 \cdot \sin(c + d \cdot x) - 5), x)$$

output

$$\operatorname{atan}\left(\frac{5 \cdot \tan(c/2 + (d \cdot x)/2)}{4} - \frac{3}{4}\right) / (2 \cdot d) - \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{d \cdot x}{2}\right)\right) - \frac{d \cdot x}{2}\right) / (2 \cdot d)$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.58

$$\int \frac{1}{5 - 3 \sin(c + dx)} dx = \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}}{4}\right)}{2d}$$

input

$$\operatorname{int}(1/(5 - 3 \cdot \sin(d \cdot x + c)), x)$$

output

$$\operatorname{atan}\left(\frac{5 \cdot \tan((c + d \cdot x)/2) - 3}{4}\right) / (2 \cdot d)$$

3.49 $\int \frac{1}{(5-3\sin(c+dx))^2} dx$

Optimal result	343
Mathematica [A] (verified)	343
Rubi [A] (verified)	344
Maple [A] (verified)	345
Fricas [A] (verification not implemented)	346
Sympy [C] (verification not implemented)	347
Maxima [A] (verification not implemented)	347
Giac [A] (verification not implemented)	348
Mupad [B] (verification not implemented)	348
Reduce [B] (verification not implemented)	349

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{1}{(5-3\sin(c+dx))^2} dx = \frac{5x}{64} - \frac{5 \arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{32d} - \frac{3 \cos(c+dx)}{16d(5-3\sin(c+dx))}$$

output

5/64*x-5/32*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d-3/16*cos(d*x+c)/d/(5-3*sin(d*x+c))

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.57

$$\int \frac{1}{(5-3\sin(c+dx))^2} dx = \frac{-25 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right) + \frac{6(-5+5\cos(c+dx)+3\sin(c+dx))}{-5+3\sin(c+dx)}}{160d}$$

input

Integrate[(5 - 3*Sin[c + d*x])^(-2), x]

output

```
(-25*ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])] + (6*(-5 + 5*Cos[c + d*x] + 3*Sin[c + d*x]))/(-5 + 3*Sin[c + d*x]))/(160*d)
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 - 3 \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 - 3 \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{16} \int -\frac{5}{5 - 3 \sin(c + dx)} dx - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \\
 & \quad \downarrow \text{3136} \\
 & \frac{5}{16} \left(\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d} \right) - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))}
 \end{aligned}$$

input

```
Int[(5 - 3*Sin[c + d*x])^(-2),x]
```

output $(5*(x/4 - \text{ArcTan}[\text{Cos}[c + d*x]/(3 - \text{Sin}[c + d*x])]/(2*d)))/16 - (3*\text{Cos}[c + d*x])/(16*d*(5 - 3*\text{Sin}[c + d*x]))$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3136 $\text{Int}[(a_ + (b_)*\text{sin}[(c_.) + (d_)*(x_)])^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2 - b^2, 2]\}, \text{Simp}[x/q, x] + \text{Simp}[(2/(d*q))*\text{ArcTan}[b*(\text{Cos}[c + d*x]/(a + q + b*\text{Sin}[c + d*x]))], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{GtQ}[a^2 - b^2, 0] \ \&\& \ \text{PosQ}[a]$

rule 3143 $\text{Int}[(a_ + (b_)*\text{sin}[(c_.) + (d_)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^{(n + 1)}/(d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\text{Sin}[c + d*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
default	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
risch	$-\frac{5 e^{i(dx+c)} - 3i}{8d(3 e^{2i(dx+c)} - 3 - 10i e^{i(dx+c)})} + \frac{5i \ln(e^{i(dx+c)} - 3i)}{64d} - \frac{5i \ln(e^{i(dx+c)} - \frac{i}{3})}{64d}$
parallelrisch	$\frac{(125i - 75i \sin(dx+c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + (75i \sin(dx+c) - 125i) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 + 4i\right) + 60 \cos(dx+c) - 36}{960 \sin(dx+c)d - 1600d}$

input `int(1/(5-3*sin(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(2*(9/400*tan(1/2*d*x+1/2*c)-3/80)/(tan(1/2*d*x+1/2*c)^2-6/5*tan(1/2*d*x+1/2*c)+1)+5/32*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02

$$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx$$

$$= \frac{5(3 \sin(dx + c) - 5) \arctan\left(\frac{5 \sin(dx+c) - 3}{4 \cos(dx+c)}\right) + 12 \cos(dx + c)}{64(3d \sin(dx + c) - 5d)}$$

input `integrate(1/(5-3*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/64*(5*(3*sin(d*x + c) - 5)*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c)) + 12*cos(d*x + c))/(3*d*sin(d*x + c) - 5*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.92 (sec) , antiderivative size = 384, normalized size of antiderivative = 6.62

$$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(5-3*sin(d*x+c))**2,x)`

output

```
Piecewise((x/(5 - 3*sin(2*atan(3/5 - 4*I/5)))**2, Eq(c, -d*x + 2*atan(3/5 - 4*I/5))), (x/(5 - 3*sin(2*atan(3/5 + 4*I/5)))**2, Eq(c, -d*x + 2*atan(3/5 + 4*I/5))), (x/(5 - 3*sin(c))**2, Eq(d, 0)), (125*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) - 150*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) + 125*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) + 36*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) - 60/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.60

$$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx = -\frac{12 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 5 \right)}{\frac{6 \sin(dx+c)}{\cos(dx+c)+1} - \frac{5 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 5} - 25 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} - \frac{3}{4} \right) \frac{1}{160 d}$$

input `integrate(1/(5-3*sin(d*x+c))^2,x, algorithm="maxima")`

output

```
-1/160*(12*(3*sin(d*x + c)/(cos(d*x + c) + 1) - 5)/(6*sin(d*x + c)/(cos(d*x + c) + 1) - 5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 5) - 25*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx$$

$$= \frac{25 dx + 25 c + \frac{24 (3 \tan(\frac{1}{2} dx + \frac{1}{2} c) - 5)}{5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 6 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5} + 50 \arctan\left(\frac{3 \cos(dx+c) - \sin(dx+c) + 3}{\cos(dx+c) + 3 \sin(dx+c) - 9}\right)}{320 d}$$

input `integrate(1/(5-3*sin(d*x+c))^2,x, algorithm="giac")`

output

```
1/320*(25*d*x + 25*c + 24*(3*tan(1/2*d*x + 1/2*c) - 5)/(5*tan(1/2*d*x + 1/2*c)^2 - 6*tan(1/2*d*x + 1/2*c) + 5) + 50*arctan((3*cos(d*x + c) - sin(d*x + c) + 3)/(cos(d*x + c) + 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 26.01 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.43

$$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 3}{4}\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d}$$

$$+ \frac{\frac{9 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{200} - \frac{3}{40}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1\right)}$$

input `int(1/(3*sin(c + d*x) - 5)^2,x)`

output

```
(5*atan((5*tan(c/2 + (d*x)/2))/4 - 3/4))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) + ((9*tan(c/2 + (d*x)/2))/200 - 3/40)/(d*(tan(c/2 + (d*x)/2)^2 - (6*tan(c/2 + (d*x)/2))/5 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

$$\int \frac{1}{(5 - 3 \sin(c + dx))^2} dx$$

$$= \frac{15 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) \sin(dx+c) - 25 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) + 6 \cos(dx+c)}{32d(3 \sin(dx+c) - 5)}$$

input

```
int(1/(5-3*sin(d*x+c))^2,x)
```

output

```
(15*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x) - 25*atan((5*tan((c + d*x)/2) - 3)/4) + 6*cos(c + d*x))/(32*d*(3*sin(c + d*x) - 5))
```

3.50 $\int \frac{1}{(5-3 \sin(c+dx))^3} dx$

Optimal result	350
Mathematica [A] (verified)	350
Rubi [A] (verified)	351
Maple [A] (verified)	353
Fricas [A] (verification not implemented)	354
Sympy [C] (verification not implemented)	354
Maxima [B] (verification not implemented)	355
Giac [A] (verification not implemented)	356
Mupad [B] (verification not implemented)	356
Reduce [B] (verification not implemented)	357

Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \frac{1}{(5-3 \sin(c+dx))^3} dx = \frac{59x}{2048} - \frac{59 \arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{1024d} - \frac{3 \cos(c+dx)}{32d(5-3 \sin(c+dx))^2} - \frac{45 \cos(c+dx)}{512d(5-3 \sin(c+dx))}$$

output `59/2048*x-59/1024*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d-3/32*cos(d*x+c)/d/(5-3*sin(d*x+c))^2-45/512*cos(d*x+c)/d/(5-3*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.36

$$\int \frac{1}{(5-3 \sin(c+dx))^3} dx = \frac{59 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right) + \frac{546 \cos(c+dx)+9(-59+9 \cos(2(c+dx))+60 \sin(c+dx))-15 \sin(2(c+dx))}{(5-3 \sin(c+dx))^2}}{1024d}$$

input `Integrate[(5 - 3*Sin[c + d*x])^(-3),x]`

output

```
-1/1024*(59*ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])] + (546*Cos[c + d*x] + 9*(-59 + 9*Cos[2*(c + d*x)] + 60*Sin[c + d*x] - 15*Sin[2*(c + d*x)]))/(5 - 3*Sin[c + d*x])^2)/d
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 - 3 \sin(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 - 3 \sin(c + dx))^3} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{32} \int -\frac{3 \sin(c + dx) + 10}{(5 - 3 \sin(c + dx))^2} dx - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{32} \int \frac{3 \sin(c + dx) + 10}{(5 - 3 \sin(c + dx))^2} dx - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \int \frac{3 \sin(c + dx) + 10}{(5 - 3 \sin(c + dx))^2} dx - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left(-\frac{1}{16} \int -\frac{59}{5 - 3 \sin(c + dx)} dx - \frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{32} \left(\frac{59}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{1}{32} \left(\frac{59}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \\ & \downarrow 3136 \\ & \frac{1}{32} \left(\frac{59}{16} \left(\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d} \right) - \frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \end{aligned}$$

input `Int[(5 - 3*Sin[c + d*x])^(-3),x]`

output `((59*(x/4 - ArcTan[Cos[c + d*x]/(3 - Sin[c + d*x]])/(2*d)))/16 - (45*Cos[c + d*x])/(16*d*(5 - 3*Sin[c + d*x]))/32 - (3*Cos[c + d*x])/(32*d*(5 - 3*Sin[c + d*x])^2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

rule 3233

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.10

method	result
derivativdivides	$\frac{\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{1280} - \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1280} - \frac{273}{256} + \frac{59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} - \frac{3}{4}\right)}{1024}}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5} d$
default	$\frac{\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{1280} - \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{1280} - \frac{273}{256} + \frac{59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} - \frac{3}{4}\right)}{1024}}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5} d$
risch	$-\frac{3(-295ie^{2i(dx+c)} + 59e^{3i(dx+c)} - 241e^{i(dx+c)} + 45i)}{256(3e^{2i(dx+c)} - 3 - 10ie^{i(dx+c)})^2 d} - \frac{59i \ln(e^{i(dx+c)} - \frac{i}{3})}{2048d} + \frac{59i \ln(e^{i(dx+c)} - 3i)}{2048d}$
parallelrisch	$\frac{1062 + 59i(59 - 9 \cos(2dx + 2c) - 60 \sin(dx + c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + 59i(9 \cos(2dx + 2c) - 59 + 60 \sin(dx + c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right)}{2048d(9 \cos(2dx + 2c) - 59 + 60 \sin(dx + c))}$

input

```
int(1/(5-3*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(50*(963/64000*tan(1/2*d*x+1/2*c)^3-11739/320000*tan(1/2*d*x+1/2*c)^2+
2313/64000*tan(1/2*d*x+1/2*c)-273/12800)/(5*tan(1/2*d*x+1/2*c)^2-6*tan(1/2
*d*x+1/2*c)+5)^2+59/1024*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.13

$$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx$$

$$= \frac{59 (9 \cos(dx + c)^2 + 30 \sin(dx + c) - 34) \arctan\left(\frac{5 \sin(dx+c)-3}{4 \cos(dx+c)}\right) - 540 \cos(dx + c) \sin(dx + c) + 1092}{2048 (9 d \cos(dx + c)^2 + 30 d \sin(dx + c) - 34 d)}$$

input `integrate(1/(5-3*sin(d*x+c))^3,x, algorithm="fricas")`

output `1/2048*(59*(9*cos(d*x + c)^2 + 30*sin(d*x + c) - 34)*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c)) - 540*cos(d*x + c)*sin(d*x + c) + 1092*cos(d*x + c))/(9*d*cos(d*x + c)^2 + 30*d*sin(d*x + c) - 34*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.04 (sec) , antiderivative size = 915, normalized size of antiderivative = 11.02

$$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx = \text{Too large to display}$$

input `integrate(1/(5-3*sin(d*x+c))**3,x)`

output

```
Piecewise((x/(5 - 3*sin(2*atan(3/5 - 4*I/5)))**3, Eq(c, -d*x + 2*atan(3/5
- 4*I/5))), (x/(5 - 3*sin(2*atan(3/5 + 4*I/5)))**3, Eq(c, -d*x + 2*atan(3/
5 + 4*I/5))), (x/(5 - 3*sin(c))**3, Eq(d, 0)), (36875*(atan(5*tan(c/2 + d*
x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(64
0000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan
(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2) + 640000*d) - 88500*(atan(5*
tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d
*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2 + d*x/2)**3 + 2
201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 12
6850*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi)
)*tan(c/2 + d*x/2)**2/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2 +
d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2) + 6
40000*d) - 88500*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2
- pi/2)/pi))*tan(c/2 + d*x/2)/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*t
an(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d
*x/2) + 640000*d) + 36875*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/
2 + d*x/2 - pi/2)/pi))/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2 +
d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2) +
640000*d) + 19260*tan(c/2 + d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 - 1536
000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*t...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(74) = 148$.

Time = 0.12 (sec) , antiderivative size = 173, normalized size of antiderivative = 2.08

$$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx = \frac{12 \left(\frac{3855 \sin(dx+c)}{\cos(dx+c)+1} - \frac{3913 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1605 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - 2275 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} - \frac{86 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{25 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 25} - 1475 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} - \frac{3}{4} \right) \frac{1}{25600 d}$$

input

```
integrate(1/(5-3*sin(d*x+c))^3,x, algorithm="maxima")
```

output

```
-1/25600*(12*(3855*sin(d*x + c)/(cos(d*x + c) + 1) - 3913*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1605*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 2275)/(60*sin(d*x + c)/(cos(d*x + c) + 1) - 86*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 25*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - 25) - 1475*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.47

$$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx$$

$$= \frac{1475 dx + 1475 c + \frac{24 \left(1605 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 3913 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 3855 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2275 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^2} + 2950 \arctan\left(\frac{3 \cos(dx+c) - \sin(dx+c)}{\cos(dx+c)+3}\right)}{51200 d}$$

input

```
integrate(1/(5-3*sin(d*x+c))^3,x, algorithm="giac")
```

output

```
1/51200*(1475*d*x + 1475*c + 24*(1605*tan(1/2*d*x + 1/2*c)^3 - 3913*tan(1/2*d*x + 1/2*c)^2 + 3855*tan(1/2*d*x + 1/2*c) - 2275)/(5*tan(1/2*d*x + 1/2*c)^2 - 6*tan(1/2*d*x + 1/2*c) + 5)^2 + 2950*arctan((3*cos(d*x + c) - sin(d*x + c) + 3)/(cos(d*x + c) + 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 25.98 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.34

$$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx = \frac{59 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{3}{4}\right)}{1024 d} - \frac{59 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{1024 d}$$

$$+ \frac{\frac{963 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{1280} - \frac{11739 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{1280} - \frac{273}{256}}{d \left(5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 5 \right)^2}$$

input

```
int(-1/(3*sin(c + d*x) - 5)^3,x)
```

output

```
(59*atan((5*tan(c/2 + (d*x)/2))/4 - 3/4))/(1024*d) - (59*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) + ((2313*tan(c/2 + (d*x)/2))/1280 - (11739*atan(c/2 + (d*x)/2)^2)/6400 + (963*tan(c/2 + (d*x)/2)^3)/1280 - 273/256)/(d*(5*tan(c/2 + (d*x)/2)^2 - 6*tan(c/2 + (d*x)/2) + 5)^2)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.58

$$\int \frac{1}{(5 - 3 \sin(c + dx))^3} dx$$

$$= \frac{531 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right)}{4} - \frac{3}{4}\right) \sin(dx+c)^2 - 1770 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right)}{4} - \frac{3}{4}\right) \sin(dx+c) + 1475 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right)}{4}\right)}{1024d (9 \sin(dx+c)^2 -$$

input

```
int(1/(5-3*sin(d*x+c))^3,x)
```

output

```
(531*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x)**2 - 1770*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x) + 1475*atan((5*tan((c + d*x)/2) - 3)/4) + 270*cos(c + d*x)*sin(c + d*x) - 546*cos(c + d*x) - 81*sin(c + d*x)**2 + 270*sin(c + d*x) - 225)/(1024*d*(9*sin(c + d*x)**2 - 30*sin(c + d*x) + 25))
```

3.51 $\int \frac{1}{(5-3 \sin(c+dx))^4} dx$

Optimal result	358
Mathematica [A] (verified)	358
Rubi [A] (verified)	359
Maple [A] (verified)	362
Fricas [A] (verification not implemented)	362
Sympy [C] (verification not implemented)	363
Maxima [B] (verification not implemented)	364
Giac [A] (verification not implemented)	364
Mupad [B] (verification not implemented)	365
Reduce [B] (verification not implemented)	366

Optimal result

Integrand size = 12, antiderivative size = 108

$$\int \frac{1}{(5-3 \sin(c+dx))^4} dx = \frac{385x}{32768} - \frac{385 \arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{16384d} - \frac{\cos(c+dx)}{16d(5-3 \sin(c+dx))^3} - \frac{25 \cos(c+dx)}{512d(5-3 \sin(c+dx))^2} - \frac{311 \cos(c+dx)}{8192d(5-3 \sin(c+dx))}$$

```
output 385/32768*x-385/16384*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d-1/16*cos(d*x+c)/
d/(5-3*sin(d*x+c))^3-25/512*cos(d*x+c)/d/(5-3*sin(d*x+c))^2-311/8192*cos(d
*x+c)/d/(5-3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.23

$$\int \frac{1}{(5-3 \sin(c+dx))^4} dx = \frac{-1925 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right) + \frac{-239470+219735 \cos(c+dx)+83970 \cos(2(c+dx))-13995 \cos(3(c+dx))+305091 \cos(4(c+dx))}{2(-5+3 \sin(c+dx))^3}}{81920d}$$

```
input Integrate[(5 - 3*Sin[c + d*x])^(-4), x]
```

output

```
(-1925*ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2]
+ Sin[(c + d*x)/2])] + (-239470 + 219735*Cos[c + d*x] + 83970*Cos[2*(c + d
*x)] - 13995*Cos[3*(c + d*x)] + 305091*Sin[c + d*x] - 105300*Sin[2*(c + d*
x)] - 8397*Sin[3*(c + d*x)])/(2*(-5 + 3*Sin[c + d*x])^3)/(81920*d)
```

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{1}{(5 - 3 \sin(c + dx))^4} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{1}{(5 - 3 \sin(c + dx))^4} dx \\
& \quad \downarrow \text{3143} \\
& -\frac{1}{48} \int -\frac{3(2 \sin(c + dx) + 5)}{(5 - 3 \sin(c + dx))^3} dx - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \quad \downarrow \text{27} \\
& \frac{1}{16} \int \frac{2 \sin(c + dx) + 5}{(5 - 3 \sin(c + dx))^3} dx - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{16} \int \frac{2 \sin(c + dx) + 5}{(5 - 3 \sin(c + dx))^3} dx - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \quad \downarrow \text{3233} \\
& \frac{1}{16} \left(-\frac{1}{32} \int -\frac{25 \sin(c + dx) + 62}{(5 - 3 \sin(c + dx))^2} dx - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \quad \downarrow \text{25} \\
& \frac{1}{16} \left(\frac{1}{32} \int \frac{25 \sin(c + dx) + 62}{(5 - 3 \sin(c + dx))^2} dx - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3}
\end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \int \frac{25 \sin(c+dx) + 62}{(5 - 3 \sin(c+dx))^2} dx - \frac{25 \cos(c+dx)}{32d(5 - 3 \sin(c+dx))^2} \right) - \frac{\cos(c+dx)}{16d(5 - 3 \sin(c+dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{16} \left(\frac{1}{32} \left(-\frac{1}{16} \int -\frac{385}{5 - 3 \sin(c+dx)} dx - \frac{311 \cos(c+dx)}{16d(5 - 3 \sin(c+dx))} \right) - \frac{25 \cos(c+dx)}{32d(5 - 3 \sin(c+dx))^2} \right) - \\
& \quad \frac{\cos(c+dx)}{16d(5 - 3 \sin(c+dx))^3} \\
& \downarrow 27 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{5 - 3 \sin(c+dx)} dx - \frac{311 \cos(c+dx)}{16d(5 - 3 \sin(c+dx))} \right) - \frac{25 \cos(c+dx)}{32d(5 - 3 \sin(c+dx))^2} \right) - \\
& \quad \frac{\cos(c+dx)}{16d(5 - 3 \sin(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{5 - 3 \sin(c+dx)} dx - \frac{311 \cos(c+dx)}{16d(5 - 3 \sin(c+dx))} \right) - \frac{25 \cos(c+dx)}{32d(5 - 3 \sin(c+dx))^2} \right) - \\
& \quad \frac{\cos(c+dx)}{16d(5 - 3 \sin(c+dx))^3} \\
& \downarrow 3136 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \left(\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3 - \sin(c+dx)}\right)}{2d} \right) - \frac{311 \cos(c+dx)}{16d(5 - 3 \sin(c+dx))} \right) - \frac{25 \cos(c+dx)}{32d(5 - 3 \sin(c+dx))^2} \right) - \\
& \quad \frac{\cos(c+dx)}{16d(5 - 3 \sin(c+dx))^3}
\end{aligned}$$

input `Int[(5 - 3*Sin[c + d*x])^(-4),x]`

output `((((385*(x/4 - ArcTan[Cos[c + d*x]/(3 - Sin[c + d*x]])/(2*d)))/16 - (311*Cos[c + d*x]/(16*d*(5 - 3*Sin[c + d*x])))/32 - (25*Cos[c + d*x]/(32*d*(5 - 3*Sin[c + d*x])^2))/16 - Cos[c + d*x]/(16*d*(5 - 3*Sin[c + d*x])^3)`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3136 $\text{Int}[(\text{a}_) + (\text{b}_)*\sin[(\text{c}_) + (\text{d}_)*(x_)]^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = \text{Rt}[\text{a}^2 - \text{b}^2, 2]\}, \text{Simp}[\text{x}/\text{q}, \text{x}] + \text{Simp}[(2/(\text{d}*\text{q}))*\text{ArcTan}[\text{b}*(\text{Cos}[\text{c} + \text{d}*x]/(\text{a} + \text{q} + \text{b}*\text{Sin}[\text{c} + \text{d}*x]))], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{PosQ}[\text{a}]$
- rule 3143 $\text{Int}[(\text{a}_) + (\text{b}_)*\sin[(\text{c}_) + (\text{d}_)*(x_)]^{(\text{n}_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b})*\text{Cos}[\text{c} + \text{d}*x]*((\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(\text{n} + 1)}/(\text{d}*(\text{n} + 1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((\text{n} + 1)*(a^2 - b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(\text{n} + 1)}*\text{Simp}[\text{a}*(\text{n} + 1) - \text{b}*(\text{n} + 2)*\text{Sin}[\text{c} + \text{d}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{n}, -1] \ \&\& \ \text{IntegerQ}[2*\text{n}]$
- rule 3233 $\text{Int}[(\text{a}_) + (\text{b}_)*\sin[(\text{e}_) + (\text{f}_)*(x_)]^{(\text{m}_)}*((\text{c}_) + (\text{d}_)*\sin[(\text{e}_) + (\text{f}_)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{b}*\text{c} - \text{a}*\text{d}))*\text{Cos}[\text{e} + \text{f}*x]*((\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(\text{m} + 1)}/(\text{f}*(\text{m} + 1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((\text{m} + 1)*(a^2 - b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(\text{m} + 1)}*\text{Simp}[(\text{a}*\text{c} - \text{b}*\text{d})*(\text{m} + 1) - (\text{b}*\text{c} - \text{a}*\text{d})*(\text{m} + 2)*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntegerQ}[2*\text{m}]$

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.08

method	result
derivativedivides	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} - \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} - \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} - \frac{10287}{4096} + \frac{385}{d} \left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3$
default	$\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} - \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} - \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} - \frac{10287}{4096} + \frac{385}{d} \left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3$
risch	$-\frac{-239470 e^{3i(dx+c)} - 86625 i e^{4i(dx+c)} + 218466 i e^{2i(dx+c)} + 10395 e^{5i(dx+c)} + 73575 e^{i(dx+c)} - 8397 i}{12288(3 e^{2i(dx+c)} - 3 - 10 i e^{i(dx+c)})^3} d + \frac{385 i \ln(e^{i(dx+c)})}{32768}$
parallelrisc	$31683960 + 48125i(27 \sin(3dx+3c) - 981 \sin(dx+c) - 270 \cos(2dx+2c) + 770) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + 48125i(-27 \sin(dx+c) - 270 \cos(2dx+2c) + 770)$

input `int(1/(5-3*sin(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*(250*(39933/5120000*tan(1/2*d*x+1/2*c)^5-672723/25600000*tan(1/2*d*x+1/2*c)^4+2870073/64000000*tan(1/2*d*x+1/2*c)^3-604899/12800000*tan(1/2*d*x+1/2*c)^2+145233/5120000*tan(1/2*d*x+1/2*c)-10287/1024000)/(5*tan(1/2*d*x+1/2*c)^2-6*tan(1/2*d*x+1/2*c)+5)^3+385/16384*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.20

$$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx = \frac{11196 \cos(dx + c)^3 - 385 (135 \cos(dx + c)^2 - 9 (3 \cos(dx + c)^2 - 28) \sin(dx + c) - 260) \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i}{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i}\right)}{32768 (135 d \cos(dx + c)^2 - 9 (3 d \cos(dx + c)^2 - 28 d))}$$

input `integrate(1/(5-3*sin(d*x+c))^4,x, algorithm="fricas")`

output

```
-1/32768*(11196*cos(d*x + c)^3 - 385*(135*cos(d*x + c)^2 - 9*(3*cos(d*x +
c)^2 - 28)*sin(d*x + c) - 260)*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c
)) + 42120*cos(d*x + c)*sin(d*x + c) - 52344*cos(d*x + c))/(135*d*cos(d*x
+ c)^2 - 9*(3*d*cos(d*x + c)^2 - 28*d)*sin(d*x + c) - 260*d)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.66 (sec) , antiderivative size = 1690, normalized size of antiderivative = 15.65

$$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(5-3*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(5 - 3*sin(2*atan(3/5 - 4*I/5)))**4, Eq(c, -d*x + 2*atan(3/5
- 4*I/5))), (x/(5 - 3*sin(2*atan(3/5 + 4*I/5)))**4, Eq(c, -d*x + 2*atan(3/
5 + 4*I/5))), (x/(5 - 3*sin(c))**4, Eq(d, 0)), (6015625*(atan(5*tan(c/2 +
d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**6/(
256000000*d*tan(c/2 + d*x/2)**6 - 921600000*d*tan(c/2 + d*x/2)**5 + 187392
0000*d*tan(c/2 + d*x/2)**4 - 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000
*d*tan(c/2 + d*x/2)**2 - 921600000*d*tan(c/2 + d*x/2) + 256000000*d) - 216
56250*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi
))*tan(c/2 + d*x/2)**5/(256000000*d*tan(c/2 + d*x/2)**6 - 921600000*d*tan(
c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)**4 - 2285568000*d*tan(c/2
+ d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 - 921600000*d*tan(c/2 + d*x
/2) + 256000000*d) + 44034375*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor
((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(256000000*d*tan(c/2 + d*x/
2)**6 - 921600000*d*tan(c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)**4
- 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 - 9
21600000*d*tan(c/2 + d*x/2) + 256000000*d) - 53707500*(atan(5*tan(c/2 + d
x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**3/(25
6000000*d*tan(c/2 + d*x/2)**6 - 921600000*d*tan(c/2 + d*x/2)**5 + 18739200
00*d*tan(c/2 + d*x/2)**4 - 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000*d
*tan(c/2 + d*x/2)**2 - 921600000*d*tan(c/2 + d*x/2) + 256000000*d) + 44...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(97) = 194.

Time = 0.11 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.34

$$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx = \frac{36 \left(\frac{403425 \sin(dx+c)}{\cos(dx+c)+1} - \frac{672110 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{637794 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{373735 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{110925 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - 142875 \right) - 48125 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} \right)}{2048000 d}$$

input `integrate(1/(5-3*sin(d*x+c))^4,x, algorithm="maxima")`

output `-1/2048000*(36*(403425*sin(d*x + c)/(cos(d*x + c) + 1) - 672110*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 637794*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 373735*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 110925*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 142875)/(450*sin(d*x + c)/(cos(d*x + c) + 1) - 915*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1116*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 915*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 450*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 125*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 - 125) - 48125*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4))/d`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.37

$$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx = \frac{48125 dx + 48125 c + \frac{72 \left(110925 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 373735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 637794 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 672110 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 403425 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 142875 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^3}{4096000 d}$$

input `integrate(1/(5-3*sin(d*x+c))^4,x, algorithm="giac")`

output

```
1/4096000*(48125*d*x + 48125*c + 72*(110925*tan(1/2*d*x + 1/2*c)^5 - 37373
5*tan(1/2*d*x + 1/2*c)^4 + 637794*tan(1/2*d*x + 1/2*c)^3 - 672110*tan(1/2*
d*x + 1/2*c)^2 + 403425*tan(1/2*d*x + 1/2*c) - 142875)/(5*tan(1/2*d*x + 1/
2*c)^2 - 6*tan(1/2*d*x + 1/2*c) + 5)^3 + 96250*arctan((3*cos(d*x + c) - si
n(d*x + c) + 3)/(cos(d*x + c) + 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 26.42 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.73

$$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{385 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{3}{4}\right)}{16384 d} - \frac{385 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{16384 d}$$

$$+ \frac{\frac{39933 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{2560000} - \frac{672723 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{12800000} + \frac{2870073 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{32000000} - \frac{604899 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400000} + \frac{145233 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2560000} - \frac{10287}{512000}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{5} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{25} - \frac{1116 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{125} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{25} - \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1 \right)}$$

input

```
int(1/(3*sin(c + d*x) - 5)^4,x)
```

output

```
(385*atan((5*tan(c/2 + (d*x)/2))/4 - 3/4))/(16384*d) - (385*(atan(tan(c/2
+ (d*x)/2)) - (d*x)/2))/(16384*d) + ((145233*tan(c/2 + (d*x)/2))/2560000 -
(604899*tan(c/2 + (d*x)/2)^2)/6400000 + (2870073*tan(c/2 + (d*x)/2)^3)/32
000000 - (672723*tan(c/2 + (d*x)/2)^4)/12800000 + (39933*tan(c/2 + (d*x)/2
)^5)/2560000 - 10287/512000)/(d*((183*tan(c/2 + (d*x)/2)^2)/25 - (18*tan(c
/2 + (d*x)/2))/5 - (1116*tan(c/2 + (d*x)/2)^3)/125 + (183*tan(c/2 + (d*x)/
2)^4)/25 - (18*tan(c/2 + (d*x)/2)^5)/5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.77

$$\int \frac{1}{(5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{51975 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx + c}{2}\right) - 3}{4}\right) \sin(dx + c)^3 - 259875 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx + c}{2}\right) - 3}{4}\right) \sin(dx + c)^2 + 433125 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx + c}{2}\right) - 3}{4}\right) \sin(dx + c) - 240625 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx + c}{2}\right) - 3}{4}\right) + 27990 \cos(c + dx) \sin(c + dx)^2 - 105300 \cos(c + dx) \sin(c + dx) + 102870 \cos(c + dx) - 12636 \sin(c + dx)^3 + 63180 \sin(c + dx)^2 - 105300 \sin(c + dx) + 58500}{(81920 d (27 \sin(c + dx)^3 - 135 \sin(c + dx)^2 + 225 \sin(c + dx) - 125))}$$

input `int(1/(5-3*sin(d*x+c))^4,x)`output `(51975*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x)**3 - 259875*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x)**2 + 433125*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x) - 240625*atan((5*tan((c + d*x)/2) - 3)/4) + 27990*cos(c + d*x)*sin(c + d*x)**2 - 105300*cos(c + d*x)*sin(c + d*x) + 102870*cos(c + d*x) - 12636*sin(c + d*x)**3 + 63180*sin(c + d*x)**2 - 105300*sin(c + d*x) + 58500)/(81920*d*(27*sin(c + d*x)**3 - 135*sin(c + d*x)**2 + 225*sin(c + d*x) - 125))`

3.52 $\int \frac{1}{-5+3\sin(c+dx)} dx$

Optimal result	367
Mathematica [A] (verified)	367
Rubi [A] (verified)	368
Maple [A] (verified)	369
Fricas [A] (verification not implemented)	369
Sympy [B] (verification not implemented)	370
Maxima [A] (verification not implemented)	370
Giac [A] (verification not implemented)	370
Mupad [B] (verification not implemented)	371
Reduce [B] (verification not implemented)	371

Optimal result

Integrand size = 12, antiderivative size = 33

$$\int \frac{1}{-5+3\sin(c+dx)} dx = -\frac{x}{4} + \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d}$$

output `-1/4*x+1/2*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.70

$$\int \frac{1}{-5+3\sin(c+dx)} dx = \frac{\arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right)}{2d}$$

input `Integrate[(-5 + 3*Sin[c + d*x])^(-1),x]`

output `ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])]/(2*d)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3137}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{3 \sin(c + dx) - 5} dx$$

↓ 3042

$$\int \frac{1}{3 \sin(c + dx) - 5} dx$$

↓ 3137

$$\frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d} - \frac{x}{4}$$

input `Int[(-5 + 3*Sin[c + d*x])^(-1),x]`

output `-1/4*x + ArcTan[Cos[c + d*x]/(3 - Sin[c + d*x])]/(2*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3137 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[-x/q, x] - Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a - q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] & & NegQ[a]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.61

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3}{4}\right)}{2d}$	20
default	$-\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3}{4}\right)}{2d}$	20
risch	$-\frac{i \ln(e^{i(dx+c)} - 3i)}{4d} + \frac{i \ln(e^{i(dx+c)} - \frac{i}{3})}{4d}$	40
parallelrisch	$-\frac{i\left(-\ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 + 4i\right)\right)}{4d}$	42

input `int(1/(-5+3*sin(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/2/d*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{1}{-5 + 3 \sin(c + dx)} dx = -\frac{\arctan\left(\frac{5 \sin(dx+c) - 3}{4 \cos(dx+c)}\right)}{4d}$$

input `integrate(1/(-5+3*sin(d*x+c)),x, algorithm="fricas")`

output `-1/4*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(22) = 44$.

Time = 0.36 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.45

$$\int \frac{1}{-5 + 3 \sin(c + dx)} dx = \begin{cases} -\frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 3}{4}\right) + \pi \left\lfloor \frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{3 \sin(c) - 5} & \text{otherwise} \end{cases}$$

input `integrate(1/(-5+3*sin(d*x+c)),x)`

output `Piecewise((-atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(2*d), Ne(d, 0)), (x/(3*sin(c) - 5), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{1}{-5 + 3 \sin(c + dx)} dx = -\frac{\arctan\left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} - \frac{3}{4}\right)}{2d}$$

input `integrate(1/(-5+3*sin(d*x+c)),x, algorithm="maxima")`

output `-1/2*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4)/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.52

$$\int \frac{1}{-5 + 3 \sin(c + dx)} dx = -\frac{dx + c + 2 \arctan\left(\frac{3 \cos(dx+c) - \sin(dx+c) + 3}{\cos(dx+c) + 3 \sin(dx+c) - 9}\right)}{4d}$$

input `integrate(1/(-5+3*sin(d*x+c)),x, algorithm="giac")`

output $-1/4*(d*x + c + 2*\arctan((3*\cos(d*x + c) - \sin(d*x + c) + 3)/(\cos(d*x + c) + 3*\sin(d*x + c) - 9)))/d$

Mupad [B] (verification not implemented)

Time = 25.86 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21

$$\int \frac{1}{-5 + 3 \sin(c + dx)} dx = \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d} - \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{3}{4}}{4}\right)}{2d}$$

input `int(1/(3*sin(c + d*x) - 5),x)`

output $(\operatorname{atan}(\tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d) - \operatorname{atan}((5*\tan(c/2 + (d*x)/2))/4 - 3/4)/(2*d)$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.58

$$\int \frac{1}{-5 + 3 \sin(c + dx)} dx = -\frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}}{4}\right)}{2d}$$

input `int(1/(-5+3*sin(d*x+c)),x)`

output $(- \operatorname{atan}((5*\tan((c + d*x)/2) - 3)/4))/(2*d)$

3.53 $\int \frac{1}{(-5+3 \sin(c+dx))^2} dx$

Optimal result	372
Mathematica [A] (verified)	372
Rubi [A] (verified)	373
Maple [A] (verified)	374
Fricas [A] (verification not implemented)	375
Sympy [C] (verification not implemented)	376
Maxima [A] (verification not implemented)	376
Giac [A] (verification not implemented)	377
Mupad [B] (verification not implemented)	377
Reduce [B] (verification not implemented)	378

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx = \frac{5x}{64} - \frac{5 \arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{32d} - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))}$$

output

```
5/64*x-5/32*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d-3/16*cos(d*x+c)/d/(5-3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.57

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx = \frac{-25 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right) + \frac{6(-5+5 \cos(c+dx)+3 \sin(c+dx))}{-5+3 \sin(c+dx)}}{160d}$$

input

```
Integrate[(-5 + 3*Sin[c + d*x])^(-2), x]
```

output

```
(-25*ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])] + (6*(-5 + 5*Cos[c + d*x] + 3*Sin[c + d*x]))/(-5 + 3*Sin[c + d*x]))/(160*d)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 \sin(c + dx) - 5)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 \sin(c + dx) - 5)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & -\frac{1}{16} \int -\frac{5}{5 - 3 \sin(c + dx)} dx - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{5}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \\
 & \quad \downarrow \text{3136} \\
 & \frac{5}{16} \left(\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d} \right) - \frac{3 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))}
 \end{aligned}$$

input

```
Int[(-5 + 3*Sin[c + d*x])^(-2),x]
```

output $(5*(x/4 - \text{ArcTan}[\text{Cos}[c + d*x]/(3 - \text{Sin}[c + d*x])]/(2*d)))/16 - (3*\text{Cos}[c + d*x])/(16*d*(5 - 3*\text{Sin}[c + d*x]))$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3136 $\text{Int}[(a_ + (b_)*\text{sin}[(c_.) + (d_)*(x_)])^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2 - b^2, 2]\}, \text{Simp}[x/q, x] + \text{Simp}[(2/(d*q))*\text{ArcTan}[b*(\text{Cos}[c + d*x]/(a + q + b*\text{Sin}[c + d*x]))], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{GtQ}[a^2 - b^2, 0] \ \&\& \ \text{PosQ}[a]$

rule 3143 $\text{Int}[(a_ + (b_)*\text{sin}[(c_.) + (d_)*(x_)])^{n_}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^{n+1}/(d*(n+1)*(a^2 - b^2))), x] + \text{Simp}[1/((n+1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[c + d*x])^{n+1}*\text{Simp}[a*(n+1) - b*(n+2)*\text{Sin}[c + d*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
default	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
risch	$-\frac{5 e^{i(dx+c)} - 3i}{8d(3 e^{2i(dx+c)} - 3 - 10i e^{i(dx+c)})} + \frac{5i \ln(e^{i(dx+c)} - 3i)}{64d} - \frac{5i \ln(e^{i(dx+c)} - \frac{i}{3})}{64d}$
parallelrisch	$\frac{(125i - 75i \sin(dx+c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + (75i \sin(dx+c) - 125i) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 + 4i\right) + 60 \cos(dx+c) - 36}{960 \sin(dx+c)d - 1600d}$

input `int(1/(-5+3*sin(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(2*(9/400*tan(1/2*d*x+1/2*c)-3/80)/(tan(1/2*d*x+1/2*c)^2-6/5*tan(1/2*d*x+1/2*c)+1)+5/32*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx$$

$$= \frac{5(3 \sin(dx + c) - 5) \arctan\left(\frac{5 \sin(dx+c) - 3}{4 \cos(dx+c)}\right) + 12 \cos(dx + c)}{64(3d \sin(dx + c) - 5d)}$$

input `integrate(1/(-5+3*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/64*(5*(3*sin(d*x + c) - 5)*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c)) + 12*cos(d*x + c))/(3*d*sin(d*x + c) - 5*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.91 (sec) , antiderivative size = 384, normalized size of antiderivative = 6.62

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(-5+3*sin(d*x+c))**2,x)`

output `Piecewise((x/(-5 + 3*sin(2*atan(3/5 - 4*I/5)))**2, Eq(c, -d*x + 2*atan(3/5 - 4*I/5))), (x/(-5 + 3*sin(2*atan(3/5 + 4*I/5)))**2, Eq(c, -d*x + 2*atan(3/5 + 4*I/5))), (x/(3*sin(c) - 5)**2, Eq(d, 0)), (125*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) - 150*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) + 125*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) + 36*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d) - 60/(800*d*tan(c/2 + d*x/2)**2 - 960*d*tan(c/2 + d*x/2) + 800*d), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.60

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx = -\frac{12 \left(\frac{3 \sin(dx+c) - 5}{\cos(dx+c)+1} \right)}{\frac{6 \sin(dx+c)}{\cos(dx+c)+1} - \frac{5 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 5} - \frac{25 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} - \frac{3}{4} \right)}{160 d}$$

input `integrate(1/(-5+3*sin(d*x+c))^2,x, algorithm="maxima")`

output `-1/160*(12*(3*sin(d*x + c)/(cos(d*x + c) + 1) - 5)/(6*sin(d*x + c)/(cos(d*x + c) + 1) - 5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 5) - 25*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4))/d`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int \frac{1}{(-5 + 3\sin(c + dx))^2} dx$$

$$= \frac{25 dx + 25 c + \frac{24(3 \tan(\frac{1}{2} dx + \frac{1}{2} c) - 5)}{5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 6 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5} + 50 \arctan\left(\frac{3 \cos(dx+c) - \sin(dx+c) + 3}{\cos(dx+c) + 3 \sin(dx+c) - 9}\right)}{320 d}$$

input `integrate(1/(-5+3*sin(d*x+c))^2,x, algorithm="giac")`

output

```
1/320*(25*d*x + 25*c + 24*(3*tan(1/2*d*x + 1/2*c) - 5)/(5*tan(1/2*d*x + 1/2*c)^2 - 6*tan(1/2*d*x + 1/2*c) + 5) + 50*arctan((3*cos(d*x + c) - sin(d*x + c) + 3)/(cos(d*x + c) + 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.43

$$\int \frac{1}{(-5 + 3\sin(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 3}{4}\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d}$$

$$+ \frac{\frac{9 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{200} - \frac{3}{40}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1\right)}$$

input `int(1/(3*sin(c + d*x) - 5)^2,x)`

output

```
(5*atan((5*tan(c/2 + (d*x)/2))/4 - 3/4))/(32*d) - (5*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(32*d) + ((9*tan(c/2 + (d*x)/2))/200 - 3/40)/(d*(tan(c/2 + (d*x)/2)^2 - (6*tan(c/2 + (d*x)/2))/5 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^2} dx$$

$$= \frac{15 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) \sin(dx+c) - 25 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) + 6 \cos(dx+c)}{32d(3 \sin(dx+c) - 5)}$$

input

```
int(1/(-5+3*sin(d*x+c))^2,x)
```

output

```
(15*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x) - 25*atan((5*tan((c + d*x)/2) - 3)/4) + 6*cos(c + d*x))/(32*d*(3*sin(c + d*x) - 5))
```

3.54 $\int \frac{1}{(-5+3 \sin(c+dx))^3} dx$

Optimal result	379
Mathematica [A] (verified)	379
Rubi [A] (verified)	380
Maple [A] (verified)	382
Fricas [A] (verification not implemented)	383
Sympy [C] (verification not implemented)	383
Maxima [B] (verification not implemented)	384
Giac [A] (verification not implemented)	385
Mupad [B] (verification not implemented)	385
Reduce [B] (verification not implemented)	386

Optimal result

Integrand size = 12, antiderivative size = 83

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx = -\frac{59x}{2048} + \frac{59 \arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{1024d} + \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} + \frac{45 \cos(c + dx)}{512d(5 - 3 \sin(c + dx))}$$

output

```
-59/2048*x+59/1024*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d+3/32*cos(d*x+c)/d/(5-3*sin(d*x+c))^2+45/512*cos(d*x+c)/d/(5-3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.36

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx = \frac{59 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right) + \frac{546 \cos(c+dx)+9(-59+9 \cos(2(c+dx))+60 \sin(c+dx)-15 \sin(2(c+dx)))}{(5-3 \sin(c+dx))^2}}{1024d}$$

input

```
Integrate[(-5 + 3*Sin[c + d*x])^(-3), x]
```

output

```
(59*ArcTan[(2*(Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] + Sin[(c + d*x)/2])] + (546*Cos[c + d*x] + 9*(-59 + 9*Cos[2*(c + d*x)] + 60*Sin[c + d*x] - 15*Sin[2*(c + d*x)]))/(5 - 3*Sin[c + d*x])^2)/(1024*d)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3 \sin(c + dx) - 5)^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(3 \sin(c + dx) - 5)^3} dx$$

$$\downarrow \text{3143}$$

$$\frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} - \frac{1}{32} \int \frac{3 \sin(c + dx) + 10}{(5 - 3 \sin(c + dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} - \frac{1}{32} \int \frac{3 \sin(c + dx) + 10}{(5 - 3 \sin(c + dx))^2} dx$$

$$\downarrow \text{3233}$$

$$\frac{1}{32} \left(\frac{1}{16} \int -\frac{59}{5 - 3 \sin(c + dx)} dx + \frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) + \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2}$$

$$\downarrow \text{27}$$

$$\frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} - \frac{59}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx \right) + \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2}$$

$$\downarrow \text{3042}$$

$$\frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} - \frac{59}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx \right) + \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2}$$

$$\begin{array}{c} \downarrow \text{3136} \\ \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} - \frac{59}{16} \left(\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d} \right) \right) + \frac{3 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \end{array}$$

input `Int[(-5 + 3*Sin[c + d*x])^(-3),x]`

output `((-59*(x/4 - ArcTan[Cos[c + d*x]/(3 - Sin[c + d*x]))/(2*d)))/16 + (45*Cos[c + d*x])/(16*d*(5 - 3*Sin[c + d*x]))/32 + (3*Cos[c + d*x])/(32*d*(5 - 3*Sin[c + d*x])^2)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] :> Simp[(-*(b*c - a*d))*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{50 \left(\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{64000} - \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{320000} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64000} - \frac{273}{12800} \right) - 59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} - \frac{3}{4}\right)}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^2 d}$
default	$\frac{50 \left(\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{64000} - \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{320000} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64000} - \frac{273}{12800} \right) - 59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} - \frac{3}{4}\right)}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^2 d}$
risch	$\frac{-\frac{885ie^{2i(dx+c)}}{256} + \frac{177e^{3i(dx+c)}}{256} - \frac{723e^{i(dx+c)}}{256} + \frac{135i}{256}}{(3e^{2i(dx+c)} - 3 - 10ie^{i(dx+c)})^2 d} - \frac{59i \ln(e^{i(dx+c)} - 3i)}{2048d} + \frac{59i \ln(e^{i(dx+c)} - \frac{i}{3})}{2048d}$
parallelrisc	$\frac{-1062 + 59i(9 \cos(2dx+2c) - 59 + 60 \sin(dx+c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + 59i(59 - 9 \cos(2dx+2c) - 60 \sin(dx+c)) \ln\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3}{4}\right)}{2048d(9 \cos(2dx+2c) - 59 + 60 \sin(dx+c))}$

input

```
int(1/(-5+3*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-50*(963/64000*tan(1/2*d*x+1/2*c)^3-11739/320000*tan(1/2*d*x+1/2*c)^2
+2313/64000*tan(1/2*d*x+1/2*c)-273/12800)/(5*tan(1/2*d*x+1/2*c)^2-6*tan(1/
2*d*x+1/2*c)+5)^2-59/1024*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.13

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx = \frac{59 (9 \cos(dx + c)^2 + 30 \sin(dx + c) - 34) \arctan\left(\frac{5 \sin(dx+c)-3}{4 \cos(dx+c)}\right) - 540 \cos(dx + c) \sin(dx + c) + 1092 \cos(dx + c)}{2048 (9 d \cos(dx + c)^2 + 30 d \sin(dx + c) - 34 d)}$$

input `integrate(1/(-5+3*sin(d*x+c))^3,x, algorithm="fricas")`

output `-1/2048*(59*(9*cos(d*x + c)^2 + 30*sin(d*x + c) - 34)*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c)) - 540*cos(d*x + c)*sin(d*x + c) + 1092*cos(d*x + c))/(9*d*cos(d*x + c)^2 + 30*d*sin(d*x + c) - 34*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.95 (sec) , antiderivative size = 915, normalized size of antiderivative = 11.02

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx = \text{Too large to display}$$

input `integrate(1/(-5+3*sin(d*x+c))**3,x)`

output

```
Piecewise((x/(-5 + 3*sin(2*atan(3/5 - 4*I/5)))**3, Eq(c, -d*x + 2*atan(3/5
- 4*I/5))), (x/(-5 + 3*sin(2*atan(3/5 + 4*I/5)))**3, Eq(c, -d*x + 2*atan(
3/5 + 4*I/5))), (x/(3*sin(c) - 5)**3, Eq(d, 0)), (-36875*(atan(5*tan(c/2 +
d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/
(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*
tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2) + 640000*d) + 88500*(atan
(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2
+ d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2 + d*x/2)**3
+ 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2) + 640000*d) -
126850*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/
pi))*tan(c/2 + d*x/2)**2/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/2
+ d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2)
+ 640000*d) + 88500*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*
x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*
d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2
+ d*x/2) + 640000*d) - 36875*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor(
(c/2 + d*x/2 - pi/2)/pi))/(640000*d*tan(c/2 + d*x/2)**4 - 1536000*d*tan(c/
2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*d*tan(c/2 + d*x/2)
+ 640000*d) - 19260*tan(c/2 + d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 - 1
536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 - 1536000*...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(74) = 148$.

Time = 0.11 (sec) , antiderivative size = 173, normalized size of antiderivative = 2.08

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx$$

$$= \frac{12 \left(\frac{3855 \sin(dx+c)}{\cos(dx+c)+1} - \frac{3913 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1605 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - 2275 \right) - 1475 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} - \frac{3}{4} \right)}{25600 d}$$

input

```
integrate(1/(-5+3*sin(d*x+c))^3,x, algorithm="maxima")
```

output

```
1/25600*(12*(3855*sin(d*x + c)/(cos(d*x + c) + 1) - 3913*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1605*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 2275)/(60*sin(d*x + c)/(cos(d*x + c) + 1) - 86*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 25*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - 25) - 1475*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) - 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.47

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx = \frac{1475 dx + 1475 c + \frac{24 \left(1605 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 3913 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 3855 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2275 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^2} + 2950 \arctan\left(\frac{3 \cos(dx + c) - \sin(dx + c)}{\cos(dx + c) + 3 \sin(dx + c) - 9}\right)}{51200 d}$$

input

```
integrate(1/(-5+3*sin(d*x+c))^3,x, algorithm="giac")
```

output

```
-1/51200*(1475*d*x + 1475*c + 24*(1605*tan(1/2*d*x + 1/2*c)^3 - 3913*tan(1/2*d*x + 1/2*c)^2 + 3855*tan(1/2*d*x + 1/2*c) - 2275)/(5*tan(1/2*d*x + 1/2*c)^2 - 6*tan(1/2*d*x + 1/2*c) + 5)^2 + 2950*arctan((3*cos(d*x + c) - sin(d*x + c) + 3)/(cos(d*x + c) + 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 26.00 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.35

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx = \frac{59 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{1024 d} - \frac{59 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{3}{4}\right)}{1024 d} - \frac{\frac{963 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{1280} - \frac{11739 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{1280} - \frac{273}{256}}{d \left(5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 5 \right)^2}$$

input `int(1/(3*sin(c + d*x) - 5)^3,x)`

output `(59*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) - (59*atan((5*tan(c/2 + (d*x)/2))/4 - 3/4))/(1024*d) - ((2313*tan(c/2 + (d*x)/2))/1280 - (11739*atan(c/2 + (d*x)/2)^2)/6400 + (963*tan(c/2 + (d*x)/2)^3)/1280 - 273/256)/(d*(5*tan(c/2 + (d*x)/2)^2 - 6*tan(c/2 + (d*x)/2) + 5)^2)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.58

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^3} dx$$

$$= \frac{-531 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right)}{4} - \frac{3}{4}\right) \sin(dx+c)^2 + 1770 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right)}{4} - \frac{3}{4}\right) \sin(dx+c) - 1475 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right)}{4} - \frac{3}{4}\right)}{1024d (9 \sin(dx+c))^2}$$

input `int(1/(-5+3*sin(d*x+c))^3,x)`

output `(- 531*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x)**2 + 1770*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x) - 1475*atan((5*tan((c + d*x)/2) - 3)/4) - 270*cos(c + d*x)*sin(c + d*x) + 546*cos(c + d*x) + 81*sin(c + d*x)**2 - 270*sin(c + d*x) + 225)/(1024*d*(9*sin(c + d*x)**2 - 30*sin(c + d*x) + 25))`

3.55 $\int \frac{1}{(-5+3 \sin(c+dx))^4} dx$

Optimal result	387
Mathematica [A] (verified)	387
Rubi [A] (verified)	388
Maple [A] (verified)	391
Fricas [A] (verification not implemented)	391
Sympy [C] (verification not implemented)	392
Maxima [B] (verification not implemented)	393
Giac [A] (verification not implemented)	393
Mupad [B] (verification not implemented)	394
Reduce [B] (verification not implemented)	395

Optimal result

Integrand size = 12, antiderivative size = 108

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx = \frac{385x}{32768} - \frac{385 \arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{16384d} - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} - \frac{25 \cos(c + dx)}{512d(5 - 3 \sin(c + dx))^2} - \frac{311 \cos(c + dx)}{8192d(5 - 3 \sin(c + dx))}$$

output `385/32768*x-385/16384*arctan(cos(d*x+c)/(3-sin(d*x+c)))/d-1/16*cos(d*x+c)/d/(5-3*sin(d*x+c))^3-25/512*cos(d*x+c)/d/(5-3*sin(d*x+c))^2-311/8192*cos(d*x+c)/d/(5-3*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.23

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx = \frac{-1925 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}\right) + \frac{-239470+219735 \cos(c+dx)+83970 \cos(2(c+dx))-13995 \cos(3(c+dx))+305091 \sin(c+dx)}{2(-5+3 \sin(c+dx))^3}}{81920d}$$

input `Integrate[(-5 + 3*Sin[c + d*x])^(-4), x]`

output

$$\begin{aligned} & (-1925 \operatorname{ArcTan}[(2 \cos[(c + dx)/2] - \sin[(c + dx)/2]) / (\cos[(c + dx)/2] \\ & + \sin[(c + dx)/2])] + (-239470 + 219735 \cos[c + dx] + 83970 \cos[2(c + d \\ & *x)] - 13995 \cos[3(c + dx)] + 305091 \sin[c + dx] - 105300 \sin[2(c + d * \\ & x)] - 8397 \sin[3(c + dx)]) / (2(-5 + 3 \sin[c + dx])^3) / (81920d) \end{aligned}$$
Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(3 \sin(c + dx) - 5)^4} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(3 \sin(c + dx) - 5)^4} dx \\ & \quad \downarrow \text{3143} \\ & -\frac{1}{48} \int -\frac{3(2 \sin(c + dx) + 5)}{(5 - 3 \sin(c + dx))^3} dx - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\ & \quad \downarrow \text{27} \\ & \frac{1}{16} \int \frac{2 \sin(c + dx) + 5}{(5 - 3 \sin(c + dx))^3} dx - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{16} \int \frac{2 \sin(c + dx) + 5}{(5 - 3 \sin(c + dx))^3} dx - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\ & \quad \downarrow \text{3233} \\ & \frac{1}{16} \left(-\frac{1}{32} \int -\frac{25 \sin(c + dx) + 62}{(5 - 3 \sin(c + dx))^2} dx - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\ & \quad \downarrow \text{25} \\ & \frac{1}{16} \left(\frac{1}{32} \int \frac{25 \sin(c + dx) + 62}{(5 - 3 \sin(c + dx))^2} dx - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \int \frac{25 \sin(c + dx) + 62}{(5 - 3 \sin(c + dx))^2} dx - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{16} \left(\frac{1}{32} \left(-\frac{1}{16} \int -\frac{385}{5 - 3 \sin(c + dx)} dx - \frac{311 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \\
& \quad \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \downarrow 27 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{311 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \\
& \quad \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{5 - 3 \sin(c + dx)} dx - \frac{311 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \\
& \quad \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3} \\
& \downarrow 3136 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \left(\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3-\sin(c+dx)}\right)}{2d} \right) - \frac{311 \cos(c + dx)}{16d(5 - 3 \sin(c + dx))} \right) - \frac{25 \cos(c + dx)}{32d(5 - 3 \sin(c + dx))^2} \right) - \\
& \quad \frac{\cos(c + dx)}{16d(5 - 3 \sin(c + dx))^3}
\end{aligned}$$

input `Int[(-5 + 3*Sin[c + d*x])^(-4),x]`

output `((((385*(x/4 - ArcTan[Cos[c + d*x]/(3 - Sin[c + d*x]])/(2*d)))/16 - (311*Cos[c + d*x])/(16*d*(5 - 3*Sin[c + d*x])))/32 - (25*Cos[c + d*x])/(32*d*(5 - 3*Sin[c + d*x])^2))/16 - Cos[c + d*x]/(16*d*(5 - 3*Sin[c + d*x])^3)`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \text{ Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3136 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(-1)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = \text{Rt}[\text{a}^2 - \text{b}^2, 2]\}, \text{Simp}[\text{x}/\text{q}, \text{x}] + \text{Simp}[(2/(d*q))*\text{ArcTan}[\text{b}*(\text{Cos}[\text{c} + \text{d}*x]/(\text{a} + \text{q} + \text{b}*\text{Sin}[\text{c} + \text{d}*x]))], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{PosQ}[\text{a}]$
- rule 3143 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(n)}, \text{x_Symbol}] \rightarrow \text{Simp}[-\text{b})*\text{Cos}[\text{c} + \text{d}*x]*((\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(n+1)}/(\text{d}*(n+1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((n+1)*(a^2 - b^2)) \text{ Int}[(\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(n+1)}*\text{Simp}[\text{a}*(n+1) - \text{b}*(n+2)*\text{Sin}[\text{c} + \text{d}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{n}, -1] \ \&\& \ \text{IntegerQ}[2*\text{n}]$
- rule 3233 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]^{(m)}*((\text{c}_.) + (\text{d}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[-(\text{b}*c - \text{a}*d)*\text{Cos}[\text{e} + \text{f}*x]*((\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(m+1)}/(\text{f}*(m+1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((m+1)*(a^2 - b^2)) \text{ Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(m+1)}*\text{Simp}[(\text{a}*c - \text{b}*d)*(m+1) - (\text{b}*c - \text{a}*d)*(m+2)*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntegerQ}[2*\text{m}]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.08

method	result
derivativedivides	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} - \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} - \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} - \frac{10287}{4096} + \frac{385}{d} \left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3$
default	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} - \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} - \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} - \frac{10287}{4096} + \frac{385}{d} \left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 - 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3$
risch	$-\frac{-239470 e^{3i(dx+c)} - 86625 i e^{4i(dx+c)} + 218466 i e^{2i(dx+c)} + 10395 e^{5i(dx+c)} + 73575 e^{i(dx+c)} - 8397 i}{12288(3 e^{2i(dx+c)} - 3 - 10 i e^{i(dx+c)})^3} d + \frac{385 i \ln(e^{i(dx+c)})}{32768}$
parallelrisc	$31683960 + 48125i(27 \sin(3dx+3c) - 981 \sin(dx+c) - 270 \cos(2dx+2c) + 770) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3 - 4i\right) + 48125i(-27 \sin(dx+c) - 270 \cos(2dx+2c) + 770)$

input `int(1/(-5+3*sin(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*(250*(39933/5120000*tan(1/2*d*x+1/2*c)^5-672723/25600000*tan(1/2*d*x+1/2*c)^4+2870073/64000000*tan(1/2*d*x+1/2*c)^3-604899/12800000*tan(1/2*d*x+1/2*c)^2+145233/5120000*tan(1/2*d*x+1/2*c)-10287/1024000)/(5*tan(1/2*d*x+1/2*c)^2-6*tan(1/2*d*x+1/2*c)+5)^3+385/16384*arctan(5/4*tan(1/2*d*x+1/2*c)-3/4))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.20

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx = \frac{11196 \cos(dx + c)^3 - 385 (135 \cos(dx + c)^2 - 9 (3 \cos(dx + c)^2 - 28) \sin(dx + c) - 260) \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3}{4}\right)}{32768 (135 d \cos(dx + c)^2 - 9 (3 d \cos(dx + c)^2 - 28 d))}$$

input `integrate(1/(-5+3*sin(d*x+c))^4,x, algorithm="fricas")`

output

```
-1/32768*(11196*cos(d*x + c)^3 - 385*(135*cos(d*x + c)^2 - 9*(3*cos(d*x +
c)^2 - 28)*sin(d*x + c) - 260)*arctan(1/4*(5*sin(d*x + c) - 3)/cos(d*x + c
)) + 42120*cos(d*x + c)*sin(d*x + c) - 52344*cos(d*x + c))/(135*d*cos(d*x
+ c)^2 - 9*(3*d*cos(d*x + c)^2 - 28*d)*sin(d*x + c) - 260*d)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.48 (sec) , antiderivative size = 1690, normalized size of antiderivative = 15.65

$$\int \frac{1}{(-5 + 3\sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(-5+3*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(-5 + 3*sin(2*atan(3/5 - 4*I/5)))**4, Eq(c, -d*x + 2*atan(3/5
- 4*I/5))), (x/(-5 + 3*sin(2*atan(3/5 + 4*I/5)))**4, Eq(c, -d*x + 2*atan(
3/5 + 4*I/5))), (x/(3*sin(c) - 5)**4, Eq(d, 0)), (6015625*(atan(5*tan(c/2
+ d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**6
/(256000000*d*tan(c/2 + d*x/2)**6 - 921600000*d*tan(c/2 + d*x/2)**5 + 1873
920000*d*tan(c/2 + d*x/2)**4 - 2285568000*d*tan(c/2 + d*x/2)**3 + 18739200
00*d*tan(c/2 + d*x/2)**2 - 921600000*d*tan(c/2 + d*x/2) + 256000000*d) - 2
1656250*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/
pi))*tan(c/2 + d*x/2)**5/(256000000*d*tan(c/2 + d*x/2)**6 - 921600000*d*ta
n(c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)**4 - 2285568000*d*tan(c/
2 + d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 - 921600000*d*tan(c/2 + d
*x/2) + 256000000*d) + 44034375*(atan(5*tan(c/2 + d*x/2)/4 - 3/4) + pi*flo
or((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(256000000*d*tan(c/2 + d*
x/2)**6 - 921600000*d*tan(c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)
**4 - 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 -
921600000*d*tan(c/2 + d*x/2) + 256000000*d) - 53707500*(atan(5*tan(c/2 +
d*x/2)/4 - 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**3/(
256000000*d*tan(c/2 + d*x/2)**6 - 921600000*d*tan(c/2 + d*x/2)**5 + 187392
0000*d*tan(c/2 + d*x/2)**4 - 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000
*d*tan(c/2 + d*x/2)**2 - 921600000*d*tan(c/2 + d*x/2) + 256000000*d) + ...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(97) = 194.

Time = 0.11 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.34

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx = \frac{36 \left(\frac{403425 \sin(dx+c)}{\cos(dx+c)+1} - \frac{672110 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{637794 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{373735 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{110925 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - 142875 \right) - 48125 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} \right)}{2048000 d}$$

input `integrate(1/(-5+3*sin(d*x+c))^4,x, algorithm="maxima")`

output

$$\frac{-1/2048000*(36*(403425*\sin(d*x + c)/(\cos(d*x + c) + 1) - 672110*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 637794*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 - 373735*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 + 110925*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 - 142875)/(450*\sin(d*x + c)/(\cos(d*x + c) + 1) - 915*\sin(d*x + c)^2/(\cos(d*x + c) + 1)^2 + 1116*\sin(d*x + c)^3/(\cos(d*x + c) + 1)^3 - 915*\sin(d*x + c)^4/(\cos(d*x + c) + 1)^4 + 450*\sin(d*x + c)^5/(\cos(d*x + c) + 1)^5 - 125*\sin(d*x + c)^6/(\cos(d*x + c) + 1)^6 - 125) - 48125*\arctan(5/4*\sin(d*x + c)/(\cos(d*x + c) + 1) - 3/4))/d}$$
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.37

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx = \frac{48125 dx + 48125 c + \frac{72 \left(110925 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 373735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 637794 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 672110 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 403425 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 142875 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^3}{4096000 d}$$

input `integrate(1/(-5+3*sin(d*x+c))^4,x, algorithm="giac")`

output

```
1/4096000*(48125*d*x + 48125*c + 72*(110925*tan(1/2*d*x + 1/2*c)^5 - 37373
5*tan(1/2*d*x + 1/2*c)^4 + 637794*tan(1/2*d*x + 1/2*c)^3 - 672110*tan(1/2*
d*x + 1/2*c)^2 + 403425*tan(1/2*d*x + 1/2*c) - 142875)/(5*tan(1/2*d*x + 1/
2*c)^2 - 6*tan(1/2*d*x + 1/2*c) + 5)^3 + 96250*arctan((3*cos(d*x + c) - si
n(d*x + c) + 3)/(cos(d*x + c) + 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.73

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{385 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{3}{4}}{4}\right)}{16384 d} - \frac{385 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{16384 d}$$

$$+ \frac{\frac{39933 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{2560000} - \frac{672723 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{12800000} + \frac{2870073 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{32000000} - \frac{604899 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400000} + \frac{145233 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2560000} - \frac{10287}{512000}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{5} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{25} - \frac{1116 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{125} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{25} - \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1 \right)}$$

input

```
int(1/(3*sin(c + d*x) - 5)^4,x)
```

output

```
(385*atan((5*tan(c/2 + (d*x)/2))/4 - 3/4))/(16384*d) - (385*(atan(tan(c/2
+ (d*x)/2)) - (d*x)/2))/(16384*d) + ((145233*tan(c/2 + (d*x)/2))/2560000 -
(604899*tan(c/2 + (d*x)/2)^2)/6400000 + (2870073*tan(c/2 + (d*x)/2)^3)/32
000000 - (672723*tan(c/2 + (d*x)/2)^4)/12800000 + (39933*tan(c/2 + (d*x)/2
)^5)/2560000 - 10287/512000)/(d*((183*tan(c/2 + (d*x)/2)^2)/25 - (18*tan(c
/2 + (d*x)/2))/5 - (1116*tan(c/2 + (d*x)/2)^3)/125 + (183*tan(c/2 + (d*x)/
2)^4)/25 - (18*tan(c/2 + (d*x)/2)^5)/5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.77

$$\int \frac{1}{(-5 + 3 \sin(c + dx))^4} dx$$

$$= \frac{51975 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) \sin(dx+c)^3 - 259875 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) \sin(dx+c)^2 + 433125 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) \sin(dx+c) - 240625 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) - 3}{4}\right) + 27990 \cos(c+dx) \sin(c+dx)^2 - 105300 \cos(c+dx) \sin(c+dx) + 102870 \cos(c+dx) - 12636 \sin(c+dx)^3 + 63180 \sin(c+dx)^2 - 105300 \sin(c+dx) + 58500}{(81920 d (27 \sin(c+dx)^3 - 135 \sin(c+dx)^2 + 225 \sin(c+dx) - 125))}$$

input `int(1/(-5+3*sin(d*x+c))^4,x)`output `(51975*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x)**3 - 259875*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x)**2 + 433125*atan((5*tan((c + d*x)/2) - 3)/4)*sin(c + d*x) - 240625*atan((5*tan((c + d*x)/2) - 3)/4) + 27990*cos(c + d*x)*sin(c + d*x)**2 - 105300*cos(c + d*x)*sin(c + d*x) + 102870*cos(c + d*x) - 12636*sin(c + d*x)**3 + 63180*sin(c + d*x)**2 - 105300*sin(c + d*x) + 58500)/(81920*d*(27*sin(c + d*x)**3 - 135*sin(c + d*x)**2 + 225*sin(c + d*x) - 125))`

3.56 $\int \frac{1}{-5-3\sin(c+dx)} dx$

Optimal result	396
Mathematica [A] (verified)	396
Rubi [A] (verified)	397
Maple [A] (verified)	398
Fricas [A] (verification not implemented)	398
Sympy [B] (verification not implemented)	399
Maxima [A] (verification not implemented)	399
Giac [A] (verification not implemented)	399
Mupad [B] (verification not implemented)	400
Reduce [B] (verification not implemented)	400

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{-5-3\sin(c+dx)} dx = -\frac{x}{4} - \frac{\arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{2d}$$

output `-1/4*x-1/2*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.81

$$\int \frac{1}{-5-3\sin(c+dx)} dx = -\frac{\arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right)}{2d}$$

input `Integrate[(-5 - 3*Sin[c + d*x])^(-1),x]`

output `-1/2*ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])]/d`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 3137}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{-3 \sin(c + dx) - 5} dx$$

↓ 3042

$$\int \frac{1}{-3 \sin(c + dx) - 5} dx$$

↓ 3137

$$-\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} - \frac{x}{4}$$

input `Int[(-5 - 3*Sin[c + d*x])^(-1),x]`

output `-1/4*x - ArcTan[Cos[c + d*x]/(3 + Sin[c + d*x])]/(2*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3137 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[-x/q, x] - Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a - q + b*Sin[c + d*x]))], x] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] & & NegQ[a]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.65

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}}{4}\right)}{2d}$	20
default	$-\frac{\arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}}{4}\right)}{2d}$	20
risch	$\frac{i \ln(e^{i(dx+c)} + \frac{i}{3})}{4d} - \frac{i \ln(e^{i(dx+c)} + 3i)}{4d}$	40
parallelrisc	$\frac{i \left(\ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) - \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 + 4i\right) \right)}{4d}$	42

input `int(1/(-5-3*sin(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/2/d*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{-5 - 3 \sin(c + dx)} dx = -\frac{\arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right)}{4d}$$

input `integrate(1/(-5-3*sin(d*x+c)),x, algorithm="fricas")`

output `-1/4*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(24) = 48$.

Time = 0.38 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{1}{-5 - 3 \sin(c + dx)} dx = \begin{cases} -\frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + \frac{3}{4}}{4}\right) + \pi \left\lfloor \frac{\frac{c}{2} + \frac{dx}{2} - \frac{\pi}{2}}{\pi} \right\rfloor}{2d} & \text{for } d \neq 0 \\ \frac{x}{-3 \sin(c) - 5} & \text{otherwise} \end{cases}$$

input `integrate(1/(-5-3*sin(d*x+c)),x)`

output `Piecewise((-atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(2*d), Ne(d, 0)), (x/(-3*sin(c) - 5), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{1}{-5 - 3 \sin(c + dx)} dx = -\frac{\arctan\left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4}\right)}{2d}$$

input `integrate(1/(-5-3*sin(d*x+c)),x, algorithm="maxima")`

output `-1/2*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4)/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{1}{-5 - 3 \sin(c + dx)} dx = -\frac{dx + c + 2 \arctan\left(\frac{-3 \cos(dx+c) + \sin(dx+c) + 3}{\cos(dx+c) - 3 \sin(dx+c) - 9}\right)}{4d}$$

input `integrate(1/(-5-3*sin(d*x+c)),x, algorithm="giac")`

output

$$-1/4*(d*x + c + 2*\arctan(-(3*\cos(d*x + c) + \sin(d*x + c) + 3)/(\cos(d*x + c) - 3*\sin(d*x + c) - 9)))/d$$
Mupad [B] (verification not implemented)

Time = 25.86 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int \frac{1}{-5 - 3 \sin(c + dx)} dx = \frac{\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}}{2d} - \frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{2d}$$

input

$$\operatorname{int}(-1/(3*\sin(c + d*x) + 5),x)$$

output

$$(\operatorname{atan}(\tan(c/2 + (d*x)/2)) - (d*x)/2)/(2*d) - \operatorname{atan}((5*\tan(c/2 + (d*x)/2))/4 + 3/4)/(2*d)$$
Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.61

$$\int \frac{1}{-5 - 3 \sin(c + dx)} dx = -\frac{\operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right)}{2d}$$

input

$$\operatorname{int}(1/(-5-3*\sin(d*x+c)),x)$$

output

$$(-\operatorname{atan}((5*\tan((c + d*x)/2) + 3)/4))/(2*d)$$

3.57 $\int \frac{1}{(-5-3\sin(c+dx))^2} dx$

Optimal result	401
Mathematica [A] (verified)	401
Rubi [A] (verified)	402
Maple [A] (verified)	403
Fricas [A] (verification not implemented)	404
Sympy [C] (verification not implemented)	405
Maxima [A] (verification not implemented)	405
Giac [A] (verification not implemented)	406
Mupad [B] (verification not implemented)	406
Reduce [B] (verification not implemented)	407

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(-5-3\sin(c+dx))^2} dx = \frac{5x}{64} + \frac{5 \arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{32d} + \frac{3 \cos(c+dx)}{16d(5+3\sin(c+dx))}$$

output

```
5/64*x+5/32*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d+3/16*cos(d*x+c)/d/(5+3*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.62

$$\int \frac{1}{(-5-3\sin(c+dx))^2} dx = \frac{25 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right) + \frac{6(-5+5\cos(c+dx)-3\sin(c+dx))}{5+3\sin(c+dx)}}{160d}$$

input

```
Integrate[(-5 - 3*Sin[c + d*x])^(-2), x]
```

output

```
(25*ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])] + (6*(-5 + 5*Cos[c + d*x] - 3*Sin[c + d*x]))/(5 + 3*Sin[c + d*x]))/(160*d)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 3143, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-3 \sin(c + dx) - 5)^2} dx$$

↓ 3042

$$\int \frac{1}{(-3 \sin(c + dx) - 5)^2} dx$$

↓ 3143

$$\frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} - \frac{1}{16} \int -\frac{5}{3 \sin(c + dx) + 5} dx$$

↓ 27

$$\frac{5}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)}$$

↓ 3042

$$\frac{5}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)}$$

↓ 3136

$$\frac{5}{16} \left(\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4} \right) + \frac{3 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)}$$

input

```
Int[(-5 - 3*Sin[c + d*x])^(-2),x]
```

output

$$\frac{(5*(x/4 + \text{ArcTan}[\text{Cos}[c + d*x]/(3 + \text{Sin}[c + d*x])])/(2*d))}{16} + \frac{(3*\text{Cos}[c + d*x])}{(16*d*(5 + 3*\text{Sin}[c + d*x]))}$$
Defintions of rubi rules used

rule 27

$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_) \text{ ; FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3136

$$\text{Int}[(a_) + (b_)*\text{sin}[(c_) + (d_)*(x_)]^{(-1)}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2 - b^2, 2]\}, \text{Simp}[x/q, x] + \text{Simp}[(2/(d*q))*\text{ArcTan}[b*(\text{Cos}[c + d*x]/(a + q + b*\text{Sin}[c + d*x]))], x]] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{GtQ}[a^2 - b^2, 0] \ \&\& \ \text{PosQ}[a]$$

rule 3143

$$\text{Int}[(a_) + (b_)*\text{sin}[(c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^{(n + 1)})/(d*(n + 1)*(a^2 - b^2)), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\text{Sin}[c + d*x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$$
Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
default	$\frac{\frac{9 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{40}}{200} + \frac{5 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + \frac{3}{4}\right)}{32}}{\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + \frac{6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{5} + 1} d$
risch	$\frac{5 e^{i(dx+c)} + 3i}{8d(3 e^{2i(dx+c)} - 3 + 10i e^{i(dx+c)})} - \frac{5i \ln(e^{i(dx+c)} + \frac{i}{3})}{64d} + \frac{5i \ln(e^{i(dx+c)} + 3i)}{64d}$
parallelrisch	$\frac{(-125i - 75i \sin(dx+c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) + (75i \sin(dx+c) + 125i) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 + 4i\right) + 60 \cos(dx+c) + 3}{960 \sin(dx+c)d + 1600d}$

input `int(1/(-5-3*sin(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(2*(9/400*tan(1/2*d*x+1/2*c)+3/80)/(tan(1/2*d*x+1/2*c)^2+6/5*tan(1/2*d*x+1/2*c)+1)+5/32*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.05

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx$$

$$= \frac{5(3 \sin(dx + c) + 5) \arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right) + 12 \cos(dx + c)}{64(3d \sin(dx + c) + 5d)}$$

input `integrate(1/(-5-3*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/64*(5*(3*sin(d*x + c) + 5)*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c)) + 12*cos(d*x + c))/(3*d*sin(d*x + c) + 5*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.91 (sec) , antiderivative size = 389, normalized size of antiderivative = 6.95

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(-5-3*sin(d*x+c))**2,x)`

output `Piecewise((x/(-5 + 3*sin(2*atan(3/5 - 4*I/5)))**2, Eq(c, -d*x - 2*atan(3/5 - 4*I/5))), (x/(-5 + 3*sin(2*atan(3/5 + 4*I/5)))**2, Eq(c, -d*x - 2*atan(3/5 + 4*I/5))), (x/(-3*sin(c) - 5)**2, Eq(d, 0)), (125*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**2/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 150*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 125*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 36*tan(c/2 + d*x/2)/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d) + 60/(800*d*tan(c/2 + d*x/2)**2 + 960*d*tan(c/2 + d*x/2) + 800*d), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.66

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx = \frac{12 \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 5 \right)}{\frac{6 \sin(dx+c)}{\cos(dx+c)+1} + \frac{5 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 5} + 25 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4} \right) \frac{1}{160 d}$$

input `integrate(1/(-5-3*sin(d*x+c))^2,x, algorithm="maxima")`

output `1/160*(12*(3*sin(d*x + c)/(cos(d*x + c) + 1) + 5)/(6*sin(d*x + c)/(cos(d*x + c) + 1) + 5*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 5) + 25*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4))/d`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.70

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx$$

$$= \frac{25 dx + 25 c + \frac{24 (3 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5)}{5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 6 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5} + 50 \arctan\left(-\frac{3 \cos(dx+c) + \sin(dx+c) + 3}{\cos(dx+c) - 3 \sin(dx+c) - 9}\right)}{320 d}$$

input `integrate(1/(-5-3*sin(d*x+c))^2,x, algorithm="giac")`

output

$$\frac{1}{320} \cdot \frac{(25 dx + 25 c + 24 (3 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5) / (5 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + 6 \tan(\frac{1}{2} dx + \frac{1}{2} c) + 5) + 50 \arctan(-\frac{3 \cos(dx+c) + \sin(dx+c) + 3}{\cos(dx+c) - 3 \sin(dx+c) - 9}))}{d}$$
Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.48

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx = \frac{5 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{32 d} - \frac{5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{32 d}$$

$$+ \frac{\frac{9 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{200} + \frac{3}{40}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \frac{6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1\right)}$$

input `int(1/(3*sin(c + d*x) + 5)^2,x)`

output

$$\frac{(5 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{d x}{2}\right)}{4} + \frac{3}{4}\right))}{(32 d)} - \frac{(5 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{d x}{2}\right)\right) - \frac{d x}{2}\right))}{(32 d)} + \frac{\left(\frac{9 \tan\left(\frac{c}{2} + \frac{d x}{2}\right)}{200} + \frac{3}{40}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{d x}{2}\right)^2 + \frac{6 \tan\left(\frac{c}{2} + \frac{d x}{2}\right)}{5} + 1\right)}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.14

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^2} dx$$

$$= \frac{15 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) + 3}{4}\right) \sin(dx+c) + 25 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx+c}{2}\right) + 3}{4}\right) + 6 \cos(dx+c)}{32d(3 \sin(dx+c) + 5)}$$

input

```
int(1/(-5-3*sin(d*x+c))^2,x)
```

output

```
(15*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x) + 25*atan((5*tan((c + d*x)/2) + 3)/4) + 6*cos(c + d*x))/(32*d*(3*sin(c + d*x) + 5))
```


3.58 $\int \frac{1}{(-5-3\sin(c+dx))^3} dx$

Optimal result	408
Mathematica [A] (verified)	408
Rubi [A] (verified)	409
Maple [A] (verified)	411
Fricas [A] (verification not implemented)	412
Sympy [C] (verification not implemented)	412
Maxima [B] (verification not implemented)	413
Giac [A] (verification not implemented)	414
Mupad [B] (verification not implemented)	414
Reduce [B] (verification not implemented)	415

Optimal result

Integrand size = 12, antiderivative size = 81

$$\int \frac{1}{(-5-3\sin(c+dx))^3} dx = -\frac{59x}{2048} - \frac{59 \arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{1024d} - \frac{3 \cos(c+dx)}{32d(5+3\sin(c+dx))^2} - \frac{45 \cos(c+dx)}{512d(5+3\sin(c+dx))}$$

output `-59/2048*x-59/1024*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d-3/32*cos(d*x+c)/d/(5+3*sin(d*x+c))^2-45/512*cos(d*x+c)/d/(5+3*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.41

$$\int \frac{1}{(-5-3\sin(c+dx))^3} dx = \frac{-59 \arctan\left(\frac{2\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right) + \frac{3(-182 \cos(c+dx)+3(59-9 \cos(2(c+dx))+60 \sin(c+dx)-15 \sin(2(c+dx))))}{(5+3 \sin(c+dx))^2}}{1024d}$$

input `Integrate[(-5 - 3*Sin[c + d*x])^(-3), x]`

output

$$\frac{(-59 \operatorname{ArcTan}[(2 \cos((c + dx)/2) + \sin((c + dx)/2)) / (\cos((c + dx)/2) - \sin((c + dx)/2))] + (3(-182 \cos[c + dx] + 3(59 - 9 \cos[2(c + dx)] + 60 \sin[c + dx] - 15 \sin[2(c + dx)]))) / (5 + 3 \sin[c + dx])^2}{1024d}$$
Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(-3 \sin(c + dx) - 5)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(-3 \sin(c + dx) - 5)^3} dx \\ & \quad \downarrow \text{3143} \\ & -\frac{1}{32} \int \frac{10 - 3 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx - \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{3042} \\ & -\frac{1}{32} \int \frac{10 - 3 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx - \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{3233} \\ & \frac{1}{32} \left(\frac{1}{16} \int -\frac{59}{3 \sin(c + dx) + 5} dx - \frac{45 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) - \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{27} \\ & \frac{1}{32} \left(-\frac{59}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx - \frac{45 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) - \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{32} \left(-\frac{59}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx - \frac{45 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) - \frac{3 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \end{aligned}$$

$$\frac{1}{32} \left(-\frac{59}{16} \left(\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4} \right) - \frac{45 \cos(c+dx)}{16d(3 \sin(c+dx)+5)} \right) - \frac{3 \cos(c+dx)}{32d(3 \sin(c+dx)+5)^2}$$

↓ 3136

input `Int[(-5 - 3*Sin[c + d*x])^(-3),x]`

output `(-3*Cos[c + d*x])/(32*d*(5 + 3*Sin[c + d*x])^2) + ((-59*(x/4 + ArcTan[Cos[c + d*x]/(3 + Sin[c + d*x]])/(2*d)))/16 - (45*Cos[c + d*x])/(16*d*(5 + 3*Sin[c + d*x]))) / 32`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3136 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{q = Rt[a^2 - b^2, 2]}, Simp[x/q, x] + Simp[(2/(d*q))*ArcTan[b*(Cos[c + d*x]/(a + q + b*Sin[c + d*x]))], x]] /; FreeQ[{a, b, c, d}, x] && GtQ[a^2 - b^2, 0] && PosQ[a]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e +
f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2))
Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(
m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{50 \left(\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{64000} + \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{320000} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64000} + \frac{273}{12800} \right) - 59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right)}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^2} \frac{1}{d}$
default	$\frac{50 \left(\frac{963 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{64000} + \frac{11739 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{320000} + \frac{2313 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{64000} + \frac{273}{12800} \right) - 59 \arctan\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right)}{\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^2} \frac{1}{d}$
risch	$-\frac{3(295ie^{2i(dx+c)} + 59e^{3i(dx+c)} - 241e^{i(dx+c)} - 45i)}{256(3e^{2i(dx+c)} - 3 + 10ie^{i(dx+c)})^2} d - \frac{59i \ln(e^{i(dx+c)} + 3i)}{2048d} + \frac{59i \ln(e^{i(dx+c)} + \frac{i}{3})}{2048d}$
parallelrisc	$\frac{1062 + 59i(-59 + 9 \cos(2dx + 2c) - 60 \sin(dx + c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) + 59i(59 - 9 \cos(2dx + 2c) + 60 \sin(dx + c)) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 + 4i\right)}{2048d(-59 + 9 \cos(2dx + 2c) - 60 \sin(dx + c))}$

input

```
int(1/(-5-3*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-50*(963/64000*tan(1/2*d*x+1/2*c)^3+11739/320000*tan(1/2*d*x+1/2*c)^2
+2313/64000*tan(1/2*d*x+1/2*c)+273/12800)/(5*tan(1/2*d*x+1/2*c)^2+6*tan(1/
2*d*x+1/2*c)+5)^2-59/1024*arctan(5/4*tan(1/2*d*x+1/2*c)+3/4))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.16

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx =$$

$$\frac{59 (9 \cos(dx + c)^2 - 30 \sin(dx + c) - 34) \arctan\left(\frac{5 \sin(dx+c)+3}{4 \cos(dx+c)}\right) - 540 \cos(dx + c) \sin(dx + c) - 1092 \cos(dx + c)}{2048 (9 d \cos(dx + c)^2 - 30 d \sin(dx + c) - 34 d)}$$

input `integrate(1/(-5-3*sin(d*x+c))^3,x, algorithm="fricas")`

output `-1/2048*(59*(9*cos(d*x + c)^2 - 30*sin(d*x + c) - 34)*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c)) - 540*cos(d*x + c)*sin(d*x + c) - 1092*cos(d*x + c))/(9*d*cos(d*x + c)^2 - 30*d*sin(d*x + c) - 34*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.96 (sec) , antiderivative size = 921, normalized size of antiderivative = 11.37

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx = \text{Too large to display}$$

input `integrate(1/(-5-3*sin(d*x+c))**3,x)`

output

```
Piecewise((x/(-5 + 3*sin(2*atan(3/5 - 4*I/5)))**3, Eq(c, -d*x - 2*atan(3/5
- 4*I/5))), (x/(-5 + 3*sin(2*atan(3/5 + 4*I/5)))**3, Eq(c, -d*x - 2*atan(
3/5 + 4*I/5))), (x/(-3*sin(c) - 5)**3, Eq(d, 0)), (-36875*(atan(5*tan(c/2
+ d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4
/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d
*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d) - 88500*(ata
n(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2
+ d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/2 + d*x/2)**3
+ 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2) + 640000*d)
- 126850*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)
/pi))*tan(c/2 + d*x/2)**2/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c/
2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2)
+ 640000*d) - 88500*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d
*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)/(640000*d*tan(c/2 + d*x/2)**4 + 1536000
*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2
+ d*x/2) + 640000*d) - 36875*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor
((c/2 + d*x/2 - pi/2)/pi))/(640000*d*tan(c/2 + d*x/2)**4 + 1536000*d*tan(c
/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000*d*tan(c/2 + d*x/2)
) + 640000*d) - 19260*tan(c/2 + d*x/2)**3/(640000*d*tan(c/2 + d*x/2)**4 +
1536000*d*tan(c/2 + d*x/2)**3 + 2201600*d*tan(c/2 + d*x/2)**2 + 1536000...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. $2(73) = 146$.

Time = 0.11 (sec) , antiderivative size = 173, normalized size of antiderivative = 2.14

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx = \frac{12 \left(\frac{3855 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3913 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1605 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + 2275 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} + \frac{86 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{25 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 25} + 1475 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} + \frac{3}{4} \right)$$

25600 d

input

```
integrate(1/(-5-3*sin(d*x+c))^3,x, algorithm="maxima")
```

output

```
-1/25600*(12*(3855*sin(d*x + c)/(cos(d*x + c) + 1) + 3913*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1605*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 2275)/(60*sin(d*x + c)/(cos(d*x + c) + 1) + 86*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 25*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 25) + 1475*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.49

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx = \frac{1475 dx + 1475 c + \frac{24 \left(1605 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 3913 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 3855 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2275 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^2} + 2950 \arctan\left(-\frac{3 \cos(dx + c)}{\cos(dx + c) - 3 \sin(dx + c) - 9}\right)}{51200 d}$$

input

```
integrate(1/(-5-3*sin(d*x+c))^3,x, algorithm="giac")
```

output

```
-1/51200*(1475*d*x + 1475*c + 24*(1605*tan(1/2*d*x + 1/2*c)^3 + 3913*tan(1/2*d*x + 1/2*c)^2 + 3855*tan(1/2*d*x + 1/2*c) + 2275)/(5*tan(1/2*d*x + 1/2*c)^2 + 6*tan(1/2*d*x + 1/2*c) + 5)^2 + 2950*arctan(-(3*cos(d*x + c) + sin(d*x + c) + 3)/(cos(d*x + c) - 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 26.05 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.38

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx = \frac{59 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2} \right)}{1024 d} - \frac{59 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{1024 d} - \frac{\frac{963 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{1280} + \frac{11739 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400} + \frac{2313 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{1280} + \frac{273}{256}}{d \left(5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + 6 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 5 \right)^2}$$

input `int(-1/(3*sin(c + d*x) + 5)^3,x)`

output `(59*(atan(tan(c/2 + (d*x)/2)) - (d*x)/2))/(1024*d) - (59*atan((5*tan(c/2 + (d*x)/2))/4 + 3/4))/(1024*d) - ((2313*tan(c/2 + (d*x)/2))/1280 + (11739*atan(c/2 + (d*x)/2)^2)/6400 + (963*tan(c/2 + (d*x)/2)^3)/1280 + 273/256)/(d*(6*tan(c/2 + (d*x)/2) + 5*tan(c/2 + (d*x)/2)^2 + 5)^2)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.62

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^3} dx$$

$$= \frac{-531 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c)^2 - 1770 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c) - 1475 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right)}{1024d (9 \sin(dx + c))^2}$$

input `int(1/(-5-3*sin(d*x+c))^3,x)`

output `(- 531*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x)**2 - 1770*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x) - 1475*atan((5*tan((c + d*x)/2) + 3)/4) - 270*cos(c + d*x)*sin(c + d*x) - 546*cos(c + d*x) - 81*sin(c + d*x)**2 - 270*sin(c + d*x) - 225)/(1024*d*(9*sin(c + d*x)**2 + 30*sin(c + d*x) + 25))`

3.59 $\int \frac{1}{(-5-3\sin(c+dx))^4} dx$

Optimal result	416
Mathematica [A] (verified)	416
Rubi [A] (verified)	417
Maple [A] (verified)	420
Fricas [A] (verification not implemented)	420
Sympy [C] (verification not implemented)	421
Maxima [B] (verification not implemented)	422
Giac [A] (verification not implemented)	422
Mupad [B] (verification not implemented)	423
Reduce [B] (verification not implemented)	424

Optimal result

Integrand size = 12, antiderivative size = 106

$$\int \frac{1}{(-5-3\sin(c+dx))^4} dx = \frac{385x}{32768} + \frac{385 \arctan\left(\frac{\cos(c+dx)}{3+\sin(c+dx)}\right)}{16384d} + \frac{\cos(c+dx)}{16d(5+3\sin(c+dx))^3} + \frac{25\cos(c+dx)}{512d(5+3\sin(c+dx))^2} + \frac{311\cos(c+dx)}{8192d(5+3\sin(c+dx))}$$

output `385/32768*x+385/16384*arctan(cos(d*x+c)/(3+sin(d*x+c)))/d+1/16*cos(d*x+c)/d/(5+3*sin(d*x+c))^3+25/512*cos(d*x+c)/d/(5+3*sin(d*x+c))^2+311/8192*cos(d*x+c)/d/(5+3*sin(d*x+c))`

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.25

$$\int \frac{1}{(-5-3\sin(c+dx))^4} dx = \frac{1925 \arctan\left(\frac{2(\cos(\frac{1}{2}(c+dx))+\sin(\frac{1}{2}(c+dx)))}{\cos(\frac{1}{2}(c+dx))-\sin(\frac{1}{2}(c+dx))}\right) + \frac{-239470+219735\cos(c+dx)+83970\cos(2(c+dx))-13995\cos(3(c+dx))-305091\sin(c+dx)}{2(5+3\sin(c+dx))^3}}{81920d}$$

input `Integrate[(-5 - 3*Sin[c + d*x])^(-4), x]`

output

```
(1925*ArcTan[(2*(Cos[(c + d*x)/2] + Sin[(c + d*x)/2]))/(Cos[(c + d*x)/2] - Sin[(c + d*x)/2])] + (-239470 + 219735*Cos[c + d*x] + 83970*Cos[2*(c + d*x)] - 13995*Cos[3*(c + d*x)] - 305091*Sin[c + d*x] + 105300*Sin[2*(c + d*x)] + 8397*Sin[3*(c + d*x)])/(2*(5 + 3*Sin[c + d*x])^3)/(81920*d)
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {3042, 3143, 27, 3042, 3233, 25, 3042, 3233, 27, 3042, 3136}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-3 \sin(c + dx) - 5)^4} dx$$

↓ 3042

$$\int \frac{1}{(-3 \sin(c + dx) - 5)^4} dx$$

↓ 3143

$$\frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} - \frac{1}{48} \int -\frac{3(5 - 2 \sin(c + dx))}{(3 \sin(c + dx) + 5)^3} dx$$

↓ 27

$$\frac{1}{16} \int \frac{5 - 2 \sin(c + dx)}{(3 \sin(c + dx) + 5)^3} dx + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

↓ 3042

$$\frac{1}{16} \int \frac{5 - 2 \sin(c + dx)}{(3 \sin(c + dx) + 5)^3} dx + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

↓ 3233

$$\frac{1}{16} \left(\frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} - \frac{1}{32} \int -\frac{62 - 25 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx \right) + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

↓ 25

$$\frac{1}{16} \left(\frac{1}{32} \int \frac{62 - 25 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \int \frac{62 - 25 \sin(c + dx)}{(3 \sin(c + dx) + 5)^2} dx + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 3233 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} - \frac{1}{16} \int -\frac{385}{3 \sin(c + dx) + 5} dx \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 27 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 3042 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \int \frac{1}{3 \sin(c + dx) + 5} dx + \frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3} \\
& \downarrow 3136 \\
& \frac{1}{16} \left(\frac{1}{32} \left(\frac{385}{16} \left(\frac{\arctan\left(\frac{\cos(c+dx)}{\sin(c+dx)+3}\right)}{2d} + \frac{x}{4} \right) + \frac{311 \cos(c + dx)}{16d(3 \sin(c + dx) + 5)} \right) + \frac{25 \cos(c + dx)}{32d(3 \sin(c + dx) + 5)^2} \right) + \\
& \quad \frac{\cos(c + dx)}{16d(3 \sin(c + dx) + 5)^3}
\end{aligned}$$

input `Int[(-5 - 3*Sin[c + d*x])^(-4),x]`

output `Cos[c + d*x]/(16*d*(5 + 3*Sin[c + d*x])^3) + ((25*Cos[c + d*x])/(32*d*(5 + 3*Sin[c + d*x])^2) + ((385*(x/4 + ArcTan[Cos[c + d*x]/(3 + Sin[c + d*x]))/(2*d)))/16 + (311*Cos[c + d*x])/(16*d*(5 + 3*Sin[c + d*x]))/32)/16`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \text{ Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3136 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(-1)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = \text{Rt}[\text{a}^2 - \text{b}^2, 2]\}, \text{Simp}[\text{x}/\text{q}, \text{x}] + \text{Simp}[(2/(d*q))*\text{ArcTan}[\text{b}*(\text{Cos}[\text{c} + \text{d}*x]/(\text{a} + \text{q} + \text{b}*\text{Sin}[\text{c} + \text{d}*x]))], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{GtQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{PosQ}[\text{a}]$
- rule 3143 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(n_)}, \text{x_Symbol}] \rightarrow \text{Simp}[-\text{b})*\text{Cos}[\text{c} + \text{d}*x]*((\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(n+1)}/(\text{d}*(n+1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((n+1)*(a^2 - b^2)) \text{ Int}[(\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(n+1)}*\text{Simp}[\text{a}*(n+1) - \text{b}*(n+2)*\text{Sin}[\text{c} + \text{d}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{n}, -1] \ \&\& \ \text{IntegerQ}[2*\text{n}]$
- rule 3233 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]^{(m_)}*((\text{c}_.) + (\text{d}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[-(\text{b}*c - \text{a}*d)*\text{Cos}[\text{e} + \text{f}*x]*((\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(m+1)}/(\text{f}*(m+1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((m+1)*(a^2 - b^2)) \text{ Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(m+1)}*\text{Simp}[(\text{a}*c - \text{b}*d)*(m+1) - (\text{b}*c - \text{a}*d)*(m+2)*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntegerQ}[2*\text{m}]$

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} + \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} + \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} + \frac{10287}{4096} + \frac{385}{d} \left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3$
default	$\frac{\frac{39933 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{20480} + \frac{672723 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{102400} + \frac{2870073 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{256000} + \frac{604899 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{51200} + \frac{145233 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{20480} + \frac{10287}{4096} + \frac{385}{d} \left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 6 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 5\right)^3$
risch	$\frac{-239470 e^{3i(dx+c)} + 86625ie^{4i(dx+c)} - 218466ie^{2i(dx+c)} + 10395 e^{5i(dx+c)} + 73575 e^{i(dx+c)} + 8397i}{12288(3 e^{2i(dx+c)} - 3 + 10ie^{i(dx+c)})^3} d - \frac{385i \ln(e^{i(dx+c)})}{32768d}$
parallelrisch	$-31683960 + 48125i(-27 \sin(3dx+3c) + 981 \sin(dx+c) - 270 \cos(2dx+2c) + 770) \ln\left(5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3 - 4i\right) + 48125i(-7$

input `int(1/(-5-3*sin(d*x+c))^4,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{250 \cdot (39933/5120000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c))^5 + 672723/25600000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^4 + 2870073/64000000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^3 + 604899/12800000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 145233/5120000 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + 10287/1024000}{(5 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c)^2 + 6 \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + 5)^3} + \frac{385}{16384} \cdot \arctan\left(\frac{5}{4} \cdot \tan(1/2 \cdot d \cdot x + 1/2 \cdot c) + \frac{3}{4}\right) \right)$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.23

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{11196 \cos(dx + c)^3 + 385 (135 \cos(dx + c)^2 + 9 (3 \cos(dx + c)^2 - 28) \sin(dx + c) - 260) \arctan\left(\frac{5 \sin(dx + c)}{4 \cos(dx + c)}\right)}{32768 (135 d \cos(dx + c)^2 + 9 (3 d \cos(dx + c)^2 - 28 d) \sin(dx + c) - 260 d)}$$

input `integrate(1/(-5-3*sin(d*x+c))^4,x, algorithm="fricas")`

output

```
1/32768*(11196*cos(d*x + c)^3 + 385*(135*cos(d*x + c)^2 + 9*(3*cos(d*x + c)
)^2 - 28)*sin(d*x + c) - 260)*arctan(1/4*(5*sin(d*x + c) + 3)/cos(d*x + c)
) - 42120*cos(d*x + c)*sin(d*x + c) - 52344*cos(d*x + c))/(135*d*cos(d*x +
c)^2 + 9*(3*d*cos(d*x + c)^2 - 28*d)*sin(d*x + c) - 260*d)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.50 (sec) , antiderivative size = 1695, normalized size of antiderivative = 15.99

$$\int \frac{1}{(-5 - 3\sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(-5-3*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(-5 + 3*sin(2*atan(3/5 - 4*I/5)))**4, Eq(c, -d*x - 2*atan(3/5
- 4*I/5))), (x/(-5 + 3*sin(2*atan(3/5 + 4*I/5)))**4, Eq(c, -d*x - 2*atan(
3/5 + 4*I/5))), (x/(-3*sin(c) - 5)**4, Eq(d, 0)), (6015625*(atan(5*tan(c/2
+ d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**
6/(256000000*d*tan(c/2 + d*x/2)**6 + 921600000*d*tan(c/2 + d*x/2)**5 + 187
3920000*d*tan(c/2 + d*x/2)**4 + 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920
000*d*tan(c/2 + d*x/2)**2 + 921600000*d*tan(c/2 + d*x/2) + 256000000*d) +
21656250*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)
/pi))*tan(c/2 + d*x/2)**5/(256000000*d*tan(c/2 + d*x/2)**6 + 921600000*d*t
an(c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)**4 + 2285568000*d*tan(
c/2 + d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2 + 921600000*d*tan(c/2 +
d*x/2) + 256000000*d) + 44034375*(atan(5*tan(c/2 + d*x/2)/4 + 3/4) + pi*fl
oor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**4/(256000000*d*tan(c/2 + d
*x/2)**6 + 921600000*d*tan(c/2 + d*x/2)**5 + 1873920000*d*tan(c/2 + d*x/2)
**4 + 2285568000*d*tan(c/2 + d*x/2)**3 + 1873920000*d*tan(c/2 + d*x/2)**2
+ 921600000*d*tan(c/2 + d*x/2) + 256000000*d) + 53707500*(atan(5*tan(c/2 +
d*x/2)/4 + 3/4) + pi*floor((c/2 + d*x/2 - pi/2)/pi))*tan(c/2 + d*x/2)**3/
(256000000*d*tan(c/2 + d*x/2)**6 + 921600000*d*tan(c/2 + d*x/2)**5 + 18739
20000*d*tan(c/2 + d*x/2)**4 + 2285568000*d*tan(c/2 + d*x/2)**3 + 187392000
0*d*tan(c/2 + d*x/2)**2 + 921600000*d*tan(c/2 + d*x/2) + 256000000*d) + ...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. $2(96) = 192$.

Time = 0.13 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.39

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{36 \left(\frac{403425 \sin(dx+c)}{\cos(dx+c)+1} + \frac{672110 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{637794 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{373735 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{110925 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + 142875 \right) + 48125 \arctan \left(\frac{5 \sin(dx+c)}{4(\cos(dx+c)+1)} \right)}{2048000 d}$$

input `integrate(1/(-5-3*sin(d*x+c))^4,x, algorithm="maxima")`

output

```
1/2048000*(36*(403425*sin(d*x + c)/(cos(d*x + c) + 1) + 672110*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 637794*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 373735*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 110925*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 142875)/(450*sin(d*x + c)/(cos(d*x + c) + 1) + 915*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1116*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 915*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 450*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 125*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 125) + 48125*arctan(5/4*sin(d*x + c)/(cos(d*x + c) + 1) + 3/4))/d
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.39

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{48125 dx + 48125 c + \frac{72 \left(110925 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 373735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 637794 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 672110 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 403425 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 142875 \right)}{\left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 6 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 5 \right)^3}{4096000 d}$$

input `integrate(1/(-5-3*sin(d*x+c))^4,x, algorithm="giac")`

output

```
1/4096000*(48125*d*x + 48125*c + 72*(110925*tan(1/2*d*x + 1/2*c)^5 + 37373
5*tan(1/2*d*x + 1/2*c)^4 + 637794*tan(1/2*d*x + 1/2*c)^3 + 672110*tan(1/2*
d*x + 1/2*c)^2 + 403425*tan(1/2*d*x + 1/2*c) + 142875)/(5*tan(1/2*d*x + 1/
2*c)^2 + 6*tan(1/2*d*x + 1/2*c) + 5)^3 + 96250*arctan(-(3*cos(d*x + c) + s
in(d*x + c) + 3)/(cos(d*x + c) - 3*sin(d*x + c) - 9)))/d
```

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.76

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{385 \operatorname{atan}\left(\frac{5 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{3}{4}\right)}{16384 d} - \frac{385 \left(\operatorname{atan}\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)\right) - \frac{dx}{2}\right)}{16384 d}$$

$$+ \frac{\frac{39933 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{2560000} + \frac{672723 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{12800000} + \frac{2870073 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{32000000} + \frac{604899 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{6400000} + \frac{145233 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{2560000} + \frac{10287}{512000}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{5} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{25} + \frac{1116 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{125} + \frac{183 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{25} + \frac{18 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{5} + 1 \right)}$$

input

```
int(1/(3*sin(c + d*x) + 5)^4,x)
```

output

```
(385*atan((5*tan(c/2 + (d*x)/2))/4 + 3/4))/(16384*d) - (385*(atan(tan(c/2
+ (d*x)/2)) - (d*x)/2))/(16384*d) + ((145233*tan(c/2 + (d*x)/2))/2560000 +
(604899*tan(c/2 + (d*x)/2)^2)/6400000 + (2870073*tan(c/2 + (d*x)/2)^3)/32
000000 + (672723*tan(c/2 + (d*x)/2)^4)/12800000 + (39933*tan(c/2 + (d*x)/2
)^5)/2560000 + 10287/512000)/(d*((18*tan(c/2 + (d*x)/2))/5 + (183*tan(c/2
+ (d*x)/2)^2)/25 + (1116*tan(c/2 + (d*x)/2)^3)/125 + (183*tan(c/2 + (d*x)/
2)^4)/25 + (18*tan(c/2 + (d*x)/2)^5)/5 + tan(c/2 + (d*x)/2)^6 + 1))
```


Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.80

$$\int \frac{1}{(-5 - 3 \sin(c + dx))^4} dx$$

$$= \frac{51975 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c)^3 + 259875 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c)^2 + 433125 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) \sin(dx + c) + 240625 \operatorname{atan}\left(\frac{5 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{4} + \frac{3}{4}\right) + 27990 \cos(c + dx) \sin(c + dx)^2 + 105300 \cos(c + dx) \sin(c + dx) + 102870 \cos(c + dx) + 12636 \sin(c + dx)^3 + 63180 \sin(c + dx)^2 + 105300 \sin(c + dx) + 58500}{(81920 d (27 \sin(c + dx)^3 + 135 \sin(c + dx)^2 + 225 \sin(c + dx) + 125))}$$

input `int(1/(-5-3*sin(d*x+c))^4,x)`output `(51975*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x)**3 + 259875*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x)**2 + 433125*atan((5*tan((c + d*x)/2) + 3)/4)*sin(c + d*x) + 240625*atan((5*tan((c + d*x)/2) + 3)/4) + 27990*cos(c + d*x)*sin(c + d*x)**2 + 105300*cos(c + d*x)*sin(c + d*x) + 102870*cos(c + d*x) + 12636*sin(c + d*x)**3 + 63180*sin(c + d*x)**2 + 105300*sin(c + d*x) + 58500)/(81920*d*(27*sin(c + d*x)**3 + 135*sin(c + d*x)**2 + 225*sin(c + d*x) + 125))`

3.60 $\int \frac{1}{3+5 \sin(c+dx)} dx$

Optimal result	425
Mathematica [A] (verified)	425
Rubi [A] (verified)	426
Maple [A] (verified)	427
Fricas [A] (verification not implemented)	428
Sympy [A] (verification not implemented)	428
Maxima [A] (verification not implemented)	428
Giac [A] (verification not implemented)	429
Mupad [B] (verification not implemented)	429
Reduce [B] (verification not implemented)	430

Optimal result

Integrand size = 12, antiderivative size = 63

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = -\frac{\log \left(3 \cos \left(\frac{1}{2}(c + dx) \right) + \sin \left(\frac{1}{2}(c + dx) \right) \right)}{4d} + \frac{\log \left(\cos \left(\frac{1}{2}(c + dx) \right) + 3 \sin \left(\frac{1}{2}(c + dx) \right) \right)}{4d}$$

output

```
-1/4*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+1/4*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = -\frac{\log \left(3 \cos \left(\frac{1}{2}(c + dx) \right) + \sin \left(\frac{1}{2}(c + dx) \right) \right)}{4d} + \frac{\log \left(\cos \left(\frac{1}{2}(c + dx) \right) + 3 \sin \left(\frac{1}{2}(c + dx) \right) \right)}{4d}$$

input

```
Integrate[(3 + 5*Sin[c + d*x])^(-1),x]
```

output

$$-1/4*\text{Log}[3*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]]/d + \text{Log}[\text{Cos}[(c + d*x)/2] + 3*\text{Sin}[(c + d*x)/2]]/(4*d)$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{5 \sin(c + dx) + 3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{5 \sin(c + dx) + 3} dx \\ & \quad \downarrow \text{3139} \\ & \frac{2 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) + 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{1081} \\ & \frac{6 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx)) + 1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx)) + 3)} \right) d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{6 \left(\frac{1}{24} \log(3 \tan(\frac{1}{2}(c + dx)) + 1) - \frac{1}{24} \log(\tan(\frac{1}{2}(c + dx)) + 3) \right)}{d} \end{aligned}$$

input

$$\text{Int}[(3 + 5*\text{Sin}[c + d*x])^{-1}, x]$$

output

$$(6*(-1/24*\text{Log}[3 + \text{Tan}[(c + d*x)/2]] + \text{Log}[1 + 3*\text{Tan}[(c + d*x)/2]]/24))/d$$

Definitions of rubi rules used

rule 1081 $\text{Int}[\{(a_)+ (b_)*(x_)+ (c_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c \text{ Int}[\text{ExpandIntegrand}[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x]] \text{ /; FreeQ}\{a, b, c\}, x] \ \&\& \ \text{NiceSqrtQ}[b^2 - 4*a*c]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3139 $\text{Int}[\{(a_)+ (b_)*\sin[(c_)+ (d_)*(x_)]\}^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] \text{ /; FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.57

method	result	size
derivativdivides	$\frac{\frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{4}}{d}$	36
default	$\frac{\frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4} - \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{4}}{d}$	36
parallelrisc	$\frac{-\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 9\right) + \ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d}$	37
norman	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{4d} + \frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d}$	38
risc	$-\frac{\ln\left(\frac{4}{5} + \frac{3i}{5} + e^{i(dx+c)}\right)}{4d} + \frac{\ln\left(e^{i(dx+c)} - \frac{4}{5} + \frac{3i}{5}\right)}{4d}$	40

input $\text{int}(1/(3+5*\sin(d*x+c)), x, \text{method}=_RETURNVERBOSE)$

output $1/d*(1/4*\ln(3*\tan(1/2*d*x+1/2*c)+1)-1/4*\ln(\tan(1/2*d*x+1/2*c)+3))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = \frac{\log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5)}{8d}$$

input `integrate(1/(3+5*sin(d*x+c)),x, algorithm="fricas")`output `-1/8*(log(4*cos(d*x + c) + 3*sin(d*x + c) + 5) - log(-4*cos(d*x + c) + 3*sin(d*x + c) + 5))/d`**Sympy [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = \begin{cases} -\frac{\log(\tan(\frac{c}{2} + \frac{dx}{2}) + 3)}{4d} + \frac{\log(3 \tan(\frac{c}{2} + \frac{dx}{2}) + 1)}{4d} & \text{for } d \neq 0 \\ \frac{x}{5 \sin(c) + 3} & \text{otherwise} \end{cases}$$

input `integrate(1/(3+5*sin(d*x+c)),x)`output `Piecewise((-log(tan(c/2 + d*x/2) + 3)/(4*d) + log(3*tan(c/2 + d*x/2) + 1)/(4*d), Ne(d, 0)), (x/(5*sin(c) + 3), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = \frac{\log\left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1\right) - \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 3\right)}{4d}$$

input `integrate(1/(3+5*sin(d*x+c)),x, algorithm="maxima")`

output $\frac{1}{4} * (\log(3 * \sin(dx + c) / (\cos(dx + c) + 1) + 1) - \log(\sin(dx + c) / (\cos(dx + c) + 1) + 3)) / d$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.57

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = \frac{\log\left(\left|3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3\right|\right)}{4d}$$

input `integrate(1/(3+5*sin(d*x+c)),x, algorithm="giac")`

output $\frac{1}{4} * (\log(\text{abs}(3 * \tan(1/2 * dx + 1/2 * c) + 1)) - \log(\text{abs}(\tan(1/2 * dx + 1/2 * c) + 3))) / d$

Mupad [B] (verification not implemented)

Time = 26.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.30

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = -\frac{\operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{2d}$$

input `int(1/(5*sin(c + d*x) + 3),x)`

output `-atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4)/(2*d)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.54

$$\int \frac{1}{3 + 5 \sin(c + dx)} dx = \frac{-\log(\tan(\frac{dx}{2} + \frac{c}{2}) + 3) + \log(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{4d}$$

input `int(1/(3+5*sin(d*x+c)),x)`

output `(- log(tan((c + d*x)/2) + 3) + log(3*tan((c + d*x)/2) + 1))/(4*d)`

3.61 $\int \frac{1}{(3+5 \sin(c+dx))^2} dx$

Optimal result	431
Mathematica [A] (verified)	431
Rubi [A] (verified)	432
Maple [A] (verified)	434
Fricas [A] (verification not implemented)	434
Sympy [B] (verification not implemented)	435
Maxima [A] (verification not implemented)	436
Giac [A] (verification not implemented)	436
Mupad [B] (verification not implemented)	437
Reduce [B] (verification not implemented)	437

Optimal result

Integrand size = 12, antiderivative size = 88

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx = \frac{3 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{3 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{5 \cos(c + dx)}{16d(3 + 5 \sin(c + dx))}$$

```
output 3/64*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-3/64*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d-5/16*cos(d*x+c)/d/(3+5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.43

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx = \frac{9(\log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) - \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))) + 20 \sin(\frac{1}{2}(c + dx))}{192d}$$

input `Integrate[(3 + 5*Sin[c + d*x])^(-2),x]`

output `(9*(Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]]) + 20*Sin[(c + d*x)/2]*((3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^(-1) + 3/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])))/(192*d)`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.80, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 3143, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 \sin(c + dx) + 3)^2} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{1}{(5 \sin(c + dx) + 3)^2} dx \\
 & \quad \downarrow 3143 \\
 & \frac{1}{16} \int -\frac{3}{5 \sin(c + dx) + 3} dx - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow 27 \\
 & -\frac{3}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow 3042 \\
 & -\frac{3}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow 3139 \\
 & -\frac{3 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) + 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow 1081
 \end{aligned}$$

$$\frac{9 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx))+1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx))+3)} \right) d \tan \left(\frac{1}{2}(c+dx) \right)}{8d} - \frac{5 \cos(c+dx)}{16d(5 \sin(c+dx)+3)}$$

↓ 2009

$$\frac{9 \left(\frac{1}{24} \log \left(3 \tan \left(\frac{1}{2}(c+dx) \right) + 1 \right) - \frac{1}{24} \log \left(\tan \left(\frac{1}{2}(c+dx) \right) + 3 \right) \right)}{8d} - \frac{5 \cos(c+dx)}{16d(5 \sin(c+dx)+3)}$$

input `Int[(3 + 5*Sin[c + d*x])^(-2),x]`

output `(-9*(-1/24*Log[3 + Tan[(c + d*x)/2]] + Log[1 + 3*Tan[(c + d*x)/2]]/24))/(8*d) - (5*Cos[c + d*x])/(16*d*(3 + 5*Sin[c + d*x]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{-\frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{64} - \frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{64}}{d}$
default	$\frac{-\frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{64} - \frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{64}}{d}$
risch	$-\frac{3 e^{i(dx+c)} + 5i}{8d(5 e^{2i(dx+c)} - 5 + 6i e^{i(dx+c)})} + \frac{3 \ln(\frac{4}{5} + \frac{3i}{5} + e^{i(dx+c)})}{64d} - \frac{3 \ln(e^{i(dx+c)} - \frac{4}{5} + \frac{3i}{5})}{64d}$
norman	$\frac{-\frac{5}{8d} - \frac{25 \tan(\frac{dx}{2} + \frac{c}{2})}{24d}}{3 \tan(\frac{dx}{2} + \frac{c}{2})^2 + 10 \tan(\frac{dx}{2} + \frac{c}{2}) + 3} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{64d} - \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{64d}$
parallelrisch	$\frac{45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3) \sin(dx+c) - 45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + \frac{1}{3}) \sin(dx+c) - 100 \sin(dx+c) - 60 \cos(dx+c) + 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{192d(3 + 5 \sin(dx+c))}$

input

```
int(1/(3+5*sin(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-5/48/(3*tan(1/2*d*x+1/2*c)+1)-3/64*ln(3*tan(1/2*d*x+1/2*c)+1)-5/16/(
tan(1/2*d*x+1/2*c)+3)+3/64*ln(tan(1/2*d*x+1/2*c)+3))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx$$

$$= \frac{3(5 \sin(dx + c) + 3) \log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - 3(5 \sin(dx + c) + 3) \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5)}{128(5d \sin(dx + c) + 3d)}$$

input `integrate(1/(3+5*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/128*(3*(5*sin(d*x + c) + 3)*log(4*cos(d*x + c) + 3*sin(d*x + c) + 5) - 3*(5*sin(d*x + c) + 3)*log(-4*cos(d*x + c) + 3*sin(d*x + c) + 5) - 40*cos(d*x + c))/(5*d*sin(d*x + c) + 3*d)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 466 vs. $2(78) = 156$.

Time = 0.72 (sec) , antiderivative size = 466, normalized size of antiderivative = 5.30

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(3+5*sin(d*x+c))**2,x)`

output `Piecewise((x/(3 - 5*sin(2*atan(1/3)))**2, Eq(c, -d*x - 2*atan(1/3))), (x/(3 - 5*sin(2*atan(3)))**2, Eq(c, -d*x - 2*atan(3))), (x/(5*sin(c) + 3)**2, Eq(d, 0)), (27*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) + 90*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) + 27*log(tan(c/2 + d*x/2) + 3)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 27*log(3*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 90*log(3*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 27*log(3*tan(c/2 + d*x/2) + 1)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 200*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 120/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.31

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx$$

$$= -\frac{\frac{40 \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} + 3 \right)}{\frac{10 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 3} + 9 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) - 9 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 3 \right)}{192 d}$$

input `integrate(1/(3+5*sin(d*x+c))^2,x, algorithm="maxima")`output `-1/192*(40*(5*sin(d*x + c)/(cos(d*x + c) + 1) + 3)/(10*sin(d*x + c)/(cos(d*x + c) + 1) + 3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3) + 9*log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - 9*log(sin(d*x + c)/(cos(d*x + c) + 1) + 3))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.92

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx =$$

$$-\frac{\frac{40 \left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)}{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3} + 9 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) - 9 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right| \right)}{192 d}$$

input `integrate(1/(3+5*sin(d*x+c))^2,x, algorithm="giac")`output `-1/192*(40*(5*tan(1/2*d*x + 1/2*c) + 3)/(3*tan(1/2*d*x + 1/2*c)^2 + 10*tan(1/2*d*x + 1/2*c) + 3) + 9*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) - 9*log(abs(tan(1/2*d*x + 1/2*c) + 3)))/d`

Mupad [B] (verification not implemented)

Time = 25.97 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.73

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx = \frac{3 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{32 d} - \frac{\frac{25 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{72} + \frac{5}{24}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \frac{10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(1/(5*sin(c + d*x) + 3)^2,x)`output `(3*atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4))/(32*d) - ((25*tan(c/2 + (d*x)/2)/72 + 5/24)/(d*((10*tan(c/2 + (d*x)/2))/3 + tan(c/2 + (d*x)/2)^2 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.11

$$\int \frac{1}{(3 + 5 \sin(c + dx))^2} dx = \frac{-20 \cos(dx + c) + 15 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) \sin(dx + c) + 9 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) - 15 \log\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64d(5 \sin(dx + c) + 3)}$$

input `int(1/(3+5*sin(d*x+c))^2,x)`output `(- 20*cos(c + d*x) + 15*log(tan((c + d*x)/2) + 3)*sin(c + d*x) + 9*log(tan((c + d*x)/2) + 3) - 15*log(3*tan((c + d*x)/2) + 1)*sin(c + d*x) - 9*log(3*tan((c + d*x)/2) + 1))/(64*d*(5*sin(c + d*x) + 3))`

3.62 $\int \frac{1}{(3+5 \sin(c+dx))^3} dx$

Optimal result	438
Mathematica [A] (verified)	438
Rubi [A] (verified)	439
Maple [A] (verified)	442
Fricas [A] (verification not implemented)	442
Sympy [B] (verification not implemented)	443
Maxima [A] (verification not implemented)	444
Giac [A] (verification not implemented)	444
Mupad [B] (verification not implemented)	445
Reduce [B] (verification not implemented)	445

Optimal result

Integrand size = 12, antiderivative size = 113

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx = -\frac{43 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{2048d} + \frac{43 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{5 \cos(c + dx)}{32d(3 + 5 \sin(c + dx))^2} + \frac{45 \cos(c + dx)}{512d(3 + 5 \sin(c + dx))}$$

```
output -43/2048*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d+43/2048*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d-5/32*cos(d*x+c)/d/(3+5*sin(d*x+c))^2+45/512*cos(d*x+c)/d/(3+5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.59

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx = -43 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) + 43 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx))) + \frac{5 \cos(c + dx)}{32d(3 + 5 \sin(c + dx))^2} + \frac{45 \cos(c + dx)}{512d(3 + 5 \sin(c + dx))}$$

input `Integrate[(3 + 5*Sin[c + d*x])^(-3),x]`

output `(-43*Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] + 43*Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]] + 40/(3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 - 40/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])^2 + Sin[(c + d*x)/2]*(-60/(3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) - 180/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])))/(2048*d)`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.88, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 \sin(c + dx) + 3)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 \sin(c + dx) + 3)^3} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{32} \int -\frac{6 - 5 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx - \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \\
 & \quad \downarrow \text{25} \\
 & -\frac{1}{32} \int \frac{6 - 5 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx - \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{32} \int \frac{6 - 5 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx - \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} - \frac{1}{16} \int -\frac{43}{5 \sin(c + dx) + 3} dx \right) - \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{32} \left(\frac{43}{16} \int \frac{1}{5 \sin(c+dx)+3} dx + \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) - \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 3042 \\
& \frac{1}{32} \left(\frac{43}{16} \int \frac{1}{5 \sin(c+dx)+3} dx + \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) - \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 3139 \\
& \frac{1}{32} \left(\frac{43 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx))+10 \tan(\frac{1}{2}(c+dx))+3} d \tan(\frac{1}{2}(c+dx))}{8d} + \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) - \\
& \quad \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 1081 \\
& \frac{1}{32} \left(\frac{129 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx))+1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx))+3)} \right) d \tan(\frac{1}{2}(c+dx))}{8d} + \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) - \\
& \quad \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 2009 \\
& \frac{1}{32} \left(\frac{129 \left(\frac{1}{24} \log(3 \tan(\frac{1}{2}(c+dx))+1) - \frac{1}{24} \log(\tan(\frac{1}{2}(c+dx))+3) \right)}{8d} + \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) - \\
& \quad \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2}
\end{aligned}$$

input `Int[(3 + 5*Sin[c + d*x])^(-3),x]`

output `(-5*Cos[c + d*x])/(32*d*(3 + 5*Sin[c + d*x])^2) + ((129*(-1/24*Log[3 + Tan[(c + d*x)/2]] + Log[1 + 3*Tan[(c + d*x)/2]]/24))/(8*d) + (45*Cos[c + d*x])/(16*d*(3 + 5*Sin[c + d*x]))/32`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 1081 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = \text{Rt}[\text{b}^2 - 4*\text{a}*c, 2]\}, \text{Simp}[\text{c} \quad \text{Int}[\text{ExpandIntegrand}[1/((\text{b}/2 - \text{q}/2 + \text{c}*x)*(\text{b}/2 + \text{q}/2 + \text{c}*x)), \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}] \ \&\& \ \text{NiceSqrtQ}[\text{b}^2 - 4*\text{a}*c]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ ; SumQ}[\text{u}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3139 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{e} = \text{FreeFactors}[\text{Tan}[(\text{c} + \text{d}*x)/2], \text{x}]\}, \text{Simp}[2*(\text{e}/\text{d}) \quad \text{Subst}[\text{Int}[1/(\text{a} + 2*\text{b}*e*x + \text{a}*e^2*x^2), \text{x}], \text{x}, \text{Tan}[(\text{c} + \text{d}*x)/2]/\text{e}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0]$
- rule 3143 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(\text{n}_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b})*\text{Cos}[\text{c} + \text{d}*x]*((\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(\text{n} + 1)}/(\text{d}*(\text{n} + 1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((\text{n} + 1)*(a^2 - b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(\text{n} + 1)}*\text{Simp}[\text{a}*(\text{n} + 1) - \text{b}*(\text{n} + 2)*\text{Sin}[\text{c} + \text{d}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{n}, -1] \ \&\& \ \text{IntegerQ}[2*\text{n}]$
- rule 3233 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]^{(\text{m}_)}*((\text{c}_.) + (\text{d}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{b}*c - \text{a}*d))*\text{Cos}[\text{e} + \text{f}*x]*((\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(\text{m} + 1)}/(\text{f}*(\text{m} + 1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((\text{m} + 1)*(a^2 - b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(\text{m} + 1)}*\text{Simp}[(\text{a}*c - \text{b}*d)*(m + 1) - (\text{b}*c - \text{a}*d)*(m + 2)*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntegerQ}[2*\text{m}]$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.88

method	result
derivativedivides	$-\frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} + \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} + \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{2048} + \frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} - \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}$
default	$-\frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} + \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} + \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{2048} + \frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} - \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}$
risch	$\frac{387ie^{2i(dx+c)} + 215e^{3i(dx+c)} - 325e^{i(dx+c)} - 225i}{256(5e^{2i(dx+c)} - 5 + 6ie^{i(dx+c)})^2} d + \frac{43 \ln(e^{i(dx+c)} - \frac{4}{5} + \frac{3i}{5})}{2048d} - \frac{43 \ln(\frac{4}{5} + \frac{3i}{5} + e^{i(dx+c)})}{2048d}$
norman	$\frac{55}{256d} + \frac{3245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{2304d} - \frac{125 \tan(\frac{dx}{2} + \frac{c}{2})^3}{768d} + \frac{1225 \tan(\frac{dx}{2} + \frac{c}{2})}{768d} - \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{2048d} + \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{2048d}$
parallelrisc	$\frac{(9675 \cos(2dx+2c) - 23220 \sin(dx+c) - 16641) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + \frac{1}{3}) + (-9675 \cos(2dx+2c) + 23220 \sin(dx+c) + 16641)}{18432d(-43 + 25 \cos(2dx+2c))}$

```
input int(1/(3+5*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-25/1152/(3*tan(1/2*d*x+1/2*c)+1)^2+155/4608/(3*tan(1/2*d*x+1/2*c)+1)
+43/2048*ln(3*tan(1/2*d*x+1/2*c)+1)+25/128/(tan(1/2*d*x+1/2*c)+3)^2-15/512
/(tan(1/2*d*x+1/2*c)+3)-43/2048*ln(tan(1/2*d*x+1/2*c)+3))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.18

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx =$$

$$-\frac{43 (25 \cos(dx + c)^2 - 30 \sin(dx + c) - 34) \log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - 43 (25 \cos(dx + c) + 3 \sin(dx + c) + 5)}{4096 (25 d \cos(dx + c) + 3 \sin(dx + c) + 5)^2}$$

```
input integrate(1/(3+5*sin(d*x+c))^3,x, algorithm="fricas")
```

output

```
-1/4096*(43*(25*cos(d*x + c)^2 - 30*sin(d*x + c) - 34)*log(4*cos(d*x + c)
+ 3*sin(d*x + c) + 5) - 43*(25*cos(d*x + c)^2 - 30*sin(d*x + c) - 34)*log(
-4*cos(d*x + c) + 3*sin(d*x + c) + 5) + 1800*cos(d*x + c)*sin(d*x + c) + 4
40*cos(d*x + c))/(25*d*cos(d*x + c)^2 - 30*d*sin(d*x + c) - 34*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1227 vs. $2(102) = 204$.

Time = 1.44 (sec) , antiderivative size = 1227, normalized size of antiderivative = 10.86

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(3+5*sin(d*x+c))**3,x)
```

output

```
Piecewise((x/(3 - 5*sin(2*atan(1/3)))**3, Eq(c, -d*x - 2*atan(1/3))), (x/(
3 - 5*sin(2*atan(3)))**3, Eq(c, -d*x - 2*atan(3))), (x/(5*sin(c) + 3)**3,
Eq(d, 0)), (-3483*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**4/(165888*d*
tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 +
d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 23220*log(tan(c/2 + d
*x/2) + 3)*tan(c/2 + d*x/2)**3/(165888*d*tan(c/2 + d*x/2)**4 + 1105920*d*t
an(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 + 1105920*d*tan(c/2 + d
*x/2) + 165888*d) - 45666*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**2/(1
65888*d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*ta
n(c/2 + d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 23220*log(tan
(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)/(165888*d*tan(c/2 + d*x/2)**4 + 110592
0*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 + 1105920*d*tan(c/
2 + d*x/2) + 165888*d) - 3483*log(tan(c/2 + d*x/2) + 3)/(165888*d*tan(c/2
+ d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**
2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 3483*log(3*tan(c/2 + d*x/2) +
1)*tan(c/2 + d*x/2)**4/(165888*d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2
+ d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) +
165888*d) + 23220*log(3*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**3/(165888
*d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2
+ d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 45666*log(3*tan...
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.73

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{40 \left(\frac{735 \sin(dx+c)}{\cos(dx+c)+1} + \frac{649 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{75 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + 99 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} + \frac{118 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{9 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 9} + 387 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) - 387 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 3 \right)$$

$$18432 d$$

input `integrate(1/(3+5*sin(d*x+c))^3,x, algorithm="maxima")`output

```
1/18432*(40*(735*sin(d*x + c)/(cos(d*x + c) + 1) + 649*sin(d*x + c)^2/(cos
(d*x + c) + 1)^2 - 75*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 99)/(60*sin(d*
x + c)/(cos(d*x + c) + 1) + 118*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*s
in(d*x + c)^3/(cos(d*x + c) + 1)^3 + 9*sin(d*x + c)^4/(cos(d*x + c) + 1)^4
+ 9) + 387*log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - 387*log(sin(d*x +
c)/(cos(d*x + c) + 1) + 3))/d
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.95

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx =$$

$$\frac{40 \left(75 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 649 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 99 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^2} - 387 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) + 387 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right| \right)$$

$$18432 d$$

input `integrate(1/(3+5*sin(d*x+c))^3,x, algorithm="giac")`output

```
-1/18432*(40*(75*tan(1/2*d*x + 1/2*c)^3 - 649*tan(1/2*d*x + 1/2*c)^2 - 735
*tan(1/2*d*x + 1/2*c) - 99)/(3*tan(1/2*d*x + 1/2*c)^2 + 10*tan(1/2*d*x + 1
/2*c) + 3)^2 - 387*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) + 387*log(abs(tan(
1/2*d*x + 1/2*c) + 3)))/d
```

Mupad [B] (verification not implemented)

Time = 27.20 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.02

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{-\frac{125 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{6912} + \frac{3245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{20736} + \frac{1225 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{6912} + \frac{55}{2304}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{3} + \frac{118 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{9} + \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

$$- \frac{43 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{1024 d}$$

input `int(1/(5*sin(c + d*x) + 3)^3,x)`output `((1225*tan(c/2 + (d*x)/2))/6912 + (3245*tan(c/2 + (d*x)/2)^2)/20736 - (125*tan(c/2 + (d*x)/2)^3)/6912 + 55/2304)/(d*((20*tan(c/2 + (d*x)/2))/3 + (118*tan(c/2 + (d*x)/2)^2)/9 + (20*tan(c/2 + (d*x)/2)^3)/3 + tan(c/2 + (d*x)/2)^4 + 1)) - (43*atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4))/(1024*d)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.65

$$\int \frac{1}{(3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{900 \cos(dx + c) \sin(dx + c) + 220 \cos(dx + c) - 1075 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) \sin(dx + c)^2 - 1290 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{(3 + 5 \sin(dx + c))^3}$$

input `int(1/(3+5*sin(d*x+c))^3,x)`

output

```
(900*cos(c + d*x)*sin(c + d*x) + 220*cos(c + d*x) - 1075*log(tan((c + d*x)/2) + 3)*sin(c + d*x)**2 - 1290*log(tan((c + d*x)/2) + 3)*sin(c + d*x) - 387*log(tan((c + d*x)/2) + 3) + 1075*log(3*tan((c + d*x)/2) + 1)*sin(c + d*x)**2 + 1290*log(3*tan((c + d*x)/2) + 1)*sin(c + d*x) + 387*log(3*tan((c + d*x)/2) + 1) + 750*sin(c + d*x)**2 + 900*sin(c + d*x) + 270)/(2048*d*(25*sin(c + d*x)**2 + 30*sin(c + d*x) + 9))
```

3.63 $\int \frac{1}{(3+5 \sin(c+dx))^4} dx$

Optimal result	447
Mathematica [A] (verified)	448
Rubi [A] (verified)	448
Maple [A] (verified)	452
Fricas [A] (verification not implemented)	452
Sympy [B] (verification not implemented)	453
Maxima [B] (verification not implemented)	454
Giac [A] (verification not implemented)	454
Mupad [B] (verification not implemented)	455
Reduce [B] (verification not implemented)	455

Optimal result

Integrand size = 12, antiderivative size = 138

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \frac{279 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{32768d} - \frac{279 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))}{32768d} - \frac{5 \cos(c + dx)}{48d(3 + 5 \sin(c + dx))^3} + \frac{25 \cos(c + dx)}{512d(3 + 5 \sin(c + dx))^2} - \frac{995 \cos(c + dx)}{24576d(3 + 5 \sin(c + dx))}$$

output

```
279/32768*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-279/32768*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d-5/48*cos(d*x+c)/d/(3+5*sin(d*x+c))^3+25/512*cos(d*x+c)/d/(3+5*sin(d*x+c))^2-995/24576*cos(d*x+c)/d/(3+5*sin(d*x+c))
```


Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.70

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx$$

$$= \frac{2511 \log \left(3 \cos \left(\frac{1}{2}(c + dx) \right) + \sin \left(\frac{1}{2}(c + dx) \right) \right) - 2511 \log \left(\cos \left(\frac{1}{2}(c + dx) \right) + 3 \sin \left(\frac{1}{2}(c + dx) \right) \right)}{(3 \cos(\frac{1}{2}(c + dx)))^2} - \frac{720}{(3 \cos(\frac{1}{2}(c + dx)))^2} + \frac{20 \sin(\frac{1}{2}(c + dx))}{(3 \cos(\frac{1}{2}(c + dx)))^2} + \frac{199}{(3 \cos(\frac{1}{2}(c + dx)))^2} + \frac{240}{(3 \cos(\frac{1}{2}(c + dx)))^2} + \frac{597}{(3 \cos(\frac{1}{2}(c + dx)))^2} + \frac{20 \sin(\frac{1}{2}(c + dx))}{(3 \cos(\frac{1}{2}(c + dx)))^2}$$

input `Integrate[(3 + 5*Sin[c + d*x])^(-4), x]`

output
$$\frac{(2511 \cdot \text{Log}[3 \cdot \text{Cos}[(c + d \cdot x)/2] + \text{Sin}[(c + d \cdot x)/2]] - 2511 \cdot \text{Log}[\text{Cos}[(c + d \cdot x)/2] + 3 \cdot \text{Sin}[(c + d \cdot x)/2]] - 2320 / (3 \cdot \text{Cos}[(c + d \cdot x)/2] + \text{Sin}[(c + d \cdot x)/2])^2 + 720 / (\text{Cos}[(c + d \cdot x)/2] + 3 \cdot \text{Sin}[(c + d \cdot x)/2])^2 + 20 \cdot \text{Sin}[(c + d \cdot x)/2] \cdot (80 / (3 \cdot \text{Cos}[(c + d \cdot x)/2] + \text{Sin}[(c + d \cdot x)/2])^3 + 199 / (3 \cdot \text{Cos}[(c + d \cdot x)/2] + \text{Sin}[(c + d \cdot x)/2]) + 240 / (\text{Cos}[(c + d \cdot x)/2] + 3 \cdot \text{Sin}[(c + d \cdot x)/2])^3 + 597 / (\text{Cos}[(c + d \cdot x)/2] + 3 \cdot \text{Sin}[(c + d \cdot x)/2])) / (294912 \cdot d)}$$

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.94, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$, Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(5 \sin(c + dx) + 3)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(5 \sin(c + dx) + 3)^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{9 - 10 \sin(c + dx)}{(5 \sin(c + dx) + 3)^3} dx - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& -\frac{1}{48} \int \frac{9 - 10 \sin(c + dx)}{(5 \sin(c + dx) + 3)^3} dx - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3042 \\
& -\frac{1}{48} \int \frac{9 - 10 \sin(c + dx)}{(5 \sin(c + dx) + 3)^3} dx - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left(\frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} - \frac{1}{32} \int -\frac{154 - 75 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx \right) - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{154 - 75 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{154 - 75 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{1}{16} \int -\frac{837}{5 \sin(c + dx) + 3} dx - \frac{995 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \\
& \quad \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left(\frac{1}{32} \left(-\frac{837}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{995 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \\
& \quad \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \left(-\frac{837}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{995 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \\
& \quad \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3139
\end{aligned}$$

$$\frac{1}{48} \left(\frac{1}{32} \left(-\frac{837 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) + 10 \tan(\frac{1}{2}(c+dx)) + 3} dx \tan(\frac{1}{2}(c+dx))}{8d} - \frac{995 \cos(c+dx)}{16d(5 \sin(c+dx) + 3)} \right) + \frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)} \right)$$

$$\frac{5 \cos(c+dx)}{48d(5 \sin(c+dx) + 3)^3}$$

↓ 1081

$$\frac{1}{48} \left(\frac{1}{32} \left(-\frac{2511 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx)) + 1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx)) + 3)} \right) dx \tan(\frac{1}{2}(c+dx))}{8d} - \frac{995 \cos(c+dx)}{16d(5 \sin(c+dx) + 3)} \right) + \frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)} \right)$$

$$\frac{5 \cos(c+dx)}{48d(5 \sin(c+dx) + 3)^3}$$

↓ 2009

$$\frac{1}{48} \left(\frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)^2} + \frac{1}{32} \left(-\frac{2511 \left(\frac{1}{24} \log(3 \tan(\frac{1}{2}(c+dx)) + 1) - \frac{1}{24} \log(\tan(\frac{1}{2}(c+dx)) + 3) \right)}{8d} - \frac{995 \cos(c+dx)}{16d(5 \sin(c+dx) + 3)} \right) + \frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)} \right)$$

$$\frac{5 \cos(c+dx)}{48d(5 \sin(c+dx) + 3)^3}$$

input `Int[(3 + 5*Sin[c + d*x])^(-4),x]`

output `(-5*Cos[c + d*x])/(48*d*(3 + 5*Sin[c + d*x])^3) + ((75*Cos[c + d*x])/(32*d*(3 + 5*Sin[c + d*x])^2) + ((-2511*(-1/24*Log[3 + Tan[(c + d*x)/2]] + Log[1 + 3*Tan[(c + d*x)/2]]/24))/(8*d) - (995*Cos[c + d*x])/(16*d*(3 + 5*Sin[c + d*x]))) / 32) / 48`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.96

method	result
derivativedivides	$-\frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^3} + \frac{275}{27648(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} - \frac{3505}{221184(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{32768} - \frac{12}{768(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}$
default	$-\frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^3} + \frac{275}{27648(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} - \frac{3505}{221184(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{32768} - \frac{12}{768(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}$
risch	$-\frac{-111042 e^{3i(dx+c)} + 62775 i e^{4i(dx+c)} - 119310 i e^{2i(dx+c)} + 20925 e^{5i(dx+c)} + 68625 e^{i(dx+c)} + 24875 i}{12288(5 e^{2i(dx+c)} - 5 + 6 i e^{i(dx+c)})^3} d - \frac{279 \ln(e^{i(dx+c)} + 3)}{32768}$
norman	$-\frac{7915}{12288d} - \frac{63425 \tan(\frac{dx}{2} + \frac{c}{2})}{12288d} - \frac{3047275 \tan(\frac{dx}{2} + \frac{c}{2})^3}{165888d} - \frac{15725 \tan(\frac{dx}{2} + \frac{c}{2})^5}{12288d} - \frac{296245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{18432d} - \frac{270245 \tan(\frac{dx}{2} + \frac{c}{2})^4}{36864d} + \frac{1}{(3 \tan(\frac{dx}{2} + \frac{c}{2})^2 + 10 \tan(\frac{dx}{2} + \frac{c}{2}) + 3)^3}$
parallelrisc	$\frac{(-10169550 \cos(2dx+2c) + 20678085 \sin(dx+c) - 2824875 \sin(3dx+3c) + 12610242) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + \frac{1}{3}) + (10169550 \cos(dx+c) - 20678085 \sin(dx+c) + 2824875 \sin(3dx+3c) - 12610242)}{12288d}$

```
input int(1/(3+5*sin(d*x+c))^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-125/20736/(3*tan(1/2*d*x+1/2*c)+1)^3+275/27648/(3*tan(1/2*d*x+1/2*c)+1)^2-3505/221184/(3*tan(1/2*d*x+1/2*c)+1)-279/32768*ln(3*tan(1/2*d*x+1/2*c)+1)-125/768/(tan(1/2*d*x+1/2*c)+3)^3+75/1024/(tan(1/2*d*x+1/2*c)+3)^2-345/8192/(tan(1/2*d*x+1/2*c)+3)+279/32768*ln(tan(1/2*d*x+1/2*c)+3))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.31

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \frac{199000 \cos(dx + c)^3 - 837(225 \cos(dx + c)^2 + 5(25 \cos(dx + c)^2 - 52) \sin(dx + c) - 252) \log(4 \cos(dx + c) + 3)}{12288d}$$

```
input integrate(1/(3+5*sin(d*x+c))^4,x, algorithm="fricas")
```

output

```
-1/196608*(199000*cos(d*x + c)^3 - 837*(225*cos(d*x + c)^2 + 5*(25*cos(d*x
+ c)^2 - 52)*sin(d*x + c) - 252)*log(4*cos(d*x + c) + 3*sin(d*x + c) + 5)
+ 837*(225*cos(d*x + c)^2 + 5*(25*cos(d*x + c)^2 - 52)*sin(d*x + c) - 252
)*log(-4*cos(d*x + c) + 3*sin(d*x + c) + 5) - 190800*cos(d*x + c)*sin(d*x
+ c) - 262320*cos(d*x + c))/(225*d*cos(d*x + c)^2 + 5*(25*d*cos(d*x + c)^2
- 52*d)*sin(d*x + c) - 252*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2356 vs. $2(126) = 252$.

Time = 3.22 (sec) , antiderivative size = 2356, normalized size of antiderivative = 17.07

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(3+5*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(3 - 5*sin(2*atan(1/3)))**4, Eq(c, -d*x - 2*atan(1/3))), (x/(
3 - 5*sin(2*atan(3)))**4, Eq(c, -d*x - 2*atan(3))), (x/(5*sin(c) + 3)**4,
Eq(d, 0)), (610173*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**6/(71663616
*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*ta
n(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/
2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 6101730*log(t
an(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**5/(71663616*d*tan(c/2 + d*x/2)**6 +
716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087
480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + 71663616
0*d*tan(c/2 + d*x/2) + 71663616*d) + 22169619*log(tan(c/2 + d*x/2) + 3)*ta
n(c/2 + d*x/2)**4/(71663616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 +
d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/
2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) +
71663616*d) + 34802460*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**3/(7166
3616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*
d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*ta
n(c/2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 22169619*
log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**2/(71663616*d*tan(c/2 + d*x/2)
**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 +
4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + ...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(124) = 248$.

Time = 0.04 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.99

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \frac{40 \left(\frac{342495 \sin(dx+c)}{\cos(dx+c)+1} + \frac{1066482 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1218910 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{486441 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{84915 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + 42741 \right) + 22599 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} \right)}{2654208 d}$$

input `integrate(1/(3+5*sin(d*x+c))^4,x, algorithm="maxima")`

output `-1/2654208*(40*(342495*sin(d*x + c)/(cos(d*x + c) + 1) + 1066482*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1218910*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 486441*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 84915*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 42741)/(270*sin(d*x + c)/(cos(d*x + c) + 1) + 981*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1540*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 981*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 270*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 27*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 27) + 22599*log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - 22599*log(sin(d*x + c)/(cos(d*x + c) + 1) + 3))/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.96

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \frac{40 \left(84915 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 486441 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 1218910 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 1066482 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 342495 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 42741 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^3} + 22599 \log \left(\frac{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1} \right) / d$$

2654208 d

input `integrate(1/(3+5*sin(d*x+c))^4,x, algorithm="giac")`

output

```
-1/2654208*(40*(84915*tan(1/2*d*x + 1/2*c)^5 + 486441*tan(1/2*d*x + 1/2*c)
^4 + 1218910*tan(1/2*d*x + 1/2*c)^3 + 1066482*tan(1/2*d*x + 1/2*c)^2 + 342
495*tan(1/2*d*x + 1/2*c) + 42741)/(3*tan(1/2*d*x + 1/2*c)^2 + 10*tan(1/2*d
*x + 1/2*c) + 3)^3 + 22599*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) - 22599*lo
g(abs(tan(1/2*d*x + 1/2*c) + 3))/d
```

Mupad [B] (verification not implemented)

Time = 28.67 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.22

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \frac{279 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{16384 d} - \frac{\frac{15725 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{331776} + \frac{270245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{995328} + \frac{3047275 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{4478976} + \frac{296245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{497664} + \frac{63425 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{331776} + 3}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{3} + \frac{1540 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{27} + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{3} + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1 \right)}$$

input

```
int(1/(5*sin(c + d*x) + 3)^4,x)
```

output

```
(279*atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4))/(16384*d) - ((63425*tan(c/2 +
(d*x)/2))/331776 + (296245*tan(c/2 + (d*x)/2)^2)/497664 + (3047275*tan(c/2
+ (d*x)/2)^3)/4478976 + (270245*tan(c/2 + (d*x)/2)^4)/995328 + (15725*tan
(c/2 + (d*x)/2)^5)/331776 + 7915/331776)/(d*(10*tan(c/2 + (d*x)/2) + (109*
tan(c/2 + (d*x)/2)^2)/3 + (1540*tan(c/2 + (d*x)/2)^3)/27 + (109*tan(c/2 +
(d*x)/2)^4)/3 + 10*tan(c/2 + (d*x)/2)^5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.95

$$\int \frac{1}{(3 + 5 \sin(c + dx))^4} dx = \frac{-298500 \cos(dx + c) \sin(dx + c)^2 - 286200 \cos(dx + c) \sin(dx + c) - 94980 \cos(dx + c) + 313875 \log}{d}$$

input `int(1/(3+5*sin(d*x+c))^4,x)`

output
$$\begin{aligned} & (-298500 \cos(c + dx) \sin(c + dx)^2 - 286200 \cos(c + dx) \sin(c + dx) \\ & - 94980 \cos(c + dx) + 313875 \log(\tan((c + dx)/2) + 3) \sin(c + dx)^3 + \\ & 564975 \log(\tan((c + dx)/2) + 3) \sin(c + dx)^2 + 338985 \log(\tan((c + dx) \\ & /2) + 3) \sin(c + dx) + 67797 \log(\tan((c + dx)/2) + 3) - 313875 \log(3 \tan \\ & ((c + dx)/2) + 1) \sin(c + dx)^3 - 564975 \log(3 \tan((c + dx)/2) + 1) \sin \\ & (c + dx)^2 - 338985 \log(3 \tan((c + dx)/2) + 1) \sin(c + dx) - 67797 \log \\ & (3 \tan((c + dx)/2) + 1) - 265000 \sin(c + dx)^3 - 477000 \sin(c + dx) \\ & ^2 - 286200 \sin(c + dx) - 57240) / (294912 d (125 \sin(c + dx)^3 + 225 \sin \\ & (c + dx)^2 + 135 \sin(c + dx) + 27)) \end{aligned}$$

3.64 $\int \frac{1}{3-5 \sin(c+dx)} dx$

Optimal result	457
Mathematica [A] (verified)	457
Rubi [A] (verified)	458
Maple [A] (verified)	459
Fricas [A] (verification not implemented)	460
Sympy [A] (verification not implemented)	460
Maxima [A] (verification not implemented)	460
Giac [A] (verification not implemented)	461
Mupad [B] (verification not implemented)	461
Reduce [B] (verification not implemented)	462

Optimal result

Integrand size = 12, antiderivative size = 65

$$\int \frac{1}{3-5 \sin(c+dx)} dx = -\frac{\log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{4d} + \frac{\log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{4d}$$

output `-1/4*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d+1/4*ln(3*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00

$$\int \frac{1}{3-5 \sin(c+dx)} dx = -\frac{\log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{4d} + \frac{\log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{4d}$$

input `Integrate[(3 - 5*Sin[c + d*x])^(-1),x]`

output

$$-1/4*\text{Log}[\text{Cos}[(c + d*x)/2] - 3*\text{Sin}[(c + d*x)/2]]/d + \text{Log}[3*\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]]/(4*d)$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{3 - 5 \sin(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{3 - 5 \sin(c + dx)} dx \\ & \quad \downarrow \text{3139} \\ & \frac{2 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{1081} \\ & \frac{6 \int \left(\frac{1}{8(1-3 \tan(\frac{1}{2}(c+dx)))} - \frac{1}{24(3-\tan(\frac{1}{2}(c+dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{6 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx))) \right)}{d} \end{aligned}$$

input

$$\text{Int}[(3 - 5*\text{Sin}[c + d*x])^(-1),x]$$

output

$$(6*(-1/24*\text{Log}[1 - 3*\text{Tan}[(c + d*x)/2]] + \text{Log}[3 - \text{Tan}[(c + d*x)/2]]/24))/d$$

Definitions of rubi rules used

rule 1081 $\text{Int}[(a_ + (b_ \cdot)x_ + (c_ \cdot)x_^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4a \cdot c, 2]\}, \text{Simp}[c \text{ Int}[\text{ExpandIntegrand}[1/((b/2 - q/2 + c \cdot x) \cdot (b/2 + q/2 + c \cdot x)), x], x], x]] \text{ /; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NiceSqrtQ}[b^2 - 4a \cdot c]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3139 $\text{Int}[(a_ + (b_ \cdot)\sin[(c_ \cdot) + (d_ \cdot)x_])^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d \cdot x)/2], x]\}, \text{Simp}[2 \cdot (e/d) \text{ Subst}[\text{Int}[1/(a + 2 \cdot b \cdot e \cdot x + a \cdot e^2 \cdot x^2), x], x, \text{Tan}[(c + d \cdot x)/2]/e], x]] \text{ /; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$\frac{\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) - \ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4}}{d}$	36
default	$\frac{\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) - \ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4}}{d}$	36
parallelrisc	$\frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 9\right) - \ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	37
norman	$\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{4d} - \frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	38
risc	$-\frac{\ln\left(-\frac{4}{5} - \frac{3i}{5} + e^{i(dx+c)}\right)}{4d} + \frac{\ln\left(e^{i(dx+c)} + \frac{4}{5} - \frac{3i}{5}\right)}{4d}$	40

input $\text{int}(1/(3-5 \cdot \sin(dx+c)), x, \text{method}=_RETURNVERBOSE)$

output $1/d \cdot (1/4 \cdot \ln(\tan(1/2 \cdot dx + 1/2 \cdot c) - 3) - 1/4 \cdot \ln(3 \cdot \tan(1/2 \cdot dx + 1/2 \cdot c) - 1))$

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.71

$$\int \frac{1}{3 - 5 \sin(c + dx)} dx = \frac{\log(4 \cos(dx + c) - 3 \sin(dx + c) + 5) - \log(-4 \cos(dx + c) - 3 \sin(dx + c) + 5)}{8d}$$

input `integrate(1/(3-5*sin(d*x+c)),x, algorithm="fricas")`output `1/8*(log(4*cos(d*x + c) - 3*sin(d*x + c) + 5) - log(-4*cos(d*x + c) - 3*sin(d*x + c) + 5))/d`**Sympy [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \frac{1}{3 - 5 \sin(c + dx)} dx = \begin{cases} \frac{\log\left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 3\right)}{4d} - \frac{\log\left(3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 1\right)}{4d} & \text{for } d \neq 0 \\ \frac{x}{3 - 5 \sin(c)} & \text{otherwise} \end{cases}$$

input `integrate(1/(3-5*sin(d*x+c)),x)`output `Piecewise((log(tan(c/2 + d*x/2) - 3)/(4*d) - log(3*tan(c/2 + d*x/2) - 1)/(4*d), Ne(d, 0)), (x/(3 - 5*sin(c)), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.75

$$\int \frac{1}{3 - 5 \sin(c + dx)} dx = -\frac{\log\left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 1\right) - \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 3\right)}{4d}$$

input `integrate(1/(3-5*sin(d*x+c)),x, algorithm="maxima")`

output
$$-1/4*(\log(3*\sin(d*x + c)/(\cos(d*x + c) + 1) - 1) - \log(\sin(d*x + c)/(\cos(d*x + c) + 1) - 3))/d$$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.55

$$\int \frac{1}{3 - 5 \sin(c + dx)} dx = -\frac{\log\left(\left|3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right|\right) - \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3\right|\right)}{4d}$$

input `integrate(1/(3-5*sin(d*x+c)),x, algorithm="giac")`

output
$$-1/4*(\log(\text{abs}(3*\tan(1/2*d*x + 1/2*c) - 1)) - \log(\text{abs}(\tan(1/2*d*x + 1/2*c) - 3)))/d$$

Mupad [B] (verification not implemented)

Time = 26.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.29

$$\int \frac{1}{3 - 5 \sin(c + dx)} dx = -\frac{\operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{5}{4}\right)}{2d}$$

input `int(-1/(5*sin(c + d*x) - 3),x)`

output
$$-\operatorname{atanh}((3*\tan(c/2 + (d*x)/2))/4 - 5/4)/(2*d)$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.52

$$\int \frac{1}{3 - 5 \sin(c + dx)} dx = \frac{\log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) - \log\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$$

input `int(1/(3-5*sin(d*x+c)),x)`

output `(log(tan((c + d*x)/2) - 3) - log(3*tan((c + d*x)/2) - 1))/(4*d)`

3.65 $\int \frac{1}{(3-5 \sin(c+dx))^2} dx$

Optimal result	463
Mathematica [A] (verified)	463
Rubi [A] (verified)	464
Maple [A] (verified)	466
Fricas [A] (verification not implemented)	466
Sympy [B] (verification not implemented)	467
Maxima [A] (verification not implemented)	468
Giac [A] (verification not implemented)	468
Mupad [B] (verification not implemented)	469
Reduce [B] (verification not implemented)	469

Optimal result

Integrand size = 12, antiderivative size = 90

$$\int \frac{1}{(3-5 \sin(c+dx))^2} dx = \frac{3 \log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{64d} - \frac{3 \log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{64d} + \frac{5 \cos(c+dx)}{16d(3-5 \sin(c+dx))}$$

output

```
3/64*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d-3/64*ln(3*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+5/16*cos(d*x+c)/d/(3-5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.44

$$\int \frac{1}{(3-5 \sin(c+dx))^2} dx = \frac{9(\log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx))) - \log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx))))}{192d} + 20 \left(\frac{1}{\cos(\frac{1}{2}(c+dx))} \right)$$

input `Integrate[(3 - 5*Sin[c + d*x])^(-2),x]`

output `(9*(Log[Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]] - Log[3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]]) + 20*(3/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]) + (3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^(-1))*Sin[(c + d*x)/2])/(192*d)`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.80, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 3143, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 - 5 \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 - 5 \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{16} \int -\frac{3}{3 - 5 \sin(c + dx)} dx + \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{3}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{3}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \\
 & \quad \downarrow \text{3139} \\
 & \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{3 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} \\
 & \quad \downarrow \text{1081}
 \end{aligned}$$

$$\frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{9 \int \left(\frac{1}{8(1 - 3 \tan(\frac{1}{2}(c + dx)))} - \frac{1}{24(3 - \tan(\frac{1}{2}(c + dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{8d}$$

↓ 2009

$$\frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{9 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx))) \right)}{8d}$$

input `Int[(3 - 5*Sin[c + d*x])^(-2),x]`

output `(-9*(-1/24*Log[1 - 3*Tan[(c + d*x)/2]] + Log[3 - Tan[(c + d*x)/2]]/24))/(8*d) + (5*Cos[c + d*x])/(16*d*(3 - 5*Sin[c + d*x]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76

method	result
derivativedivides	$\frac{-\frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{64} - \frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} + \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{64}}{d}$
default	$\frac{-\frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{64} - \frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} + \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{64}}{d}$
risch	$\frac{3 e^{i(dx+c)} - 5i}{8d(5 e^{2i(dx+c)} - 5 - 6i e^{i(dx+c)})} - \frac{3 \ln(e^{i(dx+c)} + \frac{4}{5} - \frac{3i}{5})}{64d} + \frac{3 \ln(-\frac{4}{5} - \frac{3i}{5} + e^{i(dx+c)})}{64d}$
norman	$\frac{\frac{5}{8d} - \frac{25 \tan(\frac{dx}{2} + \frac{c}{2})}{24d}}{3 \tan(\frac{dx}{2} + \frac{c}{2})^2 - 10 \tan(\frac{dx}{2} + \frac{c}{2}) + 3} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{64d} + \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{64d}$
parallelrisch	$\frac{-45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3) \sin(dx+c) + 45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{3}) \sin(dx+c) + 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3) - 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}))}{192d(-3 + 5 \sin(dx+c))}$

input

```
int(1/(3-5*sin(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-5/16/(tan(1/2*d*x+1/2*c)-3)-3/64*ln(tan(1/2*d*x+1/2*c)-3)-5/48/(3*tan(1/2*d*x+1/2*c)-1)+3/64*ln(3*tan(1/2*d*x+1/2*c)-1))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.98

$$\int \frac{1}{(3 - 5 \sin(c + dx))^2} dx = \frac{3(5 \sin(dx + c) - 3) \log(4 \cos(dx + c) - 3 \sin(dx + c) + 5) - 3(5 \sin(dx + c) - 3) \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5) + 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3) - 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}))}{128(5d \sin(dx + c) - 3d)}$$

input `integrate(1/(3-5*sin(d*x+c))^2,x, algorithm="fricas")`

output `-1/128*(3*(5*sin(d*x + c) - 3)*log(4*cos(d*x + c) - 3*sin(d*x + c) + 5) - 3*(5*sin(d*x + c) - 3)*log(-4*cos(d*x + c) - 3*sin(d*x + c) + 5) + 40*cos(d*x + c))/(5*d*sin(d*x + c) - 3*d)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 462 vs. $2(78) = 156$.

Time = 0.76 (sec) , antiderivative size = 462, normalized size of antiderivative = 5.13

$$\int \frac{1}{(3 - 5 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(3-5*sin(d*x+c))**2,x)`

output `Piecewise((x/(3 - 5*sin(2*atan(1/3)))**2, Eq(c, -d*x + 2*atan(1/3))), (x/(3 - 5*sin(2*atan(3)))**2, Eq(c, -d*x + 2*atan(3))), (x/(3 - 5*sin(c))**2, Eq(d, 0)), (-27*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 90*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) - 27*log(tan(c/2 + d*x/2) - 3)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 27*log(3*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) - 90*log(3*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 27*log(3*tan(c/2 + d*x/2) - 1)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) - 200*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 120/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.28

$$\int \frac{1}{(3 - 5 \sin(c + dx))^2} dx$$

$$= \frac{\frac{40 \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} - 3 \right)}{10 \frac{\sin(dx+c)}{\cos(dx+c)+1} - \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 3} + 9 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right) - 9 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 3 \right)}{192 d}$$

input `integrate(1/(3-5*sin(d*x+c))^2,x, algorithm="maxima")`output `1/192*(40*(5*sin(d*x + c)/(cos(d*x + c) + 1) - 3)/(10*sin(d*x + c)/(cos(d*x + c) + 1) - 3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 3) + 9*log(3*sin(d*x + c)/(cos(d*x + c) + 1) - 1) - 9*log(sin(d*x + c)/(cos(d*x + c) + 1) - 3))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.90

$$\int \frac{1}{(3 - 5 \sin(c + dx))^2} dx =$$

$$\frac{40 \left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3 \right)}{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3} - 9 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right) + 9 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3 \right| \right)$$

$$\frac{\hspace{10em}}{192 d}$$

input `integrate(1/(3-5*sin(d*x+c))^2,x, algorithm="giac")`output `-1/192*(40*(5*tan(1/2*d*x + 1/2*c) - 3)/(3*tan(1/2*d*x + 1/2*c)^2 - 10*tan(1/2*d*x + 1/2*c) + 3) - 9*log(abs(3*tan(1/2*d*x + 1/2*c) - 1)) + 9*log(abs(tan(1/2*d*x + 1/2*c) - 3)))/d`

Mupad [B] (verification not implemented)

Time = 25.98 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.71

$$\int \frac{1}{(3 - 5 \sin(c + dx))^2} dx = \frac{3 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{5}{4}}{4}\right)}{32 d} - \frac{\frac{25 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{72} - \frac{5}{24}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - \frac{10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(1/(5*sin(c + d*x) - 3)^2,x)`output `(3*atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4))/(32*d) - ((25*tan(c/2 + (d*x)/2)/72 - 5/24)/(d*(tan(c/2 + (d*x)/2)^2 - (10*tan(c/2 + (d*x)/2))/3 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.09

$$\int \frac{1}{(3 - 5 \sin(c + dx))^2} dx = \frac{-20 \cos(dx + c) - 15 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) \sin(dx + c) + 9 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) + 15 \log\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64d(5 \sin(dx + c) - 3)}$$

input `int(1/(3-5*sin(d*x+c))^2,x)`output `(- 20*cos(c + d*x) - 15*log(tan((c + d*x)/2) - 3)*sin(c + d*x) + 9*log(tan((c + d*x)/2) - 3) + 15*log(3*tan((c + d*x)/2) - 1)*sin(c + d*x) - 9*log(3*tan((c + d*x)/2) - 1))/(64*d*(5*sin(c + d*x) - 3))`

3.66 $\int \frac{1}{(3-5 \sin(c+dx))^3} dx$

Optimal result	470
Mathematica [A] (verified)	470
Rubi [A] (verified)	471
Maple [A] (verified)	474
Fricas [A] (verification not implemented)	474
Sympy [B] (verification not implemented)	475
Maxima [A] (verification not implemented)	476
Giac [A] (verification not implemented)	476
Mupad [B] (verification not implemented)	477
Reduce [B] (verification not implemented)	477

Optimal result

Integrand size = 12, antiderivative size = 115

$$\int \frac{1}{(3-5 \sin(c+dx))^3} dx = -\frac{43 \log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{2048d} + \frac{43 \log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{2048d} + \frac{5 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} - \frac{45 \cos(c+dx)}{512d(3-5 \sin(c+dx))}$$

output

```
-43/2048*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d+43/2048*ln(3*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+5/32*cos(d*x+c)/d/(3-5*sin(d*x+c))^2-45/512*cos(d*x+c)/d/(3-5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.60

$$\int \frac{1}{(3-5 \sin(c+dx))^3} dx = -43 \log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx))) + 43 \log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx))) + \frac{5 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} - \frac{45 \cos(c+dx)}{512d(3-5 \sin(c+dx))}$$

input `Integrate[(3 - 5*Sin[c + d*x])^(-3),x]`

output `(-43*Log[Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]] + 43*Log[3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] + 40/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2])^2 + (-180/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]) - 60/(3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))*Sin[(c + d*x)/2] - 40/(-3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2)/(2048*d)`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.89, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3042, 3143, 25, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3 - 5 \sin(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(3 - 5 \sin(c + dx))^3} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{32} \int -\frac{5 \sin(c + dx) + 6}{(3 - 5 \sin(c + dx))^2} dx + \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} - \frac{1}{32} \int \frac{5 \sin(c + dx) + 6}{(3 - 5 \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} - \frac{1}{32} \int \frac{5 \sin(c + dx) + 6}{(3 - 5 \sin(c + dx))^2} dx \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left(-\frac{1}{16} \int -\frac{43}{3 - 5 \sin(c + dx)} dx - \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) + \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{32} \left(\frac{43}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx - \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) + \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 3042 \\
& \frac{1}{32} \left(\frac{43}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx - \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) + \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 3139 \\
& \frac{1}{32} \left(\frac{43 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} - \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) + \\
& \quad \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 1081 \\
& \frac{1}{32} \left(\frac{129 \int \left(\frac{1}{8(1 - 3 \tan(\frac{1}{2}(c+dx)))} - \frac{1}{24(3 - \tan(\frac{1}{2}(c+dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{8d} - \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) + \\
& \quad \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 2009 \\
& \frac{1}{32} \left(\frac{129 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx))) \right)}{8d} - \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) + \\
& \quad \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2}
\end{aligned}$$

input `Int[(3 - 5*Sin[c + d*x])^(-3),x]`

output `((129*(-1/24*Log[1 - 3*Tan[(c + d*x)/2]] + Log[3 - Tan[(c + d*x)/2]]/24))/(8*d) - (45*Cos[c + d*x])/(16*d*(3 - 5*Sin[c + d*x]))/32 + (5*Cos[c + d*x])/((32*d*(3 - 5*Sin[c + d*x])^2)`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 1081 $\text{Int}[(\text{a}_) + (\text{b}_.)*(x_) + (\text{c}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = \text{Rt}[\text{b}^2 - 4*\text{a}*c, 2]\}, \text{Simp}[\text{c} \quad \text{Int}[\text{ExpandIntegrand}[1/((\text{b}/2 - \text{q}/2 + \text{c}*x)*(\text{b}/2 + \text{q}/2 + \text{c}*x)), \text{x}], \text{x}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}] \ \&\& \ \text{NiceSqrtQ}[\text{b}^2 - 4*\text{a}*c]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ ; SumQ}[\text{u}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3139 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{e} = \text{FreeFactors}[\text{Tan}[(\text{c} + \text{d}*x)/2], \text{x}]\}, \text{Simp}[2*(\text{e}/\text{d}) \quad \text{Subst}[\text{Int}[1/(\text{a} + 2*\text{b}*e*x + \text{a}*e^2*x^2), \text{x}], \text{x}, \text{Tan}[(\text{c} + \text{d}*x)/2]/\text{e}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0]$
- rule 3143 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{c}_.) + (\text{d}_.)*(x_)]^{(\text{n}_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b})*\text{Cos}[\text{c} + \text{d}*x]*((\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(\text{n} + 1)}/(\text{d}*(\text{n} + 1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((\text{n} + 1)*(a^2 - b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{c} + \text{d}*x])^{(\text{n} + 1)}*\text{Simp}[\text{a}*(\text{n} + 1) - \text{b}*(\text{n} + 2)*\text{Sin}[\text{c} + \text{d}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{n}, -1] \ \&\& \ \text{IntegerQ}[2*\text{n}]$
- rule 3233 $\text{Int}[(\text{a}_) + (\text{b}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]^{(\text{m}_)}*((\text{c}_.) + (\text{d}_.)*\sin[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{b}*c - \text{a}*d))*\text{Cos}[\text{e} + \text{f}*x]*((\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(\text{m} + 1)}/(\text{f}*(\text{m} + 1)*(a^2 - b^2))), \text{x}] + \text{Simp}[1/((\text{m} + 1)*(a^2 - b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*x])^{(\text{m} + 1)}*\text{Simp}[(\text{a}*c - \text{b}*d)*(m + 1) - (\text{b}*c - \text{a}*d)*(m + 2)*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntegerQ}[2*\text{m}]$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.87

method	result
derivativedivides	$-\frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^2} - \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} + \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{2048} + \frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} + \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}$
default	$-\frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^2} - \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} + \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{2048} + \frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} + \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}$
risch	$-\frac{-387ie^{2i(dx+c)} + 215e^{3i(dx+c)} - 325e^{i(dx+c)} + 225i}{256(5e^{2i(dx+c)} - 5 - 6ie^{i(dx+c)})^2}d - \frac{43 \ln(-\frac{4}{5} - \frac{3i}{5} + e^{i(dx+c)})}{2048d} + \frac{43 \ln(e^{i(dx+c)} + \frac{4}{5} - \frac{3i}{5})}{2048d}$
norman	$-\frac{55}{256d} - \frac{3245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{2304d} + \frac{1225 \tan(\frac{dx}{2} + \frac{c}{2})}{768d} - \frac{125 \tan(\frac{dx}{2} + \frac{c}{2})^3}{768d} + \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{2048d} - \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{2048d}$
parallelrisc	$\frac{(-9675 \cos(2dx+2c) - 23220 \sin(dx+c) + 16641) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{3}) + (9675 \cos(2dx+2c) + 23220 \sin(dx+c) - 16641)}{18432d(-43 + 25 \cos(2dx+2c))}$

```
input int(1/(3-5*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-25/128/(tan(1/2*d*x+1/2*c)-3)^2-15/512/(tan(1/2*d*x+1/2*c)-3)+43/2048*ln(tan(1/2*d*x+1/2*c)-3)+25/1152/(3*tan(1/2*d*x+1/2*c)-1)^2+155/4608/(3*tan(1/2*d*x+1/2*c)-1)-43/2048*ln(3*tan(1/2*d*x+1/2*c)-1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.16

$$\int \frac{1}{(3 - 5 \sin(c + dx))^3} dx$$

$$= \frac{43 (25 \cos(dx + c)^2 + 30 \sin(dx + c) - 34) \log(4 \cos(dx + c) - 3 \sin(dx + c) + 5) - 43 (25 \cos(dx + c) + 30 \sin(dx + c) - 34)}{4096 (25 d \cos(dx + c) + 30 d \sin(dx + c) - 34)}$$

```
input integrate(1/(3-5*sin(d*x+c))^3,x, algorithm="fricas")
```

output

```
1/4096*(43*(25*cos(d*x + c)^2 + 30*sin(d*x + c) - 34)*log(4*cos(d*x + c) -
3*sin(d*x + c) + 5) - 43*(25*cos(d*x + c)^2 + 30*sin(d*x + c) - 34)*log(-
4*cos(d*x + c) - 3*sin(d*x + c) + 5) - 1800*cos(d*x + c)*sin(d*x + c) + 44
0*cos(d*x + c))/(25*d*cos(d*x + c)^2 + 30*d*sin(d*x + c) - 34*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1224 vs. $2(102) = 204$.

Time = 1.41 (sec) , antiderivative size = 1224, normalized size of antiderivative = 10.64

$$\int \frac{1}{(3 - 5 \sin(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(3-5*sin(d*x+c))**3,x)
```

output

```
Piecewise((x/(3 - 5*sin(2*atan(1/3)))**3, Eq(c, -d*x + 2*atan(1/3))), (x/(
3 - 5*sin(2*atan(3)))**3, Eq(c, -d*x + 2*atan(3))), (x/(3 - 5*sin(c))**3,
Eq(d, 0)), (3483*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**4/(165888*d*t
an(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d
*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 23220*log(tan(c/2 + d*
x/2) - 3)*tan(c/2 + d*x/2)**3/(165888*d*tan(c/2 + d*x/2)**4 - 1105920*d*t
an(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 - 1105920*d*tan(c/2 + d*
x/2) + 165888*d) + 45666*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**2/(16
5888*d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan
(c/2 + d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 23220*log(tan(
c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)/(165888*d*tan(c/2 + d*x/2)**4 - 1105920
*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 - 1105920*d*tan(c/2
+ d*x/2) + 165888*d) + 3483*log(tan(c/2 + d*x/2) - 3)/(165888*d*tan(c/2 +
d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2
- 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 3483*log(3*tan(c/2 + d*x/2) -
1)*tan(c/2 + d*x/2)**4/(165888*d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 +
d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) +
165888*d) + 23220*log(3*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**3/(165888*
d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2
+ d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 45666*log(3*tan(...
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.70

$$\int \frac{1}{(3 - 5 \sin(c + dx))^3} dx = \frac{40 \left(\frac{735 \sin(dx+c)}{\cos(dx+c)+1} - \frac{649 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{75 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - 99 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} - \frac{118 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{9 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 9} + 387 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right) - 387 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 1 \right)$$

18432 d

input `integrate(1/(3-5*sin(d*x+c))^3,x, algorithm="maxima")`

output `-1/18432*(40*(735*sin(d*x + c)/(cos(d*x + c) + 1) - 649*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 75*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 99)/(60*sin(d*x + c)/(cos(d*x + c) + 1) - 118*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 9*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - 9) + 387*log(3*sin(d*x + c)/(cos(d*x + c) + 1) - 1) - 387*log(sin(d*x + c)/(cos(d*x + c) + 1) - 3))/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.93

$$\int \frac{1}{(3 - 5 \sin(c + dx))^3} dx = \frac{40 \left(75 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 649 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 99 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^2} + 387 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right) - 387 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3 \right| \right)$$

18432 d

input `integrate(1/(3-5*sin(d*x+c))^3,x, algorithm="giac")`

output `-1/18432*(40*(75*tan(1/2*d*x + 1/2*c)^3 + 649*tan(1/2*d*x + 1/2*c)^2 - 735*tan(1/2*d*x + 1/2*c) + 99)/(3*tan(1/2*d*x + 1/2*c)^2 - 10*tan(1/2*d*x + 1/2*c) + 3)^2 + 387*log(abs(3*tan(1/2*d*x + 1/2*c) - 1)) - 387*log(abs(tan(1/2*d*x + 1/2*c) - 3)))/d`

Mupad [B] (verification not implemented)

Time = 27.20 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.01

$$\int \frac{1}{(3 - 5 \sin(c + dx))^3} dx$$

$$= -\frac{43 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c+dx}{2}\right)}{4} - \frac{5}{4}\right)}{1024 d}$$

$$- \frac{\frac{125 \tan\left(\frac{c+dx}{2}\right)^3}{6912} + \frac{3245 \tan\left(\frac{c+dx}{2}\right)^2}{20736} - \frac{1225 \tan\left(\frac{c+dx}{2}\right)}{6912} + \frac{55}{2304}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{3} + \frac{118 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{9} - \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(-1/(5*sin(c + d*x) - 3)^3,x)`output `- (43*atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4))/(1024*d) - ((3245*tan(c/2 + (d*x)/2)^2)/20736 - (1225*tan(c/2 + (d*x)/2))/6912 + (125*tan(c/2 + (d*x)/2)^3)/6912 + 55/2304)/(d*((118*tan(c/2 + (d*x)/2)^2)/9 - (20*tan(c/2 + (d*x)/2))/3 - (20*tan(c/2 + (d*x)/2)^3)/3 + tan(c/2 + (d*x)/2)^4 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.63

$$\int \frac{1}{(3 - 5 \sin(c + dx))^3} dx$$

$$= \frac{900 \cos(dx + c) \sin(dx + c) - 220 \cos(dx + c) + 1075 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) \sin(dx + c)^2 - 1290 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{(3 - 5 \sin(dx + c))^3}$$

input `int(1/(3-5*sin(d*x+c))^3,x)`

output

```
(900*cos(c + d*x)*sin(c + d*x) - 220*cos(c + d*x) + 1075*log(tan((c + d*x)/2) - 3)*sin(c + d*x)**2 - 1290*log(tan((c + d*x)/2) - 3)*sin(c + d*x) + 387*log(tan((c + d*x)/2) - 3) - 1075*log(3*tan((c + d*x)/2) - 1)*sin(c + d*x)**2 + 1290*log(3*tan((c + d*x)/2) - 1)*sin(c + d*x) - 387*log(3*tan((c + d*x)/2) - 1) - 750*sin(c + d*x)**2 + 900*sin(c + d*x) - 270)/(2048*d*(25*sin(c + d*x)**2 - 30*sin(c + d*x) + 9))
```

3.67 $\int \frac{1}{(3-5 \sin(c+dx))^4} dx$

Optimal result	479
Mathematica [A] (verified)	480
Rubi [A] (verified)	480
Maple [A] (verified)	484
Fricas [A] (verification not implemented)	484
Sympy [B] (verification not implemented)	485
Maxima [B] (verification not implemented)	486
Giac [A] (verification not implemented)	486
Mupad [B] (verification not implemented)	487
Reduce [B] (verification not implemented)	487

Optimal result

Integrand size = 12, antiderivative size = 140

$$\int \frac{1}{(3-5 \sin(c+dx))^4} dx = \frac{279 \log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{32768d} - \frac{279 \log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{32768d} + \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} - \frac{25 \cos(c+dx)}{512d(3-5 \sin(c+dx))^2} + \frac{995 \cos(c+dx)}{24576d(3-5 \sin(c+dx))}$$

output

```
279/32768*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d-279/32768*ln(3*cos
(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+5/48*cos(d*x+c)/d/(3-5*sin(d*x+c))^3
-25/512*cos(d*x+c)/d/(3-5*sin(d*x+c))^2+995/24576*cos(d*x+c)/d/(3-5*sin(d*
x+c))
```


Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.72

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx$$

$$= \frac{2511 \log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx))) - 2511 \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) - \frac{1}{\cos(\frac{1}{2}(c + dx))}}{\dots}$$

input `Integrate[(3 - 5*Sin[c + d*x])^(-4), x]`

output `(2511*Log[Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]] - 2511*Log[3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - 720/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2])^2 + 20*(240/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2])^3 + 597/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]) + 80/(3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^3 + 199/(3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))*Sin[(c + d*x)/2] + 2320/(-3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2)/(294912*d)`

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.94, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$, Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{10 \sin(c + dx) + 9}{(3 - 5 \sin(c + dx))^3} dx + \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} - \frac{1}{48} \int \frac{10 \sin(c+dx) + 9}{(3-5 \sin(c+dx))^3} dx \\
& \downarrow 3042 \\
& \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} - \frac{1}{48} \int \frac{10 \sin(c+dx) + 9}{(3-5 \sin(c+dx))^3} dx \\
& \downarrow 3233 \\
& \frac{1}{48} \left(-\frac{1}{32} \int -\frac{75 \sin(c+dx) + 154}{(3-5 \sin(c+dx))^2} dx - \frac{75 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} \right) + \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{75 \sin(c+dx) + 154}{(3-5 \sin(c+dx))^2} dx - \frac{75 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} \right) + \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{75 \sin(c+dx) + 154}{(3-5 \sin(c+dx))^2} dx - \frac{75 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} \right) + \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{1}{16} \int -\frac{837}{3-5 \sin(c+dx)} dx + \frac{995 \cos(c+dx)}{16d(3-5 \sin(c+dx))} \right) - \frac{75 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} \right) + \\
& \quad \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c+dx)}{16d(3-5 \sin(c+dx))} - \frac{837}{16} \int \frac{1}{3-5 \sin(c+dx)} dx \right) - \frac{75 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} \right) + \\
& \quad \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c+dx)}{16d(3-5 \sin(c+dx))} - \frac{837}{16} \int \frac{1}{3-5 \sin(c+dx)} dx \right) - \frac{75 \cos(c+dx)}{32d(3-5 \sin(c+dx))^2} \right) + \\
& \quad \frac{5 \cos(c+dx)}{48d(3-5 \sin(c+dx))^3} \\
& \downarrow 3139
\end{aligned}$$

$$\frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{837 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} \right) - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))} \right) - \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3}$$

↓ 1081

$$\frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{2511 \int \left(\frac{1}{8(1 - 3 \tan(\frac{1}{2}(c+dx)))} - \frac{1}{24(3 - \tan(\frac{1}{2}(c+dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{8d} \right) - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))} \right) - \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3}$$

↓ 2009

$$\frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{2511 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx))) \right)}{8d} \right) - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))} \right) - \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} +$$

input `Int[(3 - 5*Sin[c + d*x])^(-4),x]`

output `((((-2511*(-1/24*Log[1 - 3*Tan[(c + d*x)/2]] + Log[3 - Tan[(c + d*x)/2]]/24)))/(8*d) + (995*Cos[c + d*x])/(16*d*(3 - 5*Sin[c + d*x])))/32 - (75*Cos[c + d*x])/(32*d*(3 - 5*Sin[c + d*x]^2))/48 + (5*Cos[c + d*x])/(48*d*(3 - 5*Sin[c + d*x]^3))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 1081 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3139 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.94

method	result
derivativedivides	$-\frac{125}{768(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^3} - \frac{75}{1024(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^2} - \frac{345}{8192(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{279 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{32768} - \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^3} + \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{32768} + \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} - \frac{345}{8192(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{32768} - \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}$
default	$-\frac{125}{768(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^3} - \frac{75}{1024(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^2} - \frac{345}{8192(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{279 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{32768} - \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^3} + \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{32768} + \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} - \frac{345}{8192(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{32768} - \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}$
risch	$-\frac{111042 e^{3i(dx+c)} - 62775 i e^{4i(dx+c)} + 119310 i e^{2i(dx+c)} + 20925 e^{5i(dx+c)} + 68625 e^{i(dx+c)} - 24875 i}{12288(5 e^{2i(dx+c)} - 5 - 6 i e^{i(dx+c)})^3} d + \frac{279 \ln(-\frac{4}{5} - \frac{3i}{5})}{32768}$
norman	$\frac{7915}{12288d} - \frac{15725 \tan(\frac{dx}{2} + \frac{c}{2})^5}{12288d} - \frac{3047275 \tan(\frac{dx}{2} + \frac{c}{2})^3}{165888d} - \frac{63425 \tan(\frac{dx}{2} + \frac{c}{2})}{12288d} + \frac{296245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{18432d} + \frac{270245 \tan(\frac{dx}{2} + \frac{c}{2})^4}{36864d} - \frac{279 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{32768} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{32768} - \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^3} + \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} - \frac{345}{8192(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{32768} - \frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}$
parallelrisc	$\frac{(10169550 \cos(2dx+2c) + 20678085 \sin(dx+c) - 2824875 \sin(3dx+3c) - 12610242) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{3}) + (-10169550 \cos(dx+c) + 20678085 \sin(2dx+c) - 2824875 \sin(dx+c) - 12610242) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{12288(5 e^{2i(dx+c)} - 5 - 6 i e^{i(dx+c)})^3} d$

```
input int(1/(3-5*sin(d*x+c))^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(-125/768/(tan(1/2*d*x+1/2*c)-3)^3-75/1024/(tan(1/2*d*x+1/2*c)-3)^2-34
5/8192/(tan(1/2*d*x+1/2*c)-3)-279/32768*ln(tan(1/2*d*x+1/2*c)-3)-125/20736
/(3*tan(1/2*d*x+1/2*c)-1)^3-275/27648/(3*tan(1/2*d*x+1/2*c)-1)^2-3505/2211
84/(3*tan(1/2*d*x+1/2*c)-1)+279/32768*ln(3*tan(1/2*d*x+1/2*c)-1))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.29

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx$$

$$= \frac{199000 \cos(dx + c)^3 - 837 (225 \cos(dx + c)^2 - 5 (25 \cos(dx + c)^2 - 52) \sin(dx + c) - 252) \log(4 \cos(dx + c) - 2)}{12288(5 \cos^2(dx + c) - 5 \cos(dx + c) - 2)^3}$$

```
input integrate(1/(3-5*sin(d*x+c))^4,x, algorithm="fricas")
```

output

```
1/196608*(199000*cos(d*x + c)^3 - 837*(225*cos(d*x + c)^2 - 5*(25*cos(d*x
+ c)^2 - 52)*sin(d*x + c) - 252)*log(4*cos(d*x + c) - 3*sin(d*x + c) + 5)
+ 837*(225*cos(d*x + c)^2 - 5*(25*cos(d*x + c)^2 - 52)*sin(d*x + c) - 252)
*log(-4*cos(d*x + c) - 3*sin(d*x + c) + 5) + 190800*cos(d*x + c)*sin(d*x +
c) - 262320*cos(d*x + c))/(225*d*cos(d*x + c)^2 - 5*(25*d*cos(d*x + c)^2
- 52*d)*sin(d*x + c) - 252*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2353 vs. $2(126) = 252$.

Time = 3.25 (sec) , antiderivative size = 2353, normalized size of antiderivative = 16.81

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(3-5*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(3 - 5*sin(2*atan(1/3)))**4, Eq(c, -d*x + 2*atan(1/3))), (x/(
3 - 5*sin(2*atan(3)))**4, Eq(c, -d*x + 2*atan(3))), (x/(3 - 5*sin(c))**4,
Eq(d, 0)), (-610173*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**6/(7166361
6*d*tan(c/2 + d*x/2)**6 - 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*t
an(c/2 + d*x/2)**4 - 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c
/2 + d*x/2)**2 - 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 6101730*log(
tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**5/(71663616*d*tan(c/2 + d*x/2)**6
- 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 - 408
7480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 - 7166361
60*d*tan(c/2 + d*x/2) + 71663616*d) - 22169619*log(tan(c/2 + d*x/2) - 3)*t
an(c/2 + d*x/2)**4/(71663616*d*tan(c/2 + d*x/2)**6 - 716636160*d*tan(c/2 +
d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 - 4087480320*d*tan(c/2 + d*x
/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 - 716636160*d*tan(c/2 + d*x/2) +
71663616*d) + 34802460*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**3/(716
63616*d*tan(c/2 + d*x/2)**6 - 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048
*d*tan(c/2 + d*x/2)**4 - 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*t
an(c/2 + d*x/2)**2 - 716636160*d*tan(c/2 + d*x/2) + 71663616*d) - 22169619
*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**2/(71663616*d*tan(c/2 + d*x/2
)**6 - 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4
- 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 - ...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(126) = 252$.

Time = 0.04 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.96

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx$$

$$= \frac{40 \left(\frac{342495 \sin(dx+c)}{\cos(dx+c)+1} - \frac{1066482 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1218910 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{486441 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{84915 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - 42741 \right) + 22599 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} \right)}{\frac{270 \sin(dx+c)}{\cos(dx+c)+1} - \frac{981 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1540 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{981 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{270 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - \frac{27 \sin(dx+c)^6}{(\cos(dx+c)+1)^6} - 27} 2654208 d}$$

input `integrate(1/(3-5*sin(d*x+c))^4,x, algorithm="maxima")`

output `1/2654208*(40*(342495*sin(d*x + c)/(cos(d*x + c) + 1) - 1066482*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1218910*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 486441*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 84915*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 42741)/(270*sin(d*x + c)/(cos(d*x + c) + 1) - 981*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1540*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 981*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 270*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 - 27*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 - 27) + 22599*log(3*sin(d*x + c)/(cos(d*x + c) + 1) - 1) - 22599*log(sin(d*x + c)/(cos(d*x + c) + 1) - 3))/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.95

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx =$$

$$\frac{40 \left(84915 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 486441 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 1218910 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 1066482 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 342495 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 42741 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^3} + 22599 \log \left(\frac{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1}{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3} \right)$$

2654208 d

input `integrate(1/(3-5*sin(d*x+c))^4,x, algorithm="giac")`

output

```
-1/2654208*(40*(84915*tan(1/2*d*x + 1/2*c)^5 - 486441*tan(1/2*d*x + 1/2*c)
^4 + 1218910*tan(1/2*d*x + 1/2*c)^3 - 1066482*tan(1/2*d*x + 1/2*c)^2 + 342
495*tan(1/2*d*x + 1/2*c) - 42741)/(3*tan(1/2*d*x + 1/2*c)^2 - 10*tan(1/2*d
*x + 1/2*c) + 3)^3 - 22599*log(abs(3*tan(1/2*d*x + 1/2*c) - 1)) + 22599*lo
g(abs(tan(1/2*d*x + 1/2*c) - 3)))/d
```

Mupad [B] (verification not implemented)

Time = 28.21 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.20

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx = \frac{279 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{5}{4}}{4}\right)}{16384 d} - \frac{\frac{15725 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{331776} - \frac{270245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{995328} + \frac{3047275 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{4478976} - \frac{296245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{497664} + \frac{63425 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{331776} - \frac{7915}{331776}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{3} - \frac{1540 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{27} + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{3} - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1 \right)}$$

input

```
int(1/(5*sin(c + d*x) - 3)^4,x)
```

output

```
(279*atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4))/(16384*d) - ((63425*tan(c/2 +
(d*x)/2))/331776 - (296245*tan(c/2 + (d*x)/2)^2)/497664 + (3047275*tan(c/2
+ (d*x)/2)^3)/4478976 - (270245*tan(c/2 + (d*x)/2)^4)/995328 + (15725*tan
(c/2 + (d*x)/2)^5)/331776 - 7915/331776)/(d*((109*tan(c/2 + (d*x)/2)^2)/3
- 10*tan(c/2 + (d*x)/2) - (1540*tan(c/2 + (d*x)/2)^3)/27 + (109*tan(c/2 +
(d*x)/2)^4)/3 - 10*tan(c/2 + (d*x)/2)^5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.92

$$\int \frac{1}{(3 - 5 \sin(c + dx))^4} dx = \frac{-298500 \cos(dx + c) \sin(dx + c)^2 + 286200 \cos(dx + c) \sin(dx + c) - 94980 \cos(dx + c) - 313875 \log\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{5}{4}}{4}\right)}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{3} - \frac{1540 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{27} + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{3} - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1 \right)}$$

input `int(1/(3-5*sin(d*x+c))^4,x)`

output
$$\left(-298500 \cos(c + dx) \sin(c + dx)^2 + 286200 \cos(c + dx) \sin(c + dx) - 94980 \cos(c + dx) - 313875 \log(\tan((c + dx)/2) - 3) \sin(c + dx)^3 + 564975 \log(\tan((c + dx)/2) - 3) \sin(c + dx)^2 - 338985 \log(\tan((c + dx)/2) - 3) \sin(c + dx) + 67797 \log(\tan((c + dx)/2) - 3) + 313875 \log(3 \tan((c + dx)/2) - 1) \sin(c + dx)^3 - 564975 \log(3 \tan((c + dx)/2) - 1) \sin(c + dx)^2 + 338985 \log(3 \tan((c + dx)/2) - 1) \sin(c + dx) - 67797 \log(3 \tan((c + dx)/2) - 1) + 265000 \sin(c + dx)^3 - 477000 \sin(c + dx)^2 + 286200 \sin(c + dx) - 57240 \right) / (294912 d (125 \sin(c + dx)^3 - 225 \sin(c + dx)^2 + 135 \sin(c + dx) - 27))$$

3.68 $\int \frac{1}{-3+5 \sin(c+dx)} dx$

Optimal result	489
Mathematica [A] (verified)	489
Rubi [A] (verified)	490
Maple [A] (verified)	491
Fricas [A] (verification not implemented)	492
Sympy [A] (verification not implemented)	492
Maxima [A] (verification not implemented)	492
Giac [A] (verification not implemented)	493
Mupad [B] (verification not implemented)	493
Reduce [B] (verification not implemented)	494

Optimal result

Integrand size = 12, antiderivative size = 65

$$\int \frac{1}{-3+5 \sin(c+dx)} dx = \frac{\log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{4d} - \frac{\log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{4d}$$

output

```
1/4*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d-1/4*ln(3*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00

$$\int \frac{1}{-3+5 \sin(c+dx)} dx = \frac{\log(\cos(\frac{1}{2}(c+dx)) - 3 \sin(\frac{1}{2}(c+dx)))}{4d} - \frac{\log(3 \cos(\frac{1}{2}(c+dx)) - \sin(\frac{1}{2}(c+dx)))}{4d}$$

input

```
Integrate[(-3 + 5*Sin[c + d*x])^(-1),x]
```

output

$$\frac{\text{Log}[\text{Cos}[(c + d*x)/2] - 3*\text{Sin}[(c + d*x)/2]]}{(4*d)} - \frac{\text{Log}[3*\text{Cos}[(c + d*x)/2] - \text{Sin}[(c + d*x)/2]]}{(4*d)}$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{5 \sin(c + dx) - 3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{5 \sin(c + dx) - 3} dx \\ & \quad \downarrow \text{3139} \\ & \frac{2 \int \frac{1}{-3 \tan^2(\frac{1}{2}(c+dx)) + 10 \tan(\frac{1}{2}(c+dx)) - 3} d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{1081} \\ & \frac{6 \int \left(\frac{1}{8(1-3 \tan(\frac{1}{2}(c+dx)))} - \frac{1}{24(3-\tan(\frac{1}{2}(c+dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{6(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx)))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx)))}{d} \end{aligned}$$

input

$$\text{Int}[(-3 + 5*\text{Sin}[c + d*x])^{-1}, x]$$

output

$$(-6*(-1/24*\text{Log}[1 - 3*\text{Tan}[(c + d*x)/2]] + \text{Log}[3 - \text{Tan}[(c + d*x)/2]]/24))/d$$

Definitions of rubi rules used

rule 1081 $\text{Int}[\{(a_)+(b_)*(x_)+(c_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c \text{ Int}[\text{ExpandIntegrand}[1/\{(b/2 - q/2 + c*x)\}(b/2 + q/2 + c*x)\}, x], x], x]] \text{ /; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NiceSqrtQ}[b^2 - 4*a*c]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3139 $\text{Int}[\{(a_)+(b_)*\sin[(c_)+(d_)*(x_)]\}^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] \text{ /; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{4} + \frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	36
default	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{4} + \frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	36
parallelrisc	$\frac{-\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 9\right) + \ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	37
norman	$-\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{4d} + \frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{4d}$	38
risc	$-\frac{\ln\left(e^{i(dx+c)} + \frac{4}{5} - \frac{3i}{5}\right)}{4d} + \frac{\ln\left(-\frac{4}{5} - \frac{3i}{5} + e^{i(dx+c)}\right)}{4d}$	40

input $\text{int}(1/(-3+5*\sin(d*x+c)), x, \text{method}=_RETURNVERBOSE)$

output $1/d*(-1/4*\ln(\tan(1/2*d*x+1/2*c)-3)+1/4*\ln(3*\tan(1/2*d*x+1/2*c)-1))$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.71

$$\int \frac{1}{-3 + 5 \sin(c + dx)} dx = \frac{\log(4 \cos(dx + c) - 3 \sin(dx + c) + 5) - \log(-4 \cos(dx + c) - 3 \sin(dx + c) + 5)}{8d}$$

input `integrate(1/(-3+5*sin(d*x+c)),x, algorithm="fricas")`output `-1/8*(log(4*cos(d*x + c) - 3*sin(d*x + c) + 5) - log(-4*cos(d*x + c) - 3*sin(d*x + c) + 5))/d`**Sympy [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \frac{1}{-3 + 5 \sin(c + dx)} dx = \begin{cases} -\frac{\log(\tan(\frac{c}{2} + \frac{dx}{2}) - 3)}{4d} + \frac{\log(3 \tan(\frac{c}{2} + \frac{dx}{2}) - 1)}{4d} & \text{for } d \neq 0 \\ \frac{x}{5 \sin(c) - 3} & \text{otherwise} \end{cases}$$

input `integrate(1/(-3+5*sin(d*x+c)),x)`output `Piecewise((-log(tan(c/2 + d*x/2) - 3)/(4*d) + log(3*tan(c/2 + d*x/2) - 1)/(4*d), Ne(d, 0)), (x/(5*sin(c) - 3), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.75

$$\int \frac{1}{-3 + 5 \sin(c + dx)} dx = \frac{\log\left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 1\right) - \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 3\right)}{4d}$$

input `integrate(1/(-3+5*sin(d*x+c)),x, algorithm="maxima")`

output $\frac{1}{4} * (\log(3 * \sin(d * x + c) / (\cos(d * x + c) + 1) - 1) - \log(\sin(d * x + c) / (\cos(d * x + c) + 1) - 3)) / d$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.55

$$\int \frac{1}{-3 + 5 \sin(c + dx)} dx = \frac{\log(|3 \tan(\frac{1}{2} dx + \frac{1}{2} c) - 1|) - \log(|\tan(\frac{1}{2} dx + \frac{1}{2} c) - 3|)}{4d}$$

input `integrate(1/(-3+5*sin(d*x+c)),x, algorithm="giac")`

output $\frac{1}{4} * (\log(\text{abs}(3 * \tan(1/2 * d * x + 1/2 * c) - 1)) - \log(\text{abs}(\tan(1/2 * d * x + 1/2 * c) - 3))) / d$

Mupad [B] (verification not implemented)

Time = 25.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.29

$$\int \frac{1}{-3 + 5 \sin(c + dx)} dx = \frac{\operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{5}{4}\right)}{2d}$$

input `int(1/(5*sin(c + d*x) - 3),x)`

output `atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4)/(2*d)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.52

$$\int \frac{1}{-3 + 5 \sin(c + dx)} dx = \frac{-\log(\tan(\frac{dx}{2} + \frac{c}{2}) - 3) + \log(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{4d}$$

input `int(1/(-3+5*sin(d*x+c)),x)`

output `(- log(tan((c + d*x)/2) - 3) + log(3*tan((c + d*x)/2) - 1))/(4*d)`

3.69 $\int \frac{1}{(-3+5 \sin(c+dx))^2} dx$

Optimal result	495
Mathematica [A] (verified)	495
Rubi [A] (verified)	496
Maple [A] (verified)	498
Fricas [A] (verification not implemented)	498
Sympy [B] (verification not implemented)	499
Maxima [A] (verification not implemented)	500
Giac [A] (verification not implemented)	500
Mupad [B] (verification not implemented)	501
Reduce [B] (verification not implemented)	501

Optimal result

Integrand size = 12, antiderivative size = 90

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx = \frac{3 \log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{3 \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{64d} + \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))}$$

output 3/64*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d-3/64*ln(3*cos(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+5/16*cos(d*x+c)/d/(3-5*sin(d*x+c))

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.44

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx = \frac{9(\log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx))) - \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))))}{192d} + 20 \left(\frac{1}{\cos(\frac{1}{2}(c + dx))} \right)$$

input `Integrate[(-3 + 5*Sin[c + d*x])^(-2),x]`

output `(9*(Log[Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]] - Log[3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]]) + 20*(3/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]) + (3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^(-1))*Sin[(c + d*x)/2])/(192*d)`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.80, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 3143, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 \sin(c + dx) - 3)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 \sin(c + dx) - 3)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{16} \int -\frac{3}{3 - 5 \sin(c + dx)} dx + \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{3}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{3}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \\
 & \quad \downarrow \text{3139} \\
 & \frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{3 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} \\
 & \quad \downarrow \text{1081}
 \end{aligned}$$

$$\frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{9 \int \left(\frac{1}{8(1 - 3 \tan(\frac{1}{2}(c + dx)))} - \frac{1}{24(3 - \tan(\frac{1}{2}(c + dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{8d}$$

↓ 2009

$$\frac{5 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{9 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx))) \right)}{8d}$$

input `Int[(-3 + 5*Sin[c + d*x])^(-2),x]`

output `(-9*(-1/24*Log[1 - 3*Tan[(c + d*x)/2]] + Log[3 - Tan[(c + d*x)/2]]/24))/(8*d) + (5*Cos[c + d*x])/(16*d*(3 - 5*Sin[c + d*x]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.76

method	result
derivativedivides	$\frac{-\frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{64} - \frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} + \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{64}}{d}$
default	$\frac{-\frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{64} - \frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} + \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{64}}{d}$
risch	$\frac{3 e^{i(dx+c)} - 5i}{8d(5 e^{2i(dx+c)} - 5 - 6i e^{i(dx+c)})} - \frac{3 \ln(e^{i(dx+c)} + \frac{4}{5} - \frac{3i}{5})}{64d} + \frac{3 \ln(-\frac{4}{5} - \frac{3i}{5} + e^{i(dx+c)})}{64d}$
norman	$\frac{\frac{5}{8d} - \frac{25 \tan(\frac{dx}{2} + \frac{c}{2})}{24d}}{3 \tan(\frac{dx}{2} + \frac{c}{2})^2 - 10 \tan(\frac{dx}{2} + \frac{c}{2}) + 3} - \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{64d} + \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{64d}$
parallelrisch	$\frac{-45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3) \sin(dx+c) + 45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{3}) \sin(dx+c) + 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3) - 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}))}{192d(-3 + 5 \sin(dx+c))}$

input

```
int(1/(-3+5*sin(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-5/16/(tan(1/2*d*x+1/2*c)-3)-3/64*ln(tan(1/2*d*x+1/2*c)-3)-5/48/(3*tan(1/2*d*x+1/2*c)-1)+3/64*ln(3*tan(1/2*d*x+1/2*c)-1))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.98

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx = \frac{3(5 \sin(dx + c) - 3) \log(4 \cos(dx + c) - 3 \sin(dx + c) + 5) - 3(5 \sin(dx + c) - 3) \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5)}{128(5d \sin(dx + c) - 3d)}$$

input `integrate(1/(-3+5*sin(d*x+c))^2,x, algorithm="fricas")`

output `-1/128*(3*(5*sin(d*x + c) - 3)*log(4*cos(d*x + c) - 3*sin(d*x + c) + 5) - 3*(5*sin(d*x + c) - 3)*log(-4*cos(d*x + c) - 3*sin(d*x + c) + 5) + 40*cos(d*x + c))/(5*d*sin(d*x + c) - 3*d)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 462 vs. $2(78) = 156$.

Time = 0.75 (sec) , antiderivative size = 462, normalized size of antiderivative = 5.13

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(-3+5*sin(d*x+c))**2,x)`

output `Piecewise((x/(-3 + 5*sin(2*atan(1/3)))**2, Eq(c, -d*x + 2*atan(1/3))), (x/(-3 + 5*sin(2*atan(3)))**2, Eq(c, -d*x + 2*atan(3))), (x/(5*sin(c) - 3)**2, Eq(d, 0)), (-27*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 90*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) - 27*log(tan(c/2 + d*x/2) - 3)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 27*log(3*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) - 90*log(3*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 27*log(3*tan(c/2 + d*x/2) - 1)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) - 200*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d) + 120/(576*d*tan(c/2 + d*x/2)**2 - 1920*d*tan(c/2 + d*x/2) + 576*d), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.28

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx$$

$$= \frac{\frac{40 \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} - 3 \right)}{10 \frac{\sin(dx+c)}{\cos(dx+c)+1} - \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - 3} + 9 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right) - 9 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 3 \right)}{192 d}$$

input `integrate(1/(-3+5*sin(d*x+c))^2,x, algorithm="maxima")`output `1/192*(40*(5*sin(d*x + c)/(cos(d*x + c) + 1) - 3)/(10*sin(d*x + c)/(cos(d*x + c) + 1) - 3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 3) + 9*log(3*sin(d*x + c)/(cos(d*x + c) + 1) - 1) - 9*log(sin(d*x + c)/(cos(d*x + c) + 1) - 3))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.90

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx =$$

$$\frac{40 \left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3 \right)}{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3} - 9 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right) + 9 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3 \right| \right)$$

$$\frac{\hspace{10em}}{192 d}$$

input `integrate(1/(-3+5*sin(d*x+c))^2,x, algorithm="giac")`output `-1/192*(40*(5*tan(1/2*d*x + 1/2*c) - 3)/(3*tan(1/2*d*x + 1/2*c)^2 - 10*tan(1/2*d*x + 1/2*c) + 3) - 9*log(abs(3*tan(1/2*d*x + 1/2*c) - 1)) + 9*log(abs(tan(1/2*d*x + 1/2*c) - 3)))/d`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.71

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx = \frac{3 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - \frac{5}{4}}{4}\right)}{32 d} - \frac{\frac{25 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{72} - \frac{5}{24}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - \frac{10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(1/(5*sin(c + d*x) - 3)^2,x)`output `(3*atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4))/(32*d) - ((25*tan(c/2 + (d*x)/2)/72 - 5/24)/(d*(tan(c/2 + (d*x)/2)^2 - (10*tan(c/2 + (d*x)/2))/3 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.09

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^2} dx = \frac{-20 \cos(dx + c) - 15 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) \sin(dx + c) + 9 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) + 15 \log\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)}{64d(5 \sin(dx + c) - 3)}$$

input `int(1/(-3+5*sin(d*x+c))^2,x)`output `(- 20*cos(c + d*x) - 15*log(tan((c + d*x)/2) - 3)*sin(c + d*x) + 9*log(tan((c + d*x)/2) - 3) + 15*log(3*tan((c + d*x)/2) - 1)*sin(c + d*x) - 9*log(3*tan((c + d*x)/2) - 1))/(64*d*(5*sin(c + d*x) - 3))`

3.70 $\int \frac{1}{(-3+5 \sin(c+dx))^3} dx$

Optimal result	502
Mathematica [A] (verified)	502
Rubi [A] (verified)	503
Maple [A] (verified)	506
Fricas [A] (verification not implemented)	506
Sympy [B] (verification not implemented)	507
Maxima [A] (verification not implemented)	508
Giac [A] (verification not implemented)	508
Mupad [B] (verification not implemented)	509
Reduce [B] (verification not implemented)	509

Optimal result

Integrand size = 12, antiderivative size = 115

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx = \frac{43 \log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{43 \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} + \frac{45 \cos(c + dx)}{512d(3 - 5 \sin(c + dx))}$$

```
output 43/2048*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d-43/2048*ln(3*cos(1/2
*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d-5/32*cos(d*x+c)/d/(3-5*sin(d*x+c))^2+45/
512*cos(d*x+c)/d/(3-5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.59

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx = \frac{43 \log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx))) - 43 \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} + \frac{45 \cos(c + dx)}{512d(3 - 5 \sin(c + dx))}}{\cos(\frac{1}{2}(c + dx))}$$

input `Integrate[(-3 + 5*Sin[c + d*x])^(-3),x]`

output `(43*Log[Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]] - 43*Log[3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - 40/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2])^2 + 60*(3/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]) + (3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^(-1))*Sin[(c + d*x)/2] + 40/(-3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2)/(2048*d)`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.89, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(5 \sin(c + dx) - 3)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(5 \sin(c + dx) - 3)^3} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{32} \int \frac{5 \sin(c + dx) + 6}{(3 - 5 \sin(c + dx))^2} dx - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{32} \int \frac{5 \sin(c + dx) + 6}{(3 - 5 \sin(c + dx))^2} dx - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
 & \quad \downarrow \text{3233} \\
 & \frac{1}{32} \left(\frac{1}{16} \int -\frac{43}{3 - 5 \sin(c + dx)} dx + \frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{43}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \right) - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{43}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \right) - \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 3139 \\
& \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{43 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} \right) - \\
& \quad \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 1081 \\
& \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{129 \int \left(\frac{1}{8(1 - 3 \tan(\frac{1}{2}(c+dx)))} - \frac{1}{24(3 - \tan(\frac{1}{2}(c+dx)))} \right) d \tan(\frac{1}{2}(c + dx))}{8d} \right) - \\
& \quad \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \\
& \downarrow 2009 \\
& \frac{1}{32} \left(\frac{45 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{129 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c + dx))) \right) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c + dx)))}{8d} \right) - \\
& \quad \frac{5 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2}
\end{aligned}$$

input `Int[(-3 + 5*Sin[c + d*x])^(-3),x]`

output `((-129*(-1/24*Log[1 - 3*Tan[(c + d*x)/2]] + Log[3 - Tan[(c + d*x)/2]]/24)) / (8*d) + (45*Cos[c + d*x]) / (16*d*(3 - 5*Sin[c + d*x]))) / 32 - (5*Cos[c + d*x]) / (32*d*(3 - 5*Sin[c + d*x])^2)`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 1081 $\text{Int}[((a_) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c \text{ Int}[\text{ExpandIntegrand}[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NiceSqrtQ}[b^2 - 4*a*c]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3139 $\text{Int}[((a_) + (b_*)\sin[(c_) + (d_*)(x_)])^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 3143 $\text{Int}[((a_) + (b_*)\sin[(c_) + (d_*)(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^{(n + 1)}/(d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\text{Sin}[c + d*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 3233 $\text{Int}[((a_) + (b_*)\sin[(e_) + (f_*)(x_)])^{(m_)}*((c_) + (d_*)\sin[(e_) + (f_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*\text{Cos}[e + f*x]*((a + b*\text{Sin}[e + f*x])^{(m + 1)}/(f*(m + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((m + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[e + f*x])^{(m + 1)}*\text{Simp}[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegerQ}[2*m]$

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{\frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^2} + \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{2048} - \frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} - \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} + \dots}{d}$
default	$\frac{\frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)^2} + \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)} - \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{2048} - \frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)^2} - \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)} + \dots}{d}$
risch	$\frac{-387ie^{2i(dx+c)} + 215e^{3i(dx+c)} - 325e^{i(dx+c)} + 225i}{256(5e^{2i(dx+c)} - 5 - 6ie^{i(dx+c)})^2 d} - \frac{43 \ln(e^{i(dx+c)} + \frac{4}{5} - \frac{3i}{5})}{2048d} + \frac{43 \ln(-\frac{4}{5} - \frac{3i}{5} + e^{i(dx+c)})}{2048d}$
norman	$\frac{\frac{55}{256d} + \frac{3245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{2304d} + \frac{125 \tan(\frac{dx}{2} + \frac{c}{2})^3}{768d} - \frac{1225 \tan(\frac{dx}{2} + \frac{c}{2})}{768d}}{(3 \tan(\frac{dx}{2} + \frac{c}{2})^2 - 10 \tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} - \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - 3)}{2048d} + \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) - 1)}{2048d}$
parallelrisc	$\frac{(9675 \cos(2dx+2c) + 23220 \sin(dx+c) - 16641) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{3}) + (-9675 \cos(2dx+2c) - 23220 \sin(dx+c) + 16641)}{18432d(-43 + 25 \cos(2dx+2c) - \dots)}$

```
input int(1/(-3+5*sin(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/d*(25/128/(tan(1/2*d*x+1/2*c)-3)^2+15/512/(tan(1/2*d*x+1/2*c)-3)-43/2048
*ln(tan(1/2*d*x+1/2*c)-3)-25/1152/(3*tan(1/2*d*x+1/2*c)-1)^2-155/4608/(3*t
an(1/2*d*x+1/2*c)-1)+43/2048*ln(3*tan(1/2*d*x+1/2*c)-1))
```

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.16

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx = \frac{43 (25 \cos(dx + c)^2 + 30 \sin(dx + c) - 34) \log(4 \cos(dx + c) - 3 \sin(dx + c) + 5) - 43 (25 \cos(dx + c) + 30 \sin(dx + c) - 34)}{4096 (25 d \cos(dx + c) + \dots)}$$

```
input integrate(1/(-3+5*sin(d*x+c))^3,x, algorithm="fricas")
```

output

```
-1/4096*(43*(25*cos(d*x + c)^2 + 30*sin(d*x + c) - 34)*log(4*cos(d*x + c)
- 3*sin(d*x + c) + 5) - 43*(25*cos(d*x + c)^2 + 30*sin(d*x + c) - 34)*log(
-4*cos(d*x + c) - 3*sin(d*x + c) + 5) - 1800*cos(d*x + c)*sin(d*x + c) + 4
40*cos(d*x + c))/(25*d*cos(d*x + c)^2 + 30*d*sin(d*x + c) - 34*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1224 vs. $2(102) = 204$.

Time = 1.48 (sec) , antiderivative size = 1224, normalized size of antiderivative = 10.64

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(-3+5*sin(d*x+c))**3,x)
```

output

```
Piecewise((x/(-3 + 5*sin(2*atan(1/3)))**3, Eq(c, -d*x + 2*atan(1/3))), (x/
(-3 + 5*sin(2*atan(3)))**3, Eq(c, -d*x + 2*atan(3))), (x/(5*sin(c) - 3)**3
, Eq(d, 0)), (-3483*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**4/(165888*
d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2
+ d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 23220*log(tan(c/2 +
d*x/2) - 3)*tan(c/2 + d*x/2)**3/(165888*d*tan(c/2 + d*x/2)**4 - 1105920*d
*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 - 1105920*d*tan(c/2 +
d*x/2) + 165888*d) - 45666*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**2/
(165888*d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*
tan(c/2 + d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 23220*log(t
an(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)/(165888*d*tan(c/2 + d*x/2)**4 - 1105
920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 - 1105920*d*tan(
c/2 + d*x/2) + 165888*d) - 3483*log(tan(c/2 + d*x/2) - 3)/(165888*d*tan(c/
2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)
**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 3483*log(3*tan(c/2 + d*x/2)
- 1)*tan(c/2 + d*x/2)**4/(165888*d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/
2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2)
+ 165888*d) - 23220*log(3*tan(c/2 + d*x/2) - 1)*tan(c/2 + d*x/2)**3/(1658
88*d*tan(c/2 + d*x/2)**4 - 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(
/2 + d*x/2)**2 - 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 45666*log(3*t...
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.70

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{40 \left(\frac{735 \sin(dx+c)}{\cos(dx+c)+1} - \frac{649 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{75 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - 99 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} - \frac{118 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{9 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} - 9} + 387 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} - 1 \right) - 387 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} - 3 \right)$$

$$18432 d$$

input `integrate(1/(-3+5*sin(d*x+c))^3,x, algorithm="maxima")`output `1/18432*(40*(735*sin(d*x + c)/(cos(d*x + c) + 1) - 649*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 75*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 99)/(60*sin(d*x + c)/(cos(d*x + c) + 1) - 118*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 9*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 - 9) + 387*log(3*sin(d*x + c)/(cos(d*x + c) + 1) - 1) - 387*log(sin(d*x + c)/(cos(d*x + c) + 1) - 3))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.93

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{40 \left(75 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 649 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 99 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^2} + 387 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1 \right| \right) - 387 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3 \right| \right)$$

$$18432 d$$

input `integrate(1/(-3+5*sin(d*x+c))^3,x, algorithm="giac")`output `1/18432*(40*(75*tan(1/2*d*x + 1/2*c)^3 + 649*tan(1/2*d*x + 1/2*c)^2 - 735*tan(1/2*d*x + 1/2*c) + 99)/(3*tan(1/2*d*x + 1/2*c)^2 - 10*tan(1/2*d*x + 1/2*c) + 3)^2 + 387*log(abs(3*tan(1/2*d*x + 1/2*c) - 1)) - 387*log(abs(tan(1/2*d*x + 1/2*c) - 3)))/d`

Mupad [B] (verification not implemented)

Time = 26.71 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{43 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{5}{4}\right)}{1024 d} + \frac{\frac{125 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{6912} + \frac{3245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{20736} - \frac{1225 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{6912} + \frac{55}{2304}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 - \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{3} + \frac{118 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{9} - \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(1/(5*sin(c + d*x) - 3)^3,x)`output `(43*atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4))/(1024*d) + ((3245*tan(c/2 + (d*x)/2)^2)/20736 - (1225*tan(c/2 + (d*x)/2))/6912 + (125*tan(c/2 + (d*x)/2)^3)/6912 + 55/2304)/(d*((118*tan(c/2 + (d*x)/2)^2)/9 - (20*tan(c/2 + (d*x)/2))/3 - (20*tan(c/2 + (d*x)/2)^3)/3 + tan(c/2 + (d*x)/2)^4 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.63

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^3} dx$$

$$= \frac{-900 \cos(dx + c) \sin(dx + c) + 220 \cos(dx + c) - 1075 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) \sin(dx + c)^2 + 1290 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right) \cos(dx + c)}{(-3 + 5 \sin(c + dx))^3}$$

input `int(1/(-3+5*sin(d*x+c))^3,x)`

output

```
( - 900*cos(c + d*x)*sin(c + d*x) + 220*cos(c + d*x) - 1075*log(tan((c + d
*x)/2) - 3)*sin(c + d*x)**2 + 1290*log(tan((c + d*x)/2) - 3)*sin(c + d*x)
- 387*log(tan((c + d*x)/2) - 3) + 1075*log(3*tan((c + d*x)/2) - 1)*sin(c +
d*x)**2 - 1290*log(3*tan((c + d*x)/2) - 1)*sin(c + d*x) + 387*log(3*tan((
c + d*x)/2) - 1) + 750*sin(c + d*x)**2 - 900*sin(c + d*x) + 270)/(2048*d*(
25*sin(c + d*x)**2 - 30*sin(c + d*x) + 9))
```

3.71 $\int \frac{1}{(-3+5 \sin(c+dx))^4} dx$

Optimal result	511
Mathematica [A] (verified)	512
Rubi [A] (verified)	512
Maple [A] (verified)	516
Fricas [A] (verification not implemented)	516
Sympy [B] (verification not implemented)	517
Maxima [B] (verification not implemented)	518
Giac [A] (verification not implemented)	518
Mupad [B] (verification not implemented)	519
Reduce [B] (verification not implemented)	519

Optimal result

Integrand size = 12, antiderivative size = 140

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx = \frac{279 \log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx)))}{32768d} - \frac{279 \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx)))}{32768d} + \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} - \frac{25 \cos(c + dx)}{512d(3 - 5 \sin(c + dx))^2} + \frac{995 \cos(c + dx)}{24576d(3 - 5 \sin(c + dx))}$$

output

```
279/32768*ln(cos(1/2*d*x+1/2*c)-3*sin(1/2*d*x+1/2*c))/d-279/32768*ln(3*cos
(1/2*d*x+1/2*c)-sin(1/2*d*x+1/2*c))/d+5/48*cos(d*x+c)/d/(3-5*sin(d*x+c))^3
-25/512*cos(d*x+c)/d/(3-5*sin(d*x+c))^2+995/24576*cos(d*x+c)/d/(3-5*sin(d*
x+c))
```


Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.72

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx$$

$$= \frac{2511 \log(\cos(\frac{1}{2}(c + dx)) - 3 \sin(\frac{1}{2}(c + dx))) - 2511 \log(3 \cos(\frac{1}{2}(c + dx)) - \sin(\frac{1}{2}(c + dx))) - \frac{1}{\cos(\frac{1}{2}(c + dx))}}{\dots}$$

input `Integrate[(-3 + 5*Sin[c + d*x])^(-4),x]`

output `(2511*Log[Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]] - 2511*Log[3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]] - 720/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2])^2 + 20*(240/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2])^3 + 597/(Cos[(c + d*x)/2] - 3*Sin[(c + d*x)/2]) + 80/(3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2])^3 + 199/(3*Cos[(c + d*x)/2] - Sin[(c + d*x)/2]))*Sin[(c + d*x)/2] + 2320/(-3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2)/(294912*d)`

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.94, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$, Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(5 \sin(c + dx) - 3)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(5 \sin(c + dx) - 3)^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{10 \sin(c + dx) + 9}{(3 - 5 \sin(c + dx))^3} dx + \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} - \frac{1}{48} \int \frac{10 \sin(c + dx) + 9}{(3 - 5 \sin(c + dx))^3} dx \\
& \downarrow 3042 \\
& \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} - \frac{1}{48} \int \frac{10 \sin(c + dx) + 9}{(3 - 5 \sin(c + dx))^3} dx \\
& \downarrow 3233 \\
& \frac{1}{48} \left(-\frac{1}{32} \int -\frac{75 \sin(c + dx) + 154}{(3 - 5 \sin(c + dx))^2} dx - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \right) + \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{75 \sin(c + dx) + 154}{(3 - 5 \sin(c + dx))^2} dx - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \right) + \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{75 \sin(c + dx) + 154}{(3 - 5 \sin(c + dx))^2} dx - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \right) + \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{1}{16} \int -\frac{837}{3 - 5 \sin(c + dx)} dx + \frac{995 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} \right) - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \right) + \\
& \quad \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{837}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \right) - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \right) + \\
& \quad \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c + dx)}{16d(3 - 5 \sin(c + dx))} - \frac{837}{16} \int \frac{1}{3 - 5 \sin(c + dx)} dx \right) - \frac{75 \cos(c + dx)}{32d(3 - 5 \sin(c + dx))^2} \right) + \\
& \quad \frac{5 \cos(c + dx)}{48d(3 - 5 \sin(c + dx))^3} \\
& \downarrow 3139
\end{aligned}$$

$$\frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c+dx)}{16d(3-5\sin(c+dx))} - \frac{837 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c+dx))}{8d} \right) - \frac{75 \cos(c+dx)}{32d(3-5\sin(c+dx))} \right) - \frac{5 \cos(c+dx)}{48d(3-5\sin(c+dx))^3}$$

↓ 1081

$$\frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c+dx)}{16d(3-5\sin(c+dx))} - \frac{2511 \int \left(\frac{1}{8(1-3 \tan(\frac{1}{2}(c+dx)))} - \frac{1}{24(3-\tan(\frac{1}{2}(c+dx)))} \right) d \tan(\frac{1}{2}(c+dx))}{8d} \right) - \frac{75 \cos(c+dx)}{32d(3-5\sin(c+dx))} \right) - \frac{5 \cos(c+dx)}{48d(3-5\sin(c+dx))^3}$$

↓ 2009

$$\frac{1}{48} \left(\frac{1}{32} \left(\frac{995 \cos(c+dx)}{16d(3-5\sin(c+dx))} - \frac{2511 \left(\frac{1}{24} \log(3 - \tan(\frac{1}{2}(c+dx))) - \frac{1}{24} \log(1 - 3 \tan(\frac{1}{2}(c+dx))) \right)}{8d} \right) - \frac{75 \cos(c+dx)}{32d(3-5\sin(c+dx))} \right) - \frac{5 \cos(c+dx)}{48d(3-5\sin(c+dx))^3} +$$

input `Int[(-3 + 5*Sin[c + d*x])^(-4), x]`

output `((((-2511*(-1/24*Log[1 - 3*Tan[(c + d*x)/2]] + Log[3 - Tan[(c + d*x)/2]]/24)))/(8*d) + (995*Cos[c + d*x])/(16*d*(3 - 5*Sin[c + d*x])))/32 - (75*Cos[c + d*x])/(32*d*(3 - 5*Sin[c + d*x]^2))/48 + (5*Cos[c + d*x])/(48*d*(3 - 5*Sin[c + d*x]^3))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`

rule 3233 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.94

method	result
derivativedivides	$-\frac{125}{768 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)^3} - \frac{75}{1024 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)^2} - \frac{345}{8192 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)} - \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{32768} - \frac{125}{20736 \left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3}$
default	$-\frac{125}{768 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)^3} - \frac{75}{1024 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)^2} - \frac{345}{8192 \left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)} - \frac{279 \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 3\right)}{32768} - \frac{125}{20736 \left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right)^3}$
risch	$\frac{-111042 e^{3i(dx+c)} - 62775 i e^{4i(dx+c)} + 119310 i e^{2i(dx+c)} + 20925 e^{5i(dx+c)} + 68625 e^{i(dx+c)} - 24875 i}{12288 (5 e^{2i(dx+c)} - 5 - 6 i e^{i(dx+c)})^3} d + \frac{279 \ln\left(-\frac{4}{5} - \frac{3i}{5}\right)}{32768}$
norman	$\frac{7915}{12288d} - \frac{15725 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^5}{12288d} - \frac{3047275 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^3}{165888d} - \frac{63425 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)}{12288d} + \frac{296245 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2}{18432d} + \frac{270245 \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^4}{36864d} - \dots$
parallelrisc	$\frac{(10169550 \cos(2dx+2c) + 20678085 \sin(dx+c) - 2824875 \sin(3dx+3c) - 12610242) \ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{3}\right) + (-10169550 \cos(dx+c) - 20678085 \sin(dx+c) + 2824875 \sin(3dx+3c) + 12610242)}{(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 10 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3)^3}$

input `int(1/(-3+5*sin(d*x+c))^4,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(-\frac{125}{768} \left(\tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 3\right)^{-3} - \frac{75}{1024} \left(\tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 3\right)^{-2} - \frac{345}{8192} \left(\tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 3\right)^{-1} - \frac{279}{32768} \ln\left(\tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 3\right) - \frac{125}{20736} \left(3 \tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 1\right)^{-3} - \frac{275}{27648} \left(3 \tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 1\right)^{-2} - \frac{3505}{221184} \left(3 \tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 1\right)^{-1} + \frac{279}{32768} \ln\left(3 \tan\left(\frac{1}{2}d*x + \frac{1}{2}c\right) - 1\right) \right)$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.29

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx$$

$$= \frac{199000 \cos(dx + c)^3 - 837 (225 \cos(dx + c)^2 - 5 (25 \cos(dx + c)^2 - 52) \sin(dx + c) - 252) \log(4 \cos(dx + c) - 2)}{(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 10 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3)^3}$$

input `integrate(1/(-3+5*sin(d*x+c))^4,x, algorithm="fricas")`

output

```
1/196608*(199000*cos(d*x + c)^3 - 837*(225*cos(d*x + c)^2 - 5*(25*cos(d*x
+ c)^2 - 52)*sin(d*x + c) - 252)*log(4*cos(d*x + c) - 3*sin(d*x + c) + 5)
+ 837*(225*cos(d*x + c)^2 - 5*(25*cos(d*x + c)^2 - 52)*sin(d*x + c) - 252)
*log(-4*cos(d*x + c) - 3*sin(d*x + c) + 5) + 190800*cos(d*x + c)*sin(d*x +
c) - 262320*cos(d*x + c))/(225*d*cos(d*x + c)^2 - 5*(25*d*cos(d*x + c)^2
- 52*d)*sin(d*x + c) - 252*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2353 vs. $2(126) = 252$.

Time = 3.24 (sec) , antiderivative size = 2353, normalized size of antiderivative = 16.81

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(-3+5*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(-3 + 5*sin(2*atan(1/3)))**4, Eq(c, -d*x + 2*atan(1/3))), (x/
(-3 + 5*sin(2*atan(3)))**4, Eq(c, -d*x + 2*atan(3))), (x/(5*sin(c) - 3)**4
, Eq(d, 0)), (-610173*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**6/(71663
616*d*tan(c/2 + d*x/2)**6 - 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d
*tan(c/2 + d*x/2)**4 - 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan
(c/2 + d*x/2)**2 - 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 6101730*lo
g(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**5/(71663616*d*tan(c/2 + d*x/2)**
6 - 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 - 4
087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 - 71663
6160*d*tan(c/2 + d*x/2) + 71663616*d) - 22169619*log(tan(c/2 + d*x/2) - 3)
*tan(c/2 + d*x/2)**4/(71663616*d*tan(c/2 + d*x/2)**6 - 716636160*d*tan(c/2
+ d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 - 4087480320*d*tan(c/2 + d
*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 - 716636160*d*tan(c/2 + d*x/2)
+ 71663616*d) + 34802460*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**3/(7
1663616*d*tan(c/2 + d*x/2)**6 - 716636160*d*tan(c/2 + d*x/2)**5 + 26037780
48*d*tan(c/2 + d*x/2)**4 - 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d
*tan(c/2 + d*x/2)**2 - 716636160*d*tan(c/2 + d*x/2) + 71663616*d) - 221696
19*log(tan(c/2 + d*x/2) - 3)*tan(c/2 + d*x/2)**2/(71663616*d*tan(c/2 + d*x
/2)**6 - 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**
4 - 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(126) = 252$.

Time = 0.04 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.96

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx$$

$$= \frac{40 \left(\frac{342495 \sin(dx+c)}{\cos(dx+c)+1} - \frac{1066482 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1218910 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} - \frac{486441 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{84915 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} - 42741 \right) + 22599 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} \right)}{2654208 d}$$

input `integrate(1/(-3+5*sin(d*x+c))^4,x, algorithm="maxima")`

output

```
1/2654208*(40*(342495*sin(d*x + c)/(cos(d*x + c) + 1) - 1066482*sin(d*x +
c)^2/(cos(d*x + c) + 1)^2 + 1218910*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 -
486441*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 84915*sin(d*x + c)^5/(cos(d*x
+ c) + 1)^5 - 42741)/(270*sin(d*x + c)/(cos(d*x + c) + 1) - 981*sin(d*x +
c)^2/(cos(d*x + c) + 1)^2 + 1540*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 - 98
1*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 270*sin(d*x + c)^5/(cos(d*x + c)
+ 1)^5 - 27*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 - 27) + 22599*log(3*sin(d*x
+ c)/(cos(d*x + c) + 1) - 1) - 22599*log(sin(d*x + c)/(cos(d*x + c) + 1)
- 3))/d
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.95

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx =$$

$$\frac{40 \left(84915 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 486441 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 1218910 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 1066482 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 342495 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 42741 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^3}$$

2654208 d

input `integrate(1/(-3+5*sin(d*x+c))^4,x, algorithm="giac")`

output

```
-1/2654208*(40*(84915*tan(1/2*d*x + 1/2*c)^5 - 486441*tan(1/2*d*x + 1/2*c)
^4 + 1218910*tan(1/2*d*x + 1/2*c)^3 - 1066482*tan(1/2*d*x + 1/2*c)^2 + 342
495*tan(1/2*d*x + 1/2*c) - 42741)/(3*tan(1/2*d*x + 1/2*c)^2 - 10*tan(1/2*d
*x + 1/2*c) + 3)^3 - 22599*log(abs(3*tan(1/2*d*x + 1/2*c) - 1)) + 22599*lo
g(abs(tan(1/2*d*x + 1/2*c) - 3)))/d
```

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.20

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx = \frac{279 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} - \frac{5}{4}\right)}{16384 d} - \frac{\frac{15725 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{331776} - \frac{270245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{995328} + \frac{3047275 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{4478976} - \frac{296245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{497664} + \frac{63425 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{331776} - \frac{7915}{331776}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{3} - \frac{1540 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{27} + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{3} - 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1 \right)}$$

input

```
int(1/(5*sin(c + d*x) - 3)^4,x)
```

output

```
(279*atanh((3*tan(c/2 + (d*x)/2))/4 - 5/4))/(16384*d) - ((63425*tan(c/2 +
(d*x)/2))/331776 - (296245*tan(c/2 + (d*x)/2)^2)/497664 + (3047275*tan(c/2
+ (d*x)/2)^3)/4478976 - (270245*tan(c/2 + (d*x)/2)^4)/995328 + (15725*tan
(c/2 + (d*x)/2)^5)/331776 - 7915/331776)/(d*((109*tan(c/2 + (d*x)/2)^2)/3
- 10*tan(c/2 + (d*x)/2) - (1540*tan(c/2 + (d*x)/2)^3)/27 + (109*tan(c/2 +
(d*x)/2)^4)/3 - 10*tan(c/2 + (d*x)/2)^5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.92

$$\int \frac{1}{(-3 + 5 \sin(c + dx))^4} dx = \frac{-298500 \cos(dx + c) \sin(dx + c)^2 + 286200 \cos(dx + c) \sin(dx + c) - 94980 \cos(dx + c) - 313875 \log\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) - 5}{4}\right)}{d}$$

input `int(1/(-3+5*sin(d*x+c))^4,x)`

output `(- 298500*cos(c + d*x)*sin(c + d*x)**2 + 286200*cos(c + d*x)*sin(c + d*x)
- 94980*cos(c + d*x) - 313875*log(tan((c + d*x)/2) - 3)*sin(c + d*x)**3 +
564975*log(tan((c + d*x)/2) - 3)*sin(c + d*x)**2 - 338985*log(tan((c + d*
x)/2) - 3)*sin(c + d*x) + 67797*log(tan((c + d*x)/2) - 3) + 313875*log(3*t
an((c + d*x)/2) - 1)*sin(c + d*x)**3 - 564975*log(3*tan((c + d*x)/2) - 1)*
sin(c + d*x)**2 + 338985*log(3*tan((c + d*x)/2) - 1)*sin(c + d*x) - 67797*
log(3*tan((c + d*x)/2) - 1) + 265000*sin(c + d*x)**3 - 477000*sin(c + d*x)
2 + 286200*sin(c + d*x) - 57240)/(294912*d*(125*sin(c + d*x)3 - 225*si
n(c + d*x)**2 + 135*sin(c + d*x) - 27))`

3.72 $\int \frac{1}{-3-5 \sin(c+dx)} dx$

Optimal result	521
Mathematica [A] (verified)	521
Rubi [A] (verified)	522
Maple [A] (verified)	523
Fricas [A] (verification not implemented)	524
Sympy [A] (verification not implemented)	524
Maxima [A] (verification not implemented)	524
Giac [A] (verification not implemented)	525
Mupad [B] (verification not implemented)	525
Reduce [B] (verification not implemented)	526

Optimal result

Integrand size = 12, antiderivative size = 63

$$\int \frac{1}{-3-5 \sin(c+dx)} dx = \frac{\log(3 \cos(\frac{1}{2}(c+dx)) + \sin(\frac{1}{2}(c+dx)))}{4d} - \frac{\log(\cos(\frac{1}{2}(c+dx)) + 3 \sin(\frac{1}{2}(c+dx)))}{4d}$$

output `1/4*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-1/4*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00

$$\int \frac{1}{-3-5 \sin(c+dx)} dx = \frac{\log(3 \cos(\frac{1}{2}(c+dx)) + \sin(\frac{1}{2}(c+dx)))}{4d} - \frac{\log(\cos(\frac{1}{2}(c+dx)) + 3 \sin(\frac{1}{2}(c+dx)))}{4d}$$

input `Integrate[(-3 - 5*Sin[c + d*x])^(-1),x]`

output

$$\text{Log}[3*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]]/(4*d) - \text{Log}[\text{Cos}[(c + d*x)/2] + 3*\text{Sin}[(c + d*x)/2]]/(4*d)$$
Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{-5 \sin(c + dx) - 3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{-5 \sin(c + dx) - 3} dx \\ & \quad \downarrow \text{3139} \\ & \frac{2 \int \frac{1}{-3 \tan^2(\frac{1}{2}(c+dx)) - 10 \tan(\frac{1}{2}(c+dx)) - 3} d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{1081} \\ & \frac{6 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx))+1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx))+3)} \right) d \tan(\frac{1}{2}(c + dx))}{d} \\ & \quad \downarrow \text{2009} \\ & \frac{6 \left(\frac{1}{24} \log(3 \tan(\frac{1}{2}(c + dx)) + 1) - \frac{1}{24} \log(\tan(\frac{1}{2}(c + dx)) + 3) \right)}{d} \end{aligned}$$

input

$$\text{Int}[(-3 - 5*\text{Sin}[c + d*x])^{-1}, x]$$

output

$$(-6*(-1/24*\text{Log}[3 + \text{Tan}[(c + d*x)/2]] + \text{Log}[1 + 3*\text{Tan}[(c + d*x)/2]]/24))/d$$

Definitions of rubi rules used

rule 1081 $\text{Int}[\{(a_)+(b_)*(x_)+(c_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c \text{ Int}[\text{ExpandIntegrand}[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x]] \text{ /; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NiceSqrtQ}[b^2 - 4*a*c]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3139 $\text{Int}[\{(a_)+(b_)*\sin[(c_)+(d_)*(x_)]\}^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] \text{ /; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.57

method	result	size
derivativedivides	$\frac{-\frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{4}}{d}$	36
default	$\frac{-\frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4} + \frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{4}}{d}$	36
parallelrisch	$\frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 9\right) - \ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d}$	37
norman	$\frac{\ln\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right)}{4d} - \frac{\ln\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d}$	38
risch	$\frac{\ln\left(\frac{4}{5} + \frac{3i}{5} + e^{i(dx+c)}\right)}{4d} - \frac{\ln\left(e^{i(dx+c)} - \frac{4}{5} + \frac{3i}{5}\right)}{4d}$	40

input $\text{int}(1/(-3-5*\sin(d*x+c)), x, \text{method}=_RETURNVERBOSE)$

output $1/d*(-1/4*\ln(3*\tan(1/2*d*x+1/2*c)+1)+1/4*\ln(\tan(1/2*d*x+1/2*c)+3))$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int \frac{1}{-3 - 5 \sin(c + dx)} dx = \frac{\log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5)}{8d}$$

input `integrate(1/(-3-5*sin(d*x+c)),x, algorithm="fricas")`output `1/8*(log(4*cos(d*x + c) + 3*sin(d*x + c) + 5) - log(-4*cos(d*x + c) + 3*sin(d*x + c) + 5))/d`**Sympy [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

$$\int \frac{1}{-3 - 5 \sin(c + dx)} dx = \begin{cases} \frac{\log(\tan(\frac{c}{2} + \frac{dx}{2}) + 3)}{4d} - \frac{\log(3 \tan(\frac{c}{2} + \frac{dx}{2}) + 1)}{4d} & \text{for } d \neq 0 \\ \frac{x}{-5 \sin(c) - 3} & \text{otherwise} \end{cases}$$

input `integrate(1/(-3-5*sin(d*x+c)),x)`output `Piecewise((log(tan(c/2 + d*x/2) + 3)/(4*d) - log(3*tan(c/2 + d*x/2) + 1)/(4*d), Ne(d, 0)), (x/(-5*sin(c) - 3), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.78

$$\int \frac{1}{-3 - 5 \sin(c + dx)} dx = -\frac{\log\left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1\right) - \log\left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 3\right)}{4d}$$

input `integrate(1/(-3-5*sin(d*x+c)),x, algorithm="maxima")`

output

```
-1/4*(log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - log(sin(d*x + c)/(cos(d*x + c) + 1) + 3))/d
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.57

$$\int \frac{1}{-3 - 5 \sin(c + dx)} dx = -\frac{\log\left(\left|3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right|\right) - \log\left(\left|\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3\right|\right)}{4d}$$

input

```
integrate(1/(-3-5*sin(d*x+c)),x, algorithm="giac")
```

output

```
-1/4*(log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) - log(abs(tan(1/2*d*x + 1/2*c) + 3)))/d
```

Mupad [B] (verification not implemented)

Time = 25.47 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.30

$$\int \frac{1}{-3 - 5 \sin(c + dx)} dx = \frac{\operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{2d}$$

input

```
int(-1/(5*sin(c + d*x) + 3),x)
```

output

```
atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4)/(2*d)
```

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.54

$$\int \frac{1}{-3 - 5 \sin(c + dx)} dx = \frac{\log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) - \log\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{4d}$$

input `int(1/(-3-5*sin(d*x+c)),x)`

output `(log(tan((c + d*x)/2) + 3) - log(3*tan((c + d*x)/2) + 1))/(4*d)`

3.73 $\int \frac{1}{(-3-5 \sin(c+dx))^2} dx$

Optimal result	527
Mathematica [A] (verified)	527
Rubi [A] (verified)	528
Maple [A] (verified)	530
Fricas [A] (verification not implemented)	530
Sympy [B] (verification not implemented)	531
Maxima [A] (verification not implemented)	532
Giac [A] (verification not implemented)	532
Mupad [B] (verification not implemented)	533
Reduce [B] (verification not implemented)	533

Optimal result

Integrand size = 12, antiderivative size = 88

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx = \frac{3 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{3 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))}{64d} - \frac{5 \cos(c + dx)}{16d(3 + 5 \sin(c + dx))}$$

output

```
3/64*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-3/64*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d-5/16*cos(d*x+c)/d/(3+5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.43

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx = \frac{9(\log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) - \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))) + 20 \sin(\frac{1}{2}(c + dx))}{192d}$$

input `Integrate[(-3 - 5*Sin[c + d*x])^(-2),x]`

output `(9*(Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]]) + 20*Sin[(c + d*x)/2]*((3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^(-1) + 3/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])))/(192*d)`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.80, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 3143, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(-5 \sin(c + dx) - 3)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-5 \sin(c + dx) - 3)^2} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{1}{16} \int -\frac{3}{5 \sin(c + dx) + 3} dx - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow \text{27} \\
 & -\frac{3}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{3}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow \text{3139} \\
 & -\frac{3 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) + 10 \tan(\frac{1}{2}(c+dx)) + 3} d \tan(\frac{1}{2}(c + dx))}{8d} - \frac{5 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \\
 & \quad \downarrow \text{1081}
 \end{aligned}$$

$$\frac{9 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx))+1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx))+3)} \right) d \tan \left(\frac{1}{2}(c+dx) \right)}{8d} - \frac{5 \cos(c+dx)}{16d(5 \sin(c+dx)+3)}$$

↓ 2009

$$\frac{9 \left(\frac{1}{24} \log \left(3 \tan \left(\frac{1}{2}(c+dx) \right) + 1 \right) - \frac{1}{24} \log \left(\tan \left(\frac{1}{2}(c+dx) \right) + 3 \right) \right)}{8d} - \frac{5 \cos(c+dx)}{16d(5 \sin(c+dx)+3)}$$

input `Int[(-3 - 5*Sin[c + d*x])^(-2),x]`

output `(-9*(-1/24*Log[3 + Tan[(c + d*x)/2]] + Log[1 + 3*Tan[(c + d*x)/2]]/24))/(8*d) - (5*Cos[c + d*x])/(16*d*(3 + 5*Sin[c + d*x]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`

rule 3143

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{-\frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{64} - \frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{64}}{d}$
default	$\frac{-\frac{5}{48(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{64} - \frac{5}{16(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{64}}{d}$
risch	$-\frac{3 e^{i(dx+c)} + 5i}{8d(5 e^{2i(dx+c)} - 5 + 6i e^{i(dx+c)})} + \frac{3 \ln(\frac{4}{5} + \frac{3i}{5} + e^{i(dx+c)})}{64d} - \frac{3 \ln(e^{i(dx+c)} - \frac{4}{5} + \frac{3i}{5})}{64d}$
norman	$\frac{-\frac{5}{8d} - \frac{25 \tan(\frac{dx}{2} + \frac{c}{2})}{24d}}{3 \tan(\frac{dx}{2} + \frac{c}{2})^2 + 10 \tan(\frac{dx}{2} + \frac{c}{2}) + 3} + \frac{3 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{64d} - \frac{3 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{64d}$
parallelrisch	$\frac{45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3) \sin(dx+c) - 45 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + \frac{1}{3}) \sin(dx+c) - 100 \sin(dx+c) - 60 \cos(dx+c) + 27 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{192d(3 + 5 \sin(dx+c))}$

input

```
int(1/(-3-5*sin(d*x+c))^2,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-5/48/(3*tan(1/2*d*x+1/2*c)+1)-3/64*ln(3*tan(1/2*d*x+1/2*c)+1)-5/16/(
tan(1/2*d*x+1/2*c)+3)+3/64*ln(tan(1/2*d*x+1/2*c)+3))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx$$

$$= \frac{3(5 \sin(dx + c) + 3) \log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - 3(5 \sin(dx + c) + 3) \log(-4 \cos(dx + c) + 3 \sin(dx + c) + 5)}{128(5d \sin(dx + c) + 3d)}$$

input `integrate(1/(-3-5*sin(d*x+c))^2,x, algorithm="fricas")`

output `1/128*(3*(5*sin(d*x + c) + 3)*log(4*cos(d*x + c) + 3*sin(d*x + c) + 5) - 3*(5*sin(d*x + c) + 3)*log(-4*cos(d*x + c) + 3*sin(d*x + c) + 5) - 40*cos(d*x + c))/(5*d*sin(d*x + c) + 3*d)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 468 vs. $2(78) = 156$.

Time = 0.73 (sec) , antiderivative size = 468, normalized size of antiderivative = 5.32

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(1/(-3-5*sin(d*x+c))**2,x)`

output `Piecewise((x/(-3 + 5*sin(2*atan(1/3)))**2, Eq(c, -d*x - 2*atan(1/3))), (x/(-3 + 5*sin(2*atan(3)))**2, Eq(c, -d*x - 2*atan(3))), (x/(-5*sin(c) - 3)**2, Eq(d, 0)), (27*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) + 90*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) + 27*log(tan(c/2 + d*x/2) + 3)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 27*log(3*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**2/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 90*log(3*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 27*log(3*tan(c/2 + d*x/2) + 1)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 200*tan(c/2 + d*x/2)/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d) - 120/(576*d*tan(c/2 + d*x/2)**2 + 1920*d*tan(c/2 + d*x/2) + 576*d), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.31

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx$$

$$= -\frac{\frac{40 \left(\frac{5 \sin(dx+c)}{\cos(dx+c)+1} + 3 \right)}{\frac{10 \sin(dx+c)}{\cos(dx+c)+1} + \frac{3 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + 3} + 9 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) - 9 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 3 \right)}{192 d}$$

input `integrate(1/(-3-5*sin(d*x+c))^2,x, algorithm="maxima")`output `-1/192*(40*(5*sin(d*x + c)/(cos(d*x + c) + 1) + 3)/(10*sin(d*x + c)/(cos(d*x + c) + 1) + 3*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 3) + 9*log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - 9*log(sin(d*x + c)/(cos(d*x + c) + 1) + 3))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.92

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx =$$

$$-\frac{\frac{40 \left(5 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)}{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3} + 9 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) - 9 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right| \right)}{192 d}$$

input `integrate(1/(-3-5*sin(d*x+c))^2,x, algorithm="giac")`output `-1/192*(40*(5*tan(1/2*d*x + 1/2*c) + 3)/(3*tan(1/2*d*x + 1/2*c)^2 + 10*tan(1/2*d*x + 1/2*c) + 3) + 9*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) - 9*log(abs(tan(1/2*d*x + 1/2*c) + 3)))/d`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.73

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx = \frac{3 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{32 d} - \frac{\frac{25 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{72} + \frac{5}{24}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2 + \frac{10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(1/(5*sin(c + d*x) + 3)^2,x)`output `(3*atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4))/(32*d) - ((25*tan(c/2 + (d*x)/2)/72 + 5/24)/(d*((10*tan(c/2 + (d*x)/2))/3 + tan(c/2 + (d*x)/2)^2 + 1))`**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.11

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^2} dx = \frac{-20 \cos(dx + c) + 15 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) \sin(dx + c) + 9 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) - 15 \log\left(3 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right)}{64d(5 \sin(dx + c) + 3)}$$

input `int(1/(-3-5*sin(d*x+c))^2,x)`output `(- 20*cos(c + d*x) + 15*log(tan((c + d*x)/2) + 3)*sin(c + d*x) + 9*log(tan((c + d*x)/2) + 3) - 15*log(3*tan((c + d*x)/2) + 1)*sin(c + d*x) - 9*log(3*tan((c + d*x)/2) + 1))/(64*d*(5*sin(c + d*x) + 3))`

3.74 $\int \frac{1}{(-3-5 \sin(c+dx))^3} dx$

Optimal result	534
Mathematica [A] (verified)	534
Rubi [A] (verified)	535
Maple [A] (verified)	538
Fricas [A] (verification not implemented)	538
Sympy [B] (verification not implemented)	539
Maxima [A] (verification not implemented)	540
Giac [A] (verification not implemented)	540
Mupad [B] (verification not implemented)	541
Reduce [B] (verification not implemented)	541

Optimal result

Integrand size = 12, antiderivative size = 113

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx = \frac{43 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))}{2048d} - \frac{43 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx)))}{2048d} + \frac{5 \cos(c + dx)}{32d(3 + 5 \sin(c + dx))^2} - \frac{45 \cos(c + dx)}{512d(3 + 5 \sin(c + dx))}$$

```
output 43/2048*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-43/2048*ln(cos(1/2*d
*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d+5/32*cos(d*x+c)/d/(3+5*sin(d*x+c))^2-45/
512*cos(d*x+c)/d/(3+5*sin(d*x+c))
```

Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.58

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx = \frac{43 \log(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))) - 43 \log(\cos(\frac{1}{2}(c + dx)) + 3 \sin(\frac{1}{2}(c + dx))) - \frac{45 \cos(c + dx)}{512d(3 + 5 \sin(c + dx))}}{(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^3}$$

input `Integrate[(-3 - 5*Sin[c + d*x])^(-3),x]`

output $(43*\text{Log}[3*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2]] - 43*\text{Log}[\text{Cos}[(c + d*x)/2] + 3*\text{Sin}[(c + d*x)/2]] - 40/(3*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2])^2 + 40/(\text{Cos}[(c + d*x)/2] + 3*\text{Sin}[(c + d*x)/2])^2 + 60*\text{Sin}[(c + d*x)/2]*((3*\text{Cos}[(c + d*x)/2] + \text{Sin}[(c + d*x)/2])^{-1}) + 3/(\text{Cos}[(c + d*x)/2] + 3*\text{Sin}[(c + d*x)/2]))/(2048*d)$

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.88, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 3143, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-5 \sin(c + dx) - 3)^3} dx$$

↓ 3042

$$\int \frac{1}{(-5 \sin(c + dx) - 3)^3} dx$$

↓ 3143

$$\frac{1}{32} \int \frac{6 - 5 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx + \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2}$$

↓ 3042

$$\frac{1}{32} \int \frac{6 - 5 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx + \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2}$$

↓ 3233

$$\frac{1}{32} \left(\frac{1}{16} \int -\frac{43}{5 \sin(c + dx) + 3} dx - \frac{45 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2}$$

↓ 27

$$\frac{1}{32} \left(-\frac{43}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{45 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{5 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{32} \left(-\frac{43}{16} \int \frac{1}{5 \sin(c+dx)+3} dx - \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) + \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 3139 \\
& \frac{1}{32} \left(-\frac{43 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx))+10 \tan(\frac{1}{2}(c+dx))+3} d \tan(\frac{1}{2}(c+dx))}{8d} - \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) + \\
& \quad \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 1081 \\
& \frac{1}{32} \left(-\frac{129 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx))+1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx))+3)} \right) d \tan(\frac{1}{2}(c+dx))}{8d} - \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) + \\
& \quad \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2} \\
& \downarrow 2009 \\
& \frac{1}{32} \left(-\frac{129 \left(\frac{1}{24} \log(3 \tan(\frac{1}{2}(c+dx))+1) - \frac{1}{24} \log(\tan(\frac{1}{2}(c+dx))+3) \right)}{8d} - \frac{45 \cos(c+dx)}{16d(5 \sin(c+dx)+3)} \right) + \\
& \quad \frac{5 \cos(c+dx)}{32d(5 \sin(c+dx)+3)^2}
\end{aligned}$$

input `Int[(-3 - 5*Sin[c + d*x])^(-3),x]`

output `(5*Cos[c + d*x])/(32*d*(3 + 5*Sin[c + d*x])^2) + ((-129*(-1/24*Log[3 + Tan[(c + d*x)/2]] + Log[1 + 3*Tan[(c + d*x)/2]]/24))/(8*d) - (45*Cos[c + d*x])/(16*d*(3 + 5*Sin[c + d*x]))/32`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 1081 $\text{Int}[((a_) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Simp}[c \text{ Int}[\text{ExpandIntegrand}[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NiceSqrtQ}[b^2 - 4*a*c]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3139 $\text{Int}[((a_) + (b_*)\sin[(c_) + (d_*)(x_)])^{-1}, x_Symbol] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{ Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$
- rule 3143 $\text{Int}[((a_) + (b_*)\sin[(c_) + (d_*)(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\text{Sin}[c + d*x])^{(n + 1)}/(d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\text{Sin}[c + d*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$
- rule 3233 $\text{Int}[((a_) + (b_*)\sin[(e_) + (f_*)(x_)])^{(m_)}*((c_) + (d_*)\sin[(e_) + (f_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*\text{Cos}[e + f*x]*((a + b*\text{Sin}[e + f*x])^{(m + 1)}/(f*(m + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((m + 1)*(a^2 - b^2)) \text{ Int}[(a + b*\text{Sin}[e + f*x])^{(m + 1)}*\text{Simp}[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*\text{Sin}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegerQ}[2*m]$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{\frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} - \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{2048} - \frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} + \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}}{d}$
default	$\frac{\frac{25}{1152(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} - \frac{155}{4608(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{2048} - \frac{25}{128(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} + \frac{15}{512(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}}{d}$
risch	$-\frac{387ie^{2i(dx+c)} + 215e^{3i(dx+c)} - 325e^{i(dx+c)} - 225i}{256(5e^{2i(dx+c)} - 5 + 6ie^{i(dx+c)})^2}d + \frac{43 \ln(\frac{4}{5} + \frac{3i}{5} + e^{i(dx+c)})}{2048d} - \frac{43 \ln(e^{i(dx+c)} - \frac{4}{5} + \frac{3i}{5})}{2048d}$
norman	$-\frac{55}{256d} - \frac{3245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{2304d} - \frac{1225 \tan(\frac{dx}{2} + \frac{c}{2})}{768d} + \frac{125 \tan(\frac{dx}{2} + \frac{c}{2})^3}{768d} + \frac{43 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{2048d} - \frac{43 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{2048d}$
parallelrisc	$\frac{(-9675 \cos(2dx+2c) + 23220 \sin(dx+c) + 16641) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + \frac{1}{3}) + (9675 \cos(2dx+2c) - 23220 \sin(dx+c) - 16641)}{18432d(-43 + 25 \cos(2dx+2c))}$

input `int(1/(-3-5*sin(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} * (\frac{25}{1152} / (3 * \tan(1/2 * d * x + 1/2 * c) + 1)^2 - \frac{155}{4608} / (3 * \tan(1/2 * d * x + 1/2 * c) + 1) - \frac{43}{2048} * \ln(3 * \tan(1/2 * d * x + 1/2 * c) + 1) - \frac{25}{128} / (\tan(1/2 * d * x + 1/2 * c) + 3)^2 + \frac{15}{512} / (\tan(1/2 * d * x + 1/2 * c) + 3) + \frac{43}{2048} * \ln(\tan(1/2 * d * x + 1/2 * c) + 3))$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.18

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx$$

$$= \frac{43 (25 \cos(dx + c)^2 - 30 \sin(dx + c) - 34) \log(4 \cos(dx + c) + 3 \sin(dx + c) + 5) - 43 (25 \cos(dx + c) + 3 \sin(dx + c) + 5)}{4096 (25 d \cos(dx + c) + 3 \sin(dx + c) + 5)}$$

input `integrate(1/(-3-5*sin(d*x+c))^3,x, algorithm="fricas")`

output

```
1/4096*(43*(25*cos(d*x + c)^2 - 30*sin(d*x + c) - 34)*log(4*cos(d*x + c) +
3*sin(d*x + c) + 5) - 43*(25*cos(d*x + c)^2 - 30*sin(d*x + c) - 34)*log(-
4*cos(d*x + c) + 3*sin(d*x + c) + 5) + 1800*cos(d*x + c)*sin(d*x + c) + 44
0*cos(d*x + c))/(25*d*cos(d*x + c)^2 - 30*d*sin(d*x + c) - 34*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1229 vs. $2(102) = 204$.

Time = 1.43 (sec) , antiderivative size = 1229, normalized size of antiderivative = 10.88

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(-3-5*sin(d*x+c))**3,x)
```

output

```
Piecewise((x/(-3 + 5*sin(2*atan(1/3)))**3, Eq(c, -d*x - 2*atan(1/3))), (x/
(-3 + 5*sin(2*atan(3)))**3, Eq(c, -d*x - 2*atan(3))), (x/(-5*sin(c) - 3)**
3, Eq(d, 0)), (3483*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**4/(165888*
d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2
+ d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 23220*log(tan(c/2 +
d*x/2) + 3)*tan(c/2 + d*x/2)**3/(165888*d*tan(c/2 + d*x/2)**4 + 1105920*d
*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 + 1105920*d*tan(c/2 +
d*x/2) + 165888*d) + 45666*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**2/
(165888*d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*
tan(c/2 + d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) + 23220*log(t
an(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)/(165888*d*tan(c/2 + d*x/2)**4 + 1105
920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 + 1105920*d*tan(
c/2 + d*x/2) + 165888*d) + 3483*log(tan(c/2 + d*x/2) + 3)/(165888*d*tan(c/
2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)
**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 3483*log(3*tan(c/2 + d*x/2)
+ 1)*tan(c/2 + d*x/2)**4/(165888*d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/
2 + d*x/2)**3 + 2174976*d*tan(c/2 + d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2)
+ 165888*d) - 23220*log(3*tan(c/2 + d*x/2) + 1)*tan(c/2 + d*x/2)**3/(1658
88*d*tan(c/2 + d*x/2)**4 + 1105920*d*tan(c/2 + d*x/2)**3 + 2174976*d*tan(
c/2 + d*x/2)**2 + 1105920*d*tan(c/2 + d*x/2) + 165888*d) - 45666*log(3*t...
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.73

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx = \frac{40 \left(\frac{735 \sin(dx+c)}{\cos(dx+c)+1} + \frac{649 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} - \frac{75 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + 99 \right)}{\frac{60 \sin(dx+c)}{\cos(dx+c)+1} + \frac{118 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{60 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{9 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + 9} + 387 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} + 1 \right) - 387 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right) + 387 \log \left(\frac{\sin(dx+c)}{\cos(dx+c)+1} + 1 \right)$$

input `integrate(1/(-3-5*sin(d*x+c))^3,x, algorithm="maxima")`output `-1/18432*(40*(735*sin(d*x + c)/(cos(d*x + c) + 1) + 649*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 - 75*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 99)/(60*sin(d*x + c)/(cos(d*x + c) + 1) + 118*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 60*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 9*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 9) + 387*log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - 387*log(sin(d*x + c)/(cos(d*x + c) + 1) + 3))/d`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.95

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx = \frac{40 \left(75 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 649 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 735 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 99 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^2} - 387 \log \left(\left| 3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right) + 387 \log \left(\left| \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1 \right| \right)$$

input `integrate(1/(-3-5*sin(d*x+c))^3,x, algorithm="giac")`output `1/18432*(40*(75*tan(1/2*d*x + 1/2*c)^3 - 649*tan(1/2*d*x + 1/2*c)^2 - 735*tan(1/2*d*x + 1/2*c) - 99)/(3*tan(1/2*d*x + 1/2*c)^2 + 10*tan(1/2*d*x + 1/2*c) + 3)^2 - 387*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) + 387*log(abs(tan(1/2*d*x + 1/2*c) + 3)))/d`

Mupad [B] (verification not implemented)

Time = 26.87 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.03

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx$$

$$= \frac{43 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{1024 d} - \frac{-\frac{125 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{6912} + \frac{3245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{20736} + \frac{1225 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{6912} + \frac{55}{2304}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4 + \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{3} + \frac{118 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{9} + \frac{20 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{3} + 1 \right)}$$

input `int(-1/(5*sin(c + d*x) + 3)^3,x)`output `(43*atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4))/(1024*d) - ((1225*tan(c/2 + (d*x)/2))/6912 + (3245*tan(c/2 + (d*x)/2)^2)/20736 - (125*tan(c/2 + (d*x)/2)^3)/6912 + 55/2304)/(d*((20*tan(c/2 + (d*x)/2))/3 + (118*tan(c/2 + (d*x)/2)^2)/9 + (20*tan(c/2 + (d*x)/2)^3)/3 + tan(c/2 + (d*x)/2)^4 + 1)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.65

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^3} dx$$

$$= \frac{-900 \cos(dx + c) \sin(dx + c) - 220 \cos(dx + c) + 1075 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) \sin(dx + c)^2 + 1290 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 3\right) \cos(dx + c)}{(-3 - 5 \sin(c + dx))^3}$$

input `int(1/(-3-5*sin(d*x+c))^3,x)`

output

```
( - 900*cos(c + d*x)*sin(c + d*x) - 220*cos(c + d*x) + 1075*log(tan((c + d
*x)/2) + 3)*sin(c + d*x)**2 + 1290*log(tan((c + d*x)/2) + 3)*sin(c + d*x)
+ 387*log(tan((c + d*x)/2) + 3) - 1075*log(3*tan((c + d*x)/2) + 1)*sin(c +
d*x)**2 - 1290*log(3*tan((c + d*x)/2) + 1)*sin(c + d*x) - 387*log(3*tan((
c + d*x)/2) + 1) - 750*sin(c + d*x)**2 - 900*sin(c + d*x) - 270)/(2048*d*(
25*sin(c + d*x)**2 + 30*sin(c + d*x) + 9))
```

3.75 $\int \frac{1}{(-3-5 \sin(c+dx))^4} dx$

Optimal result	543
Mathematica [A] (verified)	544
Rubi [A] (verified)	544
Maple [A] (verified)	548
Fricas [A] (verification not implemented)	548
Sympy [B] (verification not implemented)	549
Maxima [B] (verification not implemented)	550
Giac [A] (verification not implemented)	550
Mupad [B] (verification not implemented)	551
Reduce [B] (verification not implemented)	551

Optimal result

Integrand size = 12, antiderivative size = 138

$$\int \frac{1}{(-3-5 \sin(c+dx))^4} dx = \frac{279 \log(3 \cos(\frac{1}{2}(c+dx)) + \sin(\frac{1}{2}(c+dx)))}{32768d} - \frac{279 \log(\cos(\frac{1}{2}(c+dx)) + 3 \sin(\frac{1}{2}(c+dx)))}{32768d} - \frac{5 \cos(c+dx)}{48d(3+5 \sin(c+dx))^3} + \frac{25 \cos(c+dx)}{512d(3+5 \sin(c+dx))^2} - \frac{995 \cos(c+dx)}{24576d(3+5 \sin(c+dx))}$$

output

```
279/32768*ln(3*cos(1/2*d*x+1/2*c)+sin(1/2*d*x+1/2*c))/d-279/32768*ln(cos(1/2*d*x+1/2*c)+3*sin(1/2*d*x+1/2*c))/d-5/48*cos(d*x+c)/d/(3+5*sin(d*x+c))^3+25/512*cos(d*x+c)/d/(3+5*sin(d*x+c))^2-995/24576*cos(d*x+c)/d/(3+5*sin(d*x+c))
```


Mathematica [A] (verified)

Time = 0.90 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.70

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx$$

$$= \frac{2511 \log\left(3 \cos\left(\frac{1}{2}(c + dx)\right) + \sin\left(\frac{1}{2}(c + dx)\right)\right) - 2511 \log\left(\cos\left(\frac{1}{2}(c + dx)\right) + 3 \sin\left(\frac{1}{2}(c + dx)\right)\right) - \frac{1}{(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^3} + \frac{1}{(3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx)))^2} + \frac{1}{3 \cos(\frac{1}{2}(c + dx)) + \sin(\frac{1}{2}(c + dx))}}{(294912d)}$$

input

```
Integrate[(-3 - 5*Sin[c + d*x])^(-4), x]
```

output

```
(2511*Log[3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]] - 2511*Log[Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2]] - 2320/(3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^2 + 720/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])^2 + 20*Sin[(c + d*x)/2]*(80/(3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2])^3 + 199/(3*Cos[(c + d*x)/2] + Sin[(c + d*x)/2]) + 240/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])^3 + 597/(Cos[(c + d*x)/2] + 3*Sin[(c + d*x)/2])))/(294912*d)
```

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.94, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$, Rules used = {3042, 3143, 25, 3042, 3233, 25, 3042, 3233, 27, 3042, 3139, 1081, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(-5 \sin(c + dx) - 3)^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{1}{(-5 \sin(c + dx) - 3)^4} dx$$

$$\downarrow \text{3143}$$

$$\frac{1}{48} \int -\frac{9 - 10 \sin(c + dx)}{(5 \sin(c + dx) + 3)^3} dx - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3}$$

$$\begin{aligned}
& \downarrow 25 \\
& -\frac{1}{48} \int \frac{9 - 10 \sin(c + dx)}{(5 \sin(c + dx) + 3)^3} dx - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3042 \\
& -\frac{1}{48} \int \frac{9 - 10 \sin(c + dx)}{(5 \sin(c + dx) + 3)^3} dx - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left(\frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} - \frac{1}{32} \int -\frac{154 - 75 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx \right) - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 25 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{154 - 75 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \int \frac{154 - 75 \sin(c + dx)}{(5 \sin(c + dx) + 3)^2} dx + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3233 \\
& \frac{1}{48} \left(\frac{1}{32} \left(\frac{1}{16} \int -\frac{837}{5 \sin(c + dx) + 3} dx - \frac{995 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \\
& \quad \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 27 \\
& \frac{1}{48} \left(\frac{1}{32} \left(-\frac{837}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{995 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \\
& \quad \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3042 \\
& \frac{1}{48} \left(\frac{1}{32} \left(-\frac{837}{16} \int \frac{1}{5 \sin(c + dx) + 3} dx - \frac{995 \cos(c + dx)}{16d(5 \sin(c + dx) + 3)} \right) + \frac{75 \cos(c + dx)}{32d(5 \sin(c + dx) + 3)^2} \right) - \\
& \quad \frac{5 \cos(c + dx)}{48d(5 \sin(c + dx) + 3)^3} \\
& \downarrow 3139
\end{aligned}$$

$$\frac{1}{48} \left(\frac{1}{32} \left(-\frac{837 \int \frac{1}{3 \tan^2(\frac{1}{2}(c+dx)) + 10 \tan(\frac{1}{2}(c+dx)) + 3} dx \tan(\frac{1}{2}(c+dx))}{8d} - \frac{995 \cos(c+dx)}{16d(5 \sin(c+dx) + 3)} \right) + \frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)} \right)$$

$$\frac{5 \cos(c+dx)}{48d(5 \sin(c+dx) + 3)^3}$$

↓ 1081

$$\frac{1}{48} \left(\frac{1}{32} \left(-\frac{2511 \int \left(\frac{1}{8(3 \tan(\frac{1}{2}(c+dx)) + 1)} - \frac{1}{24(\tan(\frac{1}{2}(c+dx)) + 3)} \right) dx \tan(\frac{1}{2}(c+dx))}{8d} - \frac{995 \cos(c+dx)}{16d(5 \sin(c+dx) + 3)} \right) + \frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)} \right)$$

$$\frac{5 \cos(c+dx)}{48d(5 \sin(c+dx) + 3)^3}$$

↓ 2009

$$\frac{1}{48} \left(\frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)^2} + \frac{1}{32} \left(-\frac{2511 \left(\frac{1}{24} \log(3 \tan(\frac{1}{2}(c+dx)) + 1) - \frac{1}{24} \log(\tan(\frac{1}{2}(c+dx)) + 3) \right)}{8d} - \frac{995 \cos(c+dx)}{16d(5 \sin(c+dx) + 3)} \right) + \frac{75 \cos(c+dx)}{32d(5 \sin(c+dx) + 3)} \right)$$

$$\frac{5 \cos(c+dx)}{48d(5 \sin(c+dx) + 3)^3}$$

input `Int[(-3 - 5*Sin[c + d*x])^(-4),x]`

output `(-5*Cos[c + d*x])/(48*d*(3 + 5*Sin[c + d*x])^3) + ((75*Cos[c + d*x])/(32*d*(3 + 5*Sin[c + d*x])^2) + ((-2511*(-1/24*Log[3 + Tan[(c + d*x)/2]] + Log[1 + 3*Tan[(c + d*x)/2]]/24))/(8*d) - (995*Cos[c + d*x])/(16*d*(3 + 5*Sin[c + d*x]))) / 32) / 48`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 1081 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[c Int[ExpandIntegrand[1/((b/2 - q/2 + c*x)*(b/2 + q/2 + c*x)), x], x], x] /; FreeQ[{a, b, c}, x] && NiceSqrtQ[b^2 - 4*a*c]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3139 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Simp[2*(e/d) Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]`
- rule 3143 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1) - b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]`
- rule 3233 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 - b^2))), x] + Simp[1/((m + 1)*(a^2 - b^2)) Int[(a + b*Sin[e + f*x])^(m + 1)*Simp[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.96

method	result
derivativedivides	$-\frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^3} + \frac{275}{27648(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} - \frac{3505}{221184(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{32768} - \frac{125}{768(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^3} + \frac{75}{1024(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} - \frac{34}{8192(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)} + \frac{279 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{32768}$
default	$-\frac{125}{20736(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^3} + \frac{275}{27648(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)^2} - \frac{3505}{221184(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)} - \frac{279 \ln(3 \tan(\frac{dx}{2} + \frac{c}{2}) + 1)}{32768} - \frac{125}{768(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^3} + \frac{75}{1024(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)^2} - \frac{34}{8192(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)} + \frac{279 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{32768}$
risch	$-\frac{-111042 e^{3i(dx+c)} + 62775 i e^{4i(dx+c)} - 119310 i e^{2i(dx+c)} + 20925 e^{5i(dx+c)} + 68625 e^{i(dx+c)} + 24875 i}{12288(5 e^{2i(dx+c)} - 5 + 6 i e^{i(dx+c)})^3} d - \frac{279 \ln(e^{i(dx+c)} + 3)}{32768}$
norman	$-\frac{7915}{12288d} - \frac{63425 \tan(\frac{dx}{2} + \frac{c}{2})}{12288d} - \frac{3047275 \tan(\frac{dx}{2} + \frac{c}{2})^3}{165888d} - \frac{15725 \tan(\frac{dx}{2} + \frac{c}{2})^5}{12288d} - \frac{296245 \tan(\frac{dx}{2} + \frac{c}{2})^2}{18432d} - \frac{270245 \tan(\frac{dx}{2} + \frac{c}{2})^4}{36864d} + \frac{279 \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + 3)}{32768}$
parallelrisc	$\frac{(-10169550 \cos(2dx+2c) + 20678085 \sin(dx+c) - 2824875 \sin(3dx+3c) + 12610242) \ln(\tan(\frac{dx}{2} + \frac{c}{2}) + \frac{1}{3}) + (10169550 \cos(dx+c) - 20678085 \sin(dx+c) + 2824875 \sin(3dx+3c) - 12610242)}{199000 \cos(dx+c)^3 - 837(225 \cos(dx+c)^2 + 5(25 \cos(dx+c)^2 - 52) \sin(dx+c) - 252) \log(4 \cos(dx+c) + 2)}$

input

```
int(1/(-3-5*sin(d*x+c))^4,x,method=_RETURNVERBOSE)
```

output

```
1/d*(-125/20736/(3*tan(1/2*d*x+1/2*c)+1)^3+275/27648/(3*tan(1/2*d*x+1/2*c)+1)^2-3505/221184/(3*tan(1/2*d*x+1/2*c)+1)-279/32768*ln(3*tan(1/2*d*x+1/2*c)+1)-125/768/(tan(1/2*d*x+1/2*c)+3)^3+75/1024/(tan(1/2*d*x+1/2*c)+3)^2-345/8192/(tan(1/2*d*x+1/2*c)+3)+279/32768*ln(tan(1/2*d*x+1/2*c)+3))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.31

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx = \frac{199000 \cos(dx + c)^3 - 837(225 \cos(dx + c)^2 + 5(25 \cos(dx + c)^2 - 52) \sin(dx + c) - 252) \log(4 \cos(dx + c) + 2)}{199000 \cos(dx + c)^3 - 837(225 \cos(dx + c)^2 + 5(25 \cos(dx + c)^2 - 52) \sin(dx + c) - 252) \log(4 \cos(dx + c) + 2)}$$

input

```
integrate(1/(-3-5*sin(d*x+c))^4,x, algorithm="fricas")
```

output

```
-1/196608*(199000*cos(d*x + c)^3 - 837*(225*cos(d*x + c)^2 + 5*(25*cos(d*x + c)^2 - 52)*sin(d*x + c) - 252)*log(4*cos(d*x + c) + 3*sin(d*x + c) + 5) + 837*(225*cos(d*x + c)^2 + 5*(25*cos(d*x + c)^2 - 52)*sin(d*x + c) - 252)*log(-4*cos(d*x + c) + 3*sin(d*x + c) + 5) - 190800*cos(d*x + c)*sin(d*x + c) - 262320*cos(d*x + c))/(225*d*cos(d*x + c)^2 + 5*(25*d*cos(d*x + c)^2 - 52*d)*sin(d*x + c) - 252*d)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2358 vs. $2(126) = 252$.

Time = 3.24 (sec) , antiderivative size = 2358, normalized size of antiderivative = 17.09

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx = \text{Too large to display}$$

input

```
integrate(1/(-3-5*sin(d*x+c))**4,x)
```

output

```
Piecewise((x/(-3 + 5*sin(2*atan(1/3)))**4, Eq(c, -d*x - 2*atan(1/3))), (x/(-3 + 5*sin(2*atan(3)))**4, Eq(c, -d*x - 2*atan(3))), (x/(-5*sin(c) - 3)**4, Eq(d, 0)), (610173*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**6/(71663616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 6101730*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**5/(71663616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 22169619*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**4/(71663616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 34802460*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**3/(71663616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2 + 716636160*d*tan(c/2 + d*x/2) + 71663616*d) + 22169619*log(tan(c/2 + d*x/2) + 3)*tan(c/2 + d*x/2)**2/(71663616*d*tan(c/2 + d*x/2)**6 + 716636160*d*tan(c/2 + d*x/2)**5 + 2603778048*d*tan(c/2 + d*x/2)**4 + 4087480320*d*tan(c/2 + d*x/2)**3 + 2603778048*d*tan(c/2 + d*x/2)**2...
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(124) = 248$.

Time = 0.04 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.99

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx = \frac{40 \left(\frac{342495 \sin(dx+c)}{\cos(dx+c)+1} + \frac{1066482 \sin(dx+c)^2}{(\cos(dx+c)+1)^2} + \frac{1218910 \sin(dx+c)^3}{(\cos(dx+c)+1)^3} + \frac{486441 \sin(dx+c)^4}{(\cos(dx+c)+1)^4} + \frac{84915 \sin(dx+c)^5}{(\cos(dx+c)+1)^5} + 42741 \right) + 22599 \log \left(\frac{3 \sin(dx+c)}{\cos(dx+c)+1} \right)}{2654208 d}$$

input `integrate(1/(-3-5*sin(d*x+c))^4,x, algorithm="maxima")`

output `-1/2654208*(40*(342495*sin(d*x + c)/(cos(d*x + c) + 1) + 1066482*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1218910*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 486441*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 84915*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 42741)/(270*sin(d*x + c)/(cos(d*x + c) + 1) + 981*sin(d*x + c)^2/(cos(d*x + c) + 1)^2 + 1540*sin(d*x + c)^3/(cos(d*x + c) + 1)^3 + 981*sin(d*x + c)^4/(cos(d*x + c) + 1)^4 + 270*sin(d*x + c)^5/(cos(d*x + c) + 1)^5 + 27*sin(d*x + c)^6/(cos(d*x + c) + 1)^6 + 27) + 22599*log(3*sin(d*x + c)/(cos(d*x + c) + 1) + 1) - 22599*log(sin(d*x + c)/(cos(d*x + c) + 1) + 3))/d`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.96

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx = \frac{40 \left(84915 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 + 486441 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 1218910 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 1066482 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 342495 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 42741 \right)}{\left(3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 10 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3 \right)^3} + 22599 \log \left(\frac{3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1} \right) / d$$

2654208 d

input `integrate(1/(-3-5*sin(d*x+c))^4,x, algorithm="giac")`

output

```
-1/2654208*(40*(84915*tan(1/2*d*x + 1/2*c)^5 + 486441*tan(1/2*d*x + 1/2*c)
^4 + 1218910*tan(1/2*d*x + 1/2*c)^3 + 1066482*tan(1/2*d*x + 1/2*c)^2 + 342
495*tan(1/2*d*x + 1/2*c) + 42741)/(3*tan(1/2*d*x + 1/2*c)^2 + 10*tan(1/2*d
*x + 1/2*c) + 3)^3 + 22599*log(abs(3*tan(1/2*d*x + 1/2*c) + 1)) - 22599*lo
g(abs(tan(1/2*d*x + 1/2*c) + 3)))/d
```

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.22

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx = \frac{279 \operatorname{atanh}\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{4} + \frac{5}{4}\right)}{16384 d} - \frac{\frac{15725 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5}{331776} + \frac{270245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{995328} + \frac{3047275 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{4478976} + \frac{296245 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{497664} + \frac{63425 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)}{331776} + \frac{7915}{331776}}{d \left(\tan\left(\frac{c}{2} + \frac{dx}{2}\right)^6 + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^5 + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^4}{3} + \frac{1540 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^3}{27} + \frac{109 \tan\left(\frac{c}{2} + \frac{dx}{2}\right)^2}{3} + 10 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 1 \right)}$$

input

```
int(1/(5*sin(c + d*x) + 3)^4,x)
```

output

```
(279*atanh((3*tan(c/2 + (d*x)/2))/4 + 5/4))/(16384*d) - ((63425*tan(c/2 +
(d*x)/2))/331776 + (296245*tan(c/2 + (d*x)/2)^2)/497664 + (3047275*tan(c/2
+ (d*x)/2)^3)/4478976 + (270245*tan(c/2 + (d*x)/2)^4)/995328 + (15725*tan
(c/2 + (d*x)/2)^5)/331776 + 7915/331776)/(d*(10*tan(c/2 + (d*x)/2) + (109*
tan(c/2 + (d*x)/2)^2)/3 + (1540*tan(c/2 + (d*x)/2)^3)/27 + (109*tan(c/2 +
(d*x)/2)^4)/3 + 10*tan(c/2 + (d*x)/2)^5 + tan(c/2 + (d*x)/2)^6 + 1))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.95

$$\int \frac{1}{(-3 - 5 \sin(c + dx))^4} dx = \frac{-298500 \cos(dx + c) \sin(dx + c)^2 - 286200 \cos(dx + c) \sin(dx + c) - 94980 \cos(dx + c) + 313875 \log\left(\frac{3 \tan\left(\frac{c}{2} + \frac{dx}{2}\right) + 5}{4}\right)}{d}$$

input `int(1/(-3-5*sin(d*x+c))^4,x)`

output `(- 298500*cos(c + d*x)*sin(c + d*x)**2 - 286200*cos(c + d*x)*sin(c + d*x)
- 94980*cos(c + d*x) + 313875*log(tan((c + d*x)/2) + 3)*sin(c + d*x)**3 +
564975*log(tan((c + d*x)/2) + 3)*sin(c + d*x)**2 + 338985*log(tan((c + d*
x)/2) + 3)*sin(c + d*x) + 67797*log(tan((c + d*x)/2) + 3) - 313875*log(3*t
an((c + d*x)/2) + 1)*sin(c + d*x)**3 - 564975*log(3*tan((c + d*x)/2) + 1)*
sin(c + d*x)**2 - 338985*log(3*tan((c + d*x)/2) + 1)*sin(c + d*x) - 67797*
log(3*tan((c + d*x)/2) + 1) - 265000*sin(c + d*x)**3 - 477000*sin(c + d*x)
2 - 286200*sin(c + d*x) - 57240)/(294912*d*(125*sin(c + d*x)3 + 225*si
n(c + d*x)**2 + 135*sin(c + d*x) + 27))`

3.76 $\int (a + b \sin(c + dx))^{7/2} dx$

Optimal result	553
Mathematica [A] (verified)	554
Rubi [A] (verified)	554
Maple [B] (verified)	559
Fricas [C] (verification not implemented)	560
Sympy [F(-1)]	561
Maxima [F]	561
Giac [F]	562
Mupad [F(-1)]	562
Reduce [F]	562

Optimal result

Integrand size = 14, antiderivative size = 256

$$\int (a + b \sin(c + dx))^{7/2} dx = -\frac{2b(71a^2 + 25b^2) \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{105d} - \frac{24ab \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{35d} - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} + \frac{32a(11a^2 + 13b^2) E\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right) \mid \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{105d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{2(71a^4 - 46a^2b^2 - 25b^4) \text{EllipticF}\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{105d \sqrt{a + b \sin(c + dx)}}$$

output

```
-2/105*b*(71*a^2+25*b^2)*cos(d*x+c)*(a+b*sin(d*x+c))^(1/2)/d-24/35*a*b*cos
(d*x+c)*(a+b*sin(d*x+c))^(3/2)/d-2/7*b*cos(d*x+c)*(a+b*sin(d*x+c))^(5/2)/d
-32/105*a*(11*a^2+13*b^2)*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x),2^(1/2)*(b/(
a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/d/((a+b*sin(d*x+c))/(a+b))^(1/2)-2/105
*(71*a^4-46*a^2*b^2-25*b^4)*InverseJacobiAM(1/2*c-1/4*Pi+1/2*d*x,2^(1/2)*(
b/(a+b))^(1/2))*((a+b*sin(d*x+c))/(a+b))^(1/2)/d/(a+b*sin(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 1.24 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.86

$$\int (a + b \sin(c + dx))^{7/2} dx = \frac{-64a(11a^3 + 11a^2b + 13ab^2 + 13b^3) E\left(\frac{1}{4}(-2c + \pi - 2dx) \middle| \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} + 4(71a^4 - 46a^3b + 13a^2b^2 - 13ab^3) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{(210d \sqrt{a + b \sin(c + dx)})}$$

input `Integrate[(a + b*Sin[c + d*x])^(7/2),x]`

output `(-64*a*(11*a^3 + 11*a^2*b + 13*a*b^2 + 13*b^3)*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)] + 4*(71*a^4 - 46*a^3*b + 13*a^2*b^2 - 13*b^3)*Sqrt[(a + b*Sin[c + d*x])/(a + b)] - b*Cos[c + d*x]*(488*a^3 + 262*a*b^2 - 162*a*b^2*Cos[2*(c + d*x)] + b*(752*a^2 + 145*b^2)*Sin[c + d*x] - 15*b^3*Sin[3*(c + d*x)]))/(210*d*Sqrt[a + b*Sin[c + d*x]])`

Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.04, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 1.286$, Rules used = {3042, 3135, 27, 3042, 3232, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(c + dx))^{7/2} dx$$

↓ 3042

$$\int (a + b \sin(c + dx))^{7/2} dx$$

↓ 3135

$$\frac{2}{7} \int \frac{1}{2} (a + b \sin(c + dx))^{3/2} (7a^2 + 12b \sin(c + dx)a + 5b^2) dx - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d}$$

↓ 27

$$\frac{1}{7} \int (a + b \sin(c + dx))^{3/2} (7a^2 + 12b \sin(c + dx)a + 5b^2) dx - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d}$$

↓ 3042

$$\frac{1}{7} \int (a + b \sin(c + dx))^{3/2} (7a^2 + 12b \sin(c + dx)a + 5b^2) dx - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d}$$

↓ 3232

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{1}{2} \sqrt{a + b \sin(c + dx)} (a(35a^2 + 61b^2) + b(71a^2 + 25b^2) \sin(c + dx)) dx - \frac{24ab \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{5d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d}$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \int \sqrt{a + b \sin(c + dx)} (a(35a^2 + 61b^2) + b(71a^2 + 25b^2) \sin(c + dx)) dx - \frac{24ab \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{5d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \int \sqrt{a + b \sin(c + dx)} (a(35a^2 + 61b^2) + b(71a^2 + 25b^2) \sin(c + dx)) dx - \frac{24ab \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{5d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d}$$

↓ 3232

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2}{3} \int \frac{105a^4 + 254b^2a^2 + 16b(11a^2 + 13b^2) \sin(c + dx)a + 25b^4}{2\sqrt{a + b \sin(c + dx)}} dx - \frac{2b(71a^2 + 25b^2) \cos(c + dx)\sqrt{a + b \sin(c + dx)}}{3d} \right) \right. \\ \left. \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} \right) \\ \downarrow 27$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{105a^4 + 254b^2a^2 + 16b(11a^2 + 13b^2) \sin(c + dx)a + 25b^4}{\sqrt{a + b \sin(c + dx)}} dx - \frac{2b(71a^2 + 25b^2) \cos(c + dx)\sqrt{a + b \sin(c + dx)}}{3d} \right) \right. \\ \left. \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} \right) \\ \downarrow 3042$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \int \frac{105a^4 + 254b^2a^2 + 16b(11a^2 + 13b^2) \sin(c + dx)a + 25b^4}{\sqrt{a + b \sin(c + dx)}} dx - \frac{2b(71a^2 + 25b^2) \cos(c + dx)\sqrt{a + b \sin(c + dx)}}{3d} \right) \right. \\ \left. \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} \right) \\ \downarrow 3231$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(16a(11a^2 + 13b^2) \int \sqrt{a + b \sin(c + dx)} dx - (71a^4 - 46a^2b^2 - 25b^4) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) \right) \right. \\ \left. \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} \right) \\ \downarrow 3042$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(16a(11a^2 + 13b^2) \int \sqrt{a + b \sin(c + dx)} dx - (71a^4 - 46a^2b^2 - 25b^4) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) \right) \right. \\ \left. \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} \right) \\ \downarrow 3134$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{16a(11a^2 + 13b^2) \sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - (71a^4 - 46a^2b^2 - 25b^4) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) \right) \right. \\ \left. \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}{7d} \right) \\ \downarrow 3042$$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{16a(11a^2 + 13b^2) \sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\frac{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}} - (71a^4 - 46a^2b^2 - 25b^4) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) \right) \right)$$

↓ 3132

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{32a(11a^2 + 13b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \frac{1}{2b \cos(c + dx)(a + b \sin(c + dx))^{5/2}}} - (71a^4 - 46a^2b^2 - 25b^4) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) \right) \right)$$

↓ 3142

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{32a(11a^2 + 13b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \frac{(71a^4 - 46a^2b^2 - 25b^4) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{\sqrt{a + b \sin(c + dx)}}} - \right) \right) \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{32a(11a^2 + 13b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \frac{(71a^4 - 46a^2b^2 - 25b^4) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{\sqrt{a + b \sin(c + dx)}}} - \right) \right) \right)$$

↓ 3140

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{32a(11a^2 + 13b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \frac{2(71a^4 - 46a^2b^2 - 25b^4) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{d \sqrt{a + b \sin(c + dx)}}} - \right) \right) \right)$$

input `Int[(a + b*Sin[c + d*x])^(7/2),x]`

output

$$\begin{aligned} & (-2*b*\cos[c + d*x]*(a + b*\sin[c + d*x])^{5/2})/(7*d) + ((-24*a*b*\cos[c + d \\ & *x]*(a + b*\sin[c + d*x])^{3/2})/(5*d) + ((-2*b*(71*a^2 + 25*b^2)*\cos[c + d \\ & *x]*\sqrt{a + b*\sin[c + d*x]})/(3*d) + ((32*a*(11*a^2 + 13*b^2)*\text{EllipticE}[(c \\ & - \pi/2 + d*x)/2, (2*b)/(a + b)]*\sqrt{a + b*\sin[c + d*x]})/(d*\sqrt{(a + b \\ & * \sin[c + d*x])/(a + b)}) - (2*(71*a^4 - 46*a^2*b^2 - 25*b^4)*\text{EllipticF}[(c \\ & - \pi/2 + d*x)/2, (2*b)/(a + b)]*\sqrt{(a + b*\sin[c + d*x])/(a + b)})/(d*\sqrt{ \\ & t[a + b*\sin[c + d*x]})/3)/5)/7 \end{aligned}$$
Defintions of rubi rules used

rule 27

$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3132

$$\text{Int}[\sqrt{(a_)} + (b_)*\sin[(c_.) + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[2*(\sqrt{a + b}/d)*\text{EllipticE}[(1/2)*(c - \pi/2 + d*x), 2*(b/(a + b))], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[a + b, 0]$$

rule 3134

$$\text{Int}[\sqrt{(a_)} + (b_)*\sin[(c_.) + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[\sqrt{a + b*\sin[c + d*x]}/\sqrt{(a + b*\sin[c + d*x])/(a + b)} \text{ Int}[\sqrt{a/(a + b)} + (b/(a + b))*\sin[c + d*x], x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ !\text{GtQ}[a + b, 0]$$

rule 3135

$$\text{Int}[(a_ + (b_)*\sin[(c_.) + (d_)*(x_)])^{n_}, x_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((a + b*\sin[c + d*x])^{n-1}/(d*n)), x] + \text{Simp}[1/n \text{ Int}[(a + b*\sin[c + d*x])^{n-2}*\text{Simp}[a^2*n + b^2*(n-1) + a*b*(2*n-1)*\sin[c + d*x], x], x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$$

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231 `Int[((c_) + (d_)*sin[(e_) + (f_)*(x_)])/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]`

rule 3232 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-d)*Cos[e + f*x]*((a + b*Sin[e + f*x])^m/(f*(m + 1))), x] + Simp[1/(m + 1) Int[(a + b*Sin[e + f*x])^(m - 1)*Simp[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && GtQ[m, 0] && IntegerQ[2*m]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1039 vs. $2(237) = 474$.

Time = 1.51 (sec) , antiderivative size = 1040, normalized size of antiderivative = 4.06

method	result	size
default	Expression too large to display	1040

input `int((a+b*sin(d*x+c))^(7/2),x,method=_RETURNVERBOSE)`

output

```

2/105*(15*b^5*sin(d*x+c)^5+105*a^5*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin
(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*si
n(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))+71*((a+b*sin(d*x+c))/(a-b))^(1
/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*Ellipt
icF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*a^4*b+78*((a+b*sin
(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(
a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*
a^3*b^2-46*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*
(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),
(a-b)/(a+b))^(1/2))*a^2*b^3-183*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*
x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d
*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*a*b^4-25*((a+b*sin(d*x+c))/(a-b))
^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*Ell
ipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^5-176*((a+b*si
n(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))
/(a-b))^(1/2)*EllipticE(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2)
)*a^5-32*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-
b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE(((a+b*sin(d*x+c))/(a-b))^(1/2),((a
-b)/(a+b))^(1/2))*a^3*b^2+208*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+
c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE(((a+b*sin(...

```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 479, normalized size of antiderivative = 1.87

$$\int (a + b \sin(c + dx))^{7/2} dx = \text{Too large to display}$$

input

```
integrate((a+b*sin(d*x+c))^(7/2),x, algorithm="fricas")
```

output

```
-2/315*((37*a^4 - 346*a^2*b^2 - 75*b^4)*sqrt(1/2*I*b)*weierstrassPInverse(
-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*
x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b) + (37*a^4 - 346*a^2*b^2 - 75*b^4)*
sqrt(-1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a
^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b
) + 48*(11*I*a^3*b + 13*I*a*b^3)*sqrt(1/2*I*b)*weierstrassZeta(-4/3*(4*a^2
- 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(
4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c)
- 3*I*b*sin(d*x + c) - 2*I*a)/b)) + 48*(-11*I*a^3*b - 13*I*a*b^3)*sqrt(-1
/2*I*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a
b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 +
9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b)) -
3*(15*b^4*cos(d*x + c)^3 - 66*a*b^3*cos(d*x + c)*sin(d*x + c) - 2*(61*a^2*
b^2 + 20*b^4)*cos(d*x + c))*sqrt(b*sin(d*x + c) + a))/(b*d)
```

Sympy [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^{7/2} dx = \text{Timed out}$$

input

```
integrate((a+b*sin(d*x+c))**(7/2),x)
```

output

Timed out

Maxima [F]

$$\int (a + b \sin(c + dx))^{7/2} dx = \int (b \sin(dx + c) + a)^{7/2} dx$$

input

```
integrate((a+b*sin(d*x+c))^(7/2),x, algorithm="maxima")
```

output

```
integrate((b*sin(d*x + c) + a)^(7/2), x)
```

Giac [F]

$$\int (a + b \sin(c + dx))^{7/2} dx = \int (b \sin(dx + c) + a)^{7/2} dx$$

input `integrate((a+b*sin(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(7/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^{7/2} dx = \int (a + b \sin(c + dx))^{7/2} dx$$

input `int((a + b*sin(c + d*x))^(7/2),x)`

output `int((a + b*sin(c + d*x))^(7/2), x)`

Reduce [F]

$$\begin{aligned} \int (a + b \sin(c + dx))^{7/2} dx &= \left(\int \sqrt{\sin(dx + c) b + a} dx \right) a^3 \\ &+ \left(\int \sqrt{\sin(dx + c) b + a} \sin(dx + c)^3 dx \right) b^3 \\ &+ 3 \left(\int \sqrt{\sin(dx + c) b + a} \sin(dx + c)^2 dx \right) a b^2 \\ &+ 3 \left(\int \sqrt{\sin(dx + c) b + a} \sin(dx + c) dx \right) a^2 b \end{aligned}$$

input `int((a+b*sin(d*x+c))^(7/2),x)`

output

```
int(sqrt(sin(c + d*x)*b + a),x)*a**3 + int(sqrt(sin(c + d*x)*b + a)*sin(c
+ d*x)**3,x)*b**3 + 3*int(sqrt(sin(c + d*x)*b + a)*sin(c + d*x)**2,x)*a*b*
*2 + 3*int(sqrt(sin(c + d*x)*b + a)*sin(c + d*x),x)*a**2*b
```

3.77 $\int (a + b \sin(c + dx))^{5/2} dx$

Optimal result	564
Mathematica [A] (verified)	565
Rubi [A] (verified)	565
Maple [B] (verified)	570
Fricas [C] (verification not implemented)	571
Sympy [F]	571
Maxima [F]	572
Giac [F]	572
Mupad [F(-1)]	572
Reduce [F]	573

Optimal result

Integrand size = 14, antiderivative size = 207

$$\int (a + b \sin(c + dx))^{5/2} dx = -\frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{15d} - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d} + \frac{2(23a^2 + 9b^2) E\left(\frac{1}{2}(c - \frac{\pi}{2} + dx) \mid \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{15d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{16a(a^2 - b^2) \text{EllipticF}\left(\frac{1}{2}(c - \frac{\pi}{2} + dx), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{15d \sqrt{a + b \sin(c + dx)}}$$

output

```
-16/15*a*b*cos(d*x+c)*(a+b*sin(d*x+c))^(1/2)/d-2/5*b*cos(d*x+c)*(a+b*sin(d*x+c))^(3/2)/d-2/15*(23*a^2+9*b^2)*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x),2^(1/2)*(b/(a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/d/((a+b*sin(d*x+c))/(a+b))^(1/2)-16/15*a*(a^2-b^2)*InverseJacobiAM(1/2*c-1/4*Pi+1/2*d*x,2^(1/2)*(b/(a+b))^(1/2))*((a+b*sin(d*x+c))/(a+b))^(1/2)/d/(a+b*sin(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 1.15 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.89

$$\int (a + b \sin(c + dx))^{5/2} dx = \frac{-2(23a^3 + 23a^2b + 9ab^2 + 9b^3) E\left(\frac{1}{4}(-2c + \pi - 2dx) \middle| \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} + 16a(a^2 - b^2) \text{EllipticF}\left(\frac{-2c + \pi - 2dx}{4}, \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} + b \cos(c + dx) (-22a^2 - 3b^2 + 3b^2 \cos[2(c + dx)] - 28ab \sin(c + dx)) / (15d \sqrt{a + b \sin(c + dx)}}}{15d \sqrt{a + b \sin(c + dx)}}$$

input `Integrate[(a + b*Sin[c + d*x])^(5/2),x]`

output `(-2*(23*a^3 + 23*a^2*b + 9*a*b^2 + 9*b^3)*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)] + 16*a*(a^2 - b^2)*EllipticF[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)] + b*cos[c + d*x]*(-22*a^2 - 3*b^2 + 3*b^2*cos[2*(c + d*x)] - 28*a*b*sin[c + d*x]))/(15*d*Sqrt[a + b*Sin[c + d*x]])`

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$, Rules used = {3042, 3135, 27, 3042, 3232, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(c + dx))^{5/2} dx$$

↓ 3042

$$\int (a + b \sin(c + dx))^{5/2} dx$$

↓ 3135

$$\frac{2}{5} \int \frac{1}{2} \sqrt{a + b \sin(c + dx)} (5a^2 + 8b \sin(c + dx)a + 3b^2) dx - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 27

$$\frac{1}{5} \int \sqrt{a + b \sin(c + dx)} (5a^2 + 8b \sin(c + dx)a + 3b^2) dx - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \int \sqrt{a + b \sin(c + dx)} (5a^2 + 8b \sin(c + dx)a + 3b^2) dx - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3232

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{a(15a^2 + 17b^2) + b(23a^2 + 9b^2) \sin(c + dx)}{2\sqrt{a + b \sin(c + dx)}} dx - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{a(15a^2 + 17b^2) + b(23a^2 + 9b^2) \sin(c + dx)}{\sqrt{a + b \sin(c + dx)}} dx - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \int \frac{a(15a^2 + 17b^2) + b(23a^2 + 9b^2) \sin(c + dx)}{\sqrt{a + b \sin(c + dx)}} dx - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3231

$$\frac{1}{5} \left(\frac{1}{3} \left((23a^2 + 9b^2) \int \sqrt{a + b \sin(c + dx)} dx - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left((23a^2 + 9b^2) \int \sqrt{a + b \sin(c + dx)} dx - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3134

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(23a^2 + 9b^2) \sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{(23a^2 + 9b^2) \sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3132

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(23a^2 + 9b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - 8a(a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3142

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(23a^2 + 9b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{8a(a^2 - b^2) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}}} dx}{\sqrt{a + b \sin(c + dx)}} \right) - \frac{16ab \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \right) - \frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(23a^2 + 9b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{8a(a^2 - b^2) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}}} \right)}{\frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}} \right)$$

↓ 3140

$$\frac{1}{5} \left(\frac{1}{3} \left(\frac{2(23a^2 + 9b^2) \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{16a(a^2 - b^2) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c - \right)}{\frac{2b \cos(c + dx)(a + b \sin(c + dx))^{3/2}}{5d}} \right)$$

input `Int[(a + b*Sin[c + d*x])^(5/2),x]`

output `(-2*b*Cos[c + d*x]*(a + b*Sin[c + d*x])^(3/2))/(5*d) + ((-16*a*b*Cos[c + d*x]*Sqrt[a + b*Sin[c + d*x]])/(3*d) + ((2*(23*a^2 + 9*b^2)*EllipticE[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[a + b*Sin[c + d*x]])/(d*Sqrt[(a + b*Sin[c + d*x])/(a + b)]) - (16*a*(a^2 - b^2)*EllipticF[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)])/(d*Sqrt[a + b*Sin[c + d*x]]))/3)/5`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 $\text{Int}[\text{Sqrt}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]]], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[a + b*\sin[c + d*x]]/\text{Sqrt}[(a + b*\sin[c + d*x])/(a + b)] \text{ Int}[\text{Sqrt}[a/(a + b) + (b/(a + b))*\sin[c + d*x]], x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{!GtQ}[a + b, 0]$

rule 3135 $\text{Int}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((a + b*\sin[c + d*x])^{(n-1)})/(d*n)], x] + \text{Simp}[1/n \text{ Int}[(a + b*\sin[c + d*x])^{(n-2)}*\text{Simp}[a^{2*n} + b^{2*(n-1)} + a*b*(2*n-1)*\sin[c + d*x], x], x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 3140 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/(d*\text{Sqrt}[a + b]))*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2*(b/(a + b))], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[a + b, 0]$

rule 3142 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[(a + b*\sin[c + d*x])/(a + b)]/\text{Sqrt}[a + b*\sin[c + d*x]] \text{ Int}[1/\text{Sqrt}[a/(a + b) + (b/(a + b))*\sin[c + d*x]], x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x\} \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{!GtQ}[a + b, 0]$

rule 3231 $\text{Int}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]/\text{Sqrt}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)/b \text{ Int}[1/\text{Sqrt}[a + b*\sin[e + f*x]], x], x] + \text{Simp}[d/b \text{ Int}[\text{Sqrt}[a + b*\sin[e + f*x]], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

rule 3232 $\text{Int}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]^{(m_)}*((c_) + (d_)*\sin[(e_) + (f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(-d)*\cos[e + f*x]*((a + b*\sin[e + f*x])^{(m)})/(f*(m + 1)), x] + \text{Simp}[1/(m + 1) \text{ Int}[(a + b*\sin[e + f*x])^{(m-1)}*\text{Simp}[b*d*m + a*c*(m + 1) + (a*d*m + b*c*(m + 1))*\sin[e + f*x], x], x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{IntegerQ}[2*m]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 889 vs. $2(192) = 384$.

Time = 0.59 (sec) , antiderivative size = 890, normalized size of antiderivative = 4.30

method	result
default	$2a^4 \sqrt{\frac{a+b \sin(dx+c)}{a-b}} \sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}} \sqrt{-\frac{b(1+\sin(dx+c))}{a-b}} \operatorname{EllipticF}\left(\sqrt{\frac{a+b \sin(dx+c)}{a-b}}, \sqrt{\frac{a-b}{a+b}}\right) + \frac{16 \sqrt{\frac{a+b \sin(dx+c)}{a-b}} \sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}}}{\dots}$

input

```
int((a+b*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
2/15*(15*a^4*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)
)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2)
,((a-b)/(a+b))^(1/2))+8*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/
(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(
a-b))^(1/2),((a-b)/(a+b))^(1/2))*a^3*b-6*a^2*((a+b*sin(d*x+c))/(a-b))^(1/2)
)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*Elliptic
F(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^2-8*a*((a+b*sin(d*
x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b
))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^3
-9*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+s
in(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a
+b))^(1/2))*b^4-23*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b)
)^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE(((a+b*sin(d*x+c))/(a-b))
^(1/2),((a-b)/(a+b))^(1/2))*a^4+14*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin
(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE(((a+b*si
n(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*a^2*b^2+9*((a+b*sin(d*x+c))/(a
-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)
*EllipticE(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^4+3*b^4*s
in(d*x+c)^4+14*a*b^3*sin(d*x+c)^3+11*a^2*b^2*sin(d*x+c)^2-3*b^4*sin(d*x+c)
^2-14*a*b^3*sin(d*x+c)-11*b^2*a^2)/b/cos(d*x+c)/(a+b*sin(d*x+c))^(1/2)/...
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.10

$$\int (a + b \sin(c + dx))^{5/2} dx =$$

$$2 \left((a^3 - 33ab^2) \sqrt{\frac{1}{2} i b \operatorname{weierstrassPInverse} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3ib \sin(dx+c) - 2ia}{3b} \right)} + (a^3 - 33ab^2) \sqrt{-\frac{1}{2} i b \operatorname{weierstrassPInverse} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3ib \sin(dx+c) - 2ia}{3b} \right)} + (a^3 - 33ab^2) \sqrt{\frac{1}{2} i b \operatorname{weierstrassZeta} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3ib \sin(dx+c) - 2ia}{3b} \right)} + (a^3 - 33ab^2) \sqrt{-\frac{1}{2} i b \operatorname{weierstrassZeta} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3ib \sin(dx+c) - 2ia}{3b} \right)} + 11ab^2 \cos(dx+c) \sqrt{b \sin(dx+c) + a} \right) / (b*d)$$

input `integrate((a+b*sin(d*x+c))^(5/2),x, algorithm="fricas")`

output `-2/45*((a^3 - 33*a*b^2)*sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b) + (a^3 - 33*a*b^2)*sqrt(-1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b) + 3*(23*I*a^2*b + 9*I*b^3)*sqrt(1/2*I*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b)) + 3*(-23*I*a^2*b - 9*I*b^3)*sqrt(-1/2*I*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b)) + 3*(3*b^3*cos(d*x + c)*sin(d*x + c) + 11*a*b^2*cos(d*x + c))*sqrt(b*sin(d*x + c) + a)/(b*d)`

Sympy [F]

$$\int (a + b \sin(c + dx))^{5/2} dx = \int (a + b \sin(c + dx))^{5/2} dx$$

input `integrate((a+b*sin(d*x+c))**(5/2),x)`

output `Integral((a + b*sin(c + d*x))**(5/2), x)`

Maxima [F]

$$\int (a + b \sin(c + dx))^{5/2} dx = \int (b \sin(dx + c) + a)^{5/2} dx$$

input `integrate((a+b*sin(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(5/2), x)`

Giac [F]

$$\int (a + b \sin(c + dx))^{5/2} dx = \int (b \sin(dx + c) + a)^{5/2} dx$$

input `integrate((a+b*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^{5/2} dx = \int (a + b \sin(c + dx))^{5/2} dx$$

input `int((a + b*sin(c + d*x))^(5/2),x)`

output `int((a + b*sin(c + d*x))^(5/2), x)`

Reduce [F]

$$\int (a + b \sin(c + dx))^{5/2} dx = \left(\int \sqrt{\sin(dx + c)b + a} dx \right) a^2$$

$$+ \left(\int \sqrt{\sin(dx + c)b + a} \sin(dx + c)^2 dx \right) b^2$$

$$+ 2 \left(\int \sqrt{\sin(dx + c)b + a} \sin(dx + c) dx \right) ab$$

input `int((a+b*sin(d*x+c))^(5/2),x)`

output `int(sqrt(sin(c + d*x)*b + a),x)*a**2 + int(sqrt(sin(c + d*x)*b + a)*sin(c + d*x)**2,x)*b**2 + 2*int(sqrt(sin(c + d*x)*b + a)*sin(c + d*x),x)*a*b`

3.78 $\int (a + b \sin(c + dx))^{3/2} dx$

Optimal result	574
Mathematica [A] (verified)	575
Rubi [A] (verified)	575
Maple [B] (verified)	579
Fricas [C] (verification not implemented)	580
Sympy [F]	580
Maxima [F]	581
Giac [F]	581
Mupad [F(-1)]	581
Reduce [F]	582

Optimal result

Integrand size = 14, antiderivative size = 167

$$\int (a + b \sin(c + dx))^{3/2} dx = -\frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} + \frac{8aE\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right) \mid \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{3d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{2(a^2 - b^2) \operatorname{EllipticF}\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{3d \sqrt{a + b \sin(c + dx)}}$$

output

```
-2/3*b*cos(d*x+c)*(a+b*sin(d*x+c))^(1/2)/d-8/3*a*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x),2^(1/2)*(b/(a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/d/((a+b*sin(d*x+c))/(a+b))^(1/2)-2/3*(a^2-b^2)*InverseJacobiAM(1/2*c-1/4*Pi+1/2*d*x,2^(1/2)*(b/(a+b))^(1/2))*((a+b*sin(d*x+c))/(a+b))^(1/2)/d/(a+b*sin(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.85

$$\int (a + b \sin(c + dx))^{3/2} dx = \frac{-2b \cos(c + dx)(a + b \sin(c + dx)) - 8a(a + b)E\left(\frac{1}{4}(-2c + \pi - 2dx) \middle| \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}} + 2}{3d\sqrt{a + b \sin(c + dx)}}$$

input `Integrate[(a + b*Sin[c + d*x])^(3/2),x]`

output `(-2*b*Cos[c + d*x]*(a + b*Sin[c + d*x]) - 8*a*(a + b)*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)] + 2*(a^2 - b^2)*EllipticF[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)]/(3*d*Sqrt[a + b*Sin[c + d*x]])`

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {3042, 3135, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \sin(c + dx))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int (a + b \sin(c + dx))^{3/2} dx \\ & \quad \downarrow \text{3135} \\ & \frac{2}{3} \int \frac{3a^2 + 4b \sin(c + dx)a + b^2}{2\sqrt{a + b \sin(c + dx)}} dx - \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\begin{aligned}
& \frac{1}{3} \int \frac{3a^2 + 4b \sin(c + dx)a + b^2}{\sqrt{a + b \sin(c + dx)}} dx - \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{3a^2 + 4b \sin(c + dx)a + b^2}{\sqrt{a + b \sin(c + dx)}} dx - \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3231} \\
& \frac{1}{3} \left(4a \int \sqrt{a + b \sin(c + dx)} dx - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \\
& \quad \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(4a \int \sqrt{a + b \sin(c + dx)} dx - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \\
& \quad \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3134} \\
& \frac{1}{3} \left(\frac{4a \sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \\
& \quad \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(\frac{4a \sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \\
& \quad \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3132} \\
& \frac{1}{3} \left(\frac{8a \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - (a^2 - b^2) \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \right) - \\
& \quad \frac{2b \cos(c + dx) \sqrt{a + b \sin(c + dx)}}{3d} \\
& \quad \downarrow \text{3142}
\end{aligned}$$

$$\frac{1}{3} \left(\frac{8a\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}(c+dx-\frac{\pi}{2})\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - \frac{(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}}dx}{\sqrt{a+b\sin(c+dx)}} \right) - \frac{2b\cos(c+dx)\sqrt{a+b\sin(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{8a\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}(c+dx-\frac{\pi}{2})\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - \frac{(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}}\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}}dx}{\sqrt{a+b\sin(c+dx)}} \right) - \frac{2b\cos(c+dx)\sqrt{a+b\sin(c+dx)}}{3d}$$

↓ 3140

$$\frac{1}{3} \left(\frac{8a\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}(c+dx-\frac{\pi}{2})\middle|\frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - \frac{2(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}}\text{EllipticF}\left(\frac{1}{2}(c+dx-\frac{\pi}{2}),\frac{2b}{a+b}\right)}{d\sqrt{a+b\sin(c+dx)}} \right) - \frac{2b\cos(c+dx)\sqrt{a+b\sin(c+dx)}}{3d}$$

input `Int[(a + b*Sin[c + d*x])^(3/2),x]`

output `(-2*b*Cos[c + d*x]*Sqrt[a + b*Sin[c + d*x]]/(3*d) + ((8*a*EllipticE[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[a + b*Sin[c + d*x]]/(d*Sqrt[(a + b*Sin[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*EllipticF[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)])/(d*Sqrt[a + b*Sin[c + d*x]])) /3`

Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`
- rule 3135 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((a + b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[1/n Int[(a + b*Sin[c + d*x])^(n - 2)*Simp[a^2*n + b^2*(n - 1) + a*b*(2*n - 1)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`
- rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3231

```
Int[((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*sin[(e_.) + (
f_.)*(x_)]], x_Symbol] := Simp[(b*c - a*d)/b Int[1/Sqrt[a + b*Sin[e + f*x
]], x], x] + Simp[d/b Int[Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b
, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 655 vs. $2(156) = 312$.

Time = 0.42 (sec) , antiderivative size = 656, normalized size of antiderivative = 3.93

method	result
default	$\frac{2a^3 \sqrt{\frac{a+b \sin(dx+c)}{a-b}} \sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}} \sqrt{-\frac{b(1+\sin(dx+c))}{a-b}} \operatorname{EllipticF}\left(\sqrt{\frac{a+b \sin(dx+c)}{a-b}}, \sqrt{\frac{a-b}{a+b}}\right) + 2\sqrt{\frac{a+b \sin(dx+c)}{a-b}} \sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}}}{\dots}$

input

```
int((a+b*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
2/3*(3*a^3*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*
(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),
((a-b)/(a+b))^(1/2))+((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b
))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b)
)^(1/2),((a-b)/(a+b))^(1/2))*a^2*b-3*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(s
in(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*
sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^2*a-((a+b*sin(d*x+c))/(a-b
))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*E
llipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^3-4*((a+b*s
in(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))
/(a-b))^(1/2)*EllipticE(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2)
)*a^3+4*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b
*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-
b)/(a+b))^(1/2))*a*b^2+b^3*sin(d*x+c)^3+a*b^2*sin(d*x+c)^2-b^3*sin(d*x+c)-
a*b^2)/b/cos(d*x+c)/(a+b*sin(d*x+c))^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 394, normalized size of antiderivative = 2.36

$$\int (a + b \sin(c + dx))^{3/2} dx =$$

$$2 \left(3 \sqrt{b \sin(dx + c) + ab^2} \cos(dx + c) + 12i a \sqrt{\frac{1}{2} i b} \text{weierstrassZeta} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3} \right), \text{weier} \right.$$

input `integrate((a+b*sin(d*x+c))^(3/2),x, algorithm="fricas")`

output

```
-2/9*(3*sqrt(b*sin(d*x + c) + a)*b^2*cos(d*x + c) + 12*I*a*sqrt(1/2*I*b)*b
*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3
, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2
)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b)) - 12*I*a*sq
rt(-1/2*I*b)*b*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 +
9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I
*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)
/b)) - (a^2 + 3*b^2)*sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2
)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(
d*x + c) - 2*I*a)/b) - (a^2 + 3*b^2)*sqrt(-1/2*I*b)*weierstrassPInverse(-4
/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x
+ c) + 3*I*b*sin(d*x + c) + 2*I*a)/b))/(b*d)
```

Sympy [F]

$$\int (a + b \sin(c + dx))^{3/2} dx = \int (a + b \sin(c + dx))^{3/2} dx$$

input `integrate((a+b*sin(d*x+c))**(3/2),x)`

output `Integral((a + b*sin(c + d*x))**(3/2), x)`

Maxima [F]

$$\int (a + b \sin(c + dx))^{3/2} dx = \int (b \sin(dx + c) + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(3/2), x)`

Giac [F]

$$\int (a + b \sin(c + dx))^{3/2} dx = \int (b \sin(dx + c) + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^{3/2} dx = \int (a + b \sin(c + dx))^{3/2} dx$$

input `int((a + b*sin(c + d*x))^(3/2),x)`

output `int((a + b*sin(c + d*x))^(3/2), x)`

Reduce [F]

$$\int (a + b \sin(c + dx))^{3/2} dx = \left(\int \sqrt{\sin(dx + c)b + a} dx \right) a \\ + \left(\int \sqrt{\sin(dx + c)b + a} \sin(dx + c) dx \right) b$$

input `int((a+b*sin(d*x+c))^(3/2),x)`

output `int(sqrt(sin(c + d*x)*b + a),x)*a + int(sqrt(sin(c + d*x)*b + a)*sin(c + d*x),x)*b`

3.79 $\int \sqrt{a + b \sin(c + dx)} dx$

Optimal result	583
Mathematica [A] (verified)	583
Rubi [A] (verified)	584
Maple [B] (verified)	585
Fricas [C] (verification not implemented)	586
Sympy [F]	586
Maxima [F]	587
Giac [F]	587
Mupad [B] (verification not implemented)	587
Reduce [F]	588

Optimal result

Integrand size = 14, antiderivative size = 62

$$\int \sqrt{a + b \sin(c + dx)} dx = \frac{2E\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right) \middle| \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}$$

output

```
-2*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x), 2^(1/2)*(b/(a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/d/((a+b*sin(d*x+c))/(a+b))^(1/2)
```

Mathematica [A] (verified)

Time = 2.58 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.98

$$\int \sqrt{a + b \sin(c + dx)} dx = -\frac{2E\left(\frac{1}{4}(-2c + \pi - 2dx) \middle| \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}$$

input

```
Integrate[Sqrt[a + b*Sin[c + d*x]], x]
```

output

```
(-2*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[a + b*Sin[c + d*x]])/(d*Sqrt[(a + b*Sin[c + d*x])/(a + b)])
```


Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3134, 3042, 3132}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + b \sin(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + b \sin(c + dx)} dx \\
 & \quad \downarrow \text{3134} \\
 & \frac{\sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b \sin(c+dx)}{a+b}}} \\
 & \quad \downarrow \text{3132} \\
 & \frac{2\sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b \sin(c+dx)}{a+b}}}
 \end{aligned}$$

input

```
Int[Sqrt[a + b*Sin[c + d*x]],x]
```

output

```
(2*EllipticE[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[a + b*Sin[c + d*x]])/
(d*Sqrt[(a + b*Sin[c + d*x])/(a + b)])
```

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(61) = 122$.

Time = 0.80 (sec) , antiderivative size = 239, normalized size of antiderivative = 3.85

method	result
default	$\frac{2(a-b)\sqrt{\frac{a+b\sin(dx+c)}{a-b}}\sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}}\sqrt{-\frac{b(1+\sin(dx+c))}{a-b}}\left(\text{EllipticF}\left(\sqrt{\frac{a+b\sin(dx+c)}{a-b}},\sqrt{\frac{a-b}{a+b}}\right)a+\text{EllipticF}\left(\sqrt{\frac{a+b\sin(dx+c)}{a-b}},\sqrt{\frac{a-b}{a+b}}\right)\right)}{b\cos(dx+c)\sqrt{a+b\sin(dx+c)}d}$
risch	Expression too large to display

input `int((a+b*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output
$$2*(a-b)*((a+b*\sin(d*x+c))/(a-b))^(1/2)*(-b*(\sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+\sin(d*x+c))/(a-b))^(1/2)/b*(\text{EllipticF}(((a+b*\sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*a+\text{EllipticF}(((a+b*\sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b-\text{EllipticE}(((a+b*\sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*a-\text{EllipticE}(((a+b*\sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b)/\cos(d*x+c)/(a+b*\sin(d*x+c))^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 351, normalized size of antiderivative = 5.66

$$\int \sqrt{a + b \sin(c + dx)} dx$$

$$= \frac{2 \left(a \sqrt{\frac{1}{2} i b \text{weierstrassPInverse} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8i a^3 - 9i ab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3i b \sin(dx+c) - 2i a}{3b} \right)} + a \sqrt{-\frac{1}{2} i b \text{weierstrassPInverse} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8i a^3 - 9i ab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3i b \sin(dx+c) - 2i a}{3b} \right)} \right)}{b}$$

input `integrate((a+b*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output `2/3*(a*sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b) + a*sqrt(-1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b) - 3*I*sqrt(1/2*I*b)*b*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b)) + 3*I*sqrt(-1/2*I*b)*b*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b)))/(b*d)`

Sympy [F]

$$\int \sqrt{a + b \sin(c + dx)} dx = \int \sqrt{a + b \sin(c + dx)} dx$$

input `integrate((a+b*sin(d*x+c))**(1/2),x)`

output `Integral(sqrt(a + b*sin(c + d*x)), x)`

Maxima [F]

$$\int \sqrt{a + b \sin(c + dx)} dx = \int \sqrt{b \sin(dx + c) + a} dx$$

input `integrate((a+b*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*sin(d*x + c) + a), x)`

Giac [F]

$$\int \sqrt{a + b \sin(c + dx)} dx = \int \sqrt{b \sin(dx + c) + a} dx$$

input `integrate((a+b*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*sin(d*x + c) + a), x)`

Mupad [B] (verification not implemented)

Time = 25.71 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

$$\int \sqrt{a + b \sin(c + dx)} dx = \frac{2 E\left(\frac{c}{2} - \frac{\pi}{4} + \frac{dx}{2} \middle| \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}$$

input `int((a + b*sin(c + d*x))^(1/2),x)`

output `(2*ellipticE(c/2 - pi/4 + (d*x)/2, (2*b)/(a + b))*(a + b*sin(c + d*x))^(1/2))/(d*((a + b*sin(c + d*x))/(a + b))^(1/2))`

Reduce [F]

$$\int \sqrt{a + b \sin(c + dx)} dx = \int \sqrt{\sin(dx + c) b + a} dx$$

input `int((a+b*sin(d*x+c))^(1/2),x)`

output `int(sqrt(sin(c + d*x)*b + a),x)`

3.80 $\int \frac{1}{\sqrt{a+b \sin(c+dx)}} dx$

Optimal result	589
Mathematica [A] (verified)	589
Rubi [A] (verified)	590
Maple [B] (verified)	591
Fricas [C] (verification not implemented)	592
Sympy [F]	592
Maxima [F]	593
Giac [F]	593
Mupad [B] (verification not implemented)	593
Reduce [F]	594

Optimal result

Integrand size = 14, antiderivative size = 62

$$\int \frac{1}{\sqrt{a+b \sin(c+dx)}} dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{d \sqrt{a+b \sin(c+dx)}}$$

output `2*InverseJacobiAM(1/2*c-1/4*Pi+1/2*d*x,2^(1/2)*(b/(a+b))^(1/2))*((a+b*sin(d*x+c))/(a+b))^(1/2)/d/(a+b*sin(d*x+c))^(1/2)`

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.98

$$\int \frac{1}{\sqrt{a+b \sin(c+dx)}} dx = -\frac{2 \operatorname{EllipticF}\left(\frac{1}{4}(-2c + \pi - 2dx), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{d \sqrt{a+b \sin(c+dx)}}$$

input `Integrate[1/Sqrt[a + b*Sin[c + d*x]],x]`

output `(-2*EllipticF[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)])/(d*Sqrt[a + b*Sin[c + d*x]])`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx \\
 & \quad \downarrow \text{3142} \\
 & \frac{\sqrt{\frac{a+b \sin(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}}} dx}{\sqrt{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{\frac{a+b \sin(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}}} dx}{\sqrt{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{3140} \\
 & \frac{2\sqrt{\frac{a+b \sin(c+dx)}{a+b}} \text{EllipticF}\left(\frac{1}{2}(c + dx - \frac{\pi}{2}), \frac{2b}{a+b}\right)}{d\sqrt{a + b \sin(c + dx)}}
 \end{aligned}$$

input `Int[1/Sqrt[a + b*Sin[c + d*x]],x]`

output `(2*EllipticF[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)])/(d*Sqrt[a + b*Sin[c + d*x]])`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3140 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/(d*Sqrt[a + b]))*EllipticF[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3142 `Int[1/Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[(a + b*Sin[c + d*x])/(a + b)]/Sqrt[a + b*Sin[c + d*x]] Int[1/Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 125 vs. 2(60) = 120.

Time = 0.12 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.03

method	result	size
default	$\frac{2(a-b)\sqrt{\frac{a+b\sin(dx+c)}{a-b}}\sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}}\sqrt{-\frac{b(1+\sin(dx+c))}{a-b}}\operatorname{EllipticF}\left(\sqrt{\frac{a+b\sin(dx+c)}{a-b}},\sqrt{\frac{a-b}{a+b}}\right)}{b\cos(dx+c)\sqrt{a+b\sin(dx+c)}d}$	126

input `int(1/(a+b*sin(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2*(a-b)*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))/b/cos(d*x+c)/(a+b*sin(d*x+c))^(1/2)/d`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 143, normalized size of antiderivative = 2.31

$$\int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx$$

$$= \frac{2 \left(\sqrt{\frac{1}{2} i b} \operatorname{weierstrassPInverse} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3}, \frac{3b \cos(dx+c) - 3ib \sin(dx+c) - 2ia}{3b} \right) + \sqrt{-\frac{1}{2} i} b \operatorname{weierstrassPInverse} \left(-\frac{4(4a^2 - 3b^2)}{3b^2}, -\frac{8(8ia^3 - 9iab^2)}{27b^3}, \frac{3b \cos(dx+c) + 3ib \sin(dx+c) + 2ia}{3b} \right) \right)}{bd}$$

input `integrate(1/(a+b*sin(d*x+c))^(1/2),x, algorithm="fricas")`

output

```
2*(sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b) + sqrt(-1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b))/(b*d)
```

Sympy [F]

$$\int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx = \int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx$$

input `integrate(1/(a+b*sin(d*x+c))**(1/2),x)`

output

```
Integral(1/sqrt(a + b*sin(c + d*x)), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx = \int \frac{1}{\sqrt{b \sin(dx + c) + a}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*sin(d*x + c) + a), x)`

Giac [F]

$$\int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx = \int \frac{1}{\sqrt{b \sin(dx + c) + a}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(b*sin(d*x + c) + a), x)`

Mupad [B] (verification not implemented)

Time = 25.85 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx = -\frac{2 F\left(\frac{\pi}{4} - \frac{c}{2} - \frac{dx}{2} \middle| \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{d \sqrt{a + b \sin(c + dx)}}$$

input `int(1/(a + b*sin(c + d*x))^(1/2),x)`

output `-(2*ellipticF(pi/4 - c/2 - (d*x)/2, (2*b)/(a + b))*((a + b*sin(c + d*x))/(a + b))^(1/2))/(d*(a + b*sin(c + d*x))^(1/2))`

Reduce [F]

$$\int \frac{1}{\sqrt{a + b \sin(c + dx)}} dx = \int \frac{\sqrt{\sin(dx + c) b + a}}{\sin(dx + c) b + a} dx$$

input `int(1/(a+b*sin(d*x+c))^(1/2),x)`

output `int(sqrt(sin(c + d*x)*b + a)/(sin(c + d*x)*b + a),x)`

3.81 $\int \frac{1}{(a+b \sin(c+dx))^{3/2}} dx$

Optimal result	595
Mathematica [A] (verified)	595
Rubi [A] (verified)	596
Maple [B] (verified)	598
Fricas [C] (verification not implemented)	599
Sympy [F]	599
Maxima [F]	600
Giac [F]	600
Mupad [F(-1)]	600
Reduce [F]	601

Optimal result

Integrand size = 14, antiderivative size = 111

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \frac{2b \cos(c + dx)}{(a^2 - b^2) d \sqrt{a + b \sin(c + dx)}} + \frac{2E\left(\frac{1}{2}\left(c - \frac{\pi}{2} + dx\right) \middle| \frac{2b}{a+b}\right) \sqrt{a + b \sin(c + dx)}}{(a^2 - b^2) d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}$$

output

```
2*b*cos(d*x+c)/(a^2-b^2)/d/(a+b*sin(d*x+c))^(1/2)-2*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x),2^(1/2)*(b/(a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/(a^2-b^2)/d/((a+b*sin(d*x+c))/(a+b))^(1/2)
```

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \frac{2b \cos(c + dx) - 2(a + b)E\left(\frac{1}{4}(-2c + \pi - 2dx) \middle| \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{(a - b)(a + b)d \sqrt{a + b \sin(c + dx)}}$$

input

```
Integrate[(a + b*Sin[c + d*x])^(-3/2),x]
```

output

```
(2*b*cos[c + d*x] - 2*(a + b)*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*sqrt[(a + b*sin[c + d*x])/(a + b)])/((a - b)*(a + b)*d*sqrt[a + b*sin[c + d*x]])
```

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 3143, 27, 3042, 3134, 3042, 3132}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3143} \\
 & \frac{2b \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}} - \frac{2 \int -\frac{1}{2} \sqrt{a + b \sin(c + dx)} dx}{a^2 - b^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \sqrt{a + b \sin(c + dx)} dx}{a^2 - b^2} + \frac{2b \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sqrt{a + b \sin(c + dx)} dx}{a^2 - b^2} + \frac{2b \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{3134} \\
 & \frac{\sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{(a^2 - b^2) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} + \frac{2b \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{\sqrt{a + b \sin(c + dx)} \int \sqrt{\frac{a}{a+b} + \frac{b \sin(c+dx)}{a+b}} dx}{(a^2 - b^2) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} + \frac{2b \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}}$$

↓ 3132

$$\frac{2b \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}} + \frac{2\sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a+b}\right)}{d(a^2 - b^2) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}$$

input

```
Int[(a + b*Sin[c + d*x])^(-3/2),x]
```

output

```
(2*b*Cos[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Sin[c + d*x]]) + (2*EllipticE
[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[a + b*Sin[c + d*x]])/((a^2 - b^2)
*d*Sqrt[(a + b*Sin[c + d*x])/(a + b)])
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3132

```
Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[2*(Sqrt[a
+ b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]
```

rule 3134

```
Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[Sqrt[a +
b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (
b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2
, 0] && !GtQ[a + b, 0]
```

rule 3143

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos
[c + d*x]*((a + b*Sin[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 - b^2))), x] + Simp
[1/((n + 1)*(a^2 - b^2)) Int[(a + b*Sin[c + d*x])^(n + 1)*Simp[a*(n + 1)
- b*(n + 2)*Sin[c + d*x], x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 -
b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 442 vs. $2(108) = 216$.

Time = 0.18 (sec) , antiderivative size = 443, normalized size of antiderivative = 3.99

method	result
default	$\frac{2a^2 \sqrt{\frac{a+b \sin(dx+c)}{a-b}} \sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}} \sqrt{-\frac{b(1+\sin(dx+c))}{a-b}} \operatorname{EllipticF}\left(\sqrt{\frac{a+b \sin(dx+c)}{a-b}}, \sqrt{\frac{a-b}{a+b}}\right) - 2 \sqrt{\frac{a+b \sin(dx+c)}{a-b}} \sqrt{-\frac{b(\sin(dx+c)-1)}{a+b}}}{\dots}$

input

```
int(1/(a+b*sin(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

output

```
2/b*(a^2*((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-
b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a
-b)/(a+b))^(1/2))-((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x+c)-1)/(a+b))
^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(
1/2),((a-b)/(a+b))^(1/2))*b^2-((a+b*sin(d*x+c))/(a-b))^(1/2)*(-b*(sin(d*x
+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE(((a+b*sin(d*
x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*a^2+((a+b*sin(d*x+c))/(a-b))^(1/2)
*(-b*(sin(d*x+c)-1)/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)*EllipticE
(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))*b^2-b^2*sin(d*x+c)^2+
b^2)/(a^2-b^2)/cos(d*x+c)/(a+b*sin(d*x+c))^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 460, normalized size of antiderivative = 4.14

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \frac{2 \left(3 \sqrt{b \sin(dx + c) + ab^2} \cos(dx + c) + (ab \sin(dx + c) + a^2) \sqrt{\frac{1}{2} i b \text{weierst}} \right)}{(a + b \sin(c + dx))^{3/2}}$$

input `integrate(1/(a+b*sin(d*x+c))^(3/2),x, algorithm="fricas")`

output `2/3*(3*sqrt(b*sin(d*x + c) + a)*b^2*cos(d*x + c) + (a*b*sin(d*x + c) + a^2)*sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b) + (a*b*sin(d*x + c) + a^2)*sqrt(-1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b) + 3*(-I*b^2*sin(d*x + c) - I*a*b)*sqrt(1/2*I*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b)) + 3*(I*b^2*sin(d*x + c) + I*a*b)*sqrt(-1/2*I*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b)))/((a^2*b^2 - b^4)*d*sin(d*x + c) + (a^3*b - a*b^3)*d)`

Sympy [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a + b \sin(c + dx))^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))**(3/2),x)`

output `Integral((a + b*sin(c + d*x))**(-3/2), x)`

Maxima [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(-3/2), x)`

Giac [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(-3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx$$

input `int(1/(a + b*sin(c + d*x))^(3/2),x)`

output `int(1/(a + b*sin(c + d*x))^(3/2), x)`

Reduce [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{3/2}} dx = \int \frac{\sqrt{\sin(dx + c)b + a}}{\sin(dx + c)^2 b^2 + 2 \sin(dx + c) ab + a^2} dx$$

input `int(1/(a+b*sin(d*x+c))^(3/2),x)`

output `int(sqrt(sin(c + d*x)*b + a)/(sin(c + d*x)**2*b**2 + 2*sin(c + d*x)*a*b + a**2),x)`

3.82 $\int \frac{1}{(a+b \sin(c+dx))^{5/2}} dx$

Optimal result	602
Mathematica [A] (verified)	603
Rubi [A] (verified)	603
Maple [B] (verified)	607
Fricas [C] (verification not implemented)	608
Sympy [F]	609
Maxima [F]	610
Giac [F]	610
Mupad [F(-1)]	610
Reduce [F]	611

Optimal result

Integrand size = 14, antiderivative size = 231

$$\int \frac{1}{(a+b \sin(c+dx))^{5/2}} dx = \frac{2b \cos(c+dx)}{3(a^2-b^2)d(a+b \sin(c+dx))^{3/2}} + \frac{8ab \cos(c+dx)}{3(a^2-b^2)^2 d \sqrt{a+b \sin(c+dx)}} + \frac{8aE\left(\frac{1}{2}(c-\frac{\pi}{2}+dx) \mid \frac{2b}{a+b}\right) \sqrt{a+b \sin(c+dx)}}{3(a^2-b^2)^2 d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(c-\frac{\pi}{2}+dx), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{3(a^2-b^2)d \sqrt{a+b \sin(c+dx)}}$$

output

```
2/3*b*cos(d*x+c)/(a^2-b^2)/d/(a+b*sin(d*x+c))^(3/2)+8/3*a*b*cos(d*x+c)/(a^2-b^2)^2/d/(a+b*sin(d*x+c))^(1/2)-8/3*a*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x),2^(1/2)*(b/(a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/(a^2-b^2)^2/d/((a+b*sin(d*x+c))/(a+b))^(1/2)-2/3*InverseJacobiAM(1/2*c-1/4*Pi+1/2*d*x,2^(1/2)*(b/(a+b))^(1/2))*((a+b*sin(d*x+c))/(a+b))^(1/2)/(a^2-b^2)/d/(a+b*sin(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 1.15 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.72

$$\int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx = \frac{2 \left(-4a(a + b)^2 E\left(\frac{1}{4}(-2c + \pi - 2dx) \middle| \frac{2b}{a+b}\right) \left(\frac{a+b \sin(c+dx)}{a+b}\right)^{3/2} + (a - b)(a + b) \right)}{3(a - b)}$$

input `Integrate[(a + b*Sin[c + d*x])^(-5/2),x]`

output

```
(2*(-4*a*(a + b)^2*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*((a + b)*Sin[c + d*x])/(a + b))^(3/2) + (a - b)*(a + b)^2*EllipticF[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)]*((a + b*Sin[c + d*x])/(a + b))^(3/2) + b*Cos[c + d*x]*(5*a^2 - b^2 + 4*a*b*Sin[c + d*x]))/(3*(a - b)^2*(a + b)^2*d*(a + b*Sin[c + d*x])^(3/2))
```

Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$, Rules used = {3042, 3143, 27, 3042, 3233, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3143} \\ & \frac{2b \cos(c + dx)}{3d(a^2 - b^2)(a + b \sin(c + dx))^{3/2}} - \frac{2 \int -\frac{3a - b \sin(c + dx)}{2(a + b \sin(c + dx))^{3/2}} dx}{3(a^2 - b^2)} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\begin{aligned}
& \frac{\int \frac{3a-b \sin(c+dx)}{(a+b \sin(c+dx))^{3/2}} dx}{3(a^2-b^2)} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{3a-b \sin(c+dx)}{(a+b \sin(c+dx))^{3/2}} dx}{3(a^2-b^2)} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} \\
& \quad \downarrow \text{3233} \\
& \frac{\frac{8ab \cos(c+dx)}{d(a^2-b^2)\sqrt{a+b \sin(c+dx)}} - \frac{2 \int -\frac{3a^2+4b \sin(c+dx)a+b^2}{2\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2}}{3(a^2-b^2)} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\frac{\int \frac{3a^2+4b \sin(c+dx)a+b^2}{\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2} + \frac{8ab \cos(c+dx)}{d(a^2-b^2)\sqrt{a+b \sin(c+dx)}}}{3(a^2-b^2)} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{\int \frac{3a^2+4b \sin(c+dx)a+b^2}{\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2} + \frac{8ab \cos(c+dx)}{d(a^2-b^2)\sqrt{a+b \sin(c+dx)}}}{3(a^2-b^2)} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} \\
& \quad \downarrow \text{3231} \\
& \frac{\frac{4a \int \sqrt{a+b \sin(c+dx)} dx - (a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2} + \frac{8ab \cos(c+dx)}{d(a^2-b^2)\sqrt{a+b \sin(c+dx)}}}{\frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}}} \\
& \quad \downarrow \text{3042} \\
& \frac{\frac{4a \int \sqrt{a+b \sin(c+dx)} dx - (a^2-b^2) \int \frac{1}{\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2} + \frac{8ab \cos(c+dx)}{d(a^2-b^2)\sqrt{a+b \sin(c+dx)}}}{\frac{3(a^2-b^2)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} + \frac{2b \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}}} \\
& \quad \downarrow \text{3134}
\end{aligned}$$

$$\begin{aligned}
 & \frac{4a\sqrt{a+b\sin(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - (a^2-b^2) \int \frac{1}{\sqrt{a+b\sin(c+dx)}} dx \\
 & \frac{\hspace{10em}}{a^2-b^2} + \frac{8ab\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \\
 & \frac{3(a^2-b^2)}{2b\cos(c+dx)} \\
 & \frac{\hspace{10em}}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4a\sqrt{a+b\sin(c+dx)} \int \sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx)}{a+b}} dx}{\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - (a^2-b^2) \int \frac{1}{\sqrt{a+b\sin(c+dx)}} dx \\
 & \frac{\hspace{10em}}{a^2-b^2} + \frac{8ab\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \\
 & \frac{3(a^2-b^2)}{2b\cos(c+dx)} \\
 & \frac{\hspace{10em}}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3132} \\
 & \frac{8a\sqrt{a+b\sin(c+dx)} E\left(\frac{1}{2}(c+dx-\frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - (a^2-b^2) \int \frac{1}{\sqrt{a+b\sin(c+dx)}} dx \\
 & \frac{\hspace{10em}}{a^2-b^2} + \frac{8ab\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \\
 & \frac{3(a^2-b^2)}{2b\cos(c+dx)} \\
 & \frac{\hspace{10em}}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3142} \\
 & \frac{8a\sqrt{a+b\sin(c+dx)} E\left(\frac{1}{2}(c+dx-\frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - \frac{(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx)}{a+b}}} dx}{\sqrt{a+b\sin(c+dx)}} \\
 & \frac{\hspace{10em}}{a^2-b^2} + \frac{8ab\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \\
 & \frac{3(a^2-b^2)}{2b\cos(c+dx)} \\
 & \frac{\hspace{10em}}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{8a\sqrt{a+b\sin(c+dx)} E\left(\frac{1}{2}(c+dx-\frac{\pi}{2}) \middle| \frac{2b}{a+b}\right)}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - \frac{(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}} \int \frac{1}{\sqrt{\frac{a}{a+b} + \frac{b\sin(c+dx)}{a+b}}} dx}{\sqrt{a+b\sin(c+dx)}} \\
 & \frac{\hspace{10em}}{a^2-b^2} + \frac{8ab\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \\
 & \frac{3(a^2-b^2)}{2b\cos(c+dx)} \\
 & \frac{\hspace{10em}}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3140}
 \end{aligned}$$

$$\frac{\frac{2b \cos(c + dx)}{3d(a^2 - b^2)(a + b \sin(c + dx))^{3/2}} + \frac{8a \sqrt{a + b \sin(c + dx)} E\left(\frac{1}{2}(c + dx - \frac{\pi}{2}) \mid \frac{2b}{a + b}\right) - 2(a^2 - b^2) \sqrt{\frac{a + b \sin(c + dx)}{a + b}} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx - \frac{\pi}{2}), \frac{2b}{a + b}\right)}{d \sqrt{\frac{a + b \sin(c + dx)}{a + b}}} - \frac{2(a^2 - b^2)}{a^2 - b^2}}{\frac{8ab \cos(c + dx)}{d(a^2 - b^2) \sqrt{a + b \sin(c + dx)}} + \frac{2(a^2 - b^2)}{a^2 - b^2}} \frac{1}{3(a^2 - b^2)}$$

input `Int[(a + b*Sin[c + d*x])^(-5/2), x]`

output `(2*b*Cos[c + d*x])/(3*(a^2 - b^2)*d*(a + b*Sin[c + d*x])^(3/2)) + ((8*a*b*Cos[c + d*x])/((a^2 - b^2)*d*Sqrt[a + b*Sin[c + d*x]]) + ((8*a*EllipticE[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[a + b*Sin[c + d*x]])/(d*Sqrt[(a + b*Sin[c + d*x])/(a + b)]) - (2*(a^2 - b^2)*EllipticF[(c - Pi/2 + d*x)/2, (2*b)/(a + b)]*Sqrt[(a + b*Sin[c + d*x])/(a + b)])/(d*Sqrt[a + b*Sin[c + d*x]])))/(a^2 - b^2))/(3*(a^2 - b^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3132 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[2*(Sqrt[a + b]/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2*(b/(a + b))], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[a + b, 0]`

rule 3134 `Int[Sqrt[(a_) + (b_)*sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[Sqrt[a + b*Sin[c + d*x]]/Sqrt[(a + b*Sin[c + d*x])/(a + b)] Int[Sqrt[a/(a + b) + (b/(a + b))*Sin[c + d*x]], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]`

rule 3140 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]]], x_Symbol] \rightarrow \text{Simp}[(2/(d*\text{Sqrt}[a + b]))*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2*(b/(a + b))], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[a + b, 0]$

rule 3142 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]]], x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[(a + b*\sin[c + d*x])/(a + b)]/\text{Sqrt}[a + b*\sin[c + d*x]] \ \text{Int}[1/\text{Sqrt}[a/(a + b) + (b/(a + b))*\sin[c + d*x]], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ !\text{GtQ}[a + b, 0]$

rule 3143 $\text{Int}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((a + b*\sin[c + d*x])^{(n + 1)}/(d*(n + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \ \text{Int}[(a + b*\sin[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\sin[c + d*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

rule 3231 $\text{Int}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]/\text{Sqrt}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)/b \ \text{Int}[1/\text{Sqrt}[a + b*\sin[e + f*x]], x], x] + \text{Simp}[d/b \ \text{Int}[\text{Sqrt}[a + b*\sin[e + f*x]], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

rule 3233 $\text{Int}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]^{(m_)}*((c_) + (d_)*\sin[(e_) + (f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*\text{Cos}[e + f*x]*((a + b*\sin[e + f*x])^{(m + 1)}/(f*(m + 1)*(a^2 - b^2))), x] + \text{Simp}[1/((m + 1)*(a^2 - b^2)) \ \text{Int}[(a + b*\sin[e + f*x])^{(m + 1)}*\text{Simp}[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*\sin[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegerQ}[2*m]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 494 vs. $2(216) = 432$.

Time = 0.24 (sec) , antiderivative size = 495, normalized size of antiderivative = 2.14

method	result
default	$\frac{\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2} \left(\frac{2\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2}}{3(a^2-b^2)b\left(\sin(dx+c)+\frac{a}{b}\right)^2} + \frac{8b\cos(dx+c)^2a}{3(a^2-b^2)^2\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2}} + \frac{2(3a^2+b^2)\left(\frac{a}{b}-1\right)}{3(a^2-b^2)^2\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2}} \right)}{\dots}$

input `int(1/(a+b*sin(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & (-(-a-b*\sin(d*x+c))*\cos(d*x+c)^2)^{(1/2)}*(2/3/(a^2-b^2)/b*(-(-a-b*\sin(d*x+c) \\ &))*\cos(d*x+c)^2)^{(1/2)}/(\sin(d*x+c)+a/b)^2+8/3*b*\cos(d*x+c)^2/(a^2-b^2)^2*a \\ & /(-(-a-b*\sin(d*x+c))*\cos(d*x+c)^2)^{(1/2)}+2*(3*a^2+b^2)/(3*a^4-6*a^2*b^2+3* \\ & b^4)*(a/b-1)*((a+b*\sin(d*x+c))/(a-b))^{(1/2)}*(b*(1-\sin(d*x+c)))/(a+b)^{(1/2)} \\ & *(-b*(1+\sin(d*x+c))/(a-b))^{(1/2)}/(-(-a-b*\sin(d*x+c))*\cos(d*x+c)^2)^{(1/2)}*E \\ & llipticF(((a+b*\sin(d*x+c))/(a-b))^{(1/2)},((a-b)/(a+b))^{(1/2)})+8/3*a*b/(a^2- \\ & b^2)^2*(a/b-1)*((a+b*\sin(d*x+c))/(a-b))^{(1/2)}*(b*(1-\sin(d*x+c)))/(a+b)^{(1/2)} \\ & *(-b*(1+\sin(d*x+c))/(a-b))^{(1/2)}/(-(-a-b*\sin(d*x+c))*\cos(d*x+c)^2)^{(1/2)} \\ & *((-a/b-1)*EllipticE(((a+b*\sin(d*x+c))/(a-b))^{(1/2)},((a-b)/(a+b))^{(1/2)})+E \\ & llipticF(((a+b*\sin(d*x+c))/(a-b))^{(1/2)},((a-b)/(a+b))^{(1/2)}))/\cos(d*x+c)/ \\ & (a+b*\sin(d*x+c))^{(1/2)}/d \end{aligned}$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 687, normalized size of antiderivative = 2.97

$$\int \frac{1}{(a+b\sin(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(1/(a+b*sin(d*x+c))^(5/2),x, algorithm="fricas")`

output

```

-2/9*((a^4 + 4*a^2*b^2 + 3*b^4 - (a^2*b^2 + 3*b^4)*cos(d*x + c)^2 + 2*(a^3
*b + 3*a*b^3)*sin(d*x + c))*sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2
- 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I
*b*sin(d*x + c) - 2*I*a)/b) + (a^4 + 4*a^2*b^2 + 3*b^4 - (a^2*b^2 + 3*b^4)
*cos(d*x + c)^2 + 2*(a^3*b + 3*a*b^3)*sin(d*x + c))*sqrt(-1/2*I*b)*weierst
rassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1
/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b) - 12*(-I*a*b^3*cos(d
*x + c)^2 + 2*I*a^2*b^2*sin(d*x + c) + I*a^3*b + I*a*b^3)*sqrt(1/2*I*b)*we
ierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, w
eierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b
^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a)/b)) - 12*(I*a*b^3*
cos(d*x + c)^2 - 2*I*a^2*b^2*sin(d*x + c) - I*a^3*b - I*a*b^3)*sqrt(-1/2*I
*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)
/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*
a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b)) + 3*(4
*a*b^3*cos(d*x + c)*sin(d*x + c) + (5*a^2*b^2 - b^4)*cos(d*x + c))*sqrt(b*
sin(d*x + c) + a)/((a^4*b^3 - 2*a^2*b^5 + b^7)*d*cos(d*x + c)^2 - 2*(a^5*
b^2 - 2*a^3*b^4 + a*b^6)*d*sin(d*x + c) - (a^6*b - a^4*b^3 - a^2*b^5 + b^7
)*d)

```

Sympy [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a + b \sin(c + dx))^{\frac{5}{2}}} dx$$

input

```
integrate(1/(a+b*sin(d*x+c))**(5/2), x)
```

output

```
Integral((a + b*sin(c + d*x))**(-5/2), x)
```

Maxima [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{5/2}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(-5/2), x)`

Giac [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{5/2}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(-5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx = \int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx$$

input `int(1/(a + b*sin(c + d*x))^(5/2),x)`

output `int(1/(a + b*sin(c + d*x))^(5/2), x)`

Reduce [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{5/2}} dx = \int \frac{\sqrt{\sin(dx + c)b + a}}{\sin(dx + c)^3 b^3 + 3 \sin(dx + c)^2 a b^2 + 3 \sin(dx + c) a^2 b + a^3} dx$$

input `int(1/(a+b*sin(d*x+c))^(5/2),x)`

output `int(sqrt(sin(c + d*x)*b + a)/(sin(c + d*x)**3*b**3 + 3*sin(c + d*x)**2*a*b**2 + 3*sin(c + d*x)*a**2*b + a**3),x)`

3.83 $\int \frac{1}{(a+b \sin(c+dx))^{7/2}} dx$

Optimal result	612
Mathematica [A] (verified)	613
Rubi [A] (verified)	613
Maple [B] (verified)	618
Fricas [C] (verification not implemented)	619
Sympy [F]	620
Maxima [F]	621
Giac [F]	621
Mupad [F(-1)]	621
Reduce [F]	622

Optimal result

Integrand size = 14, antiderivative size = 292

$$\int \frac{1}{(a+b \sin(c+dx))^{7/2}} dx = \frac{2b \cos(c+dx)}{5(a^2-b^2)d(a+b \sin(c+dx))^{5/2}} + \frac{16ab \cos(c+dx)}{15(a^2-b^2)^2 d(a+b \sin(c+dx))^{3/2}} + \frac{2b(23a^2+9b^2) \cos(c+dx)}{15(a^2-b^2)^3 d \sqrt{a+b \sin(c+dx)}} + \frac{2(23a^2+9b^2) E\left(\frac{1}{2}(c-\frac{\pi}{2}+dx) \mid \frac{2b}{a+b}\right) \sqrt{a+b \sin(c+dx)}}{15(a^2-b^2)^3 d \sqrt{\frac{a+b \sin(c+dx)}{a+b}}} - \frac{16a \operatorname{EllipticF}\left(\frac{1}{2}(c-\frac{\pi}{2}+dx), \frac{2b}{a+b}\right) \sqrt{\frac{a+b \sin(c+dx)}{a+b}}}{15(a^2-b^2)^2 d \sqrt{a+b \sin(c+dx)}}$$

output

```
2/5*b*cos(d*x+c)/(a^2-b^2)/d/(a+b*sin(d*x+c))^(5/2)+16/15*a*b*cos(d*x+c)/(a^2-b^2)^2/d/(a+b*sin(d*x+c))^(3/2)+2/15*b*(23*a^2+9*b^2)*cos(d*x+c)/(a^2-b^2)^3/d/(a+b*sin(d*x+c))^(1/2)-2/15*(23*a^2+9*b^2)*EllipticE(cos(1/2*c+1/4*Pi+1/2*d*x),2^(1/2)*(b/(a+b))^(1/2))*(a+b*sin(d*x+c))^(1/2)/(a^2-b^2)^3/d/((a+b*sin(d*x+c))/(a+b))^(1/2)-16/15*a*InverseJacobiAM(1/2*c-1/4*Pi+1/2*d*x,2^(1/2)*(b/(a+b))^(1/2))*((a+b*sin(d*x+c))/(a+b))^(1/2)/(a^2-b^2)^2/d/(a+b*sin(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 1.55 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.68

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \frac{2 \left(-\frac{((23a^2 + 9b^2)E(\frac{1}{4}(-2c + \pi - 2dx)|\frac{2b}{a+b}) + 8a(-a+b) \text{EllipticF}(\frac{1}{4}(-2c + \pi - 2dx), \frac{2b}{a+b}))}{(a-b)^3} \right) \left(\frac{a+b \sin(c+dx)}{a+b} \right)}{15d(a + b \sin(c + dx))^{5/2}}$$

input

```
Integrate[(a + b*Sin[c + d*x])^(-7/2), x]
```

output

```
(2*(-(((23*a^2 + 9*b^2)*EllipticE[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)] + 8*a*(-a + b)*EllipticF[(-2*c + Pi - 2*d*x)/4, (2*b)/(a + b)])*(a + b*Sin[c + d*x])/(a + b))^(5/2))/(a - b)^3 + (b*Cos[c + d*x]*(34*a^4 - 5*a^2*b^2 + 3*b^4 + 2*a*b*(27*a^2 + 5*b^2)*Sin[c + d*x] + b^2*(23*a^2 + 9*b^2)*Sin[c + d*x]^2)/(a^2 - b^2)^3))/(15*d*(a + b*Sin[c + d*x])^(5/2))
```

Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.09, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 1.286$, Rules used = {3042, 3143, 27, 3042, 3233, 27, 3042, 3233, 27, 3042, 3231, 3042, 3134, 3042, 3132, 3142, 3042, 3140}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx$$

↓ 3042

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx$$

↓ 3143

$$\frac{2b \cos(c + dx)}{5d(a^2 - b^2)(a + b \sin(c + dx))^{5/2}} - \frac{2 \int -\frac{5a - 3b \sin(c + dx)}{2(a + b \sin(c + dx))^{5/2}} dx}{5(a^2 - b^2)}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{5a-3b \sin(c+dx)}{(a+b \sin(c+dx))^{5/2}} dx}{5(a^2-b^2)} + \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 3042 \\
& \frac{\int \frac{5a-3b \sin(c+dx)}{(a+b \sin(c+dx))^{5/2}} dx}{5(a^2-b^2)} + \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 3233 \\
& \frac{\frac{16ab \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}} - \frac{2 \int -\frac{3(5a^2+3b^2)-8ab \sin(c+dx)}{2(a+b \sin(c+dx))^{3/2}} dx}{3(a^2-b^2)}}{5(a^2-b^2)} + \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 27 \\
& \frac{\frac{\int \frac{3(5a^2+3b^2)-8ab \sin(c+dx)}{(a+b \sin(c+dx))^{3/2}} dx}{3(a^2-b^2)} + \frac{16ab \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}}}{5(a^2-b^2)} + \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 3042 \\
& \frac{\frac{\int \frac{3(5a^2+3b^2)-8ab \sin(c+dx)}{(a+b \sin(c+dx))^{3/2}} dx}{3(a^2-b^2)} + \frac{16ab \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}}}{5(a^2-b^2)} + \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 3233 \\
& \frac{\frac{\frac{2b(23a^2+9b^2) \cos(c+dx)}{d(a^2-b^2) \sqrt{a+b \sin(c+dx)}} - \frac{2 \int -\frac{a(15a^2+17b^2)+b(23a^2+9b^2) \sin(c+dx)}{2\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2}}{3(a^2-b^2)} + \frac{16ab \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}}}{5(a^2-b^2)} + \\
& \quad \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 27 \\
& \frac{\frac{\frac{\int \frac{a(15a^2+17b^2)+b(23a^2+9b^2) \sin(c+dx)}{\sqrt{a+b \sin(c+dx)}} dx}{a^2-b^2} + \frac{2b(23a^2+9b^2) \cos(c+dx)}{d(a^2-b^2) \sqrt{a+b \sin(c+dx)}}}{3(a^2-b^2)} + \frac{16ab \cos(c+dx)}{3d(a^2-b^2)(a+b \sin(c+dx))^{3/2}}}{5(a^2-b^2)} + \\
& \quad \frac{2b \cos(c+dx)}{5d(a^2-b^2)(a+b \sin(c+dx))^{5/2}} \\
& \downarrow 3042
\end{aligned}$$

$$\frac{\int \frac{a(15a^2+17b^2)+b(23a^2+9b^2)\sin(c+dx)}{\sqrt{a+b\sin(c+dx)} a^2-b^2} dx + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}}{3(a^2-b^2)} + \frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}} + \frac{2b\cos(c+dx)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3231

$$\frac{\frac{(23a^2+9b^2)\int\sqrt{a+b\sin(c+dx)}dx-8a(a^2-b^2)\int\frac{1}{\sqrt{a+b\sin(c+dx)}}dx}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}}{3(a^2-b^2)} + \frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}} + \frac{2b\cos(c+dx)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3042

$$\frac{\frac{(23a^2+9b^2)\int\sqrt{a+b\sin(c+dx)}dx-8a(a^2-b^2)\int\frac{1}{\sqrt{a+b\sin(c+dx)}}dx}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}}{3(a^2-b^2)} + \frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}} + \frac{2b\cos(c+dx)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3134

$$\frac{\frac{\frac{(23a^2+9b^2)\sqrt{a+b\sin(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}dx}{\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - 8a(a^2-b^2)\int\frac{1}{\sqrt{a+b\sin(c+dx)}}dx}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}}{3(a^2-b^2)} + \frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}} + \frac{2b\cos(c+dx)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3042

$$\frac{\frac{\frac{(23a^2+9b^2)\sqrt{a+b\sin(c+dx)}\int\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}dx}{\sqrt{\frac{a+b\sin(c+dx)}{a+b}}} - 8a(a^2-b^2)\int\frac{1}{\sqrt{a+b\sin(c+dx)}}dx}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}}{3(a^2-b^2)} + \frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}} + \frac{2b\cos(c+dx)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3132

$$\frac{\frac{2(23a^2+9b^2)\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}\left(c+dx-\frac{\pi}{2}\right)\middle|\frac{2b}{a+b}\right)-8a(a^2-b^2)\int\frac{1}{\sqrt{a+b\sin(c+dx)}}dx}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}$$

$$\frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3142

$$\frac{\frac{2(23a^2+9b^2)\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}\left(c+dx-\frac{\pi}{2}\right)\middle|\frac{2b}{a+b}\right)-8a(a^2-b^2)\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}}dx}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}$$

$$\frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3042

$$\frac{\frac{2(23a^2+9b^2)\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}\left(c+dx-\frac{\pi}{2}\right)\middle|\frac{2b}{a+b}\right)-8a(a^2-b^2)\int\frac{1}{\sqrt{\frac{a}{a+b}+\frac{b\sin(c+dx)}{a+b}}}dx}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}}{a^2-b^2} + \frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16ab}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}}$$

$$\frac{5(a^2-b^2)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}}$$

↓ 3140

$$\frac{2b\cos(c+dx)}{5d(a^2-b^2)(a+b\sin(c+dx))^{5/2}} + \frac{2(23a^2+9b^2)\sqrt{a+b\sin(c+dx)}E\left(\frac{1}{2}\left(c+dx-\frac{\pi}{2}\right)\middle|\frac{2b}{a+b}\right)-16a(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}}}{d\sqrt{\frac{a+b\sin(c+dx)}{a+b}}}$$

$$\frac{16ab\cos(c+dx)}{3d(a^2-b^2)(a+b\sin(c+dx))^{3/2}} + \frac{\frac{2b(23a^2+9b^2)\cos(c+dx)}{d(a^2-b^2)\sqrt{a+b\sin(c+dx)}} + \frac{16a(a^2-b^2)\sqrt{\frac{a+b\sin(c+dx)}{a+b}}}{d\sqrt{a+b\sin(c+dx)}}}{3(a^2-b^2)}$$

$$5(a^2-b^2)$$

input `Int[(a + b*Sin[c + d*x])^(-7/2),x]`

output

$$\begin{aligned} & (2*b*\cos[c + d*x]) / (5*(a^2 - b^2)*d*(a + b*\sin[c + d*x])^{5/2}) + ((16*a*b \\ & * \cos[c + d*x]) / (3*(a^2 - b^2)*d*(a + b*\sin[c + d*x])^{3/2}) + ((2*b*(23*a^2 \\ & + 9*b^2)*\cos[c + d*x]) / ((a^2 - b^2)*d*\sqrt{a + b*\sin[c + d*x]}) + ((2*(2 \\ & 3*a^2 + 9*b^2)*\text{EllipticE}[(c - \pi/2 + d*x)/2, (2*b)/(a + b)]*\sqrt{a + b*\sin \\ & [c + d*x]}) / (d*\sqrt{(a + b*\sin[c + d*x])/(a + b)})) - (16*a*(a^2 - b^2)*\text{EllipticF} \\ & [(c - \pi/2 + d*x)/2, (2*b)/(a + b)]*\sqrt{(a + b*\sin[c + d*x])/(a + b)} \\ &)) / (d*\sqrt{a + b*\sin[c + d*x]}) / (a^2 - b^2) / (3*(a^2 - b^2)) / (5*(a^2 - \\ & b^2)) \end{aligned}$$

Definitions of rubi rules used

rule 27

$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3132

$$\text{Int}[\sqrt{(a_)} + (b_)*\sin[(c_.) + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[2*(\sqrt{a + b}/d)*\text{EllipticE}[(1/2)*(c - \pi/2 + d*x), 2*(b/(a + b))], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[a + b, 0]$$

rule 3134

$$\text{Int}[\sqrt{(a_)} + (b_)*\sin[(c_.) + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[\sqrt{a + b*\sin[c + d*x]} / \sqrt{(a + b*\sin[c + d*x])/(a + b)} \text{ Int}[\sqrt{a/(a + b)} + (b/(a + b))*\sin[c + d*x], x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ !\text{GtQ}[a + b, 0]$$

rule 3140

$$\text{Int}[1/\sqrt{(a_)} + (b_)*\sin[(c_.) + (d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[(2/(d*\sqrt{a + b}))*\text{EllipticF}[(1/2)*(c - \pi/2 + d*x), 2*(b/(a + b))], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0] \ \&\& \ \text{GtQ}[a + b, 0]$$

rule 3142 $\text{Int}[1/\text{Sqrt}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]] , x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[(a + b*\sin[c + d*x])/(a + b)]/\text{Sqrt}[a + b*\sin[c + d*x]] \text{Int}[1/\text{Sqrt}[a/(a + b) + (b/(a + b))*\sin[c + d*x]], x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && !GtQ[a + b, 0]

rule 3143 $\text{Int}[(a_) + (b_)*\sin[(c_) + (d_)*(x_)]^{(n_)} , x_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((a + b*\sin[c + d*x])^{(n + 1)})/(d*(n + 1)*(a^2 - b^2)), x] + \text{Simp}[1/((n + 1)*(a^2 - b^2)) \text{Int}[(a + b*\sin[c + d*x])^{(n + 1)}*\text{Simp}[a*(n + 1) - b*(n + 2)*\sin[c + d*x], x], x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]

rule 3231 $\text{Int}[(c_) + (d_)*\sin[(e_) + (f_)*(x_)]/\text{Sqrt}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]] , x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)/b \text{Int}[1/\text{Sqrt}[a + b*\sin[e + f*x]], x], x] + \text{Simp}[d/b \text{Int}[\text{Sqrt}[a + b*\sin[e + f*x]], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0]

rule 3233 $\text{Int}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]^{(m_)}*((c_) + (d_)*\sin[(e_) + (f_)*(x_)]] , x_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*\cos[e + f*x]*((a + b*\sin[e + f*x])^{(m + 1)})/(f*(m + 1)*(a^2 - b^2)), x] + \text{Simp}[1/((m + 1)*(a^2 - b^2)) \text{Int}[(a + b*\sin[e + f*x])^{(m + 1)}*\text{Simp}[(a*c - b*d)*(m + 1) - (b*c - a*d)*(m + 2)*\sin[e + f*x], x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && LtQ[m, -1] && IntegerQ[2*m]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 581 vs. $2(273) = 546$.

Time = 0.31 (sec) , antiderivative size = 582, normalized size of antiderivative = 1.99

method	result
default	$\frac{\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2} \left(\frac{2\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2}}{5(a^2-b^2)b^2(\sin(dx+c)+\frac{c}{b})^3} + \frac{16a\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2}}{15(a^2-b^2)^2b(\sin(dx+c)+\frac{c}{b})^2} + \frac{2b\cos(dx+c)^2}{15(a^2-b^2)^3\sqrt{-(-a-b\sin(dx+c))\cos(dx+c)^2}} \right)}{1}$

input $\text{int}(1/(a+b*\sin(d*x+c))^{(7/2)},x,\text{method}=_RETURNVERBOSE)$

output

```
(-(-a-b*sin(d*x+c))*cos(d*x+c)^2)^(1/2)*(2/5/(a^2-b^2)/b^2*(-(-a-b*sin(d*x+c))*cos(d*x+c)^2)^(1/2)/(sin(d*x+c)+a/b)^3+16/15*a/(a^2-b^2)^2/b*(-(-a-b*sin(d*x+c))*cos(d*x+c)^2)^(1/2)/(sin(d*x+c)+a/b)^2+2/15*b*cos(d*x+c)^2/(a^2-b^2)^3*(23*a^2+9*b^2)/(-(-a-b*sin(d*x+c))*cos(d*x+c)^2)^(1/2)+2*(15*a^3+17*a*b^2)/(15*a^6-45*a^4*b^2+45*a^2*b^4-15*b^6)*(a/b-1)*((a+b*sin(d*x+c))/(a-b))^(1/2)*(b*(1-sin(d*x+c))/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)/(-(-a-b*sin(d*x+c))*cos(d*x+c)^2)^(1/2)*EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))+2/15*b*(23*a^2+9*b^2)/(a^2-b^2)^3*(a/b-1)*((a+b*sin(d*x+c))/(a-b))^(1/2)*(b*(1-sin(d*x+c))/(a+b))^(1/2)*(-b*(1+sin(d*x+c))/(a-b))^(1/2)/(-(-a-b*sin(d*x+c))*cos(d*x+c)^2)^(1/2)*((-a/b-1)*EllipticE(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2))+EllipticF(((a+b*sin(d*x+c))/(a-b))^(1/2),((a-b)/(a+b))^(1/2)))/cos(d*x+c)/(a+b*sin(d*x+c))^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 985, normalized size of antiderivative = 3.37

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \text{Too large to display}$$

input

```
integrate(1/(a+b*sin(d*x+c))^(7/2),x, algorithm="fricas")
```

output

```

2/45*((a^6 - 30*a^4*b^2 - 99*a^2*b^4 - 3*(a^4*b^2 - 33*a^2*b^4)*cos(d*x +
c)^2 + (3*a^5*b - 98*a^3*b^3 - 33*a*b^5 - (a^3*b^3 - 33*a*b^5)*cos(d*x + c
)^2)*sin(d*x + c))*sqrt(1/2*I*b)*weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/
b^2, -8/27*(8*I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x
+ c) - 2*I*a)/b) + (a^6 - 30*a^4*b^2 - 99*a^2*b^4 - 3*(a^4*b^2 - 33*a^2*
b^4)*cos(d*x + c)^2 + (3*a^5*b - 98*a^3*b^3 - 33*a*b^5 - (a^3*b^3 - 33*a*b
^5)*cos(d*x + c)^2)*sin(d*x + c))*sqrt(-1/2*I*b)*weierstrassPInverse(-4/3*
(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x +
c) + 3*I*b*sin(d*x + c) + 2*I*a)/b) + 3*(23*I*a^5*b + 78*I*a^3*b^3 + 27*I*
a*b^5 + 3*(-23*I*a^3*b^3 - 9*I*a*b^5)*cos(d*x + c)^2 + (69*I*a^4*b^2 + 50*
I*a^2*b^4 + 9*I*b^6 + (-23*I*a^2*b^4 - 9*I*b^6)*cos(d*x + c)^2)*sin(d*x +
c))*sqrt(1/2*I*b)*weierstrassZeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*I*a^3
- 9*I*a*b^2)/b^3, weierstrassPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(8*
I*a^3 - 9*I*a*b^2)/b^3, 1/3*(3*b*cos(d*x + c) - 3*I*b*sin(d*x + c) - 2*I*a
)/b)) + 3*(-23*I*a^5*b - 78*I*a^3*b^3 - 27*I*a*b^5 + 3*(23*I*a^3*b^3 + 9*I
*a*b^5)*cos(d*x + c)^2 + (-69*I*a^4*b^2 - 50*I*a^2*b^4 - 9*I*b^6 + (23*I*a
^2*b^4 + 9*I*b^6)*cos(d*x + c)^2)*sin(d*x + c))*sqrt(-1/2*I*b)*weierstrass
Zeta(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, weierstra
ssPInverse(-4/3*(4*a^2 - 3*b^2)/b^2, -8/27*(-8*I*a^3 + 9*I*a*b^2)/b^3, 1/3
*(3*b*cos(d*x + c) + 3*I*b*sin(d*x + c) + 2*I*a)/b)) + 3*((23*a^2*b^4 + ...

```

Sympy [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx$$

input

```
integrate(1/(a+b*sin(d*x+c))**(7/2),x)
```

output

```
Integral((a + b*sin(c + d*x))**(-7/2), x)
```

Maxima [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{7/2}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(7/2),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(-7/2), x)`

Giac [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{7/2}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(7/2),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(-7/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx$$

input `int(1/(a + b*sin(c + d*x))^(7/2),x)`

output `int(1/(a + b*sin(c + d*x))^(7/2), x)`

Reduce [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{7/2}} dx = \int \frac{\sqrt{\sin(dx + c)b + a}}{\sin(dx + c)^4 b^4 + 4 \sin(dx + c)^3 a b^3 + 6 \sin(dx + c)^2 a^2 b^2 + 4 \sin(dx + c) a^3 b + a^4} dx$$

input `int(1/(a+b*sin(d*x+c))^(7/2),x)`

output `int(sqrt(sin(c + d*x)*b + a)/(sin(c + d*x)**4*b**4 + 4*sin(c + d*x)**3*a*b**3 + 6*sin(c + d*x)**2*a**2*b**2 + 4*sin(c + d*x)*a**3*b + a**4),x)`

3.84 $\int (a + b \sin(c + dx))^{4/3} dx$

Optimal result	623
Mathematica [B] (warning: unable to verify)	623
Rubi [A] (verified)	624
Maple [F]	626
Fricas [F]	626
Sympy [F]	626
Maxima [F]	627
Giac [F]	627
Mupad [F(-1)]	627
Reduce [F]	628

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int (a + b \sin(c + dx))^{4/3} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a+b}\right) \cos(c + dx)(a + b \sin(c + dx))^{4/3}}{d \sqrt{1 + \sin(c + dx)} \left(\frac{a+b \sin(c+dx)}{a+b}\right)^{4/3}}$$

output

```
-2^(1/2)*AppellF1(1/2,-4/3,1/2,3/2,b*(1-sin(d*x+c))/(a+b),1/2-1/2*sin(d*x+c))*cos(d*x+c)*(a+b*sin(d*x+c))^(4/3)/d/(1+sin(d*x+c))^(1/2)/((a+b*sin(d*x+c))/(a+b))^(4/3)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 244 vs. 2(106) = 212.

Time = 1.88 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.30

$$\int (a + b \sin(c + dx))^{4/3} dx = 3 \sec(c + dx) \sqrt[3]{a + b \sin(c + dx)} \left(4b^2 \cos^2(c + dx) + 4(a^2 - b^2) \operatorname{AppellF1}\left(\frac{1}{3}, \frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{a+b \sin(c+dx)}{a-b}, \frac{a+b \sin(c+dx)}{a+b}\right) \right)$$

input `Integrate[(a + b*Sin[c + d*x])^(4/3),x]`

output `(-3*Sec[c + d*x]*(a + b*Sin[c + d*x])^(1/3)*(4*b^2*Cos[c + d*x]^2 + 4*(a^2 - b^2)*AppellF1[1/3, 1/2, 1/2, 4/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)] - 5*a*AppellF1[4/3, 1/2, 1/2, 7/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x]))/(16*b*d)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \sin(c + dx))^{4/3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + b \sin(c + dx))^{4/3} dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(a + b \sin(c + dx))^{4/3}}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{(a + b) \cos(c + dx) \sqrt[3]{a + b \sin(c + dx)} \int \frac{\left(\frac{a}{a+b} + \frac{b \sin(c + dx)}{a+b}\right)^{4/3}}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}}} \\
 & \quad \downarrow \text{155}
 \end{aligned}$$

$$\frac{\sqrt{2}(a+b)\cos(c+dx)\sqrt[3]{a+b\sin(c+dx)}\operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1-\sin(c+dx)), \frac{b(1-\sin(c+dx))}{a+b}\right)}{d\sqrt{\sin(c+dx)+1}\sqrt[3]{\frac{a+b\sin(c+dx)}{a+b}}}$$

input `Int[(a + b*Sin[c + d*x])^(4/3),x]`

output `-((Sqrt[2]*(a + b)*AppellF1[1/2, 1/2, -4/3, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin[c + d*x]))/(a + b)]*Cos[c + d*x]*(a + b*Sin[c + d*x])^(1/3))/(d*Sqrt[1 + Sin[c + d*x]]*((a + b*Sin[c + d*x])/(a + b))^(1/3))`

Definitions of rubi rules used

rule 155 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p])*((b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (a + b \sin(dx + c))^{\frac{4}{3}} dx$$

input

```
int((a+b*sin(d*x+c))^(4/3),x)
```

output

```
int((a+b*sin(d*x+c))^(4/3),x)
```

Fricas [F]

$$\int (a + b \sin(c + dx))^{4/3} dx = \int (b \sin(dx + c) + a)^{\frac{4}{3}} dx$$

input

```
integrate((a+b*sin(d*x+c))^(4/3),x, algorithm="fricas")
```

output

```
integral((b*sin(d*x + c) + a)^(4/3), x)
```

Sympy [F]

$$\int (a + b \sin(c + dx))^{4/3} dx = \int (a + b \sin(c + dx))^{\frac{4}{3}} dx$$

input

```
integrate((a+b*sin(d*x+c))**(4/3),x)
```

output

```
Integral((a + b*sin(c + d*x))**(4/3), x)
```

Maxima [F]

$$\int (a + b \sin(c + dx))^{4/3} dx = \int (b \sin(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(4/3), x)`

Giac [F]

$$\int (a + b \sin(c + dx))^{4/3} dx = \int (b \sin(dx + c) + a)^{\frac{4}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(4/3), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^{4/3} dx = \int (a + b \sin(c + dx))^{4/3} dx$$

input `int((a + b*sin(c + d*x))^(4/3),x)`

output `int((a + b*sin(c + d*x))^(4/3), x)`

Reduce [F]

$$\int (a + b \sin(c + dx))^{4/3} dx = \left(\int (\sin(dx + c)b + a)^{\frac{1}{3}} dx \right) a + \left(\int (\sin(dx + c)b + a)^{\frac{1}{3}} \sin(dx + c) dx \right) b$$

input `int((a+b*sin(d*x+c))^(4/3),x)`

output `int((sin(c + d*x)*b + a)**(1/3),x)*a + int((sin(c + d*x)*b + a)**(1/3)*sin(c + d*x),x)*b`

3.85 $\int (a + b \sin(c + dx))^{2/3} dx$

Optimal result	629
Mathematica [A] (warning: unable to verify)	629
Rubi [A] (verified)	630
Maple [F]	632
Fricas [F]	632
Sympy [F]	632
Maxima [F]	633
Giac [F]	633
Mupad [F(-1)]	633
Reduce [F]	634

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int (a + b \sin(c + dx))^{2/3} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a+b}\right) \cos(c + dx)(a + b \sin(c + dx))^{2/3}}{d \sqrt{1 + \sin(c + dx)} \left(\frac{a+b \sin(c+dx)}{a+b}\right)^{2/3}}$$

output

```
-2^(1/2)*AppellF1(1/2,-2/3,1/2,3/2,b*(1-sin(d*x+c))/(a+b),1/2-1/2*sin(d*x+c))*cos(d*x+c)*(a+b*sin(d*x+c))^(2/3)/d/(1+sin(d*x+c))^(1/2)/((a+b*sin(d*x+c))/(a+b))^(2/3)
```

Mathematica [A] (warning: unable to verify)

Time = 0.32 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.11

$$\int (a + b \sin(c + dx))^{2/3} dx = \frac{3 \operatorname{AppellF1}\left(\frac{5}{3}, \frac{1}{2}, \frac{1}{2}, \frac{8}{3}, \frac{a+b \sin(c+dx)}{a-b}, \frac{a+b \sin(c+dx)}{a+b}\right) \sec(c + dx) \sqrt{-\frac{b(-1+\sin(c+dx))}{a+b}} \sqrt{\frac{b(1+\sin(c+dx))}{-a+b}}}{5bd}$$

input

```
Integrate[(a + b*Sin[c + d*x])^(2/3), x]
```

output

```
(3*AppellF1[5/3, 1/2, 1/2, 8/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sec[c + d*x]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x])^(5/3))/(5*b*d)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(c + dx))^{2/3} dx$$

$$\downarrow 3042$$

$$\int (a + b \sin(c + dx))^{2/3} dx$$

$$\downarrow 3144$$

$$\frac{\cos(c + dx) \int \frac{(a + b \sin(c + dx))^{2/3}}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}$$

$$\downarrow 156$$

$$\frac{\cos(c + dx)(a + b \sin(c + dx))^{2/3} \int \frac{\left(\frac{a}{a+b} + \frac{b \sin(c + dx)}{a+b}\right)^{2/3}}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \left(\frac{a + b \sin(c + dx)}{a+b}\right)^{2/3}}$$

$$\downarrow 155$$

$$\frac{\sqrt{2} \cos(c + dx)(a + b \sin(c + dx))^{2/3} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a+b}\right)}{d \sqrt{\sin(c + dx) + 1} \left(\frac{a + b \sin(c + dx)}{a+b}\right)^{2/3}}$$

input

```
Int[(a + b*Sin[c + d*x])^(2/3), x]
```

output

```

-((Sqrt[2]*AppellF1[1/2, 1/2, -2/3, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin
[c + d*x]))/(a + b)]*Cos[c + d*x]*(a + b*Sqrt[1 + Sin[c + d*x]]^(2/3))/(d*Sqrt[1 +
Sin[c + d*x]]*((a + b*Sqrt[1 + Sin[c + d*x]]/(a + b))^(2/3)))

```

Defintions of rubi rules used

rule 155

```

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])

```

rule 156

```

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]

```

rule 3042

```

Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 3144

```

Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]

```


Maple [F]

$$\int (a + b \sin(dx + c))^{\frac{2}{3}} dx$$

input `int((a+b*sin(d*x+c))^(2/3),x)`

output `int((a+b*sin(d*x+c))^(2/3),x)`

Fricas [F]

$$\int (a + b \sin(c + dx))^{\frac{2}{3}} dx = \int (b \sin(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((b*sin(d*x + c) + a)^(2/3), x)`

Sympy [F]

$$\int (a + b \sin(c + dx))^{\frac{2}{3}} dx = \int (a + b \sin(c + dx))^{\frac{2}{3}} dx$$

input `integrate((a+b*sin(d*x+c))**(2/3),x)`

output `Integral((a + b*sin(c + d*x))**(2/3), x)`

Maxima [F]

$$\int (a + b \sin(c + dx))^{2/3} dx = \int (b \sin(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(2/3), x)`

Giac [F]

$$\int (a + b \sin(c + dx))^{2/3} dx = \int (b \sin(dx + c) + a)^{\frac{2}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(2/3), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^{2/3} dx = \int (a + b \sin(c + dx))^{2/3} dx$$

input `int((a + b*sin(c + d*x))^(2/3),x)`

output `int((a + b*sin(c + d*x))^(2/3), x)`

Reduce [F]

$$\int (a + b \sin(c + dx))^{2/3} dx = \int (\sin(dx + c) b + a)^{2/3} dx$$

input `int((a+b*sin(d*x+c))^(2/3),x)`

output `int((sin(c + d*x)*b + a)**(2/3),x)`

3.86 $\int \sqrt[3]{a + b \sin(c + dx)} dx$

Optimal result	635
Mathematica [A] (warning: unable to verify)	635
Rubi [A] (verified)	636
Maple [F]	638
Fricas [F]	638
Sympy [F]	638
Maxima [F]	639
Giac [F]	639
Mupad [F(-1)]	639
Reduce [F]	640

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right) \cos(c + dx) \sqrt[3]{a + b \sin(c + dx)}}{d \sqrt{1 + \sin(c + dx)} \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}}}$$

output

```
-2^(1/2)*AppellF1(1/2,-1/3,1/2,3/2,b*(1-sin(d*x+c))/(a+b),1/2-1/2*sin(d*x+c))*cos(d*x+c)*(a+b*sin(d*x+c))^(1/3)/d/(1+sin(d*x+c))^(1/2)/((a+b*sin(d*x+c))/(a+b))^(1/3)
```

Mathematica [A] (warning: unable to verify)

Time = 0.31 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.11

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \frac{3 \operatorname{AppellF1}\left(\frac{4}{3}, \frac{1}{2}, \frac{1}{2}, \frac{7}{3}, \frac{a + b \sin(c + dx)}{a - b}, \frac{a + b \sin(c + dx)}{a + b}\right) \sec(c + dx) \sqrt{-\frac{b(-1 + \sin(c + dx))}{a + b}} \sqrt{\frac{b(1 + \sin(c + dx))}{-a + b}} (a + b \sin(c + dx))}{4bd}$$

input `Integrate[(a + b*Sin[c + d*x])^(1/3),x]`

output `(3*AppellF1[4/3, 1/2, 1/2, 7/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sec[c + d*x]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x])^(4/3))/(4*b*d)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{a + b \sin(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{a + b \sin(c + dx)} dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{\sqrt[3]{a + b \sin(c + dx)}}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\cos(c + dx) \sqrt[3]{a + b \sin(c + dx)} \int \frac{\sqrt[3]{\frac{a}{a + b} + \frac{b \sin(c + dx)}{a + b}}}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \cos(c + dx) \sqrt[3]{a + b \sin(c + dx)} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -\frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right)}{d \sqrt{\sin(c + dx) + 1} \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}}}
 \end{aligned}$$

input `Int[(a + b*SIN[c + d*x])^(1/3),x]`

output `-((Sqrt[2]*AppellF1[1/2, 1/2, -1/3, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin[c + d*x]))/(a + b)]*Cos[c + d*x]*(a + b*SIN[c + d*x])^(1/3))/(d*Sqrt[1 + Sin[c + d*x]]*((a + b*SIN[c + d*x])/(a + b))^(1/3)))`

Defintions of rubi rules used

rule 155 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]`

Maple [F]

$$\int (a + b \sin(dx + c))^{\frac{1}{3}} dx$$

input `int((a+b*sin(d*x+c))^(1/3),x)`

output `int((a+b*sin(d*x+c))^(1/3),x)`

Fricas [F]

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \int (b \sin(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*sin(d*x + c) + a)^(1/3), x)`

Sympy [F]

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \int \sqrt[3]{a + b \sin(c + dx)} dx$$

input `integrate((a+b*sin(d*x+c))**(1/3),x)`

output `Integral((a + b*sin(c + d*x))**(1/3), x)`

Maxima [F]

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \int (b \sin(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(1/3), x)`

Giac [F]

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \int (b \sin(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+b*sin(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(1/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \int (a + b \sin(c + dx))^{1/3} dx$$

input `int((a + b*sin(c + d*x))^(1/3),x)`

output `int((a + b*sin(c + d*x))^(1/3), x)`

Reduce [F]

$$\int \sqrt[3]{a + b \sin(c + dx)} dx = \int (\sin(dx + c)b + a)^{\frac{1}{3}} dx$$

input `int((a+b*sin(d*x+c))^(1/3),x)`

output `int((sin(c + d*x)*b + a)**(1/3),x)`

$$3.87 \quad \int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx$$

Optimal result	641
Mathematica [A] (warning: unable to verify)	641
Rubi [A] (verified)	642
Maple [F]	644
Fricas [F]	644
Sympy [F]	644
Maxima [F]	645
Giac [F]	645
Mupad [F(-1)]	645
Reduce [F]	646

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right) \cos(c + dx) \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}}}{d \sqrt{1 + \sin(c + dx)} \sqrt[3]{a + b \sin(c + dx)}}$$

output

```
-2^(1/2)*AppellF1(1/2,1/3,1/2,3/2,b*(1-sin(d*x+c))/(a+b),1/2-1/2*sin(d*x+c))
)*cos(d*x+c)*((a+b*sin(d*x+c))/(a+b))^(1/3)/d/(1+sin(d*x+c))^(1/2)/(a+b*
sin(d*x+c))^(1/3)
```

Mathematica [A] (warning: unable to verify)

Time = 0.30 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.11

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \frac{3 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, \frac{1}{2}, \frac{5}{3}, \frac{a + b \sin(c + dx)}{a - b}, \frac{a + b \sin(c + dx)}{a + b}\right) \sec(c + dx) \sqrt{-\frac{b(-1 + \sin(c + dx))}{a + b}} \sqrt{\frac{b(1 + \sin(c + dx))}{-a + b}} (a + b \sin(c + dx))}{2bd}$$

input `Integrate[(a + b*Sin[c + d*x])^(-1/3),x]`

output `(3*AppellF1[2/3, 1/2, 1/2, 5/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b])*Sec[c + d*x]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x])^(2/3))/(2*b*d)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{1}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \sqrt[3]{a + b \sin(c + dx)}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\cos(c + dx) \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}} \int \frac{1}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \sqrt[3]{\frac{a}{a + b} + \frac{b \sin(c + dx)}{a + b}}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \sqrt[3]{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \cos(c + dx) \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right)}{d \sqrt{\sin(c + dx) + 1} \sqrt[3]{a + b \sin(c + dx)}}
 \end{aligned}$$

input `Int[(a + b*SIN[c + d*x])^(-1/3),x]`

output `-((Sqrt[2]*AppellF1[1/2, 1/2, 1/3, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin[c + d*x]))/(a + b)]*Cos[c + d*x]*((a + b*SIN[c + d*x])/(a + b))^(1/3))/(d*Sqrt[1 + Sin[c + d*x]]*(a + b*SIN[c + d*x])^(1/3))`

Defintions of rubi rules used

rule 155 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]`

Maple [F]

$$\int \frac{1}{(a + b \sin(dx + c))^{\frac{1}{3}}} dx$$

input `int(1/(a+b*sin(d*x+c))^(1/3),x)`

output `int(1/(a+b*sin(d*x+c))^(1/3),x)`

Fricas [F]

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(1/3),x, algorithm="fricas")`

output `integral((b*sin(d*x + c) + a)^(-1/3), x)`

Sympy [F]

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx$$

input `integrate(1/(a+b*sin(d*x+c))**(1/3),x)`

output `Integral((a + b*sin(c + d*x))**(-1/3), x)`

Maxima [F]

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(-1/3), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(-1/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \int \frac{1}{(a + b \sin(c + dx))^{1/3}} dx$$

input `int(1/(a + b*sin(c + d*x))^(1/3),x)`

output `int(1/(a + b*sin(c + d*x))^(1/3), x)`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{a + b \sin(c + dx)}} dx = \int \frac{1}{(\sin(dx + c)b + a)^{\frac{1}{3}}} dx$$

input `int(1/(a+b*sin(d*x+c))^(1/3),x)`

output `int(1/(sin(c + d*x)*b + a)**(1/3),x)`

3.88 $\int \frac{1}{(a+b \sin(c+dx))^{2/3}} dx$

Optimal result	647
Mathematica [A] (warning: unable to verify)	647
Rubi [A] (verified)	648
Maple [F]	650
Fricas [F]	650
Sympy [F]	650
Maxima [F]	651
Giac [F]	651
Mupad [F(-1)]	651
Reduce [F]	652

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{2}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right) \cos(c + dx) \left(\frac{a + b \sin(c + dx)}{a + b}\right)^{2/3}}{d \sqrt{1 + \sin(c + dx)} (a + b \sin(c + dx))^{2/3}}$$

output

```
-2^(1/2)*AppellF1(1/2,2/3,1/2,3/2,b*(1-sin(d*x+c))/(a+b),1/2-1/2*sin(d*x+c))
*cos(d*x+c)*((a+b*sin(d*x+c))/(a+b))^(2/3)/d/(1+sin(d*x+c))^(1/2)/(a+b*
sin(d*x+c))^(2/3)
```

Mathematica [A] (warning: unable to verify)

Time = 0.26 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.09

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \frac{3 \operatorname{AppellF1}\left(\frac{1}{3}, \frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{a + b \sin(c + dx)}{a - b}, \frac{a + b \sin(c + dx)}{a + b}\right) \sec(c + dx) \sqrt{-\frac{b(-1 + \sin(c + dx))}{a + b}}}{bd}$$

input

```
Integrate[(a + b*Sin[c + d*x])^(-2/3),x]
```


output

```
(3*AppellF1[1/3, 1/2, 1/2, 4/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sec[c + d*x]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x])^(1/3))/(b*d)
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx$$

↓ 3042

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx$$

↓ 3144

$$\frac{\cos(c + dx) \int \frac{1}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} (a + b \sin(c + dx))^{2/3}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}$$

↓ 156

$$\frac{\cos(c + dx) \left(\frac{a + b \sin(c + dx)}{a + b} \right)^{2/3} \int \frac{1}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \left(\frac{a}{a + b} + \frac{b \sin(c + dx)}{a + b} \right)^{2/3}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} (a + b \sin(c + dx))^{2/3}}$$

↓ 155

$$\frac{\sqrt{2} \cos(c + dx) \left(\frac{a + b \sin(c + dx)}{a + b} \right)^{2/3} \text{AppellF1} \left(\frac{1}{2}, \frac{1}{2}, \frac{2}{3}, \frac{3}{2}, \frac{1}{2} (1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b} \right)}{d \sqrt{\sin(c + dx) + 1} (a + b \sin(c + dx))^{2/3}}$$

input

```
Int[(a + b*Sin[c + d*x])^(-2/3), x]
```

output

```

-((Sqrt[2]*AppellF1[1/2, 1/2, 2/3, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin[
c + d*x]))/(a + b)]*Cos[c + d*x]*((a + b*Sin[c + d*x])/(a + b))^(2/3))/(d*
Sqrt[1 + Sin[c + d*x]]*(a + b*Sin[c + d*x])^(2/3))

```

Defintions of rubi rules used

rule 155

```

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Sim
plify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])

```

rule 156

```

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]

```

rule 3042

```

Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 3144

```

Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]

```

Maple [F]

$$\int \frac{1}{(a + b \sin(dx + c))^{\frac{2}{3}}} dx$$

input `int(1/(a+b*sin(d*x+c))^(2/3),x)`

output `int(1/(a+b*sin(d*x+c))^(2/3),x)`

Fricas [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{2}{3}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(2/3),x, algorithm="fricas")`

output `integral((b*sin(d*x + c) + a)^(-2/3), x)`

Sympy [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a + b \sin(c + dx))^{\frac{2}{3}}} dx$$

input `integrate(1/(a+b*sin(d*x+c))**(2/3),x)`

output `Integral((a + b*sin(c + d*x))**(-2/3), x)`

Maxima [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(-2/3), x)`

Giac [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{2/3}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(-2/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx$$

input `int(1/(a + b*sin(c + d*x))^(2/3),x)`

output `int(1/(a + b*sin(c + d*x))^(2/3), x)`

Reduce [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{2/3}} dx = \int \frac{1}{(\sin(dx + c) b + a)^{2/3}} dx$$

input `int(1/(a+b*sin(d*x+c))^(2/3),x)`

output `int(1/(sin(c + d*x)*b + a)**(2/3),x)`

3.89 $\int \frac{1}{(a+b \sin(c+dx))^{4/3}} dx$

Optimal result	653
Mathematica [B] (warning: unable to verify)	653
Rubi [A] (verified)	654
Maple [F]	656
Fricas [F]	656
Sympy [F]	656
Maxima [F]	657
Giac [F]	657
Mupad [F(-1)]	657
Reduce [F]	658

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right) \cos(c + dx) \left(\frac{a + b \sin(c + dx)}{a + b}\right)^{4/3}}{d \sqrt{1 + \sin(c + dx)} (a + b \sin(c + dx))^{4/3}}$$

output

```
-2^(1/2)*AppellF1(1/2,4/3,1/2,3/2,b*(1-sin(d*x+c))/(a+b),1/2-1/2*sin(d*x+c))
)*cos(d*x+c)*((a+b*sin(d*x+c))/(a+b))^(4/3)/d/(1+sin(d*x+c))^(1/2)/(a+b*
sin(d*x+c))^(4/3)
```

Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 262 vs. 2(106) = 212.

Time = 2.10 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.47

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \frac{3 \sec(c + dx) \left(5a \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, \frac{1}{2}, \frac{5}{3}, \frac{a + b \sin(c + dx)}{a - b}, \frac{a + b \sin(c + dx)}{a + b}\right) \sqrt{-\frac{b(-1 + \sin(c + dx))}{a + b}} \sqrt{-\frac{b(1 + \sin(c + dx))}{a - b}} (a + b \right)}{d \sqrt{1 + \sin(c + dx)} (a + b \sin(c + dx))^{4/3}}$$

input `Integrate[(a + b*Sin[c + d*x])^(-4/3),x]`

output `(-3*Sec[c + d*x]*(5*a*AppellF1[2/3, 1/2, 1/2, 5/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[-((b*(1 + Sin[c + d*x]))/(a - b))]*(a + b*Sin[c + d*x]) - 2*(5*b^2*Cos[c + d*x]^2 + 2*AppellF1[5/3, 1/2, 1/2, 8/3, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin[c + d*x])/(a + b)]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x]^2)))/(10*b*(a^2 - b^2)*d*(a + b*Sin[c + d*x])^(1/3))`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{1}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} (a + b \sin(c + dx))^{4/3}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\cos(c + dx) \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}} \int \frac{1}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \left(\frac{a}{a + b} + \frac{b \sin(c + dx)}{a + b}\right)^{4/3}} d \sin(c + dx)}{d(a + b) \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1} \sqrt[3]{a + b \sin(c + dx)}} \\
 & \quad \downarrow \text{155}
 \end{aligned}$$

$$\frac{\sqrt{2} \cos(c + dx) \sqrt[3]{\frac{a + b \sin(c + dx)}{a + b}} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, \frac{4}{3}, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b}\right)}{d(a + b) \sqrt{\sin(c + dx) + 1} \sqrt[3]{a + b \sin(c + dx)}}$$

input `Int[(a + b*Sin[c + d*x])^(-4/3),x]`

output `-((Sqrt[2]*AppellF1[1/2, 1/2, 4/3, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin[c + d*x]))/(a + b)]*Cos[c + d*x]*((a + b*Sin[c + d*x])/(a + b))^(1/3))/((a + b)*d*Sqrt[1 + Sin[c + d*x]]*(a + b*Sin[c + d*x])^(1/3))`

Defintions of rubi rules used

rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p])*((b*(e + f*x)/(b*e - a*f))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int \frac{1}{(a + b \sin(dx + c))^{\frac{4}{3}}} dx$$

input

```
int(1/(a+b*sin(d*x+c))^(4/3),x)
```

output

```
int(1/(a+b*sin(d*x+c))^(4/3),x)
```

Fricas [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{\frac{4}{3}}} dx$$

input

```
integrate(1/(a+b*sin(d*x+c))^(4/3),x, algorithm="fricas")
```

output

```
integral(-(b*sin(d*x + c) + a)^(2/3)/(b^2*cos(d*x + c)^2 - 2*a*b*sin(d*x +
c) - a^2 - b^2), x)
```

Sympy [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a + b \sin(c + dx))^{\frac{4}{3}}} dx$$

input

```
integrate(1/(a+b*sin(d*x+c))**(4/3),x)
```

output `Integral((a + b*sin(c + d*x))**(-4/3), x)`

Maxima [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(4/3),x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^(-4/3), x)`

Giac [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \int \frac{1}{(b \sin(dx + c) + a)^{4/3}} dx$$

input `integrate(1/(a+b*sin(d*x+c))^(4/3),x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^(-4/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx$$

input `int(1/(a + b*sin(c + d*x))^(4/3),x)`

output `int(1/(a + b*sin(c + d*x))^(4/3), x)`

Reduce [F]

$$\int \frac{1}{(a + b \sin(c + dx))^{4/3}} dx = \int \frac{1}{(\sin(dx + c)b + a)^{\frac{1}{3}} \sin(dx + c)b + (\sin(dx + c)b + a)^{\frac{1}{3}} a} dx$$

input `int(1/(a+b*sin(d*x+c))^(4/3),x)`

output `int(1/((sin(c + d*x)*b + a)**(1/3)*sin(c + d*x)*b + (sin(c + d*x)*b + a)**(1/3)*a),x)`

3.90 $\int (a + b \sin(c + dx))^n dx$

Optimal result	659
Mathematica [A] (warning: unable to verify)	659
Rubi [A] (verified)	660
Maple [F]	662
Fricas [F]	662
Sympy [F]	662
Maxima [F]	663
Giac [F]	663
Mupad [F(-1)]	663
Reduce [F]	664

Optimal result

Integrand size = 12, antiderivative size = 104

$$\int (a + b \sin(c + dx))^n dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a+b}\right) \cos(c + dx)(a + b \sin(c + dx))^n \left(\frac{a+b \sin(c + dx)}{a+b}\right)}{d \sqrt{1 + \sin(c + dx)}}$$

output

```
-2^(1/2)*AppellF1(1/2, -n, 1/2, 3/2, b*(1-sin(d*x+c))/(a+b), 1/2-1/2*sin(d*x+c)
)*cos(d*x+c)*(a+b*sin(d*x+c))^n/d/(1+sin(d*x+c))^(1/2)/(((a+b*sin(d*x+c))/
(a+b))^n)
```

Mathematica [A] (warning: unable to verify)

Time = 0.36 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.15

$$\int (a + b \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, \frac{a+b \sin(c+dx)}{a-b}, \frac{a+b \sin(c+dx)}{a+b}\right) \sec(c + dx) \sqrt{-\frac{b(-1+\sin(c+dx))}{a+b}} \sqrt{\frac{b(1+\sin(c+dx))}{-a+b}} (a + b \sin(c + dx))^n}{bd(1 + n)}$$

input

```
Integrate[(a + b*Sin[c + d*x])^n, x]
```

output

```
(AppellF1[1 + n, 1/2, 1/2, 2 + n, (a + b*Sin[c + d*x])/(a - b), (a + b*Sin
[c + d*x])/(a + b)]*Sec[c + d*x]*Sqrt[-((b*(-1 + Sin[c + d*x]))/(a + b))]*
Sqrt[(b*(1 + Sin[c + d*x]))/(-a + b)]*(a + b*Sin[c + d*x])^(1 + n))/(b*d*(
1 + n))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \sin(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + b \sin(c + dx))^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(a + b \sin(c + dx))^n}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\cos(c + dx) (a + b \sin(c + dx))^n \left(\frac{a + b \sin(c + dx)}{a + b} \right)^{-n} \int \frac{\left(\frac{a}{a + b} + \frac{b \sin(c + dx)}{a + b} \right)^n}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \cos(c + dx) (a + b \sin(c + dx))^n \left(\frac{a + b \sin(c + dx)}{a + b} \right)^{-n} \text{AppellF1} \left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2} (1 - \sin(c + dx)), \frac{b(1 - \sin(c + dx))}{a + b} \right)}{d \sqrt{\sin(c + dx) + 1}}
 \end{aligned}$$

input

```
Int[(a + b*Sin[c + d*x])^n,x]
```

output

```

-((Sqrt[2]*AppellF1[1/2, 1/2, -n, 3/2, (1 - Sin[c + d*x])/2, (b*(1 - Sin[c
+ d*x]))/(a + b)]*Cos[c + d*x]*(a + b*Sin[c + d*x])^n)/(d*Sqrt[1 + Sin[c
+ d*x]]*((a + b*Sin[c + d*x])/(a + b))^n)

```

Defintions of rubi rules used

rule 155

```

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])

```

rule 156

```

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p
]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]

```

rule 3042

```

Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 3144

```

Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]

```

Maple [F]

$$\int (a + b \sin(dx + c))^n dx$$

input `int((a+b*sin(d*x+c))^n,x)`

output `int((a+b*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (a + b \sin(c + dx))^n dx = \int (b \sin(dx + c) + a)^n dx$$

input `integrate((a+b*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((b*sin(d*x + c) + a)^n, x)`

Sympy [F]

$$\int (a + b \sin(c + dx))^n dx = \int (a + b \sin(c + dx))^n dx$$

input `integrate((a+b*sin(d*x+c))**n,x)`

output `Integral((a + b*sin(c + d*x))**n, x)`

Maxima [F]

$$\int (a + b \sin(c + dx))^n dx = \int (b \sin(dx + c) + a)^n dx$$

input `integrate((a+b*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*sin(d*x + c) + a)^n, x)`

Giac [F]

$$\int (a + b \sin(c + dx))^n dx = \int (b \sin(dx + c) + a)^n dx$$

input `integrate((a+b*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*sin(d*x + c) + a)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \sin(c + dx))^n dx = \int (a + b \sin(c + dx))^n dx$$

input `int((a + b*sin(c + d*x))^n,x)`

output `int((a + b*sin(c + d*x))^n, x)`

Reduce [F]

$$\int (a + b \sin(c + dx))^n dx = \int (\sin(dx + c)b + a)^n dx$$

input `int((a+b*sin(d*x+c))^n,x)`

output `int((sin(c + d*x)*b + a)**n,x)`

3.91 $\int (3 + 4 \sin(c + dx))^n dx$

Optimal result	665
Mathematica [A] (verified)	665
Rubi [A] (verified)	666
Maple [F]	667
Fricas [F]	668
Sympy [F]	668
Maxima [F]	668
Giac [F]	669
Mupad [F(-1)]	669
Reduce [F]	669

Optimal result

Integrand size = 12, antiderivative size = 72

$$\int (3 + 4 \sin(c + dx))^n dx$$

$$= -\frac{\sqrt{27}^n \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{4}{7}(1 - \sin(c + dx))\right) \cos(c + dx)}{d\sqrt{1 + \sin(c + dx)}}$$

output `-2^(1/2)*7^n*AppellF1(1/2,1/2,-n,3/2,1/2-1/2*sin(d*x+c),4/7-4/7*sin(d*x+c))*cos(d*x+c)/d/(1+sin(d*x+c))^(1/2)`

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.15

$$\int (3 + 4 \sin(c + dx))^n dx$$

$$= \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, -3 - 4 \sin(c + dx), \frac{1}{7}(3 + 4 \sin(c + dx))\right) \sqrt{\cos^2(c + dx)} \sec(c + dx)(3 + 4 \sin(c + dx))}{\sqrt{7}d(1 + n)}$$

input `Integrate[(3 + 4*Sin[c + d*x])^n,x]`

output

```
(AppellF1[1 + n, 1/2, 1/2, 2 + n, -3 - 4*Sin[c + d*x], (3 + 4*Sin[c + d*x])
]/7]*Sqrt[Cos[c + d*x]^2]*Sec[c + d*x]*(3 + 4*Sin[c + d*x])^(1 + n))/(Sqrt
[7]*d*(1 + n))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3144, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (4 \sin(c + dx) + 3)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (4 \sin(c + dx) + 3)^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(4 \sin(c + dx) + 3)^n}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{27}^n \cos(c + dx) \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{4}{7}(1 - \sin(c + dx))\right)}{d \sqrt{\sin(c + dx) + 1}}
 \end{aligned}$$

input

```
Int[(3 + 4*Sin[c + d*x])^n,x]
```

output

```
-((Sqrt[2]*7^n*AppellF1[1/2, 1/2, -n, 3/2, (1 - Sin[c + d*x])/2, (4*(1 - S
in[c + d*x]))/7]*Cos[c + d*x])/(d*Sqrt[1 + Sin[c + d*x]]))
```

Definitions of rubi rules used

rule 155 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3144 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]`

Maple [F]

$$\int (3 + 4 \sin(dx + c))^n dx$$

input `int((3+4*sin(d*x+c))^n,x)`

output `int((3+4*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) + 3)^n dx$$

input `integrate((3+4*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((4*sin(d*x + c) + 3)^n, x)`

Sympy [F]

$$\int (3 + 4 \sin(c + dx))^n dx = \int (4 \sin(c + dx) + 3)^n dx$$

input `integrate((3+4*sin(d*x+c))**n,x)`

output `Integral((4*sin(c + d*x) + 3)**n, x)`

Maxima [F]

$$\int (3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) + 3)^n dx$$

input `integrate((3+4*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((4*sin(d*x + c) + 3)^n, x)`

Giac [F]

$$\int (3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) + 3)^n dx$$

input `integrate((3+4*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((4*sin(d*x + c) + 3)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (3 + 4 \sin(c + dx))^n dx = \int (4 \sin(c + dx) + 3)^n dx$$

input `int((4*sin(c + d*x) + 3)^n,x)`

output `int((4*sin(c + d*x) + 3)^n, x)`

Reduce [F]

$$\int (3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) + 3)^n dx$$

input `int((3+4*sin(d*x+c))^n,x)`

output `int((4*sin(c + d*x) + 3)**n,x)`

3.92 $\int (3 - 4 \sin(c + dx))^n dx$

Optimal result	670
Mathematica [A] (verified)	670
Rubi [A] (verified)	671
Maple [F]	672
Fricas [F]	673
Sympy [F]	673
Maxima [F]	673
Giac [F]	674
Mupad [F(-1)]	674
Reduce [F]	674

Optimal result

Integrand size = 12, antiderivative size = 69

$$\int (3 - 4 \sin(c + dx))^n dx = \frac{\sqrt{2} 7^n \operatorname{AppellF1}\left(\frac{1}{2}, -n, \frac{1}{2}, \frac{3}{2}, \frac{4}{7}(1 + \sin(c + dx)), \frac{1}{2}(1 + \sin(c + dx))\right) \cos(c + dx)}{d \sqrt{1 - \sin(c + dx)}}$$

output `2^(1/2)*7^n*AppellF1(1/2,1/2,-n,3/2,1/2+1/2*sin(d*x+c),4/7+4/7*sin(d*x+c))*cos(d*x+c)/d/(1-sin(d*x+c))^(1/2)`

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.22

$$\int (3 - 4 \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, \frac{1}{7}(3 - 4 \sin(c + dx)), -3 + 4 \sin(c + dx)\right) \sqrt{\cos^2(c + dx)} \sec(c + dx)(3 - \sqrt{7}d(1 + n))}{\sqrt{7}d(1 + n)}$$

input `Integrate[(3 - 4*Sin[c + d*x])^n,x]`

output

```

-((AppellF1[1 + n, 1/2, 1/2, 2 + n, (3 - 4*Sin[c + d*x])/7, -3 + 4*Sin[c +
d*x]]*Sqrt[Cos[c + d*x]^2]*Sec[c + d*x]*(3 - 4*Sin[c + d*x])^(1 + n))/(Sq
rt[7]*d*(1 + n))

```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3144, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (3 - 4 \sin(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (3 - 4 \sin(c + dx))^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(3 - 4 \sin(c + dx))^n d \sin(c + dx)}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2}^n \cos(c + dx) \operatorname{AppellF1}\left(\frac{1}{2}, -n, \frac{1}{2}, \frac{3}{2}, \frac{4}{7}(\sin(c + dx) + 1), \frac{1}{2}(\sin(c + dx) + 1)\right)}{d \sqrt{1 - \sin(c + dx)}}
 \end{aligned}$$

input

```

Int[(3 - 4*Sin[c + d*x])^n,x]

```

output

```

(Sqrt[2]*7^n*AppellF1[1/2, -n, 1/2, 3/2, (4*(1 + Sin[c + d*x]))/7, (1 + Si
n[c + d*x])/2]*Cos[c + d*x])/(d*Sqrt[1 - Sin[c + d*x]])

```


Definitions of rubi rules used

rule 155

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3144

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)
^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (3 - 4 \sin(dx + c))^n dx$$

input

```
int((3-4*sin(d*x+c))^n,x)
```

output

```
int((3-4*sin(d*x+c))^n,x)
```

Fricas [F]

$$\int (3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) + 3)^n dx$$

input `integrate((3-4*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((-4*sin(d*x + c) + 3)^n, x)`

Sympy [F]

$$\int (3 - 4 \sin(c + dx))^n dx = \int (3 - 4 \sin(c + dx))^n dx$$

input `integrate((3-4*sin(d*x+c))**n,x)`

output `Integral((3 - 4*sin(c + d*x))**n, x)`

Maxima [F]

$$\int (3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) + 3)^n dx$$

input `integrate((3-4*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-4*sin(d*x + c) + 3)^n, x)`

Giac [F]

$$\int (3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) + 3)^n dx$$

input `integrate((3-4*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((-4*sin(d*x + c) + 3)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (3 - 4 \sin(c + dx))^n dx = \int (3 - 4 \sin(c + dx))^n dx$$

input `int((3 - 4*sin(c + d*x))^n,x)`

output `int((3 - 4*sin(c + d*x))^n, x)`

Reduce [F]

$$\int (3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) + 3)^n dx$$

input `int((3-4*sin(d*x+c))^n,x)`

output `int((- 4*sin(c + d*x) + 3)**n,x)`

3.93 $\int (4 + 3 \sin(c + dx))^n dx$

Optimal result	675
Mathematica [A] (warning: unable to verify)	675
Rubi [A] (verified)	676
Maple [F]	677
Fricas [F]	678
Sympy [F]	678
Maxima [F]	678
Giac [F]	679
Mupad [F(-1)]	679
Reduce [F]	679

Optimal result

Integrand size = 12, antiderivative size = 72

$$\int (4 + 3 \sin(c + dx))^n dx = -\frac{\sqrt{27}^n \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{3}{7}(1 - \sin(c + dx))\right) \cos(c + dx)}{d\sqrt{1 + \sin(c + dx)}}$$

output `-2^(1/2)*7^n*AppellF1(1/2,1/2,-n,3/2,1/2-1/2*sin(d*x+c),3/7-3/7*sin(d*x+c))*cos(d*x+c)/d/(1+sin(d*x+c))^(1/2)`

Mathematica [A] (warning: unable to verify)

Time = 0.16 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.38

$$\int (4 + 3 \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, 4 + 3 \sin(c + dx), \frac{1}{7}(4 + 3 \sin(c + dx))\right) \sec(c + dx) \sqrt{-1 - \sin(c + dx)} \sqrt{1 + \sin(c + dx)}}{\sqrt{7}d(1 + n)}$$

input `Integrate[(4 + 3*Sin[c + d*x])^n,x]`

output

```
(AppellF1[1 + n, 1/2, 1/2, 2 + n, 4 + 3*Sin[c + d*x], (4 + 3*Sin[c + d*x])
/7]*Sec[c + d*x]*Sqrt[-1 - Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]*(4 + 3*Sin
[c + d*x])^(1 + n))/(Sqrt[7]*d*(1 + n))
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.89, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3144, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (3 \sin(c + dx) + 4)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (3 \sin(c + dx) + 4)^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(3 \sin(c + dx) + 4)^n d \sin(c + dx)}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \cos(c + dx) \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(\sin(c + dx) + 1), -3(\sin(c + dx) + 1)\right)}{d \sqrt{1 - \sin(c + dx)}}
 \end{aligned}$$

input

```
Int[(4 + 3*Sin[c + d*x])^n,x]
```

output

```
(Sqrt[2]*AppellF1[1/2, 1/2, -n, 3/2, (1 + Sin[c + d*x])/2, -3*(1 + Sin[c +
d*x])]*Cos[c + d*x])/(d*Sqrt[1 - Sin[c + d*x]])
```

Definitions of rubi rules used

rule 155 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144 `Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]`

Maple [F]

$$\int (4 + 3 \sin(dx + c))^n dx$$

input `int((4+3*sin(d*x+c))^n,x)`

output `int((4+3*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) + 4)^n dx$$

input `integrate((4+3*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((3*sin(d*x + c) + 4)^n, x)`

Sympy [F]

$$\int (4 + 3 \sin(c + dx))^n dx = \int (3 \sin(c + dx) + 4)^n dx$$

input `integrate((4+3*sin(d*x+c))**n,x)`

output `Integral((3*sin(c + d*x) + 4)**n, x)`

Maxima [F]

$$\int (4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) + 4)^n dx$$

input `integrate((4+3*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((3*sin(d*x + c) + 4)^n, x)`

Giac [F]

$$\int (4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) + 4)^n dx$$

input `integrate((4+3*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((3*sin(d*x + c) + 4)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (4 + 3 \sin(c + dx))^n dx = \int (3 \sin(c + dx) + 4)^n dx$$

input `int((3*sin(c + d*x) + 4)^n,x)`

output `int((3*sin(c + d*x) + 4)^n, x)`

Reduce [F]

$$\int (4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) + 4)^n dx$$

input `int((4+3*sin(d*x+c))^n,x)`

output `int((3*sin(c + d*x) + 4)**n,x)`

3.94 $\int (4 - 3 \sin(c + dx))^n dx$

Optimal result	680
Mathematica [A] (verified)	680
Rubi [A] (verified)	681
Maple [F]	682
Fricas [F]	683
Sympy [F]	683
Maxima [F]	683
Giac [F]	684
Mupad [F(-1)]	684
Reduce [F]	684

Optimal result

Integrand size = 12, antiderivative size = 67

$$\int (4 - 3 \sin(c + dx))^n dx = -\frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), -3(1 - \sin(c + dx))\right) \cos(c + dx)}{d\sqrt{1 + \sin(c + dx)}}$$

output `-2^(1/2)*AppellF1(1/2,-n,1/2,3/2,-3+3*sin(d*x+c),1/2-1/2*sin(d*x+c))*cos(d*x+c)/d/(1+sin(d*x+c))^(1/2)`

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.43

$$\int (4 - 3 \sin(c + dx))^n dx = -\frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, \frac{1}{7}(4 - 3 \sin(c + dx)), 4 - 3 \sin(c + dx)\right) \sec(c + dx)(4 - 3 \sin(c + dx))^{1+n}}{\sqrt{7}d(1 + n)}$$

input `Integrate[(4 - 3*Sin[c + d*x])^n,x]`

output

```
-((AppellF1[1 + n, 1/2, 1/2, 2 + n, (4 - 3*Sin[c + d*x])/7, 4 - 3*Sin[c +
d*x]]*Sec[c + d*x]*(4 - 3*Sin[c + d*x])^(1 + n)*Sqrt[-1 + Sin[c + d*x]]*Sq
rt[1 + Sin[c + d*x]])/(Sqrt[7]*d*(1 + n))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3144, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (4 - 3 \sin(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (4 - 3 \sin(c + dx))^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(4 - 3 \sin(c + dx))^n d \sin(c + dx)}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} 7^n \cos(c + dx) \operatorname{AppellF1}\left(\frac{1}{2}, -n, \frac{1}{2}, \frac{3}{2}, \frac{3}{7}(\sin(c + dx) + 1), \frac{1}{2}(\sin(c + dx) + 1)\right)}{d \sqrt{1 - \sin(c + dx)}}
 \end{aligned}$$

input

```
Int[(4 - 3*Sin[c + d*x])^n,x]
```

output

```
(Sqrt[2]*7^n*AppellF1[1/2, -n, 1/2, 3/2, (3*(1 + Sin[c + d*x]))/7, (1 + Si
n[c + d*x])/2]*Cos[c + d*x])/(d*Sqrt[1 - Sin[c + d*x]])
```

Definitions of rubi rules used

rule 155

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3144

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)
]^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (4 - 3 \sin(dx + c))^n dx$$

input

```
int((4-3*sin(d*x+c))^n,x)
```

output

```
int((4-3*sin(d*x+c))^n,x)
```

Fricas [F]

$$\int (4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) + 4)^n dx$$

input `integrate((4-3*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((-3*sin(d*x + c) + 4)^n, x)`

Sympy [F]

$$\int (4 - 3 \sin(c + dx))^n dx = \int (4 - 3 \sin(c + dx))^n dx$$

input `integrate((4-3*sin(d*x+c))**n,x)`

output `Integral((4 - 3*sin(c + d*x))**n, x)`

Maxima [F]

$$\int (4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) + 4)^n dx$$

input `integrate((4-3*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-3*sin(d*x + c) + 4)^n, x)`

Giac [F]

$$\int (4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) + 4)^n dx$$

input `integrate((4-3*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((-3*sin(d*x + c) + 4)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (4 - 3 \sin(c + dx))^n dx = \int (4 - 3 \sin(c + dx))^n dx$$

input `int((4 - 3*sin(c + d*x))^n,x)`

output `int((4 - 3*sin(c + d*x))^n, x)`

Reduce [F]

$$\int (4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) + 4)^n dx$$

input `int((4-3*sin(d*x+c))^n,x)`

output `int((- 3*sin(c + d*x) + 4)**n,x)`

3.95 $\int (-3 + 4 \sin(c + dx))^n dx$

Optimal result	685
Mathematica [A] (verified)	685
Rubi [A] (verified)	686
Maple [F]	687
Fricas [F]	688
Sympy [F]	688
Maxima [F]	688
Giac [F]	689
Mupad [F(-1)]	689
Reduce [F]	689

Optimal result

Integrand size = 12, antiderivative size = 67

$$\int (-3 + 4 \sin(c + dx))^n dx = -\frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), 4(1 - \sin(c + dx))\right) \cos(c + dx)}{d\sqrt{1 + \sin(c + dx)}}$$

output

$$-2^{(1/2)} * \operatorname{AppellF1}(1/2, -n, 1/2, 3/2, 4 - 4 * \sin(d * x + c), 1/2 - 1/2 * \sin(d * x + c)) * \cos(d * x + c) / d / (1 + \sin(d * x + c))^{(1/2)}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.24

$$\int (-3 + 4 \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, \frac{1}{7}(3 - 4 \sin(c + dx)), -3 + 4 \sin(c + dx)\right) \sqrt{\cos^2(c + dx)} \sec(c + dx)(-3 + 4 \sin(c + dx))}{\sqrt{7}d(1 + n)}$$

input

$$\operatorname{Integrate}[(-3 + 4 * \operatorname{Sin}[c + d * x])^n, x]$$

output

```
(AppellF1[1 + n, 1/2, 1/2, 2 + n, (3 - 4*Sin[c + d*x])/7, -3 + 4*Sin[c + d*x]]*Sqrt[Cos[c + d*x]^2]*Sec[c + d*x]*(-3 + 4*Sin[c + d*x])^(1 + n))/(Sqrt[7]*d*(1 + n))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3144, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (4 \sin(c + dx) - 3)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (4 \sin(c + dx) - 3)^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(4 \sin(c + dx) - 3)^n}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & -\frac{\sqrt{2} \cos(c + dx) \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), 4(1 - \sin(c + dx))\right)}{d \sqrt{\sin(c + dx) + 1}}
 \end{aligned}$$

input

```
Int[(-3 + 4*Sin[c + d*x])^n,x]
```

output

```
-((Sqrt[2]*AppellF1[1/2, 1/2, -n, 3/2, (1 - Sin[c + d*x])/2, 4*(1 - Sin[c + d*x])]*Cos[c + d*x])/(d*Sqrt[1 + Sin[c + d*x]]))
```

Definitions of rubi rules used

rule 155

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3144

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)
]^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (-3 + 4 \sin(dx + c))^n dx$$

input

```
int((-3+4*sin(d*x+c))^n,x)
```

output

```
int((-3+4*sin(d*x+c))^n,x)
```


Fricas [F]

$$\int (-3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) - 3)^n dx$$

input `integrate((-3+4*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((4*sin(d*x + c) - 3)^n, x)`

Sympy [F]

$$\int (-3 + 4 \sin(c + dx))^n dx = \int (4 \sin(c + dx) - 3)^n dx$$

input `integrate((-3+4*sin(d*x+c))**n,x)`

output `Integral((4*sin(c + d*x) - 3)**n, x)`

Maxima [F]

$$\int (-3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) - 3)^n dx$$

input `integrate((-3+4*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((4*sin(d*x + c) - 3)^n, x)`

Giac [F]

$$\int (-3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) - 3)^n dx$$

input `integrate((-3+4*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((4*sin(d*x + c) - 3)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (-3 + 4 \sin(c + dx))^n dx = \int (4 \sin(c + dx) - 3)^n dx$$

input `int((4*sin(c + d*x) - 3)^n,x)`

output `int((4*sin(c + d*x) - 3)^n, x)`

Reduce [F]

$$\int (-3 + 4 \sin(c + dx))^n dx = \int (4 \sin(dx + c) - 3)^n dx$$

input `int((-3+4*sin(d*x+c))^n,x)`

output `int((4*sin(c + d*x) - 3)**n,x)`

3.96 $\int(-3 - 4 \sin(c + dx))^n dx$

Optimal result	690
Mathematica [A] (verified)	690
Rubi [A] (verified)	691
Maple [F]	692
Fricas [F]	693
Sympy [F]	693
Maxima [F]	693
Giac [F]	694
Mupad [F(-1)]	694
Reduce [F]	694

Optimal result

Integrand size = 12, antiderivative size = 64

$$\int(-3 - 4 \sin(c + dx))^n dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, -n, \frac{1}{2}, \frac{3}{2}, 4(1 + \sin(c + dx)), \frac{1}{2}(1 + \sin(c + dx))\right) \cos(c + dx)}{d\sqrt{1 - \sin(c + dx)}}$$

output

```
2^(1/2)*AppellF1(1/2,-n,1/2,3/2,4+4*sin(d*x+c),1/2+1/2*sin(d*x+c))*cos(d*x+c)/d/(1-sin(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.31

$$\int(-3 - 4 \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, -3 - 4 \sin(c + dx), \frac{1}{7}(3 + 4 \sin(c + dx))\right) \sqrt{\cos^2(c + dx)} \sec(c + dx)(-3 - 4 \sin(c + dx))}{\sqrt{7}d(1 + n)}$$

input

```
Integrate[(-3 - 4*Sin[c + d*x])^n,x]
```

output

```

-((AppellF1[1 + n, 1/2, 1/2, 2 + n, -3 - 4*Sin[c + d*x], (3 + 4*Sin[c + d*
x])/7]*Sqrt[Cos[c + d*x]^2]*Sec[c + d*x]*(-3 - 4*Sin[c + d*x])^(1 + n))/(S
qrt[7]*d*(1 + n))

```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3144, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (-4 \sin(c + dx) - 3)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (-4 \sin(c + dx) - 3)^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(-4 \sin(c + dx) - 3)^n d \sin(c + dx)}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{2} \cos(c + dx) \text{AppellF1}\left(\frac{1}{2}, -n, \frac{1}{2}, \frac{3}{2}, 4(\sin(c + dx) + 1), \frac{1}{2}(\sin(c + dx) + 1)\right)}{d \sqrt{1 - \sin(c + dx)}}
 \end{aligned}$$

input

```

Int[(-3 - 4*Sin[c + d*x])^n,x]

```

output

```

(Sqrt[2]*AppellF1[1/2, -n, 1/2, 3/2, 4*(1 + Sin[c + d*x]), (1 + Sin[c + d*
x])/2]*Cos[c + d*x])/(d*Sqrt[1 - Sin[c + d*x]])

```

Definitions of rubi rules used

rule 155

```
Int[((a_) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simpl
ify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3144

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)
^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (-3 - 4 \sin(dx + c))^n dx$$

input

```
int((-3-4*sin(d*x+c))^n,x)
```

output

```
int((-3-4*sin(d*x+c))^n,x)
```

Fricas [F]

$$\int (-3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) - 3)^n dx$$

input `integrate((-3-4*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((-4*sin(d*x + c) - 3)^n, x)`

Sympy [F]

$$\int (-3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(c + dx) - 3)^n dx$$

input `integrate((-3-4*sin(d*x+c))**n,x)`

output `Integral((-4*sin(c + d*x) - 3)**n, x)`

Maxima [F]

$$\int (-3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) - 3)^n dx$$

input `integrate((-3-4*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-4*sin(d*x + c) - 3)^n, x)`

Giac [F]

$$\int (-3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) - 3)^n dx$$

input `integrate((-3-4*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((-4*sin(d*x + c) - 3)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (-3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(c + dx) - 3)^n dx$$

input `int((- 4*sin(c + d*x) - 3)^n,x)`

output `int((- 4*sin(c + d*x) - 3)^n, x)`

Reduce [F]

$$\int (-3 - 4 \sin(c + dx))^n dx = \int (-4 \sin(dx + c) - 3)^n dx$$

input `int((-3-4*sin(d*x+c))^n,x)`

output `int((- 4*sin(c + d*x) - 3)**n,x)`

3.97 $\int (-4 + 3 \sin(c + dx))^n dx$

Optimal result	695
Mathematica [A] (verified)	695
Rubi [A] (verified)	696
Maple [F]	698
Fricas [F]	698
Sympy [F]	698
Maxima [F]	699
Giac [F]	699
Mupad [F(-1)]	699
Reduce [F]	700

Optimal result

Integrand size = 12, antiderivative size = 93

$$\int (-4 + 3 \sin(c + dx))^n dx = \frac{\sqrt{2} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), -3(1 - \sin(c + dx))\right) \cos(c + dx)(4 - 3 \sin(c + dx))^{-n}}{d\sqrt{1 + \sin(c + dx)}}$$

output

$$-2^{(1/2)} * \operatorname{AppellF1}(1/2, -n, 1/2, 3/2, -3+3*\sin(d*x+c), 1/2-1/2*\sin(d*x+c)) * \cos(d*x+c) * (-4+3*\sin(d*x+c))^n / d / ((4-3*\sin(d*x+c))^n / (1+\sin(d*x+c))^{(1/2)})$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.02

$$\int (-4 + 3 \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, \frac{1}{7}(4 - 3 \sin(c + dx)), 4 - 3 \sin(c + dx)\right) \sec(c + dx) \sqrt{-1 + \sin(c + dx)} \sqrt{1 + \sin(c + dx)}}{\sqrt{7}d(1 + n)}$$

input

$$\operatorname{Integrate}[(-4 + 3*\operatorname{Sin}[c + d*x])^n, x]$$

output

```
(AppellF1[1 + n, 1/2, 1/2, 2 + n, (4 - 3*Sin[c + d*x])/7, 4 - 3*Sin[c + d*
x]]*Sec[c + d*x]*Sqrt[-1 + Sin[c + d*x]]*Sqrt[1 + Sin[c + d*x]]*(-4 + 3*Si
n[c + d*x])^(1 + n))/(Sqrt[7]*d*(1 + n))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (3 \sin(c + dx) - 4)^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (3 \sin(c + dx) - 4)^n dx \\
 & \quad \downarrow \text{3144} \\
 & \frac{\cos(c + dx) \int \frac{(3 \sin(c + dx) - 4)^n}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{156} \\
 & \frac{\cos(c + dx) (4 - 3 \sin(c + dx))^{-n} (3 \sin(c + dx) - 4)^n \int \frac{(4 - 3 \sin(c + dx))^n}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} d \sin(c + dx)}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
 & \quad \downarrow \text{155} \\
 & \frac{\sqrt{27}^n \cos(c + dx) (4 - 3 \sin(c + dx))^{-n} (3 \sin(c + dx) - 4)^n \text{AppellF1}\left(\frac{1}{2}, -n, \frac{1}{2}, \frac{3}{2}, \frac{3}{7}(\sin(c + dx) + 1), \frac{1}{2}(\sin(c + dx) + 1)\right)}{d \sqrt{1 - \sin(c + dx)}}
 \end{aligned}$$

input

```
Int[(-4 + 3*Sin[c + d*x])^n,x]
```

output

```
(Sqrt[2]*7^n*AppellF1[1/2, -n, 1/2, 3/2, (3*(1 + Sin[c + d*x]))/7, (1 + Sin[c + d*x])/2]*Cos[c + d*x]*(-4 + 3*Sin[c + d*x])^n)/(d*(4 - 3*Sin[c + d*x])^n*Sqrt[1 - Sin[c + d*x]])
```

Defintions of rubi rules used

rule 155

```
Int[((a_) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 156

```
Int[((a_) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3144

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (-4 + 3 \sin(dx + c))^n dx$$

input `int((-4+3*sin(d*x+c))^n,x)`

output `int((-4+3*sin(d*x+c))^n,x)`

Fricas [F]

$$\int (-4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) - 4)^n dx$$

input `integrate((-4+3*sin(d*x+c))^n,x, algorithm="fricas")`

output `integral((3*sin(d*x + c) - 4)^n, x)`

Sympy [F]

$$\int (-4 + 3 \sin(c + dx))^n dx = \int (3 \sin(c + dx) - 4)^n dx$$

input `integrate((-4+3*sin(d*x+c))**n,x)`

output `Integral((3*sin(c + d*x) - 4)**n, x)`

Maxima [F]

$$\int (-4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) - 4)^n dx$$

input `integrate((-4+3*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((3*sin(d*x + c) - 4)^n, x)`

Giac [F]

$$\int (-4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) - 4)^n dx$$

input `integrate((-4+3*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((3*sin(d*x + c) - 4)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (-4 + 3 \sin(c + dx))^n dx = \int (3 \sin(c + dx) - 4)^n dx$$

input `int((3*sin(c + d*x) - 4)^n,x)`

output `int((3*sin(c + d*x) - 4)^n, x)`

Reduce [F]

$$\int (-4 + 3 \sin(c + dx))^n dx = \int (3 \sin(dx + c) - 4)^n dx$$

input `int((-4+3*sin(d*x+c))^n,x)`

output `int((3*sin(c + d*x) - 4)**n,x)`

3.98 $\int (-4 - 3 \sin(c + dx))^n dx$

Optimal result	701
Mathematica [A] (warning: unable to verify)	701
Rubi [A] (warning: unable to verify)	702
Maple [F]	704
Fricas [F]	704
Sympy [F]	704
Maxima [F]	705
Giac [F]	705
Mupad [F(-1)]	705
Reduce [F]	706

Optimal result

Integrand size = 12, antiderivative size = 98

$$\int (-4 - 3 \sin(c + dx))^n dx = \frac{\sqrt{27^n} \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(c + dx)), \frac{3}{7}(1 - \sin(c + dx))\right) \cos(c + dx)(-4 - 3 \sin(c + dx))^n}{d \sqrt{1 + \sin(c + dx)}}$$

output

```
-2^(1/2)*7^n*AppellF1(1/2,1/2,-n,3/2,1/2-1/2*sin(d*x+c),3/7-3/7*sin(d*x+c)
)*cos(d*x+c)*(-4-3*sin(d*x+c))^n/d/(1+sin(d*x+c))^(1/2)/((4+3*sin(d*x+c))^
n)
```

Mathematica [A] (warning: unable to verify)

Time = 0.18 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.02

$$\int (-4 - 3 \sin(c + dx))^n dx = \frac{\operatorname{AppellF1}\left(1 + n, \frac{1}{2}, \frac{1}{2}, 2 + n, 4 + 3 \sin(c + dx), \frac{1}{7}(4 + 3 \sin(c + dx))\right) \sec(c + dx)(-4 - 3 \sin(c + dx))^{n+1}}{\sqrt{7}d(1 + n)}$$

input

```
Integrate[(-4 - 3*Sin[c + d*x])^n,x]
```

output

```

-((AppellF1[1 + n, 1/2, 1/2, 2 + n, 4 + 3*Sin[c + d*x], (4 + 3*Sin[c + d*x
])/7]*Sec[c + d*x]*(-4 - 3*Sin[c + d*x])^(1 + n)*Sqrt[-1 - Sin[c + d*x]]*S
qrt[1 - Sin[c + d*x]])/(Sqrt[7]*d*(1 + n))

```

Rubi [A] (warning: unable to verify)

Time = 0.26 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.12, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 3144, 156, 27, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (-3 \sin(c + dx) - 4)^n dx \\
& \quad \downarrow \text{3042} \\
& \int (-3 \sin(c + dx) - 4)^n dx \\
& \quad \downarrow \text{3144} \\
& \frac{\cos(c + dx) \int \frac{(-3 \sin(c + dx) - 4)^n d \sin(c + dx)}{\sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}}}{d \sqrt{1 - \sin(c + dx)} \sqrt{\sin(c + dx) + 1}} \\
& \quad \downarrow \text{156} \\
& \frac{\sqrt{3} \sqrt{-\sin(c + dx) - 1} \cos(c + dx) \int \frac{(-3 \sin(c + dx) - 4)^n d \sin(c + dx)}{\sqrt{3} \sqrt{-\sin(c + dx) - 1} \sqrt{1 - \sin(c + dx)}}}{d \sqrt{1 - \sin(c + dx)} (\sin(c + dx) + 1)} \\
& \quad \downarrow \text{27} \\
& \frac{\sqrt{-\sin(c + dx) - 1} \cos(c + dx) \int \frac{(-3 \sin(c + dx) - 4)^n d \sin(c + dx)}{\sqrt{-\sin(c + dx) - 1} \sqrt{1 - \sin(c + dx)}}}{d \sqrt{1 - \sin(c + dx)} (\sin(c + dx) + 1)} \\
& \quad \downarrow \text{155} \\
& \frac{\sqrt{-\sin(c + dx) - 1} \cos(c + dx) (-3 \sin(c + dx) - 4)^{n+1} \text{AppellF1}\left(n + 1, \frac{1}{2}, \frac{1}{2}, n + 2, 3 \sin(c + dx) + 4, \frac{1}{7} (3 \sin(c + dx) + 4)\right)}{\sqrt{7} d (n + 1) \sqrt{1 - \sin(c + dx)} (\sin(c + dx) + 1)}
\end{aligned}$$

input `Int[(-4 - 3*Sin[c + d*x])^n,x]`

output `-((AppellF1[1 + n, 1/2, 1/2, 2 + n, 4 + 3*Sin[c + d*x], (4 + 3*Sin[c + d*x])/7]*Cos[c + d*x]*(-4 - 3*Sin[c + d*x])^(1 + n)*Sqrt[-1 - Sin[c + d*x]])/(Sqrt[7]*d*(1 + n)*Sqrt[1 - Sin[c + d*x]]*(1 + Sin[c + d*x]))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 155 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplifierQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplifierQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c +
d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x
)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]
```

Maple [F]

$$\int (-4 - 3 \sin(dx + c))^n dx$$

input

```
int((-4-3*sin(d*x+c))^n,x)
```

output

```
int((-4-3*sin(d*x+c))^n,x)
```

Fricas [F]

$$\int (-4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) - 4)^n dx$$

input

```
integrate((-4-3*sin(d*x+c))^n,x, algorithm="fricas")
```

output

```
integral((-3*sin(d*x + c) - 4)^n, x)
```

Sympy [F]

$$\int (-4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(c + dx) - 4)^n dx$$

input

```
integrate((-4-3*sin(d*x+c))**n,x)
```

output

```
Integral((-3*sin(c + d*x) - 4)**n, x)
```

Maxima [F]

$$\int (-4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) - 4)^n dx$$

input `integrate((-4-3*sin(d*x+c))^n,x, algorithm="maxima")`

output `integrate((-3*sin(d*x + c) - 4)^n, x)`

Giac [F]

$$\int (-4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) - 4)^n dx$$

input `integrate((-4-3*sin(d*x+c))^n,x, algorithm="giac")`

output `integrate((-3*sin(d*x + c) - 4)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (-4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(c + dx) - 4)^n dx$$

input `int((- 3*sin(c + d*x) - 4)^n,x)`

output `int((- 3*sin(c + d*x) - 4)^n, x)`

Reduce [F]

$$\int (-4 - 3 \sin(c + dx))^n dx = \int (-3 \sin(dx + c) - 4)^n dx$$

input `int((-4-3*sin(d*x+c))^n,x)`

output `int((- 3*sin(c + d*x) - 4)**n,x)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	707
4.2	Links to plain text integration problems used in this report for each CAS .	725

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."}
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal."}
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order of result is higher than in optimal."}
  ]
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
        If [Head [expn] === RootSum,
            Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
            If [Head [expn] === Integrate || Head [expn] === Int,
                Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
                9]]]]]]]]]]]]

```

```

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```



```

#      antiderivative
#      "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```



```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file