

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.1-Sine/187-4.1.4.2

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 9:53am

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22
2	detailed summary tables of results	23
2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	37
3	Listing of integrals	39
3.1	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx$.	41
3.2	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$.	52
3.3	$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$. . .	62
3.4	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$	70
3.5	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$	77
3.6	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$	85

3.7 $\int \frac{A+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)(c-c \sin(e+fx))^{3/2}}} dx \dots\dots\dots 93$

3.8 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^n (A+C \sin^2(e+fx)) dx \dots 101$

3.9 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^n (A+C \sin^2(e+fx)) dx \dots 110$

3.10 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A+C \sin^2(e+fx)) dx 119$

3.11 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A+C \sin^2(e+fx)) dx \dots 128$

3.12 $\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)} (A+C \sin^2(e+fx)) dx \dots 137$

3.13 $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx \dots\dots\dots 146$

3.14 $\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx \dots\dots\dots 155$

3.15 $\int \frac{A+B \sin(e+fx)+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)(c-c \sin(e+fx))^{3/2}}} dx \dots\dots\dots 164$

3.16 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^n (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 172$

3.17 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{5/2} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 181$

3.18 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{3/2} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 193$

3.19 $\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 203$

3.20 $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx \dots\dots\dots 212$

3.21 $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx \dots\dots\dots 220$

3.22 $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx \dots\dots\dots 228$

3.23 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{-2-m} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 236$

3.24 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 245$

3.25 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 254$

3.26 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 264$

3.27 $\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)} (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 274$

3.28 $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx \dots\dots\dots 283$

3.29 $\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx \dots\dots\dots 292$

3.30 $\int (a+b \sin(c+dx)) (A+B \sin(c+dx)+C \sin^2(c+dx)) dx \dots\dots 301$

3.31 $\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{3/2}(e+fx)} dx \dots\dots\dots 308$

3.32 $\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx)+C \sin^2(e+fx)) dx 317$

4 Appendix 323

4.1 Listing of Grading functions 323

4.2 Links to plain text integration problems used in this report for each CAS 341

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [32]. This is test number [187].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (32)	0.00 (0)
Mathematica	56.25 (18)	43.75 (14)
Fricas	28.12 (9)	71.88 (23)
Mupad	28.12 (9)	71.88 (23)
Maxima	21.88 (7)	78.12 (25)
Maple	15.62 (5)	84.38 (27)
Reduce	6.25 (2)	93.75 (30)
Giac	3.12 (1)	96.88 (31)
Sympy	3.12 (1)	96.88 (31)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	96.875	0.000	0.000	3.125
Mathematica	46.875	0.000	6.250	46.875
Fricas	15.625	6.250	3.125	75.000
Maple	3.125	9.375	0.000	87.500
Giac	3.125	0.000	0.000	96.875
Maxima	3.125	18.750	0.000	78.125
Mupad	0.000	25.000	0.000	75.000
Reduce	0.000	3.125	0.000	96.875
Sympy	0.000	3.125	0.000	96.875

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	14	100.00	0.00	0.00
Fricas	23	100.00	0.00	0.00
Mupad	23	0.00	100.00	0.00
Maxima	25	96.00	4.00	0.00
Maple	27	100.00	0.00	0.00
Giac	31	70.97	9.68	19.35
Reduce	30	100.00	0.00	0.00
Sympy	31	54.84	41.94	3.23

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Sympy	0.13
Maxima	0.16
Reduce	0.24
Fricas	0.27
Giac	0.43
Rubi	1.13
Maple	4.98
Mathematica	23.12
Mupad	59.40

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Giac	76.00	0.94	76.00	0.94
Reduce	103.50	1.78	103.50	1.78
Sympy	189.00	2.33	189.00	2.33
Mathematica	252.56	1.06	169.50	0.93
Rubi	292.19	1.06	302.50	1.08
Maple	401.20	2.81	396.00	3.38
Fricas	406.00	1.61	310.00	1.62
Mupad	541.22	2.00	510.00	2.45
Maxima	714.43	2.51	648.00	2.45

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{32}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {9, 10, 11, 12, 13, 14, 24, 25, 26, 27, 28, 29}

Mathematica {23}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	37

2.1 List of integrals sorted by grade for each CAS

Rubi	24
Mma	24
Maple	25
Fricas	25
Maxima	25
Giac	26
Mupad	26
Sympy	26
Reduce	27

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 2, 3, 4, 5, 6, 7, 15, 18, 19, 20, 21, 22, 23, 30, 31 }

B grade { }

C grade { 1, 17 }

F normal fail { 8, 9, 10, 11, 12, 13, 14, 16, 24, 25, 26, 27, 28, 29 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 30 }

B grade { 7, 15, 31 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 2, 3, 18, 19, 30 }

B grade { 1, 17 }

C grade { 31 }

F normal fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 30 }

B grade { 1, 2, 3, 17, 18, 19 }

C grade { }

F normal fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 31 }

F(-1) timedout fail { 32 }

F(-2) exception fail { }

Giac

A grade { 30 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 23, 24, 25, 26, 27, 28, 29, 31 }

F(-1) timedout fail { 4, 20, 32 }

F(-2) exception fail { 5, 6, 7, 15, 21, 22 }

Mupad

A grade { }

B grade { 1, 2, 3, 17, 18, 19, 30, 31 }

C grade { }

F normal fail { }

F(-1) timedout fail { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 }

F(-2) exception fail { }

Sympy

A grade { }

B grade { 30 }

C grade { }

F normal fail { 3, 4, 5, 7, 8, 12, 13, 14, 15, 19, 20, 21, 23, 27, 28, 29, 31 }

F(-1) timedout fail { 1, 2, 6, 9, 10, 11, 16, 17, 18, 22, 24, 25, 26 }

F(-2) exception fail { 32 }

Reduce

A grade { }

B grade { 30 }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24,
25, 26, 27, 28, 29, 31 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	B	B	F(-1)	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	384	312	899	0	892	777	0	0	215	1110
N.S.	1	0.81	2.34	0.00	2.32	2.02	0.00	0.00	0.56	2.89
time (sec)	N/A	1.475	15.587	0.000	0.182	0.146	0.000	0.000	0.322	54.808

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	B	A	F(-1)	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	285	255	264	0	648	461	0	0	141	714
N.S.	1	0.89	0.93	0.00	2.27	1.62	0.00	0.00	0.49	2.51
time (sec)	N/A	1.198	9.399	0.000	0.173	0.127	0.000	0.000	0.237	47.979

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	B	A	F	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	180	183	160	0	441	261	0	0	68	185
N.S.	1	1.02	0.89	0.00	2.45	1.45	0.00	0.00	0.38	1.03
time (sec)	N/A	0.931	2.819	0.000	0.156	0.100	0.000	0.000	0.204	42.354

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	123	128	103	0	0	0	0	0	92	0
N.S.	1	1.04	0.84	0.00	0.00	0.00	0.00	0.00	0.75	0.00
time (sec)	N/A	0.648	36.804	0.000	0.000	0.000	0.000	0.000	0.183	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	202	214	110	0	0	0	0	0	111	0
N.S.	1	1.06	0.54	0.00	0.00	0.00	0.00	0.00	0.55	0.00
time (sec)	N/A	1.099	64.316	0.000	0.000	0.000	0.000	0.000	0.191	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	207	223	131	0	0	0	0	0	132	0
N.S.	1	1.08	0.63	0.00	0.00	0.00	0.00	0.00	0.64	0.00
time (sec)	N/A	1.157	53.966	0.000	0.000	0.000	0.000	0.000	0.186	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	181	190	588	0	0	0	0	130	0
N.S.	1	1.08	1.14	3.52	0.00	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	1.120	4.783	6.955	0.000	0.000	0.000	0.000	0.189	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	256	306	0	0	0	0	0	0	69	0
N.S.	1	1.20	0.00	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	1.193	0.000	0.000	0.000	0.000	0.000	0.000	0.233	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	358	388	0	0	0	0	0	0	67	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	1.097	0.000	0.000	0.000	0.000	0.000	0.000	0.219	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	382	423	0	0	0	0	0	0	173	0
N.S.	1	1.11	0.00	0.00	0.00	0.00	0.00	0.00	0.45	0.00
time (sec)	N/A	1.348	0.000	0.000	0.000	0.000	0.000	0.000	0.226	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	372	406	0	0	0	0	0	0	140	0
N.S.	1	1.09	0.00	0.00	0.00	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	1.219	0.000	0.000	0.000	0.000	0.000	0.000	0.265	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	366	396	0	0	0	0	0	0	65	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.18	0.00
time (sec)	N/A	1.180	0.000	0.000	0.000	0.000	0.000	0.000	0.213	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	364	395	0	0	0	0	0	0	89	0
N.S.	1	1.09	0.00	0.00	0.00	0.00	0.00	0.00	0.24	0.00
time (sec)	N/A	1.182	0.000	0.000	0.000	0.000	0.000	0.000	0.195	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	405	431	0	0	0	0	0	0	121	0
N.S.	1	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	1.295	0.000	0.000	0.000	0.000	0.000	0.000	0.204	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	188	196	897	0	0	0	0	191	0
N.S.	1	1.08	1.13	5.16	0.00	0.00	0.00	0.00	1.10	0.00
time (sec)	N/A	1.151	4.956	13.925	0.000	0.000	0.000	0.000	0.187	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	268	318	0	0	0	0	0	0	105	0
N.S.	1	1.19	0.00	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	1.224	0.000	0.000	0.000	0.000	0.000	0.000	0.335	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	B	B	F(-1)	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	435	336	1029	0	1324	937	0	0	322	1253
N.S.	1	0.77	2.37	0.00	3.04	2.15	0.00	0.00	0.74	2.88
time (sec)	N/A	1.508	15.927	0.000	0.213	0.169	0.000	0.000	0.360	47.127

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	B	A	F(-1)	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	322	279	306	0	950	563	0	0	212	790
N.S.	1	0.87	0.95	0.00	2.95	1.75	0.00	0.00	0.66	2.45
time (sec)	N/A	1.241	11.047	0.000	0.192	0.135	0.000	0.000	0.252	51.634

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	B	A	F	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	197	200	177	0	644	310	0	0	102	510
N.S.	1	1.02	0.90	0.00	3.27	1.57	0.00	0.00	0.52	2.59
time (sec)	N/A	0.992	3.076	0.000	0.178	0.104	0.000	0.000	0.221	39.554

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	170	175	159	0	0	0	0	0	136	0
N.S.	1	1.03	0.94	0.00	0.00	0.00	0.00	0.00	0.80	0.00
time (sec)	N/A	0.929	40.443	0.000	0.000	0.000	0.000	0.000	0.192	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	216	228	162	0	0	0	0	0	165	0
N.S.	1	1.06	0.75	0.00	0.00	0.00	0.00	0.00	0.76	0.00
time (sec)	N/A	1.189	54.181	0.000	0.000	0.000	0.000	0.000	0.203	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	230	246	228	0	0	0	0	0	196	0
N.S.	1	1.07	0.99	0.00	0.00	0.00	0.00	0.00	0.85	0.00
time (sec)	N/A	1.246	53.123	0.000	0.000	0.000	0.000	0.000	0.192	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	236	299	196	0	0	0	0	0	247	0
N.S.	1	1.27	0.83	0.00	0.00	0.00	0.00	0.00	1.05	0.00
time (sec)	N/A	1.296	2.016	0.000	0.000	0.000	0.000	0.000	0.197	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	373	410	0	0	0	0	0	0	102	0
N.S.	1	1.10	0.00	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	1.174	0.000	0.000	0.000	0.000	0.000	0.000	0.239	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	400	441	0	0	0	0	0	0	261	0
N.S.	1	1.10	0.00	0.00	0.00	0.00	0.00	0.00	0.65	0.00
time (sec)	N/A	1.379	0.000	0.000	0.000	0.000	0.000	0.000	0.231	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	392	425	0	0	0	0	0	0	212	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.54	0.00
time (sec)	N/A	1.265	0.000	0.000	0.000	0.000	0.000	0.000	0.299	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	386	415	0	0	0	0	0	0	99	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	1.275	0.000	0.000	0.000	0.000	0.000	0.000	0.233	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	386	415	0	0	0	0	0	0	135	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	1.273	0.000	0.000	0.000	0.000	0.000	0.000	0.199	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	421	451	0	0	0	0	0	0	183	0
N.S.	1	1.07	0.00	0.00	0.00	0.00	0.00	0.00	0.43	0.00
time (sec)	N/A	1.387	0.000	0.000	0.000	0.000	0.000	0.000	0.209	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	118	92	80	102	71	189	76	105	93
N.S.	1	1.46	1.14	0.99	1.26	0.88	2.33	0.94	1.30	1.15
time (sec)	N/A	0.364	0.476	2.224	0.033	0.081	0.132	0.432	0.179	38.757

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	C	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	120	97	396	0	220	0	0	89	169
N.S.	1	1.03	0.83	3.38	0.00	1.88	0.00	0.00	0.76	1.44
time (sec)	N/A	0.673	2.912	0.846	0.000	0.105	0.000	0.000	0.185	39.255

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-1)	N/A	F(-2)	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	47	45	0	54	0	0	102	47
N.S.	1	1.00	1.04	1.00	0.00	1.20	0.00	0.00	2.27	1.04
time (sec)	N/A	0.307	40.356	0.925	0.000	1.423	0.000	0.000	0.297	173.106

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [10] had the largest ratio of [.292683000000000026]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	11	11	0.81	40	0.275
2	A	9	9	0.89	40	0.225
3	A	7	7	1.02	40	0.175
4	A	8	7	1.04	40	0.175
5	A	11	10	1.06	40	0.250
6	A	11	10	1.08	40	0.250
7	A	11	10	1.08	42	0.238
8	A	12	11	1.20	38	0.289
9	A	11	10	1.08	37	0.270
10	A	13	12	1.11	41	0.293
11	A	12	11	1.09	39	0.282
12	A	12	11	1.08	39	0.282
13	A	12	11	1.09	39	0.282
14	A	12	11	1.06	39	0.282
15	A	11	10	1.08	50	0.200
16	A	12	11	1.19	46	0.239
17	A	11	11	0.77	48	0.229
18	A	9	9	0.87	48	0.188
19	A	7	7	1.02	48	0.146
20	A	10	9	1.03	48	0.188
21	A	11	10	1.06	48	0.208

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	11	10	1.07	48	0.208
23	A	11	10	1.27	50	0.200
24	A	11	10	1.10	45	0.222
25	A	13	12	1.10	49	0.245
26	A	12	11	1.08	47	0.234
27	A	12	11	1.08	47	0.234
28	A	12	11	1.08	47	0.234
29	A	12	11	1.07	47	0.234
30	A	4	4	1.46	31	0.129
31	A	11	11	1.03	41	0.268
32	N/A	2	0	1.00	45	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx$. . . 41
3.2	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$. . . 52
3.3	$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$ 62
3.4	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$ 70
3.5	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$ 77
3.6	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$ 85
3.7	$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)(c - c \sin(e + fx))^{3/2}}} dx$ 93
3.8	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$ 101
3.9	$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$ 110
3.10	$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$. . . 119
3.11	$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$. . . 128
3.12	$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$ 137
3.13	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$ 146
3.14	$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$ 155
3.15	$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)(c - c \sin(e + fx))^{3/2}}} dx$ 164
3.16	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$	172
3.17	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$	181
3.18	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$	193
3.19	$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$	203
3.20	$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$ 212
3.21	$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$ 220
3.22	$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$ 228
3.23	$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$	236
3.24	$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$	245

3.25	$\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	254
3.26	$\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	264
3.27	$\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)} (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	274
3.28	$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$	283
3.29	$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$	292
3.30	$\int (a+b \sin(c+dx)) (A+B \sin(c+dx) + C \sin^2(c+dx)) dx$	301
3.31	$\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{3/2}(e+fx)} dx$	308
3.32	$\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$	317

3.1 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{5/2} (A + C \sin^2(e -$

Optimal result	41
Mathematica [C] (verified)	42
Rubi [A] (verified)	43
Maple [F]	46
Fricas [B] (verification not implemented)	47
Sympy [F(-1)]	48
Maxima [B] (verification not implemented)	48
Giac [F]	49
Mupad [B] (verification not implemented)	50
Reduce [F]	51

Optimal result

Integrand size = 40, antiderivative size = 384

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \frac{64c^3(C(39 - 16m + 4m^2) + A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(9 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}} + \frac{16c^2(C(39 - 16m + 4m^2) + A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)}}{f(7 + 2m)(9 + 2m)(15 + 16m + 4m^2)} + \frac{2c(C(39 - 16m + 4m^2) + A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2}}{f(5 + 2m)(7 + 2m)(9 + 2m)} - \frac{4C(1 + 2m) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{f(7 + 2m)(9 + 2m)} + \frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{7/2}}{cf(9 + 2m)}$$

output

```
64*c^3*(C*(4*m^2-16*m+39)+A*(4*m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m
/f/(5+2*m)/(7+2*m)/(9+2*m)/(4*m^2+8*m+3)/(c-c*sin(f*x+e))^(1/2)+16*c^2*(C*
(4*m^2-16*m+39)+A*(4*m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(
f*x+e))^(1/2)/f/(7+2*m)/(9+2*m)/(4*m^2+16*m+15)+2*c*(C*(4*m^2-16*m+39)+A*(
4*m^2+32*m+63))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(5+
2*m)/(7+2*m)/(9+2*m)-4*C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*
x+e))^(5/2)/f/(7+2*m)/(9+2*m)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f
*x+e))^(7/2)/c/f/(9+2*m)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 15.59 (sec) , antiderivative size = 899, normalized size of antiderivative = 2.34

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + C*Sin[e + f*x]^2),x]
```

output

```
((a*(1 + Sin[e + f*x]))^m*(c - c*Sin[e + f*x])^(5/2)*(((18900*A + 12285*C + 15648*A*m + 648*C*m + 5280*A*m^2 + 1416*C*m^2 + 896*A*m^3 + 224*C*m^3 + 64*A*m^4 + 16*C*m^4)*((1/8 + I/8)*Cos[(e + f*x)/2] + (1/8 - I/8)*Sin[(e + f*x)/2])))/((1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((18900*A + 12285*C + 15648*A*m + 648*C*m + 5280*A*m^2 + 1416*C*m^2 + 896*A*m^3 + 224*C*m^3 + 64*A*m^4 + 16*C*m^4)*((1/8 - I/8)*Cos[(e + f*x)/2] + (1/8 + I/8)*Sin[(e + f*x)/2])))/((1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((1575*A + 1575*C + 1178*A*m + 414*C*m + 292*A*m^2 + 100*C*m^2 + 24*A*m^3 + 8*C*m^3)*((1/4 - I/4)*Cos[(3*(e + f*x))/2] - (1/4 + I/4)*Sin[(3*(e + f*x))/2])))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((1575*A + 1575*C + 1178*A*m + 414*C*m + 292*A*m^2 + 100*C*m^2 + 24*A*m^3 + 8*C*m^3)*((1/4 + I/4)*Cos[(3*(e + f*x))/2] - (1/4 - I/4)*Sin[(3*(e + f*x))/2])))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((63*A + 189*C + 32*A*m + 44*C*m + 4*A*m^2 + 4*C*m^2)*((-1/4 + I/4)*Cos[(5*(e + f*x))/2] - (1/4 + I/4)*Sin[(5*(e + f*x))/2])))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((63*A + 189*C + 32*A*m + 44*C*m + 4*A*m^2 + 4*C*m^2)*((-1/4 - I/4)*Cos[(5*(e + f*x))/2] - (1/4 - I/4)*Sin[(5*(e + f*x))/2])))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((15 + 2*m)*((-3/16 - (3*I)/16)*C*Cos[(7*(e + f*x))/2] + (3/16 - (3*I)/16)*C*Sin[(7*(e + f*x))/2])))/((7 + 2*m)*(9 + 2*m)) + ((15 + 2*m)*((-3/16 + (3*I)/16)*C*Cos[(7*(e + f*x))/2] + (3/16 + (3*I)/16)*C*Sin[(7*(e + f*x))/2])))/((7 + 2*m)*(9 + ...
```

Rubi [A] (verified)

Time = 1.48 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.81, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.275$, Rules used = {3042, 3519, 27, 3042, 3452, 3042, 3219, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3519}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)} -$$

$$\frac{2 \int -\frac{1}{2} (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 9)}}$$

$$\downarrow \text{27}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 9)} +$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

$$\downarrow \text{3042}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} (ac(C(7 - 2m) + A(2m + 9)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 9)} +$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

$$\downarrow \text{3452}$$

$$\frac{ac(A(4m^2 + 32m + 63) + C(4m^2 - 16m + 39)) \int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^{5/2} dx}{2m + 7} - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m}{f(2m + 7)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2} (a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

↓ 3042

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \int (\sin(e+fx)a+a)^m (c-c \sin(e+fx))^{5/2} dx}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c \sin(e+fx))^{5/2} (a \sin(e+fx)+a)^m}{f(2m+7)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2} (a \sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3219

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left(\frac{8c \int (\sin(e+fx)a+a)^m (c-c \sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c \sin(e+fx))^{3/2} (a \sin(e+fx)+a)^m}{f(2m+5)} \right)}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c \sin(e+fx))^{3/2} (a \sin(e+fx)+a)^m}{f(2m+5)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2} (a \sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3042

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left(\frac{8c \int (\sin(e+fx)a+a)^m (c-c \sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c \sin(e+fx))^{3/2} (a \sin(e+fx)+a)^m}{f(2m+5)} \right)}{2m+7} - \frac{4acC(2m+1) \cos(e+fx)(c-c \sin(e+fx))^{3/2} (a \sin(e+fx)+a)^m}{f(2m+5)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2} (a \sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3219

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left(\frac{8c \left(\frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c \sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} \right)}{2m+7} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2} (a \sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3042

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left(\frac{8c \left(\frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c \sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} \right)}{2m+7} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)}$$

$$\frac{ac(2m+9)}{cf(2m+9)} \frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2} (a \sin(e+fx)+a)^m}{cf(2m+9)}$$

↓ 3217

$$\frac{ac(A(4m^2+32m+63)+C(4m^2-16m+39)) \left(\frac{8c \left(\frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3)\sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx)\sqrt{c-c \sin(e+fx)}(a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} + \frac{2c \cos(e+fx)(c-}{2m+7} \right)}{ac(2m+9)}$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + C*Sin[e + f*x]^2),x]`

output `(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(7/2))/(c*f*(9 + 2*m)) + ((-4*a*c*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(f*(7 + 2*m)) + (a*c*(C*(39 - 16*m + 4*m^2) + A*(63 + 32*m + 4*m^2))*((2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) + (8*c*((8*c^2*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]])/(f*(3 + 2*m))))/(5 + 2*m)))/(7 + 2*m))/(a*c*(9 + 2*m))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3217 `Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]`

rule 3219 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])`

rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

rule 3519 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]`

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{\frac{5}{2}} (A + C \sin(fx + e))^2 dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 777 vs. $2(374) = 748$.

Time = 0.15 (sec) , antiderivative size = 777, normalized size of antiderivative = 2.02

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x,
algorithm="fricas")
```

output

```
2*((16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m + 105*C*c^2)
)*cos(f*x + e)^5 + 128*(A + C)*c^2*m^2 - (16*C*c^2*m^4 + 224*C*c^2*m^3 + 7
76*C*c^2*m^2 + 904*C*c^2*m + 285*C*c^2)*cos(f*x + e)^4 + 512*(2*A - C)*c^2
*m - (16*(A + 3*C)*c^2*m^4 + 32*(5*A + 16*C)*c^2*m^3 + 8*(65*A + 253*C)*c^
2*m^2 + 8*(75*A + 328*C)*c^2*m + 3*(63*A + 289*C)*c^2)*cos(f*x + e)^3 + 96
*(21*A + 13*C)*c^2 + (16*(A + C)*c^2*m^4 + 224*(A + C)*c^2*m^3 + 8*(133*A
+ 85*C)*c^2*m^2 + 1864*(A + C)*c^2*m + 3*(231*A + 263*C)*c^2)*cos(f*x + e)
^2 + 2*(16*(A + C)*c^2*m^4 + 192*(A + C)*c^2*m^3 + 856*(A + C)*c^2*m^2 + 1
6*(109*A + 85*C)*c^2*m + 3*(483*A + 419*C)*c^2)*cos(f*x + e) + (128*(A + C)
)*c^2*m^2 + (16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m +
105*C*c^2)*cos(f*x + e)^4 + 512*(2*A - C)*c^2*m + 2*(16*C*c^2*m^4 + 176*C*
c^2*m^3 + 560*C*c^2*m^2 + 628*C*c^2*m + 195*C*c^2)*cos(f*x + e)^3 + 96*(21
*A + 13*C)*c^2 - (16*(A + C)*c^2*m^4 + 160*(A + C)*c^2*m^3 + 8*(65*A + 113
*C)*c^2*m^2 + 24*(25*A + 57*C)*c^2*m + 9*(21*A + 53*C)*c^2)*cos(f*x + e)^2
- 2*(16*(A + C)*c^2*m^4 + 192*(A + C)*c^2*m^3 + 792*(A + C)*c^2*m^2 + 16*
(77*A + 101*C)*c^2*m + 3*(147*A + 211*C)*c^2)*cos(f*x + e))*sin(f*x + e))*
sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(32*f*m^5 + 400*f*m^4 + 1
840*f*m^3 + 3800*f*m^2 + 3378*f*m + (32*f*m^5 + 400*f*m^4 + 1840*f*m^3 + 3
800*f*m^2 + 3378*f*m + 945*f)*cos(f*x + e) - (32*f*m^5 + 400*f*m^4 + 1840*
f*m^3 + 3800*f*m^2 + 3378*f*m + 945*f)*sin(f*x + e) + 945*f)
```


Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(5/2)*(A+C*sin(f*x+e)**2), x)`

output `Timed out`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 892 vs. $2(374) = 748$.

Time = 0.18 (sec) , antiderivative size = 892, normalized size of antiderivative = 2.32

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2), x, algorithm="maxima")`

output

```

-2*((4*m^2 + 24*m + 43)*a^m*c^(5/2) - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*sin(f*x + e)/(cos(f*x + e) + 1) + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + (4*m^2 + 24*m + 43)*a^m*c^(5/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^3 + 36*m^2 + 46*m + 15)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(5/2)) + 4*(2*(4*m^2 + 56*m + 219)*a^m*c^(5/2) - 4*(4*m^3 + 56*m^2 + 219*m)*a^m*c^(5/2)*sin(f*x + e)/(cos(f*x + e) + 1) + (16*m^4 + 240*m^3 + 1136*m^2 + 1380*m + 1971)*a^m*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - (48*m^4 + 496*m^3 + 1568*m^2 + 3108*m - 315)*a^m*c^(5/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 4*(8*m^4 + 68*m^3 + 290*m^2 + 111*m + 567)*a^m*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 4*(8*m^4 + 68*m^3 + 290*m^2 + 111*m + 567)*a^m*c^(5/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - (48*m^4 + 496*m^3 + 1568*m^2 + 3108*m - 315)*a^m*c^(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + (16*m^4 + 240*m^3 + 1136*m^2 + 1380*m + 1971)*a^m*c^(5/2)*sin(f*x + e)^7/(cos(f*x + e) + 1)^7 - 4*(4*m^3 + 56*m^2 + 219*m)*a^m*c^(5/2)*sin(f*x + e)^8/(cos(f*x + e) + 1)^8 + 2*(4*m^2 + 56*m + 219)*a^m*c^(5/2)*sin(f*x + e)^9/(cos(f*x + e) + 1)^9)*C*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(...

```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e)^2 + A) (-c \sin(fx + e) + c)^{5/2} (a \sin(fx + e) + a)^m dx$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x,
algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + A)*(-c*sin(f*x + e) + c)^(5/2)*(a*sin(f*x + e) + a)^m, x)

```

Mupad [B] (verification not implemented)

Time = 54.81 (sec) , antiderivative size = 1110, normalized size of antiderivative = 2.89

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(5/2),x)
```

output

```
((c - c*sin(e + f*x))^(1/2)*((C*c^2*(a + a*sin(e + f*x))^m*(m^352i + m^2*344i + m^3*128i + m^4*16i + 105i))/(8*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*5i + f*x*5i)*(a + a*sin(e + f*x))^m*(18900*A + 12285*C + 15648*A*m + 648*C*m + 5280*A*m^2 + 896*A*m^3 + 64*A*m^4 + 1416*C*m^2 + 224*C*m^3 + 16*C*m^4))/(4*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*4i + f*x*4i)*(a + a*sin(e + f*x))^m*(A*18900i + C*12285i + A*m*15648i + C*m*648i + A*m^2*5280i + A*m^3*896i + A*m^4*64i + C*m^2*1416i + C*m^3*224i + C*m^4*16i))/(4*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*3i + f*x*3i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(1575*A + 1575*C + 1178*A*m + 414*C*m + 292*A*m^2 + 24*A*m^3 + 100*C*m^2 + 8*C*m^3))/(2*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*6i + f*x*6i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(A*1575i + C*1575i + A*m*1178i + C*m*414i + A*m^2*292i + A*m^3*24i + C*m^2*100i + C*m^3*8i))/(2*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (C*c^2*exp(e*9i + f*x*9i)*(a + a*sin(e + f*x))^m*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))/(8*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) - (3*C*c^2*exp(e*1i + f*x*1i)*(a + a*sin(e + f*x))^m*(720*m + 632*m^2 + 192*m^3 + 16*m^4 + 225))/(8*f*(m^3378i + m^2*3800i + m^3*1840i + m^4*400i + m^5*32i + 945i)) - (3*C*c^2*exp(e*8i + f*x*8i)*(a + a*sin(e + f*x))^m*(m^720i + m^2*...
```

Reduce [F]

$$\begin{aligned}
& \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A \\
& + C \sin^2(e + fx)) dx = \sqrt{c} c^2 \left(\left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^4 dx \right) c \right. \\
& - 2 \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^3 dx \right) c \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) a \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) c \\
& - 2 \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) a \\
& \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} dx \right) a \right)
\end{aligned}$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+C*sin(f*x+e)^2),x)`

output `sqrt(c)*c**2*(int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**4,x)*c - 2*int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**3,x)*c + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**2,x)*a + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**2,x)*c - 2*int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x),x)*a + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1),x)*a)`

3.2 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{3/2} (A + C \sin^2(e -$

Optimal result	52
Mathematica [A] (verified)	53
Rubi [A] (verified)	53
Maple [F]	57
Fricas [A] (verification not implemented)	57
Sympy [F(-1)]	58
Maxima [B] (verification not implemented)	58
Giac [F]	59
Mupad [B] (verification not implemented)	60
Reduce [F]	61

Optimal result

Integrand size = 40, antiderivative size = 285

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \frac{8c^2(C(19 - 8m + 4m^2) + A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}} + \frac{2c(C(19 - 8m + 4m^2) + A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)}}{f(3 + 2m)(5 + 2m)(7 + 2m)} - \frac{4C(1 + 2m) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2}}{f(5 + 2m)(7 + 2m)} + \frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{cf(7 + 2m)}$$

output

```
8*c^2*(C*(4*m^2-8*m+19)+A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f
/(5+2*m)/(7+2*m)/(4*m^2+8*m+3)/(c-c*sin(f*x+e))^(1/2)+2*c*(C*(4*m^2-8*m+19)
)+A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)/
f/(3+2*m)/(5+2*m)/(7+2*m)-4*C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*s
in(f*x+e))^(3/2)/f/(5+2*m)/(7+2*m)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*
sin(f*x+e))^(5/2)/c/f/(7+2*m)
```

Mathematica [A] (verified)

Time = 9.40 (sec) , antiderivative size = 264, normalized size of antiderivative = 0.93

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \frac{c(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (700$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2),x]
```

output

```
(c*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*sqrt[c - c*Sin[e + f*x]]*(700*A + 494*C + 760*A*m + 284*C*m + 272*A*m^2 + 136*C*m^2 + 32*A*m^3 + 16*C*m^3 - 2*C*(39 + 110*m + 68*m^2 + 8*m^3)*Cos[2*(e + f*x)] - (1 + 2*m)*(4*A*(35 + 24*m + 4*m^2) + C*(253 + 80*m + 12*m^2))*Sin[e + f*x] + 15*C*Sin[3*(e + f*x)] + 46*C*m*Sin[3*(e + f*x)] + 36*C*m^2*Sin[3*(e + f*x)] + 8*C*m^3*Sin[3*(e + f*x)]))/(2*f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))
```

Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.89, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.225$, Rules used = {3042, 3519, 27, 3042, 3452, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) dx$$

↓ 3042

$$\int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) dx$$

↓ 3519

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)} - \frac{2 \int -\frac{1}{2}(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}(ac(C(5 - 2m) + A(2m + 7)) + 2acC(2m + 1) \sin(e + fx))dx}{ac(2m + 7)}$$

↓ 27

$$\frac{\int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}(ac(C(5 - 2m) + A(2m + 7)) + 2acC(2m + 1) \sin(e + fx))dx}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{\int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}(ac(C(5 - 2m) + A(2m + 7)) + 2acC(2m + 1) \sin(e + fx))dx}{ac(2m + 7)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3452

$$\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19)) \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}dx}{2m + 5} - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{f(2m + 5)}$$

$$\frac{ac(2m + 7)}{cf(2m + 7)} \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19)) \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}dx}{2m + 5} - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{f(2m + 5)}$$

$$\frac{ac(2m + 7)}{cf(2m + 7)} \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3219

$$\frac{ac(A(4m^2 + 24m + 35) + C(4m^2 - 8m + 19)) \left(\frac{4c \int(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}dx}{2m + 3} + \frac{2c \cos(e + fx) \sqrt{c - c \sin(e + fx)}(a \sin(e + fx) + a)^m}{f(2m + 3)} \right)}{2m + 5} - \frac{4acC(2m + 1) \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{f(2m + 5)}$$

$$\frac{ac(2m + 7)}{cf(2m + 7)} \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{ac(A(4m^2+24m+35)+C(4m^2-8m+19))\left(\frac{4c f(\sin(e+fx)a+a)^m \sqrt{c-c \sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)}\right)}{2m+5} - \frac{4acC(2m+1) \cos(e+fx)}{2m+5}$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{5/2}(a \sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3217

$$\frac{ac(A(4m^2+24m+35)+C(4m^2-8m+19))\left(\frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3)\sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)}\right)}{2m+5} - \frac{4acC(2m+1) \cos(e+fx)}{2m+5}$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{5/2}(a \sin(e+fx)+a)^m}{cf(2m+7)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2),x]
```

output

```
(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(c*f*(7 + 2*m)) + ((-4*a*c*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) + (a*c*(C*(19 - 8*m + 4*m^2) + A*(35 + 24*m + 4*m^2))*((8*c^2*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m))*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]])/(f*(3 + 2*m)))/(5 + 2*m)/(a*c*(7 + 2*m))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```


rule 3217

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]
```

rule 3219

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])
```

rule 3452

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]
```

rule 3519

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{\frac{3}{2}} (A + C \sin(fx + e))^2 dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)`

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.62

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx =$$

$$\frac{2((8Ccm^3 + 36Ccm^2 + 46Ccm + 15Cc) \cos(fx + e)^4 - 16(A + C)cm^2 + (8Ccm^3 + 68Ccm^2 + 110$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `-2*((8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^4 - 16*(A + C)*c*m^2 + (8*C*c*m^3 + 68*C*c*m^2 + 110*C*c*m + 39*C*c)*cos(f*x + e)^3 - 32*(3*A - C)*c*m - (8*(A + C)*c*m^3 + 4*(13*A + 5*C)*c*m^2 + 94*(A + C)*c*m + (35*A + 43*C)*c)*cos(f*x + e)^2 - 4*(35*A + 19*C)*c - (8*(A + C)*c*m^3 + 68*(A + C)*c*m^2 + 2*(95*A + 63*C)*c*m + (175*A + 143*C)*c)*cos(f*x + e) - (16*(A + C)*c*m^2 + (8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^3 + 32*(3*A - C)*c*m - 8*(4*C*c*m^2 + 8*C*c*m + 3*C*c)*cos(f*x + e)^2 + 4*(35*A + 19*C)*c - (8*(A + C)*c*m^3 + 52*(A + C)*c*m^2 + 2*(47*A + 79*C)*c*m + (35*A + 67*C)*c)*cos(f*x + e))*sin(f*x + e)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*cos(f*x + e) - (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*sin(f*x + e) + 105*f)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(3/2)*(A+C*sin(f*x+e)**2), x)`

output `Timed out`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 648 vs. $2(277) = 554$.

Time = 0.17 (sec) , antiderivative size = 648, normalized size of antiderivative = 2.27

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2), x, algorithm="maxima")`

output

```

-2*((a^m*c^(3/2)*(2*m + 5) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)/(cos(f*x +
e) + 1) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^m
*c^(3/2)*(2*m + 5)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)*A*e^(2*m*log(sin(f
*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^
2 + 1))/((4*m^2 + 8*m + 3)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2)
) + 4*(2*a^m*c^(3/2)*(2*m + 13) - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*x + e
))/(cos(f*x + e) + 1) + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)*sin(f*x +
e)^2/(cos(f*x + e) + 1)^2 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(f
*x + e)^3/(cos(f*x + e) + 1)^3 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*
sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(
3/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*si
n(f*x + e)^6/(cos(f*x + e) + 1)^6 + 2*a^m*c^(3/2)*(2*m + 13)*sin(f*x + e)^
7/(cos(f*x + e) + 1)^7)*C*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1)
- m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((16*m^4 + 128*m^3 + 344
*m^2 + 352*m + 2*(16*m^4 + 128*m^3 + 344*m^2 + 352*m + 105)*sin(f*x + e)^2
/(cos(f*x + e) + 1)^2 + (16*m^4 + 128*m^3 + 344*m^2 + 352*m + 105)*sin(f*x
+ e)^4/(cos(f*x + e) + 1)^4 + 105)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 +
1)^(3/2)))/f

```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A) (-c \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,
algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + A)*(-c*sin(f*x + e) + c)^(3/2)*(a*sin(f*x +
e) + a)^m, x)

```

Mupad [B] (verification not implemented)

Time = 47.98 (sec) , antiderivative size = 714, normalized size of antiderivative = 2.51

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(3/2),x)
```

output

```
((c - c*sin(e + f*x))^(1/2)*((C*c*(a + a*sin(e + f*x))^m*(m*46i + m^2*36i + m^3*8i + 15i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp(e*3i + f*x*3i)*(a + a*sin(e + f*x))^m*(1260*A + 735*C + 1144*A*m - 18*C*m + 336*A*m^2 + 32*A*m^3 + 100*C*m^2 + 8*C*m^3))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (c*exp(e*4i + f*x*4i)*(a + a*sin(e + f*x))^m*(A*1260i + C*735i + A*m*1144i - C*m*18i + A*m^2*336i + A*m^3*32i + C*m^2*100i + C*m^3*8i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (C*c*exp(e*7i + f*x*7i)*(a + a*sin(e + f*x))^m*(46*m + 36*m^2 + 8*m^3 + 15))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (C*c*exp(e*1i + f*x*1i)*(a + a*sin(e + f*x))^m*(174*m + 100*m^2 + 8*m^3 + 63))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (C*c*exp(e*6i + f*x*6i)*(a + a*sin(e + f*x))^m*(m*174i + m^2*100i + m^3*8i + 63i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp(e*5i + f*x*5i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(140*A + 175*C + 96*A*m + 16*C*m + 16*A*m^2 + 4*C*m^2))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (c*exp(e*2i + f*x*2i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(A*140i + C*175i + A*m*96i + C*m*16i + A*m^2*16i + C*m^2*4i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))))/(exp(e*4i + f*x*4i) - (exp(e*3i + f*x*3i)*(m*352i + m^2*344i + m^3*128i + m^4*16i + 105i))/(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))
```

Reduce [F]

$$\begin{aligned}
& \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A \\
& + C \sin^2(e + fx)) dx = \sqrt{c} c \left(- \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^3 dx \right) c \right. \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) c \\
& - \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) a \\
& \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} dx \right) a \right)
\end{aligned}$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)`

output `sqrt(c)*c*(- int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**3,x)*c + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**2,x)*c - int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x),x)*a + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1),x)*a)`

3.3 $\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+C \sin^2(e+fx)) dx$

Optimal result	62
Mathematica [A] (verified)	63
Rubi [A] (verified)	63
Maple [F]	66
Fricas [A] (verification not implemented)	66
Sympy [F]	67
Maxima [B] (verification not implemented)	67
Giac [F]	68
Mupad [B] (verification not implemented)	68
Reduce [F]	69

Optimal result

Integrand size = 40, antiderivative size = 180

$$\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+C \sin^2(e+fx)) dx$$

$$= \frac{2c(C-6Cm+A(5+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{f(1+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{4cC(1+2m) \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{af(3+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{2C \cos(e+fx)(a+a \sin(e+fx))^m(c-c \sin(e+fx))^{3/2}}{cf(5+2m)}$$

output

```
2*c*(C-6*C*m+A*(5+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(1+2*m)/(5+2*m)/(c-c*sin(f*x+e))^(1/2)+4*c*C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(3+2*m)/(5+2*m)/(c-c*sin(f*x+e))^(1/2)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/c/f/(5+2*m)
```

Mathematica [A] (verified)

Time = 2.82 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.89

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx =$$

$$\frac{(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (-30A - 19C - 32Am - 8Cm - 8Am^2 - 4Cm^2 + C(3 + 8m + 4m^2) \cos[2(e + fx)] + 8C(1 + 2m) \sin[e + fx])}{f(1 + 2m)(3 + 2m)(5 + 2m) (\cos(\frac{1}{2}(e + fx)) - \sin(\frac{1}{2}(e + fx)))}$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2),x]
```

output

```
-(((Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*Sqrt[c - c*Sin[e + f*x]]*(-30*A - 19*C - 32*A*m - 8*C*m - 8*A*m^2 - 4*C*m^2 + C*(3 + 8*m + 4*m^2)*Cos[2*(e + f*x)] + 8*C*(1 + 2*m)*Sin[e + f*x]))/(f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))
```

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {3042, 3519, 27, 3042, 3450, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3519}$$

$$\frac{2C \cos(e + fx) (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m}{cf(2m + 5)} - \frac{2 \int -\frac{1}{2} (\sin(e + fx) a + a)^m \sqrt{c - c \sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 5)}$$

↓ 27

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} (ac(C(3 - 2m) + A(2m + 5)) + 2acC(2m + 1) \sin(e + fx)) dx}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3450

$$\frac{ac(A(2m + 5) - 6Cm + C) \int (\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} dx + 2cC(2m + 1) \int (\sin(e + fx)a + a)^{m+1}}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3042

$$\frac{ac(A(2m + 5) - 6Cm + C) \int (\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} dx + 2cC(2m + 1) \int (\sin(e + fx)a + a)^{m+1}}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3217

$$\frac{\frac{2ac^2(A(2m+5)-6Cm+C) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}} + \frac{4c^2C(2m+1) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}}{f(2m+3)\sqrt{c-c \sin(e+fx)}}}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*sqrt[c - c*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2),x]
```

output

$$\frac{(2C \cos[e + fx] (a + a \sin[e + fx])^m (c - c \sin[e + fx])^{3/2}) / (c f (5 + 2m)) + ((2a^2 c^2 (C - 6Cm + A(5 + 2m)) \cos[e + fx] (a + a \sin[e + fx])^m) / (f(1 + 2m) \sqrt{c - c \sin[e + fx]}) + (4c^2 C(1 + 2m) \cos[e + fx] (a + a \sin[e + fx])^{1+m}) / (f(3 + 2m) \sqrt{c - c \sin[e + fx]})) / (a c (5 + 2m))$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_) /; \text{FreeQ}[b, x]]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3217

$$\text{Int}[\sqrt{(a_*) + (b_*) \sin[(e_*) + (f_*)(x_)]}] * ((c_*) + (d_*) \sin[(e_*) + (f_*)(x_)])^{n_}, x_Symbol] \rightarrow \text{Simp}[-2b \cos[e + fx] * ((c + d \sin[e + fx])^n / (f(2n + 1) \sqrt{a + b \sin[e + fx]})), x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[n, -2^{(-1)}]$$

rule 3450

$$\text{Int}[\sqrt{(a_*) + (b_*) \sin[(e_*) + (f_*)(x_)]}] * ((A_*) + (B_*) \sin[(e_*) + (f_*)(x_)])^{n_}, x_Symbol] \rightarrow \text{Simp}[B/d \text{ Int}[\sqrt{a + b \sin[e + fx]} * (c + d \sin[e + fx])^{n+1}, x], x] - \text{Simp}[(B*c - A*d)/d \text{ Int}[\sqrt{a + b \sin[e + fx]} * (c + d \sin[e + fx])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$$

rule 3519

$$\text{Int}[(a_*) + (b_*) \sin[(e_*) + (f_*)(x_)]^{m_*) * ((c_*) + (d_*) \sin[(e_*) + (f_*)(x_)])^{n_}, x_Symbol] \rightarrow \text{Simp}[(-C) \cos[e + fx] * (a + b \sin[e + fx])^m * ((c + d \sin[e + fx])^{n+1}) / (d*f*(m + n + 2)), x] + \text{Simp}[1/(b*d*(m + n + 2)) \text{ Int}[(a + b \sin[e + fx])^m * (c + d \sin[e + fx])^n * \text{Simp}[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) - b*c*C*(2*m + 1) \sin[e + fx], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, C, m, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ !\text{LtQ}[m, -2^{(-1)}] \ \&\& \ \text{NeQ}[m + n + 2, 0]$$

Maple [F]

$$\int (a + a \sin (fx + e))^m \sqrt{c - c \sin (fx + e)} (A + C \sin (fx + e))^2 dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.45

$$\int (a + a \sin (e + fx))^m \sqrt{c - c \sin (e + fx)} (A + C \sin^2 (e + fx)) dx =$$

$$\frac{2((4Cm^2 + 8Cm + 3C) \cos (fx + e)^3 - 4(A + C)m^2 + (4Cm^2 - C) \cos (fx + e)^2 - 16Am - (4(A + C)m^2 + 8(2A + C)m + 15A + 11C) \cos (fx + e) - (4(A + C)m^2 - (4Cm^2 + 8Cm + 3C) \cos (fx + e)^2 + 16Am - 4(2Cm + C) \cos (fx + e) + 15A + 7C) \sin (fx + e) - 15A - 7C) \sqrt{-c \sin (fx + e) + c} (a \sin (fx + e) + a)^m}{(8f^3m^3 + 36f^2m^2 + 46fm^2 + (8f^3m^3 + 36f^2m^2 + 46fm^2 + 15f) \cos (fx + e) - (8f^3m^3 + 36f^2m^2 + 46fm^2 + 15f) \sin (fx + e) + 15f)}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `-2*((4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^3 - 4*(A + C)*m^2 + (4*C*m^2 - C)*cos(f*x + e)^2 - 16*A*m - (4*(A + C)*m^2 + 8*(2*A + C)*m + 15*A + 11*C)*cos(f*x + e) - (4*(A + C)*m^2 - (4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^2 + 16*A*m - 4*(2*C*m + C)*cos(f*x + e) + 15*A + 7*C)*sin(f*x + e) - 15*A - 7*C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(8*f*m^3 + 36*f*m^2 + 46*f*m + (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*cos(f*x + e) - (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*sin(f*x + e) + 15*f)`

Sympy [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m \sqrt{-c(\sin(e + fx) - 1)} (A + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(1/2)*(A+C*sin(f*x+e)**2), x)`

output `Integral((a*(sin(e + f*x) + 1))**m*sqrt(-c*(sin(e + f*x) - 1))*(A + C*sin(e + f*x)**2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 441 vs. $2(174) = 348$.

Time = 0.16 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.45

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= 2 \left(\frac{4 \left(\frac{4 a^m \sqrt{c} m \sin(fx+e)}{\cos(fx+e)+1} - \frac{(4 m^2+4 m+5) a^m \sqrt{c} \sin(fx+e)^2}{(\cos(fx+e)+1)^2} - \frac{(4 m^2+4 m+5) a^m \sqrt{c} \sin(fx+e)^3}{(\cos(fx+e)+1)^3} + \frac{4 a^m \sqrt{c} m \sin(fx+e)^4}{(\cos(fx+e)+1)^4} - 2 a^m \sqrt{c} - \frac{2 a^m \sqrt{c} \sin(fx+e)}{(\cos(fx+e)+1)^5} \right)}{\left(8 m^3+36 m^2+46 m+ \frac{2(8 m^3+36 m^2+46 m+15) \sin(fx+e)^2}{(\cos(fx+e)+1)^2} + \frac{(8 m^3+36 m^2+46 m+15) \sin(fx+e)^4}{(\cos(fx+e)+1)^4} + 15 \right)} \right)$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2), x, algorithm="maxima")`

output

```
2*(4*(4*a^m*sqrt(c)*m*sin(f*x + e)/(cos(f*x + e) + 1) - (4*m^2 + 4*m + 5)*
a^m*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - (4*m^2 + 4*m + 5)*a^m*sq
rt(c)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 4*a^m*sqrt(c)*m*sin(f*x + e)^4
/(cos(f*x + e) + 1)^4 - 2*a^m*sqrt(c) - 2*a^m*sqrt(c)*sin(f*x + e)^5/(cos(
f*x + e) + 1)^5)*C*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log
(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^3 + 36*m^2 + 46*m + 2*(8*
m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + (8*m^3 + 3
6*m^2 + 46*m + 15)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 15)*sqrt(sin(f*x
+ e)^2/(cos(f*x + e) + 1)^2 + 1)) - (a^m*sqrt(c) + a^m*sqrt(c)*sin(f*x + e
)/(cos(f*x + e) + 1))*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) -
m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((2*m + 1)*sqrt(sin(f*x +
e)^2/(cos(f*x + e) + 1)^2 + 1)))/f
```

Giac [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) \sqrt{-c \sin(fx + e) + c(a \sin(fx + e) + a)^m} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,
algorithm="giac")
```

output

```
integrate((C*sin(f*x + e)^2 + A)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e)
+ a)^m, x)
```

Mupad [B] (verification not implemented)

Time = 42.35 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.03

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx =$$

$$\frac{(a(\sin(e + fx) + 1))^m \sqrt{-c(\sin(e + fx) - 1)} (60 A \cos(e + fx) + 35 C \cos(e + fx) - 3 C \cos(3$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(1/2),x)`

output `-((a*(sin(e + f*x) + 1))^m*(-c*(sin(e + f*x) - 1))^(1/2)*(60*A*cos(e + f*x) + 35*C*cos(e + f*x) - 3*C*cos(3*e + 3*f*x) - 8*C*sin(2*e + 2*f*x) - 4*C*m^2*cos(3*e + 3*f*x) + 64*A*m*cos(e + f*x) + 8*C*m*cos(e + f*x) + 16*A*m^2*cos(e + f*x) - 8*C*m*cos(3*e + 3*f*x) + 4*C*m^2*cos(e + f*x) - 16*C*m*sin(2*e + 2*f*x)))/(2*f*(sin(e + f*x) - 1)*(46*m + 36*m^2 + 8*m^3 + 15))`

Reduce [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + C \sin^2(e + fx)) dx \\ &= \sqrt{c} \left(\left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) c \right. \\ & \quad \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} dx \right) a \right) \end{aligned}$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

output `sqrt(c)*(int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**2,x)*c + int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1),x)*a)`

$$3.4 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$$

Optimal result	70
Mathematica [A] (verified)	71
Rubi [A] (verified)	71
Maple [F]	74
Fricas [F]	74
Sympy [F]	74
Maxima [F]	75
Giac [F(-1)]	75
Mupad [F(-1)]	76
Reduce [F]	76

Optimal result

Integrand size = 40, antiderivative size = 123

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \frac{(A + C) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) (a + a \sin(e + fx))^m}{f(1 + 2m)\sqrt{c - c \sin(e + fx)}} - \frac{2C \cos(e + fx)(a + a \sin(e + fx))^{1+m}}{af(3 + 2m)\sqrt{c - c \sin(e + fx)}}$$

```
output (A+C)*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2+m],1/2+1/2*sin(f*x+e))*(a+a*sin
(f*x+e))^m/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)-2*C*cos(f*x+e)*(a+a*sin(f*x+e)
)^(1+m)/a/f/(3+2*m)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 36.80 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.84

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx =$$

$$\frac{\cos(e + fx)(a(1 + \sin(e + fx)))^m \left(-((A + C)(3 + 2m) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) \right)}{f(1 + 2m)(3 + 2m)\sqrt{c - c \sin(e + fx)}}$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]
```

output

```
-((Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(-((A + C)*(3 + 2*m)*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]) + 2*C*(1 + 2*m)*(1 + Sin[e + f*x])))/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])
```

Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.04, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {3042, 3517, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{\sqrt{c - c \sin(e + fx)}} dx$$

$$\downarrow \text{3517}$$

$$(A + C) \int \frac{(\sin(e + fx)a + a)^m}{\sqrt{c - c \sin(e + fx)}} dx - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}}$$

$$\downarrow \text{3042}$$

$$(A + C) \int \frac{(\sin(e + fx)a + a)^m}{\sqrt{c - c\sin(e + fx)}} dx - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c\sin(e + fx)}}$$

↓ 3224

$$\frac{(A + C) \cos(e + fx) \int \sec(e + fx)(\sin(e + fx)a + a)^{m+\frac{1}{2}} dx}{\sqrt{a \sin(e + fx) + a}\sqrt{c - c\sin(e + fx)}} - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c\sin(e + fx)}}$$

↓ 3042

$$\frac{(A + C) \cos(e + fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{\sqrt{a \sin(e + fx) + a}\sqrt{c - c\sin(e + fx)}} - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c\sin(e + fx)}}$$

↓ 3146

$$\frac{a(A + C) \cos(e + fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a\sin(e+fx)} d(a \sin(e + fx))}{f\sqrt{a \sin(e + fx) + a}\sqrt{c - c\sin(e + fx)}} - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c\sin(e + fx)}}$$

↓ 78

$$\frac{(A + C) \cos(e + fx)(a \sin(e + fx) + a)^m \operatorname{Hypergeometric2F1}\left(1, m + \frac{1}{2}, m + \frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m + 1)\sqrt{c - c\sin(e + fx)}} - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c\sin(e + fx)}}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]
```

output

```
((A + C)*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) - (2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(a*f*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])
```

Definitions of rubi rules used

- rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`
- rule 3224 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`
- rule 3517 `Int[(((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2))/Sqrt[(c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*C*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(2*m + 3)*Sqrt[c + d*Sin[e + f*x]])), x] + Simp[(A + C) Int[(a + b*Sin[e + f*x])^m/Sqrt[c + d*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]`

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin(fx + e)^2)}{\sqrt{c - c \sin(fx + e)}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

Fricas [F]

$$\begin{aligned} & \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \\ &= \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(c*sin(f*x + e) - c), x)`

Sympy [F]

$$\begin{aligned} & \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx \\ &= \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{\sqrt{-c(\sin(e + fx) - 1)}} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(1/2),x)`

output

```
Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/sqrt(-c*(sin(e + f*x) - 1)), x)
```

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x,
algorithm="maxima")
```

output

```
integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/sqrt(-c*sin(f*x + e) + c), x)
```

Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx = \text{Timed out}$$

input

```
integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x,
algorithm="giac")
```

output

```
Timed out
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{\sqrt{c - c \sin(e + fx)}} dx$$

input `int(((A + C*sin(e + f*x))^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(1/2),x)`

output `int(((A + C*sin(e + f*x))^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(1/2),x)`

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx =$$

$$\frac{\sqrt{c} \left(\left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2}{\sin(fx + e) - 1} dx \right) c + \left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1}}{\sin(fx + e) - 1} dx \right) a \right)}{c}$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

output `(-sqrt(c)*int(((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**2)/(sin(e + f*x) - 1),x)*c + int(((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1))/(sin(e + f*x) - 1),x)*a)/c`

$$3.5 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$$

Optimal result	77
Mathematica [A] (verified)	78
Rubi [A] (verified)	78
Maple [F]	82
Fricas [F]	82
Sympy [F]	83
Maxima [F]	83
Giac [F(-2)]	83
Mupad [F(-1)]	84
Reduce [F]	84

Optimal result

Integrand size = 40, antiderivative size = 202

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + C) \cos(e + fx)(a + a \sin(e + fx))^{1+m}}{4af(c - c \sin(e + fx))^{3/2}} + \frac{(A + 2Am + C(9 + 2m)) \cos(e + fx)(a + a \sin(e + fx))^m}{4cf(1 + 2m)\sqrt{c - c \sin(e + fx)}} + \frac{(A(1 - 2m) - C(7 + 2m)) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right)}{4cf(1 + 2m)\sqrt{c - c \sin(e + fx)}} (a + a \sin(e + fx))$$

output

```
1/4*(A+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(3/2)+1/4
*(A+2*A*m+C*(9+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*
x+e))^(1/2)+1/4*(A*(1-2*m)-C*(7+2*m))*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2
+m],1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*x+e))^(1
/2)
```

Mathematica [A] (verified)

Time = 64.32 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.54

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx =$$

$$\frac{\cos(e + fx) \left(-4C + 4C \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx)) \right) \right) - (A + C) \operatorname{Hypergeometric2F1} \left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx)) \right)}{2cf(1 + 2m)\sqrt{c - c \sin(e + fx)}}$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(3/2),x]
```

output

```
-1/2*(Cos[e + f*x]*(-4*C + 4*C*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] - (A + C)*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2])*(a*(1 + Sin[e + f*x]))^m)/(c*f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]])
```

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3515, 27, 3042, 3452, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{3/2}} dx$$

$$\downarrow \text{3515}$$

$$\begin{aligned}
& \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} - \\
& \frac{\int -\frac{(\sin(e+fx)a+a)^m(a^2(A(3-2m)-C(2m+5))-a^2(2mA+A+C(2m+9))\sin(e+fx))}{2\sqrt{c-c\sin(e+fx)}} dx}{4a^2c} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{(\sin(e+fx)a+a)^m(a^2(A(3-2m)-C(2m+5))-a^2(2mA+A+C(2m+9))\sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx}{8a^2c} + \\
& \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(\sin(e+fx)a+a)^m(a^2(A(3-2m)-C(2m+5))-a^2(2mA+A+C(2m+9))\sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx}{8a^2c} + \\
& \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3452 \\
& \frac{2a^2(A(1-2m)-C(2m+7))\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+C(2m+9))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{8a^2c} + \\
& \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \frac{2a^2(A(1-2m)-C(2m+7))\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+C(2m+9))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{8a^2c} + \\
& \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3224 \\
& \frac{2a^2(A(1-2m)-C(2m+7))\cos(e+fx)\int \sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}} dx + \frac{2a^2(2Am+A+C(2m+9))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3042
\end{aligned}$$

$$\frac{2a^2(A(1-2m)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{\sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}} + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1) \sqrt{c-c \sin(e+fx)}} +$$

$$\frac{8a^2c}{4af(c-c \sin(e+fx))^{3/2}} (A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}$$

↓ 3146

$$\frac{2a^3(A(1-2m)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a \sin(e+fx)} d(a \sin(e+fx))}{f \sqrt{a \sin(e+fx)+a} \sqrt{c-c \sin(e+fx)}} + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1) \sqrt{c-c \sin(e+fx)}} +$$

$$\frac{8a^2c}{4af(c-c \sin(e+fx))^{3/2}} (A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}$$

↓ 78

$$\frac{2a^2(A(1-2m)-C(2m+7)) \cos(e+fx)(a \sin(e+fx)+a)^m \operatorname{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1) \sqrt{c-c \sin(e+fx)}} + \frac{2a^2(2Am+A+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1) \sqrt{c-c \sin(e+fx)}} +$$

$$\frac{8a^2c}{4af(c-c \sin(e+fx))^{3/2}} (A+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(3/2),x]
```

output

```
((A + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + ((2*a^2*(A + 2*A*m + C*(9 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*a^2*(A*(1 - 2*m) - C*(7 + 2*m))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(8*a^2*c)
```

Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3146 `Int[cos[(e_) + (f_)*(x_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`
- rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`
- rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]*(c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

rule 3515

```

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] :=
Simp[(a*A + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^
(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*
Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m
+ n + 2)) - C*(c^2*m - d^2*(n + 1)) + d*(A*c*(m + n + 2) - c*C*(3*m - n))*
Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ
[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2,
0] && NeQ[2*m + 1, 0]))

```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin(fx + e)^2)}{(c - c \sin(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{\frac{3}{2}}} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x,
algorithm="fricas")
```

output

```
integral((C*cos(f*x + e)^2 - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x +
e) + a)^m/(c^2*cos(f*x + e)^2 + 2*c^2*sin(f*x + e) - 2*c^2), x)
```

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{(-c(\sin(e + fx) - 1))^{\frac{3}{2}}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(3/2), x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/(-c*(sin(e + f*x) - 1))**(3/2), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{\frac{3}{2}}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2), x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) + c)^(3/2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2), x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,1,3,0,0,0,0]}+%%{3,[0,1,1,0,0,0,0]} / %%{16,[
0,0,0,1,1
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{(c - c \sin(e + fx))^{3/2}} dx$$

input

```
int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(
3/2),x)
```

output

```
int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(
3/2), x)
```

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \frac{\sqrt{c} \left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2}{\sin(fx + e)^2 - 2 \sin(fx + e) + 1} dx \right) c + \left(\dots \right)}{c^2}$$

input

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

output

```
(sqrt(c)*(int(((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f
*x)**2)/(sin(e + f*x)**2 - 2*sin(e + f*x) + 1),x)*c + int(((sin(e + f*x)*a
+ a)**m*sqrt(- sin(e + f*x) + 1))/(sin(e + f*x)**2 - 2*sin(e + f*x) + 1)
,x)*a))/c**2
```

3.6
$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$$

Optimal result	85
Mathematica [A] (verified)	86
Rubi [A] (verified)	86
Maple [F]	90
Fricas [F]	90
Sympy [F(-1)]	90
Maxima [F]	91
Giac [F(-2)]	91
Mupad [F(-1)]	91
Reduce [F]	92

Optimal result

Integrand size = 40, antiderivative size = 207

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \frac{(A + C) \cos(e + fx)(a + a \sin(e + fx))^{1+m}}{8af(c - c \sin(e + fx))^{5/2}} + \frac{(A(5 - 2m) - C(11 + 2m)) \cos(e + fx)(a + a \sin(e + fx))^m}{16cf(c - c \sin(e + fx))^{3/2}} + \frac{(A(3 - 8m + 4m^2) + C(19 + 24m + 4m^2)) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right)}{32c^2 f(1 + 2m) \sqrt{c - c \sin(e + fx)}}$$

output

```
1/8*(A+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(5/2)+1/16*(A*(5-2*m)-C*(11+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(c-c*sin(f*x+e))^(3/2)+1/32*(A*(4*m^2-8*m+3)+C*(4*m^2+24*m+19))*cos(f*x+e)*hypergeom([1, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c^2/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 53.97 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.63

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \frac{\cos(e + fx) (4C \text{Hypergeometric2F1}(1, \frac{1}{2} + m, \frac{3}{2} + m, \dots))}{(c - c \sin(e + fx))^{5/2}}$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(5/2), x]
```

output

```
(Cos[e + f*x]*(4*C*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] - 4*C*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] + (A + C)*Hypergeometric2F1[3, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2])*(a*(1 + Sin[e + f*x]))^m)/(4*c^2*(f + 2*f*m)*Sqrt[c - c*Sin[e + f*x]])
```

Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 3515, 27, 3042, 3451, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{5/2}} dx$$

↓ 3515

$$\frac{(A + C) \cos(e + fx) (a \sin(e + fx) + a)^{m+1}}{8af(c - c \sin(e + fx))^{5/2}} - \int \frac{(\sin(e + fx)a + a)^m ((A(9 - 2m) - C(2m + 7))a^2 + (A(1 - 2m) - C(2m + 15)) \sin(e + fx)a^2)}{2(c - c \sin(e + fx))^{3/2}} dx$$

↓ 27

$$8a^2c$$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m ((A(9-2m)-C(2m+7))a^2+(A(1-2m)-C(2m+15))\sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{\frac{16a^2c}{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}} \frac{1}{8af(c-c\sin(e+fx))^{5/2}}} +$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m ((A(9-2m)-C(2m+7))a^2+(A(1-2m)-C(2m+15))\sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{\frac{16a^2c}{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}} \frac{1}{8af(c-c\sin(e+fx))^{5/2}}} +$$

↓ 3451

$$\frac{\frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19))}{2c} \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{a^2(A(5-2m)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}}}{\frac{16a^2c}{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}} \frac{1}{8af(c-c\sin(e+fx))^{5/2}}} +$$

↓ 3042

$$\frac{\frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19))}{2c} \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{a^2(A(5-2m)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}}}{\frac{16a^2c}{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}} \frac{1}{8af(c-c\sin(e+fx))^{5/2}}} +$$

↓ 3224

$$\frac{\frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19))\cos(e+fx) \int \sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}} dx}{2c\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2(A(5-2m)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}}}{\frac{16a^2c}{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}} \frac{1}{8af(c-c\sin(e+fx))^{5/2}}}$$

↓ 3042

$$\frac{\frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19))\cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{2c\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2(A(5-2m)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}}}{\frac{16a^2c}{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}} \frac{1}{8af(c-c\sin(e+fx))^{5/2}}} +$$

↓ 3146

$$\frac{a^3(A(4m^2-8m+3)+C(4m^2+24m+19))\cos(e+fx)\int\frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}d(a\sin(e+fx))}{a-a\sin(e+fx)}}{2cf\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}+\frac{a^2(A(5-2m)-C(2m+11))\cos(e+fx)(a\sin(e+fx))}{f(c-c\sin(e+fx))^{3/2}}$$

$$\frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}}$$

↓ 78

$$\frac{a^2(A(4m^2-8m+3)+C(4m^2+24m+19))\cos(e+fx)(a\sin(e+fx)+a)^m\text{Hypergeometric2F1}\left(1,m+\frac{1}{2},m+\frac{3}{2},\frac{\sin(e+fx)a+a}{2a}\right)}{2cf(2m+1)\sqrt{c-c\sin(e+fx)}}+\frac{a^2(A(5-2m)-C(2m+11))\cos(e+fx)(a\sin(e+fx))}{f(c-c\sin(e+fx))^{3/2}}$$

$$\frac{(A+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}}\frac{16a^2c}{16a^2c}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(5/2),x]
```

output

```
((A + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(8*a*f*(c - c*Sin[e + f*x])^(5/2)) + ((a^2*(A*(5 - 2*m) - C*(11 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(c - c*Sin[e + f*x])^(3/2)) + (a^2*(A*(3 - 8*m + 4*m^2) + C*(19 + 24*m + 4*m^2))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(2*c*f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(16*a^2*c)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

rule 3224 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3451 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(a*f*(2*m + 1))), x] + Simp[(a*B*(m - n) + A*b*(m + n + 1))/(a*b*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (ILtQ[m + n, 0] && !SumSimplerQ[n, 1])) && NeQ[2*m + 1, 0]`

rule 3515 `Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*sin[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(a*A + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - C*(c^2*m - d^2*(n + 1)) + d*(A*c*(m + n + 2) - c*C*(3*m - n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] && NeQ[2*m + 1, 0]))`

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin(fx + e)^2)}{(c - c \sin(fx + e))^{\frac{5}{2}}} dx$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)`

output `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)`

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{\frac{5}{2}}} dx = \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{\frac{5}{2}}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(3*c^3*cos(f*x + e)^2 - 4*c^3 - (c^3*cos(f*x + e)^2 - 4*c^3)*sin(f*x + e)), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(5/2), x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(e + fx) + A)(a \sin(e + fx) + a)^m}{(-c \sin(e + fx) + c)^{5/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x,
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) +
c)^(5/2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x,
algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Error index.cc index_gcd Error: Bad
Argument Value`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin^2(e + fx) + A)(a + a \sin(e + fx))^m}{(c - c \sin(e + fx))^{5/2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(
5/2),x)`

output

```
int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c - c*sin(e + f*x))^(5/2), x)
```

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx =$$

$$\frac{\sqrt{c} \left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2}{\sin(fx + e)^3 - 3 \sin(fx + e)^2 + 3 \sin(fx + e) - 1} dx \right) c + \left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1}}{\sin(fx + e)^3 - 3 \sin(fx + e)^2 + 3 \sin(fx + e) - 1} dx \right) a}{c^3}$$

input

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

output

```
( - sqrt(c)*(int(((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**2)/(sin(e + f*x)**3 - 3*sin(e + f*x)**2 + 3*sin(e + f*x) - 1),x)*c + int(((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1))/(sin(e + f*x)**3 - 3*sin(e + f*x)**2 + 3*sin(e + f*x) - 1),x)*a))/c**3
```

$$3.7 \quad \int \frac{A+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$

Optimal result	93
Mathematica [A] (verified)	94
Rubi [A] (verified)	94
Maple [B] (verified)	97
Fricas [F]	98
Sympy [F]	99
Maxima [F]	99
Giac [F(-2)]	99
Mupad [F(-1)]	100
Reduce [F]	100

Optimal result

Integrand size = 42, antiderivative size = 167

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + C) \cos(e + fx) \sqrt{a + a \sin(e + fx)}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{(A - 3C) \cos(e + fx) \log(1 - \sin(e + fx))}{4cf \sqrt{a + a \sin(e + fx)} \sqrt{c - c \sin(e + fx)}} + \frac{(A + C) \cos(e + fx) \log(1 + \sin(e + fx))}{4cf \sqrt{a + a \sin(e + fx)} \sqrt{c - c \sin(e + fx)}}$$

output

```
1/4*(A+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1/2)/a/f/(c-c*sin(f*x+e))^(3/2)-1/4
*(A-3*C)*cos(f*x+e)*ln(1-sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f
*x+e))^(1/2)+1/4*(A+C)*cos(f*x+e)*ln(1+sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1
/2)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 4.78 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.14

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{\left(A + C - (A - 3C) \log\left(\cos\left(\frac{1}{2}(e + fx)\right) - \sin\left(\frac{1}{2}(e + fx)\right)\right)\right)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}}$$

input

```
Integrate[(A + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]
```

output

```
((A + C - (A - 3*C)*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2 + (A + C)*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]))/(2*f*Sqrt[a*(1 + Sin[e + f*x])])*(c - c*Sin[e + f*x])^(3/2))
```

Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 3515, 27, 3042, 3448, 3042, 3216, 3042, 3146, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a \sin(e + fx) + a}(c - c \sin(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + C \sin(e + fx)^2}{\sqrt{a \sin(e + fx) + a}(c - c \sin(e + fx))^{3/2}} dx$$

↓ 3515

$$\frac{(A + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{\int -\frac{2(a^2(A - C) - 2a^2C \sin(e + fx))}{\sqrt{\sin(e + fx)a + a}\sqrt{c - c \sin(e + fx)}} dx}{4a^2c}$$

↓ 27

$$\begin{aligned}
& \frac{\int \frac{a^2(A-C)-2a^2C \sin(e+fx)}{\sqrt{\sin(e+fx)a+a\sqrt{c-c\sin(e+fx)}}} dx}{2a^2c} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(A-C)-2a^2C \sin(e+fx)}{\sqrt{\sin(e+fx)a+a\sqrt{c-c\sin(e+fx)}}} dx}{2a^2c} + \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3448} \\
& \frac{a^2(A+C) \int \frac{\sqrt{c-c\sin(e+fx)}}{\sqrt{\sin(e+fx)a+a}} dx}{2c} + \frac{\frac{1}{2}a(A-3C) \int \frac{\sqrt{\sin(e+fx)a+a}}{\sqrt{c-c\sin(e+fx)}} dx}{2a^2c} + \\
& \quad \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a^2(A+C) \int \frac{\sqrt{c-c\sin(e+fx)}}{\sqrt{\sin(e+fx)a+a}} dx}{2c} + \frac{\frac{1}{2}a(A-3C) \int \frac{\sqrt{\sin(e+fx)a+a}}{\sqrt{c-c\sin(e+fx)}} dx}{2a^2c} + \\
& \quad \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3216} \\
& \frac{a^3(A+C) \cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2c(A-3C) \cos(e+fx) \int \frac{\cos(e+fx)}{c-c\sin(e+fx)} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \quad \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a^3(A+C) \cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2c(A-3C) \cos(e+fx) \int \frac{\cos(e+fx)}{c-c\sin(e+fx)} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \quad \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3146} \\
& \frac{a^2(A+C) \cos(e+fx) \int \frac{1}{\sin(e+fx)a+a} d(a \sin(e+fx))}{2f\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \frac{a^2(A-3C) \cos(e+fx) \int \frac{1}{c-c\sin(e+fx)} d(-c\sin(e+fx))}{2f\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \quad \frac{(A+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{16}
\end{aligned}$$

$$\frac{a^2(A+C)\cos(e+fx)\log(a\sin(e+fx)+a)}{2f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \frac{a^2(A-3C)\cos(e+fx)\log(c-c\sin(e+fx))}{2f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{2a^2c}{(A+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} \frac{1}{4af(c-c\sin(e+fx))^{3/2}}$$

input `Int[(A + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]`

output `((A + C)*Cos[e + f*x]*Sqrt[a + a*Sin[e + f*x]]/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + ((a^2*(A + C)*Cos[e + f*x]*Log[a + a*Sin[e + f*x]]/(2*f*Sqrt[a + a*Sin[e + f*x]]*Sqrt[c - c*Sin[e + f*x]]) - (a^2*(A - 3*C)*Cos[e + f*x]*Log[c - c*Sin[e + f*x]]/(2*f*Sqrt[a + a*Sin[e + f*x]]*Sqrt[c - c*Sin[e + f*x]])))/(2*a^2*c)`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2, x], x, b*Sin[e + f*x], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

rule 3216

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_)
+ (f_)*(x_)]], x_Symbol] :> Simp[a*c*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]
])*Sqrt[c + d*Sin[e + f*x]]) Int[Cos[e + f*x]/(c + d*Sin[e + f*x]), x],
x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
]
```

rule 3448

```
Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) +
(f_)*(x_)])*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] :> Simp
[(A*b + a*B)/(2*a*b) Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]
], x], x] + Simp[(B*c + A*d)/(2*c*d) Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a
+ b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 - b^2, 0]
```

rule 3515

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] :>
Simp[(a*A + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m+1)*(c + d*Sin[e + f*x])^(
n + 1)/(2*b*c*f*(2*m + 1)), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*
Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m
+ n + 2)) - C*(c^2*m - d^2*(n + 1)) + d*(A*c*(m + n + 2) - c*C*(3*m - n))*
Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ
[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2,
0] && NeQ[2*m + 1, 0]))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 587 vs. $2(149) = 298$.

Time = 6.96 (sec) , antiderivative size = 588, normalized size of antiderivative = 3.52

method	result
parts	$\frac{A\sqrt{4} \left(2 \ln \left(-\cot \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right) + \csc \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right) \right) \cos \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right)^2 - 2 \ln \left(-\cot \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right) + \csc \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right) - 1 \right) \cos \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right)}{8f \sqrt{a \sin \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right)^2} \sqrt{c \cos \left(\frac{\pi}{4} + \frac{fx}{2} + \frac{e}{2} \right)}}$

input

```
int((A+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x,met
hod=_RETURNVERBOSE)
```

output

```

1/8*A/f*4^(1/2)*(2*ln(-cot(1/4*Pi+1/2*f*x+1/2*e)+csc(1/4*Pi+1/2*f*x+1/2*e)
)*cos(1/4*Pi+1/2*f*x+1/2*e)^2-2*ln(-cot(1/4*Pi+1/2*f*x+1/2*e)+csc(1/4*Pi+1
/2*f*x+1/2*e)-1)*cos(1/4*Pi+1/2*f*x+1/2*e)^2-2*ln(-cot(1/4*Pi+1/2*f*x+1/2*
e)+csc(1/4*Pi+1/2*f*x+1/2*e)+1)*cos(1/4*Pi+1/2*f*x+1/2*e)^2+sin(1/4*Pi+1/2
*f*x+1/2*e)^2)/(a*sin(1/4*Pi+1/2*f*x+1/2*e)^2)^(1/2)/(c*cos(1/4*Pi+1/2*f*x
+1/2*e)^2)^(1/2)/c*tan(1/4*Pi+1/2*f*x+1/2*e)+1/2*C/c/f*((6*cos(1/2*f*x+1/2
*e)^2-3)*sin(1/2*f*x+1/2*e)-3*cos(1/2*f*x+1/2*e)*(-1+2*cos(1/2*f*x+1/2*e)
^2))*ln(2*(sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e))/(cos(1/2*f*x+1/2*e)+1))+
((-1+2*cos(1/2*f*x+1/2*e)^2)*sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e)*(-1+2*c
os(1/2*f*x+1/2*e)^2))*ln(-2*(cos(1/2*f*x+1/2*e)+sin(1/2*f*x+1/2*e))/(cos(1
/2*f*x+1/2*e)+1))+((-8*cos(1/2*f*x+1/2*e)^2+4)*sin(1/2*f*x+1/2*e)+4*cos(1/
2*f*x+1/2*e)*(-1+2*cos(1/2*f*x+1/2*e)^2))*ln(2/(cos(1/2*f*x+1/2*e)+1))+cos
(1/2*f*x+1/2*e)*(-2*sin(1/2*f*x+1/2*e)*cos(1/2*f*x+1/2*e)-2*sin(1/2*f*x+1/
2*e)^2))/(sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e))/(2*sin(1/2*f*x+1/2*e)*c
os(1/2*f*x+1/2*e)+1)*a^(1/2)/(-2*sin(1/2*f*x+1/2*e)*cos(1/2*f*x+1/2*e)-1)
*c)^(1/2)

```

Fricas [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin^2(fx + e) + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input

```

integrate((A+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2)
,x, algorithm="fricas")

```

output

```

integral((C*cos(f*x + e)^2 - A - C)*sqrt(a*sin(f*x + e) + a)*sqrt(-c*sin(f
*x + e) + c)/(a*c^2*cos(f*x + e)^2*sin(f*x + e) - a*c^2*cos(f*x + e)^2), x
)

```

Sympy [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{A + C \sin^2(e + fx)}{\sqrt{a(\sin(e + fx) + 1)}(-c(\sin(e + fx) - 1))^{3/2}} dx$$

input `integrate((A+C*sin(f*x+e)**2)/(a+a*sin(f*x+e))**(1/2)/(c-c*sin(f*x+e))**(3/2),x)`

output `Integral((A + C*sin(e + f*x)**2)/(sqrt(a*(sin(e + f*x) + 1))*(-c*(sin(e + f*x) - 1))**(3/2)), x)`

Maxima [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin^2(fx + e) + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((A+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)/(sqrt(a*sin(f*x + e) + a)*(-c*sin(f*x + e) + c)^(3/2)), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(e + fx)^2 + A}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx$$

input `int((A + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c - c*sin(e + f*x))
^(3/2)),x)`

output `int((A + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c - c*sin(e + f*x))
^(3/2)), x)`

Reduce [F]

$$\int \frac{A + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{\sqrt{c} \sqrt{a} \left(\int \frac{\sqrt{\sin(fx+e)+1} \sqrt{-\sin(fx+e)+1} \sin(fx+e)^2}{\sin(fx+e)^3 - \sin(fx+e)^2 - \sin(fx+e) + 1} dx \right) c + \left(\right)}{a c^2}$$

input `int((A+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x)`

output `(sqrt(c)*sqrt(a)*(int((sqrt(sin(e + f*x) + 1)*sqrt(- sin(e + f*x) + 1)*si
n(e + f*x)**2)/(sin(e + f*x)**3 - sin(e + f*x)**2 - sin(e + f*x) + 1),x)*c
+ int((sqrt(sin(e + f*x) + 1)*sqrt(- sin(e + f*x) + 1))/(sin(e + f*x)**3
- sin(e + f*x)**2 - sin(e + f*x) + 1),x)*a))/(a*c**2)`

3.8 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^n (A + C \sin^2(e +$

Optimal result	101
Mathematica [F]	102
Rubi [A] (verified)	102
Maple [F]	106
Fricas [F]	106
Sympy [F]	107
Maxima [F]	107
Giac [F]	108
Mupad [F(-1)]	108
Reduce [F]	109

Optimal result

Integrand size = 38, antiderivative size = 256

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= -\frac{C(1 + 2m) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^n}{f(1 + m + n)(2 + m + n)}$$

$$+ \frac{2^{\frac{1}{2}+n}(C(1 + 2m)(m - n) + (1 + m + n)(C(1 - m + n) + A(2 + m + n))) \cos(e + fx) \text{Hypergeometr}}{f(1 + m + n)}$$

$$+ \frac{C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{1+n}}{cf(2 + m + n)}$$

output

```
-C*(1+2*m)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n/f/(1+m+n)/(2+m
+n)+2^(1/2+n)*(C*(1+2*m)*(m-n)+(1+m+n)*(C*(1-m+n)+A*(2+m+n)))*cos(f*x+e)*h
ypergeom([1/2-n, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(1-sin(f*x+e))^(-1/2-n
)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n/f/(1+2*m)/(1+m+n)/(2+m+n)+C*cos(f*
x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1+n)/c/f/(2+m+n)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2), x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2), x]
```

Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.20, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 3519, 25, 3042, 3452, 3042, 3224, 3042, 3168, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c - c \sin(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c - c \sin(e + fx))^n dx$$

$$\downarrow \text{3519}$$

$$\frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}{cf(m + n + 2)}$$

$$\frac{\int -(\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + acC(2m + 1) \sin(e + fx)) dx}{ac(m + n + 2)}$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n (ac(C(-m+n+1)+A(m+n+2))+acC(2m+1)\sin(e+fx)) dx}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{n+1}}{cf(m+n+2)}} \frac{ac(m+n+2)}{cf(m+n+2)}$$

↓ 3042

$$\frac{\int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n (ac(C(-m+n+1)+A(m+n+2))+acC(2m+1)\sin(e+fx)) dx}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{n+1}}{cf(m+n+2)}} \frac{ac(m+n+2)}{cf(m+n+2)}$$

↓ 3452

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n))}{m+n+1} \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n dx - \frac{acC(2m+1) \cos(e+fx)(a \sin(e+fx))}{f(m+n+1)}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{n+1}}{cf(m+n+2)}} \frac{ac(m+n+2)}{cf(m+n+2)}$$

↓ 3042

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n))}{m+n+1} \int (\sin(e+fx)a+a)^m (c-c\sin(e+fx))^n dx - \frac{acC(2m+1) \cos(e+fx)(a \sin(e+fx))}{f(m+n+1)}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{n+1}}{cf(m+n+2)}} \frac{ac(m+n+2)}{cf(m+n+2)}$$

↓ 3224

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^m \int \cos^{2m}(e+fx)(c-c\sin(e+fx)) dx}{m+n+1}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{n+1}}{cf(m+n+2)}} \frac{ac(m+n+2)}{cf(m+n+2)}$$

↓ 3042

$$\frac{\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^m \int \cos(e+fx)^{2m}(c-c\sin(e+fx)) dx}{m+n+1}}{\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c\sin(e+fx))^{n+1}}{cf(m+n+2)}} \frac{ac(m+n+2)}{cf(m+n+2)}$$

↓ 3168

$$\frac{ac^3((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{\frac{1}{2}(-2m-1)+m}(c \sin(e+fx)+c)^{\frac{1}{2}(-2m-1)+m}}{f(m+n+1)}$$

$ac(m+n)$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

↓ 80

$$\frac{ac^32^{n-\frac{1}{2}}((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos(e+fx)(1-\sin(e+fx))^{\frac{1}{2}-n}(a \sin(e+fx)+a)^m(c \sin(e+fx)+c)^{\frac{1}{2}(-2m-1)+m}}{f(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

↓ 79

$$\frac{ac^22^{n+\frac{1}{2}}((m+n+1)(A(m+n+2)+C(-m+n+1))+C(2m+1)(m-n)) \cos(e+fx)(1-\sin(e+fx))^{\frac{1}{2}-n}(a \sin(e+fx)+a)^m(c \sin(e+fx)+c)^{\frac{1}{2}(-2m-1)+m}}{f(2m+1)(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2),x]
```

output

```
(C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(1 + n))/(c*f*(2 + m + n)) + (-((a*c*C*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n)/(f*(1 + m + n))) + (2^(1/2 + n)*a*c^2*(C*(1 + 2*m)*(m - n) + (1 + m + n)*(C*(1 - m + n) + A*(2 + m + n)))*Cos[e + f*x]*Hypergeometric2F1[(1 + 2*m)/2, (1 - 2*n)/2, (3 + 2*m)/2, (1 + Sin[e + f*x])/2]*(1 - Sin[e + f*x])^(1/2 - n)*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1/2 + (-1 - 2*m)/2 + m + n)*(c + c*Sin[e + f*x])^((-1 - 2*m)/2 + (1 + 2*m)/2))/(f*(1 + 2*m)*(1 + m + n)))/(a*c*(2 + m + n))
```

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 79 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 80 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3168 `Int[(cos[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[a^2*((g*Cos[e + f*x])^(p + 1)/(f*g*(a + b*Sin[e + f*x])^(p + 1)/2)*(a - b*Sin[e + f*x])^(p + 1)/2)) Subst[Int[(a + b*x)^(m + (p - 1)/2)*(a - b*x)^(p - 1)/2), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, g, m, p}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m]`
- rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_))*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3452

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(
a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d,
e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m,
-2^(-1)] && NeQ[m + n + 1, 0]
```

rule 3519

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] :=
Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1
))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*
x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1
)) - b*c*C*(2*m + 1)*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A
, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1
)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^n (A + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + A) (a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx \end{aligned}$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algo
rithm="fricas")
```

output `integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

Sympy [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (a(\sin(e + fx) + 1))^m (-c(\sin(e + fx) - 1))^n (A + C \sin^2(e + fx)) dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**n*(A+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(-c*(sin(e + f*x) - 1))**n*(A + C*sin(e + f*x)**2), x)`

Maxima [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(e + fx) + A) (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n,x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n,x)`

Reduce [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \left(\int (a + a \sin(fx + e))^m (-\sin(fx + e)c + c)^n \sin(fx + e)^2 dx \right) c \\ &+ \left(\int (a + a \sin(fx + e))^m (-\sin(fx + e)c + c)^n dx \right) a \end{aligned}$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

output

```
int((sin(e + f*x)*a + a)**m*( - sin(e + f*x)*c + c)**n*sin(e + f*x)**2,x)*
c + int((sin(e + f*x)*a + a)**m*( - sin(e + f*x)*c + c)**n,x)*a
```

3.9 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^n (A + C \sin^2(e +$

Optimal result	110
Mathematica [F]	111
Rubi [A] (warning: unable to verify)	111
Maple [F]	115
Fricas [F]	115
Sympy [F(-1)]	116
Maxima [F]	116
Giac [F]	117
Mupad [F(-1)]	117
Reduce [F]	118

Optimal result

Integrand size = 37, antiderivative size = 358

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= -\frac{C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{1+n}}{df(2 + m + n)}$$

$$-\frac{2^{\frac{1}{2}+m} a(c(C + 2Cm) + d(C(1 - m + n) + A(2 + m + n))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right)}{df(2 + m + n)}$$

$$-\frac{2^{\frac{3}{2}+m} C(dm - c(1 + m)) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right), \frac{d(1 - \sin(e + fx))}{c + d}}{df(2 + m + n)} \cos(e + fx)$$

output

```
-C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1+n)/d/f/(2+m+n)-2^(1/2+m)*a*(c*(2*C*m+C)+d*(C*(1-m+n)+A*(2+m+n)))*AppellF1(1/2,-n,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*(c+d*sin(f*x+e))^n/d/f/(2+m+n)/(((c+d*sin(f*x+e))/(c+d))^n)-2^(3/2+m)*C*(d*m-c*(1+m))*AppellF1(1/2,-n,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n/d/f/(2+m+n)/(((c+d*sin(f*x+e))/(c+d))^n)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2), x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.10 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.270$, Rules used = {3042, 3525, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c + d \sin(e + fx))^n dx$$

↓ 3525

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n (a(Ad(m + n + 2) + C(nd + d + cm)) + aC(dm - c(m + 1)) \sin(e + fx)) dx}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n (a(Ad(m + n + 2) + C(nd + d + cm)) + aC(dm - c(m + 1)) \sin(e + fx)) dx}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 3466

$$\frac{a(Ad(m + n + 2) + c(2Cm + C) + Cd(-m + n + 1)) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n dx + C(dm - c(m + 1)) \sin(e + fx)}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 3042

$$\frac{a(Ad(m + n + 2) + c(2Cm + C) + Cd(-m + n + 1)) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n dx + C(dm - c(m + 1)) \sin(e + fx)}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 3267

$$\frac{a^3 \cos(e + fx) (Ad(m + n + 2) + c(2Cm + C) + Cd(-m + n + 1)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^n d \sin(e + fx)}{\sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + \frac{a^2 C (dm - c(m + 1)) \cos(e + fx)}{f \sqrt{a - a \sin(e + fx)}}}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 157

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx) (Ad(m + n + 2) + c(2Cm + C) + Cd(-m + n + 1)) \int \frac{\sqrt{2} (\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^n d \sin(e + fx)}{\sqrt{2} f (a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} + \frac{a^2 C (dm - c(m + 1)) \cos(e + fx)}{f \sqrt{1 - \sin(e + fx)}}}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 27

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx) (Ad(m + n + 2) + c(2Cm + C) + Cd(-m + n + 1)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^n d \sin(e + fx)}{\sqrt{1 - \sin(e + fx)}} + \frac{a^2 C (dm - c(m + 1)) \cos(e + fx)}{f (a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}}}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1))(c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \left(\frac{c}{c-d} + d \sin(e+fx)\right)^{n-1}}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

↓ 155

$$\frac{\sqrt{2a^2} \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx)+a)^m (Ad(m+n+2)+c(2Cm+C)+Cd(-m+n+1))(c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \text{AppellF1}}{f(2m+1)(a-a \sin(e+fx))}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + C*Sin[e + f*x]^2),x]
```

output

```
-((C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(1 + n))/(d*f*(2 + m + n))) + ((Sqrt[2]*a^2*(c*(C + 2*C*m) + C*d*(1 - m + n) + A*d*(2 + m + n))*AppellF1[1/2 + m, 1/2, -n, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n)/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n) + (Sqrt[2]*a*C*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, -n, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*(c + d*Sin[e + f*x])^n)/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n))/(a*d*(2 + m + n))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] :> Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Sim[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Sim[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]`

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)/b Int[(a + b*SIN[e + f*x])^m*(c + d*SIN[e + f*x])^n, x
] + Simp[B/b Int[(a + b*SIN[e + f*x])^(m + 1)*(c + d*SIN[e + f*x])^n,
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ
[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3525

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2), x_Symbol] :=
Simp[(-C)*Cos[e + f*x]*(a + b*SIN[e + f*x])^m*((c + d*SIN[e + f*x])^(n + 1
))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*SIN[e + f*
x])^m*(c + d*SIN[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1
)) + C*(a*d*m - b*c*(m + 1))*SIN[e + f*x], x], x] /; FreeQ[{a, b, c, d,
e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2
- d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^n (A + C \sin(fx + e)^2) dx$$

```
input int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

```
output int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin(fx + e)^2 + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx \end{aligned}$$

```
input integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algo
rithm="fricas")
```

output `integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**n*(A+C*sin(f*x+e)**2),x)`

output Timed out

Maxima [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorith="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(e + fx) + A) (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n,x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n,x)`

Reduce [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + C \sin^2(e + fx)) dx \\ &= \left(\int (\sin(fx + e) d + c)^n (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) c \\ & \quad + \left(\int (\sin(fx + e) d + c)^n (a + a \sin(fx + e))^m dx \right) a \end{aligned}$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+C*sin(f*x+e)^2),x)
```

output

```
int((sin(e + f*x)*d + c)**n*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2,x)*c +
int((sin(e + f*x)*d + c)**n*(sin(e + f*x)*a + a)**m,x)*a
```

3.10 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A + C \sin(e+fx)) dx$

Optimal result	119
Mathematica [F]	120
Rubi [A] (warning: unable to verify)	120
Maple [F]	125
Fricas [F]	125
Sympy [F(-1)]	126
Maxima [F]	126
Giac [F]	126
Mupad [F(-1)]	127
Reduce [F]	127

Optimal result

Integrand size = 41, antiderivative size = 382

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \frac{(c^2 C + A d^2) \cos(e + fx) (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-1-m}}{d (c^2 - d^2) f (1 + m)}$$

$$\frac{2^{\frac{1}{2}+m} a (c(A + C) d (1 + m) + d^2 (C - A m + C m) - c^2 (C + 2 C m)) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2} - m, 1 + m, \frac{3}{2}, \frac{1}{2} (1 - \sin(e + fx))\right)}{(c - d) d (c + d)}$$

$$\frac{2^{\frac{3}{2}+m} C \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, 1 + m, \frac{3}{2}, \frac{1}{2} (1 - \sin(e + fx)), \frac{d(1 - \sin(e + fx))}{c + d}\right) \cos(e + fx) (1 + \sin(e + fx))}{d (c + d) f}$$

output

```
(A*d^2+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1-m)/d/(c^2-d^2)/f/(1+m)-2^(1/2+m)*a*(c*(A+C)*d*(1+m)+d^2*(-A*m+C*m+C)-c^2*(2*C*m+C))*cos(f*x+e)*hypergeom([1/2, 1/2-m],[3/2],(c-d)*(1-sin(f*x+e))/(2*c+2*d*sin(f*x+e)))*(a+a*sin(f*x+e))^(1-m)*((c+d)*(1+sin(f*x+e))/(c+d*sin(f*x+e)))^(1/2-m)/(c-d)/d/(c+d)^2/f/(1+m)/((c+d*sin(f*x+e))^m)-2^(3/2+m)*C*AppellF1(1/2,1+m,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^m*((c+d*sin(f*x+e))/(c+d))^m/d/(c+d)/f/((c+d*sin(f*x+e))^m)
```


Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + C*Sin[
e + f*x]^2),x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + C*Sin[
e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.35 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.11, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.293$, Rules used = {3042, 3523, 25, 3042, 3466, 3042, 3267, 142, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)^2) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3523}$$

$$\frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m+1)(c^2 - d^2)} -$$

$$\frac{\int -(\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + cC(md + d - cm)) + aC(c^2 - d^2)(m - ad(m+1)(c^2 - d^2))}{ad(m+1)(c^2 - d^2)}$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + cC(md + d - cm)) + aC(c^2 - d^2) (m + 1) \frac{ad(m + 1)(c^2 - d^2)}{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}}{df(m + 1)(c^2 - d^2)} \downarrow 3042$$

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + cC(md + d - cm)) + aC(c^2 - d^2) (m + 1) \frac{ad(m + 1)(c^2 - d^2)}{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}}{df(m + 1)(c^2 - d^2)} \downarrow 3466$$

$$\frac{a(cd(m + 1)(A + C) + d^2(-Am + Cm + C) - (c^2(2Cm + C))) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} \frac{ad(m + 1)(c^2 - d^2)}{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}}{df(m + 1)(c^2 - d^2)} \downarrow 3042$$

$$\frac{a(cd(m + 1)(A + C) + d^2(-Am + Cm + C) - (c^2(2Cm + C))) \int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} \frac{ad(m + 1)(c^2 - d^2)}{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}}{df(m + 1)(c^2 - d^2)} \downarrow 3267$$

$$\frac{a^3 \cos(e + fx)(cd(m + 1)(A + C) + d^2(-Am + Cm + C) - (c^2(2Cm + C))) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} (c + d \sin(e + fx))^{-m - 1}}{\sqrt{a - a \sin(e + fx)}} d \sin(e + fx)}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + \frac{a^2 C(m + 1)(c^2 - d^2)}{ad(m + 1)(c^2 - d^2)} \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m - 1}}{df(m + 1)(c^2 - d^2)} \downarrow 142$$

$$\frac{a^2 C(m + 1)(c^2 - d^2) \cos(e + fx) \int \frac{(\sin(e + fx)a + a)^{m + \frac{1}{2}} (c + d \sin(e + fx))^{-m - 1}}{\sqrt{a - a \sin(e + fx)}} d \sin(e + fx)}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} - \frac{a^2 2^{m + \frac{1}{2}} \cos(e + fx)(cd(m + 1)(A + C) + d^2(-Am + Cm + C))}{ad(m + 1)(c^2 - d^2)} \frac{(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m - 1}}{df(m + 1)(c^2 - d^2)}$$

↓ 157

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{\sqrt{2}(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{\sqrt{2}f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(cd(m+1)(A+C))}{ad}$$

$$\frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 27

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(cd(m+1)(A+C))}{ad}$$

$$\frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 156

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}\left(\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}\right)^{-m-1}d\sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(c-d)(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(cd(m+1)(A+C))}{ad}$$

$$\frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 155

$$\frac{\sqrt{2}aC(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(a\sin(e+fx)+a)^{m+1}(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\text{AppellF1}\left(m+\frac{3}{2},\frac{1}{2},m+1,m+\frac{5}{2},\frac{1}{2}\right)}{f(2m+3)(c-d)(a-a\sin(e+fx))} - \frac{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(cd(m+1)(A+C))}{ad}$$

$$\frac{(Ad^2+c^2C)\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

```
input Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + C*Sin[e + f*x]^2), x]
```

output

```
((c^2*C + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^
(-1 - m))/(d*(c^2 - d^2)*f*(1 + m)) + (-((2^(1/2 + m)*a^2*(c*(A + C)*d*(1
+ m) + d^2*(C - A*m + C*m) - c^2*(C + 2*C*m))*Cos[e + f*x]*Hypergeometric2
F1[1/2, 1/2 - m, 3/2, ((c - d)*(1 - Sin[e + f*x]))/(2*(c + d*Sin[e + f*x])
])* (a + a*Sin[e + f*x])^(-1 + m)*(((c + d)*(1 + Sin[e + f*x]))/(c + d*Sin[
e + f*x]))^(1/2 - m))/((c + d)*f*(c + d*Sin[e + f*x])^m)) + (Sqrt[2]*a*C*(
c^2 - d^2)*(1 + m)*AppellF1[3/2 + m, 1/2, 1 + m, 5/2 + m, (1 + Sin[e + f*x
])/2, -((d*(1 + Sin[e + f*x]))/(c - d))*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x
]]*(a + a*Sin[e + f*x])^(1 + m)*((c + d*Sin[e + f*x])/(c - d))^m)/((c - d)
*f*(3 + 2*m)*(a - a*Sin[e + f*x])*(c + d*Sin[e + f*x])^m))/(a*d*(c^2 - d^2
)*(1 + m))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 142

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((b*e
- a*f)*(m + 1)))*Hypergeometric2F1[m + 1, -n, m + 2, -(d*e - c*f)*((a +
b*x)/((b*c - a*d)*(e + f*x)))]/((b*e - a*f)*((c + d*x)/((b*c - a*d)*(e + f
*x))))^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m + n + p + 2,
0] && !IntegerQ[n]
```

rule 155

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*
Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/
(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f,
m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simp
lify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simpl
ify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d
*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c
- e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 156

```
Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]
)*(b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

rule 157

```
Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)
^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]
)*(b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c
- a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !Si
mplerQ[e + f*x, a + b*x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_))*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Ssin[e
+ f*x]]*Sqrt[a - b*Ssin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*
x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m
, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] &&
!IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)/b Int[(a + b*Ssin[e + f*x])^m*(c + d*Ssin[e + f*x])^n, x], x
] + Simp[B/b Int[(a + b*Ssin[e + f*x])^(m + 1)*(c + d*Ssin[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ
[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3523

```

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] :=
Simp[(-(c^2*C + A*d^2))*Cos[e + f*x]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e +
f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 -
d^2)) Int[(a + b*Ssin[e + f*x])^m*(c + d*Ssin[e + f*x])^(n + 1)*Simp[A*d*(a
*d*m + b*c*(n + 1)) + c*C*(a*c*m + b*d*(n + 1)) - b*(A*d^2*(m + n + 2) + C*
(c^2*(m + 1) + d^2*(n + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d,
e, f, A, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 -
d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])

```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{-2-m} (A + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x,
algorithm="fricas")
```

output

```
integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e)
+ c)^(-m - 2), x)
```

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(-2-m)*(A+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x,algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^(-m - 2), x)`

Giac [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx \\ &= \int (C \sin^2(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx \end{aligned}$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x,algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^(-m - 2), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{(c + d \sin(e + fx))^{m+2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(m + 2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(m + 2), x)`

Reduce [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + C \sin^2(e + fx)) dx$$

$$= \left(\int \frac{(a + a \sin(fx + e))^m}{(\sin(fx + e)d + c)^m \sin^2(fx + e) d^2 + 2(\sin(fx + e)d + c)^m \sin(fx + e) cd + (\sin(fx + e)d + c)^m} dx \right)$$

$$+ \left(\int \frac{(a + a \sin(fx + e))^m \sin^2(fx + e)}{(\sin(fx + e)d + c)^m \sin^2(fx + e) d^2 + 2(\sin(fx + e)d + c)^m \sin(fx + e) cd + (\sin(fx + e)d + c)^m} dx \right)$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+C*sin(f*x+e)^2),x)`

output `int((sin(e + f*x)*a + a)**m/((sin(e + f*x)*d + c)**m*sin(e + f*x)**2*d**2 + 2*(sin(e + f*x)*d + c)**m*sin(e + f*x)*c*d + (sin(e + f*x)*d + c)**m*c**2),x)*a + int(((sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/((sin(e + f*x)*d + c)**m*sin(e + f*x)**2*d**2 + 2*(sin(e + f*x)*d + c)**m*sin(e + f*x)*c*d + (sin(e + f*x)*d + c)**m*c**2),x)*c`

3.11 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A + C \sin^2(e+fx)) dx$

Optimal result	128
Mathematica [F]	129
Rubi [A] (warning: unable to verify)	129
Maple [F]	133
Fricas [F]	134
Sympy [F(-1)]	134
Maxima [F]	134
Giac [F]	135
Mupad [F(-1)]	135
Reduce [F]	136

Optimal result

Integrand size = 39, antiderivative size = 372

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx =$$

$$\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{5/2}}{df(7 + 2m)}$$

$$- \frac{2^{\frac{1}{2}+m} a(c + d)(2c(C + 2Cm) + d(C(5 - 2m) + A(7 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, -\frac{3}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right)}{df(7 + 2m) \sqrt{c + d}}$$

$$- \frac{2^{\frac{5}{2}+m} C(c + d)(dm - c(1 + m)) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, -\frac{3}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right), \frac{d(1 - \sin(e + fx))}{c + d}}{df(7 + 2m) \sqrt{\frac{c + d \sin(e + fx)}{c + d}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(5/2)/d/f/(7+2*m)-2^(1/2+m)*a*(c+d)*(2*c*(2*C*m+C)+d*(C*(5-2*m)+A*(7+2*m)))*AppellF1(1/2,-3/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*(c+d*sin(f*x+e))^(1/2)/d/f/(7+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)-2^(5/2+m)*C*(c+d)*(d*m-c*(1+m))*AppellF1(1/2,-3/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(7+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2), x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.22 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.09, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$, Rules used = {3042, 3525, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) (c + d \sin(e + fx))^{3/2} dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) (c + d \sin(e + fx))^{3/2} dx$$

↓ 3525

$$\frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx)) dx}{\frac{ad(2m + 7)}{df(2m + 7)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}}$$

↓ 27

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} (a(Ad(2m+7)+C(5d+2cm))+2aC(dm-c(m+1))\sin(e+fx))}{\frac{ad(2m+7)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{5/2}} df(2m+7)} \downarrow 3042$$

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} (a(Ad(2m+7)+C(5d+2cm))+2aC(dm-c(m+1))\sin(e+fx))}{\frac{ad(2m+7)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{5/2}} df(2m+7)} \downarrow 3466$$

$$\frac{a(Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} dx + 2C(dm-c(m-1))\sin(e+fx)}{\frac{ad(2m+7)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{5/2}} df(2m+7)} \downarrow 3042$$

$$\frac{a(Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} dx + 2C(dm-c(m-1))\sin(e+fx)}{\frac{ad(2m+7)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{5/2}} df(2m+7)} \downarrow 3267$$

$$\frac{a^3 \cos(e+fx)(Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^{3/2}}{\sqrt{a-a\sin(e+fx)}} d\sin(e+fx)}{\frac{ad(2m+7)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{5/2}} df(2m+7)} + \frac{2a^2C(dm-c(m+1))\cos(e+fx)}{f\sqrt{a-a\sin(e+fx)}} \downarrow 157$$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^{3/2}}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{\frac{ad(2m+7)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{5/2}} df(2m+7)} + \frac{\sqrt{2}a^2C(dm-c(m+1))\cos(e+fx)}{f\sqrt{a-a\sin(e+fx)}}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d \sin(e+fx))^{3/2} d \sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2a^2 C(dm-c(m+2))}{ad(2m+7)}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 156

$$\frac{a^3 (c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) \sqrt{c+d \sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)^{3/2} d \sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 155

$$\frac{\sqrt{2} a^2 (c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+7)+2c(2Cm+C)+Cd(5-2m)) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, -\frac{3}{2}, m+\frac{3}{2}\right)}{f(2m+1) (a-a \sin(e+fx)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + C*Sin[e + f*x]^2), x]
```

output

```
(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(5/2))/(d*f*(7 + 2*m)) + ((Sqrt[2]*a^2*(c - d)*(C*d*(5 - 2*m) + A*d*(7 + 2*m) + 2*c*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, -3/2, 3/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)]]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (2*Sqrt[2]*a*C*(c - d)*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, -3/2, 5/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)]]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(7 + 2*m))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3525

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{\frac{3}{2}} (A + C \sin(fx + e))^2 dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A) (d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,
algorithm="fricas")`

output `integral(-(C*c*cos(f*x + e)^2 - (A + C)*c + (C*d*cos(f*x + e)^2 - (A + C)*
d)*sin(f*x + e))*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(3/2)*(A+C*sin(f*x+e)**2),
x)`

output `Timed out`

Maxima [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + A) (d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(e + fx) + A) (d \sin(e + fx) + c)^{3/2} (a \sin(e + fx) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x,algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + C \sin^2(e + fx)) dx = \int (C \sin^2(e + fx) + A) (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/2),x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/2), x)`

Reduce [F]

$$\begin{aligned}
& \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A \\
& + C \sin^2(e + fx)) dx = \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^3 dx \right) cd \\
& + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) c^2 \\
& + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e) dx \right) ad \\
& + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m dx \right) ac
\end{aligned}$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+C*sin(f*x+e)^2),x)`

output `int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**3,x)*c*d + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2,x)*c**2 + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x),x)*a*d + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m,x)*a*c`

3.12 $\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+C \sin^2(e+fx)) dx$

Optimal result	137
Mathematica [F]	138
Rubi [A] (warning: unable to verify)	138
Maple [F]	142
Fricas [F]	143
Sympy [F]	143
Maxima [F]	144
Giac [F]	144
Mupad [F(-1)]	145
Reduce [F]	145

Optimal result

Integrand size = 39, antiderivative size = 366

$$\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+C \sin^2(e+fx)) dx$$

$$= -\frac{2C \cos(e+fx)(a+a \sin(e+fx))^m(c+d \sin(e+fx))^{3/2}}{df(5+2m)}$$

$$-\frac{2^{\frac{1}{2}+m}a(2c(C+2Cm)+d(C(3-2m)+A(5+2m))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}-m, -\frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1-\sin(e+fx))\right)}{df(5+2m)\sqrt{c+d \sin(e+fx)}}$$

$$-\frac{2^{\frac{5}{2}+m}C(dm-c(1+m)) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2}-m, -\frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1-\sin(e+fx))\right) \cos(e+fx)}{df(5+2m)\sqrt{\frac{c+d \sin(e+fx)}{c+d}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)/d/f/(5+2*m)-2^(1/2+m)*a*(2*c*(2*C*m+C)+d*(C*(3-2*m)+A*(5+2*m)))*AppellF1(1/2,-1/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*(c+d*sin(f*x+e))^(1/2)/d/f/(5+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)-2^(5/2+m)*C*(d*m-c*(1+m))*AppellF1(1/2,-1/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(5+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2), x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.18 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$, Rules used = {3042, 3525, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + C \sin^2(e + fx)) \sqrt{c + d \sin(e + fx)} dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2) \sqrt{c + d \sin(e + fx)} dx$$

↓ 3525

$$\frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + 2aC(dm - c(m + 1)) \sin(e + fx))}{\frac{ad(2m + 5)}{df(2m + 5)} (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}$$

↓ 27

$$\frac{\int (\sin(e+fx)a+a)^m \sqrt{c+d\sin(e+fx)} (a(Ad(2m+5)+C(3d+2cm))+2aC(dm-c(m+1))\sin(e+fx)) dx}{\frac{ad(2m+5)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{3/2}} df(2m+5)}$$

↓ 3042

$$\frac{\int (\sin(e+fx)a+a)^m \sqrt{c+d\sin(e+fx)} (a(Ad(2m+5)+C(3d+2cm))+2aC(dm-c(m+1))\sin(e+fx)) dx}{\frac{ad(2m+5)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{3/2}} df(2m+5)}$$

↓ 3466

$$\frac{a(Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \int (\sin(e+fx)a+a)^m \sqrt{c+d\sin(e+fx)} dx + 2C(dm-c(m+1))}{\frac{ad(2m+5)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{3/2}} df(2m+5)}$$

↓ 3042

$$\frac{a(Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \int (\sin(e+fx)a+a)^m \sqrt{c+d\sin(e+fx)} dx + 2C(dm-c(m+1))}{\frac{ad(2m+5)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{3/2}} df(2m+5)}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{c+d\sin(e+fx)}}{\sqrt{a-a\sin(e+fx)}} d\sin(e+fx) + \frac{2a^2C(dm-c(m+1))\cos(e+fx) \int (\sin(e+fx)a+a)^m \sqrt{c+d\sin(e+fx)} dx}{f\sqrt{a-a\sin(e+fx)}\sqrt{a\sin(e+fx)+a}}}{\frac{ad(2m+5)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{3/2}} df(2m+5)}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{c+d\sin(e+fx)}}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx) + \frac{\sqrt{2}a^2C(dm-c(m+1))\cos(e+fx) \int (\sin(e+fx)a+a)^m \sqrt{c+d\sin(e+fx)} dx}{\sqrt{2}f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}}{\frac{ad(2m+5)}{2C\cos(e+fx)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{3/2}} df(2m+5)}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{c+d \sin(e+fx)}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2a^2 C(dm-c(m+1))}{ad(2m+5)}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{3/2}}{df(2m+5)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) \sqrt{c+d \sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{\frac{c+d \sin(e+fx)}{c-d}}} + \frac{2a^2 C(dm-c(m+1))}{ad(2m+5)}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{3/2}}{df(2m+5)}$$

↓ 155

$$\frac{\sqrt{2a^2} \sqrt{1-\sin(e+fx)} \cos(e+fx) (Ad(2m+5)+2c(2Cm+C)+Cd(3-2m)) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d \sin(e+fx))^{3/2}}{df(2m+5)}$$

input `Int[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + C*Sin[e + f*x]^2), x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2))/(d*f*(5 + 2*m)) + ((Sqrt[2]*a^2*(C*d*(3 - 2*m) + A*d*(5 + 2*m) + 2*c*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, -1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (2*Sqrt[2]*a*C*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, -1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(5 + 2*m))`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3525

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c + d \sin(fx + e)} (A + C \sin(fx + e))^2 dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,
algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x +
e) + a)^m, x)`

Sympy [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx)) \sqrt{c + d \sin(e + fx)} dx$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(1/2)*(A+C*sin(f*x+e)**2),
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)*sqrt(c + d*sin(
e + f*x)), x)`

Maxima [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e)
+ a)^m, x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x,
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e)
+ a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} dx$$

input `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/2),x)`

output `int((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/2), x)`

Reduce [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + C \sin^2(e + fx)) dx$$

$$= \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) c$$

$$+ \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m dx \right) a$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+C*sin(f*x+e)^2),x)`

output `int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2,x)*c + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m,x)*a`

3.13
$$\int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$$

Optimal result	146
Mathematica [F]	147
Rubi [A] (warning: unable to verify)	147
Maple [F]	151
Fricas [F]	152
Sympy [F]	152
Maxima [F]	153
Giac [F]	153
Mupad [F(-1)]	154
Reduce [F]	154

Optimal result

Integrand size = 39, antiderivative size = 364

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= -\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)}}{df(3 + 2m)}$$

$$-\frac{2^{\frac{1}{2}+m} a(2c(C + 2Cm) + d(C - 2Cm + A(3 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx)), \frac{d(1 - \sin(e + fx))}{c+d}\right)}{df(3 + 2m) \sqrt{c + d \sin(e + fx)}}$$

$$-\frac{2^{\frac{5}{2}+m} C(dm - c(1 + m)) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx)), \frac{d(1 - \sin(e + fx))}{c+d}\right) \cos(e + fx)(1 - \sin(e + fx))}{df(3 + 2m) \sqrt{c + d \sin(e + fx)}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(3+2*m)-2^(1/2+m)*a*(2*c*(2*C*m+C)+d*(C-2*C*m+A*(3+2*m)))*AppellF1(1/2,1/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/f/(3+2*m)/(c+d*sin(f*x+e))^(1/2)-2^(5/2+m)*C*(d*m-c*(1+m))*AppellF1(1/2,1/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/f/(3+2*m)/(c+d*sin(f*x+e))^(1/2)
```

Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]],x]
```

output

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]], x]
```

Rubi [A] (warning: unable to verify)

Time = 1.18 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.09, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$, Rules used = {3042, 3525, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin(e + fx)^2)}{\sqrt{c + d \sin(e + fx)}} dx$$

$$\downarrow \text{3525}$$

$$2 \int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+2aC(dm-c(m+1)) \sin(e+fx))}{2\sqrt{c+d \sin(e+fx)}} dx$$

$$\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx) + a)^m \sqrt{c+d \sin(e+fx)}}$$

$$\frac{df(2m+3)}{\downarrow \text{27}}$$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+2aC(dm-c(m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}} - \frac{df(2m+3)}{df(2m+3)}$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+2aC(dm-c(m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}} - \frac{df(2m+3)}{df(2m+3)}$$

↓ 3466

$$\frac{a(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + 2C(dm-c(m+1)) \int \frac{(\sin(e+fx)a+a)^{m+1}}{\sqrt{c+d \sin(e+fx)}} dx}{\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}} - \frac{df(2m+3)}{df(2m+3)}$$

↓ 3042

$$\frac{a(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + 2C(dm-c(m+1)) \int \frac{(\sin(e+fx)a+a)^{m+1}}{\sqrt{c+d \sin(e+fx)}} dx}{\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}} - \frac{df(2m+3)}{df(2m+3)}$$

↓ 3267

$$\frac{\frac{a^3 \cos(e+fx)(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} + \frac{2a^2 C(dm-c(m+1)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}}}{\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}} - \frac{df(2m+3)}{df(2m+3)}$$

↓ 157

$$\frac{\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{\sqrt{2}a^2 C(dm-c(m+1)) \sqrt{1-\sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}}{\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}} - \frac{df(2m+3)}{df(2m+3)}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2a^2 C(dm-c(m+1)) \sqrt{1-\sin(e+fx)}}{ad(2m+3)}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (d(A(2m+3)-2Cm+C)+2c(2Cm+C)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{c+d \sin(e+fx)}} + \frac{2a^2 C(dm-c(m+1)) \sqrt{1-\sin(e+fx)}}{ad(2m+3)}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

↓ 155

$$\frac{\sqrt{2} a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (d(A(2m+3)-2Cm+C)+2c(2Cm+C)) (a \sin(e+fx)+a)^m \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}(\sin(e+fx))\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{c+d \sin(e+fx)}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]],x]
```

output

```
(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(d*f*(3 + 2*m)) + ((Sqrt[2]*a^2*(2*c*(C + 2*C*m) + d*(C - 2*C*m + A*(3 + 2*m)))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/(f*(1 + 2*m)*(a - a*Sin[e + f*x]) *Sqrt[c + d*Sin[e + f*x]]) + (2*Sqrt[2]*a*C*(d*m - c*(1 + m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]])/(a*d*(3 + 2*m))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*b*((e + f*x)/(b*e - a*f))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*b*((c + d*x)/(b*c - a*d))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3525

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1))/(d*f*(m + n + 2)), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + C*(a*d*m - b*c*(m + 1))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin(fx + e))^2}{\sqrt{c + d \sin(fx + e)}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)
```


Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x,
algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - A - C)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x
+ e) + c), x)`

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(1/2),
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/sqrt(c + d*sin(
e + f*x)), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x,
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e)
) + c), x)`

Giac [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x,
algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e)
) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(e + fx)^2 + A) (a + a \sin(e + fx))^m}{\sqrt{c + d \sin(e + fx)}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(1/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(1/2),x)`

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2}{\sin(fx + e) d + c} dx \right) c$$

$$+ \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m}{\sin(fx + e) d + c} dx \right) a$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)`

output `int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/(sin(e + f*x)*d + c),x)*c + int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m)/(sin(e + f*x)*d + c),x)*a`

$$3.14 \quad \int \frac{(a+a \sin(e+fx))^m (A+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$$

Optimal result	155
Mathematica [F]	156
Rubi [A] (warning: unable to verify)	156
Maple [F]	160
Fricas [F]	161
Sympy [F]	161
Maxima [F]	161
Giac [F]	162
Mupad [F(-1)]	162
Reduce [F]	163

Optimal result

Integrand size = 39, antiderivative size = 405

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \frac{2(c^2 C + A d^2) \cos(e + fx) (a + a \sin(e + fx))^m}{d(c^2 - d^2) f \sqrt{c + d \sin(e + fx)}} \\ - \frac{2^{\frac{1}{2}+m} a (c(A + C)d - d^2(A - C + 4Am) - 2c^2(C + 2Cm)) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right)}{d(c^2 - d^2) f \sqrt{c + d \sin(e + fx)}} \\ - \frac{2^{\frac{3}{2}+m} (2c^2 C(1 + m) + d^2(A - C + 2Am)) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right), \frac{d(1 - \sin(e + fx))}{c + d}}{d(c^2 - d^2) f \sqrt{c + d \sin(e + fx)}}$$

output

```
2*(A*d^2+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^(1/2)-2^(1/2+m)*a*(c*(A+C)*d-d^2*(4*A*m+A-C)-2*c^2*(2*C*m+C))*AppellF1(1/2,1/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^(1/2)-2^(3/2+m)*(2*c^2*C*(1+m)+d^2*(2*A*m+A-C))*AppellF1(1/2,1/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^(1/2)
```

Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2), x]
```

output

```
Integrate[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.29 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.282$, Rules used = {3042, 3523, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a \sin(e + fx) + a)^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

↓ 3523

$$\frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{df(c^2 - d^2) \sqrt{c + d \sin(e + fx)}} -$$

$$2 \int - \frac{(\sin(e + fx)a + a)^m (a(cC(d - 2cm) + Ad(c - 2dm)) + a(2C(m + 1)c^2 + d^2(2mA + A - C)) \sin(e + fx))}{2\sqrt{c + d \sin(e + fx)}} dx$$

$$ad(c^2 - d^2)$$

↓ 27

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(cC(d-2cm)+Ad(c-2dm))+a(2C(m+1)c^2+d^2(2mA+A-C)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(cC(d-2cm)+Ad(c-2dm))+a(2C(m+1)c^2+d^2(2mA+A-C)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3466

$$\frac{a(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (d^2(2Am+A-C)+2c^2C(m+1))}{ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3042

$$\frac{a(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (d^2(2Am+A-C)+2c^2C(m+1))}{ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} + \frac{a^2 \cos(e+fx)(d^2(2Am+A-C)+2c^2C(m+1))}{f \sqrt{a-a \sin(e+fx)}}}{ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(cd(A+C)-d^2(4Am+A-C)-2c^2(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{\sqrt{2}f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx)(d^2(2Am+A-C)+2c^2C(m+1))}{f \sqrt{a-a \sin(e+fx)}}}{ad(c^2-d^2)} + \frac{2(Ad^2+c^2C) \cos(e+fx)(a \sin(e+fx)+a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C))}{ad(c^2 - d^2)}$$

$$\frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{df(c^2 - d^2) \sqrt{c + d \sin(e + fx)}}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{c+d \sin(e+fx)}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C))}{ad(c^2 - d^2)}$$

$$\frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{df(c^2 - d^2) \sqrt{c + d \sin(e + fx)}}$$

↓ 155

$$\frac{\sqrt{2}a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (cd(A+C) - d^2(4Am+A-C) - 2c^2(2Cm+C)) (a \sin(e+fx)+a)^m \sqrt{\frac{c+d \sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, m+\frac{3}{2}, \frac{1}{2}\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{c+d \sin(e+fx)}}$$

$$\frac{2(Ad^2 + c^2C) \cos(e + fx)(a \sin(e + fx) + a)^m}{df(c^2 - d^2) \sqrt{c + d \sin(e + fx)}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2),x]`

output `(2*(c^2*C + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(d*(c^2 - d^2)*f*Sqrt[c + d*Sin[e + f*x]]) + ((Sqrt[2]*a^2*(c*(A + C)*d - d^2*(A - C + 4*A*m) - 2*c^2*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) + (Sqrt[2]*a*(2*c^2*C*(1 + m) + d^2*(A - C + 2*A*m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(c^2 - d^2))`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3523

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*sin[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-(c^2*C + A*d^2))*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + c*C*(a*c*m + b*d*(n + 1)) - b*(A*d^2*(m + n + 2) + C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])
```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + C \sin(fx + e)^2)}{(c + d \sin(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x,
algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x +
e) + a)^m/(d^2*cos(f*x + e)^2 - 2*c*d*sin(f*x + e) - c^2 - d^2), x)`

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(3/2),
x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + C*sin(e + f*x)**2)/(c + d*sin(e +
f*x))**(3/2), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x,
algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) + c)^(3/2), x)`

Giac [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(fx + e) + A)(a \sin(fx + e) + a)^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + A)*(a*sin(f*x + e) + a)^m/(d*sin(f*x + e) + c)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin^2(e + fx) + A) (a + a \sin(e + fx))^m}{(c + d \sin(e + fx))^{3/2}} dx$$

input `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(3/2),x)`

output `int(((A + C*sin(e + f*x)^2)*(a + a*sin(e + f*x))^m)/(c + d*sin(e + f*x))^(3/2), x)`

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)}{\sin(fx + e)^2 d^2 + 2 \sin(fx + e) cd + c^2} dx \right) a$$

$$+ \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m}{\sin(fx + e)^2 d^2 + 2 \sin(fx + e) cd + c^2} dx \right) a$$

input `int((a+a*sin(f*x+e))^m*(A+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)`

output `int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/(sin(e + f*x)**2*d**2 + 2*sin(e + f*x)*c*d + c**2),x)*c + int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m)/(sin(e + f*x)**2*d**2 + 2*sin(e + f*x)*c*d + c**2),x)*a`

3.15
$$\int \frac{A+B \sin(e+fx)+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx$$

Optimal result	164
Mathematica [A] (verified)	165
Rubi [A] (verified)	165
Maple [B] (verified)	168
Fricas [F]	169
Sympy [F]	170
Maxima [F]	170
Giac [F(-2)]	170
Mupad [F(-1)]	171
Reduce [F]	171

Optimal result

Integrand size = 50, antiderivative size = 174

$$\int \frac{A+B \sin(e+fx)+C \sin^2(e+fx)}{\sqrt{a+a \sin(e+fx)}(c-c \sin(e+fx))^{3/2}} dx = \frac{(A+B+C) \cos(e+fx) \sqrt{a+a \sin(e+fx)}}{4af(c-c \sin(e+fx))^{3/2}} - \frac{(A-B-3C) \cos(e+fx) \log(1-\sin(e+fx))}{4cf \sqrt{a+a \sin(e+fx)} \sqrt{c-c \sin(e+fx)}} + \frac{(A-B+C) \cos(e+fx) \log(1+\sin(e+fx))}{4cf \sqrt{a+a \sin(e+fx)} \sqrt{c-c \sin(e+fx)}}$$

output

```
1/4*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1/2)/a/f/(c-c*sin(f*x+e))^(3/2)-1/4*(A-B-3*C)*cos(f*x+e)*ln(1-sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(1/2)+1/4*(A-B+C)*cos(f*x+e)*ln(1+sin(f*x+e))/c/f/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 4.96 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.13

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + B + C + (-A + B + 3C) \log(\cos(\frac{1}{2}(e + fx))) - \dots}{\dots}$$

input

```
Integrate[(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]
```

output

```
((A + B + C + (-A + B + 3*C)*Log[Cos[(e + f*x)/2] - Sin[(e + f*x)/2]]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2 + (A - B + C)*Log[Cos[(e + f*x)/2] + Sin[(e + f*x)/2]]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^2*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2]))/(2*f*Sqrt[a*(1 + Sin[e + f*x])]*(c - c*Sin[e + f*x])^(3/2))
```

Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3514, 27, 3042, 3448, 3042, 3216, 3042, 3146, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a \sin(e + fx) + a}(c - c \sin(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \sin(e + fx) + C \sin(e + fx)^2}{\sqrt{a \sin(e + fx) + a}(c - c \sin(e + fx))^{3/2}} dx$$

↓ 3514

$$\frac{(A + B + C) \cos(e + fx) \sqrt{a \sin(e + fx) + a}}{4af(c - c \sin(e + fx))^{3/2}} - \frac{\int -\frac{2(a^2(A - B - C) - 2a^2C \sin(e + fx))}{\sqrt{\sin(e + fx)a + a}\sqrt{c - c \sin(e + fx)}} dx}{4a^2c}$$

↓ 27

$$\begin{aligned}
& \frac{\int \frac{a^2(A-B-C)-2a^2C \sin(e+fx)}{\sqrt{\sin(e+fx)a+a}\sqrt{c-c\sin(e+fx)}} dx}{2a^2c} + \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(A-B-C)-2a^2C \sin(e+fx)}{\sqrt{\sin(e+fx)a+a}\sqrt{c-c\sin(e+fx)}} dx}{2a^2c} + \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3448} \\
& \frac{a^2(A-B+C) \int \frac{\sqrt{c-c\sin(e+fx)}}{\sqrt{\sin(e+fx)a+a}} dx}{2c} + \frac{\frac{1}{2}a(A-B-3C) \int \frac{\sqrt{\sin(e+fx)a+a}}{\sqrt{c-c\sin(e+fx)}} dx}{2a^2c} + \\
& \quad \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a^2(A-B+C) \int \frac{\sqrt{c-c\sin(e+fx)}}{\sqrt{\sin(e+fx)a+a}} dx}{2c} + \frac{\frac{1}{2}a(A-B-3C) \int \frac{\sqrt{\sin(e+fx)a+a}}{\sqrt{c-c\sin(e+fx)}} dx}{2a^2c} + \\
& \quad \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3216} \\
& \frac{a^3(A-B+C) \cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2c(A-B-3C) \cos(e+fx) \int \frac{\cos(e+fx)}{c-c\sin(e+fx)} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \quad \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a^3(A-B+C) \cos(e+fx) \int \frac{\cos(e+fx)}{\sin(e+fx)a+a} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \frac{a^2c(A-B-3C) \cos(e+fx) \int \frac{\cos(e+fx)}{c-c\sin(e+fx)} dx}{2\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \quad \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{3146} \\
& \frac{a^2(A-B+C) \cos(e+fx) \int \frac{1}{\sin(e+fx)a+a} d(a \sin(e+fx))}{2f\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \frac{a^2(A-B-3C) \cos(e+fx) \int \frac{1}{c-c\sin(e+fx)} d(-c \sin(e+fx))}{2f\sqrt{a \sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} + \\
& \quad \frac{(A+B+C) \cos(e+fx) \sqrt{a \sin(e+fx) + a}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow \text{16}
\end{aligned}$$

$$\frac{a^2(A-B+C)\cos(e+fx)\log(a\sin(e+fx)+a)}{2f\sqrt{a\sin(e+fx)+a}\sqrt{c-\sin(e+fx)}} - \frac{a^2(A-B-3C)\cos(e+fx)\log(c-\sin(e+fx))}{2f\sqrt{a\sin(e+fx)+a}\sqrt{c-\sin(e+fx)}} + \frac{2a^2c}{(A+B+C)\cos(e+fx)\sqrt{a\sin(e+fx)+a}} \frac{1}{4af(c-\sin(e+fx))^{3/2}}$$

input

```
Int[(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2)/(Sqrt[a + a*Sin[e + f*x]]*(c - c*Sin[e + f*x])^(3/2)),x]
```

output

```
((A + B + C)*Cos[e + f*x]*Sqrt[a + a*Sin[e + f*x]]/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + ((a^2*(A - B + C)*Cos[e + f*x]*Log[a + a*Sin[e + f*x]]/(2*f*Sqrt[a + a*Sin[e + f*x]]*Sqrt[c - c*Sin[e + f*x]]) - (a^2*(A - B - 3*C)*Cos[e + f*x]*Log[c - c*Sin[e + f*x]]/(2*f*Sqrt[a + a*Sin[e + f*x]]*Sqrt[c - c*Sin[e + f*x]]))/(2*a^2*c))
```

Defintions of rubi rules used

rule 16

```
Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3146

```
Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2, x], x, b*Sin[e + f*x], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])
```


rule 3216

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] :> Simp[a*c*(Cos[e + f*x]/(Sqrt[a + b*Sin[e + f*x]]*Sqrt[c + d*Sin[e + f*x]])) Int[Cos[e + f*x]/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
```

rule 3448

```
Int[((A_) + (B_)*sin[(e_) + (f_)*(x_)])/(Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]]), x_Symbol] :> Simp[(A*b + a*B)/(2*a*b) Int[Sqrt[a + b*Sin[e + f*x]]/Sqrt[c + d*Sin[e + f*x]], x], x] + Simp[(B*c + A*d)/(2*c*d) Int[Sqrt[c + d*Sin[e + f*x]]/Sqrt[a + b*Sin[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0]
```

rule 3514

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_) + (f_)*(x_)])^2), x_Symbol] :> Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Simp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] && NeQ[2*m + 1, 0]))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 896 vs. $2(156) = 312$.

Time = 13.92 (sec) , antiderivative size = 897, normalized size of antiderivative = 5.16

method	result	size
parts	Expression too large to display	897

input

```
int((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

output

```

-1/8*A/f*4^(1/2)*(2*ln(-cot(1/4*Pi+1/2*f*x+1/2*e)+csc(1/4*Pi+1/2*f*x+1/2*
e)+1)*cos(1/4*Pi+1/2*f*x+1/2*e)^2+2*ln(-cot(1/4*Pi+1/2*f*x+1/2*e)+csc(1/4*P
i+1/2*f*x+1/2*e)-1)*cos(1/4*Pi+1/2*f*x+1/2*e)^2-2*ln(-cot(1/4*Pi+1/2*f*x+1
/2*e)+csc(1/4*Pi+1/2*f*x+1/2*e))*cos(1/4*Pi+1/2*f*x+1/2*e)^2-sin(1/4*Pi+1/
2*f*x+1/2*e)^2)/(a*sin(1/4*Pi+1/2*f*x+1/2*e)^2)^(1/2)/(c*cos(1/4*Pi+1/2*f*
x+1/2*e)^2)^(1/2)/c*tan(1/4*Pi+1/2*f*x+1/2*e)-1/2*B/c/f*((-1+2*cos(1/2*f*
x+1/2*e)^2)*sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e)*(-1+2*cos(1/2*f*x+1/2*
e)^2))*ln(-2*(cos(1/2*f*x+1/2*e)+sin(1/2*f*x+1/2*e))/(cos(1/2*f*x+1/2*e)+1))
+((1-2*cos(1/2*f*x+1/2*e)^2)*sin(1/2*f*x+1/2*e)+cos(1/2*f*x+1/2*e)*(-1+2*
cos(1/2*f*x+1/2*e)^2))*ln(2*(sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e))/(cos(1/
2*f*x+1/2*e)+1))+cos(1/2*f*x+1/2*e)*(2*sin(1/2*f*x+1/2*e)*cos(1/2*f*x+1/2*
e)+2*sin(1/2*f*x+1/2*e)^2))/(sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e))/((2*si
n(1/2*f*x+1/2*e)*cos(1/2*f*x+1/2*e)+1)*a)^(1/2)/(-(2*sin(1/2*f*x+1/2*e)*co
s(1/2*f*x+1/2*e)-1)*c)^(1/2)+1/2*C/c/f*((6*cos(1/2*f*x+1/2*e)^2-3)*sin(1/
2*f*x+1/2*e)-3*cos(1/2*f*x+1/2*e)*(-1+2*cos(1/2*f*x+1/2*e)^2))*ln(2*(sin(1
/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e))/(cos(1/2*f*x+1/2*e)+1))+((-1+2*cos(1/2*f
*x+1/2*e)^2)*sin(1/2*f*x+1/2*e)-cos(1/2*f*x+1/2*e)*(-1+2*cos(1/2*f*x+1/2*
e)^2))*ln(-2*(cos(1/2*f*x+1/2*e)+sin(1/2*f*x+1/2*e))/(cos(1/2*f*x+1/2*e)+1)
)+((-8*cos(1/2*f*x+1/2*e)^2+4)*sin(1/2*f*x+1/2*e)+4*cos(1/2*f*x+1/2*e)*(-1
+2*cos(1/2*f*x+1/2*e)^2))*ln(2/(cos(1/2*f*x+1/2*e)+1))+cos(1/2*f*x+1/2*...

```

Fricas [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(fx + e)^2 + B \sin(fx + e) + A}{\sqrt{a \sin(fx + e) + a(-c \sin(fx + e) + c)}^{3/2}} dx$$

input

```

integrate((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(
f*x+e))^(3/2),x, algorithm="fricas")

```

output

```

integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(a*sin(f*x + e) +
a)*sqrt(-c*sin(f*x + e) + c)/(a*c^2*cos(f*x + e)^2*sin(f*x + e) - a*c^2*c
os(f*x + e)^2), x)

```

Sympy [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a(\sin(e + fx) + 1)}(-c(\sin(e + fx) - 1))^{3/2}} dx$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(a+a*sin(f*x+e))**(1/2)/(c-c*sin(f*x+e))**(3/2),x)`

output `Integral((A + B*sin(e + f*x) + C*sin(e + f*x)**2)/(sqrt(a*(sin(e + f*x) + 1))*(-c*(sin(e + f*x) - 1))**(3/2)), x)`

Maxima [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin^2(fx + e) + B \sin(fx + e) + A}{\sqrt{a \sin(fx + e) + a}(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)/(sqrt(a*sin(f*x + e) + a))*(-c*sin(f*x + e) + c)^(3/2)), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \int \frac{C \sin(e + fx)^2 + B \sin(e + fx) + A}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx$$

input

```
int((A + B*sin(e + f*x) + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c
- c*sin(e + f*x))^(3/2)),x)
```

output

```
int((A + B*sin(e + f*x) + C*sin(e + f*x)^2)/((a + a*sin(e + f*x))^(1/2)*(c
- c*sin(e + f*x))^(3/2)), x)
```

Reduce [F]

$$\int \frac{A + B \sin(e + fx) + C \sin^2(e + fx)}{\sqrt{a + a \sin(e + fx)}(c - c \sin(e + fx))^{3/2}} dx = \frac{\sqrt{c} \sqrt{a} \left(\int \frac{\sqrt{\sin(fx+e)+1} \sqrt{-\sin(fx+e)+1} \sin(fx+e)^2}{\sin(fx+e)^3 - \sin(fx+e)^2 - \sin(fx+e) + 1} dx \right) c + \left(\int \frac{\sqrt{\sin(fx+e)+1} \sqrt{-\sin(fx+e)+1} \sin(fx+e)}{\sin(fx+e)^3 - \sin(fx+e)^2 - \sin(fx+e) + 1} dx \right) b + \int \frac{\sqrt{\sin(fx+e)+1} \sqrt{-\sin(fx+e)+1}}{\sin(fx+e)^3 - \sin(fx+e)^2 - \sin(fx+e) + 1} dx a}{(a*c**2)}$$

input

```
int((A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(a+a*sin(f*x+e))^(1/2)/(c-c*sin(f*x+e)
)^(3/2),x)
```

output

```
(sqrt(c)*sqrt(a)*(int((sqrt(sin(e + f*x) + 1)*sqrt(- sin(e + f*x) + 1)*si
n(e + f*x)**2)/(sin(e + f*x)**3 - sin(e + f*x)**2 - sin(e + f*x) + 1),x)*c
+ int((sqrt(sin(e + f*x) + 1)*sqrt(- sin(e + f*x) + 1)*sin(e + f*x))/(si
n(e + f*x)**3 - sin(e + f*x)**2 - sin(e + f*x) + 1),x)*b + int((sqrt(sin(e
+ f*x) + 1)*sqrt(- sin(e + f*x) + 1))/(sin(e + f*x)**3 - sin(e + f*x)**2
- sin(e + f*x) + 1),x)*a))/(a*c**2)
```

3.16 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^n (A + B \sin(e +$

Optimal result	172
Mathematica [F]	173
Rubi [A] (verified)	173
Maple [F]	177
Fricas [F]	178
Sympy [F(-1)]	178
Maxima [F]	178
Giac [F]	179
Mupad [F(-1)]	179
Reduce [F]	180

Optimal result

Integrand size = 46, antiderivative size = 268

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= -\frac{(C + 2Cm + B(2 + m + n)) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^n}{f(1 + m + n)(2 + m + n)}$$

$$+ \frac{2^{\frac{1}{2}+n}((1 + m + n)(C(1 - m + n) + A(2 + m + n)) + (m - n)(C + 2Cm + B(2 + m + n))) \cos(e + fx)}{f(1 + m + n)(2 + m + n)}$$

$$+ \frac{C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{1+n}}{cf(2 + m + n)}$$

output

```
-(C+2*C*m+B*(2+m+n))*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n/f/(1+m+n)/(2+m+n)+2^(1/2+n)*((1+m+n)*(C*(1-m+n)+A*(2+m+n))+(m-n)*(C+2*C*m+B*(2+m+n)))*cos(f*x+e)*hypergeom([1/2-n, 1/2+m], [3/2+m], 1/2+1/2*sin(f*x+e))*(1-sin(f*x+e))^(-1/2-n)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n/f/(1+2*m)/(1+m+n)/(2+m+n)+C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1+n)/c/f/(2+m+n)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + B*Sin[e + f*x]
] + C*Sin[e + f*x]^2), x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + B*Sin[e + f*x]
] + C*Sin[e + f*x]^2), x]
```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.19, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.239$, Rules used = {3042, 3518, 25, 3042, 3452, 3042, 3224, 3042, 3168, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3518}$$

$$\frac{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}{cf(m + n + 2)} -$$

$$\frac{\int -(\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + ac(2mC + C + B(m + n + 1))) dx}{ac(m + n + 2)}$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + ac(2mC + C + B(m + n + 2)))}{cf(m + n + 2)} \frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c - c \sin(e + fx))^n (ac(C(-m + n + 1) + A(m + n + 2)) + ac(2mC + C + B(m + n + 2)))}{cf(m + n + 2)} \frac{ac(m + n + 2)}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{n+1}}$$

↓ 3452

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))}{m+n+1} \frac{\int (\sin(e+fx)a+a)^m (c-c \sin(e+fx))^n dx}{cf(m+n+2)} \frac{ac(m+n+2)}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}} - \frac{ac(B(m+n+2)+2Cm)}{m+n+1}$$

↓ 3042

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))}{m+n+1} \frac{\int (\sin(e+fx)a+a)^m (c-c \sin(e+fx))^n dx}{cf(m+n+2)} \frac{ac(m+n+2)}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}} - \frac{ac(B(m+n+2)+2Cm)}{m+n+1}$$

↓ 3224

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))}{m+n+1} \frac{\cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^m \int \cos^{2m}(e+fx)}{cf(m+n+2)} \frac{ac(m+n+2)}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}} - \frac{ac(m+n+2)}{m+n+1}$$

↓ 3042

$$\frac{ac((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))}{m+n+1} \frac{\cos^{-2m}(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^m \int \cos(e+fx)^2}{cf(m+n+2)} \frac{ac(m+n+2)}{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}} - \frac{ac(m+n+2)}{m+n+1}$$

↓ 3168

$$\frac{ac^3 \cos(e+fx)((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{\frac{1}{2}(-2m-1)+m} (c \sin(e+fx))^{\frac{1}{2}(-2m-1)+m}}{f(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

↓ 80

$$\frac{ac^3 2^{n-\frac{1}{2}} \cos(e+fx)((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))(1-\sin(e+fx))^{\frac{1}{2}-n} (a \sin(e+fx)+a)^m (c \sin(e+fx))^{\frac{1}{2}-n}}{f(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

↓ 79

$$\frac{ac^2 2^{n+\frac{1}{2}} \cos(e+fx)((m+n+1)(A(m+n+2)+C(-m+n+1))+(m-n)(B(m+n+2)+2Cm+C))(1-\sin(e+fx))^{\frac{1}{2}-n} (a \sin(e+fx)+a)^m (c \sin(e+fx))^{\frac{1}{2}-n}}{f(2m+1)(m+n+1)}$$

$$\frac{C \cos(e+fx)(a \sin(e+fx)+a)^m (c-c \sin(e+fx))^{n+1}}{cf(m+n+2)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
(C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(1 + n))/(c*f*(2 + m + n)) + (-((a*c*(C + 2*C*m + B*(2 + m + n))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^n)/(f*(1 + m + n))) + (2^(1/2 + n)*a*c^2*((1 + m + n)*(C*(1 - m + n) + A*(2 + m + n)) + (m - n)*(C + 2*C*m + B*(2 + m + n)))*Cos[e + f*x]*Hypergeometric2F1[(1 + 2*m)/2, (1 - 2*n)/2, (3 + 2*m)/2, (1 + Sin[e + f*x])/2]*(1 - Sin[e + f*x])^(1/2 - n)*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1/2 + (-1 - 2*m)/2 + m + n)*(c + c*Sin[e + f*x])^((-1 - 2*m)/2 + (1 + 2*m)/2))/(f*(1 + 2*m)*(1 + m + n)))/(a*c*(2 + m + n))
```


Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 79 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 80 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3168 `Int[(cos[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[a^2*((g*Cos[e + f*x])^(p + 1)/(f*g*(a + b*Sin[e + f*x])^(p + 1)/2)*(a - b*Sin[e + f*x])^(p + 1)/2)) Subst[Int[(a + b*x)^(m + (p - 1)/2)*(a - b*x)^(p - 1)/2), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, g, m, p}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m]`
- rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_))*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3452

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(
a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d,
e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m,
-2^(-1)] && NeQ[m + n + 1, 0]
```

rule 3518

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_
) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x]
)^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m
+ n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*
Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] &&
EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n +
2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^n (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),
x)
```

output

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),
x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^n, x)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**n*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
-c*sin(f*x + e) + c)^n, x)
```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + B \sin(fx + e) + A) (a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^n dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+
e)^2),x, algorithm="giac")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
-c*sin(f*x + e) + c)^n, x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (C \sin^2(e + fx) + B \sin(e + fx) + A) dx$$

input

```
int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*
sin(e + f*x)^2),x)
```

output

```
int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*
sin(e + f*x)^2), x)
```

Reduce [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \left(\int (a + a \sin(fx + e))^m (-\sin(fx + e)c + c)^n \sin(fx + e)^2 dx \right) c$$

$$+ \left(\int (a + a \sin(fx + e))^m (-\sin(fx + e)c + c)^n \sin(fx + e) dx \right) b$$

$$+ \left(\int (a + a \sin(fx + e))^m (-\sin(fx + e)c + c)^n dx \right) a$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),
x)
```

output

```
int((sin(e + f*x)*a + a)**m*(- sin(e + f*x)*c + c)**n*sin(e + f*x)**2,x)*
c + int((sin(e + f*x)*a + a)**m*(- sin(e + f*x)*c + c)**n*sin(e + f*x),x)
*b + int((sin(e + f*x)*a + a)**m*(- sin(e + f*x)*c + c)**n,x)*a
```

3.17 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{5/2} (A + B \sin(e$

Optimal result	181
Mathematica [C] (verified)	182
Rubi [A] (verified)	183
Maple [F]	187
Fricas [B] (verification not implemented)	188
Sympy [F(-1)]	189
Maxima [B] (verification not implemented)	189
Giac [F]	190
Mupad [B] (verification not implemented)	191
Reduce [F]	192

Optimal result

Integrand size = 48, antiderivative size = 435

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{64c^3(B(45 - 8m - 4m^2) - C(39 - 16m + 4m^2) - A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))}{f(5 + 2m)(7 + 2m)(9 + 2m) (3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}}$$

$$\frac{16c^2(B(45 - 8m - 4m^2) - C(39 - 16m + 4m^2) - A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))}{f(7 + 2m)(9 + 2m) (15 + 16m + 4m^2)}$$

$$\frac{2c(B(45 - 8m - 4m^2) - C(39 - 16m + 4m^2) - A(63 + 32m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(9 + 2m)}$$

$$\frac{2(9B + 2C + 2Bm + 4Cm) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{f(7 + 2m)(9 + 2m)}$$

$$+ \frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{7/2}}{cf(9 + 2m)}$$

output

```
-64*c^3*(B*(-4*m^2-8*m+45)-C*(4*m^2-16*m+39)-A*(4*m^2+32*m+63))*cos(f*x+e)
*(a+a*sin(f*x+e))^m/f/(5+2*m)/(7+2*m)/(9+2*m)/(4*m^2+8*m+3)/(c-c*sin(f*x+e)
)^(1/2)-16*c^2*(B*(-4*m^2-8*m+45)-C*(4*m^2-16*m+39)-A*(4*m^2+32*m+63))*co
s(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)/f/(7+2*m)/(9+2*m)/(4*m^
2+16*m+15)-2*c*(B*(-4*m^2-8*m+45)-C*(4*m^2-16*m+39)-A*(4*m^2+32*m+63))*cos
(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(5+2*m)/(7+2*m)/(9+2*m)
)-2*(2*B*m+4*C*m+9*B+2*C)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(
5/2)/f/(7+2*m)/(9+2*m)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(
7/2)/c/f/(9+2*m)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 15.93 (sec) , antiderivative size = 1029, normalized size of antiderivative = 2.37

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + B*Sin[e +
f*x] + C*Sin[e + f*x]^2),x]
```

output

```

((a*(1 + Sin[e + f*x]))^m*(c - c*Sin[e + f*x])^(5/2)*(((18900*A - 14175*B
+ 12285*C + 15648*A*m - 4140*B*m + 648*C*m + 5280*A*m^2 - 832*B*m^2 + 1416
*C*m^2 + 896*A*m^3 - 208*B*m^3 + 224*C*m^3 + 64*A*m^4 - 16*B*m^4 + 16*C*m^
4)*((1/8 + I/8)*Cos[(e + f*x)/2] + (1/8 - I/8)*Sin[(e + f*x)/2]))/((1 + 2*
m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((18900*A - 14175*B + 12285*
C + 15648*A*m - 4140*B*m + 648*C*m + 5280*A*m^2 - 832*B*m^2 + 1416*C*m^2 +
896*A*m^3 - 208*B*m^3 + 224*C*m^3 + 64*A*m^4 - 16*B*m^4 + 16*C*m^4)*((1/8
- I/8)*Cos[(e + f*x)/2] + (1/8 + I/8)*Sin[(e + f*x)/2]))/((1 + 2*m)*(3 +
2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((3150*A - 3465*B + 3150*C + 2356*A*
m - 1706*B*m + 828*C*m + 584*A*m^2 - 316*B*m^2 + 200*C*m^2 + 48*A*m^3 - 24
*B*m^3 + 16*C*m^3)*((1/8 - I/8)*Cos[(3*(e + f*x))/2] - (1/8 + I/8)*Sin[(3*
(e + f*x))/2]))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((3150*A - 346
5*B + 3150*C + 2356*A*m - 1706*B*m + 828*C*m + 584*A*m^2 - 316*B*m^2 + 200
*C*m^2 + 48*A*m^3 - 24*B*m^3 + 16*C*m^3)*((1/8 + I/8)*Cos[(3*(e + f*x))/2]
- (1/8 - I/8)*Sin[(3*(e + f*x))/2]))/((3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(9 +
2*m)) + ((126*A - 315*B + 378*C + 64*A*m - 124*B*m + 88*C*m + 8*A*m^2 - 12
*B*m^2 + 8*C*m^2)*((-1/8 + I/8)*Cos[(5*(e + f*x))/2] - (1/8 + I/8)*Sin[(5*
(e + f*x))/2]))/((5 + 2*m)*(7 + 2*m)*(9 + 2*m)) + ((126*A - 315*B + 378*C
+ 64*A*m - 124*B*m + 88*C*m + 8*A*m^2 - 12*B*m^2 + 8*C*m^2)*((-1/8 - I/8)*
Cos[(5*(e + f*x))/2] - (1/8 - I/8)*Sin[(5*(e + f*x))/2]))/((5 + 2*m)*(7...

```

Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.77, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 3518, 27, 3042, 3452, 3042, 3219, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (c - c \sin(e + fx))^{5/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow 3518$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2}(a \sin(e + fx) + a)^m}{cf(2m + 9)} - \frac{2 \int -\frac{1}{2}(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{5/2}(ac(C(7 - 2m) + A(2m + 9)) + ac(2mB + 9B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 9)}$$

↓ 27

$$\frac{\int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{5/2}(ac(C(7 - 2m) + A(2m + 9)) + ac(2mB + 9B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 9)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2}(a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

↓ 3042

$$\frac{\int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{5/2}(ac(C(7 - 2m) + A(2m + 9)) + ac(2mB + 9B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 9)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2}(a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

↓ 3452

$$\frac{ac(-A(4m^2 + 32m + 63) + B(-4m^2 - 8m + 45) - C(4m^2 - 16m + 39)) \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{5/2} dx}{2m + 7} - \frac{2ac(2Bm + 9B + 4Cm + 2C)}{ac(2m + 9)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2}(a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

↓ 3042

$$\frac{ac(-A(4m^2 + 32m + 63) + B(-4m^2 - 8m + 45) - C(4m^2 - 16m + 39)) \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{5/2} dx}{2m + 7} - \frac{2ac(2Bm + 9B + 4Cm + 2C)}{ac(2m + 9)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2}(a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

↓ 3219

$$\frac{ac(-A(4m^2 + 32m + 63) + B(-4m^2 - 8m + 45) - C(4m^2 - 16m + 39)) \left(\frac{8c \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2} dx}{2m + 5} + \frac{2c \cos(e + fx)(c - c \sin(e + fx))^{3/2}}{f(2m + 5)} \right)}{2m + 7} - \frac{2ac(2Bm + 9B + 4Cm + 2C)}{ac(2m + 9)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{7/2}(a \sin(e + fx) + a)^m}{cf(2m + 9)}$$

↓ 3042

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45))-C(4m^2-16m+39)}{2m+7} \left(\frac{8c \int (\sin(e+fx)a+a)^m (c-c \sin(e+fx))^{3/2} dx}{2m+5} + \frac{2c \cos(e+fx)(c-c \sin(e+fx))^{3/2}}{f(2m+5)} \right)$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)} \quad ac(2m+9)$$

↓ 3219

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45))-C(4m^2-16m+39)}{2m+7} \left(\frac{8c \left(\frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c \sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} \right)$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)} \quad ac(2m+9)$$

↓ 3042

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45))-C(4m^2-16m+39)}{2m+7} \left(\frac{8c \left(\frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c \sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} \right)$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)} \quad ac(2m+9)$$

↓ 3217

$$\frac{ac(-A(4m^2+32m+63)+B(-4m^2-8m+45))-C(4m^2-16m+39)}{2m+7} \left(\frac{8c \left(\frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3) \sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)^m}{f(2m+3)} \right)}{2m+5} \right)$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{7/2}(a \sin(e+fx)+a)^m}{cf(2m+9)} \quad ac(2m+9)$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
(2*C*cos[e + f*x]*(a + a*sin[e + f*x])^m*(c - c*sin[e + f*x])^(7/2))/(c*f*(9 + 2*m)) + ((-2*a*c*(9*B + 2*C + 2*B*m + 4*C*m)*cos[e + f*x]*(a + a*sin[e + f*x])^m*(c - c*sin[e + f*x])^(5/2))/(f*(7 + 2*m)) - (a*c*(B*(45 - 8*m - 4*m^2) - C*(39 - 16*m + 4*m^2) - A*(63 + 32*m + 4*m^2))*((2*c*cos[e + f*x]*(a + a*sin[e + f*x])^m*(c - c*sin[e + f*x])^(3/2))/(f*(5 + 2*m)) + (8*c*((8*c^2*cos[e + f*x]*(a + a*sin[e + f*x])^m)/(f*(1 + 2*m)*(3 + 2*m)*sqrt[c - c*sin[e + f*x]]) + (2*c*cos[e + f*x]*(a + a*sin[e + f*x])^m*sqrt[c - c*sin[e + f*x]])/(f*(3 + 2*m))))/(5 + 2*m))/(7 + 2*m))/(a*c*(9 + 2*m))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3217

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[-2*b*cos[e + f*x]*((c + d*sin[e + f*x])^n/(f*(2*n + 1)*sqrt[a + b*sin[e + f*x]])), x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]
```

rule 3219

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*cos[e + f*x]*(a + b*sin[e + f*x])^(m - 1)*((c + d*sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n)) Int[(a + b*sin[e + f*x])^(m - 1)*(c + d*sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && IGtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])
```

rule 3452

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(
a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d,
e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m,
-2^(-1)] && NeQ[m + n + 1, 0]
```

rule 3518

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_
) + (f_)*(x_)]^2, x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x]
)^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m
+ n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*
Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] &&
EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n +
2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{\frac{5}{2}} (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)
^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)
^2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 937 vs. $2(419) = 838$.

Time = 0.17 (sec) , antiderivative size = 937, normalized size of antiderivative = 2.15

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")
```

output

```
2*((16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m + 105*C*c^2)*cos(f*x + e)^5 + 128*(A + B + C)*c^2*m^2 + (16*(B - C)*c^2*m^4 + 16*(9*B - 14*C)*c^2*m^3 + 8*(52*B - 97*C)*c^2*m^2 + 4*(111*B - 226*C)*c^2*m + 15*(9*B - 19*C)*c^2)*cos(f*x + e)^4 + 256*(4*A + B - 2*C)*c^2*m - (16*(A - 2*B + 3*C)*c^2*m^4 + 16*(10*A - 23*B + 32*C)*c^2*m^3 + 8*(65*A - 169*B + 253*C)*c^2*m^2 + 4*(150*A - 417*B + 656*C)*c^2*m + 3*(63*A - 180*B + 289*C)*c^2)*cos(f*x + e)^3 + 96*(21*A - 15*B + 13*C)*c^2 + (16*(A - B + C)*c^2*m^4 + 32*(7*A - 5*B + 7*C)*c^2*m^3 + 8*(133*A - 97*B + 85*C)*c^2*m^2 + 8*(233*A - 235*B + 233*C)*c^2*m + 3*(231*A - 255*B + 263*C)*c^2)*cos(f*x + e)^2 + 2*(16*(A - B + C)*c^2*m^4 + 192*(A - B + C)*c^2*m^3 + 8*(107*A - 99*B + 107*C)*c^2*m^2 + 16*(109*A - 89*B + 85*C)*c^2*m + 3*(483*A - 435*B + 419*C)*c^2)*cos(f*x + e) + (128*(A + B + C)*c^2*m^2 + (16*C*c^2*m^4 + 128*C*c^2*m^3 + 344*C*c^2*m^2 + 352*C*c^2*m + 105*C*c^2)*cos(f*x + e)^4 + 256*(4*A + B - 2*C)*c^2*m - (16*(B - 2*C)*c^2*m^4 + 16*(9*B - 22*C)*c^2*m^3 + 32*(13*B - 35*C)*c^2*m^2 + 4*(111*B - 314*C)*c^2*m + 15*(9*B - 26*C)*c^2)*cos(f*x + e)^3 + 96*(21*A - 15*B + 13*C)*c^2 - (16*(A - B + C)*c^2*m^4 + 32*(5*A - 7*B + 5*C)*c^2*m^3 + 8*(65*A - 117*B + 113*C)*c^2*m^2 + 24*(25*A - 51*B + 57*C)*c^2*m + 9*(21*A - 45*B + 53*C)*c^2)*cos(f*x + e)^2 - 2*(16*(A - B + C)*c^2*m^4 + 192*(A - B + C)*c^2*m^3 + 8*(99*A - 107*B + 99*C)*c^2*m^2 + 16*(77*A - 97*B + 101*C)*c^2*m + 3*(147*A - 195*B + 211*C)*c^2)*cos(...
```

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1324 vs. 2(419) = 838.

Time = 0.21 (sec) , antiderivative size = 1324, normalized size of antiderivative = 3.04

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output

```

-2*((4*m^2 + 24*m + 43)*a^m*c^(5/2) - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*sin(f*x + e)/(cos(f*x + e) + 1) + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 2*(4*m^2 + 8*m + 35)*a^m*c^(5/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 - (12*m^2 + 40*m - 15)*a^m*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + (4*m^2 + 24*m + 43)*a^m*c^(5/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)*A*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^3 + 36*m^2 + 46*m + 15)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(5/2)) - 2*((4*m^2 + 40*m + 115)*a^m*c^(5/2) - 2*(4*m^3 + 40*m^2 + 115*m)*a^m*c^(5/2)*sin(f*x + e)/(cos(f*x + e) + 1) + 2*(12*m^3 + 76*m^2 + 97*m + 175)*a^m*c^(5/2)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 - (16*m^3 + 76*m^2 + 260*m - 175)*a^m*c^(5/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 - (16*m^3 + 76*m^2 + 260*m - 175)*a^m*c^(5/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + 2*(12*m^3 + 76*m^2 + 97*m + 175)*a^m*c^(5/2)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - 2*(4*m^3 + 40*m^2 + 115*m)*a^m*c^(5/2)*sin(f*x + e)^6/(cos(f*x + e) + 1)^6 + (4*m^2 + 40*m + 115)*a^m*c^(5/2)*sin(f*x + e)^7/(cos(f*x + e) + 1)^7)*B*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((16*m^4 + 128*m^3 + 344*m^2 + 352*m + (16*m^4 + 128*m^3 + 344*m^2 + 352*m + 105)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 105)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(5/2)) + 4*(2*(4*m^2 + 56*m + 219)*a^m*c^(5/2) - 4*(4...

```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e)^2 + B \sin(fx + e) + A) (-c \sin(fx + e) + c)^{5/2} (a \sin(fx + e) + a)$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(-c*sin(f*x + e) + c)^(5/2)*(a*sin(f*x + e) + a)^m, x)

```

Mupad [B] (verification not implemented)

Time = 47.13 (sec) , antiderivative size = 1253, normalized size of antiderivative = 2.88

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(5/2)*(A + B*sin(e + f*x)
+ C*sin(e + f*x)^2),x)
```

output

```
((c - c*sin(e + f*x))^(1/2)*((C*c^2*(a + a*sin(e + f*x))^m*(m*352i + m^2*3
44i + m^3*128i + m^4*16i + 105i))/(8*f*(m*3378i + m^2*3800i + m^3*1840i +
m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*5i + f*x*5i)*(a + a*sin(e + f*x))
^m*(18900*A - 14175*B + 12285*C + 15648*A*m - 4140*B*m + 648*C*m + 5280*A*
m^2 + 896*A*m^3 + 64*A*m^4 - 832*B*m^2 - 208*B*m^3 - 16*B*m^4 + 1416*C*m^2
+ 224*C*m^3 + 16*C*m^4))/(4*f*(m*3378i + m^2*3800i + m^3*1840i + m^4*400i
+ m^5*32i + 945i)) + (c^2*exp(e*4i + f*x*4i)*(a + a*sin(e + f*x))^m*(A*18
900i - B*14175i + C*12285i + A*m*15648i - B*m*4140i + C*m*648i + A*m^2*528
0i + A*m^3*896i + A*m^4*64i - B*m^2*832i - B*m^3*208i - B*m^4*16i + C*m^2*
1416i + C*m^3*224i + C*m^4*16i))/(4*f*(m*3378i + m^2*3800i + m^3*1840i +
m^4*400i + m^5*32i + 945i)) + (C*c^2*exp(e*9i + f*x*9i)*(a + a*sin(e + f*x)
)^m*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105))/(8*f*(m*3378i + m^2*3800i
+ m^3*1840i + m^4*400i + m^5*32i + 945i)) - (c^2*exp(e*7i + f*x*7i)*(a + a
*sin(e + f*x))^m*(8*m + 4*m^2 + 3)*(126*A - 315*B + 378*C + 64*A*m - 124*B
*m + 88*C*m + 8*A*m^2 - 12*B*m^2 + 8*C*m^2))/(4*f*(m*3378i + m^2*3800i + m
^3*1840i + m^4*400i + m^5*32i + 945i)) - (c^2*exp(e*2i + f*x*2i)*(a + a*si
n(e + f*x))^m*(8*m + 4*m^2 + 3)*(A*126i - B*315i + C*378i + A*m*64i - B*m*
124i + C*m*88i + A*m^2*8i - B*m^2*12i + C*m^2*8i))/(4*f*(m*3378i + m^2*380
0i + m^3*1840i + m^4*400i + m^5*32i + 945i)) + (c^2*exp(e*3i + f*x*3i)*(2*
m + 1)*(a + a*sin(e + f*x))^m*(3150*A - 3465*B + 3150*C + 2356*A*m - 17...
```


Reduce [F]

$$\begin{aligned}
& \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2} (A + B \sin(e + fx) \\
& + C \sin^2(e + fx)) dx = \sqrt{c} c^2 \left(\left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^4 dx \right) c \right. \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^3 dx \right) b \\
& - 2 \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^3 dx \right) c \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) a \\
& - 2 \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) b \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) c \\
& - 2 \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) a \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) b \\
& \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} dx \right) a \right)
\end{aligned}$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `sqrt(c)*c**2*(int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**4,x)*c + int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**3,x)*b - 2*int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**3,x)*c + int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**2,x)*a - 2*int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**2,x)*b + int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x)**2,x)*c - 2*int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x),x)*a + int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x),x)*b + int((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1),x)*a)`

3.18 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{3/2} (A + B \sin(e$

Optimal result	193
Mathematica [A] (verified)	194
Rubi [A] (verified)	194
Maple [F]	198
Fricas [A] (verification not implemented)	198
Sympy [F(-1)]	199
Maxima [B] (verification not implemented)	199
Giac [F]	200
Mupad [B] (verification not implemented)	201
Reduce [F]	202

Optimal result

Integrand size = 48, antiderivative size = 322

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$-\frac{8c^2(B(21 - 8m - 4m^2) - C(19 - 8m + 4m^2) - A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m}{f(5 + 2m)(7 + 2m)(3 + 8m + 4m^2) \sqrt{c - c \sin(e + fx)}}$$

$$-\frac{2c(B(21 - 8m - 4m^2) - C(19 - 8m + 4m^2) - A(35 + 24m + 4m^2)) \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)}}{f(3 + 2m)(5 + 2m)(7 + 2m)}$$

$$-\frac{2(7B + 2C + 2Bm + 4Cm) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2}}{f(5 + 2m)(7 + 2m)}$$

$$+ \frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{5/2}}{cf(7 + 2m)}$$

output

```
-8*c^2*(B*(-4*m^2-8*m+21)-C*(4*m^2-8*m+19)-A*(4*m^2+24*m+35))*cos(f*x+e)*(
a+a*sin(f*x+e))^m/f/(5+2*m)/(7+2*m)/(4*m^2+8*m+3)/(c-c*sin(f*x+e))^(1/2)-2
*c*(B*(-4*m^2-8*m+21)-C*(4*m^2-8*m+19)-A*(4*m^2+24*m+35))*cos(f*x+e)*(a+a*
sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)/f/(3+2*m)/(5+2*m)/(7+2*m)-2*(2*B*m+4*
C*m+7*B+2*C)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/f/(5+2*m
)/(7+2*m)+2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(5/2)/c/f/(7+
2*m)
```

Mathematica [A] (verified)

Time = 11.05 (sec) , antiderivative size = 306, normalized size of antiderivative = 0.95

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \frac{c(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (700$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
(c*(Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*sqrt[c - c*Sin[e + f*x]]*(700*A - 546*B + 494*C + 760*A*m - 380*B*m + 284*C*m + 272*A*m^2 - 120*B*m^2 + 136*C*m^2 + 32*A*m^3 - 16*B*m^3 + 16*C*m^3 + 2*(3 + 8*m + 4*m^2)*(B*(7 + 2*m) - C*(13 + 2*m))*Cos[2*(e + f*x)] - (1 + 2*m)*(4*A*(35 + 24*m + 4*m^2) - 4*B*(63 + 32*m + 4*m^2) + C*(253 + 80*m + 12*m^2))*Sin[e + f*x] + 15*C*Sin[3*(e + f*x)] + 46*C*m*Sin[3*(e + f*x)] + 36*C*m^2*Sin[3*(e + f*x)] + 8*C*m^3*Sin[3*(e + f*x)]))/(2*f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(7 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))
```

Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.87, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 3518, 27, 3042, 3452, 3042, 3219, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

↓ 3042

$$\int (c - c \sin(e + fx))^{3/2} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

↓ 3518

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)} - \frac{2 \int -\frac{1}{2}(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}(ac(C(5 - 2m) + A(2m + 7)) + ac(2mB + 7B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 7)}$$

↓ 27

$$\frac{\int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}(ac(C(5 - 2m) + A(2m + 7)) + ac(2mB + 7B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 7)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{\int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2}(ac(C(5 - 2m) + A(2m + 7)) + ac(2mB + 7B + 2C + 4Cm) \sin(e + fx)) dx}{ac(2m + 7)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3452

$$\frac{-\frac{ac(-A(4m^2 + 24m + 35) + B(-4m^2 - 8m + 21) - C(4m^2 - 8m + 19)) \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2} dx}{2m + 5} - \frac{2ac(2Bm + 7B + 4Cm + 2C) \sin(e + fx)}{2m + 5}}{ac(2m + 7)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{-\frac{ac(-A(4m^2 + 24m + 35) + B(-4m^2 - 8m + 21) - C(4m^2 - 8m + 19)) \int(\sin(e + fx)a + a)^m(c - c \sin(e + fx))^{3/2} dx}{2m + 5} - \frac{2ac(2Bm + 7B + 4Cm + 2C) \sin(e + fx)}{2m + 5}}{ac(2m + 7)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3219

$$\frac{ac(-A(4m^2 + 24m + 35) + B(-4m^2 - 8m + 21) - C(4m^2 - 8m + 19)) \left(\frac{4c \int(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)} dx}{2m + 3} + \frac{2c \cos(e + fx) \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m}{f(2m + 3)} \right)}{2m + 5} - \frac{2ac(2Bm + 7B + 4Cm + 2C) \sin(e + fx)}{2m + 5}}{ac(2m + 7)}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{5/2}(a \sin(e + fx) + a)^m}{cf(2m + 7)}$$

↓ 3042

$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19)) \left(\frac{4c \int (\sin(e+fx)a+a)^m \sqrt{c-c \sin(e+fx)} dx}{2m+3} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)}{f(2m+3)} \right)}{2m+5} \quad ac(2m+7)$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{5/2}(a \sin(e+fx)+a)^m}{cf(2m+7)}$$

↓ 3217

$$\frac{ac(-A(4m^2+24m+35)+B(-4m^2-8m+21)-C(4m^2-8m+19)) \left(\frac{8c^2 \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)(2m+3) \sqrt{c-c \sin(e+fx)}} + \frac{2c \cos(e+fx) \sqrt{c-c \sin(e+fx)} (a \sin(e+fx)+a)}{f(2m+3)} \right)}{2m+5} \quad ac(2m+7)$$

$$\frac{2C \cos(e+fx)(c-c \sin(e+fx))^{5/2}(a \sin(e+fx)+a)^m}{cf(2m+7)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x]
+ C*Sin[e + f*x]^2),x]
```

output

```
(2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(5/2))/(c*f*
(7 + 2*m)) + ((-2*a*c*(7*B + 2*C + 2*B*m + 4*C*m)*Cos[e + f*x]*(a + a*Sin[
e + f*x])^m*(c - c*Sin[e + f*x])^(3/2))/(f*(5 + 2*m)) - (a*c*(B*(21 - 8*m
- 4*m^2) - C*(19 - 8*m + 4*m^2) - A*(35 + 24*m + 4*m^2))*((8*c^2*Cos[e +
f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]
]) + (2*c*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]])/(f
*(3 + 2*m))))/(5 + 2*m))/(a*c*(7 + 2*m))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3217

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]]*((c_) + (d_)*sin[(e_) + (f_)*
(x_)])^(n_), x_Symbol] := Simp[-2*b*Cos[e + f*x]*((c + d*Sin[e + f*x])^
n/(f*(2*n + 1)*Sqrt[a + b*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, n
}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[n, -2^(-1)]
```

rule 3219

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*(a + b*Sin[e + f*x])^
(m - 1)*((c + d*Sin[e + f*x])^n/(f*(m + n))), x] + Simp[a*((2*m - 1)/(m + n
)) Int[(a + b*Sin[e + f*x])^(m - 1)*(c + d*Sin[e + f*x])^n, x], x] /; Fre
eQ[{a, b, c, d, e, f, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && I
GtQ[m - 1/2, 0] && !LtQ[n, -1] && !(IGtQ[n - 1/2, 0] && LtQ[n, m]) && !(
ILtQ[m + n, 0] && GtQ[2*m + n + 1, 0])
```

rule 3452

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m +
n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(
a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d,
e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m,
-2^(-1)] && NeQ[m + n + 1, 0]
```

rule 3518

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x]
)^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m
+ n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*
Sin[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] &&
EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n +
2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{\frac{3}{2}} (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 563, normalized size of antiderivative = 1.75

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `-2*((8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^4 - 16*(A + B + C)*c*m^2 - (8*(B - C)*c*m^3 + 4*(11*B - 17*C)*c*m^2 + 2*(31*B - 55*C)*c*m + 3*(7*B - 13*C)*c)*cos(f*x + e)^3 - 32*(3*A + B - C)*c*m - (8*(A + C)*c*m^3 + 4*(13*A - 6*B + 5*C)*c*m^2 + 2*(47*A - 48*B + 47*C)*c*m + (35*A - 42*B + 43*C)*c)*cos(f*x + e)^2 - 4*(35*A - 21*B + 19*C)*c - (8*(A - B + C)*c*m^3 + 4*(17*A - 13*B + 17*C)*c*m^2 + 2*(95*A - 63*B + 63*C)*c*m + (175*A - 147*B + 143*C)*c)*cos(f*x + e) - (16*(A + B + C)*c*m^2 + (8*C*c*m^3 + 36*C*c*m^2 + 46*C*c*m + 15*C*c)*cos(f*x + e)^3 + 32*(3*A + B - C)*c*m + (8*B*c*m^3 + 4*(11*B - 8*C)*c*m^2 + 2*(31*B - 32*C)*c*m + 3*(7*B - 8*C)*c)*cos(f*x + e)^2 + 4*(35*A - 21*B + 19*C)*c - (8*(A - B + C)*c*m^3 + 4*(13*A - 17*B + 13*C)*c*m^2 + 2*(47*A - 79*B + 79*C)*c*m + (35*A - 63*B + 67*C)*c)*cos(f*x + e))*sin(f*x + e))*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*cos(f*x + e) - (16*f*m^4 + 128*f*m^3 + 344*f*m^2 + 352*f*m + 105*f)*sin(f*x + e) + 105*f)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 950 vs. $2(310) = 620$.

Time = 0.19 (sec) , antiderivative size = 950, normalized size of antiderivative = 2.95

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output

```

-2*((a^m*c^(3/2)*(2*m + 5) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)/(cos(f*x +
e) + 1) - a^m*c^(3/2)*(2*m - 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + a^m
*c^(3/2)*(2*m + 5)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3)*A*e^(2*m*log(sin(f
*x + e)/(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^
2 + 1))/((4*m^2 + 8*m + 3)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2)
) - 2*(a^m*c^(3/2)*(2*m + 9) - 2*(2*m^2 + 9*m)*a^m*c^(3/2)*sin(f*x + e)/(c
os(f*x + e) + 1) + (4*m^2 + 15)*a^m*c^(3/2)*sin(f*x + e)^2/(cos(f*x + e) +
1)^2 + (4*m^2 + 15)*a^m*c^(3/2)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 - 2*(
2*m^2 + 9*m)*a^m*c^(3/2)*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 + a^m*c^(3/2)
*(2*m + 9)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)*B*e^(2*m*log(sin(f*x + e)/
(cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/
((8*m^3 + 36*m^2 + 46*m + (8*m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^2/(cos
(f*x + e) + 1)^2 + 15)*(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)^(3/2)) +
4*(2*a^m*c^(3/2)*(2*m + 13) - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*x + e)/(c
os(f*x + e) + 1) + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)*sin(f*x + e)^2
/(cos(f*x + e) + 1)^2 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(f*x +
e)^3/(cos(f*x + e) + 1)^3 - (8*m^3 + 20*m^2 + 82*m - 35)*a^m*c^(3/2)*sin(
f*x + e)^4/(cos(f*x + e) + 1)^4 + (8*m^3 + 60*m^2 + 66*m + 91)*a^m*c^(3/2)
*sin(f*x + e)^5/(cos(f*x + e) + 1)^5 - 4*(2*m^2 + 13*m)*a^m*c^(3/2)*sin(f*
x + e)^6/(cos(f*x + e) + 1)^6 + 2*a^m*c^(3/2)*(2*m + 13)*sin(f*x + e)^7...

```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e)^2 + B \sin(fx + e) + A) (-c \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(-c*sin(f*x + e) + c)^(3
/2)*(a*sin(f*x + e) + a)^m, x)

```

Mupad [B] (verification not implemented)

Time = 51.63 (sec) , antiderivative size = 790, normalized size of antiderivative = 2.45

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(3/2)*(A + B*sin(e + f*x)
+ C*sin(e + f*x)^2),x)
```

output

```
((c - c*sin(e + f*x))^(1/2)*((C*c*(a + a*sin(e + f*x))^m*(m*46i + m^2*36i
+ m^3*8i + 15i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp
(e*3i + f*x*3i)*(a + a*sin(e + f*x))^m*(1260*A - 840*B + 735*C + 1144*A*m
- 128*B*m - 18*C*m + 336*A*m^2 + 32*A*m^3 + 32*B*m^2 + 100*C*m^2 + 8*C*m^3
)))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (c*exp(e*4i + f*x*4i
)*(a + a*sin(e + f*x))^m*(A*1260i - B*840i + C*735i + A*m*1144i - B*m*128i
- C*m*18i + A*m^2*336i + A*m^3*32i + B*m^2*32i + C*m^2*100i + C*m^3*8i))/
(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp(e*5i + f*x*5i)*
(2*m + 1)*(a + a*sin(e + f*x))^m*(140*A - 210*B + 175*C + 96*A*m - 88*B*m +
16*C*m + 16*A*m^2 - 8*B*m^2 + 4*C*m^2))/(4*f*(352*m + 344*m^2 + 128*m^3 +
16*m^4 + 105)) - (c*exp(e*2i + f*x*2i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(
A*140i - B*210i + C*175i + A*m*96i - B*m*88i + C*m*16i + A*m^2*16i - B*m^2
*8i + C*m^2*4i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) + (c*exp
(e*1i + f*x*1i)*(a + a*sin(e + f*x))^m*(8*m + 4*m^2 + 3)*(14*B - 21*C + 4*
B*m - 2*C*m))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (c*exp(e*
6i + f*x*6i)*(a + a*sin(e + f*x))^m*(8*m + 4*m^2 + 3)*(B*14i - C*21i + B*m
*4i - C*m*2i))/(4*f*(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)) - (C*c*exp
(e*7i + f*x*7i)*(a + a*sin(e + f*x))^m*(46*m + 36*m^2 + 8*m^3 + 15))/(4*f*
(352*m + 344*m^2 + 128*m^3 + 16*m^4 + 105)))/(exp(e*4i + f*x*4i) - (exp(e
*3i + f*x*3i)*(m*352i + m^2*344i + m^3*128i + m^4*16i + 105i))/(352*m + ...
```

Reduce [F]

$$\begin{aligned}
& \int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{3/2} (A + B \sin(e + fx) \\
& + C \sin^2(e + fx)) dx = \sqrt{c} c \left(- \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^3 dx \right) c \right. \\
& - \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) b \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) c \\
& - \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) a \\
& + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) b \\
& \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} dx \right) a \right)
\end{aligned}$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

output

```
sqrt(c)*c*( - int((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e + f*x)**3,x)*c - int((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e + f*x)**2,x)*b + int((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e + f*x)**2,x)*c - int((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e + f*x),x)*a + int((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e + f*x),x)*b + int((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1),x)*a)
```

3.19 $\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+B \sin(e$

Optimal result	203
Mathematica [A] (verified)	204
Rubi [A] (verified)	204
Maple [F]	207
Fricas [A] (verification not implemented)	207
Sympy [F]	208
Maxima [B] (verification not implemented)	208
Giac [F]	209
Mupad [B] (verification not implemented)	210
Reduce [F]	210

Optimal result

Integrand size = 48, antiderivative size = 197

$$\int (a+a \sin(e+fx))^m \sqrt{c-c \sin(e+fx)}(A+B \sin(e+fx)+C \sin^2(e+fx)) dx$$

$$= \frac{2c(C-6Cm+A(5+2m)-B(5+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{f(1+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{2c(5B+2C+2Bm+4Cm) \cos(e+fx)(a+a \sin(e+fx))^{1+m}}{af(3+2m)(5+2m)\sqrt{c-c \sin(e+fx)}} + \frac{2C \cos(e+fx)(a+a \sin(e+fx))^m(c-c \sin(e+fx))^{3/2}}{cf(5+2m)}$$

output

```
2*c*(C-6*C*m+A*(5+2*m)-B*(5+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(1+2*m)/
(5+2*m)/(c-c*sin(f*x+e))^(1/2)+2*c*(2*B*m+4*C*m+5*B+2*C)*cos(f*x+e)*(a+a*s
in(f*x+e))^(1+m)/a/f/(3+2*m)/(5+2*m)/(c-c*sin(f*x+e))^(1/2)+2*C*cos(f*x+e)
*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(3/2)/c/f/(5+2*m)
```

Mathematica [A] (verified)

Time = 3.08 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.90

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \frac{(\cos(\frac{1}{2}(e + fx)) + \sin(\frac{1}{2}(e + fx))) (a(1 + \sin(e + fx)))^m \sqrt{c - c \sin(e + fx)} (30A - 20B + 19C + 32A^2m - 8B^2m + 8C^2m + 8A^2m^2 + 4C^2m^2 - C(3 + 8m + 4m^2) \cos[2(e + fx)] + 2(1 + 2m)(5B - 4C + 2Bm) \sin[e + fx])}{f(1 + 2m)(3 + 2m)(5 + 2m)}$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c - c*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
((Cos[(e + f*x)/2] + Sin[(e + f*x)/2])*(a*(1 + Sin[e + f*x]))^m*Sqrt[c - c*Sin[e + f*x]]*(30*A - 20*B + 19*C + 32*A*m - 8*B*m + 8*C*m + 8*A*m^2 + 4*C*m^2 - C*(3 + 8*m + 4*m^2)*Cos[2*(e + f*x)] + 2*(1 + 2*m)*(5*B - 4*C + 2*B*m)*Sin[e + f*x]))/(f*(1 + 2*m)*(3 + 2*m)*(5 + 2*m)*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2]))
```

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.146$, Rules used = {3042, 3518, 27, 3042, 3450, 3042, 3217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{c - c \sin(e + fx)} (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3518}$$

$$\frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)} - \frac{2 \int -\frac{1}{2}(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}(ac(C(3 - 2m) + A(2m + 5)) + ac(2mB + 5B + 2C + 4Cm) \sin(e + fx))}{ac(2m + 5)}$$

↓ 27

$$\frac{\int(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}(ac(C(3 - 2m) + A(2m + 5)) + ac(2mB + 5B + 2C + 4Cm) \sin(e + fx))}{ac(2m + 5)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3042

$$\frac{\int(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}(ac(C(3 - 2m) + A(2m + 5)) + ac(2mB + 5B + 2C + 4Cm) \sin(e + fx))}{ac(2m + 5)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3450

$$\frac{ac(A(2m + 5) - B(2m + 5) - 6Cm + C) \int(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}dx + c(2Bm + 5B + 4Cm + ac(2m + 5))}{ac(2m + 5)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3042

$$\frac{ac(A(2m + 5) - B(2m + 5) - 6Cm + C) \int(\sin(e + fx)a + a)^m \sqrt{c - c \sin(e + fx)}dx + c(2Bm + 5B + 4Cm + ac(2m + 5))}{ac(2m + 5)} - \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

↓ 3217

$$\frac{\frac{2ac^2(A(2m+5)-B(2m+5)-6Cm+C) \cos(e+fx)(a \sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c \sin(e+fx)}} + \frac{2c^2(2Bm+5B+4Cm+2C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}}{f(2m+3)\sqrt{c-c \sin(e+fx)}}}{ac(2m + 5)} + \frac{2C \cos(e + fx)(c - c \sin(e + fx))^{3/2}(a \sin(e + fx) + a)^m}{cf(2m + 5)}$$

input $\text{Int}[(a + a*\sin[e + f*x])^m*\sqrt{c - c*\sin[e + f*x]}*(A + B*\sin[e + f*x] + C*\sin[e + f*x]^2),x]$

output $(2*C*\cos[e + f*x]*(a + a*\sin[e + f*x])^m*(c - c*\sin[e + f*x])^{3/2})/(c*f*(5 + 2*m)) + ((2*a*c^2*(C - 6*C*m + A*(5 + 2*m) - B*(5 + 2*m))*\cos[e + f*x]*(a + a*\sin[e + f*x])^m)/(f*(1 + 2*m)*\sqrt{c - c*\sin[e + f*x]}) + (2*c^2*(5*B + 2*C + 2*B*m + 4*C*m)*\cos[e + f*x]*(a + a*\sin[e + f*x])^{1+m})/(f*(3 + 2*m)*\sqrt{c - c*\sin[e + f*x]})/(a*c*(5 + 2*m))$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3217 $\text{Int}[\sqrt{(a_*) + (b_*)\sin[(e_*) + (f_*)(x_)]}] * ((c_*) + (d_*)\sin[(e_*) + (f_*)(x_)]))^n, x_Symbol] \rightarrow \text{Simp}[-2*b*\cos[e + f*x]*((c + d*\sin[e + f*x])^n/(f*(2*n + 1)*\sqrt{a + b*\sin[e + f*x]})), x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ \text{NeQ}[n, -2^{(-1)}]$

rule 3450 $\text{Int}[\sqrt{(a_*) + (b_*)\sin[(e_*) + (f_*)(x_)]}] * ((A_*) + (B_*)\sin[(e_*) + (f_*)(x_)]))^n, x_Symbol] \rightarrow \text{Simp}[B/d \text{ Int}[\sqrt{a + b*\sin[e + f*x]}*(c + d*\sin[e + f*x])^{n+1}, x], x] - \text{Simp}[(B*c - A*d)/d \text{ Int}[\sqrt{a + b*\sin[e + f*x]}*(c + d*\sin[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

rule 3518

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_) + (C_)*sin[(e_)
) + (f_)*(x_)]^2), x_Symbol] :> Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x]
)^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m +
n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m
+ n + 2) + C*(a*c*m + b*d*(n + 1)) + (b*B*d*(m + n + 2) - b*c*C*(2*m + 1))*
Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] &&
EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n +
2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c - c \sin(fx + e)} (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)
^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)
^2),x)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.57

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{-2((4Cm^2 + 8Cm + 3C) \cos(fx + e)^3 - 4(A + B + C)m^2 + (4(B + C)m^2 + 12Bm + 5B - C) \cos(fx + e))}{(a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)}}$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="fricas")
```


output

```
-2*((4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^3 - 4*(A + B + C)*m^2 + (4*(B + C)
)*m^2 + 12*B*m + 5*B - C)*cos(f*x + e)^2 - 8*(2*A + B)*m - (4*(A + C)*m^2
+ 4*(4*A - B + 2*C)*m + 15*A - 10*B + 11*C)*cos(f*x + e) - (4*(A + B + C)*
m^2 - (4*C*m^2 + 8*C*m + 3*C)*cos(f*x + e)^2 + 8*(2*A + B)*m + (4*B*m^2 +
4*(3*B - 2*C)*m + 5*B - 4*C)*cos(f*x + e) + 15*A - 5*B + 7*C)*sin(f*x + e)
- 15*A + 5*B - 7*C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(8*f
*m^3 + 36*f*m^2 + 46*f*m + (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*cos(f*x +
e) - (8*f*m^3 + 36*f*m^2 + 46*f*m + 15*f)*sin(f*x + e) + 15*f)
```

Sympy [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m \sqrt{-c(\sin(e + fx) - 1)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input

```
integrate((a+a*sin(f*x+e))**m*(c-c*sin(f*x+e))**(1/2)*(A+B*sin(f*x+e)+C*sin
n(f*x+e)**2),x)
```

output

```
Integral((a*(sin(e + f*x) + 1))**m*sqrt(-c*(sin(e + f*x) - 1))*(A + B*sin(
e + f*x) + C*sin(e + f*x)**2), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 644 vs. 2(191) = 382.

Time = 0.18 (sec) , antiderivative size = 644, normalized size of antiderivative = 3.27

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="maxima")
```

output

```

-2*(2*(2*a^m*sqrt(c)*m*sin(f*x + e)/(cos(f*x + e) + 1) + 2*a^m*sqrt(c)*m*
in(f*x + e)^2/(cos(f*x + e) + 1)^2 - a^m*sqrt(c) - a^m*sqrt(c)*sin(f*x + e
)^3/(cos(f*x + e) + 1)^3)*B*e^(2*m*log(sin(f*x + e)/(cos(f*x + e) + 1) + 1
) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((4*m^2 + 8*m + (4*m^2
+ 8*m + 3)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 3)*sqrt(sin(f*x + e)^2/(
cos(f*x + e) + 1)^2 + 1)) - 4*(4*a^m*sqrt(c)*m*sin(f*x + e)/(cos(f*x + e)
+ 1) - (4*m^2 + 4*m + 5)*a^m*sqrt(c)*sin(f*x + e)^2/(cos(f*x + e) + 1)^2 -
(4*m^2 + 4*m + 5)*a^m*sqrt(c)*sin(f*x + e)^3/(cos(f*x + e) + 1)^3 + 4*a^m
*sqrt(c)*m*sin(f*x + e)^4/(cos(f*x + e) + 1)^4 - 2*a^m*sqrt(c) - 2*a^m*sqr
t(c)*sin(f*x + e)^5/(cos(f*x + e) + 1)^5)*C*e^(2*m*log(sin(f*x + e)/(cos(f
*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/((8*m^
3 + 36*m^2 + 46*m + 2*(8*m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^2/(cos(f*x
+ e) + 1)^2 + (8*m^3 + 36*m^2 + 46*m + 15)*sin(f*x + e)^4/(cos(f*x + e) +
1)^4 + 15)*sqrt(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1)) + (a^m*sqrt(c)
+ a^m*sqrt(c)*sin(f*x + e)/(cos(f*x + e) + 1))*A*e^(2*m*log(sin(f*x + e)/(
cos(f*x + e) + 1) + 1) - m*log(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/
(2*m + 1)*sqrt(sin(f*x + e)^2/(cos(f*x + e) + 1)^2 + 1))/f

```

Giac [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{-c \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input

```

integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(
f*x+e)^2),x, algorithm="giac")

```

output

```

integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*sqrt(-c*sin(f*x + e) + c
)*(a*sin(f*x + e) + a)^m, x)

```

Mupad [B] (verification not implemented)

Time = 39.55 (sec) , antiderivative size = 510, normalized size of antiderivative = 2.59

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{\sqrt{c - c \sin(e + fx)} \left(-\frac{e^{e 3i + f x 3i} (a + a \sin(e + fx))^m (30A - 15B + 15C + 32Am + 4Bm + 8Am^2 + 4Bm^2 + 4Cm^2)}{f(m^3 8i + m^2 36i + m 46i + 15i)} - \frac{e^{e 2i + f x 2i} (a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{f(m^3 8i + m^2 36i + m 46i + 15i)} \right)}{f(m^3 8i + m^2 36i + m 46i + 15i)}$$

input

```
int((a + a*sin(e + f*x))^m*(c - c*sin(e + f*x))^(1/2)*(A + B*sin(e + f*x)
+ C*sin(e + f*x)^2),x)
```

output

```
-((c - c*sin(e + f*x))^(1/2)*((C*(a + a*sin(e + f*x))^m*(m*8i + m^2*4i + 3
i))/(2*f*(m*46i + m^2*36i + m^3*8i + 15i)) - (exp(e*2i + f*x*2i)*(a + a*si
n(e + f*x))^m*(A*30i - B*15i + C*15i + A*m*32i + B*m*4i + A*m^2*8i + B*m^2
*4i + C*m^2*4i))/(f*(m*46i + m^2*36i + m^3*8i + 15i)) - (exp(e*3i + f*x*3i
)*(a + a*sin(e + f*x))^m*(30*A - 15*B + 15*C + 32*A*m + 4*B*m + 8*A*m^2 +
4*B*m^2 + 4*C*m^2))/(f*(m*46i + m^2*36i + m^3*8i + 15i)) + (C*exp(e*5i + f
*x*5i)*(a + a*sin(e + f*x))^m*(8*m + 4*m^2 + 3))/(2*f*(m*46i + m^2*36i + m
^3*8i + 15i)) + (exp(e*1i + f*x*1i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(10*B
- 5*C + 4*B*m + 2*C*m))/(2*f*(m*46i + m^2*36i + m^3*8i + 15i)) + (exp(e*4
i + f*x*4i)*(2*m + 1)*(a + a*sin(e + f*x))^m*(B*10i - C*5i + B*m*4i + C*m*
2i))/(2*f*(m*46i + m^2*36i + m^3*8i + 15i))))/(exp(e*3i + f*x*3i) + (exp(e
*2i + f*x*2i)*(46*m + 36*m^2 + 8*m^3 + 15))/(m*46i + m^2*36i + m^3*8i + 15
i))
```

Reduce [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c - c \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \sqrt{c} \left(\left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2 dx \right) c \right.$$

$$\quad \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e) dx \right) b \right.$$

$$\quad \left. + \left(\int (a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} dx \right) a \right)$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

output

```
sqrt(c)*(int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**2,x)*c + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x),x)*b + int((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1),x)*a)
```

3.20
$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c-c \sin(e+fx)}} dx$$

Optimal result	212
Mathematica [A] (verified)	213
Rubi [A] (verified)	213
Maple [F]	216
Fricas [F]	217
Sympy [F]	217
Maxima [F]	218
Giac [F(-1)]	218
Mupad [F(-1)]	218
Reduce [F]	219

Optimal result

Integrand size = 48, antiderivative size = 170

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= -\frac{2B \cos(e + fx)(a + a \sin(e + fx))^m}{f(1 + 2m)\sqrt{c - c \sin(e + fx)}} + \frac{(A + B + C) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) (a + a \sin(e + fx))^m}{f(1 + 2m)\sqrt{c - c \sin(e + fx)}} - \frac{2C \cos(e + fx)(a + a \sin(e + fx))^{1+m}}{af(3 + 2m)\sqrt{c - c \sin(e + fx)}}$$

output

```
-2*B*cos(f*x+e)*(a+a*sin(f*x+e))^m/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)+(A+B+C)
*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2+m],1/2+1/2*sin(f*x+e))*(a+a*sin(f*x
+e))^m/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^(1
+m)/a/f/(3+2*m)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 40.44 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.94

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \frac{\cos(e + fx)(a(1 + \sin(e + fx)))^m (2(A + C)(3 + 2m) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) + B(1 + 2m) \operatorname{Hypergeometric2F1}\left(1, \frac{3}{2} + m, \frac{5}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right) - 2(3B + 2C + 2Bm + 4Cm + 2C(1 + 2m)\sin(e + fx)))}{2f(1 + 2m)(3 + 2m)\sqrt{c - c \sin(e + fx)}}$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/Sqrt[c - c*Sin[e + f*x]],x]
```

output

```
(Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(2*(A + C)*(3 + 2*m)*Hypergeometric
2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2] + B*(1 + 2*m)*Hypergeometri
c2F1[1, 3/2 + m, 5/2 + m, (1 + Sin[e + f*x])/2]*(1 + Sin[e + f*x]) - 2*(3*
B + 2*C + 2*B*m + 4*C*m + 2*C*(1 + 2*m)*Sin[e + f*x]))/(2*f*(1 + 2*m)*(3
+ 2*m)*Sqrt[c - c*Sin[e + f*x]])
```

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 3516, 3042, 3452, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{\sqrt{c - c \sin(e + fx)}} dx$$

$$\downarrow \text{3516}$$

$$\int \frac{(\sin(e + fx)a + a)^m (A + C + B \sin(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx - \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^{m+1}}{af(2m + 3)\sqrt{c - c \sin(e + fx)}}$$

$$\begin{aligned}
& \int \frac{(\sin(e+fx)a+a)^m(A+C+B\sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& (A+B+C) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx - \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \quad \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& (A+B+C) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx - \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \quad \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3224} \\
& \frac{(A+B+C)\cos(e+fx) \int \sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}} dx}{\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \\
& \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{(A+B+C)\cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \\
& \quad \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{3146} \\
& \frac{a(A+B+C)\cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a\sin(e+fx)} d(a\sin(e+fx))}{f\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} - \\
& \frac{2B\cos(e+fx)(a\sin(e+fx)+a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow \text{78}
\end{aligned}$$

$$\frac{(A + B + C) \cos(e + fx)(a \sin(e + fx) + a)^m \operatorname{Hypergeometric2F1}\left(1, m + \frac{1}{2}, m + \frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1)\sqrt{c-c\sin(e+fx)} - \frac{2B \cos(e+fx)(a \sin(e+fx) + a)^m}{f(2m+1)\sqrt{c-c\sin(e+fx)}} - \frac{2C \cos(e+fx)(a \sin(e+fx) + a)^{m+1}}{af(2m+3)\sqrt{c-c\sin(e+fx)}}$$

input `Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[c - c*Sin[e + f*x]],x]`

output `(-2*B*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + ((A + B + C)*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) - (2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(a*f*(3 + 2*m)*Sqrt[c - c*Sin[e + f*x]])`

Defintions of rubi rules used

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3146 `Int[cos[(e_.) + (f_.)*(x_)]^(p_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1)/2], x], x, b*Sin[e + f*x], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`

rule 3224

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])
```

rule 3452

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]
```

rule 3516

```
Int[(((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]) + (C_)*sin[(e_) + (f_)*(x_)])^2)/Sqrt[(c_) + (d_)*sin[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*C*Cos[e + f*x]*((a + b*Sin[e + f*x])^(m + 1)/(b*f*(2*m + 3)*Sqrt[c + d*Sin[e + f*x]]), x] + Int[(a + b*Sin[e + f*x])^m*(Simp[A + C + B*Sin[e + f*x], x]/Sqrt[c + d*Sin[e + f*x]]), x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)]
```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C \sin(fx + e)^2)}{\sqrt{c - c \sin(fx + e)}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(c*sin(f*x + e) - c), x)`

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{-c(\sin(e + fx) - 1)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(1/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/sqrt(-c*(sin(e + f*x) - 1)), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{-c \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/sqrt(-c*sin(f*x + e) + c), x)`

Giac [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{\sqrt{c - c \sin(e + fx)}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(1/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(1/2), x)`

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c - c \sin(e + fx)}} dx =$$

$$\frac{\sqrt{c} \left(\left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2}{\sin(fx + e) - 1} dx \right) c + \left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)}{\sin(fx + e) - 1} dx \right) b + \left(\int \frac{(a + a \sin(fx + e))^m}{\sin(fx + e) - 1} dx \right) a \right)}{c}$$

input `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(1/2),x)`

output `(- sqrt(c)*(int(((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x)**2)/(sin(e + f*x) - 1),x)*c + int(((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1)*sin(e + f*x))/(sin(e + f*x) - 1),x)*b + int(((sin(e + f*x)*a + a)**m*sqrt(- sin(e + f*x) + 1))/(sin(e + f*x) - 1),x)*a)/c`

3.21
$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{3/2}} dx$$

Optimal result	220
Mathematica [A] (verified)	221
Rubi [A] (verified)	221
Maple [F]	225
Fricas [F]	225
Sympy [F]	226
Maxima [F]	226
Giac [F(-2)]	226
Mupad [F(-1)]	227
Reduce [F]	227

Optimal result

Integrand size = 48, antiderivative size = 216

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \frac{(A + B + C) \cos(e + fx)(a + a \sin(e + fx))}{4af(c - c \sin(e + fx))^{3/2}} + \frac{(A + B + 2Am + 2Bm + C(9 + 2m)) \cos(e + fx)(a + a \sin(e + fx))^m}{4cf(1 + 2m)\sqrt{c - c \sin(e + fx)}} + \frac{(A(1 - 2m) - B(3 + 2m) - C(7 + 2m)) \cos(e + fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx))\right)}{4cf(1 + 2m)\sqrt{c - c \sin(e + fx)}}$$

output

```
1/4*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(3/2)+1/4*(A+B+2*A*m+2*B*m+C*(9+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)+1/4*(A*(1-2*m)-B*(3+2*m)-C*(7+2*m))*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2+m],1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 54.18 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.75

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx =$$

$$\frac{\cos(e + fx)(a(1 + \sin(e + fx)))^m ((8C + B(3 + 2m)) \operatorname{Hypergeometric2F1}(1, \frac{1}{2} + m, \frac{3}{2} + m, \frac{1}{2}(1 + \sin(e + fx)))$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/(c - c*Sin[e + f*x])^(3/2),x]
```

output

```
-1/4*(Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*((8*C + B*(3 + 2*m))*Hypergeom
etric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(-1 + Sin[e + f*x]) +
2*(B + 4*C + 2*B*m - (A + C)*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + S
in[e + f*x])/2]*(-1 + Sin[e + f*x]) - 4*C*Sin[e + f*x]))/(c*f*(1 + 2*m)*(
-1 + Sin[e + f*x])*Sqrt[c - c*Sin[e + f*x]])
```

Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {3042, 3514, 27, 3042, 3452, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)^2)}{(c - c \sin(e + fx))^{3/2}} dx$$

$$\downarrow \text{3514}$$

$$\begin{aligned}
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} - \\
& \frac{\int -\frac{(\sin(e+fx)a+a)^m(a^2(A(3-2m)-(B+C)(2m+5))-a^2(2mA+A+B+2Bm+C(2m+9))\sin(e+fx))}{2\sqrt{c-c\sin(e+fx)}} dx}{4a^2c} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{(\sin(e+fx)a+a)^m(a^2(A(3-2m)-(B+C)(2m+5))-a^2(2mA+A+B+2Bm+C(2m+9))\sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx}{8a^2c} + \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(\sin(e+fx)a+a)^m(a^2(A(3-2m)-(B+C)(2m+5))-a^2(2mA+A+B+2Bm+C(2m+9))\sin(e+fx))}{\sqrt{c-c\sin(e+fx)}} dx}{8a^2c} + \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3452 \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7))\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9))\cos(e+fx)(a\sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{8a^2c} \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7))\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9))\cos(e+fx)(a\sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{8a^2c} \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3224 \\
& \frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7))\cos(e+fx)\int \sec(e+fx)(\sin(e+fx)a+a)^{m+\frac{1}{2}} dx + \frac{2a^2(2Am+A+2Bm+B+C(2m+9))\cos(e+fx)(a\sin(e+fx))}{f(2m+1)\sqrt{c-c\sin(e+fx)}}}{\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{4af(c-c\sin(e+fx))^{3/2}} \\
& \quad \downarrow 3042
\end{aligned}$$

$$\frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}}{\cos(e+fx)} dx}{\sqrt{a \sin(e+fx)+a}\sqrt{c-c \sin(e+fx)}} + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)}{f(2m+1)\sqrt{c-c \sin(e+fx)}}$$

$$\frac{8a^2c}{4af(c-c \sin(e+fx))^{3/2}} (A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}$$

↓ 3146

$$\frac{2a^3(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{a-a \sin(e+fx)} d(a \sin(e+fx))}{f\sqrt{a \sin(e+fx)+a}\sqrt{c-c \sin(e+fx)}} + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)}{f(2m+1)\sqrt{c-c \sin(e+fx)}}$$

$$\frac{8a^2c}{4af(c-c \sin(e+fx))^{3/2}} (A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}$$

↓ 78

$$\frac{2a^2(A(1-2m)-B(2m+3)-C(2m+7)) \cos(e+fx)(a \sin(e+fx)+a)^m \text{Hypergeometric2F1}\left(1, m+\frac{1}{2}, m+\frac{3}{2}, \frac{\sin(e+fx)a+a}{2a}\right)}{f(2m+1)\sqrt{c-c \sin(e+fx)}} + \frac{2a^2(2Am+A+2Bm+B+C(2m+9)) \cos(e+fx)(a \sin(e+fx)+a)}{f(2m+1)\sqrt{c-c \sin(e+fx)}}$$

$$\frac{8a^2c}{4af(c-c \sin(e+fx))^{3/2}} (A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(3/2), x]
```

output

```
((A + B + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(4*a*f*(c - c*Sin[e + f*x])^(3/2)) + ((2*a^2*(A + B + 2*A*m + 2*B*m + C*(9 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]) + (2*a^2*(A*(1 - 2*m) - B*(3 + 2*m) - C*(7 + 2*m))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(8*a^2*c)
```


Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*(a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3146 `Int[cos[(e_) + (f_)*(x_)]^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)), x_Symbol] := Simp[1/(b^p*f) Subst[Int[(a + x)^(m + (p - 1)/2)*(a - x)^(p - 1/2), x], x, b*Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, m}, x] && IntegerQ[(p - 1)/2] && EqQ[a^2 - b^2, 0] && (GeQ[p, -1] || !IntegerQ[m + 1/2])`
- rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`
- rule 3452 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)]*(c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(f*(m + n + 1))), x] - Simp[(B*c*(m - n) - A*d*(m + n + 1))/(d*(m + n + 1)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 1, 0]`

rule 3514

```

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*
Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Si
mp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e +
f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c
^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f
*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c +
a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] &&
NeQ[2*m + 1, 0]))

```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C \sin(fx + e)^2)}{(c - c \sin(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A - C) \sqrt{-c \sin(fx + e)}}{(-c \sin(fx + e))^{3/2}} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="fricas")
```

output

```
integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(c^2*cos(f*x + e)^2 + 2*c^2*sin(f*x + e) - 2*c^2), x)
```

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(-c(\sin(e + fx) + 1))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(3/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/(-c*(sin(e + f*x) - 1))**(3/2), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) + c)^(3/2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3/2),x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,1,3,0,0,0,0]}+%%{3,[0,1,1,0,0,0,0]} / %%{16,[
0,0,0,1,1
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + A + B \sin(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx$$

input

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c -
c*sin(e + f*x))^(3/2),x)
```

output

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c -
c*sin(e + f*x))^(3/2), x)
```

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{3/2}} dx = \frac{\sqrt{c}}{c} \left(\int \frac{(a + a \sin(fx+e))^m \sqrt{-\sin(fx+e)+1} \sin(fx+e)}{\sin(fx+e)^2 - 2 \sin(fx+e) + 1} dx + \int \frac{(a + a \sin(fx+e))^m \sqrt{-\sin(fx+e)+1}}{\sin(fx+e)^2 - 2 \sin(fx+e) + 1} dx \right)$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(3
/2),x)
```

output

```
(sqrt(c)*(int(((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f
*x)**2)/(sin(e + f*x)**2 - 2*sin(e + f*x) + 1),x)*c + int(((sin(e + f*x)*a
+ a)**m*sqrt(-sin(e + f*x) + 1)*sin(e + f*x))/(sin(e + f*x)**2 - 2*sin(
e + f*x) + 1),x)*b + int(((sin(e + f*x)*a + a)**m*sqrt(-sin(e + f*x) + 1
)))/(sin(e + f*x)**2 - 2*sin(e + f*x) + 1),x)*a))/c**2
```

$$3.22 \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx$$

Optimal result	228
Mathematica [A] (verified)	229
Rubi [A] (verified)	229
Maple [F]	233
Fricas [F]	233
Sympy [F(-1)]	234
Maxima [F]	234
Giac [F(-2)]	234
Mupad [F(-1)]	235
Reduce [F]	235

Optimal result

Integrand size = 48, antiderivative size = 230

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c-c \sin(e+fx))^{5/2}} dx = \frac{(A+B+C) \cos(e+fx)(a+a \sin(e+fx))^m}{8af(c-c \sin(e+fx))^{5/2}} + \frac{(A(5-2m)-B(3+2m)-C(11+2m)) \cos(e+fx)(a+a \sin(e+fx))^m}{16cf(c-c \sin(e+fx))^{3/2}} - \frac{(B(5-8m-4m^2)-A(3-8m+4m^2)-C(19+24m+4m^2)) \cos(e+fx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}, \frac{3}{2}, \frac{a+a \sin(e+fx)}{c-c \sin(e+fx)}\right)}{32c^2 f(1+2m) \sqrt{c-c \sin(e+fx)}}$$

output

```
1/8*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)/a/f/(c-c*sin(f*x+e))^(5/2)+1/16*(A*(5-2*m)-B*(3+2*m)-C*(11+2*m))*cos(f*x+e)*(a+a*sin(f*x+e))^m/c/f/(c-c*sin(f*x+e))^(3/2)-1/32*(B*(-4*m^2-8*m+5)-A*(4*m^2-8*m+3)-C*(4*m^2+24*m+19))*cos(f*x+e)*hypergeom([1, 1/2+m],[3/2+m],1/2+1/2*sin(f*x+e))*(a+a*sin(f*x+e))^m/c^2/f/(1+2*m)/(c-c*sin(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 53.12 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.99

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \frac{\cos(e + fx)(a(1 + \sin(e + fx)))^m (8B \cos(e + fx) + 8C \sin(e + fx) + 8A \cos(e + fx) + 8A \sin(e + fx))}{(c - c \sin(e + fx))^{5/2}}$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/(c - c*Sin[e + f*x])^(5/2),x]
```

output

```
(Cos[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(8*B*(1 + 2*m) - 2*(16*C + B*(5 + 2
*m))*Hypergeometric2F1[2, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(Cos[(e
+ f*x)/2] - Sin[(e + f*x)/2])^4 + 8*(A + C)*Hypergeometric2F1[3, 1/2 + m,
3/2 + m, (1 + Sin[e + f*x])/2]*(Cos[(e + f*x)/2] - Sin[(e + f*x)/2])^4 - 1
6*C*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (1 + Sin[e + f*x])/2]*(-3 + Cos
[2*(e + f*x)] + 4*Sin[e + f*x])))/(32*c^2*(f + 2*f*m)*(-1 + Sin[e + f*x])^
2*Sqrt[c - c*Sin[e + f*x]])
```

Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {3042, 3514, 27, 3042, 3451, 3042, 3224, 3042, 3146, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{(c - c \sin(e + fx))^{5/2}} dx$$

↓ 3514

$$\begin{aligned}
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} - \\
& \frac{\int -\frac{(\sin(e+fx)a+a)^m((A(9-2m)-(B+C)(2m+7))a^2+((A+B)(1-2m)-C(2m+15))\sin(e+fx)a^2)}{2(c-c\sin(e+fx))^{3/2}} dx}{8a^2c} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{(\sin(e+fx)a+a)^m((A(9-2m)-(B+C)(2m+7))a^2+((A+B)(1-2m)-C(2m+15))\sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{16a^2c} + \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{(\sin(e+fx)a+a)^m((A(9-2m)-(B+C)(2m+7))a^2+((A+B)(1-2m)-C(2m+15))\sin(e+fx)a^2)}{(c-c\sin(e+fx))^{3/2}} dx}{16a^2c} + \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \\
& \quad \downarrow 3451 \\
& \frac{a^2(A(5-2m)-B(2m+3)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19))\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c} \\
& \quad \downarrow 3042 \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{a^2(A(5-2m)-B(2m+3)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19))\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c} \\
& \quad \downarrow 3224 \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{a^2(A(5-2m)-B(2m+3)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^2(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19))\cos(e+fx)\int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c-c\sin(e+fx)}} dx}{2c\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}} \\
& \quad \downarrow 3042 \\
& \frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}}
\end{aligned}$$

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^3(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19))\cos(e+fx)}{2c\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}}$$

↓ 3146

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^3(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19))\cos(e+fx)}{2cf\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}}$$

↓ 78

$$\frac{a^2(A(5-2m)-B(2m+3)-C(2m+11))\cos(e+fx)(a\sin(e+fx)+a)^m}{f(c-c\sin(e+fx))^{3/2}} - \frac{a^3(-A(4m^2-8m+3)+B(-4m^2-8m+5)-C(4m^2+24m+19))\cos(e+fx)}{2cf(2m+1)\sqrt{a\sin(e+fx)+a}\sqrt{c-c\sin(e+fx)}}$$

$$\frac{(A+B+C)\cos(e+fx)(a\sin(e+fx)+a)^{m+1}}{8af(c-c\sin(e+fx))^{5/2}} \quad 16a^2c$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c - c*Sin[e + f*x])^(5/2),x]
```

output

```
((A + B + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m))/(8*a*f*(c - c*Sin[e + f*x])^(5/2)) + ((a^2*(A*(5 - 2*m) - B*(3 + 2*m) - C*(11 + 2*m))*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(f*(c - c*Sin[e + f*x])^(3/2)) - (a^2*(B*(5 - 8*m - 4*m^2) - A*(3 - 8*m + 4*m^2) - C*(19 + 24*m + 4*m^2))*Cos[e + f*x]*Hypergeometric2F1[1, 1/2 + m, 3/2 + m, (a + a*Sin[e + f*x])/(2*a)]*(a + a*Sin[e + f*x])^m)/(2*c*f*(1 + 2*m)*Sqrt[c - c*Sin[e + f*x]]))/(16*a^2*c)
```


Defintions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 78 $\text{Int}[((a_) + (b_)*(x_))^{(m_)*}((c_) + (d_)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)^n * ((a + b*x)^{(m+1}) / (b^{(n+1)} * (m+1))) * \text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IntegerQ}[n]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3146 $\text{Int}[\cos[(e_.) + (f_.)*(x_)]^{(p_.)*}((a_) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/(b^p * f) \text{ Subst}[\text{Int}[(a + x)^{(m + (p - 1)/2)} * (a - x)^{-(p - 1)/2}, x], x, b*\sin[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(p - 1)/2] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ (\text{GeQ}[p, -1] \ || \ !\text{IntegerQ}[m + 1/2])$
- rule 3224 $\text{Int}[((a_) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^{(m_.)*}((c_) + (d_.)*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[a^{\text{IntPart}[m]} * c^{\text{IntPart}[m]} * (a + b*\sin[e + f*x])^{\text{FracPart}[m]} * ((c + d*\sin[e + f*x])^{\text{FracPart}[m]} / \cos[e + f*x]^{(2*\text{FracPart}[m])}) \text{ Int}[\cos[e + f*x]^{(2*m)} * (c + d*\sin[e + f*x])^{(n - m)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ (\text{FractionQ}[m] \ || \ !\text{FractionQ}[n])$
- rule 3451 $\text{Int}[((a_) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^{(m_.)*}((A_.) + (B_.)*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(A*b - a*B)*\cos[e + f*x] * (a + b*\sin[e + f*x])^m * ((c + d*\sin[e + f*x])^n / (a*f*(2*m + 1))), x] + \text{Simp}[(a*B*(m - n) + A*b*(m + n + 1)) / (a*b*(2*m + 1)) \text{ Int}[(a + b*\sin[e + f*x])^{(m + 1)} * (c + d*\sin[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 - b^2, 0] \ \&\& \ (\text{LtQ}[m, -2^{(-1)}] \ || \ (\text{ILtQ}[m + n, 0] \ \&\& \ !\text{SumSimplerQ}[n, 1])) \ \&\& \ \text{NeQ}[2*m + 1, 0]$

rule 3514

```

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*
Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Si
mp[1/(2*b*c*d*(2*m + 1)) Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e +
f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c
^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f
*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c +
a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] &&
NeQ[2*m + 1, 0]))

```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C \sin(fx + e)^2)}{(c - c \sin(fx + e))^{\frac{5}{2}}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A - C) \sqrt{-c \sin(fx + e)}}{(-c \sin(fx + e))^{5/2}}$$

input

```
integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="fricas")
```

output

```
integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(-c*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(3*c^3*cos(f*x + e)^2 - 4*c^3 - (c^3*cos(f*x + e)^2 - 4*c^3)*sin(f*x + e)), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c-c*sin(f*x+e))**(5/2),x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{(-c \sin(fx + e) + c)^{5/2}}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(-c*sin(f*x + e) + c)^(5/2), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5/2),x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Error index.cc index_gcd Error: Bad
Argument Value
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + B \sin^2(e + fx) + A \sin^3(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx$$

input

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c -
c*sin(e + f*x))^(5/2),x)
```

output

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c -
c*sin(e + f*x))^(5/2), x)
```

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c - c \sin(e + fx))^{5/2}} dx = \frac{\sqrt{c} \left(\left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)^2}{\sin(fx + e)^3 - 3 \sin(fx + e)^2 + 3 \sin(fx + e) - 1} dx \right) c + \left(\int \frac{(a + a \sin(fx + e))^m \sqrt{-\sin(fx + e) + 1} \sin(fx + e)}{\sin(fx + e)^3 - 3 \sin(fx + e)^2 + 3 \sin(fx + e) - 1} dx \right) b + \left(\int \frac{1}{\sin(fx + e)^3 - 3 \sin(fx + e)^2 + 3 \sin(fx + e) - 1} dx \right) a \right)}{c^3}$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c-c*sin(f*x+e))^(5
/2),x)
```

output

```
( - sqrt(c)*(int(((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e
+ f*x)**2)/(sin(e + f*x)**3 - 3*sin(e + f*x)**2 + 3*sin(e + f*x) - 1),x)*c
+ int(((sin(e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1)*sin(e + f*x))/(s
in(e + f*x)**3 - 3*sin(e + f*x)**2 + 3*sin(e + f*x) - 1),x)*b + int(((sin(
e + f*x)*a + a)**m*sqrt( - sin(e + f*x) + 1))/(sin(e + f*x)**3 - 3*sin(e +
f*x)**2 + 3*sin(e + f*x) - 1),x)*a))/c**3
```

3.23 $\int (a+a \sin(e+fx))^m (c-c \sin(e+fx))^{-2-m} (A + B \sin$

Optimal result	236
Mathematica [A] (warning: unable to verify)	237
Rubi [A] (verified)	237
Maple [F]	241
Fricas [F]	242
Sympy [F]	242
Maxima [F]	243
Giac [F]	243
Mupad [F(-1)]	244
Reduce [F]	244

Optimal result

Integrand size = 50, antiderivative size = 236

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$-\frac{2^{-\frac{1}{2}-m} C \cos^3(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}(3 + 2m), \frac{1}{2}(3 + 2m), \frac{1}{2}(5 + 2m), \frac{1}{2}(1 + \sin(e + fx))\right)}{f(3 + 2m)} + \frac{(A + B + C) \cos(e + fx)(a + a \sin(e + fx))^{1+m} (c - c \sin(e + fx))^{-2-m}}{2af(3 + 2m)} + \frac{(A - B + C) \cos(e + fx)(a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-1-m}}{2cf(1 + 2m)}$$

output

```
-2^(-1/2-m)*C*cos(f*x+e)^3*hypergeom([3/2+m, 3/2+m], [5/2+m], 1/2+1/2*sin(f*x+e))*(1-sin(f*x+e))^(1/2+m)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(-2-m)/f/(3+2*m)+1/2*(A+B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^(1+m)*(c-c*sin(f*x+e))^(-2-m)/a/f/(3+2*m)+1/2*(A-B+C)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(-1-m)/c/f/(1+2*m)
```

Mathematica [A] (warning: unable to verify)

Time = 2.02 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.83

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{\sec(e + fx)(1 + \sin(e + fx))^{-m}(a(1 + \sin(e + fx)))^m(c - c \sin(e + fx))^{-m} \left(2^{\frac{3}{2}+m} C(3 + 2m) \text{Hypergeometric2F1}\left[-\frac{1}{2} - m, -\frac{1}{2} - m, \frac{1}{2} - m, \frac{1 - \sin(e + fx)}{2}\right](-1 + \sin(e + fx))\sqrt{1 + \sin(e + fx)} + (1 + \sin(e + fx))^{1+m}(2A - B + 2C + 2Am + 2Cm - (A + C - 2B(1 + m))\sin(e + fx))\right)}{(c^2 f (1 + 2m) (3 + 2m) (-1 + \sin(e + fx)) (1 + \sin(e + fx))^m (c - c \sin(e + fx))^m)}$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
-((Sec[e + f*x]*(a*(1 + Sin[e + f*x]))^m*(2^(3/2 + m)*C*(3 + 2*m)*Hypergeometric2F1[-1/2 - m, -1/2 - m, 1/2 - m, (1 - Sin[e + f*x])/2]*(-1 + Sin[e + f*x])*Sqrt[1 + Sin[e + f*x]] + (1 + Sin[e + f*x])^(1 + m)*(2*A - B + 2*C + 2*A*m + 2*C*m - (A + C - 2*B*(1 + m))*Sin[e + f*x]))/(c^2*f*(1 + 2*m)*(3 + 2*m)*(-1 + Sin[e + f*x])*(1 + Sin[e + f*x])^m*(c - c*Sin[e + f*x])^m)
```

Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.27, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3514, 3042, 3451, 3042, 3224, 3042, 3168, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-2} (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

↓ 3514

$$\frac{\int (\sin(e + fx)a + a)^{m+1} (c - c \sin(e + fx))^{-m-2} ((A + B - C)(2m + 1)c^2 + 2C(2m + 1) \sin(e + fx)c^2) dx}{2ac^2(2m + 1)} + \frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^{m+1} (c - c \sin(e + fx))^{-m-2} ((A + B - C)(2m + 1)c^2 + 2C(2m + 1) \sin(e + fx)c^2) dx}{2ac^2(2m + 1)} + \frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3451

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2cC(2m + 1) \int (\sin(e + fx)a + a)^{m+1} (c - c \sin(e + fx))^{-m-2} dx}{2ac^2(2m + 1)} + \frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3042

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2cC(2m + 1) \int (\sin(e + fx)a + a)^{m+1} (c - c \sin(e + fx))^{-m-2} dx}{2ac^2(2m + 1)} + \frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3224

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2ac^2C(2m + 1) \cos^{-2m}(e + fx)(a \sin(e + fx) + a)^{m+1} (c - c \sin(e + fx))^{-m-2}}{2ac^2(2m + 1)} + \frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3042

$$\frac{\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - 2ac^2C(2m + 1) \cos^{-2m}(e + fx)(a \sin(e + fx) + a)^{m+1} (c - c \sin(e + fx))^{-m-2}}{2ac^2(2m + 1)} + \frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m (c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 3168

$$\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - \frac{2ac^4C(2m+1) \cos^3(e+fx)(a \sin(e+fx)+a)^m(c-c \sin(e+fx))^{\frac{1}{2}}}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m(c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 80

$$\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - \frac{ac^3C2^{-m-\frac{1}{2}}(2m+1) \cos^3(e+fx)(1-\sin(e+fx))^{m+\frac{1}{2}}(a \sin(e+fx))^{\frac{1}{2}}}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m(c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

↓ 79

$$\frac{c^2(2m+1)(A+B+C) \cos(e+fx)(a \sin(e+fx)+a)^{m+1}(c-c \sin(e+fx))^{-m-2}}{f(2m+3)} - \frac{ac^2C2^{\frac{1}{2}-m}(2m+1) \cos^3(e+fx)(1-\sin(e+fx))^{m+\frac{1}{2}}(a \sin(e+fx))^{\frac{1}{2}}}{2ac^2(2m+1)}$$

$$\frac{(A - B + C) \cos(e + fx)(a \sin(e + fx) + a)^m(c - c \sin(e + fx))^{-m-1}}{2cf(2m + 1)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
((A - B + C)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1 - m))/(2*c*f*(1 + 2*m)) + ((c^2*(A + B + C)*(1 + 2*m)*Cos[e + f*x]*(a + a*Sin[e + f*x])^(1 + m)*(c - c*Sin[e + f*x])^(-2 - m))/(f*(3 + 2*m)) - (2^(1/2 - m)*a*c^2*C*(1 + 2*m)*Cos[e + f*x]^3*Hypergeometric2F1[(3 + 2*m)/2, (3 + 2*m)/2, (5 + 2*m)/2, (1 + Sin[e + f*x])/2]*(1 - Sin[e + f*x])^(1/2 + m)*(a + a*Sin[e + f*x])^m*(c - c*Sin[e + f*x])^(-1/2 + (-3 - 2*m)/2)*(c + c*Sin[e + f*x])^((-3 - 2*m)/2 + (3 + 2*m)/2))/(f*(3 + 2*m)))/(2*a*c^2*(1 + 2*m))
```


Defintions of rubi rules used

- rule 79 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`
- rule 80 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3168 `Int[(cos[(e_) + (f_)*(x)]*(g_))^(p_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)), x_Symbol] := Simp[a^2*((g*Cos[e + f*x])^(p + 1)/(f*g*(a + b*Sin[e + f*x])^((p + 1)/2)*(a - b*Sin[e + f*x])^((p + 1)/2))) Subst[Int[(a + b*x)^(m + (p - 1)/2)*(a - b*x)^((p - 1)/2), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, e, f, g, m, p}, x] && EqQ[a^2 - b^2, 0] && !IntegerQ[m]`
- rule 3224 `Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[a^IntPart[m]*c^IntPart[m]*(a + b*Sin[e + f*x])^FracPart[m]*((c + d*Sin[e + f*x])^FracPart[m]/Cos[e + f*x]^(2*FracPart[m])) Int[Cos[e + f*x]^(2*m)*(c + d*Sin[e + f*x])^(n - m), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0] && (FractionQ[m] || !FractionQ[n])`

rule 3451

```

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b - a*B)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^n/(
a*f*(2*m + 1))), x] + Simp[(a*B*(m - n) + A*b*(m + n + 1))/(a*b*(2*m + 1))
  Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[
{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 - b^2, 0
] && (LtQ[m, -2^(-1)] || (ILtQ[m + n, 0] && !SumSimplerQ[n, 1])) && NeQ[2*
m + 1, 0]

```

rule 3514

```

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(a*A - b*B + a*C)*Cos[e + f*x]*(a + b*
Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(2*b*c*f*(2*m + 1))), x] - Si
mp[1/(2*b*c*d*(2*m + 1))  Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e +
f*x])^n*Simp[A*(c^2*(m + 1) + d^2*(2*m + n + 2)) - B*c*d*(m - n - 1) - C*(c
^2*m - d^2*(n + 1)) + d*((A*c + B*d)*(m + n + 2) - c*C*(3*m - n))*Sin[e + f
*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && EqQ[b*c +
a*d, 0] && EqQ[a^2 - b^2, 0] && (LtQ[m, -2^(-1)] || (EqQ[m + n + 2, 0] &&
NeQ[2*m + 1, 0]))

```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c - c \sin(fx + e))^{-2-m} (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(-c*sin(f*x + e) + c)^(-m - 2), x)`

Sympy [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (a(\sin(e + fx) + 1))^m (-c(\sin(e + fx) - 1))^{-m-2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))^m*(-c*(sin(e + f*x) - 1))^(2-m)*(A + B*sin(e + f*x) + C*sin(e + f*x)**2), x)`

Maxima [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^{-m-2} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin
(f*x+e)^2),x, algorithm="maxima")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
-c*sin(f*x + e) + c)^(-m - 2), x)
```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (-c \sin(fx + e) + c)^{-m-2} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin
(f*x+e)^2),x, algorithm="giac")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
-c*sin(f*x + e) + c)^(-m - 2), x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{(c - c \sin(e + fx))^{m+2}} dx$$

input

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(m + 2),x)
```

output

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c - c*sin(e + f*x))^(m + 2), x)
```

Reduce [F]

$$\int (a + a \sin(e + fx))^m (c - c \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \left(\int \frac{(a + a \sin(fx + e))^m}{(-\sin(fx + e)c + c)^m \sin(fx + e)^2 - 2(-\sin(fx + e)c + c)^m \sin(fx + e) + (-\sin(fx + e)c + c)^m} dx \right) a + \left(\int \frac{(a + a \sin(fx + e))^m}{(-\sin(fx + e)c + c)^m \sin(fx + e)^2} dx \right)$$

input

```
int((a+a*sin(f*x+e))^m*(c-c*sin(f*x+e))^(m-2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

output

```
(int((sin(e + f*x)*a + a)**m/((- sin(e + f*x)*c + c)**m*sin(e + f*x)**2 - 2*(- sin(e + f*x)*c + c)**m*sin(e + f*x) + (- sin(e + f*x)*c + c)**m),x)*a + int(((sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/((- sin(e + f*x)*c + c)**m*sin(e + f*x)**2 - 2*(- sin(e + f*x)*c + c)**m*sin(e + f*x) + (- sin(e + f*x)*c + c)**m),x)*c + int(((sin(e + f*x)*a + a)**m*sin(e + f*x))/((- sin(e + f*x)*c + c)**m*sin(e + f*x)**2 - 2*(- sin(e + f*x)*c + c)**m*sin(e + f*x) + (- sin(e + f*x)*c + c)**m),x)*b)/c**2
```

3.24 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^n (A + B \sin(e +$

Optimal result	245
Mathematica [F]	246
Rubi [A] (warning: unable to verify)	246
Maple [F]	250
Fricas [F]	251
Sympy [F(-1)]	251
Maxima [F]	251
Giac [F]	252
Mupad [F(-1)]	252
Reduce [F]	253

Optimal result

Integrand size = 45, antiderivative size = 373

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= -\frac{C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{1+n}}{df(2 + m + n)}$$

$$- \frac{2^{\frac{1}{2}+m} a(c(C + 2Cm) + d(C(1 - m + n) + A(2 + m + n) - B(2 + m + n))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, -n, \frac{1}{2}, \frac{1}{2} - m, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right)}{df(2 + m + n)}$$

$$+ \frac{2^{\frac{3}{2}+m} (cC(1 + m) - d(Cm + B(2 + m + n))) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, -n, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right)}{df(2 + m + n)}$$

output

```
-C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1+n)/d/f/(2+m+n)-2^(1/2+m)*a*(c*(2*C*m+C)+d*(C*(1-m+n)+A*(2+m+n)-B*(2+m+n)))*AppellF1(1/2,-n,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*(c+d*sin(f*x+e))^n/d/f/(2+m+n)/(((c+d*sin(f*x+e))/(c+d))^n+2^(3/2+m)*(c*C*(1+m)-d*(C*m+B*(2+m+n)))*AppellF1(1/2,-n,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n/d/f/(2+m+n)/(((c+d*sin(f*x+e))/(c+d))^n)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x]
] + C*Sin[e + f*x]^2), x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x]
] + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.17 (sec) , antiderivative size = 410, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 3524, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) (c + d \sin(e + fx))^n dx$$

↓ 3524

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n (a(Ad(m + n + 2) + C(nd + d + cm)) + a(Cdm - cC(m + 1) + Bd(n + 1))) dx}{\frac{ad(m + n + 2)}{df(m + n + 2)} C \cos(e + fx) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^n (a(Ad(m + n + 2) + C(nd + d + cm)) + a(Cdm - cC(m + 1) + Bd(m + n + 2)))}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}} \frac{ad(m + n + 2)}{df(m + n + 2)}$$

↓ 3466

$$\frac{(Bd(m + n + 2) - cC(m + 1) + Cdm) \int (\sin(e + fx)a + a)^{m+1} (c + d \sin(e + fx))^n dx - a(-Ad(m + n + 2) + Bd(m + n + 2))}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}} \frac{ad(m + n + 2)}{df(m + n + 2)}$$

↓ 3042

$$\frac{(Bd(m + n + 2) - cC(m + 1) + Cdm) \int (\sin(e + fx)a + a)^{m+1} (c + d \sin(e + fx))^n dx - a(-Ad(m + n + 2) + Bd(m + n + 2))}{C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{n+1}} \frac{ad(m + n + 2)}{df(m + n + 2)}$$

↓ 3267

$$\frac{a^2 \cos(e + fx)(Bd(m + n + 2) - cC(m + 1) + Cdm) \int \frac{(\sin(e + fx)a + a)^{m+\frac{1}{2}} (c + d \sin(e + fx))^n d \sin(e + fx)}{\sqrt{a - a \sin(e + fx)}}}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} - \frac{a^3 \cos(e + fx)(-Ad(m + n + 2) + Bd(m + n + 2))}{ad(m + n + 2)}$$

↓ 157

$$\frac{a^2 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(Bd(m + n + 2) - cC(m + 1) + Cdm) \int \frac{\sqrt{2}(\sin(e + fx)a + a)^{m+\frac{1}{2}} (c + d \sin(e + fx))^n d \sin(e + fx)}{\sqrt{1 - \sin(e + fx)}}}{\sqrt{2}f(a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} - \frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)}{ad(m + n + 2)}$$

↓ 27

$$\frac{a^2 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(Bd(m + n + 2) - cC(m + 1) + Cdm) \int \frac{(\sin(e + fx)a + a)^{m+\frac{1}{2}} (c + d \sin(e + fx))^n d \sin(e + fx)}{\sqrt{1 - \sin(e + fx)}}}{f(a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} - \frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)}{ad(m + n + 2)}$$

↓ 156

$$\frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (Bd(m+n+2) - cC(m+1) + Cdm) (c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)^n}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}}$$

$$\frac{C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

↓ 155

$$\frac{\sqrt{2a} \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx)+a)^{m+1} (Bd(m+n+2) - cC(m+1) + Cdm) (c+d \sin(e+fx))^n \left(\frac{c+d \sin(e+fx)}{c-d}\right)^{-n} \text{AppellF1}\left(m+\frac{3}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)}{f(2m+3)(a-a \sin(e+fx))}$$

$$\frac{C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{n+1}}{df(m+n+2)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
-((C*cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(1 + n))/(d*f*(2 + m + n))) + (-((Sqrt[2]*a^2*(C*d*m - c*C*(1 + m) - A*d*(2 + m + n) + B*d*(2 + m + n) - C*(d + c*m + d*n))*AppellF1[1/2 + m, 1/2, -n, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n)/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n) + (Sqrt[2]*a*(C*d*m - c*C*(1 + m) + B*d*(2 + m + n))*AppellF1[3/2 + m, 1/2, -n, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*(c + d*Sin[e + f*x])^n)/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*((c + d*Sin[e + f*x])/(c - d))^n))/(a*d*(2 + m + n))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3524

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^n (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**n*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
d*sin(f*x + e) + c)^n, x)
```

Giac [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin^2(fx + e) + B \sin(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^n dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+
e)^2),x, algorithm="giac")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
d*sin(f*x + e) + c)^n, x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (C \sin^2(e + fx) + B \sin(e + fx) + A) dx$$

input

```
int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*
sin(e + f*x)^2),x)
```

output

```
int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*
sin(e + f*x)^2), x)
```

Reduce [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \left(\int (\sin(fx + e) d + c)^n (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) c$$

$$+ \left(\int (\sin(fx + e) d + c)^n (a + a \sin(fx + e))^m \sin(fx + e) dx \right) b$$

$$+ \left(\int (\sin(fx + e) d + c)^n (a + a \sin(fx + e))^m dx \right) a$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),
x)
```

output

```
int((sin(e + f*x)*d + c)**n*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2,x)*c +
int((sin(e + f*x)*d + c)**n*(sin(e + f*x)*a + a)**m*sin(e + f*x),x)*b + i
nt((sin(e + f*x)*d + c)**n*(sin(e + f*x)*a + a)**m,x)*a
```

3.25 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{-2-m} (A + B \sin(e+fx) + C \sin^2(e+fx)) dx$

Optimal result	254
Mathematica [F]	255
Rubi [A] (warning: unable to verify)	255
Maple [F]	260
Fricas [F]	260
Sympy [F(-1)]	261
Maxima [F]	261
Giac [F]	261
Mupad [F(-1)]	262
Reduce [F]	262

Optimal result

Integrand size = 49, antiderivative size = 400

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \frac{(c^2 C - Bcd + Ad^2) \cos(e + fx) (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-1-m}}{d(c^2 - d^2) f(1 + m)}$$

$$- \frac{2^{\frac{1}{2}+m} a (cd(A + C + Am + Bm + Cm) - c^2(C + 2Cm) - d^2(Am + B(1 + m) - C(1 + m))) \cos(e + fx)}{d(c + d)f}$$

$$- \frac{2^{\frac{3}{2}+m} C \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, 1 + m, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx)), \frac{d(1 - \sin(e + fx))}{c + d}\right) \cos(e + fx) (1 + \sin(e + fx))}{d(c + d)f}$$

output

```
(A*d^2-B*c*d+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1-m)/
d/(c^2-d^2)/f/(1+m)-2^(1/2+m)*a*(c*d*(A*m+B*m+C*m+A+C)-c^2*(2*C*m+C)-d^2*(
A*m+B*(1+m)-C*(1+m)))*cos(f*x+e)*hypergeom([1/2, 1/2-m],[3/2],(c-d)*(1-sin
(f*x+e))/(2*c+2*d*sin(f*x+e)))*(a+a*sin(f*x+e))^(1-m)*((c+d)*(1+sin(f*x+e
)))/(c+d*sin(f*x+e))^(1/2-m)/(c-d)/d/(c+d)^2/f/(1+m)/((c+d*sin(f*x+e))^m)-
2^(3/2+m)*C*AppellF1(1/2,1+m,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin
(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^m*((c+d*sin(f
*x+e))/(c+d))^m/d/(c+d)/f/((c+d*sin(f*x+e))^m)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.38 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.10, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$, Rules used = {3042, 3522, 25, 3042, 3466, 3042, 3267, 142, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2) (c + d \sin(e + fx))^{-m-2} dx$$

$$\downarrow \text{3522}$$

$$\frac{\cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{-m-1}}{df(m+1)(c^2 - d^2)}$$

$$\int \frac{-(\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{-m-1} (a(Ad(mc + c - dm) + (cC - Bd)(md + d - cm)) + aC(c^2 - d^2))}{ad(m+1)(c^2 - d^2)} dx$$

$$\downarrow \text{25}$$

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{-m-1} (a(Ad(mc+c-dm)+(cC-Bd)(md+d-cm))+aC(c^2-d^2) \cos(e+fx) (Ad^2-Bcd+c^2C) (a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)} \frac{ad(m+1)(c^2-d^2)}{df(m+1)(c^2-d^2)}$$

↓ 3042

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{-m-1} (a(Ad(mc+c-dm)+(cC-Bd)(md+d-cm))+aC(c^2-d^2) \cos(e+fx) (Ad^2-Bcd+c^2C) (a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)} \frac{ad(m+1)(c^2-d^2)}{df(m+1)(c^2-d^2)}$$

↓ 3466

$$\frac{a(cd(Am+A+Bm+C)-d^2(Am+B(m+1)-C(m+1))-(c^2(2Cm+C))) \int (\sin(e+fx)a+a)^m \cos(e+fx) (Ad^2-Bcd+c^2C) (a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)} \frac{ad(m+1)(c^2-d^2)}{df(m+1)(c^2-d^2)}$$

↓ 3042

$$\frac{a(cd(Am+A+Bm+C)-d^2(Am+B(m+1)-C(m+1))-(c^2(2Cm+C))) \int (\sin(e+fx)a+a)^m \cos(e+fx) (Ad^2-Bcd+c^2C) (a\sin(e+fx)+a)^m (c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)} \frac{ad(m+1)(c^2-d^2)}{df(m+1)(c^2-d^2)}$$

↓ 3267

$$\frac{a^3 \cos(e+fx) (cd(Am+A+Bm+C)-d^2(Am+B(m+1)-C(m+1))-(c^2(2Cm+C))) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^{-m-1}}{\sqrt{a-a\sin(e+fx)}} d\sin(e+fx)}{f\sqrt{a-a\sin(e+fx)}\sqrt{a\sin(e+fx)+a}} \frac{ad(m+1)(c^2-d^2)}{df(m+1)(c^2-d^2)}$$

↓ 142

$$\frac{a^2 C(m+1)(c^2-d^2) \cos(e+fx) \int \frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}} (c+d\sin(e+fx))^{-m-1}}{\sqrt{a-a\sin(e+fx)}} d\sin(e+fx)}{f\sqrt{a-a\sin(e+fx)}\sqrt{a\sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}} \cos(e+fx) (a\sin(e+fx)+a)^{m-1} (cd(Am+A+Bm+C)-d^2(Am+B(m+1)-C(m+1))-(c^2(2Cm+C)))}{df(m+1)(c^2-d^2)}$$

a

↓ 157

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{\sqrt{2}(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{\sqrt{2}f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(a\sin(e+fx)+a)^m}{\sqrt{2}f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 27

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}(c+d\sin(e+fx))^{-m-1}d\sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} - \frac{a^2 2^{m+\frac{1}{2}}\cos(e+fx)(a\sin(e+fx)+a)^m}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 156

$$\frac{a^2 C(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\int\frac{(\sin(e+fx)a+a)^{m+\frac{1}{2}}\left(\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}\right)^{-m-1}d\sin(e+fx)}{\sqrt{1-\sin(e+fx)}}}{f(c-d)(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

↓ 155

$$\frac{\sqrt{2}aC(m+1)(c^2-d^2)\sqrt{1-\sin(e+fx)}\cos(e+fx)(a\sin(e+fx)+a)^{m+1}(c+d\sin(e+fx))^{-m}\left(\frac{c+d\sin(e+fx)}{c-d}\right)^m\text{AppellF1}\left(m+\frac{3}{2},\frac{1}{2},m+1,m+\frac{5}{2},\frac{1}{2}\right)}{f(2m+3)(c-d)(a-a\sin(e+fx))}$$

$$\frac{\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m(c+d\sin(e+fx))^{-m-1}}{df(m+1)(c^2-d^2)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-2 - m)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]
```

output

```
((c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(-1 - m))/(d*(c^2 - d^2)*f*(1 + m)) + (-((2^(1/2 + m)*a^2*(c*d*(A + C + A*m + B*m + C*m) - c^2*(C + 2*C*m) - d^2*(A*m + B*(1 + m) - C*(1 + m)))*Cos[e + f*x]*Hypergeometric2F1[1/2, 1/2 - m, 3/2, ((c - d)*(1 - Sin[e + f*x]))/(2*(c + d*Sin[e + f*x]))]*(a + a*Sin[e + f*x])^(-1 + m)*((c + d)*(1 + Sin[e + f*x]))/(c + d*Sin[e + f*x])^(1/2 - m))/((c + d)*f*(c + d*Sin[e + f*x])^m) + (Sqrt[2]*a*C*(c^2 - d^2)*(1 + m)*AppellF1[3/2 + m, 1/2, 1 + m, 5/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x]))/(c - d)]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*((c + d*Sin[e + f*x]))/(c - d)^m)/((c - d)*f*(3 + 2*m)*(a - a*Sin[e + f*x])*(c + d*Sin[e + f*x])^m)/(a*d*(c^2 - d^2)*(1 + m))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 142

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_)), x_] := Simp[((a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((b*e - a*f)*(m + 1)))*Hypergeometric2F1[m + 1, -n, m + 2, -(d*e - c*f)*((a + b*x)/((b*c - a*d)*(e + f*x)))]/((b*e - a*f)*((c + d*x)/((b*c - a*d)*(e + f*x))))^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m + n + p + 2, 0] && !IntegerQ[n]
```

rule 155

```
Int[((a_) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_)), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p)*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])
```

rule 156

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]
)*(b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Si
mp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]
```

rule 157

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]
)*(b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c
- a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c,
d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] &
& !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !Si
mplerQ[e + f*x, a + b*x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (
f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e
+ f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*
x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m
, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] &&
!IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) +
(f_)*(x_)])*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x], x
] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ
[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3522

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(-(c^2*C - B*c*d + A*d^2))*Cos[e + f*x
]*(a + b*Ssin[e + f*x])^m*((c + d*Ssin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 -
d^2))), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Ssin[e + f*x])^m*
(c + d*Ssin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + (c*C - B*d)*
(a*c*m + b*d*(n + 1)) + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*
(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m},
x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ
[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])
```

Maple [F]

$$\int (a + a \sin(fx + e))^m (c + d \sin(fx + e))^{-2-m} (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)
^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)
^2),x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) (a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin
(f*x+e)^2),x, algorithm="fricas")
```

output

```
integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)
^m*(d*sin(f*x + e) + c)^(-m - 2), x)
```

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(-2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))**(-2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)**(-m - 2), x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m (d \sin(fx + e) + c)^{-m-2} dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))**(-2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m*(
d*sin(f*x + e) + c)^(-m - 2), x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{(c + d \sin(e + fx))^{m+2}} dx$$

input

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c +
d*sin(e + f*x))^(m + 2),x)
```

output

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c +
d*sin(e + f*x))^(m + 2), x)
```

Reduce [F]

$$\begin{aligned} & \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{-2-m} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx \\ &= \left(\int \frac{(a + a \sin(fx + e))^m}{(\sin(fx + e)d + c)^m \sin(fx + e)^2 d^2 + 2(\sin(fx + e)d + c)^m \sin(fx + e)cd + (\sin(fx + e)d + c)} \right. \\ & \quad + \left(\int \frac{(a + a \sin(fx + e))^m \sin(fx + e)^2}{(\sin(fx + e)d + c)^m \sin(fx + e)^2 d^2 + 2(\sin(fx + e)d + c)^m \sin(fx + e)cd + (\sin(fx + e)d + c)} \right. \\ & \quad \left. \left. + \left(\int \frac{(a + a \sin(fx + e))^m \sin(fx + e)}{(\sin(fx + e)d + c)^m \sin(fx + e)^2 d^2 + 2(\sin(fx + e)d + c)^m \sin(fx + e)cd + (\sin(fx + e)d + c)} \right) \right) \right) \end{aligned}$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(2-m)*(A+B*sin(f*x+e)+C*sin(f*x+e)
)^2),x)
```

output

```
int((sin(e + f*x)*a + a)**m/((sin(e + f*x)*d + c)**m*sin(e + f*x)**2*d**2
+ 2*(sin(e + f*x)*d + c)**m*sin(e + f*x)*c*d + (sin(e + f*x)*d + c)**m*c**
2),x)*a + int(((sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/((sin(e + f*x)*d +
c)**m*sin(e + f*x)**2*d**2 + 2*(sin(e + f*x)*d + c)**m*sin(e + f*x)*c*d +
(sin(e + f*x)*d + c)**m*c**2),x)*c + int(((sin(e + f*x)*a + a)**m*sin(e +
f*x))/((sin(e + f*x)*d + c)**m*sin(e + f*x)**2*d**2 + 2*(sin(e + f*x)*d +
c)**m*sin(e + f*x)*c*d + (sin(e + f*x)*d + c)**m*c**2),x)*b
```


3.26 $\int (a+a \sin(e+fx))^m (c+d \sin(e+fx))^{3/2} (A + B \sin(e$

Optimal result	264
Mathematica [F]	265
Rubi [A] (warning: unable to verify)	265
Maple [F]	269
Fricas [F]	270
Sympy [F(-1)]	270
Maxima [F]	271
Giac [F]	271
Mupad [F(-1)]	272
Reduce [F]	272

Optimal result

Integrand size = 47, antiderivative size = 392

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx =$$

$$\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m (c + d \sin(e + fx))^{5/2}}{df(7 + 2m)}$$

$$- \frac{2^{\frac{1}{2}+m} a(c + d)(2c(C + 2Cm) - d(7B - 5C + 2Bm + 2Cm - A(7 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, -\frac{3}{2}, \frac{3}{2}, \frac{1}{2}\right)}{df(7 + 2m)}$$

$$+ \frac{2^{\frac{3}{2}+m} (c + d)(2cC(1 + m) - d(2Cm + B(7 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, -\frac{3}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right), \frac{d(1 - \sin(e + fx))}{c + d}}{df(7 + 2m) \sqrt{\frac{c + d \sin(e + fx)}{c + d}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(5/2)/d/f/(7+2*m)-2^(1/2+m)*a*(c+d)*(2*c*(2*C*m+C)-d*(7*B-5*C+2*B*m+2*C*m-A*(7+2*m)))*AppellF1(1/2,-3/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*(c+d*sin(f*x+e))^(1/2)/d/f/(7+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)+2^(3/2+m)*(c+d)*(2*c*C*(1+m)-d*(2*C*m+B*(7+2*m)))*AppellF1(1/2,-3/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(7+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.27 (sec) , antiderivative size = 425, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$, Rules used = {3042, 3524, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

↓ 3042

$$\int (a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

↓ 3524

$$\frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m (c + d \sin(e + fx))^{3/2} (a(Ad(2m + 7) + C(5d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd)) dx}{\frac{ad(2m + 7)}{df(2m + 7)} \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{5/2}}{df(2m + 7)}}$$

↓ 27

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} (a(Ad(2m+7)+C(5d+2cm)) + a(2Cdm-2cC(m+1)+Bd(2m+7)))}{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{5/2} df(2m+7)} \frac{ad(2m+7)}{df(2m+7)} \downarrow 3042$$

$$\frac{\int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} (a(Ad(2m+7)+C(5d+2cm)) + a(2Cdm-2cC(m+1)+Bd(2m+7)))}{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{5/2} df(2m+7)} \frac{ad(2m+7)}{df(2m+7)} \downarrow 3466$$

$$\frac{a(2c(2Cm+C)-d(-A(2m+7)+2Bm+7B+2Cm-5C)) \int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} dx + 2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{5/2} df(2m+7)}{ad(2m+7)} \downarrow 3042$$

$$\frac{a(2c(2Cm+C)-d(-A(2m+7)+2Bm+7B+2Cm-5C)) \int (\sin(e+fx)a+a)^m (c+d\sin(e+fx))^{3/2} dx + 2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{5/2} df(2m+7)}{ad(2m+7)} \downarrow 3267$$

$$\frac{a^3 \cos(e+fx)(2c(2Cm+C)-d(-A(2m+7)+2Bm+7B+2Cm-5C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^{3/2} d\sin(e+fx)}{f \sqrt{a-a\sin(e+fx)} \sqrt{a\sin(e+fx)+a}} + a^2 \cos(e+fx)(Bd(2m+7))}{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{5/2} df(2m+7)} \downarrow 157$$

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(2c(2Cm+C)-d(-A(2m+7)+2Bm+7B+2Cm-5C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d\sin(e+fx))^{3/2} d\sin(e+fx)}{\sqrt{2f(a-a\sin(e+fx))} \sqrt{a\sin(e+fx)+a}} + a^2 \cos(e+fx)(Bd(2m+7))}{2C \cos(e+fx)(a \sin(e+fx)+a)^m (c+d\sin(e+fx))^{5/2} df(2m+7)} \downarrow$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} (c+d \sin(e+fx))^{3/2}}{\sqrt{1-\sin(e+fx)}} d \sin(e+fx)}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{a^2}{ad(2m+7)}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 156

$$\frac{a^3 (c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C)) \sqrt{c+d \sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \left(\frac{c}{c-d} + \frac{d \sin(e+fx)}{c-d}\right)}{\sqrt{1-\sin(e+fx)}}}{f(a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a} \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

↓ 155

$$\frac{\sqrt{2} a^2 (c-d) \sqrt{1-\sin(e+fx)} \cos(e+fx) (a \sin(e+fx) + a)^m (2c(2Cm+C) - d(-A(2m+7) + 2Bm + 7B + 2Cm - 5C)) \sqrt{c+d \sin(e+fx)} \text{AppellF1}\left(m, \frac{1}{2}, -\frac{3}{2}, \frac{3}{2} + m, \frac{1 + \sin(e+fx)}{2}, -\frac{(d(1 + \sin(e+fx)))}{(c-d)} \cos(e+fx) \sqrt{1 - \sin(e+fx)}\right)}{f(2m+1)(a-a \sin(e+fx)) \sqrt{\frac{c+d \sin(e+fx)}{c-d}}}$$

$$\frac{2C \cos(e+fx) (a \sin(e+fx) + a)^m (c+d \sin(e+fx))^{5/2}}{df(2m+7)}$$

input `Int[(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2)*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]`

output `(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(5/2))/(d*f*(7 + 2*m)) + ((Sqrt[2]*a^2*(c - d)*(2*c*(C + 2*C*m) - d*(7*B - 5*C + 2*B*m + 2*C*m - A*(7 + 2*m)))*AppellF1[1/2 + m, 1/2, -3/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (Sqrt[2]*a*(c - d)*(2*C*d*m - 2*c*C*(1 + m) + B*d*(7 + 2*m))*AppellF1[3/2 + m, 1/2, -3/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(7 + 2*m))`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplifierQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplifierQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplifierQ[c + d*x, a + b*x] && !SimplifierQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3524

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin (fx + e))^m (c + d \sin (fx + e))^{\frac{3}{2}} (A + B \sin (fx + e) + C \sin (fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

Fricas [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin^2(fx + e) + B \sin(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-((C*c + B*d)*cos(f*x + e)^2 - (A + C)*c - B*d + (C*d*cos(f*x + e))^2 - B*c - (A + C)*d)*sin(f*x + e)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

Sympy [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Timed out`

Maxima [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (C \sin(fx + e)^2 + B \sin(fx + e) + A)(d \sin(fx + e) + c)^{3/2} (a \sin(fx + e) + a)^m$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(d*sin(f*x + e) + c)^(3/2)*(a*sin(f*x + e) + a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx$$

input `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)`

output `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(3/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)`

Reduce [F]

$$\int (a + a \sin(e + fx))^m (c + d \sin(e + fx))^{3/2} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx = \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^3 dx \right) cd + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) bd + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) c^2 + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e) dx \right) ad + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e) dx \right) bc + \left(\int \sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m dx \right) ac$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

output

```
int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**3,x)*c*
d + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2,x
)*b*d + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)*
*2,x)*c**2 + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e +
f*x),x)*a*d + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e +
f*x),x)*b*c + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m,x)*a*c
```

3.27 $\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+B \sin(e$

Optimal result	274
Mathematica [F]	275
Rubi [A] (warning: unable to verify)	275
Maple [F]	279
Fricas [F]	280
Sympy [F]	280
Maxima [F]	281
Giac [F]	281
Mupad [F(-1)]	282
Reduce [F]	282

Optimal result

Integrand size = 47, antiderivative size = 386

$$\int (a+a \sin(e+fx))^m \sqrt{c+d \sin(e+fx)}(A+B \sin(e+fx)+C \sin^2(e+fx)) dx$$

$$= -\frac{2C \cos(e+fx)(a+a \sin(e+fx))^m(c+d \sin(e+fx))^{3/2}}{df(5+2m)}$$

$$-\frac{2^{\frac{1}{2}+m}a(2c(C+2Cm)-d(5B-3C+2Bm+2Cm-A(5+2m))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}-m, -\frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1-\sin(e+fx))\right)}{df(5+2m)}$$

$$+\frac{2^{\frac{3}{2}+m}(2cC(1+m)-d(2Cm+B(5+2m))) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2}-m, -\frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1-\sin(e+fx))\right), \frac{d(1-\sin(e+fx))}{c+d}}{df(5+2m)\sqrt{\frac{c+d \sin(e+fx)}{c+d}}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(3/2)/d/f/(5+2*m)-2^(1/2+m)*a*(2*c*(2*C*m+C)-d*(5*B-3*C+2*B*m+2*C*m-A*(5+2*m)))*AppellF1(1/2,-1/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*(c+d*sin(f*x+e))^(1/2)/d/f/(5+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)+2^(3/2+m)*(2*c*C*(1+m)-d*(2*C*m+B*(5+2*m)))*AppellF1(1/2,-1/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(5+2*m)/((c+d*sin(f*x+e))/(c+d))^(1/2)
```

Mathematica [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input

```
Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
Integrate[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.27 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$, Rules used = {3042, 3524, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a \sin(e + fx) + a)^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin(e + fx)^2) dx$$

$$\downarrow \text{3524}$$

$$\frac{2 \int \frac{1}{2} (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 5)))}{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}} df(2m + 5)}{ad(2m + 5)}$$

$$\downarrow \text{27}$$

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 5)))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3042

$$\frac{\int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} (a(Ad(2m + 5) + C(3d + 2cm)) + a(2Cdm - 2cC(m + 1) + Bd(2m + 5)))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3466

$$\frac{a(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx + (ad(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3042

$$\frac{a(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int (\sin(e + fx)a + a)^m \sqrt{c + d \sin(e + fx)} dx + (ad(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 3267

$$\frac{a^3 \cos(e + fx)(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{(\sin(e + fx)a + a)^{m - \frac{1}{2}} \sqrt{c + d \sin(e + fx)}}{\sqrt{a - a \sin(e + fx)}} d \sin(e + fx)}{f \sqrt{a - a \sin(e + fx)} \sqrt{a \sin(e + fx) + a}} + \frac{a^2 \cos(e + fx)(Bd(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 157

$$\frac{a^3 \sqrt{1 - \sin(e + fx)} \cos(e + fx)(2c(2Cm + C) - d(-A(2m + 5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{\sqrt{2}(\sin(e + fx)a + a)^{m - \frac{1}{2}} \sqrt{c + d \sin(e + fx)}}{\sqrt{1 - \sin(e + fx)}} d \sin(e + fx)}{\sqrt{2} f (a - a \sin(e + fx)) \sqrt{a \sin(e + fx) + a}} + \frac{a^2 \cos(e + fx)(Bd(2m + 5))}{ad(2m + 5)} + \frac{2C \cos(e + fx)(a \sin(e + fx) + a)^m (c + d \sin(e + fx))^{3/2}}{df(2m + 5)}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{c+d\sin(e+fx)}}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C))}{df(2m+5)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C)) \sqrt{c+d\sin(e+fx)} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}} \sqrt{\frac{c}{c-d} + \frac{d\sin(e+fx)}{c-d}}}{\sqrt{1-\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a} \sqrt{\frac{c+d\sin(e+fx)}{c-d}}} + \frac{2C \cos(e+fx) (a\sin(e+fx) + a)^m (c+d\sin(e+fx))^{3/2}}{df(2m+5)}$$

↓ 155

$$\frac{\sqrt{2a^2} \sqrt{1-\sin(e+fx)} \cos(e+fx) (a\sin(e+fx)+a)^m (2c(2Cm+C) - d(-A(2m+5) + 2Bm + 5B + 2Cm - 3C)) \sqrt{c+d\sin(e+fx)} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, -\frac{1}{2}, \frac{3}{2}+m, \frac{1+\sin(e+fx)}{2}, -\frac{d(1+\sin(e+fx))}{c-d}\right)}{f(2m+1)(a-a\sin(e+fx))\sqrt{\frac{c+d\sin(e+fx)}{c-d}}} + \frac{2C \cos(e+fx) (a\sin(e+fx) + a)^m (c+d\sin(e+fx))^{3/2}}{df(2m+5)}$$

input

```
Int[(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]]*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(3/2))/(d*f*(5 + 2*m)) + ((Sqrt[2]*a^2*(2*c*(C + 2*C*m) - d*(5*B - 3*C + 2*B*m + 2*C*m - A*(5 + 2*m)))*AppellF1[1/2 + m, 1/2, -1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]) + (Sqrt[2]*a*(2*C*d*m - 2*c*C*(1 + m) + B*d*(5 + 2*m))*AppellF1[3/2 + m, 1/2, -1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[c + d*Sin[e + f*x]])/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[(c + d*Sin[e + f*x])/(c - d)]))/(a*d*(5 + 2*m))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*b*((e + f*x)/(b*e - a*f)))^FracPart[p] Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*b*((c + d*x)/(b*c - a*d)))^FracPart[n] Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3524

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int (a + a \sin(fx + e))^m \sqrt{c + d \sin(fx + e)} (A + B \sin(fx + e) + C \sin(fx + e)^2) dx$$

input

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```


Fricas [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m, x)`

Sympy [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a(\sin(e + fx) + 1))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

input `integrate((a+a*sin(f*x+e))**m*(c+d*sin(f*x+e))**(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*sqrt(c + d*sin(e + f*x))*(A + B*sin(e + f*x) + C*sin(e + f*x)**2), x)`

Maxima [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*sqrt(d*sin(f*x + e) + c) *(a*sin(f*x + e) + a)^m, x)`

Giac [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (C \sin(fx + e)^2 + B \sin(fx + e) + A) \sqrt{d \sin(fx + e) + c} (a \sin(fx + e) + a)^m dx$$

input `integrate((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*sqrt(d*sin(f*x + e) + c) *(a*sin(f*x + e) + a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx$$

input `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2),x)`

output `int((a + a*sin(e + f*x))^m*(c + d*sin(e + f*x))^(1/2)*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)`

Reduce [F]

$$\int (a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)} (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \left(\int \sqrt{\sin(fx + e)d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2 dx \right) c$$

$$+ \left(\int \sqrt{\sin(fx + e)d + c} (a + a \sin(fx + e))^m \sin(fx + e) dx \right) b$$

$$+ \left(\int \sqrt{\sin(fx + e)d + c} (a + a \sin(fx + e))^m dx \right) a$$

input `int((a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)`

output `int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2,x)*c + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x),x)*b + int(sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m,x)*a`

3.28
$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx$$

Optimal result	283
Mathematica [F]	284
Rubi [A] (warning: unable to verify)	284
Maple [F]	288
Fricas [F]	289
Sympy [F]	289
Maxima [F]	290
Giac [F]	290
Mupad [F(-1)]	291
Reduce [F]	291

Optimal result

Integrand size = 47, antiderivative size = 386

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= -\frac{2C \cos(e + fx)(a + a \sin(e + fx))^m \sqrt{c + d \sin(e + fx)}}{df(3 + 2m)}$$

$$- \frac{2^{\frac{1}{2}+m} a(2c(C + 2Cm) - d(3B - C + 2Bm + 2Cm - A(3 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right)}{df(3 + 2m)\sqrt{c + d \sin(e + fx)}}$$

$$+ \frac{2^{\frac{3}{2}+m} (2cC(1 + m) - d(2Cm + B(3 + 2m))) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right), \frac{d(1 - \sin(e + fx))}{c + d}}{df(3 + 2m)\sqrt{c + d \sin(e + fx)}}$$

output

```
-2*C*cos(f*x+e)*(a+a*sin(f*x+e))^m*(c+d*sin(f*x+e))^(1/2)/d/f/(3+2*m)-2^(1/2+m)*a*(2*c*(2*C*m+C)-d*(3*B-C+2*B*m+2*C*m-A*(3+2*m)))*AppellF1(1/2,1/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/f/(3+2*m)/(c+d*sin(f*x+e))^(1/2)+2^(3/2+m)*(2*c*C*(1+m)-d*(2*C*m+B*(3+2*m)))*AppellF1(1/2,1/2,-1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^m*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/f/(3+2*m)/(c+d*sin(f*x+e))^(1/2)
```

Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/Sqrt[c + d*Sin[e + f*x]],x]
```

output

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/Sqrt[c + d*Sin[e + f*x]], x]
```

Rubi [A] (warning: unable to verify)

Time = 1.27 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$, Rules used = {3042, 3524, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$\downarrow 3042$$

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{\sqrt{c + d \sin(e + fx)}} dx$$

$$\downarrow 3524$$

$$2 \int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+a(2Cdm-2cC(m+1)+Bd(2m+3)) \sin(e+fx))}{2\sqrt{c+d \sin(e+fx)}} dx$$

$$\frac{ad(2m+3)}{2C \cos(e+fx)(a \sin(e+fx) + a)^m \sqrt{c+d \sin(e+fx)}}$$

$$\frac{df(2m+3)}{\downarrow 27}$$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+a(2Cdm-2cC(m+1)+Bd(2m+3)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{\frac{ad(2m+3)}{df(2m+3)} \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}}$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a(Ad(2m+3)+C(d+2cm))+a(2Cdm-2cC(m+1)+Bd(2m+3)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{\frac{ad(2m+3)}{df(2m+3)} \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}}$$

↓ 3466

$$\frac{a(2c(2Cm+C)-d(-A(2m+3)+2Bm+3B+2Cm-C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (Bd(2m+3)-2cC(m+1))}{\frac{ad(2m+3)}{df(2m+3)} \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}}$$

↓ 3042

$$\frac{a(2c(2Cm+C)-d(-A(2m+3)+2Bm+3B+2Cm-C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx + (Bd(2m+3)-2cC(m+1))}{\frac{ad(2m+3)}{df(2m+3)} \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(2c(2Cm+C)-d(-A(2m+3)+2Bm+3B+2Cm-C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + a^2 \cos(e+fx)(Bd(2m+3)-2cC(m+1))}{\frac{ad(2m+3)}{df(2m+3)} \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(2c(2Cm+C)-d(-A(2m+3)+2Bm+3B+2Cm-C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) + a^2 \sqrt{1-\sin(e+fx)}(Bd(2m+3)-2cC(m+1))}{\frac{ad(2m+3)}{df(2m+3)} \frac{2C \cos(e+fx)(a \sin(e+fx)+a)^m \sqrt{c+d \sin(e+fx)}}{df(2m+3)}}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{c+d\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} + \frac{a^2 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C))}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}}$$

$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m \sqrt{c+d\sin(e+fx)}}{df(2m+3)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{\frac{c}{c-d} + \frac{d\sin(e+fx)}{c-d}}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m \sqrt{c+d\sin(e+fx)}}{df(2m+3)}$$

↓ 155

$$\frac{\sqrt{2a^2} \sqrt{1-\sin(e+fx)} \cos(e+fx) (a\sin(e+fx)+a)^m (2c(2Cm+C) - d(-A(2m+3) + 2Bm + 3B + 2Cm - C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \text{AppellF1}\left(m+\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, m+\frac{1}{2}\right)}{f(2m+1)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}}$$

$$\frac{2C \cos(e+fx)(a\sin(e+fx)+a)^m \sqrt{c+d\sin(e+fx)}}{df(2m+3)}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sqrt[c + d*Sin[e + f*x]],x]
```

output

```
(-2*C*Cos[e + f*x]*(a + a*Sin[e + f*x])^m*Sqrt[c + d*Sin[e + f*x]])/(d*f*(3 + 2*m)) + ((Sqrt[2]*a^2*(2*c*(C + 2*C*m) - d*(3*B - C + 2*B*m + 2*C*m - A*(3 + 2*m)))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d)])/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) + (Sqrt[2]*a*(2*C*d*m - 2*c*C*(1 + m) + B*d*(3 + 2*m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -((d*(1 + Sin[e + f*x]))/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]])*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(3 + 2*m))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3524

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-C)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(m + n + 2))), x] + Simp[1/(b*d*(m + n + 2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*Simp[A*b*d*(m + n + 2) + C*(a*c*m + b*d*(n + 1)) + (C*(a*d*m - b*c*(m + 1)) + b*B*d*(m + n + 2))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && NeQ[m + n + 2, 0]
```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C \sin(fx + e)^2)}{\sqrt{c + d \sin(fx + e)}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e))^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e) + c), x)`

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(1/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/sqrt(c + d*sin(e + f*x)), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e) + c), x)`

Giac [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a \sin(fx + e) + a)^m}{\sqrt{d \sin(fx + e) + c}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/sqrt(d*sin(f*x + e) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx)^2 + B \sin(e + fx) + A)}{\sqrt{c + d \sin(e + fx)}} dx$$

input `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(1/2),x)`

output `int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c + d*sin(e + f*x))^(1/2), x)`

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{\sqrt{c + d \sin(e + fx)}} dx$$

$$= \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)^2}{\sin(fx + e) d + c} dx \right) c$$

$$+ \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)}{\sin(fx + e) d + c} dx \right) b$$

$$+ \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m}{\sin(fx + e) d + c} dx \right) a$$

input `int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(1/2),x)`

output `int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/(sin(e + f*x)*d + c),x)*c + int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x))/(sin(e + f*x)*d + c),x)*b + int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m)/(sin(e + f*x)*d + c),x)*a`

$$3.29 \quad \int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx$$

Optimal result	292
Mathematica [F]	293
Rubi [A] (warning: unable to verify)	293
Maple [F]	297
Fricas [F]	298
Sympy [F]	298
Maxima [F]	298
Giac [F]	299
Mupad [F(-1)]	299
Reduce [F]	300

Optimal result

Integrand size = 47, antiderivative size = 421

$$\int \frac{(a+a \sin(e+fx))^m (A+B \sin(e+fx)+C \sin^2(e+fx))}{(c+d \sin(e+fx))^{3/2}} dx = \frac{2(c^2C - Bcd + Ad^2) \cos(e+fx)(a + \dots)}{d(c^2 - d^2) f \sqrt{c + d \sin(e + \dots)}} + \frac{2^{\frac{1}{2}+m} a(d^2(A+B-C+4Am) - cd(A+B+C+4Bm) + 2c^2(C+2Cm)) \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}\right)}{d(c^2 - d^2) f \sqrt{c + d \sin(e + \dots)}} + \frac{2^{\frac{3}{2}+m} (d(Bc - Ad)(1 + 2m) + C(d^2 - 2c^2(1 + m))) \operatorname{AppellF1}\left(\frac{1}{2}, -\frac{1}{2} - m, \frac{1}{2}, \frac{3}{2}, \frac{1}{2}(1 - \sin(e + fx))\right), \frac{d(1 - \sin(e + fx))}{c + d \sin(e + fx)}}{d(c^2 - d^2) f \sqrt{c + d \sin(e + \dots)}}$$

output

```
2*(A*d^2-B*c*d+C*c^2)*cos(f*x+e)*(a+a*sin(f*x+e))^m/d/(c^2-d^2)/f/(c+d*sin
(f*x+e))^(1/2)+2^(1/2+m)*a*(d^2*(4*A*m+A+B-C)-c*d*(4*B*m+A+B+C)+2*c^2*(2*C
*m+C))*AppellF1(1/2,1/2,1/2-m,3/2,d*(1-sin(f*x+e))/(c+d),1/2-1/2*sin(f*x+e
))*cos(f*x+e)*(1+sin(f*x+e))^(1/2-m)*(a+a*sin(f*x+e))^(-1+m)*((c+d*sin(f*x
+e))/(c+d))^(1/2)/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^(1/2)+2^(3/2+m)*(d*(-A*d+
B*c)*(1+2*m)+C*(d^2-2*c^2*(1+m)))*AppellF1(1/2,1/2,-1/2-m,3/2,d*(1-sin(f*x
+e))/(c+d),1/2-1/2*sin(f*x+e))*cos(f*x+e)*(1+sin(f*x+e))^(-1/2-m)*(a+a*sin
(f*x+e))^m*((c+d*sin(f*x+e))/(c+d))^(1/2)/d/(c^2-d^2)/f/(c+d*sin(f*x+e))^(
1/2)
```

Mathematica [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/(c + d*Sin[e + f*x])^(3/2),x]
```

output

```
Integrate[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))
/(c + d*Sin[e + f*x])^(3/2), x]
```

Rubi [A] (warning: unable to verify)

Time = 1.39 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.234$, Rules used = {3042, 3522, 27, 3042, 3466, 3042, 3267, 157, 27, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a \sin(e + fx) + a)^m (A + B \sin(e + fx) + C \sin(e + fx)^2)}{(c + d \sin(e + fx))^{3/2}} dx$$

↓ 3522

$$\frac{2 \cos(e + fx) (Ad^2 - Bcd + c^2C) (a \sin(e + fx) + a)^m}{df (c^2 - d^2) \sqrt{c + d \sin(e + fx)}} -$$

$$2 \int \frac{(\sin(e + fx)a + a)^m (a((cC - Bd)(d - 2cm) + Ad(c - 2dm)) - a(-2C(m + 1)c^2 + Cd^2 + d(Bc - Ad)(2m + 1)) \sin(e + fx))}{2\sqrt{c + d \sin(e + fx)}} dx$$

↓ 27

$$\frac{2 \int \frac{(\sin(e + fx)a + a)^m (a((cC - Bd)(d - 2cm) + Ad(c - 2dm)) - a(-2C(m + 1)c^2 + Cd^2 + d(Bc - Ad)(2m + 1)) \sin(e + fx))}{2\sqrt{c + d \sin(e + fx)}} dx}{ad(c^2 - d^2)}$$

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a((cC-Bd)(d-2cm)+Ad(c-2dm))-a(-2C(m+1)c^2+Cd^2+d(Bc-Ad)(2m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3042

$$\frac{\int \frac{(\sin(e+fx)a+a)^m (a((cC-Bd)(d-2cm)+Ad(c-2dm))-a(-2C(m+1)c^2+Cd^2+d(Bc-Ad)(2m+1)) \sin(e+fx))}{\sqrt{c+d \sin(e+fx)}} dx}{ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3466

$$\frac{-a(-cd(A+4Bm+B+C) + d^2(4Am+A+B-C) + 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx - (d(2m+1)(Bc-Ad))}{ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3042

$$\frac{-a(-cd(A+4Bm+B+C) + d^2(4Am+A+B-C) + 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^m}{\sqrt{c+d \sin(e+fx)}} dx - (d(2m+1)(Bc-Ad))}{ad(c^2-d^2)} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 3267

$$\frac{a^3 \cos(e+fx)(-cd(A+4Bm+B+C) + d^2(4Am+A+B-C) + 2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{a-a \sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) - a^2 \cos(e+fx)(d(Bc-Ad))}{f \sqrt{a-a \sin(e+fx)} \sqrt{a \sin(e+fx)+a}} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 157

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx)(-cd(A+4Bm+B+C) + d^2(4Am+A+B-C) + 2c^2(2Cm+C)) \int \frac{\sqrt{2}(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)} \sqrt{c+d \sin(e+fx)}} d \sin(e+fx) - a^2 \cos(e+fx)(d(Bc-Ad))}{\sqrt{2} f (a-a \sin(e+fx)) \sqrt{a \sin(e+fx)+a}} + \frac{2 \cos(e+fx) (Ad^2 - Bcd + c^2C) (a \sin(e+fx) + a)^m}{df(c^2-d^2) \sqrt{c+d \sin(e+fx)}}$$

↓ 27

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{c+d\sin(e+fx)}} d\sin(e+fx)}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}} \frac{ad(c^2-d^2)}{2\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m} \frac{df(c^2-d^2)\sqrt{c+d\sin(e+fx)}}{ad(c^2-d^2)}$$

↓ 156

$$\frac{a^3 \sqrt{1-\sin(e+fx)} \cos(e+fx) (-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C)) \sqrt{\frac{c+d\sin(e+fx)}{c-d}} \int \frac{(\sin(e+fx)a+a)^{m-\frac{1}{2}}}{\sqrt{1-\sin(e+fx)}\sqrt{\frac{c}{c-d}+\frac{d\sin(e+fx)}{c-d}}}}{f(a-a\sin(e+fx))\sqrt{a\sin(e+fx)+a}\sqrt{c+d\sin(e+fx)}} \frac{ad(c^2-d^2)}{2\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m} \frac{df(c^2-d^2)\sqrt{c+d\sin(e+fx)}}{ad(c^2-d^2)}$$

↓ 155

$$\frac{\sqrt{2a^2}\sqrt{1-\sin(e+fx)}\cos(e+fx)(a\sin(e+fx)+a)^m(-cd(A+4Bm+B+C)+d^2(4Am+A+B-C)+2c^2(2Cm+C))\sqrt{\frac{c+d\sin(e+fx)}{c-d}}\text{AppellF1}(m)}{f(2m+1)(a-a\sin(e+fx))\sqrt{c+d\sin(e+fx)}} \frac{ad(c^2-d^2)}{2\cos(e+fx)(Ad^2-Bcd+c^2C)(a\sin(e+fx)+a)^m} \frac{df(c^2-d^2)\sqrt{c+d\sin(e+fx)}}{ad(c^2-d^2)}$$

input

```
Int[((a + a*Sin[e + f*x])^m*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/(c + d*Sin[e + f*x])^(3/2),x]
```

output

```
(2*(c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + a*Sin[e + f*x])^m)/(d*(c^2 - d^2)*f*Sqrt[c + d*Sin[e + f*x]]) + (-((Sqrt[2]*a^2*(d^2*(A + B - C + 4*A*m) - c*d*(A + B + C + 4*B*m) + 2*c^2*(C + 2*C*m))*AppellF1[1/2 + m, 1/2, 1/2, 3/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x])/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^m*Sqrt[(c + d*Sin[e + f*x])/(c - d))]/(f*(1 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]) - (Sqrt[2]*a*(C*d^2 - 2*c^2*C*(1 + m) + d*(B*c - A*d)*(1 + 2*m))*AppellF1[3/2 + m, 1/2, 1/2, 5/2 + m, (1 + Sin[e + f*x])/2, -(d*(1 + Sin[e + f*x])/(c - d))]*Cos[e + f*x]*Sqrt[1 - Sin[e + f*x]]*(a + a*Sin[e + f*x])^(1 + m)*Sqrt[(c + d*Sin[e + f*x])/(c - d)]/(f*(3 + 2*m)*(a - a*Sin[e + f*x])*Sqrt[c + d*Sin[e + f*x]]))/(a*d*(c^2 - d^2))
```


Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`
- rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p]*(b*((e + f*x)/(b*e - a*f)))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`
- rule 157 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(c + d*x)^FracPart[n]/(Simplify[b/(b*c - a*d)]^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d)), x]^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && !GtQ[Simplify[b/(b*c - a*d)], 0] && !SimplerQ[c + d*x, a + b*x] && !SimplerQ[e + f*x, a + b*x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3267

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a^2*(Cos[e + f*x]/(f*Sqrt[a + b*Sin[e + f*x]]*Sqrt[a - b*Sin[e + f*x]])) Subst[Int[(a + b*x)^(m - 1/2)*((c + d*x)^n/Sqrt[a - b*x]), x], x, Sin[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !IntegerQ[m]
```

rule 3466

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)])*(c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)/b Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n, x] + Simp[B/b Int[(a + b*Sin[e + f*x])^(m + 1)*(c + d*Sin[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && NeQ[A*b + a*B, 0]
```

rule 3522

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*sin[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*sin[(e_) + (f_)*(x_)] + (C_)*sin[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-c^2*C - B*c*d + A*d^2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 - d^2)), x] + Simp[1/(b*d*(n + 1)*(c^2 - d^2)) Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^(n + 1)*Simp[A*d*(a*d*m + b*c*(n + 1)) + (c*C - B*d)*(a*c*m + b*d*(n + 1)) + b*(d*(B*c - A*d)*(m + n + 2) - C*(c^2*(m + 1) + d^2*(n + 1)))*Sin[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[m, -2^(-1)] && (LtQ[n, -1] || EqQ[m + n + 2, 0])
```

Maple [F]

$$\int \frac{(a + a \sin(fx + e))^m (A + B \sin(fx + e) + C \sin(fx + e)^2)}{(c + d \sin(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)
```

output

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)
```

Fricas [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x, algorithm="fricas")`

output `integral((C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*sqrt(d*sin(f*x + e) + c)*(a*sin(f*x + e) + a)^m/(d^2*cos(f*x + e)^2 - 2*c*d*sin(f*x + e) - c^2 - d^2), x)`

Sympy [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a(\sin(e + fx) + 1))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))**m*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/(c+d*sin(f*x+e))**(3/2),x)`

output `Integral((a*(sin(e + f*x) + 1))**m*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/(c + d*sin(e + f*x))**(3/2), x)`

Maxima [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)}{(d \sin(fx + e) + c)^{3/2}} dx$$

input `integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x, algorithm="maxima")`

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(
d*sin(f*x + e) + c)^(3/2), x)
```

Giac [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(a + a \sin(fx + e))^m}{(d \sin(fx + e) + c)^{3/2}} dx$$

input

```
integrate((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+
e))^(3/2),x, algorithm="giac")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(a*sin(f*x + e) + a)^m/(
d*sin(f*x + e) + c)^(3/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \int \frac{(a + a \sin(e + fx))^m (C \sin(e + fx) + B + C \sin(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx$$

input

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c +
d*sin(e + f*x))^(3/2),x)
```

output

```
int(((a + a*sin(e + f*x))^m*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/(c +
d*sin(e + f*x))^(3/2), x)
```

Reduce [F]

$$\int \frac{(a + a \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx))}{(c + d \sin(e + fx))^{3/2}} dx = \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))}{\sin(fx + e)^2 d^2 + 2 \sin(fx + e) cd + c^2} dx \right) b$$

$$+ \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m \sin(fx + e)}{\sin(fx + e)^2 d^2 + 2 \sin(fx + e) cd + c^2} dx \right) b$$

$$+ \left(\int \frac{\sqrt{\sin(fx + e) d + c} (a + a \sin(fx + e))^m}{\sin(fx + e)^2 d^2 + 2 \sin(fx + e) cd + c^2} dx \right) a$$

input

```
int((a+a*sin(f*x+e))^m*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/(c+d*sin(f*x+e))^(3/2),x)
```

output

```
int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x)**2)/(sin(e + f*x)**2*d**2 + 2*sin(e + f*x)*c*d + c**2),x)*c + int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m*sin(e + f*x))/(sin(e + f*x)**2*d**2 + 2*sin(e + f*x)*c*d + c**2),x)*b + int((sqrt(sin(e + f*x)*d + c)*(sin(e + f*x)*a + a)**m)/(sin(e + f*x)**2*d**2 + 2*sin(e + f*x)*c*d + c**2),x)*a
```

3.30 $\int (a+b \sin(c+dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$

Optimal result	301
Mathematica [A] (verified)	301
Rubi [A] (verified)	302
Maple [A] (verified)	303
Fricas [A] (verification not implemented)	304
Sympy [B] (verification not implemented)	305
Maxima [A] (verification not implemented)	305
Giac [A] (verification not implemented)	306
Mupad [B] (verification not implemented)	306
Reduce [B] (verification not implemented)	307

Optimal result

Integrand size = 31, antiderivative size = 81

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{1}{2}(bB + a(2A + C))x - \frac{(Ab + aB + bC) \cos(c + dx)}{d}$$

$$+ \frac{bC \cos^3(c + dx)}{3d} - \frac{(bB + aC) \cos(c + dx) \sin(c + dx)}{2d}$$

```
output 1/2*(b*B+a*(2*A+C))*x-(A*b+B*a+C*b)*cos(d*x+c)/d+1/3*b*C*cos(d*x+c)^3/d-1/
2*(B*b+C*a)*cos(d*x+c)*sin(d*x+c)/d
```

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.14

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{6bBc + 6acC + 12aAdx + 6bBdx + 6aCdx - 3(4Ab + 4aB + 3bC) \cos(c + dx) + bC \cos(3(c + dx)) - 3}{12d}$$

```
input Integrate[(a + b*Sin[c + d*x])*(A + B*Sin[c + d*x] + C*Sin[c + d*x]^2),x]
```

output

$$(6*b*B*c + 6*a*c*C + 12*a*A*d*x + 6*b*B*d*x + 6*a*C*d*x - 3*(4*A*b + 4*a*B + 3*b*C)*\text{Cos}[c + d*x] + b*C*\text{Cos}[3*(c + d*x)] - 3*b*B*\text{Sin}[2*(c + d*x)] - 3*a*C*\text{Sin}[2*(c + d*x)])/(12*d)$$
Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.46, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {3042, 3502, 3042, 3213}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin(c + dx)^2) dx$$

$$\downarrow 3502$$

$$\frac{\int (a + b \sin(c + dx))(b(3A + 2C) + (3bB - aC) \sin(c + dx)) dx}{\frac{3b}{C \cos(c + dx)(a + b \sin(c + dx))^2}}$$

$$\downarrow 3042$$

$$\frac{\int (a + b \sin(c + dx))(b(3A + 2C) + (3bB - aC) \sin(c + dx)) dx}{\frac{3b}{C \cos(c + dx)(a + b \sin(c + dx))^2}}$$

$$\downarrow 3213$$

$$\frac{-\frac{\cos(c+dx)(a(3bB-aC)+b^2(3A+2C))}{d} + \frac{3}{2}bx(a(2A+C) + bB) - \frac{b(3bB-aC) \sin(c+dx) \cos(c+dx)}{2d}}{\frac{3b}{C \cos(c + dx)(a + b \sin(c + dx))^2}}$$

input

$$\text{Int}[(a + b*\text{Sin}[c + d*x])*(A + B*\text{Sin}[c + d*x] + C*\text{Sin}[c + d*x]^2), x]$$

output

$$-1/3*(C*\text{Cos}[c + d*x]*(a + b*\text{Sin}[c + d*x])^2)/(b*d) + ((3*b*(b*B + a*(2*A + C))*x)/2 - ((b^2*(3*A + 2*C) + a*(3*b*B - a*C))*\text{Cos}[c + d*x])/d - (b*(3*b*B - a*C))*\text{Cos}[c + d*x]*\text{Sin}[c + d*x])/(2*d))/(3*b)$$
Defintions of rubi rules used

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear} \\ \text{Q}[u, x]$$

rule 3213

$$\text{Int}[(a + b*\text{sin}[e + f*x])*(c + d*\text{sin}[e + f*x]), x_Symbol] \rightarrow \text{Simp}[(2*a*c + b*d)*(x/2), x] + (-\text{Simp}[(b*c + a*d)*(\text{Cos}[e + f*x]/f), x] - \text{Simp}[b*d*\text{Cos}[e + f*x]*(\text{Sin}[e + f*x]/(2*f)), x]) \text{ ; Free} \\ \text{Q}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0]$$

rule 3502

$$\text{Int}[(a + b*\text{sin}[e + f*x])^m*(A + B*\text{sin}[e + f*x] + C*\text{sin}[e + f*x]^2), x_Symbol] \rightarrow \text{Simp}[(-C)*\text{Cos}[e + f*x]*(a + b*\text{Sin}[e + f*x])^{m+1}/(b*f*(m+2)), x] + \text{Simp}[1/(b*(m+2)) \ \text{Int}[(a + b*\text{Sin}[e + f*x])^m*\text{Simp}[A*b*(m+2) + b*C*(m+1) + (b*B*(m+2) - a*C)*\text{Sin}[e + f*x], x], x], x] \text{ ; FreeQ}\{a, b, e, f, A, B, C, m\}, x \\ \&\& \ !\text{LtQ}[m, -1]$$
Maple [A] (verified)

Time = 2.22 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.99

method	result
parts	$x A a - \frac{(A b + B a) \cos(dx+c)}{d} + \frac{(B b + C a) \left(-\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right)}{d} - \frac{C b (2 + \sin(dx+c)^2) \cos(dx+c)}{3d}$
parallelrisch	$\frac{(-3 B b - 3 C a) \sin(2 d x + 2 c) + b C \cos(3 d x + 3 c) + ((-12 A - 9 C) b - 12 B a) \cos(dx+c) + (6 B d x - 12 A - 8 C) b + 12 (A d x + \frac{1}{2} C d)}{12 d}$
risch	$x A a + \frac{x B b}{2} + \frac{x C a}{2} - \frac{\cos(dx+c) A b}{d} - \frac{\cos(dx+c) B a}{d} - \frac{3 \cos(dx+c) C b}{4 d} + \frac{b C \cos(3 d x + 3 c)}{12 d} - \frac{\sin(2 d x + 2 c)}{4 d}$
derivativdivides	$-\frac{C b (2 + \sin(dx+c)^2) \cos(dx+c)}{3} + B b \left(-\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + C a \left(-\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) - A b \cos(dx+c)$
default	$-\frac{C b (2 + \sin(dx+c)^2) \cos(dx+c)}{3} + B b \left(-\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) + C a \left(-\frac{\cos(dx+c) \sin(dx+c)}{2} + \frac{dx}{2} + \frac{c}{2} \right) - A b \cos(dx+c)$
norman	$(A a + \frac{1}{2} B b + \frac{1}{2} C a) x + (A a + \frac{1}{2} B b + \frac{1}{2} C a) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^6 + (3 A a + \frac{3}{2} B b + \frac{3}{2} C a) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + (3 A a + \frac{3}{2} B b + \frac{3}{2} C a) x \tan\left(\frac{dx}{2} + \frac{c}{2}\right)$
orering	Expression too large to display

input `int((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `x*A*a-(A*b+B*a)/d*cos(d*x+c)+(B*b+C*a)/d*(-1/2*cos(d*x+c)*sin(d*x+c)+1/2*d*x+1/2*c)-1/3*C*b/d*(2+sin(d*x+c)^2)*cos(d*x+c)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.88

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{2 C b \cos(dx + c)^3 + 3 ((2 A + C) a + B b) dx - 3 (C a + B b) \cos(dx + c) \sin(dx + c) - 6 (B a + (A + C) b) \cos(dx + c)}{6 d}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x, algorithm="fricas")`

output `1/6*(2*C*b*cos(d*x + c)^3 + 3*((2*A + C)*a + B*b)*d*x - 3*(C*a + B*b)*cos(d*x + c)*sin(d*x + c) - 6*(B*a + (A + C)*b)*cos(d*x + c))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(73) = 146$.

Time = 0.13 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.33

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \begin{cases} Aax - \frac{Ab \cos(c+dx)}{d} - \frac{Ba \cos(c+dx)}{d} + \frac{Bbx \sin^2(c+dx)}{2} + \frac{Bbx \cos^2(c+dx)}{2} - \frac{Bb \sin(c+dx) \cos(c+dx)}{2d} + \frac{Cax \sin^2(c+dx)}{2} + \\ x(a + b \sin(c)) (A + B \sin(c) + C \sin^2(c)) \end{cases}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)**2),x)`

output `Piecewise((A*a*x - A*b*cos(c + d*x)/d - B*a*cos(c + d*x)/d + B*b*x*sin(c + d*x)**2/2 + B*b*x*cos(c + d*x)**2/2 - B*b*sin(c + d*x)*cos(c + d*x)/(2*d) + C*a*x*sin(c + d*x)**2/2 + C*a*x*cos(c + d*x)**2/2 - C*a*sin(c + d*x)*cos(c + d*x)/(2*d) - C*b*sin(c + d*x)**2*cos(c + d*x)/d - 2*C*b*cos(c + d*x)**3/(3*d), Ne(d, 0)), (x*(a + b*sin(c))*(A + B*sin(c) + C*sin(c)**2), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.26

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{12(dx + c)Aa + 3(2dx + 2c - \sin(2dx + 2c))Ca + 3(2dx + 2c - \sin(2dx + 2c))Bb + 4(\cos(dx + c)^3 - 3\cos(dx + c))Cb - 12B^2a \cos(dx + c) - 12A^2b \cos(dx + c)}{12d}$$

input `integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x, algorithm="maxima")`

output `1/12*(12*(d*x + c)*A*a + 3*(2*d*x + 2*c - sin(2*d*x + 2*c))*C*a + 3*(2*d*x + 2*c - sin(2*d*x + 2*c))*B*b + 4*(cos(d*x + c)^3 - 3*cos(d*x + c))*C*b - 12*B^2*a*cos(d*x + c) - 12*A^2*b*cos(d*x + c))/d`

Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.94

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{1}{2} (2 A a + C a + B b) x + \frac{C b \cos(3 dx + 3 c)}{12 d}$$

$$- \frac{(4 B a + 4 A b + 3 C b) \cos(dx + c)}{4 d} - \frac{(C a + B b) \sin(2 dx + 2 c)}{4 d}$$

input

```
integrate((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x, algorithm="giac")
```

output

```
1/2*(2*A*a + C*a + B*b)*x + 1/12*C*b*cos(3*d*x + 3*c)/d - 1/4*(4*B*a + 4*A*b + 3*C*b)*cos(d*x + c)/d - 1/4*(C*a + B*b)*sin(2*d*x + 2*c)/d
```

Mupad [B] (verification not implemented)

Time = 38.76 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.15

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx =$$

$$\frac{6 A b \cos(c + dx) + 6 B a \cos(c + dx) + \frac{9 C b \cos(c+dx)}{2} - \frac{C b \cos(3c+3dx)}{2} + \frac{3 B b \sin(2c+2dx)}{2} + \frac{3 C a \sin(2c+2dx)}{2}}{6 d}$$

input

```
int((a + b*sin(c + d*x))*(A + B*sin(c + d*x) + C*sin(c + d*x)^2),x)
```

output

```
-(6*A*b*cos(c + d*x) + 6*B*a*cos(c + d*x) + (9*C*b*cos(c + d*x))/2 - (C*b*cos(3*c + 3*d*x))/2 + (3*B*b*sin(2*c + 2*d*x))/2 + (3*C*a*sin(2*c + 2*d*x))/2 - 6*A*a*d*x - 3*B*b*d*x - 3*C*a*d*x)/(6*d)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.30

$$\int (a + b \sin(c + dx)) (A + B \sin(c + dx) + C \sin^2(c + dx)) dx$$

$$= \frac{-2 \cos(dx + c) \sin(dx + c)^2 bc - 3 \cos(dx + c) \sin(dx + c) ac - 3 \cos(dx + c) \sin(dx + c) b^2 - 12 \cos(dx + c) \sin(dx + c) ab + 6a^2 dx + 12ab + 3ac dx + 3b^2 dx + 4bc}{6d}$$

input

```
int((a+b*sin(d*x+c))*(A+B*sin(d*x+c)+C*sin(d*x+c)^2),x)
```

output

```
( - 2*cos(c + d*x)*sin(c + d*x)**2*b*c - 3*cos(c + d*x)*sin(c + d*x)*a*c -
 3*cos(c + d*x)*sin(c + d*x)*b**2 - 12*cos(c + d*x)*a*b - 4*cos(c + d*x)*b
*c + 6*a**2*d*x + 12*a*b + 3*a*c*d*x + 3*b**2*d*x + 4*b*c)/(6*d)
```

3.31
$$\int \frac{(a+b \sin(e+fx))(A+B \sin(e+fx)+C \sin^2(e+fx))}{\sin^{\frac{3}{2}}(e+fx)} dx$$

Optimal result	308
Mathematica [A] (verified)	309
Rubi [A] (verified)	309
Maple [B] (verified)	312
Fricas [C] (verification not implemented)	313
Sympy [F]	314
Maxima [F]	314
Giac [F]	315
Mupad [B] (verification not implemented)	315
Reduce [F]	316

Optimal result

Integrand size = 41, antiderivative size = 117

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \frac{2(bB - a(A - C))E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{f}$$

$$+ \frac{2(3Ab + 3aB + bC) \operatorname{EllipticF}\left(\frac{1}{2}(e - \frac{\pi}{2} + fx), 2\right)}{3f}$$

$$- \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \cos(e + fx) \sqrt{\sin(e + fx)}}{3f}$$

output

```
-2*(b*B-a*(A-C))*EllipticE(cos(1/2*e+1/4*Pi+1/2*f*x),2^(1/2))/f+2/3*(3*A*b
+3*B*a+C*b)*InverseJacobiAM(1/2*e-1/4*Pi+1/2*f*x,2^(1/2))/f-2*a*A*cos(f*x+
e)/f/sin(f*x+e)^(1/2)-2/3*b*C*cos(f*x+e)*sin(f*x+e)^(1/2)/f
```

Mathematica [A] (verified)

Time = 2.91 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.83

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx = \frac{6(bB + a(-A + C))E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) + 2(3Ab + 3aB + bC) \operatorname{EllipticF}\left(\frac{1}{4}(-2e + \pi - 2fx), 2\right)}{3f}$$

input

```
Integrate[((a + b*Sin[e + f*x])*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sin[e + f*x]^(3/2),x]
```

output

```
-1/3*(6*(b*B + a*(-A + C))*EllipticE[(-2*e + Pi - 2*f*x)/4, 2] + 2*(3*A*b + 3*a*B + b*C)*EllipticF[(-2*e + Pi - 2*f*x)/4, 2] + (2*Cos[e + f*x]*(3*a*A + b*C*Sin[e + f*x]))/Sqrt[Sin[e + f*x]])/f
```

Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.268$, Rules used = {3042, 3510, 27, 3042, 3502, 27, 3042, 3227, 3042, 3119, 3120}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin(e + fx)^2)}{\sin(e + fx)^{3/2}} dx \\ & \quad \downarrow \text{3510} \\ & -2 \int -\frac{bC \sin^2(e + fx) + (bB - a(A - C)) \sin(e + fx) + Ab + aB}{2\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f\sqrt{\sin(e + fx)}} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\int \frac{bC \sin^2(e + fx) + (bB - a(A - C)) \sin(e + fx) + Ab + aB}{\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}}$$

↓ 3042

$$\int \frac{bC \sin(e + fx)^2 + (bB - a(A - C)) \sin(e + fx) + Ab + aB}{\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}}$$

↓ 3502

$$\frac{2}{3} \int \frac{3Ab + Cb + 3aB + 3(bB - a(A - C)) \sin(e + fx)}{2\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \sqrt{\sin(e + fx)} \cos(e + fx)}{3f}$$

↓ 27

$$\frac{1}{3} \int \frac{3Ab + Cb + 3aB + 3(bB - a(A - C)) \sin(e + fx)}{\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \sqrt{\sin(e + fx)} \cos(e + fx)}{3f}$$

↓ 3042

$$\frac{1}{3} \int \frac{3Ab + Cb + 3aB + 3(bB - a(A - C)) \sin(e + fx)}{\sqrt{\sin(e + fx)}} dx - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \sqrt{\sin(e + fx)} \cos(e + fx)}{3f}$$

↓ 3227

$$\frac{1}{3} \left((3aB + 3Ab + bC) \int \frac{1}{\sqrt{\sin(e + fx)}} dx + 3(bB - a(A - C)) \int \sqrt{\sin(e + fx)} dx \right) - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \sqrt{\sin(e + fx)} \cos(e + fx)}{3f}$$

↓ 3042

$$\frac{1}{3} \left((3aB + 3Ab + bC) \int \frac{1}{\sqrt{\sin(e + fx)}} dx + 3(bB - a(A - C)) \int \sqrt{\sin(e + fx)} dx \right) - \frac{2aA \cos(e + fx)}{f \sqrt{\sin(e + fx)}} - \frac{2bC \sqrt{\sin(e + fx)} \cos(e + fx)}{3f}$$

↓ 3119

$$\frac{1}{3} \left((3aB + 3Ab + bC) \int \frac{1}{\sqrt{\sin(e+fx)}} dx + \frac{6E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right) (bB - a(A-C))}{f} \right) - \frac{2aA \cos(e+fx)}{f\sqrt{\sin(e+fx)}} - \frac{2bC\sqrt{\sin(e+fx)} \cos(e+fx)}{3f}$$

↓ 3120

$$\frac{1}{3} \left(\frac{2\text{EllipticF}\left(\frac{1}{2}(e+fx - \frac{\pi}{2}), 2\right) (3aB + 3Ab + bC)}{f} + \frac{6E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right) (bB - a(A-C))}{f} \right) - \frac{2aA \cos(e+fx)}{f\sqrt{\sin(e+fx)}} - \frac{2bC\sqrt{\sin(e+fx)} \cos(e+fx)}{3f}$$

input `Int[((a + b*Sin[e + f*x])*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2))/Sin[e + f*x]^(3/2), x]`

output `((6*(b*B - a*(A - C))*EllipticE[(e - Pi/2 + f*x)/2, 2])/f + (2*(3*A*b + 3*a*B + b*C)*EllipticF[(e - Pi/2 + f*x)/2, 2])/f)/3 - (2*a*A*Cos[e + f*x])/(f*Sqrt[Sin[e + f*x]]) - (2*b*C*Cos[e + f*x]*Sqrt[Sin[e + f*x]])/(3*f)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3119 `Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3120 `Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

rule 3227 $\text{Int}[(b \cdot \sin(e) + f \cdot x)^m \cdot (c + d \cdot \sin(e) + f \cdot x)], x_{\text{Symbol}}] \rightarrow \text{Simp}[c \cdot \text{Int}[(b \cdot \sin(e + f \cdot x))^m, x], x] + \text{Simp}[d/b \cdot \text{Int}[(b \cdot \sin(e + f \cdot x))^{m+1}, x], x] /; \text{FreeQ}\{b, c, d, e, f, m\}, x]$

rule 3502 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (A + B \cdot \sin(e) + f \cdot x) + (C \cdot \sin(e) + f \cdot x)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-C) \cdot \text{Cos}[e + f \cdot x] \cdot (a + b \cdot \sin(e + f \cdot x))^{m+1} / (b \cdot f \cdot (m+2)), x] + \text{Simp}[1/(b \cdot (m+2)) \cdot \text{Int}[(a + b \cdot \sin(e + f \cdot x))^m \cdot \text{Simp}[A \cdot b \cdot (m+2) + b \cdot C \cdot (m+1) + (b \cdot B \cdot (m+2) - a \cdot C) \cdot \sin(e + f \cdot x)], x], x], x] /; \text{FreeQ}\{a, b, e, f, A, B, C, m\}, x] \&\& \text{!LtQ}[m, -1]$

rule 3510 $\text{Int}[(a + b \cdot \sin(e) + f \cdot x)^m \cdot (c + d \cdot \sin(e) + f \cdot x) + (f \cdot x)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-b \cdot c - a \cdot d) \cdot (A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C) \cdot \text{Cos}[e + f \cdot x] \cdot (a + b \cdot \sin(e + f \cdot x))^{m+1} / (b^2 \cdot f \cdot (m+1) \cdot (a^2 - b^2)), x] - \text{Simp}[1/(b^2 \cdot (m+1) \cdot (a^2 - b^2)) \cdot \text{Int}[(a + b \cdot \sin(e + f \cdot x))^{m+1} \cdot \text{Simp}[b \cdot (m+1) \cdot ((b \cdot B - a \cdot C) \cdot (b \cdot c - a \cdot d) - A \cdot b \cdot (a \cdot c - b \cdot d)) + (b \cdot B \cdot (a^2 \cdot d + b^2 \cdot d \cdot (m+1) - a \cdot b \cdot c \cdot (m+2)) + (b \cdot c - a \cdot d) \cdot (A \cdot b^2 \cdot (m+2) + C \cdot (a^2 + b^2 \cdot (m+1))) \cdot \sin(e + f \cdot x) - b \cdot C \cdot d \cdot (m+1) \cdot (a^2 - b^2) \cdot \sin(e + f \cdot x)^2], x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{NeQ}[a^2 - b^2, 0] \&\& \text{LtQ}[m, -1]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 395 vs. $2(108) = 216$.

Time = 0.85 (sec) , antiderivative size = 396, normalized size of antiderivative = 3.38

method	result
parts	$\frac{(Ab+Ba)\sqrt{1+\sin(fx+e)}\sqrt{-2\sin(fx+e)+2}\sqrt{-\sin(fx+e)}}{\cos(fx+e)\sqrt{\sin(fx+e)}} \text{EllipticF}\left(\sqrt{1+\sin(fx+e)}, \frac{\sqrt{2}}{2}\right) - \frac{(Bb+Ca)\sqrt{1+\sin(fx+e)}\sqrt{-2\sin(fx+e)+2}}{f}$
default	$\frac{-A\sqrt{1+\sin(fx+e)}\sqrt{-2\sin(fx+e)+2}\sqrt{-\sin(fx+e)}}{f} \text{EllipticF}\left(\sqrt{1+\sin(fx+e)}, \frac{\sqrt{2}}{2}\right) + Ab\sqrt{1+\sin(fx+e)}\sqrt{-2\sin(fx+e)+2}$

input `int((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2),x,method=_RETURNVERBOSE)`

output `(A*b+B*a)*(1+sin(f*x+e))^(1/2)*(-2*sin(f*x+e)+2)^(1/2)*(-sin(f*x+e))^(1/2)*EllipticF((1+sin(f*x+e))^(1/2),1/2*2^(1/2))/cos(f*x+e)/sin(f*x+e)^(1/2)/f-(B*b+C*a)*(1+sin(f*x+e))^(1/2)*(-2*sin(f*x+e)+2)^(1/2)*(-sin(f*x+e))^(1/2)*(2*EllipticE((1+sin(f*x+e))^(1/2),1/2*2^(1/2))-EllipticF((1+sin(f*x+e))^(1/2),1/2*2^(1/2)))/cos(f*x+e)/sin(f*x+e)^(1/2)/f+A*a*(2*(1+sin(f*x+e))^(1/2)*(-2*sin(f*x+e)+2)^(1/2)*(-sin(f*x+e))^(1/2)*EllipticE((1+sin(f*x+e))^(1/2),1/2*2^(1/2))-(1+sin(f*x+e))^(1/2)*(-2*sin(f*x+e)+2)^(1/2)*(-sin(f*x+e))^(1/2)*EllipticF((1+sin(f*x+e))^(1/2),1/2*2^(1/2))-2*cos(f*x+e)^2)/cos(f*x+e)/sin(f*x+e)^(1/2)/f+C*b*(1/3*(1+sin(f*x+e))^(1/2)*(-2*sin(f*x+e)+2)^(1/2)*(-sin(f*x+e))^(1/2)*EllipticF((1+sin(f*x+e))^(1/2),1/2*2^(1/2))-2/3*cos(f*x+e)^2*sin(f*x+e))/cos(f*x+e)/sin(f*x+e)^(1/2)/f`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.88

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \frac{2 \left(\sqrt{-\frac{1}{2}i} (3Ba + (3A + C)b) \sin(fx + e) \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e)) + \sqrt{\frac{1}{2}} \right)}{\sin^{\frac{3}{2}}(e + fx)}$$

input `integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2),x, algorithm="fricas")`

output

```
2/3*(sqrt(-1/2*I)*(3*B*a + (3*A + C)*b)*sin(f*x + e)*weierstrassPInverse(4
, 0, cos(f*x + e) + I*sin(f*x + e)) + sqrt(1/2*I)*(3*B*a + (3*A + C)*b)*si
n(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)) - 3*sq
rt(-1/2*I)*(I*(A - C)*a - I*B*b)*sin(f*x + e)*weierstrassZeta(4, 0, weiers
trassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) - 3*sqrt(1/2*I)*(-I*(A
- C)*a + I*B*b)*sin(f*x + e)*weierstrassZeta(4, 0, weierstrassPInverse(4,
0, cos(f*x + e) - I*sin(f*x + e))) - (C*b*cos(f*x + e)*sin(f*x + e) + 3*A
*a*cos(f*x + e))*sqrt(sin(f*x + e))/(f*sin(f*x + e))
```

Sympy [F]

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

input

```
integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)**2)/sin(f*x+e)**(3
/2),x)
```

output

```
Integral((a + b*sin(e + f*x))*(A + B*sin(e + f*x) + C*sin(e + f*x)**2)/sin
(e + f*x)**(3/2), x)
```

Maxima [F]

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \int \frac{(C \sin^2(fx + e) + B \sin(fx + e) + A)(b \sin(fx + e) + a)}{\sin^{\frac{3}{2}}(fx + e)} dx$$

input

```
integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2
),x, algorithm="maxima")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(b*sin(f*x + e) + a)/sin
(f*x + e)^(3/2), x)
```

Giac [F]

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \int \frac{(C \sin(fx + e)^2 + B \sin(fx + e) + A)(b \sin(fx + e) + a)}{\sin(fx + e)^{\frac{3}{2}}} dx$$

input

```
integrate((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2
),x, algorithm="giac")
```

output

```
integrate((C*sin(f*x + e)^2 + B*sin(f*x + e) + A)*(b*sin(f*x + e) + a)/sin
(f*x + e)^(3/2), x)
```

Mupad [B] (verification not implemented)

Time = 39.26 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.44

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \frac{2 B b E\left(\frac{e}{2} - \frac{\pi}{4} + \frac{fx}{2} \middle| 2\right)}{f} - \frac{2 B a F\left(\frac{\pi}{4} - \frac{e}{2} - \frac{fx}{2} \middle| 2\right)}{f}$$

$$- \frac{2 A b F\left(\frac{\pi}{4} - \frac{e}{2} - \frac{fx}{2} \middle| 2\right)}{f} + \frac{2 C a E\left(\frac{e}{2} - \frac{\pi}{4} + \frac{fx}{2} \middle| 2\right)}{f}$$

$$- \frac{A a \cos(e + fx) (\sin(e + fx)^2)^{1/4} {}_2F_1\left(\frac{1}{2}, \frac{5}{4}; \frac{3}{2}; \cos(e + fx)^2\right)}{f \sqrt{\sin(e + fx)}}$$

$$- \frac{C b \cos(e + fx) \sin(e + fx)^{5/2} {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{2}; \cos(e + fx)^2\right)}{f (\sin(e + fx)^2)^{5/4}}$$

input

```
int(((a + b*sin(e + f*x))*(A + B*sin(e + f*x) + C*sin(e + f*x)^2))/sin(e +
f*x)^(3/2),x)
```

output

```
(2*B*b*ellipticE(e/2 - pi/4 + (f*x)/2, 2))/f - (2*B*a*ellipticF(pi/4 - e/2 - (f*x)/2, 2))/f - (2*A*b*ellipticF(pi/4 - e/2 - (f*x)/2, 2))/f + (2*C*a*ellipticE(e/2 - pi/4 + (f*x)/2, 2))/f - (A*a*cos(e + f*x)*(sin(e + f*x)^2)^(1/4)*hypergeom([1/2, 5/4], 3/2, cos(e + f*x)^2))/(f*sin(e + f*x)^(1/2)) - (C*b*cos(e + f*x)*sin(e + f*x)^(5/2)*hypergeom([-1/4, 1/2], 3/2, cos(e + f*x)^2))/(f*(sin(e + f*x)^2)^(5/4))
```

Reduce [F]

$$\int \frac{(a + b \sin(e + fx))(A + B \sin(e + fx) + C \sin^2(e + fx))}{\sin^{\frac{3}{2}}(e + fx)} dx$$

$$= \left(\int \frac{\sqrt{\sin(fx + e)}}{\sin(fx + e)^2} dx \right) a^2 + 2 \left(\int \frac{\sqrt{\sin(fx + e)}}{\sin(fx + e)} dx \right) ab$$

$$+ \left(\int \sqrt{\sin(fx + e)} dx \right) ac + \left(\int \sqrt{\sin(fx + e)} dx \right) b^2$$

$$+ \left(\int \sqrt{\sin(fx + e)} \sin(fx + e) dx \right) bc$$

input

```
int((a+b*sin(f*x+e))*(A+B*sin(f*x+e)+C*sin(f*x+e)^2)/sin(f*x+e)^(3/2),x)
```

output

```
int(sqrt(sin(e + f*x))/sin(e + f*x)**2,x)*a**2 + 2*int(sqrt(sin(e + f*x))/sin(e + f*x),x)*a*b + int(sqrt(sin(e + f*x)),x)*a*c + int(sqrt(sin(e + f*x)),x)*b**2 + int(sqrt(sin(e + f*x))*sin(e + f*x),x)*b*c
```

3.32 $\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$

Optimal result	317
Mathematica [N/A]	317
Rubi [N/A]	318
Maple [N/A]	319
Fricas [N/A]	319
Sympy [F(-2)]	320
Maxima [F(-1)]	320
Giac [F(-1)]	321
Mupad [N/A]	321
Reduce [N/A]	322

Optimal result

Integrand size = 45, antiderivative size = 45

$$\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$$

$$= \text{Int}((a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)), x)$$

output

```
Defer(Int)((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x)
```

Mathematica [N/A]

Not integrable

Time = 40.36 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$$

$$= \int (a+b \sin(e+fx))^m (c+d \sin(e+fx))^n (A+B \sin(e+fx) + C \sin^2(e+fx)) dx$$

input

```
Integrate[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2),x]
```

output

```
Integrate[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x]
] + C*Sin[e + f*x]^2), x]
```

Rubi [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3042, 3544}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx) + C \sin(e + fx)^2) (c + d \sin(e + fx))^n dx$$

$$\downarrow \text{3544}$$

$$\int (a + b \sin(e + fx))^m (A + B \sin(e + fx) + C \sin^2(e + fx)) (c + d \sin(e + fx))^n dx$$

input

```
Int[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*
Sin[e + f*x]^2),x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3544 `Int[((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)]) + (C_.)*sin[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Unintegrable[(a + b*Sin[e + f*x])^m*(c + d*Sin[e + f*x])^n*(A + B*Sin[e + f*x] + C*Sin[e + f*x]^2), x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] & NeQ[c^2 - d^2, 0]`

Maple [N/A]

Not integrable

Time = 0.92 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int (a + b \sin (fx + e))^m (c + d \sin (fx + e))^n (A + B \sin (fx + e) + C \sin (fx + e)^2) dx$$

input `int((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

output `int((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2), x)`

Fricas [N/A]

Not integrable

Time = 1.42 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.20

$$\int (a + b \sin (e + fx))^m (c + d \sin (e + fx))^n (A + B \sin (e + fx) + C \sin^2 (e + fx)) dx$$

$$= \int (C \sin (fx + e)^2 + B \sin (fx + e) + A)(b \sin (fx + e) + a)^m (d \sin (fx + e) + c)^n dx$$

input `integrate((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="fricas")`

output `integral(-(C*cos(f*x + e)^2 - B*sin(f*x + e) - A - C)*(b*sin(f*x + e) + a)^m*(d*sin(f*x + e) + c)^n, x)`

Sympy [F(-2)]

Exception generated.

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Exception raised: HeuristicGCDFailed

input `integrate((a+b*sin(f*x+e))**m*(c+d*sin(f*x+e))**n*(A+B*sin(f*x+e)+C*sin(f*x+e)**2),x)`

output Exception raised: HeuristicGCDFailed >> no luck

Maxima [F(-1)]

Timed out.

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input `integrate((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="maxima")`

output Timed out

Giac [F(-1)]

Timed out.

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

= Timed out

input

```
integrate((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),x, algorithm="giac")
```

output

Timed out

Mupad [N/A]

Not integrable

Time = 173.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (C \sin(e + fx)^2 + B \sin(e + fx) + A) dx$$

input

```
int((a + b*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*sin(e + f*x)^2),x)
```

output

```
int((a + b*sin(e + f*x))^m*(c + d*sin(e + f*x))^n*(A + B*sin(e + f*x) + C*sin(e + f*x)^2), x)
```

Reduce [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.27

$$\int (a + b \sin(e + fx))^m (c + d \sin(e + fx))^n (A + B \sin(e + fx) + C \sin^2(e + fx)) dx$$

$$= \left(\int (\sin(fx + e) d + c)^n (\sin(fx + e) b + a)^m \sin(fx + e)^2 dx \right) c$$

$$+ \left(\int (\sin(fx + e) d + c)^n (\sin(fx + e) b + a)^m \sin(fx + e) dx \right) b$$

$$+ \left(\int (\sin(fx + e) d + c)^n (\sin(fx + e) b + a)^m dx \right) a$$

input

```
int((a+b*sin(f*x+e))^m*(c+d*sin(f*x+e))^n*(A+B*sin(f*x+e)+C*sin(f*x+e)^2),
x)
```

output

```
int((sin(e + f*x)*d + c)**n*(sin(e + f*x)*b + a)**m*sin(e + f*x)**2,x)*c +
int((sin(e + f*x)*d + c)**n*(sin(e + f*x)*b + a)**m*sin(e + f*x),x)*b + i
nt((sin(e + f*x)*d + c)**n*(sin(e + f*x)*b + a)**m,x)*a
```

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	323
4.2	Links to plain text integration problems used in this report for each CAS .	341

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
      Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
      If [Head [expn] === RootSum,
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
        If [Head [expn] === Integrate || Head [expn] === Int,
          Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
          9]]]]]]]]]]]

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```



```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```



```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file