

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.1-Sine/195-4.1.13

Nasser M. Abbasi

May 18, 2024 Compiled on May 18, 2024 at 12:09 Noon

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22
2	detailed summary tables of results	23
2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	38
3	Listing of integrals	40
3.1	$\int x^2 \sin(a + bx + cx^2) dx$	42
3.2	$\int x \sin(a + bx + cx^2) dx$	51
3.3	$\int \sin(a + bx + cx^2) dx$	58
3.4	$\int \frac{\sin(a+bx+cx^2)}{x} dx$	64
3.5	$\int \left(-\frac{b \cos(a+bx+cx^2)}{x} + \frac{\sin(a+bx+cx^2)}{x^2} \right) dx$	69
3.6	$\int x^2 \sin(a + bx - cx^2) dx$	74

3.7	$\int x \sin(a + bx - cx^2) dx$	83
3.8	$\int \sin(a + bx - cx^2) dx$	90
3.9	$\int \frac{\sin(a+bx-cx^2)}{x} dx$	96
3.10	$\int \left(-\frac{b \cos(a+bx-cx^2)}{x} + \frac{\sin(a+bx-cx^2)}{x^2} \right) dx$	101
3.11	$\int x^2 \sin(\frac{1}{4} + x + x^2) dx$	106
3.12	$\int x \sin(\frac{1}{4} + x + x^2) dx$	113
3.13	$\int \sin(\frac{1}{4} + x + x^2) dx$	119
3.14	$\int \frac{\sin(\frac{1}{4}+x+x^2)}{x} dx$	124
3.15	$\int \frac{\sin(\frac{1}{4}+x+x^2)}{x^2} dx$	129
3.16	$\int x^2 \sin^2(a + bx + cx^2) dx$	135
3.17	$\int x \sin^2(a + bx + cx^2) dx$	142
3.18	$\int \sin^2(a + bx + cx^2) dx$	148
3.19	$\int \frac{\sin^2(a+bx+cx^2)}{x} dx$	154
3.20	$\int x^2 \sin^2(a + bx - cx^2) dx$	159
3.21	$\int x \sin^2(a + bx - cx^2) dx$	166
3.22	$\int \sin^2(a + bx - cx^2) dx$	172
3.23	$\int \frac{\sin^2(a+bx-cx^2)}{x} dx$	178
3.24	$\int x^2 \sin^2(\frac{1}{4} + x + x^2) dx$	183
3.25	$\int x \sin^2(\frac{1}{4} + x + x^2) dx$	189
3.26	$\int \sin^2(\frac{1}{4} + x + x^2) dx$	195
3.27	$\int \frac{\sin^2(\frac{1}{4}+x+x^2)}{x} dx$	200
3.28	$\int \frac{\sin^2(\frac{1}{4}+x+x^2)}{x^2} dx$	205
3.29	$\int (d + ex)^2 \sin(a + bx + cx^2) dx$	210
3.30	$\int (d + ex) \sin(a + bx + cx^2) dx$	220
3.31	$\int \sin(a + bx + cx^2) dx$	227
3.32	$\int \frac{\sin(a+bx+cx^2)}{d+ex} dx$	233
3.33	$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx$	238
3.34	$\int (d + ex) \sin^2(a + bx + cx^2) dx$	246
3.35	$\int \sin^2(a + bx + cx^2) dx$	252
3.36	$\int \frac{\sin^2(a+bx+cx^2)}{d+ex} dx$	258
4	Appendix	263
4.1	Listing of Grading functions	263
4.2	Links to plain text integration problems used in this report for each CAS	281

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [36]. This is test number [195].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (36)	0.00 (0)
Mathematica	100.00 (36)	0.00 (0)
Fricas	100.00 (36)	0.00 (0)
Maple	94.44 (34)	5.56 (2)
Giac	94.44 (34)	5.56 (2)
Maxima	94.44 (34)	5.56 (2)
Sympy	55.56 (20)	44.44 (16)
Mupad	44.44 (16)	55.56 (20)
Reduce	27.78 (10)	72.22 (26)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	72.222	0.000	0.000	27.778
Mathematica	72.222	0.000	0.000	27.778
Fricas	72.222	0.000	0.000	27.778
Maple	66.667	0.000	0.000	33.333
Sympy	22.222	5.556	0.000	72.222
Giac	0.000	0.000	66.667	33.333
Mupad	0.000	16.667	0.000	83.333
Maxima	0.000	0.000	66.667	33.333
Reduce	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	2	100.00	0.00	0.00
Giac	2	100.00	0.00	0.00
Maxima	2	100.00	0.00	0.00
Sympy	16	100.00	0.00	0.00
Mupad	20	0.00	100.00	0.00
Reduce	26	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.08
Giac	0.15
Reduce	0.18
Rubi	0.30
Maxima	0.33
Sympy	0.46
Maple	0.95
Mathematica	2.96
Mupad	26.54

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	23.50	1.54	18.50	1.11
Mupad	37.00	1.01	19.50	1.02
Sympy	50.25	1.16	20.50	0.91
Mathematica	84.42	0.95	86.50	0.96
Fricas	90.08	0.98	94.50	0.99
Maple	94.26	0.89	72.00	0.85
Rubi	101.56	1.00	97.50	1.00
Giac	113.15	1.16	122.00	1.12
Maxima	486.00	3.22	124.00	1.96

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{4, 9, 14, 15, 19, 23, 27, 28, 32, 36}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A `time limit` of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests




Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	38

2.1 List of integrals sorted by grade for each CAS

Rubi	24
Mma	24
Maple	25
Fricas	25
Maxima	25
Giac	26
Mupad	26
Sympy	26
Reduce	27

Rubi

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31, 33, 34, 35 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31, 33, 34, 35 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31, 33, 34, 35 }

B grade { }

C grade { }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31, 33, 34, 35 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { }

B grade { }

C grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31, 33, 34, 35 }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac**A grade** { }**B grade** { }**C grade** { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31, 33, 34, 35 }**F normal fail** { 5, 10 }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Mupad****A grade** { }**B grade** { 3, 8, 11, 12, 13, 31 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { 1, 2, 5, 6, 7, 10, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 33, 34, 35 }**F(-2) exception fail** { }**Sympy****A grade** { 3, 8, 13, 18, 22, 26, 31, 35 }**B grade** { 12, 25 }**C grade** { }**F normal fail** { 1, 2, 5, 6, 7, 10, 11, 16, 17, 20, 21, 24, 29, 30, 33, 34 }**F(-1) timeout fail** { }**F(-2) exception fail** { }

Reduce

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 31,
33, 34, 35 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	249	257	157	204	1569	172	0	227	190	0
N.S.	1	1.03	0.63	0.82	6.30	0.69	0.00	0.91	0.76	0.00
time (sec)	N/A	0.721	0.794	1.649	0.776	0.101	0.000	0.151	0.155	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	124	111	100	579	118	0	183	15	0
N.S.	1	1.01	0.90	0.81	4.71	0.96	0.00	1.49	0.12	0.00
time (sec)	N/A	0.323	0.418	0.947	0.390	0.083	0.000	0.140	0.175	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	84	82	112	102	88	135	13	99
N.S.	1	1.00	0.87	0.85	1.15	1.05	0.91	1.39	0.13	1.02
time (sec)	N/A	0.264	0.206	0.572	0.044	0.074	0.207	0.135	0.155	0.063

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.159	4.221	0.171	0.166	0.075	0.360	0.128	0.157	38.781

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	110	0	0	117	0	0	45	0
N.S.	1	1.00	1.00	0.00	0.00	1.06	0.00	0.00	0.41	0.00
time (sec)	N/A	0.291	0.888	0.000	0.000	0.084	0.000	0.000	0.156	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	251	261	165	224	1570	179	0	229	194	0
N.S.	1	1.04	0.66	0.89	6.25	0.71	0.00	0.91	0.77	0.00
time (sec)	N/A	0.745	0.928	1.599	0.796	0.079	0.000	0.124	0.181	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	126	117	109	580	126	0	185	16	0
N.S.	1	1.02	0.94	0.88	4.68	1.02	0.00	1.49	0.13	0.00
time (sec)	N/A	0.336	0.345	0.921	0.404	0.121	0.000	0.148	0.185	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	89	87	112	107	94	137	14	105
N.S.	1	1.00	0.91	0.89	1.14	1.09	0.96	1.40	0.14	1.07
time (sec)	N/A	0.268	0.203	0.497	0.128	0.083	0.210	0.142	0.166	0.053

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	22	21	14	18	18	18
N.S.	1	1.00	1.12	1.00	1.38	1.31	0.88	1.12	1.12	1.12
time (sec)	N/A	0.164	4.693	0.171	0.169	0.082	0.365	0.128	0.150	38.621

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	115	0	0	124	0	0	47	0
N.S.	1	1.00	1.05	0.00	0.00	1.13	0.00	0.00	0.43	0.00
time (sec)	N/A	0.294	0.843	0.000	0.000	0.090	0.000	0.000	0.154	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	87	66	59	157	59	0	75	94	64
N.S.	1	1.06	0.80	0.72	1.91	0.72	0.00	0.91	1.15	0.78
time (sec)	N/A	0.362	0.142	1.355	0.227	0.074	0.000	0.140	0.155	0.127

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	39	30	122	32	160	65	11	32
N.S.	1	1.00	0.95	0.73	2.98	0.78	3.90	1.59	0.27	0.78
time (sec)	N/A	0.226	0.059	1.003	0.190	0.072	0.641	0.125	0.153	38.487

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	20	70	22	29	39	9	19
N.S.	1	1.00	1.00	0.83	2.92	0.92	1.21	1.62	0.38	0.79
time (sec)	N/A	0.177	0.034	0.703	0.120	0.073	0.187	0.111	0.154	0.047

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	13	13	12	13	13	13
N.S.	1	1.00	1.15	0.85	1.00	1.00	0.92	1.00	1.00	1.00
time (sec)	N/A	0.161	19.516	0.156	0.153	0.069	0.324	0.117	0.164	38.940

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	13	13	14	13	74	13
N.S.	1	1.00	1.15	0.85	1.00	1.00	1.08	1.00	5.69	1.00
time (sec)	N/A	0.288	14.377	0.146	0.155	0.070	0.301	0.128	0.156	38.840

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	248	248	170	191	1617	178	0	214	308	0
N.S.	1	1.00	0.69	0.77	6.52	0.72	0.00	0.86	1.24	0.00
time (sec)	N/A	0.462	0.659	1.405	0.806	0.086	0.000	0.164	0.167	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	118	95	611	124	0	170	56	0
N.S.	1	1.00	0.94	0.75	4.85	0.98	0.00	1.35	0.44	0.00
time (sec)	N/A	0.264	0.362	1.259	0.374	0.092	0.000	0.146	0.195	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	97	72	124	93	83	122	15	0
N.S.	1	1.00	0.97	0.72	1.24	0.93	0.83	1.22	0.15	0.00
time (sec)	N/A	0.240	0.125	1.195	0.131	0.081	0.428	0.147	0.226	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	19	17	28	22	15	19	19	19
N.S.	1	1.00	1.12	1.00	1.65	1.29	0.88	1.12	1.12	1.12
time (sec)	N/A	0.192	7.341	0.357	0.182	0.078	0.791	0.190	0.246	38.244

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	248	248	175	199	1617	187	0	216	314	0
N.S.	1	1.00	0.71	0.80	6.52	0.75	0.00	0.87	1.27	0.00
time (sec)	N/A	0.435	0.827	1.406	0.777	0.092	0.000	0.173	0.222	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	122	99	610	133	0	172	59	0
N.S.	1	1.00	0.97	0.79	4.84	1.06	0.00	1.37	0.47	0.00
time (sec)	N/A	0.257	0.281	1.264	0.391	0.087	0.000	0.195	0.169	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	102	76	124	96	88	124	16	0
N.S.	1	1.00	1.02	0.76	1.24	0.96	0.88	1.24	0.16	0.00
time (sec)	N/A	0.228	0.117	1.200	0.177	0.076	0.431	0.157	0.172	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	28	25	15	20	20	20
N.S.	1	1.00	1.11	1.00	1.56	1.39	0.83	1.11	1.11	1.11
time (sec)	N/A	0.202	7.845	0.368	0.178	0.076	0.793	0.182	0.157	38.541

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	77	64	171	57	0	64	689	0
N.S.	1	1.00	0.91	0.75	2.01	0.67	0.00	0.75	8.11	0.00
time (sec)	N/A	0.245	0.173	1.540	0.262	0.084	0.000	0.122	0.184	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	42	35	137	37	121	54	38	0
N.S.	1	1.00	0.91	0.76	2.98	0.80	2.63	1.17	0.83	0.00
time (sec)	N/A	0.199	0.078	1.426	0.194	0.093	0.801	0.138	0.167	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	20	34	19	22	26	11	0
N.S.	1	1.00	1.00	0.74	1.26	0.70	0.81	0.96	0.41	0.00
time (sec)	N/A	0.173	0.019	1.329	0.120	0.073	0.352	0.138	0.184	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	13	24	18	14	15	15	15
N.S.	1	1.00	1.13	0.87	1.60	1.20	0.93	1.00	1.00	1.00
time (sec)	N/A	0.189	10.686	0.348	0.169	0.076	0.626	0.153	0.182	38.004

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	13	26	18	15	15	15	15
N.S.	1	1.00	1.13	0.87	1.73	1.20	1.00	1.00	1.00	1.00
time (sec)	N/A	0.227	10.987	0.355	0.160	0.113	0.542	0.176	0.163	38.066

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	285	282	186	399	2269	231	0	289	505	0
N.S.	1	0.99	0.65	1.40	7.96	0.81	0.00	1.01	1.77	0.00
time (sec)	N/A	0.781	1.395	1.879	1.143	0.087	0.000	0.130	0.188	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	133	128	184	694	135	0	199	33	0
N.S.	1	0.95	0.91	1.31	4.96	0.96	0.00	1.42	0.24	0.00
time (sec)	N/A	0.342	0.936	1.013	0.393	0.077	0.000	0.163	0.165	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	84	82	112	102	88	135	13	99
N.S.	1	1.00	0.87	0.85	1.15	1.05	0.91	1.39	0.13	1.02
time (sec)	N/A	0.268	0.100	0.461	0.041	0.080	0.206	0.140	0.160	0.002

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	19	21	21	17	21	21	21
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.89	1.11	1.11	1.11
time (sec)	N/A	0.170	6.372	0.355	0.188	0.069	0.337	0.159	0.177	38.443

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	291	291	215	378	2361	257	0	296	516	0
N.S.	1	1.00	0.74	1.30	8.11	0.88	0.00	1.02	1.77	0.00
time (sec)	N/A	0.533	1.173	2.010	1.088	0.095	0.000	0.163	0.186	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	150	150	140	170	738	149	0	195	79	0
N.S.	1	1.00	0.93	1.13	4.92	0.99	0.00	1.30	0.53	0.00
time (sec)	N/A	0.290	0.660	1.534	0.464	0.091	0.000	0.154	0.175	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	97	72	124	93	83	122	15	0
N.S.	1	1.00	0.97	0.72	1.24	0.93	0.83	1.22	0.15	0.00
time (sec)	N/A	0.223	0.075	1.130	0.123	0.099	0.425	0.146	0.164	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	21	118	26	19	23	23	23
N.S.	1	1.00	1.10	1.00	5.62	1.24	0.90	1.10	1.10	1.10
time (sec)	N/A	0.215	8.642	0.514	0.201	0.081	0.778	0.197	0.239	39.445

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [6] had the largest ratio of [.6250000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	8	1.03	15	0.533
2	A	4	4	1.01	13	0.308
3	A	3	3	1.00	11	0.273
4	N/A	1	0	1.00	15	0.000
5	A	1	1	1.00	33	0.030
6	A	10	10	1.04	16	0.625
7	A	5	5	1.02	14	0.357
8	A	4	4	1.00	12	0.333
9	N/A	1	0	1.00	16	0.000
10	A	1	1	1.00	35	0.029
11	A	6	6	1.06	13	0.462
12	A	3	3	1.00	11	0.273
13	A	2	2	1.00	9	0.222
14	N/A	1	0	1.00	13	0.000
15	N/A	4	0	1.00	13	0.000
16	A	2	2	1.00	17	0.118
17	A	2	2	1.00	15	0.133
18	A	2	2	1.00	13	0.154
19	N/A	2	0	1.00	17	0.000
20	A	2	2	1.00	18	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
21	A	2	2	1.00	16	0.125
22	A	2	2	1.00	14	0.143
23	N/A	2	0	1.00	18	0.000
24	A	2	2	1.00	15	0.133
25	A	2	2	1.00	13	0.154
26	A	2	2	1.00	11	0.182
27	N/A	2	0	1.00	15	0.000
28	N/A	2	0	1.00	15	0.000
29	A	8	8	0.99	19	0.421
30	A	4	4	0.95	17	0.235
31	A	3	3	1.00	11	0.273
32	N/A	1	0	1.00	19	0.000
33	A	2	2	1.00	21	0.095
34	A	2	2	1.00	19	0.105
35	A	2	2	1.00	13	0.154
36	N/A	2	0	1.00	21	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^2 \sin(a + bx + cx^2) dx$	42
3.2	$\int x \sin(a + bx + cx^2) dx$	51
3.3	$\int \sin(a + bx + cx^2) dx$	58
3.4	$\int \frac{\sin(a+bx+cx^2)}{x} dx$	64
3.5	$\int \left(-\frac{b \cos(a+bx+cx^2)}{x} + \frac{\sin(a+bx+cx^2)}{x^2} \right) dx$	69
3.6	$\int x^2 \sin(a + bx - cx^2) dx$	74
3.7	$\int x \sin(a + bx - cx^2) dx$	83
3.8	$\int \sin(a + bx - cx^2) dx$	90
3.9	$\int \frac{\sin(a+bx-cx^2)}{x} dx$	96
3.10	$\int \left(-\frac{b \cos(a+bx-cx^2)}{x} + \frac{\sin(a+bx-cx^2)}{x^2} \right) dx$	101
3.11	$\int x^2 \sin(\frac{1}{4} + x + x^2) dx$	106
3.12	$\int x \sin(\frac{1}{4} + x + x^2) dx$	113
3.13	$\int \sin(\frac{1}{4} + x + x^2) dx$	119
3.14	$\int \frac{\sin(\frac{1}{4}+x+x^2)}{x} dx$	124
3.15	$\int \frac{\sin(\frac{1}{4}+x+x^2)}{x^2} dx$	129
3.16	$\int x^2 \sin^2(a + bx + cx^2) dx$	135
3.17	$\int x \sin^2(a + bx + cx^2) dx$	142
3.18	$\int \sin^2(a + bx + cx^2) dx$	148
3.19	$\int \frac{\sin^2(a+bx+cx^2)}{x} dx$	154
3.20	$\int x^2 \sin^2(a + bx - cx^2) dx$	159
3.21	$\int x \sin^2(a + bx - cx^2) dx$	166
3.22	$\int \sin^2(a + bx - cx^2) dx$	172
3.23	$\int \frac{\sin^2(a+bx-cx^2)}{x} dx$	178
3.24	$\int x^2 \sin^2(\frac{1}{4} + x + x^2) dx$	183
3.25	$\int x \sin^2(\frac{1}{4} + x + x^2) dx$	189

3.26	$\int \sin^2\left(\frac{1}{4} + x + x^2\right) dx$	195
3.27	$\int \frac{\sin^2(\frac{1}{4}+x+x^2)}{x} dx$	200
3.28	$\int \frac{\sin^2(\frac{1}{4}+x+x^2)}{x^2} dx$	205
3.29	$\int (d + ex)^2 \sin(a + bx + cx^2) dx$	210
3.30	$\int (d + ex) \sin(a + bx + cx^2) dx$	220
3.31	$\int \sin(a + bx + cx^2) dx$	227
3.32	$\int \frac{\sin(a+bx+cx^2)}{d+ex} dx$	233
3.33	$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx$	238
3.34	$\int (d + ex) \sin^2(a + bx + cx^2) dx$	246
3.35	$\int \sin^2(a + bx + cx^2) dx$	252
3.36	$\int \frac{\sin^2(a+bx+cx^2)}{d+ex} dx$	258

3.1 $\int x^2 \sin(a + bx + cx^2) dx$

Optimal result	42
Mathematica [A] (verified)	43
Rubi [A] (verified)	43
Maple [A] (verified)	46
Fricas [A] (verification not implemented)	47
Sympy [F]	47
Maxima [C] (verification not implemented)	48
Giac [C] (verification not implemented)	49
Mupad [F(-1)]	49
Reduce [F]	50

Optimal result

Integrand size = 15, antiderivative size = 249

$$\begin{aligned} \int x^2 \sin(a + bx + cx^2) dx &= \frac{b \cos(a + bx + cx^2)}{4c^2} - \frac{x \cos(a + bx + cx^2)}{2c} \\ &+ \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &+ \frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ &+ \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\ &- \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \end{aligned}$$

output

```
1/4*b*cos(c*x^2+b*x+a)/c^2-1/2*x*cos(c*x^2+b*x+a)/c+1/4*2^(1/2)*Pi^(1/2)*c
os(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)+1
/8*b^2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^
(1/2)/Pi^(1/2))/c^(5/2)+1/8*b^2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^
(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(5/2)-1/4*2^(1/2)*Pi^(1/2)*Fres
nelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.63

$$\int x^2 \sin(a + bx + cx^2) dx \\ = \frac{2\sqrt{c}(b - 2cx) \cos(a + x(b + cx)) + \sqrt{2\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a - \frac{b^2}{4c}\right) + b^2 \sin\left(a - \frac{b^2}{4c}\right)\right) + \sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(-2c \sin\left(a - \frac{b^2}{4c}\right) + b^2 \cos\left(a - \frac{b^2}{4c}\right)\right)}{8c^{5/2}}$$

input `Integrate[x^2*Sin[a + b*x + c*x^2], x]`

output
$$(2*\text{Sqrt}[c]*(b - 2*c*x)*\text{Cos}[a + x*(b + c*x)] + \text{Sqrt}[2*\pi]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])]*(2*c*\text{Cos}[a - b^2/(4*c)] + b^2*\text{Sin}[a - b^2/(4*c)]) + \text{Sqrt}[2*\pi]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])]*(b^2*\text{Cos}[a - b^2/(4*c)] - 2*c*\text{Sin}[a - b^2/(4*c)]))/(8*c^{(5/2)})$$

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3944, 3929, 3832, 3833, 3942, 3928, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sin(a + bx + cx^2) dx \\ \downarrow 3944 \\ -\frac{b \int x \sin(cx^2 + bx + a) dx}{2c} + \frac{\int \cos(cx^2 + bx + a) dx}{2c} - \frac{x \cos(a + bx + cx^2)}{2c} \\ \downarrow 3929 \\ \frac{\cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx - \sin\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} - \\ \frac{b \int x \sin(cx^2 + bx + a) dx}{2c} - \frac{x \cos(a + bx + cx^2)}{2c}$$

$$\begin{aligned}
& \downarrow \text{3832} \\
& \frac{\cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \\
& \frac{b \int x \sin(cx^2 + bx + a) dx}{2c} - \frac{x \cos(a + bx + cx^2)}{2c} \\
& \downarrow \text{3833} \\
& - \frac{b \int x \sin(cx^2 + bx + a) dx}{2c} + \frac{\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \\
& \frac{x \cos(a + bx + cx^2)}{2c} \\
& \downarrow \text{3942} \\
& - \frac{b \left(-\frac{b \int \sin(cx^2 + bx + a) dx}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \cos(a + bx + cx^2)}{2c} \\
& \downarrow \text{3928} \\
& - \frac{b \left(-\frac{b \left(\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx + \cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx \right)}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \cos(a + bx + cx^2)}{2c} \\
& \downarrow \text{3832} \\
& - \frac{b \left(-\frac{b \left(\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \cos(a + bx + cx^2)}{2c} \\
& \downarrow \text{3833}
\end{aligned}$$

$$\begin{aligned}
& - \frac{b \left(- \frac{b \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} - \frac{\cos(a+bx+cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \cos(a+bx+cx^2)}{2c}
\end{aligned}$$

input `Int[x^2*Sin[a + b*x + c*x^2],x]`

output
$$\begin{aligned}
& -\frac{1}{2} \left(x \cos(a + b*x + c*x^2) \right) / c + \left((\text{Sqrt}[\text{Pi}/2] * \cos(a - b^2/(4*c)) * \text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}]]) / \text{Sqrt}[c] - (\text{Sqrt}[\text{Pi}/2] * \text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] * \sin(a - b^2/(4*c))) / \text{Sqrt}[c]) / (2*c) - (b * (-1/2 * \cos(a + b*x + c*x^2) / c) - (b * ((\text{Sqrt}[\text{Pi}/2] * \cos(a - b^2/(4*c)) * \text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] / \text{Sqrt}[c] + (\text{Sqrt}[\text{Pi}/2] * \text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] * \sin(a - b^2/(4*c))) / \text{Sqrt}[c])) / (2*c))) / (2*c)
\right)
\end{aligned}$$

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3928 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] - Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3929 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3942

```

Int[((d_.) + (e_)*(x_))*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol]
:> Simp[(-e)*(Cos[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c)
Int[Sin[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d
- b*e, 0]

```

rule 3944

```

Int[((d_.) + (e_)*(x_))^(m_)*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2], x_Sym
bol] :> Simp[(-e)*(d + e*x)^(m - 1)*(Cos[a + b*x + c*x^2]/(2*c)), x] + (-Si
mp[(b*e - 2*c*d)/(2*c) Int[(d + e*x)^(m - 1)*Sin[a + b*x + c*x^2], x], x]
+ Simp[e^2*((m - 1)/(2*c)) Int[(d + e*x)^(m - 2)*Cos[a + b*x + c*x^2], x
], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]

```

Maple [A] (verified)

Time = 1.65 (sec), antiderivative size = 204, normalized size of antiderivative = 0.82

method	result
default	$-\frac{x \cos(cx^2+bx+a)}{2c} - \frac{b \left(-\frac{\cos(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2-ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) - \sin\left(\frac{b^2-ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} \right)}{2c}$
risch	$\frac{ib^2\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{16c^2\sqrt{-ic}} - \frac{\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} + \frac{ib^2\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{16c^2\sqrt{ic}}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2\cos\left(\frac{b^2-ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\sqrt{2}\sqrt{\pi}x^2\sin\left(\frac{b^2-ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\sqrt{2}\pi^{\frac{3}{2}}\left(\cos\left(\frac{4ac-b^2}{4c}\right) - \frac{\sqrt{2}\sqrt{\pi}x^2\cos\left(\frac{b^2-ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}}\right)}{2\sqrt{c}}$

input `int(x^2*sin(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -\frac{1}{2}x \cos(c*x^2 + b*x + a)/c - \frac{1}{2}b/c * (-\frac{1}{2}\cos(c*x^2 + b*x + a)/c - \frac{1}{4}b/c^{(3/2)}*2 \\ & ^{(1/2)}*\text{Pi}^{(1/2)} * (\cos((1/4*b^2 - a*c)/c) * \text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)} * (c \\ & *x + 1/2*b)) - \sin((1/4*b^2 - a*c)/c) * \text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)} * (c*x + 1/2 \\ & *b))) + \frac{1}{4}c^{(3/2)}*2^{(1/2)}*\text{Pi}^{(1/2)} * (\cos((1/4*b^2 - a*c)/c) * \text{FresnelC}(2^{(1/2)} \\ & /\text{Pi}^{(1/2)}/c^{(1/2)} * (c*x + 1/2*b)) + \sin((1/4*b^2 - a*c)/c) * \text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)} \\ & /c^{(1/2)} * (c*x + 1/2*b))) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec), antiderivative size = 172, normalized size of antiderivative = 0.69

$$\int x^2 \sin(a + bx + cx^2) dx = \frac{\sqrt{2} \left(\pi b^2 \sin\left(-\frac{b^2 - 4ac}{4c}\right) + 2\pi c \cos\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2} \left(\pi b^2 \cos\left(-\frac{b^2 - 4ac}{4c}\right) - 2\pi c \sin\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right)}{8c^3}$$

input `integrate(x^2*sin(c*x^2+b*x+a), x, algorithm="fricas")`

output

$$\begin{aligned} & \frac{1}{8} * (\sqrt{2} * (\pi * b^2 * \sin(-1/4 * (b^2 - 4*a*c)/c) + 2 * \pi * c * \cos(-1/4 * (b^2 - 4*a*c)/c)) * \sqrt{c/\pi} * \text{fresnel_cos}(1/2 * \sqrt{2} * (2*c*x + b) * \sqrt{c/\pi}/c) + \sqrt{2} * (\pi * b^2 * \cos(-1/4 * (b^2 - 4*a*c)/c) - 2 * \pi * c * \sin(-1/4 * (b^2 - 4*a*c)/c)) * \sqrt{c/\pi} * \text{fresnel_sin}(1/2 * \sqrt{2} * (2*c*x + b) * \sqrt{c/\pi}/c) - 2 * (2*c^2 * x - b*c) * \cos(c*x^2 + b*x + a))/c^3 \end{aligned}$$

Sympy [F]

$$\int x^2 \sin(a + bx + cx^2) dx = \int x^2 \sin(a + bx + cx^2) dx$$

input `integrate(x**2*sin(c*x**2+b*x+a), x)`

output `Integral(x**2*sin(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec), antiderivative size = 1569, normalized size of antiderivative = 6.30

$$\int x^2 \sin(a + bx + cx^2) dx = \text{Too large to display}$$

```
input integrate(x^2*sin(c*x^2+b*x+a),x, algorithm="maxima")
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.91

$$\int x^2 \sin(a + bx + cx^2) dx =$$

$$-\frac{i\sqrt{2}\sqrt{\pi}(b^2+2ic)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4ic}{4c}\right)}}{\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2i(c(2ix+\frac{ib}{c})-2ib)e^{(icx^2+ibx+ia)}$$

$$-\frac{i\sqrt{2}\sqrt{\pi}(b^2-2ic)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4ic}{4c}\right)}}{\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2i(c(2ix+\frac{ib}{c})-2ib)e^{(-icx^2-ibx-ia)}$$

$$\frac{16c^2}{16c^2}$$

input `integrate(x^2*sin(c*x^2+b*x+a),x, algorithm="giac")`

output

```
-1/16*(-I*sqrt(2)*sqrt(pi)*(b^2 + 2*I*c)*erf(-1/4*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*(c*(2*I*x + I*b/c) - 2*I*b)*e^(I*c*x^2 + I*b*x + I*a)/c^2 - 1/16*(I*sqrt(2)*sqrt(pi)*(b^2 - 2*I*c)*erf(-1/4*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*(c*(2*I*x + I*b/c) - 2*I*b)*e^(-I*c*x^2 - I*b*x - I*a))/c^2
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \sin(a + bx + cx^2) dx = \int x^2 \sin(cx^2 + bx + a) dx$$

input `int(x^2*sin(a + b*x + c*x^2),x)`

output `int(x^2*sin(a + b*x + c*x^2), x)`

Reduce [F]

$$\int x^2 \sin(a + bx + cx^2) dx$$

$$= \frac{6 \cos(cx^2 + bx + a) bc - 12 \cos(cx^2 + bx + a) c^2 x - 24 \left(\int \frac{x^2}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) b^2 c^2 + 6 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) b^2 c^2 + 6 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) b^2 c^2}{b^2 c^2}$$

input `int(x^2*sin(c*x^2+b*x+a),x)`

output `(6*cos(a + b*x + c*x**2)*b*c - 12*cos(a + b*x + c*x**2)*c**2*x - 24*int(x*x**2/(tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2*c**2 + 6*int(1/(tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**4 + 24*int(1/(tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**2 - 3*sin(a + b*x + c*x**2)*b**3 + 6*sin(a + b*x + c*x**2)*b**2*c*x - 3*b**4*x + 4*b**2*c**2*x**3 - 6*b*c - 12*c**2*x)/(24*c**3)`

3.2 $\int x \sin(a + bx + cx^2) dx$

Optimal result	51
Mathematica [A] (verified)	51
Rubi [A] (verified)	52
Maple [A] (verified)	53
Fricas [A] (verification not implemented)	54
Sympy [F]	55
Maxima [C] (verification not implemented)	55
Giac [C] (verification not implemented)	56
Mupad [F(-1)]	57
Reduce [F]	57

Optimal result

Integrand size = 13, antiderivative size = 123

$$\int x \sin(a + bx + cx^2) dx = -\frac{\cos(a + bx + cx^2)}{2c} - \frac{b\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ - \frac{b\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}}$$

output

```
-1/2*cos(c*x^2+b*x+a)/c-1/4*b*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1
/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)-1/4*b*2^(1/2)*Pi^(1/2)*Fres
nelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.90

$$\int x \sin(a + bx + cx^2) dx = \\ \frac{-2\sqrt{c} \cos(a + x(b + cx)) + b\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + b\sqrt{2\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{3/2}}$$

input

```
Integrate[x*Sin[a + b*x + c*x^2], x]
```

output

$$-1/4*(2* \text{Sqrt}[c]*\text{Cos}[a + x*(b + c*x)] + b*\text{Sqrt}[2*\pi]*\text{Cos}[a - b^2/(4*c)]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])] + b*\text{Sqrt}[2*\pi]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])]*\text{Sin}[a - b^2/(4*c)])/c^{(3/2)}$$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.308, Rules used = {3942, 3928, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \sin(a + bx + cx^2) dx \\ & \downarrow 3942 \\ & - \frac{b \int \sin(cx^2 + bx + a) dx}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \\ & \downarrow 3928 \\ & - \frac{b \left(\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx + \cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx \right)}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \\ & \downarrow 3832 \\ & - \frac{b \left(\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \\ & \downarrow 3833 \\ & - \frac{b \left(\frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} - \frac{\cos(a + bx + cx^2)}{2c} \end{aligned}$$

input

$$\text{Int}[x*\text{Sin}[a + b*x + c*x^2], x]$$

output

$$-1/2 \operatorname{Cos}[a + b*x + c*x^2]/c - (b*((\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{Cos}[a - b^2/(4*c)])*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])])))/\operatorname{Sqrt}[c] + (\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}]])*\operatorname{Sin}[a - b^2/(4*c)])/\operatorname{Sqrt}[c]))/(2*c)$$

Defintions of rubi rules used

rule 3832

$$\operatorname{Int}[\operatorname{Sin}[(d_.)*((e_.) + (f_.)*(x_))^2], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\operatorname{Pi}/2]/(f*\operatorname{Rt}[d, 2]))*\operatorname{FresnelS}[\operatorname{Sqrt}[2/\operatorname{Pi}]*\operatorname{Rt}[d, 2]*(e + f*x)], x] /; \operatorname{FreeQ}[\{d, e, f\}, x]$$

rule 3833

$$\operatorname{Int}[\operatorname{Cos}[(d_.)*((e_.) + (f_.)*(x_))^2], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\operatorname{Pi}/2]/(f*\operatorname{Rt}[d, 2]))*\operatorname{FresnelC}[\operatorname{Sqrt}[2/\operatorname{Pi}]*\operatorname{Rt}[d, 2]*(e + f*x)], x] /; \operatorname{FreeQ}[\{d, e, f\}, x]$$

rule 3928

$$\begin{aligned} \operatorname{Int}[\operatorname{Sin}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2], x_{\text{Symbol}}] &\rightarrow \operatorname{Simp}[\operatorname{Cos}[(b^2 - 4*a*c)/(4*c)]] \\ \operatorname{Int}[\operatorname{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \operatorname{Simp}[\operatorname{Sin}[(b^2 - 4*a*c)/(4*c)]] \\ \operatorname{Int}[\operatorname{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x] \&& \operatorname{NeQ}[b^2 - 4*a*c, 0] \end{aligned}$$

rule 3942

$$\begin{aligned} \operatorname{Int}[((d_.) + (e_.)*(x_))*\operatorname{Sin}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2], x_{\text{Symbol}}] &\rightarrow \operatorname{Simp}[-e*(\operatorname{Cos}[a + b*x + c*x^2]/(2*c)), x] + \operatorname{Simp}[(2*c*d - b*e)/(2*c)] \\ \operatorname{Int}[\operatorname{Sin}[a + b*x + c*x^2], x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e\}, x] \&& \operatorname{NeQ}[2*c*d - b*e, 0] \end{aligned}$$

Maple [A] (verified)

Time = 0.95 (sec), antiderivative size = 100, normalized size of antiderivative = 0.81

method	result
default	$-\frac{\cos(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{b^2-ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) - \sin\left(\frac{b^2-ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)\right)}{4c^{\frac{3}{2}}}$
risch	$-\frac{ib\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} - \frac{ib\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} - \frac{\cos(cx^2+bx+a)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x\cos\left(\frac{b^2-ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\sqrt{2}\sqrt{\pi}x\sin\left(\frac{b^2-ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\cos\left(\frac{b^2-ac}{4c}\right)\sqrt{2}\sqrt{\pi}}{\sqrt{c}}$

input `int(x*sin(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output `-1/2*cos(c*x^2+b*x+a)/c-1/4*b/c^(3/2)*2^(1/2)*Pi^(1/2)*(cos((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))-sin((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.96

$$\int x \sin(a + bx + cx^2) dx = \frac{\sqrt{2}\pi b \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi b \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2-4ac}{4c}\right) + 2c \cos(cx^2)}{4c^2}$$

input `integrate(x*sin(c*x^2+b*x+a),x, algorithm="fricas")`

output

```
-1/4*(sqrt(2)*pi*b*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) + sqrt(2)*pi*b*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c) + 2*c*cos(c*x^2 + b*x + a))/c^2
```

Sympy [F]

$$\int x \sin(a + bx + cx^2) dx = \int x \sin(a + bx + cx^2) dx$$

input

```
integrate(x*sin(c*x**2+b*x+a),x)
```

output

```
Integral(x*sin(a + b*x + c*x**2), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 579, normalized size of antiderivative = 4.71

$$\int x \sin(a + bx + cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*sin(c*x^2+b*x+a),x, algorithm="maxima")
```

output

```

1/16*((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x +
I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4
*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/4*(b^2 - 4*a*c)/c) + ((I - 1)*sqrt(
2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I
+ 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))
- 1))*b^2*sin(-1/4*(b^2 - 4*a*c)/c) - 2*(((I + 1)*sqrt(2)*sqrt(pi)*(erf(1
/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(
pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/
4*(b^2 - 4*a*c)/c) + (- (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2
+ 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4
*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/4*(b^2 - 4*a*c)/c))*x
- 4*(c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + e^(-1/4*(4*I*c^2*x^2
+ 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^2 - 4*a*c)/c) + c*(I*e^(1/4*(4*I*c^2*x^2
+ 4*I*b*c*x + I*b^2)/c) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sqrt(
(4*c^2*x^2 + 4*b*c*x + b^2)/c))/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.49

$$\begin{aligned}
& \int x \sin(a + bx + cx^2) dx \\
&= -\frac{\frac{i \sqrt{2} \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2x + \frac{b}{c}\right) \left(-\frac{ic}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{ib^2 - 4iac}{4c}\right)}}{\left(-\frac{ic}{|c|} + 1\right) \sqrt{|c|}} + 2 e^{(icx^2 + ibx + ia)}}{8c} \\
&\quad - \frac{-\frac{i \sqrt{2} \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2x + \frac{b}{c}\right) \left(\frac{ic}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{-ib^2 + 4iac}{4c}\right)}}{\left(\frac{ic}{|c|} + 1\right) \sqrt{|c|}} + 2 e^{(-icx^2 - ibx - ia)}}{8c}
\end{aligned}$$

input `integrate(x*sin(c*x^2+b*x+a),x, algorithm="giac")`

output

```
-1/8*(I*sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) + 2*e^(I*c*x^2 + I*b*x + I*a))/c - 1/8*(-I*sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) + 2*e^(-I*c*x^2 - I*b*x - I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \sin(a + bx + cx^2) dx = \int x \sin(cx^2 + bx + a) dx$$

input

```
int(x*sin(a + b*x + c*x^2),x)
```

output

```
int(x*sin(a + b*x + c*x^2), x)
```

Reduce [F]

$$\int x \sin(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a) x dx$$

input

```
int(x*sin(c*x^2+b*x+a),x)
```

output

```
int(sin(a + b*x + c*x**2)*x,x)
```

3.3 $\int \sin(a + bx + cx^2) dx$

Optimal result	58
Mathematica [A] (verified)	58
Rubi [A] (verified)	59
Maple [A] (verified)	60
Fricas [A] (verification not implemented)	61
Sympy [A] (verification not implemented)	61
Maxima [C] (verification not implemented)	62
Giac [C] (verification not implemented)	62
Mupad [B] (verification not implemented)	63
Reduce [F]	63

Optimal result

Integrand size = 11, antiderivative size = 97

$$\int \sin(a + bx + cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\ + \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{\sqrt{c}}$$

output

```
1/2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(1/2)+1/2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.21 (sec), antiderivative size = 84, normalized size of antiderivative = 0.87

$$\int \sin(a + bx + cx^2) dx \\ = \frac{\sqrt{\frac{\pi}{2}} \left(\cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \right)}{\sqrt{c}}$$

input `Integrate[Sin[a + b*x + c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot (\text{Cos}[a - b^2/(4c]) \cdot \text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] + \text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] \cdot \text{Sin}[a - b^2/(4c)]))}{\text{Sqrt}[c]}$$

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3928, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(a + bx + cx^2) \, dx \\
 & \downarrow 3928 \\
 & \sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) \, dx + \cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b + 2cx)^2}{4c}\right) \, dx \\
 & \downarrow 3832 \\
 & \sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) \, dx + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\
 & \downarrow 3833 \\
 & \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}
 \end{aligned}$$

input `Int[Sin[a + b*x + c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{Cos}[a - b^2/(4c]) \cdot \text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])])}{\text{Sqrt}[c]} + \frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] \cdot \text{Sin}[a - b^2/(4c)])}{\text{Sqrt}[c]}$$

Definitions of rubi rules used

rule 3832 $\text{Int}[\text{Sin}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\text{Int}[\text{Cos}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\text{Int}[\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \& \text{NeQ}[b^2 - 4*a*c, 0]$

Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{b^2-ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)-\sin\left(\frac{b^2-ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)\right)}{2\sqrt{c}}$	82
risch	$\frac{i\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}} + \frac{i\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}}$	101

input `int(sin(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2}2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}\text{cos}((1/4b^2-a*c)/c)\text{FresnelS}(2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}(c*x+1/2*b))-\text{sin}((1/4b^2-a*c)/c)\text{FresnelC}(2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}(c*x+1/2*b)))$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.05

$$\int \sin(a + bx + cx^2) dx \\ = \frac{\sqrt{2}\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi\sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2 - 4ac}{4c}\right)}{2c}$$

input `integrate(sin(c*x^2+b*x+a),x, algorithm="fricas")`

output `1/2*(sqrt(2)*pi*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) + sqrt(2)*pi*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c))/c`

Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.91

$$\int \sin(a + bx + cx^2) dx \\ = \frac{\sqrt{2}\sqrt{\pi} \left(\sin\left(a - \frac{b^2}{4c}\right) C\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) + \cos\left(a - \frac{b^2}{4c}\right) S\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) \right) \sqrt{\frac{1}{c}}}{2}$$

input `integrate(sin(c*x**2+b*x+a),x)`

output `sqrt(2)*sqrt(pi)*(sin(a - b**2/(4*c))*fresnelc(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))) + cos(a - b**2/(4*c))*fresnels(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))))*sqrt(1/c)/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.15

$$\int \sin(a + bx + cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(\left(-(i+1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i-1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) + \left(-(i-1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i+1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{-2icx-ib}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

input `integrate(sin(c*x^2+b*x+a),x, algorithm="maxima")`

output
$$\begin{aligned} & -\frac{1}{8}\sqrt{2}\sqrt{\pi}(\operatorname{erf}\left(\frac{i}{2}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4ic}{4c}\right)} \\ & + \operatorname{erf}\left(\frac{-i}{2}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4ic}{4c}\right)}) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.39

$$\int \sin(a + bx + cx^2) dx = \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) e^{\left(-\frac{ib^2-4ic}{4c}\right)}}{4\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2+4ic}{4c}\right)}}{4\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}}$$

input `integrate(sin(c*x^2+b*x+a),x, algorithm="giac")`

output
$$\begin{aligned} & \frac{1}{4}I\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) \\ & + \frac{1}{4}I\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.02

$$\int \sin(a + bx + cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} \cos\left(\frac{4ac - b^2}{4c}\right) S\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2} + \frac{\sqrt{2} \sqrt{\pi} \sin\left(\frac{4ac - b^2}{4c}\right) C\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2}$$

input `int(sin(a + b*x + c*x^2),x)`

output `(2^(1/2)*pi^(1/2)*cos((4*a*c - b^2)/(4*c))*fresnels((2^(1/2)*(b/2 + c*x)*(1/c)^(1/2))/pi^(1/2))*(1/c)^(1/2))/2 + (2^(1/2)*pi^(1/2)*sin((4*a*c - b^2)/(4*c))*fresnelc((2^(1/2)*(b/2 + c*x)*(1/c)^(1/2))/pi^(1/2))*(1/c)^(1/2))/2`

Reduce [F]

$$\int \sin(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a) dx$$

input `int(sin(c*x^2+b*x+a),x)`

output `int(sin(a + b*x + c*x**2),x)`

3.4 $\int \frac{\sin(a+bx+cx^2)}{x} dx$

Optimal result	64
Mathematica [N/A]	64
Rubi [N/A]	65
Maple [N/A]	65
Fricas [N/A]	66
Sympy [N/A]	66
Maxima [N/A]	67
Giac [N/A]	67
Mupad [N/A]	67
Reduce [N/A]	68

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \text{Int}\left(\frac{\sin(a + bx + cx^2)}{x}, x\right)$$

output `Defer(Int)(sin(c*x^2+b*x+a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 4.22 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(a + bx + cx^2)}{x} dx$$

input `Integrate[Sin[a + b*x + c*x^2]/x, x]`

output `Integrate[Sin[a + b*x + c*x^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3950}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(a + bx + cx^2)}{x} dx$$

↓ 3950

$$\int \frac{\sin(a + bx + cx^2)}{x} dx$$

input `Int[Sin[a + b*x + c*x^2]/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3950 `Int[((d_.) + (e_._)*(x_._))^((m_._))*Sin[(a_._) + (b_._)*(x_._) + (c_._)*(x_._)^2]^(n_._), x_Symbol] :> Unintegrable[(d + e*x)^m*Sin[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin(cx^2 + bx + a)}{x} dx$$

input `int(sin(c*x^2+b*x+a)/x, x)`

output `int(sin(c*x^2+b*x+a)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)}{x} dx$$

input `integrate(sin(c*x^2+b*x+a)/x,x, algorithm="fricas")`

output `integral(sin(c*x^2 + b*x + a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(a + bx + cx^2)}{x} dx$$

input `integrate(sin(c*x**2+b*x+a)/x,x)`

output `Integral(sin(a + b*x + c*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)}{x} dx$$

input `integrate(sin(c*x^2+b*x+a)/x,x, algorithm="maxima")`

output `integrate(sin(c*x^2 + b*x + a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)}{x} dx$$

input `integrate(sin(c*x^2+b*x+a)/x,x, algorithm="giac")`

output `integrate(sin(c*x^2 + b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 38.78 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)}{x} dx$$

input `int(sin(a + b*x + c*x^2)/x,x)`

output `int(sin(a + b*x + c*x^2)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)}{x} dx$$

input `int(sin(c*x^2+b*x+a)/x,x)`

output `int(sin(a + b*x + c*x**2)/x,x)`

3.5 $\int \left(-\frac{b \cos(a+bx+cx^2)}{x} + \frac{\sin(a+bx+cx^2)}{x^2} \right) dx$

Optimal result	69
Mathematica [A] (verified)	70
Rubi [A] (verified)	70
Maple [F]	71
Fricas [A] (verification not implemented)	71
Sympy [F]	72
Maxima [F]	72
Giac [F]	72
Mupad [F(-1)]	73
Reduce [F]	73

Optimal result

Integrand size = 33, antiderivative size = 110

$$\begin{aligned} & \int \left(-\frac{b \cos(a+bx+cx^2)}{x} + \frac{\sin(a+bx+cx^2)}{x^2} \right) dx \\ &= \sqrt{c}\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad - \sqrt{c}\sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) - \frac{\sin(a+bx+cx^2)}{x} \end{aligned}$$

output

```
c^(1/2)*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))-c^(1/2)*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)-sin(c*x^2+b*x+a)/x
```

Mathematica [A] (verified)

Time = 0.89 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx \\ &= \sqrt{c}\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ & - \frac{\sqrt{c}\sqrt{2\pi}x \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) + \sin(a + x(b + cx))}{x} \end{aligned}$$

input `Integrate[-((b*Cos[a + b*x + c*x^2])/x) + Sin[a + b*x + c*x^2]/x^2, x]`

output `Sqrt[c]*Sqrt[2*Pi]*Cos[a - b^2/(4*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] - (Sqrt[c]*Sqrt[2*Pi]*x*x*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] *Sin[a - b^2/(4*c)] + Sin[a + x*(b + c*x)])/x`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.030$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \left(\frac{\sin(a + bx + cx^2)}{x^2} - \frac{b \cos(a + bx + cx^2)}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \sqrt{2\pi}\sqrt{c} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \sqrt{2\pi}\sqrt{c} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \\ & \quad \frac{\sin(a + bx + cx^2)}{x} \end{aligned}$$

input `Int[-((b*Cos[a + b*x + c*x^2])/x) + Sin[a + b*x + c*x^2]/x^2, x]`

output

```
Sqrt[c]*Sqrt[2*Pi]*Cos[a - b^2/(4*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] - Sqrt[c]*Sqrt[2*Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*Sin[a - b^2/(4*c)] - Sin[a + b*x + c*x^2]/x
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [F]

$$\int \left(-\frac{b \cos(cx^2 + bx + a)}{x} + \frac{\sin(cx^2 + bx + a)}{x^2} \right) dx$$

input

```
int(-b*cos(c*x^2+b*x+a)/x+sin(c*x^2+b*x+a)/x^2,x)
```

output

```
int(-b*cos(c*x^2+b*x+a)/x+sin(c*x^2+b*x+a)/x^2,x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 117, normalized size of antiderivative = 1.06

$$\int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx \\ = \frac{\sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2-4ac}{4c}\right) - \sin(cx^2 + bx + a)}{x}$$

input

```
integrate(-b*cos(c*x^2+b*x+a)/x+sin(c*x^2+b*x+a)/x^2,x, algorithm="fricas")
```

output

```
(sqrt(2)*pi*x*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) - sqrt(2)*pi*x*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c) - sin(c*x^2 + bx + a))/x
```

Sympy [F]

$$\int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx$$

$$= - \int \left(-\frac{\sin(a + bx + cx^2)}{x^2} \right) dx - \int \frac{b \cos(a + bx + cx^2)}{x} dx$$

input `integrate(-b*cos(c*x**2+b*x+a)/x+sin(c*x**2+b*x+a)/x**2,x)`

output `-Integral(-sin(a + b*x + c*x**2)/x**2, x) - Integral(b*cos(a + b*x + c*x**2)/x, x)`

Maxima [F]

$$\int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx$$

$$= \int -\frac{b \cos(cx^2 + bx + a)}{x} + \frac{\sin(cx^2 + bx + a)}{x^2} dx$$

input `integrate(-b*cos(c*x^2+b*x+a)/x+sin(c*x^2+b*x+a)/x^2,x, algorithm="maxima")`

output `integrate(-b*cos(c*x^2 + b*x + a)/x + sin(c*x^2 + b*x + a)/x^2, x)`

Giac [F]

$$\int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx$$

$$= \int -\frac{b \cos(cx^2 + bx + a)}{x} + \frac{\sin(cx^2 + bx + a)}{x^2} dx$$

input `integrate(-b*cos(c*x^2+b*x+a)/x+sin(c*x^2+b*x+a)/x^2,x, algorithm="giac")`

output `integrate(-b*cos(c*x^2 + b*x + a)/x + sin(c*x^2 + b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx \\ &= \int \frac{\sin(cx^2 + bx + a)}{x^2} - \frac{b \cos(cx^2 + bx + a)}{x} dx \end{aligned}$$

input `int(sin(a + b*x + c*x^2)/x^2 - (b*cos(a + b*x + c*x^2))/x, x)`

output `int(sin(a + b*x + c*x^2)/x^2 - (b*cos(a + b*x + c*x^2))/x, x)`

Reduce [F]

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx + cx^2)}{x} + \frac{\sin(a + bx + cx^2)}{x^2} \right) dx \\ &= \frac{2(\int \cos(cx^2 + bx + a) dx) bcx - \sin(cx^2 + bx + a) b - 2acx}{bx} \end{aligned}$$

input `int(-b*cos(c*x^2+b*x+a)/x+sin(c*x^2+b*x+a)/x^2,x)`

output `(2*int(cos(a + b*x + c*x**2),x)*b*c*x - sin(a + b*x + c*x**2)*b - 2*a*c*x)/(b*x)`

3.6 $\int x^2 \sin(a + bx - cx^2) dx$

Optimal result	74
Mathematica [A] (verified)	75
Rubi [A] (verified)	75
Maple [A] (verified)	78
Fricas [A] (verification not implemented)	79
Sympy [F]	80
Maxima [C] (verification not implemented)	80
Giac [C] (verification not implemented)	81
Mupad [F(-1)]	82
Reduce [F]	82

Optimal result

Integrand size = 16, antiderivative size = 251

$$\begin{aligned} \int x^2 \sin(a + bx - cx^2) dx &= \frac{b \cos(a + bx - cx^2)}{4c^2} + \frac{x \cos(a + bx - cx^2)}{2c} \\ &+ \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &+ \frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ &- \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{4c^{5/2}} \\ &+ \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} \end{aligned}$$

```
output 1/4*b*cos(-c*x^2+b*x+a)/c^2+1/2*x*cos(-c*x^2+b*x+a)/c+1/4*2^(1/2)*Pi^(1/2)
*cos(a+1/4*b^2/c)*FresnelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)
)+1/8*b^2*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelS(1/2*(-2*c*x+b)/c^(1/2)
)*2^(1/2)/Pi^(1/2))/c^(5/2)-1/8*b^2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(-2*c*x+
b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)/c^(5/2)+1/4*2^(1/2)*Pi^(1/2)
*FresnelS(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)/c^(3/2
)
```

Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.66

$$\int x^2 \sin(a + bx - cx^2) dx \\ = \frac{2\sqrt{c}(b + 2cx) \cos(a + x(b - cx)) - \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a + \frac{b^2}{4c}\right) - b^2 \sin\left(a + \frac{b^2}{4c}\right)\right) - \sqrt{2\pi} c^{5/2}}{8c^{5/2}}$$

input `Integrate[x^2*Sin[a + b*x - c*x^2], x]`

output
$$(2*\operatorname{Sqrt}[c]*(b + 2*c*x)*\operatorname{Cos}[a + x*(b - c*x)] - \operatorname{Sqrt}[2*\pi]*\operatorname{FresnelC}[(-b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\pi])]*(2*c*\operatorname{Cos}[a + b^2/(4*c)] - b^2*\operatorname{Sin}[a + b^2/(4*c)]) - \operatorname{Sqrt}[2*\pi]*\operatorname{FresnelS}[(-b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\pi])]*(b^2*\operatorname{Cos}[a + b^2/(4*c)] + 2*c*\operatorname{Sin}[a + b^2/(4*c)]))/(8*c^{(5/2)})$$

Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3944, 3929, 25, 3832, 3833, 3942, 3928, 25, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sin(a + bx - cx^2) dx \\ \downarrow 3944 \\ \frac{b \int x \sin(-cx^2 + bx + a) dx}{2c} - \frac{\int \cos(-cx^2 + bx + a) dx}{2c} + \frac{x \cos(a + bx - cx^2)}{2c} \\ \downarrow 3929 \\ - \frac{\cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx - \sin\left(a + \frac{b^2}{4c}\right) \int -\sin\left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} + \\ \frac{b \int x \sin(-cx^2 + bx + a) dx}{2c} + \frac{x \cos(a + bx - cx^2)}{2c}$$

$$\begin{aligned}
& \downarrow 25 \\
& - \frac{\sin \left(a + \frac{b^2}{4c} \right) \int \sin \left(\frac{(b-2cx)^2}{4c} \right) dx + \cos \left(a + \frac{b^2}{4c} \right) \int \cos \left(\frac{(b-2cx)^2}{4c} \right) dx}{2c} + \\
& \frac{b \int x \sin (-cx^2 + bx + a) dx}{2c} + \frac{x \cos (a + bx - cx^2)}{2c} \\
& \downarrow 3832 \\
& - \frac{\cos \left(a + \frac{b^2}{4c} \right) \int \cos \left(\frac{(b-2cx)^2}{4c} \right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}}}{2c} + \\
& \frac{b \int x \sin (-cx^2 + bx + a) dx}{2c} + \frac{x \cos (a + bx - cx^2)}{2c} \\
& \downarrow 3833 \\
& \frac{b \int x \sin (-cx^2 + bx + a) dx}{2c} - \frac{-\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}}}{2c} + \\
& \frac{x \cos (a + bx - cx^2)}{2c} \\
& \downarrow 3942 \\
& - \frac{b \left(\frac{b \int \sin (-cx^2 + bx + a) dx}{2c} + \frac{\cos (a + bx - cx^2)}{2c} \right)}{2c} - \\
& \frac{-\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}}}{2c} + \frac{x \cos (a + bx - cx^2)}{2c} \\
& \downarrow 3928 \\
& - \frac{b \left(\frac{b \left(\sin \left(a + \frac{b^2}{4c} \right) \int \cos \left(\frac{(b-2cx)^2}{4c} \right) dx + \cos \left(a + \frac{b^2}{4c} \right) \int -\sin \left(\frac{(b-2cx)^2}{4c} \right) dx \right)}{2c} + \frac{\cos (a + bx - cx^2)}{2c} \right)}{2c} - \\
& \frac{-\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}}}{2c} + \frac{x \cos (a + bx - cx^2)}{2c} \\
& \downarrow 25 \\
& - \frac{b \left(\frac{b \left(\sin \left(a + \frac{b^2}{4c} \right) \int \cos \left(\frac{(b-2cx)^2}{4c} \right) dx - \cos \left(a + \frac{b^2}{4c} \right) \int \sin \left(\frac{(b-2cx)^2}{4c} \right) dx \right)}{2c} + \frac{\cos (a + bx - cx^2)}{2c} \right)}{2c} - \\
& \frac{-\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}}}{2c} + \frac{x \cos (a + bx - cx^2)}{2c}
\end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3832} \\
 & b \left(\frac{b \left(\sin \left(a + \frac{b^2}{4c} \right) \int \cos \left(\frac{(b-2cx)^2}{4c} \right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} + \frac{\cos(a+bx-cx^2)}{2c} \right) - \\
 & - \frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} + \frac{x \cos(a+bx-cx^2)}{2c} \\
 & \downarrow \text{3833} \\
 & b \left(\frac{b \left(\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} + \frac{\cos(a+bx-cx^2)}{2c} \right) - \\
 & - \frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} + \frac{x \cos(a+bx-cx^2)}{2c}
 \end{aligned}$$

input `Int[x^2*Sin[a + b*x - c*x^2], x]`

output `(x*Cos[a + b*x - c*x^2])/(2*c) - ((Sqrt[Pi/2]*Cos[a + b^2/(4*c)]*FresnelC[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/Sqrt[c]) - (Sqrt[Pi/2]*FresnelS[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a + b^2/(4*c)]/Sqrt[c]/(2*c) + (b*(Cos[a + b*x - c*x^2]/(2*c) + (b*((Sqrt[Pi/2]*Cos[a + b^2/(4*c)]*FresnelS[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/Sqrt[c] - (Sqrt[Pi/2]*FresnelC[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a + b^2/(4*c)]/Sqrt[c]))/(2*c)))/(2*c)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_._)*(x_))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 $\text{Int}[\cos[(d_.) * ((e_.) + (f_.) * (x_))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi/2}) / (f * \text{Rt}[d, 2]) * \text{FresnelC}[\sqrt{2/\pi} * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\text{Int}[\sin[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\cos[(b^2 - 4*a*c) / (4*c)] \quad \text{Int}[\sin[(b + 2*c*x)^2 / (4*c)], x], x] - \text{Simp}[\sin[(b^2 - 4*a*c) / (4*c)] \quad \text{Int}[\cos[(b + 2*c*x)^2 / (4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3929 $\text{Int}[\cos[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\cos[(b^2 - 4*a*c) / (4*c)] \quad \text{Int}[\cos[(b + 2*c*x)^2 / (4*c)], x], x] + \text{Simp}[\sin[(b^2 - 4*a*c) / (4*c)] \quad \text{Int}[\sin[(b + 2*c*x)^2 / (4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3942 $\text{Int}[((d_.) + (e_.) * (x_)) * \sin[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-e) * (\cos[a + b*x + c*x^2] / (2*c)), x] + \text{Simp}[(2*c*d - b*e) / (2*c) \quad \text{Int}[\sin[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

rule 3944 $\text{Int}[((d_.) + (e_.) * (x_))^{(m_)} * \sin[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-e) * (d + e*x)^{(m - 1)} * (\cos[a + b*x + c*x^2] / (2*c)), x] + (-\text{Simp}[(b*e - 2*c*d) / (2*c) \quad \text{Int}[(d + e*x)^{(m - 1)} * \sin[a + b*x + c*x^2], x], x] + \text{Simp}[e^{2*(m - 1) / (2*c)} \quad \text{Int}[(d + e*x)^{(m - 2)} * \cos[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[b*e - 2*c*d, 0] \&& \text{GtQ}[m, 1]$

Maple [A] (verified)

Time = 1.60 (sec), antiderivative size = 224, normalized size of antiderivative = 0.89

method	result
default	$\frac{x \cos(-cx^2+bx+a)}{2c} + b \left(\frac{\cos(-cx^2+bx+a)}{2c} + \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2+ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) + \sin\left(\frac{b^2+ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) \right)}{4c\sqrt{-c}} \right)$
risch	$\frac{ib^2\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\text{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{16c^2\sqrt{ic}} + \frac{\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\text{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} + \frac{ib^2\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\text{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{16c^2\sqrt{-ic}} -$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2\cos\left(\frac{b^2+ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} + \frac{\sqrt{2}\sqrt{\pi}x^2\sin\left(\frac{b^2+ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} + \frac{\sqrt{2}\pi^{\frac{3}{2}}\cos\left(\frac{4ac+b^2}{4c}\right)}{}$

input `int(x^2*sin(-c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output `1/2*x*cos(-c*x^2+b*x+a)/c+1/2*b/c*(1/2*cos(-c*x^2+b*x+a)/c+1/4*b/c*2^(1/2)*Pi^(1/2)/(-c)^(1/2)*(cos((1/4*b^2+a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b))+sin((1/4*b^2+a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b)))-1/4/c*2^(1/2)*Pi^(1/2)/(-c)^(1/2)*(cos((1/4*b^2+a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b))-sin((1/4*b^2+a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.71

$$\int x^2 \sin(a + bx - cx^2) dx = \frac{\sqrt{2} \left(\pi b^2 \sin\left(\frac{b^2+4ac}{4c}\right) - 2\pi c \cos\left(\frac{b^2+4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2} \left(\pi b^2 \cos\left(\frac{b^2+4ac}{4c}\right) + 2\pi c \sin\left(\frac{b^2+4ac}{4c}\right) \right)}{8c^3}$$

input `integrate(x^2*sin(-c*x^2+b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{8} \sqrt{2} (\pi b^2 \sin(\frac{1}{4}(b^2 + 4ac)/c) - 2\pi c \cos(\frac{1}{4}(b^2 + 4ac)/c)) \sqrt{c/\pi} \operatorname{fresnel_cos}(\frac{1}{2}\sqrt{2}(2cx - b)\sqrt{c/\pi}/c) - \sqrt{2} (\pi b^2 \cos(\frac{1}{4}(b^2 + 4ac)/c) + 2\pi c \sin(\frac{1}{4}(b^2 + 4ac)/c)) \sqrt{c/\pi} \operatorname{fresnel_sin}(\frac{1}{2}\sqrt{2}(2cx - b)\sqrt{c/\pi}/c) + 2(2c^2x + b^2c) \cos(cx^2 - bx - a))/c^3$$

Sympy [F]

$$\int x^2 \sin(a + bx - cx^2) dx = \int x^2 \sin(a + bx - cx^2) dx$$

input `integrate(x**2*sin(-c*x**2+b*x+a),x)`

output `Integral(x**2*sin(a + b*x - c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.80 (sec) , antiderivative size = 1570, normalized size of antiderivative = 6.25

$$\int x^2 \sin(a + bx - cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*sin(-c*x^2+b*x+a),x, algorithm="maxima")`

output

$$\begin{aligned}
 & -\frac{1}{32} * (8 * (((I + 1) * \sqrt{2}) * \sqrt{\pi}) * (\operatorname{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) - (I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(1/2 * \sqrt{-(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) * b^2 * c^3 + 4 * ((I - 1) * \sqrt{2} * \operatorname{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c) - (I + 1) * \sqrt{2} * \operatorname{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c)) * c^4) * \cos(1/4 * (b^2 + 4 * a * c) / c) + (((I - 1) * \sqrt{2} * \sqrt{\pi}) * (\operatorname{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) - (I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(1/2 * \sqrt{-(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) * b^2 * c^3 + 4 * (- (I + 1) * \sqrt{2} * \operatorname{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c) + (I - 1) * \sqrt{2} * \operatorname{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c)) * c^4) * \sin(1/4 * (b^2 + 4 * a * c) / c) * x^3 + 12 * (((-(I + 1) * \sqrt{2} * \sqrt{\pi}) * (\operatorname{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) + (I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(1/2 * \sqrt{-(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) * b^3 * c^2 + 4 * (- (I - 1) * \sqrt{2} * \operatorname{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c) + (I + 1) * \sqrt{2} * \operatorname{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c)) * b * c^3) * \cos(1/4 * (b^2 + 4 * a * c) / c) + (((-(I - 1) * \sqrt{2} * \sqrt{\pi}) * (\operatorname{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) + (I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(1/2 * \sqrt{-(4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c})) - 1) * b^3 * c^2 + 4 * ((I + 1) * \sqrt{2} * \operatorname{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c) - (I - 1) * \sqrt{2} * \operatorname{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c)) * b * c^3) * \sin(1/4 * (b^2 + 4 * a * c) / c) * x^2 - 8 * (b * c^2 * (e^{(1/4 * (4 * I * c^2 * x^2 - 4 * I * b * c * x + I * b^2) / c)} - 1))
 \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec), antiderivative size = 229, normalized size of antiderivative = 0.91

$$\begin{aligned}
 & \int x^2 \sin(a + bx - cx^2) dx = \\
 & \frac{i \sqrt{2} \sqrt{\pi} (b^2 + 2ic) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2x - \frac{b}{c}\right) \left(-\frac{ic}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{ib^2 + 4ic}{4c}\right)}}{\left(-\frac{ic}{|c|} + 1\right) \sqrt{|c|}} - 2i \left(c \left(-2ix + \frac{ib}{c}\right) - 2ib\right) e^{(icx^2 - ibx - ia)} \\
 & - \frac{i \sqrt{2} \sqrt{\pi} (b^2 - 2ic) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2x - \frac{b}{c}\right) \left(\frac{ic}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{-ib^2 - 4ic}{4c}\right)}}{\left(\frac{ic}{|c|} + 1\right) \sqrt{|c|}} - 2i \left(c \left(-2ix + \frac{ib}{c}\right) - 2ib\right) e^{(-icx^2 + ibx + ia)} \\
 & \frac{16c^2}{16c^2}
 \end{aligned}$$

input `integrate(x^2*sin(-c*x^2+b*x+a),x, algorithm="giac")`

output

$$\begin{aligned} & -\frac{1}{16} \left(I * \sqrt{2} * \sqrt{\pi} * (b^2 + 2*I*c) * \operatorname{erf}(-\frac{1}{4} * \sqrt{2} * (2*x - b/c) * (-I*c / \operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)}) * e^{-(-1/4 * (I*b^2 + 4*I*a*c)/c) / ((-I*c/\operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)})} \right. \\ & \left. - 2*I*(c*(-2*I*x + I*b/c) - 2*I*b)*e^{(I*c*x^2 - I*b*x - I*a)/c^2} - \frac{1}{16} * (-I * \sqrt{2} * \sqrt{\pi} * (b^2 - 2*I*c) * \operatorname{erf}(-\frac{1}{4} * \sqrt{2} * (2*x - b/c) * (I*c/\operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)})) * e^{-(-1/4 * (-I*b^2 - 4*I*a*c)/c) / ((I*c/\operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)})} \right. \\ & \left. - 2*I*(c*(-2*I*x + I*b/c) - 2*I*b)*e^{(-I*c*x^2 + I*b*x + I*a)/c^2} \right) \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int x^2 \sin(a + bx - cx^2) dx = \int x^2 \sin(-cx^2 + bx + a) dx$$

input

```
int(x^2*sin(a + b*x - c*x^2),x)
```

output

```
int(x^2*sin(a + b*x - c*x^2), x)
```

Reduce [F]

$$\begin{aligned} & \int x^2 \sin(a + bx - cx^2) dx \\ & = \frac{6 \cos(-cx^2 + bx + a) bc + 12 \cos(-cx^2 + bx + a) c^2 x + 24 \left(\int \frac{x^2}{\tan(-\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) b^2 c^2 - 6 \left(\int \frac{...}{\tan(...)^2 + 1} dx \right) b^3 c^3}{...} \end{aligned}$$

input

```
int(x^2*sin(-c*x^2+b*x+a),x)
```

output

$$\begin{aligned} & (6 * \cos(a + b*x - c*x**2) * b*c + 12 * \cos(a + b*x - c*x**2) * c**2*x + 24 * \int(x * \\ & * 2 / (\tan((a + b*x - c*x**2)/2)**2 + 1), x) * b**2*c**2 - 6 * \int(1 / (\tan((a + b*x - c*x**2)/2)**2 + 1), x) * b**4 - 24 * \int(1 / (\tan((a + b*x - c*x**2)/2)**2 + 1), x) * c**2 + 3 * \sin(a + b*x - c*x**2) * b**3 + 6 * \sin(a + b*x - c*x**2) * b**2*c*x + 3 * b**4*x - 4 * b**2*c**2*x**3 - 6 * b*c + 12 * c**2*x) / (24 * c**3) \end{aligned}$$

3.7 $\int x \sin(a + bx - cx^2) dx$

Optimal result	83
Mathematica [A] (verified)	83
Rubi [A] (verified)	84
Maple [A] (verified)	86
Fricas [A] (verification not implemented)	86
Sympy [F]	87
Maxima [C] (verification not implemented)	87
Giac [C] (verification not implemented)	88
Mupad [F(-1)]	89
Reduce [F]	89

Optimal result

Integrand size = 14, antiderivative size = 124

$$\int x \sin(a + bx - cx^2) dx = \frac{\cos(a + bx - cx^2)}{2c} + \frac{b\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} - \frac{b\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}}$$

output

```
1/2*cos(-c*x^2+b*x+a)/c+1/4*b*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelS(1
/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)-1/4*b*2^(1/2)*Pi^(1/2)*Fre
snelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.94

$$\int x \sin(a + bx - cx^2) dx = \frac{2\sqrt{c} \cos(a + x(b - cx)) - b\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + b\sqrt{2\pi} \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{4c^{3/2}}$$

input

```
Integrate[x*Sin[a + b*x - c*x^2], x]
```

output
$$(2*\text{Sqrt}[c]*\text{Cos}[a + x*(b - c*x)] - b*\text{Sqrt}[2*\pi]*\text{Cos}[a + b^2/(4*c)]*\text{FresnelS} [(-b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])] + b*\text{Sqrt}[2*\pi]*\text{FresnelC}[(-b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])]*\text{Sin}[a + b^2/(4*c)])/(4*c^{(3/2)})$$

Rubi [A] (verified)

Time = 0.34 (sec), antiderivative size = 126, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3942, 3928, 25, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin(a + bx - cx^2) dx \\
 & \downarrow 3942 \\
 & \frac{b \int \sin(-cx^2 + bx + a) dx}{2c} + \frac{\cos(a + bx - cx^2)}{2c} \\
 & \downarrow 3928 \\
 & \frac{b \left(\sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx + \cos\left(a + \frac{b^2}{4c}\right) \int -\sin\left(\frac{(b-2cx)^2}{4c}\right) dx \right)}{2c} + \frac{\cos(a + bx - cx^2)}{2c} \\
 & \downarrow 25 \\
 & \frac{b \left(\sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx - \cos\left(a + \frac{b^2}{4c}\right) \int \sin\left(\frac{(b-2cx)^2}{4c}\right) dx \right)}{2c} + \frac{\cos(a + bx - cx^2)}{2c} \\
 & \downarrow 3832 \\
 & \frac{b \left(\sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} + \frac{\cos(a + bx - cx^2)}{2c} \\
 & \downarrow 3833 \\
 & \frac{b \left(\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} + \frac{\cos(a + bx - cx^2)}{2c}
 \end{aligned}$$

input $\int x \sin[a + b*x - c*x^2] dx$

output $\frac{\cos[a + b*x - c*x^2]/(2*c) + (b*((\sqrt{\pi}/2)*\cos[a + b^2/(4*c)]*\text{FresnelS}[(b - 2*c*x)/(\sqrt{c}*\sqrt{2*\pi})]))/\sqrt{c} - (\sqrt{\pi}/2)*\text{FresnelC}[(b - 2*c*x)/(\sqrt{c}*\sqrt{2*\pi})]*\sin[a + b^2/(4*c)]/\sqrt{c}}{(2*c)}$

Definitions of rubi rules used

rule 25 $\int -(f_x), x \rightarrow \text{Simp}[\text{Identity}[-1] \quad \int f_x, x], x$

rule 3832 $\int \sin[d_*] * ((e_*) + (f_*) * (x_*)^2), x \rightarrow \text{Simp}[(\sqrt{\pi}/2) / (f * \text{Rt}[d, 2]) * \text{FresnelS}[\sqrt{2/\pi} * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\int \cos[d_*] * ((e_*) + (f_*) * (x_*)^2), x \rightarrow \text{Simp}[(\sqrt{\pi}/2) / (f * \text{Rt}[d, 2]) * \text{FresnelC}[\sqrt{2/\pi} * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\int \sin[(a_*) + (b_*) * (x_*) + (c_*) * (x_*)^2], x \rightarrow \text{Simp}[\cos[(b^2 - 4*a*c)/(4*c)] \quad \int \sin[(b + 2*c*x)^2/(4*c)], x], x] - \text{Simp}[\sin[(b^2 - 4*a*c)/(4*c)] \quad \int \cos[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3942 $\int ((d_*) + (e_*) * (x_*)) * \sin[(a_*) + (b_*) * (x_*) + (c_*) * (x_*)^2], x \rightarrow \text{Simp}[-e * (\cos[a + b*x + c*x^2]/(2*c)), x] + \text{Simp}[(2*c*d - b*e)/(2*c) \quad \int \sin[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.88

method	result
default	$\frac{\cos(-cx^2+bx+a)}{2c} + \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2+ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) + \sin\left(\frac{b^2+ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) \right)}{4c\sqrt{-c}}$
risch	$\frac{ib\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\text{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} + \frac{ib\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\text{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} + \frac{\cos(-cx^2+bx+a)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x\cos\left(\frac{b^2+ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} + \frac{\sqrt{2}\sqrt{\pi}x\sin\left(\frac{b^2+ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}}{2\sqrt{-c}} \left[\cos\left(\frac{b^2+ac}{4c}\right) - \frac{\sqrt{2}\sqrt{\pi}}{2\sqrt{-c}} \right]$

input `int(x*sin(-c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2}\cos(-cx^2+bx+a)/c + \frac{1}{4}b/c*2^{(1/2)}*\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(\cos((1/4)b^2+a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(-c*x+1/2*b))+\sin((1/4)b^2+a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(-c*x+1/2*b)))$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.02

$$\int x \sin(a + bx - cx^2) dx = \frac{-\sqrt{2}\pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi b \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) - 2c \cos(cx^2 - b^2/4c)}{4c^2}$$

input `integrate(x*sin(-c*x^2+b*x+a),x, algorithm="fricas")`

output

```
-1/4*(sqrt(2)*pi*b*sqrt(c/pi)*cos(1/4*(b^2 + 4*a*c)/c)*fresnel_sin(1/2*sqr
t(2)*(2*c*x - b)*sqrt(c/pi)/c) - sqrt(2)*pi*b*sqrt(c/pi)*fresnel_cos(1/2*s
qrt(2)*(2*c*x - b)*sqrt(c/pi)/c)*sin(1/4*(b^2 + 4*a*c)/c) - 2*c*cos(c*x^2
- b*x - a))/c^2
```

Sympy [F]

$$\int x \sin(a + bx - cx^2) dx = \int x \sin(a + bx - cx^2) dx$$

input

```
integrate(x*sin(-c*x**2+b*x+a),x)
```

output

```
Integral(x*sin(a + b*x - c*x**2), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 580, normalized size of antiderivative = 4.68

$$\int x \sin(a + bx - cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*sin(-c*x^2+b*x+a),x, algorithm="maxima")
```

output

```
1/16*((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(1/4*(b^2 + 4*a*c)/c) + ((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(1/4*(b^2 + 4*a*c)/c) - 2*((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(1/4*(b^2 + 4*a*c)/c) + ((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(1/4*(b^2 + 4*a*c)/c)*x + 4*(c*(e^(1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) + e^(-1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*cos(1/4*(b^2 + 4*a*c)/c) - c*(I*e^(1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*sin(1/4*(b^2 + 4*a*c)/c))*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c)/(c^2*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.49

$$\int x \sin(a + bx - cx^2) dx$$

$$= -\frac{\frac{i \sqrt{2} \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2 x - \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{i b^2 + 4 i a c}{4 c}\right)}}{\left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}} - 2 e^{(i c x^2 - i b x - i a)}}{8 c}$$

$$- \frac{-\frac{i \sqrt{2} \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2 x - \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{-i b^2 - 4 i a c}{4 c}\right)}}{\left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}} - 2 e^{(-i c x^2 + i b x + i a)}}{8 c}$$

input `integrate(x*sin(-c*x^2+b*x+a),x, algorithm="giac")`

output

```
-1/8*(I*sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 2*e^(I*c*x^2 - I*b*x - I*a))/c - 1/8*(-I*sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x - b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) - 2*e^(-I*c*x^2 + I*b*x + I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \sin(a + bx - cx^2) dx = \int x \sin(-cx^2 + bx + a) dx$$

input

```
int(x*sin(a + b*x - c*x^2),x)
```

output

```
int(x*sin(a + b*x - c*x^2), x)
```

Reduce [F]

$$\int x \sin(a + bx - cx^2) dx = \int \sin(-cx^2 + bx + a) x dx$$

input

```
int(x*sin(-c*x^2+b*x+a),x)
```

output

```
int(sin(a + b*x - c*x**2)*x,x)
```

3.8 $\int \sin(a + bx - cx^2) dx$

Optimal result	90
Mathematica [A] (verified)	90
Rubi [A] (verified)	91
Maple [A] (verified)	92
Fricas [A] (verification not implemented)	93
Sympy [A] (verification not implemented)	93
Maxima [C] (verification not implemented)	94
Giac [C] (verification not implemented)	94
Mupad [B] (verification not implemented)	95
Reduce [F]	95

Optimal result

Integrand size = 12, antiderivative size = 98

$$\int \sin(a + bx - cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{\sqrt{c}}$$

output

```
1/2*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelS(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(1/2)-1/2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.20 (sec), antiderivative size = 89, normalized size of antiderivative = 0.91

$$\int \sin(a + bx - cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \left(-\cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \right)}{\sqrt{c}}$$

input `Integrate[Sin[a + b*x - c*x^2], x]`

output
$$\left(\frac{\sqrt{\pi/2} \cdot (-\text{Cos}[a + b^2/(4c]) \cdot \text{FresnelS}[(-b + 2c*x)/(Sqrt[c]*Sqrt[2*\pi])] + \text{FresnelC}[(-b + 2c*x)/(Sqrt[c]*Sqrt[2*\pi])] \cdot \text{Sin}[a + b^2/(4c)])}{\sqrt{c}} \right) / Sqrt[c]$$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3928, 25, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(a + bx - cx^2) \, dx \\
 & \downarrow 3928 \\
 & \sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) \, dx + \cos\left(a + \frac{b^2}{4c}\right) \int -\sin\left(\frac{(b - 2cx)^2}{4c}\right) \, dx \\
 & \downarrow 25 \\
 & \sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) \, dx - \cos\left(a + \frac{b^2}{4c}\right) \int \sin\left(\frac{(b - 2cx)^2}{4c}\right) \, dx \\
 & \downarrow 3832 \\
 & \sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) \, dx + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\
 & \downarrow 3833 \\
 & \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}
 \end{aligned}$$

input `Int[Sin[a + b*x - c*x^2], x]`

output

$$\frac{(\text{Sqrt}[\text{Pi}/2]*\text{Cos}[a + b^2/(4*c)]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}]]) - \text{Sqrt}[c] - (\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}]])*\text{Sin}[a + b^2/(4*c)])/\text{Sqrt}[c]}{\text{Sqrt}[c]}$$

Defintions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 3832 $\text{Int}[\text{Sin}[(d_*)*((e_*) + (f_*)*(x_))^2], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], \text{x}] /; \text{FreeQ}[\{d, e, f\}, \text{x}]$

rule 3833 $\text{Int}[\text{Cos}[(d_*)*((e_*) + (f_*)*(x_))^2], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], \text{x}] /; \text{FreeQ}[\{d, e, f\}, \text{x}]$

rule 3928 $\text{Int}[\text{Sin}[(a_*) + (b_*)*(x_) + (c_*)*(x_)^2], \text{x_Symbol}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], \text{x}], \text{x}] - \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], \text{x}], \text{x}] /; \text{FreeQ}[\{a, b, c\}, \text{x}] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)+\sin\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)\right)}{2\sqrt{-c}}$	87
risch	$\frac{i\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\text{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}}+\frac{i\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\text{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}}$	97

input $\text{int}(\text{sin}(-c*x^2+b*x+a), \text{x}, \text{method}=\text{_RETURNVERBOSE})$

output
$$\frac{1}{2} \cdot 2^{(1/2)} \cdot \text{Pi}^{(1/2)} / (-c)^{(1/2)} \cdot (\cos((1/4 \cdot b^2 + a \cdot c) / c) \cdot \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / (-c)^{(1/2)} \cdot (-c \cdot x + 1/2 \cdot b)) + \sin((1/4 \cdot b^2 + a \cdot c) / c) \cdot \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / (-c)^{(1/2)} \cdot (-c \cdot x + 1/2 \cdot b)))$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 107, normalized size of antiderivative = 1.09

$$\int \sin(a + bx - cx^2) dx \\ = -\frac{\sqrt{2}\pi\sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi\sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right)}{2c}$$

input `integrate(sin(-c*x^2+b*x+a),x, algorithm="fricas")`

output
$$-\frac{1}{2} \cdot (\sqrt{2} \cdot \pi \cdot \sqrt{c/\pi}) \cdot \cos(1/4 \cdot (b^2 + 4 \cdot a \cdot c) / c) \cdot \text{fresnel_sin}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x - b) \cdot \sqrt{c/\pi} / c) - \sqrt{2} \cdot \pi \cdot \sqrt{c/\pi} \cdot \text{fresnel_cos}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x - b) \cdot \sqrt{c/\pi} / c) \cdot \sin(1/4 \cdot (b^2 + 4 \cdot a \cdot c) / c) / c$$

Sympy [A] (verification not implemented)

Time = 0.21 (sec), antiderivative size = 94, normalized size of antiderivative = 0.96

$$\int \sin(a + bx - cx^2) dx \\ = \frac{\sqrt{2}\sqrt{\pi}\sqrt{-\frac{1}{c}} \left(\sin\left(a + \frac{b^2}{4c}\right) C\left(\frac{\sqrt{2}(b-2cx)}{2\sqrt{\pi}\sqrt{-c}}\right) + \cos\left(a + \frac{b^2}{4c}\right) S\left(\frac{\sqrt{2}(b-2cx)}{2\sqrt{\pi}\sqrt{-c}}\right) \right)}{2}$$

input `integrate(sin(-c*x**2+b*x+a),x)`

output
$$\sqrt{2} \cdot \sqrt{\pi} \cdot \sqrt{-1/c} \cdot (\sin(a + b^{**2}/(4*c)) \cdot \text{fresnelc}(\sqrt{2} \cdot (b - 2*c) \cdot x) / (2 * \sqrt{\pi} * \sqrt{-c})) + \cos(a + b^{**2}/(4*c)) \cdot \text{fresnels}(\sqrt{2} \cdot (b - 2*c) \cdot x) / (2 * \sqrt{\pi} * \sqrt{-c})) / 2$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14

$$\int \sin(a + bx - cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(\left((i+1) \cos\left(\frac{b^2+4ac}{4c}\right) + (i-1) \sin\left(\frac{b^2+4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx-i b}{2\sqrt{ic}}\right) + \left((i-1) \cos\left(\frac{b^2+4ac}{4c}\right) + (i+1) \sin\left(\frac{b^2+4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+i b}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

input `integrate(sin(-c*x^2+b*x+a),x, algorithm="maxima")`

output
$$\begin{aligned} & -\frac{1}{8}\sqrt{2}\sqrt{\pi} \left(((I+1)\cos(1/4*(b^2+4*a*c)/c) + (I-1)\sin(1/4*(b^2+4*a*c)/c)) \operatorname{erf}\left(\frac{1/2*(2*I*c*x - I*b)}{\sqrt{I*c}}\right) + ((I-1)\cos(1/4*(b^2+4*a*c)/c) + (I+1)\sin(1/4*(b^2+4*a*c)/c)) \operatorname{erf}\left(\frac{1/2*(2*I*c*x - I*b)}{\sqrt{-I*c}}\right) \right) / \sqrt{c} \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.40

$$\begin{aligned} \int \sin(a + bx - cx^2) dx = & -\frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x - \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2+4ic}{4c}\right)}}{4\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}} \\ & + \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x - \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2-4ic}{4c}\right)}}{4\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}} \end{aligned}$$

input `integrate(sin(-c*x^2+b*x+a),x, algorithm="giac")`

output
$$\begin{aligned} & -\frac{1}{4}I*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c))\right)*e^{-\left(-\frac{1}{4}*(I*b^2 + 4*I*a*c)/c\right)} / ((-I*c/abs(c) + 1)*sqrt(abs(c))) + \\ & \frac{1}{4}I*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}*(2*x - b/c)*(I*c/abs(c) + 1)*sqrt(abs(c))\right)*e^{-\left(-\frac{1}{4}*(-I*b^2 - 4*I*a*c)/c\right)} / ((I*c/abs(c) + 1)*sqrt(abs(c))) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.07

$$\int \sin(a + bx - cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}\left(\frac{b}{2}-cx\right)\sqrt{-\frac{1}{c}}}{\sqrt{\pi}}\right) \cos\left(\frac{b^2+4ac}{4c}\right) \sqrt{-\frac{1}{c}}}{2} + \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}\left(\frac{b}{2}-cx\right)\sqrt{-\frac{1}{c}}}{\sqrt{\pi}}\right) \sin\left(\frac{b^2+4ac}{4c}\right) \sqrt{-\frac{1}{c}}}{2}$$

input `int(sin(a + b*x - c*x^2),x)`

output `(2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*(b/2 - c*x)*(-1/c)^(1/2))/pi^(1/2))*cos((4*a*c + b^2)/(4*c))*(-1/c)^(1/2))/2 + (2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*(b/2 - c*x)*(-1/c)^(1/2))/pi^(1/2))*sin((4*a*c + b^2)/(4*c))*(-1/c)^(1/2))/2`

Reduce [F]

$$\int \sin(a + bx - cx^2) dx = \int \sin(-cx^2 + bx + a) dx$$

input `int(sin(-c*x^2+b*x+a),x)`

output `int(sin(a + b*x - c*x**2),x)`

3.9 $\int \frac{\sin(a+bx-cx^2)}{x} dx$

Optimal result	96
Mathematica [N/A]	96
Rubi [N/A]	97
Maple [N/A]	97
Fricas [N/A]	98
Sympy [N/A]	98
Maxima [N/A]	99
Giac [N/A]	99
Mupad [N/A]	99
Reduce [N/A]	100

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \text{Int}\left(\frac{\sin(a + bx - cx^2)}{x}, x\right)$$

output `Defer(Int)(sin(-c*x^2+b*x+a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 4.69 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(a + bx - cx^2)}{x} dx$$

input `Integrate[Sin[a + b*x - c*x^2]/x, x]`

output `Integrate[Sin[a + b*x - c*x^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3950}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(a + bx - cx^2)}{x} dx$$

↓ 3950

$$\int \frac{\sin(a + bx - cx^2)}{x} dx$$

input `Int[Sin[a + b*x - c*x^2]/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3950 `Int[((d_.) + (e_.)*(x_))^(m_.)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_.), x_Symbol] :> Unintegrable[(d + e*x)^m*Sin[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\sin(-cx^2 + bx + a)}{x} dx$$

input `int(sin(-c*x^2+b*x+a)/x, x)`

output `int(sin(-c*x^2+b*x+a)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)}{x} dx$$

input `integrate(sin(-c*x^2+b*x+a)/x,x, algorithm="fricas")`

output `integral(-sin(c*x^2 - b*x - a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(a + bx - cx^2)}{x} dx$$

input `integrate(sin(-c*x**2+b*x+a)/x,x)`

output `Integral(sin(a + b*x - c*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)}{x} dx$$

input `integrate(sin(-c*x^2+b*x+a)/x,x, algorithm="maxima")`

output `-integrate(sin(c*x^2 - b*x - a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)}{x} dx$$

input `integrate(sin(-c*x^2+b*x+a)/x,x, algorithm="giac")`

output `integrate(sin(-c*x^2 + b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 38.62 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)}{x} dx$$

input `int(sin(a + b*x - c*x^2)/x,x)`

output `int(sin(a + b*x - c*x^2)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\sin(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)}{x} dx$$

input `int(sin(-c*x^2+b*x+a)/x,x)`

output `int(sin(a + b*x - c*x**2)/x,x)`

3.10 $\int \left(-\frac{b \cos(a+bx-cx^2)}{x} + \frac{\sin(a+bx-cx^2)}{x^2} \right) dx$

Optimal result	101
Mathematica [A] (verified)	101
Rubi [A] (verified)	102
Maple [F]	103
Fricas [A] (verification not implemented)	103
Sympy [F]	104
Maxima [F]	104
Giac [F]	104
Mupad [F(-1)]	105
Reduce [F]	105

Optimal result

Integrand size = 35, antiderivative size = 110

$$\begin{aligned} & \int \left(-\frac{b \cos(a+bx-cx^2)}{x} + \frac{\sin(a+bx-cx^2)}{x^2} \right) dx \\ &= \sqrt{c}\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &+ \sqrt{c}\sqrt{2\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) - \frac{\sin(a+bx-cx^2)}{x} \end{aligned}$$

output

```
c^(1/2)*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelC(1/2*(-2*c*x+b)/c^(1/2)*
2^(1/2)/Pi^(1/2))+c^(1/2)*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(-2*c*x+b)/c^(1/2)
*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)-sin(-c*x^2+b*x+a)/x
```

Mathematica [A] (verified)

Time = 0.84 (sec), antiderivative size = 115, normalized size of antiderivative = 1.05

$$\begin{aligned} & \int \left(-\frac{b \cos(a+bx-cx^2)}{x} + \frac{\sin(a+bx-cx^2)}{x^2} \right) dx = \\ & \frac{-\sqrt{c}\sqrt{2\pi}x \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \sqrt{c}\sqrt{2\pi}x \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) + \sin(a+x(b- \right. \end{aligned}$$

input $\text{Integrate}[-((b \cos[a + b x - c x^2])/x) + \sin[a + b x - c x^2]/x^2, x]$

output $-\left(\frac{\sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}] x \cos[a + b^2/(4 c)] \operatorname{FresnelC}\left[\frac{(-b + 2 c x)}{\sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}]}\right] + \sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}] x \operatorname{FresnelS}\left[\frac{(-b + 2 c x)}{\sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}]}\right] \sin[a + b^2/(4 c)] + \sin[a + x (b - c x)]}{x}\right)$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.029$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \left(\frac{\sin(a + bx - cx^2)}{x^2} - \frac{b \cos(a + bx - cx^2)}{x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \sqrt{2\pi} \sqrt{c} \cos\left(a + \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b - 2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \sqrt{2\pi} \sqrt{c} \sin\left(a + \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b - 2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \\ & \quad \frac{\sin(a + bx - cx^2)}{x} \end{aligned}$$

input $\text{Int}[-((b \cos[a + b x - c x^2])/x) + \sin[a + b x - c x^2]/x^2, x]$

output $\sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}] \cos[a + b^2/(4 c)] \operatorname{FresnelC}\left[\frac{(b - 2 c x)}{\sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}]}\right] + \sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}] \operatorname{FresnelS}\left[\frac{(b - 2 c x)}{\sqrt{c} \operatorname{Sqrt}[2 \operatorname{Pi}]}\right] \sin[a + b^2/(4 c)] - \sin[a + b x - c x^2]/x$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

Maple [F]

$$\int \left(-\frac{b \cos(-cx^2 + bx + a)}{x} + \frac{\sin(-cx^2 + bx + a)}{x^2} \right) dx$$

input $\text{int}(-b*\cos(-c*x^2+b*x+a)/x+\sin(-c*x^2+b*x+a)/x^2,x)$

output $\text{int}(-b*\cos(-c*x^2+b*x+a)/x+\sin(-c*x^2+b*x+a)/x^2,x)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 124, normalized size of antiderivative = 1.13

$$\int \left(-\frac{b \cos(a + bx - cx^2)}{x} + \frac{\sin(a + bx - cx^2)}{x^2} \right) dx = \\ -\frac{\sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) - \sin(cx^2 - bx - a)}{x}$$

input $\text{integrate}(-b*\cos(-c*x^2+b*x+a)/x+\sin(-c*x^2+b*x+a)/x^2,x, \text{algorithm}=\text{"fricas"})$

output $-(\sqrt{2}*\pi*x*\sqrt{c/\pi})*\cos(1/4*(b^2 + 4*a*c)/c)*\text{fresnel_cos}(1/2*\sqrt{2}*(2*c*x - b)*\sqrt{c/\pi}/c) + \sqrt{2}*\pi*x*\sqrt{c/\pi}*\text{fresnel_sin}(1/2*\sqrt{2}*(2*c*x - b)*\sqrt{c/\pi}/c)*\sin(1/4*(b^2 + 4*a*c)/c) - \sin(c*x^2 - b*x - a))/x$

Sympy [F]

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx - cx^2)}{x} + \frac{\sin(a + bx - cx^2)}{x^2} \right) dx \\ &= - \int \left(-\frac{\sin(a + bx - cx^2)}{x^2} \right) dx - \int \frac{b \cos(a + bx - cx^2)}{x} dx \end{aligned}$$

input `integrate(-b*cos(-c*x**2+b*x+a)/x+sin(-c*x**2+b*x+a)/x**2,x)`

output `-Integral(-sin(a + b*x - c*x**2)/x**2, x) - Integral(b*cos(a + b*x - c*x**2)/x, x)`

Maxima [F]

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx - cx^2)}{x} + \frac{\sin(a + bx - cx^2)}{x^2} \right) dx \\ &= \int -\frac{b \cos(-cx^2 + bx + a)}{x} + \frac{\sin(-cx^2 + bx + a)}{x^2} dx \end{aligned}$$

input `integrate(-b*cos(-c*x^2+b*x+a)/x+sin(-c*x^2+b*x+a)/x^2,x, algorithm="maxima")`

output `integrate(-b*cos(c*x^2 - b*x - a)/x - sin(c*x^2 - b*x - a)/x^2, x)`

Giac [F]

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx - cx^2)}{x} + \frac{\sin(a + bx - cx^2)}{x^2} \right) dx \\ &= \int -\frac{b \cos(-cx^2 + bx + a)}{x} + \frac{\sin(-cx^2 + bx + a)}{x^2} dx \end{aligned}$$

input `integrate(-b*cos(-c*x^2+b*x+a)/x+sin(-c*x^2+b*x+a)/x^2,x, algorithm="giac")`

output `integrate(-b*cos(-c*x^2 + b*x + a)/x + sin(-c*x^2 + b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx - cx^2)}{x} + \frac{\sin(a + bx - cx^2)}{x^2} \right) dx \\ &= \int \frac{\sin(-cx^2 + bx + a)}{x^2} - \frac{b \cos(-cx^2 + bx + a)}{x} dx \end{aligned}$$

input `int(sin(a + b*x - c*x^2)/x^2 - (b*cos(a + b*x - c*x^2))/x, x)`

output `int(sin(a + b*x - c*x^2)/x^2 - (b*cos(a + b*x - c*x^2))/x, x)`

Reduce [F]

$$\begin{aligned} & \int \left(-\frac{b \cos(a + bx - cx^2)}{x} + \frac{\sin(a + bx - cx^2)}{x^2} \right) dx \\ &= \frac{-2(\int \cos(-cx^2 + bx + a) dx) bcx - \sin(-cx^2 + bx + a) b + 2acx}{bx} \end{aligned}$$

input `int(-b*cos(-c*x^2+b*x+a)/x+sin(-c*x^2+b*x+a)/x^2,x)`

output `(- 2*int(cos(a + b*x - c*x**2),x)*b*c*x - sin(a + b*x - c*x**2)*b + 2*a*c*x)/(b*x)`

3.11 $\int x^2 \sin \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	106
Mathematica [A] (verified)	106
Rubi [A] (verified)	107
Maple [A] (verified)	109
Fricas [A] (verification not implemented)	110
Sympy [F]	110
Maxima [C] (verification not implemented)	110
Giac [C] (verification not implemented)	111
Mupad [B] (verification not implemented)	112
Reduce [F]	112

Optimal result

Integrand size = 13, antiderivative size = 82

$$\begin{aligned} \int x^2 \sin \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{1}{4} \cos \left(\frac{1}{4} + x + x^2 \right) - \frac{1}{2} x \cos \left(\frac{1}{4} + x + x^2 \right) \\ & + \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right) \end{aligned}$$

output
$$1/4*\cos(1/4+x+x^2)-1/2*x*\cos(1/4+x+x^2)+1/4*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(1+2*x)*2^(1/2)/Pi^(1/2))+1/8*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(1+2*x)*2^(1/2)/Pi^(1/2))$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.80

$$\begin{aligned} \int x^2 \sin \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{1}{8} \left(2(1-2x) \cos \left(\frac{1}{4} + x + x^2 \right) + 2\sqrt{2\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) \right. \\ & \left. + \sqrt{2\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right) \right) \end{aligned}$$

input `Integrate[x^2*Sin[1/4 + x + x^2],x]`

output
$$(2*(1 - 2*x)*Cos[1/4 + x + x^2] + 2*.Sqrt[2*Pi]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]] + Sqrt[2*Pi]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]])/8$$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3944, 3927, 3833, 3942, 3926, 3832}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \sin \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \textcolor{blue}{3944} \\
 & -\frac{1}{2} \int x \sin \left(x^2 + x + \frac{1}{4} \right) dx + \frac{1}{2} \int \cos \left(x^2 + x + \frac{1}{4} \right) dx - \frac{1}{2} x \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3927} \\
 & -\frac{1}{2} \int x \sin \left(x^2 + x + \frac{1}{4} \right) dx + \frac{1}{2} \int \cos \left(\frac{1}{4}(2x+1)^2 \right) dx - \frac{1}{2} x \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3833} \\
 & -\frac{1}{2} \int x \sin \left(x^2 + x + \frac{1}{4} \right) dx + \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right) - \frac{1}{2} x \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3942} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \sin \left(x^2 + x + \frac{1}{4} \right) dx + \frac{1}{2} \cos \left(x^2 + x + \frac{1}{4} \right) \right) + \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right) - \\
 & \quad \frac{1}{2} x \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3926} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \sin \left(\frac{1}{4}(2x+1)^2 \right) dx + \frac{1}{2} \cos \left(x^2 + x + \frac{1}{4} \right) \right) + \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right) - \\
 & \quad \frac{1}{2} x \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3832}
 \end{aligned}$$

$$\frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{2x+1}{\sqrt{2\pi}}\right) + \frac{1}{2} \left(\frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{2x+1}{\sqrt{2\pi}}\right) + \frac{1}{2} \cos\left(x^2 + x + \frac{1}{4}\right) \right) - \frac{1}{2} x \cos\left(x^2 + x + \frac{1}{4}\right)$$

input `Int[x^2*Sin[1/4 + x + x^2], x]`

output `-1/2*(x*Cos[1/4 + x + x^2]) + (Sqrt[Pi/2]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]])/2 + (Cos[1/4 + x + x^2]/2 + (Sqrt[Pi/2]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]])/2)/2`

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_))^2, x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_))^2, x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3926 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2, x_Symbol] :> Int[Sin[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

rule 3927 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2, x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

rule 3942 `Int[((d_.) + (e_.)*(x_))*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2, x_Symbol] :> Simp[(-e)*(Cos[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c)*Int[Sin[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]`

rule 3944

```
Int[((d_.) + (e_)*(x_))^(m_)*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2], x_Sym
bol] :> Simp[(-e)*(d + e*x)^(m - 1)*(Cos[a + b*x + c*x^2]/(2*c)), x] + (-Si
mp[(b*e - 2*c*d)/(2*c) Int[(d + e*x)^(m - 1)*Sin[a + b*x + c*x^2], x], x]
+ Simp[e^2*((m - 1)/(2*c)) Int[(d + e*x)^(m - 2)*Cos[a + b*x + c*x^2], x
], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Maple [A] (verified)

Time = 1.36 (sec), antiderivative size = 59, normalized size of antiderivative = 0.72

method	result
default	$-\frac{x \cos(\frac{1}{4}+x+x^2)}{2} + \frac{\cos(\frac{1}{4}+x+x^2)}{4} + \frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{8} + \frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{4}$
risch	$-\frac{i \sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{16 \sqrt{-i}} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{8 \sqrt{-i}} + \frac{(-1)^{\frac{1}{4}} \sqrt{\pi} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{16} - \frac{\sqrt{\pi} (-1)^{\frac{3}{4}} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{8}$
parts	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right) x^2}{2} - \frac{\sqrt{2} \pi^{\frac{3}{2}} \left(\frac{\operatorname{FresnelS}\left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right) \left(\left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right)^2 \sqrt{\pi}-\sqrt{2} \left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right)\right)}{\sqrt{\pi}} - \frac{\left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right) \cos\left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right)}{\sqrt{\pi}} \right)}{4}$

input `int(x^2*sin(1/4+x+x^2),x,method=_RETURNVERBOSE)`

output
$$-1/2*x*cos(1/4+x+x^2)+1/4*cos(1/4+x+x^2)+1/8*2^(1/2)*Pi^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)*(x+1/2))+1/4*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.72

$$\int x^2 \sin\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{4} (2x - 1) \cos\left(x^2 + x + \frac{1}{4}\right) + \frac{1}{4} \sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x + 1)}{2\sqrt{\pi}}\right) + \frac{1}{8} \sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}(2x + 1)}{2\sqrt{\pi}}\right)$$

input `integrate(x^2*sin(1/4+x+x^2),x, algorithm="fricas")`

output `-1/4*(2*x - 1)*cos(x^2 + x + 1/4) + 1/4*sqrt(2)*sqrt(pi)*fresnel_cos(1/2*sqrt(2)*(2*x + 1)/sqrt(pi)) + 1/8*sqrt(2)*sqrt(pi)*fresnel_sin(1/2*sqrt(2)*(2*x + 1)/sqrt(pi))`

Sympy [F]

$$\int x^2 \sin\left(\frac{1}{4} + x + x^2\right) dx = \int x^2 \sin\left(x^2 + x + \frac{1}{4}\right) dx$$

input `integrate(x**2*sin(1/4+x+x**2),x)`

output `Integral(x**2*sin(x**2 + x + 1/4), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.23 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.91

$$\int x^2 \sin\left(\frac{1}{4} + x + x^2\right) dx = \frac{16x \left(e^{(ix^2+ix+\frac{1}{4}i)} + e^{(-ix^2-ix-\frac{1}{4}i)}\right) - \sqrt{4x^2 + 4x + 1} \left(-(i+1) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\sqrt{ix^2 + ix + \frac{1}{4}i}\right) - 1\right)}{}$$

input `integrate(x^2*sin(1/4+x+x^2),x, algorithm="maxima")`

output
$$\begin{aligned} & \frac{1}{32} \left(16x \left(e^{(I*x^2 + I*x + 1/4*I)} + e^{(-I*x^2 - I*x - 1/4*I)} \right) - \sqrt{4*x^2 + 4*x + 1} \right) * (-I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{I*x^2 + I*x + 1/4*I}) - 1) \\ & + (I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{-I*x^2 - I*x - 1/4*I}) - 1) - (4*I - 4) * \sqrt{2} * \operatorname{gamma}(3/2, I*x^2 + I*x + 1/4*I) + (4*I + 4) * \sqrt{2} * \operatorname{gamma}(3/2, -I*x^2 - I*x - 1/4*I) + 8 * e^{(I*x^2 + I*x + 1/4*I)} + 8 * e^{(-I*x^2 - I*x - 1/4*I)} \end{aligned} / (2*x + 1)$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.91

$$\begin{aligned} \int x^2 \sin \left(\frac{1}{4} + x + x^2 \right) dx = & - \left(\frac{1}{32}i + \frac{3}{32} \right) \sqrt{2}\sqrt{\pi} \operatorname{erf} \left(\left(\frac{1}{4}i - \frac{1}{4} \right) \sqrt{2}(2x + 1) \right) \\ & + \left(\frac{1}{32}i - \frac{3}{32} \right) \sqrt{2}\sqrt{\pi} \operatorname{erf} \left(-\left(\frac{1}{4}i + \frac{1}{4} \right) \sqrt{2}(2x + 1) \right) \\ & - \frac{1}{8}i (-2ix + i)e^{(ix^2 + ix + \frac{1}{4}i)} - \frac{1}{8}i (-2ix + i)e^{(-ix^2 - ix - \frac{1}{4}i)} \end{aligned}$$

input `integrate(x^2*sin(1/4+x+x^2),x, algorithm="giac")`

output
$$\begin{aligned} & -(1/32*I + 3/32)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}((1/4*I - 1/4)*\sqrt{2}*(2*x + 1)) + (1/32*I - 3/32)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-(1/4*I + 1/4)*\sqrt{2}*(2*x + 1)) - 1/8*I*(-2*I*x + I)*e^{(I*x^2 + I*x + 1/4*I)} - 1/8*I*(-2*I*x + I)*e^{(-I*x^2 - I*x - 1/4*I)} \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.78

$$\int x^2 \sin \left(\frac{1}{4} + x + x^2 \right) dx = \frac{\cos \left(x^2 + x + \frac{1}{4} \right)}{4} - \frac{x \cos \left(x^2 + x + \frac{1}{4} \right)}{2} \\ + \frac{\sqrt{2} \sqrt{\pi} C \left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}} \right)}{4} + \frac{\sqrt{2} \sqrt{\pi} S \left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}} \right)}{8}$$

input `int(x^2*sin(x + x^2 + 1/4),x)`

output $\cos(x + x^2 + 1/4)/4 - (x*\cos(x + x^2 + 1/4))/2 + (2^{(1/2)}*\pi^{(1/2)}*fresne1c((2^{(1/2)}*(2*x + 1))/(2*pi^{(1/2)}))/4 + (2^{(1/2)}*\pi^{(1/2)}*fresnels((2^{(1/2)}*(2*x + 1))/(2*pi^{(1/2)}))/8$

Reduce [F]

$$\int x^2 \sin \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{\cos \left(x^2 + x + \frac{1}{4} \right) x}{2} + \frac{\cos \left(x^2 + x + \frac{1}{4} \right)}{4} \\ - \left(\int \frac{x^2}{\tan \left(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8} \right)^2 + 1} dx \right) \\ + \frac{5 \left(\int \frac{1}{\tan \left(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8} \right)^2 + 1} dx \right)}{4} + \frac{\sin \left(x^2 + x + \frac{1}{4} \right) x}{4} \\ - \frac{\sin \left(x^2 + x + \frac{1}{4} \right)}{8} + \frac{x^3}{6} - \frac{5x}{8} - \frac{1}{4}$$

input `int(x^2*sin(1/4+x+x^2),x)`

output $(- 12*\cos((4*x**2 + 4*x + 1)/4)*x + 6*\cos((4*x**2 + 4*x + 1)/4) - 24*int(x**2/(\tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) + 30*int(1/(\tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) + 6*\sin((4*x**2 + 4*x + 1)/4)*x - 3*\sin((4*x**2 + 4*x + 1)/4) + 4*x**3 - 15*x - 6)/24$

3.12 $\int x \sin \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	113
Mathematica [A] (verified)	113
Rubi [A] (verified)	114
Maple [A] (verified)	115
Fricas [A] (verification not implemented)	115
Sympy [B] (verification not implemented)	116
Maxima [C] (verification not implemented)	116
Giac [C] (verification not implemented)	117
Mupad [B] (verification not implemented)	117
Reduce [F]	118

Optimal result

Integrand size = 11, antiderivative size = 41

$$\int x \sin \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{1}{2} \cos \left(\frac{1}{4} + x + x^2 \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right)$$

output
$$-1/2*\cos(1/4+x+x^2)-1/4*2^{(1/2)}*\text{Pi}^{(1/2)}*\text{FresnelS}(1/2*(1+2*x)*2^{(1/2)}/\text{Pi}^{(1/2)})$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int x \sin \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \left(-2 \cos \left(\frac{1}{4} + x + x^2 \right) - \sqrt{2\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right) \right)$$

input
$$\text{Integrate}[x \sin[1/4 + x + x^2], x]$$

output
$$(-2*\text{Cos}[1/4 + x + x^2] - \text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[(1 + 2*x)/\text{Sqrt}[2*\text{Pi}]])/4$$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3942, 3926, 3832}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \textcolor{blue}{3942} \\
 & -\frac{1}{2} \int \sin \left(x^2 + x + \frac{1}{4} \right) dx - \frac{1}{2} \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3926} \\
 & -\frac{1}{2} \int \sin \left(\frac{1}{4}(2x+1)^2 \right) dx - \frac{1}{2} \cos \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3832} \\
 & -\frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) - \frac{1}{2} \cos \left(x^2 + x + \frac{1}{4} \right)
 \end{aligned}$$

input `Int[x*Sin[1/4 + x + x^2], x]`

output `-1/2*Cos[1/4 + x + x^2] - (Sqrt[Pi/2]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]])/2`

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3926 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Int[Sin[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

rule 3942

```

Int[((d_.) + (e_.)*(x_))*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol]
:> Simp[(-e)*(Cos[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c)
Int[Sin[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d
- b*e, 0]

```

Maple [A] (verified)

Time = 1.00 (sec), antiderivative size = 30, normalized size of antiderivative = 0.73

method	result	size
default	$-\frac{\cos\left(\frac{1}{4}+x+x^2\right)}{2}-\frac{\sqrt{2}\sqrt{\pi}\operatorname{FresnelS}\left(\frac{\sqrt{2}\left(x+\frac{1}{2}\right)}{\sqrt{\pi}}\right)}{4}$	30
risch	$\frac{i\sqrt{\pi}\operatorname{erf}\left(\sqrt{-i}x-\frac{i}{2\sqrt{-i}}\right)}{8\sqrt{-i}}-\frac{(-1)^{\frac{1}{4}}\sqrt{\pi}\operatorname{erf}\left((-1)^{\frac{1}{4}}x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{8}-\frac{\cos\left(\frac{(1+2x)^2}{4}\right)}{2}$	59
parts	$\frac{\sqrt{2}\sqrt{\pi}\operatorname{FresnelS}\left(\frac{\sqrt{2}\left(x+\frac{1}{2}\right)}{\sqrt{\pi}}\right)x}{2}-\frac{\pi\left(\operatorname{FresnelS}\left(\frac{\sqrt{2}x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2\sqrt{\pi}}\right)\left(\frac{\sqrt{2}x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2\sqrt{\pi}}\right)+\frac{\cos\left(\frac{\pi\left(\frac{\sqrt{2}x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2\sqrt{\pi}}\right)^2}{2}\right)}{\pi}\right)}{2}$	89

input `int(x*sin(1/4+x+x^2), x, method=_RETURNVERBOSE)`output `-1/2*cos(1/4+x+x^2)-1/4*2^(1/2)*Pi^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)*(x+1/2))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec), antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int x \sin\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{4} \sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right) - \frac{1}{2} \cos\left(x^2 + x + \frac{1}{4}\right)$$

input `integrate(x*sin(1/4+x+x^2), x, algorithm="fricas")`

output
$$-1/4\sqrt{2}\sqrt{\pi}\text{fresnel_sin}(1/2\sqrt{2}*(2x + 1)/\sqrt{\pi}) - 1/2\cos(x^2 + x + 1/4)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(41) = 82$.

Time = 0.64 (sec) , antiderivative size = 160, normalized size of antiderivative = 3.90

$$\int x \sin\left(\frac{1}{4} + x + x^2\right) dx = -\frac{3\sqrt{2}\sqrt{\pi}xS\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)\Gamma\left(\frac{3}{4}\right)}{8\Gamma\left(\frac{7}{4}\right)} + \frac{\sqrt{2}\sqrt{\pi}xS\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)}{2} - \frac{3\cos\left((x + \frac{1}{2})^2\right)\Gamma\left(\frac{3}{4}\right)}{8\Gamma\left(\frac{7}{4}\right)} - \frac{3\sqrt{2}\sqrt{\pi}S\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)\Gamma\left(\frac{3}{4}\right)}{16\Gamma\left(\frac{7}{4}\right)}$$

input `integrate(x*sin(1/4+x+x**2),x)`

output
$$-\frac{3\sqrt{2}\sqrt{\pi}x\text{fresnels}(\sqrt{2}x/\sqrt{\pi} + \sqrt{2}/(2\sqrt{\pi}))\gamma(3/4)/(8\gamma(7/4)) + \sqrt{2}\sqrt{\pi}x\text{fresnels}(\sqrt{2}x/\sqrt{\pi})\gamma(3/4)/(8\gamma(7/4)) + \sqrt{2}/(2\sqrt{\pi})}{8\gamma(7/4)} - \frac{3\cos((x + 1/2)^2)\gamma(3/4)/(8\gamma(7/4))}{8\gamma(7/4)} - \frac{3\sqrt{2}\sqrt{\pi}\text{fresnels}(\sqrt{2}x/\sqrt{\pi} + \sqrt{2}/(2\sqrt{\pi}))\gamma(3/4)/(16\gamma(7/4))}{8\gamma(7/4)}$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.98

$$\int x \sin\left(\frac{1}{4} + x + x^2\right) dx = -\frac{8x\left(e^{(ix^2+ix+\frac{1}{4}i)} + e^{(-ix^2-ix-\frac{1}{4}i)}\right) + \sqrt{4x^2 + 4x + 1}\left((i + 1)\sqrt{2}\sqrt{\pi}\left(\text{erf}\left(\sqrt{ix^2 + ix + \frac{1}{4}i}\right) - 1\right) - (i + 1)\sqrt{2}\sqrt{\pi}\left(\text{erf}\left(\sqrt{ix^2 + ix + \frac{1}{4}i}\right) + 1\right)\right)}{16(2x + 1)}$$

input `integrate(x*sin(1/4+x+x^2),x, algorithm="maxima")`

output

$$\begin{aligned} & -\frac{1}{16} \cdot (8x \cdot (e^{(I*x^2 + I*x + 1/4*I)} + e^{-(-I*x^2 - I*x - 1/4*I)}) + \sqrt{4*x^2 + 4*x + 1}) \cdot ((I + 1) \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot (\operatorname{erf}(\sqrt{I*x^2 + I*x + 1/4*I}) - 1) - (I - 1) \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot (\operatorname{erf}(\sqrt{-I*x^2 - I*x - 1/4*I}) - 1)) + 4 \cdot e^{(I*x^2 + I*x + 1/4*I)} + 4 \cdot e^{-(-I*x^2 - I*x - 1/4*I)}) / (2*x + 1) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\begin{aligned} \int x \sin \left(\frac{1}{4} + x + x^2 \right) dx = & - \left(\frac{1}{16}i - \frac{1}{16} \right) \sqrt{2}\sqrt{\pi} \operatorname{erf} \left(\left(\frac{1}{4}i - \frac{1}{4} \right) \sqrt{2}(2x + 1) \right) \\ & + \left(\frac{1}{16}i + \frac{1}{16} \right) \sqrt{2}\sqrt{\pi} \operatorname{erf} \left(- \left(\frac{1}{4}i + \frac{1}{4} \right) \sqrt{2}(2x + 1) \right) \\ & - \frac{1}{4} e^{(ix^2 + ix + \frac{1}{4}i)} - \frac{1}{4} e^{(-ix^2 - ix - \frac{1}{4}i)} \end{aligned}$$

input

```
integrate(x*sin(1/4+x+x^2),x, algorithm="giac")
```

output

$$\begin{aligned} & -(1/16*I - 1/16)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}((1/4*I - 1/4)*\sqrt{2}*(2*x + 1)) + (1/16*I + 1/16)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-(1/4*I + 1/4)*\sqrt{2}*(2*x + 1)) - 1/4*e^{(I*x^2 + I*x + 1/4*I)} - 1/4*e^{-(-I*x^2 - I*x - 1/4*I)} \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 38.49 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int x \sin \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{\cos(x^2 + x + \frac{1}{4})}{2} - \frac{\sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{4}$$

input

```
int(x*sin(x + x^2 + 1/4),x)
```

output

$$-\cos(x + x^2 + 1/4)/2 - (2^{(1/2)} * \pi^{(1/2)} * \operatorname{fresnels}((2^{(1/2)} * (2*x + 1)) / (2 * \pi^{(1/2)}))) / 4$$

Reduce [F]

$$\int x \sin\left(\frac{1}{4} + x + x^2\right) dx = \int \sin\left(x^2 + x + \frac{1}{4}\right) x dx$$

input `int(x*sin(1/4+x+x^2),x)`

output `int(sin((4*x**2 + 4*x + 1)/4)*x,x)`

3.13 $\int \sin\left(\frac{1}{4} + x + x^2\right) dx$

Optimal result	119
Mathematica [A] (verified)	119
Rubi [A] (verified)	120
Maple [A] (verified)	121
Fricas [A] (verification not implemented)	121
Sympy [A] (verification not implemented)	121
Maxima [C] (verification not implemented)	122
Giac [C] (verification not implemented)	122
Mupad [B] (verification not implemented)	123
Reduce [F]	123

Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right)$$

output `1/2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(1+2*x)*2^(1/2)/Pi^(1/2))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right)$$

input `Integrate[Sin[1/4 + x + x^2], x]`

output `Sqrt[Pi/2]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3926, 3832}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin \left(x^2 + x + \frac{1}{4} \right) dx \\ & \quad \downarrow \text{3926} \\ & \int \sin \left(\frac{1}{4}(2x+1)^2 \right) dx \\ & \quad \downarrow \text{3832} \\ & \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) \end{aligned}$$

input `Int[Sin[1/4 + x + x^2], x]`

output `Sqrt[Pi/2]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]]`

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_))^2, x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3926 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2, x_Symbol] :> Int[Sin[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2} \left(x+\frac{1}{2}\right)}{\sqrt{\pi}}\right)}{2}$	20
risch	$\frac{(-1)^{\frac{1}{4}} \sqrt{\pi} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{4}-\frac{i \sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{4 \sqrt{-i}}$	47

input `int(sin(1/4+x+x^2),x,method=_RETURNVERBOSE)`

output `1/2*2^(1/2)*Pi^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)*(x+1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{2} \sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)$$

input `integrate(sin(1/4+x+x^2),x, algorithm="fricas")`

output `1/2*sqrt(2)*sqrt(pi)*fresnel_sin(1/2*sqrt(2)*(2*x + 1)/sqrt(pi))`

Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{2}$$

input `integrate(sin(1/4+x+x**2),x)`

output $\sqrt{2}\sqrt{\pi} \text{fresnels}(\sqrt{2}(2x+1)/(2\sqrt{\pi}))/2$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.92

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx \\ = \frac{1}{16} \sqrt{\pi} \left((i+1) \sqrt{2} \operatorname{erf}\left(-\frac{1}{2} (-1)^{\frac{3}{4}} (2ix+i)\right) + (i+1) \sqrt{2} \operatorname{erf}\left(-\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2ix+i)\right) - (i-1) \sqrt{2} \operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2ix+i)\right) - (i-1) \sqrt{2} \operatorname{erf}\left(\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2}(2ix+i)\right) \right)$$

input `integrate(sin(1/4+x+x^2),x, algorithm="maxima")`

output $1/16\sqrt{\pi}((I+1)\sqrt{2}\operatorname{erf}(-1/2*(-1)^{(3/4)}(2Ix+I)) + (I+1)\sqrt{2}\operatorname{erf}(-(1/4I-1/4)\sqrt{2}(2Ix+I)) - (I-1)\sqrt{2}\operatorname{erf}(-(1/4I+1/4)\sqrt{2}(2Ix+I)) + (I-1)\sqrt{2}\operatorname{erf}(1/2*(2Ix+I)/\sqrt{(-I)}))$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.62

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \left(\frac{1}{8}i - \frac{1}{8}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2x+1)\right) \\ - \left(\frac{1}{8}i + \frac{1}{8}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2}(2x+1)\right)$$

input `integrate(sin(1/4+x+x^2),x, algorithm="giac")`

output $(1/8I - 1/8)\sqrt{2}\sqrt{\pi}\operatorname{erf}((1/4I - 1/4)\sqrt{2}(2x+1)) - (1/8I + 1/8)\sqrt{2}\sqrt{\pi}\operatorname{erf}(-(1/4I + 1/4)\sqrt{2}(2x+1))$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{2}$$

input `int(sin(x + x^2 + 1/4),x)`

output `(2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*(x + 1/2))/pi^(1/2)))/2`

Reduce [F]

$$\int \sin\left(\frac{1}{4} + x + x^2\right) dx = \int \sin\left(x^2 + x + \frac{1}{4}\right) dx$$

input `int(sin(1/4+x+x^2),x)`

output `int(sin((4*x**2 + 4*x + 1)/4),x)`

3.14 $\int \frac{\sin\left(\frac{1}{4}+x+x^2\right)}{x} dx$

Optimal result	124
Mathematica [N/A]	124
Rubi [N/A]	125
Maple [N/A]	125
Fricas [N/A]	126
Sympy [N/A]	126
Maxima [N/A]	127
Giac [N/A]	127
Mupad [N/A]	127
Reduce [N/A]	128

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \text{Int}\left(\frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x}, x\right)$$

output `Defer(Int)(sin(1/4+x+x^2)/x,x)`

Mathematica [N/A]

Not integrable

Time = 19.52 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

input `Integrate[Sin[1/4 + x + x^2]/x,x]`

output `Integrate[Sin[1/4 + x + x^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3950}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(x^2 + x + \frac{1}{4})}{x} dx$$

↓ 3950

$$\int \frac{\sin(x^2 + x + \frac{1}{4})}{x} dx$$

input `Int[Sin[1/4 + x + x^2]/x,x]`

output `$Aborted`

Definitions of rubi rules used

rule 3950 `Int[((d_.) + (e_.)*(x_))^(m_.)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_.), x_Symbol] :> Unintegrable[(d + e*x)^m*Sin[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{\sin(\frac{1}{4} + x + x^2)}{x} dx$$

input `int(sin(1/4+x+x^2)/x,x)`

output `int(sin(1/4+x+x^2)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(sin(1/4+x+x^2)/x,x, algorithm="fricas")`

output `integral(sin(x^2 + x + 1/4)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(sin(1/4+x+x**2)/x,x)`

output `Integral(sin(x**2 + x + 1/4)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(sin(1/4+x+x^2)/x,x, algorithm="maxima")`

output `integrate(sin(x^2 + x + 1/4)/x, x)`

Giac [N/A]

Not integrable

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(sin(1/4+x+x^2)/x,x, algorithm="giac")`

output `integrate(sin(x^2 + x + 1/4)/x, x)`

Mupad [N/A]

Not integrable

Time = 38.94 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `int(sin(x + x^2 + 1/4)/x,x)`

output `int(sin(x + x^2 + 1/4)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `int(sin(1/4+x+x^2)/x,x)`

output `int(sin((4*x**2 + 4*x + 1)/4)/x,x)`

3.15 $\int \frac{\sin\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$

Optimal result	129
Mathematica [N/A]	129
Rubi [N/A]	130
Maple [N/A]	131
Fricas [N/A]	132
Sympy [N/A]	132
Maxima [N/A]	132
Giac [N/A]	133
Mupad [N/A]	133
Reduce [N/A]	134

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right) - \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x} \\ + \operatorname{Int}\left(\frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x}, x\right)$$

output $2^{(1/2)}*\text{Pi}^{(1/2)}*\text{FresnelC}(1/2*(1+2*x)*2^{(1/2)}/\text{Pi}^{(1/2)})-\sin(1/4+x+x^2)/x+\text{D}$
 $\text{efer}(\operatorname{Int})(\cos(1/4+x+x^2)/x, x)$

Mathematica [N/A]

Not integrable

Time = 14.38 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

input $\text{Integrate}[\text{Sin}[1/4 + x + x^2]/x^2, x]$

output `Integrate[Sin[1/4 + x + x^2]/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3946, 3927, 3833, 3951}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x^2 + x + \frac{1}{4})}{x^2} dx \\
 & \quad \downarrow \textcolor{blue}{3946} \\
 & 2 \int \cos\left(x^2 + x + \frac{1}{4}\right) dx + \int \frac{\cos(x^2 + x + \frac{1}{4})}{x} dx - \frac{\sin(x^2 + x + \frac{1}{4})}{x} \\
 & \quad \downarrow \textcolor{blue}{3927} \\
 & \int \frac{\cos(x^2 + x + \frac{1}{4})}{x} dx + 2 \int \cos\left(\frac{1}{4}(2x+1)^2\right) dx - \frac{\sin(x^2 + x + \frac{1}{4})}{x} \\
 & \quad \downarrow \textcolor{blue}{3833} \\
 & \int \frac{\cos(x^2 + x + \frac{1}{4})}{x} dx + \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{2x+1}{\sqrt{2\pi}}\right) - \frac{\sin(x^2 + x + \frac{1}{4})}{x} \\
 & \quad \downarrow \textcolor{blue}{3951} \\
 & \int \frac{\cos(x^2 + x + \frac{1}{4})}{x} dx + \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{2x+1}{\sqrt{2\pi}}\right) - \frac{\sin(x^2 + x + \frac{1}{4})}{x}
 \end{aligned}$$

input `Int[Sin[1/4 + x + x^2]/x^2,x]`

output `$Aborted`

Definitions of rubi rules used

rule 3833 $\text{Int}[\cos[(d_{_}) + (e_{_})x_{_}]^2], \text{x_Symbol}] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f\sqrt{d}), \text{x}] \text{FresnelC}[\sqrt{2/\pi}\sqrt{d} \cos[e + f x], \text{x}] /; \text{FreeQ}[d, e, f], \text{x}]$

rule 3927 $\text{Int}[\cos[(a_{_}) + (b_{_})x_{_}] + (c_{_})x_{_}^2], \text{x_Symbol}] \rightarrow \text{Int}[\cos[(b + 2cx)^2/(4c)], \text{x}] /; \text{FreeQ}[a, b, c], \text{x} \&& \text{EqQ}[b^2 - 4a*c, 0]$

rule 3946 $\text{Int}[(d_{_}) + (e_{_})x_{_}]^{(m)} \sin[(a_{_}) + (b_{_})x_{_}] + (c_{_})x_{_}^2], \text{x_Symbol}] \rightarrow \text{Simp}[(d + e x)^{(m+1)} (\sin[a + b x + c x^2]/(e(m+1))), \text{x}] + (-\text{Simp}[(b e - 2 c d)/(e^{2(m+1)})] \text{Int}[(d + e x)^{(m+1)} \cos[a + b x + c x^2], \text{x}] - \text{Simp}[2(c/(e^{2(m+1)})] \text{Int}[(d + e x)^{(m+2)} \cos[a + b x + c x^2], \text{x}]) /; \text{FreeQ}[a, b, c, d, e], \text{x} \&& \text{NeQ}[b e - 2 c d, 0] \&& \text{LtQ}[m, -1]$

rule 3951 $\text{Int}[\cos[(a_{_}) + (b_{_})x_{_}] + (c_{_})x_{_}^2]^{(n)} ((d_{_}) + (e_{_})x_{_})^{(m)}, \text{x_Symbol}] \rightarrow \text{Unintegrable}[(d + e x)^m \cos[a + b x + c x^2]^n, \text{x}] /; \text{FreeQ}[a, b, c, d, e, m, n], \text{x}]$

Maple [N/A]

Not integrable

Time = 0.15 (sec), antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{\sin(\frac{1}{4} + x + x^2)}{x^2} dx$$

input `int(sin(1/4+x+x^2)/x^2,x)`

output `int(sin(1/4+x+x^2)/x^2,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(sin(1/4+x+x^2)/x^2,x, algorithm="fricas")`

output `integral(sin(x^2 + x + 1/4)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(sin(1/4+x+x**2)/x**2,x)`

output `Integral(sin(x**2 + x + 1/4)/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(sin(1/4+x+x^2)/x^2,x, algorithm="maxima")`

output `integrate(sin(x^2 + x + 1/4)/x^2, x)`

Giac [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(sin(1/4+x+x^2)/x^2,x, algorithm="giac")`

output `integrate(sin(x^2 + x + 1/4)/x^2, x)`

Mupad [N/A]

Not integrable

Time = 38.84 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `int(sin(x + x^2 + 1/4)/x^2,x)`

output `int(sin(x + x^2 + 1/4)/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 74, normalized size of antiderivative = 5.69

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x^2} dx \\ = \frac{8 \left(\int \frac{1}{\tan\left(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8}\right)^2 + 1} dx \right) x + 4 \left(\int \frac{1}{\tan\left(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8}\right)^2 x + x} dx \right) x - 2 \log(x) x - 2 \sin\left(x^2 + x + \frac{1}{4}\right) - 4x^2 - x}{2x}$$

input `int(sin(1/4+x+x^2)/x^2, x)`

output `(8*int(1/(tan((4*x**2 + 4*x + 1)/8)**2 + 1), x)*x + 4*int(1/(tan((4*x**2 + 4*x + 1)/8)**2*x + x), x)*x - 2*log(x)*x - 2*sin((4*x**2 + 4*x + 1)/4) - 4*x**2 - x)/(2*x)`

3.16 $\int x^2 \sin^2(a + bx + cx^2) dx$

Optimal result	135
Mathematica [A] (verified)	136
Rubi [A] (verified)	136
Maple [A] (verified)	137
Fricas [A] (verification not implemented)	138
Sympy [F]	138
Maxima [C] (verification not implemented)	139
Giac [C] (verification not implemented)	140
Mupad [F(-1)]	140
Reduce [F]	141

Optimal result

Integrand size = 17, antiderivative size = 248

$$\begin{aligned} \int x^2 \sin^2(a + bx + cx^2) dx = & \frac{x^3}{6} - \frac{b^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & + \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & + \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\ & + \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\ & + \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output

```
1/6*x^3-1/16*b^2*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(5/2)+1/16*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)+1/16*Pi^(1/2)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)+1/16*b^2*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(5/2)+1/16*b*sin(2*c*x^2+2*b*x+2*a)/c^2-1/8*x*sin(2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.69

$$\int x^2 \sin^2(a + bx + cx^2) dx \\ = \frac{3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a - \frac{b^2}{2c}\right) + b^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) - 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a - \frac{b^2}{2c}\right)\right)}{48c^{5/2}}$$

input `Integrate[x^2*Sin[a + b*x + c*x^2]^2, x]`

output
$$(3\sqrt{\pi} \operatorname{FresnelS}\left[\frac{(b + 2cx)}{\sqrt{c}\sqrt{\pi}}\right] * (c \cos[2a - b^2/(2c)] + b^2 \sin[2a - b^2/(2c)]) - 3\sqrt{\pi} \operatorname{FresnelC}\left[\frac{(b + 2cx)}{\sqrt{c}\sqrt{\pi}}\right] * (b^2 \cos[2a - b^2/(2c)] - c \sin[2a - b^2/(2c)]) + \sqrt{c} * (8c^2 x^3 + 3(b - 2cx) \sin[2(a + x(b + cx))])) / (48c^{5/2})$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sin^2(a + bx + cx^2) dx \\ \downarrow 3948 \\ \int \left(\frac{x^2}{2} - \frac{1}{2}x^2 \cos(2a + 2bx + 2cx^2)\right) dx \\ \downarrow 2009$$

$$\begin{aligned} & \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \frac{\sqrt{\pi}b^2 \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \\ & \frac{\sqrt{\pi}b^2 \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} + \\ & \frac{b \sin\left(2a + 2bx + 2cx^2\right)}{16c^2} - \frac{x \sin\left(2a + 2bx + 2cx^2\right)}{8c} + \frac{x^3}{6} \end{aligned}$$

input `Int[x^2*Sin[a + b*x + c*x^2]^2, x]`

output $x^{3/6} - (b^{2*}\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a - b^{2/(2*c)}]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])])/(16*c^{(5/2)}) + (\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a - b^{2/(2*c)}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])])/(16*c^{(3/2)}) + (\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*\text{Sin}[2*a - b^{2/(2*c)}])/(16*c^{(3/2)}) + (b^{2*}\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*\text{Sin}[2*a - b^{2/(2*c)}])/(16*c^{(5/2)}) + (b*\text{Sin}[2*a + 2*b*x + 2*c*x^2])/(16*c^2) - (x*\text{Sin}[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_)*(x_))^(m_)*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.40 (sec), antiderivative size = 191, normalized size of antiderivative = 0.77

method	result
default	$\frac{x^3}{6} - \frac{x \sin(2cx^2 + 2bx + 2a)}{8c} + \frac{b \left(\frac{\sin(2cx^2 + 2bx + 2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{4c^{3/2}} \right)}{4c}$
risch	$-\frac{b^2\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{2c}}\sqrt{2}\text{erf}\left(\sqrt{2}\sqrt{ic}x+\frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c^2\sqrt{ic}} + \frac{i\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{2c}}\sqrt{2}\text{erf}\left(\sqrt{2}\sqrt{ic}x+\frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c\sqrt{ic}} + \frac{b^2\sqrt{\pi}e^{\frac{i(4ac-b^2)}{2c}}\text{erf}\left(-\sqrt{-2ic}x-\frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c^2\sqrt{-2ic}}$

input `int(x^2*sin(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1/6*x^3 - 1/8*x*sin(2*c*x^2 + 2*b*x + 2*a)/c + 1/4*b/c*(1/4*sin(2*c*x^2 + 2*b*x + 2*a)/c - 1/4*b/c^{(3/2)}*Pi^{(1/2)}*(cos(1/2*(-4*a*c + b^2)/c)*FresnelC((2*c*x + b)/c^{(1/2)}/Pi^{(1/2)}) + sin(1/2*(-4*a*c + b^2)/c)*FresnelS((2*c*x + b)/c^{(1/2)}/Pi^{(1/2)})) + 1/16/c^{(3/2)}*Pi^{(1/2)}*(cos(1/2*(-4*a*c + b^2)/c)*FresnelS((2*c*x + b)/c^{(1/2)}/Pi^{(1/2)}) - sin(1/2*(-4*a*c + b^2)/c)*FresnelC((2*c*x + b)/c^{(1/2)}/Pi^{(1/2)}))}{48*c^3}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 178, normalized size of antiderivative = 0.72

$$\int x^2 \sin^2(a + bx + cx^2) dx$$

$$= \frac{8 c^3 x^3 - 6 (2 c^2 x - bc) \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) - 3 \left(\pi b^2 \cos\left(-\frac{b^2 - 4ac}{2c}\right) - \pi c \sin\left(-\frac{b^2 - 4ac}{2c}\right)\right)}{48 c^3}$$

input `integrate(x^2*sin(c*x^2+b*x+a)^2,x, algorithm="fricas")`

output
$$\frac{1/48*(8*c^3*x^3 - 6*(2*c^2*x - b*c)*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a) - 3*(pi*b^2*cos(-1/2*(b^2 - 4*a*c)/c) - pi*c*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) + 3*(pi*b^2*sin(-1/2*(b^2 - 4*a*c)/c) + pi*c*cos(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c))/c^3}{48*c^3}$$

Sympy [F]

$$\int x^2 \sin^2(a + bx + cx^2) dx = \int x^2 \sin^2(a + bx + cx^2) dx$$

input `integrate(x**2*sin(c*x**2+b*x+a)**2,x)`

output `Integral(x**2*sin(a + b*x + c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.81 (sec) , antiderivative size = 1617, normalized size of antiderivative = 6.52

$$\int x^2 \sin^2(a + bx + cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*sin(c*x^2+b*x+a)^2,x, algorithm="maxima")`

output

```
-1/384*sqrt(2)*(24*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*((I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*cos(-1/2*(b^2 - 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*sin(-1/2*(b^2 - 4*a*c)/c))*x^3 + 36*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(-1/2*(b^2 - 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*(-(I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b...
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.86

$$\int x^2 \sin^2(a + bx + cx^2) dx = \frac{1}{6} x^3 - \frac{(c(-2i x - \frac{ib}{c}) + 2i b)e^{(2icx^2+2ibx+2ia)} - \frac{\sqrt{\pi}(b^2+i c) \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{-ib^2+4ic}{2c}\right)}}{\sqrt{c}\left(-\frac{ic}{|c|}+1\right)}}{32c^2} - \frac{(c(2ix + \frac{ib}{c}) - 2i b)e^{(-2icx^2-2ibx-2ia)} - \frac{\sqrt{\pi}(b^2-i c) \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{-ib^2+4ic}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|}+1\right)}}{32c^2}$$

input `integrate(x^2*sin(c*x^2+b*x+a)^2,x, algorithm="giac")`

output $\frac{1}{6}x^3 - \frac{1}{32}((c*(-2*I*x - I*b/c) + 2*I*b)*e^{(2*I*c*x^2 + 2*I*b*x + 2*I*a)} - \sqrt{\pi}*(b^2 + I*c)*\operatorname{erf}(-1/2*\sqrt{c}*(2*x + b/c)*(-I*c/abs(c) + 1))*e^{(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1))}/c^2 - \frac{1}{32}((c*(2*I*x + I*b/c) - 2*I*b)*e^{(-2*I*c*x^2 - 2*I*b*x - 2*I*a)} - \sqrt{\pi}*(b^2 - I*c)*\operatorname{erf}(-1/2*\sqrt{c}*(2*x + b/c)*(I*c/abs(c) + 1))*e^{(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))}/c^2$

Mupad [F(-1)]

Timed out.

$$\int x^2 \sin^2(a + bx + cx^2) dx = \int x^2 \sin(cx^2 + bx + a)^2 dx$$

input `int(x^2*sin(a + b*x + c*x^2)^2,x)`

output `int(x^2*sin(a + b*x + c*x^2)^2, x)`

Reduce [F]

$$\int x^2 \sin^2(a + bx + cx^2) dx$$

$$= \frac{\cos(cx^2 + bx + a) \sin(cx^2 + bx + a) b - 2 \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) cx + 6 \left(\int \frac{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + a)}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + a)^2} dx \right)}{6}$$

input `int(x^2*sin(c*x^2+b*x+a)^2,x)`

output `(cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*b - 2*cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*c*x + 6*int(tan((a + b*x + c*x**2)/2)**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2 + 8*int(x*x**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**2 + 8*int(tan((a + b*x + c*x**2)/2)/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c - 2*int(1/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2 + sin(a + b*x + c*x**2)*b - 2*sin(a + b*x + c*x**2)*c*x)/(6*c**2)`

3.17 $\int x \sin^2(a + bx + cx^2) dx$

Optimal result	142
Mathematica [A] (verified)	142
Rubi [A] (verified)	143
Maple [A] (verified)	144
Fricas [A] (verification not implemented)	144
Sympy [F]	145
Maxima [C] (verification not implemented)	145
Giac [C] (verification not implemented)	146
Mupad [F(-1)]	147
Reduce [F]	147

Optimal result

Integrand size = 15, antiderivative size = 126

$$\begin{aligned} \int x \sin^2(a + bx + cx^2) dx &= \frac{x^2}{4} + \frac{b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ &\quad - \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} \\ &\quad - \frac{\sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output
$$1/4*x^2+1/8*b*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)-1/8*b*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)-1/8*sin(2*c*x^2+2*b*x+2*a)/c$$

Mathematica [A] (verified)

Time = 0.36 (sec), antiderivative size = 118, normalized size of antiderivative = 0.94

$$\begin{aligned} \int x \sin^2(a + bx + cx^2) dx \\ = \frac{b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 - \sin(2(a + x(b + 2cx))))}{8c^{3/2}} \end{aligned}$$

input `Integrate[x*Sin[a + b*x + c*x^2]^2,x]`

output
$$\begin{aligned} & (b \sqrt{\pi}) \cos[2a - b^2/(2c)] \text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] \\ & - b \sqrt{\pi} \text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] \sin[2a - b^2/(2c)] \\ & + \sqrt{c} (2cx^2 - \sin[2(a + x(b + cx))])/(8c^{(3/2)}) \end{aligned}$$

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \sin^2(a + bx + cx^2) dx \\ & \quad \downarrow 3948 \\ & \int \left(\frac{x}{2} - \frac{1}{2}x \cos(2a + 2bx + 2cx^2) \right) dx \\ & \quad \downarrow 2009 \\ & \frac{\sqrt{\pi}b \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} - \frac{\sqrt{\pi}b \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} - \\ & \quad \frac{\sin(2a + 2bx + 2cx^2)}{8c} + \frac{x^2}{4} \end{aligned}$$

input `Int[x*Sin[a + b*x + c*x^2]^2,x]`

output
$$\begin{aligned} & x^{2/4} + (b \sqrt{\pi}) \cos[2a - b^2/(2c)] \text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]/(8c^{(3/2)}) \\ & - (b \sqrt{\pi}) \text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] \sin[2a - b^2/(2c)]/(8c^{(3/2)}) - \sin[2a + 2bx + 2cx^2]/(8c) \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3948 $\text{Int}[(d_.) + (e_.)*(x_.))^(m_.)*\text{Sin}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^n_, x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \text{Sin}[a + b*x + c*x^2]^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{IGtQ}[n, 1]$

Maple [A] (verified)

Time = 1.26 (sec), antiderivative size = 95, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{x^2}{4} - \frac{\sin(2cx^2+2bx+2a)}{8c} + \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{8c^{3/2}}$	95
risch	$\frac{b\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c\sqrt{ic}} - \frac{b\sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \operatorname{erf}\left(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{16c\sqrt{-2ic}} + \frac{x^2}{4} - \frac{\sin(2cx^2+2bx+2a)}{8c}$	141

input $\text{int}(x*\sin(c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{4}x^2 - \frac{1}{8}\sin(2cx^2+2bx+2a)/c + \frac{1}{8}b/c^{(3/2)}\text{Pi}^{(1/2)}(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}) + \sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 124, normalized size of antiderivative = 0.98

$$\begin{aligned} & \int x \sin^2(a + bx + cx^2) dx \\ &= \frac{\pi b \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi b \sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2-4ac}{2c}\right) + 2c^2x^2 - 2c \cos(cx^2 + bx + a)}{8c^2} \end{aligned}$$

input $\text{integrate}(x*\sin(c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

output

```
1/8*(pi*b*sqrt(c/pi)*cos(-1/2*(b^2 - 4*a*c)/c)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - pi*b*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c)*sin(-1/2*(b^2 - 4*a*c)/c) + 2*c^2*x^2 - 2*c*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a))/c^2
```

Sympy [F]

$$\int x \sin^2(a + bx + cx^2) dx = \int x \sin^2(a + bx + cx^2) dx$$

input

```
integrate(x*sin(c*x**2+b*x+a)**2,x)
```

output

```
Integral(x*sin(a + b*x + c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 611, normalized size of antiderivative = 4.85

$$\int x \sin^2(a + bx + cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*sin(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

```
1/64*sqrt(2)*((-I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/2*(b^2 - 4*a*c)/c) + (-I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(-I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) + I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) - c*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c)/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c)))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.35

$$\int x \sin^2(a + bx + cx^2) dx$$

$$= \frac{1}{4} x^2 - \frac{\frac{\sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x + \frac{b}{c}\right) \left(-\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{ib^2 - 4ic}{2c}\right)}}{\sqrt{c} \left(-\frac{ic}{|c|} + 1\right)} - i e^{(2icx^2 + 2ibx + 2ia)}}{16c}$$

$$- \frac{\frac{\sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x + \frac{b}{c}\right) \left(\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{-ib^2 + 4ic}{2c}\right)}}{\sqrt{c} \left(\frac{ic}{|c|} + 1\right)} + i e^{(-2icx^2 - 2ibx - 2ia)}}{16c}$$

input `integrate(x*sin(c*x^2+b*x+a)^2,x, algorithm="giac")`

output

```
1/4*x^2 - 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))
*  
*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) - I*e^(2*I*c*x^2
+ 2*I*b*x + 2*I*a))/c - 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*
c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)) +
I*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \sin^2(a + bx + cx^2) dx = \int x \sin(cx^2 + bx + a)^2 dx$$

input

```
int(x*sin(a + b*x + c*x^2)^2,x)
```

output

```
int(x*sin(a + b*x + c*x^2)^2, x)
```

Reduce [F]

$$\begin{aligned} & \int x \sin^2(a + bx + cx^2) dx \\ &= \frac{-\cos(cx^2 + bx + a) \sin(cx^2 + bx + a) - 2 \left(\int \sin(cx^2 + bx + a)^2 dx \right) b + bx + cx^2}{4c} \end{aligned}$$

input

```
int(x*sin(c*x^2+b*x+a)^2,x)
```

output

```
( - cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2) - 2*int(sin(a + b*x + c*x**2)**2,x)*b + b*x + c*x**2)/(4*c)
```

3.18 $\int \sin^2(a + bx + cx^2) dx$

Optimal result	148
Mathematica [A] (verified)	148
Rubi [A] (verified)	149
Maple [A] (verified)	150
Fricas [A] (verification not implemented)	150
Sympy [A] (verification not implemented)	151
Maxima [C] (verification not implemented)	151
Giac [C] (verification not implemented)	152
Mupad [F(-1)]	152
Reduce [F]	153

Optimal result

Integrand size = 13, antiderivative size = 100

$$\begin{aligned} \int \sin^2(a + bx + cx^2) dx &= \frac{x}{2} - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} \\ &\quad + \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

output

```
1/2*x-1/4*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))
/c^(1/2)+1/4*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2
/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec), antiderivative size = 97, normalized size of antiderivative = 0.97

$$\begin{aligned} \int \sin^2(a + bx + cx^2) dx &= \frac{2\sqrt{c}x - \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

input `Integrate[Sin[a + b*x + c*x^2]^2, x]`

output
$$\frac{(2\sqrt{c}x - \sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] + \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])}{(4\sqrt{c})}$$

Rubi [A] (verified)

Time = 0.24 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3930, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(a + bx + cx^2) dx \\ & \quad \downarrow \text{3930} \\ & \int \left(\frac{1}{2} - \frac{1}{2} \cos(2a + 2bx + 2cx^2) \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2} \end{aligned}$$

input `Int[Sin[a + b*x + c*x^2]^2, x]`

output
$$\frac{x}{2} - \frac{(\sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] + \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])}{(4\sqrt{c})}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3930 $\text{Int}[\text{Sin}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTri}\text{gReduce}[\text{Sin}[a + b*x + c*x^2]^n, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{IGtQ}[n, 1]$

Maple [A] (verified)

Time = 1.20 (sec), antiderivative size = 72, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{x}{2} - \frac{\sqrt{\pi} (\cos(\frac{-4ac+b^2}{2c}) \text{FresnelC}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}) + \sin(\frac{-4ac+b^2}{2c}) \text{FresnelS}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}))}{4\sqrt{c}}$	72
risch	$\frac{x}{2} - \frac{\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \text{erf}(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}})}{16\sqrt{ic}} + \frac{\sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \text{erf}(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}})}{8\sqrt{-2ic}}$	111

input $\text{int}(\text{sin}(c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $1/2*x - 1/4*\text{Pi}^{(1/2)}/c^{(1/2)}*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}) + \sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 93, normalized size of antiderivative = 0.93

$$\int \sin^2(a + bx + cx^2) dx \\ = -\frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2-4ac}{2c}\right) - 2cx}{4c}$$

input $\text{integrate}(\text{sin}(c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

output

$$\begin{aligned} & -\frac{1}{4} \left(\frac{\pi \sqrt{c/\pi}}{c} \cos \left(-\frac{1}{2} (b^2 - 4ac)/c \right) \operatorname{fresnel_cos}((2cx + b)\sqrt{c/\pi}) \right. \\ & \quad \left. - \frac{\pi \sqrt{c/\pi}}{c} \operatorname{fresnel_sin}((2cx + b)\sqrt{c/\pi}) \sin \left(-\frac{1}{2} (b^2 - 4ac)/c \right) - \frac{2cx}{c} \right) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.43 (sec), antiderivative size = 83, normalized size of antiderivative = 0.83

$$\begin{aligned} & \int \sin^2(a + bx + cx^2) dx \\ &= \frac{x}{2} - \frac{\sqrt{\pi} \left(-\sin \left(2a - \frac{b^2}{2c} \right) S \left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}} \right) + \cos \left(2a - \frac{b^2}{2c} \right) C \left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}} \right) \right) \sqrt{\frac{1}{c}}}{4} \end{aligned}$$

input

```
integrate(sin(c*x**2+b*x+a)**2,x)
```

output

$$\begin{aligned} & \frac{x}{2} - \sqrt{\pi} \left(-\sin(2a - b^2/(2c)) \operatorname{fresnels}((2b + 4cx)/(2\sqrt{\pi}\sqrt{c})) + \cos(2a - b^2/(2c)) \operatorname{fresnelc}((2b + 4cx)/(2\sqrt{\pi}\sqrt{c})) \right) \sqrt{1/c}/4 \end{aligned}$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec), antiderivative size = 124, normalized size of antiderivative = 1.24

$$\begin{aligned} & \int \sin^2(a + bx + cx^2) dx \\ &= \frac{4^{1/4} \sqrt{2} \sqrt{\pi} \left(((i-1) \cos \left(-\frac{b^2-4ac}{2c} \right) + (i+1) \sin \left(-\frac{b^2-4ac}{2c} \right)) \operatorname{erf} \left(\frac{2icx+ib}{\sqrt{2i}c} \right) + ((i+1) \cos \left(-\frac{b^2-4ac}{2c} \right) + (i-1) \sin \left(-\frac{b^2-4ac}{2c} \right)) \operatorname{erfc} \left(\frac{2icx+ib}{\sqrt{2i}c} \right) \right)}{32c^2} \end{aligned}$$

input

```
integrate(sin(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

$$\begin{aligned} & \frac{1}{32} \left(4^{1/4} \sqrt{2} \sqrt{\pi} \left(((I-1) \cos(-1/2(b^2 - 4ac)/c) + (I+1) \sin(-1/2(b^2 - 4ac)/c)) \operatorname{erf}((2*I*c*x + I*b)/\sqrt{2*I*c}) + ((I+1) \cos(-1/2(b^2 - 4ac)/c) + (I-1) \sin(-1/2(b^2 - 4ac)/c)) \operatorname{erfc}((2*I*c*x + I*b)/\sqrt{-2*I*c}) \right) c^{3/2} + 16c^2 x \right) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.22

$$\int \sin^2(a + bx + cx^2) dx = \frac{1}{2}x + \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 - 4iac}{2c}\right)}}{8\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)} \\ + \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 + 4iac}{2c}\right)}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}$$

input `integrate(sin(c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/2*x + 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) + 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \sin^2(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a)^2 dx$$

input `int(sin(a + b*x + c*x^2)^2,x)`

output `int(sin(a + b*x + c*x^2)^2, x)`

Reduce [F]

$$\int \sin^2(a + bx + cx^2) dx = \int \sin(c x^2 + bx + a)^2 dx$$

input `int(sin(c*x^2+b*x+a)^2,x)`

output `int(sin(a + b*x + c*x**2)**2,x)`

3.19 $\int \frac{\sin^2(a+bx+cx^2)}{x} dx$

Optimal result	154
Mathematica [N/A]	154
Rubi [N/A]	155
Maple [N/A]	156
Fricas [N/A]	156
Sympy [N/A]	156
Maxima [N/A]	157
Giac [N/A]	157
Mupad [N/A]	158
Reduce [N/A]	158

Optimal result

Integrand size = 17, antiderivative size = 17

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \frac{\log(x)}{2} - \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx + 2cx^2)}{x}, x\right)$$

output `1/2*ln(x)-1/2*Defe r (Int)(cos(2*c*x^2+2*b*x+2*a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 7.34 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin^2(a + bx + cx^2)}{x} dx$$

input `Integrate[Sin[a + b*x + c*x^2]^2/x, x]`

output `Integrate[$\sin(a + b*x + c*x^2)^2/x$, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin^2(a + bx + cx^2)}{x} dx \\ & \quad \downarrow \textcolor{blue}{3948} \\ & \int \left(\frac{1}{2x} - \frac{\cos(2a + 2bx + 2cx^2)}{2x} \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{\log(x)}{2} - \frac{1}{2} \int \frac{\cos(2cx^2 + 2bx + 2a)}{x} dx \end{aligned}$$

input `Int[Sin[a + b*x + c*x^2]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_)*(x_))^(m_)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\sin(cx^2 + bx + a)^2}{x} dx$$

input `int(sin(c*x^2+b*x+a)^2/x,x)`

output `int(sin(c*x^2+b*x+a)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)^2}{x} dx$$

input `integrate(sin(c*x^2+b*x+a)^2/x,x, algorithm="fricas")`

output `integral(-(cos(c*x^2 + b*x + a)^2 - 1)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.79 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin^2(a + bx + cx^2)}{x} dx$$

input `integrate(sin(c*x**2+b*x+a)**2/x,x)`

output `Integral(sin(a + b*x + c*x**2)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.65

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)^2}{x} dx$$

input `integrate(sin(c*x^2+b*x+a)^2/x,x, algorithm="maxima")`

output `-1/2*integrate(cos(2*c*x^2 + 2*b*x + 2*a)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)^2}{x} dx$$

input `integrate(sin(c*x^2+b*x+a)^2/x,x, algorithm="giac")`

output `integrate(sin(c*x^2 + b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 38.24 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)^2}{x} dx$$

input `int(sin(a + b*x + c*x^2)^2/x,x)`

output `int(sin(a + b*x + c*x^2)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\sin^2(a + bx + cx^2)}{x} dx = \int \frac{\sin(cx^2 + bx + a)^2}{x} dx$$

input `int(sin(c*x^2+b*x+a)^2/x,x)`

output `int(sin(a + b*x + c*x**2)**2/x,x)`

3.20 $\int x^2 \sin^2(a + bx - cx^2) dx$

Optimal result	159
Mathematica [A] (verified)	160
Rubi [A] (verified)	160
Maple [A] (verified)	161
Fricas [A] (verification not implemented)	162
Sympy [F]	162
Maxima [C] (verification not implemented)	163
Giac [C] (verification not implemented)	164
Mupad [F(-1)]	164
Reduce [F]	165

Optimal result

Integrand size = 18, antiderivative size = 248

$$\begin{aligned} \int x^2 \sin^2(a + bx - cx^2) dx = & \frac{x^3}{6} + \frac{b^2 \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & - \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & + \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{3/2}} \\ & + \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{5/2}} \\ & + \frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx - 2cx^2)}{8c} \end{aligned}$$

output

```
1/6*x^3+1/16*b^2*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(5/2)-1/16*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)+1/16*Pi^(1/2)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)/c^(3/2)+1/16*b^2*Pi^(1/2)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)/c^(5/2)+1/16*b*sin(-2*c*x^2+2*b*x+2*a)/c^2+1/8*x*sin(-2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.71

$$\int x^2 \sin^2(a + bx - cx^2) dx \\ = \frac{3\sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a + \frac{b^2}{2c}\right) - b^2 \sin\left(2a + \frac{b^2}{2c}\right)\right) - 3\sqrt{\pi} \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a + \frac{b^2}{2c}\right) - c \sin\left(2a + \frac{b^2}{2c}\right)\right)}{48c^{5/2}}$$

input `Integrate[x^2*Sin[a + b*x - c*x^2]^2, x]`

output
$$(3*\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(-b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*(c*\text{Cos}[2*a + b^2/(2*c)] - b^2*\text{Sin}[2*a + b^2/(2*c)]) - 3*\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(-b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*(b^2*\text{Cos}[2*a + b^2/(2*c)] + c*\text{Sin}[2*a + b^2/(2*c)]) + \text{Sqrt}[c]*(8*c^2*x^3 + 3*(b + 2*c*x)*\text{Sin}[2*(a + x*(b - c*x))]))/(48*c^{(5/2)})$$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sin^2(a + bx - cx^2) dx \\ \downarrow 3948 \\ \int \left(\frac{x^2}{2} - \frac{1}{2}x^2 \cos(2a + 2bx - 2cx^2)\right) dx \\ \downarrow 2009$$

$$\begin{aligned} & \frac{\sqrt{\pi} \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} + \frac{\sqrt{\pi}b^2 \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \\ & \frac{\sqrt{\pi}b^2 \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} - \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} + \\ & \frac{b \sin\left(2a + 2bx - 2cx^2\right)}{16c^2} + \frac{x \sin\left(2a + 2bx - 2cx^2\right)}{8c} + \frac{x^3}{6} \end{aligned}$$

input `Int[x^2*Sin[a + b*x - c*x^2]^2,x]`

output $x^{3/6} + (b^{2*}\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a + b^{2/(2*c)}]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])]/(16*c^{(5/2)}) - (\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a + b^{2/(2*c)}]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])]/(16*c^{(3/2)}) + (\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a + b^{2/(2*c)}]/(16*c^{(3/2)}) + (b^{2*}\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a + b^{2/(2*c)}]/(16*c^{(5/2)}) + (b*\text{Sin}[2*a + 2*b*x - 2*c*x^2])/(16*c^2) + (x*\text{Sin}[2*a + 2*b*x - 2*c*x^2])/(8*c)$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_)*(x_))^(m_)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.41 (sec), antiderivative size = 199, normalized size of antiderivative = 0.80

method	result
default	$\frac{x^3}{6} + \frac{x \sin(-2cx^2 + 2bx + 2a)}{8c} - \frac{b \left(-\frac{\sin(-2cx^2 + 2bx + 2a)}{4c} + \frac{b\sqrt{\pi} \left(\cos\left(\frac{4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{3/2}} \right)}{4c}$
risch	$-\frac{b^2\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{2c}}\text{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{32c^2\sqrt{-2ic}} - \frac{i\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{2c}}\text{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{32c\sqrt{-2ic}} + \frac{b^2\sqrt{\pi}e^{\frac{i(4ac+b^2)}{2c}}\sqrt{2}\text{erf}\left(-\sqrt{2}\sqrt{ic}x - \frac{b}{\sqrt{2}\sqrt{ic}}\right)}{64c^2\sqrt{ic}}$

input `int(x^2*sin(-c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1/6*x^3 + 1/8*x*sin(-2*c*x^2 + 2*b*x + 2*a)/c - 1/4*b/c*(-1/4*sin(-2*c*x^2 + 2*b*x + 2*a)/c + 1/4*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)) + sin(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)))) + 1/16/c^(3/2)*Pi^(1/2)*(cos(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)) - sin(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)))}}{48*c^3}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.75

$$\int x^2 \sin^2(a + bx - cx^2) dx = \frac{8 c^3 x^3 - 6 (2 c^2 x + bc) \cos(cx^2 - bx - a) \sin(cx^2 - bx - a) - 3 \left(\pi b^2 \cos\left(\frac{b^2 + 4 ac}{2 c}\right) + \pi c \sin\left(\frac{b^2 + 4 ac}{2 c}\right)\right) \sqrt{48 c^3}}{48 c^3}$$

input `integrate(x^2*sin(-c*x^2+b*x+a)^2,x, algorithm="fricas")`

output
$$\frac{1/48*(8*c^3*x^3 - 6*(2*c^2*x + b*c)*cos(c*x^2 - b*x - a)*sin(c*x^2 - b*x - a) - 3*(pi*b^2*cos(1/2*(b^2 + 4*a*c)/c) + pi*c*sin(1/2*(b^2 + 4*a*c)/c))*sqrt(c/pi)*fresnel_cos((2*c*x - b)*sqrt(c/pi)/c) - 3*(pi*b^2*sin(1/2*(b^2 + 4*a*c)/c) - pi*c*cos(1/2*(b^2 + 4*a*c)/c))*sqrt(c/pi)*fresnel_sin((2*c*x - b)*sqrt(c/pi)/c))/c^3}{48*c^3}$$

Sympy [F]

$$\int x^2 \sin^2(a + bx - cx^2) dx = \int x^2 \sin^2(a + bx - cx^2) dx$$

input `integrate(x**2*sin(-c*x**2+b*x+a)**2,x)`

output `Integral(x**2*sin(a + b*x - c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 1617, normalized size of antiderivative = 6.52

$$\int x^2 \sin^2 (a + bx - cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*sin(-c*x^2+b*x+a)^2,x, algorithm="maxima")`

output `-1/384*sqrt(2)*(24*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*((I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*cos(1/2*(b^2 + 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*sin(1/2*(b^2 + 4*a*c)/c))*x^3 + 36*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(1/2*(b^2 + 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*(-(I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)...)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.87

$$\int x^2 \sin^2(a + bx - cx^2) dx = \frac{1}{6} x^3$$

$$-\frac{(c(-2i x + \frac{ib}{c}) - 2i b)e^{(2i cx^2 - 2i bx - 2i a)} - \frac{\sqrt{\pi}(b^2 + i c) \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x - \frac{b}{c}\right) \left(-\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{ib^2 + 4i ac}{2c}\right)}}{\sqrt{c} \left(-\frac{ic}{|c|} + 1\right)}}{32 c^2}$$

$$-\frac{(c(2i x - \frac{ib}{c}) + 2i b)e^{(-2i cx^2 + 2i bx + 2i a)} - \frac{\sqrt{\pi}(b^2 - i c) \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x - \frac{b}{c}\right) \left(\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{-ib^2 - 4i ac}{2c}\right)}}{\sqrt{c} \left(\frac{ic}{|c|} + 1\right)}}{32 c^2}$$

input `integrate(x^2*sin(-c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/6*x^3 - 1/32*((c*(-2*I*x + I*b/c) - 2*I*b)*e^(2*I*c*x^2 - 2*I*b*x - 2*I*a) - sqrt(pi)*(b^2 + I*c)*erf(-1/2*sqrt(c)*(2*x - b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)))/c^2 - 1/32*((c*(2*I*x - I*b/c) + 2*I*b)*e^(-2*I*c*x^2 + 2*I*b*x + 2*I*a) - sqrt(pi)*(b^2 - I*c)*erf(-1/2*sqrt(c)*(2*x - b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)))/c^2`

Mupad [F(-1)]

Timed out.

$$\int x^2 \sin^2(a + bx - cx^2) dx = \int x^2 \sin(-cx^2 + bx + a)^2 dx$$

input `int(x^2*sin(a + b*x - c*x^2)^2,x)`

output `int(x^2*sin(a + b*x - c*x^2)^2, x)`

Reduce [F]

$$\int x^2 \sin^2(a + bx - cx^2) dx$$

$$= \frac{\cos(-cx^2 + bx + a) \sin(-cx^2 + bx + a) b + 2 \cos(-cx^2 + bx + a) \sin(-cx^2 + bx + a) cx + 6 \left(\int \frac{1}{\tan(-cx^2 + bx + a)} dx \right)}{6}$$

input `int(x^2*sin(-c*x^2+b*x+a)^2,x)`

output `(cos(a + b*x - c*x**2)*sin(a + b*x - c*x**2)*b + 2*cos(a + b*x - c*x**2)*sin(a + b*x - c*x**2)*c*x + 6*int(tan((a + b*x - c*x**2)/2)**2/(tan((a + b*x - c*x**2)/2)**4 + 2*tan((a + b*x - c*x**2)/2)**2 + 1),x)*b**2 + 8*int(x*x**2/(tan((a + b*x - c*x**2)/2)**4 + 2*tan((a + b*x - c*x**2)/2)**2 + 1),x)*c**2 - 8*int(tan((a + b*x - c*x**2)/2)/(tan((a + b*x - c*x**2)/2)**4 + 2*tan((a + b*x - c*x**2)/2)**2 + 1),x)*c - 2*int(1/(tan((a + b*x - c*x**2)/2)**4 + 2*tan((a + b*x - c*x**2)/2)**2 + 1),x)*b**2 + sin(a + b*x - c*x**2)*b + 2*sin(a + b*x - c*x**2)*c*x)/(6*c**2)`

3.21 $\int x \sin^2(a + bx - cx^2) dx$

Optimal result	166
Mathematica [A] (verified)	166
Rubi [A] (verified)	167
Maple [A] (verified)	168
Fricas [A] (verification not implemented)	168
Sympy [F]	169
Maxima [C] (verification not implemented)	169
Giac [C] (verification not implemented)	170
Mupad [F(-1)]	171
Reduce [F]	171

Optimal result

Integrand size = 16, antiderivative size = 126

$$\begin{aligned} \int x \sin^2(a + bx - cx^2) dx &= \frac{x^2}{4} + \frac{b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ &\quad + \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{8c^{3/2}} \\ &\quad + \frac{\sin(2a + 2bx - 2cx^2)}{8c} \end{aligned}$$

output
$$\frac{1/4*x^2+1/8*b*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)+1/8*b*Pi^(1/2)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)/c^(3/2)+1/8*sin(-2*c*x^2+2*b*x+2*a)/c}{8*c^(3/2)}$$

Mathematica [A] (verified)

Time = 0.28 (sec), antiderivative size = 122, normalized size of antiderivative = 0.97

$$\begin{aligned} \int x \sin^2(a + bx - cx^2) dx \\ = \frac{-b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - b\sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 + \sin(2(a + \frac{b^2}{2c})x^2))}{8c^{3/2}} \end{aligned}$$

input `Integrate[x*Sin[a + b*x - c*x^2]^2,x]`

output
$$\frac{(-b\sqrt{\pi}\cos(2a + b^2/(2c))\text{FresnelC}((-b + 2cx)/(\sqrt{c}\sqrt{\pi})) - b\sqrt{\pi}\text{FresnelS}((-b + 2cx)/(\sqrt{c}\sqrt{\pi}))\sin(2a + b^2/(2c)) + \sqrt{c}(2cx^2 + \sin(2(a + x(b - cx))))}{(8c^{3/2})}$$

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \sin^2(a + bx - cx^2) dx \\ & \quad \downarrow 3948 \\ & \int \left(\frac{x}{2} - \frac{1}{2}x \cos(2a + 2bx - 2cx^2) \right) dx \\ & \quad \downarrow 2009 \\ & \frac{\sqrt{\pi}b \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} + \frac{\sqrt{\pi}b \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} + \\ & \quad \frac{\sin(2a + 2bx - 2cx^2)}{8c} + \frac{x^2}{4} \end{aligned}$$

input `Int[x*Sin[a + b*x - c*x^2]^2,x]`

output
$$\frac{x^2/4 + (b\sqrt{\pi}\cos(2a + b^2/(2c))\text{FresnelC}((b - 2cx)/(\sqrt{c}\sqrt{\pi}))/((8c^{3/2})) + (b\sqrt{\pi}\text{FresnelS}((b - 2cx)/(\sqrt{c}\sqrt{\pi}))\sin(2a + b^2/(2c))/((8c^{3/2})) + \sin(2a + 2bx - 2cx^2)/(8c)}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3948 $\text{Int}[(d_.) + (e_.)*(x_.))^(m_.)*\text{Sin}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \ \text{Sin}[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c, \ d, \ e, \ m\}, \ x] \ \&& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 1.26 (sec), antiderivative size = 99, normalized size of antiderivative = 0.79

method	result	S
default	$\frac{x^2}{4} + \frac{\sin(-2cx^2+2bx+2a)}{8c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) \right)}{8c^{3/2}}$	9
risch	$- \frac{b\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \text{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{16c\sqrt{-2ic}} + \frac{b\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \text{erf}\left(-\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c\sqrt{ic}} + \frac{x^2}{4} + \frac{\sin(-2cx^2+2bx+2a)}{8c}$	1

input $\text{int}(x*\sin(-c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{4}x^2 + \frac{1}{8}\sin(-2cx^2+2bx+2a)/c - \frac{1}{8}b/c^{(3/2)}\text{Pi}^{(1/2)}(\cos(1/2*(4*a*c+b^2)/c)\text{FresnelC}(1/\text{Pi}^{(1/2)}/c^{(1/2)}*(2*c*x-b)) + \sin(1/2*(4*a*c+b^2)/c)\text{FresnelS}(1/\text{Pi}^{(1/2)}/c^{(1/2)}*(2*c*x-b))$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 133, normalized size of antiderivative = 1.06

$$\int x \sin^2(a + bx - cx^2) dx = \frac{\pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{2c}\right) C\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) + \pi b \sqrt{\frac{c}{\pi}} S\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(\frac{b^2+4ac}{2c}\right) - 2c^2x^2 + 2c \cos(cx^2 - bx)}{8c^2}$$

input $\text{integrate}(x*\sin(-c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

output

```
-1/8*(pi*b*sqrt(c/pi)*cos(1/2*(b^2 + 4*a*c)/c)*fresnel_cos((2*c*x - b)*sqrt(pi/c) + pi*b*sqrt(c/pi)*fresnel_sin((2*c*x - b)*sqrt(c/pi)/c)*sin(1/2*(b^2 + 4*a*c)/c) - 2*c^2*x^2 + 2*c*cos(c*x^2 - b*x - a)*sin(c*x^2 - b*x - a))/c^2
```

Sympy [F]

$$\int x \sin^2(a + bx - cx^2) dx = \int x \sin^2(a + bx - cx^2) dx$$

input

```
integrate(x*sin(-c*x**2+b*x+a)**2,x)
```

output

```
Integral(x*sin(a + b*x - c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 610, normalized size of antiderivative = 4.84

$$\int x \sin^2(a + bx - cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*sin(-c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

```
1/64*sqrt(2)*((-I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(1/2*(b^2 + 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(1/2*(b^2 + 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(1/2*(b^2 + 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(1/2*(b^2 + 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(-I*e^(1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) + I*e^(-1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*cos(1/2*(b^2 + 4*a*c)/c) + c*(e^(1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) *sin(1/2*(b^2 + 4*a*c)/c))*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c))/(c^2*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.37

$$\int x \sin^2(a + bx - cx^2) dx$$

$$= \frac{1}{4} x^2 + \frac{\frac{\sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x - \frac{b}{c}\right) \left(-\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{ib^2 + 4ic}{2c}\right)}}{\sqrt{c} \left(-\frac{ic}{|c|} + 1\right)} + i e^{(2icx^2 - 2ibx - 2ia)}}{16c}$$

$$+ \frac{\frac{\sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x - \frac{b}{c}\right) \left(\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{-ib^2 - 4ic}{2c}\right)}}{\sqrt{c} \left(\frac{ic}{|c|} + 1\right)} - i e^{(-2icx^2 + 2ibx + 2ia)}}{16c}$$

input `integrate(x*sin(-c*x^2+b*x+a)^2,x, algorithm="giac")`

output

```
1/4*x^2 + 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x - b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) + I*e^(2*I*c*x^2 - 2*I*b*x - 2*I*a))/c + 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x - b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)) - I*e^(-2*I*c*x^2 + 2*I*b*x + 2*I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \sin^2(a + bx - cx^2) dx = \int x \sin(-cx^2 + bx + a)^2 dx$$

input

```
int(x*sin(a + b*x - c*x^2)^2,x)
```

output

```
int(x*sin(a + b*x - c*x^2)^2, x)
```

Reduce [F]

$$\int x \sin^2(a + bx - cx^2) dx = \frac{\cos(-cx^2 + bx + a) \sin(-cx^2 + bx + a) + 2 \left(\int \sin(-cx^2 + bx + a)^2 dx \right) b - bx + cx^2}{4c}$$

input

```
int(x*sin(-c*x^2+b*x+a)^2,x)
```

output

```
(cos(a + b*x - c*x**2)*sin(a + b*x - c*x**2) + 2*int(sin(a + b*x - c*x**2)**2,x)*b - b*x + c*x**2)/(4*c)
```

3.22 $\int \sin^2(a + bx - cx^2) dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [A] (verified)	174
Fricas [A] (verification not implemented)	174
Sympy [A] (verification not implemented)	175
Maxima [C] (verification not implemented)	175
Giac [C] (verification not implemented)	176
Mupad [F(-1)]	176
Reduce [F]	177

Optimal result

Integrand size = 14, antiderivative size = 100

$$\begin{aligned} \int \sin^2(a + bx - cx^2) dx &= \frac{x}{2} + \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} \\ &\quad + \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

output
$$\begin{aligned} &1/2*x+1/4*Pi^{(1/2)}*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^{(1/2)}/Pi^{(1/2)}) \\ &)/c^{(1/2)}+1/4*Pi^{(1/2)}*FresnelS((-2*c*x+b)/c^{(1/2)}/Pi^{(1/2)}))*sin(2*a+1/2*b^2/c)/c^{(1/2)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec), antiderivative size = 102, normalized size of antiderivative = 1.02

$$\begin{aligned} \int \sin^2(a + bx - cx^2) dx \\ = \frac{2\sqrt{c}x - \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - \sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

input `Integrate[Sin[a + b*x - c*x^2]^2, x]`

output
$$\frac{(2\sqrt{c}x - \sqrt{\pi}\cos[2a + b^2/(2c)]\text{FresnelC}[(-b + 2cx)/(\sqrt{c}\sqrt{\pi})] - \sqrt{\pi}\text{FresnelS}[(-b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a + b^2/(2c)])}{(4\sqrt{c})}$$

Rubi [A] (verified)

Time = 0.23 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3930, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin^2(a + bx - cx^2) dx \\
 & \quad \downarrow \text{3930} \\
 & \int \left(\frac{1}{2} - \frac{1}{2} \cos(2a + 2bx - 2cx^2) \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{\sqrt{\pi} \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2}
 \end{aligned}$$

input `Int[Sin[a + b*x - c*x^2]^2, x]`

output
$$\frac{x}{2} + \frac{(\sqrt{\pi}\cos[2a + b^2/(2c)]\text{FresnelC}[(b - 2cx)/(\sqrt{c}\sqrt{\pi})] + \sqrt{\pi}\text{FresnelS}[(b - 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a + b^2/(2c)])}{(4\sqrt{c})}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3930 $\text{Int}[\text{Sin}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTri}\text{gReduce}[\text{Sin}[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c\}, \ x] \ \&\& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 1.20 (sec), antiderivative size = 76, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x}{2} - \frac{\sqrt{\pi} (\cos(\frac{4ac+b^2}{2c}) \text{FresnelC}(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}) + \sin(\frac{4ac+b^2}{2c}) \text{FresnelS}(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}))}{4\sqrt{c}}$	76
risch	$\frac{x}{2} - \frac{\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \text{erf}(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}})}{8\sqrt{-2ic}} + \frac{\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \text{erf}(-\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}})}{16\sqrt{ic}}$	107

input $\text{int}(\text{sin}(-c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $1/2*x - 1/4*\text{Pi}^{(1/2)}/c^{(1/2)}*(\cos(1/2*(4*a*c+b^2)/c)*\text{FresnelC}(1/\text{Pi}^{(1/2)}/c^{(1/2)}*(2*c*x-b)) + \sin(1/2*(4*a*c+b^2)/c)*\text{FresnelS}(1/\text{Pi}^{(1/2)}/c^{(1/2)}*(2*c*x-b)))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 96, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \sin^2(a + bx - cx^2) \, dx \\ &= -\frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{2c}\right) C\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) + \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(\frac{b^2+4ac}{2c}\right) - 2cx}{4c} \end{aligned}$$

input $\text{integrate}(\text{sin}(-c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

```
output -1/4*(pi*sqrt(c/pi)*cos(1/2*(b^2 + 4*a*c)/c)*fresnel_cos((2*c*x - b)*sqrt(c/pi)/c) + pi*sqrt(c/pi)*fresnel_sin((2*c*x - b)*sqrt(c/pi)/c)*sin(1/2*(b^2 + 4*a*c)/c) - 2*c*x)/c
```

Sympy [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.88

$$\int \sin^2(a + bx - cx^2) dx = \frac{x}{2} - \frac{\sqrt{\pi} \sqrt{-\frac{1}{c}} \left(-\sin\left(2a + \frac{b^2}{2c}\right) S\left(\frac{2b-4cx}{2\sqrt{\pi}\sqrt{-c}}\right) + \cos\left(2a + \frac{b^2}{2c}\right) C\left(\frac{2b-4cx}{2\sqrt{\pi}\sqrt{-c}}\right) \right)}{4}$$

```
input integrate(sin(-c*x**2+b*x+a)**2,x)
```

```
output x/2 - sqrt(pi)*sqrt(-1/c)*(-sin(2*a + b**2/(2*c))*fresnels((2*b - 4*c*x)/(2*sqrt(pi)*sqrt(-c))) + cos(2*a + b**2/(2*c))*fresnelc((2*b - 4*c*x)/(2*sqrt(pi)*sqrt(-c)))))/4
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \sin^2(a + bx - cx^2) dx =$$

$$-\frac{4^{\frac{1}{4}}\sqrt{2}\sqrt{\pi}\left(\left(-(i-1)\cos\left(\frac{b^2+4ac}{2c}\right)+(i+1)\sin\left(\frac{b^2+4ac}{2c}\right)\right)\operatorname{erf}\left(\frac{2i\sqrt{c}x-i\sqrt{b}}{\sqrt{2i}c}\right)+\left(-(i+1)\cos\left(\frac{b^2+4ac}{2c}\right)+\left(i+1\right)\sin\left(\frac{b^2+4ac}{2c}\right)\right)\operatorname{erf}\left(\frac{2i\sqrt{c}x+i\sqrt{b}}{\sqrt{2i}c}\right)\right)}{32c^2}$$

```
input integrate(sin(-c*x^2+b*x+a)^2,x, algorithm="maxima")
```

```

output -1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*((-I - 1)*cos(1/2*(b^2 + 4*a*c)/c) + (I +
1)*sin(1/2*(b^2 + 4*a*c)/c))*erf((2*I*c*x - I*b)/sqrt(2*I*c)) + (-I + 1)
*cos(1/2*(b^2 + 4*a*c)/c) + (I - 1)*sin(1/2*(b^2 + 4*a*c)/c))*erf((2*I*c*x -
I*b)/sqrt(-2*I*c)))*c^(3/2) - 16*c^2*x)/c^2

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \sin^2(a + bx - cx^2) dx = \frac{1}{2}x + \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x - \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2+4iac}{2c}\right)}}{8\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)} \\ + \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x - \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2-4iac}{2c}\right)}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}$$

input `integrate(sin(-c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/2*x + 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x - b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) + 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x - b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \sin^2(a + bx - cx^2) dx = \int \sin(-cx^2 + bx + a)^2 dx$$

input `int(sin(a + b*x - c*x^2)^2,x)`

output `int(sin(a + b*x - c*x^2)^2, x)`

Reduce [F]

$$\int \sin^2(a + bx - cx^2) dx = \int \sin(-cx^2 + bx + a)^2 dx$$

input `int(sin(-c*x^2+b*x+a)^2,x)`

output `int(sin(a + b*x - c*x**2)**2,x)`

3.23 $\int \frac{\sin^2(a+bx-cx^2)}{x} dx$

Optimal result	178
Mathematica [N/A]	178
Rubi [N/A]	179
Maple [N/A]	180
Fricas [N/A]	180
Sympy [N/A]	180
Maxima [N/A]	181
Giac [N/A]	181
Mupad [N/A]	182
Reduce [N/A]	182

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \frac{\log(x)}{2} - \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx - 2cx^2)}{x}, x\right)$$

output `1/2*ln(x)-1/2*DefeR(Int)(cos(-2*c*x^2+2*b*x+2*a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 7.85 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin^2(a + bx - cx^2)}{x} dx$$

input `Integrate[Sin[a + b*x - c*x^2]^2/x, x]`

output `Integrate[Sin[a + b*x - c*x^2]^2/x, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin^2(a + bx - cx^2)}{x} dx \\ & \quad \downarrow \textcolor{blue}{3948} \\ & \int \left(\frac{1}{2x} - \frac{\cos(2a + 2bx - 2cx^2)}{2x} \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{\log(x)}{2} - \frac{1}{2} \int \frac{\cos(-2cx^2 + 2bx + 2a)}{x} dx \end{aligned}$$

input `Int[Sin[a + b*x - c*x^2]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_)*(x_))^(m_.)*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\sin(-cx^2 + bx + a)^2}{x} dx$$

input `int(sin(-c*x^2+b*x+a)^2/x,x)`

output `int(sin(-c*x^2+b*x+a)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.39

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)^2}{x} dx$$

input `integrate(sin(-c*x^2+b*x+a)^2/x,x, algorithm="fricas")`

output `integral(-(cos(c*x^2 - b*x - a)^2 - 1)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.79 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin^2(a + bx - cx^2)}{x} dx$$

input `integrate(sin(-c*x**2+b*x+a)**2/x,x)`

output `Integral(sin(a + b*x - c*x**2)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.56

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)^2}{x} dx$$

input `integrate(sin(-c*x^2+b*x+a)^2/x,x, algorithm="maxima")`

output `-1/2*integrate(cos(2*c*x^2 - 2*b*x - 2*a)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)^2}{x} dx$$

input `integrate(sin(-c*x^2+b*x+a)^2/x,x, algorithm="giac")`

output `integrate(sin(-c*x^2 + b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 38.54 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)^2}{x} dx$$

input `int(sin(a + b*x - c*x^2)^2/x,x)`

output `int(sin(a + b*x - c*x^2)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\sin^2(a + bx - cx^2)}{x} dx = \int \frac{\sin(-cx^2 + bx + a)^2}{x} dx$$

input `int(sin(-c*x^2+b*x+a)^2/x,x)`

output `int(sin(a + b*x - c*x**2)**2/x,x)`

3.24 $\int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	183
Mathematica [A] (verified)	183
Rubi [A] (verified)	184
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	185
Sympy [F]	186
Maxima [C] (verification not implemented)	186
Giac [C] (verification not implemented)	187
Mupad [F(-1)]	187
Reduce [F]	188

Optimal result

Integrand size = 15, antiderivative size = 85

$$\begin{aligned} \int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{x^3}{6} - \frac{1}{16} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) + \frac{1}{16} \sqrt{\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{\pi}} \right) \\ & + \frac{1}{16} \sin \left(\frac{1}{2} + 2x + 2x^2 \right) - \frac{1}{8} x \sin \left(\frac{1}{2} + 2x + 2x^2 \right) \end{aligned}$$

output
$$1/6*x^3-1/16*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))+1/16*Pi^(1/2)*FresnelS((1+2*x)/Pi^(1/2))+1/16*sin(1/2+2*x+2*x^2)-1/8*x*sin(1/2+2*x+2*x^2)$$

Mathematica [A] (verified)

Time = 0.17 (sec), antiderivative size = 77, normalized size of antiderivative = 0.91

$$\begin{aligned} \int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{1}{48} \left(8x^3 - 3\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) \right. \\ & + 3\sqrt{\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{\pi}} \right) + 3 \sin \left(\frac{1}{2}(1+2x)^2 \right) \\ & \left. - 6x \sin \left(\frac{1}{2}(1+2x)^2 \right) \right) \end{aligned}$$

input `Integrate[x^2*Sin[1/4 + x + x^2]^2,x]`

output
$$(8*x^3 - 3*sqrt[\pi]*FresnelC[(1 + 2*x)/sqrt[\pi]] + 3*sqrt[\pi]*FresnelS[(1 + 2*x)/sqrt[\pi]] + 3*Sin[(1 + 2*x)^2/2] - 6*x*Sin[(1 + 2*x)^2/2])/48$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \sin^2 \left(x^2 + x + \frac{1}{4} \right) dx \\ & \quad \downarrow \textcolor{blue}{3948} \\ & \int \left(\frac{x^2}{2} - \frac{1}{2}x^2 \cos \left(2x^2 + 2x + \frac{1}{2} \right) \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & -\frac{1}{16}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x^3}{6} - \frac{1}{8}x \sin \left(2x^2 + 2x + \frac{1}{2} \right) + \\ & \quad \frac{1}{16} \sin \left(2x^2 + 2x + \frac{1}{2} \right) \end{aligned}$$

input
$$\operatorname{Int}[x^2 \sin[1/4 + x + x^2]^2, x]$$

output
$$x^3/6 - (\operatorname{sqrt}[\pi]*\operatorname{FresnelC}[(1 + 2*x)/\operatorname{sqrt}[\pi]])/16 + (\operatorname{sqrt}[\pi]*\operatorname{FresnelS}[(1 + 2*x)/\operatorname{sqrt}[\pi]])/16 + \operatorname{Sin}[1/2 + 2*x + 2*x^2]/16 - (x*\operatorname{Sin}[1/2 + 2*x + 2*x^2])/8$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3948 $\text{Int}[(d_.) + (e_.)*(x_.))^(m_.)*\text{Sin}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \ \text{Sin}[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c, \ d, \ e, \ m\}, \ x] \ \&& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 1.54 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.75

method	result
default	$\frac{x^3}{6} - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{16} + \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{16} + \frac{\sin\left(\frac{1}{2}+2x+2x^2\right)}{16} - \frac{x \sin\left(\frac{1}{2}+2x+2x^2\right)}{8}$
risch	$\frac{\sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \text{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{64} + \frac{(-1)^{\frac{1}{4}} \sqrt{\pi} \sqrt{2} \text{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{64} - \frac{\sqrt{\pi} \text{erf}\left(\sqrt{-2i} x - \frac{i}{\sqrt{-2i}}\right)}{32\sqrt{-2i}} - \frac{i \sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \text{erfc}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{64}$

input $\text{int}(x^2 * \sin(1/4 + x + x^2)^2, x, \text{method}=\text{RETURNVERBOSE})$

output
$$\frac{1}{6}x^3 - \frac{1}{16}\pi^{1/2} \text{FresnelC}\left(\frac{(1+2x)\pi^{1/2}}{2}\right) + \frac{1}{16}\pi^{1/2} \text{FresnelS}\left(\frac{(1+2x)\pi^{1/2}}{2}\right) + \frac{1}{16}\sin(1/2 + 2x + 2x^2) - \frac{1}{8}x \sin(1/2 + 2x + 2x^2)$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

$$\begin{aligned} \int x^2 \sin^2\left(\frac{1}{4} + x + x^2\right) dx &= \frac{1}{6}x^3 - \frac{1}{8}(2x - 1) \cos\left(x^2 + x + \frac{1}{4}\right) \sin\left(x^2 + x + \frac{1}{4}\right) \\ &\quad - \frac{1}{16}\sqrt{\pi} C\left(\frac{2x + 1}{\sqrt{\pi}}\right) + \frac{1}{16}\sqrt{\pi} S\left(\frac{2x + 1}{\sqrt{\pi}}\right) \end{aligned}$$

input $\text{integrate}(x^2 * \sin(1/4 + x + x^2)^2, x, \text{algorithm}=\text{"fricas"})$

output $\frac{1}{6}x^3 - \frac{1}{8}(2x - 1)\cos(x^2 + x + 1/4)\sin(x^2 + x + 1/4) - \frac{1}{16}\sqrt{\pi}\text{fresnel_cos}((2x + 1)/\sqrt{\pi}) + \frac{1}{16}\sqrt{\pi}\text{fresnel_sin}((2x + 1)/\sqrt{\pi})$

Sympy [F]

$$\int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x^2 \sin^2 \left(x^2 + x + \frac{1}{4} \right) dx$$

input `integrate(x**2*sin(1/4+x+x**2)**2, x)`

output `Integral(x**2*sin(x**2 + x + 1/4)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.01

$$\int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx \\ = \frac{128x^4 + 64x^3 + 48x \left(-ie^{(2ix^2+2ix+\frac{1}{2}i)} + ie^{(-2ix^2-2ix-\frac{1}{2}i)} \right) + 3\sqrt{8x^2+8x+2} \left((i-1)\sqrt{2}\sqrt{\pi} \operatorname{erf} \left(\sqrt{\frac{x^2+x+\frac{1}{4}}{\pi}} \right) + \sqrt{\frac{2}{\pi}}e^{-\frac{(x^2+x+\frac{1}{4})^2}{2}} \right)}{192}$$

input `integrate(x^2*sin(1/4+x+x^2)^2, x, algorithm="maxima")`

output $\frac{1}{384}(128x^4 + 64x^3 + 48x(-Ie^{(2Ix^2+2Ix+1/2I)} + Ie^{-(-2Ix^2-2Ix-1/2I)}) + 3\sqrt{8x^2+8x+2}((I-1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{2Ix^2+2Ix+1/2I}) - 1) - (I+1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{-2Ix^2-2Ix-1/2I}) - 1) - (2I+2)\sqrt{2}\sqrt{\pi}\gamma(3/2, 2Ix^2+2Ix+1/2I) + (2I-2)\sqrt{2}\sqrt{\pi}\gamma(3/2, -2Ix^2-2Ix-1/2I)) - 24Ie^{(2Ix^2+2Ix+1/2I)} + 24Ie^{-(-2Ix^2-2Ix-1/2I)})/(2x+1)$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.75

$$\int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{6} x^3 - \frac{1}{32} (-2ix + i)e^{(2ix^2+2ix+\frac{1}{2}i)} \\ - \frac{1}{32} (2ix - i)e^{(-2ix^2-2ix-\frac{1}{2}i)} \\ + \frac{1}{32}i\sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) \\ - \frac{1}{32}i\sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right)$$

input `integrate(x^2*sin(1/4+x+x^2)^2,x, algorithm="giac")`

output $\frac{1}{6}x^3 - \frac{1}{32}(-2Ix + I)e^{(2Ix^2 + 2Ix + 1/2I)} - \frac{1}{32}(2Ix - I)e^{(-2Ix^2 - 2Ix - 1/2I)} + \frac{1}{32}I\sqrt{\pi}\operatorname{erf}((I-1)x + 1/2I - 1/2) - \frac{1}{32}I\sqrt{\pi}\operatorname{erf}(-(I+1)x - 1/2I - 1/2)$

Mupad [F(-1)]

Timed out.

$$\int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x^2 \sin \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(x^2*sin(x + x^2 + 1/4)^2,x)`

output `int(x^2*sin(x + x^2 + 1/4)^2, x)`

Reduce [F]

$$\int x^2 \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \text{Too large to display}$$

input `int(x^2*sin(1/4+x+x^2)^2,x)`

output

```
(3*int(tan((4*x**2 + 4*x + 1)/8)**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan(
(4*x**2 + 4*x + 1)/8)**2 + 1),x)*tan((4*x**2 + 4*x + 1)/8)**4 + 6*int(tan(
(4*x**2 + 4*x + 1)/8)**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4
*x + 1)/8)**2 + 1),x)*tan((4*x**2 + 4*x + 1)/8)**2 + 3*int(tan((4*x**2 + 4
*x + 1)/8)**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)*
*2 + 1),x) + 4*int(x**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*
x + 1)/8)**2 + 1),x)*tan((4*x**2 + 4*x + 1)/8)**4 + 8*int(x**2/(tan((4*x**2 +
4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x)*tan((4*x**2 +
4*x + 1)/8)**2 + 4*int(x**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 +
4*x + 1)/8)**2 + 1),x) + 4*int(tan((4*x**2 + 4*x + 1)/8)/(tan((4*x**2 +
4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x)*tan((4*x**2 + 4*x
+ 1)/8)**4 + 8*int(tan((4*x**2 + 4*x + 1)/8)/(tan((4*x**2 + 4*x + 1)/8)**4
+ 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x)*tan((4*x**2 + 4*x + 1)/8)**2 + 4
*int(tan((4*x**2 + 4*x + 1)/8)/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x*
*2 + 4*x + 1)/8)**2 + 1),x) - int(1/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan(
(4*x**2 + 4*x + 1)/8)**2 + 1),x)*tan((4*x**2 + 4*x + 1)/8)**4 - 2*int(1/(t
an((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x)*tan((
4*x**2 + 4*x + 1)/8)**2 - int(1/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x
**2 + 4*x + 1)/8)**2 + 1),x) - 4*tan((4*x**2 + 4*x + 1)/8)*x + 2*tan((4*x*
*2 + 4*x + 1)/8))/(3*(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x
+ 1)/8)**2 + 1))
```

3.25 $\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [A] (verified)	191
Fricas [A] (verification not implemented)	191
Sympy [B] (verification not implemented)	192
Maxima [C] (verification not implemented)	192
Giac [C] (verification not implemented)	193
Mupad [F(-1)]	193
Reduce [F]	194

Optimal result

Integrand size = 13, antiderivative size = 46

$$\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x^2}{4} + \frac{1}{8}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) - \frac{1}{8} \sin \left(\frac{1}{2} + 2x + 2x^2 \right)$$

output 1/4*x^2+1/8*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))-1/8*sin(1/2+2*x+2*x^2)

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.91

$$\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{8} \left(2x^2 + \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) - \sin \left(\frac{1}{2}(1+2x)^2 \right) \right)$$

input Integrate[x*Sin[1/4 + x + x^2]^2,x]

output (2*x^2 + Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]] - Sin[(1 + 2*x)^2/2])/8

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \sin^2 \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \text{3948} \\
 & \int \left(\frac{x}{2} - \frac{1}{2}x \cos \left(2x^2 + 2x + \frac{1}{2} \right) \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{8}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x^2}{4} - \frac{1}{8} \sin \left(2x^2 + 2x + \frac{1}{2} \right)
 \end{aligned}$$

input `Int[x*Sin[1/4 + x + x^2]^2,x]`

output `x^2/4 + (Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/8 - Sin[1/2 + 2*x + 2*x^2]/8`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_)*(x_))^(m_)*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.43 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x^2}{4} + \frac{\sqrt{\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{8} - \frac{\sin\left(\frac{1}{2}+2x+2x^2\right)}{8}$	35
risch	$-\frac{\sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \operatorname{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x+\frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{32} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-2i} x-\frac{i}{\sqrt{-2i}}\right)}{16\sqrt{-2i}} + \frac{x^2}{4} - \frac{\sin\left(\frac{(1+2x)^2}{2}\right)}{8}$	72

input `int(x*sin(1/4+x+x^2)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^2+1/8*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))-1/8*sin(1/2+2*x+2*x^2)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\begin{aligned} \int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx &= \frac{1}{4} x^2 - \frac{1}{4} \cos \left(x^2 + x + \frac{1}{4} \right) \sin \left(x^2 + x + \frac{1}{4} \right) \\ &\quad + \frac{1}{8} \sqrt{\pi} C \left(\frac{2x+1}{\sqrt{\pi}} \right) \end{aligned}$$

input `integrate(x*sin(1/4+x+x^2)^2,x, algorithm="fricas")`

output `1/4*x^2 - 1/4*cos(x^2 + x + 1/4)*sin(x^2 + x + 1/4) + 1/8*sqrt(pi)*fresnel
_cos((2*x + 1)/sqrt(pi))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(37) = 74$.

Time = 0.80 (sec), antiderivative size = 121, normalized size of antiderivative = 2.63

$$\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x^2}{4} - \frac{\sqrt{\pi} x C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right)}{4} + \frac{\sqrt{\pi} x C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right) \Gamma \left(\frac{1}{4} \right)}{16 \Gamma \left(\frac{5}{4} \right)} \\ - \frac{\sin \left(2(x + \frac{1}{2})^2 \right) \Gamma \left(\frac{1}{4} \right)}{32 \Gamma \left(\frac{5}{4} \right)} + \frac{\sqrt{\pi} C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right) \Gamma \left(\frac{1}{4} \right)}{32 \Gamma \left(\frac{5}{4} \right)}$$

input `integrate(x*sin(1/4+x+x**2)**2,x)`

output `x**2/4 - sqrt(pi)*x*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))/4 + sqrt(pi)*x*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))*gamma(1/4)/(16*gamma(5/4)) - sin(2*(x + 1/2)**2)*gamma(1/4)/(32*gamma(5/4)) + sqrt(pi)*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))*gamma(1/4)/(32*gamma(5/4))`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec), antiderivative size = 137, normalized size of antiderivative = 2.98

$$\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx \\ = \frac{32 x^3 + 16 x^2 - 8 x \left(-i e^{(2i x^2 + 2i x + \frac{1}{2}i)} + i e^{(-2i x^2 - 2i x - \frac{1}{2}i)} \right) - \sqrt{8 x^2 + 8 x + 2} \left((i - 1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{2i x^2 + 2i x + 1} \right) - \operatorname{erf} \left(\sqrt{-2i x^2 - 2i x - 1} \right) \right) + 4 \operatorname{erf} \left(\sqrt{2i x^2 + 2i x + 1} \right) - 4 \operatorname{erf} \left(\sqrt{-2i x^2 - 2i x - 1} \right) \right)}{64}$$

input `integrate(x*sin(1/4+x+x^2)^2,x, algorithm="maxima")`

output `1/64*(32*x^3 + 16*x^2 - 8*x*(-I*e^(2*I*x^2 + 2*I*x + 1/2*I) + I*e^(-2*I*x^2 - 2*I*x - 1/2*I)) - sqrt(8*x^2 + 8*x + 2)*((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(2*I*x^2 + 2*I*x + 1/2*I)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(-2*I*x^2 - 2*I*x - 1/2*I)) - 1)) + 4*I*e^(2*I*x^2 + 2*I*x + 1/2*I) - 4*I*e^(-2*I*x^2 - 2*I*x - 1/2*I))/(2*x + 1)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17

$$\begin{aligned} \int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx &= \frac{1}{4} x^2 - \left(\frac{1}{32}i + \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) \\ &\quad + \left(\frac{1}{32}i - \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right) \\ &\quad + \frac{1}{16}i e^{(2ix^2+2ix+\frac{1}{2}i)} - \frac{1}{16}i e^{(-2ix^2-2ix-\frac{1}{2}i)} \end{aligned}$$

input `integrate(x*sin(1/4+x+x^2)^2,x, algorithm="giac")`

output
$$\begin{aligned} &1/4*x^2 - (1/32*I + 1/32)*sqrt(pi)*erf((I - 1)*x + 1/2*I - 1/2) + (1/32*I \\ &- 1/32)*sqrt(pi)*erf(-(I + 1)*x - 1/2*I - 1/2) + 1/16*I*e^{(2*I*x^2 + 2*I*x \\ &+ 1/2*I)} - 1/16*I*e^{(-2*I*x^2 - 2*I*x - 1/2*I)} \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x \sin \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(x*sin(x + x^2 + 1/4)^2,x)`

output `int(x*sin(x + x^2 + 1/4)^2, x)`

Reduce [F]

$$\int x \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{\cos(x^2 + x + \frac{1}{4}) \sin(x^2 + x + \frac{1}{4})}{4} \\ - \frac{\left(\int \sin(x^2 + x + \frac{1}{4})^2 dx \right)}{2} + \frac{x^2}{4} + \frac{x}{4}$$

input `int(x*sin(1/4+x+x^2)^2,x)`

output `(- cos((4*x**2 + 4*x + 1)/4)*sin((4*x**2 + 4*x + 1)/4) - 2*int(sin((4*x**2 + 4*x + 1)/4)**2,x) + x**2 + x)/4`

3.26 $\int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	195
Mathematica [A] (verified)	195
Rubi [A] (verified)	196
Maple [A] (verified)	197
Fricas [A] (verification not implemented)	197
Sympy [A] (verification not implemented)	197
Maxima [C] (verification not implemented)	198
Giac [C] (verification not implemented)	198
Mupad [F(-1)]	199
Reduce [F]	199

Optimal result

Integrand size = 11, antiderivative size = 27

$$\int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x}{2} - \frac{1}{4}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right)$$

output `1/2*x-1/4*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \left(2x - \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) \right)$$

input `Integrate[Sin[1/4 + x + x^2]^2, x]`

output `(2*x - Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/4`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3930, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2 \left(x^2 + x + \frac{1}{4} \right) dx \\ & \quad \downarrow \text{3930} \\ & \int \left(\frac{1}{2} - \frac{1}{2} \cos \left(2x^2 + 2x + \frac{1}{2} \right) \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{x}{2} - \frac{1}{4} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) \end{aligned}$$

input `Int[Sin[1/4 + x + x^2]^2, x]`

output `x/2 - (Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/4`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3930 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTrigReduce[Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{x}{2} - \frac{\sqrt{\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{4}$	20
risch	$\frac{x}{2} + \frac{\sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \operatorname{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{16} - \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-2i} x - \frac{i}{\sqrt{-2i}}\right)}{8\sqrt{-2i}}$	58

input `int(sin(1/4+x+x^2)^2, x, method=_RETURNVERBOSE)`

output `1/2*x-1/4*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \sin^2\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{4} \sqrt{\pi} C\left(\frac{2x+1}{\sqrt{\pi}}\right) + \frac{1}{2} x$$

input `integrate(sin(1/4+x+x^2)^2, x, algorithm="fricas")`

output `-1/4*sqrt(pi)*fresnel_cos((2*x + 1)/sqrt(pi)) + 1/2*x`

Sympy [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \sin^2\left(\frac{1}{4} + x + x^2\right) dx = \frac{x}{2} - \frac{\sqrt{\pi} C\left(\frac{4x+2}{2\sqrt{\pi}}\right)}{4}$$

input `integrate(sin(1/4+x+x**2)**2, x)`

output $x/2 - \sqrt{\pi} * \text{fresnelc}((4*x + 2)/(2*\sqrt{\pi}))/4$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{16} \sqrt{\pi} \left((i-1) \operatorname{erf} \left(\frac{2ix+i}{\sqrt{2i}} \right) + (i+1) \operatorname{erf} \left(\frac{2ix+i}{\sqrt{-2i}} \right) \right) + \frac{1}{2} x$$

input `integrate(sin(1/4+x+x^2)^2,x, algorithm="maxima")`

output $\frac{1}{16}\sqrt{\pi}((I-1)\operatorname{erf}((2Ix+I)/\sqrt{2I}) + (I+1)\operatorname{erf}((2Ix+I)/\sqrt{-2I})) + 1/2*x$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\begin{aligned} \int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx &= \left(\frac{1}{16}i + \frac{1}{16} \right) \sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) \\ &\quad - \left(\frac{1}{16}i - \frac{1}{16} \right) \sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right) + \frac{1}{2}x \end{aligned}$$

input `integrate(sin(1/4+x+x^2)^2,x, algorithm="giac")`

output $(1/16*I + 1/16)*\sqrt{\pi}*\operatorname{erf}((I-1)*x + 1/2*I - 1/2) - (1/16*I - 1/16)*\sqrt{\pi}*\operatorname{erf}(-(I+1)*x - 1/2*I - 1/2) + 1/2*x$

Mupad [F(-1)]

Timed out.

$$\int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int \sin \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(sin(x + x^2 + 1/4)^2,x)`

output `int(sin(x + x^2 + 1/4)^2, x)`

Reduce [F]

$$\int \sin^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int \sin \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(sin(1/4+x+x^2)^2,x)`

output `int(sin((4*x**2 + 4*x + 1)/4)**2,x)`

3.27 $\int \frac{\sin^2\left(\frac{1}{4}+x+x^2\right)}{x} dx$

Optimal result	200
Mathematica [N/A]	200
Rubi [N/A]	201
Maple [N/A]	202
Fricas [N/A]	202
Sympy [N/A]	202
Maxima [N/A]	203
Giac [N/A]	203
Mupad [N/A]	204
Reduce [N/A]	204

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \frac{\log(x)}{2} - \frac{1}{2} \text{Int}\left(\frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{x}, x\right)$$

output `1/2*ln(x)-1/2*DefeR(Int)(cos(1/2+2*x+2*x^2)/x,x)`

Mathematica [N/A]

Not integrable

Time = 10.69 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

input `Integrate[Sin[1/4 + x + x^2]^2/x, x]`

output `Integrate[Sin[1/4 + x + x^2]^2/x, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin^2(x^2 + x + \frac{1}{4})}{x} dx \\ & \quad \downarrow \text{3948} \\ & \int \left(\frac{1}{2x} - \frac{\cos(2x^2 + 2x + \frac{1}{2})}{2x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{\log(x)}{2} - \frac{1}{2} \int \frac{\cos(2x^2 + 2x + \frac{1}{2})}{x} dx \end{aligned}$$

input `Int[Sin[1/4 + x + x^2]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_)*(x_))^(m_.)*Sin[(a_.) + (b_)*(x_) + (c_)*(x_)^2]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\sin\left(\frac{1}{4} + x + x^2\right)^2}{x} dx$$

input `int(sin(1/4+x+x^2)^2/x, x)`

output `int(sin(1/4+x+x^2)^2/x, x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

input `integrate(sin(1/4+x+x^2)^2/x, x, algorithm="fricas")`

output `integral(-(cos(x^2 + x + 1/4)^2 - 1)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.63 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin^2\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(sin(1/4+x+x**2)**2/x, x)`

output `Integral(sin(x**2 + x + 1/4)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

$$\int \frac{\sin^2 \left(\frac{1}{4} + x + x^2 \right)}{x} dx = \int \frac{\sin \left(x^2 + x + \frac{1}{4} \right)^2}{x} dx$$

input `integrate(sin(1/4+x+x^2)^2/x, x, algorithm="maxima")`

output `-1/2*integrate(cos(2*x^2 + 2*x + 1/2)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2 \left(\frac{1}{4} + x + x^2 \right)}{x} dx = \int \frac{\sin \left(x^2 + x + \frac{1}{4} \right)^2}{x} dx$$

input `integrate(sin(1/4+x+x^2)^2/x, x, algorithm="giac")`

output `integrate(sin(x^2 + x + 1/4)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 38.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

input `int(sin(x + x^2 + 1/4)^2/x,x)`

output `int(sin(x + x^2 + 1/4)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

input `int(sin(1/4+x+x^2)^2/x,x)`

output `int(sin((4*x**2 + 4*x + 1)/4)**2/x,x)`

3.28 $\int \frac{\sin^2\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$

Optimal result	205
Mathematica [N/A]	205
Rubi [N/A]	206
Maple [N/A]	207
Fricas [N/A]	207
Sympy [N/A]	208
Maxima [N/A]	208
Giac [N/A]	208
Mupad [N/A]	209
Reduce [N/A]	209

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = -\frac{1}{2x} + \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} + \sqrt{\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) \\ + \operatorname{Int}\left(\frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{x}, x\right)$$

output
$$-1/2/x+1/2*\cos(1/2+2*x+2*x^2)/x+\text{Pi}^{(1/2)}*\text{FresnelS}((1+2*x)/\text{Pi}^{(1/2)})+\text{Defer}(\text{Int})(\sin(1/2+2*x+2*x^2)/x,x)$$

Mathematica [N/A]

Not integrable

Time = 10.99 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

input
$$\text{Integrate}[\text{Sin}[1/4 + x + x^2]^2/x^2, x]$$

output `Integrate[Sin[1/4 + x + x^2]^2/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^2(x^2 + x + \frac{1}{4})}{x^2} dx \\
 & \quad \downarrow \text{3948} \\
 & \int \left(\frac{1}{2x^2} - \frac{\cos(2x^2 + 2x + \frac{1}{2})}{2x^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \int \frac{\sin(2x^2 + 2x + \frac{1}{2})}{x} dx + \sqrt{\pi} \operatorname{FresnelS}\left(\frac{2x+1}{\sqrt{\pi}}\right) + \frac{\cos(2x^2 + 2x + \frac{1}{2})}{2x} - \frac{1}{2x}
 \end{aligned}$$

input `Int[Sin[1/4 + x + x^2]^2/x^2,x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3948 $\text{Int}[(d_.) + (e_.)*(x_.))^(m_.)*\text{Sin}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \ \text{Sin}[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c, \ d, \ e, \ m\}, \ x] \ \&& \ \text{IGtQ}[n, \ 1]$

Maple [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\sin \left(\frac{1}{4} + x + x^2\right)^2}{x^2} dx$$

input `int(sin(1/4+x+x^2)^2/x^2,x)`

output `int(sin(1/4+x+x^2)^2/x^2,x)`

Fricas [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \frac{\sin^2 \left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin \left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

input `integrate(sin(1/4+x+x^2)^2/x^2,x, algorithm="fricas")`

output `integral(-(cos(x^2 + x + 1/4)^2 - 1)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.54 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin^2\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(sin(1/4+x+x**2)**2/x**2, x)`

output `Integral(sin(x**2 + x + 1/4)**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

input `integrate(sin(1/4+x+x^2)^2/x^2, x, algorithm="maxima")`

output `-1/2*(x*integrate(cos(2*x^2 + 2*x + 1/2)/x^2, x) + 1)/x`

Giac [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\sin\left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

input `integrate(sin(1/4+x+x^2)^2/x^2, x, algorithm="giac")`

output `integrate(sin(x^2 + x + 1/4)^2/x^2, x)`

Mupad [N/A]

Not integrable

Time = 38.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2 \left(\frac{1}{4} + x + x^2 \right)}{x^2} dx = \int \frac{\sin \left(x^2 + x + \frac{1}{4} \right)^2}{x^2} dx$$

input `int(sin(x + x^2 + 1/4)^2/x^2,x)`

output `int(sin(x + x^2 + 1/4)^2/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2 \left(\frac{1}{4} + x + x^2 \right)}{x^2} dx = \int \frac{\sin \left(x^2 + x + \frac{1}{4} \right)^2}{x^2} dx$$

input `int(sin(1/4+x+x^2)^2/x^2,x)`

output `int(sin((4*x**2 + 4*x + 1)/4)**2/x**2,x)`

3.29 $\int (d + ex)^2 \sin(a + bx + cx^2) dx$

Optimal result	210
Mathematica [A] (verified)	211
Rubi [A] (verified)	211
Maple [A] (verified)	215
Fricas [A] (verification not implemented)	215
Sympy [F]	216
Maxima [C] (verification not implemented)	216
Giac [C] (verification not implemented)	217
Mupad [F(-1)]	218
Reduce [F]	218

Optimal result

Integrand size = 19, antiderivative size = 285

$$\begin{aligned} \int (d + ex)^2 \sin(a + bx + cx^2) dx = & -\frac{e(2cd - be) \cos(a + bx + cx^2)}{4c^2} \\ & - \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\ & + \frac{e^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & + \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & + \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\ & - \frac{e^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \end{aligned}$$

output

```
-1/4*e*(-b*e+2*c*d)*cos(c*x^2+b*x+a)/c^2-1/2*e*(e*x+d)*cos(c*x^2+b*x+a)/c+
1/4*e^2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)+1/8*(-b*e+2*c*d)^2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(5/2)+1/8*(-b*e+2*c*d)^2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(5/2)-1/4*e^2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)
```

Mathematica [A] (verified)

Time = 1.39 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.65

$$\int (d + ex)^2 \sin(a + bx + cx^2) dx \\ = \frac{2\sqrt{c}e(be - 2c(2d + ex)) \cos(a + x(b + cx)) + \sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left((-2cd + be)^2 \cos\left(a - \frac{b^2}{4c}\right) - 2ce^2\right)}{8c^{5/2}}$$

input `Integrate[(d + e*x)^2*Sin[a + b*x + c*x^2], x]`

output
$$(2*\text{Sqrt}[c]*e*(b*e - 2*c*(2*d + e*x))*\text{Cos}[a + x*(b + c*x)] + \text{Sqrt}[2*\pi]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])]*(((-2*c*d + b*e)^2*\text{Cos}[a - b^2/(4*c)]) - 2*c*e^2*\text{Sin}[a - b^2/(4*c)]) + \text{Sqrt}[2*\pi]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\pi])]*(2*c*e^2*\text{Cos}[a - b^2/(4*c)] + (-2*c*d + b*e)^2*\text{Sin}[a - b^2/(4*c)]))/(8*c^{5/2})$$

Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 282, normalized size of antiderivative = 0.99, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3944, 3929, 3832, 3833, 3942, 3928, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 \sin(a + bx + cx^2) dx \\ \downarrow 3944 \\ \frac{(2cd - be) \int (d + ex) \sin(cx^2 + bx + a) dx}{2c} + \frac{e^2 \int \cos(cx^2 + bx + a) dx}{2c} - \\ \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\ \downarrow 3929$$

$$\begin{aligned}
& \frac{e^2 \left(\cos \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx - \sin \left(a - \frac{b^2}{4c} \right) \int \sin \left(\frac{(b+2cx)^2}{4c} \right) dx \right)}{2c} + \\
& \frac{(2cd - be) \int (d + ex) \sin(cx^2 + bx + a) dx}{2c} - \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\
& \quad \downarrow \textcolor{blue}{3832} \\
& \frac{e^2 \left(\cos \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} + \\
& \frac{(2cd - be) \int (d + ex) \sin(cx^2 + bx + a) dx}{2c} - \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\
& \quad \downarrow \textcolor{blue}{3833} \\
& \frac{(2cd - be) \int (d + ex) \sin(cx^2 + bx + a) dx}{2c} + \\
& \frac{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} - \\
& \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\
& \quad \downarrow \textcolor{blue}{3942} \\
& \frac{(2cd - be) \left(\frac{(2cd - be) \int \sin(cx^2 + bx + a) dx}{2c} - \frac{e \cos(a + bx + cx^2)}{2c} \right)}{2c} + \\
& \frac{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} - \\
& \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\
& \quad \downarrow \textcolor{blue}{3928} \\
& \frac{(2cd - be) \left(\frac{(2cd - be) \left(\sin \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx + \cos \left(a - \frac{b^2}{4c} \right) \int \sin \left(\frac{(b+2cx)^2}{4c} \right) dx \right)}{2c} - \frac{e \cos(a + bx + cx^2)}{2c} \right)}{2c} + \\
& \frac{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} - \\
& \frac{e(d + ex) \cos(a + bx + cx^2)}{2c} \\
& \quad \downarrow \textcolor{blue}{3832}
\end{aligned}$$

$$\begin{aligned}
 & \frac{(2cd-be) \left(\frac{(2cd-be) \left(\sin(a - \frac{b^2}{4c}) \int \cos(\frac{(b+2cx)^2}{4c}) dx + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} \right)}{2c} - \frac{e \cos(a+bx+cx^2)}{2c} \right)}{2c} + \\
 & \frac{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelC}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} \right)}{2c} - \\
 & \frac{e(d+ex) \cos(a+bx+cx^2)}{2c} \\
 & \quad \downarrow \text{3833} \\
 & \frac{(2cd-be) \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelC}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}) + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}}}{2c} - \frac{e \cos(a+bx+cx^2)}{2c} \right)}{2c} + \\
 & \frac{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelC}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} \right)}{2c} - \\
 & \frac{e(d+ex) \cos(a+bx+cx^2)}{2c}
 \end{aligned}$$

input `Int[(d + e*x)^2*Sin[a + b*x + c*x^2],x]`

output

$$\begin{aligned}
 & -1/2*(e*(d + e*x)*Cos[a + b*x + c*x^2])/c + (e^2*((\operatorname{Sqrt}[\operatorname{Pi}/2]*Cos[a - b^2/(4*c)]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}]])]/\operatorname{Sqrt}[c] - (\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]*)\operatorname{Sin}[a - b^2/(4*c)])/\operatorname{Sqrt}[c]))/(2*c) + ((2*c*d - b*e)*(-1/2*(e*Cos[a + b*x + c*x^2])/c + ((2*c*d - b*e)*((\operatorname{Sqrt}[\operatorname{Pi}/2]*Cos[a - b^2/(4*c)]*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}]])]/\operatorname{Sqrt}[c] + (\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]*)\operatorname{Sin}[a - b^2/(4*c)])/\operatorname{Sqrt}[c])))/(2*c))
 \end{aligned}$$

Definitions of rubi rules used

rule 3832 $\text{Int}[\text{Sin}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\text{Int}[\text{Cos}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\text{Int}[\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3929 $\text{Int}[\text{Cos}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] + \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3942 $\text{Int}[((d_{_}) + (e_{_})*(x_{_}))*\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-e)*(\text{Cos}[a + b*x + c*x^2]/(2*c)), x] + \text{Simp}[(2*c*d - b*e)/(2*c) \quad \text{Int}[\text{Sin}[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

rule 3944 $\text{Int}[((d_{_}) + (e_{_})*(x_{_}))^{(m_{_})}*\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-e)*(d + e*x)^{(m - 1)}*(\text{Cos}[a + b*x + c*x^2]/(2*c)), x] + (-\text{Simp}[(b*e - 2*c*d)/(2*c) \quad \text{Int}[(d + e*x)^{(m - 1)}*\text{Sin}[a + b*x + c*x^2], x], x] + \text{Simp}[e^{2*(m - 1)/(2*c)} \quad \text{Int}[(d + e*x)^{(m - 2)}*\text{Cos}[a + b*x + c*x^2], x], x]) /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[b*e - 2*c*d, 0] \&& \text{GtQ}[m, 1]$

Maple [A] (verified)

Time = 1.88 (sec) , antiderivative size = 399, normalized size of antiderivative = 1.40

method	result
default	$\frac{e^2 b \left(-\frac{\cos(c x^2 + b x + a)}{2 c} - \frac{b \sqrt{2} \sqrt{\pi} \left(\cos\left(\frac{b^2 - ac}{4 c}\right) \text{FresnelS}\left(\frac{\sqrt{2} \left(c x + \frac{b}{2}\right)}{\sqrt{\pi} \sqrt{c}}\right) - \sin\left(\frac{b^2 - ac}{4 c}\right) \text{FresnelC}\left(\frac{\sqrt{2} \left(c x + \frac{b}{2}\right)}{\sqrt{\pi} \sqrt{c}}\right)\right)}{4 c^{\frac{3}{2}}} \right)}{2 c}$
risch	$\frac{i \operatorname{erf}\left(-\sqrt{-i c} x+\frac{i b}{2 \sqrt{-i c}}\right) \sqrt{\pi} d^2 e^{\frac{i (4 a c-b^2)}{4 c}}}{4 \sqrt{-i c}}+\frac{i e^2 \operatorname{erf}\left(-\sqrt{-i c} x+\frac{i b}{2 \sqrt{-i c}}\right) \sqrt{\pi} b^2 e^{\frac{i (4 a c-b^2)}{4 c}}}{16 \sqrt{-i c} c^2}-\frac{e^2 \operatorname{erf}\left(-\sqrt{-i c} x+\frac{i b}{2 \sqrt{-i c}}\right) \sqrt{\pi} e^{\frac{i (4 a c-b^2)}{4 c}}}{8 \sqrt{-i c} c}$
parts	Expression too large to display

input `int((e*x+d)^2*sin(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -\frac{1}{2} e^2 c x^2 \cos(c x^2 + b x + a) - \frac{1}{2} e^2 b c (-\frac{1}{2} \cos(c x^2 + b x + a) / c - \frac{1}{4} b / c \\ & + \frac{3}{2} * 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4 * b^2 - a * c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b)) - \sin((1/4 * b^2 - a * c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b))) + \frac{1}{4} e^2 c^3 / 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4 * b^2 - a * c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b)) + \sin((1/4 * b^2 - a * c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b))) - d * e / c * \cos(c * x^2 + b * x + a) - \frac{1}{2} d * e * b / c^2 \\ & + \frac{3}{2} * 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4 * b^2 - a * c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b)) - \sin((1/4 * b^2 - a * c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b))) + \frac{1}{2} * 2^{(1/2)} * \text{Pi}^{(1/2)} / c^{(1/2)} * d^2 * (\cos((1/4 * b^2 - a * c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b)) - \sin((1/4 * b^2 - a * c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c * x + 1/2 * b))) \end{aligned}$$
Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int (d + ex)^2 \sin(a + bx + cx^2) \, dx \\ & = \frac{\sqrt{2} \left(2 \pi c e^2 \cos\left(-\frac{b^2 - 4ac}{4c}\right) + \pi (4c^2 d^2 - 4bcde + b^2 e^2) \sin\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2} \left(2 \pi c e^2 \cos\left(-\frac{b^2 - 4ac}{4c}\right) + \pi (4c^2 d^2 - 4bcde + b^2 e^2) \sin\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right)}{2c} \end{aligned}$$

input `integrate((e*x+d)^2*sin(c*x^2+b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{8} \left(\sqrt{2} \left(2\pi c e^2 \cos(-\frac{1}{4}(b^2 - 4ac)/c) + \pi (4c^2 d^2 - 4bc^2 d e + b^2 e^2) \sin(-\frac{1}{4}(b^2 - 4ac)/c) \right) \sqrt{c/\pi} \operatorname{fresnel_cos}(1/2 \sqrt{2} \sqrt{c} x) - \sqrt{2} \left(2\pi c e^2 \sin(-\frac{1}{4}(b^2 - 4ac)/c) \right) \sqrt{c/\pi} \operatorname{fresnel_sin}(1/2 \sqrt{2} \sqrt{c} x) - 2(2c^2 e^2 - 4c^2 d e - b c e^2) \cos(c x^2 + bx + a) \right) / c^3$$

Sympy [F]

$$\int (d + ex)^2 \sin(a + bx + cx^2) dx = \int (d + ex)^2 \sin(a + bx + cx^2) dx$$

input `integrate((e*x+d)**2*sin(c*x**2+b*x+a),x)`

output `Integral((d + e*x)**2*sin(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.14 (sec) , antiderivative size = 2269, normalized size of antiderivative = 7.96

$$\int (d + ex)^2 \sin(a + bx + cx^2) dx = \text{Too large to display}$$

input `integrate((e*x+d)^2*sin(c*x^2+b*x+a),x, algorithm="maxima")`

output

```

-1/8*sqrt(2)*sqrt(pi)*((-I + 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(I*c)) + (-I - 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(-I*c)))*d^2/sqrt(c) + 1/8*((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/4*(b^2 - 4*a*c)/c) + ((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/4*(b^2 - 4*a*c)/c) - 2*((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/4*(b^2 - 4*a*c)/c) + (-I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/4*(b^2 - 4*a*c)/c))*x - 4*(c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^2 - 4*a*c)/c) + c*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/4*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2/c))*d/e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2/c))) + 1/32*(8*(((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4...

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec), antiderivative size = 289, normalized size of antiderivative = 1.01

$$\begin{aligned}
 & \int (d + ex)^2 \sin(a + bx + cx^2) \, dx = \\
 & -\frac{\sqrt{2}\sqrt{\pi}(-4ic^2d^2 + 4ibcde - ib^2e^2 + 2ce^2)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x + \frac{b}{c}\right)\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2 - 4icac}{4c}\right)}}{\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}} + 2\left(ce^2\left(2x + \frac{b}{c}\right) + 4cde - 2be^2\right) \\
 & -\frac{\sqrt{2}\sqrt{\pi}(4ic^2d^2 - 4ibcde + ib^2e^2 + 2ce^2)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x + \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2 + 4icac}{4c}\right)}}{\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}} + 2\left(ce^2\left(2x + \frac{b}{c}\right) + 4cde - 2be^2\right) \\
 & \quad 16c^2
 \end{aligned}$$

input `integrate((e*x+d)^2*sin(c*x^2+b*x+a),x, algorithm="giac")`

output

```

-1/16*(sqrt(2)*sqrt(pi)*(-4*I*c^2*d^2 + 4*I*b*c*d*e - I*b^2*e^2 + 2*c*e^2)
*erf(-1/4*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b
^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) + 2*(c*e^2*(2*x + b/c) +
4*c*d*e - 2*b*e^2)*e^(I*c*x^2 + I*b*x + I*a))/c^2 - 1/16*(sqrt(2)*sqrt(pi)
)*(4*I*c^2*d^2 - 4*I*b*c*d*e + I*b^2*e^2 + 2*c*e^2)*erf(-1/4*sqrt(2)*(2*x
+ b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((I*c/
abs(c) + 1)*sqrt(abs(c))) + 2*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^(-
I*c*x^2 - I*b*x - I*a))/c^2

```

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 \sin(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a) (d + ex)^2 dx$$

input

```
int(sin(a + b*x + c*x^2)*(d + e*x)^2,x)
```

output

```
int(sin(a + b*x + c*x^2)*(d + e*x)^2, x)
```

Reduce [F]

$$\begin{aligned}
& \int (d + ex)^2 \sin(a + bx + cx^2) dx \\
&= \frac{4b^2c^2e^2x^3 - 24 \left(\int \frac{x^2}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) b^2c^2e^2 + 24 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) b^2c^2d^2 + 6 \cos(cx^2 + bx + a)}{1}
\end{aligned}$$

input

```
int((e*x+d)^2*sin(c*x^2+b*x+a),x)
```

output

```
(6*cos(a + b*x + c*x**2)*b*c*e**2 - 24*cos(a + b*x + c*x**2)*c**2*d*e - 12
*cos(a + b*x + c*x**2)*c**2*e**2*x - 24*int(x**2/(tan((a + b*x + c*x**2)/2
)**2 + 1),x)*b**2*c**2*e**2 + 96*int(x**2/(tan((a + b*x + c*x**2)/2)**2 +
1),x)*b*c**3*d*e - 96*int(x**2/(tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**4*
d**2 + 6*int(1/(tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**4*e**2 - 24*int(1/
(tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**3*c*d*e + 24*int(1/(tan((a + b*x
+ c*x**2)/2)**2 + 1),x)*b**2*c**2*d**2 + 24*int(1/(tan((a + b*x + c*x**2)/
2)**2 + 1),x)*c**2*e**2 - 3*sin(a + b*x + c*x**2)*b**3*e**2 + 12*sin(a + b
*x + c*x**2)*b**2*c*d*e + 6*sin(a + b*x + c*x**2)*b**2*c*e**2*x - 12*sin(a
+ b*x + c*x**2)*b*c**2*d**2 - 24*sin(a + b*x + c*x**2)*b*c**2*d*e*x + 24*
sin(a + b*x + c*x**2)*c**3*d**2*x - 3*b**4*e**2*x + 12*b**3*c*d*e*x - 12*b
**2*c**2*d**2*x + 4*b**2*c**2*e**2*x**3 - 16*b*c**3*d*e*x**3 - 6*b*c*e**2
+ 16*c**4*d**2*x**3 + 24*c**2*d*e - 12*c**2*e**2*x)/(24*c**3)
```

3.30 $\int (d + ex) \sin(a + bx + cx^2) dx$

Optimal result	220
Mathematica [A] (verified)	221
Rubi [A] (verified)	221
Maple [A] (verified)	223
Fricas [A] (verification not implemented)	224
Sympy [F]	224
Maxima [C] (verification not implemented)	224
Giac [C] (verification not implemented)	225
Mupad [F(-1)]	226
Reduce [F]	226

Optimal result

Integrand size = 17, antiderivative size = 140

$$\begin{aligned} \int (d + ex) \sin(a + bx + cx^2) dx &= -\frac{e \cos(a + bx + cx^2)}{2c} \\ &+ \frac{(2cd - be)\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &+ \frac{(2cd - be)\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \end{aligned}$$

output

```
-1/2*e*cos(c*x^2+b*x+a)/c+1/4*(-b*e+2*c*d)*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)+1/4*(-b*e+2*c*d)*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)
```

Mathematica [A] (verified)

Time = 0.94 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.91

$$\int (d + ex) \sin(a + bx + cx^2) dx \\ = \frac{-2\sqrt{c}e \cos(a + x(b + cx)) + (2cd - be)\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + (2cd - be)\sqrt{2\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{3/2}}$$

input `Integrate[(d + e*x)*Sin[a + b*x + c*x^2], x]`

output
$$\frac{(-2\sqrt{c}e \cos(a + x(b + cx)) + (2cd - be)\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + (2cd - be)\sqrt{2\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)) \sin\left(a - \frac{b^2}{4c}\right)}{(4c)^{3/2}}$$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3942, 3928, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) \sin(a + bx + cx^2) dx \\ \downarrow 3942 \\ \frac{(2cd - be) \int \sin(cx^2 + bx + a) dx}{2c} - \frac{e \cos(a + bx + cx^2)}{2c} \\ \downarrow 3928 \\ \frac{(2cd - be) \left(\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx + \cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx \right)}{2c} - \frac{e \cos(a + bx + cx^2)}{2c}$$

$$\begin{array}{c}
 \downarrow \text{3832} \\
 \frac{(2cd - be) \left(\sin \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{e \cos \left(a + bx + cx^2 \right)} - \\
 \frac{2c}{2c} \\
 \downarrow \text{3833} \\
 \frac{(2cd - be) \left(\frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{e \cos \left(a + bx + cx^2 \right)} - \\
 \frac{2c}{2c}
 \end{array}$$

input `Int[(d + e*x)*Sin[a + b*x + c*x^2], x]`

output `-1/2*(e*Cos[a + b*x + c*x^2])/c + ((2*c*d - b*e)*((Sqrt[Pi/2]*Cos[a - b^2/(4*c)])*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/Sqrt[c] + (Sqrt[Pi/2]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*Sin[a - b^2/(4*c)])/Sqrt[c]))/(2*c)`

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_.))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3928 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] - Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3942

```

Int[((d_) + (e_)*(x_))*Sin[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol]
:> Simp[(-e)*(Cos[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c)
Int[Sin[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d
- b*e, 0]

```

Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.31

method	result
default	$-\frac{e \cos(cx^2+bx+a)}{2c} - \frac{eb\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2-ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right) - \sin\left(\frac{b^2-ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)\right)}{4c^{3/2}} + \frac{\sqrt{2}\sqrt{\pi}d}{c}$
risch	$\frac{i \operatorname{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)\sqrt{\pi} de^{\frac{i(4ac-b^2)}{4c}}}{4\sqrt{-ic}} - \frac{ie \operatorname{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)\sqrt{\pi} b e^{\frac{i(4ac-b^2)}{4c}}}{8\sqrt{-ic}c} + \frac{i \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi} de^{-\frac{i(4ac-b^2)}{4c}}}{4\sqrt{ic}}$
parts	$\frac{\sqrt{2}\sqrt{\pi} \cos\left(\frac{b^2-ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right) ex}{2\sqrt{c}} - \frac{\sqrt{2}\sqrt{\pi} \sin\left(\frac{b^2-ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right) ex}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi} \cos\left(\frac{b^2-ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right) ex}{2\sqrt{c}} - \frac{\sqrt{2}\sqrt{\pi} \sin\left(\frac{b^2-ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right) ex}{2\sqrt{c}}$

input

```
int((e*x+d)*sin(c*x^2+b*x+a),x,method=_RETURNVERBOSE)
```

output

```

-1/2*e*cos(c*x^2+b*x+a)/c-1/4*e*b/c^(3/2)*2^(1/2)*Pi^(1/2)*(cos((1/4*b^2-a
*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))-sin((1/4*b^2-a*c)/c
)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))+1/2*2^(1/2)*Pi^(1/2)/c^(1
/2)*d*(cos((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))
-sin((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))

```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.96

$$\int (d + ex) \sin(a + bx + cx^2) dx = \frac{\sqrt{2}\pi(2cd - be)\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi(2cd - be)\sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2 - 4ac}{4c}\right)}{4c^2}$$

input `integrate((e*x+d)*sin(c*x^2+b*x+a),x, algorithm="fricas")`

output `1/4*(sqrt(2)*pi*(2*c*d - b*e)*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) + sqrt(2)*pi*(2*c*d - b*e)*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c) - 2*c*e*cos(c*x^2 + b*x + a))/c^2`

Sympy [F]

$$\int (d + ex) \sin(a + bx + cx^2) dx = \int (d + ex) \sin(a + bx + cx^2) dx$$

input `integrate((e*x+d)*sin(c*x**2+b*x+a),x)`

output `Integral((d + e*x)*sin(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 694, normalized size of antiderivative = 4.96

$$\int (d + ex) \sin(a + bx + cx^2) dx = \text{Too large to display}$$

```
input integrate((e*x+d)*sin(c*x^2+b*x+a),x, algorithm="maxima")
```

```
output -1/8*sqrt(2)*sqrt(pi)*((-I + 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(I*c)) + (-I - 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(-I*c)))*d/sqrt(c) + 1/16*((-I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/4*(b^2 - 4*a*c)/c) + ((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/4*(b^2 - 4*a*c)/c) - 2*((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/4*(b^2 - 4*a*c)/c) + (-I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/4*(b^2 - 4*a*c)/c))*x - 4*(c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^2 - 4*a*c)/c) + c*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/4*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.42

$$\int (d + ex) \sin(a + bx + cx^2) dx$$

$$= -\frac{\frac{\sqrt{2}\sqrt{\pi}(-2icd+ibe)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4iac}{4c}\right)}}{\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} + 2ee^{(icx^2+ibx+ia)}}{8c}$$

$$- \frac{\frac{\sqrt{2}\sqrt{\pi}(2icd-ibe)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4iac}{4c}\right)}}{\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}} + 2ee^{(-icx^2-ibx-ia)}}{8c}$$

input `integrate((e*x+d)*sin(c*x^2+b*x+a),x, algorithm="giac")`

output
$$\begin{aligned} & -\frac{1}{8} \sqrt{2} \sqrt{\pi} (-2I*c*d + I*b*e) \operatorname{erf}\left(\frac{-1}{4} \sqrt{2} (2*x + b/c)\right) (-I \\ & *c/abs(c) + 1) \sqrt{abs(c)} e^{-\frac{1}{4}(I*b^2 - 4*I*a*c)/c} / ((-I*c/abs(c) + \\ & 1) \sqrt{abs(c)}) + 2e^{\frac{1}{4}(I*c*x^2 + I*b*x + I*a)/c} - \frac{1}{8} \sqrt{2} \sqrt{\pi} (2*I*c*d - I*b*e) \operatorname{erf}\left(\frac{-1}{4} \sqrt{2} (2*x + b/c)\right) (I*c/abs(c) + 1) \sqrt{abs(c)} \\ & *e^{-\frac{1}{4}(-I*b^2 + 4*I*a*c)/c} / ((I*c/abs(c) + 1) \sqrt{abs(c)}) + 2e^{\frac{1}{4}(-I*c*x^2 - I*b*x - I*a)/c} \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int (d + ex) \sin(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a) (d + ex) dx$$

input `int(sin(a + b*x + c*x^2)*(d + e*x),x)`

output `int(sin(a + b*x + c*x^2)*(d + e*x), x)`

Reduce [F]

$$\begin{aligned} \int (d + ex) \sin(a + bx + cx^2) dx &= \left(\int \sin(cx^2 + bx + a) dx \right) d \\ &+ \left(\int \sin(cx^2 + bx + a) x dx \right) e \end{aligned}$$

input `int((e*x+d)*sin(c*x^2+b*x+a),x)`

output `int(sin(a + b*x + c*x**2),x)*d + int(sin(a + b*x + c*x**2)*x,x)*e`

3.31 $\int \sin(a + bx + cx^2) dx$

Optimal result	227
Mathematica [A] (verified)	227
Rubi [A] (verified)	228
Maple [A] (verified)	229
Fricas [A] (verification not implemented)	230
Sympy [A] (verification not implemented)	230
Maxima [C] (verification not implemented)	231
Giac [C] (verification not implemented)	231
Mupad [B] (verification not implemented)	232
Reduce [F]	232

Optimal result

Integrand size = 11, antiderivative size = 97

$$\int \sin(a + bx + cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\ + \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{\sqrt{c}}$$

output

```
1/2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(1/2)+1/2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec), antiderivative size = 84, normalized size of antiderivative = 0.87

$$\int \sin(a + bx + cx^2) dx \\ = \frac{\sqrt{\frac{\pi}{2}} \left(\cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \right)}{\sqrt{c}}$$

input `Integrate[Sin[a + b*x + c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot (\text{Cos}[a - b^2/(4c]) \cdot \text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] + \text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] \cdot \text{Sin}[a - b^2/(4c)]))}{\text{Sqrt}[c]}$$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3928, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sin(a + bx + cx^2) \, dx \\
 & \downarrow 3928 \\
 & \sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) \, dx + \cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b + 2cx)^2}{4c}\right) \, dx \\
 & \downarrow 3832 \\
 & \sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) \, dx + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\
 & \downarrow 3833 \\
 & \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}
 \end{aligned}$$

input `Int[Sin[a + b*x + c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{Cos}[a - b^2/(4c]) \cdot \text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])])}{\text{Sqrt}[c]} + \frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] \cdot \text{Sin}[a - b^2/(4c)])}{\text{Sqrt}[c]}$$

Definitions of rubi rules used

rule 3832 $\text{Int}[\text{Sin}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\text{Int}[\text{Cos}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\text{Int}[\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{b^2-ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)-\sin\left(\frac{b^2-ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)\right)}{2\sqrt{c}}$	82
risch	$\frac{i\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}} + \frac{i\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}}$	101

input `int(sin(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2}2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}\text{cos}((1/4b^2-a*c)/c)\text{FresnelS}(2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}(c*x+1/2*b))-\text{sin}((1/4b^2-a*c)/c)\text{FresnelC}(2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}(c*x+1/2*b)))$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.05

$$\int \sin(a + bx + cx^2) dx \\ = \frac{\sqrt{2}\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi\sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2 - 4ac}{4c}\right)}{2c}$$

input `integrate(sin(c*x^2+b*x+a),x, algorithm="fricas")`

output `1/2*(sqrt(2)*pi*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) + sqrt(2)*pi*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c))/c`

Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.91

$$\int \sin(a + bx + cx^2) dx \\ = \frac{\sqrt{2}\sqrt{\pi} \left(\sin\left(a - \frac{b^2}{4c}\right) C\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) + \cos\left(a - \frac{b^2}{4c}\right) S\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) \right) \sqrt{\frac{1}{c}}}{2}$$

input `integrate(sin(c*x**2+b*x+a),x)`

output `sqrt(2)*sqrt(pi)*(sin(a - b**2/(4*c))*fresnelc(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))) + cos(a - b**2/(4*c))*fresnels(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))))*sqrt(1/c)/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.15

$$\int \sin(a + bx + cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(\left(-(i+1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i-1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) + \left(-(i-1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i+1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{-2icx-ib}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

input `integrate(sin(c*x^2+b*x+a),x, algorithm="maxima")`

output
$$\begin{aligned} & -\frac{1}{8}\sqrt{2}\sqrt{\pi}(\operatorname{erf}\left(\frac{i}{2}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4ic}{4c}\right)} \\ & + \operatorname{erf}\left(\frac{-i}{2}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4ic}{4c}\right)}) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.39

$$\int \sin(a + bx + cx^2) dx = \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) e^{\left(-\frac{ib^2-4ic}{4c}\right)}}{4\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2+4ic}{4c}\right)}}{4\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}}$$

input `integrate(sin(c*x^2+b*x+a),x, algorithm="giac")`

output
$$\begin{aligned} & \frac{1}{4}i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) \\ & + \frac{1}{4}i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x+\frac{b}{c})\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.02

$$\int \sin(a + bx + cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} \cos\left(\frac{4ac - b^2}{4c}\right) S\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2} + \frac{\sqrt{2} \sqrt{\pi} \sin\left(\frac{4ac - b^2}{4c}\right) C\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2}$$

input `int(sin(a + b*x + c*x^2),x)`

output `(2^(1/2)*pi^(1/2)*cos((4*a*c - b^2)/(4*c))*fresnels((2^(1/2)*(b/2 + c*x)*(1/c)^(1/2))/pi^(1/2))*(1/c)^(1/2))/2 + (2^(1/2)*pi^(1/2)*sin((4*a*c - b^2)/(4*c))*fresnelc((2^(1/2)*(b/2 + c*x)*(1/c)^(1/2))/pi^(1/2))*(1/c)^(1/2))/2`

Reduce [F]

$$\int \sin(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a) dx$$

input `int(sin(c*x^2+b*x+a),x)`

output `int(sin(a + b*x + c*x**2),x)`

3.32 $\int \frac{\sin(a+bx+cx^2)}{d+ex} dx$

Optimal result	233
Mathematica [N/A]	233
Rubi [N/A]	234
Maple [N/A]	234
Fricas [N/A]	235
Sympy [N/A]	235
Maxima [N/A]	236
Giac [N/A]	236
Mupad [N/A]	236
Reduce [N/A]	237

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \text{Int}\left(\frac{\sin(a + bx + cx^2)}{d + ex}, x\right)$$

output `Defer(Int)(sin(c*x^2+b*x+a)/(e*x+d),x)`

Mathematica [N/A]

Not integrable

Time = 6.37 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(a + bx + cx^2)}{d + ex} dx$$

input `Integrate[Sin[a + b*x + c*x^2]/(d + e*x), x]`

output `Integrate[Sin[a + b*x + c*x^2]/(d + e*x), x]`

Rubi [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3950}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx$$

↓ 3950

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx$$

input `Int[Sin[a + b*x + c*x^2]/(d + e*x), x]`

output `$Aborted`

Defintions of rubi rules used

rule 3950 `Int[((d_.) + (e_.)*(x_))^(m_.)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_.), x_Symbol] :> Unintegrable[(d + e*x)^m*Sin[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\sin(cx^2 + bx + a)}{ex + d} dx$$

input `int(sin(c*x^2+b*x+a)/(e*x+d), x)`

output `int(sin(c*x^2+b*x+a)/(e*x+d),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)}{ex + d} dx$$

input `integrate(sin(c*x^2+b*x+a)/(e*x+d),x, algorithm="fricas")`

output `integral(sin(c*x^2 + b*x + a)/(e*x + d), x)`

Sympy [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(a + bx + cx^2)}{d + ex} dx$$

input `integrate(sin(c*x**2+b*x+a)/(e*x+d),x)`

output `Integral(sin(a + b*x + c*x**2)/(d + e*x), x)`

Maxima [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)}{ex + d} dx$$

input `integrate(sin(c*x^2+b*x+a)/(e*x+d),x, algorithm="maxima")`

output `integrate(sin(c*x^2 + b*x + a)/(e*x + d), x)`

Giac [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)}{ex + d} dx$$

input `integrate(sin(c*x^2+b*x+a)/(e*x+d),x, algorithm="giac")`

output `integrate(sin(c*x^2 + b*x + a)/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 38.44 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)}{d + e x} dx$$

input `int(sin(a + b*x + c*x^2)/(d + e*x),x)`

output `int(sin(a + b*x + c*x^2)/(d + e*x), x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\sin(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)}{ex + d} dx$$

input `int(sin(c*x^2+b*x+a)/(e*x+d),x)`

output `int(sin(a + b*x + c*x**2)/(d + e*x),x)`

3.33 $\int (d + ex)^2 \sin^2(a + bx + cx^2) dx$

Optimal result	238
Mathematica [A] (verified)	239
Rubi [A] (verified)	239
Maple [A] (verified)	241
Fricas [A] (verification not implemented)	241
Sympy [F]	242
Maxima [C] (verification not implemented)	242
Giac [C] (verification not implemented)	243
Mupad [F(-1)]	244
Reduce [F]	244

Optimal result

Integrand size = 21, antiderivative size = 291

$$\begin{aligned} \int (d + ex)^2 \sin^2(a + bx + cx^2) dx &= \frac{(d + ex)^3}{6e} \\ &- \frac{(2cd - be)^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ &+ \frac{e^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ &+ \frac{e^2 \sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\ &+ \frac{(2cd - be)^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\ &- \frac{e(2cd - be) \sin(2a + 2bx + 2cx^2)}{16c^2} \\ &- \frac{e(d + ex) \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output

```
1/6*(e*x+d)^3/e-1/16*(-b*e+2*c*d)^2*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(5/2)+1/16*e^2*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)+1/16*e^2*Pi^(1/2)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)+1/16*(-b*e+2*c*d)^2*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(5/2)-1/16*e*(-b*e+2*c*d)*sin(2*c*x^2+2*b*x+2*a)/c^2-1/8*e*(e*x+d)*sin(2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 1.17 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.74

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx \\ = \frac{-3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left((-2cd + be)^2 \cos\left(2a - \frac{b^2}{2c}\right) - ce^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) + 3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(ce^2 \sin\left(2a - \frac{b^2}{2c}\right) + (-2cd + be)^2 \cos\left(2a - \frac{b^2}{2c}\right)\right)}{1}$$

input

```
Integrate[(d + e*x)^2 Sin[a + b*x + c*x^2]^2, x]
```

output

```
(-3*sqrt[Pi]*FresnelC[(b + 2*c*x)/(sqrt[c]*sqrt[Pi])]*((-2*c*d + b*e)^2*Cos[2*a - b^2/(2*c)] - c*e^2*Sin[2*a - b^2/(2*c)]) + 3*sqrt[Pi]*FresnelS[(b + 2*c*x)/(sqrt[c]*sqrt[Pi])]*((c*e^2*Cos[2*a - b^2/(2*c)] + (-2*c*d + b*e)^2*Sin[2*a - b^2/(2*c)]) + sqrt[c]*(8*c^2*x*(3*d^2 + 3*d*e*x + e^2*x^2) - 3*e*(4*c*d - b*e + 2*c*e*x)*Sin[2*(a + x*(b + c*x))]))/(48*c^(5/2))
```

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx$$

$$\begin{aligned}
 & \downarrow 3948 \\
 & \int \left(\frac{1}{2}(d+ex)^2 - \frac{1}{2}(d+ex)^2 \cos(2a + 2bx + 2cx^2) \right) dx \\
 & \downarrow 2009 \\
 & -\frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) (2cd-be)^2 \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \\
 & \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) (2cd-be)^2 \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \frac{\sqrt{\pi} e^2 \sin\left(2a - \frac{b^2}{2c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} + \\
 & \frac{\sqrt{\pi} e^2 \cos\left(2a - \frac{b^2}{2c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \frac{e(2cd-be) \sin(2a + 2bx + 2cx^2)}{8c} - \\
 & \frac{16c^2}{e(d+ex) \sin(2a + 2bx + 2cx^2)} + \frac{(d+ex)^3}{6e}
 \end{aligned}$$

input `Int[(d + e*x)^2*Sin[a + b*x + c*x^2]^2, x]`

output `(d + e*x)^3/(6*e) - ((2*c*d - b*e)^2*Sqrt[Pi]*Cos[2*a - b^2/(2*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])])/(16*c^(5/2)) + (e^2*Sqrt[Pi]*Cos[2*a - b^2/(2*c)]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])])/(16*c^(3/2)) + (e^2*Sqrt[Pi]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*Sin[2*a - b^2/(2*c)])/(16*c^(3/2)) + ((2*c*d - b*e)^2*Sqrt[Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*Sin[2*a - b^2/(2*c)])/(16*c^(5/2)) - (e*(2*c*d - b*e)*Sin[2*a + 2*b*x + 2*c*x^2])/(16*c^2) - (e*(d + e*x)*Sin[2*a + 2*b*x + 2*c*x^2])/(8*c)`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_.)*(x_.))^(m_.)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 2.01 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.30

method	result
default	$-\frac{e^2 x \sin(2cx^2+2bx+2a)}{8c} + \frac{e^2 b \left(\frac{\sin(2cx^2+2bx+2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{4c^{\frac{3}{2}}} \right)}{4c}$
risch	$-\frac{\text{erf}\left(\sqrt{2}\sqrt{ic}x+\frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}d^2e^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}} - \frac{e^2 \text{erf}\left(\sqrt{2}\sqrt{ic}x+\frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}b^2e^{-\frac{i(4ac-b^2)}{2c}}}{64\sqrt{ic}c^2} + \frac{ie^2 \text{erf}\left(\sqrt{2}\sqrt{ic}x+\frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}b^2e^{-\frac{i(4ac-b^2)}{2c}}}{64\sqrt{ic}c^2}$

input `int((e*x+d)^2*sin(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -1/8 * e^2 / c * x * \sin(2*c*x^2 + 2*b*x + 2*a) + 1/4 * e^2 * b / c * (1/4 * \sin(2*c*x^2 + 2*b*x + 2*a) \\ &) / c - 1/4 * b / c^{(3/2)} * \text{Pi}^{(1/2)} * (\cos(1/2 * (-4*a*c + b^2) / c) * \text{FresnelC}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)}) + \sin(1/2 * (-4*a*c + b^2) / c) * \text{FresnelS}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)})) + 1/16 * e^2 / c^{(3/2)} * \text{Pi}^{(1/2)} * (\cos(1/2 * (-4*a*c + b^2) / c) * \text{FresnelS}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)}) - \sin(1/2 * (-4*a*c + b^2) / c) * \text{FresnelC}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)})) - 1/4 * d * e / c * \sin(2*c*x^2 + 2*b*x + 2*a) + 1/4 * d * e * b / c^{(3/2)} * \text{Pi}^{(1/2)} * (\cos(1/2 * (-4*a*c + b^2) / c) * \text{FresnelC}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)}) + \sin(1/2 * (-4*a*c + b^2) / c) * \text{FresnelS}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)})) - 1/4 * \text{Pi}^{(1/2)} / c^{(1/2)} * d^{(2)} * (\cos(1/2 * (-4*a*c + b^2) / c) * \text{FresnelC}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)}) + \sin(1/2 * (-4*a*c + b^2) / c) * \text{FresnelS}((2*c*x + b) / c^{(1/2)} / \text{Pi}^{(1/2)})) + 1/2 * d * e * x^2 + 1/2 * d^{(2)} * x + 1/6 * e^{(2)} * x^{(3)} \end{aligned}$$
Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.88

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx$$

$$= \frac{8c^3e^2x^3 + 24c^3dex^2 + 24c^3d^2x - 6(2c^2e^2x + 4c^2de - bce^2)\cos(cx^2 + bx + a)\sin(cx^2 + bx + a) + 3(d^2 + 2de)x^2 + 2dx + 1}{c^2}$$

input `integrate((e*x+d)^2*sin(c*x^2+b*x+a)^2,x, algorithm="fricas")`

output

$$\frac{1}{48} \left(8c^3e^{2x^3} + 24c^3d e x^2 + 24c^3d^2 x - 6(2c^2e^{2x} + 4c^2d e - b c e^2) \cos(c x^2 + b x + a) \sin(c x^2 + b x + a) + 3(\pi c e^2 \sin(-\frac{1}{2}(b^2 - 4a c)/c) - \pi (4c^2 d^2 - 4b c d e + b^2 e^2) \cos(-\frac{1}{2}(b^2 - 4a c)/c)) \sqrt{c/\pi} \operatorname{fresnel_cos}(2c x + b) \sqrt{c/\pi}/c + 3(\pi * c e^2 \cos(-\frac{1}{2}(b^2 - 4a c)/c) + \pi (4c^2 d^2 - 4b c d e + b^2 e^2) \sin(-\frac{1}{2}(b^2 - 4a c)/c)) \sqrt{c/\pi} \operatorname{fresnel_sin}(2c x + b) \sqrt{c/\pi}/c \right) / c^3$$

Sympy [F]

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx = \int (d + ex)^2 \sin^2(a + bx + cx^2) dx$$

input

```
integrate((e*x+d)**2*sin(c*x**2+b*x+a)**2,x)
```

output

```
Integral((d + e*x)**2*sin(a + b*x + c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.09 (sec) , antiderivative size = 2361, normalized size of antiderivative = 8.11

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx = \text{Too large to display}$$

input

```
integrate((e*x+d)^2*sin(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

```

1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I +
1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(2*I*c)) + ((I + 1)*
cos(-1/2*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*
x + I*b)/sqrt(-2*I*c)))*c^(3/2) + 16*c^2*x)*d^2/c^2 + 1/32*sqrt(2)*((-I -
1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/
c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4
*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/2*(b^2 - 4*a*c)/c) + (-I + 1)*sqrt(
2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1
) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x
+ I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*((I - 1)*sqrt(2)*sq
rt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I
+ 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b
^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(e
rf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqr
t(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) -
1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(-I*e^(1/
2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) + I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c
*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) - c*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*
c*x + I*b^2)/c)) + e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(
b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*d*e/(c^2*sqrt((4*...

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.02

$$\begin{aligned}
 \int (d + ex)^2 \sin^2(a + bx + cx^2) dx &= \frac{1}{6} e^2 x^3 + \frac{1}{2} d e x^2 + \frac{1}{2} d^2 x \\
 &\quad - \frac{-i (ce^2(2x + \frac{b}{c}) + 4cde - 2be^2)e^{(2icx^2 + 2ibx + 2ia)} + \frac{i\sqrt{\pi}(4ic^2d^2 - 4ibcde + ib^2e^2 - ce^2)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{(-4ic^2d^2 + 4ibcde - ib^2e^2 - ce^2)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2} \\
 &\quad - \frac{i (ce^2(2x + \frac{b}{c}) + 4cde - 2be^2)e^{(-2icx^2 - 2ibx - 2ia)} - \frac{i\sqrt{\pi}(-4ic^2d^2 + 4ibcde - ib^2e^2 - ce^2)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{(-4ic^2d^2 + 4ibcde - ib^2e^2 - ce^2)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2}
 \end{aligned}$$

input `integrate((e*x+d)^2*sin(c*x^2+b*x+a)^2,x, algorithm="giac")`

output

```
1/6*e^2*x^3 + 1/2*d*e*x^2 + 1/2*d^2*x - 1/32*(-I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a) + I*sqrt(pi)*(4*I*c^2*d^2 - 4*I*b*c*d*e + I*b^2*e^2 - c*e^2)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)))/c^2 - 1/32*(I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a) - I*sqrt(pi)*(-4*I*c^2*d^2 + 4*I*b*c*d*e - I*b^2*e^2 - c*e^2)*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)))/c^2
```

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a)^2 (d + ex)^2 dx$$

input

```
int(sin(a + b*x + c*x^2)^2*(d + e*x)^2, x)
```

output

```
int(sin(a + b*x + c*x^2)^2*(d + e*x)^2, x)
```

Reduce [F]

$$\int (d + ex)^2 \sin^2(a + bx + cx^2) dx$$

$$= \frac{\cos(cx^2 + bx + a) \sin(cx^2 + bx + a) b e^2 - 3 \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) c d e - 2 \cos(cx^2 + bx + a) c^2 d^2 e^2}{b^3}$$

input

```
int((e*x+d)^2*sin(c*x^2+b*x+a)^2, x)
```

output

```
(cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*b*e**2 - 3*cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*c*d*e - 2*cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*c*e**2*x + 6*int(tan((a + b*x + c*x**2)/2)**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2*e**2 - 24*int(tan((a + b*x + c*x**2)/2)**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b*c*d*e + 24*int(tan((a + b*x + c*x**2)/2)**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**2*d**2 + 8*int(x**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**2*e**2 + 8*int(tan((a + b*x + c*x**2)/2)/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c*e**2 - 2*int(1/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2*e**2 + sin(a + b*x + c*x**2)*b*e**2 - 2*sin(a + b*x + c*x**2)*c*e**2*x + 3*a*c*d*e + 3*b*c*d*e*x + 3*c**2*d*e*x**2)/(6*c**2)
```

3.34 $\int (d + ex) \sin^2(a + bx + cx^2) dx$

Optimal result	246
Mathematica [A] (verified)	247
Rubi [A] (verified)	247
Maple [A] (verified)	248
Fricas [A] (verification not implemented)	249
Sympy [F]	249
Maxima [C] (verification not implemented)	249
Giac [C] (verification not implemented)	250
Mupad [F(-1)]	251
Reduce [F]	251

Optimal result

Integrand size = 19, antiderivative size = 150

$$\begin{aligned} \int (d + ex) \sin^2(a + bx + cx^2) dx &= \frac{(d + ex)^2}{4e} \\ &- \frac{(2cd - be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ &+ \frac{(2cd - be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} \\ &- \frac{e \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output

```
1/4*(e*x+d)^2/e-1/8*(-b*e+2*c*d)*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)+1/8*(-b*e+2*c*d)*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)-1/8*e*sin(2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.93

$$\int (d + ex) \sin^2(a + bx + cx^2) dx \\ = \frac{-\left((2cd - be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + (2cd - be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right) + \sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \cos\left(2a - \frac{b^2}{2c}\right)\right)}{8c^{3/2}}$$

input `Integrate[(d + e*x)*Sin[a + b*x + c*x^2]^2, x]`

output $\frac{(-((2*c*d - b*e)*\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a - b^2/(2*c)]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]) + (2*c*d - b*e)*\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])] * \text{Sin}[2*a - b^2/(2*c)] + \text{Sqrt}[c]*(2*c*x*(2*d + e*x) - e*\text{Sin}[2*(a + x*(b + c*x))]))}{(8*c^{(3/2)})}$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) \sin^2(a + bx + cx^2) dx \\ \downarrow 3948 \\ \int \left(\frac{1}{2}(d + ex) - \frac{1}{2}(d + ex) \cos(2a + 2bx + 2cx^2) \right) dx \\ \downarrow 2009 \\ - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) (2cd - be) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} + \\ \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) (2cd - be) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} - \frac{e \sin(2a + 2bx + 2cx^2)}{8c} + \frac{(d + ex)^2}{4e}$$

input $\text{Int}[(d + e*x)*\sin[a + b*x + c*x^2]^2, x]$

output $(d + e*x)^2/(4*e) - ((2*c*d - b*e)*\sqrt{\pi}*\cos[2*a - b^2/(2*c)]*\text{FresnelC}[(b + 2*c*x)/(\sqrt{c}*\sqrt{\pi})])/(8*c^{(3/2)}) + ((2*c*d - b*e)*\sqrt{\pi}*\text{FresnelS}[(b + 2*c*x)/(\sqrt{c}*\sqrt{\pi})]*\sin[2*a - b^2/(2*c)])/(8*c^{(3/2)}) - (e*\sin[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3948 $\text{Int}[(d_.) + (e_.)*(x_.)]^m_*\sin[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^n, x] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \sin[a + b*x + c*x^2]^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{IGtQ}[n, 1]$

Maple [A] (verified)

Time = 1.53 (sec), antiderivative size = 170, normalized size of antiderivative = 1.13

method	result
default	$-\frac{e \sin(2cx^2+2bx+2a)}{8c} + \frac{eb\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{8c^{\frac{3}{2}}} - \frac{\sqrt{\pi} d \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{8c}$
risch	$-\frac{\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}de^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}} + \frac{eb\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{2c}}\sqrt{2}\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c\sqrt{ic}} + \frac{\text{erf}\left(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)\sqrt{\pi}d}{8\sqrt{-2ic}}$

input $\text{int}((e*x+d)*\sin(c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $-1/8*e*\sin(2*c*x^2+2*b*x+2*a)/c+1/8*e*b/c^{(3/2)}*\text{Pi}^{(1/2)}*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)})+\sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))-1/4*\text{Pi}^{(1/2)}/c^{(1/2)}*d*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)})+\sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))+1/2*d*x+1/4*e*x^2$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.99

$$\int (d + ex) \sin^2(a + bx + cx^2) dx$$

$$= \frac{2 c^2 e x^2 - \pi (2 c d - b e) \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4 a c}{2 c}\right) C\left(\frac{(2 c x + b) \sqrt{\frac{c}{\pi}}}{c}\right) + \pi (2 c d - b e) \sqrt{\frac{c}{\pi}} S\left(\frac{(2 c x + b) \sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2 - 4 a c}{2 c}\right)}{8 c^2}$$

input `integrate((e*x+d)*sin(c*x^2+b*x+a)^2,x, algorithm="fricas")`

output `1/8*(2*c^2*e*x^2 - pi*(2*c*d - b*e)*sqrt(c/pi)*cos(-1/2*(b^2 - 4*a*c)/c)*fresnel_resnel_cos((2*c*x + b)*sqrt(c/pi)/c) + pi*(2*c*d - b*e)*sqrt(c/pi)*fresnel_resnel_sin((2*c*x + b)*sqrt(c/pi)/c)*sin(-1/2*(b^2 - 4*a*c)/c) + 4*c^2*d*x - 2*c*e*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a))/c^2`

Sympy [F]

$$\int (d + ex) \sin^2(a + bx + cx^2) dx = \int (d + ex) \sin^2(a + bx + cx^2) dx$$

input `integrate((e*x+d)*sin(c*x**2+b*x+a)**2,x)`

output `Integral((d + e*x)*sin(a + b*x + c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 738, normalized size of antiderivative = 4.92

$$\int (d + ex) \sin^2(a + bx + cx^2) dx = \text{Too large to display}$$

input `integrate((e*x+d)*sin(c*x^2+b*x+a)^2,x, algorithm="maxima")`

output

$$\begin{aligned} & \frac{1}{32} \cdot (4^{(1/4)} \cdot \sqrt{2} \cdot \sqrt{\pi}) \cdot (((I - 1) \cdot \cos(-1/2 \cdot (b^2 - 4*a*c)/c) + (I + 1) \cdot \sin(-1/2 \cdot (b^2 - 4*a*c)/c)) \cdot \operatorname{erf}((2*I*c*x + I*b)/\sqrt{2*I*c})) + ((I + 1) \cdot \cos(-1/2 \cdot (b^2 - 4*a*c)/c) + (I - 1) \cdot \sin(-1/2 \cdot (b^2 - 4*a*c)/c)) \cdot \operatorname{erf}((2*I*c*x + I*b)/\sqrt{-2*I*c})) \cdot c^{(3/2)} + 16*c^2*x) \cdot d/c^2 + 1/64 \cdot \sqrt{2} \cdot ((-(I - 1) \cdot \sqrt{2} \cdot \sqrt{\pi}) \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c})) - 1) + (I + 1) \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))) - 1) \cdot b^2 \cdot \cos(-1/2 \cdot (b^2 - 4*a*c)/c) + (-(I + 1) \cdot \sqrt{2} \cdot \sqrt{\pi}) \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c})) - 1) + (I - 1) \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))) - 1) \cdot b^2 \cdot \sin(-1/2 \cdot (b^2 - 4*a*c)/c) - 2 \cdot ((I - 1) \cdot \sqrt{2} \cdot \sqrt{\pi}) \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c})) - 1) - (I + 1) \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))) - 1) \cdot b*c \cdot \cos(-1/2 \cdot (b^2 - 4*a*c)/c) + ((I + 1) \cdot \sqrt{2} \cdot \sqrt{\pi}) \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c})) - 1) - (I - 1) \cdot \sqrt{2} \cdot \sqrt{\pi} \cdot (\operatorname{erf}(\sqrt{1/2}) \cdot \sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))) - 1) \cdot b*c \cdot \sin(-1/2 \cdot (b^2 - 4*a*c)/c)) \cdot x + 2 \cdot \sqrt{2} \cdot (4*c^2*x^2 - c \cdot (-I \cdot e^{(1/2 \cdot (4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}) + I \cdot e^{(-1/2 \cdot (4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}) \cdot \cos(-1/2 \cdot (b^2 - 4*a*c)/c) - c \cdot (e^{(1/2 \cdot (4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}) \cdot \sin(-1/2 \cdot (b^2 - 4*a*c)/c)) \cdot \sqrt{(4*c^2*x^2 + 4*b*c*x + b^2)/c}) \cdot e/(c^2 \cdot \sqrt{(4*c^2*x^2 + 4*b*c*x + b^2)/c}) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.30

$$\begin{aligned} & \int (d + ex) \sin^2(a + bx + cx^2) dx \\ &= \frac{1}{4} ex^2 + \frac{1}{2} dx \\ & - \frac{-i e e^{(2i c x^2 + 2i b x + 2i a)} - \frac{i \sqrt{\pi} (-2i c d + i b e) \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x + \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right)\right) e^{\left(-\frac{i b^2 - 4i a c}{2c}\right)}}{\sqrt{c} \left(-\frac{i c}{|c|} + 1\right)}}{16 c} \\ & - \frac{i e e^{(-2i c x^2 - 2i b x - 2i a)} + \frac{i \sqrt{\pi} (2i c d - i b e) \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x + \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right)\right) e^{\left(-\frac{-i b^2 + 4i a c}{2c}\right)}}{\sqrt{c} \left(\frac{i c}{|c|} + 1\right)}}{16 c} \end{aligned}$$

input `integrate((e*x+d)*sin(c*x^2+b*x+a)^2,x, algorithm="giac")`

output
$$\frac{1/4*e*x^2 + 1/2*d*x - 1/16*(-I*e*e^{(2*I*c*x^2 + 2*I*b*x + 2*I*a)} - I*sqrt(pi)*(-2*I*c*d + I*b*e)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^{(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1))}/c - 1/16*(I*e*e^{(-2*I*c*x^2 - 2*I*b*x - 2*I*a)} + I*sqrt(pi)*(2*I*c*d - I*b*e)*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^{(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))})/c$$

Mupad [F(-1)]

Timed out.

$$\int (d + ex) \sin^2(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a)^2 (d + ex) dx$$

input `int(sin(a + b*x + c*x^2)^2*(d + e*x),x)`

output `int(sin(a + b*x + c*x^2)^2*(d + e*x), x)`

Reduce [F]

$$\int (d + ex) \sin^2(a + bx + cx^2) dx = \frac{-\cos(cx^2 + bx + a) \sin(cx^2 + bx + a) e - 2 \left(\int \sin(cx^2 + bx + a)^2 dx \right) be + 4 \left(\int \sin(cx^2 + bx + a)^2 dx \right)}{4c}$$

input `int((e*x+d)*sin(c*x^2+b*x+a)^2,x)`

output
$$\left(-\cos(a + b*x + c*x^{**2}) * \sin(a + b*x + c*x^{**2}) * e - 2 * \int (\sin(a + b*x + c*x^{**2})^2, x) * b*e + 4 * \int (\sin(a + b*x + c*x^{**2})^{**2}, x) * c*d + b*e*x + c*e*x^{**2} \right) / (4*c)$$

3.35 $\int \sin^2(a + bx + cx^2) dx$

Optimal result	252
Mathematica [A] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	254
Fricas [A] (verification not implemented)	254
Sympy [A] (verification not implemented)	255
Maxima [C] (verification not implemented)	255
Giac [C] (verification not implemented)	256
Mupad [F(-1)]	256
Reduce [F]	257

Optimal result

Integrand size = 13, antiderivative size = 100

$$\begin{aligned} \int \sin^2(a + bx + cx^2) dx &= \frac{x}{2} - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} \\ &\quad + \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

output

$$\begin{aligned} &1/2*x-1/4*Pi^{(1/2)}*\cos(2*a-1/2*b^2/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/Pi^{(1/2)}) \\ &/c^{(1/2)+1/4*Pi^{(1/2)}*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/Pi^{(1/2)})*\sin(2*a-1/2*b^2 \\ &/c)/c^{(1/2)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec), antiderivative size = 97, normalized size of antiderivative = 0.97

$$\begin{aligned} \int \sin^2(a + bx + cx^2) dx \\ &= \frac{2\sqrt{c}x - \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

input `Integrate[Sin[a + b*x + c*x^2]^2, x]`

output
$$\frac{(2\sqrt{c}x - \sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] + \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])}{(4\sqrt{c})}$$

Rubi [A] (verified)

Time = 0.22 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3930, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(a + bx + cx^2) dx \\ & \quad \downarrow \text{3930} \\ & \int \left(\frac{1}{2} - \frac{1}{2} \cos(2a + 2bx + 2cx^2) \right) dx \\ & \quad \downarrow \text{2009} \\ & -\frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2} \end{aligned}$$

input `Int[Sin[a + b*x + c*x^2]^2, x]`

output
$$\frac{x}{2} - \frac{(\sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] + \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])}{(4\sqrt{c})}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3930 $\text{Int}[\text{Sin}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTri}\text{gReduce}[\text{Sin}[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c\}, \ x] \ \&\& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 1.13 (sec), antiderivative size = 72, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{x}{2} - \frac{\sqrt{\pi} (\cos(\frac{-4ac+b^2}{2c}) \text{FresnelC}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}) + \sin(\frac{-4ac+b^2}{2c}) \text{FresnelS}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}))}{4\sqrt{c}}$	72
risch	$\frac{x}{2} - \frac{\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \text{erf}(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}})}{16\sqrt{ic}} + \frac{\sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \text{erf}(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}})}{8\sqrt{-2ic}}$	111

input $\text{int}(\text{sin}(c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $1/2*x - 1/4*\text{Pi}^{(1/2)}/c^{(1/2)}*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}) + \sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))$

Fricas [A] (verification not implemented)

Time = 0.10 (sec), antiderivative size = 93, normalized size of antiderivative = 0.93

$$\int \sin^2(a + bx + cx^2) dx \\ = -\frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2-4ac}{2c}\right) - 2cx}{4c}$$

input $\text{integrate}(\text{sin}(c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

output

$$\begin{aligned} & -\frac{1}{4} \left(\frac{\pi \sqrt{c/\pi}}{c} \cos \left(-\frac{1}{2} (b^2 - 4ac)/c \right) \operatorname{fresnel_cos}((2cx + b)\sqrt{c/\pi}) \right. \\ & \quad \left. - \frac{\pi \sqrt{c/\pi}}{c} \operatorname{fresnel_sin}((2cx + b)\sqrt{c/\pi}) \sin \left(-\frac{1}{2} (b^2 - 4ac)/c \right) - \frac{2cx}{c} \right) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.43 (sec), antiderivative size = 83, normalized size of antiderivative = 0.83

$$\begin{aligned} & \int \sin^2(a + bx + cx^2) dx \\ &= \frac{x}{2} - \frac{\sqrt{\pi} \left(-\sin \left(2a - \frac{b^2}{2c} \right) S \left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}} \right) + \cos \left(2a - \frac{b^2}{2c} \right) C \left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}} \right) \right) \sqrt{\frac{1}{c}}}{4} \end{aligned}$$

input

```
integrate(sin(c*x**2+b*x+a)**2,x)
```

output

$$\begin{aligned} & \frac{x}{2} - \sqrt{\pi} \left(-\sin(2a - b^2/(2c)) \operatorname{fresnels}((2b + 4cx)/(2\sqrt{\pi}\sqrt{c})) + \cos(2a - b^2/(2c)) \operatorname{fresnelc}((2b + 4cx)/(2\sqrt{\pi}\sqrt{c})) \right) \sqrt{1/c}/4 \end{aligned}$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec), antiderivative size = 124, normalized size of antiderivative = 1.24

$$\begin{aligned} & \int \sin^2(a + bx + cx^2) dx \\ &= \frac{4^{1/4} \sqrt{2} \sqrt{\pi} \left(((i-1) \cos \left(-\frac{b^2-4ac}{2c} \right) + (i+1) \sin \left(-\frac{b^2-4ac}{2c} \right)) \operatorname{erf} \left(\frac{2icx+ib}{\sqrt{2i}c} \right) + ((i+1) \cos \left(-\frac{b^2-4ac}{2c} \right) + (i-1) \sin \left(-\frac{b^2-4ac}{2c} \right)) \operatorname{erfc} \left(\frac{2icx+ib}{\sqrt{2i}c} \right) \right)}{32c^2} \end{aligned}$$

input

```
integrate(sin(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

$$\begin{aligned} & \frac{1}{32} \left(4^{1/4} \sqrt{2} \sqrt{\pi} \left(((I-1) \cos(-1/2(b^2 - 4ac)/c) + (I+1) \sin(-1/2(b^2 - 4ac)/c)) \operatorname{erf}((2*I*c*x + I*b)/\sqrt{2*I*c}) + ((I+1) \cos(-1/2(b^2 - 4ac)/c) + (I-1) \sin(-1/2(b^2 - 4ac)/c)) \operatorname{erfc}((2*I*c*x + I*b)/\sqrt{-2*I*c}) \right) c^{3/2} + 16c^2 x \right) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.22

$$\int \sin^2(a + bx + cx^2) dx = \frac{1}{2}x + \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 - 4iac}{2c}\right)}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)} \\ + \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 + 4iac}{2c}\right)}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}$$

input `integrate(sin(c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/2*x + 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) + 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \sin^2(a + bx + cx^2) dx = \int \sin(cx^2 + bx + a)^2 dx$$

input `int(sin(a + b*x + c*x^2)^2,x)`

output `int(sin(a + b*x + c*x^2)^2, x)`

Reduce [F]

$$\int \sin^2(a + bx + cx^2) dx = \int \sin(c x^2 + bx + a)^2 dx$$

input `int(sin(c*x^2+b*x+a)^2,x)`

output `int(sin(a + b*x + c*x**2)**2,x)`

3.36 $\int \frac{\sin^2(a+bx+cx^2)}{d+ex} dx$

Optimal result	258
Mathematica [N/A]	258
Rubi [N/A]	259
Maple [N/A]	260
Fricas [N/A]	260
Sympy [N/A]	260
Maxima [N/A]	261
Giac [N/A]	261
Mupad [N/A]	262
Reduce [N/A]	262

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \frac{\log(d + ex)}{2e} - \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx + 2cx^2)}{d + ex}, x\right)$$

output `1/2*ln(e*x+d)/e-1/2*DefeR(Int)(cos(2*c*x^2+2*b*x+2*a)/(e*x+d),x)`

Mathematica [N/A]

Not integrable

Time = 8.64 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx$$

input `Integrate[Sin[a + b*x + c*x^2]^2/(d + e*x), x]`

output `Integrate[Sin[a + b*x + c*x^2]^2/(d + e*x), x]`

Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3948, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx \\ & \quad \downarrow \text{3948} \\ & \int \left(\frac{1}{2(d + ex)} - \frac{\cos(2a + 2bx + 2cx^2)}{2(d + ex)} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{\log(d + ex)}{2e} - \frac{1}{2} \int \frac{\cos(2cx^2 + 2bx + 2a)}{d + ex} dx \end{aligned}$$

input `Int[Sin[a + b*x + c*x^2]^2/(d + e*x), x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3948 `Int[((d_.) + (e_.)*(x_.))^(m_.)*Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Sin[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\sin(cx^2 + bx + a)^2}{ex + d} dx$$

input `int(sin(c*x^2+b*x+a)^2/(e*x+d),x)`

output `int(sin(c*x^2+b*x+a)^2/(e*x+d),x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)^2}{ex + d} dx$$

input `integrate(sin(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="fricas")`

output `integral(-(cos(c*x^2 + b*x + a)^2 - 1)/(e*x + d), x)`

Sympy [N/A]

Not integrable

Time = 0.78 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx$$

input `integrate(sin(c*x**2+b*x+a)**2/(e*x+d),x)`

output `Integral(sin(a + b*x + c*x**2)**2/(d + e*x), x)`

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 118, normalized size of antiderivative = 5.62

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)^2}{ex + d} dx$$

input `integrate(sin(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="maxima")`

output `1/2*(2*e*integrate(-1/4*(cos(2*c*x^2 + 2*b*x)*cos(2*a) - sin(2*c*x^2 + 2*b*x)*sin(2*a))/((cos(2*a)^2 + sin(2*a)^2)*e*x + (cos(2*a)^2 + sin(2*a)^2)*d), x) - 2*e*integrate(1/4*cos(2*c*x^2 + 2*b*x + 2*a)/(e*x + d), x) + log(e*x + d))/e`

Giac [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)^2}{ex + d} dx$$

input `integrate(sin(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="giac")`

output `integrate(sin(c*x^2 + b*x + a)^2/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 39.44 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)^2}{d + ex} dx$$

input `int(sin(a + b*x + c*x^2)^2/(d + e*x),x)`

output `int(sin(a + b*x + c*x^2)^2/(d + e*x), x)`

Reduce [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\sin^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\sin(cx^2 + bx + a)^2}{ex + d} dx$$

input `int(sin(c*x^2+b*x+a)^2/(e*x+d),x)`

output `int(sin(a + b*x + c*x**2)**2/(d + e*x),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	263
4.2 Links to plain text integration problems used in this report for each CAS .	281

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```
(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*) (*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","");
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "}
  ,
]
```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```



```

ExpnType[expn_] :=
If[AtomQ[expn],
  1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn] === Power,
      If[IntegerQ[expn[[2]]],
        ExpnType[expn[[1]]],
        If[Head[expn[[2]]] === Rational,
          If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
            1,
            Max[ExpnType[expn[[1]]], 2]],
          Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
      If[Head[expn] === Plus || Head[expn] === Times,
        Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
        If[ElementaryFunctionQ[Head[expn]],
          Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], ExpnType[expn[[3]]]]]]]]]]]

```

```
Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],  
If[AppellFunctionQ[Head[expn]],  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],  
If[Head[expn]==RootSum,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],  
If[Head[expn]==Integrate || Head[expn]==Int,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],  
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=  
MemberQ[{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
} , func]
```

```
SpecialFunctionQ[func_] :=  
MemberQ[{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  

```

```
HypergeometricFunctionQ[func_] :=  
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=  
MemberQ[{AppellF1}, func]
```

Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
```

```
#      antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A", " ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
            # this assumes optimal do not as well. No check is needed here.
            if debug then
                print("result do not contain complex, this assumes optimal do not as well");
            fi;
```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                           convert(leaf_count_result,string),\"$ vs. \$2(", 
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_co
                           fi;
            fi;
        else #ExpnType(result) > ExpnType(optimal)
            if debug then
                print("ExpnType(result) > ExpnType(optimal)");
            fi;
            return "C",cat("Result contains higher order function than in optimal. Order ",
                           convert(ExpnType_result,string)," vs. order ",
                           convert(ExpnType_optimal,string),"."));
        fi;

    end proc:

    #

    # is_contains_complex(result)
    # takes expressions and returns true if it contains "I" else false
    #
    #Nasser 032417
    is_contains_complex:= proc(expression)
        return (has(expression,I));
    end proc:

    # The following summarizes the type number assigned an expression
    # based on the functions it involves
    # 1 = rational function
    # 2 = algebraic function
    # 3 = elementary function
    # 4 = special function
    # 5 = hypergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'`^`') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then
            if type(op(1,expn),'rational') then
                1
            else
                max(2,ExpnType(op(1,expn)))
            end if
        else
            max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
        end if
    elif type(expn,'`+`) or type(expn,'`*`) then
        max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif ElementaryFunctionQ(op(0,expn)) then
        max(3,ExpnType(op(1,expn)))
    elif SpecialFunctionQ(op(0,expn)) then
        max(4,apply(max,map(ExpnType,[op(expn)])))
    elif HypergeometricFunctionQ(op(0,expn)) then
        max(5,apply(max,map(ExpnType,[op(expn)])))
    elif AppellFunctionQ(op(0,expn)) then
        max(6,apply(max,map(ExpnType,[op(expn)])))
    elif op(0,expn)='int' then
        max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
```

```
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arccsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]
```

```
def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*')
```

```

m1 = expnType(expn.args[0])
m2 = expnType(list(expn.args[1:]))
return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sagemode")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(grade)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```
from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                       ]
    if debug:
```

```

if m:
    print ("func ", func , " is elementary_function")
else:
    print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral'
'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
'elliptic_pi','exp_integral_e','log_integral',
'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
'weierstrassPPrime','weierstrassSigma']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):
    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0], Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1], Integer)

```

```

    return expnType(expn.operands()[0])  #expnType(expn.args[0])
elif type(expn.operands()[1]) == Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0]) == Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))  #max(3,expnType(expn.args[0]),expnType(expn.args[1]))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(4,m1)  #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(5,m1)  #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

```

```
leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation ="none"
            else:
                grade = "B"
                grade_annotation ="Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation ="Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation ="none"
        else:
            grade = "B"
            grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(expnType_result)+"/"+str(expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file