

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.2-Cosine/202-4.2.12

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 3:20am

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24
2	detailed summary tables of results	25
2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	55
3	Listing of integrals	59
3.1	$\int x^3 \cos(a + bx^2) dx$	63
3.2	$\int x^2 \cos(a + bx^2) dx$	69
3.3	$\int x \cos(a + bx^2) dx$	76
3.4	$\int \cos(a + bx^2) dx$	81
3.5	$\int \frac{\cos(ax+bx^2)}{x} dx$	87
3.6	$\int \frac{\cos(ax+bx^2)}{x^2} dx$	92

3.7	$\int \frac{\cos(a+bx^2)}{x^3} dx$	98
3.8	$\int x^3 \cos^2(a+bx^2) dx$	105
3.9	$\int x^2 \cos^2(a+bx^2) dx$	111
3.10	$\int x \cos^2(a+bx^2) dx$	117
3.11	$\int \cos^2(a+bx^2) dx$	123
3.12	$\int \frac{\cos^2(a+bx^2)}{x} dx$	129
3.13	$\int \frac{\cos^2(a+bx^2)}{x^2} dx$	134
3.14	$\int \frac{\cos^2(a+bx^2)}{x^3} dx$	140
3.15	$\int x^3 \cos^3(a+bx^2) dx$	145
3.16	$\int x^2 \cos^3(a+bx^2) dx$	152
3.17	$\int x \cos^3(a+bx^2) dx$	160
3.18	$\int \cos^3(a+bx^2) dx$	166
3.19	$\int \frac{\cos^3(a+bx^2)}{x} dx$	172
3.20	$\int \frac{\cos^3(a+bx^2)}{x^2} dx$	177
3.21	$\int \frac{\cos^3(a+bx^2)}{x^3} dx$	183
3.22	$\int x \cos^7(a+bx^2) dx$	189
3.23	$\int x^{5/2} \cos(a+bx^2) dx$	195
3.24	$\int x^{3/2} \cos(a+bx^2) dx$	201
3.25	$\int \sqrt{x} \cos(a+bx^2) dx$	207
3.26	$\int \frac{\cos(a+bx^2)}{\sqrt{x}} dx$	212
3.27	$\int \frac{\cos(a+bx^2)}{x^{3/2}} dx$	217
3.28	$\int \frac{\cos(a+bx^2)}{x^{5/2}} dx$	223
3.29	$\int x^{5/2} \cos^2(a+bx^2) dx$	229
3.30	$\int x^{3/2} \cos^2(a+bx^2) dx$	234
3.31	$\int \sqrt{x} \cos^2(a+bx^2) dx$	239
3.32	$\int \frac{\cos^2(a+bx^2)}{\sqrt{x}} dx$	244
3.33	$\int \frac{\cos^2(a+bx^2)}{x^{3/2}} dx$	249
3.34	$\int \frac{\cos^2(a+bx^2)}{x^{5/2}} dx$	254
3.35	$\int \cos\left(a + \frac{b}{x}\right) dx$	260
3.36	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$	267
3.37	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx$	272
3.38	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx$	277
3.39	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx$	283
3.40	$\int \cos\left(a + \frac{b}{x^2}\right) dx$	289

3.41	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$	295
3.42	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$	300
3.43	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx$	306
3.44	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$	311
3.45	$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$	317
3.46	$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$	322
3.47	$\int \cos(\sqrt{x}) dx$	327
3.48	$\int \cos^2(\sqrt{x}) dx$	332
3.49	$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx$	337
3.50	$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$	373
3.51	$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx$	390
3.52	$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx$	398
3.53	$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx$	406
3.54	$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx$	417
3.55	$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx$	428
3.56	$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$	442
3.57	$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx$	451
3.58	$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx$	457
3.59	$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx$	464
3.60	$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx$	476
3.61	$\int \cos^3(\sqrt[3]{x}) dx$	487
3.62	$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx$	495
3.63	$\int (ex)^m (b \cos(c + dx^n))^p dx$	500
3.64	$\int (ex)^m (a + b \cos(c + dx^n))^p dx$	505
3.65	$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx$	510
3.66	$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$	515
3.67	$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx$	520
3.68	$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$	526
3.69	$\int \frac{\cos(ax^n)}{x} dx$	531
3.70	$\int \frac{\cos^2(ax^n)}{x} dx$	536
3.71	$\int \frac{\cos^3(ax^n)}{x} dx$	541

3.72	$\int \frac{\cos^4(a+bx^n)}{x} dx$	546
3.73	$\int \cos(a+bx^n) dx$	551
3.74	$\int \cos^2(a+bx^n) dx$	556
3.75	$\int \cos^3(a+bx^n) dx$	561
3.76	$\int x^m \cos(a+bx^n) dx$	566
3.77	$\int x^m \cos^2(a+bx^n) dx$	571
3.78	$\int x^m \cos^3(a+bx^n) dx$	576
3.79	$\int x^{-1-n} \cos(a+bx^n) dx$	581
3.80	$\int x^{-1-n} \cos^2(a+bx^n) dx$	587
3.81	$\int x^{-1-n} \cos^3(a+bx^n) dx$	592
3.82	$\int x^{-1-2n} \cos(a+bx^n) dx$	597
3.83	$\int x^{-1-2n} \cos^2(a+bx^n) dx$	604
3.84	$\int x^{-1-2n} \cos^3(a+bx^n) dx$	610
3.85	$\int x^2 \cos((a+bx)^2) dx$	616
3.86	$\int x \cos((a+bx)^2) dx$	622
3.87	$\int \cos((a+bx)^2) dx$	628
3.88	$\int \frac{\cos((a+bx)^2)}{x} dx$	633
3.89	$\int \frac{\cos((a+bx)^2)}{x^2} dx$	638
3.90	$\int x^2 \cos(a+b\sqrt{c+dx}) dx$	643
3.91	$\int x \cos(a+b\sqrt{c+dx}) dx$	651
3.92	$\int \cos(a+b\sqrt{c+dx}) dx$	657
3.93	$\int \frac{\cos(a+b\sqrt{c+dx})}{x} dx$	663
3.94	$\int \frac{\cos(a+b\sqrt{c+dx})}{x^2} dx$	669
3.95	$\int x^2 \cos(a+b\sqrt[3]{c+dx}) dx$	676
3.96	$\int x \cos(a+b\sqrt[3]{c+dx}) dx$	686
3.97	$\int \cos(a+b\sqrt[3]{c+dx}) dx$	694
3.98	$\int \frac{\cos(a+b\sqrt[3]{c+dx})}{x} dx$	701
3.99	$\int \frac{\cos(a+b\sqrt[3]{c+dx})}{x^2} dx$	708

4	Appendix	716
4.1	Listing of Grading functions	716
4.2	Links to plain text integration problems used in this report for each CAS734	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	6
1.2	Results	7
1.3	Time and leaf size Performance	11
1.4	Performance based on number of rules Rubi used	13
1.5	Performance based on number of steps Rubi used	14
1.6	Solved integrals histogram based on leaf size of result	15
1.7	Solved integrals histogram based on CPU time used	16
1.8	Leaf size vs. CPU time used	17
1.9	list of integrals with no known antiderivative	18
1.10	List of integrals solved by CAS but has no known antiderivative	18
1.11	list of integrals solved by CAS but failed verification	18
1.12	Timing	19
1.13	Verification	19
1.14	Important notes about some of the results	20
1.15	Current tree layout of integration tests	23
1.16	Design of the test system	24

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [99]. This is test number [202].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (99)	0.00 (0)
Mathematica	100.00 (99)	0.00 (0)
Fricas	91.92 (91)	8.08 (8)
Maple	87.88 (87)	12.12 (12)
Maxima	69.70 (69)	30.30 (30)
Giac	52.53 (52)	47.47 (47)
Sympy	34.34 (34)	65.66 (65)
Mupad	30.30 (30)	69.70 (69)
Reduce	29.29 (29)	70.71 (70)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	93.939	0.000	0.000	6.061
Mathematica	87.879	0.000	6.061	6.061
Fricas	81.818	0.000	4.040	14.141
Maple	62.626	5.051	14.141	18.182
Giac	24.242	7.071	15.152	53.535
Sympy	22.222	6.061	0.000	71.717
Maxima	18.182	3.030	42.424	36.364
Mupad	0.000	24.242	0.000	75.758
Reduce	0.000	23.232	0.000	76.768

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	8	100.00	0.00	0.00
Maple	12	100.00	0.00	0.00
Maxima	30	60.00	0.00	40.00
Giac	47	100.00	0.00	0.00
Sympy	65	100.00	0.00	0.00
Mupad	69	0.00	100.00	0.00
Reduce	70	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.08
Maxima	0.18
Reduce	0.18
Rubi	0.38
Mathematica	0.52
Giac	0.55
Maple	1.28
Sympy	2.37
Mupad	29.09

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	32.23	0.88	26.50	0.83
Reduce	55.38	0.99	33.00	0.82
Fricas	75.74	0.85	62.00	0.81
Sympy	87.74	1.40	45.00	1.13
Mathematica	90.46	0.93	81.00	0.96
Rubi	107.33	1.02	87.00	1.00
Maxima	109.64	1.22	73.00	1.00
Giac	114.27	1.29	65.50	1.16
Maple	145.46	1.23	64.00	0.92

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{63, 64, 66, 68, 88, 89}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {17, 22, 61}

Mathematica {67}

Maple {12, 14, 19, 21}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	26
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	55

2.1 List of integrals sorted by grade for each CAS

Rubi	26
Mma	26
Maple	27
Fricas	27
Maxima	28
Giac	28
Mupad	28
Sympy	29
Reduce	29

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 91, 92, 96, 97 }

B grade { }

C grade { 90, 93, 94, 95, 98, 99 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 15, 16, 17, 18, 20, 22, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 69, 70, 71, 72, 79, 80, 81, 82, 83, 84, 85, 86, 87, 91, 92, 97 }

B grade { 90, 93, 94, 95, 96 }

C grade { 12, 14, 19, 21, 23, 24, 25, 26, 27, 28, 73, 76, 98, 99 }

F normal fail { 29, 30, 31, 32, 33, 34, 65, 67, 74, 75, 77, 78 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 69, 70, 71, 72, 79, 80, 81, 82, 83, 84, 85, 86, 87, 90, 91, 92, 95, 96, 97 }

B grade { }

C grade { 93, 94, 98, 99 }

F normal fail { 65, 67, 73, 74, 75, 76, 77, 78 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 3, 8, 10, 15, 17, 22, 37, 43, 45, 46, 47, 48, 61, 62, 91, 92, 97 }

B grade { 90, 95, 96 }

C grade { 2, 4, 5, 6, 7, 9, 11, 12, 13, 14, 16, 18, 19, 20, 21, 35, 36, 38, 39, 40, 41, 42, 44, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 69, 70, 71, 72, 85, 86, 87 }

F normal fail { 65, 67, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 93, 94, 98, 99 }

F(-1) timedout fail { }

F(-2) exception fail { 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 }

Giac

A grade { 1, 3, 5, 8, 10, 12, 15, 17, 19, 22, 37, 38, 43, 45, 46, 47, 48, 61, 62, 90, 91, 92, 96, 97 }

B grade { 7, 14, 21, 35, 36, 39, 95 }

C grade { 2, 4, 9, 11, 16, 18, 49, 50, 51, 55, 56, 57, 85, 86, 87 }

F normal fail { 6, 13, 20, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 40, 41, 42, 44, 52, 53, 54, 58, 59, 60, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 93, 94, 98, 99 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 3, 4, 8, 10, 15, 17, 22, 37, 38, 39, 42, 43, 45, 46, 47, 48, 61, 62, 85, 86, 87, 92, 97 }

C grade { }

F normal fail { }

F(-1) timedout fail { 2, 5, 6, 7, 9, 11, 12, 13, 14, 16, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 40, 41, 44, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 90, 91, 93, 94, 95, 96, 98, 99 }

F(-2) exception fail { }

Sympy

A grade { 1, 3, 4, 8, 11, 15, 17, 18, 22, 36, 37, 38, 39, 43, 46, 47, 48, 62, 90, 91, 92, 97 }

B grade { 2, 9, 10, 16, 45, 61 }

C grade { }

F normal fail { 5, 6, 7, 12, 13, 14, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 40, 41, 42, 44, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 93, 94, 95, 96, 98, 99 }

F(-1) timedout fail { }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 3, 8, 10, 15, 17, 22, 37, 38, 39, 43, 45, 46, 47, 48, 61, 62, 90, 91, 92, 95, 96, 97 }

C grade { }

F normal fail { 2, 4, 5, 6, 7, 9, 11, 12, 13, 14, 16, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 40, 41, 42, 44, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 93, 94, 98, 99 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	32	29	31	27	27	36	45	27	27
N.S.	1	0.94	0.85	0.91	0.79	0.79	1.06	1.32	0.79	0.79
time (sec)	N/A	0.265	0.060	1.467	0.035	0.085	0.199	0.355	0.169	0.141

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	93	82	58	67	72	209	135	14	0
N.S.	1	1.02	0.90	0.64	0.74	0.79	2.30	1.48	0.15	0.00
time (sec)	N/A	0.309	0.167	1.359	0.040	0.079	0.823	0.362	0.164	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	19	13	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	1.27	0.87	0.87	0.87
time (sec)	N/A	0.201	0.006	1.076	0.032	0.076	0.077	0.346	0.161	0.090

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	57	44	48	61	61	95	10	51
N.S.	1	1.00	0.81	0.63	0.69	0.87	0.87	1.36	0.14	0.73
time (sec)	N/A	0.237	0.102	0.492	0.035	0.079	0.170	0.351	0.197	42.922

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	22	43	21	0	21	14	0
N.S.	1	1.00	0.96	0.88	1.72	0.84	0.00	0.84	0.56	0.00
time (sec)	N/A	0.236	0.060	0.686	0.103	0.073	0.000	0.351	0.175	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	86	81	57	73	70	0	0	29	0
N.S.	1	1.08	1.01	0.71	0.91	0.88	0.00	0.00	0.36	0.00
time (sec)	N/A	0.290	0.190	0.635	0.182	0.078	0.000	0.000	0.197	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	40	42	39	48	40	0	87	36	0
N.S.	1	0.95	1.00	0.93	1.14	0.95	0.00	2.07	0.86	0.00
time (sec)	N/A	0.418	0.082	0.685	0.112	0.075	0.000	0.391	0.163	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	55	40	42	42	45	78	76	48	41
N.S.	1	1.08	0.78	0.82	0.82	0.88	1.53	1.49	0.94	0.80
time (sec)	N/A	0.240	0.137	1.204	0.036	0.077	0.283	0.340	0.179	0.208

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	87	63	90	84	201	118	128	0
N.S.	1	1.00	0.96	0.69	0.99	0.92	2.21	1.30	1.41	0.00
time (sec)	N/A	0.254	0.179	1.119	0.116	0.080	1.151	0.397	0.173	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	35	27	23	23	28	60	26	28	22
N.S.	1	1.13	0.87	0.74	0.74	0.90	1.94	0.84	0.90	0.71
time (sec)	N/A	0.215	0.043	1.123	0.035	0.081	0.103	0.338	0.162	0.076

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	67	45	70	59	56	82	12	0
N.S.	1	1.00	0.96	0.64	1.00	0.84	0.80	1.17	0.17	0.00
time (sec)	N/A	0.203	0.061	0.960	0.109	0.079	0.337	0.266	0.160	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	C	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	34	68	51	31	0	35	16	0
N.S.	1	1.00	0.92	1.84	1.38	0.84	0.00	0.95	0.43	0.00
time (sec)	N/A	0.211	0.116	1.329	0.102	0.080	0.000	0.355	0.184	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	84	76	62	83	66	0	0	16	0
N.S.	1	1.11	1.00	0.82	1.09	0.87	0.00	0.00	0.21	0.00
time (sec)	N/A	0.371	0.176	0.984	0.184	0.085	0.000	0.000	0.170	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	C	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	50	97	61	48	0	107	16	0
N.S.	1	1.00	0.88	1.70	1.07	0.84	0.00	1.88	0.28	0.00
time (sec)	N/A	0.287	0.135	1.358	0.107	0.079	0.000	0.341	0.172	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	82	55	66	58	58	92	92	67	66
N.S.	1	1.04	0.70	0.84	0.73	0.73	1.16	1.16	0.85	0.84
time (sec)	N/A	0.385	0.166	1.734	0.040	0.076	0.374	0.345	0.177	0.430

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	188	188	160	130	143	148	439	259	239	0
N.S.	1	1.00	0.85	0.69	0.76	0.79	2.34	1.38	1.27	0.00
time (sec)	N/A	0.378	0.471	1.673	0.124	0.085	1.940	0.351	0.185	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	25	33	26	27	25	44	26	27	28
N.S.	1	0.76	1.00	0.79	0.82	0.76	1.33	0.79	0.82	0.85
time (sec)	N/A	0.231	0.021	2.111	0.035	0.076	0.147	0.350	0.174	41.510

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	153	153	116	101	112	121	129	185	12	0
N.S.	1	1.00	0.76	0.66	0.73	0.79	0.84	1.21	0.08	0.00
time (sec)	N/A	0.296	0.253	1.425	0.111	0.080	0.484	0.387	0.168	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	C	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	50	125	89	47	0	47	16	0
N.S.	1	1.00	0.91	2.27	1.62	0.85	0.00	0.85	0.29	0.00
time (sec)	N/A	0.257	0.163	2.387	0.138	0.073	0.000	0.349	0.183	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	172	166	128	152	136	0	0	16	0
N.S.	1	1.02	0.99	0.76	0.90	0.81	0.00	0.00	0.10	0.00
time (sec)	N/A	0.360	0.734	1.590	0.240	0.083	0.000	0.000	0.189	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	C	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	90	185	98	80	0	185	16	0
N.S.	1	1.00	0.99	2.03	1.08	0.88	0.00	2.03	0.18	0.00
time (sec)	N/A	0.400	0.210	2.298	0.139	0.073	0.000	0.364	0.163	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	37	54	50	55	51	94	52	51	55
N.S.	1	0.55	0.81	0.75	0.82	0.76	1.40	0.78	0.76	0.82
time (sec)	N/A	0.240	0.103	7.716	0.038	0.075	0.640	0.348	0.179	0.793

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	113	113	229	0	64	0	0	16	0
N.S.	1	1.02	1.02	2.06	0.00	0.58	0.00	0.00	0.14	0.00
time (sec)	N/A	0.313	0.214	0.961	0.000	0.084	0.000	0.000	0.182	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	113	111	290	0	62	0	0	14	0
N.S.	1	1.02	1.00	2.61	0.00	0.56	0.00	0.00	0.13	0.00
time (sec)	N/A	0.303	0.180	0.858	0.000	0.083	0.000	0.000	0.169	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	89	290	0	48	0	0	13	0
N.S.	1	1.00	1.10	3.58	0.00	0.59	0.00	0.00	0.16	0.00
time (sec)	N/A	0.249	0.112	0.858	0.000	0.085	0.000	0.000	0.178	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	89	338	0	48	0	0	15	0
N.S.	1	1.00	1.10	4.17	0.00	0.59	0.00	0.00	0.19	0.00
time (sec)	N/A	0.244	0.088	0.812	0.000	0.086	0.000	0.000	0.168	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	104	114	338	0	66	0	0	41	0
N.S.	1	1.06	1.16	3.45	0.00	0.67	0.00	0.00	0.42	0.00
time (sec)	N/A	0.288	0.225	0.816	0.000	0.087	0.000	0.000	0.195	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	108	117	358	0	75	0	0	49	0
N.S.	1	1.04	1.12	3.44	0.00	0.72	0.00	0.00	0.47	0.00
time (sec)	N/A	0.290	0.197	0.840	0.000	0.092	0.000	0.000	0.180	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	132	135	142	0	0	92	0	0	141	0
N.S.	1	1.02	1.08	0.00	0.00	0.70	0.00	0.00	1.07	0.00
time (sec)	N/A	0.354	0.780	0.000	0.000	0.090	0.000	0.000	0.183	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	132	135	142	0	0	91	0	0	140	0
N.S.	1	1.02	1.08	0.00	0.00	0.69	0.00	0.00	1.06	0.00
time (sec)	N/A	0.310	0.684	0.000	0.000	0.086	0.000	0.000	0.189	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	100	102	99	0	0	68	0	0	15	0
N.S.	1	1.02	0.99	0.00	0.00	0.68	0.00	0.00	0.15	0.00
time (sec)	N/A	0.308	0.570	0.000	0.000	0.086	0.000	0.000	0.195	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	96	102	99	0	0	62	0	0	17	0
N.S.	1	1.06	1.03	0.00	0.00	0.65	0.00	0.00	0.18	0.00
time (sec)	N/A	0.254	0.520	0.000	0.000	0.082	0.000	0.000	0.180	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	117	128	137	0	0	79	0	0	20	0
N.S.	1	1.09	1.17	0.00	0.00	0.68	0.00	0.00	0.17	0.00
time (sec)	N/A	0.332	0.573	0.000	0.000	0.103	0.000	0.000	0.172	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	116	122	137	0	0	85	0	0	20	0
N.S.	1	1.05	1.18	0.00	0.00	0.73	0.00	0.00	0.17	0.00
time (sec)	N/A	0.394	0.630	0.000	0.000	0.089	0.000	0.000	0.170	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	32	31	39	57	33	0	132	19	0
N.S.	1	1.03	1.00	1.26	1.84	1.06	0.00	4.26	0.61	0.00
time (sec)	N/A	0.406	0.037	1.056	0.087	0.077	0.000	0.376	0.176	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	21	43	20	15	41	26	0
N.S.	1	1.00	1.00	1.05	2.15	1.00	0.75	2.05	1.30	0.00
time (sec)	N/A	0.239	0.055	1.082	0.075	0.073	0.412	0.354	0.183	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	14	13	15	15	15	15	13
N.S.	1	1.00	1.00	1.08	1.00	1.15	1.15	1.15	1.15	1.00
time (sec)	N/A	0.198	0.005	0.600	0.025	0.070	0.257	0.322	0.198	42.482

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	29	35	51	33	31	49	34	30
N.S.	1	1.00	0.97	1.17	1.70	1.10	1.03	1.63	1.13	1.00
time (sec)	N/A	0.265	0.056	1.111	0.074	0.067	0.361	0.342	0.230	42.790

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	51	46	47	50	43	46	107	52	47
N.S.	1	1.11	1.00	1.02	1.09	0.93	1.00	2.33	1.13	1.02
time (sec)	N/A	0.355	0.007	1.123	0.085	0.070	0.483	0.392	0.257	43.466

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	87	80	57	127	73	0	0	60	0
N.S.	1	1.10	1.01	0.72	1.61	0.92	0.00	0.00	0.76	0.00
time (sec)	N/A	0.331	0.191	0.770	0.088	0.084	0.000	0.000	0.217	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	22	43	21	0	0	28	0
N.S.	1	1.00	0.96	0.88	1.72	0.84	0.00	0.00	1.12	0.00
time (sec)	N/A	0.234	0.052	0.833	0.076	0.077	0.000	0.000	0.175	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	62	48	98	65	0	0	33	55
N.S.	1	1.00	0.84	0.65	1.32	0.88	0.00	0.00	0.45	0.74
time (sec)	N/A	0.272	0.119	0.859	0.081	0.074	0.000	0.000	0.185	42.565

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	17	22	17	17	13
N.S.	1	1.00	1.00	0.93	0.87	1.13	1.47	1.13	1.13	0.87
time (sec)	N/A	0.198	0.005	0.667	0.032	0.074	0.466	0.353	0.174	42.110

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	99	88	64	74	84	0	0	18	0
N.S.	1	1.02	0.91	0.66	0.76	0.87	0.00	0.00	0.19	0.00
time (sec)	N/A	0.333	0.182	0.911	0.082	0.079	0.000	0.000	0.163	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	28	18	14	12	13	39	12	10	12
N.S.	1	1.47	0.95	0.74	0.63	0.68	2.05	0.63	0.53	0.63
time (sec)	N/A	0.196	0.037	0.340	0.031	0.078	0.097	0.369	0.170	41.018

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	5	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.62	0.75
time (sec)	N/A	0.188	0.005	0.237	0.025	0.074	0.097	0.349	0.192	0.034

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	21	22	17	16	16	20	16	13	16
N.S.	1	0.95	1.00	0.77	0.73	0.73	0.91	0.73	0.59	0.73
time (sec)	N/A	0.230	0.027	0.299	0.031	0.076	0.089	0.360	0.182	40.801

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	41	31	34	23	24	51	23	21	23
N.S.	1	1.14	0.86	0.94	0.64	0.67	1.42	0.64	0.58	0.64
time (sec)	N/A	0.212	0.033	0.300	0.028	0.091	0.090	0.323	0.172	41.691

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	235	267	165	196	136	145	0	241	112	0
N.S.	1	1.14	0.70	0.83	0.58	0.62	0.00	1.03	0.48	0.00
time (sec)	N/A	1.395	0.698	0.521	0.057	0.086	0.000	0.367	0.173	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	186	141	131	112	118	0	193	80	0
N.S.	1	1.10	0.83	0.78	0.66	0.70	0.00	1.14	0.47	0.00
time (sec)	N/A	0.947	0.400	0.479	0.043	0.085	0.000	0.369	0.170	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	102	94	64	73	78	0	143	15	0
N.S.	1	1.03	0.95	0.65	0.74	0.79	0.00	1.44	0.15	0.00
time (sec)	N/A	0.532	0.176	0.513	0.039	0.087	0.000	0.374	0.174	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	122	110	78	74	96	0	0	41	0
N.S.	1	1.11	1.00	0.71	0.67	0.87	0.00	0.00	0.37	0.00
time (sec)	N/A	0.627	0.274	0.488	0.188	0.081	0.000	0.000	0.166	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	184	196	180	129	76	134	0	0	49	0
N.S.	1	1.07	0.98	0.70	0.41	0.73	0.00	0.00	0.27	0.00
time (sec)	N/A	0.981	0.261	0.510	0.196	0.091	0.000	0.000	0.171	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	272	238	180	76	164	0	0	53	0
N.S.	1	1.09	0.95	0.72	0.30	0.66	0.00	0.00	0.21	0.00
time (sec)	N/A	1.415	0.355	0.504	0.197	0.092	0.000	0.000	0.185	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	311	328	174	219	161	184	0	224	0	0
N.S.	1	1.05	0.56	0.70	0.52	0.59	0.00	0.72	0.00	0.00
time (sec)	N/A	0.852	0.729	0.664	0.131	0.095	0.000	0.374	0.201	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	218	226	148	145	137	144	0	176	220	0
N.S.	1	1.04	0.68	0.67	0.63	0.66	0.00	0.81	1.01	0.00
time (sec)	N/A	0.613	0.491	0.638	0.127	0.095	0.000	0.381	0.195	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	109	103	67	96	90	0	124	152	0
N.S.	1	1.06	1.00	0.65	0.93	0.87	0.00	1.20	1.48	0.00
time (sec)	N/A	0.372	0.217	0.645	0.126	0.092	0.000	0.376	0.186	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	151	116	87	88	100	0	0	20	0
N.S.	1	1.30	1.00	0.75	0.76	0.86	0.00	0.00	0.17	0.00
time (sec)	N/A	0.456	0.285	0.635	0.190	0.097	0.000	0.000	0.185	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	228	234	185	146	90	154	0	0	0	0
N.S.	1	1.03	0.81	0.64	0.39	0.68	0.00	0.00	0.00	0.00
time (sec)	N/A	0.991	0.303	0.645	0.186	0.105	0.000	0.000	0.242	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	328	364	249	207	90	192	0	0	0	0
N.S.	1	1.11	0.76	0.63	0.27	0.59	0.00	0.00	0.00	0.00
time (sec)	N/A	0.979	0.442	0.643	0.196	0.102	0.000	0.000	0.259	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	100	66	58	47	48	513	47	59	62
N.S.	1	1.16	0.77	0.67	0.55	0.56	5.97	0.55	0.69	0.72
time (sec)	N/A	0.490	0.109	1.067	0.034	0.078	0.757	0.365	0.181	41.253

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	7	6	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.88	0.75	0.75	0.75
time (sec)	N/A	0.195	0.006	0.237	0.032	0.080	4.946	0.346	0.170	41.272

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	20	20	17	20	23	20
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.94	1.11	1.28	1.11
time (sec)	N/A	0.186	1.608	0.302	0.986	0.084	7.628	1.136	0.182	41.034

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	22	22	19	22	24	22
N.S.	1	1.00	1.10	1.00	1.10	1.10	0.95	1.10	1.20	1.10
time (sec)	N/A	0.190	1.688	0.276	1.522	0.092	24.433	4.846	0.172	41.097

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	89	0	0	0	0	0	29	0
N.S.	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.338	0.226	0.000	0.000	0.000	0.000	0.000	0.193	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	24	22	24	24	20	24	33	24
N.S.	1	1.00	1.09	1.00	1.09	1.09	0.91	1.09	1.50	1.09
time (sec)	N/A	0.249	1.876	0.420	1.037	0.087	7.542	1.035	0.173	40.772

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	149	0	0	0	0	0	30	0
N.S.	1	1.00	1.14	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.411	0.831	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	26	24	26	26	22	26	34	26
N.S.	1	1.00	1.08	1.00	1.08	1.08	0.92	1.08	1.42	1.08
time (sec)	N/A	0.251	1.960	0.365	1.544	0.087	23.274	4.626	0.180	40.573

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	24	25	90	24	0	0	14	0
N.S.	1	1.00	0.92	0.96	3.46	0.92	0.00	0.00	0.54	0.00
time (sec)	N/A	0.252	0.080	0.927	0.167	0.078	0.000	0.000	0.169	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	37	40	99	35	0	0	16	0
N.S.	1	1.00	0.86	0.93	2.30	0.81	0.00	0.00	0.37	0.00
time (sec)	N/A	0.230	0.165	1.104	0.273	0.078	0.000	0.000	0.171	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	53	52	180	51	0	0	16	0
N.S.	1	1.00	0.79	0.78	2.69	0.76	0.00	0.00	0.24	0.00
time (sec)	N/A	0.271	0.192	1.645	0.337	0.084	0.000	0.000	0.174	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	66	66	189	62	0	0	16	0
N.S.	1	1.00	0.84	0.84	2.39	0.78	0.00	0.00	0.20	0.00
time (sec)	N/A	0.283	0.212	2.771	0.287	0.083	0.000	0.000	0.162	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	92	75	0	0	0	0	17	0
N.S.	1	1.00	1.11	0.90	0.00	0.00	0.00	0.00	0.20	0.00
time (sec)	N/A	0.221	0.093	0.407	0.000	0.000	0.000	0.000	0.185	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	94	0	0	0	0	0	12	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	0.264	0.272	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	179	179	173	0	0	0	0	0	12	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.299	0.276	0.000	0.000	0.000	0.000	0.000	0.175	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	105	115	111	0	0	0	0	89	0
N.S.	1	1.00	1.10	1.06	0.00	0.00	0.00	0.00	0.85	0.00
time (sec)	N/A	0.284	0.224	0.798	0.000	0.000	0.000	0.000	0.178	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	141	141	129	0	0	0	0	0	16	0
N.S.	1	1.00	0.91	0.00	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	0.346	0.539	0.000	0.000	0.000	0.000	0.000	0.169	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	229	229	221	0	0	0	0	0	16	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.07	0.00
time (sec)	N/A	0.428	0.575	0.000	0.000	0.000	0.000	0.000	0.174	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	42	45	45	0	45	0	0	51	0
N.S.	1	0.89	0.96	0.96	0.00	0.96	0.00	0.00	1.09	0.00
time (sec)	N/A	0.425	0.102	1.277	0.000	0.080	0.000	0.000	0.163	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	69	53	65	0	53	0	0	21	0
N.S.	1	0.99	0.76	0.93	0.00	0.76	0.00	0.00	0.30	0.00
time (sec)	N/A	0.301	0.267	2.706	0.000	0.083	0.000	0.000	0.161	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	113	95	101	0	85	0	0	21	0
N.S.	1	0.99	0.83	0.89	0.00	0.75	0.00	0.00	0.18	0.00
time (sec)	N/A	0.407	0.246	9.819	0.000	0.092	0.000	0.000	0.183	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	65	70	65	0	70	0	0	61	0
N.S.	1	0.83	0.90	0.83	0.00	0.90	0.00	0.00	0.78	0.00
time (sec)	N/A	0.525	0.144	1.293	0.000	0.079	0.000	0.000	0.168	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	95	82	89	0	87	0	0	819	0
N.S.	1	0.98	0.85	0.92	0.00	0.90	0.00	0.00	8.44	0.00
time (sec)	N/A	0.342	0.240	2.723	0.000	0.087	0.000	0.000	0.174	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	165	141	144	0	127	0	0	0	0
N.S.	1	0.99	0.84	0.86	0.00	0.76	0.00	0.00	0.00	0.00
time (sec)	N/A	0.492	0.381	9.918	0.000	0.095	0.000	0.000	0.182	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	91	76	131	258	112	0	159	23	80
N.S.	1	0.92	0.77	1.32	2.61	1.13	0.00	1.61	0.23	0.81
time (sec)	N/A	0.244	0.295	1.559	0.381	0.079	0.000	0.346	0.181	0.171

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	45	42	63	199	63	0	119	46	39
N.S.	1	0.96	0.89	1.34	4.23	1.34	0.00	2.53	0.98	0.83
time (sec)	N/A	0.206	0.185	1.424	0.368	0.084	0.000	0.350	0.181	40.341

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	36	84	40	0	55	19	32
N.S.	1	1.00	1.00	1.24	2.90	1.38	0.00	1.90	0.66	1.10
time (sec)	N/A	0.164	0.024	0.660	0.125	0.079	0.000	0.331	0.179	0.065

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	23	20	14	23	14
N.S.	1	1.00	1.17	1.00	1.17	1.92	1.67	1.17	1.92	1.17
time (sec)	N/A	0.162	2.459	0.171	0.190	0.070	0.950	0.370	0.167	40.388

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	14	23	22	14	38	14
N.S.	1	1.00	1.17	1.00	1.17	1.92	1.83	1.17	3.17	1.17
time (sec)	N/A	0.165	4.242	0.172	0.191	0.069	0.947	0.366	0.192	40.227

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	346	348	224	825	672	103	269	474	192	0
N.S.	1	1.01	0.65	2.38	1.94	0.30	0.78	1.37	0.55	0.00
time (sec)	N/A	0.496	1.017	1.441	0.062	0.084	0.338	0.329	0.174	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	171	71	297	263	67	151	166	100	0
N.S.	1	1.02	0.43	1.78	1.57	0.40	0.90	0.99	0.60	0.00
time (sec)	N/A	0.318	0.381	1.491	0.041	0.080	0.210	0.331	0.178	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	51	48	61	60	42	65	42	39	42
N.S.	1	0.94	0.89	1.13	1.11	0.78	1.20	0.78	0.72	0.78
time (sec)	N/A	0.271	0.155	0.615	0.031	0.081	0.195	0.284	0.176	41.161

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	150	145	271	0	149	0	0	17	0
N.S.	1	1.19	1.15	2.15	0.00	1.18	0.00	0.00	0.13	0.00
time (sec)	N/A	0.424	1.371	1.079	0.000	0.089	0.000	0.000	0.187	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	184	194	240	714	0	210	0	0	17	0
N.S.	1	1.05	1.30	3.88	0.00	1.14	0.00	0.00	0.09	0.00
time (sec)	N/A	0.542	1.457	1.188	0.000	0.092	0.000	0.000	0.180	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	B	A	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	537	540	382	1809	1349	182	0	1098	357	0
N.S.	1	1.01	0.71	3.37	2.51	0.34	0.00	2.04	0.66	0.00
time (sec)	N/A	0.662	1.258	1.549	0.092	0.083	0.000	0.357	0.175	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	261	265	117	655	523	110	0	370	182	0
N.S.	1	1.02	0.45	2.51	2.00	0.42	0.00	1.42	0.70	0.00
time (sec)	N/A	0.411	0.550	1.727	0.052	0.087	0.000	0.361	0.185	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	65	131	118	57	94	81	68	68
N.S.	1	1.00	0.76	1.54	1.39	0.67	1.11	0.95	0.80	0.80
time (sec)	N/A	0.369	0.213	0.652	0.039	0.075	0.263	0.335	0.192	41.290

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	234	262	243	279	0	287	0	0	18	0
N.S.	1	1.12	1.04	1.19	0.00	1.23	0.00	0.00	0.08	0.00
time (sec)	N/A	0.656	11.091	0.947	0.000	0.097	0.000	0.000	0.195	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	332	334	138	931	0	406	0	0	18	0
N.S.	1	1.01	0.42	2.80	0.00	1.22	0.00	0.00	0.05	0.00
time (sec)	N/A	0.812	0.825	1.136	0.000	0.114	0.000	0.000	0.217	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [54] had the largest ratio of [1.7500000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	7	6	0.94	12	0.500
2	A	4	4	1.02	12	0.333
3	A	4	3	1.00	10	0.300
4	A	3	3	1.00	8	0.375
5	A	3	3	1.00	12	0.250
6	A	4	4	1.08	12	0.333
7	A	10	9	0.95	12	0.750
8	A	5	4	1.08	14	0.286
9	A	2	2	1.00	14	0.143
10	A	5	4	1.13	12	0.333
11	A	2	2	1.00	10	0.200
12	A	2	2	1.00	14	0.143
13	A	6	6	1.11	14	0.429
14	A	2	2	1.00	14	0.143
15	A	9	8	1.04	14	0.571
16	A	2	2	1.00	14	0.143
17	A	5	4	0.76	12	0.333
18	A	2	2	1.00	10	0.200
19	A	2	2	1.00	14	0.143
20	A	3	3	1.02	14	0.214
21	A	2	2	1.00	14	0.143

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	5	4	0.55	12	0.333
23	A	3	3	1.02	14	0.214
24	A	3	3	1.02	14	0.214
25	A	2	2	1.00	14	0.143
26	A	2	2	1.00	14	0.143
27	A	3	3	1.06	14	0.214
28	A	3	3	1.04	14	0.214
29	A	4	3	1.02	16	0.188
30	A	4	3	1.02	16	0.188
31	A	4	3	1.02	16	0.188
32	A	4	3	1.06	16	0.188
33	A	4	3	1.09	16	0.188
34	A	7	6	1.05	16	0.375
35	A	10	9	1.03	8	1.125
36	A	3	3	1.00	12	0.250
37	A	4	3	1.00	12	0.250
38	A	7	6	1.00	12	0.500
39	A	9	8	1.11	12	0.667
40	A	6	5	1.10	8	0.625
41	A	3	3	1.00	12	0.250
42	A	5	4	1.00	12	0.333
43	A	4	3	1.00	12	0.250
44	A	6	5	1.02	12	0.417
45	A	5	4	1.47	14	0.286
46	A	4	3	1.00	12	0.250
47	A	7	6	0.95	6	1.000
48	A	5	4	1.14	8	0.500
49	A	27	26	1.14	16	1.625
50	A	19	18	1.10	16	1.125
51	A	12	11	1.03	16	0.688
52	A	14	13	1.11	16	0.812
53	A	22	21	1.07	16	1.312

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	29	28	1.09	16	1.750
55	A	14	13	1.05	18	0.722
56	A	11	10	1.04	18	0.556
57	A	5	4	1.06	18	0.222
58	A	8	7	1.30	18	0.389
59	A	18	17	1.03	18	0.944
60	A	17	16	1.11	18	0.889
61	A	13	12	1.16	8	1.500
62	A	4	3	1.00	12	0.250
63	N/A	1	0	1.00	18	0.000
64	N/A	1	0	1.00	20	0.000
65	A	5	4	1.00	20	0.200
66	N/A	2	0	1.00	22	0.000
67	A	7	6	1.00	22	0.273
68	N/A	2	0	1.00	24	0.000
69	A	3	3	1.00	12	0.250
70	A	2	2	1.00	14	0.143
71	A	2	2	1.00	14	0.143
72	A	2	2	1.00	14	0.143
73	A	2	2	1.00	8	0.250
74	A	2	2	1.00	10	0.200
75	A	2	2	1.00	10	0.200
76	A	2	2	1.00	12	0.167
77	A	2	2	1.00	14	0.143
78	A	2	2	1.00	14	0.143
79	A	10	9	0.89	16	0.562
80	A	2	2	0.99	18	0.111
81	A	2	2	0.99	18	0.111
82	A	12	11	0.83	16	0.688
83	A	2	2	0.98	18	0.111
84	A	2	2	0.99	18	0.111
85	A	3	2	0.92	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	3	2	0.96	10	0.200
87	A	1	1	1.00	8	0.125
88	N/A	1	0	1.00	12	0.000
89	N/A	1	0	1.00	12	0.000
90	A	3	2	1.01	18	0.111
91	A	3	2	1.02	16	0.125
92	A	7	6	0.94	14	0.429
93	A	3	2	1.19	18	0.111
94	A	6	5	1.05	18	0.278
95	A	3	2	1.01	18	0.111
96	A	3	2	1.02	16	0.125
97	A	9	8	1.00	14	0.571
98	A	3	2	1.12	18	0.111
99	A	6	5	1.01	18	0.278

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \cos(a + bx^2) dx$	63
3.2	$\int x^2 \cos(a + bx^2) dx$	69
3.3	$\int x \cos(a + bx^2) dx$	76
3.4	$\int \cos(a + bx^2) dx$	81
3.5	$\int \frac{\cos(a+bx^2)}{x} dx$	87
3.6	$\int \frac{\cos(a+bx^2)}{x^2} dx$	92
3.7	$\int \frac{\cos(a+bx^2)}{x^3} dx$	98
3.8	$\int x^3 \cos^2(a + bx^2) dx$	105
3.9	$\int x^2 \cos^2(a + bx^2) dx$	111
3.10	$\int x \cos^2(a + bx^2) dx$	117
3.11	$\int \cos^2(a + bx^2) dx$	123
3.12	$\int \frac{\cos^2(a+bx^2)}{x} dx$	129
3.13	$\int \frac{\cos^2(a+bx^2)}{x^2} dx$	134
3.14	$\int \frac{\cos^2(a+bx^2)}{x^3} dx$	140
3.15	$\int x^3 \cos^3(a + bx^2) dx$	145
3.16	$\int x^2 \cos^3(a + bx^2) dx$	152
3.17	$\int x \cos^3(a + bx^2) dx$	160
3.18	$\int \cos^3(a + bx^2) dx$	166
3.19	$\int \frac{\cos^3(a+bx^2)}{x} dx$	172
3.20	$\int \frac{\cos^3(a+bx^2)}{x^2} dx$	177
3.21	$\int \frac{\cos^3(a+bx^2)}{x^3} dx$	183
3.22	$\int x \cos^7(a + bx^2) dx$	189
3.23	$\int x^{5/2} \cos(a + bx^2) dx$	195
3.24	$\int x^{3/2} \cos(a + bx^2) dx$	201
3.25	$\int \sqrt{x} \cos(a + bx^2) dx$	207

3.26	$\int \frac{\cos(ax^2)}{\sqrt{x}} dx$	212
3.27	$\int \frac{\cos(ax^2)}{x^{3/2}} dx$	217
3.28	$\int \frac{\cos(ax^2)}{x^{5/2}} dx$	223
3.29	$\int x^{5/2} \cos^2(a + bx^2) dx$	229
3.30	$\int x^{3/2} \cos^2(a + bx^2) dx$	234
3.31	$\int \sqrt{x} \cos^2(a + bx^2) dx$	239
3.32	$\int \frac{\cos^2(ax^2)}{\sqrt{x}} dx$	244
3.33	$\int \frac{\cos^2(ax^2)}{x^{3/2}} dx$	249
3.34	$\int \frac{\cos^2(ax^2)}{x^{5/2}} dx$	254
3.35	$\int \cos\left(a + \frac{b}{x}\right) dx$	260
3.36	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$	267
3.37	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx$	272
3.38	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx$	277
3.39	$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx$	283
3.40	$\int \cos\left(a + \frac{b}{x^2}\right) dx$	289
3.41	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$	295
3.42	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$	300
3.43	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx$	306
3.44	$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$	311
3.45	$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$	317
3.46	$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$	322
3.47	$\int \cos(\sqrt{x}) dx$	327
3.48	$\int \cos^2(\sqrt{x}) dx$	332
3.49	$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx$	337
3.50	$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$	373
3.51	$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx$	390
3.52	$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx$	398
3.53	$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx$	406
3.54	$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx$	417
3.55	$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx$	428
3.56	$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$	442

3.57	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}} dx$	451
3.58	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{3/2}} dx$	457
3.59	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{5/2}} dx$	464
3.60	$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{7/2}} dx$	476
3.61	$\int \cos^3(\sqrt[3]{x}) dx$	487
3.62	$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx$	495
3.63	$\int (ex)^m (b \cos(c + dx^n))^p dx$	500
3.64	$\int (ex)^m (a + b \cos(c + dx^n))^p dx$	505
3.65	$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx$	510
3.66	$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$	515
3.67	$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx$	520
3.68	$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$	526
3.69	$\int \frac{\cos(ax^n)}{x} dx$	531
3.70	$\int \frac{\cos^2(ax^n)}{x} dx$	536
3.71	$\int \frac{\cos^3(ax^n)}{x} dx$	541
3.72	$\int \frac{\cos^4(ax^n)}{x} dx$	546
3.73	$\int \cos(a + bx^n) dx$	551
3.74	$\int \cos^2(a + bx^n) dx$	556
3.75	$\int \cos^3(a + bx^n) dx$	561
3.76	$\int x^m \cos(a + bx^n) dx$	566
3.77	$\int x^m \cos^2(a + bx^n) dx$	571
3.78	$\int x^m \cos^3(a + bx^n) dx$	576
3.79	$\int x^{-1-n} \cos(a + bx^n) dx$	581
3.80	$\int x^{-1-n} \cos^2(a + bx^n) dx$	587
3.81	$\int x^{-1-n} \cos^3(a + bx^n) dx$	592
3.82	$\int x^{-1-2n} \cos(a + bx^n) dx$	597
3.83	$\int x^{-1-2n} \cos^2(a + bx^n) dx$	604
3.84	$\int x^{-1-2n} \cos^3(a + bx^n) dx$	610
3.85	$\int x^2 \cos((a + bx)^2) dx$	616
3.86	$\int x \cos((a + bx)^2) dx$	622
3.87	$\int \cos((a + bx)^2) dx$	628
3.88	$\int \frac{\cos((a+bx)^2)}{x} dx$	633
3.89	$\int \frac{\cos((a+bx)^2)}{x^2} dx$	638
3.90	$\int x^2 \cos(a + b\sqrt{c + dx}) dx$	643
3.91	$\int x \cos(a + b\sqrt{c + dx}) dx$	651
3.92	$\int \cos(a + b\sqrt{c + dx}) dx$	657

3.93	$\int \frac{\cos(a+b\sqrt{c+dx})}{x} dx$	663
3.94	$\int \frac{\cos(a+b\sqrt{c+dx})}{x^2} dx$	669
3.95	$\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx$	676
3.96	$\int x \cos(a + b\sqrt[3]{c + dx}) dx$	686
3.97	$\int \cos(a + b\sqrt[3]{c + dx}) dx$	694
3.98	$\int \frac{\cos(a+b\sqrt[3]{c+dx})}{x} dx$	701
3.99	$\int \frac{\cos(a+b\sqrt[3]{c+dx})}{x^2} dx$	708

3.1 $\int x^3 \cos(a + bx^2) dx$

Optimal result	63
Mathematica [A] (verified)	63
Rubi [A] (verified)	64
Maple [A] (verified)	66
Fricas [A] (verification not implemented)	66
Sympy [A] (verification not implemented)	67
Maxima [A] (verification not implemented)	67
Giac [A] (verification not implemented)	67
Mupad [B] (verification not implemented)	68
Reduce [B] (verification not implemented)	68

Optimal result

Integrand size = 12, antiderivative size = 34

$$\int x^3 \cos(a + bx^2) dx = \frac{\cos(a + bx^2)}{2b^2} + \frac{x^2 \sin(a + bx^2)}{2b}$$

output `1/2*cos(b*x^2+a)/b^2+1/2*x^2*sin(b*x^2+a)/b`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int x^3 \cos(a + bx^2) dx = \frac{\cos(a + bx^2) + bx^2 \sin(a + bx^2)}{2b^2}$$

input `Integrate[x^3*Cos[a + b*x^2],x]`

output `(Cos[a + b*x^2] + b*x^2*Sin[a + b*x^2])/(2*b^2)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3861, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \cos(a + bx^2) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{2} \int x^2 \cos(bx^2 + a) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int x^2 \sin\left(bx^2 + a + \frac{\pi}{2}\right) dx^2 \\
 & \quad \downarrow \text{3777} \\
 & \frac{1}{2} \left(\frac{\int -\sin(bx^2 + a) dx^2}{b} + \frac{x^2 \sin(a + bx^2)}{b} \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \left(\frac{x^2 \sin(a + bx^2)}{b} - \frac{\int \sin(bx^2 + a) dx^2}{b} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(\frac{x^2 \sin(a + bx^2)}{b} - \frac{\int \sin(bx^2 + a) dx^2}{b} \right) \\
 & \quad \downarrow \text{3118} \\
 & \frac{1}{2} \left(\frac{\cos(a + bx^2)}{b^2} + \frac{x^2 \sin(a + bx^2)}{b} \right)
 \end{aligned}$$

input `Int[x^3*Cos[a + b*x^2], x]`

output $(\text{Cos}[a + b*x^2]/b^2 + (x^2*\text{Sin}[a + b*x^2])/b)/2$

Defintions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear} \\ \text{Q}[u, x]$

rule 3118 $\text{Int}[\text{sin}[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] \text{ ; FreeQ} \\ \{c, d\}, x]$

rule 3777 $\text{Int}[(c_.) + (d_.)*(x_)^(m_.)*\text{sin}[(e_.) + (f_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[(\\ -(c + d*x)^m)*(\text{Cos}[e + f*x]/f), x] + \text{Simp}[d*(m/f) \quad \text{Int}[(c + d*x)^(m - 1)*\text{C} \\ \text{os}[e + f*x], x], x] \text{ ; FreeQ}\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$

rule 3861 $\text{Int}[(a_.) + \text{Cos}[(c_.) + (d_.)*(x_)^(n_)]*(b_.)]^(p_.)*(x_)^(m_.), x_Symbol \\] \rightarrow \text{Simp}[1/n \quad \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*\text{Cos}[c + d*x])} \\ ^p, x], x, x^n], x] \text{ ; FreeQ}\{a, b, c, d, m, n, p\}, x] \ \&\& \ \text{IntegerQ}[\text{Simplify}[\\ (m + 1)/n]] \ \&\& \ (\text{EqQ}[p, 1] \ || \ \text{EqQ}[m, n - 1] \ || \ (\text{IntegerQ}[p] \ \&\& \ \text{GtQ}[\text{Simplify}[\\ (m + 1)/n], 0]))$

Maple [A] (verified)

Time = 1.47 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

method	result
default	$\frac{\cos(bx^2+a)}{2b^2} + \frac{x^2 \sin(bx^2+a)}{2b}$
risch	$\frac{\cos(bx^2+a)}{2b^2} + \frac{x^2 \sin(bx^2+a)}{2b}$
parallelrisch	$\frac{1+\tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)x^2b}{b^2\left(1+\tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^2\right)}$
norman	$\frac{\frac{1}{b^2} + \frac{x^2 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)}{b}}{1+\tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^2}$
orering	$\frac{5 \cos(bx^2+a)}{4b^2} - \frac{3x^2 \cos(bx^2+a) - 2x^4 b \sin(bx^2+a)}{4x^2 b^2}$
meijerg	$\frac{\cos(a)\sqrt{\pi}\left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(bx^2)}{2\sqrt{\pi}} + \frac{bx^2 \sin(bx^2)}{2\sqrt{\pi}}\right)}{b^2} - \frac{\sin(a)\sqrt{\pi}\left(-\frac{x^2 b \cos(bx^2)}{2\sqrt{\pi}} + \frac{\sin(bx^2)}{2\sqrt{\pi}}\right)}{b^2}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^3 \cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\sqrt{2}\sqrt{\pi}x^3 \sin(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{3\pi^2 \left(\cos(a)\left(\frac{2 \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)b^{\frac{3}{2}}\sqrt{2}x^3}{3\pi^{\frac{3}{2}}}\right)\right)}{3\pi^{\frac{3}{2}}}$

input `int(x^3*cos(b*x^2+a),x,method=_RETURNVERBOSE)`output `1/2*cos(b*x^2+a)/b^2+1/2*x^2*sin(b*x^2+a)/b`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int x^3 \cos(a + bx^2) dx = \frac{bx^2 \sin(bx^2 + a) + \cos(bx^2 + a)}{2b^2}$$

input `integrate(x^3*cos(b*x^2+a),x, algorithm="fricas")`output `1/2*(b*x^2*sin(b*x^2 + a) + cos(b*x^2 + a))/b^2`

Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int x^3 \cos(a + bx^2) dx = \begin{cases} \frac{x^2 \sin(a + bx^2)}{2b} + \frac{\cos(a + bx^2)}{2b^2} & \text{for } b \neq 0 \\ \frac{x^4 \cos(a)}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*cos(b*x**2+a),x)`output `Piecewise((x**2*sin(a + b*x**2)/(2*b) + cos(a + b*x**2)/(2*b**2), Ne(b, 0)), (x**4*cos(a)/4, True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int x^3 \cos(a + bx^2) dx = \frac{bx^2 \sin(bx^2 + a) + \cos(bx^2 + a)}{2b^2}$$

input `integrate(x^3*cos(b*x^2+a),x, algorithm="maxima")`output `1/2*(b*x^2*sin(b*x^2 + a) + cos(b*x^2 + a))/b^2`**Giac [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.32

$$\int x^3 \cos(a + bx^2) dx = -\frac{a \sin(bx^2 + a)}{2b^2} + \frac{(bx^2 + a) \sin(bx^2 + a) + \cos(bx^2 + a)}{2b^2}$$

input `integrate(x^3*cos(b*x^2+a),x, algorithm="giac")`output `-1/2*a*sin(b*x^2 + a)/b^2 + 1/2*((b*x^2 + a)*sin(b*x^2 + a) + cos(b*x^2 + a))/b^2`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int x^3 \cos(a + bx^2) dx = \frac{\cos(bx^2 + a) + bx^2 \sin(bx^2 + a)}{2b^2}$$

input `int(x^3*cos(a + b*x^2),x)`output `(cos(a + b*x^2) + b*x^2*sin(a + b*x^2))/(2*b^2)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int x^3 \cos(a + bx^2) dx = \frac{\cos(bx^2 + a) + \sin(bx^2 + a)bx^2}{2b^2}$$

input `int(x^3*cos(b*x^2+a),x)`output `(cos(a + b*x**2) + sin(a + b*x**2)*b*x**2)/(2*b**2)`

3.2 $\int x^2 \cos(a + bx^2) dx$

Optimal result	69
Mathematica [A] (verified)	69
Rubi [A] (verified)	70
Maple [A] (verified)	71
Fricas [A] (verification not implemented)	72
Sympy [B] (verification not implemented)	72
Maxima [C] (verification not implemented)	73
Giac [C] (verification not implemented)	73
Mupad [F(-1)]	74
Reduce [F]	75

Optimal result

Integrand size = 12, antiderivative size = 91

$$\int x^2 \cos(a + bx^2) dx = -\frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{2b^{3/2}} - \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)}{2b^{3/2}} + \frac{x \sin(a + bx^2)}{2b}$$

```
output -1/4*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)/b^(3/2)-
1/4*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)*sin(a)/b^(3/2)+1
/2*x*sin(b*x^2+a)/b
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.90

$$\int x^2 \cos(a + bx^2) dx = \frac{-\sqrt{2\pi} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a) + 2\sqrt{b}x \sin(a + bx^2)}{4b^{3/2}}$$

```
input Integrate[x^2*Cos[a + b*x^2],x]
```

output

$$\frac{(-(\sqrt{2\pi} \cos[a] \operatorname{FresnelS}[\sqrt{b} \sqrt{2/\pi} x]) - \sqrt{2\pi} \operatorname{FresnelC}[\sqrt{b} \sqrt{2/\pi} x] \sin[a] + 2\sqrt{b} x \sin[a + b x^2])}{(4 b^{3/2})}$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3867, 3834, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \cos(a + bx^2) dx \\ & \quad \downarrow \text{3867} \\ & \frac{x \sin(a + bx^2)}{2b} - \frac{\int \sin(bx^2 + a) dx}{2b} \\ & \quad \downarrow \text{3834} \\ & \frac{x \sin(a + bx^2)}{2b} - \frac{\sin(a) \int \cos(bx^2) dx + \cos(a) \int \sin(bx^2) dx}{2b} \\ & \quad \downarrow \text{3832} \\ & \frac{x \sin(a + bx^2)}{2b} - \frac{\sin(a) \int \cos(bx^2) dx + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}(\sqrt{b} \sqrt{\frac{2}{\pi}} x)}{\sqrt{b}}}{2b} \\ & \quad \downarrow \text{3833} \\ & \frac{x \sin(a + bx^2)}{2b} - \frac{\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}(\sqrt{b} \sqrt{\frac{2}{\pi}} x)}{\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}(\sqrt{b} \sqrt{\frac{2}{\pi}} x)}{\sqrt{b}}}{2b} \end{aligned}$$

input

$$\operatorname{Int}[x^2 \cos[a + b x^2], x]$$

output

$$\frac{-1/2 * ((\sqrt{\pi/2} \cos[a] \operatorname{FresnelS}[\sqrt{b} \sqrt{2/\pi} x]) / \sqrt{b} + (\sqrt{\pi/2} \operatorname{FresnelC}[\sqrt{b} \sqrt{2/\pi} x] \sin[a]) / \sqrt{b})}{b} + \frac{(x \sin[a + b x^2])}{(2 * b)}$$

Defintions of rubi rules used

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3834 `Int[Sin[(c_) + (d_.)*((e_.) + (f_.)*(x_))2], x_Symbol] := Simp[Sin[c] Int[Cos[d*(e + f*x)2], x] + Simp[Cos[c] Int[Sin[d*(e + f*x)2], x] /; FreeQ[{c, d, e, f}, x]`

rule 3867 `Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_))^(m_.), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(Sin[c + d*x^n]/(d*n)), x] - Simp[e^n*((m - n + 1)/(d*n)) Int[(e*x)^(m - n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]`

Maple [A] (verified)

Time = 1.36 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.64

method	result
default	$\frac{x \sin(bx^2+a)}{2b} - \frac{\sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) \right)}{4b^{\frac{3}{2}}}$
risch	$-\frac{ie^{-ia}\sqrt{\pi} \operatorname{erf}(\sqrt{ib}x)}{8b\sqrt{ib}} + \frac{ie^{ia}\sqrt{\pi} \operatorname{erf}(\sqrt{-ib}x)}{8b\sqrt{-ib}} + \frac{x \sin(bx^2+a)}{2b}$
meijerg	$\frac{\cos(a)\sqrt{\pi}\sqrt{2} \left(\frac{x\sqrt{2}(b^2)^{\frac{3}{4}} \sin(bx^2)}{2\sqrt{\pi}b} - \frac{(b^2)^{\frac{3}{4}} \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2b^{\frac{3}{2}}} \right)}{2(b^2)^{\frac{3}{4}}} - \frac{\sin(a)\sqrt{\pi}\sqrt{2} \left(-\frac{x\sqrt{2}\sqrt{b} \cos(bx^2)}{2\sqrt{\pi}} + \frac{\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2} \right)}{2b^{\frac{3}{2}}}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2 \cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\sqrt{2}\sqrt{\pi}x^2 \sin(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\sqrt{2}\pi^{\frac{3}{2}} \left(\cos(a) \left(\frac{\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)bx^2}{\pi} - \frac{\sqrt{b}\sqrt{2}x}{2\sqrt{\pi}} \right) \right)}{2\sqrt{b}}$

input `int(x^2*cos(b*x^2+a), x, method=_RETURNVERBOSE)`

output

```
1/2*x*sin(b*x^2+a)/b-1/4/b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)
*2^(1/2)/Pi^(1/2)*x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.79

$$\int x^2 \cos(a + bx^2) dx = \frac{\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) + \sqrt{2}\pi\sqrt{\frac{b}{\pi}}C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a) - 2bx\sin(bx^2 + a)}{4b^2}$$

input

```
integrate(x^2*cos(b*x^2+a),x, algorithm="fricas")
```

output

```
-1/4*(sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x*sqrt(b/pi)) + sqrt(2)*pi*sqrt(b/pi)*fresnel_cos(sqrt(2)*x*sqrt(b/pi))*sin(a) - 2*b*x*sin(b*x^2 + a))/b^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(90) = 180.

Time = 0.82 (sec) , antiderivative size = 209, normalized size of antiderivative = 2.30

$$\int x^2 \cos(a + bx^2) dx = \frac{b^{\frac{3}{2}}x^5\sqrt{\frac{1}{b}}\sin(a)\Gamma\left(\frac{3}{4}\right)\Gamma\left(\frac{5}{4}\right) {}_2F_3\left(\frac{3}{4}, \frac{5}{4} \mid -\frac{b^2x^4}{4}\right)}{8\Gamma\left(\frac{7}{4}\right)\Gamma\left(\frac{9}{4}\right)} - \frac{\sqrt{b}x^3\sqrt{\frac{1}{b}}\cos(a)\Gamma\left(\frac{1}{4}\right)\Gamma\left(\frac{3}{4}\right) {}_2F_3\left(\frac{1}{4}, \frac{3}{4} \mid -\frac{b^2x^4}{4}\right)}{8\Gamma\left(\frac{5}{4}\right)\Gamma\left(\frac{7}{4}\right)} - \frac{\sqrt{2}\sqrt{\pi}x^2\sqrt{\frac{1}{b}}\sin(a)S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)}{2} + \frac{\sqrt{2}\sqrt{\pi}x^2\sqrt{\frac{1}{b}}\cos(a)C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)}{2}$$

input `integrate(x**2*cos(b*x**2+a),x)`

output `b**(3/2)*x**5*sqrt(1/b)*sin(a)*gamma(3/4)*gamma(5/4)*hyper((3/4, 5/4), (3/2, 7/4, 9/4), -b**2*x**4/4)/(8*gamma(7/4)*gamma(9/4)) - sqrt(b)*x**3*sqrt(1/b)*cos(a)*gamma(1/4)*gamma(3/4)*hyper((1/4, 3/4), (1/2, 5/4, 7/4), -b**2*x**4/4)/(8*gamma(5/4)*gamma(7/4)) - sqrt(2)*sqrt(pi)*x**2*sqrt(1/b)*sin(a)*fresnels(sqrt(2)*sqrt(b)*x/sqrt(pi))/2 + sqrt(2)*sqrt(pi)*x**2*sqrt(1/b)*cos(a)*fresnelc(sqrt(2)*sqrt(b)*x/sqrt(pi))/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.74

$$\int x^2 \cos(a + bx^2) dx$$

$$= \frac{8b^2x \sin(bx^2 + a) + \sqrt{2}\sqrt{\pi} \left((-i+1) \cos(a) + (i-1) \sin(a) \right) \operatorname{erf}(\sqrt{i}bx) + ((i-1) \cos(a) - (i+1) \sin(a)) \operatorname{erf}(\sqrt{-i}bx)}{16b^3}$$

input `integrate(x^2*cos(b*x^2+a),x, algorithm="maxima")`

output `1/16*(8*b^2*x*sin(b*x^2 + a) + sqrt(2)*sqrt(pi)*((-I + 1)*cos(a) + (I - 1)*sin(a))*erf(sqrt(I*b)*x) + ((I - 1)*cos(a) - (I + 1)*sin(a))*erf(sqrt(-I*b)*x))/b^3`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.48

$$\int x^2 \cos(a + bx^2) dx = -\frac{ixe^{(ibx^2+ia)}}{4b} + \frac{ixe^{(-ibx^2-ia)}}{4b} - \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{(ia)}}{8b\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}} + \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{(-ia)}}{8b\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}}$$

input `integrate(x^2*cos(b*x^2+a),x, algorithm="giac")`

output `-1/4*I*x*e^(I*b*x^2 + I*a)/b + 1/4*I*x*e^(-I*b*x^2 - I*a)/b - 1/8*I*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(-I*b/abs(b) + 1)*sqrt(abs(b)))*e^(I*a)/(b*(-I*b/abs(b) + 1)*sqrt(abs(b))) + 1/8*I*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(I*b/abs(b) + 1)*sqrt(abs(b)))*e^(-I*a)/(b*(I*b/abs(b) + 1)*sqrt(abs(b)))`

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos(a + bx^2) dx = \int x^2 \cos(bx^2 + a) dx$$

input `int(x^2*cos(a + b*x^2),x)`

output `int(x^2*cos(a + b*x^2), x)`

Reduce [F]

$$\int x^2 \cos(a + bx^2) dx = \int \cos(bx^2 + a) x^2 dx$$

input `int(x^2*cos(b*x^2+a),x)`

output `int(cos(a + b*x**2)*x**2,x)`

3.3 $\int x \cos(a + bx^2) dx$

Optimal result	76
Mathematica [A] (verified)	76
Rubi [A] (verified)	77
Maple [A] (verified)	78
Fricas [A] (verification not implemented)	79
Sympy [A] (verification not implemented)	79
Maxima [A] (verification not implemented)	79
Giac [A] (verification not implemented)	80
Mupad [B] (verification not implemented)	80
Reduce [B] (verification not implemented)	80

Optimal result

Integrand size = 10, antiderivative size = 15

$$\int x \cos(a + bx^2) dx = \frac{\sin(a + bx^2)}{2b}$$

output

```
1/2*sin(b*x^2+a)/b
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int x \cos(a + bx^2) dx = \frac{\sin(a + bx^2)}{2b}$$

input

```
Integrate[x*Cos[a + b*x^2],x]
```

output

```
Sin[a + b*x^2]/(2*b)
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3861, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \cos (a + bx^2) dx \\ & \quad \downarrow \text{3861} \\ & \frac{1}{2} \int \cos (bx^2 + a) dx^2 \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2} \int \sin \left(bx^2 + a + \frac{\pi}{2} \right) dx^2 \\ & \quad \downarrow \text{3117} \\ & \frac{\sin (a + bx^2)}{2b} \end{aligned}$$

input `Int[x*Cos[a + b*x^2],x]`

output `Sin[a + b*x^2]/(2*b)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
  := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result
derivativdivides	$\frac{\sin(bx^2+a)}{2b}$
default	$\frac{\sin(bx^2+a)}{2b}$
risch	$\frac{\sin(bx^2+a)}{2b}$
parallelrisc	$\frac{\sin(bx^2+a)}{2b}$
norman	$\frac{\tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)}{b\left(1 + \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^2\right)}$
meijerg	$\frac{\cos(a)\sin(bx^2)}{2b} - \frac{\sin(a)\sqrt{\pi}\left(\frac{1}{\sqrt{\pi}} - \frac{\cos(bx^2)}{\sqrt{\pi}}\right)}{2b}$
orering	$\frac{\cos(bx^2+a)}{4x^2b^2} - \frac{\cos(bx^2+a) - 2\sin(bx^2+a)bx^2}{4x^2b^2}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x\cos(a)\text{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\sqrt{2}\sqrt{\pi}x\sin(a)\text{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\cos(a)\sqrt{2}\sqrt{\pi}\left(\frac{\text{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{\sqrt{\pi}} - \frac{\text{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{\sqrt{\pi}}\right)}{2\sqrt{b}}$

input

```
int(x*cos(b*x^2+a),x,method=_RETURNVERBOSE)
```

output

```
1/2*sin(b*x^2+a)/b
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int x \cos(a + bx^2) dx = \frac{\sin(bx^2 + a)}{2b}$$

input `integrate(x*cos(b*x^2+a),x, algorithm="fricas")`output `1/2*sin(b*x^2 + a)/b`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.27

$$\int x \cos(a + bx^2) dx = \begin{cases} \frac{\sin(a+bx^2)}{2b} & \text{for } b \neq 0 \\ \frac{x^2 \cos(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*cos(b*x**2+a),x)`output `Piecewise((sin(a + b*x**2)/(2*b), Ne(b, 0)), (x**2*cos(a)/2, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int x \cos(a + bx^2) dx = \frac{\sin(bx^2 + a)}{2b}$$

input `integrate(x*cos(b*x^2+a),x, algorithm="maxima")`output `1/2*sin(b*x^2 + a)/b`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int x \cos (a + b x^2) d x = \frac{\sin (b x^2 + a)}{2 b}$$

input `integrate(x*cos(b*x^2+a),x, algorithm="giac")`

output `1/2*sin(b*x^2 + a)/b`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int x \cos (a + b x^2) d x = \frac{\sin (b x^2 + a)}{2 b}$$

input `int(x*cos(a + b*x^2),x)`

output `sin(a + b*x^2)/(2*b)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int x \cos (a + b x^2) d x = \frac{\sin (b x^2 + a)}{2 b}$$

input `int(x*cos(b*x^2+a),x)`

output `sin(a + b*x**2)/(2*b)`

3.4 $\int \cos(a + bx^2) dx$

Optimal result	81
Mathematica [A] (verified)	81
Rubi [A] (verified)	82
Maple [A] (verified)	83
Fricas [A] (verification not implemented)	83
Sympy [A] (verification not implemented)	84
Maxima [C] (verification not implemented)	84
Giac [C] (verification not implemented)	85
Mupad [B] (verification not implemented)	85
Reduce [F]	86

Optimal result

Integrand size = 8, antiderivative size = 70

$$\int \cos(a + bx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)}{\sqrt{b}}$$

output

```
1/2*2^(1/2)*Pi^(1/2)*cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)/b^(1/2)-1/2*2^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)*sin(a)/b^(1/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.81

$$\int \cos(a + bx^2) dx = \frac{\sqrt{\frac{\pi}{2}}\left(\cos(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)\right)}{\sqrt{b}}$$

input

```
Integrate[Cos[a + b*x^2], x]
```

output

```
(Sqrt[Pi/2]*(Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x] - FresnelS[Sqrt[b]*Sqrt[2/Pi]*x]*Sin[a]))/Sqrt[b]
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3835, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(a + bx^2) dx$$

$$\downarrow \text{3835}$$

$$\cos(a) \int \cos(bx^2) dx - \sin(a) \int \sin(bx^2) dx$$

$$\downarrow \text{3832}$$

$$\cos(a) \int \cos(bx^2) dx - \frac{\sqrt{\frac{\pi}{2}} \sin(a) \text{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}}$$

$$\downarrow \text{3833}$$

$$\frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a) \text{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}}$$

input `Int[Cos[a + b*x^2], x]`

output `(Sqrt[Pi/2]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x])/Sqrt[b] - (Sqrt[Pi/2]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x]*Sin[a])/Sqrt[b]`

Defintions of rubi rules used

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3835

```
Int[Cos[(c_) + (d_.)*((e_.) + (f_.)*(x_))2], x_Symbol] := Simp[Cos[c] Int[Cos[d*(e + f*x)2], x], x] - Simp[Sin[c] Int[Sin[d*(e + f*x)2], x], x]
/; FreeQ[{c, d, e, f}, x]
```

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.63

method	result	size
default	$\frac{\sqrt{2}\sqrt{\pi}\left(\cos(a)\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)-\sin(a)\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)\right)}{2\sqrt{b}}$	44
meijerg	$\frac{\sqrt{2}\sqrt{\pi}\cos(a)\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\sqrt{2}\sqrt{\pi}\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)\sin(a)}{2\sqrt{b}}$	52
risch	$\frac{e^{-ia}\sqrt{\pi}\operatorname{erf}\left(\sqrt{ib}x\right)}{4\sqrt{ib}} + \frac{e^{ia}\sqrt{\pi}\operatorname{erf}\left(\sqrt{-ib}x\right)}{4\sqrt{-ib}}$	52

input

```
int(cos(b*x^2+a),x,method=_RETURNVERBOSE)
```

output

```
1/2*2^(1/2)*Pi^(1/2)/b^(1/2)*(cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)-sin(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.87

$$\int \cos(a + bx^2) dx = \frac{\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) - \sqrt{2}\pi\sqrt{\frac{b}{\pi}}S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a)}{2b}$$

input

```
integrate(cos(b*x^2+a),x, algorithm="fricas")
```

output

```
1/2*(sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_cos(sqrt(2)*x*sqrt(b/pi)) - sqrt(2)*pi*sqrt(b/pi)*fresnel_sin(sqrt(2)*x*sqrt(b/pi))*sin(a))/b
```

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.87

$$\int \cos(a + bx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(-\sin(a) S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(a) C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right) \right) \sqrt{\frac{1}{b}}}{2}$$

input `integrate(cos(b*x**2+a),x)`

output `sqrt(2)*sqrt(pi)*(-sin(a)*fresnels(sqrt(2)*sqrt(b)*x/sqrt(pi)) + cos(a)*fresnelc(sqrt(2)*sqrt(b)*x/sqrt(pi)))*sqrt(1/b)/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.69

$$\int \cos(a + bx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(((i-1)\cos(a) + (i+1)\sin(a)) \operatorname{erf}(\sqrt{i}bx) + (-(i+1)\cos(a) - (i-1)\sin(a)) \operatorname{erf}(\sqrt{-i}bx) \right)}{8\sqrt{b}}$$

input `integrate(cos(b*x^2+a),x, algorithm="maxima")`

output `-1/8*sqrt(2)*sqrt(pi)*(((I - 1)*cos(a) + (I + 1)*sin(a))*erf(sqrt(I*b)*x) + (-(I + 1)*cos(a) - (I - 1)*sin(a))*erf(sqrt(-I*b)*x))/sqrt(b)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.36

$$\int \cos(a + bx^2) dx = -\frac{\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{(ia)}}{4\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}} - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{(-ia)}}{4\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}}$$

input `integrate(cos(b*x^2+a),x, algorithm="giac")`

output `-1/4*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(-I*b/abs(b) + 1)*sqrt(abs(b)))*e^(I*a)/((-I*b/abs(b) + 1)*sqrt(abs(b))) - 1/4*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(I*b/abs(b) + 1)*sqrt(abs(b)))*e^(-I*a)/((I*b/abs(b) + 1)*sqrt(abs(b)))`

Mupad [B] (verification not implemented)

Time = 42.92 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.73

$$\int \cos(a + bx^2) dx = \frac{\sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right) \cos(a)}{2\sqrt{b}} - \frac{\sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right) \sin(a)}{2\sqrt{b}}$$

input `int(cos(a + b*x^2),x)`

output `(2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*b^(1/2)*x)/pi^(1/2))*cos(a)/(2*b^(1/2)) - (2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*b^(1/2)*x)/pi^(1/2))*sin(a)/(2*b^(1/2))`

Reduce [F]

$$\int \cos(a + bx^2) dx = \int \cos(bx^2 + a) dx$$

input `int(cos(b*x^2+a),x)`

output `int(cos(a + b*x**2),x)`

3.5 $\int \frac{\cos(a+bx^2)}{x} dx$

Optimal result	87
Mathematica [A] (verified)	87
Rubi [A] (verified)	88
Maple [A] (verified)	89
Fricas [A] (verification not implemented)	89
Sympy [F]	90
Maxima [C] (verification not implemented)	90
Giac [A] (verification not implemented)	90
Mupad [F(-1)]	91
Reduce [F]	91

Optimal result

Integrand size = 12, antiderivative size = 25

$$\int \frac{\cos(a+bx^2)}{x} dx = \frac{1}{2} \cos(a) \operatorname{CosIntegral}(bx^2) - \frac{1}{2} \sin(a) \operatorname{Si}(bx^2)$$

output `1/2*cos(a)*Ci(b*x^2)-1/2*sin(a)*Si(b*x^2)`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{\cos(a+bx^2)}{x} dx = \frac{1}{2} (\cos(a) \operatorname{CosIntegral}(bx^2) - \sin(a) \operatorname{Si}(bx^2))$$

input `Integrate[Cos[a + b*x^2]/x,x]`

output `(Cos[a]*CosIntegral[b*x^2] - Sin[a]*SinIntegral[b*x^2])/2`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3859, 3856, 3857}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(a + bx^2)}{x} dx \\ & \quad \downarrow \text{3859} \\ & \cos(a) \int \frac{\cos(bx^2)}{x} dx - \sin(a) \int \frac{\sin(bx^2)}{x} dx \\ & \quad \downarrow \text{3856} \\ & \cos(a) \int \frac{\cos(bx^2)}{x} dx - \frac{1}{2} \sin(a) \text{Si}(bx^2) \\ & \quad \downarrow \text{3857} \\ & \frac{1}{2} \cos(a) \text{CosIntegral}(bx^2) - \frac{1}{2} \sin(a) \text{Si}(bx^2) \end{aligned}$$

input `Int[Cos[a + b*x^2]/x,x]`

output `(Cos[a]*CosIntegral[b*x^2])/2 - (Sin[a]*SinIntegral[b*x^2])/2`

Defintions of rubi rules used

rule 3856 `Int[Sin[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[SinIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3857 `Int[Cos[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[CosIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3859

```
Int[Cos[(c_) + (d_.)*(x_)^(n_)]/(x_), x_Symbol] := Simp[Cos[c] Int[Cos[d*
x^n]/x, x], x] - Simp[Sin[c] Int[Sin[d*x^n]/x, x], x] /; FreeQ[{c, d, n},
x]
```

Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{\cos(a) \operatorname{Ci}(bx^2)}{2} - \frac{\sin(a) \operatorname{Si}(bx^2)}{2}$	22
risch	$\frac{i\pi \operatorname{csgn}(bx^2)e^{-ia}}{4} - \frac{i \operatorname{Si}(bx^2)e^{-ia}}{2} - \frac{e^{-ia} \operatorname{expIntegral}_1(-ibx^2)}{4} - \frac{e^{ia} \operatorname{expIntegral}_1(-ibx^2)}{4}$	63
meijerg	$\frac{\cos(a)\sqrt{\pi} \left(\frac{2\gamma+4\ln(x)+\ln(b^2)}{\sqrt{\pi}} - \frac{2\gamma}{\sqrt{\pi}} - \frac{2\ln(2)}{\sqrt{\pi}} - \frac{2\ln\left(\frac{bx^2}{2}\right)}{\sqrt{\pi}} + \frac{2 \operatorname{Ci}(bx^2)}{\sqrt{\pi}} \right)}{4} - \frac{\sin(a) \operatorname{Si}(bx^2)}{2}$	72

```
input int(cos(b*x^2+a)/x,x,method=_RETURNVERBOSE)
```

```
output 1/2*cos(a)*Ci(b*x^2)-1/2*sin(a)*Si(b*x^2)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{\cos(a + bx^2)}{x} dx = \frac{1}{2} \cos(a) \operatorname{Ci}(bx^2) - \frac{1}{2} \sin(a) \operatorname{Si}(bx^2)$$

```
input integrate(cos(b*x^2+a)/x,x, algorithm="fricas")
```

```
output 1/2*cos(a)*cos_integral(b*x^2) - 1/2*sin(a)*sin_integral(b*x^2)
```

Sympy [F]

$$\int \frac{\cos(a + bx^2)}{x} dx = \int \frac{\cos(a + bx^2)}{x} dx$$

input `integrate(cos(b*x**2+a)/x,x)`

output `Integral(cos(a + b*x**2)/x, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.72

$$\int \frac{\cos(a + bx^2)}{x} dx = \frac{1}{4} (\text{Ei}(i bx^2) + \text{Ei}(-i bx^2)) \cos(a) + \frac{1}{4} (i \text{Ei}(i bx^2) - i \text{Ei}(-i bx^2)) \sin(a)$$

input `integrate(cos(b*x^2+a)/x,x, algorithm="maxima")`

output `1/4*(Ei(I*b*x^2) + Ei(-I*b*x^2))*cos(a) + 1/4*(I*Ei(I*b*x^2) - I*Ei(-I*b*x^2))*sin(a)`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{\cos(a + bx^2)}{x} dx = \frac{1}{2} \cos(a) \text{Ci}(bx^2) - \frac{1}{2} \sin(a) \text{Si}(bx^2)$$

input `integrate(cos(b*x^2+a)/x,x, algorithm="giac")`

output `1/2*cos(a)*cos_integral(b*x^2) - 1/2*sin(a)*sin_integral(b*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^2)}{x} dx = \frac{\cos(a) \operatorname{cosint}(bx^2)}{2} - \frac{\sin(a) \operatorname{sinint}(bx^2)}{2}$$

input `int(cos(a + b*x^2)/x,x)`output `(cos(a)*cosint(b*x^2))/2 - (sin(a)*sinint(b*x^2))/2`**Reduce [F]**

$$\int \frac{\cos(a + bx^2)}{x} dx = \int \frac{\cos(bx^2 + a)}{x} dx$$

input `int(cos(b*x^2+a)/x,x)`output `int(cos(a + b*x**2)/x,x)`

3.6 $\int \frac{\cos(a+bx^2)}{x^2} dx$

Optimal result	92
Mathematica [A] (verified)	92
Rubi [A] (verified)	93
Maple [A] (verified)	94
Fricas [A] (verification not implemented)	95
Sympy [F]	95
Maxima [C] (verification not implemented)	96
Giac [F]	96
Mupad [F(-1)]	96
Reduce [F]	97

Optimal result

Integrand size = 12, antiderivative size = 80

$$\int \frac{\cos(a+bx^2)}{x^2} dx = -\frac{\cos(a+bx^2)}{x} - \sqrt{b}\sqrt{2\pi} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \sqrt{b}\sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)$$

```
output -cos(b*x^2+a)/x-b^(1/2)*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)-b^(1/2)*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)*sin(a)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.01

$$\int \frac{\cos(a+bx^2)}{x^2} dx = -\frac{\cos(a)\cos(bx^2)}{x} - \sqrt{b}\sqrt{2\pi} \left(\cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) + \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a) \right) + \frac{\sin(a)\sin(bx^2)}{x}$$

input `Integrate[Cos[a + b*x^2]/x^2,x]`

output
$$-\left(\frac{\cos(a)\cos(bx^2)}{x} - \sqrt{b}\sqrt{2\pi}\left(\cos(a)\operatorname{FresnelS}\left[\sqrt{b}\sqrt{\frac{2}{\pi}}x\right] + \operatorname{FresnelC}\left[\sqrt{b}\sqrt{\frac{2}{\pi}}x\right]\sin(a)\right) + \frac{\sin(a)\sin(bx^2)}{x}\right)$$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.08, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3869, 3834, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(a + bx^2)}{x^2} dx \\ & \quad \downarrow \text{3869} \\ & -2b \int \sin(bx^2 + a) dx - \frac{\cos(a + bx^2)}{x} \\ & \quad \downarrow \text{3834} \\ & -2b \left(\sin(a) \int \cos(bx^2) dx + \cos(a) \int \sin(bx^2) dx \right) - \frac{\cos(a + bx^2)}{x} \\ & \quad \downarrow \text{3832} \\ & -2b \left(\sin(a) \int \cos(bx^2) dx + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}} \right) - \frac{\cos(a + bx^2)}{x} \\ & \quad \downarrow \text{3833} \\ & -2b \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{\sqrt{b}} \right) - \frac{\cos(a + bx^2)}{x} \end{aligned}$$

input `Int[Cos[a + b*x^2]/x^2,x]`

output $-\frac{\cos(a + bx^2)}{x} - 2b \left(\frac{\sqrt{\pi/2} \cos(a) \operatorname{FresnelS}(\sqrt{b} \sqrt{2/\pi} x)}{\sqrt{b}} + \frac{\sqrt{\pi/2} \operatorname{FresnelC}(\sqrt{b} \sqrt{2/\pi} x) \sin(a)}{\sqrt{b}} \right)$

Defintions of rubi rules used

rule 3832 $\operatorname{Int}[\sin((d_.) * ((e_.) + (f_.) * (x_.)^2)), x_Symbol] \rightarrow \operatorname{Simp}[(\sqrt{\pi/2} / (f * \operatorname{Rt}[d, 2])) * \operatorname{FresnelS}[\sqrt{2/\pi} * \operatorname{Rt}[d, 2] * (e + f * x)], x] /;$ $\operatorname{FreeQ}\{d, e, f, x\}$

rule 3833 $\operatorname{Int}[\cos((d_.) * ((e_.) + (f_.) * (x_.)^2)), x_Symbol] \rightarrow \operatorname{Simp}[(\sqrt{\pi/2} / (f * \operatorname{Rt}[d, 2])) * \operatorname{FresnelC}[\sqrt{2/\pi} * \operatorname{Rt}[d, 2] * (e + f * x)], x] /;$ $\operatorname{FreeQ}\{d, e, f, x\}$

rule 3834 $\operatorname{Int}[\sin((c_.) + (d_.) * ((e_.) + (f_.) * (x_.)^2)), x_Symbol] \rightarrow \operatorname{Simp}[\sin[c] \operatorname{Int}[\cos[d * (e + f * x)^2], x], x] + \operatorname{Simp}[\cos[c] \operatorname{Int}[\sin[d * (e + f * x)^2], x], x] /;$ $\operatorname{FreeQ}\{c, d, e, f, x\}$

rule 3869 $\operatorname{Int}[\cos((c_.) + (d_.) * (x_.)^n) * ((e_.) * (x_.)^m), x_Symbol] \rightarrow \operatorname{Simp}[(e * x)^{m+1} * (\cos[c + d * x^n] / (e * (m + 1))), x] + \operatorname{Simp}[d * (n / (e^n * (m + 1))) \operatorname{Int}[(e * x)^{m+n} * \sin[c + d * x^n], x], x] /;$ $\operatorname{FreeQ}\{c, d, e, x\} \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{LtQ}[m, -1]$

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.71

method	result
default	$-\frac{\cos(bx^2+a)}{x} - \sqrt{b} \sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) \right)$
risch	$-\frac{ie^{-ia}b\sqrt{\pi} \operatorname{erf}(\sqrt{ib}x)}{2\sqrt{ib}} + \frac{ie^{ia}b\sqrt{\pi} \operatorname{erf}(\sqrt{-ib}x)}{2\sqrt{-ib}} - \frac{\cos(bx^2+a)}{x}$
meijerg	$\frac{\cos(a)\sqrt{\pi}(b^2)^{\frac{1}{4}}\sqrt{2} \left(-\frac{4\sqrt{2}\cos(bx^2)}{\sqrt{\pi}x(b^2)^{\frac{1}{4}}} - \frac{8\sqrt{b}\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)}{(b^2)^{\frac{1}{4}}} \right)}{8} - \frac{\sin(a)\sqrt{\pi}\sqrt{b}\sqrt{2} \left(-\frac{4\sqrt{2}\sin(bx^2)}{\sqrt{b}\sqrt{\pi}x} + 8 \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) \right)}{8}$

input `int(cos(b*x^2+a)/x^2,x,method=_RETURNVERBOSE)`

output

```
-cos(b*x^2+a)/x-b^(1/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/
Pi^(1/2)*x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{\cos(a + bx^2)}{x^2} dx$$

$$= -\frac{\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} \cos(a) S\left(\sqrt{2}x \sqrt{\frac{b}{\pi}}\right) + \sqrt{2}\pi x \sqrt{\frac{b}{\pi}} C\left(\sqrt{2}x \sqrt{\frac{b}{\pi}}\right) \sin(a) + \cos(bx^2 + a)}{x}$$

input

```
integrate(cos(b*x^2+a)/x^2,x, algorithm="fricas")
```

output

```
-(sqrt(2)*pi*x*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x*sqrt(b/pi)) + sqrt(
2)*pi*x*sqrt(b/pi)*fresnel_cos(sqrt(2)*x*sqrt(b/pi))*sin(a) + cos(b*x^2 +
a))/x
```

Sympy [F]

$$\int \frac{\cos(a + bx^2)}{x^2} dx = \int \frac{\cos(a + bx^2)}{x^2} dx$$

input

```
integrate(cos(b*x**2+a)/x**2,x)
```

output

```
Integral(cos(a + b*x**2)/x**2, x)
```


Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.91

$$\int \frac{\cos(a + bx^2)}{x^2} dx$$

$$= \frac{\sqrt{bx^2} \left((-i + 1) \sqrt{2} \Gamma\left(-\frac{1}{2}, i bx^2\right) + (i - 1) \sqrt{2} \Gamma\left(-\frac{1}{2}, -i bx^2\right) \right) \cos(a) + \left((i - 1) \sqrt{2} \Gamma\left(-\frac{1}{2}, i bx^2\right) - (i + 1) \sqrt{2} \Gamma\left(-\frac{1}{2}, -i bx^2\right) \right) \sin(a)}{8x}$$

input `integrate(cos(b*x^2+a)/x^2,x, algorithm="maxima")`

output `1/8*sqrt(b*x^2)*((-I + 1)*sqrt(2)*gamma(-1/2, I*b*x^2) + (I - 1)*sqrt(2)*gamma(-1/2, -I*b*x^2))*cos(a) + ((I - 1)*sqrt(2)*gamma(-1/2, I*b*x^2) - (I + 1)*sqrt(2)*gamma(-1/2, -I*b*x^2))*sin(a)/x`

Giac [F]

$$\int \frac{\cos(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)}{x^2} dx$$

input `integrate(cos(b*x^2+a)/x^2,x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)}{x^2} dx$$

input `int(cos(a + b*x^2)/x^2,x)`

output `int(cos(a + b*x^2)/x^2, x)`

Reduce [F]

$$\int \frac{\cos(a + bx^2)}{x^2} dx = \frac{\left(\int \frac{\cos(bx^2+a)}{x^2} dx\right) x + \left(\int \frac{1}{x^2} dx\right) x + 1}{x}$$

input `int(cos(b*x^2+a)/x^2, x)`

output `(int(cos(a + b*x**2)/x**2, x)*x + int(1/x**2, x)*x + 1)/x`

3.7 $\int \frac{\cos(a+bx^2)}{x^3} dx$

Optimal result	98
Mathematica [A] (verified)	98
Rubi [A] (verified)	99
Maple [A] (verified)	101
Fricas [A] (verification not implemented)	101
Sympy [F]	102
Maxima [C] (verification not implemented)	102
Giac [B] (verification not implemented)	103
Mupad [F(-1)]	103
Reduce [F]	103

Optimal result

Integrand size = 12, antiderivative size = 42

$$\int \frac{\cos(a+bx^2)}{x^3} dx = -\frac{\cos(a+bx^2)}{2x^2} - \frac{1}{2}b \operatorname{CosIntegral}(bx^2) \sin(a) - \frac{1}{2}b \cos(a) \operatorname{Si}(bx^2)$$

output

```
-1/2*cos(b*x^2+a)/x^2-1/2*b*Ci(b*x^2)*sin(a)-1/2*b*cos(a)*Si(b*x^2)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{\cos(a+bx^2)}{x^3} dx = -\frac{\cos(a+bx^2) + bx^2 \operatorname{CosIntegral}(bx^2) \sin(a) + bx^2 \cos(a) \operatorname{Si}(bx^2)}{2x^2}$$

input

```
Integrate[Cos[a + b*x^2]/x^3,x]
```

output

```
-1/2*(Cos[a + b*x^2] + b*x^2*CosIntegral[b*x^2]*Sin[a] + b*x^2*Cos[a]*SinIntegral[b*x^2])/x^2
```

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3861, 3042, 3778, 25, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(a + bx^2)}{x^3} dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{2} \int \frac{\cos(bx^2 + a)}{x^4} dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{\sin(bx^2 + a + \frac{\pi}{2})}{x^4} dx^2 \\
 & \quad \downarrow \text{3778} \\
 & \frac{1}{2} \left(b \int -\frac{\sin(bx^2 + a)}{x^2} dx^2 - \frac{\cos(a + bx^2)}{x^2} \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \left(-b \int \frac{\sin(bx^2 + a)}{x^2} dx^2 - \frac{\cos(a + bx^2)}{x^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(-b \int \frac{\sin(bx^2 + a)}{x^2} dx^2 - \frac{\cos(a + bx^2)}{x^2} \right) \\
 & \quad \downarrow \text{3784} \\
 & \frac{1}{2} \left(-b \left(\sin(a) \int \frac{\cos(bx^2)}{x^2} dx^2 + \cos(a) \int \frac{\sin(bx^2)}{x^2} dx^2 \right) - \frac{\cos(a + bx^2)}{x^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(-b \left(\sin(a) \int \frac{\sin(bx^2 + \frac{\pi}{2})}{x^2} dx^2 + \cos(a) \int \frac{\sin(bx^2)}{x^2} dx^2 \right) - \frac{\cos(a + bx^2)}{x^2} \right)
 \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3780} \\ & \frac{1}{2} \left(-b \left(\sin(a) \int \frac{\sin(bx^2 + \frac{\pi}{2})}{x^2} dx^2 + \cos(a) \text{Si}(bx^2) \right) - \frac{\cos(a + bx^2)}{x^2} \right) \\ & \downarrow \text{3783} \\ & \frac{1}{2} \left(-b(\sin(a) \text{CosIntegral}(bx^2) + \cos(a) \text{Si}(bx^2)) - \frac{\cos(a + bx^2)}{x^2} \right) \end{aligned}$$

input `Int[Cos[a + b*x^2]/x^3,x]`

output `((-Cos[a + b*x^2]/x^2) - b*(CosIntegral[b*x^2]*Sin[a] + Cos[a]*SinIntegral[b*x^2]))/2`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3780 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

rule 3784

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*
e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*
f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x]
&& NeQ[d*e - c*f, 0]
```

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
:= Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.93

method	result
default	$-\frac{\cos(bx^2+a)}{2x^2} - b \left(\frac{\cos(a) \operatorname{Si}(bx^2)}{2} + \frac{\operatorname{Ci}(bx^2) \sin(a)}{2} \right)$
risch	$\frac{e^{-ia} \pi \operatorname{csgn}(bx^2) b}{4} - \frac{e^{-ia} \operatorname{Si}(bx^2) b}{2} + \frac{i \exp \operatorname{Integral}_1(-ibx^2) e^{-ia} b}{4} - \frac{ie^{ia} b \exp \operatorname{Integral}_1(-ibx^2)}{4} - \frac{\cos(bx^2+a)}{2x^2}$
meijerg	$\frac{\cos(a) \sqrt{\pi} \sqrt{b^2} \left(-\frac{4b^2 \cos(x^2 \sqrt{b^2})}{x^2 (b^2)^{\frac{3}{2}} \sqrt{\pi}} - \frac{4 \operatorname{Si}(x^2 \sqrt{b^2})}{\sqrt{\pi}} \right)}{8} - \frac{\sin(a) \sqrt{\pi} b \left(\frac{4\gamma - 4 + 8 \ln(x) + 4 \ln(b)}{\sqrt{\pi}} + \frac{4}{\sqrt{\pi}} - \frac{4\gamma}{\sqrt{\pi}} - \frac{4 \ln(2)}{\sqrt{\pi}} - \frac{4 \ln\left(\frac{bx^2}{2}\right)}{\sqrt{\pi}} - \frac{4 \sin(bx^2)}{\sqrt{\pi} b x^2} \right)}{8}$

input

```
int(cos(b*x^2+a)/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*cos(b*x^2+a)/x^2-b*(1/2*cos(a)*Si(b*x^2)+1/2*Ci(b*x^2)*sin(a))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.95

$$\int \frac{\cos(a + bx^2)}{x^3} dx = -\frac{bx^2 \operatorname{Ci}(bx^2) \sin(a) + bx^2 \cos(a) \operatorname{Si}(bx^2) + \cos(bx^2 + a)}{2x^2}$$

input

```
integrate(cos(b*x^2+a)/x^3,x, algorithm="fricas")
```

output `-1/2*(b*x^2*cos_integral(b*x^2)*sin(a) + b*x^2*cos(a)*sin_integral(b*x^2) + cos(b*x^2 + a))/x^2`

Sympy [F]

$$\int \frac{\cos(a + bx^2)}{x^3} dx = \int \frac{\cos(a + bx^2)}{x^3} dx$$

input `integrate(cos(b*x**2+a)/x**3,x)`

output `Integral(cos(a + b*x**2)/x**3, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.14

$$\int \frac{\cos(a + bx^2)}{x^3} dx = -\frac{1}{4} \left((i\Gamma(-1, i bx^2) - i\Gamma(-1, -i bx^2)) \cos(a) + (\Gamma(-1, i bx^2) + \Gamma(-1, -i bx^2)) \sin(a) \right) b$$

input `integrate(cos(b*x^2+a)/x^3,x, algorithm="maxima")`

output `-1/4*((I*gamma(-1, I*b*x^2) - I*gamma(-1, -I*b*x^2))*cos(a) + (gamma(-1, I*b*x^2) + gamma(-1, -I*b*x^2))*sin(a))*b`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(36) = 72$.

Time = 0.39 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.07

$$\int \frac{\cos(a + bx^2)}{x^3} dx = \frac{(bx^2 + a)b^2 \operatorname{Ci}(bx^2) \sin(a) - ab^2 \operatorname{Ci}(bx^2) \sin(a) + (bx^2 + a)b^2 \cos(a) \operatorname{Si}(bx^2) - ab^2 \cos(a) \operatorname{Si}(bx^2) + b^2 \cos(bx^2 + a)}{2b^2x^2}$$

input `integrate(cos(b*x^2+a)/x^3,x, algorithm="giac")`

output `-1/2*((b*x^2 + a)*b^2*cos_integral(b*x^2)*sin(a) - a*b^2*cos_integral(b*x^2)*sin(a) + (b*x^2 + a)*b^2*cos(a)*sin_integral(b*x^2) - a*b^2*cos(a)*sin_integral(b*x^2) + b^2*cos(b*x^2 + a))/(b^2*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^2)}{x^3} dx = \int \frac{\cos(bx^2 + a)}{x^3} dx$$

input `int(cos(a + b*x^2)/x^3,x)`

output `int(cos(a + b*x^2)/x^3, x)`

Reduce [F]

$$\int \frac{\cos(a + bx^2)}{x^3} dx = \frac{2\left(\int \frac{\cos(bx^2+a)}{x^3} dx\right) x^2 + 2\left(\int \frac{1}{x^3} dx\right) x^2 + 1}{2x^2}$$

input `int(cos(b*x^2+a)/x^3,x)`

output `(2*int(cos(a + b*x**2)/x**3,x)*x**2 + 2*int(1/x**3,x)*x**2 + 1)/(2*x**2)`

3.8 $\int x^3 \cos^2(a + bx^2) dx$

Optimal result	105
Mathematica [A] (verified)	105
Rubi [A] (verified)	106
Maple [A] (verified)	107
Fricas [A] (verification not implemented)	108
Sympy [A] (verification not implemented)	108
Maxima [A] (verification not implemented)	109
Giac [A] (verification not implemented)	109
Mupad [B] (verification not implemented)	109
Reduce [B] (verification not implemented)	110

Optimal result

Integrand size = 14, antiderivative size = 51

$$\int x^3 \cos^2(a + bx^2) dx = \frac{x^4}{8} + \frac{\cos^2(a + bx^2)}{8b^2} + \frac{x^2 \cos(a + bx^2) \sin(a + bx^2)}{4b}$$

output

```
1/8*x^4+1/8*cos(b*x^2+a)^2/b^2+1/4*x^2*cos(b*x^2+a)*sin(b*x^2+a)/b
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.78

$$\int x^3 \cos^2(a + bx^2) dx = \frac{\cos(2(a + bx^2)) + 2bx^2(bx^2 + \sin(2(a + bx^2)))}{16b^2}$$

input

```
Integrate[x^3*Cos[a + b*x^2]^2,x]
```

output

```
(Cos[2*(a + b*x^2)] + 2*b*x^2*(b*x^2 + Sin[2*(a + b*x^2)]))/(16*b^2)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3861, 3042, 3791, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \cos^2(a + bx^2) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{2} \int x^2 \cos^2(bx^2 + a) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int x^2 \sin\left(bx^2 + a + \frac{\pi}{2}\right)^2 dx^2 \\
 & \quad \downarrow \text{3791} \\
 & \frac{1}{2} \left(\frac{\int x^2 dx^2}{2} + \frac{\cos^2(a + bx^2)}{4b^2} + \frac{x^2 \sin(a + bx^2) \cos(a + bx^2)}{2b} \right) \\
 & \quad \downarrow \text{15} \\
 & \frac{1}{2} \left(\frac{\cos^2(a + bx^2)}{4b^2} + \frac{x^2 \sin(a + bx^2) \cos(a + bx^2)}{2b} + \frac{x^4}{4} \right)
 \end{aligned}$$

input `Int[x^3*Cos[a + b*x^2]^2,x]`

output `(x^4/4 + Cos[a + b*x^2]^2/(4*b^2) + (x^2*Cos[a + b*x^2]*Sin[a + b*x^2]))/(2*b)`

Definitions of rubi rules used

rule 15 $\text{Int}[(a_.)(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[a*(x^{(m+1)})/(m+1), x] /; \text{FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3791 $\text{Int}[((c_.) + (d_.)(x_))*((b_.)\sin[(e_.) + (f_.)(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[d*((b*\sin[e + f*x])^n/(f^n)), x] + (-\text{Simp}[b*(c + d*x)*\cos[e + f*x]*((b*\sin[e + f*x])^{(n-1)})/(f^n), x] + \text{Simp}[b^2*((n-1)/n) \text{Int}[(c + d*x)*(b*\sin[e + f*x])^{(n-2)}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1]$

rule 3861 $\text{Int}[(a_.) + \cos[(c_.) + (d_.)(x_)^{(n_.)}]*(b_.)]^{(p_.)}(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n) - 1)}*(a + b*\cos[c + d*x])^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, c, d, m, n, p\}, x] \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m+1)/n]] \ \&\& \ (\text{EqQ}[p, 1] \ || \ \text{EqQ}[m, n-1] \ || \ (\text{IntegerQ}[p] \ \&\& \ \text{GtQ}[\text{Simplify}[(m+1)/n], 0]))$

Maple [A] (verified)

Time = 1.20 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.82

method	result
default	$\frac{x^4}{8} + \frac{x^2 \sin(2bx^2+2a)}{8b} + \frac{\cos(2bx^2+2a)}{16b^2}$
risch	$\frac{x^4}{8} + \frac{x^2 \sin(2bx^2+2a)}{8b} + \frac{\cos(2bx^2+2a)}{16b^2}$
parallelrisch	$\frac{2b^2x^4+2x^2 \sin(2bx^2+2a)b+\cos(2bx^2+2a)-1}{16b^2}$
norman	$\frac{\frac{x^4}{8} + \frac{x^4 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^2}{4} + \frac{x^4 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^4}{8} + \frac{x^2 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)}{2b} - \frac{x^2 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^3}{2b} - \frac{\tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^2}{2b^2}}{\left(1 + \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)\right)^2}$
orering	$\frac{(16b^2x^4+35)\cos(bx^2+a)^2}{64b^2} - \frac{11(3x^2\cos(bx^2+a)^2-4x^4\cos(bx^2+a)b\sin(bx^2+a))}{64x^2b^2} + \frac{6x\cos(bx^2+a)^2-28x^3\cos(bx^2+a)}{64x^2b^2}$

input `int(x^3*cos(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output `1/8*x^4+1/8/b*x^2*sin(2*b*x^2+2*a)+1/16/b^2*cos(2*b*x^2+2*a)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.88

$$\int x^3 \cos^2(a + bx^2) dx = \frac{b^2 x^4 + 2bx^2 \cos(bx^2 + a) \sin(bx^2 + a) + \cos(bx^2 + a)^2}{8b^2}$$

input `integrate(x^3*cos(b*x^2+a)^2,x, algorithm="fricas")`

output `1/8*(b^2*x^4 + 2*b*x^2*cos(b*x^2 + a)*sin(b*x^2 + a) + cos(b*x^2 + a)^2)/b^2`

Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.53

$$\int x^3 \cos^2(a + bx^2) dx = \begin{cases} \frac{x^4 \sin^2(a+bx^2)}{8} + \frac{x^4 \cos^2(a+bx^2)}{8} + \frac{x^2 \sin(a+bx^2) \cos(a+bx^2)}{4b} + \frac{\cos^2(a+bx^2)}{8b^2} & \text{for } b \neq 0 \\ \frac{x^4 \cos^2(a)}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*cos(b*x**2+a)**2,x)`

output `Piecewise((x**4*sin(a + b*x**2)**2/8 + x**4*cos(a + b*x**2)**2/8 + x**2*sin(a + b*x**2)*cos(a + b*x**2)/(4*b) + cos(a + b*x**2)**2/(8*b**2), Ne(b, 0)), (x**4*cos(a)**2/4, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.82

$$\int x^3 \cos^2(a + bx^2) dx = \frac{2b^2x^4 + 2bx^2 \sin(2bx^2 + 2a) + \cos(2bx^2 + 2a)}{16b^2}$$

input `integrate(x^3*cos(b*x^2+a)^2,x, algorithm="maxima")`

output `1/16*(2*b^2*x^4 + 2*b*x^2*sin(2*b*x^2 + 2*a) + cos(2*b*x^2 + 2*a))/b^2`

Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.49

$$\int x^3 \cos^2(a + bx^2) dx = -\frac{(2bx^2 + 2a + \sin(2bx^2 + 2a))a}{8b^2} + \frac{2(bx^2 + a)^2 + 2(bx^2 + a)\sin(2bx^2 + 2a) + \cos(2bx^2 + 2a)}{16b^2}$$

input `integrate(x^3*cos(b*x^2+a)^2,x, algorithm="giac")`

output `-1/8*(2*b*x^2 + 2*a + sin(2*b*x^2 + 2*a))*a/b^2 + 1/16*(2*(b*x^2 + a)^2 + 2*(b*x^2 + a)*sin(2*b*x^2 + 2*a) + cos(2*b*x^2 + 2*a))/b^2`

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.80

$$\int x^3 \cos^2(a + bx^2) dx = \frac{\cos(2bx^2 + 2a)}{16b^2} + \frac{x^4}{8} + \frac{x^2 \sin(2bx^2 + 2a)}{8b}$$

input `int(x^3*cos(a + b*x^2)^2,x)`

output `cos(2*a + 2*b*x^2)/(16*b^2) + x^4/8 + (x^2*sin(2*a + 2*b*x^2))/(8*b)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.94

$$\int x^3 \cos^2(a + bx^2) dx = \frac{2 \cos(bx^2 + a) \sin(bx^2 + a) bx^2 - \sin(bx^2 + a)^2 + b^2 x^4 + 2}{8b^2}$$

input `int(x^3*cos(b*x^2+a)^2,x)`

output `(2*cos(a + b*x**2)*sin(a + b*x**2)*b*x**2 - sin(a + b*x**2)**2 + b**2*x**4 + 2)/(8*b**2)`

3.9 $\int x^2 \cos^2(a + bx^2) dx$

Optimal result	111
Mathematica [A] (verified)	111
Rubi [A] (verified)	112
Maple [A] (verified)	113
Fricas [A] (verification not implemented)	113
Sympy [B] (verification not implemented)	114
Maxima [C] (verification not implemented)	115
Giac [C] (verification not implemented)	115
Mupad [F(-1)]	116
Reduce [F]	116

Optimal result

Integrand size = 14, antiderivative size = 91

$$\int x^2 \cos^2(a + bx^2) dx = \frac{x^3}{6} - \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{16b^{3/2}} - \frac{\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a)}{16b^{3/2}} + \frac{x \sin(2a + 2bx^2)}{8b}$$

output

```
1/6*x^3-1/16*Pi^(1/2)*cos(2*a)*FresnelS(2*b^(1/2)*x/Pi^(1/2))/b^(3/2)-1/16
*Pi^(1/2)*FresnelC(2*b^(1/2)*x/Pi^(1/2))*sin(2*a)/b^(3/2)+1/8*x*sin(2*b*x^
2+2*a)/b
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.96

$$\int x^2 \cos^2(a + bx^2) dx = \frac{-3\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a) + 2\sqrt{bx}(4bx^2 + 3 \sin(2(a + bx^2)))}{48b^{3/2}}$$

input

```
Integrate[x^2*Cos[a + b*x^2]^2,x]
```


output

$$\frac{(-3\sqrt{\pi}\cos[2a]\operatorname{FresnelS}[(2\sqrt{b}x)/\sqrt{\pi}] - 3\sqrt{\pi}\operatorname{FresnelC}[(2\sqrt{b}x)/\sqrt{\pi}]\sin[2a] + 2\sqrt{b}x(4bx^2 + 3\sin[2(a + bx^2)]))}{(48b^{3/2})}$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos^2(a + bx^2) dx$$

$$\downarrow \text{3885}$$

$$\int \left(\frac{1}{2}x^2 \cos(2a + 2bx^2) + \frac{x^2}{2} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right)}{16b^{3/2}} - \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right)}{16b^{3/2}} + \frac{x \sin(2a + 2bx^2)}{8b} + \frac{x^3}{6}$$

input

$$\text{Int}[x^2 \cos[a + b x^2]^2, x]$$

output

$$\frac{x^3}{6} - \frac{(\sqrt{\pi}\cos[2a]\operatorname{FresnelS}[(2\sqrt{b}x)/\sqrt{\pi}])}{(16b^{3/2})} - \frac{(\sqrt{\pi}\operatorname{FresnelC}[(2\sqrt{b}x)/\sqrt{\pi}]\sin[2a])}{(16b^{3/2})} + \frac{(x\sin[2a + 2bx^2])}{(8b)}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.69

method	result	size
default	$\frac{x^3}{6} + \frac{x \sin(2bx^2+2a)}{8b} - \frac{\sqrt{\pi} \left(\cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right) + \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right) \right)}{16b^{\frac{3}{2}}}$	63
risch	$\frac{x^3}{6} - \frac{ie^{-2ia}\sqrt{\pi}\sqrt{2}\operatorname{erf}\left(\sqrt{2}\sqrt{ib}x\right)}{64b\sqrt{ib}} + \frac{ie^{2ia}\sqrt{\pi}\operatorname{erf}\left(\sqrt{-2ib}x\right)}{32b\sqrt{-2ib}} + \frac{x \sin(2bx^2+2a)}{8b}$	88

input `int(x^2*cos(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output `1/6*x^3+1/8*x*sin(2*b*x^2+2*a)/b-1/16/b^(3/2)*Pi^(1/2)*(cos(2*a)*FresnelS(2*b^(1/2)*x/Pi^(1/2))+sin(2*a)*FresnelC(2*b^(1/2)*x/Pi^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.92

$$\int x^2 \cos^2(a + bx^2) dx$$

$$= \frac{8b^2x^3 + 12bx \cos(bx^2 + a) \sin(bx^2 + a) - 3\pi \sqrt{\frac{b}{\pi}} \cos(2a) S\left(2x\sqrt{\frac{b}{\pi}}\right) - 3\pi \sqrt{\frac{b}{\pi}} C\left(2x\sqrt{\frac{b}{\pi}}\right) \sin(2a)}{48b^2}$$

input `integrate(x^2*cos(b*x^2+a)^2,x, algorithm="fricas")`

output

```
1/48*(8*b^2*x^3 + 12*b*x*cos(b*x^2 + a)*sin(b*x^2 + a) - 3*pi*sqrt(b/pi)*cos(2*a)*fresnel_sin(2*x*sqrt(b/pi)) - 3*pi*sqrt(b/pi)*fresnel_cos(2*x*sqrt(b/pi))*sin(2*a))/b^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(85) = 170.

Time = 1.15 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.21

$$\int x^2 \cos^2(a + bx^2) dx = \frac{b^{\frac{3}{2}} x^5 \sqrt{\frac{1}{b}} \sin(2a) \Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{5}{4}\right) {}_2F_3\left(\frac{3}{4}, \frac{5}{4} \middle| \frac{3}{2}, \frac{7}{4}, \frac{9}{4} \right) -b^2 x^4}{8 \Gamma\left(\frac{7}{4}\right) \Gamma\left(\frac{9}{4}\right)} - \frac{\sqrt{b} x^3 \sqrt{\frac{1}{b}} \cos(2a) \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right) {}_2F_3\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{1}{2}, \frac{5}{4}, \frac{7}{4} \right) -b^2 x^4}{16 \Gamma\left(\frac{5}{4}\right) \Gamma\left(\frac{7}{4}\right)} + \frac{x^3}{6} - \frac{\sqrt{\pi} x^2 \sqrt{\frac{1}{b}} \sin(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4} + \frac{\sqrt{\pi} x^2 \sqrt{\frac{1}{b}} \cos(2a) C\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4}$$

input

```
integrate(x**2*cos(b*x**2+a)**2,x)
```

output

```
b**(3/2)*x**5*sqrt(1/b)*sin(2*a)*gamma(3/4)*gamma(5/4)*hyper((3/4, 5/4), (3/2, 7/4, 9/4), -b**2*x**4)/(8*gamma(7/4)*gamma(9/4)) - sqrt(b)*x**3*sqrt(1/b)*cos(2*a)*gamma(1/4)*gamma(3/4)*hyper((1/4, 3/4), (1/2, 5/4, 7/4), -b**2*x**4)/(16*gamma(5/4)*gamma(7/4)) + x**3/6 - sqrt(pi)*x**2*sqrt(1/b)*sin(2*a)*fresnels(2*sqrt(b)*x/sqrt(pi))/4 + sqrt(pi)*x**2*sqrt(1/b)*cos(2*a)*fresnelc(2*sqrt(b)*x/sqrt(pi))/4
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.99

$$\int x^2 \cos^2(a + bx^2) dx$$

$$= \frac{64 b^3 x^3 + 48 b^2 x \sin(2bx^2 + 2a) - 3 \cdot 4^{\frac{1}{4}} \sqrt{2} \sqrt{\pi} \left((i+1) \cos(2a) - (i-1) \sin(2a) \right) \operatorname{erf}(\sqrt{2i} bx) + (-\cos(2a) + (i+1) \sin(2a)) \operatorname{erf}(\sqrt{-2i} bx)}{384 b^3}$$

input `integrate(x^2*cos(b*x^2+a)^2,x, algorithm="maxima")`

output `1/384*(64*b^3*x^3 + 48*b^2*x*sin(2*b*x^2 + 2*a) - 3*4^(1/4)*sqrt(2)*sqrt(pi)*(((I + 1)*cos(2*a) - (I - 1)*sin(2*a))*erf(sqrt(2*I*b)*x) + (- (I - 1)*cos(2*a) + (I + 1)*sin(2*a))*erf(sqrt(-2*I*b)*x))*b^(3/2))/b^3`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.30

$$\int x^2 \cos^2(a + bx^2) dx = \frac{1}{6} x^3 - \frac{i x e^{(2i b x^2 + 2i a)}}{16 b} + \frac{i x e^{(-2i b x^2 - 2i a)}}{16 b} - \frac{i \sqrt{\pi} \operatorname{erf}\left(-\sqrt{b} x \left(-\frac{i b}{|b|} + 1\right)\right) e^{(2i a)}}{32 b^{\frac{3}{2}} \left(-\frac{i b}{|b|} + 1\right)} + \frac{i \sqrt{\pi} \operatorname{erf}\left(-\sqrt{b} x \left(\frac{i b}{|b|} + 1\right)\right) e^{(-2i a)}}{32 b^{\frac{3}{2}} \left(\frac{i b}{|b|} + 1\right)}$$

input `integrate(x^2*cos(b*x^2+a)^2,x, algorithm="giac")`

output

```
1/6*x^3 - 1/16*I*x*e^(2*I*b*x^2 + 2*I*a)/b + 1/16*I*x*e^(-2*I*b*x^2 - 2*I*
a)/b - 1/32*I*sqrt(pi)*erf(-sqrt(b)*x*(-I*b/abs(b) + 1))*e^(2*I*a)/(b^(3/2
))*(-I*b/abs(b) + 1) + 1/32*I*sqrt(pi)*erf(-sqrt(b)*x*(I*b/abs(b) + 1))*e^
(-2*I*a)/(b^(3/2)*(I*b/abs(b) + 1))
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2(a + bx^2) dx = \int x^2 \cos(bx^2 + a)^2 dx$$

input

```
int(x^2*cos(a + b*x^2)^2,x)
```

output

```
int(x^2*cos(a + b*x^2)^2, x)
```

Reduce [F]

$$\int x^2 \cos^2(a + bx^2) dx$$

$$= \frac{\cos(bx^2 + a) \sin(bx^2 + a) x - 4 \left(\int \frac{x^2}{\tan\left(\frac{bx^2 + a}{2}\right)^4 + 2 \tan\left(\frac{bx^2 + a}{2}\right)^2 + 1} dx \right) b - 4 \left(\int \frac{\tan\left(\frac{bx^2 + a}{2}\right)}{\tan\left(\frac{bx^2 + a}{2}\right)^4 + 2 \tan\left(\frac{bx^2 + a}{2}\right)^2 + 1} dx \right) b}{3b}$$

input

```
int(x^2*cos(b*x^2+a)^2,x)
```

output

```
(cos(a + b*x**2)*sin(a + b*x**2)*x - 4*int(x**2/(tan((a + b*x**2)/2)**4 +
2*tan((a + b*x**2)/2)**2 + 1),x)*b - 4*int(tan((a + b*x**2)/2)/(tan((a + b
*x**2)/2)**4 + 2*tan((a + b*x**2)/2)**2 + 1),x) + sin(a + b*x**2)*x + b*x*
*3)/(3*b)
```

3.10 $\int x \cos^2(a + bx^2) dx$

Optimal result	117
Mathematica [A] (verified)	117
Rubi [A] (verified)	118
Maple [A] (verified)	119
Fricas [A] (verification not implemented)	120
Sympy [B] (verification not implemented)	120
Maxima [A] (verification not implemented)	121
Giac [A] (verification not implemented)	121
Mupad [B] (verification not implemented)	121
Reduce [B] (verification not implemented)	122

Optimal result

Integrand size = 12, antiderivative size = 31

$$\int x \cos^2(a + bx^2) dx = \frac{x^2}{4} + \frac{\cos(a + bx^2) \sin(a + bx^2)}{4b}$$

output `1/4*x^2+1/4*cos(b*x^2+a)*sin(b*x^2+a)/b`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int x \cos^2(a + bx^2) dx = \frac{2(a + bx^2) + \sin(2(a + bx^2))}{8b}$$

input `Integrate[x*Cos[a + b*x^2]^2,x]`

output `(2*(a + b*x^2) + Sin[2*(a + b*x^2)])/(8*b)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.13, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3861, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos^2(a + bx^2) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{2} \int \cos^2(bx^2 + a) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \sin\left(bx^2 + a + \frac{\pi}{2}\right)^2 dx^2 \\
 & \quad \downarrow \text{3115} \\
 & \frac{1}{2} \left(\frac{\int 1 dx^2}{2} + \frac{\sin(a + bx^2) \cos(a + bx^2)}{2b} \right) \\
 & \quad \downarrow \text{24} \\
 & \frac{1}{2} \left(\frac{\sin(a + bx^2) \cos(a + bx^2)}{2b} + \frac{x^2}{2} \right)
 \end{aligned}$$

input `Int[x*Cos[a + b*x^2]^2,x]`

output `(x^2/2 + (Cos[a + b*x^2]*Sin[a + b*x^2])/(2*b))/2`

Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`
- rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

method	result
risch	$\frac{x^2}{4} + \frac{\sin(2bx^2+2a)}{8b}$
parallelrisch	$\frac{2bx^2+\sin(2bx^2+2a)}{8b}$
derivativedivides	$\frac{\cos(bx^2+a)\sin(bx^2+a)}{2} + \frac{bx^2}{2} + \frac{a}{2}$
default	$\frac{\cos(bx^2+a)\sin(bx^2+a)}{2} + \frac{bx^2}{2} + \frac{a}{2}$
norman	$\frac{x^2}{4} + \frac{\tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)}{2b} - \frac{\tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^3}{2b} + \frac{x^2 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^2}{2} + \frac{x^2 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^4}{4}$ $\left(1 + \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^2\right)^2$
oring	$\frac{(16b^2x^4+5)\cos(bx^2+a)^2}{32x^2b^2} - \frac{5(\cos(bx^2+a)^2-4x^2\cos(bx^2+a)b\sin(bx^2+a))}{32x^2b^2} + \frac{-12\cos(bx^2+a)bx\sin(bx^2+a)}{32x^2b^2}$

input `int(x*cos(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^2+1/8*sin(2*b*x^2+2*a)/b`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int x \cos^2(a + bx^2) dx = \frac{bx^2 + \cos(bx^2 + a) \sin(bx^2 + a)}{4b}$$

input `integrate(x*cos(b*x^2+a)^2,x, algorithm="fricas")`

output `1/4*(b*x^2 + cos(b*x^2 + a)*sin(b*x^2 + a))/b`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. 2(24) = 48.

Time = 0.10 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.94

$$\int x \cos^2(a + bx^2) dx = \begin{cases} \frac{x^2 \sin^2(a + bx^2)}{4} + \frac{x^2 \cos^2(a + bx^2)}{4} + \frac{\sin(a + bx^2) \cos(a + bx^2)}{4b} & \text{for } b \neq 0 \\ \frac{x^2 \cos^2(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*cos(b*x**2+a)**2,x)`

output `Piecewise((x**2*sin(a + b*x**2)**2/4 + x**2*cos(a + b*x**2)**2/4 + sin(a + b*x**2)*cos(a + b*x**2)/(4*b), Ne(b, 0)), (x**2*cos(a)**2/2, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int x \cos^2(a + bx^2) dx = \frac{2bx^2 + \sin(2bx^2 + 2a)}{8b}$$

input `integrate(x*cos(b*x^2+a)^2,x, algorithm="maxima")`

output `1/8*(2*b*x^2 + sin(2*b*x^2 + 2*a))/b`

Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int x \cos^2(a + bx^2) dx = \frac{2bx^2 + 2a + \sin(2bx^2 + 2a)}{8b}$$

input `integrate(x*cos(b*x^2+a)^2,x, algorithm="giac")`

output `1/8*(2*b*x^2 + 2*a + sin(2*b*x^2 + 2*a))/b`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int x \cos^2(a + bx^2) dx = \frac{\sin(2bx^2 + 2a)}{8b} + \frac{x^2}{4}$$

input `int(x*cos(a + b*x^2)^2,x)`

output `sin(2*a + 2*b*x^2)/(8*b) + x^2/4`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int x \cos^2(a + bx^2) dx = \frac{\cos(bx^2 + a) \sin(bx^2 + a) + bx^2}{4b}$$

input `int(x*cos(b*x^2+a)^2,x)`

output `(cos(a + b*x**2)*sin(a + b*x**2) + b*x**2)/(4*b)`

3.11 $\int \cos^2(a + bx^2) dx$

Optimal result	123
Mathematica [A] (verified)	123
Rubi [A] (verified)	124
Maple [A] (verified)	125
Fricas [A] (verification not implemented)	125
Sympy [A] (verification not implemented)	126
Maxima [C] (verification not implemented)	126
Giac [C] (verification not implemented)	127
Mupad [F(-1)]	127
Reduce [F]	128

Optimal result

Integrand size = 10, antiderivative size = 70

$$\int \cos^2(a + bx^2) dx = \frac{x}{2} + \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} - \frac{\sqrt{\pi} \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a)}{4\sqrt{b}}$$

output

```
1/2*x+1/4*Pi^(1/2)*cos(2*a)*FresnelC(2*b^(1/2)*x/Pi^(1/2))/b^(1/2)-1/4*Pi^(1/2)*FresnelS(2*b^(1/2)*x/Pi^(1/2))*sin(2*a)/b^(1/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \cos^2(a + bx^2) dx \\ &= \frac{2\sqrt{bx} + \sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{\pi} \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a)}{4\sqrt{b}} \end{aligned}$$

input

```
Integrate[Cos[a + b*x^2]^2,x]
```

output

```
(2*Sqrt[b]*x + Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x)/Sqrt[Pi]] - Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x)/Sqrt[Pi]]*Sin[2*a])/(4*Sqrt[b])
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3839, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(a + bx^2) dx$$

$$\downarrow \text{3839}$$

$$\int \left(\frac{1}{2} \cos(2a + 2bx^2) + \frac{1}{2} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\sqrt{\pi} \cos(2a) \text{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} - \frac{\sqrt{\pi} \sin(2a) \text{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{4\sqrt{b}} + \frac{x}{2}$$

input

```
Int[Cos[a + b*x^2]^2,x]
```

output

```
x/2 + (Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x)/Sqrt[Pi]])/(4*Sqrt[b]) - (Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x)/Sqrt[Pi]]*Sin[2*a])/(4*Sqrt[b])
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3839 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_), x_Symbol] := Int[ExpandTrigReduce[(a + b*Cos[c + d*(e + f*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 1] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{\pi} \left(\cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right) - \sin(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right) \right)}{4\sqrt{b}}$	45
risch	$\frac{x}{2} + \frac{e^{-2ia} \sqrt{\pi} \sqrt{2} \operatorname{erf}\left(\sqrt{2}\sqrt{ib}x\right)}{16\sqrt{ib}} + \frac{e^{2ia} \sqrt{\pi} \operatorname{erf}\left(\sqrt{-2ib}x\right)}{8\sqrt{-2ib}}$	61

input `int(cos(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output `1/2*x+1/4*Pi^(1/2)/b^(1/2)*(cos(2*a)*FresnelC(2*b^(1/2)*x/Pi^(1/2))-sin(2*a)*FresnelS(2*b^(1/2)*x/Pi^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.84

$$\int \cos^2(a + bx^2) dx = \frac{\pi \sqrt{\frac{b}{\pi}} \cos(2a) C\left(2x\sqrt{\frac{b}{\pi}}\right) - \pi \sqrt{\frac{b}{\pi}} S\left(2x\sqrt{\frac{b}{\pi}}\right) \sin(2a) + 2bx}{4b}$$

input `integrate(cos(b*x^2+a)^2,x, algorithm="fricas")`

output `1/4*(pi*sqrt(b/pi)*cos(2*a)*fresnel_cos(2*x*sqrt(b/pi)) - pi*sqrt(b/pi)*fresnel_sin(2*x*sqrt(b/pi))*sin(2*a) + 2*b*x)/b`

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.80

$$\int \cos^2(a + bx^2) dx = \frac{x}{2} + \frac{\sqrt{\pi} \left(-\sin(2a) S\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(2a) C\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \right) \sqrt{\frac{1}{b}}}{4}$$

input `integrate(cos(b*x**2+a)**2,x)`

output `x/2 + sqrt(pi)*(-sin(2*a)*fresnels(2*sqrt(b)*x/sqrt(pi)) + cos(2*a)*fresnelc(2*sqrt(b)*x/sqrt(pi)))*sqrt(1/b)/4`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00

$$\int \cos^2(a + bx^2) dx = \frac{4^{\frac{1}{4}} \sqrt{2} \sqrt{\pi} \left((i-1) \cos(2a) + (i+1) \sin(2a) \right) \operatorname{erf}\left(\sqrt{2i} \sqrt{bx}\right) + \left(-(i+1) \cos(2a) - (i-1) \sin(2a) \right)}{32 b^2}$$

input `integrate(cos(b*x^2+a)^2,x, algorithm="maxima")`

output `-1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(2*a) + (I + 1)*sin(2*a))*erf(sqrt(2*I*b)*x) + (-(I + 1)*cos(2*a) - (I - 1)*sin(2*a))*erf(sqrt(-2*I*b)*x))*b^(3/2) - 16*b^2*x)/b^2`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.17

$$\int \cos^2(a + bx^2) dx = \frac{1}{2}x - \frac{\sqrt{\pi} \operatorname{erf}\left(-\sqrt{b}x\left(-\frac{ib}{|b|} + 1\right)\right) e^{(2ia)}}{8\sqrt{b}\left(-\frac{ib}{|b|} + 1\right)} - \frac{\sqrt{\pi} \operatorname{erf}\left(-\sqrt{b}x\left(\frac{ib}{|b|} + 1\right)\right) e^{(-2ia)}}{8\sqrt{b}\left(\frac{ib}{|b|} + 1\right)}$$

input `integrate(cos(b*x^2+a)^2,x, algorithm="giac")`

output `1/2*x - 1/8*sqrt(pi)*erf(-sqrt(b)*x*(-I*b/abs(b) + 1))*e^(2*I*a)/(sqrt(b)*(-I*b/abs(b) + 1)) - 1/8*sqrt(pi)*erf(-sqrt(b)*x*(I*b/abs(b) + 1))*e^(-2*I*a)/(sqrt(b)*(I*b/abs(b) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx^2) dx = \int \cos(bx^2 + a)^2 dx$$

input `int(cos(a + b*x^2)^2,x)`

output `int(cos(a + b*x^2)^2, x)`

Reduce [F]

$$\int \cos^2(a + bx^2) dx = \int \cos(bx^2 + a)^2 dx$$

input `int(cos(b*x^2+a)^2,x)`

output `int(cos(a + b*x**2)**2,x)`

3.12 $\int \frac{\cos^2(a+bx^2)}{x} dx$

Optimal result	129
Mathematica [A] (verified)	129
Rubi [A] (verified)	130
Maple [C] (warning: unable to verify)	131
Fricas [A] (verification not implemented)	131
Sympy [F]	131
Maxima [C] (verification not implemented)	132
Giac [A] (verification not implemented)	132
Mupad [F(-1)]	132
Reduce [F]	133

Optimal result

Integrand size = 14, antiderivative size = 37

$$\int \frac{\cos^2(a+bx^2)}{x} dx = \frac{1}{4} \cos(2a) \operatorname{CosIntegral}(2bx^2) + \frac{\log(x)}{2} - \frac{1}{4} \sin(2a) \operatorname{Si}(2bx^2)$$

output

```
1/4*cos(2*a)*Ci(2*b*x^2)+1/2*ln(x)-1/4*sin(2*a)*Si(2*b*x^2)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int \frac{\cos^2(a+bx^2)}{x} dx = \frac{1}{4} (\cos(2a) \operatorname{CosIntegral}(2bx^2) + 2 \log(x) - \sin(2a) \operatorname{Si}(2bx^2))$$

input

```
Integrate[Cos[a + b*x^2]^2/x,x]
```

output

```
(Cos[2*a]*CosIntegral[2*b*x^2] + 2*Log[x] - Sin[2*a]*SinIntegral[2*b*x^2])/4
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(a + bx^2)}{x} dx$$

↓ 3885

$$\int \left(\frac{\cos(2a + 2bx^2)}{2x} + \frac{1}{2x} \right) dx$$

↓ 2009

$$\frac{1}{4} \cos(2a) \text{CosIntegral}(2bx^2) - \frac{1}{4} \sin(2a) \text{Si}(2bx^2) + \frac{\log(x)}{2}$$

input `Int[Cos[a + b*x^2]^2/x,x]`

output `(Cos[2*a]*CosIntegral[2*b*x^2])/4 + Log[x]/2 - (Sin[2*a]*SinIntegral[2*b*x^2])/4`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.33 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.84

method	result	size
risch	$\frac{\ln(x)}{2} + \frac{i\pi \operatorname{csgn}(bx^2)e^{-2ia}}{8} - \frac{i \operatorname{Si}(2bx^2)e^{-2ia}}{4} - \frac{e^{-2ia} \operatorname{expIntegral}_1(-2ibx^2)}{8} - \frac{e^{2ia} \operatorname{expIntegral}_1(-2ibx^2)}{8}$	68

input `int(cos(b*x^2+a)^2/x,x,method=_RETURNVERBOSE)`

output $\frac{1}{2} \ln(x) + \frac{1}{8} i \pi \operatorname{csgn}(bx^2) \exp(-2Ia) - \frac{1}{4} i \operatorname{Si}(2bx^2) \exp(-2Ia) - \frac{1}{8} \exp(-2Ia) \operatorname{Ei}(1, -2Ibx^2) - \frac{1}{8} \exp(2Ia) \operatorname{Ei}(1, -2Ibx^2)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.84

$$\int \frac{\cos^2(a + bx^2)}{x} dx = \frac{1}{4} \cos(2a) \operatorname{Ci}(2bx^2) - \frac{1}{4} \sin(2a) \operatorname{Si}(2bx^2) + \frac{1}{2} \log(x)$$

input `integrate(cos(b*x^2+a)^2/x,x, algorithm="fricas")`

output $\frac{1}{4} \cos(2a) \operatorname{cos_integral}(2bx^2) - \frac{1}{4} \sin(2a) \operatorname{sin_integral}(2bx^2) + \frac{1}{2} \log(x)$

Sympy [F]

$$\int \frac{\cos^2(a + bx^2)}{x} dx = \int \frac{\cos^2(a + bx^2)}{x} dx$$

input `integrate(cos(b*x**2+a)**2/x,x)`

output `Integral(cos(a + b*x**2)**2/x, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.38

$$\int \frac{\cos^2(a + bx^2)}{x} dx = \frac{1}{8} (\operatorname{Ei}(2i bx^2) + \operatorname{Ei}(-2i bx^2)) \cos(2a) + \frac{1}{8} (i \operatorname{Ei}(2i bx^2) - i \operatorname{Ei}(-2i bx^2)) \sin(2a) + \frac{1}{2} \log(x)$$

input `integrate(cos(b*x^2+a)^2/x,x, algorithm="maxima")`

output `1/8*(Ei(2*I*b*x^2) + Ei(-2*I*b*x^2))*cos(2*a) + 1/8*(I*Ei(2*I*b*x^2) - I*Ei(-2*I*b*x^2))*sin(2*a) + 1/2*log(x)`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \frac{\cos^2(a + bx^2)}{x} dx = \frac{1}{4} \cos(2a) \operatorname{Ci}(2bx^2) + \frac{1}{4} \sin(2a) \operatorname{Si}(-2bx^2) + \frac{1}{4} \log(bx^2)$$

input `integrate(cos(b*x^2+a)^2/x,x, algorithm="giac")`

output `1/4*cos(2*a)*cos_integral(2*b*x^2) + 1/4*sin(2*a)*sin_integral(-2*b*x^2) + 1/4*log(b*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^2)}{x} dx = \int \frac{\cos(bx^2 + a)^2}{x} dx$$

input `int(cos(a + b*x^2)^2/x,x)`

output `int(cos(a + b*x^2)^2/x, x)`

Reduce [F]

$$\int \frac{\cos^2(a + bx^2)}{x} dx = \int \frac{\cos(bx^2 + a)^2}{x} dx$$

input `int(cos(b*x^2+a)^2/x, x)`

output `int(cos(a + b*x**2)**2/x, x)`

3.13 $\int \frac{\cos^2(a+bx^2)}{x^2} dx$

Optimal result	134
Mathematica [A] (verified)	134
Rubi [A] (verified)	135
Maple [A] (verified)	137
Fricas [A] (verification not implemented)	137
Sympy [F]	138
Maxima [C] (verification not implemented)	138
Giac [F]	138
Mupad [F(-1)]	139
Reduce [F]	139

Optimal result

Integrand size = 14, antiderivative size = 76

$$\int \frac{\cos^2(a+bx^2)}{x^2} dx = -\frac{\cos^2(a+bx^2)}{x} - \sqrt{b}\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) - \sqrt{b}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a)$$

output

```
-cos(b*x^2+a)^2/x-b^(1/2)*Pi^(1/2)*cos(2*a)*FresnelS(2*b^(1/2)*x/Pi^(1/2))
-b^(1/2)*Pi^(1/2)*FresnelC(2*b^(1/2)*x/Pi^(1/2))*sin(2*a)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(a+bx^2)}{x^2} dx = \frac{\cos^2(a+bx^2) + \sqrt{b}\sqrt{\pi}x \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) + \sqrt{b}\sqrt{\pi}x \operatorname{FresnelC}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right) \sin(2a)}{x}$$

input

```
Integrate[Cos[a + b*x^2]^2/x^2,x]
```

output

```

-((Cos[a + b*x^2]^2 + Sqrt[b]*Sqrt[Pi]*x*Cos[2*a]*FresnelS[(2*Sqrt[b]*x)/Sqrt[Pi]] + Sqrt[b]*Sqrt[Pi]*x*FresnelC[(2*Sqrt[b]*x)/Sqrt[Pi]]*Sin[2*a])/x
)

```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3875, 5084, 3854, 3834, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\cos^2(a + bx^2)}{x^2} dx \\
& \quad \downarrow \text{3875} \\
& -4b \int \cos(bx^2 + a) \sin(bx^2 + a) dx - \frac{\cos^2(a + bx^2)}{x} \\
& \quad \downarrow \text{5084} \\
& -2b \int \sin(2(bx^2 + a)) dx - \frac{\cos^2(a + bx^2)}{x} \\
& \quad \downarrow \text{3854} \\
& -2b \int \sin(2bx^2 + 2a) dx - \frac{\cos^2(a + bx^2)}{x} \\
& \quad \downarrow \text{3834} \\
& -2b \left(\sin(2a) \int \cos(2bx^2) dx + \cos(2a) \int \sin(2bx^2) dx \right) - \frac{\cos^2(a + bx^2)}{x} \\
& \quad \downarrow \text{3832} \\
& -2b \left(\sin(2a) \int \cos(2bx^2) dx + \frac{\sqrt{\pi} \cos(2a) \text{FresnelS}\left(\frac{2\sqrt{bx}}{\sqrt{\pi}}\right)}{2\sqrt{b}} \right) - \frac{\cos^2(a + bx^2)}{x} \\
& \quad \downarrow \text{3833}
\end{aligned}$$

$$-2b \left(\frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelC} \left(\frac{2\sqrt{bx}}{\sqrt{\pi}} \right)}{2\sqrt{b}} + \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelS} \left(\frac{2\sqrt{bx}}{\sqrt{\pi}} \right)}{2\sqrt{b}} \right) - \frac{\cos^2(a + bx^2)}{x}$$

input `Int[Cos[a + b*x^2]^2/x^2,x]`

output `-(Cos[a + b*x^2]^2/x) - 2*b*((Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x)/Sqrt[Pi]])/(2*Sqrt[b]) + (Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x)/Sqrt[Pi]]*Sin[2*a])/(2*Sqrt[b]))`

Defintions of rubi rules used

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3834 `Int[Sin[(c_) + (d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[Sin[c] Int[Cos[d*(e + f*x)^2], x], x] + Simp[Cos[c] Int[Sin[d*(e + f*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]`

rule 3854 `Int[((a_.) + (b_.)*Sin[u_])^(p_.), x_Symbol] := Int[(a + b*Sin[ExpandToSum[u, x]])^p, x] /; FreeQ[{a, b, p}, x] && BinomialQ[u, x] && !BinomialMatchQ[u, x]`

rule 3875 `Int[Cos[(a_.) + (b_.)*(x_)^(n_)]^(p_)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(Cos[a + b*x^n]^p/(m + 1)), x] + Simp[b*n*(p/(m + 1)) Int[Cos[a + b*x^n]^(p - 1)*Sin[a + b*x^n], x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 1] && EqQ[m + n, 0] && NeQ[n, 1] && IntegerQ[n]`

rule 5084

```
Int[Cos[w_]^(p_.)*(u_.)*Sin[v_]^(p_.), x_Symbol] := Simp[1/2^p Int[u*Sin[
2*v]^(p, x), x] /; EqQ[w, v] && IntegerQ[p]
```

Maple [A] (verified)

Time = 0.98 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.82

method	result	size
default	$-\frac{1}{2x} - \frac{\cos(2bx^2+2a)}{2x} - \sqrt{b}\sqrt{\pi} \left(\cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right) + \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}x}{\sqrt{\pi}}\right) \right)$	62
risch	$-\frac{1}{2x} - \frac{ie^{-2ia}b\sqrt{\pi}\sqrt{2}\operatorname{erf}(\sqrt{2}\sqrt{ib}x)}{4\sqrt{ib}} + \frac{ie^{2ia}b\sqrt{\pi}\operatorname{erf}(\sqrt{-2ib}x)}{2\sqrt{-2ib}} - \frac{\cos(2bx^2+2a)}{2x}$	83

input

```
int(cos(b*x^2+a)^2/x^2,x,method=_RETURNVERBOSE)
```

output

```
-1/2/x-1/2/x*cos(2*b*x^2+2*a)-b^(1/2)*Pi^(1/2)*(cos(2*a)*FresnelS(2*b^(1/2)
)*x/Pi^(1/2))+sin(2*a)*FresnelC(2*b^(1/2)*x/Pi^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.87

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx$$

$$= -\frac{\pi x \sqrt{\frac{b}{\pi}} \cos(2a) S\left(2x\sqrt{\frac{b}{\pi}}\right) + \pi x \sqrt{\frac{b}{\pi}} C\left(2x\sqrt{\frac{b}{\pi}}\right) \sin(2a) + \cos(bx^2 + a)^2}{x}$$

input

```
integrate(cos(b*x^2+a)^2/x^2,x, algorithm="fricas")
```

output

```
-(pi*x*sqrt(b/pi)*cos(2*a)*fresnel_sin(2*x*sqrt(b/pi)) + pi*x*sqrt(b/pi)*f
resnel_cos(2*x*sqrt(b/pi))*sin(2*a) + cos(b*x^2 + a)^2)/x
```

Sympy [F]

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx = \int \frac{\cos^2(a + bx^2)}{x^2} dx$$

input `integrate(cos(b*x**2+a)**2/x**2,x)`

output `Integral(cos(a + b*x**2)**2/x**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.09

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx = \frac{\sqrt{2}\sqrt{bx^2}((-i+1)\sqrt{2}\Gamma(-\frac{1}{2}, 2i bx^2) + (i-1)\sqrt{2}\Gamma(-\frac{1}{2}, -2i bx^2))\cos(2a) + ((i-1)\sqrt{2}\Gamma(-\frac{1}{2}, 2i bx^2) - (i+1)\sqrt{2}\Gamma(-\frac{1}{2}, -2i bx^2))\sin(2a)}{16x}$$

input `integrate(cos(b*x^2+a)^2/x^2,x, algorithm="maxima")`

output `1/16*(sqrt(2)*sqrt(b*x^2)*((-I + 1)*sqrt(2)*gamma(-1/2, 2*I*b*x^2) + (I - 1)*sqrt(2)*gamma(-1/2, -2*I*b*x^2))*cos(2*a) + ((I - 1)*sqrt(2)*gamma(-1/2, 2*I*b*x^2) - (I + 1)*sqrt(2)*gamma(-1/2, -2*I*b*x^2))*sin(2*a) - 8)/x`

Giac [F]

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)^2}{x^2} dx$$

input `integrate(cos(b*x^2+a)^2/x^2,x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)^2/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)^2}{x^2} dx$$

input `int(cos(a + b*x^2)^2/x^2,x)`output `int(cos(a + b*x^2)^2/x^2, x)`**Reduce [F]**

$$\int \frac{\cos^2(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)^2}{x^2} dx$$

input `int(cos(b*x^2+a)^2/x^2,x)`output `int(cos(a + b*x**2)**2/x**2,x)`

3.14 $\int \frac{\cos^2(a+bx^2)}{x^3} dx$

Optimal result	140
Mathematica [A] (verified)	140
Rubi [A] (verified)	141
Maple [C] (warning: unable to verify)	142
Fricas [A] (verification not implemented)	142
Sympy [F]	143
Maxima [C] (verification not implemented)	143
Giac [B] (verification not implemented)	143
Mupad [F(-1)]	144
Reduce [F]	144

Optimal result

Integrand size = 14, antiderivative size = 57

$$\int \frac{\cos^2(a+bx^2)}{x^3} dx = -\frac{1}{4x^2} - \frac{\cos(2(a+bx^2))}{4x^2} - \frac{1}{2}b \operatorname{CosIntegral}(2bx^2) \sin(2a) - \frac{1}{2}b \cos(2a) \operatorname{Si}(2bx^2)$$

output

```
-1/4/x^2-1/4*cos(2*b*x^2+2*a)/x^2-1/2*b*Ci(2*b*x^2)*sin(2*a)-1/2*b*cos(2*a)*Si(2*b*x^2)
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.88

$$\int \frac{\cos^2(a+bx^2)}{x^3} dx = -\frac{\cos^2(a+bx^2) + bx^2 \operatorname{CosIntegral}(2bx^2) \sin(2a) + bx^2 \cos(2a) \operatorname{Si}(2bx^2)}{2x^2}$$

input

```
Integrate[Cos[a + b*x^2]^2/x^3,x]
```

output

```
-1/2*(Cos[a + b*x^2]^2 + b*x^2*CosIntegral[2*b*x^2]*Sin[2*a] + b*x^2*Cos[2*a]*SinIntegral[2*b*x^2])/x^2
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx$$

$$\downarrow \text{3885}$$

$$\int \left(\frac{\cos(2a + 2bx^2)}{2x^3} + \frac{1}{2x^3} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{1}{2}b \sin(2a) \text{CosIntegral}(2bx^2) - \frac{1}{2}b \cos(2a) \text{Si}(2bx^2) - \frac{\cos(2(a + bx^2))}{4x^2} - \frac{1}{4x^2}$$

input

```
Int[Cos[a + b*x^2]^2/x^3,x]
```

output

```
-1/4*1/x^2 - Cos[2*(a + b*x^2)]/(4*x^2) - (b*CosIntegral[2*b*x^2]*Sin[2*a])/2 - (b*Cos[2*a]*SinIntegral[2*b*x^2])/2
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_.)]*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.36 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.70

method	result
risch	$-\frac{e^{-2ia}\pi \operatorname{csgn}(bx^2)bx^2 - ie^{-2ia} \operatorname{expIntegral}_1(-2ibx^2)bx^2 + ie^{2ia}b \operatorname{expIntegral}_1(-2ibx^2)x^2 + 2e^{-2ia} \operatorname{Si}(2bx^2)bx^2 + \cos(2bx^2)}$

input `int(cos(b*x^2+a)^2/x^3,x,method=_RETURNVERBOSE)`

output `-1/4*(-exp(-2*I*a)*Pi*csgn(b*x^2)*b*x^2-I*exp(-2*I*a)*Ei(1,-2*I*b*x^2)*b*x^2+I*exp(2*I*a)*b*Ei(1,-2*I*b*x^2)*x^2+2*exp(-2*I*a)*Si(2*b*x^2)*b*x^2+cos(2*b*x^2+2*a)+1)/x^2`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.84

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx = -\frac{bx^2 \operatorname{Ci}(2bx^2) \sin(2a) + bx^2 \cos(2a) \operatorname{Si}(2bx^2) + \cos(bx^2 + a)^2}{2x^2}$$

input `integrate(cos(b*x^2+a)^2/x^3,x, algorithm="fricas")`

output `-1/2*(b*x^2*cos_integral(2*b*x^2)*sin(2*a) + b*x^2*cos(2*a)*sin_integral(2*b*x^2) + cos(b*x^2 + a)^2)/x^2`

Sympy [F]

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx = \int \frac{\cos^2(a + bx^2)}{x^3} dx$$

input `integrate(cos(b*x**2+a)**2/x**3,x)`

output `Integral(cos(a + b*x**2)**2/x**3, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.07

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx = \frac{((i\Gamma(-1, 2i bx^2) - i\Gamma(-1, -2i bx^2)) \cos(2a) + (\Gamma(-1, 2i bx^2) + \Gamma(-1, -2i bx^2)) \sin(2a))bx^2 + 1}{4x^2}$$

input `integrate(cos(b*x^2+a)^2/x^3,x, algorithm="maxima")`

output `-1/4*(((I*gamma(-1, 2*I*b*x^2) - I*gamma(-1, -2*I*b*x^2))*cos(2*a) + (gamma(-1, 2*I*b*x^2) + gamma(-1, -2*I*b*x^2))*sin(2*a))*b*x^2 + 1)/x^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. 2(50) = 100.

Time = 0.34 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.88

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx = \frac{2(bx^2 + a)b^2 \operatorname{Ci}(2bx^2) \sin(2a) - 2ab^2 \operatorname{Ci}(2bx^2) \sin(2a) - 2(bx^2 + a)b^2 \cos(2a) \operatorname{Si}(-2bx^2) + 2ab^2}{4b^2x^2}$$

input `integrate(cos(b*x^2+a)^2/x^3,x, algorithm="giac")`

output `-1/4*(2*(b*x^2 + a)*b^2*cos_integral(2*b*x^2)*sin(2*a) - 2*a*b^2*cos_integ
ral(2*b*x^2)*sin(2*a) - 2*(b*x^2 + a)*b^2*cos(2*a)*sin_integral(-2*b*x^2)
+ 2*a*b^2*cos(2*a)*sin_integral(-2*b*x^2) + b^2*cos(2*b*x^2 + 2*a) + b^2)/
(b^2*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx = \int \frac{\cos(bx^2 + a)^2}{x^3} dx$$

input `int(cos(a + b*x^2)^2/x^3,x)`

output `int(cos(a + b*x^2)^2/x^3, x)`

Reduce [F]

$$\int \frac{\cos^2(a + bx^2)}{x^3} dx = \int \frac{\cos(bx^2 + a)^2}{x^3} dx$$

input `int(cos(b*x^2+a)^2/x^3,x)`

output `int(cos(a + b*x**2)**2/x**3,x)`

3.15 $\int x^3 \cos^3(a + bx^2) dx$

Optimal result	145
Mathematica [A] (verified)	145
Rubi [A] (verified)	146
Maple [A] (verified)	148
Fricas [A] (verification not implemented)	149
Sympy [A] (verification not implemented)	149
Maxima [A] (verification not implemented)	150
Giac [A] (verification not implemented)	150
Mupad [B] (verification not implemented)	151
Reduce [B] (verification not implemented)	151

Optimal result

Integrand size = 14, antiderivative size = 79

$$\int x^3 \cos^3(a + bx^2) dx = \frac{\cos(a + bx^2)}{3b^2} + \frac{\cos^3(a + bx^2)}{18b^2} + \frac{x^2 \sin(a + bx^2)}{3b} + \frac{x^2 \cos^2(a + bx^2) \sin(a + bx^2)}{6b}$$

output

```
1/3*cos(b*x^2+a)/b^2+1/18*cos(b*x^2+a)^3/b^2+1/3*x^2*sin(b*x^2+a)/b+1/6*x^2*cos(b*x^2+a)^2*sin(b*x^2+a)/b
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.70

$$\int x^3 \cos^3(a + bx^2) dx = \frac{27 \cos(a + bx^2) + \cos(3(a + bx^2)) + 3bx^2(9 \sin(a + bx^2) + \sin(3(a + bx^2)))}{72b^2}$$

input

```
Integrate[x^3*Cos[a + b*x^2]^3,x]
```

output

$$\frac{(27*\text{Cos}[a + b*x^2] + \text{Cos}[3*(a + b*x^2)] + 3*b*x^2*(9*\text{Sin}[a + b*x^2] + \text{Sin}[3*(a + b*x^2)]))}{(72*b^2)}$$
Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3861, 3042, 3791, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \cos^3(a + bx^2) dx \\ & \quad \downarrow \text{3861} \\ & \frac{1}{2} \int x^2 \cos^3(bx^2 + a) dx^2 \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2} \int x^2 \sin\left(bx^2 + a + \frac{\pi}{2}\right)^3 dx^2 \\ & \quad \downarrow \text{3791} \\ & \frac{1}{2} \left(\frac{2}{3} \int x^2 \cos(bx^2 + a) dx^2 + \frac{\cos^3(a + bx^2)}{9b^2} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{3b} \right) \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2} \left(\frac{2}{3} \int x^2 \sin\left(bx^2 + a + \frac{\pi}{2}\right) dx^2 + \frac{\cos^3(a + bx^2)}{9b^2} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{3b} \right) \\ & \quad \downarrow \text{3777} \\ & \frac{1}{2} \left(\frac{2}{3} \left(\frac{\int -\sin(bx^2 + a) dx^2}{b} + \frac{x^2 \sin(a + bx^2)}{b} \right) + \frac{\cos^3(a + bx^2)}{9b^2} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{3b} \right) \\ & \quad \downarrow \text{25} \end{aligned}$$

$$\frac{1}{2} \left(\frac{2}{3} \left(\frac{x^2 \sin(a + bx^2)}{b} - \frac{\int \sin(bx^2 + a) dx^2}{b} \right) + \frac{\cos^3(a + bx^2)}{9b^2} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{3b} \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{2}{3} \left(\frac{x^2 \sin(a + bx^2)}{b} - \frac{\int \sin(bx^2 + a) dx^2}{b} \right) + \frac{\cos^3(a + bx^2)}{9b^2} + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{3b} \right)$$

↓ 3118

$$\frac{1}{2} \left(\frac{\cos^3(a + bx^2)}{9b^2} + \frac{2}{3} \left(\frac{\cos(a + bx^2)}{b^2} + \frac{x^2 \sin(a + bx^2)}{b} \right) + \frac{x^2 \sin(a + bx^2) \cos^2(a + bx^2)}{3b} \right)$$

input `Int[x^3*Cos[a + b*x^2]^3,x]`

output `(Cos[a + b*x^2]^3/(9*b^2) + (x^2*Cos[a + b*x^2]^2*Sin[a + b*x^2])/(3*b) + (2*(Cos[a + b*x^2]/b^2 + (x^2*Sin[a + b*x^2])/b))/3)/2`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3791

```
Int[((c._) + (d._)*(x_))*((b._)*sin[(e._) + (f._)*(x_)])^(n_), x_Symbol] :=
Simp[d*((b*SIN[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]
]*((b*SIN[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n Int[(c + d*
x)*(b*SIN[e + f*x])^(n - 2), x], x)] /; FreeQ[{b, c, d, e, f}, x] && GtQ[n,
1]
```

rule 3861

```
Int[((a._) + Cos[(c._) + (d._)*(x_)^(n_)]*(b._))^(p._)*(x_)^(m_), x_Symbol
] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 1.73 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

method	result
default	$\frac{3x^2 \sin(bx^2+a)}{8b} + \frac{3 \cos(bx^2+a)}{8b^2} + \frac{x^2 \sin(3bx^2+3a)}{24b} + \frac{\cos(3bx^2+3a)}{72b^2}$
risch	$\frac{3x^2 \sin(bx^2+a)}{8b} + \frac{3 \cos(bx^2+a)}{8b^2} + \frac{x^2 \sin(3bx^2+3a)}{24b} + \frac{\cos(3bx^2+3a)}{72b^2}$
parallelrisch	$\frac{7+9 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^5 x^2 b+6 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^3 x^2 b+9 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right) x^2 b+9 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^4+12 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^2}{9b^2\left(1+\tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)\right)^3}$
norman	$\frac{x^2 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)}{b} + \frac{x^2 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^5}{b} + \frac{\tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^4}{b^2} + \frac{7}{9b^2} + \frac{2x^2 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^3}{3b} + \frac{4 \tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)^2}{3b^2}{\left(1+\tan\left(\frac{a}{2}+\frac{bx^2}{2}\right)\right)^3}$
orering	$\frac{5(40b^2x^4+27) \cos(bx^2+a)^3}{144x^4b^4} - \frac{(40b^2x^4+71)(3x^2 \cos(bx^2+a)^3-6x^4 \cos(bx^2+a)^2 b \sin(bx^2+a))}{144b^4x^6} + \frac{7x \cos(bx^2+a)^3}{12} - \frac{49x^3 \cos(bx^2+a)^3}{144b^4x^6}$

input

```
int(x^3*cos(b*x^2+a)^3,x,method=_RETURNVERBOSE)
```

output

```
3/8*x^2*sin(b*x^2+a)/b+3/8*cos(b*x^2+a)/b^2+1/24/b*x^2*sin(3*b*x^2+3*a)+1/
72/b^2*cos(3*b*x^2+3*a)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.73

$$\int x^3 \cos^3(a + bx^2) dx$$

$$= \frac{\cos(bx^2 + a)^3 + 3(bx^2 \cos(bx^2 + a)^2 + 2bx^2) \sin(bx^2 + a) + 6 \cos(bx^2 + a)}{18b^2}$$

input `integrate(x^3*cos(b*x^2+a)^3,x, algorithm="fricas")`output `1/18*(cos(b*x^2 + a)^3 + 3*(b*x^2*cos(b*x^2 + a)^2 + 2*b*x^2)*sin(b*x^2 + a) + 6*cos(b*x^2 + a))/b^2`**Sympy [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.16

$$\int x^3 \cos^3(a + bx^2) dx$$

$$= \begin{cases} \frac{x^2 \sin^3(a+bx^2)}{3b} + \frac{x^2 \sin(a+bx^2) \cos^2(a+bx^2)}{2b} + \frac{\sin^2(a+bx^2) \cos(a+bx^2)}{3b^2} + \frac{7 \cos^3(a+bx^2)}{18b^2} & \text{for } b \neq 0 \\ \frac{x^4 \cos^3(a)}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*cos(b*x**2+a)**3,x)`output `Piecewise((x**2*sin(a + b*x**2)**3/(3*b) + x**2*sin(a + b*x**2)*cos(a + b*x**2)**2/(2*b) + sin(a + b*x**2)**2*cos(a + b*x**2)/(3*b**2) + 7*cos(a + b*x**2)**3/(18*b**2), Ne(b, 0)), (x**4*cos(a)**3/4, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.73

$$\int x^3 \cos^3(a + bx^2) dx = \frac{3bx^2 \sin(3bx^2 + 3a) + 27bx^2 \sin(bx^2 + a) + \cos(3bx^2 + 3a) + 27 \cos(bx^2 + a)}{72b^2}$$

input `integrate(x^3*cos(b*x^2+a)^3,x, algorithm="maxima")`output `1/72*(3*b*x^2*sin(3*b*x^2 + 3*a) + 27*b*x^2*sin(b*x^2 + a) + cos(3*b*x^2 + 3*a) + 27*cos(b*x^2 + a))/b^2`**Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.16

$$\int x^3 \cos^3(a + bx^2) dx = \frac{(\sin(bx^2 + a)^3 - 3 \sin(bx^2 + a))a}{6b^2} + \frac{3(bx^2 + a) \sin(3bx^2 + 3a) + 27(bx^2 + a) \sin(bx^2 + a) + \cos(3bx^2 + 3a) + 27 \cos(bx^2 + a)}{72b^2}$$

input `integrate(x^3*cos(b*x^2+a)^3,x, algorithm="giac")`output `1/6*(sin(b*x^2 + a)^3 - 3*sin(b*x^2 + a))*a/b^2 + 1/72*(3*(b*x^2 + a)*sin(3*b*x^2 + 3*a) + 27*(b*x^2 + a)*sin(b*x^2 + a) + cos(3*b*x^2 + 3*a) + 27*cos(b*x^2 + a))/b^2`

Mupad [B] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

$$\int x^3 \cos^3(a + bx^2) dx$$

$$= \frac{\frac{\cos(bx^2+a)}{3} + \frac{\cos(bx^2+a)^3}{18} + b \left(\frac{x^2 \sin(bx^2+a)}{3} + \frac{x^2 \cos(bx^2+a)^2 \sin(bx^2+a)}{6} \right)}{b^2}$$

input `int(x^3*cos(a + b*x^2)^3,x)`output `(cos(a + b*x^2)/3 + cos(a + b*x^2)^3/18 + b*((x^2*sin(a + b*x^2))/3 + (x^2*cos(a + b*x^2)^2*sin(a + b*x^2))/6))/b^2`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.85

$$\int x^3 \cos^3(a + bx^2) dx$$

$$= \frac{-\cos(bx^2 + a) \sin(bx^2 + a)^2 + 7 \cos(bx^2 + a) - 3 \sin(bx^2 + a)^3 b x^2 + 9 \sin(bx^2 + a) b x^2 + 1}{18b^2}$$

input `int(x^3*cos(b*x^2+a)^3,x)`output `(- cos(a + b*x**2)*sin(a + b*x**2)**2 + 7*cos(a + b*x**2) - 3*sin(a + b*x**2)**3*b*x**2 + 9*sin(a + b*x**2)*b*x**2 + 1)/(18*b**2)`

3.16 $\int x^2 \cos^3(a + bx^2) dx$

Optimal result	152
Mathematica [A] (verified)	153
Rubi [A] (verified)	153
Maple [A] (verified)	154
Fricas [A] (verification not implemented)	155
Sympy [B] (verification not implemented)	156
Maxima [C] (verification not implemented)	157
Giac [C] (verification not implemented)	158
Mupad [F(-1)]	159
Reduce [F]	159

Optimal result

Integrand size = 14, antiderivative size = 188

$$\int x^2 \cos^3(a + bx^2) dx = -\frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \cos(3a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{24b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) \sin(3a)}{24b^{3/2}} + \frac{3x \sin(a + bx^2)}{8b} + \frac{x \sin(3a + 3bx^2)}{24b}$$

output

```
-3/16*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)/b^(3/2)
-1/144*6^(1/2)*Pi^(1/2)*cos(3*a)*FresnelS(b^(1/2)*6^(1/2)/Pi^(1/2)*x)/b^(3/2)
-3/16*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)*sin(a)/b^(3/2)
-1/144*6^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*6^(1/2)/Pi^(1/2)*x)*sin(3*a)/b^(3/2)
+3/8*x*sin(b*x^2+a)/b+1/24*x*sin(3*b*x^2+3*a)/b
```

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.85

$$\int x^2 \cos^3(a + bx^2) dx = \frac{-27\sqrt{2\pi} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \sqrt{6\pi} \cos(3a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) - 27\sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) + 27\sqrt{6\pi} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{144b^{3/2}}$$

input

```
Integrate[x^2*Cos[a + b*x^2]^3,x]
```

output

```
(-27*Sqrt[2*Pi]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x] - Sqrt[6*Pi]*Cos[3*a]*FresnelS[Sqrt[b]*Sqrt[6/Pi]*x] - 27*Sqrt[2*Pi]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x]*Sin[a] - Sqrt[6*Pi]*FresnelC[Sqrt[b]*Sqrt[6/Pi]*x]*Sin[3*a] + 54*Sqrt[b]*x*Ssin[a + b*x^2] + 6*Sqrt[b]*x*Ssin[3*(a + b*x^2)])/(144*b^(3/2))
```

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos^3(a + bx^2) dx$$

$$\downarrow \text{3885}$$

$$\int \left(\frac{3}{4}x^2 \cos(a + bx^2) + \frac{1}{4}x^2 \cos(3a + 3bx^2) \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
 & -\frac{3\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \sin(3a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{24b^{3/2}} - \\
 & \frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{8b^{3/2}} - \frac{\sqrt{\frac{\pi}{6}} \cos(3a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{24b^{3/2}} + \frac{3x \sin(a + bx^2)}{8b} + \\
 & \frac{x \sin(3a + 3bx^2)}{24b}
 \end{aligned}$$

input `Int[x^2*cos[a + b*x^2]^3,x]`

output `(-3*Sqrt[Pi/2]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x])/(8*b^(3/2)) - (Sqrt[Pi/6]*Cos[3*a]*FresnelS[Sqrt[b]*Sqrt[6/Pi]*x])/(24*b^(3/2)) - (3*Sqrt[Pi/2]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x]*Sin[a])/(8*b^(3/2)) - (Sqrt[Pi/6]*FresnelC[Sqrt[b]*Sqrt[6/Pi]*x]*Sin[3*a])/(24*b^(3/2)) + (3*x*Sin[a + b*x^2])/(8*b) + (x*Sin[3*a + 3*b*x^2])/(24*b)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_.)]*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 1.67 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.69

method	result
default	$\frac{3x \sin(bx^2+a)}{8b} - \frac{3\sqrt{2}\sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) \right)}{16b^{\frac{3}{2}}} + \frac{x \sin(3bx^2+3a)}{24b} - \frac{\sqrt{2}\sqrt{\pi}\sqrt{3} \left(\cos(3a) \right)}{24b}$
risch	$-\frac{ie^{-3ia}\sqrt{\pi}\sqrt{3} \operatorname{erf}\left(\sqrt{3}\sqrt{ib}x\right)}{288b\sqrt{ib}} - \frac{3ie^{-ia}\sqrt{\pi} \operatorname{erf}\left(\sqrt{ib}x\right)}{32b\sqrt{ib}} + \frac{3ie^{ia}\sqrt{\pi} \operatorname{erf}\left(\sqrt{-ib}x\right)}{32b\sqrt{-ib}} + \frac{ie^{3ia}\sqrt{\pi} \operatorname{erf}\left(\sqrt{-3ib}x\right)}{96b\sqrt{-3ib}} + \frac{3x \sin(bx^2+a)}{8b}$

input `int(x^2*cos(b*x^2+a)^3,x,method=_RETURNVERBOSE)`

output

```
3/8*x*sin(b*x^2+a)/b-3/16/b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)
)*2^(1/2)/Pi^(1/2)*x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x))+1/24*x*
sin(3*b*x^2+3*a)/b-1/144/b^(3/2)*2^(1/2)*Pi^(1/2)*3^(1/2)*(cos(3*a)*Fresne
lS(2^(1/2)/Pi^(1/2)*3^(1/2)*b^(1/2)*x)+sin(3*a)*FresnelC(2^(1/2)/Pi^(1/2)*
3^(1/2)*b^(1/2)*x))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.79

$$\int x^2 \cos^3(a + bx^2) dx =$$

$$\frac{\sqrt{6}\pi\sqrt{\frac{b}{\pi}}\cos(3a)S\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right) + 27\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) + \sqrt{6}\pi\sqrt{\frac{b}{\pi}}C\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right)\sin(3a) + 27\sqrt{2}\pi\sqrt{\frac{b}{\pi}}C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a) - 24(bx^2 + a)^2 + 2bx^2}{144b^2}$$

input

```
integrate(x^2*cos(b*x^2+a)^3,x, algorithm="fricas")
```

output

```
-1/144*(sqrt(6)*pi*sqrt(b/pi)*cos(3*a)*fresnel_sin(sqrt(6)*x*sqrt(b/pi)) +
27*sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x*sqrt(b/pi)) + sqrt(
6)*pi*sqrt(b/pi)*fresnel_cos(sqrt(6)*x*sqrt(b/pi))*sin(3*a) + 27*sqrt(2)*p
i*sqrt(b/pi)*fresnel_cos(sqrt(2)*x*sqrt(b/pi))*sin(a) - 24*(b*x*cos(b*x^2
+ a)^2 + 2*b*x)*sin(b*x^2 + a))/b^2
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 439 vs. $2(194) = 388$.

Time = 1.94 (sec) , antiderivative size = 439, normalized size of antiderivative = 2.34

$$\begin{aligned}
 \int x^2 \cos^3(a + bx^2) dx = & \frac{3b^{\frac{3}{2}}x^5 \sqrt{\frac{1}{b}} \sin(a) \Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{5}{4}\right) {}_2F_3\left(\begin{matrix} \frac{3}{4}, \frac{5}{4} \\ \frac{3}{2}, \frac{7}{4}, \frac{9}{4} \end{matrix} \middle| -\frac{b^2x^4}{4}\right)}{32\Gamma\left(\frac{7}{4}\right) \Gamma\left(\frac{9}{4}\right)} \\
 & + \frac{3b^{\frac{3}{2}}x^5 \sqrt{\frac{1}{b}} \sin(3a) \Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{5}{4}\right) {}_2F_3\left(\begin{matrix} \frac{3}{4}, \frac{5}{4} \\ \frac{3}{2}, \frac{7}{4}, \frac{9}{4} \end{matrix} \middle| -\frac{9b^2x^4}{4}\right)}{32\Gamma\left(\frac{7}{4}\right) \Gamma\left(\frac{9}{4}\right)} \\
 & - \frac{3\sqrt{b}x^3 \sqrt{\frac{1}{b}} \cos(a) \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right) {}_2F_3\left(\begin{matrix} \frac{1}{4}, \frac{3}{4} \\ \frac{1}{2}, \frac{5}{4}, \frac{7}{4} \end{matrix} \middle| -\frac{b^2x^4}{4}\right)}{32\Gamma\left(\frac{5}{4}\right) \Gamma\left(\frac{7}{4}\right)} \\
 & - \frac{\sqrt{b}x^3 \sqrt{\frac{1}{b}} \cos(3a) \Gamma\left(\frac{1}{4}\right) \Gamma\left(\frac{3}{4}\right) {}_2F_3\left(\begin{matrix} \frac{1}{4}, \frac{3}{4} \\ \frac{1}{2}, \frac{5}{4}, \frac{7}{4} \end{matrix} \middle| -\frac{9b^2x^4}{4}\right)}{32\Gamma\left(\frac{5}{4}\right) \Gamma\left(\frac{7}{4}\right)} \\
 & - \frac{3\sqrt{2}\sqrt{\pi}x^2 \sqrt{\frac{1}{b}} \sin(a) S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)}{8} \\
 & - \frac{\sqrt{6}\sqrt{\pi}x^2 \sqrt{\frac{1}{b}} \sin(3a) S\left(\frac{\sqrt{6}\sqrt{bx}}{\sqrt{\pi}}\right)}{24} \\
 & + \frac{3\sqrt{2}\sqrt{\pi}x^2 \sqrt{\frac{1}{b}} \cos(a) C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)}{8} \\
 & + \frac{\sqrt{6}\sqrt{\pi}x^2 \sqrt{\frac{1}{b}} \cos(3a) C\left(\frac{\sqrt{6}\sqrt{bx}}{\sqrt{\pi}}\right)}{24}
 \end{aligned}$$

input

```
integrate(x**2*cos(b*x**2+a)**3,x)
```

output

```

3*b**(3/2)*x**5*sqrt(1/b)*sin(a)*gamma(3/4)*gamma(5/4)*hyper((3/4, 5/4), (
3/2, 7/4, 9/4), -b**2*x**4/4)/(32*gamma(7/4)*gamma(9/4)) + 3*b**(3/2)*x**5
*sqrt(1/b)*sin(3*a)*gamma(3/4)*gamma(5/4)*hyper((3/4, 5/4), (3/2, 7/4, 9/4
), -9*b**2*x**4/4)/(32*gamma(7/4)*gamma(9/4)) - 3*sqrt(b)*x**3*sqrt(1/b)*c
os(a)*gamma(1/4)*gamma(3/4)*hyper((1/4, 3/4), (1/2, 5/4, 7/4), -b**2*x**4/
4)/(32*gamma(5/4)*gamma(7/4)) - sqrt(b)*x**3*sqrt(1/b)*cos(3*a)*gamma(1/4)
*gamma(3/4)*hyper((1/4, 3/4), (1/2, 5/4, 7/4), -9*b**2*x**4/4)/(32*gamma(5
/4)*gamma(7/4)) - 3*sqrt(2)*sqrt(pi)*x**2*sqrt(1/b)*sin(a)*fresnels(sqrt(2)
)*sqrt(b)*x/sqrt(pi))/8 - sqrt(6)*sqrt(pi)*x**2*sqrt(1/b)*sin(3*a)*fresnel
s(sqrt(6)*sqrt(b)*x/sqrt(pi))/24 + 3*sqrt(2)*sqrt(pi)*x**2*sqrt(1/b)*cos(a)
*fresnelc(sqrt(2)*sqrt(b)*x/sqrt(pi))/8 + sqrt(6)*sqrt(pi)*x**2*sqrt(1/b)
*cos(3*a)*fresnelc(sqrt(6)*sqrt(b)*x/sqrt(pi))/24

```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.76

$$\int x^2 \cos^3(a + bx^2) dx$$

$$= \frac{24b^2x \sin(3bx^2 + 3a) + 216b^2x \sin(bx^2 + a) + 9^{\frac{1}{4}}\sqrt{2}\sqrt{\pi} \left(-(i+1) \cos(3a) + (i-1) \sin(3a) \right) \operatorname{erf} \left(\sqrt{3Ib}x \right) + ((I-1)\cos(3a) - (I+1)\sin(3a)) \operatorname{erf}(\sqrt{-3Ib}x) b^{3/2} - 27\sqrt{2}\sqrt{\pi} \left(((I+1)\cos(a) - (I-1)\sin(a)) \operatorname{erf}(\sqrt{Ib}x) + (-(I-1)\cos(a) + (I+1)\sin(a)) \operatorname{erf}(\sqrt{-Ib}x) \right) b^{3/2}}{b^3}$$

input

```
integrate(x^2*cos(b*x^2+a)^3,x, algorithm="maxima")
```

output

```

1/576*(24*b^2*x*sin(3*b*x^2 + 3*a) + 216*b^2*x*sin(b*x^2 + a) + 9^(1/4)*sq
rt(2)*sqrt(pi)*((-I + 1)*cos(3*a) + (I - 1)*sin(3*a))*erf(sqrt(3*I*b)*x)
+ ((I - 1)*cos(3*a) - (I + 1)*sin(3*a))*erf(sqrt(-3*I*b)*x))*b^(3/2) - 27*
sqrt(2)*sqrt(pi)*(((I + 1)*cos(a) - (I - 1)*sin(a))*erf(sqrt(I*b)*x) + (-(
I - 1)*cos(a) + (I + 1)*sin(a))*erf(sqrt(-I*b)*x))*b^(3/2))/b^3

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.38

$$\int x^2 \cos^3(a + bx^2) dx = -\frac{ixe^{(3ibx^2+3ia)}}{48b} - \frac{3ixe^{(ibx^2+ia)}}{16b} + \frac{3ixe^{(-ibx^2-ia)}}{16b}$$

$$+ \frac{ixe^{(-3ibx^2-3ia)}}{48b} - \frac{i\sqrt{6}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{6}\sqrt{bx}\left(-\frac{ib}{|b|} + 1\right)\right) e^{(3ia)}}{288b^{\frac{3}{2}}\left(-\frac{ib}{|b|} + 1\right)}$$

$$- \frac{3i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{(ia)}}{32b\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}}$$

$$+ \frac{3i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{(-ia)}}{32b\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}}$$

$$+ \frac{i\sqrt{6}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{6}\sqrt{bx}\left(\frac{ib}{|b|} + 1\right)\right) e^{(-3ia)}}{288b^{\frac{3}{2}}\left(\frac{ib}{|b|} + 1\right)}$$

input `integrate(x^2*cos(b*x^2+a)^3,x, algorithm="giac")`

output `-1/48*I*x*e^(3*I*b*x^2 + 3*I*a)/b - 3/16*I*x*e^(I*b*x^2 + I*a)/b + 3/16*I*x*e^(-I*b*x^2 - I*a)/b + 1/48*I*x*e^(-3*I*b*x^2 - 3*I*a)/b - 1/288*I*sqrt(6)*sqrt(pi)*erf(-1/2*sqrt(6)*sqrt(b)*x*(-I*b/abs(b) + 1))*e^(3*I*a)/(b^(3/2)*(-I*b/abs(b) + 1)) - 3/32*I*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(-I*b/abs(b) + 1)*sqrt(abs(b)))*e^(I*a)/(b*(-I*b/abs(b) + 1)*sqrt(abs(b))) + 3/32*I*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(I*b/abs(b) + 1)*sqrt(abs(b)))*e^(-I*a)/(b*(I*b/abs(b) + 1)*sqrt(abs(b))) + 1/288*I*sqrt(6)*sqrt(pi)*erf(-1/2*sqrt(6)*sqrt(b)*x*(I*b/abs(b) + 1))*e^(-3*I*a)/(b^(3/2)*(I*b/abs(b) + 1))`

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^3(a + bx^2) dx = \int x^2 \cos(bx^2 + a)^3 dx$$

input `int(x^2*cos(a + b*x^2)^3,x)`output `int(x^2*cos(a + b*x^2)^3, x)`**Reduce [F]**

$$\int x^2 \cos^3(a + bx^2) dx$$

$$= \frac{-9 \cos(bx^2 + a) \sin(bx^2 + a) x + 60 \left(\int \frac{\tan\left(\frac{bx^2 + a}{2}\right)^3}{\tan\left(\frac{bx^2 + a}{2}\right)^6 + 3 \tan\left(\frac{bx^2 + a}{2}\right)^4 + 3 \tan\left(\frac{bx^2 + a}{2}\right)^2 + 1} dx \right) + 96 \left(\int \frac{1}{\tan\left(\frac{bx^2 + a}{2}\right)^6 + 3 \tan\left(\frac{bx^2 + a}{2}\right)^4 + 3 \tan\left(\frac{bx^2 + a}{2}\right)^2 + 1} dx \right)}$$

input `int(x^2*cos(b*x^2+a)^3,x)`output `(- 9*cos(a + b*x**2)*sin(a + b*x**2)*x + 60*int(tan((a + b*x**2)/2)**3/(tan((a + b*x**2)/2)**6 + 3*tan((a + b*x**2)/2)**4 + 3*tan((a + b*x**2)/2)**2 + 1),x) + 96*int(x**2/(tan((a + b*x**2)/2)**6 + 3*tan((a + b*x**2)/2)**4 + 3*tan((a + b*x**2)/2)**2 + 1),x)*b + 36*int(tan((a + b*x**2)/2)/(tan((a + b*x**2)/2)**6 + 3*tan((a + b*x**2)/2)**4 + 3*tan((a + b*x**2)/2)**2 + 1),x) - 3*sin(a + b*x**2)**3*x - 9*sin(a + b*x**2)*x - 10*b*x**3)/(30*b)`

3.17 $\int x \cos^3(a + bx^2) dx$

Optimal result	160
Mathematica [A] (verified)	160
Rubi [A] (warning: unable to verify)	161
Maple [A] (verified)	162
Fricas [A] (verification not implemented)	163
Sympy [A] (verification not implemented)	163
Maxima [A] (verification not implemented)	164
Giac [A] (verification not implemented)	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

Optimal result

Integrand size = 12, antiderivative size = 33

$$\int x \cos^3(a + bx^2) dx = \frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{6b}$$

output `1/2*sin(b*x^2+a)/b-1/6*sin(b*x^2+a)^3/b`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int x \cos^3(a + bx^2) dx = \frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{6b}$$

input `Integrate[x*Cos[a + b*x^2]^3,x]`

output `Sin[a + b*x^2]/(2*b) - Sin[a + b*x^2]^3/(6*b)`

Rubi [A] (warning: unable to verify)

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3861, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos^3(a + bx^2) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{2} \int \cos^3(bx^2 + a) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \sin\left(bx^2 + a + \frac{\pi}{2}\right)^3 dx^2 \\
 & \quad \downarrow \text{3113} \\
 & -\frac{\int (1 - x^4) d(-\sin(bx^2 + a))}{2b} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\sin(a + bx^2) - \frac{x^6}{3}}{2b}
 \end{aligned}$$

input `Int[x*Cos[a + b*x^2]^3,x]`

output `-1/2*(-1/3*x^6 - Sin[a + b*x^2])/b`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 2.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

method	result
derivativedivides	$\frac{(2+\cos(bx^2+a))^2 \sin(bx^2+a)}{6b}$
default	$\frac{(2+\cos(bx^2+a))^2 \sin(bx^2+a)}{6b}$
parallelrisc	$\frac{9 \sin(bx^2+a) + \sin(3bx^2+3a)}{24b}$
risc	$\frac{3 \sin(bx^2+a)}{8b} + \frac{\sin(3bx^2+3a)}{24b}$
norman	$\frac{\tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)}{b} + \frac{\tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^5}{b} + \frac{2 \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)^3}{3b}$ $\left(1 + \tan\left(\frac{a}{2} + \frac{bx^2}{2}\right)\right)^3$
orering	$\frac{5(8b^2x^4+3) \cos(bx^2+a)^3}{144b^4x^6} - \frac{5(8b^2x^4+3) (\cos(bx^2+a)^3 - 6x^2 \cos(bx^2+a)^2 b \sin(bx^2+a))}{144b^4x^6} + \frac{-18 \cos(bx^2+a)^2 b x}{144b^4x^6}$

input `int(x*cos(b*x^2+a)^3,x,method=_RETURNVERBOSE)`

output `1/6/b*(2+cos(b*x^2+a)^2)*sin(b*x^2+a)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

$$\int x \cos^3(a + bx^2) dx = \frac{(\cos(bx^2 + a)^2 + 2) \sin(bx^2 + a)}{6b}$$

input `integrate(x*cos(b*x^2+a)^3,x, algorithm="fricas")`

output `1/6*(cos(b*x^2 + a)^2 + 2)*sin(b*x^2 + a)/b`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.33

$$\int x \cos^3(a + bx^2) dx = \begin{cases} \frac{\sin^3(a+bx^2)}{3b} + \frac{\sin(a+bx^2)\cos^2(a+bx^2)}{2b} & \text{for } b \neq 0 \\ \frac{x^2 \cos^3(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*cos(b*x**2+a)**3,x)`

output `Piecewise((sin(a + b*x**2)**3/(3*b) + sin(a + b*x**2)*cos(a + b*x**2)**2/(2*b), Ne(b, 0)), (x**2*cos(a)**3/2, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int x \cos^3(a + bx^2) dx = \frac{\sin(3bx^2 + 3a) + 9 \sin(bx^2 + a)}{24b}$$

input `integrate(x*cos(b*x^2+a)^3,x, algorithm="maxima")`output `1/24*(sin(3*b*x^2 + 3*a) + 9*sin(b*x^2 + a))/b`**Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int x \cos^3(a + bx^2) dx = -\frac{\sin(bx^2 + a)^3 - 3 \sin(bx^2 + a)}{6b}$$

input `integrate(x*cos(b*x^2+a)^3,x, algorithm="giac")`output `-1/6*(sin(b*x^2 + a)^3 - 3*sin(b*x^2 + a))/b`**Mupad [B] (verification not implemented)**

Time = 41.51 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.85

$$\int x \cos^3(a + bx^2) dx = \frac{3 \sin(bx^2 + a) - \sin(bx^2 + a)^3}{6b}$$

input `int(x*cos(a + b*x^2)^3,x)`output `(3*sin(a + b*x^2) - sin(a + b*x^2)^3)/(6*b)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int x \cos^3(a + bx^2) dx = \frac{\sin(bx^2 + a) \left(-\sin(bx^2 + a)^2 + 3 \right)}{6b}$$

input `int(x*cos(b*x^2+a)^3,x)`

output `(sin(a + b*x**2)*(- sin(a + b*x**2)**2 + 3))/(6*b)`

3.18 $\int \cos^3(a + bx^2) dx$

Optimal result	166
Mathematica [A] (verified)	167
Rubi [A] (verified)	167
Maple [A] (verified)	168
Fricas [A] (verification not implemented)	169
Sympy [A] (verification not implemented)	169
Maxima [C] (verification not implemented)	170
Giac [C] (verification not implemented)	170
Mupad [F(-1)]	171
Reduce [F]	171

Optimal result

Integrand size = 10, antiderivative size = 153

$$\int \cos^3(a + bx^2) dx = \frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{6}} \cos(3a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{4\sqrt{b}} - \frac{3\sqrt{\frac{\pi}{2}} \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) \sin(a)}{4\sqrt{b}} - \frac{\sqrt{\frac{\pi}{6}} \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) \sin(3a)}{4\sqrt{b}}$$

output

```
3/8*2^(1/2)*Pi^(1/2)*cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)/b^(1/2)+1/24*6^(1/2)*Pi^(1/2)*cos(3*a)*FresnelC(b^(1/2)*6^(1/2)/Pi^(1/2)*x)/b^(1/2)-3/8*2^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)*sin(a)/b^(1/2)-1/24*6^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*6^(1/2)/Pi^(1/2)*x)*sin(3*a)/b^(1/2)
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.76

$$\int \cos^3(a + bx^2) dx$$

$$= \frac{\sqrt{\frac{\pi}{6}} \left(3\sqrt{3} \cos(a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} x \right) + \cos(3a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{6}{\pi}} x \right) - 3\sqrt{3} \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} x \right) \sin(a) \right)}{4\sqrt{b}}$$

input `Integrate[Cos[a + b*x^2]^3,x]`

output

```
(Sqrt [Pi/6]*(3*Sqrt [3]*Cos [a]*FresnelC[Sqrt [b]*Sqrt [2/Pi]*x] + Cos [3*a]*FresnelC[Sqrt [b]*Sqrt [6/Pi]*x] - 3*Sqrt [3]*FresnelS[Sqrt [b]*Sqrt [2/Pi]*x]*Sin [a] - FresnelS[Sqrt [b]*Sqrt [6/Pi]*x]*Sin [3*a]))/(4*Sqrt [b])
```

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3839, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(a + bx^2) dx$$

$$\downarrow \text{3839}$$

$$\int \left(\frac{3}{4} \cos(a + bx^2) + \frac{1}{4} \cos(3a + 3bx^2) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} x \right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{6}} \cos(3a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{6}{\pi}} x \right)}{4\sqrt{b}} - \frac{3\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} x \right)}{4\sqrt{b}} - \frac{\sqrt{\frac{\pi}{6}} \sin(3a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{6}{\pi}} x \right)}{4\sqrt{b}}$$

input `Int[Cos[a + b*x^2]^3,x]`

output $(3\sqrt{\pi/2}\cos[a]\operatorname{FresnelC}[\sqrt{b}\sqrt{2/\pi}x])/(4\sqrt{b}) + (\sqrt{\pi/6}\cos[3a]\operatorname{FresnelC}[\sqrt{b}\sqrt{6/\pi}x])/(4\sqrt{b}) - (3\sqrt{\pi/2}\operatorname{FresnelS}[\sqrt{b}\sqrt{2/\pi}x]\sin[a])/(4\sqrt{b}) - (\sqrt{\pi/6}\operatorname{FresnelS}[\sqrt{b}\sqrt{6/\pi}x]\sin[3a])/(4\sqrt{b})$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3839 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_), x_Symbol] := Int[ExpandTrigReduce[(a + b*Cos[c + d*(e + f*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 1] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.42 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.66

method	result
default	$\frac{3\sqrt{2}\sqrt{\pi}\left(\cos(a)\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) - \sin(a)\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right)\right)}{8\sqrt{b}} + \frac{\sqrt{2}\sqrt{\pi}\sqrt{3}\left(\cos(3a)\operatorname{FresnelC}\left(\frac{\sqrt{2}\sqrt{3}\sqrt{b}x}{\sqrt{\pi}}\right) - \sin(3a)\operatorname{FresnelS}\left(\frac{\sqrt{2}\sqrt{3}\sqrt{b}x}{\sqrt{\pi}}\right)\right)}{24\sqrt{b}}$
risch	$\frac{e^{-3ia}\sqrt{\pi}\sqrt{3}\operatorname{erf}\left(\sqrt{3}\sqrt{ib}x\right)}{48\sqrt{ib}} + \frac{3e^{-ia}\sqrt{\pi}\operatorname{erf}\left(\sqrt{ib}x\right)}{16\sqrt{ib}} + \frac{e^{3ia}\sqrt{\pi}\operatorname{erf}\left(\sqrt{-3ib}x\right)}{16\sqrt{-3ib}} + \frac{3e^{ia}\sqrt{\pi}\operatorname{erf}\left(\sqrt{-ib}x\right)}{16\sqrt{-ib}}$

input `int(cos(b*x^2+a)^3,x,method=_RETURNVERBOSE)`

output $3/8*2^{(1/2)}*\pi^{(1/2)}/b^{(1/2)}*(\cos(a)*\operatorname{FresnelC}(b^{(1/2)}*2^{(1/2)}/\pi^{(1/2)}*x) - \sin(a)*\operatorname{FresnelS}(b^{(1/2)}*2^{(1/2)}/\pi^{(1/2)}*x)) + 1/24*2^{(1/2)}*\pi^{(1/2)}*3^{(1/2)}/b^{(1/2)}*(\cos(3*a)*\operatorname{FresnelC}(2^{(1/2)}/\pi^{(1/2)}*3^{(1/2)}*b^{(1/2)}*x) - \sin(3*a)*\operatorname{FresnelS}(2^{(1/2)}/\pi^{(1/2)}*3^{(1/2)}*b^{(1/2)}*x))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.79

$$\int \cos^3(a + bx^2) dx = \frac{\sqrt{6}\pi\sqrt{\frac{b}{\pi}}\cos(3a)C\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right) + 9\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)C\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right) - \sqrt{6}\pi\sqrt{\frac{b}{\pi}}S\left(\sqrt{6}x\sqrt{\frac{b}{\pi}}\right)\sin(3a) - 9\sqrt{2}\pi\sqrt{\frac{b}{\pi}}S\left(\sqrt{2}x\sqrt{\frac{b}{\pi}}\right)\sin(a)}{24b}$$

input `integrate(cos(b*x^2+a)^3,x, algorithm="fricas")`output `1/24*(sqrt(6)*pi*sqrt(b/pi)*cos(3*a)*fresnel_cos(sqrt(6)*x*sqrt(b/pi)) + 9*sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_cos(sqrt(2)*x*sqrt(b/pi)) - sqrt(6)*pi*sqrt(b/pi)*fresnel_sin(sqrt(6)*x*sqrt(b/pi))*sin(3*a) - 9*sqrt(2)*pi*sqrt(b/pi)*fresnel_sin(sqrt(2)*x*sqrt(b/pi))*sin(a))/b`**Sympy [A] (verification not implemented)**

Time = 0.48 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.84

$$\int \cos^3(a + bx^2) dx = \frac{3\sqrt{2}\sqrt{\pi}\left(-\sin(a)S\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(a)C\left(\frac{\sqrt{2}\sqrt{bx}}{\sqrt{\pi}}\right)\right)\sqrt{\frac{1}{b}}}{8} + \frac{\sqrt{6}\sqrt{\pi}\left(-\sin(3a)S\left(\frac{\sqrt{6}\sqrt{bx}}{\sqrt{\pi}}\right) + \cos(3a)C\left(\frac{\sqrt{6}\sqrt{bx}}{\sqrt{\pi}}\right)\right)\sqrt{\frac{1}{b}}}{24}$$

input `integrate(cos(b*x**2+a)**3,x)`output `3*sqrt(2)*sqrt(pi)*(-sin(a)*fresnels(sqrt(2)*sqrt(b)*x/sqrt(pi)) + cos(a)*fresnelc(sqrt(2)*sqrt(b)*x/sqrt(pi)))*sqrt(1/b)/8 + sqrt(6)*sqrt(pi)*(-sin(3*a)*fresnels(sqrt(6)*sqrt(b)*x/sqrt(pi)) + cos(3*a)*fresnelc(sqrt(6)*sqrt(b)*x/sqrt(pi)))*sqrt(1/b)/24`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.73

$$\int \cos^3(a + bx^2) dx = \frac{9^{\frac{1}{4}} \sqrt{2} \sqrt{\pi} \left((i-1) \cos(3a) + (i+1) \sin(3a) \right) \operatorname{erf}(\sqrt{3i} bx) + (-i+1) \cos(3a) - (i-1) \sin(3a)}{b^2}$$

input `integrate(cos(b*x^2+a)^3,x, algorithm="maxima")`

output `-1/96*(9^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(3*a) + (I + 1)*sin(3*a))*erf(sqrt(3*I*b)*x) + (-I + 1)*cos(3*a) - (I - 1)*sin(3*a))*erf(sqrt(-3*I*b)*x))*b^(3/2) - 9*sqrt(2)*sqrt(pi)*((-I - 1)*cos(a) - (I + 1)*sin(a))*erf(sqrt(I*b)*x) + ((I + 1)*cos(a) + (I - 1)*sin(a))*erf(sqrt(-I*b)*x))*b^(3/2)/b^2`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.21

$$\int \cos^3(a + bx^2) dx = -\frac{\sqrt{6}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{6}\sqrt{bx}\left(-\frac{ib}{|b|} + 1\right)\right) e^{3ia}}{48\sqrt{b}\left(-\frac{ib}{|b|} + 1\right)} - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{ia}}{16\left(-\frac{ib}{|b|} + 1\right)\sqrt{|b|}} - \frac{3\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}\right) e^{-ia}}{16\left(\frac{ib}{|b|} + 1\right)\sqrt{|b|}} - \frac{\sqrt{6}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{6}\sqrt{bx}\left(\frac{ib}{|b|} + 1\right)\right) e^{-3ia}}{48\sqrt{b}\left(\frac{ib}{|b|} + 1\right)}$$

input `integrate(cos(b*x^2+a)^3,x, algorithm="giac")`

output `-1/48*sqrt(6)*sqrt(pi)*erf(-1/2*sqrt(6)*sqrt(b)*x*(-I*b/abs(b) + 1))*e^(3*I*a)/(sqrt(b)*(-I*b/abs(b) + 1)) - 3/16*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(-I*b/abs(b) + 1)*sqrt(abs(b)))*e^(I*a)/((-I*b/abs(b) + 1)*sqrt(abs(b))) - 3/16*sqrt(2)*sqrt(pi)*erf(-1/2*sqrt(2)*x*(I*b/abs(b) + 1)*sqrt(abs(b)))*e^(-I*a)/((I*b/abs(b) + 1)*sqrt(abs(b))) - 1/48*sqrt(6)*sqrt(pi)*erf(-1/2*sqrt(6)*sqrt(b)*x*(I*b/abs(b) + 1))*e^(-3*I*a)/(sqrt(b)*(I*b/abs(b) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \cos^3(a + bx^2) dx = \int \cos(bx^2 + a)^3 dx$$

input `int(cos(a + b*x^2)^3,x)`

output `int(cos(a + b*x^2)^3, x)`

Reduce [F]

$$\int \cos^3(a + bx^2) dx = \int \cos(bx^2 + a)^3 dx$$

input `int(cos(b*x^2+a)^3,x)`

output `int(cos(a + b*x**2)**3,x)`

3.19 $\int \frac{\cos^3(a+bx^2)}{x} dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [C] (warning: unable to verify)	174
Fricas [A] (verification not implemented)	174
Sympy [F]	175
Maxima [C] (verification not implemented)	175
Giac [A] (verification not implemented)	176
Mupad [F(-1)]	176
Reduce [F]	176

Optimal result

Integrand size = 14, antiderivative size = 55

$$\int \frac{\cos^3(a+bx^2)}{x} dx = \frac{3}{8} \cos(a) \operatorname{CosIntegral}(bx^2) + \frac{1}{8} \cos(3a) \operatorname{CosIntegral}(3bx^2) - \frac{3}{8} \sin(a) \operatorname{Si}(bx^2) - \frac{1}{8} \sin(3a) \operatorname{Si}(3bx^2)$$

output

```
3/8*cos(a)*Ci(b*x^2)+1/8*cos(3*a)*Ci(3*b*x^2)-3/8*sin(a)*Si(b*x^2)-1/8*sin(3*a)*Si(3*b*x^2)
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.91

$$\int \frac{\cos^3(a+bx^2)}{x} dx = \frac{1}{8} (3 \cos(a) \operatorname{CosIntegral}(bx^2) + \cos(3a) \operatorname{CosIntegral}(3bx^2) - 3 \sin(a) \operatorname{Si}(bx^2) - \sin(3a) \operatorname{Si}(3bx^2))$$

input

```
Integrate[Cos[a + b*x^2]^3/x,x]
```

output

```
(3*Cos[a]*CosIntegral[b*x^2] + Cos[3*a]*CosIntegral[3*b*x^2] - 3*Sin[a]*SinIntegral[b*x^2] - Sin[3*a]*SinIntegral[3*b*x^2])/8
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(a + bx^2)}{x} dx$$

$$\downarrow \text{3885}$$

$$\int \left(\frac{3 \cos(a + bx^2)}{4x} + \frac{\cos(3a + 3bx^2)}{4x} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3}{8} \cos(a) \text{CosIntegral}(bx^2) + \frac{1}{8} \cos(3a) \text{CosIntegral}(3bx^2) - \frac{3}{8} \sin(a) \text{Si}(bx^2) - \frac{1}{8} \sin(3a) \text{Si}(3bx^2)$$

input

```
Int[Cos[a + b*x^2]^3/x,x]
```

output

```
(3*Cos[a]*CosIntegral[b*x^2])/8 + (Cos[3*a]*CosIntegral[3*b*x^2])/8 - (3*Sin[a]*SinIntegral[b*x^2])/8 - (Sin[3*a]*SinIntegral[3*b*x^2])/8
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_.)]*(b_.))^p_]*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.39 (sec) , antiderivative size = 125, normalized size of antiderivative = 2.27

method	result
risch	$\frac{i\pi \operatorname{csgn}(bx^2)e^{-3ia}}{16} - \frac{i \operatorname{Si}(3bx^2)e^{-3ia}}{8} - \frac{\operatorname{expIntegral}_1(-3ibx^2)e^{-3ia}}{16} + \frac{3i\pi \operatorname{csgn}(bx^2)e^{-ia}}{16} - \frac{3ie^{-ia} \operatorname{Si}(bx^2)}{8} - \frac{3e^{-ia} \operatorname{Si}(bx^2)}{8}$

input `int(cos(b*x^2+a)^3/x,x,method=_RETURNVERBOSE)`

output `1/16*I*Pi*csgn(b*x^2)*exp(-3*I*a)-1/8*I*Si(3*b*x^2)*exp(-3*I*a)-1/16*Ei(1,-3*I*b*x^2)*exp(-3*I*a)+3/16*I*Pi*csgn(b*x^2)*exp(-I*a)-3/8*I*exp(-I*a)*Si(b*x^2)-3/16*exp(-I*a)*Ei(1,-I*b*x^2)-3/16*exp(I*a)*Ei(1,-I*b*x^2)-1/16*exp(3*I*a)*Ei(1,-3*I*b*x^2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.85

$$\int \frac{\cos^3(a + bx^2)}{x} dx = \frac{1}{8} \cos(3a) \operatorname{Ci}(3bx^2) + \frac{3}{8} \cos(a) \operatorname{Ci}(bx^2) - \frac{1}{8} \sin(3a) \operatorname{Si}(3bx^2) - \frac{3}{8} \sin(a) \operatorname{Si}(bx^2)$$

input `integrate(cos(b*x^2+a)^3/x,x, algorithm="fricas")`

output `1/8*cos(3*a)*cos_integral(3*b*x^2) + 3/8*cos(a)*cos_integral(b*x^2) - 1/8*
sin(3*a)*sin_integral(3*b*x^2) - 3/8*sin(a)*sin_integral(b*x^2)`

Sympy [F]

$$\int \frac{\cos^3(a + bx^2)}{x} dx = \int \frac{\cos^3(a + bx^2)}{x} dx$$

input `integrate(cos(b*x**2+a)**3/x,x)`

output `Integral(cos(a + b*x**2)**3/x, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.62

$$\begin{aligned} \int \frac{\cos^3(a + bx^2)}{x} dx &= \frac{1}{16} (\operatorname{Ei}(3i bx^2) + \operatorname{Ei}(-3i bx^2)) \cos(3a) \\ &+ \frac{3}{16} (\operatorname{Ei}(i bx^2) + \operatorname{Ei}(-i bx^2)) \cos(a) \\ &+ \frac{1}{16} (i \operatorname{Ei}(3i bx^2) - i \operatorname{Ei}(-3i bx^2)) \sin(3a) \\ &- \frac{3}{16} (-i \operatorname{Ei}(i bx^2) + i \operatorname{Ei}(-i bx^2)) \sin(a) \end{aligned}$$

input `integrate(cos(b*x^2+a)^3/x,x, algorithm="maxima")`

output `1/16*(Ei(3*I*b*x^2) + Ei(-3*I*b*x^2))*cos(3*a) + 3/16*(Ei(I*b*x^2) + Ei(-I
*b*x^2))*cos(a) + 1/16*(I*Ei(3*I*b*x^2) - I*Ei(-3*I*b*x^2))*sin(3*a) - 3/1
6*(-I*Ei(I*b*x^2) + I*Ei(-I*b*x^2))*sin(a)`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.85

$$\int \frac{\cos^3(a + bx^2)}{x} dx = \frac{1}{8} \cos(3a) \operatorname{Ci}(3bx^2) + \frac{3}{8} \cos(a) \operatorname{Ci}(bx^2) - \frac{3}{8} \sin(a) \operatorname{Si}(bx^2) + \frac{1}{8} \sin(3a) \operatorname{Si}(-3bx^2)$$

input `integrate(cos(b*x^2+a)^3/x,x, algorithm="giac")`

output `1/8*cos(3*a)*cos_integral(3*b*x^2) + 3/8*cos(a)*cos_integral(b*x^2) - 3/8*sin(a)*sin_integral(b*x^2) + 1/8*sin(3*a)*sin_integral(-3*b*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(a + bx^2)}{x} dx = \int \frac{\cos(bx^2 + a)^3}{x} dx$$

input `int(cos(a + b*x^2)^3/x,x)`

output `int(cos(a + b*x^2)^3/x, x)`

Reduce [F]

$$\int \frac{\cos^3(a + bx^2)}{x} dx = \int \frac{\cos(bx^2 + a)^3}{x} dx$$

input `int(cos(b*x^2+a)^3/x,x)`

output `int(cos(a + b*x**2)**3/x,x)`

3.20 $\int \frac{\cos^3(a+bx^2)}{x^2} dx$

Optimal result	177
Mathematica [A] (verified)	178
Rubi [A] (verified)	178
Maple [A] (verified)	180
Fricas [A] (verification not implemented)	180
Sympy [F]	181
Maxima [C] (verification not implemented)	181
Giac [F]	182
Mupad [F(-1)]	182
Reduce [F]	182

Optimal result

Integrand size = 14, antiderivative size = 168

$$\int \frac{\cos^3(a+bx^2)}{x^2} dx = -\frac{\cos^3(a+bx^2)}{x} - \frac{3}{2}\sqrt{b}\sqrt{\frac{\pi}{2}}\cos(a)\text{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) - \frac{1}{2}\sqrt{b}\sqrt{\frac{3\pi}{2}}\cos(3a)\text{FresnelS}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right) - \frac{3}{2}\sqrt{b}\sqrt{\frac{\pi}{2}}\text{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right)\sin(a) - \frac{1}{2}\sqrt{b}\sqrt{\frac{3\pi}{2}}\text{FresnelC}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)\sin(3a)$$

output

```
-cos(b*x^2+a)^3/x-3/4*b^(1/2)*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)-1/4*b^(1/2)*6^(1/2)*Pi^(1/2)*cos(3*a)*FresnelS(b^(1/2)*6^(1/2)/Pi^(1/2)*x)-3/4*b^(1/2)*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)*sin(a)-1/4*b^(1/2)*6^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*6^(1/2)/Pi^(1/2)*x)*sin(3*a)
```

Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.99

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \frac{3 \cos(a + bx^2) + \cos(3(a + bx^2)) + 3\sqrt{b}\sqrt{2\pi}x \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}x\right) + \sqrt{b}\sqrt{6\pi}x \cos(3a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{6}{\pi}}x\right)}{4x}$$

input

```
Integrate[Cos[a + b*x^2]^3/x^2,x]
```

output

```
-1/4*(3*Cos[a + b*x^2] + Cos[3*(a + b*x^2)] + 3*Sqrt[b]*Sqrt[2*Pi]*x*Cos[a]
]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x] + Sqrt[b]*Sqrt[6*Pi]*x*Cos[3*a]*FresnelS[
Sqrt[b]*Sqrt[6/Pi]*x] + 3*Sqrt[b]*Sqrt[2*Pi]*x*FresnelC[Sqrt[b]*Sqrt[2/Pi]
*x]*Sin[a] + Sqrt[b]*Sqrt[6*Pi]*x*FresnelC[Sqrt[b]*Sqrt[6/Pi]*x]*Sin[3*a])
/x
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3875, 5085, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^3(a + bx^2)}{x^2} dx \\ & \quad \downarrow \text{3875} \\ & -6b \int \cos^2(bx^2 + a) \sin(bx^2 + a) dx - \frac{\cos^3(a + bx^2)}{x} \\ & \quad \downarrow \text{5085} \\ & -6b \int \left(\frac{1}{4} \sin(bx^2 + a) + \frac{1}{4} \sin(3bx^2 + 3a) \right) dx - \frac{\cos^3(a + bx^2)}{x} \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$-6b \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} x \right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{6}} \sin(3a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{6}{\pi}} x \right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} x \right)}{4\sqrt{b}} + \frac{\cos^3(a + bx^2)}{x} \right)$$

input `Int[Cos[a + b*x^2]^3/x^2,x]`

output `-(Cos[a + b*x^2]^3/x) - 6*b*((Sqrt[Pi/2]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x])/(4*Sqrt[b]) + (Sqrt[Pi/6]*Cos[3*a]*FresnelS[Sqrt[b]*Sqrt[6/Pi]*x])/(4*Sqrt[b]) + (Sqrt[Pi/2]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x]*Sin[a])/(4*Sqrt[b]) + (Sqrt[Pi/6]*FresnelC[Sqrt[b]*Sqrt[6/Pi]*x]*Sin[3*a])/(4*Sqrt[b]))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3875 `Int[Cos[(a_.) + (b_.)*(x_)^(n_)]^(p_)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(Cos[a + b*x^n]^p/(m + 1)), x] + Simp[b*n*(p/(m + 1)) Int[Cos[a + b*x^n]^(p - 1)*Sin[a + b*x^n], x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 1] && EqQ[m + n, 0] && NeQ[n, 1] && IntegerQ[n]`

rule 5085 `Int[Cos[w_]^(q_.)*Sin[v_]^(p_.), x_Symbol] := Int[ExpandTrigReduce[Sin[v]^p *Cos[w]^q, x], x] /; IGtQ[p, 0] && IGtQ[q, 0] && ((PolynomialQ[v, x] && PolynomialQ[w, x]) || (BinomialQ[{v, w}, x] && IndependentQ[Cancel[v/w], x]))`

Maple [A] (verified)

Time = 1.59 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.76

method	result
default	$-\frac{3 \cos(bx^2+a)}{4x} - \frac{3\sqrt{b}\sqrt{2}\sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x}{\sqrt{\pi}}\right) \right)}{4} - \frac{\cos(3bx^2+3a)}{4x} - \frac{\sqrt{b}\sqrt{2}\sqrt{\pi}\sqrt{3} \left(\cos(3a) \operatorname{FresnelS}\left(\frac{\sqrt{3}\sqrt{b}x}{\sqrt{\pi}}\right) + \sin(3a) \operatorname{FresnelC}\left(\frac{\sqrt{3}\sqrt{b}x}{\sqrt{\pi}}\right) \right)}{4}$
risch	$-\frac{ie^{-3ia}b\sqrt{\pi}\sqrt{3} \operatorname{erf}\left(\sqrt{3}\sqrt{ib}x\right)}{8\sqrt{ib}} - \frac{3ie^{-ia}b\sqrt{\pi} \operatorname{erf}\left(\sqrt{ib}x\right)}{8\sqrt{ib}} + \frac{3ie^{ia}b\sqrt{\pi} \operatorname{erf}\left(\sqrt{-ib}x\right)}{8\sqrt{-ib}} + \frac{3ie^{3ia}b\sqrt{\pi} \operatorname{erf}\left(\sqrt{-3ib}x\right)}{8\sqrt{-3ib}} - \frac{3 \cos(bx^2+a)}{4x}$

input

```
int(cos(b*x^2+a)^3/x^2,x,method=_RETURNVERBOSE)
```

output

```
-3/4*cos(b*x^2+a)/x-3/4*b^(1/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x)-1/4*cos(3*b*x^2+3*a)/x-1/4*b^(1/2)*2^(1/2)*Pi^(1/2)*3^(1/2)*(cos(3*a)*FresnelS(2^(1/2)/Pi^(1/2)*3^(1/2)*b^(1/2)*x)+sin(3*a)*FresnelC(2^(1/2)/Pi^(1/2)*3^(1/2)*b^(1/2)*x)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.81

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \frac{\sqrt{6}\pi x \sqrt{\frac{b}{\pi}} \cos(3a) S\left(\sqrt{6}x \sqrt{\frac{b}{\pi}}\right) + 3\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} \cos(a) S\left(\sqrt{2}x \sqrt{\frac{b}{\pi}}\right) + \sqrt{6}\pi x \sqrt{\frac{b}{\pi}} C\left(\sqrt{6}x \sqrt{\frac{b}{\pi}}\right) \sin(3a) - \sqrt{2}\pi x \sqrt{\frac{b}{\pi}} C\left(\sqrt{2}x \sqrt{\frac{b}{\pi}}\right) \sin(a)}{4x}$$

input

```
integrate(cos(b*x^2+a)^3/x^2,x, algorithm="fricas")
```

output

```
-1/4*(sqrt(6)*pi*x*sqrt(b/pi)*cos(3*a)*fresnel_sin(sqrt(6)*x*sqrt(b/pi)) + 3*sqrt(2)*pi*x*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x*sqrt(b/pi)) + sqrt(6)*pi*x*sqrt(b/pi)*fresnel_cos(sqrt(6)*x*sqrt(b/pi))*sin(3*a) + 3*sqrt(2)*pi*x*sqrt(b/pi)*fresnel_cos(sqrt(2)*x*sqrt(b/pi))*sin(a) + 4*cos(b*x^2 + a)^3)/x
```

Sympy [F]

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \int \frac{\cos^3(a + bx^2)}{x^2} dx$$

input `integrate(cos(b*x**2+a)**3/x**2,x)`

output `Integral(cos(a + b*x**2)**3/x**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.90

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \frac{\sqrt{3}\sqrt{bx^2}((-i+1)\sqrt{2}\Gamma(-\frac{1}{2}, 3ibx^2) + (i-1)\sqrt{2}\Gamma(-\frac{1}{2}, -3ibx^2))\cos(3a) + ((i-1)\sqrt{2}\Gamma(-\frac{1}{2}, 3ibx^2) + (i+1)\sqrt{2}\Gamma(-\frac{1}{2}, -3ibx^2))\sin(3a)}{x}$$

input `integrate(cos(b*x^2+a)^3/x^2,x, algorithm="maxima")`

output `1/32*(sqrt(3)*sqrt(b*x^2)*((-I + 1)*sqrt(2)*gamma(-1/2, 3*I*b*x^2) + (I - 1)*sqrt(2)*gamma(-1/2, -3*I*b*x^2))*cos(3*a) + ((I - 1)*sqrt(2)*gamma(-1/2, 3*I*b*x^2) - (I + 1)*sqrt(2)*gamma(-1/2, -3*I*b*x^2))*sin(3*a)) - 3*sqrt(b*x^2)*(((I + 1)*sqrt(2)*gamma(-1/2, I*b*x^2) - (I - 1)*sqrt(2)*gamma(-1/2, -I*b*x^2))*cos(a) + (-I - 1)*sqrt(2)*gamma(-1/2, I*b*x^2) + (I + 1)*sqrt(2)*gamma(-1/2, -I*b*x^2))*sin(a))/x`

Giac [F]

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)^3}{x^2} dx$$

input `integrate(cos(b*x^2+a)^3/x^2,x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)^3/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)^3}{x^2} dx$$

input `int(cos(a + b*x^2)^3/x^2,x)`

output `int(cos(a + b*x^2)^3/x^2, x)`

Reduce [F]

$$\int \frac{\cos^3(a + bx^2)}{x^2} dx = \int \frac{\cos(bx^2 + a)^3}{x^2} dx$$

input `int(cos(b*x^2+a)^3/x^2,x)`

output `int(cos(a + b*x**2)**3/x**2,x)`

3.21 $\int \frac{\cos^3(a+bx^2)}{x^3} dx$

Optimal result	183
Mathematica [A] (verified)	183
Rubi [A] (verified)	184
Maple [C] (warning: unable to verify)	185
Fricas [A] (verification not implemented)	185
Sympy [F]	186
Maxima [C] (verification not implemented)	186
Giac [B] (verification not implemented)	187
Mupad [F(-1)]	187
Reduce [F]	188

Optimal result

Integrand size = 14, antiderivative size = 91

$$\int \frac{\cos^3(a+bx^2)}{x^3} dx = -\frac{3 \cos(a+bx^2)}{8x^2} - \frac{\cos(3(a+bx^2))}{8x^2} - \frac{3}{8}b \operatorname{CosIntegral}(bx^2) \sin(a) - \frac{3}{8}b \operatorname{CosIntegral}(3bx^2) \sin(3a) - \frac{3}{8}b \cos(a) \operatorname{Si}(bx^2) - \frac{3}{8}b \cos(3a) \operatorname{Si}(3bx^2)$$

```
output -3/8*cos(b*x^2+a)/x^2-1/8*cos(3*b*x^2+3*a)/x^2-3/8*b*Ci(b*x^2)*sin(a)-3/8*b*Ci(3*b*x^2)*sin(3*a)-3/8*b*cos(a)*Si(b*x^2)-3/8*b*cos(3*a)*Si(3*b*x^2)
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.99

$$\int \frac{\cos^3(a+bx^2)}{x^3} dx = \frac{-3 \cos(a+bx^2) + \cos(3(a+bx^2)) + 3bx^2 \operatorname{CosIntegral}(bx^2) \sin(a) + 3bx^2 \operatorname{CosIntegral}(3bx^2) \sin(3a) - 3 \cos(a) \operatorname{Si}(bx^2) - 3 \cos(3a) \operatorname{Si}(3bx^2)}{8x^2}$$

```
input Integrate[Cos[a + b*x^2]^3/x^3,x]
```


output

```
-1/8*(3*Cos[a + b*x^2] + Cos[3*(a + b*x^2)] + 3*b*x^2*CosIntegral[b*x^2]*Sin[a] + 3*b*x^2*CosIntegral[3*b*x^2]*Sin[3*a] + 3*b*x^2*Cos[a]*SinIntegral[b*x^2] + 3*b*x^2*Cos[3*a]*SinIntegral[3*b*x^2])/x^2
```

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx$$

↓ 3885

$$\int \left(\frac{3 \cos(a + bx^2)}{4x^3} + \frac{\cos(3a + 3bx^2)}{4x^3} \right) dx$$

↓ 2009

$$-\frac{3}{8}b \sin(a) \text{CosIntegral}(bx^2) - \frac{3}{8}b \sin(3a) \text{CosIntegral}(3bx^2) - \frac{3}{8}b \cos(a) \text{Si}(bx^2) - \frac{3}{8}b \cos(3a) \text{Si}(3bx^2) - \frac{3 \cos(a + bx^2)}{8x^2} - \frac{\cos(3(a + bx^2))}{8x^2}$$

input

```
Int[Cos[a + b*x^2]^3/x^3,x]
```

output

```
(-3*Cos[a + b*x^2])/(8*x^2) - Cos[3*(a + b*x^2)]/(8*x^2) - (3*b*CosIntegral[b*x^2]*Sin[a])/8 - (3*b*CosIntegral[3*b*x^2]*Sin[3*a])/8 - (3*b*Cos[a]*SinIntegral[b*x^2])/8 - (3*b*Cos[3*a]*SinIntegral[3*b*x^2])/8
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.30 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.03

method	result
risch	$-\frac{3ie^{3ia}b \operatorname{ExpIntegral}_1(-3ibx^2)x^2 - 3i \operatorname{ExpIntegral}_1(-3ibx^2)e^{-3ia}bx^2 + 3ie^{ia}b \operatorname{ExpIntegral}_1(-ibx^2)x^2 - 3i \operatorname{ExpIntegral}_1(-ibx^2)}$

input `int(cos(b*x^2+a)^3/x^3,x,method=_RETURNVERBOSE)`

output
$$-1/16*(3*I*\exp(3*I*a)*b*Ei(1,-3*I*b*x^2)*x^2-3*I*Ei(1,-3*I*b*x^2)*\exp(-3*I*a)*b*x^2+3*I*\exp(I*a)*b*Ei(1,-I*b*x^2)*x^2-3*I*Ei(1,-I*b*x^2)*\exp(-I*a)*b*x^2-3*Pi*csgn(b*x^2)*\exp(-3*I*a)*b*x^2-3*Pi*csgn(b*x^2)*\exp(-I*a)*b*x^2+6*Si(3*b*x^2)*\exp(-3*I*a)*b*x^2+6*Si(b*x^2)*\exp(-I*a)*b*x^2+6*\cos(b*x^2+a)+2*\cos(3*b*x^2+3*a))/x^2$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.88

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx = -\frac{3bx^2 \operatorname{Ci}(3bx^2) \sin(3a) + 3bx^2 \operatorname{Ci}(bx^2) \sin(a) + 3bx^2 \cos(3a) \operatorname{Si}(3bx^2) + 3bx^2 \cos(a) \operatorname{Si}(bx^2) + 4 \cos(3a) \operatorname{Si}(3bx^2) + 4 \cos(a) \operatorname{Si}(bx^2)}{8x^2}$$

input `integrate(cos(b*x^2+a)^3/x^3,x, algorithm="fricas")`

output

```
-1/8*(3*b*x^2*cos_integral(3*b*x^2)*sin(3*a) + 3*b*x^2*cos_integral(b*x^2)*sin(a) + 3*b*x^2*cos(3*a)*sin_integral(3*b*x^2) + 3*b*x^2*cos(a)*sin_integral(b*x^2) + 4*cos(b*x^2 + a)^3)/x^2
```

Sympy [F]

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx = \int \frac{\cos^3(a + bx^2)}{x^3} dx$$

input

```
integrate(cos(b*x**2+a)**3/x**3,x)
```

output

```
Integral(cos(a + b*x**2)**3/x**3, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.08

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx = \frac{3}{16} ((-i\Gamma(-1, 3i bx^2) + i\Gamma(-1, -3i bx^2)) \cos(3a) + (-i\Gamma(-1, i bx^2) + i\Gamma(-1, -i bx^2)) \cos(a) - (\Gamma(-1, 3i bx^2) + \Gamma(-1, -3i bx^2)) \sin(3a) - (\Gamma(-1, i bx^2) + \Gamma(-1, -i bx^2)) \sin(a)) * b$$

input

```
integrate(cos(b*x^2+a)^3/x^3,x, algorithm="maxima")
```

output

```
3/16*((-I*gamma(-1, 3*I*b*x^2) + I*gamma(-1, -3*I*b*x^2))*cos(3*a) + (-I*gamma(-1, I*b*x^2) + I*gamma(-1, -I*b*x^2))*cos(a) - (gamma(-1, 3*I*b*x^2) + gamma(-1, -3*I*b*x^2))*sin(3*a) - (gamma(-1, I*b*x^2) + gamma(-1, -I*b*x^2))*sin(a))*b
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(80) = 160$.

Time = 0.36 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.03

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx = \frac{3(bx^2 + a)b^2 \operatorname{Ci}(3bx^2) \sin(3a) - 3ab^2 \operatorname{Ci}(3bx^2) \sin(3a) + 3(bx^2 + a)b^2 \operatorname{Ci}(bx^2) \sin(a) - 3ab^2 \operatorname{Ci}(bx^2) \sin(a)}{b^2 x^2}$$

input `integrate(cos(b*x^2+a)^3/x^3,x, algorithm="giac")`

output `-1/8*(3*(b*x^2 + a)*b^2*cos_integral(3*b*x^2)*sin(3*a) - 3*a*b^2*cos_integral(3*b*x^2)*sin(3*a) + 3*(b*x^2 + a)*b^2*cos_integral(b*x^2)*sin(a) - 3*a*b^2*cos_integral(b*x^2)*sin(a) + 3*(b*x^2 + a)*b^2*cos(a)*sin_integral(b*x^2) - 3*a*b^2*cos(a)*sin_integral(b*x^2) - 3*(b*x^2 + a)*b^2*cos(3*a)*sin_integral(-3*b*x^2) + 3*a*b^2*cos(3*a)*sin_integral(-3*b*x^2) + b^2*cos(3*b*x^2 + 3*a) + 3*b^2*cos(b*x^2 + a))/(b^2*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx = \int \frac{\cos(bx^2 + a)^3}{x^3} dx$$

input `int(cos(a + b*x^2)^3/x^3,x)`

output `int(cos(a + b*x^2)^3/x^3, x)`

Reduce [F]

$$\int \frac{\cos^3(a + bx^2)}{x^3} dx = \int \frac{\cos(bx^2 + a)^3}{x^3} dx$$

input `int(cos(b*x^2+a)^3/x^3,x)`

output `int(cos(a + b*x**2)**3/x**3,x)`

3.22 $\int x \cos^7(a + bx^2) dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (warning: unable to verify)	190
Maple [A] (verified)	191
Fricas [A] (verification not implemented)	192
Sympy [A] (verification not implemented)	192
Maxima [A] (verification not implemented)	193
Giac [A] (verification not implemented)	193
Mupad [B] (verification not implemented)	194
Reduce [B] (verification not implemented)	194

Optimal result

Integrand size = 12, antiderivative size = 67

$$\int x \cos^7(a + bx^2) dx = \frac{\sin(a + bx^2)}{2b} - \frac{\sin^3(a + bx^2)}{2b} + \frac{3 \sin^5(a + bx^2)}{10b} - \frac{\sin^7(a + bx^2)}{14b}$$

output

$$1/2*\sin(b*x^2+a)/b-1/2*\sin(b*x^2+a)^3/b+3/10*\sin(b*x^2+a)^5/b-1/14*\sin(b*x^2+a)^7/b$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.81

$$\begin{aligned} &\int x \cos^7(a + bx^2) dx \\ &= \frac{35 \sin(a + bx^2) - 35 \sin^3(a + bx^2) + 21 \sin^5(a + bx^2) - 5 \sin^7(a + bx^2)}{70b} \end{aligned}$$

input

$$\text{Integrate}[x*\text{Cos}[a + b*x^2]^7,x]$$

output

$$(35*\text{Sin}[a + b*x^2] - 35*\text{Sin}[a + b*x^2]^3 + 21*\text{Sin}[a + b*x^2]^5 - 5*\text{Sin}[a + b*x^2]^7)/(70*b)$$

Rubi [A] (warning: unable to verify)

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.55, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3861, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos^7(a + bx^2) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{1}{2} \int \cos^7(bx^2 + a) dx^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \sin\left(bx^2 + a + \frac{\pi}{2}\right)^7 dx^2 \\
 & \quad \downarrow \text{3113} \\
 & -\frac{\int (-x^{12} + 3x^8 - 3x^4 + 1) d(-\sin(bx^2 + a))}{2b} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{-\sin(a + bx^2) - \frac{x^{14}}{7} + \frac{3x^{10}}{5} - x^6}{2b}
 \end{aligned}$$

input `Int[x*Cos[a + b*x^2]^7,x]`

output `-1/2*(-x^6 + (3*x^10)/5 - x^14/7 - Sin[a + b*x^2])/b`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3113 `Int[sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[-d^(-1) Subst[Int[Exp and[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 7.72 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.75

method	result	size
derivativdivides	$\frac{\left(\frac{16}{5} + \cos(bx^2+a)^6 + \frac{6 \cos(bx^2+a)^4}{5} + \frac{8 \cos(bx^2+a)^2}{5}\right) \sin(bx^2+a)}{14b}$	50
default	$\frac{\left(\frac{16}{5} + \cos(bx^2+a)^6 + \frac{6 \cos(bx^2+a)^4}{5} + \frac{8 \cos(bx^2+a)^2}{5}\right) \sin(bx^2+a)}{14b}$	50
parallelrisc	$\frac{1225 \sin(bx^2+a) + 5 \sin(7bx^2+7a) + 49 \sin(5bx^2+5a) + 245 \sin(3bx^2+3a)}{4480b}$	56
risc	$\frac{35 \sin(bx^2+a)}{128b} + \frac{\sin(7bx^2+7a)}{896b} + \frac{7 \sin(5bx^2+5a)}{640b} + \frac{7 \sin(3bx^2+3a)}{128b}$	63
orering	Expression too large to display	1544

input `int(x*cos(b*x^2+a)^7,x,method=_RETURNVERBOSE)`

output $1/14/b*(16/5+\cos(b*x^2+a)^6+6/5*\cos(b*x^2+a)^4+8/5*\cos(b*x^2+a)^2)*\sin(b*x^2+a)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.76

$$\int x \cos^7(a + bx^2) dx$$

$$= \frac{(5 \cos(bx^2 + a)^6 + 6 \cos(bx^2 + a)^4 + 8 \cos(bx^2 + a)^2 + 16) \sin(bx^2 + a)}{70b}$$

input `integrate(x*cos(b*x^2+a)^7,x, algorithm="fricas")`

output $1/70*(5*\cos(b*x^2 + a)^6 + 6*\cos(b*x^2 + a)^4 + 8*\cos(b*x^2 + a)^2 + 16)*\sin(b*x^2 + a)/b$

Sympy [A] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.40

$$\int x \cos^7(a + bx^2) dx$$

$$= \begin{cases} \frac{8 \sin^7(a+bx^2)}{35b} + \frac{4 \sin^5(a+bx^2) \cos^2(a+bx^2)}{5b} + \frac{\sin^3(a+bx^2) \cos^4(a+bx^2)}{b} + \frac{\sin(a+bx^2) \cos^6(a+bx^2)}{2b} & \text{for } b \neq 0 \\ \frac{x^2 \cos^7(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*cos(b*x**2+a)**7,x)`

output `Piecewise((8*sin(a + b*x**2)**7/(35*b) + 4*sin(a + b*x**2)**5*cos(a + b*x**2)**2/(5*b) + sin(a + b*x**2)**3*cos(a + b*x**2)**4/b + sin(a + b*x**2)*cos(a + b*x**2)**6/(2*b), Ne(b, 0)), (x**2*cos(a)**7/2, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.82

$$\int x \cos^7(a + bx^2) dx$$

$$= \frac{5 \sin(7bx^2 + 7a) + 49 \sin(5bx^2 + 5a) + 245 \sin(3bx^2 + 3a) + 1225 \sin(bx^2 + a)}{4480b}$$

input `integrate(x*cos(b*x^2+a)^7,x, algorithm="maxima")`output `1/4480*(5*sin(7*b*x^2 + 7*a) + 49*sin(5*b*x^2 + 5*a) + 245*sin(3*b*x^2 + 3*a) + 1225*sin(b*x^2 + a))/b`**Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.78

$$\int x \cos^7(a + bx^2) dx$$

$$= -\frac{5 \sin(bx^2 + a)^7 - 21 \sin(bx^2 + a)^5 + 35 \sin(bx^2 + a)^3 - 35 \sin(bx^2 + a)}{70b}$$

input `integrate(x*cos(b*x^2+a)^7,x, algorithm="giac")`output `-1/70*(5*sin(b*x^2 + a)^7 - 21*sin(b*x^2 + a)^5 + 35*sin(b*x^2 + a)^3 - 35*sin(b*x^2 + a))/b`

Mupad [B] (verification not implemented)

Time = 0.79 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.82

$$\int x \cos^7(a + bx^2) dx$$

$$= \frac{245 \sin(3bx^2 + 3a) + 49 \sin(5bx^2 + 5a) + 5 \sin(7bx^2 + 7a) + 1225 \sin(bx^2 + a)}{4480b}$$

input `int(x*cos(a + b*x^2)^7,x)`output `(245*sin(3*a + 3*b*x^2) + 49*sin(5*a + 5*b*x^2) + 5*sin(7*a + 7*b*x^2) + 1225*sin(a + b*x^2))/(4480*b)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.76

$$\int x \cos^7(a + bx^2) dx$$

$$= \frac{\sin(bx^2 + a) \left(-5 \sin(bx^2 + a)^6 + 21 \sin(bx^2 + a)^4 - 35 \sin(bx^2 + a)^2 + 35 \right)}{70b}$$

input `int(x*cos(b*x^2+a)^7,x)`output `(sin(a + b*x**2)*(-5*sin(a + b*x**2)**6 + 21*sin(a + b*x**2)**4 - 35*sin(a + b*x**2)**2 + 35))/(70*b)`

3.23 $\int x^{5/2} \cos(a + bx^2) dx$

Optimal result	195
Mathematica [A] (verified)	195
Rubi [A] (verified)	196
Maple [C] (verified)	197
Fricas [A] (verification not implemented)	198
Sympy [F]	198
Maxima [F(-2)]	199
Giac [F]	199
Mupad [F(-1)]	199
Reduce [F]	200

Optimal result

Integrand size = 14, antiderivative size = 111

$$\int x^{5/2} \cos(a + bx^2) dx = -\frac{3ie^{ia}x^{3/2}\Gamma(\frac{3}{4}, -ibx^2)}{16b(-ibx^2)^{3/4}} + \frac{3ie^{-ia}x^{3/2}\Gamma(\frac{3}{4}, ibx^2)}{16b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(a + bx^2)}{2b}$$

output

```
-3/16*I*exp(I*a)*x^(3/2)*GAMMA(3/4,-I*b*x^2)/b/(-I*b*x^2)^(3/4)+3/16*I*x^(3/2)*GAMMA(3/4,I*b*x^2)/b/exp(I*a)/(I*b*x^2)^(3/4)+1/2*x^(3/2)*sin(b*x^2+a)/b
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.02

$$\int x^{5/2} \cos(a + bx^2) dx = \frac{bx^{11/2} \left(3(ibx^2)^{3/4} \Gamma(\frac{3}{4}, -ibx^2) (-i \cos(a) + \sin(a)) + 3(-ibx^2)^{3/4} \Gamma(\frac{3}{4}, ibx^2) (i \cos(a) + \sin(a)) \right)}{16(b^2x^4)^{7/4}}$$

input

```
Integrate[x^(5/2)*Cos[a + b*x^2],x]
```

output

$$\frac{(b*x^{11/2}*(3*(I*b*x^2)^{3/4}*Gamma[3/4, (-I)*b*x^2]*((-I)*Cos[a] + Sin[a]) + 3*((-I)*b*x^2)^{3/4}*Gamma[3/4, I*b*x^2]*(I*Cos[a] + Sin[a]) + 8*(b^2*x^4)^{3/4}*Sin[a + b*x^2]))}{(16*(b^2*x^4)^{7/4})}$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3867, 3870, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{5/2} \cos(a + bx^2) dx \\ & \quad \downarrow \text{3867} \\ & \frac{x^{3/2} \sin(a + bx^2)}{2b} - \frac{3 \int \sqrt{x} \sin(bx^2 + a) dx}{4b} \\ & \quad \downarrow \text{3870} \\ & \frac{x^{3/2} \sin(a + bx^2)}{2b} - \frac{3 \left(\frac{1}{2} i \int e^{-ibx^2 - ia} \sqrt{x} dx - \frac{1}{2} i \int e^{ibx^2 + ia} \sqrt{x} dx \right)}{4b} \\ & \quad \downarrow \text{2648} \\ & \frac{x^{3/2} \sin(a + bx^2)}{2b} - \frac{3 \left(\frac{ie^{ia} x^{3/2} \Gamma(\frac{3}{4}, -ibx^2)}{4(-ibx^2)^{3/4}} - \frac{ie^{-ia} x^{3/2} \Gamma(\frac{3}{4}, ibx^2)}{4(ibx^2)^{3/4}} \right)}{4b} \end{aligned}$$

input

$$\text{Int}[x^{(5/2)}*\text{Cos}[a + b*x^2], x]$$

output

$$\frac{(-3*((I/4)*E^{(I*a)}*x^{(3/2)}*Gamma[3/4, (-I)*b*x^2])/((-I)*b*x^2)^{(3/4)} - ((I/4)*x^{(3/2)}*Gamma[3/4, I*b*x^2])/(E^{(I*a)}*(I*b*x^2)^{(3/4))})}{(4*b)} + \frac{(x^{(3/2)}*\text{Sin}[a + b*x^2])}{(2*b)}$$

Defintions of rubi rules used

```
rule 2648 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_
.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[
F])^((m + 1)/n)))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F
, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

```
rule 3867 Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_.), x_Symbol] := Simp[e^(n
- 1)*(e*x)^(m - n + 1)*(Sin[c + d*x^n]/(d*n)), x] - Simp[e^n*((m - n + 1)/
(d*n)) Int[(e*x)^(m - n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] &&
IGtQ[n, 0] && LtQ[n, m + 1]
```

```
rule 3870 Int[((e_.)*(x_)^(m_.)*Sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[I/2
Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] - Simp[I/2 Int[(e*x)^m*E^(c*I
+ d*I*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.96 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.06

method	result
meijerg	$\frac{2^{\frac{3}{4}} \cos(a) \sqrt{\pi} \left(\frac{2x^{\frac{3}{2}} 2^{\frac{1}{4}} (b^2)^{\frac{7}{8}} \sin(bx^2)}{7\sqrt{\pi} b} + \frac{3x^{\frac{7}{2}} (b^2)^{\frac{7}{8}} 2^{\frac{1}{4}} \sin(bx^2) \operatorname{LommelS1}\left(\frac{5}{4}, \frac{3}{2}, bx^2\right)}{14\sqrt{\pi} (bx^2)^{\frac{5}{4}}} + \frac{3x^{\frac{7}{2}} (b^2)^{\frac{7}{8}} 2^{\frac{1}{4}} (\cos(bx^2)bx^2 - \sin(bx^2)) \operatorname{LommelS1}\left(\frac{5}{4}, \frac{3}{2}, bx^2\right)}{8\sqrt{\pi} (bx^2)^{\frac{9}{4}}} \right)}{2(b^2)^{\frac{7}{8}}}$

```
input int(x^(5/2)*cos(b*x^2+a), x, method=_RETURNVERBOSE)
```

output

```
1/2*2^(3/4)/(b^2)^(7/8)*cos(a)*Pi^(1/2)*(2/7/Pi^(1/2)*x^(3/2)*2^(1/4)*(b^2)^(7/8)/b*sin(b*x^2)+3/14/Pi^(1/2)*x^(7/2)*(b^2)^(7/8)*2^(1/4)/(b*x^2)^(5/4)*sin(b*x^2)*LommelS1(5/4,3/2,b*x^2)+3/8/Pi^(1/2)*x^(7/2)*(b^2)^(7/8)*2^(1/4)/(b*x^2)^(9/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(1/4,1/2,b*x^2))-1/2*2^(3/4)/b^(7/4)*sin(a)*Pi^(1/2)*(-1/8/Pi^(1/2)*x^(7/2)*b^(7/4)*2^(1/4)/(b*x^2)^(5/4)*sin(b*x^2)*LommelS1(1/4,3/2,b*x^2)-1/2/Pi^(1/2)*x^(7/2)*b^(7/4)*2^(1/4)/(b*x^2)^(9/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(5/4,1/2,b*x^2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.58

$$\int x^{5/2} \cos(a + bx^2) dx = \frac{8bx^{3/2} \sin(bx^2 + a) + 3(ib)^{1/4} (\cos(a) - i \sin(a)) \Gamma(\frac{3}{4}, ibx^2) + 3(-ib)^{1/4} (\cos(a) + i \sin(a)) \Gamma(\frac{3}{4}, -ibx^2)}{16b^2}$$

input

```
integrate(x^(5/2)*cos(b*x^2+a),x, algorithm="fricas")
```

output

```
1/16*(8*b*x^(3/2)*sin(b*x^2 + a) + 3*(I*b)^(1/4)*(cos(a) - I*sin(a))*gamma(3/4, I*b*x^2) + 3*(-I*b)^(1/4)*(cos(a) + I*sin(a))*gamma(3/4, -I*b*x^2))/b^2
```

Sympy [F]

$$\int x^{5/2} \cos(a + bx^2) dx = \int x^{5/2} \cos(a + bx^2) dx$$

input

```
integrate(x**(5/2)*cos(b*x**2+a),x)
```

output

```
Integral(x**(5/2)*cos(a + b*x**2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int x^{5/2} \cos(a + bx^2) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^(5/2)*cos(b*x^2+a),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int x^{5/2} \cos(a + bx^2) dx = \int x^{5/2} \cos(bx^2 + a) dx$$

input `integrate(x^(5/2)*cos(b*x^2+a),x, algorithm="giac")`

output `integrate(x^(5/2)*cos(b*x^2 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{5/2} \cos(a + bx^2) dx = \int x^{5/2} \cos(bx^2 + a) dx$$

input `int(x^(5/2)*cos(a + b*x^2),x)`

output `int(x^(5/2)*cos(a + b*x^2), x)`

Reduce [F]

$$\int x^{5/2} \cos(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a) x^2 dx$$

input `int(x^(5/2)*cos(b*x^2+a),x)`

output `int(sqrt(x)*cos(a + b*x**2)*x**2,x)`

3.24 $\int x^{3/2} \cos(a + bx^2) dx$

Optimal result	201
Mathematica [A] (verified)	201
Rubi [A] (verified)	202
Maple [C] (verified)	203
Fricas [A] (verification not implemented)	204
Sympy [F]	204
Maxima [F(-2)]	205
Giac [F]	205
Mupad [F(-1)]	205
Reduce [F]	206

Optimal result

Integrand size = 14, antiderivative size = 111

$$\int x^{3/2} \cos(a + bx^2) dx = -\frac{ie^{ia}\sqrt{x}\Gamma(\frac{1}{4}, -ibx^2)}{16b\sqrt[4]{-ibx^2}} + \frac{ie^{-ia}\sqrt{x}\Gamma(\frac{1}{4}, ibx^2)}{16b\sqrt[4]{ibx^2}} + \frac{\sqrt{x} \sin(a + bx^2)}{2b}$$

output

```
-1/16*I*exp(I*a)*x^(1/2)*GAMMA(1/4, -I*b*x^2)/b/(-I*b*x^2)^(1/4)+1/16*I*x^(1/2)*GAMMA(1/4, I*b*x^2)/b/exp(I*a)/(I*b*x^2)^(1/4)+1/2*x^(1/2)*sin(b*x^2+a)/b
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00

$$\int x^{3/2} \cos(a + bx^2) dx = \frac{bx^{9/2} \left(\sqrt[4]{ibx^2} \Gamma(\frac{1}{4}, -ibx^2) (-i \cos(a) + \sin(a)) + \sqrt[4]{-ibx^2} \Gamma(\frac{1}{4}, ibx^2) (i \cos(a) + \sin(a)) + 8\sqrt[4]{b^2x^2} \right)}{16(b^2x^4)^{5/4}}$$

input

```
Integrate[x^(3/2)*Cos[a + b*x^2], x]
```

output

$$\frac{(b*x^{9/2})*((I*b*x^2)^{1/4}*Gamma[1/4, (-I)*b*x^2]*((-I)*Cos[a] + Sin[a]) + ((-I)*b*x^2)^{1/4}*Gamma[1/4, I*b*x^2]*(I*Cos[a] + Sin[a]) + 8*(b^2*x^4)^{1/4}*Sin[a + b*x^2])}{(16*(b^2*x^4)^{5/4})}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3867, 3870, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{3/2} \cos(a + bx^2) dx \\ & \quad \downarrow \text{3867} \\ & \frac{\sqrt{x} \sin(a + bx^2)}{2b} - \frac{\int \frac{\sin(bx^2+a)}{\sqrt{x}} dx}{4b} \\ & \quad \downarrow \text{3870} \\ & \frac{\sqrt{x} \sin(a + bx^2)}{2b} - \frac{\frac{1}{2}i \int \frac{e^{-ibx^2-ia}}{\sqrt{x}} dx - \frac{1}{2}i \int \frac{e^{ibx^2+ia}}{\sqrt{x}} dx}{4b} \\ & \quad \downarrow \text{2648} \\ & \frac{\sqrt{x} \sin(a + bx^2)}{2b} - \frac{\frac{ie^{ia}\sqrt{x}\Gamma(\frac{1}{4}, -ibx^2)}{4\sqrt[4]{-ibx^2}} - \frac{ie^{-ia}\sqrt{x}\Gamma(\frac{1}{4}, ibx^2)}{4\sqrt[4]{ibx^2}}}{4b} \end{aligned}$$

input

```
Int[x^(3/2)*Cos[a + b*x^2], x]
```

output

```
-1/4*(((I/4)*E^(I*a)*Sqrt[x]*Gamma[1/4, (-I)*b*x^2])/((-I)*b*x^2)^(1/4) -
((I/4)*Sqrt[x]*Gamma[1/4, I*b*x^2])/(E^(I*a)*(I*b*x^2)^(1/4))/b + (Sqrt[x]
]*Sin[a + b*x^2])/(2*b)
```

Defintions of rubi rules used

```
rule 2648 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[F])^(m + 1)/n))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

```
rule 3867 Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_.), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(Sin[c + d*x^n]/(d*n)), x] - Simp[e^n*((m - n + 1)/(d*n)) Int[(e*x)^(m - n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]
```

```
rule 3870 Int[((e_.)*(x_)^(m_.)*Sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[I/2 Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] - Simp[I/2 Int[(e*x)^m*E^(c*I + d*I*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.86 (sec) , antiderivative size = 290, normalized size of antiderivative = 2.61

method	result
meijerg	$\frac{2^{\frac{1}{4}} \cos(a) \sqrt{\pi} \left(\frac{2\sqrt{x} 2^{\frac{3}{4}} (b^2)^{\frac{5}{8}} \sin(bx^2)}{5\sqrt{\pi} b} + \frac{2\sqrt{x} 2^{\frac{3}{4}} (b^2)^{\frac{5}{8}} (\cos(bx^2)bx^2 - \sin(bx^2))}{5\sqrt{\pi} b} + \frac{x^{\frac{9}{2}} (b^2)^{\frac{5}{8}} 2^{\frac{3}{4}} b \sin(bx^2) \text{LommelS1}\left(\frac{3}{4}, \frac{3}{2}, bx^2\right)}{10\sqrt{\pi} (bx^2)^{\frac{7}{4}}} - \frac{2x^{\frac{9}{2}}}{2(b^2)^{\frac{5}{8}}} \right)}{2(b^2)^{\frac{5}{8}}}$

```
input int(x^(3/2)*cos(b*x^2+a), x, method=_RETURNVERBOSE)
```

output

```
1/2*2^(1/4)/(b^2)^(5/8)*cos(a)*Pi^(1/2)*(2/5/Pi^(1/2)*x^(1/2)*2^(3/4)*(b^2)^(5/8)/b*sin(b*x^2)+2/5/Pi^(1/2)*x^(1/2)*2^(3/4)*(b^2)^(5/8)/b*(cos(b*x^2)*b*x^2-sin(b*x^2))+1/10/Pi^(1/2)*x^(9/2)*(b^2)^(5/8)*2^(3/4)*b/(b*x^2)^(7/4)*sin(b*x^2)*LommelS1(3/4,3/2,b*x^2)-2/5/Pi^(1/2)*x^(9/2)*(b^2)^(5/8)*2^(3/4)*b/(b*x^2)^(11/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(7/4,1/2,b*x^2))-1/2*2^(1/4)/b^(5/4)*sin(a)*Pi^(1/2)*(2/9/Pi^(1/2)*x^(5/2)*2^(3/4)*b^(5/4)*sin(b*x^2)-2/9/Pi^(1/2)*x^(9/2)*b^(9/4)*2^(3/4)/(b*x^2)^(7/4)*sin(b*x^2)*LommelS1(7/4,3/2,b*x^2)-1/2/Pi^(1/2)*x^(9/2)*b^(9/4)*2^(3/4)/(b*x^2)^(11/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(3/4,1/2,b*x^2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.56

$$\int x^{3/2} \cos(a + bx^2) dx = \frac{(ib)^{3/4} (\cos(a) - i \sin(a)) \Gamma(\frac{1}{4}, ibx^2) + (-ib)^{3/4} (\cos(a) + i \sin(a)) \Gamma(\frac{1}{4}, -ibx^2) + 8b\sqrt{x} \sin(a)}{16b^2}$$

input

```
integrate(x^(3/2)*cos(b*x^2+a),x, algorithm="fricas")
```

output

```
1/16*((I*b)^(3/4)*(cos(a) - I*sin(a))*gamma(1/4, I*b*x^2) + (-I*b)^(3/4)*(cos(a) + I*sin(a))*gamma(1/4, -I*b*x^2) + 8*b*sqrt(x)*sin(b*x^2 + a))/b^2
```

Sympy [F]

$$\int x^{3/2} \cos(a + bx^2) dx = \int x^{3/2} \cos(a + bx^2) dx$$

input

```
integrate(x**(3/2)*cos(b*x**2+a),x)
```

output

```
Integral(x**(3/2)*cos(a + b*x**2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int x^{3/2} \cos(a + bx^2) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^(3/2)*cos(b*x^2+a),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int x^{3/2} \cos(a + bx^2) dx = \int x^{3/2} \cos(bx^2 + a) dx$$

input `integrate(x^(3/2)*cos(b*x^2+a),x, algorithm="giac")`

output `integrate(x^(3/2)*cos(b*x^2 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{3/2} \cos(a + bx^2) dx = \int x^{3/2} \cos(bx^2 + a) dx$$

input `int(x^(3/2)*cos(a + b*x^2),x)`

output `int(x^(3/2)*cos(a + b*x^2), x)`

Reduce [F]

$$\int x^{3/2} \cos(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a) x dx$$

input `int(x^(3/2)*cos(b*x^2+a),x)`

output `int(sqrt(x)*cos(a + b*x**2)*x,x)`

3.25 $\int \sqrt{x} \cos(a + bx^2) dx$

Optimal result	207
Mathematica [A] (verified)	207
Rubi [A] (verified)	208
Maple [C] (verified)	209
Fricas [A] (verification not implemented)	210
Sympy [F]	210
Maxima [F(-2)]	210
Giac [F]	211
Mupad [F(-1)]	211
Reduce [F]	211

Optimal result

Integrand size = 14, antiderivative size = 81

$$\int \sqrt{x} \cos(a + bx^2) dx = -\frac{e^{ia}x^{3/2}\Gamma(\frac{3}{4}, -ibx^2)}{4(-ibx^2)^{3/4}} - \frac{e^{-ia}x^{3/2}\Gamma(\frac{3}{4}, ibx^2)}{4(ibx^2)^{3/4}}$$

output `-1/4*exp(I*a)*x^(3/2)*GAMMA(3/4,-I*b*x^2)/(-I*b*x^2)^(3/4)-1/4*x^(3/2)*GAMMA(3/4,I*b*x^2)/exp(I*a)/(I*b*x^2)^(3/4)`

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.10

$$\int \sqrt{x} \cos(a + bx^2) dx = \frac{x^{3/2} \left((-ibx^2)^{3/4} \Gamma(\frac{3}{4}, ibx^2) (\cos(a) - i \sin(a)) + (ibx^2)^{3/4} \Gamma(\frac{3}{4}, -ibx^2) (\cos(a) + i \sin(a)) \right)}{4(b^2x^4)^{3/4}}$$

input `Integrate[Sqrt[x]*Cos[a + b*x^2],x]`

output

$$\frac{-1/4*(x^{3/2}*(((-I)*b*x^2)^{3/4}*Gamma[3/4, I*b*x^2]*(Cos[a] - I*Sin[a]) + (I*b*x^2)^{3/4}*Gamma[3/4, (-I)*b*x^2]*(Cos[a] + I*Sin[a])))/(b^2*x^4)^{3/4}}$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3871, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \cos(a + bx^2) dx$$

$$\downarrow \text{3871}$$

$$\frac{1}{2} \int e^{-ibx^2 - ia} \sqrt{x} dx + \frac{1}{2} \int e^{ibx^2 + ia} \sqrt{x} dx$$

$$\downarrow \text{2648}$$

$$\frac{e^{ia} x^{3/2} \Gamma\left(\frac{3}{4}, -ibx^2\right)}{4(-ibx^2)^{3/4}} - \frac{e^{-ia} x^{3/2} \Gamma\left(\frac{3}{4}, ibx^2\right)}{4(ibx^2)^{3/4}}$$

input

```
Int[Sqrt[x]*Cos[a + b*x^2], x]
```

output

$$\frac{-1/4*(E^{I*a}*x^{3/2}*Gamma[3/4, (-I)*b*x^2])/((-I)*b*x^2)^{3/4} - (x^{3/2})*Gamma[3/4, I*b*x^2]}{(4*E^{I*a}*(I*b*x^2)^{3/4})}$$

Definitions of rubi rules used

rule 2648

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[F])^(m + 1)/n))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

rule 3871

```
Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_.), x_Symbol] := Simp[1/2 Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] + Simp[1/2 Int[(e*x)^m*E^(c*I + d*I*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.86 (sec) , antiderivative size = 290, normalized size of antiderivative = 3.58

method	result
meijerg	$2^{\frac{3}{4}} \cos(a) \sqrt{\pi} \left(\frac{4 \cdot 2^{\frac{1}{4}} (b^2)^{\frac{3}{8}} \sin(bx^2)}{3\sqrt{\pi} \sqrt{x} b} + \frac{4 \cdot 2^{\frac{1}{4}} (b^2)^{\frac{3}{8}} (\cos(bx^2) b x^2 - \sin(bx^2))}{3\sqrt{\pi} \sqrt{x} b} - \frac{x^{\frac{7}{2}} (b^2)^{\frac{3}{8}} 2^{\frac{1}{4}} b \sin(bx^2) \text{LommelS1}(\frac{1}{4}, \frac{3}{2}, bx^2)}{3\sqrt{\pi} (bx^2)^{\frac{5}{4}}} - \frac{4x^{\frac{7}{2}} (b^2)^{\frac{3}{8}}}{4(b^2)^{\frac{3}{8}}} \right)$

input

```
int(x^(1/2)*cos(b*x^2+a), x, method=_RETURNVERBOSE)
```

output

```
1/4*2^(3/4)/(b^2)^(3/8)*cos(a)*Pi^(1/2)*(4/3/Pi^(1/2)/x^(1/2)*2^(1/4)*(b^2)^(3/8)/b*sin(b*x^2)+4/3/Pi^(1/2)/x^(1/2)*2^(1/4)*(b^2)^(3/8)/b*(cos(b*x^2)*b*x^2-sin(b*x^2))-1/3/Pi^(1/2)*x^(7/2)*(b^2)^(3/8)*2^(1/4)*b/(b*x^2)^(5/4)*sin(b*x^2)*LommelS1(1/4, 3/2, b*x^2)-4/3/Pi^(1/2)*x^(7/2)*(b^2)^(3/8)*2^(1/4)*b/(b*x^2)^(9/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(5/4, 1/2, b*x^2)-1/4*2^(3/4)/b^(3/4)*sin(a)*Pi^(1/2)*(4/7/Pi^(1/2)*x^(3/2)*2^(1/4)*b^(3/4)*sin(b*x^2)-4/7/Pi^(1/2)*x^(7/2)*b^(7/4)*2^(1/4)/(b*x^2)^(5/4)*sin(b*x^2)*LommelS1(5/4, 3/2, b*x^2)-1/Pi^(1/2)*x^(7/2)*b^(7/4)*2^(1/4)/(b*x^2)^(9/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(1/4, 1/2, b*x^2)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.59

$$\int \sqrt{x} \cos(a + bx^2) dx$$

$$= \frac{(ib)^{\frac{1}{4}} (i \cos(a) + \sin(a)) \Gamma(\frac{3}{4}, ibx^2) + (-ib)^{\frac{1}{4}} (-i \cos(a) + \sin(a)) \Gamma(\frac{3}{4}, -ibx^2)}{4b}$$

input `integrate(x^(1/2)*cos(b*x^2+a),x, algorithm="fricas")`

output `1/4*((I*b)^(1/4)*(I*cos(a) + sin(a))*gamma(3/4, I*b*x^2) + (-I*b)^(1/4)*(-I*cos(a) + sin(a))*gamma(3/4, -I*b*x^2))/b`

Sympy [F]

$$\int \sqrt{x} \cos(a + bx^2) dx = \int \sqrt{x} \cos(a + bx^2) dx$$

input `integrate(x**(1/2)*cos(b*x**2+a),x)`

output `Integral(sqrt(x)*cos(a + b*x**2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{x} \cos(a + bx^2) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^(1/2)*cos(b*x^2+a),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \sqrt{x} \cos(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a) dx$$

input `integrate(x^(1/2)*cos(b*x^2+a),x, algorithm="giac")`

output `integrate(sqrt(x)*cos(b*x^2 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x} \cos(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a) dx$$

input `int(x^(1/2)*cos(a + b*x^2),x)`

output `int(x^(1/2)*cos(a + b*x^2), x)`

Reduce [F]

$$\int \sqrt{x} \cos(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a) dx$$

input `int(x^(1/2)*cos(b*x^2+a),x)`

output `int(sqrt(x)*cos(a + b*x**2),x)`

3.26 $\int \frac{\cos(a+bx^2)}{\sqrt{x}} dx$

Optimal result	212
Mathematica [A] (verified)	212
Rubi [A] (verified)	213
Maple [C] (verified)	214
Fricas [A] (verification not implemented)	215
Sympy [F]	215
Maxima [F(-2)]	215
Giac [F]	216
Mupad [F(-1)]	216
Reduce [F]	216

Optimal result

Integrand size = 14, antiderivative size = 81

$$\int \frac{\cos(a+bx^2)}{\sqrt{x}} dx = -\frac{e^{ia}\sqrt{x}\Gamma(\frac{1}{4},-ibx^2)}{4\sqrt[4]{-ibx^2}} - \frac{e^{-ia}\sqrt{x}\Gamma(\frac{1}{4},ibx^2)}{4\sqrt[4]{ibx^2}}$$

output `-1/4*exp(I*a)*x^(1/2)*GAMMA(1/4,-I*b*x^2)/(-I*b*x^2)^(1/4)-1/4*x^(1/2)*GAMMA(1/4,I*b*x^2)/exp(I*a)/(I*b*x^2)^(1/4)`

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.10

$$\int \frac{\cos(a+bx^2)}{\sqrt{x}} dx = \frac{\sqrt{x}\left(\sqrt[4]{-ibx^2}\Gamma(\frac{1}{4},ibx^2)(\cos(a)-i\sin(a))+\sqrt[4]{ibx^2}\Gamma(\frac{1}{4},-ibx^2)(\cos(a)+i\sin(a))\right)}{4\sqrt[4]{b^2x^4}}$$

input `Integrate[Cos[a + b*x^2]/Sqrt[x],x]`

output

```
-1/4*(Sqrt[x]*(((I)*b*x^2)^(1/4)*Gamma[1/4, I*b*x^2]*(Cos[a] - I*Sin[a])
+ (I*b*x^2)^(1/4)*Gamma[1/4, (-I)*b*x^2]*(Cos[a] + I*Sin[a])))/(b^2*x^4)^(
1/4)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3871, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx$$

$$\downarrow \text{3871}$$

$$\frac{1}{2} \int \frac{e^{-ibx^2 - ia}}{\sqrt{x}} dx + \frac{1}{2} \int \frac{e^{ibx^2 + ia}}{\sqrt{x}} dx$$

$$\downarrow \text{2648}$$

$$-\frac{e^{ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -ibx^2\right)}{4 \sqrt[4]{-ibx^2}} - \frac{e^{-ia} \sqrt{x} \Gamma\left(\frac{1}{4}, ibx^2\right)}{4 \sqrt[4]{ibx^2}}$$

input

```
Int[Cos[a + b*x^2]/Sqrt[x],x]
```

output

```
-1/4*(E^(I*a)*Sqrt[x]*Gamma[1/4, (-I)*b*x^2])/((-I)*b*x^2)^(1/4) - (Sqrt[x]
)*Gamma[1/4, I*b*x^2])/(4*E^(I*a)*(I*b*x^2)^(1/4))
```

Defintions of rubi rules used

rule 2648

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[F])^(m + 1)/n))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

rule 3871

```
Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_.), x_Symbol] := Simp[1/2 Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] + Simp[1/2 Int[(e*x)^m*E^(c*I + d*I*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.81 (sec) , antiderivative size = 338, normalized size of antiderivative = 4.17

method	result
meijerg	$\frac{\cos(a)\sqrt{\pi}2^{\frac{1}{4}}}{4(b^2)^{\frac{1}{8}}}\left(\frac{62^{\frac{3}{4}}(b^2)^{\frac{1}{8}}\left(\frac{8b^2x^4}{27}+\frac{2}{3}\right)\sin(bx^2)}{\sqrt{\pi}x^{\frac{3}{2}}b} + \frac{42^{\frac{3}{4}}(b^2)^{\frac{1}{8}}(\cos(bx^2)bx^2 - \sin(bx^2))}{\sqrt{\pi}x^{\frac{3}{2}}b} - \frac{16x^{\frac{9}{2}}(b^2)^{\frac{1}{8}}b^22^{\frac{3}{4}}\sin(bx^2)\text{LommelS1}\left(\frac{7}{4}, \frac{3}{2}, bx^2\right)}{9\sqrt{\pi}(bx^2)^{\frac{7}{4}}}\right)$

input

```
int(cos(b*x^2+a)/x^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/4*cos(a)*Pi^(1/2)*2^(1/4)/(b^2)^(1/8)*(6/Pi^(1/2)/x^(3/2)*2^(3/4)*(b^2)^(1/8)*(8/27*b^2*x^4+2/3)/b*sin(b*x^2)+4/Pi^(1/2)/x^(3/2)*2^(3/4)*(b^2)^(1/8)/b*(cos(b*x^2)*b*x^2-sin(b*x^2))-16/9/Pi^(1/2)*x^(9/2)*(b^2)^(1/8)*b^2*2^(3/4)/(b*x^2)^(7/4)*sin(b*x^2)*LommelS1(7/4,3/2,b*x^2)-4/Pi^(1/2)*x^(9/2)*(b^2)^(1/8)*b^2*2^(3/4)/(b*x^2)^(11/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(3/4,1/2,b*x^2))-1/4*sin(a)*Pi^(1/2)*2^(1/4)/b^(1/4)*(4/5/Pi^(1/2)*x^(1/2)*2^(3/4)*b^(1/4)*sin(b*x^2)-16/5/Pi^(1/2)*x^(1/2)*2^(3/4)*b^(1/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))-4/5/Pi^(1/2)*x^(9/2)*b^(9/4)*2^(3/4)/(b*x^2)^(7/4)*sin(b*x^2)*LommelS1(3/4,3/2,b*x^2)+16/5/Pi^(1/2)*x^(9/2)*b^(9/4)*2^(3/4)/(b*x^2)^(11/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(7/4,1/2,b*x^2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.59

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx$$

$$= \frac{(ib)^{\frac{3}{4}} (i \cos(a) + \sin(a)) \Gamma(\frac{1}{4}, ibx^2) + (-ib)^{\frac{3}{4}} (-i \cos(a) + \sin(a)) \Gamma(\frac{1}{4}, -ibx^2)}{4b}$$

input `integrate(cos(b*x^2+a)/x^(1/2),x, algorithm="fricas")`

output `1/4*((I*b)^(3/4)*(I*cos(a) + sin(a))*gamma(1/4, I*b*x^2) + (-I*b)^(3/4)*(-I*cos(a) + sin(a))*gamma(1/4, -I*b*x^2))/b`

Sympy [F]

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(a + bx^2)}{\sqrt{x}} dx$$

input `integrate(cos(b*x**2+a)/x**(1/2),x)`

output `Integral(cos(a + b*x**2)/sqrt(x), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cos(b*x^2+a)/x^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(bx^2 + a)}{\sqrt{x}} dx$$

input `integrate(cos(b*x^2+a)/x^(1/2),x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)/sqrt(x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(bx^2 + a)}{\sqrt{x}} dx$$

input `int(cos(a + b*x^2)/x^(1/2),x)`

output `int(cos(a + b*x^2)/x^(1/2), x)`

Reduce [F]

$$\int \frac{\cos(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(bx^2 + a)}{\sqrt{x}} dx$$

input `int(cos(b*x^2+a)/x^(1/2),x)`

output `int(cos(a + b*x**2)/sqrt(x),x)`

3.27 $\int \frac{\cos(a+bx^2)}{x^{3/2}} dx$

Optimal result	217
Mathematica [A] (verified)	217
Rubi [A] (verified)	218
Maple [C] (verified)	219
Fricas [A] (verification not implemented)	220
Sympy [F]	220
Maxima [F(-2)]	221
Giac [F]	221
Mupad [F(-1)]	221
Reduce [F]	222

Optimal result

Integrand size = 14, antiderivative size = 98

$$\int \frac{\cos(a+bx^2)}{x^{3/2}} dx = -\frac{2 \cos(a+bx^2)}{\sqrt{x}} - \frac{ibe^{ia}x^{3/2}\Gamma(\frac{3}{4}, -ibx^2)}{(-ibx^2)^{3/4}} + \frac{ibe^{-ia}x^{3/2}\Gamma(\frac{3}{4}, ibx^2)}{(ibx^2)^{3/4}}$$

output

```
-2*cos(b*x^2+a)/x^(1/2)-I*b*exp(I*a)*x^(3/2)*GAMMA(3/4,-I*b*x^2)/(-I*b*x^2)^(3/4)+I*b*x^(3/2)*GAMMA(3/4,I*b*x^2)/exp(I*a)/(I*b*x^2)^(3/4)
```

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.16

$$\int \frac{\cos(a+bx^2)}{x^{3/2}} dx = \frac{-2(b^2x^4)^{3/4} \cos(a+bx^2) + bx^2(ibx^2)^{3/4} \Gamma(\frac{3}{4}, -ibx^2) (-i \cos(a) + \sin(a)) + i(-ibx^2)^{3/4}}{\sqrt{x} (b^2x^4)^{3/4}}$$

input

```
Integrate[Cos[a + b*x^2]/x^(3/2), x]
```

output

```
(-2*(b^2*x^4)^(3/4)*Cos[a + b*x^2] + b*x^2*(I*b*x^2)^(3/4)*Gamma[3/4, (-I)*b*x^2]*((-I)*Cos[a] + Sin[a]) + I*((-I)*b*x^2)^(7/4)*Gamma[3/4, I*b*x^2]*(I*cos[a] + Sin[a]))/(Sqrt[x]*(b^2*x^4)^(3/4))
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.06, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3869, 3870, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(a + bx^2)}{x^{3/2}} dx \\ & \quad \downarrow \text{3869} \\ & -4b \int \sqrt{x} \sin(bx^2 + a) dx - \frac{2 \cos(a + bx^2)}{\sqrt{x}} \\ & \quad \downarrow \text{3870} \\ & -\frac{2 \cos(a + bx^2)}{\sqrt{x}} - 4b \left(\frac{1}{2} i \int e^{-ibx^2 - ia} \sqrt{x} dx - \frac{1}{2} i \int e^{ibx^2 + ia} \sqrt{x} dx \right) \\ & \quad \downarrow \text{2648} \\ & -\frac{2 \cos(a + bx^2)}{\sqrt{x}} - 4b \left(\frac{ie^{ia} x^{3/2} \Gamma(\frac{3}{4}, -ibx^2)}{4(-ibx^2)^{3/4}} - \frac{ie^{-ia} x^{3/2} \Gamma(\frac{3}{4}, ibx^2)}{4(ibx^2)^{3/4}} \right) \end{aligned}$$

input `Int[Cos[a + b*x^2]/x^(3/2),x]`

output `(-2*Cos[a + b*x^2])/Sqrt[x] - 4*b*((I/4)*E^(I*a)*x^(3/2)*Gamma[3/4, (-I)*b*x^2])/((-I)*b*x^2)^(3/4) - ((I/4)*x^(3/2)*Gamma[3/4, I*b*x^2])/(E^(I*a)*(I*b*x^2)^(3/4))`

Defintions of rubi rules used

```
rule 2648 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[F])^(m + 1)/n))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

```
rule 3869 Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_)), x_Symbol] := Simp[(e*x)^(m + 1)*(Cos[c + d*x^n]/(e*(m + 1))), x] + Simp[d*(n/(e^n*(m + 1))) Int[(e*x)^(m + n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] & LtQ[m, -1]
```

```
rule 3870 Int[((e_.)*(x_)^(m_.)*Sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[I/2 Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] - Simp[I/2 Int[(e*x)^m*E^(c*I + d*I*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.82 (sec) , antiderivative size = 338, normalized size of antiderivative = 3.45

method	result
meijerg	$\frac{\cos(a)\sqrt{\pi}2^{\frac{3}{4}}(b^2)^{\frac{1}{8}}}{8} \left(-\frac{122^{\frac{1}{4}}\left(\frac{8b^2x^4}{21} + \frac{2}{3}\right)\sin(bx^2)}{\sqrt{\pi}x^{\frac{5}{2}}(b^2)^{\frac{1}{8}}b} - \frac{82^{\frac{1}{4}}(\cos(bx^2)bx^2 - \sin(bx^2))}{\sqrt{\pi}x^{\frac{5}{2}}(b^2)^{\frac{1}{8}}b} + \frac{32x^{\frac{7}{2}}b^22^{\frac{1}{4}}\sin(bx^2)\text{LommelS1}\left(\frac{5}{4}, \frac{3}{2}, bx^2\right)}{7\sqrt{\pi}(b^2)^{\frac{1}{8}}(bx^2)^{\frac{5}{4}}} + \frac{8x^{\frac{7}{2}}b^{\frac{7}{2}}}{8} \right)$

```
input int(cos(b*x^2+a)/x^(3/2),x,method=_RETURNVERBOSE)
```

output

```
1/8*cos(a)*Pi^(1/2)*2^(3/4)*(b^2)^(1/8)*(-12/Pi^(1/2)/x^(5/2)*2^(1/4)/(b^2)^(1/8)*(8/21*b^2*x^4+2/3)/b*sin(b*x^2)-8/Pi^(1/2)/x^(5/2)*2^(1/4)/(b^2)^(1/8)/b*(cos(b*x^2)*b*x^2-sin(b*x^2))+32/7/Pi^(1/2)*x^(7/2)/(b^2)^(1/8)*b^2*2^(1/4)/(b*x^2)^(5/4)*sin(b*x^2)*LommelS1(5/4,3/2,b*x^2)+8/Pi^(1/2)*x^(7/2)/(b^2)^(1/8)*b^2*2^(1/4)/(b*x^2)^(9/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(1/4,1/2,b*x^2))-1/8*sin(a)*Pi^(1/2)*2^(3/4)*b^(1/4)*(8/3/Pi^(1/2)/x^(1/2)*2^(1/4)/b^(1/4)*sin(b*x^2)+32/3/Pi^(1/2)/x^(1/2)*2^(1/4)/b^(1/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))-8/3/Pi^(1/2)*x^(7/2)*b^(7/4)*2^(1/4)/(b*x^2)^(5/4)*sin(b*x^2)*LommelS1(1/4,3/2,b*x^2)-32/3/Pi^(1/2)*x^(7/2)*b^(7/4)*2^(1/4)/(b*x^2)^(9/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))*LommelS1(5/4,1/2,b*x^2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.67

$$\int \frac{\cos(a + bx^2)}{x^{3/2}} dx = \frac{(x \cos(a) - i x \sin(a))(i b)^{\frac{1}{4}} \Gamma(\frac{3}{4}, i b x^2) + (x \cos(a) + i x \sin(a))(-i b)^{\frac{1}{4}} \Gamma(\frac{3}{4}, -i b x^2)}{x}$$

input

```
integrate(cos(b*x^2+a)/x^(3/2),x, algorithm="fricas")
```

output

```
((x*cos(a) - I*x*sin(a))*(I*b)^(1/4)*gamma(3/4, I*b*x^2) + (x*cos(a) + I*x*sin(a))*(-I*b)^(1/4)*gamma(3/4, -I*b*x^2) - 2*sqrt(x)*cos(b*x^2 + a))/x
```

Sympy [F]

$$\int \frac{\cos(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos(a + bx^2)}{x^{\frac{3}{2}}} dx$$

input

```
integrate(cos(b*x**2+a)/x**(3/2),x)
```

output

```
Integral(cos(a + b*x**2)/x**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(a + bx^2)}{x^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cos(b*x^2+a)/x^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \frac{\cos(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos(bx^2 + a)}{x^{\frac{3}{2}}} dx$$

input `integrate(cos(b*x^2+a)/x^(3/2),x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)/x^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos(bx^2 + a)}{x^{3/2}} dx$$

input `int(cos(a + b*x^2)/x^(3/2),x)`

output `int(cos(a + b*x^2)/x^(3/2), x)`

Reduce [F]

$$\int \frac{\cos(a + bx^2)}{x^{3/2}} dx = \frac{\sqrt{x} \left(\int \frac{\cos(bx^2+a)}{\sqrt{x}} dx \right) + \sqrt{x} \left(\int \frac{1}{\sqrt{x}} dx \right) + 2}{\sqrt{x}}$$

input `int(cos(b*x^2+a)/x^(3/2),x)`

output `(sqrt(x)*int(cos(a + b*x**2)/(sqrt(x)*x),x) + sqrt(x)*int(1/(sqrt(x)*x),x) + 2)/sqrt(x)`

3.28 $\int \frac{\cos(a+bx^2)}{x^{5/2}} dx$

Optimal result	223
Mathematica [A] (verified)	223
Rubi [A] (verified)	224
Maple [C] (verified)	225
Fricas [A] (verification not implemented)	226
Sympy [F]	226
Maxima [F(-2)]	227
Giac [F]	227
Mupad [F(-1)]	227
Reduce [F]	228

Optimal result

Integrand size = 14, antiderivative size = 104

$$\int \frac{\cos(a+bx^2)}{x^{5/2}} dx = -\frac{2\cos(a+bx^2)}{3x^{3/2}} - \frac{ibe^{ia}\sqrt{x}\Gamma(\frac{1}{4},-ibx^2)}{3\sqrt[4]{-ibx^2}} + \frac{ibe^{-ia}\sqrt{x}\Gamma(\frac{1}{4},ibx^2)}{3\sqrt[4]{ibx^2}}$$

output

```
-2/3*cos(b*x^2+a)/x^(3/2)-1/3*I*b*exp(I*a)*x^(1/2)*GAMMA(1/4,-I*b*x^2)/(-I
*b*x^2)^(1/4)+1/3*I*b*x^(1/2)*GAMMA(1/4,I*b*x^2)/exp(I*a)/(I*b*x^2)^(1/4)
```

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a+bx^2)}{x^{5/2}} dx = \frac{-2\sqrt[4]{b^2x^4}\cos(a+bx^2) + bx^2\sqrt[4]{ibx^2}\Gamma(\frac{1}{4},-ibx^2)(-i\cos(a) + \sin(a)) + i(-ibx^2)^{5/4}\Gamma(\frac{1}{4},ibx^2)}{3x^{3/2}\sqrt[4]{b^2x^4}}$$

input

```
Integrate[Cos[a + b*x^2]/x^(5/2),x]
```

output

```
(-2*(b^2*x^4)^(1/4)*Cos[a + b*x^2] + b*x^2*(I*b*x^2)^(1/4)*Gamma[1/4, (-I
*b*x^2)*((-I)*Cos[a] + Sin[a]) + I*((-I)*b*x^2)^(5/4)*Gamma[1/4, I*b*x^2]*
(I*Cos[a] + Sin[a]))/(3*x^(3/2)*(b^2*x^4)^(1/4))
```


Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.04, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3869, 3870, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx$$

$$\downarrow \text{3869}$$

$$-\frac{4}{3}b \int \frac{\sin(bx^2 + a)}{\sqrt{x}} dx - \frac{2 \cos(a + bx^2)}{3x^{3/2}}$$

$$\downarrow \text{3870}$$

$$-\frac{2 \cos(a + bx^2)}{3x^{3/2}} - \frac{4}{3}b \left(\frac{1}{2}i \int \frac{e^{-ibx^2 - ia}}{\sqrt{x}} dx - \frac{1}{2}i \int \frac{e^{ibx^2 + ia}}{\sqrt{x}} dx \right)$$

$$\downarrow \text{2648}$$

$$-\frac{2 \cos(a + bx^2)}{3x^{3/2}} - \frac{4}{3}b \left(\frac{ie^{ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -ibx^2\right)}{4\sqrt[4]{-ibx^2}} - \frac{ie^{-ia} \sqrt{x} \Gamma\left(\frac{1}{4}, ibx^2\right)}{4\sqrt[4]{ibx^2}} \right)$$

input `Int[Cos[a + b*x^2]/x^(5/2),x]`

output `(-2*Cos[a + b*x^2])/(3*x^(3/2)) - (4*b*(((I/4)*E^(I*a)*Sqrt[x]*Gamma[1/4, (-I)*b*x^2])/((-I)*b*x^2)^(1/4) - ((I/4)*Sqrt[x]*Gamma[1/4, I*b*x^2])/(E^(I*a)*(I*b*x^2)^(1/4))))/3`

Defintions of rubi rules used

```
rule 2648 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[F])^(m + 1)/n))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

```
rule 3869 Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_.), x_Symbol] := Simp[(e*x)^(m + 1)*(Cos[c + d*x^n]/(e*(m + 1))), x] + Simp[d*(n/(e^n*(m + 1))) Int[(e*x)^(m + n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[m, -1]
```

```
rule 3870 Int[((e_.)*(x_)^(m_.)*Sin[(c_.) + (d_.)*(x_)^(n_)], x_Symbol] := Simp[I/2 Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] - Simp[I/2 Int[(e*x)^m*E^(c*I + d*I*x^n), x], x] /; FreeQ[{c, d, e, m}, x] && IGtQ[n, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.84 (sec) , antiderivative size = 358, normalized size of antiderivative = 3.44

method	result
meijerg	$\frac{\cos(a)\sqrt{\pi}2^{\frac{1}{4}}(b^2)^{\frac{3}{8}} \left(-\frac{42^{\frac{3}{4}} \left(\frac{8b^2x^4}{15} + \frac{2}{3} \right) \sin(bx^2)}{\sqrt{\pi}x^{\frac{7}{2}}(b^2)^{\frac{3}{8}}b} - \frac{82^{\frac{3}{4}}(-16b^2x^4+5)(\cos(bx^2)bx^2-\sin(bx^2))}{15\sqrt{\pi}x^{\frac{7}{2}}(b^2)^{\frac{3}{8}}b} + \frac{32x^{\frac{9}{2}}2^{\frac{3}{4}}b^3\sin(bx^2)}{15\sqrt{\pi}(b^2)^{\frac{3}{8}}(bx^2)^{\frac{7}{4}}} \right)}{8} \text{LommelS1}\left(\frac{3}{4}, \frac{3}{2}, b\right)$

```
input int(cos(b*x^2+a)/x^(5/2),x,method=_RETURNVERBOSE)
```

output

```

1/8*cos(a)*Pi^(1/2)*2^(1/4)*(b^2)^(3/8)*(-4/Pi^(1/2)/x^(7/2)*2^(3/4)/(b^2)
^(3/8)*(8/15*b^2*x^4+2/3)/b*sin(b*x^2)-8/15/Pi^(1/2)/x^(7/2)*2^(3/4)/(b^2)
^(3/8)/b*(-16*b^2*x^4+5)*(cos(b*x^2)*b*x^2-sin(b*x^2))+32/15/Pi^(1/2)*x^(9
/2)/(b^2)^(3/8)*2^(3/4)*b^3/(b*x^2)^(7/4)*sin(b*x^2)*LommelS1(3/4,3/2,b*x^
2)-128/15/Pi^(1/2)*x^(9/2)/(b^2)^(3/8)*2^(3/4)*b^3/(b*x^2)^(11/4)*(cos(b*x
^2)*b*x^2-sin(b*x^2))*LommelS1(7/4,1/2,b*x^2))-1/8*sin(a)*Pi^(1/2)*2^(1/4)
*b^(3/4)*(12/Pi^(1/2)/x^(3/2)*2^(3/4)/b^(3/4)*(32/81*b^2*x^4+2/3)*sin(b*x^
2)+32/3/Pi^(1/2)/x^(3/2)*2^(3/4)/b^(3/4)*(cos(b*x^2)*b*x^2-sin(b*x^2))-128
/27/Pi^(1/2)*x^(9/2)*b^(9/4)*2^(3/4)/(b*x^2)^(7/4)*sin(b*x^2)*LommelS1(7/4
,3/2,b*x^2)-32/3/Pi^(1/2)*x^(9/2)*b^(9/4)*2^(3/4)/(b*x^2)^(11/4)*(cos(b*x^
2)*b*x^2-sin(b*x^2))*LommelS1(3/4,1/2,b*x^2))

```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.72

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx = \frac{(x^2 \cos(a) - i x^2 \sin(a))(i b)^{3/4} \Gamma(\frac{1}{4}, i b x^2) + (x^2 \cos(a) + i x^2 \sin(a))(-i b)^{3/4} \Gamma(\frac{1}{4}, -i b x^2)}{3 x^2}$$

input `integrate(cos(b*x^2+a)/x^(5/2),x, algorithm="fricas")`

output

```

1/3*((x^2*cos(a) - I*x^2*sin(a))*(I*b)^(3/4)*gamma(1/4, I*b*x^2) + (x^2*co
s(a) + I*x^2*sin(a))*(-I*b)^(3/4)*gamma(1/4, -I*b*x^2) - 2*sqrt(x)*cos(b*x
^2 + a))/x^2

```

Sympy [F]

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos(a + bx^2)}{x^{5/2}} dx$$

input `integrate(cos(b*x**2+a)/x**(5/2),x)`

output `Integral(cos(a + b*x**2)/x**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cos(b*x^2+a)/x^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos(bx^2 + a)}{x^{5/2}} dx$$

input `integrate(cos(b*x^2+a)/x^(5/2),x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)/x^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos(bx^2 + a)}{x^{5/2}} dx$$

input `int(cos(a + b*x^2)/x^(5/2),x)`

output `int(cos(a + b*x^2)/x^(5/2), x)`

Reduce [F]

$$\int \frac{\cos(a + bx^2)}{x^{5/2}} dx = \frac{3\sqrt{x} \left(\int \frac{\cos(bx^2+a)}{\sqrt{x}x^2} dx \right) x + 3\sqrt{x} \left(\int \frac{1}{\sqrt{x}x^2} dx \right) x + 2}{3\sqrt{x}x}$$

input `int(cos(b*x^2+a)/x^(5/2),x)`

output `(3*sqrt(x)*int(cos(a + b*x**2)/(sqrt(x)*x**2),x)*x + 3*sqrt(x)*int(1/(sqrt(x)*x**2),x)*x + 2)/(3*sqrt(x)*x)`

3.29 $\int x^{5/2} \cos^2(a + bx^2) dx$

Optimal result	229
Mathematica [A] (verified)	229
Rubi [A] (verified)	230
Maple [F]	231
Fricas [A] (verification not implemented)	231
Sympy [F]	232
Maxima [F(-2)]	232
Giac [F]	233
Mupad [F(-1)]	233
Reduce [F]	233

Optimal result

Integrand size = 16, antiderivative size = 132

$$\int x^{5/2} \cos^2(a + bx^2) dx = \frac{x^{7/2}}{7} - \frac{3ie^{2ia}x^{3/2}\Gamma(\frac{3}{4}, -2ibx^2)}{64 \cdot 2^{3/4}b(-ibx^2)^{3/4}} + \frac{3ie^{-2ia}x^{3/2}\Gamma(\frac{3}{4}, 2ibx^2)}{64 \cdot 2^{3/4}b(ibx^2)^{3/4}} + \frac{x^{3/2} \sin(2(a + bx^2))}{8b}$$

output

```
1/7*x^(7/2)-3/128*I*exp(2*I*a)*x^(3/2)*GAMMA(3/4,-2*I*b*x^2)*2^(1/4)/b/(-I
*b*x^2)^(3/4)+3/128*I*x^(3/2)*GAMMA(3/4,2*I*b*x^2)*2^(1/4)/b/exp(2*I*a)/(I
*b*x^2)^(3/4)+1/8*x^(3/2)*sin(2*b*x^2+2*a)/b
```

Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.08

$$\int x^{5/2} \cos^2(a + bx^2) dx = \frac{bx^{11/2} \left(21\sqrt[4]{2}(ibx^2)^{3/4} \Gamma(\frac{3}{4}, -2ibx^2) (-i \cos(2a) + \sin(2a)) + 21\sqrt[4]{2}(-ibx^2)^{3/4} \Gamma(\frac{3}{4}, 2ibx^2) (i \cos(2a) + \sin(2a)) \right)}{896(b^2x^4)^{7/4}}$$

input

```
Integrate[x^(5/2)*Cos[a + b*x^2]^2,x]
```

output

```
(b*x^(11/2)*(21*2^(1/4)*(I*b*x^2)^(3/4)*Gamma[3/4, (-2*I)*b*x^2]*((-I)*Cos
[2*a] + Sin[2*a]) + 21*2^(1/4)*((-I)*b*x^2)^(3/4)*Gamma[3/4, (2*I)*b*x^2]*
(I*Cos[2*a] + Sin[2*a]) + 16*(b^2*x^4)^(3/4)*(8*b*x^2 + 7*Sin[2*(a + b*x^2
)])))/(896*(b^2*x^4)^(7/4))
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3883, 3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{5/2} \cos^2(a + bx^2) dx$$

$$\downarrow \text{3883}$$

$$2 \int x^3 \cos^2(bx^2 + a) d\sqrt{x}$$

$$\downarrow \text{3885}$$

$$2 \int \left(\frac{1}{2} \cos(2bx^2 + 2a) x^3 + \frac{x^3}{2} \right) d\sqrt{x}$$

$$\downarrow \text{2009}$$

$$2 \left(\frac{x^{3/2} \sin(2a + 2bx^2)}{16b} - \frac{3ie^{2ia} x^{3/2} \Gamma(\frac{3}{4}, -2ibx^2)}{128 2^{3/4} b (-ibx^2)^{3/4}} + \frac{3ie^{-2ia} x^{3/2} \Gamma(\frac{3}{4}, 2ibx^2)}{128 2^{3/4} b (ibx^2)^{3/4}} + \frac{x^{7/2}}{14} \right)$$

input

```
Int[x^(5/2)*Cos[a + b*x^2]^2,x]
```

output

```
2*(x^(7/2)/14 - (((3*I)/128)*E^((2*I)*a)*x^(3/2)*Gamma[3/4, (-2*I)*b*x^2])
/(2^(3/4)*b*((-I)*b*x^2)^(3/4)) + (((3*I)/128)*x^(3/2)*Gamma[3/4, (2*I)*b*
x^2])/(2^(3/4)*b*E^((2*I)*a)*(I*b*x^2)^(3/4)) + (x^(3/2)*Sin[2*a + 2*b*x^2
])/((16*b))
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3883 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := With[{k = Denominator[m]}, Simp[k/e Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*(x^(k*n)/e^n])]^p, x], x, (e*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [F]

$$\int x^{\frac{5}{2}} \cos(bx^2 + a)^2 dx$$

input `int(x^(5/2)*cos(b*x^2+a)^2,x)`

output `int(x^(5/2)*cos(b*x^2+a)^2,x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.70

$$\int x^{5/2} \cos^2(a + bx^2) dx = \frac{21(2ib)^{\frac{1}{4}}(\cos(2a) - i \sin(2a))\Gamma(\frac{3}{4}, 2ibx^2) + 21(-2ib)^{\frac{1}{4}}(\cos(2a) + i \sin(2a))\Gamma(\frac{3}{4}, -2ibx^2)}{896b^2}$$

input `integrate(x^(5/2)*cos(b*x^2+a)^2,x, algorithm="fricas")`

output

```
1/896*(21*(2*I*b)^(1/4)*(cos(2*a) - I*sin(2*a))*gamma(3/4, 2*I*b*x^2) + 21
*(-2*I*b)^(1/4)*(cos(2*a) + I*sin(2*a))*gamma(3/4, -2*I*b*x^2) + 32*(4*b^2
*x^3 + 7*b*x*cos(b*x^2 + a)*sin(b*x^2 + a))*sqrt(x))/b^2
```

Sympy [F]

$$\int x^{5/2} \cos^2(a + bx^2) dx = \int x^{5/2} \cos^2(a + bx^2) dx$$

input

```
integrate(x**(5/2)*cos(b*x**2+a)**2,x)
```

output

```
Integral(x**(5/2)*cos(a + b*x**2)**2, x)
```

Maxima [F(-2)]

Exception generated.

$$\int x^{5/2} \cos^2(a + bx^2) dx = \text{Exception raised: RuntimeError}$$

input

```
integrate(x^(5/2)*cos(b*x^2+a)^2,x, algorithm="maxima")
```

output

```
Exception raised: RuntimeError >> Encountered operator mismatch in maxima-
to-sr translation
```

Giac [F]

$$\int x^{5/2} \cos^2(a + bx^2) dx = \int x^{5/2} \cos(bx^2 + a)^2 dx$$

input `integrate(x^(5/2)*cos(b*x^2+a)^2,x, algorithm="giac")`

output `integrate(x^(5/2)*cos(b*x^2 + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^{5/2} \cos^2(a + bx^2) dx = \int x^{5/2} \cos(bx^2 + a)^2 dx$$

input `int(x^(5/2)*cos(a + b*x^2)^2,x)`

output `int(x^(5/2)*cos(a + b*x^2)^2, x)`

Reduce [F]

$$\int x^{5/2} \cos^2(a + bx^2) dx = \frac{7\sqrt{x} \cos(bx^2 + a) \sin(bx^2 + a)x + 7\sqrt{x} \sin(bx^2 + a)x + 6\sqrt{x}bx^3 - 28 \left(\int \frac{\sqrt{x}x^2}{\tan\left(\frac{bx^2+a}{2}\right)^4 + 2\tan\left(\frac{bx^2+a}{2}\right)^2 + 1} dx \right)}{21b}$$

input `int(x^(5/2)*cos(b*x^2+a)^2,x)`

output `(7*sqrt(x)*cos(a + b*x**2)*sin(a + b*x**2)*x + 7*sqrt(x)*sin(a + b*x**2)*x + 6*sqrt(x)*b*x**3 - 28*int((sqrt(x)*x**2)/(tan((a + b*x**2)/2)**4 + 2*tan((a + b*x**2)/2)**2 + 1),x)*b - 42*int((sqrt(x)*tan((a + b*x**2)/2))/(tan((a + b*x**2)/2)**4 + 2*tan((a + b*x**2)/2)**2 + 1),x))/(21*b)`

3.30 $\int x^{3/2} \cos^2(a + bx^2) dx$

Optimal result	234
Mathematica [A] (verified)	234
Rubi [A] (verified)	235
Maple [F]	236
Fricas [A] (verification not implemented)	236
Sympy [F]	237
Maxima [F(-2)]	237
Giac [F]	238
Mupad [F(-1)]	238
Reduce [F]	238

Optimal result

Integrand size = 16, antiderivative size = 132

$$\int x^{3/2} \cos^2(a + bx^2) dx = \frac{x^{5/2}}{5} - \frac{ie^{2ia} \sqrt{x} \Gamma(\frac{1}{4}, -2ibx^2)}{64 \sqrt[4]{2b^4} \sqrt{-ibx^2}} + \frac{ie^{-2ia} \sqrt{x} \Gamma(\frac{1}{4}, 2ibx^2)}{64 \sqrt[4]{2b^4} \sqrt{ibx^2}} + \frac{\sqrt{x} \sin(2(a + bx^2))}{8b}$$

output

```
1/5*x^(5/2)-1/128*I*exp(2*I*a)*x^(1/2)*GAMMA(1/4,-2*I*b*x^2)*2^(3/4)/b/(-I
*b*x^2)^(1/4)+1/128*I*x^(1/2)*GAMMA(1/4,2*I*b*x^2)*2^(3/4)/b/exp(2*I*a)/(I
*b*x^2)^(1/4)+1/8*x^(1/2)*sin(2*b*x^2+2*a)/b
```

Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.08

$$\int x^{3/2} \cos^2(a + bx^2) dx = \frac{bx^{9/2} \left(5 \cdot 2^{3/4} \sqrt[4]{ibx^2} \Gamma(\frac{1}{4}, -2ibx^2) (-i \cos(2a) + \sin(2a)) + 5 \cdot 2^{3/4} \sqrt[4]{-ibx^2} \Gamma(\frac{1}{4}, 2ibx^2) (i \cos(2a) - \sin(2a)) \right) + 5x^{5/2} \sin(2(a + bx^2))}{640 (b^2 x^4)^{5/4}}$$

input

```
Integrate[x^(3/2)*Cos[a + b*x^2]^2,x]
```

output

```
(b*x^(9/2)*(5*2^(3/4)*(I*b*x^2)^(1/4)*Gamma[1/4, (-2*I)*b*x^2]*((-I)*Cos[2*a] + Sin[2*a]) + 5*2^(3/4)*((-I)*b*x^2)^(1/4)*Gamma[1/4, (2*I)*b*x^2]*(I*Cos[2*a] + Sin[2*a]) + 16*(b^2*x^4)^(1/4)*(8*b*x^2 + 5*Sin[2*(a + b*x^2)])))/(640*(b^2*x^4)^(5/4))
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3883, 3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{3/2} \cos^2(a + bx^2) dx$$

$$\downarrow \text{3883}$$

$$2 \int x^2 \cos^2(bx^2 + a) d\sqrt{x}$$

$$\downarrow \text{3885}$$

$$2 \int \left(\frac{1}{2} \cos(2bx^2 + 2a) x^2 + \frac{x^2}{2} \right) d\sqrt{x}$$

$$\downarrow \text{2009}$$

$$2 \left(\frac{\sqrt{x} \sin(2a + 2bx^2)}{16b} - \frac{ie^{2ia} \sqrt{x} \Gamma(\frac{1}{4}, -2ibx^2)}{128 \sqrt[4]{2b^4} \sqrt{-ibx^2}} + \frac{ie^{-2ia} \sqrt{x} \Gamma(\frac{1}{4}, 2ibx^2)}{128 \sqrt[4]{2b^4} \sqrt{ibx^2}} + \frac{x^{5/2}}{10} \right)$$

input

```
Int[x^(3/2)*Cos[a + b*x^2]^2,x]
```

output

```
2*(x^(5/2)/10 - ((I/128)*E^((2*I)*a)*Sqrt[x]*Gamma[1/4, (-2*I)*b*x^2])/(2^(1/4)*b*((-I)*b*x^2)^(1/4)) + ((I/128)*Sqrt[x]*Gamma[1/4, (2*I)*b*x^2])/(2^(1/4)*b*E^((2*I)*a)*(I*b*x^2)^(1/4)) + (Sqrt[x]*Sin[2*a + 2*b*x^2])/(16*b))
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3883 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := With[{k = Denominator[m]}, Simp[k/e Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*(x^(k*n)/e^n])]^p, x], x, (e*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [F]

$$\int x^{\frac{3}{2}} \cos(bx^2 + a)^2 dx$$

input `int(x^(3/2)*cos(b*x^2+a)^2,x)`

output `int(x^(3/2)*cos(b*x^2+a)^2,x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.69

$$\int x^{3/2} \cos^2(a + bx^2) dx = \frac{5(2ib)^{\frac{3}{4}} (\cos(2a) - i \sin(2a)) \Gamma(\frac{1}{4}, 2ibx^2) + 5(-2ib)^{\frac{3}{4}} (\cos(2a) + i \sin(2a)) \Gamma(\frac{1}{4}, -2ibx^2)}{640b^2}$$

input `integrate(x^(3/2)*cos(b*x^2+a)^2,x, algorithm="fricas")`

output

```
1/640*(5*(2*I*b)^(3/4)*(cos(2*a) - I*sin(2*a))*gamma(1/4, 2*I*b*x^2) + 5*(-2*I*b)^(3/4)*(cos(2*a) + I*sin(2*a))*gamma(1/4, -2*I*b*x^2) + 32*(4*b^2*x^2 + 5*b*cos(b*x^2 + a)*sin(b*x^2 + a))*sqrt(x))/b^2
```

Sympy [F]

$$\int x^{3/2} \cos^2(a + bx^2) dx = \int x^{\frac{3}{2}} \cos^2(a + bx^2) dx$$

input

```
integrate(x**(3/2)*cos(b*x**2+a)**2,x)
```

output

```
Integral(x**(3/2)*cos(a + b*x**2)**2, x)
```

Maxima [F(-2)]

Exception generated.

$$\int x^{3/2} \cos^2(a + bx^2) dx = \text{Exception raised: RuntimeError}$$

input

```
integrate(x^(3/2)*cos(b*x^2+a)^2,x, algorithm="maxima")
```

output

```
Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation
```

Giac [F]

$$\int x^{3/2} \cos^2(a + bx^2) dx = \int x^{3/2} \cos(bx^2 + a)^2 dx$$

input `integrate(x^(3/2)*cos(b*x^2+a)^2,x, algorithm="giac")`

output `integrate(x^(3/2)*cos(b*x^2 + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^{3/2} \cos^2(a + bx^2) dx = \int x^{3/2} \cos(bx^2 + a)^2 dx$$

input `int(x^(3/2)*cos(a + b*x^2)^2,x)`

output `int(x^(3/2)*cos(a + b*x^2)^2, x)`

Reduce [F]

$$\int x^{3/2} \cos^2(a + bx^2) dx = \frac{5\sqrt{x} \cos(bx^2 + a) \sin(bx^2 + a) + 5\sqrt{x} \sin(bx^2 + a) + 6\sqrt{x} bx^2 - 10 \left(\int \frac{\sqrt{x} \tan\left(\frac{bx^2}{2} + \frac{a}{2}\right)}{\tan\left(\frac{bx^2}{2} + \frac{a}{2}\right)^4 x + 2 \tan\left(\frac{bx^2}{2} + \frac{a}{2}\right)} dx \right)}{15b}$$

input `int(x^(3/2)*cos(b*x^2+a)^2,x)`

output `(5*sqrt(x)*cos(a + b*x**2)*sin(a + b*x**2) + 5*sqrt(x)*sin(a + b*x**2) + 6*sqrt(x)*b*x**2 - 10*int((sqrt(x)*tan((a + b*x**2)/2))/(tan((a + b*x**2)/2)**4*x + 2*tan((a + b*x**2)/2)**2*x + x),x) - 20*int((sqrt(x)*x)/(tan((a + b*x**2)/2)**4 + 2*tan((a + b*x**2)/2)**2 + 1),x)*b)/(15*b)`

3.31 $\int \sqrt{x} \cos^2(a + bx^2) dx$

Optimal result	239
Mathematica [A] (verified)	239
Rubi [A] (verified)	240
Maple [F]	241
Fricas [A] (verification not implemented)	241
Sympy [F]	242
Maxima [F(-2)]	242
Giac [F]	243
Mupad [F(-1)]	243
Reduce [F]	243

Optimal result

Integrand size = 16, antiderivative size = 100

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \frac{x^{3/2}}{3} - \frac{e^{2ia} x^{3/2} \Gamma(\frac{3}{4}, -2ibx^2)}{8 \cdot 2^{3/4} (-ibx^2)^{3/4}} - \frac{e^{-2ia} x^{3/2} \Gamma(\frac{3}{4}, 2ibx^2)}{8 \cdot 2^{3/4} (ibx^2)^{3/4}}$$

output

```
1/3*x^(3/2)-1/16*exp(2*I*a)*x^(3/2)*GAMMA(3/4,-2*I*b*x^2)*2^(1/4)/(-I*b*x^2)^(3/4)-1/16*x^(3/2)*GAMMA(3/4,2*I*b*x^2)*2^(1/4)/exp(2*I*a)/(I*b*x^2)^(3/4)
```

Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.99

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \frac{1}{48} x^{3/2} \left(16 - \frac{3\sqrt[4]{2} \Gamma(\frac{3}{4}, 2ibx^2) (\cos(2a) - i \sin(2a))}{(ibx^2)^{3/4}} - \frac{3\sqrt[4]{2} \Gamma(\frac{3}{4}, -2ibx^2) (\cos(2a) + i \sin(2a))}{(-ibx^2)^{3/4}} \right)$$

input

```
Integrate[Sqrt[x]*Cos[a + b*x^2]^2,x]
```


output

$$\frac{(x^{3/2}*(16 - (3*2^{1/4})*\Gamma[3/4, (2*I)*b*x^2]*(\cos[2*a] - I*\sin[2*a]))}{(I*b*x^2)^{3/4} - (3*2^{1/4})*\Gamma[3/4, (-2*I)*b*x^2]*(\cos[2*a] + I*\sin[2*a])})/((-I)*b*x^2)^{3/4}}{48}$$
Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3883, 3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{x} \cos^2(a + bx^2) dx \\ & \quad \downarrow \text{3883} \\ & 2 \int x \cos^2(bx^2 + a) d\sqrt{x} \\ & \quad \downarrow \text{3885} \\ & 2 \int \left(\frac{1}{2} \cos(2bx^2 + 2a) x + \frac{x}{2} \right) d\sqrt{x} \\ & \quad \downarrow \text{2009} \\ & 2 \left(-\frac{e^{2ia} x^{3/2} \Gamma\left(\frac{3}{4}, -2ibx^2\right)}{16 \cdot 2^{3/4} (-ibx^2)^{3/4}} - \frac{e^{-2ia} x^{3/2} \Gamma\left(\frac{3}{4}, 2ibx^2\right)}{16 \cdot 2^{3/4} (ibx^2)^{3/4}} + \frac{x^{3/2}}{6} \right) \end{aligned}$$

input

$$\text{Int}[\text{Sqrt}[x]*\text{Cos}[a + b*x^2]^2, x]$$

output

$$2*(x^{3/2}/6 - (E^{((2*I)*a)}*x^{3/2}*\Gamma[3/4, (-2*I)*b*x^2])/(16*2^{3/4}*(-I)*b*x^2)^{3/4} - (x^{3/2}*\Gamma[3/4, (2*I)*b*x^2])/(16*2^{3/4}*E^{((2*I)*a)}*(I*b*x^2)^{3/4}))$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3883 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := With[{k = Denominator[m]}, Simp[k/e Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*(x^(k*n)/e^n]])^p, x], x, (e*x)^(1/k)], x]] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [F]

$$\int \sqrt{x} \cos(bx^2 + a)^2 dx$$

input `int(x^(1/2)*cos(b*x^2+a)^2,x)`

output `int(x^(1/2)*cos(b*x^2+a)^2,x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.68

$$\int \sqrt{x} \cos^2(a + bx^2) dx$$

$$= \frac{16bx^{\frac{3}{2}} - 3(2ib)^{\frac{1}{4}}(-i \cos(2a) - \sin(2a))\Gamma(\frac{3}{4}, 2ibx^2) - 3(-2ib)^{\frac{1}{4}}(i \cos(2a) - \sin(2a))\Gamma(\frac{3}{4}, -2ibx^2)}{48b}$$

input `integrate(x^(1/2)*cos(b*x^2+a)^2,x, algorithm="fricas")`

output
$$\frac{1}{48} \cdot (16 \cdot b \cdot x^{3/2} - 3 \cdot (2 \cdot I \cdot b)^{1/4} \cdot (-I \cdot \cos(2 \cdot a) - \sin(2 \cdot a)) \cdot \text{gamma}(3/4, 2 \cdot I \cdot b \cdot x^2) - 3 \cdot (-2 \cdot I \cdot b)^{1/4} \cdot (I \cdot \cos(2 \cdot a) - \sin(2 \cdot a)) \cdot \text{gamma}(3/4, -2 \cdot I \cdot b \cdot x^2)) / b$$

Sympy [F]

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \int \sqrt{x} \cos^2(a + bx^2) dx$$

input `integrate(x**(1/2)*cos(b*x**2+a)**2,x)`

output `Integral(sqrt(x)*cos(a + b*x**2)**2, x)`

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^(1/2)*cos(b*x^2+a)^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a)^2 dx$$

input `integrate(x^(1/2)*cos(b*x^2+a)^2,x, algorithm="giac")`

output `integrate(sqrt(x)*cos(b*x^2 + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a)^2 dx$$

input `int(x^(1/2)*cos(a + b*x^2)^2,x)`

output `int(x^(1/2)*cos(a + b*x^2)^2, x)`

Reduce [F]

$$\int \sqrt{x} \cos^2(a + bx^2) dx = \int \sqrt{x} \cos(bx^2 + a)^2 dx$$

input `int(x^(1/2)*cos(b*x^2+a)^2,x)`

output `int(sqrt(x)*cos(a + b*x**2)**2,x)`

3.32 $\int \frac{\cos^2(a+bx^2)}{\sqrt{x}} dx$

Optimal result	244
Mathematica [A] (verified)	244
Rubi [A] (verified)	245
Maple [F]	246
Fricas [A] (verification not implemented)	246
Sympy [F]	247
Maxima [F(-2)]	247
Giac [F]	247
Mupad [F(-1)]	248
Reduce [F]	248

Optimal result

Integrand size = 16, antiderivative size = 96

$$\int \frac{\cos^2(a+bx^2)}{\sqrt{x}} dx = \sqrt{x} - \frac{e^{2ia}\sqrt{x}\Gamma(\frac{1}{4}, -2ibx^2)}{8\sqrt[4]{2}\sqrt[4]{-ibx^2}} - \frac{e^{-2ia}\sqrt{x}\Gamma(\frac{1}{4}, 2ibx^2)}{8\sqrt[4]{2}\sqrt[4]{ibx^2}}$$

output

```
x^(1/2)-1/16*exp(2*I*a)*x^(1/2)*GAMMA(1/4,-2*I*b*x^2)*2^(3/4)/(-I*b*x^2)^(1/4)-1/16*x^(1/2)*GAMMA(1/4,2*I*b*x^2)*2^(3/4)/exp(2*I*a)/(I*b*x^2)^(1/4)
```

Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.03

$$\int \frac{\cos^2(a+bx^2)}{\sqrt{x}} dx = \frac{1}{16}\sqrt{x} \left(16 - \frac{2^{3/4}\Gamma(\frac{1}{4}, 2ibx^2) (\cos(2a) - i \sin(2a))}{\sqrt[4]{ibx^2}} - \frac{2^{3/4}\Gamma(\frac{1}{4}, -2ibx^2) (\cos(2a) + i \sin(2a))}{\sqrt[4]{-ibx^2}} \right)$$

input

```
Integrate[Cos[a + b*x^2]^2/Sqrt[x], x]
```

output

```
(Sqrt[x]*(16 - (2^(3/4)*Gamma[1/4, (2*I)*b*x^2]*(Cos[2*a] - I*Sin[2*a]))/(
I*b*x^2)^(1/4) - (2^(3/4)*Gamma[1/4, (-2*I)*b*x^2]*(Cos[2*a] + I*Sin[2*a])
)/((-I)*b*x^2)^(1/4))/16
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.06, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3883, 3839, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx$$

↓ 3883

$$2 \int \cos^2(bx^2 + a) d\sqrt{x}$$

↓ 3839

$$2 \int \left(\frac{1}{2} \cos(2bx^2 + 2a) + \frac{1}{2} \right) d\sqrt{x}$$

↓ 2009

$$2 \left(-\frac{e^{2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, -2ibx^2\right)}{16\sqrt[4]{2}\sqrt[4]{-ibx^2}} - \frac{e^{-2ia} \sqrt{x} \Gamma\left(\frac{1}{4}, 2ibx^2\right)}{16\sqrt[4]{2}\sqrt[4]{ibx^2}} + \frac{\sqrt{x}}{2} \right)$$

input

```
Int[Cos[a + b*x^2]^2/Sqrt[x],x]
```

output

```
2*(Sqrt[x]/2 - (E^((2*I)*a)*Sqrt[x]*Gamma[1/4, (-2*I)*b*x^2])/(16*2^(1/4)*
((-I)*b*x^2)^(1/4)) - (Sqrt[x]*Gamma[1/4, (2*I)*b*x^2])/(16*2^(1/4)*E^((2*
I)*a)*(I*b*x^2)^(1/4)))
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3839 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_), x_Symbol] := Int[ExpandTrigReduce[(a + b*Cos[c + d*(e + f*x)^n]]^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 1] && IGtQ[n, 1]`

rule 3883 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.)*(x_))^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/e Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*(x^(k*n)/e^n]]^p, x], x, (e*x)^(1/k)], x]] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]`

Maple [F]

$$\int \frac{\cos(bx^2 + a)^2}{\sqrt{x}} dx$$

input `int(cos(b*x^2+a)^2/x^(1/2),x)`

output `int(cos(b*x^2+a)^2/x^(1/2),x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.65

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx$$

$$= \frac{(2ib)^{\frac{3}{4}} (i \cos(2a) + \sin(2a)) \Gamma(\frac{1}{4}, 2ibx^2) + (-2ib)^{\frac{3}{4}} (-i \cos(2a) + \sin(2a)) \Gamma(\frac{1}{4}, -2ibx^2) + 16b\sqrt{x}}{16b}$$

input `integrate(cos(b*x^2+a)^2/x^(1/2),x, algorithm="fricas")`

output $1/16*((2*I*b)^{(3/4)}*(I*\cos(2*a) + \sin(2*a))*\text{gamma}(1/4, 2*I*b*x^2) + (-2*I*b)^{(3/4)}*(-I*\cos(2*a) + \sin(2*a))*\text{gamma}(1/4, -2*I*b*x^2) + 16*b*\text{sqrt}(x))/b$

Sympy [F]

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx$$

input `integrate(cos(b*x**2+a)**2/x**(1/2), x)`

output `Integral(cos(a + b*x**2)**2/sqrt(x), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cos(b*x^2+a)^2/x^(1/2), x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(bx^2 + a)^2}{\sqrt{x}} dx$$

input `integrate(cos(b*x^2+a)^2/x^(1/2), x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)^2/sqrt(x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(bx^2 + a)^2}{\sqrt{x}} dx$$

input `int(cos(a + b*x^2)^2/x^(1/2), x)`

output `int(cos(a + b*x^2)^2/x^(1/2), x)`

Reduce [F]

$$\int \frac{\cos^2(a + bx^2)}{\sqrt{x}} dx = \int \frac{\cos(bx^2 + a)^2}{\sqrt{x}} dx$$

input `int(cos(b*x^2+a)^2/x^(1/2), x)`

output `int(cos(a + b*x**2)**2/sqrt(x), x)`

3.33 $\int \frac{\cos^2(a+bx^2)}{x^{3/2}} dx$

Optimal result	249
Mathematica [A] (verified)	249
Rubi [A] (verified)	250
Maple [F]	251
Fricas [A] (verification not implemented)	251
Sympy [F]	252
Maxima [F(-2)]	252
Giac [F]	253
Mupad [F(-1)]	253
Reduce [F]	253

Optimal result

Integrand size = 16, antiderivative size = 117

$$\int \frac{\cos^2(a+bx^2)}{x^{3/2}} dx = -\frac{1}{\sqrt{x}} - \frac{\cos(2(a+bx^2))}{\sqrt{x}} - \frac{ibe^{2ia}x^{3/2}\Gamma(\frac{3}{4}, -2ibx^2)}{2^{3/4}(-ibx^2)^{3/4}} + \frac{ibe^{-2ia}x^{3/2}\Gamma(\frac{3}{4}, 2ibx^2)}{2^{3/4}(ibx^2)^{3/4}}$$

output

```
-1/x^(1/2)-cos(2*b*x^2+2*a)/x^(1/2)-1/2*I*b*exp(2*I*a)*x^(3/2)*GAMMA(3/4,-2*I*b*x^2)*2^(1/4)/(-I*b*x^2)^(3/4)+1/2*I*b*x^(3/2)*GAMMA(3/4,2*I*b*x^2)*2^(1/4)/exp(2*I*a)/(I*b*x^2)^(3/4)
```

Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.17

$$\int \frac{\cos^2(a+bx^2)}{x^{3/2}} dx = \frac{-4(b^2x^4)^{3/4} \cos^2(a+bx^2) + \sqrt[4]{2}bx^2(ibx^2)^{3/4} \Gamma(\frac{3}{4}, -2ibx^2) (-i \cos(2a) + \sin(2a)) + 2\sqrt{x} (b^2x^4)^{3/4}}{2\sqrt{x} (b^2x^4)^{3/4}}$$

input

```
Integrate[Cos[a + b*x^2]^2/x^(3/2), x]
```

output

$$(-4*(b^2*x^4)^{(3/4)}*\text{Cos}[a + b*x^2]^2 + 2^{(1/4)}*b*x^2*(I*b*x^2)^{(3/4)}*\text{Gamma}[3/4, (-2*I)*b*x^2]*((-I)*\text{Cos}[2*a] + \text{Sin}[2*a]) + I*2^{(1/4)}*((-I)*b*x^2)^{(7/4)}*\text{Gamma}[3/4, (2*I)*b*x^2]*(I*\text{Cos}[2*a] + \text{Sin}[2*a]))/(2*\text{Sqrt}[x]*(b^2*x^4)^{(3/4)})$$
Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3883, 3885, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx \\ & \quad \downarrow \text{3883} \\ & 2 \int \frac{\cos^2(bx^2 + a)}{x} d\sqrt{x} \\ & \quad \downarrow \text{3885} \\ & 2 \int \left(\frac{\cos(2bx^2 + 2a)}{2x} + \frac{1}{2x} \right) d\sqrt{x} \\ & \quad \downarrow \text{2009} \\ & 2 \left(-\frac{\cos(2a + 2bx^2)}{2\sqrt{x}} - \frac{ie^{2ia}bx^{3/2}\Gamma(\frac{3}{4}, -2ibx^2)}{2 \cdot 2^{3/4}(-ibx^2)^{3/4}} + \frac{ie^{-2ia}bx^{3/2}\Gamma(\frac{3}{4}, 2ibx^2)}{2 \cdot 2^{3/4}(ibx^2)^{3/4}} - \frac{1}{2\sqrt{x}} \right) \end{aligned}$$

input

$$\text{Int}[\text{Cos}[a + b*x^2]^2/x^{(3/2)}, x]$$

output

$$2*(-1/2*1/\text{Sqrt}[x] - \text{Cos}[2*a + 2*b*x^2]/(2*\text{Sqrt}[x])) - ((I/2)*b*E^{((2*I)*a)}*x^{(3/2)}*\text{Gamma}[3/4, (-2*I)*b*x^2])/(2^{(3/4)}*((-I)*b*x^2)^{(3/4)}) + ((I/2)*b*x^{(3/2)}*\text{Gamma}[3/4, (2*I)*b*x^2])/(2^{(3/4)}*E^{((2*I)*a)}*(I*b*x^2)^{(3/4)})$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3883 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := With[{k = Denominator[m]}, Simp[k/e Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*(x^(k*n)/e^n]])^p, x], x, (e*x)^(1/k)], x]] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]`

rule 3885 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 1] && IGtQ[n, 0]`

Maple [F]

$$\int \frac{\cos(bx^2 + a)^2}{x^{\frac{3}{2}}} dx$$

input `int(cos(b*x^2+a)^2/x^(3/2),x)`

output `int(cos(b*x^2+a)^2/x^(3/2),x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.68

$$\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx = \frac{4\sqrt{x} \cos(bx^2 + a)^2 - (x \cos(2a) - ix \sin(2a))(2ib)^{\frac{1}{4}} \Gamma\left(\frac{3}{4}, 2ibx^2\right) - (x \cos(2a) + ix \sin(2a))(-2ib)^{\frac{1}{4}}}{2x}$$

input `integrate(cos(b*x^2+a)^2/x^(3/2),x, algorithm="fricas")`

output

```
-1/2*(4*sqrt(x)*cos(b*x^2 + a)^2 - (x*cos(2*a) - I*x*sin(2*a))*(2*I*b)^(1/4)*gamma(3/4, 2*I*b*x^2) - (x*cos(2*a) + I*x*sin(2*a))*(-2*I*b)^(1/4)*gamma(3/4, -2*I*b*x^2))/x
```

Sympy [F]

$$\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos^2(a + bx^2)}{x^{\frac{3}{2}}} dx$$

input

```
integrate(cos(b*x**2+a)**2/x**(3/2), x)
```

output

```
Integral(cos(a + b*x**2)**2/x**(3/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input

```
integrate(cos(b*x^2+a)^2/x^(3/2), x, algorithm="maxima")
```

output

```
Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation
```

Giac [F]

$$\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos(bx^2 + a)^2}{x^{3/2}} dx$$

input `integrate(cos(b*x^2+a)^2/x^(3/2),x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)^2/x^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos(bx^2 + a)^2}{x^{3/2}} dx$$

input `int(cos(a + b*x^2)^2/x^(3/2),x)`

output `int(cos(a + b*x^2)^2/x^(3/2), x)`

Reduce [F]

$$\int \frac{\cos^2(a + bx^2)}{x^{3/2}} dx = \int \frac{\cos(bx^2 + a)^2}{\sqrt{x} x} dx$$

input `int(cos(b*x^2+a)^2/x^(3/2),x)`

output `int(cos(a + b*x**2)**2/(sqrt(x)*x), x)`

3.34 $\int \frac{\cos^2(a+bx^2)}{x^{5/2}} dx$

Optimal result	254
Mathematica [A] (verified)	254
Rubi [A] (verified)	255
Maple [F]	257
Fricas [A] (verification not implemented)	257
Sympy [F]	257
Maxima [F(-2)]	258
Giac [F]	258
Mupad [F(-1)]	258
Reduce [F]	259

Optimal result

Integrand size = 16, antiderivative size = 116

$$\int \frac{\cos^2(a+bx^2)}{x^{5/2}} dx = -\frac{2 \cos^2(a+bx^2)}{3x^{3/2}} - \frac{ibe^{2ia}\sqrt{x}\Gamma(\frac{1}{4}, -2ibx^2)}{3\sqrt[4]{2}\sqrt[4]{-ibx^2}} + \frac{ibe^{-2ia}\sqrt{x}\Gamma(\frac{1}{4}, 2ibx^2)}{3\sqrt[4]{2}\sqrt[4]{ibx^2}}$$

output

```
-2/3*cos(b*x^2+a)^2/x^(3/2)-1/6*I*b*exp(2*I*a)*x^(1/2)*GAMMA(1/4,-2*I*b*x^2)*2^(3/4)/(-I*b*x^2)^(1/4)+1/6*I*b*x^(1/2)*GAMMA(1/4,2*I*b*x^2)*2^(3/4)/exp(2*I*a)/(I*b*x^2)^(1/4)
```

Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.18

$$\int \frac{\cos^2(a+bx^2)}{x^{5/2}} dx = \frac{-4\sqrt[4]{b^2x^4} \cos^2(a+bx^2) + 2^{3/4}bx^2\sqrt[4]{ibx^2}\Gamma(\frac{1}{4}, -2ibx^2) (-i \cos(2a) + \sin(2a)) + i2^{3/4}}{6x^{3/2}\sqrt[4]{b^2x^4}}$$

input

```
Integrate[Cos[a + b*x^2]^2/x^(5/2), x]
```

output

$$\left(-4*(b^2*x^4)^{(1/4)}*\text{Cos}[a + b*x^2]^2 + 2^{(3/4)}*b*x^2*(I*b*x^2)^{(1/4)}*\text{Gamma}[1/4, (-2*I)*b*x^2]*((-I)*\text{Cos}[2*a] + \text{Sin}[2*a]) + I*2^{(3/4)}*((-I)*b*x^2)^{(5/4)}*\text{Gamma}[1/4, (2*I)*b*x^2]*(I*\text{Cos}[2*a] + \text{Sin}[2*a])\right)/(6*x^{(3/2)}*(b^2*x^4)^{(1/4)})$$
Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3883, 3875, 5084, 3854, 3836, 2637}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx \\ & \quad \downarrow \text{3883} \\ & 2 \int \frac{\cos^2(bx^2 + a)}{x^2} d\sqrt{x} \\ & \quad \downarrow \text{3875} \\ & 2 \left(-\frac{8}{3}b \int \cos(bx^2 + a) \sin(bx^2 + a) d\sqrt{x} - \frac{\cos^2(a + bx^2)}{3x^{3/2}} \right) \\ & \quad \downarrow \text{5084} \\ & 2 \left(-\frac{4}{3}b \int \sin(2(bx^2 + a)) d\sqrt{x} - \frac{\cos^2(a + bx^2)}{3x^{3/2}} \right) \\ & \quad \downarrow \text{3854} \\ & 2 \left(-\frac{4}{3}b \int \sin(2bx^2 + 2a) d\sqrt{x} - \frac{\cos^2(a + bx^2)}{3x^{3/2}} \right) \\ & \quad \downarrow \text{3836} \\ & 2 \left(-\frac{\cos^2(a + bx^2)}{3x^{3/2}} - \frac{4}{3}b \left(\frac{1}{2}i \int e^{-2ibx^2 - 2ia} d\sqrt{x} - \frac{1}{2}i \int e^{2ibx^2 + 2ia} d\sqrt{x} \right) \right) \\ & \quad \downarrow \text{2637} \end{aligned}$$

$$2 \left(-\frac{\cos^2(a + bx^2)}{3x^{3/2}} - \frac{4}{3} b \left(\frac{ie^{2ia} \sqrt{x} \Gamma(\frac{1}{4}, -2ibx^2)}{8\sqrt{2}\sqrt[4]{-ibx^2}} - \frac{ie^{-2ia} \sqrt{x} \Gamma(\frac{1}{4}, 2ibx^2)}{8\sqrt{2}\sqrt[4]{ibx^2}} \right) \right)$$

input `Int[Cos[a + b*x^2]^2/x^(5/2),x]`

output `2*(-1/3*Cos[a + b*x^2]^2/x^(3/2) - (4*b*(((I/8)*E^((2*I)*a)*Sqrt[x]*Gamma[1/4, (-2*I)*b*x^2])/(2^(1/4)*((-I)*b*x^2)^(1/4)) - ((I/8)*Sqrt[x]*Gamma[1/4, (2*I)*b*x^2])/(2^(1/4)*E^((2*I)*a)*(I*b*x^2)^(1/4))))/3`

Defintions of rubi rules used

rule 2637 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(-F^a)*(c + d*x)*(Gamma[1/n, (-b)*(c + d*x)^n*Log[F]]/(d*n*((-b)*(c + d*x)^n*Log[F])^(1/n))), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]`

rule 3836 `Int[Sin[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_))], x_Symbol] := Simp[I/2 Int[E^((-c)*I - d*I*(e + f*x)^n), x], x] - Simp[I/2 Int[E^(c*I + d*I*(e + f*x)^n), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[n, 2]`

rule 3854 `Int[((a_.) + (b_.)*Sin[u_])^(p_.), x_Symbol] := Int[(a + b*Sin[ExpandToSum[u, x]])^p, x] /; FreeQ[{a, b, p}, x] && BinomialQ[u, x] && !BinomialMatchQ[u, x]`

rule 3875 `Int[Cos[(a_.) + (b_.)*(x_)^(n_)]^(p_)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*(Cos[a + b*x^n]^p/(m + 1)), x] + Simp[b*n*(p/(m + 1)) Int[Cos[a + b*x^n]^(p - 1)*Sin[a + b*x^n], x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 1] && EqQ[m + n, 0] && NeQ[n, 1] && IntegerQ[n]`

rule 3883 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := With[{k = Denominator[m]}, Simp[k/e Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*(x^(k*n))/e^n]]^p, x], x, (e*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[p] && IGtQ[n, 0] && FractionQ[m]`

rule 5084

```
Int[Cos[w_]^(p_.)*(u_.)*Sin[v_]^(p_.), x_Symbol] := Simp[1/2^p Int[u*Sin[
2*v]^(p, x), x] /; EqQ[w, v] && IntegerQ[p]
```

Maple [F]

$$\int \frac{\cos(bx^2 + a)^2}{x^{5/2}} dx$$

input

```
int(cos(b*x^2+a)^2/x^(5/2),x)
```

output

```
int(cos(b*x^2+a)^2/x^(5/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.73

$$\int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx = \frac{(x^2 \cos(2a) - i x^2 \sin(2a))(2i b)^{3/4} \Gamma(\frac{1}{4}, 2i bx^2) + (x^2 \cos(2a) + i x^2 \sin(2a))(-2i b)^{3/4} \Gamma(\frac{1}{4}, -2i bx^2)}{6 x^2}$$

input

```
integrate(cos(b*x^2+a)^2/x^(5/2),x, algorithm="fricas")
```

output

```
1/6*((x^2*cos(2*a) - I*x^2*sin(2*a))*(2*I*b)^(3/4)*gamma(1/4, 2*I*b*x^2) +
(x^2*cos(2*a) + I*x^2*sin(2*a))*(-2*I*b)^(3/4)*gamma(1/4, -2*I*b*x^2) - 4
*sqrt(x)*cos(b*x^2 + a)^2)/x^2
```

Sympy [F]

$$\int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx$$

input

```
integrate(cos(b*x**2+a)**2/x**(5/2),x)
```

output `Integral(cos(a + b*x**2)**2/x**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cos(b*x^2+a)^2/x^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> Encountered operator mismatch in maxima-to-sr translation`

Giac [F]

$$\int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos(bx^2 + a)^2}{x^{5/2}} dx$$

input `integrate(cos(b*x^2+a)^2/x^(5/2),x, algorithm="giac")`

output `integrate(cos(b*x^2 + a)^2/x^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos(bx^2 + a)^2}{x^{5/2}} dx$$

input `int(cos(a + b*x^2)^2/x^(5/2),x)`

output `int(cos(a + b*x^2)^2/x^(5/2), x)`

Reduce [F]

$$\int \frac{\cos^2(a + bx^2)}{x^{5/2}} dx = \int \frac{\cos(bx^2 + a)^2}{\sqrt{x} x^2} dx$$

input `int(cos(b*x^2+a)^2/x^(5/2),x)`

output `int(cos(a + b*x**2)**2/(sqrt(x)*x**2),x)`

3.35 $\int \cos\left(a + \frac{b}{x}\right) dx$

Optimal result	260
Mathematica [A] (verified)	260
Rubi [A] (verified)	261
Maple [A] (verified)	263
Fricas [A] (verification not implemented)	264
Sympy [F]	264
Maxima [C] (verification not implemented)	264
Giac [B] (verification not implemented)	265
Mupad [F(-1)]	265
Reduce [F]	266

Optimal result

Integrand size = 8, antiderivative size = 31

$$\int \cos\left(a + \frac{b}{x}\right) dx = x \cos\left(a + \frac{b}{x}\right) + b \operatorname{CosIntegral}\left(\frac{b}{x}\right) \sin(a) + b \cos(a) \operatorname{Si}\left(\frac{b}{x}\right)$$

output

```
x*cos(a+b/x)+b*Ci(b/x)*sin(a)+b*cos(a)*Si(b/x)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \cos\left(a + \frac{b}{x}\right) dx = x \cos\left(a + \frac{b}{x}\right) + b \operatorname{CosIntegral}\left(\frac{b}{x}\right) \sin(a) + b \cos(a) \operatorname{Si}\left(\frac{b}{x}\right)$$

input

```
Integrate[Cos[a + b/x],x]
```

output

```
x*Cos[a + b/x] + b*CosIntegral[b/x]*Sin[a] + b*Cos[a]*SinIntegral[b/x]
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$, Rules used = {3843, 3042, 3778, 25, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos\left(a + \frac{b}{x}\right) dx \\
 & \quad \downarrow \text{3843} \\
 & - \int x^2 \cos\left(a + \frac{b}{x}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{3042} \\
 & - \int x^2 \sin\left(a + \frac{b}{x} + \frac{\pi}{2}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{3778} \\
 & x \cos\left(a + \frac{b}{x}\right) - b \int -x \sin\left(a + \frac{b}{x}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{25} \\
 & b \int x \sin\left(a + \frac{b}{x}\right) d\frac{1}{x} + x \cos\left(a + \frac{b}{x}\right) \\
 & \quad \downarrow \text{3042} \\
 & b \int x \sin\left(a + \frac{b}{x}\right) d\frac{1}{x} + x \cos\left(a + \frac{b}{x}\right) \\
 & \quad \downarrow \text{3784} \\
 & b \left(\sin(a) \int x \cos\left(\frac{b}{x}\right) d\frac{1}{x} + \cos(a) \int x \sin\left(\frac{b}{x}\right) d\frac{1}{x} \right) + x \cos\left(a + \frac{b}{x}\right) \\
 & \quad \downarrow \text{3042} \\
 & b \left(\sin(a) \int x \sin\left(\frac{b}{x} + \frac{\pi}{2}\right) d\frac{1}{x} + \cos(a) \int x \sin\left(\frac{b}{x}\right) d\frac{1}{x} \right) + x \cos\left(a + \frac{b}{x}\right) \\
 & \quad \downarrow \text{3780}
 \end{aligned}$$

$$b \left(\sin(a) \int x \sin \left(\frac{b}{x} + \frac{\pi}{2} \right) d\frac{1}{x} + \cos(a) \text{Si} \left(\frac{b}{x} \right) \right) + x \cos \left(a + \frac{b}{x} \right)$$

↓ 3783

$$b \left(\sin(a) \text{CosIntegral} \left(\frac{b}{x} \right) + \cos(a) \text{Si} \left(\frac{b}{x} \right) \right) + x \cos \left(a + \frac{b}{x} \right)$$

input `Int[Cos[a + b/x], x]`

output `x*Cos[a + b/x] + b*(CosIntegral[b/x]*Sin[a] + Cos[a]*SinIntegral[b/x])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3780 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

```
rule 3784 Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*
e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*
f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x]
&& NeQ[d*e - c*f, 0]
```

```
rule 3843 Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.), x_S
ymbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x])^p, x],
x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && Intege
rQ[1/n]
```

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.26

method	result
derivativedivides	$-b \left(-\frac{\cos\left(a+\frac{b}{x}\right)x}{b} - \cos(a) \operatorname{Si}\left(\frac{b}{x}\right) - \operatorname{Ci}\left(\frac{b}{x}\right) \sin(a) \right)$
default	$-b \left(-\frac{\cos\left(a+\frac{b}{x}\right)x}{b} - \cos(a) \operatorname{Si}\left(\frac{b}{x}\right) - \operatorname{Ci}\left(\frac{b}{x}\right) \sin(a) \right)$
risch	$\frac{ib \operatorname{expIntegral}_1\left(-\frac{ib}{x}\right)e^{ia}}{2} - \frac{\pi \operatorname{csgn}\left(\frac{b}{x}\right)e^{-iab}}{2} + \operatorname{Si}\left(\frac{b}{x}\right)e^{-iab} - \frac{i \operatorname{expIntegral}_1\left(-\frac{ib}{x}\right)e^{-iab}}{2} + x \cos\left(\frac{ax+b}{x}\right)$
meijerg	$-\frac{\cos(a)\sqrt{\pi}\sqrt{b^2}\left(-\frac{4xb^2\cos\left(\frac{\sqrt{b^2}}{x}\right)}{(b^2)^{\frac{3}{2}}\sqrt{\pi}} - \frac{4\operatorname{Si}\left(\frac{\sqrt{b^2}}{x}\right)}{\sqrt{\pi}}\right)}{4} + \frac{\sin(a)\sqrt{\pi}b\left(\frac{4\gamma-4-4\ln(x)+4\ln(b)}{\sqrt{\pi}} + \frac{4}{\sqrt{\pi}} - \frac{4\gamma}{\sqrt{\pi}} - \frac{4\ln(2)}{\sqrt{\pi}} - \frac{4\ln\left(\frac{b}{2x}\right)}{\sqrt{\pi}}\right)}{4}$

```
input int(cos(a+b/x),x,method=_RETURNVERBOSE)
```

```
output -b*(-cos(a+b/x)/b*x-cos(a)*Si(b/x)-Ci(b/x)*sin(a))
```


Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \cos\left(a + \frac{b}{x}\right) dx = b \operatorname{Ci}\left(\frac{b}{x}\right) \sin(a) + b \cos(a) \operatorname{Si}\left(\frac{b}{x}\right) + x \cos\left(\frac{ax + b}{x}\right)$$

input `integrate(cos(a+b/x),x, algorithm="fricas")`

output `b*cos_integral(b/x)*sin(a) + b*cos(a)*sin_integral(b/x) + x*cos((a*x + b)/x)`

Sympy [F]

$$\int \cos\left(a + \frac{b}{x}\right) dx = \int \cos\left(a + \frac{b}{x}\right) dx$$

input `integrate(cos(a+b/x),x)`

output `Integral(cos(a + b/x), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.84

$$\begin{aligned} & \int \cos\left(a + \frac{b}{x}\right) dx \\ &= \frac{1}{2} \left(\left(-i \operatorname{Ei}\left(\frac{ib}{x}\right) + i \operatorname{Ei}\left(-\frac{ib}{x}\right) \right) \cos(a) + \left(\operatorname{Ei}\left(\frac{ib}{x}\right) + \operatorname{Ei}\left(-\frac{ib}{x}\right) \right) \sin(a) \right) b \\ & \quad + x \cos\left(\frac{ax + b}{x}\right) \end{aligned}$$

input `integrate(cos(a+b/x),x, algorithm="maxima")`

output `1/2*((-I*Ei(I*b/x) + I*Ei(-I*b/x))*cos(a) + (Ei(I*b/x) + Ei(-I*b/x))*sin(a))*b + x*cos((a*x + b)/x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(31) = 62$.

Time = 0.38 (sec) , antiderivative size = 132, normalized size of antiderivative = 4.26

$$\int \cos\left(a + \frac{b}{x}\right) dx$$

$$= \frac{ab^2 \operatorname{Ci}\left(-a + \frac{ax+b}{x}\right) \sin(a) - ab^2 \cos(a) \operatorname{Si}\left(a - \frac{ax+b}{x}\right) - \frac{(ax+b)b^2 \operatorname{Ci}\left(-a + \frac{ax+b}{x}\right) \sin(a)}{x} + \frac{(ax+b)b^2 \cos(a) \operatorname{Si}\left(a - \frac{ax+b}{x}\right)}{x}}{\left(a - \frac{ax+b}{x}\right)b}$$

input `integrate(cos(a+b/x),x, algorithm="giac")`

output `(a*b^2*cos_integral(-a + (a*x + b)/x)*sin(a) - a*b^2*cos(a)*sin_integral(a - (a*x + b)/x) - (a*x + b)*b^2*cos_integral(-a + (a*x + b)/x)*sin(a)/x + (a*x + b)*b^2*cos(a)*sin_integral(a - (a*x + b)/x)/x - b^2*cos((a*x + b)/x))/((a - (a*x + b)/x)*b)`

Mupad [F(-1)]

Timed out.

$$\int \cos\left(a + \frac{b}{x}\right) dx = \int \cos\left(a + \frac{b}{x}\right) dx$$

input `int(cos(a + b/x),x)`

output `int(cos(a + b/x), x)`

Reduce [F]

$$\int \cos\left(a + \frac{b}{x}\right) dx = \int \cos\left(\frac{ax + b}{x}\right) dx + \int 1 dx - x$$

input `int(cos(a+b/x),x)`

output `int(cos((a*x + b)/x),x) + int(1,x) - x`

$$3.36 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$$

Optimal result	267
Mathematica [A] (verified)	267
Rubi [A] (verified)	268
Maple [A] (verified)	269
Fricas [A] (verification not implemented)	269
Sympy [A] (verification not implemented)	270
Maxima [C] (verification not implemented)	270
Giac [B] (verification not implemented)	270
Mupad [F(-1)]	271
Reduce [F]	271

Optimal result

Integrand size = 12, antiderivative size = 20

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = -\cos(a) \operatorname{CosIntegral}\left(\frac{b}{x}\right) + \sin(a) \operatorname{Si}\left(\frac{b}{x}\right)$$

output `-cos(a)*Ci(b/x)+sin(a)*Si(b/x)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = -\cos(a) \operatorname{CosIntegral}\left(\frac{b}{x}\right) + \sin(a) \operatorname{Si}\left(\frac{b}{x}\right)$$

input `Integrate[Cos[a + b/x]/x,x]`

output `-(Cos[a]*CosIntegral[b/x]) + Sin[a]*SinIntegral[b/x]`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3859, 3856, 3857}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx$$

$$\downarrow \text{3859}$$

$$\cos(a) \int \frac{\cos\left(\frac{b}{x}\right)}{x} dx - \sin(a) \int \frac{\sin\left(\frac{b}{x}\right)}{x} dx$$

$$\downarrow \text{3856}$$

$$\cos(a) \int \frac{\cos\left(\frac{b}{x}\right)}{x} dx + \sin(a) \text{Si}\left(\frac{b}{x}\right)$$

$$\downarrow \text{3857}$$

$$\sin(a) \text{Si}\left(\frac{b}{x}\right) - \cos(a) \text{CosIntegral}\left(\frac{b}{x}\right)$$

input `Int[Cos[a + b/x]/x,x]`

output `-(Cos[a]*CosIntegral[b/x]) + Sin[a]*SinIntegral[b/x]`

Defintions of rubi rules used

rule 3856 `Int[Sin[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[SinIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3857 `Int[Cos[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[CosIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3859

```
Int[Cos[(c_) + (d_.)*(x_)^(n_)]/(x_), x_Symbol] := Simp[Cos[c] Int[Cos[d*
x^n]/x, x], x] - Simp[Sin[c] Int[Sin[d*x^n]/x, x], x] /; FreeQ[{c, d, n},
x]
```

Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

method	result	size
derivativdivides	$-\cos(a) \operatorname{Ci}\left(\frac{b}{x}\right) + \sin(a) \operatorname{Si}\left(\frac{b}{x}\right)$	21
default	$-\cos(a) \operatorname{Ci}\left(\frac{b}{x}\right) + \sin(a) \operatorname{Si}\left(\frac{b}{x}\right)$	21
risch	$-\frac{i\pi \operatorname{csgn}\left(\frac{b}{x}\right) e^{-ia}}{2} + i \operatorname{Si}\left(\frac{b}{x}\right) e^{-ia} + \frac{e^{-ia} \operatorname{expIntegral}_1\left(-\frac{ib}{x}\right)}{2} + \frac{e^{ia} \operatorname{expIntegral}_1\left(-\frac{ib}{x}\right)}{2}$	63
meijerg	$-\frac{\cos(a)\sqrt{\pi} \left(\frac{2\gamma - 2\ln(x) + \ln(b^2)}{\sqrt{\pi}} - \frac{2\gamma}{\sqrt{\pi}} - \frac{2\ln(2)}{\sqrt{\pi}} - \frac{2\ln\left(\frac{b}{2x}\right)}{\sqrt{\pi}} + \frac{2 \operatorname{Ci}\left(\frac{b}{x}\right)}{\sqrt{\pi}} \right)}{2} + \sin(a) \operatorname{Si}\left(\frac{b}{x}\right)$	71

input

```
int(cos(a+b/x)/x,x,method=_RETURNVERBOSE)
```

output

```
-cos(a)*Ci(b/x)+sin(a)*Si(b/x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = -\cos(a) \operatorname{Ci}\left(\frac{b}{x}\right) + \sin(a) \operatorname{Si}\left(\frac{b}{x}\right)$$

input

```
integrate(cos(a+b/x)/x,x, algorithm="fricas")
```

output

```
-cos(a)*cos_integral(b/x) + sin(a)*sin_integral(b/x)
```

Sympy [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = \sin(a) \operatorname{Si}\left(\frac{b}{x}\right) - \cos(a) \operatorname{Ci}\left(\frac{b}{x}\right)$$

input `integrate(cos(a+b/x)/x,x)`

output `sin(a)*Si(b/x) - cos(a)*Ci(b/x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.15

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = -\frac{1}{2} \left(\operatorname{Ei}\left(\frac{ib}{x}\right) + \operatorname{Ei}\left(-\frac{ib}{x}\right) \right) \cos(a) - \frac{1}{2} \left(i \operatorname{Ei}\left(\frac{ib}{x}\right) - i \operatorname{Ei}\left(-\frac{ib}{x}\right) \right) \sin(a)$$

input `integrate(cos(a+b/x)/x,x, algorithm="maxima")`

output `-1/2*(Ei(I*b/x) + Ei(-I*b/x))*cos(a) - 1/2*(I*Ei(I*b/x) - I*Ei(-I*b/x))*sin(a)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(20) = 40.

Time = 0.35 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.05

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = -\frac{b \cos(a) \operatorname{Ci}\left(-a + \frac{ax+b}{x}\right) + b \sin(a) \operatorname{Si}\left(a - \frac{ax+b}{x}\right)}{b}$$

input `integrate(cos(a+b/x)/x,x, algorithm="giac")`

output `-(b*cos(a)*cos_integral(-a + (a*x + b)/x) + b*sin(a)*sin_integral(a - (a*x + b)/x))/b`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = \sin(a) \operatorname{sinint}\left(\frac{b}{x}\right) - \cos(a) \operatorname{cosint}\left(\frac{b}{x}\right)$$

input `int(cos(a + b/x)/x,x)`

output `sin(a)*sinint(b/x) - cos(a)*cosint(b/x)`

Reduce [F]

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x} dx = \int \frac{\cos\left(\frac{ax+b}{x}\right)}{x} dx + \int \frac{1}{x} dx - \log(x)$$

input `int(cos(a+b/x)/x,x)`

output `int(cos((a*x + b)/x)/x,x) + int(1/x,x) - log(x)`

$$3.37 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx$$

Optimal result	272
Mathematica [A] (verified)	272
Rubi [A] (verified)	273
Maple [A] (verified)	274
Fricas [A] (verification not implemented)	275
Sympy [A] (verification not implemented)	275
Maxima [A] (verification not implemented)	275
Giac [A] (verification not implemented)	276
Mupad [B] (verification not implemented)	276
Reduce [B] (verification not implemented)	276

Optimal result

Integrand size = 12, antiderivative size = 13

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

output `-sin(a+b/x)/b`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

input `Integrate[Cos[a + b/x]/x^2,x]`

output `-(Sin[a + b/x]/b)`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3861, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx \\ & \quad \downarrow \text{3861} \\ & - \int \cos\left(a + \frac{b}{x}\right) d\frac{1}{x} \\ & \quad \downarrow \text{3042} \\ & - \int \sin\left(a + \frac{b}{x} + \frac{\pi}{2}\right) d\frac{1}{x} \\ & \quad \downarrow \text{3117} \\ & - \frac{\sin\left(a + \frac{b}{x}\right)}{b} \end{aligned}$$

input `Int[Cos[a + b/x]/x^2,x]`

output `-(Sin[a + b/x]/b)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
  := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$-\frac{\sin\left(a+\frac{b}{x}\right)}{b}$	14
default	$-\frac{\sin\left(a+\frac{b}{x}\right)}{b}$	14
risch	$-\frac{\sin\left(\frac{ax+b}{x}\right)}{b}$	16
parallelrisch	$-\frac{\sin\left(\frac{ax+b}{x}\right)}{b}$	16
norman	$-\frac{2 \tan\left(\frac{a}{2}+\frac{b}{2x}\right)}{b\left(1+\tan\left(\frac{a}{2}+\frac{b}{2x}\right)^2\right)}$	34
meijerg	$-\frac{\cos(a) \sin\left(\frac{b}{x}\right)}{b} + \frac{\sin(a)\sqrt{\pi}\left(\frac{1}{\sqrt{\pi}}-\frac{\cos\left(\frac{b}{x}\right)}{\sqrt{\pi}}\right)}{b}$	39
orering	$-\frac{2x \cos\left(a+\frac{b}{x}\right)}{b^2} - \frac{x^4\left(\frac{b \sin\left(a+\frac{b}{x}\right)}{x^4}-\frac{2 \cos\left(a+\frac{b}{x}\right)}{x^3}\right)}{b^2}$	51

input `int(cos(a+b/x)/x^2,x,method=_RETURNVERBOSE)`

output `-sin(a+b/x)/b`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(\frac{ax+b}{x}\right)}{b}$$

input `integrate(cos(a+b/x)/x^2,x, algorithm="fricas")`output `-sin((a*x + b)/x)/b`**Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = \begin{cases} -\frac{\sin\left(a + \frac{b}{x}\right)}{b} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{x} & \text{otherwise} \end{cases}$$

input `integrate(cos(a+b/x)/x**2,x)`output `Piecewise((-sin(a + b/x)/b, Ne(b, 0)), (-cos(a)/x, True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

input `integrate(cos(a+b/x)/x^2,x, algorithm="maxima")`output `-sin(a + b/x)/b`

Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(\frac{ax+b}{x}\right)}{b}$$

input `integrate(cos(a+b/x)/x^2,x, algorithm="giac")`output `-sin((a*x + b)/x)/b`**Mupad [B] (verification not implemented)**

Time = 42.48 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(a + \frac{b}{x}\right)}{b}$$

input `int(cos(a + b/x)/x^2,x)`output `-sin(a + b/x)/b`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} dx = -\frac{\sin\left(\frac{ax+b}{x}\right)}{b}$$

input `int(cos(a+b/x)/x^2,x)`output `(- sin((a*x + b)/x))/b`

$$3.38 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx$$

Optimal result	277
Mathematica [A] (verified)	277
Rubi [A] (verified)	278
Maple [A] (verified)	279
Fricas [A] (verification not implemented)	280
Sympy [A] (verification not implemented)	281
Maxima [C] (verification not implemented)	281
Giac [A] (verification not implemented)	281
Mupad [B] (verification not implemented)	282
Reduce [B] (verification not implemented)	282

Optimal result

Integrand size = 12, antiderivative size = 30

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = -\frac{\cos\left(a + \frac{b}{x}\right)}{b^2} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx}$$

output

```
-cos(a+b/x)/b^2-sin(a+b/x)/b/x
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = -\frac{x \cos\left(a + \frac{b}{x}\right) + b \sin\left(a + \frac{b}{x}\right)}{b^2 x}$$

input

```
Integrate[Cos[a + b/x]/x^3,x]
```

output

```
-((x*cos[a + b/x] + b*sin[a + b/x])/(b^2*x))
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3861, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx \\
 & \quad \downarrow \text{3861} \\
 & - \int \frac{\cos\left(a + \frac{b}{x}\right)}{x} d\frac{1}{x} \\
 & \quad \downarrow \text{3042} \\
 & - \int \frac{\sin\left(a + \frac{b}{x} + \frac{\pi}{2}\right)}{x} d\frac{1}{x} \\
 & \quad \downarrow \text{3777} \\
 & - \frac{\int -\sin\left(a + \frac{b}{x}\right) d\frac{1}{x}}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \sin\left(a + \frac{b}{x}\right) d\frac{1}{x}}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sin\left(a + \frac{b}{x}\right) d\frac{1}{x}}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx} \\
 & \quad \downarrow \text{3118} \\
 & - \frac{\cos\left(a + \frac{b}{x}\right)}{b^2} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx}
 \end{aligned}$$

input `Int[Cos[a + b/x]/x^3,x]`

output `-(Cos[a + b/x]/b^2) - Sin[a + b/x]/(b*x)`

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 1.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

method	result	size
risch	$-\frac{\cos\left(\frac{ax+b}{x}\right)}{b^2} - \frac{\sin\left(\frac{ax+b}{x}\right)}{bx}$	35
parallelrisch	$\frac{-b \sin\left(\frac{ax+b}{x}\right) + x - x \cos\left(\frac{ax+b}{x}\right)}{b^2 x}$	36
derivativedivides	$-\frac{\cos\left(a+\frac{b}{x}\right) + \left(a+\frac{b}{x}\right) \sin\left(a+\frac{b}{x}\right) - a \sin\left(a+\frac{b}{x}\right)}{b^2}$	42
default	$-\frac{\cos\left(a+\frac{b}{x}\right) + \left(a+\frac{b}{x}\right) \sin\left(a+\frac{b}{x}\right) - a \sin\left(a+\frac{b}{x}\right)}{b^2}$	42
orering	$-\frac{4 \cos\left(a+\frac{b}{x}\right)}{b^2} - \frac{x^4 \left(\frac{b \sin\left(a+\frac{b}{x}\right)}{x^5} - \frac{3 \cos\left(a+\frac{b}{x}\right)}{x^4}\right)}{b^2}$	50
norman	$\frac{2x^2 \tan\left(\frac{a}{2} + \frac{b}{2x}\right)^2}{b^2} - \frac{2x \tan\left(\frac{a}{2} + \frac{b}{2x}\right)}{b}$ $\frac{\left(1 + \tan\left(\frac{a}{2} + \frac{b}{2x}\right)\right)^2}{x^2}$	61
meijerg	$-\frac{2 \cos(a) \sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos\left(\frac{b}{x}\right)}{2\sqrt{\pi}} + \frac{b \sin\left(\frac{b}{x}\right)}{2\sqrt{\pi} x}\right)}{b^2} + \frac{2 \sin(a) \sqrt{\pi} \left(-\frac{b \cos\left(\frac{b}{x}\right)}{2\sqrt{\pi} x} + \frac{\sin\left(\frac{b}{x}\right)}{2\sqrt{\pi}}\right)}{b^2}$	81

input `int(cos(a+b/x)/x^3,x,method=_RETURNVERBOSE)`

output `-1/b^2*cos((a*x+b)/x)-1/b/x*sin((a*x+b)/x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.10

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = -\frac{x \cos\left(\frac{ax+b}{x}\right) + b \sin\left(\frac{ax+b}{x}\right)}{b^2 x}$$

input `integrate(cos(a+b/x)/x^3,x, algorithm="fricas")`

output `-(x*cos((a*x + b)/x) + b*sin((a*x + b)/x))/(b^2*x)`

Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = \begin{cases} -\frac{\sin\left(a + \frac{b}{x}\right)}{bx} - \frac{\cos\left(a + \frac{b}{x}\right)}{b^2} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{2x^2} & \text{otherwise} \end{cases}$$

input `integrate(cos(a+b/x)/x**3,x)`

output `Piecewise((-sin(a + b/x)/(b*x) - cos(a + b/x)/b**2, Ne(b, 0)), (-cos(a)/(2*x**2), True))`

Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.70

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = -\frac{\left(\Gamma\left(2, \frac{ib}{x}\right) + \Gamma\left(2, -\frac{ib}{x}\right)\right) \cos(a) - \left(i\Gamma\left(2, \frac{ib}{x}\right) - i\Gamma\left(2, -\frac{ib}{x}\right)\right) \sin(a)}{2b^2}$$

input `integrate(cos(a+b/x)/x^3,x, algorithm="maxima")`

output `-1/2*((gamma(2, I*b/x) + gamma(2, -I*b/x))*cos(a) - (I*gamma(2, I*b/x) - I*gamma(2, -I*b/x))*sin(a))/b^2`

Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.63

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = \frac{a \sin\left(\frac{ax+b}{x}\right) - \frac{(ax+b) \sin\left(\frac{ax+b}{x}\right)}{x} - \cos\left(\frac{ax+b}{x}\right)}{b^2}$$

input `integrate(cos(a+b/x)/x^3,x, algorithm="giac")`

output $(a*\sin((a*x + b)/x) - (a*x + b)*\sin((a*x + b)/x)/x - \cos((a*x + b)/x))/b^2$

Mupad [B] (verification not implemented)

Time = 42.79 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = -\frac{\cos\left(a + \frac{b}{x}\right)}{b^2} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx}$$

input $\text{int}(\cos(a + b/x)/x^3, x)$

output $-\cos(a + b/x)/b^2 - \sin(a + b/x)/(b*x)$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^3} dx = \frac{-\cos\left(\frac{ax+b}{x}\right)x - \sin\left(\frac{ax+b}{x}\right)b}{b^2x}$$

input $\text{int}(\cos(a+b/x)/x^3, x)$

output $(-\cos((a*x + b)/x)*x + \sin((a*x + b)/x)*b)/(b**2*x)$

$$3.39 \quad \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx$$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [A] (verified)	286
Fricas [A] (verification not implemented)	286
Sympy [A] (verification not implemented)	287
Maxima [C] (verification not implemented)	287
Giac [B] (verification not implemented)	287
Mupad [B] (verification not implemented)	288
Reduce [B] (verification not implemented)	288

Optimal result

Integrand size = 12, antiderivative size = 46

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = -\frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} + \frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{\sin\left(a + \frac{b}{x}\right)}{b x^2}$$

output

```
-2*cos(a+b/x)/b^2/x+2*sin(a+b/x)/b^3-sin(a+b/x)/b/x^2
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = -\frac{2 \cos\left(a + \frac{b}{x}\right)}{b^2 x} + \frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{\sin\left(a + \frac{b}{x}\right)}{b x^2}$$

input

```
Integrate[Cos[a + b/x]/x^4,x]
```

output

```
(-2*Cos[a + b/x])/(b^2*x) + (2*Sin[a + b/x])/b^3 - Sin[a + b/x]/(b*x^2)
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3861, 3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx \\
 & \quad \downarrow \text{3861} \\
 & - \int \frac{\cos\left(a + \frac{b}{x}\right)}{x^2} d\frac{1}{x} \\
 & \quad \downarrow \text{3042} \\
 & - \int \frac{\sin\left(a + \frac{b}{x} + \frac{\pi}{2}\right)}{x^2} d\frac{1}{x} \\
 & \quad \downarrow \text{3777} \\
 & \frac{2 \int -\frac{\sin\left(a + \frac{b}{x}\right)}{x} d\frac{1}{x}}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{2 \int \frac{\sin\left(a + \frac{b}{x}\right)}{x} d\frac{1}{x}}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \int \frac{\sin\left(a + \frac{b}{x}\right)}{x} d\frac{1}{x}}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} \\
 & \quad \downarrow \text{3777} \\
 & \frac{2 \left(\frac{\int \cos\left(a + \frac{b}{x}\right) d\frac{1}{x}}{b} - \frac{\cos\left(a + \frac{b}{x}\right)}{bx} \right)}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{2 \left(\frac{\int \sin\left(a + \frac{b}{x} + \frac{\pi}{2}\right) d\frac{1}{x}}{b} - \frac{\cos\left(a + \frac{b}{x}\right)}{bx} \right)}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2}$$

↓ 3117

$$\frac{2 \left(\frac{\sin\left(a + \frac{b}{x}\right)}{b^2} - \frac{\cos\left(a + \frac{b}{x}\right)}{bx} \right)}{b} - \frac{\sin\left(a + \frac{b}{x}\right)}{bx^2}$$

input `Int[Cos[a + b/x]/x^4,x]`

output `-(Sin[a + b/x]/(b*x^2)) + (2*(-(Cos[a + b/x]/(b*x)) + Sin[a + b/x]/b^2))/b`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

method	result
risch	$-\frac{2 \cos\left(\frac{ax+b}{x}\right)}{b^2 x} - \frac{(b^2-2x^2) \sin\left(\frac{ax+b}{x}\right)}{b^3 x^2}$
parallelrisch	$\frac{-\sin\left(\frac{ax+b}{x}\right)b^2+2\sin\left(\frac{ax+b}{x}\right)x^2-2\cos\left(\frac{ax+b}{x}\right)bx}{b^3 x^2}$
orering	$-\frac{2(3b^2-4x^2)\cos\left(a+\frac{b}{x}\right)}{x b^4} - \frac{(b^2-2x^2)x^4\left(\frac{b\sin\left(a+\frac{b}{x}\right)}{x^6}-\frac{4\cos\left(a+\frac{b}{x}\right)}{x^5}\right)}{b^4}$
norman	$\frac{-\frac{2x^2}{b^2}+\frac{4x^3\tan\left(\frac{a}{2}+\frac{b}{2x}\right)}{b^3}+\frac{2x^2\tan\left(\frac{a}{2}+\frac{b}{2x}\right)^2}{b^2}-\frac{2x\tan\left(\frac{a}{2}+\frac{b}{2x}\right)}{b}}{\left(1+\tan\left(\frac{a}{2}+\frac{b}{2x}\right)^2\right)x^3}$
derivativedivides	$-\frac{a^2\sin\left(a+\frac{b}{x}\right)-2a\left(\cos\left(a+\frac{b}{x}\right)+\left(a+\frac{b}{x}\right)\sin\left(a+\frac{b}{x}\right)\right)+\left(a+\frac{b}{x}\right)^2\sin\left(a+\frac{b}{x}\right)-2\sin\left(a+\frac{b}{x}\right)+2\left(a+\frac{b}{x}\right)\cos\left(a+\frac{b}{x}\right)}{b^3}$
default	$-\frac{a^2\sin\left(a+\frac{b}{x}\right)-2a\left(\cos\left(a+\frac{b}{x}\right)+\left(a+\frac{b}{x}\right)\sin\left(a+\frac{b}{x}\right)\right)+\left(a+\frac{b}{x}\right)^2\sin\left(a+\frac{b}{x}\right)-2\sin\left(a+\frac{b}{x}\right)+2\left(a+\frac{b}{x}\right)\cos\left(a+\frac{b}{x}\right)}{b^3}$
meijerg	$-\frac{4\cos(a)\sqrt{\pi}\sqrt{b^2}\left(\frac{(b^2)^{\frac{3}{2}}\cos\left(\frac{b}{x}\right)}{2\sqrt{\pi}x b^2}-\frac{(b^2)^{\frac{3}{2}}\left(-\frac{3b^2}{2x^2}+3\right)\sin\left(\frac{b}{x}\right)}{6\sqrt{\pi}b^3}\right)}{b^4} + \frac{4\sin(a)\sqrt{\pi}\left(-\frac{1}{2\sqrt{\pi}}+\frac{\left(-\frac{b^2}{2x^2}+1\right)\cos\left(\frac{b}{x}\right)}{2\sqrt{\pi}}+\frac{b\sin\left(\frac{b}{x}\right)}{2\sqrt{\pi}}\right)}{b^3}$

input `int(cos(a+b/x)/x^4,x,method=_RETURNVERBOSE)`

output
$$-2/b^2/x*\cos((a*x+b)/x)-(b^2-2*x^2)/b^3/x^2*\sin((a*x+b)/x)$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.93

$$\int \frac{\cos\left(a+\frac{b}{x}\right)}{x^4} dx = -\frac{2bx\cos\left(\frac{ax+b}{x}\right)+(b^2-2x^2)\sin\left(\frac{ax+b}{x}\right)}{b^3x^2}$$

input `integrate(cos(a+b/x)/x^4,x, algorithm="fricas")`

output
$$-(2*b*x*cos((a*x + b)/x) + (b^2 - 2*x^2)*sin((a*x + b)/x))/(b^3*x^2)$$

Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = \begin{cases} -\frac{\sin\left(a + \frac{b}{x}\right)}{bx^2} - \frac{2\cos\left(a + \frac{b}{x}\right)}{b^2x} + \frac{2\sin\left(a + \frac{b}{x}\right)}{b^3} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{3x^3} & \text{otherwise} \end{cases}$$

input `integrate(cos(a+b/x)/x**4,x)`

output `Piecewise((-sin(a + b/x)/(b*x**2) - 2*cos(a + b/x)/(b**2*x) + 2*sin(a + b/x)/b**3, Ne(b, 0)), (-cos(a)/(3*x**3), True))`

Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.09 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = \frac{(i\Gamma(3, \frac{ib}{x}) - i\Gamma(3, -\frac{ib}{x}))\cos(a) + (\Gamma(3, \frac{ib}{x}) + \Gamma(3, -\frac{ib}{x}))\sin(a)}{2b^3}$$

input `integrate(cos(a+b/x)/x^4,x, algorithm="maxima")`

output `1/2*((I*gamma(3, I*b/x) - I*gamma(3, -I*b/x))*cos(a) + (gamma(3, I*b/x) + gamma(3, -I*b/x))*sin(a))/b^3`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. 2(46) = 92.

Time = 0.39 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.33

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = \frac{a^2 \sin\left(\frac{ax+b}{x}\right) - 2a \cos\left(\frac{ax+b}{x}\right) - \frac{2(ax+b)a \sin\left(\frac{ax+b}{x}\right)}{x} + \frac{2(ax+b) \cos\left(\frac{ax+b}{x}\right)}{x} + \frac{(ax+b)^2 \sin\left(\frac{ax+b}{x}\right)}{x^2} - 2 \sin\left(\frac{ax+b}{x}\right)}{b^3}$$

input `integrate(cos(a+b/x)/x^4,x, algorithm="giac")`

output
$$-(a^2 \sin((a*x + b)/x) - 2*a*\cos((a*x + b)/x) - 2*(a*x + b)*a*\sin((a*x + b)/x)/x + 2*(a*x + b)*\cos((a*x + b)/x)/x + (a*x + b)^2*\sin((a*x + b)/x)/x^2 - 2*\sin((a*x + b)/x))/b^3$$

Mupad [B] (verification not implemented)

Time = 43.47 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.02

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = \frac{2 \sin\left(a + \frac{b}{x}\right)}{b^3} - \frac{b^2 \sin\left(a + \frac{b}{x}\right) + 2bx \cos\left(a + \frac{b}{x}\right)}{b^3 x^2}$$

input `int(cos(a + b/x)/x^4,x)`

output
$$(2*\sin(a + b/x))/b^3 - (b^2*\sin(a + b/x) + 2*b*x*\cos(a + b/x))/(b^3*x^2)$$

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.13

$$\int \frac{\cos\left(a + \frac{b}{x}\right)}{x^4} dx = \frac{-2 \cos\left(\frac{ax+b}{x}\right) bx - \sin\left(\frac{ax+b}{x}\right) b^2 + 2 \sin\left(\frac{ax+b}{x}\right) x^2}{b^3 x^2}$$

input `int(cos(a+b/x)/x^4,x)`

output
$$(-2*\cos((a*x + b)/x)*b*x - \sin((a*x + b)/x)*b**2 + 2*\sin((a*x + b)/x)*x**2)/(b**3*x**2)$$

3.40 $\int \cos\left(a + \frac{b}{x^2}\right) dx$

Optimal result	289
Mathematica [A] (verified)	289
Rubi [A] (verified)	290
Maple [A] (verified)	292
Fricas [A] (verification not implemented)	292
Sympy [F]	293
Maxima [C] (verification not implemented)	293
Giac [F]	294
Mupad [F(-1)]	294
Reduce [F]	294

Optimal result

Integrand size = 8, antiderivative size = 79

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = x \cos\left(a + \frac{b}{x^2}\right) + \sqrt{b}\sqrt{2\pi} \cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) + \sqrt{b}\sqrt{2\pi} \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a)$$

output

```
x*cos(a+b/x^2)+b^(1/2)*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)+b^(1/2)*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)*sin(a)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = x \cos(a) \cos\left(\frac{b}{x^2}\right) + \sqrt{b}\sqrt{2\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) + \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a) \right) - x \sin(a) \sin\left(\frac{b}{x^2}\right)$$

input `Integrate[Cos[a + b/x^2],x]`

output `x*Cos[a]*Cos[b/x^2] + Sqrt[b]*Sqrt[2*Pi]*(Cos[a]*FresnelS[(Sqrt[b]*Sqrt[2/Pi])/x] + FresnelC[(Sqrt[b]*Sqrt[2/Pi])/x]*Sin[a]) - x*Sin[a]*Sin[b/x^2]`

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3841, 3869, 3834, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos\left(a + \frac{b}{x^2}\right) dx \\
 & \quad \downarrow \text{3841} \\
 & - \int x^2 \cos\left(a + \frac{b}{x^2}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{3869} \\
 & 2b \int \sin\left(a + \frac{b}{x^2}\right) d\frac{1}{x} + x \cos\left(a + \frac{b}{x^2}\right) \\
 & \quad \downarrow \text{3834} \\
 & 2b\left(\sin(a) \int \cos\left(\frac{b}{x^2}\right) d\frac{1}{x} + \cos(a) \int \sin\left(\frac{b}{x^2}\right) d\frac{1}{x}\right) + x \cos\left(a + \frac{b}{x^2}\right) \\
 & \quad \downarrow \text{3832} \\
 & 2b\left(\sin(a) \int \cos\left(\frac{b}{x^2}\right) d\frac{1}{x} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}}\right) + x \cos\left(a + \frac{b}{x^2}\right) \\
 & \quad \downarrow \text{3833}
 \end{aligned}$$

$$2b \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC} \left(\frac{\sqrt{b} \sqrt{\frac{2}{\pi}}}{x} \right)}{\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS} \left(\frac{\sqrt{b} \sqrt{\frac{2}{\pi}}}{x} \right)}{\sqrt{b}} \right) + x \cos \left(a + \frac{b}{x^2} \right)$$

input `Int[Cos[a + b/x^2], x]`

output `x*cos[a + b/x^2] + 2*b*((Sqrt[Pi/2]*Cos[a]*FresnelS[(Sqrt[b]*Sqrt[2/Pi])/x])/Sqrt[b] + (Sqrt[Pi/2]*FresnelC[(Sqrt[b]*Sqrt[2/Pi])/x]*Sin[a])/Sqrt[b])`

Defintions of rubi rules used

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3834 `Int[Sin[(c_) + (d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[Sin[c] Int[Cos[d*(e + f*x)^2], x], x] + Simp[Cos[c] Int[Sin[d*(e + f*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]`

rule 3841 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)])*(b_.)^(p_.), x_Symbol] := Simp[-f^(-1) Subst[Int[(a + b*cos[c + d/x^n])^p/x^2, x], x, 1/(e + f*x)], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && ILtQ[n, 0] && EqQ[n, -2]`

rule 3869 `Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_))^(m_), x_Symbol] := Simp[(e*x)^(m + 1)*(Cos[c + d*x^n]/(e*(m + 1))), x] + Simp[d*(n/(e^n*(m + 1))) Int[(e*x)^(m + n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[m, -1]`

Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.72

method	result
derivativedivides	$x \cos\left(a + \frac{b}{x^2}\right) + \sqrt{b} \sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right) \right)$
default	$x \cos\left(a + \frac{b}{x^2}\right) + \sqrt{b} \sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right) \right)$
risch	$\frac{ie^{-ia}b\sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{ib}}{x}\right)}{2\sqrt{ib}} - \frac{ie^{ia}b\sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{-ib}}{x}\right)}{2\sqrt{-ib}} + x \cos\left(\frac{ax^2+b}{x^2}\right)$
meijerg	$-\frac{\cos(a)\sqrt{\pi} \sqrt{2} (b^2)^{\frac{1}{4}} \left(-\frac{4x\sqrt{2} \cos\left(\frac{b}{x^2}\right)}{\sqrt{\pi} (b^2)^{\frac{1}{4}}} - \frac{8\sqrt{b} \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)}{(b^2)^{\frac{1}{4}}} \right)}{8} + \frac{\sin(a)\sqrt{\pi} \sqrt{2} \sqrt{b} \left(-\frac{4\sqrt{2}x \sin\left(\frac{b}{x^2}\right)}{\sqrt{b}\sqrt{\pi}} + 8 \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right) \right)}{8}$

input `int(cos(a+b/x^2),x,method=_RETURNVERBOSE)`output `x*cos(a+b/x^2)+b^(1/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.92

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = \sqrt{2}\pi \sqrt{\frac{b}{\pi}} \cos(a) S\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) + \sqrt{2}\pi \sqrt{\frac{b}{\pi}} C\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) \sin(a) + x \cos\left(\frac{ax^2+b}{x^2}\right)$$

input `integrate(cos(a+b/x^2),x, algorithm="fricas")`output `sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*sqrt(b/pi)/x) + sqrt(2)*pi*sqrt(b/pi)*fresnel_cos(sqrt(2)*sqrt(b/pi)/x)*sin(a) + x*cos((a*x^2 + b)/x^2)`

Sympy [F]

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = \int \cos\left(a + \frac{b}{x^2}\right) dx$$

input `integrate(cos(a+b/x**2),x)`

output `Integral(cos(a + b/x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.61

$$\int \cos\left(a + \frac{b}{x^2}\right) dx$$

$$= \frac{\sqrt{2}\left(2\sqrt{2}bx^2\sqrt{\frac{1}{x^4}}\cos\left(\frac{ax^2+b}{x^2}\right) + \left(\left((i+1)\sqrt{\pi}\left(\operatorname{erf}\left(\sqrt{\frac{ib}{x^2}}\right) - 1\right) - (i-1)\sqrt{\pi}\left(\operatorname{erf}\left(\sqrt{-\frac{ib}{x^2}}\right) - 1\right)\right)\cos(a) + (-i-1)\sqrt{\pi}\left(\operatorname{erf}\left(\sqrt{\frac{ib}{x^2}}\right) - 1\right) + (i+1)\sqrt{\pi}\left(\operatorname{erf}\left(\sqrt{-\frac{ib}{x^2}}\right) - 1\right)\right)\sin(a)}{4bx}$$

input `integrate(cos(a+b/x^2),x, algorithm="maxima")`

output `1/4*sqrt(2)*(2*sqrt(2)*b*x^2*sqrt(x^(-4))*cos((a*x^2 + b)/x^2) + (((I + 1)*sqrt(pi)*(erf(sqrt(I*b/x^2)) - 1) - (I - 1)*sqrt(pi)*(erf(sqrt(-I*b/x^2)) - 1))*cos(a) + (-i-1)*sqrt(pi)*(erf(sqrt(I*b/x^2)) - 1) + (i+1)*sqrt(pi)*(erf(sqrt(-I*b/x^2)) - 1))*sin(a))*b*(b^2/x^4)^(1/4)*sqrt(x^4)/(b*x)`

Giac [F]

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = \int \cos\left(a + \frac{b}{x^2}\right) dx$$

input `integrate(cos(a+b/x^2),x, algorithm="giac")`

output `integrate(cos(a + b/x^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = \int \cos\left(a + \frac{b}{x^2}\right) dx$$

input `int(cos(a + b/x^2),x)`

output `int(cos(a + b/x^2), x)`

Reduce [F]

$$\int \cos\left(a + \frac{b}{x^2}\right) dx = \frac{\cos\left(\frac{ax^2+b}{x^2}\right) x^2 + 4\left(\int \frac{\cos\left(\frac{ax^2+b}{x^2}\right)}{x^4} dx\right) b^2 x + 2 \sin\left(\frac{ax^2+b}{x^2}\right) b}{x}$$

input `int(cos(a+b/x^2),x)`

output `(cos((a*x**2 + b)/x**2)*x**2 + 4*int(cos((a*x**2 + b)/x**2)/x**4,x)*b**2*x + 2*sin((a*x**2 + b)/x**2)*b)/x`

$$3.41 \quad \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

Optimal result	295
Mathematica [A] (verified)	295
Rubi [A] (verified)	296
Maple [A] (verified)	297
Fricas [A] (verification not implemented)	297
Sympy [F]	298
Maxima [C] (verification not implemented)	298
Giac [F]	298
Mupad [F(-1)]	299
Reduce [F]	299

Optimal result

Integrand size = 12, antiderivative size = 25

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = -\frac{1}{2} \cos(a) \operatorname{CosIntegral}\left(\frac{b}{x^2}\right) + \frac{1}{2} \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right)$$

output `-1/2*cos(a)*Ci(b/x^2)+1/2*sin(a)*Si(b/x^2)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = \frac{1}{2} \left(-\cos(a) \operatorname{CosIntegral}\left(\frac{b}{x^2}\right) + \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right) \right)$$

input `Integrate[Cos[a + b/x^2]/x,x]`

output `(-(Cos[a]*CosIntegral[b/x^2]) + Sin[a]*SinIntegral[b/x^2])/2`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3859, 3856, 3857}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

$$\downarrow \text{3859}$$

$$\cos(a) \int \frac{\cos\left(\frac{b}{x^2}\right)}{x} dx - \sin(a) \int \frac{\sin\left(\frac{b}{x^2}\right)}{x} dx$$

$$\downarrow \text{3856}$$

$$\cos(a) \int \frac{\cos\left(\frac{b}{x^2}\right)}{x} dx + \frac{1}{2} \sin(a) \text{Si}\left(\frac{b}{x^2}\right)$$

$$\downarrow \text{3857}$$

$$\frac{1}{2} \sin(a) \text{Si}\left(\frac{b}{x^2}\right) - \frac{1}{2} \cos(a) \text{CosIntegral}\left(\frac{b}{x^2}\right)$$

input `Int[Cos[a + b/x^2]/x,x]`

output `-1/2*(Cos[a]*CosIntegral[b/x^2]) + (Sin[a]*SinIntegral[b/x^2])/2`

Defintions of rubi rules used

rule 3856 `Int[Sin[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[SinIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3857 `Int[Cos[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[CosIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3859

```
Int[Cos[(c_) + (d_.)*(x_)^(n_)]/(x_), x_Symbol] := Simp[Cos[c] Int[Cos[d*
x^n]/x, x], x] - Simp[Sin[c] Int[Sin[d*x^n]/x, x], x] /; FreeQ[{c, d, n},
x]
```

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{\cos(a) \operatorname{Ci}\left(\frac{b}{x^2}\right)}{2} + \frac{\sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right)}{2}$	22
default	$-\frac{\cos(a) \operatorname{Ci}\left(\frac{b}{x^2}\right)}{2} + \frac{\sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right)}{2}$	22
risch	$-\frac{ie^{-ia\pi} \operatorname{csgn}\left(\frac{b}{x^2}\right)}{4} + \frac{ie^{-ia} \operatorname{Si}\left(\frac{b}{x^2}\right)}{2} + \frac{e^{-ia} \operatorname{expIntegral}_1\left(-\frac{ib}{x^2}\right)}{4} + \frac{e^{ia} \operatorname{expIntegral}_1\left(-\frac{ib}{x^2}\right)}{4}$	63
meijerg	$-\frac{\cos(a)\sqrt{\pi} \left(\frac{2\gamma - 4\ln(x) + \ln(b^2)}{\sqrt{\pi}} - \frac{2\gamma}{\sqrt{\pi}} - \frac{2\ln(2)}{\sqrt{\pi}} - \frac{2\ln\left(\frac{b}{2x^2}\right)}{\sqrt{\pi}} + \frac{2\operatorname{Ci}\left(\frac{b}{x^2}\right)}{\sqrt{\pi}} \right)}{4} + \frac{\sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right)}{2}$	72

input

```
int(cos(a+b/x^2)/x,x,method=_RETURNVERBOSE)
```

output

```
-1/2*cos(a)*Ci(b/x^2)+1/2*sin(a)*Si(b/x^2)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = -\frac{1}{2} \cos(a) \operatorname{Ci}\left(\frac{b}{x^2}\right) + \frac{1}{2} \sin(a) \operatorname{Si}\left(\frac{b}{x^2}\right)$$

input

```
integrate(cos(a+b/x^2)/x,x, algorithm="fricas")
```

output

```
-1/2*cos(a)*cos_integral(b/x^2) + 1/2*sin(a)*sin_integral(b/x^2)
```

Sympy [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

input `integrate(cos(a+b/x**2)/x,x)`

output `Integral(cos(a + b/x**2)/x, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.72

$$\begin{aligned} \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx &= -\frac{1}{4} \left(\operatorname{Ei}\left(\frac{ib}{x^2}\right) + \operatorname{Ei}\left(-\frac{ib}{x^2}\right) \right) \cos(a) \\ &\quad - \frac{1}{4} \left(i \operatorname{Ei}\left(\frac{ib}{x^2}\right) - i \operatorname{Ei}\left(-\frac{ib}{x^2}\right) \right) \sin(a) \end{aligned}$$

input `integrate(cos(a+b/x^2)/x,x, algorithm="maxima")`

output `-1/4*(Ei(I*b/x^2) + Ei(-I*b/x^2))*cos(a) - 1/4*(I*Ei(I*b/x^2) - I*Ei(-I*b/x^2))*sin(a)`

Giac [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx$$

input `integrate(cos(a+b/x^2)/x,x, algorithm="giac")`

output `integrate(cos(a + b/x^2)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = \frac{\sin(a) \operatorname{sinint}\left(\frac{b}{x^2}\right)}{2} - \frac{\cos(a) \operatorname{cosint}\left(\frac{b}{x^2}\right)}{2}$$

input `int(cos(a + b/x^2)/x,x)`output `(sin(a)*sinint(b/x^2))/2 - (cos(a)*cosint(b/x^2))/2`**Reduce [F]**

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x} dx = \int \frac{\cos\left(\frac{ax^2+b}{x^2}\right)}{x} dx + \int \frac{1}{x} dx - \log(x)$$

input `int(cos(a+b/x^2)/x,x)`output `int(cos((a*x**2 + b)/x**2)/x,x) + int(1/x,x) - log(x)`

3.42
$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$$

Optimal result	300
Mathematica [A] (verified)	300
Rubi [A] (verified)	301
Maple [A] (verified)	302
Fricas [A] (verification not implemented)	303
Sympy [F]	303
Maxima [C] (verification not implemented)	304
Giac [F]	304
Mupad [B] (verification not implemented)	305
Reduce [F]	305

Optimal result

Integrand size = 12, antiderivative size = 74

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = -\frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a)}{\sqrt{b}}$$

output

```
-1/2*2^(1/2)*Pi^(1/2)*cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)/b^(1/2)+
1/2*2^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)*sin(a)/b^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = -\frac{\sqrt{\frac{\pi}{2}} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) - \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a) \right)}{\sqrt{b}}$$

input

```
Integrate[Cos[a + b/x^2]/x^2,x]
```

output

```

-((Sqrt [Pi/2]*(Cos [a]*FresnelC[(Sqrt [b]*Sqrt [2/Pi])/x] - FresnelS[(Sqrt [b]
*Sqrt [2/Pi])/x]*Sin [a]))/Sqrt [b])

```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3865, 3835, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx \\
 & \quad \downarrow \text{3865} \\
 & - \int \cos\left(a + \frac{b}{x^2}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{3835} \\
 & \sin(a) \int \sin\left(\frac{b}{x^2}\right) d\frac{1}{x} - \cos(a) \int \cos\left(\frac{b}{x^2}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{3832} \\
 & \frac{\sqrt{\frac{\pi}{2}} \sin(a) \text{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} - \cos(a) \int \cos\left(\frac{b}{x^2}\right) d\frac{1}{x} \\
 & \quad \downarrow \text{3833} \\
 & \frac{\sqrt{\frac{\pi}{2}} \sin(a) \text{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} - \frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}}
 \end{aligned}$$

input

```

Int [Cos [a + b/x^2]/x^2,x]

```

output $-\left(\sqrt{\pi/2} \cos[a] \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2/\pi}}{x}\right) / \sqrt{b}\right) + \left(\sqrt{\pi/2} \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2/\pi}}{x}\right) \sin[a]\right) / \sqrt{b}$

Defintions of rubi rules used

rule 3832 $\operatorname{Int}[\sin[(d_.) * ((e_.) + (f_.) * (x_.)^2)], x_Symbol] \rightarrow \operatorname{Simp}[(\sqrt{\pi/2} / (f * \operatorname{Rt}[d, 2])) * \operatorname{FresnelS}[\sqrt{2/\pi} * \operatorname{Rt}[d, 2] * (e + f * x)], x] /;$ $\operatorname{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\operatorname{Int}[\cos[(d_.) * ((e_.) + (f_.) * (x_.)^2)], x_Symbol] \rightarrow \operatorname{Simp}[(\sqrt{\pi/2} / (f * \operatorname{Rt}[d, 2])) * \operatorname{FresnelC}[\sqrt{2/\pi} * \operatorname{Rt}[d, 2] * (e + f * x)], x] /;$ $\operatorname{FreeQ}[\{d, e, f\}, x]$

rule 3835 $\operatorname{Int}[\cos[(c_) + (d_.) * ((e_.) + (f_.) * (x_.)^2)], x_Symbol] \rightarrow \operatorname{Simp}[\cos[c] \operatorname{Int}[\cos[d * (e + f * x)^2], x], x] - \operatorname{Simp}[\sin[c] \operatorname{Int}[\sin[d * (e + f * x)^2], x], x] /;$ $\operatorname{FreeQ}[\{c, d, e, f\}, x]$

rule 3865 $\operatorname{Int}[\cos[(a_.) + (b_.) * (x_.)^n] * (x_.)^m, x_Symbol] \rightarrow \operatorname{Simp}[2/n \operatorname{Subst}[\operatorname{Int}[\cos[a + b * x^2], x], x, x^{(n/2)}], x] /;$ $\operatorname{FreeQ}[\{a, b, m, n\}, x] \ \&\& \ \operatorname{EqQ}[m, n/2 - 1]$

Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.65

method	result	size
derivativedivides	$-\frac{\sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2}}{\sqrt{\pi} x}\right) - \sin(a) \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2}}{\sqrt{\pi} x}\right) \right)}{2\sqrt{b}}$	48
default	$-\frac{\sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2}}{\sqrt{\pi} x}\right) - \sin(a) \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2}}{\sqrt{\pi} x}\right) \right)}{2\sqrt{b}}$	48
meijerg	$-\frac{\sqrt{2} \sqrt{\pi} \cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2}}{\sqrt{\pi} x}\right)}{2\sqrt{b}} + \frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2}}{\sqrt{\pi} x}\right) \sin(a)}{2\sqrt{b}}$	56
risch	$-\frac{e^{-ia} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{ib}}{x}\right)}{4\sqrt{ib}} - \frac{e^{ia} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{-ib}}{x}\right)}{4\sqrt{-ib}}$	56

input `int(cos(a+b/x^2)/x^2,x,method=_RETURNVERBOSE)`

output

```
-1/2*2^(1/2)*Pi^(1/2)/b^(1/2)*(cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)
-sin(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.88

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = -\frac{\sqrt{2}\pi\sqrt{\frac{b}{\pi}}\cos(a)C\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) - \sqrt{2}\pi\sqrt{\frac{b}{\pi}}S\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right)\sin(a)}{2b}$$

input

```
integrate(cos(a+b/x^2)/x^2,x, algorithm="fricas")
```

output

```
-1/2*(sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_cos(sqrt(2)*sqrt(b/pi)/x) - sqrt
(2)*pi*sqrt(b/pi)*fresnel_sin(sqrt(2)*sqrt(b/pi)/x)*sin(a))/b
```

Sympy [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$$

input

```
integrate(cos(a+b/x**2)/x**2,x)
```

output

```
Integral(cos(a + b/x**2)/x**2, x)
```


Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.32

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = \frac{\sqrt{2}\sqrt{x^4}\left(\left(-i-1\right)\sqrt{\pi}\left(\operatorname{erf}\left(\sqrt{\frac{ib}{x^2}}\right)-1\right)+\left(i+1\right)\sqrt{\pi}\left(\operatorname{erf}\left(\sqrt{-\frac{ib}{x^2}}\right)-1\right)\right)\cos(a)+\left(-i+1\right)\sqrt{\pi}\sin(a)}{8bx}$$

input `integrate(cos(a+b/x^2)/x^2,x, algorithm="maxima")`

output `-1/8*sqrt(2)*((-I - 1)*sqrt(pi)*(erf(sqrt(I*b/x^2)) - 1) + (I + 1)*sqrt(pi)*(erf(sqrt(-I*b/x^2)) - 1))*cos(a) + (-I + 1)*sqrt(pi)*(erf(sqrt(I*b/x^2)) - 1) + (I - 1)*sqrt(pi)*(erf(sqrt(-I*b/x^2)) - 1))*sin(a)*sqrt(x^4)*(b^2/x^4)^(1/4)/(b*x)`

Giac [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx$$

input `integrate(cos(a+b/x^2)/x^2,x, algorithm="giac")`

output `integrate(cos(a + b/x^2)/x^2, x)`

Mupad [B] (verification not implemented)

Time = 42.56 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.74

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = \frac{\sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}\sqrt{b}}{x\sqrt{\pi}}\right) \sin(a)}{2\sqrt{b}} - \frac{\sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}\sqrt{b}}{x\sqrt{\pi}}\right) \cos(a)}{2\sqrt{b}}$$

input `int(cos(a + b/x^2)/x^2,x)`output `(2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*b^(1/2))/(x*pi^(1/2)))*sin(a)/(2*b^(1/2)) - (2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*b^(1/2))/(x*pi^(1/2)))*cos(a)/(2*b^(1/2))`**Reduce [F]**

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} dx = \frac{\left(\int \frac{\cos\left(\frac{a x^2 + b}{x^2}\right)}{x^2} dx\right) x + \left(\int \frac{1}{x^2} dx\right) x + 1}{x}$$

input `int(cos(a+b/x^2)/x^2,x)`output `(int(cos((a*x**2 + b)/x**2)/x**2,x)*x + int(1/x**2,x)*x + 1)/x`

$$3.43 \quad \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx$$

Optimal result	306
Mathematica [A] (verified)	306
Rubi [A] (verified)	307
Maple [A] (verified)	308
Fricas [A] (verification not implemented)	309
Sympy [A] (verification not implemented)	309
Maxima [A] (verification not implemented)	309
Giac [A] (verification not implemented)	310
Mupad [B] (verification not implemented)	310
Reduce [B] (verification not implemented)	310

Optimal result

Integrand size = 12, antiderivative size = 15

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

output `-1/2*sin(a+b/x^2)/b`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

input `Integrate[Cos[a + b/x^2]/x^3,x]`

output `-1/2*Sin[a + b/x^2]/b`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3861, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx \\ & \quad \downarrow \text{3861} \\ & -\frac{1}{2} \int \cos\left(a + \frac{b}{x^2}\right) d\frac{1}{x^2} \\ & \quad \downarrow \text{3042} \\ & -\frac{1}{2} \int \sin\left(a + \frac{b}{x^2} + \frac{\pi}{2}\right) d\frac{1}{x^2} \\ & \quad \downarrow \text{3117} \\ & -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b} \end{aligned}$$

input `Int[Cos[a + b/x^2]/x^3,x]`

output `-1/2*Sin[a + b/x^2]/b`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
  := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
derivativdivides	$-\frac{\sin\left(a+\frac{b}{x^2}\right)}{2b}$	14
default	$-\frac{\sin\left(a+\frac{b}{x^2}\right)}{2b}$	14
risch	$-\frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2b}$	18
parallelrisch	$-\frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2b}$	18
norman	$-\frac{\tan\left(\frac{a}{2}+\frac{b}{2x^2}\right)}{b\left(1+\tan\left(\frac{a}{2}+\frac{b}{2x^2}\right)^2\right)}$	34
meijerg	$-\frac{\cos(a)\sin\left(\frac{b}{x^2}\right)}{2b} + \frac{\sin(a)\sqrt{\pi}\left(\frac{1}{\sqrt{\pi}}-\frac{\cos\left(\frac{b}{x^2}\right)}{\sqrt{\pi}}\right)}{2b}$	40
orering	$-\frac{3x^2\cos\left(a+\frac{b}{x^2}\right)}{4b^2} - \frac{x^6\left(\frac{2b\sin\left(a+\frac{b}{x^2}\right)}{x^6}-\frac{3\cos\left(a+\frac{b}{x^2}\right)}{x^4}\right)}{4b^2}$	54

input

```
int(cos(a+b/x^2)/x^3,x,method=_RETURNVERBOSE)
```

output

```
-1/2*sin(a+b/x^2)/b
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2b}$$

input `integrate(cos(a+b/x^2)/x^3,x, algorithm="fricas")`output `-1/2*sin((a*x^2 + b)/x^2)/b`**Sympy [A] (verification not implemented)**

Time = 0.47 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.47

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = \begin{cases} -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b} & \text{for } b \neq 0 \\ -\frac{\cos(a)}{2x^2} & \text{otherwise} \end{cases}$$

input `integrate(cos(a+b/x**2)/x**3,x)`output `Piecewise((-sin(a + b/x**2)/(2*b), Ne(b, 0)), (-cos(a)/(2*x**2), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

input `integrate(cos(a+b/x^2)/x^3,x, algorithm="maxima")`output `-1/2*sin(a + b/x^2)/b`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2b}$$

input `integrate(cos(a+b/x^2)/x^3,x, algorithm="giac")`output `-1/2*sin((a*x^2 + b)/x^2)/b`**Mupad [B] (verification not implemented)**

Time = 42.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(a + \frac{b}{x^2}\right)}{2b}$$

input `int(cos(a + b/x^2)/x^3,x)`output `-sin(a + b/x^2)/(2*b)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^3} dx = -\frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2b}$$

input `int(cos(a+b/x^2)/x^3,x)`output `(- sin((a*x**2 + b)/x**2))/(2*b)`

3.44 $\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$

Optimal result	311
Mathematica [A] (verified)	311
Rubi [A] (verified)	312
Maple [A] (verified)	314
Fricas [A] (verification not implemented)	314
Sympy [F]	315
Maxima [C] (verification not implemented)	315
Giac [F]	316
Mupad [F(-1)]	316
Reduce [F]	316

Optimal result

Integrand size = 12, antiderivative size = 97

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx = \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{2b^{3/2}} + \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a)}{2b^{3/2}} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx}$$

output

```
1/4*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)/b^(3/2)+1/4*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)*sin(a)/b^(3/2)-1/2*sin(a+b/x^2)/b/x
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.91

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx = \frac{\sqrt{2\pi}x \cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) + \sqrt{2\pi}x \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right) \sin(a) - 2\sqrt{b} \sin\left(a + \frac{b}{x^2}\right)}{4b^{3/2}x}$$

input `Integrate[Cos[a + b/x^2]/x^4,x]`

output `(Sqrt[2*Pi]*x*Cos[a]*FresnelS[(Sqrt[b]*Sqrt[2/Pi])/x] + Sqrt[2*Pi]*x*FresnelC[(Sqrt[b]*Sqrt[2/Pi])/x]*Sin[a] - 2*Sqrt[b]*Sin[a + b/x^2])/(4*b^(3/2)*x)`

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3891, 3867, 3834, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx \\
 & \quad \downarrow \text{3891} \\
 & - \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^2} d\frac{1}{x} \\
 & \quad \downarrow \text{3867} \\
 & \frac{\int \sin\left(a + \frac{b}{x^2}\right) d\frac{1}{x}}{2b} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx} \\
 & \quad \downarrow \text{3834} \\
 & \frac{\sin(a) \int \cos\left(\frac{b}{x^2}\right) d\frac{1}{x} + \cos(a) \int \sin\left(\frac{b}{x^2}\right) d\frac{1}{x}}{2b} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx} \\
 & \quad \downarrow \text{3832} \\
 & \frac{\sin(a) \int \cos\left(\frac{b}{x^2}\right) d\frac{1}{x} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \text{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}}}{2b} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx} \\
 & \quad \downarrow \text{3833}
 \end{aligned}$$

$$\frac{\frac{\sqrt{\frac{\pi}{2}} \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{\frac{2}{\pi}}}{x}\right)}{\sqrt{b}}}{2b} - \frac{\sin\left(a + \frac{b}{x^2}\right)}{2bx}$$

input `Int[Cos[a + b/x^2]/x^4,x]`

output `((Sqrt[Pi/2]*Cos[a]*FresnelS[(Sqrt[b]*Sqrt[2/Pi])/x])/Sqrt[b] + (Sqrt[Pi/2]*FresnelC[(Sqrt[b]*Sqrt[2/Pi])/x]*Sin[a])/Sqrt[b])/(2*b) - Sin[a + b/x^2]/(2*b*x)`

Defintions of rubi rules used

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_)^2), x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_)^2), x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3834 `Int[Sin[(c_) + (d_.)*((e_.) + (f_.)*(x_)^2), x_Symbol] :> Simp[Sin[c] Int[Cos[d*(e + f*x)^2], x], x] + Simp[Cos[c] Int[Sin[d*(e + f*x)^2], x], x] /; FreeQ[{c, d, e, f}, x]`

rule 3867 `Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*(e_.)*(x_)^(m_.), x_Symbol] :> Simp[e^(n - 1)*(e*x)^(m - n + 1)*(Sin[c + d*x^n]/(d*n)), x] - Simp[e^n*((m - n + 1)/(d*n)) Int[(e*x)^(m - n)*Sin[c + d*x^n], x], x] /; FreeQ[{c, d, e}, x] && IGtQ[n, 0] && LtQ[n, m + 1]`

rule 3891 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :> -Subst[Int[(a + b*Cos[c + d/x^n])^p/x^(m + 2), x], x, 1/x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m] && EqQ[n, -2]`

Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.66

method	result
derivativedivides	$-\frac{\sin\left(a+\frac{b}{x^2}\right)}{2bx} + \frac{\sqrt{2}\sqrt{\pi}\left(\cos(a)\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)+\sin(a)\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)\right)}{4b^{\frac{3}{2}}}$
default	$-\frac{\sin\left(a+\frac{b}{x^2}\right)}{2bx} + \frac{\sqrt{2}\sqrt{\pi}\left(\cos(a)\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)+\sin(a)\operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)\right)}{4b^{\frac{3}{2}}}$
risch	$\frac{ie^{-ia}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{ib}}{x}\right)}{8b\sqrt{ib}} - \frac{ie^{ia}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{-ib}}{x}\right)}{8b\sqrt{-ib}} - \frac{\sin\left(\frac{ax^2+b}{x^2}\right)}{2bx}$
meijerg	$-\frac{\cos(a)\sqrt{\pi}\sqrt{2}\left(b^2\right)^{\frac{1}{4}}\left(\frac{\sqrt{2}\left(b^2\right)^{\frac{3}{4}}\sin\left(\frac{b}{x^2}\right)}{2\sqrt{\pi}xb} - \frac{\left(b^2\right)^{\frac{3}{4}}\operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)}{2b^{\frac{3}{2}}}\right)}{2b^2} + \frac{\sin(a)\sqrt{\pi}\sqrt{2}\left(-\frac{\sqrt{2}\sqrt{b}\cos\left(\frac{b}{x^2}\right)}{2\sqrt{\pi}x} + \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}}{\sqrt{\pi}x}\right)\right)}{2b^{\frac{3}{2}}}$

input `int(cos(a+b/x^2)/x^4,x,method=_RETURNVERBOSE)`output `-1/2*sin(a+b/x^2)/b/x+1/4/b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)/x)+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)/x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.87

$$\int \frac{\cos\left(a+\frac{b}{x^2}\right)}{x^4} dx$$

$$= \frac{\sqrt{2}\pi x \sqrt{\frac{b}{\pi}} \cos(a) S\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) + \sqrt{2}\pi x \sqrt{\frac{b}{\pi}} C\left(\frac{\sqrt{2}\sqrt{\frac{b}{\pi}}}{x}\right) \sin(a) - 2b \sin\left(\frac{ax^2+b}{x^2}\right)}{4b^2x}$$

input `integrate(cos(a+b/x^2)/x^4,x, algorithm="fricas")`output `1/4*(sqrt(2)*pi*x*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*sqrt(b/pi)/x) + sqrt(2)*pi*x*sqrt(b/pi)*fresnel_cos(sqrt(2)*sqrt(b/pi)/x)*sin(a) - 2*b*sin((a*x^2 + b)/x^2))/(b^2*x)`

Sympy [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

input `integrate(cos(a+b/x**2)/x**4,x)`

output `Integral(cos(a + b/x**2)/x**4, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.76

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

$$= \frac{\sqrt{2}(x^4)^{\frac{3}{2}} \left((-i+1) \Gamma\left(\frac{3}{2}, \frac{ib}{x^2}\right) + (i-1) \Gamma\left(\frac{3}{2}, -\frac{ib}{x^2}\right) \right) \cos(a) + \left((i-1) \Gamma\left(\frac{3}{2}, \frac{ib}{x^2}\right) - (i+1) \Gamma\left(\frac{3}{2}, -\frac{ib}{x^2}\right) \right) \sin(a)}{8b^3x^3}$$

input `integrate(cos(a+b/x^2)/x^4,x, algorithm="maxima")`

output `1/8*sqrt(2)*(x^4)^(3/2)*((-I + 1)*gamma(3/2, I*b/x^2) + (I - 1)*gamma(3/2, -I*b/x^2))*cos(a) + ((I - 1)*gamma(3/2, I*b/x^2) - (I + 1)*gamma(3/2, -I*b/x^2))*sin(a)*(b^2/x^4)^(3/4)/(b^3*x^3)`

Giac [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

input `integrate(cos(a+b/x^2)/x^4,x, algorithm="giac")`

output `integrate(cos(a + b/x^2)/x^4, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx = \int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx$$

input `int(cos(a + b/x^2)/x^4,x)`

output `int(cos(a + b/x^2)/x^4, x)`

Reduce [F]

$$\int \frac{\cos\left(a + \frac{b}{x^2}\right)}{x^4} dx = \int \frac{\cos\left(\frac{ax^2+b}{x^2}\right)}{x^4} dx$$

input `int(cos(a+b/x^2)/x^4,x)`

output `int(cos((a*x**2 + b)/x**2)/x**4,x)`

3.45 $\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx$

Optimal result	317
Mathematica [A] (verified)	317
Rubi [A] (verified)	318
Maple [A] (verified)	319
Fricas [A] (verification not implemented)	320
Sympy [B] (verification not implemented)	320
Maxima [A] (verification not implemented)	320
Giac [A] (verification not implemented)	321
Mupad [B] (verification not implemented)	321
Reduce [B] (verification not implemented)	321

Optimal result

Integrand size = 14, antiderivative size = 19

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \cos(\sqrt{x}) \sin(\sqrt{x})$$

output `x^(1/2)+cos(x^(1/2))*sin(x^(1/2))`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]]^2/Sqrt[x],x]`

output `Sqrt[x] + Sin[2*Sqrt[x]]/2`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.47, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3861, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx \\
 & \quad \downarrow \text{3861} \\
 & 2 \int \cos^2(\sqrt{x}) d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sin\left(\sqrt{x} + \frac{\pi}{2}\right)^2 d\sqrt{x} \\
 & \quad \downarrow \text{3115} \\
 & 2 \left(\frac{\int 1 d\sqrt{x}}{2} + \frac{1}{2} \sin(\sqrt{x}) \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{24} \\
 & 2 \left(\frac{\sqrt{x}}{2} + \frac{1}{2} \sin(\sqrt{x}) \cos(\sqrt{x}) \right)
 \end{aligned}$$

input `Int [Cos [Sqrt [x]] ^2/Sqrt [x] ,x]`

output `2*(Sqrt [x]/2 + (Cos [Sqrt [x]] *Sin [Sqrt [x]])/2)`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3115 `Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Simp[b^2*((n - 1)/n) Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

method	result	size
derivativedivides	$\sqrt{x} + \cos(\sqrt{x}) \sin(\sqrt{x})$	14
default	$\sqrt{x} + \cos(\sqrt{x}) \sin(\sqrt{x})$	14

input `int(cos(x^(1/2))^2/x^(1/2),x,method=_RETURNVERBOSE)`

output `x^(1/2)+cos(x^(1/2))*sin(x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \cos(\sqrt{x}) \sin(\sqrt{x}) + \sqrt{x}$$

input `integrate(cos(x^(1/2))^2/x^(1/2),x, algorithm="fricas")`

output `cos(sqrt(x))*sin(sqrt(x)) + sqrt(x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(17) = 34.

Time = 0.10 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.05

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} \sin^2(\sqrt{x}) + \sqrt{x} \cos^2(\sqrt{x}) + \sin(\sqrt{x}) \cos(\sqrt{x})$$

input `integrate(cos(x**(1/2))**2/x**(1/2),x)`

output `sqrt(x)*sin(sqrt(x))**2 + sqrt(x)*cos(sqrt(x))**2 + sin(sqrt(x))*cos(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

input `integrate(cos(x^(1/2))^2/x^(1/2),x, algorithm="maxima")`

output `sqrt(x) + 1/2*sin(2*sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \sqrt{x} + \frac{1}{2} \sin(2\sqrt{x})$$

input `integrate(cos(x^(1/2))^2/x^(1/2),x, algorithm="giac")`

output `sqrt(x) + 1/2*sin(2*sqrt(x))`

Mupad [B] (verification not implemented)

Time = 41.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \frac{\sin(2\sqrt{x})}{2} + \sqrt{x}$$

input `int(cos(x^(1/2))^2/x^(1/2),x)`

output `sin(2*x^(1/2))/2 + x^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.53

$$\int \frac{\cos^2(\sqrt{x})}{\sqrt{x}} dx = \cos(\sqrt{x}) \sin(\sqrt{x}) + \sqrt{x}$$

input `int(cos(x^(1/2))^2/x^(1/2),x)`

output `cos(sqrt(x))*sin(sqrt(x)) + sqrt(x)`

3.46 $\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx$

Optimal result	322
Mathematica [A] (verified)	322
Rubi [A] (verified)	323
Maple [A] (verified)	324
Fricas [A] (verification not implemented)	324
Sympy [A] (verification not implemented)	325
Maxima [A] (verification not implemented)	325
Giac [A] (verification not implemented)	325
Mupad [B] (verification not implemented)	326
Reduce [B] (verification not implemented)	326

Optimal result

Integrand size = 12, antiderivative size = 8

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

output `2*sin(x^(1/2))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]]/Sqrt[x],x]`

output `2*Sin[Sqrt[x]]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3861, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx \\ & \quad \downarrow \text{3861} \\ & 2 \int \cos(\sqrt{x}) d\sqrt{x} \\ & \quad \downarrow \text{3042} \\ & 2 \int \sin\left(\sqrt{x} + \frac{\pi}{2}\right) d\sqrt{x} \\ & \quad \downarrow \text{3117} \\ & 2 \sin(\sqrt{x}) \end{aligned}$$

input `Int[Cos[Sqrt[x]]/Sqrt[x],x]`

output `2*Sin[Sqrt[x]]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
  :-> Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$2 \sin(\sqrt{x})$	7
default	$2 \sin(\sqrt{x})$	7
meijerg	$2 \sin(\sqrt{x})$	7

input

```
int(cos(x^(1/2))/x^(1/2),x,method=_RETURNVERBOSE)
```

output

```
2*sin(x^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input

```
integrate(cos(x^(1/2))/x^(1/2),x, algorithm="fricas")
```

output

```
2*sin(sqrt(x))
```

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `integrate(cos(x**(1/2))/x**(1/2),x)`

output `2*sin(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="maxima")`

output `2*sin(sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `integrate(cos(x^(1/2))/x^(1/2),x, algorithm="giac")`

output `2*sin(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `int(cos(x^(1/2))/x^(1/2),x)`

output `2*sin(x^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{\cos(\sqrt{x})}{\sqrt{x}} dx = 2 \sin(\sqrt{x})$$

input `int(cos(x^(1/2))/x^(1/2),x)`

output `2*sin(sqrt(x))`

3.47 $\int \cos(\sqrt{x}) dx$

Optimal result	327
Mathematica [A] (verified)	327
Rubi [A] (verified)	328
Maple [A] (verified)	329
Fricas [A] (verification not implemented)	330
Sympy [A] (verification not implemented)	330
Maxima [A] (verification not implemented)	330
Giac [A] (verification not implemented)	331
Mupad [B] (verification not implemented)	331
Reduce [B] (verification not implemented)	331

Optimal result

Integrand size = 6, antiderivative size = 22

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

output `2*cos(x^(1/2))+2*x^(1/2)*sin(x^(1/2))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `Integrate[Cos[Sqrt[x]],x]`

output `2*Cos[Sqrt[x]] + 2*Sqrt[x]*Sin[Sqrt[x]]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3843, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3843} \\
 & 2 \int \sqrt{x} \cos(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin\left(\sqrt{x} + \frac{\pi}{2}\right) \, d\sqrt{x} \\
 & \quad \downarrow \text{3777} \\
 & 2 \left(\int -\sin(\sqrt{x}) \, d\sqrt{x} + \sqrt{x} \sin(\sqrt{x}) \right) \\
 & \quad \downarrow \text{25} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3042} \\
 & 2 \left(\sqrt{x} \sin(\sqrt{x}) - \int \sin(\sqrt{x}) \, d\sqrt{x} \right) \\
 & \quad \downarrow \text{3118} \\
 & 2(\sqrt{x} \sin(\sqrt{x}) + \cos(\sqrt{x}))
 \end{aligned}$$

input

```
Int[Cos[Sqrt[x]], x]
```

output

```
2*(Cos[Sqrt[x]] + Sqrt[x]*Sin[Sqrt[x]])
```

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`
- rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`
- rule 3843 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_)])*(b_.))^(p_.), x_Symbol] :> Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x]]^p, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$	17
default	$2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$	17
meijerg	$4\sqrt{\pi} \left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos(\sqrt{x})}{2\sqrt{\pi}} + \frac{\sqrt{x} \sin(\sqrt{x})}{2\sqrt{\pi}} \right)$	33

input `int(cos(x^(1/2)),x,method=_RETURNVERBOSE)`

output `2*cos(x^(1/2))+2*x^(1/2)*sin(x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="fricas")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x**(1/2)),x)`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="maxima")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x})$$

input `integrate(cos(x^(1/2)),x, algorithm="giac")`

output `2*sqrt(x)*sin(sqrt(x)) + 2*cos(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 40.80 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*cos(x^(1/2)) + 2*x^(1/2)*sin(x^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.59

$$\int \cos(\sqrt{x}) dx = 2 \cos(\sqrt{x}) + 2\sqrt{x} \sin(\sqrt{x})$$

input `int(cos(x^(1/2)),x)`

output `2*(cos(sqrt(x)) + sqrt(x)*sin(sqrt(x)))`

3.48 $\int \cos^2(\sqrt{x}) dx$

Optimal result	332
Mathematica [A] (verified)	332
Rubi [A] (verified)	333
Maple [A] (verified)	334
Fricas [A] (verification not implemented)	335
Sympy [A] (verification not implemented)	335
Maxima [A] (verification not implemented)	335
Giac [A] (verification not implemented)	336
Mupad [B] (verification not implemented)	336
Reduce [B] (verification not implemented)	336

Optimal result

Integrand size = 8, antiderivative size = 36

$$\int \cos^2(\sqrt{x}) dx = \frac{x}{2} + \frac{1}{2} \cos^2(\sqrt{x}) + \sqrt{x} \cos(\sqrt{x}) \sin(\sqrt{x})$$

output `1/2*x+1/2*cos(x^(1/2))^2+x^(1/2)*cos(x^(1/2))*sin(x^(1/2))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int \cos^2(\sqrt{x}) dx = \frac{1}{4}(\cos(2\sqrt{x}) + 2(x + \sqrt{x} \sin(2\sqrt{x})))$$

input `Integrate[Cos[Sqrt[x]]^2,x]`

output `(Cos[2*Sqrt[x]] + 2*(x + Sqrt[x]*Sin[2*Sqrt[x]]))/4`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3843, 3042, 3791, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos^2(\sqrt{x}) \, dx \\
 & \quad \downarrow \text{3843} \\
 & 2 \int \sqrt{x} \cos^2(\sqrt{x}) \, d\sqrt{x} \\
 & \quad \downarrow \text{3042} \\
 & 2 \int \sqrt{x} \sin\left(\sqrt{x} + \frac{\pi}{2}\right)^2 \, d\sqrt{x} \\
 & \quad \downarrow \text{3791} \\
 & 2 \left(\frac{\int \sqrt{x} d\sqrt{x}}{2} + \frac{1}{4} \cos^2(\sqrt{x}) + \frac{1}{2} \sqrt{x} \sin(\sqrt{x}) \cos(\sqrt{x}) \right) \\
 & \quad \downarrow \text{15} \\
 & 2 \left(\frac{x}{4} + \frac{1}{4} \cos^2(\sqrt{x}) + \frac{1}{2} \sqrt{x} \sin(\sqrt{x}) \cos(\sqrt{x}) \right)
 \end{aligned}$$

input `Int[Cos[Sqrt[x]]^2,x]`

output `2*(x/4 + Cos[Sqrt[x]]^2/4 + (Sqrt[x]*Cos[Sqrt[x]]*Sin[Sqrt[x]])/2)`

Definitions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3791 `Int[((c_.) + (d_.)*(x_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)*(b*Sine[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1]`

rule 3843 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_))]*(b_.))^(p_.), x_Symbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x])^p, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$2\sqrt{x} \left(\frac{\cos(\sqrt{x}) \sin(\sqrt{x})}{2} + \frac{\sqrt{x}}{2} \right) - \frac{x}{2} - \frac{\sin(\sqrt{x})^2}{2}$	34
default	$2\sqrt{x} \left(\frac{\cos(\sqrt{x}) \sin(\sqrt{x})}{2} + \frac{\sqrt{x}}{2} \right) - \frac{x}{2} - \frac{\sin(\sqrt{x})^2}{2}$	34

input `int(cos(x^(1/2))^2,x,method=_RETURNVERBOSE)`

output `2*x^(1/2)*(1/2*cos(x^(1/2))*sin(x^(1/2))+1/2*x^(1/2))-1/2*x-1/2*sin(x^(1/2))^2`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.67

$$\int \cos^2(\sqrt{x}) dx = \sqrt{x} \cos(\sqrt{x}) \sin(\sqrt{x}) + \frac{1}{2} \cos(\sqrt{x})^2 + \frac{1}{2} x$$

input `integrate(cos(x^(1/2))^2,x, algorithm="fricas")`output `sqrt(x)*cos(sqrt(x))*sin(sqrt(x)) + 1/2*cos(sqrt(x))^2 + 1/2*x`**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.42

$$\int \cos^2(\sqrt{x}) dx = \sqrt{x} \sin(\sqrt{x}) \cos(\sqrt{x}) + \frac{x \sin^2(\sqrt{x})}{2} + \frac{x \cos^2(\sqrt{x})}{2} + \frac{\cos^2(\sqrt{x})}{2}$$

input `integrate(cos(x**(1/2))**2,x)`output `sqrt(x)*sin(sqrt(x))*cos(sqrt(x)) + x*sin(sqrt(x))**2/2 + x*cos(sqrt(x))**2/2 + cos(sqrt(x))**2/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \cos^2(\sqrt{x}) dx = \frac{1}{2} \sqrt{x} \sin(2\sqrt{x}) + \frac{1}{2} x + \frac{1}{4} \cos(2\sqrt{x})$$

input `integrate(cos(x^(1/2))^2,x, algorithm="maxima")`output `1/2*sqrt(x)*sin(2*sqrt(x)) + 1/2*x + 1/4*cos(2*sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \cos^2(\sqrt{x}) dx = \frac{1}{2} \sqrt{x} \sin(2\sqrt{x}) + \frac{1}{2} x + \frac{1}{4} \cos(2\sqrt{x})$$

input `integrate(cos(x^(1/2))^2,x, algorithm="giac")`

output `1/2*sqrt(x)*sin(2*sqrt(x)) + 1/2*x + 1/4*cos(2*sqrt(x))`

Mupad [B] (verification not implemented)

Time = 41.69 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.64

$$\int \cos^2(\sqrt{x}) dx = \frac{x}{2} - \frac{\sin(\sqrt{x})^2}{2} + \frac{\sqrt{x} \sin(2\sqrt{x})}{2}$$

input `int(cos(x^(1/2))^2,x)`

output `x/2 - sin(x^(1/2))^2/2 + (x^(1/2)*sin(2*x^(1/2)))/2`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.58

$$\int \cos^2(\sqrt{x}) dx = \sqrt{x} \cos(\sqrt{x}) \sin(\sqrt{x}) - \frac{\sin(\sqrt{x})^2}{2} + \frac{x}{2} + 1$$

input `int(cos(x^(1/2))^2,x)`

output `(2*sqrt(x)*cos(sqrt(x))*sin(sqrt(x)) - sin(sqrt(x))**2 + x + 2)/2`

3.49 $\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx$

Optimal result	337
Mathematica [A] (verified)	338
Rubi [A] (verified)	338
Maple [A] (verified)	367
Fricas [A] (verification not implemented)	369
Sympy [F]	369
Maxima [C] (verification not implemented)	370
Giac [C] (verification not implemented)	371
Mupad [F(-1)]	372
Reduce [F]	372

Optimal result

Integrand size = 16, antiderivative size = 235

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx = \frac{135135\sqrt{x} \cos(a + b\sqrt[3]{x})}{32b^6} - \frac{3861x^{7/6} \cos(a + b\sqrt[3]{x})}{8b^4}$$

$$+ \frac{39x^{11/6} \cos(a + b\sqrt[3]{x})}{2b^2} + \frac{405405\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{64b^{15/2}}$$

$$+ \frac{405405\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) \sin(a)}{64b^{15/2}} - \frac{405405\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{64b^7}$$

$$+ \frac{27027x^{5/6} \sin(a + b\sqrt[3]{x})}{16b^5} - \frac{429x^{3/2} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{13/6} \sin(a + b\sqrt[3]{x})}{b}$$

output

```
135135/32*x^(1/2)*cos(a+b*x^(1/3))/b^6-3861/8*x^(7/6)*cos(a+b*x^(1/3))/b^4
+39/2*x^(11/6)*cos(a+b*x^(1/3))/b^2+405405/128*2^(1/2)*Pi^(1/2)*cos(a)*Fre
snelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))/b^(15/2)+405405/128*2^(1/2)*Pi^(1/
2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))*sin(a)/b^(15/2)-405405/64*x^
(1/6)*sin(a+b*x^(1/3))/b^7+27027/16*x^(5/6)*sin(a+b*x^(1/3))/b^5-429/4*x^(
3/2)*sin(a+b*x^(1/3))/b^3+3*x^(13/6)*sin(a+b*x^(1/3))/b
```

Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.70

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx = \frac{405405\sqrt{2\pi} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + 405405\sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) \sin(a) + 6\sqrt{b} \sin(a) + 6\sqrt{b} \cos(a)}{128b^{15/2}}$$

input

```
Integrate[x^(3/2)*Cos[a + b*x^(1/3)],x]
```

output

```
(405405*Sqrt[2*Pi]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] + 405405*Sqrt[2*Pi]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] + 6*Sqrt[b]*x^(1/6)*(26*(3465*b*x^(1/3) - 396*b^3*x + 16*b^5*x^(5/3))*Cos[a + b*x^(1/3)] + (-135135 + 36036*b^2*x^(2/3) - 2288*b^4*x^(4/3) + 64*b^6*x^2)*Sin[a + b*x^(1/3)]))/(128*b^(15/2))
```

Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.14, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 1.625$, Rules used = {3897, 3042, 3777, 25, 3042, 3777, 3042, 3777, 25, 3042, 3777, 3042, 3777, 25, 3042, 3777, 3042, 3777, 25, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{3/2} \cos(a + b\sqrt[3]{x}) dx \\ & \quad \downarrow \text{3897} \\ & 3 \int x^{13/6} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x} \\ & \quad \downarrow \text{3042} \\ & 3 \int x^{13/6} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right) d\sqrt[3]{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 3777 \\
& 3 \left(\frac{13 \int -x^{11/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} + \frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} \right) \\
& \downarrow 25 \\
& 3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \int x^{11/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) \\
& \downarrow 3042 \\
& 3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \int x^{11/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) \\
& \downarrow 3777 \\
& 3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \left(\frac{11 \int x^{3/2} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \\
& \downarrow 3042 \\
& 3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \left(\frac{11 \int x^{3/2} \sin(a + b\sqrt[3]{x} + \frac{\pi}{2}) d\sqrt[3]{x}}{2b} - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \\
& \downarrow 3777 \\
& 3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \left(\frac{11 \left(\frac{9 \int -x^{7/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} + \frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} \right)}{2b} - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \\
& \downarrow 25
\end{aligned}$$

$$3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \left(\frac{11 \left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{9 \int x^{7/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)$$

↓ 3042

$$3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \left(\frac{11 \left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{9 \int x^{7/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)$$

↓ 3777

$$3 \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{13 \left(\frac{11 \left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{9 \left(\frac{7 \int x^{5/6} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} - \frac{x^{7/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)$$

↓ 3042

$$\left. \begin{aligned} & \left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{7 \int x^{5/6} \sin(a + b\sqrt[3]{x} + \frac{\pi}{2}) d\sqrt[3]{x}}{2b} - \frac{x^{7/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \right) \\ & \frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{7 \int x^{5/6} \sin(a + b\sqrt[3]{x} + \frac{\pi}{2}) d\sqrt[3]{x}}{2b} - \frac{x^{7/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} - \frac{x^{11/6} \cos(a + b\sqrt[3]{x})}{b} \end{aligned} \right)$$

↓ 3777

$$\left. \begin{aligned}
 & \left(\begin{aligned}
 & \left(\begin{aligned}
 & \left(\begin{aligned}
 & \left(\frac{5 \int -\sqrt{x} \sin(a+b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} + \frac{x^{5/6} \sin(a+b\sqrt[3]{x})}{b} \right) \\
 & \frac{x^{7/6} \cos(a+b\sqrt[3]{x})}{b} \\
 & \frac{7}{2b} \\
 & \frac{9}{2b} \\
 & \frac{11}{2b} \\
 & \frac{x^{3/2} \sin(a+b\sqrt[3]{x})}{b} \\
 & \frac{13}{2b} \\
 & \frac{3}{2b} \\
 & \frac{x^{13/6} \sin(a+b\sqrt[3]{x})}{b}
 \end{aligned}
 \end{aligned}
 \end{aligned}
 \end{aligned}
 \right)
 \end{aligned}
 \right)
 \end{aligned}
 \end{aligned}$$

$$\left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{5 \int \sqrt{x} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) - \frac{x^{7/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \frac{1}{2b}$$

↓ 3042

$$\left(\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{5 \int \sqrt{x} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) - \frac{x^{7/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \frac{1}{2b}$$

↓ 3777

3	$\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{3 \int \sqrt[6]{x} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x} - \sqrt{x} \cos(a + b\sqrt[3]{x})}{2b}$
11	$\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{3 \int \sqrt[6]{x} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x} - \sqrt{x} \cos(a + b\sqrt[3]{x})}{2b}$	$\frac{3 \int \sqrt[6]{x} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x} - \sqrt{x} \cos(a + b\sqrt[3]{x})}{2b}$
13	$\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} -$	$\frac{3 \int \sqrt[6]{x} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x} - \sqrt{x} \cos(a + b\sqrt[3]{x})}{2b}$

↓ 3042

3	$\frac{x^{13/6} \sin(a + b\sqrt[3]{x})}{b} -$	$2b$
13	$\frac{x^{3/2} \sin(a + b\sqrt[3]{x})}{b} -$	$2b$
11	$\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} -$	$2b$
9	$\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} -$	$2b$
7	$\frac{x^{5/6} \sin(a + b\sqrt[3]{x})}{b} -$	$2b$
	$\frac{3 \int \sqrt[6]{x} \sin(a + b\sqrt[3]{x} + \frac{\pi}{2}) d\sqrt[3]{x}}{2b} - \frac{\sqrt{x} \cos(a + b\sqrt[3]{x})}{b}$	$2b$

↓ 3777

		$\frac{1}{2b} \left(\frac{\int \frac{\sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}} dx \sqrt[3]{x}}{\sqrt[6]{x}} + \frac{\sqrt[6]{x} \sin(a+b\sqrt[3]{x})}{b} \right)$
	$\frac{x^{5/6} \sin(a+b\sqrt[3]{x})}{b}$	$\frac{x^{5/6} \sin(a+b\sqrt[3]{x})}{b}$
	$\frac{x^{3/2} \sin(a+b\sqrt[3]{x})}{b}$	$\frac{x^{3/2} \sin(a+b\sqrt[3]{x})}{b}$

↓ 25

↓ 3042

$$\left(\frac{\sqrt[6]{x} \sin(a+b\sqrt[3]{x})}{b} - \int \frac{\sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right)$$

$$\frac{x^{5/6} \sin(a+b\sqrt[3]{x})}{b}$$

$$\frac{x^{3/2} \sin(a+b\sqrt[3]{x})}{b}$$

$$\frac{x^{5/6} \sin(a+b\sqrt[3]{x})}{b}$$

$$\frac{x^{3/2} \sin(a+b\sqrt[3]{x})}{b}$$

↓ 3787

↓ 3042

		$\frac{\sqrt[6]{x} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right) \sin\left(\frac{\sqrt[3]{x}b+\frac{\pi}{2}}{\sqrt[6]{x}}\right) d\sqrt[3]{x}}{2b}$
		$\frac{x^{5/6} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{2b}$
		$\frac{x^{5/6} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{2b}$
		$\frac{x^{3/2} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{2b}$

↓ 3785

			$\int \frac{\sqrt[6]{x} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{x} dx = \frac{2 \sin(a) \int \cos\left(\frac{bx^{2/3}}{b}\right) d\sqrt[6]{x}}{2b}$
			$\int \frac{x^{5/6} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{x} dx = \frac{2 \sin(a) \int \cos\left(\frac{bx^{2/3}}{b}\right) d\sqrt[6]{x}}{2b}$
			$\int \frac{x^{3/2} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{x} dx = \frac{2 \sin(a) \int \cos\left(\frac{bx^{2/3}}{b}\right) d\sqrt[6]{x}}{2b}$

↓ 3786

11	$\frac{x^{3/2} \sin(a+b\sqrt[3]{x})}{b}$	$\frac{x^{5/6} \sin(a+b\sqrt[3]{x})}{b}$	$\frac{\left(\frac{\sqrt[6]{x} \sin(a+b\sqrt[3]{x})}{b} - \frac{2 \sin(a) \int \cos(bx^{2/3}) dx \sqrt[6]{x}}{2b} \right)}{2b}$
13			$2b$

↓ 3832

		$\frac{x^{5/6} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{b} - \frac{\left(\frac{\sqrt[6]{x} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{b} - \frac{2 \sin(a) \int \cos\left(\frac{bx^2/3}{b}\right) d\sqrt[6]{x}}{2b} \right)}{2b}$
	$\frac{x^{3/2} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{b}$	

↓ 3833

						$\frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\right)}{\sqrt{b}}$
			5			$\frac{\sqrt[6]{x} \sin\left(a+b\sqrt[3]{x}\right)}{b}$
			7			$\frac{x^{5/6} \sin\left(a+b\sqrt[3]{x}\right)}{b}$
			9			$\frac{x^{3/2} \sin\left(a+b\sqrt[3]{x}\right)}{b}$
			11			$\frac{x^{3/2} \sin\left(a+b\sqrt[3]{x}\right)}{b}$

input `Int[x^(3/2)*Cos[a + b*x^(1/3)],x]`

output `3*((x^(13/6)*Sin[a + b*x^(1/3)])/b - (13*(-((x^(11/6)*Cos[a + b*x^(1/3)])/b) + (11*((x^(3/2)*Sin[a + b*x^(1/3)])/b - (9*(-((x^(7/6)*Cos[a + b*x^(1/3)])/b) + (7*((x^(5/6)*Sin[a + b*x^(1/3)])/b - (5*(-((Sqrt[x]*Cos[a + b*x^(1/3)])/b) + (3*(-1/2*((Sqrt[2*Pi]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/Sqrt[b] + (Sqrt[2*Pi]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/Sqrt[b]))/b + (x^(1/6)*Sin[a + b*x^(1/3)]/b)/(2*b)))/(2*b)))/(2*b)))/(2*b)))/(2*b)))/(2*b))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3785 `Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Cos[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3786 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Sin[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3787 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3897 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)n])*(b_.)p*(x_)m], x_Symbol] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x(k*(m + 1) - 1)*(a + b*Cos[c + d*x(k*n)]]p, x], x, x(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]`

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{3x^{\frac{13}{6}} \sin\left(a+bx^{\frac{1}{3}}\right)}{b} - \left(\frac{x^{\frac{11}{6}} \cos\left(a+bx^{\frac{1}{3}}\right)}{2b} + \frac{11x^{\frac{3}{2}} \sin\left(a+bx^{\frac{1}{3}}\right)}{4b} - \left(\frac{x^{\frac{7}{6}} \cos\left(a+bx^{\frac{1}{3}}\right)}{2b} + \frac{7x^{\frac{5}{6}} \sin\left(a+bx^{\frac{1}{3}}\right)}{4b} - \dots \right) \right)$
default	$\frac{3x^{\frac{13}{6}} \sin\left(a+bx^{\frac{1}{3}}\right)}{b} - \left(\frac{x^{\frac{11}{6}} \cos\left(a+bx^{\frac{1}{3}}\right)}{2b} + \frac{11x^{\frac{3}{2}} \sin\left(a+bx^{\frac{1}{3}}\right)}{4b} - \left(\frac{x^{\frac{7}{6}} \cos\left(a+bx^{\frac{1}{3}}\right)}{2b} + \frac{7x^{\frac{5}{6}} \sin\left(a+bx^{\frac{1}{3}}\right)}{4b} - \dots \right) \right)$
meijerg	$\frac{192\sqrt{2} \cos(a)\sqrt{\pi}}{(b^2)^{\frac{15}{4}}} \left(\frac{\sqrt{x}\sqrt{2}(b^2)^{\frac{15}{4}} \left(3120x^{\frac{4}{3}}b^4 - 77220x^{\frac{2}{3}}b^2 + 675675 \right) \cos\left(bx^{\frac{1}{3}}\right)}{61440\sqrt{\pi}b^6} - \frac{x^{\frac{1}{6}}\sqrt{2}(b^2)^{\frac{15}{4}} \left(-960x^2b^6 + 34320x^{\frac{4}{3}}b^4 - 54 \dots \right)}{122880\sqrt{\pi}b^6} \right)$

input `int(x^(3/2)*cos(a+b*x^(1/3)),x,method=_RETURNVERBOSE)`

output `3*x^(13/6)*sin(a+b*x^(1/3))/b-39/b*(-1/2/b*x^(11/6)*cos(a+b*x^(1/3))+11/2/b*(1/2/b*x^(3/2)*sin(a+b*x^(1/3))-9/2/b*(-1/2/b*x^(7/6)*cos(a+b*x^(1/3))+7/2/b*(1/2/b*x^(5/6)*sin(a+b*x^(1/3))-5/2/b*(-1/2/b*x^(1/2)*cos(a+b*x^(1/3))+3/2/b*(1/2*x^(1/6)*sin(a+b*x^(1/3))/b-1/4/b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))))))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.62

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx = \frac{3 \left(135135 \sqrt{2\pi} \sqrt{\frac{b}{\pi}} \cos(a) S \left(\sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) + 135135 \sqrt{2\pi} \sqrt{\frac{b}{\pi}} C \left(\sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(a) + 52 \left(16 b^6 x^{11/6} - 396 b^4 x^{7/6} + 3465 b^2 \sqrt{x} \right) \cos(b x^{1/3} + a) - 2 \left(2288 b^5 x^{3/2} - 36036 b^3 x^{5/6} - (64 b^7 x^2 - 135135 b) x^{1/6} \right) \sin(b x^{1/3} + a) \right)}{b^8}$$

input `integrate(x^(3/2)*cos(a+b*x^(1/3)),x, algorithm="fricas")`

output `3/128*(135135*sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x^(1/6)*sqrt(b/pi)) + 135135*sqrt(2)*pi*sqrt(b/pi)*fresnel_cos(sqrt(2)*x^(1/6)*sqrt(b/pi))*sin(a) + 52*(16*b^6*x^(11/6) - 396*b^4*x^(7/6) + 3465*b^2*sqrt(x))*cos(b*x^(1/3) + a) - 2*(2288*b^5*x^(3/2) - 36036*b^3*x^(5/6) - (64*b^7*x^2 - 135135*b)*x^(1/6))*sin(b*x^(1/3) + a)/b^8`

Sympy [F]

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx = \int x^{\frac{3}{2}} \cos(a + b\sqrt[3]{x}) dx$$

input `integrate(x**(3/2)*cos(a+b*x**(1/3)),x)`

output `Integral(x**(3/2)*cos(a + b*x**(1/3)), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.06 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.58

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx = \frac{3 \left(135135 \sqrt{2} \sqrt{\pi} \left((i+1) \cos(a) - (i-1) \sin(a) \right) \operatorname{erf} \left(\sqrt{i b x^{1/6}} \right) + (-i-1) \cos(a) + (i+1) \sin(a) \right)}{b^2 \sqrt[3]{x}}$$

input `integrate(x^(3/2)*cos(a+b*x^(1/3)),x, algorithm="maxima")`

output `3/512*(135135*sqrt(2)*sqrt(pi)*(((I + 1)*cos(a) - (I - 1)*sin(a))*erf(sqrt(I*b)*x^(1/6)) + (-I - 1)*cos(a) + (I + 1)*sin(a))*erf(sqrt(-I*b)*x^(1/6)))*b^(3/2) + 208*(16*b^7*x^(11/6) - 396*b^5*x^(7/6) + 3465*b^3*sqrt(x))*cos(b*x^(1/3) + a) + 8*(64*b^8*x^(13/6) - 2288*b^6*x^(3/2) + 36036*b^4*x^(5/6) - 135135*b^2*x^(1/6))*sin(b*x^(1/3) + a))/b^9`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.03

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx =$$

$$\frac{3 \left(64i b^6 x^{\frac{13}{6}} - 416 b^5 x^{\frac{11}{6}} - 2288i b^4 x^{\frac{9}{2}} + 10296 b^3 x^{\frac{7}{6}} + 36036i b^2 x^{\frac{5}{6}} - 90090 b \sqrt{x} - 135135i x^{\frac{1}{6}} \right) e^{(i b x^{\frac{1}{3}} + i a)}}{128 b^7}$$

$$- \frac{3 \left(-64i b^6 x^{\frac{13}{6}} - 416 b^5 x^{\frac{11}{6}} + 2288i b^4 x^{\frac{9}{2}} + 10296 b^3 x^{\frac{7}{6}} - 36036i b^2 x^{\frac{5}{6}} - 90090 b \sqrt{x} + 135135i x^{\frac{1}{6}} \right) e^{(-i b x^{\frac{1}{3}} + i a)}}{128 b^7}$$

$$+ \frac{405405i \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(-\frac{1}{2} \sqrt{2} x^{\frac{1}{6}} \left(-\frac{i b}{|b|} + 1 \right) \sqrt{|b|} \right) e^{(i a)}}{256 b^7 \left(-\frac{i b}{|b|} + 1 \right) \sqrt{|b|}}$$

$$- \frac{405405i \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(-\frac{1}{2} \sqrt{2} x^{\frac{1}{6}} \left(\frac{i b}{|b|} + 1 \right) \sqrt{|b|} \right) e^{(-i a)}}{256 b^7 \left(\frac{i b}{|b|} + 1 \right) \sqrt{|b|}}$$

input `integrate(x^(3/2)*cos(a+b*x^(1/3)),x, algorithm="giac")`

output

```
-3/128*(64*I*b^6*x^(13/6) - 416*b^5*x^(11/6) - 2288*I*b^4*x^(3/2) + 10296*
b^3*x^(7/6) + 36036*I*b^2*x^(5/6) - 90090*b*sqrt(x) - 135135*I*x^(1/6))*e^
(I*b*x^(1/3) + I*a)/b^7 - 3/128*(-64*I*b^6*x^(13/6) - 416*b^5*x^(11/6) + 2
288*I*b^4*x^(3/2) + 10296*b^3*x^(7/6) - 36036*I*b^2*x^(5/6) - 90090*b*sqrt
(x) + 135135*I*x^(1/6))*e^(-I*b*x^(1/3) - I*a)/b^7 + 405405/256*I*sqrt(2)*
sqrt(pi)*erf(-1/2*sqrt(2)*x^(1/6)*(-I*b/abs(b) + 1)*sqrt(abs(b)))*e^(I*a)/
(b^7*(-I*b/abs(b) + 1)*sqrt(abs(b))) - 405405/256*I*sqrt(2)*sqrt(pi)*erf(-
1/2*sqrt(2)*x^(1/6)*(I*b/abs(b) + 1)*sqrt(abs(b)))*e^(-I*a)/(b^7*(I*b/abs(
b) + 1)*sqrt(abs(b)))
```

Mupad [F(-1)]

Timed out.

$$\int x^{3/2} \cos(a + b\sqrt[3]{x}) dx = \int x^{3/2} \cos(a + b x^{1/3}) dx$$

input `int(x^(3/2)*cos(a + b*x^(1/3)),x)`output `int(x^(3/2)*cos(a + b*x^(1/3)), x)`**Reduce [F]**

$$\int x^{3/2} \cos(a$$

$$+ b\sqrt[3]{x}) dx = \frac{39x^{\frac{11}{6}} \cos(x^{\frac{1}{3}}b+a)b^4}{2} + \frac{135135\sqrt{x} \cos(x^{\frac{1}{3}}b+a)}{32} - \frac{3861x^{\frac{7}{6}} \cos(x^{\frac{1}{3}}b+a)b^2}{8} + \frac{27027x^{\frac{5}{6}} \sin(x^{\frac{1}{3}}b+a)b}{16} - \frac{429\sqrt{x} \sin(x^{\frac{1}{3}}b+a)}{4} b^6$$

input `int(x^(3/2)*cos(a+b*x^(1/3)),x)`output `(3*(416*x**(5/6)*cos(x**(1/3)*b + a)*b**4*x + 90090*sqrt(x)*cos(x**(1/3)*b + a) - 10296*x**(1/6)*cos(x**(1/3)*b + a)*b**2*x + 36036*x**(5/6)*sin(x**(1/3)*b + a)*b - 2288*sqrt(x)*sin(x**(1/3)*b + a)*b**3*x + 64*x**(1/6)*sin(x**(1/3)*b + a)*b**5*x**2 - 45045*int((x**(1/6)*cos(x**(1/3)*b + a))/x**(2/3),x))/(64*b**6)`

3.50 $\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$

Optimal result	373
Mathematica [A] (verified)	374
Rubi [A] (verified)	374
Maple [A] (verified)	386
Fricas [A] (verification not implemented)	387
Sympy [F]	387
Maxima [C] (verification not implemented)	387
Giac [C] (verification not implemented)	388
Mupad [F(-1)]	389
Reduce [F]	389

Optimal result

Integrand size = 16, antiderivative size = 169

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx = -\frac{315\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{8b^4} + \frac{21x^{5/6} \cos(a + b\sqrt[3]{x})}{2b^2}$$

$$+ \frac{315\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{8b^{9/2}}$$

$$- \frac{315\sqrt{\frac{\pi}{2}} \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) \sin(a)}{8b^{9/2}}$$

$$- \frac{105\sqrt{x} \sin(a + b\sqrt[3]{x})}{4b^3} + \frac{3x^{7/6} \sin(a + b\sqrt[3]{x})}{b}$$

output

```
-315/8*x^(1/6)*cos(a+b*x^(1/3))/b^4+21/2*x^(5/6)*cos(a+b*x^(1/3))/b^2+315/
16*2^(1/2)*Pi^(1/2)*cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))/b^(9
/2)-315/16*2^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))*sin
(a)/b^(9/2)-105/4*x^(1/2)*sin(a+b*x^(1/3))/b^3+3*x^(7/6)*sin(a+b*x^(1/3))/
b
```

Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.83

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$$

$$= \frac{315\sqrt{2\pi} \cos(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) - 315\sqrt{2\pi} \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) \sin(a) + 6\sqrt{b}\sqrt[6]{x}(7(-15 + 4b^2x) \operatorname{in}[a + b\sqrt[3]{x}])}{16b^{9/2}}$$

input

```
Integrate[Sqrt[x]*Cos[a + b*x^(1/3)],x]
```

output

```
(315*Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] - 315*Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] + 6*Sqrt[b]*x^(1/6)*(7*(-15 + 4*b^2*x^(2/3))*Cos[a + b*x^(1/3)] + 2*b*(-35 + 4*b^2*x^(2/3))*x^(1/3)*Sin[a + b*x^(1/3)]))/(16*b^(9/2))
```

Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.10, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$, Rules used = {3897, 3042, 3777, 25, 3042, 3777, 3042, 3777, 25, 3042, 3777, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$$

$$\downarrow \text{3897}$$

$$3 \int x^{7/6} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x}$$

$$\downarrow \text{3042}$$

$$3 \int x^{7/6} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right) d\sqrt[3]{x}$$

$$\downarrow \text{3777}$$

$$\begin{aligned}
 & 3 \left(\frac{7 \int -x^{5/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} + \frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} \right) \\
 & \quad \downarrow \text{25} \\
 & 3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \int x^{5/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \int x^{5/6} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \left(\frac{5 \int \sqrt{x} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \left(\frac{5 \int \sqrt{x} \sin(a + b\sqrt[3]{x} + \frac{\pi}{2}) d\sqrt[3]{x}}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \left(\frac{5 \left(\frac{3 \int -\sqrt[6]{x} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} + \frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} \right)}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right) \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \left(\frac{5 \left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{3 \int \sqrt[6]{x} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right)}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)$$

↓ 3042

$$3 \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{7 \left(\frac{5 \left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{3 \int \sqrt[6]{x} \sin(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{2b} \right)}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)$$

↓ 3777

$$\left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} dx \sqrt[3]{x} - \frac{\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)$$

$$\left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{\sqrt[6]{x}} dx \sqrt[3]{x} - \frac{\sqrt{x} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} - \frac{x^{5/6} \cos(a + b\sqrt[3]{x})}{b} \right)$$

$$\left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\cos(a) \int \frac{\cos(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \sin(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} \right)$$

↓ 3042

$$\left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{b} - \frac{\left(\frac{\cos(a) \int \frac{\sin\left(\frac{\sqrt[3]{x}b + \frac{\pi}{2}}{\sqrt[6]{x}}\right) d\sqrt[3]{x} - \sin(a) \int \frac{\sin\left(\frac{b\sqrt[3]{x}}{\sqrt[6]{x}}\right) d\sqrt[3]{x}}{2b} - \frac{\sqrt[6]{x} \cos\left(\frac{a+b\sqrt[3]{x}}{b}\right)}{b} \right)}{2b} \right)}{2b} \right)$$

↓ 3785

$$\left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{2 \cos(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} - \sin(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} \right)$$

↓ 3786

$$\left. \begin{aligned} & \left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{3 \left(\frac{2 \cos(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} - 2 \sin(a) \int \sin(bx^{2/3}) d\sqrt[6]{x} - \frac{\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} \right) \end{aligned} \right\} 3$$

↓ 3832

$$\left. \begin{array}{l}
 \left(\begin{array}{l}
 \left(\begin{array}{l}
 \left(\frac{\sqrt{x} \sin(a + b \sqrt[3]{x})}{b} - \left(\frac{2 \cos(a) \int \cos(bx^{2/3}) d \sqrt[6]{x} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{\sqrt{b}}}{2b} - \frac{\sqrt[6]{x} \cos(a + b \sqrt[3]{x})}{b} \right) \right) \\
 \frac{\sqrt{x} \sin(a + b \sqrt[3]{x})}{b} - \frac{\left(\frac{2 \cos(a) \int \cos(bx^{2/3}) d \sqrt[6]{x} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{\sqrt{b}}}{2b} - \frac{\sqrt[6]{x} \cos(a + b \sqrt[3]{x})}{b} \right)}{2b} \\
 \frac{x^{7/6} \sin(a + b \sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b \sqrt[3]{x})}{b} - \frac{\left(\frac{2 \cos(a) \int \cos(bx^{2/3}) d \sqrt[6]{x} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{\sqrt{b}}}{2b} - \frac{\sqrt[6]{x} \cos(a + b \sqrt[3]{x})}{b} \right)}{2b} \right)}{2b}
 \end{array} \right) \\
 \frac{x^{7/6} \sin(a + b \sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b \sqrt[3]{x})}{b} - \frac{\left(\frac{2 \cos(a) \int \cos(bx^{2/3}) d \sqrt[6]{x} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right)}{\sqrt{b}}}{2b} - \frac{\sqrt[6]{x} \cos(a + b \sqrt[3]{x})}{b} \right)}{2b} \right)}{2b}
 \end{array} \right)
 \end{array} \right)$$

↓ 3833

$$\left(\frac{x^{7/6} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\left(\frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{\sqrt{b}} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{\sqrt{b}} - \frac{\sqrt[6]{x} \cos(a + b\sqrt[3]{x})}{b} \right)}{2b} \right)}{2b} \right)$$

input `Int[Sqrt[x]*Cos[a + b*x^(1/3)],x]`

output `3*((x^(7/6)*Sin[a + b*x^(1/3)])/b - (7*(-((x^(5/6)*Cos[a + b*x^(1/3)]))/b) + (5*((-3*(-((x^(1/6)*Cos[a + b*x^(1/3)]))/b) + ((Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/Sqrt[b] - (Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/Sqrt[b]))/(2*b)))/(2*b)) + (Sqrt[x]*Sin[a + b*x^(1/3)])/b)/(2*b)))/(2*b))`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3777 $\text{Int}[(\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.} * \sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m}} * (\text{Cos}[\text{e} + \text{f} * \text{x}] / \text{f}), \text{x}] + \text{Simp}[\text{d} * (\text{m} / \text{f}) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m} - 1} * \text{Cos}[\text{e} + \text{f} * \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{GtQ}[\text{m}, 0]$
- rule 3785 $\text{Int}[\sin[\text{Pi}/2 + (\text{e}_.) + (\text{f}_.) * (\text{x}_.)] / \text{Sqrt}[(\text{c}_.) + (\text{d}_.) * (\text{x}_.)], \text{x_Symbol}] \rightarrow \text{Simp}[2 / \text{d} \quad \text{Subst}[\text{Int}[\text{Cos}[\text{f} * (\text{x}^2 / \text{d})], \text{x}], \text{x}, \text{Sqrt}[\text{c} + \text{d} * \text{x}], \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{ComplexFreeQ}[\text{f}] \&\& \text{EqQ}[\text{d} * \text{e} - \text{c} * \text{f}, 0]$
- rule 3786 $\text{Int}[\sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)] / \text{Sqrt}[(\text{c}_.) + (\text{d}_.) * (\text{x}_.)], \text{x_Symbol}] \rightarrow \text{Simp}[2 / \text{d} \quad \text{Subst}[\text{Int}[\text{Sin}[\text{f} * (\text{x}^2 / \text{d})], \text{x}], \text{x}, \text{Sqrt}[\text{c} + \text{d} * \text{x}], \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{ComplexFreeQ}[\text{f}] \&\& \text{EqQ}[\text{d} * \text{e} - \text{c} * \text{f}, 0]$
- rule 3787 $\text{Int}[\sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)] / \text{Sqrt}[(\text{c}_.) + (\text{d}_.) * (\text{x}_.)], \text{x_Symbol}] \rightarrow \text{Simp}[\text{Cos}[(\text{d} * \text{e} - \text{c} * \text{f}) / \text{d}] \quad \text{Int}[\text{Sin}[\text{c} * (\text{f} / \text{d}) + \text{f} * \text{x}] / \text{Sqrt}[\text{c} + \text{d} * \text{x}], \text{x}], \text{x}] + \text{Simp}[\text{Sin}[(\text{d} * \text{e} - \text{c} * \text{f}) / \text{d}] \quad \text{Int}[\text{Cos}[\text{c} * (\text{f} / \text{d}) + \text{f} * \text{x}] / \text{Sqrt}[\text{c} + \text{d} * \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{ComplexFreeQ}[\text{f}] \&\& \text{NeQ}[\text{d} * \text{e} - \text{c} * \text{f}, 0]$
- rule 3832 $\text{Int}[\text{Sin}[(\text{d}_.) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^2)], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2] / (\text{f} * \text{Rt}[\text{d}, 2])) * \text{FresnelS}[\text{Sqrt}[2 / \text{Pi}] * \text{Rt}[\text{d}, 2] * (\text{e} + \text{f} * \text{x})], \text{x}] \text{ ; FreeQ}\{\text{d}, \text{e}, \text{f}\}, \text{x}\}$
- rule 3833 $\text{Int}[\text{Cos}[(\text{d}_.) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^2)], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2] / (\text{f} * \text{Rt}[\text{d}, 2])) * \text{FresnelC}[\text{Sqrt}[2 / \text{Pi}] * \text{Rt}[\text{d}, 2] * (\text{e} + \text{f} * \text{x})], \text{x}] \text{ ; FreeQ}\{\text{d}, \text{e}, \text{f}\}, \text{x}\}$

rule 3897

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
 := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a +
 b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x]] /; FreeQ[{a, b, c, d, m}, x]
 && IntegerQ[p] && FractionQ[n]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{3x^{7/6} \sin(a+bx^{1/3})}{b} - \frac{21 \left(-\frac{x^{5/6} \cos(a+bx^{1/3})}{2b} + \frac{5\sqrt{x} \sin(a+bx^{1/3})}{4b} - \frac{15 \left(-\frac{x^{1/6} \cos(a+bx^{1/3})}{2b} + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{2}x^{1/6}}{b}\right) \right)}{b} \right)}{4b} \right)}{b}$
default	$\frac{3x^{7/6} \sin(a+bx^{1/3})}{b} - \frac{21 \left(-\frac{x^{5/6} \cos(a+bx^{1/3})}{2b} + \frac{5\sqrt{x} \sin(a+bx^{1/3})}{4b} - \frac{15 \left(-\frac{x^{1/6} \cos(a+bx^{1/3})}{2b} + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{2}x^{1/6}}{b}\right) \right)}{b} \right)}{4b} \right)}{b}$
meijerg	$\frac{24\sqrt{2} \cos(a)\sqrt{\pi} \left(-\frac{x^{1/6} \sqrt{2} (b^2)^{9/4} (-252x^{2/3} b^2 + 945) \cos(bx^{1/3})}{1152\sqrt{\pi} b^4} - \frac{\sqrt{x} \sqrt{2} (b^2)^{9/4} (-36x^{2/3} b^2 + 315) \sin(bx^{1/3})}{576\sqrt{\pi} b^3} + \frac{105 (b^2)^{9/4} \operatorname{FresnelS}\left(\frac{\sqrt{2}x^{1/6}}{b}\right)}{b} \right)}{(b^2)^{9/4}}$

input

```
int(x^(1/2)*cos(a+b*x^(1/3)),x,method=_RETURNVERBOSE)
```

output

```
3*x^(7/6)*sin(a+b*x^(1/3))/b-21/b*(-1/2/b*x^(5/6)*cos(a+b*x^(1/3))+5/2/b*(
1/2/b*x^(1/2)*sin(a+b*x^(1/3))-3/2/b*(-1/2/b*x^(1/6)*cos(a+b*x^(1/3))+1/4/
b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6)
)-sin(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))))))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.70

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$$

$$= \frac{3 \left(105 \sqrt{2} \pi \sqrt{\frac{b}{\pi}} \cos(a) C\left(\sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) - 105 \sqrt{2} \pi \sqrt{\frac{b}{\pi}} S\left(\sqrt{2} x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(a) + 14 \left(4 b^3 x^{\frac{5}{6}} - 15 b x^{\frac{1}{6}} \right) \cos(a) \right)}{16 b^5}$$

input `integrate(x^(1/2)*cos(a+b*x^(1/3)),x, algorithm="fricas")`

output `3/16*(105*sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_cos(sqrt(2)*x^(1/6)*sqrt(b/pi)) - 105*sqrt(2)*pi*sqrt(b/pi)*fresnel_sin(sqrt(2)*x^(1/6)*sqrt(b/pi))*sin(a) + 14*(4*b^3*x^(5/6) - 15*b*x^(1/6))*cos(b*x^(1/3) + a) + 4*(4*b^4*x^(7/6) - 35*b^2*sqrt(x))*sin(b*x^(1/3) + a))/b^5`

Sympy [F]

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx = \int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$$

input `integrate(x**(1/2)*cos(a+b*x**(1/3)),x)`

output `Integral(sqrt(x)*cos(a + b*x**(1/3)), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.66

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$$

$$= \frac{3 \left(105 \sqrt{2} \sqrt{\pi} \left((-i - 1) \cos(a) - (i + 1) \sin(a) \right) \operatorname{erf}\left(\sqrt{i} b x^{\frac{1}{6}}\right) + ((i + 1) \cos(a) + (i - 1) \sin(a)) \operatorname{erf}\left(\sqrt{i} b x^{\frac{1}{6}}\right) \right)}{64 b^6}$$

input `integrate(x^(1/2)*cos(a+b*x^(1/3)),x, algorithm="maxima")`

output
$$\frac{3}{64}*(105*\sqrt{2}*\sqrt{\pi})*((-I - 1)*\cos(a) - (I + 1)*\sin(a))*\operatorname{erf}(\sqrt{I*b}*x^{(1/6)}) + ((I + 1)*\cos(a) + (I - 1)*\sin(a))*\operatorname{erf}(\sqrt{-I*b}*x^{(1/6)}) * b^{(3/2)} + 56*(4*b^4*x^{(5/6)} - 15*b^2*x^{(1/6)})*\cos(b*x^{(1/3)} + a) + 16*(4*b^5*x^{(7/6)} - 35*b^3*\sqrt{x})*\sin(b*x^{(1/3)} + a))/b^6$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.14

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx = -\frac{3 \left(8i b^3 x^{\frac{7}{6}} - 28 b^2 x^{\frac{5}{6}} - 70i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{(i b x^{\frac{1}{3}} + i a)}}{16 b^4} - \frac{3 \left(-8i b^3 x^{\frac{7}{6}} - 28 b^2 x^{\frac{5}{6}} + 70i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{(-i b x^{\frac{1}{3}} - i a)}}{16 b^4} - \frac{315 \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(-\frac{1}{2} \sqrt{2} x^{\frac{1}{6}} \left(-\frac{i b}{|b|} + 1 \right) \sqrt{|b|} \right) e^{(i a)}}{32 b^4 \left(-\frac{i b}{|b|} + 1 \right) \sqrt{|b|}} - \frac{315 \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(-\frac{1}{2} \sqrt{2} x^{\frac{1}{6}} \left(\frac{i b}{|b|} + 1 \right) \sqrt{|b|} \right) e^{(-i a)}}{32 b^4 \left(\frac{i b}{|b|} + 1 \right) \sqrt{|b|}}$$

input `integrate(x^(1/2)*cos(a+b*x^(1/3)),x, algorithm="giac")`

output
$$-3/16*(8*I*b^3*x^{(7/6)} - 28*b^2*x^{(5/6)} - 70*I*b*\sqrt{x} + 105*x^{(1/6)})*e^{(I*b*x^{(1/3)} + I*a)/b^4} - 3/16*(-8*I*b^3*x^{(7/6)} - 28*b^2*x^{(5/6)} + 70*I*b*\sqrt{x} + 105*x^{(1/6)})*e^{(-I*b*x^{(1/3)} - I*a)/b^4} - 315/32*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{2}*x^{(1/6)}*(-I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)}) * e^{(I*a)/(b^4*(-I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})} - 315/32*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-1/2*\sqrt{2}*x^{(1/6)}*(I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)}) * e^{(-I*a)/(b^4*(I*b/\operatorname{abs}(b) + 1)*\sqrt{\operatorname{abs}(b)})}$$

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx = \int \sqrt{x} \cos(a + b x^{1/3}) dx$$

input `int(x^(1/2)*cos(a + b*x^(1/3)),x)`output `int(x^(1/2)*cos(a + b*x^(1/3)), x)`**Reduce [F]**

$$\int \sqrt{x} \cos(a + b\sqrt[3]{x}) dx$$

$$= \frac{21x^{\frac{5}{6}} \cos\left(x^{\frac{1}{3}}b+a\right)b^2}{2} - \frac{315x^{\frac{1}{6}} \cos\left(x^{\frac{1}{3}}b+a\right)}{8} - \frac{105\sqrt{x} \sin\left(x^{\frac{1}{3}}b+a\right)b}{4} + 3x^{\frac{7}{6}} \sin\left(x^{\frac{1}{3}}b+a\right)b^3 + \frac{105 \left(\int \frac{\cos\left(x^{\frac{1}{3}}b+a\right)}{x^{\frac{5}{6}}} dx \right)}{16}$$

input `int(x^(1/2)*cos(a+b*x^(1/3)),x)`output `(3*(56*x**(5/6)*cos(x**(1/3)*b + a)*b**2 - 210*x**(1/6)*cos(x**(1/3)*b + a) - 140*sqrt(x)*sin(x**(1/3)*b + a)*b + 16*x**(1/6)*sin(x**(1/3)*b + a)*b**3*x + 35*int(cos(x**(1/3)*b + a)/x**(5/6),x))/(16*b**4)`

3.51
$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{\sqrt{x}} dx$$

Optimal result	390
Mathematica [A] (verified)	390
Rubi [A] (verified)	391
Maple [A] (verified)	394
Fricas [A] (verification not implemented)	395
Sympy [F]	395
Maxima [C] (verification not implemented)	395
Giac [C] (verification not implemented)	396
Mupad [F(-1)]	397
Reduce [F]	397

Optimal result

Integrand size = 16, antiderivative size = 99

$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{\sqrt{x}} dx = -\frac{3\sqrt{\frac{\pi}{2}} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{b^{3/2}} - \frac{3\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) \sin(a)}{b^{3/2}} + \frac{3\sqrt[6]{x} \sin\left(a+b\sqrt[3]{x}\right)}{b}$$

output `-3/2*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))/b^(3/2)-3/2*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))*sin(a)/b^(3/2)+3*x^(1/6)*sin(a+b*x^(1/3))/b`

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.95

$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{\sqrt{x}} dx = \frac{3\left(\sqrt{2\pi} \cos(a) \operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + \sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) \sin(a) - 2\sqrt{b}\sqrt[6]{x} \sin\left(a+b\sqrt[3]{x}\right)\right)}{2b^{3/2}}$$

input `Integrate[Cos[a + b*x^(1/3)]/Sqrt[x], x]`

output `(-3*(Sqrt[2*Pi]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] + Sqrt[2*Pi]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] - 2*Sqrt[b]*x^(1/6)*Sin[a + b*x^(1/3)]))/(2*b^(3/2))`

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {3897, 3042, 3777, 25, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx \\
 & \quad \downarrow \text{3897} \\
 & 3 \int \sqrt[6]{x} \cos(a + b\sqrt[3]{x}) d\sqrt[3]{x} \\
 & \quad \downarrow \text{3042} \\
 & 3 \int \sqrt[6]{x} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right) d\sqrt[3]{x} \\
 & \quad \downarrow \text{3777} \\
 & 3 \left(\frac{\int -\frac{\sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{2b} + \frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} \right) \\
 & \quad \downarrow \text{25} \\
 & 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\int \frac{\sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{2b} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\int \frac{\sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{2b} \right) \\
& \quad \downarrow \text{3787} \\
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\sin(a) \int \frac{\cos(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} + \cos(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\sin(a) \int \frac{\sin(\sqrt[3]{x}b + \frac{\pi}{2})}{\sqrt[6]{x}} d\sqrt[3]{x} + \cos(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{2b} \right) \\
& \quad \downarrow \text{3785} \\
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{2 \sin(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} + \cos(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{2b} \right) \\
& \quad \downarrow \text{3786} \\
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{2 \sin(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} + 2 \cos(a) \int \sin(bx^{2/3}) d\sqrt[6]{x}}{2b} \right) \\
& \quad \downarrow \text{3832} \\
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{2 \sin(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} + \frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelS}(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x})}{\sqrt{b}}}{2b} \right) \\
& \quad \downarrow \text{3833} \\
& 3 \left(\frac{\sqrt[6]{x} \sin(a + b\sqrt[3]{x})}{b} - \frac{\frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelC}(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x})}{\sqrt{b}} + \frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelS}(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x})}{\sqrt{b}}}{2b} \right)
\end{aligned}$$

input

Int[Cos[a + b*x^(1/3)]/Sqrt[x], x]

output

$$3*(-1/2*((\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a]*\text{FresnelS}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}])/\text{Sqrt}[b] + (\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[b]*\text{Sqrt}[2/\text{Pi}]*x^{(1/6)}]*\text{Sin}[a])/\text{Sqrt}[b])/b + (x^{(1/6)}*\text{Sin}[a + b*x^{(1/3)}])/b)$$
Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 3042

$$\text{Int}[\text{u}_, \text{x_Symbol}] \text{ :> } \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ;/; } \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$$

rule 3777

$$\text{Int}[((\text{c}_.) + (\text{d}_.)*(x_))^{(m_.)}*\text{sin}[(\text{e}_.) + (\text{f}_.)*(x_)], \text{x_Symbol}] \text{ :> } \text{Simp}[-(\text{c} + \text{d}*x)^m*(\text{Cos}[\text{e} + \text{f}*x]/\text{f}), \text{x}] + \text{Simp}[\text{d}*(m/\text{f}) \quad \text{Int}[(\text{c} + \text{d}*x)^{(m-1)}*\text{Cos}[\text{e} + \text{f}*x], \text{x}], \text{x}] \text{ ;/; } \text{FreeQ}\{\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \ \&\& \ \text{GtQ}[\text{m}, 0]$$

rule 3785

$$\text{Int}[\text{sin}[\text{Pi}/2 + (\text{e}_.) + (\text{f}_.)*(x_)]/\text{Sqrt}[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x_Symbol}] \text{ :> } \text{Simp}[2/\text{d} \quad \text{Subst}[\text{Int}[\text{Cos}[\text{f}*(x^2/\text{d})], \text{x}], \text{x}, \text{Sqrt}[\text{c} + \text{d}*x]], \text{x}] \text{ ;/; } \text{FreeQ}\{\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \ \&\& \ \text{ComplexFreeQ}[\text{f}] \ \&\& \ \text{EqQ}[\text{d}*e - \text{c}*f, 0]$$

rule 3786

$$\text{Int}[\text{sin}[(\text{e}_.) + (\text{f}_.)*(x_)]/\text{Sqrt}[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x_Symbol}] \text{ :> } \text{Simp}[2/\text{d} \quad \text{Subst}[\text{Int}[\text{Sin}[\text{f}*(x^2/\text{d})], \text{x}], \text{x}, \text{Sqrt}[\text{c} + \text{d}*x]], \text{x}] \text{ ;/; } \text{FreeQ}\{\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \ \&\& \ \text{ComplexFreeQ}[\text{f}] \ \&\& \ \text{EqQ}[\text{d}*e - \text{c}*f, 0]$$

rule 3787

$$\text{Int}[\text{sin}[(\text{e}_.) + (\text{f}_.)*(x_)]/\text{Sqrt}[(\text{c}_.) + (\text{d}_.)*(x_)], \text{x_Symbol}] \text{ :> } \text{Simp}[\text{Cos}[(\text{d}*e - \text{c}*f)/\text{d}] \quad \text{Int}[\text{Sin}[\text{c}*(\text{f}/\text{d}) + \text{f}*x]/\text{Sqrt}[\text{c} + \text{d}*x], \text{x}], \text{x}] + \text{Simp}[\text{Sin}[(\text{d}*e - \text{c}*f)/\text{d}] \quad \text{Int}[\text{Cos}[\text{c}*(\text{f}/\text{d}) + \text{f}*x]/\text{Sqrt}[\text{c} + \text{d}*x], \text{x}], \text{x}] \text{ ;/; } \text{FreeQ}\{\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \ \&\& \ \text{ComplexFreeQ}[\text{f}] \ \&\& \ \text{NeQ}[\text{d}*e - \text{c}*f, 0]$$

rule 3832

$$\text{Int}[\text{Sin}[(\text{d}_.)*((\text{e}_.) + (\text{f}_.)*(x_))^{(2)}], \text{x_Symbol}] \text{ :> } \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(\text{f}*Rt[\text{d}, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*Rt[\text{d}, 2]*(\text{e} + \text{f}*x)], \text{x}] \text{ ;/; } \text{FreeQ}\{\{\text{d}, \text{e}, \text{f}\}, \text{x}\}$$

```
rule 3833 Int[Cos[(d_.)*((e_.) + (f_.)*(x_)^2), x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

```
rule 3897 Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*cos[c + d*x^(k*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]
```

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

method	result
derivativedivides	$\frac{3x^{\frac{1}{6}} \sin\left(a+bx^{\frac{1}{3}}\right)}{b} - \frac{3\sqrt{2}\sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) \right)}{2b^{\frac{3}{2}}}$
default	$\frac{3x^{\frac{1}{6}} \sin\left(a+bx^{\frac{1}{3}}\right)}{b} - \frac{3\sqrt{2}\sqrt{\pi} \left(\cos(a) \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) + \sin(a) \operatorname{FresnelC}\left(\frac{\sqrt{b}\sqrt{2}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) \right)}{2b^{\frac{3}{2}}}$
meijerg	$\frac{3 \cos(a) \sqrt{\pi} \sqrt{2} \left(\frac{x^{\frac{1}{6}} \sqrt{2} (b^2)^{\frac{3}{4}} \sin\left(bx^{\frac{1}{3}}\right)}{2\sqrt{\pi} b} - \frac{(b^2)^{\frac{3}{4}} \operatorname{FresnelS}\left(\frac{\sqrt{b}\sqrt{2}x^{\frac{1}{6}}}{\sqrt{\pi}}\right)}{2b^{\frac{3}{2}}} \right)}{(b^2)^{\frac{3}{4}}} - \frac{3 \sin(a) \sqrt{\pi} \sqrt{2} \left(-\frac{x^{\frac{1}{6}} \sqrt{2} \sqrt{b} \cos\left(bx^{\frac{1}{3}}\right)}{2\sqrt{\pi}} + \dots \right)}{b^{\frac{3}{2}}}$

```
input int(cos(a+b*x^(1/3))/x^(1/2),x,method=_RETURNVERBOSE)
```

```
output 3*x^(1/6)*sin(a+b*x^(1/3))/b-3/2/b^(3/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.79

$$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \frac{3 \left(\sqrt{2}\pi \sqrt{\frac{b}{\pi}} \cos(a) S \left(\sqrt{2}x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) + \sqrt{2}\pi \sqrt{\frac{b}{\pi}} C \left(\sqrt{2}x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(a) - 2bx^{\frac{1}{6}} \sin(bx^{\frac{1}{3}} + a) \right)}{2b^2}$$

input `integrate(cos(a+b*x^(1/3))/x^(1/2),x, algorithm="fricas")`

output `-3/2*(sqrt(2)*pi*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x^(1/6)*sqrt(b/pi)) + sqrt(2)*pi*sqrt(b/pi)*fresnel_cos(sqrt(2)*x^(1/6)*sqrt(b/pi))*sin(a) - 2*b*x^(1/6)*sin(b*x^(1/3) + a))/b^2`

Sympy [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx$$

input `integrate(cos(a+b*x**(1/3))/x**(1/2),x)`

output `Integral(cos(a + b*x**(1/3))/sqrt(x), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.74

$$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \frac{3 \left(\sqrt{2}\sqrt{\pi} \left((-i+1) \cos(a) + (i-1) \sin(a) \right) \operatorname{erf} \left(\sqrt{i}bx^{\frac{1}{6}} \right) + ((i-1) \cos(a) - (i+1) \sin(a)) \operatorname{erf} \left(\sqrt{-i}bx^{\frac{1}{6}} \right) \right)}{8b^3}$$

input `integrate(cos(a+b*x^(1/3))/x^(1/2),x, algorithm="maxima")`

output
$$\frac{3}{8}(\sqrt{2}\sqrt{\pi})\left(\left(-I+1\right)\cos(a)+\left(I-1\right)\sin(a)\right)\operatorname{erf}\left(\sqrt{Ib}x^{\frac{1}{6}}\right)+\left(\left(I-1\right)\cos(a)-\left(I+1\right)\sin(a)\right)\operatorname{erf}\left(\sqrt{-Ib}x^{\frac{1}{6}}\right)+8b^2x^{\frac{1}{6}}\sin\left(bx^{\frac{1}{3}}+a\right)/b^3$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.44

$$\int \frac{\cos(a+b\sqrt[3]{x})}{\sqrt{x}} dx = -\frac{3i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x^{\frac{1}{6}}\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{ia}}{4b\left(-\frac{ib}{|b|}+1\right)\sqrt{|b|}} + \frac{3i\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x^{\frac{1}{6}}\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}\right)e^{-ia}}{4b\left(\frac{ib}{|b|}+1\right)\sqrt{|b|}} - \frac{3ix^{\frac{1}{6}}e^{ibx^{\frac{1}{3}}+ia}}{2b} + \frac{3ix^{\frac{1}{6}}e^{-ibx^{\frac{1}{3}}-ia}}{2b}$$

input `integrate(cos(a+b*x^(1/3))/x^(1/2),x, algorithm="giac")`

output
$$-\frac{3}{4}I\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x^{\frac{1}{6}}\left(-\frac{Ib}{\operatorname{abs}(b)}+1\right)\sqrt{\operatorname{abs}(b)}\right)e^{Ia}/\left(b\left(-\frac{Ib}{\operatorname{abs}(b)}+1\right)\sqrt{\operatorname{abs}(b)}\right)+\frac{3}{4}I\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\frac{1}{2}\sqrt{2}x^{\frac{1}{6}}\left(\frac{Ib}{\operatorname{abs}(b)}+1\right)\sqrt{\operatorname{abs}(b)}\right)e^{-Ia}/\left(b\left(\frac{Ib}{\operatorname{abs}(b)}+1\right)\sqrt{\operatorname{abs}(b)}\right)-\frac{3}{2}Ix^{\frac{1}{6}}e^{Ibx^{\frac{1}{3}}+Ia}/b+\frac{3}{2}Ix^{\frac{1}{6}}e^{-Ibx^{\frac{1}{3}}-Ia}/b$$

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \int \frac{\cos(a + b x^{1/3})}{\sqrt{x}} dx$$

input `int(cos(a + b*x^(1/3))/x^(1/2), x)`output `int(cos(a + b*x^(1/3))/x^(1/2), x)`**Reduce [F]**

$$\int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \int \frac{\cos(x^{1/3}b + a)}{\sqrt{x}} dx$$

input `int(cos(a+b*x^(1/3))/x^(1/2), x)`output `int(cos(x**(1/3)*b + a)/sqrt(x), x)`

3.52
$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{x^{3/2}} dx$$

Optimal result	398
Mathematica [A] (verified)	399
Rubi [A] (verified)	399
Maple [A] (verified)	402
Fricas [A] (verification not implemented)	403
Sympy [F]	403
Maxima [C] (verification not implemented)	404
Giac [F]	404
Mupad [F(-1)]	405
Reduce [F]	405

Optimal result

Integrand size = 16, antiderivative size = 110

$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{x^{3/2}} dx = -\frac{2\cos\left(a+b\sqrt[3]{x}\right)}{\sqrt{x}} - 4b^{3/2}\sqrt{2\pi}\cos(a)\operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) + 4b^{3/2}\sqrt{2\pi}\operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)\sin(a) + \frac{4b\sin\left(a+b\sqrt[3]{x}\right)}{\sqrt[6]{x}}$$

output

```
-2*cos(a+b*x^(1/3))/x^(1/2)-4*b^(3/2)*2^(1/2)*Pi^(1/2)*cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))+4*b^(3/2)*2^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))*sin(a)+4*b*sin(a+b*x^(1/3))/x^(1/6)
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = -\frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt{x}} - 4b^{3/2} \sqrt{2\pi} \cos(a) \operatorname{FresnelC}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right) + 4b^{3/2} \sqrt{2\pi} \operatorname{FresnelS}\left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x}\right) \sin(a) + \frac{4b \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}}$$

input `Integrate[Cos[a + b*x^(1/3)]/x^(3/2), x]`

output `(-2*Cos[a + b*x^(1/3)]/Sqrt[x] - 4*b^(3/2)*Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] + 4*b^(3/2)*Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] + (4*b*Sin[a + b*x^(1/3)]/x^(1/6))`

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.11, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.812$, Rules used = {3897, 3042, 3778, 25, 3042, 3778, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx \\ & \quad \downarrow \text{3897} \\ & 3 \int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} \\ & \quad \downarrow \text{3042} \\ & 3 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{5/6}} d\sqrt[3]{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 3778 \\
& 3 \left(\frac{2}{3} b \int -\frac{\sin(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 25 \\
& 3 \left(-\frac{2}{3} b \int \frac{\sin(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{3} b \int \frac{\sin(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{3} b \left(2b \int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{3} b \left(2b \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3787 \\
& 3 \left(-\frac{2}{3} b \left(2b \left(\cos(a) \int \frac{\cos(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \sin(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{3} b \left(2b \left(\cos(a) \int \frac{\sin(\sqrt[3]{x}b + \frac{\pi}{2})}{\sqrt[6]{x}} d\sqrt[3]{x} - \sin(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3785 \\
& 3 \left(-\frac{2}{3} b \left(2b \left(2 \cos(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} - \sin(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \\
& \downarrow 3786 \\
& 3 \left(-\frac{2}{3} b \left(2b \left(2 \cos(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} - 2 \sin(a) \int \sin(bx^{2/3}) d\sqrt[6]{x} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right)
\end{aligned}$$

↓ 3832

$$3 \left(-\frac{2}{3}b \left(2b \left(2 \cos(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a)}{3} \right)$$

↓ 3833

$$3 \left(-\frac{2}{3}b \left(2b \left(\frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos(a)}{3} \right)$$

input `Int[Cos[a + b*x^(1/3)]/x^(3/2),x]`

output `3*((-2*Cos[a + b*x^(1/3)])/(3*Sqrt[x]) - (2*b*(2*b*((Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/Sqrt[b] - (Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/Sqrt[b]) - (2*Sin[a + b*x^(1/3)]/x^(1/6))))/3)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3785 `Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Cos[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3786 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Sin[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3787 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3897 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.71

method	result
derivativedivides	$-\frac{2 \cos\left(a+b x^{\frac{1}{3}}\right)}{\sqrt{x}} - 4b \left(-\frac{\sin\left(a+b x^{\frac{1}{3}}\right)}{x^{\frac{1}{6}}} + \sqrt{b} \sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2} x^{\frac{1}{6}}}{\sqrt{\pi}}\right) - \sin(a) \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2} x^{\frac{1}{6}}}{\sqrt{\pi}}\right) \right) \right)$
default	$-\frac{2 \cos\left(a+b x^{\frac{1}{3}}\right)}{\sqrt{x}} - 4b \left(-\frac{\sin\left(a+b x^{\frac{1}{3}}\right)}{x^{\frac{1}{6}}} + \sqrt{b} \sqrt{2} \sqrt{\pi} \left(\cos(a) \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2} x^{\frac{1}{6}}}{\sqrt{\pi}}\right) - \sin(a) \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2} x^{\frac{1}{6}}}{\sqrt{\pi}}\right) \right) \right)$
meijerg	$\frac{3 \cos(a) \sqrt{\pi} \sqrt{2} (b^2)^{\frac{3}{4}} \left(-\frac{8 \sqrt{2} \cos\left(b x^{\frac{1}{3}}\right)}{3 \sqrt{\pi} \sqrt{x} (b^2)^{\frac{3}{4}}} + \frac{16 \sqrt{2} b \sin\left(b x^{\frac{1}{3}}\right)}{3 \sqrt{\pi} x^{\frac{1}{6}} (b^2)^{\frac{3}{4}}} - \frac{32 b^{\frac{3}{2}} \operatorname{FresnelC}\left(\frac{\sqrt{b} \sqrt{2} x^{\frac{1}{6}}}{\sqrt{\pi}}\right)}{3 (b^2)^{\frac{3}{4}}} \right)}{8} - \frac{3 \sin(a) \sqrt{\pi} \sqrt{2} b^{\frac{3}{2}} \left(-\frac{8 \sqrt{2} \sin\left(b x^{\frac{1}{3}}\right)}{3 \sqrt{\pi} \sqrt{x} (b^2)^{\frac{3}{4}}} + \frac{16 \sqrt{2} b \cos\left(b x^{\frac{1}{3}}\right)}{3 \sqrt{\pi} x^{\frac{1}{6}} (b^2)^{\frac{3}{4}}} - \frac{32 b^{\frac{3}{2}} \operatorname{FresnelS}\left(\frac{\sqrt{b} \sqrt{2} x^{\frac{1}{6}}}{\sqrt{\pi}}\right)}{3 (b^2)^{\frac{3}{4}}} \right)}{8}$

input `int(cos(a+b*x^(1/3))/x^(3/2),x,method=_RETURNVERBOSE)`

output `-2*cos(a+b*x^(1/3))/x^(1/2)-4*b*(-1/x^(1/6)*sin(a+b*x^(1/3))+b^(1/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))-sin(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.87

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = \frac{2 \left(2\sqrt{2}\pi b x \sqrt{\frac{b}{\pi}} \cos(a) C\left(\sqrt{2}x^{1/6} \sqrt{\frac{b}{\pi}}\right) - 2\sqrt{2}\pi b x \sqrt{\frac{b}{\pi}} S\left(\sqrt{2}x^{1/6} \sqrt{\frac{b}{\pi}}\right) \sin(a) - 2bx^{5/6} \sin\left(bx^{1/3} + a\right) + \sqrt{x} \right)}{x}$$

input `integrate(cos(a+b*x^(1/3))/x^(3/2),x, algorithm="fricas")`

output `-2*(2*sqrt(2)*pi*b*x*sqrt(b/pi)*cos(a)*fresnel_cos(sqrt(2)*x^(1/6)*sqrt(b/pi)) - 2*sqrt(2)*pi*b*x*sqrt(b/pi)*fresnel_sin(sqrt(2)*x^(1/6)*sqrt(b/pi))*sin(a) - 2*b*x^(5/6)*sin(b*x^(1/3) + a) + sqrt(x)*cos(b*x^(1/3) + a))/x`

Sympy [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx$$

input `integrate(cos(a+b*x**(1/3))/x**(3/2),x)`

output `Integral(cos(a + b*x**(1/3))/x**(3/2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.67

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = \frac{3 \left(\left((i-1) \sqrt{2} \Gamma\left(-\frac{3}{2}, i b x^{\frac{1}{3}}\right) - (i+1) \sqrt{2} \Gamma\left(-\frac{3}{2}, -i b x^{\frac{1}{3}}\right) \right) \cos(a) + \left((i+1) \sqrt{2} \Gamma\left(-\frac{3}{2}, i b x^{\frac{1}{3}}\right) - (i-1) \sqrt{2} \Gamma\left(-\frac{3}{2}, -i b x^{\frac{1}{3}}\right) \right) \sin(a) \right) \sqrt{b x^{\frac{1}{3}}}}{4 x^{\frac{1}{6}}}$$

input `integrate(cos(a+b*x^(1/3))/x^(3/2),x, algorithm="maxima")`

output `-3/4*((I - 1)*sqrt(2)*gamma(-3/2, I*b*x^(1/3)) - (I + 1)*sqrt(2)*gamma(-3/2, -I*b*x^(1/3)))*cos(a) + ((I + 1)*sqrt(2)*gamma(-3/2, I*b*x^(1/3)) - (I - 1)*sqrt(2)*gamma(-3/2, -I*b*x^(1/3)))*sin(a)*sqrt(b*x^(1/3))*b/x^(1/6)`

Giac [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{3}{2}}} dx$$

input `integrate(cos(a+b*x^(1/3))/x^(3/2),x, algorithm="giac")`

output `integrate(cos(b*x^(1/3) + a)/x^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos(a + bx^{1/3})}{x^{3/2}} dx$$

input `int(cos(a + b*x^(1/3))/x^(3/2), x)`output `int(cos(a + b*x^(1/3))/x^(3/2), x)`**Reduce [F]**

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} dx = \frac{\sqrt{x} \left(\int \frac{\cos(x^{1/3}b+a)}{\sqrt{x}x} dx \right) + \sqrt{x} \left(\int \frac{1}{\sqrt{x}x} dx \right) + 2}{\sqrt{x}}$$

input `int(cos(a+b*x^(1/3))/x^(3/2), x)`output `(sqrt(x)*int(cos(x**(1/3)*b + a)/(sqrt(x)*x), x) + sqrt(x)*int(1/(sqrt(x)*x), x) + 2)/sqrt(x)`

3.53 $\int \frac{\cos(a+b\sqrt[3]{x})}{x^{5/2}} dx$

Optimal result	406
Mathematica [A] (verified)	407
Rubi [A] (verified)	407
Maple [A] (verified)	412
Fricas [A] (verification not implemented)	414
Sympy [F]	414
Maxima [C] (verification not implemented)	415
Giac [F]	415
Mupad [F(-1)]	415
Reduce [F]	416

Optimal result

Integrand size = 16, antiderivative size = 184

$$\int \frac{\cos(a+b\sqrt[3]{x})}{x^{5/2}} dx = -\frac{2\cos(a+b\sqrt[3]{x})}{3x^{3/2}} + \frac{8b^2\cos(a+b\sqrt[3]{x})}{105x^{5/6}} - \frac{32b^4\cos(a+b\sqrt[3]{x})}{315\sqrt[6]{x}} - \frac{32}{315}b^{9/2}\sqrt{2\pi}\cos(a)\text{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right) - \frac{32}{315}b^{9/2}\sqrt{2\pi}\text{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)\sin(a) + \frac{4b\sin(a+b\sqrt[3]{x})}{21x^{7/6}}$$

output

```
-2/3*cos(a+b*x^(1/3))/x^(3/2)+8/105*b^2*cos(a+b*x^(1/3))/x^(5/6)-32/315*b^4*cos(a+b*x^(1/3))/x^(1/6)-32/315*b^(9/2)*2^(1/2)*Pi^(1/2)*cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))-32/315*b^(9/2)*2^(1/2)*Pi^(1/2)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))*sin(a)+4/21*b*sin(a+b*x^(1/3))/x^(7/6)-16/315*b^3*sin(a+b*x^(1/3))/x^(1/2)
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.98

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx =$$

$$2 \left(105 \cos(a + b\sqrt[3]{x}) - 12b^2 x^{2/3} \cos(a + b\sqrt[3]{x}) + 16b^4 x^{4/3} \cos(a + b\sqrt[3]{x}) + 16b^{9/2} \sqrt{2\pi} x^{3/2} \cos(a) \operatorname{FresnelS}[\sqrt{b} \sqrt{2/\pi} x^{1/6}] + 16b^{9/2} \sqrt{2\pi} x^{3/2} \operatorname{FresnelC}[\sqrt{b} \sqrt{2/\pi} x^{1/6}] \sin(a) - 30b x^{1/3} \sin(a + b\sqrt[3]{x}) + 8b^3 x \sin(a + b\sqrt[3]{x}) \right) / (315 x^{3/2})$$

input `Integrate[Cos[a + b*x^(1/3)]/x^(5/2), x]`

output `(-2*(105*Cos[a + b*x^(1/3)] - 12*b^2*x^(2/3)*Cos[a + b*x^(1/3)] + 16*b^4*x^(4/3)*Cos[a + b*x^(1/3)] + 16*b^(9/2)*Sqrt[2*Pi]*x^(3/2)*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] + 16*b^(9/2)*Sqrt[2*Pi]*x^(3/2)*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] - 30*b*x^(1/3)*Sin[a + b*x^(1/3)] + 8*b^3*x*Ssin[a + b*x^(1/3)])/(315*x^(3/2))`

Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.07, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 1.312$, Rules used = {3897, 3042, 3778, 25, 3042, 3778, 3042, 3778, 25, 3042, 3778, 3042, 3778, 25, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx$$

$$\downarrow \text{3897}$$

$$3 \int \frac{\cos(a + b\sqrt[3]{x})}{x^{11/6}} d\sqrt[3]{x}$$

$$\downarrow \text{3042}$$

$$3 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{11/6}} d\sqrt[3]{x}$$

$$\begin{aligned}
& \downarrow 3778 \\
& 3 \left(\frac{2}{9} b \int -\frac{\sin(a + b\sqrt[3]{x})}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 25 \\
& 3 \left(-\frac{2}{9} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{9} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{9} b \left(\frac{2}{7} b \int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{9} b \left(\frac{2}{7} b \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{9} b \left(\frac{2}{7} b \left(\frac{2}{5} b \int -\frac{\sin(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 25 \\
& 3 \left(-\frac{2}{9} b \left(\frac{2}{7} b \left(-\frac{2}{5} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{9} b \left(\frac{2}{7} b \left(-\frac{2}{5} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{9} b \left(\frac{2}{7} b \left(-\frac{2}{5} b \left(\frac{2}{3} b \int \frac{\cos(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{9x^{3/2}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}}
\end{aligned}$$

↓ 3042

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{7x^{7/6}} \right) \right)$$

↓ 3778

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(2b \int -\frac{\sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) \right)$$

↓ 25

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \int \frac{\sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) \right)$$

↓ 3042

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \int \frac{\sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{5x^{5/6}} \right) \right)$$

↓ 3787

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \left(\sin(a) \int \frac{\cos(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} + \cos(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) \right)$$

↓ 3042

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \left(\sin(a) \int \frac{\sin(\sqrt[3]{x}b + \frac{\pi}{2})}{\sqrt[6]{x}} d\sqrt[3]{x} + \cos(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) \right)$$

↓ 3785

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \left(2 \sin(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} + \cos(a) \int \frac{\sin(b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) \right)$$

↓ 3786

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \left(2 \sin(a) \int \cos(bx^{2/3}) d\sqrt[6]{x} + 2 \cos(a) \int \sin(bx^{2/3}) d\sqrt[6]{x} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) \right)$$

↓ 3832

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \left(2 \sin(a) \int \cos \left(bx^{2/3} \right) d\sqrt[6]{x} + \frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} \right) - \frac{2 \cos(a)}{\sqrt[6]{x}} \right) \right) \right) \right) \right)$$

↓ 3833

$$3 \left(-\frac{2}{9}b \left(\frac{2}{7}b \left(-\frac{2}{5}b \left(\frac{2}{3}b \left(-2b \left(\frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} + \frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} \right) - \frac{2 \cos(a)}{\sqrt[6]{x}} \right) \right) \right) \right) \right)$$

input `Int[Cos[a + b*x^(1/3)]/x^(5/2),x]`

output `3*((-2*Cos[a + b*x^(1/3)])/(9*x^(3/2)) - (2*b*((-2*Sin[a + b*x^(1/3)])/(7*x^(7/6)) + (2*b*((-2*Cos[a + b*x^(1/3)])/(5*x^(5/6)) - (2*b*((2*b*((-2*Cos[a + b*x^(1/3)])/x^(1/6) - 2*b*((Sqrt[2*Pi]*Cos[a]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/Sqrt[b] + (Sqrt[2*Pi]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/Sqrt[b])))/3 - (2*Sin[a + b*x^(1/3)]/(3*Sqrt[x])))/5)/7)/9)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3785 `Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := S
imp[2/d Subst[Int[Cos[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c,
d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3786 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d
Subst[Int[Sin[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f
}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3787 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cos
[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] + Simp[Sin[(
d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d
, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[
d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[
d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3897 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol
] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a +
b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]`

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.70

method	result
derivativeldivides	$\frac{2 \cos\left(a+bx^{\frac{1}{3}}\right)}{3x^{\frac{3}{2}}}$ $4b \left[-\frac{\sin\left(a+bx^{\frac{1}{3}}\right)}{7x^{\frac{7}{6}}} + \frac{\cos\left(a+bx^{\frac{1}{3}}\right)}{5x^{\frac{5}{6}}} + \frac{\sin\left(a+bx^{\frac{1}{3}}\right)}{3\sqrt{x}} + \frac{\cos\left(a+bx^{\frac{1}{3}}\right)}{x^{\frac{1}{6}}} - \sqrt{b} \sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{b} x^{\frac{1}{6}}}{\sqrt{2}}\right) \right]$
default	$\frac{2 \cos\left(a+bx^{\frac{1}{3}}\right)}{3x^{\frac{3}{2}}}$ $4b \left[-\frac{\sin\left(a+bx^{\frac{1}{3}}\right)}{7x^{\frac{7}{6}}} + \frac{\cos\left(a+bx^{\frac{1}{3}}\right)}{5x^{\frac{5}{6}}} + \frac{\sin\left(a+bx^{\frac{1}{3}}\right)}{3\sqrt{x}} + \frac{\cos\left(a+bx^{\frac{1}{3}}\right)}{x^{\frac{1}{6}}} - \sqrt{b} \sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{b} x^{\frac{1}{6}}}{\sqrt{2}}\right) \right]$
meijerg	$3 \cos(a) \sqrt{\pi} \sqrt{2} (b^2)^{\frac{9}{4}} \left(-\frac{64\sqrt{2} \left(\frac{16x^{\frac{4}{3}} b^4}{105} - \frac{4x^{\frac{2}{3}} b^2}{35} + 1 \right) \cos\left(bx^{\frac{1}{3}}\right)}{9\sqrt{\pi} x^{\frac{3}{2}} (b^2)^{\frac{9}{4}}} + \frac{128\sqrt{2} b \left(-4x^{\frac{2}{3}} b^2 + 15 \right) \sin\left(bx^{\frac{1}{3}}\right)}{945\sqrt{\pi} x^{\frac{7}{6}} (b^2)^{\frac{9}{4}}} - \frac{2048b^{\frac{9}{2}} \operatorname{FresnelS}\left(\frac{\sqrt{b} x^{\frac{1}{6}}}{\sqrt{2}}\right)}{945 (b^2)^{\frac{9}{4}}} \right)$

input `int(cos(a+b*x^(1/3))/x^(5/2),x,method=_RETURNVERBOSE)`

output `-2/3*cos(a+b*x^(1/3))/x^(3/2)-4/3*b*(-1/7/x^(7/6)*sin(a+b*x^(1/3))+2/7*b*(-1/5*cos(a+b*x^(1/3))/x^(5/6)-2/5*b*(-1/3/x^(1/2)*sin(a+b*x^(1/3))+2/3*b*(-1/x^(1/6)*cos(a+b*x^(1/3))-b^(1/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))+sin(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))))))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.73

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx = \frac{2 \left(16 \sqrt{2} \pi b^4 x^2 \sqrt{\frac{b}{\pi}} \cos(a) S \left(\sqrt{2} x^{1/6} \sqrt{\frac{b}{\pi}} \right) + 16 \sqrt{2} \pi b^4 x^2 \sqrt{\frac{b}{\pi}} C \left(\sqrt{2} x^{1/6} \sqrt{\frac{b}{\pi}} \right) \sin(a) + \left(16 b^4 x^{11/6} - 12 b^2 x^{7/6} \right) \cos(b x^{1/3} + a) + 2 \left(4 b^3 x^{3/2} - 15 b x^{5/6} \right) \sin(b x^{1/3} + a) \right)}{315 x^2}$$

input `integrate(cos(a+b*x^(1/3))/x^(5/2),x, algorithm="fricas")`

output `-2/315*(16*sqrt(2)*pi*b^4*x^2*sqrt(b/pi)*cos(a)*fresnel_sin(sqrt(2)*x^(1/6)*sqrt(b/pi)) + 16*sqrt(2)*pi*b^4*x^2*sqrt(b/pi)*fresnel_cos(sqrt(2)*x^(1/6)*sqrt(b/pi))*sin(a) + (16*b^4*x^(11/6) - 12*b^2*x^(7/6) + 105*sqrt(x))*cos(b*x^(1/3) + a) + 2*(4*b^3*x^(3/2) - 15*b*x^(5/6))*sin(b*x^(1/3) + a)/x^2`

Sympy [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx = \int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx$$

input `integrate(cos(a+b*x**(1/3))/x**(5/2),x)`

output `Integral(cos(a + b*x**(1/3))/x**(5/2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.41

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx = \frac{3 \left(\left(-(i+1) \sqrt{2} \Gamma\left(-\frac{9}{2}, i b x^{\frac{1}{3}}\right) + (i-1) \sqrt{2} \Gamma\left(-\frac{9}{2}, -i b x^{\frac{1}{3}}\right) \right) \cos(a) + \left((i-1) \sqrt{2} \Gamma\left(-\frac{9}{2}, i b x^{\frac{1}{3}}\right) - (i+1) \sqrt{2} \Gamma\left(-\frac{9}{2}, -i b x^{\frac{1}{3}}\right) \right) \sin(a) \right) \sqrt{b x^{\frac{1}{3}}}}{4 x^{\frac{1}{6}}}$$

input `integrate(cos(a+b*x^(1/3))/x^(5/2),x, algorithm="maxima")`

output `3/4*((-(I + 1)*sqrt(2)*gamma(-9/2, I*b*x^(1/3)) + (I - 1)*sqrt(2)*gamma(-9/2, -I*b*x^(1/3)))*cos(a) + ((I - 1)*sqrt(2)*gamma(-9/2, I*b*x^(1/3)) - (I + 1)*sqrt(2)*gamma(-9/2, -I*b*x^(1/3)))*sin(a))*sqrt(b*x^(1/3))*b^4/x^(1/6)`

Giac [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx = \int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{5}{2}}} dx$$

input `integrate(cos(a+b*x^(1/3))/x^(5/2),x, algorithm="giac")`

output `integrate(cos(b*x^(1/3) + a)/x^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx = \int \frac{\cos(a + b x^{1/3})}{x^{5/2}} dx$$

input `int(cos(a + b*x^(1/3))/x^(5/2),x)`

output `int(cos(a + b*x^(1/3))/x^(5/2), x)`

Reduce [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/2}} dx = \frac{3\sqrt{x} \left(\int \frac{\cos(x^{1/3}b+a)}{\sqrt{x}x^2} dx \right) x + 3\sqrt{x} \left(\int \frac{1}{\sqrt{x}x^2} dx \right) x + 2}{3\sqrt{x}x}$$

input `int(cos(a+b*x^(1/3))/x^(5/2),x)`

output `(3*sqrt(x)*int(cos(x**(1/3)*b + a)/(sqrt(x)*x**2),x)*x + 3*sqrt(x)*int(1/(sqrt(x)*x**2),x)*x + 2)/(3*sqrt(x)*x)`

3.54
$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{x^{7/2}} dx$$

Optimal result	417
Mathematica [A] (verified)	418
Rubi [A] (verified)	418
Maple [A] (verified)	423
Fricas [A] (verification not implemented)	425
Sympy [F]	425
Maxima [C] (verification not implemented)	426
Giac [F]	426
Mupad [F(-1)]	426
Reduce [F]	427

Optimal result

Integrand size = 16, antiderivative size = 250

$$\int \frac{\cos\left(a+b\sqrt[3]{x}\right)}{x^{7/2}} dx = -\frac{2\cos\left(a+b\sqrt[3]{x}\right)}{5x^{5/2}} + \frac{8b^2\cos\left(a+b\sqrt[3]{x}\right)}{715x^{11/6}}$$

$$-\frac{32b^4\cos\left(a+b\sqrt[3]{x}\right)}{45045x^{7/6}} + \frac{128b^6\cos\left(a+b\sqrt[3]{x}\right)}{675675\sqrt{x}}$$

$$+\frac{256b^{15/2}\sqrt{2\pi}\cos(a)\operatorname{FresnelC}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)}{675675}$$

$$-\frac{256b^{15/2}\sqrt{2\pi}\operatorname{FresnelS}\left(\sqrt{b}\sqrt{\frac{2}{\pi}}\sqrt[6]{x}\right)\sin(a)}{675675} + \frac{4b\sin\left(a+b\sqrt[3]{x}\right)}{65x^{13/6}}$$

$$-\frac{16b^3\sin\left(a+b\sqrt[3]{x}\right)}{6435x^{3/2}} + \frac{64b^5\sin\left(a+b\sqrt[3]{x}\right)}{225225x^{5/6}} - \frac{256b^7\sin\left(a+b\sqrt[3]{x}\right)}{675675\sqrt[6]{x}}$$

output

```
-2/5*cos(a+b*x^(1/3))/x^(5/2)+8/715*b^2*cos(a+b*x^(1/3))/x^(11/6)-32/45045
*b^4*cos(a+b*x^(1/3))/x^(7/6)+128/675675*b^6*cos(a+b*x^(1/3))/x^(1/2)+256/
675675*b^(15/2)*2^(1/2)*Pi^(1/2)*cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*
x^(1/6))-256/675675*b^(15/2)*2^(1/2)*Pi^(1/2)*FresnelS(b^(1/2)*2^(1/2)/Pi^(
1/2)*x^(1/6))*sin(a)+4/65*b*sin(a+b*x^(1/3))/x^(13/6)-16/6435*b^3*sin(a+b
*x^(1/3))/x^(3/2)+64/225225*b^5*sin(a+b*x^(1/3))/x^(5/6)-256/675675*b^7*si
n(a+b*x^(1/3))/x^(1/6)
```

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.95

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{2(-135135 \cos(a + b\sqrt[3]{x}) + 3780b^2x^{2/3} \cos(a + b\sqrt[3]{x}) - 240b^4x^{4/3} \cos(a + b\sqrt[3]{x}) + 240b^6x^{2/3} \cos(a + b\sqrt[3]{x}) - 64b^8x^{2/3} \cos(a + b\sqrt[3]{x}) + 128b^{10}x^{2/3} \cos(a + b\sqrt[3]{x}) - 128b^{15/2} \sqrt{2\pi} x^{5/2} \cos[a] \operatorname{FresnelC}[\sqrt{b} \sqrt{2\pi} x^{1/6}] - 128b^{15/2} \sqrt{2\pi} x^{5/2} \operatorname{FresnelS}[\sqrt{b} \sqrt{2\pi} x^{1/6}] \sin[a] + 20790b^2x^{1/3} \sin[a + b\sqrt[3]{x}] - 840b^3x \sin[a + b\sqrt[3]{x}] + 96b^5x^{5/3} \sin[a + b\sqrt[3]{x}] - 128b^7x^{7/3} \sin[a + b\sqrt[3]{x}])}{675675x^{5/2}}$$

input `Integrate[Cos[a + b*x^(1/3)]/x^(7/2), x]`

output `(2*(-135135*Cos[a + b*x^(1/3)] + 3780*b^2*x^(2/3)*Cos[a + b*x^(1/3)] - 240*b^4*x^(4/3)*Cos[a + b*x^(1/3)] + 64*b^6*x^2*Cos[a + b*x^(1/3)] + 128*b^(15/2)*Sqrt[2*Pi]*x^(5/2)*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)] - 128*b^(15/2)*Sqrt[2*Pi]*x^(5/2)*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a] + 20790*b*x^(1/3)*Sin[a + b*x^(1/3)] - 840*b^3*x*Ssin[a + b*x^(1/3)] + 96*b^5*x^(5/3)*Sin[a + b*x^(1/3)] - 128*b^7*x^(7/3)*Sin[a + b*x^(1/3)])/(675675*x^(5/2))`

Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.09, number of steps used = 29, number of rules used = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 1.750$, Rules used = {3897, 3042, 3778, 25, 3042, 3778, 3042, 3778, 25, 3042, 3778, 3042, 3778, 25, 3042, 3778, 3042, 3778, 25, 3042, 3778, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx \\ & \quad \downarrow \text{3897} \\ & 3 \int \frac{\cos(a + b\sqrt[3]{x})}{x^{17/6}} d\sqrt[3]{x} \\ & \quad \downarrow \text{3042} \\ & 3 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{17/6}} d\sqrt[3]{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 3778 \\
& 3 \left(\frac{2}{15} b \int -\frac{\sin(a + b\sqrt[3]{x})}{x^{5/2}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 25 \\
& 3 \left(-\frac{2}{15} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{5/2}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{15} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{5/2}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{15} b \left(\frac{2}{13} b \int \frac{\cos(a + b\sqrt[3]{x})}{x^{13/6}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{15} b \left(\frac{2}{13} b \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{13/6}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{15} b \left(\frac{2}{13} b \left(\frac{2}{11} b \int -\frac{\sin(a + b\sqrt[3]{x})}{x^{11/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 25 \\
& 3 \left(-\frac{2}{15} b \left(\frac{2}{13} b \left(-\frac{2}{11} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{11/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 3042 \\
& 3 \left(-\frac{2}{15} b \left(\frac{2}{13} b \left(-\frac{2}{11} b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{11/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{15x^{5/2}} \right) \\
& \downarrow 3778 \\
& 3 \left(-\frac{2}{15} b \left(\frac{2}{13} b \left(-\frac{2}{11} b \left(\frac{2}{9} b \int \frac{\cos(a + b\sqrt[3]{x})}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) \right)
\end{aligned}$$

↓ 3042

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{13x^{13/6}} \right) \right)$$

↓ 3778

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(\frac{2}{7}b \int -\frac{\sin(a + b\sqrt[3]{x})}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) \right)$$

↓ 25

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) \right)$$

↓ 3042

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \int \frac{\sin(a + b\sqrt[3]{x})}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{11x^{11/6}} \right) \right)$$

↓ 3778

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \left(\frac{2}{5}b \int \frac{\cos(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \right) \right)$$

↓ 3042

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \left(\frac{2}{5}b \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \sin(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{9x^{3/2}} \right) \right) \right)$$

↓ 3778

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \left(\frac{2}{5}b \left(\frac{2}{3}b \int -\frac{\sin(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) \right) \right) \right)$$

↓ 25

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \left(\frac{2}{5}b \left(-\frac{2}{3}b \int \frac{\sin(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) - \frac{2 \sin(a + b\sqrt[3]{x})}{5x^{5/6}} \right) - \frac{2 \cos(a + b\sqrt[3]{x})}{7x^{7/6}} \right) \right) \right) \right)$$

↓ 3833

$$3 \left(-\frac{2}{15}b \left(\frac{2}{13}b \left(-\frac{2}{11}b \left(\frac{2}{9}b \left(-\frac{2}{7}b \left(\frac{2}{5}b \left(-\frac{2}{3}b \left(2b \left(\frac{\sqrt{2\pi} \cos(a) \operatorname{FresnelC} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} - \frac{\sqrt{2\pi} \sin(a) \operatorname{FresnelS} \left(\sqrt{b} \sqrt{\frac{2}{\pi}} \sqrt[6]{x} \right)}{\sqrt{b}} \right) \right) \right) \right) \right) \right) \right) \right) \right)$$

input `Int[Cos[a + b*x^(1/3)]/x^(7/2),x]`

output `3*((-2*Cos[a + b*x^(1/3)])/(15*x^(5/2)) - (2*b*((-2*Sin[a + b*x^(1/3)])/(13*x^(13/6)) + (2*b*((-2*Cos[a + b*x^(1/3)])/(11*x^(11/6)) - (2*b*((-2*Sin[a + b*x^(1/3)])/(9*x^(3/2)) + (2*b*((-2*Cos[a + b*x^(1/3)])/(7*x^(7/6)) - (2*b*((-2*Sin[a + b*x^(1/3)])/(5*x^(5/6)) + (2*b*((-2*Cos[a + b*x^(1/3)])/(3*Sqrt[x]) - (2*b*(2*b*((Sqrt[2*Pi]*Cos[a]*FresnelC[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)])/Sqrt[b] - (Sqrt[2*Pi]*FresnelS[Sqrt[b]*Sqrt[2/Pi]*x^(1/6)]*Sin[a])/Sqrt[b]) - (2*Sin[a + b*x^(1/3)]/x^(1/6))))/3))/5))/7))/9))/11))/13))/15)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3785 `Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Cos[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3786 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d
Subst[Int[Sin[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f
, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]`

rule 3787 `Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cos
[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] + Simp[Sin[(
d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d
, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[
d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_)^2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[
d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3897 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol
] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a +
b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]`

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.72

method	result
	$2b \frac{\cos\left(a + b x^{\frac{1}{3}}\right)}{7x^{\frac{7}{6}}}$ $2b \frac{\sin\left(a + b x^{\frac{1}{3}}\right)}{9x^{\frac{3}{2}}} +$ $2b \frac{\cos\left(a + b x^{\frac{1}{3}}\right)}{11x^{\frac{11}{6}}}$ $2b \frac{\sin\left(a + b x^{\frac{1}{3}}\right)}{5x}$

input `int(cos(a+b*x^(1/3))/x^(7/2),x,method=_RETURNVERBOSE)`

output `-2/5*cos(a+b*x^(1/3))/x^(5/2)-4/5*b*(-1/13/x^(13/6)*sin(a+b*x^(1/3))+2/13*b*(-1/11*cos(a+b*x^(1/3))/x^(11/6)-2/11*b*(-1/9/x^(3/2)*sin(a+b*x^(1/3))+2/9*b*(-1/7/x^(7/6)*cos(a+b*x^(1/3))-2/7*b*(-1/5/x^(5/6)*sin(a+b*x^(1/3))+2/5*b*(-1/3*cos(a+b*x^(1/3))/x^(1/2)-2/3*b*(-1/x^(1/6)*sin(a+b*x^(1/3))+b^(1/2)*2^(1/2)*Pi^(1/2)*(cos(a)*FresnelC(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))-sin(a)*FresnelS(b^(1/2)*2^(1/2)/Pi^(1/2)*x^(1/6))))))))))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.66

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{2 \left(128 \sqrt{2} \pi b^7 x^3 \sqrt{\frac{b}{\pi}} \cos(a) C \left(\sqrt{2} x^{1/6} \sqrt{\frac{b}{\pi}} \right) - 128 \sqrt{2} \pi b^7 x^3 \sqrt{\frac{b}{\pi}} S \left(\sqrt{2} x^{1/6} \sqrt{\frac{b}{\pi}} \right) \sin(a) \right)}{x^{7/2}}$$

input `integrate(cos(a+b*x^(1/3))/x^(7/2),x, algorithm="fricas")`

output `2/675675*(128*sqrt(2)*pi*b^7*x^3*sqrt(b/pi)*cos(a)*fresnel_cos(sqrt(2)*x^(1/6)*sqrt(b/pi)) - 128*sqrt(2)*pi*b^7*x^3*sqrt(b/pi)*fresnel_sin(sqrt(2)*x^(1/6)*sqrt(b/pi))*sin(a) - (240*b^4*x^(11/6) - 3780*b^2*x^(7/6) - (64*b^6*x^2 - 135135)*sqrt(x))*cos(b*x^(1/3) + a) + 2*(48*b^5*x^(13/6) - 420*b^3*x^(3/2) - (64*b^7*x^2 - 10395*b)*x^(5/6))*sin(b*x^(1/3) + a))/x^3`

Sympy [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx$$

input `integrate(cos(a+b*x**(1/3))/x**(7/2),x)`

output `Integral(cos(a + b*x**(1/3))/x**(7/2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.30

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{3 \left(\left((i-1) \sqrt{2} \Gamma\left(-\frac{15}{2}, i b x^{\frac{1}{3}}\right) - (i+1) \sqrt{2} \Gamma\left(-\frac{15}{2}, -i b x^{\frac{1}{3}}\right) \right) \cos(a) + \left((i+1) \sqrt{2} \Gamma\left(-\frac{15}{2}, i b x^{\frac{1}{3}}\right) - (i-1) \sqrt{2} \Gamma\left(-\frac{15}{2}, -i b x^{\frac{1}{3}}\right) \right) \sin(a) \right)}{4 x^{\frac{1}{6}}}$$

input `integrate(cos(a+b*x^(1/3))/x^(7/2),x, algorithm="maxima")`

output `3/4*(((I - 1)*sqrt(2)*gamma(-15/2, I*b*x^(1/3)) - (I + 1)*sqrt(2)*gamma(-15/2, -I*b*x^(1/3)))*cos(a) + ((I + 1)*sqrt(2)*gamma(-15/2, I*b*x^(1/3)) - (I - 1)*sqrt(2)*gamma(-15/2, -I*b*x^(1/3)))*sin(a))*sqrt(b*x^(1/3))*b^7/x^(1/6)`

Giac [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)}{x^{\frac{7}{2}}} dx$$

input `integrate(cos(a+b*x^(1/3))/x^(7/2),x, algorithm="giac")`

output `integrate(cos(b*x^(1/3) + a)/x^(7/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \int \frac{\cos(a + b x^{1/3})}{x^{7/2}} dx$$

input `int(cos(a + b*x^(1/3))/x^(7/2),x)`

output `int(cos(a + b*x^(1/3))/x^(7/2), x)`

Reduce [F]

$$\int \frac{\cos(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{5\sqrt{x} \left(\int \frac{\cos(x^{1/3}b+a)}{\sqrt{x}x^3} dx \right) x^2 + 5\sqrt{x} \left(\int \frac{1}{\sqrt{x}x^3} dx \right) x^2 + 2}{5\sqrt{x}x^2}$$

input `int(cos(a+b*x^(1/3))/x^(7/2),x)`

output `(5*sqrt(x)*int(cos(x**(1/3)*b + a)/(sqrt(x)*x**3),x)*x**2 + 5*sqrt(x)*int(1/(sqrt(x)*x**3),x)*x**2 + 2)/(5*sqrt(x)*x**2)`

3.55 $\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx$

Optimal result	428
Mathematica [A] (verified)	429
Rubi [A] (verified)	429
Maple [A] (verified)	435
Fricas [A] (verification not implemented)	437
Sympy [F]	437
Maxima [C] (verification not implemented)	438
Giac [C] (verification not implemented)	439
Mupad [F(-1)]	440
Reduce [F]	440

Optimal result

Integrand size = 18, antiderivative size = 311

$$\begin{aligned}
 \int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = & -\frac{135135\sqrt{x}}{4096b^6} + \frac{3861x^{7/6}}{256b^4} - \frac{39x^{11/6}}{16b^2} \\
 & + \frac{x^{5/2}}{5} + \frac{135135\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{2048b^6} - \frac{3861x^{7/6} \cos^2(a + b\sqrt[3]{x})}{128b^4} \\
 & + \frac{39x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{405405\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{32768b^{15/2}} \\
 & + \frac{405405\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) \sin(2a)}{32768b^{15/2}} \\
 & + \frac{27027x^{5/6} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{512b^5} \\
 & - \frac{429x^{3/2} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{32b^3} \\
 & + \frac{3x^{13/6} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{2b} - \frac{405405\sqrt[6]{x} \sin(2a + 2b\sqrt[3]{x})}{16384b^7}
 \end{aligned}$$

output

```
-135135/4096*x^(1/2)/b^6+3861/256*x^(7/6)/b^4-39/16*x^(11/6)/b^2+1/5*x^(5/2)+135135/2048*x^(1/2)*cos(a+b*x^(1/3))^2/b^6-3861/128*x^(7/6)*cos(a+b*x^(1/3))^2/b^4+39/8*x^(11/6)*cos(a+b*x^(1/3))^2/b^2+405405/32768*Pi^(1/2)*cos(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))/b^(15/2)+405405/32768*Pi^(1/2)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))*sin(2*a)/b^(15/2)+27027/512*x^(5/6)*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/b^5-429/32*x^(3/2)*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/b^3+3/2*x^(13/6)*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/b-405405/16384*x^(1/6)*sin(2*a+2*b*x^(1/3))/b^7
```

Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.56

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \frac{2027025\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) + 2027025\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) \sin(2a) + 2\sqrt{b}\sqrt[6]{x}}{163840b^{15/2}}$$

input

```
Integrate[x^(3/2)*Cos[a + b*x^(1/3)]^2,x]
```

output

```
(2027025*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]] + 2027025*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 2*Sqrt[b]*x^(1/6)*(16384*b^7*x^(7/3) + 780*(3465*b*x^(1/3) - 1584*b^3*x + 256*b^5*x^(5/3))*Cos[2*(a + b*x^(1/3))] + 15*(-135135 + 144144*b^2*x^(2/3) - 36608*b^4*x^(4/3) + 4096*b^6*x^2)*Sin[2*(a + b*x^(1/3))]))/(163840*b^(15/2))
```

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.722$, Rules used = {3897, 3042, 3792, 15, 3042, 3792, 15, 3042, 3792, 15, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^{3/2} \cos^2(a + b\sqrt[3]{x}) \, dx \\
 & \quad \downarrow \text{3897} \\
 & 3 \int x^{13/6} \cos^2(a + b\sqrt[3]{x}) \, d\sqrt[3]{x} \\
 & \quad \downarrow \text{3042} \\
 & 3 \int x^{13/6} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right)^2 \, d\sqrt[3]{x} \\
 & \quad \downarrow \text{3792} \\
 & 3 \left(-\frac{143 \int x^{3/2} \cos^2(a + b\sqrt[3]{x}) \, d\sqrt[3]{x}}{16b^2} + \frac{1}{2} \int x^{13/6} \, d\sqrt[3]{x} + \frac{13x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{13/6} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} \right) \\
 & \quad \downarrow \text{15} \\
 & 3 \left(-\frac{143 \int x^{3/2} \cos^2(a + b\sqrt[3]{x}) \, d\sqrt[3]{x}}{16b^2} + \frac{13x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{13/6} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} + \frac{x^{5/2}}{15} \right) \\
 & \quad \downarrow \text{3042} \\
 & 3 \left(-\frac{143 \int x^{3/2} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right)^2 \, d\sqrt[3]{x}}{16b^2} + \frac{13x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{13/6} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} + \frac{x^{5/2}}{15} \right) \\
 & \quad \downarrow \text{3792} \\
 & 3 \left(-\frac{143 \left(-\frac{63 \int x^{5/6} \cos^2(a + b\sqrt[3]{x}) \, d\sqrt[3]{x}}{16b^2} + \frac{1}{2} \int x^{3/2} \, d\sqrt[3]{x} + \frac{9x^{7/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{3/2} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} \right)}{16b^2} + \frac{13x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} \right) \\
 & \quad \downarrow \text{15} \\
 & 3 \left(-\frac{143 \left(-\frac{63 \int x^{5/6} \cos^2(a + b\sqrt[3]{x}) \, d\sqrt[3]{x}}{16b^2} + \frac{9x^{7/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{3/2} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} + \frac{x^{11/6}}{11} \right)}{16b^2} + \frac{13x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$3 \left(\frac{143 \left(-\frac{63 \int x^{5/6} \sin(a+b\sqrt[3]{x+\frac{\pi}{2}})^2 d\sqrt[3]{x}}{16b^2} + \frac{9x^{7/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{x^{3/2} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} + \frac{x^{11/6}}{11} \right)}{16b^2} + \frac{13x^{11/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} \right)$$

↓ 3792

$$3 \left(\frac{143 \left(-\frac{63 \left(-\frac{15 \int \sqrt[6]{x} \cos^2(a+b\sqrt[3]{x}) d\sqrt[3]{x}}{16b^2} + \frac{1}{2} \int x^{5/6} d\sqrt[3]{x} + \frac{5\sqrt{x} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{x^{5/6} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} \right)}{16b^2} + \frac{9x^{7/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} \right)}{16b^2}$$

↓ 15

$$3 \left(\frac{143 \left(-\frac{63 \left(-\frac{15 \int \sqrt[6]{x} \cos^2(a+b\sqrt[3]{x}) d\sqrt[3]{x}}{16b^2} + \frac{5\sqrt{x} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{x^{5/6} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} + \frac{x^{7/6}}{7} \right)}{16b^2} + \frac{9x^{7/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} \right)}{16b^2}$$

↓ 3042

$$3 \left(\frac{143 \left(-\frac{63 \left(-\frac{15 \int \sqrt[6]{x} \sin(a+b\sqrt[3]{x+\frac{\pi}{2}})^2 d\sqrt[3]{x}}{16b^2} + \frac{5\sqrt{x} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{x^{5/6} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} + \frac{x^{7/6}}{7} \right)}{16b^2} + \frac{9x^{7/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} \right)}{16b^2}$$

↓ 3793

$$\left. \begin{array}{l} 143 \\ 3 \end{array} \right\} - \frac{63 \left(\frac{15 \int \left(\frac{1}{2} \sqrt[6]{x} \cos \left(2a + 2b \sqrt[3]{x} \right) + \frac{\sqrt[6]{x}}{2} \right) d \sqrt[3]{x}}{16b^2} + \frac{5\sqrt{x} \cos^2 \left(a + b \sqrt[3]{x} \right)}{8b^2} + \frac{x^{5/6} \sin \left(a + b \sqrt[3]{x} \right) \cos \left(a + b \sqrt[3]{x} \right)}{2b} + \frac{x^{7/6}}{7} \right)}{16b^2} + \frac{9x^{7/6} \cos \left(a + b \sqrt[3]{x} \right)}{16b^2}$$

↓ 2009

$$\left(\frac{13x^{11/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \left(\frac{9x^{7/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \left(\frac{5\sqrt{x} \cos^2(a + b\sqrt[3]{x})}{8b^2} - \left(\frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - \sqrt{\pi} \cos(2a)}{8b^{3/2}} \right) \right) \right) \right)$$

input `Int[x^(3/2)*Cos[a + b*x^(1/3)]^2,x]`

output `3*(x^(5/2)/15 + (13*x^(11/6)*Cos[a + b*x^(1/3)]^2)/(8*b^2) + (x^(13/6)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(2*b) - (143*(x^(11/6)/11 + (9*x^(7/6)*Cos[a + b*x^(1/3)]^2)/(8*b^2) + (x^(3/2)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(2*b) - (63*(x^(7/6)/7 + (5*Sqrt[x]*Cos[a + b*x^(1/3)]^2)/(8*b^2) + (x^(5/6)*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(2*b) - (15*(Sqrt[x]/3 - (Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/(8*b^(3/2)) - (Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/(8*b^(3/2)) + (x^(1/6)*Sin[2*a + 2*b*x^(1/3)]/(4*b))/(16*b^2)))/(16*b^2)))/(16*b^2))`

Definitions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3792 `Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[d*m*(c + d*x)^(m - 1)*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (-Simp[b*(c + d*x)^m*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^2*((n - 1)/n) Int[(c + d*x)^m*(b*Sine[e + f*x])^(n - 2), x], x] - Simp[d^2*m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sine[e + f*x])^n, x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]`
- rule 3793 `Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 3897 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]`

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.70

method	result
derivativedivides	$\frac{x^{\frac{5}{2}}}{5} + \frac{3x^{\frac{13}{6}} \sin\left(2a+2bx^{\frac{1}{3}}\right)}{4b} - \left(\frac{x^{\frac{11}{6}} \cos\left(2a+2bx^{\frac{1}{3}}\right)}{4b} + \frac{11x^{\frac{3}{2}} \sin\left(2a+2bx^{\frac{1}{3}}\right)}{16b} - \left(\frac{x^{\frac{7}{6}} \cos\left(2a+2bx^{\frac{1}{3}}\right)}{4b} + \frac{7x^{\frac{5}{6}} \sin\left(2a+2bx^{\frac{1}{3}}\right)}{4b} \right) \right)$
default	$\frac{x^{\frac{5}{2}}}{5} + \frac{3x^{\frac{13}{6}} \sin\left(2a+2bx^{\frac{1}{3}}\right)}{4b} - \left(\frac{x^{\frac{11}{6}} \cos\left(2a+2bx^{\frac{1}{3}}\right)}{4b} + \frac{11x^{\frac{3}{2}} \sin\left(2a+2bx^{\frac{1}{3}}\right)}{16b} - \left(\frac{x^{\frac{7}{6}} \cos\left(2a+2bx^{\frac{1}{3}}\right)}{4b} + \frac{7x^{\frac{5}{6}} \sin\left(2a+2bx^{\frac{1}{3}}\right)}{4b} \right) \right)$

input `int(x^(3/2)*cos(a+b*x^(1/3))^2,x,method=_RETURNVERBOSE)`

output `1/5*x^(5/2)+3/4/b*x^(13/6)*sin(2*a+2*b*x^(1/3))-39/4/b*(-1/4/b*x^(11/6)*cos(2*a+2*b*x^(1/3))+11/4/b*(1/4/b*x^(3/2)*sin(2*a+2*b*x^(1/3))-9/4/b*(-1/4/b*x^(7/6)*cos(2*a+2*b*x^(1/3))+7/4/b*(1/4/b*x^(5/6)*sin(2*a+2*b*x^(1/3))-5/4/b*(-1/4/b*x^(1/2)*cos(2*a+2*b*x^(1/3))+3/4/b*(1/4*x^(1/6)*sin(2*a+2*b*x^(1/3)))/b-1/8/b^(3/2)*Pi^(1/2)*(cos(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))+sin(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))))))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.59

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \frac{399360 b^6 x^{11/6} - 2471040 b^4 x^{7/6} - 2027025 \pi \sqrt{\frac{b}{\pi}} \cos(2a) S\left(2 x^{1/6} \sqrt{\frac{b}{\pi}}\right) - 2027025 \pi \sqrt{\frac{b}{\pi}} C\left(2 x^{1/6} \sqrt{\frac{b}{\pi}}\right) \sin(2a)}{1}$$

input `integrate(x^(3/2)*cos(a+b*x^(1/3))^2,x, algorithm="fricas")`

output `-1/163840*(399360*b^6*x^(11/6) - 2471040*b^4*x^(7/6) - 2027025*pi*sqrt(b/pi)*cos(2*a)*fresnel_sin(2*x^(1/6)*sqrt(b/pi)) - 2027025*pi*sqrt(b/pi)*fresnel_cos(2*x^(1/6)*sqrt(b/pi))*sin(2*a) - 3120*(256*b^6*x^(11/6) - 1584*b^4*x^(7/6) + 3465*b^2*sqrt(x))*cos(b*x^(1/3) + a)^2 + 60*(36608*b^5*x^(3/2) - 144144*b^3*x^(5/6) - (4096*b^7*x^2 - 135135*b)*x^(1/6))*cos(b*x^(1/3) + a)*sin(b*x^(1/3) + a) - 8*(4096*b^8*x^2 - 675675*b^2)*sqrt(x))/b^8`

Sympy [F]

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx$$

input `integrate(x**(3/2)*cos(a+b*x**(1/3))**2,x)`

output `Integral(x**(3/2)*cos(a + b*x**(1/3))**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.52

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \frac{262144 b^9 x^{5/2} + 2027025 \cdot 4^{1/4} \sqrt{2} \sqrt{\pi} \left((i+1) \cos(2a) - (i-1) \sin(2a) \right) \operatorname{erf}\left(\sqrt{2i} b x^{1/6}\right) + (-1)^{i+1} \cos(2a) - (i-1) \sin(2a)}{b^9}$$

input `integrate(x^(3/2)*cos(a+b*x^(1/3))^2,x, algorithm="maxima")`

output `1/1310720*(262144*b^9*x^(5/2) + 2027025*4^(1/4)*sqrt(2)*sqrt(pi)*(((I + 1)*cos(2*a) - (I - 1)*sin(2*a))*erf(sqrt(2*I*b)*x^(1/6)) + (-1)^{i+1}*cos(2*a) + (I + 1)*sin(2*a))*erf(sqrt(-2*I*b)*x^(1/6)))*b^(3/2) + 12480*(256*b^7*x^(11/6) - 1584*b^5*x^(7/6) + 3465*b^3*sqrt(x))*cos(2*b*x^(1/3) + 2*a) + 240*(4096*b^8*x^(13/6) - 36608*b^6*x^(3/2) + 144144*b^4*x^(5/6) - 135135*b^2*x^(1/6))*sin(2*b*x^(1/3) + 2*a))/b^9`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.72

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \frac{1}{5} x^{5/2}$$

$$\frac{3 \left(4096i b^6 x^{13/6} - 13312 b^5 x^{11/6} - 36608i b^4 x^{3/2} + 82368 b^3 x^{7/6} + 144144i b^2 x^{5/6} - 180180 b \sqrt{x} - 135135i x^{1/6} \right)}{32768 b^7}$$

$$\frac{3 \left(-4096i b^6 x^{13/6} - 13312 b^5 x^{11/6} + 36608i b^4 x^{3/2} + 82368 b^3 x^{7/6} - 144144i b^2 x^{5/6} - 180180 b \sqrt{x} + 135135i x^{1/6} \right)}{32768 b^7}$$

$$+ \frac{405405i \sqrt{\pi} \operatorname{erf} \left(-\sqrt{b} x^{1/6} \left(-\frac{ib}{|b|} + 1 \right) \right) e^{(2ia)}}{65536 b^{15/2} \left(-\frac{ib}{|b|} + 1 \right)}$$

$$- \frac{405405i \sqrt{\pi} \operatorname{erf} \left(-\sqrt{b} x^{1/6} \left(\frac{ib}{|b|} + 1 \right) \right) e^{(-2ia)}}{65536 b^{15/2} \left(\frac{ib}{|b|} + 1 \right)}$$

input `integrate(x^(3/2)*cos(a+b*x^(1/3))^2,x, algorithm="giac")`

output

```
1/5*x^(5/2) - 3/32768*(4096*I*b^6*x^(13/6) - 13312*b^5*x^(11/6) - 36608*I*
b^4*x^(3/2) + 82368*b^3*x^(7/6) + 144144*I*b^2*x^(5/6) - 180180*b*sqrt(x)
- 135135*I*x^(1/6))*e^(2*I*b*x^(1/3) + 2*I*a)/b^7 - 3/32768*(-4096*I*b^6*x
^(13/6) - 13312*b^5*x^(11/6) + 36608*I*b^4*x^(3/2) + 82368*b^3*x^(7/6) - 1
44144*I*b^2*x^(5/6) - 180180*b*sqrt(x) + 135135*I*x^(1/6))*e^(-2*I*b*x^(1/
3) - 2*I*a)/b^7 + 405405/65536*I*sqrt(pi)*erf(-sqrt(b)*x^(1/6)*(-I*b/abs(b
) + 1))*e^(2*I*a)/(b^(15/2)*(-I*b/abs(b) + 1)) - 405405/65536*I*sqrt(pi)*e
rf(-sqrt(b)*x^(1/6)*(I*b/abs(b) + 1))*e^(-2*I*a)/(b^(15/2)*(I*b/abs(b) + 1
))
```


Mupad [F(-1)]

Timed out.

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \int x^{3/2} \cos(a + b x^{1/3})^2 dx$$

input `int(x^(3/2)*cos(a + b*x^(1/3))^2,x)`output `int(x^(3/2)*cos(a + b*x^(1/3))^2, x)`**Reduce [F]**

$$\int x^{3/2} \cos^2(a + b\sqrt[3]{x}) dx = \text{too large to display}$$

input `int(x^(3/2)*cos(a+b*x^(1/3))^2,x)`

output

```
(360360*x**(5/6)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*tan((x**(1/3)*b +
a)/2)**4*b**2 + 720720*x**(5/6)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*t
an((x**(1/3)*b + a)/2)**2*b**2 + 360360*x**(5/6)*cos(x**(1/3)*b + a)*sin(x
**(1/3)*b + a)*b**2 - 68640*sqrt(x)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a
)*tan((x**(1/3)*b + a)/2)**4*b**4*x - 137280*sqrt(x)*cos(x**(1/3)*b + a)*s
in(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**2*b**4*x - 68640*sqrt(x)*cos(x
**(1/3)*b + a)*sin(x**(1/3)*b + a)*b**4*x + 900900*sqrt(x)*cos(x**(1/3)*b
+ a)*tan((x**(1/3)*b + a)/2)**4*b + 1801800*sqrt(x)*cos(x**(1/3)*b + a)*ta
n((x**(1/3)*b + a)/2)**2*b + 900900*sqrt(x)*cos(x**(1/3)*b + a)*b + 7680*x
**(1/6)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**4
*b**6*x**2 + 15360*x**(1/6)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*tan((x
**(1/3)*b + a)/2)**2*b**6*x**2 + 7680*x**(1/6)*cos(x**(1/3)*b + a)*sin(x**
(1/3)*b + a)*b**6*x**2 - 24960*x**(5/6)*sin(x**(1/3)*b + a)**2*tan((x**(1/
3)*b + a)/2)**4*b**5*x - 49920*x**(5/6)*sin(x**(1/3)*b + a)**2*tan((x**(1/
3)*b + a)/2)**2*b**5*x - 24960*x**(5/6)*sin(x**(1/3)*b + a)**2*b**5*x + 36
0360*x**(5/6)*sin(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**4*b**2 + 720720
*x**(5/6)*sin(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**2*b**2 + 360360*x**
(5/6)*sin(x**(1/3)*b + a)*b**2 + 12480*x**(5/6)*tan((x**(1/3)*b + a)/2)**4
*b**5*x - 540540*x**(5/6)*tan((x**(1/3)*b + a)/2)**3*b**2 + 24960*x**(5/6)
*tan((x**(1/3)*b + a)/2)**2*b**5*x - 900900*x**(5/6)*tan((x**(1/3)*b + ...
```

3.56 $\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$

Optimal result	442
Mathematica [A] (verified)	443
Rubi [A] (verified)	443
Maple [A] (verified)	447
Fricas [A] (verification not implemented)	447
Sympy [F]	448
Maxima [C] (verification not implemented)	448
Giac [C] (verification not implemented)	449
Mupad [F(-1)]	449
Reduce [F]	450

Optimal result

Integrand size = 18, antiderivative size = 218

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx = \frac{315\sqrt[6]{x}}{256b^4} - \frac{21x^{5/6}}{16b^2} + \frac{x^{3/2}}{3} - \frac{315\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{128b^4}$$

$$+ \frac{21x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2}$$

$$+ \frac{315\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{512b^{9/2}}$$

$$- \frac{315\sqrt{\pi} \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) \sin(2a)}{512b^{9/2}}$$

$$- \frac{105\sqrt{x} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{32b^3}$$

$$+ \frac{3x^{7/6} \cos(a + b\sqrt[3]{x}) \sin(a + b\sqrt[3]{x})}{2b}$$

output

```
315/256*x^(1/6)/b^4-21/16*x^(5/6)/b^2+1/3*x^(3/2)-315/128*x^(1/6)*cos(a+b*x^(1/3))^2/b^4+21/8*x^(5/6)*cos(a+b*x^(1/3))^2/b^2+315/512*Pi^(1/2)*cos(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))/b^(9/2)-315/512*Pi^(1/2)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))*sin(2*a)/b^(9/2)-105/32*x^(1/2)*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/b^3+3/2*x^(7/6)*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/b
```

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.68

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$$

$$= \frac{945\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 945\sqrt{\pi} \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) \sin(2a) + 2\sqrt{b}\sqrt[6]{x}(63(-15 + 16b^2x^{2/3}) \cos(2(a + b\sqrt[3]{x})) + 4b\sqrt[3]{x}(64b^3x + 9(-35 + 16b^2x^{2/3}) \sin(2(a + b\sqrt[3]{x}))))}{1536b^{9/2}}$$

input `Integrate[Sqrt[x]*Cos[a + b*x^(1/3)]^2,x]`

output `(945*Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]] - 945*Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 2*Sqrt[b]*x^(1/6)*(63*(-15 + 16*b^2*x^(2/3))*Cos[2*(a + b*x^(1/3))] + 4*b*x^(1/3)*(64*b^3*x + 9*(-35 + 16*b^2*x^(2/3))*Sin[2*(a + b*x^(1/3))])))/(1536*b^(9/2))`

Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3897, 3042, 3792, 15, 3042, 3792, 15, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$$

$$\downarrow 3897$$

$$3 \int x^{7/6} \cos^2(a + b\sqrt[3]{x}) d\sqrt[3]{x}$$

$$\downarrow 3042$$

$$3 \int x^{7/6} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right)^2 d\sqrt[3]{x}$$

$$\downarrow 3792$$

$$3 \left(-\frac{35 \int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{16b^2} + \frac{1}{2} \int x^{7/6} d\sqrt[3]{x} + \frac{7x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{7/6} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} \right)$$

↓ 15

$$3 \left(-\frac{35 \int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) d\sqrt[3]{x}}{16b^2} + \frac{7x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{7/6} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} + \frac{x^{3/2}}{9} \right)$$

↓ 3042

$$3 \left(-\frac{35 \int \sqrt{x} \sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2 d\sqrt[3]{x}}{16b^2} + \frac{7x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{x^{7/6} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} + \frac{x^{3/2}}{9} \right)$$

↓ 3792

$$3 \left(\frac{35 \left(-\frac{3 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{16b^2} + \frac{\int \sqrt{x} d\sqrt[3]{x}}{2} + \frac{3\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{\sqrt{x} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} \right)}{16b^2} + \frac{7x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} \right)$$

↓ 15

$$3 \left(\frac{35 \left(-\frac{3 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x}}{16b^2} + \frac{3\sqrt[6]{x} \cos^2(a + b\sqrt[3]{x})}{8b^2} + \frac{\sqrt{x} \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{2b} + \frac{x^{5/6}}{5} \right)}{16b^2} + \frac{7x^{5/6} \cos^2(a + b\sqrt[3]{x})}{8b^2} \right)$$

↓ 3042

$$3 \left(\frac{35 \left(-\frac{3 \int \frac{\sin(a+b\sqrt[3]{x}+\frac{\pi}{2})^2}{\sqrt[6]{x}} d\sqrt[3]{x}}{16b^2} + \frac{3\sqrt[6]{x} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{\sqrt{x} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} + \frac{x^{5/6}}{5} \right)}{16b^2} + \frac{7x^{5/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} \right)$$

↓ 3793

$$3 \left(\frac{35 \left(-\frac{3 \int \left(\frac{\cos(2a+2b\sqrt[3]{x})}{2\sqrt[6]{x}} + \frac{1}{2\sqrt[6]{x}} \right) d\sqrt[3]{x}}{16b^2} + \frac{3\sqrt[6]{x} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{\sqrt{x} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} + \frac{x^{5/6}}{5} \right)}{16b^2} + \frac{7x^{5/6} \cos^2(a+b\sqrt[3]{x})}{8b^2} \right)$$

↓ 2009

$$3 \left(\frac{35 \left(-\frac{3 \left(\frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt[6]{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{2\sqrt[6]{b}} - \frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelS}\left(\frac{2\sqrt[6]{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{2\sqrt[6]{b}} + \sqrt[6]{x} \right)}{16b^2} + \frac{3\sqrt[6]{x} \cos^2(a+b\sqrt[3]{x})}{8b^2} + \frac{\sqrt{x} \sin(a+b\sqrt[3]{x}) \cos(a+b\sqrt[3]{x})}{2b} \right)}{16b^2} \right)$$

input `Int[Sqrt[x]*Cos[a + b*x^(1/3)]^2,x]`

output

```
3*(x^(3/2)/9 + (7*x^(5/6)*Cos[a + b*x^(1/3)]^2)/(8*b^2) + (x^(7/6)*Cos[a +
b*x^(1/3)]*Sin[a + b*x^(1/3)])/(2*b) - (35*(x^(5/6)/5 + (3*x^(1/6)*Cos[a
+ b*x^(1/3)]^2)/(8*b^2) - (3*(x^(1/6) + (Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqr
t[b]*x^(1/6))/Sqrt[Pi]]))/(2*Sqrt[b]) - (Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x^(1/
6))/Sqrt[Pi]]*Sin[2*a])/(2*Sqrt[b])))/(16*b^2) + (Sqrt[x]*Cos[a + b*x^(1/3
)]*Sin[a + b*x^(1/3)])/(2*b))/(16*b^2))
```

Defintions of rubi rules used

rule 15

```
Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3792

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
:= Simp[d*m*(c + d*x)^(m - 1)*((b*Sine[e + f*x])^n/(f^2*n^2)), x] + (-Simp[
b*(c + d*x)^m*cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(f*n)), x] + Simp[b^
2*((n - 1)/n) Int[(c + d*x)^m*(b*Sine[e + f*x])^(n - 2), x], x] - Simp[d^2
*m*((m - 1)/(f^2*n^2)) Int[(c + d*x)^(m - 2)*(b*Sine[e + f*x])^n, x], x])
/; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 1]
```

rule 3793

```
Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

rule 3897

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol]
:= Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a +
b*cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]
```

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.67

method	result
derivativedivides	$\frac{x^{\frac{3}{2}}}{3} + \frac{3x^{\frac{7}{6}} \sin(2a+2bx^{\frac{1}{3}})}{4b} - \frac{21 \left(\frac{x^{\frac{5}{6}} \cos(2a+2bx^{\frac{1}{3}})}{4b} + \frac{5\sqrt{x} \sin(2a+2bx^{\frac{1}{3}})}{16b} - \frac{15 \left(\frac{x^{\frac{1}{6}} \cos(2a+2bx^{\frac{1}{3}})}{4b} + \frac{\sqrt{\pi} \cos(2a)}{b} \right)}{b} \right)}{4b}$
default	$\frac{x^{\frac{3}{2}}}{3} + \frac{3x^{\frac{7}{6}} \sin(2a+2bx^{\frac{1}{3}})}{4b} - \frac{21 \left(\frac{x^{\frac{5}{6}} \cos(2a+2bx^{\frac{1}{3}})}{4b} + \frac{5\sqrt{x} \sin(2a+2bx^{\frac{1}{3}})}{16b} - \frac{15 \left(\frac{x^{\frac{1}{6}} \cos(2a+2bx^{\frac{1}{3}})}{4b} + \frac{\sqrt{\pi} \cos(2a)}{b} \right)}{b} \right)}{4b}$

input `int(x^(1/2)*cos(a+b*x^(1/3))^2,x,method=_RETURNVERBOSE)`

output `1/3*x^(3/2)+3/4/b*x^(7/6)*sin(2*a+2*b*x^(1/3))-21/4/b*(-1/4/b*x^(5/6)*cos(2*a+2*b*x^(1/3))+5/4/b*(1/4/b*x^(1/2)*sin(2*a+2*b*x^(1/3))-3/4/b*(-1/4/b*x^(1/6)*cos(2*a+2*b*x^(1/3))+1/8/b^(3/2)*Pi^(1/2)*(cos(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))-sin(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.66

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx = \frac{512 b^5 x^{\frac{3}{2}} - 2016 b^3 x^{\frac{5}{6}} + 945 \pi \sqrt{\frac{b}{\pi}} \cos(2a) C\left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) - 945 \pi \sqrt{\frac{b}{\pi}} S\left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(2a) + 252 \left(16 b^3 x^{\frac{3}{2}} - 1536 b^5\right)}{1536 b^5}$$

input `integrate(x^(1/2)*cos(a+b*x^(1/3))^2,x, algorithm="fricas")`

output

```
1/1536*(512*b^5*x^(3/2) - 2016*b^3*x^(5/6) + 945*pi*sqrt(b/pi)*cos(2*a)*fresnel_cos(2*x^(1/6)*sqrt(b/pi)) - 945*pi*sqrt(b/pi)*fresnel_sin(2*x^(1/6)*sqrt(b/pi))*sin(2*a) + 252*(16*b^3*x^(5/6) - 15*b*x^(1/6))*cos(b*x^(1/3) + a)^2 + 144*(16*b^4*x^(7/6) - 35*b^2*sqrt(x))*cos(b*x^(1/3) + a)*sin(b*x^(1/3) + a) + 1890*b*x^(1/6))/b^5
```

Sympy [F]

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx = \int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$$

input

```
integrate(x**(1/2)*cos(a+b*x**(1/3))**2,x)
```

output

```
Integral(sqrt(x)*cos(a + b*x**(1/3))**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.63

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$$

$$= \frac{4096 b^6 x^{\frac{3}{2}} + 945 \cdot 4^{\frac{1}{4}} \sqrt{2} \sqrt{\pi} \left((-i - 1) \cos(2a) - (i + 1) \sin(2a) \right) \operatorname{erf} \left(\sqrt{2i} b x^{\frac{1}{6}} \right) + ((i + 1) \cos(2a) +$$

input

```
integrate(x^(1/2)*cos(a+b*x^(1/3))^2,x, algorithm="maxima")
```

output

```
1/12288*(4096*b^6*x^(3/2) + 945*4^(1/4)*sqrt(2)*sqrt(pi)*((-I - 1)*cos(2*a) - (I + 1)*sin(2*a))*erf(sqrt(2*I*b)*x^(1/6)) + ((I + 1)*cos(2*a) + (I - 1)*sin(2*a))*erf(sqrt(-2*I*b)*x^(1/6)))*b^(3/2) + 1008*(16*b^4*x^(5/6) - 15*b^2*x^(1/6))*cos(2*b*x^(1/3) + 2*a) + 576*(16*b^5*x^(7/6) - 35*b^3*sqrt(x))*sin(2*b*x^(1/3) + 2*a))/b^6
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.81

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$$

$$= \frac{1}{3} x^{\frac{3}{2}} - \frac{3 \left(64i b^3 x^{\frac{7}{6}} - 112 b^2 x^{\frac{5}{6}} - 140i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{(2i b x^{\frac{1}{3}} + 2i a)}}{512 b^4}$$

$$- \frac{3 \left(-64i b^3 x^{\frac{7}{6}} - 112 b^2 x^{\frac{5}{6}} + 140i b \sqrt{x} + 105 x^{\frac{1}{6}} \right) e^{(-2i b x^{\frac{1}{3}} - 2i a)}}{512 b^4}$$

$$- \frac{315 \sqrt{\pi} \operatorname{erf} \left(-\sqrt{b} x^{\frac{1}{6}} \left(-\frac{i b}{|b|} + 1 \right) \right) e^{(2i a)}}{1024 b^{\frac{9}{2}} \left(-\frac{i b}{|b|} + 1 \right)} - \frac{315 \sqrt{\pi} \operatorname{erf} \left(-\sqrt{b} x^{\frac{1}{6}} \left(\frac{i b}{|b|} + 1 \right) \right) e^{(-2i a)}}{1024 b^{\frac{9}{2}} \left(\frac{i b}{|b|} + 1 \right)}$$

input `integrate(x^(1/2)*cos(a+b*x^(1/3))^2,x, algorithm="giac")`

output `1/3*x^(3/2) - 3/512*(64*I*b^3*x^(7/6) - 112*b^2*x^(5/6) - 140*I*b*sqrt(x) + 105*x^(1/6))*e^(2*I*b*x^(1/3) + 2*I*a)/b^4 - 3/512*(-64*I*b^3*x^(7/6) - 112*b^2*x^(5/6) + 140*I*b*sqrt(x) + 105*x^(1/6))*e^(-2*I*b*x^(1/3) - 2*I*a)/b^4 - 315/1024*sqrt(pi)*erf(-sqrt(b)*x^(1/6)*(-I*b/abs(b) + 1))*e^(2*I*a)/(b^(9/2)*(-I*b/abs(b) + 1)) - 315/1024*sqrt(pi)*erf(-sqrt(b)*x^(1/6)*(I*b/abs(b) + 1))*e^(-2*I*a)/(b^(9/2)*(I*b/abs(b) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx = \int \sqrt{x} \cos(a + b x^{1/3})^2 dx$$

input `int(x^(1/2)*cos(a + b*x^(1/3))^2,x)`

output `int(x^(1/2)*cos(a + b*x^(1/3))^2, x)`

Reduce [F]

$$\int \sqrt{x} \cos^2(a + b\sqrt[3]{x}) dx$$

$$-420\sqrt{x} \cos\left(x^{\frac{1}{3}}b + a\right) \sin\left(x^{\frac{1}{3}}b + a\right) b + 144x^{\frac{7}{6}} \cos\left(x^{\frac{1}{3}}b + a\right) \sin\left(x^{\frac{1}{3}}b + a\right) b^3 - 630x^{\frac{1}{6}} \cos\left(x^{\frac{1}{3}}b + a\right)$$

=

input `int(x^(1/2)*cos(a+b*x^(1/3))^2,x)`

output

```
( - 420*sqrt(x)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*b + 144*x**(1/6)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*b**3*x - 630*x**(1/6)*cos(x**(1/3)*b + a) - 252*x**(5/6)*sin(x**(1/3)*b + a)**2*b**2 - 420*sqrt(x)*sin(x**(1/3)*b + a)*b + 32*sqrt(x)*b**4*x + 315*x**(1/6)*sin(x**(1/3)*b + a)**2 - 630*x**(1/6) + 280*int(x**(1/6)/(x**(1/3)*tan((x**(1/3)*b + a)/2)**4 + 2*x**(1/3)*tan((x**(1/3)*b + a)/2)**2 + x**(1/3)),x)*b**2 + 210*int(1/(x**(5/6)*tan((x**(1/3)*b + a)/2)**4 + 2*x**(5/6)*tan((x**(1/3)*b + a)/2)**2 + x**(5/6)),x))/(96*b**4)
```

3.57 $\int \frac{\cos^2\left(a+b\sqrt[3]{x}\right)}{\sqrt{x}} dx$

Optimal result	451
Mathematica [A] (verified)	451
Rubi [A] (verified)	452
Maple [A] (verified)	453
Fricas [A] (verification not implemented)	454
Sympy [F]	454
Maxima [C] (verification not implemented)	455
Giac [C] (verification not implemented)	455
Mupad [F(-1)]	456
Reduce [F]	456

Optimal result

Integrand size = 18, antiderivative size = 103

$$\int \frac{\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}} dx = \sqrt{x} - \frac{3\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} - \frac{3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) \sin(2a)}{8b^{3/2}} + \frac{3\sqrt[6]{x} \sin(2a+2b\sqrt[3]{x})}{4b}$$

output

```
x^(1/2)-3/8*Pi^(1/2)*cos(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))/b^(3/2)
-3/8*Pi^(1/2)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))*sin(2*a)/b^(3/2)+3/4*x^(1/6)*sin(2*a+2*b*x^(1/3))/b
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}} dx = \frac{-3\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) \sin(2a) + 2\sqrt{b}\sqrt[6]{x}(4b\sqrt[3]{x} + 3 \sin(2(a+b\sqrt[3]{x})))}{8b^{3/2}}$$

input `Integrate[Cos[a + b*x^(1/3)]^2/Sqrt[x], x]`

output `(-3*Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]] - 3*Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 2*Sqrt[b]*x^(1/6)*(4*b*x^(1/3) + 3*Sin[2*(a + b*x^(1/3))]))/(8*b^(3/2))`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3897, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx \\
 & \quad \downarrow \text{3897} \\
 & 3 \int \sqrt[6]{x} \cos^2(a + b\sqrt[3]{x}) d\sqrt[3]{x} \\
 & \quad \downarrow \text{3042} \\
 & 3 \int \sqrt[6]{x} \sin\left(a + b\sqrt[3]{x} + \frac{\pi}{2}\right)^2 d\sqrt[3]{x} \\
 & \quad \downarrow \text{3793} \\
 & 3 \int \left(\frac{1}{2}\sqrt[6]{x} \cos(2a + 2b\sqrt[3]{x}) + \frac{\sqrt[6]{x}}{2}\right) d\sqrt[3]{x} \\
 & \quad \downarrow \text{2009} \\
 & 3 \left(-\frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} - \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{8b^{3/2}} + \frac{\sqrt[6]{x} \sin(2a + 2b\sqrt[3]{x})}{4b} + \frac{\sqrt{x}}{3} \right)
 \end{aligned}$$

input `Int[Cos[a + b*x^(1/3)]^2/Sqrt[x], x]`

output

$$3*(\text{Sqrt}[x]/3 - (\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a]*\text{FresnelS}[(2*\text{Sqrt}[b]*x^{1/6})/\text{Sqrt}[\text{Pi}]])/(8*b^{3/2})) - (\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(2*\text{Sqrt}[b]*x^{1/6})/\text{Sqrt}[\text{Pi}]]*\text{Sin}[2*a])/(8*b^{3/2}) + (x^{1/6}*\text{Sin}[2*a + 2*b*x^{1/3}])/(4*b))$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; } \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3793

$$\text{Int}[(c_. + (d_.)*(x_.)^{m_.})*\text{sin}[(e_. + (f_.)*(x_.)^{n_.}], x_Symbol] \text{ :> } \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \text{Sin}[e + f*x]^n, x], x] \text{ /; } \text{FreeQ}\{c, d, e, f, m\}, x \text{ \&\& } \text{IGtQ}[n, 1] \text{ \&\& } (!\text{RationalQ}[m] \text{ || } (\text{GeQ}[m, -1] \text{ \&\& } \text{LtQ}[m, 1]))$$

rule 3897

$$\text{Int}[(a_. + \text{Cos}[c_. + (d_.)*(x_.)^{n_.}]*b_.)^{p_.}*(x_.)^{m_.}, x_Symbol] \text{ :> } \text{Module}\{k = \text{Denominator}[n]\}, \text{Simp}[k \text{ Subst}[\text{Int}[x^{k*(m+1)-1}*(a + b*\text{Cos}[c + d*x^{k*n}])^p, x], x, x^{1/k}], x] \text{ /; } \text{FreeQ}\{a, b, c, d, m\}, x \text{ \&\& } \text{IntegerQ}[p] \text{ \&\& } \text{FractionQ}[n]$$
Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.65

method	result	size
derivativedivides	$\sqrt{x} + \frac{3x^{\frac{1}{6}} \sin(2a+2bx^{\frac{1}{3}})}{4b} - \frac{3\sqrt{\pi} \left(\cos(2a) \text{FresnelS}\left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) + \sin(2a) \text{FresnelC}\left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) \right)}{8b^{\frac{3}{2}}}$	67
default	$\sqrt{x} + \frac{3x^{\frac{1}{6}} \sin(2a+2bx^{\frac{1}{3}})}{4b} - \frac{3\sqrt{\pi} \left(\cos(2a) \text{FresnelS}\left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) + \sin(2a) \text{FresnelC}\left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}}\right) \right)}{8b^{\frac{3}{2}}}$	67

input

$$\text{int}(\cos(a+b*x^{1/3})^2/x^{1/2}, x, \text{method}=_RETURNVERBOSE)$$

output

```
x^(1/2)+3/4*x^(1/6)*sin(2*a+2*b*x^(1/3))/b-3/8/b^(3/2)*Pi^(1/2)*(cos(2*a)*
FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))+sin(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/P
i^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.87

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \frac{3\pi\sqrt{\frac{b}{\pi}}\cos(2a)S\left(2x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right) + 3\pi\sqrt{\frac{b}{\pi}}C\left(2x^{\frac{1}{6}}\sqrt{\frac{b}{\pi}}\right)\sin(2a) - 12bx^{\frac{1}{6}}\cos\left(bx^{\frac{1}{3}} + a\right)\sin\left(bx^{\frac{1}{3}} + a\right) - 8b^2}{8b^2}$$

input

```
integrate(cos(a+b*x^(1/3))^2/x^(1/2),x, algorithm="fricas")
```

output

```
-1/8*(3*pi*sqrt(b/pi)*cos(2*a)*fresnel_sin(2*x^(1/6)*sqrt(b/pi)) + 3*pi*sq
rt(b/pi)*fresnel_cos(2*x^(1/6)*sqrt(b/pi))*sin(2*a) - 12*b*x^(1/6)*cos(b*x
^(1/3) + a)*sin(b*x^(1/3) + a) - 8*b^2*sqrt(x))/b^2
```

Sympy [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx$$

input

```
integrate(cos(a+b*x**(1/3))**2/x**(1/2),x)
```

output

```
Integral(cos(a + b*x**(1/3))**2/sqrt(x), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.93

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \frac{3 \cdot 4^{\frac{1}{4}} \sqrt{2} \sqrt{\pi} \left(((i+1) \cos(2a) - (i-1) \sin(2a)) \operatorname{erf}\left(\sqrt{2i} b x^{\frac{1}{6}}\right) + (-(i-1) \cos(2a) + (i+1) \sin(2a)) \operatorname{erf}\left(\sqrt{2i} b x^{\frac{1}{6}}\right) \right)}{64 b^3}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(1/2),x, algorithm="maxima")`

output `-1/64*(3*4^(1/4)*sqrt(2)*sqrt(pi)*(((I + 1)*cos(2*a) - (I - 1)*sin(2*a))*erf(sqrt(2*I*b)*x^(1/6)) + (-(I - 1)*cos(2*a) + (I + 1)*sin(2*a))*erf(sqrt(-2*I*b)*x^(1/6)))*b^(3/2) - 64*b^3*sqrt(x) - 48*b^2*x^(1/6)*sin(2*b*x^(1/3) + 2*a))/b^3`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.20

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \sqrt{x} - \frac{3i x^{\frac{1}{6}} e^{(2i b x^{\frac{1}{3}} + 2i a)}}{8b} + \frac{3i x^{\frac{1}{6}} e^{(-2i b x^{\frac{1}{3}} - 2i a)}}{8b} - \frac{3i \sqrt{\pi} \operatorname{erf}\left(-\sqrt{b} x^{\frac{1}{6}} \left(-\frac{ib}{|b|} + 1\right)\right) e^{(2i a)}}{16 b^{\frac{3}{2}} \left(-\frac{ib}{|b|} + 1\right)} + \frac{3i \sqrt{\pi} \operatorname{erf}\left(-\sqrt{b} x^{\frac{1}{6}} \left(\frac{ib}{|b|} + 1\right)\right) e^{(-2i a)}}{16 b^{\frac{3}{2}} \left(\frac{ib}{|b|} + 1\right)}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(1/2),x, algorithm="giac")`

output

```
sqrt(x) - 3/8*I*x^(1/6)*e^(2*I*b*x^(1/3) + 2*I*a)/b + 3/8*I*x^(1/6)*e^(-2*I*b*x^(1/3) - 2*I*a)/b - 3/16*I*sqrt(pi)*erf(-sqrt(b)*x^(1/6)*(-I*b/abs(b) + 1))*e^(2*I*a)/(b^(3/2)*(-I*b/abs(b) + 1)) + 3/16*I*sqrt(pi)*erf(-sqrt(b)*x^(1/6)*(I*b/abs(b) + 1))*e^(-2*I*a)/(b^(3/2)*(I*b/abs(b) + 1))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx = \int \frac{\cos(a + bx^{1/3})^2}{\sqrt{x}} dx$$

input

```
int(cos(a + b*x^(1/3))^2/x^(1/2), x)
```

output

```
int(cos(a + b*x^(1/3))^2/x^(1/2), x)
```

Reduce [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} dx$$

$$= \frac{2x^{\frac{1}{6}} \cos\left(x^{\frac{1}{3}}b + a\right) \sin\left(x^{\frac{1}{3}}b + a\right) + 2\sqrt{x}b + 2x^{\frac{1}{6}} \sin\left(x^{\frac{1}{3}}b + a\right) - \frac{4 \left(\int \frac{x^{\frac{1}{6}}}{x^{\frac{2}{3}} \tan\left(\frac{x^{\frac{1}{3}}b + a}{2}\right)^4 + 2x^{\frac{2}{3}} \tan\left(\frac{x^{\frac{1}{3}}b + a}{2}\right)^2 + x^{\frac{2}{3}} dx \right)}{3}}{b}$$

input

```
int(cos(a+b*x^(1/3))^2/x^(1/2), x)
```

output

```
(2*(3*x**(1/6)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a) + 3*sqrt(x)*b + 3*x**(1/6)*sin(x**(1/3)*b + a) - 2*int(x**(1/6)/(x**(2/3)*tan((x**(1/3)*b + a)/2)**4 + 2*x**(2/3)*tan((x**(1/3)*b + a)/2)**2 + x**(2/3)), x)*b - 2*int(tan((x**(1/3)*b + a)/2)/(x**(5/6)*tan((x**(1/3)*b + a)/2)**4 + 2*x**(5/6)*tan((x**(1/3)*b + a)/2)**2 + x**(5/6)), x))/ (3*b)
```

3.58 $\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{3/2}} dx$

Optimal result	457
Mathematica [A] (verified)	457
Rubi [A] (verified)	458
Maple [A] (verified)	460
Fricas [A] (verification not implemented)	461
Sympy [F]	461
Maxima [C] (verification not implemented)	461
Giac [F]	462
Mupad [F(-1)]	462
Reduce [F]	463

Optimal result

Integrand size = 18, antiderivative size = 116

$$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{3/2}} dx = -\frac{2\cos^2(a+b\sqrt[3]{x})}{\sqrt{x}} - 8b^{3/2}\sqrt{\pi}\cos(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) + 8b^{3/2}\sqrt{\pi}\operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)\sin(2a) + \frac{8b\cos(a+b\sqrt[3]{x})\sin(a+b\sqrt[3]{x})}{\sqrt[6]{x}}$$

output

```
-2*cos(a+b*x^(1/3))^2/x^(1/2)-8*b^(3/2)*Pi^(1/2)*cos(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))+8*b^(3/2)*Pi^(1/2)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))*sin(2*a)+8*b*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/x^(1/6)
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{3/2}} dx = -8b^{3/2}\sqrt{\pi}\cos(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - 1 - \cos(2(a+b\sqrt[3]{x})) + 8b^{3/2}\sqrt{\pi}\sqrt{x}\operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)\sin(2a) + 4b\sqrt[3]{x}\sin(2(a+b\sqrt[3]{x}))$$

\sqrt{x}

input `Integrate[Cos[a + b*x^(1/3)]^2/x^(3/2), x]`

output `-8*b^(3/2)*Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]] + (-1 - Cos[2*(a + b*x^(1/3))]) + 8*b^(3/2)*Sqrt[Pi]*Sqrt[x]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 4*b*x^(1/3)*Sin[2*(a + b*x^(1/3))]/Sqrt[x]`

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.30, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3897, 3042, 3795, 15, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx \\
 & \quad \downarrow \text{3897} \\
 & 3 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} \\
 & \quad \downarrow \text{3042} \\
 & 3 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{5/6}} d\sqrt[3]{x} \\
 & \quad \downarrow \text{3795} \\
 & 3 \left(-\frac{16}{3} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} + \frac{8}{3} b^2 \int \frac{1}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt[6]{x}} \right) \\
 & \quad \downarrow \text{15} \\
 & 3 \left(-\frac{16}{3} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt[6]{x}} + \frac{16}{3} b^2 \sqrt[6]{x} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$3 \left(-\frac{16}{3} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt[6]{x}} + \frac{16}{3} b^2 \sqrt[6]{x} \right)$$

↓ 3793

$$3 \left(-\frac{16}{3} b^2 \int \left(\frac{\cos(2a + 2b\sqrt[3]{x})}{2\sqrt[6]{x}} + \frac{1}{2\sqrt[6]{x}} \right) d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt[6]{x}} + \frac{16}{3} b^2 \sqrt[6]{x} \right)$$

↓ 2009

$$3 \left(-\frac{16}{3} b^2 \left(\frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{2\sqrt{b}} - \frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{2\sqrt{b}} + \sqrt[6]{x} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt[6]{x}} + \frac{16}{3} b^2 \sqrt[6]{x} \right)$$

input `Int[Cos[a + b*x^(1/3)]^2/x^(3/2),x]`

output `3*((16*b^2*x^(1/6))/3 - (2*Cos[a + b*x^(1/3)]^2)/(3*Sqrt[x]) - (16*b^2*(x^(1/6) + (Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/(2*Sqrt[b]) - (Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/(2*Sqrt[b])))/3 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(3*x^(1/6)))`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793

```
Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)^(n_)], x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

rule 3795

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)^(n_)])^(n_), x_Symbol] := Simp[(c + d*x)^(m + 1)*((b*Sine[e + f*x])^n/(d*(m + 1))), x] + (-Simp[b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(d^2*(m + 1)*(m + 2))), x] + Simp[b^2*f^2*n*((n - 1)/(d^2*(m + 1)*(m + 2))) Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^(n - 2), x], x] - Simp[f^2*(n^2/(d^2*(m + 1)*(m + 2))) Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^n, x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]
```

rule 3897

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)])^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]
```

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.75

method	result
derivativedivides	$-\frac{1}{\sqrt{x}} - \frac{\cos(2a+2bx^{\frac{1}{3}})}{\sqrt{x}} - 4b \left(-\frac{\sin(2a+2bx^{\frac{1}{3}})}{x^{\frac{1}{6}}} + 2\sqrt{b}\sqrt{\pi} \left(\cos(2a) \operatorname{FresnelC} \left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}} \right) - \sin(2a) \operatorname{FresnelS} \left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}} \right) \right) \right)$
default	$-\frac{1}{\sqrt{x}} - \frac{\cos(2a+2bx^{\frac{1}{3}})}{\sqrt{x}} - 4b \left(-\frac{\sin(2a+2bx^{\frac{1}{3}})}{x^{\frac{1}{6}}} + 2\sqrt{b}\sqrt{\pi} \left(\cos(2a) \operatorname{FresnelC} \left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}} \right) - \sin(2a) \operatorname{FresnelS} \left(\frac{2\sqrt{b}x^{\frac{1}{6}}}{\sqrt{\pi}} \right) \right) \right)$

input

```
int(cos(a+b*x^(1/3))^2/x^(3/2),x,method=_RETURNVERBOSE)
```

output

```
-1/x^(1/2)-1/x^(1/2)*cos(2*a+2*b*x^(1/3))-4*b*(-1/x^(1/6)*sin(2*a+2*b*x^(1/3))+2*b^(1/2)*Pi^(1/2)*(cos(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))-sin(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.86

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx = \frac{2 \left(4 \pi b x \sqrt{\frac{b}{\pi}} \cos(2a) C \left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) - 4 \pi b x \sqrt{\frac{b}{\pi}} S \left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(2a) - 4 b x^{\frac{5}{6}} \cos \left(b x^{\frac{1}{3}} + a \right) \sin \left(b x^{\frac{1}{3}} + a \right) \right)}{x}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(3/2),x, algorithm="fricas")`

output `-2*(4*pi*b*x*sqrt(b/pi)*cos(2*a)*fresnel_cos(2*x^(1/6)*sqrt(b/pi)) - 4*pi*b*x*sqrt(b/pi)*fresnel_sin(2*x^(1/6)*sqrt(b/pi))*sin(2*a) - 4*b*x^(5/6)*cos(b*x^(1/3) + a)*sin(b*x^(1/3) + a) + sqrt(x)*cos(b*x^(1/3) + a)^2)/x`

Sympy [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{\frac{3}{2}}} dx$$

input `integrate(cos(a+b*x**(1/3))**2/x**(3/2),x)`

output `Integral(cos(a + b*x**(1/3))**2/x**(3/2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.76

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx = \frac{3 \sqrt{2} \left(\left(-(i-1) \sqrt{2} \Gamma \left(-\frac{3}{2}, 2i b x^{\frac{1}{3}} \right) + (i+1) \sqrt{2} \Gamma \left(-\frac{3}{2}, -2i b x^{\frac{1}{3}} \right) \right) \cos(2a) + \left(- \right)}{4 \sqrt{2}}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(3/2),x, algorithm="maxima")`

output `1/4*(3*sqrt(2)*((-I - 1)*sqrt(2)*gamma(-3/2, 2*I*b*x^(1/3)) + (I + 1)*sqrt(2)*gamma(-3/2, -2*I*b*x^(1/3)))*cos(2*a) + (-I + 1)*sqrt(2)*gamma(-3/2, 2*I*b*x^(1/3)) + (I - 1)*sqrt(2)*gamma(-3/2, -2*I*b*x^(1/3))*sin(2*a))*sqrt(b*x^(1/3))*b*x^(1/3) - 4)/sqrt(x)`

Giac [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)^2}{x^{\frac{3}{2}}} dx$$

input `integrate(cos(a+b*x^(1/3))^2/x^(3/2),x, algorithm="giac")`

output `integrate(cos(b*x^(1/3) + a)^2/x^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos(a + bx^{1/3})^2}{x^{3/2}} dx$$

input `int(cos(a + b*x^(1/3))^2/x^(3/2),x)`

output `int(cos(a + b*x^(1/3))^2/x^(3/2), x)`

Reduce [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} dx = \int \frac{\cos(x^{1/3}b + a)^2}{\sqrt{x}x} dx$$

input `int(cos(a+b*x^(1/3))^2/x^(3/2),x)`

output `int(cos(x**(1/3)*b + a)**2/(sqrt(x)*x),x)`

3.59 $\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{5/2}} dx$

Optimal result	464
Mathematica [A] (verified)	465
Rubi [A] (verified)	465
Maple [A] (verified)	470
Fricas [A] (verification not implemented)	472
Sympy [F]	472
Maxima [C] (verification not implemented)	473
Giac [F]	473
Mupad [F(-1)]	474
Reduce [F]	474

Optimal result

Integrand size = 18, antiderivative size = 228

$$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{5/2}} dx = -\frac{16b^2}{105x^{5/6}} + \frac{256b^4}{315\sqrt[6]{x}} - \frac{2\cos^2(a+b\sqrt[3]{x})}{3x^{3/2}}$$

$$+ \frac{32b^2\cos^2(a+b\sqrt[3]{x})}{105x^{5/6}} - \frac{512b^4\cos^2(a+b\sqrt[3]{x})}{315\sqrt[6]{x}}$$

$$- \frac{512}{315}b^{9/2}\sqrt{\pi}\cos(2a)\text{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right) - \frac{512}{315}b^{9/2}\sqrt{\pi}\text{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)\sin(2a) + \frac{8b\cos(a+b\sqrt[3]{x})\sin(a+b\sqrt[3]{x})}{21x^{7/6}}$$

output

```
-16/105*b^2/x^(5/6)+256/315*b^4/x^(1/6)-2/3*cos(a+b*x^(1/3))^2/x^(3/2)+32/
105*b^2*cos(a+b*x^(1/3))^2/x^(5/6)-512/315*b^4*cos(a+b*x^(1/3))^2/x^(1/6)-
512/315*b^(9/2)*Pi^(1/2)*cos(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))-512
/315*b^(9/2)*Pi^(1/2)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))*sin(2*a)+8/21*b
*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/x^(7/6)-128/315*b^3*cos(a+b*x^(1/3))*si
n(a+b*x^(1/3))/x^(1/2)
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.81

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx = \frac{-105 - 105 \cos(2(a + b\sqrt[3]{x})) + 48b^2x^{2/3} \cos(2(a + b\sqrt[3]{x})) - 256b^4x^{4/3} \cos(2(a + b\sqrt[3]{x})) - 512b^6x^{5/3} \cos(2(a + b\sqrt[3]{x})) + 64b^8x^{2/3} \cos(2(a + b\sqrt[3]{x})) - 64b^8x^{2/3} \sin(2(a + b\sqrt[3]{x}))}{(315x^{3/2})}$$

input `Integrate[Cos[a + b*x^(1/3)]^2/x^(5/2),x]`

output

```
(-105 - 105*Cos[2*(a + b*x^(1/3))] + 48*b^2*x^(2/3)*Cos[2*(a + b*x^(1/3))]
- 256*b^4*x^(4/3)*Cos[2*(a + b*x^(1/3))] - 512*b^(9/2)*Sqrt[Pi]*x^(3/2)*C
os[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]] - 512*b^(9/2)*Sqrt[Pi]*x^(3
/2)*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 60*b*x^(1/3)*Sin[2*(
a + b*x^(1/3))] - 64*b^3*x*Ssin[2*(a + b*x^(1/3))])/(315*x^(3/2))
```

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.03, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.944$, Rules used = {3897, 3042, 3795, 15, 3042, 3795, 15, 3042, 3794, 27, 3042, 3787, 3042, 3785, 3786, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx \\ & \quad \downarrow \text{3897} \\ & 3 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{11/6}} d\sqrt[3]{x} \\ & \quad \downarrow \text{3042} \\ & 3 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{11/6}} d\sqrt[3]{x} \\ & \quad \downarrow \text{3795} \end{aligned}$$

$$3 \left(-\frac{16}{63} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/6}} d\sqrt[3]{x} + \frac{8}{63} b^2 \int \frac{1}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{9x^{3/2}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{63x^{7/6}} \right)$$

↓ 15

$$3 \left(-\frac{16}{63} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{9x^{3/2}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{63x^{7/6}} - \frac{16b^2}{315x^{5/6}} \right)$$

↓ 3042

$$3 \left(-\frac{16}{63} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{7/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{9x^{3/2}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{63x^{7/6}} - \frac{16b^2}{315x^{5/6}} \right)$$

↓ 3795

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} + \frac{8}{15} b^2 \int \frac{1}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{15\sqrt{x}} \right) \right)$$

↓ 15

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{15\sqrt{x}} - \frac{16b^2}{15\sqrt[6]{x}} \right) - \frac{2}{15\sqrt[6]{x}} \right)$$

↓ 3042

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{\sqrt{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{15\sqrt{x}} - \frac{16b^2}{15\sqrt[6]{x}} \right) \right)$$

↓ 3794

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(4b \int -\frac{\sin(2a + 2b\sqrt[3]{x})}{2\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{15\sqrt{x}} \right) \right)$$

↓ 27

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \int \frac{\sin(2a + 2b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{15\sqrt{x}} \right) \right)$$

↓ 3042

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \int \frac{\sin(2a + 2b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

↓ 3787

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \left(\sin(2a) \int \frac{\cos(2b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} + \cos(2a) \int \frac{\sin(2b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

↓ 3042

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \left(\sin(2a) \int \frac{\sin(2\sqrt[3]{xb} + \frac{\pi}{2})}{\sqrt[6]{x}} d\sqrt[3]{x} + \cos(2a) \int \frac{\sin(2b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

↓ 3785

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \left(2 \sin(2a) \int \cos(2bx^{2/3}) d\sqrt[6]{x} + \cos(2a) \int \frac{\sin(2b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

↓ 3786

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \left(2 \sin(2a) \int \cos(2bx^{2/3}) d\sqrt[6]{x} + 2 \cos(2a) \int \sin(2bx^{2/3}) d\sqrt[6]{x} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

↓ 3832

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \left(2 \sin(2a) \int \cos(2bx^{2/3}) d\sqrt[6]{x} + \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{\sqrt{b}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

↓ 3833

$$3 \left(-\frac{16}{63} b^2 \left(-\frac{16}{15} b^2 \left(-2b \left(\frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{\sqrt{b}} + \frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{\sqrt{b}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} \right) - \frac{2 \cos^2(a + b\sqrt[3]{x})}{5x^{5/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{15\sqrt[6]{x}} \right) \right)$$

input

```
Int[Cos[a + b*x^(1/3)]^2/x^(5/2), x]
```

output

```

3*((-16*b^2)/(315*x^(5/6)) - (2*Cos[a + b*x^(1/3)]^2)/(9*x^(3/2)) + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)])/(63*x^(7/6)) - (16*b^2*(-16*b^2)/(15*x^(1/6)) - (2*Cos[a + b*x^(1/3)]^2)/(5*x^(5/6)) - (16*b^2*(-2*Cos[a + b*x^(1/3)]^2)/x^(1/6) - 2*b*((Sqrt[Pi]*Cos[2*a]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/Sqrt[b] + (Sqrt[Pi]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/Sqrt[b])))/15 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]/(15*Sqrt[x])))/63)

```

Defintions of rubi rules used

rule 15

```

Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]

```

rule 27

```

Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]

```

rule 3042

```

Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]

```

rule 3785

```

Int[sin[Pi/2 + (e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Cos[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]

```

rule 3786

```

Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[Sin[f*(x^2/d)], x], x, Sqrt[c + d*x]], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && EqQ[d*e - c*f, 0]

```

rule 3787

```

Int[sin[(e_.) + (f_.)*(x_)]/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/Sqrt[c + d*x], x], x] /; FreeQ[{c, d, e, f}, x] && ComplexFreeQ[f] && NeQ[d*e - c*f, 0]

```

rule 3794 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]^n/(d*(m + 1))), x] - Simp[f*(n/(d*(m + 1))) Int[ExpandTrigReduce[(c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]^(n - 1), x], x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] && LtQ[m, -1]`

rule 3795 `Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(c + d*x)^(m + 1)*((b*Sine + f*x)^n/(d*(m + 1))), x] + (-Simp[b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x]*((b*Sine + f*x)^(n - 1)/(d^2*(m + 1)*(m + 2))), x] + Simp[b^2*f^2*n*((n - 1)/(d^2*(m + 1)*(m + 2))) Int[(c + d*x)^(m + 2)*(b*Sine + f*x)^(n - 2), x], x] - Simp[f^2*(n^2/(d^2*(m + 1)*(m + 2))) Int[(c + d*x)^(m + 2)*(b*Sine + f*x)^n, x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_))^(2)], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3897 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)]^(n_)]*(b_.)^(p_.)*(x_)]^(m_.), x_Symbol] := Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a + b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x] && IntegerQ[p] && FractionQ[n]`

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.64

method	result
derivativedivides	$-\frac{1}{3x^{\frac{3}{2}}} - \frac{\cos\left(2a+2bx^{\frac{1}{3}}\right)}{3x^{\frac{3}{2}}} - \frac{4b}{7x^{\frac{7}{6}}} + \frac{4b}{5x^{\frac{5}{6}}} \left(\frac{\sin\left(2a+2bx^{\frac{1}{3}}\right)}{3\sqrt{x}} + \frac{4b}{x^{\frac{1}{6}}} \frac{\cos\left(2a+2bx^{\frac{1}{3}}\right)}{3} \right)$
default	$-\frac{1}{3x^{\frac{3}{2}}} - \frac{\cos\left(2a+2bx^{\frac{1}{3}}\right)}{3x^{\frac{3}{2}}} - \frac{4b}{7x^{\frac{7}{6}}} + \frac{4b}{5x^{\frac{5}{6}}} \left(\frac{\sin\left(2a+2bx^{\frac{1}{3}}\right)}{3\sqrt{x}} + \frac{4b}{x^{\frac{1}{6}}} \frac{\cos\left(2a+2bx^{\frac{1}{3}}\right)}{3} \right)$

input `int(cos(a+b*x^(1/3))^2/x^(5/2),x,method=_RETURNVERBOSE)`

output `-1/3/x^(3/2)-1/3/x^(3/2)*cos(2*a+2*b*x^(1/3))-4/3*b*(-1/7/x^(7/6)*sin(2*a+2*b*x^(1/3))+4/7*b*(-1/5/x^(5/6)*cos(2*a+2*b*x^(1/3))-4/5*b*(-1/3/x^(1/2)*sin(2*a+2*b*x^(1/3))+4/3*b*(-1/x^(1/6)*cos(2*a+2*b*x^(1/3))-2*b^(1/2)*Pi^(1/2)*(cos(2*a)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))+sin(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2))))))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.68

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx =$$

$$2 \left(256 \pi b^4 x^2 \sqrt{\frac{b}{\pi}} \cos(2a) S \left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) + 256 \pi b^4 x^2 \sqrt{\frac{b}{\pi}} C \left(2 x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}} \right) \sin(2a) - 128 b^4 x^{\frac{11}{6}} + 24 b^2 x^{\frac{7}{6}} + \dots \right)$$

315

input `integrate(cos(a+b*x^(1/3))^2/x^(5/2),x, algorithm="fricas")`

output `-2/315*(256*pi*b^4*x^2*sqrt(b/pi)*cos(2*a)*fresnel_sin(2*x^(1/6)*sqrt(b/pi)) + 256*pi*b^4*x^2*sqrt(b/pi)*fresnel_cos(2*x^(1/6)*sqrt(b/pi))*sin(2*a) - 128*b^4*x^(11/6) + 24*b^2*x^(7/6) + (256*b^4*x^(11/6) - 48*b^2*x^(7/6) + 105*sqrt(x))*cos(b*x^(1/3) + a)^2 + 4*(16*b^3*x^(3/2) - 15*b*x^(5/6))*cos(b*x^(1/3) + a)*sin(b*x^(1/3) + a)/x^2`

Sympy [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx = \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{\frac{5}{2}}} dx$$

input `integrate(cos(a+b*x**(1/3))**2/x**(5/2),x)`

output `Integral(cos(a + b*x**(1/3))**2/x**(5/2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.39

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx = \frac{18\sqrt{2}\left(\left((i+1)\sqrt{2}\Gamma\left(-\frac{9}{2}, 2i bx^{\frac{1}{3}}\right) - (i-1)\sqrt{2}\Gamma\left(-\frac{9}{2}, -2i bx^{\frac{1}{3}}\right)\right)\cos(2a) + \left(-(i-1)\sqrt{2}\Gamma\left(-\frac{9}{2}, 2i bx^{\frac{1}{3}}\right) + (i+1)\sqrt{2}\Gamma\left(-\frac{9}{2}, -2i bx^{\frac{1}{3}}\right)\right)\sin(2a)\right)}{3x^{\frac{3}{2}}}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(5/2),x, algorithm="maxima")`

output `-1/3*(18*sqrt(2)*(((I + 1)*sqrt(2)*gamma(-9/2, 2*I*b*x^(1/3)) - (I - 1)*sqrt(2)*gamma(-9/2, -2*I*b*x^(1/3)))*cos(2*a) + (-(I - 1)*sqrt(2)*gamma(-9/2, 2*I*b*x^(1/3)) + (I + 1)*sqrt(2)*gamma(-9/2, -2*I*b*x^(1/3)))*sin(2*a))*sqrt(b*x^(1/3))*b^4*x^(4/3) + 1)/x^(3/2)`

Giac [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx = \int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)^2}{x^{\frac{5}{2}}} dx$$

input `integrate(cos(a+b*x^(1/3))^2/x^(5/2),x, algorithm="giac")`

output `integrate(cos(b*x^(1/3) + a)^2/x^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx = \int \frac{\cos(a + bx^{1/3})^2}{x^{5/2}} dx$$

input `int(cos(a + b*x^(1/3))^2/x^(5/2), x)`output `int(cos(a + b*x^(1/3))^2/x^(5/2), x)`**Reduce [F]**

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/2}} dx = \text{too large to display}$$

input `int(cos(a+b*x^(1/3))^2/x^(5/2), x)`

output

```
(2*(- 540*x**(1/3)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*tan((x**(1/3)*
b + a)/2)**4*b - 1080*x**(1/3)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*tan
((x**(1/3)*b + a)/2)**2*b - 540*x**(1/3)*cos(x**(1/3)*b + a)*sin(x**(1/3)*
b + a)*b + 7560*cos(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**4 + 15120*cos
(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**2 + 7560*cos(x**(1/3)*b + a) + 3
072*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**3/(sqrt(x)*tan((x**(1/3)*b + a)/2)
)**4*x + 2*sqrt(x)*tan((x**(1/3)*b + a)/2)**2*x + sqrt(x)*x),x)*tan((x**(1
/3)*b + a)/2)**4*b**3*x + 6144*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**3/(sqr
t(x)*tan((x**(1/3)*b + a)/2)**4*x + 2*sqrt(x)*tan((x**(1/3)*b + a)/2)**2*x
+ sqrt(x)*x),x)*tan((x**(1/3)*b + a)/2)**2*b**3*x + 3072*sqrt(x)*int(tan(
(x**(1/3)*b + a)/2)**3/(sqrt(x)*tan((x**(1/3)*b + a)/2)**4*x + 2*sqrt(x)*t
an((x**(1/3)*b + a)/2)**2*x + sqrt(x)*x),x)*b**3*x - 120960*sqrt(x)*int(ta
n((x**(1/3)*b + a)/2)**2/(sqrt(x)*tan((x**(1/3)*b + a)/2)**4*x**2 + 2*sqrt
(x)*tan((x**(1/3)*b + a)/2)**2*x**2 + sqrt(x)*x**2),x)*tan((x**(1/3)*b + a
)/2)**4*x - 241920*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**2/(sqrt(x)*tan((x*
*(1/3)*b + a)/2)**4*x**2 + 2*sqrt(x)*tan((x**(1/3)*b + a)/2)**2*x**2 + sqr
t(x)*x**2),x)*tan((x**(1/3)*b + a)/2)**2*x - 120960*sqrt(x)*int(tan((x**(1
/3)*b + a)/2)**2/(sqrt(x)*tan((x**(1/3)*b + a)/2)**4*x**2 + 2*sqrt(x)*tan(
(x**(1/3)*b + a)/2)**2*x**2 + sqrt(x)*x**2),x)*x - 2176*sqrt(x)*int(x**(1/
6)/(x**(1/3)*tan((x**(1/3)*b + a)/2)**4*x + 2*x**(1/3)*tan((x**(1/3)*b ...
```

3.60 $\int \frac{\cos^2\left(a+b\sqrt[3]{x}\right)}{x^{7/2}} dx$

Optimal result	476
Mathematica [A] (verified)	477
Rubi [A] (verified)	477
Maple [A] (verified)	481
Fricas [A] (verification not implemented)	483
Sympy [F]	483
Maxima [C] (verification not implemented)	484
Giac [F]	484
Mupad [F(-1)]	485
Reduce [F]	485

Optimal result

Integrand size = 18, antiderivative size = 328

$$\int \frac{\cos^2(a+b\sqrt[3]{x})}{x^{7/2}} dx = -\frac{16b^2}{715x^{11/6}} + \frac{256b^4}{45045x^{7/6}} - \frac{4096b^6}{675675\sqrt{x}}$$

$$- \frac{2\cos^2(a+b\sqrt[3]{x})}{5x^{5/2}} + \frac{32b^2\cos^2(a+b\sqrt[3]{x})}{715x^{11/6}} - \frac{512b^4\cos^2(a+b\sqrt[3]{x})}{45045x^{7/6}}$$

$$+ \frac{8192b^6\cos^2(a+b\sqrt[3]{x})}{675675\sqrt{x}} + \frac{32768b^{15/2}\sqrt{\pi}\cos(2a)\operatorname{FresnelC}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)}{675675}$$

$$- \frac{32768b^{15/2}\sqrt{\pi}\operatorname{FresnelS}\left(\frac{2\sqrt{b}\sqrt[6]{x}}{\sqrt{\pi}}\right)\sin(2a)}{675675}$$

$$+ \frac{8b\cos(a+b\sqrt[3]{x})\sin(a+b\sqrt[3]{x})}{65x^{13/6}} - \frac{128b^3\cos(a+b\sqrt[3]{x})\sin(a+b\sqrt[3]{x})}{6435x^{3/2}}$$

$$+ \frac{2048b^5\cos(a+b\sqrt[3]{x})\sin(a+b\sqrt[3]{x})}{225225x^{5/6}} - \frac{32768b^7\cos(a+b\sqrt[3]{x})\sin(a+b\sqrt[3]{x})}{675675\sqrt[3]{x}}$$

output

```
-16/715*b^2/x^(11/6)+256/45045*b^4/x^(7/6)-4096/675675*b^6/x^(1/2)-2/5*cos
(a+b*x^(1/3))^2/x^(5/2)+32/715*b^2*cos(a+b*x^(1/3))^2/x^(11/6)-512/45045*b
^4*cos(a+b*x^(1/3))^2/x^(7/6)+8192/675675*b^6*cos(a+b*x^(1/3))^2/x^(1/2)+3
2768/675675*b^(15/2)*Pi^(1/2)*cos(2*a)*FresnelC(2*b^(1/2)*x^(1/6)/Pi^(1/2)
)-32768/675675*b^(15/2)*Pi^(1/2)*FresnelS(2*b^(1/2)*x^(1/6)/Pi^(1/2))*sin(
2*a)+8/65*b*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/x^(13/6)-128/6435*b^3*cos(a+
b*x^(1/3))*sin(a+b*x^(1/3))/x^(3/2)+2048/225225*b^5*cos(a+b*x^(1/3))*sin(a
+b*x^(1/3))/x^(5/6)-32768/675675*b^7*cos(a+b*x^(1/3))*sin(a+b*x^(1/3))/x^(
1/6)
```

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.76

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{-135135 - 135135 \cos(2(a + b\sqrt[3]{x})) + 15120b^2x^{2/3} \cos(2(a + b\sqrt[3]{x})) - 3840b^4x^4}{x^{7/2}}$$

input

```
Integrate[Cos[a + b*x^(1/3)]^2/x^(7/2),x]
```

output

```
(-135135 - 135135*Cos[2*(a + b*x^(1/3))] + 15120*b^2*x^(2/3)*Cos[2*(a + b*
x^(1/3))] - 3840*b^4*x^(4/3)*Cos[2*(a + b*x^(1/3))] + 4096*b^6*x^2*Cos[2*(
a + b*x^(1/3))] + 32768*b^(15/2)*Sqrt[Pi]*x^(5/2)*Cos[2*a]*FresnelC[(2*Sqr
t[b]*x^(1/6))/Sqrt[Pi]] - 32768*b^(15/2)*Sqrt[Pi]*x^(5/2)*FresnelS[(2*Sqrt
[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a] + 41580*b*x^(1/3)*Sin[2*(a + b*x^(1/3))] -
6720*b^3*x*Sin[2*(a + b*x^(1/3))] + 3072*b^5*x^(5/3)*Sin[2*(a + b*x^(1/3)
)] - 16384*b^7*x^(7/3)*Sin[2*(a + b*x^(1/3))])/(675675*x^(5/2))
```

Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.11, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {3897, 3042, 3795, 15, 3042, 3795, 15, 3042, 3795, 15, 3042, 3795, 15, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx \\
& \quad \downarrow \text{3897} \\
& 3 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{17/6}} d\sqrt[3]{x} \\
& \quad \downarrow \text{3042} \\
& 3 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{17/6}} d\sqrt[3]{x} \\
& \quad \downarrow \text{3795} \\
& 3 \left(-\frac{16}{195} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{13/6}} d\sqrt[3]{x} + \frac{8}{195} b^2 \int \frac{1}{x^{13/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{15x^{5/2}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{195x^{13/6}} \right) \\
& \quad \downarrow \text{15} \\
& 3 \left(-\frac{16}{195} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{13/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{15x^{5/2}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{195x^{13/6}} - \frac{16b^2}{2145x^{11/6}} \right) \\
& \quad \downarrow \text{3042} \\
& 3 \left(-\frac{16}{195} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{13/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{15x^{5/2}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{195x^{13/6}} - \frac{16b^2}{2145x^{11/6}} \right) \\
& \quad \downarrow \text{3795} \\
& 3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} d\sqrt[3]{x} + \frac{8}{99} b^2 \int \frac{1}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{11x^{11/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{99x^{3/2}} \right) \right) \\
& \quad \downarrow \text{15} \\
& 3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{11x^{11/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{99x^{3/2}} - \frac{16b^2}{693x^{7/6}} \right) \right) \\
& \quad \downarrow \text{3042} \\
& 3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{3/2}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{11x^{11/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{99x^{3/2}} - \frac{16b^2}{693x^{7/6}} \right) \right)
\end{aligned}$$

↓ 3795

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} + \frac{8}{35} b^2 \int \frac{1}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{7x^{7/6}} + \frac{8b \sin(a + b\sqrt[3]{x})}{35x^{5/6}} \right) \right) \right)$$

↓ 15

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{7x^{7/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{35x^{5/6}} - \frac{1}{10} \right) \right) \right)$$

↓ 3042

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{x^{5/6}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{7x^{7/6}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{35x^{5/6}} \right) \right) \right)$$

↓ 3795

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \left(-\frac{16}{3} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} + \frac{8}{3} b^2 \int \frac{1}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \right) \right) \right)$$

↓ 15

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \left(-\frac{16}{3} b^2 \int \frac{\cos^2(a + b\sqrt[3]{x})}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \right) \right) \right)$$

↓ 3042

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \left(-\frac{16}{3} b^2 \int \frac{\sin(a + b\sqrt[3]{x} + \frac{\pi}{2})^2}{\sqrt[6]{x}} d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x}) \cos(a + b\sqrt[3]{x})}{3\sqrt[6]{x}} \right) \right) \right) \right)$$

↓ 3793

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \left(-\frac{16}{3} b^2 \int \left(\frac{\cos(2a + 2b\sqrt[3]{x})}{2\sqrt[6]{x}} + \frac{1}{2\sqrt[6]{x}} \right) d\sqrt[3]{x} - \frac{2 \cos^2(a + b\sqrt[3]{x})}{3\sqrt{x}} + \frac{8b \sin(a + b\sqrt[3]{x})}{3\sqrt{x}} \right) \right) \right) \right)$$

↓ 2009

$$3 \left(-\frac{16}{195} b^2 \left(-\frac{16}{99} b^2 \left(-\frac{16}{35} b^2 \left(-\frac{16}{3} b^2 \left(\frac{\sqrt{\pi} \cos(2a) \operatorname{FresnelC} \left(\frac{2\sqrt{b} \sqrt[6]{x}}{\sqrt{\pi}} \right)}{2\sqrt{b}} - \frac{\sqrt{\pi} \sin(2a) \operatorname{FresnelS} \left(\frac{2\sqrt{b} \sqrt[6]{x}}{\sqrt{\pi}} \right)}{2\sqrt{b}} + \sqrt[6]{x} \right. \right. \right. \right. \right. \right.$$

input `Int[Cos[a + b*x^(1/3)]^2/x^(7/2), x]`

output `3*((-16*b^2)/(2145*x^(11/6)) - (2*Cos[a + b*x^(1/3)]^2)/(15*x^(5/2)) + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]/(195*x^(13/6)) - (16*b^2*((-16*b^2)/(693*x^(7/6)) - (2*Cos[a + b*x^(1/3)]^2)/(11*x^(11/6)) + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]/(99*x^(3/2)) - (16*b^2*((-16*b^2)/(105*Sqrt[x]) - (2*Cos[a + b*x^(1/3)]^2)/(7*x^(7/6)) + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]/(35*x^(5/6)) - (16*b^2*((16*b^2*x^(1/6))/3 - (2*Cos[a + b*x^(1/3)]^2)/(3*Sqrt[x]) - (16*b^2*(x^(1/6) + (Sqrt[Pi]*Cos[2*a]*FresnelC[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]])/(2*Sqrt[b]) - (Sqrt[Pi]*FresnelS[(2*Sqrt[b]*x^(1/6))/Sqrt[Pi]]*Sin[2*a])/(2*Sqrt[b])))/3 + (8*b*Cos[a + b*x^(1/3)]*Sin[a + b*x^(1/3)]/(3*x^(1/6))))/35)/99)/195)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)^(n_)], x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 3795

```
Int[((c_.) + (d_.)*(x_)^(m_))*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol]
:> Simp[(c + d*x)^(m + 1)*((b*Sine[e + f*x])^n/(d*(m + 1))), x] + (-Simp[
b*f*n*(c + d*x)^(m + 2)*Cos[e + f*x]*((b*Sine[e + f*x])^(n - 1)/(d^2*(m + 1)
*(m + 2))), x] + Simp[b^2*f^2*n*((n - 1)/(d^2*(m + 1)*(m + 2))) Int[(c +
d*x)^(m + 2)*(b*Sine[e + f*x])^(n - 2), x], x] - Simp[f^2*(n^2/(d^2*(m + 1)*
(m + 2))) Int[(c + d*x)^(m + 2)*(b*Sine[e + f*x])^n, x], x]) /; FreeQ[{b,
c, d, e, f}, x] && GtQ[n, 1] && LtQ[m, -2]
```

rule 3897

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
:> Module[{k = Denominator[n]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*(a +
b*Cos[c + d*x^(k*n)]]^p, x], x, x^(1/k)], x] /; FreeQ[{a, b, c, d, m}, x]
&& IntegerQ[p] && FractionQ[n]
```

Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.63

method	result
	$4b \frac{\cos\left(2a+2b x^{\frac{1}{3}}\right)}{7x^{\frac{7}{6}}}$ $4b \frac{\sin\left(2a+2b x^{\frac{1}{3}}\right)}{9x^{\frac{3}{2}}} +$ $4b \frac{\cos\left(2a+2b x^{\frac{1}{3}}\right)}{11x^{\frac{11}{6}}}$

input `int(cos(a+b*x^(1/3))^2/x^(7/2),x,method=_RETURNVERBOSE)`

output
$$-1/5/x^{(5/2)}-1/5/x^{(5/2)}*\cos(2*a+2*b*x^{(1/3)})-4/5*b*(-1/13/x^{(13/6)}*\sin(2*a+2*b*x^{(1/3)})+4/13*b*(-1/11/x^{(11/6)}*\cos(2*a+2*b*x^{(1/3)})-4/11*b*(-1/9/x^{(3/2)}*\sin(2*a+2*b*x^{(1/3)})+4/9*b*(-1/7/x^{(7/6)}*\cos(2*a+2*b*x^{(1/3)})-4/7*b*(-1/5/x^{(5/6)}*\sin(2*a+2*b*x^{(1/3)})+4/5*b*(-1/3/x^{(1/2)}*\cos(2*a+2*b*x^{(1/3)})-4/3*b*(-1/x^{(1/6)}*\sin(2*a+2*b*x^{(1/3)})+2*b^{(1/2)}*Pi^{(1/2)}*(\cos(2*a)*FresnelC(2*b^{(1/2)}*x^{(1/6)}/Pi^{(1/2)})-\sin(2*a)*FresnelS(2*b^{(1/2)}*x^{(1/6)}/Pi^{(1/2)})))))))))$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.59

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{2 \left(16384 \pi b^7 x^3 \sqrt{\frac{b}{\pi}} \cos(2a) C\left(2x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) - 16384 \pi b^7 x^3 \sqrt{\frac{b}{\pi}} S\left(2x^{\frac{1}{6}} \sqrt{\frac{b}{\pi}}\right) \sin(2a) \right)}{x^{7/2}}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(7/2),x, algorithm="fricas")`

output
$$\frac{2/675675*(16384*\pi*b^7*x^3*\sqrt{b/\pi}*\cos(2*a)*fresnel_cos(2*x^{(1/6)}*\sqrt{b/\pi}) - 16384*\pi*b^7*x^3*\sqrt{b/\pi}*\fresnel_sin(2*x^{(1/6)}*\sqrt{b/\pi}))*\sin(2*a) - 2048*b^6*x^{(5/2)} + 1920*b^4*x^{(11/6)} - 7560*b^2*x^{(7/6)} - (3840*b^4*x^{(11/6)} - 15120*b^2*x^{(7/6)} - (4096*b^6*x^2 - 135135)*\sqrt{x})*\cos(b*x^{(1/3)} + a)^2 + 4*(768*b^5*x^{(13/6)} - 1680*b^3*x^{(3/2)} - (4096*b^7*x^2 - 10395*b)*x^{(5/6)})*\cos(b*x^{(1/3)} + a)*\sin(b*x^{(1/3)} + a)}{x^3}$$

Sympy [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{\frac{7}{2}}} dx$$

input `integrate(cos(a+b*x**(1/3))**2/x**(7/2),x)`

output `Integral(cos(a + b*x**(1/3))**2/x**(7/2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.27

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \frac{240\sqrt{2}\left(\left(-i-1\right)\sqrt{2}\Gamma\left(-\frac{15}{2}, 2i bx^{\frac{1}{3}}\right) + (i+1)\sqrt{2}\Gamma\left(-\frac{15}{2}, -2i bx^{\frac{1}{3}}\right)\right)\cos(2a) + \left(-i+1\right)\sqrt{2}\Gamma\left(-\frac{15}{2}, 2i bx^{\frac{1}{3}}\right)\sin(2a) + \left(i-1\right)\sqrt{2}\Gamma\left(-\frac{15}{2}, -2i bx^{\frac{1}{3}}\right)\sin(2a)}{5x^{\frac{5}{2}}}$$

input `integrate(cos(a+b*x^(1/3))^2/x^(7/2),x, algorithm="maxima")`

output `-1/5*(240*sqrt(2)*((-I - 1)*sqrt(2)*gamma(-15/2, 2*I*b*x^(1/3)) + (I + 1)*sqrt(2)*gamma(-15/2, -2*I*b*x^(1/3)))*cos(2*a) + (-I + 1)*sqrt(2)*gamma(-15/2, 2*I*b*x^(1/3)) + (I - 1)*sqrt(2)*gamma(-15/2, -2*I*b*x^(1/3))*sin(2*a))*sqrt(b*x^(1/3))*b^7*x^(7/3) + 1)/x^(5/2)`

Giac [F]

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \int \frac{\cos\left(bx^{\frac{1}{3}} + a\right)^2}{x^{\frac{7}{2}}} dx$$

input `integrate(cos(a+b*x^(1/3))^2/x^(7/2),x, algorithm="giac")`

output `integrate(cos(b*x^(1/3) + a)^2/x^(7/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \int \frac{\cos(a + bx^{1/3})^2}{x^{7/2}} dx$$

input `int(cos(a + b*x^(1/3))^2/x^(7/2), x)`output `int(cos(a + b*x^(1/3))^2/x^(7/2), x)`**Reduce [F]**

$$\int \frac{\cos^2(a + b\sqrt[3]{x})}{x^{7/2}} dx = \text{too large to display}$$

input `int(cos(a+b*x^(1/3))^2/x^(7/2), x)`

output

```
(2*( - 74844*x**(1/3)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)*tan((x**(1/3)
)*b + a)/2)**4*b - 149688*x**(1/3)*cos(x**(1/3)*b + a)*sin(x**(1/3)*b + a)
*tan((x**(1/3)*b + a)/2)**2*b - 74844*x**(1/3)*cos(x**(1/3)*b + a)*sin(x**
(1/3)*b + a)*b + 1945944*cos(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**4 +
3891888*cos(x**(1/3)*b + a)*tan((x**(1/3)*b + a)/2)**2 + 1945944*cos(x**(1
/3)*b + a) - 8192*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**3/(x**(1/6)*tan((x*
*(1/3)*b + a)/2)**4*x + 2*x**(1/6)*tan((x**(1/3)*b + a)/2)**2*x + x**(1/6)
*x),x)*tan((x**(1/3)*b + a)/2)**4*b**7*x**2 - 16384*sqrt(x)*int(tan((x**(1
/3)*b + a)/2)**3/(x**(1/6)*tan((x**(1/3)*b + a)/2)**4*x + 2*x**(1/6)*tan((
x**(1/3)*b + a)/2)**2*x + x**(1/6)*x),x)*tan((x**(1/3)*b + a)/2)**2*b**7*x
**2 - 8192*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**3/(x**(1/6)*tan((x**(1/3)*
b + a)/2)**4*x + 2*x**(1/6)*tan((x**(1/3)*b + a)/2)**2*x + x**(1/6)*x),x)*
b**7*x**2 + 193536*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**3/(sqrt(x)*tan((x*
*(1/3)*b + a)/2)**4*x**2 + 2*sqrt(x)*tan((x**(1/3)*b + a)/2)**2*x**2 + sqr
t(x)*x**2),x)*tan((x**(1/3)*b + a)/2)**4*b**3*x**2 + 387072*sqrt(x)*int(ta
n((x**(1/3)*b + a)/2)**3/(sqrt(x)*tan((x**(1/3)*b + a)/2)**4*x**2 + 2*sqrt
(x)*tan((x**(1/3)*b + a)/2)**2*x**2 + sqrt(x)*x**2),x)*tan((x**(1/3)*b + a
)/2)**2*b**3*x**2 + 193536*sqrt(x)*int(tan((x**(1/3)*b + a)/2)**3/(sqrt(x)
*tan((x**(1/3)*b + a)/2)**4*x**2 + 2*sqrt(x)*tan((x**(1/3)*b + a)/2)**2*x*
*2 + sqrt(x)*x**2),x)*b**3*x**2 + 107520*sqrt(x)*int(tan((x**(1/3)*b + ...
```

3.61 $\int \cos^3(\sqrt[3]{x}) dx$

Optimal result	487
Mathematica [A] (verified)	487
Rubi [A] (warning: unable to verify)	488
Maple [A] (verified)	491
Fricas [A] (verification not implemented)	491
Sympy [B] (verification not implemented)	492
Maxima [A] (verification not implemented)	492
Giac [A] (verification not implemented)	493
Mupad [B] (verification not implemented)	493
Reduce [B] (verification not implemented)	494

Optimal result

Integrand size = 8, antiderivative size = 86

$$\int \cos^3(\sqrt[3]{x}) dx = 4\sqrt[3]{x} \cos(\sqrt[3]{x}) + \frac{2}{3}\sqrt[3]{x} \cos^3(\sqrt[3]{x}) - \frac{14}{3} \sin(\sqrt[3]{x}) + 2x^{2/3} \sin(\sqrt[3]{x}) + x^{2/3} \cos^2(\sqrt[3]{x}) \sin(\sqrt[3]{x}) + \frac{2}{9} \sin^3(\sqrt[3]{x})$$

output

```
4*x^(1/3)*cos(x^(1/3))+2/3*x^(1/3)*cos(x^(1/3))^3-14/3*sin(x^(1/3))+2*x^(2/3)*sin(x^(1/3))+x^(2/3)*cos(x^(1/3))^2*sin(x^(1/3))+2/9*sin(x^(1/3))^3
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.77

$$\int \cos^3(\sqrt[3]{x}) dx = \frac{1}{36} (162\sqrt[3]{x} \cos(\sqrt[3]{x}) + 6\sqrt[3]{x} \cos(3\sqrt[3]{x}) + 81(-2 + x^{2/3}) \sin(\sqrt[3]{x}) + (-2 + 9x^{2/3}) \sin(3\sqrt[3]{x}))$$

input

```
Integrate[Cos[x^(1/3)]^3,x]
```

output

```
(162*x^(1/3)*Cos[x^(1/3)] + 6*x^(1/3)*Cos[3*x^(1/3)] + 81*(-2 + x^(2/3))*Sin[x^(1/3)] + (-2 + 9*x^(2/3))*Sin[3*x^(1/3)])/36
```


Rubi [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.16, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3843, 3042, 3792, 3042, 3113, 2009, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(\sqrt[3]{x}) dx$$

$$\downarrow \text{3843}$$

$$3 \int x^{2/3} \cos^3(\sqrt[3]{x}) d\sqrt[3]{x}$$

$$\downarrow \text{3042}$$

$$3 \int x^{2/3} \sin\left(\sqrt[3]{x} + \frac{\pi}{2}\right)^3 d\sqrt[3]{x}$$

$$\downarrow \text{3792}$$

$$3 \left(\frac{2}{3} \int x^{2/3} \cos(\sqrt[3]{x}) d\sqrt[3]{x} - \frac{2}{9} \int \cos^3(\sqrt[3]{x}) d\sqrt[3]{x} + \frac{1}{3} x^{2/3} \sin(\sqrt[3]{x}) \cos^2(\sqrt[3]{x}) + \frac{2}{9} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) \right)$$

$$\downarrow \text{3042}$$

$$3 \left(\frac{2}{3} \int x^{2/3} \sin\left(\sqrt[3]{x} + \frac{\pi}{2}\right) d\sqrt[3]{x} - \frac{2}{9} \int \sin\left(\sqrt[3]{x} + \frac{\pi}{2}\right)^3 d\sqrt[3]{x} + \frac{1}{3} x^{2/3} \sin(\sqrt[3]{x}) \cos^2(\sqrt[3]{x}) + \frac{2}{9} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) \right)$$

$$\downarrow \text{3113}$$

$$3 \left(\frac{2}{9} \int (1 - x^{2/3}) d(-\sin(\sqrt[3]{x})) + \frac{2}{3} \int x^{2/3} \sin\left(\sqrt[3]{x} + \frac{\pi}{2}\right) d\sqrt[3]{x} + \frac{1}{3} x^{2/3} \sin(\sqrt[3]{x}) \cos^2(\sqrt[3]{x}) + \frac{2}{9} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) \right)$$

$$\downarrow \text{2009}$$

$$3 \left(\frac{2}{3} \int x^{2/3} \sin\left(\sqrt[3]{x} + \frac{\pi}{2}\right) d\sqrt[3]{x} + \frac{1}{3} x^{2/3} \sin(\sqrt[3]{x}) \cos^2(\sqrt[3]{x}) + \frac{2}{9} \left(-\frac{x}{3} - \sin(\sqrt[3]{x}) \right) + \frac{2}{9} \sqrt[3]{x} \cos^3(\sqrt[3]{x}) \right)$$

$$\downarrow \text{3777}$$

$$3\left(\frac{2}{3}\left(2\int-\sqrt[3]{x}\sin(\sqrt[3]{x})d\sqrt[3]{x}+x^{2/3}\sin(\sqrt[3]{x})\right)+\frac{1}{3}x^{2/3}\sin(\sqrt[3]{x})\cos^2(\sqrt[3]{x})+\frac{2}{9}\left(-\frac{x}{3}-\sin(\sqrt[3]{x})\right)+\frac{2}{9}\sqrt[3]{x}\cos^3(\sqrt[3]{x})\right)$$

↓ 25

$$3\left(\frac{2}{3}\left(x^{2/3}\sin(\sqrt[3]{x})-2\int\sqrt[3]{x}\sin(\sqrt[3]{x})d\sqrt[3]{x}\right)+\frac{1}{3}x^{2/3}\sin(\sqrt[3]{x})\cos^2(\sqrt[3]{x})+\frac{2}{9}\left(-\frac{x}{3}-\sin(\sqrt[3]{x})\right)+\frac{2}{9}\sqrt[3]{x}\cos^3(\sqrt[3]{x})\right)$$

↓ 3042

$$3\left(\frac{2}{3}\left(x^{2/3}\sin(\sqrt[3]{x})-2\int\sqrt[3]{x}\sin(\sqrt[3]{x})d\sqrt[3]{x}\right)+\frac{1}{3}x^{2/3}\sin(\sqrt[3]{x})\cos^2(\sqrt[3]{x})+\frac{2}{9}\left(-\frac{x}{3}-\sin(\sqrt[3]{x})\right)+\frac{2}{9}\sqrt[3]{x}\cos^3(\sqrt[3]{x})\right)$$

↓ 3777

$$3\left(\frac{2}{3}\left(x^{2/3}\sin(\sqrt[3]{x})-2\left(\int\cos(\sqrt[3]{x})d\sqrt[3]{x}-\sqrt[3]{x}\cos(\sqrt[3]{x})\right)\right)+\frac{1}{3}x^{2/3}\sin(\sqrt[3]{x})\cos^2(\sqrt[3]{x})+\frac{2}{9}\left(-\frac{x}{3}-\sin(\sqrt[3]{x})\right)+\frac{2}{9}\sqrt[3]{x}\cos^3(\sqrt[3]{x})\right)$$

↓ 3042

$$3\left(\frac{2}{3}\left(x^{2/3}\sin(\sqrt[3]{x})-2\left(\int\sin\left(\sqrt[3]{x}+\frac{\pi}{2}\right)d\sqrt[3]{x}-\sqrt[3]{x}\cos(\sqrt[3]{x})\right)\right)+\frac{1}{3}x^{2/3}\sin(\sqrt[3]{x})\cos^2(\sqrt[3]{x})+\frac{2}{9}\left(-\frac{x}{3}-\sin(\sqrt[3]{x})\right)+\frac{2}{9}\sqrt[3]{x}\cos^3(\sqrt[3]{x})\right)$$

↓ 3117

$$3\left(\frac{1}{3}x^{2/3}\sin(\sqrt[3]{x})\cos^2(\sqrt[3]{x})+\frac{2}{3}\left(x^{2/3}\sin(\sqrt[3]{x})-2(\sin(\sqrt[3]{x})-\sqrt[3]{x}\cos(\sqrt[3]{x}))\right)+\frac{2}{9}\left(-\frac{x}{3}-\sin(\sqrt[3]{x})\right)+\frac{2}{9}\sqrt[3]{x}\cos^3(\sqrt[3]{x})\right)$$

input `Int [Cos [x^(1/3)]^3, x]`

output

```
3*((2*x^(1/3)*Cos[x^(1/3)]^3)/9 + (2*(-1/3*x - Sin[x^(1/3)]))/9 + (x^(2/3)
*Cos[x^(1/3)]^2*Sin[x^(1/3)])/3 + (2*(x^(2/3)*Sin[x^(1/3)] - 2*(-(x^(1/3))*
Cos[x^(1/3)] + Sin[x^(1/3)])))/3
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2009 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] \text{ /; SumQ}[\text{u}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3113 $\text{Int}[\sin[(\text{c}_.) + (\text{d}_.) * (\text{x}_)]^{(\text{n}_)}, \text{x_Symbol}] \rightarrow \text{Simp}[-\text{d}^{(-1)} \quad \text{Subst}[\text{Int}[\text{Exp and}[(1 - \text{x}^2)^{((\text{n} - 1)/2)}, \text{x}], \text{x}], \text{x}, \text{Cos}[\text{c} + \text{d} * \text{x}], \text{x}] \text{ /; FreeQ}\{\{\text{c}, \text{d}\}, \text{x}\} \&\& \text{IGtQ}[(\text{n} - 1)/2, 0]$
- rule 3117 $\text{Int}[\sin[\text{Pi}/2 + (\text{c}_.) + (\text{d}_.) * (\text{x}_)], \text{x_Symbol}] \rightarrow \text{Simp}[\text{Sin}[\text{c} + \text{d} * \text{x}]/\text{d}, \text{x}] \text{ /; FreeQ}\{\{\text{c}, \text{d}\}, \text{x}\}$
- rule 3777 $\text{Int}[((\text{c}_.) + (\text{d}_.) * (\text{x}_))^{(\text{m}_.)} * \sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_)], \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{c} + \text{d} * \text{x})^{\text{m}} * (\text{Cos}[\text{e} + \text{f} * \text{x}]/\text{f}), \text{x}] + \text{Simp}[\text{d} * (\text{m}/\text{f}) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{(\text{m} - 1)} * \text{Cos}[\text{e} + \text{f} * \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}\{\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{GtQ}[\text{m}, 0]$
- rule 3792 $\text{Int}[((\text{c}_.) + (\text{d}_.) * (\text{x}_))^{(\text{m}_.)} * ((\text{b}_.) * \sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_)])^{(\text{n}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{d} * \text{m} * (\text{c} + \text{d} * \text{x})^{(\text{m} - 1)} * ((\text{b} * \text{Sin}[\text{e} + \text{f} * \text{x}])^{\text{n}} / (\text{f}^2 * \text{n}^2)), \text{x}] + (-\text{Simp}[\text{b} * (\text{c} + \text{d} * \text{x})^{\text{m}} * \text{Cos}[\text{e} + \text{f} * \text{x}] * ((\text{b} * \text{Sin}[\text{e} + \text{f} * \text{x}])^{(\text{n} - 1)} / (\text{f} * \text{n})), \text{x}] + \text{Simp}[\text{b}^2 * ((\text{n} - 1)/\text{n}) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m}} * (\text{b} * \text{Sin}[\text{e} + \text{f} * \text{x}])^{(\text{n} - 2)}, \text{x}], \text{x}] - \text{Simp}[\text{d}^2 * \text{m} * ((\text{m} - 1)/(\text{f}^2 * \text{n}^2)) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{(\text{m} - 2)} * (\text{b} * \text{Sin}[\text{e} + \text{f} * \text{x}])^{\text{n}}, \text{x}], \text{x}]) \text{ /; FreeQ}\{\{\text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{GtQ}[\text{n}, 1] \&\& \text{GtQ}[\text{m}, 1]$
- rule 3843 $\text{Int}[((\text{a}_.) + \text{Cos}[(\text{c}_.) + (\text{d}_.) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_))^{(\text{n}_.)}] * (\text{b}_.))^{(\text{p}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[1/(\text{n} * \text{f}) \quad \text{Subst}[\text{Int}[\text{x}^{(1/\text{n} - 1)} * (\text{a} + \text{b} * \text{Cos}[\text{c} + \text{d} * \text{x}])^{\text{p}}, \text{x}], \text{x}, (\text{e} + \text{f} * \text{x})^{\text{n}}], \text{x}] \text{ /; FreeQ}\{\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{IGtQ}[\text{p}, 0] \&\& \text{IntegerQ}[1/\text{n}]$

Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.67

method	result
derivativedivides	$x^{\frac{2}{3}} \left(2 + \cos \left(x^{\frac{1}{3}} \right)^2 \right) \sin \left(x^{\frac{1}{3}} \right) - 4 \sin \left(x^{\frac{1}{3}} \right) + 4x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right) + \frac{2x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right)^3}{3} - \frac{2 \left(2 + \cos \left(x^{\frac{1}{3}} \right)^2 \right) \sin \left(x^{\frac{1}{3}} \right)}{3}$
default	$x^{\frac{2}{3}} \left(2 + \cos \left(x^{\frac{1}{3}} \right)^2 \right) \sin \left(x^{\frac{1}{3}} \right) - 4 \sin \left(x^{\frac{1}{3}} \right) + 4x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right) + \frac{2x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right)^3}{3} - \frac{2 \left(2 + \cos \left(x^{\frac{1}{3}} \right)^2 \right) \sin \left(x^{\frac{1}{3}} \right)}{3}$

input `int(cos(x^(1/3))^3,x,method=_RETURNVERBOSE)`output `x^(2/3)*(2+cos(x^(1/3))^2)*sin(x^(1/3))-4*sin(x^(1/3))+4*x^(1/3)*cos(x^(1/3))+2/3*x^(1/3)*cos(x^(1/3))^3-2/9*(2+cos(x^(1/3))^2)*sin(x^(1/3))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.56

$$\int \cos^3(\sqrt[3]{x}) dx = \frac{2}{3} x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right)^3 + \frac{1}{9} \left(\left(9x^{\frac{2}{3}} - 2 \right) \cos \left(x^{\frac{1}{3}} \right)^2 + 18x^{\frac{2}{3}} - 40 \right) \sin \left(x^{\frac{1}{3}} \right) + 4x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right)$$

input `integrate(cos(x^(1/3))^3,x, algorithm="fricas")`output `2/3*x^(1/3)*cos(x^(1/3))^3 + 1/9*((9*x^(2/3) - 2)*cos(x^(1/3))^2 + 18*x^(2/3) - 40)*sin(x^(1/3)) + 4*x^(1/3)*cos(x^(1/3))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 513 vs. $2(85) = 170$.

Time = 0.76 (sec) , antiderivative size = 513, normalized size of antiderivative = 5.97

$$\int \cos^3(\sqrt[3]{x}) dx = \text{Too large to display}$$

input `integrate(cos(x**(1/3))**3,x)`

output

```
54*x**(2/3)*tan(x**(1/3)/2)**5/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) + 36*x**(2/3)*tan(x**(1/3)/2)**3/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) + 54*x**(2/3)*tan(x**(1/3)/2)/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) - 42*x**(1/3)*tan(x**(1/3)/2)**6/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) - 18*x**(1/3)*tan(x**(1/3)/2)**4/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) + 18*x**(1/3)*tan(x**(1/3)/2)**2/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) + 42*x**(1/3)/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) - 84*tan(x**(1/3)/2)**5/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) - 152*tan(x**(1/3)/2)**3/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9) - 84*tan(x**(1/3)/2)/(9*tan(x**(1/3)/2)**6 + 27*tan(x**(1/3)/2)**4 + 27*tan(x**(1/3)/2)**2 + 9)
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.55

$$\int \cos^3(\sqrt[3]{x}) dx = \frac{1}{36} \left(9x^{\frac{2}{3}} - 2 \right) \sin \left(3x^{\frac{1}{3}} \right) + \frac{9}{4} \left(x^{\frac{2}{3}} - 2 \right) \sin \left(x^{\frac{1}{3}} \right) + \frac{1}{6} x^{\frac{1}{3}} \cos \left(3x^{\frac{1}{3}} \right) + \frac{9}{2} x^{\frac{1}{3}} \cos \left(x^{\frac{1}{3}} \right)$$

input `integrate(cos(x^(1/3))^3,x, algorithm="maxima")`

output $1/36*(9*x^{(2/3)} - 2)*\sin(3*x^{(1/3)}) + 9/4*(x^{(2/3)} - 2)*\sin(x^{(1/3)}) + 1/6*x^{(1/3)}*\cos(3*x^{(1/3)}) + 9/2*x^{(1/3)}*\cos(x^{(1/3)})$

Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.55

$$\int \cos^3(\sqrt[3]{x}) dx = \frac{1}{36} (9x^{2/3} - 2) \sin(3x^{1/3}) + \frac{9}{4} (x^{2/3} - 2) \sin(x^{1/3}) + \frac{1}{6} x^{1/3} \cos(3x^{1/3}) + \frac{9}{2} x^{1/3} \cos(x^{1/3})$$

input `integrate(cos(x^(1/3))^3,x, algorithm="giac")`

output $1/36*(9*x^{(2/3)} - 2)*\sin(3*x^{(1/3)}) + 9/4*(x^{(2/3)} - 2)*\sin(x^{(1/3)}) + 1/6*x^{(1/3)}*\cos(3*x^{(1/3)}) + 9/2*x^{(1/3)}*\cos(x^{(1/3)})$

Mupad [B] (verification not implemented)

Time = 41.25 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.72

$$\int \cos^3(\sqrt[3]{x}) dx = 4x^{1/3} \cos(x^{1/3}) - \frac{2 \cos(x^{1/3})^2 \sin(x^{1/3})}{9} - \frac{40 \sin(x^{1/3})}{9} + 2x^{2/3} \sin(x^{1/3}) + \frac{2x^{1/3} \cos(x^{1/3})^3}{3} + x^{2/3} \cos(x^{1/3})^2 \sin(x^{1/3})$$

input `int(cos(x^(1/3))^3,x)`

output $4*x^{(1/3)}*\cos(x^{(1/3)}) - (2*\cos(x^{(1/3)})^2*\sin(x^{(1/3)}))/9 - (40*\sin(x^{(1/3)}))/9 + 2*x^{(2/3)}*\sin(x^{(1/3)}) + (2*x^{(1/3)}*\cos(x^{(1/3)})^3)/3 + x^{(2/3)}*\cos(x^{(1/3)})^2*\sin(x^{(1/3)})$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.69

$$\int \cos^3(\sqrt[3]{x}) dx = -\frac{2x^{\frac{1}{3}} \cos(x^{\frac{1}{3}}) \sin(x^{\frac{1}{3}})^2}{3} + \frac{14x^{\frac{1}{3}} \cos(x^{\frac{1}{3}})}{3} - x^{\frac{2}{3}} \sin(x^{\frac{1}{3}})^3$$

$$+ 3x^{\frac{2}{3}} \sin(x^{\frac{1}{3}}) + \frac{2 \sin(x^{\frac{1}{3}})^3}{9} - \frac{14 \sin(x^{\frac{1}{3}})}{3}$$

input

```
int(cos(x^(1/3))^3,x)
```

output

```
( - 6*x**(1/3)*cos(x**(1/3))*sin(x**(1/3))**2 + 42*x**(1/3)*cos(x**(1/3))
- 9*x**(2/3)*sin(x**(1/3))**3 + 27*x**(2/3)*sin(x**(1/3)) + 2*sin(x**(1/3))
)**3 - 42*sin(x**(1/3)))/9
```

$$3.62 \quad \int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx$$

Optimal result	495
Mathematica [A] (verified)	495
Rubi [A] (verified)	496
Maple [A] (verified)	497
Fricas [A] (verification not implemented)	497
Sympy [A] (verification not implemented)	498
Maxima [A] (verification not implemented)	498
Giac [A] (verification not implemented)	498
Mupad [B] (verification not implemented)	499
Reduce [B] (verification not implemented)	499

Optimal result

Integrand size = 12, antiderivative size = 8

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin(\sqrt[6]{x})$$

output `6*sin(x^(1/6))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin(\sqrt[6]{x})$$

input `Integrate[Cos[x^(1/6)]/x^(5/6),x]`

output `6*Sin[x^(1/6)]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3861, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx \\ & \quad \downarrow \text{3861} \\ & 6 \int \cos(\sqrt[6]{x}) d\sqrt[6]{x} \\ & \quad \downarrow \text{3042} \\ & 6 \int \sin\left(\sqrt[6]{x} + \frac{\pi}{2}\right) d\sqrt[6]{x} \\ & \quad \downarrow \text{3117} \\ & 6 \sin(\sqrt[6]{x}) \end{aligned}$$

input `Int[Cos[x^(1/6)]/x^(5/6),x]`

output `6*Sin[x^(1/6)]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
  := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$6 \sin\left(x^{\frac{1}{6}}\right)$	7
default	$6 \sin\left(x^{\frac{1}{6}}\right)$	7
meijerg	$6 \sin\left(x^{\frac{1}{6}}\right)$	7

input

```
int(cos(x^(1/6))/x^(5/6),x,method=_RETURNVERBOSE)
```

output

```
6*sin(x^(1/6))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos\left(\sqrt[6]{x}\right)}{x^{5/6}} dx = 6 \sin\left(x^{\frac{1}{6}}\right)$$

input

```
integrate(cos(x^(1/6))/x^(5/6),x, algorithm="fricas")
```

output

```
6*sin(x^(1/6))
```

Sympy [A] (verification not implemented)

Time = 4.95 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin(\sqrt[6]{x})$$

input `integrate(cos(x**(1/6))/x**(5/6),x)`

output `6*sin(x**(1/6))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin(x^{1/6})$$

input `integrate(cos(x^(1/6))/x^(5/6),x, algorithm="maxima")`

output `6*sin(x^(1/6))`

Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin(x^{1/6})$$

input `integrate(cos(x^(1/6))/x^(5/6),x, algorithm="giac")`

output `6*sin(x^(1/6))`

Mupad [B] (verification not implemented)

Time = 41.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin(x^{1/6})$$

input `int(cos(x^(1/6))/x^(5/6),x)`

output `6*sin(x^(1/6))`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{\cos(\sqrt[6]{x})}{x^{5/6}} dx = 6 \sin\left(x^{\frac{1}{6}}\right)$$

input `int(cos(x^(1/6))/x^(5/6),x)`

output `6*sin(x**(1/6))`

3.63 $\int (ex)^m (b \cos(c + dx^n))^p dx$

Optimal result	500
Mathematica [N/A]	500
Rubi [N/A]	501
Maple [N/A]	501
Fricas [N/A]	502
Sympy [N/A]	502
Maxima [N/A]	503
Giac [N/A]	503
Mupad [N/A]	503
Reduce [N/A]	504

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \text{Int}((ex)^m (b \cos(c + dx^n))^p, x)$$

output `Defer(Int)((e*x)^m*(b*cos(c+d*x^n))^p,x)`

Mathematica [N/A]

Not integrable

Time = 1.61 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(c + dx^n))^p dx$$

input `Integrate[(e*x)^m*(b*Cos[c + d*x^n])^p,x]`

output `Integrate[(e*x)^m*(b*Cos[c + d*x^n])^p, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3909}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^m (b \cos (c + dx^n))^p dx$$

↓ 3909

$$\int (ex)^m (b \cos (c + dx^n))^p dx$$

input `Int[(e*x)^m*(b*Cos[c + d*x^n])^p,x]`

output `$Aborted`

Defintions of rubi rules used

rule 3909 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Unintegrable[(e*x)^m*(a + b*Cos[c + d*x^n])^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x]`

Maple [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (ex)^m (b \cos (c + dx^n))^p dx$$

input `int((e*x)^m*(b*cos(c+d*x^n))^p,x)`

output `int((e*x)^m*(b*cos(c+d*x^n))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^m*(b*cos(c+d*x^n))^p,x, algorithm="fricas")`

output `integral((e*x)^m*(b*cos(d*x^n + c))^p, x)`

Sympy [N/A]

Not integrable

Time = 7.63 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (b \cos(c + dx^n))^p (ex)^m dx$$

input `integrate((e*x)**m*(b*cos(c+d*x**n))**p,x)`

output `Integral((b*cos(c + d*x**n))**p*(e*x)**m, x)`

Maxima [N/A]

Not integrable

Time = 0.99 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^m*(b*cos(c+d*x^n))^p,x, algorithm="maxima")`

output `integrate((e*x)^m*(b*cos(d*x^n + c))^p, x)`

Giac [N/A]

Not integrable

Time = 1.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^m*(b*cos(c+d*x^n))^p,x, algorithm="giac")`

output `integrate((e*x)^m*(b*cos(d*x^n + c))^p, x)`

Mupad [N/A]

Not integrable

Time = 41.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (ex)^m (b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(c + dx^n))^p dx$$

input `int((e*x)^m*(b*cos(c + d*x^n))^p,x)`

output `int((e*x)^m*(b*cos(c + d*x^n))^p, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int (ex)^m (b \cos(c + dx^n))^p dx = e^m b^p \left(\int x^m \cos(x^n d + c)^p dx \right)$$

input `int((e*x)^m*(b*cos(c+d*x^n))^p,x)`

output `e**m*b**p*int(x**m*cos(x**n*d + c)**p,x)`

3.64 $\int (ex)^m (a + b \cos(c + dx^n))^p dx$

Optimal result	505
Mathematica [N/A]	505
Rubi [N/A]	506
Maple [N/A]	506
Fricas [N/A]	507
Sympy [N/A]	507
Maxima [N/A]	508
Giac [N/A]	508
Mupad [N/A]	508
Reduce [N/A]	509

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \text{Int}((ex)^m (a + b \cos(c + dx^n))^p, x)$$

output `Defer(Int)((e*x)^m*(a+b*cos(c+d*x^n))^p,x)`

Mathematica [N/A]

Not integrable

Time = 1.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (a + b \cos(c + dx^n))^p dx$$

input `Integrate[(e*x)^m*(a + b*Cos[c + d*x^n])^p,x]`

output `Integrate[(e*x)^m*(a + b*Cos[c + d*x^n])^p, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3909}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx$$

↓ 3909

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx$$

input `Int[(e*x)^m*(a + b*Cos[c + d*x^n])^p,x]`

output `$Aborted`

Defintions of rubi rules used

rule 3909 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x]`

Maple [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx$$

input `int((e*x)^m*(a+b*cos(c+d*x^n))^p,x)`

output `int((e*x)^m*(a+b*cos(c+d*x^n))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^m*(a+b*cos(c+d*x^n))^p,x, algorithm="fricas")`

output `integral((e*x)^m*(b*cos(d*x^n + c) + a)^p, x)`

Sympy [N/A]

Not integrable

Time = 24.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (a + b \cos(c + dx^n))^p dx$$

input `integrate((e*x)**m*(a+b*cos(c+d*x**n))**p,x)`

output `Integral((e*x)**m*(a + b*cos(c + d*x**n))**p, x)`

Maxima [N/A]

Not integrable

Time = 1.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^m*(a+b*cos(c+d*x^n))^p,x, algorithm="maxima")`

output `integrate((e*x)^m*(b*cos(d*x^n + c) + a)^p, x)`

Giac [N/A]

Not integrable

Time = 4.85 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^m*(a+b*cos(c+d*x^n))^p,x, algorithm="giac")`

output `integrate((e*x)^m*(b*cos(d*x^n + c) + a)^p, x)`

Mupad [N/A]

Not integrable

Time = 41.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = \int (ex)^m (a + b \cos(c + dx^n))^p dx$$

input `int((e*x)^m*(a + b*cos(c + d*x^n))^p,x)`

output `int((e*x)^m*(a + b*cos(c + d*x^n))^p, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int (ex)^m (a + b \cos(c + dx^n))^p dx = e^m \left(\int x^m (\cos(x^n d + c) b + a)^p dx \right)$$

input `int((e*x)^m*(a+b*cos(c+d*x^n))^p,x)`

output `e**m*int(x**m*(cos(x**n*d + c)*b + a)**p,x)`

3.65 $\int (ex)^{-1+n} (b \cos (c + dx^n))^p dx$

Optimal result	510
Mathematica [A] (verified)	510
Rubi [A] (verified)	511
Maple [F]	512
Fricas [F]	513
Sympy [F]	513
Maxima [F]	513
Giac [F]	514
Mupad [F(-1)]	514
Reduce [F]	514

Optimal result

Integrand size = 20, antiderivative size = 93

$$\int (ex)^{-1+n} (b \cos (c + dx^n))^p dx = \frac{x^{-n}(ex)^n (b \cos (c + dx^n))^{1+p} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+p}{2}, \frac{3+p}{2}, \cos^2(c + dx^n)\right) \sin(c + dx^n)}{bden(1+p)\sqrt{\sin^2(c + dx^n)}}$$

output `-(e*x)^n*(b*cos(c+d*x^n))^(p+1)*hypergeom([1/2, 1/2*p+1/2],[3/2+1/2*p],cos(c+d*x^n)^2)*sin(c+d*x^n)/b/d/e/n/(p+1)/(x^n)/(sin(c+d*x^n)^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.96

$$\int (ex)^{-1+n} (b \cos (c + dx^n))^p dx = \frac{x^{1-n}(ex)^{-1+n} (b \cos (c + dx^n))^p \cot (c + dx^n) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+p}{2}, \frac{3+p}{2}, \cos^2(c + dx^n)\right) \sqrt{\sin^2(c + dx^n)}}{dn(1+p)}$$

input `Integrate[(e*x)^(-1 + n)*(b*Cos[c + d*x^n])^p,x]`

output

```

-((x^(1 - n)*(e*x)^(-1 + n)*(b*cos[c + d*x^n])^p*cot[c + d*x^n]*Hypergeome
tric2F1[1/2, (1 + p)/2, (3 + p)/2, Cos[c + d*x^n]^2]*Sqrt[Sin[c + d*x^n]^2
])/ (d*n*(1 + p)))

```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3863, 3861, 3042, 3122}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (ex)^{n-1} (b \cos(c + dx^n))^p dx \\
 & \quad \downarrow \text{3863} \\
 & \frac{x^{-n}(ex)^n \int x^{n-1} (b \cos(dx^n + c))^p dx}{e} \\
 & \quad \downarrow \text{3861} \\
 & \frac{x^{-n}(ex)^n \int (b \cos(dx^n + c))^p dx^n}{en} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x^{-n}(ex)^n \int (b \sin(dx^n + c + \frac{\pi}{2}))^p dx^n}{en} \\
 & \quad \downarrow \text{3122} \\
 & \frac{x^{-n}(ex)^n \sin(c + dx^n) (b \cos(c + dx^n))^{p+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{p+1}{2}, \frac{p+3}{2}, \cos^2(dx^n + c)\right)}{bden(p+1)\sqrt{\sin^2(c + dx^n)}}
 \end{aligned}$$

input

```

Int[(e*x)^(-1 + n)*(b*cos[c + d*x^n])^p,x]

```


output $-\left(\left(e^x\right)^n \cdot \left(b \cos \left(c+d x^n\right)\right)^{\left(1+p\right)} \cdot \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{\left(1+p\right)}{2}, \frac{\left(3+p\right)}{2}, \cos \left[c+d x^n\right]^2\right] \cdot \sin \left[c+d x^n\right]\right) / \left(b \cdot d \cdot e^{n \cdot\left(1+p\right)} \cdot x^n \cdot \sqrt{\sin \left[c+d x^n\right]^2}\right)$

Definitions of rubi rules used

rule 3042 $\operatorname{Int}\left[u, x_{\text{Symbol}}\right] \rightarrow \operatorname{Int}\left[\operatorname{DeactivateTrig}\left[u, x\right], x\right] / ; \operatorname{FunctionOfTrigOfLinearQ}\left[u, x\right]$

rule 3122 $\operatorname{Int}\left[\left(\left(b_{\cdot}\right) \cdot \sin \left[\left(c_{\cdot}\right)+\left(d_{\cdot}\right) \cdot\left(x_{\cdot}\right)\right]\right)^{\left(n_{\cdot}\right)}, x_{\text{Symbol}}\right] \rightarrow \operatorname{Simp}\left[\cos \left[c+d x\right] \cdot\left(b \cdot \sin \left[c+d x\right]\right)^{\left(n+1\right)} / \left(b \cdot d \cdot\left(n+1\right) \cdot \sqrt{\cos \left[c+d x\right]^2}\right) \cdot \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{\left(n+1\right)}{2}, \frac{\left(n+3\right)}{2}, \sin \left[c+d x\right]^2\right], x\right] / ; \operatorname{FreeQ}\left[\left\{b, c, d, n\right\}, x\right] \&\& \text{!IntegerQ}\left[2 \cdot n\right]$

rule 3861 $\operatorname{Int}\left[\left(\left(a_{\cdot}\right)+\cos \left[\left(c_{\cdot}\right)+\left(d_{\cdot}\right) \cdot\left(x_{\cdot}\right)^{\left(n_{\cdot}\right)}\right]\right) \cdot\left(b_{\cdot}\right)^{\left(p_{\cdot}\right)} \cdot\left(x_{\cdot}\right)^{\left(m_{\cdot}\right)}, x_{\text{Symbol}}\right] \rightarrow \operatorname{Simp}\left[\frac{1}{n} \operatorname{Subst}\left[\operatorname{Int}\left[x^{\left(\operatorname{Simplify}\left[\frac{m+1}{n}-1\right]\right) \cdot\left(a+b \cdot \cos \left[c+d x\right]\right)^p}, x, x^n\right], x\right] / ; \operatorname{FreeQ}\left[\left\{a, b, c, d, m, n, p\right\}, x\right] \&\& \operatorname{IntegerQ}\left[\operatorname{Simplify}\left[\frac{m+1}{n}\right]\right] \&\& \left(\operatorname{EqQ}\left[p, 1\right] \mid \mid \operatorname{EqQ}\left[m, n-1\right] \mid \mid \left(\operatorname{IntegerQ}\left[p\right] \&\& \operatorname{GtQ}\left[\operatorname{Simplify}\left[\frac{m+1}{n}\right], 0\right]\right)\right)$

rule 3863 $\operatorname{Int}\left[\left(\left(a_{\cdot}\right)+\cos \left[\left(c_{\cdot}\right)+\left(d_{\cdot}\right) \cdot\left(x_{\cdot}\right)^{\left(n_{\cdot}\right)}\right]\right) \cdot\left(b_{\cdot}\right)^{\left(p_{\cdot}\right)} \cdot\left(e_{\cdot}\right) \cdot\left(x_{\cdot}\right)^{\left(m_{\cdot}\right)}, x_{\text{Symbol}}\right] \rightarrow \operatorname{Simp}\left[e^{\operatorname{IntPart}\left[m\right]} \cdot\left(e^x\right)^{\operatorname{FracPart}\left[m\right]} / x^{\operatorname{FracPart}\left[m\right]} \operatorname{Int}\left[x^m \cdot\left(a+b \cdot \cos \left[c+d x^n\right]\right)^p, x\right], x\right] / ; \operatorname{FreeQ}\left[\left\{a, b, c, d, e, m, n, p\right\}, x\right] \&\& \operatorname{IntegerQ}\left[\operatorname{Simplify}\left[\frac{m+1}{n}\right]\right]$

Maple [F]

$$\int (ex)^{-1+n} (b \cos (c+dx^n))^p dx$$

input $\operatorname{int}\left(\left(e^x\right)^{-1+n} \cdot\left(b \cdot \cos \left(c+d x^n\right)\right)^p, x\right)$

output $\operatorname{int}\left(\left(e^x\right)^{-1+n} \cdot\left(b \cdot \cos \left(c+d x^n\right)\right)^p, x\right)$

Fricas [F]

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^(-1+n)*(b*cos(c+d*x^n))^p,x, algorithm="fricas")`

output `integral((e*x)^(n - 1)*(b*cos(d*x^n + c))^p, x)`

Sympy [F]

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx = \int (b \cos(c + dx^n))^p (ex)^{n-1} dx$$

input `integrate((e*x)**(-1+n)*(b*cos(c+d*x**n))**p,x)`

output `Integral((b*cos(c + d*x**n))**p*(e*x)**(n - 1), x)`

Maxima [F]

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^(-1+n)*(b*cos(c+d*x^n))^p,x, algorithm="maxima")`

output `integrate((e*x)^(n - 1)*(b*cos(d*x^n + c))^p, x)`

Giac [F]

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^(-1+n)*(b*cos(c+d*x^n))^p,x, algorithm="giac")`

output `integrate((e*x)^(n - 1)*(b*cos(d*x^n + c))^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(c + dx^n))^p dx$$

input `int((e*x)^(n - 1)*(b*cos(c + d*x^n))^p,x)`

output `int((e*x)^(n - 1)*(b*cos(c + d*x^n))^p, x)`

Reduce [F]

$$\int (ex)^{-1+n} (b \cos(c + dx^n))^p dx = \frac{e^n b^p \left(\int \frac{x^n \cos(x^n d + c)^p}{x} dx \right)}{e}$$

input `int((e*x)^(-1+n)*(b*cos(c+d*x^n))^p,x)`

output `(e**n*b**p*int((x**n*cos(x**n*d + c)**p)/x,x))/e`

3.66 $\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$

Optimal result	515
Mathematica [N/A]	515
Rubi [N/A]	516
Maple [N/A]	517
Fricas [N/A]	517
Sympy [N/A]	517
Maxima [N/A]	518
Giac [N/A]	518
Mupad [N/A]	518
Reduce [N/A]	519

Optimal result

Integrand size = 22, antiderivative size = 22

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \frac{x^{-2n}(ex)^{2n} \text{Int}(x^{-1+2n}(b \cos(c + dx^n))^p, x)}{e}$$

output

```
(e*x)^(2*n)*Defer(Int)(x^(-1+2*n)*(b*cos(c+d*x^n))^p,x)/e/(x^(2*n))
```

Mathematica [N/A]

Not integrable

Time = 1.88 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$$

input

```
Integrate[(e*x)^(-1 + 2*n)*(b*Cos[c + d*x^n])^p,x]
```

output

```
Integrate[(e*x)^(-1 + 2*n)*(b*Cos[c + d*x^n])^p, x]
```

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3863, 3909}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^{2n-1} (b \cos(c + dx^n))^p dx$$

$$\downarrow \text{3863}$$

$$\frac{x^{-2n}(ex)^{2n} \int x^{2n-1} (b \cos(dx^n + c))^p dx}{e}$$

$$\downarrow \text{3909}$$

$$\frac{x^{-2n}(ex)^{2n} \int x^{2n-1} (b \cos(dx^n + c))^p dx}{e}$$

input `Int[(e*x)^(-1 + 2*n)*(b*Cos[c + d*x^n])^p,x]`

output `$Aborted`

Defintions of rubi rules used

rule 3863 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Simp[e^IntPart[m]*((e*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 3909 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Unintegrable[(e*x)^m*(a + b*Cos[c + d*x^n])^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x]`

Maple [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx$$

input `int((e*x)^(-1+2*n)*(b*cos(c+d*x^n))^p,x)`

output `int((e*x)^(-1+2*n)*(b*cos(c+d*x^n))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^(-1+2*n)*(b*cos(c+d*x^n))^p,x, algorithm="fricas")`

output `integral((e*x)^(2*n - 1)*(b*cos(d*x^n + c))^p, x)`

Sympy [N/A]

Not integrable

Time = 7.54 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \int (b \cos(c + dx^n))^p (ex)^{2n-1} dx$$

input `integrate((e*x)**(-1+2*n)*(b*cos(c+d*x**n))**p,x)`

output `Integral((b*cos(c + d*x**n))**p*(e*x)**(2*n - 1), x)`

Maxima [N/A]

Not integrable

Time = 1.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^(-1+2*n)*(b*cos(c+d*x^n))^p,x, algorithm="maxima")`

output `integrate((e*x)^(2*n - 1)*(b*cos(d*x^n + c))^p, x)`

Giac [N/A]

Not integrable

Time = 1.04 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(dx^n + c))^p dx$$

input `integrate((e*x)^(-1+2*n)*(b*cos(c+d*x^n))^p,x, algorithm="giac")`

output `integrate((e*x)^(2*n - 1)*(b*cos(d*x^n + c))^p, x)`

Mupad [N/A]

Not integrable

Time = 40.77 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(c + dx^n))^p dx$$

input `int((e*x)^(2*n - 1)*(b*cos(c + d*x^n))^p,x)`

output `int((e*x)^(2*n - 1)*(b*cos(c + d*x^n))^p, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.50

$$\int (ex)^{-1+2n} (b \cos(c + dx^n))^p dx = \frac{e^{2n} b^p \left(\int \frac{x^{2n} \cos(x^n d + c)^p}{x} dx \right)}{e}$$

input `int((e*x)^(-1+2*n)*(b*cos(c+d*x^n))^p,x)`

output `(e**(2*n)*b**p*int((x**(2*n)*cos(x**n*d + c)**p)/x,x))/e`

3.67 $\int (ex)^{-1+n} (a + b \cos (c + dx^n))^p dx$

Optimal result	520
Mathematica [A] (warning: unable to verify)	520
Rubi [A] (verified)	521
Maple [F]	523
Fricas [F]	523
Sympy [F]	524
Maxima [F]	524
Giac [F]	524
Mupad [F(-1)]	525
Reduce [F]	525

Optimal result

Integrand size = 22, antiderivative size = 131

$$\int (ex)^{-1+n} (a + b \cos (c + dx^n))^p dx$$

$$= \frac{\sqrt{2}x^{-n}(ex)^n \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -p, \frac{3}{2}, \frac{1}{2}(1 - \cos(c + dx^n)), \frac{b(1 - \cos(c + dx^n))}{a+b}\right) (a + b \cos(c + dx^n))^p \left(\frac{a+b \cos(c + dx^n)}{a+b}\right)}{\operatorname{den} \sqrt{1 + \cos(c + dx^n)}}$$

output

$$2^{(1/2)}*(e*x)^n*\operatorname{AppellF1}(1/2, -p, 1/2, 3/2, b*(1 - \cos(c+d*x^n))/(a+b), 1/2 - 1/2*\cos(c+d*x^n))*(a+b*\cos(c+d*x^n))^p*\sin(c+d*x^n)/d/e/n/(x^n)/(1 + \cos(c+d*x^n))^{(1/2)}/(((a+b*\cos(c+d*x^n))/(a+b))^p)$$

Mathematica [A] (warning: unable to verify)

Time = 0.83 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.14

$$\int (ex)^{-1+n} (a + b \cos (c + dx^n))^p dx =$$

$$\frac{x^{-n}(ex)^n \operatorname{AppellF1}\left(1 + p, \frac{1}{2}, \frac{1}{2}, 2 + p, \frac{a+b \cos(c+dx^n)}{a-b}, \frac{a+b \cos(c+dx^n)}{a+b}\right) \sqrt{-\frac{b(-1+\cos(c+dx^n))}{a+b}} \sqrt{\frac{b(1+\cos(c+dx^n))}{-a+b}}}{b \operatorname{den}(1 + p)}$$

input

$$\operatorname{Integrate}[(e*x)^{-1 + n}*(a + b*\operatorname{Cos}[c + d*x^n])^p, x]$$

output

```

-(((e*x)^n*AppellF1[1 + p, 1/2, 1/2, 2 + p, (a + b*Cos[c + d*x^n])/(a - b)
, (a + b*Cos[c + d*x^n])/(a + b)]*Sqrt[-((b*(-1 + Cos[c + d*x^n]))/(a + b)
)]*Sqrt[(b*(1 + Cos[c + d*x^n]))/(-a + b)]*(a + b*Cos[c + d*x^n])^(1 + p)*
Csc[c + d*x^n])/(b*d*e*n*(1 + p)*x^n)

```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3863, 3861, 3042, 3144, 156, 155}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int (ex)^{n-1} (a + b \cos(c + dx^n))^p dx \\
& \quad \downarrow \text{3863} \\
& \frac{x^{-n}(ex)^n \int x^{n-1}(a + b \cos(dx^n + c))^p dx}{e} \\
& \quad \downarrow \text{3861} \\
& \frac{x^{-n}(ex)^n \int (a + b \cos(dx^n + c))^p dx^n}{en} \\
& \quad \downarrow \text{3042} \\
& \frac{x^{-n}(ex)^n \int (a + b \sin(dx^n + c + \frac{\pi}{2}))^p dx^n}{en} \\
& \quad \downarrow \text{3144} \\
& \frac{x^{-n}(ex)^n \sin(c + dx^n) \int \frac{(a+b \cos(dx^n+c))^p}{\sqrt{1-\cos(dx^n+c)}\sqrt{\cos(dx^n+c)+1}} d \cos(dx^n + c)}{den \sqrt{1 - \cos(c + dx^n)} \sqrt{\cos(c + dx^n) + 1}} \\
& \quad \downarrow \text{156} \\
& \frac{x^{-n}(ex)^n \sin(c + dx^n) (a + b \cos(c + dx^n))^p \left(\frac{a+b \cos(c+dx^n)}{a+b}\right)^{-p} \int \frac{\left(\frac{a}{a+b} + \frac{b \cos(dx^n+c)}{a+b}\right)^p}{\sqrt{1-\cos(dx^n+c)}\sqrt{\cos(dx^n+c)+1}} d \cos(dx^n + c)}{den \sqrt{1 - \cos(c + dx^n)} \sqrt{\cos(c + dx^n) + 1}} \\
& \quad \downarrow \text{155}
\end{aligned}$$

$$\frac{\sqrt{2}x^{-n}(ex)^n \sin(c + dx^n) (a + b \cos(c + dx^n))^p \left(\frac{a+b \cos(c+dx^n)}{a+b}\right)^{-p} \text{AppellF1}\left(\frac{1}{2}, \frac{1}{2}, -p, \frac{3}{2}, \frac{1}{2}(1 - \cos(dx^n + c))\right)}{\text{den} \sqrt{\cos(c + dx^n) + 1}}$$

input `Int[(e*x)^(-1 + n)*(a + b*Cos[c + d*x^n])^p,x]`

output `(Sqrt[2]*(e*x)^n*AppellF1[1/2, 1/2, -p, 3/2, (1 - Cos[c + d*x^n])/2, (b*(1 - Cos[c + d*x^n]))/(a + b)]*(a + b*Cos[c + d*x^n])^p*Sin[c + d*x^n])/(d*e*n*x^n*Sqrt[1 + Cos[c + d*x^n]]*((a + b*Cos[c + d*x^n])/(a + b))^p)`

Defintions of rubi rules used

rule 155 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*Simplify[b/(b*c - a*d)]^n*Simplify[b/(b*e - a*f)]^p))*AppellF1[m + 1, -n, -p, m + 2, (-d)*((a + b*x)/(b*c - a*d)), (-f)*((a + b*x)/(b*e - a*f))], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && GtQ[Simplify[b/(b*e - a*f)], 0] && !(GtQ[Simplify[d/(d*a - c*b)], 0] && GtQ[Simplify[d/(d*e - c*f)], 0] && SimplerQ[c + d*x, a + b*x]) && !(GtQ[Simplify[f/(f*a - e*b)], 0] && GtQ[Simplify[f/(f*c - e*d)], 0] && SimplerQ[e + f*x, a + b*x])`

rule 156 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(e + f*x)^FracPart[p]/(Simplify[b/(b*e - a*f)]^IntPart[p])*((b*(e + f*x)/(b*e - a*f))^FracPart[p]) Int[(a + b*x)^m*(c + d*x)^n*Simp[b*(e/(b*e - a*f)) + b*f*(x/(b*e - a*f)), x]^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] && GtQ[Simplify[b/(b*c - a*d)], 0] && !GtQ[Simplify[b/(b*e - a*f)], 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3144 `Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]/(d*Sqrt[1 + Sin[c + d*x]]*Sqrt[1 - Sin[c + d*x]]) Subst[Int[(a + b*x)^n/(Sqrt[1 + x]*Sqrt[1 - x]), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0] && !IntegerQ[2*n]`

rule 3861 `Int[((a_) + Cos[(c_) + (d_)*(x_)^(n_)])*(b_)^(p_)*(x_)^(m_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

rule 3863 `Int[((a_) + Cos[(c_) + (d_)*(x_)^(n_)])*(b_)^(p_)*((e_)*(x_)^(m_)), x_Symbol] := Simp[e^IntPart[m]*((e*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Maple [F]

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx$$

input `int((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x)`

output `int((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x)`

Fricas [F]

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x, algorithm="fricas")`

output `integral((e*x)^(n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

Sympy [F]

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (a + b \cos(c + dx^n))^p dx$$

input `integrate((e*x)**(-1+n)*(a+b*cos(c+d*x**n))**p,x)`

output `Integral((e*x)**(n - 1)*(a + b*cos(c + d*x**n))**p, x)`

Maxima [F]

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x, algorithm="maxima")`

output `integrate((e*x)^(n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

Giac [F]

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x, algorithm="giac")`

output `integrate((e*x)^(n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

Mupad [F(-1)]

Timed out.

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{n-1} (a + b \cos(c + dx^n))^p dx$$

input `int((e*x)^(n - 1)*(a + b*cos(c + d*x^n))^p,x)`

output `int((e*x)^(n - 1)*(a + b*cos(c + d*x^n))^p, x)`

Reduce [F]

$$\int (ex)^{-1+n} (a + b \cos(c + dx^n))^p dx = \frac{e^n \left(\int \frac{x^n (\cos(x^n d + c) b + a)^p}{x} dx \right)}{e}$$

input `int((e*x)^(-1+n)*(a+b*cos(c+d*x^n))^p,x)`

output `(e**n*int((x**n*(cos(x**n*d + c)*b + a)**p)/x,x))/e`

3.68 $\int (ex)^{-1+2n} (a + b \cos (c + dx^n))^p dx$

Optimal result	526
Mathematica [N/A]	526
Rubi [N/A]	527
Maple [N/A]	528
Fricas [N/A]	528
Sympy [N/A]	528
Maxima [N/A]	529
Giac [N/A]	529
Mupad [N/A]	529
Reduce [N/A]	530

Optimal result

Integrand size = 24, antiderivative size = 24

$$\int (ex)^{-1+2n} (a + b \cos (c + dx^n))^p dx = \frac{x^{-2n}(ex)^{2n} \text{Int}(x^{-1+2n}(a + b \cos (c + dx^n))^p, x)}{e}$$

output `(e*x)^(2*n)*Defer(Int)(x^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x)/e/(x^(2*n))`

Mathematica [N/A]

Not integrable

Time = 1.96 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int (ex)^{-1+2n} (a + b \cos (c + dx^n))^p dx = \int (ex)^{-1+2n} (a + b \cos (c + dx^n))^p dx$$

input `Integrate[(e*x)^(-1 + 2*n)*(a + b*Cos[c + d*x^n])^p,x]`

output `Integrate[(e*x)^(-1 + 2*n)*(a + b*Cos[c + d*x^n])^p, x]`

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3863, 3909}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (ex)^{2n-1} (a + b \cos(c + dx^n))^p dx$$

$$\downarrow \text{3863}$$

$$\frac{x^{-2n}(ex)^{2n} \int x^{2n-1}(a + b \cos(dx^n + c))^p dx}{e}$$

$$\downarrow \text{3909}$$

$$\frac{x^{-2n}(ex)^{2n} \int x^{2n-1}(a + b \cos(dx^n + c))^p dx}{e}$$

input `Int[(e*x)^(-1 + 2*n)*(a + b*Cos[c + d*x^n])^p,x]`

output `$Aborted`

Defintions of rubi rules used

rule 3863 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Simp[e^IntPart[m]*((e*x)^FracPart[m]/x^FracPart[m]) Int[x^m*(a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 3909 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_.)*((e_.)*(x_)^(m_.), x_Symbol] := Unintegrable[(e*x)^m*(a + b*Cos[c + d*x^n])^p, x] /; FreeQ[{a, b, c, d, e, m, n, p}, x]`

Maple [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx$$

input `int((e*x)^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x)`

output `int((e*x)^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x, algorithm="fricas")`

output `integral((e*x)^(2*n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

Sympy [N/A]

Not integrable

Time = 23.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (a + b \cos(c + dx^n))^p dx$$

input `integrate((e*x)**(-1+2*n)*(a+b*cos(c+d*x**n))**p,x)`

output `Integral((e*x)**(2*n - 1)*(a + b*cos(c + d*x**n))**p, x)`

Maxima [N/A]

Not integrable

Time = 1.54 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x, algorithm="maxima")`

output `integrate((e*x)^(2*n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

Giac [N/A]

Not integrable

Time = 4.63 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (b \cos(dx^n + c) + a)^p dx$$

input `integrate((e*x)^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x, algorithm="giac")`

output `integrate((e*x)^(2*n - 1)*(b*cos(d*x^n + c) + a)^p, x)`

Mupad [N/A]

Not integrable

Time = 40.57 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \int (ex)^{2n-1} (a + b \cos(c + dx^n))^p dx$$

input `int((e*x)^(2*n - 1)*(a + b*cos(c + d*x^n))^p,x)`

output `int((e*x)^(2*n - 1)*(a + b*cos(c + d*x^n))^p, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.42

$$\int (ex)^{-1+2n} (a + b \cos(c + dx^n))^p dx = \frac{e^{2n} \left(\int \frac{x^{2n} (\cos(x^n d + c) b + a)^p dx}{x} \right)}{e}$$

input `int((e*x)^(-1+2*n)*(a+b*cos(c+d*x^n))^p,x)`

output `(e**(2*n)*int((x**(2*n)*(cos(x**n*d + c)*b + a)**p)/x,x))/e`

3.69 $\int \frac{\cos(a+bx^n)}{x} dx$

Optimal result	531
Mathematica [A] (verified)	531
Rubi [A] (verified)	532
Maple [A] (verified)	533
Fricas [A] (verification not implemented)	533
Sympy [F]	534
Maxima [C] (verification not implemented)	534
Giac [F]	535
Mupad [F(-1)]	535
Reduce [F]	535

Optimal result

Integrand size = 12, antiderivative size = 26

$$\int \frac{\cos(a+bx^n)}{x} dx = \frac{\cos(a) \operatorname{CosIntegral}(bx^n)}{n} - \frac{\sin(a) \operatorname{Si}(bx^n)}{n}$$

output

```
cos(a)*Ci(b*x^n)/n-sin(a)*Si(b*x^n)/n
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{\cos(a+bx^n)}{x} dx = \frac{\cos(a) \operatorname{CosIntegral}(bx^n) - \sin(a) \operatorname{Si}(bx^n)}{n}$$

input

```
Integrate[Cos[a + b*x^n]/x,x]
```

output

```
(Cos[a]*CosIntegral[b*x^n] - Sin[a]*SinIntegral[b*x^n])/n
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3859, 3856, 3857}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos(a + bx^n)}{x} dx \\ & \quad \downarrow \text{3859} \\ & \cos(a) \int \frac{\cos(bx^n)}{x} dx - \sin(a) \int \frac{\sin(bx^n)}{x} dx \\ & \quad \downarrow \text{3856} \\ & \cos(a) \int \frac{\cos(bx^n)}{x} dx - \frac{\sin(a)\text{Si}(bx^n)}{n} \\ & \quad \downarrow \text{3857} \\ & \frac{\cos(a) \text{CosIntegral}(bx^n)}{n} - \frac{\sin(a)\text{Si}(bx^n)}{n} \end{aligned}$$

input `Int[Cos[a + b*x^n]/x, x]`

output `(Cos[a]*CosIntegral[b*x^n])/n - (Sin[a]*SinIntegral[b*x^n])/n`

Defintions of rubi rules used

rule 3856 `Int[Sin[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[SinIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3857 `Int[Cos[(d_.)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[CosIntegral[d*x^n]/n, x] / ; FreeQ[{d, n}, x]`

rule 3859

```
Int[Cos[(c_) + (d_)*(x_)^(n_)]/(x_), x_Symbol] := Simp[Cos[c] Int[Cos[d*x^n]/x, x], x] - Simp[Sin[c] Int[Sin[d*x^n]/x, x], x] /; FreeQ[{c, d, n}, x]
```

Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$\frac{-\text{Si}(bx^n)\sin(a) + \text{Ci}(bx^n)\cos(a)}{n}$	25
default	$\frac{-\text{Si}(bx^n)\sin(a) + \text{Ci}(bx^n)\cos(a)}{n}$	25
risch	$\frac{ie^{-ia}\pi \text{csgn}(bx^n)}{2n} - \frac{ie^{-ia}\text{Si}(bx^n)}{n} - \frac{e^{-ia}\text{expIntegral}_1(-ibx^n)}{2n} - \frac{e^{ia}\text{expIntegral}_1(-ibx^n)}{2n}$	75
meijerg	$\frac{\sqrt{\pi} \left(\frac{2\gamma + 2n \ln(x) + \ln(b^2)}{\sqrt{\pi}} - \frac{2\gamma}{\sqrt{\pi}} - \frac{2 \ln(2)}{\sqrt{\pi}} - \frac{2 \ln\left(\frac{bx^n}{2}\right)}{\sqrt{\pi}} + \frac{2 \text{Ci}(bx^n)}{\sqrt{\pi}} \right) \cos(a)}{2n} - \frac{\sin(a) \text{Si}(bx^n)}{n}$	79

input

```
int(cos(a+b*x^n)/x,x,method=_RETURNVERBOSE)
```

output

```
1/n*(-Si(b*x^n)*sin(a)+Ci(b*x^n)*cos(a))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{\cos(a + bx^n)}{x} dx = \frac{\cos(a) \text{Ci}(bx^n) - \sin(a) \text{Si}(bx^n)}{n}$$

input

```
integrate(cos(a+b*x^n)/x,x, algorithm="fricas")
```

output

```
(cos(a)*cos_integral(b*x^n) - sin(a)*sin_integral(b*x^n))/n
```

Sympy [F]

$$\int \frac{\cos(a + bx^n)}{x} dx = \int \frac{\cos(a + bx^n)}{x} dx$$

input `integrate(cos(a+b*x**n)/x,x)`

output `Integral(cos(a + b*x**n)/x, x)`

Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.17 (sec) , antiderivative size = 90, normalized size of antiderivative = 3.46

$$\int \frac{\cos(a + bx^n)}{x} dx = \frac{\left(\operatorname{Ei}(i bx^n) + \operatorname{Ei}(-i bx^n) + \operatorname{Ei}\left(i be^{(n\overline{\log(x)})}\right) + \operatorname{Ei}\left(-i be^{(n\overline{\log(x)})}\right) \right) \cos(a) + \left(i \operatorname{Ei}(i bx^n) - i \operatorname{Ei}(-i bx^n) \right) \sin(a)}{4n}$$

input `integrate(cos(a+b*x^n)/x,x, algorithm="maxima")`

output `1/4*((Ei(I*b*x^n) + Ei(-I*b*x^n) + Ei(I*b*e^(n*conjugate(log(x)))) + Ei(-I*b*e^(n*conjugate(log(x)))))*cos(a) + (I*Ei(I*b*x^n) - I*Ei(-I*b*x^n) + I*Ei(I*b*e^(n*conjugate(log(x)))) - I*Ei(-I*b*e^(n*conjugate(log(x)))))*sin(a))/n`

Giac [F]

$$\int \frac{\cos(a + bx^n)}{x} dx = \int \frac{\cos(bx^n + a)}{x} dx$$

input `integrate(cos(a+b*x^n)/x,x, algorithm="giac")`

output `integrate(cos(b*x^n + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + bx^n)}{x} dx = \int \frac{\cos(a + b x^n)}{x} dx$$

input `int(cos(a + b*x^n)/x,x)`

output `int(cos(a + b*x^n)/x, x)`

Reduce [F]

$$\int \frac{\cos(a + bx^n)}{x} dx = \int \frac{\cos(x^n b + a)}{x} dx$$

input `int(cos(a+b*x^n)/x,x)`

output `int(cos(x**n*b + a)/x,x)`

3.70 $\int \frac{\cos^2(a+bx^n)}{x} dx$

Optimal result	536
Mathematica [A] (verified)	536
Rubi [A] (verified)	537
Maple [A] (verified)	538
Fricas [A] (verification not implemented)	538
Sympy [F]	538
Maxima [C] (verification not implemented)	539
Giac [F]	539
Mupad [F(-1)]	540
Reduce [F]	540

Optimal result

Integrand size = 14, antiderivative size = 43

$$\int \frac{\cos^2(a+bx^n)}{x} dx = \frac{\cos(2a) \operatorname{CosIntegral}(2bx^n)}{2n} + \frac{\log(x)}{2} - \frac{\sin(2a) \operatorname{Si}(2bx^n)}{2n}$$

output `1/2*cos(2*a)*Ci(2*b*x^n)/n+1/2*ln(x)-1/2*sin(2*a)*Si(2*b*x^n)/n`

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

$$\int \frac{\cos^2(a+bx^n)}{x} dx = \frac{\cos(2a) \operatorname{CosIntegral}(2bx^n) + n \log(x) - \sin(2a) \operatorname{Si}(2bx^n)}{2n}$$

input `Integrate[Cos[a + b*x^n]^2/x,x]`

output `(Cos[2*a]*CosIntegral[2*b*x^n] + n*Log[x] - Sin[2*a]*SinIntegral[2*b*x^n])/ (2*n)`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^2(a + bx^n)}{x} dx$$

↓ 3907

$$\int \left(\frac{\cos(2a + 2bx^n)}{2x} + \frac{1}{2x} \right) dx$$

↓ 2009

$$\frac{\cos(2a) \operatorname{CosIntegral}(2bx^n)}{2n} - \frac{\sin(2a) \operatorname{Si}(2bx^n)}{2n} + \frac{\log(x)}{2}$$

input

```
Int[Cos[a + b*x^n]^2/x, x]
```

output

```
(Cos[2*a]*CosIntegral[2*b*x^n])/(2*n) + Log[x]/2 - (Sin[2*a]*SinIntegral[2*b*x^n])/(2*n)
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3907

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]
```

Maple [A] (verified)

Time = 1.10 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

method	result
derivativedivides	$\frac{-\frac{\text{Si}(2bx^n)}{2} \sin(2a) + \frac{\text{Ci}(2bx^n)}{2} \cos(2a) + \frac{\ln(bx^n)}{2}}{n}$
default	$\frac{-\frac{\text{Si}(2bx^n)}{2} \sin(2a) + \frac{\text{Ci}(2bx^n)}{2} \cos(2a) + \frac{\ln(bx^n)}{2}}{n}$
risch	$\frac{\ln(x)}{2} + \frac{ie^{-2ia}\pi \text{csgn}(bx^n)}{4n} - \frac{ie^{-2ia} \text{Si}(2bx^n)}{2n} - \frac{e^{-2ia} \exp\text{Integral}_1(-2ibx^n)}{4n} - \frac{e^{2ia} \exp\text{Integral}_1(-2ibx^n)}{4n}$

input `int(cos(a+b*x^n)^2/x,x,method=_RETURNVERBOSE)`output `1/n*(-1/2*Si(2*b*x^n)*sin(2*a)+1/2*Ci(2*b*x^n)*cos(2*a)+1/2*ln(b*x^n))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.81

$$\int \frac{\cos^2(a + bx^n)}{x} dx = \frac{\cos(2a) \text{Ci}(2bx^n) + n \log(x) - \sin(2a) \text{Si}(2bx^n)}{2n}$$

input `integrate(cos(a+b*x^n)^2/x,x, algorithm="fricas")`output `1/2*(cos(2*a)*cos_integral(2*b*x^n) + n*log(x) - sin(2*a)*sin_integral(2*b*x^n))/n`**Sympy [F]**

$$\int \frac{\cos^2(a + bx^n)}{x} dx = \int \frac{\cos^2(a + bx^n)}{x} dx$$

input `integrate(cos(a+b*x**n)**2/x,x)`

output `Integral(cos(a + b*x**n)**2/x, x)`

Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.27 (sec) , antiderivative size = 99, normalized size of antiderivative = 2.30

$$\int \frac{\cos^2(a + bx^n)}{x} dx$$

$$= \frac{\left(\operatorname{Ei}(2i bx^n) + \operatorname{Ei}(-2i bx^n) + \operatorname{Ei}\left(2i be^{(n\overline{\log(x)})}\right) + \operatorname{Ei}\left(-2i be^{(n\overline{\log(x)})}\right) \right) \cos(2a) + 4n \log(x) + \left(i \operatorname{Ei}(2i bx^n) - i \operatorname{Ei}(-2i bx^n) + i \operatorname{Ei}(2i be^{(n\overline{\log(x)})}) - i \operatorname{Ei}(-2i be^{(n\overline{\log(x)})}) \right) \sin(2a)}{8n}$$

input `integrate(cos(a+b*x^n)^2/x,x, algorithm="maxima")`

output `1/8*((Ei(2*I*b*x^n) + Ei(-2*I*b*x^n) + Ei(2*I*b*e^(n*conjugate(log(x)))) + Ei(-2*I*b*e^(n*conjugate(log(x)))))*cos(2*a) + 4*n*log(x) + (I*Ei(2*I*b*x^n) - I*Ei(-2*I*b*x^n) + I*Ei(2*I*b*e^(n*conjugate(log(x)))) - I*Ei(-2*I*b*e^(n*conjugate(log(x)))))*sin(2*a))/n`

Giac [F]

$$\int \frac{\cos^2(a + bx^n)}{x} dx = \int \frac{\cos(bx^n + a)^2}{x} dx$$

input `integrate(cos(a+b*x^n)^2/x,x, algorithm="giac")`

output `integrate(cos(b*x^n + a)^2/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx^n)}{x} dx = \int \frac{\cos(a + bx^n)^2}{x} dx$$

input `int(cos(a + b*x^n)^2/x, x)`output `int(cos(a + b*x^n)^2/x, x)`**Reduce [F]**

$$\int \frac{\cos^2(a + bx^n)}{x} dx = \int \frac{\cos(x^n b + a)^2}{x} dx$$

input `int(cos(a+b*x^n)^2/x, x)`output `int(cos(x**n*b + a)**2/x, x)`

3.71 $\int \frac{\cos^3(a+bx^n)}{x} dx$

Optimal result	541
Mathematica [A] (verified)	541
Rubi [A] (verified)	542
Maple [A] (verified)	543
Fricas [A] (verification not implemented)	543
Sympy [F]	544
Maxima [C] (verification not implemented)	544
Giac [F]	545
Mupad [F(-1)]	545
Reduce [F]	545

Optimal result

Integrand size = 14, antiderivative size = 67

$$\int \frac{\cos^3(a+bx^n)}{x} dx = \frac{3 \cos(a) \operatorname{CosIntegral}(bx^n)}{4n} + \frac{\cos(3a) \operatorname{CosIntegral}(3bx^n)}{4n} - \frac{3 \sin(a) \operatorname{Si}(bx^n)}{4n} - \frac{\sin(3a) \operatorname{Si}(3bx^n)}{4n}$$

output

```
3/4*cos(a)*Ci(b*x^n)/n+1/4*cos(3*a)*Ci(3*b*x^n)/n-3/4*sin(a)*Si(b*x^n)/n-1/4*sin(3*a)*Si(3*b*x^n)/n
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.79

$$\int \frac{\cos^3(a+bx^n)}{x} dx = \frac{3 \cos(a) \operatorname{CosIntegral}(bx^n) + \cos(3a) \operatorname{CosIntegral}(3bx^n) - 3 \sin(a) \operatorname{Si}(bx^n) - \sin(3a) \operatorname{Si}(3bx^n)}{4n}$$

input

```
Integrate[Cos[a + b*x^n]^3/x,x]
```

output

```
(3*Cos[a]*CosIntegral[b*x^n] + Cos[3*a]*CosIntegral[3*b*x^n] - 3*Sin[a]*SinIntegral[b*x^n] - Sin[3*a]*SinIntegral[3*b*x^n])/(4*n)
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^3(a + bx^n)}{x} dx$$

↓ 3907

$$\int \left(\frac{3 \cos(a + bx^n)}{4x} + \frac{\cos(3a + 3bx^n)}{4x} \right) dx$$

↓ 2009

$$\frac{3 \cos(a) \text{CosIntegral}(bx^n)}{4n} + \frac{\cos(3a) \text{CosIntegral}(3bx^n)}{4n} - \frac{3 \sin(a) \text{Si}(bx^n)}{4n} - \frac{\sin(3a) \text{Si}(3bx^n)}{4n}$$

input

```
Int[Cos[a + b*x^n]^3/x,x]
```

output

```
(3*Cos[a]*CosIntegral[b*x^n])/(4*n) + (Cos[3*a]*CosIntegral[3*b*x^n])/(4*n) - (3*Sin[a]*SinIntegral[b*x^n])/(4*n) - (Sin[3*a]*SinIntegral[3*b*x^n])/(4*n)
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3907 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 1.64 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{-\frac{\text{Si}(3bx^n)\sin(3a)}{4} + \frac{\text{Ci}(3bx^n)\cos(3a)}{4} - \frac{3\text{Si}(bx^n)\sin(a)}{4} + \frac{3\text{Ci}(bx^n)\cos(a)}{4}}{n}$
default	$\frac{-\frac{\text{Si}(3bx^n)\sin(3a)}{4} + \frac{\text{Ci}(3bx^n)\cos(3a)}{4} - \frac{3\text{Si}(bx^n)\sin(a)}{4} + \frac{3\text{Ci}(bx^n)\cos(a)}{4}}{n}$
risch	$\frac{ie^{-3ia}\pi\text{csgn}(bx^n)}{8n} - \frac{ie^{-3ia}\text{Si}(3bx^n)}{4n} - \frac{e^{-3ia}\text{expIntegral}_1(-3ibx^n)}{8n} + \frac{3ie^{-ia}\pi\text{csgn}(bx^n)}{8n} - \frac{3ie^{-ia}\text{Si}(bx^n)}{4n}$

input `int(cos(a+b*x^n)^3/x,x,method=_RETURNVERBOSE)`

output `1/n*(-1/4*Si(3*b*x^n)*sin(3*a)+1/4*Ci(3*b*x^n)*cos(3*a)-3/4*Si(b*x^n)*sin(a)+3/4*Ci(b*x^n)*cos(a))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.76

$$\int \frac{\cos^3(a + bx^n)}{x} dx = \frac{\cos(3a)\text{Ci}(3bx^n) + 3\cos(a)\text{Ci}(bx^n) - \sin(3a)\text{Si}(3bx^n) - 3\sin(a)\text{Si}(bx^n)}{4n}$$

input `integrate(cos(a+b*x^n)^3/x,x, algorithm="fricas")`

output `1/4*(cos(3*a)*cos_integral(3*b*x^n) + 3*cos(a)*cos_integral(b*x^n) - sin(3*a)*sin_integral(3*b*x^n) - 3*sin(a)*sin_integral(b*x^n))/n`

Sympy [F]

$$\int \frac{\cos^3(a + bx^n)}{x} dx = \int \frac{\cos^3(a + bx^n)}{x} dx$$

input `integrate(cos(a+b*x**n)**3/x,x)`

output `Integral(cos(a + b*x**n)**3/x, x)`

Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.34 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.69

$$\int \frac{\cos^3(a + bx^n)}{x} dx = \frac{\left(\operatorname{Ei}(3i bx^n) + \operatorname{Ei}(-3i bx^n) + \operatorname{Ei}\left(3i be^{(n\overline{\log(x)})}\right) + \operatorname{Ei}\left(-3i be^{(n\overline{\log(x)})}\right) \right) \cos(3a) + 3 \left(\operatorname{Ei}(i bx^n) + \operatorname{Ei}(-i bx^n) \right) \sin(a)}{4}$$

input `integrate(cos(a+b*x^n)^3/x,x, algorithm="maxima")`

output `1/16*((Ei(3*I*b*x^n) + Ei(-3*I*b*x^n) + Ei(3*I*b*e^(n*conjugate(log(x)))) + Ei(-3*I*b*e^(n*conjugate(log(x)))))*cos(3*a) + 3*(Ei(I*b*x^n) + Ei(-I*b*x^n) + Ei(I*b*e^(n*conjugate(log(x)))) + Ei(-I*b*e^(n*conjugate(log(x)))))*cos(a) + (I*Ei(3*I*b*x^n) - I*Ei(-3*I*b*x^n) + I*Ei(3*I*b*e^(n*conjugate(log(x)))) - I*Ei(-3*I*b*e^(n*conjugate(log(x)))))*sin(3*a) - 3*(-I*Ei(I*b*x^n) + I*Ei(-I*b*x^n) - I*Ei(I*b*e^(n*conjugate(log(x)))) + I*Ei(-I*b*e^(n*conjugate(log(x)))))*sin(a))/n`

Giac [F]

$$\int \frac{\cos^3(a + bx^n)}{x} dx = \int \frac{\cos(bx^n + a)^3}{x} dx$$

input `integrate(cos(a+b*x^n)^3/x,x, algorithm="giac")`

output `integrate(cos(b*x^n + a)^3/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(a + bx^n)}{x} dx = \int \frac{\cos(a + bx^n)^3}{x} dx$$

input `int(cos(a + b*x^n)^3/x,x)`

output `int(cos(a + b*x^n)^3/x, x)`

Reduce [F]

$$\int \frac{\cos^3(a + bx^n)}{x} dx = \int \frac{\cos(x^n b + a)^3}{x} dx$$

input `int(cos(a+b*x^n)^3/x,x)`

output `int(cos(x**n*b + a)**3/x,x)`

3.72 $\int \frac{\cos^4(a+bx^n)}{x} dx$

Optimal result	546
Mathematica [A] (verified)	546
Rubi [A] (verified)	547
Maple [A] (verified)	548
Fricas [A] (verification not implemented)	548
Sympy [F]	549
Maxima [C] (verification not implemented)	549
Giac [F]	550
Mupad [F(-1)]	550
Reduce [F]	550

Optimal result

Integrand size = 14, antiderivative size = 79

$$\int \frac{\cos^4(a+bx^n)}{x} dx = \frac{\cos(2a) \operatorname{CosIntegral}(2bx^n)}{2n} + \frac{\cos(4a) \operatorname{CosIntegral}(4bx^n)}{8n} + \frac{3 \log(x)}{8} - \frac{\sin(2a) \operatorname{Si}(2bx^n)}{2n} - \frac{\sin(4a) \operatorname{Si}(4bx^n)}{8n}$$

output

```
1/2*cos(2*a)*Ci(2*b*x^n)/n+1/8*cos(4*a)*Ci(4*b*x^n)/n+3/8*ln(x)-1/2*sin(2*a)*Si(2*b*x^n)/n-1/8*sin(4*a)*Si(4*b*x^n)/n
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

$$\int \frac{\cos^4(a+bx^n)}{x} dx = \frac{3 \log(x)}{8} + \frac{4 \cos(2a) \operatorname{CosIntegral}(2bx^n) + \cos(4a) \operatorname{CosIntegral}(4bx^n) - 4 \sin(2a) \operatorname{Si}(2bx^n) - \sin(4a) \operatorname{Si}(4bx^n)}{8n}$$

input

```
Integrate[Cos[a + b*x^n]^4/x,x]
```

output

```
(3*Log[x])/8 + (4*Cos[2*a]*CosIntegral[2*b*x^n] + Cos[4*a]*CosIntegral[4*b*x^n] - 4*Sin[2*a]*SinIntegral[2*b*x^n] - Sin[4*a]*SinIntegral[4*b*x^n])/(8*n)
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos^4(a + bx^n)}{x} dx$$

$$\downarrow \text{3907}$$

$$\int \left(\frac{\cos(2a + 2bx^n)}{2x} + \frac{\cos(4a + 4bx^n)}{8x} + \frac{3}{8x} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\cos(2a) \operatorname{CosIntegral}(2bx^n)}{2n} + \frac{\cos(4a) \operatorname{CosIntegral}(4bx^n)}{8n} - \frac{\sin(2a) \operatorname{Si}(2bx^n)}{2n} - \frac{\sin(4a) \operatorname{Si}(4bx^n)}{8n} + \frac{3 \log(x)}{8}$$

input

```
Int[Cos[a + b*x^n]^4/x,x]
```

output

```
(Cos[2*a]*CosIntegral[2*b*x^n])/(2*n) + (Cos[4*a]*CosIntegral[4*b*x^n])/(8*n) + (3*Log[x])/8 - (Sin[2*a]*SinIntegral[2*b*x^n])/(2*n) - (Sin[4*a]*SinIntegral[4*b*x^n])/(8*n)
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3907 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 2.77 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{-\frac{\text{Si}(4bx^n)\sin(4a)}{8} + \frac{\text{Ci}(4bx^n)\cos(4a)}{8} - \frac{\text{Si}(2bx^n)\sin(2a)}{2} + \frac{\text{Ci}(2bx^n)\cos(2a)}{2} + \frac{3\ln(bx^n)}{8}}{n}$
default	$\frac{-\frac{\text{Si}(4bx^n)\sin(4a)}{8} + \frac{\text{Ci}(4bx^n)\cos(4a)}{8} - \frac{\text{Si}(2bx^n)\sin(2a)}{2} + \frac{\text{Ci}(2bx^n)\cos(2a)}{2} + \frac{3\ln(bx^n)}{8}}{n}$
risch	$\frac{3\ln(x)}{8} + \frac{ie^{-4ia}\pi\text{csgn}(bx^n)}{16n} - \frac{ie^{-4ia}\text{Si}(4bx^n)}{8n} - \frac{e^{-4ia}\text{expIntegral}_1(-4ibx^n)}{16n} + \frac{ie^{-2ia}\pi\text{csgn}(bx^n)}{4n} - \frac{ie^{-2ia}\text{Si}(2bx^n)}{4n}$

input `int(cos(a+b*x^n)^4/x,x,method=_RETURNVERBOSE)`

output `1/n*(-1/8*Si(4*b*x^n)*sin(4*a)+1/8*Ci(4*b*x^n)*cos(4*a)-1/2*Si(2*b*x^n)*sin(2*a)+1/2*Ci(2*b*x^n)*cos(2*a)+3/8*ln(b*x^n))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.78

$$\int \frac{\cos^4(a + bx^n)}{x} dx$$

$$= \frac{\cos(4a)\text{Ci}(4bx^n) + 4\cos(2a)\text{Ci}(2bx^n) + 3n\log(x) - \sin(4a)\text{Si}(4bx^n) - 4\sin(2a)\text{Si}(2bx^n)}{8n}$$

input `integrate(cos(a+b*x^n)^4/x,x, algorithm="fricas")`

output

```
1/8*(cos(4*a)*cos_integral(4*b*x^n) + 4*cos(2*a)*cos_integral(2*b*x^n) + 3
*n*log(x) - sin(4*a)*sin_integral(4*b*x^n) - 4*sin(2*a)*sin_integral(2*b*x
^n))/n
```

Sympy [F]

$$\int \frac{\cos^4(a + bx^n)}{x} dx = \int \frac{\cos^4(a + bx^n)}{x} dx$$

input

```
integrate(cos(a+b*x**n)**4/x,x)
```

output

```
Integral(cos(a + b*x**n)**4/x, x)
```

Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.29 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.39

$$\int \frac{\cos^4(a + bx^n)}{x} dx$$

$$= \left(\operatorname{Ei}(4i bx^n) + \operatorname{Ei}(-4i bx^n) + \operatorname{Ei}\left(4i be^{(n\overline{\log(x)})}\right) + \operatorname{Ei}\left(-4i be^{(n\overline{\log(x)})}\right) \right) \cos(4a) + 4 \left(\operatorname{Ei}(2i bx^n) + \operatorname{Ei}(-2i bx^n) + \operatorname{Ei}\left(2i be^{(n\overline{\log(x)})}\right) + \operatorname{Ei}\left(-2i be^{(n\overline{\log(x)})}\right) \right) \cos(2a) + 12n \log(x) + (I \operatorname{Ei}(4I b x^n) - I \operatorname{Ei}(-4I b x^n) + I \operatorname{Ei}(4I b e^{(n\overline{\log(x)})}) - I \operatorname{Ei}(-4I b e^{(n\overline{\log(x)})})) \sin(4a) - 4(-I \operatorname{Ei}(2I b x^n) + I \operatorname{Ei}(-2I b x^n) - I \operatorname{Ei}(2I b e^{(n\overline{\log(x)})}) + I \operatorname{Ei}(-2I b e^{(n\overline{\log(x)})})) \sin(2a) / n$$

input

```
integrate(cos(a+b*x^n)^4/x,x, algorithm="maxima")
```

output

```
1/32*((Ei(4*I*b*x^n) + Ei(-4*I*b*x^n) + Ei(4*I*b*e^(n*conjugate(log(x))))
+ Ei(-4*I*b*e^(n*conjugate(log(x)))))*cos(4*a) + 4*(Ei(2*I*b*x^n) + Ei(-2*
I*b*x^n) + Ei(2*I*b*e^(n*conjugate(log(x)))) + Ei(-2*I*b*e^(n*conjugate(lo
g(x)))))*cos(2*a) + 12*n*log(x) + (I*Ei(4*I*b*x^n) - I*Ei(-4*I*b*x^n) + I*
Ei(4*I*b*e^(n*conjugate(log(x)))) - I*Ei(-4*I*b*e^(n*conjugate(log(x)))))*
sin(4*a) - 4*(-I*Ei(2*I*b*x^n) + I*Ei(-2*I*b*x^n) - I*Ei(2*I*b*e^(n*conjug
ate(log(x)))) + I*Ei(-2*I*b*e^(n*conjugate(log(x)))))*sin(2*a))/n
```

Giac [F]

$$\int \frac{\cos^4(a + bx^n)}{x} dx = \int \frac{\cos(bx^n + a)^4}{x} dx$$

input `integrate(cos(a+b*x^n)^4/x,x, algorithm="giac")`

output `integrate(cos(b*x^n + a)^4/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(a + bx^n)}{x} dx = \int \frac{\cos(a + bx^n)^4}{x} dx$$

input `int(cos(a + b*x^n)^4/x,x)`

output `int(cos(a + b*x^n)^4/x, x)`

Reduce [F]

$$\int \frac{\cos^4(a + bx^n)}{x} dx = \int \frac{\cos(x^n b + a)^4}{x} dx$$

input `int(cos(a+b*x^n)^4/x,x)`

output `int(cos(x**n*b + a)**4/x,x)`

3.73 $\int \cos(a + bx^n) dx$

Optimal result	551
Mathematica [A] (verified)	551
Rubi [A] (verified)	552
Maple [C] (verified)	553
Fricas [F]	553
Sympy [F]	554
Maxima [F]	554
Giac [F]	554
Mupad [F(-1)]	555
Reduce [F]	555

Optimal result

Integrand size = 8, antiderivative size = 83

$$\int \cos(a + bx^n) dx = -\frac{e^{ia}x(-ibx^n)^{-1/n} \Gamma(\frac{1}{n}, -ibx^n)}{2n} - \frac{e^{-ia}x(ibx^n)^{-1/n} \Gamma(\frac{1}{n}, ibx^n)}{2n}$$

output

```
-1/2*exp(I*a)*x*GAMMA(1/n,-I*b*x^n)/n/((-I*b*x^n)^(1/n))-1/2*x*GAMMA(1/n,I
*b*x^n)/exp(I*a)/n/((I*b*x^n)^(1/n))
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.11

$$\int \cos(a + bx^n) dx = \frac{x(b^2x^{2n})^{-1/n} \left((-ibx^n)^{\frac{1}{n}} \Gamma(\frac{1}{n}, ibx^n) (\cos(a) - i \sin(a)) + (ibx^n)^{\frac{1}{n}} \Gamma(\frac{1}{n}, -ibx^n) (\cos(a) + i \sin(a)) \right)}{2n}$$

input

```
Integrate[Cos[a + b*x^n],x]
```


output

```
-1/2*(x*((-I)*b*x^n)^n^(-1)*Gamma[n^(-1), I*b*x^n]*(Cos[a] - I*Sin[a]) +
(I*b*x^n)^n^(-1)*Gamma[n^(-1), (-I)*b*x^n]*(Cos[a] + I*Sin[a]))/(n*(b^2*x
^(2*n))^n^(-1))
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3847, 2637}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos(a + bx^n) dx$$

$$\downarrow \text{3847}$$

$$\frac{1}{2} \int e^{-ibx^n - ia} dx + \frac{1}{2} \int e^{ibx^n + ia} dx$$

$$\downarrow \text{2637}$$

$$-\frac{e^{ia} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, ibx^n\right)}{2n}$$

input

```
Int[Cos[a + b*x^n], x]
```

output

```
-1/2*(E^(I*a)*x*Gamma[n^(-1), (-I)*b*x^n])/((n*((-I)*b*x^n)^n^(-1)) - (x*Gamma[n^(-1), I*b*x^n])/(2*E^(I*a)*n*(I*b*x^n)^n^(-1))
```

Defintions of rubi rules used

rule 2637 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^n)), x_Symbol] := Simp[(-F^a)*(c + d*x)*(Gamma[1/n, (-b)*(c + d*x)^n*Log[F]]/(d*n*((-b)*(c + d*x)^n*Log[F]))^(1/n)), x] /; FreeQ[{F, a, b, c, d, n}, x] && !IntegerQ[2/n]`

rule 3847 `Int[Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^n)], x_Symbol] := Simp[1/2 Int[E^((-c)*I - d*I*(e + f*x)^n), x], x] + Simp[1/2 Int[E^(c*I + d*I*(e + f*x)^n), x], x] /; FreeQ[{c, d, e, f, n}, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.41 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.90

method	result
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{2n}\right], \left[\frac{1}{2}, 1 + \frac{1}{2n}\right], -\frac{x^{2n}b^2}{4}\right) \cos(a) - \frac{b x^{1+n} \operatorname{hypergeom}\left(\left[\frac{1}{2} + \frac{1}{2n}\right], \left[\frac{3}{2}, \frac{3}{2} + \frac{1}{2n}\right], -\frac{x^{2n}b^2}{4}\right) \sin(a)}{1+n}$

input `int(cos(a+b*x^n),x,method=_RETURNVERBOSE)`

output `x*hypergeom([1/2/n],[1/2,1+1/2/n],-1/4*x^(2*n)*b^2)*cos(a)-b/(1+n)*x^(1+n)*hypergeom([1/2+1/2/n],[3/2,3/2+1/2/n],-1/4*x^(2*n)*b^2)*sin(a)`

Fricas [F]

$$\int \cos(a + bx^n) dx = \int \cos(bx^n + a) dx$$

input `integrate(cos(a+b*x^n),x, algorithm="fricas")`

output `integral(cos(b*x^n + a), x)`

Sympy [F]

$$\int \cos(a + bx^n) dx = \int \cos(a + bx^n) dx$$

input `integrate(cos(a+b*x**n),x)`

output `Integral(cos(a + b*x**n), x)`

Maxima [F]

$$\int \cos(a + bx^n) dx = \int \cos(bx^n + a) dx$$

input `integrate(cos(a+b*x^n),x, algorithm="maxima")`

output `integrate(cos(b*x^n + a), x)`

Giac [F]

$$\int \cos(a + bx^n) dx = \int \cos(bx^n + a) dx$$

input `integrate(cos(a+b*x^n),x, algorithm="giac")`

output `integrate(cos(b*x^n + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \cos(a + bx^n) dx = \int \cos(a + b x^n) dx$$

input `int(cos(a + b*x^n), x)`output `int(cos(a + b*x^n), x)`**Reduce [F]**

$$\int \cos(a + bx^n) dx = \int \cos(x^n b + a) dx + \int 1 dx - x$$

input `int(cos(a+b*x^n), x)`output `int(cos(x**n*b + a), x) + int(1, x) - x`

3.74 $\int \cos^2(a + bx^n) dx$

Optimal result	556
Mathematica [A] (verified)	556
Rubi [A] (verified)	557
Maple [F]	558
Fricas [F]	558
Sympy [F]	559
Maxima [F]	559
Giac [F]	559
Mupad [F(-1)]	560
Reduce [F]	560

Optimal result

Integrand size = 10, antiderivative size = 102

$$\int \cos^2(a + bx^n) dx = \frac{x}{2} - \frac{2^{-2-\frac{1}{n}} e^{2ia} x (-ibx^n)^{-1/n} \Gamma(\frac{1}{n}, -2ibx^n)}{n} - \frac{2^{-2-\frac{1}{n}} e^{-2ia} x (ibx^n)^{-1/n} \Gamma(\frac{1}{n}, 2ibx^n)}{n}$$

output

```
1/2*x-2^(-2-1/n)*exp(2*I*a)*x*GAMMA(1/n,-2*I*b*x^n)/n/((-I*b*x^n)^(1/n))-2^(-2-1/n)*x*GAMMA(1/n,2*I*b*x^n)/exp(2*I*a)/n/((I*b*x^n)^(1/n))
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.92

$$\int \cos^2(a + bx^n) dx = -\frac{x \left(-2n + 2^{-1/n} e^{2ia} (-ibx^n)^{-1/n} \Gamma(\frac{1}{n}, -2ibx^n) + 2^{-1/n} e^{-2ia} (ibx^n)^{-1/n} \Gamma(\frac{1}{n}, 2ibx^n) \right)}{4n}$$

input

```
Integrate[Cos[a + b*x^n]^2,x]
```

output

$$-1/4*(x*(-2*n + (E^((2*I)*a)*Gamma[n^(-1), (-2*I)*b*x^n])/(2^n^(-1)*((-I)*b*x^n)^n^(-1)) + Gamma[n^(-1), (2*I)*b*x^n]/(2^n^(-1)*E^((2*I)*a)*(I*b*x^n)^n^(-1))))/n$$
Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3849, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^2(a + bx^n) dx$$

$$\downarrow 3849$$

$$\int \left(\frac{1}{2} \cos(2a + 2bx^n) + \frac{1}{2} \right) dx$$

$$\downarrow 2009$$

$$\frac{e^{2ia} 2^{-\frac{1}{n}-2} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -2ibx^n\right)}{n} - \frac{e^{-2ia} 2^{-\frac{1}{n}-2} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, 2ibx^n\right)}{n} + \frac{x}{2}$$

input

$$\text{Int}[\text{Cos}[a + b*x^n]^2, x]$$

output

$$x/2 - (2^(-2 - n^(-1))*E^((2*I)*a)*x*Gamma[n^(-1), (-2*I)*b*x^n])/(n*((-I)*b*x^n)^n^(-1)) - (2^(-2 - n^(-1))*x*Gamma[n^(-1), (2*I)*b*x^n])/(E^((2*I)*a)*n*(I*b*x^n)^n^(-1))$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3849 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_))]*(b_.))^(p_), x_Symbol] := Int[ExpandTrigReduce[(a + b*Cos[c + d*(e + f*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 1]`

Maple [F]

$$\int \cos(a + bx^n)^2 dx$$

input `int(cos(a+b*x^n)^2,x)`

output `int(cos(a+b*x^n)^2,x)`

Fricas [F]

$$\int \cos^2(a + bx^n) dx = \int \cos(bx^n + a)^2 dx$$

input `integrate(cos(a+b*x^n)^2,x, algorithm="fricas")`

output `integral(cos(b*x^n + a)^2, x)`

Sympy [F]

$$\int \cos^2(a + bx^n) dx = \int \cos^2(a + bx^n) dx$$

input `integrate(cos(a+b*x**n)**2,x)`

output `Integral(cos(a + b*x**n)**2, x)`

Maxima [F]

$$\int \cos^2(a + bx^n) dx = \int \cos(bx^n + a)^2 dx$$

input `integrate(cos(a+b*x^n)^2,x, algorithm="maxima")`

output `1/2*x + 1/2*integrate(cos(2*b*x^n + 2*a), x)`

Giac [F]

$$\int \cos^2(a + bx^n) dx = \int \cos(bx^n + a)^2 dx$$

input `integrate(cos(a+b*x^n)^2,x, algorithm="giac")`

output `integrate(cos(b*x^n + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx^n) dx = \int \cos(a + bx^n)^2 dx$$

input `int(cos(a + b*x^n)^2,x)`output `int(cos(a + b*x^n)^2, x)`**Reduce [F]**

$$\int \cos^2(a + bx^n) dx = \int \cos(x^n b + a)^2 dx$$

input `int(cos(a+b*x^n)^2,x)`output `int(cos(x**n*b + a)**2,x)`

3.75 $\int \cos^3(a + bx^n) dx$

Optimal result	561
Mathematica [A] (verified)	562
Rubi [A] (verified)	562
Maple [F]	563
Fricas [F]	563
Sympy [F]	564
Maxima [F]	564
Giac [F]	564
Mupad [F(-1)]	565
Reduce [F]	565

Optimal result

Integrand size = 10, antiderivative size = 179

$$\int \cos^3(a + bx^n) dx = -\frac{3e^{ia}x(-ibx^n)^{-1/n} \Gamma(\frac{1}{n}, -ibx^n)}{8n} - \frac{3e^{-ia}x(ibx^n)^{-1/n} \Gamma(\frac{1}{n}, ibx^n)}{8n} - \frac{3^{-1/n}e^{3ia}x(-ibx^n)^{-1/n} \Gamma(\frac{1}{n}, -3ibx^n)}{8n} - \frac{3^{-1/n}e^{-3ia}x(ibx^n)^{-1/n} \Gamma(\frac{1}{n}, 3ibx^n)}{8n}$$

output

```
-3/8*exp(I*a)*x*GAMMA(1/n,-I*b*x^n)/n/((-I*b*x^n)^(1/n))-3/8*x*GAMMA(1/n,I
*b*x^n)/exp(I*a)/n/((I*b*x^n)^(1/n))-1/8*exp(3*I*a)*x*GAMMA(1/n,-3*I*b*x^n
)/(3^(1/n))/n/((-I*b*x^n)^(1/n))-1/8*x*GAMMA(1/n,3*I*b*x^n)/(3^(1/n))/exp(
3*I*a)/n/((I*b*x^n)^(1/n))
```

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.97

$$\int \cos^3(a + bx^n) dx = \frac{3^{-1/n} e^{-3ia} x (b^2 x^{2n})^{-1/n} \left(3^{1+\frac{1}{n}} e^{4ia} (ibx^n)^{\frac{1}{n}} \Gamma\left(\frac{1}{n}, -ibx^n\right) + 3^{1+\frac{1}{n}} e^{2ia} (-ibx^n)^{\frac{1}{n}} \Gamma\left(\frac{1}{n}, ibx^n\right) + e^{6ia} (ibx^n)^{\frac{1}{n}} \Gamma\left(\frac{1}{n}, 3ibx^n\right) \right)}{8n}$$

input

Integrate[Cos[a + b*x^n]^3, x]

output

```
-1/8*(x*(3^(1 + n^(-1))*E^((4*I)*a)*(I*b*x^n)^n^(-1)*Gamma[n^(-1), (-I)*b*x^n] + 3^(1 + n^(-1))*E^((2*I)*a)*((-I)*b*x^n)^n^(-1)*Gamma[n^(-1), I*b*x^n] + E^((6*I)*a)*(I*b*x^n)^n^(-1)*Gamma[n^(-1), (-3*I)*b*x^n] + ((-I)*b*x^n)^n^(-1)*Gamma[n^(-1), (3*I)*b*x^n]))/(3^n^(-1)*E^((3*I)*a)*n*(b^2*x^(2*n))^n^(-1))
```

Rubi [A] (verified)Time = 0.30 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3849, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos^3(a + bx^n) dx$$

$$\downarrow \text{3849}$$

$$\int \left(\frac{3}{4} \cos(a + bx^n) + \frac{1}{4} \cos(3a + 3bx^n) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3e^{ia} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -ibx^n\right)}{8n} - \frac{e^{3ia} 3^{-1/n} x (-ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, -3ibx^n\right)}{8n} - \frac{3e^{-ia} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, ibx^n\right)}{8n} - \frac{e^{-3ia} 3^{-1/n} x (ibx^n)^{-1/n} \Gamma\left(\frac{1}{n}, 3ibx^n\right)}{8n}$$

input `Int[Cos[a + b*x^n]^3,x]`

output `(-3*E^(I*a)*x*Gamma[n^(-1), (-I)*b*x^n]/(8*n*((-I)*b*x^n)^n^(-1)) - (3*x*Gamma[n^(-1), I*b*x^n]/(8*E^(I*a)*n*(I*b*x^n)^n^(-1)) - (E^((3*I)*a)*x*Gamma[n^(-1), (-3*I)*b*x^n]/(8*3^n^(-1)*n*((-I)*b*x^n)^n^(-1)) - (x*Gamma[n^(-1), (3*I)*b*x^n]/(8*3^n^(-1)*E^((3*I)*a)*n*(I*b*x^n)^n^(-1))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3849 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^p_], x_Symbol] := Int[ExpandTrigReduce[(a + b*Cos[c + d*(e + f*x)^n])^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 1]`

Maple [F]

$$\int \cos(a + bx^n)^3 dx$$

input `int(cos(a+b*x^n)^3,x)`

output `int(cos(a+b*x^n)^3,x)`

Fricas [F]

$$\int \cos^3(a + bx^n) dx = \int \cos(bx^n + a)^3 dx$$

input `integrate(cos(a+b*x^n)^3,x, algorithm="fricas")`

output `integral(cos(b*x^n + a)^3, x)`

Sympy [F]

$$\int \cos^3(a + bx^n) dx = \int \cos^3(a + bx^n) dx$$

input `integrate(cos(a+b*x**n)**3,x)`

output `Integral(cos(a + b*x**n)**3, x)`

Maxima [F]

$$\int \cos^3(a + bx^n) dx = \int \cos(bx^n + a)^3 dx$$

input `integrate(cos(a+b*x^n)^3,x, algorithm="maxima")`

output `integrate(cos(b*x^n + a)^3, x)`

Giac [F]

$$\int \cos^3(a + bx^n) dx = \int \cos(bx^n + a)^3 dx$$

input `integrate(cos(a+b*x^n)^3,x, algorithm="giac")`

output `integrate(cos(b*x^n + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \cos^3(a + bx^n) dx = \int \cos(a + bx^n)^3 dx$$

input `int(cos(a + b*x^n)^3,x)`output `int(cos(a + b*x^n)^3, x)`**Reduce [F]**

$$\int \cos^3(a + bx^n) dx = \int \cos(x^n b + a)^3 dx$$

input `int(cos(a+b*x^n)^3,x)`output `int(cos(x**n*b + a)**3,x)`

3.76 $\int x^m \cos(a + bx^n) dx$

Optimal result	566
Mathematica [A] (verified)	566
Rubi [A] (verified)	567
Maple [C] (verified)	568
Fricas [F]	568
Sympy [F]	569
Maxima [F]	569
Giac [F]	569
Mupad [F(-1)]	570
Reduce [F]	570

Optimal result

Integrand size = 12, antiderivative size = 105

$$\int x^m \cos(a + bx^n) dx = -\frac{e^{ia} x^{1+m} (-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia} x^{1+m} (ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, ibx^n\right)}{2n}$$

output

```
-1/2*exp(I*a)*x^(1+m)*GAMMA((1+m)/n,-I*b*x^n)/n/((-I*b*x^n)^((1+m)/n))-1/2
*x^(1+m)*GAMMA((1+m)/n,I*b*x^n)/exp(I*a)/n/((I*b*x^n)^((1+m)/n))
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.10

$$\int x^m \cos(a + bx^n) dx = \frac{x^{1+m} (b^n x^n)^{-\frac{1+m}{n}} \left((-ibx^n)^{\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, ibx^n\right) (\cos(a) - i \sin(a)) + (ibx^n)^{\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -ibx^n\right) (\cos(a) + i \sin(a)) \right)}{2n}$$

input

```
Integrate[x^m*Cos[a + b*x^n],x]
```

output

```
-1/2*(x^(1 + m)*((( -I)*b*x^n)^((1 + m)/n)*Gamma[(1 + m)/n, I*b*x^n]*(Cos[a]
] - I*Sin[a]) + (I*b*x^n)^((1 + m)/n)*Gamma[(1 + m)/n, (-I)*b*x^n]*(Cos[a]
+ I*Sin[a])))/(n*(b^2*x^(2*n))^((1 + m)/n))
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3905, 2648}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \cos(a + bx^n) dx$$

$$\downarrow \text{3905}$$

$$\frac{1}{2} \int e^{-ibx^n - ia} x^m dx + \frac{1}{2} \int e^{ibx^n + ia} x^m dx$$

$$\downarrow \text{2648}$$

$$-\frac{e^{ia} x^{m+1} (-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -ibx^n\right)}{2n} - \frac{e^{-ia} x^{m+1} (ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, ibx^n\right)}{2n}$$

input

```
Int[x^m*Cos[a + b*x^n], x]
```

output

```
-1/2*(E^(I*a)*x^(1 + m)*Gamma[(1 + m)/n, (-I)*b*x^n])/(n*((-I)*b*x^n)^((1
+ m)/n)) - (x^(1 + m)*Gamma[(1 + m)/n, I*b*x^n])/(2*E^(I*a)*n*(I*b*x^n)^((
1 + m)/n))
```


Definitions of rubi rules used

rule 2648

```
Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^(n_)))*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(-F^a)*((e + f*x)^(m + 1)/(f*n*((-b)*(c + d*x)^n*Log[F])^(m + 1)/n))*Gamma[(m + 1)/n, (-b)*(c + d*x)^n*Log[F]], x] /; FreeQ[{F, a, b, c, d, e, f, m, n}, x] && EqQ[d*e - c*f, 0]
```

rule 3905

```
Int[Cos[(c_.) + (d_.)*(x_)^(n_)]*((e_.)*(x_)^(m_.), x_Symbol] := Simp[1/2 Int[(e*x)^m*E^((-c)*I - d*I*x^n), x], x] + Simp[1/2 Int[(e*x)^m*E^(c*I + d*I*x^n), x], x] /; FreeQ[{c, d, e, m, n}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4.

Time = 0.80 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.06

method	result
meijerg	$\frac{x^{1+m} \operatorname{hypergeom}\left(\left[\frac{m}{2n} + \frac{1}{2n}\right], \left[\frac{1}{2}, 1 + \frac{m}{2n} + \frac{1}{2n}\right], -\frac{x^{2n}b^2}{4}\right) \cos(a)}{1+m} - \frac{b x^{1+m+n} \operatorname{hypergeom}\left(\left[\frac{1}{2} + \frac{m}{2n} + \frac{1}{2n}\right], \left[\frac{3}{2}, \frac{3}{2} + \frac{m}{2n} + \frac{1}{2n}\right], -\frac{x^{2n}b^2}{4}\right) \sin(a)}{1+m+n}$

input

```
int(x^m*cos(a+b*x^n),x,method=_RETURNVERBOSE)
```

output

```
1/(1+m)*x^(1+m)*hypergeom([1/2/n*m+1/2/n], [1/2, 1+1/2/n*m+1/2/n], -1/4*x^(2*n)*b^2)*cos(a)-b/(1+m+n)*x^(1+m+n)*hypergeom([1/2+1/2/n*m+1/2/n], [3/2, 3/2+1/2/n*m+1/2/n], -1/4*x^(2*n)*b^2)*sin(a)
```

Fricas [F]

$$\int x^m \cos(a + bx^n) dx = \int x^m \cos(bx^n + a) dx$$

input

```
integrate(x^m*cos(a+b*x^n),x, algorithm="fricas")
```

output `integral(xm*cos(b*xn + a), x)`

Sympy [F]

$$\int x^m \cos(a + bx^n) dx = \int x^m \cos(a + bx^n) dx$$

input `integrate(x**m*cos(a+b*x**n),x)`

output `Integral(x**m*cos(a + b*x**n), x)`

Maxima [F]

$$\int x^m \cos(a + bx^n) dx = \int x^m \cos(bx^n + a) dx$$

input `integrate(xm*cos(a+b*xn),x, algorithm="maxima")`

output `integrate(xm*cos(b*xn + a), x)`

Giac [F]

$$\int x^m \cos(a + bx^n) dx = \int x^m \cos(bx^n + a) dx$$

input `integrate(xm*cos(a+b*xn),x, algorithm="giac")`

output `integrate(xm*cos(b*xn + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^m \cos(a + bx^n) dx = \int x^m \cos(a + bx^n) dx$$

input `int(x^m*cos(a + b*x^n), x)`output `int(x^m*cos(a + b*x^n), x)`**Reduce [F]**

$$\int x^m \cos(a + bx^n) dx = \frac{x^m x - 2 \left(\int \frac{x^m \tan\left(\frac{x^n b + a}{2}\right)^2}{\tan\left(\frac{x^n b + a}{2}\right)^2 + 1} dx \right) m - 2 \left(\int \frac{x^m \tan\left(\frac{x^n b + a}{2}\right)^2}{\tan\left(\frac{x^n b + a}{2}\right)^2 + 1} dx \right)}{m + 1}$$

input `int(x^m*cos(a+b*x^n), x)`output `(x**m*x - 2*int((x**m*tan((x**n*b + a)/2)**2)/(tan((x**n*b + a)/2)**2 + 1), x)*m - 2*int((x**m*tan((x**n*b + a)/2)**2)/(tan((x**n*b + a)/2)**2 + 1), x))/ (m + 1)`

3.77 $\int x^m \cos^2(a + bx^n) dx$

Optimal result	571
Mathematica [A] (verified)	571
Rubi [A] (verified)	572
Maple [F]	573
Fricas [F]	573
Sympy [F]	574
Maxima [F]	574
Giac [F]	574
Mupad [F(-1)]	575
Reduce [F]	575

Optimal result

Integrand size = 14, antiderivative size = 141

$$\int x^m \cos^2(a + bx^n) dx = \frac{x^{1+m}}{2(1+m)} - \frac{2^{-\frac{1+m+2n}{n}} e^{2ia} x^{1+m} (-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -2ibx^n\right)}{n} - \frac{2^{-\frac{1+m+2n}{n}} e^{-2ia} x^{1+m} (ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, 2ibx^n\right)}{n}$$

output

```
x^(1+m)/(2+2*m)-exp(2*I*a)*x^(1+m)*GAMMA((1+m)/n,-2*I*b*x^n)/(2^((1+m+2*n)/n))/n/((-I*b*x^n)^((1+m)/n))-x^(1+m)*GAMMA((1+m)/n,2*I*b*x^n)/(2^((1+m+2*n)/n))/exp(2*I*a)/n/((I*b*x^n)^((1+m)/n))
```

Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.91

$$\int x^m \cos^2(a + bx^n) dx = \frac{x^{1+m} \left(-2n + 2^{-\frac{1+m}{n}} e^{2ia} (1+m) (-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -2ibx^n\right) + 2^{-\frac{1+m}{n}} e^{-2ia} (1+m) (ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, 2ibx^n\right) \right)}{4(1+m)n}$$

input

```
Integrate[x^m*Cos[a + b*x^n]^2,x]
```

output

```
-1/4*(x^(1 + m)*(-2*n + (E^((2*I)*a))*(1 + m)*Gamma[(1 + m)/n, (-2*I)*b*x^n
])/2^((1 + m)/n)*((-I)*b*x^n)^((1 + m)/n)) + ((1 + m)*Gamma[(1 + m)/n, (2
*I)*b*x^n])/2^((1 + m)/n)*E^((2*I)*a)*(I*b*x^n)^((1 + m)/n)))/((1 + m)*n
)
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \cos^2(a + bx^n) dx$$

↓ 3907

$$\int \left(\frac{1}{2} x^m \cos(2a + 2bx^n) + \frac{x^m}{2} \right) dx$$

↓ 2009

$$\frac{e^{2ia} 2^{-\frac{m+2n+1}{n}} x^{m+1} (-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -2ibx^n\right)}{n} - \frac{e^{-2ia} 2^{-\frac{m+2n+1}{n}} x^{m+1} (ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, 2ibx^n\right)}{n} + \frac{x^{m+1}}{2(m+1)}$$

input

```
Int[x^m*Cos[a + b*x^n]^2,x]
```

output

```
x^(1 + m)/(2*(1 + m)) - (E^((2*I)*a)*x^(1 + m)*Gamma[(1 + m)/n, (-2*I)*b*x^n
])/2^((1 + m + 2*n)/n)*n*((-I)*b*x^n)^((1 + m)/n)) - (x^(1 + m)*Gamma[(1 + m)/n, (2*I)*b*x^n])/2^((1 + m + 2*n)/n)*E^((2*I)*a)*n*(I*b*x^n)^((1 + m)/n))
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3907 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_.)]*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

Maple [F]

$$\int x^m \cos(a + b x^n)^2 dx$$

input `int(x^m*cos(a+b*x^n)^2,x)`

output `int(x^m*cos(a+b*x^n)^2,x)`

Fricas [F]

$$\int x^m \cos^2(a + b x^n) dx = \int x^m \cos(b x^n + a)^2 dx$$

input `integrate(x^m*cos(a+b*x^n)^2,x, algorithm="fricas")`

output `integral(x^m*cos(b*x^n + a)^2, x)`

Sympy [F]

$$\int x^m \cos^2(a + bx^n) dx = \int x^m \cos^2(a + bx^n) dx$$

input `integrate(x**m*cos(a+b*x**n)**2,x)`

output `Integral(x**m*cos(a + b*x**n)**2, x)`

Maxima [F]

$$\int x^m \cos^2(a + bx^n) dx = \int x^m \cos(bx^n + a)^2 dx$$

input `integrate(x^m*cos(a+b*x^n)^2,x, algorithm="maxima")`

output `1/2*(x*x^m + (m + 1)*integrate(x^m*cos(2*b*x^n + 2*a), x))/(m + 1)`

Giac [F]

$$\int x^m \cos^2(a + bx^n) dx = \int x^m \cos(bx^n + a)^2 dx$$

input `integrate(x^m*cos(a+b*x^n)^2,x, algorithm="giac")`

output `integrate(x^m*cos(b*x^n + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^m \cos^2(a + bx^n) dx = \int x^m \cos(a + bx^n)^2 dx$$

input `int(x^m*cos(a + b*x^n)^2,x)`output `int(x^m*cos(a + b*x^n)^2, x)`**Reduce [F]**

$$\int x^m \cos^2(a + bx^n) dx = \int x^m \cos(x^n b + a)^2 dx$$

input `int(x^m*cos(a+b*x^n)^2,x)`output `int(x**m*cos(x**n*b + a)**2,x)`

3.78 $\int x^m \cos^3(a + bx^n) dx$

Optimal result	576
Mathematica [A] (verified)	577
Rubi [A] (verified)	577
Maple [F]	578
Fricas [F]	578
Sympy [F]	579
Maxima [F]	579
Giac [F]	579
Mupad [F(-1)]	580
Reduce [F]	580

Optimal result

Integrand size = 14, antiderivative size = 229

$$\int x^m \cos^3(a + bx^n) dx = -\frac{3e^{ia}x^{1+m}(-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -ibx^n\right)}{8n} - \frac{3e^{-ia}x^{1+m}(ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, ibx^n\right)}{8n} - \frac{3^{-\frac{1+m}{n}} e^{3ia}x^{1+m}(-ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -3ibx^n\right)}{8n} - \frac{3^{-\frac{1+m}{n}} e^{-3ia}x^{1+m}(ibx^n)^{-\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, 3ibx^n\right)}{8n}$$

output

```
-3/8*exp(I*a)*x^(1+m)*GAMMA((1+m)/n, -I*b*x^n)/n/((-I*b*x^n)^((1+m)/n))-3/8
*x^(1+m)*GAMMA((1+m)/n, I*b*x^n)/exp(I*a)/n/((I*b*x^n)^((1+m)/n))-1/8*exp(3
*I*a)*x^(1+m)*GAMMA((1+m)/n, -3*I*b*x^n)/(3^((1+m)/n))/n/((-I*b*x^n)^((1+m)
/n))-1/8*x^(1+m)*GAMMA((1+m)/n, 3*I*b*x^n)/(3^((1+m)/n))/exp(3*I*a)/n/((I*b
*x^n)^((1+m)/n))
```

Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.97

$$\int x^m \cos^3(a + bx^n) dx = \frac{3^{-\frac{1+m}{n}} e^{-3ia} x^{1+m} (b^2 x^{2n})^{-\frac{1+m}{n}} \left(3^{\frac{1+m+n}{n}} e^{4ia} (ibx^n)^{\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, -ibx^n\right) + 3^{\frac{1+m+n}{n}} e^{2ia} (-ibx^n)^{\frac{1+m}{n}} \Gamma\left(\frac{1+m}{n}, ibx^n\right) \right)}{8n}$$

input

```
Integrate[x^m*Cos[a + b*x^n]^3,x]
```

output

```
-1/8*(x^(1 + m)*(3^((1 + m + n)/n)*E^((4*I)*a)*(I*b*x^n)^((1 + m)/n)*Gamma
[(1 + m)/n, (-I)*b*x^n] + 3^((1 + m + n)/n)*E^((2*I)*a)*((-I)*b*x^n)^((1 +
m)/n)*Gamma[(1 + m)/n, I*b*x^n] + E^((6*I)*a)*(I*b*x^n)^((1 + m)/n)*Gamma
[(1 + m)/n, (-3*I)*b*x^n] + ((-I)*b*x^n)^((1 + m)/n)*Gamma[(1 + m)/n, (3*I
)*b*x^n)))/(3^((1 + m)/n)*E^((3*I)*a)*n*(b^2*x^(2*n))^((1 + m)/n))
```

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \cos^3(a + bx^n) dx$$

$$\downarrow \text{3907}$$

$$\int \left(\frac{3}{4} x^m \cos(a + bx^n) + \frac{1}{4} x^m \cos(3a + 3bx^n) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{3e^{ia} x^{m+1} (-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -ibx^n\right)}{8n} - \frac{3e^{-ia} x^{m+1} (ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, ibx^n\right)}{8n} - \frac{e^{3ia} 3^{-\frac{m+1}{n}} x^{m+1} (-ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, -3ibx^n\right)}{8n} - \frac{e^{-3ia} 3^{-\frac{m+1}{n}} x^{m+1} (ibx^n)^{-\frac{m+1}{n}} \Gamma\left(\frac{m+1}{n}, 3ibx^n\right)}{8n}$$

input `Int[x^m*Cos[a + b*x^n]^3,x]`

output `(-3*E^(I*a)*x^(1 + m)*Gamma[(1 + m)/n, (-I)*b*x^n])/(8*n*((-I)*b*x^n)^((1 + m)/n)) - (3*x^(1 + m)*Gamma[(1 + m)/n, I*b*x^n])/(8*E^(I*a)*n*(I*b*x^n)^((1 + m)/n)) - (E^((3*I)*a)*x^(1 + m)*Gamma[(1 + m)/n, (-3*I)*b*x^n])/(8*3^((1 + m)/n)*n*((-I)*b*x^n)^((1 + m)/n)) - (x^(1 + m)*Gamma[(1 + m)/n, (3*I)*b*x^n])/(8*3^((1 + m)/n)*E^((3*I)*a)*n*(I*b*x^n)^((1 + m)/n))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3907 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^p)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

Maple [F]

$$\int x^m \cos(a + bx^n)^3 dx$$

input `int(x^m*cos(a+b*x^n)^3,x)`

output `int(x^m*cos(a+b*x^n)^3,x)`

Fricas [F]

$$\int x^m \cos^3(a + bx^n) dx = \int x^m \cos(bx^n + a)^3 dx$$

input `integrate(x^m*cos(a+b*x^n)^3,x, algorithm="fricas")`

output `integral(xm*cos(b*xn + a)3, x)`

Sympy [F]

$$\int x^m \cos^3(a + bx^n) dx = \int x^m \cos^3(a + bx^n) dx$$

input `integrate(x**m*cos(a+b*x**n)**3,x)`

output `Integral(x**m*cos(a + b*x**n)**3, x)`

Maxima [F]

$$\int x^m \cos^3(a + bx^n) dx = \int x^m \cos(bx^n + a)^3 dx$$

input `integrate(xm*cos(a+b*xn)3,x, algorithm="maxima")`

output `integrate(xm*cos(b*xn + a)3, x)`

Giac [F]

$$\int x^m \cos^3(a + bx^n) dx = \int x^m \cos(bx^n + a)^3 dx$$

input `integrate(xm*cos(a+b*xn)3,x, algorithm="giac")`

output `integrate(xm*cos(b*xn + a)3, x)`

Mupad [F(-1)]

Timed out.

$$\int x^m \cos^3(a + bx^n) dx = \int x^m \cos(a + bx^n)^3 dx$$

input `int(x^m*cos(a + b*x^n)^3,x)`output `int(x^m*cos(a + b*x^n)^3, x)`**Reduce [F]**

$$\int x^m \cos^3(a + bx^n) dx = \int x^m \cos(x^n b + a)^3 dx$$

input `int(x^m*cos(a+b*x^n)^3,x)`output `int(x**m*cos(x**n*b + a)**3,x)`

3.79 $\int x^{-1-n} \cos(a + bx^n) dx$

Optimal result	581
Mathematica [A] (verified)	581
Rubi [A] (verified)	582
Maple [A] (verified)	584
Fricas [A] (verification not implemented)	584
Sympy [F]	585
Maxima [F]	585
Giac [F]	585
Mupad [F(-1)]	586
Reduce [F]	586

Optimal result

Integrand size = 16, antiderivative size = 47

$$\int x^{-1-n} \cos(a + bx^n) dx = -\frac{x^{-n} \cos(a + bx^n)}{n} - \frac{b \operatorname{CosIntegral}(bx^n) \sin(a)}{n} - \frac{b \cos(a) \operatorname{Si}(bx^n)}{n}$$

output `-cos(a+b*x^n)/n/(x^n)-b*Ci(b*x^n)*sin(a)/n-b*cos(a)*Si(b*x^n)/n`

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int x^{-1-n} \cos(a + bx^n) dx = -\frac{x^{-n}(\cos(a + bx^n) + bx^n \operatorname{CosIntegral}(bx^n) \sin(a) + bx^n \cos(a) \operatorname{Si}(bx^n))}{n}$$

input `Integrate[x^(-1 - n)*Cos[a + b*x^n], x]`

output `-((Cos[a + b*x^n] + b*x^n*CosIntegral[b*x^n]*Sin[a] + b*x^n*Cos[a]*SinIntegral[b*x^n])/(n*x^n))`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {3861, 3042, 3778, 25, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^{-n-1} \cos(a + bx^n) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{\int x^{-2n} \cos(bx^n + a) dx^n}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int x^{-2n} \sin(bx^n + a + \frac{\pi}{2}) dx^n}{n} \\
 & \quad \downarrow \text{3778} \\
 & \frac{b \int -x^{-n} \sin(bx^n + a) dx^n - x^{-n} \cos(a + bx^n)}{n} \\
 & \quad \downarrow \text{25} \\
 & \frac{x^{-n}(-\cos(a + bx^n)) - b \int x^{-n} \sin(bx^n + a) dx^n}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x^{-n}(-\cos(a + bx^n)) - b \int x^{-n} \sin(bx^n + a) dx^n}{n} \\
 & \quad \downarrow \text{3784} \\
 & \frac{x^{-n}(-\cos(a + bx^n)) - b(\sin(a) \int x^{-n} \cos(bx^n) dx^n + \cos(a) \int x^{-n} \sin(bx^n) dx^n)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x^{-n}(-\cos(a + bx^n)) - b(\sin(a) \int x^{-n} \sin(bx^n + \frac{\pi}{2}) dx^n + \cos(a) \int x^{-n} \sin(bx^n) dx^n)}{n} \\
 & \quad \downarrow \text{3780} \\
 & \frac{x^{-n}(-\cos(a + bx^n)) - b(\sin(a) \int x^{-n} \sin(bx^n + \frac{\pi}{2}) dx^n + \cos(a) \text{Si}(bx^n))}{n}
 \end{aligned}$$

$$\frac{x^{-n}(-\cos(a + bx^n)) - b(\sin(a) \operatorname{CosIntegral}(bx^n) + \cos(a) \operatorname{Si}(bx^n))}{n}$$

input `Int[x^(-1 - n)*Cos[a + b*x^n], x]`

output `(-(Cos[a + b*x^n]/x^n) - b*(CosIntegral[b*x^n]*Sin[a] + Cos[a]*SinIntegral[b*x^n]))/n`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3778 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3780 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

rule 3784

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*
e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*
f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x]
&& NeQ[d*e - c*f, 0]
```

rule 3861

```
Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol]
:= Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*cos[c + d*x])^
p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[
(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[
(m + 1)/n], 0]))
```

Maple [A] (verified)

Time = 1.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

method	result	si
default	$b \frac{\left(-\frac{\cos(a+bx^n)x^{-n}}{b} - \text{Si}(bx^n) \cos(a) - \text{Ci}(bx^n) \sin(a) \right)}{n}$	4
risch	$\frac{be^{-ia}\pi \text{csgn}(bx^n)}{2n} - \frac{be^{-ia} \text{Si}(bx^n)}{n} + \frac{ibe^{-ia} \exp \text{Integral}_1(-ibx^n)}{2n} - \frac{ibe^{ia} \exp \text{Integral}_1(-ibx^n)}{2n} - \frac{\cos(a+bx^n)x^{-n}}{n}$	9

input

```
int(x^(-1-n)*cos(a+b*x^n),x,method=_RETURNVERBOSE)
```

output

```
1/n*b*(-cos(a+b*x^n)/b/(x^n)-Si(b*x^n)*cos(a)-Ci(b*x^n)*sin(a))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int x^{-1-n} \cos(a + bx^n) dx = -\frac{bx^n \text{Ci}(bx^n) \sin(a) + bx^n \cos(a) \text{Si}(bx^n) + \cos(bx^n + a)}{nx^n}$$

input

```
integrate(x^(-1-n)*cos(a+b*x^n),x, algorithm="fricas")
```

output `-(b*x^n*cos_integral(b*x^n)*sin(a) + b*x^n*cos(a)*sin_integral(b*x^n) + cos(b*x^n + a))/(n*x^n)`

Sympy [F]

$$\int x^{-1-n} \cos(a + bx^n) dx = \int x^{-n-1} \cos(a + bx^n) dx$$

input `integrate(x**(-1-n)*cos(a+b*x**n),x)`

output `Integral(x**(-n - 1)*cos(a + b*x**n), x)`

Maxima [F]

$$\int x^{-1-n} \cos(a + bx^n) dx = \int x^{-n-1} \cos(bx^n + a) dx$$

input `integrate(x^(-1-n)*cos(a+b*x^n),x, algorithm="maxima")`

output `integrate(x^(-n - 1)*cos(b*x^n + a), x)`

Giac [F]

$$\int x^{-1-n} \cos(a + bx^n) dx = \int x^{-n-1} \cos(bx^n + a) dx$$

input `integrate(x^(-1-n)*cos(a+b*x^n),x, algorithm="giac")`

output `integrate(x^(-n - 1)*cos(b*x^n + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{-1-n} \cos(a + bx^n) dx = \int \frac{\cos(a + bx^n)}{x^{n+1}} dx$$

input `int(cos(a + b*x^n)/x^(n + 1),x)`output `int(cos(a + b*x^n)/x^(n + 1), x)`**Reduce [F]**

$$\int x^{-1-n} \cos(a + bx^n) dx = \frac{x^n \left(\int \frac{\cos(x^n b + a)}{x^n} dx \right) n + x^n \left(\int \frac{1}{x^n} dx \right) n + 1}{x^n n}$$

input `int(x^(-1-n)*cos(a+b*x^n),x)`output `(x**n*int(cos(x**n*b + a)/(x**n*x),x)*n + x**n*int(1/(x**n*x),x)*n + 1)/(x**n*n)`

3.80 $\int x^{-1-n} \cos^2(a + bx^n) dx$

Optimal result	587
Mathematica [A] (verified)	587
Rubi [A] (verified)	588
Maple [A] (verified)	589
Fricas [A] (verification not implemented)	589
Sympy [F]	590
Maxima [F]	590
Giac [F]	590
Mupad [F(-1)]	591
Reduce [F]	591

Optimal result

Integrand size = 18, antiderivative size = 70

$$\int x^{-1-n} \cos^2(a + bx^n) dx = -\frac{x^{-n}}{2n} - \frac{x^{-n} \cos(2a + 2bx^n)}{2n} - \frac{b \operatorname{CosIntegral}(2bx^n) \sin(2a)}{n} - \frac{b \cos(2a) \operatorname{Si}(2bx^n)}{n}$$

output

```
-1/2/n/(x^n)-1/2*cos(2*a+2*b*x^n)/n/(x^n)-b*Ci(2*b*x^n)*sin(2*a)/n-b*cos(2*a)*Si(2*b*x^n)/n
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.76

$$\int x^{-1-n} \cos^2(a + bx^n) dx = -\frac{x^{-n}(\cos^2(a + bx^n) + bx^n \operatorname{CosIntegral}(2bx^n) \sin(2a) + bx^n \cos(2a) \operatorname{Si}(2bx^n))}{n}$$

input

```
Integrate[x^(-1 - n)*Cos[a + b*x^n]^2,x]
```

output

$$-\left(\frac{\cos[a + b x^n]^2 + b x^n \operatorname{CosIntegral}[2 b x^n] \sin[2 a] + b x^n \cos[2 a] \operatorname{SinIntegral}[2 b x^n]}{n x^n}\right)$$
Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.99, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{-n-1} \cos^2(a + b x^n) dx$$

$$\downarrow \text{3907}$$

$$\int \left(\frac{1}{2} x^{-n-1} \cos(2a + 2b x^n) + \frac{x^{-n-1}}{2} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{b \sin(2a) \operatorname{CosIntegral}(2b x^n)}{n} - \frac{b \cos(2a) \operatorname{Si}(2b x^n)}{n} - \frac{x^{-n} \cos(2(a + b x^n))}{2n} - \frac{x^{-n}}{2n}$$

input

$$\text{Int}[x^{(-1 - n)} \cos[a + b x^n]^2, x]$$

output

$$-1/2 * 1/(n * x^n) - \cos[2 * (a + b * x^n)] / (2 * n * x^n) - (b * \operatorname{CosIntegral}[2 * b * x^n] * \sin[2 * a]) / n - (b * \cos[2 * a] * \operatorname{SinIntegral}[2 * b * x^n]) / n$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ /; } \text{SumQ}[u]$$

rule 3907

$$\text{Int}[\left((a_.) + \cos[(c_.) + (d_.) * (x_)^(n_.)] * (b_.)\right)^(p_.) * \left((e_.) * (x_)^(m_.)\right), x_Symbol] \text{ :> } \text{Int}[\text{ExpandTrigReduce}[(e * x)^m, (a + b * \cos[c + d * x^n])^p, x], x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0]$$

Maple [A] (verified)

Time = 2.71 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.93

method	result
default	$-\frac{x^{-n}}{2n} + \frac{b \left(-\frac{\cos(2a+2bx^n)x^{-n}}{2b} - \text{Si}(2bx^n) \cos(2a) - \text{Ci}(2bx^n) \sin(2a) \right)}{n}$
risch	$\frac{(ib e^{-2ia} \exp\text{Integral}_1(-2ibx^n)x^n - ib e^{2ia} \exp\text{Integral}_1(-2ibx^n)x^n + b e^{-2ia} \pi \text{csgn}(bx^n)x^n - 2b e^{-2ia} \text{Si}(2bx^n)x^n - \cos(2a+2bx^n))}{2n}$

input `int(x^(-1-n)*cos(a+b*x^n)^2,x,method=_RETURNVERBOSE)`output
$$-1/2/n/(x^n) + 1/n*b*(-1/2*\cos(2*a+2*b*x^n)/b/(x^n) - \text{Si}(2*b*x^n)*\cos(2*a) - \text{Ci}(2*b*x^n)*\sin(2*a))$$
Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.76

$$\int x^{-1-n} \cos^2(a + bx^n) dx$$

$$= -\frac{bx^n \text{Ci}(2bx^n) \sin(2a) + bx^n \cos(2a) \text{Si}(2bx^n) + \cos(bx^n + a)^2}{nx^n}$$

input `integrate(x^(-1-n)*cos(a+b*x^n)^2,x, algorithm="fricas")`output
$$-(b*x^n*\cos_integral(2*b*x^n)*\sin(2*a) + b*x^n*\cos(2*a)*\sin_integral(2*b*x^n) + \cos(b*x^n + a)^2)/(n*x^n)$$

Sympy [F]

$$\int x^{-1-n} \cos^2(a + bx^n) dx = \int x^{-n-1} \cos^2(a + bx^n) dx$$

input `integrate(x**(-1-n)*cos(a+b*x**n)**2,x)`

output `Integral(x**(-n - 1)*cos(a + b*x**n)**2, x)`

Maxima [F]

$$\int x^{-1-n} \cos^2(a + bx^n) dx = \int x^{-n-1} \cos(bx^n + a)^2 dx$$

input `integrate(x^(-1-n)*cos(a+b*x^n)^2,x, algorithm="maxima")`

output `1/2*(n*x^n*integrate(cos(2*b*x^n + 2*a)/(x*x^n), x) - 1)/(n*x^n)`

Giac [F]

$$\int x^{-1-n} \cos^2(a + bx^n) dx = \int x^{-n-1} \cos(bx^n + a)^2 dx$$

input `integrate(x^(-1-n)*cos(a+b*x^n)^2,x, algorithm="giac")`

output `integrate(x^(-n - 1)*cos(b*x^n + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^{-1-n} \cos^2(a + bx^n) dx = \int \frac{\cos(a + bx^n)^2}{x^{n+1}} dx$$

input `int(cos(a + b*x^n)^2/x^(n + 1),x)`output `int(cos(a + b*x^n)^2/x^(n + 1), x)`**Reduce [F]**

$$\int x^{-1-n} \cos^2(a + bx^n) dx = \int \frac{\cos(x^n b + a)^2}{x^n x} dx$$

input `int(x^(-1-n)*cos(a+b*x^n)^2,x)`output `int(cos(x**n*b + a)**2/(x**n*x),x)`

3.81 $\int x^{-1-n} \cos^3(a + bx^n) dx$

Optimal result	592
Mathematica [A] (verified)	593
Rubi [A] (verified)	593
Maple [A] (verified)	594
Fricas [A] (verification not implemented)	595
Sympy [F]	595
Maxima [F]	595
Giac [F]	596
Mupad [F(-1)]	596
Reduce [F]	596

Optimal result

Integrand size = 18, antiderivative size = 114

$$\int x^{-1-n} \cos^3(a + bx^n) dx = -\frac{3x^{-n} \cos(a + bx^n)}{4n} - \frac{x^{-n} \cos(3a + 3bx^n)}{4n} - \frac{3b \operatorname{CosIntegral}(bx^n) \sin(a)}{4n} - \frac{3b \operatorname{CosIntegral}(3bx^n) \sin(3a)}{4n} - \frac{3b \cos(a) \operatorname{Si}(bx^n)}{4n} - \frac{3b \cos(3a) \operatorname{Si}(3bx^n)}{4n}$$

output

```
-3/4*cos(a+b*x^n)/n/(x^n)-1/4*cos(3*a+3*b*x^n)/n/(x^n)-3/4*b*Ci(b*x^n)*sin(a)/n-3/4*b*Ci(3*b*x^n)*sin(3*a)/n-3/4*b*cos(a)*Si(b*x^n)/n-3/4*b*cos(3*a)*Si(3*b*x^n)/n
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.83

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \frac{x^{-n}(3 \cos(a + bx^n) + \cos(3(a + bx^n))) + 3bx^n \operatorname{CosIntegral}(bx^n) \sin(a) + 3bx^n \operatorname{CosIntegral}(3bx^n) \sin(3a)}{4n}$$

input `Integrate[x^(-1 - n)*Cos[a + b*x^n]^3,x]`output `-1/4*(3*Cos[a + b*x^n] + Cos[3*(a + b*x^n)]) + 3*b*x^n*CosIntegral[b*x^n]*Sin[a] + 3*b*x^n*CosIntegral[3*b*x^n]*Sin[3*a] + 3*b*x^n*Cos[a]*SinIntegral[b*x^n] + 3*b*x^n*Cos[3*a]*SinIntegral[3*b*x^n]/(n*x^n)`**Rubi [A] (verified)**Time = 0.41 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.99, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{-n-1} \cos^3(a + bx^n) dx \\ & \quad \downarrow \text{3907} \\ & \int \left(\frac{3}{4} x^{-n-1} \cos(a + bx^n) + \frac{1}{4} x^{-n-1} \cos(3a + 3bx^n) \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{3b \sin(a) \operatorname{CosIntegral}(bx^n)}{4n} - \frac{3b \sin(3a) \operatorname{CosIntegral}(3bx^n)}{4n} - \frac{3b \cos(a) \operatorname{Si}(bx^n)}{4n} \\ & \quad - \frac{3b \cos(3a) \operatorname{Si}(3bx^n)}{4n} - \frac{3x^{-n} \cos(a + bx^n)}{4n} - \frac{x^{-n} \cos(3(a + bx^n))}{4n} \end{aligned}$$

input `Int[x^(-1 - n)*Cos[a + b*x^n]^3,x]`

output
$$\frac{(-3\cos[a + b*x^n])/(4*n*x^n) - \cos[3*(a + b*x^n)]/(4*n*x^n) - (3*b*\text{CosIntegral}[b*x^n]*\text{Sin}[a])/(4*n) - (3*b*\text{CosIntegral}[3*b*x^n]*\text{Sin}[3*a])/(4*n) - (3*b*\cos[a]*\text{SinIntegral}[b*x^n])/(4*n) - (3*b*\cos[3*a]*\text{SinIntegral}[3*b*x^n])/(4*n)}$$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3907 $\text{Int}[(a_.) + \text{Cos}[(c_.) + (d_.)*(x_)^(n_)]*(b_.)]^(p_)*((e_.)*(x_)^(m_.), x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(e*x)^m, (a + b*\text{Cos}[c + d*x^n])^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

Maple [A] (verified)

Time = 9.82 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.89

method	result
default	$\frac{3b \left(-\frac{\cos(a+bx^n)x^{-n}}{b} - \text{Si}(bx^n) \cos(a) - \text{Ci}(bx^n) \sin(a) \right)}{4n} + \frac{3b \left(-\frac{\cos(3a+3bx^n)x^{-n}}{3b} - \text{Si}(3bx^n) \cos(3a) - \text{Ci}(3bx^n) \sin(3a) \right)}{4n}$
risch	$-\frac{(-3be^{-3ia}\pi \text{csgn}(bx^n)x^n - 3be^{-ia}\pi \text{csgn}(bx^n)x^n - 3ibe^{-3ia} \exp\text{Integral}_1(-3ibx^n)x^n - 3ibe^{-ia} \exp\text{Integral}_1(-ibx^n)x^n + 3ib \dots)}{\dots}$

input $\text{int}(x^{(-1-n)}*\cos(a+b*x^n)^3,x,\text{method}=_RETURNVERBOSE)$

output
$$\frac{3}{4} \frac{1}{n} b * (-\cos(a+b*x^n)/b/(x^n) - \text{Si}(b*x^n)*\cos(a) - \text{Ci}(b*x^n)*\sin(a)) + \frac{3}{4} \frac{1}{n} b * (-\frac{1}{3} \cos(3a+3*b*x^n)/b/(x^n) - \text{Si}(3*b*x^n)*\cos(3*a) - \text{Ci}(3*b*x^n)*\sin(3*a))$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.75

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \frac{3bx^n \operatorname{Ci}(3bx^n) \sin(3a) + 3bx^n \operatorname{Ci}(bx^n) \sin(a) + 3bx^n \cos(3a) \operatorname{Si}(3bx^n) + 3bx^n \cos(a) \operatorname{Si}(bx^n) + 4}{4nx^n}$$

input `integrate(x^(-1-n)*cos(a+b*x^n)^3,x, algorithm="fricas")`output `-1/4*(3*b*x^n*cos_integral(3*b*x^n)*sin(3*a) + 3*b*x^n*cos_integral(b*x^n)*sin(a) + 3*b*x^n*cos(3*a)*sin_integral(3*b*x^n) + 3*b*x^n*cos(a)*sin_integral(b*x^n) + 4*cos(b*x^n + a)^3)/(n*x^n)`**Sympy [F]**

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \int x^{-n-1} \cos^3(a + bx^n) dx$$

input `integrate(x**(-1-n)*cos(a+b*x**n)**3,x)`output `Integral(x**(-n - 1)*cos(a + b*x**n)**3, x)`**Maxima [F]**

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \int x^{-n-1} \cos(bx^n + a)^3 dx$$

input `integrate(x^(-1-n)*cos(a+b*x^n)^3,x, algorithm="maxima")`output `integrate(x^(-n - 1)*cos(b*x^n + a)^3, x)`

Giac [F]

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \int x^{-n-1} \cos(bx^n + a)^3 dx$$

input `integrate(x^(-1-n)*cos(a+b*x^n)^3,x, algorithm="giac")`

output `integrate(x^(-n - 1)*cos(b*x^n + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \int \frac{\cos(a + bx^n)^3}{x^{n+1}} dx$$

input `int(cos(a + b*x^n)^3/x^(n + 1),x)`

output `int(cos(a + b*x^n)^3/x^(n + 1), x)`

Reduce [F]

$$\int x^{-1-n} \cos^3(a + bx^n) dx = \int \frac{\cos(x^n b + a)^3}{x^n x} dx$$

input `int(x^(-1-n)*cos(a+b*x^n)^3,x)`

output `int(cos(x**n*b + a)**3/(x**n*x),x)`

3.82 $\int x^{-1-2n} \cos(a + bx^n) dx$

Optimal result	597
Mathematica [A] (verified)	597
Rubi [A] (verified)	598
Maple [A] (verified)	600
Fricas [A] (verification not implemented)	601
Sympy [F]	601
Maxima [F]	602
Giac [F]	602
Mupad [F(-1)]	602
Reduce [F]	603

Optimal result

Integrand size = 16, antiderivative size = 78

$$\int x^{-1-2n} \cos(a + bx^n) dx = -\frac{x^{-2n} \cos(a + bx^n)}{2n} - \frac{b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{2n} + \frac{bx^{-n} \sin(a + bx^n)}{2n} + \frac{b^2 \sin(a) \operatorname{Si}(bx^n)}{2n}$$

output
$$-1/2*\cos(a+b*x^n)/n/(x^{(2*n)})-1/2*b^2*\cos(a)*\operatorname{Ci}(b*x^n)/n+1/2*b*\sin(a+b*x^n)/n/(x^n)+1/2*b^2*\sin(a)*\operatorname{Si}(b*x^n)/n$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

$$\int x^{-1-2n} \cos(a + bx^n) dx = \frac{x^{-2n}(\cos(a + bx^n) + b^2 x^{2n} \cos(a) \operatorname{CosIntegral}(bx^n) - bx^n \sin(a + bx^n) - b^2 x^{2n} \sin(a) \operatorname{Si}(bx^n))}{2n}$$

input
$$\operatorname{Integrate}[x^{(-1 - 2*n)}*\operatorname{Cos}[a + b*x^n], x]$$

output

$$-1/2*(\text{Cos}[a + b*x^n] + b^2*x^{(2*n)}*\text{Cos}[a]*\text{CosIntegral}[b*x^n] - b*x^n*\text{Sin}[a + b*x^n] - b^2*x^{(2*n)}*\text{Sin}[a]*\text{SinIntegral}[b*x^n])/ (n*x^{(2*n)})$$
Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.83, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {3861, 3042, 3778, 25, 3042, 3778, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^{-2n-1} \cos(a + bx^n) dx \\
 & \quad \downarrow \text{3861} \\
 & \frac{\int x^{-3n} \cos(bx^n + a) dx^n}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int x^{-3n} \sin(bx^n + a + \frac{\pi}{2}) dx^n}{n} \\
 & \quad \downarrow \text{3778} \\
 & \frac{\frac{1}{2}b \int -x^{-2n} \sin(bx^n + a) dx^n - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\frac{1}{2}b \int x^{-2n} \sin(bx^n + a) dx^n - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{2}b \int x^{-2n} \sin(bx^n + a) dx^n - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n} \\
 & \quad \downarrow \text{3778} \\
 & \frac{-\frac{1}{2}b(b \int x^{-n} \cos(bx^n + a) dx^n - x^{-n} \sin(a + bx^n)) - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{-\frac{1}{2}b(b \int x^{-n} \sin(bx^n + a + \frac{\pi}{2}) dx^n - x^{-n} \sin(a + bx^n)) - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n}$$

↓ 3784

$$\frac{-\frac{1}{2}b(b(\cos(a) \int x^{-n} \cos(bx^n) dx^n - \sin(a) \int x^{-n} \sin(bx^n) dx^n) - x^{-n} \sin(a + bx^n)) - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n}$$

↓ 3042

$$\frac{-\frac{1}{2}b(b(\cos(a) \int x^{-n} \sin(bx^n + \frac{\pi}{2}) dx^n - \sin(a) \int x^{-n} \sin(bx^n) dx^n) - x^{-n} \sin(a + bx^n)) - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n}$$

↓ 3780

$$\frac{-\frac{1}{2}b(b(\cos(a) \int x^{-n} \sin(bx^n + \frac{\pi}{2}) dx^n - \sin(a) \text{Si}(bx^n)) - x^{-n} \sin(a + bx^n)) - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n}$$

↓ 3783

$$\frac{-\frac{1}{2}b(b(\cos(a) \text{CosIntegral}(bx^n) - \sin(a) \text{Si}(bx^n)) - x^{-n} \sin(a + bx^n)) - \frac{1}{2}x^{-2n} \cos(a + bx^n)}{n}$$

input

```
Int[x^(-1 - 2*n)*Cos[a + b*x^n], x]
```

output

```
(-1/2*Cos[a + b*x^n]/x^(2*n) - (b*(-(Sin[a + b*x^n]/x^n) + b*(Cos[a]*CosIntegral[b*x^n] - Sin[a]*SinIntegral[b*x^n]))) / 2) / n
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```


rule 3778 `Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Simp[f/(d*(m + 1)) Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]`

rule 3780 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 3861 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.)^(p_.)*(x_)^(m_.), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*Cos[c + d*x])^p, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]] && (EqQ[p, 1] || EqQ[m, n - 1] || (IntegerQ[p] && GtQ[Simplify[(m + 1)/n], 0]))`

Maple [A] (verified)

Time = 1.29 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.83

method	result
default	$b^2 \left(-\frac{\cos(a+bx^n)x^{-2n}}{2b^2} + \frac{\sin(a+bx^n)x^{-n}}{2b} + \frac{\text{Si}(bx^n)\sin(a)}{2} - \frac{\text{Ci}(bx^n)\cos(a)}{2} \right)$
risch	$-\frac{(ib^2e^{-ia}\pi \text{csgn}(bx^n)x^{2n} - 2ib^2e^{-ia}\text{Si}(bx^n)x^{2n} - b^2e^{-ia}\text{expIntegral}_1(-ibx^n)x^{2n} - b^2e^{-ia}\text{expIntegral}_1(-ibx^n)x^{2n} - 2x^n \sin(a + \dots))}{4n}$
meijerg	$b^2\sqrt{\pi} \left(-\frac{x^2 \left(\frac{-1-2n}{2n} + \frac{1}{2n} \right) {}_2F_2 \left(\frac{-1-2n}{n}, -\frac{1}{n} \right) + (-1)^{-\frac{-1-2n}{2n} - \frac{1}{2n}} \left(-\Psi \left(1 - \frac{-1-2n}{2n} - \frac{1}{2n} \right) - \Psi \left(\frac{1}{2} - \frac{-1-2n}{2n} - \frac{1}{2n} \right) + 2n \ln(x) - 2 \ln(2) + \ln(b^2) \right) \sqrt{2}}{2\sqrt{\pi} \Gamma \left(-\frac{-1-2n}{n} - \frac{1}{n} \right)}$

input `int(x^(-1-2*n)*cos(a+b*x^n),x,method=_RETURNVERBOSE)`

output `1/n*b^2*(-1/2*cos(a+b*x^n)/b^2/(x^n)^2+1/2*sin(a+b*x^n)/b/(x^n)+1/2*Si(b*x^n)*sin(a)-1/2*Ci(b*x^n)*cos(a))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

$$\int x^{-1-2n} \cos(a + bx^n) dx$$

$$= -\frac{b^2 x^{2n} \cos(a) \operatorname{Ci}(bx^n) - b^2 x^{2n} \sin(a) \operatorname{Si}(bx^n) - bx^n \sin(bx^n + a) + \cos(bx^n + a)}{2nx^{2n}}$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n),x, algorithm="fricas")`

output `-1/2*(b^2*x^(2*n)*cos(a)*cos_integral(b*x^n) - b^2*x^(2*n)*sin(a)*sin_integral(b*x^n) - b*x^n*sin(b*x^n + a) + cos(b*x^n + a))/(n*x^(2*n))`

Sympy [F]

$$\int x^{-1-2n} \cos(a + bx^n) dx = \int x^{-2n-1} \cos(a + bx^n) dx$$

input `integrate(x**(-1-2*n)*cos(a+b*x**n),x)`

output `Integral(x**(-2*n - 1)*cos(a + b*x**n), x)`

Maxima [F]

$$\int x^{-1-2n} \cos(a + bx^n) dx = \int x^{-2n-1} \cos(bx^n + a) dx$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n),x, algorithm="maxima")`

output `integrate(x^(-2*n - 1)*cos(b*x^n + a), x)`

Giac [F]

$$\int x^{-1-2n} \cos(a + bx^n) dx = \int x^{-2n-1} \cos(bx^n + a) dx$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n),x, algorithm="giac")`

output `integrate(x^(-2*n - 1)*cos(b*x^n + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{-1-2n} \cos(a + bx^n) dx = \int \frac{\cos(a + bx^n)}{x^{2n+1}} dx$$

input `int(cos(a + b*x^n)/x^(2*n + 1),x)`

output `int(cos(a + b*x^n)/x^(2*n + 1), x)`

Reduce [F]

$$\int x^{-1-2n} \cos(a + bx^n) dx$$

$$= \frac{-\cos(x^n b + a) - x^{2n} \left(\int \frac{\cos(x^n b + a)}{x} dx \right) b^{2n} + x^n \sin(x^n b + a) b}{2x^{2n} n}$$

input `int(x^(-1-2*n)*cos(a+b*x^n),x)`

output `(- cos(x**n*b + a) - x**(2*n)*int(cos(x**n*b + a)/x,x)*b**2*n + x**n*sin(x**n*b + a)*b)/(2*x**(2*n)*n)`

3.83 $\int x^{-1-2n} \cos^2(a + bx^n) dx$

Optimal result	604
Mathematica [A] (verified)	604
Rubi [A] (verified)	605
Maple [A] (verified)	606
Fricas [A] (verification not implemented)	606
Sympy [F]	607
Maxima [F]	607
Giac [F]	607
Mupad [F(-1)]	608
Reduce [F]	608

Optimal result

Integrand size = 18, antiderivative size = 97

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = -\frac{x^{-2n}}{4n} - \frac{x^{-2n} \cos(2a + 2bx^n)}{4n} - \frac{b^2 \cos(2a) \operatorname{CosIntegral}(2bx^n)}{n} + \frac{bx^{-n} \sin(2a + 2bx^n)}{2n} + \frac{b^2 \sin(2a) \operatorname{Si}(2bx^n)}{n}$$

output -1/4/n/(x^(2*n))-1/4*cos(2*a+2*b*x^n)/n/(x^(2*n))-b^2*cos(2*a)*Ci(2*b*x^n)/n+1/2*b*sin(2*a+2*b*x^n)/n/(x^n)+b^2*sin(2*a)*Si(2*b*x^n)/n

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.85

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \frac{x^{-2n}(1 + \cos(2(a + bx^n))) + 4b^2 x^{2n} \cos(2a) \operatorname{CosIntegral}(2bx^n) - 2bx^n \sin(2(a + bx^n)) - 4b^2 x^{2n} \sin(2a) \operatorname{Si}(2bx^n)}{4n}$$

input Integrate[x^(-1 - 2*n)*Cos[a + b*x^n]^2,x]

output

```
-1/4*(1 + Cos[2*(a + b*x^n)] + 4*b^2*x^(2*n)*Cos[2*a]*CosIntegral[2*b*x^n]
- 2*b*x^n*Sin[2*(a + b*x^n)] - 4*b^2*x^(2*n)*Sin[2*a]*SinIntegral[2*b*x^n
])/ (n*x^(2*n))
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.98, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{-2n-1} \cos^2(a + bx^n) dx$$

$$\downarrow \text{3907}$$

$$\int \left(\frac{1}{2} x^{-2n-1} \cos(2a + 2bx^n) + \frac{1}{2} x^{-2n-1} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{b^2 \cos(2a) \operatorname{CosIntegral}(2bx^n)}{n} + \frac{b^2 \sin(2a) \operatorname{Si}(2bx^n)}{n} + \frac{bx^{-n} \sin(2(a + bx^n))}{2n} - \frac{x^{-2n} \cos(2(a + bx^n))}{4n} - \frac{x^{-2n}}{4n}$$

input

```
Int[x^(-1 - 2*n)*Cos[a + b*x^n]^2,x]
```

output

```
-1/4*1/(n*x^(2*n)) - Cos[2*(a + b*x^n)]/(4*n*x^(2*n)) - (b^2*Cos[2*a]*CosI
ntegral[2*b*x^n])/n + (b*Sin[2*(a + b*x^n)])/(2*n*x^n) + (b^2*Sin[2*a]*Sin
Integral[2*b*x^n])/n
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3907 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_.)]*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 2.72 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.92

method	result
default	$-\frac{x^{-2n}}{4n} + \frac{2b^2 \left(-\frac{\cos(2a+2bx^n)x^{-2n}}{8b^2} + \frac{\sin(2a+2bx^n)x^{-n}}{4b} + \frac{\text{Si}(2bx^n)\sin(2a)}{2} - \frac{\text{Ci}(2bx^n)\cos(2a)}{2} \right)}{n}$
risch	$\frac{(-2ib^2e^{-2ia}\pi \operatorname{csgn}(bx^n)x^{2n} + 4ib^2e^{-2ia}\operatorname{Si}(2bx^n)x^{2n} + 2b^2e^{-2ia}\operatorname{expIntegral}_1(-2ibx^n)x^{2n} + 2b^2e^{2ia}\operatorname{expIntegral}_1(-2ibx^n)x^{2n} + 2b^2e^{2ia}\pi \operatorname{csgn}(bx^n)x^{2n})}{4n}$

input `int(x^(-1-2*n)*cos(a+b*x^n)^2,x,method=_RETURNVERBOSE)`

output
$$-1/4/(x^n)^{2/n+2/n*b^2}*(-1/8*\cos(2*a+2*b*x^n)/b^2/(x^n)^{2+1/4*\sin(2*a+2*b*x^n)/b/(x^n)+1/2*\operatorname{Si}(2*b*x^n)*\sin(2*a)-1/2*\operatorname{Ci}(2*b*x^n)*\cos(2*a))$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.90

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \frac{2b^2x^{2n}\cos(2a)\operatorname{Ci}(2bx^n) - 2b^2x^{2n}\sin(2a)\operatorname{Si}(2bx^n) - 2bx^n\cos(bx^n + a)\sin(bx^n + a) + \cos(bx^n + a)}{2nx^{2n}}$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n)^2,x, algorithm="fricas")`

output

```
-1/2*(2*b^2*x^(2*n)*cos(2*a)*cos_integral(2*b*x^n) - 2*b^2*x^(2*n)*sin(2*a)
)*sin_integral(2*b*x^n) - 2*b*x^n*cos(b*x^n + a)*sin(b*x^n + a) + cos(b*x^n
+ a)^2)/(n*x^(2*n))
```

Sympy [F]

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \int x^{-2n-1} \cos^2(a + bx^n) dx$$

input

```
integrate(x**(-1-2*n)*cos(a+b*x**n)**2,x)
```

output

```
Integral(x**(-2*n - 1)*cos(a + b*x**n)**2, x)
```

Maxima [F]

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \int x^{-2n-1} \cos(bx^n + a)^2 dx$$

input

```
integrate(x^(-1-2*n)*cos(a+b*x^n)^2,x, algorithm="maxima")
```

output

```
1/4*(2*n*x^(2*n)*integrate(cos(2*b*x^n + 2*a)/(x*x^(2*n)), x) - 1)/(n*x^(2
*n))
```

Giac [F]

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \int x^{-2n-1} \cos(bx^n + a)^2 dx$$

input

```
integrate(x^(-1-2*n)*cos(a+b*x^n)^2,x, algorithm="giac")
```

output

```
integrate(x^(-2*n - 1)*cos(b*x^n + a)^2, x)
```


Mupad [F(-1)]

Timed out.

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \int \frac{\cos(a + bx^n)^2}{x^{2n+1}} dx$$

input `int(cos(a + b*x^n)^2/x^(2*n + 1), x)`output `int(cos(a + b*x^n)^2/x^(2*n + 1), x)`**Reduce [F]**

$$\int x^{-1-2n} \cos^2(a + bx^n) dx = \text{Too large to display}$$

input `int(x^(-1-2*n)*cos(a+b*x^n)^2, x)`

output

```
( - 2*x**n*cos(x**n*b + a)*sin(x**n*b + a)*tan((x**n*b + a)/2)**4*b - 4*x*
**n*cos(x**n*b + a)*sin(x**n*b + a)*tan((x**n*b + a)/2)**2*b - 2*x**n*cos(x
**n*b + a)*sin(x**n*b + a)*b + 8*cos(x**n*b + a)*tan((x**n*b + a)/2)**4 +
16*cos(x**n*b + a)*tan((x**n*b + a)/2)**2 + 8*cos(x**n*b + a) - 32*x**(2*n
)*int(tan((x**n*b + a)/2)/(x**n*tan((x**n*b + a)/2)**4*x + 2*x**n*tan((x**
n*b + a)/2)**2*x + x**n*x),x)*tan((x**n*b + a)/2)**4*b*n - 64*x**(2*n)*int
(tan((x**n*b + a)/2)/(x**n*tan((x**n*b + a)/2)**4*x + 2*x**n*tan((x**n*b +
a)/2)**2*x + x**n*x),x)*tan((x**n*b + a)/2)**2*b*n - 32*x**(2*n)*int(tan(
(x**n*b + a)/2)/(x**n*tan((x**n*b + a)/2)**4*x + 2*x**n*tan((x**n*b + a)/2
)**2*x + x**n*x),x)*b*n + 16*x**(2*n)*int(1/(tan((x**n*b + a)/2)**4*x + 2*
tan((x**n*b + a)/2)**2*x + x),x)*tan((x**n*b + a)/2)**4*b**2*n + 32*x**(2*
n)*int(1/(tan((x**n*b + a)/2)**4*x + 2*tan((x**n*b + a)/2)**2*x + x),x)*ta
n((x**n*b + a)/2)**2*b**2*n + 16*x**(2*n)*int(1/(tan((x**n*b + a)/2)**4*x
+ 2*tan((x**n*b + a)/2)**2*x + x),x)*b**2*n - 6*x**(2*n)*log(x)*tan((x**n*
b + a)/2)**4*b**2*n - 12*x**(2*n)*log(x)*tan((x**n*b + a)/2)**2*b**2*n - 6
*x**(2*n)*log(x)*b**2*n - 8*x**n*sin(x**n*b + a)*tan((x**n*b + a)/2)**4*b
- 16*x**n*sin(x**n*b + a)*tan((x**n*b + a)/2)**2*b - 8*x**n*sin(x**n*b + a
)*b - sin(x**n*b + a)**2*tan((x**n*b + a)/2)**4 - 2*sin(x**n*b + a)**2*tan
((x**n*b + a)/2)**2 - sin(x**n*b + a)**2 + 5*tan((x**n*b + a)/2)**4 + 10*t
an((x**n*b + a)/2)**2 - 11)/(6*x**(2*n)*n*(tan((x**n*b + a)/2)**4 + 2*t...
```

3.84 $\int x^{-1-2n} \cos^3(a + bx^n) dx$

Optimal result	610
Mathematica [A] (verified)	611
Rubi [A] (verified)	611
Maple [A] (verified)	612
Fricas [A] (verification not implemented)	613
Sympy [F]	613
Maxima [F]	613
Giac [F]	614
Mupad [F(-1)]	614
Reduce [F]	614

Optimal result

Integrand size = 18, antiderivative size = 167

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = -\frac{3x^{-2n} \cos(a + bx^n)}{8n} - \frac{x^{-2n} \cos(3a + 3bx^n)}{8n} - \frac{3b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{8n} - \frac{9b^2 \cos(3a) \operatorname{CosIntegral}(3bx^n)}{8n} + \frac{3bx^{-n} \sin(a + bx^n)}{8n} + \frac{3bx^{-n} \sin(3a + 3bx^n)}{8n} + \frac{3b^2 \sin(a) \operatorname{Si}(bx^n)}{8n} + \frac{9b^2 \sin(3a) \operatorname{Si}(3bx^n)}{8n}$$

output

```
-3/8*cos(a+b*x^n)/n/(x^(2*n))-1/8*cos(3*a+3*b*x^n)/n/(x^(2*n))-3/8*b^2*cos(a)*Ci(b*x^n)/n-9/8*b^2*cos(3*a)*Ci(3*b*x^n)/n+3/8*b*sin(a+b*x^n)/n/(x^n)+3/8*b*sin(3*a+3*b*x^n)/n/(x^n)+3/8*b^2*sin(a)*Si(b*x^n)/n+9/8*b^2*sin(3*a)*Si(3*b*x^n)/n
```

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.84

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \frac{x^{-2n}(3 \cos(a + bx^n) + \cos(3(a + bx^n)) + 3b^2 x^{2n} \cos(a) \operatorname{CosIntegral}(bx^n) + 9b^2 x^{2n} \cos(3a) \operatorname{CosIntegral}(3bx^n) - 3bx^n \sin(a + bx^n) - 3bx^n \sin(3(a + bx^n)) - 3b^2 x^{2n} \sin(a) \operatorname{SinIntegral}(bx^n) - 9b^2 x^{2n} \sin(3a) \operatorname{SinIntegral}(3bx^n))}{n x^{2n}}$$

input `Integrate[x^(-1 - 2*n)*Cos[a + b*x^n]^3,x]`

output `-1/8*(3*Cos[a + b*x^n] + Cos[3*(a + b*x^n)] + 3*b^2*x^(2*n)*Cos[a]*CosIntegral[b*x^n] + 9*b^2*x^(2*n)*Cos[3*a]*CosIntegral[3*b*x^n] - 3*b*x^n*Sin[a + b*x^n] - 3*b*x^n*Sin[3*(a + b*x^n)] - 3*b^2*x^(2*n)*Sin[a]*SinIntegral[b*x^n] - 9*b^2*x^(2*n)*Sin[3*a]*SinIntegral[3*b*x^n])/(n*x^(2*n))`

Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.99, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3907, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{-2n-1} \cos^3(a + bx^n) dx$$

$$\downarrow \text{3907}$$

$$\int \left(\frac{3}{4} x^{-2n-1} \cos(a + bx^n) + \frac{1}{4} x^{-2n-1} \cos(3a + 3bx^n) \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{-\frac{3b^2 \cos(a) \operatorname{CosIntegral}(bx^n)}{8n} - \frac{9b^2 \cos(3a) \operatorname{CosIntegral}(3bx^n)}{8n} + \frac{3b^2 \sin(a) \operatorname{Si}(bx^n)}{8n} + \frac{9b^2 \sin(3a) \operatorname{Si}(3bx^n)}{8n} + \frac{3bx^{-n} \sin(a + bx^n)}{8n} + \frac{3bx^{-n} \sin(3(a + bx^n))}{8n} - \frac{3x^{-2n} \cos(a + bx^n)}{8n} - \frac{x^{-2n} \cos(3(a + bx^n))}{8n}}{8n}$$

input `Int[x^(-1 - 2*n)*Cos[a + b*x^n]^3,x]`

output
$$\begin{aligned} & (-3*\text{Cos}[a + b*x^n])/(8*n*x^{(2*n)}) - \text{Cos}[3*(a + b*x^n)]/(8*n*x^{(2*n)}) - (3* \\ & b^2*\text{Cos}[a]*\text{CosIntegral}[b*x^n])/(8*n) - (9*b^2*\text{Cos}[3*a]*\text{CosIntegral}[3*b*x^n] \\ &)/(8*n) + (3*b*\text{Sin}[a + b*x^n])/(8*n*x^n) + (3*b*\text{Sin}[3*(a + b*x^n)]/(8*n* \\ & x^n) + (3*b^2*\text{Sin}[a]*\text{SinIntegral}[b*x^n])/(8*n) + (9*b^2*\text{Sin}[3*a]*\text{SinIntegr} \\ & \text{al}[3*b*x^n])/(8*n) \end{aligned}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3907 `Int[((a_.) + Cos[(c_.) + (d_.)*(x_)^(n_)])*(b_.))^(p_)*((e_.)*(x_)^(m_.), x_Symbol] := Int[ExpandTrigReduce[(e*x)^m, (a + b*Cos[c + d*x^n])^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && IGtQ[p, 0]`

Maple [A] (verified)

Time = 9.92 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.86

method	result
default	$\frac{3b^2 \left(-\frac{\cos(a+bx^n)x^{-2n}}{2b^2} + \frac{\sin(a+bx^n)x^{-n}}{2b} + \frac{\text{Si}(bx^n)\sin(a)}{2} - \frac{\text{Ci}(bx^n)\cos(a)}{2} \right)}{4n} + \frac{9b^2 \left(-\frac{\cos(3a+3bx^n)x^{-2n}}{18b^2} + \frac{\sin(3a+3bx^n)x^{-n}}{6b} + \frac{\text{Si}(3bx^n)\sin(3a)}{2} - \frac{\text{Ci}(3bx^n)\cos(3a)}{2} \right)}{4n}$
risch	$-\frac{(9ib^2e^{-3ia}\pi \text{csgn}(bx^n)x^{2n} + 3ib^2e^{-ia}\pi \text{csgn}(bx^n)x^{2n} - 18ib^2e^{-3ia}\text{Si}(3bx^n)x^{2n} - 6ib^2e^{-ia}\text{Si}(bx^n)x^{2n} - 3b^2e^{ia}\text{expIntegral}_1(-bx^n))}{4n}$

input `int(x^(-1-2*n)*cos(a+b*x^n)^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 3/4/n*b^2*(-1/2*\text{cos}(a+b*x^n)/b^2/(x^n)^2+1/2*\text{sin}(a+b*x^n)/b/(x^n)+1/2*\text{Si}(b \\ & *x^n)*\text{sin}(a)-1/2*\text{Ci}(b*x^n)*\text{cos}(a))+9/4/n*b^2*(-1/18*\text{cos}(3*a+3*b*x^n)/b^2/(\\ & x^n)^2+1/6*\text{sin}(3*a+3*b*x^n)/b/(x^n)+1/2*\text{Si}(3*b*x^n)*\text{sin}(3*a)-1/2*\text{Ci}(3*b*x \\ & n)*\text{cos}(3*a)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.76

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \frac{9b^2x^{2n} \cos(3a) \operatorname{Ci}(3bx^n) + 3b^2x^{2n} \cos(a) \operatorname{Ci}(bx^n) - 12bx^n \cos(bx^n + a)^2 \sin(bx^n + a) - 9b^2x^{2n} \sin(3a) \operatorname{Si}(3bx^n) - 3b^2x^{2n} \sin(a) \operatorname{Si}(bx^n) + 4 \cos(bx^n + a)^3}{8nx^{2n}}$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n)^3,x, algorithm="fricas")`

output `-1/8*(9*b^2*x^(2*n)*cos(3*a)*cos_integral(3*b*x^n) + 3*b^2*x^(2*n)*cos(a)*cos_integral(b*x^n) - 12*b*x^n*cos(b*x^n + a)^2*sin(b*x^n + a) - 9*b^2*x^(2*n)*sin(3*a)*sin_integral(3*b*x^n) - 3*b^2*x^(2*n)*sin(a)*sin_integral(b*x^n) + 4*cos(b*x^n + a)^3)/(n*x^(2*n))`

Sympy [F]

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \int x^{-2n-1} \cos^3(a + bx^n) dx$$

input `integrate(x**(-1-2*n)*cos(a+b*x**n)**3,x)`

output `Integral(x**(-2*n - 1)*cos(a + b*x**n)**3, x)`

Maxima [F]

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \int x^{-2n-1} \cos(bx^n + a)^3 dx$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n)^3,x, algorithm="maxima")`

output `integrate(x^(-2*n - 1)*cos(b*x^n + a)^3, x)`

Giac [F]

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \int x^{-2n-1} \cos(bx^n + a)^3 dx$$

input `integrate(x^(-1-2*n)*cos(a+b*x^n)^3,x, algorithm="giac")`

output `integrate(x^(-2*n - 1)*cos(b*x^n + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \int \frac{\cos(a + bx^n)^3}{x^{2n+1}} dx$$

input `int(cos(a + b*x^n)^3/x^(2*n + 1),x)`

output `int(cos(a + b*x^n)^3/x^(2*n + 1), x)`

Reduce [F]

$$\int x^{-1-2n} \cos^3(a + bx^n) dx = \text{too large to display}$$

input `int(x^(-1-2*n)*cos(a+b*x^n)^3,x)`

output

```
(54*x**n*cos(x**n*b + a)*sin(x**n*b + a)*tan((x**n*b + a)/2)**6*b + 162*x*
*n*cos(x**n*b + a)*sin(x**n*b + a)*tan((x**n*b + a)/2)**4*b + 162*x**n*cos
(x**n*b + a)*sin(x**n*b + a)*tan((x**n*b + a)/2)**2*b + 54*x**n*cos(x**n*b
+ a)*sin(x**n*b + a)*b + 4*cos(x**n*b + a)*sin(x**n*b + a)**2*tan((x**n*b
+ a)/2)**6 + 12*cos(x**n*b + a)*sin(x**n*b + a)**2*tan((x**n*b + a)/2)**4
+ 12*cos(x**n*b + a)*sin(x**n*b + a)**2*tan((x**n*b + a)/2)**2 + 4*cos(x*
*n*b + a)*sin(x**n*b + a)**2 - 136*cos(x**n*b + a)*tan((x**n*b + a)/2)**6
- 408*cos(x**n*b + a)*tan((x**n*b + a)/2)**4 - 408*cos(x**n*b + a)*tan((x*
*n*b + a)/2)**2 - 136*cos(x**n*b + a) + 840*x**(2*n)*int(tan((x**n*b + a)/
2)**3/(x**n*tan((x**n*b + a)/2)**6*x + 3*x**n*tan((x**n*b + a)/2)**4*x + 3
*x**n*tan((x**n*b + a)/2)**2*x + x**n*x),x)*tan((x**n*b + a)/2)**6*b*n + 2
520*x**(2*n)*int(tan((x**n*b + a)/2)**3/(x**n*tan((x**n*b + a)/2)**6*x + 3
*x**n*tan((x**n*b + a)/2)**4*x + 3*x**n*tan((x**n*b + a)/2)**2*x + x**n*x)
,x)*tan((x**n*b + a)/2)**4*b*n + 2520*x**(2*n)*int(tan((x**n*b + a)/2)**3/
(x**n*tan((x**n*b + a)/2)**6*x + 3*x**n*tan((x**n*b + a)/2)**4*x + 3*x**n*
tan((x**n*b + a)/2)**2*x + x**n*x),x)*tan((x**n*b + a)/2)**2*b*n + 840*x**
(2*n)*int(tan((x**n*b + a)/2)**3/(x**n*tan((x**n*b + a)/2)**6*x + 3*x**n*t
an((x**n*b + a)/2)**4*x + 3*x**n*tan((x**n*b + a)/2)**2*x + x**n*x),x)*b*n
- 288*x**(2*n)*int(1/(tan((x**n*b + a)/2)**6*x + 3*tan((x**n*b + a)/2)**4
*x + 3*tan((x**n*b + a)/2)**2*x + x),x)*tan((x**n*b + a)/2)**6*b**2*n - ...
```


3.85 $\int x^2 \cos((a + bx)^2) dx$

Optimal result	616
Mathematica [A] (verified)	616
Rubi [A] (verified)	617
Maple [A] (verified)	618
Fricas [A] (verification not implemented)	619
Sympy [F]	619
Maxima [C] (verification not implemented)	620
Giac [C] (verification not implemented)	620
Mupad [B] (verification not implemented)	621
Reduce [F]	621

Optimal result

Integrand size = 12, antiderivative size = 99

$$\int x^2 \cos((a+bx)^2) dx = \frac{a^2 \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a+bx)\right)}{b^3} - \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelS}\left(\sqrt{\frac{2}{\pi}}(a+bx)\right)}{2b^3} - \frac{a \sin((a+bx)^2)}{b^3} + \frac{(a+bx) \sin((a+bx)^2)}{2b^3}$$

output

```
1/2*a^2*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(b*x+a))/b^3-1/4*2^(1/2)
)*Pi^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)*(b*x+a))/b^3-a*sin((b*x+a)^2)/b^3+1/2
*(b*x+a)*sin((b*x+a)^2)/b^3
```

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.77

$$\int x^2 \cos((a + bx)^2) dx = \frac{-2a^2 \sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) + \sqrt{2\pi} \operatorname{FresnelS}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) + 2(a - bx) \sin((a + bx)^2)}{4b^3}$$

input

```
Integrate[x^2*Cos[(a + b*x)^2],x]
```

output

$$\frac{-1/4*(-2*a^2*\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)] + \text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)] + 2*(a - b*x)*\text{Sin}[(a + b*x)^2])/b^3$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.92, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos((a + bx)^2) dx$$

$$\downarrow \text{3915}$$

$$\frac{\int (\cos((a + bx)^2) a^2 - 2(a + bx) \cos((a + bx)^2) a + (a + bx)^2 \cos((a + bx)^2)) d(a + bx)}{b^3}$$

$$\downarrow \text{2009}$$

$$\frac{\sqrt{\frac{\pi}{2}} a^2 \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) - a \sin((a + bx)^2) + \frac{1}{2}(a + bx) \sin((a + bx)^2)}{b^3}$$

input

$$\text{Int}[x^2*\text{Cos}[(a + b*x)^2], x]$$

output

$$\frac{(a^2*\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)] - (\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*(a + b*x)]))/2 - a*\text{Sin}[(a + b*x)^2] + ((a + b*x)*\text{Sin}[(a + b*x)^2])/2}{b^3}$$

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3915 Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_.)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Module[{k = If[FractionQ[n], Denominator[n], 1]}, Simp[k/f^(m + 1) Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x^(k*n)]]^p, x^(k - 1)*(f*g - e*h + h*x^k)^m, x], x], x, (e + f*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && IGtQ[p, 0] && IGtQ[m, 0]
```

Maple [A] (verified)

Time = 1.56 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.32

method	result
default	$\frac{x \sin(x^2 b^2 + 2abx + a^2)}{2b^2} - \frac{a \left(\frac{\sin(x^2 b^2 + 2abx + a^2)}{2b^2} - \frac{a \sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(b^2 x + ab)}{\sqrt{\pi} \sqrt{b^2}}\right)}{2b \sqrt{b^2}} \right)}{b} - \frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2}(b^2 x + ab)}{\sqrt{\pi} \sqrt{b^2}}\right)}{4b^2 \sqrt{b^2}}$
risch	$-\frac{a^2 (-1)^{\frac{3}{4}} \sqrt{\pi} \operatorname{erf}\left(b(-1)^{\frac{1}{4}} x + (-1)^{\frac{1}{4}} a\right)}{4b^3} - \frac{(-1)^{\frac{1}{4}} \sqrt{\pi} \operatorname{erf}\left(b(-1)^{\frac{1}{4}} x + (-1)^{\frac{1}{4}} a\right)}{8b^3} - \frac{a^2 \sqrt{\pi} \operatorname{erf}\left(-b\sqrt{-i} x + \frac{ia}{\sqrt{-i}}\right)}{4b^3 \sqrt{-i}} - \frac{i \sqrt{\pi} \operatorname{erf}\left(-b\sqrt{-i} x + \frac{ia}{\sqrt{-i}}\right)}{8b^3}$
parts	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(b^2 x + ab)}{\sqrt{\pi} \sqrt{b^2}}\right) x^2}{2\sqrt{b^2}} - \left(\frac{\sqrt{2} \pi^{\frac{3}{2}} \operatorname{FresnelC}\left(\frac{\sqrt{2} b^2 x}{\sqrt{\pi} \sqrt{b^2}} + \frac{\sqrt{2} ab}{\sqrt{\pi} \sqrt{b^2}}\right) \operatorname{csgn}(b) \left(\frac{\operatorname{csgn}(b) \sqrt{\pi} \left(\frac{\sqrt{2} b^2 x}{\sqrt{\pi} \sqrt{b^2}} + \frac{\sqrt{2} ab}{\sqrt{\pi} \sqrt{b^2}}\right)^2}{2} - \sqrt{2} a \left(\frac{\sqrt{2} b^2 x}{\sqrt{\pi} \sqrt{b^2}} + \frac{\sqrt{2} ab}{\sqrt{\pi} \sqrt{b^2}}\right)\right)}{\sqrt{\pi}} \right)$

```
input int(x^2*cos((b*x+a)^2), x, method=_RETURNVERBOSE)
```

output

```
1/2/b^2*x*sin(b^2*x^2+2*a*b*x+a^2)-a/b*(1/2/b^2*sin(b^2*x^2+2*a*b*x+a^2)-1/2*a/b*2^(1/2)*Pi^(1/2)/(b^2)^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)/(b^2)^(1/2)*(b^2*x+a*b))-1/4/b^2*2^(1/2)*Pi^(1/2)/(b^2)^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)/(b^2)^(1/2)*(b^2*x+a*b))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.13

$$\int x^2 \cos((a + bx)^2) dx$$

$$= \frac{2\sqrt{2}\pi a^2 \sqrt{\frac{b^2}{\pi}} C\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right) - \sqrt{2}\pi \sqrt{\frac{b^2}{\pi}} S\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right) + 2(b^2x - ab) \sin(b^2x^2 + 2abx + a^2)}{4b^4}$$

input

```
integrate(x^2*cos((b*x+a)^2),x, algorithm="fricas")
```

output

```
1/4*(2*sqrt(2)*pi*a^2*sqrt(b^2/pi)*fresnel_cos(sqrt(2)*(b*x + a)*sqrt(b^2/pi)/b) - sqrt(2)*pi*sqrt(b^2/pi)*fresnel_sin(sqrt(2)*(b*x + a)*sqrt(b^2/pi)/b) + 2*(b^2*x - a*b)*sin(b^2*x^2 + 2*a*b*x + a^2))/b^4
```

Sympy [F]

$$\int x^2 \cos((a + bx)^2) dx = \int x^2 \cos(a^2 + 2abx + b^2x^2) dx$$

input

```
integrate(x**2*cos((b*x+a)**2),x)
```

output

```
Integral(x**2*cos(a**2 + 2*a*b*x + b**2*x**2), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.61

$$\int x^2 \cos((a + bx)^2) dx = \frac{4 abx \left(-i e^{(i b^2 x^2 + 2i abx + i a^2)} + i e^{(-i b^2 x^2 - 2i abx - i a^2)} \right) + 4 a^2 \left(-i e^{(i b^2 x^2 + 2i abx + i a^2)} + i e^{(-i b^2 x^2 - 2i abx - i a^2)} \right)}{1}$$

input `integrate(x^2*cos((b*x+a)^2),x, algorithm="maxima")`

output

```
-1/8*(4*a*b*x*(-I*e^(I*b^2*x^2 + 2*I*a*b*x + I*a^2) + I*e^(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)) + 4*a^2*(-I*e^(I*b^2*x^2 + 2*I*a*b*x + I*a^2) + I*e^(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)) - sqrt(b^2*x^2 + 2*a*b*x + a^2)*((-I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(I*b^2*x^2 + 2*I*a*b*x + I*a^2)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)) - 1))*a^2 + (I + 1)*sqrt(2)*gamma(3/2, I*b^2*x^2 + 2*I*a*b*x + I*a^2) - (I - 1)*sqrt(2)*gamma(3/2, -I*b^2*x^2 - 2*I*a*b*x - I*a^2))/(b^4*x + a*b^3)
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.61

$$\int x^2 \cos((a + bx)^2) dx = -\frac{\frac{(i+1)\sqrt{2}\sqrt{\pi}(2a^2+i)\operatorname{erf}\left(\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\left(x+\frac{a}{b}\right)|b\right)}{|b|} + \frac{4(i b(x+\frac{a}{b})-2i a)e^{(i b^2 x^2+2i abx+i a^2)}}{b}}{16 b^2} - \frac{\frac{(i-1)\sqrt{2}\sqrt{\pi}(2a^2-i)\operatorname{erf}\left(-\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\left(x+\frac{a}{b}\right)|b\right)}{|b|} + \frac{4(-i b(x+\frac{a}{b})+2i a)e^{(-i b^2 x^2-2i abx-i a^2)}}{b}}{16 b^2}$$

input `integrate(x^2*cos((b*x+a)^2),x, algorithm="giac")`

output

```
-1/16*((I + 1)*sqrt(2)*sqrt(pi)*(2*a^2 + I)*erf((1/2*I - 1/2)*sqrt(2)*(x +
a/b)*abs(b))/abs(b) + 4*(I*b*(x + a/b) - 2*I*a)*e^(I*b^2*x^2 + 2*I*a*b*x
+ I*a^2)/b)/b^2 - 1/16*(-(I - 1)*sqrt(2)*sqrt(pi)*(2*a^2 - I)*erf(-(1/2*I
+ 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b) + 4*(-I*b*(x + a/b) + 2*I*a)*e^(-I
*b^2*x^2 - 2*I*a*b*x - I*a^2)/b)/b^2
```

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.81

$$\int x^2 \cos((a + bx)^2) dx = \frac{x \sin((a + bx)^2)}{2b^2} - \frac{a \sin((a + bx)^2)}{2b^3} - \frac{\sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}(a+bx)}{\sqrt{\pi}}\right)}{4b^3} + \frac{\sqrt{2} a^2 \sqrt{\pi} C\left(\frac{\sqrt{2}(a+bx)}{\sqrt{\pi}}\right)}{2b^3}$$

input

```
int(x^2*cos((a + b*x)^2),x)
```

output

```
(x*sin((a + b*x)^2))/(2*b^2) - (a*sin((a + b*x)^2))/(2*b^3) - (2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*(a + b*x))/pi^(1/2)))/(4*b^3) + (2^(1/2)*a^2*pi^(1/2)*fresnelc((2^(1/2)*(a + b*x))/pi^(1/2)))/(2*b^3)
```

Reduce [F]

$$\int x^2 \cos((a + bx)^2) dx = \int \cos(b^2x^2 + 2abx + a^2) x^2 dx$$

input

```
int(x^2*cos((b*x+a)^2),x)
```

output

```
int(cos(a**2 + 2*a*b*x + b**2*x**2)*x**2,x)
```

3.86 $\int x \cos((a + bx)^2) dx$

Optimal result	622
Mathematica [A] (verified)	622
Rubi [A] (verified)	623
Maple [A] (verified)	624
Fricas [A] (verification not implemented)	624
Sympy [F]	625
Maxima [C] (verification not implemented)	625
Giac [C] (verification not implemented)	626
Mupad [B] (verification not implemented)	626
Reduce [F]	627

Optimal result

Integrand size = 10, antiderivative size = 47

$$\int x \cos((a + bx)^2) dx = -\frac{a\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^2} + \frac{\sin((a + bx)^2)}{2b^2}$$

output

$$-1/2*a*2^{(1/2)}*Pi^{(1/2)}*FresnelC(2^{(1/2)}/Pi^{(1/2)}*(b*x+a))/b^2+1/2*\sin((b*x+a)^2)/b^2$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

$$\int x \cos((a + bx)^2) dx = \frac{-a\sqrt{2\pi} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right) + \sin((a + bx)^2)}{2b^2}$$

input

`Integrate[x*Cos[(a + b*x)^2],x]`

output

$$(-(a*\sqrt{2*Pi}*FresnelC[\sqrt{2/Pi}*(a + b*x)]) + \sin[(a + b*x)^2])/(2*b^2)$$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3915, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \cos((a + bx)^2) dx$$

$$\downarrow \text{3915}$$

$$\frac{\int ((a + bx) \cos((a + bx)^2) - a \cos((a + bx)^2)) d(a + bx)}{b^2}$$

$$\downarrow \text{2009}$$

$$\frac{\frac{1}{2} \sin((a + bx)^2) - \sqrt{\frac{\pi}{2}} a \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b^2}$$

input `Int[x*Cos[(a + b*x)^2],x]`

output `(-(a*Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*(a + b*x)]) + Sin[(a + b*x)^2]/2)/b^2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3915 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Module[{k = If[FractionQ[n], Denominator[n], 1]}, Simp[k/f^(m + 1) Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x^(k*n)])^p, x^(k - 1)*(f*g - e*h + h*x^k)^m, x], x], x, (e + f*x)^(1/k)], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && IGtQ[p, 0] && IGtQ[m, 0]`

Maple [A] (verified)

Time = 1.42 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.34

method	result
default	$\frac{\sin(x^2b^2+2abx+a^2)}{2b^2} - \frac{a\sqrt{2}\sqrt{\pi}\operatorname{FresnelC}\left(\frac{\sqrt{2}(b^2x+ab)}{\sqrt{\pi}\sqrt{b^2}}\right)}{2b\sqrt{b^2}}$
risch	$\frac{(-1)^{\frac{3}{4}}a\sqrt{\pi}\operatorname{erf}\left(b(-1)^{\frac{1}{4}}x+(-1)^{\frac{1}{4}}a\right)}{4b^2} + \frac{a\sqrt{\pi}\operatorname{erf}\left(-b\sqrt{-i}x+\frac{ia}{\sqrt{-i}}\right)}{4b^2\sqrt{-i}} + \frac{\sin((bx+a)^2)}{2b^2}$
parts	$\frac{\sqrt{2}\sqrt{\pi}\operatorname{FresnelC}\left(\frac{\sqrt{2}(b^2x+ab)}{\sqrt{\pi}\sqrt{b^2}}\right)x}{2\sqrt{b^2}} - \frac{\pi\left(\operatorname{FresnelC}\left(\frac{\sqrt{2}b^2x}{\sqrt{\pi}\sqrt{b^2}}+\frac{\sqrt{2}ab}{\sqrt{\pi}\sqrt{b^2}}\right)\left(\frac{\sqrt{2}b^2x}{\sqrt{\pi}\sqrt{b^2}}+\frac{\sqrt{2}ab}{\sqrt{\pi}\sqrt{b^2}}\right) - \frac{\sin\left(\frac{\pi\left(\frac{\sqrt{2}b^2x}{\sqrt{\pi}\sqrt{b^2}}+\frac{\sqrt{2}ab}{\sqrt{\pi}\sqrt{b^2}}\right)^2}{2}\right)}{\pi}\right)}{2b^2}$

input `int(x*cos((b*x+a)^2),x,method=_RETURNVERBOSE)`

output `1/2/b^2*sin(b^2*x^2+2*a*b*x+a^2)-1/2*a/b*2^(1/2)*Pi^(1/2)/(b^2)^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)/(b^2)^(1/2)*(b^2*x+a*b))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.34

$$\int x \cos((a + bx)^2) dx = -\frac{\sqrt{2}\pi a \sqrt{\frac{b^2}{\pi}} C\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right) - b \sin(b^2x^2 + 2abx + a^2)}{2b^3}$$

input `integrate(x*cos((b*x+a)^2),x, algorithm="fricas")`

output `-1/2*(sqrt(2)*pi*a*sqrt(b^2/pi)*fresnel_cos(sqrt(2)*(b*x + a)*sqrt(b^2/pi)/b) - b*sin(b^2*x^2 + 2*a*b*x + a^2))/b^3`

Sympy [F]

$$\int x \cos((a + bx)^2) dx = \int x \cos(a^2 + 2abx + b^2x^2) dx$$

input `integrate(x*cos((b*x+a)**2),x)`

output `Integral(x*cos(a**2 + 2*a*b*x + b**2*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 199, normalized size of antiderivative = 4.23

$$\int x \cos((a + bx)^2) dx$$

$$= \frac{2bx \left(-i e^{(ib^2x^2 + 2iabx + ia^2)} + i e^{(-ib^2x^2 - 2iabx - ia^2)} \right) - \sqrt{b^2x^2 + 2abx + a^2} (-(i-1) \sqrt{2} \sqrt{\pi} (\operatorname{erf}(\sqrt{i} b^2x^2 +$$

input `integrate(x*cos((b*x+a)^2),x, algorithm="maxima")`

output `1/8*(2*b*x*(-I*e^(I*b^2*x^2 + 2*I*a*b*x + I*a^2) + I*e^(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)) - sqrt(b^2*x^2 + 2*a*b*x + a^2)*(-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(I*b^2*x^2 + 2*I*a*b*x + I*a^2)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)) - 1))*a + 2*a*(-I*e^(I*b^2*x^2 + 2*I*a*b*x + I*a^2) + I*e^(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)))/(b^3*x + a*b^2)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.53

$$\int x \cos((a + bx)^2) dx = -\frac{\frac{(i+1)\sqrt{2}\sqrt{\pi}a \operatorname{erf}\left(\frac{1}{2}i - \frac{1}{2}\right)\sqrt{2}\left(x + \frac{a}{b}\right)|b|}{|b|} + \frac{2i e^{(i b^2 x^2 + 2i a b x + i a^2)}}{b}}{8b} - \frac{\frac{(i-1)\sqrt{2}\sqrt{\pi}a \operatorname{erf}\left(-\frac{1}{2}i + \frac{1}{2}\right)\sqrt{2}\left(x + \frac{a}{b}\right)|b|}{|b|} - \frac{2i e^{(-i b^2 x^2 - 2i a b x - i a^2)}}{b}}{8b}$$

input `integrate(x*cos((b*x+a)^2),x, algorithm="giac")`

output `-1/8*(-(I + 1)*sqrt(2)*sqrt(pi)*a*erf((1/2*I - 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b) + 2*I*e^(I*b^2*x^2 + 2*I*a*b*x + I*a^2)/b)/b - 1/8*((I - 1)*sqrt(2)*sqrt(pi)*a*erf(-(1/2*I + 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b) - 2*I*e^(-I*b^2*x^2 - 2*I*a*b*x - I*a^2)/b)/b`

Mupad [B] (verification not implemented)

Time = 40.34 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int x \cos((a + bx)^2) dx = \frac{\sin((a + bx)^2)}{2b^2} - \frac{\sqrt{2}a\sqrt{\pi}C\left(\frac{\sqrt{2}(a+bx)}{\sqrt{\pi}}\right)}{2b^2}$$

input `int(x*cos((a + b*x)^2),x)`

output `sin((a + b*x)^2)/(2*b^2) - (2^(1/2)*a*pi^(1/2)*fresnelc((2^(1/2)*(a + b*x))/pi^(1/2)))/(2*b^2)`

Reduce [F]

$$\int x \cos((a + bx)^2) dx = \frac{-2(\int \cos(b^2x^2 + 2abx + a^2) dx) ab + \sin(b^2x^2 + 2abx + a^2)}{2b^2}$$

input `int(x*cos((b*x+a)^2),x)`

output `(- 2*int(cos(a**2 + 2*a*b*x + b**2*x**2),x)*a*b + sin(a**2 + 2*a*b*x + b**2*x**2))/(2*b**2)`

3.87 $\int \cos((a + bx)^2) dx$

Optimal result	628
Mathematica [A] (verified)	628
Rubi [A] (verified)	629
Maple [A] (verified)	629
Fricas [A] (verification not implemented)	630
Sympy [F]	630
Maxima [C] (verification not implemented)	631
Giac [C] (verification not implemented)	631
Mupad [B] (verification not implemented)	632
Reduce [F]	632

Optimal result

Integrand size = 8, antiderivative size = 29

$$\int \cos((a + bx)^2) dx = \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

output

```
1/2*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(b*x+a))/b
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \cos((a + bx)^2) dx = \frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

input

```
Integrate[Cos[(a + b*x)^2],x]
```

output

```
(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*(a + b*x)])/b
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos((a + bx)^2) dx$$

↓ 3833

$$\frac{\sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\sqrt{\frac{2}{\pi}}(a + bx)\right)}{b}$$

input `Int[Cos[(a + b*x)^2], x]`

output `(Sqrt[Pi/2]*FresnelC[Sqrt[2/Pi]*(a + b*x)])/b`

Defintions of rubi rules used

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.)*(x_)) ^2], x_Symbol] := Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.24

method	result	size
default	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(b^2 x + ab)}{\sqrt{\pi} \sqrt{b^2}}\right)}{2\sqrt{b^2}}$	36
risch	$-\frac{\sqrt{\pi}(-1)^{\frac{3}{4}} \operatorname{erf}\left(b(-1)^{\frac{1}{4}} x + (-1)^{\frac{1}{4}} a\right)}{4b} - \frac{\sqrt{\pi} \operatorname{erf}\left(-b\sqrt{-i} x + \frac{ia}{\sqrt{-i}}\right)}{4b\sqrt{-i}}$	56

input `int(cos((b*x+a)^2),x,method=_RETURNVERBOSE)`

output `1/2*2^(1/2)*Pi^(1/2)/(b^2)^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)/(b^2)^(1/2)*(b^2*x+a*b))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.38

$$\int \cos((a + bx)^2) dx = \frac{\sqrt{2}\pi \sqrt{\frac{b^2}{\pi}} C\left(\frac{\sqrt{2}(bx+a)\sqrt{\frac{b^2}{\pi}}}{b}\right)}{2b^2}$$

input `integrate(cos((b*x+a)^2),x, algorithm="fricas")`

output `1/2*sqrt(2)*pi*sqrt(b^2/pi)*fresnel_cos(sqrt(2)*(b*x + a)*sqrt(b^2/pi)/b)/b^2`

Sympy [F]

$$\int \cos((a + bx)^2) dx = \int \cos((a + bx)^2) dx$$

input `integrate(cos((b*x+a)**2),x)`

output `Integral(cos((a + b*x)**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.90

$$\int \cos((a + bx)^2) dx = \frac{\sqrt{\pi} \left((i - 1) \sqrt{2} \operatorname{erf} \left(-(-1)^{\frac{3}{4}} (i bx + i a) \right) - (i + 1) \sqrt{2} \operatorname{erf} \left(\left(\frac{1}{2}i + \frac{1}{2} \right) \sqrt{2}(-i bx - i a) \right) + (i - 1) \sqrt{2} \operatorname{erf} \left(\left(\frac{1}{2}i - \frac{1}{2} \right) \sqrt{2}(i bx + i a) \right) - (i + 1) \sqrt{2} \operatorname{erf} \left(\left(\frac{1}{2}i + \frac{1}{2} \right) \sqrt{2}(i bx + i a) \right) \right)}{16b}$$

input `integrate(cos((b*x+a)^2),x, algorithm="maxima")`

output `-1/16*sqrt(pi)*((I - 1)*sqrt(2)*erf(-(-1)^(3/4)*(I*b*x + I*a)) - (I + 1)*sqrt(2)*erf((1/2*I + 1/2)*sqrt(2)*(-I*b*x - I*a)) + (I - 1)*sqrt(2)*erf((1/2*I - 1/2)*sqrt(2)*(-I*b*x - I*a)) + (I + 1)*sqrt(2)*erf((I*b*x + I*a)/sqrt(-I)))/b`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.90

$$\int \cos((a + bx)^2) dx = -\frac{(i + 1) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(\left(\frac{1}{2}i - \frac{1}{2} \right) \sqrt{2} \left(x + \frac{a}{b} \right) |b| \right)}{8 |b|} + \frac{(i - 1) \sqrt{2} \sqrt{\pi} \operatorname{erf} \left(-\left(\frac{1}{2}i + \frac{1}{2} \right) \sqrt{2} \left(x + \frac{a}{b} \right) |b| \right)}{8 |b|}$$

input `integrate(cos((b*x+a)^2),x, algorithm="giac")`

output `-(1/8*I + 1/8)*sqrt(2)*sqrt(pi)*erf((1/2*I - 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b) + (1/8*I - 1/8)*sqrt(2)*sqrt(pi)*erf(-(1/2*I + 1/2)*sqrt(2)*(x + a/b)*abs(b))/abs(b)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

$$\int \cos((a + bx)^2) dx = \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2} b \sqrt{\frac{1}{b^2}} (a + bx)}{\sqrt{\pi}}\right) \sqrt{\frac{1}{b^2}}}{2}$$

input `int(cos((a + b*x)^2), x)`output `(2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*b*(1/b^2)^(1/2)*(a + b*x))/pi^(1/2))*(1/b^2)^(1/2))/2`**Reduce [F]**

$$\int \cos((a + bx)^2) dx = \int \cos(b^2x^2 + 2abx + a^2) dx$$

input `int(cos((b*x+a)^2), x)`output `int(cos(a**2 + 2*a*b*x + b**2*x**2), x)`

$$3.88 \quad \int \frac{\cos((a+bx)^2)}{x} dx$$

Optimal result	633
Mathematica [N/A]	633
Rubi [N/A]	634
Maple [N/A]	634
Fricas [N/A]	635
Sympy [N/A]	635
Maxima [N/A]	636
Giac [N/A]	636
Mupad [N/A]	636
Reduce [N/A]	637

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\cos((a+bx)^2)}{x} dx = \text{Int}\left(\frac{\cos((a+bx)^2)}{x}, x\right)$$

output `Defer(Int)(cos((b*x+a)^2)/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.46 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos((a+bx)^2)}{x} dx$$

input `Integrate[Cos[(a + b*x)^2]/x,x]`

output `Integrate[Cos[(a + b*x)^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3919}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos((a + bx)^2)}{x} dx$$

↓ 3919

$$\int \frac{\cos((a + bx)^2)}{x} dx$$

input `Int[Cos[(a + b*x)^2]/x,x]`

output `$Aborted`

Defintions of rubi rules used

rule 3919 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_.)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] :> Unintegrable[(g + h*x)^m*(a + b*Cos[c + d*(e + f*x)^n])^p, x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p}, x]`

Maple [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\cos((bx + a)^2)}{x} dx$$

input `int(cos((b*x+a)^2)/x,x)`

output `int(cos((b*x+a)^2)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.92

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos((bx+a)^2)}{x} dx$$

input `integrate(cos((b*x+a)^2)/x,x, algorithm="fricas")`

output `integral(cos(b^2*x^2 + 2*a*b*x + a^2)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.95 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos(a^2 + 2abx + b^2x^2)}{x} dx$$

input `integrate(cos((b*x+a)**2)/x,x)`

output `Integral(cos(a**2 + 2*a*b*x + b**2*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos((bx+a)^2)}{x} dx$$

input `integrate(cos((b*x+a)^2)/x,x, algorithm="maxima")`output `integrate(cos((b*x + a)^2)/x, x)`**Giac [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos((bx+a)^2)}{x} dx$$

input `integrate(cos((b*x+a)^2)/x,x, algorithm="giac")`output `integrate(cos((b*x + a)^2)/x, x)`**Mupad [N/A]**

Not integrable

Time = 40.39 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a+bx)^2)}{x} dx = \int \frac{\cos((a+bx)^2)}{x} dx$$

input `int(cos((a + b*x)^2)/x,x)`

output `int(cos((a + b*x)^2)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.92

$$\int \frac{\cos((a + bx)^2)}{x} dx = \int \frac{\cos(b^2x^2 + 2abx + a^2)}{x} dx$$

input `int(cos((b*x+a)^2)/x,x)`

output `int(cos(a**2 + 2*a*b*x + b**2*x**2)/x,x)`

3.89 $\int \frac{\cos((a+bx)^2)}{x^2} dx$

Optimal result	638
Mathematica [N/A]	638
Rubi [N/A]	639
Maple [N/A]	639
Fricas [N/A]	640
Sympy [N/A]	640
Maxima [N/A]	641
Giac [N/A]	641
Mupad [N/A]	641
Reduce [N/A]	642

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\cos((a+bx)^2)}{x^2} dx = \text{Int}\left(\frac{\cos((a+bx)^2)}{x^2}, x\right)$$

output `Defer(Int)(cos((b*x+a)^2)/x^2,x)`

Mathematica [N/A]

Not integrable

Time = 4.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a+bx)^2)}{x^2} dx = \int \frac{\cos((a+bx)^2)}{x^2} dx$$

input `Integrate[Cos[(a + b*x)^2]/x^2,x]`

output `Integrate[Cos[(a + b*x)^2]/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {3919}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos((a + bx)^2)}{x^2} dx$$

↓ 3919

$$\int \frac{\cos((a + bx)^2)}{x^2} dx$$

input `Int[Cos[(a + b*x)^2]/x^2,x]`

output `$Aborted`

Defintions of rubi rules used

rule 3919 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_.)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] :> Unintegrable[(g + h*x)^m*(a + b*Cos[c + d*(e + f*x)^n])^p, x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p}, x]`

Maple [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\cos((bx + a)^2)}{x^2} dx$$

input `int(cos((b*x+a)^2)/x^2,x)`

output `int(cos((b*x+a)^2)/x^2,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.92

$$\int \frac{\cos((a+bx)^2)}{x^2} dx = \int \frac{\cos((bx+a)^2)}{x^2} dx$$

input `integrate(cos((b*x+a)^2)/x^2,x, algorithm="fricas")`

output `integral(cos(b^2*x^2 + 2*a*b*x + a^2)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.95 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.83

$$\int \frac{\cos((a+bx)^2)}{x^2} dx = \int \frac{\cos(a^2 + 2abx + b^2x^2)}{x^2} dx$$

input `integrate(cos((b*x+a)**2)/x**2,x)`

output `Integral(cos(a**2 + 2*a*b*x + b**2*x**2)/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a + bx)^2)}{x^2} dx = \int \frac{\cos((bx + a)^2)}{x^2} dx$$

input `integrate(cos((b*x+a)^2)/x^2,x, algorithm="maxima")`

output `integrate(cos((b*x + a)^2)/x^2, x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a + bx)^2)}{x^2} dx = \int \frac{\cos((bx + a)^2)}{x^2} dx$$

input `integrate(cos((b*x+a)^2)/x^2,x, algorithm="giac")`

output `integrate(cos((b*x + a)^2)/x^2, x)`

Mupad [N/A]

Not integrable

Time = 40.23 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\cos((a + bx)^2)}{x^2} dx = \int \frac{\cos((a + bx)^2)}{x^2} dx$$

input `int(cos((a + b*x)^2)/x^2,x)`

output `int(cos((a + b*x)^2)/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 38, normalized size of antiderivative = 3.17

$$\int \frac{\cos((a + bx)^2)}{x^2} dx = \frac{\left(\int \frac{\cos(b^2x^2 + 2abx + a^2)}{x^2} dx\right) x + \left(\int \frac{1}{x^2} dx\right) x + 1}{x}$$

input `int(cos((b*x+a)^2)/x^2,x)`

output `(int(cos(a**2 + 2*a*b*x + b**2*x**2)/x**2,x)*x + int(1/x**2,x)*x + 1)/x`

3.90 $\int x^2 \cos(a + b\sqrt{c + dx}) dx$

Optimal result	643
Mathematica [C] (verified)	644
Rubi [A] (verified)	644
Maple [B] (verified)	646
Fricas [A] (verification not implemented)	647
Sympy [A] (verification not implemented)	647
Maxima [B] (verification not implemented)	648
Giac [A] (verification not implemented)	648
Mupad [F(-1)]	649
Reduce [B] (verification not implemented)	649

Optimal result

Integrand size = 18, antiderivative size = 346

$$\begin{aligned}
 \int x^2 \cos(a + b\sqrt{c + dx}) dx = & \frac{240 \cos(a + b\sqrt{c + dx})}{b^6 d^3} + \frac{24c \cos(a + b\sqrt{c + dx})}{b^4 d^3} \\
 & + \frac{2c^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} \\
 & - \frac{120(c + dx) \cos(a + b\sqrt{c + dx})}{b^4 d^3} \\
 & - \frac{12c(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^3} \\
 & + \frac{10(c + dx)^2 \cos(a + b\sqrt{c + dx})}{b^2 d^3} \\
 & + \frac{240\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^5 d^3} \\
 & + \frac{24c\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^3 d^3} \\
 & + \frac{2c^2\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd^3} \\
 & - \frac{40(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{b^3 d^3} \\
 & - \frac{4c(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{bd^3} \\
 & + \frac{2(c + dx)^{5/2} \sin(a + b\sqrt{c + dx})}{bd^3}
 \end{aligned}$$

output

$$240*\cos(a+b*(d*x+c)^(1/2))/b^6/d^3+24*c*\cos(a+b*(d*x+c)^(1/2))/b^4/d^3+2*c^2*\cos(a+b*(d*x+c)^(1/2))/b^2/d^3-120*(d*x+c)*\cos(a+b*(d*x+c)^(1/2))/b^4/d^3-12*c*(d*x+c)*\cos(a+b*(d*x+c)^(1/2))/b^2/d^3+10*(d*x+c)^2*\cos(a+b*(d*x+c)^(1/2))/b^2/d^3+240*(d*x+c)^(1/2)*\sin(a+b*(d*x+c)^(1/2))/b^5/d^3+24*c*(d*x+c)^(1/2)*\sin(a+b*(d*x+c)^(1/2))/b^3/d^3+2*c^2*(d*x+c)^(1/2)*\sin(a+b*(d*x+c)^(1/2))/b/d^3-40*(d*x+c)^(3/2)*\sin(a+b*(d*x+c)^(1/2))/b^3/d^3-4*c*(d*x+c)^(3/2)*\sin(a+b*(d*x+c)^(1/2))/b/d^3+2*(d*x+c)^(5/2)*\sin(a+b*(d*x+c)^(1/2))/b/d^3$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.02 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.65

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{e^{-i(a+b\sqrt{c+dx})} \left(120 + 120ib\sqrt{c+dx} + ib^5 d^2 x^2 \sqrt{c+dx} - 4ib^3 \sqrt{c+dx}(2c+5dx) - 12b^2(4c+5dx) + b^4 d \right)}{\dots}$$

input

```
Integrate[x^2*Cos[a + b*Sqrt[c + d*x]],x]
```

output

$$\frac{(120 + (120*I)*b*\text{Sqrt}[c + d*x] + I*b^5*d^2*x^2*\text{Sqrt}[c + d*x] - (4*I)*b^3*\text{Sqrt}[c + d*x]*(2*c + 5*d*x) - 12*b^2*(4*c + 5*d*x) + b^4*d*x*(4*c + 5*d*x) + E^{((2*I)*(a + b*\text{Sqrt}[c + d*x]))}*(120 - (120*I)*b*\text{Sqrt}[c + d*x] - I*b^5*d^2*x^2*\text{Sqrt}[c + d*x] + (4*I)*b^3*\text{Sqrt}[c + d*x]*(2*c + 5*d*x) - 12*b^2*(4*c + 5*d*x) + b^4*d*x*(4*c + 5*d*x)))/(b^6*d^3*E^{(I*(a + b*\text{Sqrt}[c + d*x]))})$$

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3913, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx$$

↓ 3913

$$\frac{2 \int \left(\frac{\cos(a+b\sqrt{c+dx})(c+dx)^{5/2}}{d^2} - \frac{2c \cos(a+b\sqrt{c+dx})(c+dx)^{3/2}}{d^2} + \frac{c^2 \cos(a+b\sqrt{c+dx})\sqrt{c+dx}}{d^2} \right) d\sqrt{c+dx}}{d}$$

↓ 2009

$$\frac{2 \left(\frac{120 \cos(a+b\sqrt{c+dx})}{b^6 d^2} + \frac{120 \sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^5 d^2} - \frac{60(c+dx) \cos(a+b\sqrt{c+dx})}{b^4 d^2} + \frac{12c \cos(a+b\sqrt{c+dx})}{b^4 d^2} - \frac{20(c+dx)^{3/2} \sin(a+b\sqrt{c+dx})}{b^3 d^2} \right)}{d}$$

input `Int[x^2*Cos[a + b*Sqrt[c + d*x]],x]`

output `(2*((120*Cos[a + b*Sqrt[c + d*x]])/(b^6*d^2) + (12*c*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^2) + (c^2*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) - (60*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^4*d^2) - (6*c*(c + d*x)*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) + (5*(c + d*x)^2*Cos[a + b*Sqrt[c + d*x]])/(b^2*d^2) + (120*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^5*d^2) + (12*c*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^2) + (c^2*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]])/(b*d^2) - (20*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b^3*d^2) - (2*c*(c + d*x)^(3/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^2) + ((c + d*x)^(5/2)*Sin[a + b*Sqrt[c + d*x]])/(b*d^2))/d`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3913 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Simp[1/(n*f) Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - e*(h/f) + h*(x^(1/n)/f))]^m, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 824 vs. $2(310) = 620$.

Time = 1.44 (sec) , antiderivative size = 825, normalized size of antiderivative = 2.38

method	result
derivativedivides	$\frac{-2ac^2 \sin(a+\sqrt{dx+c}b)+2c^2(\cos(a+\sqrt{dx+c}b)+(a+\sqrt{dx+c}b)\sin(a+\sqrt{dx+c}b))+\frac{4a^3c \sin(a+\sqrt{dx+c}b)}{b^2}-\frac{12a^2c(\cos(a+\sqrt{dx+c}b)+(a+\sqrt{dx+c}b)\sin(a+\sqrt{dx+c}b))}{b^2}}{b^2}$
default	$\frac{-2ac^2 \sin(a+\sqrt{dx+c}b)+2c^2(\cos(a+\sqrt{dx+c}b)+(a+\sqrt{dx+c}b)\sin(a+\sqrt{dx+c}b))+\frac{4a^3c \sin(a+\sqrt{dx+c}b)}{b^2}-\frac{12a^2c(\cos(a+\sqrt{dx+c}b)+(a+\sqrt{dx+c}b)\sin(a+\sqrt{dx+c}b))}{b^2}}{b^2}$
parts	$\frac{2x^2\sqrt{dx+c} \sin(a+\sqrt{dx+c}b)}{db} + \frac{2x^2 \cos(a+\sqrt{dx+c}b)}{db^2} - \frac{8\left(2ac(\sin(a+\sqrt{dx+c}b)-(a+\sqrt{dx+c}b)\cos(a+\sqrt{dx+c}b))\right)}{db^2}$

input `int(x^2*cos(a+(d*x+c)^(1/2)*b),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 2/d^3/b^2*(-a*c^2*\sin(a+(d*x+c)^(1/2)*b)+c^2*(\cos(a+(d*x+c)^(1/2)*b)+(a+(d*x+c)^(1/2)*b)*\sin(a+(d*x+c)^(1/2)*b))+2/b^2*a^3*c*\sin(a+(d*x+c)^(1/2)*b)- \\ & 6/b^2*a^2*c*(\cos(a+(d*x+c)^(1/2)*b)+(a+(d*x+c)^(1/2)*b)*\sin(a+(d*x+c)^(1/2)*b))+6/b^2*a*c*((a+(d*x+c)^(1/2)*b)^2*\sin(a+(d*x+c)^(1/2)*b)-2*\sin(a+(d*x+c)^(1/2)*b)+2*(a+(d*x+c)^(1/2)*b)*\cos(a+(d*x+c)^(1/2)*b))-2/b^2*c*((a+(d*x+c)^(1/2)*b)^3*\sin(a+(d*x+c)^(1/2)*b)+3*(a+(d*x+c)^(1/2)*b)^2*\cos(a+(d*x+c)^(1/2)*b)-6*\cos(a+(d*x+c)^(1/2)*b)-6*(a+(d*x+c)^(1/2)*b)*\sin(a+(d*x+c)^(1/2)*b))-1/b^4*a^5*\sin(a+(d*x+c)^(1/2)*b)+5/b^4*a^4*(\cos(a+(d*x+c)^(1/2)*b)+(a+(d*x+c)^(1/2)*b)*\sin(a+(d*x+c)^(1/2)*b))-10/b^4*a^3*((a+(d*x+c)^(1/2)*b)^2*\sin(a+(d*x+c)^(1/2)*b)-2*\sin(a+(d*x+c)^(1/2)*b)+2*(a+(d*x+c)^(1/2)*b)*\cos(a+(d*x+c)^(1/2)*b))+10/b^4*a^2*((a+(d*x+c)^(1/2)*b)^3*\sin(a+(d*x+c)^(1/2)*b)+3*(a+(d*x+c)^(1/2)*b)^2*\cos(a+(d*x+c)^(1/2)*b)-6*\cos(a+(d*x+c)^(1/2)*b)-6*(a+(d*x+c)^(1/2)*b)*\sin(a+(d*x+c)^(1/2)*b))-5/b^4*a*((a+(d*x+c)^(1/2)*b)^4*\sin(a+(d*x+c)^(1/2)*b)+4*(a+(d*x+c)^(1/2)*b)^3*\cos(a+(d*x+c)^(1/2)*b)-12*(a+(d*x+c)^(1/2)*b)^2*\sin(a+(d*x+c)^(1/2)*b)+24*\sin(a+(d*x+c)^(1/2)*b)-24*(a+(d*x+c)^(1/2)*b)*\cos(a+(d*x+c)^(1/2)*b))+1/b^4*((a+(d*x+c)^(1/2)*b)^5*\sin(a+(d*x+c)^(1/2)*b)+5*(a+(d*x+c)^(1/2)*b)^4*\cos(a+(d*x+c)^(1/2)*b)-20*(a+(d*x+c)^(1/2)*b)^3*\sin(a+(d*x+c)^(1/2)*b)-60*(a+(d*x+c)^(1/2)*b)^2*\cos(a+(d*x+c)^(1/2)*b)+120*\cos(a+(d*x+c)^(1/2)*b)+120*(a+(d*x+c)^(1/2)*b)*\sin(a+(d*x+c)^(1/2)*b))) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.30

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{2((b^5 d^2 x^2 - 20 b^3 dx - 8 b^3 c + 120 b)\sqrt{dx + c} \sin(\sqrt{dx + c} b + a) + (5 b^4 d^2 x^2 - 48 b^2 c + 4(b^4 c - 15 b^2) d))}{b^6 d^3}$$

input `integrate(x^2*cos(a+b*(d*x+c)^(1/2)),x, algorithm="fricas")`

output `2*((b^5*d^2*x^2 - 20*b^3*d*x - 8*b^3*c + 120*b)*sqrt(d*x + c)*sin(sqrt(d*x + c)*b + a) + (5*b^4*d^2*x^2 - 48*b^2*c + 4*(b^4*c - 15*b^2)*d*x + 120)*cos(sqrt(d*x + c)*b + a))/(b^6*d^3)`

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 269, normalized size of antiderivative = 0.78

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx$$

$$= \begin{cases} \frac{x^3 \cos(a)}{3} \\ \frac{x^3 \cos(a+b\sqrt{c})}{3} \\ \frac{2x^2 \sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{bd} + \frac{8cx \cos(a+b\sqrt{c+dx})}{b^2 d^2} + \frac{10x^2 \cos(a+b\sqrt{c+dx})}{b^2 d} - \frac{16c\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3 d^3} - \frac{40x\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3 d^2} \end{cases}$$

input `integrate(x**2*cos(a+b*(d*x+c)**(1/2)),x)`

output `Piecewise((x**3*cos(a)/3, Eq(b, 0) & (Eq(b, 0) | Eq(d, 0))), (x**3*cos(a + b*sqrt(c))/3, Eq(d, 0)), (2*x**2*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b*d) + 8*c*x*cos(a + b*sqrt(c + d*x))/(b**2*d**2) + 10*x**2*cos(a + b*sqrt(c + d*x))/(b**2*d) - 16*c*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b**3*d**3) - 40*x*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b**3*d**2) - 96*c*cos(a + b*sqrt(c + d*x))/(b**4*d**3) - 120*x*cos(a + b*sqrt(c + d*x))/(b**4*d**2) + 240*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b**5*d**3) + 240*cos(a + b*sqrt(c + d*x))/(b**6*d**3), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 672 vs. $2(310) = 620$.

Time = 0.06 (sec) , antiderivative size = 672, normalized size of antiderivative = 1.94

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx = \text{Too large to display}$$

input `integrate(x^2*cos(a+b*(d*x+c)^(1/2)),x, algorithm="maxima")`

output

```
-2*(a*c^2*sin(sqrt(d*x + c)*b + a) - ((sqrt(d*x + c)*b + a)*sin(sqrt(d*x +
c)*b + a) + cos(sqrt(d*x + c)*b + a))*c^2 - 2*a^3*c*sin(sqrt(d*x + c)*b +
a)/b^2 + 6*((sqrt(d*x + c)*b + a)*sin(sqrt(d*x + c)*b + a) + cos(sqrt(d*x
+ c)*b + a))*a^2*c/b^2 + a^5*sin(sqrt(d*x + c)*b + a)/b^4 - 5*((sqrt(d*x
+ c)*b + a)*sin(sqrt(d*x + c)*b + a) + cos(sqrt(d*x + c)*b + a))*a^4/b^4 -
6*(2*(sqrt(d*x + c)*b + a)*cos(sqrt(d*x + c)*b + a) + ((sqrt(d*x + c)*b +
a)^2 - 2)*sin(sqrt(d*x + c)*b + a))*a*c/b^2 + 10*(2*(sqrt(d*x + c)*b + a)
*cos(sqrt(d*x + c)*b + a) + ((sqrt(d*x + c)*b + a)^2 - 2)*sin(sqrt(d*x + c
)*b + a))*a^3/b^4 + 2*(3*((sqrt(d*x + c)*b + a)^2 - 2)*cos(sqrt(d*x + c)*b
+ a) + ((sqrt(d*x + c)*b + a)^3 - 6*sqrt(d*x + c)*b - 6*a)*sin(sqrt(d*x +
c)*b + a))*c/b^2 - 10*(3*((sqrt(d*x + c)*b + a)^2 - 2)*cos(sqrt(d*x + c)*
b + a) + ((sqrt(d*x + c)*b + a)^3 - 6*sqrt(d*x + c)*b - 6*a)*sin(sqrt(d*x
+ c)*b + a))*a^2/b^4 + 5*(4*((sqrt(d*x + c)*b + a)^3 - 6*sqrt(d*x + c)*b -
6*a)*cos(sqrt(d*x + c)*b + a) + ((sqrt(d*x + c)*b + a)^4 - 12*(sqrt(d*x +
c)*b + a)^2 + 24)*sin(sqrt(d*x + c)*b + a))*a/b^4 - (5*((sqrt(d*x + c)*b
+ a)^4 - 12*(sqrt(d*x + c)*b + a)^2 + 24)*cos(sqrt(d*x + c)*b + a) + ((sq
r t(d*x + c)*b + a)^5 - 20*(sqrt(d*x + c)*b + a)^3 + 120*sqrt(d*x + c)*b + 1
20*a)*sin(sqrt(d*x + c)*b + a))/b^4)/(b^2*d^3)
```

Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 474, normalized size of antiderivative = 1.37

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{2 \left(\frac{b^4 c^2 - 6(\sqrt{dx+cb+a})^2 b^2 c + 12(\sqrt{dx+cb+a}) a b^2 c - 6 a^2 b^2 c + 5(\sqrt{dx+cb+a})^4 - 20(\sqrt{dx+cb+a})^3 a + 30(\sqrt{dx+cb+a})^2 a^2 - 20(\sqrt{dx+cb+a}) a^3 + 10 a^4}{b^4} \right)}{b^4}$$

input `integrate(x^2*cos(a+b*(d*x+c)^(1/2)),x, algorithm="giac")`

output
$$2*((b^4*c^2 - 6*(\sqrt{d*x + c})*b + a)^2*b^2*c + 12*(\sqrt{d*x + c})*b + a)*a*b^2*c - 6*a^2*b^2*c + 5*(\sqrt{d*x + c})*b + a)^4 - 20*(\sqrt{d*x + c})*b + a)^3*a + 30*(\sqrt{d*x + c})*b + a)^2*a^2 - 20*(\sqrt{d*x + c})*b + a)*a^3 + 5*a^4 + 12*b^2*c - 60*(\sqrt{d*x + c})*b + a)^2 + 120*(\sqrt{d*x + c})*b + a)*a - 60*a^2 + 120)*\cos(\sqrt{d*x + c})*b + a)/b^4 + ((\sqrt{d*x + c})*b + a)*b^4*c^2 - a*b^4*c^2 - 2*(\sqrt{d*x + c})*b + a)^3*b^2*c + 6*(\sqrt{d*x + c})*b + a)^2*a*b^2*c - 6*(\sqrt{d*x + c})*b + a)*a^2*b^2*c + 2*a^3*b^2*c + (\sqrt{d*x + c})*b + a)^5 - 5*(\sqrt{d*x + c})*b + a)^4*a + 10*(\sqrt{d*x + c})*b + a)^3*a^2 - 10*(\sqrt{d*x + c})*b + a)^2*a^3 + 5*(\sqrt{d*x + c})*b + a)*a^4 - a^5 + 12*(\sqrt{d*x + c})*b + a)*b^2*c - 12*a*b^2*c - 20*(\sqrt{d*x + c})*b + a)^3 + 60*(\sqrt{d*x + c})*b + a)^2*a - 60*(\sqrt{d*x + c})*b + a)*a^2 + 20*a^3 + 120*\sqrt{d*x + c})*b)*\sin(\sqrt{d*x + c})*b + a)/b^4)/(b^2*d^3)$$

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx = \int x^2 \cos(a + b\sqrt{c + dx}) dx$$

input `int(x^2*cos(a + b*(c + d*x)^(1/2)),x)`

output `int(x^2*cos(a + b*(c + d*x)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.55

$$\int x^2 \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{8 \cos(\sqrt{dx + c}b + a) b^4 c dx + 10 \cos(\sqrt{dx + c}b + a) b^4 d^2 x^2 - 96 \cos(\sqrt{dx + c}b + a) b^2 c - 120 \cos(\sqrt{dx + c}b + a) b^4 d^2 x^2}{1}$$

input `int(x^2*cos(a+b*(d*x+c)^(1/2)),x)`

output

```
(2*(4*cos(sqrt(c + d*x)*b + a)*b**4*c*d*x + 5*cos(sqrt(c + d*x)*b + a)*b**4*d**2*x**2 - 48*cos(sqrt(c + d*x)*b + a)*b**2*c - 60*cos(sqrt(c + d*x)*b + a)*b**2*d*x + 120*cos(sqrt(c + d*x)*b + a) + sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b**5*d**2*x**2 - 8*sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b**3*c - 20*sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b**3*d*x + 120*sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b))/(b**6*d**3)
```

3.91 $\int x \cos(a + b\sqrt{c + dx}) dx$

Optimal result	651
Mathematica [A] (verified)	652
Rubi [A] (verified)	652
Maple [A] (verified)	653
Fricas [A] (verification not implemented)	654
Sympy [A] (verification not implemented)	654
Maxima [A] (verification not implemented)	655
Giac [A] (verification not implemented)	655
Mupad [F(-1)]	656
Reduce [B] (verification not implemented)	656

Optimal result

Integrand size = 16, antiderivative size = 167

$$\int x \cos(a + b\sqrt{c + dx}) dx = -\frac{12 \cos(a + b\sqrt{c + dx})}{b^4 d^2} - \frac{2c \cos(a + b\sqrt{c + dx})}{b^2 d^2} + \frac{6(c + dx) \cos(a + b\sqrt{c + dx})}{b^2 d^2} - \frac{12\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b^3 d^2} - \frac{2c\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd^2} + \frac{2(c + dx)^{3/2} \sin(a + b\sqrt{c + dx})}{bd^2}$$

output

```
-12*cos(a+b*(d*x+c)^(1/2))/b^4/d^2-2*c*cos(a+b*(d*x+c)^(1/2))/b^2/d^2+6*(d*x+c)*cos(a+b*(d*x+c)^(1/2))/b^2/d^2-12*(d*x+c)^(1/2)*sin(a+b*(d*x+c)^(1/2))/b^3/d^2-2*c*(d*x+c)^(1/2)*sin(a+b*(d*x+c)^(1/2))/b/d^2+2*(d*x+c)^(3/2)*sin(a+b*(d*x+c)^(1/2))/b/d^2
```

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.43

$$\int x \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{2((-6 + b^2(2c + 3dx)) \cos(a + b\sqrt{c + dx}) + b\sqrt{c + dx}(-6 + b^2dx) \sin(a + b\sqrt{c + dx}))}{b^4d^2}$$

input `Integrate[x*Cos[a + b*Sqrt[c + d*x]],x]`

output `(2*((-6 + b^2*(2*c + 3*d*x))*Cos[a + b*Sqrt[c + d*x]] + b*Sqrt[c + d*x]*(-6 + b^2*d*x)*Sin[a + b*Sqrt[c + d*x]]))/(b^4*d^2)`

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3913, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \cos(a + b\sqrt{c + dx}) dx$$

$$\downarrow \text{3913}$$

$$\frac{2 \int \left(\frac{(c+dx)^{3/2} \cos(a+b\sqrt{c+dx})}{d} - \frac{c\sqrt{c+dx} \cos(a+b\sqrt{c+dx})}{d} \right) d\sqrt{c+dx}}{d}$$

$$\downarrow \text{2009}$$

$$\frac{2 \left(-\frac{6 \cos(a+b\sqrt{c+dx})}{b^4d} - \frac{6\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3d} + \frac{3(c+dx) \cos(a+b\sqrt{c+dx})}{b^2d} - \frac{c \cos(a+b\sqrt{c+dx})}{b^2d} + \frac{(c+dx)^{3/2} \sin(a+b\sqrt{c+dx})}{bd} - \frac{c \sin(a+b\sqrt{c+dx})}{bd} \right)}{d}$$

input `Int[x*Cos[a + b*Sqrt[c + d*x]],x]`

output

$$\frac{(2*((-6*\cos[a + b*\sqrt{c + d*x}]))/(b^4*d) - (c*\cos[a + b*\sqrt{c + d*x}]))/(b^2*d) + (3*(c + d*x)*\cos[a + b*\sqrt{c + d*x}]))/(b^2*d) - (6*\sqrt{c + d*x})*\sin[a + b*\sqrt{c + d*x}]))/(b^3*d) - (c*\sqrt{c + d*x})*\sin[a + b*\sqrt{c + d*x}]))/(b*d) + ((c + d*x)^{(3/2)}*\sin[a + b*\sqrt{c + d*x}]))/(b*d)))/d$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> Simp[IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 3913

$$\text{Int}[(a_.) + \cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^{(n_)}]*(b_.))^{(p_.)*((g_.) + (h_.)*(x_))^{(m_.)}, x_Symbol] \text{ :> Simp}[1/(n*f) \text{ Subst}[Int[ExpandIntegrand[(a + b*\cos[c + d*x])^p, x^{(1/n - 1)}*(g - e*(h/f) + h*(x^{(1/n)}/f))^{m}, x], x, (e + f*x)^n], x] \text{ /; FreeQ}\{a, b, c, d, e, f, g, h, m\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{IntegerQ}[1/n]$$

Maple [A] (verified)

Time = 1.49 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.78

method	result
parts	$\frac{2x\sqrt{dx+c} \sin(a+\sqrt{dx+cb})}{db} + \frac{2x \cos(a+\sqrt{dx+cb})}{db^2} - 2 \left(\frac{-2(a+\sqrt{dx+cb})^2 \cos(a+\sqrt{dx+cb}) + 4 \cos(a+\sqrt{dx+cb}) + 4(a+\sqrt{dx+cb})}{b^2} \right)$
derivativedivides	$\frac{2ac \sin(a+\sqrt{dx+cb}) - 2c(\cos(a+\sqrt{dx+cb}) + (a+\sqrt{dx+cb}) \sin(a+\sqrt{dx+cb})) - \frac{2a^3 \sin(a+\sqrt{dx+cb})}{b^2} + \frac{6a^2(\cos(a+\sqrt{dx+cb}) + (a+\sqrt{dx+cb}) \sin(a+\sqrt{dx+cb}))}{b^2}}{b^2}$
default	$\frac{2ac \sin(a+\sqrt{dx+cb}) - 2c(\cos(a+\sqrt{dx+cb}) + (a+\sqrt{dx+cb}) \sin(a+\sqrt{dx+cb})) - \frac{2a^3 \sin(a+\sqrt{dx+cb})}{b^2} + \frac{6a^2(\cos(a+\sqrt{dx+cb}) + (a+\sqrt{dx+cb}) \sin(a+\sqrt{dx+cb}))}{b^2}}{b^2}$

input

$$\text{int}(x*\cos(a+(d*x+c)^{(1/2)}*b), x, \text{method}=_RETURNVERBOSE)$$

output

```
2/d/b*x*(d*x+c)^(1/2)*sin(a+(d*x+c)^(1/2)*b)+2/d/b^2*x*cos(a+(d*x+c)^(1/2)*b)-2/d/b^2*(2/d/b^2*(-(a+(d*x+c)^(1/2)*b)^2*cos(a+(d*x+c)^(1/2)*b)+2*cos(a+(d*x+c)^(1/2)*b)+2*(a+(d*x+c)^(1/2)*b)*sin(a+(d*x+c)^(1/2)*b)-a*(sin(a+(d*x+c)^(1/2)*b)-(a+(d*x+c)^(1/2)*b)*cos(a+(d*x+c)^(1/2)*b)))-2*a/d/b^2*(sin(a+(d*x+c)^(1/2)*b)-(a+(d*x+c)^(1/2)*b)*cos(a+(d*x+c)^(1/2)*b)+cos(a+(d*x+c)^(1/2)*b)*a)+2/d/b^2*(cos(a+(d*x+c)^(1/2)*b)+(a+(d*x+c)^(1/2)*b)*sin(a+(d*x+c)^(1/2)*b)-a*sin(a+(d*x+c)^(1/2)*b))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.40

$$\int x \cos \left(a + b\sqrt{c + dx} \right) dx$$

$$= \frac{2 \left((b^3 dx - 6b) \sqrt{dx + c} \sin(\sqrt{dx + c} b + a) + (3b^2 dx + 2b^2 c - 6) \cos(\sqrt{dx + c} b + a) \right)}{b^4 d^2}$$

input

```
integrate(x*cos(a+b*(d*x+c)^(1/2)),x, algorithm="fricas")
```

output

```
2*((b^3*d*x - 6*b)*sqrt(d*x + c)*sin(sqrt(d*x + c)*b + a) + (3*b^2*d*x + 2*b^2*c - 6)*cos(sqrt(d*x + c)*b + a))/(b^4*d^2)
```

Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.90

$$\int x \cos \left(a + b\sqrt{c + dx} \right) dx$$

$$= \begin{cases} \frac{x^2 \cos(a)}{2} \\ \frac{x^2 \cos(a+b\sqrt{c})}{2} \\ \frac{2x\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{bd} + \frac{4c \cos(a+b\sqrt{c+dx})}{b^2 d^2} + \frac{6x \cos(a+b\sqrt{c+dx})}{b^2 d} - \frac{12\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{b^3 d^2} - \frac{12 \cos(a+b\sqrt{c+dx})}{b^4 d^2} \end{cases}$$

input

```
integrate(x*cos(a+b*(d*x+c)**(1/2)),x)
```

for
for
otl

output

```
Piecewise((x**2*cos(a)/2, Eq(b, 0) & (Eq(b, 0) | Eq(d, 0))), (x**2*cos(a +
b*sqrt(c))/2, Eq(d, 0)), (2*x*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b*d
) + 4*c*cos(a + b*sqrt(c + d*x))/(b**2*d**2) + 6*x*cos(a + b*sqrt(c + d*x)
)/(b**2*d) - 12*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b**3*d**2) - 12*co
s(a + b*sqrt(c + d*x))/(b**4*d**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.57

$$\int x \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{2 \left(ac \sin(\sqrt{dx + cb} + a) - ((\sqrt{dx + cb} + a) \sin(\sqrt{dx + cb} + a) + \cos(\sqrt{dx + cb} + a))c - \frac{a^3 \sin(\sqrt{dx + cb}}{b^2} \right)}{b^2}$$

input

```
integrate(x*cos(a+b*(d*x+c)^(1/2)),x, algorithm="maxima")
```

output

```
2*(a*c*sin(sqrt(d*x + c)*b + a) - ((sqrt(d*x + c)*b + a)*sin(sqrt(d*x + c)
*b + a) + cos(sqrt(d*x + c)*b + a))*c - a^3*sin(sqrt(d*x + c)*b + a)/b^2 +
3*((sqrt(d*x + c)*b + a)*sin(sqrt(d*x + c)*b + a) + cos(sqrt(d*x + c)*b +
a))*a^2/b^2 - 3*(2*(sqrt(d*x + c)*b + a)*cos(sqrt(d*x + c)*b + a) + ((sqr
t(d*x + c)*b + a)^2 - 2)*sin(sqrt(d*x + c)*b + a))*a/b^2 + (3*((sqrt(d*x +
c)*b + a)^2 - 2)*cos(sqrt(d*x + c)*b + a) + ((sqrt(d*x + c)*b + a)^3 - 6*
sqrt(d*x + c)*b - 6*a)*sin(sqrt(d*x + c)*b + a))/b^2)/(b^2*d^2)
```

Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.99

$$\int x \cos(a + b\sqrt{c + dx}) dx =$$

$$\frac{2 \left(\frac{(b^2c - 3(\sqrt{dx + cb} + a)^2 + 6(\sqrt{dx + cb} + a)a - 3a^2 + 6) \cos(\sqrt{dx + cb} + a)}{b^2} + \frac{((\sqrt{dx + cb} + a)b^2c - ab^2c - (\sqrt{dx + cb} + a)^3 + 3(\sqrt{dx + cb} + a)^2 a - 6a^2) \sin(\sqrt{dx + cb} + a)}{b^2} \right)}{b^2 d^2}$$

input

```
integrate(x*cos(a+b*(d*x+c)^(1/2)),x, algorithm="giac")
```


output

```
-2*((b^2*c - 3*(sqrt(d*x + c)*b + a)^2 + 6*(sqrt(d*x + c)*b + a)*a - 3*a^2
+ 6)*cos(sqrt(d*x + c)*b + a)/b^2 + ((sqrt(d*x + c)*b + a)*b^2*c - a*b^2*
c - (sqrt(d*x + c)*b + a)^3 + 3*(sqrt(d*x + c)*b + a)^2*a - 3*(sqrt(d*x +
c)*b + a)*a^2 + a^3 + 6*sqrt(d*x + c)*b)*sin(sqrt(d*x + c)*b + a)/b^2)/(b^
2*d^2)
```

Mupad [F(-1)]

Timed out.

$$\int x \cos(a + b\sqrt{c + dx}) dx = \int x \cos(a + b\sqrt{c + dx}) dx$$

input

```
int(x*cos(a + b*(c + d*x)^(1/2)),x)
```

output

```
int(x*cos(a + b*(c + d*x)^(1/2)), x)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.60

$$\int x \cos(a + b\sqrt{c + dx}) dx$$

$$= \frac{4 \cos(\sqrt{dx + c}b + a) b^2 c + 6 \cos(\sqrt{dx + c}b + a) b^2 dx - 12 \cos(\sqrt{dx + c}b + a) + 2\sqrt{dx + c} \sin(\sqrt{dx + c}b + a)}{b^4 d^2}$$

input

```
int(x*cos(a+b*(d*x+c)^(1/2)),x)
```

output

```
(2*(2*cos(sqrt(c + d*x)*b + a)*b**2*c + 3*cos(sqrt(c + d*x)*b + a)*b**2*d*
x - 6*cos(sqrt(c + d*x)*b + a) + sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b*
*3*d*x - 6*sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b))/(b**4*d**2)
```

3.92 $\int \cos(a + b\sqrt{c + dx}) dx$

Optimal result	657
Mathematica [A] (verified)	657
Rubi [A] (verified)	658
Maple [A] (verified)	659
Fricas [A] (verification not implemented)	660
Sympy [A] (verification not implemented)	660
Maxima [A] (verification not implemented)	661
Giac [A] (verification not implemented)	661
Mupad [B] (verification not implemented)	661
Reduce [B] (verification not implemented)	662

Optimal result

Integrand size = 14, antiderivative size = 54

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2 \cos(a + b\sqrt{c + dx})}{b^2 d} + \frac{2\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{bd}$$

output

```
2*cos(a+b*(d*x+c)^(1/2))/b^2/d+2*(d*x+c)^(1/2)*sin(a+b*(d*x+c)^(1/2))/b/d
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.89

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2(\cos(a + b\sqrt{c + dx}) + b\sqrt{c + dx} \sin(a + b\sqrt{c + dx}))}{b^2 d}$$

input

```
Integrate[Cos[a + b*Sqrt[c + d*x]],x]
```

output

```
(2*(Cos[a + b*Sqrt[c + d*x]] + b*Sqrt[c + d*x]*Sin[a + b*Sqrt[c + d*x]]))/
(b^2*d)
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3843, 3042, 3777, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(a + b\sqrt{c + dx}) dx \\
 & \quad \downarrow \text{3843} \\
 & \frac{2 \int \sqrt{c + dx} \cos(a + b\sqrt{c + dx}) d\sqrt{c + dx}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \int \sqrt{c + dx} \sin(a + b\sqrt{c + dx} + \frac{\pi}{2}) d\sqrt{c + dx}}{d} \\
 & \quad \downarrow \text{3777} \\
 & \frac{2 \left(\frac{\int -\sin(a + b\sqrt{c + dx}) d\sqrt{c + dx}}{b} + \frac{\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b} \right)}{d} \\
 & \quad \downarrow \text{25} \\
 & \frac{2 \left(\frac{\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b} - \frac{\int \sin(a + b\sqrt{c + dx}) d\sqrt{c + dx}}{b} \right)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \left(\frac{\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b} - \frac{\int \sin(a + b\sqrt{c + dx}) d\sqrt{c + dx}}{b} \right)}{d} \\
 & \quad \downarrow \text{3118} \\
 & \frac{2 \left(\frac{\cos(a + b\sqrt{c + dx})}{b^2} + \frac{\sqrt{c + dx} \sin(a + b\sqrt{c + dx})}{b} \right)}{d}
 \end{aligned}$$

input `Int[Cos[a + b*Sqrt[c + d*x]],x]`

output $(2*(\cos[a + b*\sqrt{c + d*x}]/b^2 + (\sqrt{c + d*x}*\sin[a + b*\sqrt{c + d*x}])/b))/d$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3118 $\text{Int}[\sin[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\cos[c + d*x]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$

rule 3777 $\text{Int}[((c_.) + (d_.)*(x_))^{(m_.)*\sin[(e_.) + (f_.)*(x_)]}, x_Symbol] \rightarrow \text{Simp}[-(c + d*x)^m * (\cos[e + f*x]/f), x] + \text{Simp}[d*(m/f) \text{ Int}[(c + d*x)^{(m-1)} * \cos[e + f*x], x], x] \text{ ; FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{GtQ}[m, 0]$

rule 3843 $\text{Int}[((a_.) + \cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^{(n_.)}]*(b_.))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/(n*f) \text{ Subst}[\text{Int}[x^{(1/n-1)}*(a + b*\cos[c + d*x])^p, x], x, (e + f*x)^n], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{IntegerQ}[1/n]$

Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.13

method	result	size
derivativedivides	$\frac{2 \cos(a + \sqrt{dx+cb}) + 2(a + \sqrt{dx+cb}) \sin(a + \sqrt{dx+cb}) - 2a \sin(a + \sqrt{dx+cb})}{b^2 d}$	61
default	$\frac{2 \cos(a + \sqrt{dx+cb}) + 2(a + \sqrt{dx+cb}) \sin(a + \sqrt{dx+cb}) - 2a \sin(a + \sqrt{dx+cb})}{b^2 d}$	61

input $\text{int}(\cos(a+(d*x+c)^{(1/2)}*b), x, \text{method}=_RETURNVERBOSE)$

output $2/d/b^2*(\cos(a+(d*x+c)^{(1/2)*b})+(a+(d*x+c)^{(1/2)*b})*\sin(a+(d*x+c)^{(1/2)*b})-a*\sin(a+(d*x+c)^{(1/2)*b}))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2(\sqrt{dx + cb} \sin(\sqrt{dx + cb} + a) + \cos(\sqrt{dx + cb} + a))}{b^2 d}$$

input `integrate(cos(a+b*(d*x+c)^(1/2)),x, algorithm="fricas")`

output $2*(\sqrt{d*x + c}*b*\sin(\sqrt{d*x + c}*b + a) + \cos(\sqrt{d*x + c}*b + a))/(b^2*d)$

Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.20

$$\int \cos(a + b\sqrt{c + dx}) dx = \begin{cases} x \cos(a) & \text{for } b = 0 \wedge (b = 0 \vee d = 0) \\ x \cos(a + b\sqrt{c}) & \text{for } d = 0 \\ \frac{2\sqrt{c+dx} \sin(a+b\sqrt{c+dx})}{bd} + \frac{2 \cos(a+b\sqrt{c+dx})}{b^2 d} & \text{otherwise} \end{cases}$$

input `integrate(cos(a+b*(d*x+c)**(1/2)),x)`

output `Piecewise((x*cos(a), Eq(b, 0) & (Eq(b, 0) | Eq(d, 0))), (x*cos(a + b*sqrt(c)), Eq(d, 0)), (2*sqrt(c + d*x)*sin(a + b*sqrt(c + d*x))/(b*d) + 2*cos(a + b*sqrt(c + d*x))/(b**2*d), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.11

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2((\sqrt{dx + cb} + a) \sin(\sqrt{dx + cb} + a) - a \sin(\sqrt{dx + cb} + a) + \cos(\sqrt{dx + cb} + a))}{b^2 d}$$

input `integrate(cos(a+b*(d*x+c)^(1/2)),x, algorithm="maxima")`output `2*((sqrt(d*x + c)*b + a)*sin(sqrt(d*x + c)*b + a) - a*sin(sqrt(d*x + c)*b + a) + cos(sqrt(d*x + c)*b + a))/(b^2*d)`**Giac [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2(\sqrt{dx + cb} \sin(\sqrt{dx + cb} + a) + \cos(\sqrt{dx + cb} + a))}{b^2 d}$$

input `integrate(cos(a+b*(d*x+c)^(1/2)),x, algorithm="giac")`output `2*(sqrt(d*x + c)*b*sin(sqrt(d*x + c)*b + a) + cos(sqrt(d*x + c)*b + a))/(b^2*d)`**Mupad [B] (verification not implemented)**

Time = 41.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2(\cos(a + b\sqrt{c + dx}) + b \sin(a + b\sqrt{c + dx}) \sqrt{c + dx})}{b^2 d}$$

input `int(cos(a + b*(c + d*x)^(1/2)),x)`

output

```
(2*(cos(a + b*(c + d*x)^(1/2)) + b*sin(a + b*(c + d*x)^(1/2))*(c + d*x)^(1/2)))/(b^2*d)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.72

$$\int \cos(a + b\sqrt{c + dx}) dx = \frac{2 \cos(\sqrt{dx + c}b + a) + 2\sqrt{dx + c} \sin(\sqrt{dx + c}b + a) b}{b^2 d}$$

input

```
int(cos(a+b*(d*x+c)^(1/2)),x)
```

output

```
(2*(cos(sqrt(c + d*x)*b + a) + sqrt(c + d*x)*sin(sqrt(c + d*x)*b + a)*b))/(b**2*d)
```

3.93 $\int \frac{\cos(a+b\sqrt{c+dx})}{x} dx$

Optimal result	663
Mathematica [C] (verified)	664
Rubi [A] (verified)	664
Maple [B] (verified)	665
Fricas [C] (verification not implemented)	666
Sympy [F]	667
Maxima [F]	667
Giac [F]	667
Mupad [F(-1)]	668
Reduce [F]	668

Optimal result

Integrand size = 18, antiderivative size = 126

$$\int \frac{\cos(a+b\sqrt{c+dx})}{x} dx = \cos(a-b\sqrt{c}) \operatorname{CosIntegral}\left(b\left(\sqrt{c}+\sqrt{c+dx}\right)\right) + \cos(a+b\sqrt{c}) \operatorname{CosIntegral}\left(b\sqrt{c}-b\sqrt{c+dx}\right) - \sin(a-b\sqrt{c}) \operatorname{Si}\left(b\left(\sqrt{c}+\sqrt{c+dx}\right)\right) + \sin(a+b\sqrt{c}) \operatorname{Si}\left(b\sqrt{c}-b\sqrt{c+dx}\right)$$

output

```
cos(a-b*c^(1/2))*Ci(b*(c^(1/2)+(d*x+c)^(1/2)))+cos(a+b*c^(1/2))*Ci(b*c^(1/2)-b*(d*x+c)^(1/2))-sin(a-b*c^(1/2))*Si(b*(c^(1/2)+(d*x+c)^(1/2)))+sin(a+b*c^(1/2))*Si(b*c^(1/2)-b*(d*x+c)^(1/2))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.37 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.15

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \frac{1}{2} e^{-i(a+b\sqrt{c})} \left(\text{ExpIntegralEi} \left(-ib \left(-\sqrt{c} + \sqrt{c + dx} \right) \right) \right. \\ \left. + e^{2i(a+b\sqrt{c})} \text{ExpIntegralEi} \left(ib \left(-\sqrt{c} + \sqrt{c + dx} \right) \right) \right) \\ + e^{2ib\sqrt{c}} \text{ExpIntegralEi} \left(-ib \left(\sqrt{c} + \sqrt{c + dx} \right) \right) \\ \left. + e^{2ia} \text{ExpIntegralEi} \left(ib \left(\sqrt{c} + \sqrt{c + dx} \right) \right) \right)$$

input `Integrate[Cos[a + b*Sqrt[c + d*x]]/x,x]`

output `(ExpIntegralEi[(-I)*b*(-Sqrt[c] + Sqrt[c + d*x])] + E^((2*I)*(a + b*Sqrt[c]))*ExpIntegralEi[I*b*(-Sqrt[c] + Sqrt[c + d*x])] + E^((2*I)*b*Sqrt[c])*ExpIntegralEi[(-I)*b*(Sqrt[c] + Sqrt[c + d*x])] + E^((2*I)*a)*ExpIntegralEi[I*b*(Sqrt[c] + Sqrt[c + d*x])])/(2*E^(I*(a + b*Sqrt[c])))`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.19, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3913, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx \\ \downarrow \text{3913} \\ \frac{2 \int \left(\frac{d \cos(a + b\sqrt{c + dx})}{2(\sqrt{c} + \sqrt{c + dx})} - \frac{d \cos(a + b\sqrt{c + dx})}{2(\sqrt{c} - \sqrt{c + dx})} \right) d\sqrt{c + dx}}{d} \\ \downarrow \text{2009}$$

$$\frac{2(\frac{1}{2}d \cos(a + b\sqrt{c}) \operatorname{CosIntegral}(b\sqrt{c} - b\sqrt{c + dx}) + \frac{1}{2}d \cos(a - b\sqrt{c}) \operatorname{CosIntegral}(\sqrt{cb} + \sqrt{c + dx}b) + \frac{1}{2}d \sin(a + b\sqrt{c}) \operatorname{SinIntegral}(b\sqrt{c} - b\sqrt{c + dx}) - \frac{1}{2}d \sin(a - b\sqrt{c}) \operatorname{SinIntegral}(\sqrt{cb} + \sqrt{c + dx}b))}{d}$$

input `Int[Cos[a + b*Sqrt[c + d*x]]/x,x]`

output `(2*((d*Cos[a + b*Sqrt[c]]*CosIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]])/2 + (d*Cos[a - b*Sqrt[c]]*CosIntegral[b*Sqrt[c] + b*Sqrt[c + d*x]])/2 + (d*Sin[a + b*Sqrt[c]]*SinIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]])/2 - (d*Sin[a - b*Sqrt[c]]*SinIntegral[b*Sqrt[c] + b*Sqrt[c + d*x]]))/2)/d`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3913 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Simp[1/(n*f) Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - e*(h/f) + h*(x^(1/n)/f))^m, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 270 vs. 2(102) = 204.

Time = 1.08 (sec) , antiderivative size = 271, normalized size of antiderivative = 2.15

method	result
derivativedivides	$\frac{b(a+b\sqrt{c}) \left(\frac{\sin(a+b\sqrt{c}) \operatorname{Si}(b\sqrt{c}-\sqrt{dx+c}b) + \operatorname{Ci}(\sqrt{dx+c}b-b\sqrt{c}) \cos(a+b\sqrt{c})}{\sqrt{c}} \right) - b(a-b\sqrt{c}) \left(\frac{-\operatorname{Si}(\sqrt{dx+c}b+b\sqrt{c}) \sin(a-b\sqrt{c}) + \operatorname{Ci}(\sqrt{dx+c}b-b\sqrt{c}) \cos(a-b\sqrt{c})}{\sqrt{c}} \right)}{d}$
default	$\frac{b(a+b\sqrt{c}) \left(\frac{\sin(a+b\sqrt{c}) \operatorname{Si}(b\sqrt{c}-\sqrt{dx+c}b) + \operatorname{Ci}(\sqrt{dx+c}b-b\sqrt{c}) \cos(a+b\sqrt{c})}{\sqrt{c}} \right) - b(a-b\sqrt{c}) \left(\frac{-\operatorname{Si}(\sqrt{dx+c}b+b\sqrt{c}) \sin(a-b\sqrt{c}) + \operatorname{Ci}(\sqrt{dx+c}b-b\sqrt{c}) \cos(a-b\sqrt{c})}{\sqrt{c}} \right)}{d}$

input `int(cos(a+(d*x+c)^(1/2)*b)/x,x,method=_RETURNVERBOSE)`

output

```
2/b^2*(1/2*b*(a+b*c^(1/2))/c^(1/2)*(sin(a+b*c^(1/2))*Si(b*c^(1/2)-(d*x+c)^(1/2)*b)+Ci((d*x+c)^(1/2)*b-b*c^(1/2))*cos(a+b*c^(1/2)))-1/2*b*(a-b*c^(1/2))/c^(1/2)*(-Si((d*x+c)^(1/2)*b+b*c^(1/2))*sin(a-b*c^(1/2))+Ci((d*x+c)^(1/2)*b+b*c^(1/2))*cos(a-b*c^(1/2)))-a*b^2*(1/2/b/c^(1/2)*(sin(a+b*c^(1/2))*Si(b*c^(1/2)-(d*x+c)^(1/2)*b)+Ci((d*x+c)^(1/2)*b-b*c^(1/2))*cos(a+b*c^(1/2)))-1/2/b/c^(1/2)*(-Si((d*x+c)^(1/2)*b+b*c^(1/2))*sin(a-b*c^(1/2))+Ci((d*x+c)^(1/2)*b+b*c^(1/2))*cos(a-b*c^(1/2))))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.18

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \frac{1}{2} \operatorname{Ei}\left(i\sqrt{dx + cb} - \sqrt{-b^2c}\right) e^{(ia + \sqrt{-b^2c})} + \frac{1}{2} \operatorname{Ei}\left(i\sqrt{dx + cb} + \sqrt{-b^2c}\right) e^{(ia - \sqrt{-b^2c})} + \frac{1}{2} \operatorname{Ei}\left(-i\sqrt{dx + cb} - \sqrt{-b^2c}\right) e^{(-ia + \sqrt{-b^2c})} + \frac{1}{2} \operatorname{Ei}\left(-i\sqrt{dx + cb} + \sqrt{-b^2c}\right) e^{(-ia - \sqrt{-b^2c})}$$

input

```
integrate(cos(a+b*(d*x+c)^(1/2))/x,x, algorithm="fricas")
```

output

```
1/2*Ei(I*sqrt(d*x + c)*b - sqrt(-b^2*c))*e^(I*a + sqrt(-b^2*c)) + 1/2*Ei(I*sqrt(d*x + c)*b + sqrt(-b^2*c))*e^(I*a - sqrt(-b^2*c)) + 1/2*Ei(-I*sqrt(d*x + c)*b - sqrt(-b^2*c))*e^(-I*a + sqrt(-b^2*c)) + 1/2*Ei(-I*sqrt(d*x + c)*b + sqrt(-b^2*c))*e^(-I*a - sqrt(-b^2*c))
```

Sympy [F]

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \int \frac{\cos(a + b\sqrt{c + dx})}{x} dx$$

input `integrate(cos(a+b*(d*x+c)**(1/2))/x,x)`

output `Integral(cos(a + b*sqrt(c + d*x))/x, x)`

Maxima [F]

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \int \frac{\cos(\sqrt{dx + cb} + a)}{x} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/2))/x,x, algorithm="maxima")`

output `integrate(cos(sqrt(d*x + c)*b + a)/x, x)`

Giac [F]

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \int \frac{\cos(\sqrt{dx + cb} + a)}{x} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/2))/x,x, algorithm="giac")`

output `integrate(cos(sqrt(d*x + c)*b + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \int \frac{\cos(a + b\sqrt{c + dx})}{x} dx$$

input `int(cos(a + b*(c + d*x)^(1/2))/x,x)`output `int(cos(a + b*(c + d*x)^(1/2))/x, x)`**Reduce [F]**

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x} dx = \int \frac{\cos(\sqrt{dx + c}b + a)}{x} dx$$

input `int(cos(a+b*(d*x+c)^(1/2))/x,x)`output `int(cos(sqrt(c + d*x)*b + a)/x,x)`

3.94 $\int \frac{\cos(a+b\sqrt{c+dx})}{x^2} dx$

Optimal result	669
Mathematica [C] (verified)	670
Rubi [A] (verified)	670
Maple [B] (verified)	672
Fricas [C] (verification not implemented)	673
Sympy [F]	674
Maxima [F]	674
Giac [F]	674
Mupad [F(-1)]	675
Reduce [F]	675

Optimal result

Integrand size = 18, antiderivative size = 184

$$\int \frac{\cos(a+b\sqrt{c+dx})}{x^2} dx = -\frac{\cos(a+b\sqrt{c+dx})}{x} + \frac{bd \operatorname{CosIntegral}(b(\sqrt{c} + \sqrt{c+dx})) \sin(a-b\sqrt{c})}{2\sqrt{c}} - \frac{bd \operatorname{CosIntegral}(b\sqrt{c} - b\sqrt{c+dx}) \sin(a+b\sqrt{c})}{2\sqrt{c}} + \frac{bd \cos(a-b\sqrt{c}) \operatorname{Si}(b(\sqrt{c} + \sqrt{c+dx}))}{2\sqrt{c}} + \frac{bd \cos(a+b\sqrt{c}) \operatorname{Si}(b\sqrt{c} - b\sqrt{c+dx})}{2\sqrt{c}}$$

output

```
-cos(a+b*(d*x+c)^(1/2))/x+1/2*b*d*Ci(b*(c^(1/2)+(d*x+c)^(1/2)))*sin(a-b*c^(1/2))/c^(1/2)-1/2*b*d*Ci(b*c^(1/2)-b*(d*x+c)^(1/2))*sin(a+b*c^(1/2))/c^(1/2)+1/2*b*d*cos(a-b*c^(1/2))*Si(b*(c^(1/2)+(d*x+c)^(1/2)))/c^(1/2)+1/2*b*d*cos(a+b*c^(1/2))*Si(b*c^(1/2)-b*(d*x+c)^(1/2))/c^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.46 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.30

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx$$

$$= \frac{i \left(e^{-ia} \left(2i\sqrt{c} e^{-ib\sqrt{c+dx}} - b d e^{-ib\sqrt{c}} x \operatorname{ExpIntegralEi}(-ib(-\sqrt{c} + \sqrt{c+dx})) \right) + b d e^{ib\sqrt{c}} x \operatorname{ExpIntegralEi}(-ib(\sqrt{c} + \sqrt{c+dx})) \right)}{x^2}$$

input `Integrate[Cos[a + b*Sqrt[c + d*x]]/x^2,x]`

output

```
((I/4)*(((2*I)*Sqrt[c])/E^(I*b*Sqrt[c + d*x]) - (b*d*x*ExpIntegralEi[(-I)*b*(-Sqrt[c] + Sqrt[c + d*x]])/E^(I*b*Sqrt[c]) + b*d*E^(I*b*Sqrt[c])*x*ExpIntegralEi[(-I)*b*(Sqrt[c] + Sqrt[c + d*x]])/E^(I*a) + E^(I*(a - b*Sqrt[c]))*((2*I)*Sqrt[c]*E^(I*b*(Sqrt[c] + Sqrt[c + d*x])) + b*d*E^((2*I)*b*Sqrt[c])*x*ExpIntegralEi[I*b*(-Sqrt[c] + Sqrt[c + d*x])] - b*d*x*ExpIntegralEi[I*b*(Sqrt[c] + Sqrt[c + d*x])])))/(Sqrt[c]*x)
```

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3913, 27, 3823, 3814, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx$$

$$\downarrow \text{3913}$$

$$\frac{2 \int \frac{\sqrt{c+dx} \cos(a+b\sqrt{c+dx})}{x^2} d\sqrt{c+dx}}{d}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& 2d \int \frac{\sqrt{c+dx} \cos(a+b\sqrt{c+dx})}{d^2 x^2} d\sqrt{c+dx} \\
& \quad \downarrow \text{3823} \\
& 2d \left(\frac{1}{2} b \int -\frac{\sin(a+b\sqrt{c+dx})}{dx} d\sqrt{c+dx} - \frac{\cos(a+b\sqrt{c+dx})}{2dx} \right) \\
& \quad \downarrow \text{3814} \\
& 2d \left(\frac{1}{2} b \int \left(\frac{\sin(a+b\sqrt{c+dx})}{2\sqrt{c}(\sqrt{c}-\sqrt{c+dx})} + \frac{\sin(a+b\sqrt{c+dx})}{2\sqrt{c}(\sqrt{c}+\sqrt{c+dx})} \right) d\sqrt{c+dx} - \frac{\cos(a+b\sqrt{c+dx})}{2dx} \right) \\
& \quad \downarrow \text{2009} \\
& 2d \left(\frac{1}{2} b \left(\frac{\sin(a-b\sqrt{c}) \operatorname{CosIntegral}(\sqrt{cb}+\sqrt{c+dx})}{2\sqrt{c}} - \frac{\sin(a+b\sqrt{c}) \operatorname{CosIntegral}(b\sqrt{c}-b\sqrt{c+dx})}{2\sqrt{c}} + \frac{\cos(a-b\sqrt{c}) \operatorname{SiIntegral}(\sqrt{cb}+\sqrt{c+dx})}{2\sqrt{c}} - \frac{\cos(a+b\sqrt{c}) \operatorname{SiIntegral}(b\sqrt{c}-b\sqrt{c+dx})}{2\sqrt{c}} \right) \right)
\end{aligned}$$

input `Int[Cos[a + b*Sqrt[c + d*x]]/x^2,x]`

output `2*d*(-1/2*Cos[a + b*Sqrt[c + d*x]]/(d*x) + (b*((CosIntegral[b*Sqrt[c] + b*Sqrt[c + d*x]]*Sin[a - b*Sqrt[c]])/(2*Sqrt[c]) - (CosIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]]*Sin[a + b*Sqrt[c]])/(2*Sqrt[c]) + (Cos[a + b*Sqrt[c]]*SinIntegral[b*Sqrt[c] - b*Sqrt[c + d*x]])/(2*Sqrt[c]) + (Cos[a - b*Sqrt[c]]*SinIntegral[b*Sqrt[c] + b*Sqrt[c + d*x]])/(2*Sqrt[c])))/2)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3814 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)*Sin[(c_.) + (d_.)*(x_)], x_Symbol] := Int[ExpandIntegrand[Sin[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])`

rule 3823

```
Int[Cos[(c_.) + (d_.)*(x_)]*((e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_)^(n_))^(p_
), x_Symbol] :> Simp[e^m*(a + b*x^n)^(p + 1)*(Cos[c + d*x]/(b*n*(p + 1))),
x] + Simp[d*(e^m/(b*n*(p + 1))) Int[(a + b*x^n)^(p + 1)*Sin[c + d*x], x],
x] /; FreeQ[{a, b, c, d, e, m, n}, x] && ILtQ[p, -1] && EqQ[m, n - 1] && (
IntegerQ[n] || GtQ[e, 0])
```

rule 3913

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_)]*(b_.))^(p_.)*((g_
.) + (h_.)*(x_))^(m_.), x_Symbol] :> Simp[1/(n*f) Subst[Int[ExpandIntegra
nd[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - e*(h/f) + h*(x^(1/n)/f))^m, x],
x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p
, 0] && IntegerQ[1/n]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 713 vs. 2(142) = 284.

Time = 1.19 (sec) , antiderivative size = 714, normalized size of antiderivative = 3.88

method	result
derivativedivides	$2d \left(\frac{\cos(a + \sqrt{dx + c} b) \left(-\frac{a b^2 (a + \sqrt{dx + c} b)}{2c} + \frac{b^2 (-b^2 c + a^2)}{2c} \right)}{-b^2 c + a^2 - 2(a + \sqrt{dx + c} b)a + (a + \sqrt{dx + c} b)^2} - \frac{ab(\sin(a + b\sqrt{c}) \operatorname{Si}(b\sqrt{c} - \sqrt{dx + c} b) + \operatorname{Ci}(\sqrt{dx + c} b - b\sqrt{c})) \cos(a + b\sqrt{c})}{4c^{\frac{3}{2}}} \right)$
default	$2d \left(\frac{\cos(a + \sqrt{dx + c} b) \left(-\frac{a b^2 (a + \sqrt{dx + c} b)}{2c} + \frac{b^2 (-b^2 c + a^2)}{2c} \right)}{-b^2 c + a^2 - 2(a + \sqrt{dx + c} b)a + (a + \sqrt{dx + c} b)^2} - \frac{ab(\sin(a + b\sqrt{c}) \operatorname{Si}(b\sqrt{c} - \sqrt{dx + c} b) + \operatorname{Ci}(\sqrt{dx + c} b - b\sqrt{c})) \cos(a + b\sqrt{c})}{4c^{\frac{3}{2}}} \right)$

input

```
int(cos(a+(d*x+c)^(1/2)*b)/x^2,x,method=_RETURNVERBOSE)
```

output

```

2*d/b^2*(cos(a+(d*x+c)^(1/2)*b)*(-1/2*a*b^2/c*(a+(d*x+c)^(1/2)*b)+1/2*b^2*
(-b^2*c+a^2)/c)/(-b^2*c+a^2-2*(a+(d*x+c)^(1/2)*b)*a+(a+(d*x+c)^(1/2)*b)^2)
-1/4*a*b/c^(3/2)*(sin(a+b*c^(1/2))*Si(b*c^(1/2)-(d*x+c)^(1/2)*b)+Ci((d*x+c)
)^(1/2)*b-b*c^(1/2))*cos(a+b*c^(1/2)))+1/4*a*b/c^(3/2)*(-Si((d*x+c)^(1/2)*
b+b*c^(1/2))*sin(a-b*c^(1/2))+Ci((d*x+c)^(1/2)*b+b*c^(1/2))*cos(a-b*c^(1/2)
)))+1/4*b*(-b^2*c+a^2-(a+b*c^(1/2))*a)/c^(3/2)*(-Si(b*c^(1/2)-(d*x+c)^(1/2)
)*b)*cos(a+b*c^(1/2))+Ci((d*x+c)^(1/2)*b-b*c^(1/2))*sin(a+b*c^(1/2)))-1/4*
b*(-b^2*c+a^2-(a-b*c^(1/2))*a)/c^(3/2)*(Si((d*x+c)^(1/2)*b+b*c^(1/2))*cos(
a-b*c^(1/2))+Ci((d*x+c)^(1/2)*b+b*c^(1/2))*sin(a-b*c^(1/2)))-a*b^4*(cos(a+
(d*x+c)^(1/2)*b)*(-1/2/c/b^2*(a+(d*x+c)^(1/2)*b)+1/2*a/c/b^2)/(-b^2*c+a^2-
2*(a+(d*x+c)^(1/2)*b)*a+(a+(d*x+c)^(1/2)*b)^2)-1/4/c^(3/2)/b^3*(sin(a+b*c^
(1/2))*Si(b*c^(1/2)-(d*x+c)^(1/2)*b)+Ci((d*x+c)^(1/2)*b-b*c^(1/2))*cos(a+b
*c^(1/2)))+1/4/c^(3/2)/b^3*(-Si((d*x+c)^(1/2)*b+b*c^(1/2))*sin(a-b*c^(1/2)
)+Ci((d*x+c)^(1/2)*b+b*c^(1/2))*cos(a-b*c^(1/2)))-1/4/c/b^2*(-Si(b*c^(1/2)
-(d*x+c)^(1/2)*b)*cos(a+b*c^(1/2))+Ci((d*x+c)^(1/2)*b-b*c^(1/2))*sin(a+b*c
^(1/2)))-1/4/c/b^2*(Si((d*x+c)^(1/2)*b+b*c^(1/2))*cos(a-b*c^(1/2))+Ci((d*x
+c)^(1/2)*b+b*c^(1/2))*sin(a-b*c^(1/2))))

```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.14

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx$$

$$= \frac{\sqrt{-b^2cdx}\text{Ei}(i\sqrt{dx + cb} - \sqrt{-b^2c}) e^{(ia + \sqrt{-b^2c})} - \sqrt{-b^2cdx}\text{Ei}(i\sqrt{dx + cb} + \sqrt{-b^2c}) e^{(ia - \sqrt{-b^2c})} + \sqrt{-b^2c}}{c^2}$$

input

```
integrate(cos(a+b*(d*x+c)^(1/2))/x^2,x, algorithm="fricas")
```

output

```

1/4*(sqrt(-b^2*c)*d*x*Ei(I*sqrt(d*x + c)*b - sqrt(-b^2*c))*e^(I*a + sqrt(-
b^2*c)) - sqrt(-b^2*c)*d*x*Ei(I*sqrt(d*x + c)*b + sqrt(-b^2*c))*e^(I*a - s
qrt(-b^2*c)) + sqrt(-b^2*c)*d*x*Ei(-I*sqrt(d*x + c)*b - sqrt(-b^2*c))*e^(-
I*a + sqrt(-b^2*c)) - sqrt(-b^2*c)*d*x*Ei(-I*sqrt(d*x + c)*b + sqrt(-b^2*c
))*e^(-I*a - sqrt(-b^2*c)) - 4*c*cos(sqrt(d*x + c)*b + a))/(c*x)

```

Sympy [F]

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx = \int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx$$

input `integrate(cos(a+b*(d*x+c)**(1/2))/x**2,x)`

output `Integral(cos(a + b*sqrt(c + d*x))/x**2, x)`

Maxima [F]

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx = \int \frac{\cos(\sqrt{dx + cb} + a)}{x^2} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/2))/x^2,x, algorithm="maxima")`

output `integrate(cos(sqrt(d*x + c)*b + a)/x^2, x)`

Giac [F]

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx = \int \frac{\cos(\sqrt{dx + cb} + a)}{x^2} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/2))/x^2,x, algorithm="giac")`

output `integrate(cos(sqrt(d*x + c)*b + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx = \int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx$$

input `int(cos(a + b*(c + d*x)^(1/2))/x^2,x)`output `int(cos(a + b*(c + d*x)^(1/2))/x^2, x)`**Reduce [F]**

$$\int \frac{\cos(a + b\sqrt{c + dx})}{x^2} dx = \int \frac{\cos(\sqrt{dx + c}b + a)}{x^2} dx$$

input `int(cos(a+b*(d*x+c)^(1/2))/x^2,x)`output `int(cos(sqrt(c + d*x)*b + a)/x**2,x)`

3.95 $\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx$

Optimal result	677
Mathematica [C] (verified)	678
Rubi [A] (verified)	679
Maple [B] (verified)	680
Fricas [A] (verification not implemented)	681
Sympy [F]	682
Maxima [B] (verification not implemented)	682
Giac [B] (verification not implemented)	683
Mupad [F(-1)]	684
Reduce [B] (verification not implemented)	685

Optimal result

Integrand size = 18, antiderivative size = 537

$$\begin{aligned}
\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx = & -\frac{720c \cos(a + b\sqrt[3]{c + dx})}{b^6 d^3} \\
& -\frac{120960\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^8 d^3} \\
& +\frac{6c^2\sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
& +\frac{360c(c + dx)^{2/3} \cos(a + b\sqrt[3]{c + dx})}{b^4 d^3} \\
& +\frac{20160(c + dx) \cos(a + b\sqrt[3]{c + dx})}{b^6 d^3} \\
& -\frac{30c(c + dx)^{4/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
& -\frac{1008(c + dx)^{5/3} \cos(a + b\sqrt[3]{c + dx})}{b^4 d^3} \\
& +\frac{24(c + dx)^{7/3} \cos(a + b\sqrt[3]{c + dx})}{b^2 d^3} \\
& +\frac{120960 \sin(a + b\sqrt[3]{c + dx})}{b^9 d^3} - \frac{6c^2 \sin(a + b\sqrt[3]{c + dx})}{b^3 d^3} \\
& -\frac{720c\sqrt[3]{c + dx} \sin(a + b\sqrt[3]{c + dx})}{b^5 d^3} \\
& -\frac{60480(c + dx)^{2/3} \sin(a + b\sqrt[3]{c + dx})}{b^7 d^3} \\
& +\frac{3c^2(c + dx)^{2/3} \sin(a + b\sqrt[3]{c + dx})}{bd^3} \\
& +\frac{120c(c + dx) \sin(a + b\sqrt[3]{c + dx})}{b^3 d^3} \\
& +\frac{5040(c + dx)^{4/3} \sin(a + b\sqrt[3]{c + dx})}{b^5 d^3} \\
& -\frac{6c(c + dx)^{5/3} \sin(a + b\sqrt[3]{c + dx})}{bd^3} \\
& -\frac{168(c + dx)^2 \sin(a + b\sqrt[3]{c + dx})}{b^3 d^3} \\
& +\frac{3(c + dx)^{8/3} \sin(a + b\sqrt[3]{c + dx})}{bd^3}
\end{aligned}$$

output

```
-720*c*cos(a+b*(d*x+c)^(1/3))/b^6/d^3-120960*(d*x+c)^(1/3)*cos(a+b*(d*x+c)^(1/3))/b^8/d^3+6*c^2*(d*x+c)^(1/3)*cos(a+b*(d*x+c)^(1/3))/b^2/d^3+360*c*(d*x+c)^(2/3)*cos(a+b*(d*x+c)^(1/3))/b^4/d^3+20160*(d*x+c)*cos(a+b*(d*x+c)^(1/3))/b^6/d^3-30*c*(d*x+c)^(4/3)*cos(a+b*(d*x+c)^(1/3))/b^2/d^3-1008*(d*x+c)^(5/3)*cos(a+b*(d*x+c)^(1/3))/b^4/d^3+24*(d*x+c)^(7/3)*cos(a+b*(d*x+c)^(1/3))/b^2/d^3+120960*sin(a+b*(d*x+c)^(1/3))/b^9/d^3-6*c^2*sin(a+b*(d*x+c)^(1/3))/b^3/d^3-720*c*(d*x+c)^(1/3)*sin(a+b*(d*x+c)^(1/3))/b^5/d^3-60480*(d*x+c)^(2/3)*sin(a+b*(d*x+c)^(1/3))/b^7/d^3+3*c^2*(d*x+c)^(2/3)*sin(a+b*(d*x+c)^(1/3))/b/d^3+120*c*(d*x+c)*sin(a+b*(d*x+c)^(1/3))/b^3/d^3+5040*(d*x+c)^(4/3)*sin(a+b*(d*x+c)^(1/3))/b^5/d^3-6*c*(d*x+c)^(5/3)*sin(a+b*(d*x+c)^(1/3))/b/d^3-168*(d*x+c)^2*sin(a+b*(d*x+c)^(1/3))/b^3/d^3+3*(d*x+c)^(8/3)*sin(a+b*(d*x+c)^(1/3))/b/d^3
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.26 (sec) , antiderivative size = 382, normalized size of antiderivative = 0.71

$$\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx$$

$$= \frac{3e^{-i(a+b\sqrt[3]{c+dx})} \left(-40320i \left(-1 + e^{2i(a+b\sqrt[3]{c+dx})}\right) - 40320b \left(1 + e^{2i(a+b\sqrt[3]{c+dx})}\right) \sqrt[3]{c + dx} + 20160ib^2 \left(-1 + e^{2i(a+b\sqrt[3]{c+dx})}\right)\right)}{27b^3}$$

input

```
Integrate[x^2*Cos[a + b*(c + d*x)^(1/3)],x]
```

output

```
(3*((-40320*I)*(-1 + E^((2*I)*(a + b*(c + d*x)^(1/3)))) - 40320*b*(1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(c + d*x)^(1/3) + (20160*I)*b^2*(-1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(c + d*x)^(2/3) - I*b^8*d^2*(-1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*x^2*(c + d*x)^(2/3) + 2*b^7*d*(1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*x*(c + d*x)^(1/3)*(3*c + 4*d*x) - (240*I)*b^4*(-1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(c + d*x)^(1/3)*(6*c + 7*d*x) - 24*b^5*(1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(c + d*x)^(2/3)*(9*c + 14*d*x) + 240*b^3*(1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(27*c + 28*d*x) + (2*I)*b^6*(-1 + E^((2*I)*(a + b*(c + d*x)^(1/3))))*(9*c^2 + 36*c*d*x + 28*d^2*x^2))/(2*b^9*d^3*E^(I*(a + b*(c + d*x)^(1/3))))
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 540, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3913, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx$$

$$\downarrow \text{3913}$$

$$3 \int \left(\frac{\cos(a + b\sqrt[3]{c + dx})(c + dx)^{8/3}}{d^2} - \frac{2c \cos(a + b\sqrt[3]{c + dx})(c + dx)^{5/3}}{d^2} + \frac{c^2 \cos(a + b\sqrt[3]{c + dx})(c + dx)^{2/3}}{d^2} \right) d\sqrt[3]{c + dx}$$

$$\downarrow \text{2009}$$

$$3 \left(\frac{40320 \sin(a + b\sqrt[3]{c + dx})}{b^9 d^2} - \frac{40320 \sqrt[3]{c + dx} \cos(a + b\sqrt[3]{c + dx})}{b^8 d^2} - \frac{20160(c + dx)^{2/3} \sin(a + b\sqrt[3]{c + dx})}{b^7 d^2} + \frac{6720(c + dx) \cos(a + b\sqrt[3]{c + dx})}{b^6 d^2} \right)$$

input `Int[x^2*Cos[a + b*(c + d*x)^(1/3)],x]`

output

$$\begin{aligned} & (3*((-240*c*\text{Cos}[a + b*(c + d*x)^{(1/3)}])/(b^6*d^2) - (40320*(c + d*x)^{(1/3)} \\ & * \text{Cos}[a + b*(c + d*x)^{(1/3)}])/(b^8*d^2) + (2*c^2*(c + d*x)^{(1/3)}*\text{Cos}[a + b* \\ & (c + d*x)^{(1/3)}])/(b^2*d^2) + (120*c*(c + d*x)^{(2/3)}*\text{Cos}[a + b*(c + d*x)^{(1/3)}])/(b^4*d^2) + (6720*(c + d*x)*\text{Cos}[a + b*(c + d*x)^{(1/3)}])/(b^6*d^2) - \\ & (10*c*(c + d*x)^{(4/3)}*\text{Cos}[a + b*(c + d*x)^{(1/3)}])/(b^2*d^2) - (336*(c + d \\ & *x)^{(5/3)}*\text{Cos}[a + b*(c + d*x)^{(1/3)}])/(b^4*d^2) + (8*(c + d*x)^{(7/3)}*\text{Cos}[a \\ & + b*(c + d*x)^{(1/3)}])/(b^2*d^2) + (40320*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(b^9 \\ & *d^2) - (2*c^2*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(b^3*d^2) - (240*c*(c + d*x)^{(1/3)}*\text{Sin}[a + \\ & b*(c + d*x)^{(1/3)}])/(b^5*d^2) - (20160*(c + d*x)^{(2/3)}*\text{Sin}[a + \\ & b*(c + d*x)^{(1/3)}])/(b^7*d^2) + (c^2*(c + d*x)^{(2/3)}*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(b*d^2) + (40*c*(c + d*x)*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(b^3*d^2) + \\ & (1680*(c + d*x)^{(4/3)}*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(b^5*d^2) - (2*c*(c + d \\ & x)^{(5/3)}*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(b*d^2) - (56*(c + d*x)^2*\text{Sin}[a + b*(\\ & c + d*x)^{(1/3)}])/(b^3*d^2) + ((c + d*x)^{(8/3)}*\text{Sin}[a + b*(c + d*x)^{(1/3)}])/(\\ & (b*d^2)))/d \end{aligned}$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 3913

$$\begin{aligned} & \text{Int}[\{(a_.) + \text{Cos}[(c_.) + (d_.)*\{(e_.) + (f_.)*(x_.)\}^{(n_.)}]*\{(b_.)\}^{(p_.)}*\{(g_.) \\ & + (h_.)*(x_.)\}^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[1/(n*f) \text{ Subst}[\text{Int}[\text{ExpandIntegra} \\ & \text{nd}[(a + b*\text{Cos}[c + d*x])^p, x^{(1/n - 1)}*(g - e*(h/f) + h*(x^{(1/n)}/f))^m, x], \\ & x], x, (e + f*x)^n], x] \text{ ; FreeQ}\{a, b, c, d, e, f, g, h, m\}, x\} \&\& \text{IGtQ}[p \\ & , 0] \&\& \text{IntegerQ}[1/n] \end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1808 vs. $2(477) = 954$.

Time = 1.55 (sec) , antiderivative size = 1809, normalized size of antiderivative = 3.37

method	result	size
derivativeldivides	Expression too large to display	1809
default	Expression too large to display	1809
parts	Expression too large to display	2944

```
input int(x^2*cos(a+b*(d*x+c)^(1/3)),x,method=_RETURNVERBOSE)
```

```
output 3/d^3/b^3*(a^2*c^2*sin(a+b*(d*x+c)^(1/3))-2*a*c^2*(cos(a+b*(d*x+c)^(1/3))+
(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))+c^2*((a+b*(d*x+c)^(1/3))^2*sin
(a+b*(d*x+c)^(1/3))-2*sin(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*cos(a+b
*(d*x+c)^(1/3)))+2/b^3*a^5*c*sin(a+b*(d*x+c)^(1/3))-10/b^3*a^4*c*(cos(a+b*
(d*x+c)^(1/3))+(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))+20/b^3*a^3*c*((
a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1/3))-2*sin(a+b*(d*x+c)^(1/3))+2*(a+
b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3)))-20/b^3*a^2*c*((a+b*(d*x+c)^(1/3))
^3*sin(a+b*(d*x+c)^(1/3))+3*(a+b*(d*x+c)^(1/3))^2*cos(a+b*(d*x+c)^(1/3))-6
*cos(a+b*(d*x+c)^(1/3))-6*(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))+10/b
^3*a*c*((a+b*(d*x+c)^(1/3))^4*sin(a+b*(d*x+c)^(1/3))+4*(a+b*(d*x+c)^(1/3))
^3*cos(a+b*(d*x+c)^(1/3))-12*(a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1/3))+
24*sin(a+b*(d*x+c)^(1/3))-24*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3))-2
/b^3*c*((a+b*(d*x+c)^(1/3))^5*sin(a+b*(d*x+c)^(1/3))+5*(a+b*(d*x+c)^(1/3))
^4*cos(a+b*(d*x+c)^(1/3))-20*(a+b*(d*x+c)^(1/3))^3*sin(a+b*(d*x+c)^(1/3))-
60*(a+b*(d*x+c)^(1/3))^2*cos(a+b*(d*x+c)^(1/3))+120*cos(a+b*(d*x+c)^(1/3))
+120*(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))+1/b^6*a^8*sin(a+b*(d*x+c)
^(1/3))-8/b^6*a^7*(cos(a+b*(d*x+c)^(1/3))+(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x
+c)^(1/3)))+28/b^6*a^6*((a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1/3))-2*sin
(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3)))-56/b^6*a
^5*((a+b*(d*x+c)^(1/3))^3*sin(a+b*(d*x+c)^(1/3))+3*(a+b*(d*x+c)^(1/3))^...
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.34

$$\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx$$

$$= \frac{3 \left(2 \left(3360 b^3 dx + 3240 b^3 c - 12 (14 b^5 dx + 9 b^5 c)(dx + c)^{\frac{2}{3}} + (4 b^7 d^2 x^2 + 3 b^7 c dx - 20160 b)(dx + c)^{\frac{1}{3}} \right) \right)}{...}$$

```
input integrate(x^2*cos(a+b*(d*x+c)^(1/3)),x, algorithm="fricas")
```

output

```
3*(2*(3360*b^3*d*x + 3240*b^3*c - 12*(14*b^5*d*x + 9*b^5*c)*(d*x + c)^(2/3)
) + (4*b^7*d^2*x^2 + 3*b^7*c*d*x - 20160*b)*(d*x + c)^(1/3))*cos((d*x + c)
^(1/3)*b + a) - (56*b^6*d^2*x^2 + 72*b^6*c*d*x + 18*b^6*c^2 - (b^8*d^2*x^2
- 20160*b^2)*(d*x + c)^(2/3) - 240*(7*b^4*d*x + 6*b^4*c)*(d*x + c)^(1/3)
- 40320)*sin((d*x + c)^(1/3)*b + a))/(b^9*d^3)
```

Sympy [F]

$$\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx = \int x^2 \cos(a + b\sqrt[3]{c + dx}) dx$$

input

```
integrate(x**2*cos(a+b*(d*x+c)**(1/3)),x)
```

output

```
Integral(x**2*cos(a + b*(c + d*x)**(1/3)), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1349 vs. $2(477) = 954$.

Time = 0.09 (sec) , antiderivative size = 1349, normalized size of antiderivative = 2.51

$$\int x^2 \cos(a + b\sqrt[3]{c + dx}) dx = \text{Too large to display}$$

input

```
integrate(x^2*cos(a+b*(d*x+c)^(1/3)),x, algorithm="maxima")
```

output

```

3*(a^2*c^2*sin((d*x + c)^(1/3)*b + a) - 2*(((d*x + c)^(1/3)*b + a)*sin((d*x
+ c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a*c^2 + 2*a^5*c*sin((d*x
+ c)^(1/3)*b + a)/b^3 - 10*(((d*x + c)^(1/3)*b + a)*sin((d*x + c)^(1/3)*b
+ a) + cos((d*x + c)^(1/3)*b + a))*a^4*c/b^3 + (2*((d*x + c)^(1/3)*b + a)
*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c
)^(1/3)*b + a))*c^2 + a^8*sin((d*x + c)^(1/3)*b + a)/b^6 - 8*(((d*x + c)^(
1/3)*b + a)*sin((d*x + c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a^7/b
^6 + 20*(2*((d*x + c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c
)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b + a))*a^3*c/b^3 + 28*(2*((d*x
+ c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2
- 2)*sin((d*x + c)^(1/3)*b + a))*a^6/b^6 - 20*(3*(((d*x + c)^(1/3)*b + a)^
2 - 2)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^3 - 6*(d*x +
c)^(1/3)*b - 6*a)*sin((d*x + c)^(1/3)*b + a))*a^2*c/b^3 - 56*(3*(((d*x + c
)^(1/3)*b + a)^2 - 2)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a
)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*sin((d*x + c)^(1/3)*b + a))*a^5/b^6 + 10*
(4*(((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*cos((d*x + c)^(
1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^4 - 12*((d*x + c)^(1/3)*b + a)^2 +
24)*sin((d*x + c)^(1/3)*b + a))*a*c/b^3 + 70*(4*(((d*x + c)^(1/3)*b + a)^3
- 6*(d*x + c)^(1/3)*b - 6*a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/
3)*b + a)^4 - 12*((d*x + c)^(1/3)*b + a)^2 + 24)*sin((d*x + c)^(1/3)*b ...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1098 vs. $2(477) = 954$.

Time = 0.36 (sec) , antiderivative size = 1098, normalized size of antiderivative = 2.04

$$\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx = \text{Too large to display}$$

input

```
integrate(x^2*cos(a+b*(d*x+c)^(1/3)),x, algorithm="giac")
```

output

```

3*(2*((d*x + c)^(1/3)*b + a)*b^6*c^2 - a*b^6*c^2 - 5*((d*x + c)^(1/3)*b +
a)^4*b^3*c + 20*((d*x + c)^(1/3)*b + a)^3*a*b^3*c - 30*((d*x + c)^(1/3)*b
+ a)^2*a^2*b^3*c + 20*((d*x + c)^(1/3)*b + a)*a^3*b^3*c - 5*a^4*b^3*c + 4
*((d*x + c)^(1/3)*b + a)^7 - 28*((d*x + c)^(1/3)*b + a)^6*a + 84*((d*x + c)
^(1/3)*b + a)^5*a^2 - 140*((d*x + c)^(1/3)*b + a)^4*a^3 + 140*((d*x + c)
^(1/3)*b + a)^3*a^4 - 84*((d*x + c)^(1/3)*b + a)^2*a^5 + 28*((d*x + c)^(1/3)
)*b + a)*a^6 - 4*a^7 + 60*((d*x + c)^(1/3)*b + a)^2*b^3*c - 120*((d*x + c)
^(1/3)*b + a)*a*b^3*c + 60*a^2*b^3*c - 168*((d*x + c)^(1/3)*b + a)^5 + 840
*((d*x + c)^(1/3)*b + a)^4*a - 1680*((d*x + c)^(1/3)*b + a)^3*a^2 + 1680*(
(d*x + c)^(1/3)*b + a)^2*a^3 - 840*((d*x + c)^(1/3)*b + a)*a^4 + 168*a^5 -
120*b^3*c + 3360*((d*x + c)^(1/3)*b + a)^3 - 10080*((d*x + c)^(1/3)*b + a)
^2*a + 10080*((d*x + c)^(1/3)*b + a)*a^2 - 3360*a^3 - 20160*(d*x + c)^(1/
3)*b)*cos((d*x + c)^(1/3)*b + a)/b^8 + (((d*x + c)^(1/3)*b + a)^2*b^6*c^2
- 2*((d*x + c)^(1/3)*b + a)*a*b^6*c^2 + a^2*b^6*c^2 - 2*((d*x + c)^(1/3)*b
+ a)^5*b^3*c + 10*((d*x + c)^(1/3)*b + a)^4*a*b^3*c - 20*((d*x + c)^(1/3)
*b + a)^3*a^2*b^3*c + 20*((d*x + c)^(1/3)*b + a)^2*a^3*b^3*c - 10*((d*x +
c)^(1/3)*b + a)*a^4*b^3*c + 2*a^5*b^3*c + ((d*x + c)^(1/3)*b + a)^8 - 8*((
d*x + c)^(1/3)*b + a)^7*a + 28*((d*x + c)^(1/3)*b + a)^6*a^2 - 56*((d*x +
c)^(1/3)*b + a)^5*a^3 + 70*((d*x + c)^(1/3)*b + a)^4*a^4 - 56*((d*x + c)^(
1/3)*b + a)^3*a^5 + 28*((d*x + c)^(1/3)*b + a)^2*a^6 - 8*((d*x + c)^(1/...

```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx = \int x^2 \cos\left(a + b(c + dx)^{1/3}\right) dx$$

input

```
int(x^2*cos(a + b*(c + d*x)^(1/3)),x)
```

output

```
int(x^2*cos(a + b*(c + d*x)^(1/3)), x)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 357, normalized size of antiderivative = 0.66

$$\int x^2 \cos\left(a + b\sqrt[3]{c + dx}\right) dx$$

$$= \frac{-648(dx + c)^{\frac{2}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^5 c - 1008(dx + c)^{\frac{2}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^5 dx + 18(dx + c)^{\frac{1}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^5 dx + 18(dx + c)^{\frac{1}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^5 dx}{1}$$

input

```
int(x^2*cos(a+b*(d*x+c)^(1/3)),x)
```

output

```
(3*(- 216*(c + d*x)**(2/3)*cos((c + d*x)**(1/3)*b + a)*b**5*c - 336*(c +
d*x)**(2/3)*cos((c + d*x)**(1/3)*b + a)*b**5*d*x + 6*(c + d*x)**(1/3)*cos(
(c + d*x)**(1/3)*b + a)*b**7*c*d*x + 8*(c + d*x)**(1/3)*cos((c + d*x)**(1/
3)*b + a)*b**7*d**2*x**2 - 40320*(c + d*x)**(1/3)*cos((c + d*x)**(1/3)*b +
a)*b + 6480*cos((c + d*x)**(1/3)*b + a)*b**3*c + 6720*cos((c + d*x)**(1/3
)*b + a)*b**3*d*x + (c + d*x)**(2/3)*sin((c + d*x)**(1/3)*b + a)*b**8*d**2
*x**2 - 20160*(c + d*x)**(2/3)*sin((c + d*x)**(1/3)*b + a)*b**2 + 1440*(c
+ d*x)**(1/3)*sin((c + d*x)**(1/3)*b + a)*b**4*c + 1680*(c + d*x)**(1/3)*s
in((c + d*x)**(1/3)*b + a)*b**4*d*x - 18*sin((c + d*x)**(1/3)*b + a)*b**6*
c**2 - 72*sin((c + d*x)**(1/3)*b + a)*b**6*c*d*x - 56*sin((c + d*x)**(1/3
)*b + a)*b**6*d**2*x**2 + 40320*sin((c + d*x)**(1/3)*b + a)))/(b**9*d**3)
```

3.96 $\int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx$

Optimal result	686
Mathematica [A] (verified)	687
Rubi [A] (verified)	687
Maple [B] (verified)	689
Fricas [A] (verification not implemented)	690
Sympy [F]	690
Maxima [B] (verification not implemented)	690
Giac [A] (verification not implemented)	691
Mupad [F(-1)]	692
Reduce [B] (verification not implemented)	692

Optimal result

Integrand size = 16, antiderivative size = 261

$$\begin{aligned}
 \int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx = & \frac{360 \cos \left(a + b\sqrt[3]{c + dx} \right)}{b^6 d^2} - \frac{6c\sqrt[3]{c + dx} \cos \left(a + b\sqrt[3]{c + dx} \right)}{b^2 d^2} \\
 & - \frac{180(c + dx)^{2/3} \cos \left(a + b\sqrt[3]{c + dx} \right)}{b^4 d^2} \\
 & + \frac{15(c + dx)^{4/3} \cos \left(a + b\sqrt[3]{c + dx} \right)}{b^2 d^2} \\
 & + \frac{6c \sin \left(a + b\sqrt[3]{c + dx} \right)}{b^3 d^2} \\
 & + \frac{360\sqrt[3]{c + dx} \sin \left(a + b\sqrt[3]{c + dx} \right)}{b^5 d^2} \\
 & - \frac{3c(c + dx)^{2/3} \sin \left(a + b\sqrt[3]{c + dx} \right)}{bd^2} \\
 & - \frac{60(c + dx) \sin \left(a + b\sqrt[3]{c + dx} \right)}{b^3 d^2} \\
 & + \frac{3(c + dx)^{5/3} \sin \left(a + b\sqrt[3]{c + dx} \right)}{bd^2}
 \end{aligned}$$

output

```
360*cos(a+b*(d*x+c)^(1/3))/b^6/d^2-6*c*(d*x+c)^(1/3)*cos(a+b*(d*x+c)^(1/3)
)/b^2/d^2-180*(d*x+c)^(2/3)*cos(a+b*(d*x+c)^(1/3))/b^4/d^2+15*(d*x+c)^(4/3
)*cos(a+b*(d*x+c)^(1/3))/b^2/d^2+6*c*sin(a+b*(d*x+c)^(1/3))/b^3/d^2+360*(d
*x+c)^(1/3)*sin(a+b*(d*x+c)^(1/3))/b^5/d^2-3*c*(d*x+c)^(2/3)*sin(a+b*(d*x+
c)^(1/3))/b/d^2-60*(d*x+c)*sin(a+b*(d*x+c)^(1/3))/b^3/d^2+3*(d*x+c)^(5/3)*
sin(a+b*(d*x+c)^(1/3))/b/d^2
```

Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.45

$$\int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \frac{3 \left((120 - 60b^2(c + dx)^{2/3} + b^4\sqrt[3]{c + dx}(3c + 5dx)) \cos \left(a + b\sqrt[3]{c + dx} \right) + b \left(120\sqrt[3]{c + dx} + b^4 dx(c + dx) \right) \right)}{b^6 d^2}$$

input

```
Integrate[x*Cos[a + b*(c + d*x)^(1/3)],x]
```

output

```
(3*((120 - 60*b^2*(c + d*x)^(2/3) + b^4*(c + d*x)^(1/3)*(3*c + 5*d*x))*Cos
[a + b*(c + d*x)^(1/3)] + b*(120*(c + d*x)^(1/3) + b^4*d*x*(c + d*x)^(2/3)
- 2*b^2*(9*c + 10*d*x))*Sin[a + b*(c + d*x)^(1/3)]))/(b^6*d^2)
```

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.02, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3913, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

↓ 3913

$$3 \int \left(\frac{(c+dx)^{5/3} \cos\left(a+b\sqrt[3]{c+dx}\right)}{d} - \frac{c(c+dx)^{2/3} \cos\left(a+b\sqrt[3]{c+dx}\right)}{d} \right) d\sqrt[3]{c+dx}$$

↓ 2009

$$3 \left(\frac{120 \cos\left(a+b\sqrt[3]{c+dx}\right)}{b^6 d} + \frac{120 \sqrt[3]{c+dx} \sin\left(a+b\sqrt[3]{c+dx}\right)}{b^5 d} - \frac{60(c+dx)^{2/3} \cos\left(a+b\sqrt[3]{c+dx}\right)}{b^4 d} - \frac{20(c+dx) \sin\left(a+b\sqrt[3]{c+dx}\right)}{b^3 d} \right)$$

input `Int[x*Cos[a + b*(c + d*x)^(1/3)],x]`

output

```
(3*((120*Cos[a + b*(c + d*x)^(1/3)])/(b^6*d) - (2*c*(c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d) - (60*(c + d*x)^(2/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^4*d) + (5*(c + d*x)^(4/3)*Cos[a + b*(c + d*x)^(1/3)]/(b^2*d) + (2*c*Sin[a + b*(c + d*x)^(1/3)])/(b^3*d) + (120*(c + d*x)^(1/3)*Sin[a + b*(c + d*x)^(1/3)]/(b^5*d) - (c*(c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d) - (20*(c + d*x)*Sin[a + b*(c + d*x)^(1/3)]/(b^3*d) + ((c + d*x)^(5/3)*Sin[a + b*(c + d*x)^(1/3)]/(b*d)))/d
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 3913

```
Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_.)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Simp[1/(n*f) Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - e*(h/f) + h*(x^(1/n)/f))^m, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 654 vs. $2(231) = 462$.

Time = 1.73 (sec) , antiderivative size = 655, normalized size of antiderivative = 2.51

method	result
derivativedivides	$\frac{-3a^2c \sin(a+b(dx+c)^{\frac{1}{3}}) + 6ac \left(\cos(a+b(dx+c)^{\frac{1}{3}}) + (a+b(dx+c)^{\frac{1}{3}}) \sin(a+b(dx+c)^{\frac{1}{3}}) \right) - 3c \left((a+b(dx+c)^{\frac{1}{3}})^2 \sin(a+b(dx+c)^{\frac{1}{3}}) \right)}{-3a^2c \sin(a+b(dx+c)^{\frac{1}{3}}) + 6ac \left(\cos(a+b(dx+c)^{\frac{1}{3}}) + (a+b(dx+c)^{\frac{1}{3}}) \sin(a+b(dx+c)^{\frac{1}{3}}) \right) - 3c \left((a+b(dx+c)^{\frac{1}{3}})^2 \sin(a+b(dx+c)^{\frac{1}{3}}) \right)}$
default	$\frac{-3a^2c \sin(a+b(dx+c)^{\frac{1}{3}}) + 6ac \left(\cos(a+b(dx+c)^{\frac{1}{3}}) + (a+b(dx+c)^{\frac{1}{3}}) \sin(a+b(dx+c)^{\frac{1}{3}}) \right) - 3c \left((a+b(dx+c)^{\frac{1}{3}})^2 \sin(a+b(dx+c)^{\frac{1}{3}}) \right)}{-3a^2c \sin(a+b(dx+c)^{\frac{1}{3}}) + 6ac \left(\cos(a+b(dx+c)^{\frac{1}{3}}) + (a+b(dx+c)^{\frac{1}{3}}) \sin(a+b(dx+c)^{\frac{1}{3}}) \right) - 3c \left((a+b(dx+c)^{\frac{1}{3}})^2 \sin(a+b(dx+c)^{\frac{1}{3}}) \right)}$
parts	Expression too large to display

input

```
int(x*cos(a+b*(d*x+c)^(1/3)),x,method=_RETURNVERBOSE)
```

output

```
3/d^2/b^3*(-a^2*c*sin(a+b*(d*x+c)^(1/3))+2*a*c*(cos(a+b*(d*x+c)^(1/3))+(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))-c*((a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1/3))-2*sin(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3)))-1/b^3*a^5*sin(a+b*(d*x+c)^(1/3))+5/b^3*a^4*(cos(a+b*(d*x+c)^(1/3))+(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))-10/b^3*a^3*((a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1/3))-2*sin(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3)))+10/b^3*a^2*((a+b*(d*x+c)^(1/3))^3*sin(a+b*(d*x+c)^(1/3))+3*(a+b*(d*x+c)^(1/3))^2*cos(a+b*(d*x+c)^(1/3))-6*cos(a+b*(d*x+c)^(1/3))-6*(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))-5/b^3*a*((a+b*(d*x+c)^(1/3))^4*sin(a+b*(d*x+c)^(1/3))+4*(a+b*(d*x+c)^(1/3))^3*cos(a+b*(d*x+c)^(1/3))-12*(a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1/3))+24*sin(a+b*(d*x+c)^(1/3))-24*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3)))+1/b^3*((a+b*(d*x+c)^(1/3))^5*sin(a+b*(d*x+c)^(1/3))+5*(a+b*(d*x+c)^(1/3))^4*cos(a+b*(d*x+c)^(1/3))-20*(a+b*(d*x+c)^(1/3))^3*sin(a+b*(d*x+c)^(1/3))-60*(a+b*(d*x+c)^(1/3))^2*cos(a+b*(d*x+c)^(1/3))+120*cos(a+b*(d*x+c)^(1/3))+120*(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.42

$$\int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx = \frac{3 \left(\left(60 (dx + c)^{\frac{2}{3}} b^2 - (5 b^4 dx + 3 b^4 c) (dx + c)^{\frac{1}{3}} - 120 \right) \cos \left((dx + c)^{\frac{1}{3}} b + a \right) - \left((dx + c)^{\frac{2}{3}} b^5 dx - 20 b^3 c \right) \sin \left((dx + c)^{\frac{1}{3}} b + a \right) \right)}{b^6 d^2}$$

input `integrate(x*cos(a+b*(d*x+c)^(1/3)),x, algorithm="fricas")`

output `-3*((60*(d*x + c)^(2/3)*b^2 - (5*b^4*d*x + 3*b^4*c)*(d*x + c)^(1/3) - 120)*cos((d*x + c)^(1/3)*b + a) - ((d*x + c)^(2/3)*b^5*d*x - 20*b^3*d*x - 18*b^3*c + 120*(d*x + c)^(1/3)*b)*sin((d*x + c)^(1/3)*b + a))/(b^6*d^2)`

Sympy [F]

$$\int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx = \int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

input `integrate(x*cos(a+b*(d*x+c)**(1/3)),x)`

output `Integral(x*cos(a + b*(c + d*x)**(1/3)), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 523 vs. 2(231) = 462.

Time = 0.05 (sec) , antiderivative size = 523, normalized size of antiderivative = 2.00

$$\int x \cos \left(a + b\sqrt[3]{c + dx} \right) dx = \text{Too large to display}$$

input `integrate(x*cos(a+b*(d*x+c)^(1/3)),x, algorithm="maxima")`

output

```

-3*(a^2*c*sin((d*x + c)^(1/3)*b + a) - 2*(((d*x + c)^(1/3)*b + a)*sin((d*x
+ c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a*c + a^5*sin((d*x + c)^(
1/3)*b + a)/b^3 - 5*(((d*x + c)^(1/3)*b + a)*sin((d*x + c)^(1/3)*b + a) +
cos((d*x + c)^(1/3)*b + a))*a^4/b^3 + (2*(((d*x + c)^(1/3)*b + a)*cos((d*x
+ c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b
+ a))*c + 10*(2*(((d*x + c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*
x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b + a))*a^3/b^3 - 10*(3*(((
d*x + c)^(1/3)*b + a)^2 - 2)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3
)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*sin((d*x + c)^(1/3)*b + a))*a^2/b^
3 + 5*(4*(((d*x + c)^(1/3)*b + a)^3 - 6*(d*x + c)^(1/3)*b - 6*a)*cos((d*x
+ c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^4 - 12*(d*x + c)^(1/3)*b + a
)^2 + 24)*sin((d*x + c)^(1/3)*b + a))*a/b^3 - (5*(((d*x + c)^(1/3)*b + a)^
4 - 12*(d*x + c)^(1/3)*b + a)^2 + 24)*cos((d*x + c)^(1/3)*b + a) + (((d*x
+ c)^(1/3)*b + a)^5 - 20*(d*x + c)^(1/3)*b + a)^3 + 120*(d*x + c)^(1/3)*
b + 120*a)*sin((d*x + c)^(1/3)*b + a))/b^3)/(b^3*d^2)

```

Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 370, normalized size of antiderivative = 1.42

$$\int x \cos\left(a + b\sqrt[3]{c + dx}\right) dx =$$

$$\frac{3 \left(\left(2 \left((dx+c)^{\frac{1}{3}}b+a \right) b^3 c - 2 a b^3 c - 5 \left((dx+c)^{\frac{1}{3}}b+a \right)^4 + 20 \left((dx+c)^{\frac{1}{3}}b+a \right)^3 a - 30 \left((dx+c)^{\frac{1}{3}}b+a \right)^2 a^2 + 20 \left((dx+c)^{\frac{1}{3}}b+a \right) a^3 - 5 a^4 + 60 \left((dx+c)^{\frac{1}{3}}b+a \right) a^5 - 12 \left((dx+c)^{\frac{1}{3}}b+a \right)^2 + 24 \right) \sin\left((dx+c)^{\frac{1}{3}}b+a \right) a/b^3 - \left(5 \left((dx+c)^{\frac{1}{3}}b+a \right)^4 - 12 \left((dx+c)^{\frac{1}{3}}b+a \right)^2 + 24 \right) \cos\left((dx+c)^{\frac{1}{3}}b+a \right) + \left((dx+c)^{\frac{1}{3}}b+a \right)^5 - 20 \left((dx+c)^{\frac{1}{3}}b+a \right)^3 + 120 \left((dx+c)^{\frac{1}{3}}b+a \right) a \right) \sin\left((dx+c)^{\frac{1}{3}}b+a \right) / b^3}{b^5}$$

input

```
integrate(x*cos(a+b*(d*x+c)^(1/3)),x, algorithm="giac")
```

output

```
-3*((2*((d*x + c)^(1/3)*b + a)*b^3*c - 2*a*b^3*c - 5*((d*x + c)^(1/3)*b +
a)^4 + 20*((d*x + c)^(1/3)*b + a)^3*a - 30*((d*x + c)^(1/3)*b + a)^2*a^2 +
20*((d*x + c)^(1/3)*b + a)*a^3 - 5*a^4 + 60*((d*x + c)^(1/3)*b + a)^2 - 1
20*((d*x + c)^(1/3)*b + a)*a + 60*a^2 - 120)*cos((d*x + c)^(1/3)*b + a)/b^
5 + (((d*x + c)^(1/3)*b + a)^2*b^3*c - 2*((d*x + c)^(1/3)*b + a)*a*b^3*c +
a^2*b^3*c - ((d*x + c)^(1/3)*b + a)^5 + 5*((d*x + c)^(1/3)*b + a)^4*a - 1
0*((d*x + c)^(1/3)*b + a)^3*a^2 + 10*((d*x + c)^(1/3)*b + a)^2*a^3 - 5*((d
*x + c)^(1/3)*b + a)*a^4 + a^5 - 2*b^3*c + 20*((d*x + c)^(1/3)*b + a)^3 -
60*((d*x + c)^(1/3)*b + a)^2*a + 60*((d*x + c)^(1/3)*b + a)*a^2 - 20*a^3 -
120*(d*x + c)^(1/3)*b)*sin((d*x + c)^(1/3)*b + a)/b^5)/(b*d^2)
```

Mupad [F(-1)]

Timed out.

$$\int x \cos(a + b\sqrt[3]{c + dx}) dx = \int x \cos(a + b(c + dx)^{1/3}) dx$$

input

```
int(x*cos(a + b*(c + d*x)^(1/3)),x)
```

output

```
int(x*cos(a + b*(c + d*x)^(1/3)), x)
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.70

$$\int x \cos(a + b\sqrt[3]{c + dx}) dx$$

$$= \frac{-180(dx + c)^{\frac{2}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^2 + 9(dx + c)^{\frac{1}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^4 c + 15(dx + c)^{\frac{1}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^4 c + 15(dx + c)^{\frac{1}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^4 c + 15(dx + c)^{\frac{1}{3}} \cos\left((dx + c)^{\frac{1}{3}} b + a\right) b^4 c}{b^5 d^2}$$

input

```
int(x*cos(a+b*(d*x+c)^(1/3)),x)
```

output

```
(3*( - 60*(c + d*x)**(2/3)*cos((c + d*x)**(1/3)*b + a)*b**2 + 3*(c + d*x)*
*(1/3)*cos((c + d*x)**(1/3)*b + a)*b**4*c + 5*(c + d*x)**(1/3)*cos((c + d*
x)**(1/3)*b + a)*b**4*d*x + 120*cos((c + d*x)**(1/3)*b + a) + (c + d*x)**(
2/3)*sin((c + d*x)**(1/3)*b + a)*b**5*d*x + 120*(c + d*x)**(1/3)*sin((c +
d*x)**(1/3)*b + a)*b - 18*sin((c + d*x)**(1/3)*b + a)*b**3*c - 20*sin((c +
d*x)**(1/3)*b + a)*b**3*d*x))/(b**6*d**2)
```

3.97 $\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$

Optimal result	694
Mathematica [A] (verified)	694
Rubi [A] (verified)	695
Maple [A] (verified)	697
Fricas [A] (verification not implemented)	698
Sympy [A] (verification not implemented)	698
Maxima [A] (verification not implemented)	699
Giac [A] (verification not implemented)	699
Mupad [B] (verification not implemented)	700
Reduce [B] (verification not implemented)	700

Optimal result

Integrand size = 14, antiderivative size = 85

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx = \frac{6\sqrt[3]{c + dx} \cos \left(a + b\sqrt[3]{c + dx} \right)}{b^2 d} - \frac{6 \sin \left(a + b\sqrt[3]{c + dx} \right)}{b^3 d} + \frac{3(c + dx)^{2/3} \sin \left(a + b\sqrt[3]{c + dx} \right)}{bd}$$

output

```
6*(d*x+c)^(1/3)*cos(a+b*(d*x+c)^(1/3))/b^2/d-6*sin(a+b*(d*x+c)^(1/3))/b^3/d+3*(d*x+c)^(2/3)*sin(a+b*(d*x+c)^(1/3))/b/d
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.76

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx = \frac{6b\sqrt[3]{c + dx} \cos \left(a + b\sqrt[3]{c + dx} \right) + 3(-2 + b^2(c + dx)^{2/3}) \sin \left(a + b\sqrt[3]{c + dx} \right)}{b^3 d}$$

input

```
Integrate[Cos[a + b*(c + d*x)^(1/3)], x]
```

output

$$(6*b*(c + d*x)^{(1/3)}*Cos[a + b*(c + d*x)^{(1/3)}] + 3*(-2 + b^2*(c + d*x)^{(2/3)})*Sin[a + b*(c + d*x)^{(1/3)}])/(b^3*d)$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3843, 3042, 3777, 25, 3042, 3777, 3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$\downarrow \text{3843}$$

$$\frac{3 \int (c + dx)^{2/3} \cos \left(a + b\sqrt[3]{c + dx} \right) d\sqrt[3]{c + dx}}{d}$$

$$\downarrow \text{3042}$$

$$\frac{3 \int (c + dx)^{2/3} \sin \left(a + b\sqrt[3]{c + dx} + \frac{\pi}{2} \right) d\sqrt[3]{c + dx}}{d}$$

$$\downarrow \text{3777}$$

$$\frac{3 \left(\frac{2 \int -\sqrt[3]{c + dx} \sin \left(a + b\sqrt[3]{c + dx} \right) d\sqrt[3]{c + dx}}{b} + \frac{(c + dx)^{2/3} \sin \left(a + b\sqrt[3]{c + dx} \right)}{b} \right)}{d}$$

$$\downarrow \text{25}$$

$$\frac{3 \left(\frac{(c + dx)^{2/3} \sin \left(a + b\sqrt[3]{c + dx} \right)}{b} - \frac{2 \int \sqrt[3]{c + dx} \sin \left(a + b\sqrt[3]{c + dx} \right) d\sqrt[3]{c + dx}}{b} \right)}{d}$$

$$\downarrow \text{3042}$$

$$\frac{3 \left(\frac{(c + dx)^{2/3} \sin \left(a + b\sqrt[3]{c + dx} \right)}{b} - \frac{2 \int \sqrt[3]{c + dx} \sin \left(a + b\sqrt[3]{c + dx} \right) d\sqrt[3]{c + dx}}{b} \right)}{d}$$

$$\begin{array}{c}
 \downarrow \text{3777} \\
 \frac{3 \left(\frac{(c+dx)^{2/3} \sin\left(a+b\sqrt[3]{c+dx}\right)}{b} - \frac{2 \left(\frac{\int \cos\left(a+b\sqrt[3]{c+dx}\right) d\sqrt[3]{c+dx}}{b} - \frac{\sqrt[3]{c+dx} \cos\left(a+b\sqrt[3]{c+dx}\right)}{b} \right)}{b} \right)}{d} \\
 \downarrow \text{3042} \\
 \frac{3 \left(\frac{(c+dx)^{2/3} \sin\left(a+b\sqrt[3]{c+dx}\right)}{b} - \frac{2 \left(\frac{\int \sin\left(a+b\sqrt[3]{c+dx} + \frac{\pi}{2}\right) d\sqrt[3]{c+dx}}{b} - \frac{\sqrt[3]{c+dx} \cos\left(a+b\sqrt[3]{c+dx}\right)}{b} \right)}{b} \right)}{d} \\
 \downarrow \text{3117} \\
 \frac{3 \left(\frac{(c+dx)^{2/3} \sin\left(a+b\sqrt[3]{c+dx}\right)}{b} - \frac{2 \left(\frac{\sin\left(a+b\sqrt[3]{c+dx}\right)}{b^2} - \frac{\sqrt[3]{c+dx} \cos\left(a+b\sqrt[3]{c+dx}\right)}{b} \right)}{b} \right)}{d}
 \end{array}$$

input `Int[Cos[a + b*(c + d*x)^(1/3)],x]`

output `(3*(((c + d*x)^(2/3)*Sin[a + b*(c + d*x)^(1/3)])/b - (2*(-(((c + d*x)^(1/3)*Cos[a + b*(c + d*x)^(1/3)])/b) + Sin[a + b*(c + d*x)^(1/3)]/b^2))/b))/d`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

rule 3777 `Int[((c_.) + (d_.)*(x_)^(m_.))*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(-`
`(c + d*x)^m)*(Cos[e + f*x]/f), x] + Simp[d*(m/f) Int[(c + d*x)^(m - 1)*C`
`os[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]`

rule 3843 `Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_)^(n_.))]*(b_.))^(p_.), x_S`
`ymbol] := Simp[1/(n*f) Subst[Int[x^(1/n - 1)*(a + b*Cos[c + d*x]]^p, x],`
`x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && Intege`
`rQ[1/n]`

Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.54

method	result
derivativedivides	$\frac{3a^2 \sin\left(a+b(dx+c)^{\frac{1}{3}}\right) - 6a\left(\cos\left(a+b(dx+c)^{\frac{1}{3}}\right) + \left(a+b(dx+c)^{\frac{1}{3}}\right) \sin\left(a+b(dx+c)^{\frac{1}{3}}\right)\right) + 3\left(a+b(dx+c)^{\frac{1}{3}}\right)^2 \sin\left(a+b(dx+c)^{\frac{1}{3}}\right)}{db^3}$
default	$\frac{3a^2 \sin\left(a+b(dx+c)^{\frac{1}{3}}\right) - 6a\left(\cos\left(a+b(dx+c)^{\frac{1}{3}}\right) + \left(a+b(dx+c)^{\frac{1}{3}}\right) \sin\left(a+b(dx+c)^{\frac{1}{3}}\right)\right) + 3\left(a+b(dx+c)^{\frac{1}{3}}\right)^2 \sin\left(a+b(dx+c)^{\frac{1}{3}}\right)}{db^3}$

input `int(cos(a+b*(d*x+c)^(1/3)),x,method=_RETURNVERBOSE)`

output `3/d/b^3*(a^2*sin(a+b*(d*x+c)^(1/3))-2*a*(cos(a+b*(d*x+c)^(1/3))+`
`(a+b*(d*x+c)^(1/3))*sin(a+b*(d*x+c)^(1/3)))+(a+b*(d*x+c)^(1/3))^2*sin(a+b*(d*x+c)^(1`
`/3))-2*sin(a+b*(d*x+c)^(1/3))+2*(a+b*(d*x+c)^(1/3))*cos(a+b*(d*x+c)^(1/3))`
`)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \frac{3 \left(2(dx + c)^{\frac{1}{3}} b \cos \left((dx + c)^{\frac{1}{3}} b + a \right) + \left((dx + c)^{\frac{2}{3}} b^2 - 2 \right) \sin \left((dx + c)^{\frac{1}{3}} b + a \right) \right)}{b^3 d}$$

input `integrate(cos(a+b*(d*x+c)^(1/3)),x, algorithm="fricas")`output `3*(2*(d*x + c)^(1/3)*b*cos((d*x + c)^(1/3)*b + a) + ((d*x + c)^(2/3)*b^2 - 2)*sin((d*x + c)^(1/3)*b + a))/(b^3*d)`**Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.11

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \begin{cases} x \cos(a) & \text{for } b = 0 \wedge (b = 0 \vee d = 0) \\ x \cos(a + b\sqrt[3]{c}) & \text{for } d = 0 \\ \frac{3(c+dx)^{\frac{2}{3}} \sin(a+b\sqrt[3]{c+dx})}{bd} + \frac{6\sqrt[3]{c+dx} \cos(a+b\sqrt[3]{c+dx})}{b^2 d} - \frac{6 \sin(a+b\sqrt[3]{c+dx})}{b^3 d} & \text{otherwise} \end{cases}$$

input `integrate(cos(a+b*(d*x+c)**(1/3)),x)`output `Piecewise((x*cos(a), Eq(b, 0) & (Eq(b, 0) | Eq(d, 0))), (x*cos(a + b*c**(1/3)), Eq(d, 0)), (3*(c + d*x)**(2/3)*sin(a + b*(c + d*x)**(1/3))/(b*d) + 6*(c + d*x)**(1/3)*cos(a + b*(c + d*x)**(1/3))/(b**2*d) - 6*sin(a + b*(c + d*x)**(1/3))/(b**3*d), True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.39

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \frac{3 \left(a^2 \sin \left((dx + c)^{\frac{1}{3}} b + a \right) - 2 \left((dx + c)^{\frac{1}{3}} b + a \right) \sin \left((dx + c)^{\frac{1}{3}} b + a \right) + \cos \left((dx + c)^{\frac{1}{3}} b + a \right) \right) a + 2}{b^3 d}$$

input `integrate(cos(a+b*(d*x+c)^(1/3)),x, algorithm="maxima")`output `3*(a^2*sin((d*x + c)^(1/3)*b + a) - 2*((d*x + c)^(1/3)*b + a)*sin((d*x + c)^(1/3)*b + a) + cos((d*x + c)^(1/3)*b + a))*a + 2*((d*x + c)^(1/3)*b + a)*cos((d*x + c)^(1/3)*b + a) + (((d*x + c)^(1/3)*b + a)^2 - 2)*sin((d*x + c)^(1/3)*b + a))/(b^3*d)`**Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.95

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \frac{3 \left(\frac{2(dx+c)^{\frac{1}{3}} \cos \left((dx+c)^{\frac{1}{3}} b + a \right)}{b} + \frac{\left((dx+c)^{\frac{1}{3}} b + a \right)^2 - 2 \left((dx+c)^{\frac{1}{3}} b + a \right) a + a^2 - 2}{b^2} \sin \left((dx+c)^{\frac{1}{3}} b + a \right) \right)}{bd}$$

input `integrate(cos(a+b*(d*x+c)^(1/3)),x, algorithm="giac")`output `3*(2*(d*x + c)^(1/3)*cos((d*x + c)^(1/3)*b + a)/b + (((d*x + c)^(1/3)*b + a)^2 - 2*((d*x + c)^(1/3)*b + a)*a + a^2 - 2)*sin((d*x + c)^(1/3)*b + a)/b^2)/(b*d)`

Mupad [B] (verification not implemented)

Time = 41.29 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.80

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \frac{6b \cos \left(a + b(c + dx)^{1/3} \right) (c + dx)^{1/3} - 6 \sin \left(a + b(c + dx)^{1/3} \right) + 3b^2 \sin \left(a + b(c + dx)^{1/3} \right) (c + dx)^{2/3}}{b^3 d}$$

input `int(cos(a + b*(c + d*x)^(1/3)),x)`output `(6*b*cos(a + b*(c + d*x)^(1/3))*(c + d*x)^(1/3) - 6*sin(a + b*(c + d*x)^(1/3)) + 3*b^2*sin(a + b*(c + d*x)^(1/3))*(c + d*x)^(2/3))/(b^3*d)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.80

$$\int \cos \left(a + b\sqrt[3]{c + dx} \right) dx$$

$$= \frac{6(dx + c)^{1/3} \cos \left((dx + c)^{1/3} b + a \right) b + 3(dx + c)^{2/3} \sin \left((dx + c)^{1/3} b + a \right) b^2 - 6 \sin \left((dx + c)^{1/3} b + a \right)}{b^3 d}$$

input `int(cos(a+b*(d*x+c)^(1/3)),x)`output `(3*(2*(c + d*x)**(1/3)*cos((c + d*x)**(1/3)*b + a)*b + (c + d*x)**(2/3)*sin((c + d*x)**(1/3)*b + a)*b**2 - 2*sin((c + d*x)**(1/3)*b + a))/(b**3*d)`

3.98
$$\int \frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x} dx$$

Optimal result	701
Mathematica [C] (verified)	702
Rubi [A] (verified)	703
Maple [C] (verified)	704
Fricas [C] (verification not implemented)	705
Sympy [F]	706
Maxima [F]	706
Giac [F]	707
Mupad [F(-1)]	707
Reduce [F]	707

Optimal result

Integrand size = 18, antiderivative size = 234

$$\int \frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x} dx = \cos\left(a+b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \cos\left(a+(-1)^{2/3}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left((-1)^{2/3}b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \cos\left(a-\sqrt[3]{-1}b\sqrt[3]{c}\right) \operatorname{CosIntegral}\left(\sqrt[3]{-1}b\sqrt[3]{c}+b\sqrt[3]{c+dx}\right) + \sin\left(a+b\sqrt[3]{c}\right) \operatorname{Si}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) + \sin\left(a+(-1)^{2/3}b\sqrt[3]{c}\right) \operatorname{Si}\left((-1)^{2/3}b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) - \sin\left(a-\sqrt[3]{-1}b\sqrt[3]{c}\right) \operatorname{Si}\left(\sqrt[3]{-1}b\sqrt[3]{c}+b\sqrt[3]{c+dx}\right)$$

output

```
cos(a+b*c^(1/3))*Ci(b*c^(1/3)-b*(d*x+c)^(1/3))+cos(a+(-1)^(2/3)*b*c^(1/3))
*Ci((-1)^(2/3)*b*c^(1/3)-b*(d*x+c)^(1/3))+cos(a-(-1)^(1/3)*b*c^(1/3))*Ci((-1)^(1/3)*b*c^(1/3)+b*(d*x+c)^(1/3))+sin(a+b*c^(1/3))*Si(b*c^(1/3)-b*(d*x+c)^(1/3))+sin(a+(-1)^(2/3)*b*c^(1/3))*Si((-1)^(2/3)*b*c^(1/3)-b*(d*x+c)^(1/3))-sin(a-(-1)^(1/3)*b*c^(1/3))*Si((-1)^(1/3)*b*c^(1/3)+b*(d*x+c)^(1/3))
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4 in optimal.

Time = 11.09 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.04

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \frac{1}{2} \left(\text{RootSum}\left[c - \#1^3 \&, \cos(a + b\#1) \text{CosIntegral}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) - i \text{CosIntegral}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) \sin(a + b\#1) - i \cos(a + b\#1) \text{Si}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) - \sin(a + b\#1) \text{Si}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) \& \right] + \text{RootSum}\left[c - \#1^3 \&, \cos(a + b\#1) \text{CosIntegral}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) + i \text{CosIntegral}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) \sin(a + b\#1) + i \cos(a + b\#1) \text{Si}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) - \sin(a + b\#1) \text{Si}\left(b\left(\sqrt[3]{c + dx} - \#1\right)\right) \& \right] \right)$$

input `Integrate[Cos[a + b*(c + d*x)^(1/3)]/x,x]`

output

```
(RootSum[c - #1^3 & , Cos[a + b*#1]*CosIntegral[b*((c + d*x)^(1/3) - #1]]
- I*CosIntegral[b*((c + d*x)^(1/3) - #1)]*Sin[a + b*#1] - I*Cos[a + b*#1]*
SinIntegral[b*((c + d*x)^(1/3) - #1)] - Sin[a + b*#1]*SinIntegral[b*((c +
d*x)^(1/3) - #1)] & ] + RootSum[c - #1^3 & , Cos[a + b*#1]*CosIntegral[b*(
(c + d*x)^(1/3) - #1]] + I*CosIntegral[b*((c + d*x)^(1/3) - #1)]*Sin[a + b
*#1] + I*Cos[a + b*#1]*SinIntegral[b*((c + d*x)^(1/3) - #1)] - Sin[a + b*#
1]*SinIntegral[b*((c + d*x)^(1/3) - #1)] & ])/2
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.12, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3913, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx$$

↓ 3913

$$\frac{3 \int \left(-\frac{d \cos\left(a + b\sqrt[3]{c + dx}\right)}{3\left(\sqrt[3]{c} - \sqrt[3]{c + dx}\right)} + \frac{d \cos\left(a + b\sqrt[3]{c + dx}\right)}{3\left(\sqrt[3]{-1}\sqrt[3]{c} + \sqrt[3]{c + dx}\right)} + \frac{\sqrt[3]{-1}d \cos\left(a + b\sqrt[3]{c + dx}\right)}{3\left(\sqrt[3]{c} + \sqrt[3]{-1}\sqrt[3]{c + dx}\right)} \right) d\sqrt[3]{c + dx}}{d}$$

↓ 2009

$$\frac{3\left(\frac{1}{3}d \cos\left(a + b\sqrt[3]{c}\right) \text{CosIntegral}\left(b\sqrt[3]{c} - b\sqrt[3]{c + dx}\right) + \frac{1}{3}d \cos\left(a + (-1)^{2/3}b\sqrt[3]{c}\right) \text{CosIntegral}\left((-1)^{2/3}b\sqrt[3]{c} - b\sqrt[3]{c}\right)\right)}{d}$$

input `Int[Cos[a + b*(c + d*x)^(1/3)]/x,x]`

output `(3*((d*cos[a + b*c^(1/3)]*CosIntegral[b*c^(1/3) - b*(c + d*x)^(1/3)])/3 + (d*cos[a + (-1)^(2/3)*b*c^(1/3)]*CosIntegral[(-1)^(2/3)*b*c^(1/3) - b*(c + d*x)^(1/3)])/3 + (d*cos[a - (-1)^(1/3)*b*c^(1/3)]*CosIntegral[(-1)^(1/3)*b*c^(1/3) + b*(c + d*x)^(1/3)])/3 + (d*sin[a + b*c^(1/3)]*SinIntegral[b*c^(1/3) - b*(c + d*x)^(1/3)])/3 + (d*sin[a + (-1)^(2/3)*b*c^(1/3)]*SinIntegral[(-1)^(2/3)*b*c^(1/3) - b*(c + d*x)^(1/3)])/3 - (d*sin[a - (-1)^(1/3)*b*c^(1/3)]*SinIntegral[(-1)^(1/3)*b*c^(1/3) + b*(c + d*x)^(1/3)]/3))/d`

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3913 Int[((a_.) + Cos[(c_.) + (d_.)*((e_.) + (f_.)*(x_))^(n_.)]*(b_.))^(p_.)*((g_.) + (h_.)*(x_))^(m_.), x_Symbol] := Simp[1/(n*f) Subst[Int[ExpandIntegrand[(a + b*Cos[c + d*x])^p, x^(1/n - 1)*(g - e*(h/f) + h*(x^(1/n)/f))^m, x], x, (e + f*x)^n], x] /; FreeQ[{a, b, c, d, e, f, g, h, m}, x] && IGtQ[p, 0] && IntegerQ[1/n]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.95 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{a^2 b^3 \left(\sum_{R1=RootOf(-b^3 c + Z^3 - 3a Z^2 + 3a^2 Z - a^3)} \frac{\text{Si}\left(-b(dx+c)^{\frac{1}{3}} + R1 - a\right) \sin(R1) + \text{Ci}\left(b(dx+c)^{\frac{1}{3}} - R1 + a\right)}{R1^2 - 2R1a + a^2}}{a^2 b^3 \left(\sum_{R1=RootOf(-b^3 c + Z^3 - 3a Z^2 + 3a^2 Z - a^3)} \frac{\text{Si}\left(-b(dx+c)^{\frac{1}{3}} + R1 - a\right) \sin(R1) + \text{Ci}\left(b(dx+c)^{\frac{1}{3}} - R1 + a\right)}{R1^2 - 2R1a + a^2}} \right)}$
default	$\frac{a^2 b^3 \left(\sum_{R1=RootOf(-b^3 c + Z^3 - 3a Z^2 + 3a^2 Z - a^3)} \frac{\text{Si}\left(-b(dx+c)^{\frac{1}{3}} + R1 - a\right) \sin(R1) + \text{Ci}\left(b(dx+c)^{\frac{1}{3}} - R1 + a\right)}{R1^2 - 2R1a + a^2}}{a^2 b^3 \left(\sum_{R1=RootOf(-b^3 c + Z^3 - 3a Z^2 + 3a^2 Z - a^3)} \frac{\text{Si}\left(-b(dx+c)^{\frac{1}{3}} + R1 - a\right) \sin(R1) + \text{Ci}\left(b(dx+c)^{\frac{1}{3}} - R1 + a\right)}{R1^2 - 2R1a + a^2}} \right)}$

```
input int(cos(a+b*(d*x+c)^(1/3))/x,x,method=_RETURNVERBOSE)
```

```
output 3/b^3*(1/3*a^2*b^3*sum(1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^(1/3)+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^(1/3)-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))-2/3*a*b^3*sum(_R1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^(1/3)+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^(1/3)-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+1/3*b^3*sum(_R1^2/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^(1/3)+_R1-a)*sin(_R1)+Ci(b*(d*x+c)^(1/3)-_R1+a)*cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3)))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.23

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \frac{1}{2} \operatorname{Ei}\left(i(dx + c)^{\frac{1}{3}}b + \frac{1}{2}(ib^3c)^{\frac{1}{3}}(-i\sqrt{3} - 1)\right) e^{\left(\frac{1}{2}(ib^3c)^{\frac{1}{3}}(i\sqrt{3}+1)+ia\right)} + \frac{1}{2} \operatorname{Ei}\left(-i(dx + c)^{\frac{1}{3}}b + \frac{1}{2}(-ib^3c)^{\frac{1}{3}}(-i\sqrt{3} - 1)\right) e^{\left(\frac{1}{2}(-ib^3c)^{\frac{1}{3}}(i\sqrt{3}+1)-ia\right)} + \frac{1}{2} \operatorname{Ei}\left(i(dx + c)^{\frac{1}{3}}b + \frac{1}{2}(ib^3c)^{\frac{1}{3}}(i\sqrt{3} - 1)\right) e^{\left(\frac{1}{2}(ib^3c)^{\frac{1}{3}}(-i\sqrt{3}+1)+ia\right)} + \frac{1}{2} \operatorname{Ei}\left(-i(dx + c)^{\frac{1}{3}}b + \frac{1}{2}(-ib^3c)^{\frac{1}{3}}(i\sqrt{3} - 1)\right) e^{\left(\frac{1}{2}(-ib^3c)^{\frac{1}{3}}(-i\sqrt{3}+1)-ia\right)} + \frac{1}{2} \operatorname{Ei}\left(i(dx + c)^{\frac{1}{3}}b + (ib^3c)^{\frac{1}{3}}\right) e^{\left(ia-(ib^3c)^{\frac{1}{3}}\right)} + \frac{1}{2} \operatorname{Ei}\left(-i(dx + c)^{\frac{1}{3}}b + (-ib^3c)^{\frac{1}{3}}\right) e^{\left(-ia-(-ib^3c)^{\frac{1}{3}}\right)}$$

input `integrate(cos(a+b*(d*x+c)^(1/3))/x,x, algorithm="fricas")`

output

```
1/2*Ei(I*(d*x + c)^(1/3)*b + 1/2*(I*b^3*c)^(1/3)*(-I*sqrt(3) - 1))*e^(1/2*
(I*b^3*c)^(1/3)*(I*sqrt(3) + 1) + I*a) + 1/2*Ei(-I*(d*x + c)^(1/3)*b + 1/2
*(-I*b^3*c)^(1/3)*(-I*sqrt(3) - 1))*e^(1/2*(-I*b^3*c)^(1/3)*(I*sqrt(3) + 1
) - I*a) + 1/2*Ei(I*(d*x + c)^(1/3)*b + 1/2*(I*b^3*c)^(1/3)*(I*sqrt(3) - 1
))*e^(1/2*(I*b^3*c)^(1/3)*(-I*sqrt(3) + 1) + I*a) + 1/2*Ei(-I*(d*x + c)^(1
/3)*b + 1/2*(-I*b^3*c)^(1/3)*(I*sqrt(3) - 1))*e^(1/2*(-I*b^3*c)^(1/3)*(-I*
sqrt(3) + 1) - I*a) + 1/2*Ei(I*(d*x + c)^(1/3)*b + (I*b^3*c)^(1/3))*e^(I*a
- (I*b^3*c)^(1/3)) + 1/2*Ei(-I*(d*x + c)^(1/3)*b + (-I*b^3*c)^(1/3))*e^(-
I*a - (-I*b^3*c)^(1/3))
```

Sympy [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx$$

input

```
integrate(cos(a+b*(d*x+c)**(1/3))/x,x)
```

output

```
Integral(cos(a + b*(c + d*x)**(1/3))/x, x)
```

Maxima [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \int \frac{\cos\left(\left(dx + c\right)^{\frac{1}{3}}b + a\right)}{x} dx$$

input

```
integrate(cos(a+b*(d*x+c)^(1/3))/x,x, algorithm="maxima")
```

output

```
integrate(cos((d*x + c)^(1/3)*b + a)/x, x)
```

Giac [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \int \frac{\cos\left(\left(dx + c\right)^{\frac{1}{3}}b + a\right)}{x} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/3))/x,x, algorithm="giac")`

output `integrate(cos((d*x + c)^(1/3)*b + a)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \int \frac{\cos\left(a + b(c + dx)^{1/3}\right)}{x} dx$$

input `int(cos(a + b*(c + d*x)^(1/3))/x,x)`

output `int(cos(a + b*(c + d*x)^(1/3))/x, x)`

Reduce [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x} dx = \int \frac{\cos\left(\left(dx + c\right)^{\frac{1}{3}}b + a\right)}{x} dx$$

input `int(cos(a+b*(d*x+c)^(1/3))/x,x)`

output `int(cos((c + d*x)**(1/3)*b + a)/x,x)`

3.99
$$\int \frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x^2} dx$$

Optimal result	708
Mathematica [C] (verified)	709
Rubi [A] (verified)	710
Maple [C] (verified)	712
Fricas [C] (verification not implemented)	713
Sympy [F]	713
Maxima [F]	714
Giac [F]	714
Mupad [F(-1)]	714
Reduce [F]	715

Optimal result

Integrand size = 18, antiderivative size = 332

$$\begin{aligned} & \int \frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x^2} dx \\ &= -\frac{\cos\left(a+b\sqrt[3]{c+dx}\right)}{x} - \frac{bd \operatorname{CosIntegral}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) \sin\left(a+b\sqrt[3]{c}\right)}{3c^{2/3}} \\ & \quad + \frac{\sqrt[3]{-1}bd \operatorname{CosIntegral}\left(\sqrt[3]{-1}b\sqrt[3]{c}+b\sqrt[3]{c+dx}\right) \sin\left(a-\sqrt[3]{-1}b\sqrt[3]{c}\right)}{3c^{2/3}} \\ & \quad - \frac{(-1)^{2/3}bd \operatorname{CosIntegral}\left((-1)^{2/3}b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right) \sin\left(a+(-1)^{2/3}b\sqrt[3]{c}\right)}{3c^{2/3}} \\ & \quad + \frac{bd \cos\left(a+b\sqrt[3]{c}\right) \operatorname{Si}\left(b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right)}{3c^{2/3}} \\ & \quad + \frac{(-1)^{2/3}bd \cos\left(a+(-1)^{2/3}b\sqrt[3]{c}\right) \operatorname{Si}\left((-1)^{2/3}b\sqrt[3]{c}-b\sqrt[3]{c+dx}\right)}{3c^{2/3}} \\ & \quad + \frac{\sqrt[3]{-1}bd \cos\left(a-\sqrt[3]{-1}b\sqrt[3]{c}\right) \operatorname{Si}\left(\sqrt[3]{-1}b\sqrt[3]{c}+b\sqrt[3]{c+dx}\right)}{3c^{2/3}} \end{aligned}$$

output

```

-cos(a+b*(d*x+c)^(1/3))/x-1/3*b*d*Ci(b*c^(1/3)-b*(d*x+c)^(1/3))*sin(a+b*c^(1/3))/c^(2/3)+1/3*(-1)^(1/3)*b*d*Ci((-1)^(1/3)*b*c^(1/3)+b*(d*x+c)^(1/3))*sin(a-(-1)^(1/3)*b*c^(1/3))/c^(2/3)-1/3*(-1)^(2/3)*b*d*Ci((-1)^(2/3)*b*c^(1/3)-b*(d*x+c)^(1/3))*sin(a+(-1)^(2/3)*b*c^(1/3))/c^(2/3)+1/3*b*d*cos(a+b*c^(1/3))*Si(b*c^(1/3)-b*(d*x+c)^(1/3))/c^(2/3)+1/3*(-1)^(2/3)*b*d*cos(a+(-1)^(2/3)*b*c^(1/3))*Si((-1)^(2/3)*b*c^(1/3)-b*(d*x+c)^(1/3))/c^(2/3)+1/3*(-1)^(1/3)*b*d*cos(a-(-1)^(1/3)*b*c^(1/3))*Si((-1)^(1/3)*b*c^(1/3)+b*(d*x+c)^(1/3))/c^(2/3)

```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4 in optimal.

Time = 0.82 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.42

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx$$

$$= -\frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x}$$

$$-\frac{1}{6}ibd\text{RootSum}\left[c - \#1^3 \&, \frac{e^{-ia - ib\#1} \text{ExpIntegralEi}\left(-ib\left(\sqrt[3]{c + dx} - \#1\right)\right)}{\#1^2} \&\right]$$

$$+\frac{1}{6}ibd\text{RootSum}\left[c - \#1^3 \&, \frac{e^{ia + ib\#1} \text{ExpIntegralEi}\left(ib\left(\sqrt[3]{c + dx} - \#1\right)\right)}{\#1^2} \&\right]$$

input

```
Integrate[Cos[a + b*(c + d*x)^(1/3)]/x^2,x]
```

output

```

-(Cos[a + b*(c + d*x)^(1/3)]/x) - (I/6)*b*d*RootSum[c - #1^3 & , (E^((-I)*a - I*b*#1)*ExpIntegralEi[(-I)*b*((c + d*x)^(1/3) - #1)]/#1^2 & ] + (I/6)*b*d*RootSum[c - #1^3 & , (E^(I*a + I*b*#1)*ExpIntegralEi[I*b*((c + d*x)^(1/3) - #1)]/#1^2 & ]

```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3913, 27, 3823, 3814, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + b\sqrt[3]{c + dx})}{x^2} dx$$

$$\downarrow \text{3913}$$

$$\frac{3 \int \frac{(c+dx)^{2/3} \cos(a+b\sqrt[3]{c+dx})}{x^2} d\sqrt[3]{c+dx}}{d}$$

$$\downarrow \text{27}$$

$$3d \int \frac{(c+dx)^{2/3} \cos(a+b\sqrt[3]{c+dx})}{d^2 x^2} d\sqrt[3]{c+dx}$$

$$\downarrow \text{3823}$$

$$3d \left(\frac{1}{3} b \int -\frac{\sin(a+b\sqrt[3]{c+dx})}{dx} d\sqrt[3]{c+dx} - \frac{\cos(a+b\sqrt[3]{c+dx})}{3dx} \right)$$

$$\downarrow \text{3814}$$

$$3d \left(\frac{1}{3} b \int \left(\frac{\sin(a+b\sqrt[3]{c+dx})}{3c^{2/3}(\sqrt[3]{c}-\sqrt[3]{c+dx})} + \frac{\sin(a+b\sqrt[3]{c+dx})}{3c^{2/3}(\sqrt[3]{c}+\sqrt[3]{-1}\sqrt[3]{c+dx})} + \frac{\sin(a+b\sqrt[3]{c+dx})}{3c^{2/3}(\sqrt[3]{c}-(-1)^{2/3}\sqrt[3]{c+dx})} \right) d\sqrt[3]{c+dx} \right)$$

$$\downarrow \text{2009}$$

$$3d \left(\frac{1}{3} b \left(-\frac{\sin(a+b\sqrt[3]{c}) \operatorname{CosIntegral}(b\sqrt[3]{c}-b\sqrt[3]{c+dx})}{3c^{2/3}} + \frac{\sqrt[3]{-1} \sin(a-\sqrt[3]{-1}b\sqrt[3]{c}) \operatorname{CosIntegral}(\sqrt[3]{-1}\sqrt[3]{cb}+\sqrt[3]{-1}\sqrt[3]{c+dx})}{3c^{2/3}} \right) \right)$$

input

```
Int[Cos[a + b*(c + d*x)^(1/3)]/x^2,x]
```

output

$$3*d*(-1/3*\text{Cos}[a + b*(c + d*x)^{(1/3)}]/(d*x) + (b*(-1/3*(\text{CosIntegral}[b*c^{(1/3)} - b*(c + d*x)^{(1/3)}]*\text{Sin}[a + b*c^{(1/3)}])/c^{(2/3)} + ((-1)^{(1/3)}*\text{CosIntegral}[(-1)^{(1/3)}*b*c^{(1/3)} + b*(c + d*x)^{(1/3)}]*\text{Sin}[a - (-1)^{(1/3)}*b*c^{(1/3)}])/((3*c^{(2/3)}) - ((-1)^{(2/3)}*\text{CosIntegral}[(-1)^{(2/3)}*b*c^{(1/3)} - b*(c + d*x)^{(1/3)}]*\text{Sin}[a + (-1)^{(2/3)}*b*c^{(1/3)}])/((3*c^{(2/3)}) + (\text{Cos}[a + b*c^{(1/3)}]*\text{SinIntegral}[b*c^{(1/3)} - b*(c + d*x)^{(1/3)}])/((3*c^{(2/3)}) + ((-1)^{(2/3)}*\text{Cos}[a + (-1)^{(2/3)}*b*c^{(1/3)}]*\text{SinIntegral}[(-1)^{(2/3)}*b*c^{(1/3)} - b*(c + d*x)^{(1/3)}])/((3*c^{(2/3)}) + ((-1)^{(1/3)}*\text{Cos}[a - (-1)^{(1/3)}*b*c^{(1/3)}]*\text{SinIntegral}[(-1)^{(1/3)}*b*c^{(1/3)} + b*(c + d*x)^{(1/3)}])/((3*c^{(2/3)}))))/3)$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 3814

$$\text{Int}[(a_*) + (b_*)(x_)^{(n_*)})^{(p_*)} \text{Sin}[(c_*) + (d_*)(x_)], x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[\text{Sin}[c + d*x], (a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{ILtQ}[p, 0] \&\& \text{IGtQ}[n, 0] \&\& (\text{EqQ}[n, 2] \parallel \text{EqQ}[p, -1])$$

rule 3823

$$\text{Int}[\text{Cos}[(c_*) + (d_*)(x_)] * ((e_*)(x_))^{(m_*)} * ((a_*) + (b_*)(x_))^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[e^m * (a + b*x^n)^{(p+1)} * (\text{Cos}[c + d*x] / (b*n*(p+1))), x] + \text{Simp}[d * (e^m / (b*n*(p+1))) \text{Int}[(a + b*x^n)^{(p+1)} * \text{Sin}[c + d*x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \&\& \text{ILtQ}[p, -1] \&\& \text{EqQ}[m, n-1] \&\& (\text{IntegerQ}[n] \parallel \text{GtQ}[e, 0])$$

rule 3913

$$\text{Int}[(a_*) + \text{Cos}[(c_*) + (d_*)(e_*) + (f_*)(x_))^{(n_*)}] * (b_*)^{(p_*)} * ((g_*) + (h_*)(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Simp}[1/(n*f) \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(a + b*\text{Cos}[c + d*x])^p, x^{(1/n-1)} * (g - e*(h/f) + h*(x^{(1/n)}/f))^m, x], x, (e + f*x)^n], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, h, m\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{IntegerQ}[1/n]$$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.14 (sec) , antiderivative size = 931, normalized size of antiderivative = 2.80

method	result	size
derivativedivides	Expression too large to display	931
default	Expression too large to display	931

input `int(cos(a+b*(d*x+c)^(1/3))/x^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & 3*d/b^3*(b^6*a^2*(\cos(a+b*(d*x+c)^{1/3})*(1/3/c/b^3*(a+b*(d*x+c)^{1/3})-1/ \\
 & 3*a/b^3/c)/(b^3*c+a^3-3*a^2*(a+b*(d*x+c)^{1/3}))+3*a*(a+b*(d*x+c)^{1/3})^2- \\
 & (a+b*(d*x+c)^{1/3})^3)-2/9/c/b^3*\sum(1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^{1/3} \\
 & +_R1-a)*\sin(_R1)+Ci(b*(d*x+c)^{1/3}-_R1+a)*\cos(_R1)),_R1=RootOf(-b^3*c \\
 & c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+1/9/c/b^3*\sum(1/(-_RR1+a)*(-Si(-b*(d*x+c)^{1/3} \\
 & +_RR1-a)*\cos(_RR1)+Ci(b*(d*x+c)^{1/3}-_RR1+a)*\sin(_RR1)),_RR1=RootOf(- \\
 & b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2-a^3))+\cos(a+b*(d*x+c)^{1/3})*(-2/3*a*b^3/c*(\\
 & a+b*(d*x+c)^{1/3})^2+2/3*a^2*b^3/c*(a+b*(d*x+c)^{1/3}))/ (b^3*c+a^3-3*a^2*(\\
 & a+b*(d*x+c)^{1/3}))+3*a*(a+b*(d*x+c)^{1/3})^2-(a+b*(d*x+c)^{1/3})^3)+2/9*a* \\
 & b^3/c*\sum((_R1+a)/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^{1/3}+_R1-a)*\sin(_R1) \\
 & +Ci(b*(d*x+c)^{1/3}-_R1+a)*\cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z* \\
 & a^2-a^3))-2/9*a*b^3/c*\sum(_RR1/(-_RR1+a)*(-Si(-b*(d*x+c)^{1/3}+_RR1-a)*\cos \\
 & (_RR1)+Ci(b*(d*x+c)^{1/3}-_RR1+a)*\sin(_RR1)),_RR1=RootOf(-b^3*c+_Z^3-3*_Z^ \\
 & 2*a+3*_Z*a^2-a^3))+\cos(a+b*(d*x+c)^{1/3})*(2/3*a*b^3/c*(a+b*(d*x+c)^{1/3}) \\
 & ^2-a^2*b^3/c*(a+b*(d*x+c)^{1/3}))+1/3*b^3*(b^3*c+a^3)/c)/(b^3*c+a^3-3*a^2*(\\
 & a+b*(d*x+c)^{1/3}))+3*a*(a+b*(d*x+c)^{1/3})^2-(a+b*(d*x+c)^{1/3})^3)-2/9*a* \\
 & b^3/c*\sum(_R1/(_R1^2-2*_R1*a+a^2)*(Si(-b*(d*x+c)^{1/3}+_R1-a)*\sin(_R1)+Ci(\\
 & b*(d*x+c)^{1/3}-_R1+a)*\cos(_R1)),_R1=RootOf(-b^3*c+_Z^3-3*_Z^2*a+3*_Z*a^2- \\
 & a^3))-1/9*b^3/c*\sum((b^3*c+2*_RR1^2*a-3*_RR1*a^2+a^3)/(_RR1^2-2*_RR1*a+a^2) \\
 &)*(-Si(-b*(d*x+c)^{1/3}+_RR1-a)*\cos(_RR1)+Ci(b*(d*x+c)^{1/3}-_RR1+a)*si...
 \end{aligned}$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.22

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx =$$

$$\frac{2(i b^3 c)^{\frac{1}{3}} dx \operatorname{Ei}\left(i(dx + c)^{\frac{1}{3}} b + (i b^3 c)^{\frac{1}{3}}\right) e^{\left(i a - (i b^3 c)^{\frac{1}{3}}\right)} + 2(-i b^3 c)^{\frac{1}{3}} dx \operatorname{Ei}\left(-i(dx + c)^{\frac{1}{3}} b + (-i b^3 c)^{\frac{1}{3}}\right) e^{\left(-i a + (i b^3 c)^{\frac{1}{3}}\right)}}{x^2}$$

input `integrate(cos(a+b*(d*x+c)^(1/3))/x^2,x, algorithm="fricas")`

output

```
-1/12*(2*(I*b^3*c)^(1/3)*d*x*Ei(I*(d*x + c)^(1/3)*b + (I*b^3*c)^(1/3))*e^(
I*a - (I*b^3*c)^(1/3)) + 2*(-I*b^3*c)^(1/3)*d*x*Ei(-I*(d*x + c)^(1/3)*b +
(-I*b^3*c)^(1/3))*e^(-I*a - (-I*b^3*c)^(1/3)) - (I*b^3*c)^(1/3)*(I*sqrt(3)
*d*x + d*x)*Ei(I*(d*x + c)^(1/3)*b + 1/2*(I*b^3*c)^(1/3)*(-I*sqrt(3) - 1))
*e^(1/2*(I*b^3*c)^(1/3)*(I*sqrt(3) + 1) + I*a) - (-I*b^3*c)^(1/3)*(I*sqrt(3)
*d*x + d*x)*Ei(-I*(d*x + c)^(1/3)*b + 1/2*(-I*b^3*c)^(1/3)*(-I*sqrt(3) -
1))*e^(1/2*(-I*b^3*c)^(1/3)*(I*sqrt(3) + 1) - I*a) - (I*b^3*c)^(1/3)*(-I*
sqrt(3)*d*x + d*x)*Ei(I*(d*x + c)^(1/3)*b + 1/2*(I*b^3*c)^(1/3)*(I*sqrt(3)
- 1))*e^(1/2*(I*b^3*c)^(1/3)*(-I*sqrt(3) + 1) + I*a) - (-I*b^3*c)^(1/3)*(-
I*sqrt(3)*d*x + d*x)*Ei(-I*(d*x + c)^(1/3)*b + 1/2*(-I*b^3*c)^(1/3)*(I*sq
rt(3) - 1))*e^(1/2*(-I*b^3*c)^(1/3)*(-I*sqrt(3) + 1) - I*a) + 12*c*cos((d*
x + c)^(1/3)*b + a)/(c*x)
```

Sympy [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx = \int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx$$

input `integrate(cos(a+b*(d*x+c)**(1/3))/x**2,x)`

output

```
Integral(cos(a + b*(c + d*x)**(1/3))/x**2, x)
```

Maxima [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx = \int \frac{\cos\left(\left(dx + c\right)^{\frac{1}{3}}b + a\right)}{x^2} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/3))/x^2,x, algorithm="maxima")`

output `integrate(cos((d*x + c)^(1/3)*b + a)/x^2, x)`

Giac [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx = \int \frac{\cos\left(\left(dx + c\right)^{\frac{1}{3}}b + a\right)}{x^2} dx$$

input `integrate(cos(a+b*(d*x+c)^(1/3))/x^2,x, algorithm="giac")`

output `integrate(cos((d*x + c)^(1/3)*b + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx = \int \frac{\cos\left(a + b(c + dx)^{1/3}\right)}{x^2} dx$$

input `int(cos(a + b*(c + d*x)^(1/3))/x^2,x)`

output `int(cos(a + b*(c + d*x)^(1/3))/x^2, x)`

Reduce [F]

$$\int \frac{\cos\left(a + b\sqrt[3]{c + dx}\right)}{x^2} dx = \int \frac{\cos\left((dx + c)^{\frac{1}{3}} b + a\right)}{x^2} dx$$

input `int(cos(a+b*(d*x+c)^(1/3))/x^2,x)`

output `int(cos((c + d*x)**(1/3)*b + a)/x**2,x)`

CHAPTER 4

APPENDIX

4.1	Listing of Grading functions	716
4.2	Links to plain text integration problems used in this report for each CAS .	734

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
    If [Head [expn] === RootSum,
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
    If [Head [expn] === Integrate || Head [expn] === Int,
    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
    9]]]]]]]]]]]

```

```

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#                    if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#                    see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```



```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file