

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.2-Cosine/203-4.2.13

Nasser M. Abbasi

May 18, 2024 Compiled on May 18, 2024 at 4:14am

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22
2	detailed summary tables of results	23
2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	37
3	Listing of integrals	39
3.1	$\int x^2 \cos(a + bx + cx^2) dx$	41
3.2	$\int x \cos(a + bx + cx^2) dx$	50
3.3	$\int \cos(a + bx + cx^2) dx$	57
3.4	$\int \frac{\cos(a+bx+cx^2)}{x} dx$	63
3.5	$\int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx$	68
3.6	$\int x^2 \cos(a + bx - cx^2) dx$	73

3.7	$\int x \cos(a + bx - cx^2) dx$	82
3.8	$\int \cos(a + bx - cx^2) dx$	89
3.9	$\int \frac{\cos(a+bx-cx^2)}{x} dx$	95
3.10	$\int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx$	100
3.11	$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx$	105
3.12	$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx$	112
3.13	$\int \cos\left(\frac{1}{4} + x + x^2\right) dx$	118
3.14	$\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x} dx$	123
3.15	$\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$	128
3.16	$\int x^2 \cos^2(a + bx + cx^2) dx$	134
3.17	$\int x \cos^2(a + bx + cx^2) dx$	141
3.18	$\int \cos^2(a + bx + cx^2) dx$	147
3.19	$\int \frac{\cos^2(a+bx+cx^2)}{x} dx$	153
3.20	$\int x^2 \cos^2(a + bx - cx^2) dx$	158
3.21	$\int x \cos^2(a + bx - cx^2) dx$	165
3.22	$\int \cos^2(a + bx - cx^2) dx$	171
3.23	$\int \frac{\cos^2(a+bx-cx^2)}{x} dx$	177
3.24	$\int x^2 \cos^2\left(\frac{1}{4} + x + x^2\right) dx$	182
3.25	$\int x \cos^2\left(\frac{1}{4} + x + x^2\right) dx$	188
3.26	$\int \cos^2\left(\frac{1}{4} + x + x^2\right) dx$	194
3.27	$\int \frac{\cos^2\left(\frac{1}{4}+x+x^2\right)}{x} dx$	199
3.28	$\int \frac{\cos^2\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$	204
3.29	$\int (d + ex)^2 \cos(a + bx + cx^2) dx$	209
3.30	$\int (d + ex) \cos(a + bx + cx^2) dx$	218
3.31	$\int \frac{\cos(a+bx+cx^2)}{d+ex} dx$	225
3.32	$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx$	230
3.33	$\int (d + ex) \cos^2(a + bx + cx^2) dx$	238
3.34	$\int \frac{\cos^2(a+bx+cx^2)}{d+ex} dx$	244
4	Appendix	249
4.1	Listing of Grading functions	249
4.2	Links to plain text integration problems used in this report for each CAS267	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [34]. This is test number [203].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (34)	0.00 (0)
Mathematica	100.00 (34)	0.00 (0)
Fricas	100.00 (34)	0.00 (0)
Maple	94.12 (32)	5.88 (2)
Giac	94.12 (32)	5.88 (2)
Maxima	94.12 (32)	5.88 (2)
Sympy	52.94 (18)	47.06 (16)
Mupad	44.12 (15)	55.88 (19)
Reduce	29.41 (10)	70.59 (24)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	70.588	0.000	0.000	29.412
Mathematica	70.588	0.000	0.000	29.412
Fricas	70.588	0.000	0.000	29.412
Maple	64.706	0.000	0.000	35.294
Sympy	17.647	5.882	0.000	76.471
Giac	0.000	0.000	64.706	35.294
Mupad	0.000	14.706	0.000	85.294
Maxima	0.000	0.000	64.706	35.294
Reduce	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	2	100.00	0.00	0.00
Giac	2	100.00	0.00	0.00
Maxima	2	100.00	0.00	0.00
Sympy	16	100.00	0.00	0.00
Mupad	19	0.00	100.00	0.00
Reduce	24	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.08
Reduce	0.19
Rubi	0.31
Maxima	0.35
Giac	0.36
Sympy	0.69
Maple	1.15
Mathematica	2.22
Mupad	36.09

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	18.90	1.18	18.50	1.11
Mupad	32.87	1.01	19.00	1.01
Sympy	46.06	1.18	18.00	0.91
Mathematica	83.97	0.95	86.50	0.96
Fricas	89.09	0.95	94.50	0.99
Maple	95.19	0.90	68.00	0.85
Rubi	101.94	1.00	98.50	1.00
Giac	111.88	1.15	98.50	1.11
Maxima	509.28	3.35	125.00	2.46

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{4, 9, 14, 15, 19, 23, 27, 28, 31, 34}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A `time limit` of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError.`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	37

2.1 List of integrals sorted by grade for each CAS

Rubi	24
Mma	24
Maple	25
Fricas	25
Maxima	25
Giac	26
Mupad	26
Sympy	26
Reduce	27

Rubi

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }
B grade { }
C grade { }
F normal fail { }
F(-1) timeout fail { }
F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }
B grade { }
C grade { }
F normal fail { }
F(-1) timeout fail { }
F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

B grade { }

C grade { }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { }

B grade { }

C grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac**A grade** { }**B grade** { }**C grade** { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }**F normal fail** { 5, 10 }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Mupad****A grade** { }**B grade** { 3, 8, 11, 12, 13 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { 1, 2, 5, 6, 7, 10, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }**F(-2) exception fail** { }**Sympy****A grade** { 3, 8, 13, 18, 22, 26 }**B grade** { 12, 25 }**C grade** { }**F normal fail** { 1, 2, 5, 6, 7, 10, 11, 16, 17, 20, 21, 24, 29, 30, 32, 33 }**F(-1) timeout fail** { }**F(-2) exception fail** { }

Reduce

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32,
33 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	249	257	160	204	1570	173	0	225	17	0
N.S.	1	1.03	0.64	0.82	6.31	0.69	0.00	0.90	0.07	0.00
time (sec)	N/A	0.751	0.793	2.021	0.761	0.086	0.000	0.328	0.167	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	125	112	99	580	119	0	181	33	0
N.S.	1	1.02	0.91	0.80	4.72	0.97	0.00	1.47	0.27	0.00
time (sec)	N/A	0.332	0.335	1.137	0.420	0.083	0.000	0.333	0.187	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	85	81	112	103	88	135	13	99
N.S.	1	1.00	0.87	0.83	1.14	1.05	0.90	1.38	0.13	1.01
time (sec)	N/A	0.270	0.193	0.677	0.044	0.082	0.197	0.343	0.166	41.469

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.167	2.496	0.216	0.207	0.070	0.355	0.342	0.175	41.381

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	111	111	110	0	0	115	0	0	52	0
N.S.	1	1.00	0.99	0.00	0.00	1.04	0.00	0.00	0.47	0.00
time (sec)	N/A	0.304	2.962	0.000	0.000	0.090	0.000	0.000	0.163	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	251	261	164	224	1570	178	0	227	18	0
N.S.	1	1.04	0.65	0.89	6.25	0.71	0.00	0.90	0.07	0.00
time (sec)	N/A	0.723	0.756	1.731	0.906	0.082	0.000	0.366	0.180	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	127	116	110	580	125	0	183	36	0
N.S.	1	1.02	0.94	0.89	4.68	1.01	0.00	1.48	0.29	0.00
time (sec)	N/A	0.339	0.377	1.155	0.393	0.079	0.000	0.349	0.190	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	88	88	112	106	94	137	14	105
N.S.	1	1.00	0.89	0.89	1.13	1.07	0.95	1.38	0.14	1.06
time (sec)	N/A	0.277	0.192	0.698	0.044	0.077	0.208	0.339	0.194	0.051

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	20	20	14	18	18	18
N.S.	1	1.00	1.12	1.00	1.25	1.25	0.88	1.12	1.12	1.12
time (sec)	N/A	0.169	2.064	0.217	0.159	0.069	0.390	0.374	0.249	41.214

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	111	111	114	0	0	123	0	0	54	0
N.S.	1	1.00	1.03	0.00	0.00	1.11	0.00	0.00	0.49	0.00
time (sec)	N/A	0.303	4.281	0.000	0.000	0.079	0.000	0.000	0.257	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	87	67	59	160	59	0	75	31	64
N.S.	1	1.06	0.82	0.72	1.95	0.72	0.00	0.91	0.38	0.78
time (sec)	N/A	0.367	0.159	1.640	0.223	0.075	0.000	0.345	0.264	41.310

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	39	30	126	32	155	65	21	32
N.S.	1	1.00	0.95	0.73	3.07	0.78	3.78	1.59	0.51	0.78
time (sec)	N/A	0.220	0.056	1.342	0.183	0.073	0.570	0.328	0.173	0.259

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	20	70	22	29	39	9	19
N.S.	1	1.00	1.00	0.83	2.92	0.92	1.21	1.62	0.38	0.79
time (sec)	N/A	0.181	0.031	0.834	0.134	0.074	0.190	0.300	0.197	41.348

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	13	13	12	13	13	13
N.S.	1	1.00	1.15	0.85	1.00	1.00	0.92	1.00	1.00	1.00
time (sec)	N/A	0.164	13.433	0.185	0.149	0.066	0.316	0.320	0.182	41.306

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	13	13	14	13	28	13
N.S.	1	1.00	1.15	0.85	1.00	1.00	1.08	1.00	2.15	1.00
time (sec)	N/A	0.297	11.246	0.171	0.196	0.080	0.296	0.357	0.192	41.396

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	248	248	170	191	1617	178	0	212	318	0
N.S.	1	1.00	0.69	0.77	6.52	0.72	0.00	0.85	1.28	0.00
time (sec)	N/A	0.452	0.743	1.706	0.800	0.084	0.000	0.333	0.198	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	116	95	610	124	0	170	17	0
N.S.	1	1.00	0.92	0.75	4.84	0.98	0.00	1.35	0.13	0.00
time (sec)	N/A	0.267	0.284	1.515	0.419	0.091	0.000	0.385	0.210	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	97	72	124	93	83	122	15	0
N.S.	1	1.00	0.97	0.72	1.24	0.93	0.83	1.22	0.15	0.00
time (sec)	N/A	0.236	0.117	1.419	0.124	0.083	0.436	0.333	0.179	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	19	17	28	19	15	19	19	19
N.S.	1	1.00	1.12	1.00	1.65	1.12	0.88	1.12	1.12	1.12
time (sec)	N/A	0.203	2.625	0.405	0.242	0.084	0.785	0.387	0.180	40.576

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	248	248	175	199	1617	187	0	214	324	0
N.S.	1	1.00	0.71	0.80	6.52	0.75	0.00	0.86	1.31	0.00
time (sec)	N/A	0.435	0.655	1.694	0.826	0.094	0.000	0.377	0.191	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	122	99	611	133	0	172	18	0
N.S.	1	1.00	0.97	0.79	4.85	1.06	0.00	1.37	0.14	0.00
time (sec)	N/A	0.284	0.284	1.556	0.421	0.082	0.000	0.345	0.170	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	100	76	124	96	88	124	16	0
N.S.	1	1.00	1.00	0.76	1.24	0.96	0.88	1.24	0.16	0.00
time (sec)	N/A	0.246	0.121	1.355	0.121	0.079	0.418	0.358	0.178	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	28	22	15	20	20	20
N.S.	1	1.00	1.11	1.00	1.56	1.22	0.83	1.11	1.11	1.11
time (sec)	N/A	0.202	3.347	0.397	0.192	0.079	0.779	0.374	0.178	41.389

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	77	64	171	57	0	64	232	0
N.S.	1	1.00	0.91	0.75	2.01	0.67	0.00	0.75	2.73	0.00
time (sec)	N/A	0.246	0.169	2.157	0.220	0.077	0.000	0.354	0.186	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	B	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	41	35	136	37	121	54	13	0
N.S.	1	1.00	0.89	0.76	2.96	0.80	2.63	1.17	0.28	0.00
time (sec)	N/A	0.203	0.070	1.799	0.229	0.078	0.779	0.368	0.169	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	26	20	34	19	22	26	11	0
N.S.	1	1.00	0.96	0.74	1.26	0.70	0.81	0.96	0.41	0.00
time (sec)	N/A	0.185	0.018	1.660	0.112	0.073	0.342	0.386	0.171	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	13	24	15	14	15	15	15
N.S.	1	1.00	1.13	0.87	1.60	1.00	0.93	1.00	1.00	1.00
time (sec)	N/A	0.197	7.554	0.372	0.220	0.074	0.608	0.365	0.191	42.253

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	13	26	15	15	15	15	15
N.S.	1	1.00	1.13	0.87	1.73	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.232	8.606	0.375	0.151	0.078	0.511	0.391	0.177	42.600

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	285	282	187	396	2271	230	0	291	133	0
N.S.	1	0.99	0.66	1.39	7.97	0.81	0.00	1.02	0.47	0.00
time (sec)	N/A	0.791	1.381	1.977	1.059	0.110	0.000	0.377	0.175	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	134	129	182	695	136	0	201	82	0
N.S.	1	0.96	0.92	1.30	4.96	0.97	0.00	1.44	0.59	0.00
time (sec)	N/A	0.360	0.733	1.237	0.415	0.159	0.000	0.375	0.186	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	19	21	21	17	21	21	21
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.89	1.11	1.11	1.11
time (sec)	N/A	0.174	3.701	0.460	0.196	0.070	0.331	0.369	0.171	42.521

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	291	291	215	378	2360	257	0	294	532	0
N.S.	1	1.00	0.74	1.30	8.11	0.88	0.00	1.01	1.83	0.00
time (sec)	N/A	0.559	1.113	2.251	1.059	0.090	0.000	0.400	0.202	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	C	A	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	150	150	139	170	737	149	0	195	37	0
N.S.	1	1.00	0.93	1.13	4.91	0.99	0.00	1.30	0.25	0.00
time (sec)	N/A	0.296	0.549	1.805	0.435	0.085	0.000	0.383	0.181	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	21	120	23	19	23	23	23
N.S.	1	1.00	1.10	1.00	5.71	1.10	0.90	1.10	1.10	1.10
time (sec)	N/A	0.225	3.920	0.575	0.239	0.078	4.849	0.383	0.193	42.268

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [6] had the largest ratio of [.6250000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	8	1.03	15	0.533
2	A	4	4	1.02	13	0.308
3	A	3	3	1.00	11	0.273
4	N/A	1	0	1.00	15	0.000
5	A	1	1	1.00	32	0.031
6	A	10	10	1.04	16	0.625
7	A	5	5	1.02	14	0.357
8	A	4	4	1.00	12	0.333
9	N/A	1	0	1.00	16	0.000
10	A	1	1	1.00	34	0.029
11	A	6	6	1.06	13	0.462
12	A	3	3	1.00	11	0.273
13	A	2	2	1.00	9	0.222
14	N/A	1	0	1.00	13	0.000
15	N/A	4	0	1.00	13	0.000
16	A	2	2	1.00	17	0.118
17	A	2	2	1.00	15	0.133
18	A	2	2	1.00	13	0.154
19	N/A	2	0	1.00	17	0.000
20	A	2	2	1.00	18	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
21	A	2	2	1.00	16	0.125
22	A	2	2	1.00	14	0.143
23	N/A	2	0	1.00	18	0.000
24	A	2	2	1.00	15	0.133
25	A	2	2	1.00	13	0.154
26	A	2	2	1.00	11	0.182
27	N/A	2	0	1.00	15	0.000
28	N/A	2	0	1.00	15	0.000
29	A	8	8	0.99	19	0.421
30	A	4	4	0.96	17	0.235
31	N/A	1	0	1.00	19	0.000
32	A	2	2	1.00	21	0.095
33	A	2	2	1.00	19	0.105
34	N/A	2	0	1.00	21	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^2 \cos(a + bx + cx^2) dx$	41
3.2	$\int x \cos(a + bx + cx^2) dx$	50
3.3	$\int \cos(a + bx + cx^2) dx$	57
3.4	$\int \frac{\cos(a+bx+cx^2)}{x} dx$	63
3.5	$\int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b\sin(a+bx+cx^2)}{x} \right) dx$	68
3.6	$\int x^2 \cos(a + bx - cx^2) dx$	73
3.7	$\int x \cos(a + bx - cx^2) dx$	82
3.8	$\int \cos(a + bx - cx^2) dx$	89
3.9	$\int \frac{\cos(a+bx-cx^2)}{x} dx$	95
3.10	$\int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b\sin(a+bx-cx^2)}{x} \right) dx$	100
3.11	$\int x^2 \cos(\frac{1}{4} + x + x^2) dx$	105
3.12	$\int x \cos(\frac{1}{4} + x + x^2) dx$	112
3.13	$\int \cos(\frac{1}{4} + x + x^2) dx$	118
3.14	$\int \frac{\cos(\frac{1}{4}+x+x^2)}{x} dx$	123
3.15	$\int \frac{\cos(\frac{1}{4}+x+x^2)}{x^2} dx$	128
3.16	$\int x^2 \cos^2(a + bx + cx^2) dx$	134
3.17	$\int x \cos^2(a + bx + cx^2) dx$	141
3.18	$\int \cos^2(a + bx + cx^2) dx$	147
3.19	$\int \frac{\cos^2(a+bx+cx^2)}{x} dx$	153
3.20	$\int x^2 \cos^2(a + bx - cx^2) dx$	158
3.21	$\int x \cos^2(a + bx - cx^2) dx$	165
3.22	$\int \cos^2(a + bx - cx^2) dx$	171
3.23	$\int \frac{\cos^2(a+bx-cx^2)}{x} dx$	177
3.24	$\int x^2 \cos^2(\frac{1}{4} + x + x^2) dx$	182
3.25	$\int x \cos^2(\frac{1}{4} + x + x^2) dx$	188

3.26	$\int \cos^2\left(\frac{1}{4} + x + x^2\right) dx$	194
3.27	$\int \frac{\cos^2(\frac{1}{4}+x+x^2)}{x} dx$	199
3.28	$\int \frac{\cos^2(\frac{1}{4}+x+x^2)}{x^2} dx$	204
3.29	$\int (d+ex)^2 \cos(a+bx+cx^2) dx$	209
3.30	$\int (d+ex) \cos(a+bx+cx^2) dx$	218
3.31	$\int \frac{\cos(a+bx+cx^2)}{d+ex} dx$	225
3.32	$\int (d+ex)^2 \cos^2(a+bx+cx^2) dx$	230
3.33	$\int (d+ex) \cos^2(a+bx+cx^2) dx$	238
3.34	$\int \frac{\cos^2(a+bx+cx^2)}{d+ex} dx$	244

3.1 $\int x^2 \cos(a + bx + cx^2) dx$

Optimal result	41
Mathematica [A] (verified)	42
Rubi [A] (verified)	42
Maple [A] (verified)	45
Fricas [A] (verification not implemented)	46
Sympy [F]	46
Maxima [C] (verification not implemented)	47
Giac [C] (verification not implemented)	48
Mupad [F(-1)]	48
Reduce [F]	49

Optimal result

Integrand size = 15, antiderivative size = 249

$$\begin{aligned} \int x^2 \cos(a + bx + cx^2) dx &= \frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ &\quad - \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\ &\quad - \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\ &\quad - \frac{b \sin(a + bx + cx^2)}{4c^2} + \frac{x \sin(a + bx + cx^2)}{2c} \end{aligned}$$

output

```
1/8*b^2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(5/2)-1/4*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)-1/4*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)-1/8*b^2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(5/2)-1/4*b*sin(c*x^2+b*x+a)/c^2+1/2*x*sin(c*x^2+b*x+a)/c
```

Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.64

$$\int x^2 \cos(a + bx + cx^2) dx \\ = \frac{-\sqrt{2\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a - \frac{b^2}{4c}\right) + b^2 \sin\left(a - \frac{b^2}{4c}\right)\right) + \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(b^2 \cos\left(a - \frac{b^2}{4c}\right) - 2c \sin\left(a - \frac{b^2}{4c}\right)\right)}{8c^{5/2}}$$

input `Integrate[x^2*Cos[a + b*x + c*x^2], x]`

output
$$\frac{(-\operatorname{FresnelS}\left[\frac{(b+2c x)/(2 \pi)}{\sqrt{c}}\right] \left(2 c \cos \left(a-\frac{b^2}{4 c}\right)+b^2 \sin \left(a-\frac{b^2}{4 c}\right)\right)+\operatorname{FresnelC}\left[\frac{(b+2c x)/(2 \pi)}{\sqrt{c}}\right] \left(b^2 \cos \left(a-\frac{b^2}{4 c}\right)-2 c \sin \left(a-\frac{b^2}{4 c}\right)\right)}{8 c^{5/2}}$$

Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3945, 3928, 3832, 3833, 3943, 3929, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos(a + bx + cx^2) dx \\ \downarrow 3945 \\ -\frac{\int \sin(cx^2 + bx + a) dx}{2c} - \frac{b \int x \cos(cx^2 + bx + a) dx}{2c} + \frac{x \sin(a + bx + cx^2)}{2c} \\ \downarrow 3928 \\ -\frac{\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx + \cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} - \\ \frac{b \int x \cos(cx^2 + bx + a) dx}{2c} + \frac{x \sin(a + bx + cx^2)}{2c}$$

$$\begin{aligned}
& \downarrow \text{3832} \\
& - \frac{\sin(a - \frac{b^2}{4c}) \int \cos(\frac{(b+2cx)^2}{4c}) dx + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \\
& \quad \frac{b \int x \cos(cx^2 + bx + a) dx}{2c} + \frac{x \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3833} \\
& - \frac{b \int x \cos(cx^2 + bx + a) dx}{2c} - \frac{\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} + \\
& \quad \frac{x \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3943} \\
& - \frac{b \left(\frac{\sin(a + bx + cx^2)}{2c} - \frac{b \int \cos(cx^2 + bx + a) dx}{2c} \right)}{2c} - \\
& \quad \frac{\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} + \frac{x \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3929} \\
& - \frac{b \left(\frac{\sin(a + bx + cx^2)}{2c} - \frac{b \left(\cos(a - \frac{b^2}{4c}) \int \cos(\frac{(b+2cx)^2}{4c}) dx - \sin(a - \frac{b^2}{4c}) \int \sin(\frac{(b+2cx)^2}{4c}) dx \right)}{2c} \right)}{2c} - \\
& \quad \frac{\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} + \frac{x \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3832} \\
& - \frac{b \left(\frac{\sin(a + bx + cx^2)}{2c} - \frac{b \left(\cos(a - \frac{b^2}{4c}) \int \cos(\frac{(b+2cx)^2}{4c}) dx - \frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} \right)}{2c} - \\
& \quad \frac{\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} + \frac{x \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3833}
\end{aligned}$$

$$\begin{aligned}
 & - \frac{b \left(\frac{\sin(a+bx+cx^2)}{2c} - \frac{b \left(\frac{\sqrt{\frac{\pi}{2}} \cos\left(a-\frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a-\frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} \right)}{2c} - \\
 & \frac{\frac{\sqrt{\frac{\pi}{2}} \sin\left(a-\frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a-\frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} + \frac{x \sin(a+bx+cx^2)}{2c}
 \end{aligned}$$

input `Int[x^2*Cos[a + b*x + c*x^2], x]`

output
$$\begin{aligned}
 & -1/2*((\text{Sqrt}[\text{Pi}/2]*\text{Cos}[a - b^2/(4*c)]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}]])]/\text{Sqrt}[c] + (\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}]])*\text{Sin}[a - b^2/(4*c)])/\text{Sqrt}[c])/c + (x*\text{Sin}[a + b*x + c*x^2])/(2*c) - (b*(-1/2*(\text{Sqrt}[\text{Pi}/2]*\text{Cos}[a - b^2/(4*c)]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]}/\text{Sqrt}[c] - (\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*)\text{Sin}[a - b^2/(4*c)])/\text{Sqrt}[c]))/c + \text{Sin}[a + b*x + c*x^2]/(2*c))/(2*c)
 \end{aligned}$$

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3928 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] - Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3929 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3943

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.)*(x_)), x_Symbol]
 :> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c) Int
 [Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b
 *e, 0]
```

rule 3945

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.)*(x_))^(m_), x_Sym
 bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Simp[
 (b*e - 2*c*d)/(2*c) Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] -
 Simp[e^2*((m - 1)/(2*c)) Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x],
 x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Maple [A] (verified)

Time = 2.02 (sec), antiderivative size = 204, normalized size of antiderivative = 0.82

method	result
default	$\frac{x \sin(cx^2+bx+a)}{2c} - \frac{b \left(\frac{\sin(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2}{4c}-ac\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2}{4c}-ac\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} \right)}{2c}$
risch	$\frac{b^2\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{16c^2\sqrt{ic}} - \frac{i\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} - \frac{b^2\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{16c^2\sqrt{-ic}} - \frac{i\sqrt{2}\pi^{\frac{3}{2}}\left(\cos\left(\frac{4ac-b^2}{4c}\right) \left(\frac{\sin(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2}{4c}-ac\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2}{4c}-ac\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} \right)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2\cos\left(\frac{b^2}{4c}-ac\right)\text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi}x^2\sin\left(\frac{b^2}{4c}-ac\right)\text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{i\sqrt{2}\pi^{\frac{3}{2}}\left(\cos\left(\frac{4ac-b^2}{4c}\right) \left(\frac{\sin(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2}{4c}-ac\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2}{4c}-ac\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} \right)}{2c}$

input `int(x^2*cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output

```
1/2*x*sin(c*x^2+b*x+a)/c-1/2*b/c*(1/2*sin(c*x^2+b*x+a)/c-1/4*b/c^(3/2)*2^(1/2)*Pi^(1/2)*(cos((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))+sin((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))))-1/4*c^(3/2)*2^(1/2)*Pi^(1/2)*(cos((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))-sin((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.69

$$\int x^2 \cos(a + bx + cx^2) dx = \frac{\sqrt{2} \left(\pi b^2 \cos\left(-\frac{b^2 - 4ac}{4c}\right) - 2\pi c \sin\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2} \left(\pi b^2 \sin\left(-\frac{b^2 - 4ac}{4c}\right) + 2\pi c \cos\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right)}{8c^3}$$

input

```
integrate(x^2*cos(c*x^2+b*x+a),x, algorithm="fricas")
```

output

```
1/8*(sqrt(2)*(pi*b^2*cos(-1/4*(b^2 - 4*a*c)/c) - 2*pi*c*sin(-1/4*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) - sqrt(2)*(pi*b^2*sin(-1/4*(b^2 - 4*a*c)/c) + 2*pi*c*cos(-1/4*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) + 2*(2*c^2*x - b*c)*sin(c*x^2 + b*x + a))/c^3
```

Sympy [F]

$$\int x^2 \cos(a + bx + cx^2) dx = \int x^2 \cos(a + bx + cx^2) dx$$

input

```
integrate(x**2*cos(c*x**2+b*x+a),x)
```

output

```
Integral(x**2*cos(a + b*x + c*x**2), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.76 (sec) , antiderivative size = 1570, normalized size of antiderivative = 6.31

$$\int x^2 \cos(a + bx + cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*cos(c*x^2+b*x+a),x, algorithm="maxima")`

output

```
1/32*(8*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*cos(-1/4*(b^2 - 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 4*(-(I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*sin(-1/4*(b^2 - 4*a*c)/c))*x^3 + 12*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(-1/4*(b^2 - 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 4*(-(I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*sin(-1/4*(b^2 - 4*a*c)/c))*x^2 + 8*(b*c^2*(I*e^(1/4*(...)))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.90

$$\int x^2 \cos(a + bx + cx^2) dx =$$

$$-\frac{\sqrt{2}\sqrt{\pi}(b^2+2ic)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4ic}{4c}\right)}}{\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2\left(c\left(-2ix-\frac{ib}{c}\right)+2ib\right)e^{(icx^2+ibx+ia)}$$

$$-\frac{\sqrt{2}\sqrt{\pi}(b^2-2ic)\operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4ic}{4c}\right)}}{\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2\left(c\left(2ix+\frac{ib}{c}\right)-2ib\right)e^{(-icx^2-ibx-ia)}$$

$$\frac{16c^2}{16c^2}$$

input `integrate(x^2*cos(c*x^2+b*x+a),x, algorithm="giac")`

output

```
-1/16*(sqrt(2)*sqrt(pi)*(b^2 + 2*I*c)*erf(-1/4*sqrt(2)*(2*x + b/c)*(-I*c/a
bs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*s
qrt(abs(c))) - 2*(c*(-2*I*x - I*b/c) + 2*I*b)*e^(I*c*x^2 + I*b*x + I*a))/c
^2 - 1/16*(sqrt(2)*sqrt(pi)*(b^2 - 2*I*c)*erf(-1/4*sqrt(2)*(2*x + b/c)*(I*
c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1
)*sqrt(abs(c))) - 2*(c*(2*I*x + I*b/c) - 2*I*b)*e^(-I*c*x^2 - I*b*x - I*a))
/c^2
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos(a + bx + cx^2) dx = \int x^2 \cos(cx^2 + bx + a) dx$$

input `int(x^2*cos(a + b*x + c*x^2),x)`

output `int(x^2*cos(a + b*x + c*x^2), x)`

Reduce [F]

$$\int x^2 \cos(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a) x^2 dx$$

input `int(x^2*cos(c*x^2+b*x+a),x)`

output `int(cos(a + b*x + c*x**2)*x**2,x)`

3.2 $\int x \cos(a + bx + cx^2) dx$

Optimal result	50
Mathematica [A] (verified)	50
Rubi [A] (verified)	51
Maple [A] (verified)	52
Fricas [A] (verification not implemented)	53
Sympy [F]	54
Maxima [C] (verification not implemented)	54
Giac [C] (verification not implemented)	55
Mupad [F(-1)]	56
Reduce [F]	56

Optimal result

Integrand size = 13, antiderivative size = 123

$$\begin{aligned} \int x \cos(a + bx + cx^2) dx = & -\frac{b \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & + \frac{b \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} + \frac{\sin(a + bx + cx^2)}{2c} \end{aligned}$$

output

$$-1/4*b*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)+1/4*b*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)+1/2*sin(c*x^2+b*x+a)/c$$

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.91

$$\begin{aligned} & \int x \cos(a + bx + cx^2) dx \\ &= \frac{-b\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + b\sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) + 2\sqrt{c} \sin(a + x(b + cx))}{4c^{3/2}} \end{aligned}$$

input

```
Integrate[x*Cos[a + b*x + c*x^2], x]
```

output
$$\left(-\frac{b \operatorname{Sqrt}[2 \operatorname{Pi}] \cos[a - b^2/(4c)] \operatorname{FresnelC}[(b + 2cx)/(\operatorname{Sqrt}[c] \operatorname{Sqrt}[2 \operatorname{Pi}])]}{4c^{3/2}} + \frac{b \operatorname{Sqrt}[2 \operatorname{Pi}] \operatorname{FresnelS}[(b + 2cx)/(\operatorname{Sqrt}[c] \operatorname{Sqrt}[2 \operatorname{Pi})]}{\sqrt{c}} \sin[a - b^2/(4c)] + \frac{2 \operatorname{Sqrt}[c] \sin[a + x(b + cx)]}{4c^{3/2}} \right)$$

Rubi [A] (verified)

Time = 0.33 (sec), antiderivative size = 125, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3943, 3929, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(a + bx + cx^2) dx \\
 & \downarrow 3943 \\
 & \frac{\sin(a + bx + cx^2)}{2c} - \frac{b \int \cos(cx^2 + bx + a) dx}{2c} \\
 & \downarrow 3929 \\
 & \frac{\sin(a + bx + cx^2)}{2c} - \frac{b \left(\cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx - \sin\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx \right)}{2c} \\
 & \downarrow 3832 \\
 & \frac{\sin(a + bx + cx^2)}{2c} - \frac{b \left(\cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} \\
 & \downarrow 3833 \\
 & \frac{\sin(a + bx + cx^2)}{2c} - \frac{b \left(\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c}
 \end{aligned}$$

input
$$\operatorname{Int}[x \cos[a + b x + c x^2], x]$$

output

$$-1/2 * (b * ((\text{Sqrt}[\text{Pi}/2] * \text{Cos}[a - b^2/(4*c)]) * \text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]) / \text{Sqrt}[c] - (\text{Sqrt}[\text{Pi}/2] * \text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] * \text{Sin}[a - b^2/(4*c)]) / \text{Sqrt}[c])) / c + \text{Sin}[a + b*x + c*x^2] / (2*c)$$

Defintions of rubi rules used

rule 3832

$$\text{Int}[\text{Sin}[(d_.) * ((e_.) + (f_.) * (x_)) ^ 2], x_Symbol] :> \text{Simp}[(\text{Sqrt}[\text{Pi}/2] / (f * \text{Rt}[d, 2])) * \text{FresnelS}[\text{Sqrt}[2/\text{Pi}] * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$$

rule 3833

$$\text{Int}[\text{Cos}[(d_.) * ((e_.) + (f_.) * (x_)) ^ 2], x_Symbol] :> \text{Simp}[(\text{Sqrt}[\text{Pi}/2] / (f * \text{Rt}[d, 2])) * \text{FresnelC}[\text{Sqrt}[2/\text{Pi}] * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$$

rule 3929

$$\begin{aligned} \text{Int}[\text{Cos}[(a_.) + (b_.) * (x_) + (c_.) * (x_) ^ 2], x_Symbol] &:> \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \\ &\quad \text{Int}[\text{Cos}[(b + 2*c*x) ^ 2 / (4*c)], x], x] + \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \\ &\quad \text{Int}[\text{Sin}[(b + 2*c*x) ^ 2 / (4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0] \end{aligned}$$

rule 3943

$$\begin{aligned} \text{Int}[\text{Cos}[(a_.) + (b_.) * (x_) + (c_.) * (x_) ^ 2] * ((d_.) + (e_.) * (x_)), x_Symbol] &:> \text{Simp}[e * (\text{Sin}[a + b*x + c*x^2] / (2*c)), x] + \text{Simp}[(2*c*d - b*e) / (2*c) \\ &\quad \text{Int}[\text{Cos}[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b * e, 0] \end{aligned}$$

Maple [A] (verified)

Time = 1.14 (sec), antiderivative size = 99, normalized size of antiderivative = 0.80

method	result
default	$\frac{\sin(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2}{4}-ac\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2}{4}-ac\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{3/2}}$
risch	$-\frac{b\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} + \frac{b\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} + \frac{\sin(cx^2+bx+a)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x \cos\left(\frac{b^2}{4}-ac\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi}x \sin\left(\frac{b^2}{4}-ac\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\cos\left(\frac{b^2}{4}-ac\right)\sqrt{2}\sqrt{\pi}}{\sqrt{c}}$

input `int(x*cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output `1/2*sin(c*x^2+b*x+a)/c-1/4*b/c^(3/2)*2^(1/2)*Pi^(1/2)*(cos((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))+sin((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.97

$$\int x \cos(a + bx + cx^2) dx = \frac{\sqrt{2}\pi b \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi b \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2-4ac}{4c}\right) - 2c \sin(cx^2)}{4c^2}$$

input `integrate(x*cos(c*x^2+b*x+a),x, algorithm="fricas")`

output

```
-1/4*(sqrt(2)*pi*b*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) - sqrt(2)*pi*b*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c) - 2*c*sin(c*x^2 + b*x + a))/c^2
```

Sympy [F]

$$\int x \cos(a + bx + cx^2) dx = \int x \cos(a + bx + cx^2) dx$$

input

```
integrate(x*cos(c*x**2+b*x+a),x)
```

output

```
Integral(x*cos(a + b*x + c*x**2), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 580, normalized size of antiderivative = 4.72

$$\int x \cos(a + bx + cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*cos(c*x^2+b*x+a),x, algorithm="maxima")
```

output

```
1/16*((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/4*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/4*(b^2 - 4*a*c)/c) - 2*(-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/4*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/4*(b^2 - 4*a*c)/c))*x - 4*(c*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^2 - 4*a*c)/c) - c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) *sin(-1/4*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.47

$$\int x \cos(a + bx + cx^2) dx$$

$$= \frac{\frac{\sqrt{2}\sqrt{\pi}b \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(-\frac{i}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4ia}{4c}\right)}}{\left(-\frac{i}{|c|}+1\right)\sqrt{|c|}} - 2i e^{(icx^2+ibx+ia)}}{8c}$$

$$+ \frac{\frac{\sqrt{2}\sqrt{\pi}b \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}\left(2x+\frac{b}{c}\right)\left(\frac{i}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4ia}{4c}\right)}}{\left(\frac{i}{|c|}+1\right)\sqrt{|c|}} + 2i e^{(-icx^2-ibx-ia)}}{8c}$$

input `integrate(x*cos(c*x^2+b*x+a),x, algorithm="giac")`

output

```
1/8*(sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*e^(I*c*x^2 + I*b*x + I*a))/c + 1/8*(sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c))/((I*c/abs(c) + 1)*sqrt(abs(c))) + 2*I*e^(-I*c*x^2 - I*b*x - I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \cos(a + bx + cx^2) dx = \int x \cos(cx^2 + bx + a) dx$$

input

```
int(x*cos(a + b*x + c*x^2),x)
```

output

```
int(x*cos(a + b*x + c*x^2), x)
```

Reduce [F]

$$\int x \cos(a + bx + cx^2) dx = \frac{-(\int \cos(cx^2 + bx + a) dx) b + \sin(cx^2 + bx + a)}{2c}$$

input

```
int(x*cos(c*x^2+b*x+a),x)
```

output

```
( - int(cos(a + b*x + c*x**2),x)*b + sin(a + b*x + c*x**2))/(2*c)
```

3.3 $\int \cos(a + bx + cx^2) dx$

Optimal result	57
Mathematica [A] (verified)	57
Rubi [A] (verified)	58
Maple [A] (verified)	59
Fricas [A] (verification not implemented)	60
Sympy [A] (verification not implemented)	60
Maxima [C] (verification not implemented)	61
Giac [C] (verification not implemented)	61
Mupad [B] (verification not implemented)	62
Reduce [F]	62

Optimal result

Integrand size = 11, antiderivative size = 98

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{\sqrt{c}}$$

output

```
1/2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(1/2)-1/2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.87

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \left(\cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \right)}{\sqrt{c}}$$

input `Integrate[Cos[a + b*x + c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot (\text{Cos}[a - b^2/(4c]) \cdot \text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] - \text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] \cdot \text{Sin}[a - b^2/(4c)]))}{\text{Sqrt}[c]}$$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3929, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(a + bx + cx^2) dx \\
 & \downarrow 3929 \\
 & \cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) dx - \sin\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b + 2cx)^2}{4c}\right) dx \\
 & \downarrow 3832 \\
 & \cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\
 & \downarrow 3833 \\
 & \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}
 \end{aligned}$$

input `Int[Cos[a + b*x + c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{Cos}[a - b^2/(4c]) \cdot \text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]}{\text{Sqrt}[c]} - \frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] \cdot \text{Sin}[a - b^2/(4c)])}{\text{Sqrt}[c]}$$

Definitions of rubi rules used

rule 3832 $\text{Int}[\text{Sin}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\text{Int}[\text{Cos}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3929 $\text{Int}[\text{Cos}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] + \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{\frac{b^2}{4}-ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)+\sin\left(\frac{\frac{b^2}{4}-ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{c}}\right)\right)}{2\sqrt{c}}$	81
risch	$\frac{\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}\text{erf}\left(\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}}-\frac{\sqrt{\pi}e^{\frac{i(4ac-b^2)}{4c}}\text{erf}\left(-\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}}$	99

input `int(cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2}2^{(1/2)}\text{Pi}^{(1/2)}c^{(1/2)}(\cos((1/4b^2-a*c)/c)\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)})c^{(1/2)}(c*x+1/2*b))+\sin((1/4b^2-a*c)/c)\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)})c^{(1/2)}(c*x+1/2*b))$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.05

$$\int \cos(a + bx + cx^2) dx \\ = \frac{\sqrt{2}\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi\sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2 - 4ac}{4c}\right)}{2c}$$

input `integrate(cos(c*x^2+b*x+a),x, algorithm="fricas")`

output `1/2*(sqrt(2)*pi*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) - sqrt(2)*pi*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c))/c`

Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.90

$$\int \cos(a + bx + cx^2) dx \\ = \frac{\sqrt{2}\sqrt{\pi}\left(-\sin\left(a - \frac{b^2}{4c}\right)S\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) + \cos\left(a - \frac{b^2}{4c}\right)C\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right)\right)\sqrt{\frac{1}{c}}}{2}$$

input `integrate(cos(c*x**2+b*x+a),x)`

output `sqrt(2)*sqrt(pi)*(-sin(a - b**2/(4*c))*fresnels(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))) + cos(a - b**2/(4*c))*fresnelc(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))))*sqrt(1/c)/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14

$$\int \cos(a + bx + cx^2) dx = -\frac{\sqrt{2}\sqrt{\pi} \left(\left((i-1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i+1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) + \left((i+1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i-1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

input `integrate(cos(c*x^2+b*x+a),x, algorithm="maxima")`

output `-1/8*sqrt(2)*sqrt(pi)*((I - 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(I*c)) + ((I + 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(-I*c)))/sqrt(c)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.38

$$\int \cos(a + bx + cx^2) dx = -\frac{\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x + \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{ib^2-4ac}{4c}\right)}}{4\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}} - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2+4ac}{4c}\right)}}{4\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}}$$

input `integrate(cos(c*x^2+b*x+a),x, algorithm="giac")`

output `-1/4*sqrt(2)*sqrt(pi)*erf(-1/4*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 1/4*sqrt(2)*sqrt(pi)*erf(-1/4*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c)))`

Mupad [B] (verification not implemented)

Time = 41.47 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.01

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} \cos\left(\frac{4ac - b^2}{4c}\right) C\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2} - \frac{\sqrt{2} \sqrt{\pi} \sin\left(\frac{4ac - b^2}{4c}\right) S\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2}$$

input `int(cos(a + b*x + c*x^2),x)`

output `(2^(1/2)*pi^(1/2)*cos((4*a*c - b^2)/(4*c))*fresnelc((2^(1/2)*(b/2 + c*x)*(1/c)^(1/2))/pi^(1/2)*(1/c)^(1/2))/2 - (2^(1/2)*pi^(1/2)*sin((4*a*c - b^2)/(4*c))*fresnels((2^(1/2)*(b/2 + c*x)*(1/c)^(1/2))/pi^(1/2)*(1/c)^(1/2))/2`

Reduce [F]

$$\int \cos(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a) dx$$

input `int(cos(c*x^2+b*x+a),x)`

output `int(cos(a + b*x + c*x**2),x)`

3.4 $\int \frac{\cos(a+bx+cx^2)}{x} dx$

Optimal result	63
Mathematica [N/A]	63
Rubi [N/A]	64
Maple [N/A]	64
Fricas [N/A]	65
Sympy [N/A]	65
Maxima [N/A]	66
Giac [N/A]	66
Mupad [N/A]	66
Reduce [N/A]	67

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \text{Int}\left(\frac{\cos(a + bx + cx^2)}{x}, x\right)$$

output `Defer(Int)(cos(c*x^2+b*x+a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.50 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(a + bx + cx^2)}{x} dx$$

input `Integrate[Cos[a + b*x + c*x^2]/x, x]`

output `Integrate[Cos[a + b*x + c*x^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3951}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + bx + cx^2)}{x} dx$$

↓ 3951

$$\int \frac{\cos(a + bx + cx^2)}{x} dx$$

input `Int[Cos[a + b*x + c*x^2]/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3951 `Int[Cos[(a_) + (b_)*(x_) + (c_)*(x_)^2]^(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] :> Unintegrable[(d + e*x)^m*Cos[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos(cx^2 + bx + a)}{x} dx$$

input `int(cos(c*x^2+b*x+a)/x, x)`

output `int(cos(c*x^2+b*x+a)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

input `integrate(cos(c*x^2+b*x+a)/x,x, algorithm="fricas")`

output `integral(cos(c*x^2 + b*x + a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(a + bx + cx^2)}{x} dx$$

input `integrate(cos(c*x**2+b*x+a)/x,x)`

output `Integral(cos(a + b*x + c*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

input `integrate(cos(c*x^2+b*x+a)/x,x, algorithm="maxima")`

output `integrate(cos(c*x^2 + b*x + a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

input `integrate(cos(c*x^2+b*x+a)/x,x, algorithm="giac")`

output `integrate(cos(c*x^2 + b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 41.38 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

input `int(cos(a + b*x + c*x^2)/x,x)`

output `int(cos(a + b*x + c*x^2)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

input `int(cos(c*x^2+b*x+a)/x,x)`

output `int(cos(a + b*x + c*x**2)/x,x)`

3.5 $\int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx$

Optimal result	68
Mathematica [A] (verified)	68
Rubi [A] (verified)	69
Maple [F]	70
Fricas [A] (verification not implemented)	70
Sympy [F]	71
Maxima [F]	71
Giac [F]	71
Mupad [F(-1)]	72
Reduce [F]	72

Optimal result

Integrand size = 32, antiderivative size = 111

$$\begin{aligned} & \int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx \\ &= -\frac{\cos(a+bx+cx^2)}{x} - \sqrt{c}\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad - \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \end{aligned}$$

output

```
-cos(c*x^2+b*x+a)/x-c^(1/2)*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))-c^(1/2)*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)
```

Mathematica [A] (verified)

Time = 2.96 (sec), antiderivative size = 110, normalized size of antiderivative = 0.99

$$\begin{aligned} & \int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx = \\ & \frac{\cos(a+x(b+cx)) + \sqrt{c}\sqrt{2\pi}x \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \sqrt{c}\sqrt{2\pi}x \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{x} \end{aligned}$$

input $\text{Integrate}[\cos[a + b*x + c*x^2]/x^2 + (\sin[a + b*x + c*x^2])/x, x]$

output $-\left(\frac{\cos[a + x*(b + c*x)]}{x} + \sqrt{c}\sqrt{2\pi}x\cos[a - b^2/(4*c)]\text{FresnelS}\left[\frac{(b + 2*c*x)}{\sqrt{c}\sqrt{2\pi}}\right] + \sqrt{c}\sqrt{2\pi}x\text{FresnelC}\left[\frac{(b + 2*c*x)}{\sqrt{c}\sqrt{2\pi}}\right]\sin[a - b^2/(4*c)]\right)/x$

Rubi [A] (verified)

Time = 0.30 (sec), antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.031$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \left(\frac{b \sin(a + bx + cx^2)}{x} + \frac{\cos(a + bx + cx^2)}{x^2} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\sqrt{2\pi}\sqrt{c}\sin\left(a - \frac{b^2}{4c}\right)\text{FresnelC}\left(\frac{b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \sqrt{2\pi}\sqrt{c}\cos\left(a - \frac{b^2}{4c}\right)\text{FresnelS}\left(\frac{b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \\ & \quad \frac{\cos(a + bx + cx^2)}{x} \end{aligned}$$

input $\text{Int}[\cos[a + b*x + c*x^2]/x^2 + (\sin[a + b*x + c*x^2])/x, x]$

output $-(\cos[a + b*x + c*x^2]/x) - \sqrt{c}\sqrt{2\pi}\cos[a - b^2/(4*c)]\text{FresnelS}\left[\frac{(b + 2*c*x)}{\sqrt{c}\sqrt{2\pi}}\right] - \sqrt{c}\sqrt{2\pi}\text{FresnelC}\left[\frac{(b + 2*c*x)}{\sqrt{c}\sqrt{2\pi}}\right]\sin[a - b^2/(4*c)]$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

Maple [F]

$$\int \left(\frac{\cos(cx^2 + bx + a)}{x^2} + \frac{b \sin(cx^2 + bx + a)}{x} \right) dx$$

input $\text{int}(\cos(c*x^2+b*x+a)/x^2+b*\sin(c*x^2+b*x+a)/x, x)$

output $\text{int}(\cos(c*x^2+b*x+a)/x^2+b*\sin(c*x^2+b*x+a)/x, x)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 115, normalized size of antiderivative = 1.04

$$\int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx = \\ - \frac{\sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2-4ac}{4c}\right) + \cos(cx^2 + bx + a)}{x}$$

input $\text{integrate}(\cos(c*x^2+b*x+a)/x^2+b*\sin(c*x^2+b*x+a)/x, x, \text{algorithm}=\text{"fricas"})$

output $-(\sqrt{2}*\pi*x*\sqrt{c/\pi})*\cos(-1/4*(b^2 - 4*a*c)/c)*\text{fresnel_sin}(1/2*\sqrt{2})*(2*c*x + b)*\sqrt{c/\pi}/c + \sqrt{2}*\pi*x*\sqrt{c/\pi}*\text{fresnel_cos}(1/2*\sqrt{2})*(2*c*x + b)*\sqrt{c/\pi}/c)*\sin(-1/4*(b^2 - 4*a*c)/c) + \cos(c*x^2 + bx + a))/x$

Sympy [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{bx \sin(a + bx + cx^2) + \cos(a + bx + cx^2)}{x^2} dx \end{aligned}$$

input `integrate(cos(c*x**2+b*x+a)/x**2+b*sin(c*x**2+b*x+a)/x,x)`

output `Integral((b*x*sin(a + b*x + c*x**2) + cos(a + b*x + c*x**2))/x**2, x)`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(cx^2 + bx + a)}{x} + \frac{\cos(cx^2 + bx + a)}{x^2} dx \end{aligned}$$

input `integrate(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x,x, algorithm="maxima")`

output `integrate(b*sin(c*x^2 + b*x + a)/x + cos(c*x^2 + b*x + a)/x^2, x)`

Giac [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(cx^2 + bx + a)}{x} + \frac{\cos(cx^2 + bx + a)}{x^2} dx \end{aligned}$$

input `integrate(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x,x, algorithm="giac")`

output `integrate(b*sin(c*x^2 + b*x + a)/x + cos(c*x^2 + b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{\cos(cx^2 + bx + a)}{x^2} + \frac{b \sin(cx^2 + bx + a)}{x} dx \end{aligned}$$

input `int(cos(a + b*x + c*x^2)/x^2 + (b*sin(a + b*x + c*x^2))/x, x)`

output `int(cos(a + b*x + c*x^2)/x^2 + (b*sin(a + b*x + c*x^2))/x, x)`

Reduce [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \frac{\left(\int \frac{\cos(cx^2 + bx + a)}{x^2} dx \right) x + \left(\int \frac{\sin(cx^2 + bx + a)}{x} dx \right) bx + \left(\int \frac{1}{x^2} dx \right) x + 1}{x} \end{aligned}$$

input `int(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x, x)`

output `(int(cos(a + b*x + c*x**2)/x**2, x)*x + int(sin(a + b*x + c*x**2)/x, x)*b*x + int(1/x**2, x)*x + 1)/x`

3.6 $\int x^2 \cos(a + bx - cx^2) dx$

Optimal result	73
Mathematica [A] (verified)	74
Rubi [A] (verified)	74
Maple [A] (verified)	77
Fricas [A] (verification not implemented)	78
Sympy [F]	79
Maxima [C] (verification not implemented)	79
Giac [C] (verification not implemented)	80
Mupad [F(-1)]	81
Reduce [F]	81

Optimal result

Integrand size = 16, antiderivative size = 251

$$\begin{aligned} \int x^2 \cos(a + bx - cx^2) dx = & -\frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} \\ & - \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{4c^{5/2}} \\ & - \frac{b \sin(a + bx - cx^2)}{4c^2} - \frac{x \sin(a + bx - cx^2)}{2c} \end{aligned}$$

output

```
-1/8*b^2*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelC(1/2*(-2*c*x+b)/c^(1/2)
*2^(1/2)/Pi^(1/2))/c^(5/2)+1/4*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelS(
1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)-1/4*2^(1/2)*Pi^(1/2)*Fres
nelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)/c^(3/2)-1/8
*b^2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*si
n(a+1/4*b^2/c)/c^(5/2)-1/4*b*sin(-c*x^2+b*x+a)/c^2-1/2*x*sin(-c*x^2+b*x+a)
/c
```

Mathematica [A] (verified)

Time = 0.76 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.65

$$\int x^2 \cos(a + bx - cx^2) dx \\ = \frac{-\sqrt{2\pi} \operatorname{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a + \frac{b^2}{4c}\right) - b^2 \sin\left(a + \frac{b^2}{4c}\right)\right) + \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(b^2 \cos\left(a + \frac{b^2}{4c}\right) + 2c \sin\left(a + \frac{b^2}{4c}\right)\right)}{8c^{5/2}}$$

input `Integrate[x^2*Cos[a + b*x - c*x^2], x]`

output $\frac{(-\operatorname{Sqrt}[2*\operatorname{Pi}]*\operatorname{FresnelS}[(-b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]*(2*c*\operatorname{Cos}[a + b^2/(4*c)] - b^2*\operatorname{Sin}[a + b^2/(4*c)])) + \operatorname{Sqrt}[2*\operatorname{Pi}]*\operatorname{FresnelC}[(-b + 2*c*x)/(\operatorname{Sqr} t[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]*(b^2*\operatorname{Cos}[a + b^2/(4*c)] + 2*c*\operatorname{Sin}[a + b^2/(4*c)]) - 2*\operatorname{Sqr} t[c]*(b + 2*c*x)*\operatorname{Sin}[a + x*(b - c*x)]}{(8*c^{(5/2)})}$

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3945, 3928, 25, 3832, 3833, 3943, 3929, 25, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos(a + bx - cx^2) dx \\ \downarrow 3945 \\ \frac{\int \sin(-cx^2 + bx + a) dx}{2c} + \frac{b \int x \cos(-cx^2 + bx + a) dx}{2c} - \frac{x \sin(a + bx - cx^2)}{2c} \\ \downarrow 3928 \\ \frac{\sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx + \cos\left(a + \frac{b^2}{4c}\right) \int -\sin\left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} + \\ \frac{b \int x \cos(-cx^2 + bx + a) dx}{2c} - \frac{x \sin(a + bx - cx^2)}{2c}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{\sin \left(a + \frac{b^2}{4c}\right) \int \cos \left(\frac{(b-2cx)^2}{4c}\right) dx - \cos \left(a + \frac{b^2}{4c}\right) \int \sin \left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} + \\
& \frac{b \int x \cos(-cx^2 + bx + a) dx}{2c} - \frac{x \sin(a + bx - cx^2)}{2c} \\
& \downarrow 3832 \\
& \frac{\sin \left(a + \frac{b^2}{4c}\right) \int \cos \left(\frac{(b-2cx)^2}{4c}\right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos(a + \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} + \\
& \frac{b \int x \cos(-cx^2 + bx + a) dx}{2c} - \frac{x \sin(a + bx - cx^2)}{2c} \\
& \downarrow 3833 \\
& \frac{b \int x \cos(-cx^2 + bx + a) dx}{2c} + \frac{\frac{\sqrt{\frac{\pi}{2}} \cos(a + \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a + \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \\
& \frac{x \sin(a + bx - cx^2)}{2c} \\
& \downarrow 3943 \\
& \frac{b \left(\frac{b \int \cos(-cx^2 + bx + a) dx}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos(a + \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a + \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \sin(a + bx - cx^2)}{2c} \\
& \downarrow 3929 \\
& \frac{b \left(\frac{b \left(\cos(a + \frac{b^2}{4c}) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx - \sin(a + \frac{b^2}{4c}) \int -\sin\left(\frac{(b-2cx)^2}{4c}\right) dx \right)}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos(a + \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a + \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \sin(a + bx - cx^2)}{2c} \\
& \downarrow 25 \\
& \frac{b \left(\frac{b \left(\sin(a + \frac{b^2}{4c}) \int \sin\left(\frac{(b-2cx)^2}{4c}\right) dx + \cos(a + \frac{b^2}{4c}) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx \right)}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \right)}{2c} + \\
& \frac{\frac{\sqrt{\frac{\pi}{2}} \cos(a + \frac{b^2}{4c}) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin(a + \frac{b^2}{4c}) \operatorname{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}}{2c} - \frac{x \sin(a + bx - cx^2)}{2c}
\end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3832} \\
 b \left(\frac{\frac{b \left(\cos \left(a + \frac{b^2}{4c} \right) \int \cos \left(\frac{(b-2cx)^2}{4c} \right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}}}{2c} - \frac{\sin(a+bx-cx^2)}{2c} \right)}{+} \right. \\
 & \left. \frac{\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{x \sin(a+bx-cx^2)}{2c}}{2c} \right. \\
 & \downarrow \text{3833} \\
 b \left(\frac{\frac{b \left(-\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} - \frac{\sin(a+bx-cx^2)}{2c} \right) \\
 & \left. + \frac{\frac{\sqrt{\frac{\pi}{2}} \cos \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin \left(a + \frac{b^2}{4c} \right) \operatorname{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} - \frac{x \sin(a+bx-cx^2)}{2c}}{2c} \right)
 \end{aligned}$$

input `Int[x^2*Cos[a + b*x - c*x^2], x]`

output

```
((Sqrt[Pi/2]*Cos[a + b^2/(4*c)]*FresnelS[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/Sqrt[c] - (Sqrt[Pi/2]*FresnelC[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a + b^2/(4*c)]/Sqrt[c])/(2*c) - (x*Sin[a + b*x - c*x^2])/(2*c) + (b*((b*(-((Sqrt[Pi/2]*Cos[a + b^2/(4*c)]*FresnelC[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/Sqrt[c]) - (Sqrt[Pi/2]*FresnelS[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a + b^2/(4*c)])/Sqrt[c]))/(2*c) - Sin[a + b*x - c*x^2]/(2*c)))/(2*c)
```

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_._)*(x_))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 $\text{Int}[\cos[(d_*) \cdot ((e_*) + (f_*) \cdot (x_*)^2)], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi}/2) / (f \cdot \sqrt{d, 2})] * \text{FresnelC}[\sqrt{2/\pi} \cdot \sqrt{d, 2} \cdot (e + f \cdot x), x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\text{Int}[\sin[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\cos[(b^2 - 4*a*c)/(4*c)] \cdot \text{Int}[\sin[(b + 2*c*x)^2/(4*c)], x], x] - \text{Simp}[\sin[(b^2 - 4*a*c)/(4*c)] \cdot \text{Int}[\cos[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3929 $\text{Int}[\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\cos[(b^2 - 4*a*c)/(4*c)] \cdot \text{Int}[\cos[(b + 2*c*x)^2/(4*c)], x], x] + \text{Simp}[\sin[(b^2 - 4*a*c)/(4*c)] \cdot \text{Int}[\sin[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3943 $\text{Int}[\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2] * ((d_*) + (e_*) \cdot (x_*)), x_{\text{Symbol}}] \rightarrow \text{Simp}[e * (\sin[a + b*x + c*x^2]/(2*c)), x] + \text{Simp}[(2*c*d - b*e)/(2*c) \cdot \text{Int}[\cos[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

rule 3945 $\text{Int}[\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2] * ((d_*) + (e_*) \cdot (x_*)^{(m)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[e * (d + e*x)^{(m-1)} * (\sin[a + b*x + c*x^2]/(2*c)), x] + (-\text{Simp}[(b*e - 2*c*d)/(2*c) \cdot \text{Int}[(d + e*x)^{(m-1)} * \cos[a + b*x + c*x^2], x], x] - \text{Simp}[e^{2*(m-1)/(2*c)} \cdot \text{Int}[(d + e*x)^{(m-2)} * \sin[a + b*x + c*x^2], x], x]) /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[b*e - 2*c*d, 0] \&& \text{GtQ}[m, 1]$

Maple [A] (verified)

Time = 1.73 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.89

method	result
default	$-\frac{x \sin(-cx^2+bx+a)}{2c} + \frac{b \left(-\frac{\sin(-cx^2+bx+a)}{2c} + \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2+ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) - \sin\left(\frac{b^2+ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) \right)}{2c} \right)}{2c}$
risch	$\frac{b^2\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\text{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{16c^2\sqrt{-ic}} + \frac{i\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\text{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} - \frac{b^2\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\text{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{16c^2\sqrt{ic}}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2\cos\left(\frac{b^2+ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}x^2\sin\left(\frac{b^2+ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} + \frac{\sqrt{2}\pi^{\frac{3}{2}}\left(\cos\left(\frac{4ac+b^2}{4c}\right)\right)}{2\sqrt{-c}}$

input `int(x^2*cos(-c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$-1/2*x*\sin(-c*x^2+b*x+a)/c+1/2*b/c*(-1/2*\sin(-c*x^2+b*x+a)/c+1/4*b/c*2^(1/2)*Pi^(1/2)/(-c)^(1/2)*(cos((1/4*b^2+a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b)))-sin((1/4*b^2+a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b))))+1/4/c*2^(1/2)*Pi^(1/2)/(-c)^(1/2)*(cos((1/4*b^2+a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b))+sin((1/4*b^2+a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/(-c)^(1/2)*(-c*x+1/2*b))))$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.71

$$\int x^2 \cos(a + bx - cx^2) dx = \frac{\sqrt{2} \left(\pi b^2 \cos\left(\frac{b^2+4ac}{4c}\right) + 2\pi c \sin\left(\frac{b^2+4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2} \left(\pi b^2 \sin\left(\frac{b^2+4ac}{4c}\right) - 2\pi c \cos\left(\frac{b^2+4ac}{4c}\right) \right)}{8c^3}$$

input `integrate(x^2*cos(-c*x^2+b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{8} \left(\sqrt{2} \left(\pi b^2 \cos\left(\frac{1}{4} (b^2 + 4ac)/c\right) + 2\pi c \sin\left(\frac{1}{4} (b^2 + 4ac)/c\right) \right) \sqrt{\frac{c}{\pi}} \operatorname{fresnel_cos}\left(\frac{1}{2} \sqrt{2} (2cx - b) \sqrt{\frac{c}{\pi}}/c\right) + \sqrt{2} \left(\pi b^2 \sin\left(\frac{1}{4} (b^2 + 4ac)/c\right) - 2\pi c \cos\left(\frac{1}{4} (b^2 + 4ac)/c\right) \right) \sqrt{\frac{c}{\pi}} \operatorname{fresnel_sin}\left(\frac{1}{2} \sqrt{2} (2cx - b) \sqrt{\frac{c}{\pi}}/c\right) + 2(2c^2x + b^2c) \sin(cx^2 - bx - a) \right) / c^3$$

Sympy [F]

$$\int x^2 \cos(a + bx - cx^2) dx = \int x^2 \cos(a + bx - cx^2) dx$$

input `integrate(x**2*cos(-c*x**2+b*x+a),x)`

output `Integral(x**2*cos(a + b*x - c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.91 (sec) , antiderivative size = 1570, normalized size of antiderivative = 6.25

$$\int x^2 \cos(a + bx - cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*cos(-c*x^2+b*x+a),x, algorithm="maxima")`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.90

$$\int x^2 \cos(a + bx - cx^2) dx =$$

$$-\frac{\sqrt{2} \sqrt{\pi} (b^2+2 i c) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2 x-\frac{b}{c}\right) \left(-\frac{i c}{|c|}+1\right) \sqrt{|c|}\right) e^{\left(-\frac{i b^2+4 i a c}{4 c}\right)}}{\left(-\frac{i c}{|c|}+1\right) \sqrt{|c|}}-2 \left(c \left(-2 i x+\frac{i b}{c}\right)-2 i b\right) e^{(i c x^2-i b x-i a)}$$

$$-\frac{16 c^2}{\sqrt{2} \sqrt{\pi} (b^2-2 i c) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2 x-\frac{b}{c}\right) \left(\frac{i c}{|c|}+1\right) \sqrt{|c|}\right) e^{\left(-\frac{-i b^2-4 i a c}{4 c}\right)}}-\left(c \left(2 i x-\frac{i b}{c}\right)+2 i b\right) e^{(-i c x^2+i b x+i a)}$$

$$-\frac{16 c^2}{16 c^2}$$

```
input integrate(x^2*cos(-c*x^2+b*x+a),x, algorithm="giac")
```

output

```
-1/16*(sqrt(2)*sqrt(pi)*(b^2 + 2*I*c)*erf(-1/4*sqrt(2)*(2*x - b/c)*(-I*c/a
bs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((-I*c/abs(c) + 1)*s
qrt(abs(c))) - 2*(c*(-2*I*x + I*b/c) - 2*I*b)*e^(I*c*x^2 - I*b*x - I*a))/c
^2 - 1/16*(sqrt(2)*sqrt(pi)*(b^2 - 2*I*c)*erf(-1/4*sqrt(2)*(2*x - b/c)*(I*
c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/c)/((I*c/abs(c) + 1
)*sqrt(abs(c))) - 2*(c*(2*I*x - I*b/c) + 2*I*b)*e^(-I*c*x^2 + I*b*x + I*a)
)/c^2
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos(a + bx - cx^2) dx = \int x^2 \cos(-cx^2 + bx + a) dx$$

input

```
int(x^2*cos(a + b*x - c*x^2),x)
```

output

```
int(x^2*cos(a + b*x - c*x^2), x)
```

Reduce [F]

$$\int x^2 \cos(a + bx - cx^2) dx = \int \cos(-cx^2 + bx + a) x^2 dx$$

input

```
int(x^2*cos(-c*x^2+b*x+a),x)
```

output

```
int(cos(a + b*x - c*x**2)*x**2,x)
```

3.7 $\int x \cos(a + bx - cx^2) dx$

Optimal result	82
Mathematica [A] (verified)	82
Rubi [A] (verified)	83
Maple [A] (verified)	85
Fricas [A] (verification not implemented)	85
Sympy [F]	86
Maxima [C] (verification not implemented)	86
Giac [C] (verification not implemented)	87
Mupad [F(-1)]	88
Reduce [F]	88

Optimal result

Integrand size = 14, antiderivative size = 124

$$\begin{aligned} \int x \cos(a + bx - cx^2) dx = & -\frac{b\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{b\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} \\ & - \frac{\sin(a + bx - cx^2)}{2c} \end{aligned}$$

output

$$-\frac{1}{4}b^2\sqrt{\frac{\pi}{2}}\text{Pi}^{(1/2)}\cos\left(a+\frac{1}{4}b^2/c\right)\text{FresnelC}\left(\frac{1}{2}\sqrt{-2c}\sqrt{x+b}/c^{(1/2)}\right)\sqrt{2}\text{Pi}^{(1/2)}/c^{(3/2)} - \frac{1}{4}b^2\sqrt{\frac{\pi}{2}}\text{Pi}^{(1/2)}\text{FresnelS}\left(\frac{1}{2}\sqrt{-2c}\sqrt{x+b}/c^{(1/2)}\right)\sqrt{2}\text{Pi}^{(1/2)}/c^{(3/2)} - \frac{1}{2}\sin(a+bx-cx^2)/c$$

Mathematica [A] (verified)

Time = 0.38 (sec), antiderivative size = 116, normalized size of antiderivative = 0.94

$$\begin{aligned} \int x \cos(a + bx - cx^2) dx \\ = \frac{b\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + b\sqrt{2\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) - 2\sqrt{c} \sin(a + x(b - cx))}{4c^{3/2}} \end{aligned}$$

input `Integrate[x*Cos[a + b*x - c*x^2], x]`

output
$$\frac{(b \sqrt{2\pi}) \cos(a + b^2/(4c)) \operatorname{FresnelC}((-b + 2cx)/(\sqrt{c}\sqrt{2\pi}))}{\sqrt{c}} + \frac{b \sqrt{2\pi} \operatorname{FresnelS}((-b + 2cx)/(\sqrt{c}\sqrt{2\pi})) \sin(a + b^2/(4c))}{\sqrt{c}} - \frac{2 \sqrt{c} \sin(a + x(b - cx))}{c^{3/2}}$$

Rubi [A] (verified)

Time = 0.34 (sec), antiderivative size = 127, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3943, 3929, 25, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos(a + bx - cx^2) dx \\
 & \downarrow 3943 \\
 & \frac{b \int \cos(-cx^2 + bx + a) dx}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \\
 & \downarrow 3929 \\
 & \frac{b \left(\cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx - \sin\left(a + \frac{b^2}{4c}\right) \int -\sin\left(\frac{(b-2cx)^2}{4c}\right) dx \right)}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \\
 & \downarrow 25 \\
 & \frac{b \left(\sin\left(a + \frac{b^2}{4c}\right) \int \sin\left(\frac{(b-2cx)^2}{4c}\right) dx + \cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx \right)}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \\
 & \downarrow 3832 \\
 & \frac{b \left(\cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} - \frac{\sin(a + bx - cx^2)}{2c} \\
 & \downarrow 3833
 \end{aligned}$$

$$\frac{b \left(-\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} - \frac{\sin(a + bx - cx^2)}{2c}$$

input `Int[x*Cos[a + b*x - c*x^2], x]`

output `(b*(-((Sqrt[Pi/2]*Cos[a + b^2/(4*c])*FresnelC[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]))/Sqrt[c]) - (Sqrt[Pi/2]*FresnelS[(b - 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*Sin[a + b^2/(4*c)])/Sqrt[c]))/(2*c) - Sin[a + b*x - c*x^2]/(2*c)`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3832 `Int[Sin[(d_.)*((e_.) + (f_.*(x_.))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*((e_.) + (f_.*(x_.))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3929 `Int[Cos[(a_.) + (b_.*(x_) + (c_.*(x_.))^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3943 `Int[Cos[(a_.) + (b_.*(x_) + (c_.*(x_.))^2)*((d_.) + (e_.*(x_))), x_Symbol] :> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c) Int[Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]`

Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.89

method	result
default	$-\frac{\sin(-cx^2+bx+a)}{2c} + \frac{b\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)-\sin\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)\right)}{4c\sqrt{-c}}$
risch	$\frac{b\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\text{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} - \frac{b\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\text{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} - \frac{\sin(-cx^2+bx+a)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x\cos\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}x\sin\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}\left(\cos\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelC}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)-\sin\left(\frac{\frac{b^2}{4}+ac}{c}\right)\text{FresnelS}\left(\frac{\sqrt{2}\left(-cx+\frac{b}{2}\right)}{\sqrt{\pi}\sqrt{-c}}\right)\right)}{4c\sqrt{-c}}$

input `int(x*cos(-c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$-\frac{1}{2}\sin(-cx^2+b*x+a)/c + \frac{1}{4}b/c*2^{(1/2)}*\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(\cos((1/4*b^2+a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(-c*x+1/2*b)) - \sin((1/4*b^2+a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(-c*x+1/2*b)))$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.01

$$\int x \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi b \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) + 2c \sin(cx^2 - bx - a)}{4c^2}$$

input `integrate(x*cos(-c*x^2+b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{4} \sqrt{2} \pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{1}{4}(b^2 + 4ac)/c\right) \text{fresnel_cos}\left(\frac{1}{2}\sqrt{2}(2cx - b)\sqrt{\frac{c}{\pi}}/c\right) + \sqrt{2} \pi b \sqrt{\frac{c}{\pi}} \text{fresnel_sin}\left(\frac{1}{2}\sqrt{2}(2cx - b)\sqrt{\frac{c}{\pi}}/c\right) + 2c \sin(cx^2 - bx - a)/c^2$$

Sympy [F]

$$\int x \cos(a + bx - cx^2) dx = \int x \cos(a + bx - cx^2) dx$$

input `integrate(x*cos(-c*x**2+b*x+a),x)`

output `Integral(x*cos(a + b*x - c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 580, normalized size of antiderivative = 4.68

$$\int x \cos(a + bx - cx^2) dx = \text{Too large to display}$$

input `integrate(x*cos(-c*x^2+b*x+a),x, algorithm="maxima")`

output

$$\begin{aligned}
 & -\frac{1}{16}((-I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1) + (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{-(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1)) * b^2 * \cos(\frac{1}{4}(b^2 + 4*a*c)/c) + ((I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1) - (I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{-(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1)) * b^2 * \sin(\frac{1}{4}(b^2 + 4*a*c)/c) - 2 * ((-(I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{-(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1) + (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{-(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1)) * b*c * \cos(\frac{1}{4}(b^2 + 4*a*c)/c) + ((I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1) - (I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\frac{1}{2}\sqrt{-(4Ic^2x^2 - 4Ib*c*x + I*b^2)/c}) - 1)) * b*c * \sin(\frac{1}{4}(b^2 + 4*a*c)/c) * x - 4 * (c * (-I * e^{(1/4 * (4Ic^2x^2 - 4Ib*c*x + I*b^2)/c)}) * \cos(\frac{1}{4}(b^2 + 4*a*c)/c) - c * (e^{(1/4 * (4Ic^2x^2 - 4Ib*c*x + I*b^2)/c)}) * \sin(\frac{1}{4}(b^2 + 4*a*c)/c)) * \sqrt{(4c^2x^2 - 4b*c*x + b^2)/c}) / (c^2 * \sqrt{(4c^2x^2 - 4b*c*x + b^2)/c})
 \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.48

$$\begin{aligned}
 & \int x \cos(a + bx - cx^2) dx \\
 &= -\frac{\frac{\sqrt{2}\sqrt{\pi}b\operatorname{erf}(-\frac{1}{4}\sqrt{2}(2x-\frac{b}{c})(-\frac{ic}{|c|}+1)\sqrt{|c|})e^{(-\frac{ib^2+4iac}{4c})}}{(-\frac{ic}{|c|}+1)\sqrt{|c|}} + 2i e^{(icx^2-ibx-ia)}}{8c} \\
 & - \frac{\frac{\sqrt{2}\sqrt{\pi}b\operatorname{erf}(-\frac{1}{4}\sqrt{2}(2x-\frac{b}{c})(\frac{ic}{|c|}+1)\sqrt{|c|})e^{(-\frac{-ib^2-4iac}{4c})}}{(\frac{ic}{|c|}+1)\sqrt{|c|}} - 2i e^{(-icx^2+ibx+ia)}}{8c}
 \end{aligned}$$

input `integrate(x*cos(-c*x^2+b*x+a),x, algorithm="giac")`

output

```
-1/8*(sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) + 2*I*e^(I*c*x^2 - I*b*x - I*a)/c - 1/8*(sqrt(2)*sqrt(pi)*b*erf(-1/4*sqrt(2)*(2*x - b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/(I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*e^(-I*c*x^2 + I*b*x + I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \cos(a + bx - cx^2) dx = \int x \cos(-cx^2 + bx + a) dx$$

input

```
int(x*cos(a + b*x - c*x^2),x)
```

output

```
int(x*cos(a + b*x - c*x^2), x)
```

Reduce [F]

$$\int x \cos(a + bx - cx^2) dx = \frac{(\int \cos(-cx^2 + bx + a) dx) b - \sin(-cx^2 + bx + a)}{2c}$$

input

```
int(x*cos(-c*x^2+b*x+a),x)
```

output

```
(int(cos(a + b*x - c*x**2),x)*b - sin(a + b*x - c*x**2))/(2*c)
```

3.8 $\int \cos(a + bx - cx^2) dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	91
Fricas [A] (verification not implemented)	92
Sympy [A] (verification not implemented)	92
Maxima [C] (verification not implemented)	93
Giac [C] (verification not implemented)	93
Mupad [B] (verification not implemented)	94
Reduce [F]	94

Optimal result

Integrand size = 12, antiderivative size = 99

$$\int \cos(a + bx - cx^2) dx = -\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\ - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{\sqrt{c}}$$

output

```
-1/2*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(1/2)-1/2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(-2*c*x+b)/c^(1/2))*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.19 (sec), antiderivative size = 88, normalized size of antiderivative = 0.89

$$\int \cos(a + bx - cx^2) dx \\ = \frac{\sqrt{\frac{\pi}{2}} \left(\cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \right)}{\sqrt{c}}$$

input `Integrate[Cos[a + b*x - c*x^2], x]`

output
$$\frac{(\text{Sqrt}[\text{Pi}/2] \cdot (\text{Cos}[a + b^2/(4c]) \cdot \text{FresnelC}[(-b + 2cx)/(\text{Sqrt}[c] \cdot \text{Sqrt}[2\pi])] + \text{FresnelS}[(-b + 2cx)/(\text{Sqrt}[c] \cdot \text{Sqrt}[2\pi])] \cdot \text{Sin}[a + b^2/(4c)]))}{\text{Sqrt}[c]}$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3929, 25, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cos(a + bx - cx^2) dx \\
 & \downarrow 3929 \\
 & \cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) dx - \sin\left(a + \frac{b^2}{4c}\right) \int -\sin\left(\frac{(b - 2cx)^2}{4c}\right) dx \\
 & \downarrow 25 \\
 & \sin\left(a + \frac{b^2}{4c}\right) \int \sin\left(\frac{(b - 2cx)^2}{4c}\right) dx + \cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) dx \\
 & \downarrow 3832 \\
 & \cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\
 & \downarrow 3833 \\
 & -\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}
 \end{aligned}$$

input `Int[Cos[a + b*x - c*x^2], x]`

output

$$-\left(\frac{(\text{Sqrt}[\text{Pi}/2] \cdot \text{Cos}[a + b^2/(4c)] \cdot \text{FresnelC}[(b - 2c x)/(Sqrt[c] \cdot Sqrt[2 \cdot \text{Pi}])] \cdot Sqrt[c])}{Sqrt[c]} - (\text{Sqrt}[\text{Pi}/2] \cdot \text{FresnelS}[(b - 2c x)/(Sqrt[c] \cdot Sqrt[2 \cdot \text{Pi}]]) \cdot \text{Sin}[a + b^2/(4c)])\right) / Sqrt[c]$$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_{x_}), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 3832 $\text{Int}[\text{Sin}[(d_.) * ((e_.) + (f_.) * (x_))^2], x_Symbol] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f \cdot \text{Rt}[d, 2])) \cdot \text{FresnelS}[\text{Sqrt}[2/\text{Pi}] \cdot \text{Rt}[d, 2] * (e + f x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\text{Int}[\text{Cos}[(d_.) * ((e_.) + (f_.) * (x_))^2], x_Symbol] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f \cdot \text{Rt}[d, 2])) \cdot \text{FresnelC}[\text{Sqrt}[2/\text{Pi}] \cdot \text{Rt}[d, 2] * (e + f x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3929 $\text{Int}[\text{Cos}[(a_.) + (b_.) * (x_) + (c_.) * (x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4a c)/(4c)] \quad \text{Int}[\text{Cos}[(b + 2c x)^2/(4c)], x], x] + \text{Simp}[\text{Sin}[(b^2 - 4a c)/(4c)] \quad \text{Int}[\text{Sin}[(b + 2c x)^2/(4c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4a c, 0]$

Maple [A] (verified)

Time = 0.70 (sec), antiderivative size = 88, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\sqrt{2} \sqrt{\pi} \left(\cos\left(\frac{b^2+ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2} \left(-cx+\frac{b}{2}\right)}{\sqrt{\pi} \sqrt{-c}}\right) - \sin\left(\frac{b^2+ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2} \left(-cx+\frac{b}{2}\right)}{\sqrt{\pi} \sqrt{-c}}\right) \right)}{2\sqrt{-c}}$	88
risch	$\frac{\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{4c}} \text{erf}\left(\sqrt{-ic} x + \frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac+b^2)}{4c}} \text{erf}\left(-\sqrt{ic} x + \frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}}$	95

input $\text{int}(\cos(-c x^2 + b x + a), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$\frac{1}{2} \cdot 2^{(1/2)} \cdot \text{Pi}^{(1/2)} / (-c)^{(1/2)} \cdot (\cos((1/4 \cdot b^2 + a \cdot c) / c) \cdot \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / (-c)^{(1/2)} \cdot (-c \cdot x + 1/2 \cdot b)) - \sin((1/4 \cdot b^2 + a \cdot c) / c) \cdot \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / (-c)^{(1/2)} \cdot (-c \cdot x + 1/2 \cdot b)))$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 106, normalized size of antiderivative = 1.07

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\pi\sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi\sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right)}{2c}$$

input `integrate(cos(-c*x^2+b*x+a),x, algorithm="fricas")`

output
$$\frac{1}{2} \cdot (\sqrt{2} \cdot \pi \cdot \sqrt{c/\pi}) \cdot \cos(1/4 \cdot (b^2 + 4 \cdot a \cdot c) / c) \cdot \text{fresnel_cos}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x - b) \cdot \sqrt{c/\pi} / c) + \sqrt{2} \cdot \pi \cdot \sqrt{c/\pi} \cdot \text{fresnel_sin}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x - b) \cdot \sqrt{c/\pi} / c) \cdot \sin(1/4 \cdot (b^2 + 4 \cdot a \cdot c) / c) / c$$

Sympy [A] (verification not implemented)

Time = 0.21 (sec), antiderivative size = 94, normalized size of antiderivative = 0.95

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \sqrt{-\frac{1}{c}} \left(-\sin\left(a + \frac{b^2}{4c}\right) S\left(\frac{\sqrt{2}(b-2cx)}{2\sqrt{\pi}\sqrt{-c}}\right) + \cos\left(a + \frac{b^2}{4c}\right) C\left(\frac{\sqrt{2}(b-2cx)}{2\sqrt{\pi}\sqrt{-c}}\right) \right)}{2}$$

input `integrate(cos(-c*x**2+b*x+a),x)`

output
$$\frac{\sqrt{2} \cdot \sqrt{\pi} \cdot \sqrt{-1/c} \cdot (-\sin(a + b^{**2}/(4*c)) \cdot \text{fresnels}(\sqrt{2} \cdot (b - 2 \cdot c \cdot x) / (2 \cdot \sqrt{\pi} \cdot \sqrt{-c})) + \cos(a + b^{**2}/(4*c)) \cdot \text{fresnelc}(\sqrt{2} \cdot (b - 2 \cdot c \cdot x) / (2 \cdot \sqrt{\pi} \cdot \sqrt{-c}))) / 2}{2}$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.13

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(\left(-(i-1) \cos\left(\frac{b^2+4ac}{4c}\right) + (i+1) \sin\left(\frac{b^2+4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx-i b}{2\sqrt{ic}}\right) + \left(-(i+1) \cos\left(\frac{b^2+4ac}{4c}\right) + (i-1) \sin\left(\frac{b^2+4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+i b}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

input `integrate(cos(-c*x^2+b*x+a),x, algorithm="maxima")`

output `1/8*sqrt(2)*sqrt(pi)*((-I - 1)*cos(1/4*(b^2 + 4*a*c)/c) + (I + 1)*sin(1/4*(b^2 + 4*a*c)/c))*erf(1/2*(2*I*c*x - I*b)/sqrt(I*c)) + (-I + 1)*cos(1/4*(b^2 + 4*a*c)/c) + (I - 1)*sin(1/4*(b^2 + 4*a*c)/c))*erf(1/2*(2*I*c*x - I*b)/sqrt(-I*c)))/sqrt(c)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.38

$$\int \cos(a + bx - cx^2) dx = -\frac{\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x - \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{ib^2+4ic}{4c}\right)}}{4\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}} - \frac{\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}\sqrt{2}(2x - \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2-4ic}{4c}\right)}}{4\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}}$$

input `integrate(cos(-c*x^2+b*x+a),x, algorithm="giac")`

output `-1/4*sqrt(2)*sqrt(pi)*erf(-1/4*sqrt(2)*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 1/4*sqrt(2)*sqrt(pi)*erf(-1/4*sqrt(2)*(2*x - b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c)))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.06

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}\left(\frac{b}{2}-cx\right)\sqrt{-\frac{1}{c}}}{\sqrt{\pi}}\right) \cos\left(\frac{b^2+4ac}{4c}\right) \sqrt{-\frac{1}{c}}}{2} - \frac{\sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}\left(\frac{b}{2}-cx\right)\sqrt{-\frac{1}{c}}}{\sqrt{\pi}}\right) \sin\left(\frac{b^2+4ac}{4c}\right) \sqrt{-\frac{1}{c}}}{2}$$

input `int(cos(a + b*x - c*x^2),x)`

output `(2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*(b/2 - c*x)*(-1/c)^(1/2))/pi^(1/2))*cos((4*a*c + b^2)/(4*c))*(-1/c)^(1/2))/2 - (2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*(b/2 - c*x)*(-1/c)^(1/2))/pi^(1/2))*sin((4*a*c + b^2)/(4*c))*(-1/c)^(1/2))/2`

Reduce [F]

$$\int \cos(a + bx - cx^2) dx = \int \cos(-cx^2 + bx + a) dx$$

input `int(cos(-c*x^2+b*x+a),x)`

output `int(cos(a + b*x - c*x**2),x)`

3.9 $\int \frac{\cos(a+bx-cx^2)}{x} dx$

Optimal result	95
Mathematica [N/A]	95
Rubi [N/A]	96
Maple [N/A]	96
Fricas [N/A]	97
Sympy [N/A]	97
Maxima [N/A]	98
Giac [N/A]	98
Mupad [N/A]	98
Reduce [N/A]	99

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \text{Int}\left(\frac{\cos(a + bx - cx^2)}{x}, x\right)$$

output `Defer(Int)(cos(-c*x^2+b*x+a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(a + bx - cx^2)}{x} dx$$

input `Integrate[Cos[a + b*x - c*x^2]/x,x]`

output `Integrate[Cos[a + b*x - c*x^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3951}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + bx - cx^2)}{x} dx$$

↓ 3951

$$\int \frac{\cos(a + bx - cx^2)}{x} dx$$

input `Int[Cos[a + b*x - c*x^2]/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3951 `Int[Cos[(a_) + (b_)*(x_) + (c_)*(x_)^2]^(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] :> Unintegrable[(d + e*x)^m*Cos[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

input `int(cos(-c*x^2+b*x+a)/x, x)`

output `int(cos(-c*x^2+b*x+a)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

input `integrate(cos(-c*x^2+b*x+a)/x,x, algorithm="fricas")`

output `integral(cos(c*x^2 - b*x - a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(a + bx - cx^2)}{x} dx$$

input `integrate(cos(-c*x**2+b*x+a)/x,x)`

output `Integral(cos(a + b*x - c*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

input `integrate(cos(-c*x^2+b*x+a)/x,x, algorithm="maxima")`

output `integrate(cos(c*x^2 - b*x - a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

input `integrate(cos(-c*x^2+b*x+a)/x,x, algorithm="giac")`

output `integrate(cos(-c*x^2 + b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 41.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

input `int(cos(a + b*x - c*x^2)/x,x)`

output `int(cos(a + b*x - c*x^2)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

input `int(cos(-c*x^2+b*x+a)/x,x)`

output `int(cos(a + b*x - c*x**2)/x,x)`

3.10 $\int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx$

Optimal result	100
Mathematica [A] (verified)	101
Rubi [A] (verified)	101
Maple [F]	102
Fricas [A] (verification not implemented)	102
Sympy [F]	103
Maxima [F]	103
Giac [F]	103
Mupad [F(-1)]	104
Reduce [F]	104

Optimal result

Integrand size = 34, antiderivative size = 111

$$\begin{aligned} & \int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx \\ &= -\frac{\cos(a+bx-cx^2)}{x} + \sqrt{c}\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad - \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \end{aligned}$$

output

```
-cos(-c*x^2+b*x+a)/x+c^(1/2)*2^(1/2)*Pi^(1/2)*cos(a+1/4*b^2/c)*FresnelS(1/
2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))-c^(1/2)*2^(1/2)*Pi^(1/2)*FresnelC(1
/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)
```

Mathematica [A] (verified)

Time = 4.28 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.03

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= -\frac{\cos(a + x(b - cx))}{x} - \sqrt{c}\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{-b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &+ \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{-b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \end{aligned}$$

input `Integrate[Cos[a + b*x - c*x^2]/x^2 + (b*Sin[a + b*x - c*x^2])/x, x]`

output `-(Cos[a + x*(b - c*x)]/x) - Sqrt[c]*Sqrt[2*Pi]*Cos[a + b^2/(4*c)]*FresnelS[(-b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] + Sqrt[c]*Sqrt[2*Pi]*FresnelC[(-b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*Sin[a + b^2/(4*c)]`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.029$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \left(\frac{b \sin(a + bx - cx^2)}{x} + \frac{\cos(a + bx - cx^2)}{x^2} \right) dx \\ & \quad \downarrow \text{2009} \\ & -\sqrt{2\pi}\sqrt{c} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b - 2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \sqrt{2\pi}\sqrt{c} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b - 2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \\ & \quad \frac{\cos(a + bx - cx^2)}{x} \end{aligned}$$

input `Int[Cos[a + b*x - c*x^2]/x^2 + (b*Sin[a + b*x - c*x^2])/x, x]`

output

$$-(\cos[a + bx - cx^2]/x) + \sqrt{c} \cdot \sqrt{2\pi} \cdot \cos[a + b^2/(4c)] \cdot \text{FresnelS}[(b - 2cx)/(\sqrt{c} \cdot \sqrt{2\pi})] - \sqrt{c} \cdot \sqrt{2\pi} \cdot \text{FresnelC}[(b - 2cx)/(\sqrt{c} \cdot \sqrt{2\pi})] \cdot \sin[a + b^2/(4c)]$$

Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] :> \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

Maple [F]

$$\int \left(\frac{\cos(-cx^2 + bx + a)}{x^2} + \frac{b \sin(-cx^2 + bx + a)}{x} \right) dx$$

input

$$\text{int}(\cos(-c*x^2+b*x+a)/x^2+b*\sin(-c*x^2+b*x+a)/x, x)$$

output

$$\text{int}(\cos(-c*x^2+b*x+a)/x^2+b*\sin(-c*x^2+b*x+a)/x, x)$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 123, normalized size of antiderivative = 1.11

$$\int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx =$$

$$-\frac{\sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) + \cos(cx^2 - bx)}{x}$$

input

$$\text{integrate}(\cos(-c*x^2+b*x+a)/x^2+b*\sin(-c*x^2+b*x+a)/x, x, \text{algorithm}=\text{"fricas"})$$

output

$$-(\sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) \text{fresnel_sin}\left(\frac{1}{2}\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}\right) - \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \text{fresnel_cos}\left(\frac{1}{2}\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}\right) + \cos(cx^2 - bx - a))/x$$

Sympy [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{bx \sin(a + bx - cx^2) + \cos(a + bx - cx^2)}{x^2} dx \end{aligned}$$

input `integrate(cos(-c*x**2+b*x+a)/x**2+b*sin(-c*x**2+b*x+a)/x,x)`

output `Integral((b*x*sin(a + b*x - c*x**2) + cos(a + b*x - c*x**2))/x**2, x)`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(-cx^2 + bx + a)}{x} + \frac{\cos(-cx^2 + bx + a)}{x^2} dx \end{aligned}$$

input `integrate(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x,x, algorithm="maxima")`

output `integrate(-b*sin(c*x^2 - b*x - a)/x + cos(c*x^2 - b*x - a)/x^2, x)`

Giac [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(-cx^2 + bx + a)}{x} + \frac{\cos(-cx^2 + bx + a)}{x^2} dx \end{aligned}$$

input `integrate(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x,x, algorithm="giac")`

output `integrate(b*sin(-c*x^2 + b*x + a)/x + cos(-c*x^2 + b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{\cos(-cx^2 + bx + a)}{x^2} + \frac{b \sin(-cx^2 + bx + a)}{x} dx \end{aligned}$$

input `int(cos(a + b*x - c*x^2)/x^2 + (b*sin(a + b*x - c*x^2))/x, x)`

output `int(cos(a + b*x - c*x^2)/x^2 + (b*sin(a + b*x - c*x^2))/x, x)`

Reduce [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \frac{\left(\int \frac{\cos(-cx^2 + bx + a)}{x^2} dx \right) x + \left(\int \frac{\sin(-cx^2 + bx + a)}{x} dx \right) bx + \left(\int \frac{1}{x^2} dx \right) x + 1}{x} \end{aligned}$$

input `int(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x, x)`

output `(int(cos(a + b*x - c*x**2)/x**2, x)*x + int(sin(a + b*x - c*x**2)/x, x)*b*x + int(1/x**2, x)*x + 1)/x`

3.11 $\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	105
Mathematica [A] (verified)	105
Rubi [A] (verified)	106
Maple [A] (verified)	108
Fricas [A] (verification not implemented)	109
Sympy [F]	109
Maxima [C] (verification not implemented)	109
Giac [C] (verification not implemented)	110
Mupad [B] (verification not implemented)	111
Reduce [F]	111

Optimal result

Integrand size = 13, antiderivative size = 82

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right) \\ - \frac{1}{4} \sin \left(\frac{1}{4} + x + x^2 \right) + \frac{1}{2} x \sin \left(\frac{1}{4} + x + x^2 \right)$$

output $1/8*2^{(1/2)}*\text{Pi}^{(1/2)}*\text{FresnelC}(1/2*(1+2*x)*2^{(1/2)}/\text{Pi}^{(1/2)})-1/4*2^{(1/2)}*\text{Pi}^{(1/2)}*\text{FresnelS}(1/2*(1+2*x)*2^{(1/2)}/\text{Pi}^{(1/2)})-1/4*\sin(1/4+x+x^2)+1/2*x*\sin(1/4+x+x^2)$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.82

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{8} \left(\sqrt{2\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) - 2 \left(\sqrt{2\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right) + (1-2x) \sin \left(\frac{1}{4} + x + x^2 \right) \right) \right)$$

input `Integrate[x^2*Cos[1/4 + x + x^2], x]`

output
$$\frac{(\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[(1 + 2*x)/\text{Sqrt}[2*\text{Pi}]] - 2*(\text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[(1 + 2*x)/\text{Sqrt}[2*\text{Pi}]] + (1 - 2*x)*\text{Sin}[1/4 + x + x^2]))}{8}$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3945, 3926, 3832, 3943, 3927, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \cos \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \textcolor{blue}{3945} \\
 & -\frac{1}{2} \int \sin \left(x^2 + x + \frac{1}{4} \right) dx - \frac{1}{2} \int x \cos \left(x^2 + x + \frac{1}{4} \right) dx + \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3926} \\
 & -\frac{1}{2} \int x \cos \left(x^2 + x + \frac{1}{4} \right) dx - \frac{1}{2} \int \sin \left(\frac{1}{4}(2x+1)^2 \right) dx + \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3832} \\
 & -\frac{1}{2} \int x \cos \left(x^2 + x + \frac{1}{4} \right) dx - \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) + \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3943} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \cos \left(x^2 + x + \frac{1}{4} \right) dx - \frac{1}{2} \sin \left(x^2 + x + \frac{1}{4} \right) \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) + \\
 & \quad \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3927} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \cos \left(\frac{1}{4}(2x+1)^2 \right) dx - \frac{1}{2} \sin \left(x^2 + x + \frac{1}{4} \right) \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) + \\
 & \quad \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right) \\
 & \quad \downarrow \textcolor{blue}{3833}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right) - \frac{1}{2} \sin \left(x^2 + x + \frac{1}{4} \right) \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) + \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right)$$

input `Int[x^2*Cos[1/4 + x + x^2], x]`

output `-1/2*(Sqrt[Pi/2]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]]) + ((Sqrt[Pi/2]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]])/2 - Sin[1/4 + x + x^2]/2)/2 + (x*Sin[1/4 + x + x^2])/2`

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3926 `Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Int[Sin[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

rule 3927 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

rule 3943 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c) Int[Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]`

rule 3945

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.)*(x_))^(m_), x_Sym
bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Simp[
(b*e - 2*c*d)/(2*c) Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] -
Simp[e^2*((m - 1)/(2*c)) Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x],
x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Maple [A] (verified)

Time = 1.64 (sec), antiderivative size = 59, normalized size of antiderivative = 0.72

method	result
default	$\frac{x \sin(\frac{1}{4}+x+x^2)}{2} - \frac{\sin(\frac{1}{4}+x+x^2)}{4} + \frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{8} - \frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{4}$
risch	$-\frac{\sqrt{\pi} (-1)^{\frac{3}{4}} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{16} - \frac{(-1)^{\frac{1}{4}} \sqrt{\pi} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{8} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{16 \sqrt{-i}} + \frac{i \sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{8 \sqrt{-i}}$
parts	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right) x^2}{2} - \frac{\sqrt{2} \pi^{\frac{3}{2}} \left(-\frac{\operatorname{FresnelC}\left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right) \left(-\left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right)^2 \sqrt{\pi}+\sqrt{2} \left(\frac{\sqrt{2} x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2 \sqrt{\pi}}\right)\right)}{\sqrt{\pi}} + \right)}{4}$

```
input int(x^2*cos(1/4+x+x^2),x,method=_RETURNVERBOSE)
```

output

```
1/2*x*sin(1/4+x+x^2)-1/4*sin(1/4+x+x^2)+1/8*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))-1/4*2^(1/2)*Pi^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)*(x+1/2))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.72

$$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{8} \sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right) - \frac{1}{4} \sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right) + \frac{1}{4} (2x-1) \sin\left(x^2 + x + \frac{1}{4}\right)$$

input `integrate(x^2*cos(1/4+x+x^2),x, algorithm="fricas")`

output `1/8*sqrt(2)*sqrt(pi)*fresnel_cos(1/2*sqrt(2)*(2*x + 1)/sqrt(pi)) - 1/4*sqr
t(2)*sqrt(pi)*fresnel_sin(1/2*sqrt(2)*(2*x + 1)/sqrt(pi)) + 1/4*(2*x - 1)*
sin(x^2 + x + 1/4)`

Sympy [F]

$$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = \int x^2 \cos\left(x^2 + x + \frac{1}{4}\right) dx$$

input `integrate(x**2*cos(1/4+x+x**2),x)`

output `Integral(x**2*cos(x**2 + x + 1/4), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.95

$$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{-16x \left(-ie^{(ix^2+ix+\frac{1}{4})} + ie^{(-ix^2-ix-\frac{1}{4})}\right) + \sqrt{4x^2+4x+1} \left((i-1)\sqrt{2}\sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{ix^2+ix+\frac{1}{4}}\right) - \operatorname{erf}\left(\sqrt{ix^2+ix+\frac{1}{4}}\right)\right)\right)}{16}$$

input `integrate(x^2*cos(1/4+x+x^2),x, algorithm="maxima")`

output
$$\begin{aligned} & -\frac{1}{32} \left(16x(-I e^{(Ix^2 + Ix + 1/4)I} + e^{(-Ix^2 - Ix - 1/4)I}) + \sqrt{4x^2 + 4x + 1}((I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{Ix^2 + Ix + 1/4}I) - 1) - (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{-Ix^2 - Ix - 1/4}I) - 1) \right. \\ & \left. - (4I + 4)\sqrt{2}\gamma(3/2, Ix^2 + Ix + 1/4)I + (4I - 4)\sqrt{2}\gamma(3/2, -Ix^2 - Ix - 1/4)I) - 8Ie^{(Ix^2 + Ix + 1/4)I} + 8Ie^{(-Ix^2 - Ix - 1/4)I} \right) / (2x + 1) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.91

$$\begin{aligned} \int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = & -\left(\frac{3}{32}i - \frac{1}{32}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2x + 1)\right) \\ & + \left(\frac{3}{32}i + \frac{1}{32}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2}(2x + 1)\right) \\ & + \frac{1}{8} (-2ix + i)e^{(ix^2 + ix + \frac{1}{4}i)} + \frac{1}{8} (2ix - i)e^{(-ix^2 - ix - \frac{1}{4}i)} \end{aligned}$$

input `integrate(x^2*cos(1/4+x+x^2),x, algorithm="giac")`

output
$$\begin{aligned} & -(3/32*I - 1/32)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}((1/4*I - 1/4)*\sqrt{2}*(2*x + 1)) + (3/32*I + 1/32)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-(1/4*I + 1/4)*\sqrt{2}*(2*x + 1)) + 1/8*(-2*I*x + I)*e^{(Ix^2 + Ix + 1/4)I} + 1/8*(2*I*x - I)*e^{(-Ix^2 - Ix - 1/4)I} \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 41.31 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.78

$$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{x \sin(x^2 + x + \frac{1}{4})}{2} - \frac{\sin(x^2 + x + \frac{1}{4})}{4} \\ + \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{8} - \frac{\sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{4}$$

input `int(x^2*cos(x + x^2 + 1/4),x)`

output `(x*sin(x + x^2 + 1/4))/2 - sin(x + x^2 + 1/4)/4 + (2^(1/2)*pi^(1/2)*fresne1c((2^(1/2)*(2*x + 1))/(2*pi^(1/2))))/8 - (2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*(2*x + 1))/(2*pi^(1/2))))/4`

Reduce [F]

$$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = 2 \left(\int \frac{x^2}{\tan\left(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8}\right)^2 + 1} dx \right) - \frac{x^3}{3}$$

input `int(x^2*cos(1/4+x+x^2),x)`

output `(6*int(x**2/(tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) - x**3)/3`

3.12 $\int x \cos \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	112
Mathematica [A] (verified)	112
Rubi [A] (verified)	113
Maple [A] (verified)	114
Fricas [A] (verification not implemented)	114
Sympy [B] (verification not implemented)	115
Maxima [C] (verification not implemented)	115
Giac [C] (verification not implemented)	116
Mupad [B] (verification not implemented)	116
Reduce [F]	117

Optimal result

Integrand size = 11, antiderivative size = 41

$$\int x \cos \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) + \frac{1}{2} \sin \left(\frac{1}{4} + x + x^2 \right)$$

output
$$-\frac{1}{4} \cdot 2^{(1/2)} \cdot \operatorname{Pi}^{(1/2)} \cdot \operatorname{FresnelC} \left(\frac{1}{2} \cdot (1+2x) \cdot 2^{(1/2)} / \operatorname{Pi}^{(1/2)} \right) + \frac{1}{2} \sin(1/4+x+x^2)$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int x \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \left(-\sqrt{2\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) + 2 \sin \left(\frac{1}{4} + x + x^2 \right) \right)$$

input `Integrate[x*Cos[1/4 + x + x^2], x]`

output
$$(-\operatorname{Sqrt}[2 \cdot \operatorname{Pi}] \cdot \operatorname{FresnelC}[(1+2x)/\operatorname{Sqrt}[2 \cdot \operatorname{Pi}]] + 2 \cdot \operatorname{Sin}[1/4+x+x^2])/4$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3943, 3927, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \text{3943} \\
 & \frac{1}{2} \sin \left(x^2 + x + \frac{1}{4} \right) - \frac{1}{2} \int \cos \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \text{3927} \\
 & \frac{1}{2} \sin \left(x^2 + x + \frac{1}{4} \right) - \frac{1}{2} \int \cos \left(\frac{1}{4}(2x+1)^2 \right) dx \\
 & \quad \downarrow \text{3833} \\
 & \frac{1}{2} \sin \left(x^2 + x + \frac{1}{4} \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right)
 \end{aligned}$$

input `Int[x*Cos[1/4 + x + x^2], x]`

output `-1/2*(Sqrt[Pi/2]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]]) + Sin[1/4 + x + x^2]/2`

Definitions of rubi rules used

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3927 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_.)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

rule 3943

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c) Int
[Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b
*e, 0]
```

Maple [A] (verified)

Time = 1.34 (sec), antiderivative size = 30, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{\sin\left(\frac{1}{4}+x+x^2\right)}{2}-\frac{\sqrt{2}\sqrt{\pi}\operatorname{FresnelC}\left(\frac{\sqrt{2}\left(x+\frac{1}{2}\right)}{\sqrt{\pi}}\right)}{4}$	30
risch	$\frac{\sqrt{\pi}(-1)^{\frac{3}{4}}\operatorname{erf}\left((-1)^{\frac{1}{4}}x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{8}-\frac{\sqrt{\pi}\operatorname{erf}\left(\sqrt{-i}x-\frac{i}{2\sqrt{-i}}\right)}{8\sqrt{-i}}+\frac{\sin\left(\frac{(1+2x)^2}{4}\right)}{2}$	58
parts	$\frac{\sqrt{2}\sqrt{\pi}\operatorname{FresnelC}\left(\frac{\sqrt{2}\left(x+\frac{1}{2}\right)}{\sqrt{\pi}}\right)x}{2}-\frac{\pi\left(\operatorname{FresnelC}\left(\frac{\sqrt{2}x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2\sqrt{\pi}}\right)\left(\frac{\sqrt{2}x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2\sqrt{\pi}}\right)-\frac{\sin\left(\frac{\pi\left(\frac{\sqrt{2}x}{\sqrt{\pi}}+\frac{\sqrt{2}}{2\sqrt{\pi}}\right)^2}{2}\right)}{\pi}\right)}{2}$	90

input `int(x*cos(1/4+x+x^2), x, method=_RETURNVERBOSE)`output `1/2*sin(1/4+x+x^2)-1/4*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec), antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{4} \sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right) + \frac{1}{2} \sin\left(x^2 + x + \frac{1}{4}\right)$$

input `integrate(x*cos(1/4+x+x^2), x, algorithm="fricas")`

output
$$-1/4\sqrt{2}\sqrt{\pi}\text{fresnel_cos}(1/2\sqrt{2}(2x+1)/\sqrt{\pi}) + 1/2\sin(x^2+x+1/4)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(39) = 78$.

Time = 0.57 (sec), antiderivative size = 155, normalized size of antiderivative = 3.78

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = -\frac{\sqrt{2}\sqrt{\pi}xC\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)\Gamma\left(\frac{1}{4}\right)}{8\Gamma\left(\frac{5}{4}\right)} + \frac{\sqrt{2}\sqrt{\pi}xC\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)}{2} \\ + \frac{\sin\left((x + \frac{1}{2})^2\right)\Gamma\left(\frac{1}{4}\right)}{8\Gamma\left(\frac{5}{4}\right)} - \frac{\sqrt{2}\sqrt{\pi}C\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)\Gamma\left(\frac{1}{4}\right)}{16\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(x*cos(1/4+x+x**2),x)`

output
$$-\sqrt{2}\sqrt{\pi}x\text{fresnelc}(\sqrt{2}x/\sqrt{\pi} + \sqrt{2}/(2\sqrt{\pi}))\text{gamma}(1/4)/(8\text{gamma}(5/4)) + \sqrt{2}\sqrt{\pi}x\text{fresnelc}(\sqrt{2}x/\sqrt{\pi} + \sqrt{2}/(2\sqrt{\pi}))/2 + \sin((x + 1/2)^2)\text{gamma}(1/4)/(8\text{gamma}(5/4)) - \sqrt{2}\sqrt{\pi}\text{fresnelc}(\sqrt{2}x/\sqrt{\pi} + \sqrt{2}/(2\sqrt{\pi}))\text{gamma}(1/4)/(16\text{gamma}(5/4))$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.18 (sec), antiderivative size = 126, normalized size of antiderivative = 3.07

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx \\ = \frac{8x\left(-i e^{(ix^2+ix+\frac{1}{4}i)} + i e^{(-ix^2-ix-\frac{1}{4}i)}\right) + \sqrt{4x^2+4x+1}\left((i-1)\sqrt{2}\sqrt{\pi}\left(\text{erf}\left(\sqrt{ix^2+ix+\frac{1}{4}i}\right) - 1\right)\right)}{16(2x+1)}$$

input `integrate(x*cos(1/4+x+x^2),x, algorithm="maxima")`

output

$$\frac{1}{16} \left(8x(-I e^{(Ix^2 + Ix + 1/4)I} + e^{(-Ix^2 - Ix - 1/4)I}) + \sqrt{(4x^2 + 4x + 1)((I - 1)\sqrt{2}\sqrt{\pi})} \operatorname{erf}(\sqrt{Ix^2 + Ix + 1/4}I) - 1) - (I + 1)\sqrt{2}\sqrt{\pi} \operatorname{erf}(\sqrt{-Ix^2 - Ix - 1/4}I) - 1) - 4I e^{(Ix^2 + Ix + 1/4)I} + 4I e^{(-Ix^2 - Ix - 1/4)I} \right) / (2x + 1)$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\begin{aligned} \int x \cos\left(\frac{1}{4} + x + x^2\right) dx &= \left(\frac{1}{16}i + \frac{1}{16}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2x + 1)\right) \\ &\quad - \left(\frac{1}{16}i - \frac{1}{16}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2}(2x + 1)\right) \\ &\quad - \frac{1}{4}i e^{(ix^2 + ix + \frac{1}{4}i)} + \frac{1}{4}i e^{(-ix^2 - ix - \frac{1}{4}i)} \end{aligned}$$

input

```
integrate(x*cos(1/4+x+x^2),x, algorithm="giac")
```

output

$$\begin{aligned} &(1/16*I + 1/16)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}((1/4*I - 1/4)*\sqrt{2}*(2*x + 1)) - (1/16*I - 1/16)*\sqrt{2}*\sqrt{\pi}*\operatorname{erf}(-(1/4*I + 1/4)*\sqrt{2}*(2*x + 1)) - 1/4*I e^{(Ix^2 + Ix + 1/4)I} + 1/4*I e^{(-Ix^2 - Ix - 1/4)I} \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sin(x^2 + x + \frac{1}{4})}{2} - \frac{\sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{4}$$

input

```
int(x*cos(x + x^2 + 1/4),x)
```

output

$$\sin(x + x^2 + 1/4)/2 - (2^{(1/2)}\pi^{(1/2)}\operatorname{fresnelc}((2^{(1/2)}(2*x + 1))/(2\pi^{(1/2)}))/4$$

Reduce [F]

$$\int x \cos \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{\left(\int \cos \left(x^2 + x + \frac{1}{4} \right) dx \right)}{2} + \frac{\sin \left(x^2 + x + \frac{1}{4} \right)}{2}$$

input `int(x*cos(1/4+x+x^2),x)`

output `(- int(cos((4*x**2 + 4*x + 1)/4),x) + sin((4*x**2 + 4*x + 1)/4))/2`

3.13 $\int \cos\left(\frac{1}{4} + x + x^2\right) dx$

Optimal result	118
Mathematica [A] (verified)	118
Rubi [A] (verified)	119
Maple [A] (verified)	120
Fricas [A] (verification not implemented)	120
Sympy [A] (verification not implemented)	120
Maxima [C] (verification not implemented)	121
Giac [C] (verification not implemented)	121
Mupad [B] (verification not implemented)	122
Reduce [F]	122

Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right)$$

output `1/2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(1+2*x)*2^(1/2)/Pi^(1/2))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right)$$

input `Integrate[Cos[1/4 + x + x^2], x]`

output `Sqrt[Pi/2]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3927, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos \left(x^2 + x + \frac{1}{4} \right) dx \\ & \quad \downarrow \textcolor{blue}{3927} \\ & \int \cos \left(\frac{1}{4}(2x+1)^2 \right) dx \\ & \quad \downarrow \textcolor{blue}{3833} \\ & \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right) \end{aligned}$$

input `Int[Cos[1/4 + x + x^2], x]`

output `Sqrt[Pi/2]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]]`

Definitions of rubi rules used

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3927 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{2} \sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2} (x+\frac{1}{2})}{\sqrt{\pi}}\right)}{2}$	20
risch	$-\frac{\sqrt{\pi} (-1)^{\frac{3}{4}} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{4} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{4 \sqrt{-i}}$	46

input `int(cos(1/4+x+x^2),x,method=_RETURNVERBOSE)`

output `1/2*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{2} \sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)$$

input `integrate(cos(1/4+x+x^2),x, algorithm="fricas")`

output `1/2*sqrt(2)*sqrt(pi)*fresnel_cos(1/2*sqrt(2)*(2*x + 1)/sqrt(pi))`

Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{2}$$

input `integrate(cos(1/4+x+x**2),x)`

output $\sqrt{2}\sqrt{\pi} \operatorname{fresnelc}(\sqrt{2}(2x+1)/(2\sqrt{\pi}))/2$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.92

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{16}\sqrt{\pi}\left((i-1)\sqrt{2}\operatorname{erf}\left(-\frac{1}{2}(-1)^{\frac{3}{4}}(2ix+i)\right) + (i-1)\sqrt{2}\operatorname{erf}\left(-\left(\frac{1}{4}i-\frac{1}{4}\right)\sqrt{2}(2ix+i)\right) - (i+1)\sqrt{2}\operatorname{erf}\left(-\left(\frac{1}{4}i+\frac{1}{4}\right)\sqrt{2}(2ix+i)\right)\right)$$

input `integrate(cos(1/4+x+x^2),x, algorithm="maxima")`

output $-1/16\sqrt{\pi}((I-1)\sqrt{2}\operatorname{erf}(-1/2(-1)^{(3/4)}(2Ix+I)) + (I-1)\sqrt{2}\operatorname{erf}(-(1/4I-1/4)\sqrt{2}(2Ix+I)) - (I+1)\sqrt{2}\operatorname{erf}(-(1/4I+1/4)\sqrt{2}(2Ix+I)) + (I+1)\sqrt{2}\operatorname{erf}(1/2(2Ix+I)/\sqrt{-I}))$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.62

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = -\left(\frac{1}{8}i + \frac{1}{8}\right)\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right)\sqrt{2}(2x+1)\right) + \left(\frac{1}{8}i - \frac{1}{8}\right)\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right)\sqrt{2}(2x+1)\right)$$

input `integrate(cos(1/4+x+x^2),x, algorithm="giac")`

output $-(1/8I + 1/8)\sqrt{2}\sqrt{\pi}\operatorname{erf}((1/4I - 1/4)\sqrt{2}(2x+1)) + (1/8I - 1/8)\sqrt{2}\sqrt{\pi}\operatorname{erf}(-(1/4I + 1/4)\sqrt{2}(2x+1))$

Mupad [B] (verification not implemented)

Time = 41.35 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{2}$$

input `int(cos(x + x^2 + 1/4),x)`

output `(2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*(x + 1/2))/pi^(1/2)))/2`

Reduce [F]

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \int \cos\left(x^2 + x + \frac{1}{4}\right) dx$$

input `int(cos(1/4+x+x^2),x)`

output `int(cos((4*x**2 + 4*x + 1)/4),x)`

3.14 $\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x} dx$

Optimal result	123
Mathematica [N/A]	123
Rubi [N/A]	124
Maple [N/A]	124
Fricas [N/A]	125
Sympy [N/A]	125
Maxima [N/A]	126
Giac [N/A]	126
Mupad [N/A]	126
Reduce [N/A]	127

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \text{Int}\left(\frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x}, x\right)$$

output `Defer(Int)(cos(1/4+x+x^2)/x,x)`

Mathematica [N/A]

Not integrable

Time = 13.43 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

input `Integrate[Cos[1/4 + x + x^2]/x,x]`

output `Integrate[Cos[1/4 + x + x^2]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3951}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\cos(x^2 + x + \frac{1}{4})}{x} dx \\ \downarrow \text{3951} \\ \int \frac{\cos(x^2 + x + \frac{1}{4})}{x} dx \end{array}$$

input `Int[Cos[1/4 + x + x^2]/x, x]`

output `$Aborted`

Defintions of rubi rules used

rule 3951 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.*(x_)^2)^(n_.)*((d_.) + (e_.*(x_))^(m_.)), x_Symbol] :> Unintegrable[(d + e*x)^m*Cos[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{\cos(\frac{1}{4} + x + x^2)}{x} dx$$

input `int(cos(1/4+x+x^2)/x, x)`

output `int(cos(1/4+x+x^2)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(cos(1/4+x+x^2)/x,x, algorithm="fricas")`

output `integral(cos(x^2 + x + 1/4)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(cos(1/4+x+x**2)/x,x)`

output `Integral(cos(x**2 + x + 1/4)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(cos(1/4+x+x^2)/x,x, algorithm="maxima")`

output `integrate(cos(x^2 + x + 1/4)/x, x)`

Giac [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(cos(1/4+x+x^2)/x,x, algorithm="giac")`

output `integrate(cos(x^2 + x + 1/4)/x, x)`

Mupad [N/A]

Not integrable

Time = 41.31 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `int(cos(x + x^2 + 1/4)/x,x)`

output `int(cos(x + x^2 + 1/4)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `int(cos(1/4+x+x^2)/x,x)`

output `int(cos((4*x**2 + 4*x + 1)/4)/x,x)`

3.15 $\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$

Optimal result	128
Mathematica [N/A]	128
Rubi [N/A]	129
Maple [N/A]	130
Fricas [N/A]	131
Sympy [N/A]	131
Maxima [N/A]	131
Giac [N/A]	132
Mupad [N/A]	132
Reduce [N/A]	133

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = -\frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} - \sqrt{2\pi} \text{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right) \\ - \text{Int}\left(\frac{\sin\left(\frac{1}{4} + x + x^2\right)}{x}, x\right)$$

output $-\cos(1/4+x+x^2)/x-2^{(1/2)}*\text{Pi}^{(1/2)}*\text{FresnelS}(1/2*(1+2*x)*2^{(1/2)}/\text{Pi}^{(1/2)})-$
 $\text{Defer}(\text{Int})(\sin(1/4+x+x^2)/x,x)$

Mathematica [N/A]

Not integrable

Time = 11.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

input `Integrate[Cos[1/4 + x + x^2]/x^2, x]`

output `Integrate[Cos[1/4 + x + x^2]/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3947, 3926, 3832, 3950}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(x^2 + x + \frac{1}{4})}{x^2} dx \\
 & \quad \downarrow \textcolor{blue}{3947} \\
 & -2 \int \sin\left(x^2 + x + \frac{1}{4}\right) dx - \int \frac{\sin(x^2 + x + \frac{1}{4})}{x} dx - \frac{\cos(x^2 + x + \frac{1}{4})}{x} \\
 & \quad \downarrow \textcolor{blue}{3926} \\
 & - \int \frac{\sin(x^2 + x + \frac{1}{4})}{x} dx - 2 \int \sin\left(\frac{1}{4}(2x+1)^2\right) dx - \frac{\cos(x^2 + x + \frac{1}{4})}{x} \\
 & \quad \downarrow \textcolor{blue}{3832} \\
 & - \int \frac{\sin(x^2 + x + \frac{1}{4})}{x} dx - \sqrt{2\pi} \text{FresnelS}\left(\frac{2x+1}{\sqrt{2\pi}}\right) - \frac{\cos(x^2 + x + \frac{1}{4})}{x} \\
 & \quad \downarrow \textcolor{blue}{3950} \\
 & - \int \frac{\sin(x^2 + x + \frac{1}{4})}{x} dx - \sqrt{2\pi} \text{FresnelS}\left(\frac{2x+1}{\sqrt{2\pi}}\right) - \frac{\cos(x^2 + x + \frac{1}{4})}{x}
 \end{aligned}$$

input `Int[Cos[1/4 + x + x^2]/x^2,x]`

output `$Aborted`

Definitions of rubi rules used

rule 3832 $\text{Int}[\text{Sin}[(d_{_})*((e_{_}) + (f_{_})*(x_{_}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3926 $\text{Int}[\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2], x_{\text{Symbol}}] \rightarrow \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{EqQ}[b^2 - 4*a*c, 0]$

rule 3947 $\text{Int}[\text{Cos}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2]*((d_{_}) + (e_{_})*(x_{_}))^{(m_{_})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(d + e*x)^(m + 1)*(\text{Cos}[a + b*x + c*x^2]/(e*(m + 1))), x] + (\text{Simp}[(b*e - 2*c*d)/(e^2*(m + 1)) \cdot \text{Int}[(d + e*x)^(m + 1)*\text{Sin}[a + b*x + c*x^2], x] + \text{Simp}[2*(c/(e^2*(m + 1))) \cdot \text{Int}[(d + e*x)^(m + 2)*\text{Sin}[a + b*x + c*x^2], x]) /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[b*e - 2*c*d, 0] \&& \text{LtQ}[m, -1]$

rule 3950 $\text{Int}[((d_{_}) + (e_{_})*(x_{_}))^{(m_{_})}*\text{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2]^{(n_{_})}, x_{\text{Symbol}}] \rightarrow \text{Unintegrable}[(d + e*x)^m*\text{Sin}[a + b*x + c*x^2]^n, x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x]$

Maple [N/A]

Not integrable

Time = 0.17 (sec), antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

input `int(cos(1/4+x+x^2)/x^2,x)`

output `int(cos(1/4+x+x^2)/x^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(cos(1/4+x+x^2)/x^2,x, algorithm="fricas")`

output `integral(cos(x^2 + x + 1/4)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(cos(1/4+x+x**2)/x**2,x)`

output `Integral(cos(x**2 + x + 1/4)/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(cos(1/4+x+x^2)/x^2,x, algorithm="maxima")`

output `integrate(cos(x^2 + x + 1/4)/x^2, x)`

Giac [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(cos(1/4+x+x^2)/x^2,x, algorithm="giac")`

output `integrate(cos(x^2 + x + 1/4)/x^2, x)`

Mupad [N/A]

Not integrable

Time = 41.40 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `int(cos(x + x^2 + 1/4)/x^2,x)`

output `int(cos(x + x^2 + 1/4)/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.15

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \frac{\left(\int \frac{\cos(x^2+x+\frac{1}{4})}{x^2} dx\right)x + \left(\int \frac{1}{x^2} dx\right)x + 1}{x}$$

input `int(cos(1/4+x+x^2)/x^2,x)`

output `(int(cos((4*x**2 + 4*x + 1)/4)/x**2,x)*x + int(1/x**2,x)*x + 1)/x`

3.16 $\int x^2 \cos^2(a + bx + cx^2) dx$

Optimal result	134
Mathematica [A] (verified)	135
Rubi [A] (verified)	135
Maple [A] (verified)	136
Fricas [A] (verification not implemented)	137
Sympy [F]	137
Maxima [C] (verification not implemented)	138
Giac [C] (verification not implemented)	139
Mupad [F(-1)]	139
Reduce [F]	140

Optimal result

Integrand size = 17, antiderivative size = 248

$$\begin{aligned} \int x^2 \cos^2(a + bx + cx^2) dx = & \frac{x^3}{6} + \frac{b^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\ & - \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\ & - \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output

```
1/6*x^3+1/16*b^2*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(5/2)-1/16*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)-1/16*Pi^(1/2)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)-1/16*b^2*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(5/2)-1/16*b*sin(2*c*x^2+2*b*x+2*a)/c^2+1/8*x*sin(2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.69

$$\int x^2 \cos^2(a + bx + cx^2) dx \\ = \frac{-3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a - \frac{b^2}{2c}\right) + b^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) + 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a - \frac{b^2}{2c}\right) - c \sin\left(2a - \frac{b^2}{2c}\right)\right)}{48c^{5/2}}$$

input `Integrate[x^2*Cos[a + b*x + c*x^2]^2, x]`

output
$$\frac{(-3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a - \frac{b^2}{2c}\right) + b^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) + 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a - \frac{b^2}{2c}\right) - c \sin\left(2a - \frac{b^2}{2c}\right)\right))}{48c^{5/2}}$$

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos^2(a + bx + cx^2) dx \\ \downarrow 3949 \\ \int \left(\frac{1}{2}x^2 \cos(2a + 2bx + 2cx^2) + \frac{x^2}{2}\right) dx \\ \downarrow 2009$$

$$\begin{aligned}
& - \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} + \frac{\sqrt{\pi} b^2 \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} - \\
& \frac{\sqrt{\pi} b^2 \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \\
& \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} + \frac{x^3}{6}
\end{aligned}$$

input `Int[x^2*Cos[a + b*x + c*x^2]^2, x]`

output $x^{3/6} + (b^{2*}\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a - b^{2/(2*c)}]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])])/(16*c^{(5/2)}) - (\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a - b^{2/(2*c)}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])])/(16*c^{(3/2)}) - (\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*\text{Sin}[2*a - b^{2/(2*c)}])/(16*c^{(3/2)}) - (b^{2*}\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*\text{Sin}[2*a - b^{2/(2*c)}])/(16*c^{(5/2)}) - (b*\text{Sin}[2*a + 2*b*x + 2*c*x^2])/(16*c^2) + (x*\text{Sin}[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.71 (sec), antiderivative size = 191, normalized size of antiderivative = 0.77

method	result
default	$\frac{x^3}{6} + \frac{x \sin(2cx^2 + 2bx + 2a)}{8c} - \frac{b \left(\frac{\sin(2cx^2 + 2bx + 2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^{3/2}} \right)}{4c}$
risch	$\frac{x^3}{6} + \frac{b^2 \sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c^2\sqrt{ic}} - \frac{i\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c\sqrt{ic}} - \frac{b^2 \sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \operatorname{erf}\left(-\sqrt{2}\sqrt{ic}x - \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c^2\sqrt{-2ic}}$

input `int(x^2*cos(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1/6*x^3+1/8*x*sin(2*c*x^2+2*b*x+2*a)/c-1/4*b/c*(1/4*sin(2*c*x^2+2*b*x+2*a)/c-1/4*b/c^{(3/2)}*Pi^{(1/2)}*(cos(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^{(1/2)}/Pi^{(1/2)})+sin(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^{(1/2)}/Pi^{(1/2)}))-1/16/c^{(3/2)}*Pi^{(1/2)}*(cos(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^{(1/2)}/Pi^{(1/2)})-sin(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^{(1/2)}/Pi^{(1/2)}))}{48*c^3}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 178, normalized size of antiderivative = 0.72

$$\int x^2 \cos^2(a + bx + cx^2) dx$$

$$= \frac{8 c^3 x^3 + 6 (2 c^2 x - bc) \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) + 3 \left(\pi b^2 \cos\left(-\frac{b^2 - 4 ac}{2 c}\right) - \pi c \sin\left(-\frac{b^2 - 4 ac}{2 c}\right)\right)}{48 c^3}$$

input `integrate(x^2*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

output
$$\frac{1/48*(8*c^3*x^3 + 6*(2*c^2*x - b*c)*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a) + 3*(pi*b^2*cos(-1/2*(b^2 - 4*a*c)/c) - pi*c*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - 3*(pi*b^2*sin(-1/2*(b^2 - 4*a*c)/c) + pi*c*cos(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c))/c^3}{48*c^3}$$

Sympy [F]

$$\int x^2 \cos^2(a + bx + cx^2) dx = \int x^2 \cos^2(a + bx + cx^2) dx$$

input `integrate(x**2*cos(c*x**2+b*x+a)**2,x)`

output `Integral(x**2*cos(a + b*x + c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.80 (sec) , antiderivative size = 1617, normalized size of antiderivative = 6.52

$$\int x^2 \cos^2(a + bx + cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")`

output

```
-1/384*sqrt(2)*(24*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*cos(-1/2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*sin(-1/2*(b^2 - 4*a*c)/c))*x^3 + 36*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(-1/2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3...
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.85

$$\int x^2 \cos^2(a + bx + cx^2) dx = \frac{1}{6} x^3 - \frac{(c(2ix + \frac{ib}{c}) - 2ib)e^{(2icx^2 + 2ibx + 2ia)} + \frac{\sqrt{\pi}(b^2 + ic)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x + \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 - 4ic}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2} - \frac{(c(-2ix - \frac{ib}{c}) + 2ib)e^{(-2icx^2 - 2ibx - 2ia)} + \frac{\sqrt{\pi}(b^2 - ic)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x + \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 + 4ic}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2}$$

input `integrate(x^2*cos(c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/6*x^3 - 1/32*((c*(2*I*x + I*b/c) - 2*I*b)*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a) + sqrt(pi)*(b^2 + I*c)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1))/c^2 - 1/32*((c*(-2*I*x - I*b/c) + 2*I*b)*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a) + sqrt(pi)*(b^2 - I*c)*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))/c^2`

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2(a + bx + cx^2) dx = \int x^2 \cos(cx^2 + bx + a)^2 dx$$

input `int(x^2*cos(a + b*x + c*x^2)^2,x)`

output `int(x^2*cos(a + b*x + c*x^2)^2, x)`

Reduce [F]

$$\int x^2 \cos^2(a + bx + cx^2) dx$$

$$= -\cos(cx^2 + bx + a) \sin(cx^2 + bx + a) b + 2 \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) cx - 6 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}b)} dx \right)$$

input `int(x^2*cos(c*x^2+b*x+a)^2,x)`

output
$$(-\cos(a + bx + cx^2) \sin(a + bx + cx^2) * b + 2 * \cos(a + bx + cx^2) * \sin(a + bx + cx^2) * c * x - 6 * \int (\tan((a + bx + cx^2)/2))^2 / (\tan((a + bx + cx^2)/2))^4 + 2 * \tan((a + bx + cx^2)/2)^2 + 1, x) * b^2 - 8 * \int (x^2 / (\tan((a + bx + cx^2)/2))^4 + 2 * \tan((a + bx + cx^2)/2)^2 + 1, x) * c^2 - 8 * \int (\tan((a + bx + cx^2)/2) / (\tan((a + bx + cx^2)/2))^4 + 2 * \tan((a + bx + cx^2)/2)^2 + 1, x) * c + 2 * \int (1 / (\tan((a + bx + cx^2)/2))^4 + 2 * \tan((a + bx + cx^2)/2)^2 + 1, x) * b^2 - \sin(a + bx + cx^2) * b + 2 * \sin(a + bx + cx^2) * c * x + 2 * c^2 * x^3) / (6 * c^2)$$

3.17 $\int x \cos^2(a + bx + cx^2) dx$

Optimal result	141
Mathematica [A] (verified)	141
Rubi [A] (verified)	142
Maple [A] (verified)	143
Fricas [A] (verification not implemented)	143
Sympy [F]	144
Maxima [C] (verification not implemented)	144
Giac [C] (verification not implemented)	145
Mupad [F(-1)]	146
Reduce [F]	146

Optimal result

Integrand size = 15, antiderivative size = 126

$$\begin{aligned} \int x \cos^2(a + bx + cx^2) dx &= \frac{x^2}{4} - \frac{b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ &\quad + \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} \\ &\quad + \frac{\sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output
$$1/4*x^2-1/8*b*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)+1/8*b*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)+1/8*sin(2*c*x^2+2*b*x+2*a)/c$$

Mathematica [A] (verified)

Time = 0.28 (sec), antiderivative size = 116, normalized size of antiderivative = 0.92

$$\begin{aligned} \int x \cos^2(a + bx + cx^2) dx \\ = \frac{-b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 + \sin(2(a + x)))}{8c^{3/2}} \end{aligned}$$

input `Integrate[x*Cos[a + b*x + c*x^2]^2,x]`

output
$$\frac{(-b\sqrt{\pi}\cos(2a - b^2/(2c))\text{FresnelC}((b + 2cx)/(\sqrt{c}\sqrt{\pi})) + b\sqrt{\pi}\text{FresnelS}((b + 2cx)/(\sqrt{c}\sqrt{\pi}))\sin(2a - b^2/(2c)) + \sqrt{c}(2cx^2 + \sin(2(a + x(b + cx))))}{8c^{(3/2)}}$$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos^2(a + bx + cx^2) dx \\
 & \quad \downarrow \text{3949} \\
 & \int \left(\frac{1}{2}x \cos(2a + 2bx + 2cx^2) + \frac{x}{2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & - \frac{\sqrt{\pi}b \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} + \frac{\sqrt{\pi}b \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} + \\
 & \quad \frac{\sin(2a + 2bx + 2cx^2)}{8c} + \frac{x^2}{4}
 \end{aligned}$$

input `Int[x*Cos[a + b*x + c*x^2]^2,x]`

output
$$\frac{x^2}{4} - \frac{(b\sqrt{\pi}\cos(2a - b^2/(2c))\text{FresnelC}((b + 2cx)/(\sqrt{c}\sqrt{\pi})) + (b\sqrt{\pi}\text{FresnelS}((b + 2cx)/(\sqrt{c}\sqrt{\pi}))\sin(2a - b^2/(2c)))}{8c^{(3/2)}} + \frac{\sin(2a + 2bx + 2cx^2)}{8c}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3949 $\text{Int}[\cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_))^{m_.}, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \ \cos[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c, \ d, \ e, \ m\}, \ x] \ \&& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 1.52 (sec), antiderivative size = 95, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{x^2}{4} + \frac{\sin(2cx^2+2bx+2a)}{8c} - \frac{b\sqrt{\pi}(\cos(\frac{-4ac+b^2}{2c})\text{FresnelC}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}})+\sin(\frac{-4ac+b^2}{2c})\text{FresnelS}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}))}{8c^{\frac{3}{2}}}$	95
risch	$\frac{x^2}{4} - \frac{b\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{2c}}\sqrt{2}\text{erf}(\sqrt{2}\sqrt{ic}x+\frac{ib\sqrt{2}}{2\sqrt{ic}})}{32c\sqrt{ic}} + \frac{b\sqrt{\pi}e^{\frac{i(4ac-b^2)}{2c}}\text{erf}(-\sqrt{-2ic}x+\frac{ib}{\sqrt{-2ic}})}{16c\sqrt{-2ic}} + \frac{\sin(2cx^2+2bx+2a)}{8c}$	141

input $\text{int}(x*\cos(c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{4}x^2 + \frac{1}{8}\sin(2cx^2+2bx+2a)/c - \frac{1}{8}b/c^{(3/2)}\text{Pi}^{(1/2)}(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2cx+b)/c^{(1/2)}/\text{Pi}^{(1/2)})+\sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2cx+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 124, normalized size of antiderivative = 0.98

$$\int x \cos^2(a + bx + cx^2) dx = \frac{\pi b \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi b \sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2-4ac}{2c}\right) - 2c^2x^2 - 2c \cos(cx^2 + a) + C}{8c^2}$$

input $\text{integrate}(x*\cos(c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

output

```
-1/8*(pi*b*sqrt(c/pi)*cos(-1/2*(b^2 - 4*a*c)/c)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - pi*b*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c)*sin(-1/2*(b^2 - 4*a*c)/c) - 2*c^2*x^2 - 2*c*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a))/c^2
```

Sympy [F]

$$\int x \cos^2(a + bx + cx^2) dx = \int x \cos^2(a + bx + cx^2) dx$$

input

```
integrate(x*cos(c*x**2+b*x+a)**2,x)
```

output

```
Integral(x*cos(a + b*x + c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 610, normalized size of antiderivative = 4.84

$$\int x \cos^2(a + bx + cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

$$\begin{aligned} & \frac{1}{64}\sqrt{2}(((I - 1)\sqrt{2}\sqrt{\pi})(\operatorname{erf}(\sqrt{1/2})\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) - (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{1/2})\sqrt{-((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)} - 1)*b^2*\cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)\sqrt{2}\sqrt{\pi})(\operatorname{erf}(\sqrt{1/2})\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) - (I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{1/2})\sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)} - 1)) * b^2*\sin(-1/2*(b^2 - 4*a*c)/c) - 2*((-(I - 1)\sqrt{2}\sqrt{\pi})(\operatorname{erf}(\sqrt{1/2})\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) + (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{1/2})\sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)} - 1)) * b*c*\cos(-1/2*(b^2 - 4*a*c)/c) + (-(I + 1)*\sqrt{2}\sqrt{\pi})(\operatorname{erf}(\sqrt{1/2})\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) + (I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{1/2})\sqrt{(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)} - 1)) * b*c*\sin(-1/2*(b^2 - 4*a*c)/c)) * x + 2*\sqrt{2}*(4*c^2*x^2 - c*(I*e^{(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}) - I*e^{(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)})*\cos(-1/2*(b^2 - 4*a*c)/c) + c*(e^{(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}) * \sin(-1/2*(b^2 - 4*a*c)/c)) * \sqrt{(4*c^2*x^2 + 4*b*c*x + b^2)/c}) / (c^2*\sqrt{(4*c^2*x^2 + 4*b*c*x + b^2)/c})) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec), antiderivative size = 170, normalized size of antiderivative = 1.35

$$\begin{aligned} & \int x \cos^2(a + bx + cx^2) dx \\ &= \frac{1}{4}x^2 + \frac{\frac{\sqrt{\pi}b \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{ib^2-4iac}{2c}\right)}}{\sqrt{c}\left(-\frac{ic}{|c|}+1\right)} - ie^{(2icx^2+2ibx+2ia)}}{16c} \\ &+ \frac{\frac{\sqrt{\pi}b \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{-ib^2+4iac}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|}+1\right)} + ie^{(-2icx^2-2ibx-2ia)}}{16c} \end{aligned}$$

input `integrate(x*cos(c*x^2+b*x+a)^2,x, algorithm="giac")`

output

```
1/4*x^2 + 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) - I*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a))/c + 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)) + I*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \cos^2(a + bx + cx^2) dx = \int x \cos(cx^2 + bx + a)^2 dx$$

input

```
int(x*cos(a + b*x + c*x^2)^2,x)
```

output

```
int(x*cos(a + b*x + c*x^2)^2, x)
```

Reduce [F]

$$\int x \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 x dx$$

input

```
int(x*cos(c*x^2+b*x+a)^2,x)
```

output

```
int(cos(a + b*x + c*x**2)**2*x,x)
```

3.18 $\int \cos^2(a + bx + cx^2) dx$

Optimal result	147
Mathematica [A] (verified)	147
Rubi [A] (verified)	148
Maple [A] (verified)	149
Fricas [A] (verification not implemented)	149
Sympy [A] (verification not implemented)	150
Maxima [C] (verification not implemented)	150
Giac [C] (verification not implemented)	151
Mupad [F(-1)]	151
Reduce [F]	152

Optimal result

Integrand size = 13, antiderivative size = 100

$$\int \cos^2(a + bx + cx^2) dx = \frac{x}{2} + \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}}$$

output

```
1/2*x+1/4*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(1/2)-1/4*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec), antiderivative size = 97, normalized size of antiderivative = 0.97

$$\int \cos^2(a + bx + cx^2) dx = \frac{2\sqrt{c}x + \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}}$$

input `Integrate[Cos[a + b*x + c*x^2]^2, x]`

output
$$\frac{(2\sqrt{c}x + \sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] - \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])}{(4\sqrt{c})}$$

Rubi [A] (verified)

Time = 0.24 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3931, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(a + bx + cx^2) dx \\ & \quad \downarrow \text{3931} \\ & \int \left(\frac{1}{2} \cos(2a + 2bx + 2cx^2) + \frac{1}{2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2} \end{aligned}$$

input `Int[Cos[a + b*x + c*x^2]^2, x]`

output
$$\frac{x}{2} + \frac{(\sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] - \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])}{(4\sqrt{c})}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3931 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTri
gReduce[Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.42 (sec), antiderivative size = 72, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4\sqrt{c}}$	72
risch	$\frac{x}{2} + \frac{\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{16\sqrt{ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \operatorname{erf}\left(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{8\sqrt{-2ic}}$	111

input `int(cos(c*x^2+b*x+a)^2, x, method=_RETURNVERBOSE)`

output `1/2*x+1/4*Pi^(1/2)/c^(1/2)*(cos(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))+sin(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 93, normalized size of antiderivative = 0.93

$$\int \cos^2(a + bx + cx^2) dx \\ = \frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2-4ac}{2c}\right) + 2cx}{4c}$$

input `integrate(cos(c*x^2+b*x+a)^2, x, algorithm="fricas")`

output
$$\frac{1}{4} \left(\frac{\pi \sqrt{c/\pi} \cos(-1/2(b^2 - 4ac)/c) \operatorname{fresnel_cos}(2cx + b) \sqrt{c/\pi}/c - \pi \sqrt{c/\pi} \operatorname{fresnel_sin}(2cx + b) \sqrt{c/\pi}/c \sin(-1/2(b^2 - 4ac)/c) + 2cx)}{c} \right)$$

Sympy [A] (verification not implemented)

Time = 0.44 (sec), antiderivative size = 83, normalized size of antiderivative = 0.83

$$\int \cos^2(a + bx + cx^2) dx \\ = \frac{x}{2} + \frac{\sqrt{\pi} \left(-\sin\left(2a - \frac{b^2}{2c}\right) S\left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}}\right) + \cos\left(2a - \frac{b^2}{2c}\right) C\left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}}\right) \right) \sqrt{\frac{1}{c}}}{4}$$

input `integrate(cos(c*x**2+b*x+a)**2,x)`

output
$$\frac{x/2 + \sqrt{\pi} (-\sin(2a - b^2/(2c)) \operatorname{fresnels}((2b + 4cx)/(2\sqrt{\pi}\sqrt{c})) + \cos(2a - b^2/(2c)) \operatorname{fresnelc}((2b + 4cx)/(2\sqrt{\pi}\sqrt{c}))) \sqrt{1/c}}{4}$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec), antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \cos^2(a + bx + cx^2) dx = \\ -\frac{4^{1/4} \sqrt{2} \sqrt{\pi} \left(((i-1) \cos\left(-\frac{b^2-4ac}{2c}\right) + (i+1) \sin\left(-\frac{b^2-4ac}{2c}\right)) \operatorname{erf}\left(\frac{2icx+ib}{\sqrt{2}ic}\right) + ((i+1) \cos\left(-\frac{b^2-4ac}{2c}\right) - (i-1) \sin\left(-\frac{b^2-4ac}{2c}\right)) \operatorname{erfc}\left(\frac{2icx+ib}{\sqrt{2}ic}\right) \right)}{32c^2}$$

input `integrate(cos(c*x^2+b*x+a)^2,x, algorithm="maxima")`

output
$$-\frac{1}{32} \left(4^{(1/4)} \sqrt{2} \sqrt{\pi} (((I-1) \cos(-1/2(b^2 - 4ac)/c) + (I+1) \sin(-1/2(b^2 - 4ac)/c)) \operatorname{erf}((2*I*c*x + I*b)/\sqrt{2*I*c}) + ((I+1) \cos(-1/2(b^2 - 4ac)/c) + (I-1) \sin(-1/2(b^2 - 4ac)/c)) \operatorname{erfc}((2*I*c*x + I*b)/\sqrt{-2*I*c})) c^{(3/2)} - 16*c^2*x)/c^2 \right)$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.22

$$\int \cos^2(a + bx + cx^2) dx = \frac{1}{2}x - \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 - 4ic}{2c}\right)}}{8\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)} \\ - \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 + 4ic}{2c}\right)}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}$$

input `integrate(cos(c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/2*x - 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) - 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 dx$$

input `int(cos(a + b*x + c*x^2)^2,x)`

output `int(cos(a + b*x + c*x^2)^2, x)`

Reduce [F]

$$\int \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 dx$$

input `int(cos(c*x^2+b*x+a)^2,x)`

output `int(cos(a + b*x + c*x**2)**2,x)`

3.19 $\int \frac{\cos^2(a+bx+cx^2)}{x} dx$

Optimal result	153
Mathematica [N/A]	153
Rubi [N/A]	154
Maple [N/A]	155
Fricas [N/A]	155
Sympy [N/A]	155
Maxima [N/A]	156
Giac [N/A]	156
Mupad [N/A]	157
Reduce [N/A]	157

Optimal result

Integrand size = 17, antiderivative size = 17

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \frac{\log(x)}{2} + \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx + 2cx^2)}{x}, x\right)$$

output `1/2*ln(x)+1/2*Defe r (Int)(cos(2*c*x^2+2*b*x+2*a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.63 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos^2(a + bx + cx^2)}{x} dx$$

input `Integrate[$\cos[a + b*x + c*x^2]^2/x$, x]`

output `Integrate[$\cos[a + b*x + c*x^2]^2/x$, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(a + bx + cx^2)}{x} dx \\ & \quad \downarrow \textcolor{blue}{3949} \\ & \int \left(\frac{\cos(2a + 2bx + 2cx^2)}{2x} + \frac{1}{2x} \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{1}{2} \int \frac{\cos(2cx^2 + 2bx + 2a)}{x} dx + \frac{\log(x)}{2} \end{aligned}$$

input `Int[Cos[a + b*x + c*x^2]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_)*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

input `int(cos(c*x^2+b*x+a)^2/x,x)`

output `int(cos(c*x^2+b*x+a)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

input `integrate(cos(c*x^2+b*x+a)^2/x,x, algorithm="fricas")`

output `integral(cos(c*x^2 + b*x + a)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.79 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos^2(a + bx + cx^2)}{x} dx$$

input `integrate(cos(c*x**2+b*x+a)**2/x,x)`

output `Integral(cos(a + b*x + c*x**2)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.65

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

input `integrate(cos(c*x^2+b*x+a)^2/x,x, algorithm="maxima")`

output `1/2*integrate(cos(2*c*x^2 + 2*b*x + 2*a)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

input `integrate(cos(c*x^2+b*x+a)^2/x,x, algorithm="giac")`

output `integrate(cos(c*x^2 + b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 40.58 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

input `int(cos(a + b*x + c*x^2)^2/x,x)`

output `int(cos(a + b*x + c*x^2)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

input `int(cos(c*x^2+b*x+a)^2/x,x)`

output `int(cos(a + b*x + c*x**2)**2/x,x)`

3.20 $\int x^2 \cos^2(a + bx - cx^2) dx$

Optimal result	158
Mathematica [A] (verified)	159
Rubi [A] (verified)	159
Maple [A] (verified)	160
Fricas [A] (verification not implemented)	161
Sympy [F]	161
Maxima [C] (verification not implemented)	162
Giac [C] (verification not implemented)	163
Mupad [F(-1)]	163
Reduce [F]	164

Optimal result

Integrand size = 18, antiderivative size = 248

$$\begin{aligned} \int x^2 \cos^2(a + bx - cx^2) dx = & \frac{x^3}{6} - \frac{b^2 \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & + \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{3/2}} \\ & - \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{5/2}} \\ & - \frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c} \end{aligned}$$

output

```
1/6*x^3-1/16*b^2*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(5/2)+1/16*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)-1/16*Pi^(1/2)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)/c^(3/2)-1/16*b^2*Pi^(1/2)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)/c^(5/2)-1/16*b*sin(-2*c*x^2+2*b*x+2*a)/c^2-1/8*x*sin(-2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.71

$$\int x^2 \cos^2(a + bx - cx^2) dx \\ = \frac{-3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a + \frac{b^2}{2c}\right) - b^2 \sin\left(2a + \frac{b^2}{2c}\right)\right) + 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a + \frac{b^2}{2c}\right) - 3c \sin\left(2a + \frac{b^2}{2c}\right)\right)}{48c^{5/2}}$$

input `Integrate[x^2*Cos[a + b*x - c*x^2]^2, x]`

output
$$\frac{(-3\sqrt{\pi} \operatorname{FresnelS}\left[\frac{(-b + 2c*x)}{\sqrt{c}\sqrt{\pi}}\right] * (c \cos[2a + b^2/(2*c)] - b^2 \sin[2a + b^2/(2*c)]) + 3\sqrt{\pi} \operatorname{FresnelC}\left[\frac{(-b + 2c*x)}{\sqrt{c}\sqrt{\pi}}\right] * (b^2 \cos[2a + b^2/(2*c)] + c \sin[2a + b^2/(2*c)]) + \sqrt{c} * (8*c^2*x^3 - 3*(b + 2*c*x)*\sin[2*(a + x*(b - c*x))]))}{48*c^{(5/2)}}$$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \cos^2(a + bx - cx^2) dx \\ \downarrow 3949 \\ \int \left(\frac{1}{2}x^2 \cos(2a + 2bx - 2cx^2) + \frac{x^2}{2}\right) dx \\ \downarrow 2009$$

$$\begin{aligned}
& - \frac{\sqrt{\pi} \sin \left(2a + \frac{b^2}{2c}\right) \text{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \frac{\sqrt{\pi} b^2 \cos \left(2a + \frac{b^2}{2c}\right) \text{FresnelC} \left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} - \\
& \frac{\sqrt{\pi} b^2 \sin \left(2a + \frac{b^2}{2c}\right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \frac{\sqrt{\pi} \cos \left(2a + \frac{b^2}{2c}\right) \text{FresnelS} \left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \\
& \frac{b \sin \left(2a + 2bx - 2cx^2\right)}{16c^2} - \frac{x \sin \left(2a + 2bx - 2cx^2\right)}{8c} + \frac{x^3}{6}
\end{aligned}$$

input `Int[x^2*Cos[a + b*x - c*x^2]^2,x]`

output $x^{3/6} - (b^{2*}\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a + b^{2/(2*c)}]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])]/(16*c^{(5/2)}) + (\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a + b^{2/(2*c)}]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])]/(16*c^{(3/2)}) - (\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a + b^{2/(2*c)}]/(16*c^{(3/2)}) - (b^{2*}\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a + b^{2/(2*c)}]/(16*c^{(5/2)}) - (b*\text{Sin}[2*a + 2*b*x - 2*c*x^2])/(16*c^2) - (x*\text{Sin}[2*a + 2*b*x - 2*c*x^2])/(8*c)$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_)*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.69 (sec), antiderivative size = 199, normalized size of antiderivative = 0.80

method	result
default	$\frac{x^3}{6} - \frac{x \sin(-2cx^2 + 2bx + 2a)}{8c} + \frac{b \left(-\frac{\sin(-2cx^2 + 2bx + 2a)}{4c} + \frac{b\sqrt{\pi} \left(\cos\left(\frac{4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{3/2}} \right)}{4c}$
risch	$\frac{x^3}{6} + \frac{b^2\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \text{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{32c^2\sqrt{-2ic}} + \frac{i\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \text{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{32c\sqrt{-2ic}} - \frac{b^2\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \text{erf}\left(-\sqrt{2}x + \frac{b}{\sqrt{2c}}\right)}{64c^2\sqrt{ic}}$

input `int(x^2*cos(-c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1/6*x^3 - 1/8*x*sin(-2*c*x^2 + 2*b*x + 2*a)/c + 1/4*b/c*(-1/4*sin(-2*c*x^2 + 2*b*x + 2*a)/c + 1/4*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)) + sin(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)))) - 1/16/c^(3/2)*Pi^(1/2)*(cos(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)) - sin(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)))}{48*c^3}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.75

$$\int x^2 \cos^2(a + bx - cx^2) dx = \frac{8 c^3 x^3 + 6 (2 c^2 x + bc) \cos(cx^2 - bx - a) \sin(cx^2 - bx - a) + 3 \left(\pi b^2 \cos\left(\frac{b^2 + 4ac}{2c}\right) + \pi c \sin\left(\frac{b^2 + 4ac}{2c}\right)\right) \sqrt{48 c^3}}{48 c^3}$$

input `integrate(x^2*cos(-c*x^2+b*x+a)^2,x, algorithm="fricas")`

output
$$\frac{1/48*(8*c^3*x^3 + 6*(2*c^2*x + b*c)*cos(c*x^2 - b*x - a)*sin(c*x^2 - b*x - a) + 3*(pi*b^2*cos(1/2*(b^2 + 4*a*c)/c) + pi*c*sin(1/2*(b^2 + 4*a*c)/c))*sqrt(c/pi)*fresnel_cos((2*c*x - b)*sqrt(c/pi)/c) + 3*(pi*b^2*sin(1/2*(b^2 + 4*a*c)/c) - pi*c*cos(1/2*(b^2 + 4*a*c)/c))*sqrt(c/pi)*fresnel_sin((2*c*x - b)*sqrt(c/pi)/c))/c^3}{48*c^3}$$

Sympy [F]

$$\int x^2 \cos^2(a + bx - cx^2) dx = \int x^2 \cos^2(a + bx - cx^2) dx$$

input `integrate(x**2*cos(-c*x**2+b*x+a)**2,x)`

output `Integral(x**2*cos(a + b*x - c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.83 (sec) , antiderivative size = 1617, normalized size of antiderivative = 6.52

$$\int x^2 \cos^2(a + bx - cx^2) dx = \text{Too large to display}$$

input `integrate(x^2*cos(-c*x^2+b*x+a)^2,x, algorithm="maxima")`

output `-1/384*sqrt(2)*(24*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*cos(1/2*(b^2 + 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*sin(1/2*(b^2 + 4*a*c)/c))*x^3 + 36*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(1/2*(b^2 + 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)...`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.86

$$\int x^2 \cos^2(a + bx - cx^2) dx = \frac{1}{6} x^3 - \frac{(c(2ix - \frac{ib}{c}) + 2ib)e^{(2icx^2 - 2ibx - 2ia)} + \frac{\sqrt{\pi}(b^2 + ic)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x - \frac{b}{c}\right)\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 + 4ic}{2c}\right)}}{\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)}}{32c^2} - \frac{(c(-2ix + \frac{ib}{c}) - 2ib)e^{(-2icx^2 + 2ibx + 2ia)} + \frac{\sqrt{\pi}(b^2 - ic)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x - \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 - 4ic}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2}$$

input `integrate(x^2*cos(-c*x^2+b*x+a)^2,x, algorithm="giac")`

output $\frac{1}{6}x^3 - \frac{1}{32}((c(2Ix - Ib/c) + 2Ib)*e^{(2Ic*x^2 - 2Ib*x - 2Ia)} + \sqrt{\pi}*(b^2 + Ic)*\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x - b/c)(-Ic/abs(c) + 1)\right)*e^{(-1/2*(Ib^2 + 4Ia*c)/c)} / (\sqrt{c}*(-Ic/abs(c) + 1)))/c^2 - \frac{1}{32}((c(-2Ix + Ib/c) - 2Ib)*e^{(-2Ic*x^2 + 2Ib*x + 2Ia)} + \sqrt{\pi}*(b^2 - Ic)*\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x - b/c)(Ic/abs(c) + 1)\right)*e^{(-1/2*(-Ib^2 - 4Ia*c)/c)} / (\sqrt{c}(Ic/abs(c) + 1)))/c^2$

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2(a + bx - cx^2) dx = \int x^2 \cos(-cx^2 + bx + a)^2 dx$$

input `int(x^2*cos(a + b*x - c*x^2)^2,x)`

output `int(x^2*cos(a + b*x - c*x^2)^2, x)`

Reduce [F]

$$\int x^2 \cos^2(a + bx - cx^2) dx$$

$$= -\cos(-cx^2 + bx + a) \sin(-cx^2 + bx + a) b - 2\cos(-cx^2 + bx + a) \sin(-cx^2 + bx + a) cx - 6 \left(\int \tan$$

input `int(x^2*cos(-c*x^2+b*x+a)^2,x)`

output
$$(-\cos(a + bx - cx^2)*\sin(a + bx - cx^2)*b - 2*\cos(a + bx - cx^2)*\sin(a + bx - cx^2)*c*x - 6*\int(\tan((a + bx - cx^2)/2)^2/(\tan((a + bx - cx^2)/2)^4 + 2*\tan((a + bx - cx^2)/2)^2 + 1),x)*b^2 - 8*\int(x^2/(\tan((a + bx - cx^2)/2)^4 + 2*\tan((a + bx - cx^2)/2)^2 + 1),x)*c^2 + 8*\int(\tan((a + bx - cx^2)/2)/(\tan((a + bx - cx^2)/2)^4 + 2*\tan((a + bx - cx^2)/2)^2 + 1),x)*c + 2*\int(1/(\tan((a + bx - cx^2)/2)^4 + 2*\tan((a + bx - cx^2)/2)^2 + 1),x)*b^2 - \sin(a + bx - cx^2)*b - 2*\sin(a + bx - cx^2)*c*x + 2*c^2*x^3)/(6*c^2)$$

3.21 $\int x \cos^2(a + bx - cx^2) dx$

Optimal result	165
Mathematica [A] (verified)	165
Rubi [A] (verified)	166
Maple [A] (verified)	167
Fricas [A] (verification not implemented)	167
Sympy [F]	168
Maxima [C] (verification not implemented)	168
Giac [C] (verification not implemented)	169
Mupad [F(-1)]	170
Reduce [F]	170

Optimal result

Integrand size = 16, antiderivative size = 126

$$\begin{aligned} \int x \cos^2(a + bx - cx^2) dx &= \frac{x^2}{4} - \frac{b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ &\quad - \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{8c^{3/2}} \\ &\quad - \frac{\sin(2a + 2bx - 2cx^2)}{8c} \end{aligned}$$

output

```
1/4*x^2-1/8*b*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)-1/8*b*Pi^(1/2)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)/c^(3/2)-1/8*sin(-2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.28 (sec), antiderivative size = 122, normalized size of antiderivative = 0.97

$$\begin{aligned} \int x \cos^2(a + bx - cx^2) dx \\ = \frac{b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + b\sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 - \sin(2(a + x))}{8c^{3/2}} \end{aligned}$$

input `Integrate[x*Cos[a + b*x - c*x^2]^2, x]`

output
$$\begin{aligned} & (b \sqrt{\pi}) \cos[2a + b^2/(2c)] \operatorname{FresnelC}[-b + 2cx]/(\sqrt{c} \sqrt{\pi}) \\ & + b \sqrt{\pi} \operatorname{FresnelS}[-b + 2cx]/(\sqrt{c} \sqrt{\pi}) \sin[2a + b^2/(2c)] \\ & + \sqrt{c} (2cx^2 - \sin[2(a + bx - cx^2)])/(8c^{(3/2)}) \end{aligned}$$

Rubi [A] (verified)

Time = 0.28 (sec), antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \cos^2(a + bx - cx^2) dx \\ & \downarrow 3949 \\ & \int \left(\frac{1}{2}x \cos(2a + 2bx - 2cx^2) + \frac{x}{2} \right) dx \\ & \downarrow 2009 \\ & - \frac{\sqrt{\pi}b \cos\left(2a + \frac{b^2}{2c}\right) \operatorname{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} - \frac{\sqrt{\pi}b \sin\left(2a + \frac{b^2}{2c}\right) \operatorname{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} - \\ & \quad \frac{\sin(2a + 2bx - 2cx^2)}{8c} + \frac{x^2}{4} \end{aligned}$$

input `Int[x*Cos[a + b*x - c*x^2]^2, x]`

output
$$\begin{aligned} & x^2/4 - (b \sqrt{\pi}) \cos[2a + b^2/(2c)] \operatorname{FresnelC}[(b - 2cx)/(\sqrt{c} \sqrt{\pi})]/(8c^{(3/2)}) \\ & - (b \sqrt{\pi}) \operatorname{FresnelS}[(b - 2cx)/(\sqrt{c} \sqrt{\pi})] \sin[2a + b^2/(2c)]/(8c^{(3/2)}) - \sin[2a + 2bx - 2cx^2]/(8c) \end{aligned}$$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_)^m), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.56 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x^2}{4} - \frac{\sin(-2cx^2+2bx+2a)}{8c} + \frac{b\sqrt{\pi} \left(\cos\left(\frac{4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) \right)}{8c^{3/2}}$	99
risch	$\frac{x^2}{4} + \frac{b\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \text{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{16c\sqrt{-2ic}} - \frac{b\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \text{erf}\left(-\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c\sqrt{ic}} - \frac{\sin(-2cx^2+2bx+2a)}{8c}$	13

input `int(x*cos(-c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^2-1/8*sin(-2*c*x^2+2*b*x+2*a)/c+1/8*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(4*a*c+b^2)/c)*FresnelC(1/Pi^(1/2)/c^(1/2)*(2*c*x-b))+sin(1/2*(4*a*c+b^2)/c)*FresnelS(1/Pi^(1/2)/c^(1/2)*(2*c*x-b)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.06

$$\int x \cos^2(a + bx - cx^2) dx = \frac{\pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{2c}\right) C\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) + \pi b \sqrt{\frac{c}{\pi}} S\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(\frac{b^2+4ac}{2c}\right) + 2c^2x^2 + 2c \cos(cx^2 - bx - \frac{b^2+4ac}{2c})}{8c^2}$$

input `integrate(x*cos(-c*x^2+b*x+a)^2,x, algorithm="fricas")`

output

```
1/8*(pi*b*sqrt(c/pi)*cos(1/2*(b^2 + 4*a*c)/c)*fresnel_cos((2*c*x - b)*sqrt(c/pi)/c) + pi*b*sqrt(c/pi)*fresnel_sin((2*c*x - b)*sqrt(c/pi)/c)*sin(1/2*(b^2 + 4*a*c)/c) + 2*c^2*x^2 + 2*c*cos(c*x^2 - b*x - a)*sin(c*x^2 - b*x - a))/c^2
```

Sympy [F]

$$\int x \cos^2(a + bx - cx^2) dx = \int x \cos^2(a + bx - cx^2) dx$$

input

```
integrate(x*cos(-c*x**2+b*x+a)**2,x)
```

output

```
Integral(x*cos(a + b*x - c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 611, normalized size of antiderivative = 4.85

$$\int x \cos^2(a + bx - cx^2) dx = \text{Too large to display}$$

input

```
integrate(x*cos(-c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

```
1/64*sqrt(2)*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(1/2*(b^2 + 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(1/2*(b^2 + 4*a*c)/c) - 2*((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(1/2*(b^2 + 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(1/2*(b^2 + 4*a*c)/c) + 2*sqrt(2)*(4*c^2*x^2 - c*(I*e^(1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*cos(1/2*(b^2 + 4*a*c)/c) - c*(e^(1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + e^(-1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*sin(1/2*(b^2 + 4*a*c)/c))*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c))/(c^2*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c)))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.37

$$\int x \cos^2(a + bx - cx^2) dx$$

$$= \frac{1}{4} x^2 - \frac{\frac{\sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x - \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right)\right) e^{\left(-\frac{i b^2 + 4 i a c}{2 c}\right)}}{\sqrt{c} \left(-\frac{i c}{|c|} + 1\right)} + i e^{(2 i c x^2 - 2 i b x - 2 i a)}}{16 c}$$

$$- \frac{\frac{\sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} \sqrt{c} \left(2x - \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right)\right) e^{\left(-\frac{-i b^2 - 4 i a c}{2 c}\right)}}{\sqrt{c} \left(\frac{i c}{|c|} + 1\right)} - i e^{(-2 i c x^2 + 2 i b x + 2 i a)}}{16 c}$$

input `integrate(x*cos(-c*x^2+b*x+a)^2,x, algorithm="giac")`

output

```
1/4*x^2 - 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x - b/c)*(-I*c/abs(c) + 1))
 *e^(-1/2*(I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) + I*e^(2*I*c*x^2
 - 2*I*b*x - 2*I*a))/c - 1/16*(sqrt(pi)*b*erf(-1/2*sqrt(c)*(2*x - b/c)*(I*
 c/abs(c) + 1))*e^(-1/2*(-I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)) -
 I*e^(-2*I*c*x^2 + 2*I*b*x + 2*I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \cos^2(a + bx - cx^2) dx = \int x \cos(-cx^2 + bx + a)^2 dx$$

input

```
int(x*cos(a + b*x - c*x^2)^2,x)
```

output

```
int(x*cos(a + b*x - c*x^2)^2, x)
```

Reduce [F]

$$\int x \cos^2(a + bx - cx^2) dx = \int \cos(-cx^2 + bx + a)^2 x dx$$

input

```
int(x*cos(-c*x^2+b*x+a)^2,x)
```

output

```
int(cos(a + b*x - c*x**2)**2*x,x)
```

3.22 $\int \cos^2(a + bx - cx^2) dx$

Optimal result	171
Mathematica [A] (verified)	171
Rubi [A] (verified)	172
Maple [A] (verified)	173
Fricas [A] (verification not implemented)	173
Sympy [A] (verification not implemented)	174
Maxima [C] (verification not implemented)	174
Giac [C] (verification not implemented)	175
Mupad [F(-1)]	175
Reduce [F]	176

Optimal result

Integrand size = 14, antiderivative size = 100

$$\int \cos^2(a + bx - cx^2) dx = \frac{x}{2} - \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} \\ - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}}$$

output

```
1/2*x-1/4*Pi^(1/2)*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))
)/c^(1/2)-1/4*Pi^(1/2)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b
^2/c)/c^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00

$$\int \cos^2(a + bx - cx^2) dx \\ = \frac{2\sqrt{c}x + \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + \sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}}$$

input `Integrate[Cos[a + b*x - c*x^2]^2, x]`

output
$$\frac{(2\sqrt{c}x + \sqrt{\pi}\cos[2a + b^2/(2c)]\text{FresnelC}[(-b + 2cx)/(\sqrt{c}\sqrt{\pi})] + \sqrt{\pi}\text{FresnelS}[(-b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a + b^2/(2c)])}{(4\sqrt{c})}$$

Rubi [A] (verified)

Time = 0.25 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3931, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2(a + bx - cx^2) dx \\ & \quad \downarrow \text{3931} \\ & \int \left(\frac{1}{2} \cos(2a + 2bx - 2cx^2) + \frac{1}{2} \right) dx \\ & \quad \downarrow \text{2009} \\ & - \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2} \end{aligned}$$

input `Int[Cos[a + b*x - c*x^2]^2, x]`

output
$$\frac{x}{2} - \frac{(\sqrt{\pi}\cos[2a + b^2/(2c)]\text{FresnelC}[(b - 2cx)/(\sqrt{c}\sqrt{\pi})])}{(4\sqrt{c})} - \frac{(\sqrt{\pi}\text{FresnelS}[(b - 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a + b^2/(2c)])}{(4\sqrt{c})}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3931 $\text{Int}[\cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTri}\text{gReduce}[\cos[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c\}, \ x] \ \&\& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 1.36 (sec), antiderivative size = 76, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{\pi} (\cos(\frac{4ac+b^2}{2c}) \text{FresnelC}(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}) + \sin(\frac{4ac+b^2}{2c}) \text{FresnelS}(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}))}{4\sqrt{c}}$	76
risch	$\frac{x}{2} + \frac{\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \text{erf}(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}})}{8\sqrt{-2ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \text{erf}(-\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}})}{16\sqrt{ic}}$	107

input $\text{int}(\cos(-c*x^2+b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $1/2*x+1/4*\text{Pi}^{(1/2)}/c^{(1/2)}*(\cos(1/2*(4*a*c+b^2)/c)*\text{FresnelC}(1/\text{Pi}^{(1/2)}/c^{(1/2)}*(2*c*x-b))+\sin(1/2*(4*a*c+b^2)/c)*\text{FresnelS}(1/\text{Pi}^{(1/2)}/c^{(1/2)}*(2*c*x-b)))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 96, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \cos^2(a + bx - cx^2) \, dx \\ &= \frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{2c}\right) C\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) + \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(\frac{b^2+4ac}{2c}\right) + 2cx}{4c} \end{aligned}$$

input $\text{integrate}(\cos(-c*x^2+b*x+a)^2, x, \text{algorithm}=\text{"fricas"})$

output
$$\frac{1}{4} \left(\frac{\pi \sqrt{c/\pi} \cos(1/2*(b^2 + 4*a*c)/c) \operatorname{fresnel_cos}(2*c*x - b) \sqrt{c/\pi}}{c} + \frac{\pi \sqrt{c/\pi} \operatorname{fresnel_sin}(2*c*x - b) \sqrt{c/\pi}}{c} \right) + \frac{\sin(1/2*(b^2 + 4*a*c)/c) + 2*c*x}{c}$$

Sympy [A] (verification not implemented)

Time = 0.42 (sec), antiderivative size = 88, normalized size of antiderivative = 0.88

$$\int \cos^2(a + bx - cx^2) dx \\ = \frac{x}{2} + \frac{\sqrt{\pi} \sqrt{-\frac{1}{c}} \left(-\sin\left(2a + \frac{b^2}{2c}\right) S\left(\frac{2b-4cx}{2\sqrt{\pi}\sqrt{-c}}\right) + \cos\left(2a + \frac{b^2}{2c}\right) C\left(\frac{2b-4cx}{2\sqrt{\pi}\sqrt{-c}}\right) \right)}{4}$$

input `integrate(cos(-c*x**2+b*x+a)**2,x)`

output
$$\frac{x/2 + \sqrt{\pi} \sqrt{-1/c} (-\sin(2*a + b**2/(2*c)) \operatorname{fresnels}((2*b - 4*c*x)/(2*sqrt(pi)*sqrt(-c))) + \cos(2*a + b**2/(2*c)) \operatorname{fresnelc}((2*b - 4*c*x)/(2*sqrt(pi)*sqrt(-c))))}{4}$$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec), antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \cos^2(a + bx - cx^2) dx \\ = \frac{4^{1/4} \sqrt{2} \sqrt{\pi} \left(\left(-(i-1) \cos\left(\frac{b^2+4ac}{2c}\right) + (i+1) \sin\left(\frac{b^2+4ac}{2c}\right) \right) \operatorname{erf}\left(\frac{2icx-i b}{\sqrt{2}i c}\right) + \left(-(i+1) \cos\left(\frac{b^2+4ac}{2c}\right) + (i-1) \sin\left(\frac{b^2+4ac}{2c}\right) \right) \operatorname{erf}\left(\frac{2icx+i b}{\sqrt{2}i c}\right) \right)}{32c^2}$$

input `integrate(cos(-c*x^2+b*x+a)^2,x, algorithm="maxima")`

output
$$\frac{1}{32} \left(4^{1/4} \sqrt{2} \sqrt{\pi} \left(\left(-(I-1) \cos(1/2*(b^2 + 4*a*c)/c) + (I+1) \sin(1/2*(b^2 + 4*a*c)/c) \right) \operatorname{erf}\left((2*I*c*x - I*b)/\sqrt{2*I*c}\right) + \left(-(I+1) \cos(1/2*(b^2 + 4*a*c)/c) + (I-1) \sin(1/2*(b^2 + 4*a*c)/c) \right) \operatorname{erf}\left((2*I*c*x - I*b)/\sqrt{-2*I*c}\right) \right) c^{3/2} + 16*c^2*x \right) / c^2$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \cos^2(a + bx - cx^2) dx = \frac{1}{2}x - \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x - \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2+4ic}{2c}\right)}}{8\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)} \\ - \frac{\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x - \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2-4ic}{2c}\right)}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}$$

input `integrate(cos(-c*x^2+b*x+a)^2,x, algorithm="giac")`

output `1/2*x - 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x - b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) - 1/8*sqrt(pi)*erf(-1/2*sqrt(c)*(2*x - b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))`

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx - cx^2) dx = \int \cos(-cx^2 + bx + a)^2 dx$$

input `int(cos(a + b*x - c*x^2)^2,x)`

output `int(cos(a + b*x - c*x^2)^2, x)`

Reduce [F]

$$\int \cos^2(a + bx - cx^2) dx = \int \cos(-cx^2 + bx + a)^2 dx$$

input `int(cos(-c*x^2+b*x+a)^2,x)`

output `int(cos(a + b*x - c*x**2)**2,x)`

3.23 $\int \frac{\cos^2(a+bx-cx^2)}{x} dx$

Optimal result	177
Mathematica [N/A]	177
Rubi [N/A]	178
Maple [N/A]	179
Fricas [N/A]	179
Sympy [N/A]	179
Maxima [N/A]	180
Giac [N/A]	180
Mupad [N/A]	181
Reduce [N/A]	181

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \frac{\log(x)}{2} + \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx - 2cx^2)}{x}, x\right)$$

output `1/2*ln(x)+1/2*Defe r (Int)(cos(-2*c*x^2+2*b*x+2*a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 3.35 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos^2(a + bx - cx^2)}{x} dx$$

input `Integrate[$\cos[a + b*x - c*x^2]^2/x$, x]`

output `Integrate[$\cos[a + b*x - c*x^2]^2/x$, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(a + bx - cx^2)}{x} dx \\ & \quad \downarrow \textcolor{blue}{3949} \\ & \int \left(\frac{\cos(2a + 2bx - 2cx^2)}{2x} + \frac{1}{2x} \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{1}{2} \int \frac{\cos(-2cx^2 + 2bx + 2a)}{x} dx + \frac{\log(x)}{2} \end{aligned}$$

input `Int[Cos[a + b*x - c*x^2]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_)*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

input `int(cos(-c*x^2+b*x+a)^2/x,x)`

output `int(cos(-c*x^2+b*x+a)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

input `integrate(cos(-c*x^2+b*x+a)^2/x,x, algorithm="fricas")`

output `integral(cos(c*x^2 - b*x - a)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.78 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos^2(a + bx - cx^2)}{x} dx$$

input `integrate(cos(-c*x**2+b*x+a)**2/x,x)`

output `Integral(cos(a + b*x - c*x**2)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.56

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

input `integrate(cos(-c*x^2+b*x+a)^2/x,x, algorithm="maxima")`

output `1/2*integrate(cos(2*c*x^2 - 2*b*x - 2*a)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

input `integrate(cos(-c*x^2+b*x+a)^2/x,x, algorithm="giac")`

output `integrate(cos(-c*x^2 + b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 41.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

input `int(cos(a + b*x - c*x^2)^2/x,x)`

output `int(cos(a + b*x - c*x^2)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

input `int(cos(-c*x^2+b*x+a)^2/x,x)`

output `int(cos(a + b*x - c*x**2)**2/x,x)`

3.24 $\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	182
Mathematica [A] (verified)	182
Rubi [A] (verified)	183
Maple [A] (verified)	184
Fricas [A] (verification not implemented)	184
Sympy [F]	185
Maxima [C] (verification not implemented)	185
Giac [C] (verification not implemented)	186
Mupad [F(-1)]	186
Reduce [F]	187

Optimal result

Integrand size = 15, antiderivative size = 85

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x^3}{6} + \frac{1}{16}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) - \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{\pi}} \right) - \frac{1}{16} \sin \left(\frac{1}{2} + 2x + 2x^2 \right) + \frac{1}{8}x \sin \left(\frac{1}{2} + 2x + 2x^2 \right)$$

output
$$1/6*x^3+1/16*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))-1/16*Pi^(1/2)*FresnelS((1+2*x)/Pi^(1/2))-1/16*sin(1/2+2*x+2*x^2)+1/8*x*sin(1/2+2*x+2*x^2)$$

Mathematica [A] (verified)

Time = 0.17 (sec), antiderivative size = 77, normalized size of antiderivative = 0.91

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{48} \left(8x^3 + 3\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) - 3\sqrt{\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{\pi}} \right) - 3 \sin \left(\frac{1}{2}(1+2x)^2 \right) + 6x \sin \left(\frac{1}{2}(1+2x)^2 \right) \right)$$

input
$$\text{Integrate}[x^2 \cos[1/4 + x + x^2]^2, x]$$

output
$$(8*x^3 + 3*sqrt(Pi)*FresnelC[(1 + 2*x)/sqrt(Pi)] - 3*sqrt(Pi)*FresnelS[(1 + 2*x)/sqrt(Pi)] - 3*Sin[(1 + 2*x)^2/2] + 6*x*Sin[(1 + 2*x)^2/2])/48$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \cos^2 \left(x^2 + x + \frac{1}{4} \right) dx \\ & \quad \downarrow \textcolor{blue}{3949} \\ & \int \left(\frac{x^2}{2} + \frac{1}{2}x^2 \cos \left(2x^2 + 2x + \frac{1}{2} \right) \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{1}{16}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) - \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x^3}{6} + \frac{1}{8}x \sin \left(2x^2 + 2x + \frac{1}{2} \right) - \\ & \quad \frac{1}{16} \sin \left(2x^2 + 2x + \frac{1}{2} \right) \end{aligned}$$

input
$$\operatorname{Int}[x^2 \cos[1/4 + x + x^2]^2, x]$$

output
$$x^3/6 + (\sqrt{Pi}*\operatorname{FresnelC}[(1 + 2*x)/sqrt(Pi)])/16 - (\sqrt{Pi}*\operatorname{FresnelS}[(1 + 2*x)/sqrt(Pi)])/16 - \operatorname{Sin}[1/2 + 2*x + 2*x^2]/16 + (x*\operatorname{Sin}[1/2 + 2*x + 2*x^2])/8$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3949 $\text{Int}[\text{Cos}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_))^{m_.}, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \ \text{Cos}[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c, \ d, \ e, \ m\}, \ x] \ \&& \ \text{IGtQ}[n, \ 1]$

Maple [A] (verified)

Time = 2.16 (sec), antiderivative size = 64, normalized size of antiderivative = 0.75

method	result
default	$\frac{x^3}{6} + \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{16} - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{16} - \frac{\sin\left(\frac{1}{2}+2x+2x^2\right)}{16} + \frac{x \sin\left(\frac{1}{2}+2x+2x^2\right)}{8}$
risch	$\frac{x^3}{6} - \frac{\sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \text{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{64} - \frac{(-1)^{\frac{1}{4}} \sqrt{\pi} \sqrt{2} \text{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{64} + \frac{\sqrt{\pi} \text{erf}\left(\sqrt{-2i} x - \frac{i}{\sqrt{-2i}}\right)}{32\sqrt{-2i}}$

input $\text{int}(x^2 * \cos(1/4 + x + x^2)^2, x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{6}x^3 + \frac{1}{16}\pi^{1/2} \text{FresnelC}\left(\frac{(1+2x)}{\pi^{1/2}}\right) - \frac{1}{16}\pi^{1/2} \text{FresnelS}\left(\frac{(1+2x)}{\pi^{1/2}}\right) - \frac{1}{16}\sin(1/2 + 2x + 2x^2) + \frac{1}{8}x \sin(1/2 + 2x + 2x^2)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 57, normalized size of antiderivative = 0.67

$$\begin{aligned} \int x^2 \cos^2\left(\frac{1}{4} + x + x^2\right) dx &= \frac{1}{6}x^3 + \frac{1}{8}(2x - 1) \cos\left(x^2 + x + \frac{1}{4}\right) \sin\left(x^2 + x + \frac{1}{4}\right) \\ &\quad + \frac{1}{16}\sqrt{\pi} C\left(\frac{2x + 1}{\sqrt{\pi}}\right) - \frac{1}{16}\sqrt{\pi} S\left(\frac{2x + 1}{\sqrt{\pi}}\right) \end{aligned}$$

input $\text{integrate}(x^2 * \cos(1/4 + x + x^2)^2, x, \text{algorithm}=\text{"fricas"})$

output
$$\frac{1}{6}x^3 + \frac{1}{8}(2x - 1)\cos(x^2 + x + 1/4)\sin(x^2 + x + 1/4) + \frac{1}{16}\sqrt{\pi}\text{fresnel_cos}((2x + 1)/\sqrt{\pi}) - \frac{1}{16}\sqrt{\pi}\text{fresnel_sin}((2x + 1)/\sqrt{\pi})$$

Sympy [F]

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x^2 \cos^2 \left(x^2 + x + \frac{1}{4} \right) dx$$

input `integrate(x**2*cos(1/4+x+x**2)**2, x)`

output `Integral(x**2*cos(x**2 + x + 1/4)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.01

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{128x^4 + 64x^3 - 48x \left(-ie^{(2ix^2+2ix+\frac{1}{2}i)} + ie^{(-2ix^2-2ix-\frac{1}{2}i)} \right) - 3\sqrt{8x^2+8x+2} \left((i-1)\sqrt{2}\sqrt{\pi} \operatorname{erf} \left(\sqrt{\frac{x^2+x+\frac{1}{4}}{\pi}} \right) + \sqrt{\frac{x^2+x+\frac{1}{4}}{\pi}} \right)}{192}$$

input `integrate(x^2*cos(1/4+x+x^2)^2, x, algorithm="maxima")`

output
$$\begin{aligned} & \frac{1}{384} \left(128x^4 + 64x^3 - 48x \left(-Ie^{(2Ix^2+2Ix+1/2I)} + Ie^{-(-2Ix^2-2Ix-1/2I)} \right) - 3\sqrt{8x^2+8x+2} \left((I-1)\sqrt{2}\sqrt{\pi} \operatorname{erf} \left(\sqrt{\frac{x^2+x+\frac{1}{4}}{\pi}} \right) - (I+1)\sqrt{2}\sqrt{\pi} \operatorname{erf} \left(\sqrt{\frac{-x^2-x-\frac{1}{4}}{\pi}} \right) \right) - (2I+2)\sqrt{2}\sqrt{\pi} \gamma(3/2, 2Ix^2+2Ix+1/2I) + (2I-2)\sqrt{2}\sqrt{\pi} \gamma(3/2, -2Ix^2-2Ix-1/2I) \right) + 24Ie^{(2Ix^2+2Ix+1/2I)} - 24Ie^{-(-2Ix^2-2Ix-1/2I)} \right) / (2x+1) \end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.75

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{6} x^3 - \frac{1}{32} (2i x - i) e^{(2i x^2 + 2i x + \frac{1}{2}i)} \\ - \frac{1}{32} (-2i x + i) e^{(-2i x^2 - 2i x - \frac{1}{2}i)} \\ - \frac{1}{32} i \sqrt{\pi} \operatorname{erf} \left((i - 1) x + \frac{1}{2}i - \frac{1}{2} \right) \\ + \frac{1}{32} i \sqrt{\pi} \operatorname{erf} \left(-(i + 1) x - \frac{1}{2}i - \frac{1}{2} \right)$$

input `integrate(x^2*cos(1/4+x+x^2)^2,x, algorithm="giac")`

output $\frac{1}{6}x^3 - \frac{1}{32}(2\text{i}x - \text{i})e^{(2\text{i}x^2 + 2\text{i}x + \frac{1}{2}\text{i})} - \frac{1}{32}(-2\text{i}x + \text{i})e^{(-2\text{i}x^2 - 2\text{i}x - \frac{1}{2}\text{i})} - \frac{1}{32}\text{i}\sqrt{\pi}\operatorname{erf}((\text{i} - 1)x + \frac{1}{2}\text{i} - \frac{1}{2}) + \frac{1}{32}\text{i}\sqrt{\pi}\operatorname{erf}(-(\text{i} + 1)x - \frac{1}{2}\text{i} - \frac{1}{2})$

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x^2 \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(x^2*cos(x + x^2 + 1/4)^2,x)`

output `int(x^2*cos(x + x^2 + 1/4)^2, x)`

Reduce [F]

$$\begin{aligned}
 & \int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx \\
 &= \frac{\cos(x^2 + x + \frac{1}{4}) \sin(x^2 + x + \frac{1}{4}) x}{3} - \frac{\cos(x^2 + x + \frac{1}{4}) \sin(x^2 + x + \frac{1}{4})}{6} \\
 &\quad - \left(\int \frac{\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^2}{\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^4 + 2\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^2 + 1} dx \right) \\
 &\quad - \frac{4 \left(\int \frac{x^2}{\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^4 + 2\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^2 + 1} dx \right)}{3} \\
 &\quad - \frac{4 \left(\int \frac{\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})}{\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^4 + 2\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^2 + 1} dx \right)}{3} \\
 &\quad + \frac{\left(\int \frac{1}{\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^4 + 2\tan(\frac{1}{2}x^2 + \frac{1}{2}x + \frac{1}{8})^2 + 1} dx \right)}{3} \\
 &\quad + \frac{\sin(x^2 + x + \frac{1}{4}) x}{3} - \frac{\sin(x^2 + x + \frac{1}{4})}{6} + \frac{x^3}{3}
 \end{aligned}$$

input `int(x^2*cos(1/4+x+x^2)^2,x)`

output

```

(2*cos((4*x**2 + 4*x + 1)/4)*sin((4*x**2 + 4*x + 1)/4)*x - cos((4*x**2 + 4*x + 1)/4)*sin((4*x**2 + 4*x + 1)/4) - 6*int(tan((4*x**2 + 4*x + 1)/8)**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) - 8*int(x**2/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) - 8*int(tan((4*x**2 + 4*x + 1)/8)/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) + 2*int(1/(tan((4*x**2 + 4*x + 1)/8)**4 + 2*tan((4*x**2 + 4*x + 1)/8)**2 + 1),x) + 2*sin((4*x**2 + 4*x + 1)/4)*x - sin((4*x**2 + 4*x + 1)/4) + 2*x**3)/6

```

3.25 $\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	188
Mathematica [A] (verified)	188
Rubi [A] (verified)	189
Maple [A] (verified)	190
Fricas [A] (verification not implemented)	190
Sympy [B] (verification not implemented)	191
Maxima [C] (verification not implemented)	191
Giac [C] (verification not implemented)	192
Mupad [F(-1)]	192
Reduce [F]	193

Optimal result

Integrand size = 13, antiderivative size = 46

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x^2}{4} - \frac{1}{8}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) + \frac{1}{8} \sin \left(\frac{1}{2} + 2x + 2x^2 \right)$$

output 1/4*x^2-1/8*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))+1/8*sin(1/2+2*x+2*x^2)

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{8} \left(2x^2 - \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) + \sin \left(\frac{1}{2}(1+2x)^2 \right) \right)$$

input Integrate[x*Cos[1/4 + x + x^2]^2, x]

output (2*x^2 - Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]] + Sin[(1 + 2*x)^2/2])/8

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \cos^2 \left(x^2 + x + \frac{1}{4} \right) dx \\
 & \quad \downarrow \text{3949} \\
 & \int \left(\frac{1}{2}x \cos \left(2x^2 + 2x + \frac{1}{2} \right) + \frac{x}{2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\frac{1}{8}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x^2}{4} + \frac{1}{8} \sin \left(2x^2 + 2x + \frac{1}{2} \right)
 \end{aligned}$$

input `Int[x*Cos[1/4 + x + x^2]^2, x]`

output `x^2/4 - (Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/8 + Sin[1/2 + 2*x + 2*x^2]/8`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.80 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x^2}{4} - \frac{\sqrt{\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{8} + \frac{\sin\left(\frac{1}{2}+2x+2x^2\right)}{8}$	35
risch	$\frac{x^2}{4} + \frac{\sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \operatorname{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{32} - \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-2i} x - \frac{i}{\sqrt{-2i}}\right)}{16\sqrt{-2i}} + \frac{\sin\left(\frac{(1+2x)^2}{2}\right)}{8}$	72

input `int(x*cos(1/4+x+x^2)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^2-1/8*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))+1/8*sin(1/2+2*x+2*x^2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} x^2 + \frac{1}{4} \cos \left(x^2 + x + \frac{1}{4} \right) \sin \left(x^2 + x + \frac{1}{4} \right) - \frac{1}{8} \sqrt{\pi} C \left(\frac{2x+1}{\sqrt{\pi}} \right)$$

input `integrate(x*cos(1/4+x+x^2)^2,x, algorithm="fricas")`

output `1/4*x^2 + 1/4*cos(x^2 + x + 1/4)*sin(x^2 + x + 1/4) - 1/8*sqrt(pi)*fresnel_cos((2*x + 1)/sqrt(pi))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(37) = 74$.

Time = 0.78 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.63

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x^2}{4} - \frac{\sqrt{\pi} x C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right) \Gamma \left(\frac{1}{4} \right)}{16 \Gamma \left(\frac{5}{4} \right)} + \frac{\sqrt{\pi} x C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right)}{4} \\ + \frac{\sin \left(2(x + \frac{1}{2})^2 \right) \Gamma \left(\frac{1}{4} \right)}{32 \Gamma \left(\frac{5}{4} \right)} - \frac{\sqrt{\pi} C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right) \Gamma \left(\frac{1}{4} \right)}{32 \Gamma \left(\frac{5}{4} \right)}$$

input `integrate(x*cos(1/4+x+x**2)**2,x)`

output `x**2/4 - sqrt(pi)*x*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))*gamma(1/4)/(16*gamma(5/4)) + sqrt(pi)*x*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))/4 + sin(2*(x + 1/2)**2)*gamma(1/4)/(32*gamma(5/4)) - sqrt(pi)*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))*gamma(1/4)/(32*gamma(5/4))`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.23 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.96

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx \\ = \frac{32 x^3 + 16 x^2 + 8 x \left(-i e^{(2i x^2 + 2i x + \frac{1}{2}i)} + i e^{(-2i x^2 - 2i x - \frac{1}{2}i)} \right) + \sqrt{8 x^2 + 8 x + 2} \left((i - 1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{2i x^2 + 2i x + 1} \right) - \operatorname{erf} \left(\sqrt{-2i x^2 - 2i x - 1} \right) \right) + 4 \operatorname{erf} \left(\frac{\sqrt{2} \sqrt{\pi} \left(i x^2 + i x + \frac{1}{4} \right)}{\sqrt{8 x^2 + 8 x + 2}} \right) \right)}{64}$$

input `integrate(x*cos(1/4+x+x^2)^2,x, algorithm="maxima")`

output `1/64*(32*x^3 + 16*x^2 + 8*x*(-I*e^(2*I*x^2 + 2*I*x + 1/2*I) + I*e^(-2*I*x^2 - 2*I*x - 1/2*I)) + sqrt(8*x^2 + 8*x + 2)*((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(2*I*x^2 + 2*I*x + 1/2*I)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(-2*I*x^2 - 2*I*x - 1/2*I)) - 1)) - 4*I*e^(2*I*x^2 + 2*I*x + 1/2*I) + 4*I*e^(-2*I*x^2 - 2*I*x - 1/2*I))/(2*x + 1)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} x^2 + \left(\frac{1}{32}i + \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) \\ - \left(\frac{1}{32}i - \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right) \\ - \frac{1}{16}i e^{(2ix^2+2ix+\frac{1}{2}i)} + \frac{1}{16}i e^{(-2ix^2-2ix-\frac{1}{2}i)}$$

input `integrate(x*cos(1/4+x+x^2)^2,x, algorithm="giac")`

output $\frac{1}{4}x^2 + \left(\frac{1}{32}i + \frac{1}{32}\right)\sqrt{\pi}\operatorname{erf}\left((i-1)x + \frac{1}{2}i - \frac{1}{2}\right) - \left(\frac{1}{32}i - \frac{1}{32}\right)\sqrt{\pi}\operatorname{erf}\left(-(i+1)x - \frac{1}{2}i - \frac{1}{2}\right) - \frac{1}{16}i e^{(2ix^2+2ix+\frac{1}{2}i)} + \frac{1}{16}i e^{(-2ix^2-2ix-\frac{1}{2}i)}$

Mupad [F(-1)]

Timed out.

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(x*cos(x + x^2 + 1/4)^2,x)`

output `int(x*cos(x + x^2 + 1/4)^2, x)`

Reduce [F]

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int \cos \left(x^2 + x + \frac{1}{4} \right)^2 x dx$$

input `int(x*cos(1/4+x+x^2)^2,x)`

output `int(cos((4*x**2 + 4*x + 1)/4)**2*x,x)`

3.26 $\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (verified)	195
Maple [A] (verified)	196
Fricas [A] (verification not implemented)	196
Sympy [A] (verification not implemented)	196
Maxima [C] (verification not implemented)	197
Giac [C] (verification not implemented)	197
Mupad [F(-1)]	198
Reduce [F]	198

Optimal result

Integrand size = 11, antiderivative size = 27

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x}{2} + \frac{1}{4}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right)$$

output `1/2*x+1/4*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \left(2x + \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) \right)$$

input `Integrate[Cos[1/4 + x + x^2]^2,x]`

output `(2*x + Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/4`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3931, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cos^2 \left(x^2 + x + \frac{1}{4} \right) dx \\ & \quad \downarrow \text{3931} \\ & \int \left(\frac{1}{2} \cos \left(2x^2 + 2x + \frac{1}{2} \right) + \frac{1}{2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{4} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x}{2} \end{aligned}$$

input `Int[Cos[1/4 + x + x^2]^2, x]`

output `x/2 + (Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/4`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3931 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTri
gReduce[Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 1.66 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right)}{4}$	20
risch	$\frac{x}{2} - \frac{\sqrt{\pi} \sqrt{2} (-1)^{\frac{3}{4}} \operatorname{erf}\left(\sqrt{2} (-1)^{\frac{1}{4}} x + \frac{\sqrt{2} (-1)^{\frac{1}{4}}}{2}\right)}{16} + \frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-2i} x - \frac{i}{\sqrt{-2i}}\right)}{8\sqrt{-2i}}$	58

input `int(cos(1/4+x+x^2)^2, x, method=_RETURNVERBOSE)`

output `1/2*x+1/4*Pi^(1/2)*FresnelC((1+2*x)/Pi^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \sqrt{\pi} C \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{1}{2} x$$

input `integrate(cos(1/4+x+x^2)^2, x, algorithm="fricas")`

output `1/4*sqrt(pi)*fresnel_cos((2*x + 1)/sqrt(pi)) + 1/2*x`

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x}{2} + \frac{\sqrt{\pi} C \left(\frac{4x+2}{2\sqrt{\pi}} \right)}{4}$$

input `integrate(cos(1/4+x+x**2)**2, x)`

output $x/2 + \sqrt{\pi} * \text{fresnelc}((4*x + 2)/(2*\sqrt{\pi}))/4$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{1}{16} \sqrt{\pi} \left((i-1) \operatorname{erf} \left(\frac{2ix+i}{\sqrt{2i}} \right) + (i+1) \operatorname{erf} \left(\frac{2ix+i}{\sqrt{-2i}} \right) \right) + \frac{1}{2} x$$

input `integrate(cos(1/4+x+x^2)^2,x, algorithm="maxima")`

output $-1/16*\sqrt{\pi}*((I - 1)*\operatorname{erf}((2*I*x + I)/\sqrt{2*I}) + (I + 1)*\operatorname{erf}((2*I*x + I)/\sqrt{-2*I})) + 1/2*x$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = -\left(\frac{1}{16}i + \frac{1}{16} \right) \sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) + \left(\frac{1}{16}i - \frac{1}{16} \right) \sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right) + \frac{1}{2}x$$

input `integrate(cos(1/4+x+x^2)^2,x, algorithm="giac")`

output $-(1/16*I + 1/16)*\sqrt{\pi}*\operatorname{erf}((I - 1)*x + 1/2*I - 1/2) + (1/16*I - 1/16)*\sqrt{\pi}*\operatorname{erf}(-(I + 1)*x - 1/2*I - 1/2) + 1/2*x$

Mupad [F(-1)]

Timed out.

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(cos(x + x^2 + 1/4)^2,x)`

output `int(cos(x + x^2 + 1/4)^2, x)`

Reduce [F]

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

input `int(cos(1/4+x+x^2)^2,x)`

output `int(cos((4*x**2 + 4*x + 1)/4)**2,x)`

3.27 $\int \frac{\cos^2\left(\frac{1}{4}+x+x^2\right)}{x} dx$

Optimal result	199
Mathematica [N/A]	199
Rubi [N/A]	200
Maple [N/A]	201
Fricas [N/A]	201
Sympy [N/A]	201
Maxima [N/A]	202
Giac [N/A]	202
Mupad [N/A]	203
Reduce [N/A]	203

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \frac{\log(x)}{2} + \frac{1}{2} \text{Int}\left(\frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{x}, x\right)$$

output `1/2*ln(x)+1/2*Defe r (Int)(cos(1/2+2*x+2*x^2)/x,x)`

Mathematica [N/A]

Not integrable

Time = 7.55 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

input `Integrate[$\cos[\frac{1}{4} + x + x^2]^2/x$, x]`

output `Integrate[$\cos[\frac{1}{4} + x + x^2]^2/x$, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(x^2 + x + \frac{1}{4})}{x} dx \\ & \quad \downarrow \text{3949} \\ & \int \left(\frac{\cos(2x^2 + 2x + \frac{1}{2})}{2x} + \frac{1}{2x} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2} \int \frac{\cos(2x^2 + 2x + \frac{1}{2})}{x} dx + \frac{\log(x)}{2} \end{aligned}$$

input `Int[Cos[1/4 + x + x^2]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_)*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)^2}{x} dx$$

input `int(cos(1/4+x+x^2)^2/x,x)`

output `int(cos(1/4+x+x^2)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

input `integrate(cos(1/4+x+x^2)^2/x,x, algorithm="fricas")`

output `integral(cos(x^2 + x + 1/4)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.61 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos^2\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

input `integrate(cos(1/4+x+x**2)**2/x,x)`

output `Integral(cos(x**2 + x + 1/4)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2 \right)}{x} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4} \right)^2}{x} dx$$

input `integrate(cos(1/4+x+x^2)^2/x,x, algorithm="maxima")`

output `1/2*integrate(cos(2*x^2 + 2*x + 1/2)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2 \right)}{x} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4} \right)^2}{x} dx$$

input `integrate(cos(1/4+x+x^2)^2/x,x, algorithm="giac")`

output `integrate(cos(x^2 + x + 1/4)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 42.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

input `int(cos(x + x^2 + 1/4)^2/x,x)`

output `int(cos(x + x^2 + 1/4)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

input `int(cos(1/4+x+x^2)^2/x,x)`

output `int(cos((4*x**2 + 4*x + 1)/4)**2/x,x)`

3.28 $\int \frac{\cos^2\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$

Optimal result	204
Mathematica [N/A]	204
Rubi [N/A]	205
Maple [N/A]	206
Fricas [N/A]	206
Sympy [N/A]	207
Maxima [N/A]	207
Giac [N/A]	207
Mupad [N/A]	208
Reduce [N/A]	208

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = -\frac{1}{2x} - \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} - \sqrt{\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) \\ - \operatorname{Int}\left(\frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{x}, x\right)$$

output
$$-1/2/x-1/2*\cos(1/2+2*x+2*x^2)/x-\text{Pi}^{(1/2)}*\text{FresnelS}((1+2*x)/\text{Pi}^{(1/2)})-\text{Defer}(\text{Int})(\sin(1/2+2*x+2*x^2)/x,x)$$

Mathematica [N/A]

Not integrable

Time = 8.61 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

input
$$\text{Integrate}[\text{Cos}[1/4 + x + x^2]^2/x^2, x]$$

output `Integrate[Cos[1/4 + x + x^2]^2/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(x^2 + x + \frac{1}{4})}{x^2} dx \\
 & \quad \downarrow \text{3949} \\
 & \int \left(\frac{1}{2x^2} + \frac{\cos(2x^2 + 2x + \frac{1}{2})}{2x^2} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & - \int \frac{\sin(2x^2 + 2x + \frac{1}{2})}{x} dx - \sqrt{\pi} \operatorname{FresnelS}\left(\frac{2x+1}{\sqrt{\pi}}\right) - \frac{\cos(2x^2 + 2x + \frac{1}{2})}{2x} - \frac{1}{2x}
 \end{aligned}$$

input `Int[Cos[1/4 + x + x^2]^2/x^2,x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3949 $\text{Int}[\cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_))^{m_.}, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \ \cos[a + b*x + c*x^2]^n, \ x], \ x] /; \ \text{FreeQ}[\{a, \ b, \ c, \ d, \ e, \ m\}, \ x] \ \&& \ \text{IGtQ}[n, \ 1]$

Maple [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cos \left(\frac{1}{4} + x + x^2\right)^2}{x^2} dx$$

input $\text{int}(\cos(1/4+x+x^2)^2/x^2, x)$

output $\text{int}(\cos(1/4+x+x^2)^2/x^2, x)$

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

input $\text{integrate}(\cos(1/4+x+x^2)^2/x^2, x, \ \text{algorithm}=\text{"fricas"})$

output $\text{integral}(\cos(x^2 + x + 1/4)^2/x^2, x)$

Sympy [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos^2\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

input `integrate(cos(1/4+x+x**2)**2/x**2, x)`

output `Integral(cos(x**2 + x + 1/4)**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

input `integrate(cos(1/4+x+x^2)^2/x^2, x, algorithm="maxima")`

output `1/2*(x*integrate(cos(2*x^2 + 2*x + 1/2)/x^2, x) - 1)/x`

Giac [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

input `integrate(cos(1/4+x+x^2)^2/x^2, x, algorithm="giac")`

output `integrate(cos(x^2 + x + 1/4)^2/x^2, x)`

Mupad [N/A]

Not integrable

Time = 42.60 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2 \right)}{x^2} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4} \right)^2}{x^2} dx$$

input `int(cos(x + x^2 + 1/4)^2/x^2,x)`

output `int(cos(x + x^2 + 1/4)^2/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2 \right)}{x^2} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4} \right)^2}{x^2} dx$$

input `int(cos(1/4+x+x^2)^2/x^2,x)`

output `int(cos((4*x**2 + 4*x + 1)/4)**2/x**2,x)`

3.29 $\int (d + ex)^2 \cos(a + bx + cx^2) dx$

Optimal result	209
Mathematica [A] (verified)	210
Rubi [A] (verified)	210
Maple [A] (verified)	214
Fricas [A] (verification not implemented)	214
Sympy [F]	215
Maxima [C] (verification not implemented)	215
Giac [C] (verification not implemented)	216
Mupad [F(-1)]	217
Reduce [F]	217

Optimal result

Integrand size = 19, antiderivative size = 285

$$\begin{aligned} \int (d + ex)^2 \cos(a + bx + cx^2) dx = & \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & - \frac{e^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{e^2 \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\ & - \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\ & + \frac{e(2cd - be) \sin(a + bx + cx^2)}{4c^2} \\ & + \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \end{aligned}$$

output

```
1/8*(-b*e+2*c*d)^2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(5/2)-1/4*e^2*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)-1/4*e^2*2^(1/2)*Pi^(1/2)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)-1/8*(-b*e+2*c*d)^2*2^(1/2)*Pi^(1/2)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(5/2)+1/4*e*(-b*e+2*c*d)*sin(c*x^2+b*x+a)/c^2+1/2*e*(e*x+d)*sin(c*x^2+b*x+a)/c
```

Mathematica [A] (verified)

Time = 1.38 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.66

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx \\ = \frac{\sqrt{2\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left((-2cd + be)^2 \cos\left(a - \frac{b^2}{4c}\right) - 2ce^2 \sin\left(a - \frac{b^2}{4c}\right)\right) - \sqrt{2\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2ce^2 \sin\left(a - \frac{b^2}{4c}\right) + (-2cd + be)^2 \cos\left(a - \frac{b^2}{4c}\right)\right)}{8c^{5/2}}$$

input `Integrate[(d + e*x)^2*Cos[a + b*x + c*x^2], x]`

output
$$\begin{aligned} & (\text{Sqrt}[2*\text{Pi}]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*(((-2*c*d + b*e)^2*\text{Cos}[a - b^2/(4*c)] - 2*c*e^2*\text{Sin}[a - b^2/(4*c)]) - \text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*(2*c*e^2*\text{Cos}[a - b^2/(4*c)] + (-2*c*d + b*e)^2*\text{Sin}[a - b^2/(4*c)]) + 2*\text{Sqrt}[c]*e*(4*c*d - b*e + 2*c*e*x)*\text{Sin}[a + x*(b + c*x)])/(8*c^{(5/2)}) \end{aligned}$$

Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 282, normalized size of antiderivative = 0.99, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3945, 3928, 3832, 3833, 3943, 3929, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d + ex)^2 \cos(a + bx + cx^2) dx \\ & \quad \downarrow \text{3945} \\ & \frac{(2cd - be) \int (d + ex) \cos(cx^2 + bx + a) dx}{2c} - \frac{e^2 \int \sin(cx^2 + bx + a) dx}{2c} + \\ & \quad \downarrow \text{3928} \\ & \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \end{aligned}$$

$$\begin{aligned}
& - \frac{e^2 \left(\sin \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx + \cos \left(a - \frac{b^2}{4c} \right) \int \sin \left(\frac{(b+2cx)^2}{4c} \right) dx \right)}{2c} + \\
& \frac{(2cd - be) \int (d + ex) \cos(cx^2 + bx + a) dx}{2c} + \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3832} \\
& - \frac{e^2 \left(\sin \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right)}{2c} + \\
& \frac{(2cd - be) \int (d + ex) \cos(cx^2 + bx + a) dx}{2c} + \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3833} \\
& \frac{(2cd - be) \int (d + ex) \cos(cx^2 + bx + a) dx}{2c} - \\
& e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right) + \\
& \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3943} \\
& (2cd - be) \left(\frac{(2cd - be) \int \cos(cx^2 + bx + a) dx}{2c} + \frac{e \sin(a + bx + cx^2)}{2c} \right) - \\
& e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right) + \\
& \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3929} \\
& (2cd - be) \left(\frac{(2cd - be) \left(\cos \left(a - \frac{b^2}{4c} \right) \int \cos \left(\frac{(b+2cx)^2}{4c} \right) dx - \sin \left(a - \frac{b^2}{4c} \right) \int \sin \left(\frac{(b+2cx)^2}{4c} \right) dx \right)}{2c} + \frac{e \sin(a + bx + cx^2)}{2c} \right) - \\
& e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \sin \left(a - \frac{b^2}{4c} \right) \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c} \right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}} \right)}{\sqrt{c}} \right) + \\
& \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \\
& \quad \downarrow \text{3832}
\end{aligned}$$

$$\begin{aligned}
 & \frac{(2cd-be) \left(\frac{(2cd-be) \left(\cos(a - \frac{b^2}{4c}) \int \cos(\frac{(b+2cx)^2}{4c}) dx - \frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} \right)}{2c} + \frac{e \sin(a+bx+cx^2)}{2c} \right)}{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelC}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \text{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} \right)} + \\
 & \frac{2c}{e(d+ex) \sin(a+bx+cx^2)} \\
 & \downarrow \text{3833} \\
 & \frac{(2cd-be) \left(\frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \text{FresnelC}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}) - \frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}}}{2c} + \frac{e \sin(a+bx+cx^2)}{2c} \right)}{e^2 \left(\frac{\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelC}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} + \frac{\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \text{FresnelS}(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}})}{\sqrt{c}} \right)} + \\
 & \frac{2c}{e(d+ex) \sin(a+bx+cx^2)}
 \end{aligned}$$

input `Int[(d + e*x)^2*Cos[a + b*x + c*x^2],x]`

output

```

-1/2*(e^2*((Sqrt[Pi/2]*Cos[a - b^2/(4*c)]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqr
rt[2*Pi])])/Sqrt[c] + (Sqrt[Pi/2]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqr
t[2*Pi])]*Sin[a - b^2/(4*c)])/(Sqrt[c])))/c + (e*(d + e*x)*Sin[a + b*x + c*x^2])/(2
*c) + ((2*c*d - b*e)*((2*c*d - b*e)*((Sqrt[Pi/2]*Cos[a - b^2/(4*c)]*Fresn
elC[(b + 2*c*x)/(Sqrt[c]*Sqr
t[2*Pi])])/Sqr
t[c] - (Sqr
t[Pi/2]*FresnelS[(b + 2*c*x)/(Sqr
t[c]*Sqr
t[2*Pi])]*Sin[a - b^2/(4*c)])/(Sqr
t[c])))/(2*c) + (e*Sin
[a + b*x + c*x^2])/(2*c)))/(2*c)

```

Definitions of rubi rules used

rule 3832 $\text{Int}[\text{Sin}[(d_{\cdot})*((e_{\cdot}) + (f_{\cdot})*(x_{\cdot}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3833 $\text{Int}[\text{Cos}[(d_{\cdot})*((e_{\cdot}) + (f_{\cdot})*(x_{\cdot}))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

rule 3928 $\text{Int}[\text{Sin}[(a_{\cdot}) + (b_{\cdot})*(x_{\cdot}) + (c_{\cdot})*(x_{\cdot})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3929 $\text{Int}[\text{Cos}[(a_{\cdot}) + (b_{\cdot})*(x_{\cdot}) + (c_{\cdot})*(x_{\cdot})^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Cos}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] + \text{Simp}[\text{Sin}[(b^2 - 4*a*c)/(4*c)] \quad \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

rule 3943 $\text{Int}[\text{Cos}[(a_{\cdot}) + (b_{\cdot})*(x_{\cdot}) + (c_{\cdot})*(x_{\cdot})^2]*((d_{\cdot}) + (e_{\cdot})*(x_{\cdot})), x_{\text{Symbol}}] \rightarrow \text{Simp}[e*(\text{Sin}[a + b*x + c*x^2]/(2*c)), x] + \text{Simp}[(2*c*d - b*e)/(2*c) \quad \text{Int}[\text{Cos}[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

rule 3945 $\text{Int}[\text{Cos}[(a_{\cdot}) + (b_{\cdot})*(x_{\cdot}) + (c_{\cdot})*(x_{\cdot})^2]*((d_{\cdot}) + (e_{\cdot})*(x_{\cdot}))^{(m_{\cdot})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[e*(d + e*x)^(m - 1)*(\text{Sin}[a + b*x + c*x^2]/(2*c)), x] + (-\text{Simp}[(b*e - 2*c*d)/(2*c) \quad \text{Int}[(d + e*x)^(m - 1)*\text{Cos}[a + b*x + c*x^2], x], x] - \text{Simp}[e^{2*m}/(2*c)]) \quad \text{Int}[(d + e*x)^(m - 2)*\text{Sin}[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[b*e - 2*c*d, 0] \&& \text{GtQ}[m, 1]$

Maple [A] (verified)

Time = 1.98 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.39

method	result
default	$\frac{e^2 b}{2c} \left(\frac{\sin(cx^2 + bx + a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2}{4c} - ac\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2}{4c} - ac\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)\right)}{4c^{\frac{3}{2}}} \right)$
risch	$\frac{e^2 x \sin(cx^2 + bx + a)}{2c} - \frac{\text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}d^2 e^{-\frac{i(4ac-b^2)}{4c}}}{4\sqrt{ic}} + \frac{e^2 \text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}b^2 e^{-\frac{i(4ac-b^2)}{4c}}}{16\sqrt{ic}c^2} - \frac{ie^2 \text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}}{8\sqrt{ic}c} -$
parts	Expression too large to display

input `int((e*x+d)^2*cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/2*e^2/c*x*sin(c*x^2+b*x+a)-1/2*e^2*b/c*(1/2*sin(c*x^2+b*x+a)/c-1/4*b/c^{(3/2)}*2^{(1/2)}*\text{Pi}^{(1/2)}*(\cos((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b))+\sin((1/4*b^2-a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b)))-1/4*e^2/c^{(3/2)}*2^{(1/2)}*\text{Pi}^{(1/2)}*(\cos((1/4*b^2-a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b))-\sin((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b)))+d*e/c*sin(c*x^2+b*x+a)-1/2*d*e*b/c^{(3/2)}*2^{(1/2)}*\text{Pi}^{(1/2)}*(\cos((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b))+\sin((1/4*b^2-a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b)))+1/2*2^{(1/2)}*\text{Pi}^{(1/2)}/c^{(1/2)}*d^{(2)}*(\cos((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b))+\sin((1/4*b^2-a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b))) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.81

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx =$$

$$\frac{\sqrt{2} \left(2 \pi c e^2 \sin\left(-\frac{b^2 - 4ac}{4c}\right) - \pi (4c^2 d^2 - 4bcde + b^2 e^2) \cos\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2} \left(2 \pi c e^2 \cos\left(-\frac{b^2 - 4ac}{4c}\right) + \pi (4c^2 d^2 - 4bcde + b^2 e^2) \sin\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right)}{8c^{\frac{3}{2}}}$$

input `integrate((e*x+d)^2*cos(c*x^2+b*x+a),x, algorithm="fricas")`

output
$$\begin{aligned} & -\frac{1}{8} \sqrt{2} (2\pi c e^2 \sin(-\frac{1}{4}(b^2 - 4ac)/c) - \pi (4c^2 d^2 - 4b c d e + b^2 e^2) \cos(-\frac{1}{4}(b^2 - 4ac)/c)) \sqrt{c/\pi} \operatorname{fresnel_cos}(1/2 \sqrt{c} t(2) (2cx + b) \sqrt{c/\pi}) \\ & + \sqrt{2} (2\pi c e^2 \cos(-\frac{1}{4}(b^2 - 4ac)/c) + \pi (4c^2 d^2 - 4b c d e + b^2 e^2) \sin(-\frac{1}{4}(b^2 - 4ac)/c)) \sqrt{c/\pi} \operatorname{fresnel_sin}(1/2 \sqrt{2} (2cx + b) \sqrt{c/\pi}) - 2(2c^2 e^2 x^2 + 4c^2 d^2 e - b c e^2) \sin(cx^2 + bx + a)) / c^3 \end{aligned}$$

Sympy [F]

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx = \int (d + ex)^2 \cos(a + bx + cx^2) dx$$

input `integrate((e*x+d)**2*cos(c*x**2+b*x+a),x)`

output `Integral((d + e*x)**2*cos(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.06 (sec) , antiderivative size = 2271, normalized size of antiderivative = 7.97

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx = \text{Too large to display}$$

input `integrate((e*x+d)^2*cos(c*x^2+b*x+a),x, algorithm="maxima")`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.02

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx =$$

$$-\frac{i \sqrt{2} \sqrt{\pi} (-4 i c^2 d^2 + 4 i b c d e - i b^2 e^2 + 2 c e^2) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2 x + \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{i b^2 - 4 i a c}{4 c}\right)}}{\left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}} + 2 i \left(c e^2 \left(2 x + \frac{b}{c}\right) + 4 c d e - 2 b d\right)$$

$$-\frac{i \sqrt{2} \sqrt{\pi} (4 i c^2 d^2 - 4 i b c d e + i b^2 e^2 + 2 c e^2) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2 x + \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{-i b^2 + 4 i a c}{4 c}\right)}}{\left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}} - 2 i \left(c e^2 \left(2 x + \frac{b}{c}\right) + 4 c d e - 2 b d\right)$$

$$-\frac{16 c^2}{16 c^2}$$

```
input integrate((e*x+d)^2*cos(c*x^2+b*x+a),x, algorithm="giac")
```

output

$$\begin{aligned} & -\frac{1}{16} \left(I * \sqrt{2} * \sqrt{\pi} * (-4 * I * c^2 * d^2 + 4 * I * b * c * d * e - I * b^2 * e^2 + 2 * c * e^2) \right. \\ & \quad \left. * \operatorname{erf}\left(-\frac{1}{4} * \sqrt{2} * (2 * x + b / c) * (-I * c / \operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)}\right) * e^{-\frac{1}{4} * (I * b^2 - 4 * I * a * c) / c} \right) / ((-I * c / \operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)}) + 2 * I * (c * e^2 * (2 * x + b / c) + 4 * c * d * e - 2 * b * e^2) * e^{-(I * c * x^2 + I * b * x + I * a)} / c^2 - \frac{1}{16} * (-I * \sqrt{2} * \sqrt{\pi} * (4 * I * c^2 * d^2 - 4 * I * b * c * d * e + I * b^2 * e^2 + 2 * c * e^2) * \operatorname{erf}\left(-\frac{1}{4} * \sqrt{2} * (2 * x + b / c) * (I * c / \operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)}\right) * e^{-\frac{1}{4} * (-I * b^2 + 4 * I * a * c) / c}) / ((I * c / \operatorname{abs}(c) + 1) * \sqrt{\operatorname{abs}(c)}) - 2 * I * (c * e^2 * (2 * x + b / c) + 4 * c * d * e - 2 * b * e^2) * e^{-(I * c * x^2 - I * b * x - I * a)} / c^2 \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a) (d + ex)^2 dx$$

input

```
int(cos(a + b*x + c*x^2)*(d + e*x)^2,x)
```

output

```
int(cos(a + b*x + c*x^2)*(d + e*x)^2, x)
```

Reduce [F]

$$\begin{aligned} & \int (d + ex)^2 \cos(a + bx + cx^2) dx \\ &= \frac{6 \left(\int \frac{x^2}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) ce^2 - 6 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) bde + 6 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) cd^2}{3c} \end{aligned}$$

input

```
int((e*x+d)^2*cos(c*x^2+b*x+a),x)
```

output

$$\begin{aligned} & (6 * \operatorname{int}(x^{**2} / (\tan((a + b * x + c * x^{**2}) / 2)^{**2} + 1), x) * c * e^{**2} - 6 * \operatorname{int}(1 / (\tan((a + b * x + c * x^{**2}) / 2)^{**2} + 1), x) * b * d * e + 6 * \operatorname{int}(1 / (\tan((a + b * x + c * x^{**2}) / 2)^{**2} + 1), x) * c * d^{**2} + 3 * \sin(a + b * x + c * x^{**2}) * d * e + 3 * b * d * e * x - 3 * c * d^{**2} * x - c * e^{**2} * x^{**3}) / (3 * c) \end{aligned}$$

3.30 $\int (d + ex) \cos(a + bx + cx^2) dx$

Optimal result	218
Mathematica [A] (verified)	219
Rubi [A] (verified)	219
Maple [A] (verified)	221
Fricas [A] (verification not implemented)	222
Sympy [F]	222
Maxima [C] (verification not implemented)	222
Giac [C] (verification not implemented)	223
Mupad [F(-1)]	224
Reduce [F]	224

Optimal result

Integrand size = 17, antiderivative size = 140

$$\begin{aligned} \int (d + ex) \cos(a + bx + cx^2) dx &= \frac{(2cd - be)\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad - \frac{(2cd - be)\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\ &\quad + \frac{e \sin(a + bx + cx^2)}{2c} \end{aligned}$$

output

```
1/4*(-b*e+2*c*d)*2^(1/2)*Pi^(1/2)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/
c^(1/2)*2^(1/2)/Pi^(1/2))/c^(3/2)-1/4*(-b*e+2*c*d)*2^(1/2)*Pi^(1/2)*Fresne
1S(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)/c^(3/2)+1/2*e*
sin(c*x^2+b*x+a)/c
```

Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.92

$$\int (d + ex) \cos(a + bx + cx^2) dx \\ = \frac{(2cd - be)\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - (2cd - be)\sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) + 2\sqrt{ce}x \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{3/2}}$$

input `Integrate[(d + e*x)*Cos[a + b*x + c*x^2], x]`

output `((2*c*d - b*e)*Sqrt[2*Pi]*Cos[a - b^2/(4*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] - (2*c*d - b*e)*Sqrt[2*Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqr t[2*Pi])]*Sin[a - b^2/(4*c)] + 2*Sqrt[c]*e*Sin[a + x*(b + c*x)])/(4*c^(3/2))`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3943, 3929, 3832, 3833}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) \cos(a + bx + cx^2) dx \\ \downarrow 3943 \\ \frac{(2cd - be) \int \cos(cx^2 + bx + a) dx}{2c} + \frac{e \sin(a + bx + cx^2)}{2c} \\ \downarrow 3929 \\ \frac{(2cd - be) \left(\cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx - \sin\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx \right)}{2c} + \\ \frac{e \sin(a + bx + cx^2)}{2c}$$

$$\begin{aligned}
 & \downarrow \text{3832} \\
 & \frac{(2cd - be) \left(\cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} + \\
 & \quad \downarrow \text{3833} \\
 & \frac{(2cd - be) \left(\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \right)}{2c} + \\
 & \quad \frac{e \sin(a + bx + cx^2)}{2c}
 \end{aligned}$$

input `Int[(d + e*x)*Cos[a + b*x + c*x^2], x]`

output `((2*c*d - b*e)*((Sqrt[Pi/2]*Cos[a - b^2/(4*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/Sqrt[c] - (Sqrt[Pi/2]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a - b^2/(4*c)]/Sqrt[c]))/(2*c) + (e*Sin[a + b*x + c*x^2])/(2*c)`

Definitions of rubi rules used

rule 3832 `Int[Sin[(d_.)*(e_.) + (f_.)*(x_.))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3833 `Int[Cos[(d_.)*(e_.) + (f_.)*(x_.))^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

rule 3929 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Simp[Cos[(b^2 - 4*a*c)/(4*c)] Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Simp[Sin[(b^2 - 4*a*c)/(4*c)] Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

rule 3943

```
Int[Cos[(a_.) + (b_)*(x_) + (c_)*(x_)^2]*((d_.) + (e_)*(x_)), x_Symbol]
:> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Simp[(2*c*d - b*e)/(2*c) Int
[Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b
*e, 0]
```

Maple [A] (verified)

Time = 1.24 (sec), antiderivative size = 182, normalized size of antiderivative = 1.30

method	result
default	$\frac{e \sin(cx^2+bx+a)}{2c} - \frac{eb\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2-ac}{4c}\right) \text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2-ac}{4c}\right) \text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} + \frac{\sqrt{2}\sqrt{\pi}d\left(cx^2+bx+a\right)^{\frac{3}{2}}}{2c^{\frac{1}{2}}}$
risch	$\frac{\text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}de^{-\frac{i(4ac-b^2)}{4c}}}{4\sqrt{ic}} - \frac{e\text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}be^{-\frac{i(4ac-b^2)}{4c}}}{8\sqrt{ic}c} - \frac{\text{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)\sqrt{\pi}de^{\frac{i(4ac-b^2)}{4c}}}{4\sqrt{-ic}} + \frac{e\text{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)\sqrt{\pi}be^{\frac{i(4ac-b^2)}{4c}}}{8\sqrt{-ic}}$
parts	$\frac{\sqrt{2}\sqrt{\pi}\cos\left(\frac{b^2-ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)ex}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi}\sin\left(\frac{b^2-ac}{4c}\right)\text{FresnelS}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)ex}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi}\cos\left(\frac{b^2-ac}{4c}\right)\text{FresnelC}\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)ex}{2\sqrt{c}}$

input `int((e*x+d)*cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

output

```
1/2*e*sin(c*x^2+b*x+a)/c-1/4*e*b/c^(3/2)*2^(1/2)*Pi^(1/2)*(cos((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))+sin((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))+1/2*2^(1/2)*Pi^(1/2)/c^(1/2)*d*(cos((1/4*b^2-a*c)/c)*FresnelC(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b))+sin((1/4*b^2-a*c)/c)*FresnelS(2^(1/2)/Pi^(1/2)/c^(1/2)*(c*x+1/2*b)))
```

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.97

$$\int (d + ex) \cos(a + bx + cx^2) dx = \frac{\sqrt{2}\pi(2cd - be)\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi(2cd - be)\sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2 - 4ac}{4c}\right)}{4c^2}$$

input `integrate((e*x+d)*cos(c*x^2+b*x+a),x, algorithm="fricas")`

output `1/4*(sqrt(2)*pi*(2*c*d - b*e)*sqrt(c/pi)*cos(-1/4*(b^2 - 4*a*c)/c)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) - sqrt(2)*pi*(2*c*d - b*e)*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c)*sin(-1/4*(b^2 - 4*a*c)/c) + 2*c*e*sin(c*x^2 + b*x + a))/c^2`

Sympy [F]

$$\int (d + ex) \cos(a + bx + cx^2) dx = \int (d + ex) \cos(a + bx + cx^2) dx$$

input `integrate((e*x+d)*cos(c*x**2+b*x+a),x)`

output `Integral((d + e*x)*cos(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 695, normalized size of antiderivative = 4.96

$$\int (d + ex) \cos(a + bx + cx^2) dx = \text{Too large to display}$$

```
input integrate((e*x+d)*cos(c*x^2+b*x+a),x, algorithm="maxima")
```

```
output -1/8*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(I*c)) + ((I + 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(-I*c)))*d/sqrt(c) + 1/16*((((I - 1)*sqrt(2)*sqrt(pi))*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/4*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/4*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/4*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/4*(b^2 - 4*a*c)/c))*x - 4*(c*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^2 - 4*a*c)/c) - c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/4*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.44

$$\int (d + ex) \cos(a + bx + cx^2) dx$$

$$= -\frac{i \sqrt{2} \sqrt{\pi} (2i cd - i be) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2x + \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{i b^2 - 4i ac}{4c}\right)}}{\left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}} + 2i ee^{(icx^2 + ibx + ia)}$$

$$- \frac{8c}{i \sqrt{2} \sqrt{\pi} (-2i cd + i be) \operatorname{erf}\left(-\frac{1}{4} \sqrt{2} \left(2x + \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{-i b^2 + 4i ac}{4c}\right)}} - 2i ee^{(-icx^2 - ibx - ia)}$$

input `integrate((e*x+d)*cos(c*x^2+b*x+a),x, algorithm="giac")`

output
$$\begin{aligned} & -\frac{1}{8}(-I*\sqrt{2}*\sqrt{\pi}*(2*I*c*d - I*b*e)*erf(-1/4*\sqrt{2}*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^{(-1/4*(I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c)))} + 2*I*e^*e^{(I*c*x^2 + I*b*x + I*a)/c} - 1/8*(I*\sqrt{2}*\sqrt{\pi}*(-2*I*c*d + I*b*e)*erf(-1/4*\sqrt{2}*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^{(-1/4*(-I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c)))} - 2*I*e^*e^{(-I*c*x^2 - I*b*x - I*a)/c}) \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int (d + ex) \cos(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a) (d + ex) dx$$

input `int(cos(a + b*x + c*x^2)*(d + e*x),x)`

output `int(cos(a + b*x + c*x^2)*(d + e*x), x)`

Reduce [F]

$$\begin{aligned} & \int (d + ex) \cos(a + bx + cx^2) dx \\ &= \frac{-2 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) be + 4 \left(\int \frac{1}{\tan(\frac{1}{2}cx^2 + \frac{1}{2}bx + \frac{1}{2}a)^2 + 1} dx \right) cd + \sin(cx^2 + bx + a)e + bex - 2cdx}{2c} \end{aligned}$$

input `int((e*x+d)*cos(c*x^2+b*x+a),x)`

output
$$(-2*\int(1/(\tan((a + b*x + c*x**2)/2)**2 + 1),x)*b*e + 4*\int(1/(\tan((a + b*x + c*x**2)/2)**2 + 1),x)*c*d + \sin(a + b*x + c*x**2)*e + b*e*x - 2*c*d*x)/(2*c)$$

3.31 $\int \frac{\cos(a+bx+cx^2)}{d+ex} dx$

Optimal result	225
Mathematica [N/A]	225
Rubi [N/A]	226
Maple [N/A]	226
Fricas [N/A]	227
Sympy [N/A]	227
Maxima [N/A]	228
Giac [N/A]	228
Mupad [N/A]	228
Reduce [N/A]	229

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \text{Int}\left(\frac{\cos(a + bx + cx^2)}{d + ex}, x\right)$$

output `Defer(Int)(cos(c*x^2+b*x+a)/(e*x+d),x)`

Mathematica [N/A]

Not integrable

Time = 3.70 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

input `Integrate[Cos[a + b*x + c*x^2]/(d + e*x),x]`

output `Integrate[Cos[a + b*x + c*x^2]/(d + e*x), x]`

Rubi [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3951}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

↓ 3951

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

input `Int[Cos[a + b*x + c*x^2]/(d + e*x), x]`

output `$Aborted`

Definitions of rubi rules used

rule 3951 `Int[Cos[(a_) + (b_)*(x_) + (c_)*(x_)^2]^(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] :> Unintegrable[(d + e*x)^m*Cos[a + b*x + c*x^2]^n, x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.46 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

input `int(cos(c*x^2+b*x+a)/(e*x+d), x)`

output `int(cos(c*x^2+b*x+a)/(e*x+d),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

input `integrate(cos(c*x^2+b*x+a)/(e*x+d),x, algorithm="fricas")`

output `integral(cos(c*x^2 + b*x + a)/(e*x + d), x)`

Sympy [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

input `integrate(cos(c*x**2+b*x+a)/(e*x+d),x)`

output `Integral(cos(a + b*x + c*x**2)/(d + e*x), x)`

Maxima [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

input `integrate(cos(c*x^2+b*x+a)/(e*x+d),x, algorithm="maxima")`

output `integrate(cos(c*x^2 + b*x + a)/(e*x + d), x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

input `integrate(cos(c*x^2+b*x+a)/(e*x+d),x, algorithm="giac")`

output `integrate(cos(c*x^2 + b*x + a)/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 42.52 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{d + ex} dx$$

input `int(cos(a + b*x + c*x^2)/(d + e*x),x)`

output `int(cos(a + b*x + c*x^2)/(d + e*x), x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

input `int(cos(c*x^2+b*x+a)/(e*x+d),x)`

output `int(cos(a + b*x + c*x**2)/(d + e*x),x)`

3.32 $\int (d + ex)^2 \cos^2(a + bx + cx^2) dx$

Optimal result	230
Mathematica [A] (verified)	231
Rubi [A] (verified)	231
Maple [A] (verified)	233
Fricas [A] (verification not implemented)	233
Sympy [F]	234
Maxima [C] (verification not implemented)	234
Giac [C] (verification not implemented)	235
Mupad [F(-1)]	236
Reduce [F]	236

Optimal result

Integrand size = 21, antiderivative size = 291

$$\begin{aligned} \int (d + ex)^2 \cos^2(a + bx + cx^2) dx &= \frac{(d + ex)^3}{6e} \\ &+ \frac{(2cd - be)^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ &- \frac{e^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ &- \frac{e^2 \sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\ &- \frac{(2cd - be)^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\ &+ \frac{e(2cd - be) \sin(2a + 2bx + 2cx^2)}{16c^2} \\ &+ \frac{e(d + ex) \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output

```
1/6*(e*x+d)^3/e+1/16*(-b*e+2*c*d)^2*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(5/2)-1/16*e^2*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)-1/16*e^2*Pi^(1/2)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)-1/16*(-b*e+2*c*d)^2*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(5/2)+1/16*e*(-b*e+2*c*d)*sin(2*c*x^2+2*b*x+2*a)/c^2+1/8*e*(e*x+d)*sin(2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.74

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx \\ = \frac{3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left((-2cd + be)^2 \cos\left(2a - \frac{b^2}{2c}\right) - ce^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) - 3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(ce^2 \cos\left(2a - \frac{b^2}{2c}\right) - (-2cd + be)^2 \sin\left(2a - \frac{b^2}{2c}\right)\right)}{48}$$

input

```
Integrate[(d + e*x)^2*Cos[a + b*x + c*x^2]^2, x]
```

output

```
(3*.Sqrt[Pi]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*((-2*c*d + b*e)^2*Cos[2*a - b^2/(2*c)] - c*e^2*Sin[2*a - b^2/(2*c)]) - 3*.Sqrt[Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*(c*e^2*Cos[2*a - b^2/(2*c)] + (-2*c*d + b*e)^2*Sin[2*a - b^2/(2*c)]) + Sqrt[c]*(8*c^2*x*(3*d^2 + 3*d*e*x + e^2*x^2) + 3*e*(4*c*d - b*e + 2*c*e*x)*Sin[2*(a + x*(b + c*x))]))/(48*c^(5/2))
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx$$

$$\begin{aligned}
 & \downarrow \textcolor{blue}{3949} \\
 & \int \left(\frac{1}{2}(d+ex)^2 \cos(2a+2bx+2cx^2) + \frac{1}{2}(d+ex)^2 \right) dx \\
 & \downarrow \textcolor{blue}{2009} \\
 & \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) (2cd-be)^2 \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) (2cd-be)^2 \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)} - \\
 & \frac{16c^{5/2}}{\sqrt{\pi}e^2 \cos\left(2a - \frac{b^2}{2c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)} + \frac{e(2cd-be) \sin(2a+2bx+2cx^2)}{16c^{3/2}} + \\
 & \frac{16c^{3/2}}{e(d+ex) \sin(2a+2bx+2cx^2)} + \frac{(d+ex)^3}{6e}
 \end{aligned}$$

input `Int[(d + e*x)^2*Cos[a + b*x + c*x^2]^2, x]`

output `(d + e*x)^3/(6*e) + ((2*c*d - b*e)^2*Sqrt[Pi]*Cos[2*a - b^2/(2*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])])/(16*c^(5/2)) - (e^2*Sqrt[Pi]*Cos[2*a - b^2/(2*c)]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])])/(16*c^(3/2)) - (e^2*Sqrt[Pi]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*Sin[2*a - b^2/(2*c)])/(16*c^(3/2)) - ((2*c*d - b*e)^2*Sqrt[Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*Sin[2*a - b^2/(2*c)])/(16*c^(5/2)) + (e*(2*c*d - b*e)*Sin[2*a + 2*b*x + 2*c*x^2])/(16*c^2) + (e*(d + e*x)*Sin[2*a + 2*b*x + 2*c*x^2])/(8*c)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n_*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.30

method	result
default	$\frac{e^2 x \sin(2cx^2+2bx+2a)}{8c} - \frac{e^2 b \left(\frac{\sin(2cx^2+2bx+2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^{\frac{3}{2}}} \right)}$
risch	$\frac{de x^2}{2} + \frac{d^2 x}{2} + \frac{e^2 x^3}{6} + \frac{\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}d^2 e^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}} + \frac{e^2 b^2 \sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}}\sqrt{2}\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c^2\sqrt{ic}} - \frac{ie^2 b^2 \sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}}\sqrt{2}\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c^2\sqrt{ic}}$

input `int((e*x+d)^2*cos(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/8*e^2/c*x*sin(2*c*x^2+2*b*x+2*a)-1/4*e^2*b/c*(1/4*sin(2*c*x^2+2*b*x+2*a) \\ & /c-1/4*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))+sin(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))) \\ &)-1/16*e^2/c^(3/2)*Pi^(1/2)*(cos(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))-sin(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))) \\ & +1/4*d*e/c*sin(2*c*x^2+2*b*x+2*a)-1/4*d*e*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))+sin(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2)))+1/4*Pi^(1/2)/c^(1/2)*d^2*(cos(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))+sin(1/2*(-4*a*c+b^2)/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2)))+1/2*d*e*x^2+1/2*d^2*x+1/6*e^2*x^3 \end{aligned}$$
Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.88

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx$$

$$= \frac{8 c^3 e^2 x^3 + 24 c^3 d e x^2 + 24 c^3 d^2 x + 6 (2 c^2 e^2 x + 4 c^2 d e - b c e^2) \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) - 3 (2 c^2 e^2 x^2 + 6 c^2 d e x + 6 c^2 d^2 - b c e^2) \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) + 3 (8 c^3 e^2 x^3 + 24 c^3 d e x^2 + 24 c^3 d^2 x + 6 (2 c^2 e^2 x + 4 c^2 d e - b c e^2) \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) - 3 (2 c^2 e^2 x^2 + 6 c^2 d e x + 6 c^2 d^2 - b c e^2) \cos(cx^2 + bx + a) \sin(cx^2 + bx + a))}{16 c^3}$$

input `integrate((e*x+d)^2*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

output

```
1/48*(8*c^3*e^2*x^3 + 24*c^3*d*e*x^2 + 24*c^3*d^2*x + 6*(2*c^2*e^2*x + 4*c^2*d*e - b*c*e^2)*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a) - 3*(pi*c*e^2*sin(-1/2*(b^2 - 4*a*c)/c) - pi*(4*c^2*d^2 - 4*b*c*d*e + b^2*e^2)*cos(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - 3*(pi*c*e^2*cos(-1/2*(b^2 - 4*a*c)/c) + pi*(4*c^2*d^2 - 4*b*c*d*e + b^2*e^2)*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c))/c^3
```

Sympy [F]

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \int (d + ex)^2 \cos^2(a + bx + cx^2) dx$$

input

```
integrate((e*x+d)**2*cos(c*x**2+b*x+a)**2,x)
```

output

```
Integral((d + e*x)**2*cos(a + b*x + c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.06 (sec), antiderivative size = 2360, normalized size of antiderivative = 8.11

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \text{Too large to display}$$

input

```
integrate((e*x+d)^2*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

output

```

-1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(2*I*c)) + ((I + 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(-2*I*c)))*c^(3/2) - 16*c^2*x)*d^2/c^2 + 1/32*sqrt(2)*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) + c*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) + e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*d*e/(c^2*sqrt((4*...

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.01

$$\begin{aligned}
 \int (d + ex)^2 \cos^2(a + bx + cx^2) dx &= \frac{1}{6} e^2 x^3 + \frac{1}{2} d e x^2 + \frac{1}{2} d^2 x \\
 &\quad - \frac{i (ce^2(2x + \frac{b}{c}) + 4cde - 2be^2)e^{(2icx^2 + 2ibx + 2ia)} + \frac{i\sqrt{\pi}(-4ic^2d^2 + 4ibcde - ib^2e^2 + ce^2)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{-2icx^2 - 2ibx - 2ia}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)} }{32c^2} \\
 &\quad - \frac{-i (ce^2(2x + \frac{b}{c}) + 4cde - 2be^2)e^{(-2icx^2 - 2ibx - 2ia)} - \frac{i\sqrt{\pi}(4ic^2d^2 - 4ibcde + ib^2e^2 + ce^2)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right)e^{-2icx^2 + 2ibx + 2ia}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2}
 \end{aligned}$$

input `integrate((e*x+d)^2*cos(c*x^2+b*x+a)^2,x, algorithm="giac")`

output

```
1/6*e^2*x^3 + 1/2*d*e*x^2 + 1/2*d^2*x - 1/32*(I*(c*e^2*(2*x + b/c) + 4*c*d
*e - 2*b*e^2)*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a) + I*sqrt(pi)*(-4*I*c^2*d^2 +
4*I*b*c*d*e - I*b^2*e^2 + c*e^2)*erf(-1/2*sqrt(c)*(2*x + b/c)*(-I*c/abs(c
) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)))/c^2 - 1/
32*(-I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^(-2*I*c*x^2 - 2*I*b*x - 2
*I*a) - I*sqrt(pi)*(4*I*c^2*d^2 - 4*I*b*c*d*e + I*b^2*e^2 + c*e^2)*erf(-1/
2*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqr
t(c)*(I*c/abs(c) + 1)))/c^2
```

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 (d + ex)^2 dx$$

input

```
int(cos(a + b*x + c*x^2)^2*(d + e*x)^2, x)
```

output

```
int(cos(a + b*x + c*x^2)^2*(d + e*x)^2, x)
```

Reduce [F]

$$\begin{aligned} & \int (d + ex)^2 \cos^2(a + bx + cx^2) dx \\ &= \frac{-\cos(cx^2 + bx + a) \sin(cx^2 + bx + a) b e^2 + 3 \cos(cx^2 + bx + a) \sin(cx^2 + bx + a) c d e + 2 \cos(cx^2 + bx + a) c^2 d^2 e^2}{b^3} \end{aligned}$$

input

```
int((e*x+d)^2*cos(c*x^2+b*x+a)^2, x)
```

output

```
( - cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*b*e**2 + 3*cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*c*d*e + 2*cos(a + b*x + c*x**2)*sin(a + b*x + c*x**2)*c*e**2*x - 6*int(tan((a + b*x + c*x**2)/2)**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2*e**2 + 24*int(tan((a + b*x + c*x**2)/2)**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b*c*d*e - 24*int(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**2*d**2 - 8*int(x**2/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c**2*e**2 - 8*int(tan((a + b*x + c*x**2)/2)/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*c*e**2 + 2*int(1/(tan((a + b*x + c*x**2)/2)**4 + 2*tan((a + b*x + c*x**2)/2)**2 + 1),x)*b**2*e**2 - sin(a + b*x + c*x**2)*b*e**2 + 2*sin(a + b*x + c*x**2)*c*e**2*x - 3*b*c*d*e*x + 6*c**2*d**2*x + 3*c**2*d*e*x**2 + 2*c**2*e**2*x**3)/(6*c**2)
```

3.33 $\int (d + ex) \cos^2(a + bx + cx^2) dx$

Optimal result	238
Mathematica [A] (verified)	239
Rubi [A] (verified)	239
Maple [A] (verified)	240
Fricas [A] (verification not implemented)	241
Sympy [F]	241
Maxima [C] (verification not implemented)	241
Giac [C] (verification not implemented)	242
Mupad [F(-1)]	243
Reduce [F]	243

Optimal result

Integrand size = 19, antiderivative size = 150

$$\begin{aligned} \int (d + ex) \cos^2(a + bx + cx^2) dx &= \frac{(d + ex)^2}{4e} \\ &+ \frac{(2cd - be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ &- \frac{(2cd - be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} \\ &+ \frac{e \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

output

```
1/4*(e*x+d)^2/e+1/8*(-b*e+2*c*d)*Pi^(1/2)*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))/c^(3/2)-1/8*(-b*e+2*c*d)*Pi^(1/2)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)/c^(3/2)+1/8*e*sin(2*c*x^2+2*b*x+2*a)/c
```

Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.93

$$\int (d + ex) \cos^2(a + bx + cx^2) dx \\ = \frac{(2cd - be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - (2cd - be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 + b^2x^2 + d^2)}{8c^{3/2}}$$

input `Integrate[(d + e*x)*Cos[a + b*x + c*x^2]^2, x]`

output $((2*c*d - b*e)*\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a - b^2/(2*c)]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])] - (2*c*d - b*e)*\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a - b^2/(2*c)] + \text{Sqrt}[c]*(2*c*x*(2*d + e*x) + e*\text{Sin}[2*(a + x*(b + c*x))]))/(8*c^{(3/2)})$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex) \cos^2(a + bx + cx^2) dx \\ \downarrow 3949 \\ \int \left(\frac{1}{2}(d + ex) \cos(2a + 2bx + 2cx^2) + \frac{1}{2}(d + ex) \right) dx \\ \downarrow 2009 \\ \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) (2cd - be) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} - \\ \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) (2cd - be) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} + \frac{e \sin(2a + 2bx + 2cx^2)}{8c} + \frac{(d + ex)^2}{4e}$$

input $\text{Int}[(d + e*x)*\cos[a + b*x + c*x^2]^2, x]$

output $(d + e*x)^2/(4*e) + ((2*c*d - b*e)*\sqrt{\pi}*\cos[2*a - b^2/(2*c)]*\text{FresnelC}[(b + 2*c*x)/(\sqrt{c}*\sqrt{\pi})])/(8*c^{(3/2)}) - ((2*c*d - b*e)*\sqrt{\pi}*\text{FresnelS}[(b + 2*c*x)/(\sqrt{c}*\sqrt{\pi})]*\sin[2*a - b^2/(2*c)])/(8*c^{(3/2)}) + (e*\sin[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3949 $\text{Int}[\cos(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^(n_*)*((d_.) + (e_.)*(x_.))^(m_), x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \cos[a + b*x + c*x^2]^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{IGtQ}[n, 1]$

Maple [A] (verified)

Time = 1.80 (sec), antiderivative size = 170, normalized size of antiderivative = 1.13

method	result
default	$\frac{e \sin(2cx^2+2bx+2a)}{8c} - \frac{eb\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{8c^{\frac{3}{2}}} + \frac{\sqrt{\pi} d \left(\cos\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelC}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) \text{FresnelS}\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{8c^{\frac{5}{2}}}$
risch	$\frac{ex^2}{4} + \frac{dx}{2} + \frac{\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}de^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}} - \frac{e\text{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}be^{-\frac{i(4ac-b^2)}{2c}}}{32\sqrt{ic}c} - \frac{\text{erf}\left(-\sqrt{-2ic}x - \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}de^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}}$

input $\text{int}((e*x+d)*\cos(c*x^2+b*x+a)^2, x, \text{method}=\text{_RETURNVERBOSE})$

output
$$\begin{aligned} & 1/8*e*\sin(2*c*x^2+2*b*x+2*a)/c-1/8*e*b/c^{(3/2)}*\pi^{(1/2)}*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\pi^{(1/2)})+\sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\pi^{(1/2)}))+1/4*\pi^{(1/2)}/c^{(1/2)}*d*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\pi^{(1/2)})+\sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\pi^{(1/2)}))+1/2*d*x+1/4*e*x^2 \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.99

$$\int (d + ex) \cos^2(a + bx + cx^2) dx$$

$$= \frac{2c^2ex^2 + \pi(2cd - be)\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi(2cd - be)\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2 - 4ac}{2c}\right)}{8c^2}$$

input `integrate((e*x+d)*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

output `1/8*(2*c^2*e*x^2 + pi*(2*c*d - b*e)*sqrt(c/pi)*cos(-1/2*(b^2 - 4*a*c)/c)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - pi*(2*c*d - b*e)*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c)*sin(-1/2*(b^2 - 4*a*c)/c) + 4*c^2*d*x + 2*c*e*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a))/c^2`

Sympy [F]

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \int (d + ex) \cos^2(a + bx + cx^2) dx$$

input `integrate((e*x+d)*cos(c*x**2+b*x+a)**2,x)`

output `Integral((d + e*x)*cos(a + b*x + c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 737, normalized size of antiderivative = 4.91

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \text{Too large to display}$$

```
input integrate((e*x+d)*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")
```

```
output -1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(2*I*c)) + ((I + 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(-2*I*c)))*c^(3/2) - 16*c^2*x)*d/c^2 + 1/64*sqrt(2)*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) + c*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c)))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.30

$$\int (d + ex) \cos^2(a + bx + cx^2) dx$$

$$= \frac{1}{4} ex^2 + \frac{1}{2} dx - \frac{i ee^{(2icx^2+2ibx+2ia)} - \frac{i\sqrt{\pi}(2icd-ibe)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{ib^2-4ic}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|}+1\right)}}{16c}$$

$$- \frac{-i ee^{(-2icx^2-2ibx-2ia)} + \frac{i\sqrt{\pi}(-2icd+ibe)\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{-ib^2+4ic}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|}+1\right)}}{16c}$$

input `integrate((e*x+d)*cos(c*x^2+b*x+a)^2,x, algorithm="giac")`

output
$$\frac{1}{4}e x^2 + \frac{1}{2}d x - \frac{1}{16}(\text{I} e e^{(2\text{I} c x^2 + 2\text{I} b x + 2\text{I} a)} - \text{I} \sqrt{\text{pi}} (2\text{I} c d - \text{I} b e) \operatorname{erf}(-\frac{1}{2}\sqrt{c} (2x + b/c) (-\text{I} c / \text{abs}(c) + 1)) e^{-\frac{1}{2}(\text{I} b^2 - 4\text{I} a c)/c} / (\sqrt{c} (-\text{I} c / \text{abs}(c) + 1))) / c - \frac{1}{16}(-\text{I} e e^{(-2\text{I} c x^2 - 2\text{I} b x - 2\text{I} a)} + \text{I} \sqrt{\text{pi}} (-2\text{I} c d + \text{I} b e) \operatorname{erf}(-\frac{1}{2}\sqrt{c} (2x + b/c) (\text{I} c / \text{abs}(c) + 1)) e^{-\frac{1}{2}(-\text{I} b^2 + 4\text{I} a c)/c} / (\sqrt{c} (\text{I} c / \text{abs}(c) + 1))) / c$$

Mupad [F(-1)]

Timed out.

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 (d + ex) dx$$

input `int(cos(a + b*x + c*x^2)^2*(d + e*x),x)`

output `int(cos(a + b*x + c*x^2)^2*(d + e*x), x)`

Reduce [F]

$$\begin{aligned} \int (d + ex) \cos^2(a + bx + cx^2) dx &= \left(\int \cos(cx^2 + bx + a)^2 dx \right) d \\ &\quad + \left(\int \cos(cx^2 + bx + a)^2 x dx \right) e \end{aligned}$$

input `int((e*x+d)*cos(c*x^2+b*x+a)^2,x)`

output `int(cos(a + b*x + c*x**2)**2,x)*d + int(cos(a + b*x + c*x**2)**2*x,x)*e`

3.34 $\int \frac{\cos^2(a+bx+cx^2)}{d+ex} dx$

Optimal result	244
Mathematica [N/A]	244
Rubi [N/A]	245
Maple [N/A]	246
Fricas [N/A]	246
Sympy [N/A]	246
Maxima [N/A]	247
Giac [N/A]	247
Mupad [N/A]	248
Reduce [N/A]	248

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \frac{\log(d + ex)}{2e} + \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx + 2cx^2)}{d + ex}, x\right)$$

output `1/2*ln(e*x+d)/e+1/2*DefeR(Int)(cos(2*c*x^2+2*b*x+2*a)/(e*x+d),x)`

Mathematica [N/A]

Not integrable

Time = 3.92 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx$$

input `Integrate[Cos[a + b*x + c*x^2]^2/(d + e*x), x]`

output `Integrate[Cos[a + b*x + c*x^2]^2/(d + e*x), x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3949, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx \\ & \quad \downarrow \textcolor{blue}{3949} \\ & \int \left(\frac{\cos(2a + 2bx + 2cx^2)}{2(d + ex)} + \frac{1}{2(d + ex)} \right) dx \\ & \quad \downarrow \textcolor{blue}{2009} \\ & \frac{1}{2} \int \frac{\cos(2cx^2 + 2bx + 2a)}{d + ex} dx + \frac{\log(d + ex)}{2e} \end{aligned}$$

input `Int[Cos[a + b*x + c*x^2]^2/(d + e*x), x]`

output `$Aborted`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3949 `Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^(n_)*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]`

Maple [N/A]

Not integrable

Time = 0.58 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

input `int(cos(c*x^2+b*x+a)^2/(e*x+d),x)`

output `int(cos(c*x^2+b*x+a)^2/(e*x+d),x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

input `integrate(cos(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="fricas")`

output `integral(cos(c*x^2 + b*x + a)^2/(e*x + d), x)`

Sympy [N/A]

Not integrable

Time = 4.85 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx$$

input `integrate(cos(c*x**2+b*x+a)**2/(e*x+d),x)`

output `Integral(cos(a + b*x + c*x**2)**2/(d + e*x), x)`

Maxima [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 120, normalized size of antiderivative = 5.71

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

input `integrate(cos(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="maxima")`

output `-1/2*(2*e*integrate(-1/4*(cos(2*c*x^2 + 2*b*x)*cos(2*a) - sin(2*c*x^2 + 2*b*x)*sin(2*a))/((cos(2*a)^2 + sin(2*a)^2)*e*x + (cos(2*a)^2 + sin(2*a)^2)*d), x) - 2*e*integrate(1/4*cos(2*c*x^2 + 2*b*x + 2*a)/(e*x + d), x) - log(e*x + d))/e`

Giac [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

input `integrate(cos(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="giac")`

output `integrate(cos(c*x^2 + b*x + a)^2/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 42.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{d + ex} dx$$

input `int(cos(a + b*x + c*x^2)^2/(d + e*x),x)`

output `int(cos(a + b*x + c*x^2)^2/(d + e*x), x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

input `int(cos(c*x^2+b*x+a)^2/(e*x+d),x)`

output `int(cos(a + b*x + c*x**2)**2/(d + e*x),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	249
4.2 Links to plain text integration problems used in this report for each CAS .	267

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```
(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*) (*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","");
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
    ,(*ELSE*) (*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "}
      ,
      finalresult={"C","Result contains higher order function than in optimal. Order "}
    ]
  ]
]
]
```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```



```

ExpnType[expn_] :=
If[AtomQ[expn],
  1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn] === Power,
      If[IntegerQ[expn[[2]]],
        ExpnType[expn[[1]]],
        If[Head[expn[[2]]] === Rational,
          If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
            1,
            Max[ExpnType[expn[[1]]], 2]],
        Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
      If[Head[expn] === Plus || Head[expn] === Times,
        Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
        If[ElementaryFunctionQ[Head[expn]],
          Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]]]]]]
  
```

```
Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],  
If[AppellFunctionQ[Head[expn]],  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],  
If[Head[expn]==RootSum,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],  
If[Head[expn]==Integrate || Head[expn]==Int,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],  
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=  
MemberQ[{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
} , func]
```

```
SpecialFunctionQ[func_] :=  
MemberQ[{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  

```

```
HypergeometricFunctionQ[func_] :=  
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=  
MemberQ[{AppellF1}, func]
```

Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
```

```
#      antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
            # this assumes optimal do not as well. No check is needed here.
            if debug then
                print("result do not contain complex, this assumes optimal do not as well");
            fi;
```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                           convert(leaf_count_result,string),\"$ vs. \$2(", 
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_co
                           fi;
            fi;
        else #ExpnType(result) > ExpnType(optimal)
            if debug then
                print("ExpnType(result) > ExpnType(optimal)");
            fi;
            return "C",cat("Result contains higher order function than in optimal. Order ",
                           convert(ExpnType_result,string)," vs. order ",
                           convert(ExpnType_optimal,string),"."));
        fi;

    end proc:

    #

    # is_contains_complex(result)
    # takes expressions and returns true if it contains "I" else false
    #
    #Nasser 032417
    is_contains_complex:= proc(expression)
        return (has(expression,I));
    end proc:

    # The following summarizes the type number assigned an expression
    # based on the functions it involves
    # 1 = rational function
    # 2 = algebraic function
    # 3 = elementary function
    # 4 = special function
    # 5 = hypergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'`^`') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then
            if type(op(1,expn),'rational') then
                1
            else
                max(2,ExpnType(op(1,expn)))
            end if
        else
            max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
        end if
    elif type(expn,'`+`') or type(expn,'`*`') then
        max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif ElementaryFunctionQ(op(0,expn)) then
        max(3,ExpnType(op(1,expn)))
    elif SpecialFunctionQ(op(0,expn)) then
        max(4,apply(max,map(ExpnType,[op(expn)])))
    elif HypergeometricFunctionQ(op(0,expn)) then
        max(5,apply(max,map(ExpnType,[op(expn)])))
    elif AppellFunctionQ(op(0,expn)) then
        max(6,apply(max,map(ExpnType,[op(expn)])))
    elif op(0,expn)='int' then
        max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
```

```
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arccsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]
```

```
def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*')
```

```

m1 = expnType(expn.args[0])
m2 = expnType(list(expn.args[1:]))
return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sagemode")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(grade)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```
from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                       ]
    if debug:
```

```

if m:
    print ("func ", func , " is elementary_function")
else:
    print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral'
'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
'elliptic_pi','exp_integral_e','log_integral',
'veierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
'veierstrassPPrime','weierstrassSigma']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):
    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0], Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1], Integer)

```

```

    return expnType(expn.operands()[0])  #expnType(expn.args[0])
elif type(expn.operands()[1]) == Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0]) == Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))  #max(3,expnType(expn.args[0]),expnType(expn.args[1]))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(4,m1)  #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(5,m1)  #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation ="none"
            else:
                grade = "B"
                grade_annotation ="Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation ="Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation ="none"
        else:
            grade = "B"
            grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(expnType_result)+"/"+str(expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file