

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.2-Cosine/211-4.2.7.2

Nasser M. Abbasi

May 18, 2024 Compiled on May 18, 2024 at 12:28 Noon

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	43
3	Listing of integrals	46
3.1	$\int \frac{\sin^6(x)}{a-a \cos^2(x)} dx$	48
3.2	$\int \frac{\sin^5(x)}{a-a \cos^2(x)} dx$	55
3.3	$\int \frac{\sin^4(x)}{a-a \cos^2(x)} dx$	61
3.4	$\int \frac{\sin^3(x)}{a-a \cos^2(x)} dx$	67
3.5	$\int \frac{\sin^2(x)}{a-a \cos^2(x)} dx$	72
3.6	$\int \frac{\sin(x)}{a-a \cos^2(x)} dx$	77

3.7	$\int \frac{\csc(x)}{a-a\cos^2(x)} dx$	83
3.8	$\int \frac{\csc^2(x)}{a-a\cos^2(x)} dx$	89
3.9	$\int \frac{\csc^3(x)}{a-a\cos^2(x)} dx$	94
3.10	$\int \frac{\sin^7(x)}{a+b\cos^2(x)} dx$	101
3.11	$\int \frac{\sin^5(x)}{a+b\cos^2(x)} dx$	108
3.12	$\int \frac{\sin^3(x)}{a+b\cos^2(x)} dx$	114
3.13	$\int \frac{\sin(x)}{a+b\cos^2(x)} dx$	120
3.14	$\int \frac{\csc(x)}{a+b\cos^2(x)} dx$	126
3.15	$\int \frac{\csc^3(x)}{a+b\cos^2(x)} dx$	133
3.16	$\int \frac{\csc^5(x)}{a+b\cos^2(x)} dx$	141
3.17	$\int \frac{\sin^6(x)}{a+b\cos^2(x)} dx$	150
3.18	$\int \frac{\sin^4(x)}{a+b\cos^2(x)} dx$	158
3.19	$\int \frac{\sin^2(x)}{a+b\cos^2(x)} dx$	165
3.20	$\int \frac{1}{a+b\cos^2(x)} dx$	171
3.21	$\int \frac{\csc^2(x)}{a+b\cos^2(x)} dx$	177
3.22	$\int \frac{\csc^4(x)}{a+b\cos^2(x)} dx$	183
3.23	$\int \frac{\csc^6(x)}{a+b\cos^2(x)} dx$	189
3.24	$\int \frac{\sin(x)}{4-3\cos^3(x)} dx$	196
3.25	$\int \frac{\cos^7(x)}{a+b\cos^2(x)} dx$	204
3.26	$\int \frac{\cos^5(x)}{a+b\cos^2(x)} dx$	210
3.27	$\int \frac{\cos^3(x)}{a+b\cos^2(x)} dx$	216
3.28	$\int \frac{\cos(x)}{a+b\cos^2(x)} dx$	222
3.29	$\int \frac{\sec(x)}{a+b\cos^2(x)} dx$	228
3.30	$\int \frac{\sec^3(x)}{a+b\cos^2(x)} dx$	235
3.31	$\int \frac{\sec^5(x)}{a+b\cos^2(x)} dx$	243
3.32	$\int \frac{\cos^6(x)}{a+b\cos^2(x)} dx$	252
3.33	$\int \frac{\cos^4(x)}{a+b\cos^2(x)} dx$	260
3.34	$\int \frac{\cos^2(x)}{a+b\cos^2(x)} dx$	267
3.35	$\int \frac{1}{a+b\cos^2(x)} dx$	274
3.36	$\int \frac{\sec^2(x)}{a+b\cos^2(x)} dx$	280
3.37	$\int \frac{\sec^4(x)}{a+b\cos^2(x)} dx$	286
3.38	$\int \frac{\sec^6(x)}{a+b\cos^2(x)} dx$	292

3.39	$\int \frac{\cos(x)}{\sqrt{1+\cos^2(x)}} dx$	299
3.40	$\int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx$	304
3.41	$\int \frac{\cos(x)}{\sqrt{4-\cos^2(x)}} dx$	309
3.42	$\int \frac{\tan(x)}{1+\cos^2(x)} dx$	314
3.43	$\int \sqrt{a+b\cos^2(x)} \tan(x) dx$	320
3.44	$\int \sqrt{1-\cos^2(x)} \tan(x) dx$	326
3.45	$\int \frac{\tan(x)}{\sqrt{a+b\cos^2(x)}} dx$	332
3.46	$\int \frac{\tan(x)}{\sqrt{1+\cos^2(x)}} dx$	338
3.47	$\int \frac{\tan(x)}{\sqrt{1-\cos^2(x)}} dx$	344
3.48	$\int \frac{\tan^3(x)}{a+b\cos^3(x)} dx$	350
3.49	$\int \sqrt{a+b\cos^3(x)} \tan(x) dx$	358
3.50	$\int \frac{\tan(x)}{\sqrt{a+b\cos^3(x)}} dx$	364
3.51	$\int \sqrt{a+b\cos^4(x)} \tan(x) dx$	370
3.52	$\int \frac{\tan(x)}{\sqrt{a+b\cos^4(x)}} dx$	377
3.53	$\int \sqrt{a+b\cos^n(x)} \tan(x) dx$	383
3.54	$\int \frac{\tan(x)}{\sqrt{a+b\cos^n(x)}} dx$	389
4	Appendix	395
4.1	Listing of Grading functions	395
4.2	Links to plain text integration problems used in this report for each CAS413	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [54]. This is test number [211].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (54)	0.00 (0)
Mathematica	100.00 (54)	0.00 (0)
Fricas	98.15 (53)	1.85 (1)
Giac	98.15 (53)	1.85 (1)
Maxima	98.15 (53)	1.85 (1)
Maple	96.30 (52)	3.70 (2)
Mupad	74.07 (40)	25.93 (14)
Reduce	72.22 (39)	27.78 (15)
Sympy	20.37 (11)	79.63 (43)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Maple	87.037	5.556	3.704	3.704
Mathematica	83.333	14.815	1.852	0.000
Maxima	81.481	16.667	0.000	1.852
Giac	79.630	18.519	0.000	1.852
Fricas	62.963	33.333	1.852	1.852
Sympy	3.704	16.667	0.000	79.630
Mupad	0.000	74.074	0.000	25.926
Reduce	0.000	72.222	0.000	27.778

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	1	0.00	0.00	100.00
Giac	1	100.00	0.00	0.00
Maxima	1	100.00	0.00	0.00
Maple	2	100.00	0.00	0.00
Mupad	14	0.00	100.00	0.00
Reduce	15	100.00	0.00	0.00
Sympy	43	69.77	30.23	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.10
Giac	0.12
Reduce	0.17
Mathematica	0.21
Fricas	0.22
Rubi	0.27
Maple	0.44
Mupad	0.89
Sympy	10.02

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	46.69	1.19	37.50	0.96
Rubi	47.22	1.05	38.00	1.00
Giac	55.40	1.34	41.00	1.20
Maxima	55.81	1.38	48.00	1.12
Mathematica	57.26	1.23	39.00	1.00
Reduce	126.87	2.33	91.00	2.10
Fricas	160.28	3.23	113.00	2.98
Mupad	242.25	3.75	58.00	1.08
Sympy	7113.09	243.31	85.00	4.11

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A `time limit` of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError.`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	43

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 8, 10, 13, 14, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 49, 50, 51, 52, 53, 54 }

B grade { 7, 9, 11, 12, 15, 16, 30, 31 }

C grade { 48 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42, 43, 44, 45, 46, 47, 51, 52, 53, 54 }

B grade { 39, 40, 41 }

C grade { 24, 48 }

F normal fail { 49, 50 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 8, 10, 11, 12, 13, 14, 17, 18, 19, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 42, 43, 44, 45, 46, 49, 51, 52, 53, 54 }

B grade { 6, 7, 9, 15, 16, 20, 21, 22, 23, 28, 35, 36, 37, 38, 39, 40, 41, 47 }

C grade { 48 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { 50 }

Maxima

A grade { 1, 2, 3, 4, 5, 8, 9, 10, 11, 12, 13, 14, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 48, 49, 50, 51, 53, 54 }

B grade { 6, 7, 15, 16, 43, 44, 45, 46, 47 }

C grade { }

F normal fail { 52 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 2, 3, 4, 5, 8, 9, 10, 11, 12, 13, 14, 17, 18, 19, 20, 21, 22, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42, 43, 45, 48, 49, 50, 51, 52, 53, 54 }

B grade { 6, 7, 15, 16, 23, 39, 41, 44, 46, 47 }

C grade { }

F normal fail { 40 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42, 48 }

C grade { }

F normal fail { }

F(-1) timeout fail { 39, 40, 41, 43, 44, 45, 46, 47, 49, 50, 51, 52, 53, 54 }

F(-2) exception fail { }

Sympy

A grade { 5, 24 }

B grade { 1, 2, 3, 4, 6, 13, 20, 28, 35 }

C grade { }

F normal fail { 7, 8, 9, 14, 15, 16, 21, 22, 23, 29, 30, 31, 36, 37, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

F(-1) timeout fail { 10, 11, 12, 17, 18, 19, 25, 26, 27, 32, 33, 34, 38 }

F(-2) exception fail { }

Reduce

A grade { }

**B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26,
27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42 }**

C grade { }

F normal fail { 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	26	20	37	23	473	31	23	25
N.S.	1	1.00	0.79	0.61	1.12	0.70	14.33	0.94	0.70	0.76
time (sec)	N/A	0.281	0.015	0.512	0.123	0.078	1.979	0.110	0.159	0.961

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	16	19	16	14	14	78	14	19	16
N.S.	1	0.84	1.00	0.84	0.74	0.74	4.11	0.74	1.00	0.84
time (sec)	N/A	0.256	0.013	0.280	0.029	0.069	1.216	0.109	0.151	0.949

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	18	18	16	21	14	153	22	13	15
N.S.	1	0.90	0.90	0.80	1.05	0.70	7.65	1.10	0.65	0.75
time (sec)	N/A	0.232	0.013	0.167	0.108	0.095	0.742	0.110	0.152	0.967

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	7	7	12	7	10	7
N.S.	1	1.00	1.00	1.14	1.00	1.00	1.71	1.00	1.43	1.00
time (sec)	N/A	0.235	0.010	0.100	0.036	0.070	0.427	0.111	0.162	0.015

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	2	5	5	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	0.40	1.00	1.00	1.00
time (sec)	N/A	0.195	0.001	0.058	0.130	0.056	0.254	0.128	0.148	0.942

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	21	22	19	23	18	8
N.S.	1	1.00	1.00	1.12	2.62	2.75	2.38	2.88	2.25	1.00
time (sec)	N/A	0.225	0.006	0.063	0.030	0.120	0.095	0.112	0.150	0.991

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	20	51	28	37	48	0	38	24	26
N.S.	1	0.91	2.32	1.27	1.68	2.18	0.00	1.73	1.09	1.18
time (sec)	N/A	0.290	0.017	0.112	0.032	0.085	0.000	0.111	0.158	0.965

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	16	21	18	17	29	0	17	19	13
N.S.	1	0.84	1.11	0.95	0.89	1.53	0.00	0.89	1.00	0.68
time (sec)	N/A	0.260	0.005	0.108	0.027	0.064	0.000	0.115	0.150	0.939

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	75	44	51	72	0	47	33	39
N.S.	1	1.00	2.14	1.26	1.46	2.06	0.00	1.34	0.94	1.11
time (sec)	N/A	0.359	0.018	0.156	0.036	0.097	0.000	0.115	0.158	0.047

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	143	94	87	225	0	99	321	100
N.S.	1	1.00	1.83	1.21	1.12	2.88	0.00	1.27	4.12	1.28
time (sec)	N/A	0.292	0.358	1.942	0.174	0.105	0.000	0.113	0.206	0.974

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	116	57	53	152	0	59	214	65
N.S.	1	1.00	2.15	1.06	0.98	2.81	0.00	1.09	3.96	1.20
time (sec)	N/A	0.270	0.268	0.788	0.106	0.133	0.000	0.109	0.163	0.974

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	90	34	30	95	0	30	119	28
N.S.	1	1.00	2.50	0.94	0.83	2.64	0.00	0.83	3.31	0.78
time (sec)	N/A	0.230	0.248	0.308	0.140	0.151	0.000	0.114	0.160	0.981

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	18	17	73	66	17	25	18
N.S.	1	1.00	1.00	0.69	0.65	2.81	2.54	0.65	0.96	0.69
time (sec)	N/A	0.210	0.042	0.103	0.116	0.154	0.420	0.101	0.149	0.038

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	50	56	48	113	0	50	66	853
N.S.	1	1.00	1.19	1.33	1.14	2.69	0.00	1.19	1.57	20.31
time (sec)	N/A	0.252	0.079	0.181	0.108	0.166	0.000	0.114	0.157	1.171

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	77	140	95	105	274	0	103	122	1138
N.S.	1	1.24	2.26	1.53	1.69	4.42	0.00	1.66	1.97	18.35
time (sec)	N/A	0.294	0.664	0.351	0.112	0.214	0.000	0.117	0.169	1.149

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	125	204	155	200	592	0	178	197	833
N.S.	1	1.33	2.17	1.65	2.13	6.30	0.00	1.89	2.10	8.86
time (sec)	N/A	0.336	1.556	0.630	0.115	0.265	0.000	0.121	0.160	2.785

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	114	77	94	112	285	0	119	239	681
N.S.	1	1.30	0.88	1.07	1.27	3.24	0.00	1.35	2.72	7.74
time (sec)	N/A	0.356	0.253	1.235	0.112	0.234	0.000	0.116	0.277	1.099

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	74	52	61	62	211	0	80	139	126
N.S.	1	1.23	0.87	1.02	1.03	3.52	0.00	1.33	2.32	2.10
time (sec)	N/A	0.290	0.149	0.485	0.107	0.184	0.000	0.119	0.235	1.060

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	41	37	36	33	177	0	50	64	108
N.S.	1	1.02	0.92	0.90	0.82	4.42	0.00	1.25	1.60	2.70
time (sec)	N/A	0.248	0.114	0.132	0.112	0.204	0.000	0.113	0.272	1.010

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	29	21	20	163	10924	37	54	24
N.S.	1	1.00	0.97	0.70	0.67	5.43	364.13	1.23	1.80	0.80
time (sec)	N/A	0.197	0.022	0.000	0.106	0.187	19.454	0.113	0.151	0.001

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	40	39	38	228	0	55	93	34
N.S.	1	1.00	0.98	0.95	0.93	5.56	0.00	1.34	2.27	0.83
time (sec)	N/A	0.246	0.142	0.221	0.114	0.187	0.000	0.123	0.156	0.984

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	59	57	70	396	0	90	157	67
N.S.	1	1.00	0.97	0.93	1.15	6.49	0.00	1.48	2.57	1.10
time (sec)	N/A	0.291	0.267	0.432	0.114	0.199	0.000	0.122	0.193	1.002

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	90	83	127	610	0	156	238	101
N.S.	1	1.00	1.01	0.93	1.43	6.85	0.00	1.75	2.67	1.13
time (sec)	N/A	0.319	0.446	0.773	0.119	0.194	0.000	0.121	0.181	1.008

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	100	79	35	89	67	85	60	75	75
N.S.	1	1.19	0.94	0.42	1.06	0.80	1.01	0.71	0.89	0.89
time (sec)	N/A	0.350	0.200	0.125	0.116	0.128	0.594	0.122	0.167	0.172

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	111	78	91	259	0	96	148	86
N.S.	1	1.00	1.42	1.00	1.17	3.32	0.00	1.23	1.90	1.10
time (sec)	N/A	0.299	0.514	1.722	0.112	0.198	0.000	0.118	0.184	0.075

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	86	50	67	191	0	65	117	51
N.S.	1	1.00	1.54	0.89	1.20	3.41	0.00	1.16	2.09	0.91
time (sec)	N/A	0.272	0.243	0.682	0.108	0.166	0.000	0.111	0.165	0.064

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	33	50	134	0	41	91	30
N.S.	1	1.00	1.00	0.87	1.32	3.53	0.00	1.08	2.39	0.79
time (sec)	N/A	0.233	0.039	0.270	0.111	0.139	0.000	0.124	0.189	1.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	21	39	95	55508	31	69	21
N.S.	1	1.00	1.00	0.72	1.34	3.28	1914.07	1.07	2.38	0.72
time (sec)	N/A	0.210	0.017	0.100	0.105	0.129	66.630	0.117	0.159	0.986

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	38	47	64	119	0	57	116	414
N.S.	1	1.00	0.93	1.15	1.56	2.90	0.00	1.39	2.83	10.10
time (sec)	N/A	0.235	0.073	0.214	0.106	0.165	0.000	0.117	0.162	0.139

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	73	152	82	92	186	0	85	342	483
N.S.	1	1.24	2.58	1.39	1.56	3.15	0.00	1.44	5.80	8.19
time (sec)	N/A	0.284	0.483	0.521	0.109	0.205	0.000	0.119	0.183	1.130

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	117	215	137	145	270	0	127	681	969
N.S.	1	1.30	2.39	1.52	1.61	3.00	0.00	1.41	7.57	10.77
time (sec)	N/A	0.344	1.351	1.036	0.115	0.189	0.000	0.123	0.185	1.247

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	116	76	92	97	273	0	104	148	1036
N.S.	1	1.33	0.87	1.06	1.11	3.14	0.00	1.20	1.70	11.91
time (sec)	N/A	0.381	0.270	1.107	0.119	0.211	0.000	0.112	0.183	1.185

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	74	52	59	54	213	0	72	127	291
N.S.	1	1.23	0.87	0.98	0.90	3.55	0.00	1.20	2.12	4.85
time (sec)	N/A	0.287	0.145	0.418	0.115	0.200	0.000	0.111	0.156	1.146

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	34	31	183	0	48	70	425
N.S.	1	1.00	0.95	0.89	0.82	4.82	0.00	1.26	1.84	11.18
time (sec)	N/A	0.287	0.736	0.124	0.111	0.186	0.000	0.120	0.157	1.086

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	29	21	20	163	10924	37	54	24
N.S.	1	1.00	0.97	0.70	0.67	5.43	364.13	1.23	1.80	0.80
time (sec)	N/A	0.198	0.025	0.000	0.114	0.147	18.454	0.112	0.157	0.001

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	38	33	32	216	0	36	85	30
N.S.	1	1.00	1.03	0.89	0.86	5.84	0.00	0.97	2.30	0.81
time (sec)	N/A	0.241	0.468	0.335	0.113	0.180	0.000	0.118	0.169	0.978

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	55	49	48	276	0	71	211	51
N.S.	1	1.00	0.98	0.88	0.86	4.93	0.00	1.27	3.77	0.91
time (sec)	N/A	0.287	0.561	0.664	0.117	0.166	0.000	0.120	0.186	0.974

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	80	78	74	348	0	104	346	84
N.S.	1	1.00	1.01	0.99	0.94	4.41	0.00	1.32	4.38	1.06
time (sec)	N/A	0.315	0.808	1.329	0.113	0.217	0.000	0.118	0.209	1.012

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	33	8	49	0	23	20	0
N.S.	1	1.00	1.00	3.67	0.89	5.44	0.00	2.56	2.22	0.00
time (sec)	N/A	0.196	0.013	0.090	0.109	0.110	0.000	0.131	0.152	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	57	11	89	0	0	32	0
N.S.	1	1.00	1.00	3.80	0.73	5.93	0.00	0.00	2.13	0.00
time (sec)	N/A	0.204	0.031	0.247	0.108	0.132	0.000	0.000	0.163	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	53	8	39	0	29	24	0
N.S.	1	1.00	1.00	5.89	0.89	4.33	0.00	3.22	2.67	0.00
time (sec)	N/A	0.201	0.015	0.159	0.107	0.134	0.000	0.140	0.157	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	19	17	16	19	19	0	16	56	9
N.S.	1	1.12	1.00	0.94	1.12	1.12	0.00	0.94	3.29	0.53
time (sec)	N/A	0.207	0.013	0.263	0.027	0.109	0.000	0.114	0.153	0.981

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	45	40	43	95	87	0	38	14	0
N.S.	1	1.12	1.00	1.08	2.38	2.18	0.00	0.95	0.35	0.00
time (sec)	N/A	0.251	0.034	0.101	0.128	0.265	0.000	0.119	0.164	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	26	17	17	47	21	0	45	14	0
N.S.	1	1.30	0.85	0.85	2.35	1.05	0.00	2.25	0.70	0.00
time (sec)	N/A	0.266	0.023	0.089	0.112	0.121	0.000	0.114	0.151	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	30	81	64	0	24	24	0
N.S.	1	1.00	1.00	1.20	3.24	2.56	0.00	0.96	0.96	0.00
time (sec)	N/A	0.242	0.019	0.094	0.118	0.145	0.000	0.117	0.154	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	10	60	16	0	27	20	0
N.S.	1	1.00	1.00	0.91	5.45	1.45	0.00	2.45	1.82	0.00
time (sec)	N/A	0.214	0.014	0.082	0.112	0.188	0.000	0.113	0.159	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	14	8	39	17	0	33	24	0
N.S.	1	1.00	1.56	0.89	4.33	1.89	0.00	3.67	2.67	0.00
time (sec)	N/A	0.268	0.015	0.076	0.108	0.093	0.000	0.114	0.160	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	C	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	153	153	217	109	151	823	0	143	17	1281
N.S.	1	1.00	1.42	0.71	0.99	5.38	0.00	0.93	0.11	8.37
time (sec)	N/A	0.464	0.408	2.991	0.117	1.790	0.000	0.127	0.166	2.547

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	45	0	52	134	0	38	14	0
N.S.	1	1.00	1.00	0.00	1.16	2.98	0.00	0.84	0.31	0.00
time (sec)	N/A	0.261	0.033	0.000	0.121	1.743	0.000	0.120	0.164	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	F(-2)	F	A	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	0	39	0	0	24	24	0
N.S.	1	1.00	1.00	0.00	1.39	0.00	0.00	0.86	0.86	0.00
time (sec)	N/A	0.256	0.021	0.000	0.127	0.000	0.000	0.111	0.156	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	45	44	43	87	0	38	14	0
N.S.	1	1.00	1.00	0.98	0.96	1.93	0.00	0.84	0.31	0.00
time (sec)	N/A	0.255	0.041	0.365	0.045	0.250	0.000	0.112	0.184	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	28	31	0	64	0	24	24	0
N.S.	1	1.00	1.00	1.11	0.00	2.29	0.00	0.86	0.86	0.00
time (sec)	N/A	0.247	0.019	0.228	0.000	0.198	0.000	0.111	0.158	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	46	46	39	58	94	0	46	14	0
N.S.	1	0.98	0.98	0.83	1.23	2.00	0.00	0.98	0.30	0.00
time (sec)	N/A	0.269	0.040	0.286	0.117	0.113	0.000	0.135	0.159	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	24	42	71	0	27	24	0
N.S.	1	1.00	1.00	0.83	1.45	2.45	0.00	0.93	0.83	0.00
time (sec)	N/A	0.269	0.019	0.075	0.108	0.102	0.000	0.131	0.162	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [24] had the largest ratio of [.76923099999999998]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	7	7	1.00	16	0.438
2	A	8	7	0.84	16	0.438
3	A	5	5	0.90	16	0.312
4	A	6	6	1.00	16	0.375
5	A	3	3	1.00	16	0.188
6	A	6	6	1.00	14	0.429
7	A	8	8	0.91	14	0.571
8	A	6	5	0.84	16	0.312
9	A	10	10	1.00	16	0.625
10	A	6	5	1.00	15	0.333
11	A	6	5	1.00	15	0.333
12	A	6	5	1.00	15	0.333
13	A	5	4	1.00	13	0.308
14	A	7	6	1.00	13	0.462
15	A	8	7	1.24	15	0.467
16	A	9	8	1.33	15	0.533
17	A	8	7	1.30	15	0.467
18	A	7	6	1.23	15	0.400
19	A	6	5	1.02	15	0.333
20	A	4	3	1.00	10	0.300
21	A	5	4	1.00	15	0.267

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	5	4	1.00	15	0.267
23	A	5	4	1.00	15	0.267
24	A	11	10	1.19	13	0.769
25	A	5	4	1.00	15	0.267
26	A	5	4	1.00	15	0.267
27	A	5	4	1.00	15	0.267
28	A	4	3	1.00	13	0.231
29	A	6	5	1.00	13	0.385
30	A	7	6	1.24	15	0.400
31	A	8	7	1.30	15	0.467
32	A	8	7	1.33	15	0.467
33	A	7	6	1.23	15	0.400
34	A	6	5	1.00	15	0.333
35	A	4	3	1.00	10	0.300
36	A	5	4	1.00	15	0.267
37	A	5	4	1.00	15	0.267
38	A	5	4	1.00	15	0.267
39	A	4	3	1.00	13	0.231
40	A	4	3	1.00	21	0.143
41	A	4	3	1.00	15	0.200
42	A	7	6	1.12	11	0.545
43	A	7	6	1.12	15	0.400
44	A	10	9	1.30	15	0.600
45	A	6	5	1.00	15	0.333
46	A	6	5	1.00	13	0.385
47	A	9	8	1.00	15	0.533
48	A	6	5	1.00	15	0.333
49	A	8	7	1.00	15	0.467
50	A	7	6	1.00	15	0.400
51	A	8	7	1.00	15	0.467
52	A	7	6	1.00	15	0.400
53	A	8	7	0.98	15	0.467

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	7	6	1.00	15	0.400

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{\sin^6(x)}{a-a\cos^2(x)} dx$	48
3.2	$\int \frac{\sin^5(x)}{a-a\cos^2(x)} dx$	55
3.3	$\int \frac{\sin^4(x)}{a-a\cos^2(x)} dx$	61
3.4	$\int \frac{\sin^3(x)}{a-a\cos^2(x)} dx$	67
3.5	$\int \frac{\sin^2(x)}{a-a\cos^2(x)} dx$	72
3.6	$\int \frac{\sin(x)}{a-a\cos^2(x)} dx$	77
3.7	$\int \frac{\csc(x)}{a-a\cos^2(x)} dx$	83
3.8	$\int \frac{\csc^2(x)}{a-a\cos^2(x)} dx$	89
3.9	$\int \frac{\csc^3(x)}{a-a\cos^2(x)} dx$	94
3.10	$\int \frac{\sin^7(x)}{a+b\cos^2(x)} dx$	101
3.11	$\int \frac{\sin^5(x)}{a+b\cos^2(x)} dx$	108
3.12	$\int \frac{\sin^3(x)}{a+b\cos^2(x)} dx$	114
3.13	$\int \frac{\sin(x)}{a+b\cos^2(x)} dx$	120
3.14	$\int \frac{\csc(x)}{a+b\cos^2(x)} dx$	126
3.15	$\int \frac{\csc^3(x)}{a+b\cos^2(x)} dx$	133
3.16	$\int \frac{\csc^5(x)}{a+b\cos^2(x)} dx$	141
3.17	$\int \frac{\sin^6(x)}{a+b\cos^2(x)} dx$	150
3.18	$\int \frac{\sin^4(x)}{a+b\cos^2(x)} dx$	158
3.19	$\int \frac{\sin^2(x)}{a+b\cos^2(x)} dx$	165
3.20	$\int \frac{1}{a+b\cos^2(x)} dx$	171
3.21	$\int \frac{\csc^2(x)}{a+b\cos^2(x)} dx$	177
3.22	$\int \frac{\csc^4(x)}{a+b\cos^2(x)} dx$	183

3.23	$\int \frac{\csc^6(x)}{a+b\cos^2(x)} dx$	189
3.24	$\int \frac{\sin(x)}{4-3\cos^3(x)} dx$	196
3.25	$\int \frac{\cos^7(x)}{a+b\cos^2(x)} dx$	204
3.26	$\int \frac{\cos^5(x)}{a+b\cos^2(x)} dx$	210
3.27	$\int \frac{\cos^3(x)}{a+b\cos^2(x)} dx$	216
3.28	$\int \frac{\cos(x)}{a+b\cos^2(x)} dx$	222
3.29	$\int \frac{\sec(x)}{a+b\cos^2(x)} dx$	228
3.30	$\int \frac{\sec^3(x)}{a+b\cos^2(x)} dx$	235
3.31	$\int \frac{\sec^5(x)}{a+b\cos^2(x)} dx$	243
3.32	$\int \frac{\cos^6(x)}{a+b\cos^2(x)} dx$	252
3.33	$\int \frac{\cos^4(x)}{a+b\cos^2(x)} dx$	260
3.34	$\int \frac{\cos^2(x)}{a+b\cos^2(x)} dx$	267
3.35	$\int \frac{1}{a+b\cos^2(x)} dx$	274
3.36	$\int \frac{\sec^2(x)}{a+b\cos^2(x)} dx$	280
3.37	$\int \frac{\sec^4(x)}{a+b\cos^2(x)} dx$	286
3.38	$\int \frac{\sec^6(x)}{a+b\cos^2(x)} dx$	292
3.39	$\int \frac{\cos(x)}{\sqrt{1+\cos^2(x)}} dx$	299
3.40	$\int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx$	304
3.41	$\int \frac{\cos(x)}{\sqrt{4-\cos^2(x)}} dx$	309
3.42	$\int \frac{\tan(x)}{1+\cos^2(x)} dx$	314
3.43	$\int \sqrt{a+b\cos^2(x)} \tan(x) dx$	320
3.44	$\int \sqrt{1-\cos^2(x)} \tan(x) dx$	326
3.45	$\int \frac{\tan(x)}{\sqrt{a+b\cos^2(x)}} dx$	332
3.46	$\int \frac{\tan(x)}{\sqrt{1+\cos^2(x)}} dx$	338
3.47	$\int \frac{\tan(x)}{\sqrt{1-\cos^2(x)}} dx$	344
3.48	$\int \frac{\tan^3(x)}{a+b\cos^3(x)} dx$	350
3.49	$\int \sqrt{a+b\cos^3(x)} \tan(x) dx$	358
3.50	$\int \frac{\tan(x)}{\sqrt{a+b\cos^3(x)}} dx$	364
3.51	$\int \sqrt{a+b\cos^4(x)} \tan(x) dx$	370
3.52	$\int \frac{\tan(x)}{\sqrt{a+b\cos^4(x)}} dx$	377
3.53	$\int \sqrt{a+b\cos^n(x)} \tan(x) dx$	383
3.54	$\int \frac{\tan(x)}{\sqrt{a+b\cos^n(x)}} dx$	389

3.1 $\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx$

Optimal result	48
Mathematica [A] (verified)	48
Rubi [A] (verified)	49
Maple [A] (verified)	50
Fricas [A] (verification not implemented)	51
Sympy [B] (verification not implemented)	51
Maxima [A] (verification not implemented)	53
Giac [A] (verification not implemented)	53
Mupad [B] (verification not implemented)	53
Reduce [B] (verification not implemented)	54

Optimal result

Integrand size = 16, antiderivative size = 33

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{3x}{8a} - \frac{3 \cos(x) \sin(x)}{8a} - \frac{\cos(x) \sin^3(x)}{4a}$$

output 3/8*x/a-3/8*cos(x)*sin(x)/a-1/4*cos(x)*sin(x)^3/a

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{\frac{3x}{8} - \frac{1}{4} \sin(2x) + \frac{1}{32} \sin(4x)}{a}$$

input Integrate[Sin[x]^6/(a - a*Cos[x]^2), x]

output ((3*x)/8 - Sin[2*x]/4 + Sin[4*x]/32)/a

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {3042, 3654, 3042, 3115, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^6(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(x + \frac{\pi}{2})^6}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3654} \\
 & \frac{\int \sin^4(x) dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sin(x)^4 dx}{a} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\frac{3}{4} \int \sin^2(x) dx - \frac{1}{4} \sin^3(x) \cos(x)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{3}{4} \int \sin(x)^2 dx - \frac{1}{4} \sin^3(x) \cos(x)}{a} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\frac{3}{4} \left(\frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x)}{a} \\
 & \quad \downarrow \text{24} \\
 & \frac{\frac{3}{4} \left(\frac{x}{2} - \frac{1}{2} \sin(x) \cos(x) \right) - \frac{1}{4} \sin^3(x) \cos(x)}{a}
 \end{aligned}$$

input $\text{Int}[\sin[x]^6/(a - a*\cos[x]^2), x]$

output $(-1/4*(\cos[x]*\sin[x]^3) + (3*(x/2 - (\cos[x]*\sin[x])/2))/4)/a$

Definitions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3115 $\text{Int}[(b_*)\sin(c_*) + (d_*)\sin(x_*)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((b*\sin[c + d*x])^{(n - 1)/(d*n)}), x] + \text{Simp}[b^{2*((n - 1)/n)} \text{Int}[(b*\sin[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}[b, c, d, x] \&& \text{GtQ}[n, 1] \&& \text{IntegerQ}[2*n]$

rule 3654 $\text{Int}[(u_*) + (b_*)\sin(e_*) + (f_*)\sin(x_*)]^2, x_Symbol] \rightarrow \text{Simp}[a^p \text{Int}[\text{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /; \text{FreeQ}[a, b, e, f, p, x] \&& \text{EqQ}[a + b, 0] \&& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.61

method	result
parallelrisch	$\frac{12x + \sin(4x) - 8\sin(2x)}{32a}$
risch	$\frac{3x}{8a} + \frac{\sin(4x)}{32a} - \frac{\sin(2x)}{4a}$ $\frac{-\frac{5\tan(x)^3}{8} - \frac{3\tan(x)}{8}}{(1+\tan(x)^2)^2} + \frac{3\arctan(\tan(x))}{8}$
default	$\frac{a}{a}$
norman	$\frac{-\frac{3\tan(\frac{x}{2})^2}{4a} - \frac{17\tan(\frac{x}{2})^4}{4a} - \frac{7\tan(\frac{x}{2})^6}{2a} + \frac{7\tan(\frac{x}{2})^8}{2a} + \frac{17\tan(\frac{x}{2})^{10}}{4a} + \frac{3\tan(\frac{x}{2})^{12}}{4a} + \frac{3x\tan(\frac{x}{2})}{8a} + \frac{9x\tan(\frac{x}{2})^3}{4a} + \frac{45x\tan(\frac{x}{2})^5}{8a} + \frac{15x\tan(\frac{x}{2})^7}{4a}}{(1+\tan(\frac{x}{2})^2)^6} \tan(\frac{x}{2})$

input `int(sin(x)^6/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`

output `1/32*(12*x+sin(4*x)-8*sin(2*x))/a`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{(2 \cos(x)^3 - 5 \cos(x)) \sin(x) + 3 x}{8 a}$$

input `integrate(sin(x)^6/(a-a*cos(x)^2),x, algorithm="fricas")`

output `1/8*((2*cos(x)^3 - 5*cos(x))*sin(x) + 3*x)/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 473 vs. 2(29) = 58.

Time = 1.98 (sec) , antiderivative size = 473, normalized size of antiderivative = 14.33

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{3x \tan^8\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ + \frac{12x \tan^6\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ + \frac{18x \tan^4\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ + \frac{12x \tan^2\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ + \frac{3x}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ + \frac{6 \tan^7\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ + \frac{22 \tan^5\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ - \frac{22 \tan^3\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a} \\ - \frac{6 \tan\left(\frac{x}{2}\right)}{8a \tan^8\left(\frac{x}{2}\right) + 32a \tan^6\left(\frac{x}{2}\right) + 48a \tan^4\left(\frac{x}{2}\right) + 32a \tan^2\left(\frac{x}{2}\right) + 8a}$$

input `integrate(sin(x)**6/(a-a*cos(x)**2),x)`

output
$$3*x*tan(x/2)**8/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) + 12*x*tan(x/2)**6/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) + 18*x*tan(x/2)**4/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) + 12*x*tan(x/2)**2/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) + 3*x/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) + 6*tan(x/2)**7/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) + 22*tan(x/2)**5/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) - 22*tan(x/2)**3/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a) - 6*tan(x/2)/(8*a*tan(x/2)**8 + 32*a*tan(x/2)**6 + 48*a*tan(x/2)**4 + 32*a*tan(x/2)**2 + 8*a)$$

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.12

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = -\frac{5 \tan(x)^3 + 3 \tan(x)}{8(a \tan(x)^4 + 2a \tan(x)^2 + a)} + \frac{3x}{8a}$$

input `integrate(sin(x)^6/(a-a*cos(x)^2),x, algorithm="maxima")`

output `-1/8*(5*tan(x)^3 + 3*tan(x))/(a*tan(x)^4 + 2*a*tan(x)^2 + a) + 3/8*x/a`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{3x}{8a} - \frac{5 \tan(x)^3 + 3 \tan(x)}{8(\tan(x)^2 + 1)^2 a}$$

input `integrate(sin(x)^6/(a-a*cos(x)^2),x, algorithm="giac")`

output `3/8*x/a - 1/8*(5*tan(x)^3 + 3*tan(x))/((tan(x)^2 + 1)^2*a)`

Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{\sin(4x)}{32a} - \frac{\sin(2x)}{4a} + \frac{3x}{8a}$$

input `int(sin(x)^6/(a - a*cos(x)^2),x)`

output `sin(4*x)/(32*a) - sin(2*x)/(4*a) + (3*x)/(8*a)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int \frac{\sin^6(x)}{a - a \cos^2(x)} dx = \frac{-2 \cos(x) \sin(x)^3 - 3 \cos(x) \sin(x) + 3x}{8a}$$

input `int(sin(x)^6/(a-a*cos(x)^2),x)`

output `(- 2*cos(x)*sin(x)**3 - 3*cos(x)*sin(x) + 3*x)/(8*a)`

3.2 $\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx$

Optimal result	55
Mathematica [A] (verified)	55
Rubi [A] (verified)	56
Maple [A] (verified)	58
Fricas [A] (verification not implemented)	58
Sympy [B] (verification not implemented)	59
Maxima [A] (verification not implemented)	59
Giac [A] (verification not implemented)	59
Mupad [B] (verification not implemented)	60
Reduce [B] (verification not implemented)	60

Optimal result

Integrand size = 16, antiderivative size = 19

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a} + \frac{\cos^3(x)}{3a}$$

output $-\cos(x)/a+1/3*\cos(x)^3/a$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = \frac{-\frac{3 \cos(x)}{4} + \frac{1}{12} \cos(3x)}{a}$$

input `Integrate[Sin[x]^5/(a - a*Cos[x]^2), x]`

output $((-3*\cos[x])/4 + \cos[3*x]/12)/a$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {3042, 25, 3654, 25, 3042, 3113, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^5(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{\cos(x + \frac{\pi}{2})^5}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{\cos(x + \frac{\pi}{2})^5}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \textcolor{blue}{3654} \\
 & - \frac{\int -\sin^3(x) dx}{a} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & \frac{\int \sin^3(x) dx}{a} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{\int \sin(x)^3 dx}{a} \\
 & \quad \downarrow \textcolor{blue}{3113} \\
 & - \frac{\int (1 - \cos^2(x)) d \cos(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & - \frac{\cos(x) - \frac{\cos^3(x)}{3}}{a}
 \end{aligned}$$

input $\text{Int}[\sin[x]^5/(a - a*\cos[x]^2), x]$

output $-(\cos[x] - \cos[x]^3/3)/a$

Definitions of rubi rules used

rule 25 $\text{Int}[-(F_{x_}), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F_x, x], x]$

rule 2009 $\text{Int}[u_{}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_{}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3113 $\text{Int}[\sin[(c_{}) + (d_{})*(x_{})]^{(n_{})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-d^{(-1)} \text{Subst}[\text{Int}[\text{Exp} \text{and}[(1 - x^2)^{(n - 1)/2}, x], x], x, \cos[c + d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&& \text{IGtQ}[(n - 1)/2, 0]$

rule 3654 $\text{Int}[(u_{})*((a_{}) + (b_{})*\sin[(e_{}) + (f_{})*(x_{})]^{2})^{(p_{})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p \text{Int}[\text{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{EqQ}[a + b, 0] \&& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\cos(x)^3 - \cos(x)}{a}$	16
default	$\frac{\cos(x)^3 - \cos(x)}{a}$	16
parallelrisch	$\frac{\cos(3x) - 9\cos(x) - 8}{12a}$	16
risch	$-\frac{3\cos(x)}{4a} + \frac{\cos(3x)}{12a}$	18
norman	$\frac{-\frac{4\tan(\frac{x}{2})}{3a} - \frac{20\tan(\frac{x}{2})^3}{3a} - \frac{4\tan(\frac{x}{2})^7}{a} - \frac{28\tan(\frac{x}{2})^5}{3a}}{\left(1+\tan(\frac{x}{2})^2\right)^5 \tan(\frac{x}{2})}$	61

input `int(sin(x)^5/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`

output `1/a*(1/3*cos(x)^3-cos(x))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = \frac{\cos(x)^3 - 3 \cos(x)}{3a}$$

input `integrate(sin(x)^5/(a-a*cos(x)^2),x, algorithm="fricas")`

output `1/3*(cos(x)^3 - 3*cos(x))/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. $2(12) = 24$.

Time = 1.22 (sec) , antiderivative size = 78, normalized size of antiderivative = 4.11

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = -\frac{12 \tan^2(\frac{x}{2})}{3a \tan^6(\frac{x}{2}) + 9a \tan^4(\frac{x}{2}) + 9a \tan^2(\frac{x}{2}) + 3a} \\ - \frac{4}{3a \tan^6(\frac{x}{2}) + 9a \tan^4(\frac{x}{2}) + 9a \tan^2(\frac{x}{2}) + 3a}$$

input `integrate(sin(x)**5/(a-a*cos(x)**2),x)`

output `-12*tan(x/2)**2/(3*a*tan(x/2)**6 + 9*a*tan(x/2)**4 + 9*a*tan(x/2)**2 + 3*a) - 4/(3*a*tan(x/2)**6 + 9*a*tan(x/2)**4 + 9*a*tan(x/2)**2 + 3*a)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = \frac{\cos(x)^3 - 3 \cos(x)}{3a}$$

input `integrate(sin(x)^5/(a-a*cos(x)^2),x, algorithm="maxima")`

output `1/3*(cos(x)^3 - 3*cos(x))/a`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = \frac{\cos(x)^3 - 3 \cos(x)}{3a}$$

input `integrate(sin(x)^5/(a-a*cos(x)^2),x, algorithm="giac")`

output $\frac{1}{3}(\cos(x)^3 - 3\cos(x))/a$

Mupad [B] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = -\frac{3 \cos(x) - \cos(x)^3}{3a}$$

input `int(sin(x)^5/(a - a*cos(x)^2),x)`

output $-(3\cos(x) - \cos(x)^3)/(3a)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\sin^5(x)}{a - a \cos^2(x)} dx = \frac{-\cos(x) \sin(x)^2 - 2 \cos(x) + 2}{3a}$$

input `int(sin(x)^5/(a-a*cos(x)^2),x)`

output $(-\cos(x)*\sin(x)^2 - 2*\cos(x) + 2)/(3a)$

3.3 $\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx$

Optimal result	61
Mathematica [A] (verified)	61
Rubi [A] (verified)	62
Maple [A] (verified)	63
Fricas [A] (verification not implemented)	64
Sympy [B] (verification not implemented)	64
Maxima [A] (verification not implemented)	65
Giac [A] (verification not implemented)	65
Mupad [B] (verification not implemented)	65
Reduce [B] (verification not implemented)	66

Optimal result

Integrand size = 16, antiderivative size = 20

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = \frac{x}{2a} - \frac{\cos(x) \sin(x)}{2a}$$

output `1/2*x/a-1/2*cos(x)*sin(x)/a`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = \frac{\frac{x}{2} - \frac{1}{4} \sin(2x)}{a}$$

input `Integrate[Sin[x]^4/(a - a*Cos[x]^2), x]`

output `(x/2 - Sin[2*x]/4)/a`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 3654, 3042, 3115, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^4(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(x + \frac{\pi}{2})^4}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3654} \\
 & \frac{\int \sin^2(x) dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sin(x)^2 dx}{a} \\
 & \quad \downarrow \text{3115} \\
 & \frac{\frac{\int 1 dx}{2} - \frac{1}{2} \sin(x) \cos(x)}{a} \\
 & \quad \downarrow \text{24} \\
 & \frac{\frac{x}{2} - \frac{1}{2} \sin(x) \cos(x)}{a}
 \end{aligned}$$

input `Int[Sin[x]^4/(a - a*Cos[x]^2),x]`

output `(x/2 - (Cos[x]*Sin[x])/2)/a`

Definitions of rubi rules used

rule 24 $\text{Int}[a_-, x_{\text{Symbol}}] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 3042 $\text{Int}[u_-, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3115 $\text{Int}[(b_*)\sin(c_*) + (d_*)\sin(x_*)]^{(n_*)}, x_{\text{Symbol}} \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((b*\sin[c + d*x])^{(n - 1)/(d*n)}), x] + \text{Simp}[b^{2*((n - 1)/n)} \text{Int}[(b*\sin[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&& \text{GtQ}[n, 1] \&& \text{IntegerQ}[2*n]$

rule 3654 $\text{Int}[(a_*) + (b_*)\sin(e_*) + (f_*)\sin(x_*)]^2, x_{\text{Symbol}} \rightarrow \text{Simp}[a^p \text{Int}[\text{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{EqQ}[a + b, 0] \&& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
parallelrisch	$\frac{2x - \sin(2x)}{4a}$	16
risch	$\frac{x}{2a} - \frac{\sin(2x)}{4a}$	17
default	$\frac{-\frac{\tan(x)}{2(1+\tan(x)^2)} + \frac{\arctan(\tan(x))}{2}}{a}$	23
norman	$\frac{\tan(\frac{x}{2})^6 + \tan(\frac{x}{2})^8 - \tan(\frac{x}{2})^2 - \tan(\frac{x}{2})^4 + \frac{x \tan(\frac{x}{2})}{2a} + \frac{2x \tan(\frac{x}{2})^3}{a} + \frac{3x \tan(\frac{x}{2})^5}{a} + \frac{2x \tan(\frac{x}{2})^7}{a} + \frac{x \tan(\frac{x}{2})^9}{2a}}{(1+\tan(\frac{x}{2})^2)^4 \tan(\frac{x}{2})}$	119

input `int(sin(x)^4/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`

output `1/4*(2*x-sin(2*x))/a`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x) \sin(x) - x}{2a}$$

input `integrate(sin(x)^4/(a-a*cos(x)^2),x, algorithm="fricas")`

output `-1/2*(cos(x)*sin(x) - x)/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(14) = 28$.

Time = 0.74 (sec) , antiderivative size = 153, normalized size of antiderivative = 7.65

$$\begin{aligned} \int \frac{\sin^4(x)}{a - a \cos^2(x)} dx &= \frac{x \tan^4\left(\frac{x}{2}\right)}{2a \tan^4\left(\frac{x}{2}\right) + 4a \tan^2\left(\frac{x}{2}\right) + 2a} + \frac{2x \tan^2\left(\frac{x}{2}\right)}{2a \tan^4\left(\frac{x}{2}\right) + 4a \tan^2\left(\frac{x}{2}\right) + 2a} \\ &+ \frac{x}{2a \tan^4\left(\frac{x}{2}\right) + 4a \tan^2\left(\frac{x}{2}\right) + 2a} \\ &+ \frac{2 \tan^3\left(\frac{x}{2}\right)}{2a \tan^4\left(\frac{x}{2}\right) + 4a \tan^2\left(\frac{x}{2}\right) + 2a} \\ &- \frac{2 \tan\left(\frac{x}{2}\right)}{2a \tan^4\left(\frac{x}{2}\right) + 4a \tan^2\left(\frac{x}{2}\right) + 2a} \end{aligned}$$

input `integrate(sin(x)**4/(a-a*cos(x)**2),x)`

output `x*tan(x/2)**4/(2*a*tan(x/2)**4 + 4*a*tan(x/2)**2 + 2*a) + 2*x*tan(x/2)**2/(2*a*tan(x/2)**4 + 4*a*tan(x/2)**2 + 2*a) + x/(2*a*tan(x/2)**4 + 4*a*tan(x/2)**2 + 2*a) + 2*tan(x/2)**3/(2*a*tan(x/2)**4 + 4*a*tan(x/2)**2 + 2*a) - 2*tan(x/2)/(2*a*tan(x/2)**4 + 4*a*tan(x/2)**2 + 2*a)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = \frac{x}{2a} - \frac{\tan(x)}{2(a \tan(x)^2 + a)}$$

input `integrate(sin(x)^4/(a-a*cos(x)^2),x, algorithm="maxima")`

output `1/2*x/a - 1/2*tan(x)/(a*tan(x)^2 + a)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = \frac{x}{2a} - \frac{\tan(x)}{2(\tan(x)^2 + 1)a}$$

input `integrate(sin(x)^4/(a-a*cos(x)^2),x, algorithm="giac")`

output `1/2*x/a - 1/2*tan(x)/((tan(x)^2 + 1)*a)`

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = \frac{2x - \sin(2x)}{4a}$$

input `int(sin(x)^4/(a - a*cos(x)^2),x)`

output `(2*x - sin(2*x))/(4*a)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.65

$$\int \frac{\sin^4(x)}{a - a \cos^2(x)} dx = \frac{-\cos(x) \sin(x) + x}{2a}$$

input `int(sin(x)^4/(a-a*cos(x)^2),x)`

output `(- cos(x)*sin(x) + x)/(2*a)`

3.4 $\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx$

Optimal result	67
Mathematica [A] (verified)	67
Rubi [A] (verified)	68
Maple [A] (verified)	69
Fricas [A] (verification not implemented)	70
Sympy [B] (verification not implemented)	70
Maxima [A] (verification not implemented)	70
Giac [A] (verification not implemented)	71
Mupad [B] (verification not implemented)	71
Reduce [B] (verification not implemented)	71

Optimal result

Integrand size = 16, antiderivative size = 7

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a}$$

output -cos(x)/a

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a}$$

input Integrate[Sin[x]^3/(a - a*Cos[x]^2), x]

output -(Cos[x]/a)

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 25, 3654, 25, 3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^3(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\cos(x + \frac{\pi}{2})^3}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\cos(x + \frac{\pi}{2})^3}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3654} \\
 & - \frac{\int -\sin(x) dx}{a} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \sin(x) dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \sin(x) dx}{a} \\
 & \quad \downarrow \text{3118} \\
 & -\frac{\cos(x)}{a}
 \end{aligned}$$

input `Int[Sin[x]^3/(a - a*Cos[x]^2),x]`

output `-(Cos[x]/a)`

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 3042 $\text{Int}[\text{u}__, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3118 $\text{Int}[\sin[(\text{c}__.) + (\text{d}__.)*(\text{x}__)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{Cos}[\text{c} + \text{d}*\text{x}]/\text{d}, \text{x}] /; \text{FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$

rule 3654 $\text{Int}[(\text{u}__.)*((\text{a}__.) + (\text{b}__.)*\sin[(\text{e}__.) + (\text{f}__.)*(\text{x}__)]^2)^{(\text{p}__)}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{a}^{\text{p}} \quad \text{Int}[\text{ActivateTrig}[\text{u}*\cos[\text{e} + \text{f}*\text{x}]^{(2*\text{p})}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{p}\}, \text{x}] \&& \text{EqQ}[\text{a} + \text{b}, 0] \&& \text{IntegerQ}[\text{p}]$

Maple [A] (verified)

Time = 0.10 (sec), antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$-\frac{\cos(x)}{a}$	8
default	$-\frac{\cos(x)}{a}$	8
risch	$-\frac{\cos(x)}{a}$	8
parallelrisch	$\frac{-\cos(x)-1}{a}$	11
norman	$\frac{2 \tan(\frac{x}{2})^7 + 2 \tan(\frac{x}{2})^3 + 4 \tan(\frac{x}{2})^5}{a \left(1+\tan(\frac{x}{2})^2\right)^3 \tan(\frac{x}{2})}$	52

input `int(sin(x)^3/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`

output $-\cos(x)/a$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a}$$

input `integrate(sin(x)^3/(a-a*cos(x)^2),x, algorithm="fricas")`

output `-cos(x)/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(5) = 10$.

Time = 0.43 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{2}{a \tan^2\left(\frac{x}{2}\right) + a}$$

input `integrate(sin(x)**3/(a-a*cos(x)**2),x)`

output `-2/(a*tan(x/2)**2 + a)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a}$$

input `integrate(sin(x)^3/(a-a*cos(x)^2),x, algorithm="maxima")`

output `-cos(x)/a`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a}$$

input `integrate(sin(x)^3/(a-a*cos(x)^2),x, algorithm="giac")`

output `-cos(x)/a`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{a}$$

input `int(sin(x)^3/(a - a*cos(x)^2),x)`

output `-cos(x)/a`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.43

$$\int \frac{\sin^3(x)}{a - a \cos^2(x)} dx = \frac{-\cos(x) + 1}{a}$$

input `int(sin(x)^3/(a-a*cos(x)^2),x)`

output `(- cos(x) + 1)/a`

3.5 $\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx$

Optimal result	72
Mathematica [A] (verified)	72
Rubi [A] (verified)	73
Maple [A] (verified)	74
Fricas [A] (verification not implemented)	74
Sympy [A] (verification not implemented)	75
Maxima [A] (verification not implemented)	75
Giac [A] (verification not implemented)	75
Mupad [B] (verification not implemented)	76
Reduce [B] (verification not implemented)	76

Optimal result

Integrand size = 16, antiderivative size = 5

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

output x/a

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input Integrate[Sin[x]^2/(a - a*Cos[x]^2), x]

output x/a

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 3654, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^2(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(x + \frac{\pi}{2})^2}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3654} \\
 & \frac{\int 1 dx}{a} \\
 & \quad \downarrow \text{24} \\
 & \frac{x}{a}
 \end{aligned}$$

input `Int[Sin[x]^2/(a - a*Cos[x]^2),x]`

output $\frac{x}{a}$

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3654

```
Int[(u_)*(a_ + b_)*sin[e_ + f_*(x_)]^2]^p_, x_Symbol] :> Simpl[
a^p Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x]; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
risch	$\frac{x}{a}$	6
default	$\frac{\arctan(\tan(x))}{a}$	8
orering	$\frac{x \sin(x)^2}{a - a \cos(x)^2}$	18
norman	$\frac{x \tan(\frac{x}{2}) + \frac{x \tan(\frac{x}{2})^5}{a} + \frac{2x \tan(\frac{x}{2})^3}{a}}{\left(1 + \tan(\frac{x}{2})^2\right)^2 \tan(\frac{x}{2})}$	51

input `int(sin(x)^2/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`output `x/a`**Fricas [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input `integrate(sin(x)^2/(a-a*cos(x)^2),x, algorithm="fricas")`output `x/a`

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.40

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input `integrate(sin(x)**2/(a-a*cos(x)**2),x)`

output `x/a`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input `integrate(sin(x)^2/(a-a*cos(x)^2),x, algorithm="maxima")`

output `x/a`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input `integrate(sin(x)^2/(a-a*cos(x)^2),x, algorithm="giac")`

output `x/a`

Mupad [B] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input `int(sin(x)^2/(a - a*cos(x)^2),x)`

output `x/a`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{\sin^2(x)}{a - a \cos^2(x)} dx = \frac{x}{a}$$

input `int(sin(x)^2/(a-a*cos(x)^2),x)`

output `x/a`

3.6 $\int \frac{\sin(x)}{a - a \cos^2(x)} dx$

Optimal result	77
Mathematica [A] (verified)	77
Rubi [A] (verified)	78
Maple [A] (verified)	79
Fricas [B] (verification not implemented)	80
Sympy [B] (verification not implemented)	80
Maxima [B] (verification not implemented)	80
Giac [B] (verification not implemented)	81
Mupad [B] (verification not implemented)	81
Reduce [B] (verification not implemented)	81

Optimal result

Integrand size = 14, antiderivative size = 8

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = -\frac{\operatorname{arctanh}(\cos(x))}{a}$$

output -arctanh(cos(x))/a

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = -\frac{\operatorname{arctanh}(\cos(x))}{a}$$

input Integrate[Sin[x]/(a - a*Cos[x]^2), x]

output -(ArcTanh[Cos[x]]/a)

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 25, 3654, 25, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\cos(x + \frac{\pi}{2})}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\cos(x + \frac{\pi}{2})}{a - a \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3654} \\
 & - \frac{\int -\csc(x) dx}{a} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \csc(x) dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \csc(x) dx}{a} \\
 & \quad \downarrow \text{4257} \\
 & - \frac{\operatorname{arctanh}(\cos(x))}{a}
 \end{aligned}$$

input `Int[Sin[x]/(a - a*Cos[x]^2),x]`

output `-(ArcTanh[Cos[x]]/a)`

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 3042 $\text{Int}[\text{u}__, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3654 $\text{Int}[(\text{u}__.)*((\text{a}__) + (\text{b}__.)*\sin[(\text{e}__.) + (\text{f}__.)*(\text{x}__.)]^2)^{\text{p}__.}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{a}^{\text{p}} \quad \text{Int}[\text{ActivateTrig}[\text{u}*\cos[\text{e} + \text{f}*\text{x}]^{(2*\text{p})}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{p}\}, \text{x}] \&& \text{EqQ}[\text{a} + \text{b}, 0] \&& \text{IntegerQ}[\text{p}]$

rule 4257 $\text{Int}[\csc[(\text{c}__.) + (\text{d}__.)*(\text{x}__.)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[\text{c} + \text{d}*\text{x}]]/\text{d}, \text{x}] /; \text{FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$

Maple [A] (verified)

Time = 0.06 (sec), antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
derivativedivides	$-\frac{\operatorname{arctanh}(\cos(x))}{a}$	9
default	$-\frac{\operatorname{arctanh}(\cos(x))}{a}$	9
norman	$\frac{\ln(\tan(\frac{x}{2}))}{a}$	10
parallelrisch	$\frac{\ln(\tan(\frac{x}{2}))}{a}$	10
risch	$-\frac{\ln(e^{ix}+1)}{a} + \frac{\ln(e^{ix}-1)}{a}$	27

input $\text{int}(\sin(x)/(a-a*\cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output $-\operatorname{arctanh}(\cos(x))/a$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(8) = 16$.

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 2.75

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = -\frac{\log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{2a}$$

input `integrate(sin(x)/(a-a*cos(x)^2),x, algorithm="fricas")`

output `-1/2*(log(1/2*cos(x) + 1/2) - log(-1/2*cos(x) + 1/2))/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(7) = 14$.

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.38

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = \frac{\log(\cos(x) - 1)}{2a} - \frac{\log(\cos(x) + 1)}{2a}$$

input `integrate(sin(x)/(a-a*cos(x)**2),x)`

output `log(cos(x) - 1)/(2*a) - log(cos(x) + 1)/(2*a)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. $2(8) = 16$.

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.62

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = -\frac{\log(\cos(x) + 1)}{2a} + \frac{\log(\cos(x) - 1)}{2a}$$

input `integrate(sin(x)/(a-a*cos(x)^2),x, algorithm="maxima")`

output
$$-1/2\log(\cos(x) + 1)/a + 1/2\log(\cos(x) - 1)/a$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 23 vs. $2(8) = 16$.

Time = 0.11 (sec), antiderivative size = 23, normalized size of antiderivative = 2.88

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = -\frac{\log(\cos(x) + 1)}{2a} + \frac{\log(-\cos(x) + 1)}{2a}$$

input `integrate(sin(x)/(a-a*cos(x)^2),x, algorithm="giac")`

output
$$-1/2\log(\cos(x) + 1)/a + 1/2\log(-\cos(x) + 1)/a$$

Mupad [B] (verification not implemented)

Time = 0.99 (sec), antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = -\frac{\operatorname{atanh}(\cos(x))}{a}$$

input `int(sin(x)/(a - a*cos(x)^2),x)`

output
$$-\operatorname{atanh}(\cos(x))/a$$

Reduce [B] (verification not implemented)

Time = 0.15 (sec), antiderivative size = 18, normalized size of antiderivative = 2.25

$$\int \frac{\sin(x)}{a - a \cos^2(x)} dx = \frac{\log(\cos(x) - 1) - \log(\cos(x) + 1)}{2a}$$

input `int(sin(x)/(a-a*cos(x)^2),x)`

output $(\log(\cos(x) - 1) - \log(\cos(x) + 1))/(2*a)$

3.7 $\int \frac{\csc(x)}{a - a \cos^2(x)} dx$

Optimal result	83
Mathematica [B] (verified)	83
Rubi [A] (verified)	84
Maple [A] (verified)	86
Fricas [B] (verification not implemented)	86
Sympy [F]	87
Maxima [B] (verification not implemented)	87
Giac [B] (verification not implemented)	87
Mupad [B] (verification not implemented)	88
Reduce [B] (verification not implemented)	88

Optimal result

Integrand size = 14, antiderivative size = 22

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = -\frac{\operatorname{arctanh}(\cos(x))}{2a} - \frac{\cot(x) \csc(x)}{2a}$$

output -1/2*arctanh(cos(x))/a-1/2*cot(x)*csc(x)/a

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. $2(22) = 44$.

Time = 0.02 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.32

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = \frac{-\frac{1}{8} \csc^2\left(\frac{x}{2}\right) - \frac{1}{2} \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{1}{2} \log\left(\sin\left(\frac{x}{2}\right)\right) + \frac{1}{8} \sec^2\left(\frac{x}{2}\right)}{a}$$

input Integrate[Csc[x]/(a - a*Cos[x]^2), x]

output (-1/8*Csc[x/2]^2 - Log[Cos[x/2]]/2 + Log[Sin[x/2]]/2 + Sec[x/2]^2/8)/a

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3042, 25, 3654, 25, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\cos(x + \frac{\pi}{2}) \left(a - a \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\cos(x + \frac{\pi}{2}) \left(a - a \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3654} \\
 & - \frac{\int -\csc^3(x) dx}{a} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & \frac{\int \csc^3(x) dx}{a} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{\int \csc(x)^3 dx}{a} \\
 & \quad \downarrow \textcolor{blue}{4255} \\
 & \frac{\frac{\int \csc(x) dx}{2} - \frac{1}{2} \cot(x) \csc(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{\frac{\int \csc(x) dx}{2} - \frac{1}{2} \cot(x) \csc(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{4257}
 \end{aligned}$$

$$\frac{-\frac{1}{2}\operatorname{arctanh}(\cos(x)) - \frac{1}{2}\cot(x)\csc(x)}{a}$$

input `Int[Csc[x]/(a - a*Cos[x]^2), x]`

output `(-1/2*ArcTanh[Cos[x]] - (Cot[x]*Csc[x])/2)/a`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3654 `Int[(u_)*(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2^(p_), x_Symbol] :> Simp[a^p Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]`

rule 4255 `Int[(csc[(c_*) + (d_)*(x_)]*(b_*))^n_, x_Symbol] :> Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[b^2*((n - 2)/(n - 1)) Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

rule 4257 `Int[csc[(c_*) + (d_)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27

method	result	size
parallelrisch	$\frac{\tan(\frac{x}{2})^2 - \cot(\frac{x}{2})^2 + 4 \ln(\tan(\frac{x}{2}))}{8a}$	28
default	$\frac{1}{4+4 \cos(x)} - \frac{\ln(1+\cos(x))}{4} + \frac{1}{-4+4 \cos(x)} + \frac{\ln(-1+\cos(x))}{4}$	36
norman	$\frac{-\frac{1}{8a} + \frac{\tan(\frac{x}{2})^4}{8a}}{\tan(\frac{x}{2})^2} + \frac{\ln(\tan(\frac{x}{2}))}{2a}$	36
risch	$\frac{e^{3ix} + e^{ix}}{(e^{2ix}-1)^2 a} + \frac{\ln(e^{ix}-1)}{2a} - \frac{\ln(e^{ix}+1)}{2a}$	52

input `int(csc(x)/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`

output `1/8*(tan(1/2*x)^2-cot(1/2*x)^2+4*ln(tan(1/2*x)))/a`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(18) = 36$.

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.18

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx \\ = -\frac{(\cos(x)^2 - 1) \log(\frac{1}{2} \cos(x) + \frac{1}{2}) - (\cos(x)^2 - 1) \log(-\frac{1}{2} \cos(x) + \frac{1}{2}) - 2 \cos(x)}{4(a \cos(x)^2 - a)}$$

input `integrate(csc(x)/(a-a*cos(x)^2),x, algorithm="fricas")`

output `-1/4*((cos(x)^2 - 1)*log(1/2*cos(x) + 1/2) - (cos(x)^2 - 1)*log(-1/2*cos(x) + 1/2) - 2*cos(x))/(a*cos(x)^2 - a)`

Sympy [F]

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = -\frac{\int \frac{\csc(x)}{\cos^2(x)-1} dx}{a}$$

input `integrate(csc(x)/(a-a*cos(x)**2),x)`

output `-Integral(csc(x)/(cos(x)**2 - 1), x)/a`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(18) = 36$.

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = \frac{\cos(x)}{2(a \cos(x)^2 - a)} - \frac{\log(\cos(x) + 1)}{4a} + \frac{\log(\cos(x) - 1)}{4a}$$

input `integrate(csc(x)/(a-a*cos(x)^2),x, algorithm="maxima")`

output `1/2*cos(x)/(a*cos(x)^2 - a) - 1/4*log(cos(x) + 1)/a + 1/4*log(cos(x) - 1)/a`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(18) = 36$.

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.73

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = -\frac{\log(\cos(x) + 1)}{4a} + \frac{\log(-\cos(x) + 1)}{4a} + \frac{\cos(x)}{2(\cos(x)^2 - 1)a}$$

input `integrate(csc(x)/(a-a*cos(x)^2),x, algorithm="giac")`

output
$$-1/4 \log(\cos(x) + 1)/a + 1/4 \log(-\cos(x) + 1)/a + 1/2 \cos(x)/((\cos(x)^2 - 1)*a)$$

Mupad [B] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = -\frac{\cos(x)}{2(a - a \cos(x)^2)} - \frac{\operatorname{atanh}(\cos(x))}{2a}$$

input `int(1/(sin(x)*(a - a*cos(x)^2)),x)`

output
$$-\cos(x)/(2*(a - a*\cos(x)^2)) - \operatorname{atanh}(\cos(x))/(2*a)$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{\csc(x)}{a - a \cos^2(x)} dx = \frac{-\cos(x) + \log(\tan(\frac{x}{2})) \sin(x)^2}{2 \sin(x)^2 a}$$

input `int(csc(x)/(a-a*cos(x)^2),x)`

output
$$(-\cos(x) + \log(\tan(x/2)) * \sin(x)^2) / (2 * \sin(x)^2 * a)$$

3.8 $\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	91
Fricas [A] (verification not implemented)	92
Sympy [F]	92
Maxima [A] (verification not implemented)	92
Giac [A] (verification not implemented)	93
Mupad [B] (verification not implemented)	93
Reduce [B] (verification not implemented)	93

Optimal result

Integrand size = 16, antiderivative size = 19

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = -\frac{\cot(x)}{a} - \frac{\cot^3(x)}{3a}$$

output $-\cot(x)/a - 1/3*\cot(x)^3/a$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = \frac{-\frac{2 \cot(x)}{3} - \frac{1}{3} \cot(x) \csc^2(x)}{a}$$

input `Integrate[Csc[x]^2/(a - a*Cos[x]^2), x]`

output $((-2*\text{Cot}[x])/3 - (\text{Cot}[x]*\text{Csc}[x]^2)/3)/a$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 3654, 3042, 4254, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc^2(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cos(x + \frac{\pi}{2})^2 \left(a - a \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3654} \\
 & \frac{\int \csc^4(x) dx}{a} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{\int \csc(x)^4 dx}{a} \\
 & \quad \downarrow \textcolor{blue}{4254} \\
 & - \frac{\int (\cot^2(x) + 1) d \cot(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & - \frac{\frac{\cot^3(x)}{3} + \cot(x)}{a}
 \end{aligned}$$

input `Int[Csc[x]^2/(a - a*Cos[x]^2),x]`

output `-((Cot[x] + Cot[x]^3/3)/a)`

Definitions of rubi rules used

rule 2009 $\text{Int}[u_., x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_., x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3654 $\text{Int}[(u_.)*((a_) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2)^{(p_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a^p \text{Int}[\text{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{EqQ}[a + b, 0] \&& \text{IntegerQ}[p]$

rule 4254 $\text{Int}[\csc[(c_.) + (d_.)*(x_.)]^{(n_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-d^{-1} \text{Subst}[\text{Int}[\text{Exp andIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&& \text{IGtQ}[n/2, 0]$

Maple [A] (verified)

Time = 0.11 (sec), antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
default	$\frac{-\frac{1}{\tan(x)} - \frac{1}{3\tan(x)^3}}{a}$	18
risch	$\frac{4i(3e^{2ix}-1)}{3(e^{2ix}-1)^3a}$	25
parallelrisch	$\frac{\tan(\frac{x}{2})^3 - \cot(\frac{x}{2})^3 + 9\tan(\frac{x}{2}) - 9\cot(\frac{x}{2})}{24a}$	33
norman	$\frac{-\frac{1}{24a} - \frac{3\tan(\frac{x}{2})^2}{8a} + \frac{3\tan(\frac{x}{2})^4}{8a} + \frac{\tan(\frac{x}{2})^6}{24a}}{\tan(\frac{x}{2})^3}$	47

input $\text{int}(\csc(x)^2/(a - a*\cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output $1/a*(-1/\tan(x) - 1/3/\tan(x)^3)$

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.53

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = -\frac{2 \cos(x)^3 - 3 \cos(x)}{3(a \cos(x)^2 - a) \sin(x)}$$

input `integrate(csc(x)^2/(a-a*cos(x)^2),x, algorithm="fricas")`

output `-1/3*(2*cos(x)^3 - 3*cos(x))/((a*cos(x)^2 - a)*sin(x))`

Sympy [F]

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = -\frac{\int \frac{\csc^2(x)}{\cos^2(x)-1} dx}{a}$$

input `integrate(csc(x)**2/(a-a*cos(x)**2),x)`

output `-Integral(csc(x)**2/(cos(x)**2 - 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = -\frac{3 \tan(x)^2 + 1}{3 a \tan(x)^3}$$

input `integrate(csc(x)^2/(a-a*cos(x)^2),x, algorithm="maxima")`

output `-1/3*(3*tan(x)^2 + 1)/(a*tan(x)^3)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = -\frac{3 \tan(x)^2 + 1}{3 a \tan(x)^3}$$

input `integrate(csc(x)^2/(a-a*cos(x)^2),x, algorithm="giac")`

output `-1/3*(3*tan(x)^2 + 1)/(a*tan(x)^3)`

Mupad [B] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = -\frac{\cot(x) (\cot(x)^2 + 3)}{3 a}$$

input `int(1/(sin(x)^2*(a - a*cos(x)^2)),x)`

output `-(cot(x)*(cot(x)^2 + 3))/(3*a)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\csc^2(x)}{a - a \cos^2(x)} dx = \frac{\cos(x) (-2 \sin(x)^2 - 1)}{3 \sin(x)^3 a}$$

input `int(csc(x)^2/(a-a*cos(x)^2),x)`

output `(cos(x)*(- 2*sin(x)**2 - 1))/(3*sin(x)**3*a)`

3.9 $\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx$

Optimal result	94
Mathematica [B] (verified)	94
Rubi [A] (verified)	95
Maple [A] (verified)	97
Fricas [B] (verification not implemented)	97
Sympy [F]	98
Maxima [A] (verification not implemented)	98
Giac [A] (verification not implemented)	99
Mupad [B] (verification not implemented)	99
Reduce [B] (verification not implemented)	99

Optimal result

Integrand size = 16, antiderivative size = 35

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx = -\frac{3 \operatorname{arctanh}(\cos(x))}{8a} - \frac{3 \cot(x) \csc(x)}{8a} - \frac{\cot(x) \csc^3(x)}{4a}$$

output
$$-3/8*\operatorname{arctanh}(\cos(x))/a-3/8*\cot(x)*\csc(x)/a-1/4*\cot(x)*\csc(x)^3/a$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 75 vs. $2(35) = 70$.

Time = 0.02 (sec), antiderivative size = 75, normalized size of antiderivative = 2.14

$$\begin{aligned} & \int \frac{\csc^3(x)}{a - a \cos^2(x)} dx \\ &= \frac{-\frac{3}{32} \csc^2\left(\frac{x}{2}\right) - \frac{1}{64} \csc^4\left(\frac{x}{2}\right) - \frac{3}{8} \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{3}{8} \log\left(\sin\left(\frac{x}{2}\right)\right) + \frac{3}{32} \sec^2\left(\frac{x}{2}\right) + \frac{1}{64} \sec^4\left(\frac{x}{2}\right)}{a} \end{aligned}$$

input
$$\text{Integrate}[\text{Csc}[x]^3/(a - a \text{Cos}[x]^2), x]$$

output $((-3*\text{Csc}[x/2]^2)/32 - \text{Csc}[x/2]^4/64 - (3*\text{Log}[\text{Cos}[x/2]])/8 + (3*\text{Log}[\text{Sin}[x/2]])/8 + (3*\text{Sec}[x/2]^2)/32 + \text{Sec}[x/2]^4/64)/a$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3042, 25, 3654, 25, 3042, 4255, 3042, 4255, 3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc^3(x)}{a - a \cos^2(x)} dx \\
 & \quad \downarrow 3042 \\
 & \int -\frac{1}{\cos(x + \frac{\pi}{2})^3 (a - a \sin(x + \frac{\pi}{2})^2)} dx \\
 & \quad \downarrow 25 \\
 & - \int \frac{1}{\cos(x + \frac{\pi}{2})^3 (a - a \sin(x + \frac{\pi}{2})^2)} dx \\
 & \quad \downarrow 3654 \\
 & - \frac{\int -\csc^5(x) dx}{a} \\
 & \quad \downarrow 25 \\
 & \frac{\int \csc^5(x) dx}{a} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \csc(x)^5 dx}{a} \\
 & \quad \downarrow 4255 \\
 & \frac{\frac{3}{4} \int \csc^3(x) dx - \frac{1}{4} \cot(x) \csc^3(x)}{a} \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$\begin{array}{c}
 \frac{\frac{3}{4} \int \csc(x)^3 dx - \frac{1}{4} \cot(x) \csc^3(x)}{a} \\
 \downarrow \textcolor{blue}{4255} \\
 \frac{\frac{3}{4} \left(\frac{\int \csc(x) dx}{2} - \frac{1}{2} \cot(x) \csc(x) \right) - \frac{1}{4} \cot(x) \csc^3(x)}{a} \\
 \downarrow \textcolor{blue}{3042} \\
 \frac{\frac{3}{4} \left(\frac{\int \csc(x) dx}{2} - \frac{1}{2} \cot(x) \csc(x) \right) - \frac{1}{4} \cot(x) \csc^3(x)}{a} \\
 \downarrow \textcolor{blue}{4257} \\
 \frac{\frac{3}{4} \left(-\frac{1}{2} \operatorname{arctanh}(\cos(x)) - \frac{1}{2} \cot(x) \csc(x) \right) - \frac{1}{4} \cot(x) \csc^3(x)}{a}
 \end{array}$$

input `Int[Csc[x]^3/(a - a*Cos[x]^2), x]`

output `(-1/4*(Cot[x]*Csc[x]^3) + (3*(-1/2*ArcTanh[Cos[x]] - (Cot[x]*Csc[x])/2))/4)/a`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3654 `Int[(u_)*(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2^(p_), x_Symbol] :> Simp[a^p Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]`

rule 4255 $\text{Int}[(\csc(c_.) + (d_.)*(x_.))*(b_.)]^{(n_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(-b)*\cos[c + d*x]*((b*\csc[c + d*x])^{(n - 1)}/(d*(n - 1))), \text{x}] + \text{Simp}[b^{2*((n - 2)/(n - 1))}\text{Int}[(b*\csc[c + d*x])^{(n - 2)}, \text{x}], \text{x}] /; \text{FreeQ}[\{b, c, d\}, \text{x}] \&& \text{GtQ}[n, 1] \&& \text{IntegerQ}[2*n]$

rule 4257 $\text{Int}[\csc(c_.) + (d_.)*(x_.)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{ArcTanh}[\cos[c + d*x]]/d, \text{x}] /; \text{FreeQ}[\{c, d\}, \text{x}]$

Maple [A] (verified)

Time = 0.16 (sec), antiderivative size = 44, normalized size of antiderivative = 1.26

method	result	size
parallelrisch	$\frac{-\cot(\frac{x}{2})^4 + \tan(\frac{x}{2})^4 - 8\cot(\frac{x}{2})^2 + 8\tan(\frac{x}{2})^2 + 24\ln(\tan(\frac{x}{2}))}{64a}$	44
default	$\frac{\frac{1}{16(1+\cos(x))^2} + \frac{3}{16(1+\cos(x))} - \frac{3\ln(1+\cos(x))}{16} - \frac{1}{16(-1+\cos(x))^2} + \frac{3}{16(-1+\cos(x))} + \frac{3\ln(-1+\cos(x))}{16}}{a}$	52
norman	$\frac{-\frac{1}{64a} - \frac{\tan(\frac{x}{2})^2}{8a} + \frac{\tan(\frac{x}{2})^6}{8a} + \frac{\tan(\frac{x}{2})^8}{64a}}{\tan(\frac{x}{2})^4} + \frac{3\ln(\tan(\frac{x}{2}))}{8a}$	58
risch	$\frac{3e^{7ix} - 11e^{5ix} - 11e^{3ix} + 3e^{ix}}{4(e^{2ix} - 1)^4a} + \frac{3\ln(e^{ix} - 1)}{8a} - \frac{3\ln(e^{ix} + 1)}{8a}$	71

input `int(csc(x)^3/(a-a*cos(x)^2),x,method=_RETURNVERBOSE)`

output $\frac{1/64*(-\cot(1/2*x)^4 + \tan(1/2*x)^4 - 8*\cot(1/2*x)^2 + 8*\tan(1/2*x)^2 + 24*\ln(\tan(1/2*x)))/a}{a}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. $2(29) = 58$.

Time = 0.10 (sec), antiderivative size = 72, normalized size of antiderivative = 2.06

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx \\ = \frac{6 \cos(x)^3 - 3 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(\frac{1}{2} \cos(x) + \frac{1}{2}) + 3 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(-\frac{1}{2} \cos(x) + \frac{1}{2})}{16 (a \cos(x)^4 - 2 a \cos(x)^2 + a)}$$

input `integrate(csc(x)^3/(a-a*cos(x)^2),x, algorithm="fricas")`

output $\frac{1}{16} (6 \cos(x)^3 - 3 \cos(x)^4 - 2 \cos(x)^2 + 1) \log(1/2 \cos(x) + 1/2) + 3 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(-1/2 \cos(x) + 1/2) - 10 \cos(x)) / (a \cos(x)^4 - 2 a \cos(x)^2 + a)$

Sympy [F]

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx = -\frac{\int \frac{\csc^3(x)}{\cos^2(x)-1} dx}{a}$$

input `integrate(csc(x)**3/(a-a*cos(x)**2),x)`

output `-Integral(csc(x)**3/(cos(x)**2 - 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.46

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx = \frac{3 \cos(x)^3 - 5 \cos(x)}{8 (a \cos(x)^4 - 2 a \cos(x)^2 + a)} - \frac{3 \log(\cos(x) + 1)}{16 a} + \frac{3 \log(\cos(x) - 1)}{16 a}$$

input `integrate(csc(x)^3/(a-a*cos(x)^2),x, algorithm="maxima")`

output $\frac{1}{8} (3 \cos(x)^3 - 5 \cos(x)) / (a \cos(x)^4 - 2 a \cos(x)^2 + a) - \frac{3}{16} \log(\cos(x) + 1) / a + \frac{3}{16} \log(\cos(x) - 1) / a$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.34

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx = -\frac{3 \log(\cos(x) + 1)}{16a} + \frac{3 \log(-\cos(x) + 1)}{16a} + \frac{3 \cos(x)^3 - 5 \cos(x)}{8 (\cos(x)^2 - 1)^2 a}$$

input `integrate(csc(x)^3/(a-a*cos(x)^2),x, algorithm="giac")`

output `-3/16*log(cos(x) + 1)/a + 3/16*log(-cos(x) + 1)/a + 1/8*(3*cos(x)^3 - 5*cos(x))/((cos(x)^2 - 1)^2*a)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.11

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx = -\frac{3 \operatorname{atanh}(\cos(x))}{8a} - \frac{\frac{5 \cos(x)}{8} - \frac{3 \cos(x)^3}{8}}{a \cos(x)^4 - 2a \cos(x)^2 + a}$$

input `int(1/(sin(x)^3*(a - a*cos(x)^2)),x)`

output `- (3*atanh(cos(x)))/(8*a) - ((5*cos(x))/8 - (3*cos(x)^3)/8)/(a - 2*a*cos(x)^2 + a*cos(x)^4)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\csc^3(x)}{a - a \cos^2(x)} dx = \frac{-3 \cos(x) \sin(x)^2 - 2 \cos(x) + 3 \log(\tan(\frac{x}{2})) \sin(x)^4}{8 \sin(x)^4 a}$$

input `int(csc(x)^3/(a-a*cos(x)^2),x)`

output $(- 3\cos(x)\sin(x)^2 - 2\cos(x) + 3\log(\tan(x/2))\sin(x)^4)/(8\sin(x)^{**}4*a)$

3.10 $\int \frac{\sin^7(x)}{a+b\cos^2(x)} dx$

Optimal result	101
Mathematica [A] (verified)	101
Rubi [A] (verified)	102
Maple [A] (verified)	104
Fricas [A] (verification not implemented)	104
Sympy [F(-1)]	105
Maxima [A] (verification not implemented)	105
Giac [A] (verification not implemented)	105
Mupad [B] (verification not implemented)	106
Reduce [B] (verification not implemented)	106

Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{\sin^7(x)}{a+b\cos^2(x)} dx = -\frac{(a+b)^3 \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} + \frac{(a^2+3ab+3b^2)\cos(x)}{b^3} - \frac{(a+3b)\cos^3(x)}{3b^2} + \frac{\cos^5(x)}{5b}$$

output $-(a+b)^3 \arctan(b^{(1/2)} \cos(x)/a^{(1/2)})/a^{(1/2)}/b^{(7/2)} + (a^2+3*a*b+3*b^2)*\cos(x)/b^{3-1/3*(a+3*b)}*\cos(x)^3/b^{2+1/5*\cos(x)^5}/b$

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.83

$$\int \frac{\sin^7(x)}{a+b\cos^2(x)} dx = -\frac{(a+b)^3 \arctan\left(\frac{\sqrt{b}-\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} - \frac{(a+b)^3 \arctan\left(\frac{\sqrt{b}+\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} + \frac{(8a^2+22ab+19b^2)\cos(x)}{8b^3} - \frac{(4a+9b)\cos(3x)}{48b^2} + \frac{\cos(5x)}{80b}$$

input `Integrate[Sin[x]^7/(a + b*Cos[x]^2), x]`

output
$$\begin{aligned} & -((a+b)^3 \operatorname{ArcTan}\left[\frac{\sqrt{b}-\sqrt{a+b} \tan \left(\frac{x}{2}\right)}{\sqrt{a}}\right]) / (\sqrt{a} * b^{(7/2)}) \\ & -((a+b)^3 \operatorname{ArcTan}\left[\frac{\sqrt{b}+\sqrt{a+b} \tan \left(\frac{x}{2}\right)}{\sqrt{a}}\right]) / (\sqrt{a} * b^{(7/2)}) \\ & + \frac{(8 a^2+22 a b+19 b^2) \cos (x)}{8 b^3} - \frac{(4 a+9 b) \cos (3 x)}{48 b^2} + \frac{\cos (5 x)}{80 b} \end{aligned}$$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 25, 3669, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin^7(x)}{a + b \cos^2(x)} dx \\ & \quad \downarrow 3042 \\ & \int -\frac{\cos\left(x + \frac{\pi}{2}\right)^7}{a + b \sin\left(x + \frac{\pi}{2}\right)^2} dx \\ & \quad \downarrow 25 \\ & - \int \frac{\cos\left(x + \frac{\pi}{2}\right)^7}{b \sin\left(x + \frac{\pi}{2}\right)^2 + a} dx \\ & \quad \downarrow 3669 \\ & - \int \frac{(1 - \cos^2(x))^3}{b \cos^2(x) + a} d \cos(x) \\ & \quad \downarrow 300 \\ & - \int \left(-\frac{\cos^4(x)}{b} + \frac{(a+3b)\cos^2(x)}{b^2} - \frac{a^2+3ba+3b^2}{b^3} + \frac{a^3+3ba^2+3b^2a+b^3}{b^3(b \cos^2(x) + a)} \right) d \cos(x) \\ & \quad \downarrow 2009 \end{aligned}$$

$$\frac{(a^2 + 3ab + 3b^2) \cos(x)}{b^3} - \frac{(a+b)^3 \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{ab^{7/2}}} - \frac{(a+3b) \cos^3(x)}{3b^2} + \frac{\cos^5(x)}{5b}$$

input `Int[Sin[x]^7/(a + b*Cos[x]^2), x]`

output
$$-\frac{((a+b)^3 \operatorname{ArcTan}\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right))}{(b^3 \sqrt{ab^{7/2}})} + \frac{(a^2 + 3ab + 3b^2) \cos(x)}{b^3} - \frac{(a+3b) \cos^3(x)}{3b^2} + \frac{\cos^5(x)}{5b}$$

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 300 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] :> Int[PolynomialDivide[(a + b*x^2)^p, (c + d*x^2)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3669 `Int[cos[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f S ubst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]]`

Maple [A] (verified)

Time = 1.94 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{\frac{b^2 \cos(x)^5}{5} - \frac{ab \cos(x)^3}{3} - b^2 \cos(x)^3 + a^2 \cos(x) + 3 \cos(x)ab + 3 \cos(x)b^2}{b^3} + \frac{(-a^3 - 3a^2b - 3ab^2 - b^3) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{b^3 \sqrt{ab}}$
default	$\frac{\frac{b^2 \cos(x)^5}{5} - \frac{ab \cos(x)^3}{3} - b^2 \cos(x)^3 + a^2 \cos(x) + 3 \cos(x)ab + 3 \cos(x)b^2}{b^3} + \frac{(-a^3 - 3a^2b - 3ab^2 - b^3) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{b^3 \sqrt{ab}}$
risch	$\frac{e^{ix}a^2}{2b^3} + \frac{11e^{ix}a}{8b^2} + \frac{19e^{ix}}{16b} + \frac{e^{-ix}a^2}{2b^3} + \frac{11e^{-ix}a}{8b^2} + \frac{19e^{-ix}}{16b} - \frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)a^3}{2\sqrt{ab}b^3} - \frac{3i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)}{2\sqrt{ab}b}$

input `int(sin(x)^7/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{b^3} \left(\frac{1}{5} b^2 \cos(x)^5 - 1/3 a b \cos(x)^3 - b^2 \cos(x)^3 + a^2 \cos(x) + 3 \cos(x) a b + 3 \cos(x) b^2 + (-a^3 - 3 a^2 b - 3 a b^2 - b^3) / b^3 \right) / (a b)^{(1/2)} \arctan(b \cos(x)) / (a b)^{(1/2)}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.88

$$\int \frac{\sin^7(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{6 ab^3 \cos(x)^5 - 10 (a^2 b^2 + 3 ab^3) \cos(x)^3 - 15 (a^3 + 3 a^2 b + 3 ab^2 + b^3) \sqrt{-ab} \log\left(-\frac{b \cos(x)^2 + 2 \sqrt{-ab} \cos(x)}{b \cos(x)^2 + a}\right)}{30 ab^4} \right]$$

input `integrate(sin(x)^7/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[1/30 * (6*a*b^3*cos(x)^5 - 10*(a^2*b^2 + 3*a*b^3)*cos(x)^3 - 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sqrt(-a*b)*log(-(b*cos(x)^2 + 2*sqrt(-a*b)*cos(x) - a)/(b*cos(x)^2 + a)) + 30*(a^3*b + 3*a^2*b^2 + 3*a*b^3)*cos(x))/(a*b^4), 1/15 * (3*a*b^3*cos(x)^5 - 5*(a^2*b^2 + 3*a*b^3)*cos(x)^3 - 15*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sqrt(a*b)*arctan(sqrt(a*b)*cos(x)/a) + 15*(a^3*b + 3*a^2*b^2 + 3*a*b^3)*cos(x))/(a*b^4)]$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin^7(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sin(x)**7/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.12

$$\int \frac{\sin^7(x)}{a + b \cos^2(x)} dx = -\frac{(a^3 + 3 a^2 b + 3 a b^2 + b^3) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab} b^3} + \frac{3 b^2 \cos(x)^5 - 5 (ab + 3 b^2) \cos(x)^3 + 15 (a^2 + 3 ab + 3 b^2) \cos(x)}{15 b^3}$$

input `integrate(sin(x)^7/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 1/15*(3*b^2*cos(x)^5 - 5*(a*b + 3*b^2)*cos(x)^3 + 15*(a^2 + 3*a*b + 3*b^2)*cos(x))/b^3`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.27

$$\int \frac{\sin^7(x)}{a + b \cos^2(x)} dx = -\frac{(a^3 + 3 a^2 b + 3 a b^2 + b^3) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab} b^3} + \frac{3 b^4 \cos(x)^5 - 5 a b^3 \cos(x)^3 - 15 b^4 \cos(x)^3 + 15 a^2 b^2 \cos(x) + 45 a b^3 \cos(x) + 45 b^4 \cos(x)}{15 b^5}$$

input `integrate(sin(x)^7/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$\begin{aligned} & -(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\arctan(b*\cos(x)/\sqrt{a*b})/(\sqrt{a*b}*b^3) \\ & + 1/15*(3*b^4*\cos(x)^5 - 5*a*b^3*\cos(x)^3 - 15*b^4*\cos(x)^3 + 15*a^2*b^2 \\ & *\cos(x) + 45*a*b^3*\cos(x) + 45*b^4*\cos(x))/b^5 \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.28

$$\begin{aligned} \int \frac{\sin^7(x)}{a + b \cos^2(x)} dx = & \cos(x) \left(\frac{3}{b} + \frac{a \left(\frac{a}{b^2} + \frac{3}{b} \right)}{b} \right) - \cos(x)^3 \left(\frac{a}{3b^2} + \frac{1}{b} \right) \\ & + \frac{\cos(x)^5}{5b} - \frac{\operatorname{atan}\left(\frac{\sqrt{b} \cos(x) (a+b)^3}{\sqrt{a} (a^3+3 a^2 b+3 a b^2+b^3)}\right) (a+b)^3}{\sqrt{a} b^{7/2}} \end{aligned}$$

input `int(sin(x)^7/(a + b*cos(x)^2),x)`

output
$$\begin{aligned} & \cos(x)*(3/b + (a*(a/b^2 + 3/b))/b) - \cos(x)^3*(a/(3*b^2) + 1/b) + \cos(x)^5 \\ & /(5*b) - (\operatorname{atan}((b^{1/2})*\cos(x)*(a + b)^3)/(a^{1/2})*(3*a*b^2 + 3*a^2*b + a^3 + b^3)) * (a + b)^3 / (a^{1/2})*b^{(7/2)}) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 321, normalized size of antiderivative = 4.12

$$\begin{aligned} \int \frac{\sin^7(x)}{a + b \cos^2(x)} dx = & \frac{15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})-\sqrt{b}}{\sqrt{a}}\right)a^3 + 45\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})-\sqrt{b}}{\sqrt{a}}\right)a^2b + 45\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})-\sqrt{b}}{\sqrt{a}}\right)a^3b^2}{a^3b^2} \end{aligned}$$

input `int(sin(x)^7/(a+b*cos(x)^2),x)`

output

$$\begin{aligned} & (15*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a^{**3} + \\ & 45*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a^{**2}*b + \\ & 45*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a*b^{**2} \\ & + 15*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*b^{**3} - \\ & 15*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a^{**3} - \\ & 45*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a^{**2}*b - \\ & 45*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a*b^{**2} \\ & - 15*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*b^{**3} + \\ & 3*\cos(x)*\sin(x)^{**4}*a*b^{**3} + 5*\cos(x)*\sin(x)^{**2}*a^{**2}*b^{**2} + 9*\cos(x)*\sin(x) \\ &)^{**2}*a*b^{**3} + 15*\cos(x)*a^{**3}*b + 40*\cos(x)*a^{**2}*b^{**2} + 33*\cos(x)*a*b^{**3} - \\ & 15*a^{**3}*b - 40*a^{**2}*b^{**2} - 33*a*b^{**3})/(15*a*b^{**4}) \end{aligned}$$

3.11 $\int \frac{\sin^5(x)}{a+b\cos^2(x)} dx$

Optimal result	108
Mathematica [B] (verified)	108
Rubi [A] (verified)	109
Maple [A] (verified)	110
Fricas [A] (verification not implemented)	111
Sympy [F(-1)]	111
Maxima [A] (verification not implemented)	112
Giac [A] (verification not implemented)	112
Mupad [B] (verification not implemented)	112
Reduce [B] (verification not implemented)	113

Optimal result

Integrand size = 15, antiderivative size = 54

$$\int \frac{\sin^5(x)}{a+b\cos^2(x)} dx = -\frac{(a+b)^2 \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{ab^{5/2}}} + \frac{(a+2b)\cos(x)}{b^2} - \frac{\cos^3(x)}{3b}$$

output

$-(a+b)^2 \arctan(b^{(1/2)} \cos(x)/a^{(1/2)})/a^{(1/2)}/b^{(5/2)} + (a+2b)*\cos(x)/b^2 - 1/3*\cos(x)^3/b$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 116 vs. $2(54) = 108$.

Time = 0.27 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.15

$$\begin{aligned} & \int \frac{\sin^5(x)}{a+b\cos^2(x)} dx \\ &= \frac{-\frac{12(a+b)^2 \arctan\left(\frac{\sqrt{b}-\sqrt{a+b}\tan\left(\frac{x}{2}\right)}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{12(a+b)^2 \arctan\left(\frac{\sqrt{b}+\sqrt{a+b}\tan\left(\frac{x}{2}\right)}{\sqrt{a}}\right)}{\sqrt{a}} + 3\sqrt{b}(4a+7b)\cos(x) - b^{3/2}\cos(3x)}{12b^{5/2}} \end{aligned}$$

input

```
Integrate[Sin[x]^5/(a + b*Cos[x]^2), x]
```

output

$$\begin{aligned} & ((-12*(a + b)^2*\text{ArcTan}[(\text{Sqrt}[b] - \text{Sqrt}[a + b]*\text{Tan}[x/2])/\text{Sqrt}[a]])/\text{Sqrt}[a] \\ & - (12*(a + b)^2*\text{ArcTan}[(\text{Sqrt}[b] + \text{Sqrt}[a + b]*\text{Tan}[x/2])/\text{Sqrt}[a]])/\text{Sqrt}[a] \\ & + 3*\text{Sqrt}[b]*(4*a + 7*b)*\text{Cos}[x] - b^{(3/2)}*\text{Cos}[3*x])/((12*b^{(5/2)}) \end{aligned}$$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 25, 3669, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sin^5(x)}{a + b \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{\cos(x + \frac{\pi}{2})^5}{a + b \sin(x + \frac{\pi}{2})^2} dx \\ & \quad \downarrow \text{25} \\ & - \int \frac{\cos(x + \frac{\pi}{2})^5}{b \sin(x + \frac{\pi}{2})^2 + a} dx \\ & \quad \downarrow \text{3669} \\ & - \int \frac{(1 - \cos^2(x))^2}{b \cos^2(x) + a} d \cos(x) \\ & \quad \downarrow \text{300} \\ & - \int \left(\frac{\cos^2(x)}{b} - \frac{a + 2b}{b^2} + \frac{a^2 + 2ba + b^2}{b^2(b \cos^2(x) + a)} \right) d \cos(x) \\ & \quad \downarrow \text{2009} \\ & - \frac{(a + b)^2 \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{5/2}} + \frac{(a + 2b) \cos(x)}{b^2} - \frac{\cos^3(x)}{3b} \end{aligned}$$

input

$$\text{Int}[\text{Sin}[x]^5/(a + b*\text{Cos}[x]^2), x]$$

output
$$-\frac{((a+b)^2 \operatorname{ArcTan}\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right))}{\sqrt{a} b^{5/2}} + \frac{(a+2b) \cos(x)}{b^2} - \frac{\cos(x)^3}{3b}$$

Definitions of rubi rules used

rule 25
$$\operatorname{Int}[-(F_x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$$

rule 300
$$\operatorname{Int}[(a_+ + b_-)(x_-)^2)^{(p_-)} ((c_+ + d_-)(x_-)^2)^{(q_-)}, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[(a+b*x^2)^p, (c+d*x^2)^{-q}], x, x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \& \operatorname{NeQ}[b*c - a*d, 0] \& \operatorname{IGtQ}[p, 0] \& \operatorname{ILtQ}[q, 0] \& \operatorname{GeQ}[p, -q]$$

rule 2009
$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 3042
$$\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3669
$$\operatorname{Int}[\cos[(e_- + f_-)(x_-)]^{(m_-)} ((a_+ + b_-) \sin[(e_- + f_-)(x_-)]^2)^{(p_-)}, x_{\text{Symbol}}] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Sin}[e + f*x], x]\}, \operatorname{Simp}[ff/f S \operatorname{ubst}[\operatorname{Int}[(1 - ff^2 x^2)^{((m-1)/2)} ((a + b*ff^2 x^2)^p, x], x, \operatorname{Sin}[e + f*x]/ff], x]] /; \operatorname{FreeQ}[\{a, b, e, f, p\}, x] \& \operatorname{IntegerQ}[(m-1)/2]$$

Maple [A] (verified)

Time = 0.79 (sec), antiderivative size = 57, normalized size of antiderivative = 1.06

method	result
derivativedivides	$\frac{-\frac{b \cos(x)^3}{3} + \cos(x)a + 2b \cos(x)}{b^2} + \frac{(-a^2 - 2ab - b^2) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{b^2 \sqrt{ab}}$
default	$\frac{-\frac{b \cos(x)^3}{3} + \cos(x)a + 2b \cos(x)}{b^2} + \frac{(-a^2 - 2ab - b^2) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{b^2 \sqrt{ab}}$
risch	$\frac{e^{ix}a}{2b^2} + \frac{7e^{ix}}{8b} + \frac{e^{-ix}a}{2b^2} + \frac{7e^{-ix}}{8b} - \frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right) a^2}{2\sqrt{ab} b^2} - \frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right) a}{\sqrt{ab} b} - \frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)}{2\sqrt{ab}}$

input `int(sin(x)^5/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output $\frac{1/b^2*(-1/3*b*cos(x)^3+cos(x)*a+2*b*cos(x))+(-a^2-2*a*b-b^2)/b^2/(a*b)^(1/2)*arctan(b*cos(x)/(a*b)^(1/2))}{}$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.81

$$\begin{aligned} & \int \frac{\sin^5(x)}{a + b \cos^2(x)} dx \\ &= \left[-\frac{2 ab^2 \cos(x)^3 + 3 (a^2 + 2 ab + b^2) \sqrt{-ab} \log\left(\frac{-b \cos(x)^2 + 2 \sqrt{-ab} \cos(x) - a}{b \cos(x)^2 + a}\right) - 6 (a^2 b + 2 ab^2) \cos(x)}{6 ab^3} \right. \\ & \quad \left. - \frac{ab^2 \cos(x)^3 + 3 (a^2 + 2 ab + b^2) \sqrt{ab} \arctan\left(\frac{\sqrt{ab} \cos(x)}{a}\right) - 3 (a^2 b + 2 ab^2) \cos(x)}{3 ab^3} \right] \end{aligned}$$

input `integrate(sin(x)^5/(a+b*cos(x)^2),x, algorithm="fricas")`

output $[-1/6*(2*a*b^2*cos(x)^3 + 3*(a^2 + 2*a*b + b^2)*sqrt(-a*b)*log(-(b*cos(x)^2 + 2*sqrt(-a*b)*cos(x) - a)/(b*cos(x)^2 + a)) - 6*(a^2*b + 2*a*b^2)*cos(x))/(a*b^3), -1/3*(a*b^2*cos(x)^3 + 3*(a^2 + 2*a*b + b^2)*sqrt(a*b)*arctan(sqrt(a*b)*cos(x)/a) - 3*(a^2*b + 2*a*b^2)*cos(x))/(a*b^3)]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin^5(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sin(x)**5/(a+b*cos(x)**2),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.98

$$\int \frac{\sin^5(x)}{a + b \cos^2(x)} dx = -\frac{(a^2 + 2ab + b^2) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}b^2} - \frac{b \cos(x)^3 - 3(a + 2b) \cos(x)}{3b^2}$$

input `integrate(sin(x)^5/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-(a^2 + 2*a*b + b^2)*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*b^2) - 1/3*(b*c
os(x)^3 - 3*(a + 2*b)*cos(x))/b^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.09

$$\int \frac{\sin^5(x)}{a + b \cos^2(x)} dx = -\frac{(a^2 + 2ab + b^2) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}b^2} - \frac{b^2 \cos(x)^3 - 3ab \cos(x) - 6b^2 \cos(x)}{3b^3}$$

input `integrate(sin(x)^5/(a+b*cos(x)^2),x, algorithm="giac")`

output `-(a^2 + 2*a*b + b^2)*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*b^2) - 1/3*(b^2
*cos(x)^3 - 3*a*b*cos(x) - 6*b^2*cos(x))/b^3`

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.20

$$\int \frac{\sin^5(x)}{a + b \cos^2(x)} dx = \cos(x) \left(\frac{a}{b^2} + \frac{2}{b} \right) - \frac{\cos(x)^3}{3b} - \frac{\operatorname{atan}\left(\frac{\sqrt{b} \cos(x) (a+b)^2}{\sqrt{a} (a^2+2ab+b^2)}\right) (a+b)^2}{\sqrt{a} b^{5/2}}$$

input `int(sin(x)^5/(a + b*cos(x)^2),x)`

output $\cos(x) * (a/b^2 + 2/b) - \cos(x)^3/(3*b) - (\text{atan}((b^{1/2}) * \cos(x) * (a + b)^2) / (a^{1/2} * (2*a*b + a^2 + b^2))) * (a + b)^2 / (a^{1/2} * b^{(5/2)})$

Reduce [B] (verification not implemented)

Time = 0.16 (sec), antiderivative size = 214, normalized size of antiderivative = 3.96

$$\int \frac{\sin^5(x)}{a + b \cos^2(x)} dx \\ = \frac{3\sqrt{b}\sqrt{a} \text{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) a^2 + 6\sqrt{b}\sqrt{a} \text{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) ab + 3\sqrt{b}\sqrt{a} \text{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) b^2}{}$$

input `int(sin(x)^5/(a+b*cos(x)^2),x)`

output $(3*\sqrt{b}*\sqrt{a}*\text{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a^{**2} + 6*\sqrt{b}*\sqrt{a}*\text{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a*b + 3*\sqrt{b}*\sqrt{a}*\text{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*b^{**2} - 3*\sqrt{b}*\sqrt{a}*\text{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a^{**2} - 6*\sqrt{b}*\sqrt{a}*\text{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a*b - 3*\sqrt{b}*\sqrt{a}*\text{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*b^{**2} + \cos(x)*\sin(x)**2*a*b^{**2} + 3*\cos(x)*a^{**2}*b + 5*\cos(x)*a*b^{**2} - 3*a^{**2}*b - 5*a*b^{**2})/(3*a*b^{**3})$

3.12 $\int \frac{\sin^3(x)}{a+b\cos^2(x)} dx$

Optimal result	114
Mathematica [B] (verified)	114
Rubi [A] (verified)	115
Maple [A] (verified)	116
Fricas [A] (verification not implemented)	117
Sympy [F(-1)]	117
Maxima [A] (verification not implemented)	118
Giac [A] (verification not implemented)	118
Mupad [B] (verification not implemented)	118
Reduce [B] (verification not implemented)	119

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \frac{\sin^3(x)}{a+b\cos^2(x)} dx = -\frac{(a+b)\arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{3/2}} + \frac{\cos(x)}{b}$$

output $-(a+b)*\arctan(b^{(1/2)}*\cos(x)/a^{(1/2)})/a^{(1/2)}/b^{(3/2)}+\cos(x)/b$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 90 vs. $2(36) = 72$.

Time = 0.25 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.50

$$\begin{aligned} & \int \frac{\sin^3(x)}{a+b\cos^2(x)} dx \\ &= \frac{-\left((a+b)\arctan\left(\frac{\sqrt{b}-\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right)\right) - (a+b)\arctan\left(\frac{\sqrt{b}+\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right) + \sqrt{a}\sqrt{b}\cos(x)}{\sqrt{ab}^{3/2}} \end{aligned}$$

input `Integrate[Sin[x]^3/(a + b*Cos[x]^2), x]`

output
$$\left(-((a + b) \operatorname{ArcTan}\left[(\sqrt{b} - \sqrt{a + b} \tan(x/2)) / \sqrt{a} \right]) - (a + b) \operatorname{ArcTan}\left[(\sqrt{b} + \sqrt{a + b} \tan(x/2)) / \sqrt{a} \right] + \sqrt{a} \sqrt{b} \cos(x) \right) / (\sqrt{a} b^{3/2})$$

Rubi [A] (verified)

Time = 0.23 (sec), antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 25, 3669, 299, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^3(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\cos\left(x + \frac{\pi}{2}\right)^3}{a + b \sin\left(x + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\cos\left(x + \frac{\pi}{2}\right)^3}{b \sin\left(x + \frac{\pi}{2}\right)^2 + a} dx \\
 & \quad \downarrow \text{3669} \\
 & - \int \frac{1 - \cos^2(x)}{b \cos^2(x) + a} d \cos(x) \\
 & \quad \downarrow \text{299} \\
 & \frac{\cos(x)}{b} - \frac{(a + b) \int \frac{1}{b \cos^2(x) + a} d \cos(x)}{b} \\
 & \quad \downarrow \text{218} \\
 & \frac{\cos(x)}{b} - \frac{(a + b) \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{ab^{3/2}}}
 \end{aligned}$$

input
$$\operatorname{Int}[\operatorname{Sin}[x]^3 / (a + b \operatorname{Cos}[x]^2), x]$$

output $-\frac{((a+b) \operatorname{ArcTan}\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right))}{\sqrt{a} b^{3/2}} + \frac{\cos(x)}{b}$

Definitions of rubi rules used

rule 25 $\operatorname{Int}[-(F_{x_}), x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$

rule 218 $\operatorname{Int}[(a_+ + b_-)(x_-)^2, x_Symbol] \rightarrow \operatorname{Simp}[(Rt[a/b, 2]/a) \operatorname{ArcTan}[x/Rt[a/b, 2]], x]; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{PosQ}[a/b]$

rule 299 $\operatorname{Int}[(a_+ + b_-)(x_-)^2(p_+ + d_-)(c_+ + d_-)(x_-)^2, x_Symbol] \rightarrow \operatorname{Simp}[d*x * ((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - \operatorname{Simp}[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) \operatorname{Int}[(a + b*x^2)^p, x], x]; \operatorname{FreeQ}[\{a, b, c, d\}, x] \& \operatorname{NeQ}[b*c - a*d, 0] \& \operatorname{NeQ}[2*p + 3, 0]$

rule 3042 $\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x]; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3669 $\operatorname{Int}[\cos(e_+ + f_-)(x_-)^m(a_+ + b_-)\sin(e_+ + f_-)(x_-)^2, x_Symbol] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Sin}[e + f*x], x]\}, \operatorname{Simp}[ff/f S ubst[\operatorname{Int}[(1 - ff^2*x^2)^{(m - 1)/2}*(a + b*ff^2*x^2)^p, x], x, \operatorname{Sin}[e + f*x]/ff], x]]; \operatorname{FreeQ}[\{a, b, e, f, p\}, x] \& \operatorname{IntegerQ}[(m - 1)/2]$

Maple [A] (verified)

Time = 0.31 (sec), antiderivative size = 34, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\cos(x)}{b} + \frac{(-a-b) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{b \sqrt{ab}}$
default	$\frac{\cos(x)}{b} + \frac{(-a-b) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{b \sqrt{ab}}$
risch	$\frac{e^{ix}}{2b} + \frac{e^{-ix}}{2b} + \frac{i \ln\left(e^{2ix} - \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right) a}{2\sqrt{ab}} + \frac{i \ln\left(e^{2ix} - \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)}{2\sqrt{ab}} - \frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right) a}{2\sqrt{ab}} - \frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)}{2\sqrt{ab}}$

input `int(sin(x)^3/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output `cos(x)/b+(-a-b)/b/(a*b)^(1/2)*arctan(b*cos(x)/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.64

$$\int \frac{\sin^3(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{2 ab \cos(x) - \sqrt{-ab}(a + b) \log\left(\frac{-b \cos(x)^2 + 2\sqrt{-ab} \cos(x) - a}{b \cos(x)^2 + a}\right)}{2 ab^2}, \frac{ab \cos(x) - \sqrt{ab}(a + b) \arctan\left(\frac{\sqrt{ab} \cos(x)}{a}\right)}{ab^2} \right]$$

input `integrate(sin(x)^3/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[1/2*(2*a*b*cos(x) - sqrt(-a*b)*(a + b)*log(-(b*cos(x)^2 + 2*sqrt(-a*b)*cos(x) - a)/(b*cos(x)^2 + a)))/(a*b^2), (a*b*cos(x) - sqrt(a*b)*(a + b)*arctan(sqrt(a*b)*cos(x)/a))/(a*b^2)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin^3(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sin(x)**3/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \frac{\sin^3(x)}{a + b \cos^2(x)} dx = -\frac{(a + b) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}} + \frac{\cos(x)}{b}$$

input `integrate(sin(x)^3/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-(a + b)*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*b) + cos(x)/b`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int \frac{\sin^3(x)}{a + b \cos^2(x)} dx = -\frac{(a + b) \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}} + \frac{\cos(x)}{b}$$

input `integrate(sin(x)^3/(a+b*cos(x)^2),x, algorithm="giac")`

output `-(a + b)*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*b) + cos(x)/b`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

$$\int \frac{\sin^3(x)}{a + b \cos^2(x)} dx = \frac{\cos(x)}{b} - \frac{\operatorname{atan}\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right) (a + b)}{\sqrt{a} b^{3/2}}$$

input `int(sin(x)^3/(a + b*cos(x)^2),x)`

output `cos(x)/b - (atan((b^(1/2)*cos(x))/a^(1/2))*(a + b))/(a^(1/2)*b^(3/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.31

$$\int \frac{\sin^3(x)}{a + b \cos^2(x)} dx \\ = \frac{\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)-\sqrt{b}}{\sqrt{a}}\right) a + \sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)-\sqrt{b}}{\sqrt{a}}\right) b - \sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)+\sqrt{b}}{\sqrt{a}}\right) a - \sqrt{b} a^2}{a b^2}$$

input `int(sin(x)^3/(a+b*cos(x)^2),x)`

output `(sqrt(b)*sqrt(a)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*a + sqrt(b)*sqrt(a)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*b - sqrt(b)*sqrt(a)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*a - sqrt(b)*sqrt(a)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*b + cos(x)*a*b - a*b)/(a*b**2)`

3.13 $\int \frac{\sin(x)}{a+b\cos^2(x)} dx$

Optimal result	120
Mathematica [A] (verified)	120
Rubi [A] (verified)	121
Maple [A] (verified)	122
Fricas [A] (verification not implemented)	123
Sympy [B] (verification not implemented)	123
Maxima [A] (verification not implemented)	124
Giac [A] (verification not implemented)	124
Mupad [B] (verification not implemented)	124
Reduce [B] (verification not implemented)	125

Optimal result

Integrand size = 13, antiderivative size = 26

$$\int \frac{\sin(x)}{a + b\cos^2(x)} dx = -\frac{\arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

output -arctan(b^(1/2)*cos(x)/a^(1/2))/a^(1/2)/b^(1/2)

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{\sin(x)}{a + b\cos^2(x)} dx = -\frac{\arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input Integrate[Sin[x]/(a + b*Cos[x]^2), x]

output -(ArcTan[(Sqrt[b]*Cos[x])/Sqrt[a]]/(Sqrt[a]*Sqrt[b]))

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 25, 3669, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\cos(x + \frac{\pi}{2})}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\cos(x + \frac{\pi}{2})}{b \sin(x + \frac{\pi}{2})^2 + a} dx \\
 & \quad \downarrow \text{3669} \\
 & - \int \frac{1}{b \cos^2(x) + a} d \cos(x) \\
 & \quad \downarrow \text{218} \\
 & - \frac{\arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{a} \sqrt{b}}
 \end{aligned}$$

input `Int[Sin[x]/(a + b*Cos[x]^2),x]`

output `-(ArcTan[(Sqrt[b]*Cos[x])/Sqrt[a]]/(Sqrt[a]*Sqrt[b]))`

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 218 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__)^2)^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[\text{a}/\text{b}, 2]/\text{a})*\text{ArcTan}[\text{x}/\text{Rt}[\text{a}/\text{b}, 2]], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{PosQ}[\text{a}/\text{b}]$

rule 3042 $\text{Int}[\text{u}__, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3669 $\text{Int}[\cos[(\text{e}__) + (\text{f}__)*(\text{x}__)]^{(\text{m}__)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Sin}[\text{e} + \text{f*x}], \text{x}]\}, \text{Simp}[\text{ff}/\text{f} \quad \text{ubst}[\text{Int}[(1 - \text{ff}^2*\text{x}^2)^{((\text{m} - 1)/2)}*(\text{a} + \text{b}*\text{ff}^2*\text{x}^2)^\text{p}, \text{x}], \text{x}, \text{Sin}[\text{e} + \text{f*x}]/\text{ff}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{p}\}, \text{x}] \& \text{IntegerQ}[(\text{m} - 1)/2]$

Maple [A] (verified)

Time = 0.10 (sec), antiderivative size = 18, normalized size of antiderivative = 0.69

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$	18
default	$-\frac{\arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$	18
risch	$-\frac{i \ln\left(e^{2ix} + \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)}{2\sqrt{ab}} + \frac{i \ln\left(e^{2ix} - \frac{2ia e^{ix}}{\sqrt{ab}} + 1\right)}{2\sqrt{ab}}$	62

input `int(sin(x)/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output $-1/(a*b)^{(1/2)}*\arctan(b*cos(x)/(a*b)^(1/2))$

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.81

$$\int \frac{\sin(x)}{a + b \cos^2(x)} dx = \left[-\frac{\sqrt{-ab} \log \left(-\frac{b \cos(x)^2 + 2\sqrt{-ab} \cos(x) - a}{b \cos(x)^2 + a} \right)}{2ab}, -\frac{\sqrt{ab} \arctan \left(\frac{\sqrt{ab} \cos(x)}{a} \right)}{ab} \right]$$

input `integrate(sin(x)/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[-1/2*sqrt(-a*b)*log(-(b*cos(x)^2 + 2*sqrt(-a*b)*cos(x) - a)/(b*cos(x)^2 + a))/(a*b), -sqrt(a*b)*arctan(sqrt(a*b)*cos(x)/a)/(a*b)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(26) = 52$.

Time = 0.42 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.54

$$\int \frac{\sin(x)}{a + b \cos^2(x)} dx = \begin{cases} \frac{\tilde{\cos}(x)}{\cos(x)} & \text{for } a = 0 \wedge b = 0 \\ -\frac{\cos(x)}{a} & \text{for } b = 0 \\ \frac{1}{b \cos(x)} & \text{for } a = 0 \\ -\frac{\log \left(-\sqrt{-\frac{a}{b}} + \cos(x) \right)}{2b\sqrt{-\frac{a}{b}}} + \frac{\log \left(\sqrt{-\frac{a}{b}} + \cos(x) \right)}{2b\sqrt{-\frac{a}{b}}} & \text{otherwise} \end{cases}$$

input `integrate(sin(x)/(a+b*cos(x)**2),x)`

output `Piecewise((zoo/cos(x), Eq(a, 0) & Eq(b, 0)), (-cos(x)/a, Eq(b, 0)), (1/(b*cos(x)), Eq(a, 0)), (-log(-sqrt(-a/b) + cos(x))/(2*b*sqrt(-a/b)) + log(sqrt(-a/b) + cos(x))/(2*b*sqrt(-a/b)), True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.65

$$\int \frac{\sin(x)}{a + b \cos^2(x)} dx = -\frac{\arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(sin(x)/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-arctan(b*cos(x)/sqrt(a*b))/sqrt(a*b)`

Giac [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.65

$$\int \frac{\sin(x)}{a + b \cos^2(x)} dx = -\frac{\arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(sin(x)/(a+b*cos(x)^2),x, algorithm="giac")`

output `-arctan(b*cos(x)/sqrt(a*b))/sqrt(a*b)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.69

$$\int \frac{\sin(x)}{a + b \cos^2(x)} dx = -\frac{\operatorname{atan}\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{a} \sqrt{b}}$$

input `int(sin(x)/(a + b*cos(x)^2),x)`

output `-atan((b^(1/2)*cos(x))/a^(1/2))/(a^(1/2)*b^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.96

$$\int \frac{\sin(x)}{a + b \cos^2(x)} dx = -\frac{\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{\cos(x)b}{\sqrt{b} \sqrt{a}}\right)}{ab}$$

input `int(sin(x)/(a+b*cos(x)^2),x)`

output `(- sqrt(b)*sqrt(a)*atan((cos(x)*b)/(sqrt(b)*sqrt(a))))/(a*b)`

3.14 $\int \frac{\csc(x)}{a+b\cos^2(x)} dx$

Optimal result	126
Mathematica [A] (verified)	126
Rubi [A] (verified)	127
Maple [A] (verified)	129
Fricas [A] (verification not implemented)	129
Sympy [F]	130
Maxima [A] (verification not implemented)	130
Giac [A] (verification not implemented)	130
Mupad [B] (verification not implemented)	131
Reduce [B] (verification not implemented)	131

Optimal result

Integrand size = 13, antiderivative size = 42

$$\int \frac{\csc(x)}{a + b\cos^2(x)} dx = -\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a + b)} - \frac{\operatorname{arctanh}(\cos(x))}{a + b}$$

output
$$-\text{b}^{(1/2)} \cdot \arctan(\text{b}^{(1/2)} \cdot \cos(x) / \text{a}^{(1/2)}) / \text{a}^{(1/2)} / (\text{a} + \text{b}) - \operatorname{arctanh}(\cos(x)) / (\text{a} + \text{b})$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int \frac{\csc(x)}{a + b\cos^2(x)} dx = \frac{-\frac{2\sqrt{b} \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{a}} + \log(1 - \cos(x)) - \log(1 + \cos(x))}{2(a + b)}$$

input `Integrate[Csc[x]/(a + b*Cos[x]^2), x]`

output
$$(-2 \cdot \text{Sqrt}[\text{b}] \cdot \operatorname{ArcTan}[(\text{Sqrt}[\text{b}] \cdot \cos(x)) / \text{Sqrt}[\text{a}]] / \text{Sqrt}[\text{a}] + \operatorname{Log}[1 - \cos(x)] - \operatorname{Log}[1 + \cos(x)]) / (2 \cdot (\text{a} + \text{b}))$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3042, 25, 3669, 303, 218, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\cos(x + \frac{\pi}{2}) \left(a + b \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\cos(x + \frac{\pi}{2}) \left(b \sin(x + \frac{\pi}{2})^2 + a\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3669} \\
 & - \int \frac{1}{(1 - \cos^2(x)) (b \cos^2(x) + a)} d \cos(x) \\
 & \quad \downarrow \textcolor{blue}{303} \\
 & - \frac{\int \frac{1}{1 - \cos^2(x)} d \cos(x)}{a + b} - \frac{b \int \frac{1}{b \cos^2(x) + a} d \cos(x)}{a + b} \\
 & \quad \downarrow \textcolor{blue}{218} \\
 & - \frac{\int \frac{1}{1 - \cos^2(x)} d \cos(x)}{a + b} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a + b)} \\
 & \quad \downarrow \textcolor{blue}{219} \\
 & - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a + b)} - \frac{\operatorname{arctanh}(\cos(x))}{a + b}
 \end{aligned}$$

input `Int[Csc[x]/(a + b*Cos[x]^2),x]`

output $-\frac{(\sqrt{b} \operatorname{ArcTan}\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right))}{\sqrt{a}(a+b)} - \frac{\operatorname{ArcTanh}\left(\frac{\cos(x)}{a+b}\right)}{a+b}$

Definitions of rubi rules used

rule 25 $\operatorname{Int}[-(F_x), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$

rule 218 $\operatorname{Int}[(a_+ + b_-)(x_-)^2, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(Rt[a/b, 2]/a) \operatorname{ArcTan}[x/Rt[a/b, 2]], x]; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{PosQ}[a/b]$

rule 219 $\operatorname{Int}[(a_+ + b_-)(x_-)^2, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(Rt[a, 2] * Rt[-b, 2])) \operatorname{ArcTanh}[Rt[-b, 2] * (x/Rt[a, 2])], x]; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{NegQ}[a/b] \& (\operatorname{GtQ}[a, 0] \text{ || } \operatorname{LtQ}[b, 0])$

rule 303 $\operatorname{Int}[1/((a_+ + b_-)(x_-)^2)(c_+ + d_-)(x_-)^2, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[b/(b*c - a*d) \operatorname{Int}[1/(a + b*x^2), x], x] - \operatorname{Simp}[d/(b*c - a*d) \operatorname{Int}[1/(c + d*x^2), x], x]; \operatorname{FreeQ}[\{a, b, c, d\}, x] \& \operatorname{NeQ}[b*c - a*d, 0]$

rule 3042 $\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x]; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3669 $\operatorname{Int}[\cos(e_- + f_-)(x_-)^m(a_+ + b_-)\sin(e_- + f_-)(x_-)^2, x_{\text{Symbol}}] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Sin}[e + f*x], x]\}, \operatorname{Simp}[ff/f \operatorname{Subst}[\operatorname{Int}[(1 - ff^2*x^2)^{(m-1)/2} * (a + b*ff^2*x^2)^p, x], x, \operatorname{Sin}[e + f*x]/ff], x]]; \operatorname{FreeQ}[\{a, b, e, f, p\}, x] \& \operatorname{IntegerQ}[(m-1)/2]$

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.33

method	result	size
default	$-\frac{b \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a+b)\sqrt{ab}} - \frac{\ln(1+\cos(x))}{2a+2b} + \frac{\ln(-1+\cos(x))}{2a+2b}$	56
risch	$-\frac{\ln(e^{ix}+1)}{a+b} + \frac{\ln(e^{ix}-1)}{a+b} + \frac{i\sqrt{ab} \ln\left(e^{2ix} - \frac{2i\sqrt{ab}e^{ix}}{b} + 1\right)}{2a(a+b)} - \frac{i\sqrt{ab} \ln\left(e^{2ix} + \frac{2i\sqrt{ab}e^{ix}}{b} + 1\right)}{2a(a+b)}$	111

input `int(csc(x)/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -b/(a+b)/(a*b)^(1/2)*\arctan(b*cos(x)/(a*b)^(1/2))-1/(2*a+2*b)*\ln(1+\cos(x)) \\ & +1/(2*a+2*b)*\ln(-1+\cos(x)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.69

$$\begin{aligned} & \int \frac{\csc(x)}{a + b \cos^2(x)} dx \\ &= \left[\frac{\sqrt{-\frac{b}{a}} \log\left(\frac{b \cos(x)^2 - 2a\sqrt{-\frac{b}{a}} \cos(x) - a}{b \cos(x)^2 + a}\right) - \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{2(a+b)}, \right. \\ & \quad \left. - \frac{2\sqrt{\frac{b}{a}} \arctan\left(\sqrt{\frac{b}{a}} \cos(x)\right) + \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{2(a+b)} \right] \end{aligned}$$

input `integrate(csc(x)/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$\begin{aligned} & [1/2*(\sqrt{-b/a})*\log((b*cos(x)^2 - 2*a*\sqrt{-b/a}*\cos(x) - a)/(b*cos(x)^2 + a)) - \log(1/2*\cos(x) + 1/2) + \log(-1/2*\cos(x) + 1/2))/(a + b), -1/2*(2*\sqrt{b/a}*\arctan(\sqrt{b/a}*\cos(x)) + \log(1/2*\cos(x) + 1/2) - \log(-1/2*\cos(x) + 1/2))/(a + b)] \end{aligned}$$

Sympy [F]

$$\int \frac{\csc(x)}{a + b \cos^2(x)} dx = \int \frac{\csc(x)}{a + b \cos^2(x)} dx$$

input `integrate(csc(x)/(a+b*cos(x)**2),x)`

output `Integral(csc(x)/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.14

$$\int \frac{\csc(x)}{a + b \cos^2(x)} dx = -\frac{b \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}(a + b)} - \frac{\log(\cos(x) + 1)}{2(a + b)} + \frac{\log(\cos(x) - 1)}{2(a + b)}$$

input `integrate(csc(x)/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-b*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*(a + b)) - 1/2*log(cos(x) + 1)/(a + b) + 1/2*log(cos(x) - 1)/(a + b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int \frac{\csc(x)}{a + b \cos^2(x)} dx = -\frac{b \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{\sqrt{ab}(a + b)} - \frac{\log(\cos(x) + 1)}{2(a + b)} + \frac{\log(-\cos(x) + 1)}{2(a + b)}$$

input `integrate(csc(x)/(a+b*cos(x)^2),x, algorithm="giac")`

output `-b*arctan(b*cos(x)/sqrt(a*b))/(sqrt(a*b)*(a + b)) - 1/2*log(cos(x) + 1)/(a + b) + 1/2*log(-cos(x) + 1)/(a + b)`

Mupad [B] (verification not implemented)

Time = 1.17 (sec) , antiderivative size = 853, normalized size of antiderivative = 20.31

$$\int \frac{\csc(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input `int(1/(sin(x)*(a + b*cos(x)^2)),x)`

output

```
(atan((((8*a*b^3 + 4*b^4 + 4*a^2*b^2 - (cos(x)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(2*(a + b)))/(2*(a + b)) + 4*b^3*cos(x))*1i)/(2*(a + b)) - (((8*a*b^3 + 4*b^4 + 4*a^2*b^2 + (cos(x)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(2*(a + b)) - 4*b^3*cos(x))*1i)/(2*(a + b)))/(((8*a*b^3 + 4*b^4 + 4*a^2*b^2 - (cos(x)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(2*(a + b)) + 4*b^3*cos(x))/(2*(a + b)) + ((8*a*b^3 + 4*b^4 + 4*a^2*b^2 + (cos(x)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(2*(a + b)) - 4*b^3*cos(x))/(2*(a + b))))*1i)/(a + b) + (atan((( -a*b)^(1/2)*(2*b^3*cos(x) + ((-a*b)^(1/2)*(4*a*b^3 + 2*b^4 + 2*a^2*b^2 - (cos(x)*(-a*b)^(1/2)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(4*(a*b + a^2)))/(2*(a*b + a^2)))*1i)/(a*b + a^2) + ((-a*b)^(1/2)*(2*b^3*cos(x) - (( -a*b)^(1/2)*(4*a*b^3 + 2*b^4 + 2*a^2*b^2 + (cos(x)*(-a*b)^(1/2)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(4*(a*b + a^2)))/(2*(a*b + a^2)))*1i)/(a*b + a^2))/((( -a*b)^(1/2)*(2*b^3*cos(x) + ((-a*b)^(1/2)*(4*a*b^3 + 2*b^4 + 2*a^2*b^2 - (cos(x)*(-a*b)^(1/2)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(4*(a*b + a^2)))/(2*(a*b + a^2)))*1i)/(a*b + a^2))/((( -a*b)^(1/2)*(2*b^3*cos(x) - (( -a*b)^(1/2)*(4*a*b^3 + 2*b^4 + 2*a^2*b^2 + (cos(x)*(-a*b)^(1/2)*(8*a*b^4 + 8*b^5 - 8*a^2*b^3 - 8*a^3*b^2))/(4*(a*b + a^2)))/(2*(a*b + a^2)))*1i)/(a*b + a^2)))*(-a*b)^(1/2)*1i)/(a*(a + b))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.57

$$\begin{aligned} & \int \frac{\csc(x)}{a + b \cos^2(x)} dx \\ &= \frac{\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)-\sqrt{b}}{\sqrt{a}}\right)-\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)+\sqrt{b}}{\sqrt{a}}\right)+\log \left(\tan \left(\frac{x}{2}\right)\right) a}{a(a+b)} \end{aligned}$$

input `int(csc(x)/(a+b*cos(x)^2),x)`

output
$$\frac{(\sqrt{b}*\sqrt{a}*\operatorname{atan}\left(\frac{\sqrt{a+b}*\tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right) - \sqrt{b}*\sqrt{a}*\operatorname{atan}\left(\frac{\sqrt{a+b}*\tan\left(\frac{x}{2}\right) + \sqrt{b}}{\sqrt{a}}\right) + \log\left(\tan\left(\frac{x}{2}\right)\right)*a)}{a*(a+b)}$$

3.15 $\int \frac{\csc^3(x)}{a+b\cos^2(x)} dx$

Optimal result	133
Mathematica [B] (verified)	133
Rubi [A] (verified)	134
Maple [A] (verified)	136
Fricas [B] (verification not implemented)	137
Sympy [F]	137
Maxima [B] (verification not implemented)	138
Giac [B] (verification not implemented)	138
Mupad [B] (verification not implemented)	139
Reduce [B] (verification not implemented)	139

Optimal result

Integrand size = 15, antiderivative size = 62

$$\int \frac{\csc^3(x)}{a+b\cos^2(x)} dx = -\frac{b^{3/2} \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^2} - \frac{(a+3b)\operatorname{arctanh}(\cos(x))}{2(a+b)^2} - \frac{\cot(x) \csc(x)}{2(a+b)}$$

output
$$-\text{b}^{(3/2)} \arctan(\text{b}^{(1/2)} \cos(x)/\text{a}^{(1/2)})/\text{a}^{(1/2)} / (\text{a}+\text{b})^{2-1/2} * (\text{a}+3*\text{b}) * \arctan(\text{h}(\cos(x))/(\text{a}+\text{b})^{2-\cot(x)*\csc(x)/(2*a+2*b)})$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 140 vs. $2(62) = 124$.

Time = 0.66 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.26

$$\begin{aligned} & \int \frac{\csc^3(x)}{a+b\cos^2(x)} dx \\ &= \frac{-8b^{3/2} \arctan\left(\frac{\sqrt{b}-\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right) - 8b^{3/2} \arctan\left(\frac{\sqrt{b}+\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right) + \sqrt{a}(-((a+b)\csc^2(\frac{x}{2})) - 4(a+3b)(\csc(x)\cot(x)))}{8\sqrt{a}(a+b)^2} \end{aligned}$$

input
$$\text{Integrate}[\text{Csc}[x]^3/(a + b*\text{Cos}[x]^2), x]$$

output

$$(-8*b^{(3/2)}*\text{ArcTan}[(\text{Sqrt}[b] - \text{Sqrt}[a + b]*\text{Tan}[x/2])/\text{Sqrt}[a]] - 8*b^{(3/2)}*\text{ArcTan}[(\text{Sqrt}[b] + \text{Sqrt}[a + b]*\text{Tan}[x/2])/\text{Sqrt}[a]] + \text{Sqrt}[a]*(-((a + b)*\text{Csc}[x/2]^2) - 4*(a + 3*b)*(\text{Log}[\text{Cos}[x/2]] - \text{Log}[\text{Sin}[x/2]]) + (a + b)*\text{Sec}[x/2]^2))/(8*\text{Sqrt}[a]*(a + b)^2)$$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 77, normalized size of antiderivative = 1.24, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 25, 3669, 316, 397, 218, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\csc^3(x)}{a + b \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{1}{\cos(x + \frac{\pi}{2})^3 (a + b \sin(x + \frac{\pi}{2})^2)} dx \\ & \quad \downarrow \text{25} \\ & - \int \frac{1}{\cos(x + \frac{\pi}{2})^3 (b \sin(x + \frac{\pi}{2})^2 + a)} dx \\ & \quad \downarrow \text{3669} \\ & - \int \frac{1}{(1 - \cos^2(x))^2 (b \cos^2(x) + a)} d\cos(x) \\ & \quad \downarrow \text{316} \\ & - \frac{\int \frac{b \cos^2(x) + a + 2b}{(1 - \cos^2(x))(b \cos^2(x) + a)} d\cos(x)}{2(a + b)} - \frac{\cos(x)}{2(a + b)(1 - \cos^2(x))} \\ & \quad \downarrow \text{397} \\ & - \frac{\frac{2b^2 \int \frac{1}{b \cos^2(x) + a} d\cos(x)}{a + b} + \frac{(a + 3b) \int \frac{1}{1 - \cos^2(x)} d\cos(x)}{a + b}}{2(a + b)} - \frac{\cos(x)}{2(a + b)(1 - \cos^2(x))} \\ & \quad \downarrow \text{218} \end{aligned}$$

$$\begin{aligned}
 & -\frac{\frac{(a+3b) \int \frac{1}{1-\cos^2(x)} d\cos(x)}{a+b} + \frac{2b^{3/2} \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)}}{2(a+b)} - \frac{\cos(x)}{2(a+b)(1-\cos^2(x))} \\
 & \quad \downarrow 219 \\
 & -\frac{\frac{2b^{3/2} \arctan\left(\frac{\sqrt{b} \cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)} + \frac{(a+3b)\operatorname{arctanh}(\cos(x))}{a+b}}{2(a+b)} - \frac{\cos(x)}{2(a+b)(1-\cos^2(x))}
 \end{aligned}$$

input `Int[Csc[x]^3/(a + b*Cos[x]^2), x]`

output $-1/2*((2*b^{(3/2)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*\operatorname{Cos}[x])/\operatorname{Sqrt}[a]])/(\operatorname{Sqrt}[a]*(a+b)) + ((a+3*b)*\operatorname{ArcTanh}[\operatorname{Cos}[x]])/(a+b))/(a+b) - \operatorname{Cos}[x]/(2*(a+b)*(1-\operatorname{Cos}[x]^2))$

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 316 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] :> Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))), x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2, p, q, x]`

rule 397

```
Int[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*((c_) + (d_)*(x_)^2)), x_
Symbol] :> Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[
(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e
, f}, x]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3669

```
Int[cos[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^
(p_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Simp[ff/f S
ubst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]
/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.53

method	result
default	$-\frac{b^2 \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a+b)^2 \sqrt{ab}} + \frac{1}{(4a+4b)(1+\cos(x))} + \frac{(-a-3b) \ln(1+\cos(x))}{4(a+b)^2} + \frac{1}{(4a+4b)(-1+\cos(x))} + \frac{(a+3b) \ln(-1+\cos(x))}{4(a+b)^2}$
risch	$\frac{e^{3ix}+e^{ix}}{(e^{2ix}-1)^2(a+b)} + \frac{\ln(e^{ix}-1)a}{2a^2+4ab+2b^2} + \frac{3 \ln(e^{ix}-1)b}{2(a^2+2ab+b^2)} - \frac{\ln(e^{ix}+1)a}{2(a^2+2ab+b^2)} - \frac{3 \ln(e^{ix}+1)b}{2(a^2+2ab+b^2)} - \frac{i\sqrt{ab}b \ln\left(e^{2ix} + \frac{2i\sqrt{ab}e^{ix}}{b} + 1\right)}{2a(a+b)^2}$

input

```
int(csc(x)^3/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)
```

output

```
-1/(a+b)^2*b^2/(a*b)^(1/2)*arctan(b*cos(x)/(a*b)^(1/2))+1/(4*a+4*b)/(1+cos
(x))+1/4/(a+b)^2*(-a-3*b)*ln(1+cos(x))+1/(4*a+4*b)/(-1+cos(x))+1/4*(a+3*b)
/(a+b)^2*ln(-1+cos(x))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. $2(50) = 100$.

Time = 0.21 (sec), antiderivative size = 274, normalized size of antiderivative = 4.42

$$\int \frac{\csc^3(x)}{a + b \cos^2(x)} dx$$

$$= \frac{2(b \cos(x)^2 - b) \sqrt{-\frac{b}{a}} \log\left(\frac{b \cos(x)^2 - 2a \sqrt{-\frac{b}{a}} \cos(x) - a}{b \cos(x)^2 + a}\right) + 2(a + b) \cos(x) - ((a + 3b) \cos(x)^2 - a - 3b)}{4((a^2 + 2ab + b^2) \cos(x)^2 - a^2 - 2ab - b^2)}$$

$$- \frac{4(b \cos(x)^2 - b) \sqrt{\frac{b}{a}} \arctan\left(\sqrt{\frac{b}{a}} \cos(x)\right) - 2(a + b) \cos(x) + ((a + 3b) \cos(x)^2 - a - 3b) \log\left(\frac{1}{2} \cos(x)\right)}{4((a^2 + 2ab + b^2) \cos(x)^2 - a^2 - 2ab - b^2)}$$

input `integrate(csc(x)^3/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[1/4*(2*(b*cos(x)^2 - b)*sqrt(-b/a)*log((b*cos(x)^2 - 2*a*sqrt(-b/a)*cos(x) - a)/(b*cos(x)^2 + a)) + 2*(a + b)*cos(x) - ((a + 3*b)*cos(x)^2 - a - 3*b)*log(1/2*cos(x) + 1/2) + ((a + 3*b)*cos(x)^2 - a - 3*b)*log(-1/2*cos(x) + 1/2))/((a^2 + 2*a*b + b^2)*cos(x)^2 - a^2 - 2*a*b - b^2), -1/4*(4*(b*cos(x)^2 - b)*sqrt(b/a)*arctan(sqrt(b/a)*cos(x)) - 2*(a + b)*cos(x) + ((a + 3*b)*cos(x)^2 - a - 3*b)*log(1/2*cos(x) + 1/2) - ((a + 3*b)*cos(x)^2 - a - 3*b)*log(-1/2*cos(x) + 1/2))/((a^2 + 2*a*b + b^2)*cos(x)^2 - a^2 - 2*a*b - b^2)]`

Sympy [F]

$$\int \frac{\csc^3(x)}{a + b \cos^2(x)} dx = \int \frac{\csc^3(x)}{a + b \cos^2(x)} dx$$

input `integrate(csc(x)**3/(a+b*cos(x)**2),x)`

output `Integral(csc(x)**3/(a + b*cos(x)**2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(50) = 100$.

Time = 0.11 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.69

$$\int \frac{\csc^3(x)}{a + b \cos^2(x)} dx = -\frac{b^2 \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a^2 + 2ab + b^2)\sqrt{ab}} - \frac{(a + 3b) \log(\cos(x) + 1)}{4(a^2 + 2ab + b^2)} \\ + \frac{(a + 3b) \log(\cos(x) - 1)}{4(a^2 + 2ab + b^2)} + \frac{\cos(x)}{2((a + b)\cos(x)^2 - a - b)}$$

input `integrate(csc(x)^3/(a+b*cos(x)^2),x, algorithm="maxima")`

output
$$-\frac{b^2 \arctan(b \cos(x)/\sqrt{a b})}{(a^2 + 2 a b + b^2) \sqrt{a b}} - \frac{1}{4} \frac{(a + 3 b) \log(\cos(x) + 1)}{(a^2 + 2 a b + b^2)} + \frac{1}{4} \frac{(a + 3 b) \log(\cos(x) - 1)}{(a^2 + 2 a b + b^2)} + \frac{1}{2} \frac{\cos(x)}{(\cos(x)^2 - 1)(a + b)}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(50) = 100$.

Time = 0.12 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.66

$$\int \frac{\csc^3(x)}{a + b \cos^2(x)} dx = -\frac{b^2 \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a^2 + 2ab + b^2)\sqrt{ab}} - \frac{(a + 3b) \log(\cos(x) + 1)}{4(a^2 + 2ab + b^2)} \\ + \frac{(a + 3b) \log(-\cos(x) + 1)}{4(a^2 + 2ab + b^2)} + \frac{\cos(x)}{2(\cos(x)^2 - 1)(a + b)}$$

input `integrate(csc(x)^3/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$-\frac{b^2 \arctan(b \cos(x)/\sqrt{a b})}{(a^2 + 2 a b + b^2) \sqrt{a b}} - \frac{1}{4} \frac{(a + 3 b) \log(\cos(x) + 1)}{(a^2 + 2 a b + b^2)} + \frac{1}{4} \frac{(a + 3 b) \log(-\cos(x) + 1)}{(a^2 + 2 a b + b^2)} + \frac{1}{2} \frac{\cos(x)}{(\cos(x)^2 - 1)(a + b)}$$

Mupad [B] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 1138, normalized size of antiderivative = 18.35

$$\int \frac{\csc^3(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input int(1/(sin(x)^3*(a + b*cos(x)^2)),x)

```

output log(cos(x) - 1)*(b/(2*(a + b)^2) + 1/(4*(a + b))) - cos(x)/(2*sin(x)^2*(a
+ b)) - (log(cos(x) + 1)*(a + 3*b))/(4*(a + b)^2) - (atan((((-a*b^3)^(1/2)
*((cos(x)*(6*a*b^4 + 13*b^5 + a^2*b^3)))/(4*(2*a*b + a^2 + b^2)) + (((18*a*
b^6 + 4*b^7 + 32*a^2*b^5 + 28*a^3*b^4 + 12*a^4*b^3 + 2*a^5*b^2)/(2*(3*a*b^
2 + 3*a^2*b + a^3 + b^3)) - (cos(x)*(-a*b^3)^(1/2)*(48*a*b^6 + 16*b^7 + 32
*a^2*b^5 - 32*a^3*b^4 - 48*a^4*b^3 - 16*a^5*b^2))/(8*(2*a*b + a^2 + b^2)*(a
*b^2 + 2*a^2*b + a^3)))*(-a*b^3)^(1/2))/(2*(a*b^2 + 2*a^2*b + a^3)))*1i)/
(a*b^2 + 2*a^2*b + a^3) + ((-a*b^3)^(1/2)*((cos(x)*(6*a*b^4 + 13*b^5 + a^2
*b^3))/(4*(2*a*b + a^2 + b^2)) - (((18*a*b^6 + 4*b^7 + 32*a^2*b^5 + 28*a^3
*b^4 + 12*a^4*b^3 + 2*a^5*b^2)/(2*(3*a*b^2 + 3*a^2*b + a^3 + b^3)) + (cos(
x)*(-a*b^3)^(1/2)*(48*a*b^6 + 16*b^7 + 32*a^2*b^5 - 32*a^3*b^4 - 48*a^4*b^
3 - 16*a^5*b^2))/(8*(2*a*b + a^2 + b^2)*(a*b^2 + 2*a^2*b + a^3)))*(-a*b^3)
^(1/2))/(2*(a*b^2 + 2*a^2*b + a^3)))*1i)/(a*b^2 + 2*a^2*b + a^3))/(((a*b^4
)/2 + (3*b^5)/2)/(3*a*b^2 + 3*a^2*b + a^3 + b^3) - ((-a*b^3)^(1/2)*((cos(x)
)*(6*a*b^4 + 13*b^5 + a^2*b^3))/(4*(2*a*b + a^2 + b^2)) + (((18*a*b^6 + 4*
b^7 + 32*a^2*b^5 + 28*a^3*b^4 + 12*a^4*b^3 + 2*a^5*b^2)/(2*(3*a*b^2 + 3*a^
2*b + a^3 + b^3)) - (cos(x)*(-a*b^3)^(1/2)*(48*a*b^6 + 16*b^7 + 32*a^2*b^5
- 32*a^3*b^4 - 48*a^4*b^3 - 16*a^5*b^2))/(8*(2*a*b + a^2 + b^2)*(a*b^2 +
2*a^2*b + a^3)))*(-a*b^3)^(1/2))/(2*(a*b^2 + 2*a^2*b + a^3))))/(a*b^2 + 2*
a^2*b + a^3) + ((-a*b^3)^(1/2)*((cos(x)*(6*a*b^4 + 13*b^5 + a^2*b^3))/(...

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.97

$$\int \frac{\csc^3(x)}{a + b \cos^2(x)} dx \\ = \frac{2\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right) \sin(x)^2 b - 2\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) + \sqrt{b}}{\sqrt{a}}\right) \sin(x)^2 b - \cos(x) a^2 - \cos(x)}{2 \sin(x)^2 a (a^2 + 2ab + b^2)}$$

input `int(csc(x)^3/(a+b*cos(x)^2),x)`

output
$$\frac{(2\sqrt{b}\sqrt{a}\operatorname{atan}(\sqrt{a+b}\tan(x/2) - \sqrt{b})/\sqrt{a})\sin(x)^{2*2*b} - 2\sqrt{b}\sqrt{a}\operatorname{atan}(\sqrt{a+b}\tan(x/2) + \sqrt{b})/\sqrt{a})\sin(x)^{2*2*b} - \cos(x)a^{**2} - \cos(x)a*b + \log(\tan(x/2))\sin(x)^{2*a^{**2}} + 3\log(\tan(x/2))\sin(x)^{2*a*b})}{(2\sin(x)^{2*a*(a^{**2} + 2*a*b + b^{**2}))}}$$

3.16 $\int \frac{\csc^5(x)}{a+b\cos^2(x)} dx$

Optimal result	141
Mathematica [B] (verified)	141
Rubi [A] (verified)	142
Maple [A] (verified)	145
Fricas [B] (verification not implemented)	145
Sympy [F]	146
Maxima [B] (verification not implemented)	147
Giac [B] (verification not implemented)	147
Mupad [B] (verification not implemented)	148
Reduce [B] (verification not implemented)	149

Optimal result

Integrand size = 15, antiderivative size = 94

$$\int \frac{\csc^5(x)}{a+b\cos^2(x)} dx = -\frac{b^{5/2} \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^3} - \frac{(3a^2 + 10ab + 15b^2) \operatorname{arctanh}(\cos(x))}{8(a+b)^3} - \frac{(3a+7b) \cot(x) \csc(x)}{8(a+b)^2} - \frac{\cot(x) \csc^3(x)}{4(a+b)}$$

output

```
-b^(5/2)*arctan(b^(1/2)*cos(x)/a^(1/2))/a^(1/2)/(a+b)^3-1/8*(3*a^2+10*a*b+15*b^2)*arctanh(cos(x))/(a+b)^3-1/8*(3*a+7*b)*cot(x)*csc(x)/(a+b)^2-cot(x)*csc(x)^3/(4*a+4*b)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 204 vs. $2(94) = 188$.

Time = 1.56 (sec), antiderivative size = 204, normalized size of antiderivative = 2.17

$$\int \frac{\csc^5(x)}{a+b\cos^2(x)} dx = \frac{-64b^{5/2} \arctan\left(\frac{\sqrt{b}-\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right) - 64b^{5/2} \arctan\left(\frac{\sqrt{b}+\sqrt{a+b}\tan(\frac{x}{2})}{\sqrt{a}}\right) + \sqrt{a}(-2(3a^2 + 10ab + 7b^2) \csc^2(\frac{x}{2}))}{a+b\cos^2(x)}$$

input $\text{Integrate}[\text{Csc}[x]^5/(a + b \cos[x]^2), x]$

output
$$\begin{aligned} & (-64*b^{(5/2)}*\text{ArcTan}[(\text{Sqrt}[b] - \text{Sqrt}[a + b]*\tan[x/2])/\text{Sqrt}[a]] - 64*b^{(5/2)} \\ & *\text{ArcTan}[(\text{Sqrt}[b] + \text{Sqrt}[a + b]*\tan[x/2])/\text{Sqrt}[a]] + \text{Sqrt}[a]*(-2*(3*a^2 + 1 \\ & 0*a*b + 7*b^2)*\text{Csc}[x/2]^2 - (a + b)^2*\text{Csc}[x/2]^4 - 8*(3*a^2 + 10*a*b + 15* \\ & b^2)*(\log[\cos[x/2]] - \log[\sin[x/2]]) + 2*(3*a^2 + 10*a*b + 7*b^2)*\text{Sec}[x/2] \\ & ^2 + (a + b)^2*\text{Sec}[x/2]^4))/(64*\text{Sqrt}[a]*(a + b)^3) \end{aligned}$$

Rubi [A] (verified)

Time = 0.34 (sec), antiderivative size = 125, normalized size of antiderivative = 1.33, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.533, Rules used = {3042, 25, 3669, 316, 402, 397, 218, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\csc^5(x)}{a + b \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int -\frac{1}{\cos(x + \frac{\pi}{2})^5 (a + b \sin(x + \frac{\pi}{2})^2)} dx \\ & \quad \downarrow \text{25} \\ & - \int \frac{1}{\cos(x + \frac{\pi}{2})^5 (b \sin(x + \frac{\pi}{2})^2 + a)} dx \\ & \quad \downarrow \text{3669} \\ & - \int \frac{1}{(1 - \cos^2(x))^3 (b \cos^2(x) + a)} d \cos(x) \\ & \quad \downarrow \text{316} \\ & - \frac{\int \frac{3b \cos^2(x) + 3a + 4b}{(1 - \cos^2(x))^2 (b \cos^2(x) + a)} d \cos(x)}{4(a + b)} - \frac{\cos(x)}{4(a + b)(1 - \cos^2(x))^2} \\ & \quad \downarrow \text{402} \end{aligned}$$

$$\begin{aligned}
 & -\frac{\int \frac{3a^2+7ba+8b^2+b(3a+7b)\cos^2(x)}{(1-\cos^2(x))(b\cos^2(x)+a)} d\cos(x)}{2(a+b)} + \frac{(3a+7b)\cos(x)}{2(a+b)(1-\cos^2(x))} - \frac{\cos(x)}{4(a+b)(1-\cos^2(x))^2} \\
 & \quad \downarrow \text{397} \\
 & -\frac{\frac{(3a^2+10ab+15b^2)}{a+b} \int \frac{1}{1-\cos^2(x)} d\cos(x) + \frac{8b^3}{2(a+b)} \int \frac{1}{b\cos^2(x)+a} d\cos(x)}{4(a+b)} + \frac{(3a+7b)\cos(x)}{2(a+b)(1-\cos^2(x))} - \frac{\cos(x)}{4(a+b)(1-\cos^2(x))^2} \\
 & \quad \downarrow \text{218} \\
 & -\frac{\frac{(3a^2+10ab+15b^2)}{a+b} \int \frac{1}{1-\cos^2(x)} d\cos(x) + \frac{8b^{5/2}}{\sqrt{a(a+b)}} \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{2(a+b)} + \frac{(3a+7b)\cos(x)}{2(a+b)(1-\cos^2(x))} - \frac{\cos(x)}{4(a+b)(1-\cos^2(x))^2} \\
 & \quad \downarrow \text{219} \\
 & -\frac{\frac{(3a^2+10ab+15b^2)}{a+b} \operatorname{arctanh}_{(\cos(x))} + \frac{8b^{5/2}}{\sqrt{a(a+b)}} \arctan\left(\frac{\sqrt{b}\cos(x)}{\sqrt{a}}\right)}{2(a+b)} + \frac{(3a+7b)\cos(x)}{2(a+b)(1-\cos^2(x))} - \frac{\cos(x)}{4(a+b)(1-\cos^2(x))^2}
 \end{aligned}$$

input `Int[Csc[x]^5/(a + b*Cos[x]^2), x]`

output `-1/4*Cos[x]/((a + b)*(1 - Cos[x]^2)^2) - (((8*b^(5/2)*ArcTan[(Sqrt[b]*Cos[x])/Sqrt[a]])/(Sqrt[a]*(a + b)) + ((3*a^2 + 10*a*b + 15*b^2)*ArcTanh[Cos[x]])/(a + b))/(2*(a + b)) + ((3*a + 7*b)*Cos[x])/(2*(a + b)*(1 - Cos[x]^2)))/(4*(a + b))`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x]; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 219 $\text{Int}[(a_ + b_)*(x_)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(1/(Rt[a, 2]*Rt[-b, 2])) * \text{ArcTanh}[Rt[-b, 2]*(x/Rt[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{NegQ}[a/b] \&& (\text{GtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$

rule 316 $\text{Int}[(a_ + b_)*(x_)^2)^{(p_)*((c_ + d_)*(x_)^2)^{(q_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-b)*x*(a + b*x^2)^{(p + 1)*((c + d*x^2)^{(q + 1)/(2*a*(p + 1)*(b*c - a*d))}, x] + \text{Simp}[1/(2*a*(p + 1)*(b*c - a*d)) \text{Int}[(a + b*x^2)^{(p + 1)*(c + d*x^2)^q} * \text{Simp}[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x] /; \text{FreeQ}[\{a, b, c, d, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{LtQ}[p, -1] \&& !(\text{IntegerQ}[p] \&& \text{IntegerQ}[q] \&& \text{LtQ}[q, -1]) \&& \text{IntBinomialQ}[a, b, c, d, 2, p, q, x]$

rule 397 $\text{Int}[(e_ + f_)*(x_)^2)/(((a_ + b_)*(x_)^2)*(c_ + d_)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{Simp}[(b*e - a*f)/(b*c - a*d) \text{Int}[1/(a + b*x^2), x], x] - \text{Simp}[(d*e - c*f)/(b*c - a*d) \text{Int}[1/(c + d*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x]$

rule 402 $\text{Int}[(a_ + b_)*(x_)^2)^{(p_)*((c_ + d_)*(x_)^2)^{(q_)*((e_ + f_)*(x_)^2)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-(b*e - a*f)*x*(a + b*x^2)^{(p + 1)*((c + d*x^2)^{(q + 1)/(a*2*(b*c - a*d)*(p + 1))}}, x] + \text{Simp}[1/(a*2*(b*c - a*d)*(p + 1)) \text{Int}[(a + b*x^2)^{(p + 1)*(c + d*x^2)^q} * \text{Simp}[c*(b*e - a*f) + e*2*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, q\}, x] \&& \text{LtQ}[p, -1]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3669 $\text{Int}[\cos[(e_ + f_)*(x_)]^{(m_)*((a_ + b_)*\sin[(e_ + f_)*(x_)]^2)^{(p_)}, x_{\text{Symbol}}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Simp}[ff/f \text{Sbst}[\text{Int}[(1 - ff^2*x^2)^{((m - 1)/2)*(a + b*ff^2*x^2)^p}, x], x, \text{Sin}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{IntegerQ}[(m - 1)/2]$

Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.65

method	result
default	$-\frac{b^3 \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a+b)^3 \sqrt{ab}} + \frac{1}{2(8a+8b)(1+\cos(x))^2} - \frac{-3a-7b}{16(a+b)^2(1+\cos(x))} + \frac{(-3a^2-10ab-15b^2) \ln(1+\cos(x))}{16(a+b)^3} - \frac{1}{2(8a+8b)(-1+\cos(x))}$
risch	$\frac{3a e^{7ix} + 7b e^{7ix} - 11a e^{5ix} - 15b e^{5ix} - 11a e^{3ix} - 15b e^{3ix} + 3e^{ix}a + 7e^{ix}b}{4(a+b)^2(e^{2ix}-1)^4} + \frac{3 \ln(e^{ix}-1)a^2}{8(a^3+3a^2b+3ab^2+b^3)} + \frac{5 \ln(e^{ix}-1)ab}{4(a^3+3a^2b+3ab^2+b^3)} + \frac{5 \ln(e^{ix}-1)b^2}{8(a^3+3a^2b+3ab^2+b^3)}$

input `int(csc(x)^5/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/(a+b)^3*b^3/(a*b)^(1/2)*\arctan(b*\cos(x)/(a*b)^(1/2))+1/2/(8*a+8*b)/(1+\cos(x))^2-1/16*(-3*a-7*b)/(a+b)^2/(1+\cos(x))+1/16/(a+b)^3*(-3*a^2-10*a*b-15*b^2)*\ln(1+\cos(x))-1/2/(8*a+8*b)/(-1+\cos(x))^2-1/16*(-3*a-7*b)/(a+b)^2/(-1+\cos(x))+1/16*(3*a^2+10*a*b+15*b^2)/(a+b)^3*\ln(-1+\cos(x)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 283 vs. $2(80) = 160$.

Time = 0.26 (sec) , antiderivative size = 592, normalized size of antiderivative = 6.30

$$\int \frac{\csc^5(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input `integrate(csc(x)^5/(a+b*cos(x)^2),x, algorithm="fricas")`

output

```
[1/16*(2*(3*a^2 + 10*a*b + 7*b^2)*cos(x)^3 + 8*(b^2*cos(x)^4 - 2*b^2*cos(x)^2 + b^2)*sqrt(-b/a)*log((b*cos(x)^2 - 2*a*sqrt(-b/a)*cos(x) - a)/(b*cos(x)^2 + a)) - 2*(5*a^2 + 14*a*b + 9*b^2)*cos(x) - ((3*a^2 + 10*a*b + 15*b^2)*cos(x)^4 - 2*(3*a^2 + 10*a*b + 15*b^2)*cos(x)^2 + 3*a^2 + 10*a*b + 15*b^2)*log(1/2*cos(x) + 1/2) + ((3*a^2 + 10*a*b + 15*b^2)*cos(x)^4 - 2*(3*a^2 + 10*a*b + 15*b^2)*cos(x)^2 + 3*a^2 + 10*a*b + 15*b^2)*log(-1/2*cos(x) + 1/2))/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cos(x)^4 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cos(x)^2), 1/16*(2*(3*a^2 + 10*a*b + 7*b^2)*cos(x)^3 - 16*(b^2*cos(x)^4 - 2*b^2*cos(x)^2 + b^2)*sqrt(b/a)*arctan(sqrt(b/a)*cos(x)) - 2*(5*a^2 + 14*a*b + 9*b^2)*cos(x) - ((3*a^2 + 10*a*b + 15*b^2)*cos(x)^4 - 2*(3*a^2 + 10*a*b + 15*b^2)*cos(x)^2 + 3*a^2 + 10*a*b + 15*b^2)*log(1/2*cos(x) + 1/2) + ((3*a^2 + 10*a*b + 15*b^2)*cos(x)^4 - 2*(3*a^2 + 10*a*b + 15*b^2)*cos(x)^2 + 3*a^2 + 10*a*b + 15*b^2)*log(-1/2*cos(x) + 1/2))/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cos(x)^4 + a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 2*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*cos(x)^2)]
```

Sympy [F]

$$\int \frac{\csc^5(x)}{a + b \cos^2(x)} dx = \int \frac{\csc^5(x)}{a + b \cos^2(x)} dx$$

input

```
integrate(csc(x)**5/(a+b*cos(x)**2),x)
```

output

```
Integral(csc(x)**5/(a + b*cos(x)**2), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(80) = 160$.

Time = 0.11 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.13

$$\begin{aligned} & \int \frac{\csc^5(x)}{a + b \cos^2(x)} dx \\ &= -\frac{b^3 \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{ab}} - \frac{(3a^2 + 10ab + 15b^2) \log(\cos(x) + 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} \\ &+ \frac{(3a^2 + 10ab + 15b^2) \log(\cos(x) - 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} \\ &+ \frac{(3a + 7b) \cos(x)^3 - (5a + 9b) \cos(x)}{8((a^2 + 2ab + b^2) \cos(x)^4 - 2(a^2 + 2ab + b^2) \cos(x)^2 + a^2 + 2ab + b^2)} \end{aligned}$$

input `integrate(csc(x)^5/(a+b*cos(x)^2),x, algorithm="maxima")`

output
$$\begin{aligned} & -b^3 \arctan(b \cos(x)/\sqrt{a b}) / ((a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{a b}) \\ & - 1/16 * (3a^2 + 10ab + 15b^2) \log(\cos(x) + 1) / (a^3 + 3a^2b + 3ab^2 + b^3) \\ & + 1/16 * (3a^2 + 10ab + 15b^2) \log(\cos(x) - 1) / (a^3 + 3a^2b + 3ab^2 + b^3) \\ & + 1/8 * ((3a + 7b) \cos(x)^3 - (5a + 9b) \cos(x)) / ((a^2 + 2ab + b^2) \cos(x)^4 - 2(a^2 + 2ab + b^2) \cos(x)^2 + a^2 + 2ab + b^2) \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. $2(80) = 160$.

Time = 0.12 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.89

$$\begin{aligned} & \int \frac{\csc^5(x)}{a + b \cos^2(x)} dx = -\frac{b^3 \arctan\left(\frac{b \cos(x)}{\sqrt{ab}}\right)}{(a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{ab}} \\ & - \frac{(3a^2 + 10ab + 15b^2) \log(\cos(x) + 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} \\ & + \frac{(3a^2 + 10ab + 15b^2) \log(-\cos(x) + 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} \\ & + \frac{3a \cos(x)^3 + 7b \cos(x)^3 - 5a \cos(x) - 9b \cos(x)}{8(a^2 + 2ab + b^2)(\cos(x)^2 - 1)^2} \end{aligned}$$

input `integrate(csc(x)^5/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$\begin{aligned} & -b^3 \arctan(b \cos(x) / \sqrt{a b}) / ((a^3 + 3 a^2 b + 3 a b^2 + b^3) \sqrt{a b}) \\ & - 1/16 * (3 a^2 + 10 a b + 15 b^2) \log(\cos(x) + 1) / (a^3 + 3 a^2 b + 3 a b^2 + b^3) \\ & + 1/16 * (3 a^2 + 10 a b + 15 b^2) \log(-\cos(x) + 1) / (a^3 + 3 a^2 b + 3 a b^2 + b^3) \\ & + 1/8 * (3 a \cos(x)^3 + 7 b \cos(x)^3 - 5 a \cos(x) - 9 b \cos(x)) / ((a^2 + 2 a b + b^2) (\cos(x)^2 - 1)^2) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 2.78 (sec) , antiderivative size = 833, normalized size of antiderivative = 8.86

$$\int \frac{\csc^5(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input `int(1/(sin(x)^5*(a + b*cos(x)^2)),x)`

output
$$\begin{aligned} & -(\operatorname{atan}(a \cos(x) * (-a b^5)^{(3/2)} * 64 i - b \cos(x) * (-a b^5)^{(3/2)} * 64 i + a^6 b \cos(x) * (-a b^5)^{(1/2)} * 9 i + a^2 b^5 \cos(x) * (-a b^5)^{(1/2)} * 289 i + a^3 b^4 \cos(x) * (-a b^5)^{(1/2)} * 300 i + a^4 b^3 \cos(x) * (-a b^5)^{(1/2)} * 190 i + a^5 b^2 \cos(x) * (-a b^5)^{(1/2)} * 60 i) / (64 a^2 b^8 + 225 a^3 b^7 + 300 a^4 b^6 + 190 a^5 b^5 + 60 a^6 b^4 + 9 a^7 b^3) * (-a b^5)^{(1/2)} * 8 i - 3 a^3 \cos(x)^3 + 3 a^3 * \operatorname{atanh}(\cos(x)) + 5 a^3 \cos(x) - \operatorname{atan}(a \cos(x) * (-a b^5)^{(3/2)} * 64 i - b \cos(x) * (-a b^5)^{(3/2)} * 64 i + a^6 b \cos(x) * (-a b^5)^{(1/2)} * 9 i + a^2 b^5 \cos(x) * (-a b^5)^{(1/2)} * 289 i + a^3 b^4 \cos(x) * (-a b^5)^{(1/2)} * 300 i + a^4 b^3 \cos(x) * (-a b^5)^{(1/2)} * 190 i + a^5 b^2 \cos(x) * (-a b^5)^{(1/2)} * 60 i) / (64 a^2 b^8 + 225 a^3 b^7 + 300 a^4 b^6 + 190 a^5 b^5 + 60 a^6 b^4 + 9 a^7 b^3) * \cos(x)^2 * (-a b^5)^{(1/2)} * 16 i + \operatorname{atan}(a \cos(x) * (-a b^5)^{(3/2)} * 64 i - b \cos(x) * (-a b^5)^{(3/2)} * 64 i + a^6 b \cos(x) * (-a b^5)^{(1/2)} * 9 i + a^2 b^5 \cos(x) * (-a b^5)^{(1/2)} * 289 i + a^3 b^4 \cos(x) * (-a b^5)^{(1/2)} * 300 i + a^4 b^3 \cos(x) * (-a b^5)^{(1/2)} * 190 i + a^5 b^2 \cos(x) * (-a b^5)^{(1/2)} * 60 i) / (64 a^2 b^8 + 225 a^3 b^7 + 300 a^4 b^6 + 190 a^5 b^5 + 60 a^6 b^4 + 9 a^7 b^3) * \cos(x)^4 * (-a b^5)^{(1/2)} * 8 i + 9 a^2 b^2 \cos(x) + 14 a^2 b^2 \cos(x) - 6 a^3 \operatorname{atanh}(\cos(x)) * \cos(x)^2 + 3 a^3 * \operatorname{atanh}(\cos(x)) * \cos(x)^4 - 7 a^2 b^2 \cos(x)^3 - 10 a^2 b^2 \cos(x)^3 + 15 a^2 b^2 \operatorname{atanh}(\cos(x)) + 10 a^2 b^2 \operatorname{atanh}(\cos(x)) - 30 a^2 b^2 \operatorname{atanh}(\cos(x)) * \cos(x)^2 - 20 a^2 b^2 \operatorname{atanh}(\cos(x)) * \cos(x)^2 + 15 a^2 b^2 \operatorname{atanh}(\cos(x)) * \cos(x)^4 + 10 a^2 b^2 \operatorname{atanh}(\cos(x)) * \cos(x)^4) / (8 a^4 \cos(x)^4 - 16 a^4 \cos(x)^2 + 8 a^2 b^3 \dots) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 197, normalized size of antiderivative = 2.10

$$\int \frac{\csc^5(x)}{a + b \cos^2(x)} dx \\ = \frac{8\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) \sin(x)^4 b^2 - 8\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) \sin(x)^4 b^2 - 3 \cos(x) \sin(x)^2 c^2}{}$$

input `int(csc(x)^5/(a+b*cos(x)^2),x)`

output `(8*sqrt(b)*sqrt(a)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*sin(x)**4*b**2 - 8*sqrt(b)*sqrt(a)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*sin(x)**4*b**2 - 3*cos(x)*sin(x)**2*a**3 - 10*cos(x)*sin(x)**2*a**2*b - 7*cos(x)*sin(x)**2*a*b**2 - 2*cos(x)*a**3 - 4*cos(x)*a**2*b - 2*cos(x)*a*b**2 + 3*log(tan(x/2))*sin(x)**4*a**3 + 10*log(tan(x/2))*sin(x)**4*a**2*b + 15*log(tan(x/2))*sin(x)**4*a*b**2)/(8*sin(x)**4*a*(a**3 + 3*a**2*b + 3*a*b**2 + b**3))`

3.17 $\int \frac{\sin^6(x)}{a+b\cos^2(x)} dx$

Optimal result	150
Mathematica [A] (verified)	150
Rubi [A] (verified)	151
Maple [A] (verified)	153
Fricas [A] (verification not implemented)	154
Sympy [F(-1)]	155
Maxima [A] (verification not implemented)	155
Giac [A] (verification not implemented)	155
Mupad [B] (verification not implemented)	156
Reduce [B] (verification not implemented)	157

Optimal result

Integrand size = 15, antiderivative size = 88

$$\begin{aligned} \int \frac{\sin^6(x)}{a+b\cos^2(x)} dx = & -\frac{(8a^2 + 20ab + 15b^2)x}{8b^3} - \frac{(a+b)^{5/2} \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{\sqrt{ab^3}} \\ & + \frac{(4a+7b)\cos(x)\sin(x)}{8b^2} + \frac{\cos(x)\sin^3(x)}{4b} \end{aligned}$$

output
$$-1/8*(8*a^2+20*a*b+15*b^2)*x/b^3-(a+b)^(5/2)*\arctan((a+b)^(1/2)*\cot(x)/a^(1/2))/a^(1/2)/b^3+1/8*(4*a+7*b)*\cos(x)*\sin(x)/b^2+1/4*\cos(x)*\sin(x)^3/b$$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.88

$$\begin{aligned} \int \frac{\sin^6(x)}{a+b\cos^2(x)} dx \\ = \frac{-4(8a^2 + 20ab + 15b^2)x + \frac{32(a+b)^{5/2} \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}} + 8b(a+2b)\sin(2x) - b^2\sin(4x)}{32b^3} \end{aligned}$$

input `Integrate[Sin[x]^6/(a + b*Cos[x]^2), x]`

output
$$\frac{(-4*(8*a^2 + 20*a*b + 15*b^2)*x + (32*(a + b)^(5/2)*ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]])/Sqrt[a] + 8*b*(a + 2*b)*Sin[2*x] - b^2*Sin[4*x])/(32*b^3)}$$

Rubi [A] (verified)

Time = 0.36 (sec), antiderivative size = 114, normalized size of antiderivative = 1.30, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 3670, 316, 402, 397, 216, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^6(x)}{a + b \cos^2(x)} dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cos(x + \frac{\pi}{2})^6}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \downarrow \textcolor{blue}{3670} \\
 & - \int \frac{1}{(\cot^2(x) + 1)^3 ((a + b) \cot^2(x) + a)} d \cot(x) \\
 & \downarrow \textcolor{blue}{316} \\
 & \frac{\cot(x)}{4b (\cot^2(x) + 1)^2} - \frac{\int \frac{-3(a+b)\cot^2(x)+a+4b}{(\cot^2(x)+1)^2((a+b)\cot^2(x)+a)} d \cot(x)}{4b} \\
 & \downarrow \textcolor{blue}{402} \\
 & \frac{\cot(x)}{4b (\cot^2(x) + 1)^2} - \frac{\int \frac{4a^2+9ba+8b^2-(a+b)(4a+7b)\cot^2(x)}{(\cot^2(x)+1)((a+b)\cot^2(x)+a)} d \cot(x)}{2b} - \frac{(4a+7b)\cot(x)}{2b(\cot^2(x)+1)} \\
 & \downarrow \textcolor{blue}{397} \\
 & \frac{\cot(x)}{4b (\cot^2(x) + 1)^2} - \frac{\frac{8(a+b)^3 \int \frac{1}{(a+b)\cot^2(x)+a} d \cot(x)}{b} - \frac{(8a^2+20ab+15b^2) \int \frac{1}{\cot^2(x)+1} d \cot(x)}{b}}{2b} - \frac{(4a+7b)\cot(x)}{2b(\cot^2(x)+1)} \\
 & \downarrow \textcolor{blue}{216}
 \end{aligned}$$

$$\frac{\cot(x)}{4b(\cot^2(x) + 1)^2} - \frac{\frac{8(a+b)^3 \int \frac{1}{(a+b)\cot^2(x)+a} d\cot(x)}{b} - \frac{(8a^2+20ab+15b^2)}{b} \arctan(\cot(x))}{2b} - \frac{(4a+7b)\cot(x)}{2b(\cot^2(x)+1)}$$

↓ 218

$$\frac{\cot(x)}{4b(\cot^2(x) + 1)^2} - \frac{\frac{8(a+b)^{5/2} \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{\sqrt{ab}} - \frac{(8a^2+20ab+15b^2)}{b} \arctan(\cot(x))}{2b} - \frac{(4a+7b)\cot(x)}{2b(\cot^2(x)+1)}$$

input `Int[Sin[x]^6/(a + b*Cos[x]^2), x]`

output `Cot[x]/(4*b*(1 + Cot[x]^2)^2) - ((-(((8*a^2 + 20*a*b + 15*b^2)*ArcTan[Cot[x]])/b) + (8*(a + b)^(5/2)*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(Sqrt[a]*b))/(2*b) - ((4*a + 7*b)*Cot[x])/(2*b*(1 + Cot[x]^2)))/(4*b)`

Definitions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 316 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] :> Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))), x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2, p, q, x]`

rule 397

```
Int[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), x_
Symbol] :> Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[
(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e
, f}, x]
```

rule 402

```
Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x
_)^2), x_Symbol] :> Simp[(-(b*e - a*f))*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^
(q + 1)/(a*2*(b*c - a*d)*(p + 1))), x] + Simp[1/(a*2*(b*c - a*d)*(p + 1))
Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[c*(b*e - a*f) + e*2*(b*c - a*d)
*(p + 1) + d*(b*e - a*f)*(2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b
, c, d, e, f, q}, x] && LtQ[p, -1]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3670

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^
p_, x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Su
bst[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e
+ f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Maple [A] (verified)

Time = 1.24 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.07

method	result
default	$-\frac{\frac{(-\frac{1}{2}ab - \frac{9}{8}b^2)\tan(x)^3 + (-\frac{1}{2}ab - \frac{7}{8}b^2)\tan(x)}{(1+\tan(x)^2)^2}}{b^3} + \frac{(8a^2 + 20ab + 15b^2)\arctan(\tan(x))}{8} + \frac{(a+b)^3 \arctan\left(\frac{a\tan(x)}{\sqrt{(a+b)a}}\right)}{b^3 \sqrt{(a+b)a}}$
risch	$-\frac{xa^2}{b^3} - \frac{5xa}{2b^2} - \frac{15x}{8b} - \frac{ie^{2ix}a}{8b^2} - \frac{ie^{2ix}}{4b} + \frac{ie^{-2ix}a}{8b^2} + \frac{ie^{-2ix}}{4b} - \frac{a\sqrt{-(a+b)a}\ln\left(e^{2ix} + \frac{2i\sqrt{-(a+b)a} + 2a + b}{b}\right)}{2b^3} - \frac{\sqrt{-(a+b)a}}{b}$

input `int(sin(x)^6/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -1/b^3 * (((-1/2*a*b - 9/8*b^2)*tan(x)^3 + (-1/2*a*b - 7/8*b^2)*tan(x))/(1+tan(x)^2) + 1/8*(8*a^2 + 20*a*b + 15*b^2)*arctan(tan(x))) + 1/b^3*(a+b)^3/((a+b)*a)^(1/2)*arctan(a*tan(x)/((a+b)*a)^(1/2)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.23 (sec), antiderivative size = 285, normalized size of antiderivative = 3.24

$$\begin{aligned} & \int \frac{\sin^6(x)}{a + b \cos^2(x)} dx \\ &= \frac{2(a^2 + 2ab + b^2)\sqrt{-\frac{a+b}{a}} \log \left(\frac{(8a^2 + 8ab + b^2)\cos(x)^4 - 2(4a^2 + 3ab)\cos(x)^2 - 4((2a^2 + ab)\cos(x)^3 - a^2\cos(x))\sqrt{-\frac{a+b}{a}}\sin(x) + b^2\cos(x)^4 + 2ab\cos(x)^2 + a^2}{b^2\cos(x)^4 + 2ab\cos(x)^2 + a^2} \right)}{8b^3} \\ & - \frac{4(a^2 + 2ab + b^2)\sqrt{\frac{a+b}{a}} \arctan \left(\frac{((2a+b)\cos(x)^2 - a)\sqrt{\frac{a+b}{a}}}{2(a+b)\cos(x)\sin(x)} \right) + (8a^2 + 20ab + 15b^2)x + (2b^2\cos(x)^3 - (4a^2 + 20ab + 15b^2)\cos(x))\sqrt{\frac{a+b}{a}}}{8b^3} \end{aligned}$$

input

```
integrate(sin(x)^6/(a+b*cos(x)^2), x, algorithm="fricas")
```

output

$$\begin{aligned} & [1/8*(2*(a^2 + 2*a*b + b^2)*sqrt(-(a + b)/a)*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 - 4*((2*a^2 + a*b)*cos(x)^3 - a^2*cos(x))*sqrt(-(a + b)/a)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) - (8*a^2 + 20*a*b + 15*b^2)*x - (2*b^2*cos(x)^3 - (4*a*b + 9*b^2)*cos(x))*sin(x)/b^3, -1/8*(4*(a^2 + 2*a*b + b^2)*sqrt((a + b)/a)*arctan(1/2*((2*a + b)*cos(x)^2 - a)*sqrt((a + b)/a)/((a + b)*cos(x)*sin(x))) + (8*a^2 + 20*a*b + 15*b^2)*x + (2*b^2*cos(x)^3 - (4*a*b + 9*b^2)*cos(x))*sin(x))/b^3] \end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin^6(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sin(x)**6/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.27

$$\begin{aligned} \int \frac{\sin^6(x)}{a + b \cos^2(x)} dx &= \frac{(4a + 9b) \tan(x)^3 + (4a + 7b) \tan(x)}{8(b^2 \tan(x)^4 + 2b^2 \tan(x)^2 + b^2)} - \frac{(8a^2 + 20ab + 15b^2)x}{8b^3} \\ &+ \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)ab^3}} \end{aligned}$$

input `integrate(sin(x)^6/(a+b*cos(x)^2),x, algorithm="maxima")`

output `1/8*((4*a + 9*b)*tan(x)^3 + (4*a + 7*b)*tan(x))/(b^2*tan(x)^4 + 2*b^2*tan(x)^2 + b^2) - 1/8*(8*a^2 + 20*a*b + 15*b^2)*x/b^3 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*b^3)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.35

$$\begin{aligned} \int \frac{\sin^6(x)}{a + b \cos^2(x)} dx &= -\frac{(8a^2 + 20ab + 15b^2)x}{8b^3} \\ &+ \frac{(a^3 + 3a^2b + 3ab^2 + b^3)\left(\pi\left\lfloor\frac{x}{\pi} + \frac{1}{2}\right\rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right)}{\sqrt{a^2 + ab}b^3} \\ &+ \frac{4a \tan(x)^3 + 9b \tan(x)^3 + 4a \tan(x) + 7b \tan(x)}{8(\tan(x)^2 + 1)^2 b^2} \end{aligned}$$

input `integrate(sin(x)^6/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$\begin{aligned} & -\frac{1}{8}(8a^2 + 20ab + 15b^2)x/b^3 + (a^3 + 3a^2b + 3ab^2 + b^3) \cdot (\pi * \text{floor}(x/\pi + 1/2) * \text{sgn}(a) + \arctan(a \cdot \tan(x)/\sqrt{a^2 + ab})) / (\sqrt{a^2 + ab}) \cdot b^3 \\ & + \frac{1}{8}(4a \cdot \tan(x)^3 + 9b \cdot \tan(x)^3 + 4a \cdot \tan(x) + 7b \cdot \tan(x)) / ((\tan(x)^2 + 1)^{2b^2}) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 681, normalized size of antiderivative = 7.74

$$\int \frac{\sin^6(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input `int(sin(x)^6/(a + b*cos(x)^2),x)`

output
$$\begin{aligned} & \left(\frac{(\tan(x)^3(4a + 9b))/(8b^2) + (\tan(x)(4a + 7b))/(8b^2) / (2\tan(x)^2 + \tan(x)^4 + 1) + (\text{atan}((5717a^3\tan(x))/256) / (256((15ab^2)/4 + (3665a^2b)/256 + (5717a^3)/256 + (1143a^4)/(64b) + (235a^5)/(32b^2) + (5a^6)/(4b^3))) + (3665a^2\tan(x)) / (256((15ab)/4 + (3665a^2)/256 + (5717a^3)/(256b) + (1143a^4)/(64b^2) + (235a^5)/(32b^3) + (5a^6)/(4b^4))) + (1143a^4\tan(x)) / (64((15ab^3)/4 + (5717a^3b)/256 + (1143a^4)/64 + (3665a^2b^2)/256 + (235a^5b)/(32b) + (5a^6b)/(4b^2))) + (235a^5\tan(x)) / (32((15ab^4)/4 + (1143a^4b)/64 + (235a^5b)/(32b) + (3665a^2b^3)/256 + (5717a^3b^2)/256 + (5a^6b)/(4b))) + (5a^6\tan(x)) / (4((15ab^5)/4 + (235a^5b^2)/32 + (5a^6b^2)/(4b^2))) + (15ab^20i + a^28i + b^215i) \cdot \text{I} \right) / (8b^3) - (\text{atanh}((95a^2\tan(x)) * (-a^5 - 5a^5b - a^6 - 5a^2b^4 - 10a^3b^3 - 10a^4b^2)^(1/2)) / (32 * (2ab^4 + (469a^4b)/32 + (215a^5)/32 + (287a^2b^3)/32 + (517a^3b^2)/32 + (5a^6b)/(4b))) + (5a^3\tan(x)) * (-a^5 - 5a^5b - a^6 - 5a^2b^4 - 10a^3b^3 - 10a^4b^2)^(1/2)) / (4 * (2ab^5 + (215a^5b)/32 + (5a^6b)/4 + (287a^2b^4)/32 + (517a^3b^3)/32 + (469a^4b^2)/32)) + (2a\tan(x)) * (-a^5 - 5a^5b - a^6 - 5a^2b^4 - 10a^3b^3 - 10a^4b^2)^(1/2)) / (2ab^3 + (517a^3b^2)/32 + (469a^4b)/32 + (287a^2b^2)/32 + (215a^5)... \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.72

$$\int \frac{\sin^6(x)}{a + b \cos^2(x)} dx \\ = \frac{8\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right) a^2 + 16\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right) ab + 8\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right) b^2}{}$$

input `int(sin(x)^6/(a+b*cos(x)^2),x)`

output
$$(8*\sqrt{a}*\sqrt{a+b}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a^{**2} + 16*\sqrt{a}*\sqrt{a+b}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*a*b + 8*\sqrt{a}*\sqrt{a+b}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*b^{**2} + 8*\sqrt{a}*\sqrt{a+b}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a^{**2} + 16*\sqrt{a}*\sqrt{a+b}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*a*b + 8*\sqrt{a}*\sqrt{a+b}*\operatorname{atan}((\sqrt{a+b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*b^{**2} + 2*\cos(x)*\sin(x)**3*a*b^{**2} + 4*\cos(x)*\sin(x)*a^{**2}*b + 7*\cos(x)*\sin(x)*a*b^{**2} - 8*a^{**3}*x - 20*a^{**2}*b*x - 15*a*b^{**2}*x)/(8*a*b^{**3})$$

3.18 $\int \frac{\sin^4(x)}{a+b\cos^2(x)} dx$

Optimal result	158
Mathematica [A] (verified)	158
Rubi [A] (verified)	159
Maple [A] (verified)	161
Fricas [A] (verification not implemented)	161
Sympy [F(-1)]	162
Maxima [A] (verification not implemented)	162
Giac [A] (verification not implemented)	162
Mupad [B] (verification not implemented)	163
Reduce [B] (verification not implemented)	163

Optimal result

Integrand size = 15, antiderivative size = 60

$$\int \frac{\sin^4(x)}{a+b\cos^2(x)} dx = -\frac{(2a+3b)x}{2b^2} - \frac{(a+b)^{3/2} \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{\sqrt{ab^2}} + \frac{\cos(x)\sin(x)}{2b}$$

output
$$-1/2*(2*a+3*b)*x/b^2-(a+b)^(3/2)*\operatorname{arctan}((a+b)^(1/2)*\operatorname{cot}(x)/a^(1/2))/a^(1/2)/b^2+1/2*\cos(x)*\sin(x)/b$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.87

$$\int \frac{\sin^4(x)}{a+b\cos^2(x)} dx = \frac{-4ax - 6bx + \frac{4(a+b)^{3/2} \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}} + b\sin(2x)}{4b^2}$$

input `Integrate[Sin[x]^4/(a + b*Cos[x]^2), x]`

output
$$(-4*a*x - 6*b*x + (4*(a + b)^(3/2)*\operatorname{ArcTan}[(\operatorname{Sqrt}[a]*\operatorname{Tan}[x])/\operatorname{Sqrt}[a + b]])/\operatorname{Sqrt}[a] + b*\sin[2*x])/ (4*b^2)$$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 74, normalized size of antiderivative = 1.23, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3670, 316, 397, 216, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^4(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos\left(x + \frac{\pi}{2}\right)^4}{a + b \sin\left(x + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{3670} \\
 & - \int \frac{1}{(\cot^2(x) + 1)^2 ((a + b) \cot^2(x) + a)} d \cot(x) \\
 & \quad \downarrow \text{316} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\int \frac{-(a+b) \cot^2(x) + a + 2b}{(\cot^2(x) + 1)((a+b) \cot^2(x) + a)} d \cot(x)}{2b} \\
 & \quad \downarrow \text{397} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{2(a+b)^2 \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{b} - \frac{(2a+3b) \int \frac{1}{\cot^2(x) + 1} d \cot(x)}{b}}{2b} \\
 & \quad \downarrow \text{216} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{2(a+b)^2 \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{b} - \frac{(2a+3b) \arctan(\cot(x))}{b}}{2b} \\
 & \quad \downarrow \text{218} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{2(a+b)^{3/2} \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{ab}} - \frac{(2a+3b) \arctan(\cot(x))}{b}}{2b}
 \end{aligned}$$

input `Int[Sin[x]^4/(a + b*Cos[x]^2),x]`

output
$$-1/2*(-(((2*a + 3*b)*ArcTan[Cot[x]])/b) + (2*(a + b)^(3/2)*ArcTan[(Sqrt[a] + b]*Cot[x])/Sqrt[a]])/(Sqrt[a]*b))/b + Cot[x]/(2*b*(1 + Cot[x]^2))$$

Definitions of rubi rules used

rule 216
$$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(Rt[a, 2]*Rt[b, 2]))*\text{ArcTan}[Rt[b, 2]*(x/Rt[a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \mid\mid \text{GtQ}[b, 0])$$

rule 218
$$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(Rt[a/b, 2]/a)*\text{ArcTan}[x/Rt[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&& \text{PosQ}[a/b]$$

rule 316
$$\text{Int}[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), \text{x_Symbol}] \rightarrow \text{Simp}[-b*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))), \text{x}] + \text{Simp}[1/(2*a*(p + 1)*(b*c - a*d)) \text{Int}[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*\text{Simp}[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, \text{x}], \text{x}] /; \text{FreeQ}[\{a, b, c, d, q\}, \text{x}] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{LtQ}[p, -1] \&& !(\text{!IntegerQ}[p] \&& \text{IntegerQ}[q] \&& \text{LtQ}[q, -1]) \&& \text{IntBinomialQ}[a, b, c, d, 2, p, q, \text{x}]$$

rule 397
$$\text{Int}[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*((c_) + (d_)*(x_)^2)), \text{x_Symbol}] \rightarrow \text{Simp}[(b*e - a*f)/(b*c - a*d) \text{Int}[1/(a + b*x^2), \text{x}], \text{x}] - \text{Simp}[(d*e - c*f)/(b*c - a*d) \text{Int}[1/(c + d*x^2), \text{x}], \text{x}] /; \text{FreeQ}[\{a, b, c, d, e, f\}, \text{x}]$$

rule 3042
$$\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$$

rule 3670
$$\text{Int}[\cos[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*\sin[(e_) + (f_)*(x_)]^2)^(p_), \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], \text{x}]\}, \text{Simp}[ff/f \text{Sust}[\text{Int}[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), \text{x}], \text{x}, \text{Tan}[e + f*x]/ff], \text{x}]] /; \text{FreeQ}[\{a, b, e, f\}, \text{x}] \&& \text{IntegerQ}[m/2] \&& \text{IntegerQ}[p]$$

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02

method	result
default	$-\frac{\frac{b \tan(x)}{2(1+\tan(x)^2)} + \frac{(2a+3b) \arctan(\tan(x))}{2}}{b^2} + \frac{(a+b)^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{b^2 \sqrt{(a+b)a}}$
risch	$-\frac{xa}{b^2} - \frac{3x}{2b} - \frac{ie^{2ix}}{8b} + \frac{ie^{-2ix}}{8b} + \frac{\sqrt{-(a+b)a} \ln\left(e^{2ix} - \frac{2i\sqrt{-(a+b)a}-2a-b}{b}\right)}{2b^2} + \frac{\sqrt{-(a+b)a} \ln\left(e^{2ix} - \frac{2i\sqrt{-(a+b)a}-2a-b}{b}\right)}{2ab}$

input `int(sin(x)^4/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/b^2 * (-1/2*b*tan(x)/(1+tan(x)^2) + 1/2*(2*a+3*b)*arctan(tan(x))) + (a+b)^2/b \\ & ^2/((a+b)*a)^(1/2)*arctan(a*tan(x)/((a+b)*a)^(1/2)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 211, normalized size of antiderivative = 3.52

$$\begin{aligned} & \int \frac{\sin^4(x)}{a + b \cos^2(x)} dx \\ & = \frac{2 b \cos(x) \sin(x) + (a + b) \sqrt{-\frac{a+b}{a}} \log \left(\frac{(8 a^2 + 8 a b + b^2) \cos(x)^4 - 2 (4 a^2 + 3 a b) \cos(x)^2 - 4 ((2 a^2 + a b) \cos(x)^3 - a^2 \cos(x)) \sqrt{-\frac{a+b}{a}}}{b^2 \cos(x)^4 + 2 a b \cos(x)^2 + a^2} \right)}{4 b^2} \end{aligned}$$

input `integrate(sin(x)^4/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$\begin{aligned} & [1/4*(2*b*cos(x)*sin(x) + (a + b)*sqrt(-(a + b)/a)*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 - 4*((2*a^2 + a*b)*cos(x)^3 - a^2*cos(x))*sqrt(-(a + b)/a)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) - 2*(2*a + 3*b)*x]/b^2, 1/2*(b*cos(x)*sin(x) - (a + b)*sqrt((a + b)/a)*arctan(1/2*((2*a + b)*cos(x)^2 - a)*sqrt((a + b)/a)/((a + b)*cos(x)*sin(x))) - (2*a + 3*b)*x]/b^2] \end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin^4(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sin(x)**4/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.03

$$\begin{aligned} \int \frac{\sin^4(x)}{a + b \cos^2(x)} dx &= -\frac{(2a + 3b)x}{2b^2} + \frac{\tan(x)}{2(b \tan(x)^2 + b)} \\ &+ \frac{(a^2 + 2ab + b^2) \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)ab^2}} \end{aligned}$$

input `integrate(sin(x)^4/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-1/2*(2*a + 3*b)*x/b^2 + 1/2*tan(x)/(b*tan(x)^2 + b) + (a^2 + 2*a*b + b^2)*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*b^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.33

$$\begin{aligned} \int \frac{\sin^4(x)}{a + b \cos^2(x)} dx &= -\frac{(2a + 3b)x}{2b^2} \\ &+ \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right)(a^2 + 2ab + b^2)}{\sqrt{a^2 + ab^2}} \\ &+ \frac{\tan(x)}{2(\tan(x)^2 + 1)b} \end{aligned}$$

input `integrate(sin(x)^4/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$\frac{-1/2*(2*a + 3*b)*x/b^2 + (\pi*\text{floor}(x/\pi + 1/2)*\text{sgn}(a) + \arctan(a*\tan(x))/\sqrt{a^2 + a*b})*((a^2 + 2*a*b + b^2)/(\sqrt{a^2 + a*b}*b^2) + 1/2*\tan(x)/((\tan(x)^2 + 1)*b))}{a^2 + a*b}$$

Mupad [B] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.10

$$\int \frac{\sin^4(x)}{a + b \cos^2(x)} dx = \frac{\cos(x) \sin(x)}{2b} - \frac{a \operatorname{atan}\left(\frac{\sin(x)}{\cos(x)}\right)}{b^2} - \frac{3 \operatorname{atan}\left(\frac{\sin(x)}{\cos(x)}\right)}{2b} \\ - \frac{\operatorname{atanh}\left(\frac{\sin(x) \sqrt{-a^4 - 3a^3b - 3a^2b^2 - ab^3}}{\cos(x) a^2 + 2 \cos(x) a b + \cos(x) b^2}\right) \sqrt{-a^4 - 3a^3b - 3a^2b^2 - ab^3}}{ab^2}$$

input `int(sin(x)^4/(a + b*cos(x)^2),x)`

output
$$\frac{(\cos(x)*\sin(x))/(2*b) - (a*\operatorname{atan}(\sin(x)/\cos(x)))/b^2 - (3*\operatorname{atan}(\sin(x)/\cos(x)))/(2*b) - (\operatorname{atanh}((\sin(x)*(-a*b^3 - 3*a^3*b - a^4 - 3*a^2*b^2)^(1/2))/(a^2*\cos(x) + b^2*\cos(x) + 2*a*b*\cos(x)))*(-a*b^3 - 3*a^3*b - a^4 - 3*a^2*b^2)^(1/2)/(a*b^2)}{a^2 + a*b}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 139, normalized size of antiderivative = 2.32

$$\int \frac{\sin^4(x)}{a + b \cos^2(x)} dx \\ = \frac{2\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) a + 2\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) b + 2\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) b^2}{2a b^2}$$

input `int(sin(x)^4/(a+b*cos(x)^2),x)`

```
output (2*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*a +
2*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*b + 2*
sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*a + 2*
sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*b + cos
(x)*sin(x)*a*b - 2*a**2*x - 3*a*b*x)/(2*a*b**2)
```

3.19 $\int \frac{\sin^2(x)}{a+b\cos^2(x)} dx$

Optimal result	165
Mathematica [A] (verified)	165
Rubi [A] (verified)	166
Maple [A] (verified)	167
Fricas [A] (verification not implemented)	168
Sympy [F(-1)]	168
Maxima [A] (verification not implemented)	169
Giac [A] (verification not implemented)	169
Mupad [B] (verification not implemented)	170
Reduce [B] (verification not implemented)	170

Optimal result

Integrand size = 15, antiderivative size = 40

$$\int \frac{\sin^2(x)}{a + b\cos^2(x)} dx = -\frac{x}{b} - \frac{\sqrt{a+b}\arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{\sqrt{ab}}$$

output -x/b-(a+b)^(1/2)*arctan((a+b)^(1/2)*cot(x)/a^(1/2))/a^(1/2)/b

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

$$\int \frac{\sin^2(x)}{a + b\cos^2(x)} dx = \frac{-x + \frac{\sqrt{a+b}\arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}}}{b}$$

input Integrate[Sin[x]^2/(a + b*Cos[x]^2), x]

output (-x + (Sqrt[a + b]*ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]])/Sqrt[a])/b

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3670, 303, 216, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin^2(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cos(x + \frac{\pi}{2})^2}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3670} \\
 & - \int \frac{1}{(\cot^2(x) + 1) ((a + b) \cot^2(x) + a)} d \cot(x) \\
 & \quad \downarrow \text{303} \\
 & \frac{\int \frac{1}{\cot^2(x) + 1} d \cot(x)}{b} - \frac{(a + b) \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{b} \\
 & \quad \downarrow \text{216} \\
 & \frac{\arctan(\cot(x))}{b} - \frac{(a + b) \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{b} \\
 & \quad \downarrow \text{218} \\
 & \frac{\arctan(\cot(x))}{b} - \frac{\sqrt{a + b} \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{ab}}
 \end{aligned}$$

input `Int[Sin[x]^2/(a + b*Cos[x]^2),x]`

output `ArcTan[Cot[x]]/b - (Sqrt[a + b]*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(Sqr
t[a]*b)`

Definitions of rubi rules used

rule 216 $\text{Int}[(a_ + b_ \cdot) \cdot (x_ \cdot)^2 \cdot (-1), x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(Rt[a, 2] \cdot Rt[b, 2]) \cdot A \cdot \text{rcTan}[Rt[b, 2] \cdot (x/Rt[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{PosQ}[a/b] \& (\text{GtQ}[a, 0] \mid\mid \text{GtQ}[b, 0])$

rule 218 $\text{Int}[(a_ + b_ \cdot) \cdot (x_ \cdot)^2 \cdot (-1), x_{\text{Symbol}}] \rightarrow \text{Simp}[(Rt[a/b, 2]/a) \cdot \text{ArcTan}[x/Rt[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{PosQ}[a/b]$

rule 303 $\text{Int}[1/((a_ + b_ \cdot) \cdot (x_ \cdot)^2) \cdot ((c_ + d_ \cdot) \cdot (x_ \cdot)^2), x_{\text{Symbol}}] \rightarrow \text{Simp}[b/(b \cdot c - a \cdot d) \cdot \text{Int}[1/(a + b \cdot x^2), x], x] - \text{Simp}[d/(b \cdot c - a \cdot d) \cdot \text{Int}[1/(c + d \cdot x^2), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \& \text{NeQ}[b \cdot c - a \cdot d, 0]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3670 $\text{Int}[\cos[(e_ + f_ \cdot) \cdot (x_)]^m \cdot ((a_ + b_ \cdot) \cdot \sin[(e_ + f_ \cdot) \cdot (x_)]^2)^p, x_{\text{Symbol}}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f \cdot x], x]\}, \text{Simp}[ff/f \cdot \text{Subst}[\text{Int}[(a + (a + b) \cdot ff^2 \cdot x^2)^p / (1 + ff^2 \cdot x^2)^{(m/2 + p + 1)}, x], x, \text{Tan}[e + f \cdot x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f\}, x] \& \text{IntegerQ}[m/2] \& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

method	result	size
default	$-\frac{\arctan(\tan(x))}{b} + \frac{(a+b) \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{b \sqrt{(a+b)a}}$	36
risch	$-\frac{x}{b} - \frac{\sqrt{-(a+b)a} \ln\left(e^{2ix} + \frac{2i\sqrt{-(a+b)a}+2a+b}{b}\right)}{2ab} + \frac{\sqrt{-(a+b)a} \ln\left(e^{2ix} - \frac{2i\sqrt{-(a+b)a}-2a-b}{b}\right)}{2ab}$	97

input $\text{int}(\sin(x)^2/(a+b \cdot \cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$-1/b*\arctan(\tan(x))+(a+b)/b/((a+b)*a)^(1/2)*\arctan(a*\tan(x)/((a+b)*a)^(1/2))$$

Fricas [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 177, normalized size of antiderivative = 4.42

$$\int \frac{\sin^2(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{\sqrt{-\frac{a+b}{a}} \log \left(\frac{(8a^2+8ab+b^2)\cos(x)^4 - 2(4a^2+3ab)\cos(x)^2 - 4((2a^2+ab)\cos(x)^3 - a^2\cos(x))\sqrt{-\frac{a+b}{a}}\sin(x) + a^2}{b^2\cos(x)^4 + 2ab\cos(x)^2 + a^2} \right) - 4x}{4b}, \right. \\ \left. - \frac{\sqrt{\frac{a+b}{a}} \arctan \left(\frac{((2a+b)\cos(x)^2 - a)\sqrt{\frac{a+b}{a}}}{2(a+b)\cos(x)\sin(x)} \right) + 2x}{2b} \right]$$

input `integrate(sin(x)^2/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[1/4*(\sqrt{-(a+b)/a}*\log(((8*a^2+8*a*b+b^2)*\cos(x)^4 - 2*(4*a^2+3*a*b)*\cos(x)^2 - 4*((2*a^2+a*b)*\cos(x)^3 - a^2*\cos(x))*\sqrt{-(a+b)/a}*\sin(x) + a^2)/(b^2*\cos(x)^4 + 2*a*b*\cos(x)^2 + a^2)) - 4*x]/b, -1/2*(\sqrt{(a+b)/a}*\arctan(1/2*((2*a+b)*\cos(x)^2 - a)*\sqrt{(a+b)/a}/((a+b)*\cos(x)*\sin(x))) + 2*x)/b]$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin^2(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sin(x)**2/(a+b*cos(x)**2),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.82

$$\int \frac{\sin^2(x)}{a + b \cos^2(x)} dx = \frac{(a + b) \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)ab}} - \frac{x}{b}$$

input `integrate(sin(x)^2/(a+b*cos(x)^2),x, algorithm="maxima")`

output `(a + b)*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*b) - x/b`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.25

$$\int \frac{\sin^2(x)}{a + b \cos^2(x)} dx = \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2+ab}}\right)\right)(a + b)}{\sqrt{a^2 + ab}} - \frac{x}{b}$$

input `integrate(sin(x)^2/(a+b*cos(x)^2),x, algorithm="giac")`

output `(pi*floor(x/pi + 1/2)*sgn(a) + arctan(a*tan(x)/sqrt(a^2 + a*b)))*(a + b)/(sqrt(a^2 + a*b)*b) - x/b`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 108, normalized size of antiderivative = 2.70

$$\int \frac{\sin^2(x)}{a + b \cos^2(x)} dx = -\frac{\operatorname{atan}\left(\frac{2ab^2 \tan(x)}{2a^2b + 2ab^2} + \frac{2a^2b \tan(x)}{2a^2b + 2ab^2}\right)}{b}$$

$$-\frac{\operatorname{atanh}\left(\frac{2a^2b \tan(x)\sqrt{-a^2 - ba}}{2a^3b + 2a^2b^2}\right) \sqrt{-a(a + b)}}{ab}$$

input `int(sin(x)^2/(a + b*cos(x)^2),x)`

output
$$-\operatorname{atan}\left(\frac{(2ab^2 \tan(x))}{(2a^2b^2 + 2a^2b)} + \frac{(2a^2b \tan(x))}{(2a^2b^2 + 2a^2b)}\right)/b - \left(\operatorname{atanh}\left(\frac{(2a^2b \tan(x))(-a^2 - ab)}{(2a^3b + 2a^2b^2)}\right)(-a^2 - ab)^{1/2}\right)/(a^2b^2)$$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.60

$$\int \frac{\sin^2(x)}{a + b \cos^2(x)} dx$$

$$= \frac{\sqrt{a} \sqrt{a + b} \operatorname{atan}\left(\frac{\sqrt{a + b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) + \sqrt{a} \sqrt{a + b} \operatorname{atan}\left(\frac{\sqrt{a + b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) - ax}{ab}$$

input `int(sin(x)^2/(a+b*cos(x)^2),x)`

output
$$\left(\sqrt{a} \sqrt{a + b} \operatorname{atan}\left(\frac{\sqrt{a + b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) + \sqrt{a} \sqrt{a + b} \operatorname{atan}\left(\frac{\sqrt{a + b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) - a x\right)/(a b)$$

3.20 $\int \frac{1}{a+b \cos^2(x)} dx$

Optimal result	171
Mathematica [A] (verified)	171
Rubi [A] (verified)	172
Maple [A] (verified)	173
Fricas [B] (verification not implemented)	173
Sympy [B] (verification not implemented)	174
Maxima [A] (verification not implemented)	175
Giac [A] (verification not implemented)	175
Mupad [B] (verification not implemented)	175
Reduce [B] (verification not implemented)	176

Optimal result

Integrand size = 10, antiderivative size = 30

$$\int \frac{1}{a + b \cos^2(x)} dx = -\frac{\arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{a+b}}$$

output `-arctan((a+b)^(1/2)*cot(x)/a^(1/2))/a^(1/2)/(a+b)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

input `Integrate[(a + b*Cos[x]^2)^(-1), x]`

output `ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]]/(Sqrt[a]*Sqrt[a + b])`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 3660, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{a + b \sin^2(x)} dx \\
 & \quad \downarrow \text{3660} \\
 & - \int \frac{1}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \text{218} \\
 & - \frac{\arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{a+b}}
 \end{aligned}$$

input `Int[(a + b*Cos[x]^2)^(-1), x]`

output `-(ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]]/(Sqrt[a]*Sqrt[a + b]))`

Definitions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3660

```
Int[((a_) + (b_)*sin[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] :> With[{ff =
FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[1/(a + (a + b)*ff^2*x^
2), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]
```

Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}}$	21
risch	$-\frac{\ln\left(\frac{e^{2ix} + 2ia^2 + 2iab + 2a\sqrt{-a^2-ab} + b\sqrt{-a^2-ab}}{b\sqrt{-a^2-ab}}\right)}{2\sqrt{-a^2-ab}} + \frac{\ln\left(\frac{e^{2ix} - 2ia^2 + 2iab - 2a\sqrt{-a^2-ab} - b\sqrt{-a^2-ab}}{b\sqrt{-a^2-ab}}\right)}{2\sqrt{-a^2-ab}}$	160

input `int(1/(a+b*cos(x)^2), x, method=_RETURNVERBOSE)`

output `1/((a+b)*a)^(1/2)*arctan(a*tan(x)/((a+b)*a)^(1/2))`

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(22) = 44$.

Time = 0.19 (sec) , antiderivative size = 163, normalized size of antiderivative = 5.43

$$\begin{aligned} & \int \frac{1}{a + b \cos^2(x)} dx \\ &= \left[-\frac{\sqrt{-a^2 - ab} \log \left(\frac{(8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 + 4((2a+b)\cos(x)^3 - a\cos(x))\sqrt{-a^2-ab}\sin(x) + a^2}{b^2 \cos(x)^4 + 2ab \cos(x)^2 + a^2} \right)}{4(a^2 + ab)}, \right. \\ & \quad \left. - \frac{\arctan \left(\frac{(2a+b)\cos(x)^2 - a}{2\sqrt{a^2+ab}\cos(x)\sin(x)} \right)}{2\sqrt{a^2+ab}} \right] \end{aligned}$$

input `integrate(1/(a+b*cos(x)^2), x, algorithm="fricas")`

output

```
[-1/4*sqrt(-a^2 - a*b)*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 + 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2))/(a^2 + a*b), -1/2*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))/sqrt(a^2 + a*b)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10924 vs. $2(29) = 58$.

Time = 19.45 (sec) , antiderivative size = 10924, normalized size of antiderivative = 364.13

$$\int \frac{1}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input

```
integrate(1/(a+b*cos(x)**2),x)
```

output

```
Piecewise((zoo*tan(x/2)/(tan(x/2)**2 - 1), Eq(a, 0) & Eq(b, 0)), (-tan(x/2)/(2*b) + 1/(2*b*tan(x/2)), Eq(a, -b)), (-2*tan(x/2)/(b*(tan(x/2)**2 - 1)), Eq(a, 0)), (a**3*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(-sqrt(-a/(a + b)) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan(x/2))/(2*a**4*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a**3*sqrt(-a*b)*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a*b**2*sqrt(-a*b)*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - a**3*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(sqrt(-a/(a + b)) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan(x/2))/(2*a**4*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a**3*sqrt(-a*b)*sqrt(-a/(a + b)) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)...)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}}$$

input `integrate(1/(a+b*cos(x)^2),x, algorithm="maxima")`

output `arctan(a*tan(x)/sqrt((a + b)*a))/sqrt((a + b)*a)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.23

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\pi \left\lfloor \frac{x}{\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2+ab}}\right)}{\sqrt{a^2+ab}}$$

input `integrate(1/(a+b*cos(x)^2),x, algorithm="giac")`

output `(pi*floor(x/pi + 1/2)*sgn(a) + arctan(a*tan(x)/sqrt(a^2 + a*b)))/sqrt(a^2 + a*b)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\operatorname{atan}\left(\frac{a \tan(x)}{\sqrt{a^2+ba}}\right)}{\sqrt{a^2+ba}}$$

input `int(1/(a + b*cos(x)^2),x)`

output `atan((a*tan(x))/(a*b + a^2)^(1/2))/(a*b + a^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.80

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\sqrt{a} \sqrt{a+b} \left(\operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) + \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) \right)}{a(a+b)}$$

input `int(1/(a+b*cos(x)^2),x)`

output `(sqrt(a)*sqrt(a + b)*(atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a)) + atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))))/(a*(a + b))`

3.21 $\int \frac{\csc^2(x)}{a+b \cos^2(x)} dx$

Optimal result	177
Mathematica [A] (verified)	177
Rubi [A] (verified)	178
Maple [A] (verified)	179
Fricas [B] (verification not implemented)	180
Sympy [F]	180
Maxima [A] (verification not implemented)	181
Giac [A] (verification not implemented)	181
Mupad [B] (verification not implemented)	181
Reduce [B] (verification not implemented)	182

Optimal result

Integrand size = 15, antiderivative size = 41

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx = -\frac{b \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^{3/2}} - \frac{\cot(x)}{a+b}$$

output
$$-\frac{b \operatorname{arctan}\left(\frac{(a+b)^{1/2} \cot(x)}{a^{1/2}}\right)}{\sqrt{a} (a+b)^{3/2}} - \frac{\cot(x)}{a+b}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.98

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx = \frac{b \arctan\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{3/2}} - \frac{\cot(x)}{a+b}$$

input
$$\operatorname{Integrate}[\operatorname{Csc}[x]^2/(a + b \operatorname{Cos}[x]^2), x]$$

output
$$\left(\frac{b \operatorname{ArcTan}\left(\frac{\operatorname{Sqrt}[a] \operatorname{Tan}[x]}{\operatorname{Sqrt}[a+b]}\right)}{\operatorname{Sqrt}[a] (a+b)^{3/2}} - \frac{\operatorname{Cot}[x]}{a+b} \right)$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3670, 299, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc^2(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cos(x + \frac{\pi}{2})^2 \left(a + b \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3670} \\
 & - \int \frac{\cot^2(x) + 1}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \textcolor{blue}{299} \\
 & - \frac{b \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{a + b} - \frac{\cot(x)}{a + b} \\
 & \quad \downarrow \textcolor{blue}{218} \\
 & - \frac{b \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^{3/2}} - \frac{\cot(x)}{a + b}
 \end{aligned}$$

input `Int[Csc[x]^2/(a + b*Cos[x]^2),x]`

output `-((b*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(Sqrt[a]*(a + b)^(3/2))) - Cot[x]/(a + b)`

Definitions of rubi rules used

rule 218 $\text{Int}[(a_+ + b_-)(x_-)^2]^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b]$

rule 299 $\text{Int}[(a_+ + b_-)(x_-)^2]^{(p_-)} * ((c_+ + d_-)(x_-)^2), \text{x_Symbol}] \rightarrow \text{Simp}[d*x * ((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - \text{Simp}[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) * \text{Int}[(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[2*p + 3, 0]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3670 $\text{Int}[\cos[(e_- + f_-)(x_-)]^{(m_-)} * ((a_+ + b_-)*\sin[(e_- + f_-)(x_-)]^2)^{(p_-)}, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Simp}[ff/f \text{ Subst}[\text{Int}[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, \text{Tan}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f\}, x] \&& \text{IntegerQ}[m/2] \&& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.22 (sec), antiderivative size = 39, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{1}{(a+b)\tan(x)} + \frac{b \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{(a+b)\sqrt{(a+b)a}}$	39
risch	$-\frac{2i}{(e^{2ix}-1)(a+b)} + \frac{b \ln\left(\frac{e^{2ix} + \frac{-2ia^2 - 2iab + 2a\sqrt{-a^2-ab} + b\sqrt{-a^2-ab}}{\sqrt{-a^2-ab}b}}{2\sqrt{-a^2-ab}(a+b)}\right)}{2\sqrt{-a^2-ab}(a+b)} - \frac{b \ln\left(\frac{e^{2ix} + \frac{2ia^2 + 2iab + 2a\sqrt{-a^2-ab} + b\sqrt{-a^2-ab}}{b\sqrt{-a^2-ab}}}{2\sqrt{-a^2-ab}(a+b)}\right)}{2\sqrt{-a^2-ab}(a+b)}$	18

input $\text{int}(\csc(x)^2/(a+b*\cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output $-1/(a+b)/\tan(x) + b/(a+b)/((a+b)*a)^(1/2)*\arctan(a*\tan(x))/((a+b)*a)^(1/2))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(33) = 66$.

Time = 0.19 (sec) , antiderivative size = 228, normalized size of antiderivative = 5.56

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx$$

$$= \left[-\frac{\sqrt{-a^2 - ab} b \log \left(\frac{(8 a^2 + 8 a b + b^2) \cos(x)^4 - 2 (4 a^2 + 3 a b) \cos(x)^2 + 4 ((2 a + b) \cos(x)^3 - a \cos(x)) \sqrt{-a^2 - ab} \sin(x) + a^2}{b^2 \cos(x)^4 + 2 a b \cos(x)^2 + a^2} \right) \sin(x)}{4 (a^3 + 2 a^2 b + a b^2) \sin(x)} \right.$$

$$\left. - \frac{\sqrt{a^2 + ab} b \arctan \left(\frac{(2 a + b) \cos(x)^2 - a}{2 \sqrt{a^2 + ab} \cos(x) \sin(x)} \right) \sin(x) + 2 (a^2 + a b) \cos(x)}{2 (a^3 + 2 a^2 b + a b^2) \sin(x)} \right]$$

input `integrate(csc(x)^2/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[-1/4*(sqrt(-a^2 - a*b)*b*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 + 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2))*sin(x) + 4*(a^2 + a*b)*cos(x))/((a^3 + 2*a^2*b + a*b^2)*sin(x)), -1/2*(sqrt(a^2 + a*b)*b*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))*sin(x) + 2*(a^2 + a*b)*cos(x))/((a^3 + 2*a^2*b + a*b^2)*sin(x))]`

Sympy [F]

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx = \int \frac{\csc^2(x)}{a + b \cos^2(x)} dx$$

input `integrate(csc(x)**2/(a+b*cos(x)**2),x)`

output `Integral(csc(x)**2/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx = \frac{b \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}(a+b)} - \frac{1}{(a+b) \tan(x)}$$

input `integrate(csc(x)^2/(a+b*cos(x)^2),x, algorithm="maxima")`

output `b*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*(a + b)) - 1/((a + b)*tan(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.34

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx = \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2+ab}}\right)\right)b}{\sqrt{a^2+ab}(a+b)} - \frac{1}{(a+b) \tan(x)}$$

input `integrate(csc(x)^2/(a+b*cos(x)^2),x, algorithm="giac")`

output `(pi*floor(x/pi + 1/2)*sgn(a) + arctan(a*tan(x)/sqrt(a^2 + a*b)))*b/(sqrt(a^2 + a*b)*(a + b)) - 1/((a + b)*tan(x))`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx = \frac{b \operatorname{atan}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a} (a+b)^{3/2}} - \frac{1}{\tan(x) (a+b)}$$

input `int(1/(\sin(x)^2*(a + b*cos(x)^2)),x)`

output
$$\frac{(b*\operatorname{atan}((a^{1/2}*\tan(x))/(a + b)^{1/2}))/((a^{1/2}*(a + b)^{3/2}) - 1/(\tan(x)*(a + b)))}{}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.27

$$\int \frac{\csc^2(x)}{a + b \cos^2(x)} dx \\ = \frac{\sqrt{a} \sqrt{a + b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)-\sqrt{b}}{\sqrt{a}}\right) \sin(x) b + \sqrt{a} \sqrt{a + b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)+\sqrt{b}}{\sqrt{a}}\right) \sin(x) b - \cos(x) a^2 - \cos(x) a (a^2 + 2ab + b^2)}{\sin(x) a (a^2 + 2ab + b^2)}$$

input `int(csc(x)^2/(a+b*cos(x)^2),x)`

output
$$\frac{(\sqrt{a})*\sqrt{a + b}*\operatorname{atan}((\sqrt{a + b}*\tan(x/2) - \sqrt{b})/\sqrt{a})*\sin(x)*b + \sqrt{a}*\sqrt{a + b}*\operatorname{atan}((\sqrt{a + b}*\tan(x/2) + \sqrt{b})/\sqrt{a})*\sin(x)*b - \cos(x)*a^{**2} - \cos(x)*a*b}/(\sin(x)*a*(a^{**2} + 2*a*b + b^{**2}))$$

3.22 $\int \frac{\csc^4(x)}{a+b \cos^2(x)} dx$

Optimal result	183
Mathematica [A] (verified)	183
Rubi [A] (verified)	184
Maple [A] (verified)	185
Fricas [B] (verification not implemented)	186
Sympy [F]	186
Maxima [A] (verification not implemented)	187
Giac [A] (verification not implemented)	187
Mupad [B] (verification not implemented)	188
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 15, antiderivative size = 61

$$\int \frac{\csc^4(x)}{a+b \cos^2(x)} dx = -\frac{b^2 \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^{5/2}} - \frac{(a+2b) \cot(x)}{(a+b)^2} - \frac{\cot^3(x)}{3(a+b)}$$

output
$$\frac{-b^2 \arctan((a+b)^{(1/2)} \cot(x)/a^{(1/2)})/a^{(1/2)}/(a+b)^{(5/2)} - (a+2*b)*\cot(x)/(a+b)^2 - \cot^3(x)/(3*a+3*b)}{(a+b)^2 - \cot(x)^3/(3*a+3*b)}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.97

$$\int \frac{\csc^4(x)}{a+b \cos^2(x)} dx = \frac{b^2 \arctan\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{5/2}} - \frac{\cot(x) (2a + 5b + (a+b) \csc^2(x))}{3(a+b)^2}$$

input `Integrate[Csc[x]^4/(a + b*Cos[x]^2), x]`

output
$$\frac{(b^2 \text{ArcTan}[(\text{Sqrt}[a] \text{Tan}[x])/\text{Sqrt}[a+b]])/(\text{Sqrt}[a] (a+b)^{(5/2)}) - (\text{Cot}[x] ((2 a + 5 b + (a+b) \csc[x]^2))/(3 (a+b)^2))}{(a+b)^2}$$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3670, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc^4(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cos(x + \frac{\pi}{2})^4 \left(a + b \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3670} \\
 & - \int \frac{(\cot^2(x) + 1)^2}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \textcolor{blue}{300} \\
 & - \int \left(\frac{b^2}{(a + b)^2 ((a + b) \cot^2(x) + a)} + \frac{\cot^2(x)}{a + b} + \frac{a + 2b}{(a + b)^2} \right) d \cot(x) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & - \frac{b^2 \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^{5/2}} - \frac{\cot^3(x)}{3(a+b)} - \frac{(a+2b) \cot(x)}{(a+b)^2}
 \end{aligned}$$

input `Int[Csc[x]^4/(a + b*Cos[x]^2),x]`

output `-((b^2*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(Sqrt[a]*(a + b)^(5/2))) - ((a + 2*b)*Cot[x])/(a + b)^2 - Cot[x]^3/(3*(a + b))`

Definitions of rubi rules used

rule 300 $\text{Int}[(a_.) + (b_.)*(x_)^2]^{(p_.)} * ((c_.) + (d_.)*(x_)^2)]^{(q_.)}, \text{x_Symbol}] \rightarrow \text{Int}[\text{PolynomialDivide}[(a + b*x^2)^p, (c + d*x^2)^{-q}], x] /; \text{FreeQ}\{a, b, c, d\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{IGtQ}[p, 0] \& \text{ILtQ}[q, 0] \& \text{GeQ}[p, -q]$

rule 2009 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3670 $\text{Int}[\cos[(e_.) + (f_.)*(x_)]^{(m_.)} * ((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^2)]^{(p_.)}, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Simp}[ff/f \text{ Subst}[\text{Int}[(a + (a + b)*ff^2*x^2)^p / (1 + ff^2*x^2)^{(m/2 + p + 1)}, x], x, \text{Tan}[e + f*x]/ff], x]] /; \text{FreeQ}\{a, b, e, f\}, x] \& \text{IntegerQ}[m/2] \& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.93

method	result
default	$\frac{b^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{(a+b)^2 \sqrt{(a+b)a}} - \frac{1}{3(a+b) \tan(x)^3} - \frac{a+2b}{(a+b)^2 \tan(x)}$
risch	$-\frac{2i(3e^{4ix}b - 6ae^{2ix} - 12e^{2ix}b + 2a + 5b)}{3(e^{2ix} - 1)^3(a+b)^2} - \frac{b^2 \ln\left(\frac{e^{2ix} + 2ia^2 + 2iab + 2a\sqrt{-a^2 - ab} + b\sqrt{-a^2 - ab}}{b\sqrt{-a^2 - ab}}\right)}{2\sqrt{-a^2 - ab}(a+b)^2} + \frac{b^2 \ln\left(\frac{e^{2ix} - 2ia^2 + 2iab - 2a\sqrt{-a^2 - ab}}{b\sqrt{-a^2 - ab}}\right)}{2\sqrt{-a^2 - ab}(a+b)^2}$

input $\text{int}(\csc(x)^4 / (a+b*\cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$\frac{1}{(a+b)^2 b^2} \frac{((a+b)*a)^{(1/2)} \arctan(a*\tan(x)) / ((a+b)*a)^{(1/2)}}{\tan(x)^3} - \frac{1}{3(a+b)} + \frac{(a+2*b)/(a+b)^2}{\tan(x)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(51) = 102$.

Time = 0.20 (sec) , antiderivative size = 396, normalized size of antiderivative = 6.49

$$\int \frac{\csc^4(x)}{a + b \cos^2(x)} dx$$

$$= \left[\frac{4(2a^3 + 7a^2b + 5ab^2)\cos(x)^3 + 3(b^2\cos(x)^2 - b^2)\sqrt{-a^2 - ab}\log\left(\frac{(8a^2 + 8ab + b^2)\cos(x)^4 - 2(4a^2 + 3ab)\cos(x)^2}{b^2\cos(x)^2}\right)}{12(a^4 + 3a^3b + 3a^2b^2 + ab^3 - (a^4 + 3a^3b + 3a^2b^2 + ab^3)\cos(x)^2)} \right]$$

input `integrate(csc(x)^4/(a+b*cos(x)^2),x, algorithm="fricas")`

output

```
[1/12*(4*(2*a^3 + 7*a^2*b + 5*a*b^2)*cos(x)^3 + 3*(b^2*cos(x)^2 - b^2)*sqrt(-a^2 - a*b)*log((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 + 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)*sin(x) - 12*(a^3 + 3*a^2*b + 2*a*b^2)*cos(x))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cos(x)^2)*sin(x)), 1/6*(2*(2*a^3 + 7*a^2*b + 5*a*b^2)*cos(x)^3 + 3*(b^2*cos(x)^2 - b^2)*sqrt(a^2 + a*b)*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))*sin(x) - 6*(a^3 + 3*a^2*b + 2*a*b^2)*cos(x))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cos(x)^2)*sin(x))]
```

Sympy [F]

$$\int \frac{\csc^4(x)}{a + b \cos^2(x)} dx = \int \frac{\csc^4(x)}{a + b \cos^2(x)} dx$$

input `integrate(csc(x)**4/(a+b*cos(x)**2),x)`

output `Integral(csc(x)**4/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.15

$$\int \frac{\csc^4(x)}{a + b \cos^2(x)} dx = \frac{b^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}(a^2 + 2ab + b^2)} - \frac{3(a+2b)\tan(x)^2 + a + b}{3(a^2 + 2ab + b^2)\tan(x)^3}$$

input `integrate(csc(x)^4/(a+b*cos(x)^2),x, algorithm="maxima")`

output $b^{2 \cdot} \arctan(a \cdot \tan(x)) / \sqrt{(a + b) \cdot a}) / (\sqrt{(a + b) \cdot a}) \cdot (a^{2 \cdot} + 2 \cdot a \cdot b + b^{2 \cdot})$
 $- 1/3 \cdot (3 \cdot (a + 2 \cdot b) \cdot \tan(x)^{2 \cdot} + a + b) / ((a^{2 \cdot} + 2 \cdot a \cdot b + b^{2 \cdot}) \cdot \tan(x)^{3 \cdot})$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.48

$$\int \frac{\csc^4(x)}{a + b \cos^2(x)} dx = \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right) b^2}{(a^2 + 2ab + b^2) \sqrt{a^2 + ab}} - \frac{3a \tan(x)^2 + 6b \tan(x)^2 + a + b}{3(a^2 + 2ab + b^2) \tan(x)^3}$$

input `integrate(csc(x)^4/(a+b*cos(x)^2),x, algorithm="giac")`

output $(\pi * \operatorname{floor}(x/\pi + 1/2) * \operatorname{sgn}(a) + \arctan(a \cdot \tan(x)) / \sqrt{a^2 + a \cdot b})) \cdot b^{2 \cdot} / ((a^{2 \cdot} + 2 \cdot a \cdot b + b^{2 \cdot}) \cdot \sqrt{a^2 + a \cdot b}) - 1/3 \cdot (3 \cdot a \cdot \tan(x)^{2 \cdot} + 6 \cdot b \cdot \tan(x)^{2 \cdot} + a + b) / ((a^{2 \cdot} + 2 \cdot a \cdot b + b^{2 \cdot}) \cdot \tan(x)^{3 \cdot})$

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.10

$$\int \frac{\csc^4(x)}{a + b \cos^2(x)} dx = \frac{b^2 \operatorname{atan}\left(\frac{\sqrt{a} \tan(x) (a^2 + 2ab + b^2)}{(a+b)^{5/2}}\right)}{\sqrt{a} (a+b)^{5/2}} - \frac{\frac{1}{3(a+b)} + \frac{\tan(x)^2 (a+2b)}{(a+b)^2}}{\tan(x)^3}$$

input `int(1/(sin(x)^4*(a + b*cos(x)^2)),x)`

output `(b^2*atan((a^(1/2)*tan(x)*(2*a*b + a^2 + b^2))/(a + b)^(5/2)))/(a^(1/2)*(a + b)^(5/2)) - (1/(3*(a + b)) + (tan(x)^2*(a + 2*b))/(a + b)^2)/tan(x)^3`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.57

$$\begin{aligned} & \int \frac{\csc^4(x)}{a + b \cos^2(x)} dx \\ &= \frac{3\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) \sin(x)^3 b^2 + 3\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) \sin(x)^3 b^2 - 2 \cos(x) \cdot \\ & \quad 3 \sin(x)^3 a (a^3 + 3a^2b + ab^2 + b^3)}{3 \sin(x)^3 a (a^3 + 3a^2b + ab^2 + b^3)} \end{aligned}$$

input `int(csc(x)^4/(a+b*cos(x)^2),x)`

output `(3*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*sin(x)**3*b**2 + 3*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*sin(x)**3*b**2 - 2*cos(x)*sin(x)**2*a**3 - 7*cos(x)*sin(x)**2*a**2*b - 5*cos(x)*sin(x)**2*a*b**2 - cos(x)*a**3 - 2*cos(x)*a**2*b - cos(x)*a*b**2)/(3*sin(x)**3*a*(a**3 + 3*a**2*b + 3*a*b**2 + b**3))`

3.23 $\int \frac{\csc^6(x)}{a+b\cos^2(x)} dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [A] (verified)	191
Fricas [B] (verification not implemented)	192
Sympy [F]	193
Maxima [A] (verification not implemented)	193
Giac [B] (verification not implemented)	194
Mupad [B] (verification not implemented)	194
Reduce [B] (verification not implemented)	195

Optimal result

Integrand size = 15, antiderivative size = 89

$$\begin{aligned} \int \frac{\csc^6(x)}{a+b\cos^2(x)} dx = & -\frac{b^3 \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^{7/2}} - \frac{(a^2+3ab+3b^2)\cot(x)}{(a+b)^3} \\ & - \frac{(2a+3b)\cot^3(x)}{3(a+b)^2} - \frac{\cot^5(x)}{5(a+b)} \end{aligned}$$

output
$$-\frac{b^3 \arctan((a+b)^{1/2} \cot(x)/a^{1/2})}{\sqrt{a}(a+b)^{7/2}} - \frac{(a^2+3a*b+3b^2)^2 \cot(x)}{(a+b)^3} - \frac{1}{3} \frac{(2a+3b) \cot^3(x)}{(a+b)^2} - \frac{\cot^5(x)}{5(a+b)}$$

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.01

$$\begin{aligned} \int \frac{\csc^6(x)}{a+b\cos^2(x)} dx \\ = & \frac{b^3 \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{7/2}} \\ & - \frac{\cot(x) (8a^2 + 26ab + 33b^2 + (4a^2 + 13ab + 9b^2) \csc^2(x) + 3(a+b)^2 \csc^4(x))}{15(a+b)^3} \end{aligned}$$

input $\text{Integrate}[\text{Csc}[x]^6/(a + b*\text{Cos}[x]^2), x]$

output $(b^3 \text{ArcTan}[(\text{Sqrt}[a]*\text{Tan}[x])/(\text{Sqrt}[a+b])]/(\text{Sqrt}[a]*(a+b)^{(7/2)}) - (\text{Cot}[x] * (8*a^2 + 26*a*b + 33*b^2 + (4*a^2 + 13*a*b + 9*b^2)*\text{Csc}[x]^2 + 3*(a+b)^2*\text{Csc}[x]^4))/(15*(a+b)^3)$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3670, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\csc^6(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\cos(x + \frac{\pi}{2})^6 \left(a + b \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \text{3670} \\
 & - \int \frac{(\cot^2(x) + 1)^3}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \text{300} \\
 & - \int \left(\frac{\cot^4(x)}{a + b} + \frac{(2a + 3b) \cot^2(x)}{(a + b)^2} + \frac{a^2 + 3ba + 3b^2}{(a + b)^3} + \frac{b^3}{(a + b)^3 ((a + b) \cot^2(x) + a)} \right) d \cot(x) \\
 & \quad \downarrow \text{2009} \\
 & - \frac{(a^2 + 3ab + 3b^2) \cot(x)}{(a + b)^3} - \frac{b^3 \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^{7/2}} - \frac{\cot^5(x)}{5(a+b)} - \frac{(2a + 3b) \cot^3(x)}{3(a + b)^2}
 \end{aligned}$$

input $\text{Int}[\text{Csc}[x]^6/(a + b*\text{Cos}[x]^2), x]$

output

$$-\left(\frac{(b^3 \operatorname{ArcTan}\left[\frac{(a+b) \cot(x)}{\sqrt{a}}\right])}{\sqrt{a} (a+b)^{7/2}}\right) - \left(\frac{a^2+3 a b+3 b^2 \cot(x)}{(a+b)^3}\right) - \left(\frac{(2 a+3 b) \cot(x)^3}{3 (a+b)^2}\right) - \left(\frac{\cot(x)^5}{5 (a+b)}\right)$$

Definitions of rubi rules used

rule 300

$$\text{Int}[(a_.) + (b_.)*(x_)^2]^{(p_.)}*((c_.) + (d_.)*(x_)^2)^{(q_.)}, \text{x_Symbol}] \rightarrow \text{Int}[\text{PolynomialDivide}[(a+b*x^2)^p, (c+d*x^2)^{-q}], \text{x}], \text{x}] /; \text{FreeQ}[\{a, b, c, d\}, \text{x}] \& \text{NeQ}[b*c - a*d, 0] \& \text{IGtQ}[p, 0] \& \text{ILtQ}[q, 0] \& \text{GeQ}[p, -q]$$

rule 2009

$$\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \text{x}], \text{x}] /; \text{SumQ}[u]$$

rule 3042

$$\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$$

rule 3670

$$\text{Int}[\cos[(e_.) + (f_.)*(x_)]^{(m_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^2)^{(p_.)}, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\tan[e+f*x], \text{x}]\}, \text{Simp}[ff/f \text{ Subst}[\text{Int}[(a+(a+b)*ff^2*x^2)^p/(1+ff^2*x^2)^{(m/2+p+1)}, \text{x}], \text{x}, \tan[e+f*x]/ff], \text{x}]] /; \text{FreeQ}[\{a, b, e, f\}, \text{x}] \& \text{IntegerQ}[m/2] \& \text{IntegerQ}[p]$$

Maple [A] (verified)

Time = 0.77 (sec), antiderivative size = 83, normalized size of antiderivative = 0.93

method	result
default	$\frac{b^3 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{(a+b)^3 \sqrt{(a+b)a}} - \frac{1}{5(a+b) \tan(x)^5} - \frac{2a+3b}{3(a+b)^2 \tan(x)^3} - \frac{a^2+3ab+3b^2}{(a+b)^3 \tan(x)}$
risch	$-\frac{2i(15b^2e^{8ix}-30abe^{6ix}-90b^2e^{6ix}+80a^2e^{4ix}+230abe^{4ix}+240b^2e^{4ix}-40a^2e^{2ix}-130ab e^{2ix}-150e^{2ix}b^2+8a^2+26ab+33b^2)}{15(a+b)^3(e^{2ix}-1)^5} + \dots$

input

$$\text{int}(\csc(x)^6/(a+b*\cos(x)^2), \text{x}, \text{method}=\text{_RETURNVERBOSE})$$

output
$$\frac{1}{(a+b)^3 b^3} \frac{1}{((a+b)a)^{(1/2)}} \arctan(a \tan(x)) \frac{1}{((a+b)a)^{(1/2)}} - \frac{1}{5} \frac{1}{(a+b)} t \tan(x)^5 - \frac{1}{3} \frac{(2a+3b)}{(a+b)^2} \frac{1}{\tan(x)^3} - \frac{(a^2+3ab+3b^2)}{(a+b)^3} \frac{1}{\tan(x)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 273 vs. $2(77) = 154$.

Time = 0.19 (sec) , antiderivative size = 610, normalized size of antiderivative = 6.85

$$\int \frac{\csc^6(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input `integrate(csc(x)^6/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[-\frac{1}{60} (4(8a^4 + 34a^3b + 59a^2b^2 + 33ab^3)\cos(x)^5 - 20(4a^4 + 17a^3b + 28a^2b^2 + 15ab^3)\cos(x)^3 + 15(b^3\cos(x)^4 - 2b^3\cos(x)^2 + b^3)\sqrt{-a^2 - ab}\log((8a^2 + 8ab + b^2)\cos(x)^4 - 2(4a^2 + 3ab)\cos(x)^2 + 4((2a + b)\cos(x)^3 - a\cos(x))\sqrt{-a^2 - ab})\sin(x) + a^2)/(b^2\cos(x)^4 + 2ab\cos(x)^2 + a^2)\sin(x) + 60(a^4 + 4a^3b + 6a^2b^2 + 3ab^3)\cos(x))/((a^5 + 4a^4b + 6a^3b^2 + 4a^2b^3 + ab^4)\cos(x)^4 - 2(a^5 + 4a^4b + 6a^3b^2 + 4a^2b^3 + ab^4)\cos(x)^2)\sin(x)], -\frac{1}{30}(2(8a^4 + 34a^3b + 59a^2b^2 + 33ab^3)\cos(x)^5 - 10(4a^4 + 17a^3b + 28a^2b^2 + 15ab^3)\cos(x)^3 + 15(b^3\cos(x)^4 - 2b^3\cos(x)^2 + b^3)\sqrt{a^2 + ab}\arctan(\frac{1}{2}((2a + b)\cos(x)^2 - a)/(\sqrt{a^2 + ab}\cos(x)\sin(x)))\sin(x) + 30(a^4 + 4a^3b + 6a^2b^2 + 3ab^3)\cos(x))/((a^5 + 4a^4b + 6a^3b^2 + 4a^2b^3 + ab^4)\cos(x)^4 - 2(a^5 + 4a^4b + 6a^3b^2 + 4a^2b^3 + ab^4)\cos(x)^2)\sin(x)]$$

Sympy [F]

$$\int \frac{\csc^6(x)}{a + b \cos^2(x)} dx = \int \frac{\csc^6(x)}{a + b \cos^2(x)} dx$$

input `integrate(csc(x)**6/(a+b*cos(x)**2),x)`

output `Integral(csc(x)**6/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.43

$$\begin{aligned} & \int \frac{\csc^6(x)}{a + b \cos^2(x)} dx \\ &= \frac{b^3 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{(a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{(a+b)a}} \\ &\quad - \frac{15(a^2 + 3ab + 3b^2)\tan(x)^4 + 5(2a^2 + 5ab + 3b^2)\tan(x)^2 + 3a^2 + 6ab + 3b^2}{15(a^3 + 3a^2b + 3ab^2 + b^3)\tan(x)^5} \end{aligned}$$

input `integrate(csc(x)^6/(a+b*cos(x)^2),x, algorithm="maxima")`

output `b^3*arctan(a*tan(x)/sqrt((a + b)*a))/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sqrt((a + b)*a)) - 1/15*(15*(a^2 + 3*a*b + 3*b^2)*tan(x)^4 + 5*(2*a^2 + 5*a*b + 3*b^2)*tan(x)^2 + 3*a^2 + 6*a*b + 3*b^2)/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*tan(x)^5)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(77) = 154$.

Time = 0.12 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.75

$$\int \frac{\csc^6(x)}{a + b \cos^2(x)} dx = \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right) b^3}{(a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{a^2 + ab}}$$

$$- \frac{15a^2 \tan(x)^4 + 45ab \tan(x)^4 + 45b^2 \tan(x)^4 + 10a^2 \tan(x)^2 + 25ab \tan(x)^2 + 15b^2 \tan(x)^2 + 3a^2}{15(a^3 + 3a^2b + 3ab^2 + b^3)\tan(x)^5}$$

input `integrate(csc(x)^6/(a+b*cos(x)^2),x, algorithm="giac")`

output $(\pi * \operatorname{floor}(x/\pi + 1/2) * \operatorname{sgn}(a) + \arctan(a * \tan(x) / \sqrt{a^2 + a * b})) * b^3 / ((a^3 + 3 * a^2 * b + 3 * a * b^2 + b^3) * \sqrt{a^2 + a * b}) - 1/15 * (15 * a^2 * \tan(x)^4 + 45 * a * b * \tan(x)^4 + 45 * b^2 * \tan(x)^4 + 10 * a^2 * \tan(x)^2 + 25 * a * b * \tan(x)^2 + 15 * b^2 * \tan(x)^2 + 3 * a^2 + 6 * a * b + 3 * b^2) / ((a^3 + 3 * a^2 * b + 3 * a * b^2 + b^3) * \tan(x)^5)$

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.13

$$\int \frac{\csc^6(x)}{a + b \cos^2(x)} dx = \frac{b^3 \operatorname{atan}\left(\frac{\sqrt{a} \tan(x) (a^3 + 3a^2b + 3ab^2 + b^3)}{(a+b)^{7/2}}\right)}{\sqrt{a} (a+b)^{7/2}}$$

$$- \frac{\frac{1}{5(a+b)} + \frac{\tan(x)^2 (2a+3b)}{3(a+b)^2} + \frac{\tan(x)^4 (a^2+3ab+3b^2)}{(a+b)^3}}{\tan(x)^5}$$

input `int(1/(sin(x)^6*(a + b*cos(x)^2)),x)`

output $(b^3 * \operatorname{atan}((a^{1/2} * \tan(x) * (3 * a * b^2 + 3 * a^2 * b + a^3 + b^3)) / (a + b)^{7/2})) / (a^{1/2} * (a + b)^{7/2}) - (1 / (5 * (a + b))) + (\tan(x)^2 * (2 * a + 3 * b)) / (3 * (a + b)^2) + (\tan(x)^4 * (3 * a * b + a^2 + 3 * b^2)) / (a + b)^3 / \tan(x)^5$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 238, normalized size of antiderivative = 2.67

$$\int \frac{\csc^6(x)}{a + b \cos^2(x)} dx \\ = \frac{15\sqrt{a}\sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) \sin(x)^5 b^3 + 15\sqrt{a}\sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) \sin(x)^5 b^3 - 8 \cos(x)^5 b^3}{(a + b \cos^2(x))^3}$$

input `int(csc(x)^6/(a+b*cos(x)^2),x)`

output `(15*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*sin(x)**5*b**3 + 15*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*sin(x)**5*b**3 - 8*cos(x)*sin(x)**4*a**4 - 34*cos(x)*sin(x)**4*a**3*b - 59*cos(x)*sin(x)**4*a**2*b**2 - 33*cos(x)*sin(x)**4*a*b**3 - 4*cos(x)*sin(x)**2*a**4 - 17*cos(x)*sin(x)**2*a**3*b - 22*cos(x)*sin(x)**2*a**2*b**2 - 9*cos(x)*sin(x)**2*a*b**3 - 3*cos(x)*a**4 - 9*cos(x)*a**3*b - 9*cos(x)*a**2*b**2 - 3*cos(x)*a*b**3)/(15*sin(x)**5*a*(a**4 + 4*a**3*b + 6*a**2*b**2 + 4*a*b**3 + b**4))`

3.24 $\int \frac{\sin(x)}{4-3\cos^3(x)} dx$

Optimal result	196
Mathematica [A] (verified)	196
Rubi [A] (verified)	197
Maple [C] (verified)	200
Fricas [A] (verification not implemented)	201
Sympy [A] (verification not implemented)	201
Maxima [A] (verification not implemented)	202
Giac [A] (verification not implemented)	202
Mupad [B] (verification not implemented)	203
Reduce [B] (verification not implemented)	203

Optimal result

Integrand size = 13, antiderivative size = 84

$$\int \frac{\sin(x)}{4 - 3\cos^3(x)} dx = -\frac{\arctan\left(\frac{1 + \sqrt[3]{6}\cos(x)}{\sqrt{3}}\right)}{2\sqrt[3]{2}3^{5/6}} + \frac{\log(6^{2/3} - 3\cos(x))}{6\sqrt[3]{6}} \\ - \frac{\log(2\sqrt[3]{6} + 6^{2/3}\cos(x) + 3\cos^2(x))}{12\sqrt[3]{6}}$$

output
$$-1/12*\arctan(1/3*(1+6^(1/3)*\cos(x))*3^(1/2))*2^(2/3)*3^(1/6)+1/36*\ln(6^(2/3)-3*\cos(x))*6^(2/3)-1/72*\ln(2*6^(1/3)+6^(2/3)*\cos(x)+3*\cos(x)^2)*6^(2/3)$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.94

$$\int \frac{\sin(x)}{4 - 3\cos^3(x)} dx \\ = \frac{1}{72} \left(-62^{2/3}\sqrt[6]{3} \arctan\left(\frac{1 + \sqrt[3]{6}\cos(x)}{\sqrt{3}}\right) \right. \\ \left. + 6^{2/3} \left(2\log\left(2 - \sqrt[3]{6}\cos(x)\right) - \log\left(4 + 2\sqrt[3]{6}\cos(x) + 6^{2/3}\cos^2(x)\right) \right) \right)$$

input `Integrate[Sin[x]/(4 - 3*Cos[x]^3), x]`

output
$$\frac{(-6 \cdot 2^{(2/3)} \cdot 3^{(1/6)} \cdot \text{ArcTan}[(1 + 6^{(1/3)} \cdot \cos[x])/\sqrt[3]{3}] + 6^{(2/3)} \cdot (2 \cdot \log[2 - 6^{(1/3)} \cdot \cos[x]] - \log[4 + 2 \cdot 6^{(1/3)} \cdot \cos[x] + 6^{(2/3)} \cdot \cos[x]^2]))/72}{\sqrt[3]{2}}$$

Rubi [A] (verified)

Time = 0.35 (sec), antiderivative size = 100, normalized size of antiderivative = 1.19, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.769$, Rules used = {3042, 25, 3702, 750, 16, 1142, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\cos(x + \frac{\pi}{2})}{4 - 3 \sin(x + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\cos(x + \frac{\pi}{2})}{4 - 3 \sin(x + \frac{\pi}{2})^3} dx \\
 & \quad \downarrow \text{3702} \\
 & - \int \frac{1}{4 - 3 \cos^3(x)} d \cos(x) \\
 & \quad \downarrow \text{750} \\
 & - \frac{\int \frac{\sqrt[3]{3 \cos(x) + 2} \cdot 2^{2/3}}{3^{2/3} \cos^2(x) + 2^{2/3} \sqrt[3]{3 \cos(x) + 2} \sqrt[3]{2}} d \cos(x)}{6 \sqrt[3]{2}} - \frac{\int \frac{1}{2^{2/3} - \sqrt[3]{3 \cos(x)}} d \cos(x)}{6 \sqrt[3]{2}} \\
 & \quad \downarrow \text{16} \\
 & \frac{\log(2^{2/3} - \sqrt[3]{3 \cos(x)})}{6 \sqrt[3]{6}} - \frac{\int \frac{\sqrt[3]{3 \cos(x) + 2} \cdot 2^{2/3}}{3^{2/3} \cos^2(x) + 2^{2/3} \sqrt[3]{3 \cos(x) + 2} \sqrt[3]{2}} d \cos(x)}{6 \sqrt[3]{2}}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 1142 \\
& \frac{\log \left(2^{2/3} - \sqrt[3]{3} \cos(x)\right)}{6\sqrt[3]{6}} - \\
& \frac{3 \int \frac{1}{3^{2/3} \cos^2(x) + 2^{2/3}} \frac{1}{\sqrt[3]{3} \cos(x) + 2} \frac{3}{\sqrt[3]{2}} d \cos(x)}{\sqrt[3]{2}} + \frac{\int \frac{2^{2/3} \sqrt[3]{3} \left(\sqrt[3]{6} \cos(x) + 1\right)}{3^{2/3} \cos^2(x) + 2^{2/3}} \frac{3}{\sqrt[3]{3} \cos(x) + 2} \frac{3}{\sqrt[3]{2}} d \cos(x)}{2\sqrt[3]{3}} \\
& \frac{6\sqrt[3]{2}}{6\sqrt[3]{2}} \\
& \downarrow 27 \\
& \frac{\log \left(2^{2/3} - \sqrt[3]{3} \cos(x)\right)}{6\sqrt[3]{6}} - \\
& \frac{3 \int \frac{1}{3^{2/3} \cos^2(x) + 2^{2/3}} \frac{1}{\sqrt[3]{3} \cos(x) + 2} \frac{3}{\sqrt[3]{2}} d \cos(x)}{\sqrt[3]{2}} + \frac{\int \frac{\sqrt[3]{6} \cos(x) + 1}{3^{2/3} \cos^2(x) + 2^{2/3}} \frac{3}{\sqrt[3]{3} \cos(x) + 2} \frac{3}{\sqrt[3]{2}} d \cos(x)}{\sqrt[3]{2}} \\
& \frac{6\sqrt[3]{2}}{6\sqrt[3]{2}} \\
& \downarrow 1082 \\
& \frac{\log \left(2^{2/3} - \sqrt[3]{3} \cos(x)\right)}{6\sqrt[3]{6}} - \\
& \frac{\int \frac{\sqrt[3]{6} \cos(x) + 1}{3^{2/3} \cos^2(x) + 2^{2/3}} \frac{3}{\sqrt[3]{3} \cos(x) + 2} \frac{3}{\sqrt[3]{2}} d \cos(x)}{\sqrt[3]{2}} - 3^{2/3} \int \frac{1}{-\left(\frac{\sqrt[3]{6} \cos(x) + 1}{\sqrt[3]{3}}\right)^2 - 3} d\left(\frac{\sqrt[3]{6} \cos(x) + 1}{\sqrt[3]{3}}\right) \\
& \frac{6\sqrt[3]{2}}{6\sqrt[3]{2}} \\
& \downarrow 217 \\
& \frac{\log \left(2^{2/3} - \sqrt[3]{3} \cos(x)\right)}{6\sqrt[3]{6}} - \frac{\int \frac{\sqrt[3]{6} \cos(x) + 1}{3^{2/3} \cos^2(x) + 2^{2/3}} \frac{3}{\sqrt[3]{3} \cos(x) + 2} \frac{3}{\sqrt[3]{2}} d \cos(x)}{\sqrt[3]{2}} + \sqrt[6]{3} \arctan\left(\frac{\sqrt[3]{6} \cos(x) + 1}{\sqrt[3]{3}}\right) \\
& \frac{6\sqrt[3]{2}}{6\sqrt[3]{2}} \\
& \downarrow 1103 \\
& \frac{\log \left(2^{2/3} - \sqrt[3]{3} \cos(x)\right)}{6\sqrt[3]{6}} - \frac{\sqrt[6]{3} \arctan\left(\frac{\sqrt[3]{6} \cos(x) + 1}{\sqrt[3]{3}}\right) + \frac{\log\left(3^{2/3} \cos^2(x) + 2^{2/3} \sqrt[3]{3} \cos(x) + 2\sqrt[3]{2}\right)}{2\sqrt[3]{3}}}{6\sqrt[3]{2}}
\end{aligned}$$

input `Int [Sin[x]/(4 - 3*Cos[x]^3),x]`

output

$$\text{Log}[2^{(2/3)} - 3^{(1/3)} \cos[x]]/(6*6^{(1/3)}) - (3^{(1/6)} \text{ArcTan}[(1 + 6^{(1/3)} \cos[x])/\sqrt[3]{3}] + \text{Log}[2*2^{(1/3)} + 2^{(2/3)} 3^{(1/3)} \cos[x] + 3^{(2/3)} \cos[x]^2]/(2*3^{(1/3)}))/(6*2^{(1/3)})$$

Defintions of rubi rules used

rule 16

$$\text{Int}[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 25

$$\text{Int}[-(F_x_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 27

$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!Ma} \text{tchQ}[F_x, (b_)*(G_x_) /; \text{FreeQ}[b, x]]$$

rule 217

$$\text{Int}[(a_ + (b_.*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(-(Rt[-a, 2]*Rt[-b, 2])^{(-1)})*\text{ArcTan}[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$$

rule 750

$$\text{Int}[(a_ + (b_.*(x_)^3)^{-1}), x_Symbol] \rightarrow \text{Simp}[1/(3*Rt[a, 3]^2) \quad \text{Int}[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + \text{Simp}[1/(3*Rt[a, 3]^2) \quad \text{Int}[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$$

rule 1082

$$\text{Int}[(a_ + (b_.*(x_) + (c_.*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&& (\text{EqQ}[q^2, 1] \mid\mid \text{!RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}[\{a, b, c\}, x]]$$

rule 1103

$$\text{Int}[(d_ + (e_.*(x_))/((a_.) + (b_.*(x_) + (c_.*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{EqQ}[2*c*d - b*e, 0]$$

rule 1142 $\text{Int}[(d_{_}) + (e_{_})*(x_{_})/((a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2), x_{\text{Symbol}}] \Rightarrow S$
 $\text{imp}[(2*c*d - b*e)/(2*c) \quad \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c)$
 $\text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 3042 $\text{Int}[u_{_}, x_{\text{Symbol}}] \Rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinear}$
 $Q[u, x]$

rule 3702 $\text{Int}[\cos[(e_{_}) + (f_{_})*(x_{_})]^{(m_{_})}*((a_{_}) + (b_{_})*((c_{_})*\sin[(e_{_}) + (f_{_})*(x_{_})])^{(n_{_})})^{(p_{_})}, x_{\text{Symbol}}] \Rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, Si$
 $\text{mp}[ff/f \quad \text{Subst}[\text{Int}[(1 - ff^2*x^2)^{((m - 1)/2)}*(a + b*(c*ff*x)^n)^p, x], x,$
 $\text{Sin}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, c, e, f, n, p\}, x] \&& \text{IntegerQ}[(m -$
 $1)/2] \&& (\text{EqQ}[n, 4] \text{ || } \text{GtQ}[m, 0] \text{ || } \text{IGtQ}[p, 0] \text{ || } \text{IntegersQ}[m, p])$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.42

method	result
risch	$-\frac{i \left(\sum_{R=\text{RootOf}(162-Z^3+i)} R \ln(e^{2ix}+12i R e^{ix}+1) \right)}{2}$
derivativedivides	$\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cos(x)-\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}}}{3}\right)}{36} - \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cos(x)^2+\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \cos(x)}{3}+\frac{4^{\frac{2}{3}} 3^{\frac{1}{3}}}{3}\right)}{72} - \frac{4^{\frac{1}{3}} 3^{\frac{1}{6}} \arctan\left(\frac{\sqrt{3} \left(\frac{4^{\frac{2}{3}} 3^{\frac{1}{3}} \cos(x)}{2}+1\right)}{3}\right)}{12}$
default	$\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cos(x)-\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}}}{3}\right)}{36} - \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cos(x)^2+\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \cos(x)}{3}+\frac{4^{\frac{2}{3}} 3^{\frac{1}{3}}}{3}\right)}{72} - \frac{4^{\frac{1}{3}} 3^{\frac{1}{6}} \arctan\left(\frac{\sqrt{3} \left(\frac{4^{\frac{2}{3}} 3^{\frac{1}{3}} \cos(x)}{2}+1\right)}{3}\right)}{12}$

input $\text{int}(\sin(x)/(4-3*\cos(x)^3), x, \text{method}=\text{_RETURNVERBOSE})$

output $-1/2*I*\sum(_R*\ln(\exp(2*I*x)+12*I*_R*\exp(I*x)+1), _R=\text{RootOf}(162*_Z^3+I))$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.80

$$\int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx = -\frac{1}{6} \cdot 6^{\frac{1}{6}} \sqrt{\frac{1}{2}} \arctan \left(\frac{1}{3} \cdot 6^{\frac{1}{6}} \sqrt{\frac{1}{2}} \left(6^{\frac{2}{3}} \cos(x) + 6^{\frac{1}{3}} \right) \right) \\ - \frac{1}{72} \cdot 6^{\frac{2}{3}} \log \left(-3 \cos(x)^2 - 6^{\frac{2}{3}} \cos(x) - 2 \cdot 6^{\frac{1}{3}} \right) \\ + \frac{1}{36} \cdot 6^{\frac{2}{3}} \log \left(6^{\frac{2}{3}} - 3 \cos(x) \right)$$

input `integrate(sin(x)/(4-3*cos(x)^3),x, algorithm="fricas")`

output
$$-\frac{1}{6} 6^{(1/6)} \sqrt{\frac{1}{2}} \arctan \left(\frac{1}{3} 6^{(1/6)} \sqrt{\frac{1}{2}} (6^{(2/3)} \cos(x) + 6^{(1/3)}) \right) - \frac{1}{72} 6^{(2/3)} \log \left(-3 \cos(x)^2 - 6^{(2/3)} \cos(x) - 2 \cdot 6^{(1/3)} \right) + \frac{1}{36} 6^{(2/3)} \log \left(6^{(2/3)} - 3 \cos(x) \right)$$

Sympy [A] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.01

$$\int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx = \frac{6^{\frac{2}{3}} \log \left(\cos(x) - \frac{6^{\frac{2}{3}}}{3} \right)}{36} \\ - \frac{6^{\frac{2}{3}} \log \left(36 \cos^2(x) + 12 \cdot 6^{\frac{2}{3}} \cos(x) + 24 \cdot \sqrt[3]{6} \right)}{72} \\ - \frac{2^{\frac{2}{3}} \cdot \sqrt[6]{3} \operatorname{atan} \left(\frac{\sqrt[3]{2} \cdot 3^{\frac{5}{6}} \cos(x)}{3} + \frac{\sqrt{3}}{3} \right)}{12}$$

input `integrate(sin(x)/(4-3*cos(x)**3),x)`

output
$$6^{(2/3)} \log(\cos(x)) - \frac{6^{(2/3)}}{3} / 36 - 6^{(2/3)} \log(36 \cos(x)^2 + 12 \cdot 6^{(2/3)} \cos(x) + 24 \cdot 6^{(1/3)}) / 72 - 2^{(2/3)} \cdot 3^{(5/6)} \operatorname{atan} \left(2^{(1/3)} \cdot 3^{(5/6)} \cos(x) / 3 + \sqrt{3} / 3 \right) / 12$$

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.06

$$\int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx = -\frac{1}{72} \cdot 4^{\frac{1}{3}} 3^{\frac{2}{3}} \log \left(3^{\frac{2}{3}} \cos(x)^2 + 4^{\frac{1}{3}} 3^{\frac{1}{3}} \cos(x) + 4^{\frac{2}{3}} \right) \\ + \frac{1}{36} \cdot 4^{\frac{1}{3}} 3^{\frac{2}{3}} \log \left(\frac{1}{3} \cdot 3^{\frac{2}{3}} \left(3^{\frac{1}{3}} \cos(x) - 4^{\frac{1}{3}} \right) \right) - \frac{1}{12} \\ \cdot 4^{\frac{1}{3}} 3^{\frac{1}{6}} \arctan \left(\frac{1}{12} \cdot 4^{\frac{2}{3}} 3^{\frac{1}{6}} \left(2 \cdot 3^{\frac{2}{3}} \cos(x) + 4^{\frac{1}{3}} 3^{\frac{1}{3}} \right) \right)$$

input `integrate(sin(x)/(4-3*cos(x)^3),x, algorithm="maxima")`

output
$$-\frac{1}{72} 4^{(1/3)} 3^{(2/3)} \log(3^{(2/3)} \cos(x)^2 + 4^{(1/3)} 3^{(1/3)} \cos(x) + 4^{(2/3)}) + \frac{1}{36} 4^{(1/3)} 3^{(2/3)} \log(1/3 3^{(2/3)} (3^{(1/3)} \cos(x) - 4^{(1/3)})) - \frac{1}{12} 4^{(1/3)} 3^{(1/6)} \arctan(1/12 4^{(2/3)} 3^{(1/6)} (2 \cdot 3^{(2/3)} \cos(x) + 4^{(1/3)} 3^{(1/3)}))$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.71

$$\int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx = -\frac{1}{12} \sqrt{3} \left(\frac{4}{3} \right)^{\frac{1}{3}} \arctan \left(\frac{1}{4} \sqrt{3} \left(\frac{4}{3} \right)^{\frac{2}{3}} \left(\left(\frac{4}{3} \right)^{\frac{1}{3}} + 2 \cos(x) \right) \right) \\ - \frac{1}{72} \cdot 36^{\frac{1}{3}} \log \left(\cos(x)^2 + \left(\frac{4}{3} \right)^{\frac{1}{3}} \cos(x) + \left(\frac{4}{3} \right)^{\frac{2}{3}} \right) \\ + \frac{1}{12} \left(\frac{4}{3} \right)^{\frac{1}{3}} \log \left(\left(\frac{4}{3} \right)^{\frac{1}{3}} - \cos(x) \right)$$

input `integrate(sin(x)/(4-3*cos(x)^3),x, algorithm="giac")`

output
$$-\frac{1}{12} \sqrt{3} \left(\frac{4}{3} \right)^{(1/3)} \arctan \left(\frac{1}{4} \sqrt{3} \left(\frac{4}{3} \right)^{(2/3)} \left(\left(\frac{4}{3} \right)^{(1/3)} + 2 \cos(x) \right) \right) - \frac{1}{72} 36^{(1/3)} \log \left(\cos(x)^2 + \left(\frac{4}{3} \right)^{(1/3)} \cos(x) + \left(\frac{4}{3} \right)^{(2/3)} \right) + \frac{1}{12} \left(\frac{4}{3} \right)^{(1/3)} \log \left(\left(\frac{4}{3} \right)^{(1/3)} - \cos(x) \right)$$

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.89

$$\int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx = \frac{6^{2/3} \ln \left(\cos(x) - \frac{6^{2/3}}{3} \right)}{36} + \frac{6^{2/3} \ln \left(\cos(x) - \frac{6^{2/3}(-1+\sqrt{3}1i)}{6} \right) (-1+\sqrt{3}1i)}{72} - \frac{6^{2/3} \ln \left(\cos(x) + \frac{6^{2/3}(1+\sqrt{3}1i)}{6} \right) (1+\sqrt{3}1i)}{72}$$

input `int(-sin(x)/(3*cos(x)^3 - 4),x)`

output `(6^(2/3)*log(cos(x) - 6^(2/3)/3))/36 + (6^(2/3)*log(cos(x) - (6^(2/3)*(3^(1/2)*1i - 1))/6)*(3^(1/2)*1i - 1))/72 - (6^(2/3)*log(cos(x) + (6^(2/3)*(3^(1/2)*1i + 1))/6)*(3^(1/2)*1i + 1))/72`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.89

$$\int \frac{\sin(x)}{4 - 3 \cos^3(x)} dx = \frac{2^{\frac{2}{3}} \left(-2\sqrt{3} \operatorname{atan} \left(\frac{(23^{\frac{1}{3}} \cos(x) + 2^{\frac{2}{3}})^{\frac{1}{3}} \sqrt{3}}{6} \right) - \log \left(3^{\frac{2}{3}} \cos(x)^2 + 2^{\frac{2}{3}} 3^{\frac{1}{3}} \cos(x) + 2 2^{\frac{1}{3}} \right) + 2 \log \left(3^{\frac{1}{3}} \cos(x) - 2^{\frac{2}{3}} \right) \right)}{72}$$

input `int(sin(x)/(4-3*cos(x)^3),x)`

output `(2**((2/3)*(- 2*3**((1/3)*3**((1/6)*atan((2*3**((1/3)*cos(x) + 2**((2/3))/(2**((2/3)*3**((1/3)*3**((1/6)))) - log(3**((2/3)*cos(x)**2 + 2**((2/3)*3**((1/3)*cos(x) + 2*2**((1/3)) + 2*log(3**((1/3)*cos(x) - 2**((2/3))))/(24*3**((1/3)))`

3.25 $\int \frac{\cos^7(x)}{a+b\cos^2(x)} dx$

Optimal result	204
Mathematica [A] (verified)	204
Rubi [A] (verified)	205
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	207
Sympy [F(-1)]	207
Maxima [A] (verification not implemented)	208
Giac [A] (verification not implemented)	208
Mupad [B] (verification not implemented)	209
Reduce [B] (verification not implemented)	209

Optimal result

Integrand size = 15, antiderivative size = 78

$$\begin{aligned} \int \frac{\cos^7(x)}{a + b\cos^2(x)} dx = & -\frac{a^3 \operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{b^{7/2}\sqrt{a+b}} + \frac{(a^2 - ab + b^2)\sin(x)}{b^3} \\ & + \frac{(a - 2b)\sin^3(x)}{3b^2} + \frac{\sin^5(x)}{5b} \end{aligned}$$

output
$$-\text{a}^3 \operatorname{arctanh}\left(\text{b}^{(1/2)} \sin(\text{x}) / (\text{a}+\text{b})^{(1/2)}\right) / \text{b}^{(7/2)} / (\text{a}+\text{b})^{(1/2)} + (\text{a}^2 - \text{a} \cdot \text{b} + \text{b}^2)$$

$$\cdot \sin(\text{x}) / \text{b}^3 + 1/3 \cdot (\text{a} - 2 \cdot \text{b}) \cdot \sin(\text{x})^3 / \text{b}^2 + 1/5 \cdot \sin(\text{x})^5 / \text{b}$$

Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.42

$$\begin{aligned} \int \frac{\cos^7(x)}{a + b\cos^2(x)} dx = & \frac{a^3 \left(\log \left(\sqrt{a+b} - \sqrt{b}\sin(x) \right) - \log \left(\sqrt{a+b} + \sqrt{b}\sin(x) \right) \right)}{2b^{7/2}\sqrt{a+b}} \\ & + \frac{(8a^2 - 6ab + 5b^2)\sin(x)}{8b^3} + \frac{(-4a + 5b)\sin(3x)}{48b^2} + \frac{\sin(5x)}{80b} \end{aligned}$$

input `Integrate[Cos[x]^7/(a + b*Cos[x]^2), x]`

output
$$\frac{(a^3(\log[\sqrt{a+b}] - \sqrt{b}\sin[x]) - \log[\sqrt{a+b} + \sqrt{b}\sin[x]])}{(2b^{7/2})\sqrt{a+b}} + \frac{(8a^2 - 6ab + 5b^2)\sin[x]}{(8b^3)} + \frac{(-4a + 5b)\sin[3x]}{(48b^2)} + \frac{\sin[5x]}{(80b)}$$

Rubi [A] (verified)

Time = 0.30 (sec), antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.267, Rules used = {3042, 3665, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^7(x)}{a + b\cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})^7}{a + b\sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \textcolor{blue}{3665} \\
 & \int \frac{(1 - \sin^2(x))^3}{a - b\sin^2(x) + b} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{300} \\
 & \int \left(-\frac{a^3}{b^3(a - b\sin^2(x) + b)} + \frac{a^2 - ab + b^2}{b^3} + \frac{(a - 2b)\sin^2(x)}{b^2} + \frac{\sin^4(x)}{b} \right) d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{a^3 \operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{b^{7/2}\sqrt{a+b}} + \frac{(a^2 - ab + b^2)\sin(x)}{b^3} + \frac{(a - 2b)\sin^3(x)}{3b^2} + \frac{\sin^5(x)}{5b}
 \end{aligned}$$

input
$$\operatorname{Int}[\cos[x]^7/(a + b\cos[x]^2), x]$$

output

$$-((a^3 \operatorname{ArcTanh}[(\sqrt{b} \sin(x))/\sqrt{a+b}])/(b^{7/2} \sqrt{a+b})) + ((a^2 - a b + b^2) \sin(x))/b^3 + ((a - 2 b) \sin(x)^3)/(3 b^2) + \sin(x)^5/(5 b)$$

Definitions of rubi rules used

rule 300

$$\operatorname{Int}[((a_) + (b_*)*(x_)^2)^(p_*)*((c_) + (d_*)*(x_)^2)^(q_), x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[(a + b*x^2)^p, (c + d*x^2)^{-q}, x], x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \& \operatorname{NeQ}[b*c - a*d, 0] \& \operatorname{IGtQ}[p, 0] \& \operatorname{ILtQ}[q, 0] \& \operatorname{GeQ}[p, -q]$$

rule 2009

$$\operatorname{Int}[u_, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 3042

$$\operatorname{Int}[u_, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3665

$$\operatorname{Int}[\sin[(e_*) + (f_*)*(x_)]^{(m_*)}*((a_) + (b_*)*\sin[(e_*) + (f_*)*(x_)]^2)^{(p_*)}, x_{\text{Symbol}}] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Cos}[e + f*x], x]\}, \operatorname{Simp}[-ff/f \operatorname{Subst}[\operatorname{Int}[(1 - ff^2*x^2)^{(m - 1)/2}*(a + b - b*ff^2*x^2)^p, x], x, \operatorname{Cos}[e + f*x]/ff], x]] /; \operatorname{FreeQ}[\{a, b, e, f, p\}, x] \& \operatorname{IntegerQ}[(m - 1)/2]$$

Maple [A] (verified)

Time = 1.72 (sec), antiderivative size = 78, normalized size of antiderivative = 1.00

method	result
default	$\frac{\sin(x)^5 b^2}{5} + \frac{\sin(x)^3 a b}{3} - \frac{2 b^2 \sin(x)^3}{3} + \sin(x) a^2 - \sin(x) a b + \sin(x) b^2 - \frac{a^3 \operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{b^3 \sqrt{(a+b)b}}$
risch	$-\frac{i e^{i x} a^2}{2 b^3} + \frac{3 i e^{i x} a}{8 b^2} - \frac{5 i e^{i x}}{16 b} + \frac{i e^{-i x} a^2}{2 b^3} - \frac{3 i e^{-i x} a}{8 b^2} + \frac{5 i e^{-i x}}{16 b} + \frac{a^3 \ln\left(e^{2 i x} - \frac{2 i (a+b) e^{i x}}{\sqrt{ab+b^2}} - 1\right)}{2 \sqrt{ab+b^2} b^3} - \frac{a^3 \ln\left(e^{2 i x} + \frac{2 i (a+b) e^{i x}}{\sqrt{ab+b^2}} - 1\right)}{2 \sqrt{ab+b^2} b^3}$

input

$$\operatorname{int}(\cos(x)^7/(a+b*\cos(x)^2), x, \operatorname{method}=\operatorname{_RETURNVERBOSE})$$

output
$$\frac{1/b^3*(1/5*\sin(x)^5*b^2+1/3*\sin(x)^3*a*b-2/3*b^2*\sin(x)^3+\sin(x)*a^2-\sin(x)*a*b+\sin(x)*b^2)-a^3/b^3/((a+b)*b)^(1/2)*\operatorname{arctanh}(b*\sin(x)/((a+b)*b)^(1/2))}{})$$

Fricas [A] (verification not implemented)

Time = 0.20 (sec), antiderivative size = 259, normalized size of antiderivative = 3.32

$$\int \frac{\cos^7(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{15 \sqrt{ab + b^2} a^3 \log \left(-\frac{b \cos(x)^2 + 2 \sqrt{ab + b^2} \sin(x) - a - 2b}{b \cos(x)^2 + a} \right) + 2 (3 (ab^3 + b^4) \cos(x)^4 + 15 a^3 b + 5 a^2 b^2 - 2 ab^3 + 8 a^4) \sin(x)}{30 (ab^4 + b^5)} \right]$$

input `integrate(cos(x)^7/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[1/30*(15*sqrt(a*b + b^2)*a^3*log(-(b*cos(x)^2 + 2*sqrt(a*b + b^2))*sin(x) - a - 2*b)/(b*cos(x)^2 + a)) + 2*(3*(a*b^3 + b^4)*cos(x)^4 + 15*a^3*b + 5*a^2*b^2 - 2*a*b^3 + 8*b^4 - (5*a^2*b^2 + a*b^3 - 4*b^4)*cos(x)^2)*sin(x)/(a*b^4 + b^5), 1/15*(15*sqrt(-a*b - b^2)*a^3*arctan(sqrt(-a*b - b^2)*sin(x))/(a + b)) + (3*(a*b^3 + b^4)*cos(x)^4 + 15*a^3*b + 5*a^2*b^2 - 2*a*b^3 + 8*b^4 - (5*a^2*b^2 + a*b^3 - 4*b^4)*cos(x)^2)*sin(x)/(a*b^4 + b^5)]$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^7(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(cos(x)**7/(a+b*cos(x)**2),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.17

$$\int \frac{\cos^7(x)}{a + b \cos^2(x)} dx = \frac{a^3 \log \left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}} \right)}{2 \sqrt{(a+b)b} b^3} + \frac{3 b^2 \sin(x)^5 + 5 (ab - 2b^2) \sin(x)^3 + 15 (a^2 - ab + b^2) \sin(x)}{15 b^3}$$

input `integrate(cos(x)^7/(a+b*cos(x)^2),x, algorithm="maxima")`

output `1/2*a^3*log((b*sin(x) - sqrt((a + b)*b))/(b*sin(x) + sqrt((a + b)*b)))/(sqrt((a + b)*b)*b^3) + 1/15*(3*b^2*sin(x)^5 + 5*(a*b - 2*b^2)*sin(x)^3 + 15*(a^2 - a*b + b^2)*sin(x))/b^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.23

$$\int \frac{\cos^7(x)}{a + b \cos^2(x)} dx = \frac{a^3 \arctan \left(\frac{b \sin(x)}{\sqrt{-ab - b^2}} \right)}{\sqrt{-ab - b^2} b^3} + \frac{3 b^4 \sin(x)^5 + 5 ab^3 \sin(x)^3 - 10 b^4 \sin(x)^3 + 15 a^2 b^2 \sin(x) - 15 ab^3 \sin(x) + 15 b^4 \sin(x)}{15 b^5}$$

input `integrate(cos(x)^7/(a+b*cos(x)^2),x, algorithm="giac")`

output `a^3*arctan(b*sin(x)/sqrt(-a*b - b^2))/(sqrt(-a*b - b^2)*b^3) + 1/15*(3*b^4 *sin(x)^5 + 5*a*b^3*sin(x)^3 - 10*b^4*sin(x)^3 + 15*a^2*b^2*sin(x) - 15*a*b^3*sin(x) + 15*b^4*sin(x))/b^5`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.10

$$\int \frac{\cos^7(x)}{a + b \cos^2(x)} dx = \frac{\sin(x)^5}{5b} + \sin(x)^3 \left(\frac{a+b}{3b^2} - \frac{1}{b} \right) \\ + \sin(x) \left(\frac{3}{b} + \frac{(a+b) \left(\frac{a+b}{b^2} - \frac{3}{b} \right)}{b} \right) + \frac{a^3 \operatorname{atan}\left(\frac{\sqrt{b} \sin(x) 1i}{\sqrt{a+b}}\right) 1i}{b^{7/2} \sqrt{a+b}}$$

input `int(cos(x)^7/(a + b*cos(x)^2),x)`

output $\sin(x)^5/(5*b) + \sin(x)^3*((a + b)/(3*b^2) - 1/b) + \sin(x)*(3/b + ((a + b)*((a + b)/b^2 - 3/b))/b) + (a^3*\operatorname{atan}((b^(1/2)*\sin(x)*1i)/(a + b)^(1/2))*1i)/(b^(7/2)*(a + b)^(1/2))$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.90

$$\int \frac{\cos^7(x)}{a + b \cos^2(x)} dx \\ = \frac{15\sqrt{b}\sqrt{a+b}\log\left(\sqrt{a+b}\tan\left(\frac{x}{2}\right)^2 + \sqrt{a+b} - 2\sqrt{b}\tan\left(\frac{x}{2}\right)\right)a^3 - 15\sqrt{b}\sqrt{a+b}\log\left(\sqrt{a+b}\tan\left(\frac{x}{2}\right)^2 + \sqrt{a+b} - 2\sqrt{b}\tan\left(\frac{x}{2}\right)\right)b^3}{a^3(b^2 + 2ab + 3b^2)\sqrt{a+b}}$$

input `int(cos(x)^7/(a+b*cos(x)^2),x)`

output $(15*\sqrt{b}*\sqrt{a+b}*\log(\sqrt{a+b}*\tan(x/2)^2 + \sqrt{a+b}) - 2*\sqrt{b}*\tan(x/2))*a^**3 - 15*\sqrt{b}*\sqrt{a+b}*\log(\sqrt{a+b}*\tan(x/2)^2 + \sqrt{a+b}) + 2*\sqrt{b}*\tan(x/2))*a^**3 + 6*\sin(x)^**5*a*b**3 + 6*\sin(x)^**5*b**4 + 10*\sin(x)^**3*a**2*b**2 - 10*\sin(x)^**3*a*b**3 - 20*\sin(x)^**3*b**4 + 30*\sin(x)*a**3*b + 30*\sin(x)*b**4)/(30*b**4*(a + b))$

3.26 $\int \frac{\cos^5(x)}{a+b\cos^2(x)} dx$

Optimal result	210
Mathematica [A] (verified)	210
Rubi [A] (verified)	211
Maple [A] (verified)	212
Fricas [A] (verification not implemented)	213
Sympy [F(-1)]	213
Maxima [A] (verification not implemented)	214
Giac [A] (verification not implemented)	214
Mupad [B] (verification not implemented)	214
Reduce [B] (verification not implemented)	215

Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \frac{\cos^5(x)}{a+b\cos^2(x)} dx = \frac{a^2 \operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{5/2} \sqrt{a+b}} - \frac{(a-b) \sin(x)}{b^2} - \frac{\sin^3(x)}{3b}$$

output $a^2 \operatorname{arctanh}(b^{1/2} \sin(x)/(a+b)^{1/2})/b^{5/2}/(a+b)^{1/2} - (a-b) \sin(x)/b^{2-1/3} \sin(x)^3/b$

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.54

$$\begin{aligned} & \int \frac{\cos^5(x)}{a+b\cos^2(x)} dx \\ &= \frac{\frac{6a^2(-\log(\sqrt{a+b}-\sqrt{b}\sin(x))+\log(\sqrt{a+b}+\sqrt{b}\sin(x)))}{\sqrt{a+b}} + 3\sqrt{b}(-4a+3b)\sin(x) + b^{3/2}\sin(3x)}{12b^{5/2}} \end{aligned}$$

input `Integrate[Cos[x]^5/(a + b*Cos[x]^2), x]`

output $((6*a^2*(-Log[Sqrt[a + b] - Sqrt[b]*Sin[x]] + Log[Sqrt[a + b] + Sqrt[b]*Sin[x]]))/Sqrt[a + b] + 3*Sqrt[b]*(-4*a + 3*b)*Sin[x] + b^{(3/2)}*Sin[3*x])/(12*b^{(5/2)})$

Rubi [A] (verified)

Time = 0.27 (sec), antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3665, 300, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^5(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\sin\left(x + \frac{\pi}{2}\right)^5}{a + b \sin\left(x + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \textcolor{blue}{3665} \\
 & \int \frac{(1 - \sin^2(x))^2}{a - b \sin^2(x) + b} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{300} \\
 & \int \left(\frac{a^2}{b^2 (a - b \sin^2(x) + b)} - \frac{a - b}{b^2} - \frac{\sin^2(x)}{b} \right) d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{a^2 \operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{5/2} \sqrt{a+b}} - \frac{(a-b) \sin(x)}{b^2} - \frac{\sin^3(x)}{3b}
 \end{aligned}$$

input $\operatorname{Int}[\operatorname{Cos}[x]^5/(a + b*\operatorname{Cos}[x]^2), x]$

output $(a^2 \operatorname{ArcTanh}\left[\frac{(\operatorname{Sqrt}[b]*\operatorname{Sin}[x])/Sqrt[a + b]}{Sqrt[a + b]}\right])/(b^{(5/2)}*Sqrt[a + b]) - ((a - b)*\operatorname{Sin}[x])/b^2 - \operatorname{Sin}[x]^3/(3*b)$

Definitions of rubi rules used

rule 300 $\text{Int}[(a_.) + (b_.)*(x_)^2]^{(p_.)} * ((c_.) + (d_.)*(x_)^2)]^{(q_.)}, \text{x_Symbol}] \rightarrow \text{Int}[\text{PolynomialDivide}[(a + b*x^2)^p, (c + d*x^2)^{-q}], x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{IGtQ}[p, 0] \&& \text{ILtQ}[q, 0] \&& \text{GeQ}[p, -q]$

rule 2009 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3665 $\text{Int}[\sin[(e_.) + (f_.)*(x_)]^{(m_.)} * ((a_.) + (b_.) * \sin[(e_.) + (f_.)*(x_)]^2)]^{(p_.)}, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Cos}[e + f*x], x]\}, \text{Simp}[-ff/f \text{Subst}[\text{Int}[(1 - ff^2*x^2)^{((m - 1)/2)} * (a + b - b*ff^2*x^2)^p, x], x, \text{Cos}[e + f*x]/ff], x]] /; \text{FreeQ}\{a, b, e, f, p\}, x] \&& \text{IntegerQ}[(m - 1)/2]$

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.89

method	result	size
default	$-\frac{\frac{b \sin(x)^3}{3} + \sin(x)a - b \sin(x)}{b^2} + \frac{a^2 \operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{b^2 \sqrt{(a+b)b}}$	50
risch	$\frac{i e^{i x} a}{2 b^2} - \frac{3 i e^{i x}}{8 b} - \frac{i e^{-i x} a}{2 b^2} + \frac{3 i e^{-i x}}{8 b} + \frac{a^2 \ln\left(e^{2 i x} + \frac{2 i (a+b) e^{i x}}{\sqrt{ab+b^2}} - 1\right)}{2 \sqrt{ab+b^2} b^2} - \frac{a^2 \ln\left(e^{2 i x} - \frac{2 i (a+b) e^{i x}}{\sqrt{ab+b^2}} - 1\right)}{2 \sqrt{ab+b^2} b^2} + \frac{\sin(3x)}{12b}$	147

input $\text{int}(\cos(x)^5 / (a+b*\cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output $-1/b^2 * (1/3*b*\sin(x)^3 + \sin(x)*a - b*\sin(x)) + a^2/b^2 / ((a+b)*b)^{(1/2)} * \operatorname{arctanh}(b*\sin(x) / ((a+b)*b)^{(1/2)})$

Fricas [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 191, normalized size of antiderivative = 3.41

$$\int \frac{\cos^5(x)}{a + b \cos^2(x)} dx$$

$$= \left[\frac{3 \sqrt{ab + b^2} a^2 \log \left(-\frac{b \cos(x)^2 - 2 \sqrt{ab + b^2} \sin(x) - a - 2b}{b \cos(x)^2 + a} \right) - 2 (3 a^2 b + ab^2 - 2 b^3 - (ab^2 + b^3) \cos(x)^2) \sin(x)}{6 (ab^3 + b^4)}, \right.$$

$$\left. - \frac{3 \sqrt{-ab - b^2} a^2 \arctan \left(\frac{\sqrt{-ab - b^2} \sin(x)}{a + b} \right) + (3 a^2 b + ab^2 - 2 b^3 - (ab^2 + b^3) \cos(x)^2) \sin(x)}{3 (ab^3 + b^4)} \right]$$

input `integrate(cos(x)^5/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[1/6*(3*sqrt(a*b + b^2)*a^2*log(-(b*cos(x)^2 - 2*sqrt(a*b + b^2)*sin(x) - a - 2*b)/(b*cos(x)^2 + a)) - 2*(3*a^2*b + a*b^2 - 2*b^3 - (a*b^2 + b^3)*cos(x)^2)*sin(x)/(a*b^3 + b^4), -1/3*(3*sqrt(-a*b - b^2)*a^2*arctan(sqrt(-a*b - b^2)*sin(x)/(a + b)) + (3*a^2*b + a*b^2 - 2*b^3 - (a*b^2 + b^3)*cos(x)^2)*sin(x))/(a*b^3 + b^4)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^5(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(cos(x)**5/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.20

$$\int \frac{\cos^5(x)}{a + b \cos^2(x)} dx = -\frac{a^2 \log \left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}} \right)}{2 \sqrt{(a+b)b} b^2} - \frac{b \sin(x)^3 + 3(a-b) \sin(x)}{3 b^2}$$

input `integrate(cos(x)^5/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-1/2*a^2*log((b*sin(x) - sqrt((a + b)*b))/(b*sin(x) + sqrt((a + b)*b)))/(sqrt((a + b)*b)*b^2) - 1/3*(b*sin(x)^3 + 3*(a - b)*sin(x))/b^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.16

$$\int \frac{\cos^5(x)}{a + b \cos^2(x)} dx = -\frac{a^2 \arctan \left(\frac{b \sin(x)}{\sqrt{-ab - b^2}} \right)}{\sqrt{-ab - b^2} b^2} - \frac{b^2 \sin(x)^3 + 3ab \sin(x) - 3b^2 \sin(x)}{3b^3}$$

input `integrate(cos(x)^5/(a+b*cos(x)^2),x, algorithm="giac")`

output `-a^2*arctan(b*sin(x)/sqrt(-a*b - b^2))/(sqrt(-a*b - b^2)*b^2) - 1/3*(b^2*sin(x)^3 + 3*a*b*sin(x) - 3*b^2*sin(x))/b^3`

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int \frac{\cos^5(x)}{a + b \cos^2(x)} dx = \frac{a^2 \operatorname{atanh} \left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}} \right)}{b^{5/2} \sqrt{a+b}} - \frac{\sin(x)^3}{3b} - \sin(x) \left(\frac{a+b}{b^2} - \frac{2}{b} \right)$$

input `int(cos(x)^5/(a + b*cos(x)^2),x)`

output
$$(a^2 \operatorname{atanh}((b^{1/2} \sin(x))/(a + b)^{1/2})) / (b^{5/2} (a + b)^{1/2}) - \sin(x)^3 / (3*b) - \sin(x) * ((a + b)/b^2 - 2/b)$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 117, normalized size of antiderivative = 2.09

$$\int \frac{\cos^5(x)}{a + b \cos^2(x)} dx \\ = \frac{-3\sqrt{b} \sqrt{a+b} \log\left(\sqrt{a+b} \tan\left(\frac{x}{2}\right)^2 + \sqrt{a+b} - 2\sqrt{b} \tan\left(\frac{x}{2}\right)\right) a^2 + 3\sqrt{b} \sqrt{a+b} \log\left(\sqrt{a+b} \tan\left(\frac{x}{2}\right)^2 + \sqrt{a+b} - 2\sqrt{b} \tan\left(\frac{x}{2}\right)\right) b^2 + 6b^3 (a+b)}{6b^3 (a+b)}$$

input `int(cos(x)^5/(a+b*cos(x)^2),x)`

output
$$(-3*\sqrt(b)*\sqrt(a+b)*\log(\sqrt(a+b)*\tan(x/2)^2 + \sqrt(a+b) - 2*\sqrt(b)*\tan(x/2))*a^2 + 3*\sqrt(b)*\sqrt(a+b)*\log(\sqrt(a+b)*\tan(x/2)^2 + \sqrt(a+b) + 2*\sqrt(b)*\tan(x/2))*a^2 - 2*\sin(x)^3*a*b^2 - 2*\sin(x)^3*b^3 - 6*\sin(x)*a^2*b + 6*\sin(x)*b^3)/(6*b^3*(a+b))$$

3.27 $\int \frac{\cos^3(x)}{a+b\cos^2(x)} dx$

Optimal result	216
Mathematica [A] (verified)	216
Rubi [A] (verified)	217
Maple [A] (verified)	218
Fricas [A] (verification not implemented)	219
Sympy [F(-1)]	219
Maxima [A] (verification not implemented)	220
Giac [A] (verification not implemented)	220
Mupad [B] (verification not implemented)	220
Reduce [B] (verification not implemented)	221

Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \frac{\cos^3(x)}{a + b\cos^2(x)} dx = -\frac{a \operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}} + \frac{\sin(x)}{b}$$

output $-\frac{a \operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}} + \frac{\sin(x)}{b}$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{\cos^3(x)}{a + b\cos^2(x)} dx = -\frac{a \operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}} + \frac{\sin(x)}{b}$$

input $\text{Integrate}[\text{Cos}[x]^3/(a + b*\text{Cos}[x]^2), x]$

output $-\frac{a \operatorname{ArcTanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}} + \frac{\sin(x)}{b}$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3665, 299, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^3(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})^3}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \textcolor{blue}{3665} \\
 & \int \frac{1 - \sin^2(x)}{a - b \sin^2(x) + b} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{299} \\
 & \frac{\sin(x)}{b} - \frac{a \int \frac{1}{-b \sin^2(x) + a + b} d\sin(x)}{b} \\
 & \quad \downarrow \textcolor{blue}{221} \\
 & \frac{\sin(x)}{b} - \frac{a \operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}}
 \end{aligned}$$

input `Int[Cos[x]^3/(a + b*Cos[x]^2), x]`

output
$$-\frac{(a \operatorname{ArcTanh}\left[\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right])/(b^{(3/2)} \sqrt{a+b}) + \sin(x)}{b}$$

Definitions of rubi rules used

rule 221 $\text{Int}[(a_ + b_)*(x_)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{NegQ}[a/b]$

rule 299 $\text{Int}[(a_ + b_)*(x_)^2)^{(p_)*((c_ + d_)*(x_)^2)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - \text{Simp}[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) \text{Int}[(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[2*p + 3, 0]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3665 $\text{Int}[\sin[(e_ + f_)*(x_)]^{(m_)*((a_ + b_)*\sin[(e_ + f_)*(x_)]^2)^{(p_)}}, x_{\text{Symbol}}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Cos}[e + f*x], x]\}, \text{Simp}[-ff/f \text{Subst}[\text{Int}[(1 - ff^2*x^2)^{((m - 1)/2)*(a + b - b*ff^2*x^2)^p}, x], x, \text{Cos}[e + f*x]/ff], x] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{IntegerQ}[(m - 1)/2]$

Maple [A] (verified)

Time = 0.27 (sec), antiderivative size = 33, normalized size of antiderivative = 0.87

method	result	size
default	$\frac{\sin(x)}{b} - \frac{a \operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{b \sqrt{(a+b)b}}$	33
risch	$-\frac{ie^{ix}}{2b} + \frac{ie^{-ix}}{2b} + \frac{a \ln\left(e^{2ix} - \frac{2i(a+b)e^{ix}}{\sqrt{ab+b^2}} - 1\right)}{2\sqrt{ab+b^2}b} - \frac{a \ln\left(e^{2ix} + \frac{2i(a+b)e^{ix}}{\sqrt{ab+b^2}} - 1\right)}{2\sqrt{ab+b^2}b}$	110

input `int(cos(x)^3/(a+b*cos(x)^2), x, method=_RETURNVERBOSE)`

output `sin(x)/b - 1/b*a/((a+b)*b)^(1/2)*arctanh(b*sin(x)/((a+b)*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 134, normalized size of antiderivative = 3.53

$$\int \frac{\cos^3(x)}{a + b \cos^2(x)} dx$$

$$= \left[\frac{\sqrt{ab + b^2} a \log \left(-\frac{b \cos(x)^2 + 2 \sqrt{ab + b^2} \sin(x) - a - 2b}{b \cos(x)^2 + a} \right) + 2(ab + b^2) \sin(x)}{2(ab^2 + b^3)}, \frac{\sqrt{-ab - b^2} a \arctan \left(\frac{\sqrt{-ab - b^2} \sin(x)}{a + b} \right)}{ab^2 + b^3} \right]$$

input `integrate(cos(x)^3/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[1/2*(sqrt(a*b + b^2)*a*log(-(b*cos(x)^2 + 2*sqrt(a*b + b^2)*sin(x) - a - 2*b)/(b*cos(x)^2 + a)) + 2*(a*b + b^2)*sin(x))/(a*b^2 + b^3), (sqrt(-a*b - b^2)*a*arctan(sqrt(-a*b - b^2)*sin(x)/(a + b)) + (a*b + b^2)*sin(x))/(a*b^2 + b^3)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(cos(x)**3/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.32

$$\int \frac{\cos^3(x)}{a + b \cos^2(x)} dx = \frac{a \log \left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}} \right)}{2 \sqrt{(a+b)b}} + \frac{\sin(x)}{b}$$

input `integrate(cos(x)^3/(a+b*cos(x)^2),x, algorithm="maxima")`

output `1/2*a*log((b*sin(x) - sqrt((a + b)*b))/(b*sin(x) + sqrt((a + b)*b)))/(sqrt((a + b)*b)*b) + sin(x)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

$$\int \frac{\cos^3(x)}{a + b \cos^2(x)} dx = \frac{a \arctan \left(\frac{b \sin(x)}{\sqrt{-ab - b^2}} \right)}{\sqrt{-ab - b^2}} + \frac{\sin(x)}{b}$$

input `integrate(cos(x)^3/(a+b*cos(x)^2),x, algorithm="giac")`

output `a*arctan(b*sin(x)/sqrt(-a*b - b^2))/(sqrt(-a*b - b^2)*b) + sin(x)/b`

Mupad [B] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{\cos^3(x)}{a + b \cos^2(x)} dx = \frac{\sin(x)}{b} - \frac{a \operatorname{atanh} \left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}} \right)}{b^{3/2} \sqrt{a+b}}$$

input `int(cos(x)^3/(a + b*cos(x)^2),x)`

output `sin(x)/b - (a*atanh((b^(1/2)*sin(x))/(a + b)^(1/2)))/(b^(3/2)*(a + b)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.39

$$\int \frac{\cos^3(x)}{a + b \cos^2(x)} dx \\ = \frac{\sqrt{b} \sqrt{a + b} \log\left(\sqrt{a + b} \tan\left(\frac{x}{2}\right)^2 + \sqrt{a + b} - 2\sqrt{b} \tan\left(\frac{x}{2}\right)\right) a - \sqrt{b} \sqrt{a + b} \log\left(\sqrt{a + b} \tan\left(\frac{x}{2}\right)^2 + \sqrt{a + b}\right) b^2}{2b^2(a + b)}$$

input `int(cos(x)^3/(a+b*cos(x)^2),x)`

output `(sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)*tan(x/2))*a - sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan(x/2))*a + 2*sin(x)*a*b + 2*sin(x)*b**2)/(2*b**2*(a + b))`

3.28 $\int \frac{\cos(x)}{a+b\cos^2(x)} dx$

Optimal result	222
Mathematica [A] (verified)	222
Rubi [A] (verified)	223
Maple [A] (verified)	224
Fricas [B] (verification not implemented)	224
Sympy [B] (verification not implemented)	225
Maxima [A] (verification not implemented)	226
Giac [A] (verification not implemented)	226
Mupad [B] (verification not implemented)	226
Reduce [B] (verification not implemented)	227

Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{\sqrt{b} \sqrt{a+b}}$$

output arctanh(b^(1/2)*sin(x)/(a+b)^(1/2))/b^(1/2)/(a+b)^(1/2)

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{\sqrt{b} \sqrt{a+b}}$$

input Integrate[Cos[x]/(a + b*Cos[x]^2), x]

output ArcTanh[(Sqrt[b]*Sin[x])/Sqrt[a + b]]/(Sqrt[b]*Sqrt[a + b])

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 3665, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})}{a + b \sin^2(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3665} \\
 & \int \frac{1}{a - b \sin^2(x) + b} d\sin(x) \\
 & \quad \downarrow \text{221} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{\sqrt{b} \sqrt{a+b}}
 \end{aligned}$$

input `Int[Cos[x]/(a + b*Cos[x]^2), x]`

output `ArcTanh[(Sqrt[b]*Sin[x])/Sqrt[a + b]]/(Sqrt[b]*Sqrt[a + b])`

Definitions of rubi rules used

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3665

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^p_, x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Simplify[-ff/f]
Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Maple [A] (verified)

Time = 0.10 (sec), antiderivative size = 21, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{\operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{\sqrt{(a+b)b}}$	21
risch	$\frac{\ln\left(\frac{e^{2ix}+2i(a+b)e^{ix}}{\sqrt{ab+b^2}}-1\right)}{2\sqrt{ab+b^2}} - \frac{\ln\left(\frac{e^{2ix}-2i(a+b)e^{ix}}{\sqrt{ab+b^2}}-1\right)}{2\sqrt{ab+b^2}}$	80

input `int(cos(x)/(a+b*cos(x)^2), x, method=_RETURNVERBOSE)`output `1/((a+b)*b)^(1/2)*arctanh(b*sin(x)/((a+b)*b)^(1/2))`**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(21) = 42$.

Time = 0.13 (sec), antiderivative size = 95, normalized size of antiderivative = 3.28

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = \left[\frac{\log\left(\frac{-b \cos(x)^2 - 2\sqrt{ab + b^2} \sin(x) - a - 2b}{b \cos(x)^2 + a}\right)}{2\sqrt{ab + b^2}}, \right. \\ \left. - \frac{\sqrt{-ab - b^2} \arctan\left(\frac{\sqrt{-ab - b^2} \sin(x)}{a + b}\right)}{ab + b^2} \right]$$

input `integrate(cos(x)/(a+b*cos(x)^2), x, algorithm="fricas")`

output [1/2*log(-(b*cos(x)^2 - 2*sqrt(a*b + b^2)*sin(x) - a - 2*b)/(b*cos(x)^2 + a))/sqrt(a*b + b^2), -sqrt(-a*b - b^2)*arctan(sqrt(-a*b - b^2)*sin(x)/(a + b))/(a*b + b^2)]

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55508 vs. 2(27) = 54.

Time = 66.63 (sec) , antiderivative size = 55508, normalized size of antiderivative = 1914.07

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input integrate(cos(x)/(a+b*cos(x)**2),x)

output Piecewise((zoo*(-log(tan(x/2) - 1) + log(tan(x/2) + 1)), Eq(a, 0) & Eq(b, 0)), (tan(x/2)/(2*b) + 1/(2*b*tan(x/2)), Eq(a, -b)), (sin(x)/a, Eq(b, 0)), (-13*a**6*b*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b)))*log(-sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan(x/2))/(2*a**7*b*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b)))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a**6*b**2*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 24*a**6*b*sqrt(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**5*b**3*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 416*a**5*b**2*sqrt(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 858*a**4*b**4*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 1144*a**4*b**3*sqrt(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 858*a**3*b**5*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**2*b**6*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 1144*a**2*b**5*sqr t(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 858*a**2*b**4*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 416*a**2*b**3*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 858*a**2*b**2*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**2*b**1*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 1144*a**2*b**0*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)))

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.34

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = -\frac{\log\left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}}\right)}{2 \sqrt{(a+b)b}}$$

input `integrate(cos(x)/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-1/2*log((b*sin(x) - sqrt((a + b)*b))/(b*sin(x) + sqrt((a + b)*b)))/sqrt((a + b)*b)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = -\frac{\arctan\left(\frac{b \sin(x)}{\sqrt{-ab-b^2}}\right)}{\sqrt{-ab-b^2}}$$

input `integrate(cos(x)/(a+b*cos(x)^2),x, algorithm="giac")`

output `-arctan(b*sin(x)/sqrt(-a*b - b^2))/sqrt(-a*b - b^2)`

Mupad [B] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{\sqrt{b} \sqrt{a+b}}$$

input `int(cos(x)/(a + b*cos(x)^2),x)`

output `atanh((b^(1/2)*sin(x))/(a + b)^(1/2))/(b^(1/2)*(a + b)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.38

$$\int \frac{\cos(x)}{a + b \cos^2(x)} dx \\ = \frac{\sqrt{b} \sqrt{a+b} \left(-\log \left(\sqrt{a+b} \tan \left(\frac{x}{2} \right)^2 + \sqrt{a+b} - 2\sqrt{b} \tan \left(\frac{x}{2} \right) \right) + \log \left(\sqrt{a+b} \tan \left(\frac{x}{2} \right)^2 + \sqrt{a+b} + 2\sqrt{b} \tan \left(\frac{x}{2} \right) \right) \right)}{2b(a+b)}$$

input `int(cos(x)/(a+b*cos(x)^2),x)`

output `(sqrt(b)*sqrt(a + b)*(- log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqr
t(b)*tan(x/2)) + log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan
(x/2)))/(2*b*(a + b))`

3.29 $\int \frac{\sec(x)}{a+b\cos^2(x)} dx$

Optimal result	228
Mathematica [A] (verified)	228
Rubi [A] (verified)	229
Maple [A] (verified)	230
Fricas [A] (verification not implemented)	231
Sympy [F]	231
Maxima [A] (verification not implemented)	232
Giac [A] (verification not implemented)	232
Mupad [B] (verification not implemented)	233
Reduce [B] (verification not implemented)	233

Optimal result

Integrand size = 13, antiderivative size = 41

$$\int \frac{\sec(x)}{a+b\cos^2(x)} dx = \frac{\operatorname{arctanh}(\sin(x))}{a} - \frac{\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{a\sqrt{a+b}}$$

output $\operatorname{arctanh}(\sin(x))/a - b^{1/2} \operatorname{arctanh}(b^{1/2} \sin(x)/(a+b)^{1/2})/a/(a+b)^{1/2}$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

$$\int \frac{\sec(x)}{a+b\cos^2(x)} dx = \frac{\operatorname{arctanh}(\sin(x)) - \frac{\sqrt{b}\operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}}}{a}$$

input `Integrate[Sec[x]/(a + b*Cos[x]^2), x]`

output $(\operatorname{ArcTanh}[\sin(x)] - (\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*\sin(x))/\operatorname{Sqrt}[a + b]])/\operatorname{Sqrt}[a + b])/a$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 3665, 303, 219, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x + \frac{\pi}{2}) \left(a + b \sin(x + \frac{\pi}{2})^2 \right)} dx \\
 & \quad \downarrow \text{3665} \\
 & \int \frac{1}{(1 - \sin^2(x)) (a - b \sin^2(x) + b)} d \sin(x) \\
 & \quad \downarrow \text{303} \\
 & \frac{\int \frac{1}{1 - \sin^2(x)} d \sin(x)}{a} - \frac{b \int \frac{1}{-b \sin^2(x) + a + b} d \sin(x)}{a} \\
 & \quad \downarrow \text{219} \\
 & \frac{\operatorname{arctanh}(\sin(x))}{a} - \frac{b \int \frac{1}{-b \sin^2(x) + a + b} d \sin(x)}{a} \\
 & \quad \downarrow \text{221} \\
 & \frac{\operatorname{arctanh}(\sin(x))}{a} - \frac{\sqrt{b} \operatorname{barctanh}\left(\frac{\sqrt{b} \sin(x)}{\sqrt{a+b}}\right)}{a \sqrt{a+b}}
 \end{aligned}$$

input `Int[Sec[x]/(a + b*Cos[x]^2),x]`

output `ArcTanh[Sin[x]]/a - (Sqrt[b]*ArcTanh[(Sqrt[b]*Sin[x])/Sqrt[a + b]])/(a*Sqr
t[a + b])`

Definitions of rubi rules used

rule 219 $\text{Int}[(a_0 + b_0 \cdot (x_0)^2)^{-1}, x] \rightarrow \text{Simp}[1/(Rt[a, 2] * Rt[-b, 2]) * \text{ArcTanh}[Rt[-b, 2] * (x/Rt[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{NegQ}[a/b] \& (\text{GtQ}[a, 0] \text{ || } \text{LtQ}[b, 0])$

rule 221 $\text{Int}[(a_0 + b_0 \cdot (x_0)^2)^{-1}, x] \rightarrow \text{Simp}[(Rt[-a/b, 2]/a) * \text{ArcTanh}[x/Rt[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{NegQ}[a/b]$

rule 303 $\text{Int}[1/((a_0 + b_0 \cdot (x_0)^2) * (c_0 + d_0 \cdot (x_0)^2)), x] \rightarrow \text{Simp}[b/(b * c - a * d) \text{ Int}[1/(a + b * x^2), x], x] - \text{Simp}[d/(b * c - a * d) \text{ Int}[1/(c + d * x^2), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \& \text{NeQ}[b * c - a * d, 0]$

rule 3042 $\text{Int}[u, x] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3665 $\text{Int}[\sin[(e_0 + f_0 \cdot (x_0)]^{(m_0)} * (a_0 + b_0 \cdot (x_0)) * \sin[(e_0 + f_0 \cdot (x_0))^2]^p, x] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Cos}[e + f * x], x]\}, \text{Simp}[-ff/f \text{ Subst}[\text{Int}[(1 - ff^2 * x^2)^{((m - 1)/2)} * (a + b - b * ff^2 * x^2)^p, x], x, \text{Cos}[e + f * x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \& \text{IntegerQ}[(m - 1)/2]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.15

method	result	size
default	$-\frac{\ln(\sin(x)-1)}{2a} + \frac{\ln(\sin(x)+1)}{2a} - \frac{b \operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{a \sqrt{(a+b)b}}$	47
risch	$-\frac{\ln(e^{ix}-i)}{a} + \frac{\ln(e^{ix}+i)}{a} + \frac{\sqrt{(a+b)b} \ln\left(e^{2ix} - \frac{2i\sqrt{(a+b)b}e^{ix}}{b} - 1\right)}{2(a+b)a} - \frac{\sqrt{(a+b)b} \ln\left(e^{2ix} + \frac{2i\sqrt{(a+b)b}e^{ix}}{b} - 1\right)}{2(a+b)a}$	115

input $\text{int}(\sec(x)/(a+b*\cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$-1/2/a*\ln(\sin(x)-1)+1/2/a*\ln(\sin(x)+1)-b/a/((a+b)*b)^(1/2)*\operatorname{arctanh}(b*\sin(x))/((a+b)*b)^(1/2)$$

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.90

$$\int \frac{\sec(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{\sqrt{\frac{b}{a+b}} \log \left(-\frac{b \cos(x)^2 + 2(a+b)\sqrt{\frac{b}{a+b}} \sin(x) - a - 2b}{b \cos(x)^2 + a} \right) + \log(\sin(x) + 1) - \log(-\sin(x) + 1)}{2a}, \frac{2\sqrt{-\frac{b}{a+b}} \arctan \left(\frac{\sqrt{b/a+b} \sin(x)}{\sqrt{a+b}} \right)}{a} \right]$$

input `integrate(sec(x)/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[1/2*(\sqrt{b/(a+b)})*\log(-(b*\cos(x)^2 + 2*(a+b)*\sqrt{b/(a+b)}*\sin(x) - a - 2*b)/(b*\cos(x)^2 + a)) + \log(\sin(x) + 1) - \log(-\sin(x) + 1))/a, 1/2*(2*\sqrt{-b/(a+b)})*\operatorname{arctan}(\sqrt{-b/(a+b)}*\sin(x)) + \log(\sin(x) + 1) - \log(-\sin(x) + 1))/a]$$

Sympy [F]

$$\int \frac{\sec(x)}{a + b \cos^2(x)} dx = \int \frac{\sec(x)}{a + b \cos^2(x)} dx$$

input `integrate(sec(x)/(a+b*cos(x)**2),x)`

output `Integral(sec(x)/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.56

$$\int \frac{\sec(x)}{a + b \cos^2(x)} dx = \frac{b \log \left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}} \right)}{2 \sqrt{(a+b)b} a} + \frac{\log (\sin(x) + 1)}{2a} - \frac{\log (\sin(x) - 1)}{2a}$$

input `integrate(sec(x)/(a+b*cos(x)^2),x, algorithm="maxima")`

output `1/2*b*log((b*sin(x) - sqrt((a + b)*b))/(b*sin(x) + sqrt((a + b)*b)))/(sqrt((a + b)*b)*a) + 1/2*log(sin(x) + 1)/a - 1/2*log(sin(x) - 1)/a`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.39

$$\int \frac{\sec(x)}{a + b \cos^2(x)} dx = \frac{b \arctan \left(\frac{b \sin(x)}{\sqrt{-ab - b^2}} \right)}{\sqrt{-ab - b^2} a} + \frac{\log (\sin(x) + 1)}{2a} - \frac{\log (-\sin(x) + 1)}{2a}$$

input `integrate(sec(x)/(a+b*cos(x)^2),x, algorithm="giac")`

output `b*arctan(b*sin(x)/sqrt(-a*b - b^2))/(sqrt(-a*b - b^2)*a) + 1/2*log(sin(x) + 1)/a - 1/2*log(-sin(x) + 1)/a`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 414, normalized size of antiderivative = 10.10

$$\int \frac{\sec(x)}{a + b \cos^2(x)} dx = \frac{\operatorname{atanh}(\sin(x))}{a}$$

$$\operatorname{atan} \left(\frac{\left(2 b^3 \sin(x) + \frac{\left(2 a^2 b^2 - \frac{\sin(x) (8 a^3 b^2 + 16 a^2 b^3) \sqrt{b(a+b)}}{4(a^2+b a)} \right) \sqrt{b(a+b)}}{2(a^2+b a)} \right) \sqrt{b(a+b)} 1i}{\frac{a^2+b a}{\left(2 b^3 \sin(x) + \frac{\left(2 a^2 b^2 - \frac{\sin(x) (8 a^3 b^2 + 16 a^2 b^3) \sqrt{b(a+b)}}{4(a^2+b a)} \right) \sqrt{b(a+b)}}{2(a^2+b a)} \right) \sqrt{b(a+b)}}} + \frac{\left(2 b^3 \sin(x) - \frac{\left(2 a^2 b^2 + \frac{\sin(x) (8 a^3 b^2 + 16 a^2 b^3) \sqrt{b(a+b)}}{4(a^2+b a)} \right) \sqrt{b(a+b)}}{2(a^2+b a)} \right) \sqrt{b(a+b)} 1i}{\frac{a^2+b a}{\left(2 b^3 \sin(x) - \frac{\left(2 a^2 b^2 + \frac{\sin(x) (8 a^3 b^2 + 16 a^2 b^3) \sqrt{b(a+b)}}{4(a^2+b a)} \right) \sqrt{b(a+b)}}{2(a^2+b a)} \right) \sqrt{b(a+b)}}} - \frac{\left(2 b^3 \sin(x) - \frac{\left(2 a^2 b^2 + \frac{\sin(x) (8 a^3 b^2 + 16 a^2 b^3) \sqrt{b(a+b)}}{4(a^2+b a)} \right) \sqrt{b(a+b)}}{2(a^2+b a)} \right) \sqrt{b(a+b)}}{\frac{a^2+b a}{a^2+b a}}$$

input `int(1/(cos(x)*(a + b*cos(x)^2)),x)`

output `atanh(sin(x))/a + (atan(((2*b^3*sin(x) + ((2*a^2*b^2 - (sin(x)*(16*a^2*b^3 + 8*a^3*b^2)*(b*(a + b))^(1/2))/(4*(a*b + a^2)))*(b*(a + b))^(1/2))/(2*(a*b + a^2)))*(b*(a + b))^(1/2)*1i)/(a*b + a^2) + ((2*b^3*sin(x) - ((2*a^2*b^2 + (sin(x)*(16*a^2*b^3 + 8*a^3*b^2)*(b*(a + b))^(1/2))/(4*(a*b + a^2)))*(b*(a + b))^(1/2))/(2*(a*b + a^2)))*(b*(a + b))^(1/2)*1i)/(a*b + a^2))/((2*b^3*sin(x) + ((2*a^2*b^2 - (sin(x)*(16*a^2*b^3 + 8*a^3*b^2)*(b*(a + b))^(1/2))/(4*(a*b + a^2)))*(b*(a + b))^(1/2))/(2*(a*b + a^2)))*(b*(a + b))^(1/2))/(a*b + a^2) - ((2*b^3*sin(x) - ((2*a^2*b^2 + (sin(x)*(16*a^2*b^3 + 8*a^3*b^2)*(b*(a + b))^(1/2))/(4*(a*b + a^2)))*(b*(a + b))^(1/2))/(2*(a*b + a^2)))*(b*(a + b))^(1/2))/(a*b + a^2)))*(b*(a + b))^(1/2))/(a*b + a^2)))*(b*(a + b))^(1/2)*1i)/(a*b + a^2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.83

$$\int \frac{\sec(x)}{a + b \cos^2(x)} dx$$

$$= \frac{\sqrt{b} \sqrt{a+b} \log \left(\sqrt{a+b} \tan \left(\frac{x}{2} \right)^2 + \sqrt{a+b} - 2\sqrt{b} \tan \left(\frac{x}{2} \right) \right) - \sqrt{b} \sqrt{a+b} \log \left(\sqrt{a+b} \tan \left(\frac{x}{2} \right)^2 + \sqrt{a+b} + 2\sqrt{b} \tan \left(\frac{x}{2} \right) \right)}{2a}$$

input `int(sec(x)/(a+b*cos(x)^2),x)`

output

```
(sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)
*tan(x/2)) - sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b)
+ 2*sqrt(b)*tan(x/2)) - 2*log(tan(x/2) - 1)*a - 2*log(tan(x/2) - 1)*b + 2
*log(tan(x/2) + 1)*a + 2*log(tan(x/2) + 1)*b)/(2*a*(a + b))
```

3.30 $\int \frac{\sec^3(x)}{a+b\cos^2(x)} dx$

Optimal result	235
Mathematica [B] (verified)	235
Rubi [A] (verified)	236
Maple [A] (verified)	238
Fricas [A] (verification not implemented)	239
Sympy [F]	239
Maxima [A] (verification not implemented)	240
Giac [A] (verification not implemented)	240
Mupad [B] (verification not implemented)	241
Reduce [B] (verification not implemented)	241

Optimal result

Integrand size = 15, antiderivative size = 59

$$\int \frac{\sec^3(x)}{a+b\cos^2(x)} dx = \frac{(a-2b)\operatorname{arctanh}(\sin(x))}{2a^2} + \frac{b^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{a^2\sqrt{a+b}} + \frac{\sec(x)\tan(x)}{2a}$$

output
$$\frac{1/2*(a-2*b)*\operatorname{arctanh}(\sin(x))/a^2+b^{(3/2)}*\operatorname{arctanh}(b^{(1/2)}*\sin(x)/(a+b)^{(1/2)})}{a^2/(a+b)^{(1/2)}}+1/2*\sec(x)*\tan(x)/a$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 152 vs. $2(59) = 118$.

Time = 0.48 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.58

$$\begin{aligned} & \int \frac{\sec^3(x)}{a+b\cos^2(x)} dx \\ &= \frac{-2(a-2b)\log\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) + 2(a-2b)\log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right) - \frac{2b^{3/2}\log\left(\sqrt{a+b}-\sqrt{b}\sin(x)\right)}{\sqrt{a+b}} + \frac{2b^{3/2}\log\left(\sqrt{a+b}+\sqrt{b}\sin(x)\right)}{\sqrt{a+b}}}{4a^2} \end{aligned}$$

input
$$\operatorname{Integrate}[\operatorname{Sec}[x]^3/(a+b*\operatorname{Cos}[x]^2), x]$$

output

$$(-2*(a - 2*b)*\text{Log}[\cos[x/2] - \sin[x/2]] + 2*(a - 2*b)*\text{Log}[\cos[x/2] + \sin[x/2]] - (2*b^{(3/2)}*\text{Log}[\sqrt{a + b} - \sqrt{b}*\sin[x]])/\sqrt{a + b} + (2*b^{(3/2)}*\text{Log}[\sqrt{a + b} + \sqrt{b}*\sin[x]])/\sqrt{a + b} + a/(\cos[x/2] - \sin[x/2])^2 - a/(\cos[x/2] + \sin[x/2])^2)/(4*a^2)$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.24, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3665, 316, 397, 219, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^3(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x + \frac{\pi}{2})^3 (a + b \sin(x + \frac{\pi}{2})^2)} dx \\
 & \quad \downarrow \text{3665} \\
 & \int \frac{1}{(1 - \sin^2(x))^2 (a - b \sin^2(x) + b)} d\sin(x) \\
 & \quad \downarrow \text{316} \\
 & \frac{\int \frac{-b \sin^2(x) + a - b}{(1 - \sin^2(x))(-b \sin^2(x) + a + b)} d\sin(x)}{2a} + \frac{\sin(x)}{2a (1 - \sin^2(x))} \\
 & \quad \downarrow \text{397} \\
 & \frac{\frac{2b^2 \int \frac{1}{-b \sin^2(x) + a + b} d\sin(x)}{a} + \frac{(a - 2b) \int \frac{1}{1 - \sin^2(x)} d\sin(x)}{a}}{2a} + \frac{\sin(x)}{2a (1 - \sin^2(x))} \\
 & \quad \downarrow \text{219} \\
 & \frac{\frac{2b^2 \int \frac{1}{-b \sin^2(x) + a + b} d\sin(x)}{a} + \frac{(a - 2b) \text{arctanh}(\sin(x))}{a}}{2a} + \frac{\sin(x)}{2a (1 - \sin^2(x))}
 \end{aligned}$$

↓ 221

$$\frac{\frac{2b^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{a\sqrt{a+b}} + \frac{(a-2b)\operatorname{arctanh}(\sin(x))}{a}}{2a} + \frac{\sin(x)}{2a(1-\sin^2(x))}$$

input `Int[Sec[x]^3/(a + b*Cos[x]^2), x]`

output `((a - 2*b)*ArcTanh[Sin[x]])/a + (2*b^(3/2)*ArcTanh[(Sqrt[b]*Sin[x])/Sqrt[a + b]])/(a*Sqrt[a + b])/(2*a) + Sin[x]/(2*a*(1 - Sin[x]^2))`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 316 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))), x] + Simp[1/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[b*c + 2*(p + 1)*(b*c - a*d) + d*b*(2*(p + q + 2) + 1)*x^2, x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, 2, p, q, x]`

rule 397 `Int[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*((c_) + (d_)*(x_)^2)), x_Symbol] := Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3665

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^
(p_.), x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Simplify[-ff/f
Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e +
f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.39

method	result
default	$-\frac{1}{4a(\sin(x)-1)} + \frac{(-a+2b)\ln(\sin(x)-1)}{4a^2} - \frac{1}{4a(\sin(x)+1)} + \frac{(a-2b)\ln(\sin(x)+1)}{4a^2} + \frac{b^2 \operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{a^2 \sqrt{(a+b)b}}$
risch	$-\frac{i(e^{3ix}-e^{ix})}{(e^{2ix}+1)^2 a} + \frac{\ln(e^{ix}+i)}{2a} - \frac{\ln(e^{ix}+i)b}{a^2} - \frac{\ln(e^{ix}-i)}{2a} + \frac{\ln(e^{ix}-i)b}{a^2} + \frac{\sqrt{(a+b)b} b \ln\left(\frac{e^{2ix}+2i\sqrt{(a+b)b}e^{ix}}{b}-1\right)}{2(a+b)a^2} - \frac{\sqrt{(a+b)b} b \operatorname{arctanh}\left(\frac{b \sin(x)}{\sqrt{(a+b)b}}\right)}{2(a+b)a^2}$

input `int(sec(x)^3/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output

```
-1/4/a/(\sin(x)-1)+1/4/a^2*(-a+2*b)*ln(\sin(x)-1)-1/4/a/(\sin(x)+1)+1/4*(a-2*
b)/a^2*ln(\sin(x)+1)+b^2/a^2/((a+b)*b)^(1/2)*arctanh(b*sin(x)/((a+b)*b)^(1/
2))
```

Fricas [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 186, normalized size of antiderivative = 3.15

$$\int \frac{\sec^3(x)}{a + b \cos^2(x)} dx$$

$$= \frac{2 b \sqrt{\frac{b}{a+b}} \cos(x)^2 \log\left(-\frac{b \cos(x)^2 - 2(a+b)\sqrt{\frac{b}{a+b}} \sin(x) - a - 2b}{b \cos(x)^2 + a}\right) + (a - 2b) \cos(x)^2 \log(\sin(x) + 1) - (a - 2b)}{4 a^2 \cos(x)^2}$$

$$- \frac{4 b \sqrt{-\frac{b}{a+b}} \arctan\left(\sqrt{-\frac{b}{a+b}} \sin(x)\right) \cos(x)^2 - (a - 2b) \cos(x)^2 \log(\sin(x) + 1) + (a - 2b) \cos(x)^2 \log(\sin(x) + 1)}{4 a^2 \cos(x)^2}$$

input `integrate(sec(x)^3/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[1/4*(2*b*sqrt(b/(a + b))*cos(x)^2*log(-(b*cos(x)^2 - 2*(a + b)*sqrt(b/(a + b))*sin(x) - a - 2*b)/(b*cos(x)^2 + a)) + (a - 2*b)*cos(x)^2*log(sin(x) + 1) - (a - 2*b)*cos(x)^2*log(-sin(x) + 1) + 2*a*sin(x))/(a^2*cos(x)^2), - 1/4*(4*b*sqrt(-b/(a + b))*arctan(sqrt(-b/(a + b))*sin(x))*cos(x)^2 - (a - 2*b)*cos(x)^2*log(sin(x) + 1) + (a - 2*b)*cos(x)^2*log(-sin(x) + 1) - 2*a*sin(x))/(a^2*cos(x)^2)]`

Sympy [F]

$$\int \frac{\sec^3(x)}{a + b \cos^2(x)} dx = \int \frac{\sec^3(x)}{a + b \cos^2(x)} dx$$

input `integrate(sec(x)**3/(a+b*cos(x)**2),x)`

output `Integral(sec(x)**3/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.56

$$\int \frac{\sec^3(x)}{a + b \cos^2(x)} dx = -\frac{b^2 \log\left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}}\right)}{2 \sqrt{(a+b)b} a^2} + \frac{(a - 2b) \log(\sin(x) + 1)}{4 a^2} - \frac{(a - 2b) \log(\sin(x) - 1)}{4 a^2} - \frac{\sin(x)}{2(a \sin(x)^2 - a)}$$

input `integrate(sec(x)^3/(a+b*cos(x)^2),x, algorithm="maxima")`

output
$$-1/2*b^2*log((b*sin(x) - sqrt((a + b)*b))/(b*sin(x) + sqrt((a + b)*b)))/(sqr((a + b)*b)*a^2) + 1/4*(a - 2*b)*log(sin(x) + 1)/a^2 - 1/4*(a - 2*b)*log(sin(x) - 1)/a^2 - 1/2*sin(x)/(a*sin(x)^2 - a)$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.44

$$\int \frac{\sec^3(x)}{a + b \cos^2(x)} dx = -\frac{b^2 \arctan\left(\frac{b \sin(x)}{\sqrt{-ab - b^2}}\right)}{\sqrt{-ab - b^2} a^2} + \frac{(a - 2b) \log(\sin(x) + 1)}{4 a^2} - \frac{(a - 2b) \log(-\sin(x) + 1)}{4 a^2} - \frac{\sin(x)}{2(\sin(x)^2 - 1)a}$$

input `integrate(sec(x)^3/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$-b^2*arctan(b*sin(x)/sqrt(-a*b - b^2))/(sqrt(-a*b - b^2)*a^2) + 1/4*(a - 2*b)*log(sin(x) + 1)/a^2 - 1/4*(a - 2*b)*log(-sin(x) + 1)/a^2 - 1/2*sin(x)/((sin(x)^2 - 1)*a)$$

Mupad [B] (verification not implemented)

Time = 1.13 (sec) , antiderivative size = 483, normalized size of antiderivative = 8.19

$$\int \frac{\sec^3(x)}{a + b \cos^2(x)} dx =$$

$$a^2 \sin(x) + a^2 \operatorname{atanh}(\sin(x)) - 2b^2 \operatorname{atanh}(\sin(x)) + ab \sin(x) - ab \operatorname{atanh}(\sin(x)) - a^2 \operatorname{atanh}(\sin(x))$$

—

input `int(1/(cos(x)^3*(a + b*cos(x)^2)),x)`

output

$$-(a^2 \sin(x) + a^2 \operatorname{atanh}(\sin(x)) - 2b^2 \operatorname{atanh}(\sin(x)) + \operatorname{atan}((b^5 \sin(x) * (a*b^3 + b^4)^{(1/2)*8i} - a \sin(x) * (a*b^3 + b^4)^{(3/2)*4i} - b \sin(x) * (a*b^3 + b^4)^{(3/2)*8i} + a*b^4 * \sin(x) * (a*b^3 + b^4)^{(1/2)*12i} + a^4 * b * \sin(x) * (a*b^3 + b^4)^{(1/2)*1i} + a^2 * b^3 * \sin(x) * (a*b^3 + b^4)^{(1/2)*1i} - a^3 * b^2 * \sin(x) * (a*b^3 + b^4)^{(1/2)*2i}) / (3*a^2 * b^5 + 5*a^3 * b^4 + a^4 * b^3 - a^5 * b^2)) * (a * b^3 + b^4)^{(1/2)*2i} + a * b * \sin(x) - a * b * \operatorname{atanh}(\sin(x)) - a^2 * \operatorname{atanh}(\sin(x)) * \sin(x)^2 + 2 * b^2 * \operatorname{atanh}(\sin(x)) * \sin(x)^2 - \operatorname{atan}((b^5 \sin(x) * (a*b^3 + b^4)^{(1/2)*8i} - a \sin(x) * (a*b^3 + b^4)^{(3/2)*4i} - b \sin(x) * (a*b^3 + b^4)^{(3/2)*8i} + a*b^4 * \sin(x) * (a*b^3 + b^4)^{(1/2)*12i} + a^4 * b * \sin(x) * (a*b^3 + b^4)^{(1/2)*1i} + a^2 * b^3 * \sin(x) * (a*b^3 + b^4)^{(1/2)*1i} - a^3 * b^2 * \sin(x) * (a*b^3 + b^4)^{(1/2)*2i}) / (3*a^2 * b^5 + 5*a^3 * b^4 + a^4 * b^3 - a^5 * b^2)) * \sin(x)^2 * (a*b^3 + b^4)^{(1/2)*2i} + a * b * \operatorname{atanh}(\sin(x)) * \sin(x)^2) / (2*a^3 * \sin(x)^2 - 2*a^2 * b - 2*a^3 + 2*a^2 * b * \sin(x)^2)$$
Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 342, normalized size of antiderivative = 5.80

$$\int \frac{\sec^3(x)}{a + b \cos^2(x)} dx =$$

$$-\sqrt{b} \sqrt{a+b} \log\left(\sqrt{a+b} \tan\left(\frac{x}{2}\right)^2 + \sqrt{a+b} - 2\sqrt{b} \tan\left(\frac{x}{2}\right)\right) \sin(x)^2 b + \sqrt{b} \sqrt{a+b} \log\left(\sqrt{a+b} \tan\left(\frac{x}{2}\right)\right)$$

input `int(sec(x)^3/(a+b*cos(x)^2),x)`

```
output
( - sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)*tan(x/2))*sin(x)**2*b + sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)*tan(x/2))*b + sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan(x/2))*sin(x)**2*b - sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan(x/2))*b - log(tan(x/2) - 1)*sin(x)**2*a**2 + log(tan(x/2) - 1)*sin(x)**2*a*b + 2*log(tan(x/2) - 1)*sin(x)**2*b**2 + log(tan(x/2) - 1)*a**2 - log(tan(x/2) - 1)*a*b - 2*log(tan(x/2) - 1)*b**2 + log(tan(x/2) + 1)*sin(x)**2*a**2 - log(tan(x/2) + 1)*sin(x)**2*a*b - 2*log(tan(x/2) + 1)*sin(x)**2*b**2 - log(tan(x/2) + 1)*a**2 + log(tan(x/2) + 1)*a*b + 2*log(tan(x/2) + 1)*b**2 - sin(x)*a**2 - sin(x)*a*b)/(2*a**2*(sin(x)**2*a + sin(x)**2*b - a - b))
```

3.31 $\int \frac{\sec^5(x)}{a+b\cos^2(x)} dx$

Optimal result	243
Mathematica [B] (verified)	243
Rubi [A] (verified)	244
Maple [A] (verified)	247
Fricas [A] (verification not implemented)	247
Sympy [F]	248
Maxima [A] (verification not implemented)	248
Giac [A] (verification not implemented)	249
Mupad [B] (verification not implemented)	249
Reduce [B] (verification not implemented)	250

Optimal result

Integrand size = 15, antiderivative size = 90

$$\begin{aligned} \int \frac{\sec^5(x)}{a+b\cos^2(x)} dx &= \frac{(3a^2 - 4ab + 8b^2) \operatorname{arctanh}(\sin(x))}{8a^3} - \frac{b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{a^3\sqrt{a+b}} \\ &\quad + \frac{(3a - 4b) \sec(x) \tan(x)}{8a^2} + \frac{\sec^3(x) \tan(x)}{4a} \end{aligned}$$

output

```
1/8*(3*a^2-4*a*b+8*b^2)*arctanh(sin(x))/a^3-b^(5/2)*arctanh(b^(1/2)*sin(x)
/(a+b)^(1/2))/a^3/(a+b)^(1/2)+1/8*(3*a-4*b)*sec(x)*tan(x)/a^2+1/4*sec(x)^3
*tan(x)/a
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 215 vs. $2(90) = 180$.

Time = 1.35 (sec), antiderivative size = 215, normalized size of antiderivative = 2.39

$$\begin{aligned} \int \frac{\sec^5(x)}{a+b\cos^2(x)} dx &= \frac{-2(3a^2 - 4ab + 8b^2) \log\left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) + 2(3a^2 - 4ab + 8b^2) \log\left(\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)\right) + \frac{8b^{5/2} \log\left(\sqrt{a+b}\right)}{\sqrt{a+b}}}{16a^3} \end{aligned}$$

input `Integrate[Sec[x]^5/(a + b*Cos[x]^2),x]`

output
$$\begin{aligned} & (-2*(3*a^2 - 4*a*b + 8*b^2)*Log[Cos[x/2] - Sin[x/2]] + 2*(3*a^2 - 4*a*b + \\ & 8*b^2)*Log[Cos[x/2] + Sin[x/2]] + (8*b^{(5/2)}*Log[Sqrt[a + b] - Sqrt[b]*Sin[x]])/Sqrt[a + b] - (8*b^{(5/2)}*Log[Sqrt[a + b] + Sqrt[b]*Sin[x]])/Sqrt[a + b] + a^2/(Cos[x/2] - Sin[x/2])^4 - a^2/(Cos[x/2] + Sin[x/2])^4 + (a*(-3*a + 4*b))/(Cos[x/2] + Sin[x/2])^2 + (a*(-3*a + 4*b))/(-1 + Sin[x]))/(16*a^3) \end{aligned}$$

Rubi [A] (verified)

Time = 0.34 (sec), antiderivative size = 117, normalized size of antiderivative = 1.30, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 3665, 316, 402, 397, 219, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sec^5(x)}{a + b \cos^2(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{\sin(x + \frac{\pi}{2})^5 (a + b \sin(x + \frac{\pi}{2})^2)} dx \\ & \quad \downarrow \text{3665} \\ & \int \frac{1}{(1 - \sin^2(x))^3 (a - b \sin^2(x) + b)} d\sin(x) \\ & \quad \downarrow \text{316} \\ & \frac{\int \frac{-3b \sin^2(x) + 3a - b}{(1 - \sin^2(x))^2 (-b \sin^2(x) + a + b)} d\sin(x)}{4a} + \frac{\sin(x)}{4a (1 - \sin^2(x))^2} \\ & \quad \downarrow \text{402} \\ & \frac{\int \frac{3a^2 - ba + 4b^2 - (3a - 4b)b \sin^2(x)}{(1 - \sin^2(x))(-b \sin^2(x) + a + b)} d\sin(x)}{2a} + \frac{(3a - 4b)\sin(x)}{2a(1 - \sin^2(x))} + \frac{\sin(x)}{4a (1 - \sin^2(x))^2} \end{aligned}$$

$$\begin{aligned}
 & \frac{\left(3a^2 - 4ab + 8b^2\right) \int \frac{1}{1-\sin^2(x)} d\sin(x) - 8b^3 \int \frac{1}{-b\sin^2(x)+a+b} d\sin(x)}{4a} + \frac{(3a-4b)\sin(x)}{2a(1-\sin^2(x))} + \frac{\sin(x)}{4a(1-\sin^2(x))^2} \\
 & \downarrow 219 \\
 & \frac{\left(3a^2 - 4ab + 8b^2\right) \operatorname{arctanh}(\sin(x)) - 8b^3 \int \frac{1}{-b\sin^2(x)+a+b} d\sin(x)}{4a} + \frac{(3a-4b)\sin(x)}{2a(1-\sin^2(x))} + \frac{\sin(x)}{4a(1-\sin^2(x))^2} \\
 & \downarrow 221 \\
 & \frac{\left(3a^2 - 4ab + 8b^2\right) \operatorname{arctanh}(\sin(x)) - \frac{8b^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sin(x)}{\sqrt{a+b}}\right)}{a\sqrt{a+b}}}{4a} + \frac{(3a-4b)\sin(x)}{2a(1-\sin^2(x))} + \frac{\sin(x)}{4a(1-\sin^2(x))^2}
 \end{aligned}$$

input `Int[Sec[x]^5/(a + b*Cos[x]^2), x]`

output `Sin[x]/(4*a*(1 - Sin[x]^2)^2) + (((3*a^2 - 4*a*b + 8*b^2)*ArcTanh[Sin[x]])/a - (8*b^(5/2)*ArcTanh[(Sqrt[b]*Sin[x])/Sqrt[a + b]])/(a*Sqrt[a + b]))/(2*a) + ((3*a - 4*b)*Sin[x])/(2*a*(1 - Sin[x]^2))/(4*a)`

Definitions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 316 $\text{Int}[(a_ + b_*)*(x_)^2]^{(p_)}*((c_ + d_*)*(x_)^2)]^{(q_)} \rightarrow \text{Simp}[(-b)*x*(a + b*x^2)]^{(p+1)}*((c + d*x^2)]^{(q+1)}/(2*a*(p+1)*(b*c - a*d))$
 $, x] + \text{Simp}[1/(2*a*(p+1)*(b*c - a*d)) \text{ Int}[(a + b*x^2)]^{(p+1)}*(c + d*x^2)]^{(q+2)} \text{q}*\text{Simp}[b*c + 2*(p+1)*(b*c - a*d) + d*b*(2*(p+q+2) + 1)*x^2, x], x]$
 $], x] /; \text{FreeQ}[\{a, b, c, d, q\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{LtQ}[p, -1] \&& !$
 $(\text{!IntegerQ}[p] \&& \text{IntegerQ}[q] \&& \text{LtQ}[q, -1]) \&& \text{IntBinomialQ}[a, b, c, d, 2, p, q, x]$

rule 397 $\text{Int}[(e_ + f_*)*(x_)^2]/(((a_ + b_*)*(x_)^2)*((c_ + d_*)*(x_)^2)) \rightarrow \text{Simp}[(b*e - a*f)/(b*c - a*d) \text{ Int}[1/(a + b*x^2), x], x] - \text{Simp}[(d*e - c*f)/(b*c - a*d) \text{ Int}[1/(c + d*x^2), x], x]$
 $/; \text{FreeQ}[\{a, b, c, d, e, f\}, x]$

rule 402 $\text{Int}[(a_ + b_*)*(x_)^2]^{(p_)}*((c_ + d_*)*(x_)^2)]^{(q_)}*((e_ + f_*)*(x_)^2) \rightarrow \text{Simp}[(-(b*e - a*f))*x*(a + b*x^2)]^{(p+1)}*((c + d*x^2)]^{(q+1)}/(a*2*(b*c - a*d)*(p+1)), x] + \text{Simp}[1/(a*2*(b*c - a*d)*(p+1))$
 $\text{Int}[(a + b*x^2)]^{(p+1)}*(c + d*x^2)]^{(q+2)} \text{q}*\text{Simp}[c*(b*e - a*f) + e*2*(b*c - a*d)$
 $*(p+1) + d*(b*e - a*f)*(2*(p+q+2) + 1)*x^2, x], x]$
 $/; \text{FreeQ}[\{a, b, c, d, e, f, q\}, x] \&& \text{LtQ}[p, -1]$

rule 3042 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinear}$
 $Q[u, x]$

rule 3665 $\text{Int}[\sin[(e_ + f_*)*(x_)]^m*((a_ + b_*)*\sin[(e_ + f_*)*(x_)]^2)]^{(p_)} \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Cos}[e + f*x], x]\}, \text{Simp}[-ff/f$
 $\text{Subst}[\text{Int}[(1 - ff^2*x^2)]^{((m-1)/2)}*(a + b - b*ff^2*x^2)^p, x], x, \text{Cos}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{IntegerQ}[(m-1)/2]$

Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.52

method	result
default	$\frac{1}{16a(\sin(x)-1)^2} - \frac{3a-4b}{16a^2(\sin(x)-1)} + \frac{(-3a^2+4ab-8b^2)\ln(\sin(x)-1)}{16a^3} - \frac{1}{16a(\sin(x)+1)^2} - \frac{3a-4b}{16a^2(\sin(x)+1)} + \frac{(3a^2-4ab+8b^2)\ln(\sin(x)+1)}{16a^3}$
risch	$-\frac{i(3ae^{7ix}-4be^{7ix}+11ae^{5ix}-4be^{5ix}-11ae^{3ix}+4be^{3ix}-3e^{ix}a+4e^{ix}b)}{4(e^{2ix}+1)^4a^2} + \frac{3\ln(e^{ix}+i)}{8a} - \frac{\ln(e^{ix}+i)b}{2a^2} + \frac{\ln(e^{ix}+i)b^2}{a^3} - \frac{3\ln(e^{ix}+i)a^2}{16a^4}$

input `int(sec(x)^5/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/16/a/(\sin(x)-1)^2 - 1/16*(3*a-4*b)/a^2/(\sin(x)-1) + 1/16/a^3*(-3*a^2+4*a*b-8*b^2)*\ln(\sin(x)-1) - 1/16/a/(\sin(x)+1)^2 - 1/16*(3*a-4*b)/a^2/(\sin(x)+1) + 1/16*(3*a^2-4*a*b+8*b^2)/a^3*\ln(\sin(x)+1) - b^3/a^3/((a+b)*b)^(1/2)*\operatorname{arctanh}(b*\sin(x))/((a+b)*b)^(1/2) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 270, normalized size of antiderivative = 3.00

$$\begin{aligned} & \int \frac{\sec^5(x)}{a + b \cos^2(x)} dx \\ &= \frac{8 b^2 \sqrt{\frac{b}{a+b}} \cos(x)^4 \log \left(-\frac{b \cos(x)^2 + 2(a+b) \sqrt{\frac{b}{a+b}} \sin(x) - a - 2b}{b \cos(x)^2 + a} \right) + (3a^2 - 4ab + 8b^2) \cos(x)^4 \log(\sin(x) + 1)}{16a^3 \cos(x)^4} \end{aligned}$$

input `integrate(sec(x)^5/(a+b*cos(x)^2),x, algorithm="fricas")`

output

```
[1/16*(8*b^2*sqrt(b/(a + b)))*cos(x)^4*log(-(b*cos(x)^2 + 2*(a + b)*sqrt(b/(a + b))*sin(x) - a - 2*b)/(b*cos(x)^2 + a)) + (3*a^2 - 4*a*b + 8*b^2)*cos(x)^4*log(sin(x) + 1) - (3*a^2 - 4*a*b + 8*b^2)*cos(x)^4*log(-sin(x) + 1) + 2*((3*a^2 - 4*a*b)*cos(x)^2 + 2*a^2)*sin(x))/(a^3*cos(x)^4), 1/16*(16*b^2*sqrt(-b/(a + b)))*arctan(sqrt(-b/(a + b))*sin(x))*cos(x)^4 + (3*a^2 - 4*a*b + 8*b^2)*cos(x)^4*log(sin(x) + 1) - (3*a^2 - 4*a*b + 8*b^2)*cos(x)^4*log(-sin(x) + 1) + 2*((3*a^2 - 4*a*b)*cos(x)^2 + 2*a^2)*sin(x))/(a^3*cos(x)^4)]
```

Sympy [F]

$$\int \frac{\sec^5(x)}{a + b \cos^2(x)} dx = \int \frac{\sec^5(x)}{a + b \cos^2(x)} dx$$

input

```
integrate(sec(x)**5/(a+b*cos(x)**2),x)
```

output

```
Integral(sec(x)**5/(a + b*cos(x)**2), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.61

$$\int \frac{\sec^5(x)}{a + b \cos^2(x)} dx = \frac{b^3 \log\left(\frac{b \sin(x) - \sqrt{(a+b)b}}{b \sin(x) + \sqrt{(a+b)b}}\right)}{2 \sqrt{(a+b)b} a^3} - \frac{(3a - 4b) \sin(x)^3 - (5a - 4b) \sin(x)}{8 (a^2 \sin(x)^4 - 2a^2 \sin(x)^2 + a^2)} + \frac{(3a^2 - 4ab + 8b^2) \log(\sin(x) + 1)}{16 a^3} - \frac{(3a^2 - 4ab + 8b^2) \log(\sin(x) - 1)}{16 a^3}$$

input

```
integrate(sec(x)^5/(a+b*cos(x)^2),x, algorithm="maxima")
```

output

$$\frac{1/2*b^3*\log((b*\sin(x) - \sqrt{(a + b)*b}))/(\sqrt{(a + b)*b})}{\sqrt{(a + b)*b}}/(a^2*\sin(x)^4 - 2*a^2*\sin(x)^2 + a^2) + \frac{1/16*(3*a^2 - 4*a*b + 8*b^2)*\log(\sin(x) + 1)/a^3 - 1/16*(3*a^2 - 4*a*b + 8*b^2)*\log(\sin(x) - 1)/a^3}{a^2}$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.41

$$\int \frac{\sec^5(x)}{a + b \cos^2(x)} dx = \frac{b^3 \arctan\left(\frac{b \sin(x)}{\sqrt{-ab - b^2}}\right)}{\sqrt{-ab - b^2} a^3} + \frac{(3 a^2 - 4 ab + 8 b^2) \log(\sin(x) + 1)}{16 a^3} - \frac{(3 a^2 - 4 ab + 8 b^2) \log(-\sin(x) + 1)}{16 a^3} - \frac{3 a \sin(x)^3 - 4 b \sin(x)^3 - 5 a \sin(x) + 4 b \sin(x)}{8 (\sin(x)^2 - 1)^2 a^2}$$

input

```
integrate(sec(x)^5/(a+b*cos(x)^2),x, algorithm="giac")
```

output

$$\frac{b^3 \arctan(b \sin(x) / \sqrt{-a*b - b^2})}{\sqrt{-a*b - b^2} * a^3} + \frac{1/16*(3*a^2 - 4*a*b + 8*b^2)*\log(\sin(x) + 1)/a^3 - 1/16*(3*a^2 - 4*a*b + 8*b^2)*\log(-\sin(x) + 1)/a^3 - 1/8*(3*a*\sin(x)^3 - 4*b*\sin(x)^3 - 5*a*\sin(x) + 4*b*\sin(x)) / ((\sin(x)^2 - 1)^2 * a^2)}$$

Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 969, normalized size of antiderivative = 10.77

$$\int \frac{\sec^5(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input

```
int(1/(\cos(x)^5*(a + b*cos(x)^2)),x)
```

output

```
(5*a^3*sin(x) + atan((b^7*sin(x)*(a*b^5 + b^6)^(1/2)*128i - a*sin(x)*(a*b^5 + b^6)^(3/2)*64i - b*sin(x)*(a*b^5 + b^6)^(3/2)*128i + a*b^6*sin(x)*(a*b^5 + b^6)^(1/2)*192i + a^6*b*sin(x)*(a*b^5 + b^6)^(1/2)*9i + a^2*b^5*sin(x)*(a*b^5 + b^6)^(1/2)*64i + a^3*b^4*sin(x)*(a*b^5 + b^6)^(1/2)*40i + a^4*b^3*sin(x)*(a*b^5 + b^6)^(1/2)*25i - a^5*b^2*sin(x)*(a*b^5 + b^6)^(1/2)*6i)/(40*a^3*b^7 + 65*a^4*b^6 + 19*a^5*b^5 + 3*a^6*b^4 + 9*a^7*b^3))*(a*b^5 + b^6)^(1/2)*8i - 3*a^3*sin(x)^3 + 3*a^3*atanh(sin(x)) + 8*b^3*atanh(sin(x)) - 4*a*b^2*sin(x) + a^2*b*sin(x) - atan((b^7*sin(x)*(a*b^5 + b^6)^(1/2)*128i - a*sin(x)*(a*b^5 + b^6)^(3/2)*64i - b*sin(x)*(a*b^5 + b^6)^(3/2)*128i + a*b^6*sin(x)*(a*b^5 + b^6)^(1/2)*192i + a^6*b*sin(x)*(a*b^5 + b^6)^(1/2)*9i + a^2*b^5*sin(x)*(a*b^5 + b^6)^(1/2)*64i + a^3*b^4*sin(x)*(a*b^5 + b^6)^(1/2)*40i + a^4*b^3*sin(x)*(a*b^5 + b^6)^(1/2)*25i - a^5*b^2*sin(x)*(a*b^5 + b^6)^(1/2)*6i)/(40*a^3*b^7 + 65*a^4*b^6 + 19*a^5*b^5 + 3*a^6*b^4 + 9*a^7*b^3))*sin(x)^2*(a*b^5 + b^6)^(1/2)*16i + atan((b^7*sin(x)*(a*b^5 + b^6)^(1/2)*128i - a*sin(x)*(a*b^5 + b^6)^(3/2)*64i - b*sin(x)*(a*b^5 + b^6)^(3/2)*128i + a*b^6*sin(x)*(a*b^5 + b^6)^(1/2)*192i + a^6*b*sin(x)*(a*b^5 + b^6)^(1/2)*9i + a^2*b^5*sin(x)*(a*b^5 + b^6)^(1/2)*64i + a^3*b^4*sin(x)*(a*b^5 + b^6)^(1/2)*40i + a^4*b^3*sin(x)*(a*b^5 + b^6)^(1/2)*25i - a^5*b^2*sin(x)*(a*b^5 + b^6)^(1/2)*6i)/(40*a^3*b^7 + 65*a^4*b^6 + 19*a^5*b^5 + 3*a^6*b^4 + 9*a^7*b^3))*sin(x)^4*(a*b^5 + b^6)^(1/2)*8i - 6*a^3*atanh(sin(x)...)
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 681, normalized size of antiderivative = 7.57

$$\int \frac{\sec^5(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input

```
int(sec(x)^5/(a+b*cos(x)^2),x)
```

output

```
(4*sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)*tan(x/2))*sin(x)**4*b**2 - 8*sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)*tan(x/2))*sin(x)**2*b**2 + 4*sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) - 2*sqrt(b)*tan(x/2))*b**2 - 4*sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan(x/2))*sin(x)**4*b**2 + 8*sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan(x/2))*sin(x)**2*b**2 - 4*sqrt(b)*sqrt(a + b)*log(sqrt(a + b)*tan(x/2)**2 + sqrt(a + b) + 2*sqrt(b)*tan(x/2))*b**2 - 3*log(tan(x/2) - 1)*sin(x)**4*a**3 + log(tan(x/2) - 1)*sin(x)**4*a**2*b - 4*log(tan(x/2) - 1)*sin(x)**4*a*b**2 - 8*log(tan(x/2) - 1)*sin(x)**4*b**3 + 6*log(tan(x/2) - 1)*sin(x)**2*a**3 - 2*log(tan(x/2) - 1)*sin(x)**2*a**2*b + 16*log(tan(x/2) - 1)*sin(x)**2*b**3 - 3*log(tan(x/2) - 1)*a**3 + log(tan(x/2) - 1)*a**2*b - 4*log(tan(x/2) - 1)*a*b**2 - 8*log(tan(x/2) - 1)*b**3 + 3*log(tan(x/2) + 1)*sin(x)**4*a**3 - log(tan(x/2) + 1)*sin(x)**4*a**2*b + 4*log(tan(x/2) + 1)*sin(x)**4*a*b**2 + 8*log(tan(x/2) + 1)*sin(x)**4*b**3 - 6*log(tan(x/2) + 1)*sin(x)**2*a**3 + 2*log(tan(x/2) + 1)*sin(x)**2*a**2*b - 8*log(tan(x/2) + 1)*sin(x)**2*b**3 + 3*log(tan(x/2) + 1)*a**3 - log(tan(x/2) + 1)*a**2*b + 4*log(tan(x/2) + 1)*a*b**2 + 8*log(tan(x/2) + 1)*b**3 - 3*sin(x)**3*a**3 + sin(x)**3*a**2*b + 4*sin(x)**3*a*...
```

3.32 $\int \frac{\cos^6(x)}{a+b\cos^2(x)} dx$

Optimal result	252
Mathematica [A] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	255
Fricas [A] (verification not implemented)	256
Sympy [F(-1)]	256
Maxima [A] (verification not implemented)	257
Giac [A] (verification not implemented)	257
Mupad [B] (verification not implemented)	258
Reduce [B] (verification not implemented)	258

Optimal result

Integrand size = 15, antiderivative size = 87

$$\int \frac{\cos^6(x)}{a + b\cos^2(x)} dx = \frac{(8a^2 - 4ab + 3b^2)x}{8b^3} + \frac{a^{5/2}\arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{b^3\sqrt{a+b}} - \frac{(4a - 3b)\cos(x)\sin(x)}{8b^2} + \frac{\cos^3(x)\sin(x)}{4b}$$

output $1/8*(8*a^2-4*a*b+3*b^2)*x/b^3+a^(5/2)*\arctan((a+b)^(1/2)*\cot(x)/a^(1/2))/b^3/(a+b)^(1/2)-1/8*(4*a-3*b)*\cos(x)*\sin(x)/b^2+1/4*\cos(x)^3*\sin(x)/b$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.87

$$\begin{aligned} & \int \frac{\cos^6(x)}{a + b\cos^2(x)} dx \\ &= \frac{4(8a^2 - 4ab + 3b^2)x - \frac{32a^{5/2}\arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - 8(a-b)b\sin(2x) + b^2\sin(4x)}{32b^3} \end{aligned}$$

input `Integrate[Cos[x]^6/(a + b*Cos[x]^2), x]`

output
$$(4*(8*a^2 - 4*a*b + 3*b^2)*x - (32*a^{(5/2)}*\text{ArcTan}[(\text{Sqrt}[a]*\text{Tan}[x])/\text{Sqrt}[a+b]])/\text{Sqrt}[a+b] - 8*(a-b)*b*\text{Sin}[2*x] + b^2*\text{Sin}[4*x])/(32*b^3)$$

Rubi [A] (verified)

Time = 0.38 (sec), antiderivative size = 116, normalized size of antiderivative = 1.33, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 3666, 372, 440, 397, 216, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^6(x)}{a + b \cos^2(x)} dx \\
 & \downarrow \text{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})^6}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \downarrow \text{3666} \\
 & - \int \frac{\cot^6(x)}{(\cot^2(x) + 1)^3 ((a + b) \cot^2(x) + a)} d \cot(x) \\
 & \downarrow \text{372} \\
 & \frac{\cot^3(x)}{4b (\cot^2(x) + 1)^2} - \frac{\int \frac{\cot^2(x)(3a - (a - 3b) \cot^2(x))}{(\cot^2(x) + 1)^2 ((a + b) \cot^2(x) + a)} d \cot(x)}{4b} \\
 & \downarrow \text{440} \\
 & \frac{\cot^3(x)}{4b (\cot^2(x) + 1)^2} - \frac{\frac{(4a - 3b) \cot(x)}{2b (\cot^2(x) + 1)} - \frac{\int \frac{a(4a - 3b) - (4a^2 - ba + 3b^2) \cot^2(x)}{(\cot^2(x) + 1)^2 ((a + b) \cot^2(x) + a)} d \cot(x)}{2b}}{4b} \\
 & \downarrow \text{397} \\
 & \frac{\cot^3(x)}{4b (\cot^2(x) + 1)^2} - \frac{\frac{(4a - 3b) \cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{8a^3 \int \frac{1}{(a + b) \cot^2(x) + a} d \cot(x)}{b} - \frac{(8a^2 - 4ab + 3b^2) \int \frac{1}{\cot^2(x) + 1} d \cot(x)}{b}}{2b}}{4b} \\
 & \downarrow \text{216}
 \end{aligned}$$

$$\frac{\cot^3(x)}{4b(\cot^2(x) + 1)^2} - \frac{\frac{(4a-3b)\cot(x)}{2b(\cot^2(x)+1)} - \frac{\frac{8a^3 \int \frac{1}{(a+b)\cot^2(x)+a} d\cot(x)}{b} - \frac{(8a^2-4ab+3b^2)}{b} \arctan(\cot(x))}{2b}}{4b}$$

↓ 218

$$\frac{\cot^3(x)}{4b(\cot^2(x) + 1)^2} - \frac{\frac{(4a-3b)\cot(x)}{2b(\cot^2(x)+1)} - \frac{\frac{8a^{5/2} \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{b\sqrt{a+b}} - \frac{(8a^2-4ab+3b^2)}{b} \arctan(\cot(x))}{2b}}{4b}$$

input `Int[Cos[x]^6/(a + b*Cos[x]^2), x]`

output `Cot[x]^3/(4*b*(1 + Cot[x]^2)^2) - (-1/2*(-(((8*a^2 - 4*a*b + 3*b^2)*ArcTan[Cot[x]])/b) + (8*a^(5/2)*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(b*Sqrt[a + b]))/b + ((4*a - 3*b)*Cot[x])/(2*b*(1 + Cot[x]^2)))/(4*b)`

Definitions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 372 `Int[((e_)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(-a)*e^3*(e*x)^(m - 3)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*b*(b*c - a*d)*(p + 1))), x] + Simp[e^4/(2*b*(b*c - a*d)*(p + 1)) Int[(e*x)^(m - 4)*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[a*c*(m - 3) + (a*d*(m + 2*q - 1) + 2*b*c*(p + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[m, 3] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`

rule 397

```
Int[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*((c_) + (d_)*(x_)^2)), x_
Symbol] :> Simp[(b*e - a*f)/(b*c - a*d) Int[1/(a + b*x^2), x], x] - Simp[
(d*e - c*f)/(b*c - a*d) Int[1/(c + d*x^2), x], x] /; FreeQ[{a, b, c, d, e
, f}, x]
```

rule 440

```
Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)
)*((e_) + (f_)*(x_)^2), x_Symbol] :> Simp[g*(b*e - a*f)*(g*x)^(m - 1)*(a +
b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*b*(b*c - a*d)*(p + 1))), x] - Simp[
g^2/(2*b*(b*c - a*d)*(p + 1)) Int[(g*x)^(m - 2)*(a + b*x^2)^(p + 1)*(c +
d*x^2)^(q + 1)*Simp[c*(b*e - a*f)*(m - 1) + (d*(b*e - a*f)*(m + 2*q + 1) - b*2*(c
*f - d*e)*(p + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, g, q}, x] &&
LtQ[p, -1] && GtQ[m, 1]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3666

```
Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_
), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff^(m + 1
)/f Subst[Int[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1))
, x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] &
& IntegerQ[p]
```

Maple [A] (verified)

Time = 1.11 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.06

method	result
default	$\frac{\frac{(-\frac{1}{2}ab+\frac{3}{8}b^2)\tan(x)^3+(-\frac{1}{2}ab+\frac{5}{8}b^2)\tan(x)}{(1+\tan(x)^2)^2}+\frac{(8a^2-4ab+3b^2)\arctan(\tan(x))}{8}}{b^3}-\frac{a^3\arctan\left(\frac{a\tan(x)}{\sqrt{(a+b)a}}\right)}{b^3\sqrt{(a+b)a}}$
risch	$\frac{xa^2}{b^3}-\frac{xa}{2b^2}+\frac{3x}{8b}+\frac{ie^{2ix}a}{8b^2}-\frac{ie^{2ix}a}{8b}-\frac{ie^{-2ix}a}{8b^2}+\frac{ie^{-2ix}}{8b}+\frac{\sqrt{-(a+b)a}a^2\ln\left(e^{2ix}+\frac{2i\sqrt{-(a+b)a}+2a+b}{b}\right)}{2(a+b)b^3}-\frac{\sqrt{-(a+b)a}a}{b}$

input `int(cos(x)^6/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{b^3} \left(\left(-\frac{1}{2}ab + \frac{3}{8}b^2 \right) \tan(x)^3 + \left(-\frac{1}{2}ab + \frac{5}{8}b^2 \right) \tan(x) \right) / (1 + \tan(x)^2)^2 + \frac{1}{8} \left(8a^2 - 4ab + 3b^2 \right) \arctan(\tan(x)) - \frac{1}{b^3} a^3 / ((a+b)a^{(1/2)} \arctan(a \tan(x)) / ((a+b)a^{(1/2)}))$$

Fricas [A] (verification not implemented)

Time = 0.21 (sec), antiderivative size = 273, normalized size of antiderivative = 3.14

$$\int \frac{\cos^6(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{2a^2 \sqrt{-\frac{a}{a+b}} \log \left(\frac{(8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 + 4((2a^2 + 3ab + b^2) \cos(x)^3 - (a^2 + ab) \cos(x)) \sqrt{-\frac{a}{a+b}} \sin(x) + a^2}{b^2 \cos(x)^4 + 2ab \cos(x)^2 + a^2} \right)}{8b^3} \right]$$

input `integrate(cos(x)^6/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[\frac{1}{8} (2a^2 \sqrt{-a/(a+b)} \log((8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 + 4((2a^2 + 3ab + b^2) \cos(x)^3 - (a^2 + ab) \cos(x)) \sqrt{-a/(a+b)} \sin(x) + a^2) + 4((2a^2 + 3ab + b^2) \cos(x)^3 - (a^2 + ab) \cos(x)) \sqrt{-a/(a+b)} \sin(x) + a^2) / (b^2 \cos(x)^4 + 2ab \cos(x)^2 + a^2) + (8a^2 - 4ab + 3b^2)x + (2b^2 \cos(x)^3 - (4ab - 3b^2) \cos(x)) \sin(x) / b^3, \frac{1}{8} (4a^2 \sqrt{a/(a+b)} \arctan(1/2 * ((2a + b) \cos(x)^2 - a) * \sqrt{a/(a+b)}) / (a \cos(x) \sin(x)) + (8a^2 - 4ab + 3b^2)x + (2b^2 \cos(x)^3 - (4ab - 3b^2) \cos(x)) \sin(x) / b^3]$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^6(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(cos(x)**6/(a+b*cos(x)**2),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.11

$$\int \frac{\cos^6(x)}{a + b \cos^2(x)} dx = -\frac{a^3 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)ab^3}} - \frac{(4a - 3b) \tan(x)^3 + (4a - 5b) \tan(x)}{8(b^2 \tan(x)^4 + 2b^2 \tan(x)^2 + b^2)} \\ + \frac{(8a^2 - 4ab + 3b^2)x}{8b^3}$$

input `integrate(cos(x)^6/(a+b*cos(x)^2),x, algorithm="maxima")`

output
$$-\frac{a^3 \arctan(a \tan(x))}{\sqrt{(a+b)a} b^3} - \frac{1}{8}((4a - 3b) \tan(x)^3 + (4a - 5b) \tan(x)) / (b^2 \tan(x)^4 + 2b^2 \tan(x)^2 + b^2) \\ + \frac{1}{8}(8a^2 - 4ab + 3b^2)x/b^3$$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.20

$$\int \frac{\cos^6(x)}{a + b \cos^2(x)} dx = -\frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right) a^3}{\sqrt{a^2 + ab} b^3} + \frac{(8a^2 - 4ab + 3b^2)x}{8b^3} \\ - \frac{4a \tan(x)^3 - 3b \tan(x)^3 + 4a \tan(x) - 5b \tan(x)}{8(\tan(x)^2 + 1)^2 b^2}$$

input `integrate(cos(x)^6/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$-(\pi * \operatorname{floor}(x/\pi + 1/2) * \operatorname{sgn}(a) + \arctan(a \tan(x)) / \sqrt{a^2 + ab}) * a^3 / (\sqrt{a^2 + ab} b^3) + \frac{1}{8}(8a^2 - 4ab + 3b^2)x/b^3 - \frac{1}{8}(4a \tan(x)^3 - 3b \tan(x)^3 + 4a \tan(x) - 5b \tan(x)) / ((\tan(x)^2 + 1)^2 b^2)$$

Mupad [B] (verification not implemented)

Time = 1.19 (sec) , antiderivative size = 1036, normalized size of antiderivative = 11.91

$$\int \frac{\cos^6(x)}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input `int(cos(x)^6/(a + b*cos(x)^2),x)`

output

```

- ((tan(x)^3*(4*a - 3*b))/(8*b^2) + (tan(x)*(4*a - 5*b))/(8*b^2))/(2*tan(x)
 )^2 + tan(x)^4 + 1) - (atan((63*a^4*tan(x))/(64*((63*a^4)/64 - (81*a^3*b)/
 256 + (27*a^2*b^2)/256 - (35*a^5)/(32*b) + (5*a^6)/(4*b^2)))) - (81*a^3*tan
 (x))/(256*((27*a^2*b)/256 - (81*a^3)/256 + (63*a^4)/(64*b) - (35*a^5)/(32*
 b^2) + (5*a^6)/(4*b^3))) - (35*a^5*tan(x))/(32*((63*a^4*b)/64 - (35*a^5)/3
 2 + (27*a^2*b^3)/256 - (81*a^3*b^2)/256 + (5*a^6)/(4*b))) + (5*a^6*tan(x))
 /(4*((5*a^6)/4 - (35*a^5*b)/32 + (27*a^2*b^4)/256 - (81*a^3*b^3)/256 + (63
 *a^4*b^2)/64)) + (27*a^2*tan(x))/(256*((27*a^2)/256 - (81*a^3)/(256*b) + (
 63*a^4)/(64*b^2) - (35*a^5)/(32*b^3) + (5*a^6)/(4*b^4)))*(a^2*8i - a*b*4i
 + b^2*3i)*i)/(8*b^3) - (atan((((-a^5*(a + b))^(1/2)*((-a^5*(a + b))^(1/
 2)*(((3*a^2*b^8)/2 - (a^3*b^7)/2 + 2*a^4*b^6)/(2*b^6) - (tan(x)*(256*a^2*b
 ^7 + 512*a^3*b^6)*(-a^5*(a + b))^(1/2))/(128*b^4*(a*b^3 + b^4)))))/(2*(a*b^
 3 + b^4)) - (tan(x)*(128*a^7 - 64*a^6*b + 9*a^3*b^4 - 24*a^4*b^3 + 64*a^5*
 b^2))/(64*b^4))*i)/(a*b^3 + b^4) - (((-a^5*(a + b))^(1/2)*(((a^5*(a + b))^(1/2)*(((3*a^2*b^8)/2 - (a^3*b^7)/2 + 2*a^4*b^6)/(2*b^6) + (tan(x)*(256*a^2*b^7 + 512*a^3*b^6)*(-a^5*(a + b))^(1/2))/(128*b^4*(a*b^3 + b^4)))))/(2*(a*b^3 + b^4)) + (tan(x)*(128*a^7 - 64*a^6*b + 9*a^3*b^4 - 24*a^4*b^3 + 64*a^5*b^2))/(64*b^4))*i)/(a*b^3 + b^4))/(((a^5*(a + b))^(1/2)*(((3*a^2*b^8)/2 - (a^3*b^7)/2 + 2*a^4*b^6)/(2*b^6) - (tan(x)*(256*a^2*b^7 + 512*a^3*b^6)*(-a^5*(a + b))^(1/2))/(128*b^4*(a*b^3 + b^4))))/(2*(a*b^3 + b^4))...

```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.70

$$\begin{aligned}
 & \int \frac{\cos^6(x)}{a + b \cos^2(x)} dx \\
 &= \frac{-8\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)-\sqrt{b}}{\sqrt{a}}\right) a^2 - 8\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right)+\sqrt{b}}{\sqrt{a}}\right) a^2 - 2 \cos(x) \sin(x)^3 a b^2 - }{ }
 \end{aligned}$$

input `int(cos(x)^6/(a+b*cos(x)^2),x)`

output
$$\begin{aligned} & \left(-8\sqrt{a}\sqrt{a+b}\arctan\left(\frac{\sqrt{a+b}\tan(x/2) - \sqrt{b}}{\sqrt{a}}\right)a^2 \right. \\ & \left. - 8\sqrt{a}\sqrt{a+b}\arctan\left(\frac{\sqrt{a+b}\tan(x/2) + \sqrt{b}}{\sqrt{a}}\right)a^2 \right. \\ & \left. - 2\cos(x)\sin(x)^3a^2b^2 - 2\cos(x)\sin(x)^3b^3 - 4\cos(x)\sin(x)a^2b^2 + \cos(x)\sin(x)a^2b^2 + 5\cos(x)\sin(x)b^3 + 8a^3x^2 + 4a^2b^2x - a^2b^2x^2 + 3b^3x \right) / (8b^3(a+b)) \end{aligned}$$

3.33 $\int \frac{\cos^4(x)}{a+b\cos^2(x)} dx$

Optimal result	260
Mathematica [A] (verified)	260
Rubi [A] (verified)	261
Maple [A] (verified)	263
Fricas [A] (verification not implemented)	263
Sympy [F(-1)]	264
Maxima [A] (verification not implemented)	264
Giac [A] (verification not implemented)	264
Mupad [B] (verification not implemented)	265
Reduce [B] (verification not implemented)	265

Optimal result

Integrand size = 15, antiderivative size = 60

$$\int \frac{\cos^4(x)}{a + b\cos^2(x)} dx = -\frac{(2a - b)x}{2b^2} - \frac{a^{3/2} \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{b^2\sqrt{a+b}} + \frac{\cos(x)\sin(x)}{2b}$$

output
$$-1/2*(2*a-b)*x/b^2-a^(3/2)*arctan((a+b)^(1/2)*cot(x)/a^(1/2))/b^2/(a+b)^(1/2)+1/2*cos(x)*sin(x)/b$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.87

$$\int \frac{\cos^4(x)}{a + b\cos^2(x)} dx = \frac{2(-2a + b)x + \frac{4a^{3/2} \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}} + b\sin(2x)}{4b^2}$$

input `Integrate[Cos[x]^4/(a + b*Cos[x]^2), x]`

output
$$(2*(-2*a + b)*x + (4*a^(3/2)*ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]])/Sqrt[a + b] + b*Sin[2*x])/ (4*b^2)$$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 74, normalized size of antiderivative = 1.23, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 3666, 372, 397, 216, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^4(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin\left(x + \frac{\pi}{2}\right)^4}{a + b \sin\left(x + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{3666} \\
 & - \int \frac{\cot^4(x)}{\left(\cot^2(x) + 1\right)^2 ((a + b) \cot^2(x) + a)} d \cot(x) \\
 & \quad \downarrow \text{372} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\int \frac{a - (a - b) \cot^2(x)}{(\cot^2(x) + 1)((a + b) \cot^2(x) + a)} d \cot(x)}{2b} \\
 & \quad \downarrow \text{397} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{2a^2 \int \frac{1}{(a + b) \cot^2(x) + a} d \cot(x)}{b} - \frac{(2a - b) \int \frac{1}{\cot^2(x) + 1} d \cot(x)}{b}}{2b} \\
 & \quad \downarrow \text{216} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{2a^2 \int \frac{1}{(a + b) \cot^2(x) + a} d \cot(x)}{b} - \frac{(2a - b) \arctan(\cot(x))}{b}}{2b} \\
 & \quad \downarrow \text{218} \\
 & \frac{\cot(x)}{2b (\cot^2(x) + 1)} - \frac{\frac{2a^{3/2} \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{b \sqrt{a+b}} - \frac{(2a - b) \arctan(\cot(x))}{b}}{2b}
 \end{aligned}$$

input `Int[Cos[x]^4/(a + b*Cos[x]^2),x]`

output
$$-1/2*(-(((2*a - b)*ArcTan[Cot[x]])/b) + (2*a^(3/2)*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(b*Sqrt[a + b]))/b + Cot[x]/(2*b*(1 + Cot[x]^2))$$

Definitions of rubi rules used

rule 216
$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(Rt[a, 2]*Rt[b, 2]))*\text{ArcTan}[Rt[b, 2]*(x/Rt[a, 2])], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \mid\mid \text{GtQ}[b, 0])$$

rule 218
$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(Rt[a/b, 2]/a)*\text{ArcTan}[x/Rt[a/b, 2]], \text{x}] /; \text{FreeQ}[\{a, b\}, \text{x}] \&& \text{PosQ}[a/b]$$

rule 372
$$\text{Int}[((e_)*(x_)^{(m_)}*((a_) + (b_.)*(x_)^2)^{(p_)}*((c_) + (d_.)*(x_)^2)^{(q_}), \text{x_Symbol}] \rightarrow \text{Simp}[-(a)*e^3*(e*x)^(m - 3)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*b*(b*c - a*d)*(p + 1))), \text{x}] + \text{Simp}[e^4/(2*b*(b*c - a*d)*(p + 1)) \text{Int}[(e*x)^(m - 4)*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*\text{Simp}[a*c*(m - 3) + (a*d*(m + 2*q - 1) + 2*b*c*(p + 1))*x^2, \text{x}], \text{x}] /; \text{FreeQ}[\{a, b, c, d, e, q\}, \text{x}] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{LtQ}[p, -1] \&& \text{GtQ}[m, 3] \&& \text{IntBinomialQ}[a, b, c, d, e, m, 2, p, q, \text{x}]$$

rule 397
$$\text{Int}[((e_) + (f_.)*(x_)^2)/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)), \text{x_Symbol}] \rightarrow \text{Simp}[(b*e - a*f)/(b*c - a*d) \text{Int}[1/(a + b*x^2), \text{x}], \text{x}] - \text{Simp}[(d*e - c*f)/(b*c - a*d) \text{Int}[1/(c + d*x^2), \text{x}], \text{x}] /; \text{FreeQ}[\{a, b, c, d, e, f\}, \text{x}]$$

rule 3042
$$\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$$

rule 3666
$$\text{Int}[\sin[(e_*) + (f_*)*(x_*)^{(m_)}*((a_) + (b_.)*\sin[(e_*) + (f_*)*(x_*)^2])^{(p_)}, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], \text{x}]\}, \text{Simp}[ff^{(m + 1)}/f \text{Subst}[\text{Int}[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1)), \text{x}], \text{x}, \text{Tan}[e + f*x]/ff], \text{x}]] /; \text{FreeQ}[\{a, b, e, f\}, \text{x}] \&& \text{IntegerQ}[m/2] \& \& \text{IntegerQ}[p]$$

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.98

method	result
default	$-\frac{b \tan(x)}{2(1+\tan(x)^2)} + \frac{(2a-b) \arctan(\tan(x))}{2} + \frac{a^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{b^2 \sqrt{(a+b)a}}$
risch	$-\frac{xa}{b^2} + \frac{x}{2b} - \frac{ie^{2ix}}{8b} + \frac{ie^{-2ix}}{8b} + \frac{\sqrt{-(a+b)a} a \ln\left(e^{2ix} - \frac{2i\sqrt{-(a+b)a}-2a-b}{b}\right)}{2(a+b)b^2} - \frac{\sqrt{-(a+b)a} a \ln\left(e^{2ix} + \frac{2i\sqrt{-(a+b)a}+2a+b}{b}\right)}{2(a+b)b^2}$

input `int(cos(x)^4/(a+b*cos(x)^2),x,method=_RETURNVERBOSE)`

output
$$-1/b^2*(-1/2*b*tan(x)/(1+tan(x)^2)+1/2*(2*a-b)*arctan(tan(x)))+a^2/b^2/((a+b)*a)^(1/2)*arctan(a*tan(x)/((a+b)*a)^(1/2))$$

Fricas [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 213, normalized size of antiderivative = 3.55

$$\int \frac{\cos^4(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{2 b \cos(x) \sin(x) + a \sqrt{-\frac{a}{a+b}} \log\left(\frac{(8 a^2 + 8 a b + b^2) \cos(x)^4 - 2 (4 a^2 + 3 a b) \cos(x)^2 - 4 ((2 a^2 + 3 a b + b^2) \cos(x)^3 - (a^2 + a b) \cos(x))}{b^2 \cos(x)^4 + 2 a b \cos(x)^2 + a^2}\right)}{4 b^2} \right]$$

input `integrate(cos(x)^4/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[1/4*(2*b*cos(x)*sin(x) + a*sqrt(-a/(a + b))*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 - 4*((2*a^2 + 3*a*b + b^2)*cos(x)^3 - (a^2 + a*b)*cos(x))*sqrt(-a/(a + b))*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) - 2*(2*a - b)*x]/b^2, 1/2*(b*cos(x)*sin(x) - a*sqrt(a/(a + b))*arctan(1/2*((2*a + b)*cos(x)^2 - a)*sqrt(a/(a + b)))/(a*cos(x)*sin(x))) - (2*a - b)*x]/b^2]$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(cos(x)**4/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.90

$$\int \frac{\cos^4(x)}{a + b \cos^2(x)} dx = \frac{a^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)ab^2}} - \frac{(2a-b)x}{2b^2} + \frac{\tan(x)}{2(b \tan(x)^2 + b)}$$

input `integrate(cos(x)^4/(a+b*cos(x)^2),x, algorithm="maxima")`

output `a^2*arctan(a*tan(x))/sqrt((a + b)*a)/(sqrt((a + b)*a)*b^2) - 1/2*(2*a - b)*x/b^2 + 1/2*tan(x)/(b*tan(x)^2 + b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.20

$$\int \frac{\cos^4(x)}{a + b \cos^2(x)} dx = \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab^2}}\right)\right) a^2}{\sqrt{a^2 + ab^2}} - \frac{(2a-b)x}{2b^2} + \frac{\tan(x)}{2(\tan(x)^2 + 1)b}$$

input `integrate(cos(x)^4/(a+b*cos(x)^2),x, algorithm="giac")`

output
$$\frac{(\pi * \text{floor}(x/\pi + 1/2) * \text{sgn}(a) + \arctan(a * \tan(x) / \sqrt{a^2 + a * b})) * a^2 / (\sqrt{a^2 + a * b} * b^2) - 1/2 * (2 * a - b) * x / b^2 + 1/2 * \tan(x) / ((\tan(x)^2 + 1) * b)}$$

Mupad [B] (verification not implemented)

Time = 1.15 (sec), antiderivative size = 291, normalized size of antiderivative = 4.85

$$\int \frac{\cos^4(x)}{a + b \cos^2(x)} dx =$$

$$\frac{2 a^2 \operatorname{atan}\left(\frac{\sin(x)}{\cos(x)}\right) - b^2 \operatorname{atan}\left(\frac{\sin(x)}{\cos(x)}\right) - \frac{b^2 \sin(2x)}{2} + a b \operatorname{atan}\left(\frac{\sin(x)}{\cos(x)}\right) - \frac{a b \sin(2x)}{2} + \operatorname{atan}\left(\frac{a \sin(x) (-a^4 - b a^3)^3}{b^3}\right)}{a^3}$$

input $\int \cos(x)^4 / (a + b \cos(x)^2) dx$

output
$$-(2*a^2*atan(sin(x)/cos(x)) - b^2*atan(sin(x)/cos(x)) + atan((a*sin(x)*(-a^3*b - a^4)^(3/2)*8i + b*sin(x)*(-a^3*b - a^4)^(3/2)*4i + a^5*sin(x)*(-a^3*b - a^4)^(1/2)*8i - a^2*b^3*sin(x)*(-a^3*b - a^4)^(1/2)*2i + a^3*b^2*sin(x)*(-a^3*b - a^4)^(1/2)*1i + a*b^4*sin(x)*(-a^3*b - a^4)^(1/2)*1i + a^4*b*sin(x)*(-a^3*b - a^4)^(1/2)*12i)/(a^3*b^4*cos(x) - a^2*b^5*cos(x) + 5*a^4*b^3*cos(x) + 3*a^5*b^2*cos(x)))*(-a^3*b - a^4)^(1/2)*2i - (b^2*sin(2*x))/2 + a*b*atan(sin(x)/cos(x)) - (a*b*sin(2*x))/2)/(2*a*b^2 + 2*b^3)$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec), antiderivative size = 127, normalized size of antiderivative = 2.12

$$\int \frac{\cos^4(x)}{a + b \cos^2(x)} dx$$

$$= \frac{2\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) a + 2\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) a + \cos(x)^2 abx + \cos(x)^2 b^2 x}{2b^2(a+b)}$$

input $\int \cos(x)^4 / (a+b \cos(x)^2) dx$

```
output (2*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*a +
2*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*a + c
cos(x)**2*a*b*x + cos(x)**2*b**2*x + cos(x)*sin(x)*a*b + cos(x)*sin(x)*b**2
+ sin(x)**2*a*b*x + sin(x)**2*b**2*x - 2*a**2*x - 2*a*b*x)/(2*b**2*(a + b
))
```

3.34 $\int \frac{\cos^2(x)}{a+b\cos^2(x)} dx$

Optimal result	267
Mathematica [A] (verified)	267
Rubi [A] (verified)	268
Maple [A] (verified)	269
Fricas [A] (verification not implemented)	270
Sympy [F(-1)]	270
Maxima [A] (verification not implemented)	271
Giac [A] (verification not implemented)	271
Mupad [B] (verification not implemented)	272
Reduce [B] (verification not implemented)	272

Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \frac{\cos^2(x)}{a + b\cos^2(x)} dx = \frac{x}{b} + \frac{\sqrt{a} \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{b\sqrt{a+b}}$$

output x/b+a^(1/2)*arctan((a+b)^(1/2)*cot(x)/a^(1/2))/b/(a+b)^(1/2)

Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int \frac{\cos^2(x)}{a + b\cos^2(x)} dx = \frac{x - \frac{\sqrt{a} \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}}}{b}$$

input Integrate[Cos[x]^2/(a + b*Cos[x]^2), x]

output (x - (Sqrt[a]*ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]])/Sqrt[a + b])/b

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 3650, 3042, 3660, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos^2(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})^2}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3650} \\
 & \frac{x}{b} - \frac{a \int \frac{1}{b \cos^2(x) + a} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x}{b} - \frac{a \int \frac{1}{b \sin(x + \frac{\pi}{2})^2 + a} dx}{b} \\
 & \quad \downarrow \text{3660} \\
 & \frac{a \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{b} + \frac{x}{b} \\
 & \quad \downarrow \text{218} \\
 & \frac{\sqrt{a} \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{b \sqrt{a+b}} + \frac{x}{b}
 \end{aligned}$$

input `Int[Cos[x]^2/(a + b*Cos[x]^2),x]`

output `x/b + (Sqrt[a]*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(b*Sqrt[a + b])`

Definitions of rubi rules used

rule 218 $\text{Int}[(a_+ + b_-)(x_-)^2]^{-1}, x_{\text{Symbol}} \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_-, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3650 $\text{Int}[(A_- + B_-) \cdot \sin[(e_- + f_-)(x_-)]^2] / ((a_+ + b_-) \cdot \sin[(e_- + f_-)(x_-)]^2), x_{\text{Symbol}} \rightarrow \text{Simp}[B \cdot (x/b), x] + \text{Simp}[(A \cdot b - a \cdot B)/b \cdot \text{Int}[1/(a + b \cdot \sin[e + f \cdot x]^2), x], x] /; \text{FreeQ}[\{a, b, e, f, A, B\}, x]$

rule 3660 $\text{Int}[(a_+ + b_-) \cdot \sin[(e_- + f_-)(x_-)]^2]^{-1}, x_{\text{Symbol}} \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\tan[e + f \cdot x], x]\}, \text{Simp}[\text{ff}/f \cdot \text{Subst}[\text{Int}[1/(a + (a + b) \cdot \text{ff}^2 \cdot x^2), x], \tan[e + f \cdot x]/\text{ff}], x]] /; \text{FreeQ}[\{a, b, e, f\}, x]$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\arctan(\tan(x))}{b} - \frac{a \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{b \sqrt{(a+b)a}}$	34
risch	$\frac{x}{b} + \frac{\sqrt{-(a+b)a} \ln\left(e^{2ix} + \frac{2i\sqrt{-(a+b)a}+2a+b}{b}\right)}{2(a+b)b} - \frac{\sqrt{-(a+b)a} \ln\left(e^{2ix} - \frac{2i\sqrt{-(a+b)a}-2a-b}{b}\right)}{2(a+b)b}$	100

input $\text{int}(\cos(x)^2/(a+b \cdot \cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output $1/b \cdot \arctan(\tan(x)) - 1/b \cdot a / ((a+b) \cdot a)^{(1/2)} \cdot \arctan(a \cdot \tan(x)) / ((a+b) \cdot a)^{(1/2)}$

Fricas [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 183, normalized size of antiderivative = 4.82

$$\int \frac{\cos^2(x)}{a + b \cos^2(x)} dx$$

$$= \left[\frac{\sqrt{-\frac{a}{a+b}} \log \left(\frac{(8a^2+8ab+b^2)\cos(x)^4 - 2(4a^2+3ab)\cos(x)^2 + 4((2a^2+3ab+b^2)\cos(x)^3 - (a^2+ab)\cos(x))\sqrt{-\frac{a}{a+b}}\sin(x) + a^2}{b^2\cos(x)^4 + 2ab\cos(x)^2 + a^2} \right) + 4}{4b} \right]$$

input `integrate(cos(x)^2/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[1/4*(sqrt(-a/(a + b))*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 + 4*((2*a^2 + 3*a*b + b^2)*cos(x)^3 - (a^2 + a*b)*cos(x)))*sqrt(-a/(a + b))*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) + 4*x)/b, 1/2*(sqrt(a/(a + b))*arctan(1/2*((2*a + b)*cos(x)^2 - a)*sqrt(a/(a + b)))/(a*cos(x)*sin(x))) + 2*x)/b]`

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(cos(x)**2/(a+b*cos(x)**2),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{\cos^2(x)}{a + b \cos^2(x)} dx = -\frac{a \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)ab}} + \frac{x}{b}$$

input `integrate(cos(x)^2/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-a*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*b) + x/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

$$\int \frac{\cos^2(x)}{a + b \cos^2(x)} dx = -\frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2+ab}}\right)\right)a}{\sqrt{a^2 + ab}} + \frac{x}{b}$$

input `integrate(cos(x)^2/(a+b*cos(x)^2),x, algorithm="giac")`

output `-(pi*floor(x/pi + 1/2)*sgn(a) + arctan(a*tan(x)/sqrt(a^2 + a*b)))*a/(sqrt(a^2 + a*b)*b) + x/b`

Mupad [B] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 425, normalized size of antiderivative = 11.18

$$\int \frac{\cos^2(x)}{a + b \cos^2(x)} dx = \frac{x}{b} - \frac{\text{atan}\left(\frac{\sqrt{-a(a+b)}}{b^2+a^2}\right) \sqrt{-a(a+b)}}{b^2+a^2}$$

$$= \frac{\frac{b^2+a^2}{2a^3\tan(x)-\frac{(2a^2b^2-\frac{\tan(x)(16a^3b^2+8a^2b^3)\sqrt{-a(a+b)}}{4(b^2+a^2)})\sqrt{-a(a+b)}}{2(b^2+a^2)}}{\sqrt{-a(a+b)}} + \frac{\frac{b^2+a^2}{2a^3\tan(x)+\frac{(2a^2b^2+\frac{\tan(x)(16a^3b^2+8a^2b^3)\sqrt{-a(a+b)}}{4(b^2+a^2)})\sqrt{-a(a+b)}}{2(b^2+a^2)}}{\sqrt{-a(a+b)}} - \frac{\frac{b^2+a^2}{2a^3\tan(x)-\frac{(2a^2b^2-\frac{\tan(x)(16a^3b^2+8a^2b^3)\sqrt{-a(a+b)}}{4(b^2+a^2)})\sqrt{-a(a+b)}}{2(b^2+a^2)}}{\sqrt{-a(a+b)}} - \frac{\frac{b^2+a^2}{2a^3\tan(x)+\frac{(2a^2b^2+\frac{\tan(x)(16a^3b^2+8a^2b^3)\sqrt{-a(a+b)}}{4(b^2+a^2)})\sqrt{-a(a+b)}}{2(b^2+a^2)}}{\sqrt{-a(a+b)}}}{b^2+a^2}$$

input `int(cos(x)^2/(a + b*cos(x)^2),x)`

output

$$\begin{aligned} & x/b - (\text{atan}(((2*a^3*tan(x) - ((2*a^2*b^2 - (\tan(x)*(8*a^2*b^3 + 16*a^3*b^2)*(-a*(a + b))^(1/2))/(4*(a*b + b^2)))*(-a*(a + b))^(1/2)/(2*(a*b + b^2))) * (-a*(a + b))^(1/2)*1i)/(a*b + b^2) + ((2*a^3*tan(x) + ((2*a^2*b^2 - (\tan(x)*(8*a^2*b^3 + 16*a^3*b^2)*(-a*(a + b))^(1/2))/(4*(a*b + b^2)))*(-a*(a + b))^(1/2)/(2*(a*b + b^2))) * (-a*(a + b))^(1/2)*1i)/(a*b + b^2))/(((2*a^3*tan(x) - ((2*a^2*b^2 - (\tan(x)*(8*a^2*b^3 + 16*a^3*b^2)*(-a*(a + b))^(1/2))/(4*(a*b + b^2)))*(-a*(a + b))^(1/2)/(2*(a*b + b^2))) * (-a*(a + b))^(1/2)/(4*(a*b + b^2)))*(-a*(a + b))^(1/2)/(a*b + b^2) - ((2*a^3*tan(x) + ((2*a^2*b^2 + (\tan(x)*(8*a^2*b^3 + 16*a^3*b^2)*(-a*(a + b))^(1/2))/(4*(a*b + b^2)))*(-a*(a + b))^(1/2)/(2*(a*b + b^2)))*(-a*(a + b))^(1/2)/(a*b + b^2)) * (-a*(a + b))^(1/2)*1i)/(a*b + b^2))) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.84

$$\begin{aligned} & \int \frac{\cos^2(x)}{a + b \cos^2(x)} dx \\ &= \frac{-\sqrt{a}\sqrt{a+b}\text{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})-\sqrt{b}}{\sqrt{a}}\right) - \sqrt{a}\sqrt{a+b}\text{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})+\sqrt{b}}{\sqrt{a}}\right) + ax + bx}{b(a+b)} \end{aligned}$$

input `int(cos(x)^2/(a+b*cos(x)^2),x)`

output
$$\left(-\frac{\sqrt{a} \sqrt{a+b} \arctan\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\sqrt{a} \sqrt{a+b} \arctan\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) + \sqrt{b}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{a x + b x}{b(a+b)} \right)$$

3.35 $\int \frac{1}{a+b \cos^2(x)} dx$

Optimal result	274
Mathematica [A] (verified)	274
Rubi [A] (verified)	275
Maple [A] (verified)	276
Fricas [B] (verification not implemented)	276
Sympy [B] (verification not implemented)	277
Maxima [A] (verification not implemented)	278
Giac [A] (verification not implemented)	278
Mupad [B] (verification not implemented)	278
Reduce [B] (verification not implemented)	279

Optimal result

Integrand size = 10, antiderivative size = 30

$$\int \frac{1}{a + b \cos^2(x)} dx = -\frac{\arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{a+b}}$$

output `-arctan((a+b)^(1/2)*cot(x)/a^(1/2))/a^(1/2)/(a+b)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.97

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

input `Integrate[(a + b*Cos[x]^2)^(-1), x]`

output `ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]]/(Sqrt[a]*Sqrt[a + b])`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 3660, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{a + b \sin(x + \frac{\pi}{2})^2} dx \\
 & \quad \downarrow \text{3660} \\
 & - \int \frac{1}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \text{218} \\
 & - \frac{\arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{\sqrt{a} \sqrt{a+b}}
 \end{aligned}$$

input `Int[(a + b*Cos[x]^2)^(-1), x]`

output `-(ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]]/(Sqrt[a]*Sqrt[a + b]))`

Definitions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3660

```
Int[((a_) + (b_)*sin[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] :> With[{ff =
FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[1/(a + (a + b)*ff^2*x^
2), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]
```

Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}}$	21
risch	$-\frac{\ln\left(\frac{e^{2ix} + 2ia^2 + 2iab + 2a\sqrt{-a^2-ab} + b\sqrt{-a^2-ab}}{b\sqrt{-a^2-ab}}\right)}{2\sqrt{-a^2-ab}} + \frac{\ln\left(\frac{e^{2ix} - 2ia^2 + 2iab - 2a\sqrt{-a^2-ab} - b\sqrt{-a^2-ab}}{b\sqrt{-a^2-ab}}\right)}{2\sqrt{-a^2-ab}}$	160

input `int(1/(a+b*cos(x)^2), x, method=_RETURNVERBOSE)`

output `1/((a+b)*a)^(1/2)*arctan(a*tan(x)/((a+b)*a)^(1/2))`

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(22) = 44$.

Time = 0.15 (sec) , antiderivative size = 163, normalized size of antiderivative = 5.43

$$\begin{aligned} & \int \frac{1}{a + b \cos^2(x)} dx \\ &= \left[-\frac{\sqrt{-a^2 - ab} \log \left(\frac{(8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 + 4((2a+b)\cos(x)^3 - a\cos(x))\sqrt{-a^2-ab}\sin(x) + a^2}{b^2 \cos(x)^4 + 2ab \cos(x)^2 + a^2} \right)}{4(a^2 + ab)}, \right. \\ & \quad \left. - \frac{\arctan \left(\frac{(2a+b)\cos(x)^2 - a}{2\sqrt{a^2+ab}\cos(x)\sin(x)} \right)}{2\sqrt{a^2+ab}} \right] \end{aligned}$$

input `integrate(1/(a+b*cos(x)^2), x, algorithm="fricas")`

output

```
[-1/4*sqrt(-a^2 - a*b)*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 + 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2))/(a^2 + a*b), -1/2*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))/sqrt(a^2 + a*b)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10924 vs. $2(29) = 58$.

Time = 18.45 (sec) , antiderivative size = 10924, normalized size of antiderivative = 364.13

$$\int \frac{1}{a + b \cos^2(x)} dx = \text{Too large to display}$$

input

```
integrate(1/(a+b*cos(x)**2),x)
```

output

```
Piecewise((zoo*tan(x/2)/(tan(x/2)**2 - 1), Eq(a, 0) & Eq(b, 0)), (-tan(x/2)/(2*b) + 1/(2*b*tan(x/2)), Eq(a, -b)), (-2*tan(x/2)/(b*(tan(x/2)**2 - 1)), Eq(a, 0)), (a**3*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(-sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan(x/2))/(2*a**4*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b))) - 10*a**3*b*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a**3*sqrt(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a*b**2*sqrt(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - a**3*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan(x/2))/(2*a**4*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a**3*sqrt(-a*b)*sqrt(-a/(a + b) + b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(-a/(a + b) + b/(a + b) + 2*sqrt(-a*b)/(a + b)...)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}}$$

input `integrate(1/(a+b*cos(x)^2),x, algorithm="maxima")`

output `arctan(a*tan(x)/sqrt((a + b)*a))/sqrt((a + b)*a)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.23

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\pi \left\lfloor \frac{x}{\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2+ab}}\right)}{\sqrt{a^2+ab}}$$

input `integrate(1/(a+b*cos(x)^2),x, algorithm="giac")`

output `(pi*floor(x/pi + 1/2)*sgn(a) + arctan(a*tan(x)/sqrt(a^2 + a*b)))/sqrt(a^2 + a*b)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\operatorname{atan}\left(\frac{a \tan(x)}{\sqrt{a^2+ba}}\right)}{\sqrt{a^2+ba}}$$

input `int(1/(a + b*cos(x)^2),x)`

output `atan((a*tan(x))/(a*b + a^2)^(1/2))/(a*b + a^2)^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.80

$$\int \frac{1}{a + b \cos^2(x)} dx = \frac{\sqrt{a} \sqrt{a+b} \left(\operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) - \sqrt{b}}{\sqrt{a}}\right) + \operatorname{atan}\left(\frac{\sqrt{a+b} \tan(\frac{x}{2}) + \sqrt{b}}{\sqrt{a}}\right) \right)}{a(a+b)}$$

input `int(1/(a+b*cos(x)^2),x)`

output `(sqrt(a)*sqrt(a + b)*(atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a)) + atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))))/(a*(a + b))`

3.36 $\int \frac{\sec^2(x)}{a+b\cos^2(x)} dx$

Optimal result	280
Mathematica [A] (verified)	280
Rubi [A] (verified)	281
Maple [A] (verified)	282
Fricas [B] (verification not implemented)	283
Sympy [F]	283
Maxima [A] (verification not implemented)	284
Giac [A] (verification not implemented)	284
Mupad [B] (verification not implemented)	284
Reduce [B] (verification not implemented)	285

Optimal result

Integrand size = 15, antiderivative size = 37

$$\int \frac{\sec^2(x)}{a + b\cos^2(x)} dx = \frac{b \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{a^{3/2}\sqrt{a+b}} + \frac{\tan(x)}{a}$$

output `b*arctan((a+b)^(1/2)*cot(x)/a^(1/2))/a^(3/2)/(a+b)^(1/2)+tan(x)/a`

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int \frac{\sec^2(x)}{a + b\cos^2(x)} dx = -\frac{b \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{a^{3/2}\sqrt{a+b}} + \frac{\tan(x)}{a}$$

input `Integrate[Sec[x]^2/(a + b*Cos[x]^2),x]`

output `-((b*ArcTan[(Sqrt[a]*Tan[x])/Sqrt[a + b]])/(a^(3/2)*Sqrt[a + b])) + Tan[x]/a`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3666, 359, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^2(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x + \frac{\pi}{2})^2 \left(a + b \sin(x + \frac{\pi}{2})^2\right)} dx \\
 & \quad \downarrow \text{3666} \\
 & - \int \frac{(\cot^2(x) + 1) \tan^2(x)}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \text{359} \\
 & \frac{b \int \frac{1}{(a+b) \cot^2(x) + a} d \cot(x)}{a} + \frac{\tan(x)}{a} \\
 & \quad \downarrow \text{218} \\
 & \frac{b \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{a^{3/2} \sqrt{a+b}} + \frac{\tan(x)}{a}
 \end{aligned}$$

input `Int[Sec[x]^2/(a + b*Cos[x]^2),x]`

output `(b*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(a^(3/2)*Sqrt[a + b]) + Tan[x]/a`

Definitions of rubi rules used

rule 218 $\text{Int}[(a_ + b_)(x_)^2]^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{PosQ}[a/b]$

rule 359 $\text{Int}[(e_)(m_)(a_ + b_)(c_ + d_)(x_)^2]^{(p_)}, \text{x_Symbol}] \rightarrow \text{Simp}[c*(e*x)^{(m+1)}((a+b*x^2)^{(p+1)}/(a*e^{(m+1)})), x] + \text{Simp}[(a*d*(m+1) - b*c*(m+2*p+3))/(a*e^{2*(m+1)}) \text{Int}[(e*x)^{(m+2)}*(a+b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \& \text{NeQ}[b*c - a*d, 0] \& \text{LtQ}[m, -1] \& \text{ILtQ}[p, -1]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3666 $\text{Int}[\sin[(e_ + f_)(x_)]^{(m_)}((a_ + b_)*\sin[(e_ + f_)(x_)]^2)^{(p_)}, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Simp}[ff^{(m+1)}/f \text{Subst}[\text{Int}[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^{(m/2+p+1)}), x, \text{Tan}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f\}, x] \& \text{IntegerQ}[m/2] \& \text{IntegerQ}[p]$

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\tan(x)}{a} - \frac{b \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{a \sqrt{(a+b)a}}$	33
risch	$\frac{2i}{(e^{2ix}+1)a} - \frac{b \ln\left(e^{2ix} + \frac{-2ia^2 - 2iab + 2a\sqrt{-a^2-ab} + b\sqrt{-a^2-ab}}{\sqrt{-a^2-ab}b}\right)}{2\sqrt{-a^2-ab}a} + \frac{b \ln\left(e^{2ix} + \frac{2ia^2 + 2iab + 2a\sqrt{-a^2-ab} + b\sqrt{-a^2-ab}}{b\sqrt{-a^2-ab}}\right)}{2\sqrt{-a^2-ab}a}$	181

input $\text{int}(\sec(x)^2/(a+b*\cos(x)^2), x, \text{method} = \text{_RETURNVERBOSE})$

output $\tan(x)/a - b/a/((a+b)*a)^{(1/2)} * \text{arctan}(a*\tan(x)/((a+b)*a)^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(29) = 58.

Time = 0.18 (sec) , antiderivative size = 216, normalized size of antiderivative = 5.84

$$\int \frac{\sec^2(x)}{a + b \cos^2(x)} dx$$

$$= \left[-\frac{\sqrt{-a^2 - ab} \cos(x) \log \left(\frac{(8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 - 4((2a+b)\cos(x)^3 - a\cos(x))\sqrt{-a^2-ab}\sin(x)+a^2}{b^2\cos(x)^4+2ab\cos(x)^2+a^2} \right)}{4(a^3 + a^2b) \cos(x)} \right]$$

input `integrate(sec(x)^2/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[-1/4*(sqrt(-a^2 - a*b)*b*cos(x)*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 - 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) - 4*(a^2 + a*b)*sin(x)/((a^3 + a^2*b)*cos(x)), 1/2*(sqrt(a^2 + a*b)*b*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))*cos(x) + 2*(a^2 + a*b)*sin(x))/((a^3 + a^2*b)*cos(x))]`

Sympy [F]

$$\int \frac{\sec^2(x)}{a + b \cos^2(x)} dx = \int \frac{\sec^2(x)}{a + b \cos^2(x)} dx$$

input `integrate(sec(x)**2/(a+b*cos(x)**2),x)`

output `Integral(sec(x)**2/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

$$\int \frac{\sec^2(x)}{a + b \cos^2(x)} dx = -\frac{b \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a a}} + \frac{\tan(x)}{a}$$

input `integrate(sec(x)^2/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-b*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*a) + tan(x)/a`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.97

$$\int \frac{\sec^2(x)}{a + b \cos^2(x)} dx = -\frac{b \arctan\left(\frac{a \tan(x)}{\sqrt{a^2+ab}}\right)}{\sqrt{a^2+aba}} + \frac{\tan(x)}{a}$$

input `integrate(sec(x)^2/(a+b*cos(x)^2),x, algorithm="giac")`

output `-b*arctan(a*tan(x)/sqrt(a^2 + a*b))/(sqrt(a^2 + a*b)*a) + tan(x)/a`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.81

$$\int \frac{\sec^2(x)}{a + b \cos^2(x)} dx = \frac{\tan(x)}{a} - \frac{b \operatorname{atan}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{a^{3/2} \sqrt{a+b}}$$

input `int(1/(cos(x)^2*(a + b*cos(x)^2)),x)`

output `tan(x)/a - (b*atan((a^(1/2)*tan(x))/(a + b)^(1/2)))/(a^(3/2)*(a + b)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.30

$$\int \frac{\sec^2(x)}{a + b \cos^2(x)} dx \\ = \frac{-\sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) - \sqrt{b}}{\sqrt{a}}\right) \cos(x) b - \sqrt{a} \sqrt{a+b} \operatorname{atan}\left(\frac{\sqrt{a+b} \tan\left(\frac{x}{2}\right) + \sqrt{b}}{\sqrt{a}}\right) \cos(x) b + \sin(x) a^2 + s}{\cos(x) a^2 (a+b)}$$

input `int(sec(x)^2/(a+b*cos(x)^2),x)`

output `(- sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*cos(x)*b - sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*cos(x)*b + sin(x)*a**2 + sin(x)*a*b)/(cos(x)*a**2*(a + b))`

3.37 $\int \frac{\sec^4(x)}{a+b\cos^2(x)} dx$

Optimal result	286
Mathematica [A] (verified)	286
Rubi [A] (verified)	287
Maple [A] (verified)	288
Fricas [B] (verification not implemented)	289
Sympy [F]	289
Maxima [A] (verification not implemented)	290
Giac [A] (verification not implemented)	290
Mupad [B] (verification not implemented)	290
Reduce [B] (verification not implemented)	291

Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \frac{\sec^4(x)}{a+b\cos^2(x)} dx = -\frac{b^2 \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{a^{5/2}\sqrt{a+b}} + \frac{(a-b)\tan(x)}{a^2} + \frac{\tan^3(x)}{3a}$$

output
$$-\frac{b^2 \operatorname{arctan}\left(\frac{(a+b)^{1/2} \cot(x)}{\sqrt{a}}\right)}{a^{5/2} \sqrt{a+b}} + \frac{(a-b) \tan(x)}{a^2} + \frac{\tan^3(x)}{3a}$$

Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98

$$\int \frac{\sec^4(x)}{a+b\cos^2(x)} dx = \frac{b^2 \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{a^{5/2}\sqrt{a+b}} + \frac{(2a-3b+a\sec^2(x))\tan(x)}{3a^2}$$

input `Integrate[Sec[x]^4/(a + b*Cos[x]^2), x]`

output
$$\left(\frac{b^2 \operatorname{ArcTan}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{a^{5/2} \sqrt{a+b}} + \frac{(2a-3b+a \sec^2(x)) \tan(x)}{3a^2} \right)$$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3666, 364, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^4(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sin(x + \frac{\pi}{2})^4 (a + b \sin(x + \frac{\pi}{2})^2)} dx \\
 & \quad \downarrow \text{3666} \\
 & - \int \frac{(\cot^2(x) + 1)^2 \tan^4(x)}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \text{364} \\
 & - \int \left(\frac{\tan^4(x)}{a} + \frac{(a - b) \tan^2(x)}{a^2} + \frac{b^2}{a^2 ((a + b) \cot^2(x) + a)} \right) d \cot(x) \\
 & \quad \downarrow \text{2009} \\
 & - \frac{b^2 \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{a^{5/2} \sqrt{a+b}} + \frac{(a - b) \tan(x)}{a^2} + \frac{\tan^3(x)}{3a}
 \end{aligned}$$

input `Int[Sec[x]^4/(a + b*Cos[x]^2),x]`

output `-((b^2*ArcTan[(Sqrt[a + b]*Cot[x])/Sqrt[a]])/(a^(5/2)*Sqrt[a + b])) + ((a - b)*Tan[x])/a^2 + Tan[x]^3/(3*a)`

Definitions of rubi rules used

rule 364 $\text{Int}[((\text{e}_.) * (\text{x}_.)^{\text{m}_.}) * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^2)^{\text{p}_.}) / ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^2), \text{x}_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(\text{e}_* \text{x})^{\text{m}} * ((\text{a} + \text{b}_* \text{x}^2)^{\text{p}} / (\text{c} + \text{d}_* \text{x}^2)), \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{m}\}, \text{x}] \& \text{NeQ}[\text{b}_* \text{c} - \text{a}_* \text{d}, 0] \& \text{IGtQ}[\text{p}, 0] \& (\text{IntegerQ}[\text{m}] \text{ || } \text{IGtQ}[2*(\text{m} + 1), 0] \text{ || } \text{!RationalQ}[\text{m}])$

rule 2009 $\text{Int}[\text{u}_., \text{x}_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[\text{u}, \text{x}], \text{x}] /; \text{SumQ}[\text{u}]$

rule 3042 $\text{Int}[\text{u}_., \text{x}_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3666 $\text{Int}[\sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)]^{\text{m}_.} * ((\text{a}_.) + (\text{b}_.) * \sin[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)]^2)^{\text{p}_.}, \text{x}_{\text{Symbol}}] \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Tan}[\text{e} + \text{f}_* \text{x}], \text{x}]\}, \text{Simp}[\text{ff}^{\text{m} + 1} / \text{f} \text{ Subst}[\text{Int}[\text{x}^{\text{m}} * ((\text{a} + (\text{a} + \text{b}) * \text{ff}^2 * \text{x}^2)^{\text{p}} / (1 + \text{ff}^2 * \text{x}^2)^{(\text{m}/2 + \text{p} + 1)}), \text{x}], \text{x}, \text{Tan}[\text{e} + \text{f}_* \text{x}] / \text{ff}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}\}, \text{x}] \& \text{IntegerQ}[\text{m}/2] \& \text{IntegerQ}[\text{p}]$

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.88

method	result
default	$\frac{\frac{a \tan(x)^3}{3} + \tan(x)a - \tan(x)b}{a^2} + \frac{b^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{a^2 \sqrt{(a+b)a}}$
risch	$-\frac{2i(3e^{4ix}b - 6ae^{2ix} + 6e^{2ix}b - 2a + 3b)}{3(e^{2ix} + 1)^3 a^2} - \frac{b^2 \ln\left(\frac{e^{2ix} + 2ia^2 + 2iab + 2a\sqrt{-a^2 - ab} + b\sqrt{-a^2 - ab}}{b\sqrt{-a^2 - ab}}\right)}{2\sqrt{-a^2 - ab} a^2} + \frac{b^2 \ln\left(\frac{e^{2ix} - 2ia^2 + 2iab - 2a\sqrt{-a^2 - ab}}{b\sqrt{-a^2 - ab}}\right)}{2\sqrt{-a^2 - ab} a^2}$

input $\text{int}(\sec(x)^4 / (a + b * \cos(x)^2), x, \text{method}=\text{_RETURNVERBOSE})$

output $\frac{1}{a^2} \cdot \frac{1}{3} \cdot \frac{a \cdot \tan(x)^3 + \tan(x) \cdot a - \tan(x) \cdot b + b^2 / a^2}{((a + b) \cdot a)^{(1/2)}} \cdot \frac{\arctan(a \cdot \tan(x))}{((a + b) \cdot a)^{(1/2)}}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. $2(46) = 92$.

Time = 0.17 (sec) , antiderivative size = 276, normalized size of antiderivative = 4.93

$$\int \frac{\sec^4(x)}{a + b \cos^2(x)} dx$$

$$= \left[-\frac{3 \sqrt{-a^2 - ab} b^2 \cos(x)^3 \log \left(\frac{(8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 + 4((2a+b) \cos(x)^3 - a \cos(x)) \sqrt{-a^2 - ab} \sin(x) + a^2}{b^2 \cos(x)^4 + 2ab \cos(x)^2 + a^2} \right)}{12(a^4 + a^3b) \cos(x)^3} \right.$$

$$- \left. \frac{3 \sqrt{a^2 + ab} b^2 \arctan \left(\frac{(2a+b) \cos(x)^2 - a}{2\sqrt{a^2 + ab} \cos(x) \sin(x)} \right) \cos(x)^3 - 2(a^3 + a^2b + (2a^3 - a^2b - 3ab^2) \cos(x)^2) \sin(x)}{6(a^4 + a^3b) \cos(x)^3} \right]$$

input `integrate(sec(x)^4/(a+b*cos(x)^2),x, algorithm="fricas")`

output `[-1/12*(3*sqrt(-a^2 - a*b)*b^2*cos(x)^3*log((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 + 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) - 4*(a^3 + a^2*b + (2*a^3 - a^2*b - 3*a*b^2)*cos(x)^2)*sin(x))/((a^4 + a^3*b)*cos(x)^3), -1/6*(3*sqrt(a^2 + a*b)*b^2*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))*cos(x)^3 - 2*(a^3 + a^2*b + (2*a^3 - a^2*b - 3*a*b^2)*cos(x)^2)*sin(x))/((a^4 + a^3*b)*cos(x)^3)]`

Sympy [F]

$$\int \frac{\sec^4(x)}{a + b \cos^2(x)} dx = \int \frac{\sec^4(x)}{a + b \cos^2(x)} dx$$

input `integrate(sec(x)**4/(a+b*cos(x)**2),x)`

output `Integral(sec(x)**4/(a + b*cos(x)**2), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.86

$$\int \frac{\sec^4(x)}{a + b \cos^2(x)} dx = \frac{b^2 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}a^2} + \frac{a \tan(x)^3 + 3(a-b)\tan(x)}{3a^2}$$

input `integrate(sec(x)^4/(a+b*cos(x)^2),x, algorithm="maxima")`

output $b^2 \arctan(a \tan(x) / \sqrt{(a+b)a}) / (\sqrt{(a+b)a} * a^2) + 1/3 * (a * \tan(x)^3 + 3 * (a - b) * \tan(x)) / a^2$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.27

$$\int \frac{\sec^4(x)}{a + b \cos^2(x)} dx = \frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right) b^2}{\sqrt{a^2 + ab} a^2} + \frac{a^2 \tan(x)^3 + 3 a^2 \tan(x) - 3 a b \tan(x)}{3 a^3}$$

input `integrate(sec(x)^4/(a+b*cos(x)^2),x, algorithm="giac")`

output $(\pi * \operatorname{floor}(x/\pi + 1/2) * \operatorname{sgn}(a) + \arctan(a * \tan(x) / \sqrt{a^2 + a * b})) * b^2 / (\sqrt{(a^2 + a * b) * a^2}) + 1/3 * (a^2 * \tan(x)^3 + 3 * a^2 * \tan(x) - 3 * a * b * \tan(x)) / a^3$

Mupad [B] (verification not implemented)

Time = 0.97 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int \frac{\sec^4(x)}{a + b \cos^2(x)} dx = \frac{\tan(x)^3}{3a} - \tan(x) \left(\frac{a+b}{a^2} - \frac{2}{a} \right) + \frac{b^2 \operatorname{atan}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}}\right)}{a^{5/2} \sqrt{a+b}}$$

input `int(1/((cos(x)^4*(a + b*cos(x)^2)),x)`

output
$$\frac{\tan(x)^3/(3*a) - \tan(x)*((a + b)/a^2 - 2/a) + (b^2*\text{atan}((a^{1/2}*\tan(x))/(a + b)^{1/2}))/((a^{5/2}*(a + b)^{1/2}))}{(a^{5/2}*(a + b)^{1/2})}$$

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 211, normalized size of antiderivative = 3.77

input `int(sec(x)^4/(a+b*cos(x)^2),x)`

```

output (3*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*cos(x)*sin(x)**2*b**2 - 3*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*cos(x)*b**2 + 3*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*cos(x)*sin(x)**2*b**2 - 3*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*cos(x)*b**2 + 2*sin(x)**3*a**3 - sin(x)**3*a**2*b - 3*sin(x)**3*a*b**2 - 3*sin(x)*a**3 + 3*sin(x)*a*b**2)/(3*cos(x)*a**3*(sin(x)**2*a + sin(x)**2*b - a - b))

```

3.38 $\int \frac{\sec^6(x)}{a+b\cos^2(x)} dx$

Optimal result	292
Mathematica [A] (verified)	292
Rubi [A] (verified)	293
Maple [A] (verified)	294
Fricas [B] (verification not implemented)	295
Sympy [F(-1)]	295
Maxima [A] (verification not implemented)	296
Giac [A] (verification not implemented)	296
Mupad [B] (verification not implemented)	297
Reduce [B] (verification not implemented)	297

Optimal result

Integrand size = 15, antiderivative size = 79

$$\begin{aligned} \int \frac{\sec^6(x)}{a+b\cos^2(x)} dx = & \frac{b^3 \arctan\left(\frac{\sqrt{a+b}\cot(x)}{\sqrt{a}}\right)}{a^{7/2}\sqrt{a+b}} + \frac{(a^2-ab+b^2)\tan(x)}{a^3} \\ & + \frac{(2a-b)\tan^3(x)}{3a^2} + \frac{\tan^5(x)}{5a} \end{aligned}$$

output $b^3 \arctan((a+b)^{1/2} \cot(x)/a^{1/2})/a^{7/2}/(a+b)^{1/2}+(a^2-a*b+b^2)*\tan(x)/a^3+1/3*(2*a-b)*\tan(x)^3/a^2+1/5*\tan(x)^5/a$

Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\begin{aligned} \int \frac{\sec^6(x)}{a+b\cos^2(x)} dx = & -\frac{b^3 \arctan\left(\frac{\sqrt{a}\tan(x)}{\sqrt{a+b}}\right)}{a^{7/2}\sqrt{a+b}} \\ & + \frac{(8a^2-10ab+15b^2+a(4a-5b)\sec^2(x)+3a^2\sec^4(x))\tan(x)}{15a^3} \end{aligned}$$

input `Integrate[Sec[x]^6/(a + b*Cos[x]^2), x]`

output
$$-\left(\frac{(b^3 \operatorname{ArcTan}[(\sqrt{a} \tan[x])/\sqrt{a+b}])/(a^{(7/2)} \sqrt{a+b})}{a^2 - 10a*b + 15b^2 + a*(4*a - 5*b)*\sec[x]^2 + 3*a^2*\sec[x]^4*\tan[x]}\right)/(15*a^3)$$

Rubi [A] (verified)

Time = 0.32 (sec), antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 3666, 364, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sec^6(x)}{a + b \cos^2(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\sin\left(x + \frac{\pi}{2}\right)^6 \left(a + b \sin\left(x + \frac{\pi}{2}\right)^2\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3666} \\
 & - \int \frac{(\cot^2(x) + 1)^3 \tan^6(x)}{(a + b) \cot^2(x) + a} d \cot(x) \\
 & \quad \downarrow \textcolor{blue}{364} \\
 & - \int \left(\frac{\tan^6(x)}{a} + \frac{(2a - b) \tan^4(x)}{a^2} + \frac{(a^2 - ba + b^2) \tan^2(x)}{a^3} + \frac{b^3}{a^3 ((a + b) \cot^2(x) - a)} \right) d \cot(x) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{b^3 \arctan\left(\frac{\sqrt{a+b} \cot(x)}{\sqrt{a}}\right)}{a^{7/2} \sqrt{a+b}} + \frac{(2a - b) \tan^3(x)}{3a^2} + \frac{(a^2 - ab + b^2) \tan(x)}{a^3} + \frac{\tan^5(x)}{5a}
 \end{aligned}$$

input $\operatorname{Int}[\sec[x]^6/(a + b*\cos[x]^2), x]$

output
$$(b^3 \operatorname{ArcTan}[(\operatorname{Sqrt}[a+b] \operatorname{Cot}[x])/\operatorname{Sqrt}[a]])/(a^{(7/2)} \operatorname{Sqrt}[a+b]) + ((a^2 - a b + b^2) \operatorname{Tan}[x])/a^3 + ((2 a - b) \operatorname{Tan}[x]^3)/(3 a^2) + \operatorname{Tan}[x]^5/(5 a)$$

Definitions of rubi rules used

rule 364
$$\operatorname{Int}[((e_*)*(x_))^m_*((a_*) + (b_*)*(x_)^2)^p)/((c_*) + (d_*)*(x_)^2), x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(e*x)^m*((a+b*x^2)^p/(c+d*x^2)), x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, m\}, x] \& \operatorname{NeQ}[b*c - a*d, 0] \& \operatorname{IGtQ}[p, 0] \& (\operatorname{IntegerQ}[m] \mid\mid \operatorname{IGtQ}[2*(m+1), 0] \mid\mid \operatorname{!RationalQ}[m])$$

rule 2009
$$\operatorname{Int}[u_, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

rule 3042
$$\operatorname{Int}[u_, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3666
$$\operatorname{Int}[\sin[(e_*) + (f_*)*(x_)]^m_*((a_*) + (b_*)*\sin[(e_*) + (f_*)*(x_)]^2)^p, x_{\text{Symbol}}] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Tan}[e+f*x], x]\}, \operatorname{Simp}[ff^{(m+1)/f} \operatorname{Subst}[\operatorname{Int}[x^m*((a+(a+b)*ff^2*x^2)^p/(1+ff^2*x^2)^{(m/2+p+1)}, x], x, \operatorname{Tan}[e+f*x]/ff], x]] /; \operatorname{FreeQ}[\{a, b, e, f\}, x] \& \operatorname{IntegerQ}[m/2] \& \operatorname{IntegerQ}[p]$$

Maple [A] (verified)

Time = 1.33 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.99

method	result
default	$\frac{\tan(x)^5 a^2}{5} + \frac{2 a^2 \tan(x)^3}{3} - \frac{a b \tan(x)^3}{3} + \tan(x) a^2 - \tan(x) a b + \tan(x) b^2 - \frac{b^3 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{a^3 \sqrt{(a+b)a}}$
risch	$\frac{2 i (15 b^2 e^{8 i x} - 30 a b e^{6 i x} + 60 b^2 e^{6 i x} + 80 a^2 e^{4 i x} - 70 a b e^{4 i x} + 90 b^2 e^{4 i x} + 40 a^2 e^{2 i x} - 50 a b e^{2 i x} + 60 e^{2 i x} b^2 + 8 a^2 - 10 a b + 15 b^2)}{15 a^3 (e^{2 i x} + 1)^5} - \frac{b^3 \ln(e^{2 i x})}{e^{2 i x}}$

input
$$\operatorname{int}(\sec(x)^6/(a+b*\cos(x)^2), x, \operatorname{method}=\operatorname{RETURNVERBOSE})$$

output
$$\frac{1}{a^3} \left(\frac{1}{5} \tan(x)^5 a^2 + \frac{2}{3} a^2 \tan(x)^3 - \frac{1}{3} a b \tan(x)^3 + \tan(x) a^2 - \tan(x) a b + \tan(x) b^2 - b^3/a^3 \right) / ((a+b)*a)^{(1/2)} * \arctan(a \tan(x)) / ((a+b)*a)^{(1/2)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. $2(67) = 134$.

Time = 0.22 (sec) , antiderivative size = 348, normalized size of antiderivative = 4.41

$$\int \frac{\sec^6(x)}{a + b \cos^2(x)} dx \\ = \left[\frac{15 \sqrt{-a^2 - ab} b^3 \cos(x)^5 \log \left(\frac{(8a^2 + 8ab + b^2) \cos(x)^4 - 2(4a^2 + 3ab) \cos(x)^2 - 4((2a+b)\cos(x)^3 - a\cos(x))\sqrt{-a^2 - ab} \sin(x) + 15a^2b^2 \cos(x)^2 + 15a^2b^2 \cos(x)^4 + 2ab\cos(x)^2 + a^2}{b^2 \cos(x)^4 + 2ab\cos(x)^2 + a^2} \right)}{60(a^5 + a^3b^2 + a^2b^3 + ab^4 + b^5)} \right]$$

input `integrate(sec(x)^6/(a+b*cos(x)^2),x, algorithm="fricas")`

output
$$[-1/60*(15*sqrt(-a^2 - a*b)*b^3*cos(x)^5*log(((8*a^2 + 8*a*b + b^2)*cos(x)^4 - 2*(4*a^2 + 3*a*b)*cos(x)^2 - 4*((2*a + b)*cos(x)^3 - a*cos(x))*sqrt(-a^2 - a*b)*sin(x) + a^2)/(b^2*cos(x)^4 + 2*a*b*cos(x)^2 + a^2)) - 4*((8*a^4 - 2*a^3*b + 5*a^2*b^2 + 15*a*b^3)*cos(x)^4 + 3*a^4 + 3*a^3*b + (4*a^4 - a^3*b - 5*a^2*b^2)*cos(x)^2*sin(x))/((a^5 + a^4*b)*cos(x)^5), 1/30*(15*sqrt(a^2 + a*b)*b^3*arctan(1/2*((2*a + b)*cos(x)^2 - a)/(sqrt(a^2 + a*b)*cos(x)*sin(x)))*cos(x)^5 + 2*((8*a^4 - 2*a^3*b + 5*a^2*b^2 + 15*a*b^3)*cos(x)^4 + 3*a^4 + 3*a^3*b + (4*a^4 - a^3*b - 5*a^2*b^2)*cos(x)^2*sin(x))/((a^5 + a^4*b)*cos(x)^5)]$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sec^6(x)}{a + b \cos^2(x)} dx = \text{Timed out}$$

input `integrate(sec(x)**6/(a+b*cos(x)**2),x)`

output Timed out

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.94

$$\int \frac{\sec^6(x)}{a + b \cos^2(x)} dx = -\frac{b^3 \arctan\left(\frac{a \tan(x)}{\sqrt{(a+b)a}}\right)}{\sqrt{(a+b)a}a^3} + \frac{3 a^2 \tan(x)^5 + 5 (2 a^2 - ab) \tan(x)^3 + 15 (a^2 - ab + b^2) \tan(x)}{15 a^3}$$

input `integrate(sec(x)^6/(a+b*cos(x)^2),x, algorithm="maxima")`

output `-b^3*arctan(a*tan(x)/sqrt((a + b)*a))/(sqrt((a + b)*a)*a^3) + 1/15*(3*a^2*tan(x)^5 + 5*(2*a^2 - a*b)*tan(x)^3 + 15*(a^2 - a*b + b^2)*tan(x))/a^3`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.32

$$\int \frac{\sec^6(x)}{a + b \cos^2(x)} dx = -\frac{\left(\pi \lfloor \frac{x}{\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(x)}{\sqrt{a^2 + ab}}\right)\right) b^3}{\sqrt{a^2 + ab}a^3} + \frac{3 a^4 \tan(x)^5 + 10 a^4 \tan(x)^3 - 5 a^3 b \tan(x)^3 + 15 a^4 \tan(x) - 15 a^3 b \tan(x) + 15 a^2 b^2 \tan(x)}{15 a^5}$$

input `integrate(sec(x)^6/(a+b*cos(x)^2),x, algorithm="giac")`

output `-(pi*floor(x/pi + 1/2)*sgn(a) + arctan(a*tan(x)/sqrt(a^2 + a*b)))*b^3/(sqrt(a^2 + a*b)*a^3) + 1/15*(3*a^4*tan(x)^5 + 10*a^4*tan(x)^3 - 5*a^3*b*tan(x)^3 + 15*a^4*tan(x) - 15*a^3*b*tan(x) + 15*a^2*b^2*tan(x))/a^5`

Mupad [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.06

$$\int \frac{\sec^6(x)}{a + b \cos^2(x)} dx = \frac{\tan(x)^5}{5a} - \tan(x)^3 \left(\frac{a+b}{3a^2} - \frac{1}{a} \right) \\ + \tan(x) \left(\frac{3}{a} + \frac{(a+b) \left(\frac{a+b}{a^2} - \frac{3}{a} \right)}{a} \right) - \frac{b^3 \operatorname{atan} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a+b}} \right)}{a^{7/2} \sqrt{a+b}}$$

input `int(1/(cos(x)^6*(a + b*cos(x)^2)),x)`

output $\tan(x)^5/(5*a) - \tan(x)^3*((a + b)/(3*a^2) - 1/a) + \tan(x)*(3/a + ((a + b)*((a + b)/a^2 - 3/a))/a) - (b^3*\operatorname{atan}((a^(1/2)*\tan(x))/(a + b)^(1/2)))/(a^(7/2)*(a + b)^(1/2))$

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 346, normalized size of antiderivative = 4.38

$$\int \frac{\sec^6(x)}{a + b \cos^2(x)} dx \\ = \frac{-15\sqrt{a}\sqrt{a+b}\operatorname{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})-\sqrt{b}}{\sqrt{a}}\right)\cos(x)\sin(x)^4b^3 + 30\sqrt{a}\sqrt{a+b}\operatorname{atan}\left(\frac{\sqrt{a+b}\tan(\frac{x}{2})-\sqrt{b}}{\sqrt{a}}\right)\cos(x)\sin(x)^5b^2}{\sqrt{a}}$$

input `int(sec(x)^6/(a+b*cos(x)^2),x)`

```
output
( - 15*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*cos(x)*sin(x)**4*b**3 + 30*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*cos(x)*sin(x)**2*b**3 - 15*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) - sqrt(b))/sqrt(a))*cos(x)*b**3 - 15*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*cos(x)*sin(x)**4*b**3 + 30*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*cos(x)*cos(x)*sin(x)**2*b**3 - 15*sqrt(a)*sqrt(a + b)*atan((sqrt(a + b)*tan(x/2) + sqrt(b))/sqrt(a))*cos(x)*b**3 + 8*sin(x)**5*a**4 - 2*sin(x)**5*a**3*b + 5*sin(x)**5*a**2*b**2 + 15*sin(x)**5*a*b**3 - 20*sin(x)**3*a**4 + 5*sin(x)**3*a**3*b - 5*sin(x)**3*a**2*b**2 - 30*sin(x)**3*a*b**3 + 15*sin(x)*a**4 + 15*sin(x)*a*b**3)/(15*cos(x)*a**4*(sin(x)**4*a + sin(x)**4*b - 2*sin(x)**2*a - 2*sin(x)**2*b + a + b))
```

3.39 $\int \frac{\cos(x)}{\sqrt{1+\cos^2(x)}} dx$

Optimal result	299
Mathematica [A] (verified)	299
Rubi [A] (verified)	300
Maple [B] (verified)	301
Fricas [B] (verification not implemented)	301
Sympy [F]	302
Maxima [A] (verification not implemented)	302
Giac [B] (verification not implemented)	302
Mupad [F(-1)]	303
Reduce [F]	303

Optimal result

Integrand size = 13, antiderivative size = 9

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \arcsin\left(\frac{\sin(x)}{\sqrt{2}}\right)$$

output `arcsin(1/2*sin(x)*2^(1/2))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \arcsin\left(\frac{\sin(x)}{\sqrt{2}}\right)$$

input `Integrate[Cos[x]/Sqrt[1 + Cos[x]^2], x]`

output `ArcSin[Sin[x]/Sqrt[2]]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 3665, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(x)}{\sqrt{\cos^2(x) + 1}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})}{\sqrt{\sin^2(x + \frac{\pi}{2}) + 1}} dx \\
 & \quad \downarrow \textcolor{blue}{3665} \\
 & \int \frac{1}{\sqrt{2 - \sin^2(x)}} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{223} \\
 & \arcsin\left(\frac{\sin(x)}{\sqrt{2}}\right)
 \end{aligned}$$

input `Int[Cos[x]/Sqrt[1 + Cos[x]^2], x]`

output `ArcSin[Sin[x]/Sqrt[2]]`

Definitions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simplify[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3665 $\text{Int}[\sin[(e_.) + (f_.)*(x_.)]^{(m_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2)^{(p_.)}, x \text{Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Cos}[e + f*x], x]\}, \text{Simp}[-ff/f \text{Subst}[\text{Int}[(1 - ff^2*x^2)^{((m - 1)/2)}*(a + b - b*ff^2*x^2)^p, x], x, \text{Cos}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \& \text{IntegerQ}[(m - 1)/2]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 32 vs. $2(8) = 16$.

Time = 0.09 (sec), antiderivative size = 33, normalized size of antiderivative = 3.67

method	result	size
default	$-\frac{\sqrt{(1+\cos(x)^2) \sin(x)^2} \arcsin(\cos(x)^2)}{2 \sin(x) \sqrt{1+\cos(x)^2}}$	33

input `int(cos(x)/(1+cos(x)^2)^(1/2), x, method=_RETURNVERBOSE)`

output $-1/2*((1+\cos(x)^2)*\sin(x)^2)^(1/2)*\arcsin(\cos(x)^2)/\sin(x)/(1+\cos(x)^2)^(1/2)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(8) = 16$.

Time = 0.11 (sec), antiderivative size = 49, normalized size of antiderivative = 5.44

$$\begin{aligned} \int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx &= \frac{1}{2} \arctan \left(\frac{\sqrt{\cos(x)^2 + 1} \cos(x)^2 \sin(x) - \cos(x) \sin(x)}{\cos(x)^4 + \cos(x)^2 - 1} \right) \\ &\quad + \frac{1}{2} \arctan \left(\frac{\sin(x)}{\cos(x)} \right) \end{aligned}$$

input `integrate(cos(x)/(1+cos(x)^2)^(1/2), x, algorithm="fricas")`

output
$$\frac{1}{2} \operatorname{arctan}\left(\frac{\sqrt{\cos^2(x) + 1} \cdot \cos(x)^2 \sin(x) - \cos(x) \sin(x)}{\cos(x)^4 + \cos(x)^2 - 1}\right) + \frac{1}{2} \operatorname{arctan}\left(\frac{\sin(x)}{\cos(x)}\right)$$

Sympy [F]

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \int \frac{\cos(x)}{\sqrt{\cos^2(x) + 1}} dx$$

input `integrate(cos(x)/(1+cos(x)**2)**(1/2),x)`

output `Integral(cos(x)/sqrt(cos(x)**2 + 1), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \arcsin\left(\frac{1}{2} \sqrt{2} \sin(x)\right)$$

input `integrate(cos(x)/(1+cos(x)**2)**(1/2),x, algorithm="maxima")`

output `arcsin(1/2*sqrt(2)*sin(x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 23 vs. $2(8) = 16$.

Time = 0.13 (sec) , antiderivative size = 23, normalized size of antiderivative = 2.56

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \frac{1}{2} \sqrt{-\sin(x)^2 + 2 \sin(x) + \arcsin\left(\frac{1}{2} \sqrt{2} \sin(x)\right)}$$

input `integrate(cos(x)/(1+cos(x)**2)**(1/2),x, algorithm="giac")`

output $\frac{1}{2}\sqrt{-\sin(x)^2 + 2} \sin(x) + \arcsin(\frac{1}{2}\sqrt{2}\sin(x))$

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \int \frac{\cos(x)}{\sqrt{\cos(x)^2 + 1}} dx$$

input $\text{int}(\cos(x)/(cos(x)^2 + 1)^{(1/2)}, x)$

output $\text{int}(\cos(x)/(cos(x)^2 + 1)^{(1/2)}, x)$

Reduce [F]

$$\int \frac{\cos(x)}{\sqrt{1 + \cos^2(x)}} dx = \int \frac{\sqrt{\cos(x)^2 + 1} \cos(x)}{\cos(x)^2 + 1} dx$$

input $\text{int}(\cos(x)/(1+\cos(x)^2)^{(1/2)}, x)$

output $\text{int}((\sqrt{\cos(x)^2 + 1})\cos(x)/(\cos(x)^2 + 1), x)$

3.40 $\int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx$

Optimal result	304
Mathematica [A] (verified)	304
Rubi [A] (verified)	305
Maple [B] (verified)	306
Fricas [B] (verification not implemented)	306
Sympy [F]	307
Maxima [A] (verification not implemented)	307
Giac [F]	307
Mupad [F(-1)]	308
Reduce [F]	308

Optimal result

Integrand size = 21, antiderivative size = 15

$$\int \frac{\cos(5 + 3x)}{\sqrt{3 + \cos^2(5 + 3x)}} dx = \frac{1}{3} \arcsin\left(\frac{1}{2} \sin(5 + 3x)\right)$$

output `1/3*arcsin(1/2*sin(5+3*x))`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos(5 + 3x)}{\sqrt{3 + \cos^2(5 + 3x)}} dx = \frac{1}{3} \arcsin\left(\frac{1}{2} \sin(5 + 3x)\right)$$

input `Integrate[Cos[5 + 3*x]/Sqrt[3 + Cos[5 + 3*x]^2], x]`

output `ArcSin[Sin[5 + 3*x]/2]/3`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 3665, 223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(3x + 5)}{\sqrt{\cos^2(3x + 5) + 3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(3x + \frac{\pi}{2} + 5)}{\sqrt{\sin^2(3x + \frac{\pi}{2} + 5) + 3}} dx \\
 & \quad \downarrow \text{3665} \\
 & \frac{1}{3} \int \frac{1}{\sqrt{4 - \sin^2(3x + 5)}} d\sin(3x + 5) \\
 & \quad \downarrow \text{223} \\
 & \frac{1}{3} \arcsin\left(\frac{1}{2} \sin(3x + 5)\right)
 \end{aligned}$$

input `Int[Cos[5 + 3*x]/Sqrt[3 + Cos[5 + 3*x]^2], x]`

output `ArcSin[Sin[5 + 3*x]/2]/3`

Definitions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simplify[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3665

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^p_, x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Simplify[-ff/f]
Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(11) = 22$.

Time = 0.25 (sec) , antiderivative size = 57, normalized size of antiderivative = 3.80

method	result	size
default	$\frac{\sqrt{(3+\cos(5+3x)^2)\sin(5+3x)^2} \arcsin\left(-1+\frac{\sin(5+3x)^2}{2}\right)}{6\sin(5+3x)\sqrt{3+\cos(5+3x)^2}}$	57

input `int(cos(5+3*x)/(3+cos(5+3*x)^2)^(1/2), x, method=_RETURNVERBOSE)`

output
$$\frac{1}{6} \cdot \frac{(\sqrt{3+\cos(5+3x)^2}) \cdot \sin(5+3x)^2}{(\sqrt{3+\cos(5+3x)^2})^{1/2}} \cdot \arcsin\left(\frac{-1+\frac{\sin(5+3x)^2}{2}}{\sqrt{3+\cos(5+3x)^2}}\right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 89 vs. $2(11) = 22$.

Time = 0.13 (sec) , antiderivative size = 89, normalized size of antiderivative = 5.93

$$\begin{aligned} & \int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx \\ &= \frac{1}{6} \arctan \left(\frac{\sqrt{\cos(3x+5)^2 + 3} (\cos(3x+5)^2 + 1) \sin(3x+5) - 4 \cos(3x+5) \sin(3x+5)}{\cos(3x+5)^4 + 6 \cos(3x+5)^2 - 3} \right) \\ &+ \frac{1}{6} \arctan \left(\frac{\sin(3x+5)}{\cos(3x+5)} \right) \end{aligned}$$

input `integrate(cos(5+3*x)/(3+cos(5+3*x)^2)^(1/2), x, algorithm="fricas")`

output

```
1/6*arctan((sqrt(cos(3*x + 5)^2 + 3)*(cos(3*x + 5)^2 + 1)*sin(3*x + 5) - 4
*cos(3*x + 5)*sin(3*x + 5))/(cos(3*x + 5)^4 + 6*cos(3*x + 5)^2 - 3)) + 1/6
*arctan(sin(3*x + 5)/cos(3*x + 5))
```

Sympy [F]

$$\int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx = \int \frac{\cos(3x+5)}{\sqrt{\cos^2(3x+5)+3}} dx$$

input

```
integrate(cos(5+3*x)/(3+cos(5+3*x)**2)**(1/2),x)
```

output

```
Integral(cos(3*x + 5)/sqrt(cos(3*x + 5)**2 + 3), x)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.73

$$\int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx = \frac{1}{3} \arcsin\left(\frac{1}{2} \sin(3x+5)\right)$$

input

```
integrate(cos(5+3*x)/(3+cos(5+3*x)**2)^(1/2),x, algorithm="maxima")
```

output

```
1/3*arcsin(1/2*sin(3*x + 5))
```

Giac [F]

$$\int \frac{\cos(5+3x)}{\sqrt{3+\cos^2(5+3x)}} dx = \int \frac{\cos(3x+5)}{\sqrt{\cos(3x+5)^2 + 3}} dx$$

input

```
integrate(cos(5+3*x)/(3+cos(5+3*x)**2)^(1/2),x, algorithm="giac")
```

output `integrate(cos(3*x + 5)/sqrt(cos(3*x + 5)^2 + 3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(5 + 3x)}{\sqrt{3 + \cos^2(5 + 3x)}} dx = \int \frac{\cos(3x + 5)}{\sqrt{\cos(3x + 5)^2 + 3}} dx$$

input `int(cos(3*x + 5)/(cos(3*x + 5)^2 + 3)^(1/2), x)`

output `int(cos(3*x + 5)/(cos(3*x + 5)^2 + 3)^(1/2), x)`

Reduce [F]

$$\int \frac{\cos(5 + 3x)}{\sqrt{3 + \cos^2(5 + 3x)}} dx = \int \frac{\sqrt{\cos(3x + 5)^2 + 3} \cos(3x + 5)}{\cos(3x + 5)^2 + 3} dx$$

input `int(cos(5+3*x)/(3+cos(5+3*x)^2)^(1/2), x)`

output `int((sqrt(cos(3*x + 5)**2 + 3)*cos(3*x + 5))/(cos(3*x + 5)**2 + 3), x)`

3.41 $\int \frac{\cos(x)}{\sqrt{4-\cos^2(x)}} dx$

Optimal result	309
Mathematica [A] (verified)	309
Rubi [A] (verified)	310
Maple [B] (verified)	311
Fricas [B] (verification not implemented)	311
Sympy [F]	312
Maxima [A] (verification not implemented)	312
Giac [B] (verification not implemented)	312
Mupad [F(-1)]	313
Reduce [F]	313

Optimal result

Integrand size = 15, antiderivative size = 9

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \operatorname{arcsinh}\left(\frac{\sin(x)}{\sqrt{3}}\right)$$

output `arcsinh(1/3*sin(x)*3^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \operatorname{arcsinh}\left(\frac{\sin(x)}{\sqrt{3}}\right)$$

input `Integrate[Cos[x]/Sqrt[4 - Cos[x]^2], x]`

output `ArcSinh[Sin[x]/Sqrt[3]]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 3665, 222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\sin(x + \frac{\pi}{2})}{\sqrt{4 - \sin^2(x + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \textcolor{blue}{3665} \\
 & \int \frac{1}{\sqrt{\sin^2(x) + 3}} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{222} \\
 & \operatorname{arcsinh}\left(\frac{\sin(x)}{\sqrt{3}}\right)
 \end{aligned}$$

input `Int[Cos[x]/Sqrt[4 - Cos[x]^2], x]`

output `ArcSinh[Sin[x]/Sqrt[3]]`

Definitions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simplify[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3665

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^p_, x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Simplify[-ff/f]
Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(8) = 16$.

Time = 0.16 (sec), antiderivative size = 53, normalized size of antiderivative = 5.89

method	result	size
default	$-\frac{\sqrt{-\left(\cos(x)^2-4\right) \sin(x)^2} \ln \left(-\sin(x)^2+\sqrt{\sin(x)^4+3 \sin(x)^2}-\frac{3}{2}\right)}{2 \sin(x) \sqrt{4-\cos(x)^2}}$	53

input `int(cos(x)/(4-cos(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{-1/2*(-(\cos(x)^2-4)*\sin(x)^2)^(1/2)*\ln(-\sin(x)^2+(\sin(x)^4+3*\sin(x)^2)^(1/2)-3/2)/\sin(x)/(4-\cos(x)^2)^(1/2)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(8) = 16$.

Time = 0.13 (sec), antiderivative size = 39, normalized size of antiderivative = 4.33

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \frac{1}{4} \log \left(8 \cos(x)^4 - 4 (2 \cos(x)^2 - 5) \sqrt{-\cos(x)^2 + 4 \sin(x)} - 40 \cos(x)^2 + 41 \right)$$

input `integrate(cos(x)/(4-cos(x)^2)^(1/2),x, algorithm="fricas")`

output
$$\frac{1}{4} \log(8 \cos(x)^4 - 4 (2 \cos(x)^2 - 5) \sqrt{-\cos(x)^2 + 4} \sin(x) - 40 \cos(x)^2 + 41)$$

Sympy [F]

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \int \frac{\cos(x)}{\sqrt{-(\cos(x) - 2)(\cos(x) + 2)}} dx$$

input `integrate(cos(x)/(4-cos(x)**2)**(1/2),x)`

output `Integral(cos(x)/sqrt(-(cos(x) - 2)*(cos(x) + 2)), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \operatorname{arsinh}\left(\frac{1}{3} \sqrt{3} \sin(x)\right)$$

input `integrate(cos(x)/(4-cos(x)^2)^(1/2),x, algorithm="maxima")`

output `arcsinh(1/3*sqrt(3)*sin(x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(8) = 16$.

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 3.22

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \frac{1}{2} \sqrt{\sin(x)^2 + 3} \sin(x) - \frac{3}{2} \log\left(\sqrt{\sin(x)^2 + 3} - \sin(x)\right)$$

input `integrate(cos(x)/(4-cos(x)^2)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(sin(x)^2 + 3)*sin(x) - 3/2*log(sqrt(sin(x)^2 + 3) - sin(x))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = \int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx$$

input `int(cos(x)/(4 - cos(x)^2)^(1/2),x)`

output `int(cos(x)/(4 - cos(x)^2)^(1/2), x)`

Reduce [F]

$$\int \frac{\cos(x)}{\sqrt{4 - \cos^2(x)}} dx = - \left(\int \frac{\sqrt{-\cos^2(x) + 4} \cos(x)}{\cos^2(x) - 4} dx \right)$$

input `int(cos(x)/(4-cos(x)^2)^(1/2),x)`

output `- int(sqrt(-cos(x)^2 + 4)*cos(x))/(cos(x)^2 - 4),x)`

3.42 $\int \frac{\tan(x)}{1+\cos^2(x)} dx$

Optimal result	314
Mathematica [A] (verified)	314
Rubi [A] (verified)	315
Maple [A] (verified)	316
Fricas [A] (verification not implemented)	317
Sympy [F]	317
Maxima [A] (verification not implemented)	317
Giac [A] (verification not implemented)	318
Mupad [B] (verification not implemented)	318
Reduce [B] (verification not implemented)	318

Optimal result

Integrand size = 11, antiderivative size = 17

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = -\log(\cos(x)) + \frac{1}{2} \log(1 + \cos^2(x))$$

output `-ln(cos(x))+1/2*ln(1+cos(x)^2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = -\log(\cos(x)) + \frac{1}{2} \log(1 + \cos^2(x))$$

input `Integrate[Tan[x]/(1 + Cos[x]^2), x]`

output `-Log[Cos[x]] + Log[1 + Cos[x]^2]/2`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 25, 3673, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\cos^2(x) + 1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{1}{(\sin(x + \frac{\pi}{2})^2 + 1) \tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{1}{(\sin(x + \frac{\pi}{2})^2 + 1) \tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3673} \\
 & -\frac{1}{2} \int \frac{\sec^2(x)}{\cos^2(x) + 1} d\cos^2(x) \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\int \frac{1}{\cos^2(x) + 1} d\cos^2(x) - \int \sec^2(x) d\cos^2(x) \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(\int \frac{1}{\cos^2(x) + 1} d\cos^2(x) - \log(\cos^2(x)) \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(\cos^2(x) + 1) - \log(\cos^2(x)))
 \end{aligned}$$

input `Int[Tan[x]/(1 + Cos[x]^2), x]`

output `(-Log[Cos[x]^2] + Log[1 + Cos[x]^2])/2`

Definitions of rubi rules used

rule 14 $\text{Int}[(a_)/(x_), \ x_\text{Symbol}] \rightarrow \text{Simp}[a*\text{Log}[x], x] /; \ \text{FreeQ}[a, x]$

rule 16 $\text{Int}[(c_)/((a_) + (b_)*(x_)), \ x_\text{Symbol}] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \ \text{FreeQ}[\{a, b, c\}, x]$

rule 25 $\text{Int}[-(F x_), \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$

rule 47 $\text{Int}[1/(((a_) + (b_)*(x_))*(c_) + (d_)*(x_))), \ x_\text{Symbol}] \rightarrow \text{Simp}[b/(b*c - a*d) \quad \text{Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \quad \text{Int}[1/(c + d*x), x], x] /; \ \text{FreeQ}[\{a, b, c, d\}, x]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \ \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3673 $\text{Int}[(a_) + (b_)*\sin[(e_) + (f_)*(x_)]^2]^p * \tan[(e_) + (f_)*(x_)]^m, \ x_\text{Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x]^2, x]\}, \ \text{Simp}[ff^{(m + 1)/2}/(2*f) \quad \text{Subst}[\text{Int}[x^{(m - 1)/2}*((a + b*ff*x)^p/(1 - ff*x))^{(m + 1)/2}), x], x, \text{Sin}[e + f*x]^{2/ff}, x]] /; \ \text{FreeQ}[\{a, b, e, f, p\}, x] \ \&& \text{IntegerQ}[(m - 1)/2]$

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$-\ln(\cos(x)) + \frac{\ln(1+\cos(x)^2)}{2}$	16
risch	$-\ln(e^{2ix} + 1) + \frac{\ln(e^{4ix}+6e^{2ix}+1)}{2}$	29

input `int(tan(x)/(1+cos(x)^2),x,method=_RETURNVERBOSE)`

output $-\ln(\cos(x)) + 1/2 \ln(1+\cos(x)^2)$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = \frac{1}{2} \log \left(\frac{1}{2} \cos(x)^2 + \frac{1}{2} \right) - \log(-\cos(x))$$

input `integrate(tan(x)/(1+cos(x)^2),x, algorithm="fricas")`

output $1/2 \log(1/2 * \cos(x)^2 + 1/2) - \log(-\cos(x))$

Sympy [F]

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = \int \frac{\tan(x)}{\cos^2(x) + 1} dx$$

input `integrate(tan(x)/(1+cos(x)**2),x)`

output `Integral(tan(x)/(cos(x)**2 + 1), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = -\frac{1}{2} \log(\sin(x)^2 - 1) + \frac{1}{2} \log(\sin(x)^2 - 2)$$

input `integrate(tan(x)/(1+cos(x)^2),x, algorithm="maxima")`

output $-1/2 \log(\sin(x)^2 - 1) + 1/2 \log(\sin(x)^2 - 2)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = \frac{1}{2} \log(\cos(x)^2 + 1) - \log(|\cos(x)|)$$

input `integrate(tan(x)/(1+cos(x)^2),x, algorithm="giac")`

output `1/2*log(cos(x)^2 + 1) - log(abs(cos(x)))`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.53

$$\int \frac{\tan(x)}{1 + \cos^2(x)} dx = \frac{\ln(\tan(x)^2 + 2)}{2}$$

input `int(tan(x)/(cos(x)^2 + 1),x)`

output `log(tan(x)^2 + 2)/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 56, normalized size of antiderivative = 3.29

$$\begin{aligned} \int \frac{\tan(x)}{1 + \cos^2(x)} dx &= \frac{\log(-\sqrt{2} \tan(\frac{x}{2}) + \tan(\frac{x}{2})^2 + 1)}{2} - \log(\tan(\frac{x}{2}) - 1) \\ &\quad - \log(\tan(\frac{x}{2}) + 1) + \frac{\log(\sqrt{2} \tan(\frac{x}{2}) + \tan(\frac{x}{2})^2 + 1)}{2} \end{aligned}$$

input `int(tan(x)/(1+cos(x)^2),x)`

output
$$\frac{(\log(-\sqrt{2}\tan(x/2) + \tan(x/2)^2 + 1) - 2\log(\tan(x/2) - 1) - 2\log(\tan(x/2) + 1) + \log(\sqrt{2}\tan(x/2) + \tan(x/2)^2 + 1))/2}{}$$

3.43 $\int \sqrt{a + b \cos^2(x)} \tan(x) dx$

Optimal result	320
Mathematica [A] (verified)	320
Rubi [A] (verified)	321
Maple [A] (verified)	323
Fricas [A] (verification not implemented)	323
Sympy [F]	324
Maxima [B] (verification not implemented)	324
Giac [A] (verification not implemented)	325
Mupad [F(-1)]	325
Reduce [F]	325

Optimal result

Integrand size = 15, antiderivative size = 40

$$\int \sqrt{a + b \cos^2(x)} \tan(x) dx = \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^2(x)}}{\sqrt{a}} \right) - \sqrt{a + b \cos^2(x)}$$

output `a^(1/2)*arctanh((a+b*cos(x)^2)^(1/2)/a^(1/2))-(a+b*cos(x)^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos^2(x)} \tan(x) dx = \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^2(x)}}{\sqrt{a}} \right) - \sqrt{a + b \cos^2(x)}$$

input `Integrate[Sqrt[a + b*Cos[x]^2]*Tan[x],x]`

output `Sqrt[a]*ArcTanh[Sqrt[a + b*Cos[x]^2]/Sqrt[a]] - Sqrt[a + b*Cos[x]^2]`

Rubi [A] (verified)

Time = 0.25 (sec), antiderivative size = 45, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 25, 3673, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(x) \sqrt{a + b \cos^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sqrt{a + b \sin(x + \frac{\pi}{2})^2}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\sqrt{b \sin(x + \frac{\pi}{2})^2 + a}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3673} \\
 & -\frac{1}{2} \int \sqrt{b \cos^2(x) + a} \sec^2(x) d \cos^2(x) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(-a \int \frac{\sec^2(x)}{\sqrt{b \cos^2(x) + a}} d \cos^2(x) - 2 \sqrt{a + b \cos^2(x)} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(-\frac{2a \int \frac{1}{\cos^4(x) - \frac{a}{b}} d \sqrt{b \cos^2(x) + a}}{b} - 2 \sqrt{a + b \cos^2(x)} \right) \\
 & \quad \downarrow \text{221} \\
 & \frac{1}{2} \left(2\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^2(x)}}{\sqrt{a}} \right) - 2 \sqrt{a + b \cos^2(x)} \right)
 \end{aligned}$$

input `Int[Sqrt[a + b*Cos[x]^2]*Tan[x],x]`

output
$$\frac{(2\sqrt{a} \operatorname{ArcTanh}[\sqrt{a + b \cos x^2}/\sqrt{a}] - 2\sqrt{a + b \cos x^2})}{2}$$

Definitions of rubi rules used

rule 25
$$\operatorname{Int}[-(F_{x_1}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_{x_1}, x], x]$$

rule 60
$$\operatorname{Int}[(a_{\cdot} + b_{\cdot}x_{\cdot})^{m_{\cdot}} \cdot (c_{\cdot} + d_{\cdot}x_{\cdot})^{n_{\cdot}}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(a + b*x)^{m+1} \cdot ((c + d*x)^n / (b*(m+n+1))), x] + \operatorname{Simp}[n \cdot ((b*c - a*d) / (b*(m+n+1))) \operatorname{Int}[(a + b*x)^{m-1} \cdot (c + d*x)^{n-1}, x], x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \& \operatorname{GtQ}[n, 0] \& \operatorname{NeQ}[m + n + 1, 0] \& !(\operatorname{IGtQ}[m, 0] \& \operatorname{IntegerQ}[n] \mid\mid (\operatorname{GtQ}[m, 0] \& \operatorname{LtQ}[m - n, 0])) \& \operatorname{ILtQ}[m + n + 2, 0] \& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 73
$$\operatorname{Int}[(a_{\cdot} + b_{\cdot}x_{\cdot})^{m_{\cdot}} \cdot (c_{\cdot} + d_{\cdot}x_{\cdot})^{n_{\cdot}}, x_{\text{Symbol}}] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Simp}[p/b \operatorname{Subst}[\operatorname{Int}[x^{p*(m+1)-1} \cdot (c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{1/p}], x] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \& \operatorname{LtQ}[-1, m, 0] \& \operatorname{LeQ}[-1, n, 0] \& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]]]$$

rule 221
$$\operatorname{Int}[(a_{\cdot} + b_{\cdot}x_{\cdot})^2 \cdot (-1), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a) \operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{NegQ}[a/b]$$

rule 3042
$$\operatorname{Int}[u_{\cdot}, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3673
$$\operatorname{Int}[(a_{\cdot} + b_{\cdot}x_{\cdot}) \sin[(e_{\cdot} + f_{\cdot}x_{\cdot})^2]^p \tan[(e_{\cdot} + f_{\cdot}x_{\cdot})^m], x_{\text{Symbol}}] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Sin}[e + f*x]^2, x]\}, \operatorname{Simp}[ff^{(m+1)/2} / (2*f) \operatorname{Subst}[\operatorname{Int}[x^{((m-1)/2)} \cdot ((a + b*ff*x)^p / (1 - ff*x))^{(m+1)/2}], x, \operatorname{Sin}[e + f*x]^2 / ff], x] /; \operatorname{FreeQ}[\{a, b, e, f, p\}, x] \& \operatorname{IntegerQ}[(m-1)/2]]$$

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08

method	result	size
default	$-\sqrt{a + b \cos(x)^2} + \sqrt{a} \ln\left(\frac{2a+2\sqrt{a}\sqrt{a+b \cos(x)^2}}{\cos(x)}\right)$	43

input `int((a+b*cos(x)^2)^(1/2)*tan(x),x,method=_RETURNVERBOSE)`

output $-(a+b \cos(x)^2)^{(1/2)}+a^{(1/2)} \ln((2*a+2*a^{(1/2)}*(a+b \cos(x)^2)^{(1/2)})/\cos(x))$

Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.18

$$\begin{aligned} \int \sqrt{a + b \cos^2(x)} \tan(x) dx = & \left[\frac{1}{2} \sqrt{a} \log \left(\frac{b \cos(x)^2 + 2 \sqrt{b \cos(x)^2 + a} \sqrt{a} + 2a}{\cos(x)^2} \right) \right. \\ & - \sqrt{b \cos(x)^2 + a}, -\sqrt{-a} \arctan \left(\frac{\sqrt{-a}}{\sqrt{b \cos(x)^2 + a}} \right) \\ & \left. - \sqrt{b \cos(x)^2 + a} \right] \end{aligned}$$

input `integrate((a+b*cos(x)^2)^(1/2)*tan(x),x, algorithm="fricas")`

output $[1/2*\sqrt{a}*\log((b*\cos(x)^2 + 2*\sqrt{b*\cos(x)^2 + a})*\sqrt{a} + 2*a)/\cos(x)^2 - \sqrt{b*\cos(x)^2 + a}, -\sqrt{-a}*\arctan(\sqrt{-a}/\sqrt{b*\cos(x)^2 + a}) - \sqrt{b*\cos(x)^2 + a}]$

Sympy [F]

$$\int \sqrt{a + b \cos^2(x)} \tan(x) dx = \int \sqrt{a + b \cos^2(x)} \tan(x) dx$$

input `integrate((a+b*cos(x)**2)**(1/2)*tan(x),x)`

output `Integral(sqrt(a + b*cos(x)**2)*tan(x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(32) = 64$.

Time = 0.13 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.38

$$\begin{aligned} \int \sqrt{a + b \cos^2(x)} \tan(x) dx &= \frac{1}{2} \sqrt{a} \log \left(b - \frac{\sqrt{-b \sin(x)^2 + a + b} \sqrt{a}}{\sin(x) - 1} - \frac{a}{\sin(x) - 1} \right) \\ &\quad + \frac{1}{2} \sqrt{a} \log \left(-b + \frac{\sqrt{-b \sin(x)^2 + a + b} \sqrt{a}}{\sin(x) + 1} \right. \\ &\quad \left. + \frac{a}{\sin(x) + 1} \right) - \sqrt{-b \sin(x)^2 + a + b} \end{aligned}$$

input `integrate((a+b*cos(x)^2)^(1/2)*tan(x),x, algorithm="maxima")`

output `1/2*sqrt(a)*log(b - sqrt(-b*sin(x)^2 + a + b)*sqrt(a)/(sin(x) - 1) - a/(sin(x) - 1)) + 1/2*sqrt(a)*log(-b + sqrt(-b*sin(x)^2 + a + b)*sqrt(a)/(sin(x) + 1) + a/(sin(x) + 1)) - sqrt(-b*sin(x)^2 + a + b)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \sqrt{a + b \cos^2(x)} \tan(x) dx = -\frac{a \arctan\left(\frac{\sqrt{b \cos(x)^2 + a}}{\sqrt{-a}}\right)}{\sqrt{-a}} - \sqrt{b \cos(x)^2 + a}$$

input `integrate((a+b*cos(x)^2)^(1/2)*tan(x),x, algorithm="giac")`

output `-a*arctan(sqrt(b*cos(x)^2 + a)/sqrt(-a))/sqrt(-a) - sqrt(b*cos(x)^2 + a)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos^2(x)} \tan(x) dx = \int \tan(x) \sqrt{b \cos(x)^2 + a} dx$$

input `int(tan(x)*(a + b*cos(x)^2)^(1/2),x)`

output `int(tan(x)*(a + b*cos(x)^2)^(1/2), x)`

Reduce [F]

$$\int \sqrt{a + b \cos^2(x)} \tan(x) dx = \int \sqrt{\cos(x)^2 b + a} \tan(x) dx$$

input `int((a+b*cos(x)^2)^(1/2)*tan(x),x)`

output `int(sqrt(cos(x)**2*b + a)*tan(x),x)`

3.44 $\int \sqrt{1 - \cos^2(x)} \tan(x) dx$

Optimal result	326
Mathematica [A] (verified)	326
Rubi [A] (verified)	327
Maple [A] (verified)	329
Fricas [A] (verification not implemented)	330
Sympy [F]	330
Maxima [B] (verification not implemented)	330
Giac [B] (verification not implemented)	331
Mupad [F(-1)]	331
Reduce [F]	331

Optimal result

Integrand size = 15, antiderivative size = 20

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = \operatorname{arctanh}\left(\sqrt{\sin^2(x)}\right) - \sqrt{\sin^2(x)}$$

output `arctanh((sin(x)^2)^(1/2))-(sin(x)^2)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = (-1 + \operatorname{arctanh}(\sin(x)) \csc(x)) \sqrt{\sin^2(x)}$$

input `Integrate[Sqrt[1 - Cos[x]^2]*Tan[x],x]`

output `(-1 + ArcTanh[Sin[x]]*Csc[x])*Sqrt[Sin[x]^2]`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.30, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 25, 3655, 25, 3042, 3684, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{1 - \cos^2(x)} \tan(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sqrt{1 - \sin(x + \frac{\pi}{2})^2}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\sqrt{1 - \sin(x + \frac{\pi}{2})^2}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3655} \\
 & - \int -\sqrt{\sin^2(x)} \tan(x) dx \\
 & \quad \downarrow \text{25} \\
 & \int \sqrt{\sin^2(x)} \tan(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\sin(x)^2} \tan(x) dx \\
 & \quad \downarrow \text{3684} \\
 & \frac{1}{2} \int \frac{\sqrt{\sin^2(x)}}{1 - \sin^2(x)} d \sin^2(x) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{2} \left(\int \frac{1}{\sqrt{\sin^2(x)} (1 - \sin^2(x))} d \sin^2(x) - 2 \sqrt{\sin^2(x)} \right) \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{2} \left(2 \int \frac{1}{1 - \sin^4(x)} d\sqrt{\sin^2(x)} - 2\sqrt{\sin^2(x)} \right)$$

↓ 219

$$\frac{1}{2} \left(2 \operatorname{arctanh} \left(\sqrt{\sin^2(x)} \right) - 2\sqrt{\sin^2(x)} \right)$$

input `Int[Sqrt[1 - Cos[x]^2]*Tan[x], x]`

output `(2*ArcTanh[Sqrt[Sin[x]^2]] - 2*Sqrt[Sin[x]^2])/2`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^(p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 $\text{Int}[u_., x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3655 $\text{Int}[(a_.) + (b_.) \sin[(e_.) + (f_.) (x_)]^2]^p, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ActivateTrig}[a \cos[e + f x]^2]^p, x] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \& \text{EqQ}[a + b, 0]$

rule 3684 $\text{Int}[(b_.) \sin[(e_.) + (f_.) (x_)]^{(n_.)}]^p \tan[(e_.) + (f_.) (x_)]^{(m_.)}, x_{\text{Symbol}}] \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Sin}[e + f x]^2, x]\}, \text{Simp}[\text{ff}^{(m + 1)/2}/(2 f) \text{Subst}[\text{Int}[x^{((m - 1)/2)} ((b \text{ff}^{(n/2)} x^{(n/2)})^p)/(1 - \text{ff} x)^{(m + 1)/2}], x], x, \text{Sin}[e + f x]^2/\text{ff}], x] /; \text{FreeQ}[\{b, e, f, p\}, x] \& \text{IntegerQ}[(m - 1)/2] \& \text{IntegerQ}[n/2]$

Maple [A] (verified)

Time = 0.09 (sec), antiderivative size = 17, normalized size of antiderivative = 0.85

method	result
default	$-\sqrt{\frac{1}{2} - \frac{\cos(2x)}{2}} + \text{arctanh}\left(\frac{2}{\sqrt{2-2\cos(2x)}}\right)$
risch	$-\frac{\sqrt{-(e^{2ix}-1)^2 e^{-2ix}} e^{2ix}}{2(e^{2ix}-1)} + \frac{\sqrt{-(e^{2ix}-1)^2 e^{-2ix}}}{2 e^{2ix}-2} - \frac{i \sqrt{-(e^{2ix}-1)^2 e^{-2ix}} e^{ix} \ln(e^{ix}-i)}{e^{2ix}-1} + \frac{i \sqrt{-(e^{2ix}-1)^2 e^{-2ix}} e^{ix} \ln(e^{ix}+i)}{e^{2ix}-1}$

input `int((1-cos(x)^2)^(1/2)*tan(x),x,method=_RETURNVERBOSE)`

output `-(sin(x)^2)^(1/2)+arctanh(1/(sin(x)^2)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(-\sin(x) + 1) - \sin(x)$$

input `integrate((1-cos(x)^2)^(1/2)*tan(x),x, algorithm="fricas")`

output `1/2*log(sin(x) + 1) - 1/2*log(-sin(x) + 1) - sin(x)`

Sympy [F]

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = \int \sqrt{-(\cos(x) - 1)(\cos(x) + 1)} \tan(x) dx$$

input `integrate((1-cos(x)**2)**(1/2)*tan(x),x)`

output `Integral(sqrt(-(cos(x) - 1)*(cos(x) + 1))*tan(x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. $2(16) = 32$.

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\begin{aligned} \int \sqrt{1 - \cos^2(x)} \tan(x) dx &= \frac{1}{2} (-1)^{\sin(x)} \log\left(-\frac{\sin(x)}{\sin(x) + 1}\right) \\ &\quad + \frac{1}{2} (-1)^{\sin(x)} \log\left(-\frac{\sin(x)}{\sin(x) - 1}\right) - \sqrt{\sin(x)^2} \end{aligned}$$

input `integrate((1-cos(x)^2)^(1/2)*tan(x),x, algorithm="maxima")`

output `1/2*(-1)^(2*sin(x))*log(-sin(x)/(sin(x) + 1)) + 1/2*(-1)^(2*sin(x))*log(-sin(x)/(sin(x) - 1)) - sqrt(sin(x)^2)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. $2(16) = 32$.

Time = 0.11 (sec), antiderivative size = 45, normalized size of antiderivative = 2.25

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = -\sqrt{-\cos(x)^2 + 1} + \frac{1}{2} \log \left(\sqrt{-\cos(x)^2 + 1} + 1 \right) \\ - \frac{1}{2} \log \left(-\sqrt{-\cos(x)^2 + 1} + 1 \right)$$

input `integrate((1-cos(x)^2)^(1/2)*tan(x),x, algorithm="giac")`

output `-sqrt(-cos(x)^2 + 1) + 1/2*log(sqrt(-cos(x)^2 + 1) + 1) - 1/2*log(-sqrt(-cos(x)^2 + 1) + 1)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = \int \tan(x) \sqrt{1 - \cos(x)^2} dx$$

input `int(tan(x)*(1 - cos(x)^2)^(1/2),x)`

output `int(tan(x)*(1 - cos(x)^2)^(1/2), x)`

Reduce [F]

$$\int \sqrt{1 - \cos^2(x)} \tan(x) dx = \int \sqrt{-\cos(x)^2 + 1} \tan(x) dx$$

input `int((1-cos(x)^2)^(1/2)*tan(x),x)`

output `int(sqrt(-cos(x)**2 + 1)*tan(x),x)`

3.45 $\int \frac{\tan(x)}{\sqrt{a+b\cos^2(x)}} dx$

Optimal result	332
Mathematica [A] (verified)	332
Rubi [A] (verified)	333
Maple [A] (verified)	334
Fricas [A] (verification not implemented)	335
Sympy [F]	335
Maxima [B] (verification not implemented)	335
Giac [A] (verification not implemented)	336
Mupad [F(-1)]	337
Reduce [F]	337

Optimal result

Integrand size = 15, antiderivative size = 25

$$\int \frac{\tan(x)}{\sqrt{a + b\cos^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\cos^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}$$

output arctanh((a+b*cos(x)^2)^(1/2)/a^(1/2))/a^(1/2)

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{a + b\cos^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\cos^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}$$

input Integrate[Tan[x]/Sqrt[a + b*Cos[x]^2],x]

output ArcTanh[Sqrt[a + b*Cos[x]^2]/Sqrt[a]]/Sqrt[a]

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 25, 3673, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\tan(x + \frac{\pi}{2}) \sqrt{a + b \sin(x + \frac{\pi}{2})^2}} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\sqrt{b \sin(x + \frac{\pi}{2})^2 + a \tan(x + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \textcolor{blue}{3673} \\
 & -\frac{1}{2} \int \frac{\sec^2(x)}{\sqrt{b \cos^2(x) + a}} d \cos^2(x) \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & -\frac{\int \frac{1}{\frac{\cos^4(x)}{b} - \frac{a}{b}} d \sqrt{b \cos^2(x) + a}}{b} \\
 & \quad \downarrow \textcolor{blue}{221} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}
 \end{aligned}$$

input `Int[Tan[x]/Sqrt[a + b*Cos[x]^2],x]`

output `ArcTanh[Sqrt[a + b*Cos[x]^2]/Sqrt[a]]/Sqrt[a]`

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 73 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__)^{\text{m}__}*((\text{c}__) + (\text{d}__)*(\text{x}__)^{\text{n}__}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}__]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^{\text{p}/\text{b}}))^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{(1/\text{p})}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&& \text{LtQ}[-1, \text{m}, 0] \&& \text{LeQ}[-1, \text{n}, 0] \&& \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \&& \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$

rule 221 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&& \text{NegQ}[\text{a}/\text{b}]$

rule 3042 $\text{Int}[\text{u}__, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3673 $\text{Int}[(\text{a}__) + (\text{b}__)*\sin[(\text{e}__) + (\text{f}__)*(\text{x}__)]^2)^{\text{p}__}*\tan[(\text{e}__) + (\text{f}__)*(\text{x}__)]^{\text{m}__}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Sin}[\text{e} + \text{f}*\text{x}]^2, \text{x}]\}, \text{Simp}[\text{ff}^{((\text{m} + 1)/2)/(2*\text{f})} \quad \text{Subst}[\text{Int}[\text{x}^{((\text{m} - 1)/2)}*((\text{a} + \text{b}*\text{ff}*\text{x})^{\text{p}}/(1 - \text{ff}*\text{x})^{((\text{m} + 1)/2)}], \text{x}], \text{x}, \text{Sin}[\text{e} + \text{f}*\text{x}]^{2/\text{ff}}, \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{p}\}, \text{x}] \&& \text{IntegerQ}[(\text{m} - 1)/2]$

Maple [A] (verified)

Time = 0.09 (sec), antiderivative size = 30, normalized size of antiderivative = 1.20

method	result	size
default	$\frac{\ln\left(\frac{2a+2\sqrt{a}\sqrt{a+b\cos(x)^2}}{\cos(x)}\right)}{\sqrt{a}}$	30

input $\text{int}(\tan(\text{x})/(\text{a}+\text{b}*\cos(\text{x})^2)^{(1/2)}, \text{x}, \text{method}=\text{_RETURNVERBOSE})$

output $1/a^{(1/2)}*\ln((2*a+2*a^{(1/2)}*(a+b*cos(x)^2)^{(1/2)})/cos(x))$

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.56

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx \\ = \left[\frac{\log\left(\frac{b \cos(x)^2 + 2 \sqrt{b \cos(x)^2 + a} \sqrt{a + 2a}}{\cos(x)^2}\right)}{2\sqrt{a}}, -\frac{\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{b \cos(x)^2 + a}}\right)}{a} \right]$$

input `integrate(tan(x)/(a+b*cos(x)^2)^(1/2),x, algorithm="fricas")`

output `[1/2*log((b*cos(x)^2 + 2*sqrt(b*cos(x)^2 + a)*sqrt(a) + 2*a)/cos(x)^2)/sqrt(a), -sqrt(-a)*arctan(sqrt(-a)/sqrt(b*cos(x)^2 + a))/a]`

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx$$

input `integrate(tan(x)/(a+b*cos(x)**2)**(1/2),x)`

output `Integral(tan(x)/sqrt(a + b*cos(x)**2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. 2(19) = 38.

Time = 0.12 (sec) , antiderivative size = 81, normalized size of antiderivative = 3.24

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx = \frac{\log \left(b - \frac{\sqrt{-b \sin(x)^2 + a + b \sqrt{a}}}{\sin(x) - 1} - \frac{a}{\sin(x) - 1} \right)}{2 \sqrt{a}} \\ + \frac{\log \left(-b + \frac{\sqrt{-b \sin(x)^2 + a + b \sqrt{a}}}{\sin(x) + 1} + \frac{a}{\sin(x) + 1} \right)}{2 \sqrt{a}}$$

input `integrate(tan(x)/(a+b*cos(x)^2)^(1/2),x, algorithm="maxima")`

output `1/2*log(b - sqrt(-b*sin(x)^2 + a + b)*sqrt(a)/(sin(x) - 1) - a/(sin(x) - 1))/sqrt(a) + 1/2*log(-b + sqrt(-b*sin(x)^2 + a + b)*sqrt(a)/(sin(x) + 1) + a/(sin(x) + 1))/sqrt(a)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx = -\frac{\arctan \left(\frac{\sqrt{b \cos(x)^2 + a}}{\sqrt{-a}} \right)}{\sqrt{-a}}$$

input `integrate(tan(x)/(a+b*cos(x)^2)^(1/2),x, algorithm="giac")`

output `-arctan(sqrt(b*cos(x)^2 + a)/sqrt(-a))/sqrt(-a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{b \cos^2(x) + a}} dx$$

input `int(tan(x)/(a + b*cos(x)^2)^(1/2),x)`

output `int(tan(x)/(a + b*cos(x)^2)^(1/2), x)`

Reduce [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^2(x)}} dx = \int \frac{\sqrt{\cos^2(b) + a} \tan(x)}{\cos^2(b) + a} dx$$

input `int(tan(x)/(a+b*cos(x)^2)^(1/2),x)`

output `int((sqrt(cos(x)**2*b + a)*tan(x))/(cos(x)**2*b + a),x)`

3.46 $\int \frac{\tan(x)}{\sqrt{1+\cos^2(x)}} dx$

Optimal result	338
Mathematica [A] (verified)	338
Rubi [A] (verified)	339
Maple [A] (verified)	340
Fricas [A] (verification not implemented)	341
Sympy [F]	341
Maxima [B] (verification not implemented)	341
Giac [B] (verification not implemented)	342
Mupad [F(-1)]	342
Reduce [F]	343

Optimal result

Integrand size = 13, antiderivative size = 11

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \operatorname{arctanh}\left(\sqrt{1 + \cos^2(x)}\right)$$

output `arctanh((1+cos(x)^2)^(1/2))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \operatorname{arctanh}\left(\sqrt{1 + \cos^2(x)}\right)$$

input `Integrate[Tan[x]/Sqrt[1 + Cos[x]^2], x]`

output `ArcTanh[Sqrt[1 + Cos[x]^2]]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 25, 3673, 73, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\sqrt{\cos^2(x) + 1}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\sqrt{\sin\left(x + \frac{\pi}{2}\right)^2 + 1} \tan\left(x + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & -\int \frac{1}{\sqrt{\sin\left(x + \frac{\pi}{2}\right)^2 + 1} \tan\left(x + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \textcolor{blue}{3673} \\
 & -\frac{1}{2} \int \frac{\sec^2(x)}{\sqrt{\cos^2(x) + 1}} d\cos^2(x) \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & -\int \frac{1}{\cos^4(x) - 1} d\sqrt{\cos^2(x) + 1} \\
 & \quad \downarrow \textcolor{blue}{220} \\
 & \operatorname{arctanh}\left(\sqrt{\cos^2(x) + 1}\right)
 \end{aligned}$$

input `Int[Tan[x]/Sqrt[1 + Cos[x]^2], x]`

output `ArcTanh[Sqrt[1 + Cos[x]^2]]`

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 73 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__)^{\text{m}__}*(\text{c}__) + (\text{d}__)*(\text{x}__)^{\text{n}__}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}__]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^{\text{p}/\text{b}}))^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{(1/\text{p})}], \text{x}]\} /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&& \text{LtQ}[-1, \text{m}, 0] \&& \text{LeQ}[-1, \text{n}, 0] \&& \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \&& \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$

rule 220 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__)^2^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-(\text{Rt}[-\text{a}, 2]*\text{Rt}[\text{b}, 2])^{(-1)})*\text{ArcTanh}[\text{Rt}[\text{b}, 2]*(\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&& \text{NegQ}[\text{a}/\text{b}] \&& (\text{LtQ}[\text{a}, 0] \text{||} \text{GtQ}[\text{b}, 0])$

rule 3042 $\text{Int}[\text{u}__, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3673 $\text{Int}[(\text{a}__) + (\text{b}__)*\sin[(\text{e}__) + (\text{f}__)*(\text{x}__)]^2)^{\text{p}__}*\tan[(\text{e}__) + (\text{f}__)*(\text{x}__)]^{\text{m}__}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Sin}[\text{e} + \text{f}*\text{x}]^2, \text{x}]\}, \text{Simp}[\text{ff}^{((\text{m} + 1)/2)/(2*\text{f})} \quad \text{Subst}[\text{Int}[\text{x}^{((\text{m} - 1)/2)}*((\text{a} + \text{b}*\text{ff}*\text{x})^{\text{p}}/(1 - \text{ff}*\text{x})^{((\text{m} + 1)/2)}], \text{x}, \text{Sin}[\text{e} + \text{f}*\text{x}]^{2/\text{ff}], \text{x}]\} /; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{p}\}, \text{x}] \&& \text{Integ erQ}[(\text{m} - 1)/2]$

Maple [A] (verified)

Time = 0.08 (sec), antiderivative size = 10, normalized size of antiderivative = 0.91

method	result	size
default	$\text{arctanh}\left(\frac{1}{\sqrt{1+\cos(x)^2}}\right)$	10

input $\text{int}(\tan(\text{x})/(1+\cos(\text{x})^2)^{(1/2)}, \text{x}, \text{method}=\text{_RETURNVERBOSE})$

output $\operatorname{arctanh}\left(1/(1+\cos(x)^2)^{(1/2)}\right)$

Fricas [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \log\left(\frac{\sqrt{\cos(x)^2 + 1} + 1}{\cos(x)}\right)$$

input `integrate(tan(x)/(1+cos(x)^2)^{(1/2)},x, algorithm="fricas")`

output $\log((\sqrt{\cos(x)^2 + 1} + 1)/\cos(x))$

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{\cos^2(x) + 1}} dx$$

input `integrate(tan(x)/(1+cos(x)**2)**(1/2),x)`

output `Integral(tan(x)/sqrt(cos(x)**2 + 1), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(9) = 18$.

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 5.45

$$\begin{aligned} \int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx &= \frac{1}{2} \log\left(\frac{\sqrt{-\sin(x)^2 + 2}}{\sin(x) + 1} + \frac{1}{\sin(x) + 1} - 1\right) \\ &\quad + \frac{1}{2} \log\left(-\frac{\sqrt{-\sin(x)^2 + 2}}{\sin(x) - 1} - \frac{1}{\sin(x) - 1} + 1\right) \end{aligned}$$

input `integrate(tan(x)/(1+cos(x)^2)^(1/2),x, algorithm="maxima")`

output $\frac{1}{2} \log(\sqrt{-\sin(x)^2 + 2}) / (\sin(x) + 1) + \frac{1}{\sin(x) + 1 - 1} + \frac{1}{2} \log(-\sqrt{-\sin(x)^2 + 2}) / (\sin(x) - 1) - \frac{1}{\sin(x) - 1 + 1}$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(9) = 18$.

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.45

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \frac{1}{2} \log \left(\sqrt{\cos(x)^2 + 1} + 1 \right) - \frac{1}{2} \log \left(\sqrt{\cos(x)^2 + 1} - 1 \right)$$

input `integrate(tan(x)/(1+cos(x)^2)^(1/2),x, algorithm="giac")`

output $\frac{1}{2} \log(\sqrt{\cos(x)^2 + 1} + 1) - \frac{1}{2} \log(\sqrt{\cos(x)^2 + 1} - 1)$

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{\cos(x)^2 + 1}} dx$$

input `int(tan(x)/(cos(x)^2 + 1)^(1/2),x)`

output `int(tan(x)/(cos(x)^2 + 1)^(1/2), x)`

Reduce [F]

$$\int \frac{\tan(x)}{\sqrt{1 + \cos^2(x)}} dx = \int \frac{\sqrt{\cos(x)^2 + 1} \tan(x)}{\cos(x)^2 + 1} dx$$

input `int(tan(x)/(1+cos(x)^2)^(1/2),x)`

output `int((sqrt(cos(x)**2 + 1)*tan(x))/(cos(x)**2 + 1),x)`

3.47 $\int \frac{\tan(x)}{\sqrt{1-\cos^2(x)}} dx$

Optimal result	344
Mathematica [A] (verified)	344
Rubi [A] (verified)	345
Maple [A] (verified)	347
Fricas [B] (verification not implemented)	347
Sympy [F]	348
Maxima [B] (verification not implemented)	348
Giac [B] (verification not implemented)	348
Mupad [F(-1)]	349
Reduce [F]	349

Optimal result

Integrand size = 15, antiderivative size = 9

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = \operatorname{arctanh}\left(\sqrt{\sin^2(x)}\right)$$

output `arctanh((sin(x)^2)^(1/2))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.56

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = \frac{\coth^{-1}(\sin(x)) \sin(x)}{\sqrt{\sin^2(x)}}$$

input `Integrate[Tan[x]/Sqrt[1 - Cos[x]^2], x]`

output `(ArcCoth[Sin[x]]*Sin[x])/Sqrt[Sin[x]^2]`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3042, 25, 3655, 25, 3042, 3684, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\sqrt{1 - \sin(x + \frac{\pi}{2})^2} \tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\sqrt{1 - \sin(x + \frac{\pi}{2})^2} \tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \textcolor{blue}{3655} \\
 & - \int -\frac{\tan(x)}{\sqrt{\sin^2(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & \int \frac{\tan(x)}{\sqrt{\sin^2(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\tan(x)}{\sqrt{\sin(x)^2}} dx \\
 & \quad \downarrow \textcolor{blue}{3684} \\
 & \frac{1}{2} \int \frac{1}{\sqrt{\sin^2(x)} (1 - \sin^2(x))} d \sin^2(x) \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & \int \frac{1}{1 - \sin^4(x)} d \sqrt{\sin^2(x)} \\
 & \quad \downarrow \textcolor{blue}{219}
 \end{aligned}$$

$$\operatorname{arctanh}\left(\sqrt{\sin^2(x)}\right)$$

input `Int[Tan[x]/Sqrt[1 - Cos[x]^2], x]`

output `ArcTanh[Sqrt[Sin[x]^2]]`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simplify[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Simplify[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^(p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simplify[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3655 `Int[(u_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^p, x_Symbol] :> Int[ActivateTrig[u*(a*cos[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0]`

rule 3684

```
Int[((b_)*sin[(e_.) + (f_.)*(x_)]^(n_.))^(p_.*tan[(e_.) + (f_.*)(x_)]^(m_.),
), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x]^2, x]}, Simplify[ff^(m + 1)/2/(2*f) Subst[Int[x^((m - 1)/2)*((b*ff^(n/2)*x^(n/2))^p/(1 - ff*x)^(m + 1)/2)], x], x, Sin[e + f*x]^2/ff], x]] /; FreeQ[{b, e, f, p}, x] && IntegerQ[(m - 1)/2] && IntegerQ[n/2]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.89

method	result	size
default	$\operatorname{arctanh}\left(\frac{2}{\sqrt{2-2 \cos (2 x)}}\right)$	8
risch	$-\frac{2 \ln (\mathrm{e}^{i x}-i) \sin (x)}{\sqrt{-\left(\mathrm{e}^{2 i x}-1\right)^2 \mathrm{e}^{-2 i x}}}+\frac{2 \ln (\mathrm{e}^{i x}+i) \sin (x)}{\sqrt{-\left(\mathrm{e}^{2 i x}-1\right)^2 \mathrm{e}^{-2 i x}}}$	64

input `int(tan(x)/(1-cos(x)^2)^(1/2),x,method=_RETURNVERBOSE)`output `arctanh(1/(\sin(x)^2)^(1/2))`**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(7) = 14$.

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(-\sin(x) + 1)$$

input `integrate(tan(x)/(1-cos(x)^2)^(1/2),x, algorithm="fricas")`output `1/2*log(sin(x) + 1) - 1/2*log(-sin(x) + 1)`

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{-(\cos(x) - 1)(\cos(x) + 1)}} dx$$

input `integrate(tan(x)/(1-cos(x)**2)**(1/2),x)`

output `Integral(tan(x)/sqrt(-(cos(x) - 1)*(cos(x) + 1)), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(7) = 14$.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 4.33

$$\begin{aligned} & \int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx \\ &= \frac{1}{2} (-1)^{2 \sin(x)} \log \left(-\frac{\sin(x)}{\sin(x) + 1} \right) + \frac{1}{2} (-1)^{2 \sin(x)} \log \left(-\frac{\sin(x)}{\sin(x) - 1} \right) \end{aligned}$$

input `integrate(tan(x)/(1-cos(x)^2)^(1/2),x, algorithm="maxima")`

output `1/2*(-1)^(2*sin(x))*log(-sin(x)/(sin(x) + 1)) + 1/2*(-1)^(2*sin(x))*log(-sin(x)/(sin(x) - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(7) = 14$.

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 3.67

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = \frac{1}{2} \log \left(\sqrt{-\cos(x)^2 + 1} + 1 \right) - \frac{1}{2} \log \left(-\sqrt{-\cos(x)^2 + 1} + 1 \right)$$

input `integrate(tan(x)/(1-cos(x)^2)^(1/2),x, algorithm="giac")`

output $\frac{1}{2} \log(\sqrt{-\cos(x)^2 + 1}) + \frac{1}{2} \log(-\sqrt{-\cos(x)^2 + 1})$

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx$$

input `int(tan(x)/(1 - cos(x)^2)^(1/2),x)`

output `int(tan(x)/(1 - cos(x)^2)^(1/2), x)`

Reduce [F]

$$\int \frac{\tan(x)}{\sqrt{1 - \cos^2(x)}} dx = - \left(\int \frac{\sqrt{-\cos^2(x) + 1} \tan(x)}{\cos^2(x) - 1} dx \right)$$

input `int(tan(x)/(1-cos(x)^2)^(1/2),x)`

output `- int((sqrt(-cos(x)^2 + 1)*tan(x))/(cos(x)^2 - 1),x)`

3.48 $\int \frac{\tan^3(x)}{a+b\cos^3(x)} dx$

Optimal result	350
Mathematica [C] (verified)	351
Rubi [A] (verified)	351
Maple [C] (verified)	353
Fricas [C] (verification not implemented)	354
Sympy [F]	355
Maxima [A] (verification not implemented)	355
Giac [A] (verification not implemented)	356
Mupad [B] (verification not implemented)	356
Reduce [F]	357

Optimal result

Integrand size = 15, antiderivative size = 153

$$\begin{aligned} \int \frac{\tan^3(x)}{a + b\cos^3(x)} dx = & -\frac{b^{2/3} \arctan\left(\frac{\sqrt[3]{a} - \sqrt[3]{b}\cos(x)}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt{3}a^{5/3}} \\ & + \frac{\log(\cos(x))}{a} + \frac{b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{b}\cos(x)\right)}{3a^{5/3}} \\ & - \frac{b^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}\cos(x) + b^{2/3}\cos^2(x)\right)}{6a^{5/3}} \\ & - \frac{\log(a + b\cos^3(x))}{3a} + \frac{\sec^2(x)}{2a} \end{aligned}$$

output

```
-1/3*b^(2/3)*arctan(1/3*(a^(1/3)-2*b^(1/3)*cos(x))*3^(1/2)/a^(1/3))*3^(1/2)
)/a^(5/3)+ln(cos(x))/a+1/3*b^(2/3)*ln(a^(1/3)+b^(1/3)*cos(x))/a^(5/3)-1/6*
b^(2/3)*ln(a^(2/3)-a^(1/3)*b^(1/3)*cos(x)+b^(2/3)*cos(x)^2)/a^(5/3)-1/3*ln
(a+b*cos(x)^3)/a+1/2*sec(x)^2/a
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.41 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.42

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx$$

$$= \frac{6(\log(\cos(x)) + \log(\sec^2(\frac{x}{2}))) - 2\text{RootSum}\left[a + b + 3a\#1 - 3b\#1 + 3a\#1^2 + 3b\#1^2 + a\#1^3 - b\#1^3\right]}{a + b + 3a\#1 - 3b\#1 + 3a\#1^2 + 3b\#1^2 + a\#1^3 - b\#1^3}$$

input `Integrate[Tan[x]^3/(a + b*Cos[x]^3), x]`

output
$$(6*(\text{Log}[\text{Cos}[x]] + \text{Log}[\text{Sec}[x/2]^2]) - 2*\text{RootSum}[a + b + 3*a\#1 - 3*b\#1 + 3*a\#1^2 + 3*b\#1^2 + a\#1^3 - b\#1^3 \&, (\text{a}\#\text{Log}[-\#1 + \text{Tan}[x/2]^2] + b\#\text{Log}[-\#1 + \text{Tan}[x/2]^2] + 2*\text{a}\#\text{Log}[-\#1 + \text{Tan}[x/2]^2]\#\#1 + 4*\text{b}\#\text{Log}[-\#1 + \text{Tan}[x/2]^2]\#\#1 + \text{a}\#\text{Log}[-\#1 + \text{Tan}[x/2]^2]\#\#1^2 - \text{b}\#\text{Log}[-\#1 + \text{Tan}[x/2]^2]\#\#1^2)/(a - b + 2*a\#1 + 2*b\#1 + a\#1^2 - b\#1^2) \&] + 3*\text{Sec}[x]^2)/(6*a)$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 25, 3709, 2373, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx$$

$$\downarrow \text{3042}$$

$$\int -\frac{1}{\tan\left(x + \frac{\pi}{2}\right)^3 \left(a + b \sin\left(x + \frac{\pi}{2}\right)^3\right)} dx$$

$$\downarrow \text{25}$$

$$\begin{aligned}
 & - \int \frac{1}{\left(b \sin \left(x + \frac{\pi}{2}\right)^3 + a\right) \tan \left(x + \frac{\pi}{2}\right)^3} dx \\
 & \quad \downarrow \textcolor{blue}{3709} \\
 & - \int \frac{(1 - \cos^2(x)) \sec^3(x)}{b \cos^3(x) + a} d \cos(x) \\
 & \quad \downarrow \textcolor{blue}{2373} \\
 & - \int \left(\frac{\sec^3(x)}{a} - \frac{\sec(x)}{a} + \frac{b(\cos^2(x) - 1)}{a(b \cos^3(x) + a)} \right) d \cos(x) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{b^{2/3} \arctan \left(\frac{\sqrt[3]{a} - 2\sqrt[3]{b} \cos(x)}{\sqrt[3]{a}} \right)}{\sqrt[3]{3} a^{5/3}} - \frac{b^{2/3} \log \left(a^{2/3} - \sqrt[3]{a} \sqrt[3]{b} \cos(x) + b^{2/3} \cos^2(x) \right)}{6 a^{5/3}} + \\
 & \frac{b^{2/3} \log \left(\sqrt[3]{a} + \sqrt[3]{b} \cos(x) \right)}{3 a^{5/3}} - \frac{\log(a + b \cos^3(x))}{3a} + \frac{\sec^2(x)}{2a} + \frac{\log(\cos(x))}{a}
 \end{aligned}$$

input `Int[Tan[x]^3/(a + b*Cos[x]^3), x]`

output `-((b^(2/3)*ArcTan[(a^(1/3) - 2*b^(1/3)*Cos[x])/(Sqrt[3]*a^(1/3))])/(Sqrt[3]*a^(5/3))) + Log[Cos[x]]/a + (b^(2/3)*Log[a^(1/3) + b^(1/3)*Cos[x]])/(3*a^(5/3)) - (b^(2/3)*Log[a^(2/3) - a^(1/3)*b^(1/3)*Cos[x] + b^(2/3)*Cos[x]^2])/(6*a^(5/3)) - Log[a + b*Cos[x]^3]/(3*a) + Sec[x]^2/(2*a)`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2373 `Int[((Pq_)*((c_.)*(x_.))^(m_.))/((a_) + (b_)*(x_.)^(n_)), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*(Pq/(a + b*x^n)), x], x] /; FreeQ[{a, b, c, m}, x] & PolyQ[Pq, x] && IntegerQ[n] && !IGtQ[m, 0]`

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3709 $\text{Int}[(a_ + b_)*((c_)*\sin[(e_ + f_)*(x_)])^n]^p \tan[(e_ + f_)*(x_)]^m, x_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Simp}[ff^m (m + 1)/f \text{Subst}[\text{Int}[x^m ((a + b*(c*ff*x)^n)^p)/(1 - ff^2*x^2)^{(m + 1)/2}], x, \text{Sin}[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, c, e, f, n, p\}, x] \& \text{ILtQ}[(m - 1)/2, 0]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.99 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.71

method	result
risch	$\frac{2e^{2ix}}{(e^{2ix}+1)^2a} + i \left(\sum_{R=\text{RootOf}(27Z^3a^5-27ia^4Z^2-9Za^3+ia^2-ib^2)} -R \ln \left(e^{2ix} + \left(\frac{6ia^2}{b}R + \frac{2a}{b} \right) e^{ix} + 1 \right) \right)$
default	$- \frac{b \left(-\frac{\ln \left(\cos(x) + \left(\frac{a}{b} \right)^{\frac{1}{3}} \right)}{3b \left(\frac{a}{b} \right)^{\frac{2}{3}}} + \frac{\ln \left(\cos(x)^2 - \left(\frac{a}{b} \right)^{\frac{1}{3}} \cos(x) + \left(\frac{a}{b} \right)^{\frac{2}{3}} \right)}{6b \left(\frac{a}{b} \right)^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan \left(\frac{\sqrt{3} \left(\frac{2 \cos(x)}{\left(\frac{a}{b} \right)^{\frac{1}{3}}} - 1 \right)}{3} \right)}{3b \left(\frac{a}{b} \right)^{\frac{2}{3}}} + \frac{\ln(a+b \cos(x)^3)}{3b} \right)}{a} + \frac{\ln(\cos(x))}{a} +$

input $\text{int}(\tan(x)^3/(a+b*\cos(x)^3), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$2*\exp(2*I*x)/(\exp(2*I*x)+1)^2/a + I*\sum(_R*\ln(\exp(2*I*x)+(6*I/b*a^2*_R+2/b*a)*\exp(I*x)+1), _R=\text{RootOf}(27*_Z^3*a^5-27*I*a^4*_Z^2-9*_Z*a^3+I*a^2-I*b^2))+1/a*\ln(\exp(2*I*x)+1)$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.79 (sec) , antiderivative size = 823, normalized size of antiderivative = 5.38

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx = \text{Too large to display}$$

input `integrate(tan(x)^3/(a+b*cos(x)^3),x, algorithm="fricas")`

output

```
-1/12*(2*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a*cos(x)^2*log(-1/2*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a^2 + b*cos(x) + a) - 12*cos(x)^2*log(-cos(x)) - (((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a*cos(x)^2 + 3*sqrt(1/3)*a*sqrt(-(((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)^2*a^2 - 4*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a + 4)/a^2)*cos(x)^2 - 6*cos(x)^2*log(1/2*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a^2 + 3/2*sqrt(1/3)*a^2*sqrt(-((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)^2*a^2 - 4*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a + 4)/a^2) + 2*b*cos(x) - a) - (((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a*cos(x)^2 - 3*sqrt(1/3)*a*sqrt(-(((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)^2*a^2 - 4*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a + 4)/a^2)*cos(x)^2 - 6*cos(x)^2*log(-1/2*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a^2 + 3/2*sqrt(1/3)*a^2*sqrt(-(((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)^2*a^2 - 4*((1/2)^(1/3)*(I*sqrt(3) + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^(1/3) + 2/a)*a + 4)/a^2) - 2*...
```

Sympy [F]

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx = \int \frac{\tan^3(x)}{a + b \cos^3(x)} dx$$

input `integrate(tan(x)**3/(a+b*cos(x)**3),x)`

output `Integral(tan(x)**3/(a + b*cos(x)**3), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.99

$$\begin{aligned} \int \frac{\tan^3(x)}{a + b \cos^3(x)} dx &= \frac{\sqrt{3} \left(b \left(3 \left(\frac{a}{b} \right)^{\frac{1}{3}} - \frac{2a}{b} \right) + 2a \right) \arctan \left(-\frac{\sqrt{3} \left(\left(\frac{a}{b} \right)^{\frac{1}{3}} - 2 \cos(x) \right)}{3 \left(\frac{a}{b} \right)^{\frac{1}{3}}} \right)}{9a^2} \\ &\quad - \frac{\left(2 \left(\frac{a}{b} \right)^{\frac{2}{3}} + 1 \right) \log \left(\cos(x)^2 - \left(\frac{a}{b} \right)^{\frac{1}{3}} \cos(x) + \left(\frac{a}{b} \right)^{\frac{2}{3}} \right)}{6a \left(\frac{a}{b} \right)^{\frac{2}{3}}} \\ &\quad - \frac{\left(\left(\frac{a}{b} \right)^{\frac{2}{3}} - 1 \right) \log \left(\left(\frac{a}{b} \right)^{\frac{1}{3}} + \cos(x) \right)}{3a \left(\frac{a}{b} \right)^{\frac{2}{3}}} + \frac{\log(\cos(x))}{a} + \frac{1}{2a \cos(x)^2} \end{aligned}$$

input `integrate(tan(x)^3/(a+b*cos(x)^3),x, algorithm="maxima")`

output `1/9*sqrt(3)*(b*(3*(a/b)^(1/3) - 2*a/b) + 2*a)*arctan(-1/3*sqrt(3)*((a/b)^(1/3) - 2*cos(x))/(a/b)^(1/3))/a^2 - 1/6*(2*(a/b)^(2/3) + 1)*log(cos(x)^2 - (a/b)^(1/3)*cos(x) + (a/b)^(2/3))/(a*(a/b)^(2/3)) - 1/3*((a/b)^(2/3) - 1)*log((a/b)^(1/3) + cos(x))/(a*(a/b)^(2/3)) + log(cos(x))/a + 1/2/(a*cos(x)^2)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.93

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx = -\frac{b(-\frac{a}{b})^{\frac{1}{3}} \log \left(\left| -(-\frac{a}{b})^{\frac{1}{3}} + \cos(x) \right| \right)}{3a^2} - \frac{\log(|b \cos(x)^3 + a|)}{3a}$$

$$+ \frac{\log(|\cos(x)|)}{a} + \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan \left(\frac{\sqrt{3} \left((-\frac{a}{b})^{\frac{1}{3}} + 2 \cos(x) \right)}{3(-\frac{a}{b})^{\frac{1}{3}}} \right)}{3a^2}$$

$$+ \frac{(-ab^2)^{\frac{1}{3}} \log \left(\cos(x)^2 + (-\frac{a}{b})^{\frac{1}{3}} \cos(x) + (-\frac{a}{b})^{\frac{2}{3}} \right)}{6a^2} + \frac{1}{2a \cos(x)^2}$$

input `integrate(tan(x)^3/(a+b*cos(x)^3),x, algorithm="giac")`

output `-1/3*b*(-a/b)^(1/3)*log(abs(-(-a/b)^(1/3) + cos(x)))/a^2 - 1/3*log(abs(b*cos(x)^3 + a))/a + log(abs(cos(x)))/a + 1/3*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*((-a/b)^(1/3) + 2*cos(x))/(-a/b)^(1/3))/a^2 + 1/6*(-a*b^2)^(1/3)*log(cos(x)^2 + (-a/b)^(1/3)*cos(x) + (-a/b)^(2/3))/a^2 + 1/2/(a*cos(x)^2)`

Mupad [B] (verification not implemented)

Time = 2.55 (sec) , antiderivative size = 1281, normalized size of antiderivative = 8.37

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx = \text{Too large to display}$$

input `int(tan(x)^3/(a + b*cos(x)^3),x)`

output

```
(2*tan(x/2)^2)/(a - 2*a*tan(x/2)^2 + a*tan(x/2)^4) + log(tan(x/2)^2 - 1)/a
+ symsum(log((262144*(9*a*b^10 - b^11 - 37*a^2*b^9 + 85*a^3*b^8 - 107*a^4
*b^7 + 43*a^5*b^6 + 73*a^6*b^5 - 121*a^7*b^4 + 72*a^8*b^3 - 16*a^9*b^2))/a
^6 + root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*(root(27*a^
5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*(root(27*a^5*z^3 + 27*a^4*
z^2 + 9*a^3*z + a^2 - b^2, z, k)*((262144*(72*a^5*b^9 - 96*a^6*b^8 + 1428*
a^7*b^7 - 3684*a^8*b^6 + 612*a^9*b^5 + 3972*a^10*b^4 - 2112*a^11*b^3 - 192
*a^12*b^2))/a^6 + root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k
)*(root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*((262144*(518
4*a^10*b^6 - 3024*a^9*b^7 + 1728*a^11*b^5 - 6048*a^12*b^4 + 1296*a^13*b^3
+ 864*a^14*b^2))/a^6 - root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2,
z, k)*((262144*(1296*a^10*b^7 - 3888*a^11*b^6 + 2592*a^12*b^5 + 2592*a^13
*b^4 - 3888*a^14*b^3 + 1296*a^15*b^2))/a^6 - (262144*tan(x/2)^2*(1296*a^10
*b^7 - 11016*a^11*b^6 + 27216*a^12*b^5 - 28512*a^13*b^4 + 12960*a^14*b^3 -
1944*a^15*b^2))/a^6) + (262144*tan(x/2)^2*(4104*a^9*b^7 - 16740*a^10*b^6
+ 18468*a^11*b^5 - 1836*a^12*b^4 - 5292*a^13*b^3 + 1296*a^14*b^2))/a^6) +
(262144*(288*a^7*b^8 - 1836*a^8*b^7 - 1692*a^9*b^6 + 6084*a^10*b^5 + 108*a
^11*b^4 - 4248*a^12*b^3 + 1296*a^13*b^2))/a^6 + (262144*tan(x/2)^2*(4392*a
^8*b^7 - 360*a^7*b^8 + 3366*a^9*b^6 - 29934*a^10*b^5 + 35946*a^11*b^4 - 15
354*a^12*b^3 + 1944*a^13*b^2))/a^6) - (262144*tan(x/2)^2*(72*a^5*b^9 - ...)
```

Reduce [F]

$$\int \frac{\tan^3(x)}{a + b \cos^3(x)} dx = \int \frac{\tan(x)^3}{\cos(x)^3 b + a} dx$$

input

```
int(tan(x)^3/(a+b*cos(x)^3),x)
```

output

```
int(tan(x)**3/(cos(x)**3*b + a),x)
```

3.49 $\int \sqrt{a + b \cos^3(x)} \tan(x) dx$

Optimal result	358
Mathematica [A] (verified)	358
Rubi [A] (verified)	359
Maple [F]	361
Fricas [A] (verification not implemented)	361
Sympy [F]	362
Maxima [A] (verification not implemented)	362
Giac [A] (verification not implemented)	363
Mupad [F(-1)]	363
Reduce [F]	363

Optimal result

Integrand size = 15, antiderivative size = 45

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = \frac{2}{3} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^3(x)}}{\sqrt{a}} \right) - \frac{2}{3} \sqrt{a + b \cos^3(x)}$$

output `2/3*a^(1/2)*arctanh((a+b*cos(x)^3)^(1/2)/a^(1/2))-2/3*(a+b*cos(x)^3)^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = \frac{2}{3} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^3(x)}}{\sqrt{a}} \right) - \frac{2}{3} \sqrt{a + b \cos^3(x)}$$

input `Integrate[Sqrt[a + b*Cos[x]^3]*Tan[x],x]`

output `(2*.Sqrt[a]*ArcTanh[Sqrt[a + b*Cos[x]^3]/Sqrt[a]])/3 - (2*.Sqrt[a + b*Cos[x]^3])/3`

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 25, 3709, 798, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(x) \sqrt{a + b \cos^3(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sqrt{a + b \sin(x + \frac{\pi}{2})^3}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\sqrt{b \sin(x + \frac{\pi}{2})^3 + a}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3709} \\
 & - \int \sqrt{b \cos^3(x) + a} \sec(x) d \cos(x) \\
 & \quad \downarrow \text{798} \\
 & -\frac{1}{3} \int \sqrt{b \cos^3(x) + a} \sec(x) d \cos^3(x) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{3} \left(-a \int \frac{\sec(x)}{\sqrt{b \cos^3(x) + a}} d \cos^3(x) - 2\sqrt{a + b \cos^3(x)} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{3} \left(-\frac{2a \int \frac{1}{\cos^6(x) - \frac{a}{b}} d \sqrt{b \cos^3(x) + a}}{b} - 2\sqrt{a + b \cos^3(x)} \right) \\
 & \quad \downarrow \text{221} \\
 & \frac{1}{3} \left(2\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^3(x)}}{\sqrt{a}} \right) - 2\sqrt{a + b \cos^3(x)} \right)
 \end{aligned}$$

input $\text{Int}[\text{Sqrt}[a + b \cos[x]^3] \tan[x], x]$

output $\frac{(2 \sqrt{a} \operatorname{ArcTanh}[\sqrt{a + b \cos[x]^3}/\sqrt{a}] - 2 \sqrt{a + b \cos[x]^3})}{3}$

Definitions of rubi rules used

rule 25 $\text{Int}[-(F[x]), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F[x], x], x]$

rule 60 $\text{Int}[(a_{\cdot} + b_{\cdot})^{(m_{\cdot})} ((c_{\cdot} + d_{\cdot})^{(n_{\cdot})}, x_{\text{Symbol}}) \rightarrow \text{Simp}[(a + b x)^{(m + 1)} ((c + d x)^{n/(b(m + n + 1))}, x) + \text{Simp}[n*((b*c - a*d)/(b*(m + n + 1))) \text{Int}[(a + b x)^m (c + d x)^{(n - 1)}, x], x]; \text{FreeQ}[\{a, b, c, d\}, x] \& \text{GtQ}[n, 0] \& \text{NeQ}[m + n + 1, 0] \& \text{!IGtQ}[m, 0] \& (\text{!IntegerQ}[n] \text{||} (\text{GtQ}[m, 0] \& \text{LtQ}[m - n, 0])) \& \text{!ILtQ}[m + n + 2, 0] \& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 73 $\text{Int}[(a_{\cdot} + b_{\cdot})^{(m_{\cdot})} ((c_{\cdot} + d_{\cdot})^{(n_{\cdot})}, x_{\text{Symbol}}) \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{Subst}[\text{Int}[x^{(p(m + 1) - 1)} (c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b x)^{(1/p)}, x]]]; \text{FreeQ}[\{a, b, c, d\}, x] \& \text{LtQ}[-1, m, 0] \& \text{LeQ}[-1, n, 0] \& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_{\cdot} + b_{\cdot})^{(x_{\cdot})^2}^{(-1)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \operatorname{ArcTanh}[x/\text{Rt}[-a/b, 2]], x]; \text{FreeQ}[\{a, b\}, x] \& \text{NegQ}[a/b]$

rule 798 $\text{Int}[(x_{\cdot})^{(m_{\cdot})} ((a_{\cdot} + b_{\cdot})^{(n_{\cdot})})^{(p_{\cdot})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[1/n \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)} (a + b x)^p, x], x, x^n], x]; \text{FreeQ}[\{a, b, m, n, p\}, x] \& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

rule 3042 $\text{Int}[u_{\cdot}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x]; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3709

```
Int[((a_) + (b_)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_)*tan[(e_.) + (f_.)*(x_)]^(m_), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Simplify[ff^(m + 1)/f Subst[Int[x^m*((a + b*(c*ff*x)^n)^p)/(1 - ff^2*x^2)^((m + 1)/2)), x], x, Sin[e + f*x]/ff], x]] /; FreeQ[{a, b, c, e, f, n, p}, x] && ILtQ[(m - 1)/2, 0]
```

Maple [F]

$$\int \sqrt{a + b \cos(x)^3} \tan(x) dx$$

input `int((a+b*cos(x)^3)^(1/2)*tan(x),x)`

output `int((a+b*cos(x)^3)^(1/2)*tan(x),x)`

Fricas [A] (verification not implemented)

Time = 1.74 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.98

$$\begin{aligned} & \int \sqrt{a + b \cos^3(x)} \tan(x) dx \\ &= \left[\frac{1}{6} \sqrt{a} \log \left(-\frac{b^2 \cos(x)^6 + 8ab \cos(x)^3 + 4(b \cos(x)^3 + 2a) \sqrt{b \cos(x)^3 + a} \sqrt{a} + 8a^2}{\cos(x)^6} \right) \right. \\ & \quad - \frac{2}{3} \sqrt{b \cos(x)^3 + a}, -\frac{1}{3} \sqrt{-a} \arctan \left(\frac{(b \cos(x)^3 + 2a) \sqrt{b \cos(x)^3 + a} \sqrt{-a}}{2(ab \cos(x)^3 + a^2)} \right) \\ & \quad \left. - \frac{2}{3} \sqrt{b \cos(x)^3 + a} \right] \end{aligned}$$

input `integrate((a+b*cos(x)^3)^(1/2)*tan(x),x, algorithm="fricas")`

output [1/6*sqrt(a)*log(-(b^2*cos(x)^6 + 8*a*b*cos(x)^3 + 4*(b*cos(x)^3 + 2*a)*sqrt(b*cos(x)^3 + a)*sqrt(a) + 8*a^2)/cos(x)^6) - 2/3*sqrt(b*cos(x)^3 + a), -1/3*sqrt(-a)*arctan(1/2*(b*cos(x)^3 + 2*a)*sqrt(b*cos(x)^3 + a)*sqrt(-a)/(a*b*cos(x)^3 + a^2)) - 2/3*sqrt(b*cos(x)^3 + a)]

Sympy [F]

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = \int \sqrt{a + b \cos^3(x)} \tan(x) dx$$

input integrate((a+b*cos(x)**3)**(1/2)*tan(x),x)

output Integral(sqrt(a + b*cos(x)**3)*tan(x), x)

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.16

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = -\frac{1}{3} \sqrt{a} \log \left(\frac{\sqrt{b \cos(x)^3 + a} - \sqrt{a}}{\sqrt{b \cos(x)^3 + a} + \sqrt{a}} \right) - \frac{2}{3} \sqrt{b \cos(x)^3 + a}$$

input integrate((a+b*cos(x)^3)^(1/2)*tan(x),x, algorithm="maxima")

output -1/3*sqrt(a)*log((sqrt(b*cos(x)^3 + a) - sqrt(a))/(sqrt(b*cos(x)^3 + a) + sqrt(a))) - 2/3*sqrt(b*cos(x)^3 + a)

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.84

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = -\frac{2a \arctan\left(\frac{\sqrt{b \cos(x)^3 + a}}{\sqrt{-a}}\right)}{3\sqrt{-a}} - \frac{2}{3} \sqrt{b \cos(x)^3 + a}$$

input `integrate((a+b*cos(x)^3)^(1/2)*tan(x),x, algorithm="giac")`

output `-2/3*a*arctan(sqrt(b*cos(x)^3 + a)/sqrt(-a))/sqrt(-a) - 2/3*sqrt(b*cos(x)^3 + a)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = \int \tan(x) \sqrt{b \cos(x)^3 + a} dx$$

input `int(tan(x)*(a + b*cos(x)^3)^(1/2),x)`

output `int(tan(x)*(a + b*cos(x)^3)^(1/2), x)`

Reduce [F]

$$\int \sqrt{a + b \cos^3(x)} \tan(x) dx = \int \sqrt{\cos(x)^3 b + a} \tan(x) dx$$

input `int((a+b*cos(x)^3)^(1/2)*tan(x),x)`

output `int(sqrt(cos(x)**3*b + a)*tan(x),x)`

3.50 $\int \frac{\tan(x)}{\sqrt{a+b\cos^3(x)}} dx$

Optimal result	364
Mathematica [A] (verified)	364
Rubi [A] (verified)	365
Maple [F]	367
Fricas [F(-2)]	367
Sympy [F]	367
Maxima [A] (verification not implemented)	368
Giac [A] (verification not implemented)	368
Mupad [F(-1)]	368
Reduce [F]	369

Optimal result

Integrand size = 15, antiderivative size = 28

$$\int \frac{\tan(x)}{\sqrt{a + b\cos^3(x)}} dx = \frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b\cos^3(x)}}{\sqrt{a}}\right)}{3\sqrt{a}}$$

output 2/3*arctanh((a+b*cos(x)^3)^(1/2)/a^(1/2))/a^(1/2)

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{a + b\cos^3(x)}} dx = \frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b\cos^3(x)}}{\sqrt{a}}\right)}{3\sqrt{a}}$$

input Integrate[Tan[x]/Sqrt[a + b*Cos[x]^3],x]

output (2*ArcTanh[Sqrt[a + b*Cos[x]^3]/Sqrt[a]])/(3*Sqrt[a])

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 25, 3709, 798, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\tan(x + \frac{\pi}{2}) \sqrt{a + b \sin(x + \frac{\pi}{2})^3}} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\sqrt{b \sin(x + \frac{\pi}{2})^3 + a \tan(x + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \textcolor{blue}{3709} \\
 & - \int \frac{\sec(x)}{\sqrt{b \cos^3(x) + a}} d \cos(x) \\
 & \quad \downarrow \textcolor{blue}{798} \\
 & - \frac{1}{3} \int \frac{\sec(x)}{\sqrt{b \cos^3(x) + a}} d \cos^3(x) \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & - \frac{2 \int \frac{1}{\frac{\cos^6(x)}{b} - \frac{a}{b}} d \sqrt{b \cos^3(x) + a}}{3b} \\
 & \quad \downarrow \textcolor{blue}{221} \\
 & \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^3(x)}}{\sqrt{a}}\right)}{3\sqrt{a}}
 \end{aligned}$$

input `Int[Tan[x]/Sqrt[a + b*Cos[x]^3],x]`

output $(2*\text{ArcTanh}[\sqrt{a + b*\cos[x]^3}/\sqrt{a}])/(3*\sqrt{a})$

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 73 $\text{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.)^{(n_.)})^{(p_.)}, \text{x_Symbol}] \rightarrow \text{With}[p = \text{Denominator}[m], \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{LtQ}[-1, m, 0] \&& \text{LeQ}[-1, n, 0] \&& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_.) + (b_.)*(x_.)^2)^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{NegQ}[a/b]$

rule 798 $\text{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[1/n \quad \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$

rule 3709 $\text{Int}[(a_.) + (b_.)*((c_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)})^{(p_.)}*\tan[(e_.) + (f_.)*(x_.)^{(m_.)}], \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Simp}[ff^{(m + 1)/f} \quad \text{Subst}[\text{Int}[x^{m*}((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2)^{(m + 1)/2}), x], x, \text{Sin}[e + f*x]/ff], x] /; \text{FreeQ}[\{a, b, c, e, f, n, p\}, x] \&& \text{ILtQ}[(m - 1)/2, 0]$

Maple [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx$$

input `int(tan(x)/(a+b*cos(x)^3)^(1/2),x)`

output `int(tan(x)/(a+b*cos(x)^3)^(1/2),x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(x)/(a+b*cos(x)^3)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: failed of mode Union(SparseUnivariatePolynomial(Expression(Complex(Integer))), failed) cannot be coerced to mode SparseUnivariatePolynomial(Expression(Complex(Int`

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx = \int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx$$

input `integrate(tan(x)/(a+b*cos(x)**3)**(1/2),x)`

output `Integral(tan(x)/sqrt(a + b*cos(x)**3), x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.39

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx = -\frac{\log \left(\frac{\sqrt{b \cos(x)^3 + a} - \sqrt{a}}{\sqrt{b \cos(x)^3 + a} + \sqrt{a}} \right)}{3 \sqrt{a}}$$

input `integrate(tan(x)/(a+b*cos(x)^3)^(1/2),x, algorithm="maxima")`

output `-1/3*log((sqrt(b*cos(x)^3 + a) - sqrt(a))/(sqrt(b*cos(x)^3 + a) + sqrt(a)))/sqrt(a)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx = -\frac{2 \arctan \left(\frac{\sqrt{b \cos(x)^3 + a}}{\sqrt{-a}} \right)}{3 \sqrt{-a}}$$

input `integrate(tan(x)/(a+b*cos(x)^3)^(1/2),x, algorithm="giac")`

output `-2/3*arctan(sqrt(b*cos(x)^3 + a)/sqrt(-a))/sqrt(-a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx = \int \frac{\tan(x)}{\sqrt{b \cos(x)^3 + a}} dx$$

input `int(tan(x)/(a + b*cos(x)^3)^(1/2),x)`

output `int(tan(x)/(a + b*cos(x)^3)^(1/2), x)`

Reduce [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^3(x)}} dx = \int \frac{\sqrt{\cos(x)^3 b + a} \tan(x)}{\cos(x)^3 b + a} dx$$

input `int(tan(x)/(a+b*cos(x)^3)^(1/2),x)`

output `int((sqrt(cos(x)**3*b + a)*tan(x))/(cos(x)**3*b + a),x)`

3.51 $\int \sqrt{a + b \cos^4(x)} \tan(x) dx$

Optimal result	370
Mathematica [A] (verified)	370
Rubi [A] (verified)	371
Maple [A] (verified)	373
Fricas [A] (verification not implemented)	374
Sympy [F]	374
Maxima [A] (verification not implemented)	375
Giac [A] (verification not implemented)	375
Mupad [F(-1)]	375
Reduce [F]	376

Optimal result

Integrand size = 15, antiderivative size = 45

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \frac{1}{2} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^4(x)}}{\sqrt{a}} \right) - \frac{1}{2} \sqrt{a + b \cos^4(x)}$$

output `1/2*a^(1/2)*arctanh((a+b*cos(x)^4)^(1/2)/a^(1/2))-1/2*(a+b*cos(x)^4)^(1/2)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \frac{1}{2} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^4(x)}}{\sqrt{a}} \right) - \frac{1}{2} \sqrt{a + b \cos^4(x)}$$

input `Integrate[Sqrt[a + b*Cos[x]^4]*Tan[x],x]`

output `(Sqrt[a]*ArcTanh[Sqrt[a + b*Cos[x]^4]/Sqrt[a]])/2 - Sqrt[a + b*Cos[x]^4]/2`

Rubi [A] (verified)

Time = 0.25 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 25, 3708, 243, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(x) \sqrt{a + b \cos^4(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\sqrt{a + b \sin(x + \frac{\pi}{2})^4}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{25} \\
 & - \int \frac{\sqrt{b \sin(x + \frac{\pi}{2})^4 + a}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \text{3708} \\
 & -\frac{1}{2} \int \sqrt{b \cos^4(x) + a} \sec^2(x) d \cos^2(x) \\
 & \quad \downarrow \text{243} \\
 & -\frac{1}{4} \int \sqrt{b \cos^4(x) + a} \sec^2(x) d \cos^4(x) \\
 & \quad \downarrow \text{60} \\
 & \frac{1}{4} \left(-a \int \frac{\sec^2(x)}{\sqrt{b \cos^4(x) + a}} d \cos^4(x) - 2\sqrt{a + b \cos^4(x)} \right) \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{4} \left(-\frac{2a \int \frac{1}{\sqrt{b \cos^4(x) + a} - \frac{a}{b}} d \sqrt{b \cos^4(x) + a}}{b} - 2\sqrt{a + b \cos^4(x)} \right) \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

$$\frac{1}{4} \left(2\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^4(x)}}{\sqrt{a}} \right) - 2\sqrt{a + b \cos^4(x)} \right)$$

input `Int[Sqrt[a + b*Cos[x]^4]*Tan[x], x]`

output `(2*Sqrt[a]*ArcTanh[Sqrt[a + b*Cos[x]^4]/Sqrt[a]] - 2*Sqrt[a + b*Cos[x]^4])/4`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 60 `Int[((a_.) + (b_.*(x_))^(m_)*((c_.) + (d_.*(x_))^(n_)), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.*(x_))^(m_)*((c_.) + (d_.*(x_))^(n_)), x_Symbol] :> With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.*((a_) + (b_.*(x_)^2)^(-1))), x_Symbol] :> Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3708 $\text{Int}[(a_ + b_)*\sin[(e_ + f_)*(x_)]^{(n_)}*(p_)*\tan[(e_ + f_)*(x_)]^{(m_)}, x_{\text{Symbol}}] \rightarrow \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Sin}[e + f*x]^2, x]\}, \text{Simp}[\text{ff}^{((m + 1)/2)/(2*f)} \text{Subst}[\text{Int}[x^{((m - 1)/2)}*((a + b*\text{ff}^{(n/2)}*x^{(n/2)})^p/(1 - \text{ff}*x)^{((m + 1)/2}), x], x, \text{Sin}[e + f*x]^{2/\text{ff}], x]] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{IntegerQ}[n/2]$

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

method	result	size
default	$-\frac{\sqrt{a+b \cos(x)^4}}{2} + \frac{\sqrt{a} \ln\left(\frac{2a+2\sqrt{a}\sqrt{a+b \cos(x)^4}}{\cos(x)^2}\right)}{2}$	44

input `int((a+b*cos(x)^4)^(1/2)*tan(x), x, method=_RETURNVERBOSE)`

output
$$\frac{-1/2*(a+b*\cos(x)^4)^(1/2)+1/2*a^(1/2)*\ln((2*a+2*a^(1/2)*(a+b*\cos(x)^4)^(1/2))/\cos(x)^2)}$$

Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.93

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \left[\frac{1}{4} \sqrt{a} \log \left(\frac{b \cos(x)^4 + 2 \sqrt{b \cos(x)^4 + a} \sqrt{a} + 2a}{\cos(x)^4} \right) \right.$$

$$- \frac{1}{2} \sqrt{b \cos(x)^4 + a},$$

$$- \frac{1}{2} \sqrt{-a} \arctan \left(\frac{\sqrt{-a}}{\sqrt{b \cos(x)^4 + a}} \right)$$

$$\left. - \frac{1}{2} \sqrt{b \cos(x)^4 + a} \right]$$

input `integrate((a+b*cos(x)^4)^(1/2)*tan(x),x, algorithm="fricas")`

output `[1/4*sqrt(a)*log((b*cos(x)^4 + 2*sqrt(b*cos(x)^4 + a)*sqrt(a) + 2*a)/cos(x)^4) - 1/2*sqrt(b*cos(x)^4 + a), -1/2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*cos(x)^4 + a)) - 1/2*sqrt(b*cos(x)^4 + a)]`

Sympy [F]

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \int \sqrt{a + b \cos^4(x)} \tan(x) dx$$

input `integrate((a+b*cos(x)**4)**(1/2)*tan(x),x)`

output `Integral(sqrt(a + b*cos(x)**4)*tan(x), x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \frac{1}{2} \sqrt{a} \operatorname{arsinh} \left(-\frac{a}{\sqrt{ab}(\sin(x)^2 - 1)} \right) - \frac{1}{2} \sqrt{b \sin(x)^4 - 2b \sin(x)^2 + a + b}$$

input `integrate((a+b*cos(x)^4)^(1/2)*tan(x),x, algorithm="maxima")`

output `1/2*sqrt(a)*arcsinh(-a/(sqrt(a*b)*(sin(x)^2 - 1))) - 1/2*sqrt(b*sin(x)^4 - 2*b*sin(x)^2 + a + b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.84

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = -\frac{a \arctan \left(\frac{\sqrt{b \cos(x)^4 + a}}{\sqrt{-a}} \right)}{2 \sqrt{-a}} - \frac{1}{2} \sqrt{b \cos(x)^4 + a}$$

input `integrate((a+b*cos(x)^4)^(1/2)*tan(x),x, algorithm="giac")`

output `-1/2*a*arctan(sqrt(b*cos(x)^4 + a)/sqrt(-a))/sqrt(-a) - 1/2*sqrt(b*cos(x)^4 + a)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \int \tan(x) \sqrt{b \cos(x)^4 + a} dx$$

input `int(tan(x)*(a + b*cos(x)^4)^(1/2),x)`

output `int(tan(x)*(a + b*cos(x)^4)^(1/2), x)`

Reduce [F]

$$\int \sqrt{a + b \cos^4(x)} \tan(x) dx = \int \sqrt{\cos(x)^4 b + a} \tan(x) dx$$

input `int((a+b*cos(x)^4)^(1/2)*tan(x),x)`

output `int(sqrt(cos(x)**4*b + a)*tan(x),x)`

3.52 $\int \frac{\tan(x)}{\sqrt{a+b\cos^4(x)}} dx$

Optimal result	377
Mathematica [A] (verified)	377
Rubi [A] (verified)	378
Maple [A] (verified)	380
Fricas [A] (verification not implemented)	380
Sympy [F]	381
Maxima [F]	381
Giac [A] (verification not implemented)	381
Mupad [F(-1)]	382
Reduce [F]	382

Optimal result

Integrand size = 15, antiderivative size = 28

$$\int \frac{\tan(x)}{\sqrt{a + b\cos^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\cos^4(x)}}{\sqrt{a}}\right)}{2\sqrt{a}}$$

output `1/2*arctanh((a+b*cos(x)^4)^(1/2)/a^(1/2))/a^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{a + b\cos^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\cos^4(x)}}{\sqrt{a}}\right)}{2\sqrt{a}}$$

input `Integrate[Tan[x]/Sqrt[a + b*Cos[x]^4],x]`

output `ArcTanh[Sqrt[a + b*Cos[x]^4]/Sqrt[a]]/(2*Sqrt[a])`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 25, 3708, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\tan(x + \frac{\pi}{2}) \sqrt{a + b \sin(x + \frac{\pi}{2})^4}} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\sqrt{b \sin(x + \frac{\pi}{2})^4 + a \tan(x + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \textcolor{blue}{3708} \\
 & -\frac{1}{2} \int \frac{\sec^2(x)}{\sqrt{b \cos^4(x) + a}} d \cos^2(x) \\
 & \quad \downarrow \textcolor{blue}{243} \\
 & -\frac{1}{4} \int \frac{\sec^2(x)}{\sqrt{b \cos^4(x) + a}} d \cos^4(x) \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & -\frac{\int \frac{1}{\sqrt{\frac{b \cos^4(x) + a}{b} - \frac{a}{b}}} d \sqrt{b \cos^4(x) + a}}{2b} \\
 & \quad \downarrow \textcolor{blue}{221} \\
 & \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^4(x)}}{\sqrt{a}}\right)}{2\sqrt{a}}
 \end{aligned}$$

input

output $\text{ArcTanh}[\sqrt{a + b \cos[x]^4}/\sqrt{a}]/(2\sqrt{a})$

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 73 $\text{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}), \text{x_Symbol}] \rightarrow \text{With}[$
 $\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{LtQ}[-1, m, 0] \&& \text{LeQ}[-1, n, 0] \&& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{NegQ}[a/b]$

rule 243 $\text{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^2)^p, \text{x_Symbol}] \rightarrow \text{Simp}[1/2 \quad \text{Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p, x}, x, x^2], x] /; \text{FreeQ}[\{a, b, m, p\}, x] \&& \text{IntegerQ}[(m - 1)/2]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$

rule 3708 $\text{Int}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^n)^p, \text{x_Symbol}] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x]^2, x]\}, \text{Simp}[ff^{((m + 1)/2)/(2*f)} \quad \text{Subst}[\text{Int}[x^{((m - 1)/2)*((a + b*ff^{(n/2)}*x^{(n/2)})^p/(1 - ff*x)^{((m + 1)/2))}, x}, x, \text{Sin}[e + f*x]^2/ff], x]] /; \text{FreeQ}[\{a, b, e, f, p\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{IntegerQ}[n/2]$

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.11

method	result	size
default	$\frac{\ln\left(\frac{2a+2\sqrt{a}\sqrt{a+b\cos(x)^4}}{\cos(x)^2}\right)}{2\sqrt{a}}$	31

input `int(tan(x)/(a+b*cos(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2/a^(1/2)*ln((2*a+2*a^(1/2)*(a+b*cos(x)^4)^(1/2))/cos(x)^2)`

Fricas [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.29

$$\begin{aligned} & \int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx \\ &= \left[\frac{\log\left(\frac{b \cos(x)^4 + 2\sqrt{b \cos(x)^4 + a} \sqrt{a + 2a}}{\cos(x)^4}\right)}{4\sqrt{a}}, -\frac{\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{b \cos(x)^4 + a}}\right)}{2a} \right] \end{aligned}$$

input `integrate(tan(x)/(a+b*cos(x)^4)^(1/2),x, algorithm="fricas")`

output `[1/4*log((b*cos(x)^4 + 2*sqrt(b*cos(x)^4 + a)*sqrt(a) + 2*a)/cos(x)^4)/sqrt(a), -1/2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*cos(x)^4 + a))/a]`

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx = \int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx$$

input `integrate(tan(x)/(a+b*cos(x)**4)**(1/2),x)`

output `Integral(tan(x)/sqrt(a + b*cos(x)**4), x)`

Maxima [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx = \int \frac{\tan(x)}{\sqrt{b \cos(x)^4 + a}} dx$$

input `integrate(tan(x)/(a+b*cos(x)^4)^(1/2),x, algorithm="maxima")`

output `integrate(tan(x)/sqrt(b*cos(x)^4 + a), x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx = -\frac{\arctan\left(\frac{\sqrt{b \cos(x)^4 + a}}{\sqrt{-a}}\right)}{2\sqrt{-a}}$$

input `integrate(tan(x)/(a+b*cos(x)^4)^(1/2),x, algorithm="giac")`

output `-1/2*arctan(sqrt(b*cos(x)^4 + a)/sqrt(-a))/sqrt(-a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx = \int \frac{\tan(x)}{\sqrt{b \cos(x)^4 + a}} dx$$

input `int(tan(x)/(a + b*cos(x)^4)^(1/2),x)`

output `int(tan(x)/(a + b*cos(x)^4)^(1/2), x)`

Reduce [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^4(x)}} dx = \int \frac{\sqrt{\cos(x)^4 b + a} \tan(x)}{\cos(x)^4 b + a} dx$$

input `int(tan(x)/(a+b*cos(x)^4)^(1/2),x)`

output `int((sqrt(cos(x)**4*b + a)*tan(x))/(cos(x)**4*b + a),x)`

3.53 $\int \sqrt{a + b \cos^n(x)} \tan(x) dx$

Optimal result	383
Mathematica [A] (verified)	383
Rubi [A] (verified)	384
Maple [A] (verified)	386
Fricas [A] (verification not implemented)	386
Sympy [F]	387
Maxima [A] (verification not implemented)	387
Giac [A] (verification not implemented)	388
Mupad [F(-1)]	388
Reduce [F]	388

Optimal result

Integrand size = 15, antiderivative size = 47

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = \frac{2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^n(x)}}{\sqrt{a}}\right)}{n} - \frac{2\sqrt{a + b \cos^n(x)}}{n}$$

output 2*a^(1/2)*arctanh((a+b*cos(x)^n)^(1/2)/a^(1/2))/n-2*(a+b*cos(x)^n)^(1/2)/n

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = -\frac{-2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^n(x)}}{\sqrt{a}}\right) + 2\sqrt{a + b \cos^n(x)}}{n}$$

input Integrate[Sqrt[a + b*Cos[x]^n]*Tan[x],x]

output -((-2*Sqrt[a]*ArcTanh[Sqrt[a + b*Cos[x]^n]/Sqrt[a]] + 2*Sqrt[a + b*Cos[x]^n])/n)

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3042, 25, 3709, 798, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(x) \sqrt{a + b \cos^n(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{\sqrt{a + b \sin(x + \frac{\pi}{2})^n}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{\sqrt{b \sin(x + \frac{\pi}{2})^n + a}}{\tan(x + \frac{\pi}{2})} dx \\
 & \quad \downarrow \textcolor{blue}{3709} \\
 & - \int \sqrt{b \cos^n(x) + a} \sec(x) d \cos(x) \\
 & \quad \downarrow \textcolor{blue}{798} \\
 & - \frac{\int \sqrt{b \cos^n(x) + a} \sec(x) d \cos^n(x)}{n} \\
 & \quad \downarrow \textcolor{blue}{60} \\
 & - \frac{a \int \frac{\sec(x)}{\sqrt{b \cos^n(x) + a}} d \cos^n(x) + 2 \sqrt{a + b \cos^n(x)}}{n} \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & - \frac{2a \int \frac{1}{\frac{\cos^{2n}(x)}{b} - \frac{a}{b}} d \sqrt{b \cos^n(x) + a}}{n} + 2 \sqrt{a + b \cos^n(x)} \\
 & \quad \downarrow \textcolor{blue}{221} \\
 & - \frac{2 \sqrt{a + b \cos^n(x)} - 2 \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^n(x)}}{\sqrt{a}}\right)}{n}
 \end{aligned}$$

input $\text{Int}[\text{Sqrt}[a + b \cdot \text{Cos}[x]^n] \cdot \text{Tan}[x], x]$

output $-\left(\frac{(-2 \cdot \text{Sqrt}[a] \cdot \text{ArcTanh}[\text{Sqrt}[a + b \cdot \text{Cos}[x]^n]/\text{Sqrt}[a]] + 2 \cdot \text{Sqrt}[a + b \cdot \text{Cos}[x]^n])/n}{n}\right)$

Definitions of rubi rules used

rule 25 $\text{Int}[-(F_{x_}), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 60 $\text{Int}[(a_.) + (b_.) \cdot (x_)^{(m_)} \cdot ((c_.) + (d_.) \cdot (x_)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot x)^{(m + 1)} \cdot ((c + d \cdot x)^{n}/(b \cdot (m + n + 1))), x] + \text{Simp}[n \cdot ((b \cdot c - a \cdot d)/(b \cdot (m + n + 1))) \cdot \text{Int}[(a + b \cdot x)^{m} \cdot (c + d \cdot x)^{n - 1}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{GtQ}[n, 0] \&& \text{NeQ}[m + n + 1, 0] \&& !(\text{IGtQ}[m, 0] \&& (\text{!IntegerQ}[n] \text{ || } (\text{GtQ}[m, 0] \&& \text{LtQ}[m - n, 0]))) \&& \text{!ILtQ}[m + n + 2, 0] \&& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 73 $\text{Int}[(a_.) + (b_.) \cdot (x_)^{(m_)} \cdot ((c_.) + (d_.) \cdot (x_)^{(n_)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \cdot \text{Subst}[\text{Int}[x^{(p \cdot (m + 1) - 1)} \cdot (c - a \cdot (d/b) + d \cdot (x^{p/b})^n), x], x, (a + b \cdot x)^{(1/p)}], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{LtQ}[-1, m, 0] \&& \text{LeQ}[-1, n, 0] \&& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_.) + (b_.) \cdot (x_)^2 \cdot (-1), x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{NegQ}[a/b]$

rule 798 $\text{Int}[(x_)^{(m_)} \cdot ((a_.) + (b_.) \cdot (x_)^{(n_)} \cdot (p_)), x_Symbol] \rightarrow \text{Simp}[1/n \cdot \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)} \cdot (a + b \cdot x)^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3709

```
Int[((a_) + (b_)*((c_)*sin[(e_.) + (f_)*(x_)])^(n_))^(p_)*tan[(e_.) + (f_.)*(x_)]^m_, x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Si
mp[ff^(m + 1)/f Subst[Int[x^m*((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2)^(m + 1/2)), x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] &&
ILtQ[(m - 1)/2, 0]
```

Maple [A] (verified)

Time = 0.29 (sec), antiderivative size = 39, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{2\sqrt{a+b \cos(x)^n}-2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos(x)^n}}{\sqrt{a}}\right)}{n}$	39
default	$-\frac{2\sqrt{a+b \cos(x)^n}-2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos(x)^n}}{\sqrt{a}}\right)}{n}$	39

input `int((a+b*cos(x)^n)^(1/2)*tan(x), x, method=_RETURNVERBOSE)`output `-1/n*(2*(a+b*cos(x)^n)^(1/2)-2*a^(1/2)*arctanh((a+b*cos(x)^n)^(1/2)/a^(1/2)))`**Fricas [A] (verification not implemented)**

Time = 0.11 (sec), antiderivative size = 94, normalized size of antiderivative = 2.00

$$\begin{aligned} & \int \sqrt{a + b \cos^n(x)} \tan(x) dx \\ &= \left[\frac{\sqrt{a} \log \left(\frac{b \cos(x)^n + 2 \sqrt{b \cos(x)^n + a} \sqrt{a} + 2 a}{\cos(x)^n} \right) - 2 \sqrt{b \cos(x)^n + a}}{n}, \right. \\ & \quad \left. - \frac{2 \left(\sqrt{-a} \arctan \left(\frac{\sqrt{-a}}{\sqrt{b \cos(x)^n + a}} \right) + \sqrt{b \cos(x)^n + a} \right)}{n} \right] \end{aligned}$$

input `integrate((a+b*cos(x)^n)^(1/2)*tan(x), x, algorithm="fricas")`

output $[(\sqrt{a} \cdot \log((b \cdot \cos(x))^n + 2 \cdot \sqrt{b \cdot \cos(x)^n + a}) \cdot \sqrt{a} + 2 \cdot a) / \cos(x)^n - 2 \cdot \sqrt{b \cdot \cos(x)^n + a}) / n, -2 \cdot (\sqrt{-a} \cdot \arctan(\sqrt{-a} / \sqrt{b \cdot \cos(x)^n + a}) + \sqrt{b \cdot \cos(x)^n + a}) / n]$

Sympy [F]

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = \int \sqrt{a + b \cos^n(x)} \tan(x) dx$$

input `integrate((a+b*cos(x)**n)**(1/2)*tan(x),x)`

output `Integral(sqrt(a + b*cos(x)**n)*tan(x), x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.23

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = -\frac{\sqrt{a} \log \left(\frac{\sqrt{b \cos(x)^n + a} - \sqrt{a}}{\sqrt{b \cos(x)^n + a} + \sqrt{a}} \right)}{n} - \frac{2 \sqrt{b \cos(x)^n + a}}{n}$$

input `integrate((a+b*cos(x)^n)^(1/2)*tan(x),x, algorithm="maxima")`

output $-\sqrt{a} \cdot \log((\sqrt{b \cdot \cos(x)^n + a} - \sqrt{a}) / (\sqrt{b \cdot \cos(x)^n + a} + \sqrt{a})) / n - 2 \cdot \sqrt{b \cdot \cos(x)^n + a} / n$

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = -\frac{2 \left(\frac{ab \arctan\left(\frac{\sqrt{b \cos(x)^n + a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + \sqrt{b \cos(x)^n + ab} \right)}{bn}$$

input `integrate((a+b*cos(x)^n)^(1/2)*tan(x),x, algorithm="giac")`

output `-2*(a*b*arctan(sqrt(b*cos(x)^n + a)/sqrt(-a))/sqrt(-a) + sqrt(b*cos(x)^n + a)*b)/(b*n)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = \int \tan(x) \sqrt{a + b \cos(x)^n} dx$$

input `int(tan(x)*(a + b*cos(x)^n)^(1/2),x)`

output `int(tan(x)*(a + b*cos(x)^n)^(1/2), x)`

Reduce [F]

$$\int \sqrt{a + b \cos^n(x)} \tan(x) dx = \int \sqrt{\cos(x)^n b + a} \tan(x) dx$$

input `int((a+b*cos(x)^n)^(1/2)*tan(x),x)`

output `int(sqrt(cos(x)**n*b + a)*tan(x),x)`

3.54 $\int \frac{\tan(x)}{\sqrt{a+b \cos^n(x)}} dx$

Optimal result	389
Mathematica [A] (verified)	389
Rubi [A] (verified)	390
Maple [A] (verified)	392
Fricas [A] (verification not implemented)	392
Sympy [F]	393
Maxima [A] (verification not implemented)	393
Giac [A] (verification not implemented)	393
Mupad [F(-1)]	394
Reduce [F]	394

Optimal result

Integrand size = 15, antiderivative size = 29

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^n(x)}}{\sqrt{a}} \right)}{\sqrt{an}}$$

output `2*arctanh((a+b*cos(x)^n)^(1/2)/a^(1/2))/a^(1/2)/n`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a + b \cos^n(x)}}{\sqrt{a}} \right)}{\sqrt{an}}$$

input `Integrate[Tan[x]/Sqrt[a + b*Cos[x]^n],x]`

output `(2*ArcTanh[Sqrt[a + b*Cos[x]^n]/Sqrt[a]])/(Sqrt[a]*n)`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 25, 3709, 798, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int -\frac{1}{\tan(x + \frac{\pi}{2}) \sqrt{a + b \sin(x + \frac{\pi}{2})^n}} dx \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & - \int \frac{1}{\sqrt{b \sin(x + \frac{\pi}{2})^n + a \tan(x + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \textcolor{blue}{3709} \\
 & - \int \frac{\sec(x)}{\sqrt{b \cos^n(x) + a}} d \cos(x) \\
 & \quad \downarrow \textcolor{blue}{798} \\
 & - \frac{\int \frac{\sec(x)}{\sqrt{b \cos^n(x) + a}} d \cos^n(x)}{n} \\
 & \quad \downarrow \textcolor{blue}{73} \\
 & - \frac{2 \int \frac{1}{\frac{\cos^{2n}(x)}{b} - \frac{a}{b}} d \sqrt{b \cos^n(x) + a}}{bn} \\
 & \quad \downarrow \textcolor{blue}{221} \\
 & \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos^n(x)}}{\sqrt{a}}\right)}{\sqrt{an}}
 \end{aligned}$$

input `Int[Tan[x]/Sqrt[a + b*Cos[x]^n], x]`

output $(2*\text{ArcTanh}[\sqrt{a + b*\cos[x]^n}/\sqrt{a}])/(\sqrt{a}*n)$

Definitions of rubi rules used

rule 25 $\text{Int}[-(F_{x_}), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 73 $\text{Int}[(a_ + b_*(x_))^{(m_)}*((c_ + d_)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{With}[p = \text{Denominator}[m], \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^{p/b}))^n, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{LtQ}[-1, m, 0] \&& \text{LeQ}[-1, n, 0] \&& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&& \text{IntLinearQ}[a, b, c, d, m, n, x]]$

rule 221 $\text{Int}[(a_ + b_*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{NegQ}[a/b]$

rule 798 $\text{Int}[(x_)^{(m_)}*((a_ + b_)*(x_))^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/n \quad \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3709 $\text{Int}[(a_ + b_*((c_ + f_)*(x_)))^{(n_)}*(e_ + f_*(x_))^{(p_)}*\tan[e_ + f_*(x_)]^{(m_)}, x_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Simp}[ff^{(m + 1)/f} \quad \text{Subst}[\text{Int}[x^{m*((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2))^{((m + 1)/2)}, x], x, \text{Sin}[e + f*x]/ff], x] /; \text{FreeQ}[\{a, b, c, e, f, n, p\}, x] \&& \text{ILtQ}[(m - 1)/2, 0]]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos(x)^n}}{\sqrt{a}}\right)}{\sqrt{a} n}$	24
default	$\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \cos(x)^n}}{\sqrt{a}}\right)}{\sqrt{a} n}$	24

input `int(tan(x)/(a+b*cos(x)^n)^(1/2),x,method=_RETURNVERBOSE)`

output `2*arctanh((a+b*cos(x)^n)^(1/2)/a^(1/2))/a^(1/2)/n`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 71, normalized size of antiderivative = 2.45

$$\begin{aligned} & \int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx \\ &= \left[\frac{\log\left(\frac{b \cos(x)^n + 2 \sqrt{b \cos(x)^n + a} \sqrt{a} + 2 a}{\cos(x)^n}\right)}{\sqrt{a} n}, -\frac{2 \sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{b \cos(x)^n + a}}\right)}{a n} \right] \end{aligned}$$

input `integrate(tan(x)/(a+b*cos(x)^n)^(1/2),x, algorithm="fricas")`

output `[log((b*cos(x)^n + 2*sqrt(b*cos(x)^n + a)*sqrt(a) + 2*a)/cos(x)^n)/(sqrt(a)*n), -2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*cos(x)^n + a))/(a*n)]`

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = \int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx$$

input `integrate(tan(x)/(a+b*cos(x)**n)**(1/2),x)`

output `Integral(tan(x)/sqrt(a + b*cos(x)**n), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.45

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = -\frac{\log\left(\frac{\sqrt{b \cos(x)^n + a} - \sqrt{a}}{\sqrt{b \cos(x)^n + a} + \sqrt{a}}\right)}{\sqrt{a n}}$$

input `integrate(tan(x)/(a+b*cos(x)^n)^(1/2),x, algorithm="maxima")`

output `-log((sqrt(b*cos(x)^n + a) - sqrt(a))/(sqrt(b*cos(x)^n + a) + sqrt(a)))/(sqrt(a)*n)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = -\frac{2 \arctan\left(\frac{\sqrt{b \cos(x)^n + a}}{\sqrt{-a}}\right)}{\sqrt{-a n}}$$

input `integrate(tan(x)/(a+b*cos(x)^n)^(1/2),x, algorithm="giac")`

output `-2*arctan(sqrt(b*cos(x)^n + a)/sqrt(-a))/(sqrt(-a)*n)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = \int \frac{\tan(x)}{\sqrt{a + b \cos(x)^n}} dx$$

input `int(tan(x)/(a + b*cos(x)^n)^(1/2),x)`

output `int(tan(x)/(a + b*cos(x)^n)^(1/2), x)`

Reduce [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cos^n(x)}} dx = \int \frac{\sqrt{\cos(x)^n b + a} \tan(x)}{\cos(x)^n b + a} dx$$

input `int(tan(x)/(a+b*cos(x)^n)^(1/2),x)`

output `int((sqrt(cos(x)**n*b + a)*tan(x))/(cos(x)**n*b + a),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	395
4.2 Links to plain text integration problems used in this report for each CAS .	413

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```
(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*) (*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","");
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
    ,(*ELSE*) (*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "}
      ,
      finalresult={"C","Result contains higher order function than in optimal. Order "}
    ]
  ]
]
]
```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```



```

ExpnType[expn_] :=
If[AtomQ[expn],
  1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn] === Power,
      If[IntegerQ[expn[[2]]],
        ExpnType[expn[[1]]],
        If[Head[expn[[2]]] === Rational,
          If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
            1,
            Max[ExpnType[expn[[1]]], 2]],
          Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
      If[Head[expn] === Plus || Head[expn] === Times,
        Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
        If[ElementaryFunctionQ[Head[expn]],
          Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], ExpnType[expn[[3]]]]]]]]]]]

```

```
Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],  
If[AppellFunctionQ[Head[expn]],  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],  
If[Head[expn]==RootSum,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],  
If[Head[expn]==Integrate || Head[expn]==Int,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],  
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=  
MemberQ[{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
} , func]
```

```
SpecialFunctionQ[func_] :=  
MemberQ[{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  

```

```
HypergeometricFunctionQ[func_] :=  
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=  
MemberQ[{AppellF1}, func]
```

Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
```

```
#      antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
            # this assumes optimal do not as well. No check is needed here.
            if debug then
                print("result do not contain complex, this assumes optimal do not as well");
            fi;
```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                           convert(leaf_count_result,string),\"$ vs. \$2(", 
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_co
                           fi;
            fi;
        else #ExpnType(result) > ExpnType(optimal)
            if debug then
                print("ExpnType(result) > ExpnType(optimal)");
            fi;
            return "C",cat("Result contains higher order function than in optimal. Order ",
                           convert(ExpnType_result,string)," vs. order ",
                           convert(ExpnType_optimal,string),"."));
        fi;

    end proc:

    #

    # is_contains_complex(result)
    # takes expressions and returns true if it contains "I" else false
    #
    #Nasser 032417
    is_contains_complex:= proc(expression)
        return (has(expression,I));
    end proc:

    # The following summarizes the type number assigned an expression
    # based on the functions it involves
    # 1 = rational function
    # 2 = algebraic function
    # 3 = elementary function
    # 4 = special function
    # 5 = hypergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'`^`') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then
            if type(op(1,expn),'rational') then
                1
            else
                max(2,ExpnType(op(1,expn)))
            end if
        else
            max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
        end if
    elif type(expn,'`+`) or type(expn,'`*`) then
        max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif ElementaryFunctionQ(op(0,expn)) then
        max(3,ExpnType(op(1,expn)))
    elif SpecialFunctionQ(op(0,expn)) then
        max(4,apply(max,map(ExpnType,[op(expn)])))
    elif HypergeometricFunctionQ(op(0,expn)) then
        max(5,apply(max,map(ExpnType,[op(expn)])))
    elif AppellFunctionQ(op(0,expn)) then
        max(6,apply(max,map(ExpnType,[op(expn)])))
    elif op(0,expn)='int' then
        max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
```

```
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arccsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]
```

```
def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*')
```

```

m1 = expnType(expn.args[0])
m2 = expnType(list(expn.args[1:]))
return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sagemath")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(grade)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```
from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                       ]
    if debug:
```

```

if m:
    print ("func ", func , " is elementary_function")
else:
    print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral'
'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
'elliptic_pi','exp_integral_e','log_integral',
'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
'weierstrassPPrime','weierstrassSigma']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):
    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0], Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1], Integer)

```

```

    return expnType(expn.operands()[0])  #expnType(expn.args[0])
elif type(expn.operands()[1]) == Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0]) == Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))  #max(3,expnType(expn.args[0]),expnType(expn.args[1]))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(4,m1)  #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(5,m1)  #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation ="none"
            else:
                grade = "B"
                grade_annotation ="Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation ="Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation ="none"
        else:
            grade = "B"
            grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(expnType_result)+"/"+str(expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file