

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.3-Tangent/219-4.3.4.2

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 8:06am

Contents

1	Introduction	8
1.1	Listing of CAS systems tested	9
1.2	Results	10
1.3	Time and leaf size Performance	14
1.4	Performance based on number of rules Rubi used	16
1.5	Performance based on number of steps Rubi used	17
1.6	Solved integrals histogram based on leaf size of result	18
1.7	Solved integrals histogram based on CPU time used	19
1.8	Leaf size vs. CPU time used	20
1.9	list of integrals with no known antiderivative	21
1.10	List of integrals solved by CAS but has no known antiderivative	21
1.11	list of integrals solved by CAS but failed verification	21
1.12	Timing	22
1.13	Verification	22
1.14	Important notes about some of the results	23
1.15	Current tree layout of integration tests	26
1.16	Design of the test system	27
2	detailed summary tables of results	28
2.1	List of integrals sorted by grade for each CAS	29
2.2	Detailed conclusion table per each integral for all CAS systems	34
2.3	Detailed conclusion table specific for Rubi results	77
3	Listing of integrals	83
3.1	$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$	89
3.2	$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$	97
3.3	$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$.	104
3.4	$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$	111
3.5	$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$	118
3.6	$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$	125
3.7	$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$	133

3.8	$\int \cot^6(c+dx)(a+b \tan(c+dx))(B \tan(c+dx)+C \tan^2(c+dx)) dx$	143
3.9	$\int \tan(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	153
3.10	$\int (a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	163
3.11	$\int \cot(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	171
3.12	$\int \cot^2(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	179
3.13	$\int \cot^3(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	187
3.14	$\int \cot^4(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	195
3.15	$\int \cot^5(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	204
3.16	$\int \cot^6(c+dx)(a+b \tan(c+dx))^2(B \tan(c+dx)+C \tan^2(c+dx)) dx$	214
3.17	$\int (a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	225
3.18	$\int \cot(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	234
3.19	$\int \cot^2(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	243
3.20	$\int \cot^3(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	252
3.21	$\int \cot^4(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	262
3.22	$\int \cot^5(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	272
3.23	$\int \cot^6(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	283
3.24	$\int \cot^7(c+dx)(a+b \tan(c+dx))^3(B \tan(c+dx)+C \tan^2(c+dx)) dx$	295
3.25	$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	307
3.26	$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	317
3.27	$\int \frac{B \tan(c+dx)+C \tan^2(c+dx)}{a+b \tan(c+dx)} dx$	326
3.28	$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	333
3.29	$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	341
3.30	$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	349
3.31	$\int \frac{\cot^4(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	358
3.32	$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	369
3.33	$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	381
3.34	$\int \frac{B \tan(c+dx)+C \tan^2(c+dx)}{(a+b \tan(c+dx))^2} dx$	392
3.35	$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	400
3.36	$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	408
3.37	$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	419
3.38	$\int \frac{\tan^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	430
3.39	$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	443
3.40	$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	454
3.41	$\int \frac{B \tan(c+dx)+C \tan^2(c+dx)}{(a+b \tan(c+dx))^3} dx$	464

3.42	$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	473
3.43	$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	483
3.44	$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	495
3.45	$\int \tan^2(c+dx)(b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) dx$	508
3.46	$\int \tan^m(c+dx)(b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) dx$	515
3.47	$\int \tan^m(c+dx)\sqrt{b \tan(c+dx)}(A+B \tan(c+dx)+C \tan^2(c+dx)) dx$	522
3.48	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{b \tan(c+dx)}} dx$	529
3.49	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	537
3.50	$\int (a+b \tan(e+fx))^3(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	544
3.51	$\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	556
3.52	$\int (a+b \tan(e+fx))(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	567
3.53	$\int (c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	576
3.54	$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	583
3.55	$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	592
3.56	$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	603
3.57	$\int (a+b \tan(e+fx))^3(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	614
3.58	$\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	629
3.59	$\int (a+b \tan(e+fx))(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	642
3.60	$\int (c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	652
3.61	$\int \frac{(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	661
3.62	$\int \frac{(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	673
3.63	$\int \frac{(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	686
3.64	$\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	699
3.65	$\int (a+b \tan(e+fx))(c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	713
3.66	$\int (c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	725
3.67	$\int \frac{(c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	735
3.68	$\int \frac{(c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	749
3.69	$\int \frac{(c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	764
3.70	$\int \frac{(a+b \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	778
3.71	$\int \frac{(a+b \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	791
3.72	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	803
3.73	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{c+d \tan(e+fx)} dx$	812
3.74	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))} dx$	820
3.75	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))} dx$	829

3.76	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^3(c+d \tan(e+fx))} dx \dots \dots \dots$	840
3.77	$\int \frac{(a+b \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx \dots \dots \dots$	852
3.78	$\int \frac{(a+b \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx \dots \dots \dots$	867
3.79	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx \dots \dots \dots$	880
3.80	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^2} dx \dots \dots \dots$	891
3.81	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx \dots \dots \dots$	900
3.82	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^2} dx \dots \dots \dots$	911
3.83	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^3(c+d \tan(e+fx))^2} dx \dots \dots \dots$	923
3.84	$\int \frac{(a+b \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx \dots \dots \dots$	936
3.85	$\int \frac{(a+b \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx \dots \dots \dots$	950
3.86	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx \dots \dots \dots$	963
3.87	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^3} dx \dots \dots \dots$	975
3.88	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^3} dx \dots \dots \dots$	985
3.89	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^3} dx \dots \dots \dots$	997
3.90	$\int (a+b \tan(e+fx))^3 \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1010
3.91	$\int (a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1023
3.92	$\int (a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1035
3.93	$\int \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots \dots$	1046
3.94	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx \dots \dots \dots$	1057
3.95	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx \dots \dots \dots$	1068
3.96	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx \dots \dots \dots$	1079
3.97	$\int (a+b \tan(e+fx))^3(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1092
3.98	$\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1106
3.99	$\int (a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1119
3.100	$\int (c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots \dots$	1129
3.101	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx \dots \dots \dots$	1139
3.102	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx \dots \dots \dots$	1151
3.103	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx \dots \dots \dots$	1164
3.104	$\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1176
3.105	$\int (a+b \tan(e+fx))(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots$	1188
3.106	$\int (c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx \dots \dots$	1200
3.107	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx \dots \dots \dots$	1211
3.108	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx \dots \dots \dots$	1224

3.109	$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	1237
3.110	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1250
3.111	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1262
3.112	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1273
3.113	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx$	1284
3.114	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx$	1293
3.115	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}} dx$	1303
3.116	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1315
3.117	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1328
3.118	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1340
3.119	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^{3/2}} dx$	1350
3.120	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2}} dx$	1359
3.121	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{3/2}} dx$	1370
3.122	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1383
3.123	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1396
3.124	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1407
3.125	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^{5/2}} dx$	1417
3.126	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{5/2}} dx$	1426
3.127	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{5/2}} dx$	1438
3.128	$\int (a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1451
3.129	$\int (a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1463
3.130	$\int \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1474
3.131	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$	1483
3.132	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$	1491
3.133	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$	1499
3.134	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$	1509
3.135	$\int (a+b \tan(e+fx))^{3/2} (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1521
3.136	$\int \sqrt{a+b \tan(e+fx)} (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1533
3.137	$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$	1544
3.138	$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$	1553
3.139	$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$	1564

3.140	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$	1574
3.141	$\int \sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1586
3.142	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$	1598
3.143	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$	1609
3.144	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$	1620
3.145	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$	1631
3.146	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{9/2}} dx$	1641
3.147	$\int \frac{(a+b \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1654
3.148	$\int \frac{(a+b \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1664
3.149	$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1673
3.150	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$	1681
3.151	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2}\sqrt{c+d \tan(e+fx)}} dx$	1689
3.152	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{5/2}\sqrt{c+d \tan(e+fx)}} dx$	1698
3.153	$\int \frac{(a+b \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1708
3.154	$\int \frac{(a+b \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1719
3.155	$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1729
3.156	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx$	1737
3.157	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}} dx$	1746
3.158	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{5/2}(c+d \tan(e+fx))^{3/2}} dx$	1756
3.159	$\int \frac{(a+b \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1767
3.160	$\int \frac{(a+b \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1778
3.161	$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1788
3.162	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}} dx$	1798
3.163	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{5/2}} dx$	1808
3.164	$\int (a+b \tan(e+fx))^m(c+d \tan(e+fx))^n(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1819
3.165	$\int (a+b \tan(e+fx))^m(c+d \tan(e+fx))^3(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1826
3.166	$\int (a+b \tan(e+fx))^m(c+d \tan(e+fx))^2(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1839
3.167	$\int (a+b \tan(e+fx))^m(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1849
3.168	$\int (a+b \tan(e+fx))^m(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1858
3.169	$\int \frac{(a+b \tan(e+fx))^m(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	1865
3.170	$\int \frac{(a+b \tan(e+fx))^m(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$	1874

3.171	$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$	1884
4	Appendix	1897
4.1	Listing of Grading functions	1897
4.2	Links to plain text integration problems used in this report for each CAS	1915

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	9
1.2	Results	10
1.3	Time and leaf size Performance	14
1.4	Performance based on number of rules Rubi used	16
1.5	Performance based on number of steps Rubi used	17
1.6	Solved integrals histogram based on leaf size of result	18
1.7	Solved integrals histogram based on CPU time used	19
1.8	Leaf size vs. CPU time used	20
1.9	list of integrals with no known antiderivative	21
1.10	List of integrals solved by CAS but has no known antiderivative	21
1.11	list of integrals solved by CAS but failed verification	21
1.12	Timing	22
1.13	Verification	22
1.14	Important notes about some of the results	23
1.15	Current tree layout of integration tests	26
1.16	Design of the test system	27

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [171]. This is test number [219].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	98.83 (169)	1.17 (2)
Rubi	97.66 (167)	2.34 (4)
Maple	71.35 (122)	28.65 (49)
Fricas	61.40 (105)	38.60 (66)
Mupad	60.82 (104)	39.18 (67)
Giac	49.12 (84)	50.88 (87)
Maxima	49.12 (84)	50.88 (87)
Reduce	45.03 (77)	54.97 (94)
Sympy	36.84 (63)	63.16 (108)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	97.661	0.000	0.000	2.339
Maple	49.123	22.222	0.000	28.655
Maxima	46.199	2.924	0.000	50.877
Giac	39.181	9.942	0.000	50.877
Mathematica	38.596	11.696	48.538	1.170
Fricas	31.579	29.825	0.000	38.596
Sympy	5.263	15.205	16.374	63.158
Mupad	0.000	60.819	0.000	39.181
Reduce	0.000	45.029	0.000	54.971

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	2	100.00	0.00	0.00
Rubi	4	100.00	0.00	0.00
Maple	49	26.53	73.47	0.00
Fricas	66	19.70	80.30	0.00
Mupad	67	0.00	100.00	0.00
Giac	87	58.62	2.30	39.08
Maxima	87	32.18	45.98	21.84
Reduce	94	100.00	0.00	0.00
Sympy	108	74.07	4.63	21.30

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.13
Maple	0.38
Reduce	0.38
Giac	0.68
Rubi	3.33
Sympy	3.54
Mathematica	4.58
Fricas	9.22
Mupad	18.67

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	335.57	1.06	302.00	1.04
Maxima	375.69	1.35	217.50	1.20
Giac	537.92	1.72	254.50	1.53
Reduce	1997.38	5.07	614.00	3.22
Sympy	3297.92	13.67	711.00	2.70
Maple	3980.34	11.03	347.00	1.22
Fricas	8020.67	26.14	465.00	1.92
Mupad	19814.60	62.97	311.00	1.38
Mathematica	31100.89	41.01	288.00	1.16

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {49, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127}

Mathematica {109, 146}

Maple {9, 50, 51, 52, 57, 58, 59, 64, 65, 66}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	29
2.2	Detailed conclusion table per each integral for all CAS systems	34
2.3	Detailed conclusion table specific for Rubi results	77

2.1 List of integrals sorted by grade for each CAS

Rubi	29
Mma	30
Maple	30
Fricas	31
Maxima	31
Giac	32
Mupad	32
Sympy	33
Reduce	33

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170 }

B grade { }

C grade { }

F normal fail { 108, 109, 146, 171 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 28, 45, 46, 47, 48, 53, 74, 75, 76, 81, 82, 88, 91, 92, 93, 94, 95, 98, 99, 100, 101, 104, 105, 106, 107, 110, 111, 112, 113, 114, 115, 120, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 141, 142, 147, 148, 149, 150, 151, 152, 155, 156, 157, 158, 161, 162, 163, 166, 167, 168, 169, 170 }

B grade { 83, 89, 90, 96, 97, 102, 103, 108, 121, 126, 127, 138, 140, 143, 153, 154, 159, 160, 165, 171 }

C grade { 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 77, 78, 79, 80, 84, 85, 86, 87, 109, 116, 117, 118, 119, 122, 123, 124, 125, 139, 144, 145, 146 }

F normal fail { 49, 164 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89 }

B grade { 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127 }

C grade { }

F normal fail { 45, 46, 47, 48, 49, 164, 165, 166, 167, 168, 169, 170, 171 }

F(-1) timedout fail { 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163 }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 34, 35, 50, 51, 52, 53, 54, 57, 58, 59, 60, 61, 64, 65, 66, 67, 70, 71, 72, 73, 74, 79, 80 }

B grade { 32, 33, 36, 37, 38, 39, 40, 41, 42, 43, 44, 55, 56, 62, 63, 68, 69, 75, 76, 77, 78, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 98, 99, 100, 104, 105, 106, 110, 111, 112, 113, 118, 119, 125, 130, 131, 149, 150 }

C grade { }

F normal fail { 45, 46, 47, 48, 49, 164, 165, 166, 167, 168, 169, 170, 171 }

F(-1) timedout fail { 94, 95, 96, 97, 101, 102, 103, 107, 108, 109, 114, 115, 116, 117, 120, 121, 122, 123, 124, 126, 127, 128, 129, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163 }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 77, 78, 79, 80, 81, 84, 85, 86, 87 }

B grade { 76, 82, 83, 88, 89 }

C grade { }

F normal fail { 45, 46, 49, 92, 93, 100, 106, 112, 113, 128, 129, 130, 131, 135, 136, 137, 141, 148, 149, 150, 151, 156, 164, 166, 167, 168, 169, 170 }

F(-1) timedout fail { 47, 48, 90, 91, 97, 98, 99, 104, 105, 110, 111, 116, 117, 118, 119, 122, 123, 124, 125, 132, 134, 138, 139, 142, 143, 144, 145, 147, 152, 153, 154, 155, 157, 158, 159, 160, 162, 163, 165, 171 }

F(-2) exception fail { 94, 95, 96, 101, 102, 103, 107, 108, 109, 114, 115, 120, 121, 126, 127, 133, 140, 146, 161 }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 51, 52, 53, 54, 55, 59, 60, 61, 62, 65, 66, 67, 68, 70, 71, 72, 73, 74, 77, 78, 79, 80, 87 }

B grade { 50, 56, 57, 58, 63, 64, 69, 75, 76, 81, 82, 83, 84, 85, 86, 88, 89 }

C grade { }

F normal fail { 45, 46, 47, 49, 116, 117, 123, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

F(-1) timedout fail { 118, 122 }

F(-2) exception fail { 48, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 119, 120, 121, 124, 125, 126, 127 }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 92, 93, 94, 95, 100, 101, 106, 110, 111, 112, 113, 114, 115, 117, 118, 119, 120, 123, 124, 125 }

C grade { }

F normal fail { }

F(-1) timedout fail { 45, 46, 47, 48, 49, 90, 91, 96, 97, 98, 99, 102, 103, 104, 105, 107, 108, 109, 116, 121, 122, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 9, 10, 11, 18, 19, 20, 24 }

B grade { 3, 4, 5, 6, 7, 8, 12, 13, 14, 15, 16, 17, 21, 22, 23, 50, 51, 52, 53, 57, 58, 59, 60, 64, 65, 66 }

C grade { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 54, 55, 61, 62, 67, 68, 70, 71, 72, 73, 74, 77, 78, 79, 80 }

F normal fail { 45, 46, 47, 48, 49, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 165, 166, 167, 168, 169, 171 }

F(-1) timedout fail { 107, 108, 109, 145, 146 }

F(-2) exception fail { 38, 39, 40, 41, 42, 43, 44, 56, 63, 69, 75, 76, 81, 82, 83, 84, 85, 86, 87, 88, 89, 164, 170 }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 26, 27, 28, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89 }

C grade { }

F normal fail { 21, 22, 23, 24, 29, 30, 31, 45, 46, 47, 48, 49, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	124	88	86	85	139	119	106	84
N.S.	1	1.00	1.43	1.01	0.99	0.98	1.60	1.37	1.22	0.97
time (sec)	N/A	0.957	0.326	0.164	0.108	0.080	0.116	0.516	0.161	5.619

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	67	69	66	66	105	82	80	63
N.S.	1	1.00	1.02	1.05	1.00	1.00	1.59	1.24	1.21	0.95
time (sec)	N/A	0.584	0.217	0.066	0.112	0.090	0.096	0.500	0.166	5.220

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	59	46	50	50	82	53	182	58
N.S.	1	1.00	1.40	1.10	1.19	1.19	1.95	1.26	4.33	1.38
time (sec)	N/A	0.564	0.041	0.198	0.110	0.082	0.313	0.566	0.175	5.290

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	39	36	43	52	59	85	55	96	69
N.S.	1	1.05	0.97	1.16	1.41	1.59	2.30	1.49	2.59	1.86
time (sec)	N/A	0.821	0.036	0.160	0.109	0.094	0.453	0.620	0.163	5.322

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	45	62	53	68	73	116	76	134	87
N.S.	1	1.05	1.44	1.23	1.58	1.70	2.70	1.77	3.12	2.02
time (sec)	N/A	0.789	0.020	0.166	0.109	0.075	0.676	0.722	0.162	5.276

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	68	107	77	86	95	143	95	187	108
N.S.	1	1.03	1.62	1.17	1.30	1.44	2.17	1.44	2.83	1.64
time (sec)	N/A	1.028	0.029	0.217	0.114	0.079	0.993	0.725	0.168	5.214

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	89	129	95	104	121	173	112	232	127
N.S.	1	1.02	1.48	1.09	1.20	1.39	1.99	1.29	2.67	1.46
time (sec)	N/A	1.219	0.037	0.267	0.112	0.080	1.954	0.747	0.162	5.238

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	110	146	108	122	138	204	129	261	145
N.S.	1	1.02	1.35	1.00	1.13	1.28	1.89	1.19	2.42	1.34
time (sec)	N/A	1.497	0.044	0.299	0.114	0.082	2.535	0.488	0.163	5.304

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	148	160	221	148	147	146	250	211	195	151
N.S.	1	1.08	1.49	1.00	0.99	0.99	1.69	1.43	1.32	1.02
time (sec)	N/A	1.545	6.175	0.103	0.108	0.083	0.146	0.657	0.156	5.503

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	112	172	120	120	119	194	161	155	121
N.S.	1	1.00	1.54	1.07	1.07	1.06	1.73	1.44	1.38	1.08
time (sec)	N/A	0.890	1.261	0.085	0.109	0.083	0.137	0.504	0.169	5.274

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	96	87	91	91	151	105	542	91
N.S.	1	1.00	1.10	1.00	1.05	1.05	1.74	1.21	6.23	1.05
time (sec)	N/A	0.860	0.322	0.154	0.112	0.079	0.446	0.642	0.160	5.226

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	72	91	80	85	92	136	90	253	90
N.S.	1	1.03	1.30	1.14	1.21	1.31	1.94	1.29	3.61	1.29
time (sec)	N/A	0.941	0.191	0.147	0.106	0.097	0.692	0.706	0.158	5.316

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	74	100	84	93	112	158	101	232	100
N.S.	1	1.03	1.39	1.17	1.29	1.56	2.19	1.40	3.22	1.39
time (sec)	N/A	0.980	0.179	0.172	0.108	0.097	1.073	0.880	0.174	5.287

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	91	123	107	120	122	206	129	269	127
N.S.	1	1.03	1.40	1.22	1.36	1.39	2.34	1.47	3.06	1.44
time (sec)	N/A	1.287	0.238	0.211	0.113	0.080	1.938	0.517	0.170	5.297

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	118	121	152	136	149	157	252	156	338	156
N.S.	1	1.03	1.29	1.15	1.26	1.33	2.14	1.32	2.86	1.32
time (sec)	N/A	1.548	0.803	0.246	0.110	0.087	2.479	0.558	0.262	5.402

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	152	180	162	175	191	304	182	396	182
N.S.	1	1.01	1.19	1.07	1.16	1.26	2.01	1.21	2.62	1.21
time (sec)	N/A	1.884	1.988	0.288	0.114	0.094	3.838	0.554	1.012	5.384

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	165	165	209	180	179	178	313	251	244	181
N.S.	1	1.00	1.27	1.09	1.08	1.08	1.90	1.52	1.48	1.10
time (sec)	N/A	1.262	1.115	0.112	0.106	0.100	0.168	0.640	0.155	5.211

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	130	139	143	142	248	182	992	142
N.S.	1	1.00	0.93	0.99	1.02	1.01	1.77	1.30	7.09	1.01
time (sec)	N/A	1.239	0.727	0.194	0.113	0.083	0.707	0.756	0.159	5.337

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	119	113	121	124	133	211	140	670	118
N.S.	1	1.02	0.97	1.03	1.06	1.14	1.80	1.20	5.73	1.01
time (sec)	N/A	1.428	0.327	0.212	0.105	0.094	0.934	0.701	0.161	5.623

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	119	121	113	118	125	145	214	134	440	114
N.S.	1	1.02	0.95	0.99	1.05	1.22	1.80	1.13	3.70	0.96
time (sec)	N/A	1.493	0.335	0.207	0.116	0.092	1.881	0.645	0.627	5.495

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	129	126	136	142	162	253	151	42	135
N.S.	1	1.02	0.99	1.07	1.12	1.28	1.99	1.19	0.33	1.06
time (sec)	N/A	1.520	0.296	0.213	0.111	0.093	2.462	0.634	200.017	5.525

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	154	161	164	172	180	181	323	187	42	169
N.S.	1	1.05	1.06	1.12	1.17	1.18	2.10	1.21	0.27	1.10
time (sec)	N/A	1.924	0.842	0.227	0.134	0.083	3.805	0.635	200.019	5.484

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	191	204	199	209	215	225	391	221	42	204
N.S.	1	1.07	1.04	1.09	1.13	1.18	2.05	1.16	0.22	1.07
time (sec)	N/A	2.369	0.505	0.285	0.112	0.083	5.174	0.689	200.055	5.465

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	233	242	237	243	250	266	462	258	42	238
N.S.	1	1.04	1.02	1.04	1.07	1.14	1.98	1.11	0.18	1.02
time (sec)	N/A	2.841	0.757	0.339	0.108	0.110	13.571	0.710	200.020	5.512

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	144	138	127	130	190	1306	146	184	144
N.S.	1	1.13	1.09	1.00	1.02	1.50	10.28	1.15	1.45	1.13
time (sec)	N/A	1.765	1.014	0.136	0.106	0.142	0.764	0.516	0.156	5.498

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	108	118	101	109	149	1020	119	128	117
N.S.	1	1.07	1.17	1.00	1.08	1.48	10.10	1.18	1.27	1.16
time (sec)	N/A	1.218	0.425	0.116	0.110	0.097	0.609	0.520	0.178	5.351

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	95	98	87	94	110	711	100	98	100
N.S.	1	1.12	1.15	1.02	1.11	1.29	8.36	1.18	1.15	1.18
time (sec)	N/A	0.633	0.128	0.088	0.110	0.097	0.523	0.421	0.182	5.686

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	67	62	88	76	541	100	128	93
N.S.	1	1.00	1.16	1.07	1.52	1.31	9.33	1.72	2.21	1.60
time (sec)	N/A	0.762	0.094	0.138	0.110	0.081	1.257	0.508	0.258	5.616

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	82	113	95	107	118	966	118	42	115
N.S.	1	1.02	1.41	1.19	1.34	1.48	12.08	1.48	0.52	1.44
time (sec)	N/A	0.988	0.235	0.191	0.114	0.112	2.325	0.613	200.030	5.971

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	111	138	122	131	177	2067	146	42	140
N.S.	1	1.08	1.34	1.18	1.27	1.72	20.07	1.42	0.41	1.36
time (sec)	N/A	1.359	0.621	0.211	0.115	0.103	4.878	0.708	200.023	6.806

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	153	163	152	158	234	2596	178	42	175
N.S.	1	1.12	1.19	1.11	1.15	1.71	18.95	1.30	0.31	1.28
time (sec)	N/A	1.993	0.934	0.272	0.112	0.120	16.184	0.794	200.021	7.268

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	208	225	193	172	220	434	4541	242	555	210
N.S.	1	1.08	0.93	0.83	1.06	2.09	21.83	1.16	2.67	1.01
time (sec)	N/A	2.209	3.901	0.153	0.116	0.189	1.110	0.551	0.167	6.199

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	171	146	155	197	311	3497	217	446	165
N.S.	1	1.09	0.93	0.99	1.25	1.98	22.27	1.38	2.84	1.05
time (sec)	N/A	1.510	1.841	0.134	0.114	0.116	0.939	0.655	0.151	5.583

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	126	140	145	185	221	2995	213	424	163
N.S.	1	1.10	1.22	1.26	1.61	1.92	26.04	1.85	3.69	1.42
time (sec)	N/A	0.914	1.464	0.114	0.117	0.088	0.812	0.570	0.153	5.450

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	122	190	141	177	222	2895	205	623	153
N.S.	1	1.10	1.71	1.27	1.59	2.00	26.08	1.85	5.61	1.38
time (sec)	N/A	1.100	1.615	0.238	0.114	0.085	1.993	0.469	0.161	5.488

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	161	159	163	208	323	4502	238	782	180
N.S.	1	1.18	1.16	1.19	1.52	2.36	32.86	1.74	5.71	1.31
time (sec)	N/A	1.579	1.627	0.274	0.115	0.120	3.560	0.735	0.163	7.329

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	192	222	193	196	262	465	8143	296	1293	230
N.S.	1	1.16	1.01	1.02	1.36	2.42	42.41	1.54	6.73	1.20
time (sec)	N/A	2.251	2.410	0.475	0.117	0.129	6.007	0.838	0.165	8.598

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	331	363	275	263	389	890	0	385	1268	335
N.S.	1	1.10	0.83	0.79	1.18	2.69	0.00	1.16	3.83	1.01
time (sec)	N/A	3.413	3.209	0.257	0.126	0.176	0.000	0.709	0.156	6.977

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	285	222	242	366	666	0	361	1097	307
N.S.	1	1.14	0.89	0.97	1.46	2.66	0.00	1.44	4.39	1.23
time (sec)	N/A	2.378	4.275	0.195	0.117	0.144	0.000	0.650	0.154	6.022

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	189	216	288	223	333	478	0	333	974	280
N.S.	1	1.14	1.52	1.18	1.76	2.53	0.00	1.76	5.15	1.48
time (sec)	N/A	1.757	3.751	0.164	0.118	0.097	0.000	0.562	0.153	5.954

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	179	201	188	213	330	488	0	323	994	282
N.S.	1	1.12	1.05	1.19	1.84	2.73	0.00	1.80	5.55	1.58
time (sec)	N/A	1.402	2.701	0.169	0.114	0.093	0.000	0.563	0.155	5.649

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	175	197	243	208	321	482	0	315	1557	279
N.S.	1	1.13	1.39	1.19	1.83	2.75	0.00	1.80	8.90	1.59
time (sec)	N/A	1.586	3.038	0.599	0.113	0.099	0.000	0.701	0.176	5.491

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	215	257	223	243	372	683	0	373	2015	315
N.S.	1	1.20	1.04	1.13	1.73	3.18	0.00	1.73	9.37	1.47
time (sec)	N/A	2.444	2.096	0.704	0.125	0.145	0.000	0.849	0.180	7.580

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	287	333	288	289	454	917	0	466	2889	380
N.S.	1	1.16	1.00	1.01	1.58	3.20	0.00	1.62	10.07	1.32
time (sec)	N/A	3.194	6.272	1.068	0.121	0.196	0.000	0.487	0.190	10.390

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	132	136	110	0	0	0	0	0	205	0
N.S.	1	1.03	0.83	0.00	0.00	0.00	0.00	0.00	1.55	0.00
time (sec)	N/A	0.881	0.370	0.000	0.000	0.000	0.000	0.000	0.165	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	154	148	115	0	0	0	0	0	159	0
N.S.	1	0.96	0.75	0.00	0.00	0.00	0.00	0.00	1.03	0.00
time (sec)	N/A	0.904	0.267	0.000	0.000	0.000	0.000	0.000	0.167	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-1)	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	170	163	133	0	0	0	0	0	160	0
N.S.	1	0.96	0.78	0.00	0.00	0.00	0.00	0.00	0.94	0.00
time (sec)	N/A	0.926	0.338	0.000	0.000	0.000	0.000	0.000	0.190	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-1)	F	F	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	170	163	133	0	0	0	0	0	163	0
N.S.	1	0.96	0.78	0.00	0.00	0.00	0.00	0.00	0.96	0.00
time (sec)	N/A	0.914	0.336	0.000	0.000	0.000	0.000	0.000	0.178	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	289	258	0	0	0	0	0	0	680	0
N.S.	1	0.89	0.00	0.00	0.00	0.00	0.00	0.00	2.35	0.00
time (sec)	N/A	1.039	0.000	0.000	0.000	0.000	0.000	0.000	0.200	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	353	369	280	347	416	415	1001	740	614	477
N.S.	1	1.05	0.79	0.98	1.18	1.18	2.84	2.10	1.74	1.35
time (sec)	N/A	2.704	5.299	0.292	0.114	0.128	0.278	0.922	0.159	5.675

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	248	261	243	246	274	273	617	450	403	300
N.S.	1	1.05	0.98	0.99	1.10	1.10	2.49	1.81	1.62	1.21
time (sec)	N/A	1.866	2.226	0.207	0.113	0.090	0.205	0.728	0.165	5.538

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	161	174	161	147	151	150	326	223	227	153
N.S.	1	1.08	1.00	0.91	0.94	0.93	2.02	1.39	1.41	0.95
time (sec)	N/A	1.251	1.060	0.141	0.106	0.083	0.210	0.567	0.159	5.551

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	76	75	74	74	131	89	100	75
N.S.	1	1.00	1.04	1.03	1.01	1.01	1.79	1.22	1.37	1.03
time (sec)	N/A	0.629	0.426	0.090	0.115	0.080	0.099	0.516	0.158	5.368

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	156	162	148	173	183	226	2387	189	194	186
N.S.	1	1.04	0.95	1.11	1.17	1.45	15.30	1.21	1.24	1.19
time (sec)	N/A	1.359	0.754	0.187	0.111	0.181	0.785	0.534	0.165	6.352

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	265	281	216	321	338	556	9721	423	943	1875
N.S.	1	1.06	0.82	1.21	1.28	2.10	36.68	1.60	3.56	7.08
time (sec)	N/A	1.786	1.680	0.247	0.323	0.350	1.264	0.708	0.161	17.783

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	320	347	331	494	574	987	0	715	2008	502
N.S.	1	1.08	1.03	1.54	1.79	3.08	0.00	2.23	6.28	1.57
time (sec)	N/A	2.325	4.568	0.286	0.125	0.114	0.000	0.640	0.167	12.207

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	661	696	573	546	691	690	1819	1413	1122	891
N.S.	1	1.05	0.87	0.83	1.05	1.04	2.75	2.14	1.70	1.35
time (sec)	N/A	5.688	6.433	0.398	0.121	0.104	0.397	0.962	0.162	5.833

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	443	470	352	392	463	462	1134	885	748	561
N.S.	1	1.06	0.79	0.88	1.05	1.04	2.56	2.00	1.69	1.27
time (sec)	N/A	3.636	5.402	0.283	0.114	0.097	0.287	0.946	0.163	5.808

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	266	276	241	246	260	259	617	450	432	300
N.S.	1	1.04	0.91	0.92	0.98	0.97	2.32	1.69	1.62	1.13
time (sec)	N/A	1.850	1.889	0.192	0.109	0.085	0.214	0.724	0.163	5.641

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	176	141	135	134	241	191	199	141
N.S.	1	1.00	1.34	1.08	1.03	1.02	1.84	1.46	1.52	1.08
time (sec)	N/A	1.007	0.717	0.111	0.111	0.085	0.134	0.602	0.161	5.527

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	254	268	190	317	290	397	4444	344	390	325
N.S.	1	1.06	0.75	1.25	1.14	1.56	17.50	1.35	1.54	1.28
time (sec)	N/A	2.466	2.004	0.214	0.113	0.335	2.083	0.713	0.167	8.012

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	415	429	277	552	496	964	16225	701	1586	3958
N.S.	1	1.03	0.67	1.33	1.20	2.32	39.10	1.69	3.82	9.54
time (sec)	N/A	3.262	4.035	0.283	0.125	0.392	2.674	0.751	0.165	29.766

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	597	631	403	865	839	1699	0	1218	3274	807
N.S.	1	1.06	0.68	1.45	1.41	2.85	0.00	2.04	5.48	1.35
time (sec)	N/A	4.055	5.268	0.480	0.132	0.499	0.000	0.877	0.169	24.361

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	603	633	419	546	680	679	1819	1413	1163	891
N.S.	1	1.05	0.69	0.91	1.13	1.13	3.02	2.34	1.93	1.48
time (sec)	N/A	4.724	6.395	0.410	0.120	0.110	0.412	1.180	0.160	5.986

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	389	403	297	347	387	386	1001	740	684	478
N.S.	1	1.04	0.76	0.89	0.99	0.99	2.57	1.90	1.76	1.23
time (sec)	N/A	2.684	6.249	0.261	0.113	0.092	0.286	1.061	0.160	5.780

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	191	191	212	210	202	201	410	308	325	221
N.S.	1	1.00	1.11	1.10	1.06	1.05	2.15	1.61	1.70	1.16
time (sec)	N/A	1.363	1.638	0.149	0.105	0.083	0.230	0.629	0.161	5.540

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	363	387	255	501	436	623	7096	600	654	508
N.S.	1	1.07	0.70	1.38	1.20	1.72	19.55	1.65	1.80	1.40
time (sec)	N/A	4.019	2.995	0.268	0.115	0.642	22.194	0.713	0.168	9.355

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	574	589	1024	829	685	1512	24300	1042	2418	701
N.S.	1	1.03	1.78	1.44	1.19	2.63	42.33	1.82	4.21	1.22
time (sec)	N/A	5.517	6.967	0.421	0.122	0.953	28.584	0.769	0.163	13.002

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	798	830	457	1271	1119	2549	0	1758	4720	1172
N.S.	1	1.04	0.57	1.59	1.40	3.19	0.00	2.20	5.91	1.47
time (sec)	N/A	7.058	4.432	0.724	0.263	1.371	0.000	0.768	0.165	14.866

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	359	258	500	445	627	7096	601	819	508
N.S.	1	1.07	0.77	1.48	1.32	1.86	21.06	1.78	2.43	1.51
time (sec)	N/A	3.983	2.919	0.280	0.116	0.592	22.678	0.627	0.169	9.763

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	236	253	190	317	294	390	4444	344	508	325
N.S.	1	1.07	0.81	1.34	1.25	1.65	18.83	1.46	2.15	1.38
time (sec)	N/A	2.382	1.995	0.224	0.109	0.275	2.092	0.658	0.155	7.934

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	156	163	148	173	178	212	2387	190	281	186
N.S.	1	1.04	0.95	1.11	1.14	1.36	15.30	1.22	1.80	1.19
time (sec)	N/A	1.300	0.723	0.194	0.115	0.151	0.795	0.562	0.156	6.672

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	117	100	106	118	966	111	138	109
N.S.	1	1.00	1.18	1.01	1.07	1.19	9.76	1.12	1.39	1.10
time (sec)	N/A	0.759	0.144	0.129	0.105	0.100	0.569	0.533	0.157	6.310

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	165	164	313	197	243	301	24052	278	423	196
N.S.	1	0.99	1.90	1.19	1.47	1.82	145.77	1.68	2.56	1.19
time (sec)	N/A	1.023	0.982	0.252	0.118	0.256	15.741	0.572	0.161	17.560

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	281	317	543	364	520	1345	0	674	2915	393
N.S.	1	1.13	1.93	1.30	1.85	4.79	0.00	2.40	10.37	1.40
time (sec)	N/A	2.182	4.476	0.500	0.132	0.886	0.000	0.632	0.163	55.908

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	477	538	898	647	1096	3643	0	1447	8368	65819
N.S.	1	1.13	1.88	1.36	2.30	7.64	0.00	3.03	17.54	137.99
time (sec)	N/A	4.458	7.768	1.429	0.166	3.064	0.000	0.811	0.186	19.643

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	579	594	1022	829	684	1477	24300	1039	2547	701
N.S.	1	1.03	1.77	1.43	1.18	2.55	41.97	1.79	4.40	1.21
time (sec)	N/A	5.359	6.961	0.462	0.129	0.901	28.692	0.718	0.166	12.475

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	417	433	277	552	493	939	16225	701	1762	3958
N.S.	1	1.04	0.66	1.32	1.18	2.25	38.91	1.68	4.23	9.49
time (sec)	N/A	3.287	3.405	0.288	0.125	0.374	2.707	0.618	0.160	29.981

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	292	304	221	321	319	505	9721	420	1114	1875
N.S.	1	1.04	0.76	1.10	1.09	1.73	33.29	1.44	3.82	6.42
time (sec)	N/A	1.986	1.536	0.235	0.121	0.163	1.316	0.566	0.158	18.125

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	151	207	173	205	256	4396	248	603	184
N.S.	1	1.08	1.48	1.24	1.46	1.83	31.40	1.77	4.31	1.31
time (sec)	N/A	1.042	1.777	0.111	0.109	0.088	0.847	0.484	0.162	8.143

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	293	329	561	365	513	1275	0	673	2663	430
N.S.	1	1.12	1.91	1.25	1.75	4.35	0.00	2.30	9.09	1.47
time (sec)	N/A	2.244	4.955	0.491	0.129	0.911	0.000	0.627	0.159	61.570

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	509	566	984	577	1185	4174	0	1674	11926	73684
N.S.	1	1.11	1.93	1.13	2.33	8.20	0.00	3.29	23.43	144.76
time (sec)	N/A	4.768	7.927	1.500	0.172	3.335	0.000	0.856	0.224	22.174

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	841	919	1758	951	2519	9594	0	4549	27735	128667
N.S.	1	1.09	2.09	1.13	3.00	11.41	0.00	5.41	32.98	152.99
time (sec)	N/A	8.809	7.825	4.976	0.243	9.697	0.000	1.299	14.420	40.837

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	804	837	454	1271	1110	2490	0	1757	5125	1172
N.S.	1	1.04	0.56	1.58	1.38	3.10	0.00	2.19	6.37	1.46
time (sec)	N/A	6.737	4.369	0.762	0.161	1.124	0.000	0.924	0.169	15.049

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	597	631	404	865	827	1618	0	1219	3698	807
N.S.	1	1.06	0.68	1.45	1.39	2.71	0.00	2.04	6.19	1.35
time (sec)	N/A	4.097	4.754	0.486	0.141	0.455	0.000	0.691	0.168	23.270

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	352	379	331	493	543	897	0	714	2450	502
N.S.	1	1.08	0.94	1.40	1.54	2.55	0.00	2.03	6.96	1.43
time (sec)	N/A	2.507	4.502	0.296	0.128	0.164	0.000	0.619	0.167	12.168

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	209	234	261	262	367	566	0	401	1433	327
N.S.	1	1.12	1.25	1.25	1.76	2.71	0.00	1.92	6.86	1.56
time (sec)	N/A	1.630	3.532	0.197	0.124	0.101	0.000	0.569	0.165	8.121

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	487	549	912	649	1078	3496	0	1447	7603	65817
N.S.	1	1.13	1.87	1.33	2.21	7.18	0.00	2.97	15.61	135.15
time (sec)	N/A	4.198	7.841	1.367	0.166	3.655	0.000	0.791	0.178	19.798

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F(-2)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	861	932	1732	949	2537	9567	0	4552	28024	128666
N.S.	1	1.08	2.01	1.10	2.95	11.11	0.00	5.29	32.55	149.44
time (sec)	N/A	8.774	7.598	4.847	0.243	11.061	0.000	1.238	0.714	39.875

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-1)	B	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	464	476	1232	4430	0	35153	0	0	240	0
N.S.	1	1.03	2.66	9.55	0.00	75.76	0.00	0.00	0.52	0.00
time (sec)	N/A	5.823	6.324	0.727	0.000	10.117	0.000	0.000	0.211	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	325	332	314	3319	0	23984	0	0	179	0
N.S.	1	1.02	0.97	10.21	0.00	73.80	0.00	0.00	0.55	0.00
time (sec)	N/A	3.747	3.304	0.299	0.000	4.325	0.000	0.000	0.204	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	224	218	220	2197	0	12410	0	0	118	22955
N.S.	1	0.97	0.98	9.81	0.00	55.40	0.00	0.00	0.53	102.48
time (sec)	N/A	2.053	1.365	0.266	0.000	1.466	0.000	0.000	0.187	53.782

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	134	150	1298	0	2588	0	0	62	1199
N.S.	1	0.86	0.97	8.37	0.00	16.70	0.00	0.00	0.40	7.74
time (sec)	N/A	1.281	0.380	0.178	0.000	0.193	0.000	0.000	0.172	12.933

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	234	225	233	3576	0	0	0	0	99	62245
N.S.	1	0.96	1.00	15.28	0.00	0.00	0.00	0.00	0.42	266.00
time (sec)	N/A	2.996	0.485	0.210	0.000	0.000	0.000	0.000	0.199	30.395

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	317	327	362	5778	0	0	0	0	0	138318
N.S.	1	1.03	1.14	18.23	0.00	0.00	0.00	0.00	0.00	436.33
time (sec)	N/A	3.872	4.312	0.213	0.000	0.000	0.000	0.000	0.291	39.989

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-2)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	543	588	2819	9797	0	0	0	0	0	0
N.S.	1	1.08	5.19	18.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.150	6.438	0.243	0.000	0.000	0.000	0.000	0.364	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-1)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	550	566	1290	10952	0	0	0	0	510	0
N.S.	1	1.03	2.35	19.91	0.00	0.00	0.00	0.00	0.93	0.00
time (sec)	N/A	7.155	6.363	0.457	0.000	0.000	0.000	0.000	0.275	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	396	402	350	7939	0	58971	0	0	383	0
N.S.	1	1.02	0.88	20.05	0.00	148.92	0.00	0.00	0.97	0.00
time (sec)	N/A	4.690	4.218	0.293	0.000	72.225	0.000	0.000	0.255	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	273	268	260	5107	0	31081	0	0	256	0
N.S.	1	0.98	0.95	18.71	0.00	113.85	0.00	0.00	0.94	0.00
time (sec)	N/A	2.724	3.068	0.267	0.000	16.915	0.000	0.000	0.221	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	187	166	202	2500	0	6846	0	0	139	4260
N.S.	1	0.89	1.08	13.37	0.00	36.61	0.00	0.00	0.74	22.78
time (sec)	N/A	1.678	0.841	0.182	0.000	0.857	0.000	0.000	0.199	39.810

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	271	275	266	6055	0	0	0	0	212	106783
N.S.	1	1.01	0.98	22.34	0.00	0.00	0.00	0.00	0.78	394.03
time (sec)	N/A	4.418	1.692	0.208	0.000	0.000	0.000	0.000	0.225	52.297

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-2)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	372	387	1732	9865	0	0	0	0	0	0
N.S.	1	1.04	4.66	26.52	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	5.819	4.318	0.220	0.000	0.000	0.000	0.000	0.388	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-2)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	532	563	7678	14441	0	0	0	0	0	0
N.S.	1	1.06	14.43	27.14	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.250	6.728	0.240	0.000	0.000	0.000	0.000	0.702	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	503	511	564	11346	0	91140	0	0	615	0
N.S.	1	1.02	1.12	22.56	0.00	181.19	0.00	0.00	1.22	0.00
time (sec)	N/A	6.176	6.361	0.540	0.000	160.303	0.000	0.000	0.276	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	353	345	324	7338	0	48734	0	0	416	0
N.S.	1	0.98	0.92	20.79	0.00	138.06	0.00	0.00	1.18	0.00
time (sec)	N/A	3.667	3.516	0.319	0.000	36.787	0.000	0.000	0.236	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	229	208	262	3587	0	10840	0	0	231	5863
N.S.	1	0.91	1.14	15.66	0.00	47.34	0.00	0.00	1.01	25.60
time (sec)	N/A	2.152	1.405	0.187	0.000	1.799	0.000	0.000	0.200	109.008

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F(-1)	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	336	341	322	8698	0	0	0	0	340	0
N.S.	1	1.01	0.96	25.89	0.00	0.00	0.00	0.00	1.01	0.00
time (sec)	N/A	6.430	3.635	0.232	0.000	0.000	0.000	0.000	0.241	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	B	F(-2)	F(-1)	F(-1)	F(-2)	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	473	0	6112	14119	0	0	0	0	0	0
N.S.	1	0.00	12.92	29.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	6.635	0.271	0.000	0.000	0.000	0.000	0.407	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	B	F(-2)	F(-1)	F(-1)	F(-2)	F	F(-1)
verified	N/A	N/A	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	643	0	2429150	20663	0	0	0	0	0	0
N.S.	1	0.00	3777.84	32.14	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	123.155	0.374	0.000	0.000	0.000	0.000	0.557	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	407	418	392	5978	0	37247	0	0	399	28858
N.S.	1	1.03	0.96	14.69	0.00	91.52	0.00	0.00	0.98	70.90
time (sec)	N/A	4.741	6.034	0.425	0.000	11.450	0.000	0.000	0.210	104.878

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	287	287	275	5513	0	25627	0	0	310	21254
N.S.	1	1.00	0.96	19.21	0.00	89.29	0.00	0.00	1.08	74.06
time (sec)	N/A	2.929	3.963	0.193	0.000	4.714	0.000	0.000	0.194	38.709

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	194	185	192	3853	0	13473	0	0	221	16400
N.S.	1	0.95	0.99	19.86	0.00	69.45	0.00	0.00	1.14	84.54
time (sec)	N/A	1.643	0.998	0.202	0.000	1.591	0.000	0.000	0.182	18.061

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	112	129	3463	0	3194	0	0	135	4326
N.S.	1	0.84	0.97	26.04	0.00	24.02	0.00	0.00	1.02	32.53
time (sec)	N/A	0.973	0.151	0.148	0.000	0.237	0.000	0.000	0.174	9.799

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	210	195	194	13474	0	0	0	0	206	25341
N.S.	1	0.93	0.92	64.16	0.00	0.00	0.00	0.00	0.98	120.67
time (sec)	N/A	2.161	0.285	0.199	0.000	0.000	0.000	0.000	0.207	61.106

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	327	358	338	20870	0	0	0	0	0	225004
N.S.	1	1.09	1.03	63.82	0.00	0.00	0.00	0.00	0.00	688.09
time (sec)	N/A	3.788	4.914	0.227	0.000	0.000	0.000	0.000	0.722	46.475

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	511	512	920	11255	0	0	0	0	1159	0
N.S.	1	1.00	1.80	22.03	0.00	0.00	0.00	0.00	2.27	0.00
time (sec)	N/A	6.251	6.540	0.472	0.000	0.000	0.000	0.000	0.222	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	343	346	476	9399	0	0	0	0	898	54886
N.S.	1	1.01	1.39	27.40	0.00	0.00	0.00	0.00	2.62	160.02
time (sec)	N/A	3.678	6.357	0.224	0.000	0.000	0.000	0.000	0.206	62.014

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-1)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	219	290	7396	0	31879	0	0	637	40542
N.S.	1	1.09	1.44	36.80	0.00	158.60	0.00	0.00	3.17	201.70
time (sec)	N/A	1.894	1.756	0.226	0.000	62.605	0.000	0.000	0.196	36.042

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	161	218	5613	0	7982	0	0	384	8588
N.S.	1	1.03	1.39	35.75	0.00	50.84	0.00	0.00	2.45	54.70
time (sec)	N/A	1.193	0.700	0.145	0.000	1.960	0.000	0.000	0.177	15.950

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F(-2)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	262	309	296	26343	0	0	0	0	0	630337
N.S.	1	1.18	1.13	100.55	0.00	0.00	0.00	0.00	0.00	2405.87
time (sec)	N/A	3.585	3.362	0.223	0.000	0.000	0.000	0.000	0.397	92.286

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-2)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	447	515	2078	40619	0	0	0	0	0	0
N.S.	1	1.15	4.65	90.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	6.735	6.269	0.321	0.000	0.000	0.000	0.000	1.095	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F(-1)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	585	605	670	13619	0	0	0	0	0	0
N.S.	1	1.03	1.15	23.28	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.532	6.589	0.531	0.000	0.000	0.000	0.000	0.246	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	358	388	414	11386	0	0	0	0	0	88684
N.S.	1	1.08	1.16	31.80	0.00	0.00	0.00	0.00	0.00	247.72
time (sec)	N/A	4.095	4.834	0.262	0.000	0.000	0.000	0.000	0.230	110.557

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	F(-1)	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	273	300	300	8983	0	0	0	0	1243	64641
N.S.	1	1.10	1.10	32.90	0.00	0.00	0.00	0.00	4.55	236.78
time (sec)	N/A	2.507	1.998	0.251	0.000	0.000	0.000	0.000	0.209	83.068

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-2)	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	209	231	223	6803	0	13143	0	0	750	14163
N.S.	1	1.11	1.07	32.55	0.00	62.89	0.00	0.00	3.59	67.77
time (sec)	N/A	1.762	0.636	0.168	0.000	7.804	0.000	0.000	0.186	33.504

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-2)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	365	444	1948	45119	0	0	0	0	0	0
N.S.	1	1.22	5.34	123.61	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	5.917	6.250	0.280	0.000	0.000	0.000	0.000	0.555	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	F(-2)	F(-1)	F	F(-2)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	679	767	6052	67570	0	0	0	0	0	0
N.S.	1	1.13	8.91	99.51	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	11.405	6.518	0.419	0.000	0.000	0.000	0.000	1.649	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	679	707	1202	0	0	0	0	0	257	0
N.S.	1	1.04	1.77	0.00	0.00	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	9.263	8.531	0.000	0.000	0.000	0.000	0.000	0.662	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	505	523	835	0	0	0	0	0	174	0
N.S.	1	1.04	1.65	0.00	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	5.926	7.892	0.000	0.000	0.000	0.000	0.000	0.267	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	381	389	619	0	0	68078	0	0	96	0
N.S.	1	1.02	1.62	0.00	0.00	178.68	0.00	0.00	0.25	0.00
time (sec)	N/A	3.563	7.139	0.000	0.000	165.288	0.000	0.000	0.215	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	289	292	441	0	0	78051	0	0	132	0
N.S.	1	1.01	1.53	0.00	0.00	270.07	0.00	0.00	0.46	0.00
time (sec)	N/A	2.044	2.894	0.000	0.000	103.585	0.000	0.000	0.192	0.000

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	300	329	412	0	0	0	0	0	1475	0
N.S.	1	1.10	1.37	0.00	0.00	0.00	0.00	0.00	4.92	0.00
time (sec)	N/A	2.448	3.644	0.000	0.000	0.000	0.000	0.000	0.239	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-2)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	370	441	424	0	0	0	0	0	0	0
N.S.	1	1.19	1.15	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.207	6.276	0.000	0.000	0.000	0.000	0.000	0.518	0.000

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	597	693	1109	0	0	0	0	0	0	0
N.S.	1	1.16	1.86	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.456	6.930	0.000	0.000	0.000	0.000	0.000	2.132	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	683	711	1304	0	0	0	0	0	367	0
N.S.	1	1.04	1.91	0.00	0.00	0.00	0.00	0.00	0.54	0.00
time (sec)	N/A	9.585	8.155	0.000	0.000	0.000	0.000	0.000	0.811	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	510	524	867	0	0	0	0	0	206	0
N.S.	1	1.03	1.70	0.00	0.00	0.00	0.00	0.00	0.40	0.00
time (sec)	N/A	5.985	7.920	0.000	0.000	0.000	0.000	0.000	0.290	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	382	393	580	0	0	0	0	0	278	0
N.S.	1	1.03	1.52	0.00	0.00	0.00	0.00	0.00	0.73	0.00
time (sec)	N/A	3.784	6.553	0.000	0.000	0.000	0.000	0.000	0.225	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	382	420	1664	0	0	0	0	0	0	0
N.S.	1	1.10	4.36	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.249	6.802	0.000	0.000	0.000	0.000	0.000	27.646	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	402	457	519	0	0	0	0	0	0	0
N.S.	1	1.14	1.29	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.669	6.471	0.000	0.000	0.000	0.000	0.000	0.464	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-2)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	586	678	3134	0	0	0	0	0	0	0
N.S.	1	1.16	5.35	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.704	8.262	0.000	0.000	0.000	0.000	0.000	3.736	0.000

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	695	717	1261	0	0	0	0	0	331	0
N.S.	1	1.03	1.81	0.00	0.00	0.00	0.00	0.00	0.48	0.00
time (sec)	N/A	9.394	8.349	180.000	0.000	0.000	0.000	0.000	0.410	0.000

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	507	524	780	0	0	0	0	0	439	0
N.S.	1	1.03	1.54	0.00	0.00	0.00	0.00	0.00	0.87	0.00
time (sec)	N/A	5.867	7.811	0.000	0.000	0.000	0.000	0.000	0.285	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	535	562	1774	0	0	0	0	0	0	0
N.S.	1	1.05	3.32	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.247	7.609	0.000	0.000	0.000	0.000	0.000	0.396	0.000

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	545	603	802	0	0	0	0	0	47	0
N.S.	1	1.11	1.47	0.00	0.00	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	8.329	6.988	0.000	0.000	0.000	0.000	0.000	200.023	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F(-1)	F(-1)	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	590	658	641	0	0	0	0	0	0	0
N.S.	1	1.12	1.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	8.507	6.708	0.000	0.000	0.000	0.000	0.000	17.794	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	F(-1)	F(-2)	F(-1)	F(-1)	F	F	F(-1)
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	946	0	2723641	0	0	0	0	0	47	0
N.S.	1	0.00	2879.11	0.00	0.00	0.00	0.00	0.00	0.05	0.00
time (sec)	N/A	0.000	53.154	0.000	0.000	0.000	0.000	0.000	200.020	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	507	524	785	0	0	0	0	0	341	0
N.S.	1	1.03	1.55	0.00	0.00	0.00	0.00	0.00	0.67	0.00
time (sec)	N/A	5.934	7.656	0.000	0.000	0.000	0.000	0.000	0.580	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	381	392	582	0	0	0	0	0	234	0
N.S.	1	1.03	1.53	0.00	0.00	0.00	0.00	0.00	0.61	0.00
time (sec)	N/A	3.560	4.642	0.000	0.000	0.000	0.000	0.000	0.230	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	290	294	450	0	0	77916	0	0	132	0
N.S.	1	1.01	1.55	0.00	0.00	268.68	0.00	0.00	0.46	0.00
time (sec)	N/A	2.045	4.409	0.000	0.000	127.238	0.000	0.000	0.209	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	237	232	362	0	0	96324	0	0	1189	0
N.S.	1	0.98	1.53	0.00	0.00	406.43	0.00	0.00	5.02	0.00
time (sec)	N/A	1.164	1.450	0.000	0.000	128.693	0.000	0.000	0.242	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	251	303	264	0	0	0	0	0	1341	0
N.S.	1	1.21	1.05	0.00	0.00	0.00	0.00	0.00	5.34	0.00
time (sec)	N/A	2.318	1.776	0.000	0.000	0.000	0.000	0.000	0.272	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	375	458	388	0	0	0	0	0	0	0
N.S.	1	1.22	1.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.909	4.094	0.000	0.000	0.000	0.000	0.000	0.413	0.000

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	528	555	2245	0	0	0	0	0	0	0
N.S.	1	1.05	4.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.158	8.190	0.000	0.000	0.000	0.000	0.000	0.574	0.000

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	380	417	2141	0	0	0	0	0	47	0
N.S.	1	1.10	5.63	0.00	0.00	0.00	0.00	0.00	0.12	0.00
time (sec)	N/A	4.245	7.018	0.000	0.000	0.000	0.000	0.000	200.025	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	299	328	403	0	0	0	0	0	1476	0
N.S.	1	1.10	1.35	0.00	0.00	0.00	0.00	0.00	4.94	0.00
time (sec)	N/A	2.458	3.499	0.000	0.000	0.000	0.000	0.000	0.238	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	251	302	275	0	0	0	0	0	1342	0
N.S.	1	1.20	1.10	0.00	0.00	0.00	0.00	0.00	5.35	0.00
time (sec)	N/A	2.362	2.172	0.000	0.000	0.000	0.000	0.000	0.259	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	383	466	484	0	0	0	0	0	0	0
N.S.	1	1.22	1.26	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.133	6.517	0.000	0.000	0.000	0.000	0.000	0.401	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	598	704	902	0	0	0	0	0	0	0
N.S.	1	1.18	1.51	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.357	6.701	0.000	0.000	0.000	0.000	0.000	0.677	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	549	605	2650	0	0	0	0	0	47	0
N.S.	1	1.10	4.83	0.00	0.00	0.00	0.00	0.00	0.09	0.00
time (sec)	N/A	8.150	8.364	0.000	0.000	0.000	0.000	0.000	200.023	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	407	462	1135	0	0	0	0	0	0	0
N.S.	1	1.14	2.79	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.692	6.713	0.000	0.000	0.000	0.000	0.000	0.389	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-2)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	373	444	434	0	0	0	0	0	0	0
N.S.	1	1.19	1.16	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.146	6.071	0.000	0.000	0.000	0.000	0.000	0.468	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	379	461	403	0	0	0	0	0	0	0
N.S.	1	1.22	1.06	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.041	3.779	0.000	0.000	0.000	0.000	0.000	0.389	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F(-1)	F(-1)	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	651	751	903	0	0	0	0	0	0	0
N.S.	1	1.15	1.39	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	7.492	6.716	0.000	0.000	0.000	0.000	0.000	0.654	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	376	371	0	0	0	0	0	0	102	0
N.S.	1	0.99	0.00	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	1.219	0.000	0.000	0.000	0.000	0.000	0.000	45.277	0.000

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F(-1)	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	560	580	1390	0	0	0	0	0	335	0
N.S.	1	1.04	2.48	0.00	0.00	0.00	0.00	0.00	0.60	0.00
time (sec)	N/A	5.837	6.344	0.000	0.000	0.000	0.000	0.000	0.186	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	363	385	308	0	0	0	0	0	240	0
N.S.	1	1.06	0.85	0.00	0.00	0.00	0.00	0.00	0.66	0.00
time (sec)	N/A	3.196	5.035	0.000	0.000	0.000	0.000	0.000	0.170	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	247	262	202	0	0	0	0	0	145	0
N.S.	1	1.06	0.82	0.00	0.00	0.00	0.00	0.00	0.59	0.00
time (sec)	N/A	1.722	2.101	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	178	178	135	0	0	0	0	0	65	0
N.S.	1	1.00	0.76	0.00	0.00	0.00	0.00	0.00	0.37	0.00
time (sec)	N/A	0.886	0.193	0.000	0.000	0.000	0.000	0.000	0.159	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	258	265	204	0	0	0	0	0	102	0
N.S.	1	1.03	0.79	0.00	0.00	0.00	0.00	0.00	0.40	0.00
time (sec)	N/A	1.796	0.824	0.000	0.000	0.000	0.000	0.000	0.155	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	403	451	360	0	0	0	0	0	0	0
N.S.	1	1.12	0.89	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.577	6.018	0.000	0.000	0.000	0.000	0.000	0.227	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	B	F	F(-1)	F	F	F	F	F(-1)
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	702	0	2238	0	0	0	0	0	0	0
N.S.	1	0.00	3.19	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	6.202	0.000	0.000	0.000	0.000	0.000	0.277	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [24] had the largest ratio of [.50000000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	10	10	1.00	36	0.278
2	A	6	6	1.00	30	0.200
3	A	6	6	1.00	36	0.167
4	A	10	10	1.05	38	0.263
5	A	9	9	1.05	38	0.237
6	A	12	12	1.03	38	0.316
7	A	14	14	1.02	38	0.368
8	A	17	17	1.02	38	0.447
9	A	13	13	1.08	38	0.342
10	A	8	8	1.00	32	0.250
11	A	8	8	1.00	38	0.211
12	A	9	9	1.03	40	0.225
13	A	9	9	1.03	40	0.225
14	A	12	12	1.03	40	0.300
15	A	14	14	1.03	40	0.350
16	A	17	17	1.01	40	0.425
17	A	10	10	1.00	32	0.312
18	A	10	10	1.00	38	0.263
19	A	13	13	1.02	40	0.325
20	A	12	12	1.02	40	0.300
21	A	13	13	1.02	40	0.325

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	15	15	1.05	40	0.375
23	A	19	19	1.07	40	0.475
24	A	20	20	1.04	40	0.500
25	A	14	13	1.13	40	0.325
26	A	12	11	1.07	38	0.289
27	A	5	4	1.12	32	0.125
28	A	6	6	1.00	38	0.158
29	A	8	8	1.02	40	0.200
30	A	10	10	1.08	40	0.250
31	A	14	14	1.12	40	0.350
32	A	15	14	1.08	40	0.350
33	A	12	11	1.09	38	0.289
34	A	6	6	1.10	32	0.188
35	A	8	8	1.10	38	0.211
36	A	10	10	1.18	40	0.250
37	A	12	12	1.16	40	0.300
38	A	18	17	1.10	40	0.425
39	A	14	13	1.14	40	0.325
40	A	11	11	1.14	38	0.289
41	A	8	8	1.12	32	0.250
42	A	10	10	1.13	38	0.263
43	A	13	13	1.20	40	0.325
44	A	15	15	1.16	40	0.375
45	A	9	8	1.03	39	0.205
46	A	9	8	0.96	39	0.205
47	A	9	8	0.96	41	0.195
48	A	9	8	0.96	41	0.195
49	A	5	4	0.89	43	0.093
50	A	13	13	1.05	43	0.302
51	A	11	11	1.05	43	0.256
52	A	8	8	1.08	41	0.195
53	A	6	6	1.00	31	0.194

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	10	9	1.04	43	0.209
55	A	9	8	1.06	43	0.186
56	A	8	8	1.08	43	0.186
57	A	17	17	1.05	45	0.378
58	A	14	14	1.06	45	0.311
59	A	10	10	1.04	43	0.233
60	A	8	8	1.00	33	0.242
61	A	13	12	1.06	45	0.267
62	A	11	10	1.03	45	0.222
63	A	12	11	1.06	45	0.244
64	A	16	16	1.05	45	0.356
65	A	12	12	1.04	43	0.279
66	A	10	10	1.00	33	0.303
67	A	16	15	1.07	45	0.333
68	A	14	13	1.03	45	0.289
69	A	14	13	1.04	45	0.289
70	A	16	15	1.07	45	0.333
71	A	13	12	1.07	45	0.267
72	A	9	8	1.04	43	0.186
73	A	7	6	1.00	33	0.182
74	A	4	4	0.99	45	0.089
75	A	7	7	1.13	45	0.156
76	A	10	10	1.13	45	0.222
77	A	14	13	1.03	45	0.289
78	A	11	10	1.04	45	0.222
79	A	9	8	1.04	43	0.186
80	A	6	6	1.08	33	0.182
81	A	7	7	1.12	45	0.156
82	A	9	9	1.11	45	0.200
83	A	11	11	1.09	45	0.244
84	A	14	13	1.04	45	0.289
85	A	12	11	1.06	45	0.244

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	9	9	1.08	43	0.209
87	A	9	9	1.12	33	0.273
88	A	9	9	1.13	45	0.200
89	A	11	11	1.08	45	0.244
90	A	21	20	1.03	47	0.426
91	A	18	17	1.02	47	0.362
92	A	15	14	0.97	45	0.311
93	A	12	11	0.86	35	0.314
94	A	16	15	0.96	47	0.319
95	A	17	16	1.03	47	0.340
96	A	20	19	1.08	47	0.404
97	A	23	22	1.03	47	0.468
98	A	20	19	1.02	47	0.404
99	A	17	16	0.98	45	0.356
100	A	14	13	0.89	35	0.371
101	A	20	19	1.01	47	0.404
102	A	20	19	1.04	47	0.404
103	A	20	19	1.06	47	0.404
104	A	22	21	1.02	47	0.447
105	A	19	18	0.98	45	0.400
106	A	16	15	0.91	35	0.429
107	A	24	23	1.01	47	0.489
108	F	0	0	N/A	0.000	N/A
109	F	0	0	N/A	0.000	N/A
110	A	19	18	1.03	47	0.383
111	A	16	15	1.00	47	0.319
112	A	13	12	0.95	45	0.267
113	A	10	9	0.84	35	0.257
114	A	13	12	0.93	47	0.255
115	A	17	16	1.09	47	0.340
116	A	19	18	1.00	47	0.383
117	A	16	15	1.01	47	0.319

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	A	12	11	1.09	45	0.244
119	A	10	9	1.03	35	0.257
120	A	17	16	1.18	47	0.340
121	A	20	19	1.15	47	0.404
122	A	19	18	1.03	47	0.383
123	A	15	14	1.08	47	0.298
124	A	13	12	1.10	45	0.267
125	A	13	12	1.11	35	0.343
126	A	19	18	1.22	47	0.383
127	A	23	22	1.13	47	0.468
128	A	17	16	1.04	49	0.327
129	A	14	13	1.04	49	0.265
130	A	11	10	1.02	49	0.204
131	A	8	7	1.01	49	0.143
132	A	8	7	1.10	49	0.143
133	A	13	12	1.19	49	0.245
134	A	16	15	1.16	49	0.306
135	A	17	16	1.04	49	0.327
136	A	14	13	1.03	49	0.265
137	A	12	11	1.03	49	0.224
138	A	11	10	1.10	49	0.204
139	A	11	10	1.14	49	0.204
140	A	16	15	1.16	49	0.306
141	A	17	16	1.03	49	0.327
142	A	14	13	1.03	49	0.265
143	A	14	13	1.05	49	0.265
144	A	14	13	1.11	49	0.265
145	A	14	13	1.12	49	0.265
146	F	0	0	N/A	0.000	N/A
147	A	14	13	1.03	49	0.265
148	A	11	10	1.03	49	0.204
149	A	8	7	1.01	49	0.143

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	5	4	0.98	49	0.082
151	A	10	9	1.21	49	0.184
152	A	13	12	1.22	49	0.245
153	A	14	13	1.05	49	0.265
154	A	11	10	1.10	49	0.204
155	A	8	7	1.10	49	0.143
156	A	10	9	1.20	49	0.184
157	A	13	12	1.22	49	0.245
158	A	16	15	1.18	49	0.306
159	A	14	13	1.10	49	0.265
160	A	11	10	1.14	49	0.204
161	A	13	12	1.19	49	0.245
162	A	13	12	1.22	49	0.245
163	A	16	15	1.15	49	0.306
164	A	5	4	0.99	45	0.089
165	A	17	16	1.04	45	0.356
166	A	13	12	1.06	45	0.267
167	A	11	10	1.06	43	0.233
168	A	9	8	1.00	33	0.242
169	A	11	10	1.03	45	0.222
170	A	14	13	1.12	45	0.289
171	F	0	0	N/A	0.000	N/A

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	89
3.2	$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	97
3.3	$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	104
3.4	$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	111
3.5	$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	118
3.6	$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	125
3.7	$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	133
3.8	$\int \cot^6(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	143
3.9	$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	153
3.10	$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	163
3.11	$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	171
3.12	$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	179
3.13	$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	187
3.14	$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	195
3.15	$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	204
3.16	$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	214
3.17	$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	225
3.18	$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	234
3.19	$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	243
3.20	$\int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	252
3.21	$\int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	262
3.22	$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	272
3.23	$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	283
3.24	$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \dots$	295
3.25	$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx \dots$	307
3.26	$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx \dots$	317

3.27	$\int \frac{B \tan(c+dx)+C \tan^2(c+dx)}{a+b \tan(c+dx)} dx$	326
3.28	$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	333
3.29	$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	341
3.30	$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	349
3.31	$\int \frac{\cot^4(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$	358
3.32	$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	369
3.33	$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	381
3.34	$\int \frac{B \tan(c+dx)+C \tan^2(c+dx)}{(a+b \tan(c+dx))^2} dx$	392
3.35	$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	400
3.36	$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	408
3.37	$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$	419
3.38	$\int \frac{\tan^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	430
3.39	$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	443
3.40	$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	454
3.41	$\int \frac{B \tan(c+dx)+C \tan^2(c+dx)}{(a+b \tan(c+dx))^3} dx$	464
3.42	$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	473
3.43	$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	483
3.44	$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$	495
3.45	$\int \tan^2(c+dx)(b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) dx$	508
3.46	$\int \tan^m(c+dx)(b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) dx$	515
3.47	$\int \tan^m(c+dx) \sqrt{b \tan(c+dx)} (A+B \tan(c+dx)+C \tan^2(c+dx)) dx$	522
3.48	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{b \tan(c+dx)}} dx$	529
3.49	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	537
3.50	$\int (a+b \tan(e+fx))^3 (c+d \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	544
3.51	$\int (a+b \tan(e+fx))^2 (c+d \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	556
3.52	$\int (a+b \tan(e+fx)) (c+d \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	567
3.53	$\int (c+d \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	576
3.54	$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	583
3.55	$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	592
3.56	$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	603
3.57	$\int (a+b \tan(e+fx))^3 (c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	614
3.58	$\int (a+b \tan(e+fx))^2 (c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	629
3.59	$\int (a+b \tan(e+fx)) (c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	642

3.60	$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$	652
3.61	$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	661
3.62	$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	673
3.63	$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	686
3.64	$\int (a+b \tan(e+fx))^2 (c+d \tan(e+fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$	699
3.65	$\int (a+b \tan(e+fx))(c+d \tan(e+fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$	713
3.66	$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$	725
3.67	$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	735
3.68	$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	749
3.69	$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	764
3.70	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	778
3.71	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	791
3.72	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	803
3.73	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{c+d \tan(e+fx)} dx$	812
3.74	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))} dx$	820
3.75	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 (c+d \tan(e+fx))} dx$	829
3.76	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^3 (c+d \tan(e+fx))} dx$	840
3.77	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$	852
3.78	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$	867
3.79	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$	880
3.80	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^2} dx$	891
3.81	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$	900
3.82	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 (c+d \tan(e+fx))^2} dx$	911
3.83	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^3 (c+d \tan(e+fx))^2} dx$	923
3.84	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$	936
3.85	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$	950
3.86	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$	963
3.87	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^3} dx$	975
3.88	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^3} dx$	985
3.89	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 (c+d \tan(e+fx))^3} dx$	997
3.90	$\int (a+b \tan(e+fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$	1010
3.91	$\int (a+b \tan(e+fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$	1023

3.92	$\int (a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1035
3.93	$\int \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1046
3.94	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	1057
3.95	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	1068
3.96	$\int \frac{\sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	1079
3.97	$\int (a+b \tan(e+fx))^3 (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1092
3.98	$\int (a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1106
3.99	$\int (a+b \tan(e+fx)) (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1119
3.100	$\int (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1129
3.101	$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	1139
3.102	$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	1151
3.103	$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	1164
3.104	$\int (a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1176
3.105	$\int (a+b \tan(e+fx)) (c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1188
3.106	$\int (c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1200
3.107	$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$	1211
3.108	$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$	1224
3.109	$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$	1237
3.110	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1250
3.111	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1262
3.112	$\int \frac{(a+b \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1273
3.113	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx$	1284
3.114	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx$	1293
3.115	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}} dx$	1303
3.116	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1315
3.117	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1328
3.118	$\int \frac{(a+b \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1340
3.119	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^{3/2}} dx$	1350
3.120	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx)) (c+d \tan(e+fx))^{3/2}} dx$	1359
3.121	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{3/2}} dx$	1370
3.122	$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1383
3.123	$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1396

3.124	$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1407
3.125	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^{5/2}} dx$	1417
3.126	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{5/2}} dx$	1426
3.127	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{5/2}} dx$	1438
3.128	$\int (a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1451
3.129	$\int (a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1463
3.130	$\int \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1474
3.131	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$	1483
3.132	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$	1491
3.133	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$	1499
3.134	$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$	1509
3.135	$\int (a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1521
3.136	$\int \sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1533
3.137	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$	1544
3.138	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$	1553
3.139	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$	1564
3.140	$\int \frac{(c+d \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$	1574
3.141	$\int \sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1586
3.142	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$	1598
3.143	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$	1609
3.144	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$	1620
3.145	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$	1631
3.146	$\int \frac{(c+d \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{9/2}} dx$	1641
3.147	$\int \frac{(a+b \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1654
3.148	$\int \frac{(a+b \tan(e+fx))^{3/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1664
3.149	$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$	1673
3.150	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx$	1681
3.151	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}} dx$	1689
3.152	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$	1698
3.153	$\int \frac{(a+b \tan(e+fx))^{5/2}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1708

3.154	$\int \frac{(a+b \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1719
3.155	$\int \frac{\sqrt{a+b \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$	1729
3.156	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)} (c+d \tan(e+fx))^{3/2}} dx$	1737
3.157	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2} (c+d \tan(e+fx))^{3/2}} dx$	1746
3.158	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{5/2} (c+d \tan(e+fx))^{3/2}} dx$	1756
3.159	$\int \frac{(a+b \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1767
3.160	$\int \frac{(a+b \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1778
3.161	$\int \frac{\sqrt{a+b \tan(e+fx)} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$	1788
3.162	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)} (c+d \tan(e+fx))^{5/2}} dx$	1798
3.163	$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2} (c+d \tan(e+fx))^{5/2}} dx$	1808
3.164	$\int (a+b \tan(e+fx))^m (c+d \tan(e+fx))^n (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1819
3.165	$\int (a+b \tan(e+fx))^m (c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1826
3.166	$\int (a+b \tan(e+fx))^m (c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1839
3.167	$\int (a+b \tan(e+fx))^m (c+d \tan(e+fx)) (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1849
3.168	$\int (a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx)) dx$	1858
3.169	$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$	1865
3.170	$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$	1874
3.171	$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$	1884

3.1 $\int \tan(c+dx)(a+b \tan(c+dx)) (B \tan(c + dx) + C \tan^2$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	93
Fricas [A] (verification not implemented)	93
Sympy [A] (verification not implemented)	94
Maxima [A] (verification not implemented)	94
Giac [A] (verification not implemented)	95
Mupad [B] (verification not implemented)	95
Reduce [B] (verification not implemented)	96

Optimal result

Integrand size = 36, antiderivative size = 87

$$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((aB - bC)x + \frac{(bB + aC) \log(\cos(c + dx))}{d}$$

$$+ \frac{(aB - bC) \tan(c + dx)}{d} + \frac{(bB + aC) \tan^2(c + dx)}{2d} + \frac{bC \tan^3(c + dx)}{3d}$$

output

```
-(B*a-C*b)*x+(B*b+C*a)*ln(cos(d*x+c))/d+(B*a-C*b)*tan(d*x+c)/d+1/2*(B*b+C*a)*tan(d*x+c)^2/d+1/3*b*C*tan(d*x+c)^3/d
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.43

$$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{aB \arctan(\tan(c + dx))}{d} + \frac{bC \arctan(\tan(c + dx))}{d}$$

$$+ \frac{bB(2 \log(\cos(c + dx)) + \sec^2(c + dx))}{2d} + \frac{aC(2 \log(\cos(c + dx)) + \sec^2(c + dx))}{2d}$$

$$+ \frac{aB \tan(c + dx)}{d} - \frac{bC \tan(c + dx)}{d} + \frac{bC \tan^3(c + dx)}{3d}$$

input

```
Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
-((a*B*ArcTan[Tan[c + d*x]])/d) + (b*C*ArcTan[Tan[c + d*x]])/d + (b*B*(2*Log[Cos[c + d*x]] + Sec[c + d*x]^2))/(2*d) + (a*C*(2*Log[Cos[c + d*x]] + Sec[c + d*x]^2))/(2*d) + (a*B*Tan[c + d*x])/d - (b*C*Tan[c + d*x])/d + (b*C*Tan[c + d*x]^3)/(3*d)
```

Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4115, 3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan(c + dx)^2) dx \\
 & \quad \downarrow \text{4115} \\
 & \int \tan^2(c + dx)(a + b \tan(c + dx))(B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)^2(a + b \tan(c + dx))(B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \tan^2(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx + \frac{bC \tan^3(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)^2(aB - bC + (bB + aC) \tan(c + dx)) dx + \frac{bC \tan^3(c + dx)}{3d}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 4011 \\
& \int \tan(c+dx)(-bB-aC+(aB-bC)\tan(c+dx))dx + \frac{(aC+bB)\tan^2(c+dx)}{2d} + \frac{bC\tan^3(c+dx)}{3d} \\
& \downarrow 3042 \\
& \int \tan(c+dx)(-bB-aC+(aB-bC)\tan(c+dx))dx + \frac{(aC+bB)\tan^2(c+dx)}{2d} + \frac{bC\tan^3(c+dx)}{3d} \\
& \downarrow 4008 \\
& -(aC+bB)\int \tan(c+dx)dx + \frac{(aC+bB)\tan^2(c+dx)}{2d} + \frac{(aB-bC)\tan(c+dx)}{d} - x(aB-bC) + \frac{bC\tan^3(c+dx)}{3d} \\
& \downarrow 3042 \\
& -(aC+bB)\int \tan(c+dx)dx + \frac{(aC+bB)\tan^2(c+dx)}{2d} + \frac{(aB-bC)\tan(c+dx)}{d} - x(aB-bC) + \frac{bC\tan^3(c+dx)}{3d} \\
& \downarrow 3956 \\
& \frac{(aC+bB)\tan^2(c+dx)}{2d} + \frac{(aB-bC)\tan(c+dx)}{d} + \frac{(aC+bB)\log(\cos(c+dx))}{d} - x(aB-bC) + \frac{bC\tan^3(c+dx)}{3d}
\end{aligned}$$

input

```
Int[Tan[c + d*x]*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),
x]
```

output

```
-((a*B - b*C)*x) + ((b*B + a*C)*Log[Cos[c + d*x]])/d + ((a*B - b*C)*Tan[c
+ d*x])/d + ((b*B + a*C)*Tan[c + d*x]^2)/(2*d) + (b*C*Tan[c + d*x]^3)/(3*d
)
```


Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4115 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.01

method	result
norman	$(-Ba + Cb)x + \frac{(Ba - Cb)\tan(dx+c)}{d} + \frac{(Bb + Ca)\tan(dx+c)^2}{2d} + \frac{bC\tan(dx+c)^3}{3d} - \frac{(Bb + Ca)\ln(1 + \tan(dx+c))}{2d}$
parts	$\frac{(Bb + Ca)\left(\frac{\tan(dx+c)^2}{2} - \frac{\ln(1 + \tan(dx+c)^2)}{2}\right)}{d} + \frac{Ba(\tan(dx+c) - \arctan(\tan(dx+c)))}{d} + \frac{Cb\left(\frac{\tan(dx+c)^3}{3} - \tan(dx+c)\right)}{d}$
derivativdivides	$\frac{\frac{Cb\tan(dx+c)^3}{3} + \frac{Bb\tan(dx+c)^2}{2} + \frac{Ca\tan(dx+c)^2}{2} + Ba\tan(dx+c) - Cb\tan(dx+c) + \frac{(-Bb - Ca)\ln(1 + \tan(dx+c)^2)}{2}}{d} + (-Ba + Cb)$
default	$\frac{\frac{Cb\tan(dx+c)^3}{3} + \frac{Bb\tan(dx+c)^2}{2} + \frac{Ca\tan(dx+c)^2}{2} + Ba\tan(dx+c) - Cb\tan(dx+c) + \frac{(-Bb - Ca)\ln(1 + \tan(dx+c)^2)}{2}}{d} + (-Ba + Cb)$
parallelrisch	$-\frac{-2Cb\tan(dx+c)^3 + 6Badx - 3Bb\tan(dx+c)^2 - 6Cbdx - 3Ca\tan(dx+c)^2 + 3B\ln(1 + \tan(dx+c)^2)b - 6Ba\tan(dx+c)}{6d}$
risch	$-iBbx - iCax - Bax + Cbx - \frac{2iBbc}{d} - \frac{2iCac}{d} + \frac{2i(-3iBbe^{4i(dx+c)} - 3iCa e^{4i(dx+c)} + 3Ba e^{4i(dx+c)})}{6d}$

input `int (tan(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2), x, method=_RETURNVERBOSE)`

output $(-B*a+C*b)*x+(B*a-C*b)*\tan(d*x+c)/d+1/2*(B*b+C*a)*\tan(d*x+c)^2/d+1/3*b*C*\tan(d*x+c)^3/d-1/2*(B*b+C*a)/d*\ln(1+\tan(d*x+c)^2)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.98

$$\int \tan(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Cb \tan(dx + c)^3 - 6(Ba - Cb)dx + 3(Ca + Bb) \tan(dx + c)^2 + 3(Ca + Bb) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right) + 6(\dots)}{6d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2), x, algorithm="fricas")`

output

```
1/6*(2*C*b*tan(d*x + c)^3 - 6*(B*a - C*b)*d*x + 3*(C*a + B*b)*tan(d*x + c)
^2 + 3*(C*a + B*b)*log(1/(tan(d*x + c)^2 + 1)) + 6*(B*a - C*b)*tan(d*x + c
))/d
```

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.60

$$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} -Bax + \frac{Ba \tan(c+dx)}{d} - \frac{Bb \log(\tan^2(c+dx)+1)}{2d} + \frac{Bb \tan^2(c+dx)}{2d} - \frac{Ca \log(\tan^2(c+dx)+1)}{2d} + \frac{Ca \tan^2(c+dx)}{2d} + Cbx + \\ x(a + b \tan(c)) (B \tan(c) + C \tan^2(c)) \tan(c) \end{cases}$$

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((-B*a*x + B*a*tan(c + d*x)/d - B*b*log(tan(c + d*x)**2 + 1)/(2*d
) + B*b*tan(c + d*x)**2/(2*d) - C*a*log(tan(c + d*x)**2 + 1)/(2*d) + C*a*t
an(c + d*x)**2/(2*d) + C*b*x + C*b*tan(c + d*x)**3/(3*d) - C*b*tan(c + d*x
)/d, Ne(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*tan(c), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.99

$$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Cb \tan(dx + c)^3 + 3(Ca + Bb) \tan(dx + c)^2 - 6(Ba - Cb)(dx + c) - 3(Ca + Bb) \log(\tan(dx + c))^2}{6d}$$

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, alg
orithm="maxima")
```

output

```
1/6*(2*C*b*tan(d*x + c)^3 + 3*(C*a + B*b)*tan(d*x + c)^2 - 6*(B*a - C*b)*(
d*x + c) - 3*(C*a + B*b)*log(tan(d*x + c)^2 + 1) + 6*(B*a - C*b)*tan(d*x +
c))/d
```

Giac [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.37

$$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ba - Cb)(dx + c)}{d} - \frac{(Ca + Bb) \log(\tan(dx + c)^2 + 1)}{2d} + \frac{2Cbd^2 \tan(dx + c)^3 + 3Cad^2 \tan(dx + c)^2 + 3Bbd^2 \tan(dx + c)^2 + 6Bad^2 \tan(dx + c) - 6Cbd^2 \tan(dx + c)}{6d^3}$$

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, alg
orithm="giac")
```

output

```
-(B*a - C*b)*(d*x + c)/d - 1/2*(C*a + B*b)*log(tan(d*x + c)^2 + 1)/d + 1/6
*(2*C*b*d^2*tan(d*x + c)^3 + 3*C*a*d^2*tan(d*x + c)^2 + 3*B*b*d^2*tan(d*x
+ c)^2 + 6*B*a*d^2*tan(d*x + c) - 6*C*b*d^2*tan(d*x + c))/d^3
```

Mupad [B] (verification not implemented)

Time = 5.62 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.97

$$\int \tan(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\tan(c + dx) (Ba - Cb) - \ln(\tan(c + dx)^2 + 1) \left(\frac{Bb}{2} + \frac{Ca}{2}\right) + \tan(c + dx)^2 \left(\frac{Bb}{2} + \frac{Ca}{2}\right) - dx (Ba - Cb)}{d}$$

input

```
int(tan(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)),
x)
```

output

```
(tan(c + d*x)*(B*a - C*b) - log(tan(c + d*x)^2 + 1)*((B*b)/2 + (C*a)/2) +
tan(c + d*x)^2*((B*b)/2 + (C*a)/2) - d*x*(B*a - C*b) + (C*b*tan(c + d*x)^3
)/3)/d
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.22

$$\int \tan(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-3 \log(\tan(dx + c)^2 + 1) ac - 3 \log(\tan(dx + c)^2 + 1) b^2 + 2 \tan(dx + c)^3 bc + 3 \tan(dx + c)^2 ac + 3 \tan(dx + c) ab - 6 \tan(dx + c) b^2 c + 6 a^2 b dx + 6 b^2 c dx}{6d}$$

input

```
int(tan(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
( - 3*log(tan(c + d*x)**2 + 1)*a*c - 3*log(tan(c + d*x)**2 + 1)*b**2 + 2*tan(c + d*x)**3*b*c + 3*tan(c + d*x)**2*a*c + 3*tan(c + d*x)**2*b**2 + 6*tan(c + d*x)*a*b - 6*tan(c + d*x)*b*c - 6*a*b*d*x + 6*b*c*d*x)/(6*d)
```

3.2 $\int (a+b \tan(c+dx)) (B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	97
Mathematica [A] (verified)	97
Rubi [A] (verified)	98
Maple [A] (verified)	100
Fricas [A] (verification not implemented)	100
Sympy [A] (verification not implemented)	101
Maxima [A] (verification not implemented)	101
Giac [A] (verification not implemented)	102
Mupad [B] (verification not implemented)	102
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 30, antiderivative size = 66

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((bB + aC)x) - \frac{(aB - bC) \log(\cos(c + dx))}{d}$$

$$+ \frac{bB \tan(c + dx)}{d} + \frac{C(a + b \tan(c + dx))^2}{2bd}$$

output `-(B*b+C*a)*x-(B*a-C*b)*ln(cos(d*x+c))/d+b*B*tan(d*x+c)/d+1/2*C*(a+b*tan(d*x+c))^2/b/d`

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.02

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-2(bB + aC) \arctan(\tan(c + dx)) + 2(-aB + bC) \log(\cos(c + dx)) + bC \sec^2(c + dx) + 2(bB + aC) \tan(c + dx)}{2d}$$

input `Integrate[(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output

$$(-2*(b*B + a*C)*ArcTan[Tan[c + d*x]] + 2*(-(a*B) + b*C)*Log[Cos[c + d*x]] + b*C*Sec[c + d*x]^2 + 2*(b*B + a*C)*Tan[c + d*x])/(2*d)$$

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4113, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ & \quad \downarrow 3042 \\ & \int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan(c + dx)^2) dx \\ & \quad \downarrow 4113 \\ & \int (a + b \tan(c + dx))(B \tan(c + dx) - C) dx + \frac{C(a + b \tan(c + dx))^2}{2bd} \\ & \quad \downarrow 3042 \\ & \int (a + b \tan(c + dx))(B \tan(c + dx) - C) dx + \frac{C(a + b \tan(c + dx))^2}{2bd} \\ & \quad \downarrow 4008 \\ & (aB - bC) \int \tan(c + dx) dx - x(aC + bB) + \frac{C(a + b \tan(c + dx))^2}{2bd} + \frac{bB \tan(c + dx)}{d} \\ & \quad \downarrow 3042 \\ & (aB - bC) \int \tan(c + dx) dx - x(aC + bB) + \frac{C(a + b \tan(c + dx))^2}{2bd} + \frac{bB \tan(c + dx)}{d} \\ & \quad \downarrow 3956 \\ & -\frac{(aB - bC) \log(\cos(c + dx))}{d} - x(aC + bB) + \frac{C(a + b \tan(c + dx))^2}{2bd} + \frac{bB \tan(c + dx)}{d} \end{aligned}$$

input `Int[(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `-((b*B + a*C)*x) - ((a*B - b*C)*Log[Cos[c + d*x]])/d + (b*B*Tan[c + d*x])/d + (C*(a + b*Tan[c + d*x])^2)/(2*b*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :=> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :=> Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] :=> Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.05

method	result
norman	$(-Bb - Ca)x + \frac{(Bb+Ca)\tan(dx+c)}{d} + \frac{Cb\tan(dx+c)^2}{2d} + \frac{(Ba-Cb)\ln(1+\tan(dx+c)^2)}{2d}$
derivativedivides	$\frac{\frac{C\tan(dx+c)^2b}{2} + B\tan(dx+c)b + C\tan(dx+c)a + \frac{(Ba-Cb)\ln(1+\tan(dx+c)^2)}{2}}{d} + (-Bb-Ca)\arctan(\tan(dx+c))$
default	$\frac{\frac{C\tan(dx+c)^2b}{2} + B\tan(dx+c)b + C\tan(dx+c)a + \frac{(Ba-Cb)\ln(1+\tan(dx+c)^2)}{2}}{d} + (-Bb-Ca)\arctan(\tan(dx+c))$
parts	$\frac{(Bb+Ca)(\tan(dx+c) - \arctan(\tan(dx+c)))}{d} + \frac{Ba\ln(1+\tan(dx+c)^2)}{2d} + \frac{Cb\left(\frac{\tan(dx+c)^2}{2} - \frac{\ln(1+\tan(dx+c)^2)}{2}\right)}{d}$
parallelrisc	$\frac{-2Bbdx - 2Cadx + C\tan(dx+c)^2b + B\ln(1+\tan(dx+c)^2)a + 2B\tan(dx+c)b - C\ln(1+\tan(dx+c)^2)b + 2C\tan(dx+c)c}{2d}$
risc	$-Bbx - Cax + iBax - iCb x + \frac{2iBac}{d} - \frac{2iCbc}{d} + \frac{2i(-iCb e^{2i(dx+c)} + Bb e^{2i(dx+c)} + Ca e^{2i(dx+c)} + B)}{d(e^{2i(dx+c)} + 1)^2}$

input `int((a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `(-B*b-C*a)*x+(B*b+C*a)/d*tan(d*x+c)+1/2*C*b/d*tan(d*x+c)^2+1/2*(B*a-C*b)/d*ln(1+tan(d*x+c)^2)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Cb \tan(dx + c)^2 - 2(Ca + Bb)dx - (Ba - Cb) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right) + 2(Ca + Bb) \tan(dx + c)}{2d}$$

input `integrate((a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output

$$\frac{1}{2}*(C*b*\tan(dx + c)^2 - 2*(C*a + B*b)*dx - (B*a - C*b)*\log(1/(\tan(dx + c)^2 + 1)) + 2*(C*a + B*b)*\tan(dx + c))/d$$

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.59

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \frac{Ba \log(\tan^2(c+dx)+1)}{2d} - Bbx + \frac{Bb \tan(c+dx)}{d} - Cax + \frac{Ca \tan(c+dx)}{d} - \frac{Cb \log(\tan^2(c+dx)+1)}{2d} + \frac{Cb \tan^2(c+dx)}{2d} & \text{for } \\ x(a + b \tan(c)) (B \tan(c) + C \tan^2(c)) & \text{oth} \end{cases}$$

input

```
integrate((a+b*tan(dx+c))*(B*tan(dx+c)+C*tan(dx+c)**2),x)
```

output

```
Piecewise((B*a*log(tan(c + dx)**2 + 1)/(2*d) - B*b*x + B*b*tan(c + dx)/d - C*a*x + C*a*tan(c + dx)/d - C*b*log(tan(c + dx)**2 + 1)/(2*d) + C*b*tan(c + dx)**2/(2*d), Ne(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Cb \tan(dx + c)^2 - 2(Ca + Bb)(dx + c) + (Ba - Cb) \log(\tan(dx + c)^2 + 1) + 2(Ca + Bb) \tan(dx + c)}{2d}$$

input

```
integrate((a+b*tan(dx+c))*(B*tan(dx+c)+C*tan(dx+c)^2),x, algorithm="maxima")
```

output

$$\frac{1}{2}*(C*b*\tan(dx + c)^2 - 2*(C*a + B*b)*(dx + c) + (B*a - C*b)*\log(\tan(dx + c)^2 + 1) + 2*(C*a + B*b)*\tan(dx + c))/d$$

Giac [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.24

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ca + Bb)(dx + c)}{d} + \frac{(Ba - Cb) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{Cbd \tan(dx + c)^2 + 2Cad \tan(dx + c) + 2Bbd \tan(dx + c)}{2d^2}$$

input

```
integrate((a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

```
-(C*a + B*b)*(d*x + c)/d + 1/2*(B*a - C*b)*log(tan(d*x + c)^2 + 1)/d + 1/2*(C*b*d*tan(d*x + c)^2 + 2*C*a*d*tan(d*x + c) + 2*B*b*d*tan(d*x + c))/d^2
```

Mupad [B] (verification not implemented)

Time = 5.22 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.95

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\tan(c + dx) (Bb + Ca) + \ln(\tan(c + dx)^2 + 1) \left(\frac{Ba}{2} - \frac{Cb}{2}\right) - dx (Bb + Ca) + \frac{Cb \tan(c + dx)^2}{2}}{d}$$

input

```
int((B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)),x)
```

output

```
(tan(c + d*x)*(B*b + C*a) + log(tan(c + d*x)^2 + 1)*((B*a)/2 - (C*b)/2) - d*x*(B*b + C*a) + (C*b*tan(c + d*x)^2)/2)/d
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.21

$$\int (a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\log(\tan(dx + c)^2 + 1) ab - \log(\tan(dx + c)^2 + 1) bc + \tan(dx + c)^2 bc + 2 \tan(dx + c) ac + 2 \tan(dx + c) b^2 dx - 2 a c dx - 2 b^2 dx}{2d}$$

input

```
int((a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
(log(tan(c + d*x)**2 + 1)*a*b - log(tan(c + d*x)**2 + 1)*b*c + tan(c + d*x)**2*b*c + 2*tan(c + d*x)*a*c + 2*tan(c + d*x)*b**2 - 2*a*c*d*x - 2*b**2*d*x)/(2*d)
```

3.3 $\int \cot(c+dx)(a+b \tan(c+dx)) (B \tan(c + dx) + C \tan^2$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [A] (verified)	107
Fricas [A] (verification not implemented)	107
Sympy [B] (verification not implemented)	108
Maxima [A] (verification not implemented)	108
Giac [A] (verification not implemented)	109
Mupad [B] (verification not implemented)	109
Reduce [B] (verification not implemented)	110

Optimal result

Integrand size = 36, antiderivative size = 42

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (aB - bC)x - \frac{(bB + aC) \log(\cos(c + dx))}{d} + \frac{bC \tan(c + dx)}{d}$$

output

```
(B*a-C*b)*x-(B*b+C*a)*ln(cos(d*x+c))/d+b*C*tan(d*x+c)/d
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.40

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= aBx - \frac{bC \arctan(\tan(c + dx))}{d} - \frac{bB \log(\cos(c + dx))}{d}$$

$$- \frac{aC \log(\cos(c + dx))}{d} + \frac{bC \tan(c + dx)}{d}$$

input

```
Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
a*B*x - (b*C*ArcTan[Tan[c + d*x]])/d - (b*B*Log[Cos[c + d*x]])/d - (a*C*Log[Cos[c + d*x]])/d + (b*C*Tan[c + d*x])/d
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4115, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx))}{\tan(c + dx)} dx$$

$$\downarrow 4115$$

$$\int (a + b \tan(c + dx))(B + C \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(c + dx))(B + C \tan(c + dx)) dx$$

$$\downarrow 4008$$

$$(aC + bB) \int \tan(c + dx) dx + x(aB - bC) + \frac{bC \tan(c + dx)}{d}$$

$$\downarrow 3042$$

$$(aC + bB) \int \tan(c + dx) dx + x(aB - bC) + \frac{bC \tan(c + dx)}{d}$$

$$\downarrow 3956$$

$$-\frac{(aC + bB) \log(\cos(c + dx))}{d} + x(aB - bC) + \frac{bC \tan(c + dx)}{d}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]`

output `(a*B - b*C)*x - ((b*B + a*C)*Log[Cos[c + d*x]])/d + (b*C*Tan[c + d*x])/d`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4115 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]`

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.10

method	result
parallelrisc	$\frac{(Bb+Ca) \ln(\sec(dx+c)^2) + 2Cb \tan(dx+c) + 2dx(Ba-Cb)}{2d}$
norman	$(Ba - Cb)x + \frac{bC \tan(dx+c)}{d} + \frac{(Bb+Ca) \ln(1+\tan(dx+c)^2)}{2d}$
derivativedivides	$\frac{-Bb \ln(\cos(dx+c)) + Cb(\tan(dx+c) - dx - c) + Ba(dx+c) - Ca \ln(\cos(dx+c))}{d}$
default	$\frac{-Bb \ln(\cos(dx+c)) + Cb(\tan(dx+c) - dx - c) + Ba(dx+c) - Ca \ln(\cos(dx+c))}{d}$
risc	$iBbx + iCax + Bax - Cbx + \frac{2iBbc}{d} + \frac{2iCac}{d} + \frac{2iCb}{d(e^{2i(dx+c)}+1)} - \frac{\ln(e^{2i(dx+c)}+1)Bb}{d} - \frac{\ln(e^{2i(dx+c)}-1)Cb}{d}$

input `int(cot(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/2*((B*b+C*a)*ln(sec(d*x+c)^2)+2*C*b*tan(d*x+c)+2*d*x*(B*a-C*b))/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int \cot(c+dx)(a+b \tan(c+dx))(B \tan(c+dx)+C \tan^2(c+dx)) dx$$

$$= \frac{2(Ba-Cb)dx + 2Cb \tan(dx+c) - (Ca+Bb) \log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output `1/2*(2*(B*a - C*b)*d*x + 2*C*b*tan(d*x + c) - (C*a + B*b)*log(1/(tan(d*x + c)^2 + 1)))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(36) = 72.

Time = 0.31 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.95

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} Bax + \frac{Bb \log(\tan^2(c+dx)+1)}{2d} + \frac{Ca \log(\tan^2(c+dx)+1)}{2d} - Cbx + \frac{Cb \tan(c+dx)}{d} & \text{for } d \neq 0 \\ x(a + b \tan(c)) (B \tan(c) + C \tan^2(c)) \cot(c) & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((B*a*x + B*b*log(tan(c + d*x)**2 + 1)/(2*d) + C*a*log(tan(c + d*x)**2 + 1)/(2*d) - C*b*x + C*b*tan(c + d*x)/d, Ne(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*cot(c), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Cb \tan(dx + c) + 2(Ba - Cb)(dx + c) + (Ca + Bb) \log(\tan(dx + c)^2 + 1)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")`

output `1/2*(2*C*b*tan(d*x + c) + 2*(B*a - C*b)*(d*x + c) + (C*a + B*b)*log(tan(d*x + c)^2 + 1))/d`

Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.26

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Cb \tan(dx + c)}{d} + \frac{(Ba - Cb)(dx + c)}{d} + \frac{(Ca + Bb) \log(\tan(dx + c)^2 + 1)}{2d}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

```
C*b*tan(d*x + c)/d + (B*a - C*b)*(d*x + c)/d + 1/2*(C*a + B*b)*log(tan(d*x + c)^2 + 1)/d
```

Mupad [B] (verification not implemented)

Time = 5.29 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.38

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= B a x - C b x + \frac{C b \tan(c + dx)}{d}$$

$$+ \frac{B b \ln(\tan(c + dx)^2 + 1)}{2d} + \frac{C a \ln(\tan(c + dx)^2 + 1)}{2d}$$

input

```
int(cot(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)), x)
```

output

```
B*a*x - C*b*x + (C*b*tan(c + d*x))/d + (B*b*log(tan(c + d*x)^2 + 1))/(2*d) + (C*a*log(tan(c + d*x)^2 + 1))/(2*d)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 182, normalized size of antiderivative = 4.33

$$\int \cot(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\cos(dx + c) \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) ac + \cos(dx + c) \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) b^2 - \cos(dx + c) \log(\tan(dx + c))}{\cos(dx + c)}$$

input `int(cot(d*x+c)*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `(cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*a*c + cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*b**2 - cos(c + d*x)*log(tan((c + d*x)/2) - 1)*a*c - cos(c + d*x)*log(tan((c + d*x)/2) - 1)*b**2 - cos(c + d*x)*log(tan((c + d*x)/2) + 1)*a*c - cos(c + d*x)*log(tan((c + d*x)/2) + 1)*b**2 + cos(c + d*x)*a*b*d*x - cos(c + d*x)*b*c*d*x + sin(c + d*x)*b*c)/(cos(c + d*x)*d)`

3.4 $\int \cot^2(c+dx)(a+b \tan(c+dx)) (B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	111
Mathematica [A] (verified)	111
Rubi [A] (verified)	112
Maple [A] (verified)	114
Fricas [A] (verification not implemented)	115
Sympy [B] (verification not implemented)	115
Maxima [A] (verification not implemented)	116
Giac [A] (verification not implemented)	116
Mupad [B] (verification not implemented)	117
Reduce [B] (verification not implemented)	117

Optimal result

Integrand size = 38, antiderivative size = 37

$$\int \cot^2(c+dx)(a+b \tan(c+dx)) (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= (bB + aC)x - \frac{bC \log(\cos(c+dx))}{d} + \frac{aB \log(\sin(c+dx))}{d}$$

output `(B*b+C*a)*x-b*C*ln(cos(d*x+c))/d+a*B*ln(sin(d*x+c))/d`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.97

$$\int \cot^2(c+dx)(a+b \tan(c+dx)) (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= bBx + aCx - \frac{bC \log(\cos(c+dx))}{d} + \frac{aB \log(\sin(c+dx))}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `b*B*x + a*C*x - (b*C*Log[Cos[c + d*x]])/d + (a*B*Log[Sin[c + d*x]])/d`

Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3042, 4115, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c+dx)(a+b\tan(c+dx))(B\tan(c+dx)+C\tan^2(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+b\tan(c+dx))(B\tan(c+dx)+C\tan^2(c+dx))}{\tan^2(c+dx)} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \cot(c+dx)(a+b\tan(c+dx))(B+C\tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+b\tan(c+dx))(B+C\tan(c+dx))}{\tan(c+dx)} dx \\
 & \quad \downarrow \text{4072} \\
 & \int \cot(c+dx)(aB+(bB+aC)\tan(c+dx)) dx + bC \int \tan(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aB+(bB+aC)\tan(c+dx)}{\tan(c+dx)} dx + bC \int \tan(c+dx) dx \\
 & \quad \downarrow \text{3956} \\
 & \int \frac{aB+(bB+aC)\tan(c+dx)}{\tan(c+dx)} dx - \frac{bC \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{4014} \\
 & aB \int \cot(c+dx) dx + x(aC+bB) - \frac{bC \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
 & aB \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + x(aC + bB) - \frac{bC \log(\cos(c + dx))}{d} \\
 & \quad \downarrow 25 \\
 & -aB \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + x(aC + bB) - \frac{bC \log(\cos(c + dx))}{d} \\
 & \quad \downarrow 3956 \\
 & x(aC + bB) + \frac{aB \log(-\sin(c + dx))}{d} - \frac{bC \log(\cos(c + dx))}{d}
 \end{aligned}$$

input

```
Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(b*B + a*C)*x - (b*C*Log[Cos[c + d*x]])/d + (a*B*Log[-Sin[c + d*x]])/d
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]
```

rule 4072

```
Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16

method	result	size
derivativedivides	$\frac{Bb(dx+c) - Cb \ln(\cos(dx+c)) + Ba \ln(\sin(dx+c)) + Ca(dx+c)}{d}$	43
default	$\frac{Bb(dx+c) - Cb \ln(\cos(dx+c)) + Ba \ln(\sin(dx+c)) + Ca(dx+c)}{d}$	43
parallelrisch	$\frac{(-Ba+Cb) \ln(\sec(dx+c)^2) + 2Ba \ln(\tan(dx+c)) + 2x(Bb+Ca)d}{2d}$	47
norman	$(Bb + Ca)x + \frac{Ba \ln(\tan(dx+c))}{d} - \frac{(Ba-Cb) \ln(1+\tan(dx+c)^2)}{2d}$	48
risch	$Bbx + Cax - iBax + iCbx + \frac{2iCbc}{d} - \frac{2iBac}{d} - \frac{\ln(e^{2i(dx+c)}+1)Cb}{d} + \frac{Ba \ln(e^{2i(dx+c)}-1)}{d}$	77

input

```
int(cot(d*x+c)^2*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)
```

output

```
1/d*(B*b*(d*x+c)-C*b*ln(cos(d*x+c))+B*a*ln(sin(d*x+c))+C*a*(d*x+c))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59

$$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2(Ca + Bb)dx + Ba \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) - Cb \log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")`

output `1/2*(2*(C*a + B*b)*d*x + B*a*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - C*b*log(1/(tan(d*x + c)^2 + 1)))/d`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(34) = 68.

Time = 0.45 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.30

$$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} -\frac{Ba \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba \log(\tan(c+dx))}{d} + Bbx + Cax + \frac{Cb \log(\tan^2(c+dx)+1)}{2d} & \text{for } d \neq 0 \\ x(a + b \tan(c)) (B \tan(c) + C \tan^2(c)) \cot^2(c) & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((-B*a*log(tan(c + d*x)**2 + 1)/(2*d) + B*a*log(tan(c + d*x))/d + B*b*x + C*a*x + C*b*log(tan(c + d*x)**2 + 1)/(2*d), Ne(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*cot(c)**2, True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.41

$$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Ba \log(\tan(dx + c)) + 2(Ca + Bb)(dx + c) - (Ba - Cb) \log(\tan(dx + c)^2 + 1)}{2d}$$

input

```
integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
1/2*(2*B*a*log(tan(d*x + c)) + 2*(C*a + B*b)*(d*x + c) - (B*a - C*b)*log(tan(d*x + c)^2 + 1))/d
```

Giac [A] (verification not implemented)

Time = 0.62 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.49

$$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Ba \log(|\tan(dx + c)|)}{d} + \frac{(Ca + Bb)(dx + c)}{d} - \frac{(Ba - Cb) \log(\tan(dx + c)^2 + 1)}{2d}$$

input

```
integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

```
B*a*log(abs(tan(d*x + c)))/d + (C*a + B*b)*(d*x + c)/d - 1/2*(B*a - C*b)*log(tan(d*x + c)^2 + 1)/d
```

Mupad [B] (verification not implemented)

Time = 5.32 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.86

$$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{B a \ln(\tan(c + dx))}{d} - \frac{\ln(\tan(c + dx) - i) (B + C i) (a + b i)}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (B - C i) (b + a i) i}{2 d}$$

input `int(cot(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b)*1i)/(2*d) - (log(tan(c + d*x) - 1i)*(B + C*1i)*(a + b*1i))/(2*d) + (B*a*log(tan(c + d*x)))/d`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 96, normalized size of antiderivative = 2.59

$$\int \cot^2(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-\log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) ab + \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) bc - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) - 1\right) bc - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right) + 1\right) bc}{d}$$

input `int(cot(d*x+c)^2*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `(- log(tan((c + d*x)/2)**2 + 1)*a*b + log(tan((c + d*x)/2)**2 + 1)*b*c - log(tan((c + d*x)/2) - 1)*b*c - log(tan((c + d*x)/2) + 1)*b*c + log(tan((c + d*x)/2))*a*b + a*c*d*x + b**2*d*x)/d`

3.5 $\int \cot^3(c+dx)(a+b \tan(c+dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	118
Mathematica [C] (verified)	118
Rubi [A] (verified)	119
Maple [A] (verified)	121
Fricas [A] (verification not implemented)	122
Sympy [B] (verification not implemented)	122
Maxima [A] (verification not implemented)	123
Giac [A] (verification not implemented)	123
Mupad [B] (verification not implemented)	124
Reduce [B] (verification not implemented)	124

Optimal result

Integrand size = 38, antiderivative size = 43

$$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((aB - bC)x) - \frac{aB \cot(c + dx)}{d} + \frac{(bB + aC) \log(\sin(c + dx))}{d}$$

output

```
-(B*a-C*b)*x-a*B*cot(d*x+c)/d+(B*b+C*a)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.44

$$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= bCx - \frac{aB \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c + dx)\right)}{d}$$

$$+ \frac{bB \log(\sin(c + dx))}{d} + \frac{aC \log(\sin(c + dx))}{d}$$

input

```
Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
b*C*x - (a*B*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d + (b*B*Log[Sin[c + d*x]])/d + (a*C*Log[Sin[c + d*x]])/d
```

Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {3042, 4115, 3042, 4074, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx))}{\tan^3(c + dx)} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \cot^2(c + dx)(a + b \tan(c + dx))(B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))(B + C \tan(c + dx))}{\tan^2(c + dx)} dx \\
 & \quad \downarrow \text{4074} \\
 & \int \cot(c + dx)(bB + aC - (aB - bC) \tan(c + dx)) dx - \frac{aB \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{bB + aC - (aB - bC) \tan(c + dx)}{\tan(c + dx)} dx - \frac{aB \cot(c + dx)}{d} \\
 & \quad \downarrow \text{4014}
 \end{aligned}$$

$$\begin{aligned}
& (aC + bB) \int \cot(c + dx) dx - (x(aB - bC)) - \frac{aB \cot(c + dx)}{d} \\
& \quad \downarrow \text{3042} \\
& (aC + bB) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - (x(aB - bC)) - \frac{aB \cot(c + dx)}{d} \\
& \quad \downarrow \text{25} \\
& -(aC + bB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - (x(aB - bC)) - \frac{aB \cot(c + dx)}{d} \\
& \quad \downarrow \text{3956} \\
& \frac{(aC + bB) \log(-\sin(c + dx))}{d} - (x(aB - bC)) - \frac{aB \cot(c + dx)}{d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
-((a*B - b*C)*x) - (a*B*Cot[c + d*x])/d + ((b*B + a*C)*Log[-Sin[c + d*x]])/d
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

rule 4074

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e
_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{Bb \ln(\sin(dx+c)) + Cb(dx+c) + Ba(-\cot(dx+c) - dx - c) + Ca \ln(\sin(dx+c))}{d}$
default	$\frac{Bb \ln(\sin(dx+c)) + Cb(dx+c) + Ba(-\cot(dx+c) - dx - c) + Ca \ln(\sin(dx+c))}{d}$
parallelrisch	$\frac{(-Bb - Ca) \ln(\sec(dx+c)^2) + (2Bb + 2Ca) \ln(\tan(dx+c)) - 2Ba \cot(dx+c) - 2dx(Ba - Cb)}{2d}$
norman	$\frac{(-Ba + Cb)x \tan(dx+c)^2 - \frac{Ba \tan(dx+c)}{d}}{\tan(dx+c)^2} + \frac{(Bb + Ca) \ln(\tan(dx+c))}{d} - \frac{(Bb + Ca) \ln(1 + \tan(dx+c)^2)}{2d}$
risch	$-iBbx - iCax - Bax + Cbx - \frac{2iBbc}{d} - \frac{2iCac}{d} - \frac{2iBa}{d(e^{2i(dx+c)} - 1)} + \frac{\ln(e^{2i(dx+c)} - 1)Bb}{d} + \frac{\ln(e^{2i(dx+c)} - 1)Ca}{d}$

input `int(cot(d*x+c)^3*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(B*b*ln(sin(d*x+c))+C*b*(d*x+c)+B*a*(-cot(d*x+c)-d*x-c)+C*a*ln(sin(d*x+c)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.70

$$\int \cot^3(c+dx)(a+b \tan(c+dx))(B \tan(c+dx)+C \tan^2(c+dx)) dx$$

$$= -\frac{2(Ba-Cb)dx \tan(dx+c) - (Ca+Bb) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c) + 2Ba}{2d \tan(dx+c)}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output `-1/2*(2*(B*a - C*b)*d*x*tan(d*x + c) - (C*a + B*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c) + 2*B*a)/(d*tan(d*x + c))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. 2(36) = 72.

Time = 0.68 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.70

$$\int \cot^3(c+dx)(a+b \tan(c+dx))(B \tan(c+dx)+C \tan^2(c+dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a+b \tan(c))(B \tan(c)+C \tan^2(c)) \cot^3(c) \\ \text{NaN} \\ -Bax - \frac{Ba}{d \tan(c+dx)} - \frac{Bb \log(\tan^2(c+dx)+1)}{2d} + \frac{Bb \log(\tan(c+dx))}{d} - \frac{Ca \log(\tan^2(c+dx)+1)}{2d} + \frac{Ca \log(\tan(c+dx))}{d} + Cb \end{cases}$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*cot(c)**3, Eq(d, 0)), (nan, Eq(c, -d*x)), (-B*a*x - B*a/(d*tan(c + d*x)) - B*b*log(tan(c + d*x)**2 + 1)/(2*d) + B*b*log(tan(c + d*x))/d - C*a*log(tan(c + d*x)**2 + 1)/(2*d) + C*a*log(tan(c + d*x))/d + C*b*x, True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.58

$$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$-\frac{2(Ba - Cb)(dx + c) + (Ca + Bb) \log(\tan(dx + c)^2 + 1) - 2(Ca + Bb) \log(\tan(dx + c)) + \frac{2Ba}{\tan(dx + c)}}{2d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")`

output `-1/2*(2*(B*a - C*b)*(d*x + c) + (C*a + B*b)*log(tan(d*x + c)^2 + 1) - 2*(C*a + B*b)*log(tan(d*x + c)) + 2*B*a/tan(d*x + c))/d`

Giac [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.77

$$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ba - Cb)(dx + c)}{d} - \frac{(Ca + Bb) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{(Ca + Bb) \log(|\tan(dx + c)|)}{d} - \frac{Ba}{d \tan(dx + c)}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")`

output

$$-(B*a - C*b)*(d*x + c)/d - 1/2*(C*a + B*b)*\log(\tan(d*x + c)^2 + 1)/d + (C*a + B*b)*\log(\text{abs}(\tan(d*x + c)))/d - B*a/(d*\tan(d*x + c))$$
Mupad [B] (verification not implemented)

Time = 5.28 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.02

$$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (Bb + Ca)}{d} - \frac{\ln(\tan(c + dx) + 1i) (B - C 1i) (b + a 1i)}{2d}$$

$$- \frac{Ba \cot(c + dx)}{d} + \frac{\ln(\tan(c + dx) - 1i) (B + C 1i) (a + b 1i) 1i}{2d}$$

input

$$\text{int}(\cot(c + d*x)^3*(B*\tan(c + d*x) + C*\tan(c + d*x)^2)*(a + b*\tan(c + d*x)),x)$$

output

$$\frac{(\log(\tan(c + d*x))*(B*b + C*a))/d + (\log(\tan(c + d*x) - 1i)*(B + C*1i)*(a + b*1i)*1i)/(2*d) - (\log(\tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b))/(2*d) - (B*a*\cot(c + d*x))/d}$$
Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 134, normalized size of antiderivative = 3.12

$$\int \cot^3(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-\cos(dx + c)ab - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right)\sin(dx + c)ac - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right)\sin(dx + c)b^2 + \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right)\sin(dx + c)ab + \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right)\sin(dx + c)ac - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right)\sin(dx + c)b^2 + \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right)\sin(dx + c)ab}{\sin(dx + c)}$$

input

$$\text{int}(\cot(d*x+c)^3*(a+b*\tan(d*x+c))*(B*\tan(d*x+c)+C*\tan(d*x+c)^2),x)$$

output

$$\frac{(-\cos(c + d*x)*a*b - \log(\tan((c + d*x)/2)**2 + 1)*\sin(c + d*x)*a*c - \log(\tan((c + d*x)/2)**2 + 1)*\sin(c + d*x)*b**2 + \log(\tan((c + d*x)/2))*\sin(c + d*x)*a*c + \log(\tan((c + d*x)/2))*\sin(c + d*x)*b**2 - \sin(c + d*x)*a*b*d*x + \sin(c + d*x)*b*c*d*x)/(\sin(c + d*x)*d)}$$

3.6 $\int \cot^4(c+dx)(a+b \tan(c+dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	125
Mathematica [C] (verified)	126
Rubi [A] (verified)	126
Maple [A] (verified)	129
Fricas [A] (verification not implemented)	130
Sympy [B] (verification not implemented)	130
Maxima [A] (verification not implemented)	131
Giac [A] (verification not implemented)	131
Mupad [B] (verification not implemented)	132
Reduce [B] (verification not implemented)	132

Optimal result

Integrand size = 38, antiderivative size = 66

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((bB + aC)x) - \frac{(bB + aC) \cot(c + dx)}{d}$$

$$- \frac{aB \cot^2(c + dx)}{2d} - \frac{(aB - bC) \log(\sin(c + dx))}{d}$$

output

```
-(B*b+C*a)*x-(B*b+C*a)*cot(d*x+c)/d-1/2*a*B*cot(d*x+c)^2/d-(B*a-C*b)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.62

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{aB \csc^2(c + dx)}{2d} - \frac{bB \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c + dx)\right)}{d}$$

$$- \frac{aC \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c + dx)\right)}{d}$$

$$- \frac{aB \log(\sin(c + dx))}{d} + \frac{bC \log(\sin(c + dx))}{d}$$

input

```
Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
-1/2*(a*B*Csc[c + d*x]^2)/d - (b*B*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d - (a*C*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d - (a*B*Log[Sin[c + d*x]])/d + (b*C*Log[Sin[c + d*x]])/d
```

Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4115, 3042, 4074, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx))^2}{\tan^4(c + dx)} dx$$

$$\begin{aligned}
& \downarrow 4115 \\
& \int \cot^3(c+dx)(a+b \tan(c+dx))(B+C \tan(c+dx))dx \\
& \downarrow 3042 \\
& \int \frac{(a+b \tan(c+dx))(B+C \tan(c+dx))}{\tan(c+dx)^3} dx \\
& \downarrow 4074 \\
& \int \cot^2(c+dx)(bB+aC-(aB-bC) \tan(c+dx))dx - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 3042 \\
& \int \frac{bB+aC-(aB-bC) \tan(c+dx)}{\tan(c+dx)^2} dx - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 4012 \\
& \int -\cot(c+dx)(aB-bC+(bB+aC) \tan(c+dx))dx - \frac{(aC+bB) \cot(c+dx)}{d} - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 25 \\
& -\int \cot(c+dx)(aB-bC+(bB+aC) \tan(c+dx))dx - \frac{(aC+bB) \cot(c+dx)}{d} - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 3042 \\
& -\int \frac{aB-bC+(bB+aC) \tan(c+dx)}{\tan(c+dx)} dx - \frac{(aC+bB) \cot(c+dx)}{d} - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 4014 \\
& -(aB-bC) \int \cot(c+dx) dx - \frac{(aC+bB) \cot(c+dx)}{d} - x(aC+bB) - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 3042 \\
& -(aB-bC) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{(aC+bB) \cot(c+dx)}{d} - x(aC+bB) - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 25 \\
& (aB-bC) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{(aC+bB) \cot(c+dx)}{d} - x(aC+bB) - \frac{aB \cot^2(c+dx)}{2d} \\
& \downarrow 3956
\end{aligned}$$

$$-\frac{(aC + bB) \cot(c + dx)}{d} - \frac{(aB - bC) \log(-\sin(c + dx))}{d} - x(aC + bB) - \frac{aB \cot^2(c + dx)}{2d}$$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]`

output `-((b*B + a*C)*x) - ((b*B + a*C)*Cot[c + d*x])/d - (a*B*Cot[c + d*x]^2)/(2*d) - ((a*B - b*C)*Log[-Sin[c + d*x]])/d`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.17

method	result
derivativedivides	$\frac{Bb(-\cot(dx+c)-dx-c)+Cb \ln(\sin(dx+c))+Ba\left(-\frac{\cot(dx+c)^2}{2}-\ln(\sin(dx+c))\right)+Ca(-\cot(dx+c)-dx-c)}{d}$
default	$\frac{Bb(-\cot(dx+c)-dx-c)+Cb \ln(\sin(dx+c))+Ba\left(-\frac{\cot(dx+c)^2}{2}-\ln(\sin(dx+c))\right)+Ca(-\cot(dx+c)-dx-c)}{d}$
parallelrisc	$-\frac{Ba\left(-\ln(\sec(dx+c)^2)+2\ln(\tan(dx+c))\right)-Cb\left(-\ln(\sec(dx+c)^2)+2\ln(\tan(dx+c))\right)+2Bbdx+2Cadx+2Bb \cot(dx+c)}{2d}$
norman	$\frac{(-Bb-Ca)x \tan(dx+c)^3 - \frac{(Bb+Ca) \tan(dx+c)^2}{d} - \frac{Ba \tan(dx+c)}{2d}}{\tan(dx+c)^3} - \frac{(Ba-Cb) \ln(\tan(dx+c))}{d} + \frac{(Ba-Cb) \ln(1+\tan(dx+c))}{2d}$
risc	$-Bbx - Cax + iBax - iCbx + \frac{2iBac}{d} - \frac{2iCbc}{d} - \frac{2i(iBa e^{2i(dx+c)} + Bb e^{2i(dx+c)} + Ca e^{2i(dx+c)} - Bb)}{d(e^{2i(dx+c)} - 1)^2}$

input

```
int(cot(d*x+c)^4*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_
RETURNVERBOSE)
```

output

```
1/d*(B*b*(-cot(d*x+c)-d*x-c)+C*b*ln(sin(d*x+c))+B*a*(-1/2*cot(d*x+c)^2-ln(
sin(d*x+c)))+C*a*(-cot(d*x+c)-d*x-c))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.44

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{(Ba - Cb) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 + (2(Ca + Bb)dx + Ba) \tan(dx+c)^2 + Ba + 2(Ca + Bb)dx}{2d \tan(dx+c)^2}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")`

output `-1/2*((B*a - C*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^2 + (2*(C*a + B*b)*d*x + B*a)*tan(d*x + c)^2 + B*a + 2*(C*a + B*b)*tan(d*x + c))/(d*tan(d*x + c)^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(56) = 112.

Time = 0.99 (sec) , antiderivative size = 143, normalized size of antiderivative = 2.17

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c)) (B \tan(c) + C \tan^2(c)) \cot^4(c) \\ \text{NaN} \\ \frac{Ba \log(\tan^2(c+dx)+1)}{2d} - \frac{Ba \log(\tan(c+dx))}{d} - \frac{Ba}{2d \tan^2(c+dx)} - Bbx - \frac{Bb}{d \tan(c+dx)} - Cax - \frac{Ca}{d \tan(c+dx)} - \frac{Cb \log(\tan^2(c+dx)+1)}{2d} \end{cases}$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*cot(c)**4, Eq(d, 0)), (nan, Eq(c, -d*x)), (B*a*log(tan(c + d*x)**2 + 1)/(2*d) - B*a*log(tan(c + d*x))/d - B*a/(2*d*tan(c + d*x)**2) - B*b*x - B*b/(d*tan(c + d*x)) - C*a*x - C*a/(d*tan(c + d*x)) - C*b*log(tan(c + d*x)**2 + 1)/(2*d) + C*b*log(tan(c + d*x))/d, True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.30

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{2(Ca + Bb)(dx + c) - (Ba - Cb) \log(\tan(dx + c)^2 + 1) + 2(Ba - Cb) \log(\tan(dx + c)) + \frac{Ba + 2(Ca + Bb) \tan(dx + c)}{\tan(dx + c)^2}}{2d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")`

output `-1/2*(2*(C*a + B*b)*(d*x + c) - (B*a - C*b)*log(tan(d*x + c)^2 + 1) + 2*(B*a - C*b)*log(tan(d*x + c)) + (B*a + 2*(C*a + B*b)*tan(d*x + c))/tan(d*x + c)^2)/d`

Giac [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.44

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ca + Bb)(dx + c)}{d} + \frac{(Ba - Cb) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$- \frac{(Ba - Cb) \log(|\tan(dx + c)|)}{d} - \frac{Ba + 2(Ca + Bb) \tan(dx + c)}{2d \tan(dx + c)^2}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")`

output `-(C*a + B*b)*(d*x + c)/d + 1/2*(B*a - C*b)*log(tan(d*x + c)^2 + 1)/d - (B*a - C*b)*log(abs(tan(d*x + c)))/d - 1/2*(B*a + 2*(C*a + B*b)*tan(d*x + c))/(d*tan(d*x + c)^2)`

Mupad [B] (verification not implemented)

Time = 5.21 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.64

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{\ln(\tan(c + dx)) (B a - C b)}{d} - \frac{\cot(c + dx)^2 \left(\frac{B a}{2} + \tan(c + dx) (B b + C a)\right)}{d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (B + C i) (a + b i)}{2d}$$

$$- \frac{\ln(\tan(c + dx) + i) (B - C i) (b + a i) i}{2d}$$

input

```
int(cot(c + d*x)^4*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
),x)
```

output

```
(log(tan(c + d*x) - i)*(B + C*i)*(a + b*i))/(2*d) - (cot(c + d*x)^2*((B
*a)/2 + tan(c + d*x)*(B*b + C*a)))/d - (log(tan(c + d*x))*(B*a - C*b))/d -
(log(tan(c + d*x) + i)*(B - C*i)*(a*i + b)*i)/(2*d)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.83

$$\int \cot^4(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-4 \cos(dx + c) \sin(dx + c) a c - 4 \cos(dx + c) \sin(dx + c) b^2 + 4 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) \sin(dx + c)^2 a}{d}$$

input

```
int(cot(d*x+c)^4*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
( - 4*cos(c + d*x)*sin(c + d*x)*a*c - 4*cos(c + d*x)*sin(c + d*x)*b**2 + 4
*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a*b - 4*log(tan((c + d*x)/2)
**2 + 1)*sin(c + d*x)**2*b*c - 4*log(tan((c + d*x)/2))*sin(c + d*x)**2*a*b
+ 4*log(tan((c + d*x)/2))*sin(c + d*x)**2*b*c + sin(c + d*x)**2*a*b - 4*s
in(c + d*x)**2*a*c*d*x - 4*sin(c + d*x)**2*b**2*d*x - 2*a*b)/(4*sin(c + d
*x)**2*d)
```

3.7 $\int \cot^5(c+dx)(a+b \tan(c+dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	133
Mathematica [C] (verified)	134
Rubi [A] (verified)	134
Maple [A] (verified)	138
Fricas [A] (verification not implemented)	139
Sympy [B] (verification not implemented)	139
Maxima [A] (verification not implemented)	140
Giac [A] (verification not implemented)	140
Mupad [B] (verification not implemented)	141
Reduce [B] (verification not implemented)	142

Optimal result

Integrand size = 38, antiderivative size = 87

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (aB - bC)x + \frac{(aB - bC) \cot(c + dx)}{d} - \frac{(bB + aC) \cot^2(c + dx)}{2d}$$

$$- \frac{aB \cot^3(c + dx)}{3d} - \frac{(bB + aC) \log(\sin(c + dx))}{d}$$

output

```
(B*a-C*b)*x+(B*a-C*b)*cot(d*x+c)/d-1/2*(B*b+C*a)*cot(d*x+c)^2/d-1/3*a*B*cot(d*x+c)^3/d-(B*b+C*a)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.04 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.48

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{bB \csc^2(c + dx)}{2d} - \frac{aC \csc^2(c + dx)}{2d}$$

$$- \frac{aB \cot^3(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c + dx)\right)}{3d}$$

$$- \frac{bC \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c + dx)\right)}{d}$$

$$- \frac{bB \log(\sin(c + dx))}{d} - \frac{aC \log(\sin(c + dx))}{d}$$

input

```
Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
-1/2*(b*B*Csc[c + d*x]^2)/d - (a*C*Csc[c + d*x]^2)/(2*d) - (a*B*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2])/(3*d) - (b*C*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d - (b*B*Log[Sin[c + d*x]])/d - (a*C*Log[Sin[c + d*x]])/d
```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4115, 3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))(B \tan(c + dx) + C \tan(c + dx)^2)}{\tan(c + dx)^5} dx \\
& \quad \downarrow 4115 \\
& \int \cot^4(c + dx)(a + b \tan(c + dx))(B + C \tan(c + dx)) dx \\
& \quad \downarrow 3042 \\
& \int \frac{(a + b \tan(c + dx))(B + C \tan(c + dx))}{\tan(c + dx)^4} dx \\
& \quad \downarrow 4074 \\
& \int \cot^3(c + dx)(bB + aC - (aB - bC) \tan(c + dx)) dx - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& \int \frac{bB + aC - (aB - bC) \tan(c + dx)}{\tan(c + dx)^3} dx - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4012 \\
& \int -\cot^2(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 25 \\
& - \int \cot^2(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& - \int \frac{aB - bC + (bB + aC) \tan(c + dx)}{\tan(c + dx)^2} dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4012 \\
& - \int \cot(c + dx)(bB + aC - (aB - bC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} + \frac{(aB - bC) \cot(c + dx)}{d} - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& - \int \frac{bB + aC - (aB - bC) \tan(c + dx)}{\tan(c + dx)} dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} + \\
& \quad \frac{(aB - bC) \cot(c + dx)}{d} - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4014 \\
& -(aC + bB) \int \cot(c + dx) dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} + \frac{(aB - bC) \cot(c + dx)}{d} + x(aB - \\
& \quad bC) - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& -(aC + bB) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} + \frac{(aB - bC) \cot(c + dx)}{d} + \\
& \quad x(aB - bC) - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 25 \\
& (aC + bB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(aC + bB) \cot^2(c + dx)}{2d} + \frac{(aB - bC) \cot(c + dx)}{d} + \\
& \quad x(aB - bC) - \frac{aB \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3956 \\
& -\frac{(aC + bB) \cot^2(c + dx)}{2d} + \frac{(aB - bC) \cot(c + dx)}{d} - \frac{(aC + bB) \log(-\sin(c + dx))}{d} + x(aB - \\
& \quad bC) - \frac{aB \cot^3(c + dx)}{3d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(a*B - b*C)*x + ((a*B - b*C)*Cot[c + d*x])/d - ((b*B + a*C)*Cot[c + d*x]^2)/(2*d) - (a*B*Cot[c + d*x]^3)/(3*d) - ((b*B + a*C)*Log[-Sin[c + d*x]])/d
```

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinear} \\ \text{Q}[\text{u}, \text{x}]$
- rule 3956 $\text{Int}[\tan[(\text{c}_.) + (\text{d}_.) * (\text{x}_)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d} \\ * \text{x}], \text{x}]]/\text{d}, \text{x}] \text{ /; FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4012 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] \text{ } ^{\text{m}_}) * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + \\ (\text{f}_.) * (\text{x}_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{b} * \text{c} - \text{a} * \text{d}) * ((\text{a} + \text{b} * \tan[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} / \\ (\text{f} * (\text{m} + 1) * (\text{a}^2 + \text{b}^2))), \text{x}] + \text{Simp}[1 / (\text{a}^2 + \text{b}^2) \quad \text{Int}[(\text{a} + \text{b} * \tan[\text{e} + \text{f} * \text{x}]) \\ ^{\text{m} + 1} * \text{Simp}[\text{a} * \text{c} + \text{b} * \text{d} - (\text{b} * \text{c} - \text{a} * \text{d}) * \tan[\text{e} + \text{f} * \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a} \\ , \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{LtQ}[\text{m}, -1 \\]$
- rule 4014 $\text{Int}[(\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] / ((\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) \\ * (\text{x}_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{a} * \text{c} + \text{b} * \text{d}) * (\text{x} / (\text{a}^2 + \text{b}^2)), \text{x}] + \text{Simp}[(\text{b} * \text{c} - \text{a} \\ * \text{d}) / (\text{a}^2 + \text{b}^2) \quad \text{Int}[(\text{b} - \text{a} * \tan[\text{e} + \text{f} * \text{x}]) / (\text{a} + \text{b} * \tan[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] \text{ /; } \\ \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{N} \\ \text{eQ}[\text{a} * \text{c} + \text{b} * \text{d}, 0]$
- rule 4074 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)] \text{ } ^{\text{m}_}) * ((\text{A}_.) + (\text{B}_.) * \tan[(\text{e}_.) + \\ (\text{f}_.) * (\text{x}_)]) * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{b} \\ * \text{c} - \text{a} * \text{d}) * (\text{A} * \text{b} - \text{a} * \text{B}) * ((\text{a} + \text{b} * \tan[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} / (\text{b} * \text{f} * (\text{m} + 1) * (\text{a}^2 + \text{b}^2 \\))), \text{x}] + \text{Simp}[1 / (\text{a}^2 + \text{b}^2) \quad \text{Int}[(\text{a} + \text{b} * \tan[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} * \text{Simp}[\text{a} * \text{A} * \text{c} \\ + \text{b} * \text{B} * \text{c} + \text{A} * \text{b} * \text{d} - \text{a} * \text{B} * \text{d} - (\text{A} * \text{b} * \text{c} - \text{a} * \text{B} * \text{c} - \text{a} * \text{A} * \text{d} - \text{b} * \text{B} * \text{d}) * \tan[\text{e} + \text{f} * \text{x}], \text{x}], \\ \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{LtQ}[\text{m} \\ , -1] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0]$

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{Bb\left(-\frac{\cot(dx+c)^2}{2}-\ln(\sin(dx+c))\right)+Cb(-\cot(dx+c)-dx-c)+Ba\left(-\frac{\cot(dx+c)^3}{3}+\cot(dx+c)+dx+c\right)+Ca\left(-\frac{\cot(dx+c)^2}{2}\right)}{d}$
default	$\frac{Bb\left(-\frac{\cot(dx+c)^2}{2}-\ln(\sin(dx+c))\right)+Cb(-\cot(dx+c)-dx-c)+Ba\left(-\frac{\cot(dx+c)^3}{3}+\cot(dx+c)+dx+c\right)+Ca\left(-\frac{\cot(dx+c)^2}{2}\right)}{d}$
norman	$\frac{\frac{(Ba-Cb)\tan(dx+c)^3}{d}+(Ba-Cb)x\tan(dx+c)^4-\frac{(Bb+Ca)\tan(dx+c)^2}{2d}-\frac{Ba\tan(dx+c)}{3d}}{\tan(dx+c)^4}-\frac{(Bb+Ca)\ln(\tan(dx+c))}{d}+\frac{Ca}{d}$
parallelrisch	$\frac{-3Bb\left(-\ln(\sec(dx+c)^2)+2\ln(\tan(dx+c))\right)-3Ca\left(-\ln(\sec(dx+c)^2)+2\ln(\tan(dx+c))\right)-2B\cot(dx+c)^3a-3Bb\cot(dx+c)}{6d}$
risch	$iBbx + iCax + Bax - Cbx + \frac{2iBbc}{d} + \frac{2iCac}{d} - \frac{2i(3iBbe^{4i(dx+c)}+3iCa e^{4i(dx+c)}-6Ba e^{4i(dx+c)}+3Bb)}{6d}$

input

```
int(cot(d*x+c)^5*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)
```

output

```
1/d*(B*b*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+C*b*(-cot(d*x+c)-d*x-c)+B*a*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c)+C*a*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c))))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.39

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{3(Ca + Bb) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 - 3(2(Ba - Cb)dx - Ca - Bb) \tan(dx+c)^3 - 6(Ba - Cb) \tan(dx+c)^2 + 3Ca + 3Bb}{6d \tan(dx+c)^3}$$

input

```
integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")
```

output

```
-1/6*(3*(C*a + B*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^3 - 3*(2*(B*a - C*b)*d*x - C*a - B*b)*tan(d*x + c)^3 - 6*(B*a - C*b)*tan(d*x + c)^2 + 2*B*a + 3*(C*a + B*b)*tan(d*x + c))/(d*tan(d*x + c)^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. 2(75) = 150.

Time = 1.95 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.99

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c)) (B \tan(c) + C \tan^2(c)) \cot^5(c) \\ \text{NaN} \\ Bax + \frac{Ba}{d \tan(c+dx)} - \frac{Ba}{3d \tan^3(c+dx)} + \frac{Bb \log(\tan^2(c+dx)+1)}{2d} - \frac{Bb \log(\tan(c+dx))}{d} - \frac{Bb}{2d \tan^2(c+dx)} + \frac{Ca \log(\tan^2(c+dx))}{2d} \end{cases}$$

input

```
integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```


output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*cot(c)**5, Eq(d, 0)), (nan, Eq(c, -d*x)), (B*a*x + B*a/(d*tan(c + d*x)) - B*a/(3*d*tan(c + d*x)**3) + B*b*log(tan(c + d*x)**2 + 1)/(2*d) - B*b*log(tan(c + d*x))/d - B*b/(2*d*tan(c + d*x)**2) + C*a*log(tan(c + d*x)**2 + 1)/(2*d) - C*a*log(tan(c + d*x))/d - C*a/(2*d*tan(c + d*x)**2) - C*b*x - C*b/(d*tan(c + d*x)), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.20

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6(Ba - Cb)(dx + c) + 3(Ca + Bb) \log(\tan(dx + c)^2 + 1) - 6(Ca + Bb) \log(\tan(dx + c)) + \frac{6(Ba - Cb)}{6d}}{6d}$$

input

```
integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
1/6*(6*(B*a - C*b)*(d*x + c) + 3*(C*a + B*b)*log(tan(d*x + c)^2 + 1) - 6*(C*a + B*b)*log(tan(d*x + c)) + (6*(B*a - C*b)*tan(d*x + c)^2 - 2*B*a - 3*(C*a + B*b)*tan(d*x + c))/tan(d*x + c)^3)/d
```

Giac [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.29

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ba - Cb)(dx + c)}{d} + \frac{(Ca + Bb) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$- \frac{(Ca + Bb) \log(|\tan(dx + c)|)}{d}$$

$$+ \frac{6(Ba - Cb) \tan(dx + c)^2 - 2Ba - 3(Ca + Bb) \tan(dx + c)}{6d \tan(dx + c)^3}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")`

output $(B*a - C*b)*(d*x + c)/d + 1/2*(C*a + B*b)*\log(\tan(d*x + c)^2 + 1)/d - (C*a + B*b)*\log(\text{abs}(\tan(d*x + c)))/d + 1/6*(6*(B*a - C*b)*\tan(d*x + c)^2 - 2*B*a - 3*(C*a + B*b)*\tan(d*x + c))/(d*\tan(d*x + c)^3)$

Mupad [B] (verification not implemented)

Time = 5.24 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.46

$$\int \cot^5(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{\cot(c + dx)^3 ((C b - B a) \tan(c + dx)^2 + (\frac{B b}{2} + \frac{C a}{2}) \tan(c + dx) + \frac{B a}{3})}{d}$$

$$- \frac{\ln(\tan(c + dx)) (B b + C a)}{d} - \frac{\ln(\tan(c + dx) - i) (B + C i) (a + b i) i}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (B - C i) (b + a i)}{2 d}$$

input `int(cot(c + d*x)^5*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)),x)`

output $(\log(\tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b))/(2*d) - (\log(\tan(c + d*x))*(B*b + C*a))/d - (\log(\tan(c + d*x) - 1i)*(B + C*1i)*(a + b*1i)*1i)/(2*d) - (\cot(c + d*x)^3*((B*a)/3 + \tan(c + d*x)*((B*b)/2 + (C*a)/2) - \tan(c + d*x)^2*(B*a - C*b))/d$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 232, normalized size of antiderivative = 2.67

$$\int \cot^5(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{16 \cos(dx + c) \sin(dx + c)^2 ab - 12 \cos(dx + c) \sin(dx + c)^2 bc - 4 \cos(dx + c) ab + 12 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\dots}$$

input

```
int(cot(d*x+c)^5*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
(16*cos(c + d*x)*sin(c + d*x)**2*a*b - 12*cos(c + d*x)*sin(c + d*x)**2*b*c
- 4*cos(c + d*x)*a*b + 12*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**3*a*
c + 12*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**3*b**2 - 12*log(tan((c +
d*x)/2))*sin(c + d*x)**3*a*c - 12*log(tan((c + d*x)/2))*sin(c + d*x)**3*b
**2 + 12*sin(c + d*x)**3*a*b*d*x + 3*sin(c + d*x)**3*a*c + 3*sin(c + d*x)*
*3*b**2 - 12*sin(c + d*x)**3*b*c*d*x - 6*sin(c + d*x)*a*c - 6*sin(c + d*x)
*b**2)/(12*sin(c + d*x)**3*d)
```

3.8 $\int \cot^6(c+dx)(a+b \tan(c+dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	143
Mathematica [C] (verified)	144
Rubi [A] (verified)	144
Maple [A] (verified)	148
Fricas [A] (verification not implemented)	149
Sympy [B] (verification not implemented)	149
Maxima [A] (verification not implemented)	150
Giac [A] (verification not implemented)	150
Mupad [B] (verification not implemented)	151
Reduce [B] (verification not implemented)	152

Optimal result

Integrand size = 38, antiderivative size = 108

$$\int \cot^6(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (bB + aC)x + \frac{(bB + aC) \cot(c + dx)}{d} + \frac{(aB - bC) \cot^2(c + dx)}{2d}$$

$$- \frac{(bB + aC) \cot^3(c + dx)}{3d} - \frac{aB \cot^4(c + dx)}{4d} + \frac{(aB - bC) \log(\sin(c + dx))}{d}$$

output

```
(B*b+C*a)*x+(B*b+C*a)*cot(d*x+c)/d+1/2*(B*a-C*b)*cot(d*x+c)^2/d-1/3*(B*b+C
*a)*cot(d*x+c)^3/d-1/4*a*B*cot(d*x+c)^4/d+(B*a-C*b)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.04 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.35

$$\begin{aligned} & \int \cot^6(c+dx)(a+b\tan(c+dx))(B\tan(c+dx)+C\tan^2(c+dx)) dx \\ &= \frac{aB \csc^2(c+dx)}{d} - \frac{bC \csc^2(c+dx)}{2d} - \frac{aB \csc^4(c+dx)}{4d} \\ & \quad - \frac{bB \cot^3(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)\right)}{3d} \\ & \quad - \frac{aC \cot^3(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)\right)}{3d} \\ & \quad + \frac{aB \log(\sin(c+dx))}{d} - \frac{bC \log(\sin(c+dx))}{d} \end{aligned}$$

input

```
Integrate[Cot[c + d*x]^6*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(a*B*Csc[c + d*x]^2)/d - (b*C*Csc[c + d*x]^2)/(2*d) - (a*B*Csc[c + d*x]^4)/(4*d) - (b*B*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2])/(3*d) - (a*C*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2])/(3*d) + (a*B*Log[Sin[c + d*x]])/d - (b*C*Log[Sin[c + d*x]])/d
```

Rubi [A] (verified)

Time = 1.50 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.02, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4115, 3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c+dx)(a+b\tan(c+dx))(B\tan(c+dx)+C\tan^2(c+dx)) dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))(B \tan(c + dx) + C \tan(c + dx)^2)}{\tan(c + dx)^6} dx \\
& \quad \downarrow 4115 \\
& \int \cot^5(c + dx)(a + b \tan(c + dx))(B + C \tan(c + dx)) dx \\
& \quad \downarrow 3042 \\
& \int \frac{(a + b \tan(c + dx))(B + C \tan(c + dx))}{\tan(c + dx)^5} dx \\
& \quad \downarrow 4074 \\
& \int \cot^4(c + dx)(bB + aC - (aB - bC) \tan(c + dx)) dx - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 3042 \\
& \int \frac{bB + aC - (aB - bC) \tan(c + dx)}{\tan(c + dx)^4} dx - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 4012 \\
& \int -\cot^3(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 25 \\
& - \int \cot^3(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 3042 \\
& - \int \frac{aB - bC + (bB + aC) \tan(c + dx)}{\tan(c + dx)^3} dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 4012 \\
& - \int \cot^2(c + dx)(bB + aC - (aB - bC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \frac{(aB - bC) \cot^2(c + dx)}{2d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& - \int \frac{bB + aC - (aB - bC) \tan(c + dx)}{\tan(c + dx)^2} dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \\
& \quad \frac{(aB - bC) \cot^2(c + dx)}{2d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 4012 \\
& - \int -\cot(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \\
& \quad \frac{(aB - bC) \cot^2(c + dx)}{2d} + \frac{(aC + bB) \cot(c + dx)}{d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 25 \\
& \int \cot(c + dx)(aB - bC + (bB + aC) \tan(c + dx)) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \\
& \quad \frac{(aB - bC) \cot^2(c + dx)}{2d} + \frac{(aC + bB) \cot(c + dx)}{d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 3042 \\
& \int \frac{aB - bC + (bB + aC) \tan(c + dx)}{\tan(c + dx)} dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \\
& \quad \frac{(aB - bC) \cot^2(c + dx)}{2d} + \frac{(aC + bB) \cot(c + dx)}{d} - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 4014 \\
& (aB - bC) \int \cot(c + dx) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \frac{(aB - bC) \cot^2(c + dx)}{2d} + \\
& \quad \frac{(aC + bB) \cot(c + dx)}{d} + x(aC + bB) - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 3042 \\
& (aB - bC) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \frac{(aB - bC) \cot^2(c + dx)}{2d} + \\
& \quad \frac{(aC + bB) \cot(c + dx)}{d} + x(aC + bB) - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 25 \\
& -(aB - bC) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(aC + bB) \cot^3(c + dx)}{3d} + \\
& \quad \frac{(aB - bC) \cot^2(c + dx)}{2d} + \frac{(aC + bB) \cot(c + dx)}{d} + x(aC + bB) - \frac{aB \cot^4(c + dx)}{4d} \\
& \quad \downarrow 3956
\end{aligned}$$

$$-\frac{(aC + bB) \cot^3(c + dx)}{3d} + \frac{(aB - bC) \cot^2(c + dx)}{2d} + \frac{(aC + bB) \cot(c + dx)}{d} + \frac{(aB - bC) \log(-\sin(c + dx))}{d} + x(aC + bB) - \frac{aB \cot^4(c + dx)}{4d}$$

input

```
Int[Cot[c + d*x]^6*(a + b*Tan[c + d*x])*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(b*B + a*C)*x + ((b*B + a*C)*Cot[c + d*x])/d + ((a*B - b*C)*Cot[c + d*x]^2)/(2*d) - ((b*B + a*C)*Cot[c + d*x]^3)/(3*d) - (a*B*Cot[c + d*x]^4)/(4*d) + ((a*B - b*C)*Log[-Sin[c + d*x]])/d
```

Definitions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4012

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]
```

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]
```


rule 4074

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{Bb\left(-\frac{\cot(dx+c)^3}{3} + \cot(dx+c) + dx+c\right) + Cb\left(-\frac{\cot(dx+c)^2}{2} - \ln(\sin(dx+c))\right) + Ba\left(-\frac{\cot(dx+c)^4}{4} + \frac{\cot(dx+c)^2}{2} + \ln(\sin(dx+c))\right)}{d}$
default	$\frac{Bb\left(-\frac{\cot(dx+c)^3}{3} + \cot(dx+c) + dx+c\right) + Cb\left(-\frac{\cot(dx+c)^2}{2} - \ln(\sin(dx+c))\right) + Ba\left(-\frac{\cot(dx+c)^4}{4} + \frac{\cot(dx+c)^2}{2} + \ln(\sin(dx+c))\right)}{d}$
norman	$\frac{\frac{(Bb+Ca)\tan(dx+c)^4}{d} + (Bb+Ca)x\tan(dx+c)^5 - \frac{(Bb+Ca)\tan(dx+c)^2}{3d} + \frac{(Ba-Cb)\tan(dx+c)^3}{2d} - \frac{Ba\tan(dx+c)}{4d}}{\tan(dx+c)^5} + \frac{(Ba-Cb)}{d}$
parallelrisch	$-\frac{-6Ba\left(-\ln(\sec(dx+c)^2) + 2\ln(\tan(dx+c))\right) + 6Cb\left(-\ln(\sec(dx+c)^2) + 2\ln(\tan(dx+c))\right) + 3B\cot(dx+c)^4 a + 4Bb}{d}$
risch	$Bbx + Cax - iBax + iCbx - \frac{2iBac}{d} + \frac{2iCbc}{d} - \frac{2(-6iBbe^{6i(dx+c)} - 6iCa e^{6i(dx+c)} + 6Ba e^{6i(dx+c)} - \dots)}{d}$

input

```
int(cot(d*x+c)^6*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_
RETURNVERBOSE)
```

output

```
1/d*(B*b*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c)+C*b*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+B*a*(-1/4*cot(d*x+c)^4+1/2*cot(d*x+c)^2+ln(sin(d*x+c)))+C*a*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.28

$$\int \cot^6(c+dx)(a+b \tan(c+dx))(B \tan(c+dx)+C \tan^2(c+dx)) dx$$

$$= \frac{6(Ba - Cb) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 + 3(4(Ca + Bb)dx + 3Ba - 2Cb) \tan(dx+c)^4 + 12(Ca + Bb) \tan(dx+c)^4}{12d \tan(dx+c)^4}$$

input

```
integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")
```

output

```
1/12*(6*(B*a - C*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^4 + 3*(4*(C*a + B*b)*d*x + 3*B*a - 2*C*b)*tan(d*x + c)^4 + 12*(C*a + B*b)*tan(d*x + c)^3 + 6*(B*a - C*b)*tan(d*x + c)^2 - 3*B*a - 4*(C*a + B*b)*tan(d*x + c))/(d*tan(d*x + c)^4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(95) = 190.

Time = 2.53 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.89

$$\int \cot^6(c+dx)(a+b \tan(c+dx))(B \tan(c+dx)+C \tan^2(c+dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))(B \tan(c) + C \tan^2(c)) \cot^6(c) \\ \text{NaN} \end{cases}$$

$$= -\frac{Ba \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba \log(\tan(c+dx))}{d} + \frac{Ba}{2d \tan^2(c+dx)} - \frac{Ba}{4d \tan^4(c+dx)} + Bbx + \frac{Bb}{d \tan(c+dx)} - \frac{Bb}{3d \tan^3(c+dx)} + \dots$$

input

```
integrate(cot(d*x+c)**6*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))*(B*tan(c) + C*tan(c)**2)*cot(c)**6, Eq(d, 0)), (nan, Eq(c, -d*x)), (-B*a*log(tan(c + d*x)**2 + 1)/(2*d) + B*a*log(tan(c + d*x))/d + B*a/(2*d*tan(c + d*x)**2) - B*a/(4*d*tan(c + d*x)**4) + B*b*x + B*b/(d*tan(c + d*x)) - B*b/(3*d*tan(c + d*x)**3) + C*a*x + C*a/(d*tan(c + d*x)) - C*a/(3*d*tan(c + d*x)**3) + C*b*log(tan(c + d*x)**2 + 1)/(2*d) - C*b*log(tan(c + d*x))/d - C*b/(2*d*tan(c + d*x)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.13

$$\int \cot^6(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{12(Ca + Bb)(dx + c) - 6(Ba - Cb) \log(\tan(dx + c)^2 + 1) + 12(Ba - Cb) \log(\tan(dx + c)) + \frac{12(Ca + Bb)}{12d}}{12d}$$

input

```
integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
1/12*(12*(C*a + B*b)*(d*x + c) - 6*(B*a - C*b)*log(tan(d*x + c)^2 + 1) + 12*(B*a - C*b)*log(tan(d*x + c)) + (12*(C*a + B*b)*tan(d*x + c)^3 + 6*(B*a - C*b)*tan(d*x + c)^2 - 3*B*a - 4*(C*a + B*b)*tan(d*x + c))/tan(d*x + c)^4)/d
```

Giac [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.19

$$\int \cot^6(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ca + Bb)(dx + c)}{d} - \frac{(Ba - Cb) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{(Ba - Cb) \log(|\tan(dx + c)|)}{d}$$

$$+ \frac{12(Ca + Bb) \tan(dx + c)^3 + 6(Ba - Cb) \tan(dx + c)^2 - 3Ba - 4(Ca + Bb) \tan(dx + c)}{12d \tan(dx + c)^4}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")`

output `(C*a + B*b)*(d*x + c)/d - 1/2*(B*a - C*b)*log(tan(d*x + c)^2 + 1)/d + (B*a - C*b)*log(abs(tan(d*x + c)))/d + 1/12*(12*(C*a + B*b)*tan(d*x + c)^3 + 6*(B*a - C*b)*tan(d*x + c)^2 - 3*B*a - 4*(C*a + B*b)*tan(d*x + c))/(d*tan(d*x + c)^4)`

Mupad [B] (verification not implemented)

Time = 5.30 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.34

$$\int \cot^6(c + dx)(a + b \tan(c + dx))(B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx))(B a - C b)}{d}$$

$$- \frac{\cot(c + dx)^4 \left((-B b - C a) \tan(c + dx)^3 + \left(\frac{C b}{2} - \frac{B a}{2} \right) \tan(c + dx)^2 + \left(\frac{B b}{3} + \frac{C a}{3} \right) \tan(c + dx) + \frac{B}{4} \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) - i)(B + C i)(a + b i)}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) + i)(B - C i)(b + a i) i}{2 d}$$

input `int(cot(c + d*x)^6*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x))*(B*a - C*b))/d - (cot(c + d*x)^4*((B*a)/4 + tan(c + d*x))*((B*b)/3 + (C*a)/3) - tan(c + d*x)^3*(B*b + C*a) - tan(c + d*x)^2*((B*a)/2 - (C*b)/2))/d - (log(tan(c + d*x) - 1i)*(B + C*1i)*(a + b*1i))/(2*d) + (log(tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b)*1i)/(2*d)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 261, normalized size of antiderivative = 2.42

$$\int \cot^6(c + dx)(a + b \tan(c + dx)) (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{128 \cos(dx + c) \sin(dx + c)^3 ac + 128 \cos(dx + c) \sin(dx + c)^3 b^2 - 32 \cos(dx + c) \sin(dx + c) ac - 32}{}$$

input

```
int(cot(d*x+c)^6*(a+b*tan(d*x+c))*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
(128*cos(c + d*x)*sin(c + d*x)**3*a*c + 128*cos(c + d*x)*sin(c + d*x)**3*b
**2 - 32*cos(c + d*x)*sin(c + d*x)*a*c - 32*cos(c + d*x)*sin(c + d*x)*b**2
- 96*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**4*a*b + 96*log(tan((c + d
*x)/2)**2 + 1)*sin(c + d*x)**4*b*c + 96*log(tan((c + d*x)/2))*sin(c + d*x)
**4*a*b - 96*log(tan((c + d*x)/2))*sin(c + d*x)**4*b*c - 39*sin(c + d*x)**
4*a*b + 96*sin(c + d*x)**4*a*c*d*x + 96*sin(c + d*x)**4*b**2*d*x + 24*sin(
c + d*x)**4*b*c + 96*sin(c + d*x)**2*a*b - 48*sin(c + d*x)**2*b*c - 24*a*b
)/(96*sin(c + d*x)**4*d)
```

3.9 $\int \tan(c+dx)(a+b \tan(c+dx))^2 (B \tan(c + dx) + C \tan$

Optimal result	153
Mathematica [C] (verified)	154
Rubi [A] (verified)	154
Maple [A] (warning: unable to verify)	158
Fricas [A] (verification not implemented)	159
Sympy [A] (verification not implemented)	159
Maxima [A] (verification not implemented)	160
Giac [A] (verification not implemented)	160
Mupad [B] (verification not implemented)	161
Reduce [B] (verification not implemented)	161

Optimal result

Integrand size = 38, antiderivative size = 148

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((a^2B - b^2B - 2abC) x) + \frac{(2abB + a^2C - b^2C) \log(\cos(c + dx))}{d}$$

$$- \frac{b(bB + aC) \tan(c + dx)}{d} - \frac{C(a + b \tan(c + dx))^2}{d}$$

$$+ \frac{(4bB - aC)(a + b \tan(c + dx))^3}{12b^2d} + \frac{C \tan(c + dx)(a + b \tan(c + dx))^3}{4bd}$$

output

```
-(B*a^2-B*b^2-2*C*a*b)*x+(2*B*a*b+C*a^2-C*b^2)*ln(cos(d*x+c))/d-b*(B*b+C*a
)*tan(d*x+c)/d-1/2*C*(a+b*tan(d*x+c))^2/d+1/12*(4*B*b-C*a)*(a+b*tan(d*x+c)
)^3/b^2/d+1/4*C*tan(d*x+c)*(a+b*tan(d*x+c))^3/b/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.18 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.49

$$\int \tan(c+dx)(a+b\tan(c+dx))^2 (B\tan(c+dx)+C\tan^2(c+dx)) dx$$

$$= \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} + \frac{(4bB-aC)(a+b\tan(c+dx))^3}{3bd} + \frac{2((bB-aC)(i(a+ib)^2 \log(i-\tan(c+dx))-i(a-ib)^2 \log(i+\tan(c+dx))-2b^2 \tan(c+dx))-C((ia-b)^3 \log(i-\tan(c+dx))))}{4b}$$

input

```
Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(C*Tan[c + d*x]*(a + b*Tan[c + d*x])^3)/(4*b*d) + (((4*b*B - a*C)*(a + b*Tan[c + d*x])^3)/(3*b*d) + (2*((b*B - a*C)*(I*(a + I*b)^2*Log[I - Tan[c + d*x]] - I*(a - I*b)^2*Log[I + Tan[c + d*x]] - 2*b^2*Tan[c + d*x]) - C*((I*a - b)^3*Log[I - Tan[c + d*x]] - (I*a + b)^3*Log[I + Tan[c + d*x]] + 6*a*b^2*Tan[c + d*x] + b^3*Tan[c + d*x]^2)))/d)/(4*b)
```

Rubi [A] (verified)

Time = 1.54 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.342$, Rules used = {3042, 4115, 3042, 4090, 25, 3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan(c+dx)(a+b\tan(c+dx))^2 (B\tan(c+dx)+C\tan^2(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)(a+b\tan(c+dx))^2 (B\tan(c+dx)+C\tan(c+dx)^2) dx$$

$$\downarrow \text{4115}$$

$$\begin{aligned}
& \int \tan^2(c+dx)(a+b\tan(c+dx))^2(B+C\tan(c+dx))dx \\
& \quad \downarrow \text{3042} \\
& \int \tan(c+dx)^2(a+b\tan(c+dx))^2(B+C\tan(c+dx))dx \\
& \quad \downarrow \text{4090} \\
& \frac{\int -(a+b\tan(c+dx))^2(-((4bB-aC)\tan^2(c+dx))+4bC\tan(c+dx)+aC)dx}{4b} + \\
& \quad \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} \\
& \quad \downarrow \text{25} \\
& \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} - \\
& \frac{\int (a+b\tan(c+dx))^2(-((4bB-aC)\tan^2(c+dx))+4bC\tan(c+dx)+aC)dx}{4b} \\
& \quad \downarrow \text{3042} \\
& \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} - \\
& \frac{\int (a+b\tan(c+dx))^2(-((4bB-aC)\tan(c+dx)^2)+4bC\tan(c+dx)+aC)dx}{4b} \\
& \quad \downarrow \text{4113} \\
& \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} - \\
& \frac{\int (a+b\tan(c+dx))^2(4bB+4bC\tan(c+dx))dx - \frac{(4bB-aC)(a+b\tan(c+dx))^3}{3bd}}{4b} \\
& \quad \downarrow \text{3042} \\
& \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} - \\
& \frac{\int (a+b\tan(c+dx))^2(4bB+4bC\tan(c+dx))dx - \frac{(4bB-aC)(a+b\tan(c+dx))^3}{3bd}}{4b} \\
& \quad \downarrow \text{4011} \\
& \frac{C\tan(c+dx)(a+b\tan(c+dx))^3}{4bd} - \\
& \frac{\int (a+b\tan(c+dx))(4b(aB-bC)+4b(bB+aC)\tan(c+dx))dx - \frac{(4bB-aC)(a+b\tan(c+dx))^3}{3bd} + \frac{2bC(a+b\tan(c+dx))^2}{d}}{4b} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{\frac{C \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \int (a+b \tan(c+dx))(4b(aB-bC)+4b(bB+aC) \tan(c+dx))dx - \frac{(4bB-aC)(a+b \tan(c+dx))^3}{3bd} + \frac{2bC(a+b \tan(c+dx))^2}{d}}{4b}$$

↓ 4008

$$\frac{\frac{C \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - 4b(a^2C+2abB-b^2C) \int \tan(c+dx)dx + 4bx(a^2B-2abC-b^2B) + \frac{4b^2(aC+bB) \tan(c+dx)}{d} - \frac{(4bB-aC)(a+b \tan(c+dx))^3}{3bd}}{4b}$$

↓ 3042

$$\frac{\frac{C \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - 4b(a^2C+2abB-b^2C) \int \tan(c+dx)dx + 4bx(a^2B-2abC-b^2B) + \frac{4b^2(aC+bB) \tan(c+dx)}{d} - \frac{(4bB-aC)(a+b \tan(c+dx))^3}{3bd}}{4b}$$

↓ 3956

$$\frac{\frac{C \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{4b(a^2C+2abB-b^2C) \log(\cos(c+dx))}{d} + 4bx(a^2B-2abC-b^2B) + \frac{4b^2(aC+bB) \tan(c+dx)}{d} - \frac{(4bB-aC)(a+b \tan(c+dx))^3}{3bd} + \frac{2b^2C(a+b \tan(c+dx))^2}{d}}{4b}$$

input

```
Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(C*Tan[c + d*x]*(a + b*Tan[c + d*x])^3)/(4*b*d) - (4*b*(a^2*B - b^2*B - 2*a*b*C)*x - (4*b*(2*a*b*B + a^2*C - b^2*C)*Log[Cos[c + d*x]])/d + (4*b^2*(b*B + a*C)*Tan[c + d*x])/d + (2*b*C*(a + b*Tan[c + d*x])^2)/d - ((4*b*B - a*C)*(a + b*Tan[c + d*x])^3)/(3*b*d))/(4*b)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d, x\}$

rule 4008 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_.)]((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[(a*c - b*d)*x, x] + (\text{Simp}[b*d*(\text{Tan}[e + f*x]/f), x] + \text{Simp}[(b*c + a*d) \text{Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[b*c + a*d, 0]$

rule 4011 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 0]$

rule 4090 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)]^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[b*B*(a + b*\text{Tan}[e + f*x])^{(m-1)}*((c + d*\text{Tan}[e + f*x])^{(n+1)}/(d*f*(m+n))), x] + \text{Simp}[1/(d*(m+n)) \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-2)}*(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[a^2*A*d*(m+n) - b*B*(b*c*(m-1) + a*d*(n+1)) + d*(m+n)*(2*a*A*b + B*(a^2 - b^2))*\text{Tan}[e + f*x] - (b*B*(b*c - a*d)*(m-1) - b*(A*b + a*B)*d*(m+n))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, n, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[m, 1] \&\& (\text{IntegerQ}[m] \|\| \text{IntegersQ}[2*m, 2*n]) \&\& !(\text{IGtQ}[n, 1] \&\& (!\text{IntegerQ}[m] \|\| (\text{EqQ}[c, 0] \&\& \text{NeQ}[a, 0]))))$

rule 4113 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)] + (C_.)\tan[(e_.) + (f_.)(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[C*((a + b*\text{Tan}[e + f*x])^{(m+1)}/(b*f*(m+1))), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^m*\text{Simp}[A - C + B*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}\{a, b, e, f, A, B, C, m, x\} \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& !\text{LeQ}[m, -1]$

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

Maple [A] (warning: unable to verify)

Time = 0.10 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.00

method	result
parts	$\frac{(B b^2 + 2Cab) \left(\frac{\tan(dx+c)^3}{3} - \tan(dx+c) + \arctan(\tan(dx+c)) \right)}{d} + \frac{(2Bab + C a^2) \left(\frac{\tan(dx+c)^2}{2} - \frac{\ln(1+\tan(dx+c)^2)}{2} \right)}{d} + \dots$
norman	$(-B a^2 + B b^2 + 2Cab) x + \frac{(B a^2 - B b^2 - 2Cab) \tan(dx+c)}{d} + \frac{(2Bab + C a^2 - C b^2) \tan(dx+c)^2}{2d} + \frac{C b^2}{d} + \dots$
derivativedivides	$\frac{C b^2 \tan(dx+c)^4}{4} + \frac{B b^2 \tan(dx+c)^3}{3} + \frac{2Cab \tan(dx+c)^3}{3} + Bab \tan(dx+c)^2 + \frac{C a^2 \tan(dx+c)^2}{2} - \frac{C b^2 \tan(dx+c)^2}{2} + B a^2 \tan(dx+c) + \dots$
default	$\frac{C b^2 \tan(dx+c)^4}{4} + \frac{B b^2 \tan(dx+c)^3}{3} + \frac{2Cab \tan(dx+c)^3}{3} + Bab \tan(dx+c)^2 + \frac{C a^2 \tan(dx+c)^2}{2} - \frac{C b^2 \tan(dx+c)^2}{2} + B a^2 \tan(dx+c) + \dots$
parallelrisc	$-3C b^2 \tan(dx+c)^4 - 4B b^2 \tan(dx+c)^3 - 8Cab \tan(dx+c)^3 + 12B a^2 dx - 12B b^2 dx - 12Bab \tan(dx+c)^2 - 24Cab dx - \dots$
risc	$-B a^2 x + B b^2 x + 2C ab x - iC a^2 x + \frac{2i(6iC b^2 e^{6i(dx+c)} - 6iBab e^{2i(dx+c)} + 6iC b^2 e^{2i(dx+c)} + 3B a^2 e^{6i(dx+c)})}{d} + \dots$

input

```
int (tan(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2), x, method=_RETURNVERBOSE)
```

output

```
(B*b^2+2*C*a*b)/d*(1/3*tan(d*x+c)^3-tan(d*x+c)+arctan(tan(d*x+c)))+(2*B*a*b+C*a^2)/d*(1/2*tan(d*x+c)^2-1/2*ln(1+tan(d*x+c)^2))+B*a^2/d*(tan(d*x+c)-arctan(tan(d*x+c)))+C*b^2/d*(1/4*tan(d*x+c)^4-1/2*tan(d*x+c)^2+1/2*ln(1+tan(d*x+c)^2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.99

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{3Cb^2 \tan(dx + c)^4 + 4(2Cab + Bb^2) \tan(dx + c)^3 - 12(Ba^2 - 2Cab - Bb^2)dx + 6(Ca^2 + 2Bab - C$$

12 d

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, a
lgorithm="fricas")
```

output

```
1/12*(3*C*b^2*tan(d*x + c)^4 + 4*(2*C*a*b + B*b^2)*tan(d*x + c)^3 - 12*(B*
a^2 - 2*C*a*b - B*b^2)*d*x + 6*(C*a^2 + 2*B*a*b - C*b^2)*tan(d*x + c)^2 +
6*(C*a^2 + 2*B*a*b - C*b^2)*log(1/(tan(d*x + c)^2 + 1)) + 12*(B*a^2 - 2*C*
a*b - B*b^2)*tan(d*x + c))/d
```

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.69

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \left\{ \begin{array}{l} -Ba^2x + \frac{Ba^2 \tan(c+dx)}{d} - \frac{Bab \log(\tan^2(c+dx)+1)}{d} + \frac{Bab \tan^2(c+dx)}{d} + Bb^2x + \frac{Bb^2 \tan^3(c+dx)}{3d} - \frac{Bb^2 \tan(c+dx)}{d} - C \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \tan(c) \end{array} \right.$$

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((-B*a**2*x + B*a**2*tan(c + d*x)/d - B*a*b*log(tan(c + d*x)**2 +
1)/d + B*a*b*tan(c + d*x)**2/d + B*b**2*x + B*b**2*tan(c + d*x)**3/(3*d)
- B*b**2*tan(c + d*x)/d - C*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + C*a**2*t
an(c + d*x)**2/(2*d) + 2*C*a*b*x + 2*C*a*b*tan(c + d*x)**3/(3*d) - 2*C*a*b
*tan(c + d*x)/d + C*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + C*b**2*tan(c +
d*x)**4/(4*d) - C*b**2*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(a + b*tan(c))
**2*(B*tan(c) + C*tan(c)**2)*tan(c), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.99

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{3Cb^2 \tan(dx + c)^4 + 4(2Cab + Bb^2) \tan(dx + c)^3 + 6(Ca^2 + 2Bab - Cb^2) \tan(dx + c)^2 - 12(Ba^2 -$$

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
1/12*(3*C*b^2*tan(d*x + c)^4 + 4*(2*C*a*b + B*b^2)*tan(d*x + c)^3 + 6*(C*a^2 + 2*B*a*b - C*b^2)*tan(d*x + c)^2 - 12*(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c) - 6*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1) + 12*(B*a^2 - 2*C*a*b - B*b^2)*tan(d*x + c))/d
```

Giac [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.43

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ba^2 - 2Cab - Bb^2)(dx + c)}{d} - \frac{(Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{3Cb^2d^3 \tan(dx + c)^4 + 8Cabd^3 \tan(dx + c)^3 + 4Bb^2d^3 \tan(dx + c)^3 + 6Ca^2d^3 \tan(dx + c)^2 + 12B$$

input

```
integrate(tan(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

```
-(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c)/d - 1/2*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/d + 1/12*(3*C*b^2*d^3*tan(d*x + c)^4 + 8*C*a*b*d^3*tan(d*x + c)^3 + 4*B*b^2*d^3*tan(d*x + c)^3 + 6*C*a^2*d^3*tan(d*x + c)^2 + 12*B*a*b*d^3*tan(d*x + c)^2 - 6*C*b^2*d^3*tan(d*x + c)^2 + 12*B*a^2*d^3*tan(d*x + c) - 24*C*a*b*d^3*tan(d*x + c) - 12*B*b^2*d^3*tan(d*x + c))/d^4
```

Mupad [B] (verification not implemented)

Time = 5.50 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= x(-B a^2 + 2 C a b + B b^2) + \frac{\tan(c + dx)^3 \left(\frac{B b^2}{3} + \frac{2 C a b}{3}\right)}{d}$$

$$- \frac{\tan(c + dx) (-B a^2 + 2 C a b + B b^2)}{d}$$

$$- \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{C a^2}{2} + B a b - \frac{C b^2}{2}\right)}{d}$$

$$+ \frac{\tan(c + dx)^2 \left(\frac{C a^2}{2} + B a b - \frac{C b^2}{2}\right)}{d} + \frac{C b^2 \tan(c + dx)^4}{4 d}$$

input

```
int(tan(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^2,x)
```

output

```
x*(B*b^2 - B*a^2 + 2*C*a*b) + (tan(c + d*x)^3*((B*b^2)/3 + (2*C*a*b)/3))/d
- (tan(c + d*x)*(B*b^2 - B*a^2 + 2*C*a*b))/d - (log(tan(c + d*x)^2 + 1)*
(C*a^2)/2 - (C*b^2)/2 + B*a*b))/d + (tan(c + d*x)^2*((C*a^2)/2 - (C*b^2)/2
+ B*a*b))/d + (C*b^2*tan(c + d*x)^4)/(4*d)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.32

$$\int \tan(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-6 \log(\tan(dx + c)^2 + 1) a^2 c - 12 \log(\tan(dx + c)^2 + 1) a b^2 + 6 \log(\tan(dx + c)^2 + 1) b^2 c + 3 \tan(dx + c)}{d}$$

input

```
int(tan(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
( - 6*log(tan(c + d*x)**2 + 1)*a**2*c - 12*log(tan(c + d*x)**2 + 1)*a*b**2
+ 6*log(tan(c + d*x)**2 + 1)*b**2*c + 3*tan(c + d*x)**4*b**2*c + 8*tan(c
+ d*x)**3*a*b*c + 4*tan(c + d*x)**3*b**3 + 6*tan(c + d*x)**2*a**2*c + 12*t
an(c + d*x)**2*a*b**2 - 6*tan(c + d*x)**2*b**2*c + 12*tan(c + d*x)*a**2*b
- 24*tan(c + d*x)*a*b*c - 12*tan(c + d*x)*b**3 - 12*a**2*b*d*x + 24*a*b*c*
d*x + 12*b**3*d*x)/(12*d)
```

3.10 $\int (a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	163
Mathematica [C] (verified)	163
Rubi [A] (verified)	164
Maple [A] (verified)	166
Fricas [A] (verification not implemented)	167
Sympy [A] (verification not implemented)	167
Maxima [A] (verification not implemented)	168
Giac [A] (verification not implemented)	168
Mupad [B] (verification not implemented)	169
Reduce [B] (verification not implemented)	169

Optimal result

Integrand size = 32, antiderivative size = 112

$$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((2abB + a^2C - b^2C) x) - \frac{(a^2B - b^2B - 2abC) \log(\cos(c + dx))}{d}$$

$$+ \frac{b(aB - bC) \tan(c + dx)}{d} + \frac{B(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3bd}$$

output

```
-(2*B*a*b+C*a^2-C*b^2)*x-(B*a^2-B*b^2-2*C*a*b)*ln(cos(d*x+c))/d+b*(B*a-C*b)
)*tan(d*x+c)/d+1/2*B*(a+b*tan(d*x+c))^2/d+1/3*C*(a+b*tan(d*x+c))^3/b/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.26 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.54

$$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2C(a + b \tan(c + dx))^3 + 3(aB + bC) (i((a + ib)^2 \log(i - \tan(c + dx)) - (a - ib)^2 \log(i + \tan(c + dx)))$$

input `Integrate[(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `(2*C*(a + b*Tan[c + d*x])^3 + 3*(a*B + b*C)*(I*((a + I*b)^2*Log[I - Tan[c + d*x]] - (a - I*b)^2*Log[I + Tan[c + d*x]]) - 2*b^2*Tan[c + d*x]) + 3*B*((I*a - b)^3*Log[I - Tan[c + d*x]] - (I*a + b)^3*Log[I + Tan[c + d*x]] + 6*a*b^2*Tan[c + d*x] + b^3*Tan[c + d*x]^2))/(6*b*d)`

Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan(c + dx)^2) dx \\
 & \quad \downarrow \text{4113} \\
 & \int (a + b \tan(c + dx))^2 (B \tan(c + dx) - C) dx + \frac{C(a + b \tan(c + dx))^3}{3bd} \\
 & \quad \downarrow \text{3042} \\
 & \int (a + b \tan(c + dx))^2 (B \tan(c + dx) - C) dx + \frac{C(a + b \tan(c + dx))^3}{3bd} \\
 & \quad \downarrow \text{4011} \\
 & \int (a + b \tan(c + dx))(-bB - aC + (aB - bC) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^2}{2d} + \\
 & \quad \frac{C(a + b \tan(c + dx))^3}{3bd} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \int (a + b \tan(c + dx))(-bB - aC + (aB - bC) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^2}{2d} + \\
& \quad \frac{C(a + b \tan(c + dx))^3}{3bd} \\
& \quad \downarrow \text{4008} \\
& (a^2B - 2abC - b^2B) \int \tan(c + dx) dx - x(a^2C + 2abB - b^2C) + \frac{b(aB - bC) \tan(c + dx)}{d} + \\
& \quad \frac{B(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3bd} \\
& \quad \downarrow \text{3042} \\
& (a^2B - 2abC - b^2B) \int \tan(c + dx) dx - x(a^2C + 2abB - b^2C) + \frac{b(aB - bC) \tan(c + dx)}{d} + \\
& \quad \frac{B(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3bd} \\
& \quad \downarrow \text{3956} \\
& -\frac{(a^2B - 2abC - b^2B) \log(\cos(c + dx))}{d} - x(a^2C + 2abB - b^2C) + \frac{b(aB - bC) \tan(c + dx)}{d} + \\
& \quad \frac{B(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3bd}
\end{aligned}$$

input `Int[(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]`

output `-((2*a*b*B + a^2*C - b^2*C)*x) - ((a^2*B - b^2*B - 2*a*b*C)*Log[Cos[c + d*x]])/d + (b*(a*B - b*C)*Tan[c + d*x])/d + (B*(a + b*Tan[c + d*x])^2)/(2*d) + (C*(a + b*Tan[c + d*x])^3)/(3*b*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4008 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

```
rule 4011 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4113 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_)
+ (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.07

method	result
norman	$(-2Bab - C a^2 + C b^2) x + \frac{(2Bab + C a^2 - C b^2) \tan(dx+c)}{d} + \frac{C b^2 \tan(dx+c)^3}{3d} + \frac{b(Bb+2Ca) \tan(dx+c)}{2d}$
parts	$\frac{(2Bab + C a^2) (\tan(dx+c) - \arctan(\tan(dx+c)))}{d} + \frac{(B b^2 + 2C ab) \left(\frac{\tan(dx+c)^2}{2} - \frac{\ln(1 + \tan(dx+c)^2)}{2} \right)}{d} + \frac{B a^2 \ln(1 + \tan(dx+c)^2)}{2d}$
derivativedivides	$\frac{\frac{C \tan(dx+c)^3 b^2}{3} + \frac{B \tan(dx+c)^2 b^2}{2} + C \tan(dx+c)^2 ab + 2B \tan(dx+c) ab + C \tan(dx+c) a^2 - C \tan(dx+c) b^2 + \frac{(B a^2 - B b^2 - C a^2 + C b^2) \tan(dx+c)}{d}}{d}$
default	$\frac{\frac{C \tan(dx+c)^3 b^2}{3} + \frac{B \tan(dx+c)^2 b^2}{2} + C \tan(dx+c)^2 ab + 2B \tan(dx+c) ab + C \tan(dx+c) a^2 - C \tan(dx+c) b^2 + \frac{(B a^2 - B b^2 - C a^2 + C b^2) \tan(dx+c)}{d}}{d}$
parallelrisc	$\frac{2C \tan(dx+c)^3 b^2 - 12B ab dx + 3B \tan(dx+c)^2 b^2 - 6C a^2 dx + 6C b^2 dx + 6C \tan(dx+c)^2 ab + 3B \ln(1 + \tan(dx+c)^2) a^2 - 3C \tan(dx+c) b^2}{6d}$
risc	$-\frac{4iCabc}{d} - iB b^2 x + \frac{2i(-3iB b^2 e^{4i(dx+c)} - 6iCab e^{4i(dx+c)} + 6Bab e^{4i(dx+c)} + 3C a^2 e^{4i(dx+c)} - 6C b^2 e^{4i(dx+c)})}{d}$

input `int((a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output $(-2Bab-Ca^2+Cb^2)x+(2Bab+Ca^2-Cb^2)/d\tan(dx+c)+1/3Cb^2/d\tan(dx+c)^3+1/2b(Bb+2Ca)/d\tan(dx+c)^2+1/2(Ba^2-Bb^2-2Cab)/d\ln(1+\tan(dx+c)^2)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.06

$$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Cb^2 \tan(dx+c)^3 - 6(Ca^2 + 2Bab - Cb^2)dx + 3(2Cab + Bb^2) \tan(dx+c)^2 - 3(Ba^2 - 2Cab - Bb^2) \ln(1+\tan(dx+c)^2)}{6d}$$

input `integrate((a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")`

output $1/6*(2Cb^2*\tan(dx+c)^3 - 6*(Ca^2 + 2Bab - Cb^2)*dx + 3*(2Cab + Bb^2)*\tan(dx+c)^2 - 3*(Ba^2 - 2Cab - Bb^2)*\log(1/(\tan(dx+c)^2 + 1)) + 6*(Ca^2 + 2Bab - Cb^2)*\tan(dx+c))/d$

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.73

$$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \frac{Ba^2 \log(\tan^2(c+dx)+1)}{2d} - 2Babx + \frac{2Bab \tan(c+dx)}{d} - \frac{Bb^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{Bb^2 \tan^2(c+dx)}{2d} - Ca^2x + \frac{Ca^2 \tan(c+dx)}{d} \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \end{cases}$$

input `integrate((a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output

```
Piecewise((B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - 2*B*a*b*x + 2*B*a*b*tan(c + d*x)/d - B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**2*tan(c + d*x)**2/(2*d) - C*a**2*x + C*a**2*tan(c + d*x)/d - C*a*b*log(tan(c + d*x)**2 + 1)/d + C*a*b*tan(c + d*x)**2/d + C*b**2*x + C*b**2*tan(c + d*x)**3/(3*d) - C*b**2*tan(c + d*x)/d, Ne(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*tan(c)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.07

$$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2 C b^2 \tan(dx + c)^3 + 3 (2 C a b + B b^2) \tan(dx + c)^2 - 6 (C a^2 + 2 B a b - C b^2)(dx + c) + 3 (B a^2 - 2 C a b)}{6 d}$$

input

```
integrate((a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
1/6*(2*C*b^2*tan(d*x + c)^3 + 3*(2*C*a*b + B*b^2)*tan(d*x + c)^2 - 6*(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c) + 3*(B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1) + 6*(C*a^2 + 2*B*a*b - C*b^2)*tan(d*x + c))/d
```

Giac [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.44

$$\int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(C a^2 + 2 B a b - C b^2)(d x + c)}{d} + \frac{(B a^2 - 2 C a b - B b^2) \log(\tan(dx + c)^2 + 1)}{2 d}$$

$$+ \frac{2 C b^2 d^2 \tan(dx + c)^3 + 6 C a b d^2 \tan(dx + c)^2 + 3 B b^2 d^2 \tan(dx + c) + 6 C a^2 d^2 \tan(dx + c) + 12 B a b d^2}{6 d^3}$$

input

```
integrate((a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

$$-(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c)/d + 1/2*(B*a^2 - 2*C*a*b - B*b^2)*\log(\tan(d*x + c)^2 + 1)/d + 1/6*(2*C*b^2*d^2*\tan(d*x + c)^3 + 6*C*a*b*d^2*\tan(d*x + c)^2 + 3*B*b^2*d^2*\tan(d*x + c)^2 + 6*C*a^2*d^2*\tan(d*x + c) + 12*B*a*b*d^2*\tan(d*x + c) - 6*C*b^2*d^2*\tan(d*x + c))/d^3$$
Mupad [B] (verification not implemented)

Time = 5.27 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.08

$$\begin{aligned} & \int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \frac{\tan(c + dx)^2 \left(\frac{Bb^2}{2} + Cab \right)}{d} - x (Ca^2 + 2Bab - Cb^2) \\ &+ \frac{\tan(c + dx) (Ca^2 + 2Bab - Cb^2)}{d} \\ &- \frac{\ln(\tan(c + dx)^2 + 1) \left(-\frac{Ba^2}{2} + Cab + \frac{Bb^2}{2} \right)}{d} + \frac{Cb^2 \tan(c + dx)^3}{3d} \end{aligned}$$

input

$$\text{int}((B*\tan(c + d*x) + C*\tan(c + d*x)^2)*(a + b*\tan(c + d*x))^2,x)$$

output

$$\frac{(\tan(c + d*x)^2*((B*b^2)/2 + C*a*b))/d - x*(C*a^2 - C*b^2 + 2*B*a*b) + (\tan(c + d*x)*(C*a^2 - C*b^2 + 2*B*a*b))/d - (\log(\tan(c + d*x)^2 + 1)*((B*b^2)/2 - (B*a^2)/2 + C*a*b))/d + (C*b^2*\tan(c + d*x)^3)/(3*d)}$$
Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.38

$$\begin{aligned} & \int (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \frac{3 \log(\tan(dx + c)^2 + 1) a^2 b - 6 \log(\tan(dx + c)^2 + 1) abc - 3 \log(\tan(dx + c)^2 + 1) b^3 + 2 \tan(dx + c)}{d} \end{aligned}$$

input

$$\text{int}((a+b*\tan(d*x+c))^2*(B*\tan(d*x+c)+C*\tan(d*x+c)^2),x)$$

output

```
(3*log(tan(c + d*x)**2 + 1)*a**2*b - 6*log(tan(c + d*x)**2 + 1)*a*b*c - 3*  
log(tan(c + d*x)**2 + 1)*b**3 + 2*tan(c + d*x)**3*b**2*c + 6*tan(c + d*x)*  
*2*a*b*c + 3*tan(c + d*x)**2*b**3 + 6*tan(c + d*x)*a**2*c + 12*tan(c + d*x)  
)a*b**2 - 6*tan(c + d*x)*b**2*c - 6*a**2*c*d*x - 12*a*b**2*d*x + 6*b**2*c  
*d*x)/(6*d)
```

3.11 $\int \cot(c+dx)(a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	171
Mathematica [C] (verified)	171
Rubi [A] (verified)	172
Maple [A] (verified)	174
Fricas [A] (verification not implemented)	175
Sympy [A] (verification not implemented)	175
Maxima [A] (verification not implemented)	176
Giac [A] (verification not implemented)	176
Mupad [B] (verification not implemented)	177
Reduce [B] (verification not implemented)	177

Optimal result

Integrand size = 38, antiderivative size = 87

$$\int \cot(c+dx)(a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= (a^2B - b^2B - 2abC) x - \frac{(2abB + a^2C - b^2C) \log(\cos(c+dx))}{d}$$

$$+ \frac{b(bB + aC) \tan(c+dx)}{d} + \frac{C(a+b \tan(c+dx))^2}{2d}$$

output

```
(B*a^2-B*b^2-2*C*a*b)*x-(2*B*a*b+C*a^2-C*b^2)*ln(cos(d*x+c))/d+b*(B*b+C*a)*tan(d*x+c)/d+1/2*C*(a+b*tan(d*x+c))^2/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.10

$$\int \cot(c+dx)(a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \frac{(a+ib)^2(-iB+C) \log(i-\tan(c+dx)) + (a-ib)^2(iB+C) \log(i+\tan(c+dx)) + 2b(bB+2aC) \tan(c+dx)}{2d}$$

input

```
Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
((a + I*b)^2*((-I)*B + C)*Log[I - Tan[c + d*x]] + (a - I*b)^2*(I*B + C)*Log[I + Tan[c + d*x]] + 2*b*(b*B + 2*a*C)*Tan[c + d*x] + b^2*C*Tan[c + d*x]^2)/(2*d)
```

Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4115, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx))}{\tan(c + dx)} dx$$

$$\downarrow 4115$$

$$\int (a + b \tan(c + dx))^2 (B + C \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(c + dx))^2 (B + C \tan(c + dx)) dx$$

$$\downarrow 4011$$

$$\int (a + b \tan(c + dx))(aB - bC + (bB + aC) \tan(c + dx)) dx + \frac{C(a + b \tan(c + dx))^2}{2d}$$

$$\downarrow 3042$$

$$\int (a + b \tan(c + dx))(aB - bC + (bB + aC) \tan(c + dx)) dx + \frac{C(a + b \tan(c + dx))^2}{2d}$$

$$\begin{aligned}
& \downarrow 4008 \\
& (a^2C + 2abB - b^2C) \int \tan(c + dx)dx + x(a^2B - 2abC - b^2B) + \frac{b(aC + bB) \tan(c + dx)}{d} + \\
& \quad \frac{C(a + b \tan(c + dx))^2}{2d} \\
& \downarrow 3042 \\
& (a^2C + 2abB - b^2C) \int \tan(c + dx)dx + x(a^2B - 2abC - b^2B) + \frac{b(aC + bB) \tan(c + dx)}{d} + \\
& \quad \frac{C(a + b \tan(c + dx))^2}{2d} \\
& \downarrow 3956 \\
& -\frac{(a^2C + 2abB - b^2C) \log(\cos(c + dx))}{d} + x(a^2B - 2abC - b^2B) + \frac{b(aC + bB) \tan(c + dx)}{d} + \\
& \quad \frac{C(a + b \tan(c + dx))^2}{2d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(a^2*B - b^2*B - 2*a*b*C)*x - ((2*a*b*B + a^2*C - b^2*C)*Log[Cos[c + d*x]])/d + (b*(b*B + a*C)*Tan[c + d*x])/d + (C*(a + b*Tan[c + d*x])^2)/(2*d)
```

Defintions of rubi rules used

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4008

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

rule 4011

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00

method	result
parallelrisc	$\frac{\ln(\sec(dx+c)^2)(2Bab+Ca^2-Cb^2)+Cb^2 \tan(dx+c)^2+(2Bb^2+4Cab) \tan(dx+c)+2dx(Ba^2-Bb^2-2Cab)}{2d}$
norman	$(Ba^2 - Bb^2 - 2Cab)x + \frac{b(Bb+2Ca) \tan(dx+c)}{d} + \frac{Cb^2 \tan(dx+c)^2}{2d} + \frac{(2Bab+Ca^2-Cb^2) \ln(1+\tan(dx+c))}{2d}$
derivativedivides	$-\frac{\frac{(-2Bab-Ca^2+Cb^2) \ln(\cot(dx+c)^2+1)}{2} + (Ba^2-Bb^2-2Cab) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))\right) + (2Bab+Ca^2-Cb^2) \ln(\cot(dx+c))}{d}$
default	$-\frac{\frac{(-2Bab-Ca^2+Cb^2) \ln(\cot(dx+c)^2+1)}{2} + (Ba^2-Bb^2-2Cab) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))\right) + (2Bab+Ca^2-Cb^2) \ln(\cot(dx+c))}{d}$
risc	$Ba^2x - Bb^2x - 2Cabx + \frac{4iBabc}{d} + iCa^2x - iCb^2x - \frac{2iCb^2c}{d} + \frac{2iCa^2c}{d} + \frac{2ib(-iCbe^{2i(dx+c)})}{d}$

input

```
int(cot(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_
RETURNVERBOSE)
```

output

```
1/2*(ln(sec(d*x+c)^2)*(2*B*a*b+C*a^2-C*b^2)+C*b^2*tan(d*x+c)^2+(2*B*b^2+4*
C*a*b)*tan(d*x+c)+2*d*x*(B*a^2-B*b^2-2*C*a*b))/d
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Cb^2 \tan(dx + c)^2 + 2(Ba^2 - 2Cab - Bb^2)dx - (Ca^2 + 2Bab - Cb^2) \log\left(\frac{1}{\tan(dx+c)^2+1}\right) + 2(2Cab + B}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")`

output `1/2*(C*b^2*tan(d*x + c)^2 + 2*(B*a^2 - 2*C*a*b - B*b^2)*d*x - (C*a^2 + 2*B*a*b - C*b^2)*log(1/(tan(d*x + c)^2 + 1)) + 2*(2*C*a*b + B*b^2)*tan(d*x + c))/d`

Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.74

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} Ba^2x + \frac{Bab \log(\tan^2(c+dx)+1)}{d} - Bb^2x + \frac{Bb^2 \tan(c+dx)}{d} + \frac{Ca^2 \log(\tan^2(c+dx)+1)}{2d} - 2Cabx + \frac{2Cab \tan(c+dx)}{d} - \frac{CB}{d} \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \cot(c) \end{cases}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((B*a**2*x + B*a*b*log(tan(c + d*x)**2 + 1)/d - B*b**2*x + B*b**2*tan(c + d*x)/d + C*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - 2*C*a*b*x + 2*C*a*b*tan(c + d*x)/d - C*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + C*b**2*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*tan(c)**2)*cot(c), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Cb^2 \tan(dx + c)^2 + 2(Ba^2 - 2Cab - Bb^2)(dx + c) + (Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1) + 2}{2d}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
1/2*(C*b^2*tan(d*x + c)^2 + 2*(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c) + (C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1) + 2*(2*C*a*b + B*b^2)*tan(d*x + c))/d
```

Giac [A] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.21

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ba^2 - 2Cab - Bb^2)(dx + c)}{d} + \frac{(Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{Cb^2 d \tan(dx + c)^2 + 4Cab d \tan(dx + c) + 2Bb^2 d \tan(dx + c)}{2d^2}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

```
(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c)/d + 1/2*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/d + 1/2*(C*b^2*d*tan(d*x + c)^2 + 4*C*a*b*d*tan(d*x + c) + 2*B*b^2*d*tan(d*x + c))/d^2
```

Mupad [B] (verification not implemented)

Time = 5.23 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{C a^2}{2} + B a b - \frac{C b^2}{2} \right)}{d} - x(-B a^2 + 2 C a b + B b^2)$$

$$+ \frac{\tan(c + dx) (B b^2 + 2 C a b)}{d} + \frac{C b^2 \tan(c + dx)^2}{2 d}$$

input

```
int(cot(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^2,x)
```

output

```
(log(tan(c + d*x)^2 + 1)*((C*a^2)/2 - (C*b^2)/2 + B*a*b))/d - x*(B*b^2 - B*a^2 + 2*C*a*b) + (tan(c + d*x)*(B*b^2 + 2*C*a*b))/d + (C*b^2*tan(c + d*x)^2)/(2*d)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 542, normalized size of antiderivative = 6.23

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

= Too large to display

input

```
int(cot(d*x+c)*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
( - 4*cos(c + d*x)*sin(c + d*x)*a*b*c - 2*cos(c + d*x)*sin(c + d*x)*b**3 +
  2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**2*c + 4*log(tan((c + d*
x)/2)**2 + 1)*sin(c + d*x)**2*a*b**2 - 2*log(tan((c + d*x)/2)**2 + 1)*sin(
c + d*x)**2*b**2*c - 2*log(tan((c + d*x)/2)**2 + 1)*a**2*c - 4*log(tan((c
+ d*x)/2)**2 + 1)*a*b**2 + 2*log(tan((c + d*x)/2)**2 + 1)*b**2*c - 2*log(t
an((c + d*x)/2) - 1)*sin(c + d*x)**2*a**2*c - 4*log(tan((c + d*x)/2) - 1)*
sin(c + d*x)**2*a*b**2 + 2*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*b**2*
c + 2*log(tan((c + d*x)/2) - 1)*a**2*c + 4*log(tan((c + d*x)/2) - 1)*a*b**
2 - 2*log(tan((c + d*x)/2) - 1)*b**2*c - 2*log(tan((c + d*x)/2) + 1)*sin(c
+ d*x)**2*a**2*c - 4*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*a*b**2 + 2
*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*b**2*c + 2*log(tan((c + d*x)/2)
+ 1)*a**2*c + 4*log(tan((c + d*x)/2) + 1)*a*b**2 - 2*log(tan((c + d*x)/2)
+ 1)*b**2*c + 2*sin(c + d*x)**2*a**2*b*d*x - 4*sin(c + d*x)**2*a*b*c*d*x
- 2*sin(c + d*x)**2*b**3*d*x - sin(c + d*x)**2*b**2*c - 2*a**2*b*d*x + 4*a
*b*c*d*x + 2*b**3*d*x)/(2*d*(sin(c + d*x)**2 - 1))
```

3.12 $\int \cot^2(c+dx)(a+b \tan(c+dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	179
Mathematica [C] (verified)	179
Rubi [A] (verified)	180
Maple [A] (verified)	182
Fricas [A] (verification not implemented)	183
Sympy [B] (verification not implemented)	184
Maxima [A] (verification not implemented)	184
Giac [A] (verification not implemented)	185
Mupad [B] (verification not implemented)	185
Reduce [B] (verification not implemented)	186

Optimal result

Integrand size = 40, antiderivative size = 70

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (2abB + a^2C - b^2C) x - \frac{b(bB + 2aC) \log(\cos(c + dx))}{d}$$

$$+ \frac{a^2B \log(\sin(c + dx))}{d} + \frac{b^2C \tan(c + dx)}{d}$$

output

```
(2*B*a*b+C*a^2-C*b^2)*x-b*(B*b+2*C*a)*ln(cos(d*x+c))/d+a^2*B*ln(sin(d*x+c)
)/d+b^2*C*tan(d*x+c)/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.30

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{(a + ib)^2(B + iC) \log(i - \tan(c + dx)) - 2a^2B \log(\tan(c + dx)) + (a - ib)^2(B - iC) \log(i + \tan(c + dx))}{2d}$$

input

```
Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
-1/2*((a + I*b)^2*(B + I*C)*Log[I - Tan[c + d*x]] - 2*a^2*B*Log[Tan[c + d*x]] + (a - I*b)^2*(B - I*C)*Log[I + Tan[c + d*x]] - 2*b^2*C*Tan[c + d*x])/d
```

Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.225$, Rules used = {3042, 4115, 3042, 4089, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx))}{\tan^2(c + dx)} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \cot(c + dx)(a + b \tan(c + dx))^2 (B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2 (B + C \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4089} \\
 & \int \cot(c + dx) (Ba^2 + b(bB + 2aC) \tan^2(c + dx) + (Ca^2 + 2bBa - b^2C) \tan(c + dx)) dx + \\
 & \quad \frac{b^2C \tan(c + dx)}{d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\int \frac{Ba^2 + b(bB + 2aC) \tan(c + dx)^2 + (Ca^2 + 2bBa - b^2C) \tan(c + dx)}{\tan(c + dx)} dx + \frac{b^2C \tan(c + dx)}{d}$$

↓ 4107

$$a^2B \int \cot(c + dx) dx + b(2aC + bB) \int \tan(c + dx) dx + x(a^2C + 2abB - b^2C) + \frac{b^2C \tan(c + dx)}{d}$$

↓ 3042

$$a^2B \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b(2aC + bB) \int \tan(c + dx) dx + x(a^2C + 2abB - b^2C) + \frac{b^2C \tan(c + dx)}{d}$$

↓ 25

$$a^2(-B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b(2aC + bB) \int \tan(c + dx) dx + x(a^2C + 2abB - b^2C) + \frac{b^2C \tan(c + dx)}{d}$$

↓ 3956

$$x(a^2C + 2abB - b^2C) + \frac{a^2B \log(-\sin(c + dx))}{\frac{d}{b^2C \tan(c + dx)}} - \frac{b(2aC + bB) \log(\cos(c + dx))}{d} +$$

input

```
Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(2*a*b*B + a^2*C - b^2*C)*x - (b*(b*B + 2*a*C)*Log[Cos[c + d*x]])/d + (a^2*B*Log[-Sin[c + d*x]])/d + (b^2*C*Tan[c + d*x])/d
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4089 $\text{Int}[(((a_.) + (b_.)\tan[(e_.) + (f_.)(x_.)])^2((A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)])) / ((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[b^2 * B * (\text{Tan}[e + f*x] / (d*f)), x] + \text{Simp}[1/d \text{ Int}[(a^2*A*d - b^2*B*c + (2*a*A*b + B*(a^2 - b^2))*d*\text{Tan}[e + f*x] + (A*b^2*d - b*B*(b*c - 2*a*d))*\text{Tan}[e + f*x]^2) / (c + d*\text{Tan}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4107 $\text{Int}[(A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)] + (C_.)\tan[(e_.) + (f_.)(x_.)]^2 / \tan[(e_.) + (f_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[B*x, x] + (\text{Simp}[A \text{ Int}[1/\text{Tan}[e + f*x], x], x] + \text{Simp}[C \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{e, f, A, B, C\}, x] \&\& \text{NeQ}[A, C]$

rule 4115 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)} * ((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)])^{(n_.)} * ((A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)] + (C_.)\tan[(e_.) + (f_.)(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[1/b^2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^{(m+1)} * (c + d*\text{Tan}[e + f*x])^n * (b*B - a*C + b*C*\text{Tan}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[A*b^2 - a*b*B + a^2*C, 0]$

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.14

method	result
parallelrisc	$\frac{(-B a^2 + B b^2 + 2Cab) \ln(\sec(dx+c)^2) + 2B a^2 \ln(\tan(dx+c)) + 2C \tan(dx+c)b^2 + 4x(Bab + \frac{1}{2}C a^2 - \frac{1}{2}C b^2)d}{2d}$
derivativdivides	$-\frac{(B a^2 - B b^2 - 2Cab) \ln(\cot(dx+c)^2 + 1)}{2} + (2Bab + C a^2 - C b^2) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))\right) - \frac{C b^2}{\cot(dx+c)} + b(Bb + 2Ca) \ln(\cot(dx+c))$
default	$-\frac{(B a^2 - B b^2 - 2Cab) \ln(\cot(dx+c)^2 + 1)}{2} + (2Bab + C a^2 - C b^2) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))\right) - \frac{C b^2}{\cot(dx+c)} + b(Bb + 2Ca) \ln(\cot(dx+c))$
norman	$\frac{(2Bab + C a^2 - C b^2)x \tan(dx+c) + \frac{C b^2 \tan(dx+c)^2}{d}}{\tan(dx+c)} + \frac{B a^2 \ln(\tan(dx+c))}{d} - \frac{(B a^2 - B b^2 - 2Cab) \ln(1 + \tan(dx+c)^2)}{2d}$
risc	$iB b^2 x + \frac{4iCab c}{d} + \frac{2iC b^2}{d(e^{2i(dx+c)} + 1)} + 2Babx + C a^2 x - C b^2 x - iB a^2 x - \frac{2iB a^2 c}{d} + 2iCabx$

```
input int (cot (d*x+c)^2*(a+b*tan (d*x+c))^2*(B*tan (d*x+c)+C*tan (d*x+c)^2) ,x,method
=_RETURNVERBOSE)
```

```
output 1/2*((-B*a^2+B*b^2+2*C*a*b)*ln(sec (d*x+c)^2)+2*B*a^2*ln(tan (d*x+c))+2*C*ta
n(d*x+c)*b^2+4*x*(B*a*b+1/2*C*a^2-1/2*C*b^2)*d)/d
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.31

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{B a^2 \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + 2 C b^2 \tan(dx+c) + 2(C a^2 + 2 B a b - C b^2) dx - (2 C a b + B b^2) \log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2 d}$$

```
input integrate(cot (d*x+c)^2*(a+b*tan (d*x+c))^2*(B*tan (d*x+c)+C*tan (d*x+c)^2) ,x,
algorithm="fricas")
```

```
output 1/2*(B*a^2*log(tan (d*x + c)^2/(tan (d*x + c)^2 + 1)) + 2*C*b^2*tan (d*x + c)
+ 2*(C*a^2 + 2*B*a*b - C*b^2)*d*x - (2*C*a*b + B*b^2)*log(1/(tan (d*x + c)
^2 + 1)))/d
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(66) = 132$.

Time = 0.69 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.94

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} -\frac{Ba^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba^2 \log(\tan(c+dx))}{d} + 2Babx + \frac{Bb^2 \log(\tan^2(c+dx)+1)}{2d} + Ca^2x + \frac{Cab \log(\tan^2(c+dx)+1)}{d} \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \cot^2(c) \end{cases}$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((-B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**2*log(tan(c + d*x)))/d + 2*B*a*b*x + B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + C*a**2*x + C*a*b*log(tan(c + d*x)**2 + 1)/d - C*b**2*x + C*b**2*tan(c + d*x)/d, Ne(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*tan(c)**2)*cot(c)**2, True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.21

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Ba^2 \log(\tan(dx + c)) + 2Cb^2 \tan(dx + c) + 2(Ca^2 + 2Bab - Cb^2)(dx + c) - (Ba^2 - 2Cab - Bb^2)}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")`

output `1/2*(2*B*a^2*log(tan(d*x + c)) + 2*C*b^2*tan(d*x + c) + 2*(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c) - (B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1))/d`

Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.29

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Ba^2 \log(|\tan(dx + c)|)}{d} + \frac{Cb^2 \tan(dx + c)}{d} + \frac{(Ca^2 + 2 Bab - Cb^2)(dx + c)}{d}$$

$$- \frac{(Ba^2 - 2Cab - Bb^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

input

```
integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
B*a^2*log(abs(tan(d*x + c)))/d + C*b^2*tan(d*x + c)/d + (C*a^2 + 2*B*a*b -
C*b^2)*(d*x + c)/d - 1/2*(B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1
)/d
```

Mupad [B] (verification not implemented)

Time = 5.32 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.29

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Ba^2 \ln(\tan(c + dx))}{d} + \frac{\ln(\tan(c + dx) + i) (B - C i) (b + a i)^2}{2d}$$

$$+ \frac{Cb^2 \tan(c + dx)}{d} + \frac{\ln(\tan(c + dx) - i) (B + C i) (-b + a i)^2}{2d}$$

input

```
int(cot(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
)^2,x)
```

output

```
(B*a^2*log(tan(c + d*x)))/d + (log(tan(c + d*x) + i)*(B - C*i)*(a*i + b
)^2)/(2*d) + (C*b^2*tan(c + d*x))/d + (log(tan(c + d*x) - i)*(B + C*i)*(
a*i - b)^2)/(2*d)
```


3.13 $\int \cot^3(c+dx)(a+b \tan(c+dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	187
Mathematica [C] (verified)	187
Rubi [A] (verified)	188
Maple [A] (verified)	190
Fricas [A] (verification not implemented)	191
Sympy [B] (verification not implemented)	192
Maxima [A] (verification not implemented)	192
Giac [A] (verification not implemented)	193
Mupad [B] (verification not implemented)	193
Reduce [B] (verification not implemented)	194

Optimal result

Integrand size = 40, antiderivative size = 72

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((a^2B - b^2B - 2abC) x) - \frac{a^2B \cot(c + dx)}{d}$$

$$- \frac{b^2C \log(\cos(c + dx))}{d} + \frac{a(2bB + aC) \log(\sin(c + dx))}{d}$$

output

```
-(B*a^2-B*b^2-2*C*a*b)*x-a^2*B*cot(d*x+c)/d-b^2*C*ln(cos(d*x+c))/d+a*(2*B*b+C*a)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.39

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-2a^2B \cot(c + dx) + i(a + ib)^2(B + iC) \log(i - \tan(c + dx)) + 2a(2bB + aC) \log(\tan(c + dx)) - (a - ib)^2C \log(\cos(c + dx))}{2d}$$

input

```
Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(-2*a^2*B*Cot[c + d*x] + I*(a + I*b)^2*(B + I*C)*Log[I - Tan[c + d*x]] + 2*a*(2*b*B + a*C)*Log[Tan[c + d*x]] - (a - I*b)^2*(I*B + C)*Log[I + Tan[c + d*x]])/(2*d)
```

Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.225$, Rules used = {3042, 4115, 3042, 4087, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx))}{\tan^3(c + dx)} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \cot^2(c + dx)(a + b \tan(c + dx))^2 (B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2 (B + C \tan(c + dx))}{\tan^2(c + dx)} dx \\
 & \quad \downarrow \text{4087} \\
 & \int \cot(c + dx) (b^2 C \tan^2(c + dx) - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC)) dx - \\
 & \quad \frac{a^2 B \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{b^2 C \tan(c+dx)^2 - (Ba^2 - 2bCa - b^2 B) \tan(c+dx) + a(2bB + aC)}{\tan(c+dx)} dx - \frac{a^2 B \cot(c+dx)}{d} \\
& \quad \downarrow \text{4107} \\
& a(aC + 2bB) \int \cot(c+dx) dx + b^2 C \int \tan(c+dx) dx - x(a^2 B - 2abC - b^2 B) - \frac{a^2 B \cot(c+dx)}{d} \\
& \quad \downarrow \text{3042} \\
& a(aC + 2bB) \int -\tan\left(c+dx + \frac{\pi}{2}\right) dx + b^2 C \int \tan(c+dx) dx - x(a^2 B - 2abC - b^2 B) - \\
& \quad \frac{a^2 B \cot(c+dx)}{d} \\
& \quad \downarrow \text{25} \\
& -a(aC + 2bB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b^2 C \int \tan(c+dx) dx - \\
& \quad x(a^2 B - 2abC - b^2 B) - \frac{a^2 B \cot(c+dx)}{d} \\
& \quad \downarrow \text{3956} \\
& -x(a^2 B - 2abC - b^2 B) - \frac{a^2 B \cot(c+dx)}{d} + \frac{a(aC + 2bB) \log(-\sin(c+dx))}{d} - \\
& \quad \frac{b^2 C \log(\cos(c+dx))}{d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
-((a^2*B - b^2*B - 2*a*b*C)*x) - (a^2*B*Cot[c + d*x])/d - (b^2*C*Log[Cos[c + d*x]])/d + (a*(2*b*B + a*C)*Log[-Sin[c + d*x]])/d
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4087 $\text{Int}[((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^2*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[-(B*c - A*d)*(b*c - a*d)^2*((c + d*\text{Tan}[e + f*x])^{(n + 1)}/(f*d^2*(n + 1)*(c^2 + d^2))), x] + \text{Simp}[1/(d*(c^2 + d^2)) \text{Int}[(c + d*\text{Tan}[e + f*x])^{(n + 1)}*\text{Simp}[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*\text{Tan}[e + f*x] + b^2*B*(c^2 + d^2)*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{LtQ}[n, -1]$

rule 4107 $\text{Int}(((A_) + (B_.)*\tan[(e_.) + (f_.)*(x_.)] + (C_.)*\tan[(e_.) + (f_.)*(x_.)]^2)/\tan[(e_.) + (f_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[B*x, x] + (\text{Simp}[A \text{Int}[1/\text{Tan}[e + f*x], x], x] + \text{Simp}[C \text{Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{e, f, A, B, C\}, x] \&\& \text{NeQ}[A, C]$

rule 4115 $\text{Int}(((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_.)] + (C_.)*\tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[1/b^2 \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m + 1)}*(c + d*\text{Tan}[e + f*x])^n*(b*B - a*C + b*C*\text{Tan}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[A*b^2 - a*b*B + a^2*C, 0]$

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.17

method	result
derivativedivides	$\frac{B b^2(dx+c) - C b^2 \ln(\cos(dx+c)) + 2Bab \ln(\sin(dx+c)) + 2Cab(dx+c) + B a^2(-\cot(dx+c) - dx - c) + C a^2 \ln(\sin(dx+c))}{d}$
default	$\frac{B b^2(dx+c) - C b^2 \ln(\cos(dx+c)) + 2Bab \ln(\sin(dx+c)) + 2Cab(dx+c) + B a^2(-\cot(dx+c) - dx - c) + C a^2 \ln(\sin(dx+c))}{d}$
parallelrisch	$\frac{(-2Bab - C a^2 + C b^2) \ln(\sec(dx+c)^2) + (4Bab + 2C a^2) \ln(\tan(dx+c)) - 2B \cot(dx+c) a^2 - 2dx(B a^2 - B b^2 - 2Cab)}{2d}$
norman	$\frac{(-B a^2 + B b^2 + 2Cab)x \tan(dx+c)^2 - \frac{B a^2 \tan(dx+c)}{d}}{\tan(dx+c)^2} + \frac{a(2Bb + Ca) \ln(\tan(dx+c))}{d} - \frac{(2Bab + C a^2 - C b^2) \ln(1 + \tan(dx+c))}{2d}$
risch	$-B a^2 x + B b^2 x + 2Cabx + \frac{2iC b^2 c}{d} - \frac{2iC a^2 c}{d} - iC a^2 x + iC b^2 x - \frac{4iBabc}{d} - \frac{2iB a^2}{d(e^{2i(dx+c)} - 1)}$

input `int(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(B*b^2*(d*x+c)-C*b^2*ln(cos(d*x+c))+2*B*a*b*ln(sin(d*x+c))+2*C*a*b*(d*x+c)+B*a^2*(-cot(d*x+c)-d*x-c)+C*a^2*ln(sin(d*x+c)))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.56

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx =$$

$$\frac{C b^2 \log\left(\frac{1}{\tan(dx+c)^2+1}\right) \tan(dx+c) + 2(B a^2 - 2C a b - B b^2) dx \tan(dx+c) + 2B a^2 - (C a^2 + 2B a b)}{2d \tan(dx+c)}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output `-1/2*(C*b^2*log(1/(tan(d*x+c)^2+1))*tan(d*x+c)+2*(B*a^2-2*C*a*b-B*b^2)*d*x*tan(d*x+c)+2*B*a^2-(C*a^2+2*B*a*b)*log(tan(d*x+c)^2/(tan(d*x+c)^2+1))*tan(d*x+c))/(d*tan(d*x+c))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 158 vs. $2(66) = 132$.

Time = 1.07 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.19

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \cot^3(c) \\ \text{NaN} \\ -Ba^2x - \frac{Ba^2}{d \tan(c+dx)} - \frac{Bab \log(\tan^2(c+dx)+1)}{d} + \frac{2Bab \log(\tan(c+dx))}{d} + Bb^2x - \frac{Ca^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{Ca^2 \log(\tan(c+dx))}{d} \end{cases}$$

input

```
integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*tan(c)**2)*cot(c)**3, Eq(d, 0)), (nan, Eq(c, -d*x)), (-B*a**2*x - B*a**2/(d*tan(c + d*x)) - B*a*b*log(tan(c + d*x)**2 + 1)/d + 2*B*a*b*log(tan(c + d*x))/d + B*b**2*x - C*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + C*a**2*log(tan(c + d*x))/d + 2*C*a*b*x + C*b**2*log(tan(c + d*x)**2 + 1)/(2*d), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.29

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx = \frac{2(Ba^2 - 2Cab - Bb^2)(dx + c) + (Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1) - 2(Ca^2 + 2Bab) \log(\tan(c + dx))}{2d}$$

input

```
integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="maxima")
```

output

```
-1/2*(2*(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c) + (C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1) - 2*(C*a^2 + 2*B*a*b)*log(tan(d*x + c)) + 2*B*a^2/tan(d*x + c))/d
```

Giac [A] (verification not implemented)

Time = 0.88 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.40

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ba^2 - 2Cab - Bb^2)(dx + c)}{d} - \frac{(Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{(Ca^2 + 2Bab) \log(|\tan(dx + c)|)}{d} - \frac{Ba^2}{d \tan(dx + c)}$$

input

```
integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
-(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c)/d - 1/2*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/d + (C*a^2 + 2*B*a*b)*log(abs(tan(d*x + c)))/d - B*a^2/(d*tan(d*x + c))
```

Mupad [B] (verification not implemented)

Time = 5.29 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.39

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (Ca^2 + 2Bba)}{d} - \frac{\ln(\tan(c + dx) - i) (-C + B i) (-b + a i)^2}{2d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (C + B i) (b + a i)^2}{2d} - \frac{Ba^2 \cot(c + dx)}{d}$$

input

```
int(cot(c + d*x)^3*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^2,x)
```

output

$$\frac{(\log(\tan(c + dx)) * (C * a^2 + 2 * B * a * b)) / d - (\log(\tan(c + dx)) - 1i) * (B * 1i - C) * (a * 1i - b)^2 / (2 * d) + (\log(\tan(c + dx)) + 1i) * (B * 1i + C) * (a * 1i + b)^2 / (2 * d) - (B * a^2 * \cot(c + dx)) / d}$$
Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 232, normalized size of antiderivative = 3.22

$$\int \cot^3(c + dx) (a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-\cos(dx + c) a^2 b - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) \sin(dx + c) a^2 c - 2 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) \sin(dx + c) a b^2}{\sin(c + dx) * d}$$

input

$$\text{int}(\cot(dx+c)^3 * (a+b*\tan(dx+c))^2 * (B*\tan(dx+c)+C*\tan(dx+c)^2), x)$$

output

$$\frac{(-\cos(c + dx) * a^2 * b - \log(\tan((c + dx)/2)^2 + 1) * \sin(c + dx) * a^2 * c - 2 * \log(\tan((c + dx)/2)^2 + 1) * \sin(c + dx) * a * b^2 + \log(\tan((c + dx)/2)^2 + 1) * \sin(c + dx) * b^2 * c - \log(\tan((c + dx)/2) - 1) * \sin(c + dx) * b * a * c - \log(\tan((c + dx)/2) + 1) * \sin(c + dx) * b^2 * c + \log(\tan((c + dx)/2)) * \sin(c + dx) * a^2 * c + 2 * \log(\tan((c + dx)/2)) * \sin(c + dx) * a * b^2 - \sin(c + dx) * a^2 * b * dx + 2 * \sin(c + dx) * a * b * c * dx + \sin(c + dx) * b^3 * dx) / (\sin(c + dx) * d)}$$

3.14 $\int \cot^4(c+dx)(a+b \tan(c+dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	195
Mathematica [C] (verified)	195
Rubi [A] (verified)	196
Maple [A] (verified)	199
Fricas [A] (verification not implemented)	200
Sympy [B] (verification not implemented)	200
Maxima [A] (verification not implemented)	201
Giac [A] (verification not implemented)	201
Mupad [B] (verification not implemented)	202
Reduce [B] (verification not implemented)	203

Optimal result

Integrand size = 40, antiderivative size = 88

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (b^2C - a(2bB + aC)) x - \frac{a(2bB + aC) \cot(c + dx)}{d}$$

$$- \frac{a^2B \cot^2(c + dx)}{2d} - \frac{(a^2B - b^2B - 2abC) \log(\sin(c + dx))}{d}$$

output

```
(C*b^2-a*(2*B*b+C*a))*x-a*(2*B*b+C*a)*cot(d*x+c)/d-1/2*a^2*B*cot(d*x+c)^2/d-(B*a^2-B*b^2-2*C*a*b)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.40

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-2a(2bB + aC) \cot(c + dx) - a^2B \cot^2(c + dx) + (a + ib)^2(B + iC) \log(i - \tan(c + dx)) - 2(a^2B - b^2B - 2abC) \log(\sin(c + dx))}{2d}$$

input

```
Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(-2*a*(2*b*B + a*C)*Cot[c + d*x] - a^2*B*Cot[c + d*x]^2 + (a + I*b)^2*(B + I*C)*Log[I - Tan[c + d*x]] - 2*(a^2*B - b^2*B - 2*a*b*C)*Log[Tan[c + d*x]] + (a - I*b)^2*(B - I*C)*Log[I + Tan[c + d*x]])/(2*d)
```

Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 4115, 3042, 4087, 3042, 4111, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx))}{\tan^4(c + dx)} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \cot^3(c + dx)(a + b \tan(c + dx))^2 (B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2 (B + C \tan(c + dx))}{\tan^3(c + dx)} dx \\
 & \quad \downarrow \text{4087} \\
 & \int \cot^2(c + dx) (b^2 C \tan^2(c + dx) - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC)) dx - \\
 & \quad \frac{a^2 B \cot^2(c + dx)}{2d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{b^2 C \tan(c+dx)^2 - (Ba^2 - 2bCa - b^2 B) \tan(c+dx) + a(2bB + aC)}{\tan(c+dx)^2} dx - \frac{a^2 B \cot^2(c+dx)}{2d} \\
& \quad \downarrow 4111 \\
& \int -\cot(c+dx) (Ba^2 - 2bCa - b^2 B - (b^2 C - a(2bB + aC)) \tan(c+dx)) dx - \\
& \quad \frac{a^2 B \cot^2(c+dx)}{2d} - \frac{a(aC + 2bB) \cot(c+dx)}{d} \\
& \quad \downarrow 25 \\
& - \int \cot(c+dx) (Ba^2 - 2bCa - b^2 B + (Ca^2 + 2bBa - b^2 C) \tan(c+dx)) dx - \\
& \quad \frac{a^2 B \cot^2(c+dx)}{2d} - \frac{a(aC + 2bB) \cot(c+dx)}{d} \\
& \quad \downarrow 3042 \\
& - \int \frac{Ba^2 - 2bCa - b^2 B + (Ca^2 + 2bBa - b^2 C) \tan(c+dx)}{\tan(c+dx)} dx - \frac{a^2 B \cot^2(c+dx)}{2d} - \\
& \quad \frac{a(aC + 2bB) \cot(c+dx)}{d} \\
& \quad \downarrow 4014 \\
& -(a^2 B - 2abC - b^2 B) \int \cot(c+dx) dx - x(a^2 C + 2abB - b^2 C) - \frac{a^2 B \cot^2(c+dx)}{2d} - \\
& \quad \frac{a(aC + 2bB) \cot(c+dx)}{d} \\
& \quad \downarrow 3042 \\
& -(a^2 B - 2abC - b^2 B) \int -\tan\left(c+dx + \frac{\pi}{2}\right) dx - x(a^2 C + 2abB - b^2 C) - \\
& \quad \frac{a^2 B \cot^2(c+dx)}{2d} - \frac{a(aC + 2bB) \cot(c+dx)}{d} \\
& \quad \downarrow 25 \\
& (a^2 B - 2abC - b^2 B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - x(a^2 C + 2abB - b^2 C) - \\
& \quad \frac{a^2 B \cot^2(c+dx)}{2d} - \frac{a(aC + 2bB) \cot(c+dx)}{d} \\
& \quad \downarrow 3956 \\
& \frac{(a^2 B - 2abC - b^2 B) \log(-\sin(c+dx))}{d} - x(a^2 C + 2abB - b^2 C) - \frac{a^2 B \cot^2(c+dx)}{2d} - \\
& \quad \frac{a(aC + 2bB) \cot(c+dx)}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `-((2*a*b*B + a^2*C - b^2*C)*x) - (a*(2*b*B + a*C)*Cot[c + d*x])/d - (a^2*B*Cot[c + d*x]^2)/(2*d) - ((a^2*B - b^2*B - 2*a*b*C)*Log[-Sin[c + d*x]])/d`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(x_), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4087 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e
_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{B b^2 \ln(\sin(dx+c)) + C b^2(dx+c) + 2Bab(-\cot(dx+c) - dx - c) + 2Cab \ln(\sin(dx+c)) + B a^2 \left(-\frac{\cot(dx+c)^2}{2} - \ln(\sin(dx+c)) \right)}{d}$
default	$\frac{B b^2 \ln(\sin(dx+c)) + C b^2(dx+c) + 2Bab(-\cot(dx+c) - dx - c) + 2Cab \ln(\sin(dx+c)) + B a^2 \left(-\frac{\cot(dx+c)^2}{2} - \ln(\sin(dx+c)) \right)}{d}$
parallelrisc	$\frac{(B a^2 - B b^2 - 2Cab) \ln(\sec(dx+c)^2) + (-2B a^2 + 2B b^2 + 4Cab) \ln(\tan(dx+c)) - B \cot(dx+c)^2 a^2 + (-4Bab - 2C a^2) \cot(dx+c)}{2d}$
norman	$\frac{(-2Bab - C a^2 + C b^2) x \tan(dx+c)^3 - \frac{B a^2 \tan(dx+c)}{2d} - \frac{a(2Bb + Ca) \tan(dx+c)^2}{d}}{\tan(dx+c)^3} - \frac{(B a^2 - B b^2 - 2Cab) \ln(\tan(dx+c))}{d}$
risc	$-i B b^2 x - \frac{4iCab c}{d} - \frac{2ia(2Bb e^{2i(dx+c)} + Ca e^{2i(dx+c)} + iBa e^{2i(dx+c)} - 2Bb - Ca)}{d(e^{2i(dx+c)} - 1)^2} - 2Babx - C a^2 x +$

input

```
int(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method
=_RETURNVERBOSE)
```

output

```
1/d*(B*b^2*ln(sin(d*x+c))+C*b^2*(d*x+c)+2*B*a*b*(-cot(d*x+c)-d*x-c)+2*C*a*
b*ln(sin(d*x+c))+B*a^2*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+C*a^2*(-cot(d*x+
c)-d*x-c))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.39

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx =$$

$$\frac{(Ba^2 - 2Cab - Bb^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 + Ba^2 + (Ba^2 + 2(Ca^2 + 2Bab - Cb^2)dx) \tan(dx+c)}{2d \tan(dx+c)^2}$$

input

```
integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="fricas")
```

output

```
-1/2*((B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*t
an(d*x + c)^2 + B*a^2 + (B*a^2 + 2*(C*a^2 + 2*B*a*b - C*b^2)*d*x)*tan(d*x
+ c)^2 + 2*(C*a^2 + 2*B*a*b)*tan(d*x + c))/(d*tan(d*x + c)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 206 vs. 2(78) = 156.

Time = 1.94 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.34

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^2 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a+b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \cot^4(c) \\ \text{NaN} \\ \frac{Ba^2 \log(\tan^2(c+dx)+1)}{2d} - \frac{Ba^2 \log(\tan(c+dx))}{d} - \frac{Ba^2}{2d \tan^2(c+dx)} - 2Babx - \frac{2Bab}{d \tan(c+dx)} - \frac{Bb^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{Bb^2}{2d} \end{cases}$$

input

```
integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)
,x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*tan(c)**2)*cot(c)**4, Eq(d, 0)), (nan, Eq(c, -d*x)), (B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - B*a**2*log(tan(c + d*x))/d - B*a**2/(2*d*tan(c + d*x)**2) - 2*B*a*b*x - 2*B*a*b/(d*tan(c + d*x)) - B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**2*log(tan(c + d*x))/d - C*a**2*x - C*a**2/(d*tan(c + d*x)) - C*a*b*log(tan(c + d*x)**2 + 1)/d + 2*C*a*b*log(tan(c + d*x))/d + C*b**2*x, True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.36

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx = \frac{2(Ca^2 + 2Bab - Cb^2)(dx + c) - (Ba^2 - 2Cab - Bb^2) \log(\tan(dx + c)^2 + 1) + 2(Ba^2 - 2Cab - Bb^2) \log(|\tan(dx + c)|)}{2d}$$

input

```
integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

```
-1/2*(2*(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c) - (B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1) + 2*(B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)) + (B*a^2 + 2*(C*a^2 + 2*B*a*b)*tan(d*x + c))/tan(d*x + c)^2)/d
```

Giac [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.47

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx = -\frac{(Ca^2 + 2Bab - Cb^2)(dx + c)}{d} + \frac{(Ba^2 - 2Cab - Bb^2) \log(\tan(dx + c)^2 + 1)}{2d} - \frac{(Ba^2 - 2Cab - Bb^2) \log(|\tan(dx + c)|)}{d} - \frac{Ba^2 + 2(Ca^2 + 2Bab) \tan(dx + c)}{2d \tan(dx + c)^2}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")`

output `-(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c)/d + 1/2*(B*a^2 - 2*C*a*b - B*b^2)*log
(tan(d*x + c)^2 + 1)/d - (B*a^2 - 2*C*a*b - B*b^2)*log(abs(tan(d*x + c)))/
d - 1/2*(B*a^2 + 2*(C*a^2 + 2*B*a*b)*tan(d*x + c))/(d*tan(d*x + c)^2)`

Mupad [B] (verification not implemented)

Time = 5.30 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.44

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (-B a^2 + 2 C a b + B b^2)}{d}$$

$$- \frac{\cot(c + dx)^2 \left(\frac{B a^2}{2} + \tan(c + dx) (C a^2 + 2 B b a) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) + 1i) (B - C 1i) (b + a 1i)^2}{2 d}$$

$$- \frac{\ln(\tan(c + dx) - 1i) (B + C 1i) (-b + a 1i)^2}{2 d}$$

input `int(cot(c + d*x)^4*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
)^2,x)`

output `(log(tan(c + d*x))*(B*b^2 - B*a^2 + 2*C*a*b))/d - (cot(c + d*x)^2*((B*a^2)
/2 + tan(c + d*x)*(C*a^2 + 2*B*a*b)))/d - (log(tan(c + d*x) + 1i)*(B - C*1
i)*(a*1i + b)^2)/(2*d) - (log(tan(c + d*x) - 1i)*(B + C*1i)*(a*1i - b)^2)/
(2*d)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 269, normalized size of antiderivative = 3.06

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-4 \cos(dx + c) \sin(dx + c) a^2 c - 8 \cos(dx + c) \sin(dx + c) a b^2 + 4 \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) \sin(dx + c)}{\sin^2(dx + c)}$$

input

```
int(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
( - 4*cos(c + d*x)*sin(c + d*x)*a**2*c - 8*cos(c + d*x)*sin(c + d*x)*a*b**
2 + 4*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**2*b - 8*log(tan((c +
d*x)/2)**2 + 1)*sin(c + d*x)**2*a*b*c - 4*log(tan((c + d*x)/2)**2 + 1)*si
n(c + d*x)**2*b**3 - 4*log(tan((c + d*x)/2))*sin(c + d*x)**2*a**2*b + 8*lo
g(tan((c + d*x)/2))*sin(c + d*x)**2*a*b*c + 4*log(tan((c + d*x)/2))*sin(c
+ d*x)**2*b**3 + sin(c + d*x)**2*a**2*b - 4*sin(c + d*x)**2*a**2*c*d*x - 8
*sin(c + d*x)**2*a*b**2*d*x + 4*sin(c + d*x)**2*b**2*c*d*x - 2*a**2*b)/(4*
sin(c + d*x)**2*d)
```


3.15 $\int \cot^5(c+dx)(a+b \tan(c+dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	204
Mathematica [C] (verified)	205
Rubi [A] (verified)	205
Maple [A] (verified)	209
Fricas [A] (verification not implemented)	210
Sympy [B] (verification not implemented)	210
Maxima [A] (verification not implemented)	211
Giac [A] (verification not implemented)	211
Mupad [B] (verification not implemented)	212
Reduce [B] (verification not implemented)	213

Optimal result

Integrand size = 40, antiderivative size = 118

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (a^2B - b^2B - 2abC) x + \frac{(a^2B - b^2B - 2abC) \cot(c + dx)}{d}$$

$$- \frac{a(2bB + aC) \cot^2(c + dx)}{2d} - \frac{a^2B \cot^3(c + dx)}{3d}$$

$$+ \frac{(b^2C - a(2bB + aC)) \log(\sin(c + dx))}{d}$$

output

```
(B*a^2-B*b^2-2*C*a*b)*x+(B*a^2-B*b^2-2*C*a*b)*cot(d*x+c)/d-1/2*a*(2*B*b+C*a)*cot(d*x+c)^2/d-1/3*a^2*B*cot(d*x+c)^3/d+(C*b^2-a*(2*B*b+C*a))*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.80 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.29

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6(a^2B - b^2B - 2abC) \cot(c + dx) - 3a(2bB + aC) \cot^2(c + dx) - 2a^2B \cot^3(c + dx) + 3(a + ib)^2(-iB$$

input

```
Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(6*(a^2*B - b^2*B - 2*a*b*C)*Cot[c + d*x] - 3*a*(2*b*B + a*C)*Cot[c + d*x]^2 - 2*a^2*B*Cot[c + d*x]^3 + 3*(a + I*b)^2*((-I)*B + C)*Log[I - Tan[c + d*x]] - 6*(2*a*b*B + a^2*C - b^2*C)*Log[Tan[c + d*x]] + 3*(a - I*b)^2*(I*B + C)*Log[I + Tan[c + d*x]])/(6*d)
```

Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {3042, 4115, 3042, 4087, 3042, 4111, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx))^2}{\tan^5(c + dx)} dx$$

$$\downarrow \text{4115}$$

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2 (B + C \tan(c + dx)) dx$$

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^2 (B + C \tan(c + dx))}{\tan(c + dx)^4} dx \\
& \quad \downarrow 3042 \\
& \int \cot^3(c + dx) (b^2 C \tan^2(c + dx) - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC)) dx - \\
& \quad \frac{a^2 B \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4087 \\
& \int \frac{b^2 C \tan(c + dx)^2 - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC)}{\tan(c + dx)^3} dx - \frac{a^2 B \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& \int \frac{b^2 C \tan(c + dx)^2 - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC)}{\tan(c + dx)^3} dx - \frac{a^2 B \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4111 \\
& \int -\cot^2(c + dx) (Ba^2 - 2bCa - b^2 B - (b^2 C - a(2bB + aC)) \tan(c + dx)) dx - \\
& \quad \frac{a^2 B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \quad \downarrow 25 \\
& - \int \cot^2(c + dx) (Ba^2 - 2bCa - b^2 B + (Ca^2 + 2bBa - b^2 C) \tan(c + dx)) dx - \\
& \quad \frac{a^2 B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \quad \downarrow 3042 \\
& - \int \frac{Ba^2 - 2bCa - b^2 B + (Ca^2 + 2bBa - b^2 C) \tan(c + dx)}{\tan(c + dx)^2} dx - \frac{a^2 B \cot^3(c + dx)}{3d} - \\
& \quad \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \quad \downarrow 4012 \\
& - \int \cot(c + dx) (Ca^2 + 2bBa - b^2 C - (Ba^2 - 2bCa - b^2 B) \tan(c + dx)) dx + \\
& \quad \frac{(a^2 B - 2abC - b^2 B) \cot(c + dx)}{d} - \frac{a^2 B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \quad \downarrow 3042 \\
& - \int \frac{Ca^2 + 2bBa - b^2 C - (Ba^2 - 2bCa - b^2 B) \tan(c + dx)}{\tan(c + dx)} dx + \\
& \quad \frac{(a^2 B - 2abC - b^2 B) \cot(c + dx)}{d} - \frac{a^2 B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d}
\end{aligned}$$

$$\begin{aligned}
& \downarrow 4014 \\
& -(a^2C + 2abB - b^2C) \int \cot(c + dx) dx + \frac{(a^2B - 2abC - b^2B) \cot(c + dx)}{d} + \\
& \quad x(a^2B - 2abC - b^2B) - \frac{a^2B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \downarrow 3042 \\
& -(a^2C + 2abB - b^2C) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + \frac{(a^2B - 2abC - b^2B) \cot(c + dx)}{d} + \\
& \quad x(a^2B - 2abC - b^2B) - \frac{a^2B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \downarrow 25 \\
& (a^2C + 2abB - b^2C) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{(a^2B - 2abC - b^2B) \cot(c + dx)}{d} + \\
& \quad x(a^2B - 2abC - b^2B) - \frac{a^2B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d} \\
& \downarrow 3956 \\
& \frac{(a^2B - 2abC - b^2B) \cot(c + dx)}{d} - \frac{(a^2C + 2abB - b^2C) \log(-\sin(c + dx))}{d} + \\
& \quad x(a^2B - 2abC - b^2B) - \frac{a^2B \cot^3(c + dx)}{3d} - \frac{a(aC + 2bB) \cot^2(c + dx)}{2d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(a^2*B - b^2*B - 2*a*b*C)*x + ((a^2*B - b^2*B - 2*a*b*C)*Cot[c + d*x])/d - (a*(2*b*B + a*C)*Cot[c + d*x]^2)/(2*d) - (a^2*B*Cot[c + d*x]^3)/(3*d) - ((2*a*b*B + a^2*C - b^2*C)*Log[-Sin[c + d*x]])/d
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4012 $\text{Int}[\left((a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]\right)^{(m_.)}\left((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]\right), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)\left((a + b*\text{Tan}[e + f*x])^{(m + 1)} / (f*(m + 1)*(a^2 + b^2))\right), x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m + 1)}*\text{Simp}[a*c + b*d - (b*c - a*d)*\text{Tan}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{LtQ}[m, -1]$

rule 4014 $\text{Int}[\left((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]\right) / \left((a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]\right), x_Symbol] \rightarrow \text{Simp}[(a*c + b*d)*(x/(a^2 + b^2)), x] + \text{Simp}[(b*c - a*d)/(a^2 + b^2) \text{Int}[(b - a*\text{Tan}[e + f*x]) / (a + b*\text{Tan}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[a*c + b*d, 0]$

rule 4087 $\text{Int}[\left((a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]\right)^2\left((A_.) + (B_.)\tan[(e_.) + (f_.)(x_)]\right)\left((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]\right)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[\left(- (B*c - A*d)\right)*(b*c - a*d)^2\left((c + d*\text{Tan}[e + f*x])^{(n + 1)} / (f*d^2*(n + 1)*(c^2 + d^2)\right), x] + \text{Simp}[1/(d*(c^2 + d^2)) \text{Int}[(c + d*\text{Tan}[e + f*x])^{(n + 1)}*\text{Simp}[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*\text{Tan}[e + f*x] + b^2*B*(c^2 + d^2)*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{LtQ}[n, -1]$

rule 4111 $\text{Int}[\left((a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]\right)^{(m_.)}\left((A_.) + (B_.)\tan[(e_.) + (f_.)(x_)] + (C_.)\tan[(e_.) + (f_.)(x_)]^2\right), x_Symbol] \rightarrow \text{Simp}[(A*b^2 - a*b*B + a^2*C)\left((a + b*\text{Tan}[e + f*x])^{(m + 1)} / (b*f*(m + 1)*(a^2 + b^2)\right), x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m + 1)}*\text{Simp}[b*B + a*(A - C) - (A*b - a*B - b*C)*\text{Tan}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{LtQ}[m, -1] \&\& \text{NeQ}[a^2 + b^2, 0]$

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.15

method	result
derivativedivides	$\frac{B b^2(-\cot(dx+c)-dx-c)+C b^2 \ln(\sin(dx+c))+2Bab\left(-\frac{\cot(dx+c)^2}{2}-\ln(\sin(dx+c))\right)+2Cab(-\cot(dx+c)-dx-c)+1}{d}$
default	$\frac{B b^2(-\cot(dx+c)-dx-c)+C b^2 \ln(\sin(dx+c))+2Bab\left(-\frac{\cot(dx+c)^2}{2}-\ln(\sin(dx+c))\right)+2Cab(-\cot(dx+c)-dx-c)+1}{d}$
parallelrisch	$\frac{3 \ln(\sec(dx+c)^2)(2Bab+C a^2-C b^2)+6(-2Bab-C a^2+C b^2) \ln(\tan(dx+c))-2B \cot(dx+c)^3 a^2+3(-2Bab-C a^2) c}{6d}$
norman	$\frac{\left(\frac{B a^2-B b^2-2Cab}{d}\right) \tan(dx+c)^3+(B a^2-B b^2-2Cab) x \tan(dx+c)^4-\frac{B a^2 \tan(dx+c)}{3d}-\frac{a(2Bb+Ca) \tan(dx+c)^2}{2d}}{\tan(dx+c)^4} - (2Bab)$
risch	$B a^2 x - B b^2 x - 2C a b x + i C a^2 x - i C b^2 x + \frac{2i C a^2 c}{d} + \frac{4i B a b c}{d} - \frac{2i C b^2 c}{d} - \frac{2i(6i B a b e^{4i(dx+c)})}{d}$

input

```
int(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method =_RETURNVERBOSE)
```

output

```
1/d*(B*b^2*(-cot(d*x+c)-d*x-c)+C*b^2*ln(sin(d*x+c))+2*B*a*b*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+2*C*a*b*(-cot(d*x+c)-d*x-c)+B*a^2*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c)+C*a^2*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c))))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.33

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{3(Ca^2 + 2Bab - Cb^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 + 3(Ca^2 + 2Bab - 2(Ba^2 - 2Cab - Bb^2)dx - 6d \tan(dx -$$

input

```
integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="fricas")
```

output

```
-1/6*(3*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))
*tan(d*x + c)^3 + 3*(C*a^2 + 2*B*a*b - 2*(B*a^2 - 2*C*a*b - B*b^2)*d*x)*ta
n(d*x + c)^3 + 2*B*a^2 - 6*(B*a^2 - 2*C*a*b - B*b^2)*tan(d*x + c)^2 + 3*(C
*a^2 + 2*B*a*b)*tan(d*x + c))/(d*tan(d*x + c)^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(107) = 214.

Time = 2.48 (sec) , antiderivative size = 252, normalized size of antiderivative = 2.14

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \cot^5(c) \\ \text{NaN} \\ Ba^2x + \frac{Ba^2}{d \tan(c+dx)} - \frac{Ba^2}{3d \tan^3(c+dx)} + \frac{Bab \log(\tan^2(c+dx)+1)}{d} - \frac{2Bab \log(\tan(c+dx))}{d} - \frac{Bab}{d \tan^2(c+dx)} - Bb^2x - \frac{1}{d \tan} \end{cases}$$

input

```
integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)
,x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*tan(c)**2)*cot(c)**5, Eq(d, 0)), (nan, Eq(c, -d*x)), (B*a**2*x + B*a**2/(d*tan(c + d*x)) - B*a**2/(3*d*tan(c + d*x)**3) + B*a*b*log(tan(c + d*x)**2 + 1)/d - 2*B*a*b*log(tan(c + d*x))/d - B*a*b/(d*tan(c + d*x)**2) - B*b**2*x - B*b**2/(d*tan(c + d*x)) + C*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - C*a**2*log(tan(c + d*x))/d - C*a**2/(2*d*tan(c + d*x)**2) - 2*C*a*b*x - 2*C*a*b/(d*tan(c + d*x)) - C*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + C*b**2*log(tan(c + d*x))/d, True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.26

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6(Ba^2 - 2Cab - Bb^2)(dx + c) + 3(Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1) - 6(Ca^2 + 2Bab - Cb^2)}{6d}$$

input

```
integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
1/6*(6*(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c) + 3*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1) - 6*(C*a^2 + 2*B*a*b - C*b^2)*log(tan(d*x + c)) - (2*B*a^2 - 6*(B*a^2 - 2*C*a*b - B*b^2)*tan(d*x + c)^2 + 3*(C*a^2 + 2*B*a*b)*tan(d*x + c))/tan(d*x + c)^3)/d
```

Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.32

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ba^2 - 2Cab - Bb^2)(dx + c)}{d} + \frac{(Ca^2 + 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$- \frac{(Ca^2 + 2Bab - Cb^2) \log(|\tan(dx + c)|)}{d}$$

$$- \frac{2Ba^2 - 6(Ba^2 - 2Cab - Bb^2) \tan(dx + c)^2 + 3(Ca^2 + 2Bab) \tan(dx + c)}{6d \tan(dx + c)^3}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")`

output `(B*a^2 - 2*C*a*b - B*b^2)*(d*x + c)/d + 1/2*(C*a^2 + 2*B*a*b - C*b^2)*log(
tan(d*x + c)^2 + 1)/d - (C*a^2 + 2*B*a*b - C*b^2)*log(abs(tan(d*x + c)))/d
- 1/6*(2*B*a^2 - 6*(B*a^2 - 2*C*a*b - B*b^2)*tan(d*x + c)^2 + 3*(C*a^2 +
2*B*a*b)*tan(d*x + c))/(d*tan(d*x + c)^3)`

Mupad [B] (verification not implemented)

Time = 5.40 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.32

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{\cot(c + dx)^3 \left(\frac{B a^2}{3} + \tan(c + dx)^2 (-B a^2 + 2 C a b + B b^2) + \tan(c + dx) \left(\frac{C a^2}{2} + B b a \right) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx)) (C a^2 + 2 B a b - C b^2)}{d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (-C + B i) (-b + a i)^2}{2 d}$$

$$- \frac{\ln(\tan(c + dx) + i) (C + B i) (b + a i)^2}{2 d}$$

input `int(cot(c + d*x)^5*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
)^2,x)`

output `(log(tan(c + d*x) - 1i)*(B*1i - C)*(a*1i - b)^2)/(2*d) - (log(tan(c + d*x)
)*(C*a^2 - C*b^2 + 2*B*a*b))/d - (cot(c + d*x)^3*((B*a^2)/3 + tan(c + d*x)
^2*(B*b^2 - B*a^2 + 2*C*a*b) + tan(c + d*x)*((C*a^2)/2 + B*a*b))/d - (log
(tan(c + d*x) + 1i)*(B*1i + C)*(a*1i + b)^2)/(2*d)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 338, normalized size of antiderivative = 2.86

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{16 \cos(dx + c) \sin(dx + c)^2 a^2 b - 24 \cos(dx + c) \sin(dx + c)^2 abc - 12 \cos(dx + c) \sin(dx + c)^2 b^3 - 4 \dots}{\dots}$$

input

```
int(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
(16*cos(c + d*x)*sin(c + d*x)**2*a**2*b - 24*cos(c + d*x)*sin(c + d*x)**2*
a*b*c - 12*cos(c + d*x)*sin(c + d*x)**2*b**3 - 4*cos(c + d*x)*a**2*b + 12*
log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**3*a**2*c + 24*log(tan((c + d*x)
/2)**2 + 1)*sin(c + d*x)**3*a*b**2 - 12*log(tan((c + d*x)/2)**2 + 1)*sin(c
+ d*x)**3*b**2*c - 12*log(tan((c + d*x)/2))*sin(c + d*x)**3*a**2*c - 24*log
og(tan((c + d*x)/2))*sin(c + d*x)**3*a*b**2 + 12*log(tan((c + d*x)/2))*sin
(c + d*x)**3*b**2*c + 12*sin(c + d*x)**3*a**2*b*d*x + 3*sin(c + d*x)**3*a*
*2*c + 6*sin(c + d*x)**3*a*b**2 - 24*sin(c + d*x)**3*a*b*c*d*x - 12*sin(c
+ d*x)**3*b**3*d*x - 6*sin(c + d*x)*a**2*c - 12*sin(c + d*x)*a*b**2)/(12*s
in(c + d*x)**3*d)
```

3.16 $\int \cot^6(c+dx)(a+b \tan(c+dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	214
Mathematica [C] (verified)	215
Rubi [A] (verified)	215
Maple [A] (verified)	220
Fricas [A] (verification not implemented)	220
Sympy [B] (verification not implemented)	221
Maxima [A] (verification not implemented)	222
Giac [A] (verification not implemented)	222
Mupad [B] (verification not implemented)	223
Reduce [B] (verification not implemented)	224

Optimal result

Integrand size = 40, antiderivative size = 151

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (2abB + a^2C - b^2C) x - \frac{(b^2C - a(2bB + aC)) \cot(c + dx)}{d}$$

$$+ \frac{(a^2B - b^2B - 2abC) \cot^2(c + dx)}{2d} - \frac{a(2bB + aC) \cot^3(c + dx)}{3d}$$

$$- \frac{a^2B \cot^4(c + dx)}{4d} + \frac{(a^2B - b^2B - 2abC) \log(\sin(c + dx))}{d}$$

output

```
(2*B*a*b+C*a^2-C*b^2)*x-(C*b^2-a*(2*B*b+C*a))*cot(d*x+c)/d+1/2*(B*a^2-B*b^2-2*C*a*b)*cot(d*x+c)^2/d-1/3*a*(2*B*b+C*a)*cot(d*x+c)^3/d-1/4*a^2*B*cot(d*x+c)^4/d+(B*a^2-B*b^2-2*C*a*b)*ln(sin(d*x+c))/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.99 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.19

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{12(2abB + a^2C - b^2C) \cot(c + dx) + 6(a^2B - b^2B - 2abC) \cot^2(c + dx) - 4a(2bB + aC) \cot^3(c + dx)}{12d}$$

input

```
Integrate[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(12*(2*a*b*B + a^2*C - b^2*C)*Cot[c + d*x] + 6*(a^2*B - b^2*B - 2*a*b*C)*Cot[c + d*x]^2 - 4*a*(2*b*B + a*C)*Cot[c + d*x]^3 - 3*a^2*B*Cot[c + d*x]^4 - 6*((a + I*b)^2*(B + I*C)*Log[I - Tan[c + d*x]] + (-2*a^2*B + 2*b^2*B + 4*a*b*C)*Log[Tan[c + d*x]] + (a - I*b)^2*(B - I*C)*Log[I + Tan[c + d*x]]))/(12*d)
```

Rubi [A] (verified)

Time = 1.88 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.01, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.425$, Rules used = {3042, 4115, 3042, 4087, 3042, 4111, 25, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)^6} dx$$

$$\downarrow 4115$$

$$\begin{aligned}
& \int \cot^5(c+dx)(a+b \tan(c+dx))^2(B+C \tan(c+dx))dx \\
& \quad \downarrow 3042 \\
& \int \frac{(a+b \tan(c+dx))^2(B+C \tan(c+dx))}{\tan(c+dx)^5} dx \\
& \quad \downarrow 4087 \\
& \int \cot^4(c+dx) (b^2C \tan^2(c+dx) - (Ba^2 - 2bCa - b^2B) \tan(c+dx) + a(2bB + aC)) dx - \\
& \quad \quad \quad \frac{a^2B \cot^4(c+dx)}{4d} \\
& \quad \downarrow 3042 \\
& \int \frac{b^2C \tan(c+dx)^2 - (Ba^2 - 2bCa - b^2B) \tan(c+dx) + a(2bB + aC)}{\tan(c+dx)^4} dx - \frac{a^2B \cot^4(c+dx)}{4d} \\
& \quad \downarrow 4111 \\
& \int -\cot^3(c+dx) (Ba^2 - 2bCa - b^2B - (b^2C - a(2bB + aC)) \tan(c+dx)) dx - \\
& \quad \quad \quad \frac{a^2B \cot^4(c+dx)}{4d} - \frac{a(aC + 2bB) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 25 \\
& - \int \cot^3(c+dx) (Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c+dx)) dx - \\
& \quad \quad \quad \frac{a^2B \cot^4(c+dx)}{4d} - \frac{a(aC + 2bB) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& - \int \frac{Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c+dx)}{\tan(c+dx)^3} dx - \frac{a^2B \cot^4(c+dx)}{4d} - \\
& \quad \quad \quad \frac{a(aC + 2bB) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 4012 \\
& - \int \cot^2(c+dx) (Ca^2 + 2bBa - b^2C - (Ba^2 - 2bCa - b^2B) \tan(c+dx)) dx + \\
& \quad \quad \quad \frac{(a^2B - 2abC - b^2B) \cot^2(c+dx)}{2d} - \frac{a^2B \cot^4(c+dx)}{4d} - \frac{a(aC + 2bB) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& - \int \frac{Ca^2 + 2bBa - b^2C - (Ba^2 - 2bCa - b^2B) \tan(c + dx)}{\tan(c + dx)^2} dx + \\
& \frac{(a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} - \frac{a^2B \cot^4(c + dx)}{4d} - \frac{a(aC + 2bB) \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4012 \\
& - \int -\cot(c + dx) (Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c + dx)) dx + \\
& \frac{(a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \quad \frac{a(aC + 2bB) \cot^3(c + dx)}{3d} \\
& \quad \downarrow 25 \\
& \int \cot(c + dx) (Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c + dx)) dx + \\
& \frac{(a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \quad \frac{a(aC + 2bB) \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& \int \frac{Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c + dx)}{\tan(c + dx)} dx + \\
& \frac{(a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \quad \frac{a(aC + 2bB) \cot^3(c + dx)}{3d} \\
& \quad \downarrow 4014 \\
& \frac{(a^2B - 2abC - b^2B) \int \cot(c + dx) dx + (a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \\
& \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} + x(a^2C + 2abB - b^2C) - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \quad \frac{a(aC + 2bB) \cot^3(c + dx)}{3d} \\
& \quad \downarrow 3042 \\
& \frac{(a^2B - 2abC - b^2B) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + (a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \\
& \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} + x(a^2C + 2abB - b^2C) - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \quad \frac{a(aC + 2bB) \cot^3(c + dx)}{3d}
\end{aligned}$$

$$\begin{aligned}
& \downarrow 25 \\
& -(a^2B - 2abC - b^2B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{(a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \\
& \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} + x(a^2C + 2abB - b^2C) - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \frac{a(aC + 2bB) \cot^3(c + dx)}{3d} \\
& \downarrow 3956 \\
& \frac{(a^2B - 2abC - b^2B) \cot^2(c + dx)}{2d} + \frac{(a^2C + 2abB - b^2C) \cot(c + dx)}{d} + \\
& \frac{(a^2B - 2abC - b^2B) \log(-\sin(c + dx))}{d} + x(a^2C + 2abB - b^2C) - \frac{a^2B \cot^4(c + dx)}{4d} - \\
& \frac{a(aC + 2bB) \cot^3(c + dx)}{3d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(2*a*b*B + a^2*C - b^2*C)*x + ((2*a*b*B + a^2*C - b^2*C)*Cot[c + d*x])/d +
((a^2*B - b^2*B - 2*a*b*C)*Cot[c + d*x]^2)/(2*d) - (a*(2*b*B + a*C)*Cot[c
+ d*x]^3)/(3*d) - (a^2*B*Cot[c + d*x]^4)/(4*d) + ((a^2*B - b^2*B - 2*a*b*
C)*Log[-Sin[c + d*x]])/d
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4012

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

rule 4087

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f
_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[
(-(B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(
c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1
)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*
c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2
)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b
*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```


Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.07

method	result
derivativedivides	$\frac{B b^2 \left(-\frac{\cot(dx+c)^2}{2} - \ln(\sin(dx+c)) \right) + C b^2 (-\cot(dx+c) - dx - c) + 2Bab \left(-\frac{\cot(dx+c)^3}{3} + \cot(dx+c) + dx + c \right) + 2Cab \left(-\frac{\cot(dx+c)^4}{4} + \frac{2}{3} \cot(dx+c)^2 + dx + c \right)}{1}$
default	$\frac{B b^2 \left(-\frac{\cot(dx+c)^2}{2} - \ln(\sin(dx+c)) \right) + C b^2 (-\cot(dx+c) - dx - c) + 2Bab \left(-\frac{\cot(dx+c)^3}{3} + \cot(dx+c) + dx + c \right) + 2Cab \left(-\frac{\cot(dx+c)^4}{4} + \frac{2}{3} \cot(dx+c)^2 + dx + c \right)}{1}$
parallelrisc	$\frac{6(-B a^2 + B b^2 + 2Cab) \ln(\sec(dx+c)^2) + 12(B a^2 - B b^2 - 2Cab) \ln(\tan(dx+c)) - 3B \cot(dx+c)^4 a^2 + 4(-2Bab - C a^2) \tan(dx+c)}{12d}$
norman	$\frac{\frac{(2Bab + C a^2 - C b^2) \tan(dx+c)^4}{d} + (2Bab + C a^2 - C b^2) x \tan(dx+c)^5 + \frac{(B a^2 - B b^2 - 2Cab) \tan(dx+c)^3}{2d} - \frac{B a^2 \tan(dx+c)}{4d}}{\tan(dx+c)^5}$
risc	$-\frac{2iB a^2 c}{d} + iB b^2 x + \frac{4iCab c}{d} + 2Babx + C a^2 x - C b^2 x - iB a^2 x + \frac{2iB b^2 c}{d} + 2iCabx + \dots$

input `int(cot(d*x+c)^6*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/d*(B*b^2*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+C*b^2*(-cot(d*x+c)-d*x-c)+2*B*a*b*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c)+2*C*a*b*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+B*a^2*(-1/4*cot(d*x+c)^4+1/2*cot(d*x+c)^2+ln(sin(d*x+c)))+C*a^2*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.26

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6(Ba^2 - 2Cab - Bb^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 + 3(3Ba^2 - 4Cab - 2Bb^2 + 4(Ca^2 + 2Bab - Cb^2)) \tan(dx+c)^3 + \dots}{1}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output

```
1/12*(6*(B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))
*tan(d*x + c)^4 + 3*(3*B*a^2 - 4*C*a*b - 2*B*b^2 + 4*(C*a^2 + 2*B*a*b - C*
b^2)*d*x)*tan(d*x + c)^4 + 12*(C*a^2 + 2*B*a*b - C*b^2)*tan(d*x + c)^3 - 3
*B*a^2 + 6*(B*a^2 - 2*C*a*b - B*b^2)*tan(d*x + c)^2 - 4*(C*a^2 + 2*B*a*b)*
tan(d*x + c))/(d*tan(d*x + c)^4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(136) = 272$.

Time = 3.84 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.01

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^2 (B \tan(c) + C \tan^2(c)) \cot^6(c) \\ \text{NaN} \\ -\frac{Ba^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba^2 \log(\tan(c+dx))}{d} + \frac{Ba^2}{2d \tan^2(c+dx)} - \frac{Ba^2}{4d \tan^4(c+dx)} + 2Babx + \frac{2Bab}{d \tan(c+dx)} - \frac{2Bab}{3d \tan^3(c+dx)} \end{cases}$$

input

```
integrate(cot(d*x+c)**6*(a+b*tan(d*x+c))**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)
,x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**2*(B*tan(c) + C*t
an(c)**2)*cot(c)**6, Eq(d, 0)), (nan, Eq(c, -d*x)), (-B*a**2*log(tan(c + d
*x)**2 + 1)/(2*d) + B*a**2*log(tan(c + d*x))/d + B*a**2/(2*d*tan(c + d*x)*
**2) - B*a**2/(4*d*tan(c + d*x)**4) + 2*B*a*b*x + 2*B*a*b/(d*tan(c + d*x))
- 2*B*a*b/(3*d*tan(c + d*x)**3) + B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) -
B*b**2*log(tan(c + d*x))/d - B*b**2/(2*d*tan(c + d*x)**2) + C*a**2*x + C*a
**2/(d*tan(c + d*x)) - C*a**2/(3*d*tan(c + d*x)**3) + C*a*b*log(tan(c + d*
x)**2 + 1)/d - 2*C*a*b*log(tan(c + d*x))/d - C*a*b/(d*tan(c + d*x)**2) - C
*b**2*x - C*b**2/(d*tan(c + d*x)), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.16

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{12(Ca^2 + 2Bab - Cb^2)(dx + c) - 6(Ba^2 - 2Cab - Bb^2) \log(\tan(dx + c)^2 + 1) + 12(Ba^2 - 2Cab - Bb^2) \log(|\tan(dx + c)|) + 12(Ca^2 + 2Bab - Cb^2) \tan(dx + c)^3 - 3Ba^2 + 6(Ba^2 - 2Cab - Bb^2) \tan(dx + c)^2 - 4(Ca^2 + 2Bab - Cb^2) \tan(dx + c) - 4(Ba^2 - 2Cab - Bb^2) \log(|\tan(dx + c)|)}{12d \tan(dx + c)^4}$$

input

```
integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
1/12*(12*(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c) - 6*(B*a^2 - 2*C*a*b - B*b^2)
*log(tan(d*x + c)^2 + 1) + 12*(B*a^2 - 2*C*a*b - B*b^2)*log(tan(d*x + c))
+ (12*(C*a^2 + 2*B*a*b - C*b^2)*tan(d*x + c)^3 - 3*B*a^2 + 6*(B*a^2 - 2*C*
a*b - B*b^2)*tan(d*x + c)^2 - 4*(C*a^2 + 2*B*a*b)*tan(d*x + c))/tan(d*x +
c)^4)/d
```

Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.21

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ca^2 + 2Bab - Cb^2)(dx + c)}{d} - \frac{(Ba^2 - 2Cab - Bb^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{(Ba^2 - 2Cab - Bb^2) \log(|\tan(dx + c)|)}{d}$$

$$+ \frac{12(Ca^2 + 2Bab - Cb^2) \tan(dx + c)^3 - 3Ba^2 + 6(Ba^2 - 2Cab - Bb^2) \tan(dx + c)^2 - 4(Ca^2 + 2Bab - Cb^2) \tan(dx + c) - 4(Ba^2 - 2Cab - Bb^2) \log(|\tan(dx + c)|)}{12d \tan(dx + c)^4}$$

input

```
integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
(C*a^2 + 2*B*a*b - C*b^2)*(d*x + c)/d - 1/2*(B*a^2 - 2*C*a*b - B*b^2)*log(
tan(d*x + c)^2 + 1)/d + (B*a^2 - 2*C*a*b - B*b^2)*log(abs(tan(d*x + c)))/d
+ 1/12*(12*(C*a^2 + 2*B*a*b - C*b^2)*tan(d*x + c)^3 - 3*B*a^2 + 6*(B*a^2
- 2*C*a*b - B*b^2)*tan(d*x + c)^2 - 4*(C*a^2 + 2*B*a*b)*tan(d*x + c))/(d*t
an(d*x + c)^4)
```

Mupad [B] (verification not implemented)

Time = 5.38 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.21

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx =$$

$$\frac{\cot(c + dx)^4 \left(\frac{B a^2}{4} + \tan(c + dx)^2 \left(-\frac{B a^2}{2} + C a b + \frac{B b^2}{2} \right) - \tan(c + dx)^3 (C a^2 + 2 B a b - C b^2) + \tan(c + dx)^2 (C a b + B b^2) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx)) (-B a^2 + 2 C a b + B b^2)}{d}$$

$$+ \frac{\ln(\tan(c + dx) + 1i) (B - C 1i) (b + a 1i)^2}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) - 1i) (B + C 1i) (-b + a 1i)^2}{2 d}$$

input

```
int(cot(c + d*x)^6*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
)^2,x)
```

output

```
(log(tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b)^2)/(2*d) - (log(tan(c + d*x)
)*(B*b^2 - B*a^2 + 2*C*a*b))/d - (cot(c + d*x)^4*((B*a^2)/4 + tan(c + d*x)
^2*((B*b^2)/2 - (B*a^2)/2 + C*a*b) - tan(c + d*x)^3*(C*a^2 - C*b^2 + 2*B*a
*b) + tan(c + d*x)*((C*a^2)/3 + (2*B*a*b)/3))/d + (log(tan(c + d*x) - 1i)
*(B + C*1i)*(a*1i - b)^2)/(2*d)
```

Reduce [B] (verification not implemented)

Time = 1.01 (sec) , antiderivative size = 396, normalized size of antiderivative = 2.62

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^2 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{128 \cos(dx + c) \sin(dx + c)^3 a^2 c + 256 \cos(dx + c) \sin(dx + c)^3 a b^2 - 96 \cos(dx + c) \sin(dx + c)^3 b^2 c - \dots}{\dots}$$

input

```
int(cot(d*x+c)^6*(a+b*tan(d*x+c))^2*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
(128*cos(c + d*x)*sin(c + d*x)**3*a**2*c + 256*cos(c + d*x)*sin(c + d*x)**
3*a*b**2 - 96*cos(c + d*x)*sin(c + d*x)**3*b**2*c - 32*cos(c + d*x)*sin(c
+ d*x)*a**2*c - 64*cos(c + d*x)*sin(c + d*x)*a*b**2 - 96*log(tan((c + d*x)
/2)**2 + 1)*sin(c + d*x)**4*a**2*b + 192*log(tan((c + d*x)/2)**2 + 1)*sin(
c + d*x)**4*a*b*c + 96*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**4*b**3 +
96*log(tan((c + d*x)/2))*sin(c + d*x)**4*a**2*b - 192*log(tan((c + d*x)/2
))*sin(c + d*x)**4*a*b*c - 96*log(tan((c + d*x)/2))*sin(c + d*x)**4*b**3 -
39*sin(c + d*x)**4*a**2*b + 96*sin(c + d*x)**4*a**2*c*d*x + 192*sin(c + d
*x)**4*a*b**2*d*x + 48*sin(c + d*x)**4*a*b*c + 24*sin(c + d*x)**4*b**3 - 9
6*sin(c + d*x)**4*b**2*c*d*x + 96*sin(c + d*x)**2*a**2*b - 96*sin(c + d*x)
**2*a*b*c - 48*sin(c + d*x)**2*b**3 - 24*a**2*b)/(96*sin(c + d*x)**4*d)
```

3.17 $\int (a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	225
Mathematica [C] (verified)	226
Rubi [A] (verified)	226
Maple [A] (verified)	229
Fricas [A] (verification not implemented)	230
Sympy [B] (verification not implemented)	230
Maxima [A] (verification not implemented)	231
Giac [A] (verification not implemented)	232
Mupad [B] (verification not implemented)	232
Reduce [B] (verification not implemented)	233

Optimal result

Integrand size = 32, antiderivative size = 165

$$\begin{aligned} & \int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= -((3a^2bB - b^3B + a^3C - 3ab^2C) x) \\ & \quad - \frac{(a^3B - 3ab^2B - 3a^2bC + b^3C) \log(\cos(c + dx))}{d} \\ & \quad + \frac{b(a^2B - b^2B - 2abC) \tan(c + dx)}{d} + \frac{(aB - bC)(a + b \tan(c + dx))^2}{2d} \\ & \quad + \frac{B(a + b \tan(c + dx))^3}{3d} + \frac{C(a + b \tan(c + dx))^4}{4bd} \end{aligned}$$

output

```
- (3*B*a^2*b - B*b^3 + C*a^3 - 3*C*a*b^2)*x - (B*a^3 - 3*B*a*b^2 - 3*C*a^2*b + C*b^3)*ln(
cos(d*x+c))/d + b*(B*a^2 - B*b^2 - 2*C*a*b)*tan(d*x+c)/d + 1/2*(B*a - C*b)*(a + b*tan(
d*x+c))^2/d + 1/3*B*(a + b*tan(d*x+c))^3/d + 1/4*C*(a + b*tan(d*x+c))^4/b/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.12 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.27

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-6i(a + ib)^4 B \log(i - \tan(c + dx)) + 6i(a - ib)^4 B \log(i + \tan(c + dx)) - 12b^2(-6a^2 + b^2) B \tan(c + dx)}{}$$

input

```
Integrate[(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
((-6*I)*(a + I*b)^4*B*Log[I - Tan[c + d*x]] + (6*I)*(a - I*b)^4*B*Log[I + Tan[c + d*x]] - 12*b^2*(-6*a^2 + b^2)*B*Tan[c + d*x] + 24*a*b^3*B*Tan[c + d*x]^2 + 4*b^4*B*Tan[c + d*x]^3 + 3*C*(a + b*Tan[c + d*x])^4 - 6*(a*B + b*C)*((I*a - b)^3*Log[I - Tan[c + d*x]] - (I*a + b)^3*Log[I + Tan[c + d*x]] + 6*a*b^2*Tan[c + d*x] + b^3*Tan[c + d*x]^2))/(12*b*d)
```

Rubi [A] (verified)

Time = 1.26 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4113, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{4113}$$

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) - C) dx + \frac{C(a + b \tan(c + dx))^4}{4bd}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int (a + b \tan(c + dx))^3 (B \tan(c + dx) - C) dx + \frac{C(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 4011 \\
& \int (a + b \tan(c + dx))^2 (-bB - aC + (aB - bC) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{C(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3042 \\
& \int (a + b \tan(c + dx))^2 (-bB - aC + (aB - bC) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{C(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 4011 \\
& \int (a + b \tan(c + dx)) (-Ca^2 - 2bBa + b^2C + (Ba^2 - 2bCa - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{(aB - bC)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} + \frac{C(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3042 \\
& \int (a + b \tan(c + dx)) (-Ca^2 - 2bBa + b^2C + (Ba^2 - 2bCa - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{(aB - bC)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} + \frac{C(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 4008 \\
& (a^3B - 3a^2bC - 3ab^2B + b^3C) \int \tan(c + dx) dx + \frac{b(a^2B - 2abC - b^2B) \tan(c + dx)}{d} - \\
& x(a^3C + 3a^2bB - 3ab^2C - b^3B) + \frac{(aB - bC)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{C(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3042 \\
& (a^3B - 3a^2bC - 3ab^2B + b^3C) \int \tan(c + dx) dx + \frac{b(a^2B - 2abC - b^2B) \tan(c + dx)}{d} - \\
& x(a^3C + 3a^2bB - 3ab^2C - b^3B) + \frac{(aB - bC)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{C(a + b \tan(c + dx))^4}{4bd}
\end{aligned}$$

$$\begin{aligned} & \downarrow 3956 \\ & \frac{b(a^2B - 2abC - b^2B) \tan(c + dx)}{d} - \frac{(a^3B - 3a^2bC - 3ab^2B + b^3C) \log(\cos(c + dx))}{d} \\ & x(a^3C + 3a^2bB - 3ab^2C - b^3B) + \frac{(aB - bC)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} + \\ & \frac{C(a + b \tan(c + dx))^4}{4bd} \end{aligned}$$

input `Int[(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `-((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*x) - ((a^3*B - 3*a*b^2*B - 3*a^2*b*C + b^3*C)*Log[Cos[c + d*x]])/d + (b*(a^2*B - b^2*B - 2*a*b*C)*Tan[c + d*x])/d + ((a*B - b*C)*(a + b*Tan[c + d*x])^2)/(2*d) + (B*(a + b*Tan[c + d*x])^3)/(3*d) + (C*(a + b*Tan[c + d*x])^4)/(4*b*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^m)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.08

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{3Cb^3 \tan(dx + c)^4 + 4(3Cab^2 + Bb^3) \tan(dx + c)^3 - 12(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)dx + 6(3Ca^2b^2 + 3Cb^3) \tan(dx + c)^2 + 6(3Ca^2b^2 + 3Cb^3) \tan(dx + c) + 6(3Ca^2b^2 + 3Cb^3)}{d}$$

input

```
integrate((a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")
```

output

```
1/12*(3*C*b^3*tan(d*x + c)^4 + 4*(3*C*a*b^2 + B*b^3)*tan(d*x + c)^3 - 12*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*d*x + 6*(3*C*a^2*b + 3*B*a*b^2 - C*b^3)*tan(d*x + c)^2 - 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*log(1/(tan(d*x + c)^2 + 1)) + 12*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*tan(d*x + c))/d
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 313 vs. 2(151) = 302.

Time = 0.17 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.90

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \frac{Ba^3 \log(\tan^2(c+dx)+1)}{2d} - 3Ba^2bx + \frac{3Ba^2b \tan(c+dx)}{d} - \frac{3Bab^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{3Bab^2 \tan^2(c+dx)}{2d} + Bb^3x + \frac{Bb^3 \tan^2(c+dx)}{d} \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \end{cases}$$

input

```
integrate((a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a**2*b*x + 3*B*a**2
*b*tan(c + d*x)/d - 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a*b**2
*tan(c + d*x)**2/(2*d) + B*b**3*x + B*b**3*tan(c + d*x)**3/(3*d) - B*b**3*
tan(c + d*x)/d - C*a**3*x + C*a**3*tan(c + d*x)/d - 3*C*a**2*b*log(tan(c +
d*x)**2 + 1)/(2*d) + 3*C*a**2*b*tan(c + d*x)**2/(2*d) + 3*C*a*b**2*x + C*
a*b**2*tan(c + d*x)**3/d - 3*C*a*b**2*tan(c + d*x)/d + C*b**3*log(tan(c +
d*x)**2 + 1)/(2*d) + C*b**3*tan(c + d*x)**4/(4*d) - C*b**3*tan(c + d*x)**2
/(2*d), Ne(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.08

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{3 C b^3 \tan(dx + c)^4 + 4 (3 C a b^2 + B b^3) \tan(dx + c)^3 + 6 (3 C a^2 b + 3 B a b^2 - C b^3) \tan(dx + c)^2 - 12 (C$$

input

```
integrate((a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="m
axima")
```

output

```
1/12*(3*C*b^3*tan(d*x + c)^4 + 4*(3*C*a*b^2 + B*b^3)*tan(d*x + c)^3 + 6*(3
*C*a^2*b + 3*B*a*b^2 - C*b^3)*tan(d*x + c)^2 - 12*(C*a^3 + 3*B*a^2*b - 3*C
*a*b^2 - B*b^3)*(d*x + c) + 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*log(
tan(d*x + c)^2 + 1) + 12*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*tan(d*x +
c))/d
```

Giac [A] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.52

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c)}{d}$$

$$+ \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{3Cb^3d^3 \tan(dx + c)^4 + 12Cab^2d^3 \tan(dx + c)^3 + 4Bb^3d^3 \tan(dx + c)^2 + 18Ca^2bd^3 \tan(dx + c) + 18Ca^2bd^3}{d^4}$$

input

```
integrate((a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

```
-(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*(d*x + c)/d + 1/2*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^2 + 1)/d + 1/12*(3*C*b^3*d^3*tan(d*x + c)^4 + 12*C*a*b^2*d^3*tan(d*x + c)^3 + 4*B*b^3*d^3*tan(d*x + c)^2 + 18*C*a^2*b*d^3*tan(d*x + c) + 18*B*a*b^2*d^3*tan(d*x + c) - 6*C*b^3*d^3*tan(d*x + c)^2 + 12*C*a^3*d^3*tan(d*x + c) + 36*B*a^2*b*d^3*tan(d*x + c) - 36*C*a*b^2*d^3*tan(d*x + c) - 12*B*b^3*d^3*tan(d*x + c))/d^4
```

Mupad [B] (verification not implemented)

Time = 5.21 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.10

$$\int (a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= x (-Ca^3 - 3Ba^2b + 3Cab^2 + Bb^3) - \frac{\tan(c + dx)^2 \left(\frac{Cb^3}{2} - \frac{3ab(Bb + Ca)}{2} \right)}{d}$$

$$- \frac{\tan(c + dx) (-Ca^3 - 3Ba^2b + 3Cab^2 + Bb^3)}{d}$$

$$+ \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{Ba^3}{2} - \frac{3Ca^2b}{2} - \frac{3Bab^2}{2} + \frac{Cb^3}{2} \right)}{d}$$

$$+ \frac{\tan(c + dx)^3 \left(\frac{Bb^3}{3} + Cab^2 \right)}{d} + \frac{Cb^3 \tan(c + dx)^4}{4d}$$

3.18 $\int \cot(c+dx)(a+b \tan(c+dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	234
Mathematica [C] (verified)	235
Rubi [A] (verified)	235
Maple [A] (verified)	238
Fricas [A] (verification not implemented)	238
Sympy [A] (verification not implemented)	239
Maxima [A] (verification not implemented)	239
Giac [A] (verification not implemented)	240
Mupad [B] (verification not implemented)	241
Reduce [B] (verification not implemented)	241

Optimal result

Integrand size = 38, antiderivative size = 140

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (a^3 B - 3ab^2 B - 3a^2 b C + b^3 C) x - \frac{(3a^2 b B - b^3 B + a^3 C - 3ab^2 C) \log(\cos(c + dx))}{d}$$

$$+ \frac{b(2abB + a^2 C - b^2 C) \tan(c + dx)}{d}$$

$$+ \frac{(bB + aC)(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3d}$$

output

```
(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*x-(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*ln(c
os(d*x+c))/d+b*(2*B*a*b+C*a^2-C*b^2)*tan(d*x+c)/d+1/2*(B*b+C*a)*(a+b*tan(d
*x+c))^2/d+1/3*C*(a+b*tan(d*x+c))^3/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.73 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.93

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{3(a + ib)^3(-iB + C) \log(i - \tan(c + dx)) + 3(a - ib)^3(iB + C) \log(i + \tan(c + dx)) + 6b(3abB + 3a^2C)}{6d}$$

input

```
Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(3*(a + I*b)^3*((-I)*B + C)*Log[I - Tan[c + d*x]] + 3*(a - I*b)^3*(I*B + C)*Log[I + Tan[c + d*x]] + 6*b*(3*a*b*B + 3*a^2*C - b^2*C)*Tan[c + d*x] + 3*b^2*(b*B + 3*a*C)*Tan[c + d*x]^2 + 2*b^3*C*Tan[c + d*x]^3)/(6*d)
```

Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3042, 4115, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)} dx$$

$$\downarrow \text{4115}$$

$$\int (a + b \tan(c + dx))^3 (B + C \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int (a + b \tan(c + dx))^3 (B + C \tan(c + dx)) dx \\
& \quad \downarrow \text{4011} \\
& \int (a + b \tan(c + dx))^2 (aB - bC + (bB + aC) \tan(c + dx)) dx + \frac{C(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& \int (a + b \tan(c + dx))^2 (aB - bC + (bB + aC) \tan(c + dx)) dx + \frac{C(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{4011} \\
& \int (a + b \tan(c + dx)) (Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c + dx)) dx + \\
& \quad \frac{(aC + bB)(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& \int (a + b \tan(c + dx)) (Ba^2 - 2bCa - b^2B + (Ca^2 + 2bBa - b^2C) \tan(c + dx)) dx + \\
& \quad \frac{(aC + bB)(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{4008} \\
& (a^3C + 3a^2bB - 3ab^2C - b^3B) \int \tan(c + dx) dx + \frac{b(a^2C + 2abB - b^2C) \tan(c + dx)}{d} + \\
& x(a^3B - 3a^2bC - 3ab^2B + b^3C) + \frac{(aC + bB)(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& (a^3C + 3a^2bB - 3ab^2C - b^3B) \int \tan(c + dx) dx + \frac{b(a^2C + 2abB - b^2C) \tan(c + dx)}{d} + \\
& x(a^3B - 3a^2bC - 3ab^2B + b^3C) + \frac{(aC + bB)(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{3956} \\
& \frac{b(a^2C + 2abB - b^2C) \tan(c + dx)}{d} - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \log(\cos(c + dx))}{d} + \\
& x(a^3B - 3a^2bC - 3ab^2B + b^3C) + \frac{(aC + bB)(a + b \tan(c + dx))^2}{2d} + \frac{C(a + b \tan(c + dx))^3}{3d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

$$(a^3B - 3a^2bB - 3a^2bC + b^3C)x - ((3a^2bB - b^3B + a^3C - 3a^2b^2C) \cdot \text{Log}[\text{Cos}[c + dx]])/d + (b(2a^2bB + a^2C - b^2C) \cdot \text{Tan}[c + dx])/d + ((bB + aC)(a + b \cdot \text{Tan}[c + dx])^2)/(2d) + (C(a + b \cdot \text{Tan}[c + dx])^3)/(3d)$$

Defintions of rubi rules used

rule 3042

$$\text{Int}[u_, x_Symbol] \text{ :> Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\text{tan}[(c_.) + (d_.)(x_.)], x_Symbol] \text{ :> Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + dx], x]]/d, x] \text{ /; FreeQ}[\{c, d\}, x]$$

rule 4008

$$\text{Int}[(a_. + (b_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)]) \cdot ((c_.) + (d_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)]), x_Symbol] \text{ :> Simp}[(a \cdot c - b \cdot d) \cdot x, x] + (\text{Simp}[b \cdot d \cdot (\text{Tan}[e + f \cdot x]/f), x] + \text{Simp}[(b \cdot c + a \cdot d) \cdot \text{Int}[\text{Tan}[e + f \cdot x], x], x]) \text{ /; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[b \cdot c + a \cdot d, 0]$$

rule 4011

$$\text{Int}[(a_. + (b_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)])^{(m_.)} \cdot ((c_.) + (d_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)]), x_Symbol] \text{ :> Simp}[d \cdot ((a + b \cdot \text{Tan}[e + f \cdot x])^m / (f \cdot m)), x] + \text{Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^{(m-1)} \cdot \text{Simp}[a \cdot c - b \cdot d + (b \cdot c + a \cdot d) \cdot \text{Tan}[e + f \cdot x], x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$$

rule 4115

$$\text{Int}[(a_. + (b_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)])^{(m_.)} \cdot ((c_.) + (d_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)])^{(n_.)} \cdot ((A_.) + (B_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)] + (C_.) \cdot \text{tan}[(e_.) + (f_.)(x_.)]^2), x_Symbol] \text{ :> Simp}[1/b^2 \cdot \text{Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^{(m+1)} \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot (b \cdot B - a \cdot C + b \cdot C \cdot \text{Tan}[e + f \cdot x]), x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f, A, B, C, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C, 0]$$

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.99

method	result
parallelrisc	$\frac{3(3B a^2 b - B b^3 + C a^3 - 3C a b^2) \ln(\sec(dx+c)^2) + 2C b^3 \tan(dx+c)^3 + 3(B b^3 + 3C a b^2) \tan(dx+c)^2 + 6(3B a b^2 + 3C a^2 b^2) \tan(dx+c) + 6C b^3}{6d}$
norman	$(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) x + \frac{b(3B a b + 3C a^2 - C b^2) \tan(dx+c)}{d} + \frac{C b^3 \tan(dx+c)^3}{3d} + \frac{b^2(B b^3 + 3C a b^2) \tan(dx+c)^2}{3d} + \frac{b^2(B b^3 + 3C a b^2) \tan(dx+c)}{3d}$
derivativedivides	$\frac{\frac{C b^3 \tan(dx+c)^3}{3} + \frac{B b^3 \tan(dx+c)^2}{2} + \frac{3C a b^2 \tan(dx+c)^2}{2} + 3B a b^2 \tan(dx+c) + 3C a^2 b \tan(dx+c) - C b^3 \tan(dx+c) + \frac{(3B a^2 b^2 + 3C a^2 b^2) \tan(dx+c)}{d}}{d}$
default	$\frac{C b^3 \tan(dx+c)^3}{3} + \frac{B b^3 \tan(dx+c)^2}{2} + \frac{3C a b^2 \tan(dx+c)^2}{2} + 3B a b^2 \tan(dx+c) + 3C a^2 b \tan(dx+c) - C b^3 \tan(dx+c) + \frac{(3B a^2 b^2 + 3C a^2 b^2) \tan(dx+c)}{d}$
risc	$B a^3 x - 3B a b^2 x - 3C a^2 b x + C b^3 x + 3i B a^2 b x + i C a^3 x - \frac{2i B b^3 c}{d} + \frac{6i B a^2 b c}{d} - 3i C a b^2 x$

input

```
int(cot(d*x+c)*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_
RETURNVERBOSE)
```

output

```
1/6*(3*(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*ln(sec(d*x+c)^2)+2*C*b^3*tan(d*x+
c)^3+3*(B*b^3+3*C*a*b^2)*tan(d*x+c)^2+6*(3*B*a*b^2+3*C*a^2*b-C*b^3)*tan(d*
x+c)+6*d*x*(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3))/d
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.01

$$\int \cot(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \frac{2 C b^3 \tan(dx+c)^3 + 6 (B a^3 - 3 C a^2 b - 3 B a b^2 + C b^3) dx + 3 (3 C a b^2 + B b^3) \tan(dx+c)^2 - 3 (C a^3 + 3 B a^2 b + 3 C a b^2) \tan(dx+c) + 6 C b^3}{6 d}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, a
lgorithm="fricas")
```

output

```
1/6*(2*C*b^3*tan(d*x + c)^3 + 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*d*x + 3*(3*C*a*b^2 + B*b^3)*tan(d*x + c)^2 - 3*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*log(1/(tan(d*x + c)^2 + 1)) + 6*(3*C*a^2*b + 3*B*a*b^2 - C*b^3)*tan(d*x + c))/d
```

Sympy [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.77

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} Ba^3x + \frac{3Ba^2b \log(\tan^2(c+dx)+1)}{2d} - 3Bab^2x + \frac{3Bab^2 \tan(c+dx)}{d} - \frac{Bb^3 \log(\tan^2(c+dx)+1)}{2d} + \frac{Bb^3 \tan^2(c+dx)}{2d} + \frac{Ca^3 \log(\tan^2(c+dx)+1)}{2d} \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot(c) \end{cases}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((B*a**3*x + 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a*b**2*x + 3*B*a*b**2*tan(c + d*x)/d - B*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**3*tan(c + d*x)**2/(2*d) + C*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - 3*C*a**2*b*x + 3*C*a**2*b*tan(c + d*x)/d - 3*C*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*C*a*b**2*tan(c + d*x)**2/(2*d) + C*b**3*x + C*b**3*tan(c + d*x)**3/(3*d) - C*b**3*tan(c + d*x)/d, Ne(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2)*cot(c), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.02

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Cb^3 \tan(dx + c)^3 + 3(3Cab^2 + Bb^3) \tan(dx + c)^2 + 6(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx + c) + 3(Ca^3 \log(\tan^2(dx + c) + 1) - Bb^3 \log(\tan^2(dx + c) + 1))}{6d}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")
```

output

$$\frac{1}{6}*(2*C*b^3*\tan(d*x + c)^3 + 3*(3*C*a*b^2 + B*b^3)*\tan(d*x + c)^2 + 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(d*x + c) + 3*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*\log(\tan(d*x + c)^2 + 1) + 6*(3*C*a^2*b + 3*B*a*b^2 - C*b^3)*\tan(d*x + c))/d$$
Giac [A] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.30

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ba^3 - 3Ca^2b - 3Cab^2 + Cb^3)(dx + c)}{d}$$

$$+ \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{2Cb^3d^2 \tan(dx + c)^3 + 9Cab^2d^2 \tan(dx + c)^2 + 3Bb^3d^2 \tan(dx + c)^2 + 18Ca^2bd^2 \tan(dx + c) + 18Cb^3d^2}{6d^3}$$

input

```
integrate(cot(d*x+c)*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")
```

output

$$(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(d*x + c)/d + 1/2*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*\log(\tan(d*x + c)^2 + 1)/d + 1/6*(2*C*b^3*d^2*\tan(d*x + c)^3 + 9*C*a*b^2*d^2*\tan(d*x + c)^2 + 3*B*b^3*d^2*\tan(d*x + c)^2 + 18*C*a^2*b*d^2*\tan(d*x + c) + 18*B*a*b^2*d^2*\tan(d*x + c) - 6*C*b^3*d^2*\tan(d*x + c))/d^3$$

Mupad [B] (verification not implemented)

Time = 5.34 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.01

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= x (B a^3 - 3 C a^2 b - 3 B a b^2 + C b^3)$$

$$- \frac{\ln(\tan(c + dx)^2 + 1) \left(-\frac{C a^3}{2} - \frac{3 B a^2 b}{2} + \frac{3 C a b^2}{2} + \frac{B b^3}{2} \right)}{d}$$

$$+ \frac{\tan(c + dx)^2 \left(\frac{B b^3}{2} + \frac{3 C a b^2}{2} \right)}{d}$$

$$- \frac{\tan(c + dx) (C b^3 - 3 a b (B b + C a))}{d} + \frac{C b^3 \tan(c + dx)^3}{3 d}$$

input `int(cot(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^3,x)`

output `x*(B*a^3 + C*b^3 - 3*B*a*b^2 - 3*C*a^2*b) - (log(tan(c + d*x)^2 + 1)*((B*b^3)/2 - (C*a^3)/2 - (3*B*a^2*b)/2 + (3*C*a*b^2)/2))/d + (tan(c + d*x)^2*((B*b^3)/2 + (3*C*a*b^2)/2))/d - (tan(c + d*x)*(C*b^3 - 3*a*b*(B*b + C*a)))/d + (C*b^3*tan(c + d*x)^3)/(3*d)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 992, normalized size of antiderivative = 7.09

$$\int \cot(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

= Too large to display

input `int(cot(d*x+c)*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output

```
(6*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**3*c + 18*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**2*b**2 - 18*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a*b**2*c - 6*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*b**4 - 6*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*a**3*c - 18*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*a**2*b**2 + 18*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*a*b**2*c + 6*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*b**4 - 6*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*a**3*c - 18*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*a**2*b**2 + 18*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*a*b**2*c + 6*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*b**4 + 6*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*a**3*c + 18*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*a**2*b**2 - 18*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*a*b**2*c - 6*cos(c + d*x)*log(tan((c + d*x)/2) - 1)*b**4 - 6*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*a**3*c - 18*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*a**2*b**2 + 18*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*a*b**2*c + 6*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*b**4 + 6*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*a**3*c + 18*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*a**2*b**2 - 18*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*a*b**2*c - 6*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*b**4 + 6*cos(c + d*x)*sin(c + d*x)**2...
```

3.19 $\int \cot^2(c+dx)(a+b \tan(c+dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	243
Mathematica [C] (verified)	243
Rubi [A] (verified)	244
Maple [A] (verified)	248
Fricas [A] (verification not implemented)	248
Sympy [A] (verification not implemented)	249
Maxima [A] (verification not implemented)	249
Giac [A] (verification not implemented)	250
Mupad [B] (verification not implemented)	250
Reduce [B] (verification not implemented)	251

Optimal result

Integrand size = 40, antiderivative size = 117

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (3a^2bB - b^3B + a^3C - 3ab^2C) x - \frac{b(3abB + 3a^2C - b^2C) \log(\cos(c + dx))}{d}$$

$$+ \frac{a^3B \log(\sin(c + dx))}{d} + \frac{b^2(bB + 2aC) \tan(c + dx)}{d} + \frac{bC(a + b \tan(c + dx))^2}{2d}$$

output

```
(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*x-b*(3*B*a*b+3*C*a^2-C*b^2)*ln(cos(d*x+c))
)/d+a^3*B*ln(sin(d*x+c))/d+b^2*(B*b+2*C*a)*tan(d*x+c)/d+1/2*b*C*(a+b*tan(d*x+c))^2/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.97

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-(a + ib)^3(B + iC) \log(i - \tan(c + dx)) + 2a^3B \log(\tan(c + dx)) - (a - ib)^3(B - iC) \log(i + \tan(c + dx))}{2d}$$

input

```
Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(-((a + I*b)^3*(B + I*C)*Log[I - Tan[c + d*x]]) + 2*a^3*B*Log[Tan[c + d*x]] - (a - I*b)^3*(B - I*C)*Log[I + Tan[c + d*x]] + 2*b^2*(b*B + 3*a*C)*Tan[c + d*x] + b^3*C*Tan[c + d*x]^2)/(2*d)
```

Rubi [A] (verified)

Time = 1.43 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.02, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.325$, Rules used = {3042, 4115, 3042, 4090, 27, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \cot(c + dx)(a + b \tan(c + dx))^3 (B + C \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3 (B + C \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{1}{2} \int 2 \cot(c + dx)(a + b \tan(c + dx)) (Ba^2 + b(bB + 2aC) \tan^2(c + dx) + (Ca^2 + 2bBa - b^2C) \tan(c + dx)) dx + \\
 & \quad \frac{bC(a + b \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
& \int \cot(c+dx)(a+b \tan(c+dx)) (Ba^2+b(bB+2aC) \tan^2(c+dx) + (Ca^2+2bBa-b^2C) \tan(c+dx)) dx + \\
& \quad \frac{bC(a+b \tan(c+dx))^2}{2d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+b \tan(c+dx)) (Ba^2+b(bB+2aC) \tan(c+dx)^2 + (Ca^2+2bBa-b^2C) \tan(c+dx))}{\tan(c+dx) \frac{bC(a+b \tan(c+dx))^2}{2d}} dx + \\
& \quad \downarrow \text{4120} \\
& - \int -\cot(c+dx) (Ba^3+b(3Ca^2+3bBa-b^2C) \tan^2(c+dx) + (Ca^3+3bBa^2-3b^2Ca-b^3B) \tan(c+dx)) dx + \\
& \quad \frac{b^2(2aC+bB) \tan(c+dx)}{d} + \frac{bC(a+b \tan(c+dx))^2}{2d} \\
& \quad \downarrow \text{25} \\
& \int \cot(c+dx) (Ba^3+b(3Ca^2+3bBa-b^2C) \tan^2(c+dx) + (Ca^3+3bBa^2-3b^2Ca-b^3B) \tan(c+dx)) dx + \\
& \quad \frac{b^2(2aC+bB) \tan(c+dx)}{d} + \frac{bC(a+b \tan(c+dx))^2}{2d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{Ba^3+b(3Ca^2+3bBa-b^2C) \tan(c+dx)^2 + (Ca^3+3bBa^2-3b^2Ca-b^3B) \tan(c+dx)}{\tan(c+dx) \frac{b^2(2aC+bB) \tan(c+dx)}{d} + \frac{bC(a+b \tan(c+dx))^2}{2d}} dx + \\
& \quad \downarrow \text{4107} \\
& a^3B \int \cot(c+dx) dx + b(3a^2C+3abB-b^2C) \int \tan(c+dx) dx + \\
& x(a^3C+3a^2bB-3ab^2C-b^3B) + \frac{b^2(2aC+bB) \tan(c+dx)}{d} + \frac{bC(a+b \tan(c+dx))^2}{2d} \\
& \quad \downarrow \text{3042} \\
& a^3B \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + b(3a^2C+3abB-b^2C) \int \tan(c+dx) dx + \\
& x(a^3C+3a^2bB-3ab^2C-b^3B) + \frac{b^2(2aC+bB) \tan(c+dx)}{d} + \frac{bC(a+b \tan(c+dx))^2}{2d}
\end{aligned}$$

$$\begin{aligned}
 & \downarrow 25 \\
 & a^3(-B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b(3a^2C + 3abB - b^2C) \int \tan(c + dx) dx + \\
 & x(a^3C + 3a^2bB - 3ab^2C - b^3B) + \frac{b^2(2aC + bB) \tan(c + dx)}{d} + \frac{bC(a + b \tan(c + dx))^2}{2d} \\
 & \downarrow 3956 \\
 & \frac{a^3B \log(-\sin(c + dx))}{d} - \frac{b(3a^2C + 3abB - b^2C) \log(\cos(c + dx))}{d} + \\
 & x(a^3C + 3a^2bB - 3ab^2C - b^3B) + \frac{b^2(2aC + bB) \tan(c + dx)}{d} + \frac{bC(a + b \tan(c + dx))^2}{2d}
 \end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `(3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*x - (b*(3*a*b*B + 3*a^2*C - b^2*C)*Log[Cos[c + d*x]])/d + (a^3*B*Log[-Sin[c + d*x]])/d + (b^2*(b*B + 2*a*C)*Tan[c + d*x])/d + (b*C*(a + b*Tan[c + d*x])^2)/(2*d)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4090

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4107

```

Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2
)/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[
e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B,
C}, x] && NeQ[A, C]

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4120

```

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]

```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.03

method	result
parallelrisc	$(-B a^3 + 3Ba b^2 + 3C a^2 b - C b^3) \ln(\sec(dx+c)^2) + 2B a^3 \ln(\tan(dx+c)) + C b^3 \tan(dx+c)^2 + (2B b^3 + 6C a b^2) \tan(dx+c)$
derivativedivides	$\frac{C b^3 \tan(dx+c)^2 + B b^3 \tan(dx+c) + 3C a b^2 \tan(dx+c) + \frac{(-B a^3 + 3Ba b^2 + 3C a^2 b - C b^3) \ln(1 + \tan(dx+c)^2)}{2}}{d} + (3B a^2 b - B b^5)$
default	$\frac{C b^3 \tan(dx+c)^2 + B b^3 \tan(dx+c) + 3C a b^2 \tan(dx+c) + \frac{(-B a^3 + 3Ba b^2 + 3C a^2 b - C b^3) \ln(1 + \tan(dx+c)^2)}{2}}{d} + (3B a^2 b - B b^5)$
norman	$\frac{(3B a^2 b - B b^3 + C a^3 - 3C a b^2) x \tan(dx+c) + \frac{b^2 (Bb + 3Ca) \tan(dx+c)^2}{d} + \frac{C b^3 \tan(dx+c)^3}{2d}}{\tan(dx+c)} + \frac{B a^3 \ln(\tan(dx+c))}{d} - \frac{(B a^3 - B b^3)}{d}$
risc	$-iB a^3 x + 3iBa b^2 x + \frac{2ib^2 (Bb e^{2i(dx+c)} + 3Ca e^{2i(dx+c)} - iCb e^{2i(dx+c)} + Bb + 3Ca)}{d(e^{2i(dx+c)} + 1)^2} + 3iC a^2 b x + 3B$

input `int(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method =_RETURNVERBOSE)`

output $1/2*((-B*a^3+3*B*a*b^2+3*C*a^2*b-C*b^3)*\ln(\sec(d*x+c)^2)+2*B*a^3*\ln(\tan(d*x+c))+C*b^3*\tan(d*x+c)^2+(2*B*b^3+6*C*a*b^2)*\tan(d*x+c)+6*x*(B*a^2*b-1/3*B*b^3+1/3*C*a^3-C*a*b^2)*d)/d$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.14

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{C b^3 \tan(dx + c)^2 + B a^3 \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + 2(C a^3 + 3 B a^2 b - 3 C a b^2 - B b^3) dx - (3 C a^2 b + 3 B a b^2 - B b^3)}{2 d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output

```
1/2*(C*b^3*tan(d*x + c)^2 + B*a^3*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))
+ 2*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*d*x - (3*C*a^2*b + 3*B*a*b^2
- C*b^3)*log(1/(tan(d*x + c)^2 + 1)) + 2*(3*C*a*b^2 + B*b^3)*tan(d*x + c)
/d
```

Sympy [A] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.80

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} -\frac{Ba^3 \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba^3 \log(\tan(c+dx))}{d} + 3Ba^2bx + \frac{3Bab^2 \log(\tan^2(c+dx)+1)}{2d} - Bb^3x + \frac{Bb^3 \tan(c+dx)}{d} + C \dots \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot^2(c) \end{cases}$$

input

```
integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2)
,x)
```

output

```
Piecewise((-B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**3*log(tan(c + d*x)
))/d + 3*B*a**2*b*x + 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - B*b**3*x
+ B*b**3*tan(c + d*x)/d + C*a**3*x + 3*C*a**2*b*log(tan(c + d*x)**2 + 1)/
(2*d) - 3*C*a*b**2*x + 3*C*a*b**2*tan(c + d*x)/d - C*b**3*log(tan(c + d*x)
**2 + 1)/(2*d) + C*b**3*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(a + b*tan(c)
)**3*(B*tan(c) + C*tan(c)**2)*cot(c)**2, True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.06

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Cb^3 \tan(dx + c)^2 + 2Ba^3 \log(\tan(dx + c)) + 2(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c) - (Ba^3 - 3Ca^2b - 3Cab^2 - Bb^3)}{2d}$$

input

```
integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
1/2*(C*b^3*tan(d*x + c)^2 + 2*B*a^3*log(tan(d*x + c)) + 2*(C*a^3 + 3*B*a^2
*b - 3*C*a*b^2 - B*b^3)*(d*x + c) - (B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3
)*log(tan(d*x + c)^2 + 1) + 2*(3*C*a*b^2 + B*b^3)*tan(d*x + c))/d
```

Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.20

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{Ba^3 \log(|\tan(dx + c)|)}{d} + \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c)}{d}$$

$$- \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(\tan(dx + c)^2 + 1)}{2d}$$

$$+ \frac{Cb^3 d \tan(dx + c)^2 + 6Cab^2 d \tan(dx + c) + 2Bb^3 d \tan(dx + c)}{2d^2}$$

input

```
integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
B*a^3*log(abs(tan(d*x + c)))/d + (C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*(
d*x + c)/d - 1/2*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^
2 + 1)/d + 1/2*(C*b^3*d*tan(d*x + c)^2 + 6*C*a*b^2*d*tan(d*x + c) + 2*B*b^
3*d*tan(d*x + c))/d^2
```

Mupad [B] (verification not implemented)

Time = 5.62 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.01

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\tan(c + dx) (Bb^3 + 3Cab^2)}{d} + \frac{Ba^3 \ln(\tan(c + dx))}{d}$$

$$+ \frac{Cb^3 \tan(c + dx)^2}{2d} - \frac{\ln(\tan(c + dx) + 1i) (B - C 1i) (b + a 1i)^3 1i}{2d}$$

$$- \frac{\ln(\tan(c + dx) - 1i) (B + C 1i) (-b + a 1i)^3 1i}{2d}$$

input `int(cot(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^3,x)`

output `(tan(c + d*x)*(B*b^3 + 3*C*a*b^2))/d + (B*a^3*log(tan(c + d*x)))/d - (log(tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b)^3*1i)/(2*d) - (log(tan(c + d*x) - 1i)*(B + C*1i)*(a*1i - b)^3*1i)/(2*d) + (C*b^3*tan(c + d*x)^2)/(2*d)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 670, normalized size of antiderivative = 5.73

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

= Too large to display

input `int(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `(- 6*cos(c + d*x)*sin(c + d*x)*a*b**2*c - 2*cos(c + d*x)*sin(c + d*x)*b**4 - 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**3*b + 6*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**2*b*c + 6*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a*b**3 - 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*b**3*c + 2*log(tan((c + d*x)/2)**2 + 1)*a**3*b - 6*log(tan((c + d*x)/2)**2 + 1)*a**2*b*c - 6*log(tan((c + d*x)/2)**2 + 1)*a*b**3 + 2*log(tan((c + d*x)/2)**2 + 1)*b**3*c - 6*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*a**2*b*c - 6*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*a*b**3 + 2*log(tan((c + d*x)/2) - 1)*sin(c + d*x)**2*b**3*c + 6*log(tan((c + d*x)/2) - 1)*a**2*b*c + 6*log(tan((c + d*x)/2) - 1)*a*b**3 - 2*log(tan((c + d*x)/2) - 1)*b**3*c - 6*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*a**2*b*c - 6*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*a*b**3 + 2*log(tan((c + d*x)/2) + 1)*sin(c + d*x)**2*b**3*c + 6*log(tan((c + d*x)/2) + 1)*a**2*b*c + 6*log(tan((c + d*x)/2) + 1)*a*b**3 - 2*log(tan((c + d*x)/2) + 1)*b**3*c + 2*log(tan((c + d*x)/2))*sin(c + d*x)**2*a**3*b - 2*log(tan((c + d*x)/2))*a**3*b + 2*sin(c + d*x)**2*a**3*c*d*x + 6*sin(c + d*x)**2*a**2*b**2*d*x - 6*sin(c + d*x)**2*a*b**2*c*d*x - 2*sin(c + d*x)**2*b**4*d*x - sin(c + d*x)**2*b**3*c - 2*a**3*c*d*x - 6*a**2*b**2*d*x + 6*a*b**2*c*d*x + 2*b**4*d*x)/(2*d*(sin(c + d*x)**2 - 1))`

3.20 $\int \cot^3(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	252
Mathematica [C] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	257
Fricas [A] (verification not implemented)	257
Sympy [A] (verification not implemented)	258
Maxima [A] (verification not implemented)	258
Giac [A] (verification not implemented)	259
Mupad [B] (verification not implemented)	260
Reduce [B] (verification not implemented)	260

Optimal result

Integrand size = 40, antiderivative size = 119

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= -((a^3B - 3ab^2B - 3a^2bC + b^3C) x) - \frac{b^2(bB + 3aC) \log(\cos(c+dx))}{d} + \frac{a^2(3bB + aC) \log(\sin(c+dx))}{d} + \frac{b^2(aB + bC) \tan(c+dx)}{d} - \frac{aB \cot(c+dx)(a+b \tan(c+dx))^2}{d}$$

output

```
-(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*x-b^2*(B*b+3*C*a)*ln(cos(d*x+c))/d+a^2*(3*B*b+C*a)*ln(sin(d*x+c))/d+b^2*(B*a+C*b)*tan(d*x+c)/d-a*B*cot(d*x+c)*(a+b*tan(d*x+c))^2/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.95

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \frac{-2a^3B \cot(c+dx) + i(a+ib)^3(B+iC) \log(i - \tan(c+dx)) + 2a^2(3bB + aC) \log(\tan(c+dx)) + ia}{2d}$$

input

```
Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(-2*a^3*B*Cot[c + d*x] + I*(a + I*b)^3*(B + I*C)*Log[I - Tan[c + d*x]] + 2*a^2*(3*b*B + a*C)*Log[Tan[c + d*x]] + (I*a + b)^3*(B - I*C)*Log[I + Tan[c + d*x]] + 2*b^3*C*Tan[c + d*x])/(2*d)
```

Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 4115, 3042, 4088, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)^3} dx \\
 & \quad \downarrow 4115 \\
 & \int \cot^2(c + dx)(a + b \tan(c + dx))^3 (B + C \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + b \tan(c + dx))^3 (B + C \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow 4088 \\
 & \int \cot(c + dx)(a + b \tan(c + dx)) (b(aB + bC) \tan^2(c + dx) - (Ba^2 - 2bCa - b^2B) \tan(c + dx) + a(3bB + aC)) dx - \\
 & \quad \frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$\int \frac{(a + b \tan(c + dx)) (b(aB + bC) \tan(c + dx)^2 - (Ba^2 - 2bCa - b^2B) \tan(c + dx) + a(3bB + aC))}{\tan(c + dx) \frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d}} dx -$$

$$\downarrow 4120$$

$$dx) \left(\frac{(3bB + aC)a^2 + b^2(bB + 3aC) \tan^2(c + dx) - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c + dx)}{d} \right) dx + \frac{b^2(aB + bC) \tan(c + dx)}{d} - \frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d}$$

$$\downarrow 25$$

$$dx) \left(\frac{(3bB + aC)a^2 + b^2(bB + 3aC) \tan^2(c + dx) - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c + dx)}{d} \right) dx + \frac{b^2(aB + bC) \tan(c + dx)}{d} - \frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d}$$

$$\downarrow 3042$$

$$\int \frac{(3bB + aC)a^2 + b^2(bB + 3aC) \tan(c + dx)^2 - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c + dx)}{\tan(c + dx) \frac{b^2(aB + bC) \tan(c + dx)}{d} - \frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d}} dx +$$

$$\downarrow 4107$$

$$a^2(aC + 3bB) \int \cot(c + dx) dx + b^2(3aC + bB) \int \tan(c + dx) dx - x(a^3B - 3a^2bC - 3ab^2B + b^3C) + \frac{b^2(aB + bC) \tan(c + dx)}{\frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d}} -$$

$$\downarrow 3042$$

$$a^2(aC + 3bB) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b^2(3aC + bB) \int \tan(c + dx) dx - x(a^3B - 3a^2bC - 3ab^2B + b^3C) + \frac{b^2(aB + bC) \tan(c + dx)}{\frac{aB \cot(c + dx)(a + b \tan(c + dx))^2}{d}} -$$

$$\downarrow 25$$

rule 4088

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
  Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])

```

rule 4107

```

Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2
)/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A  Int[1/Tan[
e + f*x], x], x] + Simp[C  Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B,
C}, x] && NeQ[A, C]

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2  Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4120

```

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2))  Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]

```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.99

method	result
parallelrisch	$\frac{(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \ln(\sec(dx+c)^2) + (6B a^2 b + 2C a^3) \ln(\tan(dx+c)) - 2B \cot(dx+c) a^3 + 2C b^3 \tan(dx+c)}{2d}$
derivativedivides	$C b^3 \tan(dx+c) + \frac{(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \ln(1+\tan(dx+c)^2)}{2} + \frac{(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \arctan(\tan(dx+c))}{d}$
default	$C b^3 \tan(dx+c) + \frac{(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \ln(1+\tan(dx+c)^2)}{2} + \frac{(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \arctan(\tan(dx+c))}{d}$
norman	$\frac{(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) x \tan(dx+c)^2 + \frac{C b^3 \tan(dx+c)^3}{d} - \frac{B a^3 \tan(dx+c)}{d}}{\tan(dx+c)^2} + \frac{a^2(3Bb+Ca) \ln(\tan(dx+c))}{d}$
risch	$-B a^3 x + 3B a b^2 x + 3C a^2 b x - C b^3 x - 3iB a^2 b x - iC a^3 x - \frac{2i(B a^3 e^{2i(dx+c)} - b^3 e^{2i(dx+c)})}{d(e^{2i(dx+c)} - 1)}$

input `int (cot (d*x+c)^3*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2) , x, method =_RETURNVERBOSE)`

output `1/2*((-3*B*a^2*b+B*b^3-C*a^3+3*C*a*b^2)*ln(sec(d*x+c)^2)+(6*B*a^2*b+2*C*a^3)*ln(tan(d*x+c))-2*B*cot(d*x+c)*a^3+2*C*b^3*tan(d*x+c)-2*d*x*(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3))/d`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.22

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2 C b^3 \tan(dx + c)^2 - 2 B a^3 - 2 (B a^3 - 3 C a^2 b - 3 B a b^2 + C b^3) dx \tan(dx + c) + (C a^3 + 3 B a^2 b) \log\left(\frac{\tan(dx+c)}{\sec(dx+c)}\right)}{2 d \tan(dx + c)}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2) , x, algorithm="fricas")`

output

$$\frac{1}{2} * (2 * C * b^3 * \tan(dx + c)^2 - 2 * B * a^3 - 2 * (B * a^3 - 3 * C * a^2 * b - 3 * B * a * b^2 + C * b^3) * dx * \tan(dx + c) + (C * a^3 + 3 * B * a^2 * b) * \log(\tan(dx + c)^2 / (\tan(dx + c)^2 + 1)) * \tan(dx + c) - (3 * C * a * b^2 + B * b^3) * \log(1 / (\tan(dx + c)^2 + 1)) * \tan(dx + c)) / (d * \tan(dx + c))$$

Sympy [A] (verification not implemented)

Time = 1.88 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.80

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot^3(c) \\ \text{NaN} \\ -Ba^3x - \frac{Ba^3}{d \tan(c+dx)} - \frac{3Ba^2b \log(\tan^2(c+dx)+1)}{2d} + \frac{3Ba^2b \log(\tan(c+dx))}{d} + 3Bab^2x + \frac{Bb^3 \log(\tan^2(c+dx)+1)}{2d} - \frac{Ca^3}{d} \end{cases}$$

input

```
integrate(cot(dx+c)**3*(a+b*tan(dx+c))**3*(B*tan(dx+c)+C*tan(dx+c)**2),x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2)*cot(c)**3, Eq(d, 0)), (nan, Eq(c, -dx)), (-B*a**3*x - B*a**3/(d*tan(c + dx)) - 3*B*a**2*b*log(tan(c + dx)**2 + 1)/(2*d) + 3*B*a**2*b*log(tan(c + dx))/d + 3*B*a*b**2*x + B*b**3*log(tan(c + dx)**2 + 1)/(2*d) - C*a**3*log(tan(c + dx)**2 + 1)/(2*d) + C*a**3*log(tan(c + dx))/d + 3*C*a**2*b*x + 3*C*a*b**2*log(tan(c + dx)**2 + 1)/(2*d) - C*b**3*x + C*b**3*tan(c + dx)/d, True))
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.05

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2Cb^3 \tan(dx + c) - \frac{2Ba^3}{\tan(dx+c)} - 2(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx + c) - (Ca^3 + 3Ba^2b - 3Cab^2 - 3C^2ab^2)}{2d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")`

output $\frac{1}{2}*(2*C*b^3*\tan(dx + c) - 2*B*a^3/\tan(dx + c) - 2*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(dx + c) - (C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*\log(\tan(dx + c)^2 + 1) + 2*(C*a^3 + 3*B*a^2*b)*\log(\tan(dx + c)))/d$

Giac [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.13

$$\begin{aligned} & \int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \frac{Cb^3 \tan(dx + c)}{d} - \frac{Ba^3}{d \tan(dx + c)} - \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx + c)}{d} \\ & \quad - \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx + c)^2 + 1)}{2d} \\ & \quad + \frac{(Ca^3 + 3Ba^2b) \log(|\tan(dx + c)|)}{d} \end{aligned}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")`

output $C*b^3*\tan(dx + c)/d - B*a^3/(d*\tan(dx + c)) - (B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(dx + c)/d - 1/2*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*\log(\tan(dx + c)^2 + 1)/d + (C*a^3 + 3*B*a^2*b)*\log(\text{abs}(\tan(dx + c)))/d$

Mupad [B] (verification not implemented)

Time = 5.49 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.96

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (C a^3 + 3 B b a^2)}{d} - \frac{B a^3 \cot(c + dx)}{d}$$

$$+ \frac{C b^3 \tan(c + dx)}{d} + \frac{\ln(\tan(c + dx) - i) (B + C i) (a + b i)^3 i}{2 d}$$

$$- \frac{\ln(\tan(c + dx) + i) (B - C i) (a - b i)^3 i}{2 d}$$

input

```
int(cot(c + d*x)^3*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^3,x)
```

output

```
(log(tan(c + d*x))*(C*a^3 + 3*B*a^2*b))/d + (log(tan(c + d*x) - 1i)*(B + C*1i)*(a + b*1i)^3*1i)/(2*d) - (log(tan(c + d*x) + 1i)*(B - C*1i)*(a - b*1i)^3*1i)/(2*d) - (B*a^3*cot(c + d*x))/d + (C*b^3*tan(c + d*x))/d
```

Reduce [B] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 440, normalized size of antiderivative = 3.70

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-\cos(dx + c) \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) \sin(dx + c) a^3 c - 3 \cos(dx + c) \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) \sin(dx + c)}{d}$$

input

```
int(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
( - cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**3*c - 3*cos(
c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**2*b**2 + 3*cos(c + d
*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a*b**2*c + cos(c + d*x)*log(
tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*b**4 - 3*cos(c + d*x)*log(tan((c + d
*x)/2) - 1)*sin(c + d*x)*a*b**2*c - cos(c + d*x)*log(tan((c + d*x)/2) - 1)
*sin(c + d*x)*b**4 - 3*cos(c + d*x)*log(tan((c + d*x)/2) + 1)*sin(c + d*x)
*a*b**2*c - cos(c + d*x)*log(tan((c + d*x)/2) + 1)*sin(c + d*x)*b**4 + cos
(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**3*c + 3*cos(c + d*x)*log(t
an((c + d*x)/2))*sin(c + d*x)*a**2*b**2 - cos(c + d*x)*sin(c + d*x)*a**3*b
*d*x + 3*cos(c + d*x)*sin(c + d*x)*a**2*b*c*d*x + 3*cos(c + d*x)*sin(c + d
*x)*a*b**3*d*x - cos(c + d*x)*sin(c + d*x)*b**3*c*d*x + sin(c + d*x)**2*a*
*3*b + sin(c + d*x)**2*b**3*c - a**3*b)/(cos(c + d*x)*sin(c + d*x)*d)
```

3.21 $\int \cot^4(c+dx)(a+b \tan(c+dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	262
Mathematica [C] (verified)	263
Rubi [A] (verified)	263
Maple [A] (verified)	267
Fricas [A] (verification not implemented)	268
Sympy [B] (verification not implemented)	268
Maxima [A] (verification not implemented)	269
Giac [A] (verification not implemented)	270
Mupad [B] (verification not implemented)	270
Reduce [F]	271

Optimal result

Integrand size = 40, antiderivative size = 127

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -\left((3a^2bB - b^3B + a^3C - 3ab^2C) x - \frac{a^2(2bB + aC) \cot(c + dx)}{d} \right.$$

$$\left. - \frac{b^3C \log(\cos(c + dx))}{d} - \frac{a(a^2B - 3b^2B - 3abC) \log(\sin(c + dx))}{d} \right.$$

$$\left. - \frac{aB \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \right)$$

output

```
-(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*x-a^2*(2*B*b+C*a)*cot(d*x+c)/d-b^3*C*ln
(cos(d*x+c))/d-a*(B*a^2-3*B*b^2-3*C*a*b)*ln(sin(d*x+c))/d-1/2*a*B*cot(d*x+
c)^2*(a+b*tan(d*x+c))^2/d
```


$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^3 (B + C \tan(c + dx))}{\tan(c + dx)^3} dx \\
& \quad \downarrow 4088 \\
& \frac{1}{2} \int \frac{2 \cot^2(c + dx) (a + b \tan(c + dx)) (b^2 C \tan^2(c + dx) - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC))}{aB \cot^2(c + dx) (a + b \tan(c + dx))^2} dx - \\
& \quad \downarrow 27 \\
& \int \frac{\cot^2(c + dx) (a + b \tan(c + dx)) (b^2 C \tan^2(c + dx) - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC))}{aB \cot^2(c + dx) (a + b \tan(c + dx))^2} dx - \\
& \quad \downarrow 3042 \\
& \int \frac{(a + b \tan(c + dx)) (b^2 C \tan^2(c + dx)^2 - (Ba^2 - 2bCa - b^2 B) \tan(c + dx) + a(2bB + aC))}{\tan(c + dx)^2} dx - \\
& \quad \frac{aB \cot^2(c + dx) (a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 4118 \\
& \int \frac{-\cot(c + dx) (-C \tan^2(c + dx) b^3 + a(Ba^2 - 3bCa - 3b^2 B) + (Ca^3 + 3bBa^2 - 3b^2 Ca - b^3 B) \tan(c + dx))}{\frac{a^2(aC + 2bB) \cot(c + dx)}{d} - \frac{aB \cot^2(c + dx) (a + b \tan(c + dx))^2}{2d}} dx - \\
& \quad \downarrow 25 \\
& - \int \frac{\cot(c + dx) (-C \tan^2(c + dx) b^3 + a(Ba^2 - 3bCa - 3b^2 B) + (Ca^3 + 3bBa^2 - 3b^2 Ca - b^3 B) \tan(c + dx))}{\frac{a^2(aC + 2bB) \cot(c + dx)}{d} - \frac{aB \cot^2(c + dx) (a + b \tan(c + dx))^2}{2d}} dx - \\
& \quad \downarrow 3042 \\
& - \int \frac{-C \tan(c + dx)^2 b^3 + a(Ba^2 - 3bCa - 3b^2 B) + (Ca^3 + 3bBa^2 - 3b^2 Ca - b^3 B) \tan(c + dx)}{\frac{a^2(aC + 2bB) \cot(c + dx)}{d} - \frac{aB \cot^2(c + dx) (a + b \tan(c + dx))^2}{2d}} dx -
\end{aligned}$$

$$\begin{aligned}
& \downarrow 4107 \\
& -a(a^2B - 3abC - 3b^2B) \int \cot(c+dx)dx + b^3C \int \tan(c+dx)dx - \frac{a^2(aC + 2bB) \cot(c + dx)}{d} - \\
& \quad x(a^3C + 3a^2bB - 3ab^2C - b^3B) - \frac{aB \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \downarrow 3042 \\
& -a(a^2B - 3abC - 3b^2B) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b^3C \int \tan(c + dx)dx - \\
& \quad \frac{a^2(aC + 2bB) \cot(c + dx)}{d} - x(a^3C + 3a^2bB - 3ab^2C - b^3B) - \\
& \quad \frac{aB \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \downarrow 25 \\
& a(a^2B - 3abC - 3b^2B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b^3C \int \tan(c + dx)dx - \\
& \quad \frac{a^2(aC + 2bB) \cot(c + dx)}{d} - x(a^3C + 3a^2bB - 3ab^2C - b^3B) - \\
& \quad \frac{aB \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \downarrow 3956 \\
& \frac{a(a^2B - 3abC - 3b^2B) \log(-\sin(c + dx))}{d} - \frac{a^2(aC + 2bB) \cot(c + dx)}{d} - \\
& \quad x(a^3C + 3a^2bB - 3ab^2C - b^3B) - \frac{aB \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} - \frac{b^3C \log(\cos(c + dx))}{d}
\end{aligned}$$

input

```
Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
-((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*x) - (a^2*(2*b*B + a*C)*Cot[c + d*x])/d - (b^3*C*Log[Cos[c + d*x]])/d - (a*(a^2*B - 3*b^2*B - 3*a*b*C)*Log[-Sin[c + d*x]])/d - (a*B*Cot[c + d*x]^2*(a + b*Tan[c + d*x])^2)/(2*d)
```

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4107 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B, C}, x] && NeQ[A, C]`

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.07

method	result
parallelrisc	$\frac{(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \ln(\sec(dx+c)^2) + (-2B a^3 + 6B a b^2 + 6C a^2 b) \ln(\tan(dx+c)) - B \cot(dx+c)^2 a^3 + (-6B a^3 + 6B a b^2 + 6C a^2 b) \arctan(\tan(dx+c)) - \frac{B a^3}{2 \tan(dx+c)^2} - \frac{a^3}{2 \tan(dx+c)^2}}{2d}$
derivativedivides	$\frac{(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \ln(1 + \tan(dx+c)^2) + (-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \arctan(\tan(dx+c)) - \frac{B a^3}{2 \tan(dx+c)^2} - \frac{a^3}{2 \tan(dx+c)^2}}{d}$
default	$\frac{(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \ln(1 + \tan(dx+c)^2) + (-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \arctan(\tan(dx+c)) - \frac{B a^3}{2 \tan(dx+c)^2} - \frac{a^3}{2 \tan(dx+c)^2}}{d}$
norman	$\frac{(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) x \tan(dx+c)^3 - \frac{B a^3 \tan(dx+c)}{2d} - \frac{a^2(3Bb+Ca) \tan(dx+c)^2}{d}}{\tan(dx+c)^3} + \frac{(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \ln(1 + \tan(dx+c)^2)}{2d}$
risc	$i B a^3 x - \frac{2ia^2(3Bb e^{2i(dx+c)} + Ca e^{2i(dx+c)} + iBa e^{2i(dx+c)} - 3Bb - Ca)}{d(e^{2i(dx+c)} - 1)^2} - 3iBa b^2 x - 3iC a^2 b x - 3B a^3$

input

```
int(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method =_RETURNVERBOSE)
```


output

$$\frac{1}{2} \left((B^3 a^3 - 3 B^2 a^2 b - 3 C a^2 b + C^2 b^3) \ln(\sec(dx+c)^2) + (-2 B^3 a^3 + 6 B^2 a^2 b + 2 C a^2 b) \ln(\tan(dx+c)) - B \cot(dx+c)^2 a^3 + (-6 B^2 a^2 b - 2 C a^3) \cot(dx+c) - 6 x (B^2 a^2 b - 1/3 B^3 b^3 + 1/3 C a^3 - C a^2 b^2) \right) / d$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.28

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx =$$

$$\frac{C b^3 \log\left(\frac{1}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 + B a^3 + (B a^3 - 3 C a^2 b - 3 B a b^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)}{2 d \tan(dx+c)}$$

input

```
integrate(cot(dx+c)^4*(a+b*tan(dx+c))^3*(B*tan(dx+c)+C*tan(dx+c)^2),x,
algorithm="fricas")
```

output

$$\frac{-1/2(C^3 b^3 \log(1/(\tan(dx+c)^2+1)) \tan(dx+c)^2 + B^3 a^3 + (B^3 a^3 - 3 C^2 a^2 b - 3 B^2 a^2 b^2) \log(\tan(dx+c)^2/(\tan(dx+c)^2+1)) \tan(dx+c)^2 + (B^3 a^3 + 2(C^3 a^3 + 3 B^2 a^2 b - 3 C^2 a^2 b^2 - B^3 b^3) dx) \tan(dx+c)^2 + 2(C^3 a^3 + 3 B^2 a^2 b) \tan(dx+c)) / (d \tan(dx+c)^2)}$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(121) = 242.

Time = 2.46 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.99

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a+b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot^4(c) \\ \text{NaN} \\ \frac{B a^3 \log(\tan^2(c+dx)+1)}{2d} - \frac{B a^3 \log(\tan(c+dx))}{d} - \frac{B a^3}{2d \tan^2(c+dx)} - 3 B a^2 b x - \frac{3 B a^2 b}{d \tan(c+dx)} - \frac{3 B a b^2 \log(\tan^2(c+dx)+1)}{2d} + \end{cases}$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2)*cot(c)**4, Eq(d, 0)), (nan, Eq(c, -d*x)), (B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - B*a**3*log(tan(c + d*x))/d - B*a**3/(2*d*tan(c + d*x)**2) - 3*B*a**2*b*x - 3*B*a**2*b/(d*tan(c + d*x)) - 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a*b**2*log(tan(c + d*x))/d + B*b**3*x - C*a**3*x - C*a**3/(d*tan(c + d*x)) - 3*C*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*C*a**2*b*log(tan(c + d*x))/d + 3*C*a*b**2*x + C*b**3*log(tan(c + d*x)**2 + 1)/(2*d), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.12

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx = \frac{2(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c) - (Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(\tan(dx + c)^2 + 1)}{2d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")`

output `-1/2*(2*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*(d*x + c) - (B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^2 + 1) + 2*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2)*log(tan(d*x + c)) + (B*a^3 + 2*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/tan(d*x + c)^2)/d`

Giac [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.19

$$\begin{aligned}
& \int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
&= -\frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c)}{d} \\
&\quad + \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(\tan(dx + c)^2 + 1)}{2d} \\
&\quad - \frac{(Ba^3 - 3Ca^2b - 3Bab^2) \log(|\tan(dx + c)|)}{d} \\
&\quad - \frac{Ba^3 + 2(Ca^3 + 3Ba^2b) \tan(dx + c)}{2d \tan(dx + c)^2}
\end{aligned}$$

input

```
integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
-(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*(d*x + c)/d + 1/2*(B*a^3 - 3*C*a^2*b -
3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^2 + 1)/d - (B*a^3 - 3*C*a^2*b -
3*B*a*b^2)*log(abs(tan(d*x + c)))/d - 1/2*(B*a^3 + 2*(C*a^3 + 3*B*a^2*b)*t
an(d*x + c))/(d*tan(d*x + c)^2)
```

Mupad [B] (verification not implemented)

Time = 5.52 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.06

$$\begin{aligned}
& \int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\
&= \frac{\ln(\tan(c + dx)) (-Ba^3 + 3Ca^2b + 3Bab^2)}{d} \\
&\quad - \frac{\cot(c + dx)^2 \left(\tan(c + dx) (Ca^3 + 3Bba^2) + \frac{Ba^3}{2} \right)}{d} \\
&\quad + \frac{\ln(\tan(c + dx) + 1i) (B - C1i) (b + a1i)^3 1i}{2d} \\
&\quad + \frac{\ln(\tan(c + dx) - 1i) (B + C1i) (-b + a1i)^3 1i}{2d}
\end{aligned}$$

input `int(cot(c + d*x)^4*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^3,x)`

output `(log(tan(c + d*x))*(3*B*a*b^2 - B*a^3 + 3*C*a^2*b))/d - (cot(c + d*x)^2*(tan(c + d*x)*(C*a^3 + 3*B*a^2*b) + (B*a^3)/2))/d + (log(tan(c + d*x) + 1i)*(B - C*1i)*(a*1i + b)^3*1i)/(2*d) + (log(tan(c + d*x) - 1i)*(B + C*1i)*(a*1i - b)^3*1i)/(2*d)`

Reduce [F]

$$\begin{aligned} & \int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \int \cot(dx + c)^4 (a + \tan(dx + c)b)^3 (B \tan(dx + c) + C \tan(dx + c)^2) dx \end{aligned}$$

input `int(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `int(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

3.22 $\int \cot^5(c+dx)(a+b \tan(c+dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	272
Mathematica [C] (verified)	273
Rubi [A] (verified)	273
Maple [A] (verified)	278
Fricas [A] (verification not implemented)	278
Sympy [B] (verification not implemented)	279
Maxima [A] (verification not implemented)	280
Giac [A] (verification not implemented)	280
Mupad [B] (verification not implemented)	281
Reduce [F]	282

Optimal result

Integrand size = 40, antiderivative size = 154

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (a^3 B - 3ab^2 B - 3a^2 b C + b^3 C) x + \frac{a(3a^2 B - 8b^2 B - 9ab C) \cot(c + dx)}{3d}$$

$$- \frac{a^2(5bB + 3aC) \cot^2(c + dx)}{6d} - \frac{(3a^2 b B - b^3 B + a^3 C - 3ab^2 C) \log(\sin(c + dx))}{d}$$

$$- \frac{aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d}$$

output

```
(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*x+1/3*a*(3*B*a^2-8*B*b^2-9*C*a*b)*cot(d*x+c)/d-1/6*a^2*(5*B*b+3*C*a)*cot(d*x+c)^2/d-(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*ln(sin(d*x+c))/d-1/3*a*B*cot(d*x+c)^3*(a+b*tan(d*x+c))^2/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.84 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.06

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6a(a^2B - 3b^2B - 3abC) \cot(c + dx) - 3a^2(3bB + aC) \cot^2(c + dx) - 2a^3B \cot^3(c + dx) + 3(a + ib)^3(-$$

input

```
Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(6*a*(a^2*B - 3*b^2*B - 3*a*b*C)*Cot[c + d*x] - 3*a^2*(3*b*B + a*C)*Cot[c + d*x]^2 - 2*a^3*B*Cot[c + d*x]^3 + 3*(a + I*b)^3*((-I)*B + C)*Log[I - Tan[c + d*x]] - 6*(3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*Log[Tan[c + d*x]] + 3*(a - I*b)^3*(I*B + C)*Log[I + Tan[c + d*x]])/(6*d)
```

Rubi [A] (verified)

Time = 1.92 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 4115, 3042, 4088, 3042, 4118, 25, 3042, 4111, 27, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)^5} dx$$

$$\downarrow \text{4115}$$

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3 (B + C \tan(c + dx)) dx$$

$$\int \frac{(a + b \tan(c + dx))^3 (B + C \tan(c + dx))}{\tan(c + dx)^4} dx$$

↓ 3042

$$\frac{1}{3} \int \frac{\cot^3(c + dx)(a + b \tan(c + dx)) (-b(aB - 3bC) \tan^2(c + dx) - 3(Ba^2 - 2bCa - b^2B) \tan(c + dx) + a(5bB + 3aC)) dx - aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d}$$

↓ 4088

$$\frac{1}{3} \int \frac{(a + b \tan(c + dx)) (-b(aB - 3bC) \tan(c + dx)^2 - 3(Ba^2 - 2bCa - b^2B) \tan(c + dx) + a(5bB + 3aC))}{\tan(c + dx)^3} dx - \frac{aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\int -\cot^2(c + dx) (b^2(aB - 3bC) \tan^2(c + dx) + 3(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(3Ba^2 - 9bCa - 8b^2B)) dx - \frac{aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 4118

$$\frac{1}{3} \left(-\int \cot^2(c + dx) (b^2(aB - 3bC) \tan^2(c + dx) + 3(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(3Ba^2 - 9bCa - 8b^2B)) dx - \frac{aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 25

$$\frac{1}{3} \left(-\int \cot^2(c + dx) (b^2(aB - 3bC) \tan^2(c + dx) + 3(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(3Ba^2 - 9bCa - 8b^2B)) dx - \frac{aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(-\int \frac{b^2(aB - 3bC) \tan(c + dx)^2 + 3(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(3Ba^2 - 9bCa - 8b^2B)}{\tan(c + dx)^2} dx - \frac{aB \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 4111

$$\frac{1}{3} \left(- \int 3 \cot(c+dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)) dx + \frac{a(3a^2B - 9abC - 8b^3C)}{d} \right) + \frac{aB \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

↓ 27

$$\frac{1}{3} \left(-3 \int \cot(c+dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)) dx + \frac{a(3a^2B - 9abC - 8b^3C)}{d} \right) + \frac{aB \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-3 \int \frac{Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)}{\tan(c+dx)} dx + \frac{a(3a^2B - 9abC - 8b^3C)}{d} \right) + \frac{aB \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

↓ 4014

$$\frac{1}{3} \left(-3 \left((a^3C + 3a^2bB - 3ab^2C - b^3B) \int \cot(c+dx) dx - x(a^3B - 3a^2bC - 3ab^2B + b^3C) \right) + \frac{a(3a^2B - 9abC - 8b^3C)}{d} \right) + \frac{aB \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-3 \left((a^3C + 3a^2bB - 3ab^2C - b^3B) \int -\tan\left(c+dx + \frac{\pi}{2}\right) dx - x(a^3B - 3a^2bC - 3ab^2B + b^3C) \right) + \frac{a(3a^2B - 9abC - 8b^3C)}{d} \right) + \frac{aB \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

↓ 25

$$\frac{1}{3} \left(-3 \left(-(a^3C + 3a^2bB - 3ab^2C - b^3B) \int \tan\left(\frac{1}{2}(2c+\pi) + dx\right) dx - (x(a^3B - 3a^2bC - 3ab^2B + b^3C)) \right) + \frac{a(3a^2B - 9abC - 8b^3C)}{d} \right) + \frac{aB \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

↓ 3956

$$\frac{1}{3} \left(\frac{a(3a^2B - 9abC - 8b^2B) \cot(c + dx)}{d} - \frac{a^2(3aC + 5bB) \cot^2(c + dx)}{2d} - 3 \left(\frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \log(aB \cot^3(c + dx)(a + b \tan(c + dx))^2)}{3d} \right) \right)$$

input `Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]`

output `((a*(3*a^2*B - 8*b^2*B - 9*a*b*C)*Cot[c + d*x])/d - (a^2*(5*b*B + 3*a*C)*Cot[c + d*x]^2)/(2*d) - 3*(-((a^3*B - 3*a*b^2*B - 3*a^2*b*C + b^3*C)*x) + ((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*Log[-Sin[c + d*x]])/d))/3 - (a*B*Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2)/(3*d)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4088

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
  Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])

```

rule 4111

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2)  Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_
.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[1/b^2  Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4118

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_
.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f
_.)*(x_)])^(n_), x_Symbol] :> Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2))  Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]

```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.12

method	result
parallelrisc	$\frac{3(3B a^2 b - B b^3 + C a^3 - 3C a b^2) \ln(\sec(dx+c)^2) + 6(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \ln(\tan(dx+c)) - 2B \cot(dx+c)^3 a^3}{6d}$
derivativdivides	$\frac{\frac{(3B a^2 b - B b^3 + C a^3 - 3C a b^2) \ln(1+\tan(dx+c)^2)}{2} + (B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \arctan(\tan(dx+c)) + (-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \cot(dx+c)}{d}$
default	$\frac{\frac{(3B a^2 b - B b^3 + C a^3 - 3C a b^2) \ln(1+\tan(dx+c)^2)}{2} + (B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \arctan(\tan(dx+c)) + (-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \cot(dx+c)}{d}$
norman	$\frac{(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) x \tan(dx+c)^4 + \frac{a(B a^2 - 3B b^2 - 3C a b) \tan(dx+c)^3}{d} - \frac{B a^3 \tan(dx+c)}{3d} - \frac{a^2(3B b + C a) \tan(dx+c)}{2d}}{\tan(dx+c)^4}$
risc	$B a^3 x - 3B a b^2 x - 3C a^2 b x + C b^3 x + 3i B a^2 b x + i C a^3 x - \frac{6i C a b^2 c}{d} + \frac{2i C a^3 c}{d} - 3i C a b^2 c$

input `int(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method=_RETURNVERBOSE)`

output `1/6*(3*(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*ln(sec(d*x+c)^2)+6*(-3*B*a^2*b+B*b^3-C*a^3+3*C*a*b^2)*ln(tan(d*x+c))-2*B*cot(d*x+c)^3*a^3+3*(-3*B*a^2*b-C*a^3)*cot(d*x+c)^2+6*a*cot(d*x+c)*(B*a^2-3*B*b^2-3*C*a*b)+6*d*x*(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3))/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.18

$$\int \cot^5(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx = \frac{3(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 + 2Ba^3 + 3(Ca^3 + 3Ba^2b - 2(Ba^2b - Bb^3)) \cot(dx+c)}{d}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output

```
-1/6*(3*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^3 + 2*B*a^3 + 3*(C*a^3 + 3*B*a^2*b - 2*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*d*x)*tan(d*x + c)^3 - 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2)*tan(d*x + c)^2 + 3*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/(d*tan(d*x + c)^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 323 vs. $2(150) = 300$.

Time = 3.80 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.10

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot^5(c) \\ \text{NaN} \\ Ba^3x + \frac{Ba^3}{d \tan(c+dx)} - \frac{Ba^3}{3d \tan^3(c+dx)} + \frac{3Ba^2b \log(\tan^2(c+dx)+1)}{2d} - \frac{3Ba^2b \log(\tan(c+dx))}{d} - \frac{3Ba^2b}{2d \tan^2(c+dx)} - 3Bab^2x \end{cases}$$

input

```
integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2)*cot(c)**5, Eq(d, 0)), (nan, Eq(c, -d*x)), (B*a**3*x + B*a**3/(d*tan(c + d*x)) - B*a**3/(3*d*tan(c + d*x)**3) + 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a**2*b*log(tan(c + d*x))/d - 3*B*a**2*b/(2*d*tan(c + d*x)**2) - 3*B*a*b**2*x - 3*B*a*b**2/(d*tan(c + d*x)) - B*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**3*log(tan(c + d*x))/d + C*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - C*a**3*log(tan(c + d*x))/d - C*a**3/(2*d*tan(c + d*x)**2) - 3*C*a**2*b*x - 3*C*a**2*b/(d*tan(c + d*x)) - 3*C*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*C*a*b**2*log(tan(c + d*x))/d + C*b**3*x, True))
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.17

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx + c) + 3(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx + c)^2 + 1) - (2Ba^3 - 6(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \tan(dx + c)^2 + 3(Ca^3 + 3Ba^2b) \tan(dx + c)) / \tan(dx + c)^3}{d}$$

input

```
integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
1/6*(6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(d*x + c) + 3*(C*a^3 + 3*B*
a^2*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1) - 6*(C*a^3 + 3*B*a^2*b
- 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)) - (2*B*a^3 - 6*(B*a^3 - 3*C*a^2*b -
3*B*a*b^2)*tan(d*x + c)^2 + 3*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/tan(d*x +
c)^3)/d
```

Giac [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.21

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx + c)}{d} + \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx + c)^2 + 1)}{2d} - \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(|\tan(dx + c)|)}{d} - \frac{2Ba^3 - 6(Ba^3 - 3Ca^2b - 3Bab^2) \tan(dx + c)^2 + 3(Ca^3 + 3Ba^2b) \tan(dx + c)}{6d \tan(dx + c)^3}$$

input

```
integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(d*x + c)/d + 1/2*(C*a^3 + 3*B*a^2*b
*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1)/d - (C*a^3 + 3*B*a^2*b - 3
*C*a*b^2 - B*b^3)*log(abs(tan(d*x + c)))/d - 1/6*(2*B*a^3 - 6*(B*a^3 - 3*C
*a^2*b - 3*B*a*b^2)*tan(d*x + c)^2 + 3*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/(
d*tan(d*x + c)^3)
```

Mupad [B] (verification not implemented)

Time = 5.48 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.10

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (-C a^3 - 3 B a^2 b + 3 C a b^2 + B b^3)}{d}$$

$$- \frac{\cot(c + dx)^3 \left(\tan(c + dx) \left(\frac{C a^3}{2} + \frac{3 B b a^2}{2} \right) + \frac{B a^3}{3} + \tan(c + dx)^2 (-B a^3 + 3 C a^2 b + 3 B a b^2) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) - i) (B + C i) (a + b i)^3 i}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (B - C i) (a - b i)^3 i}{2 d}$$

input

```
int(cot(c + d*x)^5*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
)^3,x)
```

output

```
(log(tan(c + d*x))*(B*b^3 - C*a^3 - 3*B*a^2*b + 3*C*a*b^2))/d - (cot(c + d
*x)^3*(tan(c + d*x)*((C*a^3)/2 + (3*B*a^2*b)/2) + (B*a^3)/3 + tan(c + d*x)
^2*(3*B*a*b^2 - B*a^3 + 3*C*a^2*b))/d - (log(tan(c + d*x) - 1i)*(B + C*1i
)*(a + b*1i)^3*1i)/(2*d) + (log(tan(c + d*x) + 1i)*(B - C*1i)*(a - b*1i)^3
*1i)/(2*d)
```

Reduce [F]

$$\begin{aligned} & \int \cot^5(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \int \cot(dx + c)^5 (a + \tan(dx + c) b)^3 (B \tan(dx + c) + C \tan(dx + c)^2) dx \end{aligned}$$

input `int(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `int(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

3.23 $\int \cot^6(c+dx)(a+b \tan(c+dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	283
Mathematica [C] (verified)	284
Rubi [A] (verified)	284
Maple [A] (verified)	290
Fricas [A] (verification not implemented)	290
Sympy [B] (verification not implemented)	291
Maxima [A] (verification not implemented)	292
Giac [A] (verification not implemented)	292
Mupad [B] (verification not implemented)	293
Reduce [F]	294

Optimal result

Integrand size = 40, antiderivative size = 191

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= (3a^2bB - b^3B + a^3C - 3ab^2C) x + \frac{(3a^2bB - b^3B + a^3C - 3ab^2C) \cot(c + dx)}{d}$$

$$+ \frac{a(2a^2B - 5b^2B - 6abC) \cot^2(c + dx)}{4d} - \frac{a^2(3bB + 2aC) \cot^3(c + dx)}{6d}$$

$$+ \frac{(a^3B - 3ab^2B - 3a^2bC + b^3C) \log(\sin(c + dx))}{d}$$

$$- \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d}$$

output

```
(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*x+(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*cot(d*x+c)/d+1/4*a*(2*B*a^2-5*B*b^2-6*C*a*b)*cot(d*x+c)^2/d-1/6*a^2*(3*B*b+2*C*a)*cot(d*x+c)^3/d+(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*ln(sin(d*x+c))/d-1/4*a*B*cot(d*x+c)^4*(a+b*tan(d*x+c))^2/d
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.04

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{12(3a^2bB - b^3B + a^3C - 3ab^2C) \cot(c + dx) + 6a(a^2B - 3b^2B - 3abC) \cot^2(c + dx) - 4a^2(3bB + aC)}$$

input

```
Integrate[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(12*(3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*Cot[c + d*x] + 6*a*(a^2*B - 3*b^2*B - 3*a*b*C)*Cot[c + d*x]^2 - 4*a^2*(3*b*B + a*C)*Cot[c + d*x]^3 - 3*a^3*B*Cot[c + d*x]^4 - 6*(a + I*b)^3*(B + I*C)*Log[I - Tan[c + d*x]] + 12*(a^3*B - 3*a*b^2*B - 3*a^2*b*C + b^3*C)*Log[Tan[c + d*x]] - 6*(a - I*b)^3*(B - I*C)*Log[I + Tan[c + d*x]])/(12*d)
```

Rubi [A] (verified)

Time = 2.37 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.07, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.475$, Rules used = {3042, 4115, 3042, 4088, 27, 3042, 4118, 25, 3042, 4111, 27, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)^6} dx$$

$$\downarrow 4115$$

$$\begin{aligned}
& \int \cot^5(c+dx)(a+b \tan(c+dx))^3(B+C \tan(c+dx))dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+b \tan(c+dx))^3(B+C \tan(c+dx))}{\tan(c+dx)^5} dx \\
& \quad \downarrow \text{4088} \\
& \frac{1}{4} \int 2 \cot^4(c+dx)(a+b \tan(c+dx)) (-b(aB-2bC) \tan^2(c+dx) - 2(Ba^2-2bCa-b^2B) \tan(c+dx) + a(3bB+2aC)) dx - \\
& \quad \frac{aB \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \int \cot^4(c+dx)(a+b \tan(c+dx)) (-b(aB-2bC) \tan^2(c+dx) - 2(Ba^2-2bCa-b^2B) \tan(c+dx) + a(3bB+2aC)) dx - \\
& \quad \frac{aB \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{(a+b \tan(c+dx)) (-b(aB-2bC) \tan(c+dx)^2 - 2(Ba^2-2bCa-b^2B) \tan(c+dx) + a(3bB+2aC))}{\frac{aB \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \tan(c+dx)^4} dx \\
& \quad \downarrow \text{4118} \\
& \frac{1}{2} \left(\int -\cot^3(c+dx) (b^2(aB-2bC) \tan^2(c+dx) + 2(Ca^3+3bBa^2-3b^2Ca-b^3B) \tan(c+dx) + a(2Ba^2-6bCa-3b^2B)) dx - \right. \\
& \quad \left. \frac{aB \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \right) \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left(- \int \cot^3(c+dx) (b^2(aB-2bC) \tan^2(c+dx) + 2(Ca^3+3bBa^2-3b^2Ca-b^3B) \tan(c+dx) + a(2Ba^2-6bCa-3b^2B)) dx - \right. \\
& \quad \left. \frac{aB \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{2} \left(- \int \frac{b^2(aB - 2bC) \tan(c + dx)^2 + 2(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(2Ba^2 - 6bCa - 5b^2B)}{\tan(c + dx)^3} dx + \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right)$$

↓ 4111

$$\frac{1}{2} \left(- \int 2 \cot^2(c + dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c + dx)) dx + \frac{a(2a^2B - 6abC - 5b^2C)}{2d} + \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right)$$

↓ 27

$$\frac{1}{2} \left(-2 \int \cot^2(c + dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c + dx)) dx + \frac{a(2a^2B - 6abC - 5b^2C)}{2d} + \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right)$$

↓ 3042

$$\frac{1}{2} \left(-2 \int \frac{Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c + dx)}{\tan(c + dx)^2} dx + \frac{a(2a^2B - 6abC - 5b^2C)}{2d} + \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right)$$

↓ 4012

$$\frac{1}{2} \left(-2 \left(\int -\cot(c + dx) (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C + (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx)) dx - \frac{a^2}{2d} + \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \right)$$

↓ 25

$$\frac{1}{2} \left(-2 \left(- \int \cot(c + dx) (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C + (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx)) dx - \frac{a^2}{2d} + \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-2 \left(- \int \frac{Ba^3 - 3bCa^2 - 3b^2Ba + b^3C + (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx)}{\tan(c + dx)} dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B)}{d} \right) \right. \\ \left. \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \\ \downarrow 4014$$

$$\frac{1}{2} \left(-2 \left(-(a^3B - 3a^2bC - 3ab^2B + b^3C) \int \cot(c + dx) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot(c + dx)}{d} - (x \cot(c + dx) - \frac{x^2}{2d}) \right) \right. \\ \left. \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \\ \downarrow 3042$$

$$\frac{1}{2} \left(-2 \left(-(a^3B - 3a^2bC - 3ab^2B + b^3C) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot(c + dx)}{d} - (x \cot(c + dx) - \frac{x^2}{2d}) \right) \right. \\ \left. \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \\ \downarrow 25$$

$$\frac{1}{2} \left(-2 \left((a^3B - 3a^2bC - 3ab^2B + b^3C) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot(c + dx)}{d} - (x \cot(c + dx) - \frac{x^2}{2d}) \right) \right. \\ \left. \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \\ \downarrow 3956$$

$$\frac{1}{2} \left(\frac{a(2a^2B - 6abC - 5b^2B) \cot^2(c + dx)}{2d} - \frac{a^2(2aC + 3bB) \cot^3(c + dx)}{3d} - 2 \left(- \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B)}{d} \right) \right. \\ \left. \frac{aB \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right)$$

input

```
Int[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

$$\begin{aligned} & ((a*(2*a^2*B - 5*b^2*B - 6*a*b*C)*\text{Cot}[c + d*x]^2)/(2*d) - (a^2*(3*b*B + 2* \\ & a*C)*\text{Cot}[c + d*x]^3)/(3*d) - 2*(-((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)* \\ & x) - ((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*\text{Cot}[c + d*x])/d - ((a^3*B - \\ & 3*a*b^2*B - 3*a^2*b*C + b^3*C)*\text{Log}[-\text{Sin}[c + d*x]])/d))/2 - (a*B*\text{Cot}[c + d* \\ & x]^4*(a + b*\text{Tan}[c + d*x])^2)/(4*d) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{Ma} \\ \text{tchQ}[\text{Fx}, (b_)*(\text{Gx}_) \text{ ; FreeQ}[b, x]]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear} \\ \text{Q}[u, x]$$

rule 3956

$$\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d \\ *x], x]]/d, x] \text{ ; FreeQ}\{c, d\}, x]$$

rule 4012

$$\text{Int}[((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]^m)*((c_.) + (d_.)*\tan[(e_.) + \\ (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*((a + b*\text{Tan}[e + f*x])^{m+1}/ \\ (f*(m+1)*(a^2 + b^2))), x] + \text{Simp}[1/(a^2 + b^2) \quad \text{Int}[(a + b*\text{Tan}[e + f*x] \\)^{m+1}*\text{Simp}[a*c + b*d - (b*c - a*d)*\text{Tan}[e + f*x], x], x] \text{ ; FreeQ}\{a \\ , b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[m, -1 \\]$$

rule 4014

$$\text{Int}[((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*\tan[(e_.) + (f_. \\)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*c + b*d)*(x/(a^2 + b^2)), x] + \text{Simp}[(b*c - a \\ *d)/(a^2 + b^2) \quad \text{Int}[(b - a*\text{Tan}[e + f*x])/(a + b*\text{Tan}[e + f*x]), x], x] \text{ ;} \\ \text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{N} \\ \text{eQ}[a*c + b*d, 0]$$

rule 4088

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
  Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])

```

rule 4111

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :> Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2)  Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_
.) + (f_.)*(x_)])^2, x_Symbol] :> Simp[1/b^2  Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4118

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_
.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f
_.)*(x_)])^2, x_Symbol] :> Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2))  Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]

```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.09

method	result
parallelrisc	$6(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \ln(\sec(dx+c)^2) + 12(B a^3 - 3B a b^2 - 3C a^2 b + C b^3) \ln(\tan(dx+c)) - 3B \cot(dx+c)^4$
derivativedivides	$\frac{(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \ln(1 + \tan(dx+c)^2)}{2} + (3B a^2 b - B b^3 + C a^3 - 3C a b^2) \arctan(\tan(dx+c)) - \frac{3B a^2 b + B b^3 - C a^3}{\tan(dx+c)}$
default	$\frac{(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \ln(1 + \tan(dx+c)^2)}{2} + (3B a^2 b - B b^3 + C a^3 - 3C a b^2) \arctan(\tan(dx+c)) - \frac{3B a^2 b + B b^3 - C a^3}{\tan(dx+c)}$
norman	$\frac{(3B a^2 b - B b^3 + C a^3 - 3C a b^2) \tan(dx+c)^4}{d} + (3B a^2 b - B b^3 + C a^3 - 3C a b^2) x \tan(dx+c)^5 - \frac{B a^3 \tan(dx+c)}{4d} + \frac{a(B a^2 - 3B b^2)}{\tan(dx+c)^5}$
risc	$-iB a^3 x - \frac{2iB a^3 c}{d} + 3iB a b^2 x + 3iC a^2 b x + 3B a^2 b x - B b^3 x + C a^3 x - 3C a b^2 x + \dots$

input `int(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method =_RETURNVERBOSE)`

output `1/12*(6*(-B*a^3+3*B*a*b^2+3*C*a^2*b-C*b^3)*ln(sec(d*x+c)^2)+12*(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*ln(tan(d*x+c))-3*B*cot(d*x+c)^4*a^3+4*(-3*B*a^2*b-C*a^3)*cot(d*x+c)^3+6*a*cot(d*x+c)^2*(B*a^2-3*B*b^2-3*C*a*b)+12*cot(d*x+c)*(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)+36*x*(B*a^2*b-1/3*B*b^3+1/3*C*a^3-C*a*b^2)*d)/d`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.18

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{6(B a^3 - 3C a^2 b - 3B a b^2 + C b^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 + 3(3B a^3 - 6C a^2 b - 6B a b^2 + 4(C a^3 - 3B a b^2 + 3C a^2 b - C b^3)) \tan(dx+c)^3 + \dots}{d}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,algorithm="fricas")`

output

```
1/12*(6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^4 + 3*(3*B*a^3 - 6*C*a^2*b - 6*B*a*b^2 + 4*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*d*x)*tan(d*x + c)^4 - 3*B*a^3 + 12*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*tan(d*x + c)^3 + 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2)*tan(d*x + c)^2 - 4*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/(d*tan(d*x + c)^4)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 391 vs. $2(187) = 374$.

Time = 5.17 (sec) , antiderivative size = 391, normalized size of antiderivative = 2.05

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot^6(c) \\ \text{NaN} \\ -\frac{Ba^3 \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba^3 \log(\tan(c+dx))}{d} + \frac{Ba^3}{2d \tan^2(c+dx)} - \frac{Ba^3}{4d \tan^4(c+dx)} + 3Ba^2bx + \frac{3Ba^2b}{d \tan(c+dx)} - \frac{Ba^2b}{d \tan^3(c+dx)} \end{cases}$$

input

```
integrate(cot(d*x+c)**6*(a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2)*cot(c)**6, Eq(d, 0)), (nan, Eq(c, -d*x)), (-B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**3*log(tan(c + d*x))/d + B*a**3/(2*d*tan(c + d*x)**2) - B*a**3/(4*d*tan(c + d*x)**4) + 3*B*a**2*b*x + 3*B*a**2*b/(d*tan(c + d*x)) - B*a**2*b/(d*tan(c + d*x)**3) + 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a*b**2*log(tan(c + d*x))/d - 3*B*a*b**2/(2*d*tan(c + d*x)**2) - B*b**3*x - B*b**3/(d*tan(c + d*x)) + C*a**3*x + C*a**3/(d*tan(c + d*x)) - C*a**3/(3*d*tan(c + d*x)**3) + 3*C*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) - 3*C*a**2*b*log(tan(c + d*x))/d - 3*C*a**2*b/(2*d*tan(c + d*x)**2) - 3*C*a*b**2*x - 3*C*a*b**2/(d*tan(c + d*x)) - C*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + C*b**3*log(tan(c + d*x))/d, True))
```


Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.13

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{12(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c) - 6(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(\tan(dx + c)^2 + 1)}{d}$$

input

```
integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
1/12*(12*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*(d*x + c) - 6*(B*a^3 - 3*
C*a^2*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^2 + 1) + 12*(B*a^3 - 3*C*a^2
*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)) - (3*B*a^3 - 12*(C*a^3 + 3*B*a^2
*b - 3*C*a*b^2 - B*b^3)*tan(d*x + c)^3 - 6*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2)
*tan(d*x + c)^2 + 4*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/tan(d*x + c)^4/d
```

Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.16

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3)(dx + c)}{d}$$

$$- \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(\tan(dx + c)^2 + 1)}{d}$$

$$+ \frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \log(|\tan(dx + c)|)}{d}$$

$$- \frac{3Ba^3 - 12(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \tan(dx + c)^3 - 6(Ba^3 - 3Ca^2b - 3Bab^2) \tan(dx + c)^2}{12d \tan(dx + c)^4}$$

input

```
integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*(d*x + c)/d - 1/2*(B*a^3 - 3*C*a^2*b
*b - 3*B*a*b^2 + C*b^3)*log(tan(d*x + c)^2 + 1)/d + (B*a^3 - 3*C*a^2*b - 3
*B*a*b^2 + C*b^3)*log(abs(tan(d*x + c)))/d - 1/12*(3*B*a^3 - 12*(C*a^3 + 3
*B*a^2*b - 3*C*a*b^2 - B*b^3)*tan(d*x + c)^3 - 6*(B*a^3 - 3*C*a^2*b - 3*B*
a*b^2)*tan(d*x + c)^2 + 4*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/(d*tan(d*x + c
)^4)
```

Mupad [B] (verification not implemented)

Time = 5.47 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.07

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (B a^3 - 3 C a^2 b - 3 B a b^2 + C b^3)}{d}$$

$$- \frac{\cot(c + dx)^4 \left(\tan(c + dx) \left(\frac{C a^3}{3} + B b a^2 \right) + \frac{B a^3}{4} + \tan(c + dx)^2 \left(-\frac{B a^3}{2} + \frac{3 C a^2 b}{2} + \frac{3 B a b^2}{2} \right) + \tan(c + dx) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) + i) (B - C i) (b + a i)^3 i}{2 d}$$

$$- \frac{\ln(\tan(c + dx) - i) (B + C i) (-b + a i)^3 i}{2 d}$$

input

```
int(cot(c + d*x)^6*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x)
)^3,x)
```

output

```
(log(tan(c + d*x))*(B*a^3 + C*b^3 - 3*B*a*b^2 - 3*C*a^2*b))/d - (cot(c + d
*x)^4*(tan(c + d*x)*((C*a^3)/3 + B*a^2*b) + (B*a^3)/4 + tan(c + d*x)^2*((3
*B*a*b^2)/2 - (B*a^3)/2 + (3*C*a^2*b)/2) + tan(c + d*x)^3*(B*b^3 - C*a^3 -
3*B*a^2*b + 3*C*a*b^2))/d - (log(tan(c + d*x) + i)*(B - C*i)*(a*i + b
)^3*i)/(2*d) - (log(tan(c + d*x) - i)*(B + C*i)*(a*i - b)^3*i)/(2*d)
```

Reduce [F]

$$\begin{aligned} & \int \cot^6(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \int \cot(dx + c)^6 (a + \tan(dx + c)b)^3 (B \tan(dx + c) + C \tan(dx + c)^2) dx \end{aligned}$$

input `int(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `int(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

3.24 $\int \cot^7(c+dx)(a+b \tan(c+dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	295
Mathematica [C] (verified)	296
Rubi [A] (verified)	296
Maple [A] (verified)	302
Fricas [A] (verification not implemented)	303
Sympy [A] (verification not implemented)	303
Maxima [A] (verification not implemented)	304
Giac [A] (verification not implemented)	305
Mupad [B] (verification not implemented)	305
Reduce [F]	306

Optimal result

Integrand size = 40, antiderivative size = 233

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= -((a^3B - 3ab^2B - 3a^2bC + b^3C) x) - \frac{(a^3B - 3ab^2B - 3a^2bC + b^3C) \cot(c + dx)}{d}$$

$$+ \frac{(3a^2bB - b^3B + a^3C - 3ab^2C) \cot^2(c + dx)}{2d}$$

$$+ \frac{a(5a^2B - 12b^2B - 15abC) \cot^3(c + dx)}{15d} - \frac{a^2(7bB + 5aC) \cot^4(c + dx)}{20d}$$

$$+ \frac{(3a^2bB - b^3B + a^3C - 3ab^2C) \log(\sin(c + dx))}{d}$$

$$- \frac{aB \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d}$$

output

```

-(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*x-(B*a^3-3*B*a*b^2-3*C*a^2*b+C*b^3)*cot
(d*x+c)/d+1/2*(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*cot(d*x+c)^2/d+1/15*a*(5*B
*a^2-12*B*b^2-15*C*a*b)*cot(d*x+c)^3/d-1/20*a^2*(7*B*b+5*C*a)*cot(d*x+c)^4
/d+(3*B*a^2*b-B*b^3+C*a^3-3*C*a*b^2)*ln(sin(d*x+c))/d-1/5*a*B*cot(d*x+c)^5
*(a+b*tan(d*x+c))^2/d
    
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.76 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.02

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{-60(a^3 B - 3ab^2 B - 3a^2 b C + b^3 C) \cot(c + dx) + 30(3a^2 b B - b^3 B + a^3 C - 3ab^2 C) \cot^2(c + dx) + 20a(a^2 B - 3ab^2 B - 3a^2 b C + b^3 C) \cot^3(c + dx) + 15a^2(3b^2 B + a^2 C) \cot^4(c + dx) + 12a^3 B \cot^5(c + dx) + (30I)(a + I b)^3 (B + I C) \operatorname{Log}[I - \tan(c + dx)] + 60(3a^2 b^2 B - b^3 B + a^3 C - 3a^2 b^2 C) \operatorname{Log}[\tan(c + dx)] + 30(I a + b)^3 (B - I C) \operatorname{Log}[I + \tan(c + dx)]}{60d}$$

input

```
Integrate[Cot[c + d*x]^7*(a + b*Tan[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(-60*(a^3*B - 3*a*b^2*B - 3*a^2*b*C + b^3*C)*Cot[c + d*x] + 30*(3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*Cot[c + d*x]^2 + 20*a*(a^2*B - 3*b^2*B - 3*a*b*C)*Cot[c + d*x]^3 - 15*a^2*(3*b*B + a*C)*Cot[c + d*x]^4 - 12*a^3*B*Cot[c + d*x]^5 + (30*I)*(a + I*b)^3*(B + I*C)*Log[I - Tan[c + d*x]] + 60*(3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*Log[Tan[c + d*x]] + 30*(I*a + b)^3*(B - I*C)*Log[I + Tan[c + d*x]])/(60*d)
```

Rubi [A] (verified)

Time = 2.84 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.04, number of steps used = 20, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4115, 3042, 4088, 3042, 4118, 25, 3042, 4111, 27, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)^2)}{\tan(c + dx)^7} dx$$

$$\downarrow \text{4115}$$

$$\begin{aligned}
 & \int \cot^6(c + dx)(a + b \tan(c + dx))^3(B + C \tan(c + dx))dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3(B + C \tan(c + dx))}{\tan(c + dx)^6} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{1}{5} \int \frac{\cot^5(c + dx)(a + b \tan(c + dx) - b(3aB - 5bC) \tan^2(c + dx) - 5(Ba^2 - 2bCa - b^2B) \tan(c + dx) + a(7bB + 5aC))}{aB \cot^5(c + dx)(a + b \tan(c + dx))^2} dx - \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5} \int \frac{(a + b \tan(c + dx))(-b(3aB - 5bC) \tan(c + dx)^2 - 5(Ba^2 - 2bCa - b^2B) \tan(c + dx) + a(7bB + 5aC))}{aB \cot^5(c + dx)(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4118} \\
 & \frac{1}{5} \left(\int -\cot^4(c + dx) (b^2(3aB - 5bC) \tan^2(c + dx) + 5(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(5Ba^2 - 15bCa - 12b^2)) \right. \\
 & \quad \left. \frac{aB \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{5} \left(- \int \cot^4(c + dx) (b^2(3aB - 5bC) \tan^2(c + dx) + 5(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(5Ba^2 - 15bCa - 12b^2)) \right. \\
 & \quad \left. \frac{aB \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5} \left(- \int \frac{b^2(3aB - 5bC) \tan(c + dx)^2 + 5(Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c + dx) + a(5Ba^2 - 15bCa - 12b^2)}{\tan(c + dx)^4} \right. \\
 & \quad \left. \frac{aB \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \\
 & \quad \downarrow \text{4111}
 \end{aligned}$$

$$\frac{1}{5} \left(- \int 5 \cot^3(c+dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)) dx + \frac{a(5a^2)}{5d} \right)$$

$$\frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 27

$$\frac{1}{5} \left(-5 \int \cot^3(c+dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)) dx + \frac{a(5a^2)}{5d} \right)$$

$$\frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(-5 \int \frac{Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)}{\tan(c+dx)^3} dx + \frac{a(5a^2B - 15abC - 1)}{5d} \right)$$

$$\frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 4012

$$\frac{1}{5} \left(-5 \left(\int -\cot^2(c+dx) (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C + (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c+dx)) dx - \frac{a(5a^2)}{5d} \right) \right)$$

$$\frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 25

$$\frac{1}{5} \left(-5 \left(- \int \cot^2(c+dx) (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C + (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c+dx)) dx - \frac{a(5a^2)}{5d} \right) \right)$$

$$\frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(-5 \left(- \int \frac{Ba^3 - 3bCa^2 - 3b^2Ba + b^3C + (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B) \tan(c+dx)}{\tan(c+dx)^2} dx - \frac{(a^3C + 3a^2bB - 1)}{5d} \right) \right)$$

$$\frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 4012

$$\frac{1}{5} \left(-5 \left(- \int \cot(c+dx) (Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot^2(c+dx)}{2d} + \frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right) \right)$$

\downarrow 3042

$$\frac{1}{5} \left(-5 \left(- \int \frac{Ca^3 + 3bBa^2 - 3b^2Ca - b^3B - (Ba^3 - 3bCa^2 - 3b^2Ba + b^3C) \tan(c+dx)}{\tan(c+dx)} dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot^2(c+dx)}{2d} + \frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right) \right)$$

\downarrow 4014

$$\frac{1}{5} \left(-5 \left(-(a^3C + 3a^2bB - 3ab^2C - b^3B) \int \cot(c+dx) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot^2(c+dx)}{2d} + \frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right) \right)$$

\downarrow 3042

$$\frac{1}{5} \left(-5 \left(-(a^3C + 3a^2bB - 3ab^2C - b^3B) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot^2(c+dx)}{2d} + \frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right) \right)$$

\downarrow 25

$$\frac{1}{5} \left(-5 \left((a^3C + 3a^2bB - 3ab^2C - b^3B) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot^2(c+dx)}{2d} + \frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right) \right)$$

\downarrow 3956

$$\frac{1}{5} \left(\frac{a(5a^2B - 15abC - 12b^2B) \cot^3(c+dx)}{3d} - \frac{a^2(5aC + 7bB) \cot^4(c+dx)}{4d} - 5 \left(- \frac{(a^3C + 3a^2bB - 3ab^2C - b^3B) \cot^2(c+dx)}{2d} + \frac{aB \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right) \right)$$

input $\text{Int}[\text{Cot}[c + d*x]^7*(a + b*\text{Tan}[c + d*x])^3*(B*\text{Tan}[c + d*x] + C*\text{Tan}[c + d*x]^2), x]$

output $((a*(5*a^2*B - 12*b^2*B - 15*a*b*C)*\text{Cot}[c + d*x]^3)/(3*d) - (a^2*(7*b*B + 5*a*C)*\text{Cot}[c + d*x]^4)/(4*d) - 5*((a^3*B - 3*a*b^2*B - 3*a^2*b*C + b^3*C)*x + ((a^3*B - 3*a*b^2*B - 3*a^2*b*C + b^3*C)*\text{Cot}[c + d*x])/d - ((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*\text{Cot}[c + d*x]^2)/(2*d) - ((3*a^2*b*B - b^3*B + a^3*C - 3*a*b^2*C)*\text{Log}[-\text{Sin}[c + d*x]])/d)/5 - (a*B*\text{Cot}[c + d*x]^5*(a + b*\text{Tan}[c + d*x])^2)/(5*d)$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_*)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\text{tan}[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$

rule 4012 $\text{Int}[(a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)]^{(m_)}*((c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*((a + b*\text{Tan}[e + f*x])^{(m + 1)}/(f*(m + 1)*(a^2 + b^2))), x] + \text{Simp}[1/(a^2 + b^2) \quad \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m + 1)}*\text{Simp}[a*c + b*d - (b*c - a*d)*\text{Tan}[e + f*x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[m, -1]$

rule 4014

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)], x_Symbol] :> Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

rule 4088

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e
_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)^2], x_Symbol] :> Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.04

method	result
parallelrisch	$\frac{(-90B a^2 b + 30B b^3 - 30C a^3 + 90C a b^2) \ln(\sec(dx+c)^2) + (180B a^2 b - 60B b^3 + 60C a^3 - 180C a b^2) \ln(\tan(dx+c)) - 12B a^2 b^2 - 12B a^2 b^3 - 12C a^3 b^2 - 12C a^2 b^3}{2 \tan(dx+c)}$
derivativedivides	$\frac{(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \ln(1 + \tan(dx+c)^2)}{2} + (-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \arctan(\tan(dx+c)) - \frac{-3B a^2 b + B b^3 - 3C a^3 + 3C a b^2}{2 \tan(dx+c)}$
default	$\frac{(-3B a^2 b + B b^3 - C a^3 + 3C a b^2) \ln(1 + \tan(dx+c)^2)}{2} + (-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) \arctan(\tan(dx+c)) - \frac{-3B a^2 b + B b^3 - 3C a^3 + 3C a b^2}{2 \tan(dx+c)}$
norman	$\frac{(-B a^3 + 3B a b^2 + 3C a^2 b - C b^3) x \tan(dx+c)^6 + \frac{(3B a^2 b - B b^3 + C a^3 - 3C a b^2) \tan(dx+c)^4}{2d} - \frac{(B a^3 - 3B a b^2 - 3C a^2 b + C b^3)}{\tan(dx+c)^6}}{\tan(dx+c)^6}$
risch	$-B a^3 x + 3B a b^2 x + 3C a^2 b x - C b^3 x - 3iB a^2 b x - iC a^3 x - \frac{2i(-60C a^2 b + 15C b^3 + 23B a^3)}{d}$

input

```
int(cot(d*x+c)^7*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,method
=_RETURNVERBOSE)
```

output

```
1/60*((-90*B*a^2*b+30*B*b^3-30*C*a^3+90*C*a*b^2)*ln(sec(d*x+c)^2)+(180*B*a
^2*b-60*B*b^3+60*C*a^3-180*C*a*b^2)*ln(tan(d*x+c))-12*B*cot(d*x+c)^5*a^3+(
-45*B*a^2*b-15*C*a^3)*cot(d*x+c)^4+20*a*cot(d*x+c)^3*(B*a^2-3*B*b^2-3*C*a*
b)+(90*B*a^2*b-30*B*b^3+30*C*a^3-90*C*a*b^2)*cot(d*x+c)^2+(-60*B*a^3+180*B
*a*b^2+180*C*a^2*b-60*C*b^3)*cot(d*x+c)-60*d*x*(B*a^3-3*B*a*b^2-3*C*a^2*b+
C*b^3))/d
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.14

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{30(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^5 + 15(3Ca^3 + 9Ba^2b - 6Cab^2 - 2Bb^3) \tan(dx+c)^4 + 15(3Ca^3 + 9Ba^2b - 6Cab^2 - 2Bb^3) \tan(dx+c)^3 + 15(3Ca^3 + 9Ba^2b - 6Cab^2 - 2Bb^3) \tan(dx+c)^2 + 15(3Ca^3 + 9Ba^2b - 6Cab^2 - 2Bb^3) \tan(dx+c) + 15(3Ca^3 + 9Ba^2b - 6Cab^2 - 2Bb^3)}{d}$$

input

```
integrate(cot(d*x+c)^7*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="fricas")
```

output

```
1/60*(30*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)^2/(tan(d
*x + c)^2 + 1))*tan(d*x + c)^5 + 15*(3*C*a^3 + 9*B*a^2*b - 6*C*a*b^2 - 2*B
*b^3 - 4*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*d*x)*tan(d*x + c)^5 - 60*
(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*tan(d*x + c)^4 - 12*B*a^3 + 30*(C*
a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*tan(d*x + c)^3 + 20*(B*a^3 - 3*C*a^2*
b - 3*B*a*b^2)*tan(d*x + c)^2 - 15*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/(d*ta
n(d*x + c)^5)
```

Sympy [A] (verification not implemented)

Time = 13.57 (sec) , antiderivative size = 462, normalized size of antiderivative = 1.98

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \begin{cases} \text{NaN} \\ x(a + b \tan(c))^3 (B \tan(c) + C \tan^2(c)) \cot^7(c) \\ \text{NaN} \\ -Ba^3x - \frac{Ba^3}{d \tan(c+dx)} + \frac{Ba^3}{3d \tan^3(c+dx)} - \frac{Ba^3}{5d \tan^5(c+dx)} - \frac{3Ba^2b \log(\tan^2(c+dx)+1)}{2d} + \frac{3Ba^2b \log(\tan(c+dx))}{d} + \frac{3Ba^2b}{2d \tan^2(c+dx)} \end{cases}$$

input

```
integrate(cot(d*x+c)**7*(a+b*tan(d*x+c))**3*(B*tan(d*x+c)+C*tan(d*x+c)**2),x)
```

output

```
Piecewise((nan, Eq(c, 0) & Eq(d, 0)), (x*(a + b*tan(c))**3*(B*tan(c) + C*tan(c)**2)*cot(c)**7, Eq(d, 0)), (nan, Eq(c, -d*x)), (-B*a**3*x - B*a**3/(d*tan(c + d*x)) + B*a**3/(3*d*tan(c + d*x)**3) - B*a**3/(5*d*tan(c + d*x)**5) - 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a**2*b*log(tan(c + d*x))/d + 3*B*a**2*b/(2*d*tan(c + d*x)**2) - 3*B*a**2*b/(4*d*tan(c + d*x)**4) + 3*B*a*b**2*x + 3*B*a*b**2/(d*tan(c + d*x)) - B*a*b**2/(d*tan(c + d*x)**3) + B*b**3*log(tan(c + d*x)**2 + 1)/(2*d) - B*b**3*log(tan(c + d*x))/d - B*b**3/(2*d*tan(c + d*x)**2) - C*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + C*a**3*log(tan(c + d*x))/d + C*a**3/(2*d*tan(c + d*x)**2) - C*a**3/(4*d*tan(c + d*x)**4) + 3*C*a**2*b*x + 3*C*a**2*b/(d*tan(c + d*x)) - C*a**2*b/(d*tan(c + d*x)**3) + 3*C*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - 3*C*a*b**2*log(tan(c + d*x))/d - 3*C*a*b**2/(2*d*tan(c + d*x)**2) - C*b**3*x - C*b**3/(d*tan(c + d*x)), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.07

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx = \frac{60 (Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx + c) + 30 (Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx + c)^2 + 1) - 60 (Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx + c)) + (60 (Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \tan(dx + c)^4 + 12Ba^3 - 30 (Ca^3 + 3Ba^2b - 3Ca^2b^2 - Bb^3) \tan(dx + c)^3 - 20 (Ba^3 - 3Ca^2b - 3Bab^2) \tan(dx + c)^2 + 15 (Ca^3 + 3Ba^2b) \tan(dx + c)) / \tan(dx + c)^5}{d}$$

input

```
integrate(cot(d*x+c)^7*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
-1/60*(60*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(d*x + c) + 30*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1) - 60*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)) + (60*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*tan(d*x + c)^4 + 12*B*a^3 - 30*(C*a^3 + 3*B*a^2*b - 3*C*a*b^2 - B*b^3)*tan(d*x + c)^3 - 20*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2)*tan(d*x + c)^2 + 15*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/tan(d*x + c)^5)/d
```

Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.11

$$\int \cot^7(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= -\frac{(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3)(dx+c)}{d}$$

$$- \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(\tan(dx+c)^2 + 1)}{2d}$$

$$+ \frac{(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \log(|\tan(dx+c)|)}{d}$$

$$- \frac{60(Ba^3 - 3Ca^2b - 3Bab^2 + Cb^3) \tan(dx+c)^4 + 12Ba^3 - 30(Ca^3 + 3Ba^2b - 3Cab^2 - Bb^3) \tan(dx+c)}{60d \tan(dx+c)}$$

input

```
integrate(cot(d*x+c)^7*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
-(B*a^3 - 3*C*a^2*b - 3*B*a*b^2 + C*b^3)*(d*x + c)/d - 1/2*(C*a^3 + 3*B*a^2*b -
2*b - 3*C*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1)/d + (C*a^3 + 3*B*a^2*b -
3*C*a*b^2 - B*b^3)*log(abs(tan(d*x + c)))/d - 1/60*(60*(B*a^3 - 3*C*a^2*b
- 3*B*a*b^2 + C*b^3)*tan(d*x + c)^4 + 12*B*a^3 - 30*(C*a^3 + 3*B*a^2*b - 3
*C*a*b^2 - B*b^3)*tan(d*x + c)^3 - 20*(B*a^3 - 3*C*a^2*b - 3*B*a*b^2)*tan(
d*x + c)^2 + 15*(C*a^3 + 3*B*a^2*b)*tan(d*x + c))/(d*tan(d*x + c)^5)
```

Mupad [B] (verification not implemented)

Time = 5.51 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.02

$$\int \cot^7(c+dx)(a+b \tan(c+dx))^3 (B \tan(c+dx) + C \tan^2(c+dx)) dx =$$

$$\frac{\cot(c+dx)^5 \left(\tan(c+dx) \left(\frac{Ca^3}{4} + \frac{3Bba^2}{4} \right) + \frac{Ba^3}{5} + \tan(c+dx)^2 \left(-\frac{Ba^3}{3} + Ca^2b + Bab^2 \right) + \tan(c+dx) \right)}{d}$$

$$- \frac{\ln(\tan(c+dx)) (-Ca^3 - 3Ba^2b + 3Cab^2 + Bb^3)}{d}$$

$$+ \frac{\ln(\tan(c+dx) - i) (B + C i) (a + b i)^3 i}{2d}$$

$$- \frac{\ln(\tan(c+dx) + i) (B - C i) (a - b i)^3 i}{2d}$$

input `int(cot(c + d*x)^7*(B*tan(c + d*x) + C*tan(c + d*x)^2)*(a + b*tan(c + d*x))^3,x)`

output `(log(tan(c + d*x) - 1i)*(B + C*1i)*(a + b*1i)^3*1i)/(2*d) - (log(tan(c + d*x))*(B*b^3 - C*a^3 - 3*B*a^2*b + 3*C*a*b^2))/d - (cot(c + d*x)^5*(tan(c + d*x)*((C*a^3)/4 + (3*B*a^2*b)/4) + (B*a^3)/5 + tan(c + d*x)^2*(B*a*b^2 - (B*a^3)/3 + C*a^2*b) + tan(c + d*x)^4*(B*a^3 + C*b^3 - 3*B*a*b^2 - 3*C*a^2*b) + tan(c + d*x)^3*((B*b^3)/2 - (C*a^3)/2 - (3*B*a^2*b)/2 + (3*C*a*b^2)/2))/d - (log(tan(c + d*x) + 1i)*(B - C*1i)*(a - b*1i)^3*1i)/(2*d)`

Reduce [F]

$$\begin{aligned} & \int \cot^7(c + dx)(a + b \tan(c + dx))^3 (B \tan(c + dx) + C \tan^2(c + dx)) dx \\ &= \int \cot(dx + c)^7 (a + \tan(dx + c)b)^3 (B \tan(dx + c) + C \tan(dx + c)^2) dx \end{aligned}$$

input `int(cot(d*x+c)^7*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `int(cot(d*x+c)^7*(a+b*tan(d*x+c))^3*(B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

3.25 $\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$

Optimal result	307
Mathematica [C] (verified)	307
Rubi [A] (verified)	308
Maple [A] (verified)	312
Fricas [A] (verification not implemented)	313
Sympy [C] (verification not implemented)	313
Maxima [A] (verification not implemented)	314
Giac [A] (verification not implemented)	315
Mupad [B] (verification not implemented)	315
Reduce [B] (verification not implemented)	316

Optimal result

Integrand size = 40, antiderivative size = 127

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(bB-aC)x}{a^2+b^2} + \frac{(aB+bC) \log(\cos(c+dx))}{(a^2+b^2)d}$$

$$- \frac{a^3(bB-aC) \log(a+b \tan(c+dx))}{b^3(a^2+b^2)d} + \frac{(bB-aC) \tan(c+dx)}{b^2d} + \frac{C \tan^2(c+dx)}{2bd}$$

output

```
- (B*b-C*a)*x/(a^2+b^2)+(B*a+C*b)*ln(cos(d*x+c))/(a^2+b^2)/d-a^3*(B*b-C*a)*
ln(a+b*tan(d*x+c))/b^3/(a^2+b^2)/d+(B*b-C*a)*tan(d*x+c)/b^2/d+1/2*C*tan(d*
x+c)^2/b/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.01 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.09

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{-\frac{b(B+iC) \log(i-\tan(c+dx))}{a+ib} - \frac{b(B-iC) \log(i+\tan(c+dx))}{a-ib} + \frac{2a^3(-bB+aC) \log(a+b \tan(c+dx))}{b^2(a^2+b^2)} + \frac{2(bB-aC) \tan(c+dx)}{b} + C \tan^2(c+dx)}{2bd}$$

input

```
Integrate[(Tan[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[
c + d*x]),x]
```

output

```
((-(b*(B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)) - (b*(B - I*C)*Log[I + T
an[c + d*x]])/(a - I*b) + (2*a^3*(-(b*B) + a*C)*Log[a + b*Tan[c + d*x]])/(
b^2*(a^2 + b^2)) + (2*(b*B - a*C)*Tan[c + d*x])/b + C*Tan[c + d*x]^2)/(2*b
*d)
```

Rubi [A] (verified)

Time = 1.77 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.13, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.325$, Rules used = {3042, 4115, 3042, 4090, 27, 3042, 4130, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx) (B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2 (B \tan(c+dx) + C \tan(c+dx)^2)}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \frac{\tan^3(c+dx)(B + C \tan(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^3(B + C \tan(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{\int -\frac{2 \tan(c+dx) ((bB-aC) \tan^2(c+dx) + bC \tan(c+dx) + aC)}{a+b \tan(c+dx)} dx}{2b} + \frac{C \tan^2(c+dx)}{2bd} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\int \frac{\tan(c+dx) \left(-((bB-aC) \tan^2(c+dx)) + bC \tan(c+dx) + aC \right)}{a+b \tan(c+dx)} dx}{b} \\
& \quad \downarrow \text{3042} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\int \frac{\tan(c+dx) \left(-((bB-aC) \tan(c+dx)^2) + bC \tan(c+dx) + aC \right)}{a+b \tan(c+dx)} dx}{b} \\
& \quad \downarrow \text{4130} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\int \frac{B \tan(c+dx) b^2 + (-Ca^2 + bBa + b^2C) \tan^2(c+dx) + a(bB-aC)}{a+b \tan(c+dx)} dx}{b} - \frac{(bB-aC) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{3042} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\int \frac{B \tan(c+dx) b^2 + (-Ca^2 + bBa + b^2C) \tan(c+dx)^2 + a(bB-aC)}{a+b \tan(c+dx)} dx}{b} - \frac{(bB-aC) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{4109} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\frac{b^2(aB+bC) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^3(bB-aC) \int \frac{\tan^2(c+dx)+1}{a+b \tan(c+dx)} dx}{b} + \frac{b^2x(bB-aC)}{a^2+b^2}}{b} - \frac{(bB-aC) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{3042} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\frac{b^2(aB+bC) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^3(bB-aC) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{b} + \frac{b^2x(bB-aC)}{a^2+b^2}}{b} - \frac{(bB-aC) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{3956} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\frac{a^3(bB-aC) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{b^2(aB+bC) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2x(bB-aC)}{a^2+b^2}}{b} - \frac{(bB-aC) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{4100} \\
& \frac{C \tan^2(c+dx)}{2bd} - \frac{\frac{a^3(bB-aC) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} - \frac{b^2(aB+bC) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2x(bB-aC)}{a^2+b^2}}{b} - \frac{(bB-aC) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{16}
\end{aligned}$$

$$\frac{C \tan^2(c + dx)}{2bd} - \frac{-\frac{b^2(aB+bC) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2x(bB-aC)}{a^2+b^2} + \frac{a^3(bB-aC) \log(a+b \tan(c+dx))}{bd(a^2+b^2)}}{b} - \frac{(bB-aC) \tan(c+dx)}{bd}$$

input `Int[(Tan[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]`

output `(C*Tan[c + d*x]^2)/(2*b*d) - (((b^2*(b*B - a*C)*x)/(a^2 + b^2) - (b^2*(a*B + b*C)*Log[Cos[c + d*x]]))/(a^2 + b^2)*d) + (a^3*(b*B - a*C)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b - ((b*B - a*C)*Tan[c + d*x])/(b*d))/b`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4090

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4100

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]

```

rule 4109

```

Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2
)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_
.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{\frac{C \tan(dx+c)^2 b + B \tan(dx+c)b - C \tan(dx+c)a}{b^2} - \frac{a^3(Bb-Ca) \ln(a+b \tan(dx+c))}{b^3(a^2+b^2)} + \frac{(-Ba-Cb) \ln(1+\tan(dx+c)^2)}{2} + \frac{(-Bb+Ca) \arctan(\tan(dx+c))}{a^2+b^2}}{d}$
default	$\frac{\frac{C \tan(dx+c)^2 b + B \tan(dx+c)b - C \tan(dx+c)a}{b^2} - \frac{a^3(Bb-Ca) \ln(a+b \tan(dx+c))}{b^3(a^2+b^2)} + \frac{(-Ba-Cb) \ln(1+\tan(dx+c)^2)}{2} + \frac{(-Bb+Ca) \arctan(\tan(dx+c))}{a^2+b^2}}{d}$
norman	$\frac{(Bb-Ca) \tan(dx+c)}{b^2 d} - \frac{(Bb-Ca)x}{a^2+b^2} + \frac{C \tan(dx+c)^2}{2bd} - \frac{(Ba+Cb) \ln(1+\tan(dx+c)^2)}{2d(a^2+b^2)} - \frac{a^3(Bb-Ca) \ln(a+b \tan(dx+c))}{b^3(a^2+b^2)d}$
parallelrisc	$-\frac{2Bx b^4 d - 2Cxa b^3 d - C \tan(dx+c)^2 a^2 b^2 - C \tan(dx+c)^2 b^4 + B \ln(1+\tan(dx+c)^2) a b^3 + 2B \ln(a+b \tan(dx+c)) a^3 b - C \tan(dx+c)^2 a^2 b^2}{b^2 d}$
risch	$-\frac{2ia^4 Cx}{(a^2+b^2)b^3} - \frac{x C}{ib-a} - \frac{2iBac}{b^2 d} + \frac{2i(-iCbe^{2i(dx+c)} + Bbe^{2i(dx+c)} - Ca e^{2i(dx+c)} + Bb - Ca)}{db^2(e^{2i(dx+c)} + 1)^2} - \frac{2ia^4 Cc}{(a^2+b^2)b^3 d} + \dots$

input

```
int(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_
RETURNVERBOSE)
```

output

```
1/d*(1/b^2*(1/2*C*tan(d*x+c)^2*b+B*tan(d*x+c)*b-C*tan(d*x+c)*a)-1/b^3*a^3*
(B*b-C*a)/(a^2+b^2)*ln(a+b*tan(d*x+c))+1/(a^2+b^2)*(1/2*(-B*a-C*b)*ln(1+ta
n(d*x+c)^2)+(-B*b+C*a)*arctan(tan(d*x+c))))
```

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.50

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{2(Cab^3 - Bb^4)dx + (Ca^2b^2 + Cb^4) \tan(dx + c)^2 + (Ca^4 - Ba^3b) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) - (C}{2(a^2b^3 + b^5)}$$

input

```
integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

output

```
1/2*(2*(C*a*b^3 - B*b^4)*d*x + (C*a^2*b^2 + C*b^4)*tan(d*x + c)^2 + (C*a^4 - B*a^3*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (C*a^4 - B*a^3*b - B*a*b^3 - C*b^4)*log(1/(tan(d*x + c)^2 + 1)) - 2*(C*a^3*b - B*a^2*b^2 + C*a*b^3 - B*b^4)*tan(d*x + c))/((a^2*b^3 + b^5)*d)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.76 (sec) , antiderivative size = 1306, normalized size of antiderivative = 10.28

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx = \text{Too large to display}$$

input

```
integrate(tan(d*x+c)**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c)),x)
```

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2)*tan(c), Eq(a, 0) & Eq(b, 0) & Eq
(d, 0)), ((-B*log(tan(c + d*x)**2 + 1)/(2*d) + B*tan(c + d*x)**2/(2*d) + C
*x + C*tan(c + d*x)**3/(3*d) - C*tan(c + d*x)/d)/a, Eq(b, 0)), (-3*B*d*x*t
an(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*B*d*x/(2*b*d*tan(c + d*x)
- 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x)
) - 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) +
2*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*B/(2*b*d*tan(c + d
*x) - 2*I*b*d) - 3*I*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - 3
*C*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - 2*C*log(tan(c + d*x)**2 + 1)*tan(c
+ d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*C*log(tan(c + d*x)**2 + 1)/(2
*b*d*tan(c + d*x) - 2*I*b*d) + C*tan(c + d*x)**3/(2*b*d*tan(c + d*x) - 2*I
*b*d) + I*C*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*C/(2*b*d*
tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-3*B*d*x*tan(c + d*x)/(2*b*d*tan(c
+ d*x) + 2*I*b*d) - 3*I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B*log(ta
n(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*log(tan
(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*B*tan(c + d*x)**2/(2*
b*d*tan(c + d*x) + 2*I*b*d) + 3*B/(2*b*d*tan(c + d*x) + 2*I*b*d) + 3*I*C*d
*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*C*d*x/(2*b*d*tan(c + d*
x) + 2*I*b*d) - 2*C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d
*x) + 2*I*b*d) - 2*I*C*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.02

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{\frac{2(Ca - Bb)(dx + c)}{a^2 + b^2} + \frac{2(Ca^4 - Ba^3b) \log(b \tan(dx + c) + a)}{a^2 b^3 + b^5} - \frac{(Ba + Cb) \log(\tan(dx + c)^2 + 1)}{a^2 + b^2} + \frac{Cb \tan(dx + c)^2 - 2(Ca - Bb) \tan(dx + c)}{b^2}}{2d}$$

input

```
integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, a
lgorithm="maxima")
```

output

```
1/2*(2*(C*a - B*b)*(d*x + c)/(a^2 + b^2) + 2*(C*a^4 - B*a^3*b)*log(b*tan(d
*x + c) + a)/(a^2*b^3 + b^5) - (B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2 +
b^2) + (C*b*tan(d*x + c)^2 - 2*(C*a - B*b)*tan(d*x + c))/b^2)/d
```

Giac [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.15

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{(Ca - Bb)(dx + c)}{a^2d + b^2d} - \frac{(Ba + Cb) \log(\tan(dx + c)^2 + 1)}{2(a^2d + b^2d)}$$

$$+ \frac{(Ca^4 - Ba^3b) \log(|b \tan(dx + c) + a|)}{a^2b^3d + b^5d}$$

$$+ \frac{Cbd \tan(dx + c)^2 - 2Cad \tan(dx + c) + 2Bbd \tan(dx + c)}{2b^2d^2}$$

input

```
integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")
```

output

```
(C*a - B*b)*(d*x + c)/(a^2*d + b^2*d) - 1/2*(B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) + (C*a^4 - B*a^3*b)*log(abs(b*tan(d*x + c) + a))/(a^2*b^3*d + b^5*d) + 1/2*(C*b*d*tan(d*x + c)^2 - 2*C*a*d*tan(d*x + c) + 2*B*b*d*tan(d*x + c))/(b^2*d^2)
```

Mupad [B] (verification not implemented)

Time = 5.50 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.13

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{\tan(c+dx) \left(\frac{B}{b} - \frac{Ca}{b^2}\right)}{d} - \frac{\ln(\tan(c+dx) - i) (-C + B i)}{2d(-b + a i)}$$

$$+ \frac{\ln(a + b \tan(c+dx)) (Ca^4 - Ba^3b)}{d(a^2b^3 + b^5)}$$

$$- \frac{\ln(\tan(c+dx) + i) (B - C i)}{2d(a - b i)} + \frac{C \tan(c+dx)^2}{2bd}$$

input

```
int((tan(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)),x)
```


output

```
(tan(c + d*x)*(B/b - (C*a)/b^2))/d - (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*1i - b)) + (log(a + b*tan(c + d*x))*(C*a^4 - B*a^3*b))/(d*(b^5 + a^2*b^3)) - (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a - b*1i)) + (C*tan(c + d*x)^2)/(2*b*d)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.45

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{-\log(\tan(dx + c)^2 + 1) a b^4 - \log(\tan(dx + c)^2 + 1) b^4 c + 2 \log(a + \tan(dx + c) b) a^4 c - 2 \log(a + \tan(dx + c) b) a^3 b}{2 b^3 d (a^2 + b^2)}$$

input

```
int(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)
```

output

```
( - log(tan(c + d*x)**2 + 1)*a*b**4 - log(tan(c + d*x)**2 + 1)*b**4*c + 2*log(tan(c + d*x)*b + a)*a**4*c - 2*log(tan(c + d*x)*b + a)*a**3*b**2 + tan(c + d*x)**2*a**2*b**2*c + tan(c + d*x)**2*b**4*c - 2*tan(c + d*x)*a**3*b*c + 2*tan(c + d*x)*a**2*b**3 - 2*tan(c + d*x)*a*b**3*c + 2*tan(c + d*x)*b**5 + 2*a*b**3*c*d*x - 2*b**5*d*x)/(2*b**3*d*(a**2 + b**2))
```

3.26 $\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$

Optimal result	317
Mathematica [C] (verified)	317
Rubi [A] (verified)	318
Maple [A] (verified)	321
Fricas [A] (verification not implemented)	322
Sympy [C] (verification not implemented)	322
Maxima [A] (verification not implemented)	323
Giac [A] (verification not implemented)	324
Mupad [B] (verification not implemented)	324
Reduce [B] (verification not implemented)	325

Optimal result

Integrand size = 38, antiderivative size = 101

$$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(aB+bC)x}{a^2+b^2} - \frac{(bB-aC) \log(\cos(c+dx))}{(a^2+b^2)d}$$

$$+ \frac{a^2(bB-aC) \log(a+b \tan(c+dx))}{b^2(a^2+b^2)d} + \frac{C \tan(c+dx)}{bd}$$

output

$-(B*a+C*b)*x/(a^2+b^2)-(B*b-C*a)*\ln(\cos(d*x+c))/(a^2+b^2)/d+a^2*(B*b-C*a)*\ln(a+b*\tan(d*x+c))/b^2/(a^2+b^2)/d+C*\tan(d*x+c)/b/d$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.17

$$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{\frac{i(B+iC) \log(i-\tan(c+dx))}{a+ib} - \frac{(iB+C) \log(i+\tan(c+dx))}{a-ib} + \frac{2a^2(bB-aC) \log(a+b \tan(c+dx))}{b^2(a^2+b^2)} + \frac{2C \tan(c+dx)}{b}}{2d}$$

input

```
Integrate[(Tan[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]
```

output

```
((I*(B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b) - ((I*B + C)*Log[I + Tan[c + d*x]])/(a - I*b) + (2*a^2*(b*B - a*C)*Log[a + b*Tan[c + d*x]])/(b^2*(a^2 + b^2)) + (2*C*Tan[c + d*x])/b)/(2*d)
```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4115, 3042, 4089, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan(c+dx)^2)}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow 4115 \\
 & \int \frac{\tan^2(c+dx)(B + C \tan(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\tan(c+dx)^2(B + C \tan(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow 4089 \\
 & \frac{\int -\frac{((bB-aC) \tan^2(c+dx) + bC \tan(c+dx) + aC)}{a+b \tan(c+dx)} dx}{b} + \frac{C \tan(c+dx)}{bd} \\
 & \quad \downarrow 25 \\
 & \frac{C \tan(c+dx)}{bd} - \frac{\int -\frac{((bB-aC) \tan^2(c+dx) + bC \tan(c+dx) + aC)}{a+b \tan(c+dx)} dx}{b}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{C \tan(c+dx)}{bd} - \frac{\int \frac{-((bB-aC) \tan(c+dx)^2) + bC \tan(c+dx) + aC}{a+b \tan(c+dx)} dx}{b} \\
& \downarrow 4109 \\
& \frac{C \tan(c+dx)}{bd} - \frac{a^2(bB-aC) \int \frac{\tan^2(c+dx)+1}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{b(bB-aC) \int \tan(c+dx) dx}{a^2+b^2} + \frac{bx(aB+bC)}{a^2+b^2} \\
& \downarrow 3042 \\
& \frac{C \tan(c+dx)}{bd} - \frac{a^2(bB-aC) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{b(bB-aC) \int \tan(c+dx) dx}{a^2+b^2} + \frac{bx(aB+bC)}{a^2+b^2} \\
& \downarrow 3956 \\
& \frac{C \tan(c+dx)}{bd} - \frac{a^2(bB-aC) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{b(bB-aC) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(aB+bC)}{a^2+b^2} \\
& \downarrow 4100 \\
& \frac{C \tan(c+dx)}{bd} - \frac{a^2(bB-aC) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} + \frac{b(bB-aC) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(aB+bC)}{a^2+b^2} \\
& \downarrow 16 \\
& \frac{C \tan(c+dx)}{bd} - \frac{a^2(bB-aC) \log(a+b \tan(c+dx))}{bd(a^2+b^2)} + \frac{b(bB-aC) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(aB+bC)}{a^2+b^2}
\end{aligned}$$

input

```
Int[(Tan[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]
```

output

```
-(((b*(a*B + b*C)*x)/(a^2 + b^2) + (b*(b*B - a*C)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) - (a^2*(b*B - a*C)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b + (C*Tan[c + d*x])/(b*d)
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$
- rule 4089 $\text{Int}[(((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^2*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_.)]))/((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)]), x_Symbol] \rightarrow \text{Simp}[b^2*B*(\text{Tan}[e + f*x]/(d*f)), x] + \text{Simp}[1/d \text{ Int}[(a^2*A*d - b^2*B*c + (2*a*A*b + B*(a^2 - b^2))*d*\text{Tan}[e + f*x] + (A*b^2*d - b*B*(b*c - 2*a*d))*\text{Tan}[e + f*x]^2)/(c + d*\text{Tan}[e + f*x]), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$
- rule 4100 $\text{Int}[((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((A_.) + (C_.)*\tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] \text{ ; FreeQ}[\{a, b, e, f, A, C, m\}, x] \&\& \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_.)] + (C_.)*\tan[(e_.) + (f_.)*(x_.)]^2)/((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) \text{ ; FreeQ}[\{a, b, e, f, A, B, C\}, x] \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A*b - a*B - b*C, 0]$

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{\frac{\tan(dx+c)C}{b} + \frac{a^2(Bb-Ca)\ln(a+b\tan(dx+c))}{b^2(a^2+b^2)} + \frac{(Bb-Ca)\ln\left(\frac{1+\tan(dx+c)^2}{2}\right) + (-Ba-Cb)\arctan(\tan(dx+c))}{a^2+b^2}}{d}$
default	$\frac{\frac{\tan(dx+c)C}{b} + \frac{a^2(Bb-Ca)\ln(a+b\tan(dx+c))}{b^2(a^2+b^2)} + \frac{(Bb-Ca)\ln\left(\frac{1+\tan(dx+c)^2}{2}\right) + (-Ba-Cb)\arctan(\tan(dx+c))}{a^2+b^2}}{d}$
norman	$\frac{C \tan(dx+c)}{bd} - \frac{(Ba+Cb)x}{a^2+b^2} + \frac{a^2(Bb-Ca)\ln(a+b\tan(dx+c))}{b^2(a^2+b^2)d} + \frac{(Bb-Ca)\ln\left(\frac{1+\tan(dx+c)^2}{2}\right)}{2d(a^2+b^2)}$
parallelrisch	$\frac{-2Ba^2b^2dx - 2Cb^3dx + B\ln\left(\frac{1+\tan(dx+c)^2}{2}\right)b^3 + 2B\ln(a+b\tan(dx+c))a^2b - C\ln\left(\frac{1+\tan(dx+c)^2}{2}\right)a^2b^2 - 2C\ln(a+b\tan(dx+c))a^2b^2}{2d(a^2+b^2)b^2}$
risch	$\frac{x B}{ib-a} - \frac{ix C}{ib-a} - \frac{2ia^2 Bx}{b(a^2+b^2)} - \frac{2ia^2 Bc}{bd(a^2+b^2)} + \frac{2ia^3 Cx}{b^2(a^2+b^2)} + \frac{2ia^3 Cc}{b^2d(a^2+b^2)} + \frac{2iBx}{b} + \frac{2iBc}{bd} - \frac{2iCax}{b^2} - \frac{2iCac}{b^2d}$

input

```
int(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/d*(tan(d*x+c)*C/b+1/b^2*a^2*(B*b-C*a)/(a^2+b^2)*ln(a+b*tan(d*x+c))+1/(a^2+b^2)*(1/2*(B*b-C*a)*ln(1+tan(d*x+c)^2)+(-B*a-C*b)*arctan(tan(d*x+c))))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.48

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx =$$

$$\frac{2(Bab^2 + Cb^3)dx + (Ca^3 - Ba^2b) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) - (Ca^3 - Ba^2b + Cab^2 - Bb^3) \log(\tan(dx+c))}{2(a^2b^2 + b^4)d}$$

input

```
integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

output

```
-1/2*(2*(B*a*b^2 + C*b^3)*d*x + (C*a^3 - B*a^2*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (C*a^3 - B*a^2*b + C*a*b^2 - B*b^3)*log(1/(tan(d*x + c)^2 + 1)) - 2*(C*a^2*b + C*b^3)*tan(d*x + c))/((a^2*b^2 + b^4)*d)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 1020, normalized size of antiderivative = 10.10

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx = \text{Too large to display}$$

input

```
integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c)),x)
```

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2), Eq(a, 0) & Eq(b, 0) & Eq(d, 0))
, ((-B*x + B*tan(c + d*x)/d - C*log(tan(c + d*x)**2 + 1)/(2*d) + C*tan(c +
d*x)**2/(2*d))/a, Eq(b, 0)), (I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) -
2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*log(tan(c + d*x)**2 +
1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*log(tan(c + d*x)**2 +
1)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B/(2*b*d*tan(c + d*x) - 2*I*b*d) -
3*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*C*d*x/(2*b*d*tan
(c + d*x) - 2*I*b*d) + I*C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*ta
n(c + d*x) - 2*I*b*d) + C*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2
*I*b*d) + 2*C*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*C/(2*b*d*
tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-I*B*d*x*tan(c + d*x)/(2*b*d*tan(c
+ d*x) + 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*log(tan(c +
d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*log(tan(c +
d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B/(2*b*d*tan(c + d*x) + 2
*I*b*d) - 3*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*I*C*d*x/
(2*b*d*tan(c + d*x) + 2*I*b*d) - I*C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)
/(2*b*d*tan(c + d*x) + 2*I*b*d) + C*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c
+ d*x) + 2*I*b*d) + 2*C*tan(c + d*x)**2/(2*b*d*tan(c + d*x) + 2*I*b*d) + 3
*C/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*tan(c) + C*tan(c)**2)
)*tan(c)/(a + b*tan(c)), Eq(d, 0)), (2*B*a**2*b*log(a/b + tan(c + d*x))...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.08

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{\frac{2(Ba + Cb)(dx + c)}{a^2 + b^2} + \frac{2(Ca^3 - Ba^2b) \log(b \tan(dx + c) + a)}{a^2 b^2 + b^4} + \frac{(Ca - Bb) \log(\tan(dx + c)^2 + 1)}{a^2 + b^2} - \frac{2C \tan(dx + c)}{b}}{2d}$$

input

```
integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, alg
orithm="maxima")
```

output

```
-1/2*(2*(B*a + C*b)*(d*x + c)/(a^2 + b^2) + 2*(C*a^3 - B*a^2*b)*log(b*tan(
d*x + c) + a)/(a^2*b^2 + b^4) + (C*a - B*b)*log(tan(d*x + c)^2 + 1)/(a^2 +
b^2) - 2*C*tan(d*x + c)/b)/d
```


Giac [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.18

$$\int \frac{\tan(c+dx)(B\tan(c+dx)+C\tan^2(c+dx))}{a+b\tan(c+dx)} dx$$

$$= -\frac{(Ba+Cb)(dx+c)}{a^2d+b^2d} - \frac{(Ca-Bb)\log(\tan(dx+c)^2+1)}{2(a^2d+b^2d)}$$

$$- \frac{(Ca^3-Ba^2b)\log(|b\tan(dx+c)+a|)}{a^2b^2d+b^4d} + \frac{C\tan(dx+c)}{bd}$$

input `integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-(B*a + C*b)*(d*x + c)/(a^2*d + b^2*d) - 1/2*(C*a - B*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) - (C*a^3 - B*a^2*b)*log(abs(b*tan(d*x + c) + a))/(a^2*b^2*d + b^4*d) + C*tan(d*x + c)/(b*d)`

Mupad [B] (verification not implemented)

Time = 5.35 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.16

$$\int \frac{\tan(c+dx)(B\tan(c+dx)+C\tan^2(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{C\tan(c+dx)}{bd} + \frac{\ln(\tan(c+dx)+1i)(B-C1i)}{2d(b+a1i)}$$

$$- \frac{\ln(a+b\tan(c+dx))(Ca^3-Ba^2b)}{d(a^2b^2+b^4)} + \frac{\ln(\tan(c+dx)-i)(-C+B1i)}{2d(a+b1i)}$$

input `int((tan(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a*1i + b)) - (log(a + b*tan(c + d*x))*(C*a^3 - B*a^2*b))/(d*(b^4 + a^2*b^2)) + (C*tan(c + d*x))/(b*d) + (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a + b*1i))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.27

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{-\log(\tan(dx + c)^2 + 1) a b^2 c + \log(\tan(dx + c)^2 + 1) b^4 - 2 \log(a + \tan(dx + c) b) a^3 c + 2 \log(a + \tan(dx + c) b) a^2 b c}{2 b^2 d (a^2 + b^2)}$$

input

```
int(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)
```

output

```
( - log(tan(c + d*x)**2 + 1)*a*b**2*c + log(tan(c + d*x)**2 + 1)*b**4 - 2*
log(tan(c + d*x)*b + a)*a**3*c + 2*log(tan(c + d*x)*b + a)*a**2*b**2 + 2*t
an(c + d*x)*a**2*b*c + 2*tan(c + d*x)*b**3*c - 2*a*b**3*d*x - 2*b**3*c*d*x
)/(2*b**2*d*(a**2 + b**2))
```

3.27 $\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{a + b \tan(c+dx)} dx$

Optimal result	326
Mathematica [C] (verified)	326
Rubi [A] (verified)	327
Maple [A] (verified)	328
Fricas [A] (verification not implemented)	329
Sympy [C] (verification not implemented)	330
Maxima [A] (verification not implemented)	330
Giac [A] (verification not implemented)	331
Mupad [B] (verification not implemented)	331
Reduce [B] (verification not implemented)	332

Optimal result

Integrand size = 32, antiderivative size = 85

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx = \frac{(bB - aC)x}{a^2 + b^2} - \frac{(aB + bC) \log(\cos(c + dx))}{(a^2 + b^2) d} - \frac{a(bB - aC) \log(a + b \tan(c + dx))}{b(a^2 + b^2) d}$$

output

```
(B*b-C*a)*x/(a^2+b^2)-(B*a+C*b)*ln(cos(d*x+c))/(a^2+b^2)/d-a*(B*b-C*a)*ln(a+b*tan(d*x+c))/b/(a^2+b^2)/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.13 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.15

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx = \frac{(a - ib)b(B + iC) \log(i - \tan(c + dx)) + (a + ib)b(B - iC) \log(i + \tan(c + dx)) + 2a(-bB + aC) \log(\cos(c + dx))}{2b(a^2 + b^2) d}$$

input

```
Integrate[(B*Tan[c + d*x] + C*Tan[c + d*x]^2)/(a + b*Tan[c + d*x]),x]
```

output

$$\left((a - I*b)*b*(B + I*C)*\text{Log}[I - \text{Tan}[c + d*x]] + (a + I*b)*b*(B - I*C)*\text{Log}[I + \text{Tan}[c + d*x]] + 2*a*(-(b*B) + a*C)*\text{Log}[a + b*\text{Tan}[c + d*x]] \right) / (2*b*(a^2 + b^2)*d)$$
Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.12, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3042, 4853, 2160, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{B \tan(c + dx) + C \tan(c + dx)^2}{a + b \tan(c + dx)} dx \\ & \quad \downarrow \text{4853} \\ & \int \frac{\tan(c+dx)(B+C \tan(c+dx))}{(a+b \tan(c+dx))(\tan^2(c+dx)+1)} d \tan(c + dx) \\ & \quad \downarrow \text{2160} \\ & \int \left(\frac{a(aC-bB)}{(a^2+b^2)(a+b \tan(c+dx))} + \frac{bB-aC+(aB+bC) \tan(c+dx)}{(a^2+b^2)(\tan^2(c+dx)+1)} \right) d \tan(c + dx) \\ & \quad \downarrow \text{2009} \\ & \frac{\frac{(bB-aC) \arctan(\tan(c+dx))}{a^2+b^2} + \frac{(aB+bC) \log(\tan^2(c+dx)+1)}{2(a^2+b^2)} - \frac{a(bB-aC) \log(a+b \tan(c+dx))}{b(a^2+b^2)}}{d} \end{aligned}$$

input

$$\text{Int}[(B*\text{Tan}[c + d*x] + C*\text{Tan}[c + d*x]^2)/(a + b*\text{Tan}[c + d*x]), x]$$

output

$$\frac{((b*B - a*C)*\text{ArcTan}[\text{Tan}[c + d*x]])/(a^2 + b^2) - (a*(b*B - a*C)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(b*(a^2 + b^2)) + ((a*B + b*C)*\text{Log}[1 + \text{Tan}[c + d*x]^2])/(2*(a^2 + b^2))}{d}$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 2160

$$\text{Int}[(Pq_)*((d_) + (e_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e*x)^m*Pq*(a + b*x^2)^p, x], x] \text{ /; FreeQ}[\{a, b, d, e, m\}, x] \ \&\& \ \text{PolyQ}[Pq, x] \ \&\& \ \text{IGtQ}[p, -2]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4853

$$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{FunctionOfTrig}[u, x]\}, \text{Simp}[\text{With}[\{d = \text{FreeFactors}[\text{Tan}[v], x]\}, d/\text{Coefficient}[v, x, 1] \text{ Subst}[\text{Int}[\text{SubstFor}[1/(1 + d^2*x^2), \text{Tan}[v]/d, u, x], x, \text{Tan}[v]/d]], x] \text{ /; !FalseQ}[v] \ \&\& \ \text{FunctionOfQ}[\text{NonfreeFactors}[\text{Tan}[v], x], u, x, \text{True}] \ \&\& \ \text{TryPureTanSubst}[\text{ActivateTrig}[u], x]]]$$
Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{-\frac{a(Bb-Ca)\ln(a+b\tan(dx+c))}{(a^2+b^2)b} + \frac{(Ba+Cb)\ln(1+\tan(dx+c)^2)}{2} + (Bb-Ca)\arctan(\tan(dx+c))}{d}$
default	$\frac{-\frac{a(Bb-Ca)\ln(a+b\tan(dx+c))}{(a^2+b^2)b} + \frac{(Ba+Cb)\ln(1+\tan(dx+c)^2)}{2} + (Bb-Ca)\arctan(\tan(dx+c))}{d}$
norman	$\frac{(Bb-Ca)x}{a^2+b^2} + \frac{(Ba+Cb)\ln(1+\tan(dx+c)^2)}{2d(a^2+b^2)} - \frac{a(Bb-Ca)\ln(a+b\tan(dx+c))}{b(a^2+b^2)d}$
parallelrisch	$\frac{2Bb^2dx - 2Cabdx + B\ln(1+\tan(dx+c)^2)ab - 2B\ln(a+b\tan(dx+c))ab + C\ln(1+\tan(dx+c)^2)b^2 + 2C\ln(a+b\tan(dx+c))}{2(a^2+b^2)db}$
risch	$\frac{ixB}{ib-a} + \frac{xC}{ib-a} + \frac{2iCx}{b} + \frac{2iCc}{db} + \frac{2iaBx}{a^2+b^2} + \frac{2iaBc}{(a^2+b^2)d} - \frac{2ia^2Cx}{(a^2+b^2)b} - \frac{2ia^2Cc}{(a^2+b^2)db} - \frac{\ln(e^{2i(dx+c)}+1)C}{bd} - \frac{a}{b}$

input `int((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(-a*(B*b-C*a)/(a^2+b^2)/b*ln(a+b*tan(d*x+c))+1/(a^2+b^2)*(1/2*(B*a+C*b)*ln(1+tan(d*x+c)^2)+(B*b-C*a)*arctan(tan(d*x+c))))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.29

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx =$$

$$-\frac{2(Cab - Bb^2)dx - (Ca^2 - Bab) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) + (Ca^2 + Cb^2) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right)}{2(a^2b + b^3)d}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*(2*(C*a*b - B*b^2)*d*x - (C*a^2 - B*a*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + (C*a^2 + C*b^2)*log(1/(tan(d*x + c)^2 + 1)))/((a^2*b + b^3)*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 711, normalized size of antiderivative = 8.36

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx = \text{Too large to display}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2)/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((B*log(tan(c + d*x)**2 + 1)/(2*d) - C*x + C*tan(c + d*x)/d)/a, Eq(b, 0)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - B/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + C*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*C*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*C/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - B/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + C*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*C*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*C/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*tan(c) + C*tan(c)**2)/(a + b*tan(c)), Eq(d, 0)), (-2*B*a*b*log(a/b + tan(c + d*x))/(2*a**2*b*d + 2*b**3*d) + B*a*b*log(tan(c + d*x)**2 + 1)/(2*a**2*b*d + 2*b**3*d) + 2*B*b**2*d*x/(2*a**2*b*d + 2*b**3*d) + 2*C*a**2*log(a/b + tan(c + d*x))/(2*a**2*b*d + 2*b**3*d) - 2*C*a*b*d*x/(2*a**2*b*d + 2*b**3*d) + C*b**2*log(tan(c + d*x)**2 + 1)/(2*a**2*b*d + 2*b**3*d), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.11

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx$$

$$= -\frac{\frac{2(Ca-Bb)(dx+c)}{a^2+b^2} - \frac{2(Ca^2-Bab) \log(b \tan(dx+c)+a)}{a^2b+b^3} - \frac{(Ba+Cb) \log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(C*a - B*b)*(d*x + c)/(a^2 + b^2) - 2*(C*a^2 - B*a*b)*log(b*tan(d*x + c) + a)/(a^2*b + b^3) - (B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.18

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx = -\frac{(Ca - Bb)(dx + c)}{a^2d + b^2d} + \frac{(Ba + Cb) \log(\tan(dx + c)^2 + 1)}{2(a^2d + b^2d)} + \frac{(Ca^2 - Bab) \log(|b \tan(dx + c) + a|)}{a^2bd + b^3d}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-(C*a - B*b)*(d*x + c)/(a^2*d + b^2*d) + 1/2*(B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) + (C*a^2 - B*a*b)*log(abs(b*tan(d*x + c) + a))/(a^2*b*d + b^3*d)`

Mupad [B] (verification not implemented)

Time = 5.69 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.18

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx = \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2d (-b + a i)} + \frac{\ln(\tan(c + dx) + i) (B - C i)}{2d (a - b i)} - \frac{a \ln(a + b \tan(c + dx)) (Bb - Ca)}{bd (a^2 + b^2)}$$

input `int((B*tan(c + d*x) + C*tan(c + d*x)^2)/(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*1i - b)) + (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a - b*1i)) - (a*log(a + b*tan(c + d*x))*(B*b - C*a))/(b*d*(a^2 + b^2))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.15

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{a + b \tan(c + dx)} dx$$

$$= \frac{\log(\tan(dx + c)^2 + 1) a b^2 + \log(\tan(dx + c)^2 + 1) b^2 c + 2 \log(a + \tan(dx + c) b) a^2 c - 2 \log(a + \tan(dx + c) b) a b c}{2 b d (a^2 + b^2)}$$

input `int((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)`

output `(log(tan(c + d*x)**2 + 1)*a*b**2 + log(tan(c + d*x)**2 + 1)*b**2*c + 2*log(tan(c + d*x)*b + a)*a**2*c - 2*log(tan(c + d*x)*b + a)*a*b**2 - 2*a*b*c*d*x + 2*b**3*d*x)/(2*b*d*(a**2 + b**2))`

3.28
$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

Optimal result	333
Mathematica [A] (verified)	333
Rubi [A] (verified)	334
Maple [A] (verified)	336
Fricas [A] (verification not implemented)	336
Sympy [C] (verification not implemented)	337
Maxima [A] (verification not implemented)	338
Giac [A] (verification not implemented)	339
Mupad [B] (verification not implemented)	339
Reduce [B] (verification not implemented)	340

Optimal result

Integrand size = 38, antiderivative size = 58

$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{(aB+bC)x}{a^2+b^2} + \frac{(bB-aC) \log(a \cos(c+dx)+b \sin(c+dx))}{(a^2+b^2)d}$$

output $(B*a+C*b)*x/(a^2+b^2)+(B*b-C*a)*\ln(a*\cos(d*x+c)+b*\sin(d*x+c))/(a^2+b^2)/d$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.16

$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{-2(aB+bC) \arctan(\cot(c+dx))+(bB-aC)(2 \log(b+a \cot(c+dx))-\log(\csc^2(c+dx)))}{2(a^2+b^2)d}$$

input `Integrate[(Cot[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]`

output

$$\frac{(-2*(a*B + b*C)*\text{ArcTan}[\text{Cot}[c + d*x]] + (b*B - a*C)*(2*\text{Log}[b + a*\text{Cot}[c + d*x]] - \text{Log}[\text{Csc}[c + d*x]^2]))}{(2*(a^2 + b^2)*d)}$$
Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3042, 4115, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{\tan(c + dx)(a + b \tan(c + dx))} dx \\ & \quad \downarrow \text{4115} \\ & \int \frac{B + C \tan(c + dx)}{a + b \tan(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{B + C \tan(c + dx)}{a + b \tan(c + dx)} dx \\ & \quad \downarrow \text{4014} \\ & \frac{(bB - aC) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(aB + bC)}{a^2 + b^2} \\ & \quad \downarrow \text{3042} \\ & \frac{(bB - aC) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(aB + bC)}{a^2 + b^2} \\ & \quad \downarrow \text{4013} \\ & \frac{(bB - aC) \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)} + \frac{x(aB + bC)}{a^2 + b^2} \end{aligned}$$

input $\text{Int}[(\text{Cot}[c + d*x]*(B*\text{Tan}[c + d*x] + C*\text{Tan}[c + d*x]^2))/(a + b*\text{Tan}[c + d*x]), x]$

output $((a*B + b*C)*x)/(a^2 + b^2) + ((b*B - a*C)*\text{Log}[a*\text{Cos}[c + d*x] + b*\text{Sin}[c + d*x]])/((a^2 + b^2)*d)$

Defintions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinear } Q[u, x]$

rule 4013 $\text{Int}[(c_ + (d_)*\text{tan}[e_ + (f_)*(x_)])/((a_ + (b_)*\text{tan}[e_ + (f_)*(x_)])], x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

rule 4014 $\text{Int}[(c_ + (d_)*\text{tan}[e_ + (f_)*(x_)])/((a_ + (b_)*\text{tan}[e_ + (f_)*(x_)])*(x_)], x_Symbol] \rightarrow \text{Simp}[(a*c + b*d)*(x/(a^2 + b^2)), x] + \text{Simp}[(b*c - a*d)/(a^2 + b^2) \ \text{Int}[(b - a*\text{Tan}[e + f*x])/(a + b*\text{Tan}[e + f*x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[a*c + b*d, 0]$

rule 4115 $\text{Int}[(a_ + (b_)*\text{tan}[e_ + (f_)*(x_)])^{(m_)}*(c_ + (d_)*\text{tan}[e_ + (f_)*(x_)])^{(n_)}*((A_ + (B_)*\text{tan}[e_ + (f_)*(x_)]) + (C_)*\text{tan}[e_ + (f_)*(x_)])^2), x_Symbol] \rightarrow \text{Simp}[1/b^2 \ \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m+1)}*(c + d*\text{Tan}[e + f*x])^n*(b*B - a*C + b*C*\text{Tan}[e + f*x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, C, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[A*b^2 - a*b*B + a^2*C, 0]$

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.07

method	result
parallelrisc	$\frac{2Badx+2Cbdx+\left(-\ln\left(\sec(dx+c)^2\right)+2\ln(a+b\tan(dx+c))\right)(Bb-Ca)}{2d(a^2+b^2)}$
derivativedivides	$\frac{\frac{(Bb-Ca)\ln(a+b\tan(dx+c))}{a^2+b^2}+\frac{(-Bb+Ca)\ln\left(\frac{1+\tan(dx+c)^2}{2}\right)+(Ba+Cb)\arctan(\tan(dx+c))}{a^2+b^2}}{d}$
default	$\frac{\frac{(Bb-Ca)\ln(a+b\tan(dx+c))}{a^2+b^2}+\frac{(-Bb+Ca)\ln\left(\frac{1+\tan(dx+c)^2}{2}\right)+(Ba+Cb)\arctan(\tan(dx+c))}{a^2+b^2}}{d}$
norman	$\frac{(Ba+Cb)x}{a^2+b^2}+\frac{(Bb-Ca)\ln(a+b\tan(dx+c))}{d(a^2+b^2)}-\frac{(Bb-Ca)\ln\left(1+\tan(dx+c)^2\right)}{2d(a^2+b^2)}$
risc	$-\frac{xB}{ib-a}+\frac{ixC}{ib-a}-\frac{2iBbx}{a^2+b^2}+\frac{2iCax}{a^2+b^2}-\frac{2iBbc}{d(a^2+b^2)}+\frac{2iCac}{d(a^2+b^2)}+\frac{\ln\left(\frac{e^{2i(dx+c)}-\frac{ib+a}{ib-a}}{d(a^2+b^2)}\right)Bb}{d(a^2+b^2)}-\frac{\ln\left(\frac{e^{2i(dx+c)}-\frac{ib+a}{ib-a}}{d(a^2+b^2)}\right)}{d(a^2+b^2)}$

input `int(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2/d/(a^2+b^2)*(2*B*a*d*x+2*C*b*d*x+(-ln(sec(d*x+c)^2)+2*ln(a+b*tan(d*x+c)))*(B*b-C*a))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.31

$$\int \frac{\cot(c+dx)(B\tan(c+dx)+C\tan^2(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{2(Ba+Cb)dx - (Ca - Bb)\log\left(\frac{b^2\tan(dx+c)^2+2ab\tan(dx+c)+a^2}{\tan(dx+c)^2+1}\right)}{2(a^2+b^2)d}$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,algorithm="fricas")`

output

```
1/2*(2*(B*a + C*b)*d*x - (C*a - B*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)))/(a^2 + b^2)*d
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.26 (sec) , antiderivative size = 541, normalized size of antiderivative = 9.33

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \left\{ \begin{array}{l} \frac{\infty x (B \tan(c) + C \tan^2(c)) \cot(c)}{\tan(c)} \\ Bx + \frac{C \log(\tan^2(c + dx) + 1)}{2d} \\ \frac{iBdx \tan(c + dx)}{2bd \tan(c + dx) - 2ibd} + \frac{Bdx}{2bd \tan(c + dx) - 2ibd} + \frac{iB}{2bd \tan(c + dx) - 2ibd} + \frac{Cdx \tan(c + dx)}{2bd \tan(c + dx) - 2ibd} - \frac{iCdx}{2bd \tan(c + dx) - 2ibd} - \frac{C}{2bd \tan(c + dx) - 2ibd} \\ - \frac{iBdx \tan(c + dx)}{2bd \tan(c + dx) + 2ibd} + \frac{Bdx}{2bd \tan(c + dx) + 2ibd} - \frac{iB}{2bd \tan(c + dx) + 2ibd} + \frac{Cdx \tan(c + dx)}{2bd \tan(c + dx) + 2ibd} + \frac{iCdx}{2bd \tan(c + dx) + 2ibd} - \frac{C}{2bd \tan(c + dx) + 2ibd} \\ \frac{x (B \tan(c) + C \tan^2(c)) \cot(c)}{a + b \tan(c)} \\ \frac{2Badx}{2a^2d + 2b^2d} + \frac{2Bb \log(\frac{a}{b} + \tan(c + dx))}{2a^2d + 2b^2d} - \frac{Bb \log(\tan^2(c + dx) + 1)}{2a^2d + 2b^2d} - \frac{2Ca \log(\frac{a}{b} + \tan(c + dx))}{2a^2d + 2b^2d} + \frac{Ca \log(\tan^2(c + dx) + 1)}{2a^2d + 2b^2d} + \frac{2Cb}{2a^2d + 2b^2d} \end{array} \right.$$

input

```
integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c)),x)
```

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2)*cot(c)/tan(c), Eq(a, 0) & Eq(b,
0) & Eq(d, 0)), ((B*x + C*log(tan(c + d*x)**2 + 1)/(2*d))/a, Eq(b, 0)), (I
*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*d*x/(2*b*d*tan(c +
d*x) - 2*I*b*d) + I*B/(2*b*d*tan(c + d*x) - 2*I*b*d) + C*d*x*tan(c + d*x)/
(2*b*d*tan(c + d*x) - 2*I*b*d) - I*C*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) -
C/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-I*B*d*x*tan(c + d*x)/(2*
b*d*tan(c + d*x) + 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B/(
2*b*d*tan(c + d*x) + 2*I*b*d) + C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2
*I*b*d) + I*C*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - C/(2*b*d*tan(c + d*x) +
2*I*b*d), Eq(a, I*b)), (x*(B*tan(c) + C*tan(c)**2)*cot(c)/(a + b*tan(c)),
Eq(d, 0)), (2*B*a*d*x/(2*a**2*d + 2*b**2*d) + 2*B*b*log(a/b + tan(c + d*x
)))/(2*a**2*d + 2*b**2*d) - B*b*log(tan(c + d*x)**2 + 1)/(2*a**2*d + 2*b**2
*d) - 2*C*a*log(a/b + tan(c + d*x))/(2*a**2*d + 2*b**2*d) + C*a*log(tan(c
+ d*x)**2 + 1)/(2*a**2*d + 2*b**2*d) + 2*C*b*d*x/(2*a**2*d + 2*b**2*d), Tr
ue))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.52

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{\frac{2(Ba + Cb)(dx + c)}{a^2 + b^2} - \frac{2(Ca - Bb) \log(b \tan(dx + c) + a)}{a^2 + b^2} + \frac{(Ca - Bb) \log(\tan(dx + c)^2 + 1)}{a^2 + b^2}}{2d}$$

input

```
integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, alg
orithm="maxima")
```

output

```
1/2*(2*(B*a + C*b)*(d*x + c)/(a^2 + b^2) - 2*(C*a - B*b)*log(b*tan(d*x + c
) + a)/(a^2 + b^2) + (C*a - B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d
```

Giac [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{(Ba + Cb)(dx + c)}{a^2d + b^2d} + \frac{(Ca - Bb) \log(\tan(dx + c)^2 + 1)}{2(a^2d + b^2d)}$$

$$- \frac{(Cab - Bb^2) \log(|b \tan(dx + c) + a|)}{a^2bd + b^3d}$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `(B*a + C*b)*(d*x + c)/(a^2*d + b^2*d) + 1/2*(C*a - B*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) - (C*a*b - B*b^2)*log(abs(b*tan(d*x + c) + a))/(a^2*b*d + b^3*d)`

Mupad [B] (verification not implemented)

Time = 5.62 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.60

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{\ln(a + b \tan(c + dx)) (Bb - Ca)}{d (a^2 + b^2)} - \frac{\ln(\tan(c + dx) + 1i) (B - C 1i)}{2 d (b + a 1i)}$$

$$- \frac{\ln(\tan(c + dx) - 1i) (-C + B 1i)}{2 d (a + b 1i)}$$

input `int((cot(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)),x)`

output `(log(a + b*tan(c + d*x))*(B*b - C*a))/(d*(a^2 + b^2)) - (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a*1i + b)) - (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a + b*1i))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 128, normalized size of antiderivative = 2.21

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{\log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) ac - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 + 1\right) b^2 - \log\left(\tan\left(\frac{dx}{2} + \frac{c}{2}\right)^2 a - 2 \tan\left(\frac{dx}{2} + \frac{c}{2}\right) b - a\right) a}{d(a^2 + b^2)}$$

input `int(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)`

output `(log(tan((c + d*x)/2)**2 + 1)*a*c - log(tan((c + d*x)/2)**2 + 1)*b**2 - log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*a*c + log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*b**2 + a*b*d*x + b*c*d*x)/(d*(a**2 + b**2))`

3.29 $\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$

Optimal result	341
Mathematica [C] (verified)	341
Rubi [A] (verified)	342
Maple [A] (verified)	344
Fricas [A] (verification not implemented)	345
Sympy [C] (verification not implemented)	345
Maxima [A] (verification not implemented)	346
Giac [A] (verification not implemented)	347
Mupad [B] (verification not implemented)	347
Reduce [F]	348

Optimal result

Integrand size = 40, antiderivative size = 80

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(bB-aC)x}{a^2+b^2} + \frac{B \log(\sin(c+dx))}{ad} - \frac{b(bB-aC) \log(a \cos(c+dx)+b \sin(c+dx))}{a(a^2+b^2)d}$$

output

```
-(B*b-C*a)*x/(a^2+b^2)+B*ln(sin(d*x+c))/a/d-b*(B*b-C*a)*ln(a*cos(d*x+c)+b*
sin(d*x+c))/a/(a^2+b^2)/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.23 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.41

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx =$$

$$\frac{\frac{(B+iC) \log(i-\tan(c+dx))}{a+ib} - \frac{2B \log(\tan(c+dx))}{a} + \frac{(B-iC) \log(i+\tan(c+dx))}{a-ib} + \frac{2b(bB-aC) \log(a+b \tan(c+dx))}{a(a^2+b^2)}}{2d}$$

input

```
Integrate[(Cot[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[
c + d*x]),x]
```

output

```
-1/2*(((B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b) - (2*B*Log[Tan[c + d*x]]
)/a + ((B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b) + (2*b*(b*B - a*C)*Log[a
+ b*Tan[c + d*x]])/(a*(a^2 + b^2)))/d
```

Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4115, 3042, 4094, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^2(c+dx) (B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{\tan(c+dx) (a + b \tan(c+dx))} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \frac{\cot(c+dx) (B + C \tan(c+dx))}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B + C \tan(c+dx)}{\tan(c+dx) (a + b \tan(c+dx))} dx \\
 & \quad \downarrow \text{4094} \\
 & -\frac{b(bB - aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} + \frac{B \int \cot(c+dx) dx}{a} - \frac{x(bB - aC)}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b(bB - aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} + \frac{B \int -\tan(c+dx + \frac{\pi}{2}) dx}{a} - \frac{x(bB - aC)}{a^2 + b^2}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 25 \\
 & -\frac{b(bB - aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} - \frac{B \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx}{a} - \frac{x(bB - aC)}{a^2 + b^2} \\
 & \downarrow 3956 \\
 & -\frac{b(bB - aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} - \frac{x(bB - aC)}{a^2 + b^2} + \frac{B \log(-\sin(c + dx))}{ad} \\
 & \downarrow 4013 \\
 & -\frac{b(bB - aC) \log(a \cos(c + dx) + b \sin(c + dx))}{ad(a^2 + b^2)} - \frac{x(bB - aC)}{a^2 + b^2} + \frac{B \log(-\sin(c + dx))}{ad}
 \end{aligned}$$

input `Int[(Cot[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]`

output `-(((b*B - a*C)*x)/(a^2 + b^2)) + (B*Log[-Sin[c + d*x]])/(a*d) - (b*(b*B - a*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4094

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(B*(b*
c + a*d) + A*(a*c - b*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[b*((A*b
- a*B)/((b*c - a*d)*(a^2 + b^2))) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e
+ f*x]), x], x] + Simp[d*((B*c - A*d)/((b*c - a*d)*(c^2 + d^2))) Int[(d -
c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f,
A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e
_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.19

method	result
parallelrisch	$\frac{(-2Bb^2+2Cab) \ln(a+b \tan(dx+c)) + (-Ba^2-Cab) \ln(\sec(dx+c)^2) + 2B(a^2+b^2) \ln(\tan(dx+c)) - 2adx(Bb-Ca)}{2(a^2+b^2)ad}$
derivativedivides	$\frac{-\frac{(Bb-Ca)b \ln(a+b \tan(dx+c))}{(a^2+b^2)a} + \frac{(-Ba-Cb) \ln(1+\tan(dx+c)^2)}{2} + (-Bb+Ca) \arctan(\tan(dx+c)) + \frac{B \ln(\tan(dx+c))}{a}}{d}$
default	$\frac{-\frac{(Bb-Ca)b \ln(a+b \tan(dx+c))}{(a^2+b^2)a} + \frac{(-Ba-Cb) \ln(1+\tan(dx+c)^2)}{2} + (-Bb+Ca) \arctan(\tan(dx+c)) + \frac{B \ln(\tan(dx+c))}{a}}{d}$
norman	$-\frac{(Bb-Ca)x}{a^2+b^2} + \frac{B \ln(\tan(dx+c))}{ad} - \frac{(Ba+Cb) \ln(1+\tan(dx+c)^2)}{2d(a^2+b^2)} - \frac{(Bb-Ca)b \ln(a+b \tan(dx+c))}{(a^2+b^2)ad}$
risch	$-\frac{ixB}{ib-a} - \frac{xC}{ib-a} + \frac{2ib^2Bx}{(a^2+b^2)a} + \frac{2ib^2Bc}{(a^2+b^2)ad} - \frac{2ibCx}{a^2+b^2} - \frac{2ibCc}{(a^2+b^2)d} - \frac{2iBx}{a} - \frac{2iBc}{ad} - \frac{b^2 \ln(e^{2i(dx+c)} - \frac{ib+1}{ib-1})}{(a^2+b^2)ad}$

input

```
int(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_
RETURNVERBOSE)
```

output

$$\frac{1}{2} * ((-2 * B * b^2 + 2 * C * a * b) * \ln(a + b * \tan(dx + c)) + (-B * a^2 - C * a * b) * \ln(\sec(dx + c)^2) + 2 * B * (a^2 + b^2) * \ln(\tan(dx + c)) - 2 * a * dx * (B * b - C * a)) / (a^2 + b^2) / a / d$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.48

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{2(Ca^2 - Bab)dx + (Ba^2 + Bb^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + (Cab - Bb^2) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2+1}\right)}{2(a^3 + ab^2)d}$$

input

```
integrate(cot(dx+c)^2*(B*tan(dx+c)+C*tan(dx+c)^2)/(a+b*tan(dx+c)),x, algorithm="fricas")
```

output

$$\frac{1}{2} * (2 * (C * a^2 - B * a * b) * dx + (B * a^2 + B * b^2) * \log(\tan(dx + c)^2 / (\tan(dx + c)^2 + 1)) + (C * a * b - B * b^2) * \log((b^2 * \tan(dx + c)^2 + 2 * a * b * \tan(dx + c) + a^2) / (\tan(dx + c)^2 + 1))) / ((a^3 + a * b^2) * d)$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.32 (sec) , antiderivative size = 966, normalized size of antiderivative = 12.08

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx = \text{Too large to display}$$

input

```
integrate(cot(dx+c)**2*(B*tan(dx+c)+C*tan(dx+c)**2)/(a+b*tan(dx+c)),x)
```

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2)*cot(c)**2/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((-B*log(tan(c + d*x)**2 + 1)/(2*d) + B*log(tan(c + d*x)))/d + C*x)/a, Eq(b, 0)), ((-B*x - B/(d*tan(c + d*x)) - C*log(tan(c + d*x)**2 + 1)/(2*d) + C*log(tan(c + d*x))/d)/b, Eq(a, 0)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*B*log(tan(c + d*x))*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*B*log(tan(c + d*x))/(2*b*d*tan(c + d*x) - 2*I*b*d) + B/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + C*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*C/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*I*B*log(tan(c + d*x))*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*B*log(tan(c + d*x))/(2*b*d*tan(c + d*x) + 2*I*b*d) + B/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*C*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + C*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*C/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*tan(c) + C*tan(c)**2)*cot(c)**2/(a + b*tan(c)), Eq(d, 0)), (-B*a**2*log(tan(c + d*x)**2 + 1)/(2*a**3*d + 2*a*b**2*d) + 2*B*a**2*log(...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.34

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{\frac{2(Ca - Bb)(dx + c)}{a^2 + b^2} + \frac{2(Cab - Bb^2) \log(b \tan(dx + c) + a)}{a^3 + ab^2} - \frac{(Ba + Cb) \log(\tan(dx + c)^2 + 1)}{a^2 + b^2} + \frac{2B \log(\tan(dx + c))}{a}}{2d}$$

input

```
integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="maxima")
```

output

```
1/2*(2*(C*a - B*b)*(d*x + c)/(a^2 + b^2) + 2*(C*a*b - B*b^2)*log(b*tan(d*x + c) + a)/(a^3 + a*b^2) - (B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*B*log(tan(d*x + c))/a)/d
```

Giac [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.48

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{(Ca - Bb)(dx + c)}{a^2d + b^2d} - \frac{(Ba + Cb) \log(\tan(dx + c)^2 + 1)}{2(a^2d + b^2d)}$$

$$+ \frac{(Cab^2 - Bb^3) \log(|b \tan(dx + c) + a|)}{a^3bd + ab^3d} + \frac{B \log(|\tan(dx + c)|)}{ad}$$

input

```
integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")
```

output

```
(C*a - B*b)*(d*x + c)/(a^2*d + b^2*d) - 1/2*(B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) + (C*a*b^2 - B*b^3)*log(abs(b*tan(d*x + c) + a))/(a^3*b*d + a*b^3*d) + B*log(abs(tan(d*x + c)))/(a*d)
```

Mupad [B] (verification not implemented)

Time = 5.97 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.44

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{B \ln(\tan(c + dx))}{ad} - \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2d (-b + a i)}$$

$$- \frac{\ln(\tan(c + dx) + i) (B - C i)}{2d (a - b i)} - \frac{b \ln(a + b \tan(c + dx)) (B b - C a)}{ad (a^2 + b^2)}$$

input

```
int((cot(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)),x)
```

output

```
(B*log(tan(c + d*x)))/(a*d) - (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*1i - b)) - (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a - b*1i)) - (b*log(a + b*tan(c + d*x))*(B*b - C*a))/(a*d*(a^2 + b^2))
```


Reduce [F]

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$
$$= \int \frac{\cot(dx + c)^2 (B \tan(dx + c) + C \tan(dx + c)^2)}{a + \tan(dx + c) b} dx$$

input `int(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)`

output `int(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)`

3.30
$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

Optimal result	349
Mathematica [C] (verified)	349
Rubi [A] (verified)	350
Maple [A] (verified)	353
Fricas [A] (verification not implemented)	354
Sympy [C] (verification not implemented)	354
Maxima [A] (verification not implemented)	355
Giac [A] (verification not implemented)	356
Mupad [B] (verification not implemented)	356
Reduce [F]	357

Optimal result

Integrand size = 40, antiderivative size = 103

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(aB+bC)x}{a^2+b^2} - \frac{B \cot(c+dx)}{ad} - \frac{(bB-aC) \log(\sin(c+dx))}{a^2d}$$

$$+ \frac{b^2(bB-aC) \log(a \cos(c+dx)+b \sin(c+dx))}{a^2(a^2+b^2)d}$$

output

```
-(B*a+C*b)*x/(a^2+b^2)-B*cot(d*x+c)/a/d-(B*b-C*a)*ln(sin(d*x+c))/a^2/d+b^2
*(B*b-C*a)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^2/(a^2+b^2)/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.34

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{-\frac{2B \cot(c+dx)}{a} + \frac{i(B+iC) \log(i-\tan(c+dx))}{a+ib} + \frac{2(-bB+aC) \log(\tan(c+dx))}{a^2} - \frac{(iB+C) \log(i+\tan(c+dx))}{a-ib} + \frac{2b^2(bB-aC) \log(a+b \tan(c+dx))}{a^2(a^2+b^2)}}{2d}$$

input

```
Integrate[(Cot[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[
c + d*x]),x]
```

output

```
((-2*B*Cot[c + d*x])/a + (I*(B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b) + (
2*(-(b*B) + a*C)*Log[Tan[c + d*x]])/a^2 - ((I*B + C)*Log[I + Tan[c + d*x]]
)/(a - I*b) + (2*b^2*(b*B - a*C)*Log[a + b*Tan[c + d*x]])/(a^2*(a^2 + b^2)
))/(2*d)
```

Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.08, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4115, 3042, 4092, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \tan(c + dx) + C \tan^2(c + dx)^2}{\tan(c + dx)^3 (a + b \tan(c + dx))} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \frac{\cot^2(c + dx) (B + C \tan(c + dx))}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B + C \tan(c + dx)}{\tan(c + dx)^2 (a + b \tan(c + dx))} dx \\
 & \quad \downarrow \text{4092} \\
 & - \frac{\int \frac{\cot(c + dx) (bB \tan^2(c + dx) + aB \tan(c + dx) + bB - aC)}{a + b \tan(c + dx)} dx}{a} - \frac{B \cot(c + dx)}{ad} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{\int \frac{bB \tan(c+dx)^2 + aB \tan(c+dx) + bB - aC}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a} - \frac{B \cot(c+dx)}{ad} \\
 & \quad \downarrow 4134 \\
 & - \frac{b^2(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(bB-aC) \int \cot(c+dx) dx}{a} + \frac{ax(aB+bC)}{a^2+b^2} - \frac{B \cot(c+dx)}{ad} \\
 & \quad \downarrow 3042 \\
 & - \frac{b^2(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(bB-aC) \int -\tan(c+dx + \frac{\pi}{2}) dx}{a} + \frac{ax(aB+bC)}{a^2+b^2} - \frac{B \cot(c+dx)}{ad} \\
 & \quad \downarrow 25 \\
 & - \frac{b^2(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{(bB-aC) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} + \frac{ax(aB+bC)}{a^2+b^2} - \frac{B \cot(c+dx)}{ad} \\
 & \quad \downarrow 3956 \\
 & - \frac{b^2(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{ax(aB+bC)}{a^2+b^2} + \frac{(bB-aC) \log(-\sin(c+dx))}{ad} - \frac{B \cot(c+dx)}{ad} \\
 & \quad \downarrow 4013 \\
 & - \frac{b^2(bB-aC) \log(a \cos(c+dx) + b \sin(c+dx))}{ad(a^2+b^2)} + \frac{ax(aB+bC)}{a^2+b^2} + \frac{(bB-aC) \log(-\sin(c+dx))}{ad} - \frac{B \cot(c+dx)}{ad}
 \end{aligned}$$

input

```
Int[(Cot[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]
```

output

```
-((B*Cot[c + d*x])/(a*d)) - ((a*(a*B + b*C)*x)/(a^2 + b^2) + ((b*B - a*C)*Log[-Sin[c + d*x]])/(a*d) - (b^2*(b*B - a*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/a
```

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3956 $\text{Int}[\tan[(\text{c}_.) + (\text{d}_.)*(\text{x}_)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d}*x], \text{x}]]/\text{d}, \text{x}] \text{ ; FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4013 $\text{Int}[((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)])/((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c}/(\text{b}*f))*\text{Log}[\text{RemoveContent}[\text{a}*Cos[\text{e} + \text{f}*x] + \text{b}*Sin[\text{e} + \text{f}*x], \text{x}]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{a}*c + \text{b}*d, 0]$
- rule 4092 $\text{Int}[((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)])^{\text{m}_}*((\text{A}_.) + (\text{B}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)])^{\text{n}_}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{b}*(\text{A}*b - \text{a}*B)*(a + b*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1}*((c + d*\text{Tan}[\text{e} + \text{f}*x])^{\text{n} + 1}/(f*(\text{m} + 1)*(b*c - \text{a}*d)*(a^2 + b^2))), \text{x}] + \text{Simp}[1/((\text{m} + 1)*(b*c - \text{a}*d)*(a^2 + b^2)) \quad \text{Int}[(a + b*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1}*(c + d*\text{Tan}[\text{e} + \text{f}*x])^{\text{n}}*\text{Simp}[\text{b}*B*(b*c*(\text{m} + 1) + \text{a}*d*(\text{n} + 1)) + \text{A}*(\text{a}*(b*c - \text{a}*d)*(\text{m} + 1) - \text{b}^2*d*(\text{m} + \text{n} + 2)) - (\text{A}*b - \text{a}*B)*(b*c - \text{a}*d)*(\text{m} + 1)*\text{Tan}[\text{e} + \text{f}*x] - \text{b}*d*(\text{A}*b - \text{a}*B)*(\text{m} + \text{n} + 2)*\text{Tan}[\text{e} + \text{f}*x]^2, \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ (\text{IntegerQ}[\text{m}] \ || \ \text{IntegersQ}[2*\text{m}, 2*\text{n}]) \ \&\& \ !(\ \text{ILtQ}[\text{n}, -1] \ \&\& \ (\ !\text{IntegerQ}[\text{m}] \ || \ (\text{EqQ}[\text{c}, 0] \ \&\& \ \text{NeQ}[\text{a}, 0])))$
- rule 4115 $\text{Int}[((\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)])^{\text{m}_}*((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)])^{\text{n}_}*((\text{A}_.) + (\text{B}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)] + (\text{C}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(\text{x}_)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[1/\text{b}^2 \quad \text{Int}[(a + b*\text{Tan}[\text{e} + \text{f}*x])^{\text{m} + 1}*(c + d*\text{Tan}[\text{e} + \text{f}*x])^{\text{n}}*(\text{b}*B - \text{a}*C + \text{b}*C*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}, \text{m}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{EqQ}[\text{A}*b^2 - \text{a}*b*B + \text{a}^2*C, 0]$

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.18

method	result
parallelrisch	$\frac{(2Bb^3 - 2Ca^2b^2) \ln(a + b \tan(dx + c)) + (Ba^2b - Ca^3) \ln(\sec(dx + c)^2) - 2(a^2 + b^2)(Bb - Ca) \ln(\tan(dx + c)) - 2a(B \cot(dx + c) + C \tan(dx + c))}{2a^2d(a^2 + b^2)}$
derivativedivides	$\frac{(Bb - Ca)b^2 \ln(a + b \tan(dx + c)) + \frac{(Bb - Ca) \ln(1 + \tan(dx + c)^2)}{2} + (-Ba - Cb) \arctan(\tan(dx + c)) - \frac{B}{a \tan(dx + c)} + \frac{(-Bb + Ca) \ln(\tan(dx + c))}{a^2}}{d}$
default	$\frac{(Bb - Ca)b^2 \ln(a + b \tan(dx + c)) + \frac{(Bb - Ca) \ln(1 + \tan(dx + c)^2)}{2} + (-Ba - Cb) \arctan(\tan(dx + c)) - \frac{B}{a \tan(dx + c)} + \frac{(-Bb + Ca) \ln(\tan(dx + c))}{a^2}}{d}$
norman	$\frac{-\frac{B \tan(dx + c)}{ad} - \frac{(Ba + Cb)x \tan(dx + c)^2}{a^2 + b^2}}{\tan(dx + c)^2} + \frac{b^2(Bb - Ca) \ln(a + b \tan(dx + c))}{a^2d(a^2 + b^2)} - \frac{(Bb - Ca) \ln(\tan(dx + c))}{a^2d} + \frac{(Bb - Ca) \ln(\tan(dx + c))}{a^2}$
risch	$\frac{xB}{ib - a} - \frac{ixC}{ib - a} + \frac{2iBbx}{a^2} + \frac{2iBbc}{a^2d} - \frac{2iCx}{a} - \frac{2iCc}{ad} - \frac{2ib^3Bx}{a^2(a^2 + b^2)} - \frac{2ib^3Bc}{a^2d(a^2 + b^2)} + \frac{2ib^2Cx}{a(a^2 + b^2)} + \frac{2ib^2Cc}{ad(a^2 + b^2)}$

input

```
int(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/2*((2*B*b^3-2*C*a*b^2)*ln(a+b*tan(d*x+c))+(B*a^2*b-C*a^3)*ln(sec(d*x+c)^2)-2*(a^2+b^2)*(B*b-C*a)*ln(tan(d*x+c))-2*a*(B*cot(d*x+c)*(a^2+b^2)+a*d*x*(B*a+C*b)))/a^2/d/(a^2+b^2)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.72

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx =$$

$$\frac{2Ba^3 + 2Bab^2 + 2(Ba^3 + Ca^2b)dx \tan(dx + c) - (Ca^3 - Ba^2b + Cab^2 - Bb^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)}{2(a^4 + a^2b^2)d \tan(dx + c)}$$

input

```
integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

output

```
-1/2*(2*B*a^3 + 2*B*a*b^2 + 2*(B*a^3 + C*a^2*b)*d*x*tan(d*x + c) - (C*a^3 - B*a^2*b + C*a*b^2 - B*b^3)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c) + (C*a*b^2 - B*b^3)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1))*tan(d*x + c))/((a^4 + a^2*b^2)*d*tan(d*x + c))
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.88 (sec) , antiderivative size = 2067, normalized size of antiderivative = 20.07

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx = \text{Too large to display}$$

input

```
integrate(cot(d*x+c)**3*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c)),x)
```

output

```
Piecewise((nan, Eq(a, 0) & Eq(b, 0) & Eq(c, 0) & Eq(d, 0)), ((-B*x - B/(d*
tan(c + d*x)) - C*log(tan(c + d*x)**2 + 1)/(2*d) + C*log(tan(c + d*x))/d)/
a, Eq(b, 0)), ((B*log(tan(c + d*x)**2 + 1)/(2*d) - B*log(tan(c + d*x))/d -
B/(2*d*tan(c + d*x)**2) - C*x - C/(d*tan(c + d*x)))/b, Eq(a, 0)), (-3*B*d
*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 3*I*B*
d*x*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - I*B*log(
tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(
c + d*x)) + B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*a*d*tan(c + d*x)**2
+ 2*I*a*d*tan(c + d*x)) + 2*I*B*log(tan(c + d*x))*tan(c + d*x)**2/(2*a*d*
tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 2*B*log(tan(c + d*x))*tan(c + d*
x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 3*B*tan(c + d*x)/(2*a*
d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 2*I*B/(2*a*d*tan(c + d*x)**2 +
2*I*a*d*tan(c + d*x)) + I*C*d*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 +
2*I*a*d*tan(c + d*x)) - C*d*x*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*
d*tan(c + d*x)) - C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c
+ d*x)**2 + 2*I*a*d*tan(c + d*x)) - I*C*log(tan(c + d*x)**2 + 1)*tan(c + d
*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + 2*C*log(tan(c + d*x))
*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + 2*I*C*lo
g(tan(c + d*x))*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)
) + I*C*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)), Eq...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.27

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx =$$

$$-\frac{\frac{2(Ba+Cb)(dx+c)}{a^2+b^2} + \frac{2(Cab^2-Bb^3) \log(b \tan(dx+c)+a)}{a^4+a^2b^2} + \frac{(Ca-Bb) \log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d} - \frac{2(Ca-Bb) \log(\tan(dx+c))}{a^2} + \frac{2B}{a \tan(dx+c)}$$

input

```
integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, a
lgorithm="maxima")
```

output

```
-1/2*(2*(B*a + C*b)*(d*x + c)/(a^2 + b^2) + 2*(C*a*b^2 - B*b^3)*log(b*tan(
d*x + c) + a)/(a^4 + a^2*b^2) + (C*a - B*b)*log(tan(d*x + c)^2 + 1)/(a^2 +
b^2) - 2*(C*a - B*b)*log(tan(d*x + c))/a^2 + 2*B/(a*tan(d*x + c)))/d
```


Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.42

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= -\frac{(Ba + Cb)(dx + c)}{a^2d + b^2d} - \frac{(Ca - Bb) \log(\tan(dx + c)^2 + 1)}{2(a^2d + b^2d)}$$

$$- \frac{(Cab^3 - Bb^4) \log(|b \tan(dx + c) + a|)}{a^4bd + a^2b^3d}$$

$$+ \frac{(Ca - Bb) \log(|\tan(dx + c)|)}{a^2d} - \frac{B}{ad \tan(dx + c)}$$

input

```
integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")
```

output

```
-(B*a + C*b)*(d*x + c)/(a^2*d + b^2*d) - 1/2*(C*a - B*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) - (C*a*b^3 - B*b^4)*log(abs(b*tan(d*x + c) + a))/(a^4*b*d + a^2*b^3*d) + (C*a - B*b)*log(abs(tan(d*x + c)))/(a^2*d) - B/(a*d*tan(d*x + c))
```

Mupad [B] (verification not implemented)

Time = 6.81 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.36

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{\ln(a + b \tan(c + dx)) (B b^3 - C a b^2)}{d (a^4 + a^2 b^2)}$$

$$- \frac{\ln(\tan(c + dx)) (B b - C a)}{a^2 d} + \frac{\ln(\tan(c + dx) + i) (B - C i)}{2 d (b + a i)}$$

$$- \frac{B \cot(c + dx)}{a d} + \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2 d (a + b i)}$$

input

```
int((cot(c + d*x)^3*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)),x)
```

output

```
(log(a + b*tan(c + d*x))*(B*b^3 - C*a*b^2))/(d*(a^4 + a^2*b^2)) - (log(tan
(c + d*x))*(B*b - C*a))/(a^2*d) + (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d
*(a*1i + b)) - (B*cot(c + d*x))/(a*d) + (log(tan(c + d*x) - 1i)*(B*1i - C
))/(2*d*(a + b*1i))
```

Reduce [F]

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \int \frac{\cot(dx + c)^3 (B \tan(dx + c) + C \tan(dx + c)^2)}{a + \tan(dx + c) b} dx$$

input

```
int(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)
```

output

```
int(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)
```

3.31
$$\int \frac{\cot^4(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx$$

Optimal result	358
Mathematica [C] (verified)	359
Rubi [A] (verified)	359
Maple [A] (verified)	364
Fricas [A] (verification not implemented)	364
Sympy [C] (verification not implemented)	365
Maxima [A] (verification not implemented)	366
Giac [A] (verification not implemented)	367
Mupad [B] (verification not implemented)	367
Reduce [F]	368

Optimal result

Integrand size = 40, antiderivative size = 137

$$\begin{aligned} & \int \frac{\cot^4(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{a+b \tan(c+dx)} dx \\ &= \frac{(bB-aC)x}{a^2+b^2} + \frac{(bB-aC) \cot(c+dx)}{a^2d} - \frac{B \cot^2(c+dx)}{2ad} \\ & \quad - \frac{(a^2B-b^2B+abC) \log(\sin(c+dx))}{a^3d} \\ & \quad - \frac{b^3(bB-aC) \log(a \cos(c+dx)+b \sin(c+dx))}{a^3(a^2+b^2)d} \end{aligned}$$

output

```
(B*b-C*a)*x/(a^2+b^2)+(B*b-C*a)*cot(d*x+c)/a^2/d-1/2*B*cot(d*x+c)^2/a/d-(B
*a^2-B*b^2+C*a*b)*ln(sin(d*x+c))/a^3/d-b^3*(B*b-C*a)*ln(a*cos(d*x+c)+b*sin
(d*x+c))/a^3/(a^2+b^2)/d
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19

$$\int \frac{\cot^4(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{\frac{2(bB-aC) \cot(c+dx)}{a^2} - \frac{B \cot^2(c+dx)}{a} + \frac{(B+iC) \log(i-\tan(c+dx))}{a+ib} - \frac{2(a^2B-b^2B+abC) \log(\tan(c+dx))}{a^3} + \frac{(B-iC) \log(i+\tan(c+dx))}{a-ib}}{2d}$$

input

```
Integrate[(Cot[c + d*x]^4*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]
```

output

```
((2*(b*B - a*C)*Cot[c + d*x])/a^2 - (B*Cot[c + d*x]^2)/a + ((B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b) - (2*(a^2*B - b^2*B + a*b*C)*Log[Tan[c + d*x]])/a^3 + ((B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b) + (2*b^3*(-(b*B) + a*C)*Log[a + b*Tan[c + d*x]])/(a^3*(a^2 + b^2)))/(2*d)
```

Rubi [A] (verified)

Time = 1.99 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {3042, 4115, 3042, 4092, 27, 3042, 4132, 25, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^4(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{a + b \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \tan(c+dx) + C \tan^2(c+dx)^2}{\tan(c+dx)^4(a + b \tan(c+dx))} dx$$

$$\downarrow \text{4115}$$

$$\int \frac{\cot^3(c+dx)(B + C \tan(c+dx))}{a + b \tan(c+dx)} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{B + C \tan(c + dx)}{\tan(c + dx)^3 (a + b \tan(c + dx))} dx \\
& \downarrow 4092 \\
& \frac{\int \frac{2 \cot^2(c + dx) (bB \tan^2(c + dx) + aB \tan(c + dx) + bB - aC)}{a + b \tan(c + dx)} dx}{2a} - \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 27 \\
& \frac{\int \frac{\cot^2(c + dx) (bB \tan^2(c + dx) + aB \tan(c + dx) + bB - aC)}{a + b \tan(c + dx)} dx}{a} - \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 3042 \\
& \frac{\int \frac{bB \tan(c + dx)^2 + aB \tan(c + dx) + bB - aC}{\tan(c + dx)^2 (a + b \tan(c + dx))} dx}{a} - \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 4132 \\
& \frac{\int \frac{\cot(c + dx) (Ba^2 + C \tan(c + dx)a^2 + bCa - b(bB - aC) \tan^2(c + dx) - b^2B)}{a + b \tan(c + dx)} dx}{a} - \frac{(bB - aC) \cot(c + dx)}{ad} \\
& \frac{a}{2ad} \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 25 \\
& \frac{\int \frac{\cot(c + dx) (Ba^2 + C \tan(c + dx)a^2 + bCa - b(bB - aC) \tan^2(c + dx) - b^2B)}{a + b \tan(c + dx)} dx}{a} - \frac{(bB - aC) \cot(c + dx)}{ad} - \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 3042 \\
& \frac{\int \frac{Ba^2 + C \tan(c + dx)a^2 + bCa - b(bB - aC) \tan(c + dx)^2 - b^2B}{\tan(c + dx) (a + b \tan(c + dx))} dx}{a} - \frac{(bB - aC) \cot(c + dx)}{ad} - \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 4134 \\
& \frac{(a^2B + abC - b^2B) \int \cot(c + dx) dx}{a} + \frac{b^3(bB - aC) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a(a^2 + b^2)} - \frac{a^2x(bB - aC)}{a^2 + b^2} - \frac{(bB - aC) \cot(c + dx)}{ad} \\
& \frac{a}{2ad} \frac{B \cot^2(c + dx)}{2ad} \\
& \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
 & \frac{\frac{(a^2B+abC-b^2B) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} + \frac{b^3(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(bB-aC)}{a^2+b^2}}{a} - \frac{(bB-aC) \cot(c+dx)}{ad} \\
 & \qquad \qquad \qquad \frac{B \cot^2(c+dx)}{2ad} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & \frac{\frac{(a^2B+abC-b^2B) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} + \frac{b^3(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(bB-aC)}{a^2+b^2}}{a} - \frac{(bB-aC) \cot(c+dx)}{ad} \\
 & \qquad \qquad \qquad \frac{B \cot^2(c+dx)}{2ad} \\
 & \qquad \qquad \qquad \downarrow \text{3956} \\
 & \frac{\frac{b^3(bB-aC) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2B+abC-b^2B) \log(-\sin(c+dx))}{ad} - \frac{a^2x(bB-aC)}{a^2+b^2}}{a} - \frac{(bB-aC) \cot(c+dx)}{ad} \\
 & \qquad \qquad \qquad \frac{B \cot^2(c+dx)}{2ad} \\
 & \qquad \qquad \qquad \downarrow \text{4013} \\
 & \frac{\frac{(a^2B+abC-b^2B) \log(-\sin(c+dx))}{ad} - \frac{a^2x(bB-aC)}{a^2+b^2} + \frac{b^3(bB-aC) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)}}{a} - \frac{(bB-aC) \cot(c+dx)}{ad} \\
 & \qquad \qquad \qquad \frac{B \cot^2(c+dx)}{2ad}
 \end{aligned}$$

input `Int[(Cot[c + d*x]^4*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x]),x]`

output `-1/2*(B*Cot[c + d*x]^2)/(a*d) - (-(((b*B - a*C)*Cot[c + d*x])/(a*d)) + (-((a^2*(b*B - a*C)*x)/(a^2 + b^2)) + ((a^2*B - b^2*B + a*b*C)*Log[-Sin[c + d*x]])/(a*d) + (b^3*(b*B - a*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/a/a`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sinn[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`
- rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```


Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{-\frac{(Bb-Ca)b^3 \ln(a+b \tan(dx+c))}{a^3(a^2+b^2)} + \frac{(Ba+Cb) \ln\left(\frac{1+\tan(dx+c)^2}{2}\right) + (Bb-Ca) \arctan(\tan(dx+c))}{a^2+b^2} - \frac{B}{2a \tan(dx+c)^2} - \frac{-Bb+Ca}{a^2 \tan(dx+c)} + \frac{d}{d}}$
default	$\frac{-\frac{(Bb-Ca)b^3 \ln(a+b \tan(dx+c))}{a^3(a^2+b^2)} + \frac{(Ba+Cb) \ln\left(\frac{1+\tan(dx+c)^2}{2}\right) + (Bb-Ca) \arctan(\tan(dx+c))}{a^2+b^2} - \frac{B}{2a \tan(dx+c)^2} - \frac{-Bb+Ca}{a^2 \tan(dx+c)} + \frac{d}{d}}$
parallelrisc	$\frac{(-2Bb^4+2Ca^3b^3) \ln(a+b \tan(dx+c)) + (Ba^4+Ca^3b) \ln(\sec(dx+c)^2) + (-2Ba^4+2Bb^4-2Ca^3b-2Ca^3b^3) \ln(\tan(dx+c))}{2(a^2+b^2)a^3d}$
norman	$\frac{\frac{(Bb-Ca) \tan(dx+c)^2}{a^2d} + \frac{(Bb-Ca)x \tan(dx+c)^3}{a^2+b^2} - \frac{B \tan(dx+c)}{2ad}}{\tan(dx+c)^3} + \frac{(Ba+Cb) \ln\left(\frac{1+\tan(dx+c)^2}{2}\right)}{2d(a^2+b^2)} - \frac{(Ba^2-Bb^2+Cab) \ln(\tan(dx+c))}{a^3d}$
risc	$\frac{2iCbc}{a^2d} + \frac{x C}{ib-a} + \frac{2iBx}{a} + \frac{2iBc}{ad} + \frac{2ib^4Bc}{(a^2+b^2)a^3d} - \frac{2iBb^2c}{a^3d} - \frac{2i(iBa e^{2i(dx+c)} - Bb e^{2i(dx+c)} + Ca e^{2i(dx+c)} + E)}{a^2d(e^{2i(dx+c)} - 1)^2}$

```
input int(cot(d*x+c)^4*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*(-(B*b-C*a)*b^3/a^3/(a^2+b^2)*ln(a+b*tan(d*x+c))+1/(a^2+b^2)*(1/2*(B*a+C*b)*ln(1+tan(d*x+c)^2)+(B*b-C*a)*arctan(tan(d*x+c)))-1/2/a*B/tan(d*x+c)^2-(-B*b+C*a)/a^2/tan(d*x+c)+1/a^3*(-B*a^2+B*b^2-C*a*b)*ln(tan(d*x+c)))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.71

$$\int \frac{\cot^4(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx = \frac{Ba^4 + Ba^2b^2 + (Ba^4 + Ca^3b + Cab^3 - Bb^4) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 - (Cab^3 - Bb^4) \log\left(\frac{b^2 \tan(dx+c)}{a + b \tan(dx+c)}\right)}{d}$$

```
input integrate(cot(d*x+c)^4*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

output

```
-1/2*(B*a^4 + B*a^2*b^2 + (B*a^4 + C*a^3*b + C*a*b^3 - B*b^4)*log(tan(d*x
+ c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^2 - (C*a*b^3 - B*b^4)*log((b^2*t
an(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1))*tan(d*x +
c)^2 + (B*a^4 + B*a^2*b^2 + 2*(C*a^4 - B*a^3*b)*d*x)*tan(d*x + c)^2 + 2*(C
*a^4 - B*a^3*b + C*a^2*b^2 - B*a*b^3)*tan(d*x + c))/((a^5 + a^3*b^2)*d*tan
(d*x + c)^2)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 16.18 (sec) , antiderivative size = 2596, normalized size of antiderivative = 18.95

$$\int \frac{\cot^4(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx = \text{Too large to display}$$

input

```
integrate(cot(d*x+c)**4*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c)),x)
```


Giac [A] (verification not implemented)

Time = 0.79 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.30

$$\int \frac{\cot^4(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= -\frac{(Ca - Bb)(dx + c)}{a^2d + b^2d} + \frac{(Ba + Cb) \log(\tan(dx + c)^2 + 1)}{2(a^2d + b^2d)}$$

$$+ \frac{(Cab^4 - Bb^5) \log(|b \tan(dx + c) + a|)}{a^5bd + a^3b^3d}$$

$$- \frac{(Ba^2 + Cab - Bb^2) \log(|\tan(dx + c)|)}{a^3d} - \frac{Ba^2 + 2(Ca^2 - Bab) \tan(dx + c)}{2a^3d \tan(dx + c)^2}$$

input

```
integrate(cot(d*x+c)^4*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x, algorithm="giac")
```

output

```
-(C*a - B*b)*(d*x + c)/(a^2*d + b^2*d) + 1/2*(B*a + C*b)*log(tan(d*x + c)^2 + 1)/(a^2*d + b^2*d) + (C*a*b^4 - B*b^5)*log(abs(b*tan(d*x + c) + a))/(a^5*b*d + a^3*b^3*d) - (B*a^2 + C*a*b - B*b^2)*log(abs(tan(d*x + c)))/(a^3*d) - 1/2*(B*a^2 + 2*(C*a^2 - B*a*b)*tan(d*x + c))/(a^3*d*tan(d*x + c)^2)
```

Mupad [B] (verification not implemented)

Time = 7.27 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.28

$$\int \frac{\cot^4(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= -\frac{\cot(c + dx)^2 \left(\frac{B}{2a} - \frac{\tan(c + dx)(Bb - Ca)}{a^2} \right)}{d} + \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2d (-b + a i)}$$

$$- \frac{\ln(\tan(c + dx)) (B a^2 + C a b - B b^2)}{a^3 d}$$

$$- \frac{\ln(a + b \tan(c + dx)) (B b^4 - C a b^3)}{d (a^5 + a^3 b^2)} + \frac{\ln(\tan(c + dx) + i) (B - C i)}{2d (a - b i)}$$

input

```
int((cot(c + d*x)^4*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)),x)
```

output

```
(log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*1i - b)) - (cot(c + d*x)^2*(B/
(2*a) - (tan(c + d*x)*(B*b - C*a))/a^2))/d - (log(tan(c + d*x))*(B*a^2 - B
*b^2 + C*a*b))/(a^3*d) - (log(a + b*tan(c + d*x))*(B*b^4 - C*a*b^3))/(d*(a
^5 + a^3*b^2)) + (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a - b*1i))
```

Reduce [F]

$$\int \frac{\cot^4(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \int \frac{\cot(dx + c)^4 (B \tan(dx + c) + C \tan(dx + c)^2)}{a + \tan(dx + c) b} dx$$

input

```
int(cot(d*x+c)^4*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)
```

output

```
int(cot(d*x+c)^4*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c)),x)
```

3.32
$$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

Optimal result	369
Mathematica [C] (verified)	370
Rubi [A] (verified)	370
Maple [A] (verified)	375
Fricas [B] (verification not implemented)	375
Sympy [C] (verification not implemented)	376
Maxima [A] (verification not implemented)	377
Giac [A] (verification not implemented)	378
Mupad [B] (verification not implemented)	379
Reduce [B] (verification not implemented)	379

Optimal result

Integrand size = 40, antiderivative size = 208

$$\begin{aligned} & \int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= -\frac{(2abB - a^2C + b^2C) x}{(a^2 + b^2)^2} + \frac{(a^2B - b^2B + 2abC) \log(\cos(c+dx))}{(a^2 + b^2)^2 d} \\ &+ \frac{a^2(a^2bB + 3b^3B - 2a^3C - 4ab^2C) \log(a+b \tan(c+dx))}{b^3(a^2 + b^2)^2 d} \\ &- \frac{(abB - 2a^2C - b^2C) \tan(c+dx)}{b^2(a^2 + b^2) d} + \frac{a(bB - aC) \tan^2(c+dx)}{b(a^2 + b^2) d(a+b \tan(c+dx))} \end{aligned}$$

output

```

-(2*B*a*b-C*a^2+C*b^2)*x/(a^2+b^2)^2+(B*a^2-B*b^2+2*C*a*b)*ln(cos(d*x+c))/
(a^2+b^2)^2/d+a^2*(B*a^2*b+3*B*b^3-2*C*a^3-4*C*a*b^2)*ln(a+b*tan(d*x+c))/b
^3/(a^2+b^2)^2/d-(B*a*b-2*C*a^2-C*b^2)*tan(d*x+c)/b^2/(a^2+b^2)/d+a*(B*b-C
*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))
    
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.90 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.93

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx =$$

$$\frac{\frac{(B+iC) \log(i - \tan(c+dx))}{(a+ib)^2} + \frac{(B-iC) \log(i + \tan(c+dx))}{(a-ib)^2} + \frac{2a^2(-a^2bB - 3b^3B + 2a^3C + 4ab^2C) \log(a+b \tan(c+dx))}{b^3(a^2+b^2)^2} + \frac{2a^2(-abB + 2a^2C)}{b^3(a^2+b^2)(a+ib)}}{2d}$$

input

```
Integrate[(Tan[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]
```

output

```
-1/2*((B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)^2 + ((B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*a^2*(-(a^2*b*B) - 3*b^3*B + 2*a^3*C + 4*a*b^2*C)*Log[a + b*Tan[c + d*x]])/(b^3*(a^2 + b^2)^2) + (2*a^2*(-(a*b*B) + 2*a^2*C + b^2*C))/(b^3*(a^2 + b^2)*(a + b*Tan[c + d*x])) - (2*C*Tan[c + d*x]^2)/(b*(a + b*Tan[c + d*x]))/d
```

Rubi [A] (verified)

Time = 2.21 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.08, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {3042, 4115, 3042, 4088, 25, 3042, 4130, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c + dx)^2 (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$\downarrow 4115$$

$$\begin{aligned}
& \int \frac{\tan^3(c+dx)(B+C \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\tan(c+dx)^3(B+C \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
& \quad \downarrow \text{4088} \\
& \frac{\int -\frac{\tan(c+dx)((-2Ca^2+bBa-b^2C) \tan^2(c+dx)-b(bB-aC) \tan(c+dx)+2a(bB-aC))}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \\
& \quad \frac{a(bB-aC) \tan^2(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow \text{25} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
& \frac{\int \frac{\tan(c+dx)((-2Ca^2+bBa-b^2C) \tan^2(c+dx)-b(bB-aC) \tan(c+dx)+2a(bB-aC))}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
& \frac{\int \frac{\tan(c+dx)((-2Ca^2+bBa-b^2C) \tan(c+dx)^2-b(bB-aC) \tan(c+dx)+2a(bB-aC))}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{4130} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
& \frac{\int -\frac{((aB+bC) \tan(c+dx)b^2)+(a^2+b^2)(bB-2aC) \tan^2(c+dx)+a(-2Ca^2+bBa-b^2C)}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \frac{(-2a^2C+abB-b^2C) \tan(c+dx)}{bd} \\
& \quad \downarrow \text{25} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
& \frac{(-2a^2C+abB-b^2C) \tan(c+dx)}{bd} - \frac{\int -\frac{((aB+bC) \tan(c+dx)b^2)+(a^2+b^2)(bB-2aC) \tan^2(c+dx)+a(-2Ca^2+bBa-b^2C)}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2C + abB - b^2C) \tan(c + dx)}{bd}}{\frac{\int \frac{-(aB + bC) \tan(c + dx)b^2 + (a^2 + b^2)(bB - 2aC) \tan(c + dx)^2 + a(-2Ca^2 + bBa - b^2C)}{a + b \tan(c + dx)} dx}{b(a^2 + b^2)}}$$

↓ 4109

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2C + abB - b^2C) \tan(c + dx)}{bd}}{\frac{b^2(a^2B + 2abC - b^2B) \int \tan(c + dx) dx}{a^2 + b^2} + \frac{a^2(-2a^3C + a^2bB - 4ab^2C + 3b^3B) \int \frac{\tan^2(c + dx) + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{b^2x(a^2(-C) + 2abB)}{a^2 + b^2}}$$

↓ 3042

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2C + abB - b^2C) \tan(c + dx)}{bd}}{\frac{b^2(a^2B + 2abC - b^2B) \int \tan(c + dx) dx}{a^2 + b^2} + \frac{a^2(-2a^3C + a^2bB - 4ab^2C + 3b^3B) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{b^2x(a^2(-C) + 2abB)}{a^2 + b^2}}$$

↓ 3956

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2C + abB - b^2C) \tan(c + dx)}{bd}}{\frac{a^2(-2a^3C + a^2bB - 4ab^2C + 3b^3B) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^2(a^2B + 2abC - b^2B) \log(\cos(c + dx))}{d(a^2 + b^2)} - \frac{b^2x(a^2(-C) + 2abB)}{a^2 + b^2}}$$

↓ 4100

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2C + abB - b^2C) \tan(c + dx)}{bd}}{\frac{a^2(-2a^3C + a^2bB - 4ab^2C + 3b^3B) \int \frac{1}{a + b \tan(c + dx)} d(b \tan(c + dx))}{bd(a^2 + b^2)} + \frac{b^2(a^2B + 2abC - b^2B) \log(\cos(c + dx))}{d(a^2 + b^2)} - \frac{b^2x(a^2(-C) + 2abB + b^2C)}{a^2 + b^2}}$$

↓ 16

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2C + abB - b^2C) \tan(c + dx)}{bd}}{\frac{b^2(a^2B + 2abC - b^2B) \log(\cos(c + dx))}{d(a^2 + b^2)} - \frac{b^2x(a^2(-C) + 2abB + b^2C)}{a^2 + b^2} + \frac{a^2(-2a^3C + a^2bB - 4ab^2C + 3b^3B) \log(a + b \tan(c + dx))}{bd(a^2 + b^2)}}$$

input $\text{Int}[(\text{Tan}[c + d*x]^2*(B*\text{Tan}[c + d*x] + C*\text{Tan}[c + d*x]^2))/(a + b*\text{Tan}[c + d*x])^2, x]$

output $(a*(b*B - a*C)*\text{Tan}[c + d*x]^2)/(b*(a^2 + b^2)*d*(a + b*\text{Tan}[c + d*x])) - (-((-(b^2*(2*a*b*B - a^2*C + b^2*C)*x)/(a^2 + b^2)) + (b^2*(a^2*B - b^2*B + 2*a*b*C)*\text{Log}[\text{Cos}[c + d*x]])/(a^2 + b^2)*d + (a^2*(a^2*b*B + 3*b^3*B - 2*a^3*C - 4*a*b^2*C)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(b*(a^2 + b^2)*d))/b) + ((a*b*B - 2*a^2*C - b^2*C)*\text{Tan}[c + d*x])/(b*d))/(b*(a^2 + b^2))$

Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}\{a, b, c\}, x]$

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\text{tan}[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

rule 4088 $\text{Int}[(a_)+(b_)*\text{tan}[(e_)+(f_)*(x_)]^m*((A_)+(B_)*\text{tan}[(e_)+(f_)*(x_)]*(c_)+(d_)*\text{tan}[(e_)+(f_)*(x_)]^n), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*(B*c - A*d)*(a + b*\text{Tan}[e + f*x])^{m-1}*((c + d*\text{Tan}[e + f*x])^{n+1}/(d*f*(n+1)*(c^2 + d^2))), x] - \text{Simp}[1/(d*(n+1)*(c^2 + d^2)) \text{ Int}[(a + b*\text{Tan}[e + f*x])^{m-2}*(c + d*\text{Tan}[e + f*x])^{n+1}*\text{Simp}[a*A*d*(b*d*(m-1) - a*c*(n+1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m-1) + a*d*(n+1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n+1)*\text{Tan}[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m+n) - b*B*(c^2*(m-1) - d^2*(n+1)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegerQ}[m] || \text{IntegersQ}[2*m, 2*n])$

rule 4100

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

rule 4109

```
Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2
)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_
.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_
.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\tan(dx+c)C}{b^2} + \frac{a^2(Ba^2b+3Bb^3-2Ca^3-4Cab^2)\ln(a+b\tan(dx+c))}{b^3(a^2+b^2)^2} + \frac{a^3(Bb-Ca)}{b^3(a^2+b^2)(a+b\tan(dx+c))} + \frac{(-Ba^2+Bb^2-2Cab)\ln(1+\tan(dx+c))}{2(a^2+b^2)}$
default	$\frac{\tan(dx+c)C}{b^2} + \frac{a^2(Ba^2b+3Bb^3-2Ca^3-4Cab^2)\ln(a+b\tan(dx+c))}{b^3(a^2+b^2)^2} + \frac{a^3(Bb-Ca)}{b^3(a^2+b^2)(a+b\tan(dx+c))} + \frac{(-Ba^2+Bb^2-2Cab)\ln(1+\tan(dx+c))}{2(a^2+b^2)}$
norman	$\frac{C\tan(dx+c)^2}{bd} + \frac{(Ba^2b-2Ca^3-Cab^2)a}{db^3(a^2+b^2)} - \frac{a(2Bab-Ca^2+Cb^2)x}{a^4+2b^2a^2+b^4} - \frac{b(2Bab-Ca^2+Cb^2)x\tan(dx+c)}{a^4+2b^2a^2+b^4} + \frac{a^2(Ba^2b+3Bb^3-2Ca^3-4Cab^2)\ln(a+b\tan(dx+c))}{(a^2+b^2)^2}$
parallelrisc	$-\frac{-2Ba^5b-2Ba^3b^3+2Ca^2b^4+4Ca^6+6Ca^4b^2-2C\tan(dx+c)^2b^6+B\ln(1+\tan(dx+c)^2)\tan(dx+c)a^2b^4-2B\ln(a+b\tan(dx+c))}{(a^2+b^2)^2}$
risc	$-\frac{4iCax}{b^3} - \frac{xC}{2iab-a^2+b^2} + \frac{8ia^3Cc}{(a^4+2b^2a^2+b^4)bd} + \frac{2i(-Ba^3be^{2i(dx+c)}+2Ca^4e^{2i(dx+c)}-Cb^4e^{2i(dx+c)}-2iCa^3be^{i(dx+c)})}{(e^{2i(dx+c)}+1)(ib+a)(-ib+a)^2(-ibe^{i(dx+c)}+1)}$

input `int (tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,method =_RETURNVERBOSE)`

output `1/d*(tan(d*x+c)*C/b^2+1/b^3*a^2*(B*a^2*b+3*B*b^3-2*C*a^3-4*C*a*b^2)/(a^2+b^2)^2*ln(a+b*tan(d*x+c))+1/b^3*a^3*(B*b-C*a)/(a^2+b^2)/(a+b*tan(d*x+c))+1/(a^2+b^2)^2*(1/2*(-B*a^2+B*b^2-2*C*a*b)*ln(1+tan(d*x+c)^2)+(-2*B*a*b+C*a^2-C*b^2)*arctan(tan(d*x+c))))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 434 vs. 2(209) = 418.

Time = 0.19 (sec) , antiderivative size = 434, normalized size of antiderivative = 2.09

$$\int \frac{\tan^2(c+dx)(B\tan(c+dx)+C\tan^2(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{2Ca^4b^2-2Ba^3b^3-2(Ca^3b^3-2Ba^2b^4-Cab^5)dx-2(Ca^4b^2+2Ca^2b^4+Cb^6)\tan(dx+c)^2+(2C$$

input `integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="fricas")`

output `-1/2*(2*C*a^4*b^2 - 2*B*a^3*b^3 - 2*(C*a^3*b^3 - 2*B*a^2*b^4 - C*a*b^5)*d*x - 2*(C*a^4*b^2 + 2*C*a^2*b^4 + C*b^6)*tan(d*x + c)^2 + (2*C*a^6 - B*a^5*b + 4*C*a^4*b^2 - 3*B*a^3*b^3 + (2*C*a^5*b - B*a^4*b^2 + 4*C*a^3*b^3 - 3*B*a^2*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (2*C*a^6 - B*a^5*b + 4*C*a^4*b^2 - 2*B*a^3*b^3 + 2*C*a^2*b^4 - B*a*b^5 + (2*C*a^5*b - B*a^4*b^2 + 4*C*a^3*b^3 - 2*B*a^2*b^4 + 2*C*a*b^5 - B*b^6)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 2*(2*C*a^5*b - B*a^4*b^2 + 2*C*a^3*b^3 + C*a*b^5 + (C*a^2*b^4 - 2*B*a*b^5 - C*b^6)*d*x)*tan(d*x + c))/((a^4*b^4 + 2*a^2*b^6 + b^8)*d*tan(d*x + c) + (a^5*b^3 + 2*a^3*b^5 + a*b^7)*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.11 (sec) , antiderivative size = 4541, normalized size of antiderivative = 21.83

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**2,x)`

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2), Eq(a, 0) & Eq(b, 0) & Eq(d, 0))
, ((-B*log(tan(c + d*x)**2 + 1)/(2*d) + B*tan(c + d*x)**2/(2*d) + C*x + C*
tan(c + d*x)**3/(3*d) - C*tan(c + d*x)/d)/a**2, Eq(b, 0)), (3*I*B*d*x*tan(
c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d
) + 6*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*
x) - 4*b**2*d) - 3*I*B*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c +
d*x) - 4*b**2*d) + 2*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*b**2*d*
tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*I*B*log(tan(c +
d*x)**2 + 1)*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d
*x) - 4*b**2*d) - 2*B*log(tan(c + d*x)**2 + 1)/(4*b**2*d*tan(c + d*x)**2 -
8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 5*I*B*tan(c + d*x)/(4*b**2*d*tan(c
+ d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*B/(4*b**2*d*tan(c + d*
x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 9*C*d*x*tan(c + d*x)**2/(4*b
**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 18*I*C*d*x*t
an(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d
) + 9*C*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d
) + 4*I*C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)*
**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 8*C*log(tan(c + d*x)**2 + 1)*ta
n(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d)
- 4*I*C*log(tan(c + d*x)**2 + 1)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.06

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{2(Ca^2 - 2Bab - Cb^2)(dx+c)}{a^4 + 2a^2b^2 + b^4} - \frac{2(2Ca^5 - Ba^4b + 4Ca^3b^2 - 3Ba^2b^3) \log(b \tan(dx+c) + a)}{a^4b^3 + 2a^2b^5 + b^7} - \frac{(Ba^2 + 2Cab - Bb^2) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} - \frac{3}{a^3} - \frac{1}{2d}$$

input

```
integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="maxima")
```

output

$$\begin{aligned} & \frac{1}{2} * (2 * (C * a^2 - 2 * B * a * b - C * b^2) * (d * x + c) / (a^4 + 2 * a^2 * b^2 + b^4) - 2 * (2 * \\ & C * a^5 - B * a^4 * b + 4 * C * a^3 * b^2 - 3 * B * a^2 * b^3) * \log(b * \tan(d * x + c) + a) / (a^4 * \\ & b^3 + 2 * a^2 * b^5 + b^7) - (B * a^2 + 2 * C * a * b - B * b^2) * \log(\tan(d * x + c)^2 + 1) \\ & / (a^4 + 2 * a^2 * b^2 + b^4) - 2 * (C * a^4 - B * a^3 * b) / (a^3 * b^3 + a * b^5 + (a^2 * b^4 \\ & + b^6) * \tan(d * x + c)) + 2 * C * \tan(d * x + c) / b^2) / d \end{aligned}$$
Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.16

$$\begin{aligned} & \int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx \\ & = \frac{(Ca^2 - 2 Bab - Cb^2)(dx + c)}{a^4d + 2a^2b^2d + b^4d} - \frac{(Ba^2 + 2Cab - Bb^2) \log(\tan(dx + c)^2 + 1)}{2(a^4d + 2a^2b^2d + b^4d)} \\ & - \frac{(2Ca^5 - Ba^4b + 4Ca^3b^2 - 3Ba^2b^3) \log(|b \tan(dx + c) + a|)}{a^4b^3d + 2a^2b^5d + b^7d} \\ & + \frac{C \tan(dx + c)}{b^2d} - \frac{Ca^6 - Ba^5b + Ca^4b^2 - Ba^3b^3}{(a^2 + b^2)^2 (b \tan(dx + c) + a) b^3d} \end{aligned}$$

input

```
integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="giac")
```

output

$$\begin{aligned} & (C * a^2 - 2 * B * a * b - C * b^2) * (d * x + c) / (a^4 * d + 2 * a^2 * b^2 * d + b^4 * d) - 1 / 2 * (B \\ & * a^2 + 2 * C * a * b - B * b^2) * \log(\tan(d * x + c)^2 + 1) / (a^4 * d + 2 * a^2 * b^2 * d + b^4 \\ & * d) - (2 * C * a^5 - B * a^4 * b + 4 * C * a^3 * b^2 - 3 * B * a^2 * b^3) * \log(\text{abs}(b * \tan(d * x + \\ & c) + a)) / (a^4 * b^3 * d + 2 * a^2 * b^5 * d + b^7 * d) + C * \tan(d * x + c) / (b^2 * d) - (C * a \\ & ^6 - B * a^5 * b + C * a^4 * b^2 - B * a^3 * b^3) / ((a^2 + b^2)^2 * (b * \tan(d * x + c) + a) * \\ & b^3 * d) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 6.20 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.01

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{C \tan(c+dx)}{b^2 d} - \frac{\ln(a+b \tan(c+dx)) (2C a^5 - B a^4 b + 4C a^3 b^2 - 3B a^2 b^3)}{d (a^4 b^3 + 2a^2 b^5 + b^7)}$$

$$- \frac{\ln(\tan(c+dx) - i) (B + C i)}{2d (a^2 + a b 2i - b^2)} - \frac{\ln(\tan(c+dx) + i) (C + B i)}{2d (a^2 i + 2ab - b^2 i)}$$

$$- \frac{a^2 (C a^2 - B a b)}{b d (\tan(c+dx) b^3 + a b^2) (a^2 + b^2)}$$

input

```
int((tan(c + d*x))^2*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^2,x
```

output

```
(C*tan(c + d*x))/(b^2*d) - (log(a + b*tan(c + d*x))*(2*C*a^5 - 3*B*a^2*b^3 + 4*C*a^3*b^2 - B*a^4*b))/(d*(b^7 + 2*a^2*b^5 + a^4*b^3)) - (log(tan(c + d*x) - 1i)*(B + C*1i))/(2*d*(a*b*2i + a^2 - b^2)) - (log(tan(c + d*x) + 1i)*(B*1i + C))/(2*d*(2*a*b + a^2*1i - b^2*1i)) - (a^2*(C*a^2 - B*a*b))/(b*d*(a*b^2 + b^3*tan(c + d*x))*(a^2 + b^2))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 555, normalized size of antiderivative = 2.67

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{-4 \log(a + \tan(dx+c) b) a^6 c + 2 \log(a + \tan(dx+c) b) a^5 b^2 + 6 \log(a + \tan(dx+c) b) a^3 b^4 + 2 \tan(dx+c) a^2 b^3}{(a+b \tan(c+dx))^2}$$

input

```
int(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x
```


output

```
( - log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**5 - 2*log(tan(c + d*x)**
2 + 1)*tan(c + d*x)*a*b**5*c + log(tan(c + d*x)**2 + 1)*tan(c + d*x)*b**7
- log(tan(c + d*x)**2 + 1)*a**3*b**4 - 2*log(tan(c + d*x)**2 + 1)*a**2*b**
4*c + log(tan(c + d*x)**2 + 1)*a*b**6 - 4*log(tan(c + d*x)*b + a)*tan(c +
d*x)*a**5*b*c + 2*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**4*b**3 - 8*log(t
an(c + d*x)*b + a)*tan(c + d*x)*a**3*b**3*c + 6*log(tan(c + d*x)*b + a)*ta
n(c + d*x)*a**2*b**5 - 4*log(tan(c + d*x)*b + a)*a**6*c + 2*log(tan(c + d*
x)*b + a)*a**5*b**2 - 8*log(tan(c + d*x)*b + a)*a**4*b**2*c + 6*log(tan(c
+ d*x)*b + a)*a**3*b**4 + 2*tan(c + d*x)**2*a**4*b**2*c + 4*tan(c + d*x)**
2*a**2*b**4*c + 2*tan(c + d*x)**2*b**6*c + 4*tan(c + d*x)*a**5*b*c - 2*tan
(c + d*x)*a**4*b**3 + 6*tan(c + d*x)*a**3*b**3*c - 2*tan(c + d*x)*a**2*b**
5 + 2*tan(c + d*x)*a**2*b**4*c*d*x - 4*tan(c + d*x)*a*b**6*d*x + 2*tan(c +
d*x)*a*b**5*c - 2*tan(c + d*x)*b**6*c*d*x + 2*a**3*b**3*c*d*x - 4*a**2*b*
*5*d*x - 2*a*b**5*c*d*x)/(2*b**3*d*(tan(c + d*x)*a**4*b + 2*tan(c + d*x)*a
**2*b**3 + tan(c + d*x)*b**5 + a**5 + 2*a**3*b**2 + a*b**4))
```

3.33 $\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$

Optimal result	381
Mathematica [C] (verified)	381
Rubi [A] (verified)	382
Maple [A] (verified)	386
Fricas [B] (verification not implemented)	386
Sympy [C] (verification not implemented)	387
Maxima [A] (verification not implemented)	388
Giac [A] (verification not implemented)	389
Mupad [B] (verification not implemented)	390
Reduce [B] (verification not implemented)	390

Optimal result

Integrand size = 38, antiderivative size = 157

$$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= -\frac{(a^2B - b^2B + 2abC)x}{(a^2 + b^2)^2} - \frac{(2abB - a^2C + b^2C) \log(\cos(c+dx))}{(a^2 + b^2)^2 d}$$

$$- \frac{a(2b^3B - a^3C - 3ab^2C) \log(a+b \tan(c+dx))}{b^2(a^2 + b^2)^2 d} - \frac{a^2(bB - aC)}{b^2(a^2 + b^2)d(a+b \tan(c+dx))}$$

```
output - (B*a^2-B*b^2+2*C*a*b)*x/(a^2+b^2)^2-(2*B*a*b-C*a^2+C*b^2)*ln(cos(d*x+c))/
(a^2+b^2)^2/d-a*(2*B*b^3-C*a^3-3*C*a*b^2)*ln(a+b*tan(d*x+c))/b^2/(a^2+b^2)
^2/d-a^2*(B*b-C*a)/b^2/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.84 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.93

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\frac{i(B+iC) \log(i-\tan(c+dx))}{(a+ib)^2} - \frac{i(B-iC) \log(i+\tan(c+dx))}{(a-ib)^2} + \frac{2a \left((-2bB+3aC+\frac{a^3C}{b^2}) \log(a+b \tan(c+dx)) + \frac{a(a^2+b^2)(-bB+aC)}{b^2(a+b \tan(c+dx))} \right)}{(a^2+b^2)^2}}{2d}$$

input `Integrate[(Tan[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]`

output `((I*(B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*(B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*a*((-2*b*B + 3*a*C + (a^3*C)/b^2)*Log[a + b*Tan[c + d*x]] + (a*(a^2 + b^2)*(-(b*B) + a*C))/(b^2*(a + b*Tan[c + d*x]))))/(a^2 + b^2)^2)/(2*d)`

Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.09, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4115, 3042, 4087, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

↓ 4115

$$\int \frac{\tan^2(c+dx)(B + C \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{\tan(c+dx)^2(B+C\tan(c+dx))}{(a+b\tan(c+dx))^2} dx \\
& \quad \downarrow 4087 \\
& \int \frac{-((a^2+b^2)C\tan^2(c+dx))-b(bB-aC)\tan(c+dx)+a(bB-aC)}{a+b\tan(c+dx)} dx}{b(a^2+b^2)} - \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 25 \\
& \int \frac{-((a^2+b^2)C\tan^2(c+dx))-b(bB-aC)\tan(c+dx)+a(bB-aC)}{a+b\tan(c+dx)} dx}{b(a^2+b^2)} - \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \int \frac{-((a^2+b^2)C\tan(c+dx)^2)-b(bB-aC)\tan(c+dx)+a(bB-aC)}{a+b\tan(c+dx)} dx}{b(a^2+b^2)} - \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 4109 \\
& \frac{-\frac{b(a^2(-C)+2abB+b^2C)\int\tan(c+dx)dx}{a^2+b^2} + \frac{a(a^3(-C)-3ab^2C+2b^3B)\int\frac{\tan^2(c+dx)+1}{a+b\tan(c+dx)}dx}{a^2+b^2} + \frac{bx(a^2B+2abC-b^2B)}{a^2+b^2}}{b(a^2+b^2)} \\
& \quad \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{-\frac{b(a^2(-C)+2abB+b^2C)\int\tan(c+dx)dx}{a^2+b^2} + \frac{a(a^3(-C)-3ab^2C+2b^3B)\int\frac{\tan(c+dx)^2+1}{a+b\tan(c+dx)}dx}{a^2+b^2} + \frac{bx(a^2B+2abC-b^2B)}{a^2+b^2}}{b(a^2+b^2)} \\
& \quad \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 3956 \\
& \frac{a(a^3(-C)-3ab^2C+2b^3B)\int\frac{\tan(c+dx)^2+1}{a+b\tan(c+dx)}dx}{a^2+b^2} + \frac{b(a^2(-C)+2abB+b^2C)\log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(a^2B+2abC-b^2B)}{a^2+b^2} \\
& \quad \frac{b(a^2+b^2)}{a^2(bB-aC)} \\
& \quad \frac{b^2d(a^2+b^2)(a+b\tan(c+dx))}{4100}
\end{aligned}$$

$$\begin{aligned}
& \frac{a(a^3(-C)-3ab^2C+2b^3B) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} + \frac{b(a^2(-C)+2abB+b^2C) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(a^2B+2abC-b^2B)}{a^2+b^2} \\
& \frac{b(a^2+b^2)}{a^2(bB-aC)} \\
& \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 16 \\
& \frac{a^2(bB-aC)}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \frac{b(a^2(-C)+2abB+b^2C) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(a^2B+2abC-b^2B)}{a^2+b^2} + \frac{a(a^3(-C)-3ab^2C+2b^3B) \log(a+b \tan(c+dx))}{bd(a^2+b^2)} \\
& \frac{\quad}{b(a^2+b^2)}
\end{aligned}$$

input

```
Int[(Tan[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]
```

output

```
-(((b*(a^2*B - b^2*B + 2*a*b*C)*x)/(a^2 + b^2) + (b*(2*a*b*B - a^2*C + b^2*C)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) + (a*(2*b^3*B - a^3*C - 3*a*b^2*C)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/(b*(a^2 + b^2))) - (a^2*(b*B - a*C))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))
```

Defintions of rubi rules used

rule 16

```
Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4087

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

rule 4100

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

rule 4109

```
Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.99

method	result
derivativedivides	$-\frac{a^2(Bb-Ca)}{b^2(a^2+b^2)(a+b \tan(dx+c))} - \frac{a(2Bb^3-Ca^3-3Cab^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2 b^2} + \frac{(2Bab-Ca^2+Cb^2) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)^2} + \frac{(-Ba^2+Cb^2)}{(a^2+b^2)^2}$
default	$-\frac{a^2(Bb-Ca)}{b^2(a^2+b^2)(a+b \tan(dx+c))} - \frac{a(2Bb^3-Ca^3-3Cab^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2 b^2} + \frac{(2Bab-Ca^2+Cb^2) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)^2} + \frac{(-Ba^2+Cb^2)}{(a^2+b^2)^2}$
norman	$-\frac{a(Ba^2-Bb^2+2Cab)x}{a^4+2b^2a^2+b^4} - \frac{b(Ba^2-Bb^2+2Cab)x \tan(dx+c)}{a^4+2b^2a^2+b^4} - \frac{(Bab-Ca^2)a}{db^2(a^2+b^2)} + \frac{(2Bab-Ca^2+Cb^2) \ln(1+\tan(dx+c)^2)}{2d(a^4+2b^2a^2+b^4)}$
parallelrisch	$2Bx \tan(dx+c)b^5d - 2Bxa^3b^2d + 2Bxa^4b^4d - 4Cx a^2b^3d + 2B \ln(1+\tan(dx+c)^2) \tan(dx+c) a b^4 - 4B \ln(a+b \tan(dx+c))$
risch	$\frac{x B}{2iab-a^2+b^2} - \frac{2ia^3C}{(ib+a)db(-ib+a)^2(-ibe^{2i(dx+c)}+ae^{2i(dx+c)}+ib+a)} - \frac{ixC}{2iab-a^2+b^2} - \frac{6ia^2Cx}{a^4+2b^2a^2+b^4} - \frac{6}{d(a^4+2b^2a^2+b^4)}$

input `int (tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(-a^2*(B*b-C*a)/b^2/(a^2+b^2)/(a+b*tan(d*x+c))-a*(2*B*b^3-C*a^3-3*C*a*b^2)/(a^2+b^2)^2/b^2*ln(a+b*tan(d*x+c))+1/(a^2+b^2)^2*(1/2*(2*B*a*b-C*a^2+C*b^2)*ln(1+tan(d*x+c)^2)+(-B*a^2+B*b^2-2*C*a*b)*arctan(tan(d*x+c))))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(153) = 306.

Time = 0.12 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.98

$$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{2Ca^3b^2 - 2Ba^2b^3 - 2(Ba^3b^2 + 2Ca^2b^3 - Bab^4)dx + (Ca^5 + 3Ca^3b^2 - 2Ba^2b^3 + (Ca^4b + 3Ca^2b^3 - 2C^2ab^2) \ln(1+\tan^2(c+dx)))}{(a+b \tan(c+dx))^2}$$

input `integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output `1/2*(2*C*a^3*b^2 - 2*B*a^2*b^3 - 2*(B*a^3*b^2 + 2*C*a^2*b^3 - B*a*b^4)*d*x + (C*a^5 + 3*C*a^3*b^2 - 2*B*a^2*b^3 + (C*a^4*b + 3*C*a^2*b^3 - 2*B*a*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (C*a^5 + 2*C*a^3*b^2 + C*a*b^4 + (C*a^4*b + 2*C*a^2*b^3 + C*b^5)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 2*(C*a^4*b - B*a^3*b^2 + (B*a^2*b^3 + 2*C*a*b^4 - B*b^5)*d*x)*tan(d*x + c))/((a^4*b^3 + 2*a^2*b^5 + b^7)*d*tan(d*x + c) + (a^5*b^2 + 2*a^3*b^4 + a*b^6)*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.94 (sec) , antiderivative size = 3497, normalized size of antiderivative = 22.27

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**2,x)`

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2)/tan(c), Eq(a, 0) & Eq(b, 0) & Eq
(d, 0)), ((-B*x + B*tan(c + d*x)/d - C*log(tan(c + d*x)**2 + 1)/(2*d) + C*
tan(c + d*x)**2/(2*d))/a**2, Eq(b, 0)), (B*d*x*tan(c + d*x)**2/(4*b**2*d*t
an(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*I*B*d*x*tan(c + d
*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - B*d*x
/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 3*B*tan
(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d)
+ 2*I*B/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) +
3*I*C*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c +
d*x) - 4*b**2*d) + 6*C*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**
2*d*tan(c + d*x) - 4*b**2*d) - 3*I*C*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*
b**2*d*tan(c + d*x) - 4*b**2*d) + 2*C*log(tan(c + d*x)**2 + 1)*tan(c + d*x
)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*I
*C*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b
**2*d*tan(c + d*x) - 4*b**2*d) - 2*C*log(tan(c + d*x)**2 + 1)/(4*b**2*d*tan
(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 5*I*C*tan(c + d*x)/(
4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*C/(4*b**
2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d), Eq(a, -I*b)),
(B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x)
- 4*b**2*d) + 2*I*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.25

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx =$$

$$-\frac{\frac{2(Ba^2 + 2Cab - Bb^2)(dx+c)}{a^4 + 2a^2b^2 + b^4}}{2d} - \frac{2(Ca^4 + 3Ca^2b^2 - 2Bab^3) \log(b \tan(dx+c) + a)}{a^4b^2 + 2a^2b^4 + b^6} + \frac{(Ca^2 - 2Bab - Cb^2) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} - \frac{1}{a^3b^2 + ab}$$

input

```
integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, a
lgorithm="maxima")
```

output

```
-1/2*(2*(B*a^2 + 2*C*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - 2*(C
*a^4 + 3*C*a^2*b^2 - 2*B*a*b^3)*log(b*tan(d*x + c) + a)/(a^4*b^2 + 2*a^2*b
^4 + b^6) + (C*a^2 - 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2
*b^2 + b^4) - 2*(C*a^3 - B*a^2*b)/(a^3*b^2 + a*b^4 + (a^2*b^3 + b^5)*tan(d
*x + c)))/d
```

Giac [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.38

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= -\frac{(Ba^2 + 2Cab - Bb^2)(dx + c)}{a^4d + 2a^2b^2d + b^4d} - \frac{(Ca^2 - 2Bab - Cb^2) \log(\tan(dx + c)^2 + 1)}{2(a^4d + 2a^2b^2d + b^4d)}$$

$$+ \frac{(Ca^4 + 3Ca^2b^2 - 2Bab^3) \log(|b \tan(dx + c) + a|)}{a^4b^2d + 2a^2b^4d + b^6d}$$

$$+ \frac{Ca^5 - Ba^4b + Ca^3b^2 - Ba^2b^3}{(a^2 + b^2)^2(b \tan(dx + c) + a)b^2d}$$

input

```
integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, a
lgorithm="giac")
```

output

```
-(B*a^2 + 2*C*a*b - B*b^2)*(d*x + c)/(a^4*d + 2*a^2*b^2*d + b^4*d) - 1/2*(
C*a^2 - 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/(a^4*d + 2*a^2*b^2*d + b^
4*d) + (C*a^4 + 3*C*a^2*b^2 - 2*B*a*b^3)*log(abs(b*tan(d*x + c) + a))/(a^4
*b^2*d + 2*a^2*b^4*d + b^6*d) + (C*a^5 - B*a^4*b + C*a^3*b^2 - B*a^2*b^3)/
((a^2 + b^2)^2*(b*tan(d*x + c) + a)*b^2*d)
```

Mupad [B] (verification not implemented)

Time = 5.58 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.05

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\ln(\tan(c+dx) + 1i)(C + B 1i)}{2d(-a^2 + ab 2i + b^2)} + \frac{\ln(\tan(c+dx) - 1i)(B + C 1i)}{2d(-a^2 1i + 2ab + b^2 1i)}$$

$$- \frac{a^2(Bb - Ca)}{b^2 d(a^2 + b^2)(a + b \tan(c+dx))}$$

$$+ \frac{a \ln(a + b \tan(c+dx))(Ca^3 + 3Cab^2 - 2Bb^3)}{b^2 d(a^2 + b^2)^2}$$

input `int((tan(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^2,x)`

output `(log(tan(c + d*x) + 1i)*(B*1i + C))/(2*d*(a*b*2i - a^2 + b^2)) + (log(tan(c + d*x) - 1i)*(B + C*1i))/(2*d*(2*a*b - a^2*1i + b^2*1i)) - (a^2*(B*b - C*a))/(b^2*d*(a^2 + b^2)*(a + b*tan(c + d*x))) + (a*log(a + b*tan(c + d*x))*(C*a^3 - 2*B*b^3 + 3*C*a*b^2))/(b^2*d*(a^2 + b^2)^2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 446, normalized size of antiderivative = 2.84

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{-\log(\tan(dx+c)^2 + 1) \tan(dx+c) a^2 b^3 c + 2 \log(\tan(dx+c)^2 + 1) \tan(dx+c) a b^5 + \log(\tan(dx+c)^2 + 1) \tan(dx+c) a^3 c}{(a+b \tan(c+dx))^2}$$

input `int(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x)`

output

```
( - log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**3*c + 2*log(tan(c + d*x)
**2 + 1)*tan(c + d*x)*a*b**5 + log(tan(c + d*x)**2 + 1)*tan(c + d*x)*b**5*
c - log(tan(c + d*x)**2 + 1)*a**3*b**2*c + 2*log(tan(c + d*x)**2 + 1)*a**2
*b**4 + log(tan(c + d*x)**2 + 1)*a*b**4*c + 2*log(tan(c + d*x)*b + a)*tan(
c + d*x)*a**4*b*c + 6*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**2*b**3*c - 4
*log(tan(c + d*x)*b + a)*tan(c + d*x)*a*b**5 + 2*log(tan(c + d*x)*b + a)*a
**5*c + 6*log(tan(c + d*x)*b + a)*a**3*b**2*c - 4*log(tan(c + d*x)*b + a)*
a**2*b**4 - 2*tan(c + d*x)*a**4*b*c + 2*tan(c + d*x)*a**3*b**3 - 2*tan(c +
d*x)*a**2*b**4*d*x - 2*tan(c + d*x)*a**2*b**3*c + 2*tan(c + d*x)*a*b**5 -
4*tan(c + d*x)*a*b**4*c*d*x + 2*tan(c + d*x)*b**6*d*x - 2*a**3*b**3*d*x -
4*a**2*b**3*c*d*x + 2*a*b**5*d*x)/(2*b**2*d*(tan(c + d*x)*a**4*b + 2*tan(
c + d*x)*a**2*b**3 + tan(c + d*x)*b**5 + a**5 + 2*a**3*b**2 + a*b**4))
```

3.34 $\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{(a+b \tan(c+dx))^2} dx$

Optimal result	392
Mathematica [C] (verified)	392
Rubi [A] (verified)	393
Maple [A] (verified)	395
Fricas [A] (verification not implemented)	396
Sympy [C] (verification not implemented)	396
Maxima [A] (verification not implemented)	397
Giac [A] (verification not implemented)	398
Mupad [B] (verification not implemented)	398
Reduce [B] (verification not implemented)	399

Optimal result

Integrand size = 32, antiderivative size = 115

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{(2abB - a^2C + b^2C) x}{(a^2 + b^2)^2} - \frac{(a^2B - b^2B + 2abC) \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2)^2 d}$$

$$+ \frac{a(bB - aC)}{b(a^2 + b^2) d(a + b \tan(c + dx))}$$

output

```
(2*B*a*b-C*a^2+C*b^2)*x/(a^2+b^2)^2-(B*a^2-B*b^2+2*C*a*b)*ln(a*cos(d*x+c)+
b*sin(d*x+c))/(a^2+b^2)^2/d+a*(B*b-C*a)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.46 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.22

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{(B+iC) \log(i - \tan(c+dx))}{(a+ib)^2} + \frac{(B-iC) \log(i + \tan(c+dx))}{(a-ib)^2} + \frac{2 \left((-a^2B + b^2B - 2abC) \log(a + b \tan(c+dx)) - \frac{a(a^2+b^2)(-bB+aC)}{b(a+b \tan(c+dx))} \right)}{(a^2+b^2)^2}$$

$2d$

input `Integrate[(B*Tan[c + d*x] + C*Tan[c + d*x]^2)/(a + b*Tan[c + d*x])^2,x]`

output
$$\frac{((B + I*C)*\text{Log}[I - \text{Tan}[c + d*x]])/(a + I*b)^2 + ((B - I*C)*\text{Log}[I + \text{Tan}[c + d*x]])/(a - I*b)^2 + (2*((-(a^2*B) + b^2*B - 2*a*b*C)*\text{Log}[a + b*\text{Tan}[c + d*x]] - (a*(a^2 + b^2)*(-(b*B) + a*C))/(b*(a + b*\text{Tan}[c + d*x]))))/(a^2 + b^2)^2)/(2*d)}$$

Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4111, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{B \tan(c + dx) + C \tan(c + dx)^2}{(a + b \tan(c + dx))^2} dx \\ & \quad \downarrow \text{4111} \\ & \frac{\int \frac{bB - aC + (aB + bC) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a(bB - aC)}{bd(a^2 + b^2)(a + b \tan(c + dx))} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{bB - aC + (aB + bC) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a(bB - aC)}{bd(a^2 + b^2)(a + b \tan(c + dx))} \\ & \quad \downarrow \text{4014} \\ & \frac{x(a^2(-C) + 2abB + b^2C)}{a^2 + b^2} - \frac{(a^2B + 2abC - b^2B) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a(bB - aC)}{bd(a^2 + b^2)(a + b \tan(c + dx))} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\frac{\frac{x(a^2(-C)+2abB+b^2C)}{a^2+b^2} - \frac{(a^2B+2abC-b^2B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2}}{a^2+b^2} + \frac{a(bB-aC)}{bd(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 4013

$$\frac{\frac{a(bB-aC)}{bd(a^2+b^2)(a+b \tan(c+dx))} + \frac{\frac{x(a^2(-C)+2abB+b^2C)}{a^2+b^2} - \frac{(a^2B+2abC-b^2B) \log(a \cos(c+dx)+b \sin(c+dx))}{d(a^2+b^2)}}{a^2+b^2}}$$

input `Int[(B*Tan[c + d*x] + C*Tan[c + d*x]^2)/(a + b*Tan[c + d*x])^2,x]`

output `((((2*a*b*B - a^2*C + b^2*C)*x)/(a^2 + b^2) - ((a^2*B - b^2*B + 2*a*b*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) + (a*(b*B - a*C))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)])*(x_), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.26

method	result
derivativedivides	$\frac{\frac{a(Bb-Ca)}{(a^2+b^2)b(a+b \tan(dx+c))} - \frac{(B a^2 - B b^2 + 2Cab) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2} + \frac{(B a^2 - B b^2 + 2Cab) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)^2} + (2Bab - C a^2 + C b^2)}{d}$
default	$\frac{\frac{a(Bb-Ca)}{(a^2+b^2)b(a+b \tan(dx+c))} - \frac{(B a^2 - B b^2 + 2Cab) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2} + \frac{(B a^2 - B b^2 + 2Cab) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)^2} + (2Bab - C a^2 + C b^2)}{d}$
norman	$\frac{\frac{a(2Bab - C a^2 + C b^2)x}{a^4 + 2b^2 a^2 + b^4} + \frac{b(2Bab - C a^2 + C b^2)x \tan(dx+c)}{a^4 + 2b^2 a^2 + b^4} + \frac{a(Bb-Ca)}{(a^2+b^2)db}}{a+b \tan(dx+c)} + \frac{(B a^2 - B b^2 + 2Cab) \ln(1+\tan(dx+c)^2)}{2d(a^4 + 2b^2 a^2 + b^4)} - \dots$
parallelrisch	$\frac{-2C a^2 b^2 - 2C a^4 + 2B a^3 b + 2C x a b^3 d + 2B a b^3 - B \ln(1+\tan(dx+c)^2) a b^3 - 2B \ln(a+b \tan(dx+c)) a^3 b + 4B x \tan(dx+c) a^3 b}{d(a^4 + 2b^2 a^2 + b^4)}$
risch	$\frac{i x B}{2iab - a^2 + b^2} + \frac{x C}{2iab - a^2 + b^2} + \frac{2ia^2 B x}{a^4 + 2b^2 a^2 + b^4} - \frac{2i B b^2 x}{a^4 + 2b^2 a^2 + b^4} + \frac{4i C a b x}{a^4 + 2b^2 a^2 + b^4} + \frac{2ia^2 B c}{(a^4 + 2b^2 a^2 + b^4)d} - \frac{1}{d(a^4 + 2b^2 a^2 + b^4)}$

input

```
int((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBO
SE)
```

output

```
1/d*(a*(B*b-C*a)/(a^2+b^2)/b/(a+b*tan(d*x+c))-(B*a^2-B*b^2+2*C*a*b)/(a^2+b
^2)^2*ln(a+b*tan(d*x+c))+1/(a^2+b^2)^2*(1/2*(B*a^2-B*b^2+2*C*a*b)*ln(1+tan
(d*x+c)^2)+(2*B*a*b-C*a^2+C*b^2)*arctan(tan(d*x+c))))
```


Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.92

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{2Ca^2b - 2Bab^2 + 2(Ca^3 - 2Ba^2b - Cab^2)dx + (Ba^3 + 2Ca^2b - Bab^2 + (Ba^2b + 2Cab^2 - Bb^3) \tan(c + dx))}{2((a^4b + 2a^2b^3 + b^5)d \tan(c + dx) + (a^5 + 2a^3b^2 + ab^4)d)}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output `-1/2*(2*C*a^2*b - 2*B*a*b^2 + 2*(C*a^3 - 2*B*a^2*b - C*a*b^2)*d*x + (B*a^3 + 2*C*a^2*b - B*a*b^2 + (B*a^2*b + 2*C*a*b^2 - B*b^3)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 2*(C*a^3 - B*a^2*b - (C*a^2*b - 2*B*a*b^2 - C*b^3)*d*x)*tan(d*x + c)/((a^4*b + 2*a^2*b^3 + b^5)*d*tan(d*x + c) + (a^5 + 2*a^3*b^2 + a*b^4)*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.81 (sec) , antiderivative size = 2995, normalized size of antiderivative = 26.04

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**2,x)`

output

```
Piecewise((zoo*x*(B*tan(c) + C*tan(c)**2)/tan(c)**2, Eq(a, 0) & Eq(b, 0) &
Eq(d, 0)), ((B*log(tan(c + d*x)**2 + 1)/(2*d) - C*x + C*tan(c + d*x)/d)/a
**2, Eq(b, 0)), (I*B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b
**2*d*tan(c + d*x) - 4*b**2*d) + 2*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*
x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - I*B*d*x/(4*b**2*d*tan(c + d*
x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*B*tan(c + d*x)/(4*b**2*d*t
an(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + C*d*x*tan(c + d*x)*
*2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*I*C
*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*
b**2*d) - C*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b*
**2*d) - 3*C*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*
x) - 4*b**2*d) + 2*I*C/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x)
- 4*b**2*d), Eq(a, -I*b)), (-I*B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*
x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*B*d*x*tan(c + d*x)/(4*b**2
*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*B*d*x/(4*b**2
*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - I*B*tan(c + d*x
)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + C*d*x*
tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b*
**2*d) + 2*I*C*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(
c + d*x) - 4*b**2*d) - C*d*x/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.61

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx =$$

$$\frac{\frac{2(Ca^2 - 2Bab - Cb^2)(dx+c)}{a^4 + 2a^2b^2 + b^4} + \frac{2(Ba^2 + 2Cab - Bb^2) \log(b \tan(dx+c) + a)}{a^4 + 2a^2b^2 + b^4} - \frac{(Ba^2 + 2Cab - Bb^2) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} + \frac{2(C}{a^3b + ab^3 + (a^2b^2 + b^4) \tan(dx+c)}}{2d}$$

input

```
integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, algorithm="m
axima")
```

output

```
-1/2*(2*(C*a^2 - 2*B*a*b - C*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + 2*(B
*a^2 + 2*C*a*b - B*b^2)*log(b*tan(d*x + c) + a)/(a^4 + 2*a^2*b^2 + b^4) -
(B*a^2 + 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4)
+ 2*(C*a^2 - B*a*b)/(a^3*b + a*b^3 + (a^2*b^2 + b^4)*tan(d*x + c)))/d
```

Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.85

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= -\frac{(Ca^2 - 2 Bab - Cb^2)(dx + c)}{a^4d + 2 a^2b^2d + b^4d} + \frac{(Ba^2 + 2 Cab - Bb^2) \log(\tan(dx + c)^2 + 1)}{2(a^4d + 2 a^2b^2d + b^4d)}$$

$$- \frac{(Ba^2b + 2 Cab^2 - Bb^3) \log(|b \tan(dx + c) + a|)}{a^4bd + 2 a^2b^3d + b^5d}$$

$$- \frac{Ca^4 - Ba^3b + Ca^2b^2 - Bab^3}{(a^2 + b^2)^2(b \tan(dx + c) + a)bd}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `-(C*a^2 - 2*B*a*b - C*b^2)*(d*x + c)/(a^4*d + 2*a^2*b^2*d + b^4*d) + 1/2*(B*a^2 + 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4*d + 2*a^2*b^2*d + b^4*d) - (B*a^2*b + 2*C*a*b^2 - B*b^3)*log(abs(b*tan(d*x + c) + a))/(a^4*b*d + 2*a^2*b^3*d + b^5*d) - (C*a^4 - B*a^3*b + C*a^2*b^2 - B*a*b^3)/((a^2 + b^2)^2*(b*tan(d*x + c) + a)*b*d)`

Mupad [B] (verification not implemented)

Time = 5.45 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.42

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{a(Bb - Ca)}{bd(a^2 + b^2)(a + b \tan(c + dx))}$$

$$+ \frac{\ln(\tan(c + dx) - i)(B + Ci)}{2d(a^2 + ab2i - b^2)}$$

$$+ \frac{\ln(\tan(c + dx) + i)(C + Bi)}{2d(a^2 + 2ab - b^2i)}$$

$$- \frac{\ln(a + b \tan(c + dx)) \left(\frac{B}{a^2 + b^2} - \frac{2b(Bb - Ca)}{(a^2 + b^2)^2} \right)}{d}$$

input `int((B*tan(c + d*x) + C*tan(c + d*x)^2)/(a + b*tan(c + d*x))^2,x)`

output

```
(log(tan(c + d*x) - 1i)*(B + C*1i))/(2*d*(a*b*2i + a^2 - b^2)) - (log(a +
b*tan(c + d*x))*(B/(a^2 + b^2) - (2*b*(B*b - C*a))/(a^2 + b^2)^2))/d + (lo
g(tan(c + d*x) + 1i)*(B*1i + C))/(2*d*(2*a*b + a^2*1i - b^2*1i)) + (a*(B*b
- C*a))/(b*d*(a^2 + b^2)*(a + b*tan(c + d*x)))
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 424, normalized size of antiderivative = 3.69

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\log(\tan(dx + c)^2 + 1) \tan(dx + c) a^2 b^2 + 2 \log(\tan(dx + c)^2 + 1) \tan(dx + c) a b^2 c - \log(\tan(dx + c))}{(a + b \tan(c + dx))^2}$$

input

```
int((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x)
```

output

```
(log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**2 + 2*log(tan(c + d*x)**2 +
1)*tan(c + d*x)*a*b**2*c - log(tan(c + d*x)**2 + 1)*tan(c + d*x)*b**4 + 1
og(tan(c + d*x)**2 + 1)*a**3*b + 2*log(tan(c + d*x)**2 + 1)*a**2*b*c - log
(tan(c + d*x)**2 + 1)*a*b**3 - 2*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**2
*b**2 - 4*log(tan(c + d*x)*b + a)*tan(c + d*x)*a*b**2*c + 2*log(tan(c + d*
x)*b + a)*tan(c + d*x)*b**4 - 2*log(tan(c + d*x)*b + a)*a**3*b - 4*log(tan
(c + d*x)*b + a)*a**2*b*c + 2*log(tan(c + d*x)*b + a)*a*b**3 + 2*tan(c + d
*x)*a**3*c - 2*tan(c + d*x)*a**2*b**2 - 2*tan(c + d*x)*a**2*b*c*d*x + 4*ta
n(c + d*x)*a*b**3*d*x + 2*tan(c + d*x)*a*b**2*c - 2*tan(c + d*x)*b**4 + 2*
tan(c + d*x)*b**3*c*d*x - 2*a**3*c*d*x + 4*a**2*b**2*d*x + 2*a*b**2*c*d*x)
/(2*d*(tan(c + d*x)*a**4*b + 2*tan(c + d*x)*a**2*b**3 + tan(c + d*x)*b**5
+ a**5 + 2*a**3*b**2 + a*b**4))
```

3.35
$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

Optimal result	400
Mathematica [C] (verified)	400
Rubi [A] (verified)	401
Maple [A] (verified)	403
Fricas [A] (verification not implemented)	404
Sympy [C] (verification not implemented)	404
Maxima [A] (verification not implemented)	405
Giac [A] (verification not implemented)	406
Mupad [B] (verification not implemented)	406
Reduce [B] (verification not implemented)	407

Optimal result

Integrand size = 38, antiderivative size = 111

$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(a^2B - b^2B + 2abC)x}{(a^2 + b^2)^2} + \frac{(2abB - a^2C + b^2C) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^2 d}$$

$$- \frac{bB - aC}{(a^2 + b^2) d(a + b \tan(c+dx))}$$

output

```
(B*a^2-B*b^2+2*C*a*b)*x/(a^2+b^2)^2+(2*B*a*b-C*a^2+C*b^2)*ln(a*cos(d*x+c)+
b*sin(d*x+c))/(a^2+b^2)^2/d-(B*b-C*a)/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.62 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.71

$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{C((-ia-b) \log(i-\tan(c+dx))+i(a+ib) \log(i+\tan(c+dx))+2b \log(a+b \tan(c+dx)))}{a^2+b^2} - (bB - aC) \left(\frac{i \log(i-\tan(c+dx))}{(a+ib)^2} - \frac{i \log(i+\tan(c+dx))}{(a-ib)^2} \right)$$

$2bd$

input `Integrate[(Cot[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]`

output `((C*(((-I)*a - b)*Log[I - Tan[c + d*x]] + I*(a + I*b)*Log[I + Tan[c + d*x]] + 2*b*Log[a + b*Tan[c + d*x]]))/(a^2 + b^2) - (b*B - a*C)*((I*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b*(-2*a*Log[a + b*Tan[c + d*x]] + (a^2 + b^2)/(a + b*Tan[c + d*x])))/(a^2 + b^2)^2))/(2*b*d)`

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4115, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{\tan(c + dx) (a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \frac{B + C \tan(c + dx)}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B + C \tan(c + dx)}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{aB + bC - (bB - aC) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{bB - aC}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\int \frac{aB+bC-(bB-aC)\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} - \frac{bB-aC}{d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 4014 \\
& \frac{\frac{(a^2(-C)+2abB+b^2C) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{x(a^2B+2abC-b^2B)}{a^2+b^2}}{a^2+b^2} - \frac{bB-aC}{d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{\frac{(a^2(-C)+2abB+b^2C) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{x(a^2B+2abC-b^2B)}{a^2+b^2}}{a^2+b^2} - \frac{bB-aC}{d(a^2+b^2)(a+b\tan(c+dx))} \\
& \quad \downarrow 4013 \\
& \frac{\frac{(a^2(-C)+2abB+b^2C) \log(a\cos(c+dx)+b\sin(c+dx))}{d(a^2+b^2)} + \frac{x(a^2B+2abC-b^2B)}{a^2+b^2}}{a^2+b^2} - \frac{bB-aC}{d(a^2+b^2)(a+b\tan(c+dx))}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]`

output `((a^2*B - b^2*B + 2*a*b*C)*x)/(a^2 + b^2) + ((2*a*b*B - a^2*C + b^2*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d)/(a^2 + b^2) - (b*B - a*C)/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4013

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4014

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

rule 4115

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e
_) + (f_)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.27

method	result
derivativedivides	$-\frac{Bb-Ca}{(a^2+b^2)(a+b \tan(dx+c))} + \frac{(2Bab-Ca^2+Cb^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2} + \frac{(-2Bab+Ca^2-Cb^2) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)^2} + \frac{(Ba^2-Bb^2)}{(a^2+b^2)^2}$
default	$-\frac{Bb-Ca}{(a^2+b^2)(a+b \tan(dx+c))} + \frac{(2Bab-Ca^2+Cb^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2} + \frac{(-2Bab+Ca^2-Cb^2) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)^2} + \frac{(Ba^2-Bb^2)}{(a^2+b^2)^2}$
parallelrisc	$\frac{2(a+b \tan(dx+c))a(Bab-\frac{1}{2}Ca^2+\frac{1}{2}Cb^2) \ln(a+b \tan(dx+c)) - (a+b \tan(dx+c))a(Bab-\frac{1}{2}Ca^2+\frac{1}{2}Cb^2) \ln(\sec(dx+c))}{(a+b \tan(dx+c))da(a^2+b^2)}$
norman	$\frac{\frac{a(Ba^2-Bb^2+2Cab)x}{a^4+2b^2a^2+b^4} + \frac{b(Ba^2-Bb^2+2Cab)x \tan(dx+c)}{a^4+2b^2a^2+b^4} + \frac{(Bb-Ca)b \tan(dx+c)}{ad(a^2+b^2)}}{a+b \tan(dx+c)} + \frac{(2Bab-Ca^2+Cb^2) \ln(a+b \tan(dx+c))}{d(a^4+2b^2a^2+b^4)}$
risc	$-\frac{xB}{2iab-a^2+b^2} + \frac{ixC}{2iab-a^2+b^2} - \frac{4iabBx}{a^4+2b^2a^2+b^4} + \frac{2ia^2Cx}{a^4+2b^2a^2+b^4} - \frac{2icb^2x}{a^4+2b^2a^2+b^4} - \frac{4iabBc}{d(a^4+2b^2a^2+b^4)} + \frac{d}{d(a^4+2b^2a^2+b^4)}$

input `int(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(-\frac{(Bb - Ca)}{(a^2 + b^2)} \frac{1}{(a + b \tan(dx + c))} + \frac{(2Ba^2b - Ca^2 + Cb^2)}{(a^2 + b^2)^2} \ln(a + b \tan(dx + c)) + \frac{1}{(a^2 + b^2)^2} \left(\frac{1}{2} (-2Ba^2b + Ca^2 - Cb^2) \ln(1 + \tan(dx + c)^2) + (Ba^2 - Bb^2 + 2Ca^2b) \arctan(\tan(dx + c)) \right) \right)$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.00

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{2Cab^2 - 2Bb^3 + 2(Ba^3 + 2Ca^2b - Bab^2)dx - (Ca^3 - 2Ba^2b - Cab^2 + (Ca^2b - 2Bab^2 - Cb^3) \tan(dx))}{2((a^4b + 2a^2b^3 + b^5)d \tan(dx))}$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output
$$\frac{1}{2} \left((2Ca^2b^2 - 2Bb^3 + 2(Ba^3 + 2Ca^2b - Bb^2)) dx - (Ca^3 - 2Ba^2b - Cb^3) \tan(dx + c) \right) \log\left(\frac{b^2 \tan(dx + c)^2 + 2ab \tan(dx + c) + a^2}{\tan(dx + c)^2 + 1}\right) - \frac{2(Ca^2b - Bb^2 - (Ba^2b + 2Ca^2b - Bb^3) dx) \tan(dx + c)}{(a^4b + 2a^2b^3 + b^5) d \tan(dx + c) + (a^5 + 2a^3b^2 + ab^4) d}$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.99 (sec) , antiderivative size = 2895, normalized size of antiderivative = 26.08

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**2,x)`

Giac [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.85

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(Ba^2 + 2Cab - Bb^2)(dx+c)}{a^4d + 2a^2b^2d + b^4d} + \frac{(Ca^2 - 2Bab - Cb^2) \log(\tan(dx+c)^2 + 1)}{2(a^4d + 2a^2b^2d + b^4d)}$$

$$- \frac{(Ca^2b - 2Bab^2 - Cb^3) \log(|b \tan(dx+c) + a|)}{a^4bd + 2a^2b^3d + b^5d} + \frac{Ca^3 - Ba^2b + Cab^2 - Bb^3}{(a^2 + b^2)^2(b \tan(dx+c) + a)d}$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `(B*a^2 + 2*C*a*b - B*b^2)*(d*x + c)/(a^4*d + 2*a^2*b^2*d + b^4*d) + 1/2*(C*a^2 - 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/(a^4*d + 2*a^2*b^2*d + b^4*d) - (C*a^2*b - 2*B*a*b^2 - C*b^3)*log(abs(b*tan(d*x + c) + a))/(a^4*b*d + 2*a^2*b^3*d + b^5*d) + (C*a^3 - B*a^2*b + C*a*b^2 - B*b^3)/((a^2 + b^2)^2*(b*tan(d*x + c) + a)*d)`

Mupad [B] (verification not implemented)

Time = 5.49 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.38

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\ln(a+b \tan(c+dx))(-Ca^2 + 2Bab + Cb^2)}{d(a^2 + b^2)^2} - \frac{Bb - Ca}{d(a^2 + b^2)(a+b \tan(c+dx))}$$

$$- \frac{\ln(\tan(c+dx) + 1i)(C + B1i)}{2d(-a^2 + ab2i + b^2)} - \frac{\ln(\tan(c+dx) - 1i)(B + C1i)}{2d(-a^2 1i + 2ab + b^2 1i)}$$

input `int((cot(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^2,x)`

output

```
(log(a + b*tan(c + d*x))*(C*b^2 - C*a^2 + 2*B*a*b))/(d*(a^2 + b^2)^2) - (B
*b - C*a)/(d*(a^2 + b^2)*(a + b*tan(c + d*x))) - (log(tan(c + d*x) + 1i)*(
B*1i + C))/(2*d*(a*b*2i - a^2 + b^2)) - (log(tan(c + d*x) - 1i)*(B + C*1i)
)/(2*d*(2*a*b - a^2*1i + b^2*1i))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 623, normalized size of antiderivative = 5.61

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input

```
int(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x)
```

output

```
(cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*a**3*c - 2*cos(c + d*x)*log(tan
((c + d*x)/2)**2 + 1)*a**2*b**2 - cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1
)*a*b**2*c - cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b
- a)*a**3*c + 2*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/
2)*b - a)*a**2*b**2 + cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c +
d*x)/2)*b - a)*a*b**2*c + cos(c + d*x)*a**3*b*d*x + cos(c + d*x)*a**3*c -
cos(c + d*x)*a**2*b**2 + 2*cos(c + d*x)*a**2*b*c*d*x - cos(c + d*x)*a*b**3
*d*x + cos(c + d*x)*a*b**2*c - cos(c + d*x)*b**4 + log(tan((c + d*x)/2)**2
+ 1)*sin(c + d*x)*a**2*b*c - 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*
a*b**3 - log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*b**3*c - log(tan((c + d
*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**2*b*c + 2*log(tan(
(c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a*b**3 + log(ta
n((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*b**3*c + sin(
c + d*x)*a**2*b**2*d*x + 2*sin(c + d*x)*a*b**2*c*d*x - sin(c + d*x)*b**4*d
*x)/(d*(cos(c + d*x)*a**5 + 2*cos(c + d*x)*a**3*b**2 + cos(c + d*x)*a*b**4
+ sin(c + d*x)*a**4*b + 2*sin(c + d*x)*a**2*b**3 + sin(c + d*x)*b**5))
```

3.36
$$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

Optimal result	408
Mathematica [C] (verified)	409
Rubi [A] (verified)	409
Maple [A] (verified)	413
Fricas [B] (verification not implemented)	413
Sympy [C] (verification not implemented)	414
Maxima [A] (verification not implemented)	415
Giac [A] (verification not implemented)	416
Mupad [B] (verification not implemented)	417
Reduce [B] (verification not implemented)	417

Optimal result

Integrand size = 40, antiderivative size = 137

$$\begin{aligned} & \int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= -\frac{(2abB - a^2C + b^2C)x}{(a^2 + b^2)^2} + \frac{B \log(\sin(c+dx))}{a^2d} \\ & \quad - \frac{b(3a^2bB + b^3B - 2a^3C) \log(a \cos(c+dx) + b \sin(c+dx))}{a^2(a^2 + b^2)^2d} \\ & \quad + \frac{b(bB - aC)}{a(a^2 + b^2)d(a + b \tan(c+dx))} \end{aligned}$$

output

```
-(2*B*a*b-C*a^2+C*b^2)*x/(a^2+b^2)^2+B*ln(sin(d*x+c))/a^2/d-b*(3*B*a^2*b+B
*b^3-2*C*a^3)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^2/(a^2+b^2)^2/d+b*(B*b-C*a)/
a/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.63 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.16

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx =$$

$$\frac{\frac{(B+iC) \log(i-\tan(c+dx))}{(a+ib)^2} - \frac{2B \log(\tan(c+dx))}{a^2} + \frac{(B-iC) \log(i+\tan(c+dx))}{(a-ib)^2} + \frac{2b(3a^2bB+b^3B-2a^3C) \log(a+b \tan(c+dx))}{a^2(a^2+b^2)^2} + \frac{1}{a(a^2+b^2)}}{2d}$$

input

```
Integrate[(Cot[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]
```

output

```
-1/2*((B + I*C)*Log[I - Tan[c + d*x]]/(a + I*b)^2 - (2*B*Log[Tan[c + d*x]])/a^2 + ((B - I*C)*Log[I + Tan[c + d*x]]/(a - I*b)^2 + (2*b*(3*a^2*b*B + b^3*B - 2*a^3*C)*Log[a + b*Tan[c + d*x]]/(a^2*(a^2 + b^2)^2) + (2*b*(-(b*B) + a*C))/(a*(a^2 + b^2)*(a + b*Tan[c + d*x])))/d
```

Rubi [A] (verified)

Time = 1.58 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.18, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4115, 3042, 4092, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{\tan(c+dx)^2(a+b \tan(c+dx))^2} dx$$

$$\downarrow \text{4115}$$

$$\int \frac{\cot(c+dx)(B + C \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{B + C \tan(c + dx)}{\tan(c + dx)(a + b \tan(c + dx))^2} dx \\
& \downarrow 4092 \\
& \frac{\int \frac{\cot(c+dx)(b(bB-aC) \tan^2(c+dx) - a(bB-aC) \tan(c+dx) + (a^2+b^2)B)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(bB-aC)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{\int \frac{b(bB-aC) \tan(c+dx)^2 - a(bB-aC) \tan(c+dx) + (a^2+b^2)B}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a(a^2+b^2)} + \frac{b(bB-aC)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 4134 \\
& \frac{\frac{B(a^2+b^2) \int \cot(c+dx) dx}{a} - \frac{b(-2a^3C+3a^2bB+b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{ax(a^2(-C)+2abB+b^2C)}{a^2+b^2}}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \\
& \frac{b(bB-aC)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{\frac{B(a^2+b^2) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b(-2a^3C+3a^2bB+b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{ax(a^2(-C)+2abB+b^2C)}{a^2+b^2}}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \\
& \frac{b(bB-aC)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 25 \\
& \frac{\frac{B(a^2+b^2) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} - \frac{b(-2a^3C+3a^2bB+b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{ax(a^2(-C)+2abB+b^2C)}{a^2+b^2}}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \\
& \frac{b(bB-aC)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 3956 \\
& \frac{-\frac{b(-2a^3C+3a^2bB+b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{B(a^2+b^2) \log(-\sin(c+dx))}{ad} - \frac{ax(a^2(-C)+2abB+b^2C)}{a^2+b^2}}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \\
& \frac{b(bB-aC)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 4013
\end{aligned}$$

$$\frac{\frac{B(a^2+b^2) \log(-\sin(c+dx))}{ad} - \frac{ax(a^2(-C)+2abB+b^2C)}{a^2+b^2} - \frac{b(bB-aC)}{ad(a^2+b^2)} \log(a \cos(c+dx)+b \sin(c+dx))}{a(a^2+b^2)} + \frac{ad(a^2+b^2)(a+b \tan(c+dx))}{a(a^2+b^2)}$$

input `Int[(Cot[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]`

output `((-(a*(2*a*b*B - a^2*C + b^2*C)*x)/(a^2 + b^2)) + ((a^2 + b^2)*B*Log[-Sin[c + d*x]])/(a*d) - (b*(3*a^2*b*B + b^3*B - 2*a^3*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(b*B - a*C))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4092

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2
)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e
_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4134

```

Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]

```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19

method	result
derivativedivides	$-\frac{b(3Ba^2b+Bb^3-2Ca^3)\ln(a+b\tan(dx+c))}{(a^2+b^2)^2a^2} + \frac{(Bb-Ca)b}{(a^2+b^2)a(a+b\tan(dx+c))} + \frac{(-Ba^2+Bb^2-2Cab)\ln(1+\tan(dx+c)^2)}{2d(a^2+b^2)^2} + \frac{(-2Bab)}{(a^2+b^2)^2}$
default	$-\frac{b(3Ba^2b+Bb^3-2Ca^3)\ln(a+b\tan(dx+c))}{(a^2+b^2)^2a^2} + \frac{(Bb-Ca)b}{(a^2+b^2)a(a+b\tan(dx+c))} + \frac{(-Ba^2+Bb^2-2Cab)\ln(1+\tan(dx+c)^2)}{2d(a^2+b^2)^2} + \frac{(-2Bab)}{(a^2+b^2)^2}$
parallelrisch	$-6(Ba^2b+\frac{1}{3}Bb^3-\frac{2}{3}Ca^3)(a+b\tan(dx+c))b\ln(a+b\tan(dx+c))-a^2(a+b\tan(dx+c))(Ba^2-Bb^2+2Cab)\ln(\sec(dx+c))$
norman	$-\frac{a(2Bab-Ca^2+Cb^2)x\tan(dx+c)}{a^4+2b^2a^2+b^4} - \frac{b(2Bab-Ca^2+Cb^2)x\tan(dx+c)^2}{a^4+2b^2a^2+b^4} - \frac{(Bb^2-Cab)b\tan(dx+c)^2}{da^2(a^2+b^2)} + \frac{B\ln(\tan(dx+c))}{a^2d}$
risch	$\frac{2ib^3B}{(-ia+b)da(ia+b)^2(b e^{2i(dx+c)}+ia e^{2i(dx+c)}-b+ia)} - \frac{x C}{2iab-a^2+b^2} + \frac{6iBb^2x}{a^4+2b^2a^2+b^4} - \frac{ixB}{2iab-a^2+b^2} + \frac{2}{(a^4+2b^2a^2+b^4)}$

input `int (cot (d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,method =_RETURNVERBOSE)`

output
$$\frac{1}{d} * (-b * (3 * B * a^2 * b + B * b^3 - 2 * C * a^3) / (a^2 + b^2)^2 / a^2 * \ln(a + b * \tan(d * x + c)) + (B * b - C * a) * b / (a^2 + b^2) / a / (a + b * \tan(d * x + c)) + 1 / (a^2 + b^2)^2 * (1 / 2 * (-B * a^2 + B * b^2 - 2 * C * a * b) * \ln(1 + \tan(d * x + c)^2) + (-2 * B * a * b + C * a^2 - C * b^2) * \arctan(\tan(d * x + c)))) + 1 / a^2 * B * \ln(\tan(d * x + c))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 323 vs. 2(137) = 274.

Time = 0.12 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.36

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx =$$

$$2Ca^2b^3 - 2Bab^4 - 2(Ca^5 - 2Ba^4b - Ca^3b^2)dx - (Ba^5 + 2Ba^3b^2 + Bab^4 + (Ba^4b + 2Ba^2b^3 + Bb^5))$$

input `integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="fricas")`

output `-1/2*(2*C*a^2*b^3 - 2*B*a*b^4 - 2*(C*a^5 - 2*B*a^4*b - C*a^3*b^2)*d*x - (B
*a^5 + 2*B*a^3*b^2 + B*a*b^4 + (B*a^4*b + 2*B*a^2*b^3 + B*b^5)*tan(d*x + c
))*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - (2*C*a^4*b - 3*B*a^3*b^2 - B
*a*b^4 + (2*C*a^3*b^2 - 3*B*a^2*b^3 - B*b^5)*tan(d*x + c))*log((b^2*tan(d*
x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 2*(C*a^3*b^2
- B*a^2*b^3 + (C*a^4*b - 2*B*a^3*b^2 - C*a^2*b^3)*d*x)*tan(d*x + c))/((a^6
*b + 2*a^4*b^3 + a^2*b^5)*d*tan(d*x + c) + (a^7 + 2*a^5*b^2 + a^3*b^4)*d)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.56 (sec) , antiderivative size = 4502, normalized size of antiderivative = 32.86

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**2
,x)`

output

```
1/2*(2*(C*a^2 - 2*B*a*b - C*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + 2*(2*
C*a^3*b - 3*B*a^2*b^2 - B*b^4)*log(b*tan(d*x + c) + a)/(a^6 + 2*a^4*b^2 +
a^2*b^4) - (B*a^2 + 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*
b^2 + b^4) - 2*(C*a*b - B*b^2)/(a^4 + a^2*b^2 + (a^3*b + a*b^3)*tan(d*x +
c)) + 2*B*log(tan(d*x + c))/a^2)/d
```

Giac [A] (verification not implemented)

Time = 0.74 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.74

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{(Ca^2 - 2 Bab - Cb^2)(dx + c)}{a^4d + 2a^2b^2d + b^4d} - \frac{(Ba^2 + 2 Cab - Bb^2) \log(\tan(dx + c)^2 + 1)}{2(a^4d + 2a^2b^2d + b^4d)}$$

$$+ \frac{(2Ca^3b^2 - 3Ba^2b^3 - Bb^5) \log(|b \tan(dx + c) + a|)}{a^6bd + 2a^4b^3d + a^2b^5d}$$

$$+ \frac{B \log(|\tan(dx + c)|)}{a^2d} - \frac{Ca^4b - Ba^3b^2 + Ca^2b^3 - Bab^4}{(a^2 + b^2)^2(b \tan(dx + c) + a)a^2d}$$

input

```
integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="giac")
```

output

```
(C*a^2 - 2*B*a*b - C*b^2)*(d*x + c)/(a^4*d + 2*a^2*b^2*d + b^4*d) - 1/2*(B
*a^2 + 2*C*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4*d + 2*a^2*b^2*d + b^4
*d) + (2*C*a^3*b^2 - 3*B*a^2*b^3 - B*b^5)*log(abs(b*tan(d*x + c) + a))/(a^
6*b*d + 2*a^4*b^3*d + a^2*b^5*d) + B*log(abs(tan(d*x + c)))/(a^2*d) - (C*a
^4*b - B*a^3*b^2 + C*a^2*b^3 - B*a*b^4)/((a^2 + b^2)^2*(b*tan(d*x + c) + a
)*a^2*d)
```

Mupad [B] (verification not implemented)

Time = 7.33 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.31

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{B \ln(\tan(c + dx))}{a^2 d} - \frac{\ln(\tan(c + dx) - i) (B + C i)}{2 d (a^2 + a b 2i - b^2)}$$

$$- \frac{\ln(\tan(c + dx) + i) (C + B i)}{2 d (a^2 i + 2 a b - b^2 i)} + \frac{B b^2 - C a b}{a d (a^2 + b^2) (a + b \tan(c + dx))}$$

$$- \frac{b \ln(a + b \tan(c + dx)) (-2 C a^3 + 3 B a^2 b + B b^3)}{a^2 d (a^2 + b^2)^2}$$

input `int((cot(c + d*x))^2*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^2,x)`

output `(B*log(tan(c + d*x)))/(a^2*d) - (log(tan(c + d*x) - 1i)*(B + C*1i))/(2*d*(a*b*2i + a^2 - b^2)) - (log(tan(c + d*x) + 1i)*(B*1i + C))/(2*d*(2*a*b + a^2*1i - b^2*1i)) + (B*b^2 - C*a*b)/(a*d*(a^2 + b^2)*(a + b*tan(c + d*x))) - (b*log(a + b*tan(c + d*x))*(B*b^3 - 2*C*a^3 + 3*B*a^2*b))/(a^2*d*(a^2 + b^2)^2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 782, normalized size of antiderivative = 5.71

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `int(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x)`

output

```
( - cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*a**5*b - 2*cos(c + d*x)*log(
tan((c + d*x)/2)**2 + 1)*a**4*b*c + cos(c + d*x)*log(tan((c + d*x)/2)**2 +
1)*a**3*b**3 + 2*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)
/2)*b - a)*a**4*b*c - 3*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c
+ d*x)/2)*b - a)*a**3*b**3 - cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*ta
n((c + d*x)/2)*b - a)*a*b**5 + cos(c + d*x)*log(tan((c + d*x)/2))*a**5*b +
2*cos(c + d*x)*log(tan((c + d*x)/2))*a**3*b**3 + cos(c + d*x)*log(tan((c
+ d*x)/2))*a*b**5 + cos(c + d*x)*a**5*c*d*x - 2*cos(c + d*x)*a**4*b**2*d*x
- cos(c + d*x)*a**4*b*c + cos(c + d*x)*a**3*b**3 - cos(c + d*x)*a**3*b**2
*c*d*x - cos(c + d*x)*a**2*b**3*c + cos(c + d*x)*a*b**5 - log(tan((c + d*x)
)/2)**2 + 1)*sin(c + d*x)*a**4*b**2 - 2*log(tan((c + d*x)/2)**2 + 1)*sin(c
+ d*x)*a**3*b**2*c + log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**2*b**4
+ 2*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**
3*b**2*c - 3*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c +
d*x)*a**2*b**4 - log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*si
n(c + d*x)*b**6 + log(tan((c + d*x)/2))*sin(c + d*x)*a**4*b**2 + 2*log(tan
((c + d*x)/2))*sin(c + d*x)*a**2*b**4 + log(tan((c + d*x)/2))*sin(c + d*x)
*b**6 + sin(c + d*x)*a**4*b*c*d*x - 2*sin(c + d*x)*a**3*b**3*d*x - sin(c +
d*x)*a**2*b**3*c*d*x)/(a**2*d*(cos(c + d*x)*a**5 + 2*cos(c + d*x)*a**3*b*
**2 + cos(c + d*x)*a*b**4 + sin(c + d*x)*a**4*b + 2*sin(c + d*x)*a**2*b...
```

3.37
$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

Optimal result	419
Mathematica [C] (verified)	420
Rubi [A] (verified)	420
Maple [A] (verified)	424
Fricas [B] (verification not implemented)	425
Sympy [C] (verification not implemented)	426
Maxima [A] (verification not implemented)	427
Giac [A] (verification not implemented)	427
Mupad [B] (verification not implemented)	428
Reduce [B] (verification not implemented)	429

Optimal result

Integrand size = 40, antiderivative size = 192

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= -\frac{(a^2B - b^2B + 2abC)x}{(a^2 + b^2)^2} - \frac{(2bB - aC) \log(\sin(c+dx))}{a^3d}$$

$$+ \frac{b^2(4a^2bB + 2b^3B - 3a^3C - ab^2C) \log(a \cos(c+dx) + b \sin(c+dx))}{a^3(a^2 + b^2)^2d}$$

$$- \frac{b(a^2B + 2b^2B - abC)}{a^2(a^2 + b^2)d(a + b \tan(c+dx))} - \frac{B \cot(c+dx)}{ad(a + b \tan(c+dx))}$$

output

```
-(B*a^2-B*b^2+2*C*a*b)*x/(a^2+b^2)^2-(2*B*b-C*a)*ln(sin(d*x+c))/a^3/d+b^2*(4*B*a^2*b+2*B*b^3-3*C*a^3-C*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^3/(a^2+b^2)^2/d-b*(B*a^2+2*B*b^2-C*a*b)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))-B*cot(d*x+c)/a/d/(a+b*tan(d*x+c))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.41 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.01

$$\int \frac{\cot^3(c+dx) (B \tan(c+dx) + C \tan^2(c+dx))}{(a + b \tan(c+dx))^2} dx$$

$$= \frac{-\frac{2B \cot(c+dx)}{a^2} + \frac{i(B+iC) \log(i-\tan(c+dx))}{(a+ib)^2} + \frac{2(-2bB+aC) \log(\tan(c+dx))}{a^3} - \frac{(iB+C) \log(i+\tan(c+dx))}{(a-ib)^2} - \frac{2b^2(-4a^2bB-2b^3B+3a^2C)}{a^3}}{2d}$$

input

```
Integrate[(Cot[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]
```

output

```
((-2*B*Cot[c + d*x])/a^2 + (I*(B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)^2 + (2*(-2*b*B + a*C)*Log[Tan[c + d*x]])/a^3 - ((I*B + C)*Log[I + Tan[c + d*x]])/(a - I*b)^2 - (2*b^2*(-4*a^2*b*B - 2*b^3*B + 3*a^3*C + a*b^2*C)*Log[a + b*Tan[c + d*x]])/(a^3*(a^2 + b^2)^2) + (2*b^2*(-(b*B) + a*C))/(a^2*(a^2 + b^2)*(a + b*Tan[c + d*x])))/(2*d)
```

Rubi [A] (verified)

Time = 2.25 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.16, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 4115, 3042, 4092, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx) (B \tan(c+dx) + C \tan^2(c+dx))}{(a + b \tan(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{\tan^3(c+dx)(a + b \tan(c+dx))^2} dx$$

$$\downarrow \text{4115}$$

$$\begin{aligned}
 & \int \frac{\cot^2(c+dx)(B+C \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B+C \tan(c+dx)}{\tan(c+dx)^2(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4092} \\
 & - \frac{\int \frac{\cot(c+dx)(2bB \tan^2(c+dx)+aB \tan(c+dx)+2bB-aC)}{(a+b \tan(c+dx))^2} dx}{a} - \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{\int \frac{2bB \tan(c+dx)^2+aB \tan(c+dx)+2bB-aC}{\tan(c+dx)(a+b \tan(c+dx))^2} dx}{a} - \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{4132} \\
 & - \frac{\int \frac{\cot(c+dx)((aB+bC) \tan(c+dx)a^2+b(Ba^2-bCa+2b^2B) \tan^2(c+dx)+(a^2+b^2)(2bB-aC))}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{\int \frac{(aB+bC) \tan(c+dx)a^2+b(Ba^2-bCa+2b^2B) \tan(c+dx)^2+(a^2+b^2)(2bB-aC)}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a(a^2+b^2)} + \frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{4134} \\
 & - \frac{(\frac{a^2+b^2}{a})(2bB-aC) \int \cot(c+dx) dx}{a(a^2+b^2)} - \frac{b^2(-3a^3C+4a^2bB-ab^2C+2b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2x(a^2B+2abC-b^2B)}{a^2+b^2} + \frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{\frac{(a^2+b^2)(2bB-aC) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx}{a} - \frac{b^2(-3a^3C+4a^2bB-ab^2C+2b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2x(a^2B+2abC-b^2B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \qquad \qquad \qquad \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \qquad \qquad \qquad \downarrow 25 \\
 & \frac{\frac{(a^2+b^2)(2bB-aC) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} - \frac{b^2(-3a^3C+4a^2bB-ab^2C+2b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2x(a^2B+2abC-b^2B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \qquad \qquad \qquad \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \qquad \qquad \qquad \downarrow 3956 \\
 & \frac{\frac{b^2(-3a^3C+4a^2bB-ab^2C+2b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2+b^2)(2bB-aC) \log(-\sin(c+dx))}{ad} + \frac{a^2x(a^2B+2abC-b^2B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \qquad \qquad \qquad \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \qquad \qquad \qquad \downarrow 4013 \\
 & \frac{\frac{b(a^2B-abC+2b^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{\frac{(a^2+b^2)(2bB-aC) \log(-\sin(c+dx))}{ad} + \frac{a^2x(a^2B+2abC-b^2B)}{a^2+b^2} - \frac{b^2(-3a^3C+4a^2bB-ab^2C+2b^3B) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)}}{a(a^2+b^2)}}{a(a^2+b^2)} \\
 & \qquad \qquad \qquad \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))}
 \end{aligned}$$

input

```
Int[(Cot[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^2,x]
```

output

```
-((B*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x]))) - (((a^2*(a^2*B - b^2*B + 2*a*b*C)*x)/(a^2 + b^2) + ((a^2 + b^2)*(2*b*B - a*C)*Log[-Sin[c + d*x]])/(a*d) - (b^2*(4*a^2*b*B + 2*b^3*B - 3*a^3*C - a*b^2*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(a^2*B + 2*b^2*B - a*b*C))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/a
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_-), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 3042 $\text{Int}[\text{u}_-, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 3956 $\text{Int}[\tan[(\text{c}_-) + (\text{d}_-)(\text{x}_-)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d} * \text{x}], \text{x}]]/\text{d}, \text{x}] \text{ /; FreeQ}[\{\text{c}, \text{d}\}, \text{x}]$
- rule 4013 $\text{Int}[\frac{(\text{c}_-) + (\text{d}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)]}{(\text{a}_-) + (\text{b}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)]}, \text{x_Symbol}] \rightarrow \text{Simp}[\frac{\text{c}/(\text{b} * \text{f}) * \text{Log}[\text{RemoveContent}[\text{a} * \text{Cos}[\text{e} + \text{f} * \text{x}] + \text{b} * \text{Sin}[\text{e} + \text{f} * \text{x}], \text{x}]]}{\text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{a} * \text{c} + \text{b} * \text{d}, 0]$
- rule 4092 $\text{Int}[\frac{((\text{a}_-) + (\text{b}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)])^{\text{m}_-} * ((\text{A}_-) + (\text{B}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)]) * ((\text{c}_-) + (\text{d}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)])^{\text{n}_-}}{(\text{f}_-)(\text{x}_-)}}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{b} * (\text{A} * \text{b} - \text{a} * \text{B}) * (\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} * ((\text{c} + \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{n} + 1}) / (\text{f} * (\text{m} + 1) * (\text{b} * \text{c} - \text{a} * \text{d}) * (\text{a}^2 + \text{b}^2)), \text{x}] + \text{Simp}[1 / ((\text{m} + 1) * (\text{b} * \text{c} - \text{a} * \text{d}) * (\text{a}^2 + \text{b}^2)) \text{ Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} * (\text{c} + \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{n}} * \text{Simp}[\text{b} * \text{B} * (\text{b} * \text{c} * (\text{m} + 1) + \text{a} * \text{d} * (\text{n} + 1)) + \text{A} * (\text{a} * (\text{b} * \text{c} - \text{a} * \text{d}) * (\text{m} + 1) - \text{b}^2 * \text{d} * (\text{m} + \text{n} + 2)) - (\text{A} * \text{b} - \text{a} * \text{B}) * (\text{b} * \text{c} - \text{a} * \text{d}) * (\text{m} + 1) * \text{Tan}[\text{e} + \text{f} * \text{x}] - \text{b} * \text{d} * (\text{A} * \text{b} - \text{a} * \text{B}) * (\text{m} + \text{n} + 2) * \text{Tan}[\text{e} + \text{f} * \text{x}]^2, \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ (\text{IntegerQ}[\text{m}] \ \|\ \text{IntegersQ}[2 * \text{m}, 2 * \text{n}]) \ \&\& \ !(\text{ILtQ}[\text{n}, -1] \ \&\& \ (!\text{IntegerQ}[\text{m}] \ \|\ (\text{EqQ}[\text{c}, 0] \ \&\& \ \text{NeQ}[\text{a}, 0])))$
- rule 4115 $\text{Int}[\frac{((\text{a}_-) + (\text{b}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)])^{\text{m}_-} * ((\text{c}_-) + (\text{d}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)])^{\text{n}_-} * ((\text{A}_-) + (\text{B}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)] + (\text{C}_-)\tan[(\text{e}_-) + (\text{f}_-)(\text{x}_-)]^2}{(\text{f}_-)(\text{x}_-)}}, \text{x_Symbol}] \rightarrow \text{Simp}[1/\text{b}^2 \text{ Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} * (\text{c} + \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{n}} * (\text{b} * \text{B} - \text{a} * \text{C} + \text{b} * \text{C} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}, \text{m}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \ \&\& \ \text{EqQ}[\text{A} * \text{b}^2 - \text{a} * \text{b} * \text{B} + \text{a}^2 * \text{C}, 0]$

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{b^2(4B a^2 b + 2B b^3 - 3C a^3 - C a b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2 a^3} - \frac{(Bb-Ca)b^2}{(a^2+b^2)a^2(a+b \tan(dx+c))} + \frac{(2Bab-C a^2+C b^2) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)} + \frac{d}{(a^2+b^2)}$
default	$\frac{b^2(4B a^2 b + 2B b^3 - 3C a^3 - C a b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^2 a^3} - \frac{(Bb-Ca)b^2}{(a^2+b^2)a^2(a+b \tan(dx+c))} + \frac{(2Bab-C a^2+C b^2) \ln(1+\tan(dx+c)^2)}{2(a^2+b^2)} + \frac{d}{(a^2+b^2)}$
parallelrisc	$4(a+b \tan(dx+c))b^2(B a^2 b + \frac{1}{2}B b^3 - \frac{3}{4}C a^3 - \frac{1}{4}C a b^2) \ln(a+b \tan(dx+c)) + a^3(Bab - \frac{1}{2}C a^2 + \frac{1}{2}C b^2)(a+b \tan(dx+c))$
norman	$\frac{(B a^2 b + 2B b^3 - C a b^2) b \tan(dx+c)^3}{d a^3 (a^2+b^2)} - \frac{B \tan(dx+c)}{ad} - \frac{a(B a^2 - B b^2 + 2C ab) x \tan(dx+c)^2}{a^4 + 2b^2 a^2 + b^4} - \frac{b(B a^2 - B b^2 + 2C ab) x \tan(dx+c)^3}{a^4 + 2b^2 a^2 + b^4} + \frac{\tan(dx+c)^2(a+b \tan(dx+c))}{\tan(dx+c)^2(a+b \tan(dx+c))}$
risc	$\frac{x B}{2iab - a^2 + b^2} - \frac{i x C}{2iab - a^2 + b^2} + \frac{2ib^4 C x}{a^2(a^4 + 2b^2 a^2 + b^4)} - \frac{2i(-2iB a^3 b e^{2i(dx+c)} - 2iB a b^3 e^{2i(dx+c)} + B a^4 e^{2i(dx+c)} - 2iB a^3 b e^{2i(dx+c)} - 2iB a b^3 e^{2i(dx+c)} + B a^4 e^{2i(dx+c)} - 2iB a^3 b e^{2i(dx+c)} - 2iB a b^3 e^{2i(dx+c)} + B a^4 e^{2i(dx+c)})}{(e^{2i(dx+c)} - 1)(ib+a)(-ib+a)^2}$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 6.01 (sec) , antiderivative size = 8143, normalized size of antiderivative = 42.41

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)**3*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**2, x)`

output

```
Piecewise((( -B*x - B/(d*tan(c + d*x)) - C*log(tan(c + d*x)**2 + 1)/(2*d) +
C*log(tan(c + d*x))/d)/a**2, Eq(b, 0)), ((B*x + B/(d*tan(c + d*x)) - B/(3
*d*tan(c + d*x)**3) + C*log(tan(c + d*x)**2 + 1)/(2*d) - C*log(tan(c + d*x)
))/d - C/(2*d*tan(c + d*x)**2))/b**2, Eq(a, 0)), (-9*B*d*x*tan(c + d*x)**3
/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c +
d*x)) - 18*I*B*d*x*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d
*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) + 9*B*d*x*tan(c + d*x)/(4*a**2*d
*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 4
*I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**3 +
8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) + 8*B*log(tan(c + d*x)
**2 + 1)*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*
x)**2 - 4*a**2*d*tan(c + d*x)) + 4*I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*
x)/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c
+ d*x)) + 8*I*B*log(tan(c + d*x))*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**
3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 16*B*log(tan(c
+ d*x))*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)
**2 - 4*a**2*d*tan(c + d*x)) - 8*I*B*log(tan(c + d*x))*tan(c + d*x)/(4*a*
**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x))
- 9*B*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)
**2 - 4*a**2*d*tan(c + d*x)) - 14*I*B*tan(c + d*x)/(4*a**2*d*tan(c + d*...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.36

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx =$$

$$\frac{2(Ba^2+2Cab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} + \frac{2(3Ca^3b^2-4Ba^2b^3+Cab^4-2Bb^5) \log(b \tan(dx+c)+a)}{a^7+2a^5b^2+a^3b^4} + \frac{(Ca^2-2Bab-Cb^2) \log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} + \frac{2d}{2d}$$

input

```
integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="maxima")
```

output

```
-1/2*(2*(B*a^2 + 2*C*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + 2*(3
*C*a^3*b^2 - 4*B*a^2*b^3 + C*a*b^4 - 2*B*b^5)*log(b*tan(d*x + c) + a)/(a^7
+ 2*a^5*b^2 + a^3*b^4) + (C*a^2 - 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1
)/(a^4 + 2*a^2*b^2 + b^4) + 2*(B*a^3 + B*a*b^2 + (B*a^2*b - C*a*b^2 + 2*B*
b^3)*tan(d*x + c))/((a^4*b + a^2*b^3)*tan(d*x + c)^2 + (a^5 + a^3*b^2)*tan
(d*x + c)) - 2*(C*a - 2*B*b)*log(tan(d*x + c))/a^3)/d
```

Giac [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.54

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(Ba^2+2Cab-Bb^2)(dx+c)}{a^4d+2a^2b^2d+b^4d} - \frac{(Ca^2-2Bab-Cb^2) \log(\tan(dx+c)^2+1)}{2(a^4d+2a^2b^2d+b^4d)}$$

$$- \frac{(3Ca^3b^3-4Ba^2b^4+Cab^5-2Bb^6) \log(|b \tan(dx+c)+a|)}{a^7bd+2a^5b^3d+a^3b^5d}$$

$$+ \frac{(Ca-2Bb) \log(|\tan(dx+c)|)}{a^3d}$$

$$- \frac{Ba^5+2Ba^3b^2+Bab^4+(Ba^4b-Ca^3b^2+3Ba^2b^3-Cab^4+2Bb^5) \tan(dx+c)}{(a^2+b^2)^2(b \tan(dx+c)+a)a^2d \tan(dx+c)}$$

input

```
integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x,
algorithm="giac")
```


output

```

-(B*a^2 + 2*C*a*b - B*b^2)*(d*x + c)/(a^4*d + 2*a^2*b^2*d + b^4*d) - 1/2*(
C*a^2 - 2*B*a*b - C*b^2)*log(tan(d*x + c)^2 + 1)/(a^4*d + 2*a^2*b^2*d + b^
4*d) - (3*C*a^3*b^3 - 4*B*a^2*b^4 + C*a*b^5 - 2*B*b^6)*log(abs(b*tan(d*x +
c) + a))/(a^7*b*d + 2*a^5*b^3*d + a^3*b^5*d) + (C*a - 2*B*b)*log(abs(tan(
d*x + c)))/(a^3*d) - (B*a^5 + 2*B*a^3*b^2 + B*a*b^4 + (B*a^4*b - C*a^3*b^2
+ 3*B*a^2*b^3 - C*a*b^4 + 2*B*b^5)*tan(d*x + c))/((a^2 + b^2)^2*(b*tan(d*
x + c) + a)*a^2*d*tan(d*x + c))

```

Mupad [B] (verification not implemented)

Time = 8.60 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.20

$$\begin{aligned}
& \int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx \\
&= \frac{b^2 \ln(a + b \tan(c + dx)) (-3 C a^3 + 4 B a^2 b - C a b^2 + 2 B b^3)}{a^3 d (a^2 + b^2)^2} \\
&\quad - \frac{\ln(\tan(c + dx)) (2 B b - C a)}{a^3 d} + \frac{\ln(\tan(c + dx) + 1i) (C + B 1i)}{2 d (-a^2 + a b 2i + b^2)} \\
&\quad + \frac{\ln(\tan(c + dx) - 1i) (B + C 1i)}{2 d (-a^2 1i + 2 a b + b^2 1i)} - \frac{\frac{B}{a} + \frac{\tan(c+dx) (B a^2 b - C a b^2 + 2 B b^3)}{a^2 (a^2 + b^2)}}{d (b \tan(c + dx)^2 + a \tan(c + dx))}
\end{aligned}$$

input

```

int((cot(c + d*x)^3*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*
x))^2,x)

```

output

```

(log(tan(c + d*x) + 1i)*(B*1i + C))/(2*d*(a*b*2i - a^2 + b^2)) - (log(tan(
c + d*x))*(2*B*b - C*a))/(a^3*d) - (B/a + (tan(c + d*x)*(2*B*b^3 + B*a^2*b
- C*a*b^2))/(a^2*(a^2 + b^2)))/(d*(a*tan(c + d*x) + b*tan(c + d*x)^2)) +
(log(tan(c + d*x) - 1i)*(B + C*1i))/(2*d*(2*a*b - a^2*1i + b^2*1i)) + (b^2
*log(a + b*tan(c + d*x))*(2*B*b^3 - 3*C*a^3 + 4*B*a^2*b - C*a*b^2))/(a^3*d
*(a^2 + b^2)^2)

```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1293, normalized size of antiderivative = 6.73

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input

```
int(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^2,x)
```

output

```
( - 2*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**6*c + 4*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**5*b**2 + 2*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**4*b**2*c - 6*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**4*b**2*c + 8*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**3*b**4 - 2*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**2*b**4*c + 4*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a*b**6 + 2*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**6*c - 4*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**5*b**2 + 4*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**4*b**2*c - 8*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**3*b**4 + 2*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**2*b**4*c - 4*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a*b**6 - cos(c + d*x)*sin(c + d*x)*a**7 - 2*cos(c + d*x)*sin(c + d*x)*a**6*b*d*x - 4*cos(c + d*x)*sin(c + d*x)*a**5*b**2 - 4*cos(c + d*x)*sin(c + d*x)*a**5*b*c*d*x + 2*cos(c + d*x)*sin(c + d*x)*a**4*b**3*d*x + 2*cos(c + d*x)*sin(c + d*x)*a**4*b**2*c - 7*cos(c + d*x)*sin(c + d*x)*a**3*b**4 + 2*cos(c + d*x)*sin(c + d*x)*a**2*b**4*c - 4*cos(c + d*x)*sin(c + d*x)*a*b**6 - 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**5*b*c + 4*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**4*b**3 + 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*...
```

3.38 $\int \frac{\tan^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$

Optimal result	430
Mathematica [C] (verified)	431
Rubi [A] (verified)	431
Maple [A] (verified)	437
Fricas [B] (verification not implemented)	438
Sympy [F(-2)]	439
Maxima [A] (verification not implemented)	440
Giac [A] (verification not implemented)	440
Mupad [B] (verification not implemented)	441
Reduce [B] (verification not implemented)	442

Optimal result

Integrand size = 40, antiderivative size = 331

$$\int \frac{\tan^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(a^3B - 3ab^2B + 3a^2bC - b^3C)x}{(a^2 + b^2)^3} + \frac{(3a^2bB - b^3B - a^3C + 3ab^2C) \log(\cos(c+dx))}{(a^2 + b^2)^3 d}$$

$$+ \frac{a^2(a^4bB + 3a^2b^3B + 6b^5B - 3a^5C - 9a^3b^2C - 10ab^4C) \log(a+b \tan(c+dx))}{b^4(a^2 + b^2)^3 d}$$

$$- \frac{(a^3bB + 3ab^3B - 3a^4C - 6a^2b^2C - b^4C) \tan(c+dx)}{b^3(a^2 + b^2)^2 d}$$

$$+ \frac{a(bB - aC) \tan^3(c+dx)}{2b(a^2 + b^2) d(a+b \tan(c+dx))^2}$$

$$+ \frac{a(a^2bB + 5b^3B - 3a^3C - 7ab^2C) \tan^2(c+dx)}{2b^2(a^2 + b^2)^2 d(a+b \tan(c+dx))}$$

output

```
(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*x/(a^2+b^2)^3+(3*B*a^2*b-B*b^3-C*a^3+3*C
*a*b^2)*ln(cos(d*x+c))/(a^2+b^2)^3/d+a^2*(B*a^4*b+3*B*a^2*b^3+6*B*b^5-3*C*
a^5-9*C*a^3*b^2-10*C*a*b^4)*ln(a+b*tan(d*x+c))/b^4/(a^2+b^2)^3/d-(B*a^3*b+
3*B*a*b^3-3*C*a^4-6*C*a^2*b^2-C*b^4)*tan(d*x+c)/b^3/(a^2+b^2)^2/d+1/2*a*(B
*b-C*a)*tan(d*x+c)^3/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+1/2*a*(B*a^2*b+5*B*b
^3-3*C*a^3-7*C*a*b^2)*tan(d*x+c)^2/b^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.21 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.83

$$\int \frac{\tan^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{\frac{(B+iC) \log(i-\tan(c+dx))}{(-ia+b)^3} + \frac{(B-iC) \log(i+\tan(c+dx))}{(ia+b)^3} + \frac{2a^2(a^4bB+3a^2b^3B+6b^5B-3a^5C-9a^3b^2C-10ab^4C) \log(a+b \tan(c+dx))}{b^4(a^2+b^2)^3}}{2d} + \frac{b}{b}$$

input

```
Integrate[(Tan[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

```
((B + I*C)*Log[I - Tan[c + d*x]]/((-I)*a + b)^3 + (B - I*C)*Log[I + Tan[c + d*x]]/(I*a + b)^3 + (2*a^2*(a^4*b*B + 3*a^2*b^3*B + 6*b^5*B - 3*a^5*C - 9*a^3*b^2*C - 10*a*b^4*C)*Log[a + b*Tan[c + d*x]])/(b^4*(a^2 + b^2)^3) + (a^3*(-(a*b*B) + 3*a^2*C + 2*b^2*C))/(b^4*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) + (2*C*Tan[c + d*x]^3)/(b*(a + b*Tan[c + d*x])^2) - (2*a^2*(-2*a^3*b*B - 4*a*b^3*B + 6*a^4*C + 11*a^2*b^2*C + 3*b^4*C))/(b^4*(a^2 + b^2)^2*(a + b*Tan[c + d*x]))/(2*d)
```

Rubi [A] (verified)

Time = 3.41 (sec) , antiderivative size = 363, normalized size of antiderivative = 1.10, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.425$, Rules used = {3042, 4115, 3042, 4088, 25, 3042, 4128, 27, 3042, 4130, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^3(B \tan(c+dx) + C \tan^2(c+dx)^2)}{(a+b \tan(c+dx))^3} dx$$

$$\begin{aligned}
 & \int \frac{\tan^4(c+dx)(B+C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4115} \\
 & \int \frac{\tan(c+dx)^4(B+C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\frac{\tan^2(c+dx)((-3Ca^2+bBa-2b^2C) \tan^2(c+dx)-2b(bB-aC) \tan(c+dx)+3a(bB-aC))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{\int -\frac{\tan^2(c+dx)((-3Ca^2+bBa-2b^2C) \tan^2(c+dx)-2b(bB-aC) \tan(c+dx)+3a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} + \\
 & \quad \frac{a(bB-aC) \tan^3(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{a(bB-aC) \tan^3(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \frac{\int \frac{\tan^2(c+dx)((-3Ca^2+bBa-2b^2C) \tan^2(c+dx)-2b(bB-aC) \tan(c+dx)+3a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(bB-aC) \tan^3(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \frac{\int \frac{\tan(c+dx)^2((-3Ca^2+bBa-2b^2C) \tan(c+dx)^2-2b(bB-aC) \tan(c+dx)+3a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} \\
 & \quad \downarrow \text{4128} \\
 & \frac{a(bB-aC) \tan^3(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \frac{\int \frac{2 \tan(c+dx)((Ba^2+2bCa-b^2B) \tan(c+dx)b^2+(-3Ca^4+bBa^3-6b^2Ca^2+3b^3Ba-b^4C) \tan^2(c+dx)+a(-3Ca^3+bBa^2-7b^2Ca+5b^3B))}{(a+b \tan(c+dx))} dx}{b(a^2+b^2)} - \frac{a(-3a^3C+)}{bd} \\
 & \quad \downarrow \text{27} \\
 & \frac{a(bB-aC) \tan^3(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \frac{2 \int \frac{\tan(c+dx)((Ba^2+2bCa-b^2B) \tan(c+dx)b^2+(-3Ca^4+bBa^3-6b^2Ca^2+3b^3Ba-b^4C) \tan^2(c+dx)+a(-3Ca^3+bBa^2-7b^2Ca+5b^3B))}{(a+b \tan(c+dx))} dx}{b(a^2+b^2)} - \frac{a(-3a^3C+)}{bd} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$2 \int \frac{\frac{a(bB - aC) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{\tan(c+dx)((Ba^2+2bCa-b^2B) \tan(c+dx)b^2 + (-3Ca^4+bBa^3-6b^2Ca^2+3b^3Ba-b^4C) \tan(c+dx)^2 + a(-3Ca^3+bBa^2-7b^2Ca+5b^3B))}{a+b \tan(c+dx)}}{b(a^2+b^2)} dx - \frac{a(-3a^3C+a^3bB-6a^2b^2C+3ab^3B-b^4C)}{bd}$$

$$2b(a^2 + b^2)$$

↓ 4130

$$2 \left(\frac{\int -\frac{((-Ca^2+2bBa+b^2C) \tan(c+dx)b^3) + (a^2+b^2)^2(bB-3aC) \tan^2(c+dx) + a(-3Ca^4+bBa^3-6b^2Ca^2+3b^3Ba-b^4C)}{a+b \tan(c+dx)} dx}{b} + \frac{(-3a^4C+a^3bB-6a^2b^2C+3ab^3B-b^4C) \tan(c+dx)}{bd} \right)$$

$$2b(a^2 + b^2)$$

↓ 25

$$2 \left(\frac{(-3a^4C+a^3bB-6a^2b^2C+3ab^3B-b^4C) \tan(c+dx)}{bd} - \int \frac{((-Ca^2+2bBa+b^2C) \tan(c+dx)b^3) + (a^2+b^2)^2(bB-3aC) \tan^2(c+dx) + a(-3Ca^4+bBa^3-6b^2Ca^2+3b^3Ba-b^4C)}{a+b \tan(c+dx)} dx}{b} \right)$$

$$2b(a^2 + b^2)$$

↓ 3042

$$2 \left(\frac{(-3a^4C+a^3bB-6a^2b^2C+3ab^3B-b^4C) \tan(c+dx)}{bd} - \int \frac{((-Ca^2+2bBa+b^2C) \tan(c+dx)b^3) + (a^2+b^2)^2(bB-3aC) \tan^2(c+dx) + a(-3Ca^4+bBa^3-6b^2Ca^2+3b^3Ba-b^4C)}{a+b \tan(c+dx)} dx}{b} \right)$$

$$2b(a^2 + b^2)$$

↓ 4109

$$2 \left(\frac{(-3a^4C+a^3bB-6a^2b^2C+3ab^3B-b^4C) \tan(c+dx)}{bd} - \frac{b^3(a^3(-C)+3a^2bB+3ab^2C-b^3B) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^2(-3a^5C+a^4bB-9a^3b^2C+3a^2b^3B-10ab^4C+b^5)}{b(a^2+b^2)} \right)$$

$$2b(a^2 + b^2)$$

↓ 3042

$$2 \left(\frac{(-3a^4C + a^3bB - 6a^2b^2C + 3ab^3B - b^4C) \tan(c+dx)}{bd} - \frac{b^3(a^3(-C) + 3a^2bB + 3ab^2C - b^3B) \int \tan(c+dx) dx}{a^2 + b^2} + \frac{a^2(-3a^5C + a^4bB - 9a^3b^2C + 3a^2b^3B - 10ab^4C + 6b^5B)}{b(a^2 + b^2)} \right) - \frac{a(bB - aC) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 3956

$$2 \left(\frac{(-3a^4C + a^3bB - 6a^2b^2C + 3ab^3B - b^4C) \tan(c+dx)}{bd} - \frac{a^2(-3a^5C + a^4bB - 9a^3b^2C + 3a^2b^3B - 10ab^4C + 6b^5B) \int \frac{\tan(c+dx)^2 + 1}{a + b \tan(c+dx)} dx}{a^2 + b^2} + \frac{b^3(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{b(a^2 + b^2)} \right) - \frac{a(bB - aC) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 4100

$$2 \left(\frac{(-3a^4C + a^3bB - 6a^2b^2C + 3ab^3B - b^4C) \tan(c+dx)}{bd} - \frac{a^2(-3a^5C + a^4bB - 9a^3b^2C + 3a^2b^3B - 10ab^4C + 6b^5B) \int \frac{1}{a + b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2 + b^2)} + \frac{b^3(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{b(a^2 + b^2)} \right) - \frac{a(bB - aC) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 16

$$2 \left(\frac{(-3a^4C + a^3bB - 6a^2b^2C + 3ab^3B - b^4C) \tan(c+dx)}{bd} - \frac{b^3(a^3(-C) + 3a^2bB + 3ab^2C - b^3B) \log(\cos(c+dx))}{d(a^2 + b^2)} + \frac{b^3x(a^3B + 3a^2bC - 3ab^2B - b^3C)}{a^2 + b^2} + \frac{a^2(-3a^5C + a^4bB - 9a^3b^2C + 3a^2b^3B - 10ab^4C + 6b^5B)}{b(a^2 + b^2)} \right) - \frac{a(bB - aC) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}$$

input

```
Int[(Tan[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

$$\begin{aligned} & (a*(b*B - a*C)*\text{Tan}[c + d*x]^3)/(2*b*(a^2 + b^2)*d*(a + b*\text{Tan}[c + d*x])^2) \\ & - (-((a*(a^2*b*B + 5*b^3*B - 3*a^3*C - 7*a*b^2*C)*\text{Tan}[c + d*x]^2)/(b*(a^2 \\ & + b^2)*d*(a + b*\text{Tan}[c + d*x]))) + (2*(-(((b^3*(a^3*B - 3*a*b^2*B + 3*a^2*b \\ & *C - b^3*C)*x)/(a^2 + b^2) + (b^3*(3*a^2*b*B - b^3*B - a^3*C + 3*a*b^2*C)* \\ & \text{Log}[\text{Cos}[c + d*x]])/((a^2 + b^2)*d) + (a^2*(a^4*b*B + 3*a^2*b^3*B + 6*b^5*B \\ & - 3*a^5*C - 9*a^3*b^2*C - 10*a*b^4*C)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(b*(a^2 + \\ & b^2)*d))/b) + ((a^3*b*B + 3*a*b^3*B - 3*a^4*C - 6*a^2*b^2*C - b^4*C)*\text{Tan}[c \\ & + d*x])/(b*d)))/(b*(a^2 + b^2)))/(2*b*(a^2 + b^2)) \end{aligned}$$

Defintions of rubi rules used

rule 16

$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$$

rule 25

$$\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[Fx, x], x]$$

rule 27

$$\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] \text{ ; FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\text{tan}[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4088

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
  Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])

```

rule 4100

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]

```

rule 4109

```

Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2
)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e
_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.79

method	result
derivativedivides	$\frac{\tan(dx+c)C}{b^3} + \frac{a^2(Ba^4b+3Ba^2b^3+6Bb^5-3Ca^5-9Ca^3b^2-10Cab^4)\ln(a+b\tan(dx+c))}{b^4(a^2+b^2)^3} - \frac{a^4(Bb-Ca)}{2b^4(a^2+b^2)(a+b\tan(dx+c))^2} + \frac{a^3(2)}{b^4}$
default	$\frac{\tan(dx+c)C}{b^3} + \frac{a^2(Ba^4b+3Ba^2b^3+6Bb^5-3Ca^5-9Ca^3b^2-10Cab^4)\ln(a+b\tan(dx+c))}{b^4(a^2+b^2)^3} - \frac{a^4(Bb-Ca)}{2b^4(a^2+b^2)(a+b\tan(dx+c))^2} + \frac{a^3(2)}{b^4}$
norman	$\frac{C\tan(dx+c)^3}{db} + \frac{(Ba^3-3Bab^2+3Ca^2b-Cb^3)a^2x}{(a^4+2b^2a^2+b^4)(a^2+b^2)} + \frac{b^2(Ba^3-3Bab^2+3Ca^2b-Cb^3)x\tan(dx+c)^2}{(a^4+2b^2a^2+b^4)(a^2+b^2)} + \frac{a(2Ba^4b+4Ba^2b^3-6Ca^5)}{db^3(a^4+2b^2a^2+b^4)(a+b\tan(dx+c))^2}$
parallelrisch	Expression too large to display
risch	Expression too large to display

input `int(tan(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method
=_RETURNVERBOSE)`

output `1/d*(tan(d*x+c)*C/b^3+1/b^4*a^2*(B*a^4*b+3*B*a^2*b^3+6*B*b^5-3*C*a^5-9*C*a
^3*b^2-10*C*a*b^4)/(a^2+b^2)^3*ln(a+b*tan(d*x+c))-1/2/b^4*a^4*(B*b-C*a)/(a
^2+b^2)/(a+b*tan(d*x+c))^2+1/b^4*a^3*(2*B*a^2*b+4*B*b^3-3*C*a^3-5*C*a*b^2)
/(a^2+b^2)^2/(a+b*tan(d*x+c))+1/(a^2+b^2)^3*(1/2*(-3*B*a^2*b+B*b^3+C*a^3-3
*C*a*b^2)*ln(1+tan(d*x+c)^2)+(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*arctan(tan(
d*x+c))))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 890 vs. $2(328) = 656$.

Time = 0.18 (sec) , antiderivative size = 890, normalized size of antiderivative = 2.69

$$\int \frac{\tan^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="fricas")`

output

```

-1/2*(3*C*a^7*b^2 - B*a^6*b^3 + 9*C*a^5*b^4 - 7*B*a^4*b^5 - 2*(C*a^6*b^3 +
3*C*a^4*b^5 + 3*C*a^2*b^7 + C*b^9)*tan(d*x + c)^3 - 2*(B*a^5*b^4 + 3*C*a^
4*b^5 - 3*B*a^3*b^6 - C*a^2*b^7)*d*x - (9*C*a^7*b^2 - 3*B*a^6*b^3 + 23*C*a^
5*b^4 - 9*B*a^4*b^5 + 12*C*a^3*b^6 + 4*C*a*b^8 + 2*(B*a^3*b^6 + 3*C*a^2*b
^7 - 3*B*a*b^8 - C*b^9)*d*x)*tan(d*x + c)^2 + (3*C*a^9 - B*a^8*b + 9*C*a^7
*b^2 - 3*B*a^6*b^3 + 10*C*a^5*b^4 - 6*B*a^4*b^5 + (3*C*a^7*b^2 - B*a^6*b^3
+ 9*C*a^5*b^4 - 3*B*a^4*b^5 + 10*C*a^3*b^6 - 6*B*a^2*b^7)*tan(d*x + c)^2
+ 2*(3*C*a^8*b - B*a^7*b^2 + 9*C*a^6*b^3 - 3*B*a^5*b^4 + 10*C*a^4*b^5 - 6*
B*a^3*b^6)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^
2)/(tan(d*x + c)^2 + 1)) - (3*C*a^9 - B*a^8*b + 9*C*a^7*b^2 - 3*B*a^6*b^3
+ 9*C*a^5*b^4 - 3*B*a^4*b^5 + 3*C*a^3*b^6 - B*a^2*b^7 + (3*C*a^7*b^2 - B*a^
6*b^3 + 9*C*a^5*b^4 - 3*B*a^4*b^5 + 9*C*a^3*b^6 - 3*B*a^2*b^7 + 3*C*a*b^8
- B*b^9)*tan(d*x + c)^2 + 2*(3*C*a^8*b - B*a^7*b^2 + 9*C*a^6*b^3 - 3*B*a^
5*b^4 + 9*C*a^4*b^5 - 3*B*a^3*b^6 + 3*C*a^2*b^7 - B*a*b^8)*tan(d*x + c))*l
og(1/(tan(d*x + c)^2 + 1)) - 2*(3*C*a^8*b - B*a^7*b^2 + 6*C*a^6*b^3 - 3*B*
a^5*b^4 - 2*C*a^4*b^5 + 4*B*a^3*b^6 + C*a^2*b^7 + 2*(B*a^4*b^5 + 3*C*a^3*b
^6 - 3*B*a^2*b^7 - C*a*b^8)*d*x)*tan(d*x + c))/((a^6*b^6 + 3*a^4*b^8 + 3*a^
2*b^10 + b^12)*d*tan(d*x + c)^2 + 2*(a^7*b^5 + 3*a^5*b^7 + 3*a^3*b^9 + a^
b^11)*d*tan(d*x + c) + (a^8*b^4 + 3*a^6*b^6 + 3*a^4*b^8 + a^2*b^10)*d)

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

= Exception raised: AttributeError

input

```

integrate(tan(d*x+c)**3*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3
,x)

```

output

```

Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'

```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.18

$$\int \frac{\tan^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{2(Ba^3+3Ca^2b-3Bab^2-Cb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(3Ca^7-Ba^6b+9Ca^5b^2-3Ba^4b^3+10Ca^3b^4-6Ba^2b^5) \log(b \tan(dx+c)+a)}{a^6b^4+3a^4b^6+3a^2b^8+b^{10}} + \frac{(Ca^3-3Ba^2b-3Cab^2+Bb^3) \log(\tan(dx+c)^2+1)}{2(a^6+3a^4b^2+3a^2b^4+b^6)} + \frac{C \tan(dx+c)}{b^3d} - \frac{5Ca^9-3Ba^8b+14Ca^7b^2-10Ba^6b^3+9Ca^5b^4-7Ba^4b^5+2(3Ca^8b-2Ba^7b^2+8Ca^6b^3-6Ba^5b^4) \tan(dx+c)}{2(a^2+b^2)^3(b \tan(dx+c)+a)^2b^4d}$$

input

```
integrate(tan(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="maxima")
```

output

```
1/2*(2*(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2
+ 3*a^2*b^4 + b^6) - 2*(3*C*a^7 - B*a^6*b + 9*C*a^5*b^2 - 3*B*a^4*b^3 + 10
*C*a^3*b^4 - 6*B*a^2*b^5)*log(b*tan(d*x + c) + a)/(a^6*b^4 + 3*a^4*b^6 + 3
*a^2*b^8 + b^10) + (C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*log(tan(d*x + c
)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (5*C*a^7 - 3*B*a^6*b + 9*C*
a^5*b^2 - 7*B*a^4*b^3 + 2*(3*C*a^6*b - 2*B*a^5*b^2 + 5*C*a^4*b^3 - 4*B*a^3
*b^4)*tan(d*x + c))/(a^6*b^4 + 2*a^4*b^6 + a^2*b^8 + (a^4*b^6 + 2*a^2*b^8
+ b^10)*tan(d*x + c)^2 + 2*(a^5*b^5 + 2*a^3*b^7 + a*b^9)*tan(d*x + c)) + 2
*C*tan(d*x + c)/b^3)/d
```

Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.16

$$\int \frac{\tan^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)(dx+c)}{a^6d + 3a^4b^2d + 3a^2b^4d + b^6d} + \frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(\tan(dx+c)^2 + 1)}{2(a^6d + 3a^4b^2d + 3a^2b^4d + b^6d)} - \frac{(3Ca^7 - Ba^6b + 9Ca^5b^2 - 3Ba^4b^3 + 10Ca^3b^4 - 6Ba^2b^5) \log(|b \tan(dx+c) + a|)}{a^6b^4d + 3a^4b^6d + 3a^2b^8d + b^{10}d} + \frac{C \tan(dx+c)}{b^3d} - \frac{5Ca^9 - 3Ba^8b + 14Ca^7b^2 - 10Ba^6b^3 + 9Ca^5b^4 - 7Ba^4b^5 + 2(3Ca^8b - 2Ba^7b^2 + 8Ca^6b^3 - 6Ba^5b^4) \tan(dx+c)}{2(a^2 + b^2)^3(b \tan(dx+c) + a)^2b^4d}$$

input `integrate(tan(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="giac")`

output
$$\begin{aligned} & (B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d + 3 \\ & *a^2*b^4*d + b^6*d) + 1/2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*\log(\tan(\\ & d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) - (3*C*a^7 - B \\ & *a^6*b + 9*C*a^5*b^2 - 3*B*a^4*b^3 + 10*C*a^3*b^4 - 6*B*a^2*b^5)*\log(\text{abs}(b \\ & * \tan(d*x + c) + a))/(a^6*b^4*d + 3*a^4*b^6*d + 3*a^2*b^8*d + b^{10}*d) + C*t \\ & \text{an}(d*x + c)/(b^3*d) - 1/2*(5*C*a^9 - 3*B*a^8*b + 14*C*a^7*b^2 - 10*B*a^6*b \\ & ^3 + 9*C*a^5*b^4 - 7*B*a^4*b^5 + 2*(3*C*a^8*b - 2*B*a^7*b^2 + 8*C*a^6*b^3 \\ & - 6*B*a^5*b^4 + 5*C*a^4*b^5 - 4*B*a^3*b^6)*\tan(d*x + c))/((a^2 + b^2)^3*(b \\ & * \tan(d*x + c) + a)^2*b^4*d) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 6.98 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.01

$$\begin{aligned} & \int \frac{\tan^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \frac{C \tan(c + dx)}{b^3 d} \\ & + \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2 d (-a^3 - a^2 b 3i + 3 a b^2 + b^3 i)} + \frac{\ln(\tan(c + dx) + i) (B - C i)}{2 d (-a^3 i - 3 a^2 b + a b^2 3i + b^3)} \\ & - \frac{\frac{5 C a^7 - 3 B a^6 b + 9 C a^5 b^2 - 7 B a^4 b^3}{2 b (a^4 + 2 a^2 b^2 + b^4)} + \frac{\tan(c + dx) (3 C a^6 - 2 B a^5 b + 5 C a^4 b^2 - 4 B a^3 b^3)}{a^4 + 2 a^2 b^2 + b^4}}{d (a^2 b^3 + 2 a b^4 \tan(c + dx) + b^5 \tan(c + dx)^2)} \\ & + \frac{a^2 \ln(a + b \tan(c + dx)) (-3 C a^5 + B a^4 b - 9 C a^3 b^2 + 3 B a^2 b^3 - 10 C a b^4 + 6 B b^5)}{b^4 d (a^2 + b^2)^3} \end{aligned}$$

input `int((tan(c + d*x)^3*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*
x))^3,x)`

output

```
(log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(3*a*b^2 - a^2*b^3i - a^3 + b^3*1i)) - ((5*C*a^7 - 7*B*a^4*b^3 + 9*C*a^5*b^2 - 3*B*a^6*b)/(2*b*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(3*C*a^6 - 4*B*a^3*b^3 + 5*C*a^4*b^2 - 2*B*a^5*b))/(a^4 + b^4 + 2*a^2*b^2))/(d*(a^2*b^3 + b^5*tan(c + d*x)^2 + 2*a*b^4*tan(c + d*x))) + (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3)) + (C*tan(c + d*x))/(b^3*d) + (a^2*log(a + b*tan(c + d*x))*(6*B*b^5 - 3*C*a^5 + 3*B*a^2*b^3 - 9*C*a^3*b^2 + B*a^4*b - 10*C*a*b^4))/(b^4*d*(a^2 + b^2)^3)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1268, normalized size of antiderivative = 3.83

$$\int \frac{\tan^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input

```
int(tan(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)
```

output

```
(log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a**3*b**6*c - 3*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a**2*b**8 - 3*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a*b**8*c + log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*b**10 + 2*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**4*b**5*c - 6*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**3*b**7 - 6*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**7*c + 2*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a*b**9 + log(tan(c + d*x)**2 + 1)*a**5*b**4*c - 3*log(tan(c + d*x)**2 + 1)*a**4*b**6 - 3*log(tan(c + d*x)**2 + 1)*a**3*b**6*c + log(tan(c + d*x)**2 + 1)*a**2*b**8 - 6*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**7*b**2*c + 2*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**6*b**4 - 18*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**5*b**4*c + 6*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**4*b**6 - 20*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**3*b**6*c + 12*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**2*b**8 - 12*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**8*b*c + 4*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**7*b**3 - 36*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**6*b**3*c + 12*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**5*b**5 - 40*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**4*b**5*c + 24*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**3*b**7 - 6*log(tan(c + d*x)*b + a)*a**9*c + 2*log(tan(c + d*x)*b + a)*a**8*b**2 - 18*log(tan(c + d*x)*b + a)*a**7*b**2*c + 6*log(tan(c + d*x)*b + a)*a**6*b**4 - 20*log(tan(c + d*x)*b + a)*a**5*b**4*c + 12*log(tan(c + d*x)*b + a)*a**4*b**6 + 2*tan(c + d*x)**3*a...
```

3.39
$$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

Optimal result	443
Mathematica [C] (verified)	444
Rubi [A] (verified)	444
Maple [A] (verified)	449
Fricas [B] (verification not implemented)	449
Sympy [F(-2)]	450
Maxima [A] (verification not implemented)	451
Giac [A] (verification not implemented)	451
Mupad [B] (verification not implemented)	452
Reduce [B] (verification not implemented)	453

Optimal result

Integrand size = 40, antiderivative size = 250

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(3a^2bB - b^3B - a^3C + 3ab^2C)x}{(a^2 + b^2)^3}$$

$$+ \frac{(a^3B - 3ab^2B + 3a^2bC - b^3C) \log(\cos(c+dx))}{(a^2 + b^2)^3 d}$$

$$+ \frac{a(a^2b^3B - 3b^5B + a^5C + 3a^3b^2C + 6ab^4C) \log(a+b \tan(c+dx))}{b^3(a^2 + b^2)^3 d}$$

$$+ \frac{a(bB - aC) \tan^2(c+dx)}{2b(a^2 + b^2) d(a+b \tan(c+dx))^2} - \frac{a^2(2b^3B - a^3C - 3ab^2C)}{b^3(a^2 + b^2)^2 d(a+b \tan(c+dx))}$$

output

```
- (3*B*a^2*b-B*b^3-C*a^3+3*C*a*b^2)*x/(a^2+b^2)^3+(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*ln(cos(d*x+c))/(a^2+b^2)^3/d+a*(B*a^2*b^3-3*B*b^5+C*a^5+3*C*a^3*b^2+6*C*a*b^4)*ln(a+b*tan(d*x+c))/b^3/(a^2+b^2)^3/d+1/2*a*(B*b-C*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^2-a^2*(2*B*b^3-C*a^3-3*C*a*b^2)/b^3/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.27 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.89

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{-\frac{(B+iC) \log(i-\tan(c+dx))}{(a+ib)^3} - \frac{(B-iC) \log(i+\tan(c+dx))}{(a-ib)^3} + \frac{2a(a^2b^3B-3b^5B+a^5C+3a^3b^2C+6ab^4C) \log(a+b \tan(c+dx))}{b^3(a^2+b^2)^3} + \frac{a^3}{b^3(a^2+b^2)}}{2d}$$

input

```
Integrate[(Tan[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

```
(-(((B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)^3) - ((B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b)^3 + (2*a*(a^2*b^3*B - 3*b^5*B + a^5*C + 3*a^3*b^2*C + 6*a*b^4*C)*Log[a + b*Tan[c + d*x]])/(b^3*(a^2 + b^2)^3) + (a^3*(b*B - a*C))/(b^3*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) + (2*a^2*(-(a^2*b*B) - 3*b^3*B + 2*a^3*C + 4*a*b^2*C))/(b^3*(a^2 + b^2)^2*(a + b*Tan[c + d*x]))/(2*d)
```

Rubi [A] (verified)

Time = 2.38 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.14, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.325$, Rules used = {3042, 4115, 3042, 4088, 27, 3042, 4118, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^2(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow 4115$$

$$\begin{aligned}
& \int \frac{\tan^3(c+dx)(B+C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\tan(c+dx)^3(B+C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow \text{4088} \\
& \frac{\int -\frac{2 \tan(c+dx)((a^2+b^2)C \tan^2(c+dx)-b(bB-aC) \tan(c+dx)+a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} + \\
& \quad \frac{a(bB-aC) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow \text{27} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
& \frac{\int \frac{\tan(c+dx)((a^2+b^2)C \tan^2(c+dx)-b(bB-aC) \tan(c+dx)+a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
& \frac{\int \frac{\tan(c+dx)((a^2+b^2)C \tan^2(c+dx)-b(bB-aC) \tan(c+dx)+a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{4118} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
& \frac{\int \frac{(Ba^2+2bCa-b^2B) \tan(c+dx)b^2-(a^2+b^2)^2 C \tan^2(c+dx)+a(-Ca^3-3b^2Ca+2b^3B)}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \frac{a^2(a^3(-C)-3ab^2C+2b^3B)}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{a(bB-aC) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
& \frac{\int \frac{(Ba^2+2bCa-b^2B) \tan(c+dx)b^2-(a^2+b^2)^2 C \tan^2(c+dx)+a(-Ca^3-3b^2Ca+2b^3B)}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \frac{a^2(a^3(-C)-3ab^2C+2b^3B)}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow \text{4109}
\end{aligned}$$

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{b^2(a^3B + 3a^2bC - 3ab^2B - b^3C) \int \tan(c + dx) dx}{a^2 + b^2} - \frac{a(a^5C + 3a^3b^2C + a^2b^3B + 6ab^4C - 3b^5B) \int \frac{\tan^2(c + dx) + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^2x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2}}{b(a^2 + b^2)} + \frac{a^2}{b^2d}$$

↓ 3042

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{b^2(a^3B + 3a^2bC - 3ab^2B - b^3C) \int \tan(c + dx) dx}{a^2 + b^2} - \frac{a(a^5C + 3a^3b^2C + a^2b^3B + 6ab^4C - 3b^5B) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^2x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2}}{b(a^2 + b^2)} + \frac{a^2}{b^2d}$$

↓ 3956

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(a^5C + 3a^3b^2C + a^2b^3B + 6ab^4C - 3b^5B) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{b^2(a^3B + 3a^2bC - 3ab^2B - b^3C) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^2x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2}}{b(a^2 + b^2)} + \frac{a^2}{b^2d}$$

↓ 4100

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(a^5C + 3a^3b^2C + a^2b^3B + 6ab^4C - 3b^5B) \int \frac{1}{a + b \tan(c + dx)} d(b \tan(c + dx))}{bd(a^2 + b^2)} - \frac{b^2(a^3B + 3a^2bC - 3ab^2B - b^3C) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^2x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2}}{b(a^2 + b^2)}$$

↓ 16

$$\frac{\frac{a(bB - aC) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{b^2(a^3B + 3a^2bC - 3ab^2B - b^3C) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^2x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2} - \frac{a(a^5C + 3a^3b^2C + a^2b^3B + 6ab^4C - 3b^5B)}{bd(a^2 + b^2)}}{\frac{a^2(a^3(-C) - 3ab^2C + 2b^3B)}{b^2d(a^2 + b^2)(a + b \tan(c + dx))} + b(a^2 + b^2)}$$

input

```
Int[(Tan[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

$$\begin{aligned} & (a*(b*B - a*C)*\tan[c + d*x]^2)/(2*b*(a^2 + b^2)*d*(a + b*\tan[c + d*x])^2) \\ & - (((b^2*(3*a^2*b*B - b^3*B - a^3*C + 3*a*b^2*C)*x)/(a^2 + b^2) - (b^2*(a^3*B - 3*a*b^2*B + 3*a^2*b*C - b^3*C)*\log[\cos[c + d*x]])/(a^2 + b^2)*d) - \\ & (a*(a^2*b^3*B - 3*b^5*B + a^5*C + 3*a^3*b^2*C + 6*a*b^4*C)*\log[a + b*\tan[c + d*x]])/(b*(a^2 + b^2)*d))/(b*(a^2 + b^2)) + (a^2*(2*b^3*B - a^3*C - 3*a*b^2*C))/(b^2*(a^2 + b^2)*d*(a + b*\tan[c + d*x]))/(b*(a^2 + b^2)) \end{aligned}$$

Definitions of rubi rules used

rule 16

$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\log[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$$

rule 27

$$\text{Int}[(a_)*(F_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F, (b_)*(G_)] \text{ ; FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\log[\text{RemoveContent}[\cos[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4088

$$\begin{aligned} & \text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x_)]^{(m)}*((A_)+(B_)*\tan[(e_)+(f_)*(x_)]^{(n)}), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*(B*c - A*d)*(a + b*\tan[e + f*x])^{(m-1)}*((c + d*\tan[e + f*x])^{(n+1)})/(d*f*(n+1)*(c^2 + d^2)), x] - \text{Simp}[1/(d*(n+1)*(c^2 + d^2)) \\ & \text{Int}[(a + b*\tan[e + f*x])^{(m-2)}*(c + d*\tan[e + f*x])^{(n+1)}*\text{Simp}[a*A*d*(b*d*(m-1) - a*c*(n+1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m-1) + a*d*(n+1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n+1)*\tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m+n) - b*B*(c^2*(m-1) - d^2*(n+1)))*\tan[e + f*x]^2, x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \\ & \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \& \\ & \ \& \ \text{LtQ}[n, -1] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegersQ}[2*m, 2*n]) \end{aligned}$$

rule 4100

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

rule 4109

```
Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2
)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]
```

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_
.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_
.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f
_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.97

method	result
derivativedivides	$\frac{a(Ba^2b^3 - 3Bb^5 + Ca^5 + 3Ca^3b^2 + 6Ca^2b^4) \ln(a+b \tan(dx+c)) - \frac{a^2(Ba^2b + 3Bb^3 - 2Ca^3 - 4Ca^2b^2)}{b^3(a^2+b^2)^2(a+b \tan(dx+c))} + \frac{a^3(Bb - Ca)}{2b^3(a^2+b^2)(a+b \tan(dx+c))}}{(a^2+b^2)^3 b^3} + \frac{d}{d}$
default	$\frac{a(Ba^2b^3 - 3Bb^5 + Ca^5 + 3Ca^3b^2 + 6Ca^2b^4) \ln(a+b \tan(dx+c)) - \frac{a^2(Ba^2b + 3Bb^3 - 2Ca^3 - 4Ca^2b^2)}{b^3(a^2+b^2)^2(a+b \tan(dx+c))} + \frac{a^3(Bb - Ca)}{2b^3(a^2+b^2)(a+b \tan(dx+c))}}{(a^2+b^2)^3 b^3} + \frac{d}{d}$
norman	$-\frac{a^2(Ba^3b + 5Ba^2b^3 - 3Ca^4 - 7Ca^2b^2)}{2db^3(a^4 + 2b^2a^2 + b^4)} - \frac{(3Ba^2b - Bb^3 - Ca^3 + 3Ca^2b^2)a^2x}{(a^4 + 2b^2a^2 + b^4)(a^2 + b^2)} - \frac{b^2(3Ba^2b - Bb^3 - Ca^3 + 3Ca^2b^2)x \tan(dx+c)^2}{(a^4 + 2b^2a^2 + b^4)(a^2 + b^2)} - \frac{a}{(a+b \tan(dx+c))^2}$
risch	$-\frac{12ia^2bCc}{(a^6 + 3b^2a^4 + 3b^4a^2 + b^6)d} - \frac{x C}{3ia^2b - ib^3 - a^3 + 3ab^2} - \frac{ixB}{3ia^2b - ib^3 - a^3 + 3ab^2} + \frac{6iab^2Bc}{(a^6 + 3b^2a^4 + 3b^4a^2 + b^6)d} - \frac{a}{(a^6 + 3b^2a^4 + 3b^4a^2 + b^6)}$
parallelrisc	Expression too large to display

```
input int (tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method
=_RETURNVERBOSE)
```

```
output 1/d*(a*(B*a^2*b^3-3*B*b^5+C*a^5+3*C*a^3*b^2+6*C*a*b^4)/(a^2+b^2)^3/b^3*ln(
a+b*tan(d*x+c))-a^2*(B*a^2*b+3*B*b^3-2*C*a^3-4*C*a*b^2)/b^3/(a^2+b^2)^2/(a
+b*tan(d*x+c))+1/2*a^3*(B*b-C*a)/b^3/(a^2+b^2)/(a+b*tan(d*x+c))^2+1/(a^2+b
^2)^3*(1/2*(-B*a^3+3*B*a*b^2-3*C*a^2*b+C*b^3)*ln(1+tan(d*x+c)^2)+(-3*B*a^2
*b+B*b^3+C*a^3-3*C*a*b^2)*arctan(tan(d*x+c))))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 666 vs. 2(243) = 486.

Time = 0.14 (sec) , antiderivative size = 666, normalized size of antiderivative = 2.66

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{Ca^6b^2 + Ba^5b^3 + 7Ca^4b^4 - 5Ba^3b^5 + 2(Ca^5b^3 - 3Ba^4b^4 - 3Ca^3b^5 + Ba^2b^6)dx - (3Ca^6b^2 - Ba^5b^3 + \dots}{\dots}$$

input `integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="fricas")`

output `1/2*(C*a^6*b^2 + B*a^5*b^3 + 7*C*a^4*b^4 - 5*B*a^3*b^5 + 2*(C*a^5*b^3 - 3*B*a^4*b^4 - 3*C*a^3*b^5 + B*a^2*b^6)*d*x - (3*C*a^6*b^2 - B*a^5*b^3 + 9*C*a^4*b^4 - 7*B*a^3*b^5 - 2*(C*a^3*b^5 - 3*B*a^2*b^6 - 3*C*a*b^7 + B*b^8)*d*x)*tan(d*x + c)^2 + (C*a^8 + 3*C*a^6*b^2 + B*a^5*b^3 + 6*C*a^4*b^4 - 3*B*a^3*b^5 + (C*a^6*b^2 + 3*C*a^4*b^4 + B*a^3*b^5 + 6*C*a^2*b^6 - 3*B*a*b^7)*tan(d*x + c)^2 + 2*(C*a^7*b + 3*C*a^5*b^3 + B*a^4*b^4 + 6*C*a^3*b^5 - 3*B*a^2*b^6)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (C*a^8 + 3*C*a^6*b^2 + 3*C*a^4*b^4 + C*a^2*b^6 + (C*a^6*b^2 + 3*C*a^4*b^4 + 3*C*a^2*b^6 + C*b^8)*tan(d*x + c)^2 + 2*(C*a^7*b + 3*C*a^5*b^3 + 3*C*a^3*b^5 + C*a*b^7)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 2*(C*a^7*b + 3*C*a^5*b^3 - 3*B*a^4*b^4 - 4*C*a^3*b^5 + 3*B*a^2*b^6 - 2*(C*a^4*b^4 - 3*B*a^3*b^5 - 3*C*a^2*b^6 + B*a*b^7)*d*x)*tan(d*x + c))/((a^6*b^5 + 3*a^4*b^7 + 3*a^2*b^9 + b^11)*d*tan(d*x + c)^2 + 2*(a^7*b^4 + 3*a^5*b^6 + 3*a^3*b^8 + a*b^10)*d*tan(d*x + c) + (a^8*b^3 + 3*a^6*b^5 + 3*a^4*b^7 + a^2*b^9)*d)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

= Exception raised: AttributeError

input `integrate(tan(d*x+c)**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 366, normalized size of antiderivative = 1.46

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{2(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3)(dx + c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{2(Ca^6 + 3Ca^4b^2 + Ba^3b^3 + 6Ca^2b^4 - 3Bab^5) \log(b \tan(dx + c) + a)}{a^6b^3 + 3a^4b^5 + 3a^2b^7 + b^9} - \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} 2d$$

input

```
integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="maxima")
```

output

```
1/2*(2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2
+ 3*a^2*b^4 + b^6) + 2*(C*a^6 + 3*C*a^4*b^2 + B*a^3*b^3 + 6*C*a^2*b^4 - 3*
B*a*b^5)*log(b*tan(d*x + c) + a)/(a^6*b^3 + 3*a^4*b^5 + 3*a^2*b^7 + b^9) -
(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*
a^4*b^2 + 3*a^2*b^4 + b^6) + (3*C*a^6 - B*a^5*b + 7*C*a^4*b^2 - 5*B*a^3*b^
3 + 2*(2*C*a^5*b - B*a^4*b^2 + 4*C*a^3*b^3 - 3*B*a^2*b^4)*tan(d*x + c))/(a
^6*b^3 + 2*a^4*b^5 + a^2*b^7 + (a^4*b^5 + 2*a^2*b^7 + b^9)*tan(d*x + c)^2
+ 2*(a^5*b^4 + 2*a^3*b^6 + a*b^8)*tan(d*x + c)))/d
```

Giac [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.44

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3)(dx + c)}{a^6d + 3a^4b^2d + 3a^2b^4d + b^6d} - \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3) \log(\tan(dx + c)^2 + 1)}{2(a^6d + 3a^4b^2d + 3a^2b^4d + b^6d)} + \frac{(Ca^6 + 3Ca^4b^2 + Ba^3b^3 + 6Ca^2b^4 - 3Bab^5) \log(|b \tan(dx + c) + a|)}{a^6b^3d + 3a^4b^5d + 3a^2b^7d + b^9d} + \frac{2(2Ca^7 - Ba^6b + 6Ca^5b^2 - 4Ba^4b^3 + 4Ca^3b^4 - 3Ba^2b^5) \tan(dx + c) + \frac{3Ca^8 - Ba^7b + 10Ca^6b^2 - 6Ba^5b^3 + 3Ca^4b^4 - 3Ba^3b^5 + 3Ba^2b^6 - 3Ba^1b^7 + 3Bb^8}{b}}{2(a^2 + b^2)^3(b \tan(dx + c) + a)^2b^2d}$$

input `integrate(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="giac")`

output
$$\begin{aligned} & (C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d + 3* \\ & *a^2*b^4*d + b^6*d) - 1/2*(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*\log(\tan(\\ & d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) + (C*a^6 + 3*C \\ & *a^4*b^2 + B*a^3*b^3 + 6*C*a^2*b^4 - 3*B*a*b^5)*\log(\text{abs}(b*\tan(d*x + c) + a \\ &))/(a^6*b^3*d + 3*a^4*b^5*d + 3*a^2*b^7*d + b^9*d) + 1/2*(2*(2*C*a^7 - B*a \\ & ^6*b + 6*C*a^5*b^2 - 4*B*a^4*b^3 + 4*C*a^3*b^4 - 3*B*a^2*b^5)*\tan(d*x + c) \\ & + (3*C*a^8 - B*a^7*b + 10*C*a^6*b^2 - 6*B*a^5*b^3 + 7*C*a^4*b^4 - 5*B*a^3 \\ & *b^5)/b)/((a^2 + b^2)^3*(b*\tan(d*x + c) + a)^2*b^2*d) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 6.02 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.23

$$\begin{aligned} & \int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx \\ & = \frac{\frac{3Ca^6 - Ba^5b + 7Ca^4b^2 - 5Ba^3b^3}{2b^3(a^4 + 2a^2b^2 + b^4)} - \frac{a^2 \tan(c + dx) (-2Ca^3 + Ba^2b - 4Cab^2 + 3Bb^3)}{b^2(a^4 + 2a^2b^2 + b^4)}}{d(a^2 + 2ab \tan(c + dx) + b^2 \tan^2(c + dx))} \\ & + \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2d(-a^3 i + 3a^2b + ab^2 3i - b^3)} + \frac{\ln(\tan(c + dx) + i) (B - C i)}{2d(-a^3 + a^2b 3i + 3ab^2 - b^3 i)} \\ & + \frac{a \ln(a + b \tan(c + dx)) (Ca^5 + 3Ca^3b^2 + Ba^2b^3 + 6Cab^4 - 3Bb^5)}{b^3 d(a^2 + b^2)^3} \end{aligned}$$

input `int((tan(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*
x))^3,x)`

output
$$\begin{aligned} & ((3*C*a^6 - 5*B*a^3*b^3 + 7*C*a^4*b^2 - B*a^5*b)/(2*b^3*(a^4 + b^4 + 2*a^2 \\ & *b^2)) - (a^2*\tan(c + d*x)*(3*B*b^3 - 2*C*a^3 + B*a^2*b - 4*C*a*b^2))/(b^2 \\ & *(a^4 + b^4 + 2*a^2*b^2)))/(d*(a^2 + b^2*\tan(c + d*x)^2 + 2*a*b*\tan(c + d* \\ & x))) + (\log(\tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*b^2*3i + 3*a^2*b - a^3* \\ & 1i - b^3)) + (\log(\tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(3*a*b^2 + a^2*b*3i \\ & - a^3 - b^3*1i)) + (a*\log(a + b*\tan(c + d*x))*(C*a^5 - 3*B*b^5 + B*a^2*b^3 \\ & + 3*C*a^3*b^2 + 6*C*a*b^4))/(b^3*d*(a^2 + b^2)^3) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 1097, normalized size of antiderivative = 4.39

$$\int \frac{\tan^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input

```
int(tan(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)
```

output

```
( - log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a**3*b**6 - 3*log(tan(c + d*x)
)**2 + 1)*tan(c + d*x)**2*a**2*b**6*c + 3*log(tan(c + d*x)**2 + 1)*tan(c +
d*x)**2*a*b**8 + log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*b**8*c - 2*log(
tan(c + d*x)**2 + 1)*tan(c + d*x)*a**4*b**5 - 6*log(tan(c + d*x)**2 + 1)*t
an(c + d*x)*a**3*b**5*c + 6*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**
7 + 2*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a*b**7*c - log(tan(c + d*x)**2
+ 1)*a**5*b**4 - 3*log(tan(c + d*x)**2 + 1)*a**4*b**4*c + 3*log(tan(c + d
*x)**2 + 1)*a**3*b**6 + log(tan(c + d*x)**2 + 1)*a**2*b**6*c + 2*log(tan(c
+ d*x)*b + a)*tan(c + d*x)**2*a**6*b**2*c + 6*log(tan(c + d*x)*b + a)*tan
(c + d*x)**2*a**4*b**4*c + 2*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**3*
b**6 + 12*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a**2*b**6*c - 6*log(tan(
c + d*x)*b + a)*tan(c + d*x)**2*a*b**8 + 4*log(tan(c + d*x)*b + a)*tan(c +
d*x)*a**7*b*c + 12*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**5*b**3*c + 4*l
og(tan(c + d*x)*b + a)*tan(c + d*x)*a**4*b**5 + 24*log(tan(c + d*x)*b + a)
*tan(c + d*x)*a**3*b**5*c - 12*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**2*b
**7 + 2*log(tan(c + d*x)*b + a)*a**8*c + 6*log(tan(c + d*x)*b + a)*a**6*b
**2*c + 2*log(tan(c + d*x)*b + a)*a**5*b**4 + 12*log(tan(c + d*x)*b + a)*a
**4*b**4*c - 6*log(tan(c + d*x)*b + a)*a**3*b**6 - 2*tan(c + d*x)**2*a**6*b
**2*c + tan(c + d*x)**2*a**5*b**4 - 6*tan(c + d*x)**2*a**4*b**4*c + 4*tan(
c + d*x)**2*a**3*b**6 + 2*tan(c + d*x)**2*a**3*b**5*c*d*x - 6*tan(c + d...
```

3.40
$$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

Optimal result	454
Mathematica [C] (verified)	455
Rubi [A] (verified)	455
Maple [A] (verified)	459
Fricas [B] (verification not implemented)	459
Sympy [F(-2)]	460
Maxima [A] (verification not implemented)	460
Giac [A] (verification not implemented)	461
Mupad [B] (verification not implemented)	462
Reduce [B] (verification not implemented)	463

Optimal result

Integrand size = 38, antiderivative size = 189

$$\int \frac{\tan(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(a^3 B - 3ab^2 B + 3a^2 b C - b^3 C) x}{(a^2 + b^2)^3}$$

$$- \frac{(3a^2 b B - b^3 B - a^3 C + 3ab^2 C) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^3 d}$$

$$- \frac{a^2 (b B - a C)}{2b^2 (a^2 + b^2) d (a + b \tan(c+dx))^2} + \frac{a(2b^3 B - a^3 C - 3ab^2 C)}{b^2 (a^2 + b^2)^2 d (a + b \tan(c+dx))}$$

output

```
- (B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*x/(a^2+b^2)^3-(3*B*a^2*b-B*b^3-C*a^3+3*
C*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^3/d-1/2*a^2*(B*b-C*a)/b^2
/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+a*(2*B*b^3-C*a^3-3*C*a*b^2)/b^2/(a^2+b^2)^
2/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.75 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.52

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{bB+aC}{b(a+b \tan(c+dx))^2} - \frac{2C \tan(c+dx)}{(a+b \tan(c+dx))^2} + C \left(\frac{i \log(i-\tan(c+dx))}{(a+ib)^2} - \frac{i \log(i+\tan(c+dx))}{(a-ib)^2} + \frac{2b(-2a \log(a+b \tan(c+dx)) + \frac{a^2+b^2}{a+b \tan(c+dx)})}{(a^2+b^2)^2} \right)$$

input

```
Integrate[(Tan[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

```
(-((b*B + a*C)/(b*(a + b*Tan[c + d*x])^2)) - (2*C*Tan[c + d*x])/(a + b*Tan[c + d*x])^2 + C*((I*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b*(-2*a*Log[a + b*Tan[c + d*x]] + (a^2 + b^2)/(a + b*Tan[c + d*x])))/(a^2 + b^2)^2) + (b*B - a*C)*((I*Log[I - Tan[c + d*x]])/(a + I*b)^3 - Log[I + Tan[c + d*x]]/(I*a + b)^3 + (b*((-6*a^2 + 2*b^2)*Log[a + b*Tan[c + d*x]] + ((a^2 + b^2)*(5*a^2 + b^2 + 4*a*b*Tan[c + d*x])))/(a + b*Tan[c + d*x])^2))/(a^2 + b^2)^3)/(2*b*d)
```

Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4115, 3042, 4087, 25, 3042, 4111, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan(c+dx)^2)}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 4115 \\
& \int \frac{\tan^2(c+dx)(B + C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 3042 \\
& \int \frac{\tan(c+dx)^2(B + C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 4087 \\
& \frac{\int -\frac{((a^2+b^2)C \tan^2(c+dx) - b(bB-aC) \tan(c+dx) + a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} - \frac{a^2(bB-aC)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow 25 \\
& -\frac{\int -\frac{((a^2+b^2)C \tan^2(c+dx) - b(bB-aC) \tan(c+dx) + a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} - \frac{a^2(bB-aC)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& -\frac{\int -\frac{((a^2+b^2)C \tan(c+dx)^2 - b(bB-aC) \tan(c+dx) + a(bB-aC))}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} - \frac{a^2(bB-aC)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow 4111 \\
& -\frac{\int \frac{b(Ba^2+2bCa-b^2B) - b(-Ca^2+2bBa+b^2C) \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{a(a^3(-C)-3ab^2C+2b^3B)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \frac{b(a^2+b^2)}{a^2(bB-aC)} \\
& \quad \frac{a^2(bB-aC)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& -\frac{\int \frac{b(Ba^2+2bCa-b^2B) - b(-Ca^2+2bBa+b^2C) \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{a(a^3(-C)-3ab^2C+2b^3B)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \frac{b(a^2+b^2)}{a^2(bB-aC)} \\
& \quad \frac{a^2(bB-aC)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow 4014
\end{aligned}$$

$$\begin{aligned}
 & \frac{\frac{b(a^3(-C)+3a^2bB+3ab^2C-b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{bx(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a^2+b^2} - \frac{a(a^3(-C)-3ab^2C+2b^3B)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{b(a^2+b^2)}{a^2(bB-aC)} \\
 & \quad \frac{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2}{\downarrow 3042} \\
 & \frac{\frac{b(a^3(-C)+3a^2bB+3ab^2C-b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{bx(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a^2+b^2} - \frac{a(a^3(-C)-3ab^2C+2b^3B)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{b(a^2+b^2)}{a^2(bB-aC)} \\
 & \quad \frac{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2}{\downarrow 4013} \\
 & \quad \frac{a^2(bB-aC)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \frac{\frac{b(a^3(-C)+3a^2bB+3ab^2C-b^3B) \log(a \cos(c+dx)+b \sin(c+dx))}{d(a^2+b^2)} + \frac{bx(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a^2+b^2} - \frac{a(a^3(-C)-3ab^2C+2b^3B)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad b(a^2+b^2)
 \end{aligned}$$

input `Int[(Tan[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]`

output `-1/2*(a^2*(b*B - a*C))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2 - (((b*(a^3*B - 3*a*b^2*B + 3*a^2*b*C - b^3*C)*x)/(a^2 + b^2) + (b*(3*a^2*b*B - b^3*B - a^3*C + 3*a*b^2*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) - (a*(2*b^3*B - a^3*C - 3*a*b^2*C))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(b*(a^2 + b^2))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4013 $\text{Int}[\left(\frac{(c_.) + (d_.)\tan[e_.] + (f_.)x}{(a_.) + (b_.)\tan[e_.] + (f_.)x}\right), x_Symbol] \rightarrow \text{Simp}\left[\frac{c}{b*f}\right] * \text{Log}\left[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x], x\right] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

rule 4014 $\text{Int}\left[\frac{(c_.) + (d_.)\tan[e_.] + (f_.)x}{(a_.) + (b_.)\tan[e_.] + (f_.)x}\right], x_Symbol] \rightarrow \text{Simp}\left[\frac{(a*c + b*d)x}{a^2 + b^2}\right], x] + \text{Simp}\left[\frac{b*c - a*d}{a^2 + b^2} \text{Int}\left[\frac{b - a*\text{Tan}[e + f*x]}{a + b*\text{Tan}[e + f*x]}, x\right], x\right] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

rule 4087 $\text{Int}\left[\frac{(a_.) + (b_.)\tan[e_.] + (f_.)x}{(c_.) + (d_.)\tan[e_.] + (f_.)x}\right]^2 * \left(\frac{(A_.) + (B_.)\tan[e_.] + (f_.)x}{(c_.) + (d_.)\tan[e_.] + (f_.)x}\right)^n, x_Symbol] \rightarrow \text{Simp}\left[\frac{-(B*c - A*d)(b*c - a*d)^2 (c + d*\text{Tan}[e + f*x])^{n+1}}{(f*d^2(n+1)(c^2 + d^2))}, x\right] + \text{Simp}\left[\frac{1}{d(c^2 + d^2)} \text{Int}\left[\frac{(c + d*\text{Tan}[e + f*x])^{n+1}}{B(b*c - a*d)^2 + A*d(a^2*c - b^2*c + 2*a*b*d) + d(B(a^2*c - b^2*c + 2*a*b*d) + A(2*a*b*c - a^2*d + b^2*d))*\text{Tan}[e + f*x] + b^2*B(c^2 + d^2)*\text{Tan}[e + f*x]^2}, x\right], x\right] /;$ FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]

rule 4111 $\text{Int}\left[\frac{(a_.) + (b_.)\tan[e_.] + (f_.)x}{(c_.) + (d_.)\tan[e_.] + (f_.)x}\right]^m * \left(\frac{(A_.) + (B_.)\tan[e_.] + (f_.)x}{(c_.) + (d_.)\tan[e_.] + (f_.)x}\right)^2, x_Symbol] \rightarrow \text{Simp}\left[\frac{A*b^2 - a*b*B + a^2*C}{(a + b*\text{Tan}[e + f*x])^{m+1}}\right] / (b*f*(m+1)*(a^2 + b^2)), x] + \text{Simp}\left[\frac{1}{a^2 + b^2} \text{Int}\left[\frac{(a + b*\text{Tan}[e + f*x])^{m+1}}{b*B + a*(A - C) - (A*b - a*B - b*C)*\text{Tan}[e + f*x]}, x\right], x\right] /;$ FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]

rule 4115 $\text{Int}\left[\frac{(a_.) + (b_.)\tan[e_.] + (f_.)x}{(c_.) + (d_.)\tan[e_.] + (f_.)x}\right]^m * \left(\frac{(c_.) + (d_.)\tan[e_.] + (f_.)x}{(A_.) + (B_.)\tan[e_.] + (f_.)x}\right)^n * \left(\frac{(c_.) + (d_.)\tan[e_.] + (f_.)x}{(C_.) + (d_.)\tan[e_.] + (f_.)x}\right)^2, x_Symbol] \rightarrow \text{Simp}\left[\frac{1}{b^2} \text{Int}\left[\frac{(a + b*\text{Tan}[e + f*x])^{m+1}}{(c + d*\text{Tan}[e + f*x])^n * (b*B - a*C + b*C*\text{Tan}[e + f*x])}, x\right], x\right] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.18

method	result
derivativedivides	$-\frac{a^2(Bb-Ca)}{2b^2(a^2+b^2)(a+b \tan(dx+c))^2} - \frac{(3B a^2b - B b^3 - C a^3 + 3C a b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^3} + \frac{a(2B b^3 - C a^3 - 3C a b^2)}{(a^2+b^2)^2 b^2(a+b \tan(dx+c))} + \frac{(3B a^2b - C a^3 + 3C a b^2)}{d}$
default	$-\frac{a^2(Bb-Ca)}{2b^2(a^2+b^2)(a+b \tan(dx+c))^2} - \frac{(3B a^2b - B b^3 - C a^3 + 3C a b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^3} + \frac{a(2B b^3 - C a^3 - 3C a b^2)}{(a^2+b^2)^2 b^2(a+b \tan(dx+c))} + \frac{(3B a^2b - C a^3 + 3C a b^2)}{d}$
norman	$-\frac{(2B a b^3 - C a^4 - 3C a^2 b^2) \tan(dx+c)^2}{2ad(a^4+2b^2a^2+b^4)} - \frac{a(B a^3 - B a b^2 + 2C a^2 b)}{2db(a^4+2b^2a^2+b^4)} - \frac{(B a^3 - 3B a b^2 + 3C a^2 b - C b^3) a^2 x}{(a^4+2b^2a^2+b^4)(a^2+b^2)} - \frac{b^2(B a^3 - 3B a b^2 + 3C a^2 b - C b^3)}{(a^4+2b^2a^2+b^4)(a+b \tan(dx+c))^2}$
risch	$\frac{x B}{3ia^2b - ib^3 - a^3 + 3a b^2} - \frac{ix C}{3ia^2b - ib^3 - a^3 + 3a b^2} + \frac{6iB a^2 b x}{a^6 + 3b^2 a^4 + 3b^4 a^2 + b^6} - \frac{2iB b^3 x}{a^6 + 3b^2 a^4 + 3b^4 a^2 + b^6} - \frac{2iC a^2 x}{a^6 + 3b^2 a^4 + 3b^4 a^2 + b^6}$
parallelrisch	$-C a^7 - B a^6 b + 2B a^4 b^3 + 3B a^2 b^5 - 6C a^5 b^2 - 5C a^3 b^4 + 4B \tan(dx+c) a^3 b^4 + 4B \tan(dx+c) a b^6 - 2C \tan(dx+c) a^6 b - 8C \tan(dx+c) a^4 b^3$

input `int (tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*(-1/2*a^2*(B*b-C*a)/b^2/(a^2+b^2)/(a+b*tan(d*x+c))^2-(3*B*a^2*b-B*b^3-C*a^3+3*C*a*b^2)/(a^2+b^2)^3*ln(a+b*tan(d*x+c))+a*(2*B*b^3-C*a^3-3*C*a*b^2)/(a^2+b^2)^2/b^2/(a+b*tan(d*x+c))+1/(a^2+b^2)^3*(1/2*(3*B*a^2*b-B*b^3-C*a^3+3*C*a*b^2)*ln(1+tan(d*x+c)^2)+(-B*a^3+3*B*a*b^2-3*C*a^2*b+C*b^3)*arctan(tan(d*x+c)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 478 vs. 2(184) = 368.

Time = 0.10 (sec) , antiderivative size = 478, normalized size of antiderivative = 2.53

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{Ca^5 - 3Ba^4b - 5Ca^3b^2 + 3Ba^2b^3 - 2(Ba^5 + 3Ca^4b - 3Ba^3b^2 - Ca^2b^3)dx + (Ca^5 + Ba^4b + 7Ca^3b^2 - 5Ca^2b^3 - 3Ca^2b^2 - 3Ca^2b - 3Ca^2 - 3Ca - 3C)dx + (Ca^5 + Ba^4b + 7Ca^3b^2 - 5Ca^2b^3 - 3Ca^2b^2 - 3Ca^2b - 3Ca^2 - 3Ca - 3C)}{(a + b \tan(c + dx))^3}$$

input `integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output `1/2*(C*a^5 - 3*B*a^4*b - 5*C*a^3*b^2 + 3*B*a^2*b^3 - 2*(B*a^5 + 3*C*a^4*b - 3*B*a^3*b^2 - C*a^2*b^3)*d*x + (C*a^5 + B*a^4*b + 7*C*a^3*b^2 - 5*B*a^2*b^3 - 2*(B*a^3*b^2 + 3*C*a^2*b^3 - 3*B*a*b^4 - C*b^5)*d*x)*tan(d*x + c)^2 + (C*a^5 - 3*B*a^4*b - 3*C*a^3*b^2 + B*a^2*b^3 + (C*a^3*b^2 - 3*B*a^2*b^3 - 3*C*a*b^4 + B*b^5)*tan(d*x + c)^2 + 2*(C*a^4*b - 3*B*a^3*b^2 - 3*C*a^2*b^3 + B*a*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + 2*(B*a^5 + 3*C*a^4*b - 3*B*a^3*b^2 - 3*C*a^2*b^3 + 2*B*a*b^4 - 2*(B*a^4*b + 3*C*a^3*b^2 - 3*B*a^2*b^3 - C*a*b^4)*d*x)*tan(d*x + c)/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*d*tan(d*x + c)^2 + 2*(a^7*b + 3*a^5*b^3 + 3*a^3*b^5 + a*b^7)*d*tan(d*x + c) + (a^8 + 3*a^6*b^2 + 3*a^4*b^4 + a^2*b^6)*d)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.76

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx =$$

$$\frac{2(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)(dx+c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{2(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(b \tan(dx+c)+a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(\tan(dx+c)+\frac{a}{b})}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{2d}{2d}$$

output

```

-(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d +
3*a^2*b^4*d + b^6*d) - 1/2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*log(tan
(d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) + (C*a^3*b -
3*B*a^2*b^2 - 3*C*a*b^3 + B*b^4)*log(abs(b*tan(d*x + c) + a))/(a^6*b*d + 3
*a^4*b^3*d + 3*a^2*b^5*d + b^7*d) - 1/2*(C*a^7 + B*a^6*b + 6*C*a^5*b^2 - 2
*B*a^4*b^3 + 5*C*a^3*b^4 - 3*B*a^2*b^5 + 2*(C*a^6*b + 4*C*a^4*b^3 - 2*B*a^
3*b^4 + 3*C*a^2*b^5 - 2*B*a*b^6)*tan(d*x + c))/((a^2 + b^2)^3*(b*tan(d*x +
c) + a)^2*b^2*d)

```

Mupad [B] (verification not implemented)

Time = 5.95 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.48

$$\begin{aligned}
& \int \frac{\tan(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx \\
&= \frac{\ln(a+b \tan(c+dx))(C a^3 - 3 B a^2 b - 3 C a b^2 + B b^3)}{d(a^2 + b^2)^3} \\
&\quad - \frac{\ln(\tan(c+dx) - i)(-C + B i)}{2 d(-a^3 - a^2 b 3i + 3 a b^2 + b^3 i)} - \frac{\ln(\tan(c+dx) + i)(B - C i)}{2 d(-a^3 i - 3 a^2 b + a b^2 3i + b^3)} \\
&\quad - \frac{\frac{a(C a^4 + B a^3 b + 5 C a^2 b^2 - 3 B a b^3)}{2 b^2(a^4 + 2 a^2 b^2 + b^4)} + \frac{\tan(c+dx)(C a^4 + 3 C a^2 b^2 - 2 B a b^3)}{b(a^4 + 2 a^2 b^2 + b^4)}}{d(a^2 + 2 a b \tan(c+dx) + b^2 \tan(c+dx)^2)}
\end{aligned}$$

input

```

int((tan(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x)
)^3,x)

```

output

```

(log(a + b*tan(c + d*x))*(B*b^3 + C*a^3 - 3*B*a^2*b - 3*C*a*b^2))/(d*(a^2
+ b^2)^3) - (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(3*a*b^2 - a^2*b*3i -
a^3 + b^3*1i)) - (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a*b^2*3i - 3*a
^2*b - a^3*1i + b^3)) - ((a*(C*a^4 + 5*C*a^2*b^2 - 3*B*a*b^3 + B*a^3*b))/(
2*b^2*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(C*a^4 + 3*C*a^2*b^2 - 2*B*
a*b^3))/(b*(a^4 + b^4 + 2*a^2*b^2)))/(d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*
tan(c + d*x)))

```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 974, normalized size of antiderivative = 5.15

$$\int \frac{\tan(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input

```
int(tan(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)
```

output

```
( - log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a**3*b**2*c + 3*log(tan(c + d
*x)**2 + 1)*tan(c + d*x)**2*a**2*b**4 + 3*log(tan(c + d*x)**2 + 1)*tan(c +
d*x)**2*a*b**4*c - log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*b**6 - 2*log(
tan(c + d*x)**2 + 1)*tan(c + d*x)*a**4*b*c + 6*log(tan(c + d*x)**2 + 1)*ta
n(c + d*x)*a**3*b**3 + 6*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**3*c
- 2*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a*b**5 - log(tan(c + d*x)**2 +
1)*a**5*c + 3*log(tan(c + d*x)**2 + 1)*a**4*b**2 + 3*log(tan(c + d*x)**2 +
1)*a**3*b**2*c - log(tan(c + d*x)**2 + 1)*a**2*b**4 + 2*log(tan(c + d*x)*
b + a)*tan(c + d*x)**2*a**3*b**2*c - 6*log(tan(c + d*x)*b + a)*tan(c + d*x
)**2*a**2*b**4 - 6*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a*b**4*c + 2*lo
g(tan(c + d*x)*b + a)*tan(c + d*x)**2*b**6 + 4*log(tan(c + d*x)*b + a)*tan
(c + d*x)*a**4*b*c - 12*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**3*b**3 - 1
2*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**2*b**3*c + 4*log(tan(c + d*x)*b
+ a)*tan(c + d*x)*a*b**5 + 2*log(tan(c + d*x)*b + a)*a**5*c - 6*log(tan(c
+ d*x)*b + a)*a**4*b**2 - 6*log(tan(c + d*x)*b + a)*a**3*b**2*c + 2*log(ta
n(c + d*x)*b + a)*a**2*b**4 + tan(c + d*x)**2*a**5*c - 2*tan(c + d*x)**2*a
**3*b**3*d*x + 4*tan(c + d*x)**2*a**3*b**2*c - 2*tan(c + d*x)**2*a**2*b**4
- 6*tan(c + d*x)**2*a**2*b**3*c*d*x + 6*tan(c + d*x)**2*a*b**5*d*x + 3*ta
n(c + d*x)**2*a*b**4*c - 2*tan(c + d*x)**2*b**6 + 2*tan(c + d*x)**2*b**5*c
*d*x - 4*tan(c + d*x)*a**4*b**2*d*x - 12*tan(c + d*x)*a**3*b**2*c*d*x + ...
```

3.41
$$\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{(a+b \tan(c+dx))^3} dx$$

Optimal result	464
Mathematica [C] (verified)	465
Rubi [A] (verified)	465
Maple [A] (verified)	468
Fricas [B] (verification not implemented)	469
Sympy [F(-2)]	469
Maxima [A] (verification not implemented)	470
Giac [A] (verification not implemented)	470
Mupad [B] (verification not implemented)	471
Reduce [B] (verification not implemented)	472

Optimal result

Integrand size = 32, antiderivative size = 179

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{(3a^2bB - b^3B - a^3C + 3ab^2C) x}{(a^2 + b^2)^3}$$

$$- \frac{(a^3B - 3ab^2B + 3a^2bC - b^3C) \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2)^3 d}$$

$$+ \frac{a(bB - aC)}{2b(a^2 + b^2) d(a + b \tan(c + dx))^2} + \frac{a^2B - b^2B + 2abC}{(a^2 + b^2)^2 d(a + b \tan(c + dx))}$$

output

```
(3*B*a^2*b-B*b^3-C*a^3+3*C*a*b^2)*x/(a^2+b^2)^3-(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^3/d+1/2*a*(B*b-C*a)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+(B*a^2-B*b^2+2*C*a*b)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.70 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.05

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{\frac{(B+iC) \log(i - \tan(c+dx))}{(a+ib)^3} + \frac{(B-iC) \log(i + \tan(c+dx))}{(a-ib)^3} - \frac{2(a^3B - 3ab^2B + 3a^2bC - b^3C) \log(a+b \tan(c+dx))}{(a^2+b^2)^3} + \frac{a(bB-aC)}{b(a^2+b^2)(a+b \tan(c+dx))}}{2d}$$

input

```
Integrate[(B*Tan[c + d*x] + C*Tan[c + d*x]^2)/(a + b*Tan[c + d*x]^3,x]
```

output

```
((B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)^3 + ((B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b)^3 - (2*(a^3*B - 3*a*b^2*B + 3*a^2*b*C - b^3*C)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^3 + (a*(b*B - a*C))/(b*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) + (2*(a^2*B - b^2*B + 2*a*b*C))/((a^2 + b^2)^2*(a + b*Tan[c + d*x]))/(2*d)
```

Rubi [A] (verified)

Time = 1.40 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4111, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$\downarrow \text{4111}$$

$$\frac{\int \frac{bB - aC + (aB + bC) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} + \frac{a(bB - aC)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}$$

$$\begin{aligned}
& \int \frac{bB - aC + (aB + bC) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx + \frac{a(bB - aC)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \int \frac{-Ca^2 + 2bBa + b^2C + (Ba^2 + 2bCa - b^2B) \tan(c + dx)}{a + b \tan(c + dx)} dx + \frac{a^2B + 2abC - b^2B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
& \quad \frac{a^2 + b^2}{a(bB - aC)} \\
& \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \int \frac{-Ca^2 + 2bBa + b^2C + (Ba^2 + 2bCa - b^2B) \tan(c + dx)}{a + b \tan(c + dx)} dx + \frac{a^2B + 2abC - b^2B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
& \quad \frac{a^2 + b^2}{a(bB - aC)} \\
& \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \quad \downarrow \text{4014} \\
& \frac{x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2} - \frac{(a^3B + 3a^2bC - 3ab^2B - b^3C) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^2B + 2abC - b^2B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
& \quad \frac{a^2 + b^2}{a(bB - aC)} \\
& \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \frac{x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2} - \frac{(a^3B + 3a^2bC - 3ab^2B - b^3C) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^2B + 2abC - b^2B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
& \quad \frac{a^2 + b^2}{a(bB - aC)} \\
& \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \quad \downarrow \text{4013} \\
& \frac{a(bB - aC)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} + \\
& \quad \frac{x(a^3(-C) + 3a^2bB + 3ab^2C - b^3B)}{a^2 + b^2} - \frac{(a^3B + 3a^2bC - 3ab^2B - b^3C) \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)} \\
& \quad \frac{a^2B + 2abC - b^2B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \frac{a^2 + b^2}{a^2 + b^2}
\end{aligned}$$

input `Int[(B*Tan[c + d*x] + C*Tan[c + d*x]^2)/(a + b*Tan[c + d*x])^3,x]`

output

$$\frac{(a*(b*B - a*C))/(2*b*(a^2 + b^2)*d*(a + b*\tan[c + d*x])^2) + (((3*a^2*b*B - b^3*B - a^3*C + 3*a*b^2*C)*x)/(a^2 + b^2) - ((a^3*B - 3*a*b^2*B + 3*a^2*b*C - b^3*C)*\log[a*\cos[c + d*x] + b*\sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) + (a^2*B - b^2*B + 2*a*b*C)/((a^2 + b^2)*d*(a + b*\tan[c + d*x]))}{(a^2 + b^2)}$$

Defintions of rubi rules used

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4012

$$\text{Int}[(a_. + (b_.)*\tan[(e_.) + (f_.)*(x_)])^m*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*((a + b*\tan[e + f*x])^{m+1}/(f*(m+1)*(a^2 + b^2))), x] + \text{Simp}[1/(a^2 + b^2) \text{ Int}[(a + b*\tan[e + f*x])^{m+1}*\text{Simp}[a*c + b*d - (b*c - a*d)*\tan[e + f*x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[m, -1]$$

rule 4013

$$\text{Int}[(c_. + (d_.)*\tan[(e_.) + (f_.)*(x_)])/(a_. + (b_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\log[\text{RemoveContent}[a*\cos[e + f*x] + b*\sin[e + f*x], x]], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$$

rule 4014

$$\text{Int}[(c_. + (d_.)*\tan[(e_.) + (f_.)*(x_)])/(a_. + (b_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*c + b*d)*(x/(a^2 + b^2)), x] + \text{Simp}[(b*c - a*d)/(a^2 + b^2) \text{ Int}[(b - a*\tan[e + f*x])/(a + b*\tan[e + f*x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[a*c + b*d, 0]$$

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{\frac{a(Bb-Ca)}{2(a^2+b^2)b(a+b\tan(dx+c))^2} + \frac{Ba^2-Bb^2+2Cab}{(a^2+b^2)^2(a+b\tan(dx+c))} - \frac{(Ba^3-3Bab^2+3Ca^2b-Cb^3)\ln(a+b\tan(dx+c))}{(a^2+b^2)^3} + \frac{(Ba^3-3Bab^2+3Ca^2b-Cb^3)}{d(a^2+b^2)^3}$
default	$\frac{\frac{a(Bb-Ca)}{2(a^2+b^2)b(a+b\tan(dx+c))^2} + \frac{Ba^2-Bb^2+2Cab}{(a^2+b^2)^2(a+b\tan(dx+c))} - \frac{(Ba^3-3Bab^2+3Ca^2b-Cb^3)\ln(a+b\tan(dx+c))}{(a^2+b^2)^3} + \frac{(Ba^3-3Bab^2+3Ca^2b-Cb^3)}{d(a^2+b^2)^3}$
norman	$\frac{(Ba^2b^2-Bb^4+2Ca^2b^3)\tan(dx+c)}{db(a^4+2b^2a^2+b^4)} + \frac{(3Ba^2b-Bb^3-Ca^3+3Ca^2b^2)a^2x}{(a^4+2b^2a^2+b^4)(a^2+b^2)} + \frac{b^2(3Ba^2b-Bb^3-Ca^3+3Ca^2b^2)x\tan(dx+c)^2}{(a^4+2b^2a^2+b^4)(a^2+b^2)} + \frac{a(3Ba^2b-Bb^3-Ca^3+3Ca^2b^2)}{(a+b\tan(dx+c))^2}$
risch	$\frac{ixB}{3ia^2b-ib^3-a^3+3ab^2} + \frac{xC}{3ia^2b-ib^3-a^3+3ab^2} + \frac{2ia^3Bx}{a^6+3b^2a^4+3b^4a^2+b^6} - \frac{6iab^2Bx}{a^6+3b^2a^4+3b^4a^2+b^6} + \frac{6ia^2b^2Cx}{a^6+3b^2a^4+3b^4a^2+b^6}$
parallelrisc	$\frac{6Bx\tan(dx+c)^2a^2b^5d-2Cx\tan(dx+c)^2a^3b^4d+6Cx\tan(dx+c)^2ab^6d+12Bx\tan(dx+c)a^3b^4d-4Bx\tan(dx+c)ab^6d}{(a+b\tan(dx+c))^2}$

input

```
int((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBO
SE)
```

output

```
1/d*(1/2*a*(B*b-C*a)/(a^2+b^2)/b/(a+b*tan(d*x+c))^2+(B*a^2-B*b^2+2*C*a*b)/
(a^2+b^2)^2/(a+b*tan(d*x+c))-(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)/(a^2+b^2)^3
*ln(a+b*tan(d*x+c))+1/(a^2+b^2)^3*(1/2*(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*l
n(1+tan(d*x+c)^2)+(3*B*a^2*b-B*b^3-C*a^3+3*C*a*b^2)*arctan(tan(d*x+c))))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 488 vs. $2(176) = 352$.

Time = 0.09 (sec) , antiderivative size = 488, normalized size of antiderivative = 2.73

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx = \frac{3Ca^4b - 5Ba^3b^2 - 3Ca^2b^3 + Bab^4 + 2(Ca^5 - 3Ba^4b - 3Ca^3b^2 + Ba^2b^3)dx - (Ca^4b - 3Ba^3b^2 - 5$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output `-1/2*(3*C*a^4*b - 5*B*a^3*b^2 - 3*C*a^2*b^3 + B*a*b^4 + 2*(C*a^5 - 3*B*a^4*b - 3*C*a^3*b^2 + B*a^2*b^3)*d*x - (C*a^4*b - 3*B*a^3*b^2 - 5*C*a^2*b^3 + 3*B*a*b^4 - 2*(C*a^3*b^2 - 3*B*a^2*b^3 - 3*C*a*b^4 + B*b^5)*d*x)*tan(d*x + c)^2 + (B*a^5 + 3*C*a^4*b - 3*B*a^3*b^2 - C*a^2*b^3 + (B*a^3*b^2 + 3*C*a^2*b^3 - 3*B*a*b^4 - C*b^5)*tan(d*x + c)^2 + 2*(B*a^4*b + 3*C*a^3*b^2 - 3*B*a^2*b^3 - C*a*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 2*(C*a^5 - 2*B*a^4*b - 3*C*a^3*b^2 + 3*B*a^2*b^3 + 2*C*a*b^4 - B*b^5 - 2*(C*a^4*b - 3*B*a^3*b^2 - 3*C*a^2*b^3 + B*a*b^4)*d*x)*tan(d*x + c))/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*d*tan(d*x + c)^2 + 2*(a^7*b + 3*a^5*b^3 + 3*a^3*b^5 + a*b^7)*d*tan(d*x + c) + (a^8 + 3*a^6*b^2 + 3*a^4*b^4 + a^2*b^6)*d)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate((B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.84

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx =$$

$$\frac{2(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3)(dx + c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{2(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3) \log(b \tan(dx + c) + a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3) \log(\tan(dx + c) + a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{2d}{2d}$$

input

```
integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="maxima")
```

output

```
-1/2*(2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*log(b*tan(d*x + c) + a)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (C*a^4 - 3*B*a^3*b - 3*C*a^2*b^2 + B*a*b^3 - 2*(B*a^2*b^2 + 2*C*a*b^3 - B*b^4)*tan(d*x + c))/(a^6*b + 2*a^4*b^3 + a^2*b^5 + (a^4*b^3 + 2*a^2*b^5 + b^7)*tan(d*x + c)^2 + 2*(a^5*b^2 + 2*a^3*b^4 + a*b^6)*tan(d*x + c)))/d
```

Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.80

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx = -\frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3)(dx + c)}{a^6d + 3a^4b^2d + 3a^2b^4d + b^6d}$$

$$+ \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3) \log(\tan(dx + c)^2 + 1)}{2(a^6d + 3a^4b^2d + 3a^2b^4d + b^6d)}$$

$$- \frac{(Ba^3b + 3Ca^2b^2 - 3Bab^3 - Cb^4) \log(|b \tan(dx + c) + a|)}{a^6bd + 3a^4b^3d + 3a^2b^5d + b^7d}$$

$$- \frac{Ca^6 - 3Ba^5b - 2Ca^4b^2 - 2Ba^3b^3 - 3Ca^2b^4 + Bab^5 - 2(Ba^4b^2 + 2Ca^3b^3 + 2Cab^5 - Bb^6) \tan(dx + c)}{2(a^2 + b^2)^3(b \tan(dx + c) + a)^2bd}$$

input

```
integrate((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="giac")
```

output

```

-(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d +
3*a^2*b^4*d + b^6*d) + 1/2*(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*log(tan
(d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) - (B*a^3*b +
3*C*a^2*b^2 - 3*B*a*b^3 - C*b^4)*log(abs(b*tan(d*x + c) + a))/(a^6*b*d + 3
*a^4*b^3*d + 3*a^2*b^5*d + b^7*d) - 1/2*(C*a^6 - 3*B*a^5*b - 2*C*a^4*b^2 -
2*B*a^3*b^3 - 3*C*a^2*b^4 + B*a*b^5 - 2*(B*a^4*b^2 + 2*C*a^3*b^3 + 2*C*a*
b^5 - B*b^6)*tan(d*x + c))/((a^2 + b^2)^3*(b*tan(d*x + c) + a)^2*b*d)

```

Mupad [B] (verification not implemented)

Time = 5.65 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.58

$$\begin{aligned}
& \int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx \\
&= \frac{\frac{\tan(c+dx)(Ba^2b+2Cab^2-Bb^3)}{a^4+2a^2b^2+b^4} - \frac{Ca^4-3Ba^3b-3Ca^2b^2+Bab^3}{2b(a^4+2a^2b^2+b^4)}}{d(a^2+2ab\tan(c+dx)+b^2\tan(c+dx)^2)} \\
&\quad - \frac{\ln(a+b\tan(c+dx))\left(\frac{Ba+3Cb}{(a^2+b^2)^2} - \frac{4b^2(Ba+Cb)}{(a^2+b^2)^3}\right)}{d} \\
&\quad - \frac{\ln(\tan(c+dx)-i)(-C+Bi)}{2d(-a^3i+3a^2b+ab^23i-b^3)} - \frac{\ln(\tan(c+dx)+i)(B-Ci)}{2d(-a^3+a^2b3i+3ab^2-b^3i)}
\end{aligned}$$

input

```
int((B*tan(c + d*x) + C*tan(c + d*x)^2)/(a + b*tan(c + d*x))^3,x)
```

output

```

((tan(c + d*x)*(B*a^2*b - B*b^3 + 2*C*a*b^2))/(a^4 + b^4 + 2*a^2*b^2) - (C
*a^4 - 3*C*a^2*b^2 + B*a*b^3 - 3*B*a^3*b)/(2*b*(a^4 + b^4 + 2*a^2*b^2)))/(
d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*tan(c + d*x))) - (log(a + b*tan(c + d*
x))*(B*a + 3*C*b)/(a^2 + b^2)^2 - (4*b^2*(B*a + C*b))/(a^2 + b^2)^3))/d -
(log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*b^2*3i + 3*a^2*b - a^3*1i - b
^3)) - (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(3*a*b^2 + a^2*b*3i - a^3
- b^3*1i))

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 994, normalized size of antiderivative = 5.55

$$\int \frac{B \tan(c + dx) + C \tan^2(c + dx)}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `int((B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)`

output

```
(log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a**4*b**4 + 3*log(tan(c + d*x)**
2 + 1)*tan(c + d*x)**2*a**3*b**4*c - 3*log(tan(c + d*x)**2 + 1)*tan(c + d*
x)**2*a**2*b**6 - log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2*a*b**6*c + 2*lo
g(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**5*b**3 + 6*log(tan(c + d*x)**2 + 1)
*tan(c + d*x)*a**4*b**3*c - 6*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**3*b
**5 - 2*log(tan(c + d*x)**2 + 1)*tan(c + d*x)*a**2*b**5*c + log(tan(c + d*
x)**2 + 1)*a**6*b**2 + 3*log(tan(c + d*x)**2 + 1)*a**5*b**2*c - 3*log(tan(
c + d*x)**2 + 1)*a**4*b**4 - log(tan(c + d*x)**2 + 1)*a**3*b**4*c - 2*log(
tan(c + d*x)*b + a)*tan(c + d*x)**2*a**4*b**4 - 6*log(tan(c + d*x)*b + a)*
tan(c + d*x)**2*a**3*b**4*c + 6*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a*
**2*b**6 + 2*log(tan(c + d*x)*b + a)*tan(c + d*x)**2*a*b**6*c - 4*log(tan(c
 + d*x)*b + a)*tan(c + d*x)*a**5*b**3 - 12*log(tan(c + d*x)*b + a)*tan(c +
d*x)*a**4*b**3*c + 12*log(tan(c + d*x)*b + a)*tan(c + d*x)*a**3*b**5 + 4*
log(tan(c + d*x)*b + a)*tan(c + d*x)*a**2*b**5*c - 2*log(tan(c + d*x)*b +
a)*a**6*b**2 - 6*log(tan(c + d*x)*b + a)*a**5*b**2*c + 6*log(tan(c + d*x)*
b + a)*a**4*b**4 + 2*log(tan(c + d*x)*b + a)*a**3*b**4*c - tan(c + d*x)**2
*a**4*b**4 - 2*tan(c + d*x)**2*a**4*b**3*c*d*x + 6*tan(c + d*x)**2*a**3*b*
**5*d*x - 2*tan(c + d*x)**2*a**3*b**4*c + 6*tan(c + d*x)**2*a**2*b**5*c*d*x
 - 2*tan(c + d*x)**2*a*b**7*d*x - 2*tan(c + d*x)**2*a*b**6*c + tan(c + d*x)
)**2*b**8 - 4*tan(c + d*x)*a**5*b**2*c*d*x + 12*tan(c + d*x)*a**4*b**4*...
```

3.42
$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

Optimal result	473
Mathematica [C] (verified)	474
Rubi [A] (verified)	474
Maple [A] (verified)	477
Fricas [B] (verification not implemented)	478
Sympy [F(-2)]	479
Maxima [A] (verification not implemented)	479
Giac [A] (verification not implemented)	480
Mupad [B] (verification not implemented)	480
Reduce [B] (verification not implemented)	481

Optimal result

Integrand size = 38, antiderivative size = 175

$$\int \frac{\cot(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(a^3B - 3ab^2B + 3a^2bC - b^3C)x}{(a^2 + b^2)^3}$$

$$+ \frac{(3a^2bB - b^3B - a^3C + 3ab^2C) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^3 d}$$

$$- \frac{bB - aC}{2(a^2 + b^2) d(a + b \tan(c+dx))^2} - \frac{2abB - a^2C + b^2C}{(a^2 + b^2)^2 d(a + b \tan(c+dx))}$$

output

```
(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*x/(a^2+b^2)^3+(3*B*a^2*b-B*b^3-C*a^3+3*C
*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^3/d-1/2*(B*b-C*a)/(a^2+b^2
)/d/(a+b*tan(d*x+c))^2-(2*B*a*b-C*a^2+C*b^2)/(a^2+b^2)^2/d/(a+b*tan(d*x+c)
)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.04 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.39

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$C \left(\frac{i \log(i - \tan(c+dx))}{(a+ib)^2} - \frac{i \log(i + \tan(c+dx))}{(a-ib)^2} + \frac{2b(-2a \log(a+b \tan(c+dx)) + \frac{a^2+b^2}{a+b \tan(c+dx)})}{(a^2+b^2)^2} \right) + (bB - aC) \left(\frac{i \log(i - \tan(c+dx))}{(a+ib)^3} - \frac{i \log(i + \tan(c+dx))}{(a-ib)^3} \right)$$

$$2bd$$

input

```
Integrate[(Cot[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

```
-1/2*(C*((I*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b*(-2*a*Log[a + b*Tan[c + d*x]] + (a^2 + b^2)/(a + b*Tan[c + d*x])))/(a^2 + b^2)^2) + (b*B - a*C)*((I*Log[I - Tan[c + d*x]])/(a + I*b)^3 - Log[I + Tan[c + d*x]]/(I*a + b)^3 + (b*((-6*a^2 + 2*b^2)*Log[a + b*Tan[c + d*x]] + ((a^2 + b^2)*(5*a^2 + b^2 + 4*a*b*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2))/(a^2 + b^2)^3))/(b*d)
```

Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.13, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3042, 4115, 3042, 4012, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{\tan(c+dx)(a+b \tan(c+dx))^3} dx$$

$$\begin{aligned}
& \downarrow 4115 \\
& \int \frac{B + C \tan(c + dx)}{(a + b \tan(c + dx))^3} dx \\
& \downarrow 3042 \\
& \int \frac{B + C \tan(c + dx)}{(a + b \tan(c + dx))^3} dx \\
& \downarrow 4012 \\
& \frac{\int \frac{aB + bC - (bB - aC) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} - \frac{bB - aC}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{aB + bC - (bB - aC) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} - \frac{bB - aC}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} \\
& \downarrow 4012 \\
& \frac{\int \frac{Ba^2 + 2bCa - b^2B - (-Ca^2 + 2bBa + b^2C) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{a^2(-C) + 2abB + b^2C}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
& \frac{a^2 + b^2}{bB - aC} \\
& \frac{2d(a^2 + b^2)(a + b \tan(c + dx))^2}{\downarrow 3042} \\
& \frac{\int \frac{Ba^2 + 2bCa - b^2B - (-Ca^2 + 2bBa + b^2C) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{a^2(-C) + 2abB + b^2C}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
& \frac{a^2 + b^2}{bB - aC} \\
& \frac{2d(a^2 + b^2)(a + b \tan(c + dx))^2}{\downarrow 4014} \\
& \frac{(a^3(-C) + 3a^2bB + 3ab^2C - b^3B) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(a^3B + 3a^2bC - 3ab^2B - b^3C)}{a^2 + b^2} - \frac{a^2(-C) + 2abB + b^2C}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
& \frac{a^2 + b^2}{bB - aC} \\
& \frac{2d(a^2 + b^2)(a + b \tan(c + dx))^2}{\downarrow 3042}
\end{aligned}$$

$$\frac{\frac{(a^3(-C)+3a^2bB+3ab^2C-b^3B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a^2+b^2} - \frac{a^2(-C)+2abB+b^2C}{d(a^2+b^2)(a+b \tan(c+dx))} - \frac{a^2+b^2}{bB-aC}}{2d(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 4013

$$\frac{\frac{(a^3(-C)+3a^2bB+3ab^2C-b^3B) \log(a \cos(c+dx)+b \sin(c+dx))}{d(a^2+b^2)} + \frac{x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a^2+b^2} - \frac{a^2(-C)+2abB+b^2C}{d(a^2+b^2)(a+b \tan(c+dx))} - \frac{a^2+b^2}{bB-aC}}{2d(a^2+b^2)(a+b \tan(c+dx))^2}$$

input `Int[(Cot[c + d*x]*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]`

output `-1/2*(b*B - a*C)/((a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((a^3*B - 3*a*b^2*B + 3*a^2*b*C - b^3*C)*x)/(a^2 + b^2) + ((3*a^2*b*B - b^3*B - a^3*C + 3*a*b^2*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) - (2*a*b*B - a^2*C + b^2*C)/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a^2 + b^2)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]) , x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4013 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/(a_ + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4014 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/(a_ + (b_)*tan[(e_) + (f_
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4115 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_
+ (f_)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]
```

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{(3B a^2 b - B b^3 - C a^3 + 3C a b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^3} - \frac{Bb-Ca}{2(a^2+b^2)(a+b \tan(dx+c))^2} - \frac{2Bab-C a^2+C b^2}{(a^2+b^2)^2(a+b \tan(dx+c))} + \frac{(-3B a^2 b+B b^3)}{d}$
default	$\frac{(3B a^2 b - B b^3 - C a^3 + 3C a b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)^3} - \frac{Bb-Ca}{2(a^2+b^2)(a+b \tan(dx+c))^2} - \frac{2Bab-C a^2+C b^2}{(a^2+b^2)^2(a+b \tan(dx+c))} + \frac{(-3B a^2 b+B b^3)}{d}$
parallelrisch	$6(a+b \tan(dx+c))^2 a(B a^2 b - \frac{1}{3} B b^3 - \frac{1}{3} C a^3 + C a b^2) \ln(a+b \tan(dx+c)) - 3(a+b \tan(dx+c))^2 a(B a^2 b - \frac{1}{3} B b^3 - \frac{1}{3} C a^3)$
norman	$\frac{(B a^3 - 3B a b^2 + 3C a^2 b - C b^3) a^2 x}{(a^4 + 2b^2 a^2 + b^4)(a^2 + b^2)} + \frac{b^2 (B a^3 - 3B a b^2 + 3C a^2 b - C b^3) x \tan(dx+c)^2}{(a^4 + 2b^2 a^2 + b^4)(a^2 + b^2)} - \frac{3B a^2 b^2 + B b^4 - 2C a^3 b}{2bd(a^4 + 2b^2 a^2 + b^4)} + \frac{b(2B a b^2 - C a^2 b)}{2da(a^4 + 2b^2 a^2 + b^4)} - \frac{(-3B a^2 b + B b^3)}{(a+b \tan(dx+c))^2}$
risch	$-\frac{x B}{3i a^2 b - i b^3 - a^3 + 3a b^2} + \frac{i x C}{3i a^2 b - i b^3 - a^3 + 3a b^2} - \frac{6i B a^2 b x}{a^6 + 3b^2 a^4 + 3b^4 a^2 + b^6} + \frac{2i B b^3 x}{a^6 + 3b^2 a^4 + 3b^4 a^2 + b^6} + \frac{2i C}{a^6 + 3b^2 a^4 + 3b^4 a^2 + b^6}$

input `int(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{(3Ba^2b - Bb^3 - Ca^3 + 3C*ab^2)}{(a^2+b^2)^3} \ln(a+b\tan(dx+c)) - \frac{1}{2} \frac{(Bb - Ca)}{(a^2+b^2)} \frac{1}{(a+b\tan(dx+c))^2} - \frac{(2B*ab - Ca^2 + C*b^2)}{(a^2+b^2)^2} \frac{1}{(a+b\tan(dx+c))} + \frac{1}{(a^2+b^2)^3} \left(\frac{1}{2} (-3Ba^2b + Bb^3 + Ca^3 - 3C*ab^2) \ln(1 + \tan(dx+c)^2) + (Ba^3 - 3B*ab^2 + 3Ca^2b - C*b^3) \arctan(\tan(dx+c)) \right) \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 482 vs. $2(171) = 342$.

Time = 0.10 (sec) , antiderivative size = 482, normalized size of antiderivative = 2.75

$$\int \frac{\cot(c+dx)(B\tan(c+dx)+C\tan^2(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{5Ca^3b^2 - 7Ba^2b^3 - Cab^4 - Bb^5 + 2(Ba^5 + 3Ca^4b - 3Ba^3b^2 - Ca^2b^3)dx - (3Ca^3b^2 - 5Ba^2b^3 - 3C$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output
$$\frac{1}{2} (5Ca^3b^2 - 7Ba^2b^3 - Cab^4 - Bb^5 + 2(Ba^5 + 3Ca^4b - 3B*a^3*b^2 - C*a^2*b^3) * dx - (3Ca^3b^2 - 5Ba^2b^3 - 3C*a*b^4 + B*b^5 - 2*(Ba^3b^2 + 3Ca^2b^3 - 3B*a*b^4 - C*b^5) * dx) * \tan(dx+c)^2 - (Ca^5 - 3B*a^4*b - 3Ca^3b^2 + Ba^2b^3 + (Ca^3b^2 - 3B*a^2b^3 - 3C*a*b^4 + B*b^5) * \tan(dx+c)^2 + 2*(Ca^4b - 3B*a^3b^2 - 3Ca^2b^3 + B*a*b^4) * \tan(dx+c)) * \log((b^2 * \tan(dx+c)^2 + 2*a*b * \tan(dx+c) + a^2) / (\tan(dx+c)^2 + 1)) - 2*(2Ca^4b - 3B*a^3b^2 - 3Ca^2b^3 + 3B*a*b^4 + C*b^5 - 2*(Ba^4b + 3Ca^3b^2 - 3B*a^2b^3 - C*a*b^4) * dx) * \tan(dx+c) / ((a^6b^2 + 3a^4b^4 + 3a^2b^6 + b^8) * d * \tan(dx+c)^2 + 2*(a^7b + 3a^5b^3 + 3a^3b^5 + a*b^7) * d * \tan(dx+c) + (a^8 + 3a^6b^2 + 3a^4b^4 + a^2b^6) * d)$$

Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3,x)`

output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.83

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{2(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)(dx+c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{2(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(b \tan(dx+c)+a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(\tan(dx+c)+a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6}$$

$2d$

input `integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output $\frac{1}{2} * (2 * (B * a^3 + 3 * C * a^2 * b - 3 * B * a * b^2 - C * b^3) * (d * x + c) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) - 2 * (C * a^3 - 3 * B * a^2 * b - 3 * C * a * b^2 + B * b^3) * \log(b * \tan(d * x + c) + a) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) + (C * a^3 - 3 * B * a^2 * b - 3 * C * a * b^2 + B * b^3) * \log(\tan(d * x + c)^2 + 1) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) + (3 * C * a^3 - 5 * B * a^2 * b - C * a * b^2 - B * b^3 + 2 * (C * a^2 * b - 2 * B * a * b^2 - C * b^3) * \tan(d * x + c)) / (a^6 + 2 * a^4 * b^2 + a^2 * b^4 + (a^4 * b^2 + 2 * a^2 * b^4 + b^6) * \tan(d * x + c)^2 + 2 * (a^5 * b + 2 * a^3 * b^3 + a * b^5) * \tan(d * x + c))) / d$

Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.80

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)(dx+c)}{a^6d + 3a^4b^2d + 3a^2b^4d + b^6d}$$

$$+ \frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(\tan(dx+c)^2 + 1)}{2(a^6d + 3a^4b^2d + 3a^2b^4d + b^6d)}$$

$$- \frac{(Ca^3b - 3Ba^2b^2 - 3Cab^3 + Bb^4) \log(|b \tan(dx+c) + a|)}{a^6bd + 3a^4b^3d + 3a^2b^5d + b^7d}$$

$$+ \frac{3Ca^5 - 5Ba^4b + 2Ca^3b^2 - 6Ba^2b^3 - Cab^4 - Bb^5 + 2(Ca^4b - 2Ba^3b^2 - 2Bab^4 - Cb^5) \tan(dx+c)}{2(a^2 + b^2)^3(b \tan(dx+c) + a)^2d}$$

input

```
integrate(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x, algorithm="giac")
```

output

```
(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) + 1/2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*log(tan(d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) - (C*a^3*b - 3*B*a^2*b^2 - 3*C*a*b^3 + B*b^4)*log(abs(b*tan(d*x + c) + a))/(a^6*b*d + 3*a^4*b^3*d + 3*a^2*b^5*d + b^7*d) + 1/2*(3*C*a^5 - 5*B*a^4*b + 2*C*a^3*b^2 - 6*B*a^2*b^3 - C*a*b^4 - B*b^5 + 2*(C*a^4*b - 2*B*a^3*b^2 - 2*B*a*b^4 - C*b^5)*tan(d*x + c))/((a^2 + b^2)^3*(b*tan(d*x + c) + a)^2*d)
```

Mupad [B] (verification not implemented)

Time = 5.49 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.59

$$\int \frac{\cot(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{\ln(a+b \tan(c+dx)) \left(\frac{3Bb-Ca}{(a^2+b^2)^2} - \frac{4b^2(Bb-Ca)}{(a^2+b^2)^3} \right)}{d}$$

$$- \frac{-3Ca^3+5Ba^2b+Cab^2+Bb^3}{2(a^4+2a^2b^2+b^4)} + \frac{\tan(c+dx)(-Ca^2b+2Bab^2+Cb^3)}{a^4+2a^2b^2+b^4}$$

$$+ \frac{\ln(\tan(c+dx)-i)(-C+Bi)}{2d(-a^3-a^2b3i+3ab^2+b^31i)} + \frac{\ln(\tan(c+dx)+i)(B-C1i)}{2d(-a^31i-3a^2b+ab^23i+b^3)}$$

input `int((cot(c + d*x)*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^3,x)`

output `(log(a + b*tan(c + d*x))*((3*B*b - C*a)/(a^2 + b^2)^2 - (4*b^2*(B*b - C*a))/(a^2 + b^2)^3))/d - ((B*b^3 - 3*C*a^3 + 5*B*a^2*b + C*a*b^2)/(2*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(C*b^3 + 2*B*a*b^2 - C*a^2*b))/(a^4 + b^4 + 2*a^2*b^2))/(d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*tan(c + d*x))) + (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(3*a*b^2 - a^2*b*3i - a^3 + b^3*1i)) + (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 1557, normalized size of antiderivative = 8.90

$$\int \frac{\cot(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `int(cot(d*x+c)*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)`

output

```
(4*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**5*b*c - 12*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**4*b**3 - 12*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**3*b**3*c + 4*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**2*b**5 - 4*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**5*b*c + 12*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**4*b**3 + 12*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**3*b**3*c - 4*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**2*b**5 + 4*cos(c + d*x)*sin(c + d*x)*a**5*b**2*d*x + 12*cos(c + d*x)*sin(c + d*x)*a**4*b**2*c*d*x - 12*cos(c + d*x)*sin(c + d*x)*a**3*b**4*d*x - 4*cos(c + d*x)*sin(c + d*x)*a**2*b**4*c*d*x - 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**6*c + 6*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**5*b**2 + 8*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**4*b**2*c - 8*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**3*b**4 - 6*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**2*b**4*c + 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a*b**6 + 2*log(tan((c + d*x)/2)**2 + 1)*a**6*c - 6*log(tan((c + d*x)/2)**2 + 1)*a**5*b**2 - 6*log(tan((c + d*x)/2)**2 + 1)*a**4*b**2*c + 2*log(tan((c + d*x)/2)**2 + 1)*a**3*b**4 + 2*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)**2*a**6*c - 6*log(tan((c + d*x)/2)**2*a - 2*...
```

3.43
$$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

Optimal result	483
Mathematica [C] (verified)	484
Rubi [A] (verified)	484
Maple [A] (verified)	489
Fricas [B] (verification not implemented)	490
Sympy [F(-2)]	490
Maxima [A] (verification not implemented)	491
Giac [A] (verification not implemented)	492
Mupad [B] (verification not implemented)	493
Reduce [B] (verification not implemented)	493

Optimal result

Integrand size = 40, antiderivative size = 215

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(3a^2bB - b^3B - a^3C + 3ab^2C)x}{(a^2 + b^2)^3} + \frac{B \log(\sin(c+dx))}{a^3d}$$

$$- \frac{b(6a^4bB + 3a^2b^3B + b^5B - 3a^5C + a^3b^2C) \log(a \cos(c+dx) + b \sin(c+dx))}{a^3(a^2 + b^2)^3d}$$

$$+ \frac{b(bB - aC)}{2a(a^2 + b^2)d(a + b \tan(c+dx))^2} + \frac{b(3a^2bB + b^3B - 2a^3C)}{a^2(a^2 + b^2)^2d(a + b \tan(c+dx))}$$

output

```
- (3*B*a^2*b - B*b^3 - C*a^3 + 3*C*a*b^2)*x / (a^2+b^2)^3 + B*ln(sin(d*x+c)) / a^3/d - b*(6*B*a^4*b + 3*B*a^2*b^3 + B*b^5 - 3*C*a^5 + C*a^3*b^2)*ln(a*cos(d*x+c) + b*sin(d*x+c)) / a^3 / (a^2+b^2)^3/d + 1/2*b*(B*b - C*a) / a / (a^2+b^2) / d / (a+b*tan(d*x+c))^2 + b*(3*B*a^2*b + B*b^3 - 2*C*a^3) / a^2 / (a^2+b^2)^2/d / (a+b*tan(d*x+c))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.10 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.04

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{-\frac{(B+iC) \log(i-\tan(c+dx))}{(a+ib)^3} + \frac{2B \log(\tan(c+dx))}{a^3} - \frac{(B-iC) \log(i+\tan(c+dx))}{(a-ib)^3} - \frac{2b(6a^4bB+3a^2b^3B+b^5B-3a^5C+a^3b^2C) \log(a+b \tan(c+dx))}{a^3(a^2+b^2)^3}}{2d}$$

input

```
Integrate[(Cot[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

```
(-(((B + I*C)*Log[I - Tan[c + d*x]])/(a + I*b)^3) + (2*B*Log[Tan[c + d*x]])/a^3 - ((B - I*C)*Log[I + Tan[c + d*x]])/(a - I*b)^3 - (2*b*(6*a^4*b*B + 3*a^2*b^3*B + b^5*B - 3*a^5*C + a^3*b^2*C)*Log[a + b*Tan[c + d*x]])/(a^3*(a^2 + b^2)^3) + (b*(b*B - a*C))/(a*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) + (2*b*(3*a^2*b*B + b^3*B - 2*a^3*C))/(a^2*(a^2 + b^2)^2*(a + b*Tan[c + d*x]))/(2*d)
```

Rubi [A] (verified)

Time = 2.44 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.20, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.325$, Rules used = {3042, 4115, 3042, 4092, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{\tan^2(c+dx)(a+b \tan(c+dx))^3} dx$$

$$\downarrow \text{4115}$$

$$\begin{aligned}
& \int \frac{\cot(c+dx)(B+C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{B+C \tan(c+dx)}{\tan(c+dx)(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow \text{4092} \\
& \frac{\int \frac{2 \cot(c+dx)(b(bB-aC) \tan^2(c+dx)-a(bB-aC) \tan(c+dx)+(a^2+b^2)B)}{(a+b \tan(c+dx))^2} dx}{\frac{2a(a^2+b^2)}{b(bB-aC)}} + \\
& \quad \frac{2ad(a^2+b^2)(a+b \tan(c+dx))^2}{\downarrow \text{27}} \\
& \frac{\int \frac{\cot(c+dx)(b(bB-aC) \tan^2(c+dx)-a(bB-aC) \tan(c+dx)+(a^2+b^2)B)}{(a+b \tan(c+dx))^2} dx}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \\
& \quad \frac{2ad(a^2+b^2)(a+b \tan(c+dx))^2}{\downarrow \text{3042}} \\
& \frac{\int \frac{b(bB-aC) \tan(c+dx)^2-a(bB-aC) \tan(c+dx)+(a^2+b^2)B}{\tan(c+dx)(a+b \tan(c+dx))^2} dx}{a(a^2+b^2)} + \frac{b(bB-aC)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow \text{4132} \\
& \frac{\int \frac{\cot(c+dx)\left(-((-Ca^2+2bBa+b^2C) \tan(c+dx)a^2)+b(-2Ca^3+3bBa^2+b^3B) \tan^2(c+dx)+(a^2+b^2)^2B\right)}{\frac{a+b \tan(c+dx)}{a(a^2+b^2)}} dx}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \frac{b(-2a^3C+3a^2bB+b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \\
& \quad \frac{2ad(a^2+b^2)(a+b \tan(c+dx))^2}{\downarrow \text{3042}} \\
& \frac{\int \frac{-((-Ca^2+2bBa+b^2C) \tan(c+dx)a^2)+b(-2Ca^3+3bBa^2+b^3B) \tan(c+dx)^2+(a^2+b^2)^2B}{\frac{\tan(c+dx)(a+b \tan(c+dx))}{a(a^2+b^2)}} dx}{\frac{a(a^2+b^2)}{b(bB-aC)}} + \frac{b(-2a^3C+3a^2bB+b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \\
& \quad \frac{2ad(a^2+b^2)(a+b \tan(c+dx))^2}{\downarrow \text{4134}}
\end{aligned}$$

$$\frac{\frac{B(a^2+b^2)^2 \int \cot(c+dx) dx}{a} - \frac{b(-3a^5C+6a^4bB+a^3b^2C+3a^2b^3B+b^5B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(a^3(-C)+3a^2bB+3ab^2C-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(-2a^3C+3a^2bB+3ab^2C-b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ \frac{b(bB - aC)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

3042

$$\frac{\frac{B(a^2+b^2)^2 \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b(-3a^5C+6a^4bB+a^3b^2C+3a^2b^3B+b^5B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(a^3(-C)+3a^2bB+3ab^2C-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(-2a^3C+3a^2bB+3ab^2C-b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ \frac{b(bB - aC)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

25

$$-\frac{\frac{B(a^2+b^2)^2 \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} - \frac{b(-3a^5C+6a^4bB+a^3b^2C+3a^2b^3B+b^5B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(a^3(-C)+3a^2bB+3ab^2C-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(-2a^3C+3a^2bB+3ab^2C-b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ \frac{b(bB - aC)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

3956

$$-\frac{\frac{b(-3a^5C+6a^4bB+a^3b^2C+3a^2b^3B+b^5B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{B(a^2+b^2)^2 \log(-\sin(c+dx))}{ad} - \frac{a^2x(a^3(-C)+3a^2bB+3ab^2C-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(-2a^3C+3a^2bB+3ab^2C-b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ \frac{b(bB - aC)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

4013

$$\frac{b(bB - aC)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} + \frac{\frac{B(a^2+b^2)^2 \log(-\sin(c+dx))}{ad} - \frac{a^2x(a^3(-C)+3a^2bB+3ab^2C-b^3B)}{a^2+b^2} - \frac{b(-3a^5C+6a^4bB+a^3b^2C+3a^2b^3B+b^5B) \log(a \cos(c+dx))}{ad(a^2+b^2)}}{a(a^2+b^2)} + \frac{b(-2a^3C+3a^2bB+b^3B)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

```
input Int[(Cot[c + d*x]^2*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3, x]
```

output

$$\frac{(b(bB - aC))/(2a(a^2 + b^2)d(a + b\tan[c + dx])^2) + (((a^2(3a^2bB - b^3B - a^3C + 3ab^2C)x)/(a^2 + b^2)) + ((a^2 + b^2)^2B\text{Log}[-\sin[c + dx]]/(ad) - (b(6a^4bB + 3a^2b^3B + b^5B - 3a^5C + a^3b^2C)\text{Log}[a\cos[c + dx] + b\sin[c + dx]]/(a(a^2 + b^2)d))/(a(a^2 + b^2)) + (b(3a^2bB + b^3B - 2a^3C))/(a(a^2 + b^2)d(a + b\tan[c + dx]))))/(a(a^2 + b^2))$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[Fx, x], x]$$

rule 27

$$\text{Int}[(a_)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{!MatchQ}[Fx, (b_)(Gx_)] \text{ ; FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\cos[c + dx], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4013

$$\text{Int}[(c_ + (d_)*\tan[(e_.) + (f_.)*(x_)])/((a_ + (b_)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a\cos[e + fx] + b\sin[e + fx], x]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$$

rule 4092

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2
)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4115

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_
.) + (f_.)*(x_)]^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ
[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 -
a*b*B + a^2*C, 0]

```

rule 4132

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] :> Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.13

method	result
derivativdivides	$\frac{b(3B a^2 b + B b^3 - 2C a^3)}{(a^2 + b^2)^2 a^2 (a + b \tan(dx + c))} - \frac{b(6B a^4 b + 3B a^2 b^3 + B b^5 - 3C a^5 + C a^3 b^2) \ln(a + b \tan(dx + c))}{(a^2 + b^2)^3 a^3} + \frac{(Bb - Ca)b}{2(a^2 + b^2) a (a + b \tan(dx + c))^2} + \frac{d}{d}$
default	$\frac{b(3B a^2 b + B b^3 - 2C a^3)}{(a^2 + b^2)^2 a^2 (a + b \tan(dx + c))} - \frac{b(6B a^4 b + 3B a^2 b^3 + B b^5 - 3C a^5 + C a^3 b^2) \ln(a + b \tan(dx + c))}{(a^2 + b^2)^3 a^3} + \frac{(Bb - Ca)b}{2(a^2 + b^2) a (a + b \tan(dx + c))^2} + \frac{d}{d}$
parallelrisc	$-12(a + b \tan(dx + c))^2 b (B a^4 b + \frac{1}{2} B a^2 b^3 + \frac{1}{6} B b^5 - \frac{1}{2} C a^5 + \frac{1}{6} C a^3 b^2) \ln(a + b \tan(dx + c)) - a^3 (a + b \tan(dx + c))^2 (B a^3 - \dots)$
norman	$-\frac{b(4B a^2 b^2 + 2B b^4 - 3C a^3 b - C a b^3) \tan(dx + c)^2}{d a^2 (a^4 + 2b^2 a^2 + b^4)} - \frac{b^2 (3B a^2 b - B b^3 - C a^3 + 3C a b^2) x \tan(dx + c)^3}{(a^4 + 2b^2 a^2 + b^4) (a^2 + b^2)} - \frac{b^2 (7B a^2 b^2 + 3B b^4 - 5C a^3 b - \dots)}{2d a^3 (a^4 + 2b^2 a^2 + b^4) \tan(dx + c) (a + b \tan(dx + c))}$
risc	$\frac{2i(3iB a^2 b^4 e^{2i(dx+c)} - 2iC a^3 b^3 e^{2i(dx+c)} - 2Ba b^5 e^{2i(dx+c)} + C a^2 b^4 e^{2i(dx+c)} - 4ib^4 B a^2 + 3iC a^3 b^3 - Ba b^5 + iB b^6 e^{2i(dx+c)})}{(ib+a)^2 (-ib e^{2i(dx+c)} + a e^{2i(dx+c)} + ib+a)^2 a^2 d (-ib \dots)}$

input

```
int(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method =_RETURNVERBOSE)
```

output

```
1/d*(b*(3*B*a^2*b+B*b^3-2*C*a^3)/(a^2+b^2)^2/a^2/(a+b*tan(d*x+c))-b*(6*B*a^4*b+3*B*a^2*b^3+B*b^5-3*C*a^5+C*a^3*b^2)/(a^2+b^2)^3/a^3*ln(a+b*tan(d*x+c)))+1/2*(B*b-C*a)*b/(a^2+b^2)/a/(a+b*tan(d*x+c))^2+1/(a^2+b^2)^3*(1/2*(-B*a^3+3*B*a*b^2-3*C*a^2*b+C*b^3)*ln(1+tan(d*x+c)^2)+(-3*B*a^2*b+B*b^3+C*a^3-3*C*a*b^2)*arctan(tan(d*x+c)))+1/a^3*B*ln(tan(d*x+c)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. $2(213) = 426$.

Time = 0.15 (sec) , antiderivative size = 683, normalized size of antiderivative = 3.18

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx =$$

$$\frac{7Ca^5b^3 - 9Ba^4b^4 + Ca^3b^5 - 3Ba^2b^6 - 2(Ca^8 - 3Ba^7b - 3Ca^6b^2 + Ba^5b^3)dx - (5Ca^5b^3 - 7Ba^4b^4 - 7Ca^4b^5 + 3Ba^3b^6 - 3Ca^2b^7 + Ba^2b^8)dx - (5Ca^5b^3 - 7Ba^4b^4 - 7Ca^4b^5 + 3Ba^3b^6 - 3Ca^2b^7 + Ba^2b^8)dx}{(a + b \tan(c + dx))^3}$$

input

```
integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="fricas")
```

output

```
-1/2*(7*C*a^5*b^3 - 9*B*a^4*b^4 + C*a^3*b^5 - 3*B*a^2*b^6 - 2*(C*a^8 - 3*B
*a^7*b - 3*C*a^6*b^2 + B*a^5*b^3)*d*x - (5*C*a^5*b^3 - 7*B*a^4*b^4 - C*a^3
*b^5 - B*a^2*b^6 + 2*(C*a^6*b^2 - 3*B*a^5*b^3 - 3*C*a^4*b^4 + B*a^3*b^5)*d
*x)*tan(d*x + c)^2 - (B*a^8 + 3*B*a^6*b^2 + 3*B*a^4*b^4 + B*a^2*b^6 + (B*a
^6*b^2 + 3*B*a^4*b^4 + 3*B*a^2*b^6 + B*b^8)*tan(d*x + c)^2 + 2*(B*a^7*b +
3*B*a^5*b^3 + 3*B*a^3*b^5 + B*a*b^7)*tan(d*x + c))*log(tan(d*x + c)^2/(tan
(d*x + c)^2 + 1)) - (3*C*a^7*b - 6*B*a^6*b^2 - C*a^5*b^3 - 3*B*a^4*b^4 - B
*a^2*b^6 + (3*C*a^5*b^3 - 6*B*a^4*b^4 - C*a^3*b^5 - 3*B*a^2*b^6 - B*b^8)*t
an(d*x + c)^2 + 2*(3*C*a^6*b^2 - 6*B*a^5*b^3 - C*a^4*b^4 - 3*B*a^3*b^5 - B
*a*b^7)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/
(tan(d*x + c)^2 + 1)) - 2*(3*C*a^6*b^2 - 4*B*a^5*b^3 - 3*C*a^4*b^4 + 3*B*a
^3*b^5 + B*a*b^7 + 2*(C*a^7*b - 3*B*a^6*b^2 - 3*C*a^5*b^3 + B*a^4*b^4)*d*x
)*tan(d*x + c))/((a^9*b^2 + 3*a^7*b^4 + 3*a^5*b^6 + a^3*b^8)*d*tan(d*x + c
)^2 + 2*(a^10*b + 3*a^8*b^3 + 3*a^6*b^5 + a^4*b^7)*d*tan(d*x + c) + (a^11
+ 3*a^9*b^2 + 3*a^7*b^4 + a^5*b^6)*d)
```

Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(cot(d*x+c)**2*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3, x)`

output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.73

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{2(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3)(dx+c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{2(3Ca^5b - 6Ba^4b^2 - Ca^3b^3 - 3Ba^2b^4 - Bb^6) \log(b \tan(dx+c)+a)}{a^9 + 3a^7b^2 + 3a^5b^4 + a^3b^6} - \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} \frac{1}{2d}$$

input `integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3, x, algorithm="maxima")`

output `1/2*(2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(3*C*a^5*b - 6*B*a^4*b^2 - C*a^3*b^3 - 3*B*a^2*b^4 - B*b^6)*log(b*tan(d*x + c) + a)/(a^9 + 3*a^7*b^2 + 3*a^5*b^4 + a^3*b^6) - (B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (5*C*a^4*b - 7*B*a^3*b^2 + C*a^2*b^3 - 3*B*a*b^4 + 2*(2*C*a^3*b^2 - 3*B*a^2*b^3 - B*b^5)*tan(d*x + c))/(a^8 + 2*a^6*b^2 + a^4*b^4 + (a^6*b^2 + 2*a^4*b^4 + a^2*b^6)*tan(d*x + c)^2 + 2*(a^7*b + 2*a^5*b^3 + a^3*b^5)*tan(d*x + c)) + 2*B*log(tan(d*x + c))/a^3)/d`

Giac [A] (verification not implemented)

Time = 0.85 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.73

$$\begin{aligned}
& \int \frac{\cot^2(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx \\
&= \frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3)(dx + c)}{a^6d + 3a^4b^2d + 3a^2b^4d + b^6d} \\
&\quad - \frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3) \log(\tan(dx + c)^2 + 1)}{2(a^6d + 3a^4b^2d + 3a^2b^4d + b^6d)} \\
&\quad + \frac{(3Ca^5b^2 - 6Ba^4b^3 - Ca^3b^4 - 3Ba^2b^5 - Bb^7) \log(|b \tan(dx + c) + a|)}{a^9bd + 3a^7b^3d + 3a^5b^5d + a^3b^7d} \\
&\quad + \frac{B \log(|\tan(dx + c)|)}{a^3d} \\
&\quad - \frac{5Ca^7b - 7Ba^6b^2 + 6Ca^5b^3 - 10Ba^4b^4 + Ca^3b^5 - 3Ba^2b^6 + 2(2Ca^6b^2 - 3Ba^5b^3 + 2Ca^4b^4 - 4Ba^3b^5 - Ba^2b^6 + Cab^7) \tan(dx + c)}{2(a^2 + b^2)^3(b \tan(dx + c) + a)^2 a^3 d}
\end{aligned}$$

input

```
integrate(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="giac")
```

output

```
(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d + 3
*a^2*b^4*d + b^6*d) - 1/2*(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*log(tan(
d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) + (3*C*a^5*b^2
- 6*B*a^4*b^3 - C*a^3*b^4 - 3*B*a^2*b^5 - B*b^7)*log(abs(b*tan(d*x + c) +
a))/(a^9*b*d + 3*a^7*b^3*d + 3*a^5*b^5*d + a^3*b^7*d) + B*log(abs(tan(d*x
+ c)))/(a^3*d) - 1/2*(5*C*a^7*b - 7*B*a^6*b^2 + 6*C*a^5*b^3 - 10*B*a^4*b^
4 + C*a^3*b^5 - 3*B*a^2*b^6 + 2*(2*C*a^6*b^2 - 3*B*a^5*b^3 + 2*C*a^4*b^4 -
4*B*a^3*b^5 - B*a*b^7)*tan(d*x + c))/((a^2 + b^2)^3*(b*tan(d*x + c) + a)^
2*a^3*d)
```

Mupad [B] (verification not implemented)

Time = 7.58 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.47

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{-5C a^3 b + 7B a^2 b^2 - C a b^3 + 3B b^4}{2a(a^4 + 2a^2 b^2 + b^4)} + \frac{\tan(c+dx)(-2C a^3 b^2 + 3B a^2 b^3 + B b^5)}{a^2(a^4 + 2a^2 b^2 + b^4)} + \frac{B \ln(\tan(c+dx))}{a^3 d}$$

$$+ \frac{\ln(\tan(c+dx) - i)(-C + B i)}{2d(-a^3 i + 3a^2 b + a b^2 3i - b^3)} + \frac{\ln(\tan(c+dx) + i)(B - C i)}{2d(-a^3 + a^2 b 3i + 3a b^2 - b^3 i)}$$

$$- \frac{b \ln(a + b \tan(c+dx))(-3C a^5 + 6B a^4 b + C a^3 b^2 + 3B a^2 b^3 + B b^5)}{a^3 d(a^2 + b^2)^3}$$

input

```
int((cot(c + d*x)^2*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^3,x)
```

output

```
((3*B*b^4 + 7*B*a^2*b^2 - C*a*b^3 - 5*C*a^3*b)/(2*a*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(B*b^5 + 3*B*a^2*b^3 - 2*C*a^3*b^2))/(a^2*(a^4 + b^4 + 2*a^2*b^2)))/(d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*tan(c + d*x))) + (B*log(tan(c + d*x)))/(a^3*d) + (log(tan(c + d*x) - 1i)*(B*1i - C))/(2*d*(a*b^2*3i + 3*a^2*b - a^3*1i - b^3)) + (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(3*a*b^2 + a^2*b*3i - a^3 - b^3*1i)) - (b*log(a + b*tan(c + d*x))*(B*b^5 - 3*C*a^5 + 3*B*a^2*b^3 + C*a^3*b^2 + 6*B*a^4*b))/(a^3*d*(a^2 + b^2)^3)
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 2015, normalized size of antiderivative = 9.37

$$\int \frac{\cot^2(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Too large to display}$$

input

```
int(cot(d*x+c)^2*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)
```

output

```
( - 4*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**7*b**2 - 1
2*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**6*b**2*c + 12*
cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**5*b**4 + 4*cos(c
+ d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)*a**4*b**4*c + 12*cos(c +
d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a
**6*b**2*c - 24*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2
)*b - a)*sin(c + d*x)*a**5*b**4 - 4*cos(c + d*x)*log(tan((c + d*x)/2)**2*a
- 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**4*b**4*c - 12*cos(c + d*x)*lo
g(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)*a**3*b**6
- 4*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*si
n(c + d*x)*a*b**8 + 4*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**7
*b**2 + 12*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**5*b**4 + 12*
cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)*a**3*b**6 + 4*cos(c + d*x)
*log(tan((c + d*x)/2))*sin(c + d*x)*a*b**8 + 4*cos(c + d*x)*sin(c + d*x)*a
**7*b*c*d*x - 12*cos(c + d*x)*sin(c + d*x)*a**6*b**3*d*x - 12*cos(c + d*x)
*sin(c + d*x)*a**5*b**3*c*d*x + 4*cos(c + d*x)*sin(c + d*x)*a**4*b**5*d*x
+ 2*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**8*b + 6*log(tan((c + d
*x)/2)**2 + 1)*sin(c + d*x)**2*a**7*b*c - 8*log(tan((c + d*x)/2)**2 + 1)*s
in(c + d*x)**2*a**6*b**3 - 8*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*
a**5*b**3*c + 6*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**4*b**5 ...
```

3.44
$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

Optimal result	495
Mathematica [C] (verified)	496
Rubi [A] (verified)	496
Maple [A] (verified)	502
Fricas [B] (verification not implemented)	502
Sympy [F(-2)]	503
Maxima [A] (verification not implemented)	504
Giac [A] (verification not implemented)	504
Mupad [B] (verification not implemented)	505
Reduce [B] (verification not implemented)	506

Optimal result

Integrand size = 40, antiderivative size = 287

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(a^3B - 3ab^2B + 3a^2bC - b^3C)x}{(a^2 + b^2)^3} - \frac{(3bB - aC) \log(\sin(c+dx))}{a^4d}$$

$$+ \frac{b^2(10a^4bB + 9a^2b^3B + 3b^5B - 6a^5C - 3a^3b^2C - ab^4C) \log(a \cos(c+dx) + b \sin(c+dx))}{a^4(a^2 + b^2)^3d}$$

$$- \frac{b(2a^2B + 3b^2B - abC)}{2a^2(a^2 + b^2)d(a+b \tan(c+dx))^2} - \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

$$- \frac{b(a^4B + 6a^2b^2B + 3b^4B - 3a^3bC - ab^3C)}{a^3(a^2 + b^2)^2d(a+b \tan(c+dx))}$$

output

```
-(B*a^3-3*B*a*b^2+3*C*a^2*b-C*b^3)*x/(a^2+b^2)^3-(3*B*b-C*a)*ln(sin(d*x+c)
)/a^4/d+b^2*(10*B*a^4*b+9*B*a^2*b^3+3*B*b^5-6*C*a^5-3*C*a^3*b^2-C*a*b^4)*l
n(a*cos(d*x+c)+b*sin(d*x+c))/a^4/(a^2+b^2)^3/d-1/2*b*(2*B*a^2+3*B*b^2-C*a*
b)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^2-B*cot(d*x+c)/a/d/(a+b*tan(d*x+c))^2-
b*(B*a^4+6*B*a^2*b^2+3*B*b^4-3*C*a^3*b-C*a*b^3)/a^3/(a^2+b^2)^2/d/(a+b*tan
(d*x+c))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.27 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.00

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{B \cot(c+dx)}{a^3 d} + \frac{(B+iC) \log(i-\tan(c+dx))}{2(ia-b)^3 d}$$

$$- \frac{(3bB-aC) \log(\tan(c+dx))}{a^4 d} - \frac{(iB+C) \log(i+\tan(c+dx))}{2(a-ib)^3 d}$$

$$+ \frac{b^2(10a^4bB + 9a^2b^3B + 3b^5B - 6a^5C - 3a^3b^2C - ab^4C) \log(a+b \tan(c+dx))}{a^4(a^2+b^2)^3 d}$$

$$- \frac{b^2(bB-aC)}{2a^2(a^2+b^2)d(a+b \tan(c+dx))^2} - \frac{b^2(4a^2bB + 2b^3B - 3a^3C - ab^2C)}{a^3(a^2+b^2)^2 d(a+b \tan(c+dx))}$$

input `Integrate[(Cot[c + d*x]^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]`

output `-((B*Cot[c + d*x])/(a^3*d)) + ((B + I*C)*Log[I - Tan[c + d*x]])/(2*(I*a - b)^3*d) - ((3*b*B - a*C)*Log[Tan[c + d*x]])/(a^4*d) - ((I*B + C)*Log[I + Tan[c + d*x]])/(2*(a - I*b)^3*d) + (b^2*(10*a^4*b*B + 9*a^2*b^3*B + 3*b^5*B - 6*a^5*C - 3*a^3*b^2*C - a*b^4*C)*Log[a + b*Tan[c + d*x]])/(a^4*(a^2 + b^2)^3*d) - (b^2*(b*B - a*C))/(2*a^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (b^2*(4*a^2*b*B + 2*b^3*B - 3*a^3*C - a*b^2*C))/(a^3*(a^2 + b^2)^2*d*(a + b*Tan[c + d*x]))`

Rubi [A] (verified)

Time = 3.19 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.16, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 4115, 3042, 4092, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\cot^3(c+dx)(B \tan(c+dx) + C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 3042 \\
& \int \frac{B \tan(c+dx) + C \tan^2(c+dx)}{\tan(c+dx)^3(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 4115 \\
& \int \frac{\cot^2(c+dx)(B + C \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 3042 \\
& \int \frac{B + C \tan(c+dx)}{\tan(c+dx)^2(a+b \tan(c+dx))^3} dx \\
& \quad \downarrow 4092 \\
& - \frac{\int \frac{\cot(c+dx)(3bB \tan^2(c+dx) + aB \tan(c+dx) + 3bB - aC)}{(a+b \tan(c+dx))^3} dx}{a} - \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& - \frac{\int \frac{3bB \tan(c+dx)^2 + aB \tan(c+dx) + 3bB - aC}{\tan(c+dx)(a+b \tan(c+dx))^3} dx}{a} - \frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \\
& \quad \downarrow 4132 \\
& - \frac{\int \frac{2 \cot(c+dx)((aB+bC) \tan(c+dx)a^2 + b(2Ba^2 - bCa + 3b^2B) \tan^2(c+dx) + (a^2+b^2)(3bB - aC))}{(a+b \tan(c+dx))^2} dx}{2a(a^2+b^2)} + \frac{b(2a^2B - abC + 3b^2B)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow \\
& \frac{a}{ad(a+b \tan(c+dx))^2} \\
& \quad \downarrow 27 \\
& - \frac{\int \frac{\cot(c+dx)((aB+bC) \tan(c+dx)a^2 + b(2Ba^2 - bCa + 3b^2B) \tan^2(c+dx) + (a^2+b^2)(3bB - aC))}{(a+b \tan(c+dx))^2} dx}{a(a^2+b^2)} + \frac{b(2a^2B - abC + 3b^2B)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
& \quad \downarrow \\
& \frac{a}{ad(a+b \tan(c+dx))^2} \\
& \quad \downarrow 3042
\end{aligned}$$

$$\int \frac{(aB+bC) \tan(c+dx)a^2+b(2Ba^2-bCa+3b^2B) \tan(c+dx)^2+(a^2+b^2)(3bB-aC)}{\tan(c+dx)(a+b \tan(c+dx))^2} dx + \frac{b(2a^2B-abC+3b^2B)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

4132

$$\int \frac{\cot(c+dx) \left((Ba^2+2bCa-b^2B) \tan(c+dx)a^3+b(Ba^4-3bCa^3+6b^2Ba^2-b^3Ca+3b^4B) \tan^2(c+dx)+(a^2+b^2)^2(3bB-aC) \right)}{a+b \tan(c+dx)} dx + \frac{b(a^4B-3a^3bC+6a^2b^2B-ab^3C+3b^4B)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

3042

$$\int \frac{(Ba^2+2bCa-b^2B) \tan(c+dx)a^3+b(Ba^4-3bCa^3+6b^2Ba^2-b^3Ca+3b^4B) \tan(c+dx)^2+(a^2+b^2)^2(3bB-aC)}{\tan(c+dx)(a+b \tan(c+dx))} dx + \frac{b(a^4B-3a^3bC+6a^2b^2B-ab^3C+3b^4B)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

4134

$$\frac{(a^2+b^2)^2(3bB-aC) \int \cot(c+dx) dx}{a} - \frac{b^2(-6a^5C+10a^4bB-3a^3b^2C+9a^2b^3B-ab^4C+3b^5B)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{a^3x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2} + \frac{b(a^4B-3a^3bC+6a^2b^2B-ab^3C+3b^4B)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

3042

$$\frac{(a^2+b^2)^2(3bB-aC) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b^2(-6a^5C+10a^4bB-3a^3b^2C+9a^2b^3B-ab^4C+3b^5B)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{a^3x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

25

$$\frac{\frac{(a^2+b^2)^2(3bB-aC) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} - \frac{b^2(-6a^5C+10a^4bB-3a^3b^2C+9a^2b^3B-ab^4C+3b^5B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a(a^2+b^2)}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

↓ 3956

$$\frac{\frac{b^2(-6a^5C+10a^4bB-3a^3b^2C+9a^2b^3B-ab^4C+3b^5B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2+b^2)^2(3bB-aC) \log(-\sin(c+dx))}{ad} + \frac{a^3x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a(a^2+b^2)}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

↓ 4013

$$\frac{\frac{b(2a^2B-abC+3b^2B)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(a^4B-3a^3bC+6a^2b^2B-ab^3C+3b^4B)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{(a^2+b^2)^2(3bB-aC) \log(-\sin(c+dx))}{ad} + \frac{a^3x(a^3B+3a^2bC-3ab^2B-b^3C)}{a^2+b^2}}{a(a^2+b^2)}$$

$$\frac{B \cot(c+dx)}{ad(a+b \tan(c+dx))^2}$$

input

```
Int[((Cot[c + d*x])^3*(B*Tan[c + d*x] + C*Tan[c + d*x]^2))/(a + b*Tan[c + d*x])^3,x]
```

output

```
-((B*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^2)) - ((b*(2*a^2*B + 3*b^2*B - a*b*C))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((a^3*(a^3*B - 3*a*b^2*B + 3*a^2*b*C - b^3*C)*x)/(a^2 + b^2) + ((a^2 + b^2)^2*(3*b*B - a*C)*Log[-Sin[c + d*x]])/(a*d) - (b^2*(10*a^4*b*B + 9*a^2*b^3*B + 3*b^5*B - 6*a^5*C - 3*a^3*b^2*C - a*b^4*C)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(a^4*B + 6*a^2*b^2*B + 3*b^4*B - 3*a^3*b*C - a*b^3*C))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a*(a^2 + b^2))/a
```


Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`
- rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4115

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[1/b^2 Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*(b*B - a*C + b*C*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[A*b^2 - a*b*B + a^2*C, 0]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.01

method	result
derivativedivides	$-\frac{b^2(4B a^2 b+2B b^3-3C a^3-C a b^2)}{(a^2+b^2)^2 a^3(a+b \tan(dx+c))} + \frac{b^2(10B a^4 b+9B a^2 b^3+3B b^5-6C a^5-3C a^3 b^2-C a b^4) \ln(a+b \tan(dx+c))}{(a^2+b^2)^3 a^4} - \frac{(B b^2)}{2(a^2+b^2) a^2}$
default	$-\frac{b^2(4B a^2 b+2B b^3-3C a^3-C a b^2)}{(a^2+b^2)^2 a^3(a+b \tan(dx+c))} + \frac{b^2(10B a^4 b+9B a^2 b^3+3B b^5-6C a^5-3C a^3 b^2-C a b^4) \ln(a+b \tan(dx+c))}{(a^2+b^2)^3 a^4} - \frac{(B b^2)}{2(a^2+b^2) a^2}$
parallelrisch	$20(B a^4 b+\frac{9}{10} B a^2 b^3+\frac{3}{10} B b^5-\frac{3}{5} C a^5-\frac{3}{10} C a^3 b^2-\frac{1}{10} C a b^4)(a+b \tan(dx+c))^2 b^2 \ln(a+b \tan(dx+c))+3 a^4(a+b \tan(dx+c))$
norman	$\frac{b(3B a^4 b+11B a^2 b^3+6B b^5-4C a^3 b^2-2C a b^4) \tan(dx+c)^3}{d a^3(a^4+2b^2 a^2+b^4)} - \frac{B \tan(dx+c)}{a d} + \frac{b^2(4B a^4 b+17B a^2 b^3+9B b^5-7C a^3 b^2-3C a b^4) \tan(dx+c)}{2 a^4 d(a^4+2b^2 a^2+b^4)}$
risch	Expression too large to display

input `int(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,method =_RETURNVERBOSE)`

output `1/d*(-b^2*(4*B*a^2*b+2*B*b^3-3*C*a^3-C*a*b^2)/(a^2+b^2)^2/a^3/(a+b*tan(d*x+c))+b^2*(10*B*a^4*b+9*B*a^2*b^3+3*B*b^5-6*C*a^5-3*C*a^3*b^2-C*a*b^4)/(a^2+b^2)^3/a^4*ln(a+b*tan(d*x+c))-1/2*(B*b-C*a)*b^2/(a^2+b^2)/a^2/(a+b*tan(d*x+c))^2+1/(a^2+b^2)^3*(1/2*(3*B*a^2*b-B*b^3-C*a^3+3*C*a*b^2)*ln(1+tan(d*x+c)^2)+(-B*a^3+3*B*a*b^2-3*C*a^2*b+C*b^3)*arctan(tan(d*x+c)))-1/a^3*B/tan(d*x+c)+(-3*B*b+C*a)/a^4*ln(tan(d*x+c)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 917 vs. 2(283) = 566.

Time = 0.20 (sec) , antiderivative size = 917, normalized size of antiderivative = 3.20

$$\int \frac{\cot^3(c+dx)(B \tan(c+dx)+C \tan^2(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="fricas")`

output `-1/2*(2*B*a^9 + 6*B*a^7*b^2 + 6*B*a^5*b^4 + 2*B*a^3*b^6 + (7*C*a^5*b^4 - 9
*B*a^4*b^5 + C*a^3*b^6 - 3*B*a^2*b^7 + 2*(B*a^7*b^2 + 3*C*a^6*b^3 - 3*B*a^5
*b^4 - C*a^4*b^5)*d*x)*tan(d*x + c)^3 + 2*(B*a^7*b^2 + 4*C*a^6*b^3 - 2*B*
a^5*b^4 - 3*C*a^4*b^5 + 6*B*a^3*b^6 - C*a^2*b^7 + 3*B*a*b^8 + 2*(B*a^8*b +
3*C*a^7*b^2 - 3*B*a^6*b^3 - C*a^5*b^4)*d*x)*tan(d*x + c)^2 - ((C*a^7*b^2
- 3*B*a^6*b^3 + 3*C*a^5*b^4 - 9*B*a^4*b^5 + 3*C*a^3*b^6 - 9*B*a^2*b^7 + C*
a*b^8 - 3*B*b^9)*tan(d*x + c)^3 + 2*(C*a^8*b - 3*B*a^7*b^2 + 3*C*a^6*b^3 -
9*B*a^5*b^4 + 3*C*a^4*b^5 - 9*B*a^3*b^6 + C*a^2*b^7 - 3*B*a*b^8)*tan(d*x
+ c)^2 + (C*a^9 - 3*B*a^8*b + 3*C*a^7*b^2 - 9*B*a^6*b^3 + 3*C*a^5*b^4 - 9*
B*a^4*b^5 + C*a^3*b^6 - 3*B*a^2*b^7)*tan(d*x + c))*log(tan(d*x + c)^2/(tan
(d*x + c)^2 + 1)) + ((6*C*a^5*b^4 - 10*B*a^4*b^5 + 3*C*a^3*b^6 - 9*B*a^2*b
^7 + C*a*b^8 - 3*B*b^9)*tan(d*x + c)^3 + 2*(6*C*a^6*b^3 - 10*B*a^5*b^4 + 3
*C*a^4*b^5 - 9*B*a^3*b^6 + C*a^2*b^7 - 3*B*a*b^8)*tan(d*x + c)^2 + (6*C*a^7
*b^2 - 10*B*a^6*b^3 + 3*C*a^5*b^4 - 9*B*a^4*b^5 + C*a^3*b^6 - 3*B*a^2*b^7
)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d
*x + c)^2 + 1)) + (4*B*a^8*b + 12*B*a^6*b^3 - 9*C*a^5*b^4 + 23*B*a^4*b^5 -
3*C*a^3*b^6 + 9*B*a^2*b^7 + 2*(B*a^9 + 3*C*a^8*b - 3*B*a^7*b^2 - C*a^6*b^8
3)*d*x)*tan(d*x + c))/((a^10*b^2 + 3*a^8*b^4 + 3*a^6*b^6 + a^4*b^8)*d*tan(
d*x + c)^3 + 2*(a^11*b + 3*a^9*b^3 + 3*a^7*b^5 + a^5*b^7)*d*tan(d*x + c)^2
+ (a^12 + 3*a^10*b^2 + 3*a^8*b^4 + a^6*b^6)*d*tan(d*x + c))`

Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(cot(d*x+c)**3*(B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))**3
,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 454, normalized size of antiderivative = 1.58

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx =$$

$$-\frac{2(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)(dx + c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{2(6Ca^5b^2 - 10Ba^4b^3 + 3Ca^3b^4 - 9Ba^2b^5 + Cab^6 - 3Bb^7) \log(b \tan(dx + c) + a)}{a^{10} + 3a^8b^2 + 3a^6b^4 + a^4b^6} + \frac{(Ca^3 - 3Ba^2b)}{2(a^2 + b^2)^3 (b \tan(dx + c) + a)}$$

input

```
integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="maxima")
```

output

```
-1/2*(2*(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 +
+ 3*a^2*b^4 + b^6) + 2*(6*C*a^5*b^2 - 10*B*a^4*b^3 + 3*C*a^3*b^4 - 9*B*a^
2*b^5 + C*a*b^6 - 3*B*b^7)*log(b*tan(d*x + c) + a)/(a^10 + 3*a^8*b^2 + 3*a
^6*b^4 + a^4*b^6) + (C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*log(tan(d*x +
c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (2*B*a^6 + 4*B*a^4*b^2 + 2
*B*a^2*b^4 + 2*(B*a^4*b^2 - 3*C*a^3*b^3 + 6*B*a^2*b^4 - C*a*b^5 + 3*B*b^6)
*tan(d*x + c)^2 + (4*B*a^5*b - 7*C*a^4*b^2 + 17*B*a^3*b^3 - 3*C*a^2*b^4 +
9*B*a*b^5)*tan(d*x + c))/((a^7*b^2 + 2*a^5*b^4 + a^3*b^6)*tan(d*x + c)^3 +
2*(a^8*b + 2*a^6*b^3 + a^4*b^5)*tan(d*x + c)^2 + (a^9 + 2*a^7*b^2 + a^5*b
^4)*tan(d*x + c)) - 2*(C*a - 3*B*b)*log(tan(d*x + c))/a^4)/d
```

Giac [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 466, normalized size of antiderivative = 1.62

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= -\frac{(Ba^3 + 3Ca^2b - 3Bab^2 - Cb^3)(dx + c)}{a^6d + 3a^4b^2d + 3a^2b^4d + b^6d}$$

$$-\frac{(Ca^3 - 3Ba^2b - 3Cab^2 + Bb^3) \log(\tan(dx + c)^2 + 1)}{2(a^6d + 3a^4b^2d + 3a^2b^4d + b^6d)}$$

$$-\frac{(6Ca^5b^3 - 10Ba^4b^4 + 3Ca^3b^5 - 9Ba^2b^6 + Cab^7 - 3Bb^8) \log(|b \tan(dx + c) + a|)}{a^{10}bd + 3a^8b^3d + 3a^6b^5d + a^4b^7d}$$

$$+ \frac{(Ca - 3Bb) \log(|\tan(dx + c)|)}{a^4d}$$

$$-\frac{2Ba^9 + 6Ba^7b^2 + 6Ba^5b^4 + 2Ba^3b^6 + 2(Ba^7b^2 - 3Ca^6b^3 + 7Ba^5b^4 - 4Ca^4b^5 + 9Ba^3b^6 - Ca^2b^7 - 2(a^2 + b^2)^3 (b \tan(dx + c)))}{2(a^2 + b^2)^3 (b \tan(dx + c))}$$

input `integrate(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x,
algorithm="giac")`

output
$$-(B*a^3 + 3*C*a^2*b - 3*B*a*b^2 - C*b^3)*(d*x + c)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) - 1/2*(C*a^3 - 3*B*a^2*b - 3*C*a*b^2 + B*b^3)*\log(\tan(d*x + c)^2 + 1)/(a^6*d + 3*a^4*b^2*d + 3*a^2*b^4*d + b^6*d) - (6*C*a^5*b^3 - 10*B*a^4*b^4 + 3*C*a^3*b^5 - 9*B*a^2*b^6 + C*a*b^7 - 3*B*b^8)*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^{10}*b*d + 3*a^8*b^3*d + 3*a^6*b^5*d + a^4*b^7*d) + (C*a - 3*B*b)*\log(\text{abs}(\tan(d*x + c)))/(a^4*d) - 1/2*(2*B*a^9 + 6*B*a^7*b^2 + 6*B*a^5*b^4 + 2*B*a^3*b^6 + 2*(B*a^7*b^2 - 3*C*a^6*b^3 + 7*B*a^5*b^4 - 4*C*a^4*b^5 + 9*B*a^3*b^6 - C*a^2*b^7 + 3*B*a*b^8)*\tan(d*x + c)^2 + (4*B*a^8*b - 7*C*a^7*b^2 + 21*B*a^6*b^3 - 10*C*a^5*b^4 + 26*B*a^4*b^5 - 3*C*a^3*b^6 + 9*B*a^2*b^7)*\tan(d*x + c))/((a^2 + b^2)^3*(b*\tan(d*x + c) + a)^2*a^4*d*\tan(d*x + c))$$

Mupad [B] (verification not implemented)

Time = 10.39 (sec) , antiderivative size = 380, normalized size of antiderivative = 1.32

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{b^2 \ln(a + b \tan(c + dx)) (-6 C a^5 + 10 B a^4 b - 3 C a^3 b^2 + 9 B a^2 b^3 - C a b^4 + 3 B b^5)}{a^4 d (a^2 + b^2)^3}$$

$$- \frac{\ln(\tan(c + dx) - i) (-C + B i)}{2 d (-a^3 - a^2 b 3i + 3 a b^2 + b^3 i)}$$

$$- \frac{\ln(\tan(c + dx)) (3 B b - C a)}{a^4 d} - \frac{\ln(\tan(c + dx) + i) (B - C i)}{2 d (-a^3 i - 3 a^2 b + a b^2 3i + b^3)}$$

$$- \frac{\frac{B}{a} + \frac{\tan(c+dx)^2 (B a^4 b^2 - 3 C a^3 b^3 + 6 B a^2 b^4 - C a b^5 + 3 B b^6)}{a^3 (a^4 + 2 a^2 b^2 + b^4)}}{d (a^2 \tan(c + dx) + 2 a b \tan(c + dx)^2 + b^2 \tan(c + dx)^3)} + \frac{\tan(c+dx) (4 B a^4 b - 7 C a^3 b^2 + 17 B a^2 b^3 - 3 C a b^4 + 9 B b^5)}{2 a^2 (a^4 + 2 a^2 b^2 + b^4)}$$

input `int((cot(c + d*x)^3*(B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^3,x)`

output

```
(b^2*log(a + b*tan(c + d*x))*(3*B*b^5 - 6*C*a^5 + 9*B*a^2*b^3 - 3*C*a^3*b^2 + 10*B*a^4*b - C*a*b^4))/(a^4*d*(a^2 + b^2)^3) - (log(tan(c + d*x) - 1i) * (B*1i - C))/(2*d*(3*a*b^2 - a^2*b*3i - a^3 + b^3*1i)) - (log(tan(c + d*x) * (3*B*b - C*a))/(a^4*d) - (log(tan(c + d*x) + 1i)*(B - C*1i))/(2*d*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3)) - (B/a + (tan(c + d*x)^2*(3*B*b^6 + 6*B*a^2*b^4 + B*a^4*b^2 - 3*C*a^3*b^3 - C*a*b^5))/(a^3*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(9*B*b^5 + 17*B*a^2*b^3 - 7*C*a^3*b^2 + 4*B*a^4*b - 3*C*a*b^4))/(2*a^2*(a^4 + b^4 + 2*a^2*b^2)))/(d*(a^2*tan(c + d*x) + b^2*tan(c + d*x)^3 + 2*a*b*tan(c + d*x)^2))
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 2889, normalized size of antiderivative = 10.07

$$\int \frac{\cot^3(c + dx) (B \tan(c + dx) + C \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input

```
int(cot(d*x+c)^3*(B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^3,x)
```

output

```
( - 16*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**8*b*c
+ 48*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**7*b**3 +
48*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**6*b**3*c
- 16*cos(c + d*x)*log(tan((c + d*x)/2)**2 + 1)*sin(c + d*x)**2*a**5*b**5 -
96*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin
(c + d*x)**2*a**6*b**3*c + 160*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*
tan((c + d*x)/2)*b - a)*sin(c + d*x)**2*a**5*b**5 - 48*cos(c + d*x)*log(ta
n((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)**2*a**4*b**5*
c + 144*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)
*sin(c + d*x)**2*a**3*b**7 - 16*cos(c + d*x)*log(tan((c + d*x)/2)**2*a - 2
*tan((c + d*x)/2)*b - a)*sin(c + d*x)**2*a**2*b**7*c + 48*cos(c + d*x)*log
(tan((c + d*x)/2)**2*a - 2*tan((c + d*x)/2)*b - a)*sin(c + d*x)**2*a*b**9
+ 16*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)**2*a**8*b*c - 48*cos(
c + d*x)*log(tan((c + d*x)/2))*sin(c + d*x)**2*a**7*b**3 + 48*cos(c + d*x)
*log(tan((c + d*x)/2))*sin(c + d*x)**2*a**6*b**3*c - 144*cos(c + d*x)*log(
tan((c + d*x)/2))*sin(c + d*x)**2*a**5*b**5 + 48*cos(c + d*x)*log(tan((c +
d*x)/2))*sin(c + d*x)**2*a**4*b**5*c - 144*cos(c + d*x)*log(tan((c + d*x)
/2))*sin(c + d*x)**2*a**3*b**7 + 16*cos(c + d*x)*log(tan((c + d*x)/2))*sin
(c + d*x)**2*a**2*b**7*c - 48*cos(c + d*x)*log(tan((c + d*x)/2))*sin(c + d
*x)**2*a*b**9 + 2*cos(c + d*x)*sin(c + d*x)**2*a**9*b - 16*cos(c + d*x)...
```


3.45 $\int \tan^2(c+dx)(b \tan(c+dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	508
Mathematica [A] (verified)	509
Rubi [A] (verified)	509
Maple [F]	512
Fricas [F]	512
Sympy [F]	512
Maxima [F]	513
Giac [F]	513
Mupad [F(-1)]	514
Reduce [F]	514

Optimal result

Integrand size = 39, antiderivative size = 132

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{C(b \tan(c + dx))^{3+n}}{b^3 d(3 + n)}$$

$$+ \frac{(A - C) \operatorname{Hypergeometric2F1}\left(1, \frac{3+n}{2}, \frac{5+n}{2}, -\tan^2(c + dx)\right) (b \tan(c + dx))^{3+n}}{b^3 d(3 + n)}$$

$$+ \frac{B \operatorname{Hypergeometric2F1}\left(1, \frac{4+n}{2}, \frac{6+n}{2}, -\tan^2(c + dx)\right) (b \tan(c + dx))^{4+n}}{b^4 d(4 + n)}$$

output

```
C*(b*tan(d*x+c))^(3+n)/b^3/d/(3+n)+(A-C)*hypergeom([1, 3/2+1/2*n], [5/2+1/2*n], -tan(d*x+c)^2)*(b*tan(d*x+c))^(3+n)/b^3/d/(3+n)+B*hypergeom([1, 2+1/2*n], [3+1/2*n], -tan(d*x+c)^2)*(b*tan(d*x+c))^(4+n)/b^4/d/(4+n)
```

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.83

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\tan^3(c + dx)(b \tan(c + dx))^n (C(4 + n) + (A - C)(4 + n) \operatorname{Hypergeometric2F1}\left(1, \frac{3+n}{2}, \frac{5+n}{2}, -\tan^2(c + dx)\right) + B(3 + n) \operatorname{Hypergeometric2F1}\left(1, \frac{4+n}{2}, \frac{6+n}{2}, -\tan^2(c + dx)\right)}{d(3 + n)(4 + n)}$$

input

```
Integrate[Tan[c + d*x]^2*(b*Tan[c + d*x])^n*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(Tan[c + d*x]^3*(b*Tan[c + d*x])^n*(C*(4 + n) + (A - C)*(4 + n)*Hypergeometric2F1[1, (3 + n)/2, (5 + n)/2, -Tan[c + d*x]^2] + B*(3 + n)*Hypergeometric2F1[1, (4 + n)/2, (6 + n)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(d*(3 + n)*(4 + n))
```

Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.205$, Rules used = {2030, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow \text{2030}$$

$$\frac{\int (b \tan(c + dx))^{n+2} (C \tan^2(c + dx) + B \tan(c + dx) + A) dx}{b^2}$$

$$\downarrow \text{3042}$$

$$\frac{\int (b \tan(c + dx))^{n+2} (C \tan(c + dx)^2 + B \tan(c + dx) + A) dx}{b^2}$$

$$\downarrow \text{4113}$$

$$\begin{aligned}
 & \frac{\int (b \tan(c + dx))^{n+2} (A - C + B \tan(c + dx)) dx + \frac{C(b \tan(c + dx))^{n+3}}{bd(n+3)}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int (b \tan(c + dx))^{n+2} (A - C + B \tan(c + dx)) dx + \frac{C(b \tan(c + dx))^{n+3}}{bd(n+3)}}{b^2} \\
 & \quad \downarrow \text{4021} \\
 & \frac{(A - C) \int (b \tan(c + dx))^{n+2} dx + \frac{B \int (b \tan(c + dx))^{n+3} dx}{b} + \frac{C(b \tan(c + dx))^{n+3}}{bd(n+3)}}{b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - C) \int (b \tan(c + dx))^{n+2} dx + \frac{B \int (b \tan(c + dx))^{n+3} dx}{b} + \frac{C(b \tan(c + dx))^{n+3}}{bd(n+3)}}{b^2} \\
 & \quad \downarrow \text{3957} \\
 & \frac{\frac{b(A - C) \int \frac{(b \tan(c + dx))^{n+2}}{\tan^2(c + dx)b^2 + b^2} d(b \tan(c + dx))}{d} + \frac{B \int \frac{(b \tan(c + dx))^{n+3}}{\tan^2(c + dx)b^2 + b^2} d(b \tan(c + dx))}{d} + \frac{C(b \tan(c + dx))^{n+3}}{bd(n+3)}}{b^2} \\
 & \quad \downarrow \text{278} \\
 & \frac{(A - C)(b \tan(c + dx))^{n+3} \operatorname{Hypergeometric2F1}\left(1, \frac{n+3}{2}, \frac{n+5}{2}, -\tan^2(c + dx)\right)}{bd(n+3)} + \frac{B(b \tan(c + dx))^{n+4} \operatorname{Hypergeometric2F1}\left(1, \frac{n+4}{2}, \frac{n+6}{2}, -\tan^2(c + dx)\right)}{b^2 d(n+4)}
 \end{aligned}$$

input

```
Int[Tan[c + d*x]^2*(b*Tan[c + d*x])^n*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
((C*(b*Tan[c + d*x])^(3 + n))/(b*d*(3 + n)) + ((A - C)*Hypergeometric2F1[1, (3 + n)/2, (5 + n)/2, -Tan[c + d*x]^2]*(b*Tan[c + d*x])^(3 + n))/(b*d*(3 + n)) + (B*Hypergeometric2F1[1, (4 + n)/2, (6 + n)/2, -Tan[c + d*x]^2]*(b*Tan[c + d*x])^(4 + n))/(b^2*d*(4 + n)))/b^2
```

Definitions of rubi rules used

rule 278 $\text{Int}[\text{((c_.)*(x_))}^{\text{(m_.)}* \text{((a_) + (b_.)*(x_)^2)}^{\text{(p_)}, x_Symbol] \text{ :> Simp}[a^{\text{p}}*((c*x)^{\text{(m + 1)}}/(c*(m + 1)))*\text{Hypergeometric2F1}[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] \text{ /; FreeQ}\{a, b, c, m, p\}, x] \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

rule 2030 $\text{Int}[(\text{Fx_.}) * (\text{v_.})^{\text{(m_.)}* \text{((b_.)*(v_))}^{\text{(n_)}, x_Symbol] \text{ :> Simp}[1/b^{\text{m}} \ \text{Int}[(b*v)^{\text{(m + n)}}*Fx, x], x] \text{ /; FreeQ}\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

rule 3042 $\text{Int}[u_, x_Symbol] \text{ :> Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3957 $\text{Int}[\text{((b_.)*tan}[(c_.) + (d_.)*(x_)]\text{)}^{\text{(n_)}, x_Symbol] \text{ :> Simp}[b/d \ \text{Subst}[\text{Int}[x^{\text{n}}/(b^2 + x^2), x], x, b*\text{Tan}[c + d*x]], x] \text{ /; FreeQ}\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[n]$

rule 4021 $\text{Int}[\text{((b_.)*tan}[(e_.) + (f_.)*(x_)]\text{)}^{\text{(m_.)}* \text{((c_) + (d_.)*tan}[(e_.) + (f_.)*(x_)]\text{)}, x_Symbol] \text{ :> Simp}[c \ \text{Int}[(b*\text{Tan}[e + f*x])^{\text{m}}, x], x] + \text{Simp}[d/b \ \text{Int}[(b*\text{Tan}[e + f*x])^{\text{(m + 1)}}, x], x] \text{ /; FreeQ}\{b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ !\text{IntegerQ}[2*m]$

rule 4113 $\text{Int}[\text{((a_.) + (b_.)*tan}[(e_.) + (f_.)*(x_)]\text{)}^{\text{(m_.)}* \text{((A_.) + (B_.)*tan}[(e_.) + (f_.)*(x_)] + (C_.)*tan}[(e_.) + (f_.)*(x_)]^2\text{)}, x_Symbol] \text{ :> Simp}[C*((a + b*\text{Tan}[e + f*x])^{\text{(m + 1)}}/(b*f*(m + 1))), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{\text{m}}*\text{Simp}[A - C + B*\text{Tan}[e + f*x], x], x] \text{ /; FreeQ}\{a, b, e, f, A, B, C, m\}, x] \ \&\& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\& \ !\text{LeQ}[m, -1]$

Maple [F]

$$\int \tan(dx+c)^2 (b \tan(dx+c))^n (A+B \tan(dx+c)+C \tan(dx+c)^2) dx$$

input `int(tan(d*x+c)^2*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `int(tan(d*x+c)^2*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

Fricas [F]

$$\begin{aligned} & \int \tan^2(c+dx)(b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) dx \\ &= \int (C \tan(dx+c)^2 + B \tan(dx+c) + A)(b \tan(dx+c))^n \tan(dx+c)^2 dx \end{aligned}$$

input `integrate(tan(d*x+c)^2*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="fricas")`

output `integral((C*tan(d*x+c)^4 + B*tan(d*x+c)^3 + A*tan(d*x+c)^2)*(b*tan(d
*x+c))^n, x)`

Sympy [F]

$$\begin{aligned} & \int \tan^2(c+dx)(b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) dx \\ &= \int (b \tan(c+dx))^n (A+B \tan(c+dx)+C \tan^2(c+dx)) \tan^2(c+dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**2*(b*tan(d*x+c))**n*(A+B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Integral((b*tan(c+d*x))**n*(A+B*tan(c+d*x)+C*tan(c+d*x)**2)*tan(c+d*x)**2, x)`

Maxima [F]

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int (C \tan(dx + c)^2 + B \tan(dx + c) + A)(b \tan(dx + c))^n \tan(dx + c)^2 dx$$

input

```
integrate(tan(d*x+c)^2*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*(b*tan(d*x + c))^n*tan(d
*x + c)^2, x)
```

Giac [F]

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int (C \tan(dx + c)^2 + B \tan(dx + c) + A)(b \tan(dx + c))^n \tan(dx + c)^2 dx$$

input

```
integrate(tan(d*x+c)^2*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*(b*tan(d*x + c))^n*tan(d
*x + c)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int \tan(c + dx)^2 (b \tan(c + dx))^n (C \tan(c + dx)^2 + B \tan(c + dx) + A) dx$$

input `int(tan(c + d*x)^2*(b*tan(c + d*x))^n*(A + B*tan(c + d*x) + C*tan(c + d*x)^2),x)`

output `int(tan(c + d*x)^2*(b*tan(c + d*x))^n*(A + B*tan(c + d*x) + C*tan(c + d*x)^2), x)`

Reduce [F]

$$\int \tan^2(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{b^n \left(\tan(dx + c)^n \tan(dx + c)^2 bn - \tan(dx + c)^n bn - 2 \tan(dx + c)^n b + \left(\int \frac{\tan(dx+c)^n}{\tan(dx+c)} dx \right) bd n^2 + 2 \left(\int \right)}{}$$

input `int(tan(d*x+c)^2*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `(b**n*(tan(c + d*x)**n*tan(c + d*x)**2*b*n - tan(c + d*x)**n*b*n - 2*tan(c + d*x)**n*b + int(tan(c + d*x)**n/tan(c + d*x),x)*b*d*n**2 + 2*int(tan(c + d*x)**n/tan(c + d*x),x)*b*d*n + int(tan(c + d*x)**n*tan(c + d*x)**4,x)*c*d*n**2 + 2*int(tan(c + d*x)**n*tan(c + d*x)**4,x)*c*d*n + int(tan(c + d*x)**n*tan(c + d*x)**2,x)*a*d*n**2 + 2*int(tan(c + d*x)**n*tan(c + d*x)**2,x)*a*d*n))/(d*n*(n + 2))`

3.46 $\int \tan^m(c+dx)(b \tan(c+dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$

Optimal result	515
Mathematica [A] (verified)	516
Rubi [A] (verified)	516
Maple [F]	519
Fricas [F]	519
Sympy [F]	519
Maxima [F]	520
Giac [F]	520
Mupad [F(-1)]	521
Reduce [F]	521

Optimal result

Integrand size = 39, antiderivative size = 154

$$\int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{C \tan^{1+m}(c + dx)(b \tan(c + dx))^n}{d(1 + m + n)}$$

$$+ \frac{(A - C) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + m + n), \frac{1}{2}(3 + m + n), -\tan^2(c + dx)\right) \tan^{1+m}(c + dx)(b \tan(c + dx))^n}{d(1 + m + n)}$$

$$+ \frac{B \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(2 + m + n), \frac{1}{2}(4 + m + n), -\tan^2(c + dx)\right) \tan^{2+m}(c + dx)(b \tan(c + dx))^n}{d(2 + m + n)}$$

output

```
C*tan(d*x+c)^(1+m)*(b*tan(d*x+c))^n/d/(1+m+n)+(A-C)*hypergeom([1, 1/2+1/2*m+1/2*n], [3/2+1/2*m+1/2*n], -tan(d*x+c)^2)*tan(d*x+c)^(1+m)*(b*tan(d*x+c))^n/d/(1+m+n)+B*hypergeom([1, 1+1/2*m+1/2*n], [2+1/2*m+1/2*n], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)*(b*tan(d*x+c))^n/d/(2+m+n)
```


Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.75

$$\int \tan^m(c+dx)(b \tan(c+dx))^n (A + B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \frac{\tan^{1+m}(c+dx)(b \tan(c+dx))^n \left(\frac{C}{1+m+n} + \frac{(A-C) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1+m+n), \frac{1}{2}(3+m+n), -\tan^2(c+dx)\right)}{1+m+n} + \frac{B \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1+m+n), \frac{1}{2}(3+m+n), -\tan^2(c+dx)\right)}{1+m+n} \right)}{d}$$

input

```
Integrate[Tan[c + d*x]^m*(b*Tan[c + d*x])^n*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(Tan[c + d*x]^(1 + m)*(b*Tan[c + d*x])^n*(C/(1 + m + n) + ((A - C)*Hypergeometric2F1[1, (1 + m + n)/2, (3 + m + n)/2, -Tan[c + d*x]^2])/(1 + m + n) + (B*Hypergeometric2F1[1, (2 + m + n)/2, (4 + m + n)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m + n))/d
```

Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.205$, Rules used = {2034, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^m(c+dx)(b \tan(c+dx))^n (A + B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$\downarrow \text{2034}$$

$$\tan^{-n}(c+dx)(b \tan(c+dx))^n \int \tan^{m+n}(c+dx) (C \tan^2(c+dx) + B \tan(c+dx) + A) dx$$

$$\downarrow \text{3042}$$

$$\tan^{-n}(c+dx)(b \tan(c+dx))^n \int \tan(c+dx)^{m+n} (C \tan(c+dx)^2 + B \tan(c+dx) + A) dx$$

$$\downarrow \text{4113}$$

$$\begin{aligned}
& \tan^{-n}(c+dx)(b \tan(c+dx))^n \left(\int \tan^{m+n}(c+dx)(A-C+B \tan(c+dx))dx + \frac{C \tan^{m+n+1}(c+dx)}{d(m+n+1)} \right) \\
& \quad \downarrow \text{3042} \\
& \tan^{-n}(c+dx)(b \tan(c+dx))^n \left(\int \tan(c+dx)^{m+n}(A-C+B \tan(c+dx))dx + \frac{C \tan^{m+n+1}(c+dx)}{d(m+n+1)} \right) \\
& \quad \downarrow \text{4021} \\
& \tan^{-n}(c+dx)(b \tan(c+dx))^n \left((A-C) \int \tan^{m+n}(c+dx)dx + B \int \tan^{m+n+1}(c+dx)dx + \frac{C \tan^{m+n+1}(c+dx)}{d(m+n+1)} \right) \\
& \quad \downarrow \text{3042} \\
& \tan^{-n}(c+dx)(b \tan(c+dx))^n \left((A-C) \int \tan(c+dx)^{m+n}dx + B \int \tan(c+dx)^{m+n+1}dx + \frac{C \tan^{m+n+1}(c+dx)}{d(m+n+1)} \right) \\
& \quad \downarrow \text{3957} \\
& \tan^{-n}(c+dx)(b \tan(c+dx))^n \left(\frac{(A-C) \int \frac{\tan^{m+n}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{B \int \frac{\tan^{m+n+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{C \tan^{m+n+1}(c+dx)}{d(m+n+1)} \right) \\
& \quad \downarrow \text{278} \\
& \tan^{-n}(c+dx)(b \tan(c+dx))^n \left(\frac{(A-C) \tan^{m+n+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(m+n+1), \frac{1}{2}(m+n+3), -\tan^2(c+dx)\right)}{d(m+n+1)} + \frac{B \tan^{m+n+1}(c+dx)}{d(m+n+1)} \right)
\end{aligned}$$

input

```
Int[Tan[c + d*x]^m*(b*Tan[c + d*x])^n*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
((b*Tan[c + d*x])^n*((C*Tan[c + d*x]^(1 + m + n))/(d*(1 + m + n)) + ((A - C)*Hypergeometric2F1[1, (1 + m + n)/2, (3 + m + n)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m + n))/(d*(1 + m + n)) + (B*Hypergeometric2F1[1, (2 + m + n)/2, (4 + m + n)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m + n))/(d*(2 + m + n))))/Tan[c + d*x]^n
```

Defintions of rubi rules used

rule 278 $\text{Int}[\text{((c_)}*(x_))^{\text{(m_)}* \text{((a_)} + \text{(b_)}*(x_)^2)^{\text{(p_)}}, x_Symbol] \text{ :> Simp}[a^p*((c*x)^{(m+1))/(c*(m+1))]*\text{Hypergeometric2F1}[-p, (m+1)/2, (m+1)/2+1, (-b)*(x^2/a)], x] \text{ /; FreeQ}\{a, b, c, m, p\}, x] \&\& \text{ !IGtQ}\{p, 0\} \&\& (\text{ILtQ}\{p, 0\} \text{ || GtQ}\{a, 0\})$

rule 2034 $\text{Int}[(F x_)* \text{((a_)}*(v_))^{\text{(m_)}* \text{((b_)}*(v_))^{\text{(n_)}}, x_Symbol] \text{ :> Simp}[b^{\text{IntPart}[n]}*((b*v)^{\text{FracPart}[n]}/(a^{\text{IntPart}[n]}*(a*v)^{\text{FracPart}[n]})) \text{ Int}[(a*v)^{(m+n)}]*F x, x], x] \text{ /; FreeQ}\{a, b, m, n\}, x] \&\& \text{ !IntegerQ}\{m\} \&\& \text{ !IntegerQ}\{n\} \&\& \text{ !IntegerQ}\{m+n\}$

rule 3042 $\text{Int}[u_, x_Symbol] \text{ :> Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3957 $\text{Int}[\text{((b_)}*\text{tan}[(c_)} + \text{(d_)}*(x_))]^{\text{(n_)}}, x_Symbol] \text{ :> Simp}[b/d \text{ Subst}[\text{Int}[x^n/(b^2+x^2), x], x, b*\text{Tan}[c+d*x]], x] \text{ /; FreeQ}\{b, c, d, n\}, x] \&\& \text{ !IntegerQ}\{n\}$

rule 4021 $\text{Int}[\text{((b_)}*\text{tan}[(e_)} + \text{(f_)}*(x_))]^{\text{(m_)}* \text{((c_)} + \text{(d_)}*\text{tan}[(e_)} + \text{(f_)}*(x_)]), x_Symbol] \text{ :> Simp}[c \text{ Int}[(b*\text{Tan}[e+f*x])^m, x], x] + \text{Simp}[d/b \text{ Int}[(b*\text{Tan}[e+f*x])^{(m+1)}, x], x] \text{ /; FreeQ}\{b, c, d, e, f, m\}, x] \&\& \text{ NeQ}[c^2+d^2, 0] \&\& \text{ !IntegerQ}\{2*m\}$

rule 4113 $\text{Int}[\text{((a_)} + \text{(b_)}*\text{tan}[(e_)} + \text{(f_)}*(x_))]^{\text{(m_)}* \text{((A_)} + \text{(B_)}*\text{tan}[(e_)} + \text{(f_)}*(x_)] + \text{(C_)}*\text{tan}[(e_)} + \text{(f_)}*(x_)]^2), x_Symbol] \text{ :> Simp}[C*((a+b*\text{Tan}[e+f*x])^{(m+1)}/(b*f*(m+1))), x] + \text{Int}[(a+b*\text{Tan}[e+f*x])^m*\text{Simp}[A-C+B*\text{Tan}[e+f*x], x], x] \text{ /; FreeQ}\{a, b, e, f, A, B, C, m\}, x] \&\& \text{ NeQ}[A*b^2-a*b*B+a^2*C, 0] \&\& \text{ !LeQ}\{m, -1\}$

Maple [F]

$$\int \tan(dx + c)^m (b \tan(dx + c))^n (A + B \tan(dx + c) + C \tan(dx + c)^2) dx$$

input `int(tan(d*x+c)^m*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `int(tan(d*x+c)^m*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

Fricas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx \\ & = \int (C \tan(dx + c)^2 + B \tan(dx + c) + A)(b \tan(dx + c))^n \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="fricas")`

output `integral((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*(b*tan(d*x + c))^n*tan(d*x + c)^m, x)`

Sympy [F]

$$\begin{aligned} & \int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx \\ & = \int (b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) \tan^m(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**m*(b*tan(d*x+c))**n*(A+B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Integral((b*tan(c + d*x))**n*(A + B*tan(c + d*x) + C*tan(c + d*x)**2)*tan(c + d*x)**m, x)`

Maxima [F]

$$\int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int (C \tan(dx + c)^2 + B \tan(dx + c) + A)(b \tan(dx + c))^n \tan(dx + c)^m dx$$

input

```
integrate(tan(d*x+c)^m*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="maxima")
```

output

```
integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*(b*tan(d*x + c))^n*tan(d
*x + c)^m, x)
```

Giac [F]

$$\int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int (C \tan(dx + c)^2 + B \tan(dx + c) + A)(b \tan(dx + c))^n \tan(dx + c)^m dx$$

input

```
integrate(tan(d*x+c)^m*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x,
algorithm="giac")
```

output

```
integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*(b*tan(d*x + c))^n*tan(d
*x + c)^m, x)
```

Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int \tan(c + dx)^m (b \tan(c + dx))^n (C \tan(c + dx)^2 + B \tan(c + dx) + A) dx$$

input `int(tan(c + d*x)^m*(b*tan(c + d*x))^n*(A + B*tan(c + d*x) + C*tan(c + d*x)^2),x)`

output `int(tan(c + d*x)^m*(b*tan(c + d*x))^n*(A + B*tan(c + d*x) + C*tan(c + d*x)^2), x)`

Reduce [F]

$$\int \tan^m(c + dx)(b \tan(c + dx))^n (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{b^n \left(\tan(dx + c)^{m+n} b + \left(\int \tan(dx + c)^{m+n} dx \right) adm + \left(\int \tan(dx + c)^{m+n} dx \right) adn - \left(\int \frac{\tan(dx+c)^{m+n}}{\tan(dx+c)} dx \right) \right)}{d(m+n)}$$

input `int(tan(d*x+c)^m*(b*tan(d*x+c))^n*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)`

output `(b**n*(tan(c + d*x)**(m + n)*b + int(tan(c + d*x)**(m + n),x)*a*d*m + int(tan(c + d*x)**(m + n),x)*a*d*n - int(tan(c + d*x)**(m + n)/tan(c + d*x),x)*b*d*m - int(tan(c + d*x)**(m + n)/tan(c + d*x),x)*b*d*n + int(tan(c + d*x)**(m + n)*tan(c + d*x)**2,x)*c*d*m + int(tan(c + d*x)**(m + n)*tan(c + d*x)**2,x)*c*d*n))/(d*(m + n))`

3.47 $\int \tan^m(c+dx) \sqrt{b \tan(c+dx)} (A + B \tan(c+dx) + C \tan^2(c+dx)) dx$

Optimal result	522
Mathematica [A] (verified)	523
Rubi [A] (verified)	523
Maple [F]	526
Fricas [F]	526
Sympy [F]	527
Maxima [F(-1)]	527
Giac [F]	527
Mupad [F(-1)]	528
Reduce [F]	528

Optimal result

Integrand size = 41, antiderivative size = 170

$$\int \tan^m(c+dx) \sqrt{b \tan(c+dx)} (A + B \tan(c+dx) + C \tan^2(c+dx)) dx$$

$$= \frac{2C \tan^{1+m}(c+dx) \sqrt{b \tan(c+dx)}}{d(3+2m)}$$

$$+ \frac{2(A-C) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(3+2m), \frac{1}{4}(7+2m), -\tan^2(c+dx)\right) \tan^{1+m}(c+dx) \sqrt{b \tan(c+dx)}}{d(3+2m)}$$

$$+ \frac{2B \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(5+2m), \frac{1}{4}(9+2m), -\tan^2(c+dx)\right) \tan^{2+m}(c+dx) \sqrt{b \tan(c+dx)}}{d(5+2m)}$$

output

```
2*C*tan(d*x+c)^(1+m)*(b*tan(d*x+c))^(1/2)/d/(3+2*m)+2*(A-C)*hypergeom([1,
3/4+1/2*m],[7/4+1/2*m],-tan(d*x+c)^2)*tan(d*x+c)^(1+m)*(b*tan(d*x+c))^(1/2
)/d/(3+2*m)+2*B*hypergeom([1, 5/4+1/2*m],[9/4+1/2*m],-tan(d*x+c)^2)*tan(d*
x+c)^(2+m)*(b*tan(d*x+c))^(1/2)/d/(5+2*m)
```

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.78

$$\int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{2 \tan^{1+m}(c + dx) \sqrt{b \tan(c + dx)} (C(5 + 2m) + (A - C)(5 + 2m) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(3 + 2m), \frac{1}{4}(3 + 2m) + 1, -\tan^2(c + dx)\right))}{d(3 + 2m)(5 + 2m)}$$

input

```
Integrate[Tan[c + d*x]^m*Sqrt[b*Tan[c + d*x]]*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2),x]
```

output

```
(2*Tan[c + d*x]^(1 + m)*Sqrt[b*Tan[c + d*x]]*(C*(5 + 2*m) + (A - C)*(5 + 2*m)*Hypergeometric2F1[1, (3 + 2*m)/4, (7 + 2*m)/4, -Tan[c + d*x]^2] + B*(3 + 2*m)*Hypergeometric2F1[1, (5 + 2*m)/4, (9 + 2*m)/4, -Tan[c + d*x]^2]*Tan[c + d*x]))/(d*(3 + 2*m)*(5 + 2*m))
```

Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.195$, Rules used = {2034, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{b \tan(c + dx)} \tan^m(c + dx) (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$\downarrow 2034$$

$$\frac{\sqrt{b \tan(c + dx)} \int \tan^{m+\frac{1}{2}}(c + dx) (C \tan^2(c + dx) + B \tan(c + dx) + A) dx}{\sqrt{\tan(c + dx)}}$$

$$\downarrow 3042$$

$$\frac{\sqrt{b \tan(c + dx)} \int \tan(c + dx)^{m+\frac{1}{2}} (C \tan(c + dx)^2 + B \tan(c + dx) + A) dx}{\sqrt{\tan(c + dx)}}$$

$$\downarrow 4113$$

$$\begin{aligned}
& \frac{\sqrt{b \tan(c+dx)} \left(\int \tan^{m+\frac{1}{2}}(c+dx)(A-C+B \tan(c+dx))dx + \frac{2C \tan^{m+\frac{3}{2}}(c+dx)}{d(2m+3)} \right)}{\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{b \tan(c+dx)} \left(\int \tan(c+dx)^{m+\frac{1}{2}}(A-C+B \tan(c+dx))dx + \frac{2C \tan^{m+\frac{3}{2}}(c+dx)}{d(2m+3)} \right)}{\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{4021} \\
& \frac{\sqrt{b \tan(c+dx)} \left((A-C) \int \tan^{m+\frac{1}{2}}(c+dx)dx + B \int \tan^{m+\frac{3}{2}}(c+dx)dx + \frac{2C \tan^{m+\frac{3}{2}}(c+dx)}{d(2m+3)} \right)}{\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{b \tan(c+dx)} \left((A-C) \int \tan(c+dx)^{m+\frac{1}{2}}dx + B \int \tan(c+dx)^{m+\frac{3}{2}}dx + \frac{2C \tan^{m+\frac{3}{2}}(c+dx)}{d(2m+3)} \right)}{\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{3957} \\
& \frac{\sqrt{b \tan(c+dx)} \left(\frac{(A-C) \int \frac{\tan^{m+\frac{1}{2}}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{B \int \frac{\tan^{m+\frac{3}{2}}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{2C \tan^{m+\frac{3}{2}}(c+dx)}{d(2m+3)} \right)}{\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{278} \\
& \frac{\sqrt{b \tan(c+dx)} \left(\frac{2(A-C) \tan^{m+\frac{3}{2}}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(2m+3), \frac{1}{4}(2m+7), -\tan^2(c+dx)\right)}{d(2m+3)} + \frac{2B \tan^{m+\frac{5}{2}}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(2m+5), \frac{1}{4}(2m+7), -\tan^2(c+dx)\right)}{d(2m+3)} \right)}{\sqrt{\tan(c+dx)}}
\end{aligned}$$

input

```
Int[Tan[c + d*x]^m*sqrt[b*Tan[c + d*x]]*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2), x]
```

output

```
(Sqrt[b*Tan[c + d*x]]*((2*C*Tan[c + d*x]^(3/2 + m))/(d*(3 + 2*m)) + (2*(A
- C)*Hypergeometric2F1[1, (3 + 2*m)/4, (7 + 2*m)/4, -Tan[c + d*x]^2]*Tan[c
+ d*x]^(3/2 + m))/(d*(3 + 2*m)) + (2*B*Hypergeometric2F1[1, (5 + 2*m)/4,
(9 + 2*m)/4, -Tan[c + d*x]^2]*Tan[c + d*x]^(5/2 + m))/(d*(5 + 2*m))))/Sqrt
[Tan[c + d*x]]
```

Defintions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

rule 2034

```
Int[(Fx_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3957

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

rule 4021

```
Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x
_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^
2 + d^2, 0] && !IntegerQ[2*m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [F]

$$\int \tan(dx + c)^m \sqrt{b \tan(dx + c)} (A + B \tan(dx + c) + C \tan(dx + c)^2) dx$$

input

```
int(tan(d*x+c)^m*(b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

output

```
int(tan(d*x+c)^m*(b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x)
```

Fricas [F]

$$\begin{aligned} & \int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx \\ & = \int (C \tan(dx + c)^2 + B \tan(dx + c) + A) \sqrt{b \tan(dx + c)} \tan(dx + c)^m dx \end{aligned}$$

input

```
integrate(tan(d*x+c)^m*(b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)^2
),x, algorithm="fricas")
```

output

```
integral((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c))*tan(
d*x + c)^m, x)
```

Sympy [F]

$$\int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)**2),x)`

output `Integral(sqrt(b*tan(c + d*x))*(A + B*tan(c + d*x) + C*tan(c + d*x)**2)*tan(c + d*x)**m, x)`

Maxima [F(-1)]

Timed out.

$$\int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int (C \tan(dx + c)^2 + B \tan(dx + c) + A) \sqrt{b \tan(dx + c)} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)^2),x, algorithm="giac")`

output `integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c))*tan(d*x + c)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \int \tan(c + dx)^m \sqrt{b \tan(c + dx)} (C \tan(c + dx)^2 + B \tan(c + dx) + A) dx$$

input `int(tan(c + d*x)^m*(b*tan(c + d*x))^(1/2)*(A + B*tan(c + d*x) + C*tan(c + d*x)^2), x)`

output `int(tan(c + d*x)^m*(b*tan(c + d*x))^(1/2)*(A + B*tan(c + d*x) + C*tan(c + d*x)^2), x)`

Reduce [F]

$$\int \tan^m(c + dx) \sqrt{b \tan(c + dx)} (A + B \tan(c + dx) + C \tan^2(c + dx)) dx$$

$$= \frac{\sqrt{b} \left(2 \tan(dx + c)^{m+\frac{1}{2}} b + 2 \left(\int \tan(dx + c)^{m+\frac{1}{2}} dx \right) adm + \left(\int \tan(dx + c)^{m+\frac{1}{2}} dx \right) ad - 2 \left(\int \frac{\tan(dx+c)}{\tan(dx+c)} dx \right) \right)}{1}$$

input `int(tan(d*x+c)^m*(b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)+C*tan(d*x+c)^2), x)`

output `(sqrt(b)*(2*tan(c + d*x)**((2*m + 1)/2)*b + 2*int(tan(c + d*x)**((2*m + 1)/2), x)*a*d*m + int(tan(c + d*x)**((2*m + 1)/2), x)*a*d - 2*int(tan(c + d*x)**((2*m + 1)/2)/tan(c + d*x), x)*b*d*m - int(tan(c + d*x)**((2*m + 1)/2)/tan(c + d*x), x)*b*d + 2*int(tan(c + d*x)**((2*m + 1)/2)*tan(c + d*x)**2, x)*c*d*m + int(tan(c + d*x)**((2*m + 1)/2)*tan(c + d*x)**2, x)*c*d))/(d*(2*m + 1))`

3.48
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{b \tan(c+dx)}} dx$$

Optimal result	529
Mathematica [A] (verified)	530
Rubi [A] (verified)	530
Maple [F]	533
Fricas [F]	533
Sympy [F]	534
Maxima [F(-1)]	534
Giac [F(-2)]	534
Mupad [F(-1)]	535
Reduce [F]	535

Optimal result

Integrand size = 41, antiderivative size = 170

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{b \tan(c+dx)}} dx = \frac{2C \tan^{1+m}(c+dx)}{d(1+2m)\sqrt{b \tan(c+dx)}} + \frac{2(A-C) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(1+2m), \frac{1}{4}(5+2m), -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{d(1+2m)\sqrt{b \tan(c+dx)}} + \frac{2B \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(3+2m), \frac{1}{4}(7+2m), -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{d(3+2m)\sqrt{b \tan(c+dx)}}$$

output

```
2*C*tan(d*x+c)^(1+m)/d/(1+2*m)/(b*tan(d*x+c))^(1/2)+2*(A-C)*hypergeom([1,
1/4+1/2*m],[5/4+1/2*m],-tan(d*x+c)^2)*tan(d*x+c)^(1+m)/d/(1+2*m)/(b*tan(d*
x+c))^(1/2)+2*B*hypergeom([1, 3/4+1/2*m],[7/4+1/2*m],-tan(d*x+c)^2)*tan(d*
x+c)^(2+m)/d/(3+2*m)/(b*tan(d*x+c))^(1/2)
```

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.78

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

$$= \frac{2\tan^{1+m}(c+dx)(C(3+2m)+(A-C)(3+2m)\text{Hypergeometric2F1}\left(1, \frac{1}{4}(1+2m), \frac{1}{4}(5+2m), -\tan^2(c+dx)\right))}{d(1+2m)(3+2m)\sqrt{b\tan(c+dx)}}$$

input

```
Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2))/Sqrt[b*
Tan[c + d*x]],x]
```

output

```
(2*Tan[c + d*x]^(1 + m)*(C*(3 + 2*m) + (A - C)*(3 + 2*m)*Hypergeometric2F1
[1, (1 + 2*m)/4, (5 + 2*m)/4, -Tan[c + d*x]^2] + B*(1 + 2*m)*Hypergeometri
c2F1[1, (3 + 2*m)/4, (7 + 2*m)/4, -Tan[c + d*x]^2]*Tan[c + d*x]))/(d*(1 +
2*m)*(3 + 2*m)*Sqrt[b*Tan[c + d*x]])
```

Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.195$, Rules used = {2034, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

$$\downarrow 2034$$

$$\frac{\sqrt{\tan(c+dx)} \int \tan^{m-\frac{1}{2}}(c+dx)(C\tan^2(c+dx)+B\tan(c+dx)+A) dx}{\sqrt{b\tan(c+dx)}}$$

$$\downarrow 3042$$

$$\frac{\sqrt{\tan(c+dx)} \int \tan(c+dx)^{m-\frac{1}{2}}(C\tan(c+dx)^2+B\tan(c+dx)+A) dx}{\sqrt{b\tan(c+dx)}}$$

$$\begin{aligned}
& \downarrow 4113 \\
& \frac{\sqrt{\tan(c+dx)} \left(\int \tan^{m-\frac{1}{2}}(c+dx)(A-C+B\tan(c+dx))dx + \frac{2C \tan^{m+\frac{1}{2}}(c+dx)}{d(2m+1)} \right)}{\sqrt{b \tan(c+dx)}} \\
& \downarrow 3042 \\
& \frac{\sqrt{\tan(c+dx)} \left(\int \tan(c+dx)^{m-\frac{1}{2}}(A-C+B\tan(c+dx))dx + \frac{2C \tan^{m+\frac{1}{2}}(c+dx)}{d(2m+1)} \right)}{\sqrt{b \tan(c+dx)}} \\
& \downarrow 4021 \\
& \frac{\sqrt{\tan(c+dx)} \left((A-C) \int \tan^{m-\frac{1}{2}}(c+dx)dx + B \int \tan^{m+\frac{1}{2}}(c+dx)dx + \frac{2C \tan^{m+\frac{1}{2}}(c+dx)}{d(2m+1)} \right)}{\sqrt{b \tan(c+dx)}} \\
& \downarrow 3042 \\
& \frac{\sqrt{\tan(c+dx)} \left((A-C) \int \tan(c+dx)^{m-\frac{1}{2}}dx + B \int \tan(c+dx)^{m+\frac{1}{2}}dx + \frac{2C \tan^{m+\frac{1}{2}}(c+dx)}{d(2m+1)} \right)}{\sqrt{b \tan(c+dx)}} \\
& \downarrow 3957 \\
& \frac{\sqrt{\tan(c+dx)} \left(\frac{(A-C) \int \frac{\tan^{m-\frac{1}{2}}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{B \int \frac{\tan^{m+\frac{1}{2}}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{2C \tan^{m+\frac{1}{2}}(c+dx)}{d(2m+1)} \right)}{\sqrt{b \tan(c+dx)}} \\
& \downarrow 278 \\
& \frac{\sqrt{\tan(c+dx)} \left(\frac{2(A-C) \tan^{m+\frac{1}{2}}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(2m+1), \frac{1}{4}(2m+5), -\tan^2(c+dx)\right)}{d(2m+1)} + \frac{2B \tan^{m+\frac{3}{2}}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{1}{4}(2m+1), \frac{1}{4}(2m+5), -\tan^2(c+dx)\right)}{d(2m+1)} \right)}{\sqrt{b \tan(c+dx)}}
\end{aligned}$$

input

```
Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2))/Sqrt[b*Tan[c + d*x]],x]
```


output

```
(Sqrt[Tan[c + d*x]]*((2*C*Tan[c + d*x]^(1/2 + m))/(d*(1 + 2*m)) + (2*(A -
C)*Hypergeometric2F1[1, (1 + 2*m)/4, (5 + 2*m)/4, -Tan[c + d*x]^2]*Tan[c +
d*x]^(1/2 + m))/(d*(1 + 2*m)) + (2*B*Hypergeometric2F1[1, (3 + 2*m)/4, (7
+ 2*m)/4, -Tan[c + d*x]^2]*Tan[c + d*x]^(3/2 + m))/(d*(3 + 2*m))))/Sqrt[b
*Tan[c + d*x]]
```

Defintions of rubi rules used

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

rule 2034

```
Int[(Fx_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Simp[b^IntPart
[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n
)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] &&
!IntegerQ[m + n]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3957

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

rule 4021

```
Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x
_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^
2 + d^2, 0] && !IntegerQ[2*m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [F]

$$\int \frac{\tan(dx+c)^m (A+B\tan(dx+c)+C\tan(dx+c)^2)}{\sqrt{b\tan(dx+c)}} dx$$

input

```
int(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x)
```

output

```
int(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x)
```

Fricas [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

$$= \int \frac{(C\tan(dx+c)^2+B\tan(dx+c)+A)\tan(dx+c)^m}{\sqrt{b\tan(dx+c)}} dx$$

input

```
integrate(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(b*tan(d*x+c))^(1/2
),x, algorithm="fricas")
```

output

```
integral((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c))*tan(
d*x + c)^m/(b*tan(d*x + c)), x)
```

Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

$$= \int \frac{(A+B\tan(c+dx)+C\tan^2(c+dx))\tan^m(c+dx)}{\sqrt{b\tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c)+C*tan(d*x+c)**2)/(b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x) + C*tan(c + d*x)**2)*tan(c + d*x)**m/sqrt(b*tan(c + d*x)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Timed out`

Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

= Exception raised: TypeError

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{1, [3,9]%%}+%%{4, [3,7]%%}+%%{6, [3,5]%%}+%%{4, [3,3]%%}+%%

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

$$= \int \frac{\tan(c+dx)^m (C\tan(c+dx)^2 + B\tan(c+dx) + A)}{\sqrt{b\tan(c+dx)}} dx$$

input `int((tan(c+d*x)^m*(A+B*tan(c+d*x)+C*tan(c+d*x)^2))/(b*tan(c+d*x))^(1/2),x)`

output `int((tan(c+d*x)^m*(A+B*tan(c+d*x)+C*tan(c+d*x)^2))/(b*tan(c+d*x))^(1/2),x)`

Reduce [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{b\tan(c+dx)}} dx$$

$$= \frac{\sqrt{b} \left(2\tan(dx+c)^{m+\frac{1}{2}}c + 2 \left(\int \tan(dx+c)^{m+\frac{1}{2}} dx \right) bdm + \left(\int \tan(dx+c)^{m+\frac{1}{2}} dx \right) bd + 2 \left(\int \frac{\tan(dx+c)^m}{\tan(dx+c)} dx \right) \right)}{bd(2m+1)}$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x)`

output

```
(sqrt(b)*(2*tan(c + d*x)**((2*m + 1)/2)*c + 2*int(tan(c + d*x)**((2*m + 1)
/2),x)*b*d*m + int(tan(c + d*x)**((2*m + 1)/2),x)*b*d + 2*int(tan(c + d*x)
**((2*m + 1)/2)/tan(c + d*x),x)*a*d*m + int(tan(c + d*x)**((2*m + 1)/2)/ta
n(c + d*x),x)*a*d - 2*int(tan(c + d*x)**((2*m + 1)/2)/tan(c + d*x),x)*c*d*
m - int(tan(c + d*x)**((2*m + 1)/2)/tan(c + d*x),x)*c*d))/(b*d*(2*m + 1))
```

$$3.49 \quad \int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

Optimal result	537
Mathematica [F]	538
Rubi [A] (warning: unable to verify)	538
Maple [F]	540
Fricas [F]	540
Sympy [F]	541
Maxima [F]	541
Giac [F]	542
Mupad [F(-1)]	542
Reduce [F]	543

Optimal result

Integrand size = 43, antiderivative size = 289

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx)+C \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx =$$

$$\frac{(B+i(A-C)) \operatorname{AppellF1}\left(\frac{1}{2}, -m, 1, \frac{3}{2}, \frac{a+b \tan(c+dx)}{a}, \frac{a+b \tan(c+dx)}{a-ib}\right) \tan^m(c+dx) \left(-\frac{b \tan(c+dx)}{a}\right)^{-m} \sqrt{a+b \tan(c+dx)}}{(a-ib)d}$$

$$\frac{(A+iB-C) \operatorname{AppellF1}\left(\frac{1}{2}, -m, 1, \frac{3}{2}, \frac{a+b \tan(c+dx)}{a}, \frac{a+b \tan(c+dx)}{a+ib}\right) \tan^m(c+dx) \left(-\frac{b \tan(c+dx)}{a}\right)^{-m} \sqrt{a+b \tan(c+dx)}}{(ia-b)d}$$

$$+\frac{2C \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -m, \frac{3}{2}, \frac{a+b \tan(c+dx)}{a}\right) \tan^m(c+dx) \left(-\frac{b \tan(c+dx)}{a}\right)^{-m} \sqrt{a+b \tan(c+dx)}}{bd}$$

output

```
-(B+I*(A-C))*AppellF1(1/2,-m,1,3/2,(a+b*tan(d*x+c))/a,(a+b*tan(d*x+c))/(a-I*b))*tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)/(a-I*b)/d/((-b*tan(d*x+c)/a)^m)-(A+I*B-C)*AppellF1(1/2,-m,1,3/2,(a+b*tan(d*x+c))/a,(a+b*tan(d*x+c))/(a+I*b))*tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)/(I*a-b)/d/((-b*tan(d*x+c)/a)^m)+2*C*hypergeom([1/2, -m],[3/2],(a+b*tan(d*x+c))/a)*tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)/b/d/((-b*tan(d*x+c)/a)^m)
```

Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$= \int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input

```
Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2))/Sqrt[a + b*Tan[c + d*x]], x]
```

output

```
Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2))/Sqrt[a + b*Tan[c + d*x]], x]
```

Rubi [A] (warning: unable to verify)

Time = 1.04 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.89, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.093$, Rules used = {3042, 4138, 2353, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^m (A+B\tan(c+dx)+C\tan(c+dx)^2)}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow \text{4138}$$

$$\int \frac{\tan^m(c+dx)(C\tan^2(c+dx)+B\tan(c+dx)+A)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan(c+dx)$$

$$\downarrow \text{2353}$$

$$\int \left(\frac{C \tan^m(c+dx)}{\sqrt{a+b \tan(c+dx)}} + \frac{(i(A-C)-B) \tan^m(c+dx)}{2(i-\tan(c+dx))\sqrt{a+b \tan(c+dx)}} + \frac{(B+i(A-C)) \tan^m(c+dx)}{2(\tan(c+dx)+i)\sqrt{a+b \tan(c+dx)}} \right) d \tan(c+dx)$$

d
↓ 2009

$$\frac{(A+iB-C) \tan^{m+1}(c+dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m+1, \frac{1}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right)}{2(m+1)\sqrt{a+b \tan(c+dx)}} + \frac{(A-iB-C) \tan^{m+1}(c+dx) \sqrt{\frac{b \tan(c+dx)}{a}}}{2(m+1)\sqrt{a+b \tan(c+dx)}}$$

input

```
Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x] + C*Tan[c + d*x]^2))/Sqrt[a + b*Tan[c + d*x]],x]
```

output

```
((2*C*Hypergeometric2F1[1/2, -m, 3/2, 1 + (b*Tan[c + d*x])/a]*Tan[c + d*x]^m*Sqrt[a + b*Tan[c + d*x]]/(b*(-((b*Tan[c + d*x])/a))^m) + ((A + I*B - C)*AppellF1[1 + m, 1/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])/(2*(1 + m)*Sqrt[a + b*Tan[c + d*x]]) + ((A - I*B - C)*AppellF1[1 + m, 1/2, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])/(2*(1 + m)*Sqrt[a + b*Tan[c + d*x]]))/d
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2353

```
Int[(Px_)*((e_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0]))
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```


rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F]

$$\int \frac{\tan(dx+c)^m (A+B\tan(dx+c)+C\tan(dx+c)^2)}{\sqrt{a+b\tan(dx+c)}} dx$$

input

```
int(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^(1/2),x)
```

output

```
int(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^(1/2),x)
```

Fricas [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx)+C\tan^2(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$= \int \frac{(C\tan(dx+c)^2+B\tan(dx+c)+A)\tan(dx+c)^m}{\sqrt{b\tan(dx+c)+a}} dx$$

input

```
integrate(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^(1
/2),x, algorithm="fricas")
```

output

```
integral((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*tan(d*x + c)^m/sqrt(b*tan
(d*x + c) + a), x)
```

Sympy [F]

$$\int \frac{\tan^m(c + dx) (A + B \tan(c + dx) + C \tan^2(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx) + C \tan^2(c + dx)) \tan^m(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c)+C*tan(d*x+c)**2)/(a+b*tan(d*x+c))*
*(1/2),x)`

output `Integral((A + B*tan(c + d*x) + C*tan(c + d*x)**2)*tan(c + d*x)**m/sqrt(a +
b*tan(c + d*x)), x)`

Maxima [F]

$$\int \frac{\tan^m(c + dx) (A + B \tan(c + dx) + C \tan^2(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \int \frac{(C \tan(dx + c)^2 + B \tan(dx + c) + A) \tan(dx + c)^m}{\sqrt{b \tan(dx + c) + a}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^(1
/2),x, algorithm="maxima")`

output `integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*tan(d*x + c)^m/sqrt(b*ta
n(d*x + c) + a), x)`

Giac [F]

$$\int \frac{\tan^m(c + dx) (A + B \tan(c + dx) + C \tan^2(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \int \frac{(C \tan(dx + c)^2 + B \tan(dx + c) + A) \tan(dx + c)^m}{\sqrt{b \tan(dx + c) + a}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((C*tan(d*x + c)^2 + B*tan(d*x + c) + A)*tan(d*x + c)^m/sqrt(b*tan(d*x + c) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c + dx) (A + B \tan(c + dx) + C \tan^2(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \int \frac{\tan(c + dx)^m (C \tan(c + dx)^2 + B \tan(c + dx) + A)}{\sqrt{a + b \tan(c + dx)}} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^(1/2),x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x) + C*tan(c + d*x)^2))/(a + b*tan(c + d*x))^(1/2), x)`

Reduce [F]

$$\int \frac{\tan^m(c + dx) (A + B \tan(c + dx) + C \tan^2(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c)+C*tan(d*x+c)^2)/(a+b*tan(d*x+c))^(1/2),x)`

output

```
(2*tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*a + 2*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x)**2)/(tan(c + d*x)*b + a),x)*b*c*d*m + int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x)**2)/(tan(c + d*x)*b + a),x)*b*c*d - 4*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x)**2)/(2*tan(c + d*x)*b*m + tan(c + d*x)*b + 2*a*m + a),x)*a*b*d*m**2 - 4*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x)**2)/(2*tan(c + d*x)*b*m + tan(c + d*x)*b + 2*a*m + a),x)*a*b*d*m - int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x)**2)/(2*tan(c + d*x)*b*m + tan(c + d*x)*b + 2*a*m + a),x)*a*b*d + 2*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x))/(tan(c + d*x)*b + a),x)*b**2*d*m + int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x))/(tan(c + d*x)*b + a),x)*b**2*d - 4*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x))/(2*tan(c + d*x)*b*m + tan(c + d*x)*b + 2*a*m + a),x)*a**2*d*m**2 - 2*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a)*tan(c + d*x))/(2*tan(c + d*x)*b*m + tan(c + d*x)*b + 2*a*m + a),x)*a**2*d*m - 4*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a))/(2*tan(c + d*x)**2*b*m + tan(c + d*x)**2*b + 2*tan(c + d*x)*a*m + tan(c + d*x)*a),x)*a**2*d*m**2 - 2*int((tan(c + d*x)**m*sqrt(tan(c + d*x)*b + a))/(2*tan(c + d*x)**2*b*m + tan(c + d*x)**2*b + 2*tan(c + d*x)*a*m + tan(c + d*x)*a),x)*a**2*d*m)/(b*d*(2*m + 1))
```

3.50 $\int (a+b \tan(e+fx))^3 (c+d \tan(e+fx)) (A + B \tan(e +$

Optimal result	544
Mathematica [C] (verified)	545
Rubi [A] (verified)	545
Maple [A] (warning: unable to verify)	549
Fricas [A] (verification not implemented)	550
Sympy [B] (verification not implemented)	551
Maxima [A] (verification not implemented)	552
Giac [B] (verification not implemented)	552
Mupad [B] (verification not implemented)	554
Reduce [B] (verification not implemented)	555

Optimal result

Integrand size = 43, antiderivative size = 353

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(a^3(Ac - cC - Bd) - 3ab^2(Ac - cC - Bd) - 3a^2b(Bc + (A - C)d) + b^3(Bc + (A - C)d)) x}{f} - \frac{(3a^2b(Ac - cC - Bd) - b^3(Ac - cC - Bd) + a^3(Bc + (A - C)d) - 3ab^2(Bc + (A - C)d)) \log(\cos(e + fx))}{f}$$

$$+ \frac{b(2ab(Ac - cC - Bd) + a^2(Bc + (A - C)d) - b^2(Bc + (A - C)d)) \tan(e + fx)}{f}$$

$$+ \frac{(Abc + aBc - bcC + aAd - bBd - aCd)(a + b \tan(e + fx))^2}{2f}$$

$$+ \frac{(Bc + (A - C)d)(a + b \tan(e + fx))^3}{3f}$$

$$- \frac{(aCd - 5b(cC + Bd))(a + b \tan(e + fx))^4}{20b^2f} + \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

output

```
(a^3*(A*c-B*d-C*c)-3*a*b^2*(A*c-B*d-C*c)-3*a^2*b*(B*c+(A-C)*d)+b^3*(B*c+(A-C)*d))*x-(3*a^2*b*(A*c-B*d-C*c)-b^3*(A*c-B*d-C*c)+a^3*(B*c+(A-C)*d)-3*a*b^2*(B*c+(A-C)*d))*ln(cos(f*x+e))/f+b*(2*a*b*(A*c-B*d-C*c)+a^2*(B*c+(A-C)*d)-b^2*(B*c+(A-C)*d))*tan(f*x+e)/f+1/2*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*(a+b*tan(f*x+e))^2/f+1/3*(B*c+(A-C)*d)*(a+b*tan(f*x+e))^3/f-1/20*(C*a*d-5*b*(B*d+C*c))*(a+b*tan(f*x+e))^4/b^2/f+1/5*C*d*tan(f*x+e)*(a+b*tan(f*x+e))^4/b/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.30 (sec) , antiderivative size = 280, normalized size of antiderivative = 0.79

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3(-aCd + 5b(cC + Bd))(a + b \tan(e + fx))^4}{b} + 12Cd \tan(e + fx)(a + b \tan(e + fx))^4 + 30(ABC - aBc - bcC - aAd - \dots)$$

input

```
Integrate[(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x]
+ C*Tan[e + f*x]^2),x]
```

output

```
((3*(-(a*C*d) + 5*b*(c*C + B*d))*(a + b*Tan[e + f*x])^4)/b + 12*C*d*Tan[e
+ f*x]*(a + b*Tan[e + f*x])^4 + 30*(A*b*c - a*B*c - b*c*C - a*A*d - b*B*d
+ a*C*d)*((I*a - b)^3*Log[I - Tan[e + f*x]] - (I*a + b)^3*Log[I + Tan[e +
f*x]] + 6*a*b^2*Tan[e + f*x] + b^3*Tan[e + f*x]^2) - 10*(B*c + (A - C)*d)*
((3*I)*(a + I*b)^4*Log[I - Tan[e + f*x]] - (3*I)*(a - I*b)^4*Log[I + Tan[e
+ f*x]] + 6*b^2*(-6*a^2 + b^2)*Tan[e + f*x] - 12*a*b^3*Tan[e + f*x]^2 - 2
*b^4*Tan[e + f*x]^3))/(60*b*f)
```

Rubi [A] (verified)

Time = 2.70 (sec) , antiderivative size = 369, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.302$, Rules used = {3042, 4120, 25, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4120

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf} - \frac{\int -(a + b \tan(e + fx))^3 (-(aCd - 5b(cC + Bd)) \tan^2(e + fx) + 5b(Bc + (A - C)d) \tan(e + fx) + 5Abc - aCd) dx}{5b}$$

↓ 25

$$\frac{\int (a + b \tan(e + fx))^3 (-(aCd - 5b(cC + Bd)) \tan^2(e + fx) + 5b(Bc + (A - C)d) \tan(e + fx) + 5Abc - aCd) dx}{5bf} - \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))^3 (-(aCd - 5b(cC + Bd)) \tan(e + fx)^2 + 5b(Bc + (A - C)d) \tan(e + fx) + 5Abc - aCd) dx}{5bf} - \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 4113

$$\frac{\int (a + b \tan(e + fx))^3 (5b(Ac - Cc - Bd) + 5b(Bc + (A - C)d) \tan(e + fx)) dx - \frac{(aCd - 5b(Bd + cC))(a + b \tan(e + fx))^4}{4bf}}{5bf} - \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))^3 (5b(Ac - Cc - Bd) + 5b(Bc + (A - C)d) \tan(e + fx)) dx - \frac{(aCd - 5b(Bd + cC))(a + b \tan(e + fx))^4}{4bf}}{5bf} - \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 4011

$$\frac{\int (a + b \tan(e + fx))^2 (5b(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx) - 5b(bBc + b(A - C)d - a(Ac - aAd + bBd))) dx}{5bf} - \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))^2 (5b(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx) - 5b(bBc + b(A - C)d - a(Ac - Cc - Bd)))}{5b}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 4011

$$\frac{\int (a + b \tan(e + fx)) (5b((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd)) + 5b((Bc + (A - C)d) \tan(e + fx)))}{5b}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx)) (5b((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd)) + 5b((Bc + (A - C)d) \tan(e + fx)))}{5b}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 4008

$$\frac{5b(a^3(d(A - C) + Bc) + 3a^2b(Ac - Bd - cC) - 3ab^2(d(A - C) + Bc) - b^3(Ac - Bd - cC)) \int \tan(e + fx) dx + \int (a + b \tan(e + fx))^2 (5b(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx) - 5b(bBc + b(A - C)d - a(Ac - Cc - Bd)))}{5b}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 3042

$$\frac{5b(a^3(d(A - C) + Bc) + 3a^2b(Ac - Bd - cC) - 3ab^2(d(A - C) + Bc) - b^3(Ac - Bd - cC)) \int \tan(e + fx) dx + \int (a + b \tan(e + fx))^2 (5b(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx) - 5b(bBc + b(A - C)d - a(Ac - Cc - Bd)))}{5b}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

↓ 3956

$$\frac{5b^2 \tan(e + fx) (a^2(d(A - C) + Bc) + 2ab(Ac - Bd - cC) - b^2(d(A - C) + Bc))}{f} - \frac{5b \log(\cos(e + fx)) (a^3(d(A - C) + Bc) + 3a^2b(Ac - Bd - cC) - 3ab^2(d(A - C) + Bc) - b^3(Ac - Bd - cC))}{f}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^4}{5bf}$$

input `Int[(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `(C*d*Tan[e + f*x]*(a + b*Tan[e + f*x])^4)/(5*b*f) + (5*b*(a^3*(A*c - c*C - B*d) - 3*a*b^2*(A*c - c*C - B*d) - 3*a^2*b*(B*c + (A - C)*d) + b^3*(B*c + (A - C)*d))*x - (5*b*(3*a^2*b*(A*c - c*C - B*d) - b^3*(A*c - c*C - B*d) + a^3*(B*c + (A - C)*d) - 3*a*b^2*(B*c + (A - C)*d))*Log[Cos[e + f*x]]/f + (5*b^2*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d) - b^2*(B*c + (A - C)*d))*Tan[e + f*x]/f + (5*b*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d)*(a + b*Tan[e + f*x])^2)/(2*f) + (5*b*(B*c + (A - C)*d)*(a + b*Tan[e + f*x])^3)/(3*f) - ((a*C*d - 5*b*(c*C + B*d))*(a + b*Tan[e + f*x])^4)/(4*b*f)/(5*b)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

Maple [A] (warning: unable to verify)

Time = 0.29 (sec) , antiderivative size = 347, normalized size of antiderivative = 0.98

method	result
parts	$\frac{(Aa^3d+3Aa^2bc+B a^3c) \ln(1+\tan(fx+e)^2)}{2f} + \frac{(Bb^3d+3Ca b^2d+C b^3c) \left(\frac{\tan(fx+e)^4}{4} - \frac{\tan(fx+e)^2}{2} + \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f}$
norman	$(Aa^3c - 3Aa^2bd - 3Aa b^2c + Ab^3d - Ba^3d - 3Ba^2bc + 3Ba b^2d + Bb^3c - Ca^3c +$
derivativedivides	$\frac{(Aa^3d+3Aa^2bc-3Aa b^2d-A b^3c+B a^3c-3Ba^2bd-3Ba b^2c+B b^3d-a^3Cd-3Ca^2bc+3Ca b^2d+C b^3c) \ln(1+\tan(fx+e)^2)}{2} + (Aa$
default	$\frac{(Aa^3d+3Aa^2bc-3Aa b^2d-A b^3c+B a^3c-3Ba^2bd-3Ba b^2c+B b^3d-a^3Cd-3Ca^2bc+3Ca b^2d+C b^3c) \ln(1+\tan(fx+e)^2)}{2} + (Aa$
parallelrisc	$\frac{12Cb^3d \tan(fx+e)^5 - 20Cb^3d \tan(fx+e)^3 + 15Bb^3d \tan(fx+e)^4 + 15Cb^3c \tan(fx+e)^4 + 20Ab^3d \tan(fx+e)^3 + 20Bb^3c \tan(fx+e)^3}{2}$
risc	Expression too large to display

input

```
int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x,
method=_RETURNVERBOSE)
```

output

```
1/2*(A*a^3*d+3*A*a^2*b*c+B*a^3*c)/f*ln(1+tan(f*x+e)^2)+(B*b^3*d+3*C*a*b^2*
d+C*b^3*c)/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(f*x+e)^2))+(A
*b^3*d+3*B*a*b^2*d+B*b^3*c+3*C*a^2*b*d+3*C*a*b^2*c)/f*(1/3*tan(f*x+e)^3-ta
n(f*x+e)+arctan(tan(f*x+e)))+(3*A*a^2*b*d+3*A*a*b^2*c+B*a^3*d+3*B*a^2*b*c+
C*a^3*c)/f*(tan(f*x+e)-arctan(tan(f*x+e)))+(3*A*a*b^2*d+A*b^3*c+3*B*a^2*b*
d+3*B*a*b^2*c+C*a^3*d+3*C*a^2*b*c)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)
^2))+A*a^3*c*x+C*b^3*d/f*(1/5*tan(f*x+e)^5-1/3*tan(f*x+e)^3+tan(f*x+e)-arc
tan(tan(f*x+e)))
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.18

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12Cb^3d \tan^5(fx + e) + 15(Cb^3c + (3Cab^2 + Bb^3)d) \tan^4(fx + e) + 20((3Cab^2 + Bb^3)c + (3Ca^2b +$$

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x, algorithm="fricas")
```

output

```
1/60*(12*C*b^3*d*tan(f*x + e)^5 + 15*(C*b^3*c + (3*C*a*b^2 + B*b^3)*d)*tan
(f*x + e)^4 + 20*((3*C*a*b^2 + B*b^3)*c + (3*C*a^2*b + 3*B*a*b^2 + (A - C)
*b^3)*d)*tan(f*x + e)^3 + 60*(((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 +
B*b^3)*c - (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d)*f*x + 3
0*((3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*c + (C*a^3 + 3*B*a^2*b + 3*(A - C)
)*a*b^2 - B*b^3)*d)*tan(f*x + e)^2 - 30*((B*a^3 + 3*(A - C)*a^2*b - 3*B*a*
b^2 - (A - C)*b^3)*c + ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)
*d)*log(1/(tan(f*x + e)^2 + 1)) + 60*((C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2
- B*b^3)*c + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d)*tan(f
*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1001 vs. $2(316) = 632$.

Time = 0.28 (sec) , antiderivative size = 1001, normalized size of antiderivative = 2.84

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))**3*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A**3*c*x + A**3*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*A**3*b*c*log(tan(e + f*x)**2 + 1)/(2*f) - 3*A**3*b*d*x + 3*A**3*b*d*tan(e + f*x)/f - 3*A**3*b**2*c*x + 3*A**3*b**2*c*tan(e + f*x)/f - 3*A**3*b**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*A**3*b**2*d*tan(e + f*x)**2/(2*f) - A**3*b**3*c*log(tan(e + f*x)**2 + 1)/(2*f) + A**3*b**3*c*tan(e + f*x)**2/(2*f) + A**3*b**3*d*x + A**3*b**3*d*tan(e + f*x)**3/(3*f) - A**3*b**3*d*tan(e + f*x)/f + B**3*c*log(tan(e + f*x)**2 + 1)/(2*f) - B**3*d*x + B**3*d*tan(e + f*x)/f - 3*B**3*b*c*x + 3*B**3*b*c*tan(e + f*x)/f - 3*B**3*b*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B**3*b*d*tan(e + f*x)**2/(2*f) - 3*B**3*b**2*c*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B**3*b**2*c*tan(e + f*x)**2/(2*f) + 3*B**3*b**2*d*x + B**3*b**2*d*tan(e + f*x)**3/f - 3*B**3*b**2*d*tan(e + f*x)/f + B**3*b**3*c*x + B**3*b**3*c*tan(e + f*x)**3/(3*f) - B**3*b**3*c*tan(e + f*x)/f + B**3*b**3*d*log(tan(e + f*x)**2 + 1)/(2*f) + B**3*b**3*d*tan(e + f*x)**4/(4*f) - B**3*b**3*d*tan(e + f*x)**2/(2*f) - C**3*c*x + C**3*c*tan(e + f*x)/f - C**3*d*log(tan(e + f*x)**2 + 1)/(2*f) + C**3*d*tan(e + f*x)**2/(2*f) - 3*C**3*b*c*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C**3*b*c*tan(e + f*x)**2/(2*f) + 3*C**3*b*d*x + C**3*b*d*tan(e + f*x)**3/f - 3*C**3*b*d*tan(e + f*x)/f + 3*C**3*b**2*c*x + C**3*b**2*c*tan(e + f*x)**3/f - 3*C**3*b**2*c*tan(e + f*x)/f + 3*C**3*b**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C**3*b**2*d*tan(e + f*x)**4/(4*f) - 3*C**3*b**2*d*tan(e + f*x)**2/(2*f) + C**3*b**3*c*log(tan...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.18

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12 C b^3 d \tan^5(fx + e) + 15 (C b^3 c + (3 C a b^2 + B b^3) d) \tan^4(fx + e) + 20 ((3 C a b^2 + B b^3) c + (3 C a^2 b + B b^3) d) \tan^3(fx + e) + 15 (3 C a^2 c + 3 C a b^2 + B b^3) \tan^2(fx + e) + 10 (3 C a^2 d + 3 C a b^2 + B b^3) \tan(fx + e) + 10 (3 C a^2 c + 3 C a b^2 + B b^3) \tan^2(fx + e) + 10 (3 C a^2 d + 3 C a b^2 + B b^3) \tan^3(fx + e) + 10 (3 C a^2 c + 3 C a b^2 + B b^3) \tan^4(fx + e) + 10 (3 C a^2 d + 3 C a b^2 + B b^3) \tan^5(fx + e)}{1}$$

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
1/60*(12*C*b^3*d*tan(f*x + e)^5 + 15*(C*b^3*c + (3*C*a*b^2 + B*b^3)*d)*tan(f*x + e)^4 + 20*((3*C*a*b^2 + B*b^3)*c + (3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*d)*tan(f*x + e)^3 + 30*((3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*c + (C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*d)*tan(f*x + e)^2 + 60*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c - (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d*(f*x + e) + 30*((B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c + ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*d)*log(tan(f*x + e)^2 + 1) + 60*((C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*c + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d)*tan(f*x + e))/f
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 740 vs. 2(345) = 690.

Time = 0.92 (sec) , antiderivative size = 740, normalized size of antiderivative = 2.10

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
(A*a^3*c - C*a^3*c - 3*B*a^2*b*c - 3*A*a*b^2*c + 3*C*a*b^2*c + B*b^3*c - B
*a^3*d - 3*A*a^2*b*d + 3*C*a^2*b*d + 3*B*a*b^2*d + A*b^3*d - C*b^3*d)*(f*x
+ e)/f + 1/2*(B*a^3*c + 3*A*a^2*b*c - 3*C*a^2*b*c - 3*B*a*b^2*c - A*b^3*c
+ C*b^3*c + A*a^3*d - C*a^3*d - 3*B*a^2*b*d - 3*A*a*b^2*d + 3*C*a*b^2*d +
B*b^3*d)*log(tan(f*x + e)^2 + 1)/f + 1/60*(12*C*b^3*d*f^4*tan(f*x + e)^5
+ 15*C*b^3*c*f^4*tan(f*x + e)^4 + 45*C*a*b^2*d*f^4*tan(f*x + e)^4 + 15*B*b
^3*d*f^4*tan(f*x + e)^4 + 60*C*a*b^2*c*f^4*tan(f*x + e)^3 + 20*B*b^3*c*f^4
*tan(f*x + e)^3 + 60*C*a^2*b*d*f^4*tan(f*x + e)^3 + 60*B*a*b^2*d*f^4*tan(f
*x + e)^3 + 20*A*b^3*d*f^4*tan(f*x + e)^3 - 20*C*b^3*d*f^4*tan(f*x + e)^3
+ 90*C*a^2*b*c*f^4*tan(f*x + e)^2 + 90*B*a*b^2*c*f^4*tan(f*x + e)^2 + 30*A
*b^3*c*f^4*tan(f*x + e)^2 - 30*C*b^3*c*f^4*tan(f*x + e)^2 + 30*C*a^3*d*f^4
*tan(f*x + e)^2 + 90*B*a^2*b*d*f^4*tan(f*x + e)^2 + 90*A*a*b^2*d*f^4*tan(f
*x + e)^2 - 90*C*a*b^2*d*f^4*tan(f*x + e)^2 - 30*B*b^3*d*f^4*tan(f*x + e)^
2 + 60*C*a^3*c*f^4*tan(f*x + e) + 180*B*a^2*b*c*f^4*tan(f*x + e) + 180*A*a
*b^2*c*f^4*tan(f*x + e) - 180*C*a*b^2*c*f^4*tan(f*x + e) - 60*B*b^3*c*f^4*
tan(f*x + e) + 60*B*a^3*d*f^4*tan(f*x + e) + 180*A*a^2*b*d*f^4*tan(f*x + e
) - 180*C*a^2*b*d*f^4*tan(f*x + e) - 180*B*a*b^2*d*f^4*tan(f*x + e) - 60*A
*b^3*d*f^4*tan(f*x + e) + 60*C*b^3*d*f^4*tan(f*x + e))/f^5
```

Mupad [B] (verification not implemented)

Time = 5.68 (sec) , antiderivative size = 477, normalized size of antiderivative = 1.35

$$\begin{aligned}
& \int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= x (Aa^3c + Ab^3d - Ba^3d + Bb^3c - Ca^3c - Cb^3d - 3Aab^2c - 3Aa^2bd - 3Ba^2bc \\
&\quad + 3Bab^2d + 3Cab^2c + 3Ca^2bd) + \frac{\tan(e + fx)^4 \left(\frac{Bb^3d}{4} + \frac{Cb^3c}{4} + \frac{3Cab^2d}{4} \right)}{f} \\
&\quad + \frac{\tan(e + fx)^3 \left(\frac{Ab^3d}{3} + \frac{Bb^3c}{3} - \frac{Cb^3d}{3} + Bab^2d + Cab^2c + Ca^2bd \right)}{f} \\
&\quad + \frac{\tan(e + fx)^2 \left(\frac{Ab^3c}{2} - \frac{Bb^3d}{2} + \frac{Ca^3d}{2} - \frac{Cb^3c}{2} + \frac{3Aab^2d}{2} + \frac{3Bab^2c}{2} + \frac{3Ba^2bd}{2} + \frac{3Ca^2bc}{2} - \frac{3Cab^2d}{2} \right)}{f} \\
&\quad + \frac{\ln(\tan(e + fx)^2 + 1) \left(\frac{Aa^3d}{2} - \frac{Ab^3c}{2} + \frac{Ba^3c}{2} + \frac{Bb^3d}{2} - \frac{Ca^3d}{2} + \frac{Cb^3c}{2} + \frac{3Aa^2bc}{2} - \frac{3Aab^2d}{2} - \frac{3Ba^2bc}{2} - \dots \right)}{f} \\
&\quad + \frac{\tan(e + fx) (Ba^3d - Ab^3d - Bb^3c + Ca^3c + Cb^3d + 3Aab^2c + 3Aa^2bd + 3Ba^2bc - 3Bab^2c)}{f} \\
&\quad + \frac{Cb^3d \tan(e + fx)^5}{5f}
\end{aligned}$$

input

```
int((a + b*tan(e + f*x))^3*(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
x*(A*a^3*c + A*b^3*d - B*a^3*d + B*b^3*c - C*a^3*c - C*b^3*d - 3*A*a*b^2*c
- 3*A*a^2*b*d - 3*B*a^2*b*c + 3*B*a*b^2*d + 3*C*a*b^2*c + 3*C*a^2*b*d) +
(tan(e + f*x)^4*((B*b^3*d)/4 + (C*b^3*c)/4 + (3*C*a*b^2*d)/4))/f + (tan(e
+ f*x)^3*((A*b^3*d)/3 + (B*b^3*c)/3 - (C*b^3*d)/3 + B*a*b^2*d + C*a*b^2*c
+ C*a^2*b*d))/f + (tan(e + f*x)^2*((A*b^3*c)/2 - (B*b^3*d)/2 + (C*a^3*d)/2
- (C*b^3*c)/2 + (3*A*a*b^2*d)/2 + (3*B*a*b^2*c)/2 + (3*B*a^2*b*d)/2 + (3*
C*a^2*b*c)/2 - (3*C*a*b^2*d)/2))/f + (log(tan(e + f*x)^2 + 1)*((A*a^3*d)/2
- (A*b^3*c)/2 + (B*a^3*c)/2 + (B*b^3*d)/2 - (C*a^3*d)/2 + (C*b^3*c)/2 + (
3*A*a^2*b*c)/2 - (3*A*a*b^2*d)/2 - (3*B*a*b^2*c)/2 - (3*B*a^2*b*d)/2 - (3*
C*a^2*b*c)/2 + (3*C*a*b^2*d)/2))/f + (tan(e + f*x)*(B*a^3*d - A*b^3*d - B*
b^3*c + C*a^3*c + C*b^3*d + 3*A*a*b^2*c + 3*A*a^2*b*d + 3*B*a^2*b*c - 3*B*
a*b^2*d - 3*C*a*b^2*c - 3*C*a^2*b*d))/f + (C*b^3*d*tan(e + f*x)^5)/(5*f)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 614, normalized size of antiderivative = 1.74

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{180a^2bcdfx + 120 \log(\tan(fx + e)^2 + 1) a^3bc - 30 \log(\tan(fx + e)^2 + 1) a^3cd - 180 \log(\tan(fx + e)^2 + 1) a^2b^2cd + 120 \log(\tan(fx + e)^2 + 1) a^2b^3cd - 90 \log(\tan(fx + e)^2 + 1) a^2b^2c^2d - 120 \log(\tan(fx + e)^2 + 1) a^2b^3c^2d + 30 \log(\tan(fx + e)^2 + 1) b^4cd + 45 \tan(fx + e) a^4b^2cd + 15 \tan(fx + e) a^4b^3cd + 15 \tan(fx + e) a^4b^3c^2d + 60 \tan(fx + e) a^3a^2b^2cd + 80 \tan(fx + e) a^3a^2b^3cd + 60 \tan(fx + e) a^3a^2b^2c^2d + 20 \tan(fx + e) a^3b^4cd - 20 \tan(fx + e) a^3b^3c^2d + 30 \tan(fx + e) a^2a^3cd + 180 \tan(fx + e) a^2a^2b^2cd + 90 \tan(fx + e) a^2a^2b^3cd + 120 \tan(fx + e) a^2a^2b^3c^2d - 90 \tan(fx + e) a^2a^2b^2c^2d - 30 \tan(fx + e) a^2b^4cd - 30 \tan(fx + e) a^2b^3c^2d + 240 \tan(fx + e) a^3b^2cd + 60 \tan(fx + e) a^3b^3cd + 360 \tan(fx + e) a^2b^2cd - 180 \tan(fx + e) a^2b^2c^2d - 240 \tan(fx + e) a^2b^3cd - 180 \tan(fx + e) a^2b^2c^2d - 60 \tan(fx + e) b^4cd + 60 \tan(fx + e) b^3cd + 60 a^4c^2fx - 240 a^3b^2d^2fx - 60 a^3c^2d^2fx - 360 a^2b^2c^2fx + 180 a^2b^2cd^2fx + 240 a^2b^3d^2fx + 180 a^2b^2c^2d^2fx + 60 b^4c^2fx - 60 b^3c^2d^2fx}{(60f)}$$

input

```
int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
(30*log(tan(e + f*x)**2 + 1)*a**4*d + 120*log(tan(e + f*x)**2 + 1)*a**3*b*c - 30*log(tan(e + f*x)**2 + 1)*a**3*c*d - 180*log(tan(e + f*x)**2 + 1)*a**2*b**2*d - 90*log(tan(e + f*x)**2 + 1)*a**2*b*c**2 - 120*log(tan(e + f*x)**2 + 1)*a*b**3*c + 90*log(tan(e + f*x)**2 + 1)*a*b**2*c*d + 30*log(tan(e + f*x)**2 + 1)*b**4*d + 30*log(tan(e + f*x)**2 + 1)*b**3*c**2 + 12*tan(e + f*x)**5*b**3*c*d + 45*tan(e + f*x)**4*a*b**2*c*d + 15*tan(e + f*x)**4*b**4*d + 15*tan(e + f*x)**4*b**3*c**2 + 60*tan(e + f*x)**3*a**2*b*c*d + 80*tan(e + f*x)**3*a*b**3*d + 60*tan(e + f*x)**3*a*b**2*c**2 + 20*tan(e + f*x)**3*b**4*c - 20*tan(e + f*x)**3*b**3*c*d + 30*tan(e + f*x)**2*a**3*c*d + 180*tan(e + f*x)**2*a**2*b**2*d + 90*tan(e + f*x)**2*a**2*b*c**2 + 120*tan(e + f*x)**2*a*b**3*c - 90*tan(e + f*x)**2*a*b**2*c*d - 30*tan(e + f*x)**2*b**4*d - 30*tan(e + f*x)**2*b**3*c**2 + 240*tan(e + f*x)*a**3*b*d + 60*tan(e + f*x)*a**3*c**2 + 360*tan(e + f*x)*a**2*b**2*c - 180*tan(e + f*x)*a**2*b*c*d - 240*tan(e + f*x)*a*b**3*d - 180*tan(e + f*x)*a*b**2*c**2 - 60*tan(e + f*x)*b**4*c + 60*tan(e + f*x)*b**3*c*d + 60*a**4*c*f*x - 240*a**3*b*d*f*x - 60*a**3*c**2*f*x - 360*a**2*b**2*c*f*x + 180*a**2*b*c*d*f*x + 240*a**2*b**3*d*f*x + 180*a*b**2*c**2*f*x + 60*b**4*c*f*x - 60*b**3*c*d*f*x)/(60*f)
```


3.51 $\int (a+b \tan(e+fx))^2(c+d \tan(e+fx)) (A + B \tan(e +$

Optimal result	556
Mathematica [C] (verified)	557
Rubi [A] (verified)	557
Maple [A] (warning: unable to verify)	561
Fricas [A] (verification not implemented)	562
Sympy [B] (verification not implemented)	562
Maxima [A] (verification not implemented)	563
Giac [A] (verification not implemented)	564
Mupad [B] (verification not implemented)	565
Reduce [B] (verification not implemented)	565

Optimal result

Integrand size = 43, antiderivative size = 248

$$\int (a + b \tan(e + fx))^2(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(a^2(Ac - cC - Bd) - b^2(Ac - cC - Bd) - 2ab(Bc + (A - C)d)) x}{f} - \frac{(2ab(Ac - cC - Bd) + a^2(Bc + (A - C)d) - b^2(Bc + (A - C)d)) \log(\cos(e + fx))}{f}$$

$$+ \frac{b(ABC + aBc - bcC + aAd - bBd - aCd) \tan(e + fx)}{f}$$

$$+ \frac{(Bc + (A - C)d)(a + b \tan(e + fx))^2}{2f}$$

$$- \frac{(aCd - 4b(cC + Bd))(a + b \tan(e + fx))^3}{12b^2f} + \frac{Cd \tan(e + fx)(a + b \tan(e + fx))^3}{4bf}$$

output

```
(a^2*(A*c-B*d-C*c)-b^2*(A*c-B*d-C*c)-2*a*b*(B*c+(A-C)*d))*x-(2*a*b*(A*c-B*d-C*c)+a^2*(B*c+(A-C)*d)-b^2*(B*c+(A-C)*d))*ln(cos(f*x+e))/f+b*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*tan(f*x+e)/f+1/2*(B*c+(A-C)*d)*(a+b*tan(f*x+e))^2/f-1/12*(C*a*d-4*b*(B*d+C*c))*(a+b*tan(f*x+e))^3/b^2/f+1/4*C*d*tan(f*x+e)*(a+b*tan(f*x+e))^3/b/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.23 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.98

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(-aCd + 4b(cC + Bd))(a + b \tan(e + fx))^3}{b} + 3Cd \tan(e + fx)(a + b \tan(e + fx))^3 - 6(ABC - aBc - bcC - aAd - b$$

input

```
Integrate[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x]
+ C*Tan[e + f*x]^2),x]
```

output

```
(((-(a*C*d) + 4*b*(c*C + B*d))*(a + b*Tan[e + f*x])^3)/b + 3*C*d*Tan[e + f
*x]*(a + b*Tan[e + f*x])^3 - 6*(A*b*c - a*B*c - b*c*C - a*A*d - b*B*d + a*
C*d)*(I*((a + I*b)^2*Log[I - Tan[e + f*x]] - (a - I*b)^2*Log[I + Tan[e + f
*x]]) - 2*b^2*Tan[e + f*x]) + 6*(B*c + (A - C)*d)*((I*a - b)^3*Log[I - Tan
[e + f*x]] - (I*a + b)^3*Log[I + Tan[e + f*x]] + 6*a*b^2*Tan[e + f*x] + b^
3*Tan[e + f*x]^2))/(12*b*f)
```

Rubi [A] (verified)

Time = 1.87 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.256$, Rules used = {3042, 4120, 25, 3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4120

$$\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^3}{4bf} - \frac{\int -(a+b \tan(e+fx))^2 (-((aCd-4b(cC+Bd)) \tan^2(e+fx)) + 4b(Bc+(A-C)d) \tan(e+fx) + 4Abc - aCd)}{4b}$$

↓ 25

$$\frac{\int (a+b \tan(e+fx))^2 (-((aCd-4b(cC+Bd)) \tan^2(e+fx)) + 4b(Bc+(A-C)d) \tan(e+fx) + 4Abc - aCd)}{4b} - \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^3}{4bf}$$

↓ 3042

$$\frac{\int (a+b \tan(e+fx))^2 (-((aCd-4b(cC+Bd)) \tan(e+fx)^2) + 4b(Bc+(A-C)d) \tan(e+fx) + 4Abc - aCd)}{4b} - \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^3}{4bf}$$

↓ 4113

$$\frac{\int (a+b \tan(e+fx))^2 (4b(Ac-Cc-Bd) + 4b(Bc+(A-C)d) \tan(e+fx)) dx - \frac{(aCd-4b(Bd+cC))(a+b \tan(e+fx))^3}{3bf}}{4b} - \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^3}{4bf}$$

↓ 3042

$$\frac{\int (a+b \tan(e+fx))^2 (4b(Ac-Cc-Bd) + 4b(Bc+(A-C)d) \tan(e+fx)) dx - \frac{(aCd-4b(Bd+cC))(a+b \tan(e+fx))^3}{3bf}}{4b} - \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^3}{4bf}$$

↓ 4011

$$\frac{\int (a+b \tan(e+fx))(4b(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx) - 4b(bBc+b(A-C)d - a(Ac-Cc-Bd))) dx}{4b} - \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^3}{4bf}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))(4b(abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx) - 4b(bBc + b(A - C)d - a(Ac - C))}{4b}$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^3}{4bf}$$

↓ 4008

$$4b(a^2(d(A - C) + Bc) + 2ab(Ac - Bd - cC) - b^2(d(A - C) + Bc)) \int \tan(e + fx)dx + 4bx(a^2(Ac - Bd - cC))$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^3}{4bf}$$

↓ 3042

$$4b(a^2(d(A - C) + Bc) + 2ab(Ac - Bd - cC) - b^2(d(A - C) + Bc)) \int \tan(e + fx)dx + 4bx(a^2(Ac - Bd - cC))$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^3}{4bf}$$

↓ 3956

$$\frac{-4b \log(\cos(e + fx))(a^2(d(A - C) + Bc) + 2ab(Ac - Bd - cC) - b^2(d(A - C) + Bc))}{f} + 4bx(a^2(Ac - Bd - cC) - 2ab(d(A - C) + Bc))$$

$$\frac{Cd \tan(e + fx)(a + b \tan(e + fx))^3}{4bf}$$

input

```
Int[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(C*d*Tan[e + f*x]*(a + b*Tan[e + f*x])^3)/(4*b*f) + (4*b*(a^2*(A*c - c*C - B*d) - b^2*(A*c - c*C - B*d) - 2*a*b*(B*c + (A - C)*d))*x - (4*b*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d) - b^2*(B*c + (A - C)*d))*Log[Cos[e + f*x]]/f + (4*b^2*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d)*Tan[e + f*x])/f + (2*b*(B*c + (A - C)*d)*(a + b*Tan[e + f*x])^2)/f - ((a*C*d - 4*b*(c*C + B*d))*(a + b*Tan[e + f*x])^3)/(3*b*f)/(4*b)
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)])^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

Maple [A] (warning: unable to verify)

Time = 0.21 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.99

method	result
parts	$\frac{(A a^2 d + 2 A a b c + B a^2 c) \ln(1 + \tan(f x + e)^2)}{2 f} + \frac{(B b^2 d + 2 C a b d + C b^2 c) \left(\frac{\tan(f x + e)^3}{3} - \tan(f x + e) + \arctan(\tan(f x + e)) \right)}{f}$
norman	$(A a^2 c - 2 A a b d - A b^2 c - B a^2 d - 2 B a b c + B b^2 d - C a^2 c + 2 C a b d + C b^2 c) x + \frac{(2 A a^2 d + 2 A a b c + B a^2 c) \ln(1 + \tan(f x + e)^2)}{2 f} + \frac{(B b^2 d + 2 C a b d + C b^2 c) \left(\frac{\tan(f x + e)^3}{3} - \tan(f x + e) + \arctan(\tan(f x + e)) \right)}{f}$
derivativedivides	$\frac{\frac{d C b^2 \tan(f x + e)^4}{4} + \frac{B b^2 d \tan(f x + e)^3}{3} + \frac{2 C a b d \tan(f x + e)^3}{3} + \frac{C b^2 c \tan(f x + e)^3}{3} + \frac{A b^2 d \tan(f x + e)^2}{2} + B a b d \tan(f x + e)^2 + \frac{B b^2 c}{2}}{\frac{d C b^2 \tan(f x + e)^4}{4} + \frac{B b^2 d \tan(f x + e)^3}{3} + \frac{2 C a b d \tan(f x + e)^3}{3} + \frac{C b^2 c \tan(f x + e)^3}{3} + \frac{A b^2 d \tan(f x + e)^2}{2} + B a b d \tan(f x + e)^2 + \frac{B b^2 c}{2}}$
default	$\frac{d C b^2 \tan(f x + e)^4}{4} + \frac{B b^2 d \tan(f x + e)^3}{3} + \frac{2 C a b d \tan(f x + e)^3}{3} + \frac{C b^2 c \tan(f x + e)^3}{3} + \frac{A b^2 d \tan(f x + e)^2}{2} + B a b d \tan(f x + e)^2 + \frac{B b^2 c}{2}$
parallelrisc	$\frac{3 d C b^2 \tan(f x + e)^4 + 4 B b^2 d \tan(f x + e)^3 + 4 C b^2 c \tan(f x + e)^3 + 6 A b^2 d \tan(f x + e)^2 + 6 B b^2 c \tan(f x + e)^2 + 6 C a^2 d \tan(f x + e)}{\frac{d C b^2 \tan(f x + e)^4}{4} + \frac{B b^2 d \tan(f x + e)^3}{3} + \frac{2 C a b d \tan(f x + e)^3}{3} + \frac{C b^2 c \tan(f x + e)^3}{3} + \frac{A b^2 d \tan(f x + e)^2}{2} + B a b d \tan(f x + e)^2 + \frac{B b^2 c}{2}}$
risc	Expression too large to display

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,
method=_RETURNVERBOSE)
```

output

```
1/2*(A*a^2*d+2*A*a*b*c+B*a^2*c)/f*ln(1+tan(f*x+e)^2)+(B*b^2*d+2*C*a*b*d+C*
b^2*c)/f*(1/3*tan(f*x+e)^3-tan(f*x+e)+arctan(tan(f*x+e)))+(A*b^2*d+2*B*a*b
*d+B*b^2*c+C*a^2*d+2*C*a*b*c)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)^2))+
(2*A*a*b*d+A*b^2*c+B*a^2*d+2*B*a*b*c+C*a^2*c)/f*(tan(f*x+e)-arctan(tan(f*x
+e)))+A*a^2*c*x+d*C*b^2/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(
f*x+e)^2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.10

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3Cb^2d \tan(fx + e)^4 + 4(Cb^2c + (2Cab + Bb^2)d) \tan(fx + e)^3 + 12(((A - C)a^2 - 2Bab - (A - C)b^2$$

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

```
1/12*(3*C*b^2*d*tan(f*x + e)^4 + 4*(C*b^2*c + (2*C*a*b + B*b^2)*d)*tan(f*x + e)^3 + 12*(((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c - (B*a^2 + 2*(A - C)*a*b - B*b^2)*d)*f*x + 6*((2*C*a*b + B*b^2)*c + (C*a^2 + 2*B*a*b + (A - C)*b^2)*d)*tan(f*x + e)^2 - 6*((B*a^2 + 2*(A - C)*a*b - B*b^2)*c + ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d)*log(1/(tan(f*x + e)^2 + 1)) + 12*((C*a^2 + 2*B*a*b + (A - C)*b^2)*c + (B*a^2 + 2*(A - C)*a*b - B*b^2)*d)*tan(f*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 617 vs. 2(218) = 436.

Time = 0.21 (sec) , antiderivative size = 617, normalized size of antiderivative = 2.49

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))**2*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A**2*c*x + A**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + A*b*c*
log(tan(e + f*x)**2 + 1)/f - 2*A*b*d*x + 2*A*b*d*tan(e + f*x)/f - A*b*
**2*c*x + A*b**2*c*tan(e + f*x)/f - A*b**2*d*log(tan(e + f*x)**2 + 1)/(2*f)
+ A*b**2*d*tan(e + f*x)**2/(2*f) + B**2*c*log(tan(e + f*x)**2 + 1)/(2*f)
) - B**2*d*x + B**2*d*tan(e + f*x)/f - 2*B*b*c*x + 2*B*b*c*tan(e +
f*x)/f - B*a*b*d*log(tan(e + f*x)**2 + 1)/f + B*a*b*d*tan(e + f*x)**2/f -
B*b**2*c*log(tan(e + f*x)**2 + 1)/(2*f) + B*b**2*c*tan(e + f*x)**2/(2*f)
+ B*b**2*d*x + B*b**2*d*tan(e + f*x)**3/(3*f) - B*b**2*d*tan(e + f*x)/f -
C**2*c*x + C**2*c*tan(e + f*x)/f - C**2*d*log(tan(e + f*x)**2 + 1)/(
2*f) + C**2*d*tan(e + f*x)**2/(2*f) - C*a*b*c*log(tan(e + f*x)**2 + 1)/f
+ C*a*b*c*tan(e + f*x)**2/f + 2*C*a*b*d*x + 2*C*a*b*d*tan(e + f*x)**3/(3*
f) - 2*C*a*b*d*tan(e + f*x)/f + C*b**2*c*x + C*b**2*c*tan(e + f*x)**3/(3*f)
) - C*b**2*c*tan(e + f*x)/f + C*b**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + C*
b**2*d*tan(e + f*x)**4/(4*f) - C*b**2*d*tan(e + f*x)**2/(2*f), Ne(f, 0)),
(x*(a + b*tan(e))**2*(c + d*tan(e))*(A + B*tan(e) + C*tan(e)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.10

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3Cb^2d \tan(fx + e)^4 + 4(Cb^2c + (2Cab + Bb^2)d) \tan(fx + e)^3 + 6((2Cab + Bb^2)c + (Ca^2 + 2Bab +$$

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x, algorithm="maxima")
```

output

```
1/12*(3*C*b^2*d*tan(f*x + e)^4 + 4*(C*b^2*c + (2*C*a*b + B*b^2)*d)*tan(f*x
+ e)^3 + 6*((2*C*a*b + B*b^2)*c + (C*a^2 + 2*B*a*b + (A - C)*b^2)*d)*tan(
f*x + e)^2 + 12*(((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c - (B*a^2 + 2*(A -
C)*a*b - B*b^2)*d)*(f*x + e) + 6*((B*a^2 + 2*(A - C)*a*b - B*b^2)*c + ((A
- C)*a^2 - 2*B*a*b - (A - C)*b^2)*d)*log(tan(f*x + e)^2 + 1) + 12*(((C*a^2
+ 2*B*a*b + (A - C)*b^2)*c + (B*a^2 + 2*(A - C)*a*b - B*b^2)*d)*tan(f*x +
e))/f
```


Giac [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.81

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(Aa^2c - Ca^2c - 2Babc - Ab^2c + Cb^2c - Ba^2d - 2Aabd + 2Cabd + Bb^2d)(fx + e)}{f}$$

$$+ \frac{(Ba^2c + 2Aabc - 2Cabc - Bb^2c + Aa^2d - Ca^2d - 2Babd - Ab^2d + Cb^2d) \log(\tan(fx + e)^2 + 1)}{2f}$$

$$+ \frac{3Cb^2df^3 \tan(fx + e)^4 + 4Cb^2cf^3 \tan(fx + e)^3 + 8Cabdf^3 \tan(fx + e)^3 + 4Bb^2df^3 \tan(fx + e)^3 + 12C^2abdf^3 \tan(fx + e)^2 + 12C^2abdf^3 \tan(fx + e) + 12C^2abdf^3 \tan(fx + e) - 12C^2abdf^3 \tan(fx + e) - 12C^2abdf^3 \tan(fx + e) - 12C^2abdf^3 \tan(fx + e)}{f^4}$$

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
(A*a^2*c - C*a^2*c - 2*B*a*b*c - A*b^2*c + C*b^2*c - B*a^2*d - 2*A*a*b*d + 2*C*a*b*d + B*b^2*d)*(f*x + e)/f + 1/2*(B*a^2*c + 2*A*a*b*c - 2*C*a*b*c - B*b^2*c + A*a^2*d - C*a^2*d - 2*B*a*b*d - A*b^2*d + C*b^2*d)*log(tan(f*x + e)^2 + 1)/f + 1/12*(3*C*b^2*d*f^3*tan(f*x + e)^4 + 4*C*b^2*c*f^3*tan(f*x + e)^3 + 8*C*a*b*d*f^3*tan(f*x + e)^3 + 4*B*b^2*d*f^3*tan(f*x + e)^3 + 12*C*a*b*c*f^3*tan(f*x + e)^2 + 6*B*b^2*c*f^3*tan(f*x + e)^2 + 6*C*a^2*d*f^3*tan(f*x + e)^2 + 12*B*a*b*d*f^3*tan(f*x + e)^2 + 6*A*b^2*d*f^3*tan(f*x + e)^2 - 6*C*b^2*d*f^3*tan(f*x + e)^2 + 12*C*a^2*c*f^3*tan(f*x + e) + 24*B*a*b*c*f^3*tan(f*x + e) + 12*A*b^2*c*f^3*tan(f*x + e) - 12*C*b^2*c*f^3*tan(f*x + e) + 12*B*a^2*d*f^3*tan(f*x + e) + 24*A*a*b*d*f^3*tan(f*x + e) - 24*C*a*b*d*f^3*tan(f*x + e) - 12*B*b^2*d*f^3*tan(f*x + e))/f^4
```

Mupad [B] (verification not implemented)

Time = 5.54 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.21

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{\tan(e + fx)^2 \left(\frac{Ab^2d}{2} + \frac{Bb^2c}{2} + \frac{Ca^2d}{2} - \frac{Cb^2d}{2} + Babd + Cabc \right)}{f} - \frac{x (Ab^2c - Aa^2c + Ba^2d + Ca^2c - Bb^2d - Cb^2c + 2Aabd + 2Babc - 2Cabd)}{\ln(\tan(e + fx)^2 + 1) \left(\frac{Ab^2d}{2} - \frac{Ba^2c}{2} - \frac{Aa^2d}{2} + \frac{Bb^2c}{2} + \frac{Ca^2d}{2} - \frac{Cb^2d}{2} - Aabc + Babd + Cabc \right)} + \frac{\tan(e + fx) (Ab^2c + Ba^2d + Ca^2c - Bb^2d - Cb^2c + 2Aabd + 2Babc - 2Cabd)}{f} + \frac{\tan(e + fx)^3 \left(\frac{Bb^2d}{3} + \frac{Cb^2c}{3} + \frac{2Cabd}{3} \right)}{f} + \frac{Cb^2d \tan(e + fx)^4}{4f}$$

input

```
int((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
(tan(e + f*x)^2*((A*b^2*d)/2 + (B*b^2*c)/2 + (C*a^2*d)/2 - (C*b^2*d)/2 + B*a*b*d + C*a*b*c))/f - x*(A*b^2*c - A*a^2*c + B*a^2*d + C*a^2*c - B*b^2*d - C*b^2*c + 2*A*a*b*d + 2*B*a*b*c - 2*C*a*b*d) - (log(tan(e + f*x)^2 + 1)*((A*b^2*d)/2 - (B*a^2*c)/2 - (A*a^2*d)/2 + (B*b^2*c)/2 + (C*a^2*d)/2 - (C*b^2*d)/2 - A*a*b*c + B*a*b*d + C*a*b*c))/f + (tan(e + f*x)*(A*b^2*c + B*a^2*d + C*a^2*c - B*b^2*d - C*b^2*c + 2*A*a*b*d + 2*B*a*b*c - 2*C*a*b*d))/f + (tan(e + f*x)^3*((B*b^2*d)/3 + (C*b^2*c)/3 + (2*C*a*b*d)/3))/f + (C*b^2*d*tan(e + f*x)^4)/(4*f)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.62

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{-6 \tan^2(fx + e) b^2 cd + 24 abcd fx - 12 \tan(fx + e) b^2 c^2 - 24 \tan(fx + e) abcd + 18 \log(\tan(fx + e))^2 - \dots}{f}$$

input `int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `(6*log(tan(e + f*x)**2 + 1)*a**3*d + 18*log(tan(e + f*x)**2 + 1)*a**2*b*c - 6*log(tan(e + f*x)**2 + 1)*a**2*c*d - 18*log(tan(e + f*x)**2 + 1)*a*b**2*d - 12*log(tan(e + f*x)**2 + 1)*a*b*c**2 - 6*log(tan(e + f*x)**2 + 1)*b**3*c + 6*log(tan(e + f*x)**2 + 1)*b**2*c*d + 3*tan(e + f*x)**4*b**2*c*d + 8*tan(e + f*x)**3*a*b*c*d + 4*tan(e + f*x)**3*b**3*d + 4*tan(e + f*x)**3*b**2*c**2 + 6*tan(e + f*x)**2*a**2*c*d + 18*tan(e + f*x)**2*a*b**2*d + 12*tan(e + f*x)**2*a*b*c**2 + 6*tan(e + f*x)**2*b**3*c - 6*tan(e + f*x)**2*b**2*c*d + 36*tan(e + f*x)*a**2*b*d + 12*tan(e + f*x)*a**2*c**2 + 36*tan(e + f*x)*a*b**2*c - 24*tan(e + f*x)*a*b*c*d - 12*tan(e + f*x)*b**3*d - 12*tan(e + f*x)*b**2*c**2 + 12*a**3*c*f*x - 36*a**2*b*d*f*x - 12*a**2*c**2*f*x - 36*a*b**2*c*f*x + 24*a*b*c*d*f*x + 12*b**3*d*f*x + 12*b**2*c**2*f*x)/(12*f)`

3.52 $\int (a+b \tan(e+fx))(c+d \tan(e+fx)) (A + B \tan(e +$

Optimal result	567
Mathematica [C] (verified)	568
Rubi [A] (verified)	568
Maple [A] (warning: unable to verify)	571
Fricas [A] (verification not implemented)	572
Sympy [B] (verification not implemented)	572
Maxima [A] (verification not implemented)	573
Giac [A] (verification not implemented)	574
Mupad [B] (verification not implemented)	574
Reduce [B] (verification not implemented)	575

Optimal result

Integrand size = 41, antiderivative size = 161

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= (a(Ac - cC - Bd) - b(Bc + (A - C)d))x$$

$$- \frac{(Abc + aBc - bcC + aAd - bBd - aCd) \log(\cos(e + fx))}{f}$$

$$+ \frac{(Ab + aB - bC)d \tan(e + fx)}{f} - \frac{(bcC - 3bBd - 3aCd)(c + d \tan(e + fx))^2}{6d^2 f}$$

$$+ \frac{bC \tan(e + fx)(c + d \tan(e + fx))^2}{3df}$$

output

```
(a*(A*c-B*d-C*c)-b*(B*c+(A-C)*d))*x-(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*
ln(cos(f*x+e))/f+(A*b+B*a-C*b)*d*tan(f*x+e)/f-1/6*(-3*B*b*d-3*C*a*d+C*b*c)
*(c+d*tan(f*x+e))^2/d^2/f+1/3*b*C*tan(f*x+e)*(c+d*tan(f*x+e))^2/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.06 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.00

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3(a + ib)(A + iB - C)(-ic + d) \log(i - \tan(e + fx)) + 3(a - ib)(A - iB - C)(ic + d) \log(i + \tan(e + fx))}{6f}$$

input

```
Integrate[(a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(3*(a + I*b)*(A + I*B - C)*((-I)*c + d)*Log[I - Tan[e + f*x]] + 3*(a - I*b)*(A - I*B - C)*(I*c + d)*Log[I + Tan[e + f*x]] + 6*(A*b + a*B - b*C)*d*Tan[e + f*x] + ((-(b*c*C) + 3*b*B*d + 3*a*C*d)*(c + d*Tan[e + f*x])^2)/d^2 + (2*b*C*Tan[e + f*x]*(c + d*Tan[e + f*x])^2)/d)/(6*f)
```

Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.195$, Rules used = {3042, 4120, 3042, 4113, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow \text{4120}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{\int (c+d \tan(e+fx)) \left((bcC - 3adC - 3bBd) \tan^2(e+fx) - 3(Ab - Cb + aB)d \tan(e+fx) + bcC - 3aAd \right) dx} \frac{3d}{3d}$$

↓ 3042

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{\int (c+d \tan(e+fx)) \left((bcC - 3adC - 3bBd) \tan(e+fx)^2 - 3(Ab - Cb + aB)d \tan(e+fx) + bcC - 3aAd \right) dx} \frac{3d}{3d}$$

↓ 4113

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{\int (c+d \tan(e+fx)) (3(bB - a(A - C))d - 3(Ab - Cb + aB)d \tan(e+fx)) dx + \frac{(-3aCd - 3bBd + bcC)(c+d \tan(e+fx))}{2df}} \frac{3d}{3d}$$

↓ 3042

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{\int (c+d \tan(e+fx)) (3(bB - a(A - C))d - 3(Ab - Cb + aB)d \tan(e+fx)) dx + \frac{(-3aCd - 3bBd + bcC)(c+d \tan(e+fx))}{2df}} \frac{3d}{3d}$$

↓ 4008

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{-3d(aAd + aBc - aCd + Abc - bBd - bcC) \int \tan(e+fx) dx + 3dx(-a(Ac - Bd - cC) + bd(A - C) + bBc)} \frac{3d}{3d}$$

↓ 3042

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{-3d(aAd + aBc - aCd + Abc - bBd - bcC) \int \tan(e+fx) dx + 3dx(-a(Ac - Bd - cC) + bd(A - C) + bBc)} \frac{3d}{3d}$$

↓ 3956

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^2}{3df}}{\frac{3d \log(\cos(e+fx))(aAd + aBc - aCd + Abc - bBd - bcC)}{f} + 3dx(-a(Ac - Bd - cC) + bd(A - C) + bBc) - \frac{3d^2 \tan(e+fx)(aB + Ab)}{f}} \frac{3d}{3d}$$

input `Int[(a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `(b*C*Tan[e + f*x]*(c + d*Tan[e + f*x])^2)/(3*d*f) - (3*d*(b*B*c + b*(A - C)*d - a*(A*c - c*C - B*d))*x + (3*d*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d)*Log[Cos[e + f*x]])/f - (3*(A*b + a*B - b*C)*d^2*Tan[e + f*x])/f + ((b*c*C - 3*b*B*d - 3*a*C*d)*(c + d*Tan[e + f*x])^2)/(2*d*f)/(3*d)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :=> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :=> Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=> Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)])^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

Maple [A] (warning: unable to verify)

Time = 0.14 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.91

method	result
parts	$\frac{(Aad+Abc+Bac) \ln(1+\tan(fx+e)^2)}{2f} + \frac{(Bbd+Cad+Cbc) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f} + \frac{(Abd+Bad+Bbc)}{f}$
norman	$(Aac - Abd - Bad - Bbc - Cac + Cbd) x + \frac{(Abd+Bad+Bbc+Cac-Cbd) \tan(fx+e)}{f} + \frac{(Bbd+Cad+Cbc)}{f}$
derivativedivides	$\frac{C \tan(fx+e)^3 bd}{3} + \frac{B \tan(fx+e)^2 bd}{2} + \frac{C \tan(fx+e)^2 ad}{2} + \frac{C \tan(fx+e)^2 bc}{2} + A \tan(fx+e) bd + B \tan(fx+e) ad + B \tan(fx+e) b$
default	$\frac{C \tan(fx+e)^3 bd}{3} + \frac{B \tan(fx+e)^2 bd}{2} + \frac{C \tan(fx+e)^2 ad}{2} + \frac{C \tan(fx+e)^2 bc}{2} + A \tan(fx+e) bd + B \tan(fx+e) ad + B \tan(fx+e) b$
parallelrisch	$\frac{2C \tan(fx+e)^3 bd + 6Aacfx - 6Abdfx - 6Badfx - 6Bbcfx + 3B \tan(fx+e)^2 bd - 6Cacfx + 6Cbdfx + 3C \tan(fx+e)^2 ad + 3C \tan(fx+e) b}{f}$
risch	$-Badx + Cbd x + \frac{2i(-3iCbc e^{4i(fx+e)} - 3iBbd e^{2i(fx+e)} - 3iCad e^{4i(fx+e)} + 3Abd e^{4i(fx+e)} + 3Bade^{4i(fx+e)})}{f}$

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,me
thod=_RETURNVERBOSE)
```

output

```
1/2*(A*a*d+A*b*c+B*a*c)/f*ln(1+tan(f*x+e)^2)+(B*b*d+C*a*d+C*b*c)/f*(1/2*ta
n(f*x+e)^2-1/2*ln(1+tan(f*x+e)^2))+(A*b*d+B*a*d+B*b*c+C*a*c)/f*(tan(f*x+e)
-arctan(tan(f*x+e)))+A*a*c*x+C*b*d/f*(1/3*tan(f*x+e)^3-tan(f*x+e)+arctan(t
an(f*x+e)))
```


Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.93

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2 C b d \tan (f x + e)^3 + 6 (((A - C) a - B b) c - (B a + (A - C) b) d) f x + 3 (C b c + (C a + B b) d) \tan (f x + e)}{f}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

```
1/6*(2*C*b*d*tan(f*x + e)^3 + 6*(((A - C)*a - B*b)*c - (B*a + (A - C)*b)*d)*f*x + 3*(C*b*c + (C*a + B*b)*d)*tan(f*x + e)^2 - 3*((B*a + (A - C)*b)*c + ((A - C)*a - B*b)*d)*log(1/(tan(f*x + e)^2 + 1)) + 6*((C*a + B*b)*c + (B*a + (A - C)*b)*d)*tan(f*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(148) = 296.

Time = 0.21 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.02

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \begin{cases} A a c x + \frac{A a d \log(\tan^2(e + fx) + 1)}{2f} + \frac{A b c \log(\tan^2(e + fx) + 1)}{2f} - A b d x + \frac{A b d \tan(e + fx)}{f} + \frac{B a c \log(\tan^2(e + fx) + 1)}{2f} - B a d x \\ x(a + b \tan(e)) (c + d \tan(e)) (A + B \tan(e) + C \tan^2(e)) \end{cases}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A*a*c*x + A*a*d*log(tan(e + f*x)**2 + 1)/(2*f) + A*b*c*log(tan(
e + f*x)**2 + 1)/(2*f) - A*b*d*x + A*b*d*tan(e + f*x)/f + B*a*c*log(tan(e
+ f*x)**2 + 1)/(2*f) - B*a*d*x + B*a*d*tan(e + f*x)/f - B*b*c*x + B*b*c*ta
n(e + f*x)/f - B*b*d*log(tan(e + f*x)**2 + 1)/(2*f) + B*b*d*tan(e + f*x)**
2/(2*f) - C*a*c*x + C*a*c*tan(e + f*x)/f - C*a*d*log(tan(e + f*x)**2 + 1)/
(2*f) + C*a*d*tan(e + f*x)**2/(2*f) - C*b*c*log(tan(e + f*x)**2 + 1)/(2*f)
+ C*b*c*tan(e + f*x)**2/(2*f) + C*b*d*x + C*b*d*tan(e + f*x)**3/(3*f) - C
*b*d*tan(e + f*x)/f, Ne(f, 0)), (x*(a + b*tan(e))*(c + d*tan(e))*(A + B*ta
n(e) + C*tan(e)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.94

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2Cbd \tan(fx + e)^3 + 3(Cbc + (Ca + Bb)d) \tan(fx + e)^2 + 6(((A - C)a - Bb)c - (Ba + (A - C)b)d)}{f}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2
),x, algorithm="maxima")
```

output

```
1/6*(2*C*b*d*tan(f*x + e)^3 + 3*(C*b*c + (C*a + B*b)*d)*tan(f*x + e)^2 + 6
*(((A - C)*a - B*b)*c - (B*a + (A - C)*b)*d)*(f*x + e) + 3*((B*a + (A - C)
*b)*c + ((A - C)*a - B*b)*d)*log(tan(f*x + e)^2 + 1) + 6*((C*a + B*b)*c +
(B*a + (A - C)*b)*d)*tan(f*x + e))/f
```

Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.39

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(Aac - Cac - Bbc - Bad - Abd + Cbd)(fx + e)}{f}$$

$$+ \frac{(Bac + Abc - Cbc + Aad - Cad - Bbd) \log(\tan(fx + e)^2 + 1)}{2f}$$

$$+ \frac{2Cbd f^2 \tan(fx + e)^3 + 3Cbc f^2 \tan(fx + e)^2 + 3Cad f^2 \tan(fx + e)^2 + 3Bbd f^2 \tan(fx + e)^2 + 6Cbd f^2 \tan(fx + e)}{3f^3}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
(A*a*c - C*a*c - B*b*c - B*a*d - A*b*d + C*b*d)*(f*x + e)/f + 1/2*(B*a*c + A*b*c - C*b*c + A*a*d - C*a*d - B*b*d)*log(tan(f*x + e)^2 + 1)/f + 1/6*(2*C*b*d*f^2*tan(f*x + e)^3 + 3*C*b*c*f^2*tan(f*x + e)^2 + 3*C*a*d*f^2*tan(f*x + e)^2 + 3*B*b*d*f^2*tan(f*x + e)^2 + 6*C*a*c*f^2*tan(f*x + e) + 6*B*b*c*f^2*tan(f*x + e) + 6*B*a*d*f^2*tan(f*x + e) + 6*A*b*d*f^2*tan(f*x + e) - 6*C*b*d*f^2*tan(f*x + e))/f^3
```

Mupad [B] (verification not implemented)

Time = 5.55 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.95

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{\ln(\tan(e + fx)^2 + 1) \left(\frac{Aad}{2} + \frac{Abc}{2} + \frac{Bac}{2} - \frac{Bbd}{2} - \frac{Cad}{2} - \frac{Cbc}{2} \right)}{f}$$

$$- x(Abd - Aac + Bad + Bbc + Cac - Cbd) + \frac{\tan(e + fx)^2 \left(\frac{Bbd}{2} + \frac{Cad}{2} + \frac{Cbc}{2} \right)}{f}$$

$$+ \frac{\tan(e + fx) (Abd + Bad + Bbc + Cac - Cbd)}{f} + \frac{Cbd \tan(e + fx)^3}{3f}$$

input

```
int((a + b*tan(e + f*x))*(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```


3.53 $\int (c+d \tan(e+fx)) (A + B \tan(e + fx) + C \tan^2(e +$

Optimal result	576
Mathematica [A] (verified)	576
Rubi [A] (verified)	577
Maple [A] (verified)	579
Fricas [A] (verification not implemented)	579
Sympy [B] (verification not implemented)	580
Maxima [A] (verification not implemented)	580
Giac [A] (verification not implemented)	581
Mupad [B] (verification not implemented)	581
Reduce [B] (verification not implemented)	582

Optimal result

Integrand size = 31, antiderivative size = 73

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= (Ac - cC - Bd)x - \frac{(Bc + (A - C)d) \log(\cos(e + fx))}{f}$$

$$+ \frac{Bd \tan(e + fx)}{f} + \frac{C(c + d \tan(e + fx))^2}{2df}$$

output

```
(A*c-B*d-C*c)*x-(B*c+(A-C)*d)*ln(cos(f*x+e))/f+B*d*tan(f*x+e)/f+1/2*C*(c+d
*tan(f*x+e))^2/d/f
```

Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.04

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2Acfx - 2(cC + Bd) \arctan(\tan(e + fx)) - 2(Bc + (A - C)d) \log(\cos(e + fx)) + Cd \sec^2(e + fx) + 2}{2f}$$

input

```
Integrate[(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(2*A*c*f*x - 2*(c*C + B*d)*ArcTan[Tan[e + f*x]] - 2*(B*c + (A - C)*d)*Log[
Cos[e + f*x]] + C*d*Sec[e + f*x]^2 + 2*(c*C + B*d)*Tan[e + f*x])/(2*f)
```

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4113, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4113

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx)) dx + \frac{C(c + d \tan(e + fx))^2}{2df}$$

↓ 3042

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx)) dx + \frac{C(c + d \tan(e + fx))^2}{2df}$$

↓ 4008

$$(d(A - C) + Bc) \int \tan(e + fx) dx + x(Ac - Bd - cC) + \frac{Bd \tan(e + fx)}{f} + \frac{C(c + d \tan(e + fx))^2}{2df}$$

↓ 3042

$$(d(A - C) + Bc) \int \tan(e + fx) dx + x(Ac - Bd - cC) + \frac{Bd \tan(e + fx)}{f} + \frac{C(c + d \tan(e + fx))^2}{2df}$$

↓ 3956

$$-\frac{(d(A-C) + Bc)\log(\cos(e+fx))}{f} + x(Ac - Bd - cC) + \frac{Bd\tan(e+fx)}{f} + \frac{C(c + d\tan(e+fx))^2}{2df}$$

input `Int[(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `(A*c - c*C - B*d)*x - ((B*c + (A - C)*d)*Log[Cos[e + f*x]])/f + (B*d*Tan[e + f*x])/f + (C*(c + d*Tan[e + f*x])^2)/(2*d*f)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

output

$$\frac{1}{2} * (C * d * \tan(f * x + e)^2 + 2 * ((A - C) * c - B * d) * f * x - (B * c + (A - C) * d) * \log\left(\frac{1}{\tan(f * x + e)^2 + 1}\right) + 2 * (C * c + B * d) * \tan(f * x + e)) / f$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(60) = 120$.

Time = 0.10 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.79

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \begin{cases} A c x + \frac{A d \log(\tan^2(e + fx) + 1)}{2f} + \frac{B c \log(\tan^2(e + fx) + 1)}{2f} - B d x + \frac{B d \tan(e + fx)}{f} - C c x + \frac{C c \tan(e + fx)}{f} - \frac{C d \log(\tan^2(e + fx) + 1)}{2f} \\ x(c + d \tan(e)) (A + B \tan(e) + C \tan^2(e)) \end{cases}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A*c*x + A*d*log(tan(e + f*x)**2 + 1)/(2*f) + B*c*log(tan(e + f*x)**2 + 1)/(2*f) - B*d*x + B*d*tan(e + f*x)/f - C*c*x + C*c*tan(e + f*x)/f - C*d*log(tan(e + f*x)**2 + 1)/(2*f) + C*d*tan(e + f*x)**2/(2*f), Ne(f, 0)), (x*(c + d*tan(e))*(A + B*tan(e) + C*tan(e)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.01

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{C d \tan(fx + e)^2 + 2((A - C)c - B d)(fx + e) + (B c + (A - C)d) \log(\tan(fx + e)^2 + 1) + 2(Cc + B d) \tan(fx + e)}{2f}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

$$\frac{1}{2} * (C * d * \tan(f * x + e)^2 + 2 * ((A - C) * c - B * d) * (f * x + e) + (B * c + (A - C) * d) * \log(\tan(f * x + e)^2 + 1) + 2 * (C * c + B * d) * \tan(f * x + e)) / f$$

Giac [A] (verification not implemented)

Time = 0.52 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.22

$$\begin{aligned} & \int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \frac{(Ac - Cc - Bd)(fx + e)}{f} + \frac{(Bc + Ad - Cd) \log(\tan(fx + e)^2 + 1)}{2f} \\ &+ \frac{Cdf \tan(fx + e)^2 + 2Ccf \tan(fx + e) + 2Bdf \tan(fx + e)}{2f^2} \end{aligned}$$

input `integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `(A*c - C*c - B*d)*(f*x + e)/f + 1/2*(B*c + A*d - C*d)*log(tan(f*x + e)^2 + 1)/f + 1/2*(C*d*f*tan(f*x + e)^2 + 2*C*c*f*tan(f*x + e) + 2*B*d*f*tan(f*x + e))/f^2`

Mupad [B] (verification not implemented)

Time = 5.37 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.03

$$\begin{aligned} & \int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \frac{\tan(e + fx) (Bd + Cc)}{f} - x(Bd - Ac + Cc) \\ &+ \frac{\ln(\tan(e + fx)^2 + 1) \left(\frac{Ad}{2} + \frac{Bc}{2} - \frac{Cd}{2}\right)}{f} + \frac{Cd \tan(e + fx)^2}{2f} \end{aligned}$$

input `int((c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `(tan(e + f*x)*(B*d + C*c))/f - x*(B*d - A*c + C*c) + (log(tan(e + f*x)^2 + 1)*((A*d)/2 + (B*c)/2 - (C*d)/2))/f + (C*d*tan(e + f*x)^2)/(2*f)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.37

$$\int (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{\log(\tan(fx + e)^2 + 1) ad + \log(\tan(fx + e)^2 + 1) bc - \log(\tan(fx + e)^2 + 1) cd + \tan(fx + e)^2 cd + 2 \tan(e + fx) b^2 d + 2 \tan(e + fx) c^2 d + 2 a^2 c f x - 2 b^2 d f x - 2 c^2 f x}{2f}$$

input `int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `(log(tan(e + f*x)**2 + 1)*a*d + log(tan(e + f*x)**2 + 1)*b*c - log(tan(e + f*x)**2 + 1)*c*d + tan(e + f*x)**2*c*d + 2*tan(e + f*x)*b*d + 2*tan(e + f*x)*c**2 + 2*a*c*f*x - 2*b*d*f*x - 2*c**2*f*x)/(2*f)`

3.54
$$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$$

Optimal result	583
Mathematica [C] (verified)	584
Rubi [A] (verified)	584
Maple [A] (verified)	587
Fricas [A] (verification not implemented)	588
Sympy [C] (verification not implemented)	588
Maxima [A] (verification not implemented)	589
Giac [A] (verification not implemented)	590
Mupad [B] (verification not implemented)	590
Reduce [B] (verification not implemented)	591

Optimal result

Integrand size = 43, antiderivative size = 156

$$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$$

$$= \frac{(a(Ac - cC - Bd) + b(Bc + (A - C)d))x}{a^2 + b^2}$$

$$+ \frac{(Abc - aBc - bcC - aAd - bBd + aCd) \log(\cos(e+fx))}{(a^2 + b^2) f}$$

$$+ \frac{(Ab^2 - a(bB - aC))(bc - ad) \log(a + b \tan(e+fx))}{b^2 (a^2 + b^2) f} + \frac{Cd \tan(e+fx)}{bf}$$

output

```
(a*(A*c-B*d-C*c)+b*(B*c+(A-C)*d))*x/(a^2+b^2)+(-A*a*d+A*b*c-B*a*c-B*b*d+C*
a*d-C*b*c)*ln(cos(f*x+e))/(a^2+b^2)/f+(A*b^2-a*(B*b-C*a))*(-a*d+b*c)*ln(a+
b*tan(f*x+e))/b^2/(a^2+b^2)/f+C*d*tan(f*x+e)/b/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.95

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{\frac{(A+iB-C)(-ic+d) \log(i-\tan(e+fx))}{a+ib} + \frac{(A-iB-C)(ic+d) \log(i+\tan(e+fx))}{a-ib} + \frac{2(Ab^2+a(-bB+aC))(bc-ad) \log(a+b \tan(e+fx))}{b^2(a^2+b^2)}}{2f}$$

input

```
Integrate[((c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]
```

output

```
((A + I*B - C)*((-I)*c + d)*Log[I - Tan[e + f*x]]/(a + I*b) + ((A - I*B - C)*(I*c + d)*Log[I + Tan[e + f*x]]/(a - I*b) + (2*(A*b^2 + a*(-(b*B) + a*C))*(b*c - a*d)*Log[a + b*Tan[e + f*x]]/(b^2*(a^2 + b^2)) + (2*C*d*Tan[e + f*x])/b)/(2*f)
```

Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4120, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan(e + fx)^2)}{a + b \tan(e + fx)} dx$$

↓ 4120

$$\frac{Cd \tan(e + fx)}{bf} - \frac{\int -\frac{(bcC - adC + bBd) \tan^2(e + fx) + b(Bc + (A - C)d) \tan(e + fx) + Abc - aCd}{a + b \tan(e + fx)} dx}{b}$$

↓ 25

$$\frac{\int \frac{(bcC - adC + bBd) \tan^2(e + fx) + b(Bc + (A - C)d) \tan(e + fx) + Abc - aCd}{a + b \tan(e + fx)} dx}{b} + \frac{Cd \tan(e + fx)}{bf}$$

↓ 3042

$$\frac{\int \frac{(bcC - adC + bBd) \tan(e + fx)^2 + b(Bc + (A - C)d) \tan(e + fx) + Abc - aCd}{a + b \tan(e + fx)} dx}{b} + \frac{Cd \tan(e + fx)}{bf}$$

↓ 4109

$$\frac{\frac{(bc - ad)(Ab^2 - a(bB - aC)) \int \frac{\tan^2(e + fx) + 1}{a + b \tan(e + fx)} dx}{a^2 + b^2} - \frac{b(-aAd - aBc + aCd + Abc - bBd - bcC) \int \tan(e + fx) dx}{a^2 + b^2} + \frac{bx(a(Ac - Bd - cC) + bd(A - C) + bB)}{a^2 + b^2}}{b}$$

$\frac{Cd \tan(e + fx)}{bf}$

↓ 3042

$$\frac{-\frac{b(-aAd - aBc + aCd + Abc - bBd - bcC) \int \tan(e + fx) dx}{a^2 + b^2} + \frac{(bc - ad)(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{a + b \tan(e + fx)} dx}{a^2 + b^2} + \frac{bx(a(Ac - Bd - cC) + bd(A - C) + bB)}{a^2 + b^2}}{b}$$

$\frac{Cd \tan(e + fx)}{bf}$

↓ 3956

$$\frac{\frac{(bc - ad)(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{a + b \tan(e + fx)} dx}{a^2 + b^2} + \frac{b \log(\cos(e + fx))(-aAd - aBc + aCd + Abc - bBd - bcC)}{f(a^2 + b^2)} + \frac{bx(a(Ac - Bd - cC) + bd(A - C) + bB)}{a^2 + b^2}}{b}$$

$\frac{Cd \tan(e + fx)}{bf}$

↓ 4100

$$\frac{\frac{(bc - ad)(Ab^2 - a(bB - aC)) \int \frac{1}{a + b \tan(e + fx)} d(b \tan(e + fx))}{bf(a^2 + b^2)} + \frac{b \log(\cos(e + fx))(-aAd - aBc + aCd + Abc - bBd - bcC)}{f(a^2 + b^2)} + \frac{bx(a(Ac - Bd - cC) + bd(A - C) + bB)}{a^2 + b^2}}{b}$$

$\frac{Cd \tan(e + fx)}{bf}$

↓ 16

$$\frac{\frac{(bc-ad)(Ab^2-a(bB-aC)) \log(a+b \tan(e+fx))}{bf(a^2+b^2)} + \frac{b \log(\cos(e+fx))(-aAd-aBc+aCd+Abc-bBd-bcC)}{f(a^2+b^2)} + \frac{bx(a(Ac-Bd-cC)+bd(A-C)+b^2)}{a^2+b^2}}{\frac{C d \tan(e+fx)}{bf}}$$

input

```
Int[((c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*
Tan[e + f*x]),x]
```

output

```
((b*(b*B*c + b*(A - C)*d + a*(A*c - c*C - B*d))*x)/(a^2 + b^2) + (b*(A*b*c
- a*B*c - b*c*C - a*A*d - b*B*d + a*C*d)*Log[Cos[e + f*x]])/((a^2 + b^2)*
f) + ((A*b^2 - a*(b*B - a*C))*(b*c - a*d)*Log[a + b*Tan[e + f*x]])/(b*(a^2
+ b^2)*f))/b + (C*d*Tan[e + f*x])/(b*f)
```

Defintions of rubi rules used

rule 16

```
Int[(c.)/((a.) + (b.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3956

```
Int[tan[(c.) + (d.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4100

```
Int[((a.) + (b.)*tan[(e.) + (f.)*(x_)])^(m.)*((A_) + (C.)*tan[(e.) +
(f.)*(x_)^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

rule 4109

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2
)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]
```

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_
)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{\frac{\tan(fx+e)Cd}{b} + \frac{(Aad - Abc + Bac + Bbd - Cad + Cbc) \ln(1 + \tan(fx+e)^2)}{2} + (Aac + Abd - Bad + Bbc - Cac - Cbd) \arctan(\tan(fx+e))}{a^2 + b^2} + \frac{f}{f}$
default	$\frac{\frac{\tan(fx+e)Cd}{b} + \frac{(Aad - Abc + Bac + Bbd - Cad + Cbc) \ln(1 + \tan(fx+e)^2)}{2} + (Aac + Abd - Bad + Bbc - Cac - Cbd) \arctan(\tan(fx+e))}{a^2 + b^2} + \frac{f}{f}$
norman	$\frac{(Aac + Abd - Bad + Bbc - Cac - Cbd)x}{a^2 + b^2} + \frac{Cd \tan(fx+e)}{bf} + \frac{(Aad - Abc + Bac + Bbd - Cad + Cbc) \ln(1 + \tan(fx+e)^2)}{2(a^2 + b^2)f}$
parallelrisc	$\frac{2Aa b^2 c f x + 2A b^3 d f x - 2B a b^2 d f x + 2B b^3 c f x - 2C a b^2 c f x - 2C b^3 d f x + A \ln(1 + \tan(fx+e)^2) a b^2 d - A \ln(1 + \tan(fx+e)^2)}{f(a^2 + b^2)}$
risc	$\frac{2iCd}{fb(e^{2i(fx+e)} + 1)} - \frac{\ln(e^{2i(fx+e)} - \frac{ib+a}{ib-a}) Aad}{f(a^2 + b^2)} + \frac{b \ln(e^{2i(fx+e)} - \frac{ib+a}{ib-a}) Ac}{f(a^2 + b^2)} - \frac{\ln(e^{2i(fx+e)} - \frac{ib+a}{ib-a}) Bac}{f(a^2 + b^2)} + \frac{2iBd}{bf}$

input

```
int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x,me
thod=_RETURNVERBOSE)
```


output

```
1/f*(tan(f*x+e)*C*d/b+1/(a^2+b^2)*(1/2*(A*a*d-A*b*c+B*a*c+B*b*d-C*a*d+C*b*c)*ln(1+tan(f*x+e)^2)+(A*a*c+A*b*d-B*a*d+B*b*c-C*a*c-C*b*d)*arctan(tan(f*x+e)))+(-A*a*b^2*d+A*b^3*c+B*a^2*b*d-B*a*b^2*c-C*a^3*d+C*a^2*b*c)/b^2/(a^2+b^2)*ln(a+b*tan(f*x+e)))
```

Fricas [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.45

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{2(((A - C)ab^2 + Bb^3)c - (Bab^2 - (A - C)b^3)d)fx + 2(Ca^2b + Cb^3)d \tan(fx + e) + ((Ca^2b - Bab^2 +$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="fricas")
```

output

```
1/2*(2*(((A - C)*a*b^2 + B*b^3)*c - (B*a*b^2 - (A - C)*b^3)*d)*f*x + 2*(C*a^2*b + C*b^3)*d*tan(f*x + e) + ((C*a^2*b - B*a*b^2 + A*b^3)*c - (C*a^3 - B*a^2*b + A*a*b^2)*d)*log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) - ((C*a^2*b + C*b^3)*c - (C*a^3 - B*a^2*b + C*a*b^2 - B*b^3)*d)*log(1/(tan(f*x + e)^2 + 1)))/((a^2*b^2 + b^4)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 2387, normalized size of antiderivative = 15.30

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)),x)
```

output

```
Piecewise((zoo*x*(c + d*tan(e))*(A + B*tan(e) + C*tan(e)**2)/tan(e), Eq(a,
0) & Eq(b, 0) & Eq(f, 0)), ((A*c*x + A*d*log(tan(e + f*x)**2 + 1)/(2*f) +
B*c*log(tan(e + f*x)**2 + 1)/(2*f) - B*d*x + B*d*tan(e + f*x)/f - C*c*x +
C*c*tan(e + f*x)/f - C*d*log(tan(e + f*x)**2 + 1)/(2*f) + C*d*tan(e + f*x)
)**2/(2*f))/a, Eq(b, 0)), (I*A*c*f*x*tan(e + f*x)/(2*b*f*tan(e + f*x) - 2*
I*b*f) + A*c*f*x/(2*b*f*tan(e + f*x) - 2*I*b*f) + I*A*c/(2*b*f*tan(e + f*x)
) - 2*I*b*f) + A*d*f*x*tan(e + f*x)/(2*b*f*tan(e + f*x) - 2*I*b*f) - I*A*d
*f*x/(2*b*f*tan(e + f*x) - 2*I*b*f) - A*d/(2*b*f*tan(e + f*x) - 2*I*b*f) +
B*c*f*x*tan(e + f*x)/(2*b*f*tan(e + f*x) - 2*I*b*f) - I*B*c*f*x/(2*b*f*tan
(e + f*x) - 2*I*b*f) - B*c/(2*b*f*tan(e + f*x) - 2*I*b*f) + I*B*d*f*x*tan
(e + f*x)/(2*b*f*tan(e + f*x) - 2*I*b*f) + B*d*f*x/(2*b*f*tan(e + f*x) - 2
*I*b*f) + B*d*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*b*f*tan(e + f*x) -
2*I*b*f) - I*B*d*log(tan(e + f*x)**2 + 1)/(2*b*f*tan(e + f*x) - 2*I*b*f) -
I*B*d/(2*b*f*tan(e + f*x) - 2*I*b*f) + I*C*c*f*x*tan(e + f*x)/(2*b*f*tan(
e + f*x) - 2*I*b*f) + C*c*f*x/(2*b*f*tan(e + f*x) - 2*I*b*f) + C*c*log(tan
(e + f*x)**2 + 1)*tan(e + f*x)/(2*b*f*tan(e + f*x) - 2*I*b*f) - I*C*c*log(
tan(e + f*x)**2 + 1)/(2*b*f*tan(e + f*x) - 2*I*b*f) - I*C*c/(2*b*f*tan(e +
f*x) - 2*I*b*f) - 3*C*d*f*x*tan(e + f*x)/(2*b*f*tan(e + f*x) - 2*I*b*f) +
3*I*C*d*f*x/(2*b*f*tan(e + f*x) - 2*I*b*f) + I*C*d*log(tan(e + f*x)**2 +
1)*tan(e + f*x)/(2*b*f*tan(e + f*x) - 2*I*b*f) + C*d*log(tan(e + f*x)**...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.17

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{2Cd \tan(fx+e)}{b} + \frac{2(((A-C)a+Bb)c - (Ba - (A-C)b)d)(fx+e)}{a^2+b^2} + \frac{2((Ca^2b - Bab^2 + Ab^3)c - (Ca^3 - Ba^2b + Aab^2)d) \log(b \tan(fx+e) + a)}{a^2b^2 + b^4} + \frac{2f}{2f}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
),x, algorithm="maxima")
```

output

```
1/2*(2*C*d*tan(f*x + e)/b + 2*(((A - C)*a + B*b)*c - (B*a - (A - C)*b)*d)*
(f*x + e)/(a^2 + b^2) + 2*((C*a^2*b - B*a*b^2 + A*b^3)*c - (C*a^3 - B*a^2*b
+ A*a*b^2)*d)*log(b*tan(f*x + e) + a)/(a^2*b^2 + b^4) + ((B*a - (A - C)*
b)*c + ((A - C)*a + B*b)*d)*log(tan(f*x + e)^2 + 1)/(a^2 + b^2))/f
```

Giac [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.21

$$\begin{aligned}
& \int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\
&= \frac{(Aac - Cac + Bbc - Bad + Abd - Cbd)(fx + e)}{a^2 f + b^2 f} \\
&\quad + \frac{(Bac - Abc + Cbc + Aad - Cad + Bbd) \log(\tan(fx + e)^2 + 1)}{2(a^2 f + b^2 f)} \\
&\quad + \frac{(Ca^2 bc - Bab^2 c + Ab^3 c - Ca^3 d + Ba^2 bd - Aab^2 d) \log(|b \tan(fx + e) + a|)}{a^2 b^2 f + b^4 f} \\
&\quad + \frac{Cd \tan(fx + e)}{bf}
\end{aligned}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="giac")
```

output

```
(A*a*c - C*a*c + B*b*c - B*a*d + A*b*d - C*b*d)*(f*x + e)/(a^2*f + b^2*f) + 1/2*(B*a*c - A*b*c + C*b*c + A*a*d - C*a*d + B*b*d)*log(tan(f*x + e)^2 + 1)/(a^2*f + b^2*f) + (C*a^2*b*c - B*a*b^2*c + A*b^3*c - C*a^3*d + B*a^2*b*d - A*a*b^2*d)*log(abs(b*tan(f*x + e) + a))/(a^2*b^2*f + b^4*f) + C*d*tan(f*x + e)/(b*f)
```

Mupad [B] (verification not implemented)

Time = 6.35 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.19

$$\begin{aligned}
& \int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\
&= \frac{\ln(\tan(e + fx) - i)(Ad + Bc - Cd - Acli + Bcli + Ccli)}{2f(a + bli)} \\
&\quad + \frac{\ln(\tan(e + fx) + i)(Bd + Adli + Bcli - Ac + Cc - Ccli)}{2f(b + ali)} \\
&\quad - \frac{\ln(a + b \tan(e + fx))(b^2(Aad + Bac) - b(Ba^2 d + Ca^2 c) - Ab^3 c + Ca^3 d)}{f(a^2 b^2 + b^4)} \\
&\quad + \frac{Cd \tan(e + fx)}{bf}
\end{aligned}$$

input `int(((c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x)),x)`

output `(log(tan(e + f*x) - 1i)*(A*d - A*c*1i + B*c + B*d*1i + C*c*1i - C*d))/(2*f*(a + b*1i)) + (log(tan(e + f*x) + 1i)*(A*d*1i - A*c + B*c*1i + B*d + C*c - C*d*1i))/(2*f*(a*1i + b)) - (log(a + b*tan(e + f*x))*(b^2*(A*a*d + B*a*c) - b*(B*a^2*d + C*a^2*c) - A*b^3*c + C*a^3*d))/(f*(b^4 + a^2*b^2)) + (C*d*tan(e + f*x))/(b*f)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.24

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{\log(\tan(fx + e)^2 + 1) a^2 b^2 d - \log(\tan(fx + e)^2 + 1) a b^2 c d + \log(\tan(fx + e)^2 + 1) b^4 d + \log(\tan(fx + e)^2 + 1) a^2 b^2 c d}{a^2 b^2 d - a b^2 c d + b^4 d + a^2 b^2 c d}$$

input `int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x)`

output `(log(tan(e + f*x)**2 + 1)*a**2*b**2*d - log(tan(e + f*x)**2 + 1)*a*b**2*c*d + log(tan(e + f*x)**2 + 1)*b**4*d + log(tan(e + f*x)**2 + 1)*b**3*c**2 - 2*log(tan(e + f*x)*b + a)*a**3*c*d + 2*log(tan(e + f*x)*b + a)*a**2*b*c**2 + 2*tan(e + f*x)*a**2*b*c*d + 2*tan(e + f*x)*b**3*c*d + 2*a**2*b**2*c*f*x - 2*a*b**2*c**2*f*x + 2*b**4*c*f*x - 2*b**3*c*d*f*x)/(2*b**2*f*(a**2 + b**2))`

3.55
$$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$$

Optimal result	592
Mathematica [C] (verified)	593
Rubi [A] (verified)	593
Maple [A] (verified)	596
Fricas [B] (verification not implemented)	597
Sympy [C] (verification not implemented)	598
Maxima [A] (verification not implemented)	599
Giac [A] (verification not implemented)	599
Mupad [B] (verification not implemented)	600
Reduce [B] (verification not implemented)	601

Optimal result

Integrand size = 43, antiderivative size = 265

$$\begin{aligned} & \int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx \\ &= \frac{(a^2(Ac - cC - Bd) - b^2(Ac - cC - Bd) + 2ab(Bc + (A - C)d)) x}{(a^2 + b^2)^2} \\ & \quad + \frac{(2ab(Ac - cC - Bd) - a^2(Bc + (A - C)d) + b^2(Bc + (A - C)d)) \log(\cos(e+fx))}{(a^2 + b^2)^2 f} \\ & \quad + \frac{(a^4Cd + b^4(Bc + Ad) + 2ab^3(Ac - cC - Bd) - a^2b^2(Bc + (A - 3C)d)) \log(a+b \tan(e+fx))}{b^2 (a^2 + b^2)^2 f} \\ & \quad - \frac{(Ab^2 - a(bB - aC))(bc - ad)}{b^2 (a^2 + b^2) f(a+b \tan(e+fx))} \end{aligned}$$

output

```
(a^2*(A*c-B*d-C*c)-b^2*(A*c-B*d-C*c)+2*a*b*(B*c+(A-C)*d))*x/(a^2+b^2)^2+(2
*a*b*(A*c-B*d-C*c)-a^2*(B*c+(A-C)*d)+b^2*(B*c+(A-C)*d))*ln(cos(f*x+e))/(a^
2+b^2)^2/f+(a^4*C*d+b^4*(A*d+B*c)+2*a*b^3*(A*c-B*d-C*c)-a^2*b^2*(B*c+(A-3*
C)*d))*ln(a+b*tan(f*x+e))/b^2/(a^2+b^2)^2/f-(A*b^2-a*(B*b-C*a))*(-a*d+b*c)
/b^2/(a^2+b^2)/f/(a+b*tan(f*x+e))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.68 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.82

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{\frac{(A+iB-C)(-ic+d) \log(i-\tan(e+fx))}{(a+ib)^2} + \frac{(A-iB-C)(ic+d) \log(i+\tan(e+fx))}{(a-ib)^2} + \frac{2(a^4Cd+b^4(Bc+Ad)+2ab^3(Ac-cC-Bd)-a^2b^2(Bc+(A+ib)^2))}{b^2(a^2+b^2)^2}}{2f}$$

input

```
Integrate[((c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^2,x]
```

output

```
((A + I*B - C)*((-I)*c + d)*Log[I - Tan[e + f*x]]/(a + I*b)^2 + ((A - I*B - C)*(I*c + d)*Log[I + Tan[e + f*x]]/(a - I*b)^2 + (2*(a^4*C*d + b^4*(B*c + A*d) + 2*a*b^3*(A*c - c*C - B*d) - a^2*b^2*(B*c + (A - 3*C)*d))*Log[a + b*Tan[e + f*x]]/(b^2*(a^2 + b^2)^2) - (2*(A*b^2 + a*(-(b*B) + a*C))*(b*c - a*d))/(b^2*(a^2 + b^2)*(a + b*Tan[e + f*x])))/(2*f)
```

Rubi [A] (verified)

Time = 1.79 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.06, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4118, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^2} dx$$

$$\downarrow \text{4118}$$

$$\int \frac{Cda^2+b(Ac-Cc-Bd)a+(a^2+b^2)Cd \tan^2(e+fx)+b^2(Bc+Ad)-b(Abc-aBc-bCc-aAd-bBd+aCd) \tan(e+fx)}{a+b \tan(e+fx)} dx$$

$$\frac{b(a^2+b^2)}{b^2 f(a^2+b^2)(a+b \tan(e+fx))} \frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

3042

$$\int \frac{Cda^2+b(Ac-Cc-Bd)a+(a^2+b^2)Cd \tan(e+fx)^2+b^2(Bc+Ad)-b(Abc-aBc-bCc-aAd-bBd+aCd) \tan(e+fx)}{a+b \tan(e+fx)} dx$$

$$\frac{b(a^2+b^2)}{b^2 f(a^2+b^2)(a+b \tan(e+fx))} \frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

4109

$$-\frac{b(-(a^2(d(A-C)+Bc))+2ab(Ac-Bd-cC)+b^2(d(A-C)+Bc)) \int \tan(e+fx) dx}{a^2+b^2} + \frac{(a^4Cd-a^2b^2(d(A-3C)+Bc)+2ab^3(Ac-Bd-cC)+b^4(Ad+bc)) \int \tan(e+fx) dx}{a^2+b^2}$$

$$\frac{b(a^2+b^2)}{b^2 f(a^2+b^2)(a+b \tan(e+fx))} \frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

3042

$$-\frac{b(-(a^2(d(A-C)+Bc))+2ab(Ac-Bd-cC)+b^2(d(A-C)+Bc)) \int \tan(e+fx) dx}{a^2+b^2} + \frac{(a^4Cd-a^2b^2(d(A-3C)+Bc)+2ab^3(Ac-Bd-cC)+b^4(Ad+bc)) \int \tan(e+fx) dx}{a^2+b^2}$$

$$\frac{b(a^2+b^2)}{b^2 f(a^2+b^2)(a+b \tan(e+fx))} \frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

3956

$$\frac{(a^4Cd-a^2b^2(d(A-3C)+Bc)+2ab^3(Ac-Bd-cC)+b^4(Ad+bc)) \int \frac{\tan(e+fx)^2+1}{a+b \tan(e+fx)} dx}{a^2+b^2} + \frac{b \log(\cos(e+fx))(-a^2(d(A-C)+Bc))+2ab(Ac-Bd-cC)}{f(a^2+b^2)}$$

$$\frac{b(a^2+b^2)}{b^2 f(a^2+b^2)(a+b \tan(e+fx))} \frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

4100

$$\frac{(a^4Cd-a^2b^2(d(A-3C)+Bc)+2ab^3(Ac-Bd-cC)+b^4(Ad+bc)) \int \frac{1}{a+b \tan(e+fx)} d(b \tan(e+fx))}{bf(a^2+b^2)} + \frac{b \log(\cos(e+fx))(-a^2(d(A-C)+Bc))+2ab(Ac-Bd-cC)}{f(a^2+b^2)}$$

$$\frac{b(a^2+b^2)}{b^2 f(a^2+b^2)(a+b \tan(e+fx))} \frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 16

$$\frac{b \log(\cos(e+fx))(-a^2(d(A-C)+Bc)+2ab(Ac-Bd-cC)+b^2(d(A-C)+Bc))}{f(a^2+b^2)} + \frac{bx(a^2(Ac-Bd-cC)+2ab(d(A-C)+Bc)-b^2(Ac-Bd-cC))}{a^2+b^2}}{b(a^2+b^2)}$$

$$\frac{(bc-ad)(Ab^2-a(bB-aC))}{b^2 f(a^2+b^2)(a+b \tan(e+fx))}$$

input

```
Int[((c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*
Tan[e + f*x])^2,x]
```

output

```
((b*(a^2*(A*c - c*C - B*d) - b^2*(A*c - c*C - B*d) + 2*a*b*(B*c + (A - C)*
d))*x)/(a^2 + b^2) + (b*(2*a*b*(A*c - c*C - B*d) - a^2*(B*c + (A - C)*d) +
b^2*(B*c + (A - C)*d))*Log[Cos[e + f*x]]/((a^2 + b^2)*f) + ((a^4*C*d + b
^4*(B*c + A*d) + 2*a*b^3*(A*c - c*C - B*d) - a^2*b^2*(B*c + (A - 3*C)*d))*
Log[a + b*Tan[e + f*x]]/(b*(a^2 + b^2)*f)/(b*(a^2 + b^2)) - ((A*b^2 - a*
(b*B - a*C))*(b*c - a*d))/(b^2*(a^2 + b^2)*f*(a + b*Tan[e + f*x]))
```

Defintions of rubi rules used

rule 16

```
Int[(c.)/((a.) + (b.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3956

```
Int[tan[(c.) + (d.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4100

```
Int[((a.) + (b.)*tan[(e.) + (f.)*(x_)])^(m.)*((A.) + (C.)*tan[(e.) +
(f.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```


rule 4109

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]
```

rule 4118

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^n*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{(A a^2 d - 2 A a b c - A b^2 d + B a^2 c + 2 B a b d - B b^2 c - C a^2 d + 2 C a b c + d C b^2) \ln(1 + \tan(fx + e))^2}{2(a^2 + b^2)^2} + \frac{(A a^2 c + 2 A a b d - A b^2 c - B a^2 d + 2 B a b c - B b^2 d + C a^2 c + 2 C a b d - C b^2 c)}{(a^2 + b^2)^2}$
default	$\frac{(A a^2 d - 2 A a b c - A b^2 d + B a^2 c + 2 B a b d - B b^2 c - C a^2 d + 2 C a b c + d C b^2) \ln(1 + \tan(fx + e))^2}{2(a^2 + b^2)^2} + \frac{(A a^2 c + 2 A a b d - A b^2 c - B a^2 d + 2 B a b c - B b^2 d + C a^2 c + 2 C a b d - C b^2 c)}{(a^2 + b^2)^2}$
norman	$\frac{a(A a^2 c + 2 A a b d - A b^2 c - B a^2 d + 2 B a b c + B b^2 d - C a^2 c - 2 C a b d + C b^2 c)x}{a^4 + 2 b^2 a^2 + b^4} + \frac{A a b^2 d - A b^3 c - B a^2 b d + B a b^2 c + a^3 C d - C a^2 b c}{b^2 f(a^2 + b^2)} + \frac{b(A a^2 c + 2 A a b d - A b^2 c - B a^2 d + 2 B a b c - B b^2 d + C a^2 c + 2 C a b d - C b^2 c)}{a + b \tan(fx + e)}$
parallelrisch	Expression too large to display
risch	Expression too large to display

input

```
int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, method=_RETURNVERBOSE)
```

output

```
1/f*(1/(a^2+b^2)^2*(1/2*(A*a^2*d-2*A*a*b*c-A*b^2*d+B*a^2*c+2*B*a*b*d-B*b^2*c-C*a^2*d+2*C*a*b*c+C*b^2*d)*ln(1+tan(f*x+e)^2)+(A*a^2*c+2*A*a*b*d-A*b^2*c-B*a^2*d+2*B*a*b*c+B*b^2*d-C*a^2*c-2*C*a*b*d+C*b^2*c)*arctan(tan(f*x+e)))-(-A*a*b^2*d+A*b^3*c+B*a^2*b*d-B*a*b^2*c-C*a^3*d+C*a^2*b*c)/b^2/(a^2+b^2)/(a+b*tan(f*x+e))+1/(a^2+b^2)^2*(-A*a^2*b^2*d+2*A*a*b^3*c+A*b^4*d-B*a^2*b^2*c-2*B*a*b^3*d+B*b^4*c+C*a^4*d+3*C*a^2*b^2*d-2*C*a*b^3*c)/b^2*ln(a+b*tan(f*x+e)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 556 vs. $2(265) = 530$.

Time = 0.35 (sec) , antiderivative size = 556, normalized size of antiderivative = 2.10

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{2(((A - C)a^3b^2 + 2Ba^2b^3 - (A - C)ab^4)c - (Ba^3b^2 - 2(A - C)a^2b^3 - Bab^4)d)fx - 2(Ca^2b^3 - Bab^4)}{...}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")
```

output

```
1/2*(2*(((A - C)*a^3*b^2 + 2*B*a^2*b^3 - (A - C)*a*b^4)*c - (B*a^3*b^2 - 2*(A - C)*a^2*b^3 - B*a*b^4)*d)*f*x - 2*(C*a^2*b^3 - B*a*b^4 + A*b^5)*c + 2*(C*a^3*b^2 - B*a^2*b^3 + A*a*b^4)*d - ((B*a^3*b^2 - 2*(A - C)*a^2*b^3 - B*a*b^4)*c - (C*a^5 - (A - 3*C)*a^3*b^2 - 2*B*a^2*b^3 + A*a*b^4)*d + ((B*a^2*b^3 - 2*(A - C)*a*b^4 - B*b^5)*c - (C*a^4*b - (A - 3*C)*a^2*b^3 - 2*B*a*b^4 + A*b^5)*d)*tan(f*x + e))*log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) - ((C*a^4*b + 2*C*a^2*b^3 + C*b^5)*d*tan(f*x + e) + (C*a^5 + 2*C*a^3*b^2 + C*a*b^4)*d)*log(1/(tan(f*x + e)^2 + 1)) + 2*(((A - C)*a^2*b^3 + 2*B*a*b^4 - (A - C)*b^5)*c - (B*a^2*b^3 - 2*(A - C)*a*b^4 - B*b^5)*d)*f*x + (C*a^3*b^2 - B*a^2*b^3 + A*a*b^4)*c - (C*a^4*b - B*a^3*b^2 + A*a^2*b^3)*d)*tan(f*x + e))/((a^4*b^3 + 2*a^2*b^5 + b^7)*f*tan(f*x + e) + (a^5*b^2 + 2*a^3*b^4 + a*b^6)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.26 (sec) , antiderivative size = 9721, normalized size of antiderivative = 36.68

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**2,x)`

output

```
Piecewise((zoo*x*(c + d*tan(e))*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2, Eq
(a, 0) & Eq(b, 0) & Eq(f, 0)), ((A*c*x + A*d*log(tan(e + f*x)**2 + 1)/(2*f
) + B*c*log(tan(e + f*x)**2 + 1)/(2*f) - B*d*x + B*d*tan(e + f*x)/f - C*c*
x + C*c*tan(e + f*x)/f - C*d*log(tan(e + f*x)**2 + 1)/(2*f) + C*d*tan(e +
f*x)**2/(2*f))/a**2, Eq(b, 0)), (-A*c*f*x*tan(e + f*x)**2/(4*b**2*f*tan(e
+ f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + 2*I*A*c*f*x*tan(e + f*x)
/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + A*c*f*x
/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f)
+ 2*I*A*c/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f)
+ I*A*d*f*x*tan(e + f*x)**2/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e +
f*x) - 4*b**2*f) + 2*A*d*f*x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 - 8*I
*b**2*f*tan(e + f*x) - 4*b**2*f) - I*A*d*f*x/(4*b**2*f*tan(e + f*x)**2 - 8
*I*b**2*f*tan(e + f*x) - 4*b**2*f) + I*A*d*tan(e + f*x)/(4*b**2*f*tan(e +
f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + I*B*c*f*x*tan(e + f*x)**2/
(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + 2*B*c*f*
x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**
2*f) - I*B*c*f*x/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b
**2*f) + I*B*c*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e +
f*x) - 4*b**2*f) + B*d*f*x*tan(e + f*x)**2/(4*b**2*f*tan(e + f*x)**2 - ...
```

Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.28

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{2(((A-C)a^2+2Bab-(A-C)b^2)c-(Ba^2-2(A-C)ab-Bb^2)d)(fx+e)}{a^4+2a^2b^2+b^4} - \frac{2((Ba^2b^2-2(A-C)ab^3-Bb^4)c-(Ca^4-(A-3C)a^2b^2-2Bab^3+Ab^4)d)\log(b\tan(fx+e)+a)}{a^4b^2+2a^2b^4+b^6} + \frac{((A-C)a^2+2Bab-(A-C)b^2)d\log(\tan(fx+e)^2+1)}{a^4+2a^2b^2+b^4} - \frac{2((Ca^3-3Ab^2c+Ab^3)d)\log(\tan(fx+e))}{(a^2+b^2)^2(b\tan(fx+e)+a)b^2f}$$

input `integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/2*(2*(((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c - (B*a^2 - 2*(A - C)*a*b - B*b^2)*d)*(f*x + e)/(a^4 + 2*a^2*b^2 + b^4) - 2*((B*a^2*b^2 - 2*(A - C)*a*b^3 - B*b^4)*c - (C*a^4 - (A - 3*C)*a^2*b^2 - 2*B*a*b^3 + A*b^4)*d)*log(b*tan(f*x + e) + a)/(a^4*b^2 + 2*a^2*b^4 + b^6) + ((B*a^2 - 2*(A - C)*a*b - B*b^2)*c + ((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*d)*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*((C*a^2*b - B*a*b^2 + A*b^3)*c - (C*a^3 - B*a^2*b + A*a*b^2)*d)/(a^3*b^2 + a*b^4 + (a^2*b^3 + b^5)*tan(f*x + e))/f`

Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.60

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{(Aa^2c - Ca^2c + 2Babc - Ab^2c + Cb^2c - Ba^2d + 2Aabd - 2Cabd + Bb^2d)(fx + e)}{a^4f + 2a^2b^2f + b^4f} + \frac{(Ba^2c - 2Aabc + 2Cabc - Bb^2c + Aa^2d - Ca^2d + 2Babd - Ab^2d + Cb^2d) \log(\tan(fx + e)^2 + 1)}{2(a^4f + 2a^2b^2f + b^4f)} - \frac{(Ba^2b^2c - 2Aab^3c + 2Cab^3c - Bb^4c - Ca^4d + Aa^2b^2d - 3Ca^2b^2d + 2Bab^3d - Ab^4d) \log(|b \tan(fx + e)|)}{a^4b^2f + 2a^2b^4f + b^6f} - \frac{Ca^4bc - Ba^3b^2c + Aa^2b^3c + Ca^2b^3c - Bab^4c + Ab^5c - Ca^5d + Ba^4bd - Aa^3b^2d - Ca^3b^2d + Ba^2b^3d}{(a^2 + b^2)^2(b \tan(fx + e) + a)b^2f}$$

input `integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output

```
(A*a^2*c - C*a^2*c + 2*B*a*b*c - A*b^2*c + C*b^2*c - B*a^2*d + 2*A*a*b*d -
2*C*a*b*d + B*b^2*d)*(f*x + e)/(a^4*f + 2*a^2*b^2*f + b^4*f) + 1/2*(B*a^2
*c - 2*A*a*b*c + 2*C*a*b*c - B*b^2*c + A*a^2*d - C*a^2*d + 2*B*a*b*d - A*b
^2*d + C*b^2*d)*log(tan(f*x + e)^2 + 1)/(a^4*f + 2*a^2*b^2*f + b^4*f) - (B
*a^2*b^2*c - 2*A*a*b^3*c + 2*C*a*b^3*c - B*b^4*c - C*a^4*d + A*a^2*b^2*d -
3*C*a^2*b^2*d + 2*B*a*b^3*d - A*b^4*d)*log(abs(b*tan(f*x + e) + a))/(a^4*
b^2*f + 2*a^2*b^4*f + b^6*f) - (C*a^4*b*c - B*a^3*b^2*c + A*a^2*b^3*c + C*
a^2*b^3*c - B*a*b^4*c + A*b^5*c - C*a^5*d + B*a^4*b*d - A*a^3*b^2*d - C*a^
3*b^2*d + B*a^2*b^3*d - A*a*b^4*d)/((a^2 + b^2)^2*(b*tan(f*x + e) + a)*b^2
*f)
```

Mupad [B] (verification not implemented)

Time = 17.78 (sec) , antiderivative size = 1875, normalized size of antiderivative = 7.08

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int(((c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*
tan(e + f*x))^2,x)
```

output

```
(log(a + b*tan(e + f*x))*(b^4*(A*d + B*c) - b^3*(2*B*a*d - 2*A*a*c + 2*C*a*c) - b^2*(A*a^2*d + B*a^2*c - 3*C*a^2*d) + C*a^4*d))/(f*(b^6 + 2*a^2*b^4 + a^4*b^2)) - (log((A*B*b^4*d^2 - A*B*b^4*c^2 + B*C*a^4*d^2 + B*C*b^4*c^2 - A^2*b^4*c*d + B^2*b^4*c*d + C^2*a^4*c*d - A^2*a*b^3*c^2 + A^2*a*b^3*d^2 + B^2*a*b^3*c^2 - B^2*a*b^3*d^2 - C^2*a*b^3*c^2 + C^2*a*b^3*d^2 + A*B*a^2*b^2*c^2 - A*B*a^2*b^2*d^2 - B*C*a^2*b^2*c^2 + 3*B*C*a^2*b^2*d^2 + A^2*a^2*b^2*c*d - B^2*a^2*b^2*c*d + 3*C^2*a^2*b^2*c*d - A*C*a^4*c*d + A*C*b^4*c*d + 2*A*C*a*b^3*c^2 - 2*A*C*a*b^3*d^2 - 4*A*C*a^2*b^2*c*d + 4*A*B*a*b^3*c*d - 4*B*C*a*b^3*c*d))/(b*(a^2 + b^2)^2) + (tan(e + f*x)*(A^2*b^4*c^2 + B^2*b^4*d^2 + C^2*a^4*d^2 + C^2*b^4*c^2 + C^2*b^4*d^2 + A^2*a^2*b^2*d^2 + B^2*a^2*b^2*c^2 + 3*C^2*a^2*b^2*d^2 - A*C*a^4*d^2 - 2*A*C*b^4*c^2 - A*C*b^4*d^2 - 4*A*C*a^2*b^2*d^2 - 2*A*B*b^4*c*d - B*C*a^4*c*d + B*C*b^4*c*d - 2*A*B*a*b^3*c^2 + 2*A*B*a*b^3*d^2 + 2*B*C*a*b^3*c^2 - 2*B*C*a*b^3*d^2 - 2*A^2*a*b^3*c*d + 2*B^2*a*b^3*c*d - 2*C^2*a*b^3*c*d + 2*A*B*a^2*b^2*c*d - 4*B*C*a^2*b^2*c*d + 4*A*C*a*b^3*c*d))/(b*(a^2 + b^2)^2) + ((c + d*1i)*(A + B*1i - C)*(A*b*c - B*b*d - 4*C*a*d - C*b*c + (tan(e + f*x))*(3*A*b^4*d + 3*B*b^4*c + 2*C*a^4*d - 5*C*b^4*d + 4*A*a*b^3*c - 4*B*a*b^3*d - 4*C*a*b^3*c - A*a^2*b^2*d - B*a^2*b^2*c + C*a^2*b^2*d))/(b*(a^2 + b^2)) + (b*(c + d*1i)*(4*a*b - a^2*tan(e + f*x) + 3*b^2*tan(e + f*x))*(A + B*1i - C)*1i)/(a*1i - b)^2)*1i)/(2*(a*1i - b)^2)*(A*c + A*d*1i + B*c*1i - B*d - C*c - C*d*1i))/(2*...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 943, normalized size of antiderivative = 3.56

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x)
```

output

```
(log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**3*d - log(tan(e + f*x)**2 +
1)*tan(e + f*x)*a**2*b**4*c - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*
b**3*c*d + log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**5*d + 2*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a*b**4*c**2 - log(tan(e + f*x)**2 + 1)*tan(e + f
*x)*b**6*c + log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**5*c*d + log(tan(e +
f*x)**2 + 1)*a**4*b**2*d - log(tan(e + f*x)**2 + 1)*a**3*b**3*c - log(tan(
e + f*x)**2 + 1)*a**3*b**2*c*d + log(tan(e + f*x)**2 + 1)*a**2*b**4*d + 2*
log(tan(e + f*x)**2 + 1)*a**2*b**3*c**2 - log(tan(e + f*x)**2 + 1)*a*b**5*
c + log(tan(e + f*x)**2 + 1)*a*b**4*c*d + 2*log(tan(e + f*x)*b + a)*tan(e
+ f*x)*a**4*b*c*d - 2*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**3*b**3*d + 2
*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b**4*c + 6*log(tan(e + f*x)*b +
a)*tan(e + f*x)*a**2*b**3*c*d - 2*log(tan(e + f*x)*b + a)*tan(e + f*x)*a*
b**5*d - 4*log(tan(e + f*x)*b + a)*tan(e + f*x)*a*b**4*c**2 + 2*log(tan(e
+ f*x)*b + a)*tan(e + f*x)*b**6*c + 2*log(tan(e + f*x)*b + a)*a**5*c*d - 2
*log(tan(e + f*x)*b + a)*a**4*b**2*d + 2*log(tan(e + f*x)*b + a)*a**3*b**3
*c + 6*log(tan(e + f*x)*b + a)*a**3*b**2*c*d - 2*log(tan(e + f*x)*b + a)*a
**2*b**4*d - 4*log(tan(e + f*x)*b + a)*a**2*b**3*c**2 + 2*log(tan(e + f*x)
*b + a)*a*b**5*c - 2*tan(e + f*x)*a**4*b*c*d + 2*tan(e + f*x)*a**3*b**3*c*
f*x + 2*tan(e + f*x)*a**3*b**2*c**2 + 2*tan(e + f*x)*a**2*b**4*d*f*x - 2*t
an(e + f*x)*a**2*b**3*c**2*f*x - 2*tan(e + f*x)*a**2*b**3*c*d + 2*tan(e...
```

3.56
$$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

Optimal result	603
Mathematica [C] (verified)	604
Rubi [A] (verified)	604
Maple [A] (verified)	607
Fricas [B] (verification not implemented)	608
Sympy [F(-2)]	609
Maxima [A] (verification not implemented)	610
Giac [B] (verification not implemented)	611
Mupad [B] (verification not implemented)	612
Reduce [B] (verification not implemented)	613

Optimal result

Integrand size = 43, antiderivative size = 320

$$\int \frac{(c+d \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

$$= \frac{(a^3(Ac - cC - Bd) - 3ab^2(Ac - cC - Bd) + 3a^2b(Bc + (A - C)d) - b^3(Bc + (A - C)d))x}{(a^2 + b^2)^3} + \frac{(3a^2b(Ac - cC - Bd) - b^3(Ac - cC - Bd) - a^3(Bc + (A - C)d) + 3ab^2(Bc + (A - C)d)) \log(a \cos(fx+e) + b \sin(fx+e))}{(a^2 + b^2)^3 f}$$

$$- \frac{(Ab^2 - a(bB - aC))(bc - ad)}{2b^2(a^2 + b^2)f(a + b \tan(e+fx))^2}$$

$$- \frac{a^4Cd + b^4(Bc + Ad) + 2ab^3(Ac - cC - Bd) - a^2b^2(Bc + (A - 3C)d)}{b^2(a^2 + b^2)^2 f(a + b \tan(e+fx))}$$

output

```
(a^3*(A*c-B*d-C*c)-3*a*b^2*(A*c-B*d-C*c)+3*a^2*b*(B*c+(A-C)*d)-b^3*(B*c+(A-C)*d))*x/(a^2+b^2)^3+(3*a^2*b*(A*c-B*d-C*c)-b^3*(A*c-B*d-C*c)-a^3*(B*c+(A-C)*d)+3*a*b^2*(B*c+(A-C)*d))*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)^3/f-1/2*(A*b^2-a*(B*b-C*a))*(-a*d+b*c)/b^2/(a^2+b^2)/f/(a+b*tan(f*x+e))^2-(a^4*C*d+b^4*(A*d+B*c)+2*a*b^3*(A*c-B*d-C*c)-a^2*b^2*(B*c+(A-3*C)*d))/b^2/(a^2+b^2)^2/f/(a+b*tan(f*x+e))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.57 (sec) , antiderivative size = 331, normalized size of antiderivative = 1.03

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

$$\frac{bcC - bBd - aCd}{(a + b \tan(e + fx))^2} - \frac{2bC(c + d \tan(e + fx))}{(a + b \tan(e + fx))^2} + 2b(Bc + (A - C)d) \left(-\frac{i \log(i - \tan(e + fx))}{2(a + ib)^2} + \frac{i \log(i + \tan(e + fx))}{2(a - ib)^2} + \frac{b(2a \log(a + b \tan(e + fx)))}{(a + b \tan(e + fx))^3} \right)$$

=

input

```
Integrate[((c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

```
((b*c*C - b*B*d - a*C*d)/(a + b*Tan[e + f*x])^2 - (2*b*C*(c + d*Tan[e + f*x]))/(a + b*Tan[e + f*x])^2 + 2*b*(B*c + (A - C)*d)*((-1/2*I)*Log[I - Tan[e + f*x]])/(a + I*b)^2 + ((I/2)*Log[I + Tan[e + f*x]])/(a - I*b)^2 + (b*(2*a*Log[a + b*Tan[e + f*x]] - (a^2 + b^2)/(a + b*Tan[e + f*x])))/(a^2 + b^2)^2 - b*(-(A*b*c) + a*B*c + b*c*C + a*A*d + b*B*d - a*C*d)*(Log[I - Tan[e + f*x]]/((-I)*a + b)^3 + Log[I + Tan[e + f*x]]/(I*a + b)^3 + (b*((6*a^2 - 2*b^2)*Log[a + b*Tan[e + f*x]] - ((a^2 + b^2)*(5*a^2 + b^2 + 4*a*b*Tan[e + f*x]))/(a + b*Tan[e + f*x])^2))/(a^2 + b^2)^3)/(2*b^2*f)
```

Rubi [A] (verified)

Time = 2.32 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4118, 3042, 4111, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^3} dx$$

↓ 4118

$$\int \frac{Cda^2 + b(Ac - Cc - Bd)a + (a^2 + b^2)Cd \tan^2(e + fx) + b^2(Bc + Ad) - b(Abc - aBc - bCc - aAd - bBd + aCd) \tan(e + fx)}{(a + b \tan(e + fx))^2} dx$$

$$\frac{b(a^2 + b^2)}{2b^2 f (a^2 + b^2) (a + b \tan(e + fx))^2} (bc - ad) (Ab^2 - a(bB - aC))$$

↓ 3042

$$\int \frac{Cda^2 + b(Ac - Cc - Bd)a + (a^2 + b^2)Cd \tan^2(e + fx) + b^2(Bc + Ad) - b(Abc - aBc - bCc - aAd - bBd + aCd) \tan(e + fx)}{(a + b \tan(e + fx))^2} dx$$

$$\frac{b(a^2 + b^2)}{2b^2 f (a^2 + b^2) (a + b \tan(e + fx))^2} (bc - ad) (Ab^2 - a(bB - aC))$$

↓ 4111

$$\int \frac{b((Ac - Cc - Bd)a^2 + 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd)) - b(-((Bc + (A - C)d)a^2) + 2b(Ac - Cc - Bd)a + b^2(Bc + (A - C)d)) \tan(e + fx)}{a + b \tan(e + fx)} dx$$

$$\frac{b(a^2 + b^2)}{2b^2 f (a^2 + b^2) (a + b \tan(e + fx))^2} (bc - ad) (Ab^2 - a(bB - aC))$$

↓ 3042

$$\int \frac{b((Ac - Cc - Bd)a^2 + 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd)) - b(-((Bc + (A - C)d)a^2) + 2b(Ac - Cc - Bd)a + b^2(Bc + (A - C)d)) \tan(e + fx)}{a^2 + b^2} dx$$

$$\frac{b(a^2 + b^2)}{2b^2 f (a^2 + b^2) (a + b \tan(e + fx))^2} (bc - ad) (Ab^2 - a(bB - aC))$$

↓ 4014

$$\frac{b(- (a^3(d(A - C) + Bc)) + 3a^2b(Ac - Bd - cC) + 3ab^2(d(A - C) + Bc) - b^3(Ac - Bd - cC)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx + bx(a^3(Ac - Bd - cC) + 3a^2b(d(A - C) + Bc) - 3ab^2(Ac - Bd - cC))}{a^2 + b^2}}{a^2 + b^2}$$

$$\frac{b(a^2 + b^2)}{2b^2 f (a^2 + b^2) (a + b \tan(e + fx))^2} (bc - ad) (Ab^2 - a(bB - aC))$$

↓ 3042

$$\frac{b \left(- \left(a^3 (d(A-C)+Bc) \right) + 3a^2 b (Ac-Bd-cC) + 3ab^2 (d(A-C)+Bc) - b^3 (Ac-Bd-cC) \right) \int \frac{b-a \tan(e+fx)}{a+b \tan(e+fx)} dx + bx \left(a^3 (Ac-Bd-cC) + 3a^2 b (d(A-C)+Bc) - 3ab^2 (Ac-Bd-cC) \right)}{a^2+b^2} + \frac{bx \left(a^3 (Ac-Bd-cC) + 3a^2 b (d(A-C)+Bc) - 3ab^2 (Ac-Bd-cC) \right)}{a^2+b^2}$$

$$\frac{(bc-ad)(Ab^2-a(bB-aC))}{2b^2 f (a^2+b^2) (a+b \tan(e+fx))^2}$$

↓ 4013

$$\frac{b \left(- \left(a^3 (d(A-C)+Bc) \right) + 3a^2 b (Ac-Bd-cC) + 3ab^2 (d(A-C)+Bc) - b^3 (Ac-Bd-cC) \right) \log(a \cos(e+fx) + b \sin(e+fx)) + bx \left(a^3 (Ac-Bd-cC) + 3a^2 b (d(A-C)+Bc) - 3ab^2 (Ac-Bd-cC) \right)}{f(a^2+b^2)} + \frac{bx \left(a^3 (Ac-Bd-cC) + 3a^2 b (d(A-C)+Bc) - 3ab^2 (Ac-Bd-cC) \right)}{a^2+b^2}$$

$$\frac{(bc-ad)(Ab^2-a(bB-aC))}{2b^2 f (a^2+b^2) (a+b \tan(e+fx))^2}$$

input

```
Int[((c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

```
-1/2*((A*b^2 - a*(b*B - a*C))*(b*c - a*d))/(b^2*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^2) + (((b*(a^3*(A*c - c*C - B*d) - 3*a*b^2*(A*c - c*C - B*d) + 3*a^2*b*(B*c + (A - C)*d) - b^3*(B*c + (A - C)*d))*x)/(a^2 + b^2) + (b*(3*a^2*b*(A*c - c*C - B*d) - b^3*(A*c - c*C - B*d) - a^3*(B*c + (A - C)*d) + 3*a*b^2*(B*c + (A - C)*d))*Log[a*Cos[e + f*x] + b*Sin[e + f*x]]/((a^2 + b^2)*f))/(a^2 + b^2) - (a^4*C*d + b^4*(B*c + A*d) + 2*a*b^3*(A*c - c*C - B*d) - a^2*b^2*(B*c + (A - 3*C)*d))/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x]))/(b*(a^2 + b^2))
```

Defintions of rubi rules used

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4013

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;`
`FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /;`
`FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4118 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 494, normalized size of antiderivative = 1.54

method	result
derivativedivides	$\frac{(A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c) \ln(1 + \tan(fx + e))}{2} + \frac{(A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c)}{(a^2 + b^2)^3}$
default	$\frac{(A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c) \ln(1 + \tan(fx + e))}{2} + \frac{(A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c)}{(a^2 + b^2)^3}$
norman	$\frac{(A a^2 b^2 d - 2A a b^3 c - A b^4 d + B a^2 b^2 c + 2B a b^3 d - B b^4 c - a^4 C d - 3C a^2 b^2 d + 2C a b^3 c) \tan(fx + e)}{fb(a^4 + 2b^2 a^2 + b^4)} + \frac{(A a^3 c + 3A a^2 b d - 3A a b^2 c - A b^3 d)}{(a^2 + b^2)^3}$
risch	Expression too large to display
parallelrisch	Expression too large to display

input `int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, method=_RETURNVERBOSE)`

output
$$\frac{1}{f} \left(\frac{1}{(a^2 + b^2)^3} \left(\frac{1}{2} (A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c) \ln(1 + \tan(fx + e)) + (A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c) \right) \right. \\ \left. - \frac{1}{2} \frac{(-A a^2 b^2 d + 2A a b^3 c - A b^4 d + B a^2 b^2 c + 2B a b^3 d - B b^4 c - a^4 C d - 3C a^2 b^2 d + 2C a b^3 c) \arctan(\tan(fx + e))}{b^2 (a^2 + b^2)} - \frac{(-A a^2 b^2 d + 2A a b^3 c + A b^4 d - B a^2 b^2 c - 2B a^2 b^3 d + B b^4 c + C a^4 d + 3C a^2 b^2 d - 2C a^2 b^3 c)}{(a^2 + b^2)^2} \frac{1}{b^2} \right. \\ \left. - \frac{(A a^3 d - 3A a^2 b c - 3A a b^2 d + A b^3 c + B a^3 c + 3B a^2 b d - 3B a b^2 c - B b^3 d - a^3 C d + 3C a^2 b c + 3C a b^2 d - C b^3 c)}{(a^2 + b^2)^3} \ln(a + b \tan(fx + e)) \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 987 vs. 2(316) = 632.

Time = 0.11 (sec) , antiderivative size = 987, normalized size of antiderivative = 3.08

$$\int \frac{(c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

output

```

1/2*(2*(((A - C)*a^5 + 3*B*a^4*b - 3*(A - C)*a^3*b^2 - B*a^2*b^3)*c - (B*a
^5 - 3*(A - C)*a^4*b - 3*B*a^3*b^2 + (A - C)*a^2*b^3)*d)*f*x + (2*(((A - C
)*a^3*b^2 + 3*B*a^2*b^3 - 3*(A - C)*a*b^4 - B*b^5)*c - (B*a^3*b^2 - 3*(A -
C)*a^2*b^3 - 3*B*a*b^4 + (A - C)*b^5)*d)*f*x + (C*a^4*b - 3*B*a^3*b^2 + 5
*(A - C)*a^2*b^3 + 3*B*a*b^4 - A*b^5)*c + (C*a^5 + B*a^4*b - (3*A - 7*C)*a
^3*b^2 - 5*B*a^2*b^3 + 3*A*a*b^4)*d)*tan(f*x + e)^2 - (3*C*a^4*b - 5*B*a^3
*b^2 + (7*A - 3*C)*a^2*b^3 + B*a*b^4 + A*b^5)*c + (C*a^5 - 3*B*a^4*b + 5*(
A - C)*a^3*b^2 + 3*B*a^2*b^3 - A*a*b^4)*d - (((B*a^3*b^2 - 3*(A - C)*a^2*b
^3 - 3*B*a*b^4 + (A - C)*b^5)*c + ((A - C)*a^3*b^2 + 3*B*a^2*b^3 - 3*(A -
C)*a*b^4 - B*b^5)*d)*tan(f*x + e)^2 + (B*a^5 - 3*(A - C)*a^4*b - 3*B*a^3*b
^2 + (A - C)*a^2*b^3)*c + ((A - C)*a^5 + 3*B*a^4*b - 3*(A - C)*a^3*b^2 - B
*a^2*b^3)*d + 2*((B*a^4*b - 3*(A - C)*a^3*b^2 - 3*B*a^2*b^3 + (A - C)*a*b^
4)*c + ((A - C)*a^4*b + 3*B*a^3*b^2 - 3*(A - C)*a^2*b^3 - B*a*b^4)*d)*tan(
f*x + e))*log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e
)^2 + 1)) + 2*(2*(((A - C)*a^4*b + 3*B*a^3*b^2 - 3*(A - C)*a^2*b^3 - B*a*b
^4)*c - (B*a^4*b - 3*(A - C)*a^3*b^2 - 3*B*a^2*b^3 + (A - C)*a*b^4)*d)*f*x
+ (C*a^5 - 2*B*a^4*b + 3*(A - C)*a^3*b^2 + 3*B*a^2*b^3 - (3*A - 2*C)*a*b^
4 - B*b^5)*c + (B*a^5 - (2*A - 3*C)*a^4*b - 3*B*a^3*b^2 + 3*(A - C)*a^2*b^
3 + 2*B*a*b^4 - A*b^5)*d)*tan(f*x + e))/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6
+ b^8)*f*tan(f*x + e)^2 + 2*(a^7*b + 3*a^5*b^3 + 3*a^3*b^5 + a*b^7)*f*t...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

= Exception raised: AttributeError

input

```

integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e
))**3,x)

```

output

```

Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'

```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 574, normalized size of antiderivative = 1.79

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

$$= \frac{2(((A-C)a^3 + 3Ba^2b - 3(A-C)ab^2 - Bb^3)c - (Ba^3 - 3(A-C)a^2b - 3Bab^2 + (A-C)b^3)d)(fx+e)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{2((Ba^3 - 3(A-C)a^2b - 3Bab^2 + (A-C)b^3)d)(fx+e)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6}$$

input

```
integrate((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^3,x, algorithm="maxima")
```

output

```
1/2*(2*(((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c - (B*a^3 - 3
*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*d)*(f*x + e)/(a^6 + 3*a^4*b^2 +
3*a^2*b^4 + b^6) - 2*((B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*
c + ((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*d)*log(b*tan(f*x +
e) + a)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + ((B*a^3 - 3*(A - C)*a^2*b -
3*B*a*b^2 + (A - C)*b^3)*c + ((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 -
B*b^3)*d)*log(tan(f*x + e)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (
(C*a^4*b - 3*B*a^3*b^2 + (5*A - 3*C)*a^2*b^3 + B*a*b^4 + A*b^5)*c + (C*a^5
+ B*a^4*b - (3*A - 5*C)*a^3*b^2 - 3*B*a^2*b^3 + A*a*b^4)*d - 2*((B*a^2*b^
3 - 2*(A - C)*a*b^4 - B*b^5)*c - (C*a^4*b - (A - 3*C)*a^2*b^3 - 2*B*a*b^4
+ A*b^5)*d)*tan(f*x + e))/(a^6*b^2 + 2*a^4*b^4 + a^2*b^6 + (a^4*b^4 + 2*a^
2*b^6 + b^8)*tan(f*x + e)^2 + 2*(a^5*b^3 + 2*a^3*b^5 + a*b^7)*tan(f*x + e
))/f
```


Mupad [B] (verification not implemented)

Time = 12.21 (sec) , antiderivative size = 502, normalized size of antiderivative = 1.57

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx =$$

$$\frac{Ab^5c + Ca^5d + Aab^4d + Bab^4c + Ba^4bd + Ca^4bc + 5Aa^2b^3c - 3Aa^3b^2d - 3Ba^3b^2c - 3Ba^2b^3d - 3Ca^2b^3c + 5Ca^3b^2d}{2b^2(a^4 + 2a^2b^2 + b^4)} + \frac{\tan(e + fx)}{f(a^2 + 2ab \tan(e + fx) + b^2 \tan^2(e + fx))}$$

$$- \frac{\ln(\tan(e + fx) + 1i)(Bd + Ad1i + Bc1i - Ac + Cc - Cd1i)}{2f(-a^31i - 3a^2b + ab^23i + b^3)}$$

$$- \frac{\ln(\tan(e + fx) - 1i)(Ad + Bc - Cd - Ac1i + Bd1i + Cc1i)}{2f(-a^3 - a^2b3i + 3ab^2 + b^31i)}$$

$$- \frac{\ln(a + b \tan(e + fx))((Ad + Bc - Cd)a^3 + (3Bd - 3Ac + 3Cc)a^2b + (3Cd - 3Bc - 3Ad)b)}{f(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)}$$

input

```
int(((c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*
tan(e + f*x))^3,x)
```

output

```
- ((A*b^5*c + C*a^5*d + A*a*b^4*d + B*a*b^4*c + B*a^4*b*d + C*a^4*b*c + 5*
A*a^2*b^3*c - 3*A*a^3*b^2*d - 3*B*a^3*b^2*c - 3*B*a^2*b^3*d - 3*C*a^2*b^3*
c + 5*C*a^3*b^2*d)/(2*b^2*(a^4 + b^4 + 2*a^2*b^2)) + (tan(e + f*x)*(A*b^4*
d + B*b^4*c + C*a^4*d + 2*A*a*b^3*c - 2*B*a*b^3*d - 2*C*a*b^3*c - A*a^2*b^
2*d - B*a^2*b^2*c + 3*C*a^2*b^2*d))/(b*(a^4 + b^4 + 2*a^2*b^2)))/(f*(a^2 +
b^2*tan(e + f*x)^2 + 2*a*b*tan(e + f*x))) - (log(tan(e + f*x) + 1i)*(A*d*
1i - A*c + B*c*1i + B*d + C*c - C*d*1i))/(2*f*(a*b^2*3i - 3*a^2*b - a^3*1i
+ b^3)) - (log(tan(e + f*x) - 1i)*(A*d - A*c*1i + B*c + B*d*1i + C*c*1i -
C*d))/(2*f*(3*a*b^2 - a^2*b*3i - a^3 + b^3*1i)) - (log(a + b*tan(e + f*x)
)*(a^3*(A*d + B*c - C*d) - b^3*(B*d - A*c + C*c) + a^2*b*(3*B*d - 3*A*c +
3*C*c) - a*b^2*(3*A*d + 3*B*c - 3*C*d)))/(f*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4
*b^2))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2008, normalized size of antiderivative = 6.28

$$\int \frac{(c + d \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input `int((c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x)`

output

```
(log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**3*d - 2*log(tan(e + f*x)
**2 + 1)*tan(e + f*x)**2*a**4*b**4*c - log(tan(e + f*x)**2 + 1)*tan(e + f*
x)**2*a**4*b**3*c*d + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**4
*c**2 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**6*c + 3*log(tan
(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**5*c*d - log(tan(e + f*x)**2 + 1)
*tan(e + f*x)**2*a*b**7*d - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**
6*c**2 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**6*b**2*d - 4*log(tan(e
 + f*x)**2 + 1)*tan(e + f*x)*a**5*b**3*c - 2*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)*a**5*b**2*c*d + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**4*b**3
*c**2 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**5*c + 6*log(tan(e
 + f*x)**2 + 1)*tan(e + f*x)*a**3*b**4*c*d - 2*log(tan(e + f*x)**2 + 1)*tan
(e + f*x)*a**2*b**6*d - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**5*
c**2 + log(tan(e + f*x)**2 + 1)*a**7*b*d - 2*log(tan(e + f*x)**2 + 1)*a**6
*b**2*c - log(tan(e + f*x)**2 + 1)*a**6*b*c*d + 3*log(tan(e + f*x)**2 + 1)
*a**5*b**2*c**2 - 2*log(tan(e + f*x)**2 + 1)*a**4*b**4*c + 3*log(tan(e + f
*x)**2 + 1)*a**4*b**3*c*d - log(tan(e + f*x)**2 + 1)*a**3*b**5*d - log(tan
(e + f*x)**2 + 1)*a**3*b**4*c**2 - 2*log(tan(e + f*x)*b + a)*tan(e + f*x)*
**2*a**5*b**3*d + 4*log(tan(e + f*x)*b + a)*tan(e + f*x)**2*a**4*b**4*c + 2
*log(tan(e + f*x)*b + a)*tan(e + f*x)**2*a**4*b**3*c*d - 6*log(tan(e + f*x)
)*b + a)*tan(e + f*x)**2*a**3*b**4*c**2 + 4*log(tan(e + f*x)*b + a)*tan...
```

3.57 $\int (a+b \tan(e+fx))^3 (c+d \tan(e+fx))^2 (A+B \tan(e+fx) + C \tan^2(e+fx)) dx$

Optimal result	614
Mathematica [C] (verified)	615
Rubi [A] (verified)	616
Maple [A] (warning: unable to verify)	622
Fricas [A] (verification not implemented)	623
Sympy [B] (verification not implemented)	623
Maxima [A] (verification not implemented)	624
Giac [B] (verification not implemented)	625
Mupad [B] (verification not implemented)	626
Reduce [B] (verification not implemented)	627

Optimal result

Integrand size = 45, antiderivative size = 661

$$\begin{aligned}
& \int (a+b \tan(e+fx))^3 (c+d \tan(e+fx))^2 (A+B \tan(e+fx) + C \tan^2(e+fx)) dx \\
&= -((a^3(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) - 3ab^2(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) \\
&\quad + 3a^2b(2c(A - C)d + B(c^2 - d^2)) - b^3(2c(A - C)d + B(c^2 - d^2))) x \\
&\quad + \frac{(3a^2b(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) - b^3(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) - a^3(2c(A - C)d + B(c^2 - d^2)))}{f} \\
&\quad + \frac{d(3a^2b(Ac - cC - Bd) - b^3(Ac - cC - Bd) + a^3(Bc + (A - C)d) - 3ab^2(Bc + (A - C)d)) \tan(e+fx)}{f} \\
&\quad + \frac{(a^3B - 3ab^2B + 3a^2b(A - C) - b^3(A - C))(c + d \tan(e+fx))^2}{2f} \\
&\quad + \frac{(4a^3Cd^3 - 3a^2bd^2(3cC - 16Bd) + 3ab^2d(2c^2C - 5Bcd + 20(A - C)d^2) - b^3(c^3C - 2Bc^2d + 5c(A - C)d^2))}{60d^4f} \\
&\quad + \frac{b(5b(Ab + aB - bC)d^2 + (bc - ad)(bcC - 2bBd - aCd)) \tan(e+fx)(c + d \tan(e+fx))^3}{20d^3f} \\
&\quad - \frac{(bcC - 2bBd - aCd)(a + b \tan(e+fx))^2(c + d \tan(e+fx))^3}{10d^2f} \\
&\quad + \frac{C(a + b \tan(e+fx))^3(c + d \tan(e+fx))^3}{6df}
\end{aligned}$$

output

```

-(a^3*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-3*a*b^2*(c^2*C+2*B*c*d-C*d^2-A*(c^
2-d^2))+3*a^2*b*(2*c*(A-C)*d+B*(c^2-d^2))-b^3*(2*c*(A-C)*d+B*(c^2-d^2)))*x
+(3*a^2*b*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-b^3*(c^2*C+2*B*c*d-C*d^2-A*(c^
2-d^2))-a^3*(2*c*(A-C)*d+B*(c^2-d^2))+3*a*b^2*(2*c*(A-C)*d+B*(c^2-d^2)))*l
n(cos(f*x+e))/f+d*(3*a^2*b*(A*c-B*d-C*c)-b^3*(A*c-B*d-C*c)+a^3*(B*c+(A-C)*
d)-3*a*b^2*(B*c+(A-C)*d))*tan(f*x+e)/f+1/2*(B*a^3-3*B*a*b^2+3*a^2*b*(A-C)-
b^3*(A-C))*(c+d*tan(f*x+e))^2/f+1/60*(4*a^3*C*d^3-3*a^2*b*d^2*(-16*B*d+3*C
*c)+3*a*b^2*d*(2*c^2*C-5*B*c*d+20*(A-C)*d^2)-b^3*(c^3*C-2*B*c^2*d+5*c*(A-C
)*d^2+20*B*d^3))*(c+d*tan(f*x+e))^3/d^4/f+1/20*b*(5*b*(A*b+B*a-C*b)*d^2+(-
a*d+b*c)*(-2*B*b*d-C*a*d+C*b*c))*tan(f*x+e)*(c+d*tan(f*x+e))^3/d^3/f-1/10*
(-2*B*b*d-C*a*d+C*b*c)*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3/d^2/f+1/6*C*(
a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^3/d/f

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.43 (sec) , antiderivative size = 573, normalized size of antiderivative = 0.87

$$\begin{aligned}
 & \int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 &= \frac{C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3}{6df} \\
 &+ \frac{3(bcC - 2bBd - aCd)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3}{5df} + \frac{3b(5b(Ab + aB - bC)d^2 + (bc - ad)(bcC - 2bBd - aCd) \tan(e + fx)(c + d \tan(e + fx))^3)}{2df}
 \end{aligned}$$

input

```

Integrate[(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x
] + C*Tan[e + f*x]^2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^3)/(6*d*f) + ((-3*(b*c*C -
2*b*B*d - a*C*d)*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^3)/(5*d*f) +
((3*b*(5*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(b*c*C - 2*b*B*d - a*C*d))*
Tan[e + f*x]*(c + d*Tan[e + f*x])^3)/(2*d*f) - (((-24*a*d*(5*b*(A*b + a*B
- b*C)*d^2 + (b*c - a*d)*(b*c*C - 2*b*B*d - a*C*d)) + b*(-120*(a^2*B - b^2
*B + 2*a*b*(A - C))*d^3 + 6*c*(5*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(b*
c*C - 2*b*B*d - a*C*d))))*(c + d*Tan[e + f*x])^3)/(3*d*f) - (60*(d^2*(3*a^
2*b*(A*c - c*C + B*d) - b^3*(A*c - c*C + B*d) + a^3*(B*c - (A - C)*d) - 3*
a*b^2*(B*c - (A - C)*d))*(I*(c + I*d)^2*Log[I - Tan[e + f*x]] - I*(c - I*d
)^2*Log[I + Tan[e + f*x]] - 2*d^2*Tan[e + f*x]) + (a^3*B - 3*a*b^2*B + 3*a
^2*b*(A - C) - b^3*(A - C))*d^2*((I*c - d)^3*Log[I - Tan[e + f*x]] - (I*c
+ d)^3*Log[I + Tan[e + f*x]] + 6*c*d^2*Tan[e + f*x] + d^3*Tan[e + f*x]^2))
)/f)/(4*d))/(5*d))/(6*d)
```

Rubi [A] (verified)

Time = 5.69 (sec) , antiderivative size = 696, normalized size of antiderivative = 1.05, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.378$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4120, 25, 3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\frac{\int -3(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 ((bcC - adC - 2bBd) \tan^2(e + fx) - 2(Ab - Cb + aB)d \tan(e + fx) + C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3) dx}{6df}$$

$$\downarrow 27$$

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{\int (a + b \tan(e + fx))^2(c + d \tan(e + fx))^2 ((bcC - adC - 2bBd) \tan^2(e + fx) - 2(Ab - Cb + aB)d \tan(e + fx))}{2d}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{\int (a + b \tan(e + fx))^2(c + d \tan(e + fx))^2 ((bcC - adC - 2bBd) \tan(e + fx)^2 - 2(Ab - Cb + aB)d \tan(e + fx))}{2d}$$

↓ 4130

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{\int -2(a + b \tan(e + fx))(c + d \tan(e + fx))^2(c(cC - 2Bd)b^2 - ad(2cC + 3Bd)b + a^2(5A - 4C)d^2 + (5b(Ab - Cb + aB)d^2 + (bc - ad)(bcC - adC - 2bBd)) \tan(e + fx))}{5d}}{2d}$$

↓ 27

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \int (a + b \tan(e + fx))(c + d \tan(e + fx))^2(c(cC - 2Bd)b^2 - ad(2cC + 3Bd)b + a^2(5A - 4C)d^2)}{2d}}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \int (a + b \tan(e + fx))(c + d \tan(e + fx))^2(c(cC - 2Bd)b^2 - ad(2cC + 3Bd)b + a^2(5A - 4C)d^2)}{2d}}$$

↓ 4120

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{b \tan(e + fx)(c + d \tan(e + fx))^3 (5bd^2(aB + Ab - bC) + (bc - ad)(-aCd - 2bBd + bcC))}{4df} \right)}{2d}}$$

↓ 25

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{\int (c + d \tan(e + fx))^2 (-c(Cc^2 - 2Bdc + 5(A - C)d^2)b^3 + 3acd(2cC - 5Bd)b^2 - 3a^2d^2(3cC + 4ad))}{4df} \right)}{2d}}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3}{6df} - \\ & \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{\int (c + d \tan(e + fx))^2 (-c(Cc^2 - 2Bdc + 5(A-C)d^2)b^3 + 3acd(2cC - 5Bd)b^2 - 3a^2d^2(3cC + 4 \end{aligned}$$

$$\begin{aligned} & \downarrow 4113 \\ & \frac{C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3}{6df} - \\ & \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{\int (c + d \tan(e + fx))^2 (20(Ba^3 + 3b(A-C)a^2 - 3b^2Ba - b^3(A-C))d^3 \tan(e + fx) - 20(-((A \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3}{6df} - \\ & \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{\int (c + d \tan(e + fx))^2 (20(Ba^3 + 3b(A-C)a^2 - 3b^2Ba - b^3(A-C))d^3 \tan(e + fx) - 20(-((A \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3}{6df} - \\ & \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{\int (c + d \tan(e + fx)) (20((Ac - Cc - Bd)a^3 - 3b(Bc + (A - C)d)a^2 - 3b^2(Ac - Cc - Bd)a + b^3(B \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^3}{6df} - \\ & \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{\int (c + d \tan(e + fx)) (20((Ac - Cc - Bd)a^3 - 3b(Bc + (A - C)d)a^2 - 3b^2(Ac - Cc - Bd)a + b^3(B \end{aligned}$$

$$\downarrow 4008$$

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{-20d^3(-a^3(2cd(A-C) + B(c^2 - d^2))) + 3a^2b(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) + 3ab^2}{f} \right)}{2}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{-20d^3(-a^3(2cd(A-C) + B(c^2 - d^2))) + 3a^2b(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) + 3ab^2}{f} \right)}{2}$$

↓ 3956

$$\frac{C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^3}{6df} - \frac{(-aCd - 2bBd + bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{2 \left(\frac{20d^3 \log(\cos(e + fx))(-a^3(2cd(A-C) + B(c^2 - d^2))) + 3a^2b(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) + 3ab^2}{f} \right)}{2}$$

input

```
Int[(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*
Tan[e + f*x]^2), x]
```


output

```
(C*(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^3)/(6*d*f) - (((b*c*C - 2*b
*B*d - a*C*d)*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^3)/(5*d*f) - (2*
((b*(5*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(b*c*C - 2*b*B*d - a*C*d))*Ta
n[e + f*x]*(c + d*Tan[e + f*x])^3)/(4*d*f) + (-20*d^3*(a^3*(c^2*C + 2*B*c*
d - C*d^2 - A*(c^2 - d^2)) - 3*a*b^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d
^2)) + 3*a^2*b*(2*c*(A - C)*d + B*(c^2 - d^2)) - b^3*(2*c*(A - C)*d + B*(c
^2 - d^2)))*x + (20*d^3*(3*a^2*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2))
- b^3*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - a^3*(2*c*(A - C)*d + B*
(c^2 - d^2)) + 3*a*b^2*(2*c*(A - C)*d + B*(c^2 - d^2))*Log[Cos[e + f*x]])
/f + (20*d^4*(3*a^2*b*(A*c - c*C - B*d) - b^3*(A*c - c*C - B*d) + a^3*(B*c
+ (A - C)*d) - 3*a*b^2*(B*c + (A - C)*d))*Tan[e + f*x])/f + (10*(a^3*B -
3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C))*d^3*(c + d*Tan[e + f*x])^2)/f +
((4*a^3*C*d^3 - 3*a^2*b*d^2*(3*c*C - 16*B*d) + 3*a*b^2*d*(2*c^2*C - 5*B*c
*d + 20*(A - C)*d^2) - b^3*(c^3*C - 2*B*c^2*d + 5*c*(A - C)*d^2 + 20*B*d^3
))*(c + d*Tan[e + f*x])^3)/(3*d*f)/(4*d))/(5*d))/(2*d)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4008

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

rule 4011

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp
[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp
[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e.
_) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (warning: unable to verify)

Time = 0.40 (sec) , antiderivative size = 546, normalized size of antiderivative = 0.83

method	result
parts	$\frac{(2Aa^3cd+3Aa^2bc^2+Ba^3c^2)\ln(1+\tan(fx+e))}{2f} + \frac{(Bb^3d^2+3Ca^2b^2d^2+2Cb^3cd)\left(\frac{\tan(fx+e)^5}{5} - \frac{\tan(fx+e)^3}{3} + \tan(fx+e)\right)}{f}$
norman	$(Aa^3c^2 - Aa^3d^2 - 6Aa^2bcd - 3Aa^2b^2c^2 + 3Aa^2b^2d^2 + 2Ab^3cd - 2Ba^3cd - 3Ba^2bc^2)$
derivativedivides	Expression too large to display
default	Expression too large to display
parallelrisch	Expression too large to display
risch	Expression too large to display

input

```
int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x,method=_RETURNVERBOSE)
```

output

```
1/2*(2*A*a^3*c*d+3*A*a^2*b*c^2+B*a^3*c^2)/f*ln(1+tan(f*x+e)^2)+(B*b^3*d^2+
3*C*a*b^2*d^2+2*C*b^3*c*d)/f*(1/5*tan(f*x+e)^5-1/3*tan(f*x+e)^3+tan(f*x+e)
-arctan(tan(f*x+e)))+(A*b^3*d^2+3*B*a*b^2*d^2+2*B*b^3*c*d+3*C*a^2*b*d^2+6*
C*a*b^2*c*d+C*b^3*c^2)/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(f
*x+e)^2))+(A*a^3*d^2+6*A*a^2*b*c*d+3*A*a*b^2*c^2+2*B*a^3*c*d+3*B*a^2*b*c^2
+C*a^3*c^2)/f*(tan(f*x+e)-arctan(tan(f*x+e)))+(3*A*a*b^2*d^2+2*A*b^3*c*d+3
*B*a^2*b*d^2+6*B*a*b^2*c*d+B*b^3*c^2+C*a^3*d^2+6*C*a^2*b*c*d+3*C*a*b^2*c^2
)/f*(1/3*tan(f*x+e)^3-tan(f*x+e)+arctan(tan(f*x+e)))+(3*A*a^2*b*d^2+6*A*a*
b^2*c*d+A*b^3*c^2+B*a^3*d^2+6*B*a^2*b*c*d+3*B*a*b^2*c^2+2*C*a^3*c*d+3*C*a^
2*b*c^2)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)^2))+A*a^3*c^2*x+C*b^3*d^2
/f*(1/6*tan(f*x+e)^6-1/4*tan(f*x+e)^4+1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)
^2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 690, normalized size of antiderivative = 1.04

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

```
1/60*(10*C*b^3*d^2*tan(f*x + e)^6 + 12*(2*C*b^3*c*d + (3*C*a*b^2 + B*b^3)*d^2)*tan(f*x + e)^5 + 15*(C*b^3*c^2 + 2*(3*C*a*b^2 + B*b^3)*c*d + (3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*d^2)*tan(f*x + e)^4 + 20*((3*C*a*b^2 + B*b^3)*c^2 + 2*(3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*c*d + (C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*d^2)*tan(f*x + e)^3 + 60*((A - C)*a^3 - 3*B*a^2*b + b - 3*(A - C)*a*b^2 + B*b^3)*c^2 - 2*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c*d - ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*d^2)*f*x + 30*((3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*c^2 + 2*(C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*c*d + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^2)*tan(f*x + e)^2 - 30*((B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c^2 + 2*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c*d - (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^2)*log(1/(tan(f*x + e)^2 + 1)) + 60*((C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*c^2 + 2*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c*d + ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*d^2)*tan(f*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1819 vs. 2(604) = 1208.

Time = 0.40 (sec) , antiderivative size = 1819, normalized size of antiderivative = 2.75

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))**3*(c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A**3*c**2*x + A**3*c*d*log(tan(e + f*x)**2 + 1)/f - A**3*
d**2*x + A**3*d**2*tan(e + f*x)/f + 3*A**2*b*c**2*log(tan(e + f*x)**2
+ 1)/(2*f) - 6*A**2*b*c*d*x + 6*A**2*b*c*d*tan(e + f*x)/f - 3*A**2*b
*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*A**2*b*d**2*tan(e + f*x)**2/(2*
f) - 3*A*a*b**2*c**2*x + 3*A*a*b**2*c**2*tan(e + f*x)/f - 3*A*a*b**2*c*d*log(tan(e + f*x)**2 + 1)/f + 3*A*a*b**2*c*d*tan(e + f*x)**2/f + 3*A*a*b**2*
d**2*x + A*a*b**2*d**2*tan(e + f*x)**3/f - 3*A*a*b**2*d**2*tan(e + f*x)/f
- A*b**3*c**2*log(tan(e + f*x)**2 + 1)/(2*f) + A*b**3*c**2*tan(e + f*x)**2
/(2*f) + 2*A*b**3*c*d*x + 2*A*b**3*c*d*tan(e + f*x)**3/(3*f) - 2*A*b**3*c*
d*tan(e + f*x)/f + A*b**3*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + A*b**3*d**
2*tan(e + f*x)**4/(4*f) - A*b**3*d**2*tan(e + f*x)**2/(2*f) + B*a**3*c**2*
log(tan(e + f*x)**2 + 1)/(2*f) - 2*B*a**3*c*d*x + 2*B*a**3*c*d*tan(e + f*x)
)/f - B*a**3*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + B*a**3*d**2*tan(e + f*x)
)**2/(2*f) - 3*B*a**2*b*c**2*x + 3*B*a**2*b*c**2*tan(e + f*x)/f - 3*B*a**2
*b*c*d*log(tan(e + f*x)**2 + 1)/f + 3*B*a**2*b*c*d*tan(e + f*x)**2/f + 3*B
*a**2*b*d**2*x + B*a**2*b*d**2*tan(e + f*x)**3/f - 3*B*a**2*b*d**2*tan(e +
f*x)/f - 3*B*a*b**2*c**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*a*b**2*c**2
*tan(e + f*x)**2/(2*f) + 6*B*a*b**2*c*d*x + 2*B*a*b**2*c*d*tan(e + f*x)**3
/f - 6*B*a*b**2*c*d*tan(e + f*x)/f + 3*B*a*b**2*d**2*log(tan(e + f*x)**2 +
1)/(2*f) + 3*B*a*b**2*d**2*tan(e + f*x)**4/(4*f) - 3*B*a*b**2*d**2*tan...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 691, normalized size of antiderivative = 1.05

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+
e)^2),x, algorithm="maxima")
```

output

```

1/60*(10*C*b^3*d^2*tan(f*x + e)^6 + 12*(2*C*b^3*c*d + (3*C*a*b^2 + B*b^3)*
d^2)*tan(f*x + e)^5 + 15*(C*b^3*c^2 + 2*(3*C*a*b^2 + B*b^3)*c*d + (3*C*a^2
*b + 3*B*a*b^2 + (A - C)*b^3)*d^2)*tan(f*x + e)^4 + 20*((3*C*a*b^2 + B*b^3
)*c^2 + 2*(3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*c*d + (C*a^3 + 3*B*a^2*b +
3*(A - C)*a*b^2 - B*b^3)*d^2)*tan(f*x + e)^3 + 30*((3*C*a^2*b + 3*B*a*b^2
+ (A - C)*b^3)*c^2 + 2*(C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*c*d
+ (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^2)*tan(f*x + e)^2
+ 60*(((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c^2 - 2*(B*a^3 +
3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c*d - ((A - C)*a^3 - 3*B*a^2*b
- 3*(A - C)*a*b^2 + B*b^3)*d^2)*(f*x + e) + 30*((B*a^3 + 3*(A - C)*a^2*b
- 3*B*a*b^2 - (A - C)*b^3)*c^2 + 2*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*
b^2 + B*b^3)*c*d - (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^2
)*log(tan(f*x + e)^2 + 1) + 60*((C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b
^3)*c^2 + 2*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c*d + ((A
- C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*d^2)*tan(f*x + e))/f

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1413 vs. $2(650) = 1300$.

Time = 0.96 (sec) , antiderivative size = 1413, normalized size of antiderivative = 2.14

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```

integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+
e)^2),x, algorithm="giac")

```

output

```
(A*a^3*c^2 - C*a^3*c^2 - 3*B*a^2*b*c^2 - 3*A*a*b^2*c^2 + 3*C*a*b^2*c^2 + B
*b^3*c^2 - 2*B*a^3*c*d - 6*A*a^2*b*c*d + 6*C*a^2*b*c*d + 6*B*a*b^2*c*d + 2
*A*b^3*c*d - 2*C*b^3*c*d - A*a^3*d^2 + C*a^3*d^2 + 3*B*a^2*b*d^2 + 3*A*a*b
^2*d^2 - 3*C*a*b^2*d^2 - B*b^3*d^2)*(f*x + e)/f + 1/2*(B*a^3*c^2 + 3*A*a^2
*b*c^2 - 3*C*a^2*b*c^2 - 3*B*a*b^2*c^2 - A*b^3*c^2 + C*b^3*c^2 + 2*A*a^3*c
*d - 2*C*a^3*c*d - 6*B*a^2*b*c*d - 6*A*a*b^2*c*d + 6*C*a*b^2*c*d + 2*B*b^3
*c*d - B*a^3*d^2 - 3*A*a^2*b*d^2 + 3*C*a^2*b*d^2 + 3*B*a*b^2*d^2 + A*b^3*d
^2 - C*b^3*d^2)*log(tan(f*x + e)^2 + 1)/f + 1/60*(10*C*b^3*d^2*f^5*tan(f*x
+ e)^6 + 24*C*b^3*c*d*f^5*tan(f*x + e)^5 + 36*C*a*b^2*d^2*f^5*tan(f*x + e
)^5 + 12*B*b^3*d^2*f^5*tan(f*x + e)^5 + 15*C*b^3*c^2*f^5*tan(f*x + e)^4 +
90*C*a*b^2*c*d*f^5*tan(f*x + e)^4 + 30*B*b^3*c*d*f^5*tan(f*x + e)^4 + 45*C
*a^2*b*d^2*f^5*tan(f*x + e)^4 + 45*B*a*b^2*d^2*f^5*tan(f*x + e)^4 + 15*A*b
^3*d^2*f^5*tan(f*x + e)^4 - 15*C*b^3*d^2*f^5*tan(f*x + e)^4 + 60*C*a*b^2*c
^2*f^5*tan(f*x + e)^3 + 20*B*b^3*c^2*f^5*tan(f*x + e)^3 + 120*C*a^2*b*c*d*
f^5*tan(f*x + e)^3 + 120*B*a*b^2*c*d*f^5*tan(f*x + e)^3 + 40*A*b^3*c*d*f^5
*tan(f*x + e)^3 - 40*C*b^3*c*d*f^5*tan(f*x + e)^3 + 20*C*a^3*d^2*f^5*tan(f
*x + e)^3 + 60*B*a^2*b*d^2*f^5*tan(f*x + e)^3 + 60*A*a*b^2*d^2*f^5*tan(f*x
+ e)^3 - 60*C*a*b^2*d^2*f^5*tan(f*x + e)^3 - 20*B*b^3*d^2*f^5*tan(f*x + e
)^3 + 90*C*a^2*b*c^2*f^5*tan(f*x + e)^2 + 90*B*a*b^2*c^2*f^5*tan(f*x + e)^
2 + 30*A*b^3*c^2*f^5*tan(f*x + e)^2 - 30*C*b^3*c^2*f^5*tan(f*x + e)^2 + ...
```

Mupad [B] (verification not implemented)

Time = 5.83 (sec) , antiderivative size = 891, normalized size of antiderivative = 1.35

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
int((a + b*tan(e + f*x))^3*(c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*
tan(e + f*x)^2),x)
```

output

```

x*(A*a^3*c^2 - A*a^3*d^2 + B*b^3*c^2 - C*a^3*c^2 - B*b^3*d^2 + C*a^3*d^2 +
  2*A*b^3*c*d - 2*B*a^3*c*d - 2*C*b^3*c*d - 3*A*a*b^2*c^2 + 3*A*a*b^2*d^2 -
  3*B*a^2*b*c^2 + 3*B*a^2*b*d^2 + 3*C*a*b^2*c^2 - 3*C*a*b^2*d^2 - 6*A*a^2*b
*c*d + 6*B*a*b^2*c*d + 6*C*a^2*b*c*d) - (tan(e + f*x)*(B*b^3*c^2 - A*a^3*d
^2 - b^2*d*(B*b*d + 3*C*a*d + 2*C*b*c) - C*a^3*c^2 + C*a^3*d^2 + 2*A*b^3*c
*d - 2*B*a^3*c*d - 3*A*a*b^2*c^2 + 3*A*a*b^2*d^2 - 3*B*a^2*b*c^2 + 3*B*a^2
*b*d^2 + 3*C*a*b^2*c^2 - 6*A*a^2*b*c*d + 6*B*a*b^2*c*d + 6*C*a^2*b*c*d))/f
- (log(tan(e + f*x)^2 + 1)*((A*b^3*c^2)/2 - (B*a^3*c^2)/2 - (A*b^3*d^2)/2
+ (B*a^3*d^2)/2 - (C*b^3*c^2)/2 + (C*b^3*d^2)/2 - A*a^3*c*d - B*b^3*c*d +
C*a^3*c*d - (3*A*a^2*b*c^2)/2 + (3*A*a^2*b*d^2)/2 + (3*B*a*b^2*c^2)/2 - (
3*B*a*b^2*d^2)/2 + (3*C*a^2*b*c^2)/2 - (3*C*a^2*b*d^2)/2 + 3*A*a*b^2*c*d +
3*B*a^2*b*c*d - 3*C*a*b^2*c*d))/f + (tan(e + f*x)^4*((A*b^3*d^2)/4 + (C*b
^3*c^2)/4 - (C*b^3*d^2)/4 + (B*b^3*c*d)/2 + (3*B*a*b^2*d^2)/4 + (3*C*a^2*b
*d^2)/4 + (3*C*a*b^2*c*d)/2))/f + (tan(e + f*x)^3*((B*b^3*c^2)/3 - (b^2*d*
(B*b*d + 3*C*a*d + 2*C*b*c))/3 + (C*a^3*d^2)/3 + (2*A*b^3*c*d)/3 + A*a*b^2
*d^2 + B*a^2*b*d^2 + C*a*b^2*c^2 + 2*B*a*b^2*c*d + 2*C*a^2*b*c*d))/f + (ta
n(e + f*x)^2*((A*b^3*c^2)/2 - (A*b^3*d^2)/2 + (B*a^3*d^2)/2 - (C*b^3*c^2)/
2 + (C*b^3*d^2)/2 - B*b^3*c*d + C*a^3*c*d + (3*A*a^2*b*d^2)/2 + (3*B*a*b^2
*c^2)/2 - (3*B*a*b^2*d^2)/2 + (3*C*a^2*b*c^2)/2 - (3*C*a^2*b*d^2)/2 + 3*A*
a*b^2*c*d + 3*B*a^2*b*c*d - 3*C*a*b^2*c*d))/f + (b^2*d*tan(e + f*x)^5*(...

```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1122, normalized size of antiderivative = 1.70

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```

int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x)

```


output

```
(60*log(tan(e + f*x)**2 + 1)*a**4*c*d + 120*log(tan(e + f*x)**2 + 1)*a**3*
b*c**2 - 120*log(tan(e + f*x)**2 + 1)*a**3*b*d**2 - 60*log(tan(e + f*x)**2
+ 1)*a**3*c**2*d - 360*log(tan(e + f*x)**2 + 1)*a**2*b**2*c*d - 90*log(ta
n(e + f*x)**2 + 1)*a**2*b*c**3 + 90*log(tan(e + f*x)**2 + 1)*a**2*b*c*d**2
- 120*log(tan(e + f*x)**2 + 1)*a*b**3*c**2 + 120*log(tan(e + f*x)**2 + 1)
*a*b**3*d**2 + 180*log(tan(e + f*x)**2 + 1)*a*b**2*c**2*d + 60*log(tan(e +
f*x)**2 + 1)*b**4*c*d + 30*log(tan(e + f*x)**2 + 1)*b**3*c**3 - 30*log(ta
n(e + f*x)**2 + 1)*b**3*c*d**2 + 10*tan(e + f*x)**6*b**3*c*d**2 + 36*tan(e
+ f*x)**5*a*b**2*c*d**2 + 12*tan(e + f*x)**5*b**4*d**2 + 24*tan(e + f*x)*
*5*b**3*c**2*d + 45*tan(e + f*x)**4*a**2*b*c*d**2 + 60*tan(e + f*x)**4*a*b
**3*d**2 + 90*tan(e + f*x)**4*a*b**2*c**2*d + 30*tan(e + f*x)**4*b**4*c*d
+ 15*tan(e + f*x)**4*b**3*c**3 - 15*tan(e + f*x)**4*b**3*c*d**2 + 20*tan(e
+ f*x)**3*a**3*c*d**2 + 120*tan(e + f*x)**3*a**2*b**2*d**2 + 120*tan(e +
f*x)**3*a**2*b*c**2*d + 160*tan(e + f*x)**3*a*b**3*c*d + 60*tan(e + f*x)**
3*a*b**2*c**3 - 60*tan(e + f*x)**3*a*b**2*c*d**2 + 20*tan(e + f*x)**3*b**4
*c**2 - 20*tan(e + f*x)**3*b**4*d**2 - 40*tan(e + f*x)**3*b**3*c**2*d + 12
0*tan(e + f*x)**2*a**3*b*d**2 + 60*tan(e + f*x)**2*a**3*c**2*d + 360*tan(e
+ f*x)**2*a**2*b**2*c*d + 90*tan(e + f*x)**2*a**2*b*c**3 - 90*tan(e + f*x)
)**2*a**2*b*c*d**2 + 120*tan(e + f*x)**2*a*b**3*c**2 - 120*tan(e + f*x)**2
*a*b**3*d**2 - 180*tan(e + f*x)**2*a*b**2*c**2*d - 60*tan(e + f*x)**2*b...
```

3.58 $\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^2 (A + B \tan(e +$

Optimal result	629
Mathematica [C] (verified)	630
Rubi [A] (verified)	631
Maple [A] (warning: unable to verify)	635
Fricas [A] (verification not implemented)	636
Sympy [B] (verification not implemented)	637
Maxima [A] (verification not implemented)	638
Giac [B] (verification not implemented)	638
Mupad [B] (verification not implemented)	640
Reduce [B] (verification not implemented)	641

Optimal result

Integrand size = 45, antiderivative size = 443

$$\begin{aligned}
 & \int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) \, dx \\
 &= -((a^2(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) - b^2(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) \\
 &\quad + 2ab(2c(A - C)d + B(c^2 - d^2))) x \\
 &\quad + \frac{(2ab(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) - a^2(2c(A - C)d + B(c^2 - d^2)) + b^2(2c(A - C)d + B(c^2 - d^2)))}{f} \\
 &\quad + \frac{d(2ab(Ac - cC - Bd) + a^2(Bc + (A - C)d) - b^2(Bc + (A - C)d)) \tan(e + fx)}{f} \\
 &\quad + \frac{(a^2B - b^2B + 2ab(A - C))(c + d \tan(e + fx))^2}{2f} \\
 &\quad + \frac{(8a^2Cd^2 - 10abd(cC - 4Bd) + b^2(2c^2C - 5Bcd + 20(A - C)d^2))(c + d \tan(e + fx))^3}{60d^3f} \\
 &\quad - \frac{b(2bcC - 5bBd - 2aCd) \tan(e + fx)(c + d \tan(e + fx))^3}{20d^2f} \\
 &\quad + \frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df}
 \end{aligned}$$

output

```

-(a^2*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-b^2*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))
+2*a*b*(2*c*(A-C)*d+B*(c^2-d^2))*x+(2*a*b*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))
-a^2*(2*c*(A-C)*d+B*(c^2-d^2))+b^2*(2*c*(A-C)*d+B*(c^2-d^2))*ln(cos(f*x+e))
/f+d*(2*a*b*(A*c-B*d-C*c)+a^2*(B*c+(A-C)*d)-b^2*(B*c+(A-C)*d))*tan(f*x+e)
/f+1/2*(B*a^2-B*b^2+2*a*b*(A-C))*(c+d*tan(f*x+e))^2/f+1/60*(8*a^2*C*d^2-10*a*b*d*(-4*B*d+C*c)
+b^2*(2*c^2*C-5*B*c*d+20*(A-C)*d^2))*(c+d*tan(f*x+e))^3/d^3/f-1/20*b*(-5*B*b*d-2*C*a*d+2*C*b*c)
*tan(f*x+e)*(c+d*tan(f*x+e))^3/d^2/f+1/5*C*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3/d/f

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.40 (sec) , antiderivative size = 352, normalized size of antiderivative = 0.79

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(8a^2Cd^2 + 10abd(-cC + 4Bd) + b^2(2c^2C - 5Bcd + 20(A - C)d^2)) (c + d \tan(e + fx))^3 + 3bd(-2bcC - 3bd^2) \tan(e + fx) + 3bd^2 \tan^2(e + fx)}{60d^3f}$$

input

```

Integrate[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x]
+ C*Tan[e + f*x]^2),x]

```

output

```

((8*a^2*C*d^2 + 10*a*b*d*(-(c*C) + 4*B*d) + b^2*(2*c^2*C - 5*B*c*d + 20*(A
- C)*d^2))*(c + d*Tan[e + f*x])^3 + 3*b*d*(-2*b*c*C + 5*b*B*d + 2*a*C*d)*
Tan[e + f*x]*(c + d*Tan[e + f*x])^3 + 12*C*d^2*(a + b*Tan[e + f*x])^2*(c +
d*Tan[e + f*x])^3 + 30*d*(d*(2*a*b*(A*c - c*C + B*d) + a^2*(B*c + (-A + C)
*d) - b^2*(B*c + (-A + C)*d))*(I*((c + I*d)^2*Log[I - Tan[e + f*x]] - (c
- I*d)^2*Log[I + Tan[e + f*x]]) - 2*d^2*Tan[e + f*x]) + (a^2*B - b^2*B + 2
*a*b*(A - C))*d*((I*c - d)^3*Log[I - Tan[e + f*x]] - (I*c + d)^3*Log[I + T
an[e + f*x]] + 6*c*d^2*Tan[e + f*x] + d^3*Tan[e + f*x]^2))/(60*d^3*f)

```

Rubi [A] (verified)

Time = 3.64 (sec) , antiderivative size = 470, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.311$, Rules used = {3042, 4130, 25, 3042, 4120, 25, 3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4130

$$\frac{\int -((a + b \tan(e + fx))(c + d \tan(e + fx))^2 ((2bcC - 2adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3) dx}{5d}$$

↓ 25

$$\frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 ((2bcC - 2adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3) dx}{5df}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 ((2bcC - 2adC - 5bBd) \tan(e + fx)^2 - 5(Ab - Cb + aB)d \tan(e + fx) + C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3) dx}{5df}$$

↓ 4120

$$\frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 ((2bcC - 2adC - 5bBd) \tan(e + fx)^2 - 5(Ab - Cb + aB)d \tan(e + fx) + C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3) dx}{5df}$$

↓ 25

$$\frac{b \tan(e + fx)(-2aCd - 5bBd + 2bcC)(c + d \tan(e + fx))^3}{4df} - \frac{\int -(c + d \tan(e + fx))^2 (-c(2cC - 5Bd)b^2 + 10acCdb - 4a^2(5A - 3C)d^2 - ((2Cc^2 - 5Bdc + 5d^2)(a + b \tan(e + fx))) dx}{5d}$$

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{\int (c + d \tan(e + fx))^2 (-c(2cC - 5Bd)b^2 + 10acCdb - 4a^2(5A - 3C)d^2 - ((2Cc^2 - 5Bdc + 20(A - C)d^2)b^2 - 10ad(cC - 4Bd)b + 8a^2Cd^2) \tan^2(e + fx) - 2c^2d) dx}{4d}$$

5d

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{\int (c + d \tan(e + fx))^2 (-c(2cC - 5Bd)b^2 + 10acCdb - 4a^2(5A - 3C)d^2 - ((2Cc^2 - 5Bdc + 20(A - C)d^2)b^2 - 10ad(cC - 4Bd)b + 8a^2Cd^2) \tan(e + fx)^2 - 2c^2d) dx}{4d}$$

5d

↓ 4113

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{\int (c + d \tan(e + fx))^2 (20(-((A - C)a^2) + 2bBa + b^2(A - C))d^2 - 20(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)) dx - \frac{(c + d \tan(e + fx))^3 (8a^2Cd^2 - 10abd(cC - 4Bd) - 2c^2d)}{3c}}{4d}$$

5d

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{\int (c + d \tan(e + fx))^2 (20(-((A - C)a^2) + 2bBa + b^2(A - C))d^2 - 20(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)) dx - \frac{(c + d \tan(e + fx))^3 (8a^2Cd^2 - 10abd(cC - 4Bd) - 2c^2d)}{3c}}{4d}$$

5d

↓ 4011

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{\int (c + d \tan(e + fx)) (-20((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd))d^2 - 20((Bc + (A - C)d)a^2 + 2b(Ac - Cc - Bd)a - b^2(Bc + (A - C)d))d^2 \tan(e + fx)) dx}{4d}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{\int (c + d \tan(e + fx)) (-20((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd))d^2 - 20((Bc + (A - C)d)a^2 + 2b(Ac - Cc - Bd)a - b^2(Bc + (A - C)d))d^2 \tan(e + fx)) dx}{4d}$$

↓ 4008

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{20d^2(-a^2(2cd(A-C)+B(c^2-d^2))+2ab(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)+b^2(2cd(A-C)+B(c^2-d^2))) \int \tan(e+fx)dx - \frac{(c+d \tan(e+fx))^3(8a^2}{f}}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{20d^2(-a^2(2cd(A-C)+B(c^2-d^2))+2ab(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)+b^2(2cd(A-C)+B(c^2-d^2))) \int \tan(e+fx)dx - \frac{(c+d \tan(e+fx))^3(8a^2}{f}}$$

↓ 3956

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3}{5df} - \frac{(c+d \tan(e+fx))^3(8a^2Cd^2-10abd(cC-4Bd)+b^2(20d^2(A-C)-5Bcd+2c^2C))}{3df} - \frac{20d^2 \log(\cos(e+fx))(-a^2(2cd(A-C)+B(c^2-d^2))+2ab(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)+b^2(2cd(A-C)+B(c^2-d^2)))}{f}$$

input

```
Int[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
(C*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^3)/(5*d*f) - ((b*(2*b*c*C - 5*b*B*d - 2*a*C*d)*Tan[e + f*x]*(c + d*Tan[e + f*x])^3)/(4*d*f) + (20*d^2*(a^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + 2*a*b*(2*c*(A - C)*d + B*(c^2 - d^2)))*x - (20*d^2*(2*a*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - a^2*(2*c*(A - C)*d + B*(c^2 - d^2)) + b^2*(2*c*(A - C)*d + B*(c^2 - d^2)))*Log[Cos[e + f*x]]/f - (20*d^3*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d) - b^2*(B*c + (A - C)*d))*Tan[e + f*x])/f - (10*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2*(c + d*Tan[e + f*x])^2)/f - ((8*a^2*C*d^2 - 10*a*b*d*(c*C - 4*B*d) + b^2*(2*c^2*C - 5*B*c*d + 20*(A - C)*d^2))*(c + d*Tan[e + f*x])^3)/(3*d*f)/(4*d)/(5*d)
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] :> Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (warning: unable to verify)

Time = 0.28 (sec) , antiderivative size = 392, normalized size of antiderivative = 0.88

method	result
parts	$\frac{(2A^2cd + 2Aabc^2 + B^2a^2c^2) \ln(1 + \tan^2(fx + e))}{2f} + \frac{(B^2b^2d^2 + 2Cab^2d^2 + 2Cb^2cd)}{f} \left(\frac{\tan^4(fx + e)}{4} - \frac{\tan^2(fx + e)}{2} + \frac{\ln(1 + \tan^2(fx + e))}{2} \right)$
norman	$(A^2a^2c^2 - A^2a^2d^2 - 4Aabcd - Ab^2c^2 + Ab^2d^2 - 2Ba^2cd - 2Babc^2 + 2Babd^2 + 2Bb^2cd)$
derivativedivides	$\frac{(2A^2cd + 2Aabc^2 - 2Aabd^2 - 2Ab^2cd + B^2a^2c^2 - B^2a^2d^2 - 4Aabcd - B^2b^2c^2 + B^2b^2d^2 - 2Ca^2cd - 2Cab^2c^2 + 2Cab^2d^2 + 2Cb^2cd) \ln(1 + \tan^2(fx + e))}{2}$
default	$\frac{(2A^2cd + 2Aabc^2 - 2Aabd^2 - 2Ab^2cd + B^2a^2c^2 - B^2a^2d^2 - 4Aabcd - B^2b^2c^2 + B^2b^2d^2 - 2Ca^2cd - 2Cab^2c^2 + 2Cab^2d^2 + 2Cb^2cd) \ln(1 + \tan^2(fx + e))}{2}$
parallelrisch	$-240 \tan(fx + e) Cabcd + 120 Babcd \tan^2(fx + e) + 80 Cabcd \tan^3(fx + e) + 240 \tan(fx + e) Aabcd + 20C a^2 d^2 \tan^3(fx + e)$
risch	Expression too large to display

input `int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2} * (2 * A * a^2 * c * d + 2 * A * a * b * c^2 + B * a^2 * c^2) / f * \ln(1 + \tan(f * x + e)^2) + (B * b^2 * d^2 + 2 * C * a * b * d^2 + 2 * C * b^2 * c * d) / f * (1/4 * \tan(f * x + e)^4 - 1/2 * \tan(f * x + e)^2 + 1/2 * \ln(1 + \tan(f * x + e)^2)) + (A * b^2 * d^2 + 2 * B * a * b * d^2 + 2 * B * b^2 * c * d + C * a^2 * d^2 + 4 * C * a * b * c * d + C * b^2 * c^2) / f * (1/3 * \tan(f * x + e)^3 - \tan(f * x + e) + \arctan(\tan(f * x + e))) + (A * a^2 * d^2 + 4 * A * a * b * c * d + A * b^2 * c^2 + 2 * B * a^2 * c * d + 2 * B * a * b * c^2 + C * a^2 * c^2) / f * (\tan(f * x + e) - \arctan(\tan(f * x + e))) + (2 * A * a * b * d^2 + 2 * A * b^2 * c * d + B * a^2 * d^2 + 4 * B * a * b * c * d + B * b^2 * c^2 + 2 * C * a^2 * c * d + 2 * C * a * b * c^2) / f * (1/2 * \tan(f * x + e)^2 - 1/2 * \ln(1 + \tan(f * x + e)^2)) + A * a^2 * c^2 * x + b^2 * d^2 * C / f * (1/5 * \tan(f * x + e)^5 - 1/3 * \tan(f * x + e)^3 + \tan(f * x + e) - \arctan(\tan(f * x + e)))$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 462, normalized size of antiderivative = 1.04

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12 C b^2 d^2 \tan(fx + e)^5 + 15 (2 C b^2 c d + (2 C a b + B b^2) d^2) \tan(fx + e)^4 + 20 (C b^2 c^2 + 2 (2 C a b + B b^2) c d + (A - C) b^2 d^2) \tan(fx + e)^3 + 60 ((A - C) a^2 - 2 B a b - (A - C) b^2) c^2 - 2 (B a^2 + 2 (A - C) a b - B b^2) c d - ((A - C) a^2 - 2 B a b - (A - C) b^2) d^2 * f * x + 30 ((2 C a b + B b^2) c^2 + 2 (C a^2 + 2 B a b + (A - C) b^2) c d + (B a^2 + 2 (A - C) a b - B b^2) d^2) * \tan(fx + e)^2 - 30 ((B a^2 + 2 (A - C) a b - B b^2) c^2 + 2 ((A - C) a^2 - 2 B a b - (A - C) b^2) c d - (B a^2 + 2 (A - C) a b - B b^2) d^2) * \log(1 / (\tan(fx + e)^2 + 1)) + 60 ((C a^2 + 2 B a b + (A - C) b^2) c^2 + 2 (B a^2 + 2 (A - C) a b - B b^2) c d + ((A - C) a^2 - 2 B a b - (A - C) b^2) d^2) * \tan(fx + e)}{f}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output
$$\frac{1}{60} * (12 * C * b^2 * d^2 * \tan(f * x + e)^5 + 15 * (2 * C * b^2 * c * d + (2 * C * a * b + B * b^2) * d^2) * \tan(f * x + e)^4 + 20 * (C * b^2 * c^2 + 2 * (2 * C * a * b + B * b^2) * c * d + (C * a^2 + 2 * B * a * b + (A - C) * b^2) * d^2) * \tan(f * x + e)^3 + 60 * (((A - C) * a^2 - 2 * B * a * b - (A - C) * b^2) * c^2 - 2 * (B * a^2 + 2 * (A - C) * a * b - B * b^2) * c * d - ((A - C) * a^2 - 2 * B * a * b - (A - C) * b^2) * d^2) * f * x + 30 * ((2 * C * a * b + B * b^2) * c^2 + 2 * (C * a^2 + 2 * B * a * b + (A - C) * b^2) * c * d + (B * a^2 + 2 * (A - C) * a * b - B * b^2) * d^2) * \tan(f * x + e)^2 - 30 * ((B * a^2 + 2 * (A - C) * a * b - B * b^2) * c^2 + 2 * ((A - C) * a^2 - 2 * B * a * b - (A - C) * b^2) * c * d - (B * a^2 + 2 * (A - C) * a * b - B * b^2) * d^2) * \log(1 / (\tan(f * x + e)^2 + 1)) + 60 * ((C * a^2 + 2 * B * a * b + (A - C) * b^2) * c^2 + 2 * (B * a^2 + 2 * (A - C) * a * b - B * b^2) * c * d + ((A - C) * a^2 - 2 * B * a * b - (A - C) * b^2) * d^2) * \tan(f * x + e)) / f$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1134 vs. $2(396) = 792$.

Time = 0.29 (sec) , antiderivative size = 1134, normalized size of antiderivative = 2.56

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))**2*(c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A**2*c**2*x + A**2*c*d*log(tan(e + f*x)**2 + 1)/f - A**2*d**2*x + A**2*d**2*tan(e + f*x)/f + A*b*c**2*log(tan(e + f*x)**2 + 1)/f - 4*A*b*c*d*x + 4*A*b*c*d*tan(e + f*x)/f - A*b*d**2*log(tan(e + f*x)**2 + 1)/f + A*b*d**2*tan(e + f*x)**2/f - A*b**2*c**2*x + A*b**2*c**2*tan(e + f*x)/f - A*b**2*c*d*log(tan(e + f*x)**2 + 1)/f + A*b**2*c*d*tan(e + f*x)**2/f + A*b**2*d**2*x + A*b**2*d**2*tan(e + f*x)**3/(3*f) - A*b**2*d**2*tan(e + f*x)/f + B*a**2*c**2*log(tan(e + f*x)**2 + 1)/(2*f) - 2*B*a**2*c*d*x + 2*B*a**2*c*d*tan(e + f*x)/f - B*a**2*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + B*a**2*d**2*tan(e + f*x)**2/(2*f) - 2*B*a*b*c**2*x + 2*B*a*b*c**2*tan(e + f*x)/f - 2*B*a*b*c*d*log(tan(e + f*x)**2 + 1)/f + 2*B*a*b*c*d*tan(e + f*x)**2/f + 2*B*a*b*d**2*x + 2*B*a*b*d**2*tan(e + f*x)**3/(3*f) - 2*B*a*b*d**2*tan(e + f*x)/f - B*b**2*c**2*log(tan(e + f*x)**2 + 1)/(2*f) + B*b**2*c**2*tan(e + f*x)**2/(2*f) + 2*B*b**2*c*d*x + 2*B*b**2*c*d*tan(e + f*x)**3/(3*f) - 2*B*b**2*c*d*tan(e + f*x)/f + B*b**2*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + B*b**2*d**2*tan(e + f*x)**4/(4*f) - B*b**2*d**2*tan(e + f*x)**2/(2*f) - C*a**2*c**2*x + C*a**2*c**2*tan(e + f*x)/f - C*a**2*c*d*log(tan(e + f*x)**2 + 1)/f + C*a**2*c*d*tan(e + f*x)**2/f + C*a**2*d**2*x + C*a**2*d**2*tan(e + f*x)**3/(3*f) - C*a**2*d**2*tan(e + f*x)/f - C*a*b*c**2*log(tan(e + f*x)**2 + 1)/f + C*a*b*c**2*tan(e + f*x)**2/f + 4*C*a*b*c*d*x + 4*C*a*b*c*d*tan(e + f*x)**3/(3*f) - 4*C*a*b*c*d*tan(e + f*x)/f + C*a...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 463, normalized size of antiderivative = 1.05

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12 C b^2 d^2 \tan(fx + e)^5 + 15 (2 C b^2 c d + (2 C a b + B b^2) d^2) \tan(fx + e)^4 + 20 (C b^2 c^2 + 2 (2 C a b + B b^2) c d + (A - C) b^2 d^2) \tan(fx + e)^3 + 30 ((2 C a b + B b^2) c^2 + 2 (C a^2 + 2 B a b + (A - C) b^2) c d + (B a^2 + 2 (A - C) a b - B b^2) d^2) \tan(fx + e)^2 + 60 (((A - C) a^2 - 2 B a b - (A - C) b^2) c^2 - 2 (B a^2 + 2 (A - C) a b - B b^2) c d - ((A - C) a^2 - 2 B a b - (A - C) b^2) d^2) \tan(fx + e) + 30 ((B a^2 + 2 (A - C) a b - B b^2) c^2 + 2 ((A - C) a^2 - 2 B a b - (A - C) b^2) c d - (B a^2 + 2 (A - C) a b - B b^2) d^2) \log(\tan(fx + e)^2 + 1) + 60 ((C a^2 + 2 B a b + (A - C) b^2) c^2 + 2 (B a^2 + 2 (A - C) a b - B b^2) c d + ((A - C) a^2 - 2 B a b - (A - C) b^2) d^2) \tan(fx + e)}{f}$$

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
1/60*(12*C*b^2*d^2*tan(f*x + e)^5 + 15*(2*C*b^2*c*d + (2*C*a*b + B*b^2)*d^2)*tan(f*x + e)^4 + 20*(C*b^2*c^2 + 2*(2*C*a*b + B*b^2)*c*d + (C*a^2 + 2*B*a*b + (A - C)*b^2)*d^2)*tan(f*x + e)^3 + 30*((2*C*a*b + B*b^2)*c^2 + 2*(C*a^2 + 2*B*a*b + (A - C)*b^2)*c*d + (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^2)*tan(f*x + e)^2 + 60*(((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^2 - 2*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d - ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^2)*(f*x + e) + 30*((B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2 + 2*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d - (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^2)*log(tan(f*x + e)^2 + 1) + 60*((C*a^2 + 2*B*a*b + (A - C)*b^2)*c^2 + 2*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d + ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^2)*tan(f*x + e))/f
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 885 vs. 2(436) = 872.

Time = 0.95 (sec) , antiderivative size = 885, normalized size of antiderivative = 2.00

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
(A*a^2*c^2 - C*a^2*c^2 - 2*B*a*b*c^2 - A*b^2*c^2 + C*b^2*c^2 - 2*B*a^2*c*d
- 4*A*a*b*c*d + 4*C*a*b*c*d + 2*B*b^2*c*d - A*a^2*d^2 + C*a^2*d^2 + 2*B*a
*b*d^2 + A*b^2*d^2 - C*b^2*d^2)*(f*x + e)/f + 1/2*(B*a^2*c^2 + 2*A*a*b*c^2
- 2*C*a*b*c^2 - B*b^2*c^2 + 2*A*a^2*c*d - 2*C*a^2*c*d - 4*B*a*b*c*d - 2*A
*b^2*c*d + 2*C*b^2*c*d - B*a^2*d^2 - 2*A*a*b*d^2 + 2*C*a*b*d^2 + B*b^2*d^2
)*log(tan(f*x + e)^2 + 1)/f + 1/60*(12*C*b^2*d^2*f^4*tan(f*x + e)^5 + 30*C
*b^2*c*d*f^4*tan(f*x + e)^4 + 30*C*a*b*d^2*f^4*tan(f*x + e)^4 + 15*B*b^2*d
^2*f^4*tan(f*x + e)^4 + 20*C*b^2*c^2*f^4*tan(f*x + e)^3 + 80*C*a*b*c*d*f^4
*tan(f*x + e)^3 + 40*B*b^2*c*d*f^4*tan(f*x + e)^3 + 20*C*a^2*d^2*f^4*tan(f
*x + e)^3 + 40*B*a*b*d^2*f^4*tan(f*x + e)^3 + 20*A*b^2*d^2*f^4*tan(f*x +
e)^3 - 20*C*b^2*d^2*f^4*tan(f*x + e)^3 + 60*C*a*b*c^2*f^4*tan(f*x + e)^2 +
30*B*b^2*c^2*f^4*tan(f*x + e)^2 + 60*C*a^2*c*d*f^4*tan(f*x + e)^2 + 120*B*
a*b*c*d*f^4*tan(f*x + e)^2 + 60*A*b^2*c*d*f^4*tan(f*x + e)^2 - 60*C*b^2*c*
d*f^4*tan(f*x + e)^2 + 30*B*a^2*d^2*f^4*tan(f*x + e)^2 + 60*A*a*b*d^2*f^4*
tan(f*x + e)^2 - 60*C*a*b*d^2*f^4*tan(f*x + e)^2 - 30*B*b^2*d^2*f^4*tan(f*
x + e)^2 + 60*C*a^2*c^2*f^4*tan(f*x + e) + 120*B*a*b*c^2*f^4*tan(f*x + e)
+ 60*A*b^2*c^2*f^4*tan(f*x + e) - 60*C*b^2*c^2*f^4*tan(f*x + e) + 120*B*a^
2*c*d*f^4*tan(f*x + e) + 240*A*a*b*c*d*f^4*tan(f*x + e) - 240*C*a*b*c*d*f^
4*tan(f*x + e) - 120*B*b^2*c*d*f^4*tan(f*x + e) + 60*A*a^2*d^2*f^4*tan(f*x
+ e) - 60*C*a^2*d^2*f^4*tan(f*x + e) - 120*B*a*b*d^2*f^4*tan(f*x + e) ...
```

Mupad [B] (verification not implemented)

Time = 5.81 (sec) , antiderivative size = 561, normalized size of antiderivative = 1.27

$$\begin{aligned}
& \int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= x (A a^2 c^2 - A a^2 d^2 - A b^2 c^2 + A b^2 d^2 - C a^2 c^2 + C a^2 d^2 + C b^2 c^2 - C b^2 d^2 \\
&\quad - 2 B a b c^2 + 2 B a b d^2 - 2 B a^2 c d + 2 B b^2 c d - 4 A a b c d + 4 C a b c d) \\
&\quad \ln(\tan(e + fx)^2 + 1) \left(\frac{B a^2 d^2}{2} - \frac{B a^2 c^2}{2} + \frac{B b^2 c^2}{2} - \frac{B b^2 d^2}{2} - A a b c^2 + A a b d^2 - A a^2 c d + C a b c^2 + A \right. \\
&\quad \left. - \frac{\tan(e + fx)^2 \left(\frac{B a^2 d^2}{2} + \frac{B b^2 c^2}{2} - \frac{b d (B b d + 2 C a d + 2 C b c)}{2} + A a b d^2 + C a b c^2 + A b^2 c d + C a^2 c d + 2 B a \right)}{f} \right. \\
&\quad \left. + \frac{\tan(e + fx)^3 \left(\frac{A b^2 d^2}{3} + \frac{C a^2 d^2}{3} + \frac{C b^2 c^2}{3} - \frac{C b^2 d^2}{3} + \frac{2 B a b d^2}{3} + \frac{2 B b^2 c d}{3} + \frac{4 C a b c d}{3} \right)}{f} \right. \\
&\quad \left. + \frac{\tan(e + fx) (A a^2 d^2 + A b^2 c^2 - A b^2 d^2 + C a^2 c^2 - C a^2 d^2 - C b^2 c^2 + C b^2 d^2 + 2 B a b c^2 - 2 B a b d^2)}{f} \right. \\
&\quad \left. + \frac{b d \tan(e + fx)^4 (B b d + 2 C a d + 2 C b c)}{4 f} + \frac{C b^2 d^2 \tan(e + fx)^5}{5 f} \right)
\end{aligned}$$

input

```
int((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*
tan(e + f*x)^2),x)
```

output

```
x*(A*a^2*c^2 - A*a^2*d^2 - A*b^2*c^2 + A*b^2*d^2 - C*a^2*c^2 + C*a^2*d^2 +
C*b^2*c^2 - C*b^2*d^2 - 2*B*a*b*c^2 + 2*B*a*b*d^2 - 2*B*a^2*c*d + 2*B*b^2
*c*d - 4*A*a*b*c*d + 4*C*a*b*c*d) - (log(tan(e + f*x)^2 + 1)*((B*a^2*d^2)/
2 - (B*a^2*c^2)/2 + (B*b^2*c^2)/2 - (B*b^2*d^2)/2 - A*a*b*c^2 + A*a*b*d^2
- A*a^2*c*d + C*a*b*c^2 + A*b^2*c*d - C*a*b*d^2 + C*a^2*c*d - C*b^2*c*d +
2*B*a*b*c*d))/f + (tan(e + f*x)^2*((B*a^2*d^2)/2 + (B*b^2*c^2)/2 - (b*d*(B
*b*d + 2*C*a*d + 2*C*b*c))/2 + A*a*b*d^2 + C*a*b*c^2 + A*b^2*c*d + C*a^2*c
*d + 2*B*a*b*c*d))/f + (tan(e + f*x)^3*((A*b^2*d^2)/3 + (C*a^2*d^2)/3 + (C
*b^2*c^2)/3 - (C*b^2*d^2)/3 + (2*B*a*b*d^2)/3 + (2*B*b^2*c*d)/3 + (4*C*a*b
*c*d)/3))/f + (tan(e + f*x)*(A*a^2*d^2 + A*b^2*c^2 - A*b^2*d^2 + C*a^2*c^2
- C*a^2*d^2 - C*b^2*c^2 + C*b^2*d^2 + 2*B*a*b*c^2 - 2*B*a*b*d^2 + 2*B*a^2
*c*d - 2*B*b^2*c*d + 4*A*a*b*c*d - 4*C*a*b*c*d))/f + (b*d*tan(e + f*x)^4*(
B*b*d + 2*C*a*d + 2*C*b*c))/(4*f) + (C*b^2*d^2*tan(e + f*x)^5)/(5*f)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 748, normalized size of antiderivative = 1.69

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x)
```

output

```
(60*log(tan(e + f*x)**2 + 1)*a**3*c*d + 90*log(tan(e + f*x)**2 + 1)*a**2*b
*c**2 - 90*log(tan(e + f*x)**2 + 1)*a**2*b*d**2 - 60*log(tan(e + f*x)**2 +
1)*a**2*c**2*d - 180*log(tan(e + f*x)**2 + 1)*a*b**2*c*d - 60*log(tan(e +
f*x)**2 + 1)*a*b*c**3 + 60*log(tan(e + f*x)**2 + 1)*a*b*c*d**2 - 30*log(t
an(e + f*x)**2 + 1)*b**3*c**2 + 30*log(tan(e + f*x)**2 + 1)*b**3*d**2 + 60
*log(tan(e + f*x)**2 + 1)*b**2*c**2*d + 12*tan(e + f*x)**5*b**2*c*d**2 + 3
0*tan(e + f*x)**4*a*b*c*d**2 + 15*tan(e + f*x)**4*b**3*d**2 + 30*tan(e + f
*x)**4*b**2*c**2*d + 20*tan(e + f*x)**3*a**2*c*d**2 + 60*tan(e + f*x)**3*a
*b**2*d**2 + 80*tan(e + f*x)**3*a*b*c**2*d + 40*tan(e + f*x)**3*b**3*c*d +
20*tan(e + f*x)**3*b**2*c**3 - 20*tan(e + f*x)**3*b**2*c*d**2 + 90*tan(e
+ f*x)**2*a**2*b*d**2 + 60*tan(e + f*x)**2*a**2*c**2*d + 180*tan(e + f*x)*
**2*a*b**2*c*d + 60*tan(e + f*x)**2*a*b*c**3 - 60*tan(e + f*x)**2*a*b*c*d**
2 + 30*tan(e + f*x)**2*b**3*c**2 - 30*tan(e + f*x)**2*b**3*d**2 - 60*tan(e
+ f*x)**2*b**2*c**2*d + 60*tan(e + f*x)*a**3*d**2 + 360*tan(e + f*x)*a**2
*b*c*d + 60*tan(e + f*x)*a**2*c**3 - 60*tan(e + f*x)*a**2*c*d**2 + 180*tan
(e + f*x)*a*b**2*c**2 - 180*tan(e + f*x)*a*b**2*d**2 - 240*tan(e + f*x)*a*
b*c**2*d - 120*tan(e + f*x)*b**3*c*d - 60*tan(e + f*x)*b**2*c**3 + 60*tan(
e + f*x)*b**2*c*d**2 + 60*a**3*c**2*f*x - 60*a**3*d**2*f*x - 360*a**2*b*c*
d*f*x - 60*a**2*c**3*f*x + 60*a**2*c*d**2*f*x - 180*a*b**2*c**2*f*x + 180*
a*b**2*d**2*f*x + 240*a*b*c**2*d*f*x + 120*b**3*c*d*f*x + 60*b**2*c**3*...
```

3.59 $\int (a+b \tan(e+fx))(c+d \tan(e+fx))^2 (A + B \tan(e +$

Optimal result	642
Mathematica [C] (verified)	643
Rubi [A] (verified)	643
Maple [A] (warning: unable to verify)	646
Fricas [A] (verification not implemented)	647
Sympy [B] (verification not implemented)	648
Maxima [A] (verification not implemented)	649
Giac [A] (verification not implemented)	649
Mupad [B] (verification not implemented)	650
Reduce [B] (verification not implemented)	651

Optimal result

Integrand size = 43, antiderivative size = 266

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= -((a(c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) + b(2c(A - C)d + B(c^2 - d^2))) x$$

$$- \frac{(a(Bc^2 - 2cCd - Bd^2) - b(c^2C + 2Bcd - Cd^2) + A(2acd + b(c^2 - d^2))) \log(\cos(e + fx))}{f}$$

$$+ \frac{d(abc + aBc - bcC + aAd - bBd - aCd) \tan(e + fx)}{f}$$

$$+ \frac{(Ab + aB - bC)(c + d \tan(e + fx))^2}{2f}$$

$$- \frac{(bcC - 4bBd - 4aCd)(c + d \tan(e + fx))^3}{12d^2 f} + \frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df}$$

output

```
- (a*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))+b*(2*c*(A-C)*d+B*(c^2-d^2)))*x-(a*(B*c^2-B*d^2-2*C*c*d)-b*(2*B*c*d+C*c^2-C*d^2)+A*(2*a*c*d+b*(c^2-d^2)))*ln(cos(f*x+e))/f+d*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*tan(f*x+e)/f+1/2*(A*b+B*a-C*b)*(c+d*tan(f*x+e))^2/f-1/12*(-4*B*b*d-4*C*a*d+C*b*c)*(c+d*tan(f*x+e))^3/d^2/f+1/4*b*C*tan(f*x+e)*(c+d*tan(f*x+e))^3/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.89 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.91

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(-bcC + 4bBd + 4aCd)(c + d \tan(e + fx))^3}{d} + 3bC \tan(e + fx)(c + d \tan(e + fx))^3 + 6(ABC + aBc - bcC - aAd + b$$

input

```
Integrate[(a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x]
+ C*Tan[e + f*x]^2),x]
```

output

```
(((-(b*c*C) + 4*b*B*d + 4*a*C*d)*(c + d*Tan[e + f*x])^3)/d + 3*b*C*Tan[e +
f*x]*(c + d*Tan[e + f*x])^3 + 6*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d +
a*C*d)*(I*((c + I*d)^2*Log[I - Tan[e + f*x]] - (c - I*d)^2*Log[I + Tan[e +
f*x]]) - 2*d^2*Tan[e + f*x]) + 6*(A*b + a*B - b*C)*((I*c - d)^3*Log[I - T
an[e + f*x]] - (I*c + d)^3*Log[I + Tan[e + f*x]] + 6*c*d^2*Tan[e + f*x] +
d^3*Tan[e + f*x]^2))/(12*d*f)
```

Rubi [A] (verified)

Time = 1.85 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {3042, 4120, 3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4120

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \int (c + d \tan(e + fx))^2 ((bcC - 4adC - 4bBd) \tan^2(e + fx) - 4(Ab - Cb + aB)d \tan(e + fx) + bcC - 4aAd) dx}{4d}$$

↓ 3042

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \int (c + d \tan(e + fx))^2 ((bcC - 4adC - 4bBd) \tan(e + fx)^2 - 4(Ab - Cb + aB)d \tan(e + fx) + bcC - 4aAd) dx}{4d}$$

↓ 4113

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \int (c + d \tan(e + fx))^2 (4(bB - a(A - C))d - 4(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{(-4aCd - 4bBd + bcC)(c + d \tan(e + fx))}{3df}}{4d}$$

↓ 3042

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \int (c + d \tan(e + fx))^2 (4(bB - a(A - C))d - 4(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{(-4aCd - 4bBd + bcC)(c + d \tan(e + fx))}{3df}}{4d}$$

↓ 4011

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \int (c + d \tan(e + fx))(4d(bBc + b(A - C)d - a(Ac - Cc - Bd)) - 4d(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx)) dx}{4d}$$

↓ 3042

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \int (c + d \tan(e + fx))(4d(bBc + b(A - C)d - a(Ac - Cc - Bd)) - 4d(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx)) dx}{4d}$$

↓ 4008

$$\frac{\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - 4d(2aAcd + aB(c^2 - d^2) - 2acCd + Ab(c^2 - d^2) - b(2Bcd + c^2C - Cd^2)) \int \tan(e + fx) dx + 4dx(a(-A(c^2 - d^2) + bC) \tan(e + fx) + bcC - 4aAd)}{4d}$$

↓ 3042

$$\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \frac{-4d(2aAcd + aB(c^2 - d^2) - 2acCd + Ab(c^2 - d^2) - b(2Bcd + c^2C - Cd^2)) \int \tan(e + fx) dx + 4dx(a(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2))}{4df}$$

↓ 3956

$$\frac{bC \tan(e + fx)(c + d \tan(e + fx))^3}{4df} - \frac{4d \log(\cos(e + fx))(2aAcd + aB(c^2 - d^2) - 2acCd + Ab(c^2 - d^2) - b(2Bcd + c^2C - Cd^2))}{f} + 4dx(a(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2))$$

input

```
Int[(a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
(b*C*Tan[e + f*x]*(c + d*Tan[e + f*x])^3)/(4*d*f) - (4*d*(a*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + b*(2*c*(A - C)*d + B*(c^2 - d^2)))*x + (4*d*(2*a*A*c*d - 2*a*c*C*d + A*b*(c^2 - d^2) + a*B*(c^2 - d^2) - b*(c^2*C + 2*B*c*d - C*d^2))*Log[Cos[e + f*x]])/f - (4*d^2*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d)*Tan[e + f*x])/f - (2*(A*b + a*B - b*C)*d*(c + d*Tan[e + f*x])^2)/f + ((b*c*C - 4*b*B*d - 4*a*C*d)*(c + d*Tan[e + f*x])^3)/(3*d*f))/ (4*d)
```

Defintions of rubi rules used

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4008

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

rule 4011

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

Maple [A] (warning: unable to verify)

Time = 0.19 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.92

method	result
parts	$\frac{(2Aacd+Abc^2+Ba c^2) \ln(1+\tan(fx+e))}{2f} + \frac{(Bbd^2+Ca d^2+2Cbcd) \left(\frac{\tan(fx+e)^3}{3} - \tan(fx+e) + \arctan(\tan(fx+e)) \right)}{f}$
norman	$(Aa c^2 - Aa d^2 - 2Abcd - 2Bacd - Bb c^2 + Bb d^2 - Ca c^2 + Ca d^2 + 2Cbcd) x + \frac{(Aa c^2 - Aa d^2 - 2Abcd - 2Bacd - Bb c^2 + Bb d^2 - Ca c^2 + Ca d^2 + 2Cbcd) \tan(fx+e)}{2}$
derivativedivides	$\frac{Cb d^2 \tan(fx+e)^4}{4} + \frac{Bb d^2 \tan(fx+e)^3}{3} + \frac{Ca d^2 \tan(fx+e)^3}{3} + \frac{2Cbcd \tan(fx+e)^3}{3} + \frac{Ab d^2 \tan(fx+e)^2}{2} + \frac{Ba d^2 \tan(fx+e)^2}{2} + Bbcd$
default	$\frac{Cb d^2 \tan(fx+e)^4}{4} + \frac{Bb d^2 \tan(fx+e)^3}{3} + \frac{Ca d^2 \tan(fx+e)^3}{3} + \frac{2Cbcd \tan(fx+e)^3}{3} + \frac{Ab d^2 \tan(fx+e)^2}{2} + \frac{Ba d^2 \tan(fx+e)^2}{2} + Bbcd$
parallelrisch	$3Cb d^2 \tan(fx+e)^4 + 4Bb d^2 \tan(fx+e)^3 + 4Ca d^2 \tan(fx+e)^3 + 6Ab d^2 \tan(fx+e)^2 + 6Ba d^2 \tan(fx+e)^2 + 6Cb c^2 \tan(fx+e)$
risch	Expression too large to display

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,
method=_RETURNVERBOSE)
```

output

```
1/2*(2*A*a*c*d+A*b*c^2+B*a*c^2)/f*ln(1+tan(f*x+e)^2)+(B*b*d^2+C*a*d^2+2*C*
b*c*d)/f*(1/3*tan(f*x+e)^3-tan(f*x+e)+arctan(tan(f*x+e)))+(A*b*d^2+B*a*d^2
+2*B*b*c*d+2*C*a*c*d+C*b*c^2)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)^2))+
(A*a*d^2+2*A*b*c*d+2*B*a*c*d+B*b*c^2+C*a*c^2)/f*(tan(f*x+e)-arctan(tan(f*x
+e)))+A*a*c^2*x+C*b*d^2/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(
f*x+e)^2))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.97

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3Cb d^2 \tan(fx + e)^4 + 4(2Cbcd + (Ca + Bb)d^2) \tan(fx + e)^3 + 12(((A - C)a - Bb)c^2 - 2(Ba + (A$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x, algorithm="fricas")
```

output

```
1/12*(3*C*b*d^2*tan(f*x + e)^4 + 4*(2*C*b*c*d + (C*a + B*b)*d^2)*tan(f*x +
e)^3 + 12*(((A - C)*a - B*b)*c^2 - 2*(B*a + (A - C)*b)*c*d - ((A - C)*a -
B*b)*d^2)*f*x + 6*(C*b*c^2 + 2*(C*a + B*b)*c*d + (B*a + (A - C)*b)*d^2)*t
an(f*x + e)^2 - 6*((B*a + (A - C)*b)*c^2 + 2*((A - C)*a - B*b)*c*d - (B*a
+ (A - C)*b)*d^2)*log(1/(tan(f*x + e)^2 + 1)) + 12*((C*a + B*b)*c^2 + 2*(B
*a + (A - C)*b)*c*d + ((A - C)*a - B*b)*d^2)*tan(f*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 617 vs. $2(246) = 492$.

Time = 0.21 (sec) , antiderivative size = 617, normalized size of antiderivative = 2.32

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)
)**2),x)
```

output

```
Piecewise(((A*a*c**2*x + A*a*c*d*log(tan(e + f*x)**2 + 1)/f - A*a*d**2*x +
A*a*d**2*tan(e + f*x)/f + A*b*c**2*log(tan(e + f*x)**2 + 1)/(2*f) - 2*A*b*
c*d*x + 2*A*b*c*d*tan(e + f*x)/f - A*b*d**2*log(tan(e + f*x)**2 + 1)/(2*f)
+ A*b*d**2*tan(e + f*x)**2/(2*f) + B*a*c**2*log(tan(e + f*x)**2 + 1)/(2*f)
) - 2*B*a*c*d*x + 2*B*a*c*d*tan(e + f*x)/f - B*a*d**2*log(tan(e + f*x)**2
+ 1)/(2*f) + B*a*d**2*tan(e + f*x)**2/(2*f) - B*b*c**2*x + B*b*c**2*tan(e
+ f*x)/f - B*b*c*d*log(tan(e + f*x)**2 + 1)/f + B*b*c*d*tan(e + f*x)**2/f
+ B*b*d**2*x + B*b*d**2*tan(e + f*x)**3/(3*f) - B*b*d**2*tan(e + f*x)/f -
C*a*c**2*x + C*a*c**2*tan(e + f*x)/f - C*a*c*d*log(tan(e + f*x)**2 + 1)/f
+ C*a*c*d*tan(e + f*x)**2/f + C*a*d**2*x + C*a*d**2*tan(e + f*x)**3/(3*f)
- C*a*d**2*tan(e + f*x)/f - C*b*c**2*log(tan(e + f*x)**2 + 1)/(2*f) + C*b*
c**2*tan(e + f*x)**2/(2*f) + 2*C*b*c*d*x + 2*C*b*c*d*tan(e + f*x)**3/(3*f)
- 2*C*b*c*d*tan(e + f*x)/f + C*b*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + C*
b*d**2*tan(e + f*x)**4/(4*f) - C*b*d**2*tan(e + f*x)**2/(2*f), Ne(f, 0)),
(x*(a + b*tan(e))*(c + d*tan(e))**2*(A + B*tan(e) + C*tan(e)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.98

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3 C b d^2 \tan^4(fx + e) + 4(2 C b c d + (C a + B b) d^2) \tan^3(fx + e) + 6(C b c^2 + 2(C a + B b) c d + (B a + (A - C) b) d^2) \tan^2(fx + e) + 12(((A - C) a - B b) c^2 - 2(B a + (A - C) b) c d - ((A - C) a - B b) d^2) (fx + e) + 6((B a + (A - C) b) c^2 + 2((A - C) a - B b) c d - (B a + (A - C) b) d^2) \log(\tan^2(fx + e) + 1) + 12((C a + B b) c^2 + 2(B a + (A - C) b) c d + ((A - C) a - B b) d^2) \tan(fx + e)}{f}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
1/12*(3*C*b*d^2*tan(f*x + e)^4 + 4*(2*C*b*c*d + (C*a + B*b)*d^2)*tan(f*x + e)^3 + 6*(C*b*c^2 + 2*(C*a + B*b)*c*d + (B*a + (A - C)*b)*d^2)*tan(f*x + e)^2 + 12*(((A - C)*a - B*b)*c^2 - 2*(B*a + (A - C)*b)*c*d - ((A - C)*a - B*b)*d^2)*(f*x + e) + 6*((B*a + (A - C)*b)*c^2 + 2*((A - C)*a - B*b)*c*d - (B*a + (A - C)*b)*d^2)*log(tan(f*x + e)^2 + 1) + 12*((C*a + B*b)*c^2 + 2*(B*a + (A - C)*b)*c*d + ((A - C)*a - B*b)*d^2)*tan(f*x + e)/f
```

Giac [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.69

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(A a c^2 - C a c^2 - B b c^2 - 2 B a c d - 2 A b c d + 2 C b c d - A a d^2 + C a d^2 + B b d^2)(f x + e)}{f} + \frac{(B a c^2 + A b c^2 - C b c^2 + 2 A a c d - 2 C a c d - 2 B b c d - B a d^2 - A b d^2 + C b d^2) \log(\tan^2(f x + e) + 1)}{2 f} + \frac{3 C b d^2 f^3 \tan^4(f x + e) + 8 C b c d f^3 \tan^3(f x + e) + 4 C a d^2 f^3 \tan^2(f x + e) + 4 B b d^2 f^3 \tan(f x + e)}{f}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
(A*a*c^2 - C*a*c^2 - B*b*c^2 - 2*B*a*c*d - 2*A*b*c*d + 2*C*b*c*d - A*a*d^2
+ C*a*d^2 + B*b*d^2)*(f*x + e)/f + 1/2*(B*a*c^2 + A*b*c^2 - C*b*c^2 + 2*A
*a*c*d - 2*C*a*c*d - 2*B*b*c*d - B*a*d^2 - A*b*d^2 + C*b*d^2)*log(tan(f*x
+ e)^2 + 1)/f + 1/12*(3*C*b*d^2*f^3*tan(f*x + e)^4 + 8*C*b*c*d*f^3*tan(f*x
+ e)^3 + 4*C*a*d^2*f^3*tan(f*x + e)^3 + 4*B*b*d^2*f^3*tan(f*x + e)^3 + 6*
C*b*c^2*f^3*tan(f*x + e)^2 + 12*C*a*c*d*f^3*tan(f*x + e)^2 + 12*B*b*c*d*f^
3*tan(f*x + e)^2 + 6*B*a*d^2*f^3*tan(f*x + e)^2 + 6*A*b*d^2*f^3*tan(f*x +
e)^2 - 6*C*b*d^2*f^3*tan(f*x + e)^2 + 12*C*a*c^2*f^3*tan(f*x + e) + 12*B*b
*c^2*f^3*tan(f*x + e) + 24*B*a*c*d*f^3*tan(f*x + e) + 24*A*b*c*d*f^3*tan(f
*x + e) - 24*C*b*c*d*f^3*tan(f*x + e) + 12*A*a*d^2*f^3*tan(f*x + e) - 12*C
*a*d^2*f^3*tan(f*x + e) - 12*B*b*d^2*f^3*tan(f*x + e))/f^4
```

Mupad [B] (verification not implemented)

Time = 5.64 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.13

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{\tan(e + fx)^2 \left(\frac{A b d^2}{2} + \frac{B a d^2}{2} + \frac{C b c^2}{2} - \frac{C b d^2}{2} + B b c d + C a c d \right)}{f}$$

$$- \frac{x (A a d^2 - A a c^2 + B b c^2 + C a c^2 - B b d^2 - C a d^2 + 2 A b c d + 2 B a c d - 2 C b c d)}{\ln(\tan(e + fx)^2 + 1) \left(\frac{A b d^2}{2} - \frac{B a c^2}{2} - \frac{A b c^2}{2} + \frac{B a d^2}{2} + \frac{C b c^2}{2} - \frac{C b d^2}{2} - A a c d + B b c d + C a c d \right)}$$

$$+ \frac{\tan(e + fx) (A a d^2 + B b c^2 + C a c^2 - B b d^2 - C a d^2 + 2 A b c d + 2 B a c d - 2 C b c d)}{f}$$

$$+ \frac{\tan(e + fx)^3 \left(\frac{B b d^2}{3} + \frac{C a d^2}{3} + \frac{2 C b c d}{3} \right)}{f} + \frac{C b d^2 \tan(e + fx)^4}{4 f}$$

input

```
int((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*ta
n(e + f*x)^2),x)
```

output

```
(tan(e + f*x)^2*((A*b*d^2)/2 + (B*a*d^2)/2 + (C*b*c^2)/2 - (C*b*d^2)/2 + B
*b*c*d + C*a*c*d))/f - x*(A*a*d^2 - A*a*c^2 + B*b*c^2 + C*a*c^2 - B*b*d^2
- C*a*d^2 + 2*A*b*c*d + 2*B*a*c*d - 2*C*b*c*d) - (log(tan(e + f*x)^2 + 1)*
((A*b*d^2)/2 - (B*a*c^2)/2 - (A*b*c^2)/2 + (B*a*d^2)/2 + (C*b*c^2)/2 - (C*
b*d^2)/2 - A*a*c*d + B*b*c*d + C*a*c*d))/f + (tan(e + f*x)*(A*a*d^2 + B*b*
c^2 + C*a*c^2 - B*b*d^2 - C*a*d^2 + 2*A*b*c*d + 2*B*a*c*d - 2*C*b*c*d))/f
+ (tan(e + f*x)^3*((B*b*d^2)/3 + (C*a*d^2)/3 + (2*C*b*c*d)/3))/f + (C*b*d^
2*tan(e + f*x)^4)/(4*f)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 432, normalized size of antiderivative = 1.62

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12acd^2fx + 24b^2c^2dfx + 12 \tan^2(fx + e) ab d^2 + 12 \tan^2(fx + e) b^2 cd - 6 \log(\tan^2(fx + e) + 1) bc^3 + \dots}{\dots}$$

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
(12*log(tan(e + f*x)**2 + 1)*a**2*c*d + 12*log(tan(e + f*x)**2 + 1)*a*b*c*
*2 - 12*log(tan(e + f*x)**2 + 1)*a*b*d**2 - 12*log(tan(e + f*x)**2 + 1)*a*
c**2*d - 12*log(tan(e + f*x)**2 + 1)*b**2*c*d - 6*log(tan(e + f*x)**2 + 1)
*b*c**3 + 6*log(tan(e + f*x)**2 + 1)*b*c*d**2 + 3*tan(e + f*x)**4*b*c*d**2
+ 4*tan(e + f*x)**3*a*c*d**2 + 4*tan(e + f*x)**3*b**2*d**2 + 8*tan(e + f*
x)**3*b*c**2*d + 12*tan(e + f*x)**2*a*b*d**2 + 12*tan(e + f*x)**2*a*c**2*d
+ 12*tan(e + f*x)**2*b**2*c*d + 6*tan(e + f*x)**2*b*c**3 - 6*tan(e + f*x)
**2*b*c*d**2 + 12*tan(e + f*x)*a**2*d**2 + 48*tan(e + f*x)*a*b*c*d + 12*ta
n(e + f*x)*a*c**3 - 12*tan(e + f*x)*a*c*d**2 + 12*tan(e + f*x)*b**2*c**2 -
12*tan(e + f*x)*b**2*d**2 - 24*tan(e + f*x)*b*c**2*d + 12*a**2*c**2*f*x -
12*a**2*d**2*f*x - 48*a*b*c*d*f*x - 12*a*c**3*f*x + 12*a*c*d**2*f*x - 12*
b**2*c**2*f*x + 12*b**2*d**2*f*x + 24*b*c**2*d*f*x)/(12*f)
```


3.60 $\int (c+d \tan(e+fx))^2 (A + B \tan(e + fx) + C \tan^2(e -$

Optimal result	652
Mathematica [C] (verified)	653
Rubi [A] (verified)	653
Maple [A] (verified)	656
Fricas [A] (verification not implemented)	656
Sympy [B] (verification not implemented)	657
Maxima [A] (verification not implemented)	657
Giac [A] (verification not implemented)	658
Mupad [B] (verification not implemented)	659
Reduce [B] (verification not implemented)	659

Optimal result

Integrand size = 33, antiderivative size = 131

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= -((c^2C + 2Bcd - Cd^2 - A(c^2 - d^2)) x$$

$$- \frac{(2c(A - C)d + B(c^2 - d^2)) \log(\cos(e + fx))}{f} + \frac{d(Bc + (A - C)d) \tan(e + fx)}{f}$$

$$+ \frac{B(c + d \tan(e + fx))^2}{2f} + \frac{C(c + d \tan(e + fx))^3}{3df}$$

output

```
-(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))*x-(2*c*(A-C)*d+B*(c^2-d^2))*ln(cos(f*x+
e))/f+d*(B*c+(A-C)*d)*tan(f*x+e)/f+1/2*B*(c+d*tan(f*x+e))^2/f+1/3*C*(c+d*t
an(f*x+e))^3/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.72 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.34

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2C(c + d \tan(e + fx))^3 + 3(Bc + (-A + C)d) (i((c + id)^2 \log(i - \tan(e + fx)) - (c - id)^2 \log(i + \tan(e + fx)))}{6df}$$

input

```
Integrate[(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(2*C*(c + d*Tan[e + f*x])^3 + 3*(B*c + (-A + C)*d)*(I*((c + I*d)^2*Log[I - Tan[e + f*x]] - (c - I*d)^2*Log[I + Tan[e + f*x]]) - 2*d^2*Tan[e + f*x]) + 3*B*((I*c - d)^3*Log[I - Tan[e + f*x]] - (I*c + d)^3*Log[I + Tan[e + f*x]]) + 6*c*d^2*Tan[e + f*x] + d^3*Tan[e + f*x]^2)/(6*d*f)
```

Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow \text{4113}$$

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^2 dx + \frac{C(c + d \tan(e + fx))^3}{3df}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^2 dx + \frac{C(c + d \tan(e + fx))^3}{3df} \\
& \downarrow 4011 \\
& \int (c + d \tan(e + fx))(Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \\
& \quad \frac{B(c + d \tan(e + fx))^2}{2f} + \frac{C(c + d \tan(e + fx))^3}{3df} \\
& \downarrow 3042 \\
& \int (c + d \tan(e + fx))(Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \\
& \quad \frac{B(c + d \tan(e + fx))^2}{2f} + \frac{C(c + d \tan(e + fx))^3}{3df} \\
& \downarrow 4008 \\
& (2cd(A - C) + B(c^2 - d^2)) \int \tan(e + fx) dx - x(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) + \\
& \quad \frac{d \tan(e + fx)(d(A - C) + Bc)}{f} + \frac{B(c + d \tan(e + fx))^2}{2f} + \frac{C(c + d \tan(e + fx))^3}{3df} \\
& \downarrow 3042 \\
& (2cd(A - C) + B(c^2 - d^2)) \int \tan(e + fx) dx - x(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) + \\
& \quad \frac{d \tan(e + fx)(d(A - C) + Bc)}{f} + \frac{B(c + d \tan(e + fx))^2}{2f} + \frac{C(c + d \tan(e + fx))^3}{3df} \\
& \downarrow 3956 \\
& -\frac{(2cd(A - C) + B(c^2 - d^2)) \log(\cos(e + fx))}{f} - x(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) + \\
& \quad \frac{d \tan(e + fx)(d(A - C) + Bc)}{f} + \frac{B(c + d \tan(e + fx))^2}{2f} + \frac{C(c + d \tan(e + fx))^3}{3df}
\end{aligned}$$

input

```
Int[(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
-((c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2))*x) - ((2*c*(A - C)*d + B*(c^2 - d^2))*Log[Cos[e + f*x]]/f + (d*(B*c + (A - C)*d)*Tan[e + f*x])/f + (B*(c + d*Tan[e + f*x])^2)/(2*f) + (C*(c + d*Tan[e + f*x])^3)/(3*d*f)
```

Definitions of rubi rules used

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}\{c, d\}, x]$

rule 4008 $\text{Int}[((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*c - b*d)*x, x] + (\text{Simp}[b*d*(\text{Tan}[e + f*x]/f), x] + \text{Simp}[(b*c + a*d) \text{ Int}[\text{Tan}[e + f*x], x], x]) \text{ ; FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[b*c + a*d, 0]$

rule 4011 $\text{Int}[((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)])^m*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{m-1}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

rule 4113 $\text{Int}[((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)])^m*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_)] + (C_.)*\tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] \rightarrow \text{Simp}[C*((a + b*\text{Tan}[e + f*x])^{m+1}/(b*f*(m+1))), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^m*\text{Simp}[A - C + B*\text{Tan}[e + f*x], x], x] \text{ ; FreeQ}\{a, b, e, f, A, B, C, m\}, x \ \&\& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\& \ !\text{LeQ}[m, -1]$

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.08

method	result
norman	$(A c^2 - A d^2 - 2Bcd - c^2 C + C d^2) x + \frac{(A d^2 + 2Bcd + c^2 C - C d^2) \tan(fx+e)}{f} + \frac{C d^2 \tan(fx+e)^3}{3f}$
parts	$A c^2 x + \frac{(2Acd + B c^2) \ln(1 + \tan(fx+e)^2)}{2f} + \frac{(B d^2 + 2Ccd) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1 + \tan(fx+e)^2)}{2} \right)}{f} + \frac{(A d^2 + 2Bcd + c^2 C - C d^2) \tan(fx+e)}{f}$
derivativdivides	$\frac{C d^2 \tan(fx+e)^3}{3} + \frac{B d^2 \tan(fx+e)^2}{2} + Ccd \tan(fx+e)^2 + \tan(fx+e) A d^2 + 2 \tan(fx+e) Bcd + \tan(fx+e) c^2 C - \tan(fx+e) C d^2}{f}$
default	$\frac{C d^2 \tan(fx+e)^3}{3} + \frac{B d^2 \tan(fx+e)^2}{2} + Ccd \tan(fx+e)^2 + \tan(fx+e) A d^2 + 2 \tan(fx+e) Bcd + \tan(fx+e) c^2 C - \tan(fx+e) C d^2}{f}$
parallelrisc	$\frac{2C d^2 \tan(fx+e)^3 + 6A x c^2 f - 6A d^2 f x - 12Bcdfx + 3B d^2 \tan(fx+e)^2 - 6C c^2 f x + 6C d^2 f x + 6Ccd \tan(fx+e)^2 + 6A \ln(1 + \tan(fx+e)^2)}{f}$
risc	$iB c^2 x - \frac{2iB d^2 e}{f} + 2iAcdx + \frac{4iAcde}{f} + A c^2 x - A d^2 x - 2Bcdx - C c^2 x + C d^2 x + \frac{2iBc}{f}$

input `int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETURNVERBOSE)`

output $(A*c^2-A*d^2-2*B*c*d-C*c^2+C*d^2)*x+(A*d^2+2*B*c*d+C*c^2-C*d^2)/f*\tan(f*x+e)+1/3*C*d^2/f*\tan(f*x+e)^3+1/2*d*(B*d+2*C*c)/f*\tan(f*x+e)^2+1/2*(2*A*c*d+B*c^2-B*d^2-2*C*c*d)/f*\ln(1+\tan(f*x+e)^2)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.02

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2C d^2 \tan(fx+e)^3 + 6((A-C)c^2 - 2Bcd - (A-C)d^2)fx + 3(2Ccd + Bd^2) \tan(fx+e)^2 - 3(BCd^2 - B^2d^2 - 2C^2cd)}{6f}$$

input `integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output

```
1/6*(2*C*d^2*tan(f*x + e)^3 + 6*((A - C)*c^2 - 2*B*c*d - (A - C)*d^2)*f*x
+ 3*(2*C*c*d + B*d^2)*tan(f*x + e)^2 - 3*(B*c^2 + 2*(A - C)*c*d - B*d^2)*l
og(1/(tan(f*x + e)^2 + 1)) + 6*(C*c^2 + 2*B*c*d + (A - C)*d^2)*tan(f*x + e
))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 241 vs. $2(107) = 214$.

Time = 0.13 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.84

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \begin{cases} Ac^2x + \frac{Acd \log(\tan^2(e+fx)+1)}{f} - Ad^2x + \frac{Ad^2 \tan(e+fx)}{f} + \frac{Bc^2 \log(\tan^2(e+fx)+1)}{2f} - 2Bcdx + \frac{2Bcd \tan(e+fx)}{f} - \frac{Bd^2 \tan^2(e+fx)}{2f} \\ x(c + d \tan(e))^2 (A + B \tan(e) + C \tan^2(e)) \end{cases}$$

input

```
integrate((c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A*c**2*x + A*c*d*log(tan(e + f*x)**2 + 1)/f - A*d**2*x + A*d**2
*tan(e + f*x)/f + B*c**2*log(tan(e + f*x)**2 + 1)/(2*f) - 2*B*c*d*x + 2*B*
c*d*tan(e + f*x)/f - B*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + B*d**2*tan(e
+ f*x)**2/(2*f) - C*c**2*x + C*c**2*tan(e + f*x)/f - C*c*d*log(tan(e + f*x
)**2 + 1)/f + C*c*d*tan(e + f*x)**2/f + C*d**2*x + C*d**2*tan(e + f*x)**3/
(3*f) - C*d**2*tan(e + f*x)/f, Ne(f, 0)), (x*(c + d*tan(e))**2*(A + B*tan(
e) + C*tan(e)**2), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.03

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2Cd^2 \tan^3(fx + e) + 3(2Ccd + Bd^2) \tan^2(fx + e) + 6((A - C)c^2 - 2Bcd - (A - C)d^2)(fx + e) + 3(Ac^2 + 2Bcd + Cc^2) \tan(fx + e) + 3(Ac^2 + 2Bcd + Cc^2) \tan^2(fx + e) + 3(Ac^2 + 2Bcd + Cc^2) \tan^3(fx + e)}{6f}$$

input `integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output $\frac{1}{6}*(2*C*d^2*\tan(f*x + e)^3 + 3*(2*C*c*d + B*d^2)*\tan(f*x + e)^2 + 6*((A - C)*c^2 - 2*B*c*d - (A - C)*d^2)*(f*x + e) + 3*(B*c^2 + 2*(A - C)*c*d - B*d^2)*\log(\tan(f*x + e)^2 + 1) + 6*(C*c^2 + 2*B*c*d + (A - C)*d^2)*\tan(f*x + e))/f$

Giac [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.46

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(Ac^2 - Cc^2 - 2Bcd - Ad^2 + Cd^2)(fx + e)}{f} + \frac{(Bc^2 + 2Acd - 2Ccd - Bd^2) \log(\tan(fx + e)^2 + 1)}{2f} + \frac{2Cd^2f^2 \tan(fx + e)^3 + 6Ccdf^2 \tan(fx + e)^2 + 3Bd^2f^2 \tan(fx + e)^2 + 6Cc^2f^2 \tan(fx + e) + 12Bcd^2f^2}{6f^3}$$

input `integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output $(A*c^2 - C*c^2 - 2*B*c*d - A*d^2 + C*d^2)*(f*x + e)/f + 1/2*(B*c^2 + 2*A*c*d - 2*C*c*d - B*d^2)*\log(\tan(f*x + e)^2 + 1)/f + 1/6*(2*C*d^2*f^2*\tan(f*x + e)^3 + 6*C*c*d*f^2*\tan(f*x + e)^2 + 3*B*d^2*f^2*\tan(f*x + e)^2 + 6*C*c^2*f^2*\tan(f*x + e) + 12*B*c*d*f^2*\tan(f*x + e) + 6*A*d^2*f^2*\tan(f*x + e) - 6*C*d^2*f^2*\tan(f*x + e))/f^3$

Mupad [B] (verification not implemented)

Time = 5.53 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.08

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{\tan(e + fx)^2 \left(\frac{Bd^2}{2} + Ccd \right)}{f} - x (Ad^2 - Ac^2 + Cc^2 - Cd^2 + 2Bcd)$$

$$+ \frac{\tan(e + fx) (Ad^2 + Cc^2 - Cd^2 + 2Bcd)}{f}$$

$$+ \frac{\ln(\tan(e + fx)^2 + 1) \left(\frac{Bc^2}{2} - \frac{Bd^2}{2} + Acd - Ccd \right)}{f} + \frac{Cd^2 \tan(e + fx)^3}{3f}$$

input

```
int((c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
(tan(e + f*x)^2*((B*d^2)/2 + C*c*d))/f - x*(A*d^2 - A*c^2 + C*c^2 - C*d^2 + 2*B*c*d) + (tan(e + f*x)*(A*d^2 + C*c^2 - C*d^2 + 2*B*c*d))/f + (log(tan(e + f*x)^2 + 1)*((B*c^2)/2 - (B*d^2)/2 + A*c*d - C*c*d))/f + (C*d^2*tan(e + f*x)^3)/(3*f)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.52

$$\int (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{6 \log(\tan(fx + e)^2 + 1) acd + 3 \log(\tan(fx + e)^2 + 1) bc^2 - 3 \log(\tan(fx + e)^2 + 1) bd^2 - 6 \log(\tan(fx + e)^2 + 1) cd^2}{3f}$$

input

```
int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```


output

```
(6*log(tan(e + f*x)**2 + 1)*a*c*d + 3*log(tan(e + f*x)**2 + 1)*b*c**2 - 3*
log(tan(e + f*x)**2 + 1)*b*d**2 - 6*log(tan(e + f*x)**2 + 1)*c**2*d + 2*ta
n(e + f*x)**3*c*d**2 + 3*tan(e + f*x)**2*b*d**2 + 6*tan(e + f*x)**2*c**2*d
+ 6*tan(e + f*x)*a*d**2 + 12*tan(e + f*x)*b*c*d + 6*tan(e + f*x)*c**3 - 6
*tan(e + f*x)*c*d**2 + 6*a*c**2*f*x - 6*a*d**2*f*x - 12*b*c*d*f*x - 6*c**3
*f*x + 6*c*d**2*f*x)/(6*f)
```

3.61
$$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$$

Optimal result	661
Mathematica [C] (verified)	662
Rubi [A] (verified)	662
Maple [A] (verified)	667
Fricas [A] (verification not implemented)	667
Sympy [C] (verification not implemented)	668
Maxima [A] (verification not implemented)	669
Giac [A] (verification not implemented)	670
Mupad [B] (verification not implemented)	671
Reduce [B] (verification not implemented)	672

Optimal result

Integrand size = 45, antiderivative size = 254

$$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$$

$$= -\frac{(a(c^2C+2Bcd-Cd^2-A(c^2-d^2))-b(2c(A-C)d+B(c^2-d^2)))x}{a^2+b^2}$$

$$-\frac{(a(Bc^2-2cCd-Bd^2)+b(c^2C+2Bcd-Cd^2)+A(2acd-b(c^2-d^2)))\log(\cos(e+fx))}{(a^2+b^2)f}$$

$$+\frac{(Ab^2-a(bB-aC))(bc-ad)^2\log(a+b \tan(e+fx))}{b^3(a^2+b^2)f}$$

$$+\frac{d(bcC+bBd-aCd)\tan(e+fx)}{b^2f}+\frac{C(c+d \tan(e+fx))^2}{2bf}$$

output

```
-(a*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-b*(2*c*(A-C)*d+B*(c^2-d^2)))*x/(a^2+b^2)-(a*(B*c^2-B*d^2-2*C*c*d)+b*(2*B*c*d+C*c^2-C*d^2)+A*(2*a*c*d-b*(c^2-d^2)))*ln(cos(f*x+e))/(a^2+b^2)/f+(A*b^2-a*(B*b-C*a))*(-a*d+b*c)^2*ln(a+b*tan(f*x+e))/b^3/(a^2+b^2)/f+d*(B*b*d-C*a*d+C*b*c)*tan(f*x+e)/b^2/f+1/2*C*(c+d*tan(f*x+e))^2/b/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.00 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.75

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{\frac{b(-iA+B+iC)(c+id)^2 \log(i-\tan(e+fx))}{a+ib} + \frac{b(iA+B-iC)(c-id)^2 \log(i+\tan(e+fx))}{a-ib} + \frac{2(Ab^2+a(-bB+aC))(bc-ad)^2 \log(a+b \tan(e+fx))}{b^2(a^2+b^2)}}{2bf}$$

input

```
Integrate[((c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(a + b*Tan[e + f*x]),x]
```

output

```
((b*((-I)*A + B + I*C)*(c + I*d)^2*Log[I - Tan[e + f*x]])/(a + I*b) + (b*(
I*A + B - I*C)*(c - I*d)^2*Log[I + Tan[e + f*x]])/(a - I*b) + (2*(A*b^2 +
a*(-(b*B) + a*C))*(b*c - a*d)^2*Log[a + b*Tan[e + f*x]])/(b^2*(a^2 + b^2))
+ (2*d*(b*c*C + b*B*d - a*C*d)*Tan[e + f*x])/b + C*(c + d*Tan[e + f*x])^2
)/(2*b*f)
```

Rubi [A] (verified)

Time = 2.47 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4130, 27, 3042, 4120, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$\downarrow 3042$$

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{a + b \tan(e + fx)} dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \frac{\int \frac{2(c+d \tan(e+fx))((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx}{\frac{2b}{C(c+d \tan(e+fx))^2}} + \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{\int \frac{(c+d \tan(e+fx))((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx}{\frac{b}{C(c+d \tan(e+fx))^2}} + \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{\int \frac{(c+d \tan(e+fx))((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx}{\frac{b}{C(c+d \tan(e+fx))^2}} + \\
 & \qquad \qquad \qquad \downarrow 4120 \\
 & \frac{\frac{d \tan(e+fx)(-aCd+bBd+bcC)}{bf} - \int -\frac{Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+((Cc^2+2Bdc+(A-C)d^2)b^2-ad(2cC+Bd)b+a^2Cd^2) \tan^2(e+fx)}{a+b \tan(e+fx)} dx}{\frac{b}{C(c+d \tan(e+fx))^2}} + \\
 & \qquad \qquad \qquad \downarrow 25 \\
 & \frac{\int \frac{Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+((Cc^2+2Bdc+(A-C)d^2)b^2-ad(2cC+Bd)b+a^2Cd^2) \tan^2(e+fx)+ad(aCd-b(2cC+Bd))}{a+b \tan(e+fx)} dx}{\frac{b}{C(c+d \tan(e+fx))^2}} + \frac{d \tan(e+fx)}{b} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{\int \frac{Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+((Cc^2+2Bdc+(A-C)d^2)b^2-ad(2cC+Bd)b+a^2Cd^2) \tan(e+fx)^2+ad(aCd-b(2cC+Bd))}{a+b \tan(e+fx)} dx}{\frac{b}{C(c+d \tan(e+fx))^2}} + \frac{d \tan(e+fx)}{b} \\
 & \qquad \qquad \qquad \downarrow 4109
 \end{aligned}$$

$$\frac{b^2(2aAcd+aB(c^2-d^2)-2acCd-Ab(c^2-d^2)+b(2Bcd+c^2C-Cd^2)) \int \tan(e+fx) dx}{a^2+b^2} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan^2(e+fx)+1}{a+b \tan(e+fx)} dx}{b(a^2+b^2)} - \frac{b^2x(a(-A(c^2-d^2)+$$

$$\frac{C(c+d \tan(e+fx))^2}{2bf} \qquad b$$

↓ 3042

$$\frac{b^2(2aAcd+aB(c^2-d^2)-2acCd-Ab(c^2-d^2)+b(2Bcd+c^2C-Cd^2)) \int \tan(e+fx) dx}{a^2+b^2} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{a+b \tan(e+fx)} dx}{b(a^2+b^2)} - \frac{b^2x(a(-A(c^2-d^2)+$$

$$\frac{C(c+d \tan(e+fx))^2}{2bf} \qquad b$$

↓ 3956

$$\frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{a+b \tan(e+fx)} dx}{a^2+b^2} - \frac{b^2 \log(\cos(e+fx))(2aAcd+aB(c^2-d^2)-2acCd-Ab(c^2-d^2)+b(2Bcd+c^2C-Cd^2))}{f(a^2+b^2)} - \frac{b^2x(a(-A(c^2-d^2)+$$

$$\frac{C(c+d \tan(e+fx))^2}{2bf} \qquad b$$

↓ 4100

$$\frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{1}{a+b \tan(e+fx)} d(b \tan(e+fx))}{bf(a^2+b^2)} - \frac{b^2 \log(\cos(e+fx))(2aAcd+aB(c^2-d^2)-2acCd-Ab(c^2-d^2)+b(2Bcd+c^2C-Cd^2))}{f(a^2+b^2)} - \frac{b^2x(a(-A(c^2-d^2)+$$

$$\frac{C(c+d \tan(e+fx))^2}{2bf} \qquad b$$

↓ 16

$$\frac{b^2 \log(\cos(e+fx))(2aAcd+aB(c^2-d^2)-2acCd-Ab(c^2-d^2)+b(2Bcd+c^2C-Cd^2))}{f(a^2+b^2)} - \frac{b^2x(a(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)-b(2cd(A-C)+B(c^2-d^2)))}{a^2+b^2}$$

$$\frac{C(c+d \tan(e+fx))^2}{2bf} \qquad b$$

input `Int[((c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]`

output

$$\begin{aligned} & (C*(c + d*\text{Tan}[e + f*x])^2)/(2*b*f) + (((-(b^2*(a*(c^2*C + 2*B*c*d - C*d^2 \\ & - A*(c^2 - d^2)) - b*(2*c*(A - C)*d + B*(c^2 - d^2))) * x)/(a^2 + b^2)) - (b \\ & ^2*(2*a*A*c*d - 2*a*c*C*d - A*b*(c^2 - d^2) + a*B*(c^2 - d^2) + b*(c^2*C + \\ & 2*B*c*d - C*d^2))*\text{Log}[\text{Cos}[e + f*x]])/(a^2 + b^2)*f) + ((A*b^2 - a*(b*B - \\ & a*C))*(b*c - a*d)^2*\text{Log}[a + b*\text{Tan}[e + f*x]])/(b*(a^2 + b^2)*f)/b + (d*(b \\ & *c*C + b*B*d - a*C*d)*\text{Tan}[e + f*x])/(b*f))/b \end{aligned}$$

Defintions of rubi rules used

rule 16

$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$$

rule 25

$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$$

rule 27

$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$$

rule 3042

$$\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\text{tan}[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4100

$$\text{Int}[(a_)+(b_)*\text{tan}[(e_)+(f_)*(x_)]^{(m_)}*((A_)+(C_)*\text{tan}[(e_)+(f_)*(x_)]^2), x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] \text{ ; FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$$

rule 4109

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2
)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]
```

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_
)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(GtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.25

method	result
derivativedivides	$\frac{d\left(\frac{\tan(fx+e)^2 Cbd + B \tan(fx+e)bd - d \tan(fx+e)Ca + 2 \tan(fx+e)Cbc}{b^2}\right)}{b^2} + \frac{(2Aacd - Abc^2 + Abd^2 + Bac^2 - B ad^2 + 2Bbcd - 2Cacd + 2C^2d^2)}{2}$
default	$\frac{d\left(\frac{\tan(fx+e)^2 Cbd + B \tan(fx+e)bd - d \tan(fx+e)Ca + 2 \tan(fx+e)Cbc}{b^2}\right)}{b^2} + \frac{(2Aacd - Abc^2 + Abd^2 + Bac^2 - B ad^2 + 2Bbcd - 2Cacd + 2C^2d^2)}{2}$
norman	$\frac{(Aa^2c^2 - Aad^2 + 2Abcd - 2Bacd + Bb^2c^2 - Bbd^2 - Ca^2c^2 + Cad^2 - 2Cbcd)x}{a^2 + b^2} + \frac{d(Bbd - Cad + 2Cbc) \tan(fx+e)}{b^2 f} + \frac{C^2 d^2}{2}$
parallelrisc	$C \tan(fx+e)^2 a^2 b^2 d^2 + 2B \tan(fx+e) a^2 b^2 d^2 - 2C \tan(fx+e) a^3 b d^2 - 2C \tan(fx+e) a b^3 d^2 + 4C \tan(fx+e) b^4 cd + 2Bx$
risc	Expression too large to display

```
input int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x,
method=_RETURNVERBOSE)
```

```
output 1/f*(d/b^2*(1/2*tan(f*x+e)^2*C*b*d+B*tan(f*x+e)*b*d-d*tan(f*x+e)*C*a+2*tan
(f*x+e)*C*b*c)+1/(a^2+b^2)*(1/2*(2*A*a*c*d-A*b*c^2+A*b*d^2+B*a*c^2-B*a*d^2
+2*B*b*c*d-2*C*a*c*d+C*b*c^2-C*b*d^2)*ln(1+tan(f*x+e)^2)+(A*a*c^2-A*a*d^2+
2*A*b*c*d-2*B*a*c*d+B*b*c^2-B*b*d^2-C*a*c^2+C*a*d^2-2*C*b*c*d)*arctan(tan(
f*x+e)))+1/b^3*(A*a^2*b^2*d^2-2*A*a*b^3*c*d+A*b^4*c^2-B*a^3*b*d^2+2*B*a^2*
b^2*c*d-B*a*b^3*c^2+C*a^4*d^2-2*C*a^3*b*c*d+C*a^2*b^2*c^2)/(a^2+b^2)*ln(a+
b*tan(f*x+e))
```

Fricas [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 397, normalized size of antiderivative = 1.56

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{(Ca^2b^2 + Cb^4)d^2 \tan^2(fx + e) + 2(((A - C)ab^3 + Bb^4)c^2 - 2(Bab^3 - (A - C)b^4)cd - ((A - C)ab^3 +$$

input `integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output `1/2*((C*a^2*b^2 + C*b^4)*d^2*tan(f*x + e)^2 + 2*((A - C)*a*b^3 + B*b^4)*c^2 - 2*(B*a*b^3 - (A - C)*b^4)*c*d - ((A - C)*a*b^3 + B*b^4)*d^2)*f*x + ((C*a^2*b^2 - B*a*b^3 + A*b^4)*c^2 - 2*(C*a^3*b - B*a^2*b^2 + A*a*b^3)*c*d + (C*a^4 - B*a^3*b + A*a^2*b^2)*d^2)*log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) - ((C*a^2*b^2 + C*b^4)*c^2 - 2*(C*a^3*b - B*a^2*b^2 + C*a*b^3 - B*b^4)*c*d + (C*a^4 - B*a^3*b + A*a^2*b^2 - B*a*b^3 + (A - C)*b^4)*d^2)*log(1/(tan(f*x + e)^2 + 1)) + 2*(2*(C*a^2*b^2 + C*b^4)*c*d - (C*a^3*b - B*a^2*b^2 + C*a*b^3 - B*b^4)*d^2)*tan(f*x + e))/((a^2*b^3 + b^5)*f)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.08 (sec) , antiderivative size = 4444, normalized size of antiderivative = 17.50

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)),x)`

output

```

1/2*(2*((A - C)*a + B*b)*c^2 - 2*(B*a - (A - C)*b)*c*d - ((A - C)*a + B*b
)*d^2)*(f*x + e)/(a^2 + b^2) + 2*((C*a^2*b^2 - B*a*b^3 + A*b^4)*c^2 - 2*(C
*a^3*b - B*a^2*b^2 + A*a*b^3)*c*d + (C*a^4 - B*a^3*b + A*a^2*b^2)*d^2)*log
(b*tan(f*x + e) + a)/(a^2*b^3 + b^5) + ((B*a - (A - C)*b)*c^2 + 2*((A - C)
*a + B*b)*c*d - (B*a - (A - C)*b)*d^2)*log(tan(f*x + e)^2 + 1)/(a^2 + b^2)
+ (C*b*d^2*tan(f*x + e)^2 + 2*(2*C*b*c*d - (C*a - B*b)*d^2)*tan(f*x + e))
/b^2)/f

```

Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.35

$$\begin{aligned}
& \int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\
&= \frac{(Aac^2 - Cac^2 + Bbc^2 - 2 Bacd + 2 Abcd - 2 Cbcd - Aad^2 + Cad^2 - Bbd^2)(fx + e)}{a^2 f + b^2 f} \\
&+ \frac{(Bac^2 - Abc^2 + Cbc^2 + 2 Aacd - 2 Cacd + 2 Bbcd - Bad^2 + Abd^2 - Cbd^2) \log(\tan(fx + e)^2 + 1)}{2(a^2 f + b^2 f)} \\
&+ \frac{(Ca^2 b^2 c^2 - Bab^3 c^2 + Ab^4 c^2 - 2 Ca^3 bcd + 2 Ba^2 b^2 cd - 2 Aab^3 cd + Ca^4 d^2 - Ba^3 bd^2 + Aa^2 b^2 d^2) \log(\tan(fx + e)^2 + 1)}{a^2 b^3 f + b^5 f} \\
&+ \frac{Cbd^2 f \tan(fx + e)^2 + 4 Cbcdf \tan(fx + e) - 2 Cad^2 f \tan(fx + e) + 2 Bbd^2 f \tan(fx + e)}{2 b^2 f^2}
\end{aligned}$$

input

```

integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e)),x, algorithm="giac")

```

output

```

(A*a*c^2 - C*a*c^2 + B*b*c^2 - 2*B*a*c*d + 2*A*b*c*d - 2*C*b*c*d - A*a*d^2
+ C*a*d^2 - B*b*d^2)*(f*x + e)/(a^2*f + b^2*f) + 1/2*(B*a*c^2 - A*b*c^2 +
C*b*c^2 + 2*A*a*c*d - 2*C*a*c*d + 2*B*b*c*d - B*a*d^2 + A*b*d^2 - C*b*d^2
)*log(tan(f*x + e)^2 + 1)/(a^2*f + b^2*f) + (C*a^2*b^2*c^2 - B*a*b^3*c^2 +
A*b^4*c^2 - 2*C*a^3*b*c*d + 2*B*a^2*b^2*c*d - 2*A*a*b^3*c*d + C*a^4*d^2 -
B*a^3*b*d^2 + A*a^2*b^2*d^2)*log(abs(b*tan(f*x + e) + a))/(a^2*b^3*f + b^
5*f) + 1/2*(C*b*d^2*f*tan(f*x + e)^2 + 4*C*b*c*d*f*tan(f*x + e) - 2*C*a*d^
2*f*tan(f*x + e) + 2*B*b*d^2*f*tan(f*x + e))/(b^2*f^2)

```

Mupad [B] (verification not implemented)

Time = 8.01 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.28

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{\tan(e + fx) \left(\frac{Bd^2 + 2Ccd}{b} - \frac{Ca^2d^2}{b^2} \right)}{f}$$

$$+ \frac{\ln(a + b \tan(e + fx)) (b^2 (Ca^2c^2 + 2Ba^2cd + Aa^2d^2) - b(Ba^3d^2 + 2Cca^3d) - b^3(Bac^2 + 2Aa^2cd))}{f(a^2b^3 + b^5)}$$

$$+ \frac{\ln(\tan(e + fx) + 1i) (Ad^2 - Ac^2 + Bc^21i - Bd^21i + Cc^2 - Cd^2 + Acd2i + 2Bcd - Ccd2i)}{2f(b + a1i)}$$

$$+ \frac{\ln(\tan(e + fx) - 1i) (Bc^2 - Bd^2 + 2Acd - 2Ccd - Ac^21i + Ad^21i + Cc^21i - Cd^21i + Bcd2i)}{2f(a + b1i)}$$

$$+ \frac{Cd^2 \tan(e + fx)^2}{2bf}$$

input

```
int(((c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x)),x)
```

output

```
(tan(e + f*x)*((B*d^2 + 2*C*c*d)/b - (C*a*d^2)/b^2))/f + (log(a + b*tan(e + f*x))*(b^2*(A*a^2*d^2 + C*a^2*c^2 + 2*B*a^2*c*d) - b*(B*a^3*d^2 + 2*C*a^3*c*d) - b^3*(B*a*c^2 + 2*A*a*c*d) + A*b^4*c^2 + C*a^4*d^2))/(f*(b^5 + a^2*b^3)) + (log(tan(e + f*x) + 1i)*(A*d^2 - A*c^2 + B*c^2*1i - B*d^2*1i + C*c^2 - C*d^2 + A*c*d*2i + 2*B*c*d - C*c*d*2i))/(2*f*(a*1i + b)) + (log(tan(e + f*x) - 1i)*(A*d^2*1i - A*c^2*1i + B*c^2 - B*d^2 + C*c^2*1i - C*d^2*1i + 2*A*c*d + B*c*d*2i - 2*C*c*d))/(2*f*(a + b*1i)) + (C*d^2*tan(e + f*x)^2)/(2*b*f)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.54

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{2 \log(\tan(fx + e)^2 + 1) a^2 b^3 cd - 2 \log(\tan(fx + e)^2 + 1) a b^3 c^2 d + 2 \log(\tan(fx + e)^2 + 1) b^5 cd + \log$$

input

```
int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x)
```

output

```
(2*log(tan(e + f*x)**2 + 1)*a**2*b**3*c*d - 2*log(tan(e + f*x)**2 + 1)*a*b
**3*c**2*d + 2*log(tan(e + f*x)**2 + 1)*b**5*c*d + log(tan(e + f*x)**2 + 1
)*b**4*c**3 - log(tan(e + f*x)**2 + 1)*b**4*c*d**2 + 2*log(tan(e + f*x)*b
+ a)*a**4*c*d**2 - 4*log(tan(e + f*x)*b + a)*a**3*b*c**2*d + 2*log(tan(e +
f*x)*b + a)*a**2*b**2*c**3 + tan(e + f*x)**2*a**2*b**2*c*d**2 + tan(e + f
*x)**2*b**4*c*d**2 - 2*tan(e + f*x)*a**3*b*c*d**2 + 2*tan(e + f*x)*a**2*b*
*3*d**2 + 4*tan(e + f*x)*a**2*b**2*c**2*d - 2*tan(e + f*x)*a*b**3*c*d**2 +
2*tan(e + f*x)*b**5*d**2 + 4*tan(e + f*x)*b**4*c**2*d + 2*a**2*b**3*c**2*
f*x - 2*a**2*b**3*d**2*f*x - 2*a*b**3*c**3*f*x + 2*a*b**3*c*d**2*f*x + 2*b
**5*c**2*f*x - 2*b**5*d**2*f*x - 4*b**4*c**2*d*f*x)/(2*b**3*f*(a**2 + b**2
))
```

3.62
$$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$$

Optimal result	673
Mathematica [C] (verified)	674
Rubi [A] (verified)	674
Maple [A] (verified)	678
Fricas [B] (verification not implemented)	679
Sympy [C] (verification not implemented)	680
Maxima [A] (verification not implemented)	681
Giac [A] (verification not implemented)	682
Mupad [B] (verification not implemented)	683
Reduce [B] (verification not implemented)	684

Optimal result

Integrand size = 45, antiderivative size = 415

$$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx =$$

$$\frac{(a^2(c^2C+2Bcd-Cd^2-A(c^2-d^2))-b^2(c^2C+2Bcd-Cd^2-A(c^2-d^2))-2ab(2c(A-C)d+B(c^2-d^2))}{(a^2+b^2)^2}$$

$$\frac{(2ab(c^2C+2Bcd-Cd^2-A(c^2-d^2))+a^2(2c(A-C)d+B(c^2-d^2))-b^2(2c(A-C)d+B(c^2-d^2))}{(a^2+b^2)^2} f$$

$$\frac{(bc-ad)(a^3bBd-2a^4Cd-b^4(Bc+2Ad))-ab^3(2Ac-2cC-3Bd)+a^2b^2(Bc-4Cd) \log(a+b \tan(e+fx))}{b^3(a^2+b^2)^2} f$$

$$+ \frac{(Ab^2-abB+2a^2C+b^2C)d^2 \tan(e+fx)}{b^2(a^2+b^2)f}$$

$$- \frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}{b(a^2+b^2)f(a+b \tan(e+fx))}$$

output

```
-(a^2*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-b^2*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-2*a*b*(2*c*(A-C)*d+B*(c^2-d^2))*x/(a^2+b^2)^2-(2*a*b*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))+a^2*(2*c*(A-C)*d+B*(c^2-d^2))-b^2*(2*c*(A-C)*d+B*(c^2-d^2))*ln(cos(f*x+e))/(a^2+b^2)^2/f-(-a*d+b*c)*(a^3*b*B*d-2*a^4*C*d-b^4*(2*A*d+B*c)-a*b^3*(2*A*c-3*B*d-2*C*c)+a^2*b^2*(B*c-4*C*d))*ln(a+b*tan(f*x+e))/b^3/(a^2+b^2)^2/f+(A*b^2-B*a*b+2*C*a^2+C*b^2)*d^2*tan(f*x+e)/b^2/(a^2+b^2)/f-(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^2/b/(a^2+b^2)/f/(a+b*tan(f*x+e))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.04 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.67

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{\frac{(-iA+B+iC)(c+id)^2 \log(i-\tan(e+fx))}{(a+ib)^2} + \frac{(iA+B-iC)(c-id)^2 \log(i+\tan(e+fx))}{(a-ib)^2} + \frac{2(bc-ad)(-a^3bBd+2a^4Cd+b^4(Bc+2Ad)+ab^3(2Ac-b^3(a^2-2f))}{b^3(a^2-2f)}}{b^3(a^2-2f)}$$

input

```
Integrate[((c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(a + b*Tan[e + f*x])^2,x]
```

output

```
((((-I)*A + B + I*C)*(c + I*d)^2*Log[I - Tan[e + f*x]])/(a + I*b)^2 + ((I*
A + B - I*C)*(c - I*d)^2*Log[I + Tan[e + f*x]])/(a - I*b)^2 + (2*(b*c - a*
d)*(-(a^3*b*B*d) + 2*a^4*C*d + b^4*(B*c + 2*A*d) + a*b^3*(2*A*c - 2*c*C -
3*B*d) + a^2*b^2*(-(B*c) + 4*C*d))*Log[a + b*Tan[e + f*x]])/(b^3*(a^2 + b^
2)^2) - (2*(A*b^2 - a*b*B + 2*a^2*C + b^2*C)*(b*c - a*d)^2)/(b^3*(a^2 + b^
2)*(a + b*Tan[e + f*x])) + (2*C*(c + d*Tan[e + f*x])^2)/(b*(a + b*Tan[e +
f*x]))) / (2*f)
```

Rubi [A] (verified)

Time = 3.26 (sec) , antiderivative size = 429, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4128, 3042, 4120, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^2} dx$$

↓ 4128

$$\int \frac{(c+d \tan(e+fx))((2Ca^2-bBa+Ab^2+b^2C)d \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-2ad)+Ab(ac+2bd))}{a+b \tan(e+fx)} dx$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))((2Ca^2-bBa+Ab^2+b^2C)d \tan(e+fx)^2-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-2ad)+Ab(ac+2bd))}{a+b \tan(e+fx)} dx$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 4120

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \int \frac{-((2aAcd-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2)-c((bB-aC)(bc-2ad)+Ab(ac+2bd))}{a+b \tan(e+fx)} dx$$

$$\frac{b(a^2+b^2)}{b(a^2+b^2)}$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \int \frac{-((2aAcd-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2)-c((bB-aC)(bc-2ad)+Ab(ac+2bd))}{a+b \tan(e+fx)} dx$$

$$\frac{b(a^2+b^2)}{b(a^2+b^2)}$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 4109

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \frac{b^2(a^2(2cd(A-C)+B(c^2-d^2))+2ab(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)-b^2(2cd(A-C)+B(c^2-d^2)))}{a^2+b^2} \int \tan(e+fx) dx$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \frac{b^2(a^2(2cd(A-C)+B(c^2-d^2))+2ab(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)-b^2(2cd(A-C)+B(c^2-d^2)))}{a^2+b^2} \int \tan(e+fx) dx$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 3956

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \frac{(bc-ad)(-2a^4Cd+a^3bBd+a^2b^2(Bc-4Cd)-ab^3(2Ac-3Bd-2cC)-b^4(2Ad+Bc))}{a^2+b^2} \int \frac{\tan(e+fx)^2+1}{a+b \tan(e+fx)} dx + b^2 \log(\cos(e+fx))$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 4100

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \frac{(bc-ad)(-2a^4Cd+a^3bBd+a^2b^2(Bc-4Cd)-ab^3(2Ac-3Bd-2cC)-b^4(2Ad+Bc))}{bf(a^2+b^2)} \int \frac{1}{a+b \tan(e+fx)} d(b \tan(e+fx))$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 16

$$\frac{d^2 \tan(e+fx)(2a^2C-abB+Ab^2+b^2C)}{bf} - \frac{b^2 \log(\cos(e+fx))(a^2(2cd(A-C)+B(c^2-d^2))+2ab(-A(c^2-d^2)+2Bcd+c^2C-Cd^2)-b^2(2cd(A-C)+B(c^2-d^2)))}{f(a^2+b^2)}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

input

```
Int[((c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^2,x]
```

output

$$\begin{aligned}
& -(((A*b^2 - a*(b*B - a*C))*(c + d*\text{Tan}[e + f*x])^2)/(b*(a^2 + b^2)*f*(a + b \\
& * \text{Tan}[e + f*x])) + (-(((b^2*(a^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) \\
& - b^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - 2*a*b*(2*c*(A - C)*d + \\
& B*(c^2 - d^2)))*x)/(a^2 + b^2) + (b^2*(2*a*b*(c^2*C + 2*B*c*d - C*d^2 - A \\
& (c^2 - d^2)) + a^2*(2*c*(A - C)*d + B*(c^2 - d^2)) - b^2*(2*c*(A - C)*d + \\
& B*(c^2 - d^2)))*\text{Log}[\text{Cos}[e + f*x]])/(a^2 + b^2)*f) + ((b*c - a*d)*(a^3*b*B \\
& *d - 2*a^4*C*d - b^4*(B*c + 2*A*d) - a*b^3*(2*A*c - 2*c*C - 3*B*d) + a^2*b \\
& ^2*(B*c - 4*C*d))*\text{Log}[a + b*\text{Tan}[e + f*x]])/(b*(a^2 + b^2)*f)/b) + ((A*b^2 \\
& - a*b*B + 2*a^2*C + b^2*C)*d^2*\text{Tan}[e + f*x])/(b*f))/(b*(a^2 + b^2))
\end{aligned}$$

Defintions of rubi rules used

rule 16

$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\text{tan}[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4100

$$\text{Int}[(a_)+(b_)*\text{tan}[(e_)+(f_)*(x_)]^m, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], b*\text{Tan}[e + f*x], x] \text{ ; FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$$

rule 4109

$$\begin{aligned}
& \text{Int}[(A_)+(B_)*\text{tan}[(e_)+(f_)*(x_)] + (C_)*\text{tan}[(e_)+(f_)*(x_)]^2 \\
&)/((a_)+(b_)*\text{tan}[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a \\
& *C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 \\
& + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(\\
& a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) \text{ ; FreeQ}[\{a, b, e, f, A, B, C\}, x] \ \&\ \\
& \ \& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\ \ \text{NeQ}[a^2 + b^2, 0] \ \&\ \ \text{NeQ}[A*b - a*B - b*C \\
& , 0]
\end{aligned}$$

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4128

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 552, normalized size of antiderivative = 1.33

method	result
derivativedivides	$\frac{\tan(fx+e)C d^2}{b^2} + \frac{(2A a^2 cd - 2A ab c^2 + 2A ab d^2 - 2A b^2 cd + B a^2 c^2 - B a^2 d^2 + 4B abcd - B b^2 c^2 + B b^2 d^2 - 2C a^2 cd + 2C ab c^2 - 2C ab d^2)}{2}$
default	$\frac{\tan(fx+e)C d^2}{b^2} + \frac{(2A a^2 cd - 2A ab c^2 + 2A ab d^2 - 2A b^2 cd + B a^2 c^2 - B a^2 d^2 + 4B abcd - B b^2 c^2 + B b^2 d^2 - 2C a^2 cd + 2C ab c^2 - 2C ab d^2)}{2}$
norman	$\frac{a(A a^2 c^2 - A a^2 d^2 + 4A abcd - A b^2 c^2 + A b^2 d^2 - 2B a^2 cd + 2B ab c^2 - 2B ab d^2 + 2B b^2 cd - C a^2 c^2 + a^2 C d^2 - 4C abcd + C b^2 c^2 - b^2 d^2 C)}{a^4 + 2b^2 a^2 + b^4} x$
parallelrisc	Expression too large to display
risc	Expression too large to display

input `int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2, x,method=_RETURNVERBOSE)`

output
$$\frac{1}{f} \left(\frac{\tan(fx+e) C d^2}{b^2} + \frac{1}{(a^2+b^2)^2} \left(\frac{1}{2} (2Aa^2cd - 2Aab^2c^2 + 2Aab^2d^2 - 2Aab^2cd + B^2a^2c^2 - B^2a^2d^2 + 4Bab^2cd - B^2b^2c^2 + B^2b^2d^2 - 2C^2a^2cd + 2C^2ab^2c^2 - 2C^2ab^2d^2 + 2C^2b^2cd) \ln(1+\tan(fx+e)^2) + (Aa^2c^2 - Aa^2d^2 + 4Aab^2cd - Ab^2c^2 + Ab^2d^2 - 2B^2a^2cd + 2B^2ab^2c^2 - 2B^2ab^2d^2 + 2B^2b^2cd - Ca^2c^2 + Ca^2d^2 - 4Cab^2cd + C^2b^2c^2 - C^2b^2d^2) \arctan(\tan(fx+e)) \right) - \frac{1}{b^3} (Aa^2b^2d^2 - 2Aab^3cd + Ab^4c^2 - Ba^3b^2d^2 + 2B^2a^2b^2cd - Ba^2b^3c^2 + Ca^4d^2 - 2Ca^3b^2cd + Ca^2b^2c^2) / (a^2+b^2) / (a+b\tan(fx+e)) + (-2Aa^2b^3cd + 2Aa^2b^4c^2 - 2Aa^2b^4d^2 + 2Aa^2b^5cd + B^2a^4b^2d^2 - Ba^2b^3c^2 + 3B^2a^2b^3d^2 - 4B^2a^2b^4cd + B^2b^5c^2 - 2Ca^5d^2 + 2Ca^4b^2cd - 4Ca^3b^2d^2 + 6Ca^2b^3cd - 2Ca^2b^4c^2) / b^3 / (a^2+b^2)^2 \ln(a+b\tan(fx+e)) \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 964 vs. $2(413) = 826$.

Time = 0.39 (sec) , antiderivative size = 964, normalized size of antiderivative = 2.32

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

```

1/2*(2*(C*a^4*b^2 + 2*C*a^2*b^4 + C*b^6)*d^2*tan(f*x + e)^2 - 2*(C*a^2*b^4
- B*a*b^5 + A*b^6)*c^2 + 4*(C*a^3*b^3 - B*a^2*b^4 + A*a*b^5)*c*d - 2*(C*a
^4*b^2 - B*a^3*b^3 + A*a^2*b^4)*d^2 + 2*((A - C)*a^3*b^3 + 2*B*a^2*b^4 -
(A - C)*a*b^5)*c^2 - 2*(B*a^3*b^3 - 2*(A - C)*a^2*b^4 - B*a*b^5)*c*d - ((A
- C)*a^3*b^3 + 2*B*a^2*b^4 - (A - C)*a*b^5)*d^2)*f*x - ((B*a^3*b^3 - 2*(A
- C)*a^2*b^4 - B*a*b^5)*c^2 - 2*(C*a^5*b - (A - 3*C)*a^3*b^3 - 2*B*a^2*b^
4 + A*a*b^5)*c*d + (2*C*a^6 - B*a^5*b + 4*C*a^4*b^2 - 3*B*a^3*b^3 + 2*A*a^
2*b^4)*d^2 + ((B*a^2*b^4 - 2*(A - C)*a*b^5 - B*b^6)*c^2 - 2*(C*a^4*b^2 - (
A - 3*C)*a^2*b^4 - 2*B*a*b^5 + A*b^6)*c*d + (2*C*a^5*b - B*a^4*b^2 + 4*C*a
^3*b^3 - 3*B*a^2*b^4 + 2*A*a*b^5)*d^2)*tan(f*x + e))*log((b^2*tan(f*x + e)
^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) - (2*(C*a^5*b + 2*C*a
^3*b^3 + C*a*b^5)*c*d - (2*C*a^6 - B*a^5*b + 4*C*a^4*b^2 - 2*B*a^3*b^3 + 2
*C*a^2*b^4 - B*a*b^5)*d^2 + (2*(C*a^4*b^2 + 2*C*a^2*b^4 + C*b^6)*c*d - (2*
C*a^5*b - B*a^4*b^2 + 4*C*a^3*b^3 - 2*B*a^2*b^4 + 2*C*a*b^5 - B*b^6)*d^2)*
tan(f*x + e))*log(1/(tan(f*x + e)^2 + 1)) + 2*((C*a^3*b^3 - B*a^2*b^4 + A*
a*b^5)*c^2 - 2*(C*a^4*b^2 - B*a^3*b^3 + A*a^2*b^4)*c*d + (2*C*a^5*b - B*a^
4*b^2 + (A + 2*C)*a^3*b^3 + C*a*b^5)*d^2 + (((A - C)*a^2*b^4 + 2*B*a*b^5 -
(A - C)*b^6)*c^2 - 2*(B*a^2*b^4 - 2*(A - C)*a*b^5 - B*b^6)*c*d - ((A - C)
*a^2*b^4 + 2*B*a*b^5 - (A - C)*b^6)*d^2)*f*x)*tan(f*x + e))/((a^4*b^4 + 2*
a^2*b^6 + b^8)*f*tan(f*x + e) + (a^5*b^3 + 2*a^3*b^5 + a*b^7)*f)

```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.67 (sec) , antiderivative size = 16225, normalized size of antiderivative = 39.10

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*
x+e))**2,x)

```

output

```
Piecewise((zoo*x*(c + d*tan(e))**2*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2,
Eq(a, 0) & Eq(b, 0) & Eq(f, 0)), ((A*c**2*x + A*c*d*log(tan(e + f*x)**2 +
1)/f - A*d**2*x + A*d**2*tan(e + f*x)/f + B*c**2*log(tan(e + f*x)**2 + 1)
/(2*f) - 2*B*c*d*x + 2*B*c*d*tan(e + f*x)/f - B*d**2*log(tan(e + f*x)**2 +
1)/(2*f) + B*d**2*tan(e + f*x)**2/(2*f) - C*c**2*x + C*c**2*tan(e + f*x)/
f - C*c*d*log(tan(e + f*x)**2 + 1)/f + C*c*d*tan(e + f*x)**2/f + C*d**2*x
+ C*d**2*tan(e + f*x)**3/(3*f) - C*d**2*tan(e + f*x)/f)/a**2, Eq(b, 0)), (
-A*c**2*f*x*tan(e + f*x)**2/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e +
f*x) - 4*b**2*f) + 2*I*A*c**2*f*x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2
- 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + A*c**2*f*x/(4*b**2*f*tan(e + f*x)*
*2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) - A*c**2*tan(e + f*x)/(4*b**2*f*t
an(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + 2*I*A*c**2/(4*b**2*
f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + 2*I*A*c*d*f*x*ta
n(e + f*x)**2/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2
*f) + 4*A*c*d*f*x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(
e + f*x) - 4*b**2*f) - 2*I*A*c*d*f*x/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*
f*tan(e + f*x) - 4*b**2*f) + 2*I*A*c*d*tan(e + f*x)/(4*b**2*f*tan(e + f*x)
**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + A*d**2*f*x*tan(e + f*x)**2/(4*
b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) - 2*I*A*d**2*
f*x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - ...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 496, normalized size of antiderivative = 1.20

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{2Cd^2 \tan(fx+e)}{b^2} + \frac{2(((A-C)a^2+2Bab-(A-C)b^2)c^2-2(Ba^2-2(A-C)ab-Bb^2)cd-((A-C)a^2+2Bab-(A-C)b^2)d^2)(fx+e)}{a^4+2a^2b^2+b^4} - \frac{2((Ba^2-2(A-C)ab-Bb^2)c^2-2(Ba^2-2(A-C)ab-Bb^2)cd-((A-C)a^2+2Bab-(A-C)b^2)d^2)(fx+e)}{a^4+2a^2b^2+b^4}$$

input

```
integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e))^2,x, algorithm="maxima")
```

output

```

1/2*(2*C*d^2*tan(f*x + e)/b^2 + 2*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c
^2 - 2*(B*a^2 - 2*(A - C)*a*b - B*b^2)*c*d - ((A - C)*a^2 + 2*B*a*b - (A -
C)*b^2)*d^2)*(f*x + e)/(a^4 + 2*a^2*b^2 + b^4) - 2*((B*a^2*b^3 - 2*(A - C
)*a*b^4 - B*b^5)*c^2 - 2*(C*a^4*b - (A - 3*C)*a^2*b^3 - 2*B*a*b^4 + A*b^5)
*c*d + (2*C*a^5 - B*a^4*b + 4*C*a^3*b^2 - 3*B*a^2*b^3 + 2*A*a*b^4)*d^2)*lo
g(b*tan(f*x + e) + a)/(a^4*b^3 + 2*a^2*b^5 + b^7) + ((B*a^2 - 2*(A - C)*a*
b - B*b^2)*c^2 + 2*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c*d - (B*a^2 - 2*
(A - C)*a*b - B*b^2)*d^2)*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4)
- 2*((C*a^2*b^2 - B*a*b^3 + A*b^4)*c^2 - 2*(C*a^3*b - B*a^2*b^2 + A*a*b^3)
*c*d + (C*a^4 - B*a^3*b + A*a^2*b^2)*d^2)/(a^3*b^3 + a*b^5 + (a^2*b^4 + b
6)*tan(f*x + e))/f

```

Giac [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 701, normalized size of antiderivative = 1.69

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{(Aa^2c^2 - Ca^2c^2 + 2Babc^2 - Ab^2c^2 + Cb^2c^2 - 2Ba^2cd + 4Aabcd - 4Cabcd + 2Bb^2cd - Aa^2d^2 + Ca^2d^2 + Cb^2d^2)}{a^4f + 2a^2b^2f + b^4f}$$

$$+ \frac{(Ba^2c^2 - 2Aabc^2 + 2Cabc^2 - Bb^2c^2 + 2Aa^2cd - 2Ca^2cd + 4Babcd - 2Ab^2cd + 2Cb^2cd - Ba^2d^2 + Cb^2d^2)}{2(a^4f + 2a^2b^2f + b^4f)}$$

$$- \frac{(Ba^2b^3c^2 - 2Aab^4c^2 + 2Cab^4c^2 - Bb^5c^2 - 2Ca^4bcd + 2Aa^2b^3cd - 6Ca^2b^3cd + 4Bab^4cd - 2Ab^5cd - 2Ba^2b^3d^2 + 2Aab^4d^2 - 2Cab^4d^2 - Bb^5d^2)}{a^4b^3f + 2a^2b^5f + b^7f}$$

$$+ \frac{Cd^2 \tan(fx + e)}{b^2f}$$

$$- \frac{Ca^4b^2c^2 - Ba^3b^3c^2 + Aa^2b^4c^2 + Ca^2b^4c^2 - Bab^5c^2 + Ab^6c^2 - 2Ca^5bcd + 2Ba^4b^2cd - 2Aa^3b^3cd - 2Ba^2b^4d^2 + 2Aab^4d^2 - 2Cab^4d^2 - Bb^5d^2}{(a^2 + b^2)^2(b \tan(fx + e) + a)}$$

input

```

integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e))^2,x, algorithm="giac")

```

output

```
(A*a^2*c^2 - C*a^2*c^2 + 2*B*a*b*c^2 - A*b^2*c^2 + C*b^2*c^2 - 2*B*a^2*c*d
+ 4*A*a*b*c*d - 4*C*a*b*c*d + 2*B*b^2*c*d - A*a^2*d^2 + C*a^2*d^2 - 2*B*a
*b*d^2 + A*b^2*d^2 - C*b^2*d^2)*(f*x + e)/(a^4*f + 2*a^2*b^2*f + b^4*f) +
1/2*(B*a^2*c^2 - 2*A*a*b*c^2 + 2*C*a*b*c^2 - B*b^2*c^2 + 2*A*a^2*c*d - 2*C
*a^2*c*d + 4*B*a*b*c*d - 2*A*b^2*c*d + 2*C*b^2*c*d - B*a^2*d^2 + 2*A*a*b*d
^2 - 2*C*a*b*d^2 + B*b^2*d^2)*log(tan(f*x + e)^2 + 1)/(a^4*f + 2*a^2*b^2*f
+ b^4*f) - (B*a^2*b^3*c^2 - 2*A*a*b^4*c^2 + 2*C*a*b^4*c^2 - B*b^5*c^2 - 2
*C*a^4*b*c*d + 2*A*a^2*b^3*c*d - 6*C*a^2*b^3*c*d + 4*B*a*b^4*c*d - 2*A*b^5
*c*d + 2*C*a^5*d^2 - B*a^4*b*d^2 + 4*C*a^3*b^2*d^2 - 3*B*a^2*b^3*d^2 + 2*A
*a*b^4*d^2)*log(abs(b*tan(f*x + e) + a))/(a^4*b^3*f + 2*a^2*b^5*f + b^7*f)
+ C*d^2*tan(f*x + e)/(b^2*f) - (C*a^4*b^2*c^2 - B*a^3*b^3*c^2 + A*a^2*b^4
*c^2 + C*a^2*b^4*c^2 - B*a*b^5*c^2 + A*b^6*c^2 - 2*C*a^5*b*c*d + 2*B*a^4*b
^2*c*d - 2*A*a^3*b^3*c*d - 2*C*a^3*b^3*c*d + 2*B*a^2*b^4*c*d - 2*A*a*b^5*c
*d + C*a^6*d^2 - B*a^5*b*d^2 + A*a^4*b^2*d^2 + C*a^4*b^2*d^2 - B*a^3*b^3*d
^2 + A*a^2*b^4*d^2)/((a^2 + b^2)^2*(b*tan(f*x + e) + a)*b^3*f)
```

Mupad [B] (verification not implemented)

Time = 29.77 (sec) , antiderivative size = 3958, normalized size of antiderivative = 9.54

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int(((c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a +
b*tan(e + f*x))^2,x)
```


output

```
(log((2*C^2*a^5*d^4 + 4*C^2*a^3*b^2*d^4 - 2*C^2*a^5*c^2*d^2 - A*B*b^5*c^4
- 2*A*C*a^5*d^4 + B*C*b^5*c^4 - A^2*a*b^4*c^4 - A^2*a*b^4*d^4 + B^2*a*b^4*
c^4 + B^2*a*b^4*d^4 - C^2*a*b^4*c^4 + 2*A^2*b^5*c*d^3 - 2*A^2*b^5*c^3*d +
C^2*a*b^4*d^4 + 2*B^2*b^5*c^3*d - 4*C^2*a^3*b^2*c^2*d^2 + A*B*a^2*b^3*c^4
+ 3*A*B*a^2*b^3*d^4 - 4*A*C*a^3*b^2*d^4 - B*C*a^2*b^3*c^4 + 5*A*B*b^5*c^2*
d^2 + 2*A*C*a^5*c^2*d^2 - 3*B*C*a^2*b^3*d^4 - B*C*b^5*c^2*d^2 + 2*B^2*a^4*
b*c*d^3 - 2*C^2*a^4*b*c*d^3 + 2*C^2*a^4*b*c^3*d + 6*A^2*a*b^4*c^2*d^2 - 2*
A^2*a^2*b^3*c*d^3 + 2*A^2*a^2*b^3*c^3*d - 6*B^2*a*b^4*c^2*d^2 + 6*B^2*a^2*
b^3*c*d^3 - 2*B^2*a^2*b^3*c^3*d + 4*C^2*a*b^4*c^2*d^2 - 6*C^2*a^2*b^3*c*d^
3 + 6*C^2*a^2*b^3*c^3*d + A*B*a^4*b*d^4 + 2*A*C*a*b^4*c^4 - B*C*a^4*b*d^4
- 2*A*C*b^5*c*d^3 + 2*A*C*b^5*c^3*d - 4*B*C*a^5*c*d^3 - 8*A*B*a*b^4*c*d^3
+ 8*A*B*a*b^4*c^3*d + 2*A*C*a^4*b*c*d^3 - 2*A*C*a^4*b*c^3*d + 4*B*C*a*b^4*
c*d^3 - 8*B*C*a*b^4*c^3*d - A*B*a^4*b*c^2*d^2 - 10*A*C*a*b^4*c^2*d^2 + 8*A
*C*a^2*b^3*c*d^3 - 8*A*C*a^2*b^3*c^3*d - 8*B*C*a^3*b^2*c*d^3 + 5*B*C*a^4*b
*c^2*d^2 - 8*A*B*a^2*b^3*c^2*d^2 + 4*A*C*a^3*b^2*c^2*d^2 + 16*B*C*a^2*b^3*
c^2*d^2)/(b^2*(a^2 + b^2)^2) + ((c*i1 + d)^2*((tan(e + f*x))*(3*B*b^5*c^2 -
5*B*b^5*d^2 - 4*C*a^5*d^2 + 6*A*b^5*c*d - 10*C*b^5*c*d + 4*A*a*b^4*c^2 -
4*A*a*b^4*d^2 + 2*B*a^4*b*d^2 - 4*C*a*b^4*c^2 + 8*C*a*b^4*d^2 - B*a^2*b^3*
c^2 + B*a^2*b^3*d^2 - 8*B*a*b^4*c*d + 4*C*a^4*b*c*d - 2*A*a^2*b^3*c*d + 2*
C*a^2*b^3*c*d))/(b^2*(a^2 + b^2)) - (A*b^2*d^2 - A*b^2*c^2 - 8*C*a^2*d^...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1586, normalized size of antiderivative = 3.82

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int(((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,
x)
```

output

```
(2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**4*c*d - log(tan(e + f*x)*
**2 + 1)*tan(e + f*x)*a**2*b**5*c**2 + log(tan(e + f*x)**2 + 1)*tan(e + f*x
)*a**2*b**5*d**2 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**4*c**2*
d + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**6*c*d + 2*log(tan(e + f*x
)**2 + 1)*tan(e + f*x)*a*b**5*c**3 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*
x)*a*b**5*c*d**2 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**7*c**2 + log(t
an(e + f*x)**2 + 1)*tan(e + f*x)*b**7*d**2 + 2*log(tan(e + f*x)**2 + 1)*ta
n(e + f*x)*b**6*c**2*d + 2*log(tan(e + f*x)**2 + 1)*a**4*b**3*c*d - log(ta
n(e + f*x)**2 + 1)*a**3*b**4*c**2 + log(tan(e + f*x)**2 + 1)*a**3*b**4*d**
2 - 2*log(tan(e + f*x)**2 + 1)*a**3*b**3*c**2*d + 2*log(tan(e + f*x)**2 +
1)*a**2*b**5*c*d + 2*log(tan(e + f*x)**2 + 1)*a**2*b**4*c**3 - 2*log(tan(e
 + f*x)**2 + 1)*a**2*b**4*c*d**2 - log(tan(e + f*x)**2 + 1)*a*b**6*c**2 +
log(tan(e + f*x)**2 + 1)*a*b**6*d**2 + 2*log(tan(e + f*x)**2 + 1)*a*b**5*c
**2*d - 4*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**5*b*c*d**2 + 2*log(tan(e
 + f*x)*b + a)*tan(e + f*x)*a**4*b**3*d**2 + 4*log(tan(e + f*x)*b + a)*tan
(e + f*x)*a**4*b**2*c**2*d - 4*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**3*b
**4*c*d - 8*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**3*b**3*c*d**2 + 2*log(
tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b**5*c**2 + 2*log(tan(e + f*x)*b + a
)*tan(e + f*x)*a**2*b**5*d**2 + 12*log(tan(e + f*x)*b + a)*tan(e + f*x)*a*
**2*b**4*c**2*d - 4*log(tan(e + f*x)*b + a)*tan(e + f*x)*a*b**6*c*d - 4*...
```

3.63
$$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

Optimal result	686
Mathematica [C] (verified)	687
Rubi [A] (verified)	688
Maple [A] (verified)	692
Fricas [B] (verification not implemented)	693
Sympy [F(-2)]	694
Maxima [A] (verification not implemented)	695
Giac [B] (verification not implemented)	695
Mupad [B] (verification not implemented)	696
Reduce [B] (verification not implemented)	697

Optimal result

Integrand size = 45, antiderivative size = 597

$$\int \frac{(c+d \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx =$$

$$\frac{(a^3(c^2C+2Bcd-Cd^2-A(c^2-d^2))-3ab^2(c^2C+2Bcd-Cd^2-A(c^2-d^2))-3a^2b(2c(A-C)d+E)}{(a^2+b^2)^3}$$

$$-\frac{(3a^2b(c^2C+2Bcd-Cd^2-A(c^2-d^2))-b^3(c^2C+2Bcd-Cd^2-A(c^2-d^2))+a^3(2c(A-C)d+E)}{(a^2+b^2)^3 f}$$

$$+\frac{(a^6Cd^2+3a^4b^2Cd^2-3a^2b^4(c^2C+2Bcd-2Cd^2-A(c^2-d^2))+b^6(c(cC+2Bd)-A(c^2-d^2))-a^5)}{b^3(a^2+b^2)^3 f}$$

$$-\frac{(bc-ad)(a^4Cd+b^4(Bc+Ad)+2ab^3(Ac-cC-Bd)-a^2b^2(Bc+(A-3C)d))}{b^3(a^2+b^2)^2 f(a+b \tan(e+fx))}$$

$$-\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^2}{2b(a^2+b^2) f(a+b \tan(e+fx))^2}$$

output

$$\begin{aligned}
& -(a^3(c^2C+2Bcd-Cd^2-A(c^2-d^2))-3ab^2(c^2C+2Bcd-Cd^2-A(c^2-d^2))-3a^2b(2c(A-C)d+B(c^2-d^2))+b^3(2c(A-C)d+B(c^2-d^2))) \\
& / (a^2+b^2)^3 - (3a^2b(c^2C+2Bcd-Cd^2-A(c^2-d^2))-b^3(c^2C+2Bcd-Cd^2-A(c^2-d^2))+a^3(2c(A-C)d+B(c^2-d^2))-3ab^2(2c(A-C)d+B(c^2-d^2))) \\
& * \ln(\cos(fx+e)) / (a^2+b^2)^3 / f + (a^6Cd^2+3a^4b^2Cd^2-3a^2b^4(c^2C+2Bcd-2Cd^2-A(c^2-d^2))+b^6(c(2Bd+Cc)-A(c^2-d^2))-a^3 \\
& * b^3(2c(A-C)d+B(c^2-d^2))+3ab^5(2c(A-C)d+B(c^2-d^2))) * \ln(a+b \tan(fx+e)) / b^3 / (a^2+b^2)^3 / f - (-ad+bc) * (a^4Cd+b^4(A+Bc)+2ab^3(Ac-Bd-Cc) \\
& - a^2b^2(Bc+(A-3C)d)) / b^3 / (a^2+b^2)^2 / f / (a+b \tan(fx+e)) - 1/2 * (Ab^2-a(Bb-Ca)) * (c+d \tan(fx+e))^2 / b / (a^2+b^2) / f / (a+b \tan(fx+e))^2
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.27 (sec) , antiderivative size = 403, normalized size of antiderivative = 0.68

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

$$= \frac{(a - ib)^3 (-iA + B + iC)(c + id)^2 \log(i - \tan(e + fx)) + (a + ib)^3 (iA + B - iC)(c - id)^2 \log(i + \tan(e + fx))}{(a^2 + b^2)^3}$$

input

```
Integrate[((c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)) / (a + b*Tan[e + f*x])^3, x]
```

output

$$\begin{aligned}
& ((a - I*b)^3 * ((-I)*A + B + I*C) * (c + I*d)^2 * \text{Log}[I - \text{Tan}[e + f*x]] + (a + I*b)^3 * (I*A + B - I*C) * (c - I*d)^2 * \text{Log}[I + \text{Tan}[e + f*x]] + (2*(a^6*C*d^2 + 3*a^4*b^2*C*d^2 + 3*a*b^5*(2*c*(A - C)*d + B*(c^2 - d^2)) - 3*a^2*b^4*(c^2*C + 2*B*c*d - 2*C*d^2 + A*(-c^2 + d^2)) + b^6*(c*(c*C + 2*B*d) + A*(-c^2 + d^2)) + a^3*b^3*(2*c*(-A + C)*d + B*(-c^2 + d^2))) * \text{Log}[a + b*\text{Tan}[e + f*x]]) / b^3 - ((a^2 + b^2)^2 * (A*b^2 + a*(-(b*B) + a*C)) * (b*c - a*d)^2) / (b^3 * (a + b*\text{Tan}[e + f*x])^2) + (2*(a^2 + b^2) * (-(b*c) + a*d) * (-(a^3*b*B*d) + 2*a^4*C*d + b^4*(B*c + 2*A*d) + a*b^3*(2*A*c - 2*c*C - 3*B*d) + a^2*b^2*(-(B*c) + 4*C*d))) / (b^3 * (a + b*\text{Tan}[e + f*x])) / (2*(a^2 + b^2)^3 * f)
\end{aligned}$$

Rubi [A] (verified)

Time = 4.06 (sec) , antiderivative size = 631, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.244$, Rules used = {3042, 4128, 27, 3042, 4118, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{2(c + d \tan(e + fx))((a^2 + b^2)Cd \tan^2(e + fx) - b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(bc - ad) + Ab(ac + bd))}{(a + b \tan(e + fx))^2} dx$$

$$\frac{2b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^2} \frac{1}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 27

$$\int \frac{(c + d \tan(e + fx))((a^2 + b^2)Cd \tan^2(e + fx) - b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(bc - ad) + Ab(ac + bd))}{(a + b \tan(e + fx))^2} dx$$

$$\frac{b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^2} \frac{1}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))((a^2 + b^2)Cd \tan(e + fx)^2 - b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(bc - ad) + Ab(ac + bd))}{(a + b \tan(e + fx))^2} dx$$

$$\frac{b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^2} \frac{1}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 4118

$$\int \frac{Cd^2a^4 - b^2(Cc^2 + 2Bdc - 3Cd^2 - A(c^2 - d^2))a^2 + 2b^3(2c(A-C)d + B(c^2 - d^2))a + (a^2 + b^2)^2 Cd^2 \tan^2(e + fx) + b^4(c(cC + 2Bd) - A(c^2 - d^2)) + b^2((2c(A-C)d + B(c^2 - d^2))a + (a^2 + b^2)^2 Cd^2 \tan^2(e + fx) + b^4(c(cC + 2Bd) - A(c^2 - d^2)))}{\frac{a + b \tan(e + fx)}{b(a^2 + b^2)}} dx$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{Cd^2a^4 - b^2(Cc^2 + 2Bdc - 3Cd^2 - A(c^2 - d^2))a^2 + 2b^3(2c(A-C)d + B(c^2 - d^2))a + (a^2 + b^2)^2 Cd^2 \tan^2(e + fx) + b^4(c(cC + 2Bd) - A(c^2 - d^2)) + b^2((2c(A-C)d + B(c^2 - d^2))a + (a^2 + b^2)^2 Cd^2 \tan^2(e + fx) + b^4(c(cC + 2Bd) - A(c^2 - d^2)))}{\frac{a + b \tan(e + fx)}{b(a^2 + b^2)}} dx$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 4109

$$\frac{b^2(a^3(2cd(A-C) + B(c^2 - d^2)) + 3a^2b(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) - 3ab^2(2cd(A-C) + B(c^2 - d^2)) - b^3(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2)) \int \tan(e + fx) dx}{a^2 + b^2}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3042

$$\frac{b^2(a^3(2cd(A-C) + B(c^2 - d^2)) + 3a^2b(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2) - 3ab^2(2cd(A-C) + B(c^2 - d^2)) - b^3(-A(c^2 - d^2) + 2Bcd + c^2C - Cd^2)) \int \tan(e + fx) dx}{a^2 + b^2}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3956

$$\frac{(a^6Cd^2 + 3a^4b^2Cd^2 - a^3b^3(2cd(A-C) + B(c^2 - d^2)) - 3a^2b^4(-A(c^2 - d^2) + 2Bcd + c^2C - 2Cd^2) + 3ab^5(2cd(A-C) + B(c^2 - d^2)) + b^6(c(2Bd + cC) - A(c^2 - d^2))) \int \tan(e + fx) dx}{a^2 + b^2}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 4100

$$\frac{(a^6 C d^2 + 3 a^4 b^2 C d^2 - a^3 b^3 (2 c d (A - C) + B (c^2 - d^2)) - 3 a^2 b^4 (-A (c^2 - d^2) + 2 B c d + c^2 C - 2 C d^2) + 3 a b^5 (2 c d (A - C) + B (c^2 - d^2)) + b^6 (c (2 B d + c C) - A (c^2 - d^2)))}{b f (a^2 + b^2)}$$

$$\frac{(A b^2 - a (b B - a C)) (c + d \tan(e + f x))^2}{2 b f (a^2 + b^2) (a + b \tan(e + f x))^2}$$

↓ 16

$$-\frac{b^2 \log(\cos(e + f x)) (a^3 (2 c d (A - C) + B (c^2 - d^2)) + 3 a^2 b (-A (c^2 - d^2) + 2 B c d + c^2 C - C d^2) - 3 a b^2 (2 c d (A - C) + B (c^2 - d^2)) - b^3 (-A (c^2 - d^2) + 2 B c d + c^2 C - C d^2))}{f (a^2 + b^2)}$$

$$\frac{(A b^2 - a (b B - a C)) (c + d \tan(e + f x))^2}{2 b f (a^2 + b^2) (a + b \tan(e + f x))^2}$$

input

```
Int[((c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

```
-1/2*((A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^2)/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^2) + (((-((b^2*(a^3*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - 3*a*b^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - 3*a^2*b*(2*c*(A - C)*d + B*(c^2 - d^2)) + b^3*(2*c*(A - C)*d + B*(c^2 - d^2))))*x)/(a^2 + b^2)) - (b^2*(3*a^2*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b^3*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + a^3*(2*c*(A - C)*d + B*(c^2 - d^2)) - 3*a*b^2*(2*c*(A - C)*d + B*(c^2 - d^2)))*Log[Cos[e + f*x]]/((a^2 + b^2)*f) + ((a^6*C*d^2 + 3*a^4*b^2*C*d^2 - 3*a^2*b^4*(c^2*C + 2*B*c*d - 2*C*d^2 - A*(c^2 - d^2)) + b^6*(c*(c*C + 2*B*d) - A*(c^2 - d^2)) - a^3*b^3*(2*c*(A - C)*d + B*(c^2 - d^2)) + 3*a*b^5*(2*c*(A - C)*d + B*(c^2 - d^2)))*Log[a + b*Tan[e + f*x]]/(b*(a^2 + b^2)*f))/(b*(a^2 + b^2)) - ((b*c - a*d)*(a^4*C*d + b^4*(B*c + A*d) + 2*a*b^3*(A*c - c*C - B*d) - a^2*b^2*(B*c + (A - 3*C)*d)))/(b^2*(a^2 + b^2)*f*(a + b*Tan[e + f*x]))/(b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4100 $\text{Int}[((a_)+(b_)*\tan[(e_)+(f_)*(x_)]^{(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[((A_)+(B_)*\tan[(e_)+(f_)*(x_)] + (C_)*\tan[(e_)+(f_)*(x_)]^2)/((a_)+(b_)*\tan[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A*b - a*B - b*C, 0]$

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 865, normalized size of antiderivative = 1.45

method	result
derivativedivides	$\frac{(2Aa^3cd - 3Aa^2b^2c^2 + 3Aa^2bd^2 - 6Aab^2cd + Ab^3c^2 - Ab^3d^2 + Ba^3c^2 - Ba^3d^2 + 6Ba^2bcd - 3Ba^2c^2 + 3Ba^2d^2 - 2Bb^3cd - 2Ca^3cd)}{2}$
default	$\frac{(2Aa^3cd - 3Aa^2b^2c^2 + 3Aa^2bd^2 - 6Aab^2cd + Ab^3c^2 - Ab^3d^2 + Ba^3c^2 - Ba^3d^2 + 6Ba^2bcd - 3Ba^2c^2 + 3Ba^2d^2 - 2Bb^3cd - 2Ca^3cd)}{2}$
norman	Expression too large to display
risch	Expression too large to display
parallelrisc	Expression too large to display

input `int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3, x,method=_RETURNVERBOSE)`

output `1/f*(1/(a^2+b^2)^3*(1/2*(2*A*a^3*c*d-3*A*a^2*b*c^2+3*A*a^2*b*d^2-6*A*a*b^2*c*d+A*b^3*c^2-A*b^3*d^2+B*a^3*c^2-B*a^3*d^2+6*B*a^2*b*c*d-3*B*a*b^2*c^2+3*B*a*b^2*d^2-2*B*b^3*c*d-2*C*a^3*c*d+3*C*a^2*b*c^2-3*C*a^2*b*d^2+6*C*a*b^2*c*d-C*b^3*c^2+C*b^3*d^2)*ln(1+tan(f*x+e)^2)+(A*a^3*c^2-A*a^3*d^2+6*A*a^2*b*c*d-3*A*a*b^2*c^2+3*A*a*b^2*d^2-2*A*b^3*c*d-2*B*a^3*c*d+3*B*a^2*b*c^2-3*B*a^2*b*d^2+6*B*a*b^2*c*d-B*b^3*c^2+B*b^3*d^2-C*a^3*c^2+C*a^3*d^2-6*C*a^2*b*c*d+3*C*a*b^2*c^2-3*C*a*b^2*d^2+2*C*b^3*c*d)*arctan(tan(f*x+e)))-1/2*(A*a^2*b^2*d^2-2*A*a*b^3*c*d+A*b^4*c^2-B*a^3*b*d^2+2*B*a^2*b^2*c*d-B*a*b^3*c^2+C*a^4*d^2-2*C*a^3*b*c*d+C*a^2*b^2*c^2)/b^3/(a^2+b^2)/(a+b*tan(f*x+e))^2-(-2*A*a^2*b^3*c*d+2*A*a*b^4*c^2-2*A*a*b^4*d^2+2*A*b^5*c*d+B*a^4*b*d^2-B*a^2*b^3*c^2+3*B*a^2*b^3*d^2-4*B*a*b^4*c*d+B*b^5*c^2-2*C*a^5*d^2+2*C*a^4*b*c*d-4*C*a^3*b^2*d^2+6*C*a^2*b^3*c*d-2*C*a*b^4*c^2)/b^3/(a^2+b^2)^2/(a+b*tan(f*x+e))+1/(a^2+b^2)^3*(-2*A*a^3*b^3*c*d+3*A*a^2*b^4*c^2-3*A*a^2*b^4*d^2+6*A*a*b^5*c*d-A*b^6*c^2+A*b^6*d^2-B*a^3*b^3*c^2+B*a^3*b^3*d^2-6*B*a^2*b^4*c*d+3*B*a*b^5*c^2-3*B*a*b^5*d^2+2*B*b^6*c*d+C*a^6*d^2+3*C*a^4*b^2*d^2+2*C*a^3*b^3*c*d-3*C*a^2*b^4*c^2+6*C*a^2*b^4*d^2-6*C*a*b^5*c*d+C*b^6*c^2)/b^3*ln(a+b*tan(f*x+e))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1699 vs. 2(595) = 1190.

Time = 0.50 (sec) , antiderivative size = 1699, normalized size of antiderivative = 2.85

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

output

```

-1/2*((3*C*a^4*b^4 - 5*B*a^3*b^5 + (7*A - 3*C)*a^2*b^6 + B*a*b^7 + A*b^8)*
c^2 - 2*(C*a^5*b^3 - 3*B*a^4*b^4 + 5*(A - C)*a^3*b^5 + 3*B*a^2*b^6 - A*a*b
^7)*c*d - (C*a^6*b^2 + B*a^5*b^3 - (3*A - 7*C)*a^4*b^4 - 5*B*a^3*b^5 + 3*A
*a^2*b^6)*d^2 - 2*(((A - C)*a^5*b^3 + 3*B*a^4*b^4 - 3*(A - C)*a^3*b^5 - B*
a^2*b^6)*c^2 - 2*(B*a^5*b^3 - 3*(A - C)*a^4*b^4 - 3*B*a^3*b^5 + (A - C)*a
^2*b^6)*c*d - ((A - C)*a^5*b^3 + 3*B*a^4*b^4 - 3*(A - C)*a^3*b^5 - B*a^2*b
^6)*d^2)*f*x - ((C*a^4*b^4 - 3*B*a^3*b^5 + 5*(A - C)*a^2*b^6 + 3*B*a*b^7 -
A*b^8)*c^2 + 2*(C*a^5*b^3 + B*a^4*b^4 - (3*A - 7*C)*a^3*b^5 - 5*B*a^2*b^6
+ 3*A*a*b^7)*c*d - (3*C*a^6*b^2 - B*a^5*b^3 - (A - 9*C)*a^4*b^4 - 7*B*a^3*
b^5 + 5*A*a^2*b^6)*d^2 + 2*(((A - C)*a^3*b^5 + 3*B*a^2*b^6 - 3*(A - C)*a*b
^7 - B*b^8)*c^2 - 2*(B*a^3*b^5 - 3*(A - C)*a^2*b^6 - 3*B*a*b^7 + (A - C)*b
^8)*c*d - ((A - C)*a^3*b^5 + 3*B*a^2*b^6 - 3*(A - C)*a*b^7 - B*b^8)*d^2)*f
*x)*tan(f*x + e)^2 + ((B*a^5*b^3 - 3*(A - C)*a^4*b^4 - 3*B*a^3*b^5 + (A -
C)*a^2*b^6)*c^2 + 2*(((A - C)*a^5*b^3 + 3*B*a^4*b^4 - 3*(A - C)*a^3*b^5 - B
*a^2*b^6)*c*d - (C*a^8 + 3*C*a^6*b^2 + B*a^5*b^3 - 3*(A - 2*C)*a^4*b^4 - 3
*B*a^3*b^5 + A*a^2*b^6)*d^2 + ((B*a^3*b^5 - 3*(A - C)*a^2*b^6 - 3*B*a*b^7
+ (A - C)*b^8)*c^2 + 2*(((A - C)*a^3*b^5 + 3*B*a^2*b^6 - 3*(A - C)*a*b^7 -
B*b^8)*c*d - (C*a^6*b^2 + 3*C*a^4*b^4 + B*a^3*b^5 - 3*(A - 2*C)*a^2*b^6 -
3*B*a*b^7 + A*b^8)*d^2)*tan(f*x + e)^2 + 2*(((B*a^4*b^4 - 3*(A - C)*a^3*b^5
- 3*B*a^2*b^6 + (A - C)*a*b^7)*c^2 + 2*(((A - C)*a^4*b^4 + 3*B*a^3*b^5 ...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

= Exception raised: AttributeError

input

```

integrate((c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*
x+e))**3,x)

```

output

```

Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'

```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 839, normalized size of antiderivative = 1.41

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="maxima")
```

output

```
1/2*(2*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c^2 - 2*(B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c*d - ((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*d^2)*(f*x + e)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 2*((B*a^3*b^3 - 3*(A - C)*a^2*b^4 - 3*B*a*b^5 + (A - C)*b^6)*c^2 + 2*((A - C)*a^3*b^3 + 3*B*a^2*b^4 - 3*(A - C)*a*b^5 - B*b^6)*c*d - (C*a^6 + 3*C*a^4*b^2 + B*a^3*b^3 - 3*(A - 2*C)*a^2*b^4 - 3*B*a*b^5 + A*b^6)*d^2)*log(b*tan(f*x + e) + a)/(a^6*b^3 + 3*a^4*b^5 + 3*a^2*b^7 + b^9) + ((B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c^2 + 2*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c*d - (B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*d^2)*log(tan(f*x + e)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((C*a^4*b^2 - 3*B*a^3*b^3 + (5*A - 3*C)*a^2*b^4 + B*a*b^5 + A*b^6)*c^2 + 2*(C*a^5*b + B*a^4*b^2 - (3*A - 5*C)*a^3*b^3 - 3*B*a^2*b^4 + A*a*b^5)*c*d - (3*C*a^6 - B*a^5*b - (A - 7*C)*a^4*b^2 - 5*B*a^3*b^3 + 3*A*a^2*b^4)*d^2 - 2*((B*a^2*b^4 - 2*(A - C)*a*b^5 - B*b^6)*c^2 - 2*(C*a^4*b^2 - (A - 3*C)*a^2*b^4 - 2*B*a*b^5 + A*b^6)*c*d + (2*C*a^5*b - B*a^4*b^2 + 4*C*a^3*b^3 - 3*B*a^2*b^4 + 2*A*a*b^5)*d^2)*tan(f*x + e))/(a^6*b^3 + 2*a^4*b^5 + a^2*b^7 + (a^4*b^5 + 2*a^2*b^7 + b^9)*tan(f*x + e)^2 + 2*(a^5*b^4 + 2*a^3*b^6 + a*b^8)*tan(f*x + e))/f
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1218 vs. 2(595) = 1190.

Time = 0.88 (sec) , antiderivative size = 1218, normalized size of antiderivative = 2.04

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="giac")
```

output

```
(A*a^3*c^2 - C*a^3*c^2 + 3*B*a^2*b*c^2 - 3*A*a*b^2*c^2 + 3*C*a*b^2*c^2 - B*b^3*c^2 - 2*B*a^3*c*d + 6*A*a^2*b*c*d - 6*C*a^2*b*c*d + 6*B*a*b^2*c*d - 2*A*b^3*c*d + 2*C*b^3*c*d - A*a^3*d^2 + C*a^3*d^2 - 3*B*a^2*b*d^2 + 3*A*a*b^2*d^2 - 3*C*a*b^2*d^2 + B*b^3*d^2)*(f*x + e)/(a^6*f + 3*a^4*b^2*f + 3*a^2*b^4*f + b^6*f) + 1/2*(B*a^3*c^2 - 3*A*a^2*b*c^2 + 3*C*a^2*b*c^2 - 3*B*a*b^2*c^2 + A*b^3*c^2 - C*b^3*c^2 + 2*A*a^3*c*d - 2*C*a^3*c*d + 6*B*a^2*b*c*d - 6*A*a*b^2*c*d + 6*C*a*b^2*c*d - 2*B*b^3*c*d - B*a^3*d^2 + 3*A*a^2*b*d^2 - 3*C*a^2*b*d^2 + 3*B*a*b^2*d^2 - A*b^3*d^2 + C*b^3*d^2)*log(tan(f*x + e)^2 + 1)/(a^6*f + 3*a^4*b^2*f + 3*a^2*b^4*f + b^6*f) - (B*a^3*b^3*c^2 - 3*A*a^2*b^4*c^2 + 3*C*a^2*b^4*c^2 - 3*B*a*b^5*c^2 + A*b^6*c^2 - C*b^6*c^2 + 2*A*a^3*b^3*c*d - 2*C*a^3*b^3*c*d + 6*B*a^2*b^4*c*d - 6*A*a*b^5*c*d + 6*C*a*b^5*c*d - 2*B*b^6*c*d - C*a^6*d^2 - 3*C*a^4*b^2*d^2 - B*a^3*b^3*d^2 + 3*A*a^2*b^4*d^2 - 6*C*a^2*b^4*d^2 + 3*B*a*b^5*d^2 - A*b^6*d^2)*log(abs(b*tan(f*x + e) + a))/(a^6*b^3*f + 3*a^4*b^5*f + 3*a^2*b^7*f + b^9*f) + 1/2*(2*(B*a^4*b^3*c^2 - 2*A*a^3*b^4*c^2 + 2*C*a^3*b^4*c^2 - 2*A*a*b^6*c^2 + 2*C*a*b^6*c^2 - B*b^7*c^2 - 2*C*a^6*b*c*d + 2*A*a^4*b^3*c*d - 8*C*a^4*b^3*c*d + 4*B*a^3*b^4*c*d - 6*C*a^2*b^5*c*d + 4*B*a*b^6*c*d - 2*A*b^7*c*d + 2*C*a^7*d^2 - B*a^6*b*d^2 + 6*C*a^5*b^2*d^2 - 4*B*a^4*b^3*d^2 + 2*A*a^3*b^4*d^2 + 4*C*a^3*b^4*d^2 - 3*B*a^2*b^5*d^2 + 2*A*a*b^6*d^2)*tan(f*x + e) - (C*a^6*b^2*c^2 - 3*B*a^5*b^3*c^2 + 5*A*a^4*b^4*c^2 - 2*C*a^4*b^4*c^2 - 2*B*a^3*b...
```

Mupad [B] (verification not implemented)

Time = 24.36 (sec) , antiderivative size = 807, normalized size of antiderivative = 1.35

$$\int \frac{(c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx =$$

$$\frac{\ln(a + b \tan(e + fx)) \left(\frac{a^2 (b^4 (3A d^2 - 3A c^2 + 3C c^2 - 6C d^2 + 6B c d) + 3C b^4 d^2) - b^6 (A d^2 - A c^2 + C c^2 + 2B c d) + C b^6 d^2 - a b^5}{a^6 b^3 + 3a^4 b^5 + 3a^2 b^7 + b^9} \right)}{f}$$

$$\frac{A b^6 c^2 - 3C a^6 d^2 + B a b^5 c^2 + B a^5 b d^2 + 5A a^2 b^4 c^2 - 3A a^2 b^4 d^2 + A a^4 b^2 d^2 - 3B a^3 b^3 c^2 + 5B a^3 b^3 d^2 - 3C a^2 b^4 c^2 + C a^4 b^2 c^2 - 7C a^4 b^2 d^2}{2b^3 (a^4 + 2a^2 b^2 + b^4)}$$

$$\frac{\ln(\tan(e + fx) - i) (B c^2 - B d^2 + 2A c d - 2C c d - A c^2 \operatorname{li} + A d^2 \operatorname{li} + C c^2 \operatorname{li} - C d^2 \operatorname{li} + B c d 2i)}{2f (-a^3 - a^2 b 3i + 3a b^2 + b^3 \operatorname{li})}$$

$$\frac{\ln(\tan(e + fx) + i) (A d^2 - A c^2 + B c^2 \operatorname{li} - B d^2 \operatorname{li} + C c^2 - C d^2 + A c d 2i + 2B c d - C c d 2i)}{2f (-a^3 \operatorname{li} - 3a^2 b + a b^2 3i + b^3)}$$

input `int(((c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^3,x)`

output `- (log(a + b*tan(e + f*x))*((a^2*(b^4*(3*A*d^2 - 3*A*c^2 + 3*C*c^2 - 6*C*d^2 + 6*B*c*d) + 3*C*b^4*d^2) - b^6*(A*d^2 - A*c^2 + C*c^2 + 2*B*c*d) + C*b^6*d^2 - a*b^5*(3*B*c^2 - 3*B*d^2 + 6*A*c*d - 6*C*c*d) + a^3*b^3*(B*c^2 - B*d^2 + 2*A*c*d - 2*C*c*d)))/(b^9 + 3*a^2*b^7 + 3*a^4*b^5 + a^6*b^3) - (C*d^2)/b^3)/f - ((A*b^6*c^2 - 3*C*a^6*d^2 + B*a*b^5*c^2 + B*a^5*b*d^2 + 5*A*a^2*b^4*c^2 - 3*A*a^2*b^4*d^2 + A*a^4*b^2*d^2 - 3*B*a^3*b^3*c^2 + 5*B*a^3*b^3*d^2 - 3*C*a^2*b^4*c^2 + C*a^4*b^2*c^2 - 7*C*a^4*b^2*d^2 + 2*A*a*b^5*c*d + 2*C*a^5*b*c*d - 6*A*a^3*b^3*c*d - 6*B*a^2*b^4*c*d + 2*B*a^4*b^2*c*d + 10*C*a^3*b^3*c*d)/(2*b^3*(a^4 + b^4 + 2*a^2*b^2)) + (tan(e + f*x)*(B*b^5*c^2 - 2*C*a^5*d^2 + 2*A*b^5*c*d + 2*A*a*b^4*c^2 - 2*A*a*b^4*d^2 + B*a^4*b*d^2 - 2*C*a*b^4*c^2 - B*a^2*b^3*c^2 + 3*B*a^2*b^3*d^2 - 4*C*a^3*b^2*d^2 - 4*B*a*b^4*c*d + 2*C*a^4*b*c*d - 2*A*a^2*b^3*c*d + 6*C*a^2*b^3*c*d))/(b^2*(a^4 + b^4 + 2*a^2*b^2)))/(f*(a^2 + b^2*tan(e + f*x)^2 + 2*a*b*tan(e + f*x))) - (log(tan(e + f*x) - 1)*(A*d^2*1i - A*c^2*1i + B*c^2 - B*d^2 + C*c^2*1i - C*d^2*1i + 2*A*c*d + B*c*d*2i - 2*C*c*d))/(2*f*(3*a*b^2 - a^2*b*3i - a^3 + b^3*1i)) - (log(tan(e + f*x) + 1)*(A*d^2 - A*c^2 + B*c^2*1i - B*d^2*1i + C*c^2 - C*d^2 + A*c*d*2i + 2*B*c*d - C*c*d*2i))/(2*f*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3))`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 3274, normalized size of antiderivative = 5.48

$$\int \frac{(c + d \tan(e + f x))^2 (A + B \tan(e + f x) + C \tan^2(e + f x))}{(a + b \tan(e + f x))^3} dx = \text{Too large to display}$$

input `int((c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x)`

output

```
(2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**5*c*d - 2*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)**2*a**4*b**6*c**2 + 2*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**2*a**4*b**6*d**2 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2
*a**4*b**5*c**2*d + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**6*c
**3 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**6*c*d**2 - 2*log(
tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**8*c**2 + 2*log(tan(e + f*x)**
2 + 1)*tan(e + f*x)**2*a**2*b**8*d**2 + 6*log(tan(e + f*x)**2 + 1)*tan(e +
f*x)**2*a**2*b**7*c**2*d - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b
**9*c*d - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**8*c**3 + log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a*b**8*c*d**2 + 4*log(tan(e + f*x)**2 + 1)
*tan(e + f*x)*a**6*b**4*c*d - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5
*b**5*c**2 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*b**5*d**2 - 4*lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*b**4*c**2*d + 6*log(tan(e + f*x)*
**2 + 1)*tan(e + f*x)*a**4*b**5*c**3 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f
*x)*a**4*b**5*c*d**2 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**7*c
**2 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**7*d**2 + 12*log(tan(
e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**6*c**2*d - 4*log(tan(e + f*x)**2 + 1
)*tan(e + f*x)*a**2*b**8*c*d - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**
2*b**7*c**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**7*c*d**2 + 2
*log(tan(e + f*x)**2 + 1)*a**7*b**3*c*d - 2*log(tan(e + f*x)**2 + 1)*a...
```

3.64 $\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^3 (A + B \tan(e +$

Optimal result	699
Mathematica [C] (verified)	700
Rubi [A] (verified)	701
Maple [A] (warning: unable to verify)	706
Fricas [A] (verification not implemented)	707
Sympy [B] (verification not implemented)	707
Maxima [A] (verification not implemented)	708
Giac [B] (verification not implemented)	709
Mupad [B] (verification not implemented)	710
Reduce [B] (verification not implemented)	711

Optimal result

Integrand size = 45, antiderivative size = 603

$$\begin{aligned}
 & \int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) \, dx \\
 &= (a^2 (Ac^3 - c^3 C - 3Bc^2 d - 3Acd^2 + 3cCd^2 + Bd^3) \\
 &\quad + b^2 (c^3 C + 3Bc^2 d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2)) \\
 &\quad - 2ab((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))) x \\
 &+ \frac{(2ab(c^3 C + 3Bc^2 d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2)) - a^2((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) + b^2((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)))}{f} \\
 &- \frac{d(2ab(c^2 C + 2Bcd - Cd^2 - A(c^2 - d^2)) - a^2(2c(A - C)d + B(c^2 - d^2)) + b^2(2c(A - C)d + B(c^2 - d^2)))}{f} \\
 &+ \frac{(2ab(Ac - cC - Bd) + a^2(Bc + (A - C)d) - b^2(Bc + (A - C)d)) (c + d \tan(e + fx))^2}{2f} \\
 &+ \frac{(a^2 B - b^2 B + 2ab(A - C)) (c + d \tan(e + fx))^3}{3f} \\
 &+ \frac{(5a^2 C d^2 - 6abd(cC - 5Bd) + b^2(c^2 C - 3Bcd + 15(A - C)d^2)) (c + d \tan(e + fx))^4}{60d^3 f} \\
 &- \frac{b(bcC - 3bBd - aCd) \tan(e + fx) (c + d \tan(e + fx))^4}{15d^2 f} \\
 &+ \frac{C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^4}{6df}
 \end{aligned}$$

output

```
(a^2*(A*c^3-3*A*c*d^2-3*B*c^2*d+B*d^3-C*c^3+3*C*c*d^2)+b^2*(c^3*C+3*B*c^2*d-3*C*c*d^2-B*d^3-A*(c^3-3*c*d^2))-2*a*b*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*d^2)))*x+(2*a*b*(c^3*C+3*B*c^2*d-3*C*c*d^2-B*d^3-A*(c^3-3*c*d^2))-a^2*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*d^2))+b^2*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*d^2)))*ln(cos(f*x+e))/f-d*(2*a*b*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-a^2*(2*c*(A-C)*d+B*(c^2-d^2))+b^2*(2*c*(A-C)*d+B*(c^2-d^2)))*tan(f*x+e)/f+1/2*(2*a*b*(A*c-B*d-C*c)+a^2*(B*c+(A-C)*d)-b^2*(B*c+(A-C)*d))*(c+d*tan(f*x+e))^2/f+1/3*(B*a^2-B*b^2+2*a*b*(A-C))*(c+d*tan(f*x+e))^3/f+1/60*(5*a^2*C*d^2-6*a*b*d*(-5*B*d+C*c)+b^2*(c^2*C-3*B*c*d+15*(A-C)*d^2))*(c+d*tan(f*x+e))^4/d^3/f-1/15*b*(-3*B*b*d-C*a*d+C*b*c)*tan(f*x+e)*(c+d*tan(f*x+e))^4/d^2/f+1/6*C*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^4/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.40 (sec) , antiderivative size = 419, normalized size of antiderivative = 0.69

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^4}{6df}$$

$$+ \frac{-\frac{2b(bcC-3bBd-aCd)\tan(e+fx)(c+d\tan(e+fx))^4}{5df} - \frac{(5a^2C^2d^2-6abd(cC-5Bd)+b^2(c^2C-3Bcd+15(A-C)d^2))(c+d\tan(e+fx))^4}{2df}}{5(3d(2a$$

input

```
Integrate[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(C*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^4)/(6*d*f) + ((-2*b*(b*c*C - 3*b*B*d - a*C*d)*Tan[e + f*x]*(c + d*Tan[e + f*x])^4)/(5*d*f) - (-1/2*((5*a^2*C*d^2 - 6*a*b*d*(c*C - 5*B*d) + b^2*(c^2*C - 3*B*c*d + 15*(A - C)*d^2))*(c + d*Tan[e + f*x])^4)/(d*f) + (5*(3*d*(2*a*b*(A*c - c*C + B*d) + a^2*(B*c - (A - C)*d) - b^2*(B*c - (A - C)*d))*((I*c - d)^3*Log[I - Tan[e + f*x]] - (I*c + d)^3*Log[I + Tan[e + f*x]] + 6*c*d^2*Tan[e + f*x] + d^3*Tan[e + f*x]^2) + (a^2*B - b^2*B + 2*a*b*(A - C))*d*((3*I)*(c + I*d)^4*Log[I - Tan[e + f*x]] - (3*I)*(c - I*d)^4*Log[I + Tan[e + f*x]] - 6*d^2*(6*c^2 - d^2)*Tan[e + f*x] - 12*c*d^3*Tan[e + f*x]^2 - 2*d^4*Tan[e + f*x]^3)))/f)/(5*d))/(6*d)
```

Rubi [A] (verified)

Time = 4.72 (sec) , antiderivative size = 633, normalized size of antiderivative = 1.05, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.356$, Rules used = {3042, 4130, 27, 3042, 4120, 25, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4130

$$\frac{\int -2(a + b \tan(e + fx))(c + d \tan(e + fx))^3 ((bcC - adC - 3bBd) \tan^2(e + fx) - 3(Ab - Cb + aB)d \tan(e + fx)) dx}{\frac{C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^4}{6df} - 6d}$$

↓ 27

$$\frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 ((bcC - adC - 3bBd) \tan^2(e + fx) - 3(Ab - Cb + aB)d \tan(e + fx)) dx}{\frac{C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^4}{6df} - 3d}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 ((bcC - adC - 3bBd) \tan(e + fx)^2 - 3(Ab - Cb + aB)d \tan(e + fx)) dx}{\frac{C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^4}{6df} - 3d}$$

↓ 4120

$$\frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 ((bcC - adC - 3bBd) \tan(e + fx)^2 - 3(Ab - Cb + aB)d \tan(e + fx)) dx}{\frac{C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^4}{6df} - 3d}$$

↓ 25

$$\frac{b \tan(e + fx)(-aCd - 3bBd + bcC)(c + d \tan(e + fx))^4}{5df} - \frac{\int -(c + d \tan(e + fx))^3 (-c(cC - 3Bd)b^2 + 6acCdb - 5a^2(3A - 2C)d^2 - ((Cc^2 - 3Bdc + 15(A$$

3d

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx))^3 (-c(cC - 3Bd)b^2 + 6acCdb - 5a^2(3A - 2C)d^2 - ((C^2 - 3Bdc + 15(A - C)d^2)b^2 - 6ad(cC - 5Bd)b + 5a^2Cd^2) \tan^2(e + fx) - 15(Bc - 5Ad)a^2) dx}{5d}$$

3d

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx))^3 (-c(cC - 3Bd)b^2 + 6acCdb - 5a^2(3A - 2C)d^2 - ((C^2 - 3Bdc + 15(A - C)d^2)b^2 - 6ad(cC - 5Bd)b + 5a^2Cd^2) \tan(e + fx)^2 - 15(Bc - 5Ad)a^2) dx}{5d}$$

3d

↓ 4113

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx))^3 (15(-((A - C)a^2) + 2bBa + b^2(A - C))d^2 - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)) dx - \frac{(c + d \tan(e + fx))^4 (5a^2Cd^2 - 6abd(cC - 5Ad))}{4df}}{5d}$$

3d

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx))^3 (15(-((A - C)a^2) + 2bBa + b^2(A - C))d^2 - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)) dx - \frac{(c + d \tan(e + fx))^4 (5a^2Cd^2 - 6abd(cC - 5Ad))}{4df}}{5d}$$

3d

↓ 4011

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx))^2 (-15((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd))d^2 - 15((Bc + (A - C)d)a^2 + 2b(Ac - Cc - Bd)a - b^2(Bc + (A - C)d))) dx}{5d}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx))^2 (-15((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd))d^2 - 15((Bc + (A - C)d)a^2 + 2b(Ac - Cc - Bd)a - b^2(Bc + (A - C)d))) dx}{5d}$$

↓ 4011

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx)) (15((Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a^2 + 2b(2c(A - C)d + B(c^2 - d^2))a - b^2(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))d^2 + 15(-((2c(A - C) - b^2d)(c^2 - d^2))a^2 + 2b(2c(A - C)d + B(c^2 - d^2))a - b^2(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))d}{6df}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{\int (c + d \tan(e + fx)) (15((Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a^2 + 2b(2c(A - C)d + B(c^2 - d^2))a - b^2(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))d^2 + 15(-((2c(A - C) - b^2d)(c^2 - d^2))a^2 + 2b(2c(A - C)d + B(c^2 - d^2))a - b^2(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))d}{6df}$$

↓ 4008

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{15d^2(-a^2(d(A - C)(3c^2 - d^2) + B(c^3 - 3cd^2))) + 2ab(-A(c^3 - 3cd^2) + 3Bc^2d - Bd^3 + c^3C - 3cCd^2) + b^2(d(A - C)(3c^2 - d^2) + B(c^3 - 3cd^2)) \int \tan(e + fx)}{6df}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{15d^2(-a^2(d(A - C)(3c^2 - d^2) + B(c^3 - 3cd^2))) + 2ab(-A(c^3 - 3cd^2) + 3Bc^2d - Bd^3 + c^3C - 3cCd^2) + b^2(d(A - C)(3c^2 - d^2) + B(c^3 - 3cd^2)) \int \tan(e + fx)}{6df}$$

↓ 3956

$$\frac{C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^4}{6df} - \frac{(c + d \tan(e + fx))^4(5a^2Cd^2 - 6abd(cC - 5Bd) + b^2(15d^2(A - C) - 3Bcd + c^2C))}{4df} + \frac{15d^3 \tan(e + fx)(-a^2(2cd(A - C) + B(c^2 - d^2))) + 2ab(-A(c^2 - d^2) + 2Bcd + c^3)}{f}$$

input

```
Int[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*
Tan[e + f*x]^2),x]
```

output

```
(C*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^4)/(6*d*f) - ((b*(b*c*C - 3
*b*B*d - a*C*d)*Tan[e + f*x]*(c + d*Tan[e + f*x])^4)/(5*d*f) + (-15*d^2*(a
^2*(A*c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3) + b^2*(c^3*
C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) - 2*a*b*((A - C)*d*
(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)))*x - (15*d^2*(2*a*b*(c^3*C + 3*B*c^2*d
- 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) - a^2*((A - C)*d*(3*c^2 - d^2) +
B*(c^3 - 3*c*d^2)) + b^2*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)))*Lo
g[Cos[e + f*x]])/f + (15*d^3*(2*a*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d
^2)) - a^2*(2*c*(A - C)*d + B*(c^2 - d^2)) + b^2*(2*c*(A - C)*d + B*(c^2 -
d^2)))*Tan[e + f*x])/f - (15*d^2*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A
- C)*d) - b^2*(B*c + (A - C)*d))*(c + d*Tan[e + f*x])^2)/(2*f) - (5*(a^2*B
- b^2*B + 2*a*b*(A - C))*d^2*(c + d*Tan[e + f*x])^3)/f - ((5*a^2*C*d^2 -
6*a*b*d*(c*C - 5*B*d) + b^2*(c^2*C - 3*B*c*d + 15*(A - C)*d^2))*(c + d*Tan
[e + f*x])^4)/(4*d*f)/(5*d)/(3*d)
```

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x])/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp
[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp
[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e.
_) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (warning: unable to verify)

Time = 0.41 (sec) , antiderivative size = 546, normalized size of antiderivative = 0.91

method	result
parts	$\frac{(3Aa^2c^2d+2Aabc^3+Ba^2c^3)\ln(1+\tan(fx+e)^2)}{2f} + \frac{(Bb^2d^3+2Cab^2d^3+3Cb^2cd^2)\left(\frac{\tan(fx+e)^5}{5} - \frac{\tan(fx+e)^3}{3} + \tan(fx+e)\right)}{f}$
norman	$(Aa^2c^3 - 3Aa^2cd^2 - 6Aabc^2d + 2Aabd^3 - Ab^2c^3 + 3Ab^2cd^2 - 3Ba^2c^2d + Ba^2d^3)$
derivativedivides	Expression too large to display
default	Expression too large to display
parallelrisch	Expression too large to display
risch	Expression too large to display

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x,method=_RETURNVERBOSE)
```

output

```
1/2*(3*A*a^2*c^2*d+2*A*a*b*c^3+B*a^2*c^3)/f*ln(1+tan(f*x+e)^2)+(B*b^2*d^3+
2*C*a*b*d^3+3*C*b^2*c*d^2)/f*(1/5*tan(f*x+e)^5-1/3*tan(f*x+e)^3+tan(f*x+e)
-arctan(tan(f*x+e)))+(A*b^2*d^3+2*B*a*b*d^3+3*B*b^2*c*d^2+C*a^2*d^3+6*C*a*
b*c*d^2+3*C*b^2*c^2*d)/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(f
*x+e)^2))+(3*A*a^2*c*d^2+6*A*a*b*c^2*d+A*b^2*c^3+3*B*a^2*c^2*d+2*B*a*b*c^3
+C*a^2*c^3)/f*(tan(f*x+e)-arctan(tan(f*x+e)))+(2*A*a*b*d^3+3*A*b^2*c*d^2+B
*a^2*d^3+6*B*a*b*c*d^2+3*B*b^2*c^2*d+3*C*a^2*c*d^2+6*C*a*b*c^2*d+C*b^2*c^3
)/f*(1/3*tan(f*x+e)^3-tan(f*x+e)+arctan(tan(f*x+e)))+(A*a^2*d^3+6*A*a*b*c*
d^2+3*A*b^2*c^2*d+3*B*a^2*c*d^2+6*B*a*b*c^2*d+B*b^2*c^3+3*C*a^2*c^2*d+2*C*
a*b*c^3)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)^2))+A*a^2*c^3*x+C*b^2*d^3
/f*(1/6*tan(f*x+e)^6-1/4*tan(f*x+e)^4+1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)
^2))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 679, normalized size of antiderivative = 1.13

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

```
1/60*(10*C*b^2*d^3*tan(f*x + e)^6 + 12*(3*C*b^2*c*d^2 + (2*C*a*b + B*b^2)*d^3)*tan(f*x + e)^5 + 15*(3*C*b^2*c^2*d + 3*(2*C*a*b + B*b^2)*c*d^2 + (C*a^2 + 2*B*a*b + (A - C)*b^2)*d^3)*tan(f*x + e)^4 + 20*(C*b^2*c^3 + 3*(2*C*a*b + B*b^2)*c^2*d + 3*(C*a^2 + 2*B*a*b + (A - C)*b^2)*c*d^2 + (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^3)*tan(f*x + e)^3 + 60*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^3 - 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2*d - 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^2 + (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^3)*f*x + 30*((2*C*a*b + B*b^2)*c^3 + 3*(C*a^2 + 2*B*a*b + (A - C)*b^2)*c^2*d + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d^2 + ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^3)*tan(f*x + e)^2 - 30*((B*a^2 + 2*(A - C)*a*b - B*b^2)*c^3 + 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^2*d - 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d^2 - ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^3)*log(1/(tan(f*x + e)^2 + 1)) + 60*((C*a^2 + 2*B*a*b + (A - C)*b^2)*c^3 + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2*d + 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^2 - (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^3)*tan(f*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1819 vs. 2(547) = 1094.

Time = 0.41 (sec) , antiderivative size = 1819, normalized size of antiderivative = 3.02

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))**2*(c+d*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```


output

```
Piecewise((A**2*c**3*x + 3*A**2*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f)
- 3*A**2*c*d**2*x + 3*A**2*c*d**2*tan(e + f*x)/f - A**2*d**3*log(tan
(e + f*x)**2 + 1)/(2*f) + A**2*d**3*tan(e + f*x)**2/(2*f) + A*b*c**3*log
(tan(e + f*x)**2 + 1)/f - 6*A*b*c**2*d*x + 6*A*b*c**2*d*tan(e + f*x)
/f - 3*A*b*c*d**2*log(tan(e + f*x)**2 + 1)/f + 3*A*b*c*d**2*tan(e + f*
x)**2/f + 2*A*b*d**3*x + 2*A*b*d**3*tan(e + f*x)**3/(3*f) - 2*A*b*d*
**3*tan(e + f*x)/f - A*b**2*c**3*x + A*b**2*c**3*tan(e + f*x)/f - 3*A*b**2*
c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*A*b**2*c**2*d*tan(e + f*x)**2/(2
*f) + 3*A*b**2*c*d**2*x + A*b**2*c*d**2*tan(e + f*x)**3/f - 3*A*b**2*c*d**
2*tan(e + f*x)/f + A*b**2*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + A*b**2*d**
3*tan(e + f*x)**4/(4*f) - A*b**2*d**3*tan(e + f*x)**2/(2*f) + B**2*c**3*
log(tan(e + f*x)**2 + 1)/(2*f) - 3*B**2*c**2*d*x + 3*B**2*c**2*d*tan(e
+ f*x)/f - 3*B**2*c*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B**2*c*d*
**2*tan(e + f*x)**2/(2*f) + B**2*d**3*x + B**2*d**3*tan(e + f*x)**3/(3*
f) - B**2*d**3*tan(e + f*x)/f - 2*B*a*b*c**3*x + 2*B*a*b*c**3*tan(e + f*
x)/f - 3*B*a*b*c**2*d*log(tan(e + f*x)**2 + 1)/f + 3*B*a*b*c**2*d*tan(e +
f*x)**2/f + 6*B*a*b*c*d**2*x + 2*B*a*b*c*d**2*tan(e + f*x)**3/f - 6*B*a*b*
c*d**2*tan(e + f*x)/f + B*a*b*d**3*log(tan(e + f*x)**2 + 1)/f + B*a*b*d**3
*tan(e + f*x)**4/(2*f) - B*a*b*d**3*tan(e + f*x)**2/f - B*b**2*c**3*log(ta
n(e + f*x)**2 + 1)/(2*f) + B*b**2*c**3*tan(e + f*x)**2/(2*f) + 3*B*b**2...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 680, normalized size of antiderivative = 1.13

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+
e)^2),x, algorithm="maxima")
```

output

```

1/60*(10*C*b^2*d^3*tan(f*x + e)^6 + 12*(3*C*b^2*c*d^2 + (2*C*a*b + B*b^2)*
d^3)*tan(f*x + e)^5 + 15*(3*C*b^2*c^2*d + 3*(2*C*a*b + B*b^2)*c*d^2 + (C*a
^2 + 2*B*a*b + (A - C)*b^2)*d^3)*tan(f*x + e)^4 + 20*(C*b^2*c^3 + 3*(2*C*a
*b + B*b^2)*c^2*d + 3*(C*a^2 + 2*B*a*b + (A - C)*b^2)*c*d^2 + (B*a^2 + 2*(
A - C)*a*b - B*b^2)*d^3)*tan(f*x + e)^3 + 30*((2*C*a*b + B*b^2)*c^3 + 3*(C
*a^2 + 2*B*a*b + (A - C)*b^2)*c^2*d + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*
d^2 + ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^3)*tan(f*x + e)^2 + 60*((A
- C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^3 - 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*
c^2*d - 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^2 + (B*a^2 + 2*(A - C)
*a*b - B*b^2)*d^3)*(f*x + e) + 30*((B*a^2 + 2*(A - C)*a*b - B*b^2)*c^3 + 3
*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^2*d - 3*(B*a^2 + 2*(A - C)*a*b -
B*b^2)*c*d^2 - ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^3)*log(tan(f*x + e)
^2 + 1) + 60*((C*a^2 + 2*B*a*b + (A - C)*b^2)*c^3 + 3*(B*a^2 + 2*(A - C)*a
*b - B*b^2)*c^2*d + 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^2 - (B*a^2
+ 2*(A - C)*a*b - B*b^2)*d^3)*tan(f*x + e))/f

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1413 vs. 2(593) = 1186.

Time = 1.18 (sec) , antiderivative size = 1413, normalized size of antiderivative = 2.34

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```

integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+
e)^2),x, algorithm="giac")

```

output

```
(A*a^2*c^3 - C*a^2*c^3 - 2*B*a*b*c^3 - A*b^2*c^3 + C*b^2*c^3 - 3*B*a^2*c^2*d - 6*A*a*b*c^2*d + 6*C*a*b*c^2*d + 3*B*b^2*c^2*d - 3*A*a^2*c*d^2 + 3*C*a^2*c*d^2 + 6*B*a*b*c*d^2 + 3*A*b^2*c*d^2 - 3*C*b^2*c*d^2 + B*a^2*d^3 + 2*A*a*b*d^3 - 2*C*a*b*d^3 - B*b^2*d^3)*(f*x + e)/f + 1/2*(B*a^2*c^3 + 2*A*a*b*c^3 - 2*C*a*b*c^3 - B*b^2*c^3 + 3*A*a^2*c^2*d - 3*C*a^2*c^2*d - 6*B*a*b*c^2*d - 3*A*b^2*c^2*d + 3*C*b^2*c^2*d - 3*B*a^2*c*d^2 - 6*A*a*b*c*d^2 + 6*C*a*b*c*d^2 + 3*B*b^2*c*d^2 - A*a^2*d^3 + C*a^2*d^3 + 2*B*a*b*d^3 + A*b^2*d^3 - C*b^2*d^3)*log(tan(f*x + e)^2 + 1)/f + 1/60*(10*C*b^2*d^3*f^5*tan(f*x + e)^6 + 36*C*b^2*c*d^2*f^5*tan(f*x + e)^5 + 24*C*a*b*d^3*f^5*tan(f*x + e)^5 + 12*B*b^2*d^3*f^5*tan(f*x + e)^5 + 45*C*b^2*c^2*d*f^5*tan(f*x + e)^4 + 90*C*a*b*c*d^2*f^5*tan(f*x + e)^4 + 45*B*b^2*c*d^2*f^5*tan(f*x + e)^4 + 15*C*a^2*d^3*f^5*tan(f*x + e)^4 + 30*B*a*b*d^3*f^5*tan(f*x + e)^4 + 15*A*b^2*d^3*f^5*tan(f*x + e)^4 - 15*C*b^2*d^3*f^5*tan(f*x + e)^4 + 20*C*b^2*c^3*f^5*tan(f*x + e)^3 + 120*C*a*b*c^2*d*f^5*tan(f*x + e)^3 + 60*B*b^2*c^2*d*f^5*tan(f*x + e)^3 + 60*C*a^2*c*d^2*f^5*tan(f*x + e)^3 + 120*B*a*b*c*d^2*f^5*tan(f*x + e)^3 + 60*A*b^2*c*d^2*f^5*tan(f*x + e)^3 - 60*C*b^2*c*d^2*f^5*tan(f*x + e)^3 + 20*B*a^2*d^3*f^5*tan(f*x + e)^3 + 40*A*a*b*d^3*f^5*tan(f*x + e)^3 - 40*C*a*b*d^3*f^5*tan(f*x + e)^3 - 20*B*b^2*d^3*f^5*tan(f*x + e)^3 + 60*C*a*b*c^3*f^5*tan(f*x + e)^2 + 30*B*b^2*c^3*f^5*tan(f*x + e)^2 + 90*C*a^2*c^2*d*f^5*tan(f*x + e)^2 + 180*B*a*b*c^2*d*f^5*tan(f*x + e)^2 ...
```

Mupad [B] (verification not implemented)

Time = 5.99 (sec) , antiderivative size = 891, normalized size of antiderivative = 1.48

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
int((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
x*(A*a^2*c^3 - A*b^2*c^3 + B*a^2*d^3 - C*a^2*c^3 - B*b^2*d^3 + C*b^2*c^3 +
2*A*a*b*d^3 - 2*B*a*b*c^3 - 2*C*a*b*d^3 - 3*A*a^2*c*d^2 + 3*A*b^2*c*d^2 -
3*B*a^2*c^2*d + 3*B*b^2*c^2*d + 3*C*a^2*c*d^2 - 3*C*b^2*c*d^2 - 6*A*a*b*c
^2*d + 6*B*a*b*c*d^2 + 6*C*a*b*c^2*d) - (tan(e + f*x)*(B*a^2*d^3 - A*b^2*c
^3 - b*d^2*(B*b*d + 2*C*a*d + 3*C*b*c) - C*a^2*c^3 + C*b^2*c^3 + 2*A*a*b*d
^3 - 2*B*a*b*c^3 - 3*A*a^2*c*d^2 + 3*A*b^2*c*d^2 - 3*B*a^2*c^2*d + 3*B*b^2
*c^2*d + 3*C*a^2*c*d^2 - 6*A*a*b*c^2*d + 6*B*a*b*c*d^2 + 6*C*a*b*c^2*d))/f
- (log(tan(e + f*x)^2 + 1)*((A*a^2*d^3)/2 - (B*a^2*c^3)/2 - (A*b^2*d^3)/2
+ (B*b^2*c^3)/2 - (C*a^2*d^3)/2 + (C*b^2*d^3)/2 - A*a*b*c^3 - B*a*b*d^3 +
C*a*b*c^3 - (3*A*a^2*c^2*d)/2 + (3*A*b^2*c^2*d)/2 + (3*B*a^2*c*d^2)/2 - (
3*B*b^2*c*d^2)/2 + (3*C*a^2*c^2*d)/2 - (3*C*b^2*c^2*d)/2 + 3*A*a*b*c*d^2 +
3*B*a*b*c^2*d - 3*C*a*b*c*d^2))/f + (tan(e + f*x)^4*((A*b^2*d^3)/4 + (C*a
^2*d^3)/4 - (C*b^2*d^3)/4 + (B*a*b*d^3)/2 + (3*B*b^2*c*d^2)/4 + (3*C*b^2*c
^2*d)/4 + (3*C*a*b*c*d^2)/2))/f + (tan(e + f*x)^3*((B*a^2*d^3)/3 - (b*d^2*
(B*b*d + 2*C*a*d + 3*C*b*c))/3 + (C*b^2*c^3)/3 + (2*A*a*b*d^3)/3 + A*b^2*c
*d^2 + B*b^2*c^2*d + C*a^2*c*d^2 + 2*B*a*b*c*d^2 + 2*C*a*b*c^2*d))/f + (ta
n(e + f*x)^2*((A*a^2*d^3)/2 - (A*b^2*d^3)/2 + (B*b^2*c^3)/2 - (C*a^2*d^3)/
2 + (C*b^2*d^3)/2 - B*a*b*d^3 + C*a*b*c^3 + (3*A*b^2*c^2*d)/2 + (3*B*a^2*c
*d^2)/2 - (3*B*b^2*c*d^2)/2 + (3*C*a^2*c^2*d)/2 - (3*C*b^2*c^2*d)/2 + 3*A*
a*b*c*d^2 + 3*B*a*b*c^2*d - 3*C*a*b*c*d^2))/f + (b*d^2*tan(e + f*x)^5*(...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1163, normalized size of antiderivative = 1.93

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x)
```

output

```
(90*log(tan(e + f*x)**2 + 1)*a**3*c**2*d - 30*log(tan(e + f*x)**2 + 1)*a**
3*d**3 + 90*log(tan(e + f*x)**2 + 1)*a**2*b*c**3 - 270*log(tan(e + f*x)**2
+ 1)*a**2*b*c*d**2 - 90*log(tan(e + f*x)**2 + 1)*a**2*c**3*d + 30*log(tan
(e + f*x)**2 + 1)*a**2*c*d**3 - 270*log(tan(e + f*x)**2 + 1)*a*b**2*c**2*d
+ 90*log(tan(e + f*x)**2 + 1)*a*b**2*d**3 - 60*log(tan(e + f*x)**2 + 1)*a
*b*c**4 + 180*log(tan(e + f*x)**2 + 1)*a*b*c**2*d**2 - 30*log(tan(e + f*x)
**2 + 1)*b**3*c**3 + 90*log(tan(e + f*x)**2 + 1)*b**3*c*d**2 + 90*log(tan(
e + f*x)**2 + 1)*b**2*c**3*d - 30*log(tan(e + f*x)**2 + 1)*b**2*c*d**3 + 1
0*tan(e + f*x)**6*b**2*c*d**3 + 24*tan(e + f*x)**5*a*b*c*d**3 + 12*tan(e +
f*x)**5*b**3*d**3 + 36*tan(e + f*x)**5*b**2*c**2*d**2 + 15*tan(e + f*x)**
4*a**2*c*d**3 + 45*tan(e + f*x)**4*a*b**2*d**3 + 90*tan(e + f*x)**4*a*b*c*
*2*d**2 + 45*tan(e + f*x)**4*b**3*c*d**2 + 45*tan(e + f*x)**4*b**2*c**3*d
- 15*tan(e + f*x)**4*b**2*c*d**3 + 60*tan(e + f*x)**3*a**2*b*d**3 + 60*tan
(e + f*x)**3*a**2*c**2*d**2 + 180*tan(e + f*x)**3*a*b**2*c*d**2 + 120*tan(
e + f*x)**3*a*b*c**3*d - 40*tan(e + f*x)**3*a*b*c*d**3 + 60*tan(e + f*x)**
3*b**3*c**2*d - 20*tan(e + f*x)**3*b**3*d**3 + 20*tan(e + f*x)**3*b**2*c**
4 - 60*tan(e + f*x)**3*b**2*c**2*d**2 + 30*tan(e + f*x)**2*a**3*d**3 + 270
*tan(e + f*x)**2*a**2*b*c*d**2 + 90*tan(e + f*x)**2*a**2*c**3*d - 30*tan(e
+ f*x)**2*a**2*c*d**3 + 270*tan(e + f*x)**2*a*b**2*c**2*d - 90*tan(e + f*
x)**2*a*b**2*d**3 + 60*tan(e + f*x)**2*a*b*c**4 - 180*tan(e + f*x)**2*a...
```

3.65 $\int (a+b \tan(e+fx))(c+d \tan(e+fx))^3 (A + B \tan(e +$

Optimal result	713
Mathematica [C] (verified)	714
Rubi [A] (verified)	715
Maple [A] (warning: unable to verify)	718
Fricas [A] (verification not implemented)	719
Sympy [B] (verification not implemented)	720
Maxima [A] (verification not implemented)	721
Giac [A] (verification not implemented)	721
Mupad [B] (verification not implemented)	723
Reduce [B] (verification not implemented)	724

Optimal result

Integrand size = 43, antiderivative size = 389

$$\begin{aligned}
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 &= (a(Ac^3 - c^3C - 3Bc^2d - 3Acd^2 + 3cCd^2 + Bd^3) \\
 &\quad - b((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))) x \\
 &\quad - \frac{(A(bc^3 + 3ac^2d - 3bcd^2 - ad^3) - b(c^3C + 3Bc^2d - 3cCd^2 - Bd^3) + a(Bc^3 - 3c^2Cd - 3Bcd^2 + Cd^3))}{f} \\
 &\quad + \frac{d(a(Bc^2 - 2cCd - Bd^2) - b(c^2C + 2Bcd - Cd^2) + A(2acd + b(c^2 - d^2))) \tan(e + fx)}{f} \\
 &\quad + \frac{(Abc + aBc - bcC + aAd - bBd - aCd)(c + d \tan(e + fx))^2}{2f} \\
 &\quad + \frac{(Ab + aB - bC)(c + d \tan(e + fx))^3}{3f} \\
 &\quad - \frac{(bcC - 5bBd - 5aCd)(c + d \tan(e + fx))^4}{20d^2f} + \frac{bC \tan(e + fx)(c + d \tan(e + fx))^4}{5df}
 \end{aligned}$$

output

```
(a*(A*c^3-3*A*c*d^2-3*B*c^2*d+B*d^3-C*c^3+3*C*c*d^2)-b*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*d^2)))*x-(A*(3*a*c^2*d-a*d^3+b*c^3-3*b*c*d^2)-b*(3*B*c^2*d-B*d^3+C*c^3-3*C*c*d^2)+a*(B*c^3-3*B*c*d^2-3*C*c^2*d+C*d^3))*ln(cos(f*x+e))/f+d*(a*(B*c^2-B*d^2-2*C*c*d)-b*(2*B*c*d+C*c^2-C*d^2)+A*(2*a*c*d+b*(c^2-d^2)))*tan(f*x+e)/f+1/2*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*(c+d*tan(f*x+e))^2/f+1/3*(A*b+B*a-C*b)*(c+d*tan(f*x+e))^3/f-1/20*(-5*B*b*d-5*C*a*d+C*b*c)*(c+d*tan(f*x+e))^4/d^2/f+1/5*b*C*tan(f*x+e)*(c+d*tan(f*x+e))^4/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.25 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.76

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{bC \tan(e + fx)(c + d \tan(e + fx))^4}{5df} + \frac{(bcC - 5bBd - 5aCd)(c + d \tan(e + fx))^4}{4df} + \frac{5(3(abc + aBc - bcC - aAd + bBd + aCd)((ic - d)^3 \log(i - \tan(e + fx)) - (ic + d)^3 \log(i + \tan(e + fx)))}{df}$$

input

```
Integrate[(a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(b*C*Tan[e + f*x]*(c + d*Tan[e + f*x])^4)/(5*d*f) - (((b*c*C - 5*b*B*d - 5*a*C*d)*(c + d*Tan[e + f*x])^4)/(4*d*f) + (5*(3*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d))*((I*c - d)^3*Log[I - Tan[e + f*x]] - (I*c + d)^3*Log[I + Tan[e + f*x]] + 6*c*d^2*Tan[e + f*x] + d^3*Tan[e + f*x]^2) + (A*b + a*B - b*C))*((3*I)*(c + I*d)^4*Log[I - Tan[e + f*x]] - (3*I)*(c - I*d)^4*Log[I + Tan[e + f*x]] - 6*d^2*(6*c^2 - d^2)*Tan[e + f*x] - 12*c*d^3*Tan[e + f*x]^2 - 2*d^4*Tan[e + f*x]^3)))/(6*f))/(5*d)
```

Rubi [A] (verified)

Time = 2.68 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.279$, Rules used = {3042, 4120, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 & \quad \downarrow 3042 \\
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx \\
 & \quad \downarrow 4120 \\
 & \frac{bC \tan(e + fx)(c + d \tan(e + fx))^4}{5df} - \\
 & \frac{\int (c + d \tan(e + fx))^3 ((bcC - 5adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + bcC - 5aAd) dx}{5d} \\
 & \quad \downarrow 3042 \\
 & \frac{bC \tan(e + fx)(c + d \tan(e + fx))^4}{5df} - \\
 & \frac{\int (c + d \tan(e + fx))^3 ((bcC - 5adC - 5bBd) \tan(e + fx)^2 - 5(Ab - Cb + aB)d \tan(e + fx) + bcC - 5aAd) dx}{5d} \\
 & \quad \downarrow 4113 \\
 & \frac{bC \tan(e + fx)(c + d \tan(e + fx))^4}{5df} - \\
 & \frac{\int (c + d \tan(e + fx))^3 (5(bB - a(A - C))d - 5(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{(-5aCd - 5bBd + bcC)(c + d \tan(e + fx))}{4df}}{5d} \\
 & \quad \downarrow 3042 \\
 & \frac{bC \tan(e + fx)(c + d \tan(e + fx))^4}{5df} - \\
 & \frac{\int (c + d \tan(e + fx))^3 (5(bB - a(A - C))d - 5(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{(-5aCd - 5bBd + bcC)(c + d \tan(e + fx))}{4df}}{5d} \\
 & \quad \downarrow 4011
 \end{aligned}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - \int (c+d \tan(e+fx))^2 (5d(bBc+b(A-C)d-a(Ac-Cc-Bd)) - 5d(Abc+aBc-bCc+aAd-bBd-aCd) t}{5d}}{\downarrow 3042}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - \int (c+d \tan(e+fx))^2 (5d(bBc+b(A-C)d-a(Ac-Cc-Bd)) - 5d(Abc+aBc-bCc+aAd-bBd-aCd) t}{5d}}{\downarrow 4011}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - \int (c+d \tan(e+fx)) (5d(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2))+b(2c(A-C)d+B(c^2-d^2))) - 5d(2aAc d-2}}{\downarrow 3042}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - \int (c+d \tan(e+fx)) (5d(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2))+b(2c(A-C)d+B(c^2-d^2))) - 5d(2aAc d-2}}{\downarrow 4008}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - 5d(A(3ac^2d-ad^3+bc^3-3bcd^2)+a(Bc^3-3Bcd^2-3c^2Cd+Cd^3))-b(3Bc^2d-Bd^3+c^3C-3cCd^2)) \int \tan}{\downarrow 3042}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - 5d(A(3ac^2d-ad^3+bc^3-3bcd^2)+a(Bc^3-3Bcd^2-3c^2Cd+Cd^3))-b(3Bc^2d-Bd^3+c^3C-3cCd^2)) \int \tan}{\downarrow 3956}$$

$$\frac{\frac{bC \tan(e+fx)(c+d \tan(e+fx))^4}{5df} - \frac{5d^2 \tan(e+fx)(2aAc d+aB(c^2-d^2)-2acCd+Ab(c^2-d^2)-b(2Bcd+c^2C-Cd^2))}{f} + \frac{5d \log(\cos(e+fx))(A(3ac^2d-ad^3+bc^3-3bcd^2)+a(Bc^3-3Bcd^2-3c^2Cd+Cd^3))-b(3Bc^2d-Bd^3+c^3C-3cCd^2))}{f}}{f}$$

input $\text{Int}[(a + b*\text{Tan}[e + f*x])*(c + d*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2),x]$

output $(b*C*\text{Tan}[e + f*x]*(c + d*\text{Tan}[e + f*x])^4)/(5*d*f) - (5*d*(a*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) + b*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)))*x + (5*d*(A*(b*c^3 + 3*a*c^2*d - 3*b*c*d^2 - a*d^3) - b*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3) + a*(B*c^3 - 3*c^2*C*d - 3*B*c*d^2 + C*d^3))*\text{Log}[\text{Cos}[e + f*x]])/f - (5*d^2*(2*a*A*c*d - 2*a*c*C*d + A*b*(c^2 - d^2) + a*B*(c^2 - d^2) - b*(c^2*C + 2*B*c*d - C*d^2))*\text{Tan}[e + f*x])/f - (5*d*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d)*(c + d*\text{Tan}[e + f*x])^2)/(2*f) - (5*(A*b + a*B - b*C)*d*(c + d*\text{Tan}[e + f*x])^3)/(3*f) + ((b*c*C - 5*b*B*d - 5*a*C*d)*(c + d*\text{Tan}[e + f*x])^4)/(4*d*f)/(5*d)$

Defintions of rubi rules used

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\text{tan}[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

rule 4008 $\text{Int}[((a_) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*c - b*d)*x, x] + (\text{Simp}[b*d*(\text{Tan}[e + f*x]/f), x] + \text{Simp}[(b*c + a*d) \text{Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[b*c + a*d, 0]$

rule 4011 $\text{Int}[((a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^m*((c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{m-1}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

Maple [A] (warning: unable to verify)

Time = 0.26 (sec) , antiderivative size = 347, normalized size of antiderivative = 0.89

method	result
parts	$\frac{(3Aa^2d + Abc^3 + Ba^3c^3) \ln(1 + \tan(fx + e))}{2f} + \frac{(Bbd^3 + Cad^3 + 3Cbc d^2) \left(\frac{\tan(fx + e)^4}{4} - \frac{\tan(fx + e)^2}{2} + \frac{\ln(1 + \tan(fx + e))}{2} \right)}{f}$
norman	$(Aa^3c^3 - 3Aac^2d^2 - 3Abc^2d + Abd^3 - 3Ba^2c^2d + Ba^3d^3 - Bbc^3 + 3Bbcd^2 - Ca^3c^3 +$
derivativedivides	$\frac{(3Aa^2d - Aa^3d^3 + Abc^3 - 3Abcd^2 + Ba^3c^3 - 3Bacd^2 - 3Bbc^2d + Bbd^3 - 3Ca^2c^2d + Ca^3d^3 - Cbc^3 + 3Cbcd^2) \ln(1 + \tan(fx + e)^2)}{2} + (Aa^3c^3 +$
default	$\frac{(3Aa^2d - Aa^3d^3 + Abc^3 - 3Abcd^2 + Ba^3c^3 - 3Bacd^2 - 3Bbc^2d + Bbd^3 - 3Ca^2c^2d + Ca^3d^3 - Cbc^3 + 3Cbcd^2) \ln(1 + \tan(fx + e)^2)}{2} + (Aa^3c^3 +$
parallelrisc	$180Cb^2c^2dx - 180Aac^2d^2fx - 180Abc^2dfx - 180Ba^2c^2dfx + 180Bbcd^2fx + 180Cac^2d^2fx - 20Cb^3d^3 \tan(fx + e)^3 + 30Aa^3d^3$
risc	Expression too large to display

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,
method=_RETURNVERBOSE)
```

output

```
1/2*(3*A*a*c^2*d+A*b*c^3+B*a*c^3)/f*ln(1+tan(f*x+e)^2)+(B*b*d^3+C*a*d^3+3*
C*b*c*d^2)/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(f*x+e)^2))+(A
*b*d^3+B*a*d^3+3*B*b*c*d^2+3*C*a*c*d^2+3*C*b*c^2*d)/f*(1/3*tan(f*x+e)^3-ta
n(f*x+e)+arctan(tan(f*x+e)))+(3*A*a*c*d^2+3*A*b*c^2*d+3*B*a*c^2*d+B*b*c^3+
C*a*c^3)/f*(tan(f*x+e)-arctan(tan(f*x+e)))+(A*a*d^3+3*A*b*c*d^2+3*B*a*c*d^
2+3*B*b*c^2*d+3*C*a*c^2*d+C*b*c^3)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)
^2))+A*a*c^3*x+C*b*d^3/f*(1/5*tan(f*x+e)^5-1/3*tan(f*x+e)^3+tan(f*x+e)-arc
tan(tan(f*x+e)))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 386, normalized size of antiderivative = 0.99

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12 C b d^3 \tan^5(fx + e) + 15 (3 C b c d^2 + (C a + B b) d^3) \tan^4(fx + e) + 20 (3 C b c^2 d + 3 (C a + B b) c d^2 + (A + B \tan(e + fx) + C \tan^2(e + fx)) d^3) \tan^3(fx + e) + 60 ((A - C) a - B b) c^3 - 3 (B a + (A - C) b) c^2 d - 3 ((A - C) a - B b) c d^2 + (B a + (A - C) b) d^3) f x + 30 (C b c^3 + 3 (C a + B b) c^2 d + 3 (B a + (A - C) b) c d^2 + ((A - C) a - B b) d^3) \tan^2(fx + e) - 30 ((B a + (A - C) b) c^3 + 3 ((A - C) a - B b) c^2 d - 3 (B a + (A - C) b) c d^2 - ((A - C) a - B b) d^3) \log(1 / (\tan(fx + e)^2 + 1)) + 60 ((C a + B b) c^3 + 3 (B a + (A - C) b) c^2 d + 3 ((A - C) a - B b) c d^2 - (B a + (A - C) b) d^3) \tan(fx + e) / f$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x, algorithm="fricas")
```

output

```
1/60*(12*C*b*d^3*tan(f*x + e)^5 + 15*(3*C*b*c*d^2 + (C*a + B*b)*d^3)*tan(f
*x + e)^4 + 20*(3*C*b*c^2*d + 3*(C*a + B*b)*c*d^2 + (B*a + (A - C)*b)*d^3)
*tan(f*x + e)^3 + 60*(((A - C)*a - B*b)*c^3 - 3*(B*a + (A - C)*b)*c^2*d -
3*((A - C)*a - B*b)*c*d^2 + (B*a + (A - C)*b)*d^3)*f*x + 30*(C*b*c^3 + 3*(
C*a + B*b)*c^2*d + 3*(B*a + (A - C)*b)*c*d^2 + ((A - C)*a - B*b)*d^3)*tan(
f*x + e)^2 - 30*((B*a + (A - C)*b)*c^3 + 3*((A - C)*a - B*b)*c^2*d - 3*(B*
a + (A - C)*b)*c*d^2 - ((A - C)*a - B*b)*d^3)*log(1/(tan(f*x + e)^2 + 1))
+ 60*((C*a + B*b)*c^3 + 3*(B*a + (A - C)*b)*c^2*d + 3*((A - C)*a - B*b)*c*
d^2 - (B*a + (A - C)*b)*d^3)*tan(f*x + e))/f
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1001 vs. $2(379) = 758$.

Time = 0.29 (sec) , antiderivative size = 1001, normalized size of antiderivative = 2.57

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Piecewise((A*a*c**3*x + 3*A*a*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) - 3*A*a*c*d**2*x + 3*A*a*c*d**2*tan(e + f*x)/f - A*a*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + A*a*d**3*tan(e + f*x)**2/(2*f) + A*b*c**3*log(tan(e + f*x)**2 + 1)/(2*f) - 3*A*b*c**2*d*x + 3*A*b*c**2*d*tan(e + f*x)/f - 3*A*b*c*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*A*b*c*d**2*tan(e + f*x)**2/(2*f) + A*b*d**3*x + A*b*d**3*tan(e + f*x)**3/(3*f) - A*b*d**3*tan(e + f*x)/f + B*a*c**3*log(tan(e + f*x)**2 + 1)/(2*f) - 3*B*a*c**2*d*x + 3*B*a*c**2*d*tan(e + f*x)/f - 3*B*a*c*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*a*c*d**2*tan(e + f*x)**2/(2*f) + B*a*d**3*x + B*a*d**3*tan(e + f*x)**3/(3*f) - B*a*d**3*tan(e + f*x)/f - B*b*c**3*x + B*b*c**3*tan(e + f*x)/f - 3*B*b*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*b*c**2*d*tan(e + f*x)**2/(2*f) + 3*B*b*c*d**2*x + B*b*c*d**2*tan(e + f*x)**3/f - 3*B*b*c*d**2*tan(e + f*x)/f + B*b*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + B*b*d**3*tan(e + f*x)**4/(4*f) - B*b*d**3*tan(e + f*x)**2/(2*f) - C*a*c**3*x + C*a*c**3*tan(e + f*x)/f - 3*C*a*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C*a*c**2*d*tan(e + f*x)**2/(2*f) + 3*C*a*c*d**2*x + C*a*c*d**2*tan(e + f*x)**3/f - 3*C*a*c*d**2*tan(e + f*x)/f + C*a*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*a*d**3*tan(e + f*x)**4/(4*f) - C*a*d**3*tan(e + f*x)**2/(2*f) - C*b*c**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*b*c**3*tan(e + f*x)**2/(2*f) + 3*C*b*c**2*d*x + C*b*c**2*d*tan(e + f*x)**3/f - 3*C*b*c**2*d*tan(e + f*x)/f + 3*C*b*c*d**2*log(tan(e + f*x...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 387, normalized size of antiderivative = 0.99

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{12 C b d^3 \tan^5(fx + e) + 15 (3 C b c d^2 + (Ca + Bb)d^3) \tan^4(fx + e) + 20 (3 C b c^2 d + 3 (Ca + Bb) c d^2 + (A + B) c^2 d^2) \tan^3(fx + e) + 15 (3 C b c^2 d + 3 (Ca + Bb) c d^2 + (A + B) c^2 d^2) \tan^2(fx + e) + 6 C b c^2 d \tan(fx + e) + 3 C b c^2 d^2}{f}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
1/60*(12*C*b*d^3*tan(f*x + e)^5 + 15*(3*C*b*c*d^2 + (C*a + B*b)*d^3)*tan(f*x + e)^4 + 20*(3*C*b*c^2*d + 3*(C*a + B*b)*c*d^2 + (B*a + (A - C)*b)*d^3)*tan(f*x + e)^3 + 30*(C*b*c^3 + 3*(C*a + B*b)*c^2*d + 3*(B*a + (A - C)*b)*c*d^2 + ((A - C)*a - B*b)*d^3)*tan(f*x + e)^2 + 60*(((A - C)*a - B*b)*c^3 - 3*(B*a + (A - C)*b)*c^2*d - 3*((A - C)*a - B*b)*c*d^2 + (B*a + (A - C)*b)*d^3)*(f*x + e) + 30*((B*a + (A - C)*b)*c^3 + 3*((A - C)*a - B*b)*c^2*d - 3*(B*a + (A - C)*b)*c*d^2 - ((A - C)*a - B*b)*d^3)*log(tan(f*x + e)^2 + 1) + 60*((C*a + B*b)*c^3 + 3*(B*a + (A - C)*b)*c^2*d + 3*((A - C)*a - B*b)*c*d^2 - (B*a + (A - C)*b)*d^3)*tan(f*x + e))/f
```

Giac [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 740, normalized size of antiderivative = 1.90

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
(A*a*c^3 - C*a*c^3 - B*b*c^3 - 3*B*a*c^2*d - 3*A*b*c^2*d + 3*C*b*c^2*d - 3
*A*a*c*d^2 + 3*C*a*c*d^2 + 3*B*b*c*d^2 + B*a*d^3 + A*b*d^3 - C*b*d^3)*(f*x
+ e)/f + 1/2*(B*a*c^3 + A*b*c^3 - C*b*c^3 + 3*A*a*c^2*d - 3*C*a*c^2*d - 3
*B*b*c^2*d - 3*B*a*c*d^2 - 3*A*b*c*d^2 + 3*C*b*c*d^2 - A*a*d^3 + C*a*d^3 +
B*b*d^3)*log(tan(f*x + e)^2 + 1)/f + 1/60*(12*C*b*d^3*f^4*tan(f*x + e)^5
+ 45*C*b*c*d^2*f^4*tan(f*x + e)^4 + 15*C*a*d^3*f^4*tan(f*x + e)^4 + 15*B*b
*d^3*f^4*tan(f*x + e)^4 + 60*C*b*c^2*d*f^4*tan(f*x + e)^3 + 60*C*a*c*d^2*f
^4*tan(f*x + e)^3 + 60*B*b*c*d^2*f^4*tan(f*x + e)^3 + 20*B*a*d^3*f^4*tan(f
*x + e)^3 + 20*A*b*d^3*f^4*tan(f*x + e)^3 - 20*C*b*d^3*f^4*tan(f*x + e)^3
+ 30*C*b*c^3*f^4*tan(f*x + e)^2 + 90*C*a*c^2*d*f^4*tan(f*x + e)^2 + 90*B*b
*c^2*d*f^4*tan(f*x + e)^2 + 90*B*a*c*d^2*f^4*tan(f*x + e)^2 + 90*A*b*c*d^2
*f^4*tan(f*x + e)^2 - 90*C*b*c*d^2*f^4*tan(f*x + e)^2 + 30*A*a*d^3*f^4*tan
(f*x + e)^2 - 30*C*a*d^3*f^4*tan(f*x + e)^2 - 30*B*b*d^3*f^4*tan(f*x + e)^
2 + 60*C*a*c^3*f^4*tan(f*x + e) + 60*B*b*c^3*f^4*tan(f*x + e) + 180*B*a*c^
2*d*f^4*tan(f*x + e) + 180*A*b*c^2*d*f^4*tan(f*x + e) - 180*C*b*c^2*d*f^4*
tan(f*x + e) + 180*A*a*c*d^2*f^4*tan(f*x + e) - 180*C*a*c*d^2*f^4*tan(f*x
+ e) - 180*B*b*c*d^2*f^4*tan(f*x + e) - 60*B*a*d^3*f^4*tan(f*x + e) - 60*A
*b*d^3*f^4*tan(f*x + e) + 60*C*b*d^3*f^4*tan(f*x + e))/f^5
```

Mupad [B] (verification not implemented)

Time = 5.78 (sec) , antiderivative size = 478, normalized size of antiderivative = 1.23

$$\begin{aligned}
& \int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
& = x (A a c^3 + A b d^3 + B a d^3 - B b c^3 - C a c^3 - C b d^3 - 3 A a c d^2 - 3 A b c^2 d - 3 B a c^2 d \\
& \quad + 3 B b c d^2 + 3 C a c d^2 + 3 C b c^2 d) + \frac{\tan(e + fx)^4 \left(\frac{B b d^3}{4} + \frac{C a d^3}{4} + \frac{3 C b c d^2}{4} \right)}{f} \\
& \quad + \frac{\tan(e + fx)^3 \left(\frac{A b d^3}{3} + \frac{B a d^3}{3} - \frac{C b d^3}{3} + B b c d^2 + C a c d^2 + C b c^2 d \right)}{f} \\
& \quad + \frac{\tan(e + fx)^2 \left(\frac{A a d^3}{2} - \frac{B b d^3}{2} - \frac{C a d^3}{2} + \frac{C b c^3}{2} + \frac{3 A b c d^2}{2} + \frac{3 B a c d^2}{2} + \frac{3 B b c^2 d}{2} + \frac{3 C a c^2 d}{2} - \frac{3 C b c d^2}{2} \right)}{f} \\
& \quad - \frac{\ln(\tan(e + fx)^2 + 1) \left(\frac{A a d^3}{2} - \frac{A b c^3}{2} - \frac{B a c^3}{2} - \frac{B b d^3}{2} - \frac{C a d^3}{2} + \frac{C b c^3}{2} - \frac{3 A a c^2 d}{2} + \frac{3 A b c d^2}{2} + \frac{3 B a c d^2}{2} + \frac{3 B b c^2 d}{2} - \frac{3 C a c^2 d}{2} - \frac{3 C b c d^2}{2} \right)}{f} \\
& \quad + \frac{\tan(e + fx) (B b c^3 - B a d^3 - A b d^3 + C a c^3 + C b d^3 + 3 A a c d^2 + 3 A b c^2 d + 3 B a c^2 d - 3 B b c d^2)}{f} \\
& \quad + \frac{C b d^3 \tan(e + fx)^5}{5 f}
\end{aligned}$$

input

```
int((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
x*(A*a*c^3 + A*b*d^3 + B*a*d^3 - B*b*c^3 - C*a*c^3 - C*b*d^3 - 3*A*a*c*d^2 - 3*A*b*c^2*d - 3*B*a*c^2*d + 3*B*b*c*d^2 + 3*C*a*c*d^2 + 3*C*b*c^2*d) + (tan(e + f*x)^4*((B*b*d^3)/4 + (C*a*d^3)/4 + (3*C*b*c*d^2)/4))/f + (tan(e + f*x)^3*((A*b*d^3)/3 + (B*a*d^3)/3 - (C*b*d^3)/3 + B*b*c*d^2 + C*a*c*d^2 + C*b*c^2*d))/f + (tan(e + f*x)^2*((A*a*d^3)/2 - (B*b*d^3)/2 - (C*a*d^3)/2 + (C*b*c^3)/2 + (3*A*b*c*d^2)/2 + (3*B*a*c*d^2)/2 + (3*B*b*c^2*d)/2 + (3*C*a*c^2*d)/2 - (3*C*b*c*d^2)/2))/f - (log(tan(e + f*x)^2 + 1)*((A*a*d^3)/2 - (A*b*c^3)/2 - (B*a*c^3)/2 - (B*b*d^3)/2 - (C*a*d^3)/2 + (C*b*c^3)/2 - (3*A*a*c^2*d)/2 + (3*A*b*c*d^2)/2 + (3*B*a*c*d^2)/2 + (3*B*b*c^2*d)/2 + (3*C*a*c^2*d)/2 - (3*C*b*c*d^2)/2))/f + (tan(e + f*x)*(B*b*c^3 - B*a*d^3 - A*b*d^3 + C*a*c^3 + C*b*d^3 + 3*A*a*c*d^2 + 3*A*b*c^2*d + 3*B*a*c^2*d - 3*B*b*c*d^2 - 3*C*a*c*d^2 - 3*C*b*c^2*d))/f + (C*b*d^3*tan(e + f*x)^5)/(5*f)
```


Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 684, normalized size of antiderivative = 1.76

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
(90*log(tan(e + f*x)**2 + 1)*a**2*c**2*d - 30*log(tan(e + f*x)**2 + 1)*a**
2*d**3 + 60*log(tan(e + f*x)**2 + 1)*a*b*c**3 - 180*log(tan(e + f*x)**2 +
1)*a*b*c*d**2 - 90*log(tan(e + f*x)**2 + 1)*a*c**3*d + 30*log(tan(e + f*x)
**2 + 1)*a*c*d**3 - 90*log(tan(e + f*x)**2 + 1)*b**2*c**2*d + 30*log(tan(e
+ f*x)**2 + 1)*b**2*d**3 - 30*log(tan(e + f*x)**2 + 1)*b*c**4 + 90*log(ta
n(e + f*x)**2 + 1)*b*c**2*d**2 + 12*tan(e + f*x)**5*b*c*d**3 + 15*tan(e +
f*x)**4*a*c*d**3 + 15*tan(e + f*x)**4*b**2*d**3 + 45*tan(e + f*x)**4*b*c**
2*d**2 + 40*tan(e + f*x)**3*a*b*d**3 + 60*tan(e + f*x)**3*a*c**2*d**2 + 60
*tan(e + f*x)**3*b**2*c*d**2 + 60*tan(e + f*x)**3*b*c**3*d - 20*tan(e + f*
x)**3*b*c*d**3 + 30*tan(e + f*x)**2*a**2*d**3 + 180*tan(e + f*x)**2*a*b*c*
d**2 + 90*tan(e + f*x)**2*a*c**3*d - 30*tan(e + f*x)**2*a*c*d**3 + 90*tan(
e + f*x)**2*b**2*c**2*d - 30*tan(e + f*x)**2*b**2*d**3 + 30*tan(e + f*x)**
2*b*c**4 - 90*tan(e + f*x)**2*b*c**2*d**2 + 180*tan(e + f*x)*a**2*c*d**2 +
360*tan(e + f*x)*a*b*c**2*d - 120*tan(e + f*x)*a*b*d**3 + 60*tan(e + f*x)
*a*c**4 - 180*tan(e + f*x)*a*c**2*d**2 + 60*tan(e + f*x)*b**2*c**3 - 180*t
an(e + f*x)*b**2*c*d**2 - 180*tan(e + f*x)*b*c**3*d + 60*tan(e + f*x)*b*c*
d**3 + 60*a**2*c**3*f*x - 180*a**2*c*d**2*f*x - 360*a*b*c**2*d*f*x + 120*a
*b*d**3*f*x - 60*a*c**4*f*x + 180*a*c**2*d**2*f*x - 60*b**2*c**3*f*x + 180
*b**2*c*d**2*f*x + 180*b*c**3*d*f*x - 60*b*c*d**3*f*x)/(60*f)
```

3.66 $\int (c+d \tan(e+fx))^3 (A + B \tan(e + fx) + C \tan^2(e -$

Optimal result	725
Mathematica [C] (verified)	726
Rubi [A] (verified)	726
Maple [A] (warning: unable to verify)	729
Fricas [A] (verification not implemented)	730
Sympy [B] (verification not implemented)	730
Maxima [A] (verification not implemented)	731
Giac [A] (verification not implemented)	732
Mupad [B] (verification not implemented)	733
Reduce [B] (verification not implemented)	733

Optimal result

Integrand size = 33, antiderivative size = 191

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= -((c^3C + 3Bc^2d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2))x$$

$$- \frac{((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) \log(\cos(e + fx))}{f}$$

$$+ \frac{d(2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)}{f} + \frac{(Bc + (A - C)d)(c + d \tan(e + fx))^2}{2f}$$

$$+ \frac{B(c + d \tan(e + fx))^3}{3f} + \frac{C(c + d \tan(e + fx))^4}{4df}$$

output

```
-(c^3*C+3*B*c^2*d-3*C*c*d^2-B*d^3-A*(c^3-3*c*d^2))*x-((A-C)*d*(3*c^2-d^2)+
B*(c^3-3*c*d^2))*ln(cos(f*x+e))/f+d*(2*c*(A-C)*d+B*(c^2-d^2))*tan(f*x+e)/f
+1/2*(B*c+(A-C)*d)*(c+d*tan(f*x+e))^2/f+1/3*B*(c+d*tan(f*x+e))^3/f+1/4*C*(
c+d*tan(f*x+e))^4/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.64 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.11

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3C(c + d \tan(e + fx))^4 - 6(Bc + (-A + C)d)((ic - d)^3 \log(i - \tan(e + fx)) - (ic + d)^3 \log(i + \tan(e + fx)))}{12df}$$

input

```
Integrate[(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(3*C*(c + d*Tan[e + f*x])^4 - 6*(B*c + (-A + C)*d)*((I*c - d)^3*Log[I - Tan[e + f*x]] - (I*c + d)^3*Log[I + Tan[e + f*x]] + 6*c*d^2*Tan[e + f*x] + d^3*Tan[e + f*x]^2) + 2*B*((-3*I)*(c + I*d)^4*Log[I - Tan[e + f*x]] + (3*I)*(c - I*d)^4*Log[I + Tan[e + f*x]] - 6*d^2*(-6*c^2 + d^2)*Tan[e + f*x] + 12*c*d^3*Tan[e + f*x]^2 + 2*d^4*Tan[e + f*x]^3)/(12*d*f)
```

Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3042, 4113, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow \text{4113}$$

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^3 dx + \frac{C(c + d \tan(e + fx))^4}{4df}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^3 dx + \frac{C(c + d \tan(e + fx))^4}{4df} \\
& \downarrow 4011 \\
& \int (c + d \tan(e + fx))^2 (Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \\
& \quad \frac{B(c + d \tan(e + fx))^3}{3f} + \frac{C(c + d \tan(e + fx))^4}{4df} \\
& \downarrow 3042 \\
& \int (c + d \tan(e + fx))^2 (Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \\
& \quad \frac{B(c + d \tan(e + fx))^3}{3f} + \frac{C(c + d \tan(e + fx))^4}{4df} \\
& \downarrow 4011 \\
& \int (c + d \tan(e + \\
& fx)) (-Cc^2 - 2Bdc + Cd^2 + A(c^2 - d^2) + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)) dx + \\
& \quad \frac{(d(A - C) + Bc)(c + d \tan(e + fx))^2}{2f} + \frac{B(c + d \tan(e + fx))^3}{3f} + \frac{C(c + d \tan(e + fx))^4}{4df} \\
& \downarrow 3042 \\
& \int (c + d \tan(e + \\
& fx)) (-Cc^2 - 2Bdc + Cd^2 + A(c^2 - d^2) + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)) dx + \\
& \quad \frac{(d(A - C) + Bc)(c + d \tan(e + fx))^2}{2f} + \frac{B(c + d \tan(e + fx))^3}{3f} + \frac{C(c + d \tan(e + fx))^4}{4df} \\
& \downarrow 4008 \\
& \quad (d(A - C)(3c^2 - d^2) + B(c^3 - 3cd^2)) \int \tan(e + fx) dx + \\
& \quad \frac{d \tan(e + fx) (2cd(A - C) + B(c^2 - d^2))}{f} - \\
& x(-A(c^3 - 3cd^2) + 3Bc^2d - Bd^3 + c^3C - 3cCd^2) + \frac{(d(A - C) + Bc)(c + d \tan(e + fx))^2}{2f} + \\
& \quad \frac{B(c + d \tan(e + fx))^3}{3f} + \frac{C(c + d \tan(e + fx))^4}{4df} \\
& \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& \frac{(d(A-C)(3c^2-d^2) + B(c^3-3cd^2)) \int \tan(e+fx) dx +}{f} \\
& \frac{d \tan(e+fx) (2cd(A-C) + B(c^2-d^2))}{f} - \\
& x(-A(c^3-3cd^2) + 3Bc^2d - Bd^3 + c^3C - 3cCd^2) + \frac{(d(A-C) + Bc)(c + d \tan(e+fx))^2}{2f} + \\
& \frac{B(c + d \tan(e+fx))^3}{3f} + \frac{C(c + d \tan(e+fx))^4}{4df} \\
& \quad \downarrow \text{3956} \\
& \frac{d \tan(e+fx) (2cd(A-C) + B(c^2-d^2))}{f} - \\
& \frac{(d(A-C)(3c^2-d^2) + B(c^3-3cd^2)) \log(\cos(e+fx))}{f} - \\
& x(-A(c^3-3cd^2) + 3Bc^2d - Bd^3 + c^3C - 3cCd^2) + \frac{(d(A-C) + Bc)(c + d \tan(e+fx))^2}{2f} + \\
& \frac{B(c + d \tan(e+fx))^3}{3f} + \frac{C(c + d \tan(e+fx))^4}{4df}
\end{aligned}$$

input `Int[(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `-((c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2))*x) - (((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2))*Log[Cos[e + f*x]])/f + (d*(2*c*(A - C)*d + B*(c^2 - d^2))*Tan[e + f*x])/f + ((B*c + (A - C)*d)*(c + d*Tan[e + f*x])^2)/(2*f) + (B*(c + d*Tan[e + f*x])^3)/(3*f) + (C*(c + d*Tan[e + f*x])^4)/(4*d*f)`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4008 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

```
rule 4011 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4113 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_)
+ (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [A] (warning: unable to verify)

Time = 0.15 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.10

method	result
parts	$A c^3 x + \frac{(3A c^2 d + B c^3) \ln(1 + \tan(fx + e)^2)}{2f} + \frac{(B d^3 + 3C c d^2) \left(\frac{\tan(fx + e)^3}{3} - \tan(fx + e) + \arctan(\tan(fx + e)) \right)}{f}$
norman	$(A c^3 - 3A c d^2 - 3B c^2 d + B d^3 - c^3 C + 3C c d^2) x + \frac{(3A c d^2 + 3B c^2 d - B d^3 + c^3 C - 3C c d^2) \tan(fx + e)}{f}$
derivativdivides	$\frac{C d^3 \tan(fx + e)^4}{4} + \frac{B d^3 \tan(fx + e)^3}{3} + C c d^2 \tan(fx + e)^3 + \frac{A d^3 \tan(fx + e)^2}{2} + \frac{3B c d^2 \tan(fx + e)^2}{2} + \frac{3C c^2 d \tan(fx + e)^2}{2} - \frac{C d^3}{2}$
default	$\frac{C d^3 \tan(fx + e)^4}{4} + \frac{B d^3 \tan(fx + e)^3}{3} + C c d^2 \tan(fx + e)^3 + \frac{A d^3 \tan(fx + e)^2}{2} + \frac{3B c d^2 \tan(fx + e)^2}{2} + \frac{3C c^2 d \tan(fx + e)^2}{2} - \frac{C d^3}{2}$
parallelrisc	$3C d^3 \tan(fx + e)^4 + 4B d^3 \tan(fx + e)^3 + 6A d^3 \tan(fx + e)^2 - 6C d^3 \tan(fx + e)^2 - 12 \tan(fx + e) B d^3 + 12 \tan(fx + e) c^3 C$
risc	$-3A c d^2 x - 3B c^2 dx + 3C c d^2 x + 3i A c^2 dx - \frac{2i A d^3 e}{f} + \frac{2i B c^3 e}{f} + \frac{2i C d^3 e}{f} - \frac{3 \ln(e^{2i(fx + e)} + 1)}{f}$

input `int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `A*c^3*x+1/2*(3*A*c^2*d+B*c^3)/f*ln(1+tan(f*x+e)^2)+(B*d^3+3*C*c*d^2)/f*(1/3*tan(f*x+e)^3-tan(f*x+e)+arctan(tan(f*x+e)))+(A*d^3+3*B*c*d^2+3*C*c^2*d)/f*(1/2*tan(f*x+e)^2-1/2*ln(1+tan(f*x+e)^2))+(3*A*c*d^2+3*B*c^2*d+C*c^3)/f*(tan(f*x+e)-arctan(tan(f*x+e)))+C*d^3/f*(1/4*tan(f*x+e)^4-1/2*tan(f*x+e)^2+1/2*ln(1+tan(f*x+e)^2))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.05

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3Cd^3 \tan^4(fx + e) + 4(3Ccd^2 + Bd^3) \tan^3(fx + e) + 12((A - C)c^3 - 3Bc^2d - 3(A - C)cd^2 + Bd^3)}{f}$$

input `integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `1/12*(3*C*d^3*tan(f*x + e)^4 + 4*(3*C*c*d^2 + B*d^3)*tan(f*x + e)^3 + 12*((A - C)*c^3 - 3*B*c^2*d - 3*(A - C)*c*d^2 + B*d^3)*f*x + 6*(3*C*c^2*d + 3*B*c*d^2 + (A - C)*d^3)*tan(f*x + e)^2 - 6*(B*c^3 + 3*(A - C)*c^2*d - 3*B*c*d^2 - (A - C)*d^3)*log(1/(tan(f*x + e)^2 + 1)) + 12*(C*c^3 + 3*B*c^2*d + 3*(A - C)*c*d^2 - B*d^3)*tan(f*x + e))/f`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. 2(163) = 326.

Time = 0.23 (sec) , antiderivative size = 410, normalized size of antiderivative = 2.15

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \begin{cases} Ac^3x + \frac{3Ac^2d \log(\tan^2(e+fx)+1)}{2f} - 3Acd^2x + \frac{3Acd^2 \tan(e+fx)}{f} - \frac{Ad^3 \log(\tan^2(e+fx)+1)}{2f} + \frac{Ad^3 \tan^2(e+fx)}{2f} + \frac{Bc^3 \log(\tan^2(e+fx)+1)}{2f} \\ x(c + d \tan(e))^3 (A + B \tan(e) + C \tan^2(e)) \end{cases}$$

input `integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Piecewise((A*c**3*x + 3*A*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) - 3*A*c*d**2*x + 3*A*c*d**2*tan(e + f*x)/f - A*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + A*d**3*tan(e + f*x)**2/(2*f) + B*c**3*log(tan(e + f*x)**2 + 1)/(2*f) - 3*B*c**2*d*x + 3*B*c**2*d*tan(e + f*x)/f - 3*B*c*d**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*c*d**2*tan(e + f*x)**2/(2*f) + B*d**3*x + B*d**3*tan(e + f*x)**3/(3*f) - B*d**3*tan(e + f*x)/f - C*c**3*x + C*c**3*tan(e + f*x)/f - 3*C*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C*c**2*d*tan(e + f*x)**2/(2*f) + 3*C*c*d**2*x + C*c*d**2*tan(e + f*x)**3/f - 3*C*c*d**2*tan(e + f*x)/f + C*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*d**3*tan(e + f*x)**4/(4*f) - C*d**3*tan(e + f*x)**2/(2*f), Ne(f, 0)), (x*(c + d*tan(e))^3*(A + B*tan(e) + C*tan(e)**2), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.06

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{3Cd^3 \tan^4(fx + e) + 4(3Ccd^2 + Bd^3) \tan^3(fx + e) + 6(3Cc^2d + 3Bcd^2 + (A - C)d^3) \tan^2(fx + e) + \dots}{\dots}$$

input `integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output

```
1/12*(3*C*d^3*tan(f*x + e)^4 + 4*(3*C*c*d^2 + B*d^3)*tan(f*x + e)^3 + 6*(3
*C*c^2*d + 3*B*c*d^2 + (A - C)*d^3)*tan(f*x + e)^2 + 12*((A - C)*c^3 - 3*B
*c^2*d - 3*(A - C)*c*d^2 + B*d^3)*(f*x + e) + 6*(B*c^3 + 3*(A - C)*c^2*d -
3*B*c*d^2 - (A - C)*d^3)*log(tan(f*x + e)^2 + 1) + 12*(C*c^3 + 3*B*c^2*d
+ 3*(A - C)*c*d^2 - B*d^3)*tan(f*x + e))/f
```

Giac [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.61

$$\int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(Ac^3 - Cc^3 - 3Bc^2d - 3Acd^2 + 3Ccd^2 + Bd^3)(fx + e)}{f}$$

$$+ \frac{(Bc^3 + 3Ac^2d - 3Cc^2d - 3Bcd^2 - Ad^3 + Cd^3) \log(\tan(fx + e)^2 + 1)}{2f}$$

$$+ \frac{3Cd^3f^3 \tan(fx + e)^4 + 12Ccd^2f^3 \tan(fx + e)^3 + 4Bd^3f^3 \tan(fx + e)^3 + 18Cc^2df^3 \tan(fx + e)^2 -$$

input

```
integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm=
"giac")
```

output

```
(A*c^3 - C*c^3 - 3*B*c^2*d - 3*A*c*d^2 + 3*C*c*d^2 + B*d^3)*(f*x + e)/f +
1/2*(B*c^3 + 3*A*c^2*d - 3*C*c^2*d - 3*B*c*d^2 - A*d^3 + C*d^3)*log(tan(f*
x + e)^2 + 1)/f + 1/12*(3*C*d^3*f^3*tan(f*x + e)^4 + 12*C*c*d^2*f^3*tan(f*
x + e)^3 + 4*B*d^3*f^3*tan(f*x + e)^3 + 18*C*c^2*d*f^3*tan(f*x + e)^2 + 18
*B*c*d^2*f^3*tan(f*x + e)^2 + 6*A*d^3*f^3*tan(f*x + e)^2 - 6*C*d^3*f^3*tan
(f*x + e)^2 + 12*C*c^3*f^3*tan(f*x + e) + 36*B*c^2*d*f^3*tan(f*x + e) + 36
*A*c*d^2*f^3*tan(f*x + e) - 36*C*c*d^2*f^3*tan(f*x + e) - 12*B*d^3*f^3*tan
(f*x + e))/f^4
```

Mupad [B] (verification not implemented)

Time = 5.54 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.16

$$\begin{aligned}
& \int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= x (A c^3 + B d^3 - C c^3 - 3 A c d^2 - 3 B c^2 d + 3 C c d^2) \\
&+ \frac{\tan(e + fx) (C c^3 - B d^3 + 3 A c d^2 + 3 B c^2 d - 3 C c d^2)}{f} \\
&+ \frac{\tan(e + fx)^3 \left(\frac{B d^3}{3} + C c d^2 \right)}{f} \\
&- \frac{\ln(\tan(e + fx)^2 + 1) \left(\frac{A d^3}{2} - \frac{B c^3}{2} - \frac{C d^3}{2} - \frac{3 A c^2 d}{2} + \frac{3 B c d^2}{2} + \frac{3 C c^2 d}{2} \right)}{f} \\
&+ \frac{\tan(e + fx)^2 \left(\frac{A d^3}{2} - \frac{C d^3}{2} + \frac{3 B c d^2}{2} + \frac{3 C c^2 d}{2} \right)}{f} + \frac{C d^3 \tan(e + fx)^4}{4 f}
\end{aligned}$$

input `int((c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `x*(A*c^3 + B*d^3 - C*c^3 - 3*A*c*d^2 - 3*B*c^2*d + 3*C*c*d^2) + (tan(e + f*x)*(C*c^3 - B*d^3 + 3*A*c*d^2 + 3*B*c^2*d - 3*C*c*d^2))/f + (tan(e + f*x)^3*((B*d^3)/3 + C*c*d^2))/f - (log(tan(e + f*x)^2 + 1)*((A*d^3)/2 - (B*c^3)/2 - (C*d^3)/2 - (3*A*c^2*d)/2 + (3*B*c*d^2)/2 + (3*C*c^2*d)/2))/f + (tan(e + f*x)^2*((A*d^3)/2 - (C*d^3)/2 + (3*B*c*d^2)/2 + (3*C*c^2*d)/2))/f + (C*d^3*tan(e + f*x)^4)/(4*f)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.70

$$\begin{aligned}
& \int (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= \frac{18 \log(\tan(fx + e)^2 + 1) a c^2 d - 6 \log(\tan(fx + e)^2 + 1) a d^3 + 6 \log(\tan(fx + e)^2 + 1) b c^3 - 18 \log(\tan(fx + e)^2 + 1) b c^2 d - 6 \log(\tan(fx + e)^2 + 1) b c d^2 + 6 \log(\tan(fx + e)^2 + 1) b d^3}{f}
\end{aligned}$$

input `int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output

```
(18*log(tan(e + f*x)**2 + 1)*a*c**2*d - 6*log(tan(e + f*x)**2 + 1)*a*d**3
+ 6*log(tan(e + f*x)**2 + 1)*b*c**3 - 18*log(tan(e + f*x)**2 + 1)*b*c*d**2
- 18*log(tan(e + f*x)**2 + 1)*c**3*d + 6*log(tan(e + f*x)**2 + 1)*c*d**3
+ 3*tan(e + f*x)**4*c*d**3 + 4*tan(e + f*x)**3*b*d**3 + 12*tan(e + f*x)**3
*c**2*d**2 + 6*tan(e + f*x)**2*a*d**3 + 18*tan(e + f*x)**2*b*c*d**2 + 18*t
an(e + f*x)**2*c**3*d - 6*tan(e + f*x)**2*c*d**3 + 36*tan(e + f*x)*a*c*d**
2 + 36*tan(e + f*x)*b*c**2*d - 12*tan(e + f*x)*b*d**3 + 12*tan(e + f*x)*c*
*4 - 36*tan(e + f*x)*c**2*d**2 + 12*a*c**3*f*x - 36*a*c*d**2*f*x - 36*b*c*
*2*d*f*x + 12*b*d**3*f*x - 12*c**4*f*x + 36*c**2*d**2*f*x)/(12*f)
```

3.67
$$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$$

Optimal result	735
Mathematica [C] (verified)	736
Rubi [A] (verified)	736
Maple [A] (verified)	741
Fricas [A] (verification not implemented)	742
Sympy [C] (verification not implemented)	743
Maxima [A] (verification not implemented)	744
Giac [A] (verification not implemented)	745
Mupad [B] (verification not implemented)	746
Reduce [B] (verification not implemented)	747

Optimal result

Integrand size = 45, antiderivative size = 363

$$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx =$$

$$\frac{(a(c^3C + 3Bc^2d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2)) - b((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))) x}{a^2 + b^2}$$

$$\frac{(b(c^3C + 3Bc^2d - 3cCd^2 - Bd^3) + a(Bc^3 - 3c^2Cd - 3Bcd^2 + Cd^3) + A(ad(3c^2 - d^2) - b(c^3 - 3cd^2)))}{(a^2 + b^2) f}$$

$$+ \frac{(Ab^2 - a(bB - aC)) (bc - ad)^3 \log(a + b \tan(e + fx))}{b^4 (a^2 + b^2) f}$$

$$+ \frac{d(b^2d(Bc + (A - C)d) + (bc - ad)(bcC + bBd - aCd)) \tan(e + fx)}{b^3 f}$$

$$+ \frac{(bcC + bBd - aCd)(c + d \tan(e + fx))^2}{2b^2 f} + \frac{C(c + d \tan(e + fx))^3}{3bf}$$

output

```
-(a*(c^3*C+3*B*c^2*d-3*C*c*d^2-B*d^3-A*(c^3-3*c*d^2))-b*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*d^2)))*x/(a^2+b^2)-(b*(3*B*c^2*d-B*d^3+C*c^3-3*C*c*d^2)+a*(B*c^3-3*B*c*d^2-3*C*c^2*d+C*d^3)+A*(a*d*(3*c^2-d^2)-b*(c^3-3*c*d^2)))*ln(cos(f*x+e))/(a^2+b^2)/f+(A*b^2-a*(B*b-C*a))*(-a*d+b*c)^3*ln(a+b*tan(f*x+e))/b^4/(a^2+b^2)/f+d*(b^2*d*(B*c+(A-C)*d)+(-a*d+b*c)*(B*b*d-C*a*d+C*b*c))*tan(f*x+e)/b^3/f+1/2*(B*b*d-C*a*d+C*b*c)*(c+d*tan(f*x+e))^2/b^2/f+1/3*C*(c+d*tan(f*x+e))^3/b/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.00 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.70

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{3b^2(-iA+B+iC)(c+id)^3 \log(i-\tan(e+fx))}{a+ib} - \frac{3b^2(A-iB-C)(ic+d)^3 \log(i+\tan(e+fx))}{a-ib} + \frac{6(Ab^2+a(-bB+aC))(bc-ad)^3 \log(a+b \tan(e+fx))}{b^2(a^2+b^2)}$$

input

```
Integrate[((c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(a + b*Tan[e + f*x]),x]
```

output

```
((3*b^2*((-I)*A + B + I*C)*(c + I*d)^3*Log[I - Tan[e + f*x]])/(a + I*b) -
(3*b^2*(A - I*B - C)*(I*c + d)^3*Log[I + Tan[e + f*x]])/(a - I*b) + (6*(A*
b^2 + a*(-(b*B) + a*C))*(b*c - a*d)^3*Log[a + b*Tan[e + f*x]])/(b^2*(a^2 +
b^2)) + 6*b*d^2*(B*c + (A - C)*d)*Tan[e + f*x] + (6*d*(b*c - a*d)*(b*c*C
+ b*B*d - a*C*d)*Tan[e + f*x])/b + 3*(b*c*C + b*B*d - a*C*d)*(c + d*Tan[e
+ f*x])^2 + 2*b*C*(c + d*Tan[e + f*x])^3)/(6*b^2*f)
```

Rubi [A] (verified)

Time = 4.02 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4120, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{a + b \tan(e + fx)} dx$$

$$\begin{aligned}
 & \int \frac{3(c+d \tan(e+fx))^2((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{3b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 4130 \\
 & \int \frac{(c+d \tan(e+fx))^2((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 27 \\
 & \int \frac{(c+d \tan(e+fx))^2((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 3042 \\
 & \int \frac{(c+d \tan(e+fx))^2((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 4130 \\
 & \int \frac{2(c+d \tan(e+fx))(Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+(d(Bc+(A-C)d)b^2+(bc-ad)(bcC-adC+bBd) \tan^2(e+fx)+ad(aCd-b(2cC+Bd)))}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 27 \\
 & \int \frac{(c+d \tan(e+fx))(Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+(d(Bc+(A-C)d)b^2+(bc-ad)(bcC-adC+bBd) \tan^2(e+fx)+ad(aCd-b(2cC+Bd)))}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 3042 \\
 & \int \frac{(c+d \tan(e+fx))(Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+(d(Bc+(A-C)d)b^2+(bc-ad)(bcC-adC+bBd) \tan^2(e+fx)+ad(aCd-b(2cC+Bd)))}{a+b \tan(e+fx)} dx + \\
 & \quad \frac{b}{3bf} C(c+d \tan(e+fx))^3 \\
 & \quad \downarrow 4120
 \end{aligned}$$

$$\frac{d \tan(e+fx) \left((bc-ad)(-aCd+bBd+bcC)+b^2d(d(A-C)+Bc) \right)}{bf} \int - \frac{\left((A-C)d(3c^2-d^2)+B(c^3-3cd^2) \right) \tan(e+fx)b^3+A(bc^3-ad^3)b^2-\left((Cc^3+3Bdc^2+3(A-C)d^2c-Bd^3) \right) b^3}{b}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 25

$$\int \frac{\left((A-C)d(3c^2-d^2)+B(c^3-3cd^2) \right) \tan(e+fx)b^3+A(bc^3-ad^3)b^2-\left((Cc^3+3Bdc^2+3(A-C)d^2c-Bd^3) \right) b^3+ad(3Cc^2+3Bdc+(A-C)d^2)b^2-a^2d^2(3cC+Bd)}{a+b \tan(e+fx)} \frac{1}{b}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 3042

$$\int \frac{\left((A-C)d(3c^2-d^2)+B(c^3-3cd^2) \right) \tan(e+fx)b^3+A(bc^3-ad^3)b^2-\left((Cc^3+3Bdc^2+3(A-C)d^2c-Bd^3) \right) b^3+ad(3Cc^2+3Bdc+(A-C)d^2)b^2-a^2d^2(3cC+Bd)}{a+b \tan(e+fx)} \frac{1}{b}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 4109

$$\frac{(bc-ad)^3(Ab^2-a(bB-aC)) \int \frac{\tan^2(e+fx)+1}{a+b \tan(e+fx)} dx}{a^2+b^2} + \frac{b^3(aAd(3c^2-d^2)-a(Cd(3c^2-d^2)-B(c^3-3cd^2))-Ab(c^3-3cd^2)+b(3Bc^2d-Bd^3+c^3C-3cCd^2)) \int \tan(e+fx)}{a^2+b^2} \frac{1}{b}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 3042

$$\frac{(bc-ad)^3(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{a+b \tan(e+fx)} dx}{a^2+b^2} + \frac{b^3(aAd(3c^2-d^2)-a(Cd(3c^2-d^2)-B(c^3-3cd^2))-Ab(c^3-3cd^2)+b(3Bc^2d-Bd^3+c^3C-3cCd^2)) \int \tan(e+fx)}{a^2+b^2} \frac{1}{b}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 3956

$$\frac{(bc-ad)^3(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{a+b \tan(e+fx)} dx}{a^2+b^2} - \frac{b^3 \log(\cos(e+fx))(aAd(3c^2-d^2)-a(Cd(3c^2-d^2)-B(c^3-3cd^2))-Ab(c^3-3cd^2))+b(3Bc^2d-Bd^3+c^3C-3)}{f(a^2+b^2)}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 4100

$$\frac{(bc-ad)^3(Ab^2-a(bB-aC)) \int \frac{1}{a+b \tan(e+fx)} d(b \tan(e+fx))}{bf(a^2+b^2)} - \frac{b^3 \log(\cos(e+fx))(aAd(3c^2-d^2)-a(Cd(3c^2-d^2)-B(c^3-3cd^2))-Ab(c^3-3cd^2))+b(3Bc^2d-Bd^3+c^3C-3)}{f(a^2+b^2)}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

↓ 16

$$\frac{(bc-ad)^3(Ab^2-a(bB-aC)) \log(a+b \tan(e+fx))}{bf(a^2+b^2)} - \frac{b^3 \log(\cos(e+fx))(aAd(3c^2-d^2)-a(Cd(3c^2-d^2)-B(c^3-3cd^2))-Ab(c^3-3cd^2))+b(3Bc^2d-Bd^3+c^3C-3)}{f(a^2+b^2)}$$

$$\frac{C(c+d \tan(e+fx))^3}{3bf}$$

input

```
Int[((c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]
```

output

```
(C*(c + d*Tan[e + f*x])^3)/(3*b*f) + (((b*c*C + b*B*d - a*C*d)*(c + d*Tan[e + f*x])^2)/(2*b*f) + (((-(b^3*(a*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) - b*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2))))*x)/(a^2 + b^2)) - (b^3*(a*A*d*(3*c^2 - d^2) - A*b*(c^3 - 3*c*d^2) + b*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3) - a*(C*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2))))*Log[Cos[e + f*x]]/((a^2 + b^2)*f) + ((A*b^2 - a*(b*B - a*C))*(b*c - a*d)^3*Log[a + b*Tan[e + f*x]]/(b*(a^2 + b^2)*f))/b + (d*(b^2*d*(B*c + (A - C)*d) + (b*c - a*d)*(b*c*C + b*B*d - a*C*d))*Tan[e + f*x]/(b*f))/b/b
```


Definitions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`
- rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 501, normalized size of antiderivative = 1.38

method	result
norman	$\frac{(Aa^3c^3 - 3Aac^2d^2 + 3Ab^2c^2d - Ab^2d^3 - 3Bac^2d + Ba^2d^3 + Bb^2c^3 - 3Bbc^2d^2 - Ca^3c^3 + 3Cac^2d^2 - 3Cb^2c^2d + Cb^2d^3)x}{a^2 + b^2} + \frac{(Ab^2d^2 + 3Ab^2c^2d - 3Ab^2d^3 - 3Bac^2d + Ba^2d^3 + Bb^2c^3 - 3Bbc^2d^2 - Ca^3c^3 + 3Cac^2d^2 - 3Cb^2c^2d + Cb^2d^3)}{b^3}$
derivativedivides	$d \left(\frac{Cb^2d^2 \tan^3(fx+e)}{3} + \frac{Bb^2d^2 \tan^2(fx+e)}{2} - \frac{Cab^2d^2 \tan(fx+e)}{2} + \frac{3Cb^2cd \tan(fx+e)}{2} + \tan(fx+e)Ab^2d^2 - \tan(fx+e)Bab^2d^2 + 3 \right)$
default	$d \left(\frac{Cb^2d^2 \tan^3(fx+e)}{3} + \frac{Bb^2d^2 \tan^2(fx+e)}{2} - \frac{Cab^2d^2 \tan(fx+e)}{2} + \frac{3Cb^2cd \tan(fx+e)}{2} + \tan(fx+e)Ab^2d^2 - \tan(fx+e)Bab^2d^2 + 3 \right)$
parallelrisch	Expression too large to display
risch	Expression too large to display

output

```

1/6*(2*(C*a^2*b^3 + C*b^5)*d^3*tan(f*x + e)^3 + 6*(((A - C)*a*b^4 + B*b^5)
*c^3 - 3*(B*a*b^4 - (A - C)*b^5)*c^2*d - 3*((A - C)*a*b^4 + B*b^5)*c*d^2 +
(B*a*b^4 - (A - C)*b^5)*d^3)*f*x + 3*(3*(C*a^2*b^3 + C*b^5)*c*d^2 - (C*a^
3*b^2 - B*a^2*b^3 + C*a*b^4 - B*b^5)*d^3)*tan(f*x + e)^2 + 3*((C*a^2*b^3 -
B*a*b^4 + A*b^5)*c^3 - 3*(C*a^3*b^2 - B*a^2*b^3 + A*a*b^4)*c^2*d + 3*(C*a
^4*b - B*a^3*b^2 + A*a^2*b^3)*c*d^2 - (C*a^5 - B*a^4*b + A*a^3*b^2)*d^3)*l
og((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) -
3*((C*a^2*b^3 + C*b^5)*c^3 - 3*(C*a^3*b^2 - B*a^2*b^3 + C*a*b^4 - B*b^5)*
c^2*d + 3*(C*a^4*b - B*a^3*b^2 + A*a^2*b^3 - B*a*b^4 + (A - C)*b^5)*c*d^2
- (C*a^5 - B*a^4*b + A*a^3*b^2 + (A - C)*a*b^4 + B*b^5)*d^3)*log(1/(tan(f*
x + e)^2 + 1)) + 6*(3*(C*a^2*b^3 + C*b^5)*c^2*d - 3*(C*a^3*b^2 - B*a^2*b^3
+ C*a*b^4 - B*b^5)*c*d^2 + (C*a^4*b - B*a^3*b^2 + A*a^2*b^3 - B*a*b^4 + (
A - C)*b^5)*d^3)*tan(f*x + e))/((a^2*b^4 + b^6)*f)

```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 22.19 (sec) , antiderivative size = 7096, normalized size of antiderivative = 19.55

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input

```

integrate((c+d*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*
x+e)),x)

```

output

```
Piecewise((zoo*x*(c + d*tan(e))**3*(A + B*tan(e) + C*tan(e)**2)/tan(e), Eq
(a, 0) & Eq(b, 0) & Eq(f, 0)), ((A*c**3*x + 3*A*c**2*d*log(tan(e + f*x)**2
+ 1)/(2*f) - 3*A*c*d**2*x + 3*A*c*d**2*tan(e + f*x)/f - A*d**3*log(tan(e
+ f*x)**2 + 1)/(2*f) + A*d**3*tan(e + f*x)**2/(2*f) + B*c**3*log(tan(e + f
*x)**2 + 1)/(2*f) - 3*B*c**2*d*x + 3*B*c**2*d*tan(e + f*x)/f - 3*B*c*d**2*
log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*c*d**2*tan(e + f*x)**2/(2*f) + B*d**3
*x + B*d**3*tan(e + f*x)**3/(3*f) - B*d**3*tan(e + f*x)/f - C*c**3*x + C*c
**3*tan(e + f*x)/f - 3*C*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C*c**2*
d*tan(e + f*x)**2/(2*f) + 3*C*c*d**2*x + C*c*d**2*tan(e + f*x)**3/f - 3*C*
c*d**2*tan(e + f*x)/f + C*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*d**3*tan
(e + f*x)**4/(4*f) - C*d**3*tan(e + f*x)**2/(2*f))/a, Eq(b, 0)), (3*I*A*c*
**3*f*x*tan(e + f*x)/(6*b*f*tan(e + f*x) - 6*I*b*f) + 3*A*c**3*f*x/(6*b*f*t
an(e + f*x) - 6*I*b*f) + 3*I*A*c**3/(6*b*f*tan(e + f*x) - 6*I*b*f) + 9*A*c
**2*d*f*x*tan(e + f*x)/(6*b*f*tan(e + f*x) - 6*I*b*f) - 9*I*A*c**2*d*f*x/(
6*b*f*tan(e + f*x) - 6*I*b*f) - 9*A*c**2*d/(6*b*f*tan(e + f*x) - 6*I*b*f)
+ 9*I*A*c*d**2*f*x*tan(e + f*x)/(6*b*f*tan(e + f*x) - 6*I*b*f) + 9*A*c*d**
2*f*x/(6*b*f*tan(e + f*x) - 6*I*b*f) + 9*A*c*d**2*log(tan(e + f*x)**2 + 1)
*tan(e + f*x)/(6*b*f*tan(e + f*x) - 6*I*b*f) - 9*I*A*c*d**2*log(tan(e + f*
x)**2 + 1)/(6*b*f*tan(e + f*x) - 6*I*b*f) - 9*I*A*c*d**2/(6*b*f*tan(e + f*
x) - 6*I*b*f) - 9*A*d**3*f*x*tan(e + f*x)/(6*b*f*tan(e + f*x) - 6*I*b*f...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.20

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{6(((A-C)a+Bb)c^3-3(Ba-(A-C)b)c^2d-3((A-C)a+Bb)cd^2+(Ba-(A-C)b)d^3)(fx+e)}{a^2+b^2} + \frac{6((Ca^2b^3-Bab^4+Ab^5)c^3-3(Ca^3b^2-Ba^2b^3+...$$

input

```
integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e)),x, algorithm="maxima")
```

output

```

1/6*(6*((A - C)*a + B*b)*c^3 - 3*(B*a - (A - C)*b)*c^2*d - 3*((A - C)*a +
B*b)*c*d^2 + (B*a - (A - C)*b)*d^3)*(f*x + e)/(a^2 + b^2) + 6*((C*a^2*b^3
- B*a*b^4 + A*b^5)*c^3 - 3*(C*a^3*b^2 - B*a^2*b^3 + A*a*b^4)*c^2*d + 3*(C
*a^4*b - B*a^3*b^2 + A*a^2*b^3)*c*d^2 - (C*a^5 - B*a^4*b + A*a^3*b^2)*d^3)
*log(b*tan(f*x + e) + a)/(a^2*b^4 + b^6) + 3*((B*a - (A - C)*b)*c^3 + 3*((
A - C)*a + B*b)*c^2*d - 3*(B*a - (A - C)*b)*c*d^2 - ((A - C)*a + B*b)*d^3)
*log(tan(f*x + e)^2 + 1)/(a^2 + b^2) + (2*C*b^2*d^3*tan(f*x + e)^3 + 3*(3*
C*b^2*c*d^2 - (C*a*b - B*b^2)*d^3)*tan(f*x + e)^2 + 6*(3*C*b^2*c^2*d - 3*(
C*a*b - B*b^2)*c*d^2 + (C*a^2 - B*a*b + (A - C)*b^2)*d^3)*tan(f*x + e))/b^
3)/f

```

Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 600, normalized size of antiderivative = 1.65

$$\begin{aligned}
& \int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\
&= \frac{(Aac^3 - Cac^3 + Bbc^3 - 3Bac^2d + 3Abc^2d - 3Cbc^2d - 3Aacd^2 + 3Cacd^2 - 3Bbcd^2 + Bad^3 - Abd^3 + Cad^3)}{a^2f + b^2f} \\
&+ \frac{(Bac^3 - Abc^3 + Cbc^3 + 3Aac^2d - 3Cac^2d + 3Bbc^2d - 3Bacd^2 + 3Abcd^2 - 3Cbcd^2 - Aad^3 + Cad^3)}{2(a^2f + b^2f)} \\
&+ \frac{(Ca^2b^3c^3 - Bab^4c^3 + Ab^5c^3 - 3Ca^3b^2c^2d + 3Ba^2b^3c^2d - 3Aab^4c^2d + 3Ca^4bcd^2 - 3Ba^3b^2cd^2 + 3Aab^3cd^2)}{a^2b^4f + b^6f} \\
&+ \frac{2Cb^2d^3f^2 \tan(fx + e)^3 + 9Cb^2cd^2f^2 \tan(fx + e)^2 - 3Cabd^3f^2 \tan(fx + e)^2 + 3Bb^2d^3f^2 \tan(fx + e)}{a^2b^4f + b^6f}
\end{aligned}$$

input

```

integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e)),x, algorithm="giac")

```

output

```
(A*a*c^3 - C*a*c^3 + B*b*c^3 - 3*B*a*c^2*d + 3*A*b*c^2*d - 3*C*b*c^2*d - 3
*A*a*c*d^2 + 3*C*a*c*d^2 - 3*B*b*c*d^2 + B*a*d^3 - A*b*d^3 + C*b*d^3)*(f*x
+ e)/(a^2*f + b^2*f) + 1/2*(B*a*c^3 - A*b*c^3 + C*b*c^3 + 3*A*a*c^2*d - 3
*C*a*c^2*d + 3*B*b*c^2*d - 3*B*a*c*d^2 + 3*A*b*c*d^2 - 3*C*b*c*d^2 - A*a*d
^3 + C*a*d^3 - B*b*d^3)*log(tan(f*x + e)^2 + 1)/(a^2*f + b^2*f) + (C*a^2*b
^3*c^3 - B*a*b^4*c^3 + A*b^5*c^3 - 3*C*a^3*b^2*c^2*d + 3*B*a^2*b^3*c^2*d -
3*A*a*b^4*c^2*d + 3*C*a^4*b*c*d^2 - 3*B*a^3*b^2*c*d^2 + 3*A*a^2*b^3*c*d^2
- C*a^5*d^3 + B*a^4*b*d^3 - A*a^3*b^2*d^3)*log(abs(b*tan(f*x + e) + a))/(
a^2*b^4*f + b^6*f) + 1/6*(2*C*b^2*d^3*f^2*tan(f*x + e)^3 + 9*C*b^2*c*d^2*f
^2*tan(f*x + e)^2 - 3*C*a*b*d^3*f^2*tan(f*x + e)^2 + 3*B*b^2*d^3*f^2*tan(f
*x + e)^2 + 18*C*b^2*c^2*d*f^2*tan(f*x + e) - 18*C*a*b*c*d^2*f^2*tan(f*x +
e) + 18*B*b^2*c*d^2*f^2*tan(f*x + e) + 6*C*a^2*d^3*f^2*tan(f*x + e) - 6*B
*a*b*d^3*f^2*tan(f*x + e) + 6*A*b^2*d^3*f^2*tan(f*x + e) - 6*C*b^2*d^3*f^2
*tan(f*x + e))/(b^3*f^3)
```

Mupad [B] (verification not implemented)

Time = 9.35 (sec) , antiderivative size = 508, normalized size of antiderivative = 1.40

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{\tan(e + fx)^2 \left(\frac{B d^3 + 3 C c d^2}{2b} - \frac{C a d^3}{2b^2} \right)}{\tan(e + fx) \left(\frac{a \left(\frac{B d^3 + 3 C c d^2}{b} - \frac{C a d^3}{b^2} \right)}{b} - \frac{3 C c^2 d + 3 B c d^2 + A d^3}{b} + \frac{C d^3}{b} \right)}$$

$$- \frac{\ln(a + b \tan(e + fx)) (b^4 (B a c^3 + 3 A a d c^2) - b^3 (C a^2 c^3 + 3 B a^2 c^2 d + 3 A a^2 c d^2) + b^2 (3 C a^3 c^2 d + 3 B a^2 c^2 d^2 + 3 A a^2 c d^3) - b (3 C a^3 c^2 d^2 + 3 B a^2 c^2 d^3 + 3 A a^2 c d^3) - 3 C a^3 c^2 d^3 + 3 B a^2 c^2 d^3 + 3 A a^2 c d^3)}{f (a^2 b^4 + b^6)}$$

$$- \frac{\ln(\tan(e + fx) + 1i) (A c^3 + A d^3 1i - B c^3 1i + B d^3 - C c^3 - C d^3 1i - 3 A c d^2 - A c^2 d 3i + B c d^2)}{2 f (b + a 1i)}$$

$$- \frac{\ln(\tan(e + fx) - 1i) (A d^3 - B c^3 - C d^3 - 3 A c^2 d + 3 B c d^2 + 3 C c^2 d + A c^3 1i + B d^3 1i - C c^3 1i)}{2 f (a + b 1i)}$$

$$+ \frac{C d^3 \tan(e + fx)^3}{3 b f}$$

input

```
int(((c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a +
b*tan(e + f*x)),x)
```

output

```
(tan(e + f*x)^2*((B*d^3 + 3*C*c*d^2)/(2*b) - (C*a*d^3)/(2*b^2)))/f - (tan(
e + f*x)*((a*((B*d^3 + 3*C*c*d^2)/b - (C*a*d^3)/b^2))/b - (A*d^3 + 3*B*c*d
^2 + 3*C*c^2*d)/b + (C*d^3)/b))/f - (log(a + b*tan(e + f*x))*(b^4*(B*a*c^3
+ 3*A*a*c^2*d) - b^3*(C*a^2*c^3 + 3*A*a^2*c*d^2 + 3*B*a^2*c^2*d) + b^2*(A
*a^3*d^3 + 3*B*a^3*c*d^2 + 3*C*a^3*c^2*d) - b*(B*a^4*d^3 + 3*C*a^4*c*d^2)
- A*b^5*c^3 + C*a^5*d^3))/(f*(b^6 + a^2*b^4)) - (log(tan(e + f*x) + 1i)*(A
*c^3 + A*d^3*1i - B*c^3*1i + B*d^3 - C*c^3 - C*d^3*1i - 3*A*c*d^2 - A*c^2*d
^3i + B*c*d^2*3i - 3*B*c^2*d + 3*C*c*d^2 + C*c^2*d*3i))/(2*f*(a*1i + b))
- (log(tan(e + f*x) - 1i)*(A*c^3*1i + A*d^3 - B*c^3 + B*d^3*1i - C*c^3*1i
- C*d^3 - A*c*d^2*3i - 3*A*c^2*d + 3*B*c*d^2 - B*c^2*d*3i + C*c*d^2*3i + 3
*C*c^2*d))/(2*f*(a + b*1i)) + (C*d^3*tan(e + f*x)^3)/(3*b*f)
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 654, normalized size of antiderivative = 1.80

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{-18a^2b^4cd^2fx + 18ab^4c^2d^2fx - 3 \log(\tan(fx + e)^2 + 1) a^2b^4d^3 + 9 \log(\tan(fx + e)^2 + 1) b^6c^2d - 9 \log(\tan(fx + e)^2 + 1) b^6c^2d - 9 \log(\tan(fx + e)^2 + 1) b^6c^2d}{}$$

input

```
int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x)
```


output

```
(9*log(tan(e + f*x)**2 + 1)*a**2*b**4*c**2*d - 3*log(tan(e + f*x)**2 + 1)*
a**2*b**4*d**3 - 9*log(tan(e + f*x)**2 + 1)*a*b**4*c**3*d + 3*log(tan(e +
f*x)**2 + 1)*a*b**4*c*d**3 + 9*log(tan(e + f*x)**2 + 1)*b**6*c**2*d - 3*lo
g(tan(e + f*x)**2 + 1)*b**6*d**3 + 3*log(tan(e + f*x)**2 + 1)*b**5*c**4 -
9*log(tan(e + f*x)**2 + 1)*b**5*c**2*d**2 - 6*log(tan(e + f*x)*b + a)*a**5
*c*d**3 + 18*log(tan(e + f*x)*b + a)*a**4*b*c**2*d**2 - 18*log(tan(e + f*x
)*b + a)*a**3*b**2*c**3*d + 6*log(tan(e + f*x)*b + a)*a**2*b**3*c**4 + 2*t
an(e + f*x)**3*a**2*b**3*c*d**3 + 2*tan(e + f*x)**3*b**5*c*d**3 - 3*tan(e
+ f*x)**2*a**3*b**2*c*d**3 + 3*tan(e + f*x)**2*a**2*b**4*d**3 + 9*tan(e +
f*x)**2*a**2*b**3*c**2*d**2 - 3*tan(e + f*x)**2*a*b**4*c*d**3 + 3*tan(e +
f*x)**2*b**6*d**3 + 9*tan(e + f*x)**2*b**5*c**2*d**2 + 6*tan(e + f*x)*a**4
*b*c*d**3 - 18*tan(e + f*x)*a**3*b**2*c**2*d**2 + 18*tan(e + f*x)*a**2*b**
4*c*d**2 + 18*tan(e + f*x)*a**2*b**3*c**3*d - 18*tan(e + f*x)*a*b**4*c**2*
d**2 + 18*tan(e + f*x)*b**6*c*d**2 + 18*tan(e + f*x)*b**5*c**3*d - 6*tan(e
+ f*x)*b**5*c*d**3 + 6*a**2*b**4*c**3*f*x - 18*a**2*b**4*c*d**2*f*x - 6*a
*b**4*c**4*f*x + 18*a*b**4*c**2*d**2*f*x + 6*b**6*c**3*f*x - 18*b**6*c*d**
2*f*x - 18*b**5*c**3*d*f*x + 6*b**5*c*d**3*f*x)/(6*b**4*f*(a**2 + b**2))
```

3.68 $\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$

Optimal result	749
Mathematica [C] (verified)	750
Rubi [A] (verified)	751
Maple [A] (verified)	756
Fricas [B] (verification not implemented)	757
Sympy [C] (verification not implemented)	758
Maxima [A] (verification not implemented)	759
Giac [A] (verification not implemented)	760
Mupad [B] (verification not implemented)	761
Reduce [B] (verification not implemented)	762

Optimal result

Integrand size = 45, antiderivative size = 574

$$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx =$$

$$\frac{(b^2(Ac^3 - c^3C - 3Bc^2d - 3Acd^2 + 3cCd^2 + Bd^3) + a^2(c^3C + 3Bc^2d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2))}{(a^2 + b^2)^2}$$

$$+ \frac{(2ab(Ac^3 - c^3C - 3Bc^2d - 3Acd^2 + 3cCd^2 + Bd^3) - a^2((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) + b^2((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))}{(a^2 + b^2)^2 f}$$

$$- \frac{(bc - ad)^2 (2a^3bBd - 3a^4Cd - b^4(Bc + 3Ad) - 2ab^3(Ac - cC - 2Bd) + a^2b^2(Bc - (A + 5C)d)) \log\left(\frac{a + b \tan(e + fx)}{a + b \tan(e + fx)}\right)}{b^4 (a^2 + b^2)^2 f}$$

$$- \frac{d^2(3a^3Cd - Ab^2(bc - ad) - b^3(2cC + Bd) - a^2b(3cC + 2Bd) + ab^2(Bc + 2Cd)) \tan(e + fx)}{b^3 (a^2 + b^2) f}$$

$$+ \frac{(2Ab^2 - 2abB + 3a^2C + b^2C) d(c + d \tan(e + fx))^2}{2b^2 (a^2 + b^2) f}$$

$$- \frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^3}{b (a^2 + b^2) f (a + b \tan(e + fx))}$$

output

```

-(b^2*(A*c^3-3*A*c*d^2-3*B*c^2*d+B*d^3-C*c^3+3*C*c*d^2)+a^2*(c^3*C+3*B*c^2
*d-3*C*c*d^2-B*d^3-A*(c^3-3*c*d^2))-2*a*b*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*
d^2)))*x/(a^2+b^2)^2+(2*a*b*(A*c^3-3*A*c*d^2-3*B*c^2*d+B*d^3-C*c^3+3*C*c*d
^2)-a^2*((A-C)*d*(3*c^2-d^2)+B*(c^3-3*c*d^2))+b^2*((A-C)*d*(3*c^2-d^2)+B*(
c^3-3*c*d^2)))*ln(cos(f*x+e))/(a^2+b^2)^2/f-(-a*d+b*c)^2*(2*a^3*b*B*d-3*a^
4*C*d-b^4*(3*A*d+B*c)-2*a*b^3*(A*c-2*B*d-C*c)+a^2*b^2*(B*c-(A+5*C)*d))*ln(
a+b*tan(f*x+e))/b^4/(a^2+b^2)^2/f-d^2*(3*a^3*C*d-A*b^2*(-a*d+b*c)-b^3*(B*d
+2*C*c)-a^2*b*(2*B*d+3*C*c)+a*b^2*(B*c+2*C*d))*tan(f*x+e)/b^3/(a^2+b^2)/f+
1/2*(2*A*b^2-2*B*a*b+3*C*a^2+C*b^2)*d*(c+d*tan(f*x+e))^2/b^2/(a^2+b^2)/f-(
A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^3/b/(a^2+b^2)/f/(a+b*tan(f*x+e))

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.97 (sec) , antiderivative size = 1024, normalized size of antiderivative = 1.78

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{C(c + d \tan(e + fx))^3}{2bf(a + b \tan(e + fx))} + \frac{(3bcC + 2bBd - 3aCd)(c + d \tan(e + fx))^2}{bf(a + b \tan(e + fx))} + \frac{2 \left(-\frac{b^2(2aAbc^3 - a^2Bc^3 + b^2Bc^3 - 2abc^3C - 3a^2Ac^2d + 3Ab^2c^2d - 6abBc^2d + 3a^2c^2Cd - 3b^2c^2Cd - 6aAbcd^2)}{bf(a + b \tan(e + fx))} \right)}{bf(a + b \tan(e + fx))}$$

input

```

Integrate[((c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(a + b*Tan[e + f*x])^2,x]

```

output

```
(C*(c + d*Tan[e + f*x])^3)/(2*b*f*(a + b*Tan[e + f*x])) + (((3*b*c*C + 2*b
*B*d - 3*a*C*d)*(c + d*Tan[e + f*x])^2)/(b*f*(a + b*Tan[e + f*x])) + (2*(-
1/2*(b^2*(2*a*A*b*c^3 - a^2*B*c^3 + b^2*B*c^3 - 2*a*b*c^3*C - 3*a^2*A*c^2*
d + 3*A*b^2*c^2*d - 6*a*b*B*c^2*d + 3*a^2*c^2*C*d - 3*b^2*c^2*C*d - 6*a*A*
b*c*d^2 + 3*a^2*B*c*d^2 - 3*b^2*B*c*d^2 + 6*a*b*c*C*d^2 + a^2*A*d^3 - A*b^
2*d^3 + 2*a*b*B*d^3 - a^2*C*d^3 + b^2*C*d^3 + I*(a^2*A*c^3 - A*b^2*c^3 + 2
*a*b*B*c^3 - a^2*c^3*C + b^2*c^3*C + 6*a*A*b*c^2*d - 3*a^2*B*c^2*d + 3*b^2
*B*c^2*d - 6*a*b*c^2*C*d - 3*a^2*A*c*d^2 + 3*A*b^2*c*d^2 - 6*a*b*B*c*d^2 +
3*a^2*c*C*d^2 - 3*b^2*c*C*d^2 - 2*a*A*b*d^3 + a^2*B*d^3 - b^2*B*d^3 + 2*a
*b*C*d^3))*Log[I - Tan[e + f*x]])/((a^2 + b^2)^2*f) + (b^2*(-2*a*A*b*c^3 +
a^2*B*c^3 - b^2*B*c^3 + 2*a*b*c^3*C + 3*a^2*A*c^2*d - 3*A*b^2*c^2*d + 6*a
*b*B*c^2*d - 3*a^2*c^2*C*d + 3*b^2*c^2*C*d + 6*a*A*b*c*d^2 - 3*a^2*B*c*d^2
+ 3*b^2*B*c*d^2 - 6*a*b*c*C*d^2 - a^2*A*d^3 + A*b^2*d^3 - 2*a*b*B*d^3 + a
^2*C*d^3 - b^2*C*d^3 + I*(a^2*A*c^3 - A*b^2*c^3 + 2*a*b*B*c^3 - a^2*c^3*C
+ b^2*c^3*C + 6*a*A*b*c^2*d - 3*a^2*B*c^2*d + 3*b^2*B*c^2*d - 6*a*b*c^2*C*
d - 3*a^2*A*c*d^2 + 3*A*b^2*c*d^2 - 6*a*b*B*c*d^2 + 3*a^2*c*C*d^2 - 3*b^2*
c*C*d^2 - 2*a*A*b*d^3 + a^2*B*d^3 - b^2*B*d^3 + 2*a*b*C*d^3))*Log[I + Tan[
e + f*x]])/(2*(a^2 + b^2)^2*f) - ((b*c - a*d)^2*(2*a^3*b*B*d - 3*a^4*C*d -
b^4*(B*c + 3*A*d) - 2*a*b^3*(A*c - c*C - 2*B*d) + a^2*b^2*(B*c - (A + 5*C
)*d))*Log[a + b*Tan[e + f*x]])/(b^2*(a^2 + b^2)^2*f) + ((b*c - a*d)^2*(...
```

Rubi [A] (verified)

Time = 5.52 (sec) , antiderivative size = 589, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4128, 3042, 4130, 27, 3042, 4120, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^2} dx$$

↓ 4128

$$\int \frac{(c+d \tan(e+fx))^2 ((3Ca^2-2bBa+2Ab^2+b^2C)d \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-3ad)+Ab(ac+3bd))}{a+b \tan(e+fx)} d$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^2 ((3Ca^2-2bBa+2Ab^2+b^2C)d \tan(e+fx)^2-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-3ad)+Ab(ac+3bd))}{a+b \tan(e+fx)} d$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{}$$

↓ 4130

$$\int -\frac{2(c+d \tan(e+fx))(-((2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2)-c((bB-aC)(bc-3ad)+Ab(ac+3bd))b+a(3Ca^2-2bBa+2Ab^2+b^2C))}{a+b \tan(e+fx)} d$$

$$\frac{2b}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{}$$

↓ 27

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \int \frac{(c+d \tan(e+fx))(-((2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2)-c((bB-aC)(bc-3ad)+Ab(ac+3bd))b+a(3Ca^2-2bBa+2Ab^2+b^2C))}{a+b \tan(e+fx)} d$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{}$$

↓ 3042

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \int \frac{(c+d \tan(e+fx))(-((2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2)-c((bB-aC)(bc-3ad)+Ab(ac+3bd))b+a(3Ca^2-2bBa+2Ab^2+b^2C))}{a+b \tan(e+fx)} d$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}$$

$$\frac{bf(a^2+b^2)(a+b \tan(e+fx))}{}$$

↓ 4120

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \frac{d^2 \tan(e+fx)(3a^3Cd-a^2b(2Bd+3cC)-Ab^2(bc-ad)+ab^2(Bc+2Cd)-b^3(Bd+2cC))}{bf} - \int \frac{3Cd^3a^4-2bd^2}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 3042

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \frac{d^2 \tan(e+fx)(3a^3Cd-a^2b(2Bd+3cC)-Ab^2(bc-ad)+ab^2(Bc+2Cd)-b^3(Bd+2cC))}{bf} - \int \frac{3Cd^3a^4-2bd^2}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 4109

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \frac{d^2 \tan(e+fx)(3a^3Cd-a^2b(2Bd+3cC)-Ab^2(bc-ad)+ab^2(Bc+2Cd)-b^3(Bd+2cC))}{bf} - \frac{b^3(a^2(d(A-C))}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 3042

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \frac{d^2 \tan(e+fx)(3a^3Cd-a^2b(2Bd+3cC)-Ab^2(bc-ad)+ab^2(Bc+2Cd)-b^3(Bd+2cC))}{bf} - \frac{b^3(a^2(d(A-C))}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 3956

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)(c+d \tan(e+fx))^2}{2bf} - \frac{d^2 \tan(e+fx)(3a^3Cd-a^2b(2Bd+3cC)-Ab^2(bc-ad)+ab^2(Bc+2Cd)-b^3(Bd+2cC))}{bf} - \frac{(bc-ad)^2(-3}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 4100

$$\frac{d(3a^2C - 2abB + 2Ab^2 + b^2C)(c + d \tan(e + fx))^2}{2bf} - \frac{d^2 \tan(e + fx)(3a^3Cd - a^2b(2Bd + 3cC) - Ab^2(bc - ad) + ab^2(Bc + 2Cd) - b^3(Bd + 2cC))}{bf} - \frac{(bc - ad)^2(-3a^2c + b^2c^2)}{b^3}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 16

$$\frac{d(3a^2C - 2abB + 2Ab^2 + b^2C)(c + d \tan(e + fx))^2}{2bf} - \frac{d^2 \tan(e + fx)(3a^3Cd - a^2b(2Bd + 3cC) - Ab^2(bc - ad) + ab^2(Bc + 2Cd) - b^3(Bd + 2cC))}{bf} - \frac{b^3 \log(\cos(e + fx))}{b^3}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

input

```
Int[((c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^2,x]
```

output

```
-(((A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^3)/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x]))) + (((2*A*b^2 - 2*a*b*B + 3*a^2*C + b^2*C)*d*(c + d*Tan[e + f*x])^2)/(2*b*f) - (-(((b^3*(b^2*(A*c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3) + a^2*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) - 2*a*b*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)))*x)/(a^2 + b^2) - (b^3*(2*a*b*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) + a^2*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)) - b^2*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)))*Log[Cos[e + f*x]])/(a^2 + b^2)*f) - ((b*c - a*d)^2*(2*a^3*b*B*d - 3*a^4*C*d - b^4*(B*c + 3*A*d) - 2*a*b^3*(A*c - c*C - 2*B*d) + a^2*b^2*(B*c - (A + 5*C)*d))*Log[a + b*Tan[e + f*x]])/(b*(a^2 + b^2)*f))/b + (d^2*(3*a^3*C*d - A*b^2*(b*c - a*d) - b^3*(2*c*C + B*d) - a^2*b*(3*c*C + 2*B*d) + a*b^2*(B*c + 2*C*d))*Tan[e + f*x])/(b*f))/b)/(b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4100 $\text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x_)]^{(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[(A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)/((a_)+(b_)*\tan[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A*b - a*B - b*C, 0]$
- rule 4120 $\text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x_)]*(c_)+(d_)*\tan[(e_)+(f_)*(x_)]^{(n_)*((A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[b*C*\text{Tan}[e + f*x]*((c + d*\text{Tan}[e + f*x])^{(n + 1)/(d*f*(n + 2))}), x] - \text{Simp}[1/(d*(n + 2)) \text{ Int}[(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*\text{Tan}[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ !\text{LtQ}[n, -1]$

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 829, normalized size of antiderivative = 1.44

method	result
derivativedivides	$\frac{d^2 \left(\frac{\tan(fx+e)^2 C b d}{2} + B \tan(fx+e) b d - 2 d \tan(fx+e) C a + 3 \tan(fx+e) C b c \right)}{b^3} + \frac{(3 A a^2 c^2 d - A a^2 d^3 - 2 A a b c^3 + 6 A a b c d^2 - 3 A b^2 c^2 d + \dots)}{b^3}$
default	$\frac{d^2 \left(\frac{\tan(fx+e)^2 C b d}{2} + B \tan(fx+e) b d - 2 d \tan(fx+e) C a + 3 \tan(fx+e) C b c \right)}{b^3} + \frac{(3 A a^2 c^2 d - A a^2 d^3 - 2 A a b c^3 + 6 A a b c d^2 - 3 A b^2 c^2 d + \dots)}{b^3}$
norman	Expression too large to display
parallelrisc	Expression too large to display
risc	Expression too large to display

input `int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2, x,method=_RETURNVERBOSE)`

output
$$\frac{1}{f} \left(\frac{d^2}{b^3} \left(\frac{1}{2} \tan(fx+e)^2 C b d + B \tan(fx+e) b d - 2 d \tan(fx+e) C a + 3 \tan(fx+e) C b c \right) + \frac{1}{(a^2+b^2)^2} \left(\frac{1}{2} (3 A a^2 c^2 d - A a^2 d^3 - 2 A a b c^3 + 6 A a b c d^2 - 3 A b^2 c^2 d + A b^2 d^3 + B a^2 c^3 - 3 B a^2 c d^2 + 6 B a b c^2 d - 2 B a b d^3 - B b^2 c^3 + 3 B b^2 c d^2 - 3 C a^2 c^2 d + C a^2 d^3 + 2 C a b c^3 - 6 C a b c d^2 + 3 C b^2 c^2 d - C b^2 d^3) \right) \ln(1 + \tan(fx+e)^2) + (A a^2 c^3 - 3 A a^2 c d^2 + 6 A a b c^2 d - 2 A a b d^3 - A b^2 c^3 + 3 A b^2 c d^2 - 3 B a^2 c^2 d + B a^2 d^3 + 2 B a b c^3 - 6 B a b c d^2 + 3 B b^2 c^2 d - B b^2 d^3 - C a^2 c^3 + 3 C a^2 c d^2 - 6 C a b c^2 d + 2 C a b d^3 + C b^2 c^3 - 3 C b^2 c d^2) \arctan(\tan(fx+e)) - (-A a^3 b^2 d^3 + 3 A a^2 b^3 c d^2 - 3 A a b^4 c^2 d + A b^5 c^3 + B a^4 b d^3 - 3 B a^3 b^2 c d^2 + 3 B a^2 b^3 c^2 d - B a b^4 c^3 - C a^5 d^3 + 3 C a^4 b c d^2 - 3 C a^3 b^2 c^2 d + C a^2 b^3 c^3) / b^4 / (a^2 + b^2) / (a + b \tan(fx+e)) + 1/b^4 * (A a^4 b^2 d^3 - 3 A a^2 b^4 c^2 d + 3 A a^2 b^4 d^3 + 2 A a b^5 c^3 - 6 A a b^5 c d^2 + 3 A b^6 c^2 d - 2 B a^5 b d^3 + 3 B a^4 b^2 c d^2 - 4 B a^3 b^3 d^3 - B a^2 b^4 c^3 + 9 B a^2 b^4 c d^2 - 6 B a b^5 c^2 d + B b^6 c^3 + 3 C a^6 d^3 - 6 C a^5 b c d^2 + 3 C a^4 b^2 c^2 d + 5 C a^4 b^2 d^3 - 12 C a^3 b^3 c d^2 + 9 C a^2 b^4 c^2 d - 2 C a b^5 c^3) / (a^2 + b^2)^2 \ln(a + b \tan(fx+e)) \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1512 vs. $2(571) = 1142$.

Time = 0.95 (sec) , antiderivative size = 1512, normalized size of antiderivative = 2.63

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

```

1/2*((C*a^4*b^3 + 2*C*a^2*b^5 + C*b^7)*d^3*tan(f*x + e)^3 - 2*(C*a^2*b^5 -
B*a*b^6 + A*b^7)*c^3 + 6*(C*a^3*b^4 - B*a^2*b^5 + A*a*b^6)*c^2*d - 6*(C*a
^4*b^3 - B*a^3*b^4 + A*a^2*b^5)*c*d^2 + (3*C*a^5*b^2 - 2*B*a^4*b^3 + 2*(A
+ C)*a^3*b^4 + C*a*b^6)*d^3 + 2*((A - C)*a^3*b^4 + 2*B*a^2*b^5 - (A - C)*
a*b^6)*c^3 - 3*(B*a^3*b^4 - 2*(A - C)*a^2*b^5 - B*a*b^6)*c^2*d - 3*((A - C
)*a^3*b^4 + 2*B*a^2*b^5 - (A - C)*a*b^6)*c*d^2 + (B*a^3*b^4 - 2*(A - C)*a^
2*b^5 - B*a*b^6)*d^3)*f*x + (6*(C*a^4*b^3 + 2*C*a^2*b^5 + C*b^7)*c*d^2 - (
3*C*a^5*b^2 - 2*B*a^4*b^3 + 6*C*a^3*b^4 - 4*B*a^2*b^5 + 3*C*a*b^6 - 2*B*b^
7)*d^3)*tan(f*x + e)^2 - ((B*a^3*b^4 - 2*(A - C)*a^2*b^5 - B*a*b^6)*c^3 -
3*(C*a^5*b^2 - (A - 3*C)*a^3*b^4 - 2*B*a^2*b^5 + A*a*b^6)*c^2*d + 3*(2*C*a
^6*b - B*a^5*b^2 + 4*C*a^4*b^3 - 3*B*a^3*b^4 + 2*A*a^2*b^5)*c*d^2 - (3*C*a
^7 - 2*B*a^6*b + (A + 5*C)*a^5*b^2 - 4*B*a^4*b^3 + 3*A*a^3*b^4)*d^3 + ((B*
a^2*b^5 - 2*(A - C)*a*b^6 - B*b^7)*c^3 - 3*(C*a^4*b^3 - (A - 3*C)*a^2*b^5
- 2*B*a*b^6 + A*b^7)*c^2*d + 3*(2*C*a^5*b^2 - B*a^4*b^3 + 4*C*a^3*b^4 - 3*
B*a^2*b^5 + 2*A*a*b^6)*c*d^2 - (3*C*a^6*b - 2*B*a^5*b^2 + (A + 5*C)*a^4*b^
3 - 4*B*a^3*b^4 + 3*A*a^2*b^5)*d^3)*tan(f*x + e))*log((b^2*tan(f*x + e)^2
+ 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) - (3*(C*a^5*b^2 + 2*C*a^
3*b^4 + C*a*b^6)*c^2*d - 3*(2*C*a^6*b - B*a^5*b^2 + 4*C*a^4*b^3 - 2*B*a^3*
b^4 + 2*C*a^2*b^5 - B*a*b^6)*c*d^2 + (3*C*a^7 - 2*B*a^6*b + (A + 5*C)*a^5*
b^2 - 4*B*a^4*b^3 + (2*A + C)*a^3*b^4 - 2*B*a^2*b^5 + (A - C)*a*b^6)*d^...

```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 28.58 (sec) , antiderivative size = 24300, normalized size of antiderivative = 42.33

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((c+d*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*
x+e))**2,x)

```

output

```
Piecewise((zoo*x*(c + d*tan(e))**3*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2,
Eq(a, 0) & Eq(b, 0) & Eq(f, 0)), ((A*c**3*x + 3*A*c**2*d*log(tan(e + f*x)
**2 + 1)/(2*f) - 3*A*c*d**2*x + 3*A*c*d**2*tan(e + f*x)/f - A*d**3*log(tan
(e + f*x)**2 + 1)/(2*f) + A*d**3*tan(e + f*x)**2/(2*f) + B*c**3*log(tan(e
+ f*x)**2 + 1)/(2*f) - 3*B*c**2*d*x + 3*B*c**2*d*tan(e + f*x)/f - 3*B*c*d*
**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*c*d**2*tan(e + f*x)**2/(2*f) + B*d
**3*x + B*d**3*tan(e + f*x)**3/(3*f) - B*d**3*tan(e + f*x)/f - C*c**3*x +
C*c**3*tan(e + f*x)/f - 3*C*c**2*d*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C*c*
**2*d*tan(e + f*x)**2/(2*f) + 3*C*c*d**2*x + C*c*d**2*tan(e + f*x)**3/f - 3
*C*c*d**2*tan(e + f*x)/f + C*d**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*d**3*
tan(e + f*x)**4/(4*f) - C*d**3*tan(e + f*x)**2/(2*f))/a**2, Eq(b, 0)), (-A
*c**3*f*x*tan(e + f*x)**2/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f
*x) - 4*b**2*f) + 2*I*A*c**3*f*x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 -
8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + A*c**3*f*x/(4*b**2*f*tan(e + f*x)**2
- 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) - A*c**3*tan(e + f*x)/(4*b**2*f*tan
(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + 2*I*A*c**3/(4*b**2*f*
tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**2*f) + 3*I*A*c**2*d*f*x*t
an(e + f*x)**2/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*tan(e + f*x) - 4*b**
2*f) + 6*A*c**2*d*f*x*tan(e + f*x)/(4*b**2*f*tan(e + f*x)**2 - 8*I*b**2*f*
tan(e + f*x) - 4*b**2*f) - 3*I*A*c**2*d*f*x/(4*b**2*f*tan(e + f*x)**2 - ...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 685, normalized size of antiderivative = 1.19

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{2(((A-C)a^2 + 2Bab - (A-C)b^2)c^3 - 3(Ba^2 - 2(A-C)ab - Bb^2)c^2 d - 3((A-C)a^2 + 2Bab - (A-C)b^2)cd^2 + (Ba^2 - 2(A-C)ab - Bb^2)d^3)(fx + e)}{a^4 + 2a^2b^2 + b^4}$$

input

```
integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e))^2,x, algorithm="maxima")
```

output

```

1/2*(2*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c^3 - 3*(B*a^2 - 2*(A - C)*a
*b - B*b^2)*c^2*d - 3*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c*d^2 + (B*a^2
- 2*(A - C)*a*b - B*b^2)*d^3)*(f*x + e)/(a^4 + 2*a^2*b^2 + b^4) - 2*((B*a
^2*b^4 - 2*(A - C)*a*b^5 - B*b^6)*c^3 - 3*(C*a^4*b^2 - (A - 3*C)*a^2*b^4 -
2*B*a*b^5 + A*b^6)*c^2*d + 3*(2*C*a^5*b - B*a^4*b^2 + 4*C*a^3*b^3 - 3*B*a
^2*b^4 + 2*A*a*b^5)*c*d^2 - (3*C*a^6 - 2*B*a^5*b + (A + 5*C)*a^4*b^2 - 4*B
*a^3*b^3 + 3*A*a^2*b^4)*d^3)*log(b*tan(f*x + e) + a)/(a^4*b^4 + 2*a^2*b^6
+ b^8) + ((B*a^2 - 2*(A - C)*a*b - B*b^2)*c^3 + 3*((A - C)*a^2 + 2*B*a*b -
(A - C)*b^2)*c^2*d - 3*(B*a^2 - 2*(A - C)*a*b - B*b^2)*c*d^2 - ((A - C)*a
^2 + 2*B*a*b - (A - C)*b^2)*d^3)*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2
+ b^4) - 2*((C*a^2*b^3 - B*a*b^4 + A*b^5)*c^3 - 3*(C*a^3*b^2 - B*a^2*b^3 +
A*a*b^4)*c^2*d + 3*(C*a^4*b - B*a^3*b^2 + A*a^2*b^3)*c*d^2 - (C*a^5 - B*a
^4*b + A*a^3*b^2)*d^3)/(a^3*b^4 + a*b^6 + (a^2*b^5 + b^7)*tan(f*x + e)) +
(C*b*d^3*tan(f*x + e)^2 + 2*(3*C*b*c*d^2 - (2*C*a - B*b)*d^3)*tan(f*x + e)
)/b^3)/f

```

Giac [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 1042, normalized size of antiderivative = 1.82

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e))^2,x, algorithm="giac")

```

output

```
(A*a^2*c^3 - C*a^2*c^3 + 2*B*a*b*c^3 - A*b^2*c^3 + C*b^2*c^3 - 3*B*a^2*c^2
*d + 6*A*a*b*c^2*d - 6*C*a*b*c^2*d + 3*B*b^2*c^2*d - 3*A*a^2*c*d^2 + 3*C*a
^2*c*d^2 - 6*B*a*b*c*d^2 + 3*A*b^2*c*d^2 - 3*C*b^2*c*d^2 + B*a^2*d^3 - 2*A
*a*b*d^3 + 2*C*a*b*d^3 - B*b^2*d^3)*(f*x + e)/(a^4*f + 2*a^2*b^2*f + b^4*f
) + 1/2*(B*a^2*c^3 - 2*A*a*b*c^3 + 2*C*a*b*c^3 - B*b^2*c^3 + 3*A*a^2*c^2*d
- 3*C*a^2*c^2*d + 6*B*a*b*c^2*d - 3*A*b^2*c^2*d + 3*C*b^2*c^2*d - 3*B*a^2
*c*d^2 + 6*A*a*b*c*d^2 - 6*C*a*b*c*d^2 + 3*B*b^2*c*d^2 - A*a^2*d^3 + C*a^2
*d^3 - 2*B*a*b*d^3 + A*b^2*d^3 - C*b^2*d^3)*log(tan(f*x + e)^2 + 1)/(a^4*f
+ 2*a^2*b^2*f + b^4*f) - (B*a^2*b^4*c^3 - 2*A*a*b^5*c^3 + 2*C*a*b^5*c^3 -
B*b^6*c^3 - 3*C*a^4*b^2*c^2*d + 3*A*a^2*b^4*c^2*d - 9*C*a^2*b^4*c^2*d + 6
*B*a*b^5*c^2*d - 3*A*b^6*c^2*d + 6*C*a^5*b*c*d^2 - 3*B*a^4*b^2*c*d^2 + 12*
C*a^3*b^3*c*d^2 - 9*B*a^2*b^4*c*d^2 + 6*A*a*b^5*c*d^2 - 3*C*a^6*d^3 + 2*B*
a^5*b*d^3 - A*a^4*b^2*d^3 - 5*C*a^4*b^2*d^3 + 4*B*a^3*b^3*d^3 - 3*A*a^2*b^
4*d^3)*log(abs(b*tan(f*x + e) + a))/(a^4*b^4*f + 2*a^2*b^6*f + b^8*f) + 1/
2*(C*b^2*d^3*f*tan(f*x + e)^2 + 6*C*b^2*c*d^2*f*tan(f*x + e) - 4*C*a*b*d^3
*f*tan(f*x + e) + 2*B*b^2*d^3*f*tan(f*x + e))/(b^4*f^2) - (C*a^4*b^3*c^3 -
B*a^3*b^4*c^3 + A*a^2*b^5*c^3 + C*a^2*b^5*c^3 - B*a*b^6*c^3 + A*b^7*c^3 -
3*C*a^5*b^2*c^2*d + 3*B*a^4*b^3*c^2*d - 3*A*a^3*b^4*c^2*d - 3*C*a^3*b^4*c
^2*d + 3*B*a^2*b^5*c^2*d - 3*A*a*b^6*c^2*d + 3*C*a^6*b*c*d^2 - 3*B*a^5*b^2
*c*d^2 + 3*A*a^4*b^3*c*d^2 + 3*C*a^4*b^3*c*d^2 - 3*B*a^3*b^4*c*d^2 + 3*...
```

Mupad [B] (verification not implemented)

Time = 13.00 (sec) , antiderivative size = 701, normalized size of antiderivative = 1.22

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{\tan(e + fx) \left(\frac{Bd^3 + 3Ccd^2}{b^2} - \frac{2Cad^3}{b^3} \right)}{f}$$

$$- \frac{\ln(\tan(e + fx) + 1i) (Bc^3 - Ad^3 + Cd^3 + 3Ac^2d - 3Bcd^2 - 3C^2d + Ac^3 1i + Bd^3 1i - Cc^3 1i)}{2f(-a^2 + ab2i + b^2)}$$

$$+ \frac{\ln(a + b \tan(e + fx)) (b^4 (3Aa^2d^3 - Ba^2c^3 - 3Aa^2c^2d + 9Ba^2cd^2 + 9Ca^2c^2d) - b^5 (2Cac^3 - 3C^2d))}{2f(-a^2 + ab2i + b^2)}$$

$$- \frac{\ln(\tan(e + fx) - 1i) (Ac^3 - Ad^3 1i + Bc^3 1i + Bd^3 - Cc^3 + Cd^3 1i - 3Acd^2 + Ac^2d3i - Bcd^2 3i)}{2f(-a^2 1i + 2ab + b^2 1i)}$$

$$- \frac{-Ca^5d^3 + 3Ca^4bcd^2 + Ba^4bd^3 - 3Ca^3b^2c^2d - 3Ba^3b^2cd^2 - Aa^3b^2d^3 + Ca^2b^3c^3 + 3Ba^2b^3c^3}{bf(\tan(e + fx) b^4 + ab^3)(a^2 + b^2)}$$

$$+ \frac{Cd^3 \tan(e + fx)^2}{2b^2 f}$$

input `int(((c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^2,x)`

output
$$\begin{aligned} & (\tan(e + f*x)*((B*d^3 + 3*C*c*d^2)/b^2 - (2*C*a*d^3)/b^3))/f - (\log(\tan(e + f*x) + 1i)*(A*c^3*1i - A*d^3 + B*c^3 + B*d^3*1i - C*c^3*1i + C*d^3 - A*c*d^2*3i + 3*A*c^2*d - 3*B*c*d^2 - B*c^2*d*3i + C*c*d^2*3i - 3*C*c^2*d))/(2*f*(a*b^2i - a^2 + b^2)) + (\log(a + b*tan(e + f*x))*(b^4*(3*A*a^2*d^3 - B*a^2*c^3 - 3*A*a^2*c^2*d + 9*B*a^2*c*d^2 + 9*C*a^2*c^2*d) - b^5*(2*C*a*c^3 - 2*A*a*c^3 + 6*A*a*c*d^2 + 6*B*a*c^2*d) - b^3*(4*B*a^3*d^3 + 12*C*a^3*c*d^2) + b^6*(B*c^3 + 3*A*c^2*d) - b*(2*B*a^5*d^3 + 6*C*a^5*c*d^2) + b^2*(A*a^4*d^3 + 5*C*a^4*d^3 + 3*B*a^4*c*d^2 + 3*C*a^4*c^2*d) + 3*C*a^6*d^3))/(f*(b^8 + 2*a^2*b^6 + a^4*b^4)) - (\log(\tan(e + f*x) - 1i)*(A*c^3 - A*d^3*1i + B*c^3*1i + B*d^3 - C*c^3 + C*d^3*1i - 3*A*c*d^2 + A*c^2*d*3i - B*c*d^2*3i - 3*B*c^2*d + 3*C*c*d^2 - C*c^2*d*3i))/(2*f*(2*a*b - a^2*1i + b^2*1i)) - (A*b^5*c^3 - C*a^5*d^3 - B*a*b^4*c^3 + B*a^4*b*d^3 - A*a^3*b^2*d^3 + C*a^2*b^3*c^3 + 3*A*a^2*b^3*c*d^2 + 3*B*a^2*b^3*c^2*d - 3*B*a^3*b^2*c*d^2 - 3*C*a^3*b^2*c^2*d - 3*A*a*b^4*c^2*d + 3*C*a^4*b*c*d^2)/(b*f*(a*b^3 + b^4*tan(e + f*x))*(a^2 + b^2)) + (C*d^3*tan(e + f*x)^2)/(2*b^2*f) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 2418, normalized size of antiderivative = 4.21

$$\int \frac{(c + d \tan(e + f x))^3 (A + B \tan(e + f x) + C \tan^2(e + f x))}{(a + b \tan(e + f x))^2} dx = \text{Too large to display}$$

input `int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x)`

output

```
(3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**5*c**2*d - log(tan(e + f*
x)**2 + 1)*tan(e + f*x)*a**3*b**5*d**3 - log(tan(e + f*x)**2 + 1)*tan(e +
f*x)*a**2*b**6*c**3 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**6*c
d**2 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**5*c**3*d + log(tan(
e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**5*c*d**3 + 3*log(tan(e + f*x)**2 + 1
)*tan(e + f*x)*a*b**7*c**2*d - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**
7*d**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**6*c**4 - 6*log(tan(e
 + f*x)**2 + 1)*tan(e + f*x)*a*b**6*c**2*d**2 - log(tan(e + f*x)**2 + 1)*t
an(e + f*x)*b**8*c**3 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**8*c*d**
2 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**7*c**3*d - log(tan(e + f*x)
**2 + 1)*tan(e + f*x)*b**7*c*d**3 + 3*log(tan(e + f*x)**2 + 1)*a**4*b**4*c
**2*d - log(tan(e + f*x)**2 + 1)*a**4*b**4*d**3 - log(tan(e + f*x)**2 + 1)
*a**3*b**5*c**3 + 3*log(tan(e + f*x)**2 + 1)*a**3*b**5*c*d**2 - 3*log(tan(
e + f*x)**2 + 1)*a**3*b**4*c**3*d + log(tan(e + f*x)**2 + 1)*a**3*b**4*c*d
**3 + 3*log(tan(e + f*x)**2 + 1)*a**2*b**6*c**2*d - log(tan(e + f*x)**2 +
1)*a**2*b**6*d**3 + 2*log(tan(e + f*x)**2 + 1)*a**2*b**5*c**4 - 6*log(tan(
e + f*x)**2 + 1)*a**2*b**5*c**2*d**2 - log(tan(e + f*x)**2 + 1)*a*b**7*c**
3 + 3*log(tan(e + f*x)**2 + 1)*a*b**7*c*d**2 + 3*log(tan(e + f*x)**2 + 1)*
a*b**6*c**3*d - log(tan(e + f*x)**2 + 1)*a*b**6*c*d**3 + 6*log(tan(e + f*x
)*b + a)*tan(e + f*x)*a**6*b*c*d**3 - 2*log(tan(e + f*x)*b + a)*tan(e +...
```


3.69
$$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

Optimal result	764
Mathematica [C] (verified)	765
Rubi [A] (verified)	766
Maple [A] (verified)	771
Fricas [B] (verification not implemented)	772
Sympy [F(-2)]	773
Maxima [A] (verification not implemented)	774
Giac [B] (verification not implemented)	775
Mupad [B] (verification not implemented)	776
Reduce [B] (verification not implemented)	777

Optimal result

Integrand size = 45, antiderivative size = 798

$$\int \frac{(c+d \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx =$$

$$\frac{(3ab^2(Ac^3 - c^3C - 3Bc^2d - 3Acd^2 + 3cCd^2 + Bd^3) + a^3(c^3C + 3Bc^2d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2 - 3cd^2 + b^2)))}{(a^2 + b^2)}$$

$$\frac{(b^3(Ac^3 - c^3C - 3Bc^2d - 3Acd^2 + 3cCd^2 + Bd^3) + 3a^2b(c^3C + 3Bc^2d - 3cCd^2 - Bd^3 - A(c^3 - 3cd^2 - 3cd^2 + b^2)))}{(a^2 + b^2)}$$

$$\frac{(bc - ad)(a^5bBd^2 - 3a^6Cd^2 + a^4b^2d(Bc - 9Cd) + a^3b^3B(c^2 + 3d^2) - b^6(c(cC + 3Bd) - A(c^2 - 3d^2)))}{b^4(a^2 + b^2)}$$

$$\frac{d^2(a^3bBd - 3a^4Cd - ab^3(2Ac - 2cC - 3Bd) + a^2b^2(Bc - 6Cd) - b^4(Bc + (2A + C)d)) \tan(e+fx)}{b^3(a^2 + b^2)^2 f}$$

$$+ \frac{(a^3bBd - 3a^4Cd - b^4(2Bc + 3Ad) - ab^3(4Ac - 4cC - 5Bd) + a^2b^2(2Bc + (A - 7C)d))(c + d \tan(e+fx))}{2b^2(a^2 + b^2)^2 f(a + b \tan(e+fx))}$$

$$- \frac{(Ab^2 - a(bB - aC))(c + d \tan(e+fx))^3}{2b(a^2 + b^2) f(a + b \tan(e+fx))^2}$$

output

```

(((A + I*B - C)*(c + I*d)^3*Log[I - Tan[e + f*x]])/((-I)*a + b)^3 + ((A -
I*B - C)*(c - I*d)^3*Log[I + Tan[e + f*x]])/(I*a + b)^3 - (2*(b*c - a*d)*(
a^5*b*B*d^2 - 3*a^6*C*d^2 + a^4*b^2*d*(B*c - 9*C*d) + a^3*b^3*B*(c^2 + 3*d
^2) + b^6*(-(c*(c*C + 3*B*d)) + A*(c^2 - 3*d^2)) + a*b^5*(8*c*(-A + C)*d -
3*B*(c^2 - 2*d^2)) + a^2*b^4*(3*c^2*C + 6*B*c*d - 10*C*d^2 + A*(-3*c^2 +
d^2))) * Log[a + b*Tan[e + f*x]])/(b^4*(a^2 + b^2)^3) - ((A*b^2 - a*b*B + 3*
a^2*C + 2*b^2*C)*(b*c - a*d)^3)/(b^4*(a^2 + b^2)*(a + b*Tan[e + f*x])^2) -
(2*(b*c - a*d)^2*(-2*a^3*b*B*d + 6*a^4*C*d + 2*a*b^3*(A*c - c*C - 2*B*d)
+ b^4*(B*c + 3*(A + C)*d) + a^2*b^2*(-(B*c) + (A + 11*C)*d))/(b^4*(a^2 +
b^2)^2*(a + b*Tan[e + f*x])) + (2*C*(c + d*Tan[e + f*x])^3)/(b*(a + b*Tan[
e + f*x])^2))/(2*f)

```

Rubi [A] (verified)

Time = 7.06 (sec) , antiderivative size = 830, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4128, 3042, 4128, 3042, 4120, 27, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{(c + d \tan(e + fx))^2 ((3Ca^2 - bBa + Ab^2 + 2b^2C)d \tan^2(e + fx) - 2b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(2bc - 3ad) + Ab(2ac + 3bd))}{(a + b \tan(e + fx))^2}$$

$$\frac{2b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^3}$$

$$\frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^3}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^2 ((3Ca^2-bBa+Ab^2+2b^2C)d \tan(e+fx)^2-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(2bc-3ad)+Ab(2ac+3bd))}{(a+b \tan(e+fx))^2} \\ \frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3} \\ \frac{2bf(a^2+b^2)(a+b \tan(e+fx))^2}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 4128

$$\int \frac{(c+d \tan(e+fx))(-2((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+2bd)((bB-aC)(2bc-3ad)+Ab(2ac+3bd))}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))(-2((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+2bd)((bB-aC)(2bc-3ad)+Ab(2ac+3bd))}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 4120

$$\int \frac{2(3Cd^3a^5-bd^2(3cC+Bd)a^4+6b^2Cd^3a^3-b^3(Ac^3-Cc^3-3Bdc^2-3Ad^2c+9Cd^2c+3Bd^3)a^2-b^4(2Bc^3+6Adc^2-6Cdc^2-6Bd^2c-2Ad^3-Cd^3)a-(a^2+b^2)^2d^2)}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 27

$$2 \int \frac{3Cd^3a^5-bd^2(3cC+Bd)a^4+6b^2Cd^3a^3-b^3(Ac^3-Cc^3-3Bdc^2-3Ad^2c+9Cd^2c+3Bd^3)a^2-b^4(2Bc^3+6Adc^2-6Cdc^2-6Bd^2c-2Ad^3-Cd^3)a-(a^2+b^2)^2d^2}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^3}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 3042

$$2 \int \frac{3Cda^3a^5 - bd^2(3cC + Bd)a^4 + 6b^2Cd^3a^3 - b^3(Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2c + 9Cd^2c + 3Bd^3)a^2 - b^4(2Bc^3 + 6Adc^2 - 6Cdc^2 - 6Bd^2c - 2Ad^3 - Cd^3)a - (a^2 + b^2)^2 d^2}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 4109

$$\frac{(-3Cda^4 + bBda^3 + b^2(2Bc + (A - 7C)d)a^2 - b^3(4Ac - 4Cc - 5Bd)a - b^4(2Bc + 3Ad))(c + d \tan(e + fx))^2}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2(-3Cda^4 + bBda^3 + b^2(Bc - 6Cd)a^2 - b^3(2Bc + 3Ad))}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 3042

$$\frac{(-3Cda^4 + bBda^3 + b^2(2Bc + (A - 7C)d)a^2 - b^3(4Ac - 4Cc - 5Bd)a - b^4(2Bc + 3Ad))(c + d \tan(e + fx))^2}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2(-3Cda^4 + bBda^3 + b^2(Bc - 6Cd)a^2 - b^3(2Bc + 3Ad))}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 3956

$$\frac{(-3Cda^4 + bBda^3 + b^2(2Bc + (A - 7C)d)a^2 - b^3(4Ac - 4Cc - 5Bd)a - b^4(2Bc + 3Ad))(c + d \tan(e + fx))^2}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2(-3Cda^4 + bBda^3 + b^2(Bc - 6Cd)a^2 - b^3(2Bc + 3Ad))}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 4100

$$\frac{(-3Cda^4 + bBda^3 + b^2(2Bc + (A - 7C)d)a^2 - b^3(4Ac - 4Cc - 5Bd)a - b^4(2Bc + 3Ad))(c + d \tan(e + fx))^2}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2(-3Cda^4 + bBda^3 + b^2(Bc - 6Cd)a^2 - b^3(2Bc + 3Ad))}{\dots}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 16

$$\frac{(-3Cda^4 + bBda^3 + b^2(2Bc + (A - 7C)d)a^2 - b^3(4Ac - 4Cc - 5Bd)a - b^4(2Bc + 3Ad))(c + d \tan(e + fx))^2}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2(-3Cda^4 + bBda^3 + b^2(Bc - 6Cd)a^2 - b^3(2Bc + 3Ad))}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^3}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

input

```
Int[((c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

```
-1/2*((A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^3)/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^2) + (((a^3*b*B*d - 3*a^4*C*d - b^4*(2*B*c + 3*A*d) - a*b^3*(4*A*c - 4*c*C - 5*B*d) + a^2*b^2*(2*B*c + (A - 7*C)*d))*(c + d*Tan[e + f*x])^2)/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])) + ((-2*(-((b^3*(a^3*(A*c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3) + 3*a*b^2*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) + 3*a^2*b*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2))) - b^3*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2))))*x)/(a^2 + b^2)) - (b^3*(3*a^2*b*(A*c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3) + b^3*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3 - A*(c^3 - 3*c*d^2)) - a^3*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)) + 3*a*b^2*((A - C)*d*(3*c^2 - d^2) + B*(c^3 - 3*c*d^2)))*Log[Cos[e + f*x]])/((a^2 + b^2)*f) + ((b*c - a*d)*(a^5*b*B*d^2 - 3*a^6*C*d^2 + a^4*b^2*d*(B*c - 9*C*d) + a^3*b^3*B*(c^2 + 3*d^2) - b^6*(c*(c*C + 3*B*d) - A*(c^2 - 3*d^2)) - a*b^5*(8*c*(A - C)*d + 3*B*(c^2 - 2*d^2)) + a^2*b^4*(3*c^2*C + 6*B*c*d - 10*C*d^2 - A*(3*c^2 - d^2)))*Log[a + b*Tan[e + f*x]])/(b*(a^2 + b^2)*f)))/b - (2*d^2*(a^3*b*B*d - 3*a^4*C*d - a*b^3*(2*A*c - 2*c*C - 3*B*d) + a^2*b^2*(B*c - 6*C*d) - b^4*(B*c + (2*A + C)*d))*Tan[e + f*x])/(b*f))/(b*(a^2 + b^2)))/(2*b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4100 $\text{Int}[((a_)+(b_)*\tan[(e_)+(f_)*(x_)])^{(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x_)])^2}, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[((A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)/((a_)+(b_)*\tan[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A*b - a*B - b*C, 0]$
- rule 4120 $\text{Int}[((a_)+(b_)*\tan[(e_)+(f_)*(x_)])^{(c_)+(d_)*\tan[(e_)+(f_)*(x_)]^{(n_)*((A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[b*C*\text{Tan}[e + f*x]*((c + d*\text{Tan}[e + f*x])^{(n + 1)/(d*f*(n + 2))}), x] - \text{Simp}[1/(d*(n + 2)) \text{ Int}[(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*\text{Tan}[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ !\text{LtQ}[n, -1]$

rule 4128

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]

```

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 1271, normalized size of antiderivative = 1.59

method	result	size
derivativdivides	Expression too large to display	1271
default	Expression too large to display	1271
norman	Expression too large to display	2076
parallelrisch	Expression too large to display	6687
risch	Expression too large to display	6825

input

```

int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,
x,method=_RETURNVERBOSE)

```


output

```

1/f*(tan(f*x+e)*C*d^3/b^3+1/(a^2+b^2)^3*(1/2*(3*A*a^3*c^2*d-A*a^3*d^3-3*A*
a^2*b*c^3+9*A*a^2*b*c*d^2-9*A*a*b^2*c^2*d+3*A*a*b^2*d^3+A*b^3*c^3-3*A*b^3*
c*d^2+B*a^3*c^3-3*B*a^3*c*d^2+9*B*a^2*b*c^2*d-3*B*a^2*b*d^3-3*B*a*b^2*c^3+
9*B*a*b^2*c*d^2-3*B*b^3*c^2*d+B*b^3*d^3-3*C*a^3*c^2*d+C*a^3*d^3+3*C*a^2*b*
c^3-9*C*a^2*b*c*d^2+9*C*a*b^2*c^2*d-3*C*a*b^2*d^3-C*b^3*c^3+3*C*b^3*c*d^2)
*ln(1+tan(f*x+e)^2)+(A*a^3*c^3-3*A*a^3*c*d^2+9*A*a^2*b*c^2*d-3*A*a^2*b*d^3
-3*A*a*b^2*c^3+9*A*a*b^2*c*d^2-3*A*b^3*c^2*d+A*b^3*d^3-3*B*a^3*c^2*d+B*a^3
*d^3+3*B*a^2*b*c^3-9*B*a^2*b*c*d^2+9*B*a*b^2*c^2*d-3*B*a*b^2*d^3-B*b^3*c^3
+3*B*b^3*c*d^2-C*a^3*c^3+3*C*a^3*c*d^2-9*C*a^2*b*c^2*d+3*C*a^2*b*d^3+3*C*a
*b^2*c^3-9*C*a*b^2*c*d^2+3*C*b^3*c^2*d-C*b^3*d^3)*arctan(tan(f*x+e)))-1/2*
(-A*a^3*b^2*d^3+3*A*a^2*b^3*c*d^2-3*A*a*b^4*c^2*d+A*b^5*c^3+B*a^4*b*d^3-3*
B*a^3*b^2*c*d^2+3*B*a^2*b^3*c^2*d-B*a*b^4*c^3-C*a^5*d^3+3*C*a^4*b*c*d^2-3*
C*a^3*b^2*c^2*d+C*a^2*b^3*c^3)/b^4/(a^2+b^2)/(a+b*tan(f*x+e))^2-1/b^4*(A*a
^4*b^2*d^3-3*A*a^2*b^4*c^2*d+3*A*a^2*b^4*d^3+2*A*a*b^5*c^3-6*A*a*b^5*c*d^2
+3*A*b^6*c^2*d-2*B*a^5*b*d^3+3*B*a^4*b^2*c*d^2-4*B*a^3*b^3*d^3-B*a^2*b^4*c
^3+9*B*a^2*b^4*c*d^2-6*B*a*b^5*c^2*d+B*b^6*c^3+3*C*a^6*d^3-6*C*a^5*b*c*d^2
+3*C*a^4*b^2*c^2*d+5*C*a^4*b^2*d^3-12*C*a^3*b^3*c*d^2+9*C*a^2*b^4*c^2*d-2*
C*a*b^5*c^3)/(a^2+b^2)^2/(a+b*tan(f*x+e))+(-3*A*a^3*b^4*c^2*d+A*a^3*b^4*d^
3+3*A*a^2*b^5*c^3-9*A*a^2*b^5*c*d^2+9*A*a*b^6*c^2*d-3*A*a*b^6*d^3-A*b^7*c^
3+3*A*b^7*c*d^2+B*a^6*b*d^3+3*B*a^4*b^3*d^3-B*a^3*b^4*c^3+3*B*a^3*b^4*c...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2549 vs. $2(794) = 1588$.

Time = 1.37 (sec) , antiderivative size = 2549, normalized size of antiderivative = 3.19

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```

integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+
e))^3,x, algorithm="fricas")

```

output

```

1/2*(2*(C*a^6*b^3 + 3*C*a^4*b^5 + 3*C*a^2*b^7 + C*b^9)*d^3*tan(f*x + e)^3
- (3*C*a^4*b^5 - 5*B*a^3*b^6 + (7*A - 3*C)*a^2*b^7 + B*a*b^8 + A*b^9)*c^3
+ 3*(C*a^5*b^4 - 3*B*a^4*b^5 + 5*(A - C)*a^3*b^6 + 3*B*a^2*b^7 - A*a*b^8)*
c^2*d + 3*(C*a^6*b^3 + B*a^5*b^4 - (3*A - 7*C)*a^4*b^5 - 5*B*a^3*b^6 + 3*A
*a^2*b^7)*c*d^2 - (3*C*a^7*b^2 - B*a^6*b^3 - (A - 9*C)*a^5*b^4 - 7*B*a^4*b
^5 + 5*A*a^3*b^6)*d^3 + 2*(((A - C)*a^5*b^4 + 3*B*a^4*b^5 - 3*(A - C)*a^3*
b^6 - B*a^2*b^7)*c^3 - 3*(B*a^5*b^4 - 3*(A - C)*a^4*b^5 - 3*B*a^3*b^6 + (A
- C)*a^2*b^7)*c^2*d - 3*((A - C)*a^5*b^4 + 3*B*a^4*b^5 - 3*(A - C)*a^3*b
^6 - B*a^2*b^7)*c*d^2 + (B*a^5*b^4 - 3*(A - C)*a^4*b^5 - 3*B*a^3*b^6 + (A
- C)*a^2*b^7)*d^3)*f*x + ((C*a^4*b^5 - 3*B*a^3*b^6 + 5*(A - C)*a^2*b^7 + 3*
B*a*b^8 - A*b^9)*c^3 + 3*(C*a^5*b^4 + B*a^4*b^5 - (3*A - 7*C)*a^3*b^6 - 5*
B*a^2*b^7 + 3*A*a*b^8)*c^2*d - 3*(3*C*a^6*b^3 - B*a^5*b^4 - (A - 9*C)*a^4*
b^5 - 7*B*a^3*b^6 + 5*A*a^2*b^7)*c*d^2 + (9*C*a^7*b^2 - 3*B*a^6*b^3 + (A +
23*C)*a^5*b^4 - 9*B*a^4*b^5 + (7*A + 12*C)*a^3*b^6 + 4*C*a*b^8)*d^3 + 2*(
((A - C)*a^3*b^6 + 3*B*a^2*b^7 - 3*(A - C)*a*b^8 - B*b^9)*c^3 - 3*(B*a^3*b
^6 - 3*(A - C)*a^2*b^7 - 3*B*a*b^8 + (A - C)*b^9)*c^2*d - 3*((A - C)*a^3*b
^6 + 3*B*a^2*b^7 - 3*(A - C)*a*b^8 - B*b^9)*c*d^2 + (B*a^3*b^6 - 3*(A - C)
*a^2*b^7 - 3*B*a*b^8 + (A - C)*b^9)*d^3)*f*x)*tan(f*x + e)^2 - ((B*a^5*b^4
- 3*(A - C)*a^4*b^5 - 3*B*a^3*b^6 + (A - C)*a^2*b^7)*c^3 + 3*((A - C)*a^5
*b^4 + 3*B*a^4*b^5 - 3*(A - C)*a^3*b^6 - B*a^2*b^7)*c^2*d - 3*(C*a^8*b ...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

= Exception raised: AttributeError

input

```

integrate((c+d*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*
x+e))**3,x)

```

output

```

Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'

```

Maxima [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 1119, normalized size of antiderivative = 1.40

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="maxima")
```

output

```
1/2*(2*C*d^3*tan(f*x + e)/b^3 + 2*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c^3 - 3*(B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c^2*d - 3*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c*d^2 + (B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*d^3)*(f*x + e)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 2*((B*a^3*b^4 - 3*(A - C)*a^2*b^5 - 3*B*a*b^6 + (A - C)*b^7)*c^3 + 3*((A - C)*a^3*b^4 + 3*B*a^2*b^5 - 3*(A - C)*a*b^6 - B*b^7)*c^2*d - 3*(C*a^6*b + 3*C*a^4*b^3 + B*a^3*b^4 - 3*(A - 2*C)*a^2*b^5 - 3*B*a*b^6 + A*b^7)*c*d^2 + (3*C*a^7 - B*a^6*b + 9*C*a^5*b^2 - 3*B*a^4*b^3 - (A - 10*C)*a^3*b^4 - 6*B*a^2*b^5 + 3*A*a*b^6)*d^3)*log(b*tan(f*x + e) + a)/(a^6*b^4 + 3*a^4*b^6 + 3*a^2*b^8 + b^10) + ((B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c^3 + 3*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c^2*d - 3*(B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c*d^2 - ((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*d^3)*log(tan(f*x + e)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((C*a^4*b^3 - 3*B*a^3*b^4 + (5*A - 3*C)*a^2*b^5 + B*a*b^6 + A*b^7)*c^3 + 3*(C*a^5*b^2 + B*a^4*b^3 - (3*A - 5*C)*a^3*b^4 - 3*B*a^2*b^5 + A*a*b^6)*c^2*d - 3*(3*C*a^6*b - B*a^5*b^2 - (A - 7*C)*a^4*b^3 - 5*B*a^3*b^4 + 3*A*a^2*b^5)*c*d^2 + (5*C*a^7 - 3*B*a^6*b + (A + 9*C)*a^5*b^2 - 7*B*a^4*b^3 + 5*A*a^3*b^4)*d^3 - 2*((B*a^2*b^5 - 2*(A - C)*a*b^6 - B*b^7)*c^3 - 3*(C*a^4*b^3 - (A - 3*C)*a^2*b^5 - 2*B*a*b^6 + A*b^7)*c^2*d + 3*(2*C*a^5*b^2 - B*a^4*b^3 + 4*C*a^3*b^4 - ...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1758 vs. $2(794) = 1588$.

Time = 0.77 (sec) , antiderivative size = 1758, normalized size of antiderivative = 2.20

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="giac")
```

output

```
(A*a^3*c^3 - C*a^3*c^3 + 3*B*a^2*b*c^3 - 3*A*a*b^2*c^3 + 3*C*a*b^2*c^3 - B*b^3*c^3 - 3*B*a^3*c^2*d + 9*A*a^2*b*c^2*d - 9*C*a^2*b*c^2*d + 9*B*a*b^2*c^2*d - 3*A*b^3*c^2*d + 3*C*b^3*c^2*d - 3*A*a^3*c*d^2 + 3*C*a^3*c*d^2 - 9*B*a^2*b*c*d^2 + 9*A*a*b^2*c*d^2 - 9*C*a*b^2*c*d^2 + 3*B*b^3*c*d^2 + B*a^3*d^3 - 3*A*a^2*b*d^3 + 3*C*a^2*b*d^3 - 3*B*a*b^2*d^3 + A*b^3*d^3 - C*b^3*d^3)*(f*x + e)/(a^6*f + 3*a^4*b^2*f + 3*a^2*b^4*f + b^6*f) + 1/2*(B*a^3*c^3 - 3*A*a^2*b*c^3 + 3*C*a^2*b*c^3 - 3*B*a*b^2*c^3 + A*b^3*c^3 - C*b^3*c^3 + 3*A*a^3*c^2*d - 3*C*a^3*c^2*d + 9*B*a^2*b*c^2*d - 9*A*a*b^2*c^2*d + 9*C*a*b^2*c^2*d - 3*B*b^3*c^2*d - 3*B*a^3*c*d^2 + 9*A*a^2*b*c*d^2 - 9*C*a^2*b*c*d^2 + 9*B*a*b^2*c*d^2 - 3*A*b^3*c*d^2 + 3*C*b^3*c*d^2 - A*a^3*d^3 + C*a^3*d^3 - 3*B*a^2*b*d^3 + 3*A*a*b^2*d^3 - 3*C*a*b^2*d^3 + B*b^3*d^3)*log(tan(f*x + e)^2 + 1)/(a^6*f + 3*a^4*b^2*f + 3*a^2*b^4*f + b^6*f) - (B*a^3*b^4*c^3 - 3*A*a^2*b^5*c^3 + 3*C*a^2*b^5*c^3 - 3*B*a*b^6*c^3 + A*b^7*c^3 - C*b^7*c^3 + 3*A*a^3*b^4*c^2*d - 3*C*a^3*b^4*c^2*d + 9*B*a^2*b^5*c^2*d - 9*A*a*b^6*c^2*d + 9*C*a*b^6*c^2*d - 3*B*b^7*c^2*d - 3*C*a^6*b*c*d^2 - 9*C*a^4*b^3*c*d^2 - 3*B*a^3*b^4*c*d^2 + 9*A*a^2*b^5*c*d^2 - 18*C*a^2*b^5*c*d^2 + 9*B*a*b^6*c*d^2 - 3*A*b^7*c*d^2 + 3*C*a^7*d^3 - B*a^6*b*d^3 + 9*C*a^5*b^2*d^3 - 3*B*a^4*b^3*d^3 - A*a^3*b^4*d^3 + 10*C*a^3*b^4*d^3 - 6*B*a^2*b^5*d^3 + 3*A*a*b^6*d^3)*log(abs(b*tan(f*x + e) + a))/(a^6*b^4*f + 3*a^4*b^6*f + 3*a^2*b^8*f + b^10*f) + C*d^3*tan(f*x + e)/(b^3*f) - 1/2*(C*a^6*b^3*c^3 - 3*B...
```

Mupad [B] (verification not implemented)

Time = 14.87 (sec) , antiderivative size = 1172, normalized size of antiderivative = 1.47

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int(((c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a +
b*tan(e + f*x))^3,x)
```

output

```
(log(tan(e + f*x) + 1i)*(A*c^3 + A*d^3*1i - B*c^3*1i + B*d^3 - C*c^3 - C*d
^3*1i - 3*A*c*d^2 - A*c^2*d^3*1i + B*c*d^2*3i - 3*B*c^2*d + 3*C*c*d^2 + C*c^
2*d^3*1i))/(2*f*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3)) - ((tan(e + f*x)*(B*b^6
*c^3 + 3*C*a^6*d^3 + 2*A*a*b^5*c^3 - 2*B*a^5*b*d^3 - 2*C*a*b^5*c^3 + 3*A*b
^6*c^2*d + 3*A*a^2*b^4*d^3 + A*a^4*b^2*d^3 - B*a^2*b^4*c^3 - 4*B*a^3*b^3*d
^3 + 5*C*a^4*b^2*d^3 - 3*A*a^2*b^4*c^2*d + 9*B*a^2*b^4*c*d^2 + 3*B*a^4*b^2
*c*d^2 + 9*C*a^2*b^4*c^2*d - 12*C*a^3*b^3*c*d^2 + 3*C*a^4*b^2*c^2*d - 6*A
a*b^5*c*d^2 - 6*B*a*b^5*c^2*d - 6*C*a^5*b*c*d^2))/(a^4 + b^4 + 2*a^2*b^2)
+ (A*b^7*c^3 + 5*C*a^7*d^3 + B*a*b^6*c^3 - 3*B*a^6*b*d^3 + 5*A*a^2*b^5*c^3
+ 5*A*a^3*b^4*d^3 + A*a^5*b^2*d^3 - 3*B*a^3*b^4*c^3 - 7*B*a^4*b^3*d^3 - 3
*C*a^2*b^5*c^3 + C*a^4*b^3*c^3 + 9*C*a^5*b^2*d^3 - 9*A*a^2*b^5*c*d^2 - 9*A
a^3*b^4*c^2*d + 3*A*a^4*b^3*c*d^2 - 9*B*a^2*b^5*c^2*d + 15*B*a^3*b^4*c*d^
2 + 3*B*a^4*b^3*c^2*d + 3*B*a^5*b^2*c*d^2 + 15*C*a^3*b^4*c^2*d - 21*C*a^4*
b^3*c*d^2 + 3*C*a^5*b^2*c^2*d + 3*A*a*b^6*c^2*d - 9*C*a^6*b*c*d^2)/(2*b*(a
^4 + b^4 + 2*a^2*b^2)))/(f*(a^2*b^3 + b^5*tan(e + f*x)^2 + 2*a*b^4*tan(e +
f*x))) + (log(a + b*tan(e + f*x))*(b^3*(3*B*a^4*d^3 + 9*C*a^4*c*d^2) - b^
6*(3*A*a*d^3 - 3*B*a*c^3 - 9*A*a*c^2*d + 9*B*a*c*d^2 + 9*C*a*c^2*d) + b^5*
(3*A*a^2*c^3 + 6*B*a^2*d^3 - 3*C*a^2*c^3 - 9*A*a^2*c*d^2 - 9*B*a^2*c^2*d +
18*C*a^2*c*d^2) + b^4*(A*a^3*d^3 - B*a^3*c^3 - 10*C*a^3*d^3 - 3*A*a^3*c^2
*d + 3*B*a^3*c*d^2 + 3*C*a^3*c^2*d) + b*(B*a^6*d^3 + 3*C*a^6*c*d^2) + b...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 4720, normalized size of antiderivative = 5.91

$$\int \frac{(c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,
x)
```

output

```
(3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**6*c**2*d - log(tan(e +
f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**6*d**3 - 2*log(tan(e + f*x)**2 + 1)*
tan(e + f*x)**2*a**4*b**7*c**3 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**
2*a**4*b**7*c*d**2 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*b**6*
c**3*d + log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*b**6*c*d**3 + 3*log
(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**7*c**4 - 9*log(tan(e + f*x)*
**2 + 1)*tan(e + f*x)**2*a**3*b**7*c**2*d**2 - 2*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**2*a**2*b**9*c**3 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2
*a**2*b**9*c*d**2 + 9*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**8*c
**3*d - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**8*c*d**3 - 3*lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**10*c**2*d + log(tan(e + f*x)**
2 + 1)*tan(e + f*x)**2*a*b**10*d**3 - log(tan(e + f*x)**2 + 1)*tan(e + f*x
)**2*a*b**9*c**4 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**9*c**2*
d**2 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**6*b**5*c**2*d - 2*log(ta
n(e + f*x)**2 + 1)*tan(e + f*x)*a**6*b**5*d**3 - 4*log(tan(e + f*x)**2 + 1
)*tan(e + f*x)*a**5*b**6*c**3 + 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a
**5*b**6*c*d**2 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*b**5*c**3*d
+ 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*b**5*c*d**3 + 6*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)*a**4*b**6*c**4 - 18*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)*a**4*b**6*c**2*d**2 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*...
```

3.70
$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

Optimal result	778
Mathematica [C] (verified)	779
Rubi [A] (verified)	779
Maple [A] (verified)	784
Fricas [A] (verification not implemented)	785
Sympy [C] (verification not implemented)	786
Maxima [A] (verification not implemented)	787
Giac [A] (verification not implemented)	787
Mupad [B] (verification not implemented)	788
Reduce [B] (verification not implemented)	789

Optimal result

Integrand size = 45, antiderivative size = 337

$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

$$= \frac{(a^3(Ac - cC + Bd) - 3ab^2(Ac - cC + Bd) - 3a^2b(Bc - (A - C)d) + b^3(Bc - (A - C)d)) x}{c^2 + d^2}$$

$$- \frac{(3a^2b(Ac - cC + Bd) - b^3(Ac - cC + Bd) + a^3(Bc - (A - C)d) - 3ab^2(Bc - (A - C)d)) \log(\cos(e+fx))}{(c^2 + d^2) f}$$

$$- \frac{(bc - ad)^3 (c^2C - Bcd + Ad^2) \log(c + d \tan(e + fx))}{d^4 (c^2 + d^2) f}$$

$$+ \frac{b(b(Ab + aB - bC)d^2 + (bc - ad)(bcC - bBd - aCd)) \tan(e + fx)}{d^3 f}$$

$$- \frac{(bcC - bBd - aCd)(a + b \tan(e + fx))^2}{2d^2 f} + \frac{C(a + b \tan(e + fx))^3}{3df}$$

output

```
(a^3*(A*c+B*d-C*c)-3*a*b^2*(A*c+B*d-C*c)-3*a^2*b*(B*c-(A-C)*d)+b^3*(B*c-(A-C)*d))*x/(c^2+d^2)-(3*a^2*b*(A*c+B*d-C*c)-b^3*(A*c+B*d-C*c)+a^3*(B*c-(A-C)*d)-3*a*b^2*(B*c-(A-C)*d))*ln(cos(f*x+e))/(c^2+d^2)/f-(-a*d+b*c)^3*(A*d^2-B*c*d+C*c^2)*ln(c+d*tan(f*x+e))/d^4/(c^2+d^2)/f+b*(b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-B*b*d-C*a*d+C*b*c))*tan(f*x+e)/d^3/f-1/2*(-B*b*d-C*a*d+C*b*c)*(a+b*tan(f*x+e))^2/d^2/f+1/3*C*(a+b*tan(f*x+e))^3/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.92 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.77

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{3(a+ib)^3(-iA+B+iC)d^2 \log(i-\tan(e+fx))}{c+id} + \frac{3(a-ib)^3(iA+B-iC)d^2 \log(i+\tan(e+fx))}{c-id} + \frac{6(-bc+ad)^3(c^2C-Bcd+Ad^2) \log(c+d \tan(e+fx))}{d^2(c^2+d^2)}$$

input

```
Integrate[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x]),x]
```

output

```
((3*(a + I*b)^3*((-I)*A + B + I*C)*d^2*Log[I - Tan[e + f*x]])/(c + I*d) +
(3*(a - I*b)^3*(I*A + B - I*C)*d^2*Log[I + Tan[e + f*x]])/(c - I*d) + (6*(
-(b*c) + a*d)^3*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]])/(d^2*(c^2
+ d^2)) + 6*b^2*(A*b + a*B - b*C)*d*Tan[e + f*x] - (6*b*(b*c - a*d)*(-(b*
c*C) + b*B*d + a*C*d)*Tan[e + f*x])/d - 3*(b*c*C - b*B*d - a*C*d)*(a + b*T
an[e + f*x])^2 + 2*C*d*(a + b*Tan[e + f*x])^3)/(6*d^2*f)
```

Rubi [A] (verified)

Time = 3.98 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4120, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{c + d \tan(e + fx)} dx$$

$$\begin{aligned}
 & \int \frac{-3(a+b \tan(e+fx))^2((bcC-adC-bBd) \tan^2(e+fx)-(Ab-Cb+aB)d \tan(e+fx)+bcC-aAd)}{c+d \tan(e+fx)} dx + \\
 & \quad \frac{3d}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 4130 \\
 & \int \frac{(a+b \tan(e+fx))^2((bcC-adC-bBd) \tan^2(e+fx)-(Ab-Cb+aB)d \tan(e+fx)+bcC-aAd)}{c+d \tan(e+fx)} dx - \\
 & \quad \frac{3df}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 27 \\
 & \int \frac{(a+b \tan(e+fx))^2((bcC-adC-bBd) \tan^2(e+fx)-(Ab-Cb+aB)d \tan(e+fx)+bcC-aAd)}{c+d \tan(e+fx)} dx \\
 & \quad \frac{d}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a+b \tan(e+fx))^2((bcC-adC-bBd) \tan^2(e+fx)-(Ab-Cb+aB)d \tan(e+fx)+bcC-aAd)}{c+d \tan(e+fx)} dx \\
 & \quad \frac{d}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 4130 \\
 & \int \frac{2(a+b \tan(e+fx))(-c(cC-Bd)b^2+2acCdb-a^2Ad^2-(b(Ab-Cb+aB)d^2+(bc-ad)(bcC-adC-bBd)) \tan^2(e+fx)-(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx))}{c+d \tan(e+fx)} dx \\
 & \quad \frac{2d}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 27 \\
 & \int \frac{(a+b \tan(e+fx))(-c(cC-Bd)b^2+2acCdb-a^2Ad^2-(b(Ab-Cb+aB)d^2+(bc-ad)(bcC-adC-bBd)) \tan^2(e+fx)-(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx))}{c+d \tan(e+fx)} dx \\
 & \quad \frac{d}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a+b \tan(e+fx))(-c(cC-Bd)b^2+2acCdb-a^2Ad^2-(b(Ab-Cb+aB)d^2+(bc-ad)(bcC-adC-bBd)) \tan^2(e+fx)-(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx))}{c+d \tan(e+fx)} dx \\
 & \quad \frac{d}{C(a+b \tan(e+fx))^3} \\
 & \quad \downarrow 4120
 \end{aligned}$$

$$\frac{C(a + b \tan(e + fx))^3}{3df} - \frac{\int -\frac{c(Cc^2 - Bdc + (A-C)d^2)b^3 - 3acd(cC - Bd)b^2 + 3a^2cCd^2b - a^3Ad^3 - \left(-\left((Cc^3 - Bdc^2 + (A-C)d^2c + Bd^3)b^3\right) + 3ad(Cc^2 - Bdc + (A-C)d^2)b^2 - 3a^2d^2(cC - B) \right)}{c + d \tan(e + fx)}}{d}$$

25

$$\frac{C(a + b \tan(e + fx))^3}{3df} - \frac{\int \frac{c(Cc^2 - Bdc + (A-C)d^2)b^3 - 3acd(cC - Bd)b^2 + 3a^2cCd^2b - a^3Ad^3 - \left(-\left((Cc^3 - Bdc^2 + (A-C)d^2c + Bd^3)b^3\right) + 3ad(Cc^2 - Bdc + (A-C)d^2)b^2 - 3a^2d^2(cC - Bd)b \right)}{c + d \tan(e + fx)}}{d}$$

3042

$$\frac{C(a + b \tan(e + fx))^3}{3df} - \frac{\int \frac{c(Cc^2 - Bdc + (A-C)d^2)b^3 - 3acd(cC - Bd)b^2 + 3a^2cCd^2b - a^3Ad^3 - \left(-\left((Cc^3 - Bdc^2 + (A-C)d^2c + Bd^3)b^3\right) + 3ad(Cc^2 - Bdc + (A-C)d^2)b^2 - 3a^2d^2(cC - Bd)b \right)}{c + d \tan(e + fx)}}{d}$$

4109

$$\frac{C(a + b \tan(e + fx))^3}{3df} - \frac{d^3(a^3(Bc - d(A - C)) + 3a^2b(Ac + Bd - cC) - 3ab^2(Bc - d(A - C)) - b^3(Ac + Bd - cC)) \int \tan(e + fx) dx}{c^2 + d^2} + \frac{(bc - ad)^3(Ad^2 - Bcd + c^2C) \int \frac{\tan^2(e + fx) + 1}{c + d \tan(e + fx)} dx}{c^2 + d^2} - \frac{d^3x(a^3(Bc - d(A - C)) + 3a^2b(Ac + Bd - cC) - 3ab^2(Bc - d(A - C)) - b^3(Ac + Bd - cC))}{d}$$

3042

$$\frac{C(a + b \tan(e + fx))^3}{3df} - \frac{d^3(a^3(Bc - d(A - C)) + 3a^2b(Ac + Bd - cC) - 3ab^2(Bc - d(A - C)) - b^3(Ac + Bd - cC)) \int \tan(e + fx) dx}{c^2 + d^2} + \frac{(bc - ad)^3(Ad^2 - Bcd + c^2C) \int \frac{\tan(e + fx)^2 + 1}{c + d \tan(e + fx)} dx}{c^2 + d^2} - \frac{d^3x(a^3(Bc - d(A - C)) + 3a^2b(Ac + Bd - cC) - 3ab^2(Bc - d(A - C)) - b^3(Ac + Bd - cC))}{d}$$

3956

$$\frac{C(a + b \tan(e + fx))^3}{3df} - \frac{(bc - ad)^3(Ad^2 - Bcd + c^2C) \int \frac{\tan(e + fx)^2 + 1}{c + d \tan(e + fx)} dx}{c^2 + d^2} + \frac{d^3 \log(\cos(e + fx))(a^3(Bc - d(A - C)) + 3a^2b(Ac + Bd - cC) - 3ab^2(Bc - d(A - C)) - b^3(Ac + Bd - cC))}{f(c^2 + d^2)} - \frac{d^3x(a^3(Bc - d(A - C)) + 3a^2b(Ac + Bd - cC) - 3ab^2(Bc - d(A - C)) - b^3(Ac + Bd - cC))}{d}$$

$$\begin{aligned} & \downarrow 4100 \\ & \frac{C(a + b \tan(e + fx))^3}{3df} - \\ & \frac{(bc-ad)^3 (Ad^2 - Bcd + c^2 C) \int \frac{1}{c+d \tan(e+fx)} d(d \tan(e+fx))}{df(c^2+d^2)} + \frac{d^3 \log(\cos(e+fx)) (a^3(Bc-d(A-C)) + 3a^2b(Ac+Bd-cC) - 3ab^2(Bc-d(A-C)) - b^3(Ac+Bd-cC))}{f(c^2+d^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 16 \\ & \frac{C(a + b \tan(e + fx))^3}{3df} - \\ & \frac{d^3 \log(\cos(e+fx)) (a^3(Bc-d(A-C)) + 3a^2b(Ac+Bd-cC) - 3ab^2(Bc-d(A-C)) - b^3(Ac+Bd-cC))}{f(c^2+d^2)} - \frac{d^3 x (a^3(Ac+Bd-cC) - 3a^2b(Bc-d(A-C)) - 3ab^2(Ac+Bd-cC))}{c^2+d^2} \end{aligned}$$

input

```
Int[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]),x]
```

output

```
(C*(a + b*Tan[e + f*x])^3)/(3*d*f) - (((b*c*C - b*B*d - a*C*d)*(a + b*Tan[e + f*x])^2)/(2*d*f) + (((-((d^3*(a^3*(A*c - c*C + B*d) - 3*a*b^2*(A*c - c*C + B*d) - 3*a^2*b*(B*c - (A - C)*d) + b^3*(B*c - (A - C)*d))*x)/(c^2 + d^2)) + (d^3*(3*a^2*b*(A*c - c*C + B*d) - b^3*(A*c - c*C + B*d) + a^3*(B*c - (A - C)*d) - 3*a*b^2*(B*c - (A - C)*d))*Log[Cos[e + f*x]]/(c^2 + d^2)*f) + ((b*c - a*d)^3*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]]/(d*(c^2 + d^2)*f))/d - (b*(b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(b*c*C - b*B*d - a*C*d))*Tan[e + f*x])/(d*f))/d
```

Defintions of rubi rules used

rule 16

```
Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`
- rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2] / ((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`
- rule 4120 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 500, normalized size of antiderivative = 1.48

method	result
norman	$\frac{(A a^3 c + 3 A a^2 b d - 3 A a b^2 c - A b^3 d + B a^3 d - 3 B a^2 b c - 3 B a b^2 d + B b^3 c - C a^3 c - 3 C a^2 b d + 3 C a b^2 c + C b^3 d) x}{c^2 + d^2} + \frac{(A b^2 d^2 - b \left(\frac{C b^2 d^2 \tan^3(fx+e)}{3} + \frac{B b^2 d^2 \tan^2(fx+e)}{2} + \frac{3 C a b d^2 \tan(fx+e)}{2} - \frac{C b^2 c d \tan(fx+e)}{2} + \tan(fx+e) A b^2 d^2 + 3 \tan(fx+e) B a b d^2 - \right)}{d^3}$
derivativdivides	
default	$b \left(\frac{C b^2 d^2 \tan^3(fx+e)}{3} + \frac{B b^2 d^2 \tan^2(fx+e)}{2} + \frac{3 C a b d^2 \tan(fx+e)}{2} - \frac{C b^2 c d \tan(fx+e)}{2} + \tan(fx+e) A b^2 d^2 + 3 \tan(fx+e) B a b d^2 - \right) / d^3$
parallelrisc	Expression too large to display
risc	Expression too large to display

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, method=_RETURNVERBOSE)
```

output

```
(A*a^3*c+3*A*a^2*b*d-3*A*a*b^2*c-A*b^3*d+B*a^3*d-3*B*a^2*b*c-3*B*a*b^2*d+B
*b^3*c-C*a^3*c-3*C*a^2*b*d+3*C*a*b^2*c+C*b^3*d)/(c^2+d^2)*x+(A*b^2*d^2+3*B
*a*b*d^2-B*b^2*c*d+3*C*a^2*d^2-3*C*a*b*c*d+C*b^2*c^2-C*b^2*d^2)*b/f/d^3*ta
n(f*x+e)+1/3*C*b^3/f/d*tan(f*x+e)^3+1/2*b^2*(B*b*d+3*C*a*d-C*b*c)/d^2/f*ta
n(f*x+e)^2+(A*a^3*d^5-3*A*a^2*b*c*d^4+3*A*a*b^2*c^2*d^3-A*b^3*c^3*d^2-B*a^
3*c*d^4+3*B*a^2*b*c^2*d^3-3*B*a*b^2*c^3*d^2+B*b^3*c^4*d+C*a^3*c^2*d^3-3*C*
a^2*b*c^3*d^2+3*C*a*b^2*c^4*d-C*b^3*c^5)/(c^2+d^2)/d^4/f*ln(c+d*tan(f*x+e)
)-1/2*(A*a^3*d-3*A*a^2*b*c-3*A*a*b^2*d+A*b^3*c-B*a^3*c-3*B*a^2*b*d+3*B*a*b
^2*c+B*b^3*d-C*a^3*d+3*C*a^2*b*c+3*C*a*b^2*d-C*b^3*c)/f/(c^2+d^2)*ln(1+tan
(f*x+e)^2)
```

Fricas [A] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 627, normalized size of antiderivative = 1.86

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{2(Cb^3c^2d^3 + Cb^3d^5) \tan(fx + e)^3 + 6(((A - C)a^3 - 3Ba^2b - 3(A - C)ab^2 + Bb^3)cd^4 + (Ba^3 + 3(A$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e)),x, algorithm="fricas")
```

output

```
1/6*(2*(C*b^3*c^2*d^3 + C*b^3*d^5)*tan(f*x + e)^3 + 6*(((A - C)*a^3 - 3*B*
a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c*d^4 + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*
b^2 - (A - C)*b^3)*d^5)*f*x - 3*(C*b^3*c^3*d^2 + C*b^3*c*d^4 - (3*C*a*b^2
+ B*b^3)*c^2*d^3 - (3*C*a*b^2 + B*b^3)*d^5)*tan(f*x + e)^2 - 3*(C*b^3*c^5
- A*a^3*d^5 - (3*C*a*b^2 + B*b^3)*c^4*d + (3*C*a^2*b + 3*B*a*b^2 + A*b^3)*
c^3*d^2 - (C*a^3 + 3*B*a^2*b + 3*A*a*b^2)*c^2*d^3 + (B*a^3 + 3*A*a^2*b)*c
d^4)*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 +
1)) + 3*(C*b^3*c^5 - (3*C*a*b^2 + B*b^3)*c^4*d + (3*C*a^2*b + 3*B*a*b^2 +
A*b^3)*c^3*d^2 - (C*a^3 + 3*B*a^2*b + 3*A*a*b^2)*c^2*d^3 + (3*C*a^2*b + 3
*B*a*b^2 + (A - C)*b^3)*c*d^4 - (C*a^3 + 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b
^3)*d^5)*log(1/(tan(f*x + e)^2 + 1)) + 6*(C*b^3*c^4*d - (3*C*a*b^2 + B*b^3
)*c^3*d^2 + (3*C*a^2*b + 3*B*a*b^2 + A*b^3)*c^2*d^3 - (3*C*a*b^2 + B*b^3)*
c*d^4 + (3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*d^5)*tan(f*x + e))/((c^2*d^4
+ d^6)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 22.68 (sec) , antiderivative size = 7096, normalized size of antiderivative = 21.06

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)),x)
```

output

```
Piecewise((zoo*x*(a + b*tan(e))**3*(A + B*tan(e) + C*tan(e)**2)/tan(e), Eq(c, 0) & Eq(d, 0) & Eq(f, 0)), ((A*a**3*x + 3*A*a**2*b*log(tan(e + f*x)**2 + 1)/(2*f) - 3*A*a*b**2*x + 3*A*a*b**2*tan(e + f*x)/f - A*b**3*log(tan(e + f*x)**2 + 1)/(2*f) + A*b**3*tan(e + f*x)**2/(2*f) + B*a**3*log(tan(e + f*x)**2 + 1)/(2*f) - 3*B*a**2*b*x + 3*B*a**2*b*tan(e + f*x)/f - 3*B*a*b**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*a*b**2*tan(e + f*x)**2/(2*f) + B*b**3*x + B*b**3*tan(e + f*x)**3/(3*f) - B*b**3*tan(e + f*x)/f - C*a**3*x + C*a**3*tan(e + f*x)/f - 3*C*a**2*b*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C*a**2*b*tan(e + f*x)**2/(2*f) + 3*C*a*b**2*x + C*a*b**2*tan(e + f*x)**3/f - 3*C*a*b**2*tan(e + f*x)/f + C*b**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*b**3*tan(e + f*x)**4/(4*f) - C*b**3*tan(e + f*x)**2/(2*f))/c, Eq(d, 0)), (3*I*A*a**3*f*x*tan(e + f*x)/(6*d*f*tan(e + f*x) - 6*I*d*f) + 3*A*a**3*f*x/(6*d*f*tan(e + f*x) - 6*I*d*f) + 3*I*A*a**3/(6*d*f*tan(e + f*x) - 6*I*d*f) + 9*A*a**2*b*f*x*tan(e + f*x)/(6*d*f*tan(e + f*x) - 6*I*d*f) - 9*I*A*a**2*b*f*x/(6*d*f*tan(e + f*x) - 6*I*d*f) - 9*A*a**2*b/(6*d*f*tan(e + f*x) - 6*I*d*f) + 9*I*A*a*b**2*f*x*tan(e + f*x)/(6*d*f*tan(e + f*x) - 6*I*d*f) + 9*A*a*b**2*f*x/(6*d*f*tan(e + f*x) - 6*I*d*f) + 9*A*a*b**2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(6*d*f*tan(e + f*x) - 6*I*d*f) - 9*I*A*a*b**2*log(tan(e + f*x)**2 + 1)/(6*d*f*tan(e + f*x) - 6*I*d*f) - 9*I*A*a*b**2/(6*d*f*tan(e + f*x) - 6*I*d*f) - 9*A*b**3*f*x*tan(e + f*x)/(6*d*f*tan(e + f*x) - 6*I*d*f)...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.32

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{6(((A-C)a^3 - 3Ba^2b - 3(A-C)ab^2 + Bb^3)c + (Ba^3 + 3(A-C)a^2b - 3Bab^2 - (A-C)b^3)d)(fx+e)}{c^2+d^2} - \frac{6(Cb^3c^5 - Aa^3d^5 - (3Cab^2 + Bb^3)c^4d + (3$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="maxima")`

output `1/6*(6*(((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d)*(f*x + e)/(c^2 + d^2) - 6*(C*b^3*c^5 - A*a^3*d^5 - (3*C*a*b^2 + B*b^3)*c^4*d + (3*C*a^2*b + 3*B*a*b^2 + A*b^3)*c^3*d^2 - (C*a^3 + 3*B*a^2*b + 3*A*a*b^2)*c^2*d^3 + (B*a^3 + 3*A*a^2*b)*c*d^4)*log(d*tan(f*x + e) + c)/(c^2*d^4 + d^6) + 3*((B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c - ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*d)*log(tan(f*x + e)^2 + 1)/(c^2 + d^2) + (2*C*b^3*d^2*tan(f*x + e)^3 - 3*(C*b^3*c*d - (3*C*a*b^2 + B*b^3)*d^2)*tan(f*x + e)^2 + 6*(C*b^3*c^2 - (3*C*a*b^2 + B*b^3)*c*d + (3*C*a^2*b + 3*B*a*b^2 + (A - C)*b^3)*d^2)*tan(f*x + e))/d^3)/f`

Giac [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 601, normalized size of antiderivative = 1.78

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{(Aa^3c - Ca^3c - 3Ba^2bc - 3Aab^2c + 3Cab^2c + Bb^3c + Ba^3d + 3Aa^2bd - 3Ca^2bd - 3Bab^2d - Ab^3d + (Ba^3c + 3Aa^2bc - 3Ca^2bc - 3Bab^2c - Ab^3c + Cb^3c - Aa^3d + Ca^3d + 3Ba^2bd + 3Aab^2d - 3Cab^2c) + (Cb^3c^5 - 3Cab^2c^4d - Bb^3c^4d + 3Ca^2bc^3d^2 + 3Bab^2c^3d^2 + Ab^3c^3d^2 - Ca^3c^2d^3 - 3Ba^2bc^2d^3 - 3Aab^2c^2d^3) + 2Cb^3d^2f^2 \tan(fx + e)^3 - 3Cb^3cdf^2 \tan(fx + e)^2 + 9Cab^2d^2f^2 \tan(fx + e)^2 + 3Bb^3d^2f^2 \tan(fx + e))}{c^2f + d^2f}$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="giac")
```

output

```
(A*a^3*c - C*a^3*c - 3*B*a^2*b*c - 3*A*a*b^2*c + 3*C*a*b^2*c + B*b^3*c + B*a^3*d + 3*A*a^2*b*d - 3*C*a^2*b*d - 3*B*a*b^2*d - A*b^3*d + C*b^3*d)*(f*x + e)/(c^2*f + d^2*f) + 1/2*(B*a^3*c + 3*A*a^2*b*c - 3*C*a^2*b*c - 3*B*a*b^2*c - A*b^3*c + C*b^3*c - A*a^3*d + C*a^3*d + 3*B*a^2*b*d + 3*A*a*b^2*d - 3*C*a*b^2*d - B*b^3*d)*log(tan(f*x + e)^2 + 1)/(c^2*f + d^2*f) - (C*b^3*c^5 - 3*C*a*b^2*c^4*d - B*b^3*c^4*d + 3*C*a^2*b*c^3*d^2 + 3*B*a*b^2*c^3*d^2 + A*b^3*c^3*d^2 - C*a^3*c^2*d^3 - 3*B*a^2*b*c^2*d^3 - 3*A*a*b^2*c^2*d^3 + B*a^3*c*d^4 + 3*A*a^2*b*c*d^4 - A*a^3*d^5)*log(abs(d*tan(f*x + e) + c))/(c^2*d^4*f + d^6*f) + 1/6*(2*C*b^3*d^2*f^2*tan(f*x + e)^3 - 3*C*b^3*c*d*f^2*tan(f*x + e)^2 + 9*C*a*b^2*d^2*f^2*tan(f*x + e)^2 + 3*B*b^3*d^2*f^2*tan(f*x + e)^2 + 6*C*b^3*c^2*f^2*tan(f*x + e) - 18*C*a*b^2*c*d*f^2*tan(f*x + e) - 6*B*b^3*c*d*f^2*tan(f*x + e) + 18*C*a^2*b*d^2*f^2*tan(f*x + e) + 18*B*a*b^2*d^2*f^2*tan(f*x + e) + 6*A*b^3*d^2*f^2*tan(f*x + e) - 6*C*b^3*d^2*f^2*tan(f*x + e))/(d^3*f^3)
```

Mupad [B] (verification not implemented)

Time = 9.76 (sec) , antiderivative size = 508, normalized size of antiderivative = 1.51

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{\tan(e + fx)^2 \left(\frac{Bb^3 + 3Cab^2}{2d} - \frac{Cb^3c}{2d^2} \right)}{f \left(\frac{c \left(\frac{Bb^3 + 3Cab^2}{d} - \frac{Cb^3e}{d^2} \right)}{d} - \frac{3Ca^2b + 3Bab^2 + Ab^3}{d} + \frac{Cb^3}{d} \right)}$$

$$- \frac{\ln(c + d \tan(e + fx)) (d^4 (Bca^3 + 3Abca^2) - d^3 (Ca^3c^2 + 3Ba^2bc^2 + 3Aab^2c^2) + d^2 (3Ca^2bc^2 + 3Bab^2c^2) + d (3Cab^2c^2 + 3Aab^2c^2) + 3Ab^3c^2)}{f (c^2d^4 + d^6)}$$

$$- \frac{\ln(\tan(e + fx) + 1i) (Aa^3 + Ab^31i - Ba^31i + Bb^3 - Ca^3 - Cb^31i - 3Aab^2 - Aa^2b3i + Bab^2)}{2f (d + c1i)}$$

$$- \frac{\ln(\tan(e + fx) - 1i) (Ab^3 - Ba^3 - Cb^3 - 3Aa^2b + 3Bab^2 + 3Ca^2b + Aa^31i + Bb^31i - Ca^31i)}{2f (c + d1i)}$$

$$+ \frac{Cb^3 \tan(e + fx)^3}{3df}$$

input `int(((a + b*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x)),x)`

output `(tan(e + f*x)^2*((B*b^3 + 3*C*a*b^2)/(2*d) - (C*b^3*c)/(2*d^2)))/f - (tan(e + f*x)*((c*((B*b^3 + 3*C*a*b^2)/d - (C*b^3*c)/d^2))/d - (A*b^3 + 3*B*a*b^2 + 3*C*a^2*b)/d + (C*b^3)/d))/f - (log(c + d*tan(e + f*x))*(d^4*(B*a^3*c + 3*A*a^2*b*c) - d^3*(C*a^3*c^2 + 3*A*a*b^2*c^2 + 3*B*a^2*b*c^2) + d^2*(A*b^3*c^3 + 3*B*a*b^2*c^3 + 3*C*a^2*b*c^3) - d*(B*b^3*c^4 + 3*C*a*b^2*c^4) - A*a^3*d^5 + C*b^3*c^5))/(f*(d^6 + c^2*d^4)) - (log(tan(e + f*x) + 1i)*(A*a^3 + A*b^3*1i - B*a^3*1i + B*b^3 - C*a^3 - C*b^3*1i - 3*A*a*b^2 - A*a^2*b*3i + B*a*b^2*3i - 3*B*a^2*b + 3*C*a*b^2 + C*a^2*b*3i))/(2*f*(c*1i + d)) - (log(tan(e + f*x) - 1i)*(A*a^3*1i + A*b^3 - B*a^3 + B*b^3*1i - C*a^3*1i - C*b^3 - A*a*b^2*3i - 3*A*a^2*b + 3*B*a*b^2 - B*a^2*b*3i + C*a*b^2*3i + 3*C*a^2*b))/(2*f*(c + d*1i)) + (C*b^3*tan(e + f*x)^3)/(3*d*f)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 819, normalized size of antiderivative = 2.43

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx = \text{Too large to display}$$

input `int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)`

output

```
( - 3*log(tan(e + f*x)**2 + 1)*a**4*d**5 + 12*log(tan(e + f*x)**2 + 1)*a**
3*b*c*d**4 + 3*log(tan(e + f*x)**2 + 1)*a**3*c*d**5 + 18*log(tan(e + f*x)*
**2 + 1)*a**2*b**2*d**5 - 9*log(tan(e + f*x)**2 + 1)*a**2*b*c**2*d**4 - 12*
log(tan(e + f*x)**2 + 1)*a*b**3*c*d**4 - 9*log(tan(e + f*x)**2 + 1)*a*b**2
*c*d**5 - 3*log(tan(e + f*x)**2 + 1)*b**4*d**5 + 3*log(tan(e + f*x)**2 + 1
)*b**3*c**2*d**4 + 6*log(tan(e + f*x)*d + c)*a**4*d**5 - 24*log(tan(e + f*
x)*d + c)*a**3*b*c*d**4 + 6*log(tan(e + f*x)*d + c)*a**3*c**3*d**3 + 36*lo
g(tan(e + f*x)*d + c)*a**2*b**2*c**2*d**3 - 18*log(tan(e + f*x)*d + c)*a**
2*b*c**4*d**2 - 24*log(tan(e + f*x)*d + c)*a*b**3*c**3*d**2 + 18*log(tan(e
+ f*x)*d + c)*a*b**2*c**5*d + 6*log(tan(e + f*x)*d + c)*b**4*c**4*d - 6*1
og(tan(e + f*x)*d + c)*b**3*c**6 + 2*tan(e + f*x)**3*b**3*c**3*d**3 + 2*ta
n(e + f*x)**3*b**3*c*d**5 + 9*tan(e + f*x)**2*a*b**2*c**3*d**3 + 9*tan(e +
f*x)**2*a*b**2*c*d**5 + 3*tan(e + f*x)**2*b**4*c**2*d**3 + 3*tan(e + f*x)
**2*b**4*d**5 - 3*tan(e + f*x)**2*b**3*c**4*d**2 - 3*tan(e + f*x)**2*b**3*
c**2*d**4 + 18*tan(e + f*x)*a**2*b*c**3*d**3 + 18*tan(e + f*x)*a**2*b*c*d*
**5 + 24*tan(e + f*x)*a*b**3*c**2*d**3 + 24*tan(e + f*x)*a*b**3*d**5 - 18*t
an(e + f*x)*a*b**2*c**4*d**2 - 18*tan(e + f*x)*a*b**2*c**2*d**4 - 6*tan(e
+ f*x)*b**4*c**3*d**2 - 6*tan(e + f*x)*b**4*c*d**4 + 6*tan(e + f*x)*b**3*c
**5*d - 6*tan(e + f*x)*b**3*c*d**5 + 6*a**4*c*d**4*f*x + 24*a**3*b*d**5*f*
x - 6*a**3*c**2*d**4*f*x - 36*a**2*b**2*c*d**4*f*x - 18*a**2*b*c*d**5*f...
```

3.71
$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

Optimal result	791
Mathematica [C] (verified)	792
Rubi [A] (verified)	792
Maple [A] (verified)	797
Fricas [A] (verification not implemented)	797
Sympy [C] (verification not implemented)	798
Maxima [A] (verification not implemented)	799
Giac [A] (verification not implemented)	800
Mupad [B] (verification not implemented)	801
Reduce [B] (verification not implemented)	802

Optimal result

Integrand size = 45, antiderivative size = 236

$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

$$= \frac{(a^2(Ac - cC + Bd) - b^2(Ac - cC + Bd) - 2ab(Bc - (A - C)d)) x}{c^2 + d^2}$$

$$- \frac{(2ab(Ac - cC + Bd) + a^2(Bc - (A - C)d) - b^2(Bc - (A - C)d)) \log(\cos(e+fx))}{(c^2 + d^2) f}$$

$$+ \frac{(bc - ad)^2 (c^2 C - Bcd + Ad^2) \log(c + d \tan(e+fx))}{d^3 (c^2 + d^2) f}$$

$$- \frac{b(bcC - bBd - aCd) \tan(e+fx)}{d^2 f} + \frac{C(a + b \tan(e+fx))^2}{2df}$$

output

```
(a^2*(A*c+B*d-C*c)-b^2*(A*c+B*d-C*c)-2*a*b*(B*c-(A-C)*d))*x/(c^2+d^2)-(2*a
*b*(A*c+B*d-C*c)+a^2*(B*c-(A-C)*d)-b^2*(B*c-(A-C)*d))*ln(cos(f*x+e))/(c^2+
d^2)/f+(-a*d+b*c)^2*(A*d^2-B*c*d+C*c^2)*ln(c+d*tan(f*x+e))/d^3/(c^2+d^2)/f
-b*(-B*b*d-C*a*d+C*b*c)*tan(f*x+e)/d^2/f+1/2*C*(a+b*tan(f*x+e))^2/d/f
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.99 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.81

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{\frac{(a+ib)^2(-iA+B+iC)d \log(i-\tan(e+fx))}{c+id} + \frac{(a-ib)^2(iA+B-iC)d \log(i+\tan(e+fx))}{c-id} + \frac{2(bc-ad)^2(c^2C-Bcd+Ad^2) \log(c+d \tan(e+fx))}{d^2(c^2+d^2)}}{2df}$$

input

```
Integrate[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x]),x]
```

output

```
((a + I*b)^2*((-I)*A + B + I*C)*d*Log[I - Tan[e + f*x]]/(c + I*d) + ((a
- I*b)^2*(I*A + B - I*C)*d*Log[I + Tan[e + f*x]]/(c - I*d) + (2*(b*c - a*
d)^2*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]]/(d^2*(c^2 + d^2)) +
(2*b*(-(b*c*C) + b*B*d + a*C*d)*Tan[e + f*x])/d + C*(a + b*Tan[e + f*x])^2
)/(2*d*f)
```

Rubi [A] (verified)

Time = 2.38 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4130, 27, 3042, 4120, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{c + d \tan(e + fx)} dx$$

$$\downarrow 4130$$

$$\frac{C(a + b \tan(e + fx))^2}{2df} - \frac{d^2(a^2(Bc - d(A - C)) + 2ab(Ac + Bd - cC) - b^2(Bc - d(A - C))) \int \tan(e + fx) dx}{c^2 + d^2} - \frac{(bc - ad)^2(A d^2 - Bcd + c^2 C) \int \frac{\tan^2(e + fx) + 1}{c + d \tan(e + fx)} dx}{d(c^2 + d^2)} - \frac{d^2 x(a^2(Ac + Bd - cC) - 2ab(Bc - d(A - C)) + b^2(Bc - d(A - C)))}{c^2 + d^2}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^2}{2df} - \frac{d^2(a^2(Bc - d(A - C)) + 2ab(Ac + Bd - cC) - b^2(Bc - d(A - C))) \int \tan(e + fx) dx}{c^2 + d^2} - \frac{(bc - ad)^2(A d^2 - Bcd + c^2 C) \int \frac{\tan(e + fx)^2 + 1}{c + d \tan(e + fx)} dx}{d(c^2 + d^2)} - \frac{d^2 x(a^2(Ac + Bd - cC) - 2ab(Bc - d(A - C)) + b^2(Bc - d(A - C)))}{c^2 + d^2}$$

↓ 3956

$$\frac{C(a + b \tan(e + fx))^2}{2df} - \frac{(bc - ad)^2(A d^2 - Bcd + c^2 C) \int \frac{\tan(e + fx)^2 + 1}{c + d \tan(e + fx)} dx}{c^2 + d^2} + \frac{d^2 \log(\cos(e + fx))(a^2(Bc - d(A - C)) + 2ab(Ac + Bd - cC) - b^2(Bc - d(A - C)))}{f(c^2 + d^2)} - \frac{d^2 x(a^2(Ac + Bd - cC) - 2ab(Bc - d(A - C)) + b^2(Bc - d(A - C)))}{c^2 + d^2}$$

↓ 4100

$$\frac{C(a + b \tan(e + fx))^2}{2df} - \frac{(bc - ad)^2(A d^2 - Bcd + c^2 C) \int \frac{1}{c + d \tan(e + fx)} d(d \tan(e + fx))}{df(c^2 + d^2)} + \frac{d^2 \log(\cos(e + fx))(a^2(Bc - d(A - C)) + 2ab(Ac + Bd - cC) - b^2(Bc - d(A - C)))}{f(c^2 + d^2)} - \frac{d^2 x(a^2(Ac + Bd - cC) - 2ab(Bc - d(A - C)) + b^2(Bc - d(A - C)))}{c^2 + d^2}$$

↓ 16

$$\frac{C(a + b \tan(e + fx))^2}{2df} - \frac{d^2 \log(\cos(e + fx))(a^2(Bc - d(A - C)) + 2ab(Ac + Bd - cC) - b^2(Bc - d(A - C)))}{f(c^2 + d^2)} - \frac{d^2 x(a^2(Ac + Bd - cC) - 2ab(Bc - d(A - C)) - b^2(Ac + Bd - cC))}{c^2 + d^2} - \frac{(bc - ad)^2(A d^2 - Bcd + c^2 C)}{d(c^2 + d^2)}$$

input

```
Int[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]),x]
```

output
$$\frac{(C*(a + b*\text{Tan}[e + f*x])^2)/(2*d*f) - (((d^2*(a^2*(A*c - c*C + B*d) - b^2*(A*c - c*C + B*d) - 2*a*b*(B*c - (A - C)*d))*x)/(c^2 + d^2)) + (d^2*(2*a*b*(A*c - c*C + B*d) + a^2*(B*c - (A - C)*d) - b^2*(B*c - (A - C)*d))*\text{Log}[\text{Cos}[e + f*x]]/(c^2 + d^2)*f - ((b*c - a*d)^2*(c^2*C - B*c*d + A*d^2))*\text{Log}[c + d*\text{Tan}[e + f*x]]/(d*(c^2 + d^2)*f))/d + (b*(b*c*C - b*B*d - a*C*d)*\text{Tan}[e + f*x])/(d*f))/d$$

Defintions of rubi rules used

rule 16
$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$$

rule 25
$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$$

rule 27
$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$$

rule 3042
$$\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956
$$\text{Int}[\tan[(c_)+(d_)*(x)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] \text{ ; FreeQ}[\{c, d\}, x]$$

rule 4100
$$\text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x)]^(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x)]^2), x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] \text{ ; FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$$

rule 4109

```

Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2
)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]

```

rule 4120

```

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_
)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]

```

rule 4130

```

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.34

method	result
derivativedivides	$\frac{b\left(\frac{\tan(fx+e)^2 C b d}{2} + B \tan(fx+e) b d + 2 d \tan(fx+e) C a - \tan(fx+e) C b c\right)}{d^2} + \frac{(-A a^2 d + 2 A a b c + A b^2 d + B a^2 c + 2 B a b d - B b^2 c + C a^2 d)}{2}$
default	$\frac{b\left(\frac{\tan(fx+e)^2 C b d}{2} + B \tan(fx+e) b d + 2 d \tan(fx+e) C a - \tan(fx+e) C b c\right)}{d^2} + \frac{(-A a^2 d + 2 A a b c + A b^2 d + B a^2 c + 2 B a b d - B b^2 c + C a^2 d)}{2}$
norman	$\frac{(A a^2 c + 2 A a b d - A b^2 c + B a^2 d - 2 B a b c - B b^2 d - C a^2 c - 2 C a b d + C b^2 c) x}{c^2 + d^2} + \frac{b(B b d + 2 C a d - C b c) \tan(fx+e)}{d^2 f} + \frac{C b^2}{2}$
parallelrisc	$-\frac{A \ln(1 + \tan(fx+e)^2) a^2 d^4 - A \ln(1 + \tan(fx+e)^2) b^2 d^4 - 2 A \ln(c + d \tan(fx+e)) a^2 d^4 - C \ln(1 + \tan(fx+e)^2) a^2 d^4 + \dots}{c^2 + d^2}$
risc	Expression too large to display

```
input int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x,
method=_RETURNVERBOSE)
```

```
output 1/f*(b/d^2*(1/2*tan(f*x+e)^2*C*b*d+B*tan(f*x+e)*b*d+2*d*tan(f*x+e)*C*a-tan
(f*x+e)*C*b*c)+1/(c^2+d^2)*(1/2*(-A*a^2*d+2*A*a*b*c+A*b^2*d+B*a^2*c+2*B*a*
b*d-B*b^2*c+C*a^2*d-2*C*a*b*c-C*b^2*d)*ln(1+tan(f*x+e)^2)+(A*a^2*c+2*A*a*b
*d-A*b^2*c+B*a^2*d-2*B*a*b*c-B*b^2*d-C*a^2*c-2*C*a*b*d+C*b^2*c)*arctan(tan
(f*x+e)))+1/d^3*(A*a^2*d^4-2*A*a*b*c*d^3+A*b^2*c^2*d^2-B*a^2*c*d^3+2*B*a*b
*c^2*d^2-B*b^2*c^3*d+C*a^2*c^2*d^2-2*C*a*b*c^3*d+C*b^2*c^4)/(c^2+d^2)*ln(c
+d*tan(f*x+e))
```

Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.65

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{2(((A - C)a^2 - 2 Bab - (A - C)b^2)cd^3 + (Ba^2 + 2(A - C)ab - Bb^2)d^4)fx + (Cb^2c^2d^2 + Cb^2d^4) \tan(e + fx)}{c^2 + d^2}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="fricas")`

output `1/2*(2*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^3 + (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^4)*f*x + (C*b^2*c^2*d^2 + C*b^2*d^4)*tan(f*x + e)^2 + (C*b^2*c^4 + A*a^2*d^4 - (2*C*a*b + B*b^2)*c^3*d + (C*a^2 + 2*B*a*b + A*b^2)*c^2*d^2 - (B*a^2 + 2*A*a*b)*c*d^3)*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - (C*b^2*c^4 - (2*C*a*b + B*b^2)*c^3*d + (C*a^2 + 2*B*a*b + A*b^2)*c^2*d^2 - (2*C*a*b + B*b^2)*c*d^3 + (C*a^2 + 2*B*a*b + (A - C)*b^2)*d^4)*log(1/(tan(f*x + e)^2 + 1)) - 2*(C*b^2*c^3*d + C*b^2*c*d^3 - (2*C*a*b + B*b^2)*c^2*d^2 - (2*C*a*b + B*b^2)*d^4)*tan(f*x + e)/((c^2*d^3 + d^5)*f)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.09 (sec) , antiderivative size = 4444, normalized size of antiderivative = 18.83

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)),x)`

output

```
Piecewise((zoo*x*(a + b*tan(e))**2*(A + B*tan(e) + C*tan(e)**2)/tan(e), Eq
(c, 0) & Eq(d, 0) & Eq(f, 0)), ((A**2*x + A*b*log(tan(e + f*x)**2 + 1)
/f - A**2*x + A**2*tan(e + f*x)/f + B**2*log(tan(e + f*x)**2 + 1)/(2
*f) - 2*B*a*b*x + 2*B*a*b*tan(e + f*x)/f - B**2*log(tan(e + f*x)**2 + 1)
/(2*f) + B**2*tan(e + f*x)**2/(2*f) - C**2*x + C**2*tan(e + f*x)/f -
C*a*b*log(tan(e + f*x)**2 + 1)/f + C*a*b*tan(e + f*x)**2/f + C**2*x + C
**2*tan(e + f*x)**3/(3*f) - C**2*tan(e + f*x)/f)/c, Eq(d, 0)), (I*A*a
**2*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) + A**2*f*x/(2*d*f*tan
(e + f*x) - 2*I*d*f) + I*A**2/(2*d*f*tan(e + f*x) - 2*I*d*f) + 2*A*a*b*f
*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) - 2*I*A*a*b*f*x/(2*d*f*tan(
e + f*x) - 2*I*d*f) - 2*A*a*b/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*A*b**2*f*x
*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) + A**2*f*x/(2*d*f*tan(e +
f*x) - 2*I*d*f) + A**2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*d*f*tan(
e + f*x) - 2*I*d*f) - I*A*b**2*log(tan(e + f*x)**2 + 1)/(2*d*f*tan(e + f*x)
) - 2*I*d*f) - I*A*b**2/(2*d*f*tan(e + f*x) - 2*I*d*f) + B**2*f*x*tan(e
+ f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*B**2*f*x/(2*d*f*tan(e + f*x) -
2*I*d*f) - B**2/(2*d*f*tan(e + f*x) - 2*I*d*f) + 2*I*B*a*b*f*x*tan(e +
f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) + 2*B*a*b*f*x/(2*d*f*tan(e + f*x) - 2*
I*d*f) + 2*B*a*b*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*d*f*tan(e + f*x)
- 2*I*d*f) - 2*I*B*a*b*log(tan(e + f*x)**2 + 1)/(2*d*f*tan(e + f*x) - ...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.25

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{2(((A-C)a^2 - 2Bab - (A-C)b^2)c + (Ba^2 + 2(A-C)ab - Bb^2)d)(fx+e)}{c^2+d^2} + \frac{2(Cb^2c^4 + Aa^2d^4 - (2Cab + Bb^2)c^3d + (Ca^2 + 2Bab + Ab^2)c^2d^2 - (B$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e)),x, algorithm="maxima")
```

output

```

1/2*(2*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c + (B*a^2 + 2*(A - C)*a*b -
B*b^2)*d)*(f*x + e)/(c^2 + d^2) + 2*(C*b^2*c^4 + A*a^2*d^4 - (2*C*a*b + B
*b^2)*c^3*d + (C*a^2 + 2*B*a*b + A*b^2)*c^2*d^2 - (B*a^2 + 2*A*a*b)*c*d^3)
*log(d*tan(f*x + e) + c)/(c^2*d^3 + d^5) + ((B*a^2 + 2*(A - C)*a*b - B*b^2
)*c - ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d)*log(tan(f*x + e)^2 + 1)/(c^
2 + d^2) + (C*b^2*d*tan(f*x + e)^2 - 2*(C*b^2*c - (2*C*a*b + B*b^2)*d)*tan
(f*x + e))/d^2)/f

```

Giac [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.46

$$\begin{aligned}
& \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx \\
&= \frac{(Aa^2c - Ca^2c - 2Babc - Ab^2c + Cb^2c + Ba^2d + 2Aabd - 2Cabd - Bb^2d)(fx + e)}{c^2f + d^2f} \\
&+ \frac{(Ba^2c + 2Aabc - 2Cabc - Bb^2c - Aa^2d + Ca^2d + 2Babd + Ab^2d - Cb^2d) \log(\tan(fx + e)^2 + 1)}{2(c^2f + d^2f)} \\
&+ \frac{(Cb^2c^4 - 2Cabc^3d - Bb^2c^3d + Ca^2c^2d^2 + 2Babc^2d^2 + Ab^2c^2d^2 - Ba^2cd^3 - 2Aabcd^3 + Aa^2d^4) \log(|c + d \tan(fx + e)|)}{c^2d^3f + d^5f} \\
&+ \frac{Cb^2df \tan(fx + e)^2 - 2Cb^2cf \tan(fx + e) + 4Cabdf \tan(fx + e) + 2Bb^2df \tan(fx + e)}{2d^2f^2}
\end{aligned}$$

input

```

integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e)),x, algorithm="giac")

```

output

```

(A*a^2*c - C*a^2*c - 2*B*a*b*c - A*b^2*c + C*b^2*c + B*a^2*d + 2*A*a*b*d -
2*C*a*b*d - B*b^2*d)*(f*x + e)/(c^2*f + d^2*f) + 1/2*(B*a^2*c + 2*A*a*b*c
- 2*C*a*b*c - B*b^2*c - A*a^2*d + C*a^2*d + 2*B*a*b*d + A*b^2*d - C*b^2*d
)*log(tan(f*x + e)^2 + 1)/(c^2*f + d^2*f) + (C*b^2*c^4 - 2*C*a*b*c^3*d - B
*b^2*c^3*d + C*a^2*c^2*d^2 + 2*B*a*b*c^2*d^2 + A*b^2*c^2*d^2 - B*a^2*c*d^3
- 2*A*a*b*c*d^3 + A*a^2*d^4)*log(abs(d*tan(f*x + e) + c))/(c^2*d^3*f + d^
5*f) + 1/2*(C*b^2*d*f*tan(f*x + e)^2 - 2*C*b^2*c*f*tan(f*x + e) + 4*C*a*b*
d*f*tan(f*x + e) + 2*B*b^2*d*f*tan(f*x + e))/(d^2*f^2)

```

Mupad [B] (verification not implemented)

Time = 7.93 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.38

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{\tan(e + fx) \left(\frac{Bb^2 + 2Cab}{d} - \frac{Cb^2c}{d^2} \right)}{f}$$

$$+ \frac{\ln(c + d \tan(e + fx)) (d^2 (Ca^2c^2 + 2Babc^2 + Ab^2c^2) - d(Bb^2c^3 + 2Cabc^3) - d^3(Bca^2 + 2Abc^2))}{f(c^2d^3 + d^5)}$$

$$+ \frac{\ln(\tan(e + fx) + 1i) (Ab^2 - Aa^2 + Ba^21i - Bb^21i + Ca^2 - Cb^2 + Aab2i + 2Bab - Cab2i)}{2f(d + c1i)}$$

$$+ \frac{\ln(\tan(e + fx) - 1i) (Ba^2 - Bb^2 + 2Aab - 2Cab - Aa^21i + Ab^21i + Ca^21i - Cb^21i + Bab2i)}{2f(c + d1i)}$$

$$+ \frac{Cb^2 \tan(e + fx)^2}{2df}$$

input

```
int(((a + b*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x)),x)
```

output

```
(tan(e + f*x)*((B*b^2 + 2*C*a*b)/d - (C*b^2*c)/d^2))/f + (log(c + d*tan(e + f*x))*(d^2*(A*b^2*c^2 + C*a^2*c^2 + 2*B*a*b*c^2) - d*(B*b^2*c^3 + 2*C*a*b*c^3) - d^3*(B*a^2*c + 2*A*a*b*c) + A*a^2*d^4 + C*b^2*c^4))/(f*(d^5 + c^2*d^3)) + (log(tan(e + f*x) + 1i)*(A*b^2 - A*a^2 + B*a^2*1i - B*b^2*1i + C*a^2 - C*b^2 + A*a*b*2i + 2*B*a*b - C*a*b*2i))/(2*f*(c*1i + d)) + (log(tan(e + f*x) - 1i)*(A*b^2*1i - A*a^2*1i + B*a^2 - B*b^2 + C*a^2*1i - C*b^2*1i + 2*A*a*b + B*a*b*2i - 2*C*a*b))/(2*f*(c + d*1i)) + (C*b^2*tan(e + f*x)^2)/(2*d*f)
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 508, normalized size of antiderivative = 2.15

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{-\log(\tan(fx + e)^2 + 1) a^3 d^4 + \log(\tan(fx + e)^2 + 1) a^2 c d^4 - \log(\tan(fx + e)^2 + 1) b^3 c d^3 - \log(\tan$$

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)
```

output

```
( - log(tan(e + f*x)**2 + 1)*a**3*d**4 + 3*log(tan(e + f*x)**2 + 1)*a**2*b
*c*d**3 + log(tan(e + f*x)**2 + 1)*a**2*c*d**4 + 3*log(tan(e + f*x)**2 + 1
)*a*b**2*d**4 - 2*log(tan(e + f*x)**2 + 1)*a*b*c**2*d**3 - log(tan(e + f*x
)**2 + 1)*b**3*c*d**3 - log(tan(e + f*x)**2 + 1)*b**2*c*d**4 + 2*log(tan(e
+ f*x)*d + c)*a**3*d**4 - 6*log(tan(e + f*x)*d + c)*a**2*b*c*d**3 + 2*log
(tan(e + f*x)*d + c)*a**2*c**3*d**2 + 6*log(tan(e + f*x)*d + c)*a*b**2*c**
2*d**2 - 4*log(tan(e + f*x)*d + c)*a*b*c**4*d - 2*log(tan(e + f*x)*d + c)*
b**3*c**3*d + 2*log(tan(e + f*x)*d + c)*b**2*c**5 + tan(e + f*x)**2*b**2*c
**3*d**2 + tan(e + f*x)**2*b**2*c*d**4 + 4*tan(e + f*x)*a*b*c**3*d**2 + 4*
tan(e + f*x)*a*b*c*d**4 + 2*tan(e + f*x)*b**3*c**2*d**2 + 2*tan(e + f*x)*b
**3*d**4 - 2*tan(e + f*x)*b**2*c**4*d - 2*tan(e + f*x)*b**2*c**2*d**3 + 2*
a**3*c*d**3*f*x + 6*a**2*b*d**4*f*x - 2*a**2*c**2*d**3*f*x - 6*a*b**2*c*d
**3*f*x - 4*a*b*c*d**4*f*x - 2*b**3*d**4*f*x + 2*b**2*c**2*d**3*f*x)/(2*d**
3*f*(c**2 + d**2))
```

3.72
$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

Optimal result	803
Mathematica [C] (verified)	804
Rubi [A] (verified)	804
Maple [A] (verified)	807
Fricas [A] (verification not implemented)	808
Sympy [C] (verification not implemented)	808
Maxima [A] (verification not implemented)	809
Giac [A] (verification not implemented)	810
Mupad [B] (verification not implemented)	810
Reduce [B] (verification not implemented)	811

Optimal result

Integrand size = 43, antiderivative size = 156

$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

$$= \frac{(a(Ac - cC + Bd) - b(Bc - (A - C)d))x}{c^2 + d^2}$$

$$- \frac{(Abc + aBc - bcC - aAd + bBd + aCd) \log(\cos(e+fx))}{(c^2 + d^2) f}$$

$$- \frac{(bc - ad)(c^2C - Bcd + Ad^2) \log(c+d \tan(e+fx))}{d^2(c^2 + d^2) f} + \frac{bC \tan(e+fx)}{df}$$

output

```
(a*(A*c+B*d-C*c)-b*(B*c-(A-C)*d))*x/(c^2+d^2)-(-A*a*d+A*b*c+B*a*c+B*b*d+C*
a*d-C*b*c)*ln(cos(f*x+e))/(c^2+d^2)/f-(-a*d+b*c)*(A*d^2-B*c*d+C*c^2)*ln(c+
d*tan(f*x+e))/d^2/(c^2+d^2)/f+b*C*tan(f*x+e)/d/f
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.72 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{\frac{(-ia+b)(A+iB-C) \log(i-\tan(e+fx))}{c+id} + \frac{(ia+b)(A-iB-C) \log(i+\tan(e+fx))}{c-id} + \frac{2(-bc+ad)(c^2C-Bcd+Ad^2) \log(c+d \tan(e+fx))}{d^2(c^2+d^2)} + 2b}{2f}$$

input

```
Integrate[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]),x]
```

output

```
((((-I)*a + b)*(A + I*B - C)*Log[I - Tan[e + f*x]])/(c + I*d) + ((I*a + b)*(A - I*B - C)*Log[I + Tan[e + f*x]])/(c - I*d) + (2*(-(b*c) + a*d)*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]])/(d^2*(c^2 + d^2)) + (2*b*C*Tan[e + f*x])/d)/(2*f)
```

Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4120, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan(e + fx)^2)}{c + d \tan(e + fx)} dx$$

↓ 4120

$$\frac{bC \tan(e + fx)}{df} - \frac{\int \frac{(bcC - adC - bBd) \tan^2(e + fx) - (Ab - Cb + aB)d \tan(e + fx) + bcC - aAd}{c + d \tan(e + fx)} dx}{d}$$

↓ 3042

$$\frac{bC \tan(e + fx)}{df} - \frac{\int \frac{(bcC - adC - bBd) \tan(e + fx)^2 - (Ab - Cb + aB)d \tan(e + fx) + bcC - aAd}{c + d \tan(e + fx)} dx}{d}$$

↓ 4109

$$\frac{bC \tan(e + fx)}{df} - \frac{(bc - ad)(Ad^2 - Bcd + c^2C) \int \frac{\tan^2(e + fx) + 1}{c + d \tan(e + fx)} dx - \frac{d(-aAd + aBc + aCd + Abc + bBd - bcC) \int \tan(e + fx) dx}{c^2 + d^2} - \frac{dx(a(Ac + Bd - cC) - b(Bc - d(A - C)))}{c^2 + d^2}}{d}$$

↓ 3042

$$\frac{bC \tan(e + fx)}{df} - \frac{-\frac{d(-aAd + aBc + aCd + Abc + bBd - bcC) \int \tan(e + fx) dx}{c^2 + d^2} + \frac{(bc - ad)(Ad^2 - Bcd + c^2C) \int \frac{\tan(e + fx)^2 + 1}{c + d \tan(e + fx)} dx}{c^2 + d^2} - \frac{dx(a(Ac + Bd - cC) - b(Bc - d(A - C)))}{c^2 + d^2}}{d}$$

↓ 3956

$$\frac{bC \tan(e + fx)}{df} - \frac{(bc - ad)(Ad^2 - Bcd + c^2C) \int \frac{\tan(e + fx)^2 + 1}{c + d \tan(e + fx)} dx + \frac{d \log(\cos(e + fx))(-aAd + aBc + aCd + Abc + bBd - bcC)}{f(c^2 + d^2)} - \frac{dx(a(Ac + Bd - cC) - b(Bc - d(A - C)))}{c^2 + d^2}}{d}$$

↓ 4100

$$\frac{bC \tan(e + fx)}{df} - \frac{(bc - ad)(Ad^2 - Bcd + c^2C) \int \frac{1}{c + d \tan(e + fx)} d(d \tan(e + fx)) + \frac{d \log(\cos(e + fx))(-aAd + aBc + aCd + Abc + bBd - bcC)}{f(c^2 + d^2)} - \frac{dx(a(Ac + Bd - cC) - b(Bc - d(A - C)))}{c^2 + d^2}}{d}$$

↓ 16

$$\frac{bC \tan(e + fx)}{df} - \frac{(bc - ad)(Ad^2 - Bcd + c^2C) \log(c + d \tan(e + fx)) + \frac{d \log(\cos(e + fx))(-aAd + aBc + aCd + Abc + bBd - bcC)}{f(c^2 + d^2)} - \frac{dx(a(Ac + Bd - cC) - b(Bc - d(A - C)))}{c^2 + d^2}}{d}$$

input $\text{Int}[(a + b \tan[e + f x]) (A + B \tan[e + f x] + C \tan[e + f x]^2) / (c + d \tan[e + f x]), x]$

output $-\left(-\left(\left(d(a(Ac - cC + Bd) - b(Bc - (A - C)d))\right)x\right)/(c^2 + d^2)\right) + (d(A^2b^2c + a^2B^2c - b^2c^2C - a^2Ad + b^2Bd + a^2Cd) \log[\cos[e + fx]]) / ((c^2 + d^2)f) + ((b^2c - a^2d)(c^2C - B^2cd + A^2d^2) \log[c + d \tan[e + fx]]) / (d(c^2 + d^2)f) + (b^2C \tan[e + fx]) / (df)$

Definitions of rubi rules used

rule 16 $\text{Int}[(c_./((a_.) + (b_.)x)), x_Symbol] \rightarrow \text{Simp}[c(\log[\text{RemoveContent}[a + bx, x]/b]), x] /; \text{FreeQ}\{a, b, c\}, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)x], x_Symbol] \rightarrow \text{Simp}[-\log[\text{RemoveContent}[\cos[c + dx], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

rule 4100 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m, x_Symbol] \rightarrow \text{Simp}[A/(b^m f) \text{Subst}[\text{Int}[(a + x)^m, x], x, b \tan[e + fx]], x] /; \text{FreeQ}\{a, b, e, f, A, C, m\}, x] \&\& \text{EqQ}[A, C]$

rule 4109 $\text{Int}[(A_.) + (B_.)\tan[(e_.) + (f_.)x] + (C_.)\tan[(e_.) + (f_.)x]^2 / ((a_.) + (b_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[(a^2A + b^2B - a^2C)(x/(a^2 + b^2)), x] + (\text{Simp}[(A^2b^2 - a^2b^2B + a^2^2C)/(a^2 + b^2) \text{Int}[(1 + \tan[e + fx]^2)/(a + b \tan[e + fx]), x], x] - \text{Simp}[(A^2b - a^2B - b^2C)/(a^2 + b^2) \text{Int}[\tan[e + fx], x], x]) /; \text{FreeQ}\{a, b, e, f, A, B, C\}, x] \&\& \text{NeQ}[A^2b^2 - a^2b^2B + a^2^2C, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A^2b - a^2B - b^2C, 0]$

rule 4120

```
Int[((a_) + (b_) * tan[(e_) + (f_) * (x_)]) * ((c_) + (d_) * tan[(e_) + (f_) * (x_)])^(n_) * ((A_) + (B_) * tan[(e_) + (f_) * (x_)]) + (C_) * tan[(e_) + (f_) * (x_)])^2, x_Symbol] :> Simp[b * C * Tan[e + f * x] * ((c + d * Tan[e + f * x])^(n + 1) / (d * f * (n + 2))), x] - Simp[1 / (d * (n + 2)) Int[(c + d * Tan[e + f * x])^n * Simp[b * c * C - a * A * d * (n + 2) - (A * b + a * B - b * C) * d * (n + 2) * Tan[e + f * x] - (a * C * d * (n + 2) - b * (c * C - B * d * (n + 2))) * Tan[e + f * x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b * c - a * d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{\tan(fx+e)Cb}{d} + \frac{(-Aad+Abc+Bac+Bbd+Cad-Cbc) \ln(1+\tan(fx+e)^2)}{2} + \frac{(Aac+Abd+Bad-Bbc-Cac-Cbd) \arctan(\tan(fx+e))}{c^2+d^2} + \frac{f}{c^2+d^2}$
default	$\frac{\tan(fx+e)Cb}{d} + \frac{(-Aad+Abc+Bac+Bbd+Cad-Cbc) \ln(1+\tan(fx+e)^2)}{2} + \frac{(Aac+Abd+Bad-Bbc-Cac-Cbd) \arctan(\tan(fx+e))}{c^2+d^2} + \frac{f}{c^2+d^2}$
norman	$\frac{(Aac+Abd+Bad-Bbc-Cac-Cbd)x}{c^2+d^2} + \frac{bC \tan(fx+e)}{df} + \frac{(Aa d^3 - Abc d^2 - Bac d^2 + Bb c^2 d + Ca c^2 d - Cb c^3) \ln(c+d)}{d^2 f (c^2+d^2)}$
parallelrisc	$-\frac{2Aac d^2 fx - 2Ab d^3 fx - 2Ba d^3 fx + 2Bbc d^2 fx + 2Cac d^2 fx + 2Cb d^3 fx + A \ln(1+\tan(fx+e)^2) a d^3 - A \ln(1+\tan(fx+e))}{c^2+d^2}$
risc	$-\frac{x A a}{i d - c} + \frac{x B b}{i d - c} + \frac{x C a}{i d - c} + \frac{i x A b}{i d - c} + \frac{i x B a}{i d - c} - \frac{i x C b}{i d - c} + \frac{2 i B b x}{d} + \frac{2 i C a x}{d} - \frac{\ln(e^{2 i (f x + e)} + 1) B b}{d f} - \frac{\ln(e^{2 i (f x + e)} + 1) C b}{d f}$

input

```
int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

output

```
1/f*(tan(f*x+e)*C*b/d+1/(c^2+d^2)*(1/2*(-A*a*d+A*b*c+B*a*c+B*b*d+C*a*d-C*b*c)*ln(1+tan(f*x+e)^2)+(A*a*c+A*b*d+B*a*d-B*b*c-C*a*c-C*b*d)*arctan(tan(f*x+e)))+1/d^2*(A*a*d^3-A*b*c*d^2-B*a*c*d^2+B*b*c^2*d+C*a*c^2*d-C*b*c^3)/(c^2+d^2)*ln(c+d*tan(f*x+e))
```

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.36

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{2(((A - C)a - Bb)cd^2 + (Ba + (A - C)b)d^3)fx - (Cbc^3 - Aad^3 - (Ca + Bb)c^2d + (Ba + Ab)cd^2) \log \dots}{\dots}$$

input

```
integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="fricas")
```

output

```
1/2*(2*(((A - C)*a - B*b)*c*d^2 + (B*a + (A - C)*b)*d^3)*f*x - (C*b*c^3 - A*a*d^3 - (C*a + B*b)*c^2*d + (B*a + A*b)*c*d^2)*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) + (C*b*c^3 + C*b*c*d^2 - (C*a + B*b)*c^2*d - (C*a + B*b)*d^3)*log(1/(tan(f*x + e)^2 + 1)) + 2*(C*b*c^2*d + C*b*d^3)*tan(f*x + e))/((c^2*d^2 + d^4)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.79 (sec) , antiderivative size = 2387, normalized size of antiderivative = 15.30

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)),x)
```

output

```
Piecewise((zoo*x*(a + b*tan(e))*(A + B*tan(e) + C*tan(e)**2)/tan(e), Eq(c,
0) & Eq(d, 0) & Eq(f, 0)), ((A*a*x + A*b*log(tan(e + f*x)**2 + 1)/(2*f) +
B*a*log(tan(e + f*x)**2 + 1)/(2*f) - B*b*x + B*b*tan(e + f*x)/f - C*a*x +
C*a*tan(e + f*x)/f - C*b*log(tan(e + f*x)**2 + 1)/(2*f) + C*b*tan(e + f*x)
)**2/(2*f))/c, Eq(d, 0)), (I*A*a*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*
I*d*f) + A*a*f*x/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*A*a/(2*d*f*tan(e + f*x)
) - 2*I*d*f) + A*b*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*A*b
*f*x/(2*d*f*tan(e + f*x) - 2*I*d*f) - A*b/(2*d*f*tan(e + f*x) - 2*I*d*f) +
B*a*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*B*a*f*x/(2*d*f*ta
n(e + f*x) - 2*I*d*f) - B*a/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*B*b*f*x*tan
(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) + B*b*f*x/(2*d*f*tan(e + f*x) - 2
*I*d*f) + B*b*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*d*f*tan(e + f*x) -
2*I*d*f) - I*B*b*log(tan(e + f*x)**2 + 1)/(2*d*f*tan(e + f*x) - 2*I*d*f) -
I*B*b/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*C*a*f*x*tan(e + f*x)/(2*d*f*tan(
e + f*x) - 2*I*d*f) + C*a*f*x/(2*d*f*tan(e + f*x) - 2*I*d*f) + C*a*log(tan
(e + f*x)**2 + 1)*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*C*a*log(
tan(e + f*x)**2 + 1)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*C*a/(2*d*f*tan(e +
f*x) - 2*I*d*f) - 3*C*b*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) +
3*I*C*b*f*x/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*C*b*log(tan(e + f*x)**2 +
1)*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) + C*b*log(tan(e + f*x)**...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.14

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{\frac{2Cb \tan(fx+e)}{d} + \frac{2(((A-C)a - Bb)c + (Ba + (A-C)b)d)(fx+e)}{c^2 + d^2} - \frac{2(Cbc^3 - Aad^3 - (Ca + Bb)c^2d + (Ba + Ab)cd^2) \log(d \tan(fx+e) + c)}{c^2d^2 + d^4}}{2f} + \dots$$

input

```
integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
),x, algorithm="maxima")
```

output

```
1/2*(2*C*b*tan(f*x + e)/d + 2*(((A - C)*a - B*b)*c + (B*a + (A - C)*b)*d)*
(f*x + e)/(c^2 + d^2) - 2*(C*b*c^3 - A*a*d^3 - (C*a + B*b)*c^2*d + (B*a +
A*b)*c*d^2)*log(d*tan(f*x + e) + c)/(c^2*d^2 + d^4) + ((B*a + (A - C)*b)*c
- ((A - C)*a - B*b)*d)*log(tan(f*x + e)^2 + 1)/(c^2 + d^2))/f
```

Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.22

$$\begin{aligned}
& \int \frac{(a + b \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx \\
&= \frac{(Aac - Cac - Bbc + Bad + Abd - Cbd)(fx + e)}{c^2 f + d^2 f} \\
&\quad + \frac{(Bac + Abc - Cbc - Aad + Cad + Bbd) \log(\tan(fx + e)^2 + 1)}{2(c^2 f + d^2 f)} \\
&\quad - \frac{(Cbc^3 - Cac^2 d - Bbc^2 d + Bacd^2 + Abcd^2 - Aad^3) \log(|d \tan(fx + e) + c|)}{c^2 d^2 f + d^4 f} \\
&\quad + \frac{Cb \tan(fx + e)}{df}
\end{aligned}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="giac")`

output `(A*a*c - C*a*c - B*b*c + B*a*d + A*b*d - C*b*d)*(f*x + e)/(c^2*f + d^2*f) + 1/2*(B*a*c + A*b*c - C*b*c - A*a*d + C*a*d + B*b*d)*log(tan(f*x + e)^2 + 1)/(c^2*f + d^2*f) - (C*b*c^3 - C*a*c^2*d - B*b*c^2*d + B*a*c*d^2 + A*b*c*d^2 - A*a*d^3)*log(abs(d*tan(f*x + e) + c))/(c^2*d^2*f + d^4*f) + C*b*tan(f*x + e)/(d*f)`

Mupad [B] (verification not implemented)

Time = 6.67 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.19

$$\begin{aligned}
& \int \frac{(a + b \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx \\
&= \frac{\ln(\tan(e + fx) - i) (Ab + Ba - Cb - Aali + Bbli + Cali)}{2f(c + dli)} \\
&\quad + \frac{\ln(\tan(e + fx) + li) (Bb + Abli + Bali - Aa + Ca - Cbli)}{2f(d + cli)} \\
&\quad - \frac{\ln(c + d \tan(e + fx)) (d^2 (Abc + Bac) - d(Bbc^2 + Cac^2) - Aad^3 + Cbc^3)}{f(c^2 d^2 + d^4)} \\
&\quad + \frac{Cb \tan(e + fx)}{df}
\end{aligned}$$

input `int(((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x)),x)`

output `(log(tan(e + f*x) - 1i)*(A*b - A*a*1i + B*a + B*b*1i + C*a*1i - C*b))/(2*f*(c + d*1i)) + (log(tan(e + f*x) + 1i)*(A*b*1i - A*a + B*a*1i + B*b + C*a - C*b*1i))/(2*f*(c*1i + d)) - (log(c + d*tan(e + f*x))*(d^2*(A*b*c + B*a*c) - d*(B*b*c^2 + C*a*c^2) - A*a*d^3 + C*b*c^3))/(f*(d^4 + c^2*d^2)) + (C*b*tan(e + f*x))/(d*f)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{-\log(\tan(fx + e)^2 + 1) a^2 d^3 + 2 \log(\tan(fx + e)^2 + 1) abc d^2 + \log(\tan(fx + e)^2 + 1) ac d^3 + \log(\tan(fx + e)^2 + 1) a^2 d^3 + 2 \log(\tan(fx + e)^2 + 1) abc d^2 + \log(\tan(fx + e)^2 + 1) ac d^3 + \log(\tan(fx + e)^2 + 1) a^2 d^3}{(c + d \tan(e + fx))^2}$$

input `int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)`

output `(- log(tan(e + f*x)**2 + 1)*a**2*d**3 + 2*log(tan(e + f*x)**2 + 1)*a*b*c*d**2 + log(tan(e + f*x)**2 + 1)*a*c*d**3 + log(tan(e + f*x)**2 + 1)*b**2*d**3 - log(tan(e + f*x)**2 + 1)*b*c**2*d**2 + 2*log(tan(e + f*x)*d + c)*a**2*d**3 - 4*log(tan(e + f*x)*d + c)*a*b*c*d**2 + 2*log(tan(e + f*x)*d + c)*a*c**3*d + 2*log(tan(e + f*x)*d + c)*b**2*c**2*d - 2*log(tan(e + f*x)*d + c)*b*c**4 + 2*tan(e + f*x)*b*c**3*d + 2*tan(e + f*x)*b*c*d**3 + 2*a**2*c*d**2*f*x + 4*a*b*d**3*f*x - 2*a*c**2*d**2*f*x - 2*b**2*c*d**2*f*x - 2*b*c*d**3*f*x)/(2*d**2*f*(c**2 + d**2))`

3.73 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{c+d \tan(e+fx)} dx$

Optimal result	812
Mathematica [C] (verified)	812
Rubi [A] (verified)	813
Maple [A] (verified)	815
Fricas [A] (verification not implemented)	816
Sympy [C] (verification not implemented)	816
Maxima [A] (verification not implemented)	817
Giac [A] (verification not implemented)	818
Mupad [B] (verification not implemented)	818
Reduce [B] (verification not implemented)	819

Optimal result

Integrand size = 33, antiderivative size = 99

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{(Ac - cC + Bd)x}{c^2 + d^2} - \frac{(Bc - (A - C)d) \log(\cos(e + fx))}{(c^2 + d^2) f}$$

$$+ \frac{(c^2C - Bcd + Ad^2) \log(c + d \tan(e + fx))}{d(c^2 + d^2) f}$$

output

$(A*c+B*d-C*c)*x/(c^2+d^2)-(B*c-(A-C)*d)*\ln(\cos(f*x+e))/(c^2+d^2)/f+(A*d^2-B*c*d+C*c^2)*\ln(c+d*\tan(f*x+e))/d/(c^2+d^2)/f$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.18

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{\frac{(-iA+B+iC) \log(i-\tan(e+fx))}{c+id} + \frac{(iA+B-iC) \log(i+\tan(e+fx))}{c-id} + \frac{2(c^2C-Bcd+Ad^2) \log(c+d \tan(e+fx))}{d(c^2+d^2)}}{2f}$$

input `Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x]),x]`

output `((((-I)*A + B + I*C)*Log[I - Tan[e + f*x]])/(c + I*d) + ((I*A + B - I*C)*Log[I + Tan[e + f*x]])/(c - I*d) + (2*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]])/(d*(c^2 + d^2)))/(2*f)`

Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)^2}{c + d \tan(e + fx)} dx$$

↓ 4109

$$\frac{(Ad^2 - Bcd + c^2C) \int \frac{\tan^2(e+fx)+1}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \frac{(Bc - d(A - C)) \int \tan(e + fx) dx}{c^2 + d^2} + \frac{x(Ac + Bd - cC)}{c^2 + d^2}$$

↓ 3042

$$\frac{(Bc - d(A - C)) \int \tan(e + fx) dx}{c^2 + d^2} + \frac{(Ad^2 - Bcd + c^2C) \int \frac{\tan(e+fx)^2+1}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \frac{x(Ac + Bd - cC)}{c^2 + d^2}$$

↓ 3956

$$\frac{(Ad^2 - Bcd + c^2C) \int \frac{\tan(e+fx)^2+1}{c+d \tan(e+fx)} dx}{c^2 + d^2} - \frac{(Bc - d(A - C)) \log(\cos(e + fx))}{f(c^2 + d^2)} + \frac{x(Ac + Bd - cC)}{c^2 + d^2}$$

↓ 4100

$$\frac{(Ad^2 - Bcd + c^2C) \int \frac{1}{c+d \tan(e+fx)} d(d \tan(e+fx))}{df(c^2 + d^2)} - \frac{(Bc - d(A - C)) \log(\cos(e+fx))}{f(c^2 + d^2)} + \frac{x(Ac + Bd - cC)}{c^2 + d^2}$$

↓ 16

$$\frac{(Ad^2 - Bcd + c^2C) \log(c + d \tan(e+fx))}{df(c^2 + d^2)} - \frac{(Bc - d(A - C)) \log(\cos(e+fx))}{f(c^2 + d^2)} + \frac{x(Ac + Bd - cC)}{c^2 + d^2}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x]),x]`

output `((A*c - c*C + B*d)*x)/(c^2 + d^2) - ((B*c - (A - C)*d)*Log[Cos[e + f*x]])/((c^2 + d^2)*f) + ((c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]])/(d*(c^2 + d^2)*f)`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2
)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.01

method	result
derivativedivides	$\frac{\frac{(-Ad+Bc+Cd)\ln(1+\tan(fx+e))^2}{2} + (Ac+Bd-Ce)\arctan(\tan(fx+e))}{c^2+d^2} + \frac{(Ad^2-Bcd+c^2C)\ln(c+d\tan(fx+e))}{(c^2+d^2)d}$
default	$\frac{\frac{(-Ad+Bc+Cd)\ln(1+\tan(fx+e))^2}{2} + (Ac+Bd-Ce)\arctan(\tan(fx+e))}{c^2+d^2} + \frac{(Ad^2-Bcd+c^2C)\ln(c+d\tan(fx+e))}{(c^2+d^2)d}$
norman	$\frac{(Ac+Bd-Cc)x}{c^2+d^2} + \frac{(Ad^2-Bcd+c^2C)\ln(c+d\tan(fx+e))}{d(c^2+d^2)f} - \frac{(Ad-Bc-Cd)\ln(1+\tan(fx+e)^2)}{2f(c^2+d^2)}$
parallelrisch	$-\frac{2Axcdf-2Bxd^2f+2Cxdf+A\ln(1+\tan(fx+e)^2)d^2-2A\ln(c+d\tan(fx+e))d^2-B\ln(1+\tan(fx+e)^2)cd+2B\ln(c+d\tan(fx+e))d}{2(c^2+d^2)fd}$
risch	$\frac{ixB}{id-c} - \frac{xA}{id-c} + \frac{xC}{id-c} - \frac{2idAx}{c^2+d^2} - \frac{2idAe}{(c^2+d^2)f} + \frac{2iBcx}{c^2+d^2} + \frac{2iBce}{(c^2+d^2)f} - \frac{2ic^2Cx}{(c^2+d^2)d} - \frac{2ic^2Ce}{(c^2+d^2)fd} + \frac{2iCx}{d} +$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x,method=_RETURNVERBO
SE)
```

output

```
1/f*(1/(c^2+d^2)*(1/2*(-A*d+B*c+C*d)*ln(1+tan(f*x+e)^2)+(A*c+B*d-C*c)*arct
an(tan(f*x+e)))+(A*d^2-B*c*d+C*c^2)/(c^2+d^2)/d*ln(c+d*tan(f*x+e)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.19

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{2((A - C)cd + Bd^2)fx + (Cc^2 - Bcd + Ad^2) \log\left(\frac{d^2 \tan^2(fx+e) + 2cd \tan(fx+e) + c^2}{\tan^2(fx+e) + 1}\right) - (Cc^2 + Cd^2) \log\left(\frac{1}{\tan^2(fx+e) + 1}\right)}{2(c^2d + d^3)f}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="fricas")`

output `1/2*(2*((A - C)*c*d + B*d^2)*f*x + (C*c^2 - B*c*d + A*d^2)*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - (C*c^2 + C*d^2)*log(1/(tan(f*x + e)^2 + 1)))/((c^2*d + d^3)*f)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 966, normalized size of antiderivative = 9.76

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)),x)`

output

```
Piecewise((zoo*x*(A + B*tan(e) + C*tan(e)**2)/tan(e), Eq(c, 0) & Eq(d, 0)
& Eq(f, 0)), ((A*x + B*log(tan(e + f*x)**2 + 1)/(2*f) - C*x + C*tan(e + f*
x)/f)/c, Eq(d, 0)), (I*A*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) +
A*f*x/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*A/(2*d*f*tan(e + f*x) - 2*I*d*f)
+ B*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*B*f*x/(2*d*f*tan(
e + f*x) - 2*I*d*f) - B/(2*d*f*tan(e + f*x) - 2*I*d*f) + I*C*f*x*tan(e + f
*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) + C*f*x/(2*d*f*tan(e + f*x) - 2*I*d*f)
+ C*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*d*f*tan(e + f*x) - 2*I*d*f) -
I*C*log(tan(e + f*x)**2 + 1)/(2*d*f*tan(e + f*x) - 2*I*d*f) - I*C/(2*d*f*
tan(e + f*x) - 2*I*d*f), Eq(c, -I*d)), (-I*A*f*x*tan(e + f*x)/(2*d*f*tan(e
+ f*x) + 2*I*d*f) + A*f*x/(2*d*f*tan(e + f*x) + 2*I*d*f) - I*A/(2*d*f*tan
(e + f*x) + 2*I*d*f) + B*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) + 2*I*d*f) +
I*B*f*x/(2*d*f*tan(e + f*x) + 2*I*d*f) - B/(2*d*f*tan(e + f*x) + 2*I*d*f)
- I*C*f*x*tan(e + f*x)/(2*d*f*tan(e + f*x) + 2*I*d*f) + C*f*x/(2*d*f*tan(
e + f*x) + 2*I*d*f) + C*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*d*f*tan(e
+ f*x) + 2*I*d*f) + I*C*log(tan(e + f*x)**2 + 1)/(2*d*f*tan(e + f*x) + 2*
I*d*f) + I*C/(2*d*f*tan(e + f*x) + 2*I*d*f), Eq(c, I*d)), (x*(A + B*tan(e)
+ C*tan(e)**2)/(c + d*tan(e)), Eq(f, 0)), (2*A*c*d*f*x/(2*c**2*d*f + 2*d*
*3*f) + 2*A*d**2*log(c/d + tan(e + f*x))/(2*c**2*d*f + 2*d**3*f) - A*d**2*
log(tan(e + f*x)**2 + 1)/(2*c**2*d*f + 2*d**3*f) - 2*B*c*d*log(c/d + ta...
```

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.07

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{2((A-C)c + Bd)(fx+e)}{c^2+d^2} + \frac{2(Cc^2 - Bcd + Ad^2) \log(d \tan(fx+e) + c)}{c^2 d + d^3} + \frac{(Bc - (A-C)d) \log(\tan(fx+e)^2 + 1)}{c^2 + d^2}$$

2 f

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="m
axima")
```

output

```
1/2*(2*((A - C)*c + B*d)*(f*x + e)/(c^2 + d^2) + 2*(C*c^2 - B*c*d + A*d^2)
*log(d*tan(f*x + e) + c)/(c^2*d + d^3) + (B*c - (A - C)*d)*log(tan(f*x + e
)^2 + 1)/(c^2 + d^2))/f
```

Giac [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{(Ac - Cc + Bd)(fx + e)}{c^2 f + d^2 f} + \frac{(Bc - Ad + Cd) \log(\tan(fx + e)^2 + 1)}{2(c^2 f + d^2 f)}$$

$$+ \frac{(Cc^2 - Bcd + Ad^2) \log(|d \tan(fx + e) + c|)}{c^2 d f + d^3 f}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="giac")`

output `(A*c - C*c + B*d)*(f*x + e)/(c^2*f + d^2*f) + 1/2*(B*c - A*d + C*d)*log(tan(f*x + e)^2 + 1)/(c^2*f + d^2*f) + (C*c^2 - B*c*d + A*d^2)*log(abs(d*tan(f*x + e) + c))/(c^2*d*f + d^3*f)`

Mupad [B] (verification not implemented)

Time = 6.31 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.10

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{\ln(\tan(e + fx) + 1i) (C - A + B 1i)}{2 f (d + c 1i)} + \frac{\ln(\tan(e + fx) - i) (B - A 1i + C 1i)}{2 f (c + d 1i)}$$

$$+ \frac{\ln(c + d \tan(e + fx)) (C c^2 - B c d + A d^2)}{d f (c^2 + d^2)}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/(c + d*tan(e + f*x)),x)`

output `(log(tan(e + f*x) + 1i)*(B*1i - A + C))/(2*f*(c*1i + d)) + (log(tan(e + f*x) - 1i)*(B - A*1i + C*1i))/(2*f*(c + d*1i)) + (log(c + d*tan(e + f*x))*(A*d^2 + C*c^2 - B*c*d))/(d*f*(c^2 + d^2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.39

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{c + d \tan(e + fx)} dx$$

$$= \frac{-\log(\tan(fx + e)^2 + 1) a d^2 + \log(\tan(fx + e)^2 + 1) bcd + \log(\tan(fx + e)^2 + 1) c d^2 + 2 \log(d \tan(fx + e))}{2df}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)
```

output

```
( - log(tan(e + f*x)**2 + 1)*a*d**2 + log(tan(e + f*x)**2 + 1)*b*c*d + log
(tan(e + f*x)**2 + 1)*c*d**2 + 2*log(tan(e + f*x)*d + c)*a*d**2 - 2*log(ta
n(e + f*x)*d + c)*b*c*d + 2*log(tan(e + f*x)*d + c)*c**3 + 2*a*c*d*f*x + 2
*b*d**2*f*x - 2*c**2*d*f*x)/(2*d*f*(c**2 + d**2))
```


3.74 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))} dx$

Optimal result	820
Mathematica [A] (verified)	821
Rubi [A] (verified)	821
Maple [A] (verified)	823
Fricas [A] (verification not implemented)	824
Sympy [C] (verification not implemented)	824
Maxima [A] (verification not implemented)	825
Giac [A] (verification not implemented)	826
Mupad [B] (verification not implemented)	827
Reduce [B] (verification not implemented)	827

Optimal result

Integrand size = 45, antiderivative size = 165

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$= \frac{(a(Ac - cC + Bd) + b(Bc - (A - C)d))x}{(a^2 + b^2)(c^2 + d^2)}$$

$$+ \frac{(Ab^2 - a(bB - aC)) \log(a \cos(e + fx) + b \sin(e + fx))}{(a^2 + b^2)(bc - ad)f}$$

$$- \frac{(c^2C - Bcd + Ad^2) \log(c \cos(e + fx) + d \sin(e + fx))}{(bc - ad)(c^2 + d^2)f}$$

output

```
(a*(A*c+B*d-C*c)+b*(B*c-(A-C)*d))*x/(a^2+b^2)/(c^2+d^2)+(A*b^2-a*(B*b-C*a)
)*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)/(-a*d+b*c)/f-(A*d^2-B*c*d+C*c^2)
*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)/(c^2+d^2)/f
```

Mathematica [A] (verified)

Time = 0.98 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.90

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx =$$

$$\frac{\left(\frac{Abc - aBc - bcC + aAd + bBd - aCd + \sqrt{-b^2}(bBc + b(-A + C)d + a(Ac - cC + Bd))}{b} \right) \log(\sqrt{-b^2} - b \tan(e + fx))}{(a^2 + b^2)(c^2 + d^2)} + \frac{2(Ab^2 + a(-bB + aC)) \log(a + b \tan(e + fx))}{(a^2 + b^2)(-bc + ad)}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])),x]
```

output

```
-1/2*(((A*b*c - a*B*c - b*c*C + a*A*d + b*B*d - a*C*d + (Sqrt[-b^2]*(b*B*c + b*(-A + C)*d + a*(A*c - c*C + B*d)))/b)*Log[Sqrt[-b^2] - b*Tan[e + f*x]])/((a^2 + b^2)*(c^2 + d^2)) + (2*(A*b^2 + a*(-b*B) + a*C))*Log[a + b*Tan[e + f*x]]/((a^2 + b^2)*(-b*c) + a*d) + ((A*b*c - a*B*c - b*c*C + a*A*d + b*B*d - a*C*d + (b*(b*B*c + b*(-A + C)*d + a*(A*c - c*C + B*d)))/Sqrt[-b^2])*Log[Sqrt[-b^2] + b*Tan[e + f*x]]/((a^2 + b^2)*(c^2 + d^2)) + (2*(c^2*C - B*c*d + A*d^2))*Log[c + d*Tan[e + f*x]]/((b*c - a*d)*(c^2 + d^2))/f
```

Rubi [A] (verified)Time = 1.02 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$\downarrow \text{4134}$$

$$\frac{(Ab^2 - a(bB - aC)) \int \frac{b-a \tan(e+fx)}{a+b \tan(e+fx)} dx}{(a^2 + b^2)(bc - ad)} - \frac{(Ad^2 - Bcd + c^2C) \int \frac{d-c \tan(e+fx)}{c+d \tan(e+fx)} dx}{(c^2 + d^2)(bc - ad)} +$$

$$\frac{x(a(Ac + Bd - cC) - bd(A - C) + bBc)}{(a^2 + b^2)(c^2 + d^2)}$$

↓ 3042

$$\frac{(Ab^2 - a(bB - aC)) \int \frac{b-a \tan(e+fx)}{a+b \tan(e+fx)} dx}{(a^2 + b^2)(bc - ad)} - \frac{(Ad^2 - Bcd + c^2C) \int \frac{d-c \tan(e+fx)}{c+d \tan(e+fx)} dx}{(c^2 + d^2)(bc - ad)} +$$

$$\frac{x(a(Ac + Bd - cC) - bd(A - C) + bBc)}{(a^2 + b^2)(c^2 + d^2)}$$

↓ 4013

$$\frac{x(a(Ac + Bd - cC) - bd(A - C) + bBc)}{(a^2 + b^2)(c^2 + d^2)} +$$

$$\frac{(Ab^2 - a(bB - aC)) \log(a \cos(e + fx) + b \sin(e + fx))}{f(a^2 + b^2)(bc - ad)} -$$

$$\frac{(Ad^2 - Bcd + c^2C) \log(c \cos(e + fx) + d \sin(e + fx))}{f(c^2 + d^2)(bc - ad)}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])),x]
```

output

```
((b*B*c - b*(A - C)*d + a*(A*c - c*C + B*d))*x)/((a^2 + b^2)*(c^2 + d^2)) + ((A*b^2 - a*(b*B - a*C))*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f) - ((c^2*C - B*c*d + A*d^2)*Log[c*Cos[e + f*x] + d*Sin[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f)
```

Defintions of rubi rules used

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4013

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_.)])), x_Symbol] :> Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{(-Aad - Abc + Bac - Bbd + Cad + Cbc) \ln(1 + \tan(fx + e))^2}{2} + (Aac - Abd + Bad + Bbc - Cac + Cbd) \arctan(\tan(fx + e)) - \frac{(Ab^2 - Bab + C)}{(a^2 + b^2)(c^2 + d^2)}$
default	$\frac{(-Aad - Abc + Bac - Bbd + Cad + Cbc) \ln(1 + \tan(fx + e))^2}{2} + (Aac - Abd + Bad + Bbc - Cac + Cbd) \arctan(\tan(fx + e)) - \frac{(Ab^2 - Bab + C)}{(a^2 + b^2)(c^2 + d^2)}$
norman	$\frac{(Aac - Abd + Bad + Bbc - Cac + Cbd)x}{(a^2 + b^2)(c^2 + d^2)} + \frac{(Ad^2 - Bcd + c^2C) \ln(c + d \tan(fx + e))}{f(a^2c^2d + a^2d^3 - bc^3 - bcd^2)} - \frac{(Ab^2 - Bab + Ca^2) \ln(a + b \tan(fx + e))}{(ad - bc)(a^2 + b^2)f}$
parallelrisc	$-2Ax^2cdf + 2Axabc^2f + 2Axabd^2f - 2Axb^2cdf + 2Cxa^2cdf - 2Cxab^2f - 2Cxabd^2f + 2Cxb^2cdf + A \ln(1 + \tan(fx + e))$
risc	$-\frac{2iBabx}{a^3d - a^2bc + ab^2d - b^3c} - \frac{xA}{iad + ibc - ac + bd} + \frac{xC}{iad + ibc - ac + bd} + \frac{2iAb^2x}{a^3d - a^2bc + ab^2d - b^3c} + \frac{2iCa^2x}{a^3d - a^2bc + ab^2d - b^3c}$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)),x,me
thod=_RETURNVERBOSE)
```

output

```
1/f*(1/(a^2+b^2)/(c^2+d^2)*(1/2*(-A*a*d-A*b*c+B*a*c-B*b*d+C*a*d+C*b*c)*ln(
1+tan(f*x+e)^2)+(A*a*c-A*b*d+B*a*d+B*b*c-C*a*c+C*b*d)*arctan(tan(f*x+e)))-
(A*b^2-B*a*b+C*a^2)/(a*d-b*c)/(a^2+b^2)*ln(a+b*tan(f*x+e)+(A*d^2-B*c*d+C*
c^2)/(c^2+d^2)/(a*d-b*c)*ln(c+d*tan(f*x+e)))
```

Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.82

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$= \frac{2(((A - C)ab + Bb^2)c^2 - ((A - C)a^2 + (A - C)b^2)cd - (Ba^2 - (A - C)ab)d^2)fx + ((Ca^2 - Bab + A$$

$$2((a^2b + b^3)c^3 - (a^3 + a*b^2)*c^2*d + (a^2*b + b^3)*c*d^2 - (a^3 + a*b^2)*d^3)*f)$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)),x, algorithm="fricas")
```

output

```
1/2*(2*(((A - C)*a*b + B*b^2)*c^2 - ((A - C)*a^2 + (A - C)*b^2)*c*d - (B*a^2 - (A - C)*a*b)*d^2)*f*x + ((C*a^2 - B*a*b + A*b^2)*c^2 + (C*a^2 - B*a*b + A*b^2)*d^2)*log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1)) - ((C*a^2 + C*b^2)*c^2 - (B*a^2 + B*b^2)*c*d + (A*a^2 + A*b^2)*d^2)*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)))/(((a^2*b + b^3)*c^3 - (a^3 + a*b^2)*c^2*d + (a^2*b + b^3)*c*d^2 - (a^3 + a*b^2)*d^3)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 15.74 (sec) , antiderivative size = 24052, normalized size of antiderivative = 145.77

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx = \text{Too large to display}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)),x)
```

output

```
Piecewise(((2*A*c*d*f*x/(2*c**2*d*f + 2*d**3*f) + 2*A*d**2*log(c/d + tan(e
+ f*x)))/(2*c**2*d*f + 2*d**3*f) - A*d**2*log(tan(e + f*x)**2 + 1)/(2*c**2
*d*f + 2*d**3*f) - 2*B*c*d*log(c/d + tan(e + f*x))/(2*c**2*d*f + 2*d**3*f)
+ B*c*d*log(tan(e + f*x)**2 + 1)/(2*c**2*d*f + 2*d**3*f) + 2*B*d**2*f*x/(
2*c**2*d*f + 2*d**3*f) + 2*C*c**2*log(c/d + tan(e + f*x))/(2*c**2*d*f + 2*
d**3*f) - 2*C*c*d*f*x/(2*c**2*d*f + 2*d**3*f) + C*d**2*log(tan(e + f*x)**2
+ 1)/(2*c**2*d*f + 2*d**3*f))/a, Eq(b, 0)), ((2*A*a*b*f*x/(2*a**2*b*f + 2
*b**3*f) + 2*A*b**2*log(a/b + tan(e + f*x))/(2*a**2*b*f + 2*b**3*f) - A*b*
**2*log(tan(e + f*x)**2 + 1)/(2*a**2*b*f + 2*b**3*f) - 2*B*a*b*log(a/b + ta
n(e + f*x))/(2*a**2*b*f + 2*b**3*f) + B*a*b*log(tan(e + f*x)**2 + 1)/(2*a*
**2*b*f + 2*b**3*f) + 2*B*b**2*f*x/(2*a**2*b*f + 2*b**3*f) + 2*C*a**2*log(a
/b + tan(e + f*x))/(2*a**2*b*f + 2*b**3*f) - 2*C*a*b*f*x/(2*a**2*b*f + 2*b
**3*f) + C*b**2*log(tan(e + f*x)**2 + 1)/(2*a**2*b*f + 2*b**3*f))/c, Eq(d,
0)), (I*A*c**2*f*x*tan(e + f*x)/(2*b*c**3*f*tan(e + f*x) - 2*I*b*c**3*f +
2*I*b*c**2*d*f*tan(e + f*x) + 2*b*c**2*d*f + 2*b*c*d**2*f*tan(e + f*x) -
2*I*b*c*d**2*f + 2*I*b*d**3*f*tan(e + f*x) + 2*b*d**3*f) + A*c**2*f*x/(2*b
*c**3*f*tan(e + f*x) - 2*I*b*c**3*f + 2*I*b*c**2*d*f*tan(e + f*x) + 2*b*c*
**2*d*f + 2*b*c*d**2*f*tan(e + f*x) - 2*I*b*c*d**2*f + 2*I*b*d**3*f*tan(e +
f*x) + 2*b*d**3*f) + I*A*c**2/(2*b*c**3*f*tan(e + f*x) - 2*I*b*c**3*f + 2
*I*b*c**2*d*f*tan(e + f*x) + 2*b*c**2*d*f + 2*b*c*d**2*f*tan(e + f*x) -...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.47

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$= \frac{2(((A-C)a+Bb)c+(Ba-(A-C)b)d)(fx+e)}{(a^2+b^2)c^2+(a^2+b^2)d^2} + \frac{2(Ca^2-Bab+Ab^2) \log(b \tan(fx+e)+a)}{(a^2b+b^3)c-(a^3+ab^2)d} - \frac{2(Cc^2-Bcd+Ad^2) \log(d \tan(fx+e)+c)}{bc^3-ac^2d+bcd^2-ad^3} + \frac{((I$$

2 f

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)
),x, algorithm="maxima")
```

output

```
1/2*(2*((A - C)*a + B*b)*c + (B*a - (A - C)*b)*d)*(f*x + e)/((a^2 + b^2)*
c^2 + (a^2 + b^2)*d^2) + 2*(C*a^2 - B*a*b + A*b^2)*log(b*tan(f*x + e) + a)
/((a^2*b + b^3)*c - (a^3 + a*b^2)*d) - 2*(C*c^2 - B*c*d + A*d^2)*log(d*tan
(f*x + e) + c)/(b*c^3 - a*c^2*d + b*c*d^2 - a*d^3) + ((B*a - (A - C)*b)*c
- ((A - C)*a + B*b)*d)*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*c^2 + (a^2 + b
^2)*d^2))/f
```

Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.68

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$= \frac{(Aac - Cac + Bbc + Bad - Abd + Cbd)(fx + e)}{a^2c^2f + b^2c^2f + a^2d^2f + b^2d^2f}$$

$$+ \frac{(Bac - Abc + Cbc - Aad + Cad - Bbd) \log(\tan(fx + e)^2 + 1)}{2(a^2c^2f + b^2c^2f + a^2d^2f + b^2d^2f)}$$

$$+ \frac{(Ca^2b - Bab^2 + Ab^3) \log(|b \tan(fx + e) + a|)}{a^2b^2cf + b^4cf - a^3bdf - ab^3df}$$

$$- \frac{(Cc^2d - Bcd^2 + Ad^3) \log(|d \tan(fx + e) + c|)}{bc^3df - ac^2d^2f + bcd^3f - ad^4f}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)
),x, algorithm="giac")
```

output

```
(A*a*c - C*a*c + B*b*c + B*a*d - A*b*d + C*b*d)*(f*x + e)/(a^2*c^2*f + b^2
*c^2*f + a^2*d^2*f + b^2*d^2*f) + 1/2*(B*a*c - A*b*c + C*b*c - A*a*d + C*a
*d - B*b*d)*log(tan(f*x + e)^2 + 1)/(a^2*c^2*f + b^2*c^2*f + a^2*d^2*f + b
^2*d^2*f) + (C*a^2*b - B*a*b^2 + A*b^3)*log(abs(b*tan(f*x + e) + a))/(a^2*
b^2*c*f + b^4*c*f - a^3*b*d*f - a*b^3*d*f) - (C*c^2*d - B*c*d^2 + A*d^3)*l
og(abs(d*tan(f*x + e) + c))/(b*c^3*d*f - a*c^2*d^2*f + b*c*d^3*f - a*d^4*f
)
```

Mupad [B] (verification not implemented)

Time = 17.56 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.19

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$= \frac{\ln(c + d \tan(e + fx)) (C c^2 - B c d + A d^2)}{f (a d - b c) (c^2 + d^2)}$$

$$+ \frac{\ln(\tan(e + fx) + 1i) (C - A + B 1i)}{2 f (a c 1i + a d + b c - b d 1i)}$$

$$- \frac{\ln(a + b \tan(e + fx)) (C a^2 - B a b + A b^2)}{f (d a^3 - c a^2 b + d a b^2 - c b^3)}$$

$$- \frac{\ln(\tan(e + fx) - i) (A - C + B 1i)}{2 f (a d - a c 1i + b c + b d 1i)}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))),x)
```

output

```
(log(tan(e + f*x) + 1i)*(B*1i - A + C))/(2*f*(a*c*1i + a*d + b*c - b*d*1i)) - (log(tan(e + f*x) - 1i)*(A + B*1i - C))/(2*f*(a*d - a*c*1i + b*c + b*d*1i)) - (log(a + b*tan(e + f*x))*(A*b^2 + C*a^2 - B*a*b))/(f*(a^3*d - b^3*c - a^2*b*c + a*b^2*d)) + (log(c + d*tan(e + f*x))*(A*d^2 + C*c^2 - B*c*d))/(f*(a*d - b*c)*(c^2 + d^2))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.56

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx$$

$$= \frac{-\log(\tan(fx + e)^2 + 1) a^3 d^2 + \log(\tan(fx + e)^2 + 1) a^2 b c d + \log(\tan(fx + e)^2 + 1) a^2 c d^2 - \log(\tan$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)),x)
```


output

```
( - log(tan(e + f*x)**2 + 1)*a**3*d**2 + log(tan(e + f*x)**2 + 1)*a**2*b*c
*d + log(tan(e + f*x)**2 + 1)*a**2*c*d**2 - log(tan(e + f*x)**2 + 1)*a*b**
2*d**2 + log(tan(e + f*x)**2 + 1)*b**3*c*d - log(tan(e + f*x)**2 + 1)*b**2
*c**3 - 2*log(tan(e + f*x)*b + a)*a**2*c**3 - 2*log(tan(e + f*x)*b + a)*a
*2*c*d**2 + 2*log(tan(e + f*x)*d + c)*a**3*d**2 - 2*log(tan(e + f*x)*d + c
)*a**2*b*c*d + 2*log(tan(e + f*x)*d + c)*a**2*c**3 + 2*log(tan(e + f*x)*d
+ c)*a*b**2*d**2 - 2*log(tan(e + f*x)*d + c)*b**3*c*d + 2*log(tan(e + f*x)
*d + c)*b**2*c**3 + 2*a**3*c*d*f*x - 2*a**2*b*c**2*f*x - 2*a**2*c**2*d*f*x
+ 2*a*b**2*c*d*f*x + 2*a*b*c**3*f*x + 2*a*b*c*d**2*f*x - 2*b**3*c**2*f*x
- 2*b**2*c**2*d*f*x)/(2*f*(a**3*c**2*d + a**3*d**3 - a**2*b*c**3 - a**2*b*
c*d**2 + a*b**2*c**2*d + a*b**2*d**3 - b**3*c**3 - b**3*c*d**2))
```

3.75 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))} dx$

Optimal result	829
Mathematica [A] (verified)	830
Rubi [A] (verified)	830
Maple [A] (verified)	833
Fricas [B] (verification not implemented)	834
Sympy [F(-2)]	835
Maxima [A] (verification not implemented)	836
Giac [B] (verification not implemented)	837
Mupad [B] (verification not implemented)	838
Reduce [B] (verification not implemented)	839

Optimal result

Integrand size = 45, antiderivative size = 281

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2(c + d \tan(e + fx))} dx$$

$$= \frac{(a^2(Ac - cC + Bd) - b^2(Ac - cC + Bd) + 2ab(Bc - (A - C)d))x}{(a^2 + b^2)^2(c^2 + d^2)}$$

$$+ \frac{(2ab^3c(A - C) + 2a^3bBd - a^4Cd + b^4(Bc - Ad) - a^2b^2(Bc + 3Ad - Cd)) \log(a \cos(e + fx) + b \sin(e + fx))}{(a^2 + b^2)^2(bc - ad)^2 f}$$

$$+ \frac{d(c^2C - Bcd + Ad^2) \log(c \cos(e + fx) + d \sin(e + fx))}{(bc - ad)^2(c^2 + d^2) f}$$

$$- \frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))}$$

output

```
(a^2*(A*c+B*d-C*c)-b^2*(A*c+B*d-C*c)+2*a*b*(B*c-(A-C)*d))*x/(a^2+b^2)^2/(c^2+d^2)+(2*a*b^3*c*(A-C)+2*a^3*b*B*d-a^4*C*d+b^4*(-A*d+B*c)-a^2*b^2*(3*A*d+B*c-C*d))*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)^2/(-a*d+b*c)^2/f+d*(A*d^2-B*c*d+C*c^2)*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^2/(c^2+d^2)/f-(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))
```

Mathematica [A] (verified)

Time = 4.48 (sec) , antiderivative size = 543, normalized size of antiderivative = 1.93

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx$$

$$= \frac{(bc-ad) \left(2aAbc - a^2Bc + b^2Bc - 2abcC + a^2Ad - Ab^2d + 2abBd - a^2Cd + b^2Cd + \frac{\sqrt{-b^2} (a^2(Ac - cC + Bd) - b^2(Ac - cC + Bd) + 2ab(Bc + (-A + C)d))}{b} \right)}{2(a^2 + b^2)(c^2 + d^2)}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])),x]
```

output

```
(-1/2*((b*c - a*d)*(2*a*A*b*c - a^2*B*c + b^2*B*c - 2*a*b*c*C + a^2*A*d - A*b^2*d + 2*a*b*B*d - a^2*C*d + b^2*C*d + (Sqrt[-b^2]*(a^2*(A*c - c*C + B*d) - b^2*(A*c - c*C + B*d) + 2*a*b*(B*c + (-A + C)*d)))/b)*Log[Sqrt[-b^2] - b*Tan[e + f*x]]/((a^2 + b^2)*(c^2 + d^2)) + ((2*a*b^3*c*(-A + C) - 2*a^3*b*B*d + a^4*C*d + b^4*(-(B*c) + A*d) + a^2*b^2*(B*c + 3*A*d - C*d))*Log[a + b*Tan[e + f*x]]/((a^2 + b^2)*(-(b*c) + a*d)) - ((b*c - a*d)*(2*a*A*b*c - a^2*B*c + b^2*B*c - 2*a*b*c*C + a^2*A*d - A*b^2*d + 2*a*b*B*d - a^2*C*d + b^2*C*d + (Sqrt[-b^2]*(-(a^2*(A*c - c*C + B*d)) + b^2*(A*c - c*C + B*d) - 2*a*b*(B*c + (-A + C)*d)))/b)*Log[Sqrt[-b^2] + b*Tan[e + f*x]]/(2*(a^2 + b^2)*(c^2 + d^2)) + ((a^2 + b^2)*d*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]]/((b*c - a*d)*(c^2 + d^2)) - (A*b^2)/(a + b*Tan[e + f*x]) + (a*(b*B - a*C))/(a + b*Tan[e + f*x]))/((a^2 + b^2)*(b*c - a*d)*f)
```

Rubi [A] (verified)Time = 2.18 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.13, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4132, 25, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx \\
& \downarrow 4132 \\
& \int \frac{-Ada^2 + bc(A-C)a - (Ab^2 - a(bB - aC))d \tan^2(e + fx) + b^2(Bc - Ad) - (Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx \\
& \frac{(a^2 + b^2)(bc - ad)}{Ab^2 - a(bB - aC)} \\
& \frac{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}{} \\
& \downarrow 25 \\
& \int \frac{-Ada^2 + bc(A-C)a - (Ab^2 - a(bB - aC))d \tan^2(e + fx) + b^2(Bc - Ad) - (Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx \\
& \frac{(a^2 + b^2)(bc - ad)}{Ab^2 - a(bB - aC)} \\
& \frac{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}{} \\
& \downarrow 3042 \\
& \int \frac{-Ada^2 + bc(A-C)a - (Ab^2 - a(bB - aC))d \tan(e + fx)^2 + b^2(Bc - Ad) - (Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx \\
& \frac{(a^2 + b^2)(bc - ad)}{Ab^2 - a(bB - aC)} \\
& \frac{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}{} \\
& \downarrow 4134 \\
& \frac{d(a^2 + b^2)(Ad^2 - Bcd + c^2C) \int \frac{d - c \tan(e + fx)}{c + d \tan(e + fx)} dx}{(c^2 + d^2)(bc - ad)} + \frac{(a^4(-C)d + 2a^3bBd - a^2b^2(3Ad + Bc - Cd) + 2ab^3c(A - C) + b^4(Bc - Ad)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx}{(a^2 + b^2)(bc - ad)} + \\
& \frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} \\
& \downarrow 3042 \\
& \frac{d(a^2 + b^2)(Ad^2 - Bcd + c^2C) \int \frac{d - c \tan(e + fx)}{c + d \tan(e + fx)} dx}{(c^2 + d^2)(bc - ad)} + \frac{(a^4(-C)d + 2a^3bBd - a^2b^2(3Ad + Bc - Cd) + 2ab^3c(A - C) + b^4(Bc - Ad)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx}{(a^2 + b^2)(bc - ad)} + \\
& \frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} \\
& \downarrow 4013
\end{aligned}$$

$$\frac{\frac{d(a^2+b^2)(Ad^2-Bcd+c^2C) \log(c \cos(e+fx)+d \sin(e+fx))}{f(c^2+d^2)(bc-ad)} + \frac{x(bc-ad)(a^2(Ac+Bd-cC)+2ab(Bc-d(A-C))-b^2(Ac+Bd-cC))}{(a^2+b^2)(c^2+d^2)} + \frac{(a^4(-C))}{(a^2+b^2)(bc-ad)}}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} \frac{Ab^2 - a(bB - aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])),x]`

output `((((b*c - a*d)*(a^2*(A*c - c*C + B*d) - b^2*(A*c - c*C + B*d) + 2*a*b*(B*c - (A - C)*d))*x)/((a^2 + b^2)*(c^2 + d^2)) + ((2*a*b^3*c*(A - C) + 2*a^3*b*B*d - a^4*C*d + b^4*(B*c - A*d) - a^2*b^2*(B*c + 3*A*d - C*d))*Log[a*Cos[e + f*x] + b*Sin[e + f*x]]/((a^2 + b^2)*(b*c - a*d)*f) + ((a^2 + b^2)*d*(c^2*C - B*c*d + A*d^2)*Log[c*Cos[e + f*x] + d*Sin[e + f*x]]/((b*c - a*d)*(c^2 + d^2)*f))/((a^2 + b^2)*(b*c - a*d)) - (A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x]))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
(!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.30

method	result
derivativedivides	$\frac{(-A a^2 d - 2A a b c + A b^2 d + B a^2 c - 2B a b d - B b^2 c + C a^2 d + 2C a b c - d C b^2) \ln(1 + \tan(fx + e)^2)}{2(a^2 + b^2)^2(c^2 + d^2)} + \frac{(A a^2 c - 2A a b d - A b^2 c + B a^2 d + 2B a b c - B b^2 d - C a^2 c + 2C a b d + C b^2 c)x}{(a^2 + b^2)^2(c^2 + d^2)}$
default	$\frac{(-A a^2 d - 2A a b c + A b^2 d + B a^2 c - 2B a b d - B b^2 c + C a^2 d + 2C a b c - d C b^2) \ln(1 + \tan(fx + e)^2)}{2(a^2 + b^2)^2(c^2 + d^2)} + \frac{(A a^2 c - 2A a b d - A b^2 c + B a^2 d + 2B a b c - B b^2 d - C a^2 c + 2C a b d + C b^2 c)x}{(a^2 + b^2)^2(c^2 + d^2)}$
norman	$\frac{a(A a^2 c - 2A a b d - A b^2 c + B a^2 d + 2B a b c - B b^2 d - C a^2 c + 2C a b d + C b^2 c)x}{(a^4 + 2b^2 a^2 + b^4)(c^2 + d^2)} + \frac{b^3 A - B a b^2 + C a^2 b}{b f(a d - b c)(a^2 + b^2)} + \frac{(A a^2 c - 2A a b d - A b^2 c + B a^2 d + 2B a b c - B b^2 d - C a^2 c + 2C a b d + C b^2 c)}{(a^4 + 2b^2 a^2 + b^4)(c^2 + d^2)}$
parallelrisc	Expression too large to display
risc	Expression too large to display

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e)),x,
method=_RETURNVERBOSE)`

output `1/f*(1/(a^2+b^2)^2/(c^2+d^2)*(1/2*(-A*a^2*d-2*A*a*b*c+A*b^2*d+B*a^2*c-2*B*
a*b*d-B*b^2*c+C*a^2*d+2*C*a*b*c-C*b^2*d)*ln(1+tan(f*x+e)^2)+(A*a^2*c-2*A*a
*b*d-A*b^2*c+B*a^2*d+2*B*a*b*c-B*b^2*d-C*a^2*c+2*C*a*b*d+C*b^2*c)*arctan(t
an(f*x+e)))-(3*A*a^2*b^2*d-2*A*a*b^3*c+A*b^4*d-2*B*a^3*b*d+B*a^2*b^2*c-B*b
^4*c+C*a^4*d-C*a^2*b^2*d+2*C*a*b^3*c)/(a*d-b*c)^2/(a^2+b^2)^2*ln(a+b*tan(f
*x+e))+(A*b^2-B*a*b+C*a^2)/(a*d-b*c)/(a^2+b^2)/(a+b*tan(f*x+e))+(A*d^2-B*c
*d+C*c^2)*d/(c^2+d^2)/(a*d-b*c)^2*ln(c+d*tan(f*x+e)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1345 vs. $2(280) = 560$.

Time = 0.89 (sec) , antiderivative size = 1345, normalized size of antiderivative = 4.79

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+
e)),x, algorithm="fricas")`

output

```

-1/2*(2*(C*a^2*b^3 - B*a*b^4 + A*b^5)*c^3 - 2*(C*a^3*b^2 - B*a^2*b^3 + A*a
*b^4)*c^2*d + 2*(C*a^2*b^3 - B*a*b^4 + A*b^5)*c*d^2 - 2*(C*a^3*b^2 - B*a^2
*b^3 + A*a*b^4)*d^3 - 2*((A - C)*a^3*b^2 + 2*B*a^2*b^3 - (A - C)*a*b^4)*c
^3 - (2*(A - C)*a^4*b + 3*B*a^3*b^2 + B*a*b^4)*c^2*d + ((A - C)*a^5 + 3*(A
- C)*a^3*b^2 + 2*B*a^2*b^3)*c*d^2 + (B*a^5 - 2*(A - C)*a^4*b - B*a^3*b^2)
*d^3)*f*x + ((B*a^3*b^2 - 2*(A - C)*a^2*b^3 - B*a*b^4)*c^3 + (C*a^5 - 2*B*
a^4*b + (3*A - C)*a^3*b^2 + A*a*b^4)*c^2*d + (B*a^3*b^2 - 2*(A - C)*a^2*b^
3 - B*a*b^4)*c*d^2 + (C*a^5 - 2*B*a^4*b + (3*A - C)*a^3*b^2 + A*a*b^4)*d^3
+ ((B*a^2*b^3 - 2*(A - C)*a*b^4 - B*b^5)*c^3 + (C*a^4*b - 2*B*a^3*b^2 + (
3*A - C)*a^2*b^3 + A*b^5)*c^2*d + (B*a^2*b^3 - 2*(A - C)*a*b^4 - B*b^5)*c*
d^2 + (C*a^4*b - 2*B*a^3*b^2 + (3*A - C)*a^2*b^3 + A*b^5)*d^3)*tan(f*x + e
))*log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1
)) - ((C*a^5 + 2*C*a^3*b^2 + C*a*b^4)*c^2*d - (B*a^5 + 2*B*a^3*b^2 + B*a*b
^4)*c*d^2 + (A*a^5 + 2*A*a^3*b^2 + A*a*b^4)*d^3 + ((C*a^4*b + 2*C*a^2*b^3
+ C*b^5)*c^2*d - (B*a^4*b + 2*B*a^2*b^3 + B*b^5)*c*d^2 + (A*a^4*b + 2*A*a^
2*b^3 + A*b^5)*d^3)*tan(f*x + e))*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x
+ e) + c^2)/(tan(f*x + e)^2 + 1)) - 2*((C*a^3*b^2 - B*a^2*b^3 + A*a*b^4)*c
^3 - (C*a^4*b - B*a^3*b^2 + A*a^2*b^3)*c^2*d + (C*a^3*b^2 - B*a^2*b^3 + A*
a*b^4)*c*d^2 - (C*a^4*b - B*a^3*b^2 + A*a^2*b^3)*d^3 + (((A - C)*a^2*b^3 +
2*B*a*b^4 - (A - C)*b^5)*c^3 - (2*(A - C)*a^3*b^2 + 3*B*a^2*b^3 + B*b^...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx$$

= Exception raised: NotImplementedError

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**2/(c+d*tan(f*
x+e)),x)

```

output

```

Exception raised: NotImplementedError >> no valid subset found

```


Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 520, normalized size of antiderivative = 1.85

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx$$

$$= \frac{2(((A-C)a^2 + 2Bab - (A-C)b^2)c + (Ba^2 - 2(A-C)ab - Bb^2)d)(fx+e)}{(a^4 + 2a^2b^2 + b^4)c^2 + (a^4 + 2a^2b^2 + b^4)d^2} - \frac{2((Ba^2b^2 - 2(A-C)ab^3 - Bb^4)c + (Ca^4 - 2Ba^3b + (3A-C)a^2b^2 + A$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e)),x, algorithm="maxima")`

output `1/2*(2*(((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c + (B*a^2 - 2*(A - C)*a*b - B*b^2)*d)*(f*x + e)/((a^4 + 2*a^2*b^2 + b^4)*c^2 + (a^4 + 2*a^2*b^2 + b^4)*d^2) - 2*((B*a^2*b^2 - 2*(A - C)*a*b^3 - B*b^4)*c + (C*a^4 - 2*B*a^3*b + (3*A - C)*a^2*b^2 + A*b^4)*d)*log(b*tan(f*x + e) + a)/((a^4*b^2 + 2*a^2*b^4 + b^6)*c^2 - 2*(a^5*b + 2*a^3*b^3 + a*b^5)*c*d + (a^6 + 2*a^4*b^2 + a^2*b^4)*d^2) + 2*(C*c^2*d - B*c*d^2 + A*d^3)*log(d*tan(f*x + e) + c)/(b^2*c^4 - 2*a*b*c^3*d - 2*a*b*c*d^3 + a^2*d^4 + (a^2 + b^2)*c^2*d^2) + ((B*a^2 - 2*(A - C)*a*b - B*b^2)*c - ((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*d)*log(tan(f*x + e)^2 + 1)/((a^4 + 2*a^2*b^2 + b^4)*c^2 + (a^4 + 2*a^2*b^2 + b^4)*d^2) - 2*(C*a^2 - B*a*b + A*b^2)/((a^3*b + a*b^3)*c - (a^4 + a^2*b^2)*d + ((a^2*b^2 + b^4)*c - (a^3*b + a*b^3)*d)*tan(f*x + e))/f`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 674 vs. $2(280) = 560$.

Time = 0.63 (sec) , antiderivative size = 674, normalized size of antiderivative = 2.40

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx$$

$$= \frac{(Aa^2c - Ca^2c + 2Babc - Ab^2c + Cb^2c + Ba^2d - 2Aabd + 2Cabd - Bb^2d)(fx + e)}{a^4c^2f + 2a^2b^2c^2f + b^4c^2f + a^4d^2f + 2a^2b^2d^2f + b^4d^2f}$$

$$+ \frac{(Ba^2c - 2Aabc + 2Cabc - Bb^2c - Aa^2d + Ca^2d - 2Babd + Ab^2d - Cb^2d) \log(\tan(fx + e)^2 + 1)}{2(a^4c^2f + 2a^2b^2c^2f + b^4c^2f + a^4d^2f + 2a^2b^2d^2f + b^4d^2f)}$$

$$- \frac{(Ba^2b^3c - 2Aab^4c + 2Cab^4c - Bb^5c + Ca^4bd - 2Ba^3b^2d + 3Aa^2b^3d - Ca^2b^3d + Ab^5d) \log(|b \tan(fx + e)|)}{a^4b^3c^2f + 2a^2b^5c^2f + b^7c^2f - 2a^5b^2cdf - 4a^3b^4cdf - 2ab^6cdf + a^6bd^2f + 2a^4b^3d^2f + a^2b^5d^2f}$$

$$+ \frac{(C^2d^2 - Bcd^3 + Ad^4) \log(|d \tan(fx + e) + c|)}{b^2c^4df - 2abc^3d^2f + a^2c^2d^3f + b^2c^2d^3f - 2abcd^4f + a^2d^5f}$$

$$- \frac{Ca^4bc - Ba^3b^2c + Aa^2b^3c + Ca^2b^3c - Bab^4c + Ab^5c - Ca^5d + Ba^4bd - Aa^3b^2d - Ca^3b^2d + Ba^2b^3d}{(a^2 + b^2)^2(bc - ad)^2(b \tan(fx + e) + a)f}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e)),x, algorithm="giac")
```

output

```
(A*a^2*c - C*a^2*c + 2*B*a*b*c - A*b^2*c + C*b^2*c + B*a^2*d - 2*A*a*b*d + 2*C*a*b*d - B*b^2*d)*(f*x + e)/(a^4*c^2*f + 2*a^2*b^2*c^2*f + b^4*c^2*f + a^4*d^2*f + 2*a^2*b^2*d^2*f + b^4*d^2*f) + 1/2*(B*a^2*c - 2*A*a*b*c + 2*C*a*b*c - B*b^2*c - A*a^2*d + C*a^2*d - 2*B*a*b*d + A*b^2*d - C*b^2*d)*log(tan(f*x + e)^2 + 1)/(a^4*c^2*f + 2*a^2*b^2*c^2*f + b^4*c^2*f + a^4*d^2*f + 2*a^2*b^2*d^2*f + b^4*d^2*f) - (B*a^2*b^3*c - 2*A*a*b^4*c + 2*C*a*b^4*c - B*b^5*c + C*a^4*b*d - 2*B*a^3*b^2*d + 3*A*a^2*b^3*d - C*a^2*b^3*d + A*b^5*d)*log(abs(b*tan(f*x + e) + a))/(a^4*b^3*c^2*f + 2*a^2*b^5*c^2*f + b^7*c^2*f - 2*a^5*b^2*c*d*f - 4*a^3*b^4*c*d*f - 2*a*b^6*c*d*f + a^6*b*d^2*f + 2*a^4*b^3*d^2*f + a^2*b^5*d^2*f) + (C*c^2*d^2 - B*c*d^3 + A*d^4)*log(abs(d*tan(f*x + e) + c))/(b^2*c^4*d*f - 2*a*b*c^3*d^2*f + a^2*c^2*d^3*f + b^2*c^2*d^3*f - 2*a*b*c*d^4*f + a^2*d^5*f) - (C*a^4*b*c - B*a^3*b^2*c + A*a^2*b^3*c + C*a^2*b^3*c - B*a*b^4*c + A*b^5*c - C*a^5*d + B*a^4*b*d - A*a^3*b^2*d - C*a^3*b^2*d + B*a^2*b^3*d - A*a*b^4*d)/((a^2 + b^2)^2*(b*c - a*d)^2*(b*tan(f*x + e) + a)*f)
```

Mupad [B] (verification not implemented)

Time = 55.91 (sec) , antiderivative size = 393, normalized size of antiderivative = 1.40

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx$$

$$= \frac{\ln(\tan(e + fx) - i) (B - A i + C i)}{2 f (a^2 c - b^2 c - 2 a b d + a^2 d i - b^2 d i + a b c 2 i)}$$

$$- \frac{\ln(\tan(e + fx) + i) (A i + B - C i)}{2 f (b^2 c - a^2 c + 2 a b d + a^2 d i - b^2 d i + a b c 2 i)}$$

$$- \frac{\ln(a + b \tan(e + fx)) (C d a^4 - 2 B d a^3 b + (3 A d + B c - C d) a^2 b^2 + (2 C c - 2 A c) a b^3 + (A d - C c) a^2 b^4 - 2 a b^5 c d + b^6 c^2)}{f (a^6 d^2 - 2 a^5 b c d + a^4 b^2 c^2 + 2 a^4 b^2 d^2 - 4 a^3 b^3 c d + 2 a^2 b^4 c^2 + a^2 b^4 d^2 - 2 a b^5 c d + b^6 c^2)}$$

$$+ \frac{C a^2 - B a b + A b^2}{f (a d - b c) (a^2 + b^2) (a + b \tan(e + fx))}$$

$$+ \frac{d \ln(c + d \tan(e + fx)) (C c^2 - B c d + A d^2)}{f (a d - b c)^2 (c^2 + d^2)}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))),x)`

output `(log(tan(e + f*x) - i)*(B - A*i + C*i))/(2*f*(a^2*c + a^2*d*i - b^2*c - b^2*d*i + a*b*c*2i - 2*a*b*d)) - (log(tan(e + f*x) + i)*(A*i + B - C*i))/(2*f*(a^2*d*i - a^2*c + b^2*c - b^2*d*i + a*b*c*2i + 2*a*b*d)) - (log(a + b*tan(e + f*x))*(b^4*(A*d - B*c) + a^2*b^2*(3*A*d + B*c - C*d) + C*a^4*d - a*b^3*(2*A*c - 2*C*c) - 2*B*a^3*b*d))/(f*(a^6*d^2 + b^6*c^2 + 2*a^2*b^4*c^2 + a^4*b^2*c^2 + a^2*b^4*d^2 + 2*a^4*b^2*d^2 - 2*a*b^5*c*d - 2*a^5*b*c*d - 4*a^3*b^3*c*d)) + (A*b^2 + C*a^2 - B*a*b)/(f*(a*d - b*c)*(a^2 + b^2)*(a + b*tan(e + f*x))) + (d*log(c + d*tan(e + f*x))*(A*d^2 + C*c^2 - B*c*d))/(f*(a*d - b*c)^2*(c^2 + d^2))`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 2915, normalized size of antiderivative = 10.37

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e)),x)`

output

```
( - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*b*d**3 + log(tan(e + f*x)**
2 + 1)*tan(e + f*x)*a**4*b**2*c*d**2 + log(tan(e + f*x)**2 + 1)*tan(e + f
*x)*a**4*b*c*d**3 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**3*c**2*d
- log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b**3*d**3 - log(tan(e + f*x)*
**2 + 1)*tan(e + f*x)*a**2*b**4*c**3 + log(tan(e + f*x)**2 + 1)*tan(e + f*x
)*a**2*b**4*c*d**2 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**3*c**
3*d - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**3*c*d**3 + log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a*b**5*c**2*d + 2*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)*a*b**4*c**4 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**4*c**2
*d**2 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**6*c**3 - log(tan(e + f*x)
**2 + 1)*tan(e + f*x)*b**5*c**3*d - log(tan(e + f*x)**2 + 1)*a**6*d**3 + 1
og(tan(e + f*x)**2 + 1)*a**5*b*c*d**2 + log(tan(e + f*x)**2 + 1)*a**5*c*d*
*3 + log(tan(e + f*x)**2 + 1)*a**4*b**2*c**2*d - log(tan(e + f*x)**2 + 1)*
a**4*b**2*d**3 - log(tan(e + f*x)**2 + 1)*a**3*b**3*c**3 + log(tan(e + f*x
)**2 + 1)*a**3*b**3*c*d**2 - 3*log(tan(e + f*x)**2 + 1)*a**3*b**2*c**3*d -
log(tan(e + f*x)**2 + 1)*a**3*b**2*c*d**3 + log(tan(e + f*x)**2 + 1)*a**2
*b**4*c**2*d + 2*log(tan(e + f*x)**2 + 1)*a**2*b**3*c**4 + 2*log(tan(e + f
*x)**2 + 1)*a**2*b**3*c**2*d**2 - log(tan(e + f*x)**2 + 1)*a*b**5*c**3 - l
og(tan(e + f*x)**2 + 1)*a*b**4*c**3*d - 2*log(tan(e + f*x)*b + a)*tan(e +
f*x)*a**4*b*c*d...
```

3.76 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^3(c+d \tan(e+fx))} dx$

Optimal result	840
Mathematica [A] (verified)	841
Rubi [A] (verified)	842
Maple [A] (verified)	846
Fricas [B] (verification not implemented)	847
Sympy [F(-2)]	847
Maxima [B] (verification not implemented)	847
Giac [B] (verification not implemented)	848
Mupad [B] (verification not implemented)	849
Reduce [B] (verification not implemented)	850

Optimal result

Integrand size = 45, antiderivative size = 477

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3(c + d \tan(e + fx))} dx$$

$$= \frac{(a^3(Ac - cC + Bd) - 3ab^2(Ac - cC + Bd) + 3a^2b(Bc - (A - C)d) - b^3(Bc - (A - C)d))x}{(a^2 + b^2)^3(c^2 + d^2)}$$

$$+ \frac{(3ab^5Bc^2 - 3a^5bBd^2 + a^6Cd^2 + 3a^4b^2d(Bc + 2Ad - Cd) + b^6(c(cC - Bd) - A(c^2 - d^2)) - a^3b^3(8c(A - C) - 3Bd))}{(a^2 + b^2)^3(bc - ad)}$$

$$- \frac{d^2(c^2C - Bcd + Ad^2) \log(c \cos(e + fx) + d \sin(e + fx))}{(bc - ad)^3(c^2 + d^2)f}$$

$$- \frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2}$$

$$- \frac{2ab^3c(A - C) + 2a^3bBd - a^4Cd + b^4(Bc - Ad) - a^2b^2(Bc + 3Ad - Cd)}{(a^2 + b^2)^2(bc - ad)^2f(a + b \tan(e + fx))}$$

output

```
(a^3*(A*c+B*d-C*c)-3*a*b^2*(A*c+B*d-C*c)+3*a^2*b*(B*c-(A-C)*d)-b^3*(B*c-(A-C)*d))*x/(a^2+b^2)^3/(c^2+d^2)+(3*a*b^5*B*c^2-3*a^5*b*B*d^2+a^6*C*d^2+3*a^4*b^2*d*(2*A*d+B*c-C*d)+b^6*(c*(-B*d+C*c)-A*(c^2-d^2))-a^3*b^3*(8*c*(A-C)*d+B*(c^2-d^2))-3*a^2*b^4*(c*(2*B*d+C*c)-A*(c^2+d^2)))*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)^3/(-a*d+b*c)^3/f-d^2*(A*d^2-B*c*d+C*c^2)*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^3/(c^2+d^2)/f-1/2*(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))^2-(2*a*b^3*c*(A-C)+2*a^3*b*B*d-a^4*C*d+b^4*(-A*d+B*c)-a^2*b^2*(3*A*d+B*c-C*d))/(a^2+b^2)^2/(-a*d+b*c)^2/f/(a+b*tan(f*x+e))
```

Mathematica [A] (verified)

Time = 7.77 (sec) , antiderivative size = 898, normalized size of antiderivative = 1.88

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx$$

$$= -\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2}$$

$$-\frac{b(bc - ad)^2 \left(3a^2 Abc - Ab^3 c - a^3 Bc + 3ab^2 Bc - 3a^2 bcC + b^3 cC + a^3 Ad - 3aAb^2 d + 3a^2 bBd - b^3 Bd - a^3 Cd + 3ab^2 Cd + \sqrt{-b^2} (a^3 (Ac - cC + Bd) - 3ab^2 (Ac - cC - Bd)) \right)}{(a^2 + b^2)(c^2 + d^2)}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])),x]
```

output

```

-1/2*(A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x
])^2) - (((-((b*(b*c - a*d)^2*(3*a^2*A*b*c - A*b^3*c - a^3*B*c + 3*a*b^2*
B*c - 3*a^2*b*c*C + b^3*c*C + a^3*A*d - 3*a*A*b^2*d + 3*a^2*b*B*d - b^3*B*
d - a^3*C*d + 3*a*b^2*C*d + (Sqrt[-b^2]*(a^3*(A*c - c*C + B*d) - 3*a*b^2*(
A*c - c*C + B*d) + 3*a^2*b*(B*c - (A - C)*d) - b^3*(B*c - (A - C)*d)))/b)*
Log[Sqrt[-b^2] - b*Tan[e + f*x]])/((a^2 + b^2)*(c^2 + d^2))) + (2*b*(3*a*b
^5*B*c^2 - 3*a^5*b*B*d^2 + a^6*C*d^2 + 3*a^4*b^2*d*(B*c + 2*A*d - C*d) + b
^6*(c*(c*C - B*d) - A*(c^2 - d^2)) - a^3*b^3*(8*c*(A - C)*d + B*(c^2 - d^2
)) - 3*a^2*b^4*(c*(c*C + 2*B*d) - A*(c^2 + d^2))*Log[a + b*Tan[e + f*x]])
/((a^2 + b^2)*(b*c - a*d)) - (b*(b*c - a*d)^2*(3*a^2*A*b*c - A*b^3*c - a^3
*B*c + 3*a*b^2*B*c - 3*a^2*b*c*C + b^3*c*C + a^3*A*d - 3*a*A*b^2*d + 3*a^2
*b*B*d - b^3*B*d - a^3*C*d + 3*a*b^2*C*d - (Sqrt[-b^2]*(a^3*(A*c - c*C + B
*d) - 3*a*b^2*(A*c - c*C + B*d) + 3*a^2*b*(B*c - (A - C)*d) - b^3*(B*c - (
A - C)*d)))/b)*Log[Sqrt[-b^2] + b*Tan[e + f*x]])/((a^2 + b^2)*(c^2 + d^2))
- (2*b*(a^2 + b^2)^2*d^2*(c^2*C - B*c*d + A*d^2)*Log[c + d*Tan[e + f*x]])
/((b*c - a*d)*(c^2 + d^2))/((b*(a^2 + b^2)*(b*c - a*d)*f)) - (-a*(-2*a*(A
*b^2 - a*(b*B - a*C))*d + 2*b*(A*b - a*B - b*C)*(b*c - a*d)) - 2*b^2*(a*b
*c*(A - C) - a^2*A*d + b^2*(B*c - A*d))/((a^2 + b^2)*(b*c - a*d)*f*(a + b
*Tan[e + f*x]))/(2*(a^2 + b^2)*(b*c - a*d))

```

Rubi [A] (verified)

Time = 4.46 (sec) , antiderivative size = 538, normalized size of antiderivative = 1.13, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4132, 27, 3042, 4132, 25, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx$$

↓ 4132

$$\frac{-\frac{d^2(a^2+b^2)^2(Ad^2-Bcd+c^2C)\int\frac{d-c\tan(e+fx)}{c+d\tan(e+fx)}dx}{(c^2+d^2)(bc-ad)}+\frac{(a^6Cd^2-3a^5bBd^2+3a^4b^2d(2Ad+Bc-Cd)-a^3b^3(8cd(A-C)+B(c^2-d^2))-3a^2b^4(c(2Bd+cC)-A(c^2-d^2))}{(a^2+b^2)(bc-ad)}}{(a^2+b^2)^2}}$$

$$\frac{Ab^2 - a(bB - aC)}{2f(a^2 + b^2)(bc - ad)(a + b\tan(e + fx))^2}$$

↓ 3042

$$\frac{-\frac{d^2(a^2+b^2)^2(Ad^2-Bcd+c^2C)\int\frac{d-c\tan(e+fx)}{c+d\tan(e+fx)}dx}{(c^2+d^2)(bc-ad)}+\frac{(a^6Cd^2-3a^5bBd^2+3a^4b^2d(2Ad+Bc-Cd)-a^3b^3(8cd(A-C)+B(c^2-d^2))-3a^2b^4(c(2Bd+cC)-A(c^2-d^2))}{(a^2+b^2)(bc-ad)}}{(a^2+b^2)^2}}$$

$$\frac{Ab^2 - a(bB - aC)}{2f(a^2 + b^2)(bc - ad)(a + b\tan(e + fx))^2}$$

↓ 4013

$$\frac{-\frac{d^2(a^2+b^2)^2(Ad^2-Bcd+c^2C)\log(c\cos(e+fx)+d\sin(e+fx))}{f(c^2+d^2)(bc-ad)}+\frac{x(bc-ad)^2(a^3(Ac+Bd-cC)+3a^2b(Bc-d(A-C))-3ab^2(Ac+Bd-cC)-b^3(Bc-d(A-C)))}{(a^2+b^2)(c^2+d^2)}}{(a^2+b^2)^2}}$$

$$\frac{Ab^2 - a(bB - aC)}{2f(a^2 + b^2)(bc - ad)(a + b\tan(e + fx))^2}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])),x]
```

output

```
-1/2*(A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])^2) + (((b*c - a*d)^2*(a^3*(A*c - c*C + B*d) - 3*a*b^2*(A*c - c*C + B*d) + 3*a^2*b*(B*c - (A - C)*d) - b^3*(B*c - (A - C)*d))*x)/((a^2 + b^2)*(c^2 + d^2)) + ((3*a*b^5*B*c^2 - 3*a^5*b*B*d^2 + a^6*C*d^2 + 3*a^4*b^2*d*(B*c + 2*A*d - C*d) + b^6*(c*(c*C - B*d) - A*(c^2 - d^2)) - a^3*b^3*(8*c*(A - C)*d + B*(c^2 - d^2)) - 3*a^2*b^4*(c*(c*C + 2*B*d) - A*(c^2 + d^2)))*Log[a*cos[e + f*x] + b*sin[e + f*x]]/((a^2 + b^2)*(b*c - a*d)*f) - ((a^2 + b^2)^2*d^2*(c^2*C - B*c*d + A*d^2)*Log[c*cos[e + f*x] + d*sin[e + f*x]]/((b*c - a*d)*(c^2 + d^2)*f))/((a^2 + b^2)*(b*c - a*d)) - (2*a*b^3*c*(A - C) + 2*a^3*b*B*d - a^4*C*d + b^4*(B*c - A*d) - a^2*b^2*(B*c + 3*A*d - C*d))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x]))/((a^2 + b^2)*(b*c - a*d))
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4013 $\text{Int}[((\text{c}_) + (\text{d}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])/((\text{a}_) + (\text{b}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c}/(\text{b}*\text{f}))*\text{Log}[\text{RemoveContent}[\text{a}*\text{Cos}[\text{e} + \text{f}*x] + \text{b}*\text{Sin}[\text{e} + \text{f}*x], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{a}*c + \text{b}*d, 0]$
- rule 4132 $\text{Int}[((\text{a}_) + (\text{b}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])^m*((\text{c}_) + (\text{d}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])^n*((\text{A}_) + (\text{B}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)]) + (\text{C}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])^2), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{A}*b^2 - \text{a}*(\text{b}*B - \text{a}*C))*(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{m+1}*((\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^{n+1}/(\text{f}*(m+1)*(b*c - a*d)*(a^2 + b^2))), \text{x}] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{m+1}*(\text{c} + \text{d}*\text{Tan}[\text{e} + \text{f}*x])^n*\text{Simp}[\text{A}*(\text{a}*(\text{b}*c - \text{a}*d)*(m+1) - \text{b}^2*\text{d}*(m+n+2)) + (\text{b}*B - \text{a}*C)*(b*c*(m+1) + \text{a}*d*(n+1)) - (m+1)*(b*c - \text{a}*d)*(A*b - \text{a}*B - \text{b}*C)*\text{Tan}[\text{e} + \text{f}*x] - \text{d}*(\text{A}*b^2 - \text{a}*(\text{b}*B - \text{a}*C))*(m+n+2)*\text{Tan}[\text{e} + \text{f}*x]^2, \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}, \text{n}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{!(ILtQ}[\text{n}, -1] \ \&\& \ (\text{!IntegerQ}[\text{m}] \ || \ (\text{EqQ}[\text{c}, 0] \ \&\& \ \text{NeQ}[\text{a}, 0])))$
- rule 4134 $\text{Int}[((\text{A}_) + (\text{B}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)]) + (\text{C}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])^2/(((\text{a}_) + (\text{b}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])*((\text{c}_) + (\text{d}_)*\text{tan}[(\text{e}_) + (\text{f}_)*(x_)])*(\text{x}_))), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{a}*(\text{A}*c - \text{c}*C + \text{B}*d) + \text{b}*(\text{B}*c - \text{A}*d + \text{C}*d))*(\text{x}/((\text{a}^2 + \text{b}^2)*(c^2 + d^2))), \text{x}] + (\text{Simp}[(\text{A}*b^2 - \text{a}*b*B + \text{a}^2*C)/((b*c - \text{a}*d)*(a^2 + b^2)) \quad \text{Int}[(\text{b} - \text{a}*\text{Tan}[\text{e} + \text{f}*x])/(a + \text{b}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] - \text{Simp}[(\text{c}^2*C - \text{B}*c*d + \text{A}*d^2)/((b*c - \text{a}*d)*(c^2 + d^2)) \quad \text{Int}[(\text{d} - \text{c}*\text{Tan}[\text{e} + \text{f}*x])/(c + \text{d}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0]$

Maple [A] (verified)

Time = 1.43 (sec) , antiderivative size = 647, normalized size of antiderivative = 1.36

method	result
derivativedivides	$\frac{(-A a^3 d - 3A a^2 b c + 3A a b^2 d + A b^3 c + B a^3 c - 3B a^2 b d - 3B a b^2 c + B b^3 d + a^3 C d + 3C a^2 b c - 3C a b^2 d - C b^3 c) \ln(1 + \tan(fx + e)^2)}{2} + \frac{(A a^2 + b^2)^3}{(a^2 + b^2)^3 (c^2)}$
default	$\frac{(-A a^3 d - 3A a^2 b c + 3A a b^2 d + A b^3 c + B a^3 c - 3B a^2 b d - 3B a b^2 c + B b^3 d + a^3 C d + 3C a^2 b c - 3C a b^2 d - C b^3 c) \ln(1 + \tan(fx + e)^2)}{2} + \frac{(A a^2 + b^2)^3}{(a^2 + b^2)^3 (c^2)}$
norman	Expression too large to display
parallelrisc	Expression too large to display
risc	Expression too large to display

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e)),x,
method=_RETURNVERBOSE)
```

output

```
1/f*(1/(a^2+b^2)^3/(c^2+d^2))*(1/2*(-A*a^3*d-3*A*a^2*b*c+3*A*a*b^2*d+A*b^3*c+
B*a^3*c-3*B*a^2*b*d-3*B*a*b^2*c+B*b^3*d+C*a^3*d+3*C*a^2*b*c-3*C*a*b^2*d-
C*b^3*c)*ln(1+tan(f*x+e)^2)+(A*a^3*c-3*A*a^2*b*d-3*A*a*b^2*c+A*b^3*d+B*a^3
*d+3*B*a^2*b*c-3*B*a*b^2*d-B*b^3*c-C*a^3*c+3*C*a^2*b*d+3*C*a*b^2*c-C*b^3*d
)*arctan(tan(f*x+e)))+(3*A*a^2*b^2*d-2*A*a*b^3*c+A*b^4*d-2*B*a^3*b*d+B*a^2
*b^2*c-B*b^4*c+C*a^4*d-C*a^2*b^2*d+2*C*a*b^3*c)/(a*d-b*c)^2/(a^2+b^2)^2/(a
+b*tan(f*x+e))-(6*A*a^4*b^2*d^2-8*A*a^3*b^3*c*d+3*A*a^2*b^4*c^2+3*A*a^2*b^
4*d^2-A*b^6*c^2+A*b^6*d^2-3*B*a^5*b*d^2+3*B*a^4*b^2*c*d-B*a^3*b^3*c^2+B*a^
3*b^3*d^2-6*B*a^2*b^4*c*d+3*B*a*b^5*c^2-B*b^6*c*d+C*a^6*d^2-3*C*a^4*b^2*d^
2+8*C*a^3*b^3*c*d-3*C*a^2*b^4*c^2+C*b^6*c^2)/(a*d-b*c)^3/(a^2+b^2)^3*ln(a+
b*tan(f*x+e))+1/2*(A*b^2-B*a*b+C*a^2)/(a*d-b*c)/(a^2+b^2)/(a+b*tan(f*x+e))
^2+(A*d^2-B*c*d+C*c^2)*d^2/(c^2+d^2)/(a*d-b*c)^3*ln(c+d*tan(f*x+e))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3643 vs. $2(475) = 950$.

Time = 3.06 (sec) , antiderivative size = 3643, normalized size of antiderivative = 7.64

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e)),x, algorithm="fricas")`

output Too large to include

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx$$

= Exception raised: NotImplementedError

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**3/(c+d*tan(f*x+e)),x)`

output Exception raised: NotImplementedError >> no valid subset found

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1096 vs. $2(475) = 950$.

Time = 0.17 (sec) , antiderivative size = 1096, normalized size of antiderivative = 2.30

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e)),x, algorithm="maxima")
```

output

```
1/2*(2*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c + (B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*d)*(f*x + e)/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2) - 2*((B*a^3*b^3 - 3*(A - C)*a^2*b^4 - 3*B*a*b^5 + (A - C)*b^6)*c^2 - (3*B*a^4*b^2 - 8*(A - C)*a^3*b^3 - 6*B*a^2*b^4 - B*b^6)*c*d - (C*a^6 - 3*B*a^5*b + 3*(2*A - C)*a^4*b^2 + B*a^3*b^3 + 3*A*a^2*b^4 + A*b^6)*d^2)*log(b*tan(f*x + e) + a)/((a^6*b^3 + 3*a^4*b^5 + 3*a^2*b^7 + b^9)*c^3 - 3*(a^7*b^2 + 3*a^5*b^4 + 3*a^3*b^6 + a*b^8)*c^2*d + 3*(a^8*b + 3*a^6*b^3 + 3*a^4*b^5 + a^2*b^7)*c*d^2 - (a^9 + 3*a^7*b^2 + 3*a^5*b^4 + a^3*b^6)*d^3) - 2*(C*c^2*d^2 - B*c*d^3 + A*d^4)*log(d*tan(f*x + e) + c)/(b^3*c^5 - 3*a*b^2*c^4*d + 3*a^2*b*c*d^4 - a^3*d^5 + (3*a^2*b + b^3)*c^3*d^2 - (a^3 + 3*a*b^2)*c^2*d^3) + ((B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c - ((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*d)*log(tan(f*x + e)^2 + 1)/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2) - ((C*a^4*b - 3*B*a^3*b^2 + (5*A - 3*C)*a^2*b^3 + B*a*b^4 + A*b^5)*c - (3*C*a^5 - 5*B*a^4*b + (7*A - C)*a^3*b^2 - B*a^2*b^3 + 3*A*a*b^4)*d - 2*((B*a^2*b^3 - 2*(A - C)*a*b^4 - B*b^5)*c + (C*a^4*b - 2*B*a^3*b^2 + (3*A - C)*a^2*b^3 + A*b^5)*d)*tan(f*x + e))/((a^6*b^2 + 2*a^4*b^4 + a^2*b^6)*c^2 - 2*(a^7*b + 2*a^5*b^3 + a^3*b^5)*c*d + (a^8 + 2*a^6*b^2 + a^4*b^4)*d^2 + ((a^4*b^4 + 2*a^2*b^6 + b^8)*c^2 - 2*(a^5*b^3 + 2*a^3*b^5 + a*b^7)*c*d + (a^6*b^2 ...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1447 vs. $2(475) = 950$.

Time = 0.81 (sec) , antiderivative size = 1447, normalized size of antiderivative = 3.03

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e)),x, algorithm="giac")
```

output

```
(A*a^3*c - C*a^3*c + 3*B*a^2*b*c - 3*A*a*b^2*c + 3*C*a*b^2*c - B*b^3*c + B
*a^3*d - 3*A*a^2*b*d + 3*C*a^2*b*d - 3*B*a*b^2*d + A*b^3*d - C*b^3*d)*(f*x
+ e)/(a^6*c^2*f + 3*a^4*b^2*c^2*f + 3*a^2*b^4*c^2*f + b^6*c^2*f + a^6*d^2
*f + 3*a^4*b^2*d^2*f + 3*a^2*b^4*d^2*f + b^6*d^2*f) + 1/2*(B*a^3*c - 3*A*a
^2*b*c + 3*C*a^2*b*c - 3*B*a*b^2*c + A*b^3*c - C*b^3*c - A*a^3*d + C*a^3*d
- 3*B*a^2*b*d + 3*A*a*b^2*d - 3*C*a*b^2*d + B*b^3*d)*log(tan(f*x + e)^2 +
1)/(a^6*c^2*f + 3*a^4*b^2*c^2*f + 3*a^2*b^4*c^2*f + b^6*c^2*f + a^6*d^2*f
+ 3*a^4*b^2*d^2*f + 3*a^2*b^4*d^2*f + b^6*d^2*f) - (B*a^3*b^4*c^2 - 3*A*a
^2*b^5*c^2 + 3*C*a^2*b^5*c^2 - 3*B*a*b^6*c^2 + A*b^7*c^2 - C*b^7*c^2 - 3*B
*a^4*b^3*c*d + 8*A*a^3*b^4*c*d - 8*C*a^3*b^4*c*d + 6*B*a^2*b^5*c*d + B*b^7
*c*d - C*a^6*b*d^2 + 3*B*a^5*b^2*d^2 - 6*A*a^4*b^3*d^2 + 3*C*a^4*b^3*d^2 -
B*a^3*b^4*d^2 - 3*A*a^2*b^5*d^2 - A*b^7*d^2)*log(abs(b*tan(f*x + e) + a))
/(a^6*b^4*c^3*f + 3*a^4*b^6*c^3*f + 3*a^2*b^8*c^3*f + b^10*c^3*f - 3*a^7*b
^3*c^2*d*f - 9*a^5*b^5*c^2*d*f - 9*a^3*b^7*c^2*d*f - 3*a*b^9*c^2*d*f + 3*a
^8*b^2*c*d^2*f + 9*a^6*b^4*c*d^2*f + 9*a^4*b^6*c*d^2*f + 3*a^2*b^8*c*d^2*f
- a^9*b*d^3*f - 3*a^7*b^3*d^3*f - 3*a^5*b^5*d^3*f - a^3*b^7*d^3*f) - (C*c
^2*d^3 - B*c*d^4 + A*d^5)*log(abs(d*tan(f*x + e) + c))/(b^3*c^5*d*f - 3*a*
b^2*c^4*d^2*f + 3*a^2*b*c^3*d^3*f + b^3*c^3*d^3*f - a^3*c^2*d^4*f - 3*a*b^
2*c^2*d^4*f + 3*a^2*b*c*d^5*f - a^3*d^6*f) - 1/2*(C*a^6*b^2*c^2 - 3*B*a^5*
b^3*c^2 + 5*A*a^4*b^4*c^2 - 2*C*a^4*b^4*c^2 - 2*B*a^3*b^5*c^2 + 6*A*a^2...
```

Mupad [B] (verification not implemented)

Time = 19.64 (sec) , antiderivative size = 65819, normalized size of antiderivative = 137.99

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^3*(c + d
*tan(e + f*x))),x)
```

output

```

-(((A*b^5*c - 3*C*a^5*d - 3*A*a*b^4*d + B*a*b^4*c + 5*B*a^4*b*d + C*a^4*b*c + 5*A*a^2*b^3*c - 7*A*a^3*b^2*d - 3*B*a^3*b^2*c + B*a^2*b^3*d - 3*C*a^2*b^3*c + C*a^3*b^2*d)/(2*(a^2*d^2 + b^2*c^2 - 2*a*b*c*d)*(a^4 + b^4 + 2*a^2*b^2)) - (tan(e + f*x)*(A*b^5*d - B*b^5*c - 2*A*a*b^4*c + 2*C*a*b^4*c + C*a^4*b*d + 3*A*a^2*b^3*d + B*a^2*b^3*c - 2*B*a^3*b^2*d - C*a^2*b^3*d))/((a^2*d^2 + b^2*c^2 - 2*a*b*c*d)*(a^4 + b^4 + 2*a^2*b^2)))/(a^2 + b^2*tan(e + f*x)^2 + 2*a*b*tan(e + f*x)) - symsum(log(- (A^3*b^8*c^2*d^4 - 4*A^3*a^2*b^6*d^6 - 7*A^3*a^4*b^4*d^6 - A^3*b^8*d^6 + A^2*C*b^8*d^6 - 3*A^3*a^2*b^6*c^2*d^4 - B^3*a^3*b^5*c^2*d^4 - C^3*a^2*b^6*c^2*d^4 - 2*C^3*a^3*b^5*c^3*d^3 + 7*C^3*a^4*b^4*c^2*d^4 + A^2*B*a*b^7*d^6 + A^2*B*b^8*c*d^5 + A^3*a*b^7*c*d^5 + C^3*a^7*b*c*d^5 - A*B^2*a^2*b^6*d^6 - 3*A*B^2*a^6*b^2*d^6 + 2*A^2*B*a^3*b^5*d^6 + 9*A^2*B*a^5*b^3*d^6 - A*C^2*a^2*b^6*d^6 - 4*A*C^2*a^4*b^4*d^6 + A*C^2*a^6*b^2*d^6 + 5*A^2*C*a^2*b^6*d^6 + 11*A^2*C*a^4*b^4*d^6 - A^2*C*a^6*b^2*d^6 + A*C^2*b^8*c^2*d^4 - 2*A^2*C*b^8*c^2*d^4 - B*C^2*b^8*c^3*d^3 + B^2*C*b^8*c^2*d^4 + 9*A^3*a^3*b^5*c*d^5 - B^3*a*b^7*c^2*d^4 + B^3*a^2*b^6*c*d^5 + B^3*a^4*b^4*c*d^5 + 2*C^3*a*b^7*c^3*d^3 - 3*C^3*a^5*b^3*c*d^5 + A*B*C*a^7*b*d^6 - 2*A*B*C*b^8*c*d^5 + 3*A*B^2*a^2*b^6*c^2*d^4 - A*B^2*a^4*b^4*c^2*d^4 + 3*A^2*B*a^3*b^5*c^2*d^4 - A*C^2*a^2*b^6*c^2*d^4 + 4*A*C^2*a^3*b^5*c^3*d^3 - 14*A*C^2*a^4*b^4*c^2*d^4 + 5*A^2*C*a^2*b^6*c^2*d^4 - 2*A^2*C*a^3*b^5*c^3*d^3 + 7*A^2*C*a^4*b^4*c^2*d^4 + 6*B*C^2*a^2*b^6*c^3*d^...

```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 8368, normalized size of antiderivative = 17.54

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e)),x)
```

output

```
( - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**8*b**2*d**4 + log(tan(e +
f*x)**2 + 1)*tan(e + f*x)**2*a**7*b**3*c*d**3 + log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**2*a**7*b**2*c*d**4 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*
**2*a**6*b**4*c**2*d**2 - 5*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b
**5*c**3*d - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**5*c*d**3 -
6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**4*c**3*d**2 - 3*log(ta
n(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**4*c*d**4 + 2*log(tan(e + f*x)**
2 + 1)*tan(e + f*x)**2*a**4*b**6*c**4 + 6*log(tan(e + f*x)**2 + 1)*tan(e +
f*x)**2*a**4*b**6*c**2*d**2 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*
**4*b**6*d**4 + 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*b**5*c**4*d
+ 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*b**5*c**2*d**3 - 6*log(
tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**7*c**3*d - 3*log(tan(e + f*x)
**2 + 1)*tan(e + f*x)**2*a**3*b**7*c*d**3 - 3*log(tan(e + f*x)**2 + 1)*tan
(e + f*x)**2*a**3*b**6*c**5 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a
**3*b**6*c**3*d**2 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**8*
c**4 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**8*c**2*d**2 - lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**9*c**3*d + log(tan(e + f*x)**2
+ 1)*tan(e + f*x)**2*a*b**8*c**5 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x
)*a**9*b*d**4 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**8*b**2*c*d**3 +
2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**8*b*c*d**4 + 6*log(tan(e + ...
```


3.77
$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$$

Optimal result	852
Mathematica [C] (verified)	853
Rubi [A] (verified)	854
Maple [A] (verified)	859
Fricas [B] (verification not implemented)	860
Sympy [C] (verification not implemented)	861
Maxima [A] (verification not implemented)	862
Giac [A] (verification not implemented)	863
Mupad [B] (verification not implemented)	864
Reduce [B] (verification not implemented)	865

Optimal result

Integrand size = 45, antiderivative size = 579

$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx =$$

$$- \frac{(a^3(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - 3ab^2(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - 3a^2b(2c(A - C)d - B(c^2 - d^2)))}{(c^2 + d^2)^2}$$

$$+ \frac{(3a^2b(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - b^3(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) + a^3(2c(A - C)d - B(c^2 - d^2)))}{(c^2 + d^2)^2 f}$$

$$+ \frac{(bc - ad)^2 (b(3c^4C - 2Bc^3d + c^2(A + 5C)d^2 - 4Bcd^3 + 3Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2))) \log}{d^4 (c^2 + d^2)^2 f}$$

$$+ \frac{b^2(ad(3c^2C - Bcd + (A + 2C)d^2) - b(3c^3C - 2Bc^2d + c(A + 2C)d^2 - Bd^3)) \tan(e+fx)}{d^3 (c^2 + d^2) f}$$

$$+ \frac{b(3c^2C - 2Bcd + (2A + C)d^2) (a + b \tan(e+fx))^2}{2d^2 (c^2 + d^2) f}$$

$$- \frac{(c^2C - Bcd + Ad^2) (a + b \tan(e+fx))^3}{d (c^2 + d^2) f (c + d \tan(e+fx))}$$

output

```

-(a^3*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-3*a*b^2*(c^2*C-2*B*c*d-C*d^2-A*(c^
2-d^2))-3*a^2*b*(2*c*(A-C)*d-B*(c^2-d^2))+b^3*(2*c*(A-C)*d-B*(c^2-d^2)))*x
/(c^2+d^2)^2+(3*a^2*b*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-b^3*(c^2*C-2*B*c*d
-C*d^2-A*(c^2-d^2))+a^3*(2*c*(A-C)*d-B*(c^2-d^2))-3*a*b^2*(2*c*(A-C)*d-B*(
c^2-d^2))*ln(cos(f*x+e))/(c^2+d^2)^2/f+(-a*d+b*c)^2*(b*(3*c^4*C-2*B*c^3*d
+c^2*(A+5*C)*d^2-4*B*c*d^3+3*A*d^4)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*ln(c+
d*tan(f*x+e))/d^4/(c^2+d^2)^2/f+b^2*(a*d*(3*c^2*C-B*c*d+(A+2*C)*d^2)-b*(3*
c^3*C-2*B*c^2*d+c*(A+2*C)*d^2-B*d^3))*tan(f*x+e)/d^3/(c^2+d^2)/f+1/2*b*(3*
c^2*C-2*B*c*d+(2*A+C)*d^2)*(a+b*tan(f*x+e))^2/d^2/(c^2+d^2)/f-(A*d^2-B*c*d
+C*c^2)*(a+b*tan(f*x+e))^3/d/(c^2+d^2)/f/(c+d*tan(f*x+e))

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.96 (sec) , antiderivative size = 1022, normalized size of antiderivative = 1.77

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{C(a + b \tan(e + fx))^3}{2df(c + d \tan(e + fx))} + \frac{(-3bcC + 2bBd + 3aCd)(a + b \tan(e + fx))^2}{df(c + d \tan(e + fx))} + \frac{2 \left(-\frac{d^2(-3a^2Abc^2 + Ab^3c^2 - a^3Bc^2 + 3ab^2Bc^2 + 3a^2bc^2C - b^3c^2C + 2a^3Acd - 6aAb^2cd - 6a^2bBcd + 2b^3Bcd)}{df(c + d \tan(e + fx))} \right)}{df(c + d \tan(e + fx))}$$

input

```

Integrate[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^2,x]

```

output

```
(C*(a + b*Tan[e + f*x])^3)/(2*d*f*(c + d*Tan[e + f*x])) + (((-3*b*c*C + 2*
b*B*d + 3*a*C*d)*(a + b*Tan[e + f*x])^2)/(d*f*(c + d*Tan[e + f*x])) + (2*(
-1/2*(d^2*(-3*a^2*A*b*c^2 + A*b^3*c^2 - a^3*B*c^2 + 3*a*b^2*B*c^2 + 3*a^2*
b*c^2*C - b^3*c^2*C + 2*a^3*A*c*d - 6*a*A*b^2*c*d - 6*a^2*b*B*c*d + 2*b^3*
B*c*d - 2*a^3*c*C*d + 6*a*b^2*c*C*d + 3*a^2*A*b*d^2 - A*b^3*d^2 + a^3*B*d^
2 - 3*a*b^2*B*d^2 - 3*a^2*b*C*d^2 + b^3*C*d^2 + I*(a^3*A*c^2 - 3*a*A*b^2*c
^2 - 3*a^2*b*B*c^2 + b^3*B*c^2 - a^3*c^2*C + 3*a*b^2*c^2*C + 6*a^2*A*b*c*d
- 2*A*b^3*c*d + 2*a^3*B*c*d - 6*a*b^2*B*c*d - 6*a^2*b*c*C*d + 2*b^3*c*C*d
- a^3*A*d^2 + 3*a*A*b^2*d^2 + 3*a^2*b*B*d^2 - b^3*B*d^2 + a^3*C*d^2 - 3*a
*b^2*C*d^2))*Log[I - Tan[e + f*x]])/(c^2 + d^2)^2*f) + (d^2*(3*a^2*A*b*c^
2 - A*b^3*c^2 + a^3*B*c^2 - 3*a*b^2*B*c^2 - 3*a^2*b*c^2*C + b^3*c^2*C - 2*
a^3*A*c*d + 6*a*A*b^2*c*d + 6*a^2*b*B*c*d - 2*b^3*B*c*d + 2*a^3*c*C*d - 6*
a*b^2*c*C*d - 3*a^2*A*b*d^2 + A*b^3*d^2 - a^3*B*d^2 + 3*a*b^2*B*d^2 + 3*a^
2*b*C*d^2 - b^3*C*d^2 + I*(a^3*A*c^2 - 3*a*A*b^2*c^2 - 3*a^2*b*B*c^2 + b^3
*B*c^2 - a^3*c^2*C + 3*a*b^2*c^2*C + 6*a^2*A*b*c*d - 2*A*b^3*c*d + 2*a^3*B
*c*d - 6*a*b^2*B*c*d - 6*a^2*b*c*C*d + 2*b^3*c*C*d - a^3*A*d^2 + 3*a*A*b^2
*d^2 + 3*a^2*b*B*d^2 - b^3*B*d^2 + a^3*C*d^2 - 3*a*b^2*C*d^2))*Log[I + Tan
[e + f*x]])/(2*(c^2 + d^2)^2*f) + ((b*c - a*d)^2*(b*(3*c^4*C - 2*B*c^3*d +
c^2*(A + 5*C)*d^2 - 4*B*c*d^3 + 3*A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2
- d^2)))*Log[c + d*Tan[e + f*x]])/(d^2*(c^2 + d^2)^2*f) - ((b*c - a*d)^...
```

Rubi [A] (verified)

Time = 5.36 (sec) , antiderivative size = 594, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4128, 3042, 4130, 27, 3042, 4120, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^2} dx$$

↓ 4128

$$\int \frac{(a+b \tan(e+fx))^2 (b(3C^2-2Bdc+(2A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{c+d \tan(e+fx)} d$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^2 (b(3C^2-2Bdc+(2A+C)d^2) \tan(e+fx)^2+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{c+d \tan(e+fx)} d$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{}$$

↓ 4130

$$\int -\frac{2(a+b \tan(e+fx))(c(3C^2-2Bdc+(2A+C)d^2)b^2-(ad(3C^2-Bdc+(A+2C)d^2)-b(3C^3-2Bdc^2+(A+2C)d^2c-Bd^3)) \tan^2(e+fx)b-ad(Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{c+d \tan(e+fx)} d$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{}$$

↓ 27

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \int \frac{(a+b \tan(e+fx))(c(3C^2-2Bdc+(2A+C)d^2)b^2-(ad(3C^2-Bdc+(A+2C)d^2)-b(3C^3-2Bdc^2+(A+2C)d^2c-Bd^3)) \tan^2(e+fx)b-ad(Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{c+d \tan(e+fx)} d$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{}$$

↓ 3042

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \int \frac{(a+b \tan(e+fx))(c(3C^2-2Bdc+(2A+C)d^2)b^2-(ad(3C^2-Bdc+(A+2C)d^2)-b(3C^3-2Bdc^2+(A+2C)d^2c-Bd^3)) \tan^2(e+fx)b-ad(Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{c+d \tan(e+fx)} d$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{}$$

↓ 4120

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{\int \frac{c(3Cc^3-2Bdc^2+(A+2C)d^2c-Bd^3)b^3-3acd(2Cc^2-Bdc+(A+C)d^2)b^2+(c^2+d^2)((3Cc^2-2Bdc+3c^2C)(a+b \tan(e+fx))^2)}{c^2+d^2} dx}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 3042

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{\int \frac{c(3Cc^3-2Bdc^2+(A+2C)d^2c-Bd^3)b^3-3acd(2Cc^2-Bdc+(A+C)d^2)b^2+(c^2+d^2)((3Cc^2-2Bdc+3c^2C)(a+b \tan(e+fx))^2)}{c^2+d^2} dx}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 4109

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{d^3(a^3(2cd(A-C)-B(c^2-d^2))+3a^2b(-A(c^2-d^2)-2Bcd+c^2C-Cd^2))-3ab^2(2cd(A-C)-B(c^2-d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 3042

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{d^3(a^3(2cd(A-C)-B(c^2-d^2))+3a^2b(-A(c^2-d^2)-2Bcd+c^2C-Cd^2))-3ab^2(2cd(A-C)-B(c^2-d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 3956

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{(bc-ad)^2(ad^2(2cd(A-C)-B(c^2-d^2))+b(c^2d^2(A+5C)+3Ad^4-2Bc^3d-4Bcd^3+3c^4C)) \int \frac{\tan(e+fx)}{c+d \tan(e+fx)} dx}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 4100

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{(bc-ad)^2(ad^2(2cd(A-C)-B(c^2-d^2))+b(c^2d^2(A+5C)+3Ad^4-2Bc^3d-4Bcd^3+3c^4C))f}{df(c^2+d^2)} - \frac{1}{c+d \tan(e+fx)}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 16

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)(a+b \tan(e+fx))^2}{2df} - \frac{d^3 \log(\cos(e+fx))(a^3(2cd(A-C)-B(c^2-d^2))+3a^2b(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)-3ab^2(2cd(A-C)-B(c^2-d^2)+c^2d^2))}{f(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

input

```
Int[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]^2,x]
```

output

```
-(((c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^3)/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))) + ((b*(3*c^2*C - 2*B*c*d + (2*A + C)*d^2)*(a + b*Tan[e + f*x])^2)/(2*d*f) - (((-((d^3*(a^3*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - 3*a*b^2*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - 3*a^2*b*(2*c*(A - C)*d - B*(c^2 - d^2)) + b^3*(2*c*(A - C)*d - B*(c^2 - d^2))))*x)/(c^2 + d^2)) + (d^3*(3*a^2*b*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b^3*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + a^3*(2*c*(A - C)*d - B*(c^2 - d^2)) - 3*a*b^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Log[Cos[e + f*x]]/(c^2 + d^2)*f) + ((b*c - a*d)^2*(b*(3*c^4*C - 2*B*c^3*d + c^2*(A + 5*C)*d^2 - 4*B*c*d^3 + 3*A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Log[c + d*Tan[e + f*x]]/(d*(c^2 + d^2)*f))/d - (b^2*(a*d*(3*c^2*C - B*c*d + (A + 2*C)*d^2) - b*(3*c^3*C - 2*B*c^2*d + c*(A + 2*C)*d^2 - B*d^3))*Tan[e + f*x]/(d*f))/d)/(d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4100 $\text{Int}[((a_)+(b_)*\tan[(e_)+(f_)*(x_)])^{(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x_)])^2}, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \&\& \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[((A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)/((a_)+(b_)*\tan[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A*b - a*B - b*C, 0]$
- rule 4120 $\text{Int}[((a_)+(b_)*\tan[(e_)+(f_)*(x_)])^{(c_)+(d_)*\tan[(e_)+(f_)*(x_)]^{(n_)*((A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[b*C*\text{Tan}[e + f*x]*((c + d*\text{Tan}[e + f*x])^{(n + 1)/(d*f*(n + 2))}), x] - \text{Simp}[1/(d*(n + 2)) \text{ Int}[(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*\text{Tan}[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!LtQ}[n, -1]$

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 829, normalized size of antiderivative = 1.43

method	result
derivativedivides	$\frac{b^2 \left(\frac{\tan^2(fx+e) C b d}{2} + B \tan(fx+e) b d + 3 d \tan(fx+e) C a - 2 \tan(fx+e) C b c \right)}{d^3} + \frac{(-2 A a^3 c d + 3 A a^2 b c^2 - 3 A a^2 b d^2 + 6 A a b^2 c d - A b^3 c^2)}{d^3}$
default	$\frac{b^2 \left(\frac{\tan^2(fx+e) C b d}{2} + B \tan(fx+e) b d + 3 d \tan(fx+e) C a - 2 \tan(fx+e) C b c \right)}{d^3} + \frac{(-2 A a^3 c d + 3 A a^2 b c^2 - 3 A a^2 b d^2 + 6 A a b^2 c d - A b^3 c^2)}{d^3}$
norman	Expression too large to display
parallelrisc	Expression too large to display
risc	Expression too large to display

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,
x,method=_RETURNVERBOSE)
```

output

```
1/f*(b^2/d^3*(1/2*tan(f*x+e)^2*C*b*d+B*tan(f*x+e)*b*d+3*d*tan(f*x+e)*C*a-2
*tan(f*x+e)*C*b*c)+1/(c^2+d^2)^2*(1/2*(-2*A*a^3*c*d+3*A*a^2*b*c^2-3*A*a^2*
b*d^2+6*A*a*b^2*c*d-A*b^3*c^2+A*b^3*d^2+B*a^3*c^2-B*a^3*d^2+6*B*a^2*b*c*d-
3*B*a*b^2*c^2+3*B*a*b^2*d^2-2*B*b^3*c*d+2*C*a^3*c*d-3*C*a^2*b*c^2+3*C*a^2*
b*d^2-6*C*a*b^2*c*d+C*b^3*c^2-C*b^3*d^2)*ln(1+tan(f*x+e)^2)+(A*a^3*c^2-A*a
^3*d^2+6*A*a^2*b*c*d-3*A*a*b^2*c^2+3*A*a*b^2*d^2-2*A*b^3*c*d+2*B*a^3*c*d-3
*B*a^2*b*c^2+3*B*a^2*b*d^2-6*B*a*b^2*c*d+B*b^3*c^2-B*b^3*d^2-C*a^3*c^2+C*a
^3*d^2-6*C*a^2*b*c*d+3*C*a*b^2*c^2-3*C*a*b^2*d^2+2*C*b^3*c*d)*arctan(tan(f
*x+e))-1/d^4*(A*a^3*d^5-3*A*a^2*b*c*d^4+3*A*a*b^2*c^2*d^3-A*b^3*c^3*d^2-B
*a^3*c*d^4+3*B*a^2*b*c^2*d^3-3*B*a*b^2*c^3*d^2+B*b^3*c^4*d+C*a^3*c^2*d^3-3
*C*a^2*b*c^3*d^2+3*C*a*b^2*c^4*d-C*b^3*c^5)/(c^2+d^2)/(c+d*tan(f*x+e))+1/d
^4*(2*A*a^3*c*d^5-3*A*a^2*b*c^2*d^4+3*A*a^2*b*d^6-6*A*a*b^2*c*d^5+A*b^3*c^
4*d^2+3*A*b^3*c^2*d^4-B*a^3*c^2*d^4+B*a^3*d^6-6*B*a^2*b*c*d^5+3*B*a*b^2*c^
4*d^2+9*B*a*b^2*c^2*d^4-2*B*b^3*c^5*d-4*B*b^3*c^3*d^3-2*C*a^3*c*d^5+3*C*a^
2*b*c^4*d^2+9*C*a^2*b*c^2*d^4-6*C*a*b^2*c^5*d-12*C*a*b^2*c^3*d^3+3*C*b^3*c
^6+5*C*b^3*c^4*d^2)/(c^2+d^2)^2*ln(c+d*tan(f*x+e)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1477 vs. $2(577) = 1154$.

Time = 0.90 (sec) , antiderivative size = 1477, normalized size of antiderivative = 2.55

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^2,x, algorithm="fricas")
```

output

```

1/2*(3*C*b^3*c^5*d^2 - 2*A*a^3*d^7 - 2*(3*C*a*b^2 + B*b^3)*c^4*d^3 + 2*(3*
C*a^2*b + 3*B*a*b^2 + (A + C)*b^3)*c^3*d^4 - 2*(C*a^3 + 3*B*a^2*b + 3*A*a*
b^2)*c^2*d^5 + (2*B*a^3 + 6*A*a^2*b + C*b^3)*c*d^6 + (C*b^3*c^4*d^3 + 2*C*
b^3*c^2*d^5 + C*b^3*d^7)*tan(f*x + e)^3 + 2*(((A - C)*a^3 - 3*B*a^2*b - 3*
(A - C)*a*b^2 + B*b^3)*c^3*d^4 + 2*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 -
(A - C)*b^3)*c^2*d^5 - ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)
*c*d^6)*f*x - (3*C*b^3*c^5*d^2 + 6*C*b^3*c^3*d^4 + 3*C*b^3*c*d^6 - 2*(3*C*
a*b^2 + B*b^3)*c^4*d^3 - 4*(3*C*a*b^2 + B*b^3)*c^2*d^5 - 2*(3*C*a*b^2 + B*
b^3)*d^7)*tan(f*x + e)^2 + (3*C*b^3*c^7 - 2*(3*C*a*b^2 + B*b^3)*c^6*d + (3
*C*a^2*b + 3*B*a*b^2 + (A + 5*C)*b^3)*c^5*d^2 - 4*(3*C*a*b^2 + B*b^3)*c^4*
d^3 - (B*a^3 + 3*(A - 3*C)*a^2*b - 9*B*a*b^2 - 3*A*b^3)*c^3*d^4 + 2*((A -
C)*a^3 - 3*B*a^2*b - 3*A*a*b^2)*c^2*d^5 + (B*a^3 + 3*A*a^2*b)*c*d^6 + (3*C
*b^3*c^6*d - 2*(3*C*a*b^2 + B*b^3)*c^5*d^2 + (3*C*a^2*b + 3*B*a*b^2 + (A +
5*C)*b^3)*c^4*d^3 - 4*(3*C*a*b^2 + B*b^3)*c^3*d^4 - (B*a^3 + 3*(A - 3*C)*
a^2*b - 9*B*a*b^2 - 3*A*b^3)*c^2*d^5 + 2*((A - C)*a^3 - 3*B*a^2*b - 3*A*a*
b^2)*c*d^6 + (B*a^3 + 3*A*a^2*b)*d^7)*tan(f*x + e))*log((d^2*tan(f*x + e)^
2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - (3*C*b^3*c^7 - 2*(3*
C*a*b^2 + B*b^3)*c^6*d + (3*C*a^2*b + 3*B*a*b^2 + (A + 5*C)*b^3)*c^5*d^2 -
4*(3*C*a*b^2 + B*b^3)*c^4*d^3 + (6*C*a^2*b + 6*B*a*b^2 + (2*A + C)*b^3)*c
^3*d^4 - 2*(3*C*a*b^2 + B*b^3)*c^2*d^5 + (3*C*a^2*b + 3*B*a*b^2 + (A - ...

```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 28.69 (sec) , antiderivative size = 24300, normalized size of antiderivative = 41.97

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((a+b*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*
x+e))**2,x)

```

output

```
Piecewise((zoo*x*(a + b*tan(e))**3*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2,
Eq(c, 0) & Eq(d, 0) & Eq(f, 0)), ((A*a**3*x + 3*A*a**2*b*log(tan(e + f*x)
**2 + 1)/(2*f) - 3*A*a*b**2*x + 3*A*a*b**2*tan(e + f*x)/f - A*b**3*log(tan
(e + f*x)**2 + 1)/(2*f) + A*b**3*tan(e + f*x)**2/(2*f) + B*a**3*log(tan(e
+ f*x)**2 + 1)/(2*f) - 3*B*a**2*b*x + 3*B*a**2*b*tan(e + f*x)/f - 3*B*a*b*
**2*log(tan(e + f*x)**2 + 1)/(2*f) + 3*B*a*b**2*tan(e + f*x)**2/(2*f) + B*b
**3*x + B*b**3*tan(e + f*x)**3/(3*f) - B*b**3*tan(e + f*x)/f - C*a**3*x +
C*a**3*tan(e + f*x)/f - 3*C*a**2*b*log(tan(e + f*x)**2 + 1)/(2*f) + 3*C*a*
**2*b*tan(e + f*x)**2/(2*f) + 3*C*a*b**2*x + C*a*b**2*tan(e + f*x)**3/f - 3
*C*a*b**2*tan(e + f*x)/f + C*b**3*log(tan(e + f*x)**2 + 1)/(2*f) + C*b**3*
tan(e + f*x)**4/(4*f) - C*b**3*tan(e + f*x)**2/(2*f))/c**2, Eq(d, 0)), (-A
*a**3*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f
*x) - 4*d**2*f) + 2*I*A*a**3*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 -
8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + A*a**3*f*x/(4*d**2*f*tan(e + f*x)**2
- 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - A*a**3*tan(e + f*x)/(4*d**2*f*tan
(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*A*a**3/(4*d**2*f*
tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 3*I*A*a**2*b*f*x*t
an(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**
2*f) + 6*A*a**2*b*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*
tan(e + f*x) - 4*d**2*f) - 3*I*A*a**2*b*f*x/(4*d**2*f*tan(e + f*x)**2 - ...
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 684, normalized size of antiderivative = 1.18

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^2,x, algorithm="maxima")
```

output

```

1/2*(2*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c^2 + 2*(B*a^3
+ 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c*d - ((A - C)*a^3 - 3*B*a^2
*b - 3*(A - C)*a*b^2 + B*b^3)*d^2)*(f*x + e)/(c^4 + 2*c^2*d^2 + d^4) + 2*(
3*C*b^3*c^6 - 2*(3*C*a*b^2 + B*b^3)*c^5*d + (3*C*a^2*b + 3*B*a*b^2 + (A +
5*C)*b^3)*c^4*d^2 - 4*(3*C*a*b^2 + B*b^3)*c^3*d^3 - (B*a^3 + 3*(A - 3*C)*a
^2*b - 9*B*a*b^2 - 3*A*b^3)*c^2*d^4 + 2*((A - C)*a^3 - 3*B*a^2*b - 3*A*a*b
^2)*c*d^5 + (B*a^3 + 3*A*a^2*b)*d^6)*log(d*tan(f*x + e) + c)/(c^4*d^4 + 2*
c^2*d^6 + d^8) + ((B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c^2
- 2*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c*d - (B*a^3 + 3*(
A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^2)*log(tan(f*x + e)^2 + 1)/(c^4
+ 2*c^2*d^2 + d^4) + 2*(C*b^3*c^5 - A*a^3*d^5 - (3*C*a*b^2 + B*b^3)*c^4*d
+ (3*C*a^2*b + 3*B*a*b^2 + A*b^3)*c^3*d^2 - (C*a^3 + 3*B*a^2*b + 3*A*a*b^2
)*c^2*d^3 + (B*a^3 + 3*A*a^2*b)*c*d^4)/(c^3*d^4 + c*d^6 + (c^2*d^5 + d^7)*
tan(f*x + e)) + (C*b^3*d*tan(f*x + e)^2 - 2*(2*C*b^3*c - (3*C*a*b^2 + B*b
^3)*d)*tan(f*x + e))/d^3)/f

```

Giac [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 1039, normalized size of antiderivative = 1.79

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^2,x, algorithm="giac")

```

output

```
(A*a^3*c^2 - C*a^3*c^2 - 3*B*a^2*b*c^2 - 3*A*a*b^2*c^2 + 3*C*a*b^2*c^2 + B
*b^3*c^2 + 2*B*a^3*c*d + 6*A*a^2*b*c*d - 6*C*a^2*b*c*d - 6*B*a*b^2*c*d - 2
*A*b^3*c*d + 2*C*b^3*c*d - A*a^3*d^2 + C*a^3*d^2 + 3*B*a^2*b*d^2 + 3*A*a*b
^2*d^2 - 3*C*a*b^2*d^2 - B*b^3*d^2)*(f*x + e)/(c^4*f + 2*c^2*d^2*f + d^4*f
) + 1/2*(B*a^3*c^2 + 3*A*a^2*b*c^2 - 3*C*a^2*b*c^2 - 3*B*a*b^2*c^2 - A*b^3
*c^2 + C*b^3*c^2 - 2*A*a^3*c*d + 2*C*a^3*c*d + 6*B*a^2*b*c*d + 6*A*a*b^2*c
*d - 6*C*a*b^2*c*d - 2*B*b^3*c*d - B*a^3*d^2 - 3*A*a^2*b*d^2 + 3*C*a^2*b*d
^2 + 3*B*a*b^2*d^2 + A*b^3*d^2 - C*b^3*d^2)*log(tan(f*x + e)^2 + 1)/(c^4*f
+ 2*c^2*d^2*f + d^4*f) + (3*C*b^3*c^6 - 6*C*a*b^2*c^5*d - 2*B*b^3*c^5*d +
3*C*a^2*b*c^4*d^2 + 3*B*a*b^2*c^4*d^2 + A*b^3*c^4*d^2 + 5*C*b^3*c^4*d^2 -
12*C*a*b^2*c^3*d^3 - 4*B*b^3*c^3*d^3 - B*a^3*c^2*d^4 - 3*A*a^2*b*c^2*d^4
+ 9*C*a^2*b*c^2*d^4 + 9*B*a*b^2*c^2*d^4 + 3*A*b^3*c^2*d^4 + 2*A*a^3*c*d^5
- 2*C*a^3*c*d^5 - 6*B*a^2*b*c*d^5 - 6*A*a*b^2*c*d^5 + B*a^3*d^6 + 3*A*a^2*
b*d^6)*log(abs(d*tan(f*x + e) + c))/(c^4*d^4*f + 2*c^2*d^6*f + d^8*f) + 1/
2*(C*b^3*d^2*f*tan(f*x + e)^2 - 4*C*b^3*c*d*f*tan(f*x + e) + 6*C*a*b^2*d^2
*f*tan(f*x + e) + 2*B*b^3*d^2*f*tan(f*x + e))/(d^4*f^2) + (C*b^3*c^7 - 3*C
*a*b^2*c^6*d - B*b^3*c^6*d + 3*C*a^2*b*c^5*d^2 + 3*B*a*b^2*c^5*d^2 + A*b^3
*c^5*d^2 + C*b^3*c^5*d^2 - C*a^3*c^4*d^3 - 3*B*a^2*b*c^4*d^3 - 3*A*a*b^2*c
^4*d^3 - 3*C*a*b^2*c^4*d^3 - B*b^3*c^4*d^3 + B*a^3*c^3*d^4 + 3*A*a^2*b*c^3
*d^4 + 3*C*a^2*b*c^3*d^4 + 3*B*a*b^2*c^3*d^4 + A*b^3*c^3*d^4 - A*a^3*c^...
```

Mupad [B] (verification not implemented)

Time = 12.48 (sec) , antiderivative size = 701, normalized size of antiderivative = 1.21

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{\tan(e + fx) \left(\frac{Bb^3 + 3Cab^2}{d^2} - \frac{2Cb^3c}{d^3} \right)}{f}$$

$$- \frac{\ln(\tan(e + fx) + 1i) (Ba^3 - Ab^3 + Cb^3 + 3Aa^2b - 3Ba^2b^2 - 3Ca^2b + Aa^3 1i + Bb^3 1i - Ca^3 1i)}{2f(-c^2 + cd 2i + d^2)}$$

$$+ \frac{\ln(c + d \tan(e + fx)) (d^4 (3Ab^3c^2 - Ba^3c^2 - 3Aa^2bc^2 + 9Bab^2c^2 + 9Ca^2bc^2) - d^5 (2Ca^3c - 3Ab^3c^2))}{d^4 (3Ab^3c^2 - Ba^3c^2 - 3Aa^2bc^2 + 9Bab^2c^2 + 9Ca^2bc^2) - d^5 (2Ca^3c - 3Ab^3c^2)}$$

$$- \frac{\ln(\tan(e + fx) - 1i) (Aa^3 - Ab^3 1i + Ba^3 1i + Bb^3 - Ca^3 + Cb^3 1i - 3Aab^2 + Aa^2b 3i - Bab^2 3i)}{2f(-c^2 1i + 2cd + d^2 1i)}$$

$$- \frac{Ca^3c^2d^3 - Ba^3cd^4 + Aa^3d^5 - 3Ca^2bc^3d^2 + 3Ba^2bc^2d^3 - 3Aa^2bcd^4 + 3Cab^2c^4d - 3Bab^2c^4d}{df(\tan(e + fx)d^4 + cd^3)(c^2 + d^2)}$$

$$+ \frac{Cb^3 \tan(e + fx)^2}{2d^2 f}$$

input `int(((a + b*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^2,x)`

output
$$\begin{aligned} & (\tan(e + f*x)*((B*b^3 + 3*C*a*b^2)/d^2 - (2*C*b^3*c)/d^3))/f - (\log(\tan(e + f*x) + 1i)*(A*a^3*1i - A*b^3 + B*a^3 + B*b^3*1i - C*a^3*1i + C*b^3 - A*a*b^2*3i + 3*A*a^2*b - 3*B*a*b^2 - B*a^2*b*3i + C*a*b^2*3i - 3*C*a^2*b))/(2*f*(c*d*2i - c^2 + d^2)) + (\log(c + d*tan(e + f*x))*(d^4*(3*A*b^3*c^2 - B*a^3*c^2 - 3*A*a^2*b*c^2 + 9*B*a*b^2*c^2 + 9*C*a^2*b*c^2) - d^5*(2*C*a^3*c - 2*A*a^3*c + 6*A*a*b^2*c + 6*B*a^2*b*c) - d^3*(4*B*b^3*c^3 + 12*C*a*b^2*c^3) + d^6*(B*a^3 + 3*A*a^2*b) - d*(2*B*b^3*c^5 + 6*C*a*b^2*c^5) + d^2*(A*b^3*c^4 + 5*C*b^3*c^4 + 3*B*a*b^2*c^4 + 3*C*a^2*b*c^4) + 3*C*b^3*c^6))/(f*(d^8 + 2*c^2*d^6 + c^4*d^4)) - (\log(\tan(e + f*x) - 1i)*(A*a^3 - A*b^3*1i + B*a^3*1i + B*b^3 - C*a^3 + C*b^3*1i - 3*A*a*b^2 + A*a^2*b*3i - B*a*b^2*3i - 3*B*a^2*b + 3*C*a*b^2 - C*a^2*b*3i))/(2*f*(2*c*d - c^2*1i + d^2*1i)) - (A*a^3*d^5 - C*b^3*c^5 - B*a^3*c*d^4 + B*b^3*c^4*d - A*b^3*c^3*d^2 + C*a^3*c^2*d^3 + 3*A*a*b^2*c^2*d^3 - 3*B*a*b^2*c^3*d^2 + 3*B*a^2*b*c^2*d^3 - 3*C*a^2*b*c^3*d^2 - 3*A*a^2*b*c*d^4 + 3*C*a*b^2*c^4*d)/(d*f*(c*d^3 + d^4*tan(e + f*x))*(c^2 + d^2)) + (C*b^3*tan(e + f*x)^2)/(2*d^2*f) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2547, normalized size of antiderivative = 4.40

$$\int \frac{(a + b \tan(e + f x))^3 (A + B \tan(e + f x) + C \tan^2(e + f x))}{(c + d \tan(e + f x))^2} dx = \text{Too large to display}$$

input `int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x)`

output

```
( - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**4*c**2*d**6 + 4*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a**3*b*c**3*d**5 - 4*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)*a**3*b*c*d**7 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*c
**3*d**6 + 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**2*c**2*d**6 -
3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b*c**4*d**5 + 3*log(tan(e + f
*x)**2 + 1)*tan(e + f*x)*a**2*b*c**2*d**7 - 4*log(tan(e + f*x)**2 + 1)*tan
(e + f*x)*a*b**3*c**3*d**5 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**
3*c*d**7 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**2*c**3*d**6 - 2*lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**4*c**2*d**6 + log(tan(e + f*x)**2 +
1)*tan(e + f*x)*b**3*c**4*d**5 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b*
**3*c**2*d**7 - 2*log(tan(e + f*x)**2 + 1)*a**4*c**3*d**5 + 4*log(tan(e + f
*x)**2 + 1)*a**3*b*c**4*d**4 - 4*log(tan(e + f*x)**2 + 1)*a**3*b*c**2*d**6
+ 2*log(tan(e + f*x)**2 + 1)*a**3*c**4*d**5 + 12*log(tan(e + f*x)**2 + 1)
*a**2*b**2*c**3*d**5 - 3*log(tan(e + f*x)**2 + 1)*a**2*b*c**5*d**4 + 3*log
(tan(e + f*x)**2 + 1)*a**2*b*c**3*d**6 - 4*log(tan(e + f*x)**2 + 1)*a*b**3
*c**4*d**4 + 4*log(tan(e + f*x)**2 + 1)*a*b**3*c**2*d**6 - 6*log(tan(e + f
*x)**2 + 1)*a*b**2*c**4*d**5 - 2*log(tan(e + f*x)**2 + 1)*b**4*c**3*d**5 +
log(tan(e + f*x)**2 + 1)*b**3*c**5*d**4 - log(tan(e + f*x)**2 + 1)*b**3*c
**3*d**6 + 4*log(tan(e + f*x)*d + c)*tan(e + f*x)*a**4*c**2*d**6 - 8*log(t
an(e + f*x)*d + c)*tan(e + f*x)*a**3*b*c**3*d**5 + 8*log(tan(e + f*x)*d...
```

3.78
$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$$

Optimal result	867
Mathematica [C] (verified)	868
Rubi [A] (verified)	868
Maple [A] (verified)	872
Fricas [B] (verification not implemented)	873
Sympy [C] (verification not implemented)	874
Maxima [A] (verification not implemented)	875
Giac [A] (verification not implemented)	876
Mupad [B] (verification not implemented)	877
Reduce [B] (verification not implemented)	878

Optimal result

Integrand size = 45, antiderivative size = 417

$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx =$$

$$-\frac{(a^2(c^2C-2Bcd-Cd^2-A(c^2-d^2))-b^2(c^2C-2Bcd-Cd^2-A(c^2-d^2))-2ab(2c(A-C)d-B(c^2-d^2)))}{(c^2+d^2)^2}$$

$$+\frac{(2ab(c^2C-2Bcd-Cd^2-A(c^2-d^2))+a^2(2c(A-C)d-B(c^2-d^2))-b^2(2c(A-C)d-B(c^2-d^2)))}{(c^2+d^2)^2 f}$$

$$-\frac{(bc-ad)(b(2c^4C-Bc^3d+4c^2Cd^2-3Bcd^3+2Ad^4)+ad^2(2c(A-C)d-B(c^2-d^2))) \log(c+d \tan(e+fx))}{d^3(c^2+d^2)^2 f}$$

$$+\frac{b^2(2c^2C-Bcd+(A+C)d^2) \tan(e+fx)}{d^2(c^2+d^2) f}$$

$$-\frac{(c^2C-Bcd+Ad^2)(a+b \tan(e+fx))^2}{d(c^2+d^2) f(c+d \tan(e+fx))}$$

output

```

-(a^2*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-b^2*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-2*a*b*(2*c*(A-C)*d-B*(c^2-d^2)))*x/(c^2+d^2)^2+(2*a*b*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))+a^2*(2*c*(A-C)*d-B*(c^2-d^2))-b^2*(2*c*(A-C)*d-B*(c^2-d^2)))*ln(cos(f*x+e))/(c^2+d^2)^2/f-((a*d+b*c)*(b*(2*A*d^4-B*c^3*d-3*B*c*d^3+2*C*c^4+4*C*c^2*d^2)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*ln(c+d*tan(f*x+e)))/d^3/(c^2+d^2)^2/f+b^2*(2*c^2*C-B*c*d+(A+C)*d^2)*tan(f*x+e)/d^2/(c^2+d^2)/f-(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^2/d/(c^2+d^2)/f/(c+d*tan(f*x+e))
    
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.40 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.66

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{\frac{(a+ib)^2(-iA+B+iC) \log(i-\tan(e+fx))}{(c+id)^2} + \frac{(a-ib)^2(iA+B-iC) \log(i+\tan(e+fx))}{(c-id)^2} + \frac{2(-bc+ad)(b(2c^4C-Bc^3d+4c^2Cd^2-3Bcd^3+2Ad^4))}{d^3(c^2+2cd+d^2)}}{2f}$$

input

```
Integrate[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^2,x]
```

output

```
((a + I*b)^2*(-I)*A + B + I*C)*Log[I - Tan[e + f*x]]/(c + I*d)^2 + ((a
- I*b)^2*(I*A + B - I*C)*Log[I + Tan[e + f*x]]/(c - I*d)^2 + (2*(-b*c) +
a*d)*(b*(2*c^4*C - B*c^3*d + 4*c^2*C*d^2 - 3*B*c*d^3 + 2*A*d^4) + a*d^2*(
2*c*(A - C)*d + B*(-c^2 + d^2)))*Log[c + d*Tan[e + f*x]]/(d^3*(c^2 + d^2)
^2) - (2*(b*c - a*d)^2*(2*c^2*C - B*c*d + (A + C)*d^2))/(d^3*(c^2 + d^2)*(
c + d*Tan[e + f*x])) + (2*C*(a + b*Tan[e + f*x])^2)/(d*(c + d*Tan[e + f*x]
)))/(2*f)
```

Rubi [A] (verified)

Time = 3.29 (sec) , antiderivative size = 433, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4128, 3042, 4120, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^2} dx$$

↓ 4128

$$\int \frac{(a+b \tan(e+fx))(b(2Cc^2-Bdc+(A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+2bd)+(2bc-ad)(cC-Bd))}{c+d \tan(e+fx)} dx$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^2}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{df(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))(b(2Cc^2-Bdc+(A+C)d^2) \tan(e+fx)^2+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+2bd)+(2bc-ad)(cC-Bd))}{c+d \tan(e+fx)} dx$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^2}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{df(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 4120

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \int \frac{c(2Cc^2-Bdc+(A+C)d^2)b^2+(2bcC-2adC-bBd)(c^2+d^2) \tan^2(e+fx)b-ad(Ad(ac+2bd)+(2bc-ad)(cC-Bd))}{c+d \tan(e+fx)} dx$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^2}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{df(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 3042

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \int \frac{c(2Cc^2-Bdc+(A+C)d^2)b^2+(2bcC-2adC-bBd)(c^2+d^2) \tan(e+fx)^2b-ad(Ad(ac+2bd)+(2bc-ad)(cC-Bd))}{c+d \tan(e+fx)} dx$$

$$\frac{d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^2}$$

$$\frac{df(c^2+d^2)(c+d \tan(e+fx))}{df(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 4109

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \frac{d^2(a^2(2cd(A-C)-B(c^2-d^2))+2ab(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)-b^2(2cd(A-C)-B(c^2-d^2)))}{c^2+d^2} \int \tan(e+fx)$$

$$\frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^2}{df(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 3042

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \frac{d^2(a^2(2cd(A-C)-B(c^2-d^2))+2ab(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)-b^2(2cd(A-C)-B(c^2-d^2))) \int \tan(e+fx)}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 3956

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \frac{(bc-ad)(ad^2(2cd(A-C)-B(c^2-d^2))+b(2Ad^4-Bc^3d-3Bcd^3+2c^4C+4c^2Cd^2)) \int \frac{\tan(e+fx)^2+1}{c+d \tan(e+fx)} dx - d^2 \log(\cos(e+fx))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 4100

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \frac{(bc-ad)(ad^2(2cd(A-C)-B(c^2-d^2))+b(2Ad^4-Bc^3d-3Bcd^3+2c^4C+4c^2Cd^2)) \int \frac{1}{c+d \tan(e+fx)} d(d \tan(e+fx))}{df(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

↓ 16

$$\frac{b^2 \tan(e+fx)(d^2(A+C)-Bcd+2c^2C)}{df} - \frac{d^2 \log(\cos(e+fx))(a^2(2cd(A-C)-B(c^2-d^2))+2ab(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)-b^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)(c + d \tan(e + fx))}$$

input

```
Int[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^2,x]
```

output

$$\begin{aligned}
& -(((c^2C - Bcd + Ad^2)(a + b\tan[e + fx])^2)/(d(c^2 + d^2)f(c + d \\
& * \tan[e + fx])) + (-(((d^2(a^2(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) \\
& - b^2(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - 2ab(2c(A - C)d - \\
& B(c^2 - d^2))x)/(c^2 + d^2) - (d^2(2ab(c^2C - 2Bcd - Cd^2 - A \\
& (c^2 - d^2)) + a^2(2c(A - C)d - B(c^2 - d^2)) - b^2(2c(A - C)d - \\
& B(c^2 - d^2)))\text{Log}[\text{Cos}[e + fx]])/(c^2 + d^2)f) + ((b^2c - ad)(b(2c^4C \\
& - Bc^3d + 4c^2Cd^2 - 3Bcd^3 + 2Ad^4) + ad^2(2c(A - C)d \\
& - B(c^2 - d^2)))\text{Log}[c + d\tan[e + fx]]/(d(c^2 + d^2)f))/d) + (b^2(2 \\
& * c^2C - Bcd + (A + C)d^2)\text{Tan}[e + fx]/(df))/(d(c^2 + d^2))
\end{aligned}$$

Defintions of rubi rules used

rule 16

$$\text{Int}[(c_)/((a_)+(b_)(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3956

$$\text{Int}[\tan[(c_)+(d_)(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$$

rule 4100

$$\text{Int}[(a_)+(b_)\tan[(e_)+(f_)(x_)]^{(m_)}((A_)+(C_)\tan[(e_)+(f_)(x_)]^2), x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], b*\text{Tan}[e + fx]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \&\& \text{EqQ}[A, C]$$

rule 4109

$$\begin{aligned}
& \text{Int}[(A_)+(B_)\tan[(e_)+(f_)(x_)] + (C_)\tan[(e_)+(f_)(x_)]^2 \\
&)/((a_)+(b_)\tan[(e_)+(f_)(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a \\
& *C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 \\
& + \text{Tan}[e + fx]^2)/(a + b*\text{Tan}[e + fx]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(\\
& a^2 + b^2) \text{ Int}[\text{Tan}[e + fx], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \& \\
& \& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A*b - a*B - b*C \\
& , 0]
\end{aligned}$$

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4128

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) +
(f_)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 552, normalized size of antiderivative = 1.32

method	result
derivativdivides	$\frac{\tan(fx+e)C b^2}{d^2} + \frac{(-2A a^2 cd + 2A ab c^2 - 2A ab d^2 + 2A b^2 cd + B a^2 c^2 - B a^2 d^2 + 4B abcd - B b^2 c^2 + B b^2 d^2 + 2C a^2 cd - 2Cab c^2 + 2Cab d^2)}{2}$
default	$\frac{\tan(fx+e)C b^2}{d^2} + \frac{(-2A a^2 cd + 2A ab c^2 - 2A ab d^2 + 2A b^2 cd + B a^2 c^2 - B a^2 d^2 + 4B abcd - B b^2 c^2 + B b^2 d^2 + 2C a^2 cd - 2Cab c^2 + 2Cab d^2)}{2}$
norman	$\frac{c(A a^2 c^2 - A a^2 d^2 + 4A abcd - A b^2 c^2 + A b^2 d^2 + 2B a^2 cd - 2B ab c^2 + 2B ab d^2 - 2B b^2 cd - C a^2 c^2 + a^2 C d^2 - 4Cab cd + C b^2 c^2 - b^2 d^2 C)}{c^4 + 2c^2 d^2 + d^4}$
parallelrish	Expression too large to display
rish	Expression too large to display

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,
x,method=_RETURNVERBOSE)
```

output

```
1/f*(tan(f*x+e)*C*b^2/d^2+1/(c^2+d^2)^2*(1/2*(-2*A*a^2*c*d+2*A*a*b*c^2-2*A
*a*b*d^2+2*A*b^2*c*d+B*a^2*c^2-B*a^2*d^2+4*B*a*b*c*d-B*b^2*c^2+B*b^2*d^2+2
*C*a^2*c*d-2*C*a*b*c^2+2*C*a*b*d^2-2*C*b^2*c*d)*ln(1+tan(f*x+e)^2)+(A*a^2*c
c^2-A*a^2*d^2+4*A*a*b*c*d-A*b^2*c^2+A*b^2*d^2+2*B*a^2*c*d-2*B*a*b*c^2+2*B*
a*b*d^2-2*B*b^2*c*d-C*a^2*c^2+C*a^2*d^2-4*C*a*b*c*d+C*b^2*c^2-C*b^2*d^2)*a
rctan(tan(f*x+e))-1/d^3*(A*a^2*d^4-2*A*a*b*c*d^3+A*b^2*c^2*d^2-B*a^2*c*d^
3+2*B*a*b*c^2*d^2-B*b^2*c^3*d+C*a^2*c^2*d^2-2*C*a*b*c^3*d+C*b^2*c^4)/(c^2+
d^2)/(c+d*tan(f*x+e))+1/d^3*(2*A*a^2*c*d^4-2*A*a*b*c^2*d^3+2*A*a*b*d^5-2*A
*b^2*c*d^4-B*a^2*c^2*d^3+B*a^2*d^5-4*B*a*b*c*d^4+B*b^2*c^4*d+3*B*b^2*c^2*d
^3-2*C*a^2*c*d^4+2*C*a*b*c^4*d+6*C*a*b*c^2*d^3-2*C*b^2*c^5-4*C*b^2*c^3*d^2
)/(c^2+d^2)^2*ln(c+d*tan(f*x+e)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 939 vs. $2(417) = 834$.

Time = 0.37 (sec) , antiderivative size = 939, normalized size of antiderivative = 2.25

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^2,x, algorithm="fricas")
```

output

```

-1/2*(2*C*b^2*c^4*d^2 + 2*A*a^2*d^6 - 2*(2*C*a*b + B*b^2)*c^3*d^3 + 2*(C*a
^2 + 2*B*a*b + A*b^2)*c^2*d^4 - 2*(B*a^2 + 2*A*a*b)*c*d^5 - 2*(((A - C)*a^
2 - 2*B*a*b - (A - C)*b^2)*c^3*d^3 + 2*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2
*d^4 - ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^5)*f*x - 2*(C*b^2*c^4*d^2
+ 2*C*b^2*c^2*d^4 + C*b^2*d^6)*tan(f*x + e)^2 + (2*C*b^2*c^6 + 4*C*b^2*c^
4*d^2 - (2*C*a*b + B*b^2)*c^5*d + (B*a^2 + 2*(A - 3*C)*a*b - 3*B*b^2)*c^3*
d^3 - 2*((A - C)*a^2 - 2*B*a*b - A*b^2)*c^2*d^4 - (B*a^2 + 2*A*a*b)*c*d^5
+ (2*C*b^2*c^5*d + 4*C*b^2*c^3*d^3 - (2*C*a*b + B*b^2)*c^4*d^2 + (B*a^2 +
2*(A - 3*C)*a*b - 3*B*b^2)*c^2*d^4 - 2*((A - C)*a^2 - 2*B*a*b - A*b^2)*c*d
^5 - (B*a^2 + 2*A*a*b)*d^6)*tan(f*x + e))*log((d^2*tan(f*x + e)^2 + 2*c*d*
tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - (2*C*b^2*c^6 + 4*C*b^2*c^4*d^2
+ 2*C*b^2*c^2*d^4 - (2*C*a*b + B*b^2)*c^5*d - 2*(2*C*a*b + B*b^2)*c^3*d^3
- (2*C*a*b + B*b^2)*c*d^5 + (2*C*b^2*c^5*d + 4*C*b^2*c^3*d^3 + 2*C*b^2*c*
d^5 - (2*C*a*b + B*b^2)*c^4*d^2 - 2*(2*C*a*b + B*b^2)*c^2*d^4 - (2*C*a*b +
B*b^2)*d^6)*tan(f*x + e))*log(1/(tan(f*x + e)^2 + 1)) - 2*(2*C*b^2*c^5*d
- (2*C*a*b + B*b^2)*c^4*d^2 + (C*a^2 + 2*B*a*b + (A + 2*C)*b^2)*c^3*d^3 -
(B*a^2 + 2*A*a*b)*c^2*d^4 + (A*a^2 + C*b^2)*c*d^5 + (((A - C)*a^2 - 2*B*a*
b - (A - C)*b^2)*c^2*d^4 + 2*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d^5 - ((A -
C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^6)*f*x)*tan(f*x + e))/((c^4*d^4 + 2*c^2
*d^6 + d^8)*f*tan(f*x + e) + (c^5*d^3 + 2*c^3*d^5 + c*d^7)*f)

```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.71 (sec) , antiderivative size = 16225, normalized size of antiderivative = 38.91

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((a+b*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*
x+e))**2,x)

```

output

```
Piecewise((zoo*x*(a + b*tan(e))**2*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2,
Eq(c, 0) & Eq(d, 0) & Eq(f, 0)), ((A*a**2*x + A*a*b*log(tan(e + f*x)**2 +
1)/f - A*b**2*x + A*b**2*tan(e + f*x)/f + B*a**2*log(tan(e + f*x)**2 + 1)
/(2*f) - 2*B*a*b*x + 2*B*a*b*tan(e + f*x)/f - B*b**2*log(tan(e + f*x)**2 +
1)/(2*f) + B*b**2*tan(e + f*x)**2/(2*f) - C*a**2*x + C*a**2*tan(e + f*x)/
f - C*a*b*log(tan(e + f*x)**2 + 1)/f + C*a*b*tan(e + f*x)**2/f + C*b**2*x
+ C*b**2*tan(e + f*x)**3/(3*f) - C*b**2*tan(e + f*x)/f)/c**2, Eq(d, 0)), (
-A*a**2*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e +
f*x) - 4*d**2*f) + 2*I*A*a**2*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2
- 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + A*a**2*f*x/(4*d**2*f*tan(e + f*x)*
*2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - A*a**2*tan(e + f*x)/(4*d**2*f*t
an(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*A*a**2/(4*d**2*
f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*A*a*b*f*x*ta
n(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2
*f) + 4*A*a*b*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(
e + f*x) - 4*d**2*f) - 2*I*A*a*b*f*x/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*
f*tan(e + f*x) - 4*d**2*f) + 2*I*A*a*b*tan(e + f*x)/(4*d**2*f*tan(e + f*x)
**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + A*b**2*f*x*tan(e + f*x)**2/(4*
d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - 2*I*A*b**2*
f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - ...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 493, normalized size of antiderivative = 1.18

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{2Cb^2 \tan(fx+e)}{d^2} + \frac{2(((A-C)a^2 - 2Bab - (A-C)b^2)c^2 + 2(Ba^2 + 2(A-C)ab - Bb^2)cd - ((A-C)a^2 - 2Bab - (A-C)b^2)d^2)(fx+e)}{c^4 + 2c^2d^2 + d^4} - \frac{2(2C$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^2,x, algorithm="maxima")
```


output

```

1/2*(2*C*b^2*tan(f*x + e)/d^2 + 2*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c
^2 + 2*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d - ((A - C)*a^2 - 2*B*a*b - (A -
C)*b^2)*d^2)*(f*x + e)/(c^4 + 2*c^2*d^2 + d^4) - 2*(2*C*b^2*c^5 + 4*C*b^2
*c^3*d^2 - (2*C*a*b + B*b^2)*c^4*d + (B*a^2 + 2*(A - 3*C)*a*b - 3*B*b^2)*c
^2*d^3 - 2*((A - C)*a^2 - 2*B*a*b - A*b^2)*c*d^4 - (B*a^2 + 2*A*a*b)*d^5)*
log(d*tan(f*x + e) + c)/(c^4*d^3 + 2*c^2*d^5 + d^7) + ((B*a^2 + 2*(A - C)*
a*b - B*b^2)*c^2 - 2*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d - (B*a^2 +
2*(A - C)*a*b - B*b^2)*d^2)*log(tan(f*x + e)^2 + 1)/(c^4 + 2*c^2*d^2 + d^4
) - 2*(C*b^2*c^4 + A*a^2*d^4 - (2*C*a*b + B*b^2)*c^3*d + (C*a^2 + 2*B*a*b
+ A*b^2)*c^2*d^2 - (B*a^2 + 2*A*a*b)*c*d^3)/(c^3*d^3 + c*d^5 + (c^2*d^4 +
d^6)*tan(f*x + e))/f

```

Giac [A] (verification not implemented)

Time = 0.62 (sec) , antiderivative size = 701, normalized size of antiderivative = 1.68

$$\begin{aligned}
& \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx \\
&= \frac{(Aa^2c^2 - Ca^2c^2 - 2Babc^2 - Ab^2c^2 + Cb^2c^2 + 2Ba^2cd + 4Aabcd - 4Cabcd - 2Bb^2cd - Aa^2d^2 + Ca^2c^2)}{c^4f + 2c^2d^2f + d^4f} \\
&+ \frac{(Ba^2c^2 + 2Aabc^2 - 2Cabc^2 - Bb^2c^2 - 2Aa^2cd + 2Ca^2cd + 4Babcd + 2Ab^2cd - 2Cb^2cd - Ba^2d^2 - 2Aa^2c^2)}{2(c^4f + 2c^2d^2f + d^4f)} \\
&- \frac{(2Cb^2c^5 - 2Cabc^4d - Bb^2c^4d + 4Cb^2c^3d^2 + Ba^2c^2d^3 + 2Aabc^2d^3 - 6Cabc^2d^3 - 3Bb^2c^2d^3 - 2Aa^2c^2d^3)}{c^4d^3f + 2c^2d^5f + d^7f} \\
&+ \frac{Cb^2 \tan(fx + e)}{d^2f} \\
&- \frac{Cb^2c^6 - 2Cabc^5d - Bb^2c^5d + Ca^2c^4d^2 + 2Babc^4d^2 + Ab^2c^4d^2 + Cb^2c^4d^2 - Ba^2c^3d^3 - 2Aabc^3d^3 - 2Aa^2c^3d^3}{(c^2 + d^2)^2(d \tan(fx + e))}
\end{aligned}$$

input

```

integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^2,x, algorithm="giac")

```

output

```
(A*a^2*c^2 - C*a^2*c^2 - 2*B*a*b*c^2 - A*b^2*c^2 + C*b^2*c^2 + 2*B*a^2*c*d
+ 4*A*a*b*c*d - 4*C*a*b*c*d - 2*B*b^2*c*d - A*a^2*d^2 + C*a^2*d^2 + 2*B*a
*b*d^2 + A*b^2*d^2 - C*b^2*d^2)*(f*x + e)/(c^4*f + 2*c^2*d^2*f + d^4*f) +
1/2*(B*a^2*c^2 + 2*A*a*b*c^2 - 2*C*a*b*c^2 - B*b^2*c^2 - 2*A*a^2*c*d + 2*C
*a^2*c*d + 4*B*a*b*c*d + 2*A*b^2*c*d - 2*C*b^2*c*d - B*a^2*d^2 - 2*A*a*b*d
^2 + 2*C*a*b*d^2 + B*b^2*d^2)*log(tan(f*x + e)^2 + 1)/(c^4*f + 2*c^2*d^2*f
+ d^4*f) - (2*C*b^2*c^5 - 2*C*a*b*c^4*d - B*b^2*c^4*d + 4*C*b^2*c^3*d^2 +
B*a^2*c^2*d^3 + 2*A*a*b*c^2*d^3 - 6*C*a*b*c^2*d^3 - 3*B*b^2*c^2*d^3 - 2*A
*a^2*c*d^4 + 2*C*a^2*c*d^4 + 4*B*a*b*c*d^4 + 2*A*b^2*c*d^4 - B*a^2*d^5 - 2
*A*a*b*d^5)*log(abs(d*tan(f*x + e) + c))/(c^4*d^3*f + 2*c^2*d^5*f + d^7*f)
+ C*b^2*tan(f*x + e)/(d^2*f) - (C*b^2*c^6 - 2*C*a*b*c^5*d - B*b^2*c^5*d +
C*a^2*c^4*d^2 + 2*B*a*b*c^4*d^2 + A*b^2*c^4*d^2 + C*b^2*c^4*d^2 - B*a^2*c
^3*d^3 - 2*A*a*b*c^3*d^3 - 2*C*a*b*c^3*d^3 - B*b^2*c^3*d^3 + A*a^2*c^2*d^4
+ C*a^2*c^2*d^4 + 2*B*a*b*c^2*d^4 + A*b^2*c^2*d^4 - B*a^2*c*d^5 - 2*A*a*b
*c*d^5 + A*a^2*d^6)/((c^2 + d^2)^2*(d*tan(f*x + e) + c)*d^3*f)
```

Mupad [B] (verification not implemented)

Time = 29.98 (sec) , antiderivative size = 3958, normalized size of antiderivative = 9.49

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int(((a + b*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c +
d*tan(e + f*x))^2,x)
```

output

```
(log((2*C^2*b^4*c^5 - 2*C^2*a^2*b^2*c^5 + 4*C^2*b^4*c^3*d^2 - A*B*a^4*d^5
- 2*A*C*b^4*c^5 + B*C*a^4*d^5 + 2*A^2*a*b^3*d^5 - 2*A^2*a^3*b*d^5 - A^2*a^
4*c*d^4 + 2*B^2*a^3*b*d^5 - A^2*b^4*c*d^4 + B^2*a^4*c*d^4 + B^2*b^4*c*d^4
- C^2*a^4*c*d^4 + C^2*b^4*c*d^4 - 4*C^2*a^2*b^2*c^3*d^2 + 5*A*B*a^2*b^2*d^
5 + 2*A*C*a^2*b^2*c^5 + A*B*a^4*c^2*d^3 + 3*A*B*b^4*c^2*d^3 - B*C*a^2*b^2*
d^5 - 4*A*C*b^4*c^3*d^2 - B*C*a^4*c^2*d^3 - 3*B*C*b^4*c^2*d^3 + 2*B^2*a*b^
3*c^4*d - 2*C^2*a*b^3*c^4*d + 2*C^2*a^3*b*c^4*d - 2*A^2*a*b^3*c^2*d^3 + 6*
A^2*a^2*b^2*c*d^4 + 2*A^2*a^3*b*c^2*d^3 + 6*B^2*a*b^3*c^2*d^3 - 6*B^2*a^2*
b^2*c*d^4 - 2*B^2*a^3*b*c^2*d^3 - 6*C^2*a*b^3*c^2*d^3 + 4*C^2*a^2*b^2*c*d^
4 + 6*C^2*a^3*b*c^2*d^3 - 2*A*C*a*b^3*d^5 + 2*A*C*a^3*b*d^5 - 4*B*C*a*b^3*
c^5 + A*B*b^4*c^4*d + 2*A*C*a^4*c*d^4 - B*C*b^4*c^4*d - 8*A*B*a*b^3*c*d^4
+ 8*A*B*a^3*b*c*d^4 + 2*A*C*a*b^3*c^4*d - 2*A*C*a^3*b*c^4*d + 4*B*C*a*b^3*
c*d^4 - 8*B*C*a^3*b*c*d^4 - A*B*a^2*b^2*c^4*d + 8*A*C*a*b^3*c^2*d^3 - 10*A
*C*a^2*b^2*c*d^4 - 8*A*C*a^3*b*c^2*d^3 - 8*B*C*a*b^3*c^3*d^2 + 5*B*C*a^2*b
^2*c^4*d - 8*A*B*a^2*b^2*c^2*d^3 + 4*A*C*a^2*b^2*c^3*d^2 + 16*B*C*a^2*b^2*
c^2*d^3)/(d^2*(c^2 + d^2)^2) + ((a+1i - b)^2*((A*b^2*d^2 - A*a^2*d^2 + C*a
^2*d^2 - 8*C*b^2*c^2 - C*b^2*d^2 + 2*B*a*b*d^2 + 4*B*b^2*c*d + 8*C*a*b*c*d
)/d - (tan(e + f*x))*(3*B*a^2*d^5 - 5*B*b^2*d^5 - 4*C*b^2*c^5 + 6*A*a*b*d^5
- 10*C*a*b*d^5 + 4*A*a^2*c*d^4 - 4*A*b^2*c*d^4 + 2*B*b^2*c^4*d - 4*C*a^2*
c*d^4 + 8*C*b^2*c*d^4 - B*a^2*c^2*d^3 + B*b^2*c^2*d^3 - 8*B*a*b*c*d^4 + ...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1762, normalized size of antiderivative = 4.23

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,
x)
```

output

```
( - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*c**2*d**5 + 3*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a**2*b*c**3*d**4 - 3*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)*a**2*b*c*d**6 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*c
**3*d**5 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**2*c**2*d**5 - 2*lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b*c**4*d**4 + 2*log(tan(e + f*x)**2
+ 1)*tan(e + f*x)*a*b*c**2*d**6 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b*
**3*c**3*d**4 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**3*c*d**6 - 2*log(t
an(e + f*x)**2 + 1)*tan(e + f*x)*b**2*c**3*d**5 - 2*log(tan(e + f*x)**2 +
1)*a**3*c**3*d**4 + 3*log(tan(e + f*x)**2 + 1)*a**2*b*c**4*d**3 - 3*log(ta
n(e + f*x)**2 + 1)*a**2*b*c**2*d**5 + 2*log(tan(e + f*x)**2 + 1)*a**2*c**4
*d**4 + 6*log(tan(e + f*x)**2 + 1)*a*b**2*c**3*d**4 - 2*log(tan(e + f*x)**
2 + 1)*a*b*c**5*d**3 + 2*log(tan(e + f*x)**2 + 1)*a*b*c**3*d**5 - log(tan(
e + f*x)**2 + 1)*b**3*c**4*d**3 + log(tan(e + f*x)**2 + 1)*b**3*c**2*d**5
- 2*log(tan(e + f*x)**2 + 1)*b**2*c**4*d**4 + 4*log(tan(e + f*x)*d + c)*ta
n(e + f*x)*a**3*c**2*d**5 - 6*log(tan(e + f*x)*d + c)*tan(e + f*x)*a**2*b*
c**3*d**4 + 6*log(tan(e + f*x)*d + c)*tan(e + f*x)*a**2*b*c*d**6 - 4*log(t
an(e + f*x)*d + c)*tan(e + f*x)*a**2*c**3*d**5 - 12*log(tan(e + f*x)*d + c
)*tan(e + f*x)*a*b**2*c**2*d**5 + 4*log(tan(e + f*x)*d + c)*tan(e + f*x)*a
*b*c**6*d**2 + 12*log(tan(e + f*x)*d + c)*tan(e + f*x)*a*b*c**4*d**4 + 2*l
og(tan(e + f*x)*d + c)*tan(e + f*x)*b**3*c**5*d**2 + 6*log(tan(e + f*x)...
```

3.79
$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$$

Optimal result	880
Mathematica [C] (verified)	881
Rubi [A] (verified)	881
Maple [A] (verified)	884
Fricas [A] (verification not implemented)	885
Sympy [C] (verification not implemented)	886
Maxima [A] (verification not implemented)	887
Giac [A] (verification not implemented)	887
Mupad [B] (verification not implemented)	888
Reduce [B] (verification not implemented)	889

Optimal result

Integrand size = 43, antiderivative size = 292

$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$$

$$= -\frac{(a(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - b(2c(A - C)d - B(c^2 - d^2)))x}{(c^2 + d^2)^2}$$

$$- \frac{(a(Bc^2 + 2cCd - Bd^2) - b(c^2C - 2Bcd - Cd^2) - A(2acd - b(c^2 - d^2))) \log(\cos(e+fx))}{(c^2 + d^2)^2 f}$$

$$+ \frac{(b(c^4C - c^2(A - 3C)d^2 - 2Bcd^3 + Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2))) \log(c+d \tan(e+fx))}{d^2 (c^2 + d^2)^2 f}$$

$$+ \frac{(bc - ad)(c^2C - Bcd + Ad^2)}{d^2 (c^2 + d^2) f(c+d \tan(e+fx))}$$

output

```
- (a*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-b*(2*c*(A-C)*d-B*(c^2-d^2)) *x / (c^2+d^2)^2 - (a*(B*c^2-B*d^2+2*C*c*d)-b*(-2*B*c*d+C*c^2-C*d^2)-A*(2*a*c*d-b*(c^2-d^2))) *ln(cos(f*x+e)) / (c^2+d^2)^2 / f + (b*(c^4*C-c^2*(A-3*C)*d^2-2*B*c*d^3+Ad^4)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2))) *ln(c+d*tan(f*x+e)) / d^2 / (c^2+d^2)^2 / f + (-a*d+b*c)*(A*d^2-B*c*d+C*c^2) / d^2 / (c^2+d^2) / f / (c+d*tan(f*x+e))
```


$$\int \frac{bC(c^2+d^2) \tan^2(e+fx) + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{c+d \tan(e+fx)} dx + \frac{d(c^2+d^2)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))} + \frac{(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 3042

$$\int \frac{bC(c^2+d^2) \tan(e+fx)^2 + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{c+d \tan(e+fx)} dx + \frac{d(c^2+d^2)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))} + \frac{(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 4109

$$\frac{-\frac{d(2aAc-d-aB(c^2-d^2))-2acCd-Ab(c^2-d^2)+b(-2Bcd+c^2C-Cd^2)}{c^2+d^2} \int \tan(e+fx) dx + \frac{(ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-3C)+Ad^4-2Bcd^3+c^4C))}{c^2+d^2}}{d(c^2+d^2)} + \frac{(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 3042

$$\frac{-\frac{d(2aAc-d-aB(c^2-d^2))-2acCd-Ab(c^2-d^2)+b(-2Bcd+c^2C-Cd^2)}{c^2+d^2} \int \tan(e+fx) dx + \frac{(ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-3C)+Ad^4-2Bcd^3+c^4C))}{c^2+d^2}}{d(c^2+d^2)} + \frac{(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 3956

$$\frac{(ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-3C)+Ad^4-2Bcd^3+c^4C)) \int \frac{\tan(e+fx)^2+1}{c+d \tan(e+fx)} dx + \frac{d \log(\cos(e+fx))(2aAc-d-aB(c^2-d^2))-2acCd-A}{f(c^2+d^2)}}{d(c^2+d^2)} + \frac{(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 4100

$$\frac{(ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-3C)+Ad^4-2Bcd^3+c^4C)) \int \frac{1}{c+d \tan(e+fx)} d(d \tan(e+fx)) + \frac{d \log(\cos(e+fx))(2aAc-d-aB(c^2-d^2))-2acCd-A}{f(c^2+d^2)}}{d(c^2+d^2)} + \frac{(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2)(c+d \tan(e+fx))}$$

↓ 16

$$\frac{(bc - ad)(Ad^2 - Bcd + c^2C)}{d^2 f(c^2 + d^2)(c + d \tan(e + fx))} + \frac{d \log(\cos(e + fx))(2aAc d - aB(c^2 - d^2) - 2acCd - Ab(c^2 - d^2) + b(-2Bcd + c^2C - Cd^2))}{f(c^2 + d^2)} - \frac{dx(a(-A(c^2 - d^2) - 2Bcd + c^2C - Cd^2) - b(2cd(A - C) - c^2 + d^2))}{d(c^2 + d^2)}$$

input `Int[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^2,x]`

output `((-((d*(a*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b*(2*c*(A - C)*d - B*(c^2 - d^2)))*x)/(c^2 + d^2)) + (d*(2*a*A*c*d - 2*a*c*C*d - A*b*(c^2 - d^2) - a*B*(c^2 - d^2) + b*(c^2*C - 2*B*c*d - C*d^2))*Log[Cos[e + f*x]])/((c^2 + d^2)*f) + ((b*(c^4*C - c^2*(A - 3*C)*d^2 - 2*B*c*d^3 + A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Log[c + d*Tan[e + f*x]])/(d*(c^2 + d^2)*f))/((d*(c^2 + d^2)) + ((b*c - a*d)*(c^2*C - B*c*d + A*d^2))/(d^2*(c^2 + d^2))*f*(c + d*Tan[e + f*x]))`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]
```

rule 4118

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{(-2Aacd+Abc^2-Abd^2+Ba c^2-Ba d^2+2Bbcd+2Cacd-Cb c^2+Cb d^2) \ln(1+\tan(fx+e)^2)}{2} + \frac{(Aa c^2-Aa d^2+2Abcd+2Bacd-Bb c^2+Bb d^2)}{(c^2+d^2)^2}$
default	$\frac{(-2Aacd+Abc^2-Abd^2+Ba c^2-Ba d^2+2Bbcd+2Cacd-Cb c^2+Cb d^2) \ln(1+\tan(fx+e)^2)}{2} + \frac{(Aa c^2-Aa d^2+2Abcd+2Bacd-Bb c^2+Bb d^2)}{(c^2+d^2)^2}$
norman	$\frac{c(Aa c^2-Aa d^2+2Abcd+2Bacd-Bb c^2+Bb d^2-Ca c^2+Ca d^2-2Cbcd)}{c^4+2c^2d^2+d^4}x + \frac{d(Aa c^2-Aa d^2+2Abcd+2Bacd-Bb c^2+Bb d^2-Ca c^2+Ca d^2-2Cbcd)}{c^4+2c^2d^2+d^4} + \frac{c+d \tan(fx+e)}{c+d \tan(fx+e)}$
parallelrisch	Expression too large to display
risch	Expression too large to display

input

```
int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, method=_RETURNVERBOSE)
```

output

```
1/f*(1/(c^2+d^2)^2*(1/2*(-2*A*a*c*d+A*b*c^2-A*b*d^2+B*a*c^2-B*a*d^2+2*B*b*
c*d+2*C*a*c*d-C*b*c^2+C*b*d^2)*ln(1+tan(f*x+e)^2)+(A*a*c^2-A*a*d^2+2*A*b*c
*d+2*B*a*c*d-B*b*c^2+B*b*d^2-C*a*c^2+C*a*d^2-2*C*b*c*d)*arctan(tan(f*x+e))
)-(A*a*d^3-A*b*c*d^2-B*a*c*d^2+B*b*c^2*d+C*a*c^2*d-C*b*c^3)/d^2/(c^2+d^2)/
(c+d*tan(f*x+e))+(2*A*a*c*d^3-A*b*c^2*d^2+A*b*d^4-B*a*c^2*d^2+B*a*d^4-2*B*
b*c*d^3-2*C*a*c*d^3+C*b*c^4+3*C*b*c^2*d^2)/(c^2+d^2)^2/d^2*ln(c+d*tan(f*x+
e)))
```

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 505, normalized size of antiderivative = 1.73

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{2Cb^3d^2 - 2Aad^5 - 2(Ca + Bb)c^2d^3 + 2(Ba + Ab)cd^4 + 2(((A - C)a - Bb)c^3d^2 + 2(Ba + (A - C)b$$

input

```
integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^2,x, algorithm="fricas")
```

output

```
1/2*(2*C*b*c^3*d^2 - 2*A*a*d^5 - 2*(C*a + B*b)*c^2*d^3 + 2*(B*a + A*b)*c*d
^4 + 2*(((A - C)*a - B*b)*c^3*d^2 + 2*(B*a + (A - C)*b)*c^2*d^3 - ((A - C)
*a - B*b)*c*d^4)*f*x + (C*b*c^5 - (B*a + (A - 3*C)*b)*c^3*d^2 + 2*((A - C)
*a - B*b)*c^2*d^3 + (B*a + A*b)*c*d^4 + (C*b*c^4*d - (B*a + (A - 3*C)*b)*c
^2*d^3 + 2*((A - C)*a - B*b)*c*d^4 + (B*a + A*b)*d^5)*tan(f*x + e))*log((d
^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - (C*b
*c^5 + 2*C*b*c^3*d^2 + C*b*c*d^4 + (C*b*c^4*d + 2*C*b*c^2*d^3 + C*b*d^5)*t
an(f*x + e))*log(1/(tan(f*x + e)^2 + 1)) - 2*(C*b*c^4*d - A*a*c*d^4 - (C*a
+ B*b)*c^3*d^2 + (B*a + A*b)*c^2*d^3 - (((A - C)*a - B*b)*c^2*d^3 + 2*(B*
a + (A - C)*b)*c*d^4 - ((A - C)*a - B*b)*d^5)*f*x)*tan(f*x + e))/((c^4*d^3
+ 2*c^2*d^5 + d^7)*f*tan(f*x + e) + (c^5*d^2 + 2*c^3*d^4 + c*d^6)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.32 (sec) , antiderivative size = 9721, normalized size of antiderivative = 33.29

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**2,x)`

output

```
Piecewise((zoo*x*(a + b*tan(e))*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2, Eq
(c, 0) & Eq(d, 0) & Eq(f, 0)), ((A*a*x + A*b*log(tan(e + f*x)**2 + 1)/(2*f
) + B*a*log(tan(e + f*x)**2 + 1)/(2*f) - B*b*x + B*b*tan(e + f*x)/f - C*a*
x + C*a*tan(e + f*x)/f - C*b*log(tan(e + f*x)**2 + 1)/(2*f) + C*b*tan(e +
f*x)**2/(2*f))/c**2, Eq(d, 0)), (-A*a*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e
+ f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*A*a*f*x*tan(e + f*x)
/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + A*a*f*x
/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f)
+ 2*I*A*a/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f)
+ I*A*b*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e +
f*x) - 4*d**2*f) + 2*A*b*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I
*d**2*f*tan(e + f*x) - 4*d**2*f) - I*A*b*f*x/(4*d**2*f*tan(e + f*x)**2 - 8
*I*d**2*f*tan(e + f*x) - 4*d**2*f) + I*A*b*tan(e + f*x)/(4*d**2*f*tan(e +
f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + I*B*a*f*x*tan(e + f*x)**2/
(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*B*a*f*
x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**
2*f) - I*B*a*f*x/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d
**2*f) + I*B*a*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e +
f*x) - 4*d**2*f) + B*b*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 - ...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.09

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{2(((A-C)a-Bb)c^2+2(Ba+(A-C)b)cd-((A-C)a-Bb)d^2)(fx+e)}{c^4+2c^2d^2+d^4} + \frac{2(Cbc^4-(Ba+(A-3C)b)c^2d^2+2((A-C)a-Bb)cd^3+(Ba+Ab)d^4)}{c^4d^2+2c^2d^4+d^6} \log(\tan(fx+e)^2+1)$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/2*(2*((A - C)*a - B*b)*c^2 + 2*(B*a + (A - C)*b)*c*d - ((A - C)*a - B*b)*d^2)*(f*x + e)/(c^4 + 2*c^2*d^2 + d^4) + 2*(C*b*c^4 - (B*a + (A - 3*C)*b)*c^2*d^2 + 2*((A - C)*a - B*b)*c*d^3 + (B*a + A*b)*d^4)*log(d*tan(f*x + e) + c)/(c^4*d^2 + 2*c^2*d^4 + d^6) + ((B*a + (A - C)*b)*c^2 - 2*((A - C)*a - B*b)*c*d - (B*a + (A - C)*b)*d^2)*log(tan(f*x + e)^2 + 1)/(c^4 + 2*c^2*d^2 + d^4) + 2*(C*b*c^3 - A*a*d^3 - (C*a + B*b)*c^2*d + (B*a + A*b)*c*d^2)/(c^3*d^2 + c*d^4 + (c^2*d^3 + d^5)*tan(f*x + e))/f`

Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.44

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{(Aac^2 - Cac^2 - Bbc^2 + 2 Bacd + 2 Abcd - 2 Cbcd - Aad^2 + Cad^2 + Bbd^2)(fx + e)}{c^4f + 2c^2d^2f + d^4f} + \frac{(Bac^2 + Abc^2 - Cbc^2 - 2 Aacd + 2 Cacd + 2 Bbcd - Bad^2 - Abd^2 + Cbd^2) \log(\tan(fx + e)^2 + 1)}{2(c^4f + 2c^2d^2f + d^4f)} + \frac{(Cbc^4 - Bac^2d^2 - Abc^2d^2 + 3Cbc^2d^2 + 2Aacd^3 - 2Cacd^3 - 2Bbcd^3 + Bad^4 + Abd^4) \log(|d \tan(fx + e) + c|)}{c^4d^2f + 2c^2d^4f + d^6f} + \frac{Cbc^5 - Cac^4d - Bbc^4d + Bac^3d^2 + Abc^3d^2 + Cbc^3d^2 - Aac^2d^3 - Cac^2d^3 - Bbc^2d^3 + Bacd^4 + Abcd^4}{(c^2 + d^2)^2(d \tan(fx + e) + c)d^2f}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="giac")`

output

```
(A*a*c^2 - C*a*c^2 - B*b*c^2 + 2*B*a*c*d + 2*A*b*c*d - 2*C*b*c*d - A*a*d^2
+ C*a*d^2 + B*b*d^2)*(f*x + e)/(c^4*f + 2*c^2*d^2*f + d^4*f) + 1/2*(B*a*c
^2 + A*b*c^2 - C*b*c^2 - 2*A*a*c*d + 2*C*a*c*d + 2*B*b*c*d - B*a*d^2 - A*b
*d^2 + C*b*d^2)*log(tan(f*x + e)^2 + 1)/(c^4*f + 2*c^2*d^2*f + d^4*f) + (C
*b*c^4 - B*a*c^2*d^2 - A*b*c^2*d^2 + 3*C*b*c^2*d^2 + 2*A*a*c*d^3 - 2*C*a*c
*d^3 - 2*B*b*c*d^3 + B*a*d^4 + A*b*d^4)*log(abs(d*tan(f*x + e) + c))/(c^4*
d^2*f + 2*c^2*d^4*f + d^6*f) + (C*b*c^5 - C*a*c^4*d - B*b*c^4*d + B*a*c^3*
d^2 + A*b*c^3*d^2 + C*b*c^3*d^2 - A*a*c^2*d^3 - C*a*c^2*d^3 - B*b*c^2*d^3
+ B*a*c*d^4 + A*b*c*d^4 - A*a*d^5)/((c^2 + d^2)^2*(d*tan(f*x + e) + c)*d^2
*f)
```

Mupad [B] (verification not implemented)

Time = 18.12 (sec) , antiderivative size = 1875, normalized size of antiderivative = 6.42

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int(((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*
tan(e + f*x))^2,x)
```

output

```
(log(c + d*tan(e + f*x))*(d^4*(A*b + B*a) - d^3*(2*B*b*c - 2*A*a*c + 2*C*a*c) - d^2*(A*b*c^2 + B*a*c^2 - 3*C*b*c^2) + C*b*c^4))/(f*(d^6 + 2*c^2*d^4 + c^4*d^2)) - (log((A*B*b^2*d^4 - A*B*a^2*d^4 + B*C*a^2*d^4 + B*C*b^2*c^4 - A^2*a*b*d^4 + B^2*a*b*d^4 + C^2*a*b*c^4 - A^2*a^2*c*d^3 + A^2*b^2*c*d^3 + B^2*a^2*c*d^3 - B^2*b^2*c*d^3 - C^2*a^2*c*d^3 + C^2*b^2*c*d^3 + A*B*a^2*c^2*d^2 - A*B*b^2*c^2*d^2 - B*C*a^2*c^2*d^2 + 3*B*C*b^2*c^2*d^2 + A^2*a*b*c^2*d^2 - B^2*a*b*c^2*d^2 + 3*C^2*a*b*c^2*d^2 - A*C*a*b*c^4 + A*C*a*b*d^4 + 2*A*C*a^2*c*d^3 - 2*A*C*b^2*c*d^3 - 4*A*C*a*b*c^2*d^2 + 4*A*B*a*b*c*d^3 - 4*B*C*a*b*c*d^3))/(d*(c^2 + d^2)^2) + (tan(e + f*x)*(A^2*a^2*d^4 + B^2*b^2*d^4 + C^2*a^2*d^4 + C^2*b^2*c^4 + C^2*b^2*d^4 + A^2*b^2*c^2*d^2 + B^2*a^2*c^2*d^2 + 3*C^2*b^2*c^2*d^2 - 2*A*C*a^2*d^4 - A*C*b^2*c^4 - A*C*b^2*d^4 - 4*A*C*b^2*c^2*d^2 - 2*A*B*a*b*d^4 - B*C*a*b*c^4 + B*C*a*b*d^4 - 2*A*B*a^2*c*d^3 + 2*A*B*b^2*c*d^3 + 2*B*C*a^2*c*d^3 - 2*B*C*b^2*c*d^3 - 2*A^2*a*b*c*d^3 + 2*B^2*a*b*c*d^3 - 2*C^2*a*b*c*d^3 + 2*A*B*a*b*c^2*d^2 - 4*B*C*a*b*c^2*d^2 + 4*A*C*a*b*c*d^3))/(d*(c^2 + d^2)^2) + ((a*1i + b)*(B*1i - A + C)*(A*a*d - B*b*d - C*a*d - 4*C*b*c + (tan(e + f*x))*(3*A*b*d^4 + 3*B*a*d^4 + 2*C*b*c^4 - 5*C*b*d^4 + 4*A*a*c*d^3 - 4*B*b*c*d^3 - 4*C*a*c*d^3 - A*b*c^2*d^2 - B*a*c^2*d^2 + C*b*c^2*d^2))/(d*(c^2 + d^2)) + (d*(a*1i + b)*(4*c*d - c^2*tan(e + f*x) + 3*d^2*tan(e + f*x))*(B*1i - A + C))/(c*1i + d)^2))/(2*(c*1i + d)^2)*(A*a*1i + A*b + B*a - B*b*1i - C*a*1i - C*b))/(2*f*(c*d...
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1114, normalized size of antiderivative = 3.82

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x)
```

output

```
( - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*c**2*d**4 + 2*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a*b*c**3*d**3 - 2*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)*a*b*c*d**5 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*c**3*d**4
+ 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**2*c**2*d**4 - log(tan(e + f*x)
)**2 + 1)*tan(e + f*x)*b*c**4*d**3 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)
*b*c**2*d**5 - 2*log(tan(e + f*x)**2 + 1)*a**2*c**3*d**3 + 2*log(tan(e + f
*x)**2 + 1)*a*b*c**4*d**2 - 2*log(tan(e + f*x)**2 + 1)*a*b*c**2*d**4 + 2*l
og(tan(e + f*x)**2 + 1)*a*c**4*d**3 + 2*log(tan(e + f*x)**2 + 1)*b**2*c**3
*d**3 - log(tan(e + f*x)**2 + 1)*b*c**5*d**2 + log(tan(e + f*x)**2 + 1)*b*
c**3*d**4 + 4*log(tan(e + f*x)*d + c)*tan(e + f*x)*a**2*c**2*d**4 - 4*log(
tan(e + f*x)*d + c)*tan(e + f*x)*a*b*c**3*d**3 + 4*log(tan(e + f*x)*d + c)
*tan(e + f*x)*a*b*c*d**5 - 4*log(tan(e + f*x)*d + c)*tan(e + f*x)*a*c**3*d
**4 - 4*log(tan(e + f*x)*d + c)*tan(e + f*x)*b**2*c**2*d**4 + 2*log(tan(e
+ f*x)*d + c)*tan(e + f*x)*b*c**6*d + 6*log(tan(e + f*x)*d + c)*tan(e + f*
x)*b*c**4*d**3 + 4*log(tan(e + f*x)*d + c)*a**2*c**3*d**3 - 4*log(tan(e +
f*x)*d + c)*a*b*c**4*d**2 + 4*log(tan(e + f*x)*d + c)*a*b*c**2*d**4 - 4*lo
g(tan(e + f*x)*d + c)*a*c**4*d**3 - 4*log(tan(e + f*x)*d + c)*b**2*c**3*d*
**3 + 2*log(tan(e + f*x)*d + c)*b*c**7 + 6*log(tan(e + f*x)*d + c)*b*c**5*d
**2 + 2*tan(e + f*x)*a**2*c**3*d**3*f*x + 2*tan(e + f*x)*a**2*c**2*d**4 -
2*tan(e + f*x)*a**2*c*d**5*f*x + 2*tan(e + f*x)*a**2*d**6 - 4*tan(e + f...
```

3.80
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^2} dx$$

Optimal result	891
Mathematica [C] (verified)	892
Rubi [A] (verified)	892
Maple [A] (verified)	894
Fricas [A] (verification not implemented)	895
Sympy [C] (verification not implemented)	896
Maxima [A] (verification not implemented)	897
Giac [A] (verification not implemented)	897
Mupad [B] (verification not implemented)	898
Reduce [B] (verification not implemented)	898

Optimal result

Integrand size = 33, antiderivative size = 140

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx$$

$$= -\frac{(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2))x}{(c^2 + d^2)^2}$$

$$+ \frac{(2c(A - C)d - B(c^2 - d^2)) \log(c \cos(e + fx) + d \sin(e + fx))}{(c^2 + d^2)^2 f}$$

$$- \frac{c^2C - Bcd + Ad^2}{d(c^2 + d^2) f(c + d \tan(e + fx))}$$

output

```
-(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))*x/(c^2+d^2)^2+(2*c*(A-C)*d-B*(c^2-d^2))
*ln(c*cos(f*x+e)+d*sin(f*x+e))/(c^2+d^2)^2/f-(A*d^2-B*c*d+C*c^2)/d/(c^2+d^
2)/f/(c+d*tan(f*x+e))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.78 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.48

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{B((-ic-d) \log(i - \tan(e+fx)) + i(c+id) \log(i + \tan(e+fx)) + 2d \log(c+d \tan(e+fx)))}{c^2+d^2} - \frac{2C}{c+d \tan(e+fx)} + (Bc + (-A + C)d) \left(\frac{i \log}{2df} \right)$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x]^2,x]
```

output

```
((B*(((-I)*c - d)*Log[I - Tan[e + f*x]] + I*(c + I*d)*Log[I + Tan[e + f*x]] + 2*d*Log[c + d*Tan[e + f*x]]))/(c^2 + d^2) - (2*C)/(c + d*Tan[e + f*x]) + (B*c + (-A + C)*d)*((I*Log[I - Tan[e + f*x]])/(c + I*d)^2 - (I*Log[I + Tan[e + f*x]])/(c - I*d)^2 + (2*d*(-2*c*Log[c + d*Tan[e + f*x]] + (c^2 + d^2)/(c + d*Tan[e + f*x])))/(c^2 + d^2)^2))/(2*d*f)
```

Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4111, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(c + d \tan(e + fx))^2} dx$$

↓ 4111

$$\begin{aligned}
 & \frac{\int \frac{Ac-Cc+Bd+(Bc-(A-C)d)\tan(e+fx)}{c+d\tan(e+fx)} dx}{c^2+d^2} - \frac{Ad^2-Bcd+c^2C}{df(c^2+d^2)(c+d\tan(e+fx))} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{Ac-Cc+Bd+(Bc-(A-C)d)\tan(e+fx)}{c+d\tan(e+fx)} dx}{c^2+d^2} - \frac{Ad^2-Bcd+c^2C}{df(c^2+d^2)(c+d\tan(e+fx))} \\
 & \quad \downarrow 4014 \\
 & \frac{(2cd(A-C)-B(c^2-d^2)) \int \frac{d-c\tan(e+fx)}{c+d\tan(e+fx)} dx}{c^2+d^2} - \frac{x(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)}{c^2+d^2} - \\
 & \quad \frac{c^2+d^2}{df(c^2+d^2)(c+d\tan(e+fx))} \frac{Ad^2-Bcd+c^2C}{df(c^2+d^2)(c+d\tan(e+fx))} \\
 & \quad \downarrow 3042 \\
 & \frac{(2cd(A-C)-B(c^2-d^2)) \int \frac{d-c\tan(e+fx)}{c+d\tan(e+fx)} dx}{c^2+d^2} - \frac{x(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)}{c^2+d^2} - \\
 & \quad \frac{c^2+d^2}{df(c^2+d^2)(c+d\tan(e+fx))} \frac{Ad^2-Bcd+c^2C}{df(c^2+d^2)(c+d\tan(e+fx))} \\
 & \quad \downarrow 4013 \\
 & \frac{(2cd(A-C)-B(c^2-d^2)) \log(c\cos(e+fx)+d\sin(e+fx))}{f(c^2+d^2)} - \frac{x(-A(c^2-d^2)-2Bcd+c^2C-Cd^2)}{c^2+d^2} - \\
 & \quad \frac{c^2+d^2}{df(c^2+d^2)(c+d\tan(e+fx))} \frac{Ad^2-Bcd+c^2C}{df(c^2+d^2)(c+d\tan(e+fx))}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^2,x]`

output `(-(((c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2))*x)/(c^2 + d^2)) + ((2*c*(A - C)*d - B*(c^2 - d^2))*Log[c*Cos[e + f*x] + d*Sin[e + f*x]])/(c^2 + d^2)*f)/(c^2 + d^2) - (c^2*C - B*c*d + A*d^2)/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)])*(x_), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4111 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*(a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2)), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.24

method	result
derivativedivides	$\frac{\frac{(-2Acd+Bc^2-Bd^2+2Ccd)\ln(1+\tan(fx+e))^2}{2} + (Ac^2-A d^2+2Bcd-c^2C+C d^2)\arctan(\tan(fx+e))}{(c^2+d^2)^2} - \frac{A d^2-Bcd+c^2C}{(c^2+d^2)d(c+d\tan(fx+e))}$
default	$\frac{\frac{(-2Acd+Bc^2-Bd^2+2Ccd)\ln(1+\tan(fx+e))^2}{2} + (Ac^2-A d^2+2Bcd-c^2C+C d^2)\arctan(\tan(fx+e))}{(c^2+d^2)^2} - \frac{A d^2-Bcd+c^2C}{(c^2+d^2)d(c+d\tan(fx+e))}$
norman	$\frac{c(Ac^2-A d^2+2Bcd-c^2C+C d^2)x}{c^4+2c^2d^2+d^4} + \frac{d(Ac^2-A d^2+2Bcd-c^2C+C d^2)x\tan(fx+e)}{c^4+2c^2d^2+d^4} - \frac{A d^2-Bcd+c^2C}{(c^2+d^2)fd} + \frac{(2Acd-Bc^2+Bd^2)}{f(c^4)}$
parallelrisch	$-\frac{2Ac^2d^2-2Bc^3d-2Bcd^3+2C^2d^2+2c^4C+2Ad^4-2Ax\tan(fx+e)c^2d^2f-4Bx\tan(fx+e)c^3f+2Cx\tan(fx+e)c^4}{(c^2+d^2)^2}$
risch	$\frac{2ic^2C}{(id+c)f(-id+c)^2(-ide^{2i(fx+e)}+ce^{2i(fx+e)}+id+c)} - \frac{x A}{2icd-c^2+d^2} + \frac{x C}{2icd-c^2+d^2} + \frac{ix B}{2icd-c^2+d^2} - \frac{2iBx}{c^4+2c^2d^2+d^4}$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{f} * \left(\frac{1}{(c^2+d^2)^2} * \left(\frac{1}{2} * (-2*A*c*d+B*c^2-B*d^2+2*C*c*d) * \ln(1+\tan(f*x+e))^2 + (A*c^2-A*d^2+2*B*c*d-C*c^2+C*d^2) * \arctan(\tan(f*x+e)) \right) - (A*d^2-B*c*d+C*c^2) / (c^2+d^2) / d / (c+d*\tan(f*x+e)) + (2*A*c*d-B*c^2+B*d^2-2*C*c*d) / (c^2+d^2)^2 * \ln(c+d*\tan(f*x+e)) \right)$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.83

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx = \frac{2 C c^2 d - 2 B c d^2 + 2 A d^3 - 2 ((A - C) c^3 + 2 B c^2 d - (A - C) c d^2) f x + (B c^3 - 2 (A - C) c^2 d - B c d^2)}{(c + d \tan(e + fx))^2}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="fricas")`

output

```
-1/2*(2*C*c^2*d - 2*B*c*d^2 + 2*A*d^3 - 2*((A - C)*c^3 + 2*B*c^2*d - (A -
C)*c*d^2)*f*x + (B*c^3 - 2*(A - C)*c^2*d - B*c*d^2 + (B*c^2*d - 2*(A - C)*
c*d^2 - B*d^3)*tan(f*x + e))*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e)
+ c^2)/(tan(f*x + e)^2 + 1)) - 2*(C*c^3 - B*c^2*d + A*c*d^2 + ((A - C)*c^2
*d + 2*B*c*d^2 - (A - C)*d^3)*f*x)*tan(f*x + e))/((c^4*d + 2*c^2*d^3 + d^5
)*f*tan(f*x + e) + (c^5 + 2*c^3*d^2 + c*d^4)*f)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.85 (sec) , antiderivative size = 4396, normalized size of antiderivative = 31.40

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**2,x)
```

output

```
Piecewise((zoo*x*(A + B*tan(e) + C*tan(e)**2)/tan(e)**2, Eq(c, 0) & Eq(d,
0) & Eq(f, 0)), ((A*x + B*log(tan(e + f*x)**2 + 1)/(2*f) - C*x + C*tan(e +
f*x)/f)/c**2, Eq(d, 0)), (-A*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**
2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*A*f*x*tan(e + f*x)/(4*d**2*f
*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + A*f*x/(4*d**2*f*t
an(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - A*tan(e + f*x)/(4*d
**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*A/(4*d**
2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + I*B*f*x*tan(e
+ f*x)**2/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f)
+ 2*B*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x)
- 4*d**2*f) - I*B*f*x/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x)
- 4*d**2*f) + I*B*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan
(e + f*x) - 4*d**2*f) + C*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 -
8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - 2*I*C*f*x*tan(e + f*x)/(4*d**2*f*tan
(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - C*f*x/(4*d**2*f*tan(e
+ f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) - 3*C*tan(e + f*x)/(4*d**
2*f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f) + 2*I*C/(4*d**2*
f*tan(e + f*x)**2 - 8*I*d**2*f*tan(e + f*x) - 4*d**2*f), Eq(c, -I*d)), (-A
*f*x*tan(e + f*x)**2/(4*d**2*f*tan(e + f*x)**2 + 8*I*d**2*f*tan(e + f*x) -
4*d**2*f) - 2*I*A*f*x*tan(e + f*x)/(4*d**2*f*tan(e + f*x)**2 + 8*I*d**...
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.46

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{2((A-C)c^2 + 2Bcd - (A-C)d^2)(fx+e)}{c^4 + 2c^2d^2 + d^4} - \frac{2(Bc^2 - 2(A-C)cd - Bd^2) \log(d \tan(fx+e) + c)}{c^4 + 2c^2d^2 + d^4} + \frac{(Bc^2 - 2(A-C)cd - Bd^2) \log(\tan(fx+e)^2 + 1)}{c^4 + 2c^2d^2 + d^4}$$

$$\frac{\hspace{10em}}{2f}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/2*(2*((A - C)*c^2 + 2*B*c*d - (A - C)*d^2)*(f*x + e)/(c^4 + 2*c^2*d^2 + d^4) - 2*(B*c^2 - 2*(A - C)*c*d - B*d^2)*log(d*tan(f*x + e) + c)/(c^4 + 2*c^2*d^2 + d^4) + (B*c^2 - 2*(A - C)*c*d - B*d^2)*log(tan(f*x + e)^2 + 1)/(c^4 + 2*c^2*d^2 + d^4) - 2*(C*c^2 - B*c*d + A*d^2)/(c^3*d + c*d^3 + (c^2*d^2 + d^4)*tan(f*x + e)))/f`

Giac [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.77

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{(Ac^2 - Cc^2 + 2Bcd - Ad^2 + Cd^2)(fx + e)}{c^4f + 2c^2d^2f + d^4f}$$

$$+ \frac{(Bc^2 - 2Acd + 2Ccd - Bd^2) \log(\tan(fx + e)^2 + 1)}{2(c^4f + 2c^2d^2f + d^4f)}$$

$$- \frac{(Bc^2d - 2Acd^2 + 2Ccd^2 - Bd^3) \log(|d \tan(fx + e) + c|)}{c^4df + 2c^2d^3f + d^5f}$$

$$- \frac{Cc^4 - Bc^3d + Ac^2d^2 + Cc^2d^2 - Bcd^3 + Ad^4}{(c^2 + d^2)^2(d \tan(fx + e) + c)df}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="giac")`

output

```
(A*c^2 - C*c^2 + 2*B*c*d - A*d^2 + C*d^2)*(f*x + e)/(c^4*f + 2*c^2*d^2*f +
d^4*f) + 1/2*(B*c^2 - 2*A*c*d + 2*C*c*d - B*d^2)*log(tan(f*x + e)^2 + 1)/
(c^4*f + 2*c^2*d^2*f + d^4*f) - (B*c^2*d - 2*A*c*d^2 + 2*C*c*d^2 - B*d^3)*
log(abs(d*tan(f*x + e) + c))/(c^4*d*f + 2*c^2*d^3*f + d^5*f) - (C*c^4 - B*
c^3*d + A*c^2*d^2 + C*c^2*d^2 - B*c*d^3 + A*d^4)/((c^2 + d^2)^2*(d*tan(f*x
+ e) + c)*d*f)
```

Mupad [B] (verification not implemented)

Time = 8.14 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.31

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{\ln(c + d \tan(e + fx)) (-B c^2 + (2A - 2C) cd + B d^2)}{f (c^4 + 2c^2 d^2 + d^4)}$$

$$- \frac{\ln(\tan(e + fx) - i) (A - C + B i)}{2 f (-c^2 i + 2cd + d^2 i)}$$

$$- \frac{\ln(\tan(e + fx) + i) (A i + B - C i)}{2 f (-c^2 + cd 2i + d^2)} - \frac{C c^2 - B cd + A d^2}{d f (c^2 + d^2) (c + d \tan(e + fx))}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/(c + d*tan(e + f*x))^2,x)
```

output

```
(log(c + d*tan(e + f*x))*(B*d^2 - B*c^2 + c*d*(2*A - 2*C))/(f*(c^4 + d^4
+ 2*c^2*d^2)) - (log(tan(e + f*x) - 1i)*(A + B*1i - C))/(2*f*(2*c*d - c^2*
1i + d^2*1i)) - (log(tan(e + f*x) + 1i)*(A*1i + B - C*1i))/(2*f*(c*d*2i -
c^2 + d^2)) - (A*d^2 + C*c^2 - B*c*d)/(d*f*(c^2 + d^2)*(c + d*tan(e + f*x)
))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 603, normalized size of antiderivative = 4.31

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x)
```

output

```
( - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*c**2*d**2 + log(tan(e + f*x)
**2 + 1)*tan(e + f*x)*b*c**3*d - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b*c
*d**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*c**3*d**2 - 2*log(tan(e +
f*x)**2 + 1)*a*c**3*d + log(tan(e + f*x)**2 + 1)*b*c**4 - log(tan(e + f*x)
**2 + 1)*b*c**2*d**2 + 2*log(tan(e + f*x)**2 + 1)*c**4*d + 4*log(tan(e + f
*x)*d + c)*tan(e + f*x)*a*c**2*d**2 - 2*log(tan(e + f*x)*d + c)*tan(e + f
*x)*b*c**3*d + 2*log(tan(e + f*x)*d + c)*tan(e + f*x)*b*c*d**3 - 4*log(tan(
e + f*x)*d + c)*tan(e + f*x)*c**3*d**2 + 4*log(tan(e + f*x)*d + c)*a*c**3*
d - 2*log(tan(e + f*x)*d + c)*b*c**4 + 2*log(tan(e + f*x)*d + c)*b*c**2*d*
*2 - 4*log(tan(e + f*x)*d + c)*c**4*d + 2*tan(e + f*x)*a*c**3*d*f*x + 2*ta
n(e + f*x)*a*c**2*d**2 - 2*tan(e + f*x)*a*c*d**3*f*x + 2*tan(e + f*x)*a*d*
*4 - 2*tan(e + f*x)*b*c**3*d + 4*tan(e + f*x)*b*c**2*d**2*f*x - 2*tan(e +
f*x)*b*c*d**3 + 2*tan(e + f*x)*c**5 - 2*tan(e + f*x)*c**4*d*f*x + 2*tan(e
+ f*x)*c**3*d**2 + 2*tan(e + f*x)*c**2*d**3*f*x + 2*a*c**4*f*x - 2*a*c**2*
d**2*f*x + 4*b*c**3*d*f*x - 2*c**5*f*x + 2*c**3*d**2*f*x)/(2*c*f*(tan(e +
f*x)*c**4*d + 2*tan(e + f*x)*c**2*d**3 + tan(e + f*x)*d**5 + c**5 + 2*c**3
*d**2 + c*d**4))
```


3.81 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$

Optimal result	900
Mathematica [A] (verified)	901
Rubi [A] (verified)	901
Maple [A] (verified)	904
Fricas [B] (verification not implemented)	905
Sympy [F(-2)]	906
Maxima [A] (verification not implemented)	907
Giac [B] (verification not implemented)	908
Mupad [B] (verification not implemented)	909
Reduce [B] (verification not implemented)	910

Optimal result

Integrand size = 45, antiderivative size = 293

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

$$= -\frac{(a(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) + b(2c(A - C)d - B(c^2 - d^2)))x}{(a^2 + b^2)(c^2 + d^2)^2}$$

$$+ \frac{b(Ab^2 - a(bB - aC)) \log(a \cos(e + fx) + b \sin(e + fx))}{(a^2 + b^2)(bc - ad)^2 f}$$

$$- \frac{(b(c^4C - 2Bc^3d + c^2(3A - C)d^2 + Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2))) \log(c \cos(e + fx) + d \sin(e + fx))}{(bc - ad)^2 (c^2 + d^2)^2 f}$$

$$+ \frac{c^2C - Bcd + Ad^2}{(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))}$$

output

```
- (a*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))+b*(2*c*(A-C)*d-B*(c^2-d^2)))*x/(a^2+b^2)/(c^2+d^2)^2+b*(A*b^2-a*(B*b-C*a))*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)/(-a*d+b*c)^2/f-(b*(c^4*C-2*B*c^3*d+c^2*(3*A-C)*d^2+Ad^4)-ad^2*(2*c*(A-C)*d-B*(c^2-d^2)))*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^2/(c^2+d^2)^2/f+(A*d^2-B*c*d+C*c^2)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))
```

Mathematica [A] (verified)

Time = 4.95 (sec) , antiderivative size = 561, normalized size of antiderivative = 1.91

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

$$\frac{(bc-ad) \left(Abc^2 - aBc^2 - bc^2C + 2aAcd + 2bBcd - 2acCd - Abd^2 + aBd^2 + bCd^2 + \frac{\sqrt{-b^2} (a(-c^2C + 2Bcd + Cd^2 + A(c^2 - d^2)) + b(2c(-A+C)d + B(c^2 - d^2)))}{b} \right)}{2(a^2 + b^2)(c^2 + d^2)}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^2),x]
```

output

```
((b*c - a*d)*(A*b*c^2 - a*B*c^2 - b*c^2*C + 2*a*A*c*d + 2*b*B*c*d - 2*a*c*C*d - A*b*d^2 + a*B*d^2 + b*C*d^2 + (Sqrt[-b^2]*(a*(-c^2*C) + 2*B*c*d + C*d^2 + A*(c^2 - d^2)) + b*(2*c*(-A + C)*d + B*(c^2 - d^2))))/b)*Log[Sqrt[-b^2] - b*Tan[e + f*x]]/(2*(a^2 + b^2)*(c^2 + d^2)) - (b*(A*b^2 + a*(-(b*B) + a*C))*(c^2 + d^2)*Log[a + b*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d)) + ((b*c - a*d)*(A*b*c^2 - a*B*c^2 - b*c^2*C + 2*a*A*c*d + 2*b*B*c*d - 2*a*c*C*d - A*b*d^2 + a*B*d^2 + b*C*d^2 + (Sqrt[-b^2]*(a*(c^2*C - 2*B*c*d - C*d^2 + A*(-c^2 + d^2)) + b*(2*c*(A - C)*d + B*(-c^2 + d^2)))))/b)*Log[Sqrt[-b^2] + b*Tan[e + f*x]]/(2*(a^2 + b^2)*(c^2 + d^2)) + ((b*(c^4*C - 2*B*c^3*d + c^2*(3*A - C)*d^2 + A*d^4) + a*d^2*(2*c*(-A + C)*d + B*(c^2 - d^2)))*Log[c + d*Tan[e + f*x]])/((b*c - a*d)*(c^2 + d^2)) - (A*d^2)/(c + d*Tan[e + f*x]) + (c*(-(c*C) + B*d))/(c + d*Tan[e + f*x])/((-b*c) + a*d)*(c^2 + d^2)*f)
```

Rubi [A] (verified)

Time = 2.24 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4132, 25, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx \\
& \quad \downarrow \text{4132} \\
& \int \frac{-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx + \\
& \quad \frac{(c^2 + d^2)(bc - ad)}{Ad^2 - Bcd + c^2C} \\
& \quad \frac{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} \\
& \quad \downarrow \text{25} \\
& \frac{Ad^2 - Bcd + c^2C}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} - \\
& \int \frac{-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx \\
& \quad \frac{(c^2 + d^2)(bc - ad)}{(c^2 + d^2)(bc - ad)} \\
& \quad \downarrow \text{3042} \\
& \frac{Ad^2 - Bcd + c^2C}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} - \\
& \int \frac{-b(Cc^2 - Bdc + Ad^2) \tan(e + fx)^2 - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))} dx \\
& \quad \frac{(c^2 + d^2)(bc - ad)}{(c^2 + d^2)(bc - ad)} \\
& \quad \downarrow \text{4134} \\
& \frac{Ad^2 - Bcd + c^2C}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} - \\
& - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx}{(a^2 + b^2)(bc - ad)} + \frac{(b(c^2 d^2(3A - C) + Ad^4 - 2Bc^3 d + c^4 C) - ad^2(2cd(A - C) - B(c^2 - d^2))) \int \frac{d - c \tan(e + fx)}{c + d \tan(e + fx)} dx}{(c^2 + d^2)(bc - ad)} \\
& \quad \frac{(c^2 + d^2)(bc - ad)}{(c^2 + d^2)(bc - ad)} \\
& \quad \downarrow \text{3042} \\
& \frac{Ad^2 - Bcd + c^2C}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} - \\
& - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx}{(a^2 + b^2)(bc - ad)} + \frac{(b(c^2 d^2(3A - C) + Ad^4 - 2Bc^3 d + c^4 C) - ad^2(2cd(A - C) - B(c^2 - d^2))) \int \frac{d - c \tan(e + fx)}{c + d \tan(e + fx)} dx}{(c^2 + d^2)(bc - ad)} \\
& \quad \frac{(c^2 + d^2)(bc - ad)}{(c^2 + d^2)(bc - ad)} \\
& \quad \downarrow \text{4013}
\end{aligned}$$

$$\frac{Ad^2 - Bcd + c^2C}{(c^2 + d^2)(bc - ad)} - \frac{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))}{f(a^2 + b^2)(bc - ad)} + \frac{x(bc - ad)(-A(c^2 - d^2) - 2Bcd + c^2C - Cd^2) + b(2cd(A - C) - B(c^2 - d^2))}{(a^2 + b^2)(c^2 + d^2)} + \frac{-\frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \log(a \cos(e + fx) + b \sin(e + fx))}{f(a^2 + b^2)(bc - ad)}}{(c^2 + d^2)(bc - ad)}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^2), x]
```

output

```
-((((b*c - a*d)*(a*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + b*(2*c*(A - C)*d - B*(c^2 - d^2)))*x)/((a^2 + b^2)*(c^2 + d^2)) - (b*(A*b^2 - a*(b*B - a*C))*(c^2 + d^2)*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f) + ((b*(c^4*C - 2*B*c^3*d + c^2*(3*A - C)*d^2 + A*d^4) - a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Log[c*Cos[e + f*x] + d*Sin[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f))/((b*c - a*d)*(c^2 + d^2)) + (c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4013

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.25

method	result
derivativedivides	$\frac{(-2Aacd - Abc^2 + Abd^2 + Baa^2 - Baa^2 - 2Bbcd + 2Cacd + Cbc^2 - Cbd^2) \ln(1 + \tan(fx + e)^2)}{2} + \frac{(Aa^2c^2 - Aa^2d^2 - 2Abcd + 2Bacd + Bbc^2 - Bbd^2 - Ca^2c^2 + Ca^2d^2 + 2Cbcd)cx}{(a^2 + b^2)(c^2 + d^2)^2}$
default	$\frac{(-2Aacd - Abc^2 + Abd^2 + Baa^2 - Baa^2 - 2Bbcd + 2Cacd + Cbc^2 - Cbd^2) \ln(1 + \tan(fx + e)^2)}{2} + \frac{(Aa^2c^2 - Aa^2d^2 - 2Abcd + 2Bacd + Bbc^2 - Bbd^2 - Ca^2c^2 + Ca^2d^2 + 2Cbcd)cx}{(a^2 + b^2)(c^2 + d^2)^2}$
norman	$\frac{(Aa^2c^2 - Aa^2d^2 - 2Abcd + 2Bacd + Bbc^2 - Bbd^2 - Ca^2c^2 + Ca^2d^2 + 2Cbcd)cx}{(a^2 + b^2)(c^4 + 2c^2d^2 + d^4)} + \frac{(Aa^2c^2 - Aa^2d^2 - 2Abcd + 2Bacd + Bbc^2 - Bbd^2 - Ca^2c^2 + Ca^2d^2 + 2Cbcd)cx}{(a^2 + b^2)(c^4 + 2c^2d^2 + d^4)}$
parallelrisc	Expression too large to display
risc	Expression too large to display

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^2,x,
method=_RETURNVERBOSE)`

output `1/f*(1/(a^2+b^2)/(c^2+d^2)^2*(1/2*(-2*A*a*c*d-A*b*c^2+A*b*d^2+B*a*c^2-B*a*d^2-2*B*b*c*d+2*C*a*c*d+C*b*c^2-C*b*d^2)*ln(1+tan(f*x+e)^2)+(A*a*c^2-A*a*d^2-2*A*b*c*d+2*B*a*c*d+B*b*c^2-B*b*d^2-C*a*c^2+C*a*d^2+2*C*b*c*d)*arctan(tan(f*x+e)))+(A*b^2-B*a*b+C*a^2)*b/(a*d-b*c)^2/(a^2+b^2)*ln(a+b*tan(f*x+e))+(2*A*a*c*d^3-3*A*b*c^2*d^2-A*b*d^4-B*a*c^2*d^2+B*a*d^4+2*B*b*c^3*d-2*C*a*c*d^3-C*b*c^4+C*b*c^2*d^2)/(c^2+d^2)^2/(a*d-b*c)^2*ln(c+d*tan(f*x+e))-(A*d^2-B*c*d+C*c^2)/(c^2+d^2)/(a*d-b*c)/(c+d*tan(f*x+e)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1275 vs. $2(291) = 582$.

Time = 0.91 (sec) , antiderivative size = 1275, normalized size of antiderivative = 4.35

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^2,x, algorithm="fricas")`

output

```

1/2*(2*(C*a^2*b + C*b^3)*c^3*d^2 - 2*(C*a^3 + B*a^2*b + C*a*b^2 + B*b^3)*c
^2*d^3 + 2*(B*a^3 + A*a^2*b + B*a*b^2 + A*b^3)*c*d^4 - 2*(A*a^3 + A*a*b^2)
*d^5 + 2*(((A - C)*a*b^2 + B*b^3)*c^5 - 2*((A - C)*a^2*b + (A - C)*b^3)*c^
4*d + ((A - C)*a^3 - 3*B*a^2*b + 3*(A - C)*a*b^2 - B*b^3)*c^3*d^2 + 2*(B*a
^3 + B*a*b^2)*c^2*d^3 - ((A - C)*a^3 + B*a^2*b)*c*d^4)*f*x + ((C*a^2*b - B
*a*b^2 + A*b^3)*c^5 + 2*(C*a^2*b - B*a*b^2 + A*b^3)*c^3*d^2 + (C*a^2*b - B
*a*b^2 + A*b^3)*c*d^4 + ((C*a^2*b - B*a*b^2 + A*b^3)*c^4*d + 2*(C*a^2*b -
B*a*b^2 + A*b^3)*c^2*d^3 + (C*a^2*b - B*a*b^2 + A*b^3)*d^5)*tan(f*x + e))*
log((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(tan(f*x + e)^2 + 1))
- ((C*a^2*b + C*b^3)*c^5 - 2*(B*a^2*b + B*b^3)*c^4*d + (B*a^3 + (3*A - C)*
a^2*b + B*a*b^2 + (3*A - C)*b^3)*c^3*d^2 - 2*((A - C)*a^3 + (A - C)*a*b^2)
*c^2*d^3 - (B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*c*d^4 + ((C*a^2*b + C*b^3)*
c^4*d - 2*(B*a^2*b + B*b^3)*c^3*d^2 + (B*a^3 + (3*A - C)*a^2*b + B*a*b^2 +
(3*A - C)*b^3)*c^2*d^3 - 2*((A - C)*a^3 + (A - C)*a*b^2)*c*d^4 - (B*a^3 -
A*a^2*b + B*a*b^2 - A*b^3)*d^5)*tan(f*x + e))*log((d^2*tan(f*x + e)^2 + 2
*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - 2*((C*a^2*b + C*b^3)*c^4*
d - (C*a^3 + B*a^2*b + C*a*b^2 + B*b^3)*c^3*d^2 + (B*a^3 + A*a^2*b + B*a*b
^2 + A*b^3)*c^2*d^3 - (A*a^3 + A*a*b^2)*c*d^4 - (((A - C)*a*b^2 + B*b^3)*c
^4*d - 2*((A - C)*a^2*b + (A - C)*b^3)*c^3*d^2 + ((A - C)*a^3 - 3*B*a^2*b
+ 3*(A - C)*a*b^2 - B*b^3)*c^2*d^3 + 2*(B*a^3 + B*a*b^2)*c*d^4 - ((A - ...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

= Exception raised: NotImplementedError

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+
e))**2,x)

```

output

```

Exception raised: NotImplementedError >> no valid subset found

```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 513, normalized size of antiderivative = 1.75

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

$$= \frac{2(((A-C)a+Bb)c^2+2(Ba-(A-C)b)cd-((A-C)a+Bb)d^2)(fx+e)}{(a^2+b^2)c^4+2(a^2+b^2)c^2d^2+(a^2+b^2)d^4} + \frac{2(Ca^2b-Bab^2+Ab^3)\log(b\tan(fx+e)+a)}{(a^2b^2+b^4)c^2-2(a^3b+ab^3)cd+(a^4+a^2b^2)d^2} - \frac{2(Cbc^4-2Bbc^3d-2abc^3)}{b^2c^6-2abc^3}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/2*(2*(((A - C)*a + B*b)*c^2 + 2*(B*a - (A - C)*b)*c*d - ((A - C)*a + B*b)*d^2)*(f*x + e)/((a^2 + b^2)*c^4 + 2*(a^2 + b^2)*c^2*d^2 + (a^2 + b^2)*d^4) + 2*(C*a^2*b - B*a*b^2 + A*b^3)*log(b*tan(f*x + e) + a)/((a^2*b^2 + b^4)*c^2 - 2*(a^3*b + a*b^3)*c*d + (a^4 + a^2*b^2)*d^2) - 2*(C*b*c^4 - 2*B*b*c^3*d - 2*(A - C)*a*c*d^3 + (B*a + (3*A - C)*b)*c^2*d^2 - (B*a - A*b)*d^4)*log(d*tan(f*x + e) + c)/(b^2*c^6 - 2*a*b*c^5*d - 4*a*b*c^3*d^3 - 2*a*b*c*d^5 + a^2*d^6 + (a^2 + 2*b^2)*c^4*d^2 + (2*a^2 + b^2)*c^2*d^4) + ((B*a - (A - C)*b)*c^2 - 2*((A - C)*a + B*b)*c*d - (B*a - (A - C)*b)*d^2)*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*c^4 + 2*(a^2 + b^2)*c^2*d^2 + (a^2 + b^2)*d^4) + 2*(C*c^2 - B*c*d + A*d^2)/(b*c^4 - a*c^3*d + b*c^2*d^2 - a*c*d^3 + (b*c^3*d - a*c^2*d^2 + b*c*d^3 - a*d^4)*tan(f*x + e))/f`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 673 vs. $2(291) = 582$.

Time = 0.63 (sec) , antiderivative size = 673, normalized size of antiderivative = 2.30

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

$$= \frac{(Aac^2 - Cac^2 + Bbc^2 + 2 Bacd - 2 Abcd + 2 Cbcd - Aad^2 + Cad^2 - Bbd^2)(fx + e)}{a^2c^4f + b^2c^4f + 2a^2c^2d^2f + 2b^2c^2d^2f + a^2d^4f + b^2d^4f}$$

$$+ \frac{(Bac^2 - Abc^2 + Cbc^2 - 2 Aacd + 2 Cacd - 2 Bbcd - Bad^2 + Abd^2 - Cbd^2) \log(\tan(fx + e)^2 + 1)}{2(a^2c^4f + b^2c^4f + 2a^2c^2d^2f + 2b^2c^2d^2f + a^2d^4f + b^2d^4f)}$$

$$+ \frac{(Ca^2b^2 - Bab^3 + Ab^4) \log(|b \tan(fx + e) + a|)}{a^2b^3c^2f + b^5c^2f - 2a^3b^2cdf - 2ab^4cdf + a^4bd^2f + a^2b^3d^2f}$$

$$- \frac{(Cbc^4d - 2Bbc^3d^2 + Bac^2d^3 + 3Abc^2d^3 - Cbc^2d^3 - 2Aacd^4 + 2Cacd^4 - Bad^5 + Abd^5) \log(|d \tan(fx + e) + c|)}{b^2c^6df - 2abc^5d^2f + a^2c^4d^3f + 2b^2c^4d^3f - 4abc^3d^4f + 2a^2c^2d^5f + b^2c^2d^5f - 2abcd^6f + a^2Cbc^5 - Cac^4d - Bbc^4d + Bac^3d^2 + Abc^3d^2 + Cbc^3d^2 - Aac^2d^3 - Cac^2d^3 - Bbc^2d^3 + Bacd^4 + Abcd^4}$$

$$+ \frac{Cbc^5 - Cac^4d - Bbc^4d + Bac^3d^2 + Abc^3d^2 + Cbc^3d^2 - Aac^2d^3 - Cac^2d^3 - Bbc^2d^3 + Bacd^4 + Abcd^4}{(bc - ad)^2(c^2 + d^2)^2(d \tan(fx + e) + c)f}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^2,x, algorithm="giac")
```

output

```
(A*a*c^2 - C*a*c^2 + B*b*c^2 + 2*B*a*c*d - 2*A*b*c*d + 2*C*b*c*d - A*a*d^2 + C*a*d^2 - B*b*d^2)*(f*x + e)/(a^2*c^4*f + b^2*c^4*f + 2*a^2*c^2*d^2*f + 2*b^2*c^2*d^2*f + a^2*d^4*f + b^2*d^4*f) + 1/2*(B*a*c^2 - A*b*c^2 + C*b*c^2 - 2*A*a*c*d + 2*C*a*c*d - 2*B*b*c*d - B*a*d^2 + A*b*d^2 - C*b*d^2)*log(tan(f*x + e)^2 + 1)/(a^2*c^4*f + b^2*c^4*f + 2*a^2*c^2*d^2*f + 2*b^2*c^2*d^2*f + a^2*d^4*f + b^2*d^4*f) + (C*a^2*b^2 - B*a*b^3 + A*b^4)*log(abs(b*tan(f*x + e) + a))/(a^2*b^3*c^2*f + b^5*c^2*f - 2*a^3*b^2*c*d*f - 2*a*b^4*c*d*f + a^4*b*d^2*f + a^2*b^3*d^2*f) - (C*b*c^4*d - 2*B*b*c^3*d^2 + B*a*c^2*d^3 + 3*A*b*c^2*d^3 - C*b*c^2*d^3 - 2*A*a*c*d^4 + 2*C*a*c*d^4 - B*a*d^5 + A*b*d^5)*log(abs(d*tan(f*x + e) + c))/(b^2*c^6*d*f - 2*a*b*c^5*d^2*f + a^2*c^4*d^3*f + 2*b^2*c^4*d^3*f - 4*a*b*c^3*d^4*f + 2*a^2*c^2*d^5*f + b^2*c^2*d^5*f - 2*a*b*c*d^6*f + a^2*d^7*f) + (C*b*c^5 - C*a*c^4*d - B*b*c^4*d + B*a*c^3*d^2 + A*b*c^3*d^2 + C*b*c^3*d^2 - A*a*c^2*d^3 - C*a*c^2*d^3 - B*b*c^2*d^3 + B*a*c*d^4 + A*b*c*d^4 - A*a*d^5)/((b*c - a*d)^2*(c^2 + d^2)^2*(d*tan(f*x + e) + c)*f)
```

Mupad [B] (verification not implemented)

Time = 61.57 (sec) , antiderivative size = 430, normalized size of antiderivative = 1.47

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

$$= \frac{\ln(\tan(e + fx) - i) (B - A i + C i)}{2 f (a c^2 - a d^2 - 2 b c d + b c^2 i - b d^2 i + a c d 2i)}$$

$$- \frac{\ln(\tan(e + fx) + i) (A i + B - C i)}{2 f (a d^2 - a c^2 + 2 b c d + b c^2 i - b d^2 i + a c d 2i)}$$

$$+ \frac{\ln(a + b \tan(e + fx)) (C a^2 b - B a b^2 + A b^3)}{f (a^4 d^2 - 2 a^3 b c d + a^2 b^2 c^2 + a^2 b^2 d^2 - 2 a b^3 c d + b^4 c^2)}$$

$$- \frac{\ln(c + d \tan(e + fx)) (C b c^4 - 2 B b c^3 d + (3 A b + B a - C b) c^2 d^2 + (2 C a - 2 A a) c d^3 + (A b - 3 A^2 a + 2 C a) d^4)}{f (a^2 c^4 d^2 + 2 a^2 c^2 d^4 + a^2 d^6 - 2 a b c^5 d - 4 a b c^3 d^3 - 2 a b c d^5 + b^2 c^6 + 2 b^2 c^4 d^2 + b^2 c^2 d^4)}$$

$$- \frac{C c^2 - B c d + A d^2}{f (a d - b c) (c^2 + d^2) (c + d \tan(e + fx))}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^2),x)
```

output

```
(log(tan(e + f*x) - 1i)*(B - A*1i + C*1i))/(2*f*(a*c^2 - a*d^2 + b*c^2*1i - b*d^2*1i + a*c*d*2i - 2*b*c*d)) - (log(tan(e + f*x) + 1i)*(A*1i + B - C*1i))/(2*f*(a*d^2 - a*c^2 + b*c^2*1i - b*d^2*1i + a*c*d*2i + 2*b*c*d)) + (log(a + b*tan(e + f*x))*(A*b^3 - B*a*b^2 + C*a^2*b))/(f*(a^4*d^2 + b^4*c^2 + a^2*b^2*c^2 + a^2*b^2*d^2 - 2*a*b^3*c*d - 2*a^3*b*c*d)) - (log(c + d*tan(e + f*x))*(d^4*(A*b - B*a) + c^2*d^2*(3*A*b + B*a - C*b) + C*b*c^4 - c*d^3*(2*A*a - 2*C*a) - 2*B*b*c^3*d))/(f*(a^2*d^6 + b^2*c^6 + 2*a^2*c^2*d^4 + a^2*c^4*d^2 + b^2*c^2*d^4 + 2*b^2*c^4*d^2 - 2*a*b*c*d^5 - 2*a*b*c^5*d - 4*a*b*c^3*d^3)) - (A*d^2 + C*c^2 - B*c*d)/(f*(a*d - b*c)*(c^2 + d^2)*(c + d*tan(e + f*x)))
```

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 2663, normalized size of antiderivative = 9.09

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^2,x)`

output

```
( - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**4*c**2*d**4 + 4*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a**3*b*c**3*d**3 + 2*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)*a**3*c**3*d**4 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*
b**2*c**4*d**2 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**2*c**2*d*
**4 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b*c**4*d**3 - log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)*a**2*b*c**2*d**5 + 4*log(tan(e + f*x)**2 + 1)*
tan(e + f*x)*a*b**3*c**3*d**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*
b**2*c**3*d**4 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**4*c**4*d**2 +
log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**3*c**6*d - log(tan(e + f*x)**2 +
1)*tan(e + f*x)*b**3*c**4*d**3 - 2*log(tan(e + f*x)**2 + 1)*a**4*c**3*d**3
+ 4*log(tan(e + f*x)**2 + 1)*a**3*b*c**4*d**2 + 2*log(tan(e + f*x)**2 + 1
)*a**3*c**4*d**3 - 2*log(tan(e + f*x)**2 + 1)*a**2*b**2*c**5*d - 2*log(tan
(e + f*x)**2 + 1)*a**2*b**2*c**3*d**3 - 3*log(tan(e + f*x)**2 + 1)*a**2*b*
c**5*d**2 - log(tan(e + f*x)**2 + 1)*a**2*b*c**3*d**4 + 4*log(tan(e + f*x)
**2 + 1)*a*b**3*c**4*d**2 + 2*log(tan(e + f*x)**2 + 1)*a*b**2*c**4*d**3 -
2*log(tan(e + f*x)**2 + 1)*b**4*c**5*d + log(tan(e + f*x)**2 + 1)*b**3*c**
7 - log(tan(e + f*x)**2 + 1)*b**3*c**5*d**2 + 2*log(tan(e + f*x)*b + a)*ta
n(e + f*x)*a**2*b*c**6*d + 4*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b*c
**4*d**3 + 2*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b*c**2*d**5 + 2*log
(tan(e + f*x)*b + a)*a**2*b*c**7 + 4*log(tan(e + f*x)*b + a)*a**2*b*c**...
```

3.82
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^2} dx$$

Optimal result	911
Mathematica [A] (verified)	912
Rubi [A] (verified)	913
Maple [A] (verified)	917
Fricas [B] (verification not implemented)	917
Sympy [F(-2)]	918
Maxima [B] (verification not implemented)	918
Giac [B] (verification not implemented)	919
Mupad [B] (verification not implemented)	920
Reduce [B] (verification not implemented)	921

Optimal result

Integrand size = 45, antiderivative size = 509

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2(c + d \tan(e + fx))^2} dx =$$

$$\frac{(a^2(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - b^2(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) + 2ab(2c(A - C)d - Bcd)}{(a^2 + b^2)^2(c^2 + d^2)^2}$$

$$+ \frac{b(3a^3bBd - 2a^4Cd + b^4(Bc - 2Ad) - a^2b^2(Bc + 4Ad) + ab^3(2Ac - 2cC + Bd)) \log(a \cos(e + fx) + c \tan(e + fx))}{(a^2 + b^2)^2(bc - ad)^3 f}$$

$$+ \frac{d(b(2c^4C - 3Bc^3d + 4Ac^2d^2 - Bcd^3 + 2Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2))) \log(c \cos(e + fx) + d \tan(e + fx))}{(bc - ad)^3(c^2 + d^2)^2 f}$$

$$- \frac{d(b^2c(cC - Bd) - abB(c^2 + d^2) + a^2(2c^2C - Bcd + Cd^2) + A(a^2d^2 + b^2(c^2 + 2d^2)))}{(a^2 + b^2)(bc - ad)^2(c^2 + d^2) f(c + d \tan(e + fx))}$$

$$- \frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad) f(a + b \tan(e + fx))(c + d \tan(e + fx))}$$

output

```

-(a^2*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-b^2*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))
+2*a*b*(2*c*(A-C)*d-B*(c^2-d^2)))*x/(a^2+b^2)^2/(c^2+d^2)^2+b*(3*a^3*b*B*d-2*a^4*C*d+b^4*(-2*A*d+B*c)-a^2*b^2*(4*A*d+B*c)+a*b^3*(2*A*c+B*d-2*C*c)
)*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)^2/(-a*d+b*c)^3/f+d*(b*(4*A*c^2*d^2+2*A*d^4-3*B*c^3*d-B*c*d^3+2*C*c^4)-a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^3/(c^2+d^2)^2/f-d*(b^2*c*(-B*d+C*c)-a*b*B*(c^2+d^2)+a^2*(-B*c*d+2*C*c^2+C*d^2)+A*(a^2*d^2+b^2*(c^2+2*d^2)))/(a^2+b^2)/(-a*d+b*c)^2/(c^2+d^2)/f/(c+d*tan(f*x+e))-(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))
    
```

Mathematica [A] (verified)

Time = 7.93 (sec) , antiderivative size = 984, normalized size of antiderivative = 1.93

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx$$

$$= - \frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))}$$

$$\frac{b(bc - ad)^2 \left(2aAbc^2 - a^2Bc^2 + b^2Bc^2 - 2abc^2C + 2a^2Acd - 2Ab^2cd + 4abBcd - 2a^2cCd + 2b^2cCd - 2aAbd^2 + a^2Bd^2 - b^2Bd^2 + 2abCd^2 - \frac{\sqrt{-b^2}(a^2(c^2C - 2Bcd))}{2(a^2 + b^2)(c^2 + d^2)} \right)}{2(a^2 + b^2)(c^2 + d^2)}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x]^2),x]
    
```

output

```

-((A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])*
(c + d*Tan[e + f*x]))) - (-(((b*(b*c - a*d)^2*(2*a*A*b*c^2 - a^2*B*c^2 + b
^2*B*c^2 - 2*a*b*c^2*C + 2*a^2*A*c*d - 2*A*b^2*c*d + 4*a*b*B*c*d - 2*a^2*c
*C*d + 2*b^2*c*C*d - 2*a*A*b*d^2 + a^2*B*d^2 - b^2*B*d^2 + 2*a*b*C*d^2 - (
Sqrt[-b^2]*(a^2*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b^2*(c^2*C - 2
*B*c*d - C*d^2 - A*(c^2 - d^2)) + 2*a*b*(2*c*(A - C)*d - B*(c^2 - d^2)))))/
b)*Log[Sqrt[-b^2] - b*Tan[e + f*x]]/(2*(a^2 + b^2)*(c^2 + d^2)) - (b^2*(c
^2 + d^2)*(3*a^3*b*B*d - 2*a^4*C*d + b^4*(B*c - 2*A*d) - a^2*b^2*(B*c + 4*
A*d) + a*b^3*(2*A*c - 2*c*C + B*d))*Log[a + b*Tan[e + f*x]]/((a^2 + b^2)*
(b*c - a*d)) + (b*(b*c - a*d)^2*(2*a*A*b*c^2 - a^2*B*c^2 + b^2*B*c^2 - 2*a
*b*c^2*C + 2*a^2*A*c*d - 2*A*b^2*c*d + 4*a*b*B*c*d - 2*a^2*c*C*d + 2*b^2*c
*C*d - 2*a*A*b*d^2 + a^2*B*d^2 - b^2*B*d^2 + 2*a*b*C*d^2 + (Sqrt[-b^2]*(a
^2*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b^2*(c^2*C - 2*B*c*d - C*d^2
- A*(c^2 - d^2)) + 2*a*b*(2*c*(A - C)*d - B*(c^2 - d^2)))))/b)*Log[Sqrt[-b
^2] + b*Tan[e + f*x]]/(2*(a^2 + b^2)*(c^2 + d^2)) - (b*(a^2 + b^2)*d*(b*(
2*c^4*C - 3*B*c^3*d + 4*A*c^2*d^2 - B*c*d^3 + 2*A*d^4) - a*d^2*(2*c*(A - C
)*d - B*(c^2 - d^2)))*Log[c + d*Tan[e + f*x]]/((b*c - a*d)*(c^2 + d^2)))/
(b*(-(b*c) + a*d)*(c^2 + d^2)*f)) - (-((c*(-2*c*(A*b^2 - a*(b*B - a*C))*d +
(A*b - a*B - b*C)*d*(b*c - a*d))) + d^2*(2*A*b^2*d - a*A*(b*c - a*d) - (b
*B - a*C)*(b*c + a*d)))/((- (b*c) + a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*...

```

Rubi [A] (verified)

Time = 4.77 (sec) , antiderivative size = 566, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4132, 3042, 4132, 25, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx$$

↓ 4132

$$\int \frac{2Adb^2+2(Ab^2-a(bB-aC))d \tan^2(e+fx)-aA(bc-ad)-(bB-aC)(bc+ad)+(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$$

$$\frac{(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$\frac{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))}{}$$

3042

$$\int \frac{2Adb^2+2(Ab^2-a(bB-aC))d \tan(e+fx)^2-aA(bc-ad)-(bB-aC)(bc+ad)+(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$$

$$\frac{(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$\frac{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))}{}$$

4132

$$\int -\frac{d^2(Ac-Cc+Bd)a^3-2Abd(c^2+d^2)a^2-b^2(Cc^3+2Cd^2c-Bd^3-A(c^3+2d^2c))a-bd(Ad^2a^2+(2Cc^2-Bdc+Cd^2)a^2-bB(c^2+d^2)a+b^2c(cC-Bd)+Ab^2(c^2+d^2)(bc-ad))}{(a+b \tan(e+fx))(c+d \tan(e+fx))} dx$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))}$$

25

$$\frac{d(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))} - \int \frac{d^2(Ac-Cc+Bd)a^3-2Abd(c^2+d^2)a^2-b^2(Cc^3+2Cd^2c-Bd^3)}{(a+b \tan(e+fx))(c+d \tan(e+fx))} dx$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))}$$

3042

$$\frac{d(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))} - \int \frac{d^2(Ac-Cc+Bd)a^3-2Abd(c^2+d^2)a^2-b^2(Cc^3+2Cd^2c-Bd^3)}{(a+b \tan(e+fx))(c+d \tan(e+fx))} dx$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))}$$

4134

$$\frac{d(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))} - \frac{d(a^2+b^2)(b(4Ac^2d^2+2Ad^4-3Bc^3d-Bcd^3+2c^4C)-ad^2(2cd))}{(c^2+d^2)(bc-ad)}$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))}$$

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4013 $\text{Int}[\text{((c}_.) + (\text{d}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) / ((\text{a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c}/(\text{b*f}))\text{Log}[\text{RemoveContent}[\text{a*Cos}[\text{e} + \text{f*x}] + \text{b*Sin}[\text{e} + \text{f*x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{NeQ}[\text{b*c} - \text{a*d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{EqQ}[\text{a*c} + \text{b*d}, 0]$
- rule 4132 $\text{Int}[\text{((a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])^{\text{m}} * ((\text{c}_.) + (\text{d}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])^{\text{n}} * ((\text{A}_.) + (\text{B}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) + (\text{C}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]^2}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{A*b}^2 - \text{a*(b*B} - \text{a*C})) * (\text{a} + \text{b*Tan}[\text{e} + \text{f*x}])^{\text{m} + 1} * ((\text{c} + \text{d*Tan}[\text{e} + \text{f*x}])^{\text{n} + 1} / (\text{f*(m} + 1) * (\text{b*c} - \text{a*d}) * (\text{a}^2 + \text{b}^2))), \text{x}] + \text{Simp}[1 / ((\text{m} + 1) * (\text{b*c} - \text{a*d}) * (\text{a}^2 + \text{b}^2)) \quad \text{Int}[(\text{a} + \text{b*Tan}[\text{e} + \text{f*x}])^{\text{m} + 1} * (\text{c} + \text{d*Tan}[\text{e} + \text{f*x}])^{\text{n}} * \text{Simp}[\text{A*(a*(b*c} - \text{a*d)*(m} + 1) - \text{b}^2 * \text{d*(m} + \text{n} + 2) + (\text{b*B} - \text{a*C}) * (\text{b*c*(m} + 1) + \text{a*d*(n} + 1)) - (\text{m} + 1) * (\text{b*c} - \text{a*d}) * (\text{A*b} - \text{a*B} - \text{b*C}) * \text{Tan}[\text{e} + \text{f*x}] - \text{d*(A*b}^2 - \text{a*(b*B} - \text{a*C})) * (\text{m} + \text{n} + 2) * \text{Tan}[\text{e} + \text{f*x}]^2}, \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}, \text{n}\}, \text{x}] \&\& \text{NeQ}[\text{b*c} - \text{a*d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{LtQ}[\text{m}, -1] \&\& !(\text{ILtQ}[\text{n}, -1] \&\& (!\text{IntegerQ}[\text{m}] || (\text{EqQ}[\text{c}, 0] \&\& \text{NeQ}[\text{a}, 0])))$
- rule 4134 $\text{Int}[\text{((A}_.) + (\text{B}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) + (\text{C}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]^2} / \text{((a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) * ((\text{c}_.) + (\text{d}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) * (\text{x}_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{a*(A*c} - \text{c*C} + \text{B*d}) + \text{b*(B*c} - \text{A*d} + \text{C*d})) * (\text{x} / ((\text{a}^2 + \text{b}^2) * (\text{c}^2 + \text{d}^2))), \text{x}] + (\text{Simp}[(\text{A*b}^2 - \text{a*b*B} + \text{a}^2 * \text{C}) / ((\text{b*c} - \text{a*d}) * (\text{a}^2 + \text{b}^2)) \quad \text{Int}[(\text{b} - \text{a*Tan}[\text{e} + \text{f*x}]) / (\text{a} + \text{b*Tan}[\text{e} + \text{f*x}]), \text{x}], \text{x}] - \text{Simp}[(\text{c}^2 * \text{C} - \text{B*c*d} + \text{A*d}^2) / ((\text{b*c} - \text{a*d}) * (\text{c}^2 + \text{d}^2)) \quad \text{Int}[(\text{d} - \text{c*Tan}[\text{e} + \text{f*x}]) / (\text{c} + \text{d*Tan}[\text{e} + \text{f*x}]), \text{x}], \text{x}]) \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}] \&\& \text{NeQ}[\text{b*c} - \text{a*d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0]$

Maple [A] (verified)

Time = 1.50 (sec) , antiderivative size = 577, normalized size of antiderivative = 1.13

method	result
derivativedivides	$\frac{(-2Aa^2cd - 2Aabc^2 + 2Aabd^2 + 2Ab^2cd + Ba^2c^2 - Ba^2d^2 - 4Babcd - Bb^2c^2 + Bb^2d^2 + 2Ca^2cd + 2Cab c^2 - 2Cab d^2 - 2Cb^2cd) \ln(1 + \frac{c+d \tan(fx+e)}{a+b \tan(fx+e)})}{2}$
default	$\frac{(-2Aa^2cd - 2Aabc^2 + 2Aabd^2 + 2Ab^2cd + Ba^2c^2 - Ba^2d^2 - 4Babcd - Bb^2c^2 + Bb^2d^2 + 2Ca^2cd + 2Cab c^2 - 2Cab d^2 - 2Cb^2cd) \ln(1 + \frac{c+d \tan(fx+e)}{a+b \tan(fx+e)})}{2}$
norman	Expression too large to display
risch	Expression too large to display
parallelrisc	Expression too large to display

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^2,
x,method=_RETURNVERBOSE)
```

output

```
1/f*(1/(a^2+b^2)^2/(c^2+d^2)^2*(1/2*(-2*A*a^2*c*d-2*A*a*b*c^2+2*A*a*b*d^2+
2*A*b^2*c*d+B*a^2*c^2-B*a^2*d^2-4*B*a*b*c*d-B*b^2*c^2+B*b^2*d^2+2*C*a^2*c*
d+2*C*a*b*c^2-2*C*a*b*d^2-2*C*b^2*c*d)*ln(1+tan(f*x+e)^2)+(A*a^2*c^2-A*a^2
*d^2-4*A*a*b*c*d-A*b^2*c^2+A*b^2*d^2+2*B*a^2*c*d+2*B*a*b*c^2-2*B*a*b*d^2-2
*B*b^2*c*d-C*a^2*c^2+C*a^2*d^2+4*C*a*b*c*d+C*b^2*c^2-C*b^2*d^2)*arctan(tan
(f*x+e)))+b*(4*A*a^2*b^2*d-2*A*a*b^3*c+2*A*b^4*d-3*B*a^3*b*d+B*a^2*b^2*c-B
*a*b^3*d-B*b^4*c+2*C*a^4*d+2*C*a*b^3*c)/(a*d-b*c)^3/(a^2+b^2)^2*ln(a+b*tan
(f*x+e))-(A*b^2-B*a*b+C*a^2)*b/(a*d-b*c)^2/(a^2+b^2)/(a+b*tan(f*x+e))+d*(2
*A*a*c*d^3-4*A*b*c^2*d^2-2*A*b*d^4-B*a*c^2*d^2+B*a*d^4+3*B*b*c^3*d+B*b*c*d
^3-2*C*a*c*d^3-2*C*b*c^4)/(c^2+d^2)^2/(a*d-b*c)^3*ln(c+d*tan(f*x+e))-(A*d^
2-B*c*d+C*c^2)*d/(c^2+d^2)/(a*d-b*c)^2/(c+d*tan(f*x+e)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4174 vs. 2(512) = 1024.

Time = 3.33 (sec) , antiderivative size = 4174, normalized size of antiderivative = 8.20

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^2,x, algorithm="fricas")`

output Too large to include

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx$$

= Exception raised: NotImplementedError

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**2/(c+d*tan(f*x+e))**2,x)`

output Exception raised: NotImplementedError >> no valid subset found

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1185 vs. $2(512) = 1024$.

Time = 0.17 (sec) , antiderivative size = 1185, normalized size of antiderivative = 2.33

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^2,x, algorithm="maxima")`

output

```

1/2*(2*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c^2 + 2*(B*a^2 - 2*(A - C)*a
*b - B*b^2)*c*d - ((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*d^2)*(f*x + e)/((a
^4 + 2*a^2*b^2 + b^4)*c^4 + 2*(a^4 + 2*a^2*b^2 + b^4)*c^2*d^2 + (a^4 + 2*a
^2*b^2 + b^4)*d^4) - 2*((B*a^2*b^3 - 2*(A - C)*a*b^4 - B*b^5)*c + (2*C*a^4
*b - 3*B*a^3*b^2 + 4*A*a^2*b^3 - B*a*b^4 + 2*A*b^5)*d)*log(b*tan(f*x + e)
+ a)/((a^4*b^3 + 2*a^2*b^5 + b^7)*c^3 - 3*(a^5*b^2 + 2*a^3*b^4 + a*b^6)*c^
2*d + 3*(a^6*b + 2*a^4*b^3 + a^2*b^5)*c*d^2 - (a^7 + 2*a^5*b^2 + a^3*b^4)*
d^3) + 2*(2*C*b*c^4*d - 3*B*b*c^3*d^2 + (B*a + 4*A*b)*c^2*d^3 - (2*(A - C)
*a + B*b)*c*d^4 - (B*a - 2*A*b)*d^5)*log(d*tan(f*x + e) + c)/(b^3*c^7 - 3*
a*b^2*c^6*d + 3*a^2*b*c*d^6 - a^3*d^7 + (3*a^2*b + 2*b^3)*c^5*d^2 - (a^3 +
6*a*b^2)*c^4*d^3 + (6*a^2*b + b^3)*c^3*d^4 - (2*a^3 + 3*a*b^2)*c^2*d^5) +
((B*a^2 - 2*(A - C)*a*b - B*b^2)*c^2 - 2*((A - C)*a^2 + 2*B*a*b - (A - C)
*b^2)*c*d - (B*a^2 - 2*(A - C)*a*b - B*b^2)*d^2)*log(tan(f*x + e)^2 + 1)/((
a^4 + 2*a^2*b^2 + b^4)*c^4 + 2*(a^4 + 2*a^2*b^2 + b^4)*c^2*d^2 + (a^4 + 2
*a^2*b^2 + b^4)*d^4) - 2*((C*a^2*b - B*a*b^2 + A*b^3)*c^3 + (C*a^3 + C*a*b
^2)*c^2*d - (B*a^3 - C*a^2*b + 2*B*a*b^2 - A*b^3)*c*d^2 + (A*a^3 + A*a*b^2)
*d^3 + ((2*C*a^2*b - B*a*b^2 + (A + C)*b^3)*c^2*d - (B*a^2*b + B*b^3)*c*d
^2 + ((A + C)*a^2*b - B*a*b^2 + 2*A*b^3)*d^3)*tan(f*x + e))/((a^3*b^2 + a
b^4)*c^5 - 2*(a^4*b + a^2*b^3)*c^4*d + (a^5 + 2*a^3*b^2 + a*b^4)*c^3*d^2 -
2*(a^4*b + a^2*b^3)*c^2*d^3 + (a^5 + a^3*b^2)*c*d^4 + ((a^2*b^3 + b^5)...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1674 vs. $2(512) = 1024$.

Time = 0.86 (sec) , antiderivative size = 1674, normalized size of antiderivative = 3.29

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+
e))^2,x, algorithm="giac")

```

output

```
(A*a^2*c^2 - C*a^2*c^2 + 2*B*a*b*c^2 - A*b^2*c^2 + C*b^2*c^2 + 2*B*a^2*c*d
- 4*A*a*b*c*d + 4*C*a*b*c*d - 2*B*b^2*c*d - A*a^2*d^2 + C*a^2*d^2 - 2*B*a
*b*d^2 + A*b^2*d^2 - C*b^2*d^2)*(f*x + e)/(a^4*c^4*f + 2*a^2*b^2*c^4*f + b
^4*c^4*f + 2*a^4*c^2*d^2*f + 4*a^2*b^2*c^2*d^2*f + 2*b^4*c^2*d^2*f + a^4*d
^4*f + 2*a^2*b^2*d^4*f + b^4*d^4*f) + 1/2*(B*a^2*c^2 - 2*A*a*b*c^2 + 2*C*a
*b*c^2 - B*b^2*c^2 - 2*A*a^2*c*d + 2*C*a^2*c*d - 4*B*a*b*c*d + 2*A*b^2*c*d
- 2*C*b^2*c*d - B*a^2*d^2 + 2*A*a*b*d^2 - 2*C*a*b*d^2 + B*b^2*d^2)*log(ta
n(f*x + e)^2 + 1)/(a^4*c^4*f + 2*a^2*b^2*c^4*f + b^4*c^4*f + 2*a^4*c^2*d^2
*f + 4*a^2*b^2*c^2*d^2*f + 2*b^4*c^2*d^2*f + a^4*d^4*f + 2*a^2*b^2*d^4*f +
b^4*d^4*f) - (B*a^2*b^4*c - 2*A*a*b^5*c + 2*C*a*b^5*c - B*b^6*c + 2*C*a^4
*b^2*d - 3*B*a^3*b^3*d + 4*A*a^2*b^4*d - B*a*b^5*d + 2*A*b^6*d)*log(abs(b*
tan(f*x + e) + a))/(a^4*b^4*c^3*f + 2*a^2*b^6*c^3*f + b^8*c^3*f - 3*a^5*b^
3*c^2*d*f - 6*a^3*b^5*c^2*d*f - 3*a*b^7*c^2*d*f + 3*a^6*b^2*c*d^2*f + 6*a^
4*b^4*c*d^2*f + 3*a^2*b^6*c*d^2*f - a^7*b*d^3*f - 2*a^5*b^3*d^3*f - a^3*b^
5*d^3*f) + (2*C*b*c^4*d^2 - 3*B*b*c^3*d^3 + B*a*c^2*d^4 + 4*A*b*c^2*d^4 -
2*A*a*c*d^5 + 2*C*a*c*d^5 - B*b*c*d^5 - B*a*d^6 + 2*A*b*d^6)*log(abs(d*tan
(f*x + e) + c))/(b^3*c^7*d*f - 3*a*b^2*c^6*d^2*f + 3*a^2*b*c^5*d^3*f + 2*b
^3*c^5*d^3*f - a^3*c^4*d^4*f - 6*a*b^2*c^4*d^4*f + 6*a^2*b*c^3*d^5*f + b^3
*c^3*d^5*f - 2*a^3*c^2*d^6*f - 3*a*b^2*c^2*d^6*f + 3*a^2*b*c*d^7*f - a^3*d
^8*f) - (C*a^4*b*c^5 - B*a^3*b^2*c^5 + A*a^2*b^3*c^5 + C*a^2*b^3*c^5 - ...
```

Mupad [B] (verification not implemented)

Time = 22.17 (sec) , antiderivative size = 73684, normalized size of antiderivative = 144.76

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^2*(c + d
*tan(e + f*x))^2),x)
```

output

```
(symsum(log((tan(e + f*x))*(4*A^3*a^3*b^4*d^7 + B^3*a^2*b^5*d^7 + 4*A^3*b^7*c^3*d^4 + 2*C^3*a^5*b^2*d^7 + B^3*b^7*c^2*d^5 + 2*C^3*b^7*c^5*d^2 + 4*A^2*B*b^7*d^7 - 4*B^3*a^2*b^5*c^2*d^5 - 3*B^3*a^2*b^5*c^4*d^3 + 10*B^3*a^3*b^4*c^3*d^4 - 3*B^3*a^4*b^3*c^2*d^5 - 4*A*B^2*a*b^6*d^7 - 4*A*B^2*b^7*c*d^6 + 2*B^3*a*b^6*c*d^6 - 6*A*B^2*a^3*b^4*d^7 + 8*A^2*B*a^2*b^5*d^7 - 3*A^2*B*a^4*b^3*d^7 + 4*A*C^2*a^3*b^4*d^7 - 4*A*C^2*a^5*b^2*d^7 - 8*A^2*C*a^3*b^4*d^7 + 2*A^2*C*a^5*b^2*d^7 - 6*A*B^2*b^7*c^3*d^4 - 3*B*C^2*a^4*b^3*d^7 + 8*A^2*B*b^7*c^2*d^5 - 3*A^2*B*b^7*c^4*d^3 + 4*A*C^2*b^7*c^3*d^4 - 4*A*C^2*b^7*c^5*d^2 - 8*A^2*C*b^7*c^3*d^4 + 2*A^2*C*b^7*c^5*d^2 - 3*B*C^2*b^7*c^4*d^3 - 4*A^3*a*b^6*c^2*d^5 - 4*A^3*a^2*b^5*c*d^6 + 6*B^3*a*b^6*c^3*d^4 + 6*B^3*a^3*b^4*c*d^6 - 2*C^3*a*b^6*c^4*d^3 - 2*C^3*a^4*b^3*c*d^6 - 10*A*B^2*a^2*b^5*c^3*d^4 - 10*A*B^2*a^3*b^4*c^2*d^5 + 18*A^2*B*a^2*b^5*c^2*d^5 + 2*B*C^2*a^2*b^5*c^2*d^5 + 4*B*C^2*a^4*b^3*c^4*d^3 + 2*B^2*C*a^2*b^5*c^3*d^4 + 2*B^2*C*a^2*b^5*c^5*d^2 + 2*B^2*C*a^3*b^4*c^2*d^5 - 6*B^2*C*a^3*b^4*c^4*d^3 - 6*B^2*C*a^4*b^3*c^3*d^4 + 2*B^2*C*a^5*b^2*c^2*d^5 + 10*A*B*C*a^4*b^3*d^7 + 10*A*B*C*b^7*c^4*d^3 - 8*A^2*B*a*b^6*c*d^6 - 2*A*B^2*a*b^6*c^2*d^5 + 6*A*B^2*a*b^6*c^4*d^3 - 2*A*B^2*a^2*b^5*c*d^6 + 6*A*B^2*a^4*b^3*c*d^6 - 4*A^2*B*a*b^6*c^3*d^4 - 4*A^2*B*a^3*b^4*c*d^6 - 4*A*C^2*a*b^6*c^2*d^5 + 4*A*C^2*a*b^6*c^4*d^3 - 4*A*C^2*a^2*b^5*c*d^6 + 4*A*C^2*a^4*b^3*c*d^6 + 8*A^2*C*a*b^6*c^2*d^5 - 2*A^2*C*a*b^6*c^4*d^3 + 8*A^2*C*a^2*b^5*c*d^6 - 2*A^2*...
```

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 11926, normalized size of antiderivative = 23.43

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^2, x)
```

output

```
( - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**7*b*c*d**6 + 3*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a**6*b**2*c**2*d**5 + log(tan(e + f*x)**2
+ 1)*tan(e + f*x)**2*a**6*b**2*d**7 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f
*x)**2*a**6*b*c**2*d**6 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*
b**3*c**3*d**4 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**3*c*d*
*6 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**2*c**3*d**5 - 2*lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*b**2*c*d**7 - 4*log(tan(e + f*
x)**2 + 1)*tan(e + f*x)**2*a**4*b**4*c**4*d**3 + 3*log(tan(e + f*x)**2 + 1
)*tan(e + f*x)**2*a**4*b**4*c**2*d**5 + log(tan(e + f*x)**2 + 1)*tan(e + f
*x)**2*a**4*b**4*d**7 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*b*
*3*c**4*d**4 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*b**3*c**2*d
**6 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**5*c**3*d**4 - 2*l
og(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*b**5*c*d**6 + 4*log(tan(e + f
*x)**2 + 1)*tan(e + f*x)**2*a**3*b**4*c**5*d**3 + 4*log(tan(e + f*x)**2 +
1)*tan(e + f*x)**2*a**3*b**4*c**3*d**5 + log(tan(e + f*x)**2 + 1)*tan(e +
f*x)**2*a**2*b**6*c**6*d - 5*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2
*b**6*c**4*d**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**5*c**
6*d**2 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**5*c**4*d**4 +
2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**7*c**3*d**4 - 2*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a*b**6*c**7*d - 2*log(tan(e + f*x)**2 + ...
```

3.83
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^3(c+d \tan(e+fx))^2} dx$$

Optimal result	923
Mathematica [B] (verified)	924
Rubi [A] (verified)	925
Maple [A] (verified)	929
Fricas [B] (verification not implemented)	930
Sympy [F(-2)]	931
Maxima [B] (verification not implemented)	931
Giac [B] (verification not implemented)	932
Mupad [B] (verification not implemented)	933
Reduce [B] (verification not implemented)	934

Optimal result

Integrand size = 45, antiderivative size = 841

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3(c + d \tan(e + fx))^2} dx =$$

$$\frac{(a^3(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) - 3ab^2(c^2C - 2Bcd - Cd^2 - A(c^2 - d^2)) + 3a^2b(2c(A - C)d - (a^2 + b^2)^3(c^2 + d^2)^2)}{b(6a^5bBd^2 - 3a^6Cd^2 - a^4b^2d(4Bc + (10A - C)d) - b^6(c(cC - 2Bd) - A(c^2 - 3d^2)) + ab^5(2c(A - C) - b^2(c^2 + d^2)))}$$

$$\frac{d^2(b(3c^4C - 4Bc^3d + c^2(5A + C)d^2 - 2Bcd^3 + 3Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2))) \log(c \cos(e + fx))}{(bc - ad)^4(c^2 + d^2)^2 f}$$

$$\frac{d(3a^3bBd(c^2 + d^2) + ab^3(2Ac - 2cC + Bd)(c^2 + d^2) - a^4d(3c^2C - Bcd + (A + 2C)d^2) - a^2b^2(Bc^3 + Bd^3))}{(a^2 + b^2)^2(bc - ad)^3(c^2 + d^2)f(c + d \tan(e + fx))}$$

$$\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2(c + d \tan(e + fx))}$$

$$\frac{5a^3bBd - 3a^4Cd + b^4(2Bc - 3Ad) + ab^3(4Ac - 4cC + Bd) - a^2b^2(2Bc + (7A - C)d)}{2(a^2 + b^2)^2(bc - ad)^2f(a + b \tan(e + fx))(c + d \tan(e + fx))}$$

output

```

-(a^3*(c^2*C-2*B*c*d-C*d^2-A*(c^2-d^2))-3*a*b^2*(c^2*C-2*B*c*d-C*d^2-A*(c^
2-d^2))+3*a^2*b*(2*c*(A-C)*d-B*(c^2-d^2))-b^3*(2*c*(A-C)*d-B*(c^2-d^2)))*x
/(a^2+b^2)^3/(c^2+d^2)^2-b*(6*a^5*b*B*d^2-3*a^6*C*d^2-a^4*b^2*d*(4*B*c+(10
*A-C)*d)-b^6*(c*(-2*B*d+C*c)-A*(c^2-3*d^2))+a*b^5*(2*c*(A-C)*d-B*(3*c^2-d^
2))+3*a^2*b^4*(c*(2*B*d+C*c)-A*(c^2+3*d^2))+a^3*b^3*(10*c*(A-C)*d+B*(c^2+3
*d^2)))*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)^3/(-a*d+b*c)^4/f-d^2*(b*(3
*c^4*C-4*B*c^3*d+c^2*(5*A+C)*d^2-2*B*c*d^3+3*A*d^4)-a*d^2*(2*c*(A-C)*d-B*(
c^2-d^2)))*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^4/(c^2+d^2)^2/f-d*(3*a
^3*b*B*d*(c^2+d^2)+a*b^3*(2*A*c+B*d-2*C*c)*(c^2+d^2)-a^4*d*(3*c^2*C-B*c*d+
(A+2*C)*d^2)-a^2*b^2*(4*A*c^2*d+6*A*d^3+B*c^3-B*c*d^2+2*C*c^2*d)-b^4*(d*(2
*A*c^2+3*A*d^2+C*c^2)-B*(c^3+2*c*d^2)))/(a^2+b^2)^2/(-a*d+b*c)^3/(c^2+d^2)
/f/(c+d*tan(f*x+e))-1/2*(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*ta
n(f*x+e))^2/(c+d*tan(f*x+e))-1/2*(5*a^3*b*B*d-3*a^4*C*d+b^4*(-3*A*d+2*B*c)
+a*b^3*(4*A*c+B*d-4*C*c)-a^2*b^2*(2*B*c+(7*A-C)*d))/(a^2+b^2)^2/(-a*d+b*c)
^2/f/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1758 vs. $2(841) = 1682$.

Time = 7.83 (sec) , antiderivative size = 1758, normalized size of antiderivative = 2.09

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^3*
(c + d*Tan[e + f*x])^2),x]

```

output

```

-1/2*(A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x
])^2*(c + d*Tan[e + f*x])) - (((-((a*(-3*a*(A*b^2 - a*(b*B - a*C))*d + 2*b
*(A*b - a*B - b*C)*(b*c - a*d))) + b^2*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b
*B - a*C)*(2*b*c + a*d)))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])*
(c + d*Tan[e + f*x]))) - (((-(((b*c - a*d)^3*(-(b^2*(-3*a^2*A*b*c^2 + A*b
^3*c^2 + a^3*B*c^2 - 3*a*b^2*B*c^2 + 3*a^2*b*c^2*C - b^3*c^2*C - 2*a^3*A*c
*d + 6*a*A*b^2*c*d - 6*a^2*b*B*c*d + 2*b^3*B*c*d + 2*a^3*c*C*d - 6*a*b^2*c
*C*d + 3*a^2*A*b*d^2 - A*b^3*d^2 - a^3*B*d^2 + 3*a*b^2*B*d^2 - 3*a^2*b*C*d
^2 + b^3*C*d^2)) + Sqrt[-b^2]*(a^3*A*b*c^2 - 3*a*A*b^3*c^2 + 3*a^2*b^2*B*c
^2 - b^4*B*c^2 - a^3*b*c^2*C + 3*a*b^3*c^2*C - 6*a^2*A*b^2*c*d + 2*A*b^4*c
*d + 2*a^3*b*B*c*d - 6*a*b^3*B*c*d + 6*a^2*b^2*c*C*d - 2*b^4*c*C*d - a^3*A
*b*d^2 + 3*a*A*b^3*d^2 - 3*a^2*b^2*B*d^2 + b^4*B*d^2 + a^3*b*C*d^2 - 3*a*b
^3*C*d^2))*Log[Sqrt[-b^2] - b*Tan[e + f*x]]/(b*(a^2 + b^2)*(c^2 + d^2)))
- (2*b^2*(c^2 + d^2)*(6*a^5*b*B*d^2 - 3*a^6*C*d^2 - a^4*b^2*d*(4*B*c + (10
*A - C)*d) - b^6*(c*(c*C - 2*B*d) - A*(c^2 - 3*d^2)) + a*b^5*(2*c*(A - C)*
d - B*(3*c^2 - d^2)) + 3*a^2*b^4*(c*(c*C + 2*B*d) - A*(c^2 + 3*d^2)) + a^3
*b^3*(10*c*(A - C)*d + B*(c^2 + 3*d^2))*Log[a + b*Tan[e + f*x]]/((a^2 +
b^2)*(b*c - a*d)) + ((b*c - a*d)^3*(b^2*(-3*a^2*A*b*c^2 + A*b^3*c^2 + a^3*
B*c^2 - 3*a*b^2*B*c^2 + 3*a^2*b*c^2*C - b^3*c^2*C - 2*a^3*A*c*d + 6*a*A*b^
2*c*d - 6*a^2*b*B*c*d + 2*b^3*B*c*d + 2*a^3*c*C*d - 6*a*b^2*c*C*d + 3*a...

```

Rubi [A] (verified)

Time = 8.81 (sec) , antiderivative size = 919, normalized size of antiderivative = 1.09, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.244$, Rules used = {3042, 4132, 3042, 4132, 3042, 4132, 27, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx$$

↓ 4132

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan^2(e+fx)-2aA(bc-ad)-(bB-aC)(2bc+ad)+2(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^2} dx$$

$$\frac{2(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$\frac{2f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))^2(c+d \tan(e+fx))}{}$$

↓ 3042

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan(e+fx)^2-2aA(bc-ad)-(bB-aC)(2bc+ad)+2(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^2} dx$$

$$\frac{2(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$\frac{2f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))^2(c+d \tan(e+fx))}{}$$

↓ 4132

$$\frac{-3a^4Cd+5a^3bBd-a^2b^2(7Ad+2Bc-Cd)+ab^3(4Ac+Bd-4cC)+b^4(2Bc-3Ad)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))} - \int \frac{2(Ba^2-2b(A-C)a-b^2B) \tan(e+fx)(bc-ad)^2-2d(-3Cda^4+}$$

$$\frac{Ab^2-a(bB-aC)}{2f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))^2(c+d \tan(e+fx))}$$

↓ 3042

$$\frac{-3a^4Cd+5a^3bBd-a^2b^2(7Ad+2Bc-Cd)+ab^3(4Ac+Bd-4cC)+b^4(2Bc-3Ad)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))} - \int \frac{2(Ba^2-2b(A-C)a-b^2B) \tan(e+fx)(bc-ad)^2-2d(-3Cda^4+}$$

$$\frac{Ab^2-a(bB-aC)}{2f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))^2(c+d \tan(e+fx))}$$

↓ 4132

$$\frac{Ab^2-a(bB-aC)}{2(a^2+b^2)(bc-ad)f(a+b \tan(e+fx))^2(c+d \tan(e+fx))} -$$

$$\int \frac{2(d^3(Ac-Cc+Bd)a^5-3Abd^2(c^2+d^2)a^4+b^2d(3Ac^3-3Cc^3+5$$

$$\frac{-3Cda^4+5bBda^3-b^2(2Bc+7Ad-Cd)a^2+b^3(4Ac-4Cc+Bd)a+b^4(2Bc-3Ad)}{(a^2+b^2)(bc-ad)f(a+b \tan(e+fx))(c+d \tan(e+fx))} -$$

↓ 27

$$\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

$$\frac{-3Cda^4 + 5bBda^3 - b^2(2Bc + 7Ad - Cd)a^2 + b^3(4Ac - 4Cc + Bd)a + b^4(2Bc - 3Ad)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

$$\frac{-3Cda^4 + 5bBda^3 - b^2(2Bc + 7Ad - Cd)a^2 + b^3(4Ac - 4Cc + Bd)a + b^4(2Bc - 3Ad)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

↓ 4134

$$\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

$$\frac{-3Cda^4 + 5bBda^3 - b^2(2Bc + 7Ad - Cd)a^2 + b^3(4Ac - 4Cc + Bd)a + b^4(2Bc - 3Ad)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

$$\frac{-3Cda^4 + 5bBda^3 - b^2(2Bc + 7Ad - Cd)a^2 + b^3(4Ac - 4Cc + Bd)a + b^4(2Bc - 3Ad)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

↓ 4013

$$\frac{Ab^2 - a(bB - aC)}{2(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^2(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

$$\frac{-3Cda^4 + 5bBda^3 - b^2(2Bc + 7Ad - Cd)a^2 + b^3(4Ac - 4Cc + Bd)a + b^4(2Bc - 3Ad)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))} - \frac{2d(-d(3Cc^2 - Bdc + (A+2C)d^2)a^4 + 3bBd(c^2 + d^2)a^3 - b^2(Bc^3 +$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2)/((a + b*\text{Tan}[e + f*x])^3*(c + d*\text{Tan}[e + f*x])^2), x]$

output
$$\begin{aligned} & -1/2*(A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*\text{Tan}[e + f*x])^2*(c + d*\text{Tan}[e + f*x])) - ((5*a^3*b*B*d - 3*a^4*C*d + b^4*(2*B*c - 3*A*d) + a*b^3*(4*A*c - 4*c*C + B*d) - a^2*b^2*(2*B*c + 7*A*d - C*d))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*\text{Tan}[e + f*x])*(c + d*\text{Tan}[e + f*x])) - ((-2*((b*c - a*d)^3*(a^3*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - 3*a*b^2*(c^2*C - 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + 3*a^2*b*(2*c*(A - C)*d - B*(c^2 - d^2)) - b^3*(2*c*(A - C)*d - B*(c^2 - d^2)))*x)/((a^2 + b^2)*(c^2 + d^2)) + (b*(c^2 + d^2)*(6*a^5*b*B*d^2 - 3*a^6*C*d^2 - a^4*b^2*d*(4*B*c + (10*A - C)*d) - b^6*(c*(c*C - 2*B*d) - A*(c^2 - 3*d^2)) + a*b^5*(2*c*(A - C)*d - B*(3*c^2 - d^2)) + 3*a^2*b^4*(c*(c*C + 2*B*d) - A*(c^2 + 3*d^2)) + a^3*b^3*(10*c*(A - C)*d + B*(c^2 + 3*d^2)))*Log[a*Cos[e + f*x] + b*Sin[e + f*x]]/((a^2 + b^2)*(b*c - a*d)*f) + ((a^2 + b^2)^2*d^2*(b*(3*c^4*C - 4*B*c^3*d + c^2*(5*A + C)*d^2 - 2*B*c*d^3 + 3*A*d^4) - a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Log[c*Cos[e + f*x] + d*Sin[e + f*x]]/((b*c - a*d)*(c^2 + d^2)*f)))/((b*c - a*d)*(c^2 + d^2)) - (2*d*(3*a^3*b*B*d*(c^2 + d^2) + a*b^3*(2*A*c - 2*c*C + B*d)*(c^2 + d^2) - a^4*d*(3*c^2*C - B*c*d + (A + 2*C)*d^2) - a^2*b^2*(B*c^3 + 4*A*c^2*d + 2*c^2*C*d - B*c*d^2 + 6*A*d^3) - b^4*(d*(2*A*c^2 + c^2*C + 3*A*d^2) - B*(c^3 + 2*c*d^2)))/((b*c - a*d)*(c^2 + d^2)*f*(c + d*\text{Tan}[e + f*x]))/((a^2 + b^2)*(b*c - a*d))/((2*(a^2 + b^2)*(b*c - a*d)) \end{aligned}$$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x_)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4013 $\text{Int}[((c_) + (d_)*\text{tan}[(e_) + (f_)*(x_)])/((a_) + (b_)*\text{tan}[(e_) + (f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 4.98 (sec) , antiderivative size = 951, normalized size of antiderivative = 1.13

method	result
derivativedivides	$\frac{(-2A a^3 cd - 3A a^2 b c^2 + 3A a^2 b d^2 + 6A a b^2 cd + A b^3 c^2 - b^3 A d^2 + B a^3 c^2 - B a^3 d^2 - 6B a^2 bcd - 3B a b^2 c^2 + 3B a b^2 d^2 + 2B b^3 cd + 2C a^3)}{2}$
default	$\frac{(-2A a^3 cd - 3A a^2 b c^2 + 3A a^2 b d^2 + 6A a b^2 cd + A b^3 c^2 - b^3 A d^2 + B a^3 c^2 - B a^3 d^2 - 6B a^2 bcd - 3B a b^2 c^2 + 3B a b^2 d^2 + 2B b^3 cd + 2C a^3)}{2}$
norman	Expression too large to display
risch	Expression too large to display
parallelrisc	Expression too large to display

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e))^2, x,method=_RETURNVERBOSE)`

output
$$\frac{1}{f} \left(\frac{1}{(a^2+b^2)^3} \frac{1}{(c^2+d^2)^2} \left(\frac{1}{2} (-2Aa^3cd-3Aa^2b^2c^2+3Aa^2b^2d^2+6Aa^2b^2cd+Aa^3c^2-Aa^3d^2+B^3a^3c^2-B^3a^3d^2-6B^2a^2b^2cd-3B^2a^2b^2c^2+3B^2a^2b^2d^2+2B^2b^3cd+2C^3a^3cd+3C^3a^2b^2c^2-3C^3a^2b^2d^2-6C^3a^2b^2cd-C^3b^3c^2+C^3b^3d^2) \ln(1+\tan(fx+e)^2) + (Aa^3c^2-Aa^3d^2-6Aa^2b^2cd-3Aa^2b^2c^2+3Aa^2b^2d^2+2Aa^2b^3cd+2B^3a^3cd+3B^3a^2b^2c^2-3B^3a^2b^2d^2-6B^2a^2b^2cd-B^2b^3c^2+B^2b^3d^2-C^3a^3c^2+C^3a^3d^2+6C^3a^2b^2cd+3C^3a^2b^2c^2-3C^3a^2b^2d^2-2C^3b^3cd) \arctan(\tan(fx+e)) \right) - b(4Aa^2b^2d-2Aa^2b^3c+2Aa^2b^4d-3B^3a^3b^2d+B^3a^2b^2c-B^3a^2b^3d-B^3b^4c+2C^3a^4d+2C^3a^2b^3c) / (a^2+b^2)^2 / (a+b\tan(fx+e))^3 + b(10Aa^4b^2d^2-10Aa^3b^3cd+3Aa^2b^4c^2+9Aa^2b^4d^2-2Aa^2b^5cd-Aa^2b^6c^2+3Aa^2b^6d^2-6B^3a^5b^2d+4B^3a^4b^2cd-B^3a^3b^3c^2-3B^3a^3b^3d^2-6B^3a^2b^4cd+3B^3a^2b^5c^2-B^3a^2b^5d^2-2B^3b^6cd+3C^3a^6d^2-C^3a^4b^2d^2+10C^3a^3b^3cd-3C^3a^2b^4c^2+2C^3a^2b^5cd+C^3b^6c^2) / (a^2+b^2)^3 \ln(a+b\tan(fx+e)) - \frac{1}{2} (A^2b^2-B^2a^2+B^2C^2) \frac{b}{(a^2+b^2)^2} \frac{1}{(a+b\tan(fx+e))^2} \frac{1}{(c^2+d^2)^2} (2Aa^3cd-5A^2b^2c^2d^2-3A^2b^2d^4-B^2a^3c^2d^2+B^2a^3d^4+4B^2b^3c^3d+2B^2b^3cd^3-2C^3a^3cd^3-3C^3b^3c^4-C^3b^3c^2d^2) / (c^2+d^2)^2 / (a^2+b^2)^4 \ln(c+d\tan(fx+e)) - (A^2d^2-B^2cd+C^2c^2) \frac{d^2}{(c^2+d^2)^2} \frac{1}{(a^2+b^2)^3} \frac{1}{(c+d\tan(fx+e))^3} \right)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9594 vs. $2(835) = 1670$.

Time = 9.70 (sec) , antiderivative size = 9594, normalized size of antiderivative = 11.41

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e))^2,x, algorithm="fricas")`

output Too large to include

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx$$

= Exception raised: NotImplementedError

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**3/(c+d*tan(f*x+e))**2,x)`

output `Exception raised: NotImplementedError >> no valid subset found`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2519 vs. 2(835) = 1670.

Time = 0.24 (sec) , antiderivative size = 2519, normalized size of antiderivative = 3.00

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e))^2,x, algorithm="maxima")`

output

```

1/2*(2*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c^2 + 2*(B*a^3
- 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c*d - ((A - C)*a^3 + 3*B*a^2
*b - 3*(A - C)*a*b^2 - B*b^3)*d^2)*(f*x + e)/((a^6 + 3*a^4*b^2 + 3*a^2*b^4
+ b^6)*c^4 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d^2 + (a^6 + 3*a^4
*b^2 + 3*a^2*b^4 + b^6)*d^4) - 2*((B*a^3*b^4 - 3*(A - C)*a^2*b^5 - 3*B*a*b
^6 + (A - C)*b^7)*c^2 - 2*(2*B*a^4*b^3 - 5*(A - C)*a^3*b^4 - 3*B*a^2*b^5 -
(A - C)*a*b^6 - B*b^7)*c*d - (3*C*a^6*b - 6*B*a^5*b^2 + (10*A - C)*a^4*b^
3 - 3*B*a^3*b^4 + 9*A*a^2*b^5 - B*a*b^6 + 3*A*b^7)*d^2)*log(b*tan(f*x + e)
+ a)/((a^6*b^4 + 3*a^4*b^6 + 3*a^2*b^8 + b^10)*c^4 - 4*(a^7*b^3 + 3*a^5*b
^5 + 3*a^3*b^7 + a*b^9)*c^3*d + 6*(a^8*b^2 + 3*a^6*b^4 + 3*a^4*b^6 + a^2*b
^8)*c^2*d^2 - 4*(a^9*b + 3*a^7*b^3 + 3*a^5*b^5 + a^3*b^7)*c*d^3 + (a^10 +
3*a^8*b^2 + 3*a^6*b^4 + a^4*b^6)*d^4) - 2*(3*C*b*c^4*d^2 - 4*B*b*c^3*d^3 +
(B*a + (5*A + C)*b)*c^2*d^4 - 2*((A - C)*a + B*b)*c*d^5 - (B*a - 3*A*b)*d
^6)*log(d*tan(f*x + e) + c)/(b^4*c^8 - 4*a*b^3*c^7*d - 4*a^3*b*c^6*d^2 + a^
4*d^8 + 2*(3*a^2*b^2 + b^4)*c^6*d^2 - 4*(a^3*b + 2*a*b^3)*c^5*d^3 + (a^4 +
12*a^2*b^2 + b^4)*c^4*d^4 - 4*(2*a^3*b + a*b^3)*c^3*d^5 + 2*(a^4 + 3*a^2*b
^2)*c^2*d^6) + ((B*a^3 - 3*(A - C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*c^2 -
2*((A - C)*a^3 + 3*B*a^2*b - 3*(A - C)*a*b^2 - B*b^3)*c*d - (B*a^3 - 3*(A
- C)*a^2*b - 3*B*a*b^2 + (A - C)*b^3)*d^2)*log(tan(f*x + e)^2 + 1)/((a^6 +
3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^4 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4549 vs. $2(835) = 1670$.

Time = 1.30 (sec) , antiderivative size = 4549, normalized size of antiderivative = 5.41

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+
e))^2,x, algorithm="giac")

```

output

```
(A*a^3*c^2 - C*a^3*c^2 + 3*B*a^2*b*c^2 - 3*A*a*b^2*c^2 + 3*C*a*b^2*c^2 - B
*b^3*c^2 + 2*B*a^3*c*d - 6*A*a^2*b*c*d + 6*C*a^2*b*c*d - 6*B*a*b^2*c*d + 2
*A*b^3*c*d - 2*C*b^3*c*d - A*a^3*d^2 + C*a^3*d^2 - 3*B*a^2*b*d^2 + 3*A*a*b
^2*d^2 - 3*C*a*b^2*d^2 + B*b^3*d^2)*(f*x + e)/(a^6*c^4*f + 3*a^4*b^2*c^4*f
+ 3*a^2*b^4*c^4*f + b^6*c^4*f + 2*a^6*c^2*d^2*f + 6*a^4*b^2*c^2*d^2*f + 6
*a^2*b^4*c^2*d^2*f + 2*b^6*c^2*d^2*f + a^6*d^4*f + 3*a^4*b^2*d^4*f + 3*a^2
*b^4*d^4*f + b^6*d^4*f) + 1/2*(B*a^3*c^2 - 3*A*a^2*b*c^2 + 3*C*a^2*b*c^2 -
3*B*a*b^2*c^2 + A*b^3*c^2 - C*b^3*c^2 - 2*A*a^3*c*d + 2*C*a^3*c*d - 6*B*a
^2*b*c*d + 6*A*a*b^2*c*d - 6*C*a*b^2*c*d + 2*B*b^3*c*d - B*a^3*d^2 + 3*A*a
^2*b*d^2 - 3*C*a^2*b*d^2 + 3*B*a*b^2*d^2 - A*b^3*d^2 + C*b^3*d^2)*log(tan(
f*x + e)^2 + 1)/(a^6*c^4*f + 3*a^4*b^2*c^4*f + 3*a^2*b^4*c^4*f + b^6*c^4*f
+ 2*a^6*c^2*d^2*f + 6*a^4*b^2*c^2*d^2*f + 6*a^2*b^4*c^2*d^2*f + 2*b^6*c^2
*d^2*f + a^6*d^4*f + 3*a^4*b^2*d^4*f + 3*a^2*b^4*d^4*f + b^6*d^4*f) - (B*a
^3*b^5*c^2 - 3*A*a^2*b^6*c^2 + 3*C*a^2*b^6*c^2 - 3*B*a*b^7*c^2 + A*b^8*c^2
- C*b^8*c^2 - 4*B*a^4*b^4*c*d + 10*A*a^3*b^5*c*d - 10*C*a^3*b^5*c*d + 6*B
*a^2*b^6*c*d + 2*A*a*b^7*c*d - 2*C*a*b^7*c*d + 2*B*b^8*c*d - 3*C*a^6*b^2*d
^2 + 6*B*a^5*b^3*d^2 - 10*A*a^4*b^4*d^2 + C*a^4*b^4*d^2 + 3*B*a^3*b^5*d^2
- 9*A*a^2*b^6*d^2 + B*a*b^7*d^2 - 3*A*b^8*d^2)*log(abs(b*tan(f*x + e) + a)
)/(a^6*b^5*c^4*f + 3*a^4*b^7*c^4*f + 3*a^2*b^9*c^4*f + b^11*c^4*f - 4*a^7*
b^4*c^3*d*f - 12*a^5*b^6*c^3*d*f - 12*a^3*b^8*c^3*d*f - 4*a*b^10*c^3*d*...
```

Mupad [B] (verification not implemented)

Time = 40.84 (sec) , antiderivative size = 128667, normalized size of antiderivative = 152.99

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^3*(c + d
*tan(e + f*x))^2),x)
```

output

```
(symsum(log((24*A^3*a^3*b^7*d^9 + 27*A^3*a^5*b^5*d^9 + B^3*a^2*b^8*d^9 + 4
*B^3*a^4*b^6*d^9 + 7*B^3*a^6*b^4*d^9 + 3*A^3*b^10*c^3*d^6 - A^3*b^10*c^5*d
^4 + 4*B^3*b^10*c^2*d^7 + 6*B^3*b^10*c^4*d^5 + C^3*b^10*c^5*d^4 + 9*A^2*B*
b^10*d^9 + 9*A^3*a*b^9*d^9 + 16*A^3*a^2*b^8*c^3*d^6 + 3*A^3*a^2*b^8*c^5*d^
4 + 26*A^3*a^3*b^7*c^2*d^7 - 6*A^3*a^3*b^7*c^4*d^5 - 11*A^3*a^4*b^6*c^3*d^
6 + 31*A^3*a^5*b^5*c^2*d^7 + 5*B^3*a^2*b^8*c^2*d^7 + 6*B^3*a^2*b^8*c^4*d^5
+ 28*B^3*a^3*b^7*c^3*d^6 + 7*B^3*a^3*b^7*c^5*d^4 - 14*B^3*a^4*b^6*c^2*d^7
- 20*B^3*a^4*b^6*c^4*d^5 + 19*B^3*a^5*b^5*c^3*d^6 + 9*B^3*a^6*b^4*c^2*d^7
- 7*C^3*a^2*b^8*c^3*d^6 - 3*C^3*a^2*b^8*c^5*d^4 + C^3*a^3*b^7*c^2*d^7 + 1
5*C^3*a^3*b^7*c^4*d^5 + 6*C^3*a^3*b^7*c^6*d^3 - 28*C^3*a^4*b^6*c^3*d^6 - 2
4*C^3*a^4*b^6*c^5*d^4 - 4*C^3*a^5*b^5*c^2*d^7 + 3*C^3*a^6*b^4*c^3*d^6 - 9*
C^3*a^7*b^3*c^2*d^7 - 9*C^3*a^7*b^3*c^4*d^5 - 6*A*B^2*a*b^9*d^9 - 9*A^2*C*
a*b^9*d^9 - 12*A*B^2*b^10*c*d^8 + 4*B^3*a*b^9*c*d^8 - 20*A*B^2*a^3*b^7*d^9
- 28*A*B^2*a^5*b^5*d^9 + 6*A*B^2*a^7*b^3*d^9 + 21*A^2*B*a^2*b^8*d^9 + 13*
A^2*B*a^4*b^6*d^9 - 27*A^2*B*a^6*b^4*d^9 - 3*A*C^2*a^3*b^7*d^9 - 9*A*C^2*a
^7*b^3*d^9 - 21*A^2*C*a^3*b^7*d^9 - 27*A^2*C*a^5*b^5*d^9 + 9*A^2*C*a^7*b^3
*d^9 - 17*A*B^2*b^10*c^3*d^6 + 3*A*B^2*b^10*c^5*d^4 + B*C^2*a^4*b^6*d^9 +
3*B*C^2*a^8*b^2*d^9 + 12*A^2*B*b^10*c^2*d^7 - 7*A^2*B*b^10*c^4*d^5 - B^2*C
*a^3*b^7*d^9 - 2*B^2*C*a^5*b^5*d^9 - 9*B^2*C*a^7*b^3*d^9 + 3*A*C^2*b^10*c^
3*d^6 - 3*A*C^2*b^10*c^5*d^4 - 6*A^2*C*b^10*c^3*d^6 + 3*A^2*C*b^10*c^5*...
```

Reduce [B] (verification not implemented)

Time = 14.42 (sec) , antiderivative size = 27735, normalized size of antiderivative = 32.98

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3/(c+d*tan(f*x+e))^2,
x)
```

output

```
( - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**9*b**2*c*d**7 + 10*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**8*b**3*c**2*d**6 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**8*b**3*d**8 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**8*b**2*c**2*d**7 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**7*b**4*c**3*d**5 - 14*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**7*b**4*c*d**7 - 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**7*b**3*c**3*d**6 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**7*b**3*c*d**8 - 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**5*c**4*d**4 + 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**5*c**2*d**6 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**5*d**8 - 5*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**4*c**4*d**5 + 9*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**4*c**2*d**7 + 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**6*c**5*d**3 + 10*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**6*c**3*d**5 - 10*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**6*c*d**7 + 20*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**5*c**5*d**4 + 16*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**5*c**3*d**6 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**5*c*d**8 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**7*c**6*d**2 - 20*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**7*c**4*d**4 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**7*c**2*d**6 - 10*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**6*c**6*...
```

3.84
$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$$

Optimal result	936
Mathematica [C] (verified)	937
Rubi [A] (verified)	938
Maple [A] (verified)	943
Fricas [B] (verification not implemented)	944
Sympy [F(-2)]	945
Maxima [A] (verification not implemented)	946
Giac [B] (verification not implemented)	947
Mupad [B] (verification not implemented)	948
Reduce [B] (verification not implemented)	949

Optimal result

Integrand size = 45, antiderivative size = 804

$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx =$$

$$\frac{(3ab^2(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) + a^3(c^3C - 3Bc^2d - 3cCd^2 + Bd^3 - A(c^3 - 3ca^2d + c^2d^2)) - (3a^2b(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) - b^3(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) - (bc - ad)(b^2(3c^6C - Bc^5d + 9c^4Cd^2 - 3Bc^3d^3 - c^2(A - 10C)d^4 - 6Bcd^5 + 3Ad^6) + a^2d^3((A - C)d^4 - B(c^2 - d^2))) \tan(e+fx))}{d^3(c^2 + d^2)^2 f}$$

$$+ \frac{b^2(b(3c^4C - Bc^3d + 6c^2Cd^2 - 3Bcd^3) + (2A + C)d^4) + ad^2(2c(A - C)d - B(c^2 - d^2)) \tan(e+fx)}{d^3(c^2 + d^2)^2 f}$$

$$- \frac{(c^2C - Bcd + Ad^2)(a+b \tan(e+fx))^3}{2d(c^2 + d^2) f(c+d \tan(e+fx))^2}$$

$$- \frac{(b(3c^4C - Bc^3d - c^2(A - 7C)d^2 - 5Bcd^3 + 3Ad^4) + 2ad^2(2c(A - C)d - B(c^2 - d^2))) (a+b \tan(e+fx))}{2d^2(c^2 + d^2)^2 f(c+d \tan(e+fx))}$$

output

```

-(3*a*b^2*(A*c^3-3*A*c*d^2+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)+a^3*(c^3*C-3*B
*c^2*d-3*C*c*d^2+B*d^3-A*(c^3-3*c*d^2))-3*a^2*b*((A-C)*d*(3*c^2-d^2)-B*(c^
3-3*c*d^2))+b^3*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2)))*x/(c^2+d^2)^3-(3*a^
2*b*(A*c^3-3*A*c*d^2+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)-b^3*(A*c^3-3*A*c*d^2
+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)-a^3*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2)
)+3*a*b^2*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2)))*ln(cos(f*x+e))/(c^2+d^2)^
3/f-(-a*d+b*c)*(b^2*(3*c^6*C-B*c^5*d+9*c^4*C*d^2-3*B*c^3*d^3-c^2*(A-10*C)*
d^4-6*B*c*d^5+3*A*d^6)+a^2*d^3*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2))+a*b*d
^2*(8*c*(A-C)*d^3-B*(c^4+6*c^2*d^2-3*d^4)))*ln(c+d*tan(f*x+e))/d^4/(c^2+d^
2)^3/f+b^2*(b*(3*c^4*C-B*c^3*d+6*C*c^2*d^2-3*B*c*d^3+(2*A+C)*d^4)+a*d^2*(2
*c*(A-C)*d-B*(c^2-d^2)))*tan(f*x+e)/d^3/(c^2+d^2)^2/f-1/2*(A*d^2-B*c*d+C*c
^2)*(a+b*tan(f*x+e))^3/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^2-1/2*(b*(3*c^4*C-B
c^3*d-c^2*(A-7*C)*d^2-5*B*c*d^3+3*A*d^4)+2*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)
))*(a+b*tan(f*x+e))^2/d^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.37 (sec) , antiderivative size = 454, normalized size of antiderivative = 0.56

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

$$= \frac{(a+ib)^3(A+iB-C) \log(i-\tan(e+fx))}{(-ic+d)^3} + \frac{(a-ib)^3(A-iB-C) \log(i+\tan(e+fx))}{(ic+d)^3} + \frac{2(-bc+ad)(b^2(3c^6C-Bc^5d+9c^4Cd^2-3Bc^3d^3-c^2(A-10C)d^4-6Bcd^5+3Ad^6)+a^2d^3((A-C)d(3c^2-d^2)-B(c^3-3cd^2))+ab d^2(8c(A-C)d^3-B(c^4+6c^2d^2-3d^4)))*\ln(c+d \tan(fx+e))}{d^4(c^2+d^2)^3} + \frac{b^2(b(3c^4C-Bc^3d+6C c^2d^2-3Bcd^3+(2A+C)d^4)+a d^2(2c(A-C)d-B(c^2-d^2)))*\tan(fx+e)}{d^3(c^2+d^2)^2} - \frac{1}{2} \frac{(A d^2-B c d+C c^2)(a+b \tan(fx+e))^3}{d(c^2+d^2)} - \frac{1}{2} \frac{(b(3c^4C-Bc^3d-c^2(A-7C)d^2-5Bcd^3+3Ad^4)+2ad^2(2c(A-C)d-B(c^2-d^2)))*(a+b \tan(fx+e))^2}{d^2(c^2+d^2)^2} - \frac{1}{2} \frac{(a+b \tan(fx+e))^2}{d(c^2+d^2)^2} - \frac{1}{2} \frac{(a+b \tan(fx+e))^2}{d(c+d \tan(fx+e))}$$

input

```

Integrate[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^3,x]

```

output

```

(((a + I*b)^3*(A + I*B - C)*Log[I - Tan[e + f*x]])/((-I)*c + d)^3 + ((a -
I*b)^3*(A - I*B - C)*Log[I + Tan[e + f*x]])/(I*c + d)^3 + (2*(-(b*c) + a*d
)*(b^2*(3*c^6*C - B*c^5*d + 9*c^4*C*d^2 - 3*B*c^3*d^3 - c^2*(A - 10*C)*d^4
- 6*B*c*d^5 + 3*A*d^6) + a^2*d^3*(-((A - C)*d*(-3*c^2 + d^2)) - B*(c^3 -
3*c*d^2)) - a*b*d^2*(8*c*(-A + C)*d^3 + B*(c^4 + 6*c^2*d^2 - 3*d^4)))*Log[
c + d*Tan[e + f*x]]/(d^4*(c^2 + d^2)^3) + ((b*c - a*d)^3*(3*c^2*C - B*c*d
+ (A + 2*C)*d^2))/(d^4*(c^2 + d^2)*(c + d*Tan[e + f*x])^2) + (2*C*(a + b*
Tan[e + f*x])^3)/(d*(c + d*Tan[e + f*x])^2) - (2*(b*c - a*d)^2*(b*(6*c^4*C
- 2*B*c^3*d + c^2*(A + 11*C)*d^2 - 4*B*c*d^3 + 3*(A + C)*d^4) + a*d^2*(2*
c*(A - C)*d + B*(-c^2 + d^2)))/(d^4*(c^2 + d^2)^2*(c + d*Tan[e + f*x]))/(
2*f)

```

Rubi [A] (verified)

Time = 6.74 (sec) , antiderivative size = 837, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4128, 3042, 4128, 3042, 4120, 27, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{(a + b \tan(e + fx))^2 (b(3Cc^2 - Bdc + (A + 2C)d^2) \tan^2(e + fx) + 2d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(2ac + 3bd) + (3bc - 2ad)(cC - Bd))}{(c + d \tan(e + fx))^2} \frac{2d(c^2 + d^2)}{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^2 (b(3Cc^2-Bdc+(A+2C)d^2) \tan(e+fx)^2+2d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(2ac+3bd)+(3bc-2ad)(cC-Bd))}{(c+d \tan(e+fx))^2} \\ \frac{2d(c^2+d^2)}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3} \\ \frac{2df(c^2+d^2)(c+d \tan(e+fx))^2}{2df(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 4128

$$\int \frac{(a+b \tan(e+fx)) (2((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+2bd)(Ad(2ac+3bd)+(3bc-2ad)(cC-Bd))}{(c+d \tan(e+fx))^2}$$

$$\frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}{2df(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx)) (2((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+2bd)(Ad(2ac+3bd)+(3bc-2ad)(cC-Bd))}{(c+d \tan(e+fx))^2}$$

$$\frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}{2df(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 4120

$$2b^2 \tan(e+fx) \frac{(ad^2(2cd(A-C)-B(c^2-d^2))+b(d^4(2A+C)-Bc^3d-3Bcd^3+3e^4C+6c^2Cd^2))}{df} - \int \frac{2(-c(3Cc^4-Bdc^3+6Cd^2c^2-3Bd^3c+(2A+C)d^4)b^3-(3bcC-3adC-bBd)(c^2+d^2)^2 \tan^2(e+fx)b^2+3ad(Cc^4-(A-3C)d^2c^2-2Bd^3c+Ad^4)b^2+3a^2d^3(2c(A-C)-Bd))}{(c+d \tan(e+fx))^2}$$

$$\frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}{2df(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 27

$$2 \int \frac{-c(3Cc^4-Bdc^3+6Cd^2c^2-3Bd^3c+(2A+C)d^4)b^3-(3bcC-3adC-bBd)(c^2+d^2)^2 \tan^2(e+fx)b^2+3ad(Cc^4-(A-3C)d^2c^2-2Bd^3c+Ad^4)b^2+3a^2d^3(2c(A-C)-Bd)}{(c+d \tan(e+fx))^2}$$

$$\frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^3}{2df(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 3042

$$2 \int \frac{-c(3Ce^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4)b^3 - (3bcC - 3adC - bBd)(c^2 + d^2)^2 \tan(e+fx)^2 b^2 + 3ad(Cc^4 - (A-3C)d^2c^2 - 2Bd^3c + Ad^4)b^2 + 3a^2d^3(2c(A-C) - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 4109

$$\frac{2(a(2c(A-C)d - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} + 2 \left(- \frac{((Cc^3 - 3Bdc^2 - 3Cd^2c + Bd^3 - A(c^3 - 3cd^2))a^3 - 3b((A-C)d^2 - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} \right)$$

$$\frac{(Cc^2 - Bdc + Ad^2)(a + b \tan(e + fx))^3}{2d(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 3042

$$\frac{2(a(2c(A-C)d - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} + 2 \left(- \frac{((Cc^3 - 3Bdc^2 - 3Cd^2c + Bd^3 - A(c^3 - 3cd^2))a^3 - 3b((A-C)d^2 - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} \right)$$

$$\frac{(Cc^2 - Bdc + Ad^2)(a + b \tan(e + fx))^3}{2d(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 3956

$$\frac{2(a(2c(A-C)d - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} + 2 \left(- \frac{((Cc^3 - 3Bdc^2 - 3Cd^2c + Bd^3 - A(c^3 - 3cd^2))a^3 - 3b((A-C)d^2 - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} \right)$$

$$\frac{(Cc^2 - Bdc + Ad^2)(a + b \tan(e + fx))^3}{2d(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 4100

$$\frac{2(a(2c(A-C)d - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} + 2 \left(- \frac{((Cc^3 - 3Bdc^2 - 3Cd^2c + Bd^3 - A(c^3 - 3cd^2))a^3 - 3b((A-C)d^2 - B(c^2 - d^2))d^2 + b(3Cc^4 - Bdc^3 + 6Cd^2c^2 - 3Bd^3c + (2A+C)d^4) \tan(e+fx)b^2}{df} \right)$$

$$\frac{(Cc^2 - Bdc + Ad^2)(a + b \tan(e + fx))^3}{2d(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 16

$$\frac{2(a(2c(A-C)d-B(c^2-d^2))d^2+b(3Cc^4-Bdc^3+6Cd^2e^2-3Bd^3c+(2A+C)d^4))\tan(e+fx)b^2}{df} + \frac{2\left(\frac{((Cc^3-3Bdc^2-3Cd^2c+Bd^3-A(c^3-3cd^2))a^3-3b(A-C))}{(C^2-Bdc+Ad^2)}\right)}{2d(c^2+d^2)f(c+d\tan(e+fx))^2}$$

$$\frac{(C^2 - Bdc + Ad^2)(a + b\tan(e + fx))^3}{2d(c^2 + d^2)f(c + d\tan(e + fx))^2}$$

input

```
Int[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^3,x]
```

output

```
-1/2*((c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^3)/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^2) + (-(((b*(3*c^4*C - B*c^3*d - c^2*(A - 7*C)*d^2 - 5*B*c*d^3 + 3*A*d^4) + 2*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2))))*(a + b*Tan[e + f*x])^2)/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])) + ((2*(-((d^3*(3*a*b^2*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) + a^3*(c^3*C - 3*B*c^2*d - 3*c*C*d^2 + B*d^3 - A*(c^3 - 3*c*d^2)) - 3*a^2*b*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2))) * x)/(c^2 + d^2)) - (d^3*(3*a^2*b*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) - b^3*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) - a^3*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)) + 3*a*b^2*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)))*Log[Cos[e + f*x]])/((c^2 + d^2)*f) - ((b*c - a*d)*(b^2*(3*c^6*C - B*c^5*d + 9*c^4*C*d^2 - 3*B*c^3*d^3 - c^2*(A - 10*C)*d^4 - 6*B*c*d^5 + 3*A*d^6) + a^2*d^3*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)) + a*b*d^2*(8*c*(A - C)*d^3 - B*(c^4 + 6*c^2*d^2 - 3*d^4))*Log[c + d*Tan[e + f*x]])/(d*(c^2 + d^2)*f))/d + (2*b^2*(b*(3*c^4*C - B*c^3*d + 6*c^2*C*d^2 - 3*B*c*d^3 + (2*A + C)*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2))*Tan[e + f*x])/(d*f))/(d*(c^2 + d^2)))/(2*d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4100 $\text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x_)]^{(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \ \&\& \ \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[(A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)/((a_)+(b_)*\tan[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A*b - a*B - b*C, 0]$
- rule 4120 $\text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x_)]*(c_)+(d_)*\tan[(e_)+(f_)*(x_)]^{(n_)*((A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[b*C*\text{Tan}[e + f*x]*((c + d*\text{Tan}[e + f*x])^{(n + 1)/(d*f*(n + 2))}), x] - \text{Simp}[1/(d*(n + 2)) \text{ Int}[(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*\text{Tan}[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ !\text{LtQ}[n, -1]$

rule 4128

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]

```

Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 1271, normalized size of antiderivative = 1.58

method	result	size
derivativdivides	Expression too large to display	1271
default	Expression too large to display	1271
norman	Expression too large to display	2076
parallelrisc	Expression too large to display	6687
risc	Expression too large to display	6825

input

```

int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,
x,method=_RETURNVERBOSE)

```

output

```

1/f*(tan(f*x+e)*C*b^3/d^3+1/(c^2+d^2)^3*(1/2*(-3*A*a^3*c^2*d+A*a^3*d^3+3*A
*a^2*b*c^3-9*A*a^2*b*c*d^2+9*A*a*b^2*c^2*d-3*A*a*b^2*d^3-A*b^3*c^3+3*A*b^3
*c*d^2+B*a^3*c^3-3*B*a^3*c*d^2+9*B*a^2*b*c^2*d-3*B*a^2*b*d^3-3*B*a*b^2*c^3
+9*B*a*b^2*c*d^2-3*B*b^3*c^2*d+B*b^3*d^3+3*C*a^3*c^2*d-C*a^3*d^3-3*C*a^2*b
*c^3+9*C*a^2*b*c*d^2-9*C*a*b^2*c^2*d+3*C*a*b^2*d^3+C*b^3*c^3-3*C*b^3*c*d^2
)*ln(1+tan(f*x+e)^2)+(A*a^3*c^3-3*A*a^3*c*d^2+9*A*a^2*b*c^2*d-3*A*a^2*b*d^
3-3*A*a*b^2*c^3+9*A*a*b^2*c*d^2-3*A*b^3*c^2*d+A*b^3*d^3+3*B*a^3*c^2*d-B*a^
3*d^3-3*B*a^2*b*c^3+9*B*a^2*b*c*d^2-9*B*a*b^2*c^2*d+3*B*a*b^2*d^3+B*b^3*c^
3-3*B*b^3*c*d^2-C*a^3*c^3+3*C*a^3*c*d^2-9*C*a^2*b*c^2*d+3*C*a^2*b*d^3+3*C*
a*b^2*c^3-9*C*a*b^2*c*d^2+3*C*b^3*c^2*d-C*b^3*d^3)*arctan(tan(f*x+e))-1/2
/d^4*(A*a^3*d^5-3*A*a^2*b*c*d^4+3*A*a*b^2*c^2*d^3-A*b^3*c^3*d^2-B*a^3*c*d^
4+3*B*a^2*b*c^2*d^3-3*B*a*b^2*c^3*d^2+B*b^3*c^4*d+C*a^3*c^2*d^3-3*C*a^2*b*
c^3*d^2+3*C*a*b^2*c^4*d-C*b^3*c^5)/(c^2+d^2)/(c+d*tan(f*x+e))^2-1/d^4*(2*A
*a^3*c*d^5-3*A*a^2*b*c^2*d^4+3*A*a^2*b*d^6-6*A*a*b^2*c*d^5+A*b^3*c^4*d^2+3
*A*b^3*c^2*d^4-B*a^3*c^2*d^4+B*a^3*d^6-6*B*a^2*b*c*d^5+3*B*a*b^2*c^4*d^2+9
*B*a*b^2*c^2*d^4-2*B*b^3*c^5*d-4*B*b^3*c^3*d^3-2*C*a^3*c*d^5+3*C*a^2*b*c^4
*d^2+9*C*a^2*b*c^2*d^4-6*C*a*b^2*c^5*d-12*C*a*b^2*c^3*d^3+3*C*b^3*c^6+5*C*
b^3*c^4*d^2)/(c^2+d^2)^2/(c+d*tan(f*x+e))+1/d^4*(3*A*a^3*c^2*d^5-A*a^3*d^7
-3*A*a^2*b*c^3*d^4+9*A*a^2*b*c*d^6-9*A*a*b^2*c^2*d^5+3*A*a*b^2*d^7+A*b^3*c
^3*d^4-3*A*b^3*c*d^6-B*a^3*c^3*d^4+3*B*a^3*c*d^6-9*B*a^2*b*c^2*d^5+3*B*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2490 vs. $2(797) = 1594$.

Time = 1.12 (sec) , antiderivative size = 2490, normalized size of antiderivative = 3.10

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```

integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^3,x, algorithm="fricas")

```

output

```

-1/2*(3*C*b^3*c^7*d^2 + A*a^3*d^9 - (3*C*a*b^2 + B*b^3)*c^6*d^3 - (3*C*a^2
*b + 3*B*a*b^2 + (A - 9*C)*b^3)*c^5*d^4 + (3*C*a^3 + 9*B*a^2*b + 3*(3*A -
7*C)*a*b^2 - 7*B*b^3)*c^4*d^5 - 5*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - A
*b^3)*c^3*d^6 + ((7*A - 3*C)*a^3 - 9*B*a^2*b - 9*A*a*b^2)*c^2*d^7 + (B*a^3
+ 3*A*a^2*b)*c*d^8 - 2*(C*b^3*c^6*d^3 + 3*C*b^3*c^4*d^5 + 3*C*b^3*c^2*d^7
+ C*b^3*d^9)*tan(f*x + e)^3 - 2*(((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b
^2 + B*b^3)*c^5*d^4 + 3*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3
)*c^4*d^5 - 3*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c^3*d^6
- (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c^2*d^7)*f*x - (9*C*
b^3*c^7*d^2 - A*a^3*d^9 - 3*(3*C*a*b^2 + B*b^3)*c^6*d^3 + (3*C*a^2*b + 3*B
*a*b^2 + (A + 23*C)*b^3)*c^5*d^4 + (C*a^3 + 3*B*a^2*b + 3*(A - 9*C)*a*b^2
- 9*B*b^3)*c^4*d^5 - (3*B*a^3 + 3*(3*A - 7*C)*a^2*b - 21*B*a*b^2 - (7*A +
12*C)*b^3)*c^3*d^6 + 5*((A - C)*a^3 - 3*B*a^2*b - 3*A*a*b^2)*c^2*d^7 + (3*
B*a^3 + 9*A*a^2*b + 4*C*b^3)*c*d^8 + 2*(((A - C)*a^3 - 3*B*a^2*b - 3*(A -
C)*a*b^2 + B*b^3)*c^3*d^6 + 3*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A -
C)*b^3)*c^2*d^7 - 3*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c
*d^8 - (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^9)*f*x)*tan(f*
x + e)^2 + (3*C*b^3*c^9 + 9*C*b^3*c^7*d^2 - (3*C*a*b^2 + B*b^3)*c^8*d - 3*
(3*C*a*b^2 + B*b^3)*c^6*d^3 + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A -
10*C)*b^3)*c^5*d^4 - 3*((A - C)*a^3 - 3*B*a^2*b - 3*(A - 2*C)*a*b^2 + 2...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

= Exception raised: AttributeError

input

```

integrate((a+b*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*
x+e))**3,x)

```

output

```

Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'

```

Maxima [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 1110, normalized size of antiderivative = 1.38

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="maxima")
```

output

```
1/2*(2*C*b^3*tan(f*x + e)/d^3 + 2*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c^3 + 3*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c^2*d - 3*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c*d^2 - (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*d^3)*(f*x + e)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) - 2*(3*C*b^3*c^7 + 9*C*b^3*c^5*d^2 - (3*C*a*b^2 + B*b^3)*c^6*d - 3*(3*C*a*b^2 + B*b^3)*c^4*d^3 + (B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - 10*C)*b^3)*c^3*d^4 - 3*((A - C)*a^3 - 3*B*a^2*b - 3*(A - 2*C)*a*b^2 + 2*B*b^3)*c^2*d^5 - 3*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - A*b^3)*c*d^6 + ((A - C)*a^3 - 3*B*a^2*b - 3*A*a*b^2)*d^7)*log(d*tan(f*x + e) + c)/(c^6*d^4 + 3*c^4*d^6 + 3*c^2*d^8 + d^10) + ((B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c^3 - 3*((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*c^2*d - 3*(B*a^3 + 3*(A - C)*a^2*b - 3*B*a*b^2 - (A - C)*b^3)*c*d^2 + ((A - C)*a^3 - 3*B*a^2*b - 3*(A - C)*a*b^2 + B*b^3)*d^3)*log(tan(f*x + e)^2 + 1)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) - (5*C*b^3*c^7 + A*a^3*d^7 - 3*(3*C*a*b^2 + B*b^3)*c^6*d + (3*C*a^2*b + 3*B*a*b^2 + (A + 9*C)*b^3)*c^5*d^2 + (C*a^3 + 3*B*a^2*b + 3*(A - 7*C)*a*b^2 - 7*B*b^3)*c^4*d^3 - (3*B*a^3 + 3*(3*A - 5*C)*a^2*b - 15*B*a*b^2 - 5*A*b^3)*c^3*d^4 + ((5*A - 3*C)*a^3 - 9*B*a^2*b - 9*A*a*b^2)*c^2*d^5 + (B*a^3 + 3*A*a^2*b)*c*d^6 + 2*(3*C*b^3*c^6*d - 2*(3*C*a*b^2 + B*b^3)*c^5*d^2 + (3*C*a^2*b + 3*B*a*b^2 + (A + 5*C)*b^3)*c^4*d^3 - 4*(3*C*a*b^2 + B*b^3)*c^3*d^4 - (B*a^3 + 3*(A ...
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1757 vs. $2(797) = 1594$.

Time = 0.92 (sec) , antiderivative size = 1757, normalized size of antiderivative = 2.19

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="giac")
```

output

```
(A*a^3*c^3 - C*a^3*c^3 - 3*B*a^2*b*c^3 - 3*A*a*b^2*c^3 + 3*C*a*b^2*c^3 + B*b^3*c^3 + 3*B*a^3*c^2*d + 9*A*a^2*b*c^2*d - 9*C*a^2*b*c^2*d - 9*B*a*b^2*c^2*d - 3*A*b^3*c^2*d + 3*C*b^3*c^2*d - 3*A*a^3*c*d^2 + 3*C*a^3*c*d^2 + 9*B*a^2*b*c*d^2 + 9*A*a*b^2*c*d^2 - 9*C*a*b^2*c*d^2 - 3*B*b^3*c*d^2 - B*a^3*d^3 - 3*A*a^2*b*d^3 + 3*C*a^2*b*d^3 + 3*B*a*b^2*d^3 + A*b^3*d^3 - C*b^3*d^3)*(f*x + e)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) + 1/2*(B*a^3*c^3 + 3*A*a^2*b*c^3 - 3*C*a^2*b*c^3 - 3*B*a*b^2*c^3 - A*b^3*c^3 + C*b^3*c^3 - 3*A*a^3*c^2*d + 3*C*a^3*c^2*d + 9*B*a^2*b*c^2*d + 9*A*a*b^2*c^2*d - 9*C*a*b^2*c^2*d - 3*B*b^3*c^2*d - 3*B*a^3*c*d^2 - 9*A*a^2*b*c*d^2 + 9*C*a^2*b*c*d^2 + 9*B*a*b^2*c*d^2 + 3*A*b^3*c*d^2 - 3*C*b^3*c*d^2 + A*a^3*d^3 - C*a^3*d^3 - 3*B*a^2*b*d^3 - 3*A*a*b^2*d^3 + 3*C*a*b^2*d^3 + B*b^3*d^3)*log(tan(f*x + e)^2 + 1)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) - (3*C*b^3*c^7 - 3*C*a*b^2*c^6*d - B*b^3*c^6*d + 9*C*b^3*c^5*d^2 - 9*C*a*b^2*c^4*d^3 - 3*B*b^3*c^4*d^3 + B*a^3*c^3*d^4 + 3*A*a^2*b*c^3*d^4 - 3*C*a^2*b*c^3*d^4 - 3*B*a*b^2*c^3*d^4 - A*b^3*c^3*d^4 + 10*C*b^3*c^3*d^4 - 3*A*a^3*c^2*d^5 + 3*C*a^3*c^2*d^5 + 9*B*a^2*b*c^2*d^5 + 9*A*a*b^2*c^2*d^5 - 18*C*a*b^2*c^2*d^5 - 6*B*b^3*c^2*d^5 - 3*B*a^3*c*d^6 - 9*A*a^2*b*c*d^6 + 9*C*a^2*b*c*d^6 + 9*B*a*b^2*c*d^6 + 3*A*b^3*c*d^6 + A*a^3*d^7 - C*a^3*d^7 - 3*B*a^2*b*d^7 - 3*A*a*b^2*d^7)*log(abs(d*tan(f*x + e) + c))/(c^6*d^4*f + 3*c^4*d^6*f + 3*c^2*d^8*f + d^10*f) + C*b^3*tan(f*x + e)/(d^3*f) - 1/2*(5*C*b^3*c^9 - 9*C*a...
```


Mupad [B] (verification not implemented)

Time = 15.05 (sec) , antiderivative size = 1172, normalized size of antiderivative = 1.46

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int(((a + b*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c +
d*tan(e + f*x))^3,x)
```

output

```
(log(tan(e + f*x) + 1i)*(A*a^3 + A*b^3*1i - B*a^3*1i + B*b^3 - C*a^3 - C*b
^3*1i - 3*A*a*b^2 - A*a^2*b*3i + B*a*b^2*3i - 3*B*a^2*b + 3*C*a*b^2 + C*a^
2*b*3i))/(2*f*(c*d^2*3i - 3*c^2*d - c^3*1i + d^3)) - ((A*a^3*d^7 + 5*C*b^3
*c^7 + B*a^3*c*d^6 - 3*B*b^3*c^6*d + 5*A*a^3*c^2*d^5 + 5*A*b^3*c^3*d^4 + A
*b^3*c^5*d^2 - 3*B*a^3*c^3*d^4 - 7*B*b^3*c^4*d^3 - 3*C*a^3*c^2*d^5 + C*a^3
*c^4*d^3 + 9*C*b^3*c^5*d^2 - 9*A*a*b^2*c^2*d^5 + 3*A*a*b^2*c^4*d^3 - 9*A*a
^2*b*c^3*d^4 + 15*B*a*b^2*c^3*d^4 + 3*B*a*b^2*c^5*d^2 - 9*B*a^2*b*c^2*d^5
+ 3*B*a^2*b*c^4*d^3 - 21*C*a*b^2*c^4*d^3 + 15*C*a^2*b*c^3*d^4 + 3*C*a^2*b*
c^5*d^2 + 3*A*a^2*b*c*d^6 - 9*C*a*b^2*c^6*d)/(2*d*(c^4 + d^4 + 2*c^2*d^2))
+ (tan(e + f*x)*(B*a^3*d^6 + 3*C*b^3*c^6 + 3*A*a^2*b*d^6 + 2*A*a^3*c*d^5
- 2*B*b^3*c^5*d - 2*C*a^3*c*d^5 + 3*A*b^3*c^2*d^4 + A*b^3*c^4*d^2 - B*a^3*
c^2*d^4 - 4*B*b^3*c^3*d^3 + 5*C*b^3*c^4*d^2 - 3*A*a^2*b*c^2*d^4 + 9*B*a*b^
2*c^2*d^4 + 3*B*a*b^2*c^4*d^2 - 12*C*a*b^2*c^3*d^3 + 9*C*a^2*b*c^2*d^4 + 3
*C*a^2*b*c^4*d^2 - 6*A*a*b^2*c*d^5 - 6*B*a^2*b*c*d^5 - 6*C*a*b^2*c^5*d))/(
c^4 + d^4 + 2*c^2*d^2))/(f*(c^2*d^3 + d^5*tan(e + f*x)^2 + 2*c*d^4*tan(e +
f*x))) + (log(c + d*tan(e + f*x))*(d^3*(3*B*b^3*c^4 + 9*C*a*b^2*c^4) - d^
6*(3*A*b^3*c - 3*B*a^3*c - 9*A*a^2*b*c + 9*B*a*b^2*c + 9*C*a^2*b*c) + d^5*
(3*A*a^3*c^2 + 6*B*b^3*c^2 - 3*C*a^3*c^2 - 9*A*a*b^2*c^2 - 9*B*a^2*b*c^2 +
18*C*a*b^2*c^2) + d^4*(A*b^3*c^3 - B*a^3*c^3 - 10*C*b^3*c^3 - 3*A*a^2*b*c
^3 + 3*B*a*b^2*c^3 + 3*C*a^2*b*c^3) + d^7*(C*a^3 - A*a^3 + 3*A*a*b^2 + ...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 5125, normalized size of antiderivative = 6.37

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,
x)
```

output

```
( - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*c**3*d**7 + log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a**4*c*d**9 + 4*log(tan(e + f*x)**2 + 1)*ta
n(e + f*x)**2*a**3*b*c**4*d**6 - 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*
**2*a**3*b*c**2*d**8 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*c**4
*d**7 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*c**2*d**9 + 18*log(t
an(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**2*c**3*d**7 - 6*log(tan(e + f*
x)**2 + 1)*tan(e + f*x)**2*a**2*b**2*c*d**9 - 3*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**2*a**2*b*c**5*d**6 + 9*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*
**2*a**2*b*c**3*d**8 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**3*c*
**4*d**6 + 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**3*c**2*d**8 - 9
*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**2*c**4*d**7 + 3*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)**2*a*b**2*c**2*d**9 - 3*log(tan(e + f*x)**2 + 1)
)*tan(e + f*x)**2*b**4*c**3*d**7 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)**
2*b**4*c*d**9 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b**3*c**5*d**6 -
3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b**3*c**3*d**8 - 6*log(tan(e +
f*x)**2 + 1)*tan(e + f*x)*a**4*c**4*d**6 + 2*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)*a**4*c**2*d**8 + 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b*c
**5*d**5 - 24*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*b*c**3*d**7 + 6*log
(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*c**5*d**6 - 2*log(tan(e + f*x)**
2 + 1)*tan(e + f*x)*a**3*c**3*d**8 + 36*log(tan(e + f*x)**2 + 1)*tan(e ...
```

3.85
$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$$

Optimal result	950
Mathematica [C] (verified)	951
Rubi [A] (verified)	952
Maple [A] (verified)	956
Fricas [B] (verification not implemented)	957
Sympy [F(-2)]	958
Maxima [A] (verification not implemented)	959
Giac [B] (verification not implemented)	959
Mupad [B] (verification not implemented)	960
Reduce [B] (verification not implemented)	961

Optimal result

Integrand size = 45, antiderivative size = 597

$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx =$$

$$\frac{(b^2(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) + a^2(c^3C - 3Bc^2d - 3cCd^2 + Bd^3 - A(c^3 - 3cd^2))}{(c^2 + d^2)^3}$$

$$\frac{(2ab(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) - a^2((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2)) + b^2((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2))}{(c^2 + d^2)^3 f}$$

$$\frac{(2abd^3(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) - b^2(c^6C + 3c^4Cd^2 + Bc^3d^3 - 3c^2(A - 2C)d^4 - B(c^3 - 3cd^2)))}{d^3 (c^2 + d^2)^3 f}$$

$$\frac{(c^2C - Bcd + Ad^2)(a+b \tan(e+fx))^2}{2d(c^2 + d^2) f(c+d \tan(e+fx))^2}$$

$$+ \frac{(bc - ad)(b(c^4C - c^2(A - 3C)d^2 - 2Bcd^3 + Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2)))}{d^3 (c^2 + d^2)^2 f(c+d \tan(e+fx))}$$

output

```

-(b^2*(A*c^3-3*A*c*d^2+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)+a^2*(c^3*C-3*B*c^2
*d-3*C*c*d^2+B*d^3-A*(c^3-3*c*d^2))-2*a*b*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*
d^2)))*x/(c^2+d^2)^3-(2*a*b*(A*c^3-3*A*c*d^2+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d
^2)-a^2*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2))+b^2*((A-C)*d*(3*c^2-d^2)-B*(
c^3-3*c*d^2))*ln(cos(f*x+e))/(c^2+d^2)^3/f-(2*a*b*d^3*(A*c^3-3*A*c*d^2+3*
B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)-b^2*(c^6*C+3*c^4*C*d^2+B*c^3*d^3-3*c^2*(A-2
*C)*d^4-3*B*c*d^5+A*d^6)-a^2*d^3*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2)))*ln
(c+d*tan(f*x+e))/d^3/(c^2+d^2)^3/f-1/2*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e)
)^2/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^2+(-a*d+b*c)*(b*(c^4*C-c^2*(A-3*C)*d^2-
2*B*c*d^3+A*d^4)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))/d^3/(c^2+d^2)^2/f/(c+d*t
an(f*x+e))

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.75 (sec) , antiderivative size = 404, normalized size of antiderivative = 0.68

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

$$= \frac{(a + ib)^2 (-iA + B + iC)(c - id)^3 \log(i - \tan(e + fx)) + (a - ib)^2 (iA + B - iC)(c + id)^3 \log(i + \tan(e + fx))}{(c + d \tan(e + fx))^3}$$

input

```

Integrate[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^3,x]

```

output

```

((a + I*b)^2*((-I)*A + B + I*C)*(c - I*d)^3*Log[I - Tan[e + f*x]] + (a - I
*b)^2*(I*A + B - I*C)*(c + I*d)^3*Log[I + Tan[e + f*x]] + (2*(2*a*b*d^3*(-
(A*c^3) + c^3*C - 3*B*c^2*d + 3*A*c*d^2 - 3*c*C*d^2 + B*d^3) + b^2*(c^6*C
+ 3*c^4*C*d^2 + B*c^3*d^3 - 3*c^2*(A - 2*C)*d^4 - 3*B*c*d^5 + A*d^6) + a^2
*d^3*(-((A - C)*d*(-3*c^2 + d^2)) - B*(c^3 - 3*c*d^2)))*Log[c + d*Tan[e +
f*x]])/d^3 - ((b*c - a*d)^2*(c^2 + d^2)^2*(c^2*C - B*c*d + A*d^2))/(d^3*(c
+ d*Tan[e + f*x])^2) + (2*(b*c - a*d)*(c^2 + d^2)*(b*(2*c^4*C - B*c^3*d +
4*c^2*C*d^2 - 3*B*c*d^3 + 2*A*d^4) + a*d^2*(2*c*(A - C)*d + B*(-c^2 + d^2
))))/(d^3*(c + d*Tan[e + f*x]))/(2*(c^2 + d^2)^3*f)

```

Rubi [A] (verified)

Time = 4.10 (sec) , antiderivative size = 631, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.244$, Rules used = {3042, 4128, 27, 3042, 4118, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{2(a + b \tan(e + fx))(bC(c^2 + d^2) \tan^2(e + fx) + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + bd) + (bc - ad)(cC - Bd))}{(c + d \tan(e + fx))^2} dx$$

$$\frac{2d(c^2 + d^2)}{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}$$

$$\frac{2df(c^2 + d^2)(c + d \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 27

$$\int \frac{(a + b \tan(e + fx))(bC(c^2 + d^2) \tan^2(e + fx) + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + bd) + (bc - ad)(cC - Bd))}{(c + d \tan(e + fx))^2} dx$$

$$\frac{d(c^2 + d^2)}{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}$$

$$\frac{2df(c^2 + d^2)(c + d \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))(bC(c^2 + d^2) \tan(e + fx)^2 + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + bd) + (bc - ad)(cC - Bd))}{(c + d \tan(e + fx))^2} dx$$

$$\frac{d(c^2 + d^2)}{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}$$

$$\frac{2df(c^2 + d^2)(c + d \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 4118

$$\int \frac{C(c^2+d^2)^2 \tan^2(e+fx)b^2 + (Cc^4 - (A-3C)d^2c^2 - 2Bd^3c + Ad^4)b^2 + 2ad^2(2c(A-C)d - B(c^2-d^2))b - a^2d^2(Cc^2 - 2Bdc - Cd^2 - A(c^2-d^2)) - d^2((2c(A-C)d - B(c^2-d^2)) - d^2((2c(A-C)d - B(c^2-d^2)))}{d(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{C(c^2+d^2)^2 \tan(e+fx)^2b^2 + (Cc^4 - (A-3C)d^2c^2 - 2Bd^3c + Ad^4)b^2 + 2ad^2(2c(A-C)d - B(c^2-d^2))b - a^2d^2(Cc^2 - 2Bdc - Cd^2 - A(c^2-d^2)) - d^2((2c(A-C)d - B(c^2-d^2)) - d^2((2c(A-C)d - B(c^2-d^2)))}{d(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 4109

$$\frac{d^2(-a^2(d(A-C)(3c^2-d^2) - B(c^3-3cd^2))) + 2ab(Ac^3 - 3Acd^2 + 3Bc^2d - Bd^3 - c^3C + 3cCd^2) + b^2(d(A-C)(3c^2-d^2) - B(c^3-3cd^2)) \int \tan(e+fx)dx}{c^2+d^2} - \frac{(-a^2d^2 \int \tan(e+fx)dx)}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 3042

$$\frac{d^2(-a^2(d(A-C)(3c^2-d^2) - B(c^3-3cd^2))) + 2ab(Ac^3 - 3Acd^2 + 3Bc^2d - Bd^3 - c^3C + 3cCd^2) + b^2(d(A-C)(3c^2-d^2) - B(c^3-3cd^2)) \int \tan(e+fx)dx}{c^2+d^2} - \frac{(-a^2d^2 \int \tan(e+fx)dx)}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 3956

$$\frac{(-a^2d^3(d(A-C)(3c^2-d^2) - B(c^3-3cd^2))) + 2abd^3(Ac^3 - 3Acd^2 + 3Bc^2d - Bd^3 - c^3C + 3cCd^2) - b^2(-3c^2d^4(A-2C) + Ad^6 + Bc^3d^3 - 3Bcd^5 + c^6C + 3c^4Cd^2)}{c^2+d^2} \int \tan(e+fx)dx$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{2df(c^2 + d^2)(c + d \tan(e + fx))^2}$$

↓ 4100

$$\frac{(-a^2 d^3 (d(A-C)(3c^2-d^2)-B(c^3-3cd^2))+2abd^3 (Ac^3-3Acd^2+3Bc^2d-Bd^3-c^3C+3cCd^2))-b^2 (-3c^2d^4(A-2C)+Ad^6+Bc^3d^3-3Bcd^5+c^6C+3c^4Cd^2)}{df(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2}$$

↓ 16

$$\frac{d^2 \log(\cos(e+fx))(-a^2(d(A-C)(3c^2-d^2)-B(c^3-3cd^2)))+2ab(Ac^3-3Acd^2+3Bc^2d-Bd^3-c^3C+3cCd^2)+b^2(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))}{f(c^2+d^2)}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2}$$

input

```
Int[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^3,x]
```

output

```
-1/2*((c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^2)/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^2) + (((-((d^2*(b^2*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) + a^2*(c^3*C - 3*B*c^2*d - 3*c*C*d^2 + B*d^3 - A*(c^3 - 3*c*d^2)) - 2*a*b*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2))))*x)/(c^2 + d^2)) - (d^2*(2*a*b*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) - a^2*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)) + b^2*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)))*Log[Cos[e + f*x]])/(c^2 + d^2)*f) - ((2*a*b*d^3*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) - b^2*(c^6*C + 3*c^4*C*d^2 + B*c^3*d^3 - 3*c^2*(A - 2*C)*d^4 - 3*B*c*d^5 + A*d^6) - a^2*d^3*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)))*Log[c + d*Tan[e + f*x]])/(d*(c^2 + d^2)*f))/(d*(c^2 + d^2)) + ((b*c - a*d)*(b*(c^4*C - c^2*(A - 3*C)*d^2 - 2*B*c*d^3 + A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))/(d^2*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))/(d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 3956 $\text{Int}[\tan[(c_)+(d_)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$
- rule 4100 $\text{Int}[(a_)+(b_)*\tan[(e_)+(f_)*(x_)]^{(m_)*((A_)+(C_)*\tan[(e_)+(f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[A/(b*f) \text{ Subst}[\text{Int}[(a + x)^m, x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, A, C, m\}, x] \&\& \text{EqQ}[A, C]$
- rule 4109 $\text{Int}[(A_)+(B_)*\tan[(e_)+(f_)*(x_)]+(C_)*\tan[(e_)+(f_)*(x_)]^2)/((a_)+(b_)*\tan[(e_)+(f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(1 + \text{Tan}[e + f*x]^2)/(a + b*\text{Tan}[e + f*x]), x], x] - \text{Simp}[(A*b - a*B - b*C)/(a^2 + b^2) \text{ Int}[\text{Tan}[e + f*x], x], x]) /; \text{FreeQ}[\{a, b, e, f, A, B, C\}, x] \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A*b - a*B - b*C, 0]$

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 865, normalized size of antiderivative = 1.45

method	result
derivativedivides	$\frac{(-3A^2a^2c^2d + A^2a^2d^3 + 2Aabc^3 - 6Aabc d^2 + 3Ab^2c^2d - Ab^2d^3 + Ba^2c^3 - 3Ba^2cd^2 + 6Babc^2d - 2Babd^3 - Bb^2c^3 + 3Bb^2cd^2 + 3Ca^2d^2)}{2}$
default	$\frac{(-3A^2a^2c^2d + A^2a^2d^3 + 2Aabc^3 - 6Aabc d^2 + 3Ab^2c^2d - Ab^2d^3 + Ba^2c^3 - 3Ba^2cd^2 + 6Babc^2d - 2Babd^3 - Bb^2c^3 + 3Bb^2cd^2 + 3Ca^2d^2)}{2}$
norman	Expression too large to display
risch	Expression too large to display
parallelrisc	Expression too large to display

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,
x,method=_RETURNVERBOSE)
```

output

```
1/f*(1/(c^2+d^2)^3*(1/2*(-3*A*a^2*c^2*d+A*a^2*d^3+2*A*a*b*c^3-6*A*a*b*c*d^
2+3*A*b^2*c^2*d-A*b^2*d^3+B*a^2*c^3-3*B*a^2*c*d^2+6*B*a*b*c^2*d-2*B*a*b*d^
3-B*b^2*c^3+3*B*b^2*c*d^2+3*C*a^2*c^2*d-C*a^2*d^3-2*C*a*b*c^3+6*C*a*b*c*d^
2-3*C*b^2*c^2*d+C*b^2*d^3)*ln(1+tan(f*x+e)^2)+(A*a^2*c^3-3*A*a^2*c*d^2+6*A
*a*b*c^2*d-2*A*a*b*d^3-A*b^2*c^3+3*A*b^2*c*d^2+3*B*a^2*c^2*d-B*a^2*d^3-2*B
*a*b*c^3+6*B*a*b*c*d^2-3*B*b^2*c^2*d+B*b^2*d^3-C*a^2*c^3+3*C*a^2*c*d^2-6*C
*a*b*c^2*d+2*C*a*b*d^3+C*b^2*c^3-3*C*b^2*c*d^2)*arctan(tan(f*x+e)))-1/2*(A
*a^2*d^4-2*A*a*b*c*d^3+A*b^2*c^2*d^2-B*a^2*c*d^3+2*B*a*b*c^2*d^2-B*b^2*c^3
*d+C*a^2*c^2*d^2-2*C*a*b*c^3*d+C*b^2*c^4)/d^3/(c^2+d^2)/(c+d*tan(f*x+e))^2
-(2*A*a^2*c*d^4-2*A*a*b*c^2*d^3+2*A*a*b*d^5-2*A*b^2*c*d^4-B*a^2*c^2*d^3+B*
a^2*d^5-4*B*a*b*c*d^4+B*b^2*c^4*d+3*B*b^2*c^2*d^3-2*C*a^2*c*d^4+2*C*a*b*c^
4*d+6*C*a*b*c^2*d^3-2*C*b^2*c^5-4*C*b^2*c^3*d^2)/d^3/(c^2+d^2)^2/(c+d*tan(
f*x+e)))+(3*A*a^2*c^2*d^4-A*a^2*d^6-2*A*a*b*c^3*d^3+6*A*a*b*c*d^5-3*A*b^2*c
^2*d^4+A*b^2*d^6-B*a^2*c^3*d^3+3*B*a^2*c*d^5-6*B*a*b*c^2*d^4+2*B*a*b*d^6+B
*b^2*c^3*d^3-3*B*b^2*c*d^5-3*C*a^2*c^2*d^4+C*a^2*d^6+2*C*a*b*c^3*d^3-6*C*a
*b*c*d^5+C*b^2*c^6+3*C*b^2*c^4*d^2+6*C*b^2*c^2*d^4)/(c^2+d^2)^3/d^3*ln(c+d
*tan(f*x+e))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1618 vs. 2(591) = 1182.

Time = 0.45 (sec) , antiderivative size = 1618, normalized size of antiderivative = 2.71

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e))^3,x, algorithm="fricas")
```

output

```

1/2*(C*b^2*c^6*d^2 - A*a^2*d^8 + (2*C*a*b + B*b^2)*c^5*d^3 - (3*C*a^2 + 6*
B*a*b + (3*A - 7*C)*b^2)*c^4*d^4 + 5*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^3*d
^5 - ((7*A - 3*C)*a^2 - 6*B*a*b - 3*A*b^2)*c^2*d^6 - (B*a^2 + 2*A*a*b)*c*d
^7 + 2*(((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^5*d^3 + 3*(B*a^2 + 2*(A -
C)*a*b - B*b^2)*c^4*d^4 - 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^3*d^5
- (B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2*d^6)*f*x - (3*C*b^2*c^6*d^2 + A*a^2*
d^8 - (2*C*a*b + B*b^2)*c^5*d^3 - (C*a^2 + 2*B*a*b + (A - 9*C)*b^2)*c^4*d^
4 + (3*B*a^2 + 2*(3*A - 7*C)*a*b - 7*B*b^2)*c^3*d^5 - 5*((A - C)*a^2 - 2*B
*a*b - A*b^2)*c^2*d^6 - 3*(B*a^2 + 2*A*a*b)*c*d^7 - 2*(((A - C)*a^2 - 2*B*
a*b - (A - C)*b^2)*c^3*d^5 + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2*d^6 - 3
*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^7 - (B*a^2 + 2*(A - C)*a*b - B*
b^2)*d^8)*f*x)*tan(f*x + e)^2 + (C*b^2*c^8 + 3*C*b^2*c^6*d^2 - (B*a^2 + 2*
(A - C)*a*b - B*b^2)*c^5*d^3 + 3*((A - C)*a^2 - 2*B*a*b - (A - 2*C)*b^2)*c
^4*d^4 + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^3*d^5 - ((A - C)*a^2 - 2*B*a*
b - A*b^2)*c^2*d^6 + (C*b^2*c^6*d^2 + 3*C*b^2*c^4*d^4 - (B*a^2 + 2*(A - C)
*a*b - B*b^2)*c^3*d^5 + 3*((A - C)*a^2 - 2*B*a*b - (A - 2*C)*b^2)*c^2*d^6
+ 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d^7 - ((A - C)*a^2 - 2*B*a*b - A*b^2
)*d^8)*tan(f*x + e)^2 + 2*(C*b^2*c^7*d + 3*C*b^2*c^5*d^3 - (B*a^2 + 2*(A -
C)*a*b - B*b^2)*c^4*d^4 + 3*((A - C)*a^2 - 2*B*a*b - (A - 2*C)*b^2)*c^3*d
^5 + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2*d^6 - ((A - C)*a^2 - 2*B*a*b...

```

Sympy [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

= Exception raised: AttributeError

input

```

integrate((a+b*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*
x+e))**3,x)

```

output

```

Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'

```

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 827, normalized size of antiderivative = 1.39

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="maxima")
```

output

```
1/2*(2*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^3 + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c^2*d - 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c*d^2 - (B*a^2 + 2*(A - C)*a*b - B*b^2)*d^3)*(f*x + e)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) + 2*(C*b^2*c^6 + 3*C*b^2*c^4*d^2 - (B*a^2 + 2*(A - C)*a*b - B*b^2)*c^3*d^3 + 3*((A - C)*a^2 - 2*B*a*b - (A - 2*C)*b^2)*c^2*d^4 + 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d^5 - ((A - C)*a^2 - 2*B*a*b - A*b^2)*d^6)*log(d*tan(f*x + e) + c)/(c^6*d^3 + 3*c^4*d^5 + 3*c^2*d^7 + d^9) + ((B*a^2 + 2*(A - C)*a*b - B*b^2)*c^3 - 3*((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*c^2*d - 3*(B*a^2 + 2*(A - C)*a*b - B*b^2)*c*d^2 + ((A - C)*a^2 - 2*B*a*b - (A - C)*b^2)*d^3)*log(tan(f*x + e)^2 + 1)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) + (3*C*b^2*c^6 - A*a^2*d^6 - (2*C*a*b + B*b^2)*c^5*d - (C*a^2 + 2*B*a*b + (A - 7*C)*b^2)*c^4*d^2 + (3*B*a^2 + 2*(3*A - 5*C)*a*b - 5*B*b^2)*c^3*d^3 - ((5*A - 3*C)*a^2 - 6*B*a*b - 3*A*b^2)*c^2*d^4 - (B*a^2 + 2*A*a*b)*c*d^5 + 2*(2*C*b^2*c^5*d + 4*C*b^2*c^3*d^3 - (2*C*a*b + B*b^2)*c^4*d^2 + (B*a^2 + 2*(A - 3*C)*a*b - 3*B*b^2)*c^2*d^4 - 2*((A - C)*a^2 - 2*B*a*b - A*b^2)*c*d^5 - (B*a^2 + 2*A*a*b)*d^6)*tan(f*x + e))/(c^6*d^3 + 2*c^4*d^5 + c^2*d^7 + (c^4*d^5 + 2*c^2*d^7 + d^9)*tan(f*x + e)^2 + 2*(c^5*d^4 + 2*c^3*d^6 + c*d^8)*tan(f*x + e))/f
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1219 vs. 2(591) = 1182.

Time = 0.69 (sec) , antiderivative size = 1219, normalized size of antiderivative = 2.04

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="giac")
```

output

```
(A*a^2*c^3 - C*a^2*c^3 - 2*B*a*b*c^3 - A*b^2*c^3 + C*b^2*c^3 + 3*B*a^2*c^2*d + 6*A*a*b*c^2*d - 6*C*a*b*c^2*d - 3*B*b^2*c^2*d - 3*A*a^2*c*d^2 + 3*C*a^2*c*d^2 + 6*B*a*b*c*d^2 + 3*A*b^2*c*d^2 - 3*C*b^2*c*d^2 - B*a^2*d^3 - 2*A*a*b*d^3 + 2*C*a*b*d^3 + B*b^2*d^3)*(f*x + e)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) + 1/2*(B*a^2*c^3 + 2*A*a*b*c^3 - 2*C*a*b*c^3 - B*b^2*c^3 - 3*A*a^2*c^2*d + 3*C*a^2*c^2*d + 6*B*a*b*c^2*d + 3*A*b^2*c^2*d - 3*C*b^2*c^2*d - 3*B*a^2*c*d^2 - 6*A*a*b*c*d^2 + 6*C*a*b*c*d^2 + 3*B*b^2*c*d^2 + A*a^2*d^3 - C*a^2*d^3 - 2*B*a*b*d^3 - A*b^2*d^3 + C*b^2*d^3)*log(tan(f*x + e)^2 + 1)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) + (C*b^2*c^6 + 3*C*b^2*c^4*d^2 - B*a^2*c^3*d^3 - 2*A*a*b*c^3*d^3 + 2*C*a*b*c^3*d^3 + B*b^2*c^3*d^3 + 3*A*a^2*c^2*d^4 - 3*C*a^2*c^2*d^4 - 6*B*a*b*c^2*d^4 - 3*A*b^2*c^2*d^4 + 6*C*b^2*c^2*d^4 + 3*B*a^2*c*d^5 + 6*A*a*b*c*d^5 - 6*C*a*b*c*d^5 - 3*B*b^2*c*d^5 - A*a^2*d^6 + C*a^2*d^6 + 2*B*a*b*d^6 + A*b^2*d^6)*log(abs(d*tan(f*x + e) + c))/(c^6*d^3*f + 3*c^4*d^5*f + 3*c^2*d^7*f + d^9*f) + 1/2*(2*(2*C*b^2*c^7 - 2*C*a*b*c^6*d - B*b^2*c^6*d + 6*C*b^2*c^5*d^2 + B*a^2*c^4*d^3 + 2*A*a*b*c^4*d^3 - 8*C*a*b*c^4*d^3 - 4*B*b^2*c^4*d^3 - 2*A*a^2*c^3*d^4 + 2*C*a^2*c^3*d^4 + 4*B*a*b*c^3*d^4 + 2*A*b^2*c^3*d^4 + 4*C*b^2*c^3*d^4 - 6*C*a*b*c^2*d^5 - 3*B*b^2*c^2*d^5 - 2*A*a^2*c*d^6 + 2*C*a^2*c*d^6 + 4*B*a*b*c*d^6 + 2*A*b^2*c*d^6 - B*a^2*d^7 - 2*A*a*b*d^7)*tan(f*x + e) + (3*C*b^2*c^8 - 2*C*a*b*c^7*d - B*b^2*c^7*d - C*a^2*c^6*d^2 - 2*B*a*b*c^6*d^2 - A...
```

Mupad [B] (verification not implemented)

Time = 23.27 (sec) , antiderivative size = 807, normalized size of antiderivative = 1.35

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx =$$

$$\frac{Aa^2 d^6 - 3Cb^2 c^6 + Ba^2 c d^5 + Bb^2 c^5 d + 5Aa^2 c^2 d^4 - 3Ab^2 c^2 d^4 + Ab^2 c^4 d^2 - 3Ba^2 c^3 d^3 + 5Bb^2 c^3 d^3 - 3Ca^2 c^2 d^4 + Ca^2 c^4 d^2 - 7Cb^2 c^4}{2d^3(c^4 + 2c^2 d^2 + d^4)}$$

$$\frac{\ln(c + d \tan(e + fx)) \left(\frac{c^2(d^4(3Ab^2 - 3Aa^2 + 3Ca^2 - 6Cb^2 + 6Bab) + 3Cb^2 d^4) - d^6(Ab^2 - Aa^2 + Ca^2 + 2Bab) + Cb^2 d^6 - c d^5}{c^6 d^3 + 3c^4 d^5 + 3c^2 d^7 + d^9} \right)}{\ln(\tan(e + fx) - i) \frac{(Ba^2 - Bb^2 + 2Aab - 2Cab - Aa^2 li + Ab^2 li + Ca^2 li - Cb^2 li + Bab 2i)}{2f(-c^3 - c^2 d 3i + 3cd^2 + d^3 li)}} +$$

$$\frac{\ln(\tan(e + fx) + i) (Ab^2 - Aa^2 + Ba^2 li - Bb^2 li + Ca^2 - Cb^2 + Aab 2i + 2Bab - Cab 2i)}{2f(-c^3 li - 3c^2 d + cd^2 3i + d^3)}$$

input `int(((a + b*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^3,x)`

output

$$\begin{aligned}
 & - ((Aa^2d^6 - 3Cb^2c^6 + Ba^2cd^5 + Bb^2c^5d + 5Aa^2c^2d^4 \\
 & - 3Ab^2c^2d^4 + Ab^2c^4d^2 - 3Ba^2c^3d^3 + 5Bb^2c^3d^3 - 3C \\
 & Ca^2c^2d^4 + Ca^2c^4d^2 - 7Cb^2c^4d^2 + 2Aa*b*c*d^5 + 2Ca*b* \\
 & c^5d - 6Aa*b*c^3d^3 - 6Ba*b*c^2d^4 + 2Ba*b*c^4d^2 + 10Ca*b*c^3 \\
 & *d^3)/(2d^3*(c^4 + d^4 + 2c^2d^2)) + (\tan(e + f*x)*(Ba^2d^5 - 2Cb^2 \\
 & *c^5 + 2Aa*b*d^5 + 2Aa^2*c*d^4 - 2Ab^2*c*d^4 + Bb^2*c^4*d - 2Ca^2 \\
 & *c*d^4 - Ba^2*c^2*d^3 + 3Bb^2*c^2*d^3 - 4Cb^2*c^3*d^2 - 4Ba*b*c*d^4 \\
 & + 2Ca*b*c^4*d - 2Aa*b*c^2*d^3 + 6Ca*b*c^2*d^3))/(d^2*(c^4 + d^4 + 2 \\
 & *c^2d^2)))/(f*(c^2 + d^2*tan(e + f*x)^2 + 2*c*d*tan(e + f*x))) - (\log(c + \\
 & d*tan(e + f*x))*((c^2*(d^4*(3Ab^2 - 3Aa^2 + 3Ca^2 - 6Cb^2 + 6Ba \\
 & *b) + 3Cb^2*d^4) - d^6*(Ab^2 - Aa^2 + Ca^2 + 2Ba*b) + Cb^2*d^6 - c \\
 & *d^5*(3Ba^2 - 3Bb^2 + 6Aa*b - 6Ca*b) + c^3*d^3*(Ba^2 - Bb^2 + 2A \\
 & Aa*b - 2Ca*b))/(d^9 + 3c^2*d^7 + 3c^4*d^5 + c^6*d^3) - (Cb^2/d^3))/ \\
 & f - (\log(\tan(e + f*x) - 1i)*(Ab^2*1i - Aa^2*1i + Ba^2 - Bb^2 + Ca^2*1 \\
 & i - Cb^2*1i + 2Aa*b + Ba*b*2i - 2Ca*b))/(2*f*(3*c*d^2 - c^2*d*3i - c \\
 & ^3 + d^3*1i)) - (\log(\tan(e + f*x) + 1i)*(Ab^2 - Aa^2 + Ba^2*1i - Bb^2* \\
 & 1i + Ca^2 - Cb^2 + Aa*b*2i + 2Ba*b - Ca*b*2i))/(2*f*(c*d^2*3i - 3*c^ \\
 & 2*d - c^3*1i + d^3))
 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 3698, normalized size of antiderivative = 6.19

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x)`

output

```
( - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**3*c**3*d**6 + log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a**3*c*d**8 + 3*log(tan(e + f*x)**2 + 1)*ta
n(e + f*x)**2*a**2*b*c**4*d**5 - 9*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**
2*a**2*b*c**2*d**7 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*c**4*
d**6 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*c**2*d**8 + 9*log(tan
(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**2*c**3*d**6 - 3*log(tan(e + f*x)**2
+ 1)*tan(e + f*x)**2*a*b**2*c*d**8 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f
*x)**2*a*b*c**5*d**5 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b*c**3
*d**7 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b**3*c**4*d**5 + 3*log(ta
n(e + f*x)**2 + 1)*tan(e + f*x)**2*b**3*c**2*d**7 - 3*log(tan(e + f*x)**2
+ 1)*tan(e + f*x)**2*b**2*c**4*d**6 + log(tan(e + f*x)**2 + 1)*tan(e + f*x
)**2*b**2*c**2*d**8 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*c**4*d*
*5 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**3*c**2*d**7 + 6*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)*a**2*b*c**5*d**4 - 18*log(tan(e + f*x)**2 + 1)
*tan(e + f*x)*a**2*b*c**3*d**6 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a
**2*c**5*d**5 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*c**3*d**7 + 1
8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b**2*c**4*d**5 - 6*log(tan(e + f
*x)**2 + 1)*tan(e + f*x)*a*b**2*c**2*d**7 - 4*log(tan(e + f*x)**2 + 1)*tan
(e + f*x)*a*b*c**6*d**4 + 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b*c**
4*d**6 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**3*c**5*d**4 + 6*log...
```

3.86 $\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$

Optimal result	963
Mathematica [C] (verified)	964
Rubi [A] (verified)	964
Maple [A] (verified)	968
Fricas [B] (verification not implemented)	969
Sympy [F(-2)]	970
Maxima [A] (verification not implemented)	970
Giac [B] (verification not implemented)	971
Mupad [B] (verification not implemented)	972
Reduce [B] (verification not implemented)	973

Optimal result

Integrand size = 43, antiderivative size = 352

$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx =$$

$$\frac{(a(c^3C - 3Bc^2d - 3cCd^2 + Bd^3 - A(c^3 - 3cd^2)) - b((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2))) x}{(c^2 + d^2)^3}$$

$$+ \frac{(b(c^3C - 3Bc^2d - 3cCd^2 + Bd^3) - a(Bc^3 + 3c^2Cd - 3Bcd^2 - Cd^3) + A(ad(3c^2 - d^2) - b(c^3 - 3cd^2)))}{(c^2 + d^2)^3 f}$$

$$+ \frac{(bc - ad)(c^2C - Bcd + Ad^2)}{2d^2(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

$$- \frac{b(c^4C - c^2(A - 3C)d^2 - 2Bcd^3 + Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2))}{d^2(c^2 + d^2)^2 f(c + d \tan(e + fx))}$$

output

```
-(a*(c^3*C-3*B*c^2*d-3*C*c*d^2+B*d^3-A*(c^3-3*c*d^2))-b*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2)))*x/(c^2+d^2)^3+(b*(-3*B*c^2*d+B*d^3+C*c^3-3*C*c*d^2)-a*(B*c^3-3*B*c*d^2+3*C*c^2*d-C*d^3)+A*(a*d*(3*c^2-d^2)-b*(c^3-3*c*d^2)))*ln((c*cos(f*x+e)+d*sin(f*x+e))/(c^2+d^2)^3/f+1/2*(-a*d+b*c)*(A*d^2-B*c*d+C*c^2)/d^2/(c^2+d^2)/f/(c+d*tan(f*x+e))^2-(b*(c^4*C-c^2*(A-3*C)*d^2-2*B*c*d^3+A*d^4)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))/d^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))
```


Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.50 (sec) , antiderivative size = 331, normalized size of antiderivative = 0.94

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

$$\frac{aCd - b(cC + Bd)}{(c + d \tan(e + fx))^2} - \frac{2Cd(a + b \tan(e + fx))}{(c + d \tan(e + fx))^2} + 2(Ab + aB - bC)d \left(-\frac{i \log(i - \tan(e + fx))}{2(c + id)^2} + \frac{i \log(i + \tan(e + fx))}{2(c - id)^2} + \frac{d(2c \log(c + d \tan(e + fx)))}{(c + d \tan(e + fx))^3} \right)$$

=

input

```
Integrate[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^3,x]
```

output

```
((a*C*d - b*(c*C + B*d))/(c + d*Tan[e + f*x])^2 - (2*C*d*(a + b*Tan[e + f*x]))/(c + d*Tan[e + f*x])^2 + 2*(A*b + a*B - b*C)*d*(((-1/2*I)*Log[I - Tan[e + f*x]])/(c + I*d)^2 + ((I/2)*Log[I + Tan[e + f*x]])/(c - I*d)^2 + (d*(2*c*Log[c + d*Tan[e + f*x]] - (c^2 + d^2)/(c + d*Tan[e + f*x])))/(c^2 + d^2)^2) - d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*(Log[I - Tan[e + f*x]]/((-I)*c + d)^3 + Log[I + Tan[e + f*x]]/(I*c + d)^3 + (d*((6*c^2 - 2*d^2)*Log[c + d*Tan[e + f*x]] - ((c^2 + d^2)*(5*c^2 + d^2 + 4*c*d*Tan[e + f*x]))/(c + d*Tan[e + f*x])^2))/(c^2 + d^2)^3)/(2*d^2*f)
```

Rubi [A] (verified)

Time = 2.51 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.08, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4118, 3042, 4111, 25, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx)) (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^3} dx$$

↓ 4118

$$\int \frac{bC(c^2+d^2) \tan^2(e+fx) + d(ABC+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{(c+d \tan(e+fx))^2} dx + \frac{d(c^2+d^2) (bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f (c^2+d^2) (c+d \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{bC(c^2+d^2) \tan(e+fx)^2 + d(ABC+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{(c+d \tan(e+fx))^2} dx + \frac{d(c^2+d^2) (bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f (c^2+d^2) (c+d \tan(e+fx))^2}$$

↓ 4111

$$\int \frac{d(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2))) + d(2aAc d-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{c+d \tan(e+fx)} dx - \frac{d(c^2+d^2) (bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f (c^2+d^2) (c+d \tan(e+fx))^2}$$

↓ 25

$$\int \frac{d(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2))) + d(2aAc d-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{c+d \tan(e+fx)} dx - \frac{d(c^2+d^2) (bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f (c^2+d^2) (c+d \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{d(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2))) + d(2aAc d-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{c+d \tan(e+fx)} dx - \frac{d(c^2+d^2) (bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f (c^2+d^2) (c+d \tan(e+fx))^2}$$

↓ 4014

$$\frac{\frac{d(aAd(3c^2-d^2)-a(Bc^3-3Bcd^2+3c^2Cd-Cd^3))-Ab(c^3-3cd^2)+b(-3Bc^2d+Bd^3+c^3C-3cCd^2)}{c^2+d^2} \int \frac{d-c \tan(e+fx)}{c+d \tan(e+fx)} dx - dx(a(Ac^3-3Acd^2+3Bc^2d-Bd^3))}{c^2+d^2} \frac{d(c^2+d^2)}{d(c^2+d^2)}$$

$$\frac{(bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 3042

$$\frac{\frac{d(aAd(3c^2-d^2)-a(Bc^3-3Bcd^2+3c^2Cd-Cd^3))-Ab(c^3-3cd^2)+b(-3Bc^2d+Bd^3+c^3C-3cCd^2)}{c^2+d^2} \int \frac{d-c \tan(e+fx)}{c+d \tan(e+fx)} dx - dx(a(Ac^3-3Acd^2+3Bc^2d-Bd^3))}{c^2+d^2} \frac{d(c^2+d^2)}{d(c^2+d^2)}$$

$$\frac{(bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f(c^2+d^2)(c+d \tan(e+fx))^2}$$

↓ 4013

$$\frac{(bc-ad)(Ad^2-Bcd+c^2C)}{2d^2 f(c^2+d^2)(c+d \tan(e+fx))^2} + \frac{ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-3C)+Ad^4-2Bcd^3+c^4C)}{df(c^2+d^2)(c+d \tan(e+fx))} - \frac{d(aAd(3c^2-d^2)-a(Bc^3-3Bcd^2+3c^2Cd-Cd^3))-Ab(c^3-3cd^2)+b(-3Bc^2d+Bd^3+c^3C-3cCd^2)}{f(c^2+d^2)}$$

```
input Int[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]^3,x]
```

```
output ((b*c - a*d)*(c^2*C - B*c*d + A*d^2))/(2*d^2*(c^2 + d^2)*f*(c + d*Tan[e + f*x]^2) + (-( -(d*(b*(A - C)*d*(3*c^2 - d^2) - b*B*(c^3 - 3*c*d^2) + a*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3))*x)/(c^2 + d^2) ) - (d*(a*A*d*(3*c^2 - d^2) - A*b*(c^3 - 3*c*d^2) + b*(c^3*C - 3*B*c^2*d - 3*c*C*d^2 + B*d^3) - a*(B*c^3 + 3*c^2*C*d - 3*B*c*d^2 - C*d^3))*Log[c*Cos[e + f*x] + d*Sin[e + f*x]]/(c^2 + d^2)*f)/(c^2 + d^2) - (b*(c^4*C - c^2*(A - 3*C)*d^2 - 2*B*c*d^3 + A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))/(d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_.), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 3042 $\text{Int}[\text{u}_., \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinear} \\ \text{Q}[\text{u}, \text{x}]$
- rule 4013 $\text{Int}[\text{((c}_.) + (\text{d}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) / ((\text{a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c}/(\text{b*f}))\text{Log}[\text{RemoveContent}[\text{a*Cos}[\text{e} + \text{f*x}] + \text{b*Si} \\ \text{n}[\text{e} + \text{f*x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{NeQ}[\text{b*c} - \text{a*d}, 0] \&\& \\ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{EqQ}[\text{a*c} + \text{b*d}, 0]$
- rule 4014 $\text{Int}[\text{((c}_.) + (\text{d}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]) / ((\text{a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{a*c} + \text{b*d})\text{(x}/(\text{a}^2 + \text{b}^2)), \text{x}] + \text{Simp}[(\text{b*c} - \text{a} \\ \text{*d})/(\text{a}^2 + \text{b}^2) \quad \text{Int}[(\text{b} - \text{a*Tan}[\text{e} + \text{f*x}]) / (\text{a} + \text{b*Tan}[\text{e} + \text{f*x}]), \text{x}], \text{x}] \text{ ;} \\ \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{NeQ}[\text{b*c} - \text{a*d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{N} \\ \text{eq}[\text{a*c} + \text{b*d}, 0]$
- rule 4111 $\text{Int}[\text{((a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])}^{\text{(m)}}\text{((A}_.) + (\text{B}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])} \\ + (\text{C}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{A*b}^2 - \\ \text{a*b*B} + \text{a}^2\text{C})\text{((a} + \text{b*Tan}[\text{e} + \text{f*x}])^{\text{(m} + 1)} / (\text{b*f}^{\text{(m} + 1)}\text{(a}^2 + \text{b}^2))), \text{x} \\] + \text{Simp}[1/(\text{a}^2 + \text{b}^2) \quad \text{Int}[(\text{a} + \text{b*Tan}[\text{e} + \text{f*x}])^{\text{(m} + 1)}\text{Simp}[\text{b*B} + \text{a*(A} - \\ \text{C)} - (\text{A*b} - \text{a*B} - \text{b*C})\text{Tan}[\text{e} + \text{f*x}], \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{A}, \text{B} \\ , \text{C}\}, \text{x}] \&\& \text{NeQ}[\text{A*b}^2 - \text{a*b*B} + \text{a}^2\text{C}, 0] \&\& \text{LtQ}[\text{m}, -1] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \\]$
- rule 4118 $\text{Int}[\text{((a}_.) + (\text{b}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])}^{\text{(n)}}\text{((A}_.) + (\text{B}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)])} \\ + (\text{C}_.)\text{tan}[(\text{e}_.) + (\text{f}_.)\text{(x}_.)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{b*c} - \text{a*d})\text{(c}^2\text{C} - \text{B*c*d} + \text{A*d}^2)\text{((c} \\ + \text{d*Tan}[\text{e} + \text{f*x}])^{\text{(n} + 1)} / (\text{d}^2\text{f}^{\text{(n} + 1)}\text{(c}^2 + \text{d}^2))), \text{x}] + \text{Simp}[1/(\text{d}^2\text{(c}^2 \\ + \text{d}^2)) \quad \text{Int}[(\text{c} + \text{d*Tan}[\text{e} + \text{f*x}])^{\text{(n} + 1)}\text{Simp}[\text{a*d}^2\text{(A*c} - \text{c*C} + \text{B*d}) + \text{b*} \\ (\text{c}^2\text{C} - \text{B*c*d} + \text{A*d}^2) + \text{d}^2\text{(A*b*c} + \text{a*B*c} - \text{b*c*C} - \text{a*A*d} + \text{b*B*d} + \text{a*C*d}) \\ \text{*Tan}[\text{e} + \text{f*x}] + \text{b*C}^2\text{(c}^2 + \text{d}^2)\text{Tan}[\text{e} + \text{f*x}]^2, \text{x}], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \\ \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}] \&\& \text{NeQ}[\text{b*c} - \text{a*d}, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{LtQ}[\text{n} \\ , -1]$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 493, normalized size of antiderivative = 1.40

method	result
derivativedivides	$\frac{(-3Aa^2c^2d + Aa^3d^3 + Ab^2c^3 - 3Abcd^2 + Ba^2c^3 - 3Bacd^2 + 3Bb^2c^2d - Bbd^3 + 3Ca^2c^2d - Ca^3d^3 - Cb^2c^3 + 3Cbc^2d^2) \ln(1 + \tan(fx+e))}{2(c^2+d^2)^3} + \frac{Aa^2c^2d + Aa^3d^3 + Ab^2c^3 - 3Abcd^2 + Ba^2c^3 - 3Bacd^2 + 3Bb^2c^2d - Bbd^3 + 3Ca^2c^2d - Ca^3d^3 - Cb^2c^3 + 3Cbc^2d^2}{(c^2+d^2)^3}$
default	$\frac{(-3Aa^2c^2d + Aa^3d^3 + Ab^2c^3 - 3Abcd^2 + Ba^2c^3 - 3Bacd^2 + 3Bb^2c^2d - Bbd^3 + 3Ca^2c^2d - Ca^3d^3 - Cb^2c^3 + 3Cbc^2d^2) \ln(1 + \tan(fx+e))}{2(c^2+d^2)^3} + \frac{Aa^2c^2d + Aa^3d^3 + Ab^2c^3 - 3Abcd^2 + Ba^2c^3 - 3Bacd^2 + 3Bb^2c^2d - Bbd^3 + 3Ca^2c^2d - Ca^3d^3 - Cb^2c^3 + 3Cbc^2d^2}{(c^2+d^2)^3}$
norman	$\frac{(Aa^2c^3 - 3Aacd^2 + 3Ab^2c^2d - Ab^3d^3 + 3Ba^2c^2d - Ba^3d^3 - Bb^2c^3 + 3Bbc^2d^2 - Ca^2c^3 + 3Cacd^2 - 3Cb^2c^2d + Cbd^3)c^2x}{(c^4 + 2c^2d^2 + d^4)(c^2 + d^2)} + \frac{d^2(Aa^2c^3 - 3Aacd^2 + 3Ab^2c^2d - Ab^3d^3 + 3Ba^2c^2d - Ba^3d^3 - Bb^2c^3 + 3Bbc^2d^2 - Ca^2c^3 + 3Cacd^2 - 3Cb^2c^2d + Cbd^3)}{(c^2 + d^2)^3}$
risch	Expression too large to display
parallelrisk	Expression too large to display

input `int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, method=_RETURNVERBOSE)`

output `1/f*(1/(c^2+d^2)^3*(1/2*(-3*A*a*c^2*d+A*a*d^3+A*b*c^3-3*A*b*c*d^2+B*a*c^3-3*B*a*c*d^2+3*B*b*c^2*d-B*b*d^3+3*C*a*c^2*d-C*a*d^3-C*b*c^3+3*C*b*c*d^2)*ln(1+tan(f*x+e)^2)+(A*a*c^3-3*A*a*c*d^2+3*A*b*c^2*d-A*b*d^3+3*B*a*c^2*d-B*a*d^3-B*b*c^3+3*B*b*c*d^2-C*a*c^3+3*C*a*c*d^2-3*C*b*c^2*d+C*b*d^3)*arctan(tan(f*x+e)))+(3*A*a*c^2*d-A*a*d^3-A*b*c^3+3*A*b*c*d^2-B*a*c^3+3*B*a*c*d^2-3*B*b*c^2*d+B*b*d^3-3*C*a*c^2*d+C*a*d^3+C*b*c^3-3*C*b*c*d^2)/(c^2+d^2)^3*ln(c+d*tan(f*x+e))-1/2*(A*a*d^3-A*b*c*d^2-B*a*c*d^2+B*b*c^2*d+C*a*c^2*d-C*b*c^3)/d^2/(c^2+d^2)/(c+d*tan(f*x+e))^2-(2*A*a*c*d^3-A*b*c^2*d^2+A*b*d^4-B*a*c^2*d^2+B*a*d^4-2*B*b*c*d^3-2*C*a*c*d^3+C*b*c^4+3*C*b*c^2*d^2)/(c^2+d^2)^2/d^2/(c+d*tan(f*x+e))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 897 vs. $2(350) = 700$.

Time = 0.16 (sec) , antiderivative size = 897, normalized size of antiderivative = 2.55

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="fricas")`

output

```
1/2*(C*b*c^5 - A*a*d^5 - 3*(C*a + B*b)*c^4*d + 5*(B*a + (A - C)*b)*c^3*d^2
- ((7*A - 3*C)*a - 3*B*b)*c^2*d^3 - (B*a + A*b)*c*d^4 + 2*(((A - C)*a - B
*b)*c^5 + 3*(B*a + (A - C)*b)*c^4*d - 3*((A - C)*a - B*b)*c^3*d^2 - (B*a +
(A - C)*b)*c^2*d^3)*f*x + (C*b*c^5 - A*a*d^5 + (C*a + B*b)*c^4*d - (3*B*a
+ (3*A - 7*C)*b)*c^3*d^2 + 5*((A - C)*a - B*b)*c^2*d^3 + 3*(B*a + A*b)*c*
d^4 + 2*(((A - C)*a - B*b)*c^3*d^2 + 3*(B*a + (A - C)*b)*c^2*d^3 - 3*((A -
C)*a - B*b)*c*d^4 - (B*a + (A - C)*b)*d^5)*f*x)*tan(f*x + e)^2 - ((B*a +
(A - C)*b)*c^5 - 3*((A - C)*a - B*b)*c^4*d - 3*(B*a + (A - C)*b)*c^3*d^2 +
((A - C)*a - B*b)*c^2*d^3 + ((B*a + (A - C)*b)*c^3*d^2 - 3*((A - C)*a - B
*b)*c^2*d^3 - 3*(B*a + (A - C)*b)*c*d^4 + ((A - C)*a - B*b)*d^5)*tan(f*x +
e)^2 + 2*((B*a + (A - C)*b)*c^4*d - 3*((A - C)*a - B*b)*c^3*d^2 - 3*(B*a
+ (A - C)*b)*c^2*d^3 + ((A - C)*a - B*b)*c*d^4)*tan(f*x + e))*log((d^2*tan
(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) + 2*((C*a +
B*b)*c^5 - (2*B*a + (2*A - 3*C)*b)*c^4*d + 3*((A - C)*a - B*b)*c^3*d^2 + 3
*(B*a + (A - C)*b)*c^2*d^3 - ((3*A - 2*C)*a - 2*B*b)*c*d^4 - (B*a + A*b)*d
^5 + 2*(((A - C)*a - B*b)*c^4*d + 3*(B*a + (A - C)*b)*c^3*d^2 - 3*((A - C)
*a - B*b)*c^2*d^3 - (B*a + (A - C)*b)*c*d^4)*f*x)*tan(f*x + e))/((c^6*d^2
+ 3*c^4*d^4 + 3*c^2*d^6 + d^8)*f*tan(f*x + e)^2 + 2*(c^7*d + 3*c^5*d^3 + 3
*c^3*d^5 + c*d^7)*f*tan(f*x + e) + (c^8 + 3*c^6*d^2 + 3*c^4*d^4 + c^2*d^6)
*f)
```

Sympy [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

= Exception raised: AttributeError

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**3,x)`

output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 543, normalized size of antiderivative = 1.54

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

$$= \frac{2(((A-C)a-Bb)c^3+3(Ba+(A-C)b)c^2d-3((A-C)a-Bb)cd^2-(Ba+(A-C)b)d^3)(fx+e)}{c^6+3c^4d^2+3c^2d^4+d^6} - \frac{2((Ba+(A-C)b)c^3-3((A-C)a-Bb)c^2d-3((A-C)a-Bb)cd^2-(Ba+(A-C)b)d^3)(fx+e)}{c^6+3c^4d^2+3c^2d^4+d^6}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="maxima")`

output

```

1/2*(2*((A - C)*a - B*b)*c^3 + 3*(B*a + (A - C)*b)*c^2*d - 3*((A - C)*a -
B*b)*c*d^2 - (B*a + (A - C)*b)*d^3)*(f*x + e)/(c^6 + 3*c^4*d^2 + 3*c^2*d^
4 + d^6) - 2*((B*a + (A - C)*b)*c^3 - 3*((A - C)*a - B*b)*c^2*d - 3*(B*a +
(A - C)*b)*c*d^2 + ((A - C)*a - B*b)*d^3)*log(d*tan(f*x + e) + c)/(c^6 +
3*c^4*d^2 + 3*c^2*d^4 + d^6) + ((B*a + (A - C)*b)*c^3 - 3*((A - C)*a - B*b
)*c^2*d - 3*(B*a + (A - C)*b)*c*d^2 + ((A - C)*a - B*b)*d^3)*log(tan(f*x +
e)^2 + 1)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) - (C*b*c^5 + A*a*d^5 + (C*a
+ B*b)*c^4*d - (3*B*a + (3*A - 5*C)*b)*c^3*d^2 + ((5*A - 3*C)*a - 3*B*b)*
c^2*d^3 + (B*a + A*b)*c*d^4 + 2*(C*b*c^4*d - (B*a + (A - 3*C)*b)*c^2*d^3 +
2*((A - C)*a - B*b)*c*d^4 + (B*a + A*b)*d^5)*tan(f*x + e))/(c^6*d^2 + 2*c
^4*d^4 + c^2*d^6 + (c^4*d^4 + 2*c^2*d^6 + d^8)*tan(f*x + e)^2 + 2*(c^5*d^3
+ 2*c^3*d^5 + c*d^7)*tan(f*x + e))/f

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 714 vs. $2(350) = 700$.

Time = 0.62 (sec) , antiderivative size = 714, normalized size of antiderivative = 2.03

$$\begin{aligned}
& \int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx \\
&= \frac{(Aac^3 - Cac^3 - Bbc^3 + 3Bac^2d + 3Abc^2d - 3Cbc^2d - 3Aacd^2 + 3Cacd^2 + 3Bbcd^2 - Bad^3 - Abd^3 + \\
&\quad c^6f + 3c^4d^2f + 3c^2d^4f + d^6f}{c^6f + 3c^4d^2f + 3c^2d^4f + d^6f} \\
&+ \frac{(Bac^3 + Abc^3 - Cbc^3 - 3Aac^2d + 3Cac^2d + 3Bbc^2d - 3Bacd^2 - 3Abcd^2 + 3Cbcd^2 + Aad^3 - Cad^3 - \\
&\quad 2(c^6f + 3c^4d^2f + 3c^2d^4f + d^6f))}{2(c^6f + 3c^4d^2f + 3c^2d^4f + d^6f)} \\
&- \frac{(Bac^3d + Abc^3d - Cbc^3d - 3Aac^2d^2 + 3Cac^2d^2 + 3Bbc^2d^2 - 3Bacd^3 - 3Abcd^3 + 3Cbcd^3 + Aad^4 - \\
&\quad c^6df + 3c^4d^3f + 3c^2d^5f + d^7f)}{c^6df + 3c^4d^3f + 3c^2d^5f + d^7f} \\
&- \frac{Cbc^7 + Cac^6d + Bbc^6d - 3Bac^5d^2 - 3Abc^5d^2 + 6Cbc^5d^2 + 5Aac^4d^3 - 2Cac^4d^3 - 2Bbc^4d^3 - 2Bacd^4 - \\
&\quad 2Abcd^4 + 2Cbcd^4 + Aad^5 - c^6d^2f + 3c^4d^4f + 3c^2d^6f + d^8f)}{c^6d^2f + 3c^4d^4f + 3c^2d^6f + d^8f}
\end{aligned}$$

input

```

integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^3,x, algorithm="giac")

```


output

```
(A*a*c^3 - C*a*c^3 - B*b*c^3 + 3*B*a*c^2*d + 3*A*b*c^2*d - 3*C*b*c^2*d - 3
*A*a*c*d^2 + 3*C*a*c*d^2 + 3*B*b*c*d^2 - B*a*d^3 - A*b*d^3 + C*b*d^3)*(f*x
+ e)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) + 1/2*(B*a*c^3 + A*b*c^3
- C*b*c^3 - 3*A*a*c^2*d + 3*C*a*c^2*d + 3*B*b*c^2*d - 3*B*a*c*d^2 - 3*A*b
*c*d^2 + 3*C*b*c*d^2 + A*a*d^3 - C*a*d^3 - B*b*d^3)*log(tan(f*x + e)^2 + 1
)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) - (B*a*c^3*d + A*b*c^3*d - C
*b*c^3*d - 3*A*a*c^2*d^2 + 3*C*a*c^2*d^2 + 3*B*b*c^2*d^2 - 3*B*a*c*d^3 - 3
*A*b*c*d^3 + 3*C*b*c*d^3 + A*a*d^4 - C*a*d^4 - B*b*d^4)*log(abs(d*tan(f*x
+ e) + c))/(c^6*d*f + 3*c^4*d^3*f + 3*c^2*d^5*f + d^7*f) - 1/2*(C*b*c^7 +
C*a*c^6*d + B*b*c^6*d - 3*B*a*c^5*d^2 - 3*A*b*c^5*d^2 + 6*C*b*c^5*d^2 + 5*
A*a*c^4*d^3 - 2*C*a*c^4*d^3 - 2*B*b*c^4*d^3 - 2*B*a*c^3*d^4 - 2*A*b*c^3*d^
4 + 5*C*b*c^3*d^4 + 6*A*a*c^2*d^5 - 3*C*a*c^2*d^5 - 3*B*b*c^2*d^5 + B*a*c*
d^6 + A*b*c*d^6 + A*a*d^7 + 2*(C*b*c^6*d - B*a*c^4*d^3 - A*b*c^4*d^3 + 4*C
*b*c^4*d^3 + 2*A*a*c^3*d^4 - 2*C*a*c^3*d^4 - 2*B*b*c^3*d^4 + 3*C*b*c^2*d^5
+ 2*A*a*c*d^6 - 2*C*a*c*d^6 - 2*B*b*c*d^6 + B*a*d^7 + A*b*d^7)*tan(f*x +
e))/((c^2 + d^2)^3*(d*tan(f*x + e) + c)^2*d^2*f)
```

Mupad [B] (verification not implemented)

Time = 12.17 (sec) , antiderivative size = 502, normalized size of antiderivative = 1.43

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx =$$

$$\frac{Aad^5 + Cbc^5 + Abcd^4 + Bacd^4 + Bbc^4d + Cac^4d + 5Aac^2d^3 - 3Abc^3d^2 - 3Bac^3d^2 - 3Bbc^2d^3 - 3Cac^2d^3 + 5Cbc^3d^2}{2d^2(c^4 + 2c^2d^2 + d^4)} + \frac{\tan(e + fx)}{f(c^2 + 2cd \tan(e + fx) + d^2 \tan^2(e + fx))}$$

$$- \frac{\ln(\tan(e + fx) + i)(Bb + Abli + Ba li - Aa + Ca - Cbli)}{2f(-c^3 li - 3c^2d + cd^2 3i + d^3)}$$

$$- \frac{\ln(\tan(e + fx) - i)(Ab + Ba - Cb - Aa li + Bbli + Ca li)}{2f(-c^3 - c^2d 3i + 3cd^2 + d^3 li)}$$

$$- \frac{\ln(c + d \tan(e + fx))((Ab + Ba - Cb)c^3 + (3Bb - 3Aa + 3Ca)c^2d + (3Cb - 3Ba - 3Ab)c)}{f(c^6 + 3c^4d^2 + 3c^2d^4 + d^6)}$$

input

```
int(((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*
tan(e + f*x))^3,x)
```

output

```
- ((A*a*d^5 + C*b*c^5 + A*b*c*d^4 + B*a*c*d^4 + B*b*c^4*d + C*a*c^4*d + 5*
A*a*c^2*d^3 - 3*A*b*c^3*d^2 - 3*B*a*c^3*d^2 - 3*B*b*c^2*d^3 - 3*C*a*c^2*d^
3 + 5*C*b*c^3*d^2)/(2*d^2*(c^4 + d^4 + 2*c^2*d^2)) + (tan(e + f*x)*(A*b*d^
4 + B*a*d^4 + C*b*c^4 + 2*A*a*c*d^3 - 2*B*b*c*d^3 - 2*C*a*c*d^3 - A*b*c^2*
d^2 - B*a*c^2*d^2 + 3*C*b*c^2*d^2))/(d*(c^4 + d^4 + 2*c^2*d^2)))/(f*(c^2 +
d^2*tan(e + f*x)^2 + 2*c*d*tan(e + f*x))) - (log(tan(e + f*x) + 1i)*(A*b*
1i - A*a + B*a*1i + B*b + C*a - C*b*1i))/(2*f*(c*d^2*3i - 3*c^2*d - c^3*1i
+ d^3)) - (log(tan(e + f*x) - 1i)*(A*b - A*a*1i + B*a + B*b*1i + C*a*1i -
C*b))/(2*f*(3*c*d^2 - c^2*d*3i - c^3 + d^3*1i)) - (log(c + d*tan(e + f*x)
))*(c^3*(A*b + B*a - C*b) - d^3*(B*b - A*a + C*a) + c^2*d*(3*B*b - 3*A*a +
3*C*a) - c*d^2*(3*A*b + 3*B*a - 3*C*b)))/(f*(c^6 + d^6 + 3*c^2*d^4 + 3*c^4
*d^2))
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 2450, normalized size of antiderivative = 6.96

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x)
```

output

```
( - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*c**3*d**4 + log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a**2*c*d**6 + 2*log(tan(e + f*x)**2 + 1)*ta
n(e + f*x)**2*a*b*c**4*d**3 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a
*b*c**2*d**5 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*c**4*d**4 - lo
g(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*c**2*d**6 + 3*log(tan(e + f*x)**2
+ 1)*tan(e + f*x)**2*b**2*c**3*d**4 - log(tan(e + f*x)**2 + 1)*tan(e + f*
x)**2*b**2*c*d**6 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b*c**5*d**3 +
3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b*c**3*d**5 - 6*log(tan(e + f*
x)**2 + 1)*tan(e + f*x)*a**2*c**4*d**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e
+ f*x)*a**2*c**2*d**5 + 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b*c**5*d
**2 - 12*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*b*c**3*d**4 + 6*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)*a*c**5*d**3 - 2*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)*a*c**3*d**5 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**2*c**4*d
**3 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*b**2*c**2*d**5 - 2*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)*b*c**6*d**2 + 6*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)*b*c**4*d**4 - 3*log(tan(e + f*x)**2 + 1)*a**2*c**5*d**2 + log(tan
(e + f*x)**2 + 1)*a**2*c**3*d**4 + 2*log(tan(e + f*x)**2 + 1)*a*b*c**6*d -
6*log(tan(e + f*x)**2 + 1)*a*b*c**4*d**3 + 3*log(tan(e + f*x)**2 + 1)*a*c
**6*d**2 - log(tan(e + f*x)**2 + 1)*a*c**4*d**4 + 3*log(tan(e + f*x)**2 +
1)*b**2*c**5*d**2 - log(tan(e + f*x)**2 + 1)*b**2*c**3*d**4 - log(tan(e...
```

3.87 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^3} dx$

Optimal result	975
Mathematica [C] (verified)	976
Rubi [A] (verified)	976
Maple [A] (verified)	979
Fricas [B] (verification not implemented)	980
Sympy [F(-2)]	981
Maxima [A] (verification not implemented)	981
Giac [A] (verification not implemented)	982
Mupad [B] (verification not implemented)	983
Reduce [B] (verification not implemented)	983

Optimal result

Integrand size = 33, antiderivative size = 209

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx$$

$$= -\frac{(c^3 C - 3Bc^2 d - 3cCd^2 + Bd^3 - A(c^3 - 3cd^2)) x}{(c^2 + d^2)^3}$$

$$+ \frac{((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2)) \log(c \cos(e + fx) + d \sin(e + fx))}{(c^2 + d^2)^3 f}$$

$$- \frac{c^2 C - Bcd + Ad^2}{2d(c^2 + d^2) f(c + d \tan(e + fx))^2} - \frac{2c(A - C)d - B(c^2 - d^2)}{(c^2 + d^2)^2 f(c + d \tan(e + fx))}$$

output

```
-(c^3*C-3*B*c^2*d-3*C*c*d^2+B*d^3-A*(c^3-3*c*d^2))*x/(c^2+d^2)^3+((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2))*ln(c*cos(f*x+e)+d*sin(f*x+e))/(c^2+d^2)^3/f-1/2*(A*d^2-B*c*d+C*c^2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^2-(2*c*(A-C)*d-B*(c^2-d^2))/(c^2+d^2)^2/f/(c+d*tan(f*x+e))
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.53 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.25

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx =$$

$$\frac{C}{(c + d \tan(e + fx))^2} + B \left(\frac{i \log(i - \tan(e + fx))}{(c + id)^2} - \frac{i \log(i + \tan(e + fx))}{(c - id)^2} + \frac{2d(-2c \log(c + d \tan(e + fx)) + \frac{c^2 + d^2}{c + d \tan(e + fx)})}{(c^2 + d^2)^2} \right) - (Bc +$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^3,x]
```

output

```
-1/2*(C/(c + d*Tan[e + f*x])^2 + B*((I*Log[I - Tan[e + f*x]])/(c + I*d)^2 - (I*Log[I + Tan[e + f*x]])/(c - I*d)^2 + (2*d*(-2*c*Log[c + d*Tan[e + f*x]] + (c^2 + d^2)/(c + d*Tan[e + f*x])))/(c^2 + d^2)^2) - (B*c + (-A + C)*d)*((I*Log[I - Tan[e + f*x]])/(c + I*d)^3 - Log[I + Tan[e + f*x]]/(I*c + d)^3 + (d*((-6*c^2 + 2*d^2)*Log[c + d*Tan[e + f*x]] + ((c^2 + d^2)*(5*c^2 + d^2 + 4*c*d*Tan[e + f*x]))/(c + d*Tan[e + f*x])^2))/(c^2 + d^2)^3)/(d*f)
```

Rubi [A] (verified)

Time = 1.63 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4111, 3042, 4012, 25, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(c + d \tan(e + fx))^3} dx$$

$$\begin{aligned}
& \int \frac{Ac - Cc + Bd + (Bc - (A - C)d) \tan(e + fx)}{(c + d \tan(e + fx))^2} dx \quad \downarrow \text{4111} \\
& \frac{Ad^2 - Bcd + c^2 C}{2df (c^2 + d^2) (c + d \tan(e + fx))^2} \\
& \int \frac{Ac - Cc + Bd + (Bc - (A - C)d) \tan(e + fx)}{(c + d \tan(e + fx))^2} dx \quad \downarrow \text{3042} \\
& \frac{Ad^2 - Bcd + c^2 C}{2df (c^2 + d^2) (c + d \tan(e + fx))^2} \\
& \int - \frac{Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2) + (2c(A - C)d - B(c^2 - d^2)) \tan(e + fx)}{c + d \tan(e + fx)} dx \quad \downarrow \text{4012} \\
& \frac{2cd(A - C) - B(c^2 - d^2)}{f(c^2 + d^2)(c + d \tan(e + fx))} \\
& \frac{c^2 + d^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2} \\
& \int - \frac{Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2) + (2c(A - C)d - B(c^2 - d^2)) \tan(e + fx)}{c + d \tan(e + fx)} dx \quad \downarrow \text{25} \\
& \frac{2cd(A - C) - B(c^2 - d^2)}{f(c^2 + d^2)(c + d \tan(e + fx))} \\
& \frac{c^2 + d^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2} \\
& \int \frac{Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2) + (2c(A - C)d - B(c^2 - d^2)) \tan(e + fx)}{c + d \tan(e + fx)} dx \quad \downarrow \text{3042} \\
& \frac{2cd(A - C) - B(c^2 - d^2)}{f(c^2 + d^2)(c + d \tan(e + fx))} \\
& \frac{c^2 + d^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2} \\
& \int - \frac{(d(A - C)(3c^2 - d^2) - B(c^3 - 3cd^2)) \int \frac{d - c \tan(e + fx)}{c + d \tan(e + fx)} dx - x(Ac^3 - 3Ac^2d + 3Bc^2d - Bd^3 - c^3C + 3cCd^2)}{c^2 + d^2} dx \quad \downarrow \text{4014} \\
& \frac{2cd(A - C) - B(c^2 - d^2)}{f(c^2 + d^2)(c + d \tan(e + fx))} \\
& \frac{c^2 + d^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2} \\
& \int \frac{Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2) + (2c(A - C)d - B(c^2 - d^2)) \tan(e + fx)}{c + d \tan(e + fx)} dx \quad \downarrow \text{3042} \\
& \frac{2cd(A - C) - B(c^2 - d^2)}{f(c^2 + d^2)(c + d \tan(e + fx))} \\
& \frac{c^2 + d^2}{2df (c^2 + d^2) (c + d \tan(e + fx))^2}
\end{aligned}$$

$$\begin{aligned}
& - \frac{(d(A-C)(3c^2-d^2) - B(c^3-3cd^2)) \int \frac{d-c \tan(e+fx)}{c+d \tan(e+fx)} dx - x \frac{(Ac^3-3Acd^2+3Bc^2d-Bd^3-c^3C+3cCd^2)}{c^2+d^2}}{c^2+d^2} - \frac{2cd(A-C) - B(c^2-d^2)}{f(c^2+d^2)(c+d \tan(e+fx))} \\
& \frac{c^2+d^2}{2df(c^2+d^2)(c+d \tan(e+fx))^2} \\
& \quad \downarrow \text{4013} \\
& - \frac{2cd(A-C) - B(c^2-d^2)}{f(c^2+d^2)(c+d \tan(e+fx))} - \frac{(d(A-C)(3c^2-d^2) - B(c^3-3cd^2)) \log(c \cos(e+fx) + d \sin(e+fx))}{f(c^2+d^2)} - \frac{x(Ac^3-3Acd^2+3Bc^2d-Bd^3-c^3C+3cCd^2)}{c^2+d^2} \\
& \frac{c^2+d^2}{2df(c^2+d^2)(c+d \tan(e+fx))^2}
\end{aligned}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^3,x]`

output `-1/2*(c^2*C - B*c*d + A*d^2)/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^2) + (-(((A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3)*x)/(c^2 + d^2)) - (((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2))*Log[c*Cos[e + f*x] + d*Sin[e + f*x]])/((c^2 + d^2)*f))/(c^2 + d^2) - (2*c*(A - C)*d - B*(c^2 - d^2))/((c^2 + d^2)*f*(c + d*Tan[e + f*x]))/(c^2 + d^2)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4013 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4014 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4111 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.25

method	result
derivativedivides	$\frac{(-3A c^2 d + A d^3 + B c^3 - 3B c d^2 + 3C c^2 d - C d^3) \ln(1 + \tan(fx + e)^2)}{2} + \frac{(A c^3 - 3A c d^2 + 3B c^2 d - B d^3 - c^3 C + 3C c d^2) \arctan(\tan(fx + e))}{(c^2 + d^2)^3}$
default	$\frac{(-3A c^2 d + A d^3 + B c^3 - 3B c d^2 + 3C c^2 d - C d^3) \ln(1 + \tan(fx + e)^2)}{2} + \frac{(A c^3 - 3A c d^2 + 3B c^2 d - B d^3 - c^3 C + 3C c d^2) \arctan(\tan(fx + e))}{(c^2 + d^2)^3}$
norman	$\frac{(A c^3 - 3A c d^2 + 3B c^2 d - B d^3 - c^3 C + 3C c d^2) c^2 x}{(c^4 + 2c^2 d^2 + d^4)(c^2 + d^2)} + \frac{d^2 (A c^3 - 3A c d^2 + 3B c^2 d - B d^3 - c^3 C + 3C c d^2) x \tan(fx + e)^2}{(c^4 + 2c^2 d^2 + d^4)(c^2 + d^2)} - \frac{5A e^2 d^3 + A d^5}{2(c^2 + d^2)}$
risch	Expression too large to display
parallelrisch	Expression too large to display

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/f*(1/(c^2+d^2)^3*(1/2*(-3*A*c^2*d+A*d^3+B*c^3-3*B*c*d^2+3*C*c^2*d-C*d^3)*ln(1+tan(f*x+e)^2)+(A*c^3-3*A*c*d^2+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)*arctan(tan(f*x+e)))-1/2*(A*d^2-B*c*d+C*c^2)/(c^2+d^2)/d/(c+d*tan(f*x+e))^2-(2*A*c*d-B*c^2+B*d^2-2*C*c*d)/(c^2+d^2)^2/(c+d*tan(f*x+e))+(3*A*c^2*d-A*d^3-B*c^3+3*B*c*d^2-3*C*c^2*d+C*d^3)/(c^2+d^2)^3*ln(c+d*tan(f*x+e)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 566 vs. $2(207) = 414$.

Time = 0.10 (sec) , antiderivative size = 566, normalized size of antiderivative = 2.71

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx =$$

$$\frac{3Cc^4d - 5Bc^3d^2 + (7A - 3C)c^2d^3 + Bcd^4 + Ad^5 - 2((A - C)c^5 + 3Bc^4d - 3(A - C)c^3d^2 - Bc^2d^3 - Bc^2d^3)}{(c + d \tan(e + fx))^3}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="fricas")`

output `-1/2*(3*C*c^4*d - 5*B*c^3*d^2 + (7*A - 3*C)*c^2*d^3 + B*c*d^4 + A*d^5 - 2*((A - C)*c^5 + 3*B*c^4*d - 3*(A - C)*c^3*d^2 - B*c^2*d^3)*f*x - (C*c^4*d - 3*B*c^3*d^2 + 5*(A - C)*c^2*d^3 + 3*B*c*d^4 - A*d^5 + 2*((A - C)*c^3*d^2 + 3*B*c^2*d^3 - 3*(A - C)*c*d^4 - B*d^5)*f*x)*tan(f*x + e)^2 + (B*c^5 - 3*(A - C)*c^4*d - 3*B*c^3*d^2 + (A - C)*c^2*d^3 + (B*c^3*d^2 - 3*(A - C)*c^2*d^3 - 3*B*c*d^4 + (A - C)*d^5)*tan(f*x + e)^2 + 2*(B*c^4*d - 3*(A - C)*c^3*d^2 - 3*B*c^2*d^3 + (A - C)*c*d^4)*tan(f*x + e))*log((d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2)/(tan(f*x + e)^2 + 1)) - 2*(C*c^5 - 2*B*c^4*d + 3*(A - C)*c^3*d^2 + 3*B*c^2*d^3 - (3*A - 2*C)*c*d^4 - B*d^5 + 2*((A - C)*c^4*d + 3*B*c^3*d^2 - 3*(A - C)*c^2*d^3 - B*c*d^4)*f*x)*tan(f*x + e))/((c^6*d^2 + 3*c^4*d^4 + 3*c^2*d^6 + d^8)*f*tan(f*x + e)^2 + 2*(c^7*d + 3*c^5*d^3 + 3*c^3*d^5 + c*d^7)*f*tan(f*x + e) + (c^8 + 3*c^6*d^2 + 3*c^4*d^4 + c^2*d^6)*f)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**3,x)`

output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.76

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx$$

$$= \frac{2((A-C)c^3+3Bc^2d-3(A-C)cd^2-Bd^3)(fx+e)}{c^6+3c^4d^2+3c^2d^4+d^6} - \frac{2(Bc^3-3(A-C)c^2d-3Bcd^2+(A-C)d^3) \log(d \tan(fx+e)+c)}{c^6+3c^4d^2+3c^2d^4+d^6} + \frac{(Bc^3-3(A-C)c^2d-3Bcd^2+(A-C)d^3)}{2f}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="maxima")`

output `1/2*(2*((A - C)*c^3 + 3*B*c^2*d - 3*(A - C)*c*d^2 - B*d^3)*(f*x + e)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) - 2*(B*c^3 - 3*(A - C)*c^2*d - 3*B*c*d^2 + (A - C)*d^3)*log(d*tan(f*x + e) + c)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) + (B*c^3 - 3*(A - C)*c^2*d - 3*B*c*d^2 + (A - C)*d^3)*log(tan(f*x + e)^2 + 1)/(c^6 + 3*c^4*d^2 + 3*c^2*d^4 + d^6) - (C*c^4 - 3*B*c^3*d + (5*A - 3*C)*c^2*d^2 + B*c*d^3 + A*d^4 - 2*(B*c^2*d^2 - 2*(A - C)*c*d^3 - B*d^4)*tan(f*x + e))/(c^6*d + 2*c^4*d^3 + c^2*d^5 + (c^4*d^3 + 2*c^2*d^5 + d^7)*tan(f*x + e)^2 + 2*(c^5*d^2 + 2*c^3*d^4 + c*d^6)*tan(f*x + e))/f`

Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 401, normalized size of antiderivative = 1.92

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx$$

$$= \frac{(Ac^3 - Cc^3 + 3Bc^2d - 3Acd^2 + 3Ccd^2 - Bd^3)(fx + e)}{c^6f + 3c^4d^2f + 3c^2d^4f + d^6f}$$

$$+ \frac{(Bc^3 - 3Ac^2d + 3Cc^2d^2 - 3Bcd^3 + Ad^3 - Cd^3) \log(\tan(fx + e)^2 + 1)}{2(c^6f + 3c^4d^2f + 3c^2d^4f + d^6f)}$$

$$- \frac{(Bc^3d - 3Ac^2d^2 + 3Cc^2d^2 - 3Bcd^3 + Ad^4 - Cd^4) \log(|d \tan(fx + e) + c|)}{c^6df + 3c^4d^3f + 3c^2d^5f + d^7f}$$

$$- \frac{Cc^6 - 3Bc^5d + 5Ac^4d^2 - 2Cc^4d^2 - 2Bc^3d^3 + 6Ac^2d^4 - 3Cc^2d^4 + Bcd^5 + Ad^6 - 2(Bc^4d^2 - 2Ac^3d^2)}{2(c^2 + d^2)^3(d \tan(fx + e) + c)^2df}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="giac")
```

output

```
(A*c^3 - C*c^3 + 3*B*c^2*d - 3*A*c*d^2 + 3*C*c*d^2 - B*d^3)*(f*x + e)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) + 1/2*(B*c^3 - 3*A*c^2*d + 3*C*c^2*d - 3*B*c*d^2 + A*d^3 - C*d^3)*log(tan(f*x + e)^2 + 1)/(c^6*f + 3*c^4*d^2*f + 3*c^2*d^4*f + d^6*f) - (B*c^3*d - 3*A*c^2*d^2 + 3*C*c^2*d^2 - 3*B*c*d^3 + A*d^4 - C*d^4)*log(abs(d*tan(f*x + e) + c))/(c^6*d*f + 3*c^4*d^3*f + 3*c^2*d^5*f + d^7*f) - 1/2*(C*c^6 - 3*B*c^5*d + 5*A*c^4*d^2 - 2*C*c^4*d^2 - 2*B*c^3*d^3 + 6*A*c^2*d^4 - 3*C*c^2*d^4 + B*c*d^5 + A*d^6 - 2*(B*c^4*d^2 - 2*A*c^3*d^2 + 2*C*c^3*d^3 - 2*A*c*d^5 + 2*C*c*d^5 - B*d^6)*tan(f*x + e))/((c^2 + d^2)^3*(d*tan(f*x + e) + c)^2*d*f)
```

Mupad [B] (verification not implemented)

Time = 8.12 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.56

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx$$

$$= - \frac{\frac{\tan(e+fx)(Bd^3+2Ac d^2-Bc^2d-2Ccd^2)}{c^4+2c^2d^2+d^4} + \frac{Ad^4+Cc^4+5Ac^2d^2-3Cc^2d^2+Bcd^3-3Bc^3d}{2d(c^4+2c^2d^2+d^4)}}{f(c^2+2cd \tan(e+fx)+d^2 \tan(e+fx)^2)}$$

$$- \frac{\ln(\tan(e+fx)-i)(B-A1i+C1i)}{2f(-c^3-c^2d3i+3cd^2+d^31i)}$$

$$- \frac{\ln(c+d \tan(e+fx))(Bc^3+(3C-3A)c^2d-3Bcd^2+(A-C)d^3)}{f(c^6+3c^4d^2+3c^2d^4+d^6)}$$

$$- \frac{\ln(\tan(e+fx)+1i)(C-A+B1i)}{2f(-c^31i-3c^2d+cd^23i+d^3)}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/(c + d*tan(e + f*x))^3,x)`

output `- ((tan(e + f*x)*(B*d^3 + 2*A*c*d^2 - B*c^2*d - 2*C*c*d^2))/(c^4 + d^4 + 2*c^2*d^2) + (A*d^4 + C*c^4 + 5*A*c^2*d^2 - 3*C*c^2*d^2 + B*c*d^3 - 3*B*c^3*d)/(2*d*(c^4 + d^4 + 2*c^2*d^2)))/(f*(c^2 + d^2*tan(e + f*x)^2 + 2*c*d*tan(e + f*x))) - (log(tan(e + f*x) - 1i)*(B - A*1i + C*1i))/(2*f*(3*c*d^2 - c^2*d*3i - c^3 + d^3*1i)) - (log(c + d*tan(e + f*x))*(B*c^3 + d^3*(A - C) - c^2*d*(3*A - 3*C) - 3*B*c*d^2))/(f*(c^6 + d^6 + 3*c^2*d^4 + 3*c^4*d^2)) - (log(tan(e + f*x) + 1i)*(B*1i - A + C))/(2*f*(c*d^2*3i - 3*c^2*d - c^3*1i + d^3))`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 1433, normalized size of antiderivative = 6.86

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x)`

output

```
( - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*c**3*d**4 + log(tan(e + f
*x)**2 + 1)*tan(e + f*x)**2*a*c*d**6 + log(tan(e + f*x)**2 + 1)*tan(e + f*
*x)**2*b*c**4*d**3 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b*c**2*d**5
+ 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*c**4*d**4 - log(tan(e + f*x)
**2 + 1)*tan(e + f*x)**2*c**2*d**6 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*
*x)*a*c**4*d**3 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a*c**2*d**5 + 2*l
og(tan(e + f*x)**2 + 1)*tan(e + f*x)*b*c**5*d**2 - 6*log(tan(e + f*x)**2 +
1)*tan(e + f*x)*b*c**3*d**4 + 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*c**
5*d**3 - 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*c**3*d**5 - 3*log(tan(e +
f*x)**2 + 1)*a*c**5*d**2 + log(tan(e + f*x)**2 + 1)*a*c**3*d**4 + log(tan
(e + f*x)**2 + 1)*b*c**6*d - 3*log(tan(e + f*x)**2 + 1)*b*c**4*d**3 + 3*lo
g(tan(e + f*x)**2 + 1)*c**6*d**2 - log(tan(e + f*x)**2 + 1)*c**4*d**4 + 6*
log(tan(e + f*x)*d + c)*tan(e + f*x)**2*a*c**3*d**4 - 2*log(tan(e + f*x)*d
+ c)*tan(e + f*x)**2*a*c*d**6 - 2*log(tan(e + f*x)*d + c)*tan(e + f*x)**2
*b*c**4*d**3 + 6*log(tan(e + f*x)*d + c)*tan(e + f*x)**2*b*c**2*d**5 - 6*l
og(tan(e + f*x)*d + c)*tan(e + f*x)**2*c**4*d**4 + 2*log(tan(e + f*x)*d +
c)*tan(e + f*x)**2*c**2*d**6 + 12*log(tan(e + f*x)*d + c)*tan(e + f*x)*a*c
**4*d**3 - 4*log(tan(e + f*x)*d + c)*tan(e + f*x)*a*c**2*d**5 - 4*log(tan(
e + f*x)*d + c)*tan(e + f*x)*b*c**5*d**2 + 12*log(tan(e + f*x)*d + c)*tan(
e + f*x)*b*c**3*d**4 - 12*log(tan(e + f*x)*d + c)*tan(e + f*x)*c**5*d**...
```

3.88 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^3} dx$

Optimal result	985
Mathematica [A] (verified)	986
Rubi [A] (verified)	987
Maple [A] (verified)	990
Fricas [B] (verification not implemented)	991
Sympy [F(-2)]	992
Maxima [B] (verification not implemented)	992
Giac [B] (verification not implemented)	993
Mupad [B] (verification not implemented)	994
Reduce [B] (verification not implemented)	995

Optimal result

Integrand size = 45, antiderivative size = 487

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx =$$

$$\frac{(a(c^3C - 3Bc^2d - 3cCd^2 + Bd^3 - A(c^3 - 3cd^2)) + b((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2))) x}{(a^2 + b^2)(c^2 + d^2)^3}$$

$$+ \frac{b^2(Ab^2 - a(bB - aC)) \log(a \cos(e + fx) + b \sin(e + fx))}{(a^2 + b^2)(bc - ad)^3 f}$$

$$- \frac{(b^2(c^6C - 3Bc^5d + 3c^4(2A - C)d^2 + Bc^3d^3 + 3Ac^2d^4 + Ad^6) + a^2d^3((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2)))}{(bc - ad)^3(c^2 + d^2)}$$

$$+ \frac{c^2C - Bcd + Ad^2}{2(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

$$+ \frac{b(c^4C - 2Bc^3d + c^2(3A - C)d^2 + Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2))}{(bc - ad)^2(c^2 + d^2)^2 f(c + d \tan(e + fx))}$$

output

```

-(a*(c^3*C-3*B*c^2*d-3*C*c*d^2+B*d^3-A*(c^3-3*c*d^2))+b*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2)))*x/(a^2+b^2)/(c^2+d^2)^3+b^2*(A*b^2-a*(B*b-C*a))*ln(a*cos(f*x+e)+b*sin(f*x+e))/(a^2+b^2)/(-a*d+b*c)^3/f-(b^2*(c^6*C-3*B*c^5*d+3*c^4*(2*A-C)*d^2+B*c^3*d^3+3*A*c^2*d^4+A*d^6)+a^2*d^3*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2))-a*b*d^2*(8*c^3*(A-C)*d-B*(3*c^4-6*c^2*d^2-d^4)))*ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^3/(c^2+d^2)^3/f+1/2*(A*d^2-B*c*d+C*c^2)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))^2+(b*(c^4*C-2*B*c^3*d+c^2*(3*A-C)*d^2+A*d^4)-a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))/(-a*d+b*c)^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))
    
```

Mathematica [A] (verified)

Time = 7.84 (sec) , antiderivative size = 912, normalized size of antiderivative = 1.87

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx$$

$$= -\frac{Ad^2 - c(-cC + Bd)}{2(-bc + ad)(c^2 + d^2) f(c + d \tan(e + fx))^2}$$

$$-\frac{b(bc - ad)^2 \left(Abc^3 - aBc^3 - bc^3C + 3aAc^2d + 3bBc^2d - 3ac^2Cd - 3Abcd^2 + 3aBcd^2 + 3bcCd^2 - aAd^3 - bBd^3 + aCd^3 - \sqrt{-b^2} \left(a(c^3C - 3Bc^2d - 3cCd^2 + Bd^3 - \dots) \right) \right)}{(a^2 + b^2)(c^2 + d^2)}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^3),x]
    
```

output

```

-1/2*(A*d^2 - c*(-(c*C) + B*d))/((-b*c) + a*d)*(c^2 + d^2)*f*(c + d*Tan[e
+ f*x])^2) - (-((-(b*(b*c - a*d)^2*(A*b*c^3 - a*B*c^3 - b*c^3*C + 3*a*A*
c^2*d + 3*b*B*c^2*d - 3*a*c^2*C*d - 3*A*b*c*d^2 + 3*a*B*c*d^2 + 3*b*c*C*d^
2 - a*A*d^3 - b*B*d^3 + a*C*d^3 - (Sqrt[-b^2]*(a*(c^3*C - 3*B*c^2*d - 3*c*
C*d^2 + B*d^3 - A*(c^3 - 3*c*d^2)) + b*((A - C)*d*(3*c^2 - d^2) - B*(c^3 -
3*c*d^2))))/b)*Log[Sqrt[-b^2] - b*Tan[e + f*x]])/((a^2 + b^2)*(c^2 + d^2)
)) + (2*b^3*(A*b^2 - a*(b*B - a*C))*(c^2 + d^2)^2*Log[a + b*Tan[e + f*x]])
/((a^2 + b^2)*(b*c - a*d)) - (b*(b*c - a*d)^2*(A*b*c^3 - a*B*c^3 - b*c^3*C
+ 3*a*A*c^2*d + 3*b*B*c^2*d - 3*a*c^2*C*d - 3*A*b*c*d^2 + 3*a*B*c*d^2 + 3
*b*c*C*d^2 - a*A*d^3 - b*B*d^3 + a*C*d^3 + (Sqrt[-b^2]*(b*(A - C)*d*(3*c^2
- d^2) - b*B*(c^3 - 3*c*d^2) - a*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 +
3*c*C*d^2 - B*d^3)))/b)*Log[Sqrt[-b^2] + b*Tan[e + f*x]])/((a^2 + b^2)*(c
^2 + d^2)) - (2*b*(b^2*(c^6*C - 3*B*c^5*d + 3*c^4*(2*A - C)*d^2 + B*c^3*d^
3 + 3*A*c^2*d^4 + A*d^6) + a^2*d^3*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c
*d^2)) - a*b*d^2*(8*c^3*(A - C)*d - B*(3*c^4 - 6*c^2*d^2 - d^4)))*Log[c +
d*Tan[e + f*x]])/((b*c - a*d)*(c^2 + d^2))/((b*(-b*c) + a*d)*(c^2 + d^2)*
f) - (-2*d^2*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)) - c*(2*d*(b*c
- a*d)*(B*c - (A - C)*d) - 2*b*c*(c^2*C - B*c*d + A*d^2)))/((-b*c) + a*d)
*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))/(2*(-b*c) + a*d)*(c^2 + d^2))

```

Rubi [A] (verified)

Time = 4.20 (sec) , antiderivative size = 549, normalized size of antiderivative = 1.13, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx$$

↓ 4132

$$\int \frac{-2(-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2))}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx + \frac{2(c^2 + d^2)(bc - ad)}{Ad^2 - Bcd + c^2 C} \frac{1}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

27

$$\frac{Ad^2 - Bcd + c^2 C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} \int \frac{-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

3042

$$\frac{Ad^2 - Bcd + c^2 C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} \int \frac{-b(Cc^2 - Bdc + Ad^2) \tan(e + fx)^2 - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^2} dx$$

4132

$$\frac{Ad^2 - Bcd + c^2 C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} \int \frac{(2c(A - C)d - B(c^2 - d^2)) \tan(e + fx)(bc - ad)^2 - b(b(Cc^4 - 2Bdc^3 + (3A - C)d^2 c^2 + Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2))) \tan^2(e + fx) + A(2abdc^3 - b^2(c^2 + d^2)^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))(c^2 + d^2)(bc - ad)} dx$$

3042

$$\frac{Ad^2 - Bcd + c^2 C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} \int \frac{(2c(A - C)d - B(c^2 - d^2)) \tan(e + fx)(bc - ad)^2 - b(b(Cc^4 - 2Bdc^3 + (3A - C)d^2 c^2 + Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2))) \tan(e + fx)^2 + A(2abdc^3 - b^2(c^2 + d^2)^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))(c^2 + d^2)(bc - ad)} dx$$

4134

$$\frac{Ad^2 - Bcd + c^2 C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} \int \frac{b^2(c^2 + d^2)^2 (Ab^2 - a(bB - aC)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx + (a^2 d^3 (d(A - C)(3c^2 - d^2) - B(c^3 - 3cd^2)) - ab d^2 (8c^3 d(A - C) - B(3c^4 - 6c^2 d^2 - d^4)) + b^2 (3c^4 d^2 (2A - C))}{(a^2 + b^2)(bc - ad)} dx}{(c^2 + d^2)(bc - ad)}$$

3042

$$\frac{Ad^2 - Bcd + c^2C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} - \frac{b^2(c^2 + d^2)^2 (Ab^2 - a(bB - aC)) \int \frac{b - a \tan(e + fx)}{a + b \tan(e + fx)} dx}{(a^2 + b^2)(bc - ad)} + \frac{(a^2 d^3 (d(A - C)(3c^2 - d^2) - B(c^3 - 3cd^2)) - abd^2 (8c^3 d(A - C) - B(3c^4 - 6c^2 d^2 - d^4)) + b^2 (3c^4 d^2 (2A - C) - B(c^3 - 3cd^2)))}{(c^2 + d^2)(bc - ad)}$$

↓ 4013

$$\frac{Ad^2 - Bcd + c^2C}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2} - \frac{b^2(c^2 + d^2)^2 (Ab^2 - a(bB - aC)) \log(a \cos(e + fx) + b \sin(e + fx))}{f(a^2 + b^2)(bc - ad)} + \frac{x(bc - ad)^2 (a(-A(c^3 - 3cd^2) - 3Bc^2 d + Bd^3 + c^3 C - 3cCd^2) + b(d(A - C)(3c^2 - d^2) - B(c^3 - 3cd^2)))}{(a^2 + b^2)(c^2 + d^2)}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^3), x]
```

output

```
(c^2*C - B*c*d + A*d^2)/(2*(b*c - a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^2) - (((b*c - a*d)^2*(a*(c^3*C - 3*B*c^2*d - 3*c*C*d^2 + B*d^3 - A*(c^3 - 3*c*d^2)) + b*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)))*x)/((a^2 + b^2)*(c^2 + d^2)) - (b^2*(A*b^2 - a*(b*B - a*C))*(c^2 + d^2)^2*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f) + ((b^2*(c^6*C - 3*B*c^5*d + 3*c^4*(2*A - C)*d^2 + B*c^3*d^3 + 3*A*c^2*d^4 + A*d^6) + a^2*d^3*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)) - a*b*d^2*(8*c^3*(A - C)*d - B*(3*c^4 - 6*c^2*d^2 - d^4)))*Log[c*Cos[e + f*x] + d*Sin[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f)/((b*c - a*d)*(c^2 + d^2)) - (b*(c^4*C - 2*B*c^3*d + c^2*(3*A - C)*d^2 + A*d^4) - a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))/((b*c - a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x]))/((b*c - a*d)*(c^2 + d^2))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4013

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4132

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^
2)/(((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 1.37 (sec) , antiderivative size = 649, normalized size of antiderivative = 1.33

method	result
derivativedivides	$\frac{(-3Aa^2c^2d + Aa^2d^3 - Abc^3 + 3Abcd^2 + Ba^2c^3 - 3Bac^2d - 3Bbc^2d + Bbd^3 + 3Ca^2c^2d - Ca^2d^3 + Cbc^3 - 3Cbc^2d^2) \ln(1 + \tan(fx + e))}{2(a^2 + b^2)(c^2 + d^2)} + \frac{A}{(a^2 + b^2)(c^2 + d^2)}$
default	$\frac{(-3Aa^2c^2d + Aa^2d^3 - Abc^3 + 3Abcd^2 + Ba^2c^3 - 3Bac^2d - 3Bbc^2d + Bbd^3 + 3Ca^2c^2d - Ca^2d^3 + Cbc^3 - 3Cbc^2d^2) \ln(1 + \tan(fx + e))}{2(a^2 + b^2)(c^2 + d^2)} + \frac{A}{(a^2 + b^2)(c^2 + d^2)}$
norman	Expression too large to display
parallelrisch	Expression too large to display
risch	Expression too large to display

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^3,x,
method=_RETURNVERBOSE)
```

output

```
1/f*(1/(a^2+b^2)/(c^2+d^2)^3*(1/2*(-3*A*a*c^2*d+A*a*d^3-A*b*c^3+3*A*b*c*d^2+B*a*c^3-3*B*a*c*d^2-3*B*b*c^2*d+B*b*d^3+3*C*a*c^2*d-C*a*d^3+C*b*c^3-3*C*b*c*d^2)*ln(1+tan(f*x+e)^2)+(A*a*c^3-3*A*a*c*d^2-3*A*b*c^2*d+A*b*d^3+3*B*a*c^2*d-B*a*d^3+B*b*c^3-3*B*b*c*d^2-C*a*c^3+3*C*a*c*d^2+3*C*b*c^2*d-C*b*d^3)*arctan(tan(f*x+e)))-(A*b^2-B*a*b+C*a^2)*b^2/(a*d-b*c)^3/(a^2+b^2)*ln(a+b*tan(f*x+e))-(2*A*a*c*d^3-3*A*b*c^2*d^2-A*b*d^4-B*a*c^2*d^2+B*a*d^4+2*B*b*c^3*d-2*C*a*c*d^3-C*b*c^4+C*b*c^2*d^2)/(c^2+d^2)^2/(a*d-b*c)^2/(c+d*tan(f*x+e)))+(3*A*a^2*c^2*d^4-A*a^2*d^6-8*A*a*b*c^3*d^3+6*A*b^2*c^4*d^2+3*A*b^2*c^2*d^4+A*b^2*d^6-B*a^2*c^3*d^3+3*B*a^2*c*d^5+3*B*a*b*c^4*d^2-6*B*a*b*c^2*d^4-B*a*b*d^6-3*B*b^2*c^5*d+B*b^2*c^3*d^3-3*C*a^2*c^2*d^4+C*a^2*d^6+8*C*a*b*c^3*d^3+C*b^2*c^6-3*C*b^2*c^4*d^2)/(c^2+d^2)^3/(a*d-b*c)^3*ln(c+d*tan(f*x+e))-1/2*(A*d^2-B*c*d+C*c^2)/(c^2+d^2)/(a*d-b*c)/(c+d*tan(f*x+e))^2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3496 vs. 2(485) = 970.

Time = 3.65 (sec) , antiderivative size = 3496, normalized size of antiderivative = 7.18

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^3,x, algorithm="fricas")`

output Too large to include

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx$$

= Exception raised: NotImplementedError

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))**3,x)`

output Exception raised: NotImplementedError >> no valid subset found

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1078 vs. 2(485) = 970.

Time = 0.17 (sec) , antiderivative size = 1078, normalized size of antiderivative = 2.21

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^3,x, algorithm="maxima")`

output

```

1/2*(2*((A - C)*a + B*b)*c^3 + 3*(B*a - (A - C)*b)*c^2*d - 3*((A - C)*a +
B*b)*c*d^2 - (B*a - (A - C)*b)*d^3)*(f*x + e)/((a^2 + b^2)*c^6 + 3*(a^2 +
b^2)*c^4*d^2 + 3*(a^2 + b^2)*c^2*d^4 + (a^2 + b^2)*d^6) + 2*(C*a^2*b^2 -
B*a*b^3 + A*b^4)*log(b*tan(f*x + e) + a)/((a^2*b^3 + b^5)*c^3 - 3*(a^3*b^2
+ a*b^4)*c^2*d + 3*(a^4*b + a^2*b^3)*c*d^2 - (a^5 + a^3*b^2)*d^3) - 2*(C*
b^2*c^6 - 3*B*b^2*c^5*d + 3*B*a^2*c*d^5 + 3*(B*a*b + (2*A - C)*b^2)*c^4*d^
2 - (B*a^2 + 8*(A - C)*a*b - B*b^2)*c^3*d^3 + 3*((A - C)*a^2 - 2*B*a*b + A
*b^2)*c^2*d^4 - ((A - C)*a^2 + B*a*b - A*b^2)*d^6)*log(d*tan(f*x + e) + c)
/(b^3*c^9 - 3*a*b^2*c^8*d + 3*a^2*b*c*d^8 - a^3*d^9 + 3*(a^2*b + b^3)*c^7*
d^2 - (a^3 + 9*a*b^2)*c^6*d^3 + 3*(3*a^2*b + b^3)*c^5*d^4 - 3*(a^3 + 3*a*b
^2)*c^4*d^5 + (9*a^2*b + b^3)*c^3*d^6 - 3*(a^3 + a*b^2)*c^2*d^7) + ((B*a -
(A - C)*b)*c^3 - 3*((A - C)*a + B*b)*c^2*d - 3*(B*a - (A - C)*b)*c*d^2 +
((A - C)*a + B*b)*d^3)*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*c^6 + 3*(a^2 +
b^2)*c^4*d^2 + 3*(a^2 + b^2)*c^2*d^4 + (a^2 + b^2)*d^6) + (3*C*b*c^5 - A*
a*d^5 - (C*a + 5*B*b)*c^4*d + (3*B*a + (7*A - C)*b)*c^3*d^2 - ((5*A - 3*C)
*a + B*b)*c^2*d^3 - (B*a - 3*A*b)*c*d^4 + 2*(C*b*c^4*d - 2*B*b*c^3*d^2 - 2
*(A - C)*a*c*d^4 + (B*a + (3*A - C)*b)*c^2*d^3 - (B*a - A*b)*d^5)*tan(f*x
+ e))/(b^2*c^8 - 2*a*b*c^7*d - 4*a*b*c^5*d^3 - 2*a*b*c^3*d^5 + a^2*c^2*d^6
+ (a^2 + 2*b^2)*c^6*d^2 + (2*a^2 + b^2)*c^4*d^4 + (b^2*c^6*d^2 - 2*a*b*c^
5*d^3 - 4*a*b*c^3*d^5 - 2*a*b*c*d^7 + a^2*d^8 + (a^2 + 2*b^2)*c^4*d^4 + ...

```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1447 vs. $2(485) = 970$.

Time = 0.79 (sec) , antiderivative size = 1447, normalized size of antiderivative = 2.97

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)
)^3,x, algorithm="giac")

```

output

```
(A*a*c^3 - C*a*c^3 + B*b*c^3 + 3*B*a*c^2*d - 3*A*b*c^2*d + 3*C*b*c^2*d - 3
*A*a*c*d^2 + 3*C*a*c*d^2 - 3*B*b*c*d^2 - B*a*d^3 + A*b*d^3 - C*b*d^3)*(f*x
+ e)/(a^2*c^6*f + b^2*c^6*f + 3*a^2*c^4*d^2*f + 3*b^2*c^4*d^2*f + 3*a^2*c
^2*d^4*f + 3*b^2*c^2*d^4*f + a^2*d^6*f + b^2*d^6*f) + 1/2*(B*a*c^3 - A*b*c
^3 + C*b*c^3 - 3*A*a*c^2*d + 3*C*a*c^2*d - 3*B*b*c^2*d - 3*B*a*c*d^2 + 3*A
*b*c*d^2 - 3*C*b*c*d^2 + A*a*d^3 - C*a*d^3 + B*b*d^3)*log(tan(f*x + e)^2 +
1)/(a^2*c^6*f + b^2*c^6*f + 3*a^2*c^4*d^2*f + 3*b^2*c^4*d^2*f + 3*a^2*c^2
*d^4*f + 3*b^2*c^2*d^4*f + a^2*d^6*f + b^2*d^6*f) + (C*a^2*b^3 - B*a*b^4 +
A*b^5)*log(abs(b*tan(f*x + e) + a))/(a^2*b^4*c^3*f + b^6*c^3*f - 3*a^3*b^
3*c^2*d*f - 3*a*b^5*c^2*d*f + 3*a^4*b^2*c*d^2*f + 3*a^2*b^4*c*d^2*f - a^5*
b*d^3*f - a^3*b^3*d^3*f) - (C*b^2*c^6*d - 3*B*b^2*c^5*d^2 + 3*B*a*b*c^4*d^
3 + 6*A*b^2*c^4*d^3 - 3*C*b^2*c^4*d^3 - B*a^2*c^3*d^4 - 8*A*a*b*c^3*d^4 +
8*C*a*b*c^3*d^4 + B*b^2*c^3*d^4 + 3*A*a^2*c^2*d^5 - 3*C*a^2*c^2*d^5 - 6*B*
a*b*c^2*d^5 + 3*A*b^2*c^2*d^5 + 3*B*a^2*c*d^6 - A*a^2*d^7 + C*a^2*d^7 - B*
a*b*d^7 + A*b^2*d^7)*log(abs(d*tan(f*x + e) + c))/(b^3*c^9*d*f - 3*a*b^2*c
^8*d^2*f + 3*a^2*b*c^7*d^3*f + 3*b^3*c^7*d^3*f - a^3*c^6*d^4*f - 9*a*b^2*c
^6*d^4*f + 9*a^2*b*c^5*d^5*f + 3*b^3*c^5*d^5*f - 3*a^3*c^4*d^6*f - 9*a*b^2
*c^4*d^6*f + 9*a^2*b*c^3*d^7*f + b^3*c^3*d^7*f - 3*a^3*c^2*d^8*f - 3*a*b^2
*c^2*d^8*f + 3*a^2*b*c*d^9*f - a^3*d^10*f) + 1/2*(3*C*b^2*c^8 - 4*C*a*b*c^
7*d - 5*B*b^2*c^7*d + C*a^2*c^6*d^2 + 8*B*a*b*c^6*d^2 + 7*A*b^2*c^6*d^2...
```

Mupad [B] (verification not implemented)

Time = 19.80 (sec) , antiderivative size = 65817, normalized size of antiderivative = 135.15

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))*(c + d*t
an(e + f*x))^3),x)
```

output

```
(symsum(log(- root(480*a^9*b*c^7*d^11*f^4 + 480*a*b^9*c^11*d^7*f^4 + 360*a^9*b*c^9*d^9*f^4 + 360*a^9*b*c^5*d^13*f^4 + 360*a*b^9*c^13*d^5*f^4 + 360*a*b^9*c^9*d^9*f^4 + 144*a^9*b*c^11*d^7*f^4 + 144*a^9*b*c^3*d^15*f^4 + 144*a*b^9*c^15*d^3*f^4 + 144*a*b^9*c^7*d^11*f^4 + 48*a^7*b^3*c*d^17*f^4 + 48*a^3*b^7*c^17*d*f^4 + 24*a^9*b*c^13*d^5*f^4 + 24*a^5*b^5*c^17*d*f^4 + 24*a^5*b^5*c*d^17*f^4 + 24*a*b^9*c^5*d^13*f^4 + 24*a^9*b*c*d^17*f^4 + 24*a*b^9*c^17*d*f^4 + 3920*a^5*b^5*c^9*d^9*f^4 - 3360*a^6*b^4*c^8*d^10*f^4 - 3360*a^4*b^6*c^10*d^8*f^4 - 3024*a^6*b^4*c^10*d^8*f^4 + 3024*a^5*b^5*c^11*d^7*f^4 + 3024*a^5*b^5*c^7*d^11*f^4 - 3024*a^4*b^6*c^8*d^10*f^4 + 2320*a^7*b^3*c^9*d^9*f^4 + 2320*a^3*b^7*c^9*d^9*f^4 - 2240*a^6*b^4*c^6*d^12*f^4 - 2240*a^4*b^6*c^12*d^6*f^4 + 2160*a^7*b^3*c^7*d^11*f^4 + 2160*a^3*b^7*c^11*d^7*f^4 - 1624*a^6*b^4*c^12*d^6*f^4 - 1624*a^4*b^6*c^6*d^12*f^4 + 1488*a^7*b^3*c^11*d^7*f^4 + 1488*a^3*b^7*c^7*d^11*f^4 + 1344*a^5*b^5*c^13*d^5*f^4 + 1344*a^5*b^5*c^5*d^13*f^4 - 1320*a^8*b^2*c^8*d^10*f^4 - 1320*a^2*b^8*c^10*d^8*f^4 + 1200*a^7*b^3*c^5*d^13*f^4 + 1200*a^3*b^7*c^13*d^5*f^4 - 1060*a^8*b^2*c^6*d^12*f^4 - 1060*a^2*b^8*c^12*d^6*f^4 - 948*a^8*b^2*c^10*d^8*f^4 - 948*a^2*b^8*c^8*d^10*f^4 - 840*a^6*b^4*c^4*d^14*f^4 - 840*a^4*b^6*c^14*d^4*f^4 + 528*a^7*b^3*c^13*d^5*f^4 + 528*a^3*b^7*c^5*d^13*f^4 - 480*a^8*b^2*c^4*d^14*f^4 - 480*a^6*b^4*c^14*d^4*f^4 - 480*a^4*b^6*c^4*d^14*f^4 - 480*a^2*b^8*c^14*d^4*f^4 - 368*a^8*b^2*c^12*d^6*f^4 + 368*a^7*b^3*c^3*d^15*f^4 + 3...
```

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 7603, normalized size of antiderivative = 15.61

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^3,x)
```


output

```
( - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**5*c**2*d**6 + log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)**2*a**5*d**8 + 9*log(tan(e + f*x)**2 + 1)*tan(
e + f*x)**2*a**4*b*c**3*d**5 - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*
a**4*b*c*d**7 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*c**3*d**6
- log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**4*c*d**8 - 9*log(tan(e + f*x
)**2 + 1)*tan(e + f*x)**2*a**3*b**2*c**4*d**4 + log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**2*a**3*b**2*d**8 - 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2
*a**3*b*c**4*d**5 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**3*c
**5*d**3 + 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**3*c**3*d**5
- 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**3*c*d**7 + 6*log(tan(
e + f*x)**2 + 1)*tan(e + f*x)**2*a**2*b**2*c**5*d**4 + 6*log(tan(e + f*x)*
**2 + 1)*tan(e + f*x)**2*a**2*b**2*c**3*d**6 - 9*log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**2*a*b**4*c**4*d**4 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*
**2*a*b**4*c**2*d**6 - 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*a*b**3*c
**4*d**5 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b**5*c**5*d**3 - log(
tan(e + f*x)**2 + 1)*tan(e + f*x)**2*b**5*c**3*d**5 - log(tan(e + f*x)**2
+ 1)*tan(e + f*x)**2*b**4*c**7*d**2 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f
*x)**2*b**4*c**5*d**4 - 6*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*c**3*
d**5 + 2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**5*c*d**7 + 18*log(tan(e
+ f*x)**2 + 1)*tan(e + f*x)*a**4*b*c**4*d**4 - 6*log(tan(e + f*x)**2 + ...
```

3.89
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^3} dx$$

Optimal result	997
Mathematica [B] (verified)	998
Rubi [A] (verified)	999
Maple [A] (verified)	1003
Fricas [B] (verification not implemented)	1005
Sympy [F(-2)]	1005
Maxima [B] (verification not implemented)	1005
Giac [B] (verification not implemented)	1006
Mupad [B] (verification not implemented)	1007
Reduce [B] (verification not implemented)	1008

Optimal result

Integrand size = 45, antiderivative size = 861

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2(c + d \tan(e + fx))^3} dx =$$

$$\frac{(b^2(Ac^3 - c^3C + 3Bc^2d - 3Acd^2 + 3cCd^2 - Bd^3) + a^2(c^3C - 3Bc^2d - 3cCd^2 + Bd^3 - A(c^3 - 3cd^2))}{(a^2 + b^2)^2 (c^2 + d^2)^3}$$

$$+ \frac{b^2(4a^3bBd - 3a^4Cd + b^4(Bc - 3Ad) + 2ab^3(Ac - cC + Bd) - a^2b^2(Bc + (5A + C)d)) \log(a \cos(e + fx))}{(a^2 + b^2)^2 (bc - ad)^4 f}$$

$$+ \frac{d(b^2(3c^6C - 6Bc^5d + c^4(10A - C)d^2 - 3Bc^3d^3 + 9Ac^2d^4 - Bcd^5 + 3Ad^6) + a^2d^3((A - C)d(3c^2 - d^2) + (bc - ad)(c^2 + d^2))}{(bc - ad)^2}$$

$$- \frac{d(b^2c(cC - Bd) - 2abB(c^2 + d^2) + a^2(3c^2C - Bcd + 2Cd^2) + A(a^2d^2 + b^2(2c^2 + 3d^2)))}{2(a^2 + b^2)(bc - ad)^2(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

$$- \frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^2}$$

$$- \frac{d(b^3c(2c^3C - 3Bc^2d - Bd^3) + a^2b(3c^4C - 3Bc^3d + 2c^2Cd^2 - Bcd^3 + Cd^4) + a^3d^2(2cCd + B(c^2 - d^2)))}{(a^2 + b^2)(bc - ad)^3(c^2 + d^2)}$$

output

```

-(b^2*(A*c^3-3*A*c*d^2+3*B*c^2*d-B*d^3-C*c^3+3*C*c*d^2)+a^2*(c^3*C-3*B*c^2
*d-3*C*c*d^2+B*d^3-A*(c^3-3*c*d^2)))+2*a*b*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*
d^2)))*x/(a^2+b^2)^2/(c^2+d^2)^3+b^2*(4*a^3*b*B*d-3*a^4*C*d+b^4*(-3*A*d+B*
c)+2*a*b^3*(A*c+B*d-C*c)-a^2*b^2*(B*c+(5*A+C)*d))*ln(a*cos(f*x+e)+b*sin(f*
x+e))/(a^2+b^2)^2/(-a*d+b*c)^4/f+d*(b^2*(3*c^6*C-6*B*c^5*d+c^4*(10*A-C)*d^
2-3*B*c^3*d^3+9*A*c^2*d^4-B*c*d^5+3*A*d^6)+a^2*d^3*((A-C)*d*(3*c^2-d^2)-B*
(c^3-3*c*d^2))-2*a*b*d^2*(c*(A-C)*d*(5*c^2+d^2)-B*(2*c^4-3*c^2*d^2-d^4)))*
ln(c*cos(f*x+e)+d*sin(f*x+e))/(-a*d+b*c)^4/(c^2+d^2)^3/f-1/2*d*(b^2*c*(-B*
d+C*c)-2*a*b*B*(c^2+d^2)+a^2*(-B*c*d+3*C*c^2+2*C*d^2)+A*(a^2*d^2+b^2*(2*c^
2+3*d^2)))/(a^2+b^2)/(-a*d+b*c)^2/(c^2+d^2)/f/(c+d*tan(f*x+e))^2-(A*b^2-a*
(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^2-d*(b
^3*c*(-3*B*c^2*d-B*d^3+2*C*c^3)+a^2*b*(-3*B*c^3*d-B*c*d^3+3*C*c^4+2*C*c^2*
d^2+C*d^4)+a^3*d^2*(2*C*c*d+B*(c^2-d^2))+a*b^2*(2*c*C*d^3-B*(c^4+c^2*d^2+2
*d^4))-A*(2*a^3*c*d^3+2*a*b^2*c*d^3-2*a^2*b*d^2*(2*c^2+d^2)-b^3*(c^4+6*c^2
*d^2+3*d^4)))/(a^2+b^2)/(-a*d+b*c)^3/(c^2+d^2)^2/f/(c+d*tan(f*x+e))

```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1732 vs. $2(861) = 1722$.

Time = 7.60 (sec) , antiderivative size = 1732, normalized size of antiderivative = 2.01

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*
(c + d*Tan[e + f*x])^3),x]

```

output

```

-((A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])*
(c + d*Tan[e + f*x])^2)) - (-1/2*(-(c*(-3*c*(A*b^2 - a*(b*B - a*C))*d + (A
*b - a*B - b*C)*d*(b*c - a*d))) + d^2*(3*A*b^2*d - a*A*(b*c - a*d) - (b*B
- a*C)*(b*c + 2*a*d)))/((-b*c) + a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^
2) - (((-(((b*c - a*d)^3*(-b^2*(2*a*A*b*c^3 - a^2*B*c^3 + b^2*B*c^3 - 2*
a*b*c^3*C + 3*a^2*A*c^2*d - 3*A*b^2*c^2*d + 6*a*b*B*c^2*d - 3*a^2*c^2*C*d
+ 3*b^2*c^2*C*d - 6*a*A*b*c*d^2 + 3*a^2*B*c*d^2 - 3*b^2*B*c*d^2 + 6*a*b*c*
C*d^2 - a^2*A*d^3 + A*b^2*d^3 - 2*a*b*B*d^3 + a^2*C*d^3 - b^2*C*d^3)) + Sq
rt[-b^2]*(-(a^2*A*b*c^3) + A*b^3*c^3 - 2*a*b^2*B*c^3 + a^2*b*c^3*C - b^3*c
^3*C + 6*a*A*b^2*c^2*d - 3*a^2*b*B*c^2*d + 3*b^3*B*c^2*d - 6*a*b^2*c^2*C*d
+ 3*a^2*A*b*c*d^2 - 3*A*b^3*c*d^2 + 6*a*b^2*B*c*d^2 - 3*a^2*b*c*C*d^2 + 3
*b^3*c*C*d^2 - 2*a*A*b^2*d^3 + a^2*b*B*d^3 - b^3*B*d^3 + 2*a*b^2*C*d^3))*L
og[Sqrt[-b^2] - b*Tan[e + f*x]]/(b*(a^2 + b^2)*(c^2 + d^2))) - (2*b^3*(c^
2 + d^2)^2*(4*a^3*b*B*d - 3*a^4*C*d + b^4*(B*c - 3*A*d) + 2*a*b^3*(A*c -
c*C + B*d) - a^2*b^2*(B*c + (5*A + C)*d))*Log[a + b*Tan[e + f*x]]/((a^2 +
b^2)*(b*c - a*d)) + ((b*c - a*d)^3*(b^2*(2*a*A*b*c^3 - a^2*B*c^3 + b^2*B*c
^3 - 2*a*b*c^3*C + 3*a^2*A*c^2*d - 3*A*b^2*c^2*d + 6*a*b*B*c^2*d - 3*a^2*c
^2*C*d + 3*b^2*c^2*C*d - 6*a*A*b*c*d^2 + 3*a^2*B*c*d^2 - 3*b^2*B*c*d^2 + 6
*a*b*c*C*d^2 - a^2*A*d^3 + A*b^2*d^3 - 2*a*b*B*d^3 + a^2*C*d^3 - b^2*C*d^3
) + Sqrt[-b^2]*(-(a^2*A*b*c^3) + A*b^3*c^3 - 2*a*b^2*B*c^3 + a^2*b*c^3*...

```

Rubi [A] (verified)

Time = 8.77 (sec) , antiderivative size = 932, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.244$, Rules used = {3042, 4132, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx$$

↓ 4132

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan^2(e+fx)-aA(bc-ad)-(bB-aC)(bc+2ad)+(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^3} dx$$

$$\frac{(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))^2$$

↓ 3042

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan(e+fx)^2-aA(bc-ad)-(bB-aC)(bc+2ad)+(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^3} dx$$

$$\frac{(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))^2$$

↓ 4132

$$\int -\frac{2(d^2(Ac-Cc+Bd)a^3-b(2A+C)d(c^2+d^2)a^2+b^2(Ac-Cc+Bd)(c^2+2d^2)a-bd(Ad^2a^2+(3Cc^2-Bdc+2Cd^2)a^2-2bB(c^2+d^2)a+b^2c(cC-Bd)+Ab^2(2c^2+3d^2)+b^2c(cC-Bd))}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$$

$$\frac{2(c^2+d^2)(bc-ad)}{2(c^2+d^2)(bc-ad)}$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))^2}$$

↓ 27

$$\frac{d(a^2Ad^2+a^2(-Bcd+3c^2C+2Cd^2)-2abB(c^2+d^2)+Ab^2(2c^2+3d^2)+b^2c(cC-Bd))}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2} - \int \frac{d^2(Ac-Cc+Bd)a^3-b(2A+C)d(c^2+d^2)a^2+b^2(Ac-Cc+Bd)(c^2+2d^2)a-bd(Ad^2a^2+(3Cc^2-Bdc+2Cd^2)a^2-2bB(c^2+d^2)a+b^2c(cC-Bd)+Ab^2(2c^2+3d^2)+b^2c(cC-Bd))}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))^2}$$

↓ 3042

$$\frac{d(a^2Ad^2+a^2(-Bcd+3c^2C+2Cd^2)-2abB(c^2+d^2)+Ab^2(2c^2+3d^2)+b^2c(cC-Bd))}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2} - \int \frac{d^2(Ac-Cc+Bd)a^3-b(2A+C)d(c^2+d^2)a^2+b^2(Ac-Cc+Bd)(c^2+2d^2)a-bd(Ad^2a^2+(3Cc^2-Bdc+2Cd^2)a^2-2bB(c^2+d^2)a+b^2c(cC-Bd)+Ab^2(2c^2+3d^2)+b^2c(cC-Bd))}{(a+b \tan(e+fx))(c+d \tan(e+fx))^2} dx$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))(c+d \tan(e+fx))^2}$$

↓ 4132

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^2} - \frac{d^3(Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2))a^4 - bd^2(3Cc^3 - 4Bdc^2 + Cd^3)}{f}$$

$$\frac{d(Ad^2a^2 + (3Cc^2 - Bdc + 2Cd^2)a^2 - 2bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(2c^2 + 3d^2))}{2(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^2} - \frac{d^3(Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2))a^4 - bd^2(3Cc^3 - 4Bdc^2 + Cd^3)}{f}$$

$$\frac{d(Ad^2a^2 + (3Cc^2 - Bdc + 2Cd^2)a^2 - 2bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(2c^2 + 3d^2))}{2(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 4134

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^2} - \frac{((Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3)a^2 - 2b((A - C)d(3c^2 + 3d^2) - b^2))}{f}$$

$$\frac{d(Ad^2a^2 + (3Cc^2 - Bdc + 2Cd^2)a^2 - 2bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(2c^2 + 3d^2))}{2(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^2} - \frac{((Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3)a^2 - 2b((A - C)d(3c^2 + 3d^2) - b^2))}{f}$$

$$\frac{d(Ad^2a^2 + (3Cc^2 - Bdc + 2Cd^2)a^2 - 2bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(2c^2 + 3d^2))}{2(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

↓ 4013

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^2} - \frac{((Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3)a^2 - 2b((A - C)d(3c^2 + 3d^2) - b^2))}{f}$$

$$\frac{d(Ad^2a^2 + (3Cc^2 - Bdc + 2Cd^2)a^2 - 2bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(2c^2 + 3d^2))}{2(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^2}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2)/((a + b*\text{Tan}[e + f*x])^2*(c + d*\text{Tan}[e + f*x])^3), x]$

output
$$-\frac{((A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*\text{Tan}[e + f*x])*(c + d*\text{Tan}[e + f*x]^2)) - ((d*(a^2*A*d^2 + b^2*c*(c*C - B*d) - 2*a*b*B*(c^2 + d^2) + A*b^2*(2*c^2 + 3*d^2) + a^2*(3*c^2*C - B*c*d + 2*C*d^2)))/(2*(b*c - a*d)*(c^2 + d^2)*f*(c + d*\text{Tan}[e + f*x]^2) - (((b*c - a*d)^3*(a^2*(A*c^3 - c^3*C + 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 - B*d^3) + b^2*(c^3*C - 3*B*c^2*d - 3*c*C*d^2 + B*d^3 - A*(c^3 - 3*c*d^2)) - 2*a*b*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2))))*x)/((a^2 + b^2)*(c^2 + d^2)) + (b^2*(c^2 + d^2)^2*(4*a^3*b*B*d - 3*a^4*C*d + b^4*(B*c - 3*A*d) + 2*a*b^3*(A*c - c*C + B*d) - a^2*b^2*(B*c + (5*A + C)*d))*\text{Log}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f) + ((a^2 + b^2)*d*(b^2*(3*c^6*C - 6*B*c^5*d + c^4*(10*A - C)*d^2 - 3*B*c^3*d^3 + 9*A*c^2*d^4 - B*c*d^5 + 3*A*d^6) + a^2*d^3*((A - C)*d*(3*c^2 - d^2) - B*(c^3 - 3*c*d^2)) - 2*a*b*d^2*(c*(A - C)*d*(5*c^2 + d^2) - B*(2*c^4 - 3*c^2*d^2 - d^4)))*\text{Log}[c*\text{Cos}[e + f*x] + d*\text{Sin}[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f))/((b*c - a*d)*(c^2 + d^2)) - (d*(b^3*c*(2*c^3*C - 3*B*c^2*d - B*d^3) + a^2*b*(3*c^4*C - 3*B*c^3*d + 2*c^2*C*d^2 - B*c*d^3 + C*d^4) + a^3*d^2*(2*c*C*d + B*(c^2 - d^2)) + a*b^2*(2*c*C*d^3 - B*(c^4 + c^2*d^2 + 2*d^4)) - A*(2*a^3*c*d^3 + 2*a*b^2*c*d^3 - 2*a^2*b*d^2*(2*c^2 + d^2) - b^3*(c^4 + 6*c^2*d^2 + 3*d^4)))/((b*c - a*d)*(c^2 + d^2)*f*(c + d*\text{Tan}[e + f*x]))/((b*c - a*d)*(c^2 + d^2))/((a^2 + b^2)*(b*c - a*d))$$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)*(G_x)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4013

```
Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4132

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4134

```
Int[((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^
2)/(((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [A] (verified)

Time = 4.85 (sec) , antiderivative size = 949, normalized size of antiderivative = 1.10

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9567 vs. $2(862) = 1724$.

Time = 11.06 (sec) , antiderivative size = 9567, normalized size of antiderivative = 11.11

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^3,x, algorithm="fricas")
```

output

Too large to include

Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx$$

= Exception raised: NotImplementedError

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**2/(c+d*tan(f*x+e))**3,x)
```

output

Exception raised: NotImplementedError >> no valid subset found

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2537 vs. $2(862) = 1724$.

Time = 0.24 (sec) , antiderivative size = 2537, normalized size of antiderivative = 2.95

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^3,x, algorithm="maxima")`

output

$$\begin{aligned} & 1/2*(2*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c^3 + 3*(B*a^2 - 2*(A - C)*a \\ & *b - B*b^2)*c^2*d - 3*((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*c*d^2 - (B*a^2 \\ & - 2*(A - C)*a*b - B*b^2)*d^3)*(f*x + e)/((a^4 + 2*a^2*b^2 + b^4)*c^6 + 3* \\ & (a^4 + 2*a^2*b^2 + b^4)*c^4*d^2 + 3*(a^4 + 2*a^2*b^2 + b^4)*c^2*d^4 + (a^4 \\ & + 2*a^2*b^2 + b^4)*d^6) - 2*((B*a^2*b^4 - 2*(A - C)*a*b^5 - B*b^6)*c + (3 \\ & *C*a^4*b^2 - 4*B*a^3*b^3 + (5*A + C)*a^2*b^4 - 2*B*a*b^5 + 3*A*b^6)*d)*\log \\ & (b*\tan(f*x + e) + a)/((a^4*b^4 + 2*a^2*b^6 + b^8)*c^4 - 4*(a^5*b^3 + 2*a^3 \\ & *b^5 + a*b^7)*c^3*d + 6*(a^6*b^2 + 2*a^4*b^4 + a^2*b^6)*c^2*d^2 - 4*(a^7*b \\ & + 2*a^5*b^3 + a^3*b^5)*c*d^3 + (a^8 + 2*a^6*b^2 + a^4*b^4)*d^4) + 2*(3*C* \\ & b^2*c^6*d - 6*B*b^2*c^5*d^2 + (4*B*a*b + (10*A - C)*b^2)*c^4*d^3 - (B*a^2 \\ & + 10*(A - C)*a*b + 3*B*b^2)*c^3*d^4 + 3*((A - C)*a^2 - 2*B*a*b + 3*A*b^2)* \\ & c^2*d^5 + (3*B*a^2 - 2*(A - C)*a*b - B*b^2)*c*d^6 - ((A - C)*a^2 + 2*B*a*b \\ & - 3*A*b^2)*d^7)*\log(d*\tan(f*x + e) + c)/(b^4*c^10 - 4*a*b^3*c^9*d - 4*a^3 \\ & *b*c*d^9 + a^4*d^10 + 3*(2*a^2*b^2 + b^4)*c^8*d^2 - 4*(a^3*b + 3*a*b^3)*c^ \\ & 7*d^3 + (a^4 + 18*a^2*b^2 + 3*b^4)*c^6*d^4 - 12*(a^3*b + a*b^3)*c^5*d^5 + \\ & (3*a^4 + 18*a^2*b^2 + b^4)*c^4*d^6 - 4*(3*a^3*b + a*b^3)*c^3*d^7 + 3*(a^4 \\ & + 2*a^2*b^2)*c^2*d^8) + ((B*a^2 - 2*(A - C)*a*b - B*b^2)*c^3 - 3*((A - C)* \\ & a^2 + 2*B*a*b - (A - C)*b^2)*c^2*d - 3*(B*a^2 - 2*(A - C)*a*b - B*b^2)*c*d \\ & ^2 + ((A - C)*a^2 + 2*B*a*b - (A - C)*b^2)*d^3)*\log(\tan(f*x + e)^2 + 1)/((\\ & a^4 + 2*a^2*b^2 + b^4)*c^6 + 3*(a^4 + 2*a^2*b^2 + b^4)*c^4*d^2 + 3*(a^4\dots \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4552 vs. $2(862) = 1724$.

Time = 1.24 (sec) , antiderivative size = 4552, normalized size of antiderivative = 5.29

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^3,x, algorithm="giac")`

output

```
(A*a^2*c^3 - C*a^2*c^3 + 2*B*a*b*c^3 - A*b^2*c^3 + C*b^2*c^3 + 3*B*a^2*c^2
*d - 6*A*a*b*c^2*d + 6*C*a*b*c^2*d - 3*B*b^2*c^2*d - 3*A*a^2*c*d^2 + 3*C*a
^2*c*d^2 - 6*B*a*b*c*d^2 + 3*A*b^2*c*d^2 - 3*C*b^2*c*d^2 - B*a^2*d^3 + 2*A
*a*b*d^3 - 2*C*a*b*d^3 + B*b^2*d^3)*(f*x + e)/(a^4*c^6*f + 2*a^2*b^2*c^6*f
+ b^4*c^6*f + 3*a^4*c^4*d^2*f + 6*a^2*b^2*c^4*d^2*f + 3*b^4*c^4*d^2*f + 3
*a^4*c^2*d^4*f + 6*a^2*b^2*c^2*d^4*f + 3*b^4*c^2*d^4*f + a^4*d^6*f + 2*a^2
*b^2*d^6*f + b^4*d^6*f) + 1/2*(B*a^2*c^3 - 2*A*a*b*c^3 + 2*C*a*b*c^3 - B*b
^2*c^3 - 3*A*a^2*c^2*d + 3*C*a^2*c^2*d - 6*B*a*b*c^2*d + 3*A*b^2*c^2*d - 3
*C*b^2*c^2*d - 3*B*a^2*c*d^2 + 6*A*a*b*c*d^2 - 6*C*a*b*c*d^2 + 3*B*b^2*c*d
^2 + A*a^2*d^3 - C*a^2*d^3 + 2*B*a*b*d^3 - A*b^2*d^3 + C*b^2*d^3)*log(tan(
f*x + e)^2 + 1)/(a^4*c^6*f + 2*a^2*b^2*c^6*f + b^4*c^6*f + 3*a^4*c^4*d^2*f
+ 6*a^2*b^2*c^4*d^2*f + 3*b^4*c^4*d^2*f + 3*a^4*c^2*d^4*f + 6*a^2*b^2*c^2
*d^4*f + 3*b^4*c^2*d^4*f + a^4*d^6*f + 2*a^2*b^2*d^6*f + b^4*d^6*f) - (B*a
^2*b^5*c - 2*A*a*b^6*c + 2*C*a*b^6*c - B*b^7*c + 3*C*a^4*b^3*d - 4*B*a^3*b
^4*d + 5*A*a^2*b^5*d + C*a^2*b^5*d - 2*B*a*b^6*d + 3*A*b^7*d)*log(abs(b*ta
n(f*x + e) + a))/(a^4*b^5*c^4*f + 2*a^2*b^7*c^4*f + b^9*c^4*f - 4*a^5*b^4*c
^3*d*f - 8*a^3*b^6*c^3*d*f - 4*a*b^8*c^3*d*f + 6*a^6*b^3*c^2*d^2*f + 12*a
^4*b^5*c^2*d^2*f + 6*a^2*b^7*c^2*d^2*f - 4*a^7*b^2*c*d^3*f - 8*a^5*b^4*c*d
^3*f - 4*a^3*b^6*c*d^3*f + a^8*b*d^4*f + 2*a^6*b^3*d^4*f + a^4*b^5*d^4*f)
+ (3*C*b^2*c^6*d^2 - 6*B*b^2*c^5*d^3 + 4*B*a*b*c^4*d^4 + 10*A*b^2*c^4*d...
```

Mupad [B] (verification not implemented)

Time = 39.88 (sec) , antiderivative size = 128666, normalized size of antiderivative = 149.44

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^2*(c + d
*tan(e + f*x))^3),x)
```

output

```
((2*A*b^4*c^6 - A*a^4*d^6 - 2*B*a*b^3*c^6 - B*a^4*c*d^5 - A*a^2*b^2*d^6 -
5*A*a^4*c^2*d^4 + 2*C*a^2*b^2*c^6 + 2*A*b^4*c^2*d^4 + 4*A*b^4*c^4*d^2 + 3
*B*a^4*c^3*d^3 + 3*C*a^4*c^2*d^4 - C*a^4*c^4*d^2 + 9*A*a*b^3*c^3*d^3 + 9*A
*a^3*b*c^3*d^3 - 5*B*a*b^3*c^2*d^4 - 11*B*a*b^3*c^4*d^2 - B*a^2*b^2*c*d^5
- 3*B*a^3*b*c^2*d^4 - 7*B*a^3*b*c^4*d^2 + C*a*b^3*c^3*d^3 + C*a^3*b*c^3*d^
3 - 5*A*a^2*b^2*c^2*d^4 + 3*B*a^2*b^2*c^3*d^3 + 5*C*a^2*b^2*c^2*d^4 + 3*C*
a^2*b^2*c^4*d^2 + 5*A*a*b^3*c*d^5 + 5*A*a^3*b*c*d^5 + 5*C*a*b^3*c^5*d + 5*
C*a^3*b*c^5*d)/(2*(a^3*d^3 - b^3*c^3 + 3*a*b^2*c^2*d - 3*a^2*b*c*d^2)*(a^2
*c^4 + a^2*d^4 + b^2*c^4 + b^2*d^4 + 2*a^2*c^2*d^2 + 2*b^2*c^2*d^2)) + (ta
n(e + f*x)*(3*A*a*b^3*d^6 - 2*B*a^4*d^6 + 3*A*a^3*b*d^6 - 4*A*a^4*c*d^5 +
9*A*b^4*c*d^5 + 4*A*b^4*c^5*d + 4*C*a^4*c*d^5 + 5*C*b^4*c^5*d - 2*B*a^2*b^
2*d^6 + 17*A*b^4*c^3*d^3 + 2*B*a^4*c^2*d^4 - 3*B*b^4*c^2*d^4 - 7*B*b^4*c^4
*d^2 + C*b^4*c^3*d^3 + 3*A*a*b^3*c^2*d^4 + A*a^2*b^2*c*d^5 + 3*A*a^3*b*c^2
*d^4 - 11*B*a*b^3*c^3*d^3 - 3*B*a^3*b*c^3*d^3 + 3*C*a*b^3*c^2*d^4 + 3*C*a*
b^3*c^4*d^2 + 8*C*a^2*b^2*c*d^5 + 9*C*a^2*b^2*c^5*d + 3*C*a^3*b*c^2*d^4 +
3*C*a^3*b*c^4*d^2 + 9*A*a^2*b^2*c^3*d^3 - B*a^2*b^2*c^2*d^4 - 7*B*a^2*b^2*
c^4*d^2 + 9*C*a^2*b^2*c^3*d^3 - 7*B*a*b^3*c*d^5 - 4*B*a*b^3*c^5*d - 3*B*a^
3*b*c*d^5))/(2*(a^3*d^3 - b^3*c^3 + 3*a*b^2*c^2*d - 3*a^2*b*c*d^2)*(a^2*c^
4 + a^2*d^4 + b^2*c^4 + b^2*d^4 + 2*a^2*c^2*d^2 + 2*b^2*c^2*d^2)) + (tan(e
+ f*x)^2*(3*A*b^4*d^6 - 2*B*a*b^3*d^6 - B*a^3*b*d^6 - B*b^4*c*d^5 + 2*...
```

Reduce [B] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 28024, normalized size of antiderivative = 32.55

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^3,
x)
```

output

```
( - 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**8*b*c**2*d**8 + log(tan(
e + f*x)**2 + 1)*tan(e + f*x)**3*a**8*b*d**10 + 5*log(tan(e + f*x)**2 + 1)
*tan(e + f*x)**3*a**7*b**2*c**3*d**7 + log(tan(e + f*x)**2 + 1)*tan(e + f*
x)**3*a**7*b**2*c*d**9 + 3*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**7*b
*c**3*d**8 - log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**7*b*c*d**10 + 8*l
og(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**3*c**4*d**6 - 11*log(tan(e
 + f*x)**2 + 1)*tan(e + f*x)**3*a**6*b**3*c**2*d**8 + log(tan(e + f*x)**2
 + 1)*tan(e + f*x)**3*a**6*b**3*d**10 - 4*log(tan(e + f*x)**2 + 1)*tan(e +
f*x)**3*a**6*b**2*c**4*d**7 - 4*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a
**6*b**2*c**2*d**9 - 22*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**4
*c**5*d**5 + 7*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**4*c**3*d**
7 + log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**4*c*d**9 - 10*log(tan
(e + f*x)**2 + 1)*tan(e + f*x)**3*a**5*b**3*c**5*d**6 + 11*log(tan(e + f*x)
)**2 + 1)*tan(e + f*x)**3*a**5*b**3*c**3*d**8 + log(tan(e + f*x)**2 + 1)*t
an(e + f*x)**3*a**5*b**3*c*d**10 + 13*log(tan(e + f*x)**2 + 1)*tan(e + f*x)
)**3*a**4*b**5*c**6*d**4 + 25*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**
4*b**5*c**4*d**6 - 8*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**5*c*
**2*d**8 + 20*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**4*c**6*d**5
 + 10*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**4*c**4*d**7 - 2*log(
tan(e + f*x)**2 + 1)*tan(e + f*x)**3*a**4*b**4*c**2*d**9 + log(tan(e + ...
```

3.90 $\int (a+b \tan(e+fx))^3 \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)) dx$

Optimal result	1010
Mathematica [B] (verified)	1011
Rubi [A] (warning: unable to verify)	1012
Maple [B] (verified)	1018
Fricas [B] (verification not implemented)	1019
Sympy [F]	1020
Maxima [F(-1)]	1020
Giac [F(-2)]	1021
Mupad [F(-1)]	1021
Reduce [F]	1022

Optimal result

Integrand size = 47, antiderivative size = 464

$$\begin{aligned}
 & \int (a+b \tan(e+fx))^3 \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx) + C \tan^2(e+fx)) dx \\
 = & -\frac{(a-ib)^3(iA+B-iC)\sqrt{c-id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f} \\
 & + \frac{(a+ib)^3(iA-B-iC)\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f} \\
 & + \frac{2(a^3B-3ab^2B+3a^2b(A-C)-b^3(A-C))\sqrt{c+d \tan(e+fx)}}{f} \\
 & + \frac{2(40a^3Cd^3-6a^2bd^2(16cC-45Bd)+9ab^2d(8c^2C-14Bcd+35(A-C)d^2)-b^3(16c^3C-24Bc^2d+315d^4f)}{315d^4f} \\
 & + \frac{2b(21b(Ab+aB-bC)d^2+4(bc-ad)(2bcC-3bBd-2aCd))\tan(e+fx)(c+d \tan(e+fx))^{3/2}}{105d^3f} \\
 & - \frac{2(2bcC-3bBd-2aCd)(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{3/2}}{21d^2f} \\
 & + \frac{2C(a+b \tan(e+fx))^3(c+d \tan(e+fx))^{3/2}}{9df}
 \end{aligned}$$

output

```

-(a-I*b)^3*(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)
)^(1/2))/f+(a+I*b)^3*(I*A-B-I*C)*(c+I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1
/2)/(c+I*d)^(1/2))/f+2*(B*a^3-3*B*a*b^2+3*a^2*b*(A-C)-b^3*(A-C))*(c+d*tan(
f*x+e))^(1/2)/f+2/315*(40*a^3*C*d^3-6*a^2*b*d^2*(-45*B*d+16*C*c)+9*a*b^2*d
*(8*c^2*C-14*B*c*d+35*(A-C)*d^2)-b^3*(16*c^3*C-24*B*c^2*d+42*c*(A-C)*d^2+1
05*B*d^3))*(c+d*tan(f*x+e))^(3/2)/d^4/f+2/105*b*(21*b*(A*b+B*a-C*b)*d^2+4*
(-a*d+b*c)*(-3*B*b*d-2*C*a*d+2*C*b*c))*tan(f*x+e)*(c+d*tan(f*x+e))^(3/2)/d
^3/f-2/21*(-3*B*b*d-2*C*a*d+2*C*b*c)*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(
3/2)/d^2/f+2/9*C*(a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(3/2)/d/f

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1232 vs. $2(464) = 928$.

Time = 6.32 (sec) , antiderivative size = 1232, normalized size of antiderivative = 2.66

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```

Integrate[(a + b*Tan[e + f*x])^3*sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f
*x] + C*Tan[e + f*x]^2),x]

```


output

```
(2*C*(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^(3/2))/(9*d*f) + (2*((-3*
(2*b*c*C - 3*b*B*d - 2*a*C*d)*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^
(3/2))/(7*d*f) + (2*((3*b*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b
*c*C - 3*b*B*d - 2*a*C*d))*Tan[e + f*x]*(c + d*Tan[e + f*x])^(3/2))/(10*d*
f) - (2*((2*((-15*a*d*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*C
- 3*b*B*d - 2*a*C*d)))/8 + b*((-315*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3)/
8 + (3*c*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*C - 3*b*B*d -
2*a*C*d)))/4))*(c + d*Tan[e + f*x])^(3/2))/(3*d*f) + ((I/2)*((-15*a*d*(a^2
*(21*A - 13*C)*d^2 + 4*b^2*c*(2*c*C - 3*B*d) - a*b*d*(16*c*C + 9*B*d)))/8
+ (3*b*c*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*C - 3*b*B*d -
2*a*C*d)))/4 + (15*a*d*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*
C - 3*b*B*d - 2*a*C*d)))/8 + ((5*I)/2)*d*((63*a*(a^2*B - b^2*B + 2*a*b*(A
- C))*d^2)/4 + (3*b*(a^2*(21*A - 13*C)*d^2 + 4*b^2*c*(2*c*C - 3*B*d) - a*b
*d*(16*c*C + 9*B*d)))/4 - (3*b*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)
*(2*b*c*C - 3*b*B*d - 2*a*C*d)))/4) - b*((-315*(a^2*B - b^2*B + 2*a*b*(A -
C))*d^3)/8 + (3*c*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*C -
3*b*B*d - 2*a*C*d)))/4))*((2*(c - I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*
x]]/Sqrt[c - I*d]])/(-c + I*d) + 2*Sqrt[c + d*Tan[e + f*x]]))/f - ((I/2)*
(-15*a*d*(a^2*(21*A - 13*C)*d^2 + 4*b^2*c*(2*c*C - 3*B*d) - a*b*d*(16*c*C
+ 9*B*d)))/8 + (3*b*c*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*...
```

Rubi [A] (warning: unable to verify)

Time = 5.82 (sec) , antiderivative size = 476, normalized size of antiderivative = 1.03, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.426$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \frac{2 \int -\frac{3}{2}(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} ((2bcC - 2adC - 3bBd) \tan^2(e + fx) - 3(Ab - Cb + aB)d \tan(e + fx))}{9df} \frac{9d}{27} \\
 & \quad \downarrow 27 \\
 & \frac{2C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2}}{9df} - \\
 & \frac{\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} ((2bcC - 2adC - 3bBd) \tan^2(e + fx) - 3(Ab - Cb + aB)d \tan(e + fx))}{3d} \\
 & \quad \downarrow 3042 \\
 & \frac{2C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2}}{9df} - \\
 & \frac{\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} ((2bcC - 2adC - 3bBd) \tan(e + fx)^2 - 3(Ab - Cb + aB)d \tan(e + fx))}{3d} \\
 & \quad \downarrow 4130 \\
 & \frac{2C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2}}{9df} - \\
 & \frac{2 \int -\frac{1}{2}(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (4c(2cC - 3Bd)b^2 - ad(16cC + 9Bd)b + a^2(21A - 13C)d^2 + (21b(Ab - Cb + aB)d^2 + 4(bc - ad)(2bcC - 2adC))}{7d}}{3d} \\
 & \quad \downarrow 27 \\
 & \frac{2C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2}}{9df} - \\
 & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (4c(2cC - 3Bd)b^2 - ad(16cC + 9Bd)b + a^2)}{7d}}{3d} \\
 & \quad \downarrow 3042 \\
 & \frac{2C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2}}{9df} - \\
 & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (4c(2cC - 3Bd)b^2 - ad(16cC + 9Bd)b + a^2)}{7d}}{3d} \\
 & \quad \downarrow 4120 \\
 & \frac{2C(a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2}}{9df} - \\
 & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}}{7df} - \frac{\frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2} (21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}}{5df}}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} - \\ & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c + d \tan(e + fx)} (-2c(8Cc^2 - 12Bdc + 21(A - C)d^2)b^3 + 18acd(4cC - 7Bd)b^2 - 3a^2d^2)}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} - \\ & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c + d \tan(e + fx)} (-2c(8Cc^2 - 12Bdc + 21(A - C)d^2)b^3 + 18acd(4cC - 7Bd)b^2 - 3a^2d^2)}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 4113 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} - \\ & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c + d \tan(e + fx)} (105(Ba^3 + 3b(A - C)a^2 - 3b^2Ba - b^3(A - C))d^3 \tan(e + fx) - 105(-}}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} - \\ & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c + d \tan(e + fx)} (105(Ba^3 + 3b(A - C)a^2 - 3b^2Ba - b^3(A - C))d^3 \tan(e + fx) - 105(-}}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} - \\ & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int 105((Ac - Cc - Bd)a^3 - 3b(Bc + (A - C)d)a^2 - 3b^2(Ac - Cc - Bd)a + b^3(Bc + (A - C)d))}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} - \\ & \frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int 105((Ac - Cc - Bd)a^3 - 3b(Bc + (A - C)d)a^2 - 3b^2(Ac - Cc - Bd)a + b^3(Bc + (A - C)d))}{7df} \end{aligned}$$

↓ 4022

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} -$$

$$\frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2}(21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} -$$

$$\frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2}(21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}$$

↓ 4020

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} -$$

$$\frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2}(21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}$$

↓ 25

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} -$$

$$\frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2}(21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}$$

↓ 73

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df} -$$

$$\frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2}(21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}$$

↓ 221

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{3/2}}{9df}$$

$$\frac{2(-2aCd - 3bBd + 2bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{3/2}(21bd^2(aB + Ab - bC) + 4(bc - ad)(-2aCd - 3bBd + 2bcC))}{5df}$$

input `Int[(a + b*Tan[e + f*x])^3*sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `(2*C*(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^(3/2))/(9*d*f) - ((2*(2*b*c*C - 3*b*B*d - 2*a*C*d)*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(3/2))/(7*d*f) - ((2*b*(21*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*C - 3*b*B*d - 2*a*C*d))*Tan[e + f*x]*(c + d*Tan[e + f*x])^(3/2))/(5*d*f) + ((105*(a - I*b)^3*(A - I*B - C)*sqrt[c - I*d]*d^3*ArcTan[Tan[e + f*x]/sqrt[c - I*d]])/f + (105*(a + I*b)^3*(A + I*B - C)*sqrt[c + I*d]*d^3*ArcTan[Tan[e + f*x]/sqrt[c + I*d]])/f + (210*(a^3*B - 3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C))*d^3*sqrt[c + d*Tan[e + f*x]])/f + (2*(40*a^3*C*d^3 - 6*a^2*b*d^2*(16*c*C - 45*B*d) + 9*a*b^2*d*(8*c^2*C - 14*B*c*d + 35*(A - C)*d^2) - b^3*(16*c^3*C - 24*B*c^2*d + 42*c*(A - C)*d^2 + 105*B*d^3))*(c + d*Tan[e + f*x])^(3/2))/(3*d*f)/(5*d)/(7*d)/(3*d)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 $\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4011 $\text{Int}[(a + (b \cdot \tan[e + f \cdot x]) + (c + (d \cdot \tan[e + f \cdot x]) + (f \cdot x))^m], x_Symbol] \rightarrow \text{Simp}[d \cdot (a + b \cdot \tan[e + f \cdot x])^m / (f \cdot m), x] + \text{Int}[(a + b \cdot \tan[e + f \cdot x])^{m-1} \cdot \text{Simp}[a \cdot c - b \cdot d + (b \cdot c + a \cdot d) \cdot \tan[e + f \cdot x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

rule 4020 $\text{Int}[(a + (b \cdot \tan[e + f \cdot x]) + (c + (d \cdot \tan[e + f \cdot x]) + (f \cdot x))^m], x_Symbol] \rightarrow \text{Simp}[c \cdot (d/f) \ \text{Subst}[\text{Int}[(a + (b/d) \cdot x)^m / (d^2 + c \cdot x), x], x, d \cdot \tan[e + f \cdot x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[(a + (b \cdot \tan[e + f \cdot x]) + (c + (d \cdot \tan[e + f \cdot x]) + (f \cdot x))^m], x_Symbol] \rightarrow \text{Simp}[(c + I \cdot d)/2 \ \text{Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (1 - I \cdot \tan[e + f \cdot x]), x], x] + \text{Simp}[(c - I \cdot d)/2 \ \text{Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (1 + I \cdot \tan[e + f \cdot x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{!IntegerQ}[m]$

rule 4113 $\text{Int}[(a + (b \cdot \tan[e + f \cdot x]) + (c + (d \cdot \tan[e + f \cdot x]) + (f \cdot x))^m \cdot (A + (B \cdot \tan[e + f \cdot x]) + (C \cdot \tan[e + f \cdot x]) + (f \cdot x))^2], x_Symbol] \rightarrow \text{Simp}[C \cdot (a + b \cdot \tan[e + f \cdot x])^{m+1} / (b \cdot f \cdot (m+1)), x] + \text{Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot \text{Simp}[A - C + B \cdot \tan[e + f \cdot x], x], x] /; \text{FreeQ}[\{a, b, e, f, A, B, C, m\}, x] \ \&\& \ \text{NeQ}[A \cdot b^2 - a \cdot b \cdot B + a^2 \cdot C, 0] \ \&\& \ \text{!LeQ}[m, -1]$

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4429 vs. $2(424) = 848$.

Time = 0.73 (sec) , antiderivative size = 4430, normalized size of antiderivative = 9.55

method	result	size
parts	Expression too large to display	4430
derivativedivides	Expression too large to display	6624
default	Expression too large to display	6624

input

```
int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x,method=_RETURNVERBOSE)
```

output

```

2*C*b^3/f/d^4*(1/9*(c+d*tan(f*x+e))^(9/2)-3/7*c*(c+d*tan(f*x+e))^(7/2)+3/5
*c^2*(c+d*tan(f*x+e))^(5/2)-1/5*(c+d*tan(f*x+e))^(5/2)*d^2-1/3*c^3*(c+d*ta
n(f*x+e))^(3/2)+1/3*c*d^2*(c+d*tan(f*x+e))^(3/2)+(c+d*tan(f*x+e))^(1/2)*d^
4+d^4*(-1/8*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e
))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+1/2*(-(c^2+d^2)^(1
/2)+c)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(
c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))+1/8*(2*(c^2+d^2)
^(1/2)+2*c)^(1/2)*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1
/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+1/2*(-(c^2+d^2)^(1/2)+c)/(2*(c^2+d^2)^(1/2
)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)-(2*(c^2+d^2)^(1/2)+2*c)^(1/2
)))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))) -3/4/f/d*ln(d*tan(f*x+e)+c-(c+d*tan(f*x
+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(
1/2)+2*c)^(1/2)*a*b^2*c-3/4/f/d*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(
2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*B*(c^2+d^2)^(1/2)*(2*(c^2+d^
2)^(1/2)+2*c)^(1/2)*a^2*b+3/4/f/d*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)
*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(
1/2)*a^2*b*c+3/4/f/d*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)
^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(c^2+d^2)^(1/2)*(2*(c^2+d^2)^(1/2)+2
*c)^(1/2)*a*b^2+3/4/f/d*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d
^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*a...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35153 vs. $2(414) = 828$.

Time = 10.12 (sec) , antiderivative size = 35153, normalized size of antiderivative = 75.76

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```

integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(
f*x+e)^2),x, algorithm="fricas")

```

output

Too large to include

Sympy [F]

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**3*(c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**3*sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `Timed out`

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Exception raised: TypeError

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,29,11]}%%}+%%{12,[0,27,11]}%%}+%%{66,[0,25,11]}%%}+%%{2
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Hanged

input

```
int((a + b*tan(e + f*x))^3*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^4 + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^5 dx \right) b^3 c \\
&\quad + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) a b^2 c \\
&\quad + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b^4 \\
&\quad + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a^2 b c \\
&\quad + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a b^3 \\
&\quad + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^3 c \\
&\quad + 6 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^2 b^2 \\
&\quad + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^3 b
\end{aligned}$$

input

```
int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**4 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5,x)*b**3*c + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a*b**2*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**4 + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a**2*b*c + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*b**3 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**3*c + 6*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*b**2 + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**3*b
```

3.91 $\int (a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}(A+B \tan(e$

Optimal result	1023
Mathematica [A] (verified)	1024
Rubi [A] (warning: unable to verify)	1025
Maple [B] (verified)	1030
Fricas [B] (verification not implemented)	1031
Sympy [F]	1032
Maxima [F(-1)]	1032
Giac [F(-2)]	1033
Mupad [F(-1)]	1033
Reduce [F]	1034

Optimal result

Integrand size = 47, antiderivative size = 325

$$\int (a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx) + C \tan^2(e+fx)) dx$$

$$= -\frac{(a-ib)^2(B+i(A-C))\sqrt{c-id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f}$$

$$- \frac{(a+ib)^2(B-i(A-C))\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f}$$

$$+ \frac{2(a^2B-b^2B+2ab(A-C))\sqrt{c+d \tan(e+fx)}}{f}$$

$$+ \frac{2(20a^2Cd^2-14abd(2cC-5Bd)+b^2(8c^2C-14Bcd+35(A-C)d^2))(c+d \tan(e+fx))^{3/2}}{105d^3f}$$

$$- \frac{2b(4bcC-7bBd-4aCd)\tan(e+fx)(c+d \tan(e+fx))^{3/2}}{35d^2f}$$

$$+ \frac{2C(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{3/2}}{7df}$$

output

$$\begin{aligned} & -(a-I*b)^2*(B+I*(A-C))*(c-I*d)^{(1/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)/(c-I*d)} \\ &)^{(1/2)}/f-(a+I*b)^2*(B-I*(A-C))*(c+I*d)^{(1/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)} \\ &)^{(1/2)/(c+I*d)^{(1/2)}/f+2*(B*a^2-B*b^2+2*a*b*(A-C))*(c+d*\tan(f*x+e))^{(1/2)}/f+ \\ & 2/105*(20*a^2*C*d^2-14*a*b*d*(-5*B*d+2*C*c)+b^2*(8*c^2*C-14*B*c*d+35*(A-C) \\ & *d^2))*(c+d*\tan(f*x+e))^{(3/2)}/d^3/f-2/35*b*(-7*B*b*d-4*C*a*d+4*C*b*c)*\tan(\\ & f*x+e)*(c+d*\tan(f*x+e))^{(3/2)}/d^2/f+2/7*C*(a+b*\tan(f*x+e))^2*(c+d*\tan(f*x+ \\ & e))^{(3/2)}/d/f \end{aligned}$$
Mathematica [A] (verified)

Time = 3.30 (sec) , antiderivative size = 314, normalized size of antiderivative = 0.97

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2 \left((20a^2Cd^2 + 14abd(-2cC + 5Bd) + b^2(8c^2C - 14Bcd + 35(A - C)d^2)) (c + d \tan(e + fx))^{3/2} + 3bd \right)}{105d^3f}$$

input

$$\text{Integrate}[(a + b*\text{Tan}[e + f*x])^2*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2),x]$$

output

$$\begin{aligned} & (2*((20*a^2*C*d^2 + 14*a*b*d*(-2*c*C + 5*B*d) + b^2*(8*c^2*C - 14*B*c*d + \\ & 35*(A - C)*d^2))*(c + d*\text{Tan}[e + f*x])^{(3/2)} + 3*b*d*(-4*b*c*C + 7*b*B*d + \\ & 4*a*C*d)*\text{Tan}[e + f*x]*(c + d*\text{Tan}[e + f*x])^{(3/2)} + 15*C*d^2*(a + b*\text{Tan}[e + \\ & f*x])^2*(c + d*\text{Tan}[e + f*x])^{(3/2)} + (105*(a - I*b)^2*(I*A + B - I*C)*d^3 \\ & *(-(\text{Sqrt}[c - I*d]*\text{ArcTanh}[\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[c - I*d]]) + \text{Sqrt}[\\ & c + d*\text{Tan}[e + f*x]]))/2 + (105*(a + I*b)^2*((-I)*A + B + I*C)*d^3*(-(\text{Sqrt}[\\ & c + I*d]*\text{ArcTanh}[\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[c + I*d]]) + \text{Sqrt}[c + d*\text{Tan} \\ & [e + f*x]]))/2))/(105*d^3*f) \end{aligned}$$

Rubi [A] (warning: unable to verify)

Time = 3.75 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.02, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.362$, Rules used = {3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4130

$$\frac{2 \int -\frac{1}{2} (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} ((4bcC - 4adC - 7bBd) \tan^2(e + fx) - 7(Ab - Cb + aB)d \tan(e + fx) + 2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}) dx}{7df}$$

↓ 27

$$\frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} ((4bcC - 4adC - 7bBd) \tan^2(e + fx) - 7(Ab - Cb + aB)d \tan(e + fx) + 2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}) dx}{7d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} ((4bcC - 4adC - 7bBd) \tan(e + fx)^2 - 7(Ab - Cb + aB)d \tan(e + fx) + 2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}) dx}{7d}$$

↓ 4120

$$\frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx) (-4aCd - 7bBd + 4bcC) (c + d \tan(e + fx))^{3/2}}{5df} - \frac{2 \int -\frac{1}{2} \sqrt{c + d \tan(e + fx)} (-2c(4cC - 7Bd)b^2 + 28acCdb - 5a^2(7A - 3C)d^2 - ((8Cc^2 - 7d^2) \tan^2(e + fx) - 7(Ab - Cb + aB)d \tan(e + fx) + 2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2})) dx}{7d}}$$

↓ 27

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c+d \tan(e+fx)}(-2c(4cC-7Bd)b^2+28acCdb-5a^2(7A-3C)d^2-((8Cc^2-14Bdc+35(A-C)d^2)b^2-14ad(2cC-5Bd)b+20a^2Cd^2) \tan^2(e+fx)}{5d} dx}{7d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c+d \tan(e+fx)}(-2c(4cC-7Bd)b^2+28acCdb-5a^2(7A-3C)d^2-((8Cc^2-14Bdc+35(A-C)d^2)b^2-14ad(2cC-5Bd)b+20a^2Cd^2) \tan(e+fx)}{5d} dx}{7d}$$

↓ 4113

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c+d \tan(e+fx)}(35(-((A-C)a^2)+2bBa+b^2(A-C))d^2-35(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx))dx - \frac{2(c+d \tan(e+fx))^{3/2}(20a^2Cd^2-14abd)}{5d}}{5d} dx}{7d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \sqrt{c+d \tan(e+fx)}(35(-((A-C)a^2)+2bBa+b^2(A-C))d^2-35(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx))dx - \frac{2(c+d \tan(e+fx))^{3/2}(20a^2Cd^2-14abd)}{5d}}{5d} dx}{7d}$$

↓ 4011

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \frac{-35((Ac-Cc-Bd)a^2-2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd))d^2-35((Bc+(A-C)d)a^2+2b(Ac-Cc-Bd)a-b^2(Bc+(A-C)d)) \tan(e+fx)d^2}{\sqrt{c+d \tan(e+fx)}} dx - \frac{2(c+d \tan(e+fx))^{3/2}}{5d}}{5d} dx}{7d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{\int \frac{-35((Ac-Cc-Bd)a^2-2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd))d^2-35((Bc+(A-C)d)a^2+2b(Ac-Cc-Bd)a-b^2(Bc+(A-C)d)) \tan(e+fx)d^2}{\sqrt{c+d \tan(e+fx)}} dx - \frac{2(c+d \tan(e+fx))^{3/2}}{5d}}{5d} dx}{7d}$$

↓ 4022

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(-4aCd - 7bBd + 4bcC)(c + d \tan(e + fx))^{3/2}}{5df} + \frac{-\frac{35}{2}d^2(a+ib)^2(c+id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{35}{2}d^2(a-ib)^2(c-id)(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{7df}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(-4aCd - 7bBd + 4bcC)(c + d \tan(e + fx))^{3/2}}{5df} + \frac{-\frac{35}{2}d^2(a+ib)^2(c+id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{35}{2}d^2(a-ib)^2(c-id)(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{7df}$$

↓ 4020

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(-4aCd - 7bBd + 4bcC)(c + d \tan(e + fx))^{3/2}}{5df} + \frac{35id^2(a-ib)^2(c-id)(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - 35id^2(a+ib)^2(c+id)(A+iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{7df}$$

↓ 25

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(-4aCd - 7bBd + 4bcC)(c + d \tan(e + fx))^{3/2}}{5df} + \frac{35id^2(a-ib)^2(c-id)(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - 35id^2(a+ib)^2(c+id)(A+iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{7df}$$

↓ 73

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(-4aCd - 7bBd + 4bcC)(c + d \tan(e + fx))^{3/2}}{5df} + \frac{35d(a-ib)^2(c-id)(A-iB-C) \int \frac{1}{i \tan^2(e+fx) + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)} - 35d(a+ib)^2(c+id)(A+iB-C) \int \frac{1}{i \tan^2(e+fx) + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{7df}$$

↓ 221

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}}{7df} - \frac{2b \tan(e + fx)(-4aCd - 7bBd + 4bcC)(c + d \tan(e + fx))^{3/2}}{5df} + \frac{2(c+d \tan(e+fx))^{3/2}(20a^2Cd^2 - 14abd(2cC - 5Bd) + b^2(35d^2(A-C) - 14Bcd + 8c^2C))}{3df}$$

input $\text{Int}[(a + b \cdot \tan[e + f \cdot x])^2 \sqrt{c + d \cdot \tan[e + f \cdot x]} \cdot (A + B \cdot \tan[e + f \cdot x] + C \cdot \tan[e + f \cdot x]^2), x]$

output $(2 \cdot C \cdot (a + b \cdot \tan[e + f \cdot x])^2 \cdot (c + d \cdot \tan[e + f \cdot x])^{3/2}) / (7 \cdot d \cdot f) - ((2 \cdot b \cdot (4 \cdot b \cdot c \cdot C - 7 \cdot b \cdot B \cdot d - 4 \cdot a \cdot C \cdot d) \cdot \tan[e + f \cdot x] \cdot (c + d \cdot \tan[e + f \cdot x])^{3/2}) / (5 \cdot d \cdot f) + ((-35 \cdot (a - I \cdot b)^2 \cdot (A - I \cdot B - C) \cdot \sqrt{c - I \cdot d} \cdot d^2 \cdot \text{ArcTan}[\tan[e + f \cdot x] / \sqrt{c - I \cdot d}]) / f - (35 \cdot (a + I \cdot b)^2 \cdot (A + I \cdot B - C) \cdot \sqrt{c + I \cdot d} \cdot d^2 \cdot \text{ArcTan}[\tan[e + f \cdot x] / \sqrt{c + I \cdot d}]) / f - (70 \cdot (a^2 \cdot B - b^2 \cdot B + 2 \cdot a \cdot b \cdot (A - C)) \cdot d^2 \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]}) / f - (2 \cdot (20 \cdot a^2 \cdot C \cdot d^2 - 14 \cdot a \cdot b \cdot d \cdot (2 \cdot c \cdot C - 5 \cdot B \cdot d) + b^2 \cdot (8 \cdot c^2 \cdot C - 14 \cdot B \cdot c \cdot d + 35 \cdot (A - C) \cdot d^2)) \cdot (c + d \cdot \tan[e + f \cdot x])^{3/2}) / (3 \cdot d \cdot f)) / (5 \cdot d)) / (7 \cdot d)$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_)(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)(G_x)] /; \text{FreeQ}[b, x]$

rule 73 $\text{Int}[(a_ + (b_)(x_))^{(m_)} \cdot ((c_ + (d_)(x_))^{(n_)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p \cdot (m + 1) - 1)} \cdot (c - a \cdot (d/b) + d \cdot (x^p/b))^{(n)}, x], x, (a + b \cdot x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x / \text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4011 $\text{Int}[\left((a_{.}) + (b_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right), x_Symbol] \rightarrow \text{Simp}[d\left((a + b\tan[e + f*x])^m/(f*m)\right), x] + \text{Int}[(a + b\tan[e + f*x])^{(m-1)}\text{Simp}[a*c - b*d + (b*c + a*d)\tan[e + f*x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

rule 4020 $\text{Int}[\left((a_{.}) + (b_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\tan[e + f*x]], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]

rule 4022 $\text{Int}[\left((a_{.}) + (b_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{Int}[(a + b*\tan[e + f*x])^m*(1 - I*\tan[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{Int}[(a + b*\tan[e + f*x])^m*(1 + I*\tan[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4113 $\text{Int}[\left((a_{.}) + (b_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right)^{(m_{.})}\left((A_{.}) + (B_{.})\tan[(e_{.}) + (f_{.})(x_{.})] + (C_{.})\tan[(e_{.}) + (f_{.})(x_{.})]^2\right), x_Symbol] \rightarrow \text{Simp}[C\left((a + b*\tan[e + f*x])^{(m+1)}/(b*f*(m+1))\right), x] + \text{Int}[(a + b*\tan[e + f*x])^m\text{Simp}[A - C + B*\tan[e + f*x], x], x] /;$ FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]

rule 4120 $\text{Int}[\left((a_{.}) + (b_{.})\tan[(e_{.}) + (f_{.})(x_{.})]\right)^{(n_{.})}\left((c_{.}) + (d_{.})\tan[(e_{.}) + (f_{.})(x_{.})] + (A_{.}) + (B_{.})\tan[(e_{.}) + (f_{.})(x_{.})] + (C_{.})\tan[(e_{.}) + (f_{.})(x_{.})]^2\right), x_Symbol] \rightarrow \text{Simp}[b*C*\tan[e + f*x]\left((c + d*\tan[e + f*x])^{(n+1)}/(d*f*(n+2))\right), x] - \text{Simp}[1/(d*(n+2)) \text{Int}[(c + d*\tan[e + f*x])^n\text{Simp}[b*c*C - a*A*d*(n+2) - (A*b + a*B - b*C)*d*(n+2)*\tan[e + f*x] - (a*C*d*(n+2) - b*(c*C - B*d*(n+2))]*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]

rule 4130

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3318 vs. $2(291) = 582$.

Time = 0.30 (sec) , antiderivative size = 3319, normalized size of antiderivative = 10.21

method	result	size
parts	Expression too large to display	3319
derivativeldivides	Expression too large to display	4775
default	Expression too large to display	4775

input

```

int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x,method=_RETURNVERBOSE)

```

output

```
A*a^2/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)-(
2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))-1/4/f/d*ln(d*
tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2
)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*b^2*c+1/4/f/d*ln(d*tan(f*x+e)+c+(
c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*
(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*a^2-1/4/f/d*ln(d*tan(f*x+e)+c+(
c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*
(c^2+d^2)^(1/2)+2*c)^(1/2)*a^2*c-2/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arcta
n((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1
/2)-2*c)^(1/2))*B*a*b-1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*
(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2
)*(c^2+d^2)^(1/2)*a^2+1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*
(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2
)*(c^2+d^2)^(1/2)*b^2+1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*
(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2
)*a^2*c+1/2/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2
)+2*c)^(1/2)+(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1
/2)*a*b-1/2/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2
)+2*c)^(1/2)+(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*a*b*c+2/f*b*
(B*b+2*C*a)/d^2*(1/5*(c+d*tan(f*x+e))^(5/2)-1/3*(c+d*tan(f*x+e))^(3/2))*...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 23984 vs. $2(281) = 562$.

Time = 4.33 (sec) , antiderivative size = 23984, normalized size of antiderivative = 73.80

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(
f*x+e)^2),x, algorithm="fricas")
```

output

Too large to include

Sympy [F]

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**2*(c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**2*sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `Timed out`

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Exception raised: TypeError

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,24,9]}%%}+%%{10,[0,22,9]}%%}+%%{45,[0,20,9]}%%}+%%{120,
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Hanged

input

```
int((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^3 + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b^2 c \\
&\quad + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) abc \\
&\quad + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) b^3 \\
&\quad + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^2 c \\
&\quad + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a b^2 \\
&\quad + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^2 b
\end{aligned}$$

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**3 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**2*c + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*b*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b**3 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*c + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*b**2 + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**2*b
```

3.92 $\int (a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)} (A+B \tan(e$

Optimal result	1035
Mathematica [A] (verified)	1036
Rubi [A] (warning: unable to verify)	1036
Maple [B] (verified)	1040
Fricas [B] (verification not implemented)	1041
Sympy [F]	1042
Maxima [F]	1042
Giac [F(-2)]	1043
Mupad [B] (verification not implemented)	1043
Reduce [F]	1044

Optimal result

Integrand size = 45, antiderivative size = 224

$$\int (a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx) + C \tan^2(e+fx)) dx$$

$$= -\frac{(ia+b)(A-iB-C)\sqrt{c-id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f}$$

$$+ \frac{(ia-b)(A+iB-C)\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f}$$

$$+ \frac{2(Ab+aB-bC)\sqrt{c+d \tan(e+fx)}}{f}$$

$$- \frac{2(2bcC-5bBd-5aCd)(c+d \tan(e+fx))^{3/2}}{15d^2 f}$$

$$+ \frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{3/2}}{5df}$$

output

```

-(I*a+b)*(A-I*B-C)*(c-I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1
/2))/f+(I*a-b)*(A+I*B-C)*(c+I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I
*d)^(1/2))/f+2*(A*b+B*a-C*b)*(c+d*tan(f*x+e))^(1/2)/f-2/15*(-5*B*b*d-5*C*a
*d+2*C*b*c)*(c+d*tan(f*x+e))^(3/2)/d^2/f+2/5*b*C*tan(f*x+e)*(c+d*tan(f*x+e
))^3/2/d/f
    
```


Mathematica [A] (verified)

Time = 1.36 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.98

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{2(-2bcC + 5bBd + 5aCd)(c + d \tan(e + fx))^{3/2}}{d} + 6bC \tan(e + fx)(c + d \tan(e + fx))^{3/2} + 15(ia + b)(A - iB - C)d$$

input

```
Integrate[(a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x]
] + C*Tan[e + f*x]^2), x]
```

output

```
((2*(-2*b*c*C + 5*b*B*d + 5*a*C*d)*(c + d*Tan[e + f*x])^(3/2))/d + 6*b*C*T
an[e + f*x]*(c + d*Tan[e + f*x])^(3/2) + 15*(I*a + b)*(A - I*B - C)*d*(-(S
qrt[c - I*d]*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]) + Sqrt[c + d
*Tan[e + f*x]]) + 15*((-I)*a + b)*(A + I*B - C)*d*(-(Sqrt[c + I*d]*ArcTanh
[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]]) + Sqrt[c + d*Tan[e + f*x]])))/(15
*d*f)
```

Rubi [A] (warning: unable to verify)

Time = 2.05 (sec) , antiderivative size = 218, normalized size of antiderivative = 0.97, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.311$, Rules used = {3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow \text{4120}$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{2 \int \frac{1}{2} \sqrt{c + d \tan(e + fx)} ((2bcC - 5adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 5aAd) dx}{5d}$$

↓ 27

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{\int \sqrt{c + d \tan(e + fx)} ((2bcC - 5adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 5aAd) dx}{5d}$$

↓ 3042

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{\int \sqrt{c + d \tan(e + fx)} ((2bcC - 5adC - 5bBd) \tan(e + fx)^2 - 5(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 5aAd) dx}{5d}$$

↓ 4113

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{\int \sqrt{c + d \tan(e + fx)} (5(bB - a(A - C))d - 5(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{2(-5aCd - 5bBd + 2bcC)(c + d \tan(e + fx))}{3df}}{5d}$$

↓ 3042

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{\int \sqrt{c + d \tan(e + fx)} (5(bB - a(A - C))d - 5(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{2(-5aCd - 5bBd + 2bcC)(c + d \tan(e + fx))}{3df}}{5d}$$

↓ 4011

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{\int \frac{5d(bBc + b(A - C)d - a(Ac - Cc - Bd)) - 5d(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{10d(aB + Ab - bC) \sqrt{c + d \tan(e + fx)}}{f} + \frac{2(-5aCd - 5bBd + 2bcC)(c + d \tan(e + fx))}{3df}}{5d}$$

↓ 3042

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \frac{\int \frac{5d(bBc + b(A - C)d - a(Ac - Cc - Bd)) - 5d(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{10d(aB + Ab - bC) \sqrt{c + d \tan(e + fx)}}{f} + \frac{2(-5aCd - 5bBd + 2bcC)(c + d \tan(e + fx))}{3df}}{5d}$$

↓ 4022

$$\begin{aligned}
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \\
 & \frac{-\frac{5}{2}d(a + ib)(c + id)(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{5}{2}d(a - ib)(c - id)(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx - 10}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \\
 & \frac{-\frac{5}{2}d(a + ib)(c + id)(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{5}{2}d(a - ib)(c - id)(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx - 10}{5d} \\
 & \quad \downarrow \text{4020} \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \\
 & \frac{5id(a - ib)(c - id)(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} + \frac{5id(a + ib)(c + id)(A + iB - C) \int -\frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \\
 & \quad \downarrow \text{25} \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \\
 & \frac{5id(a - ib)(c - id)(A - iB - C) \int \frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \frac{5id(a + ib)(c + id)(A + iB - C) \int \frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \\
 & \quad \downarrow \text{73} \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \\
 & \frac{5(a + ib)(c + id)(A + iB - C) \int \frac{1}{-\frac{i \tan^2(e + fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{f} - \frac{5(a - ib)(c - id)(A - iB - C) \int \frac{1}{\frac{i \tan^2(e + fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{f} - \\
 & \quad \downarrow \text{221} \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{3/2}}{5df} - \\
 & \frac{5d(a - ib)\sqrt{c - id}(A - iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f} - \frac{5d(a + ib)\sqrt{c + id}(A + iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c + id}}\right)}{f} - \frac{10d(aB + Ab - bC)\sqrt{c + d \tan(e + fx)}}{f} -
 \end{aligned}$$

input

```
Int[(a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*
Tan[e + f*x]^2), x]
```

output $(2*b*C*\text{Tan}[e + f*x]*(c + d*\text{Tan}[e + f*x])^{(3/2)})/(5*d*f) - ((-5*(a - I*b)*(A - I*B - C)*\text{Sqrt}[c - I*d]*d*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]])/f - (5*(a + I*b)*(A + I*B - C)*\text{Sqrt}[c + I*d]*d*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c + I*d]])/f - (10*(A*b + a*B - b*C)*d*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/f + (2*(2*b*c*C - 5*b*B*d - 5*a*C*d)*(c + d*\text{Tan}[e + f*x])^{(3/2)})/(3*d*f)/(5*d)$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$

rule 73 $\text{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4011 $\text{Int}[(a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)]^{(m_)}*((c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$

rule 4020 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x], x_Symbol] \rightarrow \text{Simp}[c(d/f) \text{Subst}[\text{Int}[(a + (b/d)x]^m/(d^2 + cx), x], x, d\tan[e + fx]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x], x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{Int}[(a + b\tan[e + fx])^m(1 - I\tan[e + fx]), x], x] + \text{Simp}[(c - I*d)/2 \text{Int}[(a + b\tan[e + fx])^m(1 + I\tan[e + fx]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!IntegerQ}[m]$

rule 4113 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x] + (C_.)\tan[(e_.) + (f_.)x]^2, x_Symbol] \rightarrow \text{Simp}[C*((a + b\tan[e + fx])^{m+1}/(b*f*(m+1))), x] + \text{Int}[(a + b\tan[e + fx])^m \text{Simp}[A - C + B\tan[e + fx], x], x] /; \text{FreeQ}\{a, b, e, f, A, B, C, m\}, x] \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{!LeQ}[m, -1]$

rule 4120 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x] * ((c_.) + (d_.)\tan[(e_.) + (f_.)x])^{n_.*} * ((A_.) + (B_.)\tan[(e_.) + (f_.)x] + (C_.)\tan[(e_.) + (f_.)x]^2), x_Symbol] \rightarrow \text{Simp}[b*C\tan[e + fx] * ((c + d\tan[e + fx])^{n+1}/(d*f*(n+2))), x] - \text{Simp}[1/(d*(n+2)) \text{Int}[(c + d\tan[e + fx])^n \text{Simp}[b*c*C - a*A*d*(n+2) - (A*b + a*B - b*C)*d*(n+2)*\tan[e + fx] - (a*C*d*(n+2) - b*(c*C - B*d*(n+2)))*\tan[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!LtQ}[n, -1]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2196 vs. $2(194) = 388$.

Time = 0.27 (sec) , antiderivative size = 2197, normalized size of antiderivative = 9.81

method	result	size
parts	Expression too large to display	2197
derivativedivides	Expression too large to display	3028
default	Expression too large to display	3028

input `int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `1/f*(A*b+B*a)*(2*(c+d*tan(f*x+e))^(1/2)-1/4*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+(-(c^2+d^2)^(1/2)+c)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))+1/4*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+(-(c^2+d^2)^(1/2)+c)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)-(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)))+2/3/f/d*B*b*(c+d*tan(f*x+e))^(3/2)+2/3/f/d*C*a*(c+d*tan(f*x+e))^(3/2)+1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*b+1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*a-1/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*B*b-1/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*C*a-1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*b*c-1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*a*c-1/4/...`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12410 vs. $2(187) = 374$.

Time = 1.47 (sec) , antiderivative size = 12410, normalized size of antiderivative = 55.40

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))*sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan^2(fx + e) + B \tan(fx + e) + A) (b \tan(fx + e) + a) \sqrt{d \tan(fx + e) + c} dx$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)*sqrt(d*tan(f*x + e) + c), x)`

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Exception raised: TypeError

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1,[0,19,7]}%%}+%%{8,[0,17,7]}%%}+%%{28,[0,15,7]}%%}+%%{56,[0
```

Mupad [B] (verification not implemented)

Time = 53.78 (sec) , antiderivative size = 22955, normalized size of antiderivative = 102.48

$$\int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
int((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```


output

```

((2*B*a*d - 4*C*a*c)/(d*f) + (4*C*a*c)/(d*f))*(c + d*tan(e + f*x))^(1/2) +
((2*B*b*d - 6*C*b*c)/(3*d^2*f) + (4*C*b*c)/(3*d^2*f))*(c + d*tan(e + f*x)
)^(3/2) + (c + d*tan(e + f*x))^(1/2)*(2*c*((2*B*b*d - 6*C*b*c)/(d^2*f) + (
4*C*b*c)/(d^2*f)) + (2*A*b*d^2 + 6*C*b*c^2 - 4*B*b*c*d)/(d^2*f) - (2*C*b*(
d^4*f + c^2*d^2*f))/(d^4*f^2)) - atan((((8*(4*A*b*d^4*f^2 - 4*C*b*d^4*f^2
+ 4*A*b*c^2*d^2*f^2 - 4*C*b*c^2*d^2*f^2))/f^3 - 64*c*d^2*(c + d*tan(e + f
*x))^(1/2)*((A^2*b^2*c)/(4*f^2) - (4*A*C^3*b^4*d^2*f^4 - B^4*b^4*d^2*f^4 -
C^4*b^4*d^2*f^4 - A^4*b^4*d^2*f^4 + 4*A^3*C*b^4*d^2*f^4 - 4*A^2*B^2*b^4*c
^2*f^4 + 2*A^2*B^2*b^4*d^2*f^4 - 6*A^2*C^2*b^4*d^2*f^4 - 4*B^2*C^2*b^4*c^2
*f^4 + 2*B^2*C^2*b^4*d^2*f^4 + 4*A*B^3*b^4*c*d*f^4 - 4*A^3*B*b^4*c*d*f^4 +
4*B*C^3*b^4*c*d*f^4 - 4*B^3*C*b^4*c*d*f^4 + 8*A*B^2*C*b^4*c^2*f^4 - 4*A*B
^2*C*b^4*d^2*f^4 - 12*A*B*C^2*b^4*c*d*f^4 + 12*A^2*B*C*b^4*c*d*f^4)^(1/2)/
(4*f^4) - (B^2*b^2*c)/(4*f^2) + (C^2*b^2*c)/(4*f^2) - (A*B*b^2*d)/(2*f^2)
- (A*C*b^2*c)/(2*f^2) + (B*C*b^2*d)/(2*f^2))^(1/2))*((A^2*b^2*c)/(4*f^2) -
(4*A*C^3*b^4*d^2*f^4 - B^4*b^4*d^2*f^4 - C^4*b^4*d^2*f^4 - A^4*b^4*d^2*f^
4 + 4*A^3*C*b^4*d^2*f^4 - 4*A^2*B^2*b^4*c^2*f^4 + 2*A^2*B^2*b^4*d^2*f^4 -
6*A^2*C^2*b^4*d^2*f^4 - 4*B^2*C^2*b^4*c^2*f^4 + 2*B^2*C^2*b^4*d^2*f^4 + 4*
A*B^3*b^4*c*d*f^4 - 4*A^3*B*b^4*c*d*f^4 + 4*B*C^3*b^4*c*d*f^4 - 4*B^3*C*b^
4*c*d*f^4 + 8*A*B^2*C*b^4*c^2*f^4 - 4*A*B^2*C*b^4*d^2*f^4 - 12*A*B*C^2*b^4
*c*d*f^4 + 12*A^2*B*C*b^4*c*d*f^4)^(1/2)/(4*f^4) - (B^2*b^2*c)/(4*f^2) ...

```

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^2 + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) bc \\
&\quad + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) ac \\
&\quad + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) b^2 \\
&\quad + 2 \left(\int \sqrt{d \tan(fx + e) + c} dx \right) ab
\end{aligned}$$

input

```

int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2
),x)

```

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e
+ f*x)**3,x)*b*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*c + i
nt(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*b**2 + 2*int(sqrt(tan(e + f
*x)*d + c)*tan(e + f*x),x)*a*b
```

3.93 $\int \sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx)) dx$

Optimal result	1046
Mathematica [A] (verified)	1047
Rubi [A] (warning: unable to verify)	1047
Maple [B] (verified)	1051
Fricas [B] (verification not implemented)	1052
Sympy [F]	1053
Maxima [F]	1054
Giac [F(-2)]	1054
Mupad [B] (verification not implemented)	1055
Reduce [F]	1055

Optimal result

Integrand size = 35, antiderivative size = 155

$$\int \sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= -\frac{(iA + B - iC)\sqrt{c - id} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c - id}}\right)}{f}$$

$$- \frac{(B - i(A - C))\sqrt{c + id} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c + id}}\right)}{f}$$

$$+ \frac{2B\sqrt{c + d \tan(e + fx)}}{f} + \frac{2C(c + d \tan(e + fx))^{3/2}}{3df}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/f
-(B-I*(A-C))*(c+I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/f
+2*B*(c+d*tan(f*x+e))^(1/2)/f+2/3*C*(c+d*tan(f*x+e))^(3/2)/d/f
    
```

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.97

$$\int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{-3i(A - iB - C)\sqrt{c - id} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c - id}}\right) + 3i(A + iB - C)\sqrt{c + id} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c + id}}\right)}{3df}$$

input

```
Integrate[Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
((-3*I)*(A - I*B - C)*Sqrt[c - I*d]*d*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]] + (3*I)*(A + I*B - C)*Sqrt[c + I*d]*d*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]] + 2*Sqrt[c + d*Tan[e + f*x]]*(c*C + 3*B*d + C*d*Tan[e + f*x]))/(3*d*f)
```

Rubi [A] (warning: unable to verify)

Time = 1.28 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.86, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4113, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4113$$

$$\int (A - C + B \tan(e + fx)) \sqrt{c + d \tan(e + fx)} dx + \frac{2C(c + d \tan(e + fx))^{3/2}}{3df}$$

$$\downarrow 3042$$

$$\begin{aligned}
& \int (A - C + B \tan(e + fx)) \sqrt{c + d \tan(e + fx)} dx + \frac{2C(c + d \tan(e + fx))^{3/2}}{3df} \\
& \quad \downarrow 4011 \\
& \int \frac{Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2B\sqrt{c + d \tan(e + fx)}}{f} + \\
& \quad \frac{2C(c + d \tan(e + fx))^{3/2}}{3df} \\
& \quad \downarrow 3042 \\
& \int \frac{Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2B\sqrt{c + d \tan(e + fx)}}{f} + \\
& \quad \frac{2C(c + d \tan(e + fx))^{3/2}}{3df} \\
& \quad \downarrow 4022 \\
& \frac{1}{2}(c + id)(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(c - id)(A - iB - \\
C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2B\sqrt{c + d \tan(e + fx)}}{f} + \frac{2C(c + d \tan(e + fx))^{3/2}}{3df} \\
& \quad \downarrow 3042 \\
& \frac{1}{2}(c + id)(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(c - id)(A - iB - \\
C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2B\sqrt{c + d \tan(e + fx)}}{f} + \frac{2C(c + d \tan(e + fx))^{3/2}}{3df} \\
& \quad \downarrow 4020 \\
& \frac{i(c - id)(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \\
& \frac{i(c + id)(A + iB - C) \int -\frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}} d(-i \tan(e + fx))}{2f} + \\
& \frac{2B\sqrt{c + d \tan(e + fx)}}{f} + \frac{2C(c + d \tan(e + fx))^{3/2}}{3df} \\
& \quad \downarrow 25
\end{aligned}$$

$$\begin{aligned}
& \frac{i(c-id)(A-iB-C) \int \frac{1}{(1-i \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} + \\
& \frac{i(c+id)(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1) \sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} + \\
& \frac{2B \sqrt{c+d \tan(e+fx)}}{f} + \frac{2C(c+d \tan(e+fx))^{3/2}}{3df} \\
& \quad \downarrow 73 \\
& \frac{(c+id)(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d \sqrt{c+d \tan(e+fx)}}{df} + \\
& \frac{(c-id)(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d \sqrt{c+d \tan(e+fx)}}{df} + \frac{2B \sqrt{c+d \tan(e+fx)}}{f} + \\
& \frac{2C(c+d \tan(e+fx))^{3/2}}{3df} \\
& \quad \downarrow 221 \\
& \frac{\sqrt{c-id}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{\sqrt{c+id}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f} + \\
& \frac{2B \sqrt{c+d \tan(e+fx)}}{f} + \frac{2C(c+d \tan(e+fx))^{3/2}}{3df}
\end{aligned}$$

input `Int[Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `((A - I*B - C)*Sqrt[c - I*d]*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + ((A + I*B - C)*Sqrt[c + I*d]*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f + (2*B*Sqrt[c + d*Tan[e + f*x]])/f + (2*C*(c + d*Tan[e + f*x])^(3/2))/(3*d*f)`

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{\text{m}_}) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{n}_}), \text{x_Symbol}] \rightarrow \text{With}[\{p = \text{Denominator}[\text{m}]\}, \text{Simp}[p/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{(p*(\text{m} + 1) - 1)} * (\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^p/\text{b}))^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{(1/p)}], \text{x}]] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[-1, \text{m}, 0] \&\& \text{LeQ}[-1, \text{n}, 0] \&\& \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \&\& \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a}) * \text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4011 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_}) * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])), \text{x_Symbol}] \rightarrow \text{Simp}[\text{d} * ((\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} / (\text{f} * \text{m})), \text{x}] + \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m} - 1} * \text{Simp}[\text{a} * \text{c} - \text{b} * \text{d} + (\text{b} * \text{c} + \text{a} * \text{d}) * \text{Tan}[\text{e} + \text{f} * \text{x}], \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{GtQ}[\text{m}, 0]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_}) * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c} * (\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d}) * \text{x})^{\text{m}} / (\text{d}^2 + \text{c} * \text{x}), \text{x}], \text{x}, \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_}) * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I} * \text{d})/2 \quad \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 - \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I} * \text{d})/2 \quad \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 + \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{!IntegerQ}[\text{m}]$

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1297 vs. $2(130) = 260$.

Time = 0.18 (sec) , antiderivative size = 1298, normalized size of antiderivative = 8.37

method	result	size
parts	Expression too large to display	1298
derivativedivides	Expression too large to display	1472
default	Expression too large to display	1472

input

```
int(((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETUR
NVERBOSE)
```


output

```

-1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(
(1/2)+(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)+1/f
*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+
d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*A+1/4/f/d*ln(d*tan(f
*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/
2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c+1/4*A/f/d*(2*(c^2+d^2)^(1/2)+2*c)^(1
/2)*(c^2+d^2)^(1/2)*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(
1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+A/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arct
an((2*(c+d*tan(f*x+e))^(1/2)-(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(
1/2)-2*c)^(1/2))-1/4*A/f/d*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c*ln(d*tan(f*x+e)
+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+B
/f*(2*(c+d*tan(f*x+e))^(1/2)-1/4*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*ln(d*tan(f*
x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2
)))+(-(c^2+d^2)^(1/2)+c)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f
*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))
+1/4*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)
)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))+(-(c^2+d^2)^(1/2)+c)/(2*(
c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)-(2*(c^2+d^2)^(1
/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)))+C*(2/3/f/d*(c+d*tan(f*x+e)
)^(3/2)+1/4/f/d*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*ln(d*tan(...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2588 vs. $2(123) = 246$.

Time = 0.19 (sec) , antiderivative size = 2588, normalized size of antiderivative = 16.70

$$\int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```

integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algori
thm="fricas")

```

output

```

-1/6*(3*d*f*sqrt(-(f^2*sqrt(-(4*(A^2*B^2 - 2*A*B^2*C + B^2*C^2)*c^2 + 4*(A
^3*B - A*B^3 + 3*A*B*C^2 - B*C^3 - (3*A^2*B - B^3)*C)*c*d + (A^4 - 2*A^2*B
^2 + B^4 - 4*A*C^3 + C^4 + 2*(3*A^2 - B^2)*C^2 - 4*(A^3 - A*B^2)*C)*d^2)/f
^4) + (A^2 - B^2 - 2*A*C + C^2)*c - 2*(A*B - B*C)*d)/f^2)*log((2*(A^3*B +
A*B^3 + 3*A*B*C^2 - B*C^3 - (3*A^2*B + B^3)*C)*c + (A^4 - B^4 - 4*A^3*C +
6*A^2*C^2 - 4*A*C^3 + C^4)*d)*sqrt(d*tan(f*x + e) + c) + ((A - C)*f^3*sqrt
(-(4*(A^2*B^2 - 2*A*B^2*C + B^2*C^2)*c^2 + 4*(A^3*B - A*B^3 + 3*A*B*C^2 -
B*C^3 - (3*A^2*B - B^3)*C)*c*d + (A^4 - 2*A^2*B^2 + B^4 - 4*A*C^3 + C^4 +
2*(3*A^2 - B^2)*C^2 - 4*(A^3 - A*B^2)*C)*d^2)/f^4) + (2*(A*B^2 - B^2*C)*c
+ (A^2*B - B^3 - 2*A*B*C + B*C^2)*d)*f)*sqrt(-(f^2*sqrt(-(4*(A^2*B^2 - 2*A
*B^2*C + B^2*C^2)*c^2 + 4*(A^3*B - A*B^3 + 3*A*B*C^2 - B*C^3 - (3*A^2*B -
B^3)*C)*c*d + (A^4 - 2*A^2*B^2 + B^4 - 4*A*C^3 + C^4 + 2*(3*A^2 - B^2)*C^2
- 4*(A^3 - A*B^2)*C)*d^2)/f^4) + (A^2 - B^2 - 2*A*C + C^2)*c - 2*(A*B - B
*C)*d)/f^2)) - 3*d*f*sqrt(-(f^2*sqrt(-(4*(A^2*B^2 - 2*A*B^2*C + B^2*C^2)*c
^2 + 4*(A^3*B - A*B^3 + 3*A*B*C^2 - B*C^3 - (3*A^2*B - B^3)*C)*c*d + (A^4
- 2*A^2*B^2 + B^4 - 4*A*C^3 + C^4 + 2*(3*A^2 - B^2)*C^2 - 4*(A^3 - A*B^2)*
C)*d^2)/f^4) + (A^2 - B^2 - 2*A*C + C^2)*c - 2*(A*B - B*C)*d)/f^2)*log((2*
(A^3*B + A*B^3 + 3*A*B*C^2 - B*C^3 - (3*A^2*B + B^3)*C)*c + (A^4 - B^4 - 4
*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)*d)*sqrt(d*tan(f*x + e) + c) - ((A - C)
*f^3*sqrt(-(4*(A^2*B^2 - 2*A*B^2*C + B^2*C^2)*c^2 + 4*(A^3*B - A*B^3 + ...

```

Sympy [F]

$$\begin{aligned}
 & \int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 &= \int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx
 \end{aligned}$$

input

```
integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)
, x)
```

Maxima [F]

$$\int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan^2(fx + e) + B \tan(fx + e) + A) \sqrt{d \tan(fx + e) + c} dx$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(d*tan(f*x + e) + c), x)`

Giac [F(-2)]

Exception generated.

$$\int \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \text{Exception raised: TypeError}$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{1, [0,14,5]%%}+%%{6, [0,12,5]%%}+%%{15, [0,10,5]%%}+%%{20, [0`

input `int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int(sqrt(tan(e + f*x)*d + c),x)*a + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*b`

3.94 $\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$

Optimal result	1057
Mathematica [A] (verified)	1058
Rubi [A] (warning: unable to verify)	1058
Maple [B] (verified)	1063
Fricas [F(-1)]	1064
Sympy [F]	1064
Maxima [F(-2)]	1064
Giac [F(-2)]	1065
Mupad [B] (verification not implemented)	1065
Reduce [F]	1066

Optimal result

Integrand size = 47, antiderivative size = 234

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$$

$$= -\frac{(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a-ib)f}$$

$$+ \frac{(iA-B-iC)\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a+ib)f}$$

$$- \frac{2(Ab^2-a(bB-aC))\sqrt{bc-ad}\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{b^{3/2}(a^2+b^2)f}$$

$$+ \frac{2C\sqrt{c+d \tan(e+fx)}}{bf}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(
a-I*b)/f+(I*A-B-I*C)*(c+I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(
1/2))/(a+I*b)/f-2*(A*b^2-a*(B*b-C*a))*(-a*d+b*c)^(1/2)*arctanh(b^(1/2)*(c
+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/b^(3/2)/(a^2+b^2)/f+2*C*(c+d*tan(f*
x+e))^(1/2)/b/f
    
```

Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \frac{b^{3/2}(-ia + b)(A - iB - C)\sqrt{c - id} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c - id}}\right) + b^{3/2}(ia + b)(A + iB - C)\sqrt{c + id} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c + id}}\right)}{b^2}$$

input

```
Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2
))/(a + b*Tan[e + f*x]),x]
```

output

```
(b^(3/2)*((-I)*a + b)*(A - I*B - C)*Sqrt[c - I*d]*ArcTanh[Sqrt[c + d*Tan[e
+ f*x]]/Sqrt[c - I*d]] + b^(3/2)*(I*a + b)*(A + I*B - C)*Sqrt[c + I*d]*Ar
cTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]] - 2*(A*b^2 + a*(-(b*B) + a*C
))*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a
*d]] + 2*Sqrt[b]*(a^2 + b^2)*C*Sqrt[c + d*Tan[e + f*x]]/(b^(3/2)*(a^2 + b
^2)*f)
```

Rubi [A] (warning: unable to verify)

Time = 3.00 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.96, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.319$, Rules used = {3042, 4130, 27, 3042, 4136, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan(e + fx)^2)}{a + b \tan(e + fx)} dx$$

↓ 4130

$$\frac{2 \int \frac{(bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd}{2(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{b} + \frac{2C\sqrt{c+d \tan(e+fx)}}{bf}$$

↓ 27

$$\frac{\int \frac{(bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{b} + \frac{2C\sqrt{c+d \tan(e+fx)}}{bf}$$

↓ 3042

$$\frac{\int \frac{(bcC-adC+bBd) \tan(e+fx)^2+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{b} + \frac{2C\sqrt{c+d \tan(e+fx)}}{bf}$$

↓ 4136

$$\frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan^2(e+fx)+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{\int \frac{b(Bc+b(A-C)d+a(Ac-Cc-Bd))-b(Abc-aBc-bCc-aAd-bBd+aCd) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2}$$

$$\frac{2C\sqrt{c+d \tan(e+fx)}}{bf}$$

↓ 3042

$$\frac{\int \frac{b(Bc+b(A-C)d+a(Ac-Cc-Bd))-b(Abc-aBc-bCc-aAd-bBd+aCd) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2}$$

$$\frac{2C\sqrt{c+d \tan(e+fx)}}{bf}$$

↓ 4022

$$\frac{2C\sqrt{c+d \tan(e+fx)}}{bf} +$$

$$\frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{\frac{1}{2}b(a-ib)(c+id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}b(a+ib)(c-id)(A-iB-C)}{a^2+b^2}$$

$$\frac{2C\sqrt{c+d \tan(e+fx)}}{bf} +$$

$$\frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{\frac{1}{2}b(a-ib)(c+id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}b(a+ib)(c-id)(A-iB-C)}{a^2+b^2}$$

↓ 4020

$$\frac{\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}} dx}{a^2+b^2} + \frac{ib(a+ib)(c-id)(A-iB-C) \int -\frac{1}{(1-i\tan(e+fx))\sqrt{c+d\tan(e+fx)}} d(i\tan(e+fx))}{2f} - \frac{ib(a+ib)(c-id)(A-iB-C)}{a^2+b^2}}{b}$$

25

$$\frac{\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}} dx}{a^2+b^2} + \frac{ib(a-ib)(c+id)(A+iB-C) \int \frac{1}{(i\tan(e+fx)+1)\sqrt{c+d\tan(e+fx)}} d(-i\tan(e+fx))}{2f} - \frac{ib(a-ib)(c+id)(A+iB-C)}{a^2+b^2}}{b}$$

73

$$\frac{\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}} dx}{a^2+b^2} + \frac{b(a-ib)(c+id)(A+iB-C) \int -\frac{1}{i\tan^2(e+fx) - \frac{ic}{d} + 1} d\sqrt{c+d\tan(e+fx)}}{df} - \frac{b(a+ib)(c-id)(A-iB-C)}{a^2+b^2}}{b}$$

221

$$\frac{\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}} dx}{a^2+b^2} + \frac{b(a+ib)\sqrt{c-id}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{b(a-ib)\sqrt{c+id}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f}}{a^2+b^2}}$$

4117

$$\frac{\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{(bc-ad)(Ab^2-a(bB-aC)) \int \frac{1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}} d\tan(e+fx)}{f(a^2+b^2)} + \frac{b(a+ib)\sqrt{c-id}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{b(a-ib)\sqrt{c+id}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f}}{a^2+b^2}}$$

73

$$\frac{\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2(bc-ad)(Ab^2-a(bB-aC)) \int \frac{1}{a + \frac{b(c+d\tan(e+fx))}{d} - \frac{bc}{d}} d\sqrt{c+d\tan(e+fx)}}{df(a^2+b^2)} + \frac{b(a+ib)\sqrt{c-id}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{b(a-ib)\sqrt{c+id}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f}}{a^2+b^2}}$$

221

$$\frac{2C\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2\sqrt{bc-ad}(Ab^2-a(bB-aC))\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d\tan(e+fx)}}{\sqrt{bc-ad}}\right)}{\sqrt{b}f(a^2+b^2)} + \frac{b(a+ib)\sqrt{c-id}(A-iB-C)\operatorname{arctan}\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{b(a-ib)\sqrt{c+id}(A+iB-C)\operatorname{arctan}\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f}$$

input `Int[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]`

output `((((a + I*b)*b*(A - I*B - C)*Sqrt[c - I*d]*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + ((a - I*b)*b*(A + I*B - C)*Sqrt[c + I*d]*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f)/(a^2 + b^2) - (2*(A*b^2 - a*(b*B - a*C))*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/(Sqrt[b]*(a^2 + b^2)*f))/b + (2*C*Sqrt[c + d*Tan[e + f*x]])/(b*f)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^(m)/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^(m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^(m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4117

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^(m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^(m*(c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3575 vs. $2(200) = 400$.

Time = 0.21 (sec) , antiderivative size = 3576, normalized size of antiderivative = 15.28

method	result	size
derivativelimit	Expression too large to display	3576
default	Expression too large to display	3576

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

output

```
-2/f*b/(a^2+b^2)/((a*d-b*c)*b)^(1/2)*arctan(b*(c+d*tan(f*x+e))^(1/2)/((a*d-b*c)*b)^(1/2))*A*a*d+1/4/f/(a^2+b^2)/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*a-1/4/f/(a^2+b^2)/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*a*c+1/4/f/(a^2+b^2)/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*a-1/4/f/(a^2+b^2)/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*A*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*a*c+1/4/f/(a^2+b^2)/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*b-1/4/f/(a^2+b^2)/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*B*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*b*c-1/4/f/(a^2+b^2)/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*a+1/4/f/(a^2+b^2)/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*C*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*a*c-2/f*b/(a^2+b^2)/((a*d-b*c)*b)^(1/2)*arctan(b*(c+d*tan(f*x+e))^(1/2)/((a*d-b*c)*b)^(1/2))*B*a*c-2/f/b/(a^2+b^2)/((a*d-b*c)*b)^(1/2)*arctan(b*(c+d*tan(f*x+e))^(1/2)/((a*d-b*c)*b)^(1/2))...
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\begin{aligned} & \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\ &= \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \end{aligned}$$

input `integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)),x)`

output `Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x)), x)`

Maxima [F(-2)]

Exception generated.

$$\begin{aligned} & \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\ &= \text{Exception raised: ValueError} \end{aligned}$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail

Giac [**F(-2)**]

Exception generated.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

= Exception raised: TypeError

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [**B**] (verification not implemented)

Time = 30.40 (sec) , antiderivative size = 62245, normalized size of antiderivative = 266.00

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x)),x)`

output

```
atan(((((((32*(4*C*a*b^8*d^11*f^4 - 4*C*b^9*c*d^10*f^4 + 8*C*a^3*b^6*d^11*
f^4 + 4*C*a^5*b^4*d^11*f^4 - 4*C*b^9*c^3*d^8*f^4 + 4*C*a*b^8*c^2*d^9*f^4 -
8*C*a^2*b^7*c*d^10*f^4 - 4*C*a^4*b^5*c*d^10*f^4 - 8*C*a^2*b^7*c^3*d^8*f^4
+ 8*C*a^3*b^6*c^2*d^9*f^4 - 4*C*a^4*b^5*c^3*d^8*f^4 + 4*C*a^5*b^4*c^2*d^9
*f^4)))/(b*f^5) - (32*(c + d*tan(e + f*x))^(1/2)*(((8*C^2*a^2*c*f^2 - 8*C^
2*b^2*c*f^2 + 16*C^2*a*b*d*f^2)^2/4 - (C^4*c^2 + C^4*d^2)*(16*a^4*f^4 + 16
*b^4*f^4 + 32*a^2*b^2*f^4))^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 - 8*
C^2*a*b*d*f^2)/(16*(a^4*f^4 + b^4*f^4 + 2*a^2*b^2*f^4)))^(1/2)*(16*b^10*d^
10*f^4 + 16*a^2*b^8*d^10*f^4 - 16*a^4*b^6*d^10*f^4 - 16*a^6*b^4*d^10*f^4 +
24*b^10*c^2*d^8*f^4 + 40*a^2*b^8*c^2*d^8*f^4 + 8*a^4*b^6*c^2*d^8*f^4 - 8*
a^6*b^4*c^2*d^8*f^4 + 8*a*b^9*c*d^9*f^4 + 24*a^3*b^7*c*d^9*f^4 + 24*a^5*b^
5*c*d^9*f^4 + 8*a^7*b^3*c*d^9*f^4))/(b*f^4))*(((8*C^2*a^2*c*f^2 - 8*C^2*b
^2*c*f^2 + 16*C^2*a*b*d*f^2)^2/4 - (C^4*c^2 + C^4*d^2)*(16*a^4*f^4 + 16*b^
4*f^4 + 32*a^2*b^2*f^4))^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 - 8*C^2
*a*b*d*f^2)/(16*(a^4*f^4 + b^4*f^4 + 2*a^2*b^2*f^4)))^(1/2) - (32*(c + d*t
an(e + f*x))^(1/2)*(14*C^2*a*b^7*d^11*f^2 - 2*C^2*a^5*b^3*d^11*f^2 - 10*C^
2*b^8*c^3*d^8*f^2 - 4*C^2*a^3*b^5*d^11*f^2 - 16*C^2*a^7*b*d^11*f^2 + 8*C^2
*a^8*c*d^10*f^2 - 6*C^2*b^8*c*d^10*f^2 + 18*C^2*a*b^7*c^2*d^9*f^2 + 12*C^2
*a^2*b^6*c*d^10*f^2 + 2*C^2*a^4*b^4*c*d^10*f^2 + 24*C^2*a^6*b^2*c*d^10*f^2
- 16*C^2*a^7*b*c^2*d^9*f^2 + 4*C^2*a^2*b^6*c^3*d^8*f^2 + 4*C^2*a^3*b^5...
```

Reduce [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

$$= \left(\int \frac{\sqrt{d \tan(fx + e) + c}}{\tan(fx + e) b + a} dx \right) a + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) c$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) b$$

input

```
int(((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)*b + a),x)*a + int((sqrt(tan(e +
f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*c + int((sqrt(tan(e
+ f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*b
```


3.95
$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$$

Optimal result	1068
Mathematica [A] (verified)	1069
Rubi [A] (warning: unable to verify)	1069
Maple [B] (verified)	1075
Fricas [F(-1)]	1075
Sympy [F]	1075
Maxima [F(-2)]	1076
Giac [F(-2)]	1076
Mupad [B] (verification not implemented)	1077
Reduce [F]	1078

Optimal result

Integrand size = 47, antiderivative size = 317

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$$

$$= -\frac{(iA+B-iC)\sqrt{c-id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a-ib)^2 f}$$

$$-\frac{(B-i(A-C))\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a+ib)^2 f}$$

$$-\frac{(a^3 b B d+a^4 C d+b^4(2 B c+A d)+a b^3(4 A c-4 c C-3 B d)-a^2 b^2(2 B c+3 A d-5 C d)) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{c}}{\sqrt{a+b \tan(e+fx)}}\right)}{b^{3/2}(a^2+b^2)^2 \sqrt{bc-ad} f}$$

$$-\frac{(A b^2-a(b B-a C)) \sqrt{c+d \tan(e+fx)}}{b(a^2+b^2) f(a+b \tan(e+fx))}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(
a-I*b)^2/f-(B-I*(A-C))*(c+I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d
)^(1/2))/(a+I*b)^2/f-(a^3*b*B*d+a^4*C*d+b^4*(A*d+2*B*c)+a*b^3*(4*A*c-3*B*d
-4*C*c)-a^2*b^2*(3*A*d+2*B*c-5*C*d))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2
))/(-a*d+b*c)^(1/2)/b^(3/2)/(a^2+b^2)^2/(-a*d+b*c)^(1/2)/f-(A*b^2-a*(B*b-C
*a))*(c+d*tan(f*x+e))^(1/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))
    
```

Mathematica [A] (verified)

Time = 4.31 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

$$= \frac{2 \left(\frac{ib \left(-(a+ib)^2(A-iB-C)\sqrt{c-id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right) + (a-ib)^2(A+iB-C)\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right) \right)}{2f} - \frac{(a^3bBd+a^4Cd+b^4(2Bc+d^2))}{(a^2+b^2)^2} \right)}{(a^2+b^2)^2}$$

input

```
Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2
))/(a + b*Tan[e + f*x])^2,x]
```

output

```
(2*(((I/2)*b*(-((a + I*b)^2*(A - I*B - C)*Sqrt[c - I*d]*ArcTanh[Sqrt[c +
d*Tan[e + f*x]]/Sqrt[c - I*d]]) + (a - I*b)^2*(A + I*B - C)*Sqrt[c + I*d]*
ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]]))/f - ((a^3*b*B*d + a^4*C*
d + b^4*(2*B*c + A*d) + a*b^3*(4*A*c - 4*c*C - 3*B*d) + a^2*b^2*(-2*B*c -
3*A*d + 5*C*d))*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]
])/(2*Sqrt[b]*Sqrt[b*c - a*d]*f))/(a^2 + b^2)^2 - (C*Sqrt[c + d*Tan[e + f
x]])/(f*(a + b*Tan[e + f*x])) + (((-A*b^2) + a*b*B + a^2*C + 2*b^2*C)*Sqrt
[c + d*Tan[e + f*x]])/(2*(a^2 + b^2)*f*(a + b*Tan[e + f*x])))/b
```

Rubi [A] (warning: unable to verify)

Time = 3.87 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.03, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.340$, Rules used = {3042, 4128, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan(e+fx)^2)}{(a+b \tan(e+fx))^2} dx$$

↓ 4128

$$\int \frac{-((-Ca^2-bBa+Ab^2-2b^2C)d \tan^2(e+fx))-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+2(bB-aC)\left(bc-\frac{ad}{2}\right)+2Ab\left(ac+\frac{bd}{2}\right)}{2(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}} \frac{1}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 27

$$\int \frac{-((-Ca^2-bBa+Ab^2-2b^2C)d \tan^2(e+fx))-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(2bc-ad)+Ab(2ac+bd)}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}} \frac{1}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\int \frac{-((-Ca^2-bBa+Ab^2-2b^2C)d \tan^2(e+fx))-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(2bc-ad)+Ab(2ac+bd)}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}} \frac{1}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 4136

$$\int \frac{2\left(b\left((Ac-Cc-Bd)a^2+2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd)\right)-b\left(-\left((Bc+(A-C)d)a^2\right)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)\right) \tan(e+fx)\right)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{(a^4Cd+a^3b^2)}{a^2+b^2}$$

$$\frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}} \frac{1}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 27

$$2 \int \frac{b\left((Ac-Cc-Bd)a^2+2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd)\right)-b\left(-\left((Bc+(A-C)d)a^2\right)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)\right) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{(a^4Cd+a^3b^2)}{a^2+b^2}$$

$$\frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}} \frac{1}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\frac{2 \int \frac{b((Ac-Cc-Bd)a^2+2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd))-b(-(Bc+(A-C)d)a^2)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)}{\sqrt{c+d \tan(e+fx)}} \tan(e+fx) dx}{a^2+b^2} + \frac{(a^4Cd+a^3bB}{2b(a^2+b^2)}$$

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

4022

$$- \frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} +$$

$$\frac{(a^4Cd+a^3bBd-a^2b^2(3Ad+2Bc-5Cd)+ab^3(4Ac-3Bd-4cC)+b^4(Ad+2Bc)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{2\left(\frac{1}{2}b(a-ib)^2(c+id)(A+iB}\right)}{2b(a^2+b^2)}$$

3042

$$- \frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} +$$

$$\frac{(a^4Cd+a^3bBd-a^2b^2(3Ad+2Bc-5Cd)+ab^3(4Ac-3Bd-4cC)+b^4(Ad+2Bc)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{2\left(\frac{1}{2}b(a-ib)^2(c+id)(A+iB}\right)}{2b(a^2+b^2)}$$

4020

$$- \frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} +$$

$$\frac{(a^4Cd+a^3bBd-a^2b^2(3Ad+2Bc-5Cd)+ab^3(4Ac-3Bd-4cC)+b^4(Ad+2Bc)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{2\left(\frac{ib(a+ib)^2(c-id)(A-iB}\right)}{2b(a^2+b^2)}$$

25

$$- \frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} +$$

$$\frac{(a^4Cd+a^3bBd-a^2b^2(3Ad+2Bc-5Cd)+ab^3(4Ac-3Bd-4cC)+b^4(Ad+2Bc)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} + \frac{2\left(\frac{ib(a-ib)^2(c+id)(A+iB}\right)}{2b(a^2+b^2)}$$

73

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{(a^4Cd + a^3bBd - a^2b^2(3Ad + 2Bc - 5Cd) + ab^3(4Ac - 3Bd - 4cC) + b^4(Ad + 2Bc)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \frac{2 \left(\frac{b(a-ib)^2(c+id)(A+iB-c)}{f} \right)}{2b(a^2 + b^2)}$$

221

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{(a^4Cd + a^3bBd - a^2b^2(3Ad + 2Bc - 5Cd) + ab^3(4Ac - 3Bd - 4cC) + b^4(Ad + 2Bc)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \frac{2 \left(\frac{b(a-ib)^2\sqrt{c+id}(A+iB-c)}{f} \right)}{2b(a^2 + b^2)}$$

4117

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{(a^4Cd + a^3bBd - a^2b^2(3Ad + 2Bc - 5Cd) + ab^3(4Ac - 3Bd - 4cC) + b^4(Ad + 2Bc)) \int \frac{1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d \tan(e+fx)}{f(a^2 + b^2)} + \frac{2 \left(\frac{b(a-ib)^2\sqrt{c+id}(A+iB-c)}{f} \right)}{2b(a^2 + b^2)}$$

73

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{2(a^4Cd + a^3bBd - a^2b^2(3Ad + 2Bc - 5Cd) + ab^3(4Ac - 3Bd - 4cC) + b^4(Ad + 2Bc)) \int \frac{1}{a + \frac{b(c+d \tan(e+fx))}{d} - \frac{bc}{d}} d \sqrt{c+d \tan(e+fx)}}{df(a^2 + b^2)} + \frac{2 \left(\frac{b(a-ib)^2\sqrt{c+id}(A+iB-c)}{f} \right)}{2b(a^2 + b^2)}$$

221

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{2(a^4Cd + a^3bBd - a^2b^2(3Ad + 2Bc - 5Cd) + ab^3(4Ac - 3Bd - 4cC) + b^4(Ad + 2Bc)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{\sqrt{b}f(a^2 + b^2)\sqrt{bc-ad}} + \frac{2 \left(\frac{b(a-ib)^2\sqrt{c+id}(A+iB-c)}{f} \right)}{2b(a^2 + b^2)}$$

input

```
Int[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^2,x]
```

output
$$\frac{((2*((a + I*b)^2*b*(A - I*B - C)*\text{Sqrt}[c - I*d]*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]])/f + ((a - I*b)^2*b*(A + I*B - C)*\text{Sqrt}[c + I*d]*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c + I*d]])/f)/(a^2 + b^2) - (2*(a^3*b*B*d + a^4*C*d + b^4*(2*B*c + A*d) + a*b^3*(4*A*c - 4*c*C - 3*B*d) - a^2*b^2*(2*B*c + 3*A*d - 5*C*d))*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/\text{Sqrt}[b*c - a*d]])/(\text{Sqrt}[b]*(a^2 + b^2)*\text{Sqrt}[b*c - a*d]*f))/(2*b*(a^2 + b^2)) - ((A*b^2 - a*(b*B - a*C))*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/(b*(a^2 + b^2)*f*(a + b*\text{Tan}[e + f*x]))$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 27
$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] \text{ ; FreeQ}[b, x]$$

rule 73
$$\text{Int}[(a_.) + (b_.)*(x_)^(m_)*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)*(c - a*(d/b) + d*(x^p/b))}^n, x], x, (a + b*x)^{(1/p)}, x]] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 221
$$\text{Int}[(a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$$

rule 3042
$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4020
$$\text{Int}[(a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \quad \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\text{Tan}[e + f*x]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$$

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4117

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5777 vs. $2(284) = 568$.

Time = 0.21 (sec) , antiderivative size = 5778, normalized size of antiderivative = 18.23

method	result	size
derivativeldivides	Expression too large to display	5778
default	Expression too large to display	5778

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")
```

output

```
Timed out
```

Sympy [F]

$$\begin{aligned} & \int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx \\ &= \int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx \end{aligned}$$

input `integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**2,x)`

output `Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x)**2, x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

= Exception raised: ValueError

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

= Exception raised: TypeError

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,14,6]}%%+%%{6,[0,12,6]}%%+%%{15,[0,10,6]}%%+
%%{20,[0
```

Mupad [B] (verification not implemented)

Time = 39.99 (sec) , antiderivative size = 138318, normalized size of antiderivative = 436.33

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x))^2,x)
```

output

```
atan((((8*(156*B^3*a^2*b^9*d^12*f^2 - 16*B^3*a^4*b^7*d^12*f^2 - 120*B^3*a
^6*b^5*d^12*f^2 + 48*B^3*a^8*b^3*d^12*f^2 + 12*B^3*b^11*c^2*d^10*f^2 + 12*
B^3*b^11*c^4*d^8*f^2 - 4*B^3*a^10*b*d^12*f^2 - 124*B^3*a*b^10*c*d^11*f^2 -
124*B^3*a*b^10*c^3*d^9*f^2 + 224*B^3*a^3*b^8*c*d^11*f^2 + 200*B^3*a^5*b^6
*c*d^11*f^2 - 128*B^3*a^7*b^4*c*d^11*f^2 + 20*B^3*a^9*b^2*c*d^11*f^2 - 4*B
^3*a^10*b*c^2*d^10*f^2 + 44*B^3*a^2*b^9*c^2*d^10*f^2 - 112*B^3*a^2*b^9*c^4
*d^8*f^2 + 224*B^3*a^3*b^8*c^3*d^9*f^2 - 40*B^3*a^4*b^7*c^2*d^10*f^2 - 24*
B^3*a^4*b^7*c^4*d^8*f^2 + 200*B^3*a^5*b^6*c^3*d^9*f^2 - 40*B^3*a^6*b^5*c^2
*d^10*f^2 + 80*B^3*a^6*b^5*c^4*d^8*f^2 - 128*B^3*a^7*b^4*c^3*d^9*f^2 + 28*
B^3*a^8*b^3*c^2*d^10*f^2 - 20*B^3*a^8*b^3*c^4*d^8*f^2 + 20*B^3*a^9*b^2*c^3
*d^9*f^2)))/(a^8*f^5 + b^8*f^5 + 4*a^2*b^6*f^5 + 6*a^4*b^4*f^5 + 4*a^6*b^2*
f^5) + (((8*(80*B*a*b^14*d^11*f^4 - 48*B*b^15*c*d^10*f^4 + 384*B*a^3*b^12*
d^11*f^4 + 720*B*a^5*b^10*d^11*f^4 + 640*B*a^7*b^8*d^11*f^4 + 240*B*a^9*b^
6*d^11*f^4 - 16*B*a^13*b^2*d^11*f^4 - 48*B*b^15*c^3*d^8*f^4 + 80*B*a*b^14*
c^2*d^9*f^4 - 224*B*a^2*b^13*c*d^10*f^4 - 400*B*a^4*b^11*c*d^10*f^4 - 320*
B*a^6*b^9*c*d^10*f^4 - 80*B*a^8*b^7*c*d^10*f^4 + 32*B*a^10*b^5*c*d^10*f^4
+ 16*B*a^12*b^3*c*d^10*f^4 - 224*B*a^2*b^13*c^3*d^8*f^4 + 384*B*a^3*b^12*c
^2*d^9*f^4 - 400*B*a^4*b^11*c^3*d^8*f^4 + 720*B*a^5*b^10*c^2*d^9*f^4 - 320
*B*a^6*b^9*c^3*d^8*f^4 + 640*B*a^7*b^8*c^2*d^9*f^4 - 80*B*a^8*b^7*c^3*d^8*
f^4 + 240*B*a^9*b^6*c^2*d^9*f^4 + 32*B*a^10*b^5*c^3*d^8*f^4 + 16*B*a^12...
```

Reduce [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^2,x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2*d + 4*sqrt(tan(e + f*x)*d + c)*a*b*c +
 2*sqrt(tan(e + f*x)*d + c)*b**2*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e +
 f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e
 + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*a**3*b*d**
 2*f - 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f
*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*
x)*a*b*c + a**2*c),x)*tan(e + f*x)*a**2*b**2*c*d*f + 2*int(sqrt(tan(e + f*
x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)
**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e
 + f*x)*a*b**3*c**2*f + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2
*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*
d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a**4*d**2*f - 3*int(sqrt(tan(e + f*x)
)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)*
**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a**3*b
*c*d*f + 2*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e
 + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e +
 f*x)*a*b*c + a**2*c),x)*a**2*b**2*c**2*f - int(sqrt(tan(e + f*x)*d + c)/(
tan(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)*
**2*a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**
 2 + tan(e + f*x)*a**3*d**2 - 4*tan(e + f*x)*a*b**2*c**2 + a**3*c*d - 2*...
```

3.96
$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

Optimal result	1079
Mathematica [B] (verified)	1080
Rubi [A] (warning: unable to verify)	1081
Maple [B] (verified)	1088
Fricas [F(-1)]	1089
Sympy [F]	1089
Maxima [F(-2)]	1089
Giac [F(-2)]	1090
Mupad [F(-1)]	1090
Reduce [F]	1091

Optimal result

Integrand size = 47, antiderivative size = 543

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

$$= -\frac{(A-iB-C)\sqrt{c-id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(ia+b)^3 f}$$

$$+ \frac{(A+iB-C)\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(ia-b)^3 f}$$

$$+ \frac{(3a^5 b B d^2 + a^6 C d^2 - 3a^4 b^2 d(4Bc + 5Ad - 6Cd) - 3a^2 b^4(8Ac^2 - 8c^2 C - 16Bcd - 6Ad^2 + 5Cd^2) + 2a^3 b^2 C d)}{4b(a^2 + b^2)^2 (bc - ad)f(a + b \tan(e + fx))}$$

$$- \frac{(Ab^2 - a(bB - aC))\sqrt{c+d \tan(e+fx)}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

$$- \frac{(3a^3 b B d + a^4 C d + b^4(4Bc + Ad) + ab^3(8Ac - 8cC - 5Bd) - a^2 b^2(4Bc + 7Ad - 9Cd))\sqrt{c+d \tan(e+fx)}}{4b(a^2 + b^2)^2 (bc - ad)f(a + b \tan(e + fx))}$$

output

```

-(A-I*B-C)*(c-I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(I*
a+b)^3/f+(A+I*B-C)*(c+I*d)^(1/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1
/2))/(I*a-b)^3/f+1/4*(3*a^5*b*B*d^2+a^6*C*d^2-3*a^4*b^2*d*(5*A*d+4*B*c-6*C
*d)-3*a^2*b^4*(8*A*c^2-6*A*d^2-16*B*c*d-8*C*c^2+5*C*d^2)+2*a^3*b^3*(20*c*(
A-C)*d+B*(4*c^2-13*d^2))-3*a*b^5*(8*c*(A-C)*d+B*(8*c^2-d^2))-b^6*(4*c*(B*d
+2*C*c)-A*(8*c^2+d^2))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^
(1/2))/b^(3/2)/(a^2+b^2)^3/(-a*d+b*c)^(3/2)/f-1/2*(A*b^2-a*(B*b-C*a))*(c+d
*tan(f*x+e))^(1/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^2-1/4*(3*a^3*b*B*d+a^4*C
*d+b^4*(A*d+4*B*c)+a*b^3*(8*A*c-5*B*d-8*C*c)-a^2*b^2*(7*A*d+4*B*c-9*C*d))*
(c+d*tan(f*x+e))^(1/2)/b/(a^2+b^2)^2/(-a*d+b*c)/f/(a+b*tan(f*x+e))

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2819 vs. $2(543) = 1086$.

Time = 6.44 (sec) , antiderivative size = 2819, normalized size of antiderivative = 5.19

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

= Result too large to show

input

```

Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2
))/(a + b*Tan[e + f*x])^3,x]

```

output

```
(-2*C*sqrt[c + d*Tan[e + f*x]])/(3*b*f*(a + b*Tan[e + f*x])^2) - (2*(-1/2*
(((b^2*(-3*A*b*c + 4*b*c*C - a*C*d))/2 - a*((-3*b^2*(B*c + (A - C)*d))/2 -
(a*(b*c*C - 3*b*B*d - a*C*d))/2))*sqrt[c + d*Tan[e + f*x]]/((a^2 + b^2)*
(b*c - a*d)*f*(a + b*Tan[e + f*x])^2) - (-(I*sqrt[c - I*d]*(b*(b*c - a*
d)*((3*b*(3*A*b^2 - 3*a*b*B - a^2*C - 4*b^2*C)*d*(b*c - a*d))/4 + 3*a*b*(b
*c - a*d)*(A*b*c - a*B*c - b*c*C - a*A*d - b*B*d + a*C*d) + (3*b*(b*c - a*
d)*(a^2*C*d + b^2*(4*B*c + A*d) + a*b*(4*A*c - 4*c*C - B*d)))/4) + a*((3*(
b*c - a*d)*((b^2*d)/2 - a*(b*c - a*d))*((a^2*C*d + b^2*(4*B*c + A*d) + a*b*
(4*A*c - 4*c*C - B*d)))/4 + (-(b*c) + (a*d)/2)*((3*a*(3*A*b^2 - 3*a*b*B -
a^2*C - 4*b^2*C)*d*(b*c - a*d))/4 - 3*b^2*(b*c - a*d)*(A*b*c - a*B*c - b*c
*C - a*A*d - b*B*d + a*C*d)) - (d*((3*b^2*(b*c - a*d)*(a^2*C*d + b^2*(4*B*
c + A*d) + a*b*(4*A*c - 4*c*C - B*d)))/4 - a*((3*a*(3*A*b^2 - 3*a*b*B - a^
2*C - 4*b^2*C)*d*(b*c - a*d))/4 - 3*b^2*(b*c - a*d)*(A*b*c - a*B*c - b*c*C
- a*A*d - b*B*d + a*C*d)))/2) - I*(a*(b*c - a*d)*((3*b*(3*A*b^2 - 3*a*b*B
- a^2*C - 4*b^2*C)*d*(b*c - a*d))/4 + 3*a*b*(b*c - a*d)*(A*b*c - a*B*c -
b*c*C - a*A*d - b*B*d + a*C*d) + (3*b*(b*c - a*d)*(a^2*C*d + b^2*(4*B*c +
A*d) + a*b*(4*A*c - 4*c*C - B*d)))/4) - b*((3*(b*c - a*d)*((b^2*d)/2 - a*
(b*c - a*d))*((a^2*C*d + b^2*(4*B*c + A*d) + a*b*(4*A*c - 4*c*C - B*d)))/4
+ (-(b*c) + (a*d)/2)*((3*a*(3*A*b^2 - 3*a*b*B - a^2*C - 4*b^2*C)*d*(b*c -
a*d))/4 - 3*b^2*(b*c - a*d)*(A*b*c - a*B*c - b*c*C - a*A*d - b*B*d + a*...
```

Rubi [A] (warning: unable to verify)

Time = 7.15 (sec) , antiderivative size = 588, normalized size of antiderivative = 1.08, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.404$, Rules used = {3042, 4128, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx)^2)}{(a + b \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{-((-Ca^2-3bBa+3Ab^2-4b^2C)d \tan^2(e+fx))-4b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+2(bB-aC)\left(2bc-\frac{ad}{2}\right)+Ab(4ac+bd)}{2(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC)) \sqrt{c+d \tan(e+fx)}} \frac{1}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 27

$$\int \frac{-((-Ca^2-3bBa+3Ab^2-4b^2C)d \tan^2(e+fx))-4b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(4bc-ad)+Ab(4ac+bd)}{(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{4b(a^2+b^2)}{(Ab^2-a(bB-aC)) \sqrt{c+d \tan(e+fx)}} \frac{1}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{-((-Ca^2-3bBa+3Ab^2-4b^2C)d \tan(e+fx)^2)-4b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(4bc-ad)+Ab(4ac+bd)}{(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{4b(a^2+b^2)}{(Ab^2-a(bB-aC)) \sqrt{c+d \tan(e+fx)}} \frac{1}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 4132

$$\int \frac{-d(Cda^4+3bBda^3-b^2(4Bc+7Ad-9Cd)a^2+b^3(8Ac-8Cc-5Bd)a+b^4(4Bc+Ad)) \tan^2(e+fx)-8b(bc-ad)\left(-((Bc+(A-C)d)a^2)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)\right)}{2(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}$$

$$\frac{(Ab^2-a(bB-aC)) \sqrt{c+d \tan(e+fx)}}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 27

$$\int \frac{-d(Cda^4+3bBda^3-b^2(4Bc+7Ad-9Cd)a^2+b^3(8Ac-8Cc-5Bd)a+b^4(4Bc+Ad)) \tan^2(e+fx)-8b(bc-ad)\left(-((Bc+(A-C)d)a^2)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)\right)}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} \frac{1}{2(a^2+b^2)}$$

$$\frac{(Ab^2-a(bB-aC)) \sqrt{c+d \tan(e+fx)}}{2bf(a^2+b^2)(a+b \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{-d(Cda^4+3bBda^3-b^2(4Bc+7Ad-9Cd)a^2+b^3(8Ac-8Cc-5Bd)a+b^4(4Bc+Ad)) \tan(e+fx)^2-8b(bc-ad)\left(-((Bc+(A-C)d)a^2)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)\right) \sqrt{c+d \tan(e+fx)}}{2(a^2+b^2)(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

4136

$$8 \int \frac{b(bc-ad)\left((Ac-Cc-Bd)a^3+3b(Bc+(A-C)d)a^2-3b^2(Ac-Cc-Bd)a-b^3(Bc+(A-C)d)\right)-b(bc-ad)\left(-((Bc+(A-C)d)a^3)+3b(Ac-Cc-Bd)a^2+3b^2(Bc+(A-C)d)\right) \sqrt{c+d \tan(e+fx)}}{\sqrt{c+d \tan(e+fx)} a^2+b^2}$$

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

27

$$8 \int \frac{b(bc-ad)\left((Ac-Cc-Bd)a^3+3b(Bc+(A-C)d)a^2-3b^2(Ac-Cc-Bd)a-b^3(Bc+(A-C)d)\right)-b(bc-ad)\left(-((Bc+(A-C)d)a^3)+3b(Ac-Cc-Bd)a^2+3b^2(Bc+(A-C)d)\right) \sqrt{c+d \tan(e+fx)}}{\sqrt{c+d \tan(e+fx)} a^2+b^2}$$

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

3042

$$8 \int \frac{b(bc-ad)\left((Ac-Cc-Bd)a^3+3b(Bc+(A-C)d)a^2-3b^2(Ac-Cc-Bd)a-b^3(Bc+(A-C)d)\right)-b(bc-ad)\left(-((Bc+(A-C)d)a^3)+3b(Ac-Cc-Bd)a^2+3b^2(Bc+(A-C)d)\right) \sqrt{c+d \tan(e+fx)}}{\sqrt{c+d \tan(e+fx)} a^2+b^2}$$

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

4022

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2} +$$

$$-\frac{\sqrt{c+d \tan(e+fx)}(a^4Cd+3a^3bBd-a^2b^2(7Ad+4Bc-9Cd)+ab^3(8Ac-5Bd-8cC)+b^4(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} + \frac{(a^6Cd^2+3a^5bBd^2-3a^4b^2d(5Ad+4Bc-6Cc)+3a^3b^3Bd^2+3a^2b^4(5Ad+4Bc-6Cc)+3ab^5(8Ac-5Bd-8cC)+b^6(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

3042

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} +$$

$$-\frac{\sqrt{c+d \tan(e+fx)}(a^4Cd+3a^3bBd-a^2b^2(7Ad+4Bc-9Cd)+ab^3(8Ac-5Bd-8cC)+b^4(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} + \frac{(a^6Cd^2+3a^5bBd^2-3a^4b^2d(5Ad+4Bc-6C))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

4020

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} +$$

$$-\frac{\sqrt{c+d \tan(e+fx)}(a^4Cd+3a^3bBd-a^2b^2(7Ad+4Bc-9Cd)+ab^3(8Ac-5Bd-8cC)+b^4(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} + \frac{(a^6Cd^2+3a^5bBd^2-3a^4b^2d(5Ad+4Bc-6C))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

25

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} +$$

$$-\frac{\sqrt{c+d \tan(e+fx)}(a^4Cd+3a^3bBd-a^2b^2(7Ad+4Bc-9Cd)+ab^3(8Ac-5Bd-8cC)+b^4(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} + \frac{(a^6Cd^2+3a^5bBd^2-3a^4b^2d(5Ad+4Bc-6C))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

73

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} +$$

$$-\frac{\sqrt{c+d \tan(e+fx)}(a^4Cd+3a^3bBd-a^2b^2(7Ad+4Bc-9Cd)+ab^3(8Ac-5Bd-8cC)+b^4(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} + \frac{(a^6Cd^2+3a^5bBd^2-3a^4b^2d(5Ad+4Bc-6C))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

221

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} +$$

$$-\frac{\sqrt{c+d \tan(e+fx)}(a^4Cd+3a^3bBd-a^2b^2(7Ad+4Bc-9Cd)+ab^3(8Ac-5Bd-8cC)+b^4(Ad+4Bc))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} + \frac{(a^6Cd^2+3a^5bBd^2-3a^4b^2d(5Ad+4Bc-6C))}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))}$$

4117

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{(a^6Cd^2 + 3a^5bBd^2 - 3a^4b^2d(5Ad + 4Bc - 6C)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} + \dots$$

73

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{2(a^6Cd^2 + 3a^5bBd^2 - 3a^4b^2d(5Ad + 4Bc - 6C)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} + \dots$$

221

$$-\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{2(a^6Cd^2 + 3a^5bBd^2 - 3a^4b^2d(5Ad + 4Bc - 6C)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} + \dots$$

input

```
Int[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

```
-1/2*((A*b^2 - a*(b*B - a*C))*Sqrt[c + d*Tan[e + f*x]])/(b*(a^2 + b^2)*f*(
a + b*Tan[e + f*x])^2) + (((8*((a + I*b)^3*b*(A - I*B - C)*Sqrt[c - I*d]*
(b*c - a*d)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + ((a - I*b)^3*b*(A + I*
B - C)*Sqrt[c + I*d]*(b*c - a*d)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f))/
(a^2 + b^2) + (2*(3*a^5*b*B*d^2 + a^6*C*d^2 - 3*a^4*b^2*d*(4*B*c + 5*A*d -
6*C*d) - 3*a^2*b^4*(8*A*c^2 - 8*c^2*C - 16*B*c*d - 6*A*d^2 + 5*C*d^2) + 2*
a^3*b^3*(20*c*(A - C)*d + B*(4*c^2 - 13*d^2)) - 3*a*b^5*(8*c*(A - C)*d + B
*(8*c^2 - d^2)) - b^6*(4*c*(2*c*C + B*d) - A*(8*c^2 + d^2)))*ArcTanh[(Sqrt
[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]/(Sqrt[b]*(a^2 + b^2)*Sqrt[
b*c - a*d]*f))/(2*(a^2 + b^2)*(b*c - a*d)) - ((3*a^3*b*B*d + a^4*C*d + b^4
*(4*B*c + A*d) + a*b^3*(8*A*c - 8*c*C - 5*B*d) - a^2*b^2*(4*B*c + 7*A*d -
9*C*d))*Sqrt[c + d*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e
+ f*x]))/(4*b*(a^2 + b^2))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4020

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4117

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 9796 vs. $2(503) = 1006$.

Time = 0.24 (sec) , antiderivative size = 9797, normalized size of antiderivative = 18.04

method	result	size
derivativedivides	Expression too large to display	9797
default	Expression too large to display	9797

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^3,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\begin{aligned} & \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx \\ &= \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx \end{aligned}$$

input

```
integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**3,x)
```

output

```
Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x)**3, x)
```

Maxima [F(-2)]

Exception generated.

$$\begin{aligned} & \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx \\ &= \text{Exception raised: ValueError} \end{aligned}$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail)

Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

= Exception raised: TypeError

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0, 19, 8]}%%}+%%{8, [0, 17, 8]}%%}+%%{28, [0, 15, 8]}%%}+%%{56, [0

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Hanged}$$

input `int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^3,x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{too large to display}$$

input `int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x)`

output `(- 2*sqrt(tan(e + f*x)*d + c)*a**2*d + 8*sqrt(tan(e + f*x)*d + c)*a*b*c + 6*sqrt(tan(e + f*x)*d + c)*b**2*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a**3*b**2*d**2*f - 5*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a**2*b**3*c*d*f + 4*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a*b**4*c**2*f + 2*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)*a**4*b*d**2*f - 10*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)*a**3*b**2*c*d*f + 8*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + ...`

3.97 $\int (a+b \tan(e+fx))^3 (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)) dx$

Optimal result	1092
Mathematica [B] (verified)	1093
Rubi [A] (warning: unable to verify)	1094
Maple [B] (verified)	1101
Fricas [F(-1)]	1101
Sympy [F]	1102
Maxima [F(-1)]	1102
Giac [F(-2)]	1103
Mupad [F(-1)]	1103
Reduce [F]	1104

Optimal result

Integrand size = 47, antiderivative size = 550

$$\begin{aligned}
 & \int (a+b \tan(e+fx))^3 (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)) \\
 & + C \tan^2(e+fx) dx = \frac{(ia+b)^3(A-iB-C)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f} \\
 & + \frac{(a+ib)^3(iA-B-iC)(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f} \\
 & + \frac{2(3a^2b(AC-cC-Bd) - b^3(AC-cC-Bd) + a^3(Bc+(A-C)d) - 3ab^2(Bc+(A-C)d)) \sqrt{c+d \tan(e+fx)}}{f} \\
 & + \frac{2(a^3B - 3ab^2B + 3a^2b(A-C) - b^3(A-C))(c+d \tan(e+fx))^{3/2}}{3f} \\
 & + \frac{2(168a^3Cd^3 - 2a^2bd^2(192cC - 847Bd) + 33ab^2d(8c^2C - 18Bcd + 63(A-C)d^2) - b^3(48c^3C - 88Bc^2d - 2b(99b(Ab+aB-bC)d^2 + 4(bc-ad)(6bcC - 11bBd - 6aCd)) \tan(e+fx)(c+d \tan(e+fx))^{5/2}}{3465d^4f} \\
 & - \frac{2(6bcC - 11bBd - 6aCd)(a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{5/2}}{99d^2f} \\
 & + \frac{2C(a+b \tan(e+fx))^3 (c+d \tan(e+fx))^{5/2}}{11df}
 \end{aligned}$$

output

```
(I*a+b)^3*(A-I*B-C)*(c-I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/f+(a+I*b)^3*(I*A-B-I*C)*(c+I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/f+2*(3*a^2*b*(A*c-B*d-C*c)-b^3*(A*c-B*d-C*c)+a^3*(B*c+(A-C)*d)-3*a*b^2*(B*c+(A-C)*d))*(c+d*tan(f*x+e))^(1/2)/f+2/3*(B*a^3-3*B*a*b^2+3*a^2*b*(A-C)-b^3*(A-C))*(c+d*tan(f*x+e))^(3/2)/f+2/3465*(168*a^3*C*d^3-2*a^2*b*d^2*(-847*B*d+192*C*c)+33*a*b^2*d*(8*c^2*C-18*B*c*d+63*(A-C)*d^2)-b^3*(48*c^3*C-88*B*c^2*d+198*c*(A-C)*d^2+693*B*d^3))*(c+d*tan(f*x+e))^(5/2)/d^4/f+2/693*b*(99*b*(A*b+B*a-C*b)*d^2+4*(-a*d+b*c)*(-11*B*b*d-6*C*a*d+6*C*b*c))*tan(f*x+e)*(c+d*tan(f*x+e))^(5/2)/d^3/f-2/99*(-11*B*b*d-6*C*a*d+6*C*b*c)*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(5/2)/d^2/f+2/11*C*(a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(5/2)/d/f
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1290 vs. $2(550) = 1100$.

Time = 6.36 (sec) , antiderivative size = 1290, normalized size of antiderivative = 2.35

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
Integrate[(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
(2*C*(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^(5/2))/(11*d*f) + (2*(((-6*b*c*C + 11*b*B*d + 6*a*C*d)*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(5/2))/(9*d*f) + (2*((b*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d))*Tan[e + f*x]*(c + d*Tan[e + f*x])^(5/2))/(14*d*f) - (2*((2*((-7*a*d*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d)))/8 + b*((-693*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3)/8 + (c*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d)))/4))*(c + d*Tan[e + f*x])^(5/2))/(5*d*f) + ((I/2)*((-7*a*d*(3*a^2*(33*A - 25*C)*d^2 + 4*b^2*c*(6*c*C - 11*B*d) - a*b*d*(48*c*C + 55*B*d)))/8 + (b*c*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d)))/4 + (7*a*d*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d)))/8 + ((7*I)/2)*d*((99*a*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2)/4 + (b*(3*a^2*(33*A - 25*C)*d^2 + 4*b^2*c*(6*c*C - 11*B*d) - a*b*d*(48*c*C + 55*B*d)))/4 - (b*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d)))/4) - b*((-693*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3)/8 + (c*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d)))/4))*((2*(c + d*Tan[e + f*x])^(3/2))/3 + (c - I*d)*((2*(c - I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]])/(-c + I*d) + 2*Sqrt[c + d*Tan[e + f*x]]))/f - ((I/2)*((-7*a*d*(3*a^2*(33*A - 25*C)*d^2 + 4*b^2*c*(6*c*C - 11*B*d) - a*b*d*(48*c*C + 55*B*d)))/8 + (b*c*(...
```

Rubi [A] (warning: unable to verify)

Time = 7.16 (sec) , antiderivative size = 566, normalized size of antiderivative = 1.03, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.468$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \frac{2 \int -\frac{1}{2}(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2} ((6bcC - 6adC - 11bBd) \tan^2(e + fx) - 11(Ab - Cb + aB)d \tan(e + fx) + 11d)}{11df} \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \quad \downarrow 27 \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \frac{\int (a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2} ((6bcC - 6adC - 11bBd) \tan^2(e + fx) - 11(Ab - Cb + aB)d \tan(e + fx) + 11d)}{11d} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \frac{\int (a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2} ((6bcC - 6adC - 11bBd) \tan(e + fx)^2 - 11(Ab - Cb + aB)d \tan(e + fx) + 11d)}{11d} \\
 & \quad \downarrow 4130 \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \frac{2 \int -\frac{1}{2}(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (4c(6cC - 11Bd)b^2 - ad(48cC + 55Bd)b + 3a^2(33A - 25C)d^2 + (99b(Ab - Cb + aB)d^2 + 4(bc - ad)(6bcC - 11d^2)))}{9d}}{11d} \\
 & \quad \downarrow 27 \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2} - \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (4c(6cC - 11Bd)b^2 - ad(48cC + 55Bd)b + 3a^2(33A - 25C)d^2 + (99b(Ab - Cb + aB)d^2 + 4(bc - ad)(6bcC - 11d^2)))}{9df}}{11d} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2} - \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (4c(6cC - 11Bd)b^2 - ad(48cC + 55Bd)b + 3a^2(33A - 25C)d^2 + (99b(Ab - Cb + aB)d^2 + 4(bc - ad)(6bcC - 11d^2)))}{9df}}{11d} \\
 & \quad \downarrow 4120 \\
 & \quad \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} \\
 & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2} (99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df}}{9df}}{11d}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ \hline & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c + d \tan(e + fx))^{3/2} (-2c(24C^2 - 44Bdc + 99(A - C)d^2)b^3 + 66acd(4cC - 9Bd)b^2 - \dots)}{9df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ \hline & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c + d \tan(e + fx))^{3/2} (-2c(24C^2 - 44Bdc + 99(A - C)d^2)b^3 + 66acd(4cC - 9Bd)b^2 - \dots)}{9df} \end{aligned}$$

$$\begin{aligned} & \downarrow 4113 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ \hline & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c + d \tan(e + fx))^{3/2} (693(Ba^3 + 3b(A - C)a^2 - 3b^2Ba - b^3(A - C))d^3 \tan(e + fx) - 69 \dots)}{9df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ \hline & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c + d \tan(e + fx))^{3/2} (693(Ba^3 + 3b(A - C)a^2 - 3b^2Ba - b^3(A - C))d^3 \tan(e + fx) - 69 \dots)}{9df} \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ \hline & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int \sqrt{c + d \tan(e + fx)} (693((Ac - Cc - Bd)a^3 - 3b(Bc + (A - C)d)a^2 - 3b^2(Ac - Cc - Bd)a - \dots))}{9df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ \hline & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int \sqrt{c + d \tan(e + fx)} (693((Ac - Cc - Bd)a^3 - 3b(Bc + (A - C)d)a^2 - 3b^2(Ac - Cc - Bd)a - \dots))}{9df} \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \int \frac{-693((Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a^3 + 3b(2c(A - C)d + B(c^2 - d^2))a^2 - 3b^2(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a + 3b^3(B(c^2 - d^2) - Cc^2))}{(a + b \tan(e + fx))^5} dx \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \int \frac{-693((Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a^3 + 3b(2c(A - C)d + B(c^2 - d^2))a^2 - 3b^2(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a + 3b^3(B(c^2 - d^2) - Cc^2))}{(a + b \tan(e + fx))^5} dx \end{aligned}$$

$$\begin{aligned} & \downarrow 4022 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2}(99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2}(99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df} \end{aligned}$$

$$\begin{aligned} & \downarrow 4020 \\ & \frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \\ & \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2}(99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df} \end{aligned}$$

\downarrow 25

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2}(99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df}$$

73

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2}(99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df}$$

221

$$\frac{2C(a + b \tan(e + fx))^3(c + d \tan(e + fx))^{5/2}}{11df} - \frac{2(-6aCd - 11bBd + 6bcC)(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e + fx)(c + d \tan(e + fx))^{5/2}(99bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 11bBd + 6bcC))}{7df}$$

input

```
Int[(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
(2*C*(a + b*Tan[e + f*x])^3*(c + d*Tan[e + f*x])^(5/2))/(11*d*f) - ((2*(6*b*c*C - 11*b*B*d - 6*a*C*d)*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(5/2))/(9*d*f) - ((2*b*(99*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 11*b*B*d - 6*a*C*d))*Tan[e + f*x]*(c + d*Tan[e + f*x])^(5/2))/(7*d*f) + ((693*(a - I*b)^3*(A - I*B - C)*(c - I*d)^(3/2)*d^3*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + (693*(a + I*b)^3*(A + I*B - C)*(c + I*d)^(3/2)*d^3*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f + (1386*d^3*(3*a^2*b*(A*c - c*C - B*d) - b^3*(A*c - c*C - B*d) + a^3*(B*c + (A - C)*d) - 3*a*b^2*(B*c + (A - C)*d))*Sqrt[c + d*Tan[e + f*x]]/f + (462*(a^3*B - 3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C))*d^3*(c + d*Tan[e + f*x])^(3/2))/f + (2*(168*a^3*C*d^3 - 2*a^2*b*d^2*(192*c*C - 847*B*d) + 33*a*b^2*d*(8*c^2*C - 18*B*c*d + 63*(A - C)*d^2) - b^3*(48*c^3*C - 88*B*c^2*d + 198*c*(A - C)*d^2 + 693*B*d^3))*(c + d*Tan[e + f*x])^(5/2))/(5*d*f)/(7*d)/(9*d)/(11*d)
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e.
) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 10951 vs. $2(507) = 1014$.

Time = 0.46 (sec) , antiderivative size = 10952, normalized size of antiderivative = 19.91

method	result	size
parts	Expression too large to display	10952
derivativedivides	Expression too large to display	11056
default	Expression too large to display	11056

input `int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input

```
integrate((a+b*tan(f*x+e))**3*(c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Integral((a + b*tan(e + f*x))**3*(c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)
```

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
Timed out
```

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%}{1, [0,34,13]%%}+%%{14, [0,32,13]%%}+%%{91, [0,30,13]%%}+%%{3`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Hanged}$$

input `int((a + b*tan(e + f*x))^3*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `\text{Hanged}`

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))^3 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) \\
& + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^4 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^6 dx \right) b^3 cd \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^5 dx \right) a b^2 cd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^5 dx \right) b^4 d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^5 dx \right) b^3 c^2 \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) a^2 bcd \\
& + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) a b^3 d \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) a b^2 c^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b^4 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a^3 cd \\
& + 6 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a^2 b^2 d \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a^2 b c^2 \\
& + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a b^3 c \\
& + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^3 b d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^3 c^2 \\
& + 6 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^2 b^2 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^4 d \\
& + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^3 b c
\end{aligned}$$

input

```
int((a+b*tan(f*x+e))^3*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**4*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**6,x)*b**3*c*d + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5,x)*a*b**2*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5,x)*b**4*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5,x)*b**3*c**2 + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a**2*b*c*d + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a*b**3*d + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a*b**2*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**4*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a**3*c*d + 6*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a**2*b**2*d + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a**2*b*c**2 + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*b**3*c + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**3*b*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**3*c**2 + 6*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*b**2*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**4*d + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**3*b*c
```

3.98 $\int (a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{3/2} (A+B \tan(e$

Optimal result	1106
Mathematica [A] (verified)	1107
Rubi [A] (warning: unable to verify)	1108
Maple [B] (verified)	1113
Fricas [B] (verification not implemented)	1114
Sympy [F]	1114
Maxima [F(-1)]	1115
Giac [F(-2)]	1115
Mupad [F(-1)]	1116
Reduce [F]	1117

Optimal result

Integrand size = 47, antiderivative size = 396

$$\begin{aligned}
 & \int (a+b \tan(e+fx))^2 (c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx)) dx = \\
 & \frac{(a-ib)^2(B+i(A-C))(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f} \\
 & + \frac{(a+ib)^2(iA-B-iC)(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f} \\
 & + \frac{2(2ab(Ac-cC-Bd)+a^2(Bc+(A-C)d)-b^2(Bc+(A-C)d)) \sqrt{c+d \tan(e+fx)}}{f} \\
 & + \frac{2(a^2B-b^2B+2ab(A-C))(c+d \tan(e+fx))^{3/2}}{3f} \\
 & + \frac{2(28a^2Cd^2-18abd(2cC-7Bd)+b^2(8c^2C-18Bcd+63(A-C)d^2))(c+d \tan(e+fx))^{5/2}}{315d^3f} \\
 & - \frac{2b(4bcC-9bBd-4aCd) \tan(e+fx)(c+d \tan(e+fx))^{5/2}}{63d^2f} \\
 & + \frac{2C(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{5/2}}{9df}
 \end{aligned}$$

output

```

-(a-I*b)^2*(B+I*(A-C))*(c-I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)
)^(1/2))/f+(a+I*b)^2*(I*A-B-I*C)*(c+I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1
/2)/(c+I*d)^(1/2))/f+2*(2*a*b*(A*c-B*d-C*c)+a^2*(B*c+(A-C)*d)-b^2*(B*c+(A-
C)*d))*(c+d*tan(f*x+e))^(1/2)/f+2/3*(B*a^2-B*b^2+2*a*b*(A-C))*(c+d*tan(f*x
+e))^(3/2)/f+2/315*(28*a^2*C*d^2-18*a*b*d*(-7*B*d+2*C*c)+b^2*(8*c^2*C-18*B
*c*d+63*(A-C)*d^2))*(c+d*tan(f*x+e))^(5/2)/d^3/f-2/63*b*(-9*B*b*d-4*C*a*d+
4*C*b*c)*tan(f*x+e)*(c+d*tan(f*x+e))^(5/2)/d^2/f+2/9*C*(a+b*tan(f*x+e))^2*
(c+d*tan(f*x+e))^(5/2)/d/f

```

Mathematica [A] (verified)

Time = 4.22 (sec) , antiderivative size = 350, normalized size of antiderivative = 0.88

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{2 \left((28a^2Cd^2 + 18abd(-2cC + 7Bd) + b^2(8c^2C - 18Bcd + 63(A - C)d^2)) (c + d \tan(e + fx))^{5/2} + 5b^2d(-4b^2cC + 9b^2Bd + 4a^2Cd) \tan(e + fx) (c + d \tan(e + fx))^{5/2} + 35C^2d^2(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} + (105(a - I*b)^2(I*A + B - I*C)*d^3 * (-3*(c - I*d)^{3/2} * \text{ArcTanh}[\text{Sqrt}[c + d \tan(e + fx)]/\text{Sqrt}[c - I*d]] + \text{Sqrt}[c + d \tan(e + fx)] * (4*c - (3*I)*d + d \tan(e + fx))) / 2 + (105*(a + I*b)^2 * ((-I)*A + B + I*C)*d^3 * (-3*(c + I*d)^{3/2} * \text{ArcTanh}[\text{Sqrt}[c + d \tan(e + fx)]/\text{Sqrt}[c + I*d]] + \text{Sqrt}[c + d \tan(e + fx)] * (4*c + (3*I)*d + d \tan(e + fx))) / 2 \right)}{(315*d^3*f)}$$

input

```

Integrate[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e +
f*x] + C*Tan[e + f*x]^2),x]

```

output

```

(2*((28*a^2*C*d^2 + 18*a*b*d*(-2*c*C + 7*B*d) + b^2*(8*c^2*C - 18*B*c*d +
63*(A - C)*d^2))*(c + d*Tan[e + f*x])^(5/2) + 5*b*d*(-4*b^2*c*C + 9*b^2*B*d +
4*a^2*C*d)*Tan[e + f*x]*(c + d*Tan[e + f*x])^(5/2) + 35*C*d^2*(a + b*Tan[e +
f*x])^2*(c + d*Tan[e + f*x])^(5/2) + (105*(a - I*b)^2*(I*A + B - I*C)*d^3
*(-3*(c - I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]] + Sqr
t[c + d*Tan[e + f*x]]*(4*c - (3*I)*d + d*Tan[e + f*x]))) / 2 + (105*(a + I*b
)^2*((-I)*A + B + I*C)*d^3*(-3*(c + I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e +
f*x]]/Sqrt[c + I*d]] + Sqrt[c + d*Tan[e + f*x]]*(4*c + (3*I)*d + d*Tan[e +
f*x]))) / 2) / (315*d^3*f)

```


Rubi [A] (warning: unable to verify)

Time = 4.69 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.02, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.404$, Rules used = {3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

↓ 4130

$$\frac{2 \int -\frac{1}{2} (a + b \tan(e + fx)) (c + d \tan(e + fx))^{3/2} ((4bcC - 4adC - 9bBd) \tan^2(e + fx) - 9(Ab - Cb + aB) d \tan(e + fx)) dx + \frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}}{9df}}{9d}$$

↓ 27

$$\frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (a + b \tan(e + fx)) (c + d \tan(e + fx))^{3/2} ((4bcC - 4adC - 9bBd) \tan^2(e + fx) - 9(Ab - Cb + aB) d \tan(e + fx)) dx}{9d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (a + b \tan(e + fx)) (c + d \tan(e + fx))^{3/2} ((4bcC - 4adC - 9bBd) \tan(e + fx)^2 - 9(Ab - Cb + aB) d \tan(e + fx)) dx}{9d}$$

↓ 4120

$$\frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}}{9df} - \frac{2 \int -\frac{1}{2} (c + d \tan(e + fx))^{3/2} (-2c(4cC - 9Bd)b^2 + 36acCdb - 7a^2(9A - 5C)d^2 - ((8Cc - 2b \tan(e + fx))(-4aCd - 9bBd + 4bcC)(c + d \tan(e + fx))^{5/2})}{7df} dx}{9d}$$

↓ 27

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c+d \tan(e+fx))^{3/2} (-2c(4cC-9Bd)b^2+36acCdb-7a^2(9A-5C)d^2 - ((8Cc^2-18Bdc+63(A-C)d^2)b^2-18ad(2cC-7Bd)b+28a^2Cd^2) \tan^2(e+fx)) dx}{7d}$$

9d

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c+d \tan(e+fx))^{3/2} (-2c(4cC-9Bd)b^2+36acCdb-7a^2(9A-5C)d^2 - ((8Cc^2-18Bdc+63(A-C)d^2)b^2-18ad(2cC-7Bd)b+28a^2Cd^2) \tan(e+fx)) dx}{7d}$$

9d

↓ 4113

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c+d \tan(e+fx))^{3/2} (63(-((A-C)a^2)+2bBa+b^2(A-C))d^2-63(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx)) dx - \frac{2(c+d \tan(e+fx))^{5/2} (28a^2Cd^2-18ad)}{7d}}{7d}$$

9d

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int (c+d \tan(e+fx))^{3/2} (63(-((A-C)a^2)+2bBa+b^2(A-C))d^2-63(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx)) dx - \frac{2(c+d \tan(e+fx))^{5/2} (28a^2Cd^2-18ad)}{7d}}{7d}$$

9d

↓ 4011

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int \sqrt{c+d \tan(e+fx)} (-63((Ac-Cc-Bd)a^2-2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd))d^2-63((Bc+(A-C)d)a^2+2b(Ac-Cc-Bd)a-b^2(Bc+(A-C)d))) dx}{7d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{\int \sqrt{c+d \tan(e+fx)} (-63((Ac-Cc-Bd)a^2-2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd))d^2-63((Bc+(A-C)d)a^2+2b(Ac-Cc-Bd)a-b^2(Bc+(A-C)d))) dx}{7d}$$

↓ 4011

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \int \frac{63((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^2+2b(2c(A-C)d+B(c^2-d^2))a-b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2)))d^2+63(-((2c(A-C)d+B(c^2-d^2))a^2)+2b(Cc^2+2Bdc-Cd^2-A(c^2-d^2)))}{\sqrt{c+d \tan(e+fx)}} dx$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \int \frac{63((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^2+2b(2c(A-C)d+B(c^2-d^2))a-b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2)))d^2+63(-((2c(A-C)d+B(c^2-d^2))a^2)+2b(Cc^2+2Bdc-Cd^2-A(c^2-d^2)))}{\sqrt{c+d \tan(e+fx)}} dx$$

↓ 4022

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e+fx)(-4aCd-9bBd+4bcC)(c+d \tan(e+fx))^{5/2}}{7df} + \frac{-\frac{63}{2}d^2(a+ib)^2(c+id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{63}{2}d^2(a-ib)^2(c-id)^2(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{2f}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e+fx)(-4aCd-9bBd+4bcC)(c+d \tan(e+fx))^{5/2}}{7df} + \frac{-\frac{63}{2}d^2(a+ib)^2(c+id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{63}{2}d^2(a-ib)^2(c-id)^2(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{2f}$$

↓ 4020

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e+fx)(-4aCd-9bBd+4bcC)(c+d \tan(e+fx))^{5/2}}{7df} + \frac{63id^2(a-ib)^2(c-id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - 63id^2(a+ib)^2(c+id)^2(A+iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f}$$

↓ 25

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2b \tan(e+fx)(-4aCd-9bBd+4bcC)(c+d \tan(e+fx))^{5/2}}{7df} + \frac{63id^2(a-ib)^2(c-id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - 63id^2(a+ib)^2(c+id)^2(A+iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f}$$

↓ 73

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{63d(a+ib)^2(c+id)^2(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{i}{d} + 1} d\sqrt{c+d \tan(e+fx)} - 63d(a-ib)^2}{f} - \frac{2b \tan(e+fx)(-4aCd-9bBd+4bcC)(c+d \tan(e+fx))^{5/2}}{7df} + \dots$$

↓ 221

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}}{9df} - \frac{2(c+d \tan(e+fx))^{5/2}(28a^2Cd^2-18abd(2cC-7Bd)+b^2(63d^2(A-C)-18Bcd+8c^2C))}{5df} - \frac{2b \tan(e+fx)(-4aCd-9bBd+4bcC)(c+d \tan(e+fx))^{5/2}}{7df} + \dots$$

input `Int[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]`

output `(2*C*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(5/2))/(9*d*f) - ((2*b*(4*b*c*C - 9*b*B*d - 4*a*C*d)*Tan[e + f*x]*(c + d*Tan[e + f*x])^(5/2))/(7*d*f) + ((-63*(a - I*b)^2*(A - I*B - C)*(c - I*d)^(3/2)*d^2*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f - (63*(a + I*b)^2*(A + I*B - C)*(c + I*d)^(3/2)*d^2*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f - (126*d^2*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d) - b^2*(B*c + (A - C)*d))*Sqrt[c + d*Tan[e + f*x]])/f - (42*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2*(c + d*Tan[e + f*x])^(3/2))/f - (2*(28*a^2*C*d^2 - 18*a*b*d*(2*c*C - 7*B*d) + b^2*(8*c^2*C - 18*B*c*d + 63*(A - C)*d^2))*(c + d*Tan[e + f*x])^(5/2)/(5*d*f))/(7*d))/(9*d)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p(m+1)-1)}(c - a(d/b) + d(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 221 $\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) * \text{ArcTanh}[x / \text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4011 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 0]$
- rule 4020 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \text{ Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$
- rule 4022 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 - I*\text{Tan}[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 + I*\text{Tan}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!IntegerQ}[m]$
- rule 4113 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)(x_)] + (C_.)\tan[(e_.) + (f_.)(x_)]^2), x_Symbol] \rightarrow \text{Simp}[C*((a + b*\text{Tan}[e + f*x])^{(m+1)}/(b*f*(m+1))), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^m*\text{Simp}[A - C + B*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}[\{a, b, e, f, A, B, C, m\}, x] \&\& \text{NeQ}[A*b^2 - a*b*B + a^2*C, 0] \&\& \text{!LeQ}[m, -1]$

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 7938 vs. $2(357) = 714$.

Time = 0.29 (sec) , antiderivative size = 7939, normalized size of antiderivative = 20.05

method	result	size
parts	Expression too large to display	7939
derivativedivides	Expression too large to display	8031
default	Expression too large to display	8031

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58971 vs. $2(347) = 694$.

Time = 72.23 (sec) , antiderivative size = 58971, normalized size of antiderivative = 148.92

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

Too large to include

Sympy [F]

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input

```
integrate((a+b*tan(f*x+e))**2*(c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Integral((a + b*tan(e + f*x))**2*(c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)
```

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output Timed out

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%[1, [0,29,11]%%]}+%%{12, [0,27,11]%%]}+%%{66, [0,25,11]%%]}+%%{2

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Hanged}$$

input

```
int((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) \\
& + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^3 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^5 dx \right) b^2 c d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) a b c d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b^3 d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b^2 c^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a^2 c d \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a b^2 d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a b c^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) b^3 c \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^2 b d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^2 c^2 \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a b^2 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^3 d \\
& + 3 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^2 b c
\end{aligned}$$

input `int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**3*c + int(sqrt(tan(e + f*x)*d + c)*tan(
e + f*x)**5,x)*b**2*c*d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x
)*a*b*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**3*d + int(s
qrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**2*c**2 + int(sqrt(tan(e + f*
x)*d + c)*tan(e + f*x)**3,x)*a**2*c*d + 3*int(sqrt(tan(e + f*x)*d + c)*tan
(e + f*x)**3,x)*a*b**2*d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,
x)*a*b*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b**3*c + 3*i
nt(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*b*d + int(sqrt(tan(e +
f*x)*d + c)*tan(e + f*x)**2,x)*a**2*c**2 + 3*int(sqrt(tan(e + f*x)*d + c)
*tan(e + f*x)**2,x)*a*b**2*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x
)*a**3*d + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**2*b*c
```

3.99 $\int (a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2} (A + B \tan(e$

Optimal result	1119
Mathematica [A] (verified)	1120
Rubi [A] (warning: unable to verify)	1120
Maple [B] (verified)	1125
Fricas [B] (verification not implemented)	1125
Sympy [F]	1126
Maxima [F(-1)]	1126
Giac [F(-2)]	1127
Mupad [F(-1)]	1127
Reduce [F]	1128

Optimal result

Integrand size = 45, antiderivative size = 273

$$\begin{aligned}
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \\
 & \frac{(ia + b)(A - iB - C)(c - id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f} \\
 & + \frac{(ia - b)(A + iB - C)(c + id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f} \\
 & + \frac{2(Abc + aBc - bcC + aAd - bBd - aCd) \sqrt{c + d \tan(e + fx)}}{f} \\
 & + \frac{2(Ab + aB - bC)(c + d \tan(e + fx))^{3/2}}{3f} \\
 & - \frac{2(2bcC - 7bBd - 7aCd)(c + d \tan(e + fx))^{5/2}}{35d^2 f} \\
 & + \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df}
 \end{aligned}$$

output

$$\begin{aligned} & -(I*a+b)*(A-I*B-C)*(c-I*d)^{(3/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c-I*d)^{(1/2)})/f+(I*a-b)*(A+I*B-C)*(c+I*d)^{(3/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c+I*d)^{(1/2)})/f+2*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*(c+d*\tan(f*x+e))^{(1/2)}/f+2/3*(A*b+B*a-C*b)*(c+d*\tan(f*x+e))^{(3/2)}/f-2/35*(-7*B*b*d-7*C*a*d+2*C*b*c)*(c+d*\tan(f*x+e))^{(5/2)}/d^2/f+2/7*b*C*\tan(f*x+e)*(c+d*\tan(f*x+e))^{(5/2)}/d/f \end{aligned}$$
Mathematica [A] (verified)

Time = 3.07 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.95

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{2(-2bcC+7bBd+7aCd)(c+d \tan(e+fx))^{5/2}}{d} + 10bC \tan(e + fx)(c + d \tan(e + fx))^{5/2} + \frac{35}{3}(i$$

input

$$\text{Integrate}[(a + b*\text{Tan}[e + f*x])*(c + d*\text{Tan}[e + f*x])^{(3/2)}*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2), x]$$

output

$$\begin{aligned} & ((2*(-2*b*c*C + 7*b*B*d + 7*a*C*d)*(c + d*\text{Tan}[e + f*x])^{(5/2)})/d + 10*b*C*\text{Tan}[e + f*x]*(c + d*\text{Tan}[e + f*x])^{(5/2)} + (35*(I*a + b)*(A - I*B - C)*d*(-3*(c - I*d)^{(3/2)}*\text{ArcTanh}[\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[c - I*d]] + \text{Sqrt}[c + d*\text{Tan}[e + f*x]]*(4*c - (3*I)*d + d*\text{Tan}[e + f*x]))) / 3 + (35*((-I)*a + b)*(A + I*B - C)*d*(-3*(c + I*d)^{(3/2)}*\text{ArcTanh}[\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[c + I*d]] + \text{Sqrt}[c + d*\text{Tan}[e + f*x]]*(4*c + (3*I)*d + d*\text{Tan}[e + f*x]))) / 3) / (35*d*f) \end{aligned}$$
Rubi [A] (warning: unable to verify)

Time = 2.72 (sec) , antiderivative size = 268, normalized size of antiderivative = 0.98, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.356$, Rules used = {3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 & \quad \downarrow 3042 \\
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx \\
 & \quad \downarrow 4120 \\
 & \frac{2bcC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \\
 & \frac{2 \int \frac{1}{2}(c + d \tan(e + fx))^{3/2} ((2bcC - 7adC - 7bBd) \tan^2(e + fx) - 7(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 7aAa)}{7d} dx \\
 & \quad \downarrow 27 \\
 & \frac{2bcC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \\
 & \frac{\int (c + d \tan(e + fx))^{3/2} ((2bcC - 7adC - 7bBd) \tan^2(e + fx) - 7(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 7aAa)}{7d} dx \\
 & \quad \downarrow 3042 \\
 & \frac{2bcC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \\
 & \frac{\int (c + d \tan(e + fx))^{3/2} ((2bcC - 7adC - 7bBd) \tan(e + fx)^2 - 7(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 7aAa)}{7d} dx \\
 & \quad \downarrow 4113 \\
 & \frac{2bcC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \\
 & \frac{\int (c + d \tan(e + fx))^{3/2} (7(bB - a(A - C))d - 7(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{2(-7aCd - 7bBd + 2bcC)(c + d \tan(e + fx))}{5df}}{7d} \\
 & \quad \downarrow 3042 \\
 & \frac{2bcC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \\
 & \frac{\int (c + d \tan(e + fx))^{3/2} (7(bB - a(A - C))d - 7(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{2(-7aCd - 7bBd + 2bcC)(c + d \tan(e + fx))}{5df}}{7d} \\
 & \quad \downarrow 4011 \\
 & \frac{2bcC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \\
 & \frac{\int \sqrt{c + d \tan(e + fx)} (7d(bBc + b(A - C)d - a(Ac - Cc - Bd)) - 7d(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx))}{7d} dx
 \end{aligned}$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{\int \sqrt{c + d \tan(e + fx)}(7d(bBc + b(A - C)d - a(Ac - Cc - Bd)) - 7d(Abc + aBc - bCc + aAd - bBd - aCd) \tan(e + fx))}{7d}$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{\int \frac{7d(a(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2)) + b(2c(A - C)d + B(c^2 - d^2))) - 7d(2aAcd - 2acCd + Ab(c^2 - d^2) + aB(c^2 - d^2) - b(Cc^2 + 2Bdc - Cd^2)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}}$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{\int \frac{7d(a(Cc^2 + 2Bdc - Cd^2 - A(c^2 - d^2)) + b(2c(A - C)d + B(c^2 - d^2))) - 7d(2aAcd - 2acCd + Ab(c^2 - d^2) + aB(c^2 - d^2) - b(Cc^2 + 2Bdc - Cd^2)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}}$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{-\frac{7}{2}d(a + ib)(c + id)^2(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{7}{2}d(a - ib)(c - id)^2(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx -$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{-\frac{7}{2}d(a + ib)(c + id)^2(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{7}{2}d(a - ib)(c - id)^2(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx -$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{7id(a - ib)(c - id)^2(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} + \frac{7id(a + ib)(c + id)^2(A + iB - C) \int -\frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}} dx}{2f}$$

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{7id(a - ib)(c - id)^2(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} + \frac{7id(a + ib)(c + id)^2(A + iB - C) \int -\frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}} dx}{2f}$$

↓ 25

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{7id(a-ib)(c-id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} - \frac{7id(a+ib)(c+id)^2(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}}}{2f}$$

↓ 73

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{7(a+ib)(c+id)^2(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{f} - \frac{7(a-ib)(c-id)^2(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{f}$$

↓ 221

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{5/2}}{7df} - \frac{7d(a-ib)(c-id)^{3/2}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} - \frac{7d(a+ib)(c+id)^{3/2}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f} - \frac{14d(aB+Ab-bC)(c+d \tan(e+fx))}{3f}$$

7d

input

```
Int[(a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
(2*b*C*Tan[e + f*x]*(c + d*Tan[e + f*x])^(5/2))/(7*d*f) - ((-7*(a - I*b)*(A - I*B - C)*(c - I*d)^(3/2)*d*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f - (7*(a + I*b)*(A + I*B - C)*(c + I*d)^(3/2)*d*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f - (14*d*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d)*Sqrt[c + d*Tan[e + f*x]])/f - (14*(A*b + a*B - b*C)*d*(c + d*Tan[e + f*x])^(3/2))/(3*f) + (2*(2*b*c*C - 7*b*B*d - 7*a*C*d)*(c + d*Tan[e + f*x])^(5/2))/(5*d*f))/(7*d)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```


- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
 [(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
 , x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
 0] && GtQ[m, 0]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
 c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
 b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
 1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
 *(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
 - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`
- rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
 + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
 b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
 mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
 NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120

```

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f
_)*(x_)])^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5106 vs. $2(239) = 478$.

Time = 0.27 (sec) , antiderivative size = 5107, normalized size of antiderivative = 18.71

method	result	size
parts	Expression too large to display	5107
derivativedivides	Expression too large to display	5149
default	Expression too large to display	5149

input

```

int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2
),x,method=_RETURNVERBOSE)

```

output

```

result too large to display

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31081 vs. $2(232) = 464$.

Time = 16.92 (sec) , antiderivative size = 31081, normalized size of antiderivative = 113.85

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))*(c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output Timed out

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0,24,9]%%}+%%{10, [0,22,9]%%}+%%{45, [0,20,9]%%}+%%{120,`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Hanged}$$

input `int((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `\text{Hanged}`

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) \\
& + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^2 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) bcd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) acd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) b^2 d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) bc^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) abd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) ac^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) b^2 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^2 d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) abc
\end{aligned}$$

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**2*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b**2*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b*c**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*b*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*b**2*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**2*d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a*b*c
```

3.100 $\int (c+d \tan(e+fx))^{3/2} (A + B \tan(e + fx) + C \tan^2$

Optimal result	1129
Mathematica [A] (verified)	1130
Rubi [A] (warning: unable to verify)	1130
Maple [B] (verified)	1134
Fricas [B] (verification not implemented)	1135
Sympy [F]	1136
Maxima [F]	1136
Giac [F(-2)]	1136
Mupad [B] (verification not implemented)	1137
Reduce [F]	1138

Optimal result

Integrand size = 35, antiderivative size = 187

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx =$$

$$\frac{(iA + B - iC)(c - id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f}$$

$$- \frac{(B - i(A - C))(c + id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f}$$

$$+ \frac{2(Bc + (A - C)d)\sqrt{c + d \tan(e + fx)}}{f}$$

$$+ \frac{2B(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df}$$

output

```
-(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/f
-(B-I*(A-C))*(c+I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/f
+2*(B*c+(A-C)*d)*(c+d*tan(f*x+e))^(1/2)/f+2/3*B*(c+d*tan(f*x+e))^(3/2)/f+2
/5*C*(c+d*tan(f*x+e))^(5/2)/d/f
```

Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.08

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{6C(c + d \tan(e + fx))^{5/2}}{d} + 5(iA + B - iC) \left(-3(c - id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c + d \tan(e + fx)}}{\sqrt{c - id}}\right) + \sqrt{c} \right)$$

input

```
Integrate[(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
((6*C*(c + d*Tan[e + f*x])^(5/2))/d + 5*(I*A + B - I*C)*(-3*(c - I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]] + Sqrt[c + d*Tan[e + f*x]]*(4*c - (3*I)*d + d*Tan[e + f*x])) + 5*((-I)*A + B + I*C)*(-3*(c + I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]] + Sqrt[c + d*Tan[e + f*x]]*(4*c + (3*I)*d + d*Tan[e + f*x])))/(15*f)
```

Rubi [A] (warning: unable to verify)

Time = 1.68 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.89, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4113, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow \text{4113}$$

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^{3/2} dx + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df}$$

$$\begin{aligned} & \downarrow 3042 \\ & \int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^{3/2} dx + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df} \\ & \downarrow 4011 \\ & \int \sqrt{c + d \tan(e + fx)}(Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \\ & \quad \frac{2B(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df} \\ & \downarrow 3042 \\ & \int \sqrt{c + d \tan(e + fx)}(Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \\ & \quad \frac{2B(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df} \\ & \downarrow 4011 \\ & \int \frac{-Cc^2 - 2Bdc + Cd^2 + A(c^2 - d^2) + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \\ & \frac{2(d(A - C) + Bc)\sqrt{c + d \tan(e + fx)}}{f} + \frac{2B(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df} \\ & \downarrow 3042 \\ & \int \frac{-Cc^2 - 2Bdc + Cd^2 + A(c^2 - d^2) + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \\ & \frac{2(d(A - C) + Bc)\sqrt{c + d \tan(e + fx)}}{f} + \frac{2B(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df} \\ & \downarrow 4022 \\ & \frac{1}{2}(c + id)^2(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(c - id)^2(A - iB - \\ & C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2(d(A - C) + Bc)\sqrt{c + d \tan(e + fx)}}{f} + \\ & \quad \frac{2B(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5df} \\ & \downarrow 3042 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2}(c+id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(c-id)^2(A-iB-C) \\
& \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2(d(A-C)+Bc)\sqrt{c+d \tan(e+fx)}}{f} + \\
& \frac{2B(c+d \tan(e+fx))^{3/2}}{3f} + \frac{2C(c+d \tan(e+fx))^{5/2}}{5df} \\
& \quad \downarrow 4020 \\
& \frac{i(c-id)^2(A-iB-C) \int -\frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} - \\
& \frac{i(c+id)^2(A+iB-C) \int -\frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} + \\
& \frac{2(d(A-C)+Bc)\sqrt{c+d \tan(e+fx)}}{f} + \frac{2B(c+d \tan(e+fx))^{3/2}}{3f} + \frac{2C(c+d \tan(e+fx))^{5/2}}{5df} \\
& \quad \downarrow 25 \\
& -\frac{i(c-id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} + \\
& \frac{i(c+id)^2(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} + \\
& \frac{2(d(A-C)+Bc)\sqrt{c+d \tan(e+fx)}}{f} + \frac{2B(c+d \tan(e+fx))^{3/2}}{3f} + \frac{2C(c+d \tan(e+fx))^{5/2}}{5df} \\
& \quad \downarrow 73 \\
& \frac{(c+id)^2(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} + \\
& \frac{(c-id)^2(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} + \\
& \frac{2(d(A-C)+Bc)\sqrt{c+d \tan(e+fx)}}{f} + \frac{2B(c+d \tan(e+fx))^{3/2}}{3f} + \frac{2C(c+d \tan(e+fx))^{5/2}}{5df} \\
& \quad \downarrow 221 \\
& \frac{(c-id)^{3/2}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{(c+id)^{3/2}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f} + \\
& \frac{2(d(A-C)+Bc)\sqrt{c+d \tan(e+fx)}}{f} + \frac{2B(c+d \tan(e+fx))^{3/2}}{3f} + \frac{2C(c+d \tan(e+fx))^{5/2}}{5df}
\end{aligned}$$

input

```
Int[(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output
$$\begin{aligned} & ((A - I*B - C)*(c - I*d)^{(3/2)}*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + ((A \\ & + I*B - C)*(c + I*d)^{(3/2)}*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f + (2*(B* \\ & c + (A - C)*d)*Sqrt[c + d*Tan[e + f*x]])/f + (2*B*(c + d*Tan[e + f*x])^{(3/ \\ & 2)})/(3*f) + (2*C*(c + d*Tan[e + f*x])^{(5/2)})/(5*d*f) \end{aligned}$$

Defintions of rubi rules used

rule 25
$$Int[-(Fx_), x_Symbol] \rightarrow Simp[Identity[-1] \quad Int[Fx, x], x]$$

rule 73
$$Int[((a_.) + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow With[\\ \{p = Denominator[m]\}, Simp[p/b \quad Subst[Int[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + \\ d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; FreeQ[\{a, b, c, d\}, x] \&\& Lt \\ Q[-1, m, 0] \&\& LeQ[-1, n, 0] \&\& LeQ[Denominator[n], Denominator[m]] \&\& IntL \\ inearQ[a, b, c, d, m, n, x]$$

rule 221
$$Int[((a_) + (b_.)*(x_)^2)^{(-1)}, x_Symbol] \rightarrow Simp[(Rt[-a/b, 2]/a)*ArcTanh[x \\ /Rt[-a/b, 2]], x] /; FreeQ[\{a, b\}, x] \&\& NegQ[a/b]$$

rule 3042
$$Int[u_, x_Symbol] \rightarrow Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear \\ Q[u, x]$$

rule 4011
$$Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^{(m_)}*((c_.) + (d_.)*tan[(e_.) + \\ (f_.)*(x_)]), x_Symbol] \rightarrow Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int \\ [(a + b*Tan[e + f*x])^{(m - 1)}*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x] \\ , x] /; FreeQ[\{a, b, c, d, e, f\}, x] \&\& NeQ[b*c - a*d, 0] \&\& NeQ[a^2 + b^2, \\ 0] \&\& GtQ[m, 0]$$

rule 4020
$$Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^{(m_)}*((c_) + (d_.)*tan[(e_.) + \\ (f_.)*(x_)]), x_Symbol] \rightarrow Simp[c*(d/f) \quad Subst[Int[(a + (b/d)*x)^m/(d^2 + \\ c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[\{a, b, c, d, e, f, m\}, x] \&\& NeQ[\\ b*c - a*d, 0] \&\& NeQ[a^2 + b^2, 0] \&\& EqQ[c^2 + d^2, 0]$$

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2499 vs. $2(158) = 316$.

Time = 0.18 (sec) , antiderivative size = 2500, normalized size of antiderivative = 13.37

method	result	size
parts	Expression too large to display	2500
derivativedivides	Expression too large to display	2517
default	Expression too large to display	2517

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETURN
VERBOSE)
```

output

```
A*(2/f*d*(c+d*tan(f*x+e))^(1/2)-1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))
^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2
*c)^(1/2)*(c^2+d^2)^(1/2)*c+1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/
2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(
1/2)*c^2-1/4/f*d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1
/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+2/f*d/(2*(c^
2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)
)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c-1/f*d/(2*(c^2+d^2)^(1/2)-2*
c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(
2*(c^2+d^2)^(1/2)-2*c)^(1/2))*(c^2+d^2)^(1/2)+1/4/f/d*ln(-d*tan(f*x+e)-c+(
c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*(2*(c
^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*c-1/4/f/d*ln(-d*tan(f*x+e)-c+(c+d
*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*(2*(c^2+
d^2)^(1/2)+2*c)^(1/2)*c^2+1/4/f*d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)
)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(
1/2)-2/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((-2*(c+d*tan(f*x+e))^(1/2)
+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c+1/f*d/(2*
(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((-2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(
1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*(c^2+d^2)^(1/2))+B*(2/3/f
*(c+d*tan(f*x+e))^(3/2)+2/f*(c+d*tan(f*x+e))^(1/2)*c+1/4/f*ln(d*tan(f*x...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6846 vs. $2(151) = 302$.

Time = 0.86 (sec) , antiderivative size = 6846, normalized size of antiderivative = 36.61

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algori
thm="fricas")
```

output

Too large to include

Sympy [F]

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(d \tan(fx + e) + c)^{3/2} dx$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(3/2), x)`

Giac [F(-2)]

Exception generated.

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,19,7]}%%+%%{8,[0,17,7]}%%+%%{28,[0,15,7]}%%+
%%{56,[0
```

Mupad [B] (verification not implemented)

Time = 39.81 (sec) , antiderivative size = 4260, normalized size of antiderivative = 22.78

$$\int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
((2*C*c^2)/(d*f) - (2*C*(d^3*f + c^2*d*f))/(d^2*f^2))*(c + d*tan(e + f*x))
^(1/2) - log((((16*c*d^2*((-B^4*d^2*f^4*(3*c^2 - d^2)^2)^(1/2) + B^2*c^3*
f^2 - 3*B^2*c*d^2*f^2)/f^4)^(1/2)*(B*c^2 + B*d^2 + f*((-B^4*d^2*f^4*(3*c^
2 - d^2)^2)^(1/2) + B^2*c^3*f^2 - 3*B^2*c*d^2*f^2)/f^4)^(1/2)*(c + d*tan(e
+ f*x))^(1/2)))/f - (16*B^2*d^2*(c + d*tan(e + f*x))^(1/2)*(c^4 + d^4 - 6
*c^2*d^2))/f^2)*(((B^4*d^2*f^4*(3*c^2 - d^2)^2)^(1/2) + B^2*c^3*f^2 - 3*B
^2*c*d^2*f^2)/f^4)^(1/2))/2 - (8*B^3*d^2*(c^2 - d^2)*(c^2 + d^2)^2)/f^3)*(
((6*B^4*c^2*d^4*f^4 - B^4*d^6*f^4 - 9*B^4*c^4*d^2*f^4)^(1/2) + B^2*c^3*f^2
- 3*B^2*c*d^2*f^2)/(4*f^4))^(1/2) - log((((16*c*d^2*((-B^4*d^2*f^4*(3*c
^2 - d^2)^2)^(1/2) - B^2*c^3*f^2 + 3*B^2*c*d^2*f^2)/f^4)^(1/2)*(B*c^2 + B*
d^2 + f*((-B^4*d^2*f^4*(3*c^2 - d^2)^2)^(1/2) - B^2*c^3*f^2 + 3*B^2*c*d^
2*f^2)/f^4)^(1/2)*(c + d*tan(e + f*x))^(1/2)))/f - (16*B^2*d^2*(c + d*tan(
e + f*x))^(1/2)*(c^4 + d^4 - 6*c^2*d^2))/f^2)*(((B^4*d^2*f^4*(3*c^2 - d^
2)^2)^(1/2) - B^2*c^3*f^2 + 3*B^2*c*d^2*f^2)/f^4)^(1/2))/2 - (8*B^3*d^2*(c
^2 - d^2)*(c^2 + d^2)^2)/f^3)*(((6*B^4*c^2*d^4*f^4 - B^4*d^6*f^4 - 9*B^4*
c^4*d^2*f^4)^(1/2) - B^2*c^3*f^2 + 3*B^2*c*d^2*f^2)/(4*f^4))^(1/2) + log((
((16*c*d^2*((-B^4*d^2*f^4*(3*c^2 - d^2)^2)^(1/2) + B^2*c^3*f^2 - 3*B^2*c*
d^2*f^2)/f^4)^(1/2)*(B*c^2 + B*d^2 - f*((-B^4*d^2*f^4*(3*c^2 - d^2)^2)^(1
/2) + B^2*c^3*f^2 - 3*B^2*c*d^2*f^2)/f^4)^(1/2)*(c + d*tan(e + f*x))^(1/2)
))/f + (16*B^2*d^2*(c + d*tan(e + f*x))^(1/2)*(c^4 + d^4 - 6*c^2*d^2))/...
```

Reduce [F]

$$\begin{aligned}
& \int (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) \\
& + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} dx \right) ac \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) cd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) bd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) c^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) ad \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) bc
\end{aligned}$$

input `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int(sqrt(tan(e + f*x)*d + c),x)*a*c + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*b*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*b*c`

3.101 $\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$

Optimal result	1139
Mathematica [A] (verified)	1140
Rubi [A] (warning: unable to verify)	1140
Maple [B] (verified)	1146
Fricas [F(-1)]	1147
Sympy [F]	1147
Maxima [F(-2)]	1147
Giac [F(-2)]	1148
Mupad [B] (verification not implemented)	1148
Reduce [F]	1149

Optimal result

Integrand size = 47, antiderivative size = 271

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx =$$

$$\frac{(iA+B-iC)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a-ib)f} - \frac{(A+iB-C)(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(ia-b)f}$$

$$- \frac{2(Ab^2-a(bB-aC))(bc-ad)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{b^{5/2}(a^2+b^2)f}$$

$$+ \frac{2(bcC+bBd-aCd)\sqrt{c+d \tan(e+fx)}}{b^2f} + \frac{2C(c+d \tan(e+fx))^{3/2}}{3bf}$$

output

```
-(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(a-I*b)/f-(A+I*B-C)*(c+I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(I*a-b)/f-2*(A*b^2-a*(B*b-C*a))*(-a*d+b*c)^(3/2)*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/b^(5/2)/(a^2+b^2)/f+2*(B*b*d-C*a*d+C*b*c)*(c+d*tan(f*x+e))^(1/2)/b^2/f+2/3*C*(c+d*tan(f*x+e))^(3/2)/b/f
```


Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.98

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \frac{3ib \left(-((a+ib)(A-iB-C)(c-id)^{3/2} \operatorname{arctanh} \left(\frac{c + d \tan(e + fx)}{\sqrt{c - id}} \right) + (a - ib)(A + iB - C)(c + id)^{3/2} \operatorname{arctanh} \left(\frac{c + d \tan(e + fx)}{\sqrt{c + id}} \right) \right)}{(a^2 + b^2) - (6(Ab^2 + a(-bB) + aC))(b^2c - a^2d)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{b} \sqrt{c + d \tan(e + fx)}}{\sqrt{b^2c - a^2d}} \right) + (6(b^2cC + bB^2d - a^2Cd) \sqrt{c + d \tan(e + fx)})/b + 2C(c + d \tan(e + fx))^{3/2}}{(3b^2f)}$$

input

```
Integrate[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]
```

output

```
((((3*I)*b*(-((a + I*b)*(A - I*B - C)*(c - I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]) + (a - I*b)*(A + I*B - C)*(c + I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]]))/(a^2 + b^2) - (6*(A*b^2 + a*(-(b*B) + a*C))*(b*c - a*d)^(3/2)*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/(b^(3/2)*(a^2 + b^2)) + (6*(b*c*C + b*B*d - a*C*d)*Sqrt[c + d*Tan[e + f*x]]/b + 2*C*(c + d*Tan[e + f*x])^(3/2))/(3*b*f)
```

Rubi [A] (warning: unable to verify)Time = 4.42 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.01, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.404$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{a + b \tan(e + fx)} dx$$

↓ 4130

$$\begin{aligned}
 & \frac{2 \int \frac{3\sqrt{c+d \tan(e+fx)}((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{2(a+b \tan(e+fx))} dx}{\frac{3b}{2C(c+d \tan(e+fx))^{3/2}}} + \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{\int \frac{\sqrt{c+d \tan(e+fx)}((bcC-adC+bBd) \tan^2(e+fx)+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx}{\frac{b}{2C(c+d \tan(e+fx))^{3/2}}} + \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{\int \frac{\sqrt{c+d \tan(e+fx)}((bcC-adC+bBd) \tan(e+fx)^2+b(Bc+(A-C)d) \tan(e+fx)+Abc-aCd)}{a+b \tan(e+fx)} dx}{\frac{b}{2C(c+d \tan(e+fx))^{3/2}}} + \\
 & \qquad \qquad \qquad \downarrow 4130 \\
 & \frac{2 \int \frac{Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+(d(Bc+(A-C)d)b^2+(bc-ad)(bcC-adC+bBd)) \tan^2(e+fx)+ad(aCd-b(2cC+Bd))}{2(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{\frac{b}{2C(c+d \tan(e+fx))^{3/2}}} + \frac{2(-aCd+bBd)}{b} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{\int \frac{Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+(d(Bc+(A-C)d)b^2+(bc-ad)(bcC-adC+bBd)) \tan^2(e+fx)+ad(aCd-b(2cC+Bd))}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{\frac{b}{2C(c+d \tan(e+fx))^{3/2}}} + \frac{2(-aCd+bBd)}{b} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{\int \frac{Ac^2b^2+(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2+(d(Bc+(A-C)d)b^2+(bc-ad)(bcC-adC+bBd)) \tan(e+fx)^2+ad(aCd-b(2cC+Bd))}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{\frac{b}{2C(c+d \tan(e+fx))^{3/2}}} + \frac{2(-aCd+bBd)}{b} \\
 & \qquad \qquad \qquad \downarrow 4136
 \end{aligned}$$

$$\int \frac{b^2(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d+B(c^2-d^2)))-b^2(2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2))\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}\frac{a^2+b^2}{b}} dx + \dots$$

$$\frac{2C(c+d\tan(e+fx))^{3/2}}{3bf}$$

↓ 25

$$\frac{(bc-ad)^2(Ab^2-a(bB-aC))\int\frac{\tan^2(e+fx)+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}}dx}{\frac{a^2+b^2}{b}} - \frac{b^2(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d+B(c^2-d^2)))-b^2(2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2))\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}\frac{a^2+b^2}{b}}$$

$$\frac{2C(c+d\tan(e+fx))^{3/2}}{3bf}$$

↓ 3042

$$\frac{(bc-ad)^2(Ab^2-a(bB-aC))\int\frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}}dx}{\frac{a^2+b^2}{b}} - \frac{b^2(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d+B(c^2-d^2)))-b^2(2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2))\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}\frac{a^2+b^2}{b}}$$

$$\frac{2C(c+d\tan(e+fx))^{3/2}}{3bf}$$

↓ 4022

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2C(c+d\tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC))\int\frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}}dx}{\frac{a^2+b^2}{b}} - \frac{\frac{1}{2}b^2(a+ib)(c-id)^2(A-iB-C)\int\frac{i\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}}dx}{b}$$

↓ 3042

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2C(c+d\tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC))\int\frac{\tan(e+fx)^2+1}{(a+b\tan(e+fx))\sqrt{c+d\tan(e+fx)}}dx}{\frac{a^2+b^2}{b}} - \frac{\frac{1}{2}b^2(a+ib)(c-id)^2(A-iB-C)\int\frac{i\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}}dx}{b}$$

↓ 4020

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\frac{2C(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{ib^2(a-ib)(c+id)^2(A+iB-C) \int \frac{1}{(i \tan(e+fx))^2} dx}{b}}{b}$$

25

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\frac{2C(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{ib^2(a+ib)(c-id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))^2} dx}{b}}{b}$$

73

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\frac{2C(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{b^2(a+ib)(c-id)^2(A-iB-C) \int \frac{i \tan^2(e+fx)}{df} dx}{b}}{b}$$

221

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\frac{2C(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{b^2(a+ib)(c-id)^{3/2}(A-iB-C) \arctan\left(\frac{1}{f}\right)}{b}}{b}$$

4117

$$\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\frac{2C(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{(bc-ad)^2(Ab^2-a(bB-aC)) \int \frac{1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d \tan(e+fx)}{f(a^2+b^2)} - \frac{b^2(a+ib)(c-id)^{3/2}(A-iB-C)}{f}}{b}$$

73

$$\frac{\frac{2C(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2(bc-ad)^2(Ab^2 - a(bB - aC)) \int \frac{1}{a + \frac{b(c+d \tan(e+fx))}{d} - \frac{bc}{d}} d \sqrt{c+d \tan(e+fx)} - \frac{b^2(a+ib)(c-id)^{3/2}(A-iB-C)}{f}}{\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\dots}{df(a^2+b^2)}} + \frac{\dots}{b}$$

↓ 221

$$\frac{\frac{2C(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2(bc-ad)^{3/2}(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right) - \frac{b^2(a+ib)(c-id)^{3/2}(A-iB-C) \operatorname{arctan}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{f}}{\frac{2(-aCd+bBd+bcC)\sqrt{c+d \tan(e+fx)}}{bf} + \frac{\dots}{\sqrt{b}f(a^2+b^2)}} + \frac{\dots}{b}$$

input

```
Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]
```

output

```
(2*C*(c + d*Tan[e + f*x])^(3/2))/(3*b*f) + (((-(((a + I*b)*b^2*(A - I*B - C)*(c - I*d)^(3/2)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f) - ((a - I*b)*b^2*(A + I*B - C)*(c + I*d)^(3/2)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f)/(a^2 + b^2)) - (2*(A*b^2 - a*(b*B - a*C))*(b*c - a*d)^(3/2)*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]/(Sqrt[b]*(a^2 + b^2)*f))/b + (2*(b*c*C + b*B*d - a*C*d)*Sqrt[c + d*Tan[e + f*x]])/(b*f))/b
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

- rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p*(m+1)-1)}(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] \text{ /}; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 221 $\text{Int}[(a_.) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ /}; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /}; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4020 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)*\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \text{ Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\text{Tan}[e + f*x]], x] \text{ /}; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$
- rule 4022 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)*\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 - I*\text{Tan}[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 + I*\text{Tan}[e + f*x]), x], x] \text{ /}; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{!IntegerQ}[m]$
- rule 4117 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)(x_)]^{(m_.)}((c_.) + (d_.)*\tan[(e_.) + (f_.)(x_)]^{(n_.)}((A_.) + (C_.)*\tan[(e_.) + (f_.)(x_)]^2), x_Symbol] \rightarrow \text{Simp}[A/f \text{ Subst}[\text{Int}[(a + b*x)^m*(c + d*x)^n, x], x, \text{Tan}[e + f*x]], x] \text{ /}; \text{FreeQ}[\{a, b, c, d, e, f, A, C, m, n\}, x] \ \&\& \ \text{EqQ}[A, C]$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6054 vs. $2(234) = 468$.

Time = 0.21 (sec) , antiderivative size = 6055, normalized size of antiderivative = 22.34

method	result	size
derivativeldivides	Expression too large to display	6055
default	Expression too large to display	6055

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)),x)`

output `Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Exception raised: ValueError}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for m
ore detail
```

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Exception raised: TypeError}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e)),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [B] (verification not implemented)

Time = 52.30 (sec) , antiderivative size = 106783, normalized size of antiderivative = 394.03

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input

```
int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x)),x)
```

output

```
atan(((((((32*(4*B*a*b^8*d^12*f^4 - 4*B*b^9*c*d^11*f^4 + 8*B*a^3*b^6*d^12*
f^4 + 4*B*a^5*b^4*d^12*f^4 - 4*B*b^9*c^3*d^9*f^4 + 8*B*a*b^8*c^2*d^10*f^4
+ 4*B*a*b^8*c^4*d^8*f^4 - 12*B*a^2*b^7*c*d^11*f^4 - 12*B*a^4*b^5*c*d^11*f^
4 - 4*B*a^6*b^3*c*d^11*f^4 - 12*B*a^2*b^7*c^3*d^9*f^4 + 16*B*a^3*b^6*c^2*d
^10*f^4 + 8*B*a^3*b^6*c^4*d^8*f^4 - 12*B*a^4*b^5*c^3*d^9*f^4 + 8*B*a^5*b^4
*c^2*d^10*f^4 + 4*B*a^5*b^4*c^4*d^8*f^4 - 4*B*a^6*b^3*c^3*d^9*f^4)))/(b*f^5
) - (32*(c + d*tan(e + f*x))^(1/2)*(-(((8*B^2*a^2*c^3*f^2 - 8*B^2*b^2*c^3*
f^2 - 16*B^2*a*b*d^3*f^2 - 24*B^2*a^2*c*d^2*f^2 + 24*B^2*b^2*c*d^2*f^2 + 4
8*B^2*a*b*c^2*d*f^2)^2/4 - (16*a^4*f^4 + 16*b^4*f^4 + 32*a^2*b^2*f^4)*(B^4
*c^6 + B^4*d^6 + 3*B^4*c^2*d^4 + 3*B^4*c^4*d^2)))^(1/2) - 4*B^2*a^2*c^3*f^2
+ 4*B^2*b^2*c^3*f^2 + 8*B^2*a*b*d^3*f^2 + 12*B^2*a^2*c*d^2*f^2 - 12*B^2*b
^2*c*d^2*f^2 - 24*B^2*a*b*c^2*d*f^2)/(16*(a^4*f^4 + b^4*f^4 + 2*a^2*b^2*f^
4)))^(1/2)*(16*b^10*d^10*f^4 + 16*a^2*b^8*d^10*f^4 - 16*a^4*b^6*d^10*f^4 -
16*a^6*b^4*d^10*f^4 + 24*b^10*c^2*d^8*f^4 + 40*a^2*b^8*c^2*d^8*f^4 + 8*a^
4*b^6*c^2*d^8*f^4 - 8*a^6*b^4*c^2*d^8*f^4 + 8*a*b^9*c*d^9*f^4 + 24*a^3*b^7
*c*d^9*f^4 + 24*a^5*b^5*c*d^9*f^4 + 8*a^7*b^3*c*d^9*f^4))/(b*f^4))*(-(((8*
B^2*a^2*c^3*f^2 - 8*B^2*b^2*c^3*f^2 - 16*B^2*a*b*d^3*f^2 - 24*B^2*a^2*c*d^
2*f^2 + 24*B^2*b^2*c*d^2*f^2 + 48*B^2*a*b*c^2*d*f^2)^2/4 - (16*a^4*f^4 + 1
6*b^4*f^4 + 32*a^2*b^2*f^4)*(B^4*c^6 + B^4*d^6 + 3*B^4*c^2*d^4 + 3*B^4*c^4
*d^2)))^(1/2) - 4*B^2*a^2*c^3*f^2 + 4*B^2*b^2*c^3*f^2 + 8*B^2*a*b*d^3*f^2...
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \left(\int \frac{\sqrt{d \tan(fx + e) + c}}{\tan(fx + e) b + a} dx \right) ac$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^3}{\tan(fx + e) b + a} dx \right) cd$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) bd$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) c^2$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) ad$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) bc$$

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)*b + a),x)*a*c + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*b + a),x)*c*d + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*b*d + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*c**2 + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*a*d + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*b*c
```

3.102
$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$$

Optimal result	1151
Mathematica [B] (verified)	1152
Rubi [A] (warning: unable to verify)	1153
Maple [B] (verified)	1160
Fricas [F(-1)]	1160
Sympy [F]	1160
Maxima [F(-2)]	1161
Giac [F(-2)]	1161
Mupad [F(-1)]	1162
Reduce [F]	1162

Optimal result

Integrand size = 47, antiderivative size = 372

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx =$$

$$\frac{(iA+B-iC)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a-ib)^2 f}$$

$$-\frac{(B-i(A-C))(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a+ib)^2 f}$$

$$+\frac{\sqrt{bc-ad}(a^3bBd-3a^4Cd-b^4(2Bc+3Ad)-ab^3(4Ac-4cC-5Bd)+a^2b^2(2Bc+(A-7C)d)) \operatorname{arctan}\left(\frac{b \tan(e+fx)}{a}\right)}{b^{5/2}(a^2+b^2)^2 f}$$

$$+\frac{(Ab^2-abB+3a^2C+2b^2C)d\sqrt{c+d \tan(e+fx)}}{b^2(a^2+b^2)f}$$

$$-\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{b(a^2+b^2)f(a+b \tan(e+fx))}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(
a-I*b)^2/f-(B-I*(A-C))*(c+I*d)^(3/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)
)^(1/2))/(a+I*b)^2/f+(-a*d+b*c)^(1/2)*(a^3*b*B*d-3*a^4*C*d-b^4*(3*A*d+2*B*
c)-a*b^3*(4*A*c-5*B*d-4*C*c)+a^2*b^2*(2*B*c+(A-7*C)*d))*arctanh(b^(1/2)*(c
+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/b^(5/2)/(a^2+b^2)^2/f+(A*b^2-B*a*b+
3*C*a^2+2*C*b^2)*d*(c+d*tan(f*x+e))^(1/2)/b^2/(a^2+b^2)/f-(A*b^2-a*(B*b-C*
a))*(c+d*tan(f*x+e))^(3/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1732 vs. $2(372) = 744$.

Time = 4.32 (sec) , antiderivative size = 1732, normalized size of antiderivative = 4.66

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^2,x]

```

output

```
(-4*a^2*A*b^3*c*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] + 2*a^3*b^2*B*c*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] - 2*a*b^4*B*c*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] + 4*a^2*b^3*c*C*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] + a^3*A*b^2*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] - 3*a*A*b^4*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] + a^4*b*B*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] + 5*a^2*b^3*B*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] - 3*a^5*C*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] - 7*a^3*b^2*C*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]] - 4*a*A*b^4*c*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]*Tan[e + f*x] + 2*a^2*b^3*B*c*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]*Tan[e + f*x] - 2*b^5*B*c*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]*Tan[e + f*x] + 4*a*b^4*c*C*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]*Tan[e + f*x] + a^2*A*b^3*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]*Tan[e + f*x] - 3*A*b^5*d*Sqrt[b*c - a*d]*ArcTanh[(Sqrt[b]*Sqrt[c + d*...
```

Rubi [A] (warning: unable to verify)

Time = 5.82 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.04, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.404$, Rules used = {3042, 4128, 27, 3042, 4130, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^2} dx$$

↓ 4128

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left((3Ca^2 - bBa + Ab^2 + 2b^2C) d \tan^2(e+fx) - 2b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + 2(bB - aC) \left(bc - \frac{3ad}{2} \right) + 2Ab \left(ac + \frac{3bd}{2} \right) \right)}{2(a+b \tan(e+fx))} \frac{b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}} \frac{1}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left((3Ca^2 - bBa + Ab^2 + 2b^2C) d \tan^2(e+fx) - 2b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB - aC)(2bc - 3ad) + Ab(2ac + 3bd) \right)}{a+b \tan(e+fx)} \frac{2b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}} \frac{1}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left((3Ca^2 - bBa + Ab^2 + 2b^2C) d \tan(e+fx)^2 - 2b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB - aC)(2bc - 3ad) + Ab(2ac + 3bd) \right)}{a+b \tan(e+fx)} \frac{2b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}} \frac{1}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 4130

$$2 \int - \frac{-2(2aAc d - 2acC d - Ab(c^2 - d^2) + aB(c^2 - d^2) + b(Cc^2 + 2Bdc - Cd^2)) \tan(e+fx)b^2 - c((bB - aC)(2bc - 3ad) + Ab(2ac + 3bd))b + a(3Ca^2 - bBa + Ab^2 + 2b^2C) d}{2(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} \frac{1}{b}$$

$2b(a^2 + b^2)$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 27

$$\frac{2d(3a^2C - abB + Ab^2 + 2b^2C) \sqrt{c+d \tan(e+fx)}}{bf} - \int \frac{-2(2aAc d - 2acC d - Ab(c^2 - d^2) + aB(c^2 - d^2) + b(Cc^2 + 2Bdc - Cd^2)) \tan(e+fx)b^2 - c((bB - aC)(2bc - 3ad) + Ab(2ac + 3bd))b + a(3Ca^2 - bBa + Ab^2 + 2b^2C) d}{2(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} \frac{1}{b}$$

$2b(a^2 + b^2)$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 3042

$$\frac{2d(3a^2C-abB+Ab^2+2b^2C)\sqrt{c+d\tan(e+fx)}}{bf} - \int \frac{-2(2aAcd-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2))\tan(e+fx)b^2-c((bB-aC)(2bc-2b^2))}{2b(a^2+b^2)}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 4136

$$\frac{2d(3a^2C-abB+Ab^2+2b^2C)\sqrt{c+d\tan(e+fx)}}{bf} - \int \frac{2(b^2((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^2-2b(2c(A-C)d+B(c^2-d^2))a-b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))))}{2b(a^2+b^2)}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 27

$$\frac{2d(3a^2C-abB+Ab^2+2b^2C)\sqrt{c+d\tan(e+fx)}}{bf} - \int \frac{b^2((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^2-2b(2c(A-C)d+B(c^2-d^2))a-b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))))}{2b(a^2+b^2)}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 3042

$$\frac{2d(3a^2C-abB+Ab^2+2b^2C)\sqrt{c+d\tan(e+fx)}}{bf} - \int \frac{b^2((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^2-2b(2c(A-C)d+B(c^2-d^2))a-b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))))}{2b(a^2+b^2)}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 4022

$$- \frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))} +$$

$$\frac{2d(3a^2C-abB+Ab^2+2b^2C)\sqrt{c+d\tan(e+fx)}}{bf} - \frac{(bc-ad)(-3a^4Cd+a^3bBd+a^2b^2(d(A-7C)+2Bc)-ab^3(4Ac-5Bd-4cC)-b^4(3Ad+2Bc))\int \frac{\tan(e+fx)}{a^2+b^2}}{a^2+b^2}$$

$2b(a^2 + b^2)$

↓ 3042

$$\begin{aligned}
 & -\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \\
 & \frac{2d(3a^2C - abB + Ab^2 + 2b^2C)\sqrt{c + d \tan(e + fx)}}{bf} - \frac{(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{\tan(e + fx)}{a + b \tan(e + fx)}}{a^2 + b^2}
 \end{aligned}$$

$2b(a^2 + b^2)$

↓ 4020

$$\begin{aligned}
 & -\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \\
 & \frac{2d(3a^2C - abB + Ab^2 + 2b^2C)\sqrt{c + d \tan(e + fx)}}{bf} - \frac{(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{\tan(e + fx)}{a + b \tan(e + fx)}}{a^2 + b^2}
 \end{aligned}$$

↓ 25

$$\begin{aligned}
 & -\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \\
 & \frac{2d(3a^2C - abB + Ab^2 + 2b^2C)\sqrt{c + d \tan(e + fx)}}{bf} - \frac{(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{\tan(e + fx)}{a + b \tan(e + fx)}}{a^2 + b^2}
 \end{aligned}$$

↓ 73

$$\begin{aligned}
 & -\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \\
 & \frac{2d(3a^2C - abB + Ab^2 + 2b^2C)\sqrt{c + d \tan(e + fx)}}{bf} - \frac{(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{\tan(e + fx)}{a + b \tan(e + fx)}}{a^2 + b^2}
 \end{aligned}$$

↓ 221

$$\begin{aligned}
 & -\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \\
 & \frac{2d(3a^2C - abB + Ab^2 + 2b^2C)\sqrt{c + d \tan(e + fx)}}{bf} - \frac{(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{\tan(e + fx)}{a + b \tan(e + fx)}}{a^2 + b^2}
 \end{aligned}$$

$2b(a^2 + b^2)$

↓ 4117

$$-\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{2d(3a^2C - abB + Ab^2 + 2b^2C)\sqrt{c + d \tan(e + fx)}}{bf} - \frac{(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{dx}{(a + b \tan(e + fx))}}{f(a^2 + b^2)}$$

2b(a^2 +

73

$$-\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{2(bc - ad)(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \int \frac{dx}{a + b \tan(e + fx)}}{df(a^2 + b^2)}$$

2b(a^2 +

221

$$-\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{3/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))} + \frac{2\sqrt{bc - ad}(-3a^4Cd + a^3bBd + a^2b^2(d(A - 7C) + 2Bc) - ab^3(4Ac - 5Bd - 4cC) - b^4(3Ad + 2Bc)) \operatorname{arctanh}\left(\frac{c + d \tan(e + fx)}{a + b \tan(e + fx)}\right)}{\sqrt{bf}(a^2 + b^2)}$$

2b(a^2 + b^2)

input `Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^2,x]`

output `-(((A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(3/2))/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x]))) + (-(((2*(-(((a + I*b)^2*b^2*(A - I*B - C)*(c - I*d)^(3/2)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f) - ((a - I*b)^2*b^2*(A + I*B - C)*(c + I*d)^(3/2)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f)))/(a^2 + b^2) - (2*Sqrt[b*c - a*d]*(a^3*b*B*d - 3*a^4*C*d - b^4*(2*B*c + 3*A*d) - a*b^3*(4*A*c - 4*c*C - 5*B*d) + a^2*b^2*(2*B*c + (A - 7*C)*d))*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/(Sqrt[b]*(a^2 + b^2)*f)/b) + (2*(A*b^2 - a*b*B + 3*a^2*C + 2*b^2*C)*d*Sqrt[c + d*Tan[e + f*x]]/(b*f))/(2*b*(a^2 + b^2))`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{\text{n}_}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{1/\text{p}}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{LtQ}[-1, \text{m}, 0] \ \&\& \ \text{LeQ}[-1, \text{n}, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \ \&\& \ \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)]), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d})*\text{x})^{\text{m}}/(\text{d}^2 + \text{c}*\text{x}), \text{x}], \text{x}, \text{d}*\text{Tan}[\text{e} + \text{f}*\text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 - \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 + \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!IntegerQ}[\text{m}]$

rule 4117

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :>
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)])], x_Symbol] :> Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 9864 vs. $2(337) = 674$.

Time = 0.22 (sec) , antiderivative size = 9865, normalized size of antiderivative = 26.52

method	result	size
derivativeldivides	Expression too large to display	9865
default	Expression too large to display	9865

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")
```

output

```
Timed out
```

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

input

```
integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**2,x)
```

output

```
Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)*
*2)/(a + b*tan(e + f*x))**2, x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for m
ore detail
```

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Exception raised: TypeError}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^2,x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{%%{-1, [0,19,8]%%}+%%{-8, [0,17,8]%%}+%%{-28, [0,15,8]%%
%}+%%{-5
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^2,x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2*c*d + 4*sqrt(tan(e + f*x)*d + c)*a*b*c
**2 + 2*sqrt(tan(e + f*x)*d + c)*a*b*d**2 + 2*sqrt(tan(e + f*x)*d + c)*b**
2*c*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f
*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*
x)*a*b*c + a**2*c),x)*tan(e + f*x)*a**3*b*c*d**2*f - 3*int(sqrt(tan(e + f*
x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)
**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e
 + f*x)*a**2*b**2*c**2*d*f + 2*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)*
**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)
)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*a*b**3*c**3*f +
int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a
*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c
 + a**2*c),x)*a**4*c*d**2*f - 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)
**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*
x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a**3*b*c**2*d*f + 2*int(sqrt
(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + t
an(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*
c),x)*a**2*b**2*c**3*f - int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*a*b
**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*a**2*b*d**2 - 3*
tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**2 + tan(e + f*x)...
```


3.103
$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

Optimal result	1164
Mathematica [B] (verified)	1165
Rubi [A] (warning: unable to verify)	1165
Maple [B] (verified)	1172
Fricas [F(-1)]	1172
Sympy [F]	1173
Maxima [F(-2)]	1173
Giac [F(-2)]	1173
Mupad [F(-1)]	1174
Reduce [F]	1174

Optimal result

Integrand size = 47, antiderivative size = 532

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx =$$

$$\frac{(A-iB-C)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(ia+b)^3 f}$$

$$+ \frac{(A+iB-C)(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(ia-b)^3 f}$$

$$\frac{(a^5 b B d^2 + 3 a^6 C d^2 + a^4 b^2 d (4 B c + 3 (A + 2 C) d) - b^6 (8 A c^2 - 8 c^2 C - 12 B c d - 3 A d^2) + a^2 b^4 (24 A c^2 - 24 A c B d + 3 a^4 C d + b^4 (4 B c + 3 A d) + a b^3 (8 A c - 8 c C - 7 B d) - a^2 b^2 (4 B c + 5 A d - 11 C d)) \sqrt{c+d \tan(e+fx)}}{4 b^2 (a^2 + b^2)^2 f (a + b \tan(e+fx))}$$

$$\frac{(A b^2 - a (b B - a C)) (c + d \tan(e+fx))^{3/2}}{2 b (a^2 + b^2) f (a + b \tan(e+fx))^2}$$

output

$$\begin{aligned}
& -(A-I*B-C)*(c-I*d)^{(3/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c-I*d)^{(1/2)})/(I* \\
& a+b)^3/f+(A+I*B-C)*(c+I*d)^{(3/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c+I*d)^{(1/2)})/ \\
& (I*a-b)^3/f-1/4*(a^5*b*B*d^2+3*a^6*C*d^2+a^4*b^2*d*(4*B*c+3*(A+2*C)*d) \\
& -b^6*(8*A*c^2-3*A*d^2-12*B*c*d-8*C*c^2)+a^2*b^4*(24*A*c^2-26*A*d^2-48*B*c \\
& *d-24*C*c^2+35*C*d^2)-2*a^3*b^3*(12*c*(A-C)*d+B*(4*c^2-9*d^2))+a*b^5*(40*c \\
& *(A-C)*d+3*B*(8*c^2-5*d^2))*\operatorname{arctanh}(b^{(1/2)}*(c+d*\tan(f*x+e))^{(1/2)}/(-a*d+ \\
& b*c)^{(1/2)})/b^{(5/2)}/(a^2+b^2)^3/(-a*d+b*c)^{(1/2)}/f-1/4*(a^3*b*B*d+3*a^4*C* \\
& d+b^4*(3*A*d+4*B*c)+a*b^3*(8*A*c-7*B*d-8*C*c)-a^2*b^2*(5*A*d+4*B*c-11*C*d) \\
&)*(c+d*\tan(f*x+e))^{(1/2)}/b^2/(a^2+b^2)^2/f/(a+b*\tan(f*x+e))-1/2*(A*b^2-a*(\\
& B*b-C*a))*(c+d*\tan(f*x+e))^{(3/2)}/b/(a^2+b^2)/f/(a+b*\tan(f*x+e))^2
\end{aligned}$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 7678 vs. $2(532) = 1064$.

Time = 6.73 (sec) , antiderivative size = 7678, normalized size of antiderivative = 14.43

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Result too large to show}$$

input

```
Integrate[(((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

Result too large to show

Rubi [A] (warning: unable to verify)

Time = 7.25 (sec) , antiderivative size = 563, normalized size of antiderivative = 1.06, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.404$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-((-3Ca^2 - bBa + Ab^2 - 4b^2C) d \tan^2(e+fx)) - 4b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + 2(bB - aC) \left(2bc - \frac{3ad}{2} \right) + 2Ab \right)}{2(a+b \tan(e+fx))^2} dx$$

$$\frac{2b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}} \frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-((-3Ca^2 - bBa + Ab^2 - 4b^2C) d \tan^2(e+fx)) - 4b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB - aC)(4bc - 3ad) + Ab(4ac + 3bd) \right)}{(a+b \tan(e+fx))^2} dx$$

$$\frac{4b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}} \frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-((-3Ca^2 - bBa + Ab^2 - 4b^2C) d \tan(e+fx)^2) - 4b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB - aC)(4bc - 3ad) + Ab(4ac + 3bd) \right)}{(a+b \tan(e+fx))^2} dx$$

$$\frac{4b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}} \frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 4128

$$\int \frac{-8(ac+bd)((A-C)(bc-ad) - B(ac+bd)) + (bc-ad)(bBc + b(A-C)d + a(Ac - Cc - Bd)) \tan(e+fx) b^2 + 2 \left(ac + \frac{bd}{2} \right) ((bB - aC)(4bc - 3ad) + Ab(4ac + 3bd)) b + d(3Cda^4 + 3Cdb^4)}{2(a+b \tan(e+fx))^2} dx$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 27

$$\int \frac{-8((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+2\left(ac+\frac{bd}{2}\right)((bB-aC)(4bc-3ad)+Ab(4ac+3bd))b+d(3Cda^4)}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{-8((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+2\left(ac+\frac{bd}{2}\right)((bB-aC)(4bc-3ad)+Ab(4ac+3bd))b+d(3Cda^4)}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 4136

$$\int \frac{8\left(b^2\left((C^2+2Bdc-Cd^2-A(c^2-d^2))a^3-3b(2c(A-C)d+B(c^2-d^2))a^2-3b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))a+b^3(2c(A-C)d+B(c^2-d^2))\right)-b^2\left(2c(A-C)d+B(c^2-d^2)\right)\right)}{(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 27

$$\frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bc)-2a^3b^3(12cd(A-C)+B(4c^2-9d^2))+a^2b^4(24Ac^2-26Ad^2-48Bcd-24c^2C+35Cd^2)+ab^5(40cd(A-C)+3B(8c^2-5d^2)))}{a^2+b^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 3042

$$\frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bc)-2a^3b^3(12cd(A-C)+B(4c^2-9d^2))+a^2b^4(24Ac^2-26Ad^2-48Bcd-24c^2C+35Cd^2)+ab^5(40cd(A-C)+3B(8c^2-5d^2)))}{a^2+b^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 4022

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bd))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bd))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 4020

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bd))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 25

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bd))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 73

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bd))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 221

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

4117

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

73

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{2(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

221

$$-\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2} + \frac{\sqrt{c+d \tan(e+fx)}(3a^4Cd+a^3bBd-a^2b^2(5Ad+4Bc-11Cd)+ab^3(8Ac-7Bd-8cC)+b^4(3Ad+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))} + \frac{2(3a^6Cd^2+a^5bBd^2+a^4b^2d(3d(A+2C)+4Bc))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

input

```
Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]
```

output

```

-1/2*((A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(3/2))/(b*(a^2 + b^2)*f
*(a + b*Tan[e + f*x])^2) + (((-8*(-((a + I*b)^3*b^2*(A - I*B - C)*(c - I*
d)^(3/2)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f) - ((a - I*b)^3*b^2*(A + I*
B - C)*(c + I*d)^(3/2)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f))/(a^2 + b^2)
- (2*(a^5*b*B*d^2 + 3*a^6*C*d^2 + a^4*b^2*d*(4*B*c + 3*(A + 2*C)*d) - b^6
*(8*A*c^2 - 8*c^2*C - 12*B*c*d - 3*A*d^2) + a^2*b^4*(24*A*c^2 - 24*c^2*C -
48*B*c*d - 26*A*d^2 + 35*C*d^2) - 2*a^3*b^3*(12*c*(A - C)*d + B*(4*c^2 -
9*d^2)) + a*b^5*(40*c*(A - C)*d + 3*B*(8*c^2 - 5*d^2)))*ArcTanh[(Sqrt[b]*S
qrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]/(Sqrt[b]*(a^2 + b^2)*Sqrt[b*c -
a*d]*f))/(2*b*(a^2 + b^2)) - ((a^3*b*B*d + 3*a^4*C*d + b^4*(4*B*c + 3*A*d
) + a*b^3*(8*A*c - 8*c*C - 7*B*d) - a^2*b^2*(4*B*c + 5*A*d - 11*C*d))*Sqrt
[c + d*Tan[e + f*x]])/(b*(a^2 + b^2)*f*(a + b*Tan[e + f*x]))/(4*b*(a^2 +
b^2))

```

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4020 $\text{Int}[(a_.) + (b_.)\tan[e_.] + (f_.)x_])^m((c_.) + (d_.)\tan[e_.] + (f_.)x_])$, x_Symbol] $\rightarrow \text{Simp}[c(d/f) \text{Subst}[\text{Int}[(a + (b/d)x]^m/(d^2 + cx), x], x, d\tan[e + fx]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[(a_.) + (b_.)\tan[e_.] + (f_.)x_])^m((c_.) + (d_.)\tan[e_.] + (f_.)x_])$, x_Symbol] $\rightarrow \text{Simp}[(c + I*d)/2 \text{Int}[(a + b\tan[e + fx])^m(1 - I\tan[e + fx]), x], x] + \text{Simp}[(c - I*d)/2 \text{Int}[(a + b\tan[e + fx])^m(1 + I\tan[e + fx]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{IntegerQ}[m]$

rule 4117 $\text{Int}[(a_.) + (b_.)\tan[e_.] + (f_.)x_])^m((c_.) + (d_.)\tan[e_.] + (f_.)x_])^n((A_.) + (C_.)\tan[e_.] + (f_.)x_])^2$, x_Symbol] $\rightarrow \text{Simp}[A/f \text{Subst}[\text{Int}[(a + b*x)^m(c + d*x)^n, x], x, \tan[e + fx]], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, C, m, n\}, x] \&\& \text{EqQ}[A, C]$

rule 4128 $\text{Int}[(a_.) + (b_.)\tan[e_.] + (f_.)x_])^m((c_.) + (d_.)\tan[e_.] + (f_.)x_])^n((A_.) + (B_.)\tan[e_.] + (f_.)x_]) + (C_.)\tan[e_.] + (f_.)x_])^2$, x_Symbol] $\rightarrow \text{Simp}[(A*d^2 + c(c*C - B*d))(a + b\tan[e + fx])^m((c + d\tan[e + fx])^{n+1}/(d*f*(n+1)*(c^2 + d^2))), x] - \text{Simp}[1/(d*(n+1)*(c^2 + d^2)) \text{Int}[(a + b\tan[e + fx])^{m-1}(c + d\tan[e + fx])^{n+1} \text{Simp}[A*d*(b*d*m - a*c*(n+1)) + (c*C - B*d)*(b*c*m + a*d*(n+1)) - d*(n+1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*\tan[e + fx] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n+1)))*\tan[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[m, 0] \&\& \text{LtQ}[n, -1]$

rule 4136 $\text{Int}[(c_.) + (d_.)\tan[e_.] + (f_.)x_])^n((A_.) + (B_.)\tan[e_.] + (f_.)x_]) + (C_.)\tan[e_.] + (f_.)x_])^2)/((a_.) + (b_.)\tan[e_.] + (f_.)x_])$, x_Symbol] $\rightarrow \text{Simp}[1/(a^2 + b^2) \text{Int}[(c + d\tan[e + fx])^n \text{Simp}[b*B + a*(A - C) + (a*B - b*(A - C))*\tan[e + fx], x], x], x] + \text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{Int}[(c + d\tan[e + fx])^n((1 + \tan[e + fx]^2)/(a + b\tan[e + fx])), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[n, 0] \&\& \text{LeQ}[n, -1]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 14440 vs. $2(492) = 984$.

Time = 0.24 (sec) , antiderivative size = 14441, normalized size of antiderivative = 27.14

method	result	size
derivativedivides	Expression too large to display	14441
default	Expression too large to display	14441

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)
```

output

result too large to display

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**3,x)`

output `Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x))**3, x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Exception raised: TypeError}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1,[0,19,8]}%%+%%{8,[0,17,8]}%%+%%{28,[0,15,8]}%%+
%%{56,[0
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x))^3,x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^3,x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2*c*d + 8*sqrt(tan(e + f*x)*d + c)*a*b*c
**2 + 6*sqrt(tan(e + f*x)*d + c)*a*b*d**2 + 6*sqrt(tan(e + f*x)*d + c)*b**
2*c*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f
*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*
tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c +
a**3*c),x)*tan(e + f*x)**2*a**3*b**2*c*d**2*f - 5*int(sqrt(tan(e + f*x)*d
+ c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)*
**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(
e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a**
2*b**3*c**2*d*f + 4*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d +
3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a
**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x
)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a*b**4*c**3*f + 2*int(sqrt(tan(e +
f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e
+ f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c
+ tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)
*a**4*b*c*d**2*f - 10*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d
+ 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2
*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f
*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)*a**3*b**2*c**2*d*f + 8*int(sqrt(...
```

3.104 $\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^{5/2} (A + B \tan(e+fx) + C \tan^2(e+fx)) dx =$

Optimal result	1176
Mathematica [A] (verified)	1177
Rubi [A] (warning: unable to verify)	1178
Maple [B] (verified)	1184
Fricas [B] (verification not implemented)	1184
Sympy [F]	1185
Maxima [F(-1)]	1185
Giac [F(-2)]	1186
Mupad [F(-1)]	1186
Reduce [F]	1187

Optimal result

Integrand size = 47, antiderivative size = 503

$$\int (a+b \tan(e+fx))^2(c+d \tan(e+fx))^{5/2} (A + B \tan(e+fx) + C \tan^2(e+fx)) dx =$$

$$\frac{(a-ib)^2(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f}$$

$$+ \frac{(a+ib)^2(iA-B-iC)(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f}$$

$$- \frac{2(2ab(c^2C+2Bcd-Cd^2-A(c^2-d^2))-a^2(2c(A-C)d+B(c^2-d^2))+b^2(2c(A-C)d+B(c^2-d^2)))}{f}$$

$$+ \frac{2(2ab(Ac-cC-Bd)+a^2(Bc+(A-C)d)-b^2(Bc+(A-C)d))(c+d \tan(e+fx))^{3/2}}{3f}$$

$$+ \frac{2(a^2B-b^2B+2ab(A-C))(c+d \tan(e+fx))^{5/2}}{5f}$$

$$+ \frac{2(36a^2Cd^2-22abd(2cC-9Bd)+b^2(8c^2C-22Bcd+99(A-C)d^2))(c+d \tan(e+fx))^{7/2}}{693d^3f}$$

$$- \frac{2b(4bcC-11bBd-4aCd) \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{99d^2f}$$

$$+ \frac{2C(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{7/2}}{11df}$$

output

```

-(a-I*b)^2*(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)
)^(1/2))/f+(a+I*b)^2*(I*A-B-I*C)*(c+I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1
/2)/(c+I*d)^(1/2))/f-2*(2*a*b*(c^2*C+2*B*c*d-C*d^2-A*(c^2-d^2))-a^2*(2*c*(
A-C)*d+B*(c^2-d^2))+b^2*(2*c*(A-C)*d+B*(c^2-d^2)))*(c+d*tan(f*x+e))^(1/2)/
f+2/3*(2*a*b*(A*c-B*d-C*c)+a^2*(B*c+(A-C)*d)-b^2*(B*c+(A-C)*d))*(c+d*tan(f
*x+e))^(3/2)/f+2/5*(B*a^2-B*b^2+2*a*b*(A-C))*(c+d*tan(f*x+e))^(5/2)/f+2/69
3*(36*a^2*C*d^2-22*a*b*d*(-9*B*d+2*C*c)+b^2*(8*c^2*C-22*B*c*d+99*(A-C)*d^2
))*(c+d*tan(f*x+e))^(7/2)/d^3/f-2/99*b*(-11*B*b*d-4*C*a*d+4*C*b*c)*tan(f*x
+e)*(c+d*tan(f*x+e))^(7/2)/d^2/f+2/11*C*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e)
)^(7/2)/d/f
    
```

Mathematica [A] (verified)

Time = 6.36 (sec) , antiderivative size = 564, normalized size of antiderivative = 1.12

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{2C(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{7/2}}{11df}$$

$$+ 2 \left(\frac{b(-4bcC + 11bBd + 4aCd) \tan(e + fx) (c + d \tan(e + fx))^{7/2}}{9df} - \frac{2 \left(\frac{(-36a^2Cd^2 + 22abd(2cC - 9Bd) - b^2(8c^2C - 22Bcd + 99(A-C)d^2)) (c + d \tan(e + fx))^{7/2}}{14df} \right)}{11df} \right)$$

input

```

Integrate[(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e +
f*x] + C*Tan[e + f*x]^2),x]
    
```

output

```
(2*C*(a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(7/2))/(11*d*f) + (2*((b*
(-4*b*c*C + 11*b*B*d + 4*a*C*d)*Tan[e + f*x]*(c + d*Tan[e + f*x])^(7/2))/(
9*d*f) - (2*(((-36*a^2*C*d^2 + 22*a*b*d*(2*c*C - 9*B*d) - b^2*(8*c^2*C - 2
2*B*c*d + 99*(A - C)*d^2))*(c + d*Tan[e + f*x])^(7/2))/(14*d*f) + ((I/2)*(
((99*I)/4)*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2 + (99*(2*a*b*B - a^2*(A - C
) + b^2*(A - C))*d^2)/4)*((2*(c + d*Tan[e + f*x])^(5/2))/5 + (c - I*d)*((2
*(c + d*Tan[e + f*x])^(3/2))/3 + (c - I*d)*((2*(c - I*d)^(3/2)*ArcTanh[Sqr
t[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]))/(-c + I*d) + 2*Sqrt[c + d*Tan[e + f*
x]])))/f - ((I/2)*((-99*I)/4)*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2 + (99*
(2*a*b*B - a^2*(A - C) + b^2*(A - C))*d^2)/4)*((2*(c + d*Tan[e + f*x])^(5/
2))/5 + (c + I*d)*((2*(c + d*Tan[e + f*x])^(3/2))/3 + (c + I*d)*((2*(c + I
*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]]))/(-c - I*d) + 2*
Sqrt[c + d*Tan[e + f*x]])))/f))/(9*d))/(11*d)
```

Rubi [A] (warning: unable to verify)

Time = 6.18 (sec) , antiderivative size = 511, normalized size of antiderivative = 1.02, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\frac{2 \int -\frac{1}{2} (a + b \tan(e + fx)) (c + d \tan(e + fx))^{5/2} ((4bcC - 4adC - 11bBd) \tan^2(e + fx) - 11(Ab - Cb + aB) d \tan(e + fx) + 11d)}{11df} dx}{11df}$$

$$\downarrow 27$$

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} ((4bcC - 4adC - 11bBd) \tan^2(e + fx) - 11(Ab - Cb + aB)d \tan(e + fx)) dx}{11d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} ((4bcC - 4adC - 11bBd) \tan(e + fx)^2 - 11(Ab - Cb + aB)d \tan(e + fx)) dx}{11d}$$

↓ 4120

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{2 \int -\frac{1}{2}(c + d \tan(e + fx))^{5/2} (-2c(4cC - 11Bd)b^2 + 44acCdb - 9a^2(11A - 7C)d^2 - ((8c^2C - 22Bcd)b^2 + 44acCdb - 9a^2(11A - 7C)d^2 - ((8C^2 - 22Bdc + 99(A - C)d^2)b^2 - 22ad(2cC - 9Bd)b + 36a^2Cd^2) \tan(e + fx)) dx}{9d}}{11d}$$

↓ 27

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{\int (c + d \tan(e + fx))^{5/2} (-((8c^2C - 22Bcd)b^2 + 44acCdb - 9a^2(11A - 7C)d^2 - ((8C^2 - 22Bdc + 99(A - C)d^2)b^2 - 22ad(2cC - 9Bd)b + 36a^2Cd^2) \tan(e + fx)) dx}{9d}}{11d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{\int (c + d \tan(e + fx))^{5/2} (-((8c^2C - 22Bcd)b^2 + 44acCdb - 9a^2(11A - 7C)d^2 - ((8C^2 - 22Bdc + 99(A - C)d^2)b^2 - 22ad(2cC - 9Bd)b + 36a^2Cd^2) \tan(e + fx)) dx}{9d}}{11d}$$

↓ 4113

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{\int (c + d \tan(e + fx))^{5/2} (99(-((A - C)a^2 + 2bBa + b^2(A - C))d^2 - 99(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)) dx - \frac{2(c + d \tan(e + fx))^{7/2} (36a^2Cd^2 - 22ad(2cC - 9Bd)b + 36a^2Cd^2)}{9d}}{9d}}{11d}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{\int (c + d \tan(e + fx))^{5/2} (99(-((A - C)a^2 + 2bBa + b^2(A - C))d^2 - 99(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)) dx - \frac{2(c + d \tan(e + fx))^{7/2} (36a^2Cd^2 - 22ad(2cC - 9Bd)b + 36a^2Cd^2)}{9d}}{9d}}{11d}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \\ & \int (c + d \tan(e + fx))^{3/2} (-99((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd))d^2 - 99((Bc + (A - C)d)a^2 + 2b(Ac - Cc - Bd)a - b^2(Bc + (A - C)d))d) \sqrt{c + d \tan(e + fx)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \\ & \int (c + d \tan(e + fx))^{3/2} (-99((Ac - Cc - Bd)a^2 - 2b(Bc + (A - C)d)a - b^2(Ac - Cc - Bd))d^2 - 99((Bc + (A - C)d)a^2 + 2b(Ac - Cc - Bd)a - b^2(Bc + (A - C)d))d) \sqrt{c + d \tan(e + fx)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \\ & \int \sqrt{c + d \tan(e + fx)} (99((C^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a^2 + 2b(2c(A - C)d + B(c^2 - d^2))a - b^2(C^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))d^2 + 99(-((2c(A - C)d + B(c^2 - d^2))a - b^2(C^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))) \sqrt{c + d \tan(e + fx)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \\ & \int \sqrt{c + d \tan(e + fx)} (99((C^2 + 2Bdc - Cd^2 - A(c^2 - d^2))a^2 + 2b(2c(A - C)d + B(c^2 - d^2))a - b^2(C^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))d^2 + 99(-((2c(A - C)d + B(c^2 - d^2))a - b^2(C^2 + 2Bdc - Cd^2 - A(c^2 - d^2)))) \sqrt{c + d \tan(e + fx)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \\ & \int \frac{99d^2(-((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))a^2) + 2b(Cc^3 + 3Bdc^2 - 3Cd^2c - Bd^3 - A(c^3 - 3cd^2))a + b^2((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) \tan(e + fx) - 99d^2}{\sqrt{c + d \tan(e + fx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \\ & \int \frac{99d^2(-((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))a^2) + 2b(Cc^3 + 3Bdc^2 - 3Cd^2c - Bd^3 - A(c^3 - 3cd^2))a + b^2((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) \tan(e + fx) - 99d^2}{\sqrt{c + d \tan(e + fx)}} \end{aligned}$$

$$\downarrow 4022$$

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} + \frac{-\frac{99}{2}d^2(a+ib)^2(c+id)^3(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{99}{2}d^2(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{11df}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} + \frac{-\frac{99}{2}d^2(a+ib)^2(c+id)^3(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{99}{2}d^2(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{11df}$$

↓ 4020

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} + \frac{99id^2(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - \frac{99i}{2}d^2(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{11df}$$

↓ 25

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} + \frac{99id^2(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - \frac{99i}{2}d^2(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{11df}$$

↓ 73

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} + \frac{99d(a+ib)^2(c+id)^3(A+iB-C) \int \frac{1}{f \frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)} - \frac{99d(a-ib)^2(c-id)^3(A-iB-C) \int \frac{1}{f \frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{11df}$$

↓ 221

$$\frac{2C(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{7/2}}{11df} - \frac{2b \tan(e + fx)(-4aCd - 11bBd + 4bcC)(c + d \tan(e + fx))^{7/2}}{9df} + \frac{2(c+d \tan(e+fx))^{7/2}(36a^2Cd^2 - 22abd(2cC - 9Bd) + b^2(99d^2(A-C) - 22Bcd + 8c^2C))}{7df}$$

input $\text{Int}[(a + b*\text{Tan}[e + f*x])^2*(c + d*\text{Tan}[e + f*x])^{(5/2)}*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2), x]$

output $(2*C*(a + b*\text{Tan}[e + f*x])^2*(c + d*\text{Tan}[e + f*x])^{(7/2)})/(11*d*f) - ((2*b*(4*b*c*C - 11*b*B*d - 4*a*C*d)*\text{Tan}[e + f*x]*(c + d*\text{Tan}[e + f*x])^{(7/2)})/(9*d*f) + ((-99*(a - I*b)^2*(A - I*B - C)*(c - I*d)^{(5/2)}*d^2*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]])/f - (99*(a + I*b)^2*(A + I*B - C)*(c + I*d)^{(5/2)}*d^2*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c + I*d]])/f + (198*d^2*(2*a*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - a^2*(2*c*(A - C)*d + B*(c^2 - d^2)) + b^2*(2*c*(A - C)*d + B*(c^2 - d^2)))*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/f - (66*d^2*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d) - b^2*(B*c + (A - C)*d))*(c + d*\text{Tan}[e + f*x])^{(3/2)})/f - (198*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2*(c + d*\text{Tan}[e + f*x])^{(5/2)})/(5*f) - (2*(36*a^2*C*d^2 - 22*a*b*d*(2*c*C - 9*B*d) + b^2*(8*c^2*C - 22*B*c*d + 99*(A - C)*d^2))*(c + d*\text{Tan}[e + f*x])^{(7/2)})/(7*d*f))/(9*d))/(11*d)$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_*)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$

rule 73 $\text{Int}[((a_*) + (b_*)*(x_*)^{(m_*)})*((c_*) + (d_*)*(x_*)^{(n_*)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)}*(c - a*(d/b) + d*(x^p/b)^n], x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_*) + (b_*)*(x_*)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^(m*(1 - I*Tan[e + f*x])), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^(m*(1 + I*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

rule 4130

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 11345 vs. $2(459) = 918$.

Time = 0.54 (sec) , antiderivative size = 11346, normalized size of antiderivative = 22.56

method	result	size
parts	Expression too large to display	11346
derivativedivides	Expression too large to display	11478
default	Expression too large to display	11478

input

```

int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x,method=_RETURNVERBOSE)

```

output

result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 91140 vs. $2(449) = 898$.

Time = 160.30 (sec) , antiderivative size = 91140, normalized size of antiderivative = 181.19

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**2*(c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**2*(c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output Timed out

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%}{1, [0,34,13]%%}+%%{14, [0,32,13]%%}+%%{91, [0,30,13]%%}+%%{3`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Hanged}$$

input `int((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int (a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c),x)*a**3*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**6,x)*b**2*c*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5,x)*a*b*c*d**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5,x)*b**3*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a**2*c*d**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a**2*c*d**2 + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a*b**2*d**2 + 4*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*a*b*c**2*d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**3*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**2*c**3 + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a**2*b*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a**2*c**2*d + 6*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*b**2*c*d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*b*c**3 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b**3*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**3*d**2 + 6*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*b*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*c**3 + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*b**2*c**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**3*c*d + 3*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**2*b*c**2
```


3.105 $\int (a+b \tan(e+fx))(c+d \tan(e+fx))^{5/2} (A + B \tan(e+fx)) dx$

Optimal result	1188
Mathematica [A] (verified)	1189
Rubi [A] (warning: unable to verify)	1190
Maple [B] (verified)	1195
Fricas [B] (verification not implemented)	1195
Sympy [F]	1196
Maxima [F(-1)]	1196
Giac [F(-2)]	1197
Mupad [F(-1)]	1197
Reduce [F]	1198

Optimal result

Integrand size = 45, antiderivative size = 353

$$\begin{aligned}
 & \int (a+b \tan(e+fx))(c+d \tan(e+fx))^{5/2} (A + B \tan(e+fx) + C \tan^2(e+fx)) dx = \\
 & \frac{(ia+b)(A-iB-C)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f} \\
 & + \frac{(ia-b)(A+iB-C)(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f} \\
 & + \frac{2(a(Bc^2 - 2cCd - Bd^2) - b(c^2C + 2Bcd - Cd^2) + A(2acd + b(c^2 - d^2))) \sqrt{c+d \tan(e+fx)}}{f} \\
 & + \frac{2(Abc + aBc - bcC + aAd - bBd - aCd)(c+d \tan(e+fx))^{3/2}}{3f} \\
 & + \frac{2(Ab + aB - bC)(c+d \tan(e+fx))^{5/2}}{5f} \\
 & - \frac{2(2bcC - 9bBd - 9aCd)(c+d \tan(e+fx))^{7/2}}{63d^2 f} \\
 & + \frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df}
 \end{aligned}$$

output

$$\begin{aligned}
& -(I*a+b)*(A-I*B-C)*(c-I*d)^{(5/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c-I*d)^{(1/2)})/f+(I*a-b)*(A+I*B-C)*(c+I*d)^{(5/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c+I*d)^{(1/2)})/f+2*(a*(B*c^2-B*d^2-2*C*c*d)-b*(2*B*c*d+C*c^2-C*d^2)+A*(2*a*c*d+b*(c^2-d^2)))*(c+d*\tan(f*x+e))^{(1/2)}/f+2/3*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)*(c+d*\tan(f*x+e))^{(3/2)}/f+2/5*(A*b+B*a-C*b)*(c+d*\tan(f*x+e))^{(5/2)}/f-2/63*(-9*B*b*d-9*C*a*d+2*C*b*c)*(c+d*\tan(f*x+e))^{(7/2)}/d^2/f+2/9*b*C*\tan(f*x+e)*(c+d*\tan(f*x+e))^{(7/2)}/d/f
\end{aligned}$$
Mathematica [A] (verified)

Time = 3.52 (sec) , antiderivative size = 324, normalized size of antiderivative = 0.92

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{2(-2bcC+9bBd+9aCd)(c+d \tan(e+fx))^{7/2}}{d} + 14bC \tan(e + fx)(c + d \tan(e + fx))^{7/2} + \frac{63}{2}i(d$$

input

$$\text{Integrate}[(a + b*\text{Tan}[e + f*x])*(c + d*\text{Tan}[e + f*x])^{(5/2)}*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2),x]$$

output

$$\begin{aligned}
& ((2*(-2*b*c*C + 9*b*B*d + 9*a*C*d)*(c + d*\text{Tan}[e + f*x])^{(7/2)})/d + 14*b*C*\text{Tan}[e + f*x]*(c + d*\text{Tan}[e + f*x])^{(7/2)} + ((63*I)/2)*(a - I*b)*(A - I*B - C)*d*((2*(c + d*\text{Tan}[e + f*x])^{(5/2)})/5 + (2*(c - I*d)*(-3*(c - I*d)^{(3/2)}*\text{ArcTanh}[\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[c - I*d]] + \text{Sqrt}[c + d*\text{Tan}[e + f*x]]*(4*c - (3*I)*d + d*\text{Tan}[e + f*x])))/3) - ((63*I)/2)*(a + I*b)*(A + I*B - C)*d*((2*(c + d*\text{Tan}[e + f*x])^{(5/2)})/5 + (2*(c + I*d)*(-3*(c + I*d)^{(3/2)}*\text{ArcTanh}[\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[c + I*d]] + \text{Sqrt}[c + d*\text{Tan}[e + f*x]]*(4*c + (3*I)*d + d*\text{Tan}[e + f*x])))/3)/(63*d*f)
\end{aligned}$$

Rubi [A] (warning: unable to verify)

Time = 3.67 (sec) , antiderivative size = 345, normalized size of antiderivative = 0.98, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 & \quad \downarrow 3042 \\
 & \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx \\
 & \quad \downarrow 4120 \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \\
 & \frac{2 \int \frac{1}{2}(c + d \tan(e + fx))^{5/2} ((2bcC - 9adC - 9bBd) \tan^2(e + fx) - 9(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 9aAa)}{9d} dx \\
 & \quad \downarrow 27 \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \\
 & \frac{\int (c + d \tan(e + fx))^{5/2} ((2bcC - 9adC - 9bBd) \tan^2(e + fx) - 9(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 9aAa)}{9d} dx \\
 & \quad \downarrow 3042 \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \\
 & \frac{\int (c + d \tan(e + fx))^{5/2} ((2bcC - 9adC - 9bBd) \tan(e + fx)^2 - 9(Ab - Cb + aB)d \tan(e + fx) + 2bcC - 9aAa)}{9d} dx \\
 & \quad \downarrow 4113 \\
 & \frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \\
 & \frac{\int (c + d \tan(e + fx))^{5/2} (9(bB - a(A - C))d - 9(Ab - Cb + aB)d \tan(e + fx)) dx + \frac{2(-9aCd - 9bBd + 2bcC)(c + d \tan(e + fx))^{7/2}}{7df}}{9d} \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int (c+d \tan(e+fx))^{5/2} (9(bB-a(A-C))d - 9(Ab-Cb+aB)d \tan(e+fx)) dx + \frac{2(-9aCd-9bBd+2bcC)(c+d \tan(e+fx))^{3/2}}{7df}}{9d}$$

↓ 4011

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int (c+d \tan(e+fx))^{3/2} (9d(bBc+b(A-C)d - a(Ac-Cc-Bd)) - 9d(Abc+aBc-bCc+aAd-bBd-aCd)) dx}{9d}$$

↓ 3042

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int (c+d \tan(e+fx))^{3/2} (9d(bBc+b(A-C)d - a(Ac-Cc-Bd)) - 9d(Abc+aBc-bCc+aAd-bBd-aCd)) dx}{9d}$$

↓ 4011

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int \sqrt{c+d \tan(e+fx)} (9d(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2)) + b(2c(A-C)d + B(c^2-d^2))) - 9d(2aAc-d)) dx}{9d}$$

↓ 3042

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int \sqrt{c+d \tan(e+fx)} (9d(a(Cc^2+2Bdc-Cd^2-A(c^2-d^2)) + b(2c(A-C)d + B(c^2-d^2))) - 9d(2aAc-d)) dx}{9d}$$

↓ 4011

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int \frac{9d(a(Cc^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2)) + b((A-C)d(3c^2-d^2) + B(c^3-3cd^2))) - 9d(A(bc^3+3adc^2-3bd^2c-ad^3)) - b(Cc^3+3Bdc^2-3Ca)}{\sqrt{c+d \tan(e+fx)}} dx}{9d}$$

↓ 3042

$$\frac{\frac{2bC \tan(e+fx)(c+d \tan(e+fx))^{7/2}}{9df} - \int \frac{9d(a(Cc^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2)) + b((A-C)d(3c^2-d^2) + B(c^3-3cd^2))) - 9d(A(bc^3+3adc^2-3bd^2c-ad^3)) - b(Cc^3+3Bdc^2-3Ca)}{\sqrt{c+d \tan(e+fx)}} dx}{9d}$$

↓ 4022

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{-\frac{9}{2}d(a + ib)(c + id)^3(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{9}{2}d(a - ib)(c - id)^3(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx}{-}$$

↓ 3042

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{-\frac{9}{2}d(a + ib)(c + id)^3(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{9}{2}d(a - ib)(c - id)^3(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx}{-}$$

↓ 4020

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{-\frac{9id(a - ib)(c - id)^3(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} + \frac{9id(a + ib)(c + id)^3(A + iB - C) \int -\frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}}}{2f}}{-}$$

↓ 25

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{9id(a - ib)(c - id)^3(A - iB - C) \int \frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \frac{9id(a + ib)(c + id)^3(A + iB - C) \int \frac{1}{(i \tan(e + fx) + 1)\sqrt{c + d \tan(e + fx)}}}{2f}}{-}$$

↓ 73

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{9(a + ib)(c + id)^3(A + iB - C) \int \frac{1}{-i \tan^2(e + fx) - \frac{ic}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{f} - \frac{9(a - ib)(c - id)^3(A - iB - C) \int \frac{1}{i \tan^2(e + fx) + \frac{id}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{f}}{-}$$

↓ 221

$$\frac{2bC \tan(e + fx)(c + d \tan(e + fx))^{7/2}}{9df} - \frac{9d(a - ib)(c - id)^{5/2}(A - iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f} - \frac{9d(a + ib)(c + id)^{5/2}(A + iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c + id}}\right)}{f} - \frac{18d\sqrt{c + d \tan(e + fx)}(2aAc + d)}{-}$$

input $\text{Int}[(a + b \cdot \tan[e + f \cdot x]) \cdot (c + d \cdot \tan[e + f \cdot x])^{5/2} \cdot (A + B \cdot \tan[e + f \cdot x] + C \cdot \tan[e + f \cdot x]^2), x]$

output $(2 \cdot b \cdot C \cdot \tan[e + f \cdot x] \cdot (c + d \cdot \tan[e + f \cdot x])^{7/2}) / (9 \cdot d \cdot f) - ((-9 \cdot (a - I \cdot b) \cdot (A - I \cdot B - C) \cdot (c - I \cdot d)^{5/2} \cdot d \cdot \text{ArcTan}[\tan[e + f \cdot x] / \text{Sqrt}[c - I \cdot d]]) / f - (9 \cdot (a + I \cdot b) \cdot (A + I \cdot B - C) \cdot (c + I \cdot d)^{5/2} \cdot d \cdot \text{ArcTan}[\tan[e + f \cdot x] / \text{Sqrt}[c + I \cdot d]]) / f - (18 \cdot d \cdot (2 \cdot a \cdot A \cdot c \cdot d - 2 \cdot a \cdot c \cdot C \cdot d + A \cdot b \cdot (c^2 - d^2) + a \cdot B \cdot (c^2 - d^2) - b \cdot (c^2 \cdot C + 2 \cdot B \cdot c \cdot d - C \cdot d^2)) \cdot \text{Sqrt}[c + d \cdot \tan[e + f \cdot x]]) / f - (6 \cdot d \cdot (A \cdot b \cdot c + a \cdot B \cdot c - b \cdot c \cdot C + a \cdot A \cdot d - b \cdot B \cdot d - a \cdot C \cdot d) \cdot (c + d \cdot \tan[e + f \cdot x])^{3/2}) / f - (18 \cdot (A \cdot b + a \cdot B - b \cdot C) \cdot d \cdot (c + d \cdot \tan[e + f \cdot x])^{5/2}) / (5 \cdot f) + (2 \cdot (2 \cdot b \cdot c \cdot C - 9 \cdot b \cdot B \cdot d - 9 \cdot a \cdot C \cdot d) \cdot (c + d \cdot \tan[e + f \cdot x])^{7/2}) / (7 \cdot d \cdot f)) / (9 \cdot d)$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F x_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$

rule 27 $\text{Int}[(a_)(F x_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F x, (b_)(G x_)] /; \text{FreeQ}[b, x]$

rule 73 $\text{Int}[(a_ + (b_)(x_))^{(m_)} \cdot ((c_) + (d_)(x_))^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p \cdot (m + 1) - 1)} \cdot (c - a \cdot (d/b) + d \cdot (x^p/b))^{(n_)}], x, (a + b \cdot x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_ + (b_)(x_)^2)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x / \text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4011 $\text{Int}[\left((a_{\cdot}) + (b_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right)^{(m_{\cdot})}\left((c_{\cdot}) + (d_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[d\left((a + b\tan[e + f*x])^m/(f*m)\right), x] + \text{Int}[(a + b\tan[e + f*x])^{(m-1)}\text{Simp}[a*c - b*d + (b*c + a*d)\tan[e + f*x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

rule 4020 $\text{Int}[\left((a_{\cdot}) + (b_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right)^{(m_{\cdot})}\left((c_{\cdot}) + (d_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[c*(d/f) \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\tan[e + f*x]], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]

rule 4022 $\text{Int}[\left((a_{\cdot}) + (b_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right)^{(m_{\cdot})}\left((c_{\cdot}) + (d_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[(c + I*d)/2 \text{Int}[(a + b*\tan[e + f*x])^m*(1 - I*\tan[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{Int}[(a + b*\tan[e + f*x])^m*(1 + I*\tan[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4113 $\text{Int}[\left((a_{\cdot}) + (b_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right)^{(m_{\cdot})}\left((A_{\cdot}) + (B_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})] + (C_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]^2\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[C\left((a + b*\tan[e + f*x])^{(m+1)}/(b*f*(m+1))\right), x] + \text{Int}[(a + b*\tan[e + f*x])^m\text{Simp}[A - C + B*\tan[e + f*x], x], x] /;$ FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]

rule 4120 $\text{Int}[\left((a_{\cdot}) + (b_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]\right)^{(n_{\cdot})}\left((A_{\cdot}) + (B_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})] + (C_{\cdot})\tan[(e_{\cdot}) + (f_{\cdot})(x_{\cdot})]^2\right), x_{\text{Symbol}}] \rightarrow \text{Simp}[b*C*\tan[e + f*x]\left((c + d*\tan[e + f*x])^{(n+1)}/(d*f*(n+2))\right), x] - \text{Simp}[1/(d*(n+2)) \text{Int}[(c + d*\tan[e + f*x])^n\text{Simp}[b*c*C - a*A*d*(n+2) - (A*b + a*B - b*C)*d*(n+2)*\tan[e + f*x] - (a*C*d*(n+2) - b*(c*C - B*d*(n+2))]*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 7337 vs. $2(315) = 630$.

Time = 0.32 (sec) , antiderivative size = 7338, normalized size of antiderivative = 20.79

method	result	size
parts	Expression too large to display	7338
derivativeldivides	Expression too large to display	7402
default	Expression too large to display	7402

input `int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48734 vs. $2(308) = 616$.

Time = 36.79 (sec) , antiderivative size = 48734, normalized size of antiderivative = 138.06

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `Too large to include`

Sympy [F]

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Integral((a + b*tan(e + f*x))*(c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)
```

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
Timed out
```

Giac [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%}{1, [0,29,11]}%%}+%%{12, [0,27,11]}%%}+%%{66, [0,25,11]}%%}+%%{2`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Hanged}$$

input `int((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `\text{Hanged}`

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) \\
& + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a^2 c^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^5 dx \right) b c d^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) a c d^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b^2 d^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) b c^2 d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a b d^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) a c^2 d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) b^2 c d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) b c^3 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a^2 d^2 \\
& + 4 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a b c d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a c^3 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) b^2 c^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a^2 c d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a b c^2
\end{aligned}$$

input

```
int((a+b*tan(f*x+e))*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```

int(sqrt(tan(e + f*x)*d + c),x)*a**2*c**2 + int(sqrt(tan(e + f*x)*d + c)*t
an(e + f*x)**5,x)*b*c*d**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,
x)*a*c*d**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b**2*d**2 +
2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*b*c**2*d + 2*int(sqrt(ta
n(e + f*x)*d + c)*tan(e + f*x)**3,x)*a*b*d**2 + 2*int(sqrt(tan(e + f*x)*d
+ c)*tan(e + f*x)**3,x)*a*c**2*d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e +
f*x)**3,x)*b**2*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b*c*
*3 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a**2*d**2 + 4*int(sqr
t(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*b*c*d + int(sqrt(tan(e + f*x)*d
+ c)*tan(e + f*x)**2,x)*a*c**3 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x
)**2,x)*b**2*c**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a**2*c*
d + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a*b*c**2

```

3.106 $\int (c+d \tan(e+fx))^{5/2} (A + B \tan(e + fx) + C \tan^2$

Optimal result	1200
Mathematica [A] (verified)	1201
Rubi [A] (warning: unable to verify)	1201
Maple [B] (verified)	1206
Fricas [B] (verification not implemented)	1207
Sympy [F]	1208
Maxima [F]	1208
Giac [F(-2)]	1208
Mupad [B] (verification not implemented)	1209
Reduce [F]	1210

Optimal result

Integrand size = 35, antiderivative size = 229

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx =$$

$$\frac{(iA + B - iC)(c - id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{f}$$

$$- \frac{(B - i(A - C))(c + id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{f}$$

$$+ \frac{2(2c(A - C)d + B(c^2 - d^2)) \sqrt{c + d \tan(e + fx)}}{f}$$

$$+ \frac{2(Bc + (A - C)d)(c + d \tan(e + fx))^{3/2}}{3f}$$

$$+ \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

output

```
-(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/f
-(B-I*(A-C))*(c+I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/f
+2*(2*c*(A-C)*d+B*(c^2-d^2))*(c+d*tan(f*x+e))^(1/2)/f+2/3*(B*c+(A-C)*d)*(c
+d*tan(f*x+e))^(3/2)/f+2/5*B*(c+d*tan(f*x+e))^(5/2)/f+2/7*C*(c+d*tan(f*x+e
))^(7/2)/d/f
```

Mathematica [A] (verified)

Time = 1.40 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.14

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{4C(c + d \tan(e + fx))^{7/2}}{d} + 7i(A - iB - C) \left(\frac{2}{5}(c + d \tan(e + fx))^{5/2} + \frac{2}{3}(c - id) \left(-3(c - id) \right) \right)$$

input

```
Integrate[(c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
((4*C*(c + d*Tan[e + f*x])^(7/2))/d + (7*I)*(A - I*B - C)*((2*(c + d*Tan[e + f*x])^(5/2))/5 + (2*(c - I*d)*(-3*(c - I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]] + Sqrt[c + d*Tan[e + f*x]]*(4*c - (3*I)*d + d*Tan[e + f*x])))/3) - (7*I)*(A + I*B - C)*((2*(c + d*Tan[e + f*x])^(5/2))/5 + (2*(c + I*d)*(-3*(c + I*d)^(3/2)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]] + Sqrt[c + d*Tan[e + f*x]]*(4*c + (3*I)*d + d*Tan[e + f*x])))/3)))/(14*f)
```

Rubi [A] (warning: unable to verify)

Time = 2.15 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.91, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4113, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow \text{4113}$$

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^{5/2} dx + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 3042

$$\int (A - C + B \tan(e + fx))(c + d \tan(e + fx))^{5/2} dx + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 4011

$$\int (c + d \tan(e + fx))^{3/2} (Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 3042

$$\int (c + d \tan(e + fx))^{3/2} (Ac - Cc - Bd + (Bc + (A - C)d) \tan(e + fx)) dx + \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 4011

$$\int \sqrt{c + d \tan(e + fx)} (-Cc^2 - 2Bdc + Cd^2 + A(c^2 - d^2) + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)) dx + \frac{2(d(A - C) + Bc)(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 3042

$$\int \sqrt{c + d \tan(e + fx)} (-Cc^2 - 2Bdc + Cd^2 + A(c^2 - d^2) + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)) dx + \frac{2(d(A - C) + Bc)(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 4011

$$\int \frac{-Cc^3 - 3Bdc^2 + 3Cd^2c + Bd^3 + A(c^3 - 3cd^2) + ((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx +$$

$$\frac{2(2cd(A - C) + B(c^2 - d^2)) \sqrt{c + d \tan(e + fx)}}{f} + \frac{2(d(A - C) + Bc)(c + d \tan(e + fx))^{3/2}}{3f} +$$

$$\frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 3042

$$\int \frac{-Cc^3 - 3Bdc^2 + 3Cd^2c + Bd^3 + A(c^3 - 3cd^2) + ((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx +$$

$$\frac{2(2cd(A - C) + B(c^2 - d^2)) \sqrt{c + d \tan(e + fx)}}{f} + \frac{2(d(A - C) + Bc)(c + d \tan(e + fx))^{3/2}}{3f} +$$

$$\frac{2B(c + d \tan(e + fx))^{5/2}}{5f} + \frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 4022

$$C) \int \frac{1}{2}(c + id)^3(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(c - id)^3(A - iB -$$

$$\frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2(2cd(A - C) + B(c^2 - d^2)) \sqrt{c + d \tan(e + fx)}}{f} +$$

$$\frac{2(d(A - C) + Bc)(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} +$$

$$\frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 3042

$$C) \int \frac{1}{2}(c + id)^3(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(c - id)^3(A - iB -$$

$$\frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2(2cd(A - C) + B(c^2 - d^2)) \sqrt{c + d \tan(e + fx)}}{f} +$$

$$\frac{2(d(A - C) + Bc)(c + d \tan(e + fx))^{3/2}}{3f} + \frac{2B(c + d \tan(e + fx))^{5/2}}{5f} +$$

$$\frac{2C(c + d \tan(e + fx))^{7/2}}{7df}$$

↓ 4020

$$\begin{aligned}
& \frac{i(c-id)^3(A-iB-C) \int -\frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} - \\
& \frac{i(c+id)^3(A+iB-C) \int -\frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} + \\
& \frac{2(2cd(A-C) + B(c^2 - d^2)) \sqrt{c+d \tan(e+fx)}}{f} + \frac{2(d(A-C) + Bc)(c+d \tan(e+fx))^{3/2}}{3f} + \\
& \frac{2B(c+d \tan(e+fx))^{5/2}}{5f} + \frac{2C(c+d \tan(e+fx))^{7/2}}{7df} \\
& \quad \downarrow 25 \\
& -\frac{i(c-id)^3(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} + \\
& \frac{i(c+id)^3(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} + \\
& \frac{2(2cd(A-C) + B(c^2 - d^2)) \sqrt{c+d \tan(e+fx)}}{f} + \frac{2(d(A-C) + Bc)(c+d \tan(e+fx))^{3/2}}{3f} + \\
& \frac{2B(c+d \tan(e+fx))^{5/2}}{5f} + \frac{2C(c+d \tan(e+fx))^{7/2}}{7df} \\
& \quad \downarrow 73 \\
& \frac{(c+id)^3(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} + \\
& \frac{(c-id)^3(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} + \\
& \frac{2(2cd(A-C) + B(c^2 - d^2)) \sqrt{c+d \tan(e+fx)}}{f} + \frac{2(d(A-C) + Bc)(c+d \tan(e+fx))^{3/2}}{3f} + \\
& \frac{2B(c+d \tan(e+fx))^{5/2}}{5f} + \frac{2C(c+d \tan(e+fx))^{7/2}}{7df} \\
& \quad \downarrow 221 \\
& \frac{(c-id)^{5/2}(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f} + \frac{(c+id)^{5/2}(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f} + \\
& \frac{2(2cd(A-C) + B(c^2 - d^2)) \sqrt{c+d \tan(e+fx)}}{f} + \frac{2(d(A-C) + Bc)(c+d \tan(e+fx))^{3/2}}{3f} + \\
& \frac{2B(c+d \tan(e+fx))^{5/2}}{5f} + \frac{2C(c+d \tan(e+fx))^{7/2}}{7df}
\end{aligned}$$

input `Int[(c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output
$$\begin{aligned} & ((A - I*B - C)*(c - I*d)^{(5/2)}*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + ((A \\ & + I*B - C)*(c + I*d)^{(5/2)}*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f + (2*(2* \\ & c*(A - C)*d + B*(c^2 - d^2))*Sqrt[c + d*Tan[e + f*x]])/f + (2*(B*c + (A - \\ & C)*d)*(c + d*Tan[e + f*x])^{(3/2)})/(3*f) + (2*B*(c + d*Tan[e + f*x])^{(5/2)}) \\ & / (5*f) + (2*C*(c + d*Tan[e + f*x])^{(7/2)})/(7*d*f) \end{aligned}$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(F x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F x, x], x]$$

rule 73
$$\text{Int}[(a + b x)^m (c + d x)^n, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{p(m+1)-1} (c - a(d/b) + d(x^p/b))^n, x], x, (a + b x)^{1/p}], x]] \text{ /; FreeQ}\{a, b, c, d, x\} \&\& \text{Lt} Q[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 221
$$\text{Int}[(a + b x^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ /; FreeQ}\{a, b, x\} \&\& \text{NegQ}[a/b]$$

rule 3042
$$\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinear} Q[u, x]$$

rule 4011
$$\text{Int}[(a + b \tan[e + f x])^m (c + d \tan[e + f x] + (f x)), x_Symbol] \rightarrow \text{Simp}[d*((a + b \tan[e + f x])^m / (f m)), x] + \text{Int}[(a + b \tan[e + f x])^{m-1} * \text{Simp}[a c - b d + (b c + a d) \tan[e + f x], x], x] \text{ /; FreeQ}\{a, b, c, d, e, f, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 0]$$

rule 4020
$$\text{Int}[(a + b \tan[e + f x])^m (c + d \tan[e + f x] + (f x)), x_Symbol] \rightarrow \text{Simp}[c(d/f) \quad \text{Subst}[\text{Int}[(a + (b/d)x)^m / (d^2 + c x), x], x, d \tan[e + f x]], x] \text{ /; FreeQ}\{a, b, c, d, e, f, m, x\} \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$$

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3586 vs. $2(196) = 392$.

Time = 0.19 (sec) , antiderivative size = 3587, normalized size of antiderivative = 15.66

method	result	size
parts	Expression too large to display	3587
derivativedivides	Expression too large to display	3614
default	Expression too large to display	3614

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x,method=_RETUR
NVERBOSE)
```

output

```
A*(2/3/f*d*(c+d*tan(f*x+e))^(3/2)+4/f*d*(c+d*tan(f*x+e))^(1/2)*c-1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*c^2+1/4/f*d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)+1/4/f/d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^3-3/4/f*d*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c+3/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^2-1/f*d^3/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))-2/f*d/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*(c^2+d^2)^(1/2)*c+1/4/f/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)*c^2-1/4/f*d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*(c^2+d^2)^(1/2)-1/4/f/d*ln(-d*tan(f*x+e)-c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^3+3/4/f*d*ln(-d*tan(f*x+e)-c+(...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10840 vs. $2(189) = 378$.

Time = 1.80 (sec) , antiderivative size = 10840, normalized size of antiderivative = 47.34

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

Too large to include

Sympy [F]

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(d \tan(fx + e) + c)^{5/2} dx$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(5/2), x)`

Giac [F(-2)]

Exception generated.

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Exception raised: TypeError}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{%%[1,[0,24,9]%%]}+%%{10,[0,22,9]%%]}+%%{45,[0,20,9]%%}
+%%{120,
```

Mupad [B] (verification not implemented)

Time = 109.01 (sec) , antiderivative size = 5863, normalized size of antiderivative = 25.60

$$\int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```
int((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)
```

output

```
((2*C*c^2)/(3*d*f) - (2*C*(d^3*f + c^2*d*f))/(3*d^2*f^2))*(c + d*tan(e + f
*x))^(3/2) - log(((((-B^4*d^2*f^4*(5*c^4 + d^4 - 10*c^2*d^2)^2)^(1/2) + B^
2*c^5*f^2 - 10*B^2*c^3*d^2*f^2 + 5*B^2*c*d^4*f^2)/f^4)^(1/2)*(((((-B^4*d^2
*f^4*(5*c^4 + d^4 - 10*c^2*d^2)^2)^(1/2) + B^2*c^5*f^2 - 10*B^2*c^3*d^2*f^
2 + 5*B^2*c*d^4*f^2)/f^4)^(1/2)*(32*B*c^4*d^2 - 32*B*d^6 + 32*c*d^2*f*((-
B^4*d^2*f^4*(5*c^4 + d^4 - 10*c^2*d^2)^2)^(1/2) + B^2*c^5*f^2 - 10*B^2*c^3
*d^2*f^2 + 5*B^2*c*d^4*f^2)/f^4)^(1/2)*(c + d*tan(e + f*x))^(1/2)))/(2*f)
- (16*B^2*d^2*(c + d*tan(e + f*x))^(1/2)*(c^6 - d^6 + 15*c^2*d^4 - 15*c^4*
d^2))/f^2))/2 - (8*B^3*c*d^2*(c^2 - 3*d^2)*(c^2 + d^2)^3)/f^3)*((((20*B^4*c
^2*d^8*f^4 - B^4*d^10*f^4 - 110*B^4*c^4*d^6*f^4 + 100*B^4*c^6*d^4*f^4 - 25
*B^4*c^8*d^2*f^4)^(1/2) + B^2*c^5*f^2 - 10*B^2*c^3*d^2*f^2 + 5*B^2*c*d^4*f
^2)/(4*f^4))^(1/2) + log(- ((((-B^4*d^2*f^4*(5*c^4 + d^4 - 10*c^2*d^2)^2)^(
1/2) + B^2*c^5*f^2 - 10*B^2*c^3*d^2*f^2 + 5*B^2*c*d^4*f^2)/f^4)^(1/2)*(((
((-B^4*d^2*f^4*(5*c^4 + d^4 - 10*c^2*d^2)^2)^(1/2) + B^2*c^5*f^2 - 10*B^2*
c^3*d^2*f^2 + 5*B^2*c*d^4*f^2)/f^4)^(1/2)*(32*B*d^6 - 32*B*c^4*d^2 + 32*c*
d^2*f*(((-B^4*d^2*f^4*(5*c^4 + d^4 - 10*c^2*d^2)^2)^(1/2) + B^2*c^5*f^2 -
10*B^2*c^3*d^2*f^2 + 5*B^2*c*d^4*f^2)/f^4)^(1/2)*(c + d*tan(e + f*x))^(1/2
)))/(2*f) - (16*B^2*d^2*(c + d*tan(e + f*x))^(1/2)*(c^6 - d^6 + 15*c^2*d^4
- 15*c^4*d^2))/f^2))/2 - (8*B^3*c*d^2*(c^2 - 3*d^2)*(c^2 + d^2)^3)/f^3)*((
20*B^4*c^2*d^8*f^4 - B^4*d^10*f^4 - 110*B^4*c^4*d^6*f^4 + 100*B^4*c^6*...
```

Reduce [F]

$$\begin{aligned}
& \int (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) \\
& + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} dx \right) a c^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^4 dx \right) c d^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) b d^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^3 dx \right) c^2 d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) a d^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) b c d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e)^2 dx \right) c^3 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) a c d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \tan(fx + e) dx \right) b c^2
\end{aligned}$$

input `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int(sqrt(tan(e + f*x)*d + c),x)*a*c**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4,x)*c*d**2 + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*b*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3,x)*c**2*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*a*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*b*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2,x)*c**3 + 2*int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*a*c*d + int(sqrt(tan(e + f*x)*d + c)*tan(e + f*x),x)*b*c**2`

3.107 $\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx$

Optimal result	1211
Mathematica [A] (verified)	1212
Rubi [A] (warning: unable to verify)	1212
Maple [B] (verified)	1219
Fricas [F(-1)]	1220
Sympy [F(-1)]	1220
Maxima [F(-2)]	1220
Giac [F(-2)]	1221
Mupad [F(-1)]	1221
Reduce [F]	1222

Optimal result

Integrand size = 47, antiderivative size = 336

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{a+b \tan(e+fx)} dx =$$

$$\frac{(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a-ib)f}$$

$$+ \frac{(iA-B-iC)(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a+ib)f}$$

$$- \frac{2(Ab^2-a(bB-aC))(bc-ad)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{b^{7/2}(a^2+b^2)f}$$

$$+ \frac{2(b^2d(Bc+(A-C)d)+(bc-ad)(bcC+bBd-aCd))\sqrt{c+d \tan(e+fx)}}{b^3f}$$

$$+ \frac{2(bcC+bBd-aCd)(c+d \tan(e+fx))^{3/2}}{3b^2f} + \frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

output

$$-(I*A+B-I*C)*(c-I*d)^{(5/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c-I*d)^{(1/2)})/(a-I*b)/f+(I*A-B-I*C)*(c+I*d)^{(5/2)}*\operatorname{arctanh}((c+d*\tan(f*x+e))^{(1/2)}/(c+I*d)^{(1/2)})/(a+I*b)/f-2*(A*b^2-a*(B*b-C*a))*(-a*d+b*c)^{(5/2)}*\operatorname{arctanh}(b^{(1/2)}*(c+d*\tan(f*x+e))^{(1/2)}/(-a*d+b*c)^{(1/2)})/b^{(7/2)}/(a^2+b^2)/f+2*(b^2*d*(B*c+(A-C)*d)+(-a*d+b*c)*(B*b*d-C*a*d+C*b*c))*(c+d*\tan(f*x+e))^{(1/2)}/b^3/f+2/3*(B*b*d-C*a*d+C*b*c)*(c+d*\tan(f*x+e))^{(3/2)}/b^2/f+2/5*C*(c+d*\tan(f*x+e))^{(5/2)}/b/f$$
Mathematica [A] (verified)

Time = 3.63 (sec) , antiderivative size = 322, normalized size of antiderivative = 0.96

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \frac{15 \left(b^{7/2} (-ia+b)(A-iB-C)(c-id)^{5/2} \operatorname{arctanh} \right)}{\dots}$$

input

```
Integrate[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]
```

output

$$\left((15*b^{(7/2)}*((-I)*a + b)*(A - I*B - C)*(c - I*d)^{(5/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[c + d*\tan[e + f*x]]/\operatorname{Sqrt}[c - I*d]] + b^{(7/2)}*(I*a + b)*(A + I*B - C)*(c + I*d)^{(5/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[c + d*\tan[e + f*x]]/\operatorname{Sqrt}[c + I*d]] - 2*(A*b^2 + a*(-b*B) + a*C))*(b*c - a*d)^{(5/2)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[c + d*\tan[e + f*x]])/\operatorname{Sqrt}[b*c - a*d]] \right) / (b^{(5/2)}*(a^2 + b^2)) + (30*(b^2*d*(B*c + (A - C)*d) + (b*c - a*d)*(b*c*C + b*B*d - a*C*d))*\operatorname{Sqrt}[c + d*\tan[e + f*x]]/b^2 + (10*(b*c*C + b*B*d - a*C*d)*(c + d*\tan[e + f*x])^{(3/2)})/b + 6*C*(c + d*\tan[e + f*x])^{(5/2)})/(15*b*f)$$
Rubi [A] (warning: unable to verify)

Time = 6.43 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.01, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.489$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4136, 25, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{a + b \tan(e + fx)} dx \\
 & \quad \downarrow \text{4130} \\
 & \frac{2 \int \frac{5(c + d \tan(e + fx))^{3/2} ((bcC - adC + bBd) \tan^2(e + fx) + b(Bc + (A - C)d) \tan(e + fx) + Abc - aCd)}{2(a + b \tan(e + fx))} dx}{\frac{5b}{2C(c + d \tan(e + fx))^{5/2}} + \frac{5bf}{5bf}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(c + d \tan(e + fx))^{3/2} ((bcC - adC + bBd) \tan^2(e + fx) + b(Bc + (A - C)d) \tan(e + fx) + Abc - aCd)}{a + b \tan(e + fx)} dx}{\frac{b}{2C(c + d \tan(e + fx))^{5/2}} + \frac{5bf}{5bf}} + \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(c + d \tan(e + fx))^{3/2} ((bcC - adC + bBd) \tan(e + fx)^2 + b(Bc + (A - C)d) \tan(e + fx) + Abc - aCd)}{a + b \tan(e + fx)} dx}{\frac{b}{2C(c + d \tan(e + fx))^{5/2}} + \frac{5bf}{5bf}} + \\
 & \quad \downarrow \text{4130} \\
 & \frac{2 \int \frac{3\sqrt{c + d \tan(e + fx)} (Ac^2b^2 + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)b^2 + (d(Bc + (A - C)d)b^2 + (bc - ad)(bcC - adC + bBd)) \tan^2(e + fx) + ad(aCd - b(2cC + Bd)))}{2(a + b \tan(e + fx))} dx}{\frac{3b}{b} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5bf}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{c + d \tan(e + fx)} (Ac^2b^2 + (2c(A - C)d + B(c^2 - d^2)) \tan(e + fx)b^2 + (d(Bc + (A - C)d)b^2 + (bc - ad)(bcC - adC + bBd)) \tan^2(e + fx) + ad(aCd - b(2cC + Bd)))}{a + b \tan(e + fx)} dx}{\frac{b}{b} + \frac{2C(c + d \tan(e + fx))^{5/2}}{5bf}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(Ac^2b^2 + (2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2 + (d(Bc+(A-C)d)b^2 + (bc-ad)(bcC-adC+bBd)) \tan(e+fx)^2 + ad(aCd-b(2cC+Bd)) \right)}{a+b \tan(e+fx)} dx$$

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

b

↓ 4130

$$2 \int \frac{\left((A-C)d(3c^2-d^2) + B(c^3-3cd^2) \right) \tan(e+fx)b^3 + A(bc^3-ad^3)b^2 + (d(2c(A-C)d+B(c^2-d^2))b^3 + (bc-ad)(d(Bc+(A-C)d)b^2 + (bc-ad)(bcC-adC+bBd))}{2(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

b

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 27

$$\int \frac{\left((A-C)d(3c^2-d^2) + B(c^3-3cd^2) \right) \tan(e+fx)b^3 + A(bc^3-ad^3)b^2 + (d(2c(A-C)d+B(c^2-d^2))b^3 + (bc-ad)(d(Bc+(A-C)d)b^2 + (bc-ad)(bcC-adC+bBd))}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

b

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 3042

$$\int \frac{\left((A-C)d(3c^2-d^2) + B(c^3-3cd^2) \right) \tan(e+fx)b^3 + A(bc^3-ad^3)b^2 + (d(2c(A-C)d+B(c^2-d^2))b^3 + (bc-ad)(d(Bc+(A-C)d)b^2 + (bc-ad)(bcC-adC+bBd))}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

b

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 4136

$$(bc-ad)^3(Ab^2-a(bB-aC)) \int \frac{\tan^2(e+fx)+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx + \int -\frac{b^3(a(Cc^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))-b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))}{a^2+b^2} dx$$

b

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 25

$$\frac{(bc-ad)^3 (Ab^2 - a(bB - aC)) \int \frac{\tan^2(e+fx)+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx - \int \frac{(b(A-C)d(3c^2-d^2)+bB(c^3-3cd^2)+a(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3))b^3}{a^2+b^2}}{b}$$

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 25

$$\frac{(bc-ad)^3 (Ab^2 - a(bB - aC)) \int \frac{\tan^2(e+fx)+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx + \int \frac{(b(A-C)d(3c^2-d^2)+bB(c^3-3cd^2)-a(Cc^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2)))b^3}{a^2+b^2}}{b}$$

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 3042

$$\frac{(bc-ad)^3 (Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx + \int \frac{(b(A-C)d(3c^2-d^2)+bB(c^3-3cd^2)-a(Cc^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2)))b^3}{a^2+b^2}}{b}$$

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf}$$

↓ 4022

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf} +$$

$$\frac{2(-aCd+bBd+bcC)(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{2\sqrt{c+d \tan(e+fx)}((bc-ad)(-aCd+bBd+bcC)+b^2d(d(A-C)+Bc))}{bf} + \frac{(bc-ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a+b \tan(e+fx))}{a^2+b^2}}{b}$$

↓ 3042

$$\frac{2C(c+d \tan(e+fx))^{5/2}}{5bf} +$$

$$\frac{2(-aCd+bBd+bcC)(c+d \tan(e+fx))^{3/2}}{3bf} + \frac{2\sqrt{c+d \tan(e+fx)}((bc-ad)(-aCd+bBd+bcC)+b^2d(d(A-C)+Bc))}{bf} + \frac{(bc-ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a+b \tan(e+fx))}{a^2+b^2}}{b}$$

↓ 4020

$$\frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c + d \tan(e + fx)}((bc - ad)(-aCd + bBd + bcC) + b^2 d(d(A - C) + Bc))}{bf} + \frac{(bc - ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a + b \tan(e + fx))}{a^2 + b^2} dx}{a^2 + b^2}$$

↓ 25

$$\frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c + d \tan(e + fx)}((bc - ad)(-aCd + bBd + bcC) + b^2 d(d(A - C) + Bc))}{bf} + \frac{(bc - ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a + b \tan(e + fx))}{a^2 + b^2} dx}{a^2 + b^2}$$

↓ 73

$$\frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c + d \tan(e + fx)}((bc - ad)(-aCd + bBd + bcC) + b^2 d(d(A - C) + Bc))}{bf} + \frac{(bc - ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a + b \tan(e + fx))}{a^2 + b^2} dx}{a^2 + b^2}$$

↓ 221

$$\frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c + d \tan(e + fx)}((bc - ad)(-aCd + bBd + bcC) + b^2 d(d(A - C) + Bc))}{bf} + \frac{(bc - ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a + b \tan(e + fx))}{a^2 + b^2} dx}{a^2 + b^2}$$

b

↓ 4117

$$\frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c + d \tan(e + fx)}((bc - ad)(-aCd + bBd + bcC) + b^2 d(d(A - C) + Bc))}{bf} + \frac{(bc - ad)^3 (Ab^2 - a(bB - aC)) \int \frac{(a + b \tan(e + fx))}{a^2 + b^2} dx}{a^2 + b^2}$$

b

↓ 73

$$\begin{aligned}
 & \frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(bc-ad)^3 (Ab^2 - a(bB - aC)) f \frac{b(c}{a+}}{df(a^2} \\
 & \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c+d \tan(e+fx)}((bc-ad)(-aCd + bBd + bcC) + b^2 d(d(A-C) + Bc))}{bf} + \frac{2(bc-ad)^{5/2} (Ab^2 - a(bB - aC)) \arctan\left(\frac{b(c}{a+}}{df(a^2} \right)}{\sqrt{bf}(a^2 + b^2)} \\
 & \quad \downarrow \text{221} \\
 & \frac{2C(c + d \tan(e + fx))^{5/2}}{5bf} + \frac{2(-aCd + bBd + bcC)(c + d \tan(e + fx))^{3/2}}{3bf} + \frac{2\sqrt{c+d \tan(e+fx)}((bc-ad)(-aCd + bBd + bcC) + b^2 d(d(A-C) + Bc))}{bf} + \frac{2(bc-ad)^{5/2} (Ab^2 - a(bB - aC)) \arctan\left(\frac{b(c}{a+}}{df(a^2} \right)}{\sqrt{bf}(a^2 + b^2)}
 \end{aligned}$$

input

```
Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x]),x]
```

output

```
(2*C*(c + d*Tan[e + f*x])^(5/2))/(5*b*f) + ((2*(b*c*C + b*B*d - a*C*d)*(c + d*Tan[e + f*x])^(3/2))/(3*b*f) + (((((a + I*b)*b^3*(A - I*B - C)*(c - I*d)^(5/2)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/f + ((a - I*b)*b^3*(A + I*B - C)*(c + I*d)^(5/2)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/f)/(a^2 + b^2) - (2*(A*b^2 - a*(b*B - a*C))*(b*c - a*d)^(5/2)*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/(Sqrt[b]*(a^2 + b^2)*f))/b + (2*(b^2*d*(B*c + (A - C)*d) + (b*c - a*d)*(b*c*C + b*B*d - a*C*d))*Sqrt[c + d*Tan[e + f*x]])/(b*f))/b
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

- rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p(m+1)-1)}(c - a(d/b) + d(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}, x]] \text{ /}; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 221 $\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) * \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ /}; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /}; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4020 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \text{ Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\text{Tan}[e + f*x]], x] \text{ /}; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$
- rule 4022 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 - I*\text{Tan}[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 + I*\text{Tan}[e + f*x]), x], x] \text{ /}; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!IntegerQ}[m]$
- rule 4117 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)(x_)]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_)]^{(n_.)}((A_) + (C_.)\tan[(e_.) + (f_.)(x_)]^2), x_Symbol] \rightarrow \text{Simp}[A/f \text{ Subst}[\text{Int}[(a + b*x)^m*(c + d*x)^n, x], x, \text{Tan}[e + f*x]], x] \text{ /}; \text{FreeQ}[\{a, b, c, d, e, f, A, C, m, n\}, x] \&\& \text{EqQ}[A, C]$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2], x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && ( !IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8697 vs. $2(294) = 588$.

Time = 0.23 (sec) , antiderivative size = 8698, normalized size of antiderivative = 25.89

method	result	size
derivativedivides	Expression too large to display	8698
default	Expression too large to display	8698

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```


Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Exception raised: ValueError}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for m
ore detail
```

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Exception raised: TypeError}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e)),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\begin{aligned}
& \int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{a + b \tan(e + fx)} dx = \left(\int \frac{\sqrt{d \tan(fx + e) + c}}{\tan(fx + e) b + a} dx \right) a c^2 \\
& + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^4}{\tan(fx + e) b + a} dx \right) c d^2 \\
& + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^3}{\tan(fx + e) b + a} dx \right) b d^2 \\
& + 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^3}{\tan(fx + e) b + a} dx \right) c^2 d \\
& + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) a d^2 \\
& + 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) b c d \\
& + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) c^3 \\
& + 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) a c d \\
& + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) b c^2
\end{aligned}$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)*b + a),x)*a*c**2 + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)*b + a),x)*c*d**2 + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*b + a),x)*b*d**2 + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*b + a),x)*c**2*d + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*a*d**2 + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*b*c*d + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*c**3 + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*a*c*d + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*b*c**2
```

3.108
$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx$$

Optimal result	1224
Mathematica [B] (verified)	1225
Rubi [F]	1225
Maple [B] (verified)	1233
Fricas [F(-1)]	1234
Sympy [F(-1)]	1234
Maxima [F(-2)]	1234
Giac [F(-2)]	1235
Mupad [F(-1)]	1235
Reduce [F]	1236

Optimal result

Integrand size = 47, antiderivative size = 473

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^2} dx =$$

$$\frac{(iA+B- iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a-ib)^2 f}$$

$$- \frac{(B-i(A-C))(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a+ib)^2 f}$$

$$+ \frac{(bc-ad)^{3/2} (3a^3 b B d - 5a^4 C d - b^4 (2Bc + 5Ad) - ab^3 (4Ac - 4cC - 7Bd) + a^2 b^2 (2Bc - (A + 9C)d)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{b^{7/2} (a^2 + b^2)^2 f}$$

$$- \frac{d(5a^3 C d - Ab^2 (bc - ad) - 2b^3 (2cC + Bd) - a^2 b (5cC + 3Bd) + ab^2 (Bc + 4Cd)) \sqrt{c+d \tan(e+fx)}}{b^3 (a^2 + b^2) f}$$

$$+ \frac{(3Ab^2 - 3abB + 5a^2 C + 2b^2 C) d (c+d \tan(e+fx))^{3/2}}{3b^2 (a^2 + b^2) f}$$

$$- \frac{(Ab^2 - a(bB - aC)) (c+d \tan(e+fx))^{5/2}}{b (a^2 + b^2) f (a+b \tan(e+fx))}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(
a-I*b)^2/f-(B-I*(A-C))*(c+I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)
)^(1/2))/(a+I*b)^2/f+(-a*d+b*c)^(3/2)*(3*a^3*b*B*d-5*a^4*C*d-b^4*(5*A*d+2*
B*c)-a*b^3*(4*A*c-7*B*d-4*C*c)+a^2*b^2*(2*B*c-(A+9*C)*d))*arctanh(b^(1/2)*
(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/b^(7/2)/(a^2+b^2)^2/f-d*(5*a^3*C*
d-A*b^2*(-a*d+b*c)-2*b^3*(B*d+2*C*c)-a^2*b*(3*B*d+5*C*c)+a*b^2*(B*c+4*C*d)
)*(c+d*tan(f*x+e))^(1/2)/b^3/(a^2+b^2)/f+1/3*(3*A*b^2-3*B*a*b+5*C*a^2+2*C*
b^2)*d*(c+d*tan(f*x+e))^(3/2)/b^2/(a^2+b^2)/f-(A*b^2-a*(B*b-C*a))*(c+d*tan
(f*x+e))^(5/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 6112 vs. $2(473) = 946$.

Time = 6.63 (sec) , antiderivative size = 6112, normalized size of antiderivative = 12.92

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Result too large to show}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^2,x]

```

output

Result too large to show

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^2} dx$$

↓ 4128

$$\int \frac{(c+d \tan(e+fx))^{3/2} \left((5Ca^2-3bBa+3Ab^2+2b^2C)d \tan^2(e+fx)-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+2(bB-aC) \left(bc-\frac{5ad}{2} \right) +2Ab(ac+bd) \right)}{2(a+b \tan(e+fx))} \\ \frac{b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}} \\ \frac{bf(a^2+b^2)(a+b \tan(e+fx))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 27

$$\int \frac{(c+d \tan(e+fx))^{3/2} \left((5Ca^2-3bBa+3Ab^2+2b^2C)d \tan^2(e+fx)-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(2bc-5ad)+Ab(2ac+bd) \right)}{a+b \tan(e+fx)} \\ \frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}} \\ \frac{bf(a^2+b^2)(a+b \tan(e+fx))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^{3/2} \left((5Ca^2-3bBa+3Ab^2+2b^2C)d \tan^2(e+fx)^2-2b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(2bc-5ad)+Ab(2ac+bd) \right)}{a+b \tan(e+fx)} \\ \frac{2b(a^2+b^2)}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}} \\ \frac{bf(a^2+b^2)(a+b \tan(e+fx))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 4130

$$2 \int \frac{3\sqrt{c+d \tan(e+fx)} \left(-2(2aAc d-2acC d-Ab(c^2-d^2))+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2) \right) \tan(e+fx)b^2-c((bB-aC)(2bc-5ad)+Ab(2ac+5bd))b+a(5Ca^2-3bBa+3Ab^2+2b^2C)}{2(a+b \tan(e+fx))} \\ \frac{3b}{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}} \\ \frac{bf(a^2+b^2)(a+b \tan(e+fx))}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 27

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \int \frac{\sqrt{c+d \tan(e+fx)} \left(-2(2aAc d-2acC d-Ab(c^2-d^2))+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2) \right) \tan(e+fx)}{2(a+b \tan(e+fx))} \\ \frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{bf(a^2+b^2)(a+b \tan(e+fx))}$$

↓ 3042

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \int \frac{\sqrt{c+d \tan(e+fx)}(-2(2aAc d-2acC d-Ab(c^2-d^2))+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)}{bf} dx$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 4130

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d \int \frac{5Cd^3a^4 - bd^2(10cC+3Bd)a^3 + b^2d(5Cc^2+4Bdc+(A+4C)d^2)a^2 + b^3(2Ac^3-2Cc^3-5Bdc^2-4Ccd^2)}{bf} dx}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 27

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d \int \frac{\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf} dx}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 3042

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d \int \frac{\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf} dx}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 4136

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d \int \frac{\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf} dx}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 27

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C-3abB+3Ab^2+2b^2C)(c+d \tan(e+fx))^{3/2}}{3bf} - \frac{2d\sqrt{c+d \tan(e+fx)}(5a^3Cd-a^2b(3Bd+5cC)-Ab^2(bc-ad)+ab^2(Bc+4Cd)-2b^3(Bd+2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) (a + b \tan(e + fx))}$$

↓ 25

$$\frac{2d(5a^2C - 3abB + 3Ab^2 + 2b^2C)(c + d \tan(e + fx))^{3/2}}{3bf} - \frac{2d\sqrt{c + d \tan(e + fx)}(5a^3Cd - a^2b(3Bd + 5cC) - Ab^2(bc - ad) + ab^2(Bc + 4Cd) - 2b^3(Bd + 2cC))}{bf}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{bf(a^2 + b^2)(a + b \tan(e + fx))}$$

input

```
Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^2,x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 14118 vs. $2(434) = 868$.

Time = 0.27 (sec) , antiderivative size = 14119, normalized size of antiderivative = 29.85

method	result	size
derivativedivides	Expression too large to display	14119
default	Expression too large to display	14119

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^2,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")
```

output

Timed out

Sympy [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**2,x)
```

output

Timed out

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Exception raised: ValueError}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for m
ore detail
```

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Exception raised: TypeError}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^2,x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{%%}{1, [0,22,10]%%}+%%{10, [0,20,10]%%}+%%{45, [0,18,10]%%
%%}+%%{1
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x))^2,x)
```

output

```
\text{Hanged}
```


Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^2} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^2,x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2*c**2*d + 4*sqrt(tan(e + f*x)*d + c)*a*
b*c**3 + 4*sqrt(tan(e + f*x)*d + c)*a*b*c*d**2 + 2*sqrt(tan(e + f*x)*d + c
)*b**2*c**2*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*t
an(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*ta
n(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*a**3*b*c**2*d**2*f - 3*int(sqrt
(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + t
an(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*
c),x)*tan(e + f*x)*a**2*b**2*c**3*d*f + 2*int(sqrt(tan(e + f*x)*d + c)/(ta
n(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c +
tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*a*b**
3*c**4*f + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e
+ f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e +
f*x)*a*b*c + a**2*c),x)*a**4*c**2*d**2*f - 3*int(sqrt(tan(e + f*x)*d + c)
/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*
c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a**3*b*c**3*d*
f + 2*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)
)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)
*a*b*c + a**2*c),x)*a**2*b**2*c**4*f - 2*int(sqrt(tan(e + f*x)*d + c)/(tan
(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*
a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**...
```

3.109
$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx$$

Optimal result	1237
Mathematica [C] (warning: unable to verify)	1238
Rubi [F]	1238
Maple [B] (verified)	1246
Fricas [F(-1)]	1247
Sympy [F(-1)]	1247
Maxima [F(-2)]	1247
Giac [F(-2)]	1248
Mupad [F(-1)]	1248
Reduce [F]	1249

Optimal result

Integrand size = 47, antiderivative size = 643

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^3} dx =$$

$$-\frac{(A-iB-C)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(ia+b)^3 f}$$

$$+\frac{(A+iB-C)(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(ia-b)^3 f}$$

$$+\frac{\sqrt{bc-ad}(3a^5bBd^2-15a^6Cd^2+a^4b^2d(4Bc+(A-46C)d)-3a^2b^4(8Ac^2-8c^2C-16Bcd-6Ad^2+21C^2d))}{4b^3(a^2+b^2)^2 f}$$

$$-\frac{d(3a^3bBd-15a^4Cd-ab^3(8Ac-8cC-11Bd)+a^2b^2(4Bc+(A-31C)d)-b^4(4Bc+7Ad+8Cd))}{4b^3(a^2+b^2)^2 f}$$

$$+\frac{(a^3bBd-5a^4Cd-b^4(4Bc+5Ad)-ab^3(8Ac-8cC-9Bd)+a^2b^2(4Bc+3Ad-13Cd))(c+d \tan(e+fx))}{4b^2(a^2+b^2)^2 f(a+b \tan(e+fx))}$$

$$-\frac{(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{2b(a^2+b^2) f(a+b \tan(e+fx))^2}$$

output

```

-(A-I*B-C)*(c-I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(I*
a+b)^3/f+(A+I*B-C)*(c+I*d)^(5/2)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1
/2))/(I*a-b)^3/f+1/4*(-a*d+b*c)^(1/2)*(3*a^5*b*B*d^2-15*a^6*C*d^2+a^4*b^2*
d*(4*B*c+(A-46*C)*d)-3*a^2*b^4*(8*A*c^2-6*A*d^2-16*B*c*d-8*C*c^2+21*C*d^2)
-a*b^5*(56*c*(A-C)*d+B*(24*c^2-35*d^2))-b^6*(4*c*(5*B*d+2*C*c)-A*(8*c^2-15
*d^2))+2*a^3*b^3*(4*c*(A-C)*d+B*(4*c^2+3*d^2))*arctanh(b^(1/2)*(c+d*tan(f
*x+e))^(1/2)/(-a*d+b*c)^(1/2))/b^(7/2)/(a^2+b^2)^3/f-1/4*d*(3*a^3*b*B*d-15
*a^4*C*d-a*b^3*(8*A*c-11*B*d-8*C*c)+a^2*b^2*(4*B*c+(A-31*C)*d)-b^4*(7*A*d+
4*B*c+8*C*d))*(c+d*tan(f*x+e))^(1/2)/b^3/(a^2+b^2)^2/f+1/4*(a^3*b*B*d-5*a^
4*C*d-b^4*(5*A*d+4*B*c)-a*b^3*(8*A*c-9*B*d-8*C*c)+a^2*b^2*(3*A*d+4*B*c-13*
C*d))*(c+d*tan(f*x+e))^(3/2)/b^2/(a^2+b^2)^2/f/(a+b*tan(f*x+e))-1/2*(A*b^2
-a*(B*b-C*a))*(c+d*tan(f*x+e))^(5/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^2

```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 123.16 (sec) , antiderivative size = 2429150, normalized size of antiderivative = 3777.84

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Result too large to show}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^3,x]

```

output

Result too large to show

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^3} dx$$

↓ 4128

$$\int \frac{(c+d \tan(e+fx))^{3/2} \left((5Ca^2 - bBa + Ab^2 + 4b^2C) d \tan^2(e+fx) - 4b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + 2(bB - aC) \left(2bc - \frac{5ad}{2} \right) + 2Ab \left(2ac + \frac{5ad}{2} \right) \right)}{2(a+b \tan(e+fx))^2}$$

$$\frac{2b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}} \frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 27

$$\int \frac{(c+d \tan(e+fx))^{3/2} \left((5Ca^2 - bBa + Ab^2 + 4b^2C) d \tan^2(e+fx) - 4b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB - aC)(4bc - 5ad) + Ab(4ac + 5bd) \right)}{(a+b \tan(e+fx))^2}$$

$$\frac{4b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}} \frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^{3/2} \left((5Ca^2 - bBa + Ab^2 + 4b^2C) d \tan^2(e+fx) - 4b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB - aC)(4bc - 5ad) + Ab(4ac + 5bd) \right)}{(a+b \tan(e+fx))^2}$$

$$\frac{4b(a^2 + b^2)}{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}} \frac{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 4128

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-8((ac+bd)((A-C)(bc-ad) - B(ac+bd)) + (bc-ad)(bBc + b(A-C)d + a(Ac - Cc - Bd))) \tan(e+fx)b^2 + 2 \left(ac + \frac{3bd}{2} \right) ((bB - aC)(4bc - 5ad) + Ab(4ac + 5bd)) \right)}{2(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-8((ac+bd)((A-C)(bc-ad) - B(ac+bd)) + (bc-ad)(bBc + b(A-C)d + a(Ac - Cc - Bd))) \tan(e+fx)b^2 + 2 \left(ac + \frac{3bd}{2} \right) ((bB - aC)(4bc - 5ad) + Ab(4ac + 5bd)) \right)}{2(a+b \tan(e+fx))^2}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2bf(a^2 + b^2)(a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-8((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+2\left(ac+\frac{3bd}{2}\right)((bB-aC)(4bc-5ad)+Ab(4a$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 4130

$$2 \int -\frac{15Cd^3a^5-3bd^2(5cC+Bd)a^4-b^2d^2(Bc+(A-31C)d)a^3-b^3(8Ac^3-8Cc^3-20Bdc^2-17Ad^2c+47Cd^2c+11Bd^3)a^2-b^4(16Bc^3+40Adc^2-40Cdc^2-31Bd^2c-7$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 27

$$\int \frac{15Cd^3a^5-3bd^2(5cC+Bd)a^4-b^2d^2(Bc+(A-31C)d)a^3-b^3(8Ac^3-8Cc^3-20Bdc^2-17Ad^2c+47Cd^2c+11Bd^3)a^2-b^4(16Bc^3+40Adc^2-40Cdc^2-31Bd^2c-7$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{15Cd^3a^5-3bd^2(5cC+Bd)a^4-b^2d^2(Bc+(A-31C)d)a^3-b^3(8Ac^3-8Cc^3-20Bdc^2-17Ad^2c+47Cd^2c+11Bd^3)a^2-b^4(16Bc^3+40Adc^2-40Cdc^2-31Bd^2c-7$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2bf (a^2 + b^2) (a + b \tan(e + fx))^2}$$

↓ 4136

$$\frac{(-5Cda^4+bBda^3+b^2(4Bc+3Ad-13Cd)a^2-b^3(8Ac-8Cc-9Bd)a-b^4(4Bc+5Ad))(c+d \tan(e+fx))^{3/2}}{b(a^2+b^2)f(a+b \tan(e+fx))} + \frac{2d\sqrt{c+d \tan(e+fx)}(-15Cda^4+3bBda^3+b^2(4Bc+3Ad-13Cd)a^2-b^3(8Ac-8Cc-9Bd)a-b^4(4Bc+5Ad))}{b(a^2+b^2)f(a+b \tan(e+fx))}$$

$$\frac{(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{2b (a^2 + b^2) f(a + b \tan(e + fx))^2}$$

↓ 27

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

↓ 25

$$\frac{(-5Cda^4 + bBda^3 + b^2(4Bc + 3Ad - 13Cd)a^2 - b^3(8Ac - 8Cc - 9Bd)a - b^4(4Bc + 5Ad))(c + d \tan(e + fx))^{3/2}}{b(a^2 + b^2)f(a + b \tan(e + fx))} + \frac{-2d\sqrt{c + d \tan(e + fx)}(-15Cda^4 + 3bBda^3)}{b(a^2 + b^2)f(a + b \tan(e + fx))}$$

$$\frac{(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{2b(a^2 + b^2)f(a + b \tan(e + fx))^2}$$

input `Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^3,x]`

output `$Aborted`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 20662 vs. 2(599) = 1198.

Time = 0.37 (sec) , antiderivative size = 20663, normalized size of antiderivative = 32.14

method	result	size
derivativedivides	Expression too large to display	20663
default	Expression too large to display	20663

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^3,x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**3,x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Exception raised: ValueError}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for m
ore detail
```

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Exception raised: TypeError}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^3,x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{%%{-1, [0,21,6]%%}+%%{-10, [0,19,6]%%}+%%{-45, [0,17,6]%%
%%}+%%{-
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x))^3,x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^3} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^3,x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2*c**2*d + 8*sqrt(tan(e + f*x)*d + c)*a*
b*c**3 + 12*sqrt(tan(e + f*x)*d + c)*a*b*c*d**2 + 6*sqrt(tan(e + f*x)*d +
c)*b**2*c**2*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*
tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2
*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a
**2*b*c + a**3*c),x)*tan(e + f*x)**2*a**3*b**2*c**2*d**2*f - 5*int(sqrt(ta
n(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + t
an(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b
**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e +
f*x)**2*a**2*b**3*c**3*d*f + 4*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)
**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e
+ f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3
*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a*b**4*c**4*f + 2*int(
sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**
2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)
**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*
tan(e + f*x)*a**4*b*c**2*d**2*f - 10*int(sqrt(tan(e + f*x)*d + c)/(tan(e +
f*x)**4*b**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*
tan(e + f*x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*
d + 3*tan(e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)*a**3*b**2*c**3*d...
```

3.110 $\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$

Optimal result	1250
Mathematica [A] (verified)	1251
Rubi [A] (warning: unable to verify)	1252
Maple [B] (verified)	1257
Fricas [B] (verification not implemented)	1258
Sympy [F]	1258
Maxima [F(-1)]	1259
Giac [F(-2)]	1259
Mupad [B] (verification not implemented)	1260
Reduce [F]	1261

Optimal result

Integrand size = 47, antiderivative size = 407

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{(ia + b)^3 (A - iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}f}$$

$$- \frac{(ia - b)^3 (A + iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}f}$$

$$+ \frac{2(72a^3Cd^3 - 6a^2bd^2(32cC - 49Bd) + 21ab^2d(8c^2C - 10Bcd + 15(A - C)d^2) - b^3(48c^3C - 56Bc^2d - 105d^4f)}{105d^4f}$$

$$+ \frac{2b(35b(Ab + aB - bC)d^2 + 4(bc - ad)(6bcC - 7bBd - 6aCd)) \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{105d^3f}$$

$$- \frac{2(6bcC - 7bBd - 6aCd)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{35d^2f}$$

$$+ \frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df}$$

output

$$\begin{aligned} & (I*a+b)^3*(A-I*B-C)*\operatorname{arctanh}((c+d*\tan(f*x+e))^{1/2}/(c-I*d)^{1/2})/(c-I*d)^{1/2}/f - (I*a-b)^3*(A+I*B-C)*\operatorname{arctanh}((c+d*\tan(f*x+e))^{1/2}/(c+I*d)^{1/2})/ \\ & (c+I*d)^{1/2}/f + 2/105*(72*a^3*C*d^3 - 6*a^2*b*d^2*(-49*B*d + 32*C*c) + 21*a*b^2*d \\ & * (8*c^2*C - 10*B*c*d + 15*(A-C)*d^2) - b^3*(48*c^3*C - 56*B*c^2*d + 70*c*(A-C)*d^2 + \\ & 105*B*d^3)) * (c+d*\tan(f*x+e))^{1/2} / d^4 / f + 2/105*b*(35*b*(A*b+B*a-C*b)*d^2 + 4 \\ & * (-a*d+b*c)*(-7*B*b*d - 6*C*a*d + 6*C*b*c)) * \tan(f*x+e) * (c+d*\tan(f*x+e))^{1/2} / \\ & d^3 / f - 2/35*(-7*B*b*d - 6*C*a*d + 6*C*b*c) * (a+b*\tan(f*x+e))^2 * (c+d*\tan(f*x+e))^{1/2} / \\ & d^2 / f + 2/7*C*(a+b*\tan(f*x+e))^3 * (c+d*\tan(f*x+e))^{1/2} / d / f \end{aligned}$$

Mathematica [A] (verified)

Time = 6.03 (sec) , antiderivative size = 392, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\ & = \frac{105(a-ib)^3(iA+B-iC)\operatorname{darctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} + \frac{105i(a+ib)^3(A+iB-C)\operatorname{darctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} - \frac{2(-72a^3Cd^3+6a^2b} \end{aligned}$$

input

```
Integrate[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/Sqrt[c + d*Tan[e + f*x]],x]
```

output

```
((-105*(a - I*b)^3*(I*A + B - I*C)*d*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt
[c - I*d]])/Sqrt[c - I*d] + ((105*I)*(a + I*b)^3*(A + I*B - C)*d*ArcTanh[S
qrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d] - (2*(-72*a^3*C*d^3
+ 6*a^2*b*d^2*(32*c*C - 49*B*d) - 21*a*b^2*d*(8*c^2*C - 10*B*c*d + 15*(A -
C)*d^2) + b^3*(48*c^3*C - 56*B*c^2*d + 70*c*(A - C)*d^2 + 105*B*d^3))*Sqr
t[c + d*Tan[e + f*x]]/d^3 + (2*b*(35*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a
*d)*(6*b*c*C - 7*b*B*d - 6*a*C*d))*Tan[e + f*x]*Sqrt[c + d*Tan[e + f*x]]/
d^2 + (6*(-6*b*c*C + 7*b*B*d + 6*a*C*d)*(a + b*Tan[e + f*x])^2*Sqrt[c + d*
Tan[e + f*x]])/d + 30*C*(a + b*Tan[e + f*x])^3*Sqrt[c + d*Tan[e + f*x]]/(
105*d*f)
```


Rubi [A] (warning: unable to verify)

Time = 4.74 (sec) , antiderivative size = 418, normalized size of antiderivative = 1.03, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.383$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4130} \\
 & \frac{2 \int -\frac{(a+b \tan(e+fx))^2 ((6bcC-6adC-7bBd) \tan^2(e+fx) - 7(Ab-Cb+aB)d \tan(e+fx) + 6bcC - a(7A-C)d)}{2\sqrt{c+d \tan(e+fx)}} dx}{\frac{7d}{7df} \frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df}} + \\
 & \quad \downarrow \text{27} \\
 & \frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \\
 & \frac{\int \frac{(a+b \tan(e+fx))^2 ((6bcC-6adC-7bBd) \tan^2(e+fx) - 7(Ab-Cb+aB)d \tan(e+fx) + 6bcC - a(7A-C)d)}{\sqrt{c+d \tan(e+fx)}} dx}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \\
 & \frac{\int \frac{(a+b \tan(e+fx))^2 ((6bcC-6adC-7bBd) \tan^2(e+fx) - 7(Ab-Cb+aB)d \tan(e+fx) + 6bcC - a(7A-C)d)}{\sqrt{c+d \tan(e+fx)}} dx}{7d} \\
 & \quad \downarrow \text{4130}
 \end{aligned}$$

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2 \int - \frac{(a+b \tan(e+fx))(4c(6cC-7Bd)b^2 - ad(48cC+7Bd)b+a^2(35A-11C)d^2 + (35b(Ab-Cb+aB)d^2 + 4(bc-ad)(6bcC-6adC-7bBd)) \tan^2(e+fx) + 35(Ba^2+2b(Ab-Cb+aB)d^2 + 4(bc-ad)(6bcC-6adC-7bBd)))}{2\sqrt{c+d \tan(e+fx)}}}{5d} dx$$

7d

↓ 27

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd-7bBd+6bcC)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \int \frac{(a+b \tan(e+fx))(4c(6cC-7Bd)b^2 - ad(48cC+7Bd)b+a^2(35A-11C)d^2 + (35b(Ab-Cb+aB)d^2 + 4(bc-ad)(6bcC-6adC-7bBd)))}{\sqrt{c+d \tan(e+fx)}} dx$$

7d

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd-7bBd+6bcC)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \int \frac{(a+b \tan(e+fx))(4c(6cC-7Bd)b^2 - ad(48cC+7Bd)b+a^2(35A-11C)d^2 + (35b(Ab-Cb+aB)d^2 + 4(bc-ad)(6bcC-6adC-7bBd)))}{\sqrt{c+d \tan(e+fx)}} dx$$

7d

↓ 4120

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd-7bBd+6bcC)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b \tan(e+fx) \sqrt{c+d \tan(e+fx)} (35bd^2(aB+Ab-bC)+4(bc-ad)(-6aCd-7bBd+6bcC))}{3df}$$

↓ 27

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd-7bBd+6bcC)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \int \frac{-2c(24C^2-28Bdc+35(A-C)d^2)b^3+42acd(4cC-5Bd)b^2-3a^2d^2(64cC+7Bd)b+3a^3d^2}{\sqrt{c+d \tan(e+fx)}} dx$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd-7bBd+6bcC)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \int \frac{-2c(24C^2-28Bdc+35(A-C)d^2)b^3+42acd(4cC-5Bd)b^2-3a^2d^2(64cC+7Bd)b+3a^3d^2}{\sqrt{c+d \tan(e+fx)}} dx$$

↓ 4113

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{\int \frac{105(Ba^3 + 3b(A-C)a^2 - 3b^2Ba - b^3(A-C))d^3 \tan(e + fx) - 105(-(A-C)a^3) + 3bBa^2}{\sqrt{c + d \tan(e + fx)}}}{\sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{\int \frac{105(Ba^3 + 3b(A-C)a^2 - 3b^2Ba - b^3(A-C))d^3 \tan(e + fx) - 105(-(A-C)a^3) + 3bBa^2}{\sqrt{c + d \tan(e + fx)}}}{\sqrt{c + d \tan(e + fx)}}$$

↓ 4022

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx) \sqrt{c + d \tan(e + fx)} (35bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 7bBd + 6bcC))}{3df}$$

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx) \sqrt{c + d \tan(e + fx)} (35bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 7bBd + 6bcC))}{3df}$$

↓ 4020

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx) \sqrt{c + d \tan(e + fx)} (35bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 7bBd + 6bcC))}{3df}$$

↓ 25

$$\frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx) \sqrt{c + d \tan(e + fx)} (35bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 7bBd + 6bcC))}{3df}$$

$$\begin{aligned}
 & \downarrow 73 \\
 & \frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \\
 & \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx) \sqrt{c + d \tan(e + fx)} (35bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 7bBd + 6bcC))}{3df} + \\
 & \downarrow 221 \\
 & \frac{2C(a + b \tan(e + fx))^3 \sqrt{c + d \tan(e + fx)}}{7df} - \\
 & \frac{2(-6aCd - 7bBd + 6bcC)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx) \sqrt{c + d \tan(e + fx)} (35bd^2(aB + Ab - bC) + 4(bc - ad)(-6aCd - 7bBd + 6bcC))}{3df} +
 \end{aligned}$$

input

```
Int[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[
c + d*Tan[e + f*x]],x]
```

output

```
(2*C*(a + b*Tan[e + f*x])^3*Sqrt[c + d*Tan[e + f*x]]/(7*d*f) - ((2*(6*b*c
*C - 7*b*B*d - 6*a*C*d)*(a + b*Tan[e + f*x])^2*Sqrt[c + d*Tan[e + f*x]]/(
5*d*f) - ((2*b*(35*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 7*b*
B*d - 6*a*C*d))*Tan[e + f*x]*Sqrt[c + d*Tan[e + f*x]])/(3*d*f) + ((105*(a
- I*b)^3*(A - I*B - C)*d^3*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]]/(Sqrt[c - I
*d]*f) + (105*(a + I*b)^3*(A + I*B - C)*d^3*ArcTan[Tan[e + f*x]/Sqrt[c + I
*d]]/(Sqrt[c + I*d]*f) + (2*(72*a^3*C*d^3 - 6*a^2*b*d^2*(32*c*C - 49*B*d)
+ 21*a*b^2*d*(8*c^2*C - 10*B*c*d + 15*(A - C)*d^2) - b^3*(48*c^3*C - 56*B
*c^2*d + 70*c*(A - C)*d^2 + 105*B*d^3))*Sqrt[c + d*Tan[e + f*x]]/(d*f))/(
3*d))/(5*d))/(7*d)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
 c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
 b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
 1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
 *(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
 - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`
- rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
 + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
 b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
 mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
 NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)]^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] :> Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5977 vs. $2(371) = 742$.

Time = 0.42 (sec) , antiderivative size = 5978, normalized size of antiderivative = 14.69

method	result	size
parts	Expression too large to display	5978
derivativedivides	Expression too large to display	25426
default	Expression too large to display	25426

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1
/2),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37247 vs. $2(363) = 726$.

Time = 11.45 (sec) , antiderivative size = 37247, normalized size of antiderivative = 91.52

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\ &= \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((a + b*tan(e + f*x))**3*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

output

Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

= Exception raised: TypeError

input

```
integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{%%}{1, [0,24,9]%%}+%%{10, [0,22,9]%%}+%%{45, [0,20,9]%%}+%%{120,
```


Mupad [B] (verification not implemented)

Time = 104.88 (sec) , antiderivative size = 28858, normalized size of antiderivative = 70.90

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input

```
int(((a + b*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c +
d*tan(e + f*x))^(1/2),x)
```

output

```
atan((((8*(4*A*a^3*d^3*f^2 - 12*A*a*b^2*d^3*f^2 + 4*A*b^3*c*d^2*f^2 - 12*
A*a^2*b*c*d^2*f^2))/f^3 - 64*c*d^2*(c + d*tan(e + f*x))^(1/2)*(((8*A^2*a^
6*c*f^2 - 8*A^2*b^6*c*f^2 + 48*A^2*a*b^5*d*f^2 + 48*A^2*a^5*b*d*f^2 + 120*
A^2*a^2*b^4*c*f^2 - 120*A^2*a^4*b^2*c*f^2 - 160*A^2*a^3*b^3*d*f^2)^2/4 - (
16*c^2*f^4 + 16*d^2*f^4)*(A^4*a^12 + A^4*b^12 + 6*A^4*a^2*b^10 + 15*A^4*a^
4*b^8 + 20*A^4*a^6*b^6 + 15*A^4*a^8*b^4 + 6*A^4*a^10*b^2)))^(1/2) - 4*A^2*a
^6*c*f^2 + 4*A^2*b^6*c*f^2 - 24*A^2*a*b^5*d*f^2 - 24*A^2*a^5*b*d*f^2 - 60*
A^2*a^2*b^4*c*f^2 + 60*A^2*a^4*b^2*c*f^2 + 80*A^2*a^3*b^3*d*f^2)/(16*(c^2*
f^4 + d^2*f^4)))^(1/2))*(((8*A^2*a^6*c*f^2 - 8*A^2*b^6*c*f^2 + 48*A^2*a*b
^5*d*f^2 + 48*A^2*a^5*b*d*f^2 + 120*A^2*a^2*b^4*c*f^2 - 120*A^2*a^4*b^2*c*
f^2 - 160*A^2*a^3*b^3*d*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)*(A^4*a^12 + A
^4*b^12 + 6*A^4*a^2*b^10 + 15*A^4*a^4*b^8 + 20*A^4*a^6*b^6 + 15*A^4*a^8*b^
4 + 6*A^4*a^10*b^2)))^(1/2) - 4*A^2*a^6*c*f^2 + 4*A^2*b^6*c*f^2 - 24*A^2*a*
b^5*d*f^2 - 24*A^2*a^5*b*d*f^2 - 60*A^2*a^2*b^4*c*f^2 + 60*A^2*a^4*b^2*c*f
^2 + 80*A^2*a^3*b^3*d*f^2)/(16*(c^2*f^4 + d^2*f^4)))^(1/2) - (16*(c + d*ta
n(e + f*x))^(1/2)*(A^2*a^6*d^2 - A^2*b^6*d^2 + 15*A^2*a^2*b^4*d^2 - 15*A^2
*a^4*b^2*d^2))/f^2)*(((8*A^2*a^6*c*f^2 - 8*A^2*b^6*c*f^2 + 48*A^2*a*b^5*d
*f^2 + 48*A^2*a^5*b*d*f^2 + 120*A^2*a^2*b^4*c*f^2 - 120*A^2*a^4*b^2*c*f^2
- 160*A^2*a^3*b^3*d*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)*(A^4*a^12 + A^4*b
^12 + 6*A^4*a^2*b^10 + 15*A^4*a^4*b^8 + 20*A^4*a^6*b^6 + 15*A^4*a^8*b^4...
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{2\sqrt{d \tan(fx + e) + c} a^4 + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^5}{d \tan(fx + e) + c} dx \right) b^3 c d f + 3 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^4}{d \tan(fx + e) + c} dx \right) a b^2 c d f}{1}$$

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```
(2*sqrt(tan(e + f*x)*d + c)*a**4 + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5)/(tan(e + f*x)*d + c),x)*b**3*c*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)*d + c),x)*a*b**2*c*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)*d + c),x)*b**4*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*d + c),x)*a**2*b*c*d*f + 4*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*d + c),x)*a*b**3*d*f - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**4*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**3*c*d*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**2*b**2*d*f + 4*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*d + c),x)*a**3*b*d*f)/(d*f)
```

3.111
$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1262
Mathematica [A] (verified)	1263
Rubi [A] (warning: unable to verify)	1263
Maple [B] (verified)	1268
Fricas [B] (verification not implemented)	1269
Sympy [F]	1269
Maxima [F(-1)]	1270
Giac [F(-2)]	1270
Mupad [B] (verification not implemented)	1271
Reduce [F]	1272

Optimal result

Integrand size = 47, antiderivative size = 287

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= -\frac{(a - ib)^2 (B + i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c - id} f}$$

$$+ \frac{(a + ib)^2 (iA - B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c + id} f}$$

$$+ \frac{2(12a^2Cd^2 - 10abd(2cC - 3Bd) + b^2(8c^2C - 10Bcd + 15(A - C)d^2)) \sqrt{c + d \tan(e + fx)}}{15d^3 f}$$

$$- \frac{2b(4bcC - 5bBd - 4aCd) \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{15d^2 f}$$

$$+ \frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df}$$

output

$$-(a-I*b)^2*(B+I*(A-C))*\operatorname{arctanh}((c+d*\tan(f*x+e))^{1/2}/(c-I*d)^{1/2})/(c-I*d)^{1/2}/f+(a+I*b)^2*(I*A-B-I*C)*\operatorname{arctanh}((c+d*\tan(f*x+e))^{1/2}/(c+I*d)^{1/2})/(c+I*d)^{1/2}/f+2/15*(12*a^2*C*d^2-10*a*b*d*(-3*B*d+2*C*c)+b^2*(8*c^2*C-10*B*c*d+15*(A-C)*d^2))*(c+d*\tan(f*x+e))^{1/2}/d^3/f-2/15*b*(-5*B*b*d-4*C*a*d+4*C*b*c)*\tan(f*x+e)*(c+d*\tan(f*x+e))^{1/2}/d^2/f+2/5*C*(a+b*\tan(f*x+e))^2*(c+d*\tan(f*x+e))^{1/2}/d/f$$
Mathematica [A] (verified)

Time = 3.96 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.96

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{15(a-ib)^2(iA+B-iC)d \operatorname{darctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} + \frac{15i(a+ib)^2(A+iB-C)d \operatorname{darctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} + \frac{2(12a^2Cd^2+10abd(-$$

input

```
Integrate[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]
```

output

$$((-15*(a - I*b)^2*(I*A + B - I*C)*d*\operatorname{ArcTanh}[\operatorname{Sqrt}[c + d*\operatorname{Tan}[e + f*x]]/\operatorname{Sqrt}[c - I*d]])/\operatorname{Sqrt}[c - I*d] + ((15*I)*(a + I*b)^2*(A + I*B - C)*d*\operatorname{ArcTanh}[\operatorname{Sqrt}[c + d*\operatorname{Tan}[e + f*x]]/\operatorname{Sqrt}[c + I*d]])/\operatorname{Sqrt}[c + I*d] + (2*(12*a^2*C*d^2 + 10*a*b*d*(-2*c*C + 3*B*d) + b^2*(8*c^2*C - 10*B*c*d + 15*(A - C)*d^2))*\operatorname{Sqrt}[c + d*\operatorname{Tan}[e + f*x]]/d^2 + (2*b*(-4*b*c*C + 5*b*B*d + 4*a*C*d)*\operatorname{Tan}[e + f*x]*\operatorname{Sqrt}[c + d*\operatorname{Tan}[e + f*x]])/d + 6*C*(a + b*\operatorname{Tan}[e + f*x])^2*\operatorname{Sqrt}[c + d*\operatorname{Tan}[e + f*x]]/(15*d*f)$$
Rubi [A] (warning: unable to verify)Time = 2.93 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.319$, Rules used = {3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4130} \\
 & 2 \int - \frac{(a + b \tan(e + fx))((4bcC - 4adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + 4bcC - a(5A - C)d)}{2\sqrt{c + d \tan(e + fx)}} dx + \\
 & \quad \frac{5d}{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \\
 & \int \frac{(a + b \tan(e + fx))((4bcC - 4adC - 5bBd) \tan^2(e + fx) - 5(Ab - Cb + aB)d \tan(e + fx) + 4bcC - a(5A - C)d)}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \frac{5d}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \\
 & \int \frac{(a + b \tan(e + fx))((4bcC - 4adC - 5bBd) \tan(e + fx)^2 - 5(Ab - Cb + aB)d \tan(e + fx) + 4bcC - a(5A - C)d)}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \frac{5d}{5d} \\
 & \quad \downarrow \text{4120} \\
 & \frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \\
 & \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} - \frac{2 \int - \frac{-2c(4cC - 5Bd)b^2 + 20acCdb - 3a^2(5A - C)d^2 - ((8Cc^2 - 10Bdc + 15(A - C)d^2)b^2 - 10ad(2c - 3Bd)b + 12a^2Cd^2) \tan^2(e + fx) - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan^2(e + fx)}{2\sqrt{c + d \tan(e + fx)}} dx}{3d} \\
 & \quad \frac{5d}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \\
 & \int \frac{-2c(4cC - 5Bd)b^2 + 20acCdb - 3a^2(5A - C)d^2 - ((8Cc^2 - 10Bdc + 15(A - C)d^2)b^2 - 10ad(2c - 3Bd)b + 12a^2Cd^2) \tan^2(e + fx) - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \frac{5d}{3d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{\int \frac{-2c(4cC - 5Bd)b^2 + 20acCdb - 3a^2(5A - C)d^2 - ((8Cc^2 - 10Bdc + 15(A - C)d^2)b^2 - 10ad(2cC - 3Bd)b + 12a^2Cd^2) \tan(e + fx)^2 - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx}{3d}$$

5d

↓ 4113

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{\int \frac{15(-((A - C)a^2) + 2bBa + b^2(A - C))d^2 - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{2\sqrt{c + d \tan(e + fx)}(12a^2Cd^2 - 10abd(2cC - 3Bd) + b^2(15d^2(A - C) - 10Bd))}{df}}{3d}$$

5d

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{\int \frac{15(-((A - C)a^2) + 2bBa + b^2(A - C))d^2 - 15(Ba^2 + 2b(A - C)a - b^2B)d^2 \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{2\sqrt{c + d \tan(e + fx)}(12a^2Cd^2 - 10abd(2cC - 3Bd) + b^2(15d^2(A - C) - 10Bd))}{df}}{3d}$$

5d

↓ 4022

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} + \frac{-\frac{15}{2}d^2(a + ib)^2(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{15}{2}d^2(a - ib)^2(A - iB - C) \int \frac{i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx}{3}$$

5d

↓ 3042

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} + \frac{-\frac{15}{2}d^2(a + ib)^2(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx - \frac{15}{2}d^2(a - ib)^2(A - iB - C) \int \frac{i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx}{3}$$

5d

↓ 4020

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} + \frac{15id^2(a - ib)^2(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} + \frac{15id^2(a + ib)^2(A + iB - C) \int \frac{1}{(1 + i \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f}$$

5d

↓ 25

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} + \frac{15id^2(a - ib)^2(A - iB - C) \int \frac{1}{(1 - i \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \frac{15id^2(a + ib)^2(A + iB - C) \int \frac{1}{(1 + i \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f}$$

5d

↓ 73

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} + \frac{15d(a - ib)^2(A - iB - C) \int \frac{1}{i \tan^2 \frac{(e + fx)}{d} + \frac{ic}{d} + 1} d \sqrt{c + d \tan(e + fx)}}{f} - \frac{15d(a + ib)^2(A + iB - C) \int \frac{1}{i \tan^2 \frac{(e + fx)}{d} + \frac{ic}{d} + 1} d \sqrt{c + d \tan(e + fx)}}{f}$$

5d

↓ 221

$$\frac{2C(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b \tan(e + fx)(-4aCd - 5bBd + 4bcC) \sqrt{c + d \tan(e + fx)}}{3df} + \frac{2\sqrt{c + d \tan(e + fx)}(12a^2Cd^2 - 10abd(2cC - 3Bd) + b^2(15d^2(A - C) - 10Bcd + 8c^2C))}{df} - \frac{15d^2(a - ib)^2(A - iB - C) \int \frac{1}{i \tan^2 \frac{(e + fx)}{d} + \frac{ic}{d} + 1} d \sqrt{c + d \tan(e + fx)}}{f} - \frac{15d^2(a + ib)^2(A + iB - C) \int \frac{1}{i \tan^2 \frac{(e + fx)}{d} + \frac{ic}{d} + 1} d \sqrt{c + d \tan(e + fx)}}{f}$$

5d

input

```
Int[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[
c + d*Tan[e + f*x]],x]
```

output

```
(2*C*(a + b*Tan[e + f*x])^2*Sqrt[c + d*Tan[e + f*x]]/(5*d*f) - ((2*b*(4*b
*c*C - 5*b*B*d - 4*a*C*d)*Tan[e + f*x]*Sqrt[c + d*Tan[e + f*x]]/(3*d*f) +
((-15*(a - I*b)^2*(A - I*B - C)*d^2*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(
Sqrt[c - I*d]*f) - (15*(a + I*b)^2*(A + I*B - C)*d^2*ArcTan[Tan[e + f*x]/S
qrt[c + I*d]]/(Sqrt[c + I*d]*f) - (2*(12*a^2*C*d^2 - 10*a*b*d*(2*c*C - 3*
B*d) + b^2*(8*c^2*C - 10*B*c*d + 15*(A - C)*d^2))*Sqrt[c + d*Tan[e + f*x]]
)/(d*f))/(3*d))/(5*d)
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{\text{n}_}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{1/\text{p}}, \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{LtQ}[-1, \text{m}, 0] \ \&\& \ \text{LeQ}[-1, \text{n}, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \ \&\& \ \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)]), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d})*\text{x})^{\text{m}}/(\text{d}^2 + \text{c}*\text{x}), \text{x}], \text{x}, \text{d}*\text{Tan}[\text{e} + \text{f}*\text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 - \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 + \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!IntegerQ}[\text{m}]$

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e
_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5512 vs. $2(254) = 508$.

Time = 0.19 (sec) , antiderivative size = 5513, normalized size of antiderivative = 19.21

method	result	size
parts	Expression too large to display	5513
derivativedivides	Expression too large to display	18289
default	Expression too large to display	18289

input `int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25627 vs. 2(244) = 488.

Time = 4.71 (sec) , antiderivative size = 25627, normalized size of antiderivative = 89.29

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Too large to include`

Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\ &= \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((a + b*tan(e + f*x))**2*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

output

Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

= Exception raised: TypeError

input

```
integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0,19,7]}%%+%%{8, [0,17,7]}%%+%%{28, [0,15,7]}%%+%%{56, [0
```

Mupad [B] (verification not implemented)

Time = 38.71 (sec) , antiderivative size = 21254, normalized size of antiderivative = 74.06

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input

```
int(((a + b*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2),x)
```

output

```
atan((((16*(2*C*b^2*d^3*f^2 - 2*C*a^2*d^3*f^2 + 4*C*a*b*c*d^2*f^2))/f^3 - 64*c*d^2*(c + d*tan(e + f*x))^(1/2)*(((8*C^2*a^4*c*f^2 + 8*C^2*b^4*c*f^2 - 32*C^2*a*b^3*d*f^2 + 32*C^2*a^3*b*d*f^2 - 48*C^2*a^2*b^2*c*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)*(C^4*a^8 + C^4*b^8 + 4*C^4*a^2*b^6 + 6*C^4*a^4*b^4 + 4*C^4*a^6*b^2))^(1/2) - 4*C^2*a^4*c*f^2 - 4*C^2*b^4*c*f^2 + 16*C^2*a*b^3*d*f^2 - 16*C^2*a^3*b*d*f^2 + 24*C^2*a^2*b^2*c*f^2)/(16*(c^2*f^4 + d^2*f^4))^(1/2))*(((8*C^2*a^4*c*f^2 + 8*C^2*b^4*c*f^2 - 32*C^2*a*b^3*d*f^2 + 32*C^2*a^3*b*d*f^2 - 48*C^2*a^2*b^2*c*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)*(C^4*a^8 + C^4*b^8 + 4*C^4*a^2*b^6 + 6*C^4*a^4*b^4 + 4*C^4*a^6*b^2))^(1/2) - 4*C^2*a^4*c*f^2 - 4*C^2*b^4*c*f^2 + 16*C^2*a*b^3*d*f^2 - 16*C^2*a^3*b*d*f^2 + 24*C^2*a^2*b^2*c*f^2)/(16*(c^2*f^4 + d^2*f^4))^(1/2) - (16*(c + d*tan(e + f*x))^(1/2)*(C^2*a^4*d^2 + C^2*b^4*d^2 - 6*C^2*a^2*b^2*d^2))/f^2)*(((8*C^2*a^4*c*f^2 + 8*C^2*b^4*c*f^2 - 32*C^2*a*b^3*d*f^2 + 32*C^2*a^3*b*d*f^2 - 48*C^2*a^2*b^2*c*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)*(C^4*a^8 + C^4*b^8 + 4*C^4*a^2*b^6 + 6*C^4*a^4*b^4 + 4*C^4*a^6*b^2))^(1/2) - 4*C^2*a^4*c*f^2 - 4*C^2*b^4*c*f^2 + 16*C^2*a*b^3*d*f^2 - 16*C^2*a^3*b*d*f^2 + 24*C^2*a^2*b^2*c*f^2)/(16*(c^2*f^4 + d^2*f^4))^(1/2)*1i - (((16*(2*C*b^2*d^3*f^2 - 2*C*a^2*d^3*f^2 + 4*C*a*b*c*d^2*f^2))/f^3 + 64*c*d^2*(c + d*tan(e + f*x))^(1/2)*(((8*C^2*a^4*c*f^2 + 8*C^2*b^4*c*f^2 - 32*C^2*a*b^3*d*f^2 + 32*C^2*a^3*b*d*f^2 - 48*C^2*a^2*b^2*c*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)...
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{2\sqrt{d \tan(fx + e) + c} a^3 + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^4}{d \tan(fx + e) + c} dx \right) b^2 c d f + 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^3}{d \tan(fx + e) + c} dx \right) a b c d f \cdot$$

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```
(2*sqrt(tan(e + f*x)*d + c)*a**3 + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)*d + c),x)*b**2*c*d*f + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*d + c),x)*a*b*c*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*d + c),x)*b**3*d*f - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**3*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**2*c*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a*b**2*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*d + c),x)*a**2*b*d*f)/(d*f)
```

3.112
$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1273
Mathematica [A] (verified)	1274
Rubi [A] (warning: unable to verify)	1274
Maple [B] (verified)	1278
Fricas [B] (verification not implemented)	1279
Sympy [F]	1280
Maxima [F]	1280
Giac [F(-2)]	1281
Mupad [B] (verification not implemented)	1281
Reduce [F]	1282

Optimal result

Integrand size = 45, antiderivative size = 194

$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

$$= -\frac{(ia+b)(A-iB-C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}f}$$

$$+ \frac{(ia-b)(A+iB-C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}f}$$

$$- \frac{2(2bcC-3bBd-3aCd)\sqrt{c+d \tan(e+fx)}}{3d^2f}$$

$$+ \frac{2bC \tan(e+fx)\sqrt{c+d \tan(e+fx)}}{3df}$$

output

```
-(I*a+b)*(A-I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(1/2)/f+(I*a-b)*(A+I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(1/2)/f-2/3*(-3*B*b*d-3*C*a*d+2*C*b*c)*(c+d*tan(f*x+e))^(1/2)/d^2/f+2/3*b*C*tan(f*x+e)*(c+d*tan(f*x+e))^(1/2)/d/f
```

Mathematica [A] (verified)

Time = 1.00 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.99

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{2 \left(-\frac{3i(a-ib)(A-iB-C)d \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{2\sqrt{c-id}} + \frac{3i(a+ib)(A+iB-C)d \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{2\sqrt{c+id}} + \frac{(-2bcC+3bBd+3aCd)}{d} \right)}{3df}$$

input

```
Integrate[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]
```

output

```
(2*((( (-3*I)/2)*(a - I*b)*(A - I*B - C)*d*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + (((3*I)/2)*(a + I*b)*(A + I*B - C)*d*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d] + ((-2*b*c*C + 3*b*B*d + 3*a*C*d)*Sqrt[c + d*Tan[e + f*x]])/d + b*C*Tan[e + f*x]*Sqrt[c + d*Tan[e + f*x]]))/(3*d*f)
```

Rubi [A] (warning: unable to verify)

Time = 1.64 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.95, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4120, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan(e + fx)^2)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$\downarrow 4120$$

$$\begin{aligned}
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{2 \int \frac{(2bcC-3adC-3bBd) \tan^2(e+fx) - 3(Ab-Cb+aB)d \tan(e+fx) + 2bcC-3aAd}{2\sqrt{c+d \tan(e+fx)}} dx}{3d} \\
 & \quad \downarrow 27 \\
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{\int \frac{(2bcC-3adC-3bBd) \tan^2(e+fx) - 3(Ab-Cb+aB)d \tan(e+fx) + 2bcC-3aAd}{\sqrt{c+d \tan(e+fx)}} dx}{3d} \\
 & \quad \downarrow 3042 \\
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{\int \frac{(2bcC-3adC-3bBd) \tan(e+fx)^2 - 3(Ab-Cb+aB)d \tan(e+fx) + 2bcC-3aAd}{\sqrt{c+d \tan(e+fx)}} dx}{3d} \\
 & \quad \downarrow 4113 \\
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{\int \frac{3(bB-a(A-C))d - 3(Ab-Cb+aB)d \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2(-3aCd-3bBd+2bcC)\sqrt{c+d \tan(e+fx)}}{df}}{3d} \\
 & \quad \downarrow 3042 \\
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{\int \frac{3(bB-a(A-C))d - 3(Ab-Cb+aB)d \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2(-3aCd-3bBd+2bcC)\sqrt{c+d \tan(e+fx)}}{df}}{3d} \\
 & \quad \downarrow 4022 \\
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{-\frac{3}{2}d(a+ib)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{3}{2}d(a-ib)(A-iB-C) \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2(-3aCd-3bBd+2bcC)\sqrt{c+d \tan(e+fx)}}{df}}{3d} \\
 & \quad \downarrow 3042 \\
 & \frac{2bC \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3df} - \\
 & \frac{-\frac{3}{2}d(a+ib)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{3}{2}d(a-ib)(A-iB-C) \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2(-3aCd-3bBd+2bcC)\sqrt{c+d \tan(e+fx)}}{df}}{3d}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 4020 \\ & \frac{2bC \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{3id(a-ib)(A-iB-C) \int -\frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} + \frac{3id(a+ib)(A+iB-C) \int -\frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} \\ & \hline & 3d \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{2bC \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{3id(a-ib)(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} - \frac{3id(a+ib)(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} \\ & \hline & 3d \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & \frac{2bC \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{3(a+ib)(A+iB-C) \int \frac{1}{-i \frac{\tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{f} - \frac{3(a-ib)(A-iB-C) \int \frac{1}{i \frac{\tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{f} + \frac{2(-3aCd - 3bBd + 2bcC)\sqrt{c+d \tan(e+fx)}}{3d} \\ & \hline & 3d \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & \frac{2bC \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{3d(a-ib)(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} - \frac{3d(a+ib)(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}} + \frac{2(-3aCd - 3bBd + 2bcC)\sqrt{c+d \tan(e+fx)}}{df} \\ & \hline & 3d \end{aligned}$$

input `Int[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]`

output `(2*b*C*Tan[e + f*x]*Sqrt[c + d*Tan[e + f*x]])/(3*d*f) - ((-3*(a - I*b)*(A - I*B - C)*d*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f) - (3*(a + I*b)*(A + I*B - C)*d*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f) + (2*(2*b*c*C - 3*b*B*d - 3*a*C*d)*Sqrt[c + d*Tan[e + f*x]]/(d*f))/(3*d)`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^{(\text{m}_)}*((\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{(\text{n}_)}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{(\text{p}*(\text{m} + 1) - 1)}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{(1/\text{p})}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{LtQ}[-1, \text{m}, 0] \ \&\& \ \text{LeQ}[-1, \text{n}, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \ \&\& \ \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^2)^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{(\text{m}_)}*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])], \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d})*\text{x})^{\text{m}}/(\text{d}^2 + \text{c}*\text{x}), \text{x}], \text{x}, \text{d}*\text{Tan}[\text{e} + \text{f}*\text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{(\text{m}_)}*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 - \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 + \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!IntegerQ}[\text{m}]$

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] :> Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3852 vs. $2(166) = 332$.

Time = 0.20 (sec) , antiderivative size = 3853, normalized size of antiderivative = 19.86

method	result	size
parts	Expression too large to display	3853
derivativeldivides	Expression too large to display	4138
default	Expression too large to display	4138

input

```
int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
A*a*(1/4/f/d/(c^2+d^2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*
(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^2+1/4/f*d/(c^2+d^2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+
(c^2+d^2)^(1/2))*
(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-1/4/f/d/(c^2+d^2)^(3/2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+
(c^2+d^2)^(1/2))*
(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^3-1/4/f*d/(c^2+d^2)^(3/2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+
(c^2+d^2)^(1/2))*
(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c-1/f/d/(c^2+d^2)^(1/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^2-1/f*d/(c^2+d^2)^(1/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))+1/f/d/(c^2+d^2)^(3/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^4+3/f*d/(c^2+d^2)^(3/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^2+2/f*d^3/(c^2+d^2)^(3/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))-1/4/f/d/(c^2+d^2)*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2)*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*
(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^2-1/4/f*d/(c^2+d^2)...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13473 vs. $2(159) = 318$.

Time = 1.59 (sec) , antiderivative size = 13473, normalized size of antiderivative = 69.45

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)^(1/2),x, algorithm="fricas")
```

output

Too large to include

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)}{\sqrt{d \tan(fx + e) + c}} dx$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)/sqrt(d*tan(f*x + e) + c), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

= Exception raised: TypeError

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [0,14,5]%%}+%%{6, [0,12,5]%%}+%%{15, [0,10,5]%%}+%%{20, [0`

Mupad [B] (verification not implemented)

Time = 18.06 (sec) , antiderivative size = 16400, normalized size of antiderivative = 84.54

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `int(((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2),x)`

output

```

((2*B*b*d - 6*C*b*c)/(d^2*f) + (4*C*b*c)/(d^2*f))*(c + d*tan(e + f*x))^(1/2)
- atan((((8*(4*C*a*d^3*f^2 - 4*A*a*d^3*f^2 + 4*B*a*c*d^2*f^2))/f^3 - 6
4*c*d^2*(c + d*tan(e + f*x))^(1/2)*(((8*A^2*a^2*c*f^2 - 8*B^2*a^2*c*f^2 +
8*C^2*a^2*c*f^2 + 16*A*B*a^2*d*f^2 - 16*A*C*a^2*c*f^2 - 16*B*C*a^2*d*f^2)
^2/4 - (16*c^2*f^4 + 16*d^2*f^4)*(A^4*a^4 + B^4*a^4 + C^4*a^4 - 4*A*C^3*a^
4 - 4*A^3*C*a^4 + 2*A^2*B^2*a^4 + 6*A^2*C^2*a^4 + 2*B^2*C^2*a^4 - 4*A*B^2*
C*a^4))^(1/2) - 4*A^2*a^2*c*f^2 + 4*B^2*a^2*c*f^2 - 4*C^2*a^2*c*f^2 - 8*A*
B*a^2*d*f^2 + 8*A*C*a^2*c*f^2 + 8*B*C*a^2*d*f^2)/(16*(c^2*f^4 + d^2*f^4)))
^(1/2))*(((8*A^2*a^2*c*f^2 - 8*B^2*a^2*c*f^2 + 8*C^2*a^2*c*f^2 + 16*A*B*a
^2*d*f^2 - 16*A*C*a^2*c*f^2 - 16*B*C*a^2*d*f^2)^2/4 - (16*c^2*f^4 + 16*d^2
*f^4)*(A^4*a^4 + B^4*a^4 + C^4*a^4 - 4*A*C^3*a^4 - 4*A^3*C*a^4 + 2*A^2*B^2
*a^4 + 6*A^2*C^2*a^4 + 2*B^2*C^2*a^4 - 4*A*B^2*C*a^4))^(1/2) - 4*A^2*a^2*c
*f^2 + 4*B^2*a^2*c*f^2 - 4*C^2*a^2*c*f^2 - 8*A*B*a^2*d*f^2 + 8*A*C*a^2*c*f
^2 + 8*B*C*a^2*d*f^2)/(16*(c^2*f^4 + d^2*f^4)))^(1/2) - (16*(c + d*tan(e +
f*x))^(1/2)*(A^2*a^2*d^2 - B^2*a^2*d^2 + C^2*a^2*d^2 - 2*A*C*a^2*d^2))/f^
2)*((((8*A^2*a^2*c*f^2 - 8*B^2*a^2*c*f^2 + 8*C^2*a^2*c*f^2 + 16*A*B*a^2*d*
f^2 - 16*A*C*a^2*c*f^2 - 16*B*C*a^2*d*f^2)^2/4 - (16*c^2*f^4 + 16*d^2*f^4)
*(A^4*a^4 + B^4*a^4 + C^4*a^4 - 4*A*C^3*a^4 - 4*A^3*C*a^4 + 2*A^2*B^2*a^4
+ 6*A^2*C^2*a^4 + 2*B^2*C^2*a^4 - 4*A*B^2*C*a^4))^(1/2) - 4*A^2*a^2*c*f^2
+ 4*B^2*a^2*c*f^2 - 4*C^2*a^2*c*f^2 - 8*A*B*a^2*d*f^2 + 8*A*C*a^2*c*f^2...

```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{2\sqrt{d \tan(fx + e) + c} a^2 + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^3}{d \tan(fx + e) + c} dx \right) bcd f - \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) a^2 d f + \dots$$

input

```

int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2)
),x)

```

output

```
(2*sqrt(tan(e + f*x)*d + c)*a**2 + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)*d + c),x)*b*c*d*f - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**2*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a*c*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*b**2*d*f + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)*d + c),x)*a*b*d*f)/(d*f)
```


3.113
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1284
Mathematica [A] (verified)	1285
Rubi [A] (warning: unable to verify)	1285
Maple [B] (verified)	1288
Fricas [B] (verification not implemented)	1289
Sympy [F]	1290
Maxima [F]	1290
Giac [F(-2)]	1291
Mupad [B] (verification not implemented)	1291
Reduce [F]	1292

Optimal result

Integrand size = 35, antiderivative size = 133

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx = -\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}f} - \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}f} + \frac{2C \sqrt{c + d \tan(e + fx)}}{df}$$

output

```
-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(1/2)/f
-(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(1/2)/f
+2*C*(c+d*tan(f*x+e))^(1/2)/d/f
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.97

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{-\frac{i(A-iB-C)\operatorname{arctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} + \frac{i(A+iB-C)\operatorname{arctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} + \frac{2C\sqrt{c+d\tan(e+fx)}}{d}}{f}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/Sqrt[c + d*Tan[e + f*x]], x]
```

output

```
(((-I)*(A - I*B - C)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + (I*(A + I*B - C)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d] + (2*C*Sqrt[c + d*Tan[e + f*x]]/d)/f
```

Rubi [A] (warning: unable to verify)

Time = 0.97 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.84, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$\downarrow \text{4113}$$

$$\int \frac{A - C + B \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2C\sqrt{c + d \tan(e + fx)}}{df}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{A - C + B \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{2C \sqrt{c + d \tan(e + fx)}}{df} \\
& \downarrow 4022 \\
& \frac{1}{2}(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \\
& \quad \frac{2C \sqrt{c + d \tan(e + fx)}}{df} \\
& \downarrow 3042 \\
& \frac{1}{2}(A + iB - C) \int \frac{1 - i \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx + \frac{1}{2}(A - iB - C) \int \frac{i \tan(e + fx) + 1}{\sqrt{c + d \tan(e + fx)}} dx + \\
& \quad \frac{2C \sqrt{c + d \tan(e + fx)}}{df} \\
& \downarrow 4020 \\
& \frac{i(A - iB - C) \int -\frac{1}{(1 - i \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \\
& \frac{i(A + iB - C) \int -\frac{1}{(i \tan(e + fx) + 1) \sqrt{c + d \tan(e + fx)}} d(-i \tan(e + fx))}{2f} + \frac{2C \sqrt{c + d \tan(e + fx)}}{df} \\
& \downarrow 25 \\
& -\frac{i(A - iB - C) \int \frac{1}{(1 - i \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} + \\
& \frac{i(A + iB - C) \int \frac{1}{(i \tan(e + fx) + 1) \sqrt{c + d \tan(e + fx)}} d(-i \tan(e + fx))}{2f} + \frac{2C \sqrt{c + d \tan(e + fx)}}{df} \\
& \downarrow 73 \\
& \frac{(A + iB - C) \int \frac{1}{-\frac{i \tan^2(e + fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{df} + \\
& \frac{(A - iB - C) \int \frac{1}{\frac{i \tan^2(e + fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{df} + \frac{2C \sqrt{c + d \tan(e + fx)}}{df} \\
& \downarrow 221 \\
& \frac{(A - iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f\sqrt{c - id}} + \frac{(A + iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c + id}}\right)}{f\sqrt{c + id}} + \frac{2C \sqrt{c + d \tan(e + fx)}}{df}
\end{aligned}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2)/\text{Sqrt}[c + d*\text{Tan}[e + f*x]],x]$

output $((A - I*B - C)*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]]/(\text{Sqrt}[c - I*d]*f) + ((A + I*B - C)*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c + I*d]]/(\text{Sqrt}[c + I*d]*f) + (2*C*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/(d*f)$

Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$

rule 73 $\text{Int}[(a + b*(x))^m*((c + d*(x))^n), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{p*(m+1)-1}*(c - a*(d/b) + d*(x^p/b))^n], x], x, (a + b*x)^{1/p}], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a + b*(x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4020 $\text{Int}[(a + b*\text{tan}[e + f*x] + (f*(x)))^m*((c + d*\text{tan}[e + f*x] + (f*(x))))], x_Symbol] \rightarrow \text{Simp}[c*(d/f) \text{ Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[(a + b*\text{tan}[e + f*x] + (f*(x)))^m*((c + d*\text{tan}[e + f*x] + (f*(x))))], x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 - I*\text{Tan}[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 + I*\text{Tan}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!IntegerQ}[m]$

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3462 vs. $2(112) = 224$.

Time = 0.15 (sec) , antiderivative size = 3463, normalized size of antiderivative = 26.04

method	result	size
parts	Expression too large to display	3463
derivativedivides	Expression too large to display	5570
default	Expression too large to display	5570

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x,method=_RETUR
NVERBOSE)
```

output

```

A*(1/4/f/d/(c^2+d^2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2))*(2*(c^2+d^2)
^(1/2)+2*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^2+1/4/f
*d/(c^2+d^2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2))*(2*(c^2+d^2)^(1/2)+2
*c)^(1/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)-1/4/f/d/(c^2+d^2)
^(3/2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1
/2)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^3-1/4/f*d/(c^2+d^2)^(
3/2)*ln(d*tan(f*x+e)+c+(c+d*tan(f*x+e))^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2
)+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c-1/f/d/(c^2+d^2)^(1/2)/(
2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d^2)
^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^2-1/f*d/(c^2+d^2)^(1/2)
)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d
^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))+1/f/d/(c^2+d^2)^(3/2)
/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2+d
^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^4+3/f*d/(c^2+d^2)^(3
/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2*(c^2
+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))*c^2+2/f*d^3/(c^2+d
^2)^(3/2)/(2*(c^2+d^2)^(1/2)-2*c)^(1/2)*arctan((2*(c+d*tan(f*x+e))^(1/2)+(2
*(c^2+d^2)^(1/2)+2*c)^(1/2))/(2*(c^2+d^2)^(1/2)-2*c)^(1/2))-1/4/f/d/(c^2+d
^2)*ln(d*tan(f*x+e)+c-(c+d*tan(f*x+e))^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)
+(c^2+d^2)^(1/2))*(2*(c^2+d^2)^(1/2)+2*c)^(1/2)*c^2-1/4/f*d/(c^2+d^2)*1...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3194 vs. $2(105) = 210$.

Time = 0.24 (sec) , antiderivative size = 3194, normalized size of antiderivative = 24.02

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algori
thm="fricas")

```

output

Too large to include

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{\sqrt{d \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/sqrt(d*tan(f*x + e) + c), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{%%}{1, [0,9,3]%%}+%%{4, [0,7,3]%%}+%%{6, [0,5,3]%%}+%%{4, [0,3,3]`

Mupad [B] (verification not implemented)

Time = 9.80 (sec) , antiderivative size = 4326, normalized size of antiderivative = 32.53

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/(c + d*tan(e + f*x))^(1/2),x)`

output

```

2*atanh((32*C^2*d^2*((-16*C^4*d^2*f^4)^(1/2)/(16*(c^2*f^4 + d^2*f^4)) - (C
^2*c*f^2)/(4*(c^2*f^4 + d^2*f^4)))^(1/2)*(c + d*tan(e + f*x))^(1/2))/((16*
C^3*c*d^3*f^3)/(c^2*f^4 + d^2*f^4) - (4*C*d^3*f^2*(-16*C^4*d^2*f^4)^(1/2))
/(c^2*f^5 + d^2*f^5)) + (8*c*d^2*((-16*C^4*d^2*f^4)^(1/2)/(16*(c^2*f^4 + d
^2*f^4)) - (C^2*c*f^2)/(4*(c^2*f^4 + d^2*f^4)))^(1/2)*(c + d*tan(e + f*x))
^(1/2)*(-16*C^4*d^2*f^4)^(1/2))/((16*C^3*c*d^5*f^5)/(c^2*f^4 + d^2*f^4) -
(4*C*d^5*f^4*(-16*C^4*d^2*f^4)^(1/2))/(c^2*f^5 + d^2*f^5) + (16*C^3*c^3*d^
3*f^5)/(c^2*f^4 + d^2*f^4) - (4*C*c^2*d^3*f^4*(-16*C^4*d^2*f^4)^(1/2))/(c^
2*f^5 + d^2*f^5)) - (32*C^2*c^2*d^2*f^2*((-16*C^4*d^2*f^4)^(1/2)/(16*(c^2*
f^4 + d^2*f^4)) - (C^2*c*f^2)/(4*(c^2*f^4 + d^2*f^4)))^(1/2)*(c + d*tan(e
+ f*x))^(1/2))/((16*C^3*c*d^5*f^5)/(c^2*f^4 + d^2*f^4) - (4*C*d^5*f^4*(-16
*C^4*d^2*f^4)^(1/2))/(c^2*f^5 + d^2*f^5) + (16*C^3*c^3*d^3*f^5)/(c^2*f^4 +
d^2*f^4) - (4*C*c^2*d^3*f^4*(-16*C^4*d^2*f^4)^(1/2))/(c^2*f^5 + d^2*f^5))
)*((-16*C^4*d^2*f^4)^(1/2)/(16*(c^2*f^4 + d^2*f^4)) - (C^2*c*f^2)/(4*(c^2*
f^4 + d^2*f^4)))^(1/2) - 2*atanh((8*c*d^2*(- (-16*C^4*d^2*f^4)^(1/2)/(16*(
c^2*f^4 + d^2*f^4)) - (C^2*c*f^2)/(4*(c^2*f^4 + d^2*f^4)))^(1/2)*(c + d*ta
n(e + f*x))^(1/2)*(-16*C^4*d^2*f^4)^(1/2))/((16*C^3*c*d^5*f^5)/(c^2*f^4 +
d^2*f^4) + (4*C*d^5*f^4*(-16*C^4*d^2*f^4)^(1/2))/(c^2*f^5 + d^2*f^5) + (16
*C^3*c^3*d^3*f^5)/(c^2*f^4 + d^2*f^4) + (4*C*c^2*d^3*f^4*(-16*C^4*d^2*f^4)
^(1/2))/(c^2*f^5 + d^2*f^5)) - (32*C^2*d^2*(- (-16*C^4*d^2*f^4)^(1/2)/(...

```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{2\sqrt{d \tan(fx + e) + c} a - \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) adf + \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{d \tan(fx + e) + c} dx \right) cdf + \left(\int \frac{\sqrt{d \tan(fx + e) + c}}{d \tan(fx + e) + c} dx \right) df$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```

(2*sqrt(tan(e + f*x)*d + c)*a - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)
**2)/(tan(e + f*x)*d + c),x)*a*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e +
f*x)**2)/(tan(e + f*x)*d + c),x)*c*d*f + int((sqrt(tan(e + f*x)*d + c)*ta
n(e + f*x))/(tan(e + f*x)*d + c),x)*b*d*f)/(d*f)

```

3.114 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$

Optimal result	1293
Mathematica [A] (verified)	1294
Rubi [A] (warning: unable to verify)	1294
Maple [B] (verified)	1298
Fricas [F(-1)]	1298
Sympy [F]	1299
Maxima [F(-2)]	1299
Giac [F(-2)]	1300
Mupad [B] (verification not implemented)	1300
Reduce [F]	1301

Optimal result

Integrand size = 47, antiderivative size = 210

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx$$

$$= -\frac{(iA + B - iC)\operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a - ib)\sqrt{c - id}f} - \frac{(A + iB - C)\operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(ia - b)\sqrt{c + id}f}$$

$$- \frac{2(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{\sqrt{b}(a^2 + b^2)\sqrt{bc - ad}f}$$

output

```
-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(a-I*b)/(c-I*d)
^(1/2)/f-(A+I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(I*a-b)/(
c+I*d)^(1/2)/f-2*(A*b^2-a*(B*b-C*a))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)
)/(-a*d+b*c)^(1/2)/b^(1/2)/(a^2+b^2)/(-a*d+b*c)^(1/2)/f
```

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.92

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{(-ia+b)(A-iB-C)\operatorname{arctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} + \frac{(ia+b)(A+iB-C)\operatorname{arctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} - \frac{2(Ab^2+a(-bB+aC))\operatorname{arctanh}\left(\frac{\sqrt{c+d\tan(e+fx)}}{\sqrt{b\sqrt{bc-ad}}}\right)}{(a^2 + b^2) f}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]),x]
```

output

```
((((-I)*a + b)*(A - I*B - C)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + ((I*a + b)*(A + I*B - C)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d] - (2*(A*b^2 + a*(-b*B) + a*C))*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]/(Sqrt[b]*Sqrt[b*c - a*d]))/((a^2 + b^2)*f)
```

Rubi [A] (warning: unable to verify)

Time = 2.16 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.93, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.255$, Rules used = {3042, 4136, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx$$

$$\downarrow \text{4136}$$

$$\begin{aligned}
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan^2(e+fx)+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx + \int \frac{bB+a(A-C)-(Ab-Cb-aB) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{bB+a(A-C)-(Ab-Cb-aB) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{4022} \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \\
 & \frac{\frac{1}{2}(a - ib)(A + iB - C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(a + ib)(A - iB - C) \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \\
 & \frac{\frac{1}{2}(a - ib)(A + iB - C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(a + ib)(A - iB - C) \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{4020} \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \\
 & \frac{i(a+ib)(A-iB-C) \int -\frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - \frac{i(a-ib)(A+iB-C) \int -\frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f}}{a^2 + b^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \\
 & \frac{i(a-ib)(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx)) - \frac{i(a+ib)(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f}}{a^2 + b^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \\
 & \frac{(a-ib)(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)} + (a+ib)(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} \\
 & \quad \downarrow \\
 & \frac{\hspace{10em}}{a^2 + b^2}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 221 \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} + \\
 & \frac{(a+ib)(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} + \frac{(a-ib)(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}} \\
 & \downarrow 4117 \\
 & \frac{(Ab^2 - a(bB - aC)) \int \frac{1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d \tan(e + fx)}{f(a^2 + b^2)} + \\
 & \frac{(a+ib)(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} + \frac{(a-ib)(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}} \\
 & \downarrow 73 \\
 & \frac{2(Ab^2 - a(bB - aC)) \int \frac{1}{a+\frac{b(c+d \tan(e+fx))}{d}-\frac{bc}{d}} d\sqrt{c+d \tan(e+fx)}}{df(a^2 + b^2)} + \\
 & \frac{(a+ib)(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} + \frac{(a-ib)(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}} \\
 & \downarrow 221 \\
 & \frac{2(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{\sqrt{b}f(a^2 + b^2)\sqrt{bc-ad}} + \\
 & \frac{(a+ib)(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} + \frac{(a-ib)(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]),x]`

output `((a + I*b)*(A - I*B - C)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]]/(Sqrt[c - I*d]*f) + ((a - I*b)*(A + I*B - C)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]]/(Sqrt[c + I*d]*f))/(a^2 + b^2) - (2*(A*b^2 - a*(b*B - a*C))*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]]/(Sqrt[b]*(a^2 + b^2)*Sqrt[b*c - a*d]*f)`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{\text{m}_.}) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{n}_.}), \text{x_Symbol}] \rightarrow \text{With}[\{p = \text{Denominator}[\text{m}]\}, \text{Simp}[p/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{(p*(\text{m} + 1) - 1)} * (\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^p/\text{b}))^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{(1/p)}], \text{x}]] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[-1, \text{m}, 0] \&\& \text{LeQ}[-1, \text{n}, 0] \&\& \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \&\& \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a}) * \text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_., \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])], \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d}) * \text{x})^{\text{m}} / (\text{d}^2 + \text{c} * \text{x}), \text{x}], \text{x}, \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I} * \text{d}) / 2 \quad \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 - \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I} * \text{d}) / 2 \quad \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 + \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{!IntegerQ}[\text{m}]$
- rule 4117 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{n}_.} * ((\text{A}_.) + (\text{C}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{A}/\text{f} \quad \text{Subst}[\text{Int}[(\text{a} + \text{b} * \text{x})^{\text{m}} * (\text{c} + \text{d} * \text{x})^{\text{n}}, \text{x}], \text{x}, \text{Tan}[\text{e} + \text{f} * \text{x}]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{C}, \text{m}, \text{n}\}, \text{x}] \&\& \text{EqQ}[\text{A}, \text{C}]$

rule 4136

```

Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 13473 vs. $2(179) = 358$.

Time = 0.20 (sec) , antiderivative size = 13474, normalized size of antiderivative = 64.16

method	result	size
derivativedivides	Expression too large to display	13474
default	Expression too large to display	13474

input

```

int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(1/2
),x,method=_RETURNVERBOSE)

```

output

result too large to display

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input

```

integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e)
)^(1/2),x, algorithm="fricas")

```

output

Timed out

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))*sqrt(c + d*tan(e + f*x))), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

Mupad [B] (verification not implemented)

Time = 61.11 (sec) , antiderivative size = 25341, normalized size of antiderivative = 120.67

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^(1/2)),x)`

output

```
(log((((((((((128*C*b^2*d^8*(a*d + b*c)^2*(a^2 + b^2)^2)/f - 64*b^2*d^8*(a^2 + b^2)^2*(c + d*tan(e + f*x))^(1/2)*((4*(-C^4*f^4*(a^2*d - b^2*d + 2*a*b*c)^2)^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 + 8*C^2*a*b*d*f^2)/(f^4*(a^2 + b^2)^2*(c^2 + d^2))))^(1/2)*(3*b^3*c^2 + 2*b^3*d^2 - a^2*b*c^2 - 2*a^2*b*d^2 + a^3*c*d + a*b^2*c*d))*((4*(-C^4*f^4*(a^2*d - b^2*d + 2*a*b*c)^2)^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 + 8*C^2*a*b*d*f^2)/(f^4*(a^2 + b^2)^2*(c^2 + d^2))))^(1/2))/4 + (64*C^2*b*d^8*(c + d*tan(e + f*x))^(1/2)*(5*b^6*c - 4*a^6*c - 2*a^2*b^4*c + 5*a^4*b^2*c - 2*a^3*b^3*d + 7*a*b^5*d + 7*a^5*b*d))/f^2)*((4*(-C^4*f^4*(a^2*d - b^2*d + 2*a*b*c)^2)^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 + 8*C^2*a*b*d*f^2)/(f^4*(a^2 + b^2)^2*(c^2 + d^2))))^(1/2))/4 + (32*C^3*b*d^8*(4*a^5*d - b^5*c - 9*a^2*b^3*c - 15*a^3*b^2*d + 12*a^4*b*c + a*b^4*d))/f^3)*((4*(-C^4*f^4*(a^2*d - b^2*d + 2*a*b*c)^2)^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 + 8*C^2*a*b*d*f^2)/(f^4*(a^2 + b^2)^2*(c^2 + d^2))))^(1/2))/4 - (32*C^4*b*d^8*(2*a^4 + b^4)*(c + d*tan(e + f*x))^(1/2))/f^4)*((4*(-C^4*f^4*(a^2*d - b^2*d + 2*a*b*c)^2)^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 + 8*C^2*a*b*d*f^2)/(f^4*(a^2 + b^2)^2*(c^2 + d^2))))^(1/2))/4 + (32*C^5*a^2*b^2*d^8)/f^5)*(((32*C^4*a^2*b^2*d^2*f^4 - 16*C^4*b^4*d^2*f^4 - 64*C^4*a^2*b^2*c^2*f^4 - 16*C^4*a^4*d^2*f^4 + 64*C^4*a*b^3*c*d*f^4 - 64*C^4*a^3*b*c*d*f^4)^(1/2) - 4*C^2*a^2*c*f^2 + 4*C^2*b^2*c*f^2 + 8*C^2*a*b*d*f^2)/(a^4*c^2*f^4 + a^4*d^2*f^4 + b^4*c^2*f^4 + b^...
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{2\sqrt{d \tan(fx + e) + c} - \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^3}{\tan(fx + e)^2 b d + \tan(fx + e) a d + \tan(fx + e) b c + a c} dx \right) b d f - \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)}{\tan(fx + e)^2 b d + \tan(fx + e) a d + \tan(fx + e) b c + a c} dx \right) d f}{d f}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(1/2),x)
```

output

```
(2*sqrt(tan(e + f*x)*d + c) - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**
3)/(tan(e + f*x)**2*b*d + tan(e + f*x)*a*d + tan(e + f*x)*b*c + a*c),x)*b*
d*f - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*b*d
+ tan(e + f*x)*a*d + tan(e + f*x)*b*c + a*c),x)*a*d*f + int((sqrt(tan(e +
f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*b*d + tan(e + f*x)*a*d + tan
(e + f*x)*b*c + a*c),x)*c*d*f)/(d*f)
```

3.115
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1303
Mathematica [A] (verified)	1304
Rubi [A] (warning: unable to verify)	1304
Maple [B] (verified)	1310
Fricas [F(-1)]	1310
Sympy [F]	1311
Maxima [F(-2)]	1311
Giac [F(-2)]	1312
Mupad [B] (verification not implemented)	1312
Reduce [F]	1313

Optimal result

Integrand size = 47, antiderivative size = 327

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx$$

$$= -\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(a - ib)^2 \sqrt{c - id} f} - \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a + ib)^2 \sqrt{c + id} f}$$

$$- \frac{(3a^3 b B d - a^4 C d + b^4 (2Bc - Ad) + ab^3 (4Ac - 4cC - Bd) - a^2 b^2 (2Bc + 5Ad - 3Cd)) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{c}}{\sqrt{a^2 + b^2}}\right)}{\sqrt{b} (a^2 + b^2)^2 (bc - ad)^{3/2} f}$$

$$- \frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{(a^2 + b^2) (bc - ad) f (a + b \tan(e + fx))}$$

output

```
-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(a-I*b)^2/(c-I*d)^(1/2)/f-(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(a+I*b)^2/(c+I*d)^(1/2)/f-(3*a^3*b*B*d-a^4*C*d+b^4*(-A*d+2*B*c)+a*b^3*(4*A*c-B*d-4*C*c)-a^2*b^2*(5*A*d+2*B*c-3*C*d))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/b^(1/2)/(a^2+b^2)^2/(-a*d+b*c)^(3/2)/f-(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(1/2)/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))
```

Mathematica [A] (verified)

Time = 4.91 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.03

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{i \left(\frac{(a+ib)^2(A-iB-C)(bc-ad) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} + \frac{(a-ib)^2(A+iB-C)(-bc+ad) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} \right)}{a^2+b^2} + \frac{(-3a^3bBd+a^4C)}{(a^2+b^2)}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*
Sqrt[c + d*Tan[e + f*x]]),x]
```

output

```
(((-I)*(((a + I*b)^2*(A - I*B - C)*(b*c - a*d)*ArcTanh[Sqrt[c + d*Tan[e +
f*x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + ((a - I*b)^2*(A + I*B - C)*(-(b*c) +
a*d)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d]))/(a^
2 + b^2) + ((-3*a^3*b*B*d + a^4*C*d + b^4*(-2*B*c + A*d) + a*b^3*(-4*A*c +
4*c*C + B*d) + a^2*b^2*(2*B*c + 5*A*d - 3*C*d))*ArcTanh[(Sqrt[b]*Sqrt[c +
d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/(Sqrt[b]*(a^2 + b^2)*Sqrt[b*c - a*d])
- ((A*b^2 + a*(-(b*B) + a*C))*Sqrt[c + d*Tan[e + f*x]]/(a + b*Tan[e + f*x
]))/((a^2 + b^2)*(b*c - a*d)*f)
```

Rubi [A] (warning: unable to verify)Time = 3.79 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.09, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.340$, Rules used = {3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx$$

↓ 4132

$$\frac{\int \frac{(2A-C)da^2 - b(2Ac - 2Cc - Bd)a + (Ab^2 - a(bB - aC))d \tan^2(e+fx) - b^2(2Bc - Ad) + 2(Ab - Cb - aB)(bc - ad) \tan(e+fx)}{2(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx}{\frac{(a^2 + b^2)(bc - ad)}{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}} = \frac{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}$$

↓ 27

$$\frac{\int \frac{(2A-C)da^2 - b(2Ac - 2Cc - Bd)a + (Ab^2 - a(bB - aC))d \tan^2(e+fx) - b^2(2Bc - Ad) + 2(Ab - Cb - aB)(bc - ad) \tan(e+fx)}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx}{\frac{2(a^2 + b^2)(bc - ad)}{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}} = \frac{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}$$

↓ 3042

$$\frac{\int \frac{(2A-C)da^2 - b(2Ac - 2Cc - Bd)a + (Ab^2 - a(bB - aC))d \tan(e+fx)^2 - b^2(2Bc - Ad) + 2(Ab - Cb - aB)(bc - ad) \tan(e+fx)}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx}{\frac{2(a^2 + b^2)(bc - ad)}{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}} = \frac{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}$$

↓ 4136

$$\frac{\int -\frac{2((A-C)a^2 + 2bBa - b^2(A-C))(bc - ad) + (Ba^2 - 2b(A-C)a - b^2B) \tan(e+fx)(bc - ad)}{\sqrt{c+d \tan(e+fx)} a^2 + b^2} dx}{\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}} = \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4a^2 + b^2)(bc - ad))}{2(a^2 + b^2)(bc - ad)}$$

↓ 27

$$\frac{2 \int \frac{((A-C)a^2 + 2bBa - b^2(A-C))(bc - ad) + (Ba^2 - 2b(A-C)a - b^2B) \tan(e+fx)(bc - ad)}{\sqrt{c+d \tan(e+fx)} a^2 + b^2} dx}{\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))}} = \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4a^2 + b^2)(bc - ad))}{2(a^2 + b^2)(bc - ad)}$$

↓ 3042

$$\begin{aligned}
 & \frac{2 \int \frac{((A-C)a^2 + 2bBa - b^2(A-C))(bc-ad) + (Ba^2 - 2b(A-C)a - b^2B) \tan(e+fx)(bc-ad)}{\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad+2Bc-3Cd) + ab^3(4A+3B))}{2(a^2+b^2)(bc-ad)} \\
 & \frac{(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} \\
 & \quad \downarrow 4022 \\
 & \frac{(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad+2Bc-3Cd) + ab^3(4Ac-Bd-4cC) + b^4(2Bc-Ad)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{2\left(\frac{1}{2}(a-ib)^2(A+iB)\right)}{2(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 3042 \\
 & \frac{(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad+2Bc-3Cd) + ab^3(4Ac-Bd-4cC) + b^4(2Bc-Ad)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{2\left(\frac{1}{2}(a-ib)^2(A+iB)\right)}{2(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 4020 \\
 & \frac{(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad+2Bc-3Cd) + ab^3(4Ac-Bd-4cC) + b^4(2Bc-Ad)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{2\left(\frac{i(a+ib)^2(A-ib)}{2(a^2+b^2)(bc-ad)}\right)}{2(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 25 \\
 & \frac{(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad+2Bc-3Cd) + ab^3(4Ac-Bd-4cC) + b^4(2Bc-Ad)) \int \frac{\tan(e+fx)^2+1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2+b^2} - \frac{2\left(\frac{i(a-ib)^2(A+iB-i)}{2(a^2+b^2)(bc-ad)}\right)}{2(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 73
 \end{aligned}$$

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4Ac - Bd - 4cC) + b^4(2Bc - Ad)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{2 \left(\frac{(a-ib)^2(A+iB-C)}{f \sqrt{bc-ad}} \right)}{2(a^2 + b^2)(bc - ad)}$$

221

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4Ac - Bd - 4cC) + b^4(2Bc - Ad)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{2 \left(\frac{(a-ib)^2(A+iB-C)}{f \sqrt{bc-ad}} \right)}{2(a^2 + b^2)(bc - ad)}$$

4117

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} - \frac{(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4Ac - Bd - 4cC) + b^4(2Bc - Ad)) \int \frac{1}{(a+b \tan(e+fx)) \sqrt{c+d \tan(e+fx)}} d \tan(e+fx)}{f(a^2 + b^2)} - \frac{2 \left(\frac{(a-ib)^2(A+iB-C)}{f \sqrt{bc-ad}} \right)}{2(a^2 + b^2)(bc - ad)}$$

73

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} - \frac{2(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4Ac - Bd - 4cC) + b^4(2Bc - Ad)) \int \frac{1}{a + \frac{b(c+d \tan(e+fx))}{d} - \frac{bc}{d}} d \sqrt{c+d \tan(e+fx)}}{df(a^2 + b^2)} - \frac{2 \left(\frac{(a-ib)^2(A+iB-C)}{f \sqrt{bc-ad}} \right)}{2(a^2 + b^2)(bc - ad)}$$

221

$$\frac{(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))} - \frac{2(a^4(-C)d + 3a^3bBd - a^2b^2(5Ad + 2Bc - 3Cd) + ab^3(4Ac - Bd - 4cC) + b^4(2Bc - Ad)) \operatorname{arctanh} \left(\frac{\sqrt{b} \sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}} \right)}{\sqrt{bf(a^2 + b^2)} \sqrt{bc-ad}} - \frac{2 \left(\frac{(a-ib)^2(A+iB-C)(bc - ad)}{f \sqrt{bc-ad}} \right)}{2(a^2 + b^2)(bc - ad)}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*Sqrt[c + d*Tan[e + f*x]]),x]
```


output

```
-1/2*((-2*((a + I*b)^2*(A - I*B - C)*(b*c - a*d)*ArcTan[Tan[e + f*x]/Sqrt
[c - I*d]])/(Sqrt[c - I*d]*f) + ((a - I*b)^2*(A + I*B - C)*(b*c - a*d)*Arc
Tan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f)))/(a^2 + b^2) + (2*(3*a
^3*b*B*d - a^4*C*d + b^4*(2*B*c - A*d) + a*b^3*(4*A*c - 4*c*C - B*d) - a^2
*b^2*(2*B*c + 5*A*d - 3*C*d))*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/S
qrt[b*c - a*d]])/(Sqrt[b]*(a^2 + b^2)*Sqrt[b*c - a*d]*f)/((a^2 + b^2)*(b*
c - a*d)) - ((A*b^2 - a*(b*B - a*C))*Sqrt[c + d*Tan[e + f*x]])/((a^2 + b^2
)*(b*c - a*d)*f*(a + b*Tan[e + f*x]))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4020

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4117

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2]), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2]), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)^2]))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 20869 vs. $2(294) = 588$.

Time = 0.23 (sec) , antiderivative size = 20870, normalized size of antiderivative = 63.82

method	result	size
derivativedivides	Expression too large to display	20870
default	Expression too large to display	20870

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")
```

output

```
Timed out
```

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**2/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**2 *sqrt(c + d*tan(e + f*x))), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to rounding error%%{%%}{1, [0,14,6]%%}+%%{6, [0,12,6]%%}+%%{15, [0,10,6]%%}+%%{20, [0`

Mupad [B] (verification not implemented)

Time = 46.47 (sec) , antiderivative size = 225004, normalized size of antiderivative = 688.09

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^(1/2)),x)`

output

```
(atan(((((((16*(8*C^3*a^6*b^7*d^11*f^2 - 78*C^3*a^4*b^9*d^11*f^2 + 60*C^3*
a^8*b^5*d^11*f^2 - 24*C^3*a^10*b^3*d^11*f^2 + 2*C^3*a^12*b*d^11*f^2 - 32*C
^3*a*b^12*c^3*d^8*f^2 + 152*C^3*a^3*b^10*c*d^10*f^2 + 128*C^3*a^5*b^8*c*d
^10*f^2 - 64*C^3*a^7*b^6*c*d^10*f^2 - 32*C^3*a^9*b^4*c*d^10*f^2 + 8*C^3*a^1
1*b^2*c*d^10*f^2 - 40*C^3*a^2*b^11*c^2*d^9*f^2 + 64*C^3*a^3*b^10*c^3*d^8*f
^2 - 216*C^3*a^4*b^9*c^2*d^9*f^2 + 96*C^3*a^5*b^8*c^3*d^8*f^2 - 120*C^3*a^
6*b^7*c^2*d^9*f^2 + 56*C^3*a^8*b^5*c^2*d^9*f^2)))/((a^10*d^2*f^5 + b^10*c^2*
f^5 + 4*a^2*b^8*c^2*f^5 + 6*a^4*b^6*c^2*f^5 + 4*a^6*b^4*c^2*f^5 + a^8*b^2*
c^2*f^5 + a^2*b^8*d^2*f^5 + 4*a^4*b^6*d^2*f^5 + 6*a^6*b^4*d^2*f^5 + 4*a^8*
b^2*d^2*f^5 - 2*a*b^9*c*d*f^5 - 2*a^9*b*c*d*f^5 - 8*a^3*b^7*c*d*f^5 - 12*a
^5*b^5*c*d*f^5 - 8*a^7*b^3*c*d*f^5) - (((((16*(40*C*a^3*b^14*d^12*f^4 + 19
2*C*a^5*b^12*d^12*f^4 + 360*C*a^7*b^10*d^12*f^4 + 320*C*a^9*b^8*d^12*f^4 +
120*C*a^11*b^6*d^12*f^4 - 8*C*a^15*b^2*d^12*f^4 + 8*C*b^17*c^3*d^9*f^4 +
40*C*a*b^16*c^2*d^10*f^4 + 32*C*a*b^16*c^4*d^8*f^4 - 88*C*a^2*b^15*c*d^11*
f^4 - 448*C*a^4*b^13*c*d^11*f^4 - 920*C*a^6*b^11*c*d^11*f^4 - 960*C*a^8*b^
9*c*d^11*f^4 - 520*C*a^10*b^7*c*d^11*f^4 - 128*C*a^12*b^5*c*d^11*f^4 - 8*C
*a^14*b^3*c*d^11*f^4 - 32*C*a^2*b^15*c^3*d^9*f^4 + 256*C*a^3*b^14*c^2*d^10
*f^4 + 160*C*a^3*b^14*c^4*d^8*f^4 - 280*C*a^4*b^13*c^3*d^9*f^4 + 680*C*a^5
*b^12*c^2*d^10*f^4 + 320*C*a^5*b^12*c^4*d^8*f^4 - 640*C*a^6*b^11*c^3*d^9*f
^4 + 960*C*a^7*b^10*c^2*d^10*f^4 + 320*C*a^7*b^10*c^4*d^8*f^4 - 680*C*a...
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(1
/2),x)
```

output

```
(4*sqrt(tan(e + f*x)*d + c)*b*c - 2*sqrt(tan(e + f*x)*d + c)*c*d + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**2 + tan(e + f*x)*a**3*d**2 - 4*tan(e + f*x)*a*b**2*c**2 + a**3*c*d - 2*a**2*b*c**2),x)*tan(e + f*x)*a**3*b*d**3*f - 6*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**2 + tan(e + f*x)*a**3*d**2 - 4*tan(e + f*x)*a*b**2*c**2 + a**3*c*d - 2*a**2*b*c**2),x)*tan(e + f*x)*a**2*b**2*c*d**2*f + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**2 + tan(e + f*x)*a**3*d**2 - 4*tan(e + f*x)*a*b**2*c**2 + a**3*c*d - 2*a**2*b*c**2),x)*tan(e + f*x)*a**2*b*c*d**3*f + 12*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**2 + tan(e + f*x)*a**3*d**2 - 4*tan(e + f*x)*a*b**2*c**2 + a**3*c*d - 2*a**2*b*c**2),x)*tan(e + f*x)*a*b**3*c**2*d*f - 4*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*a*b**2*d**2 - 2*tan(e + f*x)**3*b**3*c*d + 2*tan(e + f*x)**2*a**2*b*d**2 - 3*tan(e + f*x)**2*a*b**2*c*d - 2*tan(e + f*x)**2*b**3*c**2 + tan(e + f*x)*a**3*d**2 - 4*tan(e + f...
```

3.116
$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1315
Mathematica [C] (verified)	1316
Rubi [A] (warning: unable to verify)	1317
Maple [B] (verified)	1324
Fricas [F(-1)]	1324
Sympy [F]	1325
Maxima [F(-1)]	1325
Giac [F]	1325
Mupad [F(-1)]	1326
Reduce [F]	1326

Optimal result

Integrand size = 47, antiderivative size = 511

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(a - ib)^3 (iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{3/2} f}$$

$$- \frac{(ia - b)^3 (A + iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{3/2} f}$$

$$- \frac{2(c^2 C - Bcd + Ad^2) (a + b \tan(e + fx))^3}{d (c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

$$+ \frac{2b(6a^2 d^3 (12c^2 C - 5Bcd + (5A + 7C)d^2) - 15abd(8c^3 C - 6Bc^2 d + c(3A + 5C)d^2 - 3Bd^3) + b^2(48c^4 C - 15d^4 (c^2 + d^2) f)}{15d^4 (c^2 + d^2) f}$$

$$- \frac{2b^2(4(bc - ad) (6c^2 C - 5Bcd + (5A + C)d^2) - 5d^2((A - C)(bc - ad) + B(ac + bd))) \tan(e + fx) \sqrt{c + d \tan(e + fx)}}{15d^3 (c^2 + d^2) f}$$

$$+ \frac{2b(6c^2 C - 5Bcd + (5A + C)d^2) (a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5d^2 (c^2 + d^2) f}$$

output

```

-(a-I*b)^3*(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(3/2)/f-(I*a-b)^3*(A+I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(3/2)/f-2*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^3/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)+2/15*b*(6*a^2*d^2*(12*c^2*C-5*B*c*d+(5*A+7*C)*d^2)-15*a*b*d*(8*c^3*C-6*B*c^2*d+c*(3*A+5*C)*d^2-3*B*d^3)+b^2*(48*c^4*C-40*B*c^3*d+6*c^2*(5*A+3*C)*d^2-25*B*c*d^3+15*(A-C)*d^4))*(c+d*tan(f*x+e))^(1/2)/d^4/(c^2+d^2)/f-2/15*b^2*(4*(-a*d+b*c)*(6*c^2*C-5*B*c*d+(5*A+C)*d^2)-5*d^2*((A-C)*(-a*d+b*c)+B*(a*c+b*d))*tan(f*x+e)*(c+d*tan(f*x+e))^(1/2)/d^3/(c^2+d^2)/f+2/5*b*(6*c^2*C-5*B*c*d+(5*A+C)*d^2)*(a+b*tan(f*x+e))^2*(c+d*tan(f*x+e))^(1/2)/d^2/(c^2+d^2)/f
    
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.54 (sec) , antiderivative size = 920, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \frac{2C(a + b \tan(e + fx))^3}{5df \sqrt{c + d \tan(e + fx)}}$$

$$\begin{aligned}
 & 2 \left(\frac{(-6bcC+5bBd+6aCd)(a+b \tan(e+fx))^2}{3df \sqrt{c+d \tan(e+fx)}} + \right. \\
 & \left. 2 \left(\frac{(15b(Ab+aB-bC)d^2+4(bc-ad)(6bcC-5bBd-6aCd))(a+b \tan(e+fx))}{2df \sqrt{c+d \tan(e+fx)}} + \frac{2(-48b^3c^3C+40b^3Bc^2c}{\dots} \right) \right)
 \end{aligned}$$

input

```
Integrate[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^(3/2),x]
```

output

```
(2*C*(a + b*Tan[e + f*x])^3)/(5*d*f*Sqrt[c + d*Tan[e + f*x]]) + (2*((( -6*b
*c*C + 5*b*B*d + 6*a*C*d)*(a + b*Tan[e + f*x])^2)/(3*d*f*Sqrt[c + d*Tan[e
+ f*x]]) + (2*(((15*b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(6*b*c*C - 5*b
*B*d - 6*a*C*d))*(a + b*Tan[e + f*x]))/(2*d*f*Sqrt[c + d*Tan[e + f*x]]) +
((-2*(-48*b^3*c^3*C + 40*b^3*B*c^2*d + 144*a*b^2*c^2*C*d - 30*A*b^3*c*d^2
- 110*a*b^2*B*c*d^2 - 144*a^2*b*c*C*d^2 + 30*b^3*c*C*d^2 + 60*a*A*b^2*d^3
+ 85*a^2*b*B*d^3 - 15*b^3*B*d^3 + 48*a^3*C*d^3 - 60*a*b^2*C*d^3))/(d*Sqrt[
c + d*Tan[e + f*x]]) + (2*(((45*a^2*A*b*d^3 - 15*A*b^3*d^3 + 15*a^3*B*d^3
- 45*a*b^2*B*d^3 - 45*a^2*b*C*d^3 + 15*b^3*C*d^3)*((-I)*ArcTanh[Sqrt[c +
d*Tan[e + f*x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + (I*ArcTanh[Sqrt[c + d*Tan[
e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d]))/2 + ((-1/2*(c*d*(45*a^2*A*b*d^3
- 15*A*b^3*d^3 + 15*a^3*B*d^3 - 45*a*b^2*B*d^3 - 45*a^2*b*C*d^3 + 15*b^3*C
*d^3)) + d^2*(-48*b^3*c^3*C + 40*b^3*B*c^2*d + 144*a*b^2*c^2*C*d - 30*A*b
^3*c*d^2 - 110*a*b^2*B*c*d^2 - 144*a^2*b*c*C*d^2 + 30*b^3*c*C*d^2 + 15*a^3
*A*d^3 + 15*a*A*b^2*d^3 + 40*a^2*b*B*d^3 + 33*a^3*C*d^3 - 15*a*b^2*C*d^3)/
2 + (48*b^3*c^3*C - 40*b^3*B*c^2*d - 144*a*b^2*c^2*C*d + 30*A*b^3*c*d^2 +
110*a*b^2*B*c*d^2 + 144*a^2*b*c*C*d^2 - 30*b^3*c*C*d^2 - 60*a*A*b^2*d^3 -
85*a^2*b*B*d^3 + 15*b^3*B*d^3 - 48*a^3*C*d^3 + 60*a*b^2*C*d^3)/2))*(-Hype
rgeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)]/((I*c + d)*Sqr
t[c + d*Tan[e + f*x]])) + Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e ...
```

Rubi [A] (warning: unable to verify)

Time = 6.25 (sec) , antiderivative size = 512, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.383$, Rules used = {3042, 4128, 27, 3042, 4130, 27, 3042, 4120, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$2 \int \frac{(a + b \tan(e + fx))^2 (b(6C^2 - 5Bdc + (5A + C)d^2) \tan^2(e + fx) + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + 6bd) + 2(3bc - \frac{ad}{2})(cC - Bd))}{2\sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{d(c^2 + d^2)}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\int \frac{(a + b \tan(e + fx))^2 (b(6C^2 - 5Bdc + (5A + C)d^2) \tan^2(e + fx) + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + 6bd) + (6bc - ad)(cC - Bd))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{d(c^2 + d^2)}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2 (b(6C^2 - 5Bdc + (5A + C)d^2) \tan^2(e + fx) + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + 6bd) + (6bc - ad)(cC - Bd))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{d(c^2 + d^2)}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4130

$$2 \int \frac{(a + b \tan(e + fx)) (-5((Bc - (A - C)d)a^2 + 2b(Ac - Cc + Bd)a - b^2(Bc - (A - C)d)) \tan(e + fx) + d^2 - 5a(Ad(ac + 6bd) + (6bc - ad)(cC - Bd)) + b(4(bc - ad)(6C^2 - 5Bdc + (5A + C)d^2)))}{2\sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{d(c^2 + d^2)}{5df} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{2b(d^2(5A + C) - 5Bcd + 6c^2C)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \int \frac{(a + b \tan(e + fx)) (-5((Bc - (A - C)d)a^2 + 2b(Ac - Cc + Bd)a - b^2(Bc - (A - C)d)) \tan(e + fx) + d^2 - 5a(Ad(ac + 6bd) + (6bc - ad)(cC - Bd)) + b(4(bc - ad)(6C^2 - 5Bdc + (5A + C)d^2)))}{5df} dx$$

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \int \frac{(a+b \tan(e+fx))(-5((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d))}{3df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4120

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4113

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4022

$$- \frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

↓ 3042

$$- \frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

↓ 4020

$$- \frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

↓ 25

$$- \frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b(d^2(5A+C)-5Bcd+6c^2C)(a+b \tan(e+fx))^2 \sqrt{c+d \tan(e+fx)}}{5df} - \frac{2b^2 \tan(e+fx) \sqrt{c+d \tan(e+fx)} (4(bc-ad)(d^2(5A+C)-5Bcd+6c^2C)-5d^2((A-C))}{3df}$$

↓ 73

$$-\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b(d^2(5A+C) - 5Bcd + 6c^2C)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b^2 \tan(e + fx) \sqrt{c + d \tan(e + fx)} (4(bc - ad)(d^2(5A+C) - 5Bcd + 6c^2C) - 5d^2((A-C)))}{3df}$$

↓ 221

$$-\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b(d^2(5A+C) - 5Bcd + 6c^2C)(a + b \tan(e + fx))^2 \sqrt{c + d \tan(e + fx)}}{5df} - \frac{2b^2 \tan(e + fx) \sqrt{c + d \tan(e + fx)} (4(bc - ad)(d^2(5A+C) - 5Bcd + 6c^2C) - 5d^2((A-C)))}{3df}$$

input

```
Int[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x]
```

output

```
(-2*(c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^3)/(d*(c^2 + d^2)*f*sqrt[c + d*Tan[e + f*x]]) + ((2*b*(6*c^2*C - 5*B*c*d + (5*A + C)*d^2)*(a + b*Tan[e + f*x])^2*sqrt[c + d*Tan[e + f*x]])/(5*d*f) - ((2*b^2*(4*(b*c - a*d)*(6*c^2*C - 5*B*c*d + (5*A + C)*d^2) - 5*d^2*((A - C)*(b*c - a*d) + B*(a*c + b*d)))*Tan[e + f*x]*sqrt[c + d*Tan[e + f*x]])/(3*d*f) - ((15*(a - I*b)^3*(A - I*B - C)*(c + I*d)*d^3*ArcTan[Tan[e + f*x]/sqrt[c - I*d]])/(sqrt[c - I*d]*f) + (15*(a + I*b)^3*(A + I*B - C)*(c - I*d)*d^3*ArcTan[Tan[e + f*x]/sqrt[c + I*d]])/(sqrt[c + I*d]*f) + (2*b*(6*a^2*d^2*(12*c^2*C - 5*B*c*d + (5*A + 7*C)*d^2) - 15*a*b*d*(8*c^3*C - 6*B*c^2*d + c*(3*A + 5*C)*d^2 - 3*B*d^3) + b^2*(48*c^4*C - 40*B*c^3*d + 6*c^2*(5*A + 3*C)*d^2 - 25*B*c*d^3 + 15*(A - C)*d^4))*sqrt[c + d*Tan[e + f*x]])/(d*f)/(3*d)/(5*d)/(d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```


Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 11254 vs. $2(476) = 952$.

Time = 0.47 (sec) , antiderivative size = 11255, normalized size of antiderivative = 22.03

method	result	size
parts	Expression too large to display	11255
derivatividivides	Expression too large to display	49725
default	Expression too large to display	49725

input `int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((a+b*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((a + b*tan(e + f*x))**3*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(3/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3/2,x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^3}{(d \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3/2,x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^3/(d*tan(f*x + e) + c)^3/2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input

```
int(((a + b*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c +
d*tan(e + f*x))^(3/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3
/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**4 + int((sqrt(tan(e + f*x)*d + c)*tan(e
+ f*x)**5)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f
*x)**b**3*c*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5)/(tan(e
+ f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*b**3*c**2*d*f + 3*int((sqrt
(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*
x)*c*d + c**2),x)*tan(e + f*x)*a*b**2*c*d**2*f + int((sqrt(tan(e + f*x)*d
+ c)*tan(e + f*x)**4)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x
)*tan(e + f*x)*b**4*d**2*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)*
**4)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*b**2*c**2*d*f
+ int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**2*d**2 + 2
*tan(e + f*x)*c*d + c**2),x)*b**4*c*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*
tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan
(e + f*x)*a**2*b*c*d**2*f + 4*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**
3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a*b
**3*d**2*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)
**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a**2*b*c**2*d*f + 4*int((sqrt(tan
(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c
*d + c**2),x)*a*b**3*c*d*f - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2
)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a**4*
d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**...
```

$$3.117 \quad \int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1328
Mathematica [C] (verified)	1329
Rubi [A] (warning: unable to verify)	1330
Maple [B] (verified)	1335
Fricas [F(-1)]	1336
Sympy [F]	1336
Maxima [F(-1)]	1337
Giac [F]	1337
Mupad [B] (verification not implemented)	1337
Reduce [F]	1338

Optimal result

Integrand size = 47, antiderivative size = 343

$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx =$$

$$\frac{(a-ib)^2 (iA+B-iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c-id)^{3/2} f}$$

$$- \frac{(a+ib)^2 (B-i(A-C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c+id)^{3/2} f}$$

$$- \frac{2(c^2 C - Bcd + Ad^2) (a+b \tan(e+fx))^2}{d(c^2 + d^2) f \sqrt{c+d \tan(e+fx)}}$$

$$+ \frac{2b(6ad(2c^2 C - Bcd + (A+C)d^2) - b(8c^3 C - 6Bc^2 d + c(3A + 5C)d^2 - 3Bd^3)) \sqrt{c+d \tan(e+fx)}}{3d^3 (c^2 + d^2) f}$$

$$+ \frac{2b^2(4c^2 C - 3Bcd + (3A + C)d^2) \tan(e+fx) \sqrt{c+d \tan(e+fx)}}{3d^2 (c^2 + d^2) f}$$

output

```

-(a-I*b)^2*(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(3/2)/f-(a+I*b)^2*(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(3/2)/f-2*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^2/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)+2/3*b*(6*a*d*(2*c^2*C-B*c*d+(A+C)*d^2)-b*(8*c^3*C-6*B*c^2*d+c*(3*A+5*C)*d^2-3*B*d^3))*(c+d*tan(f*x+e))^(1/2)/d^3/(c^2+d^2)/f+2/3*b^2*(4*c^2*C-3*B*c*d+(3*A+C)*d^2)*tan(f*x+e)*(c+d*tan(f*x+e))^(1/2)/d^2/(c^2+d^2)/f
    
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.36 (sec) , antiderivative size = 476, normalized size of antiderivative = 1.39

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \frac{2C(a + b \tan(e + fx))^2}{3df \sqrt{c + d \tan(e + fx)}}$$

$$2 \left(\frac{(-4bcC+3bBd+4aCd)(a+b \tan(e+fx))}{df \sqrt{c+d \tan(e+fx)}} + \frac{2(8b^2c^2C-6b^2Bcd-16abcCd+3Ab^2d^2+9abBd^2+8a^2Cd^2-3b^2Cd^2)}{d \sqrt{c+d \tan(e+fx)}} + \frac{\frac{3}{2}(a^2B-b^2B+2ab(A-C))d}{\dots} \right) +$$

input

```

Integrate[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x]
    
```

output

```
(2*C*(a + b*Tan[e + f*x])^2)/(3*d*f*Sqrt[c + d*Tan[e + f*x]]) + (2*(((4*b
*c*C + 3*b*B*d + 4*a*C*d)*(a + b*Tan[e + f*x]))/(d*f*Sqrt[c + d*Tan[e + f
x]]) + ((-2*(8*b^2*c^2*C - 6*b^2*B*c*d - 16*a*b*c*C*d + 3*A*b^2*d^2 + 9*a*
b*B*d^2 + 8*a^2*C*d^2 - 3*b^2*C*d^2))/(d*Sqrt[c + d*Tan[e + f*x])) + (2*((
3*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2*((-I)*ArcTanh[Sqrt[c + d*Tan[e + f*
x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + (I*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sq
rt[c + I*d]])/Sqrt[c + I*d]))/2 + (((-3*c*(a^2*B - b^2*B + 2*a*b*(A - C))*
d^3)/2 - (3*(2*a*b*B - a^2*(A - C) + b^2*(A - C))*d^4)/2)*(-(Hypergeometri
c2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)]/((I*c + d)*Sqrt[c + d*T
an[e + f*x]])) + Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c +
I*d)]/((I*c - d)*Sqrt[c + d*Tan[e + f*x]])))/d)/d)/(2*d*f))/(3*d)
```

Rubi [A] (warning: unable to verify)

Time = 3.68 (sec) , antiderivative size = 346, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.319$, Rules used = {3042, 4128, 27, 3042, 4120, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$2 \int \frac{(a + b \tan(e + fx)) (b(4C^2 - 3Bdc + (3A + C)d^2) \tan^2(e + fx) + d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + Ad(ac + 4bd) + 2(2bc - \frac{ad}{2})(cC - Bd))}{2\sqrt{c + d \tan(e + fx)} d(c^2 + d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{\int \frac{(a+b \tan(e+fx))(b(4Cc^2-3Bdc+(3A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+4bd)+(4bc-ad)(cC-Bd))}{\sqrt{c+d \tan(e+fx)}} dx}{\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{\int \frac{(a+b \tan(e+fx))(b(4Cc^2-3Bdc+(3A+C)d^2) \tan(e+fx)^2+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+4bd)+(4bc-ad)(cC-Bd))}{\sqrt{c+d \tan(e+fx)}} dx}{\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4120

$$\frac{\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C) \sqrt{c+d \tan(e+fx)}}{3df} - \int \frac{2c(4Cc^2-3Bdc+(3A+C)d^2)b^2 - (6ad(2Cc^2-Bdc+(A+C)d^2) - b(8Cc^3-6Bdc^2+(3A+C)d^2))}{d(c^2+d^2)}}{\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C) \sqrt{c+d \tan(e+fx)}}{3df} - \int \frac{2c(4Cc^2-3Bdc+(3A+C)d^2)b^2 - (6ad(2Cc^2-Bdc+(A+C)d^2) - b(8Cc^3-6Bdc^2+(3A+C)d^2))}{d(c^2+d^2)}}{\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C) \sqrt{c+d \tan(e+fx)}}{3df} - \int \frac{2c(4Cc^2-3Bdc+(3A+C)d^2)b^2 - (6ad(2Cc^2-Bdc+(A+C)d^2) - b(8Cc^3-6Bdc^2+(3A+C)d^2))}{d(c^2+d^2)}}{\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4113

$$\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C)\sqrt{c+d \tan(e+fx)}}{3df} - \frac{\int \frac{-3((Ac-Cc+Bd)a^2-2b(Bc-(A-C)d)a-b^2(Ac-Cc+Bd))d^2-3((Bc-(A-C)d)a^2+2b(Bc-(A-C)d)a-b^2(Ac-Cc+Bd))}{\sqrt{c+d \tan(e+fx)}} dx}{d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C)\sqrt{c+d \tan(e+fx)}}{3df} - \frac{\int \frac{-3((Ac-Cc+Bd)a^2-2b(Bc-(A-C)d)a-b^2(Ac-Cc+Bd))d^2-3((Bc-(A-C)d)a^2+2b(Bc-(A-C)d)a-b^2(Ac-Cc+Bd))}{\sqrt{c+d \tan(e+fx)}} dx}{d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}}$$

↓ 4022

$$-\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C)\sqrt{c+d \tan(e+fx)}}{3df} - \frac{-\frac{3}{2}d^2(a+ib)^2(c-id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{3}{2}d^2(a-ib)^2(c+id)(A-ib-C)}{d(c^2+d^2)}$$

↓ 3042

$$-\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C)\sqrt{c+d \tan(e+fx)}}{3df} - \frac{-\frac{3}{2}d^2(a+ib)^2(c-id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{3}{2}d^2(a-ib)^2(c+id)(A-ib-C)}{d(c^2+d^2)}$$

↓ 4020

$$-\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} +$$

$$\frac{2b^2 \tan(e+fx)(d^2(3A+C)-3Bcd+4c^2C)\sqrt{c+d \tan(e+fx)}}{3df} - \frac{3id^2(a-ib)^2(c+id)(A-ib-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} +$$

↓ 25

$$\begin{aligned}
 & \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \\
 & \frac{2b^2 \tan(e + fx)(d^2(3A + C) - 3Bcd + 4c^2C)\sqrt{c + d \tan(e + fx)}}{3df} - \frac{3id^2(a - ib)^2(c + id)(A - iB - C) \int \frac{1}{(1 - i \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d(i \tan(e + fx))}{2f} - \frac{3id^2}{d} \\
 & \quad \downarrow 73 \\
 & \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \\
 & \frac{2b^2 \tan(e + fx)(d^2(3A + C) - 3Bcd + 4c^2C)\sqrt{c + d \tan(e + fx)}}{3df} - \frac{3d(a - ib)^2(c + id)(A - iB - C) \int \frac{1}{\frac{i \tan^2(e + fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c + d \tan(e + fx)}}{f} - \frac{3d(a + ib)^2(c - id)(A + iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f\sqrt{c - id}} - \frac{3d^2(a + ib)^2(c - id)(A + iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c + id}}\right)}{f\sqrt{c + id}} \\
 & \quad \downarrow 221 \\
 & \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \\
 & \frac{2b^2 \tan(e + fx)(d^2(3A + C) - 3Bcd + 4c^2C)\sqrt{c + d \tan(e + fx)}}{3df} - \frac{3d^2(a - ib)^2(c + id)(A - iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f\sqrt{c - id}} - \frac{3d^2(a + ib)^2(c - id)(A + iB - C) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c + id}}\right)}{f\sqrt{c + id}} \\
 & \hspace{20em} d(c^2 + d^2)
 \end{aligned}$$

input

```
Int[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x]
```

output

```
(-2*(c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^2)/(d*(c^2 + d^2)*f*sqrt[c + d*Tan[e + f*x]]) + ((2*b^2*(4*c^2*C - 3*B*c*d + (3*A + C)*d^2)*Tan[e + f*x]*sqrt[c + d*Tan[e + f*x]])/(3*d*f) - ((-3*(a - I*b)^2*(A - I*B - C)*(c + I*d)*d^2*ArcTan[Tan[e + f*x]/sqrt[c - I*d]])/(sqrt[c - I*d]*f) - (3*(a + I*b)^2*(A + I*B - C)*(c - I*d)*d^2*ArcTan[Tan[e + f*x]/sqrt[c + I*d]])/(sqrt[c + I*d]*f) - (2*b*(6*a*d*(2*c^2*C - B*c*d + (A + C)*d^2) - b*(8*c^3*C - 6*B*c^2*d + c*(3*A + 5*C)*d^2 - 3*B*d^3))*sqrt[c + d*Tan[e + f*x]])/(d*f))/(3*d))/(d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 9398 vs. $2(312) = 624$.

Time = 0.22 (sec) , antiderivative size = 9399, normalized size of antiderivative = 27.40

method	result	size
parts	Expression too large to display	9399
derivativedivides	Expression too large to display	36710
default	Expression too large to display	36710

input `int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((a+b*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((a + b*tan(e + f*x))**2*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(3/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e)}{(d \tan(fx + e))^{3/2}}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^2/(d*tan(f*x + e) + c)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 62.01 (sec) , antiderivative size = 54886, normalized size of antiderivative = 160.02

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `int(((a + b*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2),x)`

output

```
(2*(B*b^2*c^3 + B*a^2*c*d^2 - 2*B*a*b*c^2*d))/(d^2*f*(c^2 + d^2)*(c + d*tan(e + f*x))^(1/2)) - atan((((-(8*B^2*a^4*c^3*f^2 + 8*B^2*b^4*c^3*f^2 - 4*8*B^2*a^2*b^2*c^3*f^2 + 32*B^2*a*b^3*d^3*f^2 - 32*B^2*a^3*b*d^3*f^2 - 24*B^2*a^4*c*d^2*f^2 - 24*B^2*b^4*c*d^2*f^2 - 96*B^2*a*b^3*c^2*d*f^2 + 96*B^2*a^3*b*c^2*d*f^2 + 144*B^2*a^2*b^2*c*d^2*f^2)^2/4 - (16*c^6*f^4 + 16*d^6*f^4 + 48*c^2*d^4*f^4 + 48*c^4*d^2*f^4)*(B^4*a^8 + B^4*b^8 + 4*B^4*a^2*b^6 + 6*B^4*a^4*b^4 + 4*B^4*a^6*b^2)))^(1/2) - 4*B^2*a^4*c^3*f^2 - 4*B^2*b^4*c^3*f^2 + 24*B^2*a^2*b^2*c^3*f^2 - 16*B^2*a*b^3*d^3*f^2 + 16*B^2*a^3*b*d^3*f^2 + 12*B^2*a^4*c*d^2*f^2 + 12*B^2*b^4*c*d^2*f^2 + 48*B^2*a*b^3*c^2*d*f^2 - 48*B^2*a^3*b*c^2*d*f^2 - 72*B^2*a^2*b^2*c*d^2*f^2)/(16*(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*f^4)))^(1/2)*((c + d*tan(e + f*x))^(1/2))*(-(((8*B^2*a^4*c^3*f^2 + 8*B^2*b^4*c^3*f^2 - 48*B^2*a^2*b^2*c^3*f^2 + 32*B^2*a*b^3*d^3*f^2 - 32*B^2*a^3*b*d^3*f^2 - 24*B^2*a^4*c*d^2*f^2 - 24*B^2*b^4*c*d^2*f^2 - 96*B^2*a*b^3*c^2*d*f^2 + 96*B^2*a^3*b*c^2*d*f^2 + 144*B^2*a^2*b^2*c*d^2*f^2)^2/4 - (16*c^6*f^4 + 16*d^6*f^4 + 48*c^2*d^4*f^4 + 48*c^4*d^2*f^4)*(B^4*a^8 + B^4*b^8 + 4*B^4*a^2*b^6 + 6*B^4*a^4*b^4 + 4*B^4*a^6*b^2)))^(1/2) - 4*B^2*a^4*c^3*f^2 - 4*B^2*b^4*c^3*f^2 + 24*B^2*a^2*b^2*c^3*f^2 - 16*B^2*a*b^3*d^3*f^2 + 16*B^2*a^3*b*d^3*f^2 + 12*B^2*a^4*c*d^2*f^2 + 12*B^2*b^4*c*d^2*f^2 + 48*B^2*a*b^3*c^2*d*f^2 - 48*B^2*a^3*b*c^2*d*f^2 - 72*B^2*a^2*b^2*c*d^2*f^2)/(16*(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*f^4...
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**3 + int((sqrt(tan(e + f*x)*d + c)*tan(e
+ f*x)**4)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f
*x)*b**2*c*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e
+ f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*b**2*c**2*d*f + 2*int((sqrt
(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*
x)*c*d + c**2),x)*tan(e + f*x)*a*b*c*d**2*f + int((sqrt(tan(e + f*x)*d + c
)*tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*t
an(e + f*x)*b**3*d**2*f + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)
/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*b*c**2*d*f + int(
(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e
+ f*x)*c*d + c**2),x)*b**3*c*d*f - int((sqrt(tan(e + f*x)*d + c)*tan(e +
f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x
)*a**3*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*
x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a**2*c*d**2*f + 3*
int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*t
an(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a*b**2*d**2*f - int((sqrt(tan(e +
f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d +
c**2),x)*a**3*c*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(
e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a**2*c**2*d*f + 3*int((sq
rt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e...
```


3.118
$$\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1340
Mathematica [C] (verified)	1341
Rubi [A] (warning: unable to verify)	1341
Maple [B] (verified)	1345
Fricas [B] (verification not implemented)	1345
Sympy [F]	1346
Maxima [F(-1)]	1346
Giac [F(-1)]	1347
Mupad [B] (verification not implemented)	1347
Reduce [F]	1348

Optimal result

Integrand size = 45, antiderivative size = 201

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx =$$

$$-\frac{(ia + b)(A - iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{3/2} f}$$

$$+ \frac{(ia - b)(A + iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{3/2} f}$$

$$+ \frac{2(bc - ad)(c^2 C - Bcd + Ad^2)}{d^2 (c^2 + d^2) f \sqrt{c + d \tan(e + fx)}} + \frac{2bC \sqrt{c + d \tan(e + fx)}}{d^2 f}$$

output

```
-(I*a+b)*(A-I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(3/2)/f+(I*a-b)*(A+I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(3/2)/f+2*(-a*d+b*c)*(A*d^2-B*c*d+C*c^2)/d^2/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)+2*b*C*(c+d*tan(f*x+e))^(1/2)/d^2/f
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.76 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.44

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \frac{(Ab + aB - bC) \left(-\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} \right)}{(c + d \tan(e + fx))^{3/2}}$$

input

```
Integrate[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x]
```

output

```
((A*b + a*B - b*C)*((-I)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]/Sqrt[c - I*d] + (I*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]]/Sqrt[c + I*d]) - (2*(-2*b*c*C + b*B*d + 2*a*C*d))/(d*Sqrt[c + d*Tan[e + f*x]]) + ((A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*((-I)*c + d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)] + (I*c + d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c + I*d)]))/((c^2 + d^2)*Sqrt[c + d*Tan[e + f*x]]) + (2*C*(a + b*Tan[e + f*x]))/Sqrt[c + d*Tan[e + f*x]]/(d*f)
```

Rubi [A] (warning: unable to verify)

Time = 1.89 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.09, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.244$, Rules used = {3042, 4118, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{3/2}} dx$$

$$\begin{aligned}
& \downarrow 4118 \\
& \frac{\int \frac{bC(c^2+d^2) \tan^2(e+fx) + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{\sqrt{c+d \tan(e+fx)}} dx}{\frac{d(c^2+d^2)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}} + \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}}} \\
& \downarrow 3042 \\
& \frac{\int \frac{bC(c^2+d^2) \tan(e+fx)^2 + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{\sqrt{c+d \tan(e+fx)}} dx}{\frac{d(c^2+d^2)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}} + \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}}} \\
& \downarrow 4113 \\
& \frac{\int \frac{d(a(Ac-Cc+Bd) - b(Bc - (A-C)d)) + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2bC(c^2+d^2) \sqrt{c+d \tan(e+fx)}}{df}}{\frac{d(c^2+d^2)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}} + \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}}} \\
& \downarrow 3042 \\
& \frac{\int \frac{d(a(Ac-Cc+Bd) - b(Bc - (A-C)d)) + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2bC(c^2+d^2) \sqrt{c+d \tan(e+fx)}}{df}}{\frac{d(c^2+d^2)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}} + \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}}} \\
& \downarrow 4022 \\
& \frac{\frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}} + \frac{\frac{1}{2}d(a+ib)(c-id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}d(a-ib)(c+id)(A-iB-C) \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2bC}{d(c^2+d^2)}}{d(c^2+d^2)}}{d(c^2+d^2)} \\
& \downarrow 3042 \\
& \frac{\frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{d^2 f(c^2+d^2) \sqrt{c+d \tan(e+fx)}} + \frac{\frac{1}{2}d(a+ib)(c-id)(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}d(a-ib)(c+id)(A-iB-C) \int \frac{i \tan(e+fx)+1}{\sqrt{c+d \tan(e+fx)}} dx + \frac{2bC}{d(c^2+d^2)}}{d(c^2+d^2)}}{d(c^2+d^2)}
\end{aligned}$$

$$\begin{aligned} & \downarrow 4020 \\ & \frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{d^2 f (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} + \\ & \frac{id(a-ib)(c+id)(A-iB-C) \int -\frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} - \frac{id(a+ib)(c-id)(A+iB-C) \int -\frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}}}{2f} \\ & \hline & d(c^2 + d^2) \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{d^2 f (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} + \\ & \frac{id(a-ib)(c+id)(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} + \frac{id(a+ib)(c-id)(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} \\ & \hline & d(c^2 + d^2) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & \frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{d^2 f (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} + \\ & \frac{(a+ib)(c-id)(A+iB-C) \int \frac{1}{-\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{f} + \frac{(a-ib)(c+id)(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{f} \\ & \hline & d(c^2 + d^2) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & \frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{d^2 f (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} + \\ & \frac{d(a-ib)(c+id)(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} + \frac{d(a+ib)(c-id)(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}} + \frac{2bC(c^2+d^2)\sqrt{c+d \tan(e+fx)}}{df} \\ & \hline & d(c^2 + d^2) \end{aligned}$$

input `Int[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2), x]`

output `(2*(b*c - a*d)*(c^2*C - B*c*d + A*d^2))/(d^2*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]]) + (((a - I*b)*(A - I*B - C)*(c + I*d)*d*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f) + ((a + I*b)*(A + I*B - C)*(c - I*d)*d*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f) + (2*b*C*(c^2 + d^2)*Sqrt[c + d*Tan[e + f*x]])/(d*f))/(d*(c^2 + d^2))`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{\text{m}_.}) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{n}_.}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \text{ Subst}[\text{Int}[\text{x}^{\text{p} * (\text{m} + 1) - 1} * (\text{c} - \text{a} * (\text{d}/\text{b}) + \text{d} * (\text{x}^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b} * \text{x})^{(1/\text{p})}], \text{x}]] \text{ /}; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&\& \text{LtQ}[-1, \text{m}, 0] \&\& \text{LeQ}[-1, \text{n}, 0] \&\& \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \&\& \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a}) * \text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ /}; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&\& \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_., \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /}; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])], \text{x_Symbol}] \rightarrow \text{Simp}[\text{c} * (\text{d}/\text{f}) \text{ Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d}) * \text{x})^{\text{m}} / (\text{d}^2 + \text{c} * \text{x}), \text{x}], \text{x}, \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}]], \text{x}] \text{ /}; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_.} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I} * \text{d})/2 \text{ Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 - \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I} * \text{d})/2 \text{ Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 + \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] \text{ /}; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{!IntegerQ}[\text{m}]$
- rule 4113 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)])^{\text{m}_.} * ((\text{A}_.) + (\text{B}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)]) + (\text{C}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_.)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{C} * ((\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m} + 1} / (\text{b} * \text{f} * (\text{m} + 1))), \text{x}] + \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * \text{Simp}[\text{A} - \text{C} + \text{B} * \text{Tan}[\text{e} + \text{f} * \text{x}], \text{x}], \text{x}] \text{ /}; \text{FreeQ}[\{\text{a}, \text{b}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}, \text{m}\}, \text{x}] \&\& \text{NeQ}[\text{A} * \text{b}^2 - \text{a} * \text{b} * \text{B} + \text{a}^2 * \text{C}, 0] \&\& \text{!LeQ}[\text{m}, -1]$

rule 4118

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)^2), x_Symbol] :> Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 7395 vs. $2(177) = 354$.

Time = 0.23 (sec) , antiderivative size = 7396, normalized size of antiderivative = 36.80

method	result	size
parts	Expression too large to display	7396
derivativedivides	Expression too large to display	23472
default	Expression too large to display	23472

input

```

int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2
),x,method=_RETURNVERBOSE)

```

output

```

result too large to display

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31879 vs. $2(170) = 340$.

Time = 62.60 (sec) , antiderivative size = 31879, normalized size of antiderivative = 158.60

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(3/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output Timed out

Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3/2,x, algorithm="giac")`

output `Timed out`

Mupad [B] (verification not implemented)

Time = 36.04 (sec) , antiderivative size = 40542, normalized size of antiderivative = 201.70

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `int(((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^3/2,x)`

output

```
atan((((c + d*tan(e + f*x))^(1/2)*(16*A^2*a^2*d^10*f^3 - 16*B^2*a^2*d^10*f^3 + 16*C^2*a^2*d^10*f^3 + 32*A^2*a^2*c^2*d^8*f^3 - 32*A^2*a^2*c^6*d^4*f^3 - 16*A^2*a^2*c^8*d^2*f^3 - 32*B^2*a^2*c^2*d^8*f^3 + 32*B^2*a^2*c^6*d^4*f^3 + 16*B^2*a^2*c^8*d^2*f^3 + 32*C^2*a^2*c^2*d^8*f^3 - 32*C^2*a^2*c^6*d^4*f^3 - 16*C^2*a^2*c^8*d^2*f^3 - 32*A*C*a^2*d^10*f^3 - 64*A*B*a^2*c*d^9*f^3 + 64*B*C*a^2*c*d^9*f^3 - 192*A*B*a^2*c^3*d^7*f^3 - 192*A*B*a^2*c^5*d^5*f^3 - 64*A*B*a^2*c^7*d^3*f^3 - 64*A*C*a^2*c^2*d^8*f^3 + 64*A*C*a^2*c^6*d^4*f^3 + 32*A*C*a^2*c^8*d^2*f^3 + 192*B*C*a^2*c^3*d^7*f^3 + 192*B*C*a^2*c^5*d^5*f^3 + 64*B*C*a^2*c^7*d^3*f^3) - (((8*A^2*a^2*c^3*f^2 - 8*B^2*a^2*c^3*f^2 + 8*C^2*a^2*c^3*f^2 - 16*A*B*a^2*d^3*f^2 - 16*A*C*a^2*c^3*f^2 + 16*B*C*a^2*d^3*f^2 - 24*A^2*a^2*c*d^2*f^2 + 24*B^2*a^2*c*d^2*f^2 - 24*C^2*a^2*c*d^2*f^2 + 48*A*B*a^2*c^2*d*f^2 + 48*A*C*a^2*c^2*d*f^2 - 48*B*C*a^2*c^2*d*f^2)^2/4 - (16*c^6*f^4 + 16*d^6*f^4 + 48*c^2*d^4*f^4 + 48*c^4*d^2*f^4)*(A^4*a^4 + B^4*a^4 + C^4*a^4 - 4*A*C^3*a^4 - 4*A^3*C*a^4 + 2*A^2*B^2*a^4 + 6*A^2*C^2*a^4 + 2*B^2*C^2*a^4 - 4*A*B^2*C*a^4))^(1/2) - 4*A^2*a^2*c^3*f^2 + 4*B^2*a^2*c^3*f^2 - 4*C^2*a^2*c^3*f^2 + 8*A*B*a^2*d^3*f^2 + 8*A*C*a^2*c^3*f^2 - 8*B*C*a^2*d^3*f^2 + 12*A^2*a^2*c*d^2*f^2 - 12*B^2*a^2*c*d^2*f^2 + 12*C^2*a^2*c*d^2*f^2 - 24*A*B*a^2*c^2*d*f^2 - 24*A*C*a^2*c*d^2*f^2 + 24*B*C*a^2*c^2*d*f^2)/(16*(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*f^4)))^(1/2))*((c + d*tan(e + f*x))^(1/2)*(((8*A^2*a^2*c^3*f^2 - 8*B^2*a^2*c^3*f^2 + ...
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```
int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2 + int((sqrt(tan(e + f*x)*d + c)*tan(e
+ f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f
*x)*b*c*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f
*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*b*c**2*d*f - int((sqrt(tan(e +
f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d +
c**2),x)*tan(e + f*x)*a**2*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e +
f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*
x)*a*c*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*
x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*b**2*d**2*f - int(
(sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e
+ f*x)*c*d + c**2),x)*a**2*c*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e +
f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*c**2*d*f
+ int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2
*tan(e + f*x)*c*d + c**2),x)*b**2*c*d*f + 2*int((sqrt(tan(e + f*x)*d + c)*
tan(e + f*x))/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e
+ f*x)*a*b*d**2*f + 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e +
f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*b*c*d*f)/(d*f*(tan(e + f*x
)*d + c))
```

3.119 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^{3/2}} dx$

Optimal result	1350
Mathematica [C] (verified)	1351
Rubi [A] (warning: unable to verify)	1351
Maple [B] (verified)	1354
Fricas [B] (verification not implemented)	1355
Sympy [F]	1355
Maxima [F(-1)]	1355
Giac [F(-2)]	1356
Mupad [B] (verification not implemented)	1356
Reduce [F]	1357

Optimal result

Integrand size = 35, antiderivative size = 157

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{3/2} f} - \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{3/2} f} - \frac{2(c^2 C - Bcd + Ad^2)}{d(c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

output

```
-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(3/2)/f
-(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(3/2)/f
-2*(A*d^2-B*c*d+C*c^2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.70 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.39

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \frac{-iB \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} \right)}{d}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^(3/2), x]
```

output

```
((-I)*B*(ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]/Sqrt[c - I*d] - ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]]/Sqrt[c + I*d]) - (2*C)/Sqrt[c + d*Tan[e + f*x]] + ((B*c + (-A + C)*d)*((( -I)*c + d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)] + (I*c + d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c + I*d)]))/((c^2 + d^2)*Sqrt[c + d*Tan[e + f*x]]))/(d*f)
```

Rubi [A] (warning: unable to verify)

Time = 1.19 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 4111, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4111

$$\begin{aligned}
 & \frac{\int \frac{Ac-Cc+Bd+(Bc-(A-C)d)\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} - \frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Ac-Cc+Bd+(Bc-(A-C)d)\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} - \frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} \\
 & \quad \downarrow \text{4022} \\
 & \frac{\frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} + \frac{\frac{1}{2}(c-id)(A+iB-C) \int \frac{1-i\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx + \frac{1}{2}(c+id)(A-iB-C) \int \frac{i\tan(e+fx)+1}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2}}{c^2+d^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} + \frac{\frac{1}{2}(c-id)(A+iB-C) \int \frac{1-i\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx + \frac{1}{2}(c+id)(A-iB-C) \int \frac{i\tan(e+fx)+1}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2}}{c^2+d^2} \\
 & \quad \downarrow \text{4020} \\
 & \frac{\frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} + \frac{i(c+id)(A-iB-C) \int -\frac{1}{(1-i\tan(e+fx))\sqrt{c+d\tan(e+fx)}} d(i\tan(e+fx))}{2f} - \frac{i(c-id)(A+iB-C) \int -\frac{1}{(i\tan(e+fx)+1)\sqrt{c+d\tan(e+fx)}} d(-i\tan(e+fx))}{2f}}{c^2+d^2}}{c^2+d^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{\frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} + \frac{i(c-id)(A+iB-C) \int \frac{1}{(i\tan(e+fx)+1)\sqrt{c+d\tan(e+fx)}} d(-i\tan(e+fx))}{2f} - \frac{i(c+id)(A-iB-C) \int \frac{1}{(1-i\tan(e+fx))\sqrt{c+d\tan(e+fx)}} d(i\tan(e+fx))}{2f}}{c^2+d^2}}{c^2+d^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{\frac{2(Ad^2-Bcd+c^2C)}{df(c^2+d^2)\sqrt{c+d\tan(e+fx)}} + \frac{(c-id)(A+iB-C) \int \frac{1}{-i\tan^2(e+fx) - \frac{ic}{d} + 1} d\sqrt{c+d\tan(e+fx)}}{df} + \frac{(c+id)(A-iB-C) \int \frac{1}{i\tan^2(e+fx) + \frac{ic}{d} + 1} d\sqrt{c+d\tan(e+fx)}}{df}}{c^2+d^2}}{c^2+d^2} \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

$$\frac{2(Ad^2 - Bcd + c^2C)}{df(c^2 + d^2)\sqrt{c + d\tan(e + fx)}} + \frac{(c+id)(A-iB-C)\arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} + \frac{(c-id)(A+iB-C)\arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}}$$

$$c^2 + d^2$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^(3/2),x]`

output `((A - I*B - C)*(c + I*d)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]]/(Sqrt[c - I*d]*f) + ((A + I*B - C)*(c - I*d)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]]/(Sqrt[c + I*d]*f))/(c^2 + d^2) - (2*(c^2*C - B*c*d + A*d^2))/(d*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5612 vs. $2(136) = 272$.

Time = 0.14 (sec) , antiderivative size = 5613, normalized size of antiderivative = 35.75

method	result	size
parts	Expression too large to display	5613
derivativedivides	Expression too large to display	11427
default	Expression too large to display	11427

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x,method=_RETUR
NVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7982 vs. $2(129) = 258$.

Time = 1.96 (sec) , antiderivative size = 7982, normalized size of antiderivative = 50.84

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(3/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{1, [3,9,3]}+%%{4, [3,7,3]}+%%{6, [3,5,3]}+%%{
4, [3,3,3]}
```

Mupad [B] (verification not implemented)

Time = 15.95 (sec) , antiderivative size = 8588, normalized size of antiderivative = 54.70

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/(c + d*tan(e + f*x))^(3/2),x)
```

output

```
(log(((((((96*C^4*c^2*d^4*f^4 - 16*C^4*d^6*f^4 - 144*C^4*c^4*d^2*f^4)^(1/2)
) - 4*C^2*c^3*f^2 + 12*C^2*c*d^2*f^2)/(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 +
3*c^4*d^2*f^4))^(1/2)*(64*C*c*d^11*f^4 - ((c + d*tan(e + f*x))^(1/2))*((9
6*C^4*c^2*d^4*f^4 - 16*C^4*d^6*f^4 - 144*C^4*c^4*d^2*f^4)^(1/2) - 4*C^2*c^
3*f^2 + 12*C^2*c*d^2*f^2)/(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*f
^4))^(1/2)*(64*c*d^12*f^5 + 320*c^3*d^10*f^5 + 640*c^5*d^8*f^5 + 640*c^7*d
^6*f^5 + 320*c^9*d^4*f^5 + 64*c^11*d^2*f^5))/4 + 256*C*c^3*d^9*f^4 + 384*C
*c^5*d^7*f^4 + 256*C*c^7*d^5*f^4 + 64*C*c^9*d^3*f^4))/4 + (c + d*tan(e + f
*x))^(1/2)*(16*C^2*d^10*f^3 + 32*C^2*c^2*d^8*f^3 - 32*C^2*c^6*d^4*f^3 - 16
*C^2*c^8*d^2*f^3))*(((96*C^4*c^2*d^4*f^4 - 16*C^4*d^6*f^4 - 144*C^4*c^4*d^
2*f^4)^(1/2) - 4*C^2*c^3*f^2 + 12*C^2*c*d^2*f^2)/(c^6*f^4 + d^6*f^4 + 3*c^
2*d^4*f^4 + 3*c^4*d^2*f^4))^(1/2))/4 - 8*C^3*d^9*f^2 - 24*C^3*c^2*d^7*f^2
- 24*C^3*c^4*d^5*f^2 - 8*C^3*c^6*d^3*f^2))*(((96*C^4*c^2*d^4*f^4 - 16*C^4*d
^6*f^4 - 144*C^4*c^4*d^2*f^4)^(1/2) - 4*C^2*c^3*f^2 + 12*C^2*c*d^2*f^2)/(c
^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*f^4))^(1/2))/4 + (log(((((-(
96*C^4*c^2*d^4*f^4 - 16*C^4*d^6*f^4 - 144*C^4*c^4*d^2*f^4)^(1/2) + 4*C^2*c
^3*f^2 - 12*C^2*c*d^2*f^2)/(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*
f^4))^(1/2)*(64*C*c*d^11*f^4 - ((c + d*tan(e + f*x))^(1/2))*(-(96*C^4*c^2*
d^4*f^4 - 16*C^4*d^6*f^4 - 144*C^4*c^4*d^2*f^4)^(1/2) + 4*C^2*c^3*f^2 - 12
*C^2*c*d^2*f^2)/(c^6*f^4 + d^6*f^4 + 3*c^2*d^4*f^4 + 3*c^4*d^2*f^4))^(1...
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{3/2}} dx = \frac{-2\sqrt{d \tan(fx + e) + c} a - \left(\int \frac{\sqrt{d \tan(fx + e) + c} \tan(fx + e)^2}{\tan(fx + e)^2 d^2 + 2 \tan(fx + e) cd + c^2} dx \right)}{1}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*c*d**2*f - int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*c*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*c**2*d*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*b*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*b*c*d*f)/(d*f*(tan(e + f*x)*d + c))
```

3.120 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2}} dx$

Optimal result	1359
Mathematica [A] (verified)	1360
Rubi [A] (warning: unable to verify)	1360
Maple [B] (verified)	1365
Fricas [F(-1)]	1366
Sympy [F]	1366
Maxima [F(-2)]	1367
Giac [F(-2)]	1367
Mupad [B] (verification not implemented)	1368
Reduce [F]	1369

Optimal result

Integrand size = 47, antiderivative size = 262

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \frac{(A - iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(ia + b)(c - id)^{3/2} f}$$

$$+ \frac{(iA - B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a + ib)(c + id)^{3/2} f}$$

$$- \frac{2\sqrt{b}(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{(a^2 + b^2)(bc - ad)^{3/2} f}$$

$$+ \frac{2(c^2C - Bcd + Ad^2)}{(bc - ad)(c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

output

```
(A-I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(I*a+b)/(c-I*d)^(3/2)/f+(I*A-B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(a+I*b)/(c+I*d)^(3/2)/f-2*b^(1/2)*(A*b^2-a*(B*b-C*a))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/(a^2+b^2)/(-a*d+b*c)^(3/2)/f+2*(A*d^2-B*c*d+C*c^2)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 3.36 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.13

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \frac{i \left(\frac{(a+ib)(A-iB-C)(c+id)(-bc+ad) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{\sqrt{c-id}} + \frac{(a-ib)(A-iB-C)(c-id)(bc-ad) \operatorname{arctanh}\left(\frac{\sqrt{c-d \tan(e+fx)}}{\sqrt{c+id}}\right)}{\sqrt{c+id}} \right)}{a^2+b^2}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

```
((-I)*(((a + I*b)*(A - I*B - C)*(c + I*d)*(-(b*c) + a*d)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]])/Sqrt[c - I*d] + ((a - I*b)*(A + I*B - C)*(c - I*d)*(b*c - a*d)*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c + I*d]])/Sqrt[c + I*d]))/(a^2 + b^2) + (2*Sqrt[b]*(A*b^2 + a*(-(b*B) + a*C))*(c^2 + d^2)*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/((a^2 + b^2)*Sqrt[b*c - a*d]) - (2*(c^2*C - B*c*d + A*d^2))/Sqrt[c + d*Tan[e + f*x]]/((- (b*c) + a*d)*(c^2 + d^2)*f)
```

Rubi [A] (warning: unable to verify)

Time = 3.59 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.18, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.340$, Rules used = {3042, 4132, 27, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4132

$$\begin{aligned}
 & \frac{2 \int -\frac{-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAcd - ad(cC - Bd) - Ab(c^2 + d^2)}{2(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx}{\frac{(c^2 + d^2)(bc - ad)}{2(Ad^2 - Bcd + c^2C)}} + \\
 & \frac{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}}{\downarrow 27} \\
 & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}} - \\
 & \frac{\int -\frac{-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAcd - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx}{(c^2 + d^2)(bc - ad)} \\
 & \frac{\downarrow 3042}{2(Ad^2 - Bcd + c^2C)} - \\
 & \frac{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}}{\int -\frac{-b(Cc^2 - Bdc + Ad^2) \tan(e + fx)^2 - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAcd - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx} \\
 & \frac{(c^2 + d^2)(bc - ad)}{\downarrow 4136} \\
 & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}} - \\
 & \frac{\int -\frac{(bc - ad)(bBc - b(A - C)d + a(Ac - Cc + Bd)) + (bc - ad)(aBc + bCc - bBd + aCd - A(bc + ad)) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan^2(e + fx)}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2} \\
 & \frac{(c^2 + d^2)(bc - ad)}{\downarrow 25} \\
 & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}} - \\
 & \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan^2(e + fx) + 1}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2} - \frac{\int \frac{(bc - ad)(bBc - b(A - C)d + a(Ac - Cc + Bd)) + (bc - ad)(aBc + bCc - bBd + aCd - A(bBc - b(A - C)d + a(Ac - Cc + Bd))) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2} \\
 & \frac{(c^2 + d^2)(bc - ad)}{\downarrow 3042} \\
 & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}} - \\
 & \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2} - \frac{\int \frac{(bc - ad)(bBc - b(A - C)d + a(Ac - Cc + Bd)) + (bc - ad)(aBc + bCc - bBd + aCd - A(bBc - b(A - C)d + a(Ac - Cc + Bd))) \tan(e + fx)}{\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2} \\
 & \frac{(c^2 + d^2)(bc - ad)}{\downarrow 4022}
 \end{aligned}$$

$$\frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{\frac{1}{2}(a-ib)(c-id)(A+iB-C)(bc-ad) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(a+ib)(c+id)}{a^2 + b^2}$$

$$(c^2 + d^2)(bc - ad)$$

↓ 3042

$$\frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{\frac{1}{2}(a-ib)(c-id)(A+iB-C)(bc-ad) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(a+ib)(c+id)}{a^2 + b^2}$$

$$(c^2 + d^2)(bc - ad)$$

↓ 4020

$$\frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{i(a+ib)(c+id)(A-iB-C)(bc-ad) \int -\frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f}$$

$$(c^2 + d^2)(bc - ad)$$

↓ 25

$$\frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{i(a-ib)(c-id)(A+iB-C)(bc-ad) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f}$$

$$(c^2 + d^2)(bc - ad)$$

↓ 73

$$\frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{(a-ib)(c-id)(A+iB-C)(bc-ad) \int -\frac{1}{d} \frac{d\sqrt{c+d \tan(e+fx)}}{i \tan^2(e+fx) - \frac{ic}{d} + 1}}{df}$$

$$(c^2 + d^2)(bc - ad)$$

↓ 221

$$\frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{\tan(e+fx)^2 + 1}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx}{a^2 + b^2} - \frac{(a+ib)(c+id)(A-iB-C)(bc-ad) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right) + (a-ib)(c-id)(A+iB-C)}{f\sqrt{c-id} a^2 + b^2}$$

$$(c^2 + d^2)(bc - ad)$$

$$\begin{aligned} & \downarrow 4117 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \\ & \frac{b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} d \tan(e + fx)}{f(a^2 + b^2)} - \frac{(a + ib)(c + id)(A - iB - C)(bc - ad) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f\sqrt{c - id}} + \frac{(a - ib)(c - id)}{a^2 + b^2} \\ & \hline & (c^2 + d^2)(bc - ad) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \\ & \frac{2b(c^2 + d^2)(Ab^2 - a(bB - aC)) \int \frac{1}{a + \frac{b(c + d \tan(e + fx))}{d} - \frac{bc}{d}} d \sqrt{c + d \tan(e + fx)}}{df(a^2 + b^2)} - \frac{(a + ib)(c + id)(A - iB - C)(bc - ad) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f\sqrt{c - id}} + \frac{(a - ib)(c - id)}{a^2 + b^2} \\ & \hline & (c^2 + d^2)(bc - ad) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \\ & \frac{2\sqrt{b}(c^2 + d^2)(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c + d \tan(e + fx)}}{\sqrt{bc - ad}}\right)}{f(a^2 + b^2)\sqrt{bc - ad}} - \frac{(a + ib)(c + id)(A - iB - C)(bc - ad) \arctan\left(\frac{\tan(e + fx)}{\sqrt{c - id}}\right)}{f\sqrt{c - id}} + \frac{(a - ib)(c - id)(A + iB - C)(bc - ad)}{f\sqrt{c - id}} \\ & \hline & (c^2 + d^2)(bc - ad) \end{aligned}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

```
-(((a + I*b)*(A - I*B - C)*(c + I*d)*(b*c - a*d)*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f) + ((a - I*b)*(A + I*B - C)*(c - I*d)*(b*c - a*d)*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f)/(a^2 + b^2) + (2*Sqrt[b]*(A*b^2 - a*(b*B - a*C))*(c^2 + d^2)*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/(a^2 + b^2)*Sqrt[b*c - a*d]*f)/((b*c - a*d)*(c^2 + d^2)) + (2*(c^2*C - B*c*d + A*d^2))/((b*c - a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])
```


Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(x_)^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*(x_)^{\text{n}_}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[x^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(x^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*x)^{1/\text{p}}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{LtQ}[-1, \text{m}, 0] \ \&\& \ \text{LeQ}[-1, \text{n}, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \ \&\& \ \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d})*x)^{\text{m}}/(\text{d}^2 + \text{c}*x), \text{x}], \text{x}, \text{d}*\text{Tan}[\text{e} + \text{f}*x]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I}*d)/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m}}*(1 - \text{I}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I}*d)/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m}}*(1 + \text{I}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!IntegerQ}[\text{m}]$

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] +
Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :=
Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] +
Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 26342 vs. $2(229) = 458$.

Time = 0.22 (sec) , antiderivative size = 26343, normalized size of antiderivative = 100.55

method	result	size
derivativedivides	Expression too large to display	26343
default	Expression too large to display	26343

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))**(3/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

Mupad [B] (verification not implemented)

Time = 92.29 (sec) , antiderivative size = 630337, normalized size of antiderivative = 2405.87

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^(3/2)),x)
```

output

```
(atan((((c + d*tan(e + f*x))^(1/2)*(32*A^4*a^9*b^5*d^31*f^5 - 64*A^4*a^7*b^7*d^31*f^5 + 64*A^4*b^14*c^7*d^24*f^5 + 352*A^4*b^14*c^9*d^22*f^5 + 672*A^4*b^14*c^11*d^20*f^5 + 224*A^4*b^14*c^13*d^18*f^5 - 1120*A^4*b^14*c^15*d^16*f^5 - 2016*A^4*b^14*c^17*d^14*f^5 - 1568*A^4*b^14*c^19*d^12*f^5 - 608*A^4*b^14*c^21*d^10*f^5 - 96*A^4*b^14*c^23*d^8*f^5 - 448*A^4*a*b^13*c^6*d^25*f^5 - 2400*A^4*a*b^13*c^8*d^23*f^5 - 4256*A^4*a*b^13*c^10*d^21*f^5 - 224*A^4*a*b^13*c^12*d^19*f^5 + 10080*A^4*a*b^13*c^14*d^17*f^5 + 16352*A^4*a*b^13*c^16*d^15*f^5 + 12320*A^4*a*b^13*c^18*d^13*f^5 + 4704*A^4*a*b^13*c^20*d^11*f^5 + 736*A^4*a*b^13*c^22*d^9*f^5 + 448*A^4*a^6*b^8*c*d^30*f^5 - 288*A^4*a^8*b^6*c*d^30*f^5 + 1344*A^4*a^2*b^12*c^5*d^26*f^5 + 6912*A^4*a^2*b^12*c^7*d^24*f^5 + 10752*A^4*a^2*b^12*c^9*d^22*f^5 - 5376*A^4*a^2*b^12*c^11*d^20*f^5 - 40320*A^4*a^2*b^12*c^13*d^18*f^5 - 59136*A^4*a^2*b^12*c^15*d^16*f^5 - 43008*A^4*a^2*b^12*c^17*d^14*f^5 - 16128*A^4*a^2*b^12*c^19*d^12*f^5 - 2496*A^4*a^2*b^12*c^21*d^10*f^5 - 2240*A^4*a^3*b^11*c^4*d^27*f^5 - 10752*A^4*a^3*b^11*c^6*d^25*f^5 - 12544*A^4*a^3*b^11*c^8*d^23*f^5 + 25088*A^4*a^3*b^11*c^10*d^21*f^5 + 94080*A^4*a^3*b^11*c^12*d^19*f^5 + 125440*A^4*a^3*b^11*c^14*d^17*f^5 + 87808*A^4*a^3*b^11*c^16*d^15*f^5 + 32256*A^4*a^3*b^11*c^18*d^13*f^5 + 4928*A^4*a^3*b^11*c^20*d^11*f^5 + 2240*A^4*a^4*b^10*c^3*d^28*f^5 + 9408*A^4*a^4*b^10*c^5*d^26*f^5 + 3136*A^4*a^4*b^10*c^7*d^24*f^5 - 53312*A^4*a^4*b^10*c^9*d^22*f^5 - 141120*A^4*a^4*b^10*c^11*d^20*f^5 ...
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 4*sqrt(tan(e + f*x)*d + c)*a + 2*sqrt(tan(e + f*x)*d + c)*c - int(sqrt
(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*
tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**
2),x)*tan(e + f*x)*a**2*d**2*f + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)
)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e +
f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*tan(e + f*x)*a*c*d**2*f - i
nt(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d*
*2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2
+ a*c**2),x)*a**2*c*d*f + int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**3*b*
d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a
*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*a*c**2*d*f - 2*int((sqrt(tan(e + f
*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d*
*2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2
+ a*c**2),x)*tan(e + f*x)*a*b*d**2*f + int((sqrt(tan(e + f*x)*d + c)*tan(e
+ f*x)**3)/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f
*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*tan
(e + f*x)*b*c*d**2*f - 2*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(t
an(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d +
2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*a*b*c*d*f + int((
sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*b*d**2 + tan...
```

3.121
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1370
Mathematica [B] (verified)	1371
Rubi [A] (warning: unable to verify)	1372
Maple [B] (verified)	1378
Fricas [F(-1)]	1379
Sympy [F]	1379
Maxima [F(-2)]	1380
Giac [F(-2)]	1380
Mupad [F(-1)]	1381
Reduce [F]	1381

Optimal result

Integrand size = 47, antiderivative size = 447

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right) - (B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a - ib)^2(c - id)^{3/2} f - (a + ib)^2(c + id)^{3/2} f}$$

$$- \frac{\sqrt{b}(5a^3bBd - 3a^4Cd + b^4(2Bc - 3Ad) + ab^3(4Ac - 4cC + Bd) - a^2b^2(2Bc + (7A - C)d)) \operatorname{arctanh}\left(\frac{\sqrt{b}}{\sqrt{c+d \tan(e+fx)}}\right)}{(a^2 + b^2)^2 (bc - ad)^{5/2} f}$$

$$- \frac{d(2b^2c(cC - Bd) - abB(c^2 + d^2) + a^2(3c^2C - 2Bcd + Cd^2) + A(2a^2d^2 + b^2(c^2 + 3d^2)))}{(a^2 + b^2) (bc - ad)^2 (c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

$$- \frac{Ab^2 - a(bB - aC)}{(a^2 + b^2) (bc - ad) f (a + b \tan(e + fx)) \sqrt{c + d \tan(e + fx)}}$$

output

```

-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(a-I*b)^(2/(c-I*d)^(3/2)/f-(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(a+I*b)^(2/(c+I*d)^(3/2)/f-b^(1/2)*(5*a^3*b*B*d-3*a^4*C*d+b^4*(-3*A*d+2*B*c)+a*b^3*(4*A*c+B*d-4*C*c)-a^2*b^2*(2*B*c+(7*A-C)*d))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/(a^2+b^2)^(2/(-a*d+b*c)^(5/2)/f-d*(2*b^2*c*(-B*d+C*c)-a*b*B*(c^2+d^2)+a^2*(-2*B*c*d+3*C*c^2+C*d^2)+A*(2*a^2*d^2+b^2*(c^2+3*d^2)))/(a^2+b^2)/(-a*d+b*c)^(2/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)-(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(1/2)

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2078 vs. $2(447) = 894$.

Time = 6.27 (sec) , antiderivative size = 2078, normalized size of antiderivative = 4.65

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \text{Result too large to show}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(3/2)),x]

```


output

```

-((A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])*
Sqrt[c + d*Tan[e + f*x]])) - ((-2*((I*Sqrt[c - I*d]*((b*(-(b*c) + a*d))*((-3*(A*b^2 - a*(b*B - a*C))*d^2)/2 - c*(A*b - a*B - b*C)*(b*c - a*d) + (d*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2))/2 + a*(-1/2*(a*d*((-3*c*(A*b^2 - a*(b*B - a*C))*d)/2 + (A*b - a*B - b*C)*d*(b*c - a*d))) + (((b*d^2)/2 - (c*(-(b*c) + a*d))/2)*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2 - (b*(-(c*((-3*c*(A*b^2 - a*(b*B - a*C))*d)/2 + (A*b - a*B - b*C)*d*(b*c - a*d))) + (d^2*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2))/2 - I*((a*(-(b*c) + a*d)*((-3*(A*b^2 - a*(b*B - a*C))*d^2)/2 - c*(A*b - a*B - b*C)*(b*c - a*d) + (d*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2))/2 - b*(-1/2*(a*d*((-3*c*(A*b^2 - a*(b*B - a*C))*d)/2 + (A*b - a*B - b*C)*d*(b*c - a*d))) + (((b*d^2)/2 - (c*(-(b*c) + a*d))/2)*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2 - (b*(-(c*((-3*c*(A*b^2 - a*(b*B - a*C))*d)/2 + (A*b - a*B - b*C)*d*(b*c - a*d))) + (d^2*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2))/2)))*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]/((-c + I*d)*f) - (I*Sqrt[c + I*d]*((b*(-(b*c) + a*d)*((-3*(A*b^2 - a*(b*B - a*C))*d^2)/2 - c*(A*b - a*B - b*C)*(b*c - a*d) + (d*(3*A*b^2*d - 2*a*A*(b*c - a*d) - (b*B - a*C)*(2*b*c + a*d)))/2))/2 + a*(-1/2*(a*d*((-3*c*(A*b^2 - a*(b*B - a*C))*d)/2 + (A*b - a*B - b*C)*d*(b*c - a*d))) + ...

```

Rubi [A] (warning: unable to verify)

Time = 6.73 (sec) , antiderivative size = 515, normalized size of antiderivative = 1.15, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.404$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx$$

↓ 4132

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan^2(e+fx)-2aA(bc-ad)-(bB-aC)(2bc+ad)+2(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{2(a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}$$

↓ 27

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan^2(e+fx)-2aA(bc-ad)-(bB-aC)(2bc+ad)+2(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{2(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}$$

↓ 3042

$$\int \frac{3Adb^2+3(Ab^2-a(bB-aC))d \tan(e+fx)^2-2aA(bc-ad)-(bB-aC)(2bc+ad)+2(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{2(a^2+b^2)(bc-ad)}{Ab^2-a(bB-aC)}$$

$$f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}$$

↓ 4132

$$2 \int -\frac{2d^2(Ac-Cc+Bd)a^3-b(4A-C)d(c^2+d^2)a^2-b^2(2Cc^3+Bdc^2+4Cd^2c-Bd^3-2A(c^3+2d^2c))a-bd(2Ad^2a^2+(3Cc^2-2Bdc+Cd^2)a^2-bB(c^2+d^2)a+2b^2c^2)}{2(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}}$$

↓ 27

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+3c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+3d^2)+2b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} \int \frac{2d^2(Ac-Cc+Bd)a^3-b(4A-C)d(c^2+d^2)a^2-b^2(2Cc^3+2b^2c^2)}{(c^2+d^2)(bc-ad)}$$

$$\frac{Ab^2-a(bB-aC)}{f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}}$$

↓ 3042

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+3c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+3d^2)+2b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \int \frac{2d^2(Ac-Cc+Bd)a^3-b(4A-C)d(c^2+d^2)a^2-b^2(2Cc^3+}$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}}$$

↓ 4136

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+3c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+3d^2)+2b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \int \frac{2((bc-ad)^2((Ac-Cc+Bd)a^2+2b(Bc-(A-C)d)a-b^2(Ac$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+3c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+3d^2)+2b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \int \frac{(bc-ad)^2((Ac-Cc+Bd)a^2+2b(Bc-(A-C)d)a-b^2(Ac$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+3c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+3d^2)+2b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \int \frac{(bc-ad)^2((Ac-Cc+Bd)a^2+2b(Bc-(A-C)d)a-b^2(Ac$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}}$$

↓ 4022

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}}$$

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+3c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+3d^2)+2b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \int \frac{b(c^2+d^2)(-3a^4Cd+5a^3bBd-a^2b^2(d(7A-C)+2Bc)+ab^3(4$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}$$

↓ 4020

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}$$

↓ 25

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}$$

↓ 73

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}$$

↓ 221

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - C))}$$

↓ 4117

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - 3B))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}}$$

73

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{2b(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - 3B))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}}$$

221

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 3c^2C + Cd^2) - abB(c^2 + d^2) + Ab^2(c^2 + 3d^2) + 2b^2c(cC - Bd))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{2\sqrt{b}(c^2 + d^2)(-3a^4Cd + 5a^3bBd - a^2b^2(d(7A - C) + 2Bc) + ab^3(4A - 3B))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}}$$

```
input Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(3/2)),x]
```

```
output -((A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]) - (-(((2*((a + I*b)^2*(A - I*B - C)*(c + I*d)*(b*c - a*d)^2*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f) + ((a - I*b)^2*(A + I*B - C)*(c - I*d)*(b*c - a*d)^2*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f)))/(a^2 + b^2) - (2*Sqrt[b]*(c^2 + d^2)*(5*a^3*b*B*d - 3*a^4*C*d + b^4*(2*B*c - 3*A*d) + a*b^3*(4*A*c - 4*c*C + B*d) - a^2*b^2*(2*B*c + (7*A - C)*d))*ArcTanh[(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])/Sqrt[b*c - a*d]])/((a^2 + b^2)*Sqrt[b*c - a*d]*f)/((b*c - a*d)*(c^2 + d^2)) + (2*d*(2*a^2*A*d^2 + 2*b^2*c*(c*C - B*d) - a*b*B*(c^2 + d^2) + A*b^2*(c^2 + 3*d^2) + a^2*(3*c^2*C - 2*B*c*d + C*d^2)))/((b*c - a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]]))/(2*(a^2 + b^2)*(b*c - a*d))
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{\text{n}_}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \quad \text{Subst}[\text{Int}[\text{x}^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(\text{x}^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*\text{x})^{1/\text{p}}, \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{LtQ}[-1, \text{m}, 0] \ \&\& \ \text{LeQ}[-1, \text{n}, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \ \&\& \ \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(\text{x}_.)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)]]), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d})*\text{x})^{\text{m}}/(\text{d}^2 + \text{c}*\text{x}), \text{x}], \text{x}, \text{d}*\text{Tan}[\text{e} + \text{f}*\text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)])^{\text{m}_})*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)]]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 - \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I}*\text{d})/2 \quad \text{Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(1 + \text{I}*\text{Tan}[\text{e} + \text{f}*\text{x}]), \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*\text{c} - \text{a}*\text{d}, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!IntegerQ}[\text{m}]$

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] +
Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :=
Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] +
Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 40618 vs. $2(412) = 824$.

Time = 0.32 (sec) , antiderivative size = 40619, normalized size of antiderivative = 90.87

method	result	size
derivativedivides	Expression too large to display	40619
default	Expression too large to display	40619

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**2/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**2*(c + d*tan(e + f*x))**(3/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [3, 19, 8]}%%}+%%{8, [3, 17, 8]}%%}+%%{28, [3, 15, 8]}%%}+%%{56, [3`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^2*(c + d
*tan(e + f*x))^(3/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(3
/2),x)
```

output

```
( - 8*sqrt(tan(e + f*x)*d + c)*a*d - 4*sqrt(tan(e + f*x)*d + c)*b*c + 6*sqrt(tan(e + f*x)*d + c)*c*d - int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*a*b**2*d**3 + 2*tan(e + f*x)**4*b**3*c*d**2 + 2*tan(e + f*x)**3*a**2*b*d**3 + 6*tan(e + f*x)**3*a*b**2*c*d**2 + 4*tan(e + f*x)**3*b**3*c**2*d + tan(e + f*x)**2*a**3*d**3 + 6*tan(e + f*x)**2*a**2*b*c*d**2 + 9*tan(e + f*x)**2*a*b**2*c**2*d + 2*tan(e + f*x)**2*b**3*c**3 + 2*tan(e + f*x)*a**3*c*d**2 + 6*tan(e + f*x)*a**2*b*c**2*d + 4*tan(e + f*x)*a*b**2*c**3 + a**3*c**2*d + 2*a**2*b*c**3),x)*tan(e + f*x)**2*a**3*b*d**4*f - 6*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*a*b**2*d**3 + 2*tan(e + f*x)**4*b**3*c*d**2 + 2*tan(e + f*x)**3*a**2*b*d**3 + 6*tan(e + f*x)**3*a*b**2*c*d**2 + 4*tan(e + f*x)**3*b**3*c**2*d + tan(e + f*x)**2*a**3*d**3 + 6*tan(e + f*x)**2*a**2*b*c*d**2 + 9*tan(e + f*x)**2*a*b**2*c**2*d + 2*tan(e + f*x)**2*b**3*c**3 + 2*tan(e + f*x)*a**3*c*d**2 + 6*tan(e + f*x)*a**2*b*c**2*d + 4*tan(e + f*x)*a*b**2*c**3 + a**3*c**2*d + 2*a**2*b*c**3),x)*tan(e + f*x)**2*a**2*b**2*c*d**3*f + 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*a*b**2*d**3 + 2*tan(e + f*x)**4*b**3*c*d**2 + 2*tan(e + f*x)**3*a**2*b*d**3 + 6*tan(e + f*x)**3*a*b**2*c*d**2 + 4*tan(e + f*x)**3*b**3*c**2*d + tan(e + f*x)**2*a**3*d**3 + 6*tan(e + f*x)**2*a**2*b*c*d**2 + 9*tan(e + f*x)**2*a*b**2*c**2*d + 2*tan(e + f*x)**2*b**3*c**3 + 2*tan(e + f*x)*a**3*c*d**2 + 6*tan(e + f*x)*a**2*b*c**2*d + 4*tan(e + f*x)*a*b**2*c**3 + a**3*c**2*d + 2*a**2*b...
```

3.122
$$\int \frac{(a+b \tan(e+fx))^3 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1383
Mathematica [C] (verified)	1384
Rubi [A] (warning: unable to verify)	1386
Maple [B] (verified)	1392
Fricas [F(-1)]	1393
Sympy [F]	1393
Maxima [F(-1)]	1394
Giac [F(-1)]	1394
Mupad [F(-1)]	1394
Reduce [F]	1395

Optimal result

Integrand size = 47, antiderivative size = 585

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(a - ib)^3 (iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{5/2} f}$$

$$- \frac{(ia - b)^3 (A + iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{5/2} f}$$

$$- \frac{2(c^2 C - Bcd + Ad^2) (a + b \tan(e + fx))^3}{3d (c^2 + d^2) f (c + d \tan(e + fx))^{3/2}}$$

$$- \frac{2(b(2c^4 C - Bc^3 d + 4c^2 C d^2 - 3Bcd^3 + 2Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2))) (a + b \tan(e + fx))^2}{d^2 (c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

$$+ \frac{2b(3abd(8c^4 C - 2Bc^3 d - c^2(A - 17C)d^2 - 8Bcd^3 + (5A + 3C)d^4) - b^2(16c^5 C - 8Bc^4 d + 2c^3(A + 15C)d^2))}{3d^4 (c^2 + d^2)}$$

$$+ \frac{2b^2(b(8c^4 C - 4Bc^3 d + c^2(A + 15C)d^2 - 10Bcd^3 + (7A + C)d^4) + 3ad^2(2c(A - C)d - B(c^2 - d^2))) \tan(e + fx)}{3d^3 (c^2 + d^2)^2 f}$$

output

```

-(a-I*b)^3*(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*
d)^(5/2)/f-(I*a-b)^3*(A+I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2
))/((c+I*d)^(5/2)/f-2/3*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^3/d/(c^2+d^2)/
f/(c+d*tan(f*x+e))^(3/2)-2*(b*(2*A*d^4-B*c^3*d-3*B*c*d^3+2*C*c^4+4*C*c^2*d
^2)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*(a+b*tan(f*x+e))^2/d^2/(c^2+d^2)^2/f/
(c+d*tan(f*x+e))^(1/2)+2/3*b*(3*a*b*d*(8*c^4*C-2*B*c^3*d-c^2*(A-17*C)*d^2-
8*B*c*d^3+(5*A+3*C)*d^4)-b^2*(16*c^5*C-8*B*c^4*d+2*c^3*(A+15*C)*d^2-17*B*c
^2*d^3+8*c*(A+C)*d^4-3*B*d^5)+6*a^2*d^3*(2*c*(A-C)*d-B*(c^2-d^2))*(c+d*ta
n(f*x+e))^(1/2)/d^4/(c^2+d^2)^2/f+2/3*b^2*(b*(8*c^4*C-4*B*c^3*d+c^2*(A+15*
C)*d^2-10*B*c*d^3+(7*A+C)*d^4)+3*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*tan(f*x+
e)*(c+d*tan(f*x+e))^(1/2)/d^3/(c^2+d^2)^2/f

```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.59 (sec) , antiderivative size = 670, normalized size of antiderivative = 1.15

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \frac{2C(a + b \tan(e + fx))^3}{3df(c + d \tan(e + fx))^{3/2}}$$

$$\begin{aligned}
 & \left(\frac{3(2bcC - bBd - 2aCd)(a + b \tan(e + fx))^2}{df(c + d \tan(e + fx))^{3/2}} + \right. \\
 & \quad \left. \frac{3(b(Ab + aB - bC)d^2 + 4(bc - ad)(2bcC - bBd - 2aCd))(a + b \tan(e + fx))}{2df(c + d \tan(e + fx))^{3/2}} + \right. \\
 & \quad \left. \frac{2(-16b^3e^3C + 8b^3Bc)}{3} \right)
 \end{aligned}$$

input

```
Integrate[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^(5/2),x]
```

output

```
(2*C*(a + b*Tan[e + f*x])^3)/(3*d*f*(c + d*Tan[e + f*x])^(3/2)) + (2*((-3*(2*b*c*C - b*B*d - 2*a*C*d)*(a + b*Tan[e + f*x])^2)/(d*f*(c + d*Tan[e + f*x])^(3/2)) + (2*((-3*(b*(A*b + a*B - b*C)*d^2 + 4*(b*c - a*d)*(2*b*c*C - b*B*d - 2*a*C*d))*(a + b*Tan[e + f*x]))/(2*d*f*(c + d*Tan[e + f*x])^(3/2)) - (3*((-2*(-16*b^3*c^3*C + 8*b^3*B*c^2*d + 48*a*b^2*c^2*C*d - 2*A*b^3*c*d^2 - 18*a*b^2*B*c*d^2 - 48*a^2*b*c*C*d^2 + 2*b^3*c*C*d^2 + 9*a^2*b*B*d^3 + b^3*B*d^3 + 16*a^3*C*d^3))/(3*d*(c + d*Tan[e + f*x])^(3/2)) + (2((((3*c*(a^3*B - 3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C))*d^4)/2 + (3*(3*a^2*b*B - b^3*B - a^3*(A - C) + 3*a*b^2*(A - C))*d^5)/2)*(-1/3*Hypergeometric2F1[-3/2, 1, -1/2, (c + d*Tan[e + f*x])/(c - I*d)]/((I*c + d)*(c + d*Tan[e + f*x])^(3/2)) + Hypergeometric2F1[-3/2, 1, -1/2, (c + d*Tan[e + f*x])/(c + I*d)]/(3*(I*c - d)*(c + d*Tan[e + f*x])^(3/2))))/d - (3*(a^3*B - 3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C))*d^3*(-(Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)]/((I*c + d)*Sqrt[c + d*Tan[e + f*x]])) + Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c + I*d)]/((I*c - d)*Sqrt[c + d*Tan[e + f*x]])))/2))/(3*d)))/(4*d*f))/d)/(3*d)
```

Rubi [A] (warning: unable to verify)

Time = 7.53 (sec) , antiderivative size = 605, normalized size of antiderivative = 1.03, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.383$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4120, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$2 \int \frac{3(a+b \tan(e+fx))^2 (b(2Cc^2 - Bdc + (A+C)d^2) \tan^2(e+fx) + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+2bd) + \frac{2}{3} (3bc - \frac{3ad}{2}) (cC - B))}{2(c+d \tan(e+fx))^{3/2}} \frac{3d(c^2 + d^2)}{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3} \frac{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{(a+b \tan(e+fx))^2 (b(2Cc^2 - Bdc + (A+C)d^2) \tan^2(e+fx) + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+2bd) + (2bc-ad)(cC - Bd))}{(c+d \tan(e+fx))^{3/2}} \frac{d(c^2 + d^2)}{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3} \frac{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^2 (b(2Cc^2 - Bdc + (A+C)d^2) \tan^2(e+fx) + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+2bd) + (2bc-ad)(cC - Bd))}{(c+d \tan(e+fx))^{3/2}} \frac{d(c^2 + d^2)}{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3} \frac{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$2 \int \frac{(a+b \tan(e+fx)) \left(((ac+bd)((A-C)(bc-ad) + B(ac+bd)) - (bc-ad)(bBc - b(A-C)d - a(Ac - Cc + Bd))) \tan(e+fx)d^2 + 2 \left(\frac{ac}{2} + 2bd \right) (Ad(ac+2bd) + (2bc-ad)(cC - B)) \right)}{2\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{(a+b \tan(e+fx)) \left(((ac+bd)((A-C)(bc-ad) + B(ac+bd)) - (bc-ad)(bBc - b(A-C)d - a(Ac - Cc + Bd))) \tan(e+fx)d^2 + 2 \left(\frac{ac}{2} + 2bd \right) (Ad(ac+2bd) + (2bc-ad)(cC - B)) \right)}{\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{-3((Cc^2-2Bdc-Cd^2-A(c^2-d^2))a^3-3b(2c(A-C)d-B(c^2-d^2))a^2-3b^2(Cc^2-2Bdc-Cd^2-A(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))d^3-3((2c(A-C)d-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))d^3-3((2c(A-C)d-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))d^3-3((2c(A-C)d-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))d^3}{\sqrt{c+d \tan(e+fx)}}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 4022

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$-\frac{2(a+b \tan(e+fx))^2(ad^2(2cd(A-C)-B(c^2-d^2))+b(2Ad^4-Bc^3d-3Bcd^3+2c^4C+4c^2Cd^2))}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{2b^2 \tan(e+fx)\sqrt{c+d \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))}{\sqrt{c+d \tan(e+fx)}}$$

↓ 3042

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$-\frac{2(a+b \tan(e+fx))^2(ad^2(2cd(A-C)-B(c^2-d^2))+b(2Ad^4-Bc^3d-3Bcd^3+2c^4C+4c^2Cd^2))}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{2b^2 \tan(e+fx)\sqrt{c+d \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))}{\sqrt{c+d \tan(e+fx)}}$$

↓ 4020

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$-\frac{2(a+b \tan(e+fx))^2(ad^2(2cd(A-C)-B(c^2-d^2))+b(2Ad^4-Bc^3d-3Bcd^3+2c^4C+4c^2Cd^2))}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{2b^2 \tan(e+fx)\sqrt{c+d \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))}{\sqrt{c+d \tan(e+fx)}}$$

↓ 25

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^3}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$-\frac{2(a+b \tan(e+fx))^2(ad^2(2cd(A-C)-B(c^2-d^2))+b(2Ad^4-Bc^3d-3Bcd^3+2c^4C+4c^2Cd^2))}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{2b^2 \tan(e+fx)\sqrt{c+d \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))a+b^3(2c(A-C)d-B(c^2-d^2)))}{\sqrt{c+d \tan(e+fx)}}$$

↓ 73

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(a + b \tan(e + fx))^2(ad^2(2cd(A - C) - B(c^2 - d^2)) + b(2Ad^4 - Bc^3d - 3Bcd^3 + 2c^4C + 4c^2Cd^2))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{2b^2 \tan(e + fx)\sqrt{c + d \tan(e + fx)}(3ad^2(2cd(A - C) - B(c^2 - d^2)) + b(2Ad^4 - Bc^3d - 3Bcd^3 + 2c^4C + 4c^2Cd^2))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}}$$

↓ 221

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^3}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(a + b \tan(e + fx))^2(ad^2(2cd(A - C) - B(c^2 - d^2)) + b(2Ad^4 - Bc^3d - 3Bcd^3 + 2c^4C + 4c^2Cd^2))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{2b^2 \tan(e + fx)\sqrt{c + d \tan(e + fx)}(3ad^2(2cd(A - C) - B(c^2 - d^2)) + b(2Ad^4 - Bc^3d - 3Bcd^3 + 2c^4C + 4c^2Cd^2))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}}$$

input

```
Int[((a + b*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2), x]
```

output

```
(-2*(c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^3)/(3*d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) + ((-2*(b*(2*c^4*C - B*c^3*d + 4*c^2*C*d^2 - 3*B*c*d^3 + 2*A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*(a + b*Tan[e + f*x])^2)/(d*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]]) + ((2*b^2*(b*(8*c^4*C - 4*B*c^3*d + c^2*(A + 15*C)*d^2 - 10*B*c*d^3 + (7*A + C)*d^4) + 3*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Tan[e + f*x]*Sqrt[c + d*Tan[e + f*x]])/(3*d*f) + ((3*(a - I*b)^3*(A - I*B - C)*(c + I*d)^2*d^3*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f) + (3*(a + I*b)^3*(A + I*B - C)*(c - I*d)^2*d^3*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f) + (2*b*(3*a*b*d*(8*c^4*C - 2*B*c^3*d - c^2*(A - 17*C)*d^2 - 8*B*c*d^3 + (5*A + 3*C)*d^4) - b^2*(16*c^5*C - 8*B*c^4*d + 2*c^3*(A + 15*C)*d^2 - 17*B*c^2*d^3 + 8*c*(A + C)*d^4 - 3*B*d^5) + 6*a^2*d^3*(2*c*(A - C)*d - B*(c^2 - d^2)))*Sqrt[c + d*Tan[e + f*x]])/(d*f)/(3*d)/(d*(c^2 + d^2))/(d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \text{ Int}[\text{Fx}, \text{x}], \text{x}] \text{ /; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ /; FreeQ}[\text{b}, \text{x}]$
- rule 73 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(x_)^{\text{m}_}*((\text{c}_.) + (\text{d}_.)*(x_)^{\text{n}_}), \text{x_Symbol}] \rightarrow \text{With}[\{\text{p} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{p}/\text{b} \text{ Subst}[\text{Int}[x^{\text{p}*(\text{m} + 1) - 1}*(\text{c} - \text{a}*(\text{d}/\text{b}) + \text{d}*(x^{\text{p}}/\text{b})^{\text{n}}, \text{x}], \text{x}, (\text{a} + \text{b}*x)^{1/\text{p}}], \text{x}]] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \ \&\& \ \text{LtQ}[-1, \text{m}, 0] \ \&\& \ \text{LeQ}[-1, \text{n}, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[\text{n}], \text{Denominator}[\text{m}]] \ \&\& \ \text{IntLinearQ}[\text{a}, \text{b}, \text{c}, \text{d}, \text{m}, \text{n}, \text{x}]$
- rule 221 $\text{Int}[(\text{a}_.) + (\text{b}_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-\text{a}/\text{b}, 2]/\text{a})*\text{ArcTanh}[\text{x}/\text{Rt}[-\text{a}/\text{b}, 2]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{NegQ}[\text{a}/\text{b}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ /; FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{m}_}*((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c}*(\text{d}/\text{f}) \text{ Subst}[\text{Int}[(\text{a} + (\text{b}/\text{d})*x)^{\text{m}}/(\text{d}^2 + \text{c}*x), \text{x}], \text{x}, \text{d}*\text{Tan}[\text{e} + \text{f}*x]], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]^{\text{m}_}*((\text{c}_.) + (\text{d}_.)*\tan[(\text{e}_.) + (\text{f}_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I}*d)/2 \text{ Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m}}*(1 - \text{I}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I}*d)/2 \text{ Int}[(\text{a} + \text{b}*\text{Tan}[\text{e} + \text{f}*x])^{\text{m}}*(1 + \text{I}*\text{Tan}[\text{e} + \text{f}*x]), \text{x}], \text{x}] \text{ /; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \ \&\& \ \text{NeQ}[\text{b}*c - \text{a}*d, 0] \ \&\& \ \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \ \&\& \ \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \ \&\& \ \text{!IntegerQ}[\text{m}]$

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)]^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)]^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 13618 vs. 2(550) = 1100.

Time = 0.53 (sec) , antiderivative size = 13619, normalized size of antiderivative = 23.28

method	result	size
parts	Expression too large to display	13619
derivativedivides	Expression too large to display	85156
default	Expression too large to display	85156

input `int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((a+b*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(5/2),x)`

output `Integral((a + b*tan(e + f*x))**3*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(5/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output Timed out

Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input `int(((a + b*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((a+b*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**4 + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*b**3*c*d**3*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b**3*c**2*d**2*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**5)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b**3*c**3*d*f + 9*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*a*b**2*c*d**3*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*b**4*d**3*f + 18*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*a*b**2*c**2*d**2*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b**4*c*d**2*f + 9*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*a*b**2*c**3*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*b**4*c**2*d*f + ...
```


3.123
$$\int \frac{(a+b \tan(e+fx))^2 (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1396
Mathematica [C] (verified)	1397
Rubi [A] (warning: unable to verify)	1398
Maple [B] (verified)	1403
Fricas [F(-1)]	1403
Sympy [F]	1404
Maxima [F(-1)]	1404
Giac [F]	1404
Mupad [B] (verification not implemented)	1405
Reduce [F]	1406

Optimal result

Integrand size = 47, antiderivative size = 358

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(a - ib)^2 (iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{5/2} f}$$

$$- \frac{(a + ib)^2 (B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{5/2} f}$$

$$- \frac{2(c^2 C - Bcd + Ad^2) (a + b \tan(e + fx))^2}{3d (c^2 + d^2) f (c + d \tan(e + fx))^{3/2}}$$

$$+ \frac{2(bc - ad) (b(4c^4 C - Bc^3 d - 2c^2(A - 5C)d^2 - 7Bcd^3 + 4Ad^4) + 3ad^2(2c(A - C)d - B(c^2 - d^2)))}{3d^3 (c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

$$+ \frac{2b^2(4c^2 C - Bcd + (A + 3C)d^2) \sqrt{c + d \tan(e + fx)}}{3d^3 (c^2 + d^2) f}$$

output

$$\frac{-(a-I*b)^2*(I*A+B-I*C)*\operatorname{arctanh}\left(\frac{(c+d*\tan(f*x+e))^{1/2}}{(c-I*d)^{1/2}}\right)/(c-I*d)^{5/2}/f-(a+I*b)^2*(B-I*(A-C))*\operatorname{arctanh}\left(\frac{(c+d*\tan(f*x+e))^{1/2}}{(c+I*d)^{1/2}}\right)/(c+I*d)^{5/2}/f-2/3*(A*d^2-B*c*d+C*c^2)*(a+b*\tan(f*x+e))^2/d/(c^2+d^2)/f/(c+d*\tan(f*x+e))^{3/2}+2/3*(-a*d+b*c)*(b*(4*c^4*C-B*c^3*d-2*c^2*(A-5*C)*d^2-7*B*c*d^3+4*A*d^4)+3*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))/d^3/(c^2+d^2)^{2/2}/f/(c+d*\tan(f*x+e))^{1/2}+2/3*b^2*(4*c^2*C-B*c*d+(A+3*C)*d^2)*(c+d*\tan(f*x+e))^{1/2}/d^3/(c^2+d^2)/f}{1}$$
Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 4.83 (sec) , antiderivative size = 414, normalized size of antiderivative = 1.16

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{-2(c - id)(c + id)(8a^2Cd^2 + abd(-16cC + Bd) + b^2(8c^2C - 2Bcd + (-A + C)d^2)) - d^2(-2ab(Ac - c$$

input

```
Integrate[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/
(c + d*Tan[e + f*x])^(5/2),x]
```

output

$$\frac{-1/3*(-2*(c - I*d)*(c + I*d)*(8*a^2*C*d^2 + a*b*d*(-16*c*C + B*d) + b^2*(8*c^2*C - 2*B*c*d + (-A + C)*d^2)) - d^2*(-2*a*b*(A*c - c*C + B*d) - a^2*(B*c + (-A + C)*d) + b^2*(B*c + (-A + C)*d))*(I*(c + I*d)*\operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, (c + d*\tan[e + f*x])/(c - I*d)] - (I*c + d)*\operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, (c + d*\tan[e + f*x])/(c + I*d)]) - 6*(c - I*d)*(c + I*d)*d*(4*b*c*C - b*B*d - 4*a*C*d)*(a + b*\tan[e + f*x]) - 6*C*(c - I*d)*(c + I*d)*d^2*(a + b*\tan[e + f*x])^2 - 3*(a^2*B - b^2*B + 2*a*b*(A - C))*d^2*(I*(c + I*d)*\operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, (c + d*\tan[e + f*x])/(c - I*d)] - (I*c + d)*\operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, (c + d*\tan[e + f*x])/(c + I*d)])*(c + d*\tan[e + f*x])}{d^3*(c^2 + d^2)*f*(c + d*\tan[e + f*x])^{3/2}}$$

Rubi [A] (warning: unable to verify)

Time = 4.09 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.08, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.298$, Rules used = {3042, 4128, 27, 3042, 4118, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$2 \int \frac{(a + b \tan(e + fx))(b(4Cc^2 - Bdc + (A + 3C)d^2) \tan^2(e + fx) + 3d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + 3ad(Ac - Cc + Bd) + 4b(Cc^2 - Bdc + Ad^2 - Bcd + c^2C))}{2(c + d \tan(e + fx))^{3/2}}$$

$$\frac{3d(c^2 + d^2)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{(a + b \tan(e + fx))(b(4Cc^2 - Bdc + (A + 3C)d^2) \tan^2(e + fx) + 3d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + 3ad(Ac - Cc + Bd) + 4b(Cc^2 - Bdc + Ad^2 - Bcd + c^2C))}{(c + d \tan(e + fx))^{3/2}}$$

$$\frac{3d(c^2 + d^2)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))(b(4Cc^2 - Bdc + (A + 3C)d^2) \tan(e + fx)^2 + 3d((A - C)(bc - ad) + B(ac + bd)) \tan(e + fx) + 3ad(Ac - Cc + Bd) + 4b(Cc^2 - Bdc + Ad^2 - Bcd + c^2C))}{(c + d \tan(e + fx))^{3/2}}$$

$$\frac{3d(c^2 + d^2)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} \frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}}$$

↓ 4118

$$\int \frac{(c^2+d^2)(4Cc^2-Bdc+(A+3C)d^2) \tan^2(e+fx)b^2+(4Cc^4-Bdc^3-2(A-5C)d^2c^2-7Bd^3c+4Ad^4)b^2+6ad^2(2c(A-C)d-B(c^2-d^2))b-3a^2d^2(Cc^2-2Bdc-Cd^2)}{\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{(c^2+d^2)(4Cc^2-Bdc+(A+3C)d^2) \tan(e+fx)^2b^2+(4Cc^4-Bdc^3-2(A-5C)d^2c^2-7Bd^3c+4Ad^4)b^2+6ad^2(2c(A-C)d-B(c^2-d^2))b-3a^2d^2(Cc^2-2Bdc-Cd^2)}{\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 4113

$$\int \frac{-3((Cc^2-2Bdc-Cd^2-A(c^2-d^2))a^2-2b(2c(A-C)d-B(c^2-d^2))a-b^2(Cc^2-2Bdc-Cd^2-A(c^2-d^2)))d^2-3((2c(A-C)d-B(c^2-d^2))a^2+2b(Cc^2-2Bdc-Cd^2))}{\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{-3((Cc^2-2Bdc-Cd^2-A(c^2-d^2))a^2-2b(2c(A-C)d-B(c^2-d^2))a-b^2(Cc^2-2Bdc-Cd^2-A(c^2-d^2)))d^2-3((2c(A-C)d-B(c^2-d^2))a^2+2b(Cc^2-2Bdc-Cd^2))}{\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 4022

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^2}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2(bc-ad)(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-2c^2d^2(A-5C)+4Ad^4-Bc^3d-7Bcd^3+4c^4C))}{d^2f(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{\frac{3}{2}d^2(a+ib)^2(c-id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}}}{3d(c^2+d^2)}$$

↓ 3042

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(bc-ad)(3ad^2(2cd(A-C) - B(c^2 - d^2)) + b(-2c^2d^2(A-5C) + 4Ad^4 - Bc^3d - 7Bcd^3 + 4c^4C))}{d^2f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{\frac{3}{2}d^2(a+ib)^2(c-id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}}}{3d(c^2 + d^2)}$$

4020

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(bc-ad)(3ad^2(2cd(A-C) - B(c^2 - d^2)) + b(-2c^2d^2(A-5C) + 4Ad^4 - Bc^3d - 7Bcd^3 + 4c^4C))}{d^2f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{3id^2(a-ib)^2(c+id)^2(A-iB-C) \int -\frac{1-i \tan(e+fx)}{2f}}{3d(c^2 + d^2)}$$

25

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(bc-ad)(3ad^2(2cd(A-C) - B(c^2 - d^2)) + b(-2c^2d^2(A-5C) + 4Ad^4 - Bc^3d - 7Bcd^3 + 4c^4C))}{d^2f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{3id^2(a-ib)^2(c+id)^2(A-iB-C) \int \frac{1-i \tan(e+fx)}{2f}}{3d(c^2 + d^2)}$$

73

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(bc-ad)(3ad^2(2cd(A-C) - B(c^2 - d^2)) + b(-2c^2d^2(A-5C) + 4Ad^4 - Bc^3d - 7Bcd^3 + 4c^4C))}{d^2f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{3d(a+ib)^2(c-id)^2(A+iB-C) \int \frac{1-i \tan^2(e+fx) - \frac{ic}{d}}{f}}{3d(c^2 + d^2)}$$

221

$$\frac{2(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^2}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \frac{2(bc-ad)(3ad^2(2cd(A-C) - B(c^2 - d^2)) + b(-2c^2d^2(A-5C) + 4Ad^4 - Bc^3d - 7Bcd^3 + 4c^4C))}{d^2f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} + \frac{3d^2(a-ib)^2(c+id)^2(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}}}{3d(c^2 + d^2)}$$

input

```
Int[((a + b*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2), x]
```

output

$$\begin{aligned} & (-2*(c^2*C - B*c*d + A*d^2)*(a + b*\text{Tan}[e + f*x])^2)/(3*d*(c^2 + d^2)*f*(c \\ & + d*\text{Tan}[e + f*x])^{3/2}) + ((2*(b*c - a*d)*(b*(4*c^4*C - B*c^3*d - 2*c^2*(\\ & A - 5*C)*d^2 - 7*B*c*d^3 + 4*A*d^4) + 3*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^ \\ & 2))))/(d^2*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]) + ((3*(a - I*b)^2*(A - \\ & I*B - C)*(c + I*d)^2*d^2*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]])/(\text{Sqrt}[c - I*d \\ &]*f) + (3*(a + I*b)^2*(A + I*B - C)*(c - I*d)^2*d^2*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqr} \\ & \text{rt}[c + I*d]])/(\text{Sqrt}[c + I*d]*f) + (2*b^2*(c^2 + d^2)*(4*c^2*C - B*c*d + (A \\ & + 3*C)*d^2)*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/(d*f)/(d*(c^2 + d^2))/(3*d*(c^2 + \\ & d^2)) \end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[a, \text{x}] \ \&\& \ \text{!Ma} \\ \text{tchQ}[\text{Fx}, (b_)*(\text{Gx}_) \text{ ; FreeQ}[b, \text{x}]$$

rule 73

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_)^{(n_)}, \text{x_Symbol}] \rightarrow \text{With} [\\ \{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + \\ d*(x^p/b)^n, \text{x}], \text{x}, (a + b*x)^{(1/p)}, \text{x}]] \text{ ; FreeQ}[\{a, b, c, d\}, \text{x}] \ \&\& \ \text{Lt} \\ \text{Q}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntL} \\ \text{inearQ}[a, b, c, d, m, n, \text{x}]$$

rule 221

$$\text{Int}[(a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x \\ / \text{Rt}[-a/b, 2]], \text{x}] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{NegQ}[a/b]$$

rule 3042

$$\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinear} \\ \text{Q}[u, \text{x}]$$

rule 4020

$$\text{Int}[(a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^{(m_)}*((c_.) + (d_.)*\text{tan}[(e_.) + \\ (f_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[c*(d/f) \quad \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + \\ c*x), \text{x}], \text{x}, d*\text{Tan}[e + f*x], \text{x}] \text{ ; FreeQ}[\{a, b, c, d, e, f, m\}, \text{x}] \ \&\& \ \text{NeQ} [\\ b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$$

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 11385 vs. $2(325) = 650$.

Time = 0.26 (sec) , antiderivative size = 11386, normalized size of antiderivative = 31.80

method	result	size
parts	Expression too large to display	11386
derivativeldivides	Expression too large to display	61833
default	Expression too large to display	61833

input `int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((a+b*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(5/2),x)`

output `Integral((a + b*tan(e + f*x))**2*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(5/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^2}{(d \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^2/(d*tan(f*x + e) + c)^(5/2), x)`

Mupad [B] (verification not implemented)

Time = 110.56 (sec) , antiderivative size = 88684, normalized size of antiderivative = 247.72

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input

```
int(((a + b*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c +
d*tan(e + f*x))^(5/2),x)
```

output

```
atan((((c + d*tan(e + f*x))^(1/2)*(96*A^2*a^2*b^2*d^18*f^3 - 16*A^2*b^4*d^
18*f^3 - 16*A^2*a^4*d^18*f^3 + 320*A^2*a^4*c^4*d^14*f^3 + 1024*A^2*a^4*c^6
*d^12*f^3 + 1440*A^2*a^4*c^8*d^10*f^3 + 1024*A^2*a^4*c^10*d^8*f^3 + 320*A^
2*a^4*c^12*d^6*f^3 - 16*A^2*a^4*c^16*d^2*f^3 + 320*A^2*b^4*c^4*d^14*f^3 +
1024*A^2*b^4*c^6*d^12*f^3 + 1440*A^2*b^4*c^8*d^10*f^3 + 1024*A^2*b^4*c^10*
d^8*f^3 + 320*A^2*b^4*c^12*d^6*f^3 - 16*A^2*b^4*c^16*d^2*f^3 - 256*A^2*a*b
^3*c*d^17*f^3 + 256*A^2*a^3*b*c*d^17*f^3 - 1280*A^2*a*b^3*c^3*d^15*f^3 - 2
304*A^2*a*b^3*c^5*d^13*f^3 - 1280*A^2*a*b^3*c^7*d^11*f^3 + 1280*A^2*a*b^3*
c^9*d^9*f^3 + 2304*A^2*a*b^3*c^11*d^7*f^3 + 1280*A^2*a*b^3*c^13*d^5*f^3 +
256*A^2*a*b^3*c^15*d^3*f^3 + 1280*A^2*a^3*b*c^3*d^15*f^3 + 2304*A^2*a^3*b*
c^5*d^13*f^3 + 1280*A^2*a^3*b*c^7*d^11*f^3 - 1280*A^2*a^3*b*c^9*d^9*f^3 -
2304*A^2*a^3*b*c^11*d^7*f^3 - 1280*A^2*a^3*b*c^13*d^5*f^3 - 256*A^2*a^3*b*
c^15*d^3*f^3 - 1920*A^2*a^2*b^2*c^4*d^14*f^3 - 6144*A^2*a^2*b^2*c^6*d^12*f
^3 - 8640*A^2*a^2*b^2*c^8*d^10*f^3 - 6144*A^2*a^2*b^2*c^10*d^8*f^3 - 1920*
A^2*a^2*b^2*c^12*d^6*f^3 + 96*A^2*a^2*b^2*c^16*d^2*f^3) + (((8*A^2*a^4*c^
5*f^2 + 8*A^2*b^4*c^5*f^2 - 48*A^2*a^2*b^2*c^5*f^2 - 80*A^2*a^4*c^3*d^2*f^
2 - 80*A^2*b^4*c^3*d^2*f^2 - 32*A^2*a*b^3*d^5*f^2 + 32*A^2*a^3*b*d^5*f^2 +
40*A^2*a^4*c*d^4*f^2 + 40*A^2*b^4*c*d^4*f^2 - 160*A^2*a*b^3*c^4*d*f^2 + 1
60*A^2*a^3*b*c^4*d*f^2 + 320*A^2*a*b^3*c^2*d^3*f^2 - 240*A^2*a^2*b^2*c*d^4
*f^2 - 320*A^2*a^3*b*c^2*d^3*f^2 + 480*A^2*a^2*b^2*c^3*d^2*f^2)^2/4 - (...
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((a+b*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**3 + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*b**2*c*d**3*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b**2*c**2*d**2*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**4)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*b**2*c**3*d*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*a*b*c*d**3*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*b**3*d**3*f + 12*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*a*b*c**2*d**2*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b**3*c*d**2*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*a*b*c**3*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*b**3*c**2*d*f - 3*int((sq...
```

3.124 $\int \frac{(a+b \tan(e+fx))(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$

Optimal result	1407
Mathematica [C] (verified)	1408
Rubi [A] (warning: unable to verify)	1408
Maple [B] (verified)	1412
Fricas [F(-1)]	1413
Sympy [F]	1413
Maxima [F(-1)]	1414
Giac [F(-2)]	1414
Mupad [B] (verification not implemented)	1414
Reduce [F]	1415

Optimal result

Integrand size = 45, antiderivative size = 273

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(a - ib)(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{5/2} f}$$

$$+ \frac{(ia - b)(A + iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{5/2} f}$$

$$+ \frac{2(bc - ad)(c^2 C - Bcd + Ad^2)}{3d^2 (c^2 + d^2) f (c + d \tan(e + fx))^{3/2}}$$

$$- \frac{2(b(c^4 C - c^2(A - 3C)d^2 - 2Bcd^3 + Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2)))}{d^2 (c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

output

```
- (a-I*b)*(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(5/2)/f+(I*a-b)*(A+I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(5/2)/f+2/3*(-a*d+b*c)*(A*d^2-B*c*d+C*c^2)/d^2/(c^2+d^2)/f/(c+d*tan(f*x+e))^(3/2)-2*(b*(c^4*C-c^2*(A-3*C)*d^2-2*B*c*d^3+A*d^4)+a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))/d^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.00 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{2(c - id)(c + id)(2bcC + bBd - 2aCd) + d(abc + aBc - bcC - aAd + bBd + aCd)}{(c + id)} \text{Hypergeometric2F1}[-3/2, 1, -1/2, (c + d \tan(e + fx))/(c + id)] + 6C(c - id)(c + id)d(a + b \tan(e + fx)) - 3(Ab + aB - bC)d(I(c + id) \text{Hypergeometric2F1}[-1/2, 1, 1/2, (c + d \tan(e + fx))/(c - id)] - (Ic + d) \text{Hypergeometric2F1}[-1/2, 1, 1/2, (c + d \tan(e + fx))/(c + id)]) * (c + d \tan(e + fx)) / (d^2 * (c^2 + d^2) * f * (c + d \tan(e + fx))^{3/2})$$

input

```
Integrate[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2),x]
```

output

```
-1/3*(2*(c - I*d)*(c + I*d)*(2*b*c*C + b*B*d - 2*a*C*d) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*(I*(c + I*d)*Hypergeometric2F1[-3/2, 1, -1/2, (c + d*Tan[e + f*x])/(c - I*d)] - (I*c + d)*Hypergeometric2F1[-3/2, 1, -1/2, (c + d*Tan[e + f*x])/(c + I*d)]) + 6*C*(c - I*d)*(c + I*d)*d*(a + b*Tan[e + f*x]) - 3*(A*b + a*B - b*C)*d*(I*(c + I*d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)] - (I*c + d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c + I*d)])*(c + d*Tan[e + f*x])/(d^2*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2))
```

Rubi [A] (warning: unable to verify)

Time = 2.51 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.10, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4118, 3042, 4111, 25, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{5/2}} dx$$

4118

$$\int \frac{bC(c^2+d^2) \tan^2(e+fx) + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{(c+d \tan(e+fx))^{3/2}} dx + \frac{d(c^2+d^2)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}} \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

3042

$$\int \frac{bC(c^2+d^2) \tan(e+fx)^2 + d(abc+aBc-bCc-aAd+bBd+aCd) \tan(e+fx) + ad(Ac-Cc+Bd) + b(Cc^2-Bdc+Ad^2)}{(c+d \tan(e+fx))^{3/2}} dx + \frac{d(c^2+d^2)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}} \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

4111

$$\int \frac{d(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2))) + d(2aAcd-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)} c^2+d^2} dx - \frac{d(c^2+d^2)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}} \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

25

$$\int \frac{d(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2))) + d(2aAcd-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)} c^2+d^2} dx - \frac{d(c^2+d^2)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}} \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

3042

$$\int \frac{d(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2))) + d(2aAcd-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{\sqrt{c+d \tan(e+fx)} c^2+d^2} dx - \frac{d(c^2+d^2)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}} \frac{2(bc-ad)(Ad^2-Bcd+c^2C)}{3d^2 f(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

4022

$$\frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{3d^2 f (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2(ad^2 (2cd(A-C) - B(c^2 - d^2)) + b(-c^2 d^2(A-3C) + Ad^4 - 2Bcd^3 + c^4 C))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{-\frac{1}{2}d(a+ib)(c-id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{1}{2}d(a-ib)(c+id)^2(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{c^2 + d^2}$$

$$d(c^2 + d^2)$$

↓ 3042

$$\frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{3d^2 f (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2(ad^2 (2cd(A-C) - B(c^2 - d^2)) + b(-c^2 d^2(A-3C) + Ad^4 - 2Bcd^3 + c^4 C))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{-\frac{1}{2}d(a+ib)(c-id)^2(A+iB-C) \int \frac{1-i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx - \frac{1}{2}d(a-ib)(c+id)^2(A-iB-C) \int \frac{1+i \tan(e+fx)}{\sqrt{c+d \tan(e+fx)}} dx}{c^2 + d^2}$$

$$d(c^2 + d^2)$$

↓ 4020

$$\frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{3d^2 f (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2(ad^2 (2cd(A-C) - B(c^2 - d^2)) + b(-c^2 d^2(A-3C) + Ad^4 - 2Bcd^3 + c^4 C))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{id(a+ib)(c-id)^2(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) - id(a-ib)(c+id)^2(A-iB-C) \int \frac{1}{(-i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f}$$

$$d(c^2 + d^2)$$

↓ 25

$$\frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{3d^2 f (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2(ad^2 (2cd(A-C) - B(c^2 - d^2)) + b(-c^2 d^2(A-3C) + Ad^4 - 2Bcd^3 + c^4 C))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{id(a-ib)(c+id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx)) + id(a+ib)(c-id)^2(A+iB-C) \int \frac{1}{(1+i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f}$$

$$d(c^2 + d^2)$$

↓ 73

$$\frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{3d^2 f (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2(ad^2 (2cd(A-C) - B(c^2 - d^2)) + b(-c^2 d^2(A-3C) + Ad^4 - 2Bcd^3 + c^4 C))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{(a+ib)(c-id)^2(A+iB-C) \int \frac{1}{f} \frac{d\sqrt{c+d \tan(e+fx)}}{d} - \frac{ic}{d} + 1}{f}$$

$$d(c^2 + d^2)$$

↓ 221

$$\frac{2(bc - ad) (Ad^2 - Bcd + c^2C)}{3d^2 f (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2(ad^2 (2cd(A-C) - B(c^2 - d^2)) + b(-c^2 d^2(A-3C) + Ad^4 - 2Bcd^3 + c^4 C))}{df(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{d(a+ib)(c-id)^2(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right) - d(a-ib)(c+id)^2(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c+id}}$$

$$d(c^2 + d^2)$$

input `Int[((a + b*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2), x]`

output `(2*(b*c - a*d)*(c^2*C - B*c*d + A*d^2))/(3*d^2*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) + (-(((a - I*b)*(A - I*B - C)*(c + I*d)^2*d*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f)) - ((a + I*b)*(A + I*B - C)*(c - I*d)^2*d*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f))/(c^2 + d^2) - (2*(b*(c^4*C - c^2*(A - 3*C)*d^2 - 2*B*c*d^3 + A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))/(d*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]]))/(d*(c^2 + d^2))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

rule 4118

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8982 vs. $2(246) = 492$.

Time = 0.25 (sec) , antiderivative size = 8983, normalized size of antiderivative = 32.90

method	result	size
parts	Expression too large to display	8983
derivativedivides	Expression too large to display	40201
default	Expression too large to display	40201

input `int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(5/2),x)`

output `Integral((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(5/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^5/2,x, algorithm="maxima")`

output `Timed out`

Giac [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^5/2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to rounding error%%{1, [6,14,5]%%}+%%{6, [6,12,5]%%}+%%{15, [6,10,5]%%}+%%{20, [6`

Mupad [B] (verification not implemented)

Time = 83.07 (sec) , antiderivative size = 64641, normalized size of antiderivative = 236.78

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `int(((a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2),x)`

output `((2*(C*b*c^3 + A*b*c*d^2 - B*b*c^2*d))/(3*(c^2 + d^2)) - (2*(c + d*tan(e + f*x))*(A*b*d^4 + C*b*c^4 - 2*B*b*c*d^3 - A*b*c^2*d^2 + 3*C*b*c^2*d^2))/(c^2 + d^2)^2)/(d^2*f*(c + d*tan(e + f*x))^(3/2)) - atan(-(((c + d*tan(e + f*x))^(1/2)*(16*A^2*b^2*d^18*f^3 - 16*B^2*b^2*d^18*f^3 + 16*C^2*b^2*d^18*f^3 - 320*A^2*b^2*c^4*d^14*f^3 - 1024*A^2*b^2*c^6*d^12*f^3 - 1440*A^2*b^2*c^8*d^10*f^3 - 1024*A^2*b^2*c^10*d^8*f^3 - 320*A^2*b^2*c^12*d^6*f^3 + 16*A^2*b^2*c^16*d^2*f^3 + 320*B^2*b^2*c^4*d^14*f^3 + 1024*B^2*b^2*c^6*d^12*f^3 + 1440*B^2*b^2*c^8*d^10*f^3 + 1024*B^2*b^2*c^10*d^8*f^3 + 320*B^2*b^2*c^12*d^6*f^3 - 16*B^2*b^2*c^16*d^2*f^3 - 320*C^2*b^2*c^4*d^14*f^3 - 1024*C^2*b^2*c^6*d^12*f^3 - 1440*C^2*b^2*c^8*d^10*f^3 - 1024*C^2*b^2*c^10*d^8*f^3 - 320*C^2*b^2*c^12*d^6*f^3 + 16*C^2*b^2*c^16*d^2*f^3 - 32*A*C*b^2*d^18*f^3 - 128*A*B*b^2*c*d^17*f^3 + 128*B*C*b^2*c*d^17*f^3 - 640*A*B*b^2*c^3*d^15*f^3 - 1152*A*B*b^2*c^5*d^13*f^3 - 640*A*B*b^2*c^7*d^11*f^3 + 640*A*B*b^2*c^9*d^9*f^3 + 1152*A*B*b^2*c^11*d^7*f^3 + 640*A*B*b^2*c^13*d^5*f^3 + 128*A*B*b^2*c^15*d^3*f^3 + 640*A*C*b^2*c^4*d^14*f^3 + 2048*A*C*b^2*c^6*d^12*f^3 + 2880*A*C*b^2*c^8*d^10*f^3 + 2048*A*C*b^2*c^10*d^8*f^3 + 640*A*C*b^2*c^12*d^6*f^3 - 32*A*C*b^2*c^16*d^2*f^3 + 640*B*C*b^2*c^3*d^15*f^3 + 1152*B*C*b^2*c^5*d^13*f^3 + 640*B*C*b^2*c^7*d^11*f^3 - 640*B*C*b^2*c^9*d^9*f^3 - 1152*B*C*b^2*c^11*d^7*f^3 - 640*B*C*b^2*c^13*d^5*f^3 - 128*B*C*b^2*c^15*d^3*f^3) + (((8*A^2*b^2*c^5*f^2 - 8*B^2*b^2*c^5*f^2 + 8*C^2*b^2*c^5*f^2 - 80*A...`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `int((a+b*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)`

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*a**2 + 3*int((sqrt(tan(e + f*x)*d + c)*tan(
e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e +
f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*b*c*d**3*f + 6*int((sqrt(tan(e + f*
x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**
2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b*c**2*d**2*f + 3*int((s
qrt(tan(e + f*x)*d + c)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e +
f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*b*c**3*d*f - 3*int((sqr
t(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f
*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*a**2*d**3
*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**
3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*
x)**2*a*c*d**3*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e
 + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3)
,x)*tan(e + f*x)**2*b**2*d**3*f - 6*int((sqrt(tan(e + f*x)*d + c)*tan(e +
f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)
*c**2*d + c**3),x)*tan(e + f*x)*a**2*c*d**2*f + 6*int((sqrt(tan(e + f*x)*d
 + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 +
3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*a*c**2*d**2*f + 6*int((sqrt(
tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)
)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b**2*c*d**2...
```

3.125
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1417
Mathematica [C] (verified)	1418
Rubi [A] (warning: unable to verify)	1418
Maple [B] (verified)	1422
Fricas [B] (verification not implemented)	1422
Sympy [F]	1423
Maxima [F(-1)]	1423
Giac [F(-2)]	1423
Mupad [B] (verification not implemented)	1424
Reduce [F]	1425

Optimal result

Integrand size = 35, antiderivative size = 209

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(c - id)^{5/2} f} - \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(c + id)^{5/2} f} - \frac{2(c^2 C - Bcd + Ad^2)}{3d(c^2 + d^2) f (c + d \tan(e + fx))^{3/2}} - \frac{2(2c(A - C)d - B(c^2 - d^2))}{(c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

output

```

-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(c-I*d)^(5/2)/f
-(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(c+I*d)^(5/2)/f
-2/3*(A*d^2-B*c*d+C*c^2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(3/2)-2*(2*c*(A-C)
*d-B*(c^2-d^2))/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)
    
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.64 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.07

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{2C(c^2 + d^2) + (Bc + (-A + C)d) \left(i(c + id) \operatorname{Hypergeometric2F1} \left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{c+d \tan(e+fx)}{c-id} \right) - (ic + d) \operatorname{Hypergeometric2F1} \left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{c+d \tan(e+fx)}{c+id} \right) \right)}{(c + d \tan(e + fx))^{5/2}}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^(5/2), x]
```

output

```
-1/3*(2*C*(c^2 + d^2) + (B*c + (-A + C)*d)*(I*(c + I*d)*Hypergeometric2F1[-3/2, 1, -1/2, (c + d*Tan[e + f*x])/(c - I*d)] - (I*c + d)*Hypergeometric2F1[-3/2, 1, -1/2, (c + d*Tan[e + f*x])/(c + I*d)]) - 3*B*(I*(c + I*d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c - I*d)] - (I*c + d)*Hypergeometric2F1[-1/2, 1, 1/2, (c + d*Tan[e + f*x])/(c + I*d)]*(c + d*Tan[e + f*x]))/(d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2))
```

Rubi [A] (warning: unable to verify)

Time = 1.76 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.11, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4111, 3042, 4012, 25, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(c + d \tan(e + fx))^{5/2}} dx$$

$$\downarrow \text{4111}$$

$$\begin{aligned}
 & \frac{\int \frac{Ac-Cc+Bd+(Bc-(A-C)d)\tan(e+fx)}{(c+d\tan(e+fx))^{3/2}} dx}{c^2+d^2} - \frac{2(Ad^2-Bcd+c^2C)}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Ac-Cc+Bd+(Bc-(A-C)d)\tan(e+fx)}{(c+d\tan(e+fx))^{3/2}} dx}{c^2+d^2} - \frac{2(Ad^2-Bcd+c^2C)}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int -\frac{Ce^2-2Bdc-Cd^2-A(c^2-d^2)+(2c(A-C)d-B(c^2-d^2))\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} - \frac{2(2cd(A-C)-B(c^2-d^2))}{f(c^2+d^2)\sqrt{c+d\tan(e+fx)}} \\
 & \quad \frac{c^2+d^2}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int -\frac{Ce^2-2Bdc-Cd^2-A(c^2-d^2)+(2c(A-C)d-B(c^2-d^2))\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} - \frac{2(2cd(A-C)-B(c^2-d^2))}{f(c^2+d^2)\sqrt{c+d\tan(e+fx)}} \\
 & \quad \frac{c^2+d^2}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -\frac{Ce^2-2Bdc-Cd^2-A(c^2-d^2)+(2c(A-C)d-B(c^2-d^2))\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} - \frac{2(2cd(A-C)-B(c^2-d^2))}{f(c^2+d^2)\sqrt{c+d\tan(e+fx)}} \\
 & \quad \frac{c^2+d^2}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} \\
 & \quad \downarrow \text{4022} \\
 & \frac{2(Ad^2-Bcd+c^2C)}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} + \\
 & \frac{2(2cd(A-C)-B(c^2-d^2))}{f(c^2+d^2)\sqrt{c+d\tan(e+fx)}} - \frac{-\frac{1}{2}(c-id)^2(A+iB-C)\int \frac{1-i\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx - \frac{1}{2}(c+id)^2(A-iB-C)\int \frac{i\tan(e+fx)+1}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2(Ad^2-Bcd+c^2C)}{3df(c^2+d^2)(c+d\tan(e+fx))^{3/2}} + \\
 & \frac{2(2cd(A-C)-B(c^2-d^2))}{f(c^2+d^2)\sqrt{c+d\tan(e+fx)}} - \frac{-\frac{1}{2}(c-id)^2(A+iB-C)\int \frac{1-i\tan(e+fx)}{\sqrt{c+d\tan(e+fx)}} dx - \frac{1}{2}(c+id)^2(A-iB-C)\int \frac{i\tan(e+fx)+1}{\sqrt{c+d\tan(e+fx)}} dx}{c^2+d^2} \\
 & \quad \frac{c^2+d^2}{c^2+d^2}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 4020 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \\ & \frac{2(2cd(A-C) - B(c^2 - d^2))}{f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{i(c-id)^2(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} - \frac{i(c+id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} \\ & \frac{c^2 + d^2}{c^2 + d^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \\ & \frac{2(2cd(A-C) - B(c^2 - d^2))}{f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{i(c+id)^2(A-iB-C) \int \frac{1}{(1-i \tan(e+fx))\sqrt{c+d \tan(e+fx)}} d(i \tan(e+fx))}{2f} - \frac{i(c-id)^2(A+iB-C) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c+d \tan(e+fx)}} d(-i \tan(e+fx))}{2f} \\ & \frac{c^2 + d^2}{c^2 + d^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \\ & \frac{2(2cd(A-C) - B(c^2 - d^2))}{f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{(c-id)^2(A+iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} - \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} - \frac{(c+id)^2(A-iB-C) \int \frac{1}{\frac{i \tan^2(e+fx)}{d} + \frac{ic}{d} + 1} d\sqrt{c+d \tan(e+fx)}}{df} \\ & \frac{c^2 + d^2}{c^2 + d^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & \frac{2(Ad^2 - Bcd + c^2C)}{3df(c^2 + d^2)(c + d \tan(e + fx))^{3/2}} + \\ & \frac{2(2cd(A-C) - B(c^2 - d^2))}{f(c^2 + d^2)\sqrt{c + d \tan(e + fx)}} - \frac{(c-id)^2(A+iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c+id}}\right)}{f\sqrt{c+id}} - \frac{(c+id)^2(A-iB-C) \arctan\left(\frac{\tan(e+fx)}{\sqrt{c-id}}\right)}{f\sqrt{c-id}} \\ & \frac{c^2 + d^2}{c^2 + d^2} \end{aligned}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(c + d*Tan[e + f*x])^(5/2),x]`

output `(-2*(c^2*C - B*c*d + A*d^2))/(3*d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2) + (-(((A - I*B - C)*(c + I*d)^2*ArcTan[Tan[e + f*x]/Sqrt[c - I*d]])/(Sqrt[c - I*d]*f)) - ((A + I*B - C)*(c - I*d)^2*ArcTan[Tan[e + f*x]/Sqrt[c + I*d]])/(Sqrt[c + I*d]*f))/(c^2 + d^2) - (2*(2*c*(A - C)*d - B*(c^2 - d^2)))/((c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])/(c^2 + d^2)`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4111

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6802 vs. $2(184) = 368$.

Time = 0.17 (sec) , antiderivative size = 6803, normalized size of antiderivative = 32.55

method	result	size
parts	Expression too large to display	6803
derivativedivides	Expression too large to display	20647
default	Expression too large to display	20647

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x,method=_RETUR
NVERBOSE)
```

output

result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13143 vs. $2(177) = 354$.

Time = 7.80 (sec) , antiderivative size = 13143, normalized size of antiderivative = 62.89

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algori
thm="fricas")
```

output Too large to include

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)**(5/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x)**(5/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output

```
Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to divide, perhaps due to ro
unding error%%{%%{1,[6,14,5]%%}+%%{6,[6,12,5]%%}+%%{15,[6,10,5]%%}+
%%{20,[6
```

Mupad [B] (verification not implemented)

Time = 33.50 (sec) , antiderivative size = 14163, normalized size of antiderivative = 67.77

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/(c + d*tan(e + f*x))^(5/2),x)
```

output

```
(log(96*A^3*c^3*d^13*f^2 - ((((((320*A^4*c^2*d^8*f^4 - 16*A^4*d^10*f^4 - 1
760*A^4*c^4*d^6*f^4 + 1600*A^4*c^6*d^4*f^4 - 400*A^4*c^8*d^2*f^4)^(1/2) -
4*A^2*c^5*f^2 + 40*A^2*c^3*d^2*f^2 - 20*A^2*c*d^4*f^2)/(c^10*f^4 + d^10*f^
4 + 5*c^2*d^8*f^4 + 10*c^4*d^6*f^4 + 10*c^6*d^4*f^4 + 5*c^8*d^2*f^4))^(1/2)
)*((((((320*A^4*c^2*d^8*f^4 - 16*A^4*d^10*f^4 - 1760*A^4*c^4*d^6*f^4 + 1600
*A^4*c^6*d^4*f^4 - 400*A^4*c^8*d^2*f^4)^(1/2) - 4*A^2*c^5*f^2 + 40*A^2*c^3
*d^2*f^2 - 20*A^2*c*d^4*f^2)/(c^10*f^4 + d^10*f^4 + 5*c^2*d^8*f^4 + 10*c^4
*d^6*f^4 + 10*c^6*d^4*f^4 + 5*c^8*d^2*f^4))^(1/2)*(c + d*tan(e + f*x))^(1/
2)*(64*c*d^22*f^5 + 640*c^3*d^20*f^5 + 2880*c^5*d^18*f^5 + 7680*c^7*d^16*f
^5 + 13440*c^9*d^14*f^5 + 16128*c^11*d^12*f^5 + 13440*c^13*d^10*f^5 + 7680
*c^15*d^8*f^5 + 2880*c^17*d^6*f^5 + 640*c^19*d^4*f^5 + 64*c^21*d^2*f^5))/4
- 32*A*d^21*f^4 - 160*A*c^2*d^19*f^4 - 128*A*c^4*d^17*f^4 + 896*A*c^6*d^1
5*f^4 + 3136*A*c^8*d^13*f^4 + 4928*A*c^10*d^11*f^4 + 4480*A*c^12*d^9*f^4 +
2432*A*c^14*d^7*f^4 + 736*A*c^16*d^5*f^4 + 96*A*c^18*d^3*f^4))/4 - (c + d
*tan(e + f*x))^(1/2)*(320*A^2*c^4*d^14*f^3 - 16*A^2*d^18*f^3 + 1024*A^2*c^
6*d^12*f^3 + 1440*A^2*c^8*d^10*f^3 + 1024*A^2*c^10*d^8*f^3 + 320*A^2*c^12*
d^6*f^3 - 16*A^2*c^16*d^2*f^3))*((((320*A^4*c^2*d^8*f^4 - 16*A^4*d^10*f^4 -
1760*A^4*c^4*d^6*f^4 + 1600*A^4*c^6*d^4*f^4 - 400*A^4*c^8*d^2*f^4)^(1/2)
- 4*A^2*c^5*f^2 + 40*A^2*c^3*d^2*f^2 - 20*A^2*c*d^4*f^2)/(c^10*f^4 + d^10*
f^4 + 5*c^2*d^8*f^4 + 10*c^4*d^6*f^4 + 10*c^6*d^4*f^4 + 5*c^8*d^2*f^4))...
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)`

output `(- 2*sqrt(tan(e + f*x)*d + c)*a - 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*a*d**3*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*c*d**3*f - 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*a*c*d**2*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*c**2*d**2*f - 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*a*c**2*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x)**2)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*c**3*d*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*b*d**3*f + 6*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*b*c*d**2*f + 3*int((sqrt(tan(e + f*x)*d + c)*tan(e + f*x))/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*b*c**2*d*f)/(3*d*f*(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2))`

3.126
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1426
Mathematica [B] (verified)	1427
Rubi [A] (warning: unable to verify)	1428
Maple [B] (verified)	1434
Fricas [F(-1)]	1434
Sympy [F]	1434
Maxima [F(-2)]	1435
Giac [F(-2)]	1435
Mupad [F(-1)]	1436
Reduce [F]	1436

Optimal result

Integrand size = 47, antiderivative size = 365

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \frac{(A - iB - C) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right)}{(ia + b)(c - id)^{5/2} f}$$

$$+ \frac{(iA - B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a + ib)(c + id)^{5/2} f}$$

$$- \frac{2b^{3/2}(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c+d \tan(e+fx)}}{\sqrt{bc-ad}}\right)}{(a^2 + b^2)(bc - ad)^{5/2} f}$$

$$+ \frac{2(c^2C - Bcd + Ad^2)}{3(bc - ad)(c^2 + d^2) f(c + d \tan(e + fx))^{3/2}}$$

$$+ \frac{2(b(c^4C - 2Bc^3d + c^2(3A - C)d^2 + Ad^4) - ad^2(2c(A - C)d - B(c^2 - d^2)))}{(bc - ad)^2 (c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

output

```
(A-I*B-C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(I*a+b)/(c-I*d)^(5/2)/f+(I*A-B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(a+I*b)/(c+I*d)^(5/2)/f-2*b^(3/2)*(A*b^2-a*(B*b-C*a))*arctanh(b^(1/2)*(c+d*tan(f*x+e)))^(1/2)/(-a*d+b*c)^(1/2)/(a^2+b^2)/(-a*d+b*c)^(5/2)/f+2/3*(A*d^2-B*c*d+C*c^2)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))^(3/2)+2*(b*(c^4*C-2*B*c^3*d+c^2*(3*A-C)*d^2+A*d^4)-a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))/(-a*d+b*c)^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1948 vs. $2(365) = 730$.

Time = 6.25 (sec) , antiderivative size = 1948, normalized size of antiderivative = 5.34

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^(5/2)),x]
```

output

```
(-2*(A*d^2 - c*(-(c*C) + B*d)))/(3*(-(b*c) + a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) - (2*((-2*((I*Sqrt[c - I*d]*((b*(-(b*c) + a*d)*((-3*c*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*d*(c^2*C - B*c*d + A*d^2))/2 - (3*d*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2))/2 + a*((-3*((b*d^2)/2 - (c*(-(b*c) + a*d))/2)*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2 - (a*d*((3*d*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*c*(c^2*C - B*c*d + A*d^2))/2))/2 - (b*((-3*d^2*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2 - c*((3*d*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*c*(c^2*C - B*c*d + A*d^2))/2))/2 - I*((a*(-(b*c) + a*d)*((-3*c*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*d*(c^2*C - B*c*d + A*d^2))/2 - (3*d*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2))/2 - b*((-3*((b*d^2)/2 - (c*(-(b*c) + a*d))/2)*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2 - (a*d*((3*d*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*c*(c^2*C - B*c*d + A*d^2))/2))/2 - (b*((-3*d^2*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2 - c*((3*d*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*c*(c^2*C - B*c*d + A*d^2))/2))/2))*ArcTanh[Sqrt[c + d*Tan[e + f*x]]/Sqrt[c - I*d]]/((-c + I*d)*f) - (I*Sqrt[c + I*d]*((b*(-(b*c) + a*d)*((-3*c*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*d*(c^2*C - B*c*d + A*d^2))/2 - (3*d*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2))/2 + a*((-3*((b*d^2)/2 - (c*(-(b*c) + a*d))/2)*(a*A*c*d - a*d*(c*C - B*d) - A*b*(c^2 + d^2)))/2 - (a*d*((3*d*(b*c - a*d)*(B*c - (A - C)*d))/2 - (3*b*c*...
```


Rubi [A] (warning: unable to verify)

Time = 5.92 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.22, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.383$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4132} \\
 & 2 \int \frac{-\frac{3(-b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2))}{2(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx}{\frac{3(c^2 + d^2)(bc - ad)}{2(Ad^2 - Bcd + c^2C)}} + \\
 & \quad \downarrow \text{27} \\
 & \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \\
 & \int \frac{-\frac{b(Cc^2 - Bdc + Ad^2) \tan^2(e + fx) - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx}{(c^2 + d^2)(bc - ad)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \\
 & \int \frac{-\frac{b(Cc^2 - Bdc + Ad^2) \tan(e + fx)^2 - (bc - ad)(Bc - (A - C)d) \tan(e + fx) + aAc d - ad(cC - Bd) - Ab(c^2 + d^2)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} dx}{(c^2 + d^2)(bc - ad)} \\
 & \quad \downarrow \text{4132}
 \end{aligned}$$

$$2 \int \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} \frac{(2c(A-C)d - B(c^2 - d^2)) \tan(e+fx)(bc-ad)^2 - b(b(Cc^4 - 2Bdc^3 + (3A-C)d^2c^2 + Ad^4) - ad^2(2c(A-C)d - B(c^2 - d^2))) \tan^2(e+fx) + A(2abdc^3 - b^2(c^2 + d^2))}{2(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} \frac{1}{(c^2 + d^2)(bc - ad)} dx$$

↓ 27

$$2 \int \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} \frac{(2c(A-C)d - B(c^2 - d^2)) \tan(e+fx)(bc-ad)^2 - b(b(Cc^4 - 2Bdc^3 + (3A-C)d^2c^2 + Ad^4) - ad^2(2c(A-C)d - B(c^2 - d^2))) \tan^2(e+fx) + A(2abdc^3 - b^2(c^2 + d^2))}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} \frac{1}{(c^2 + d^2)(bc - ad)} dx$$

↓ 3042

$$2 \int \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} \frac{(2c(A-C)d - B(c^2 - d^2)) \tan(e+fx)(bc-ad)^2 - b(b(Cc^4 - 2Bdc^3 + (3A-C)d^2c^2 + Ad^4) - ad^2(2c(A-C)d - B(c^2 - d^2))) \tan^2(e+fx) + A(2abdc^3 - b^2(c^2 + d^2))}{(a+b \tan(e+fx))\sqrt{c+d \tan(e+fx)}} \frac{1}{(c^2 + d^2)(bc - ad)} dx$$

↓ 4136

$$2 \int \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} \frac{(a(Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2)) + b(2c(A-C)d - B(c^2 - d^2)))(bc - ad)^2 + (2aAcd - 2acCd + Ab(c^2 - d^2) - aB(c^2 - d^2) - b(Cc^2 - 2Bdc - Cd^2)) \tan(e+fx)(bc - ad)}{\frac{\sqrt{c+d \tan(e+fx)}}{a^2 + b^2}} \frac{1}{(c^2 + d^2)(bc - ad)} dx$$

↓ 3042

$$2 \int \frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} \frac{(a(Cc^2 - 2Bdc - Cd^2 - A(c^2 - d^2)) + b(2c(A-C)d - B(c^2 - d^2)))(bc - ad)^2 + (2aAcd - 2acCd + Ab(c^2 - d^2) - aB(c^2 - d^2) - b(Cc^2 - 2Bdc - Cd^2)) \tan(e+fx)(bc - ad)}{\frac{\sqrt{c+d \tan(e+fx)}}{a^2 + b^2}} \frac{1}{(c^2 + d^2)(bc - ad)} dx$$

↓ 4022

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2}$$

$(c^2 + d^2)(bc - ad)$

↓ 3042

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2}$$

$(c^2 + d^2)(bc - ad)$

↓ 4020

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2}$$

↓ 25

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2}$$

$(c^2 + d^2)(bc - ad)$

↓ 73

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2}$$

$(c^2 + d^2)(bc - ad)$

↓ 221

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{\tan(e + fx)^2 + 1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{a^2 + b^2}$$

$(c^2 + d^2)(bc - ad)$

↓ 4117

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{1}{(a + b \tan(e + fx))\sqrt{c + d \tan(e + fx)}} dx}{f(a^2 + b^2)}$$

$(c^2 + d^2)(bc - ad)$

↓ 73

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{2b^2(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \int \frac{1}{\frac{a + b(c + d \tan(e + fx))}{d} - \frac{bc}{d}} d\sqrt{c + d \tan(e + fx)}}{df(a^2 + b^2)}$$

$(c^2 + d^2)(bc - ad)$

↓ 221

$$\frac{2(Ad^2 - Bcd + c^2C)}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2(b(c^2d^2(3A-C) + Ad^4 - 2Bc^3d + c^4C) - ad^2(2cd(A-C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} + \frac{2b^{3/2}(c^2 + d^2)^2(Ab^2 - a(bB - aC)) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{c + d \tan(e + fx)}}{\sqrt{bc - ad}}\right)}{f(a^2 + b^2)\sqrt{bc - ad}}$$

$(c^2 + d^2)(bc - ad)$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*(c + d*Tan[e + f*x])^(5/2)),x]
```

output
$$\frac{(2*(c^2*C - B*c*d + A*d^2))/(3*(b*c - a*d)*(c^2 + d^2)*f*(c + d*\text{Tan}[e + f*x])^{(3/2)}) - (((-(((a + I*b)*(A - I*B - C)*(c + I*d)^2*(b*c - a*d)^2*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]])/(\text{Sqrt}[c - I*d]*f)) - ((a - I*b)*(A + I*B - C)*(c - I*d)^2*(b*c - a*d)^2*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c + I*d]])/(\text{Sqrt}[c + I*d]*f)))/(a^2 + b^2) + (2*b^{(3/2)}*(A*b^2 - a*(b*B - a*C))*(c^2 + d^2)^2*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/\text{Sqrt}[b*c - a*d]])/((a^2 + b^2)*\text{Sqrt}[b*c - a*d]*f))/((b*c - a*d)*(c^2 + d^2)) - (2*(b*(c^4*C - 2*B*c^3*d + c^2*(3*A - C)*d^2 + A*d^4) - a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))/((b*c - a*d)*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]))/((b*c - a*d)*(c^2 + d^2))$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27
$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[a, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (b_)*(Gx_)] \text{ ; FreeQ}[b, \text{x}]$$

rule 73
$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_)*((c_.) + (d_.)*(x_)^{(n_)}, \text{x_Symbol}] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b)^n}, \text{x}], \text{x}, (a + b*x)^{(1/p)}, \text{x}]] \text{ ; FreeQ}[\{a, b, c, d\}, \text{x}] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, \text{x}]$$

rule 221
$$\text{Int}[(a_) + (b_.)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], \text{x}] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{NegQ}[a/b]$$

rule 3042
$$\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinearQ}[u, \text{x}]$$

rule 4020
$$\text{Int}[(a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^{(m_)*((c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]), \text{x_Symbol}] \rightarrow \text{Simp}[c*(d/f) \quad \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), \text{x}], \text{x}, d*\text{Tan}[e + f*x], \text{x}] \text{ ; FreeQ}[\{a, b, c, d, e, f, m\}, \text{x}] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$$

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4117

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)
*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 45118 vs. $2(328) = 656$.

Time = 0.28 (sec) , antiderivative size = 45119, normalized size of antiderivative = 123.61

method	result	size
derivativedivides	Expression too large to display	45119
default	Expression too large to display	45119

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))**(5/2),x)`

output

```
Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))**(5/2)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail
```

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")
```

output

```
Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```


Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))*(c + d*tan(e + f*x))^(5/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(5/2),x)
```

output

```
( - 4*sqrt(tan(e + f*x)*d + c)*a + 2*sqrt(tan(e + f*x)*d + c)*c - 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3 + a*c**3),x)*tan(e + f*x)**2*a**2*d**3*f + 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3 + a*c**3),x)*tan(e + f*x)**2*a*c*d**3*f - 6*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3 + a*c**3),x)*tan(e + f*x)*a**2*c*d**2*f + 6*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3 + a*c**3),x)*tan(e + f*x)*a*c**2*d**2*f - 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3 + a*c**3),x)*a**2*c**2*d*f + 3*int(sqrt(tan(e + f*x)*d + c)/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 ...
```

3.127
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^2(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1438
Mathematica [B] (verified)	1439
Rubi [A] (warning: unable to verify)	1439
Maple [B] (verified)	1447
Fricas [F(-1)]	1447
Sympy [F]	1448
Maxima [F(-2)]	1448
Giac [F(-2)]	1449
Mupad [F(-1)]	1449
Reduce [F]	1449

Optimal result

Integrand size = 47, antiderivative size = 679

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right) - (B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c+id}}\right)}{(a - ib)^2(c - id)^{5/2} f - (a + ib)^2(c + id)^{5/2} f}$$

$$- \frac{b^{3/2}(7a^3bBd - 5a^4Cd + b^4(2Bc - 5Ad) + ab^3(4Ac - 4cC + 3Bd) - a^2b^2(2Bc + (9A + C)d)) \operatorname{arctanh}\left(\frac{\sqrt{c+d \tan(e+fx)}}{\sqrt{c-id}}\right) + (a^2 + b^2)^2(bc - ad)^{7/2} f}{d(2b^2c(cC - Bd) - 3abB(c^2 + d^2) + a^2(5c^2C - 2Bcd + 3Cd^2) + A(2a^2d^2 + b^2(3c^2 + 5d^2))) - 3(a^2 + b^2)(bc - ad)^2(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}}$$

$$- \frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2} + d(2a^3d^2(Bc^2 + 2cCd - Bd^2) + 2b^3c(2c^3C - 3Bc^2d - Bd^3) - ab^2(Bc^4 - 4cCd^3 + 3Bd^4) + a^2b(5c^4C - 6c^3Cd - 3Bc^3d - Bd^4) - ab^2(5c^4C - 6c^3Cd - 3Bc^3d - Bd^4)) - (a^2 + b^2)(bc - ad)^3(c^2 + d^2)}$$

output

```

-(I*A+B-I*C)*arctanh((c+d*tan(f*x+e))^(1/2)/(c-I*d)^(1/2))/(a-I*b)^(2/(c-I*d)^(5/2)/f-(B-I*(A-C))*arctanh((c+d*tan(f*x+e))^(1/2)/(c+I*d)^(1/2))/(a+I*b)^(2/(c+I*d)^(5/2)/f-b^(3/2)*(7*a^3*b*B*d-5*a^4*C*d+b^4*(-5*A*d+2*B*c)+a*b^3*(4*A*c+3*B*d-4*C*c)-a^2*b^2*(2*B*c+(9*A+C)*d))*arctanh(b^(1/2)*(c+d*tan(f*x+e))^(1/2)/(-a*d+b*c)^(1/2))/(a^2+b^2)^(2/(-a*d+b*c)^(7/2)/f-1/3*d*(2*b^2*c*(-B*d+C*c)-3*a*b*B*(c^2+d^2)+a^2*(-2*B*c*d+5*C*c^2+3*C*d^2)+A*(2*a^2*d^2+b^2*(3*c^2+5*d^2)))/(a^2+b^2)/(-a*d+b*c)^(2/(c^2+d^2)/f/(c+d*tan(f*x+e))^(3/2)-(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))/(c+d*tan(f*x+e))^(3/2)-d*(2*a^3*d^2*(B*c^2-B*d^2+2*C*c*d)+2*b^3*c*(-3*B*c^2*d-B*d^3+2*C*c^3)-a*b^2*(B*c^4+3*B*d^4-4*C*c*d^3)+a^2*b*(-6*B*c^3*d-2*B*c*d^3+5*C*c^4+2*C*c^2*d^2+C*d^4)-A*(4*a^3*c*d^3+4*a*b^2*c*d^3-4*a^2*b*d^2*(2*c^2+d^2)-b^3*(c^4+10*c^2*d^2+5*d^4)))/(a^2+b^2)/(-a*d+b*c)^3/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 6052 vs. $2(679) = 1358$.

Time = 6.52 (sec) , antiderivative size = 6052, normalized size of antiderivative = 8.91

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \text{Result too large to show}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d*Tan[e + f*x])^(5/2)),x]

```

output

Result too large to show

Rubi [A] (warning: unable to verify)

Time = 11.40 (sec) , antiderivative size = 767, normalized size of antiderivative = 1.13, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.468$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed

below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx$$

↓ 4132

$$\int \frac{5Adb^2 + 5(Ab^2 - a(bB - aC))d \tan^2(e + fx) - 2aA(bc - ad) - (bB - aC)(2bc + 3ad) + 2(Ab - Cb - aB)(bc - ad) \tan(e + fx)}{2(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx$$

$$\frac{\frac{(a^2 + b^2)(bc - ad)}{Ab^2 - a(bB - aC)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{5Adb^2 + 5(Ab^2 - a(bB - aC))d \tan^2(e + fx) - 2aA(bc - ad) - (bB - aC)(2bc + 3ad) + 2(Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx$$

$$\frac{\frac{2(a^2 + b^2)(bc - ad)}{Ab^2 - a(bB - aC)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{5Adb^2 + 5(Ab^2 - a(bB - aC))d \tan(e + fx)^2 - 2aA(bc - ad) - (bB - aC)(2bc + 3ad) + 2(Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2}} dx$$

$$\frac{\frac{2(a^2 + b^2)(bc - ad)}{Ab^2 - a(bB - aC)}}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}}$$

↓ 4132

$$2 \int \frac{3(2d^2(Ac - Cc + Bd)a^3 - b(4A + C)d(c^2 + d^2)a^2 - b^2(2Cc^3 - Bdc^2 + 4Cd^2c - 3Bd^3 - 2A(c^3 + 2d^2c))a - bd(2Ad^2a^2 + (5Cc^2 - 2Bdc + 3Cd^2)a^2 - 3bB(c^2 + d^2))}{2(a + b \tan(e + fx))(c + d \tan(e + fx))^{5/2} \cdot 3(c^2 + d^2)(bc - ad)}$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+5c^2C+3Cd^2)-3abB(c^2+d^2)+Ab^2(3c^2+5d^2)+2b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d\tan(e+fx))^{3/2}} - \int \frac{2d^2(Ac-Cc+Bd)a^3-b(4A+C)d(c^2+d^2)a^2-b^2(2C}{$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b\tan(e + fx))(c + d\tan(e + fx))^{3/2}}$$

↓ 3042

$$\frac{2d(2a^2Ad^2+a^2(-2Bcd+5c^2C+3Cd^2)-3abB(c^2+d^2)+Ab^2(3c^2+5d^2)+2b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d\tan(e+fx))^{3/2}} - \int \frac{2d^2(Ac-Cc+Bd)a^3-b(4A+C)d(c^2+d^2)a^2-b^2(2C}{$$

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b\tan(e + fx))(c + d\tan(e + fx))^{3/2}}$$

↓ 4132

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b\tan(e + fx))(c + d\tan(e + fx))^{3/2}} -$$

$$2 \int \frac{2d^3(Cc^2-2Bdc-Cd^2-A(c^2-d^2))a^4-2bd^2(3Cc^3-4Bc}{$$

$$\frac{2d(2Ad^2a^2+(5Cc^2-2Bdc+3Cd^2)a^2-3bB(c^2+d^2)a+2b^2c(cC-Bd)+Ab^2(3c^2+5d^2))}{3(bc-ad)(c^2+d^2)f(c+d\tan(e+fx))^{3/2}} -$$

↓ 27

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b\tan(e + fx))(c + d\tan(e + fx))^{3/2}} -$$

$$\int \frac{2d^3(Cc^2-2Bdc-Cd^2-A(c^2-d^2))a^4-2bd^2(3Cc^3-4Bc}{$$

$$\frac{2d(2Ad^2a^2+(5Cc^2-2Bdc+3Cd^2)a^2-3bB(c^2+d^2)a+2b^2c(cC-Bd)+Ab^2(3c^2+5d^2))}{3(bc-ad)(c^2+d^2)f(c+d\tan(e+fx))^{3/2}} -$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b\tan(e + fx))(c + d\tan(e + fx))^{3/2}} -$$

$$\int \frac{2d^3(Cc^2-2Bdc-Cd^2-A(c^2-d^2))a^4-2bd^2(3Cc^3-4Bc}{$$

$$\frac{2d(2Ad^2a^2+(5Cc^2-2Bdc+3Cd^2)a^2-3bB(c^2+d^2)a+2b^2c(cC-Bd)+Ab^2(3c^2+5d^2))}{3(bc-ad)(c^2+d^2)f(c+d\tan(e+fx))^{3/2}} -$$

↓ 4136

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{b^2(-5Cda^4 + 7bBda^3 - b^2(2Bc + (9A + C)d)a^2 + b^3(4Ac - 4C))}{3(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}} - \frac{2d(2Ad^2a^2 + (5Cc^2 - 2Bdc + 3Cd^2)a^2 - 3bB(c^2 + d^2)a + 2b^2c(cC - Bd) + Ab^2(3c^2 + 5d^2))}{3(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{b^2(c^2 + d^2)^2(-5Cda^4 + 7bBda^3 - b^2(2Bc + (9A + C)d)a^2 + b^3(4Ac - 4C))}{3(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}} - \frac{2d(2Ad^2a^2 + (5Cc^2 - 2Bdc + 3Cd^2)a^2 - 3bB(c^2 + d^2)a + 2b^2c(cC - Bd) + Ab^2(3c^2 + 5d^2))}{3(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{b^2(c^2 + d^2)^2(-5Cda^4 + 7bBda^3 - b^2(2Bc + (9A + C)d)a^2 + b^3(4Ac - 4C))}{3(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}} - \frac{2d(2Ad^2a^2 + (5Cc^2 - 2Bdc + 3Cd^2)a^2 - 3bB(c^2 + d^2)a + 2b^2c(cC - Bd) + Ab^2(3c^2 + 5d^2))}{3(bc - ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}}$$

↓ 4022

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5Ccd^2))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 5c^2C + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5Ccd^2))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 5c^2C + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

↓ 4020

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5Bcd^2 - 2Bcd^2 + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

↓ 25

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5Bcd^2 - 2Bcd^2 + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

↓ 73

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5Bcd^2 - 2Bcd^2 + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

↓ 221

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5Bcd^2 - 2Bcd^2 + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

↓ 4117

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5C^2d^2))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 5c^2C + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

73

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5C^2d^2))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 5c^2C + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

221

$$\frac{Ab^2 - a(bB - aC)}{f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^3d^2(Bc^2 - Bd^2 + 2cCd) + a^2b(-6Bc^3d - 2Bcd^3 + 5C^2d^2))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} - \frac{2d(2a^2Ad^2 + a^2(-2Bcd + 5c^2C + 3Cd^2) - 3abB(c^2 + d^2) + Ab^2(3c^2 + 5d^2) + 2b^2c(cC - Bd))}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^2*(c + d
*Tan[e + f*x])^(5/2)),x]
```

output

$$\begin{aligned}
& -((A*b^2 - a*(b*B - a*C))/((a^2 + b^2)*(b*c - a*d)*f*(a + b*\text{Tan}[e + f*x])*(c + d*\text{Tan}[e + f*x])^{(3/2)})) - ((2*d*(2*a^2*A*d^2 + 2*b^2*c*(c*C - B*d) - 3*a*b*B*(c^2 + d^2) + A*b^2*(3*c^2 + 5*d^2) + a^2*(5*c^2*C - 2*B*c*d + 3*C*d^2)))/(3*(b*c - a*d)*(c^2 + d^2)*f*(c + d*\text{Tan}[e + f*x])^{(3/2)}) - (((-2*(-((a + I*b)^2*(A - I*B - C)*(c + I*d)^2*(b*c - a*d)^3*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c - I*d]])/(\text{Sqrt}[c - I*d]*f)) - ((a - I*b)^2*(A + I*B - C)*(c - I*d)^2*(b*c - a*d)^3*\text{ArcTan}[\text{Tan}[e + f*x]/\text{Sqrt}[c + I*d]])/(\text{Sqrt}[c + I*d]*f)))/(a^2 + b^2) - (2*b^{(3/2)}*(c^2 + d^2)^2*(7*a^3*b*B*d - 5*a^4*C*d + b^4*(2*B*c - 5*A*d) + a*b^3*(4*A*c - 4*c*C + 3*B*d) - a^2*b^2*(2*B*c + (9*A + C)*d))*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/\text{Sqrt}[b*c - a*d]])/((a^2 + b^2)*\text{Sqrt}[b*c - a*d]*f))/((b*c - a*d)*(c^2 + d^2)) - (2*d*(2*a^3*d^2*(B*c^2 + 2*c*C*d - B*d^2) + 2*b^3*c*(2*c^3*C - 3*B*c^2*d - B*d^3) - a*b^2*(B*c^4 - 4*c*C*d^3 + 3*B*d^4) + a^2*b*(5*c^4*C - 6*B*c^3*d + 2*c^2*C*d^2 - 2*B*c*d^3 + C*d^4) - A*(4*a^3*c*d^3 + 4*a*b^2*c*d^3 - 4*a^2*b*d^2*(2*c^2 + d^2) - b^3*(c^4 + 10*c^2*d^2 + 5*d^4)))/((b*c - a*d)*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]))/((b*c - a*d)*(c^2 + d^2)))/(2*(a^2 + b^2)*(b*c - a*d))
\end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 27

$$\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] \text{ ; FreeQ}[b, x]$$

rule 73

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)}*(c - a*(d/b) + d*(x^{(p/b)})^n), x], x, (a + b*x)^{(1/p)}], x]] \text{ ; FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 221

$$\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 67569 vs. 2(640) = 1280.

Time = 0.42 (sec) , antiderivative size = 67570, normalized size of antiderivative = 99.51

method	result	size
derivativedivides	Expression too large to display	67570
default	Expression too large to display	67570

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(5
/2),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+
e))^(5/2),x, algorithm="fricas")
```

output Timed out

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**2/(c+d*tan(f*x+e))**(5/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**2*(c + d*tan(e + f*x))**(5/2)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a*d-b*c>0)', see `assume?` for more detail`

Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:Unable to divide, perhaps due to rounding error%%{%%{-1,[6,22,10]%%}+%%{-10,[6,20,10]%%}+%%{-45,[6,18,10]%%}+%%`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^2*(c + d*tan(e + f*x))^(5/2)),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^2 (c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^2/(c+d*tan(f*x+e))^(5/2),x)`

output

```
( - 16*sqrt(tan(e + f*x)*d + c)*a*d - 4*sqrt(tan(e + f*x)*d + c)*b*c + 10*
sqrt(tan(e + f*x)*d + c)*c*d - 27*int(sqrt(tan(e + f*x)*d + c)/(3*tan(e +
f*x)**5*a*b**2*d**4 + 2*tan(e + f*x)**5*b**3*c*d**3 + 6*tan(e + f*x)**4*a*
*2*b*d**4 + 13*tan(e + f*x)**4*a*b**2*c*d**3 + 6*tan(e + f*x)**4*b**3*c**2
*d**2 + 3*tan(e + f*x)**3*a**3*d**4 + 20*tan(e + f*x)**3*a**2*b*c*d**3 + 2
1*tan(e + f*x)**3*a*b**2*c**2*d**2 + 6*tan(e + f*x)**3*b**3*c**3*d + 9*tan
(e + f*x)**2*a**3*c*d**3 + 24*tan(e + f*x)**2*a**2*b*c**2*d**2 + 15*tan(e
+ f*x)**2*a*b**2*c**3*d + 2*tan(e + f*x)**2*b**3*c**4 + 9*tan(e + f*x)*a**
3*c**2*d**2 + 12*tan(e + f*x)*a**2*b*c**3*d + 4*tan(e + f*x)*a*b**2*c**4 +
3*a**3*c**3*d + 2*a**2*b*c**4),x)*tan(e + f*x)**3*a**3*b*d**5*f - 54*int(
sqrt(tan(e + f*x)*d + c)/(3*tan(e + f*x)**5*a*b**2*d**4 + 2*tan(e + f*x)**
5*b**3*c*d**3 + 6*tan(e + f*x)**4*a**2*b*d**4 + 13*tan(e + f*x)**4*a*b**2*
c*d**3 + 6*tan(e + f*x)**4*b**3*c**2*d**2 + 3*tan(e + f*x)**3*a**3*d**4 +
20*tan(e + f*x)**3*a**2*b*c*d**3 + 21*tan(e + f*x)**3*a*b**2*c**2*d**2 + 6
*tan(e + f*x)**3*b**3*c**3*d + 9*tan(e + f*x)**2*a**3*c*d**3 + 24*tan(e +
f*x)**2*a**2*b*c**2*d**2 + 15*tan(e + f*x)**2*a*b**2*c**3*d + 2*tan(e + f*
x)**2*b**3*c**4 + 9*tan(e + f*x)*a**3*c**2*d**2 + 12*tan(e + f*x)*a**2*b*c
**3*d + 4*tan(e + f*x)*a*b**2*c**4 + 3*a**3*c**3*d + 2*a**2*b*c**4),x)*tan
(e + f*x)**3*a**2*b**2*c*d**4*f + 45*int(sqrt(tan(e + f*x)*d + c)/(3*tan(e
+ f*x)**5*a*b**2*d**4 + 2*tan(e + f*x)**5*b**3*c*d**3 + 6*tan(e + f*x)...
```

3.128 $\int (a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)) dx$

Optimal result	1451
Mathematica [A] (verified)	1452
Rubi [A] (verified)	1453
Maple [F(-1)]	1458
Fricas [F(-1)]	1459
Sympy [F]	1459
Maxima [F]	1460
Giac [F]	1460
Mupad [F(-1)]	1461
Reduce [F]	1461

Optimal result

Integrand size = 49, antiderivative size = 679

$$\begin{aligned}
 & \int (a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx) + C \tan^2(e+fx)) dx = \\
 & \frac{(a-ib)^{5/2}(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{f} \\
 & - \frac{(a+ib)^{5/2}(B-i(A-C))\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{f} \\
 & - \frac{(5a^4Cd^4 - 20a^3bd^3(cC + 2Bd) + 30a^2b^2d^2(c^2C - 4Bcd - 8(A-C)d^2) - 20ab^3d(c^3C - 2Bc^2d + 8c(A-C) - 64ad^2) + (64b(a^2B - b^2B + 2ab(A-C))d^3 - (bc-ad)(16b(Ab+aB-bC)d^2 + (bc-ad)(5bcC - 8bBd - 5aCd))\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{64bd^3f} \\
 & + \frac{(16b(Ab+aB-bC)d^2 + (bc-ad)(5bcC - 8bBd - 5aCd))\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{32d^3f} \\
 & - \frac{(5bcC - 8bBd - 5aCd)(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}}{24d^2f} \\
 & + \frac{C(a+b \tan(e+fx))^{5/2}(c+d \tan(e+fx))^{3/2}}{4df}
 \end{aligned}$$

output

```

-(a-I*b)^(5/2)*(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-(a+I*b)^(5/2)*(B-I*(A-C))*(c+I*d)^(1/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-1/64*(5*a^4*C*d^4-20*a^3*b*d^3*(2*B*d+C*c)+30*a^2*b^2*d^2*(c^2*C-4*B*c*d-8*(A-C)*d^2)-20*a*b^3*d*(c^3*C-2*B*c^2*d+8*c*(A-C)*d^2-16*B*d^3)+b^4*(5*c^4*C-8*B*c^3*d+16*c^2*(A-C)*d^2+64*B*c*d^3+128*(A-C)*d^4)*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(3/2)/d^(7/2)/f+1/64*(64*b*(B*a^2-B*b^2+2*a*b*(A-C))*d^3-(-a*d+b*c)*(16*b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-8*B*b*d-5*C*a*d+5*C*b*c)))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b/d^3/f+1/32*(16*b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-8*B*b*d-5*C*a*d+5*C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)/d^3/f-1/24*(-8*B*b*d-5*C*a*d+5*C*b*c)*(a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)/d^2/f+1/4*C*(a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(3/2)/d/f

```

Mathematica [A] (verified)

Time = 8.53 (sec) , antiderivative size = 1202, normalized size of antiderivative = 1.77

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```

Integrate[(a + b*Tan[e + f*x])^(5/2)*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^(5/2)*(c + d*Tan[e + f*x])^(3/2))/(4*d*f) + (((-5*
b*c*C + 8*b*B*d + 5*a*C*d)*(a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])
^(3/2))/(6*d*f) + ((3*(16*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(5*b*c*C -
8*b*B*d - 5*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/
(8*d*f) + (((24*b*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3 - (3*(b*c - a*d)*(16
*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(5*b*c*C - 8*b*B*d - 5*a*C*d)))/8)*
Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f) + ((24*b*d^3*(b*(
3*a^2*b*(A*c - c*C - B*d) - b^3*(A*c - c*C - B*d) + a^3*(B*c + (A - C)*d)
- 3*a*b^2*(B*c + (A - C)*d)) + Sqrt[-b^2]*(a^3*(A*c - c*C - B*d) - 3*a*b^2
*(A*c - c*C - B*d) - 3*a^2*b*(B*c + (A - C)*d) + b^3*(B*c + (A - C)*d))*A
rcTanh[(Sqrt[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + S
qrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (S
qrt[-b^2]*d)/b]) - (24*b*d^3*(b*(3*a^2*b*(A*c - c*C - B*d) - b^3*(A*c - c*
C - B*d) + a^3*(B*c + (A - C)*d) - 3*a*b^2*(B*c + (A - C)*d)) - Sqrt[-b^2]
*(a^3*(A*c - c*C - B*d) - 3*a*b^2*(A*c - c*C - B*d) - 3*a^2*b*(B*c + (A -
C)*d) + b^3*(B*c + (A - C)*d))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a
+ b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqr
t[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) - (3*Sqrt[b]*Sqrt[c - (a*d)/
b]*Sqrt[(c/(c - (a*d)/b) - (a*d)/(b*(c - (a*d)/b)))^(-1)]*Sqrt[c/(c - (a*d)
)/b] - (a*d)/(b*(c - (a*d)/b))]*(5*a^4*C*d^4 - 20*a^3*b*d^3*(c*C + 2*B*...
```

Rubi [A] (verified)

Time = 9.26 (sec) , antiderivative size = 707, normalized size of antiderivative = 1.04, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.327$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \frac{\int -\frac{1}{2}(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} ((5bcC - 5adC - 8bBd) \tan^2(e + fx) - 8(Ab - Cb + aB)d \tan(e + fx) + 4d)}{4df} \\
 & \quad \downarrow 27 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}}{4df} - \frac{\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} ((5bcC - 5adC - 8bBd) \tan^2(e + fx) - 8(Ab - Cb + aB)d \tan(e + fx) + 8d)}{8d} \\
 & \quad \downarrow 3042 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}}{4df} - \frac{\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} ((5bcC - 5adC - 8bBd) \tan(e + fx)^2 - 8(Ab - Cb + aB)d \tan(e + fx) + 8d)}{8d} \\
 & \quad \downarrow 4130 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}}{4df} - \frac{\int -\frac{3}{2} \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (c(5cC - 8Bd)b^2 - 2ad(5cC + 4Bd)b + a^2(16A - 11C)d^2 + (16b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 8bBd) + 3d))}{3d}}{8d} \\
 & \quad \downarrow 27 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}}{4df} - \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} - \int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (c(5cC - 8Bd)b^2 - 2ad(5cC + 4Bd)b + a^2(16A - 11C)d^2 + (16b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 8bBd) + 3d))}{3df}}{8d} \\
 & \quad \downarrow 3042 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}}{4df} - \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} - \int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (c(5cC - 8Bd)b^2 - 2ad(5cC + 4Bd)b + a^2(16A - 11C)d^2 + (16b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 8bBd) + 3d))}{3df}}{8d} \\
 & \quad \downarrow 4130 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}}{4df} - \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} - \int \sqrt{c + d \tan(e + fx)} (-c(5Cc^2 - 8Bdc + 16(A - C)d^2)b^3 + ad(15Cc^2 - 32Bdc - 48(A - C)d^2 + (bc - ad)(5bcC - 5adC - 8bBd) + 3d))}{3df}}{8d}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} - \\ & \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\int \frac{\sqrt{c+d \tan(e+fx)}(-c(5Cc^2 - 8Bdc + 16(A-C)d^2)b^3 + ad(15Cc^2 - 32Bdc - 48(A-C)d^2))}{\sqrt{c+d \tan(e+fx)}} dx}{\sqrt{c+d \tan(e+fx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} - \\ & \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\int \frac{\sqrt{c+d \tan(e+fx)}(-c(5Cc^2 - 8Bdc + 16(A-C)d^2)b^3 + ad(15Cc^2 - 32Bdc - 48(A-C)d^2))}{\sqrt{c+d \tan(e+fx)}} dx}{\sqrt{c+d \tan(e+fx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 4130 \\ & \frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} - \\ & \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\int \frac{c(5Cc^3 - 8Bdc^2 + 16(A-C)d^2c - 64Bd^3)b^4 - 4ad(5Cc^3 - 10Bdc^2 - 56(A-C)d^2c)}{\sqrt{c+d \tan(e+fx)}} dx}{\sqrt{c+d \tan(e+fx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} - \\ & \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\int \frac{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(64bd^3(a^2B + 2ab(A-C) - b^2B) - (bc - ad)(16bd^2))}{\sqrt{c+d \tan(e+fx)}} dx}{\sqrt{a+b \tan(e+fx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} - \\ & \frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\int \frac{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(64bd^3(a^2B + 2ab(A-C) - b^2B) - (bc - ad)(16bd^2))}{\sqrt{c+d \tan(e+fx)}} dx}{\sqrt{a+b \tan(e+fx)}} \end{aligned}$$

$$\downarrow 4138$$

$$\frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} -$$

$$\frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (64bd^3 (a^2 B + 2ab(A - C) - b^2 B) - (bc - ad)(16bd^2 - 2df))}{bf}$$

↓ 2348

$$\frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} -$$

$$\frac{(5bcC - 5adC - 8bBd)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{(16b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 8bBd)) \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}}{2df}$$

↓ 2009

$$\frac{C(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}}{4df} -$$

$$\frac{(-5aCd - 8bBd + 5bcC)(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (16bd^2 (aB + Ab - bC) + (bc - ad)(-5aCd - 8bBd + 5bcC))}{2df}$$

input

```
Int[(a + b*Tan[e + f*x])^(5/2)*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x]
] + C*Tan[e + f*x]^2),x]
```

output

```
(C*(a + b*Tan[e + f*x])^(5/2)*(c + d*Tan[e + f*x])^(3/2))/(4*d*f) - (((5*b*c*C - 8*b*B*d - 5*a*C*d)*(a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(3/2))/(3*d*f) - (((16*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(5*b*c*C - 8*b*B*d - 5*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(2*d*f) + (-1/2*(128*(a - I*b)^(5/2)*b*(B + I*(A - C))*Sqrt[c - I*d]*d^3*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])] - 128*(a + I*b)^(5/2)*b*(I*A - B - I*C)*Sqrt[c + I*d]*d^3*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])] + (2*(5*a^4*C*d^4 - 20*a^3*b*d^3*(c*C + 2*B*d) + 30*a^2*b^2*d^2*(c^2*C - 4*B*c*d - 8*(A - C)*d^2) - 20*a*b^3*d*(c^3*C - 2*B*c^2*d + 8*c*(A - C)*d^2 - 16*B*d^3) + b^4*(5*c^4*C - 8*B*c^3*d + 16*c^2*(A - C)*d^2 + 64*B*c*d^3 + 128*(A - C)*d^4))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[b]*Sqrt[d]))/(b*f) + ((64*b*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3 - (b*c - a*d)*(16*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(5*b*c*C - 8*b*B*d - 5*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f))/(4*d))/(2*d))/(8*d)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int (a + b \tan(fx + e))^{5/2} \sqrt{c + d \tan(fx + e)} (A + B \tan(fx + e) + C \tan(fx + e)^2) dx$$

input

```
int((a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

output

```
int((a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

Fricas [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*
tan(f*x+e)^2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input

```
integrate((a+b*tan(f*x+e))**(5/2)*(c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+
C*tan(f*x+e)**2),x)
```

output

```
Integral((a + b*tan(e + f*x))**(5/2)*sqrt(c + d*tan(e + f*x))*(A + B*tan(e
+ f*x) + C*tan(e + f*x)**2), x)
```


Maxima [F]

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^{5/2} \sqrt{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(5/2)*sqrt(d*tan(f*x + e) + c), x)
```

Giac [F]

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^{5/2} \sqrt{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(5/2)*sqrt(d*tan(f*x + e) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (C \tan(e + fx)^2 + B \tan(e + fx) + A$$

input

```
int((a + b*tan(e + f*x))^(5/2)*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)
```

output

```
int((a + b*tan(e + f*x))^(5/2)*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)
```

Reduce [F]

$$\int (a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^4 dx \right) b^2 c$$

$$+ 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) abc$$

$$+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) b^3$$

$$+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) a^2 c$$

$$+ 3 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) a b^2$$

$$+ 3 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) a^2 b$$

$$+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) a^3$$

input

```
int((a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)
```

output

```
int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4,x)*b
**2*c + 2*int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*
x)**3,x)*a*b*c + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan
(e + f*x)**3,x)*b**3 + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b +
a)*tan(e + f*x)**2,x)*a**2*c + 3*int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*tan(e + f*x)**2,x)*a*b**2 + 3*int(sqrt(tan(e + f*x)*d + c)*sq
rt(tan(e + f*x)*b + a)*tan(e + f*x),x)*a**2*b + int(sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a),x)*a**3
```

3.129 $\int (a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx)) dx$

Optimal result	1463
Mathematica [A] (verified)	1464
Rubi [A] (verified)	1465
Maple [F(-1)]	1469
Fricas [F(-1)]	1470
Sympy [F]	1470
Maxima [F]	1471
Giac [F]	1471
Mupad [F(-1)]	1472
Reduce [F]	1472

Optimal result

Integrand size = 49, antiderivative size = 505

$$\begin{aligned}
 & \int (a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)} (A+B \tan(e+fx) + C \tan^2(e+fx)) dx = \\
 & \frac{(a-ib)^{3/2}(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{f} \\
 & + \frac{(a+ib)^{3/2}(iA-B-iC)\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{f} \\
 & - \frac{(a^3Cd^3 - 3a^2bd^2(cC + 2Bd) + 3ab^2d(c^2C - 4Bcd - 8(A - C)d^2) - b^3(c^3C - 2Bc^2d + 8c(A - C)d^2 - 1)}{8b^{3/2}d^{5/2}f}}{f} \\
 & + \frac{(8b(Ab + aB - bC)d^2 + (bc - ad)(bcC - 2bBd - aCd)) \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}}{8bd^2f} \\
 & - \frac{(bcC - 2bBd - aCd)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{4d^2f} \\
 & + \frac{C(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}}{3df}
 \end{aligned}$$

output

```

-(a-I*b)^(3/2)*(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f+(a+I*b)^(3/2)*(I*A-B-I*C)*(c+I*d)^(1/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-1/8*(a^3*C*d^3-3*a^2*b*d^2*(2*B*d+C*c)+3*a*b^2*d*(c^2*C-4*B*c*d-8*(A-C)*d^2)-b^3*(c^3*C-2*B*c^2*d+8*c*(A-C)*d^2-16*B*d^3))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(3/2)/d^(5/2)/f+1/8*(8*b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-2*B*b*d-C*a*d+C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b/d^2/f-1/4*(-2*B*b*d-C*a*d+C*b*c)*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)/d^2/f+1/3*C*(a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)/d/f

```

Mathematica [A] (verified)

Time = 7.89 (sec) , antiderivative size = 835, normalized size of antiderivative = 1.65

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{C(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}}{3df}$$

$$+ \frac{-3(bcC - 2bBd - aCd) \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}}{4df} + \frac{3(8b(Ab + aB - bC)d^2 + (bc - ad)(bcC - 2bBd - aCd)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{4bf} +$$

input

```

Integrate[(a + b*Tan[e + f*x])^(3/2)*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(3/2))/(3*d*f) + ((-3*(
b*c*C - 2*b*B*d - a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/
2))/(4*d*f) + ((3*(8*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(b*c*C - 2*b*B*
d - a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(4*b*f) + (
(6*b*d^2*(Sqrt[-b^2]*(a^2*(A*c - c*C - B*d) - b^2*(A*c - c*C - B*d) - 2*a*
b*(B*c + (A - C)*d)) + b*(2*a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d)
- b^2*(B*c + (A - C)*d))*ArcTanh[(Sqrt[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*
Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[-a
+ Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) + (6*b*d^2*(Sqrt[-b^2]*(a^2*(A
*c - c*C - B*d) - b^2*(A*c - c*C - B*d) - 2*a*b*(B*c + (A - C)*d) - b*(2*
a*b*(A*c - c*C - B*d) + a^2*(B*c + (A - C)*d) - b^2*(B*c + (A - C)*d))*Ar
cTanh[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt
[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-
b^2]*d)/b]) - (3*Sqrt[b]*Sqrt[c - (a*d)/b])*(a^3*C*d^3 - 3*a^2*b*d^2*(c*C +
2*B*d) + 3*a*b^2*d*(c^2*C - 4*B*c*d - 8*(A - C)*d^2) - b^3*(c^3*C - 2*B*c
^2*d + 8*c*(A - C)*d^2 - 16*B*d^3))*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*
x]])/(Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*c + b*d*Tan[e + f*x])/(b*c - a*d
)]/(4*Sqrt[d]*Sqrt[c + d*Tan[e + f*x]])/(b^2*f))/(2*d))/(3*d)
```

Rubi [A] (verified)

Time = 5.93 (sec) , antiderivative size = 523, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \frac{\int -\frac{3}{2}\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}((bcC-adC-2bBd)\tan^2(e+fx)-2(Ab-Cb+aB)d\tan(e+fx))}{\frac{3d}{3df}C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{3/2}}{3df} - \\
 & \frac{\int \sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}((bcC-adC-2bBd)\tan^2(e+fx)-2(Ab-Cb+aB)d\tan(e+fx))}{2d} \\
 & \quad \downarrow 3042 \\
 & \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{3/2}}{3df} - \\
 & \frac{\int \sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}((bcC-adC-2bBd)\tan(e+fx)^2-2(Ab-Cb+aB)d\tan(e+fx))}{2d} \\
 & \quad \downarrow 4130 \\
 & \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{3/2}}{3df} - \\
 & \frac{\int -\frac{\sqrt{c+d\tan(e+fx)}(c(cC-2Bd)b^2-2ad(cC+3Bd)b+a^2(8A-7C)d^2+(8b(Ab-Cb+aB)d^2+(bc-ad)(bcC-adC-2bBd)\tan^2(e+fx)+8(Ba^2+2b(A-C)a-b^2B))}{2\sqrt{a+b\tan(e+fx)}}}{2d}}{2d} \\
 & \quad \downarrow 27 \\
 & \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{3/2}}{3df} - \\
 & \frac{(-aCd-2bBd+bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \frac{\int \frac{\sqrt{c+d\tan(e+fx)}(c(cC-2Bd)b^2-2ad(cC+3Bd)b+a^2(8A-7C)d^2+(8b(Ab-Cb+aB)d^2+(bc-ad)(bcC-adC-2bBd)\tan^2(e+fx)+8(Ba^2+2b(A-C)a-b^2B))}{\sqrt{a+b\tan(e+fx)}}}{4d}}{2d} \\
 & \quad \downarrow 3042 \\
 & \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{3/2}}{3df} - \\
 & \frac{(-aCd-2bBd+bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \frac{\int \frac{\sqrt{c+d\tan(e+fx)}(c(cC-2Bd)b^2-2ad(cC+3Bd)b+a^2(8A-7C)d^2+(8b(Ab-Cb+aB)d^2+(bc-ad)(bcC-adC-2bBd)\tan^2(e+fx)+8(Ba^2+2b(A-C)a-b^2B))}{\sqrt{a+b\tan(e+fx)}}}{4d}}{2d} \\
 & \quad \downarrow 4130
 \end{aligned}$$

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{-c(Cc^2 - 2Bdc - 8(A-C)d^2)b^3 + ad(3Cc^2 + 20Bdc + 8(A-C)d^2)b^2 - a^2d^2(16Ac - 13Cc)}{f} - \frac{(-aCd - 2bBd + bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df}$$

27

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{(-aCd - 2bBd + bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df} - \frac{\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}(8bd^2(aB + Ab - bC) + (bc - ad)(-aCd - 2bBd + bcC))}{bf}$$

3042

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{(-aCd - 2bBd + bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df} - \frac{\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}(8bd^2(aB + Ab - bC) + (bc - ad)(-aCd - 2bBd + bcC))}{bf}$$

4138

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{(-aCd - 2bBd + bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df} - \frac{\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}(8bd^2(aB + Ab - bC) + (bc - ad)(-aCd - 2bBd + bcC))}{bf}$$

2348

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{(bcC - adC - 2bBd)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df} - \frac{(8b(Ab - Cb + aB)d^2 + (bc - ad)(bcC - adC - 2bBd))\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}}{bf}$$

2009

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}}{3df} - \frac{(-aCd - 2bBd + bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df} - \frac{\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}(8bd^2(aB + Ab - bC) + (bc - ad)(-aCd - 2bBd + bcC))}{bf}$$

input $\text{Int}[(a + b \cdot \tan[e + f \cdot x])^{3/2} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]} \cdot (A + B \cdot \tan[e + f \cdot x] + C \cdot \tan[e + f \cdot x]^2), x]$

output $(C \cdot (a + b \cdot \tan[e + f \cdot x])^{3/2} \cdot (c + d \cdot \tan[e + f \cdot x])^{3/2}) / (3 \cdot d \cdot f) - ((b \cdot c \cdot C - 2 \cdot b \cdot B \cdot d - a \cdot C \cdot d) \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]} \cdot (c + d \cdot \tan[e + f \cdot x])^{3/2}) / (2 \cdot d \cdot f) - (-1/2 \cdot (16 \cdot (a - I \cdot b)^{3/2} \cdot b \cdot (B + I \cdot (A - C)) \cdot \sqrt{c - I \cdot d} \cdot d^2 \cdot \text{ArcTanh}[(\sqrt{c - I \cdot d} \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]}) / (\sqrt{a - I \cdot b} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]})]) - 16 \cdot (a + I \cdot b)^{3/2} \cdot b \cdot (I \cdot A - B - I \cdot C) \cdot \sqrt{c + I \cdot d} \cdot d^2 \cdot \text{ArcTanh}[(\sqrt{c + I \cdot d} \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]}) / (\sqrt{a + I \cdot b} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]})]) + (2 \cdot (a^3 \cdot C \cdot d^3 - 3 \cdot a^2 \cdot b \cdot d^2 \cdot (c \cdot C + 2 \cdot B \cdot d) + 3 \cdot a \cdot b^2 \cdot d \cdot (c^2 \cdot C - 4 \cdot B \cdot c \cdot d - 8 \cdot (A - C) \cdot d^2) - b^3 \cdot (c^3 \cdot C - 2 \cdot B \cdot c^2 \cdot d + 8 \cdot c \cdot (A - C) \cdot d^2 - 16 \cdot B \cdot d^3)) \cdot \text{ArcTanh}[(\sqrt{d} \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]}) / (\sqrt{b} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]})]) / (\sqrt{b} \cdot \sqrt{d})) / (b \cdot f) + ((8 \cdot b \cdot (A \cdot b + a \cdot B - b \cdot C) \cdot d^2 + (b \cdot c - a \cdot d) \cdot (b \cdot c \cdot C - 2 \cdot b \cdot B \cdot d - a \cdot C \cdot d)) \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]}) / (b \cdot f)) / (4 \cdot d)) / (2 \cdot d)$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)(F_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F, (b_)(G_)] /; \text{FreeQ}[b, x]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2348 $\text{Int}[(P_)((c_)+(d_)(x_))^{(m_)}((e_)+(f_)(x_))^{(n_)}((a_)+(b_)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P \cdot (c + d \cdot x)^m \cdot (e + f \cdot x)^n \cdot (a + b \cdot x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2 \cdot p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2], x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2], x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int (a + b \tan(fx + e))^{\frac{3}{2}} \sqrt{c + d \tan(fx + e)} (A + B \tan(fx + e) + C \tan(fx + e)^2) dx$$

input

```
int((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

output

```
int((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

Fricas [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*
tan(f*x+e)^2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input

```
integrate((a+b*tan(f*x+e))**(3/2)*(c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+
C*tan(f*x+e)**2),x)
```

output

```
Integral((a + b*tan(e + f*x))**(3/2)*sqrt(c + d*tan(e + f*x))*(A + B*tan(e
+ f*x) + C*tan(e + f*x)**2), x)
```

Maxima [F]

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) (b \tan(fx + e) + a)^{3/2} \sqrt{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)*sqrt(d*tan(f*x + e) + c), x)
```

Giac [F]

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) (b \tan(fx + e) + a)^{3/2} \sqrt{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)*sqrt(d*tan(f*x + e) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (C \tan(e + fx)^2 + B \tan(e + fx) + A)$$

input `int((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `int((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\begin{aligned} & \int (a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) \\ & + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) bc \\ & + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) ac \\ & + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) b^2 \\ & + 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) ab \\ & + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) a^2 \end{aligned}$$

input `int((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output

```
int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3,x)*b
*c + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2
,x)*a*c + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*
x)**2,x)*b**2 + 2*int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*ta
n(e + f*x),x)*a*b + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a),
x)*a**2
```

3.130 $\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$

Optimal result	1474
Mathematica [A] (verified)	1475
Rubi [A] (verified)	1476
Maple [F(-1)]	1479
Fricas [B] (verification not implemented)	1480
Sympy [F]	1480
Maxima [F]	1481
Giac [F]	1481
Mupad [F(-1)]	1482
Reduce [F]	1482

Optimal result

Integrand size = 49, antiderivative size = 381

$$\begin{aligned}
 & \int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 = & -\frac{\sqrt{a - ib}(iA + B - iC)\sqrt{c - id} \operatorname{arctanh}\left(\frac{\sqrt{c - id}\sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib}\sqrt{c + d \tan(e + fx)}}\right)}{f} \\
 & -\frac{\sqrt{a + ib}(B - i(A - C))\sqrt{c + id} \operatorname{arctanh}\left(\frac{\sqrt{c + id}\sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib}\sqrt{c + d \tan(e + fx)}}\right)}{f} \\
 & -\frac{(a^2Cd^2 - 2abd(cC + 2Bd) + b^2(c^2C - 4Bcd - 8(A - C)d^2)) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a + b \tan(e + fx)}}{\sqrt{b}\sqrt{c + d \tan(e + fx)}}\right)}{4b^{3/2}d^{3/2}f} \\
 & -\frac{(bcC - 4bBd - aCd)\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}}{4bdf} \\
 & +\frac{C\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df}
 \end{aligned}$$

output

```

-(a-I*b)^(1/2)*(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*
x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-(a+I*b)^(1/2)*(B-I*(A-
C))*(c+I*d)^(1/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/
2)/(c+d*tan(f*x+e))^(1/2))/f-1/4*(a^2*C*d^2-2*a*b*d*(2*B*d+C*c)+b^2*(c^2*C
-4*B*c*d-8*(A-C)*d^2))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d
*tan(f*x+e))^(1/2))/b^(3/2)/d^(3/2)/f-1/4*(-4*B*b*d-C*a*d+C*b*c)*(a+b*tan(
f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b/d/f+1/2*C*(a+b*tan(f*x+e))^(1/2)*(c
+d*tan(f*x+e))^(3/2)/d/f
    
```

Mathematica [A] (verified)

Time = 7.14 (sec) , antiderivative size = 619, normalized size of antiderivative = 1.62

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{C \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}}{2df}$$

$$+ \frac{(-bcC + 4bBd + aCd) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{2bf} + \frac{2bd(b(ABC + aBc - bcC + aAd - bBd - aCd) - \sqrt{-b^2}(bBc + b(A - C)d - a(Ac - cC - Bd))) \sqrt{-a + \sqrt{-b^2}} \sqrt{-c + \frac{\sqrt{-b^2}d}{b}}}{\dots}$$

input

```

Integrate[Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e +
f*x] + C*Tan[e + f*x]^2),x]
    
```


output

```
(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(2*d*f) + (((-(b*c
*C) + 4*b*B*d + a*C*d)*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/
(2*b*f) + ((2*b*d*(b*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d) - Sqr
t[-b^2]*(b*B*c + b*(A - C)*d - a*(A*c - c*C - B*d)))*ArcTanh[(Sqrt[-c + (S
qrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c +
d*Tan[e + f*x]])])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) - (
2*b*d*(b*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d) + Sqrt[-b^2]*(b*B
*c + b*(A - C)*d - a*(A*c - c*C - B*d)))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*d)/
b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]
]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) - (Sqrt[b]*Sqrt[c -
(a*d)/b]*(a^2*C*d^2 - 2*a*b*d*(c*C + 2*B*d) + b^2*(c^2*C - 4*B*c*d - 8*(A
- C)*d^2))*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c - (
a*d)/b])]*Sqrt[(b*c + b*d*Tan[e + f*x])/(b*c - a*d)]/(2*Sqrt[d]*Sqrt[c +
d*Tan[e + f*x]))/(b^2*f))/(2*d)
```

Rubi [A] (verified)

Time = 3.56 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.204$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) \, dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan(e + fx)^2) \, dx \\
 & \quad \downarrow \text{4130} \\
 & \int -\frac{\sqrt{c + d \tan(e + fx)} ((bcC - adC - 4bBd) \tan^2(e + fx) - 4(Ab - Cb + aB)d \tan(e + fx) + bcC - a(4A - 3C)d)}{2\sqrt{a + b \tan(e + fx)}} \, dx + \\
 & \quad \frac{2d}{2df} \frac{C\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2df} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \int \frac{\sqrt{c+d\tan(e+fx)}((bcC-adC-4bBd)\tan^2(e+fx)-4(Ab-Cb+aB)d\tan(e+fx)+bcC-a(4A-3C)d)}{\sqrt{a+b\tan(e+fx)}} dx$$

4d

↓ 3042

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \int \frac{\sqrt{c+d\tan(e+fx)}((bcC-adC-4bBd)\tan(e+fx)^2-4(Ab-Cb+aB)d\tan(e+fx)+bcC-a(4A-3C)d)}{\sqrt{a+b\tan(e+fx)}} dx$$

4d

↓ 4130

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \int \frac{c(cC+4Bd)b^2-2ad(4Ac-3Cc-2Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd)\tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC-4bBd))\tan^2(e+fx)}{2\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx$$

4d

↓ 27

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \int \frac{c(cC+4Bd)b^2-2ad(4Ac-3Cc-2Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd)\tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC-4bBd))\tan^2(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx$$

4d

↓ 3042

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \int \frac{c(cC+4Bd)b^2-2ad(4Ac-3Cc-2Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd)\tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC-4bBd))\tan(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx$$

4d

↓ 4138

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \int \frac{c(cC+4Bd)b^2-2ad(4Ac-3Cc-2Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd)\tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC-4bBd))\tan^2(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(\tan^2(e+fx)+1)} dx$$

4d

↓ 2348

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \frac{(-aCd-4bBd+bcC)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{bf} + \frac{\int\left(\frac{-8Ad^2b^2+8Cd^2b^2+c^2Cb^2-4Bcdb^2-4aBd^2b-2acCdb+a^2Cd^2}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} + \frac{-8Bd^2b^2+8Acdb^2-8c}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{b}\sqrt{d}}$$

↓ 2009

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2df} - \frac{(-aCd-4bBd+bcC)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2(a^2Cd^2-2abd(2Bd+cC)+b^2(-8d^2(A-C)-4Bcd+c^2C))\operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b\tan(e+fx)}}{\sqrt{b}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{b}\sqrt{d}}$$

4d

input `Int[Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]`

output `(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(2*d*f) - ((8*Sqrt[a - I*b]*b*(B + I*(A - C))*Sqrt[c - I*d]*d*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])] - 8*Sqrt[a + I*b]*b*(I*A - B - I*C)*Sqrt[c + I*d]*d*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])] + (2*(a^2*C*d^2 - 2*a*b*d*(c*C + 2*B*d) + b^2*(c^2*C - 4*B*c*d - 8*(A - C)*d^2))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[b]*Sqrt[d]))/(2*b*f) + ((b*c*C - 4*b*B*d - a*C*d)*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f))/(4*d)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_
)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^
n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P
x, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) &&
!(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan (fx + e)} \sqrt{c + d \tan (fx + e)} (A + B \tan (fx + e) + C \tan (fx + e)^2) dx$$

input `int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34031 vs. $2(308) = 616$.

Time = 165.29 (sec) , antiderivative size = 68078, normalized size of antiderivative = 178.68

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**(1/2)*(c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output

```
Integral(sqrt(a + b*tan(e + f*x))*sqrt(c + d*tan(e + f*x))*(A + B*tan(e +
f*x) + C*tan(e + f*x)**2), x)
```

Maxima [F]

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a} \sqrt{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*
tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)
*sqrt(d*tan(f*x + e) + c), x)
```

Giac [F]

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a} \sqrt{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*
tan(f*x+e)^2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)
*sqrt(d*tan(f*x + e) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Hanged

input `int((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) c$$

$$+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) b$$

$$+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) a$$

input `int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2,x)*c + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x),x)*b + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a),x)*a`

3.131
$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$$

Optimal result	1483
Mathematica [A] (verified)	1484
Rubi [A] (verified)	1484
Maple [F(-1)]	1487
Fricas [B] (verification not implemented)	1488
Sympy [F]	1488
Maxima [F]	1489
Giac [F]	1489
Mupad [F(-1)]	1490
Reduce [F]	1490

Optimal result

Integrand size = 49, antiderivative size = 289

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$$

$$= -\frac{(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a-ib}f}$$

$$+ \frac{(iA-B-iC)\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a+ib}f}$$

$$+ \frac{(bcC+2bBd-aCd)\operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{b^{3/2}\sqrt{d}f}$$

$$+ \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf}$$

output

```
-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(1/2)/f+(I*A-B-I*C)*(c+I*d)^(1/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a+I*b)^(1/2)/f+(2*B*b*d-C*a*d+C*b*c)*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(3/2)/d^(1/2)/f+C*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b/f
```


Mathematica [A] (verified)

Time = 2.89 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.53

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

$$= \frac{b(bBc + b(A - C)d + \sqrt{-b^2}(Ac - cC - Bd)) \operatorname{arctanh}\left(\frac{\sqrt{-c + \frac{\sqrt{-b^2}d}{b}} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{-a + \sqrt{-b^2}} \sqrt{-c + \frac{\sqrt{-b^2}d}{b}}} + \frac{b(\sqrt{-b^2}(Ac - cC - Bd) - b(Bc + (A - C)d)) \operatorname{arctanh}\left(\frac{\sqrt{-c + \frac{\sqrt{-b^2}d}{b}} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a + \sqrt{-b^2}} \sqrt{c + \frac{\sqrt{-b^2}d}{b}}}$$

input `Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[a + b*Tan[e + f*x]],x]`

output `((b*(b*B*c + b*(A - C)*d + Sqrt[-b^2]*(A*c - c*C - B*d))*ArcTanh[(Sqrt[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) + (b*(Sqrt[-b^2]*(A*c - c*C - B*d) - b*(B*c + (A - C)*d))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) + b*C*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]] + (Sqrt[b]*Sqrt[c - (a*d)/b]*(b*c*C + 2*b*B*d - a*C*d)*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)])/(Sqrt[d]*Sqrt[c + d*Tan[e + f*x]])/(b^2*f)`

Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

$$\begin{aligned}
 & \int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan(e+fx)^2)}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(bcC-adC+2bBd) \tan^2(e+fx)+2b(Bc+(A-C)d) \tan(e+fx)+2Abc-C(bc+ad)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx \\
 & \quad \downarrow \text{4130} \\
 & \frac{b}{bf} + \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(bcC-adC+2bBd) \tan^2(e+fx)+2b(Bc+(A-C)d) \tan(e+fx)+2Abc-C(bc+ad)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2b}{bf} + \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf} \\
 & \quad \downarrow \text{4138} \\
 & \int \frac{(bcC-adC+2bBd) \tan^2(e+fx)+2b(Bc+(A-C)d) \tan(e+fx)+2Abc-C(bc+ad)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(\tan^2(e+fx)+1)} d \tan(e+fx) \\
 & \quad \downarrow \text{2348} \\
 & \frac{2bf}{bf} + \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf} + \\
 & \int \left(\frac{bcC-adC+2bBd}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} + \frac{-2bBc-2Abd+2bCd+i(2Abc-2bCc-2bBd)}{2(i-\tan(e+fx))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} + \frac{2bBc+2Abd-2bCd+i(2Abc-2bCc-2bBd)}{2(\tan(e+fx)+i)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{C\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2b\sqrt{c-id}(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b\tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{a-ib}} - \frac{2b\sqrt{c+id}(B-i(A-C))\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b\tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{a+ib}} + \frac{2(-aCd+2bBd+...)}{2bf}$$

input

```
Int[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[a + b*Tan[e + f*x]],x]
```

output

```
((-2*b*(I*A + B - I*C)*Sqrt[c - I*d]*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b] - (2*b*(B - I*(A - C))*Sqrt[c + I*d]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] + (2*(b*c*C + 2*b*B*d - a*C*d)*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[b]*Sqrt[d])/(2*b*f) + (C*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(fx + e)} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{\sqrt{a + b \tan(fx + e)}} dx$$

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(1/2),x)
```

output

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(1/2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39018 vs. $2(226) = 452$.

Time = 103.58 (sec) , antiderivative size = 78051, normalized size of antiderivative = 270.07

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Too large to display}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="fricas")
```

output

Too large to include

Sympy [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

$$= \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

input

```
integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(1/2),x)
```

output

```
Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(a + b*tan(e + f*x)), x)
```

Maxima [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{d \tan(fx + e) + c}}{\sqrt{b \tan(fx + e) + a}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(d*tan(f*x + e) + c) /sqrt(b*tan(f*x + e) + a), x)
```

Giac [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{d \tan(fx + e) + c}}{\sqrt{b \tan(fx + e) + a}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(d*tan(f*x + e) + c) /sqrt(b*tan(f*x + e) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Hanged}$$

input `int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(1/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\begin{aligned} & \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx \\ &= \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) c \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) b \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a}}{\tan(fx + e) b + a} dx \right) a \end{aligned}$$

input `int(((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x)`

output `int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*c + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*b + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)*b + a),x)*a`

3.132
$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$$

Optimal result	1491
Mathematica [A] (verified)	1492
Rubi [A] (verified)	1492
Maple [F(-1)]	1495
Fricas [F(-1)]	1496
Sympy [F]	1496
Maxima [F(-1)]	1496
Giac [F]	1497
Mupad [F(-1)]	1497
Reduce [F]	1497

Optimal result

Integrand size = 49, antiderivative size = 300

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx =$$

$$-\frac{(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{3/2}f}$$

$$-\frac{(B-i(A-C))\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{3/2}f}$$

$$+\frac{2C\sqrt{d}\operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{b^{3/2}f}-\frac{2(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}}{b(a^2+b^2)f\sqrt{a+b \tan(e+fx)}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(3/2)/f-(B-I*(A-C))*(c+I*d)^(1
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(3/2)/f+2*C*d^(1/2)*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(
1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(3/2)/f-2*(A*b^2-a*(B*b-C*a))*(c+d
*tan(f*x+e))^(1/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^(1/2)
    
```


Mathematica [A] (verified)

Time = 3.64 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.37

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx = \frac{(iA+B-iC)\sqrt{-c+id} \operatorname{arctanh}\left(\frac{\sqrt{-c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{-a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(-a+ib)^{3/2}}$$

input `Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(3/2),x]`

output `(((-I*A + B - I*C)*Sqrt[-c + I*d]*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(-a + I*b)^(3/2) + (I*(A + I*B - C)*Sqrt[c + I*d]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((a + I*b)^(3/2) + ((B + I*(A - C))*Sqrt[c + d*Tan[e + f*x]])/((a - I*b)*Sqrt[a + b*Tan[e + f*x]]) + ((-I)*A + B + I*C)*Sqrt[c + d*Tan[e + f*x]])/((a + I*b)*Sqrt[a + b*Tan[e + f*x]]) + (2*C*(-((b*(c + d*Tan[e + f*x]))/Sqrt[a + b*Tan[e + f*x]]) + Sqrt[d]*Sqrt[b*c - a*d]*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/Sqrt[b*c - a*d]])*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d))]/(b^2*Sqrt[c + d*Tan[e + f*x]]))/f`

Rubi [A] (verified)

Time = 2.45 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4128, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan(e+fx)^2)}{(a+b \tan(e+fx))^{3/2}} dx$$

$$\begin{aligned}
 & \downarrow 4128 \\
 & \frac{2 \int \frac{(a^2+b^2)Cd \tan^2(e+fx) - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd)}{2\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{b(a^2+b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}} \\
 & \frac{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}}{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}} \\
 & \downarrow 27 \\
 & \frac{\int \frac{(a^2+b^2)Cd \tan^2(e+fx) - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{b(a^2+b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}} \\
 & \frac{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}}{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}} \\
 & \downarrow 3042 \\
 & \frac{\int \frac{(a^2+b^2)Cd \tan(e+fx)^2 - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{b(a^2+b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}} \\
 & \frac{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}}{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}} \\
 & \downarrow 4138 \\
 & \frac{\int \frac{(a^2+b^2)Cd \tan^2(e+fx) - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)} (\tan^2(e+fx)+1)} d \tan(e+fx)}{\frac{bf(a^2+b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}} \\
 & \frac{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}}{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}} \\
 & \downarrow 2348 \\
 & \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}} + \\
 & \frac{\int \left(\frac{(a^2+b^2)Cd}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} + \frac{Acb^2 - cCb^2 - Bdb^2 - aBcb - aAdb + aCdb + i(Bcb^2 + Adb^2 - Cdb^2 + aAcb - acCb - aBdb)}{2(i - \tan(e+fx)) \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} + \frac{-Acb^2 + cCb^2}{bf(a^2+b^2)} \right)}{bf(a^2+b^2)} \\
 & \downarrow 2009 \\
 & \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{bf(a^2+b^2) \sqrt{a+b \tan(e+fx)}} + \\
 & \frac{2C\sqrt{d}(a^2+b^2) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{b}} - \frac{b(a+ib)\sqrt{c-id}(iA+B-iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a-ib}} + \frac{b(b+ia)\sqrt{c+id}(A+iB-C)}{bf(a^2+b^2)}
 \end{aligned}$$

input $\text{Int}[(\text{Sqrt}[c + d*\text{Tan}[e + f*x]]*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2))/(a + b*\text{Tan}[e + f*x])^{3/2}, x]$

output $(-(((a + I*b)*b*(I*A + B - I*C)*\text{Sqrt}[c - I*d]*\text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/ \text{Sqrt}[a - I*b]) + (b*(I*a + b)*(A + I*B - C)*\text{Sqrt}[c + I*d]*\text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/ \text{Sqrt}[a + I*b] + (2*(a^2 + b^2)*C*\text{Sqrt}[d]*\text{ArcTanh}[(\text{Sqrt}[d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/ \text{Sqrt}[b]) / (b*(a^2 + b^2)*f) - (2*(A*b^2 - a*(b*B - a*C))*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]) / (b*(a^2 + b^2)*f*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2348 $\text{Int}[(P_x)*((c_) + (d_)*(x_))^{(m_)}*((e_) + (f_)*(x_))^{(n_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P_x, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan (fx + e)} (A + B \tan (fx + e) + C \tan (fx + e)^2)}{(a + b \tan (fx + e))^{\frac{3}{2}}} dx$$

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(3/2),x)
```

output

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(3/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(3/2),x)
```

output

```
Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x))**(3/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e))}{(b \tan(fx + e))^{3/2}} dx$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(d*tan(f*x + e) + c)/(b*tan(f*x + e) + a)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input `int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(3/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x)`

output

```
(2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b*d + int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*
b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b*c*d*f - int((sqrt(ta
n(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*
**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*b**2*c**2*f + int((sq
rt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e +
f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*c*d*f - int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*b
**2 + 2*tan(e + f*x)*a*b + a**2),x)*a*b*c**2*f + int((sqrt(tan(e + f*x)*d
+ c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b**2*d + 2*ta
n(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan
(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*a*b**2*c*d*f - int((sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b**2*
d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d
+ 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*b**3*c**2*f + int((sqrt(
tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**
3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)
*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a**2*b*c*d*f - int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b**
2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a...
```

3.133
$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$$

Optimal result	1499
Mathematica [A] (verified)	1500
Rubi [A] (verified)	1500
Maple [F(-1)]	1505
Fricas [F(-1)]	1505
Sympy [F]	1506
Maxima [F(-2)]	1506
Giac [F]	1506
Mupad [F(-1)]	1507
Reduce [F]	1507

Optimal result

Integrand size = 49, antiderivative size = 370

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx =$$

$$\frac{(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{5/2}f}$$

$$-\frac{(B-i(A-C))\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{5/2}f}$$

$$-\frac{2(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}}{3b(a^2+b^2)f(a+b \tan(e+fx))^{3/2}}$$

$$-\frac{2(2a^3bBd+a^4Cd+b^4(3Bc+Ad)+2ab^3(3Ac-3cC-2Bd)-a^2b^2(3Bc+5Ad-7Cd))\sqrt{c+d \tan(e+fx)}}{3b(a^2+b^2)^2(bc-ad)f\sqrt{a+b \tan(e+fx)}}$$

output

```
-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(5/2)/f-(B-I*(A-C))*(c+I*d)^(1/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a+I*b)^(5/2)/f-2/3*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(1/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^(3/2)-2/3*(2*a^3*b*B*d+a^4*C*d+b^4*(A*d+3*B*c)+2*a*b^3*(3*A*c-2*B*d-3*C*c)-a^2*b^2*(5*A*d+3*B*c-7*C*d))*(c+d*tan(f*x+e))^(1/2)/b/(a^2+b^2)^2/(-a*d+b*c)/f/(a+b*tan(f*x+e))^(1/2)
```


Mathematica [A] (verified)

Time = 6.28 (sec) , antiderivative size = 424, normalized size of antiderivative = 1.15

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \frac{-\frac{3C\sqrt{c+d \tan(e+fx)}}{(a+b \tan(e+fx))^{3/2}} + \frac{(-2Ab^2+2abB+a^2C+3)}{(a^2+b^2)(a+b \tan(e+fx))}}{(a+b \tan(e+fx))^{3/2}}$$

input

```
Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2
))/(a + b*Tan[e + f*x])^(5/2),x]
```

output

```
((-3*C*Sqrt[c + d*Tan[e + f*x]])/(a + b*Tan[e + f*x])^(3/2) + ((-2*A*b^2 +
2*a*b*B + a^2*C + 3*b^2*C)*Sqrt[c + d*Tan[e + f*x]]/((a^2 + b^2)*(a + b*
Tan[e + f*x])^(3/2)) + (-3*b*((a + I*b)^2*(I*A + B - I*C)*Sqrt[-c + I*d]*
ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c +
d*Tan[e + f*x]])])/Sqrt[-a + I*b] + ((a - I*b)^2*((-I)*A + B + I*C)*Sqrt[
c + I*d]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*S
qrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b]) + (2*(2*a^3*b*B*d + a^4*C*d + b^
4*(3*B*c + A*d) + 2*a*b^3*(3*A*c - 3*c*C - 2*B*d) + a^2*b^2*(-3*B*c - 5*A*
d + 7*C*d))*Sqrt[c + d*Tan[e + f*x]]/((-b*c) + a*d)*Sqrt[a + b*Tan[e + f
*x]]))/(a^2 + b^2)^2)/(3*b*f)
```

Rubi [A] (verified)

Time = 4.21 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.19, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$, Rules used = {3042, 4128, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$\frac{2 \int \frac{-((-Ca^2 - 2bBa + 2Ab^2 - 3b^2C)d \tan^2(e+fx)) - 3b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(3bc-ad) + Ab(3ac+bd)}{2(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}} dx}{\frac{3b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}} \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{3bf(a^2 + b^2)(a+b \tan(e+fx))^{3/2}}}$$

↓ 27

$$\frac{\int \frac{-((-Ca^2 - 2bBa + 2Ab^2 - 3b^2C)d \tan^2(e+fx)) - 3b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(3bc-ad) + Ab(3ac+bd)}{(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}} dx}{\frac{3b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}} \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{3bf(a^2 + b^2)(a+b \tan(e+fx))^{3/2}}}$$

↓ 3042

$$\frac{\int \frac{-((-Ca^2 - 2bBa + 2Ab^2 - 3b^2C)d \tan^2(e+fx)^2) - 3b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(3bc-ad) + Ab(3ac+bd)}{(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}} dx}{\frac{3b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}} \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{3bf(a^2 + b^2)(a+b \tan(e+fx))^{3/2}}}$$

↓ 4132

$$\frac{2 \int \frac{3(b(bc-ad))((Ac-Cc-Bd)a^2 + 2b(Bc+(A-C)d)a - b^2(Ac-Cc-Bd)) - b(bc-ad)((Bc+(A-C)d)a^2) + 2b(Ac-Cc-Bd)a + b^2(Bc+(A-C)d) \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{3b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}} \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{3bf(a^2 + b^2)(a+b \tan(e+fx))^{3/2}}}$$

↓ 27

$$\frac{3 \int \frac{b(bc-ad)((Ac-Cc-Bd)a^2 + 2b(Bc+(A-C)d)a - b^2(Ac-Cc-Bd)) - b(bc-ad)((Bc+(A-C)d)a^2) + 2b(Ac-Cc-Bd)a + b^2(Bc+(A-C)d) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{3b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}} \frac{2(Ab^2 - a(bB - aC)) \sqrt{c+d \tan(e+fx)}}{3bf(a^2 + b^2)(a+b \tan(e+fx))^{3/2}}}$$

↓ 3042

$$3 \int \frac{b(bc-ad)((Ac-Cc-Bd)a^2+2b(Bc+(A-C)d)a-b^2(Ac-Cc-Bd))-b(bc-ad)(-(Bc+(A-C)d)a^2)+2b(Ac-Cc-Bd)a+b^2(Bc+(A-C)d)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \tan(e+fx) dx$$

$3b(a^2 + b^2)$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}}$$

↓ 4099

$$-\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}} +$$

$$-\frac{2\sqrt{c+d \tan(e+fx)}(a^4Cd+2a^3bBd-a^2b^2(5Ad+3Bc-7Cd)+2ab^3(3Ac-2Bd-3cC)+b^4(Ad+3Bc))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}} + \frac{3\left(\frac{1}{2}b(a-ib)^2(c+id)(A+iB-C)(bc-ad)\right)}{3b(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 3042

$$-\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}} +$$

$$-\frac{2\sqrt{c+d \tan(e+fx)}(a^4Cd+2a^3bBd-a^2b^2(5Ad+3Bc-7Cd)+2ab^3(3Ac-2Bd-3cC)+b^4(Ad+3Bc))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}} + \frac{3\left(\frac{1}{2}b(a-ib)^2(c+id)(A+iB-C)(bc-ad)\right)}{3b(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 4098

$$-\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}} +$$

$$-\frac{2\sqrt{c+d \tan(e+fx)}(a^4Cd+2a^3bBd-a^2b^2(5Ad+3Bc-7Cd)+2ab^3(3Ac-2Bd-3cC)+b^4(Ad+3Bc))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}} + \frac{3\left(\frac{b(a-ib)^2(c+id)(A+iB-C)(bc-ad)}{f}\right)}{3b(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 104

$$-\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}} +$$

$$-\frac{2\sqrt{c+d \tan(e+fx)}(a^4Cd+2a^3bBd-a^2b^2(5Ad+3Bc-7Cd)+2ab^3(3Ac-2Bd-3cC)+b^4(Ad+3Bc))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}} + \frac{3\left(\frac{b(a-ib)^2(c+id)(A+iB-C)(bc-ad)}{f}\right)}{3b(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 221

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}} + \frac{-2\sqrt{c+d \tan(e+fx)}(a^4Cd+2a^3bBd-a^2b^2(5Ad+3Bc-7Cd)+2ab^3(3Ac-2Bd-3cC)+b^4(Ad+3Bc))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}} + \frac{3 \left(\frac{ib(a-ib)^2 \sqrt{c+id}(A+iB-C)(bc-ad)}{f\sqrt{a}} \right)}{3b(a^2 + b^2)}$$

input `Int[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(5/2),x]`

output `(-2*(A*b^2 - a*(b*B - a*C))*Sqrt[c + d*Tan[e + f*x]]/(3*b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^(3/2)) + ((3*((-I)*(a + I*b)^2*b*(A - I*B - C)*Sqrt[c - I*d]*(b*c - a*d)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a - I*b]*f) + (I*(a - I*b)^2*b*(A + I*B - C)*Sqrt[c + I*d]*(b*c - a*d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*f)))/((a^2 + b^2)*(b*c - a*d)) - (2*(2*a^3*b*B*d + a^4*C*d + b^4*(3*B*c + A*d) + 2*a*b^3*(3*A*c - 3*c*C - 2*B*d) - a^2*b^2*(3*B*c + 5*A*d - 7*C*d))*Sqrt[c + d*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e + f*x]]))/(3*b*(a^2 + b^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 221 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan (fx + e)} (A + B \tan (fx + e) + C \tan (fx + e)^2)}{(a + b \tan (fx + e))^{\frac{5}{2}}} dx$$

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(5/2),x)
```

output

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(5/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan (e + fx)} (A + B \tan (e + fx) + C \tan^2 (e + fx))}{(a + b \tan (e + fx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(5/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx$$

input `integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**(5/2),x)`

output `Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(((2*b*d+2*a*c)^2>0)', see `assume?` for more)`

Giac [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan^2(fx + e) + B \tan(fx + e) + A) \sqrt{c + d \tan(fx + e)}}{(b \tan(fx + e) + a)^{5/2}} dx$$

input `integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x, algorithm="giac")`

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(d*tan(f*x + e) + c)
/(b*tan(f*x + e) + a)^(5/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(
a + b*tan(e + f*x))^(5/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(5/2),x)
```


output

```
(2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*c*d**2
+ 4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**2*d
**2 - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b*c
**2*d + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b*d**2 + 2*s
qrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*c**2*d - 2*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b**2*c*d - 2*sqrt(tan(e + f*x)*d + c
)*sqrt(tan(e + f*x)*b + a)*b*c**3 + 3*int((sqrt(tan(e + f*x)*d + c)*sqrt(t
an(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**3*d + 3*tan(e + f*x)*
**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d + 3*tan(
e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*c + a**
3*c),x)*tan(e + f*x)**2*a**3*b**2*d**3*f - 3*int((sqrt(tan(e + f*x)*d + c)
*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**3*d + 3*tan(e
+ f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b*d +
3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**2*b*
c + a**3*c),x)*tan(e + f*x)**2*a**2*b**3*c*d**2*f - 3*int((sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**3*d +
3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a
**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x
)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a**2*b**2*c*d**3*f - 3*int((sqrt(t
an(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)...
```

3.134
$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$$

Optimal result	1509
Mathematica [A] (verified)	1510
Rubi [A] (verified)	1511
Maple [F(-1)]	1517
Fricas [F(-1)]	1517
Sympy [F]	1517
Maxima [F(-1)]	1518
Giac [F]	1518
Mupad [F(-1)]	1519
Reduce [F]	1519

Optimal result

Integrand size = 49, antiderivative size = 597

$$\int \frac{\sqrt{c+d \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx =$$

$$-\frac{(iA+B-iC)\sqrt{c-id}\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{7/2}f}$$

$$-\frac{(B-i(A-C))\sqrt{c+id}\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{7/2}f}$$

$$-\frac{2(Ab^2-a(bB-aC))\sqrt{c+d \tan(e+fx)}}{5b(a^2+b^2)f(a+b \tan(e+fx))^{5/2}}$$

$$-\frac{2(4a^3bBd+a^4Cd+b^4(5Bc+Ad)+2ab^3(5Ac-5cC-3Bd)-a^2b^2(5Bc+9Ad-11Cd))\sqrt{c+d \tan(e+fx)}}{15b(a^2+b^2)^2(bc-ad)f(a+b \tan(e+fx))^{3/2}}$$

$$+\frac{2(8a^5bBd^2+2a^6Cd^2-a^4b^2d(25Bc+33Ad-39Cd)-a^2b^4(45Ac^2-45c^2C-90Bcd-29Ad^2+23Cd^2))\sqrt{c+d \tan(e+fx)}}{15b(a^2+b^2)^2(bc-ad)f(a+b \tan(e+fx))^{3/2}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(1/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(7/2)/f-(B-I*(A-C))*(c+I*d)^(1
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(7/2)/f-2/5*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(1/2
)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^(5/2)-2/15*(4*a^3*b*B*d+a^4*C*d+b^4*(A*d+
5*B*c)+2*a*b^3*(5*A*c-3*B*d-5*C*c)-a^2*b^2*(9*A*d+5*B*c-11*C*d))*(c+d*tan(
f*x+e))^(1/2)/b/(a^2+b^2)^2/(-a*d+b*c)/f/(a+b*tan(f*x+e))^(3/2)+2/15*(8*a^
5*b*B*d^2+2*a^6*C*d^2-a^4*b^2*d*(33*A*d+25*B*c-39*C*d)-a^2*b^4*(45*A*c^2-2
9*A*d^2-90*B*c*d-45*C*c^2+23*C*d^2)+a^3*b^3*(80*c*(A-C)*d+B*(15*c^2-49*d^2
))-a*b^5*(40*c*(A-C)*d+B*(45*c^2-3*d^2))-b^6*(5*c*(B*d+3*C*c)-A*(15*c^2+2*
d^2))*(c+d*tan(f*x+e))^(1/2)/b/(a^2+b^2)^3/(-a*d+b*c)^2/f/(a+b*tan(f*x+e)
)^(1/2)
    
```

Mathematica [A] (verified)

Time = 6.93 (sec) , antiderivative size = 1109, normalized size of antiderivative = 1.86

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx =$$

$$\frac{C \sqrt{c + d \tan(e + fx)}}{2bf(a + b \tan(e + fx))^{5/2}}$$

$$2 \left[\frac{2(b^2(bc-ad)(a^2Cd+b^2(5Bc+Ad)+ab^2))}{\dots} \right]$$

$$\frac{2(\frac{1}{2}b^2(-4Abc+5bcC-aCd)-a(-2b^2(Bc+(A-C)d)-\frac{1}{2}a(bcC-4bBd-aCd))\sqrt{c+d \tan(e+fx)}}{5(a^2+b^2)(bc-ad)f(a+b \tan(e+fx))^{5/2}}}$$

input

```

Integrate[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2
))/ (a + b*Tan[e + f*x])^(7/2), x]
    
```

output

```

-1/2*(C*Sqrt[c + d*Tan[e + f*x]])/(b*f*(a + b*Tan[e + f*x])^(5/2)) - ((-2*
((b^2*(-4*A*b*c + 5*b*c*C - a*C*d))/2 - a*(-2*b^2*(B*c + (A - C)*d) - (a*(
b*c*C - 4*b*B*d - a*C*d))/2))*Sqrt[c + d*Tan[e + f*x]])/(5*(a^2 + b^2)*(b*
c - a*d)*f*(a + b*Tan[e + f*x])^(5/2)) - (2*((-2*(b^2*(b*c - a*d)*(a^2*C*d
+ b^2*(5*B*c + A*d) + a*b*(5*A*c - 5*c*C - B*d)) - a*(a*(4*A*b^2 - 4*a*b*
B - a^2*C - 5*b^2*C)*d*(b*c - a*d) - 5*b^2*(b*c - a*d)*(A*b*c - a*B*c - b*
c*C - a*A*d - b*B*d + a*C*d)))*Sqrt[c + d*Tan[e + f*x]])/(3*(a^2 + b^2)*(b
*c - a*d)*f*(a + b*Tan[e + f*x])^(3/2)) - (2*((-15*b*(b*c - a*d)^2*((I*a
- b)^3*(A - I*B - C)*Sqrt[-c + I*d]*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan
[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[-a + I*b] - (
(I*a + b)^3*(A + I*B - C)*Sqrt[c + I*d]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*
Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b]))/
(2*(a^2 + b^2)*f) - (2*(b^2*((b*c - a*d)*(b^2*d - (3*a*(b*c - a*d))/2)*(a^
2*C*d + b^2*(5*B*c + A*d) + a*b*(5*A*c - 5*c*C - B*d)) + ((-3*b*c)/2 + (a*
d)/2)*(a*(4*A*b^2 - 4*a*b*B - a^2*C - 5*b^2*C)*d*(b*c - a*d) - 5*b^2*(b*c
- a*d)*(A*b*c - a*B*c - b*c*C - a*A*d - b*B*d + a*C*d))) - a*((3*b*(b*c -
a*d)*(b*(4*A*b^2 - 4*a*b*B - a^2*C - 5*b^2*C)*d*(b*c - a*d) + 5*a*b*(b*c -
a*d)*(A*b*c - a*B*c - b*c*C - a*A*d - b*B*d + a*C*d) + b*(b*c - a*d)*(a^2
*C*d + b^2*(5*B*c + A*d) + a*b*(5*A*c - 5*c*C - B*d)))/2 - a*d*(b^2*(b*c
- a*d)*(a^2*C*d + b^2*(5*B*c + A*d) + a*b*(5*A*c - 5*c*C - B*d)) - a*(a...

```

Rubi [A] (verified)

Time = 7.46 (sec) , antiderivative size = 693, normalized size of antiderivative = 1.16, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx)^2)}{(a + b \tan(e + fx))^{7/2}} dx$$

↓ 4128

$$2 \int \frac{-((-Ca^2 - 4bBa + 4Ab^2 - 5b^2C)d \tan^2(e+fx)) - 5b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(5bc-ad) + Ab(5ac+bd)}{2(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{5b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \frac{1}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}}$$

↓ 27

$$\int \frac{-((-Ca^2 - 4bBa + 4Ab^2 - 5b^2C)d \tan^2(e+fx)) - 5b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(5bc-ad) + Ab(5ac+bd)}{(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{5b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \frac{1}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}}$$

↓ 3042

$$\int \frac{-((-Ca^2 - 4bBa + 4Ab^2 - 5b^2C)d \tan^2(e+fx)) - 5b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(5bc-ad) + Ab(5ac+bd)}{(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{5b(a^2 + b^2)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \frac{1}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}}$$

↓ 4132

$$2 \int \frac{2d(Cda^4 + 4bBda^3 - b^2(5Bc + 9Ad - 11Cd)a^2 + 2b^3(5Ac - 5Cc - 3Bd)a + b^4(5Bc + Ad)) \tan^2(e+fx) + 15b(bc-ad) \left(-((Bc + (A-C)d)a^2) + 2b(Ac - Cc - Bd)a + b^2 \right)}{2(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}}$$

↓ 27

$$\int \frac{2d(Cda^4 + 4bBda^3 - b^2(5Bc + 9Ad - 11Cd)a^2 + 2b^3(5Ac - 5Cc - 3Bd)a + b^4(5Bc + Ad)) \tan^2(e+fx) + 15b(bc-ad) \left(-((Bc + (A-C)d)a^2) + 2b(Ac - Cc - Bd)a + b^2 \right)}{(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}}$$

↓ 3042

$$\int \frac{2d(Cda^4+4bBda^3-b^2(5Bc+9Ad-11Cd)a^2+2b^3(5Ac-5Cc-3Bd)a+b^4(5Bc+Ad)) \tan(e+fx)^2+15b(bc-ad) \left(-((Bc+(A-C)d)a^2)+2b(Ac-Cc-Bd)a+b^2 \right)}{(a+b \tan(e+fx))^5} \frac{1}{3(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 4132

$$2 \int \frac{15(b(bc-ad)^2((Ac-Cc-Bd)a^3+3b(Bc+(A-C)d)a^2-3b^2(Ac-Cc-Bd)a-b^3(Bc+(A-C)d))-b(bc-ad)^2 \left(-((Bc+(A-C)d)a^3)+3b(Ac-Cc-Bd)a^2+3b^2 \right))}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \frac{1}{(a^2+b^2)(bc-ad)}$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 27

$$15 \int \frac{b(bc-ad)^2((Ac-Cc-Bd)a^3+3b(Bc+(A-C)d)a^2-3b^2(Ac-Cc-Bd)a-b^3(Bc+(A-C)d))-b(bc-ad)^2 \left(-((Bc+(A-C)d)a^3)+3b(Ac-Cc-Bd)a^2+3b^2 \right)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \frac{1}{(a^2+b^2)(bc-ad)}$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 3042

$$15 \int \frac{b(bc-ad)^2((Ac-Cc-Bd)a^3+3b(Bc+(A-C)d)a^2-3b^2(Ac-Cc-Bd)a-b^3(Bc+(A-C)d))-b(bc-ad)^2 \left(-((Bc+(A-C)d)a^3)+3b(Ac-Cc-Bd)a^2+3b^2 \right)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \frac{1}{(a^2+b^2)(bc-ad)}$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 4099

$$-\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}} +$$

$$-\frac{2\sqrt{c+d \tan(e+fx)}(a^4Cd+4a^3bBd-a^2b^2(9Ad+5Bc-11Cd)+2ab^3(5Ac-3Bd-5Cc)+b^4(Ad+5Bc))}{3f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))^{3/2}} - \frac{2\sqrt{c+d \tan(e+fx)}(2a^6Cd^2+8a^5bBd^2)}{3f(a^2+b^2)(bc-ad)(a+b \tan(e+fx))^{3/2}}$$

↓ 3042

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c + d \tan(e + fx)}(2a^6Cd^2 + 8a^5bBd^2)}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 4098

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c + d \tan(e + fx)}(2a^6Cd^2 + 8a^5bBd^2)}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 104

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c + d \tan(e + fx)}(2a^6Cd^2 + 8a^5bBd^2)}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 221

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{5bf(a^2 + b^2)(a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c + d \tan(e + fx)}(2a^6Cd^2 + 8a^5bBd^2)}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

input

```
Int[(Sqrt[c + d*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(7/2),x]
```

output

$$\begin{aligned} & (-2*(A*b^2 - a*(b*B - a*C))*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/(5*b*(a^2 + b^2)*f*(\\ & a + b*\text{Tan}[e + f*x])^{(5/2)}) + ((-2*(4*a^3*b*B*d + a^4*C*d + b^4*(5*B*c + A* \\ & d) + 2*a*b^3*(5*A*c - 5*c*C - 3*B*d) - a^2*b^2*(5*B*c + 9*A*d - 11*C*d))*\text{S} \\ & \text{qrt}[c + d*\text{Tan}[e + f*x]])/(3*(a^2 + b^2)*(b*c - a*d)*f*(a + b*\text{Tan}[e + f*x]) \\ & ^{(3/2)}) - ((-15*(((-I)*(a + I*b)^3*b*(A - I*B - C))*\text{Sqrt}[c - I*d]*(b*c - a* \\ & d)^2*\text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a - I*b]*\text{Sqrt}[\\ & c + d*\text{Tan}[e + f*x]])])]/(\text{Sqrt}[a - I*b]*f) + (I*(a - I*b)^3*b*(A + I*B - C)* \\ & \text{Sqrt}[c + I*d]*(b*c - a*d)^2*\text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x] \\ &])/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])]/(\text{Sqrt}[a + I*b]*f)))/((a^2 + \\ & b^2)*(b*c - a*d)) - (2*(8*a^5*b*B*d^2 + 2*a^6*C*d^2 - a^4*b^2*d*(25*B*c + \\ & 33*A*d - 39*C*d) - a^2*b^4*(45*A*c^2 - 45*c^2*C - 90*B*c*d - 29*A*d^2 + 23 \\ & *C*d^2) + a^3*b^3*(80*c*(A - C)*d + B*(15*c^2 - 49*d^2)) - a*b^5*(40*c*(A \\ & - C)*d + B*(45*c^2 - 3*d^2)) - b^6*(5*c*(3*c*C + B*d) - A*(15*c^2 + 2*d^2) \\ &))*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f*\text{Sqrt}[a + b*\text{Tan}[e + \\ & f*x]])/(3*(a^2 + b^2)*(b*c - a*d))/(5*b*(a^2 + b^2)) \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$$

rule 104

$$\text{Int}[(((a_.) + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)})/((e_.) + (f_.)*(x_)), x_] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Simp}[q \text{ Subst}[\text{Int}[x^{(q*(m+1) - 1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$$

rule 221

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4098

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

rule 4099

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
(!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(fx + e)} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(a + b \tan(fx + e))^{\frac{7}{2}}} dx$$

input

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(7/2),x)
```

output

```
int((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(7/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(7/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \int \frac{\sqrt{c + d \tan(e + fx)} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*ta
n(f*x+e))**(7/2),x)
```

output

```
Integral(sqrt(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)
/(a + b*tan(e + f*x))**(7/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(7/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e))}{(b \tan(fx + e))^{7/2}}$$

input

```
integrate((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(7/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(d*tan(f*x + e) + c)
/(b*tan(f*x + e) + a)^(7/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(7/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{\sqrt{c + d \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{too large to display}$$

input

```
int(((c+d*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x)
```

output

```
(4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*a*b*c
*d**3 + 16*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*
*2*b**3*d**3 - 4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)**2*b**2*c**2*d**2 + 10*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b
+ a)*tan(e + f*x)*a**2*c*d**3 + 40*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f
*x)*b + a)*tan(e + f*x)*a*b**2*d**3 - 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan
(e + f*x)*b + a)*tan(e + f*x)*a*b*c**2*d**2 - 8*sqrt(tan(e + f*x)*d + c)*s
qrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**3*c*d**2 + 2*sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**2*c**3*d + 30*sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**2*b*d**3 + 10*sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a)*a**2*c**2*d**2 - 20*sqrt(tan(e + f*x)*d + c)*s
qrt(tan(e + f*x)*b + a)*a*b**2*c*d**2 - 16*sqrt(tan(e + f*x)*d + c)*sqrt(t
an(e + f*x)*b + a)*a*b*c**3*d + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*
x)*b + a)*b**3*c**2*d + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a
)*b**2*c**4 + 15*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*ta
n(e + f*x))/(tan(e + f*x)**5*b**4*d + 4*tan(e + f*x)**4*a*b**3*d + tan(e +
f*x)**4*b**4*c + 6*tan(e + f*x)**3*a**2*b**2*d + 4*tan(e + f*x)**3*a*b**3
*c + 4*tan(e + f*x)**2*a**3*b*d + 6*tan(e + f*x)**2*a**2*b**2*c + tan(e +
f*x)*a**4*d + 4*tan(e + f*x)*a**3*b*c + a**4*c),x)*tan(e + f*x)**3*a**4*b*
*3*d**4*f - 30*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*t...
```


output

```

-(a-I*b)^(3/2)*(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-(a+I*b)^(3/2)*(B-I*(A-C))*(c+I*d)^(3/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f+1/64*(3*a^4*C*d^4-4*a^3*b*d^3*(2*B*d+3*C*c)+6*a^2*b^2*d^2*(3*c^2*C+12*B*c*d+8*(A-C)*d^2)-12*a*b^3*d*(c^3*C-6*B*c^2*d-24*c*(A-C)*d^2+16*B*d^3)+b^4*(3*c^4*C-8*B*c^3*d+48*c^2*(A-C)*d^2-192*B*c*d^3-128*(A-C)*d^4))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(5/2)/d^(5/2)/f+1/64*(64*b*(B*a^2-B*b^2+2*a*b*(A-C))*d^3+(-a*d+b*c)*(48*b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-8*B*b*d-3*C*a*d+3*C*b*c)))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b^2/d^2/f+1/96*(48*b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-8*B*b*d-3*C*a*d+3*C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)/b/d^2/f-1/24*(-8*B*b*d-3*C*a*d+3*C*b*c)*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)/d^2/f+1/4*C*(a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(5/2)/d/f

```

Mathematica [A] (verified)

Time = 8.15 (sec) , antiderivative size = 1304, normalized size of antiderivative = 1.91

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```

Integrate[(a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(5/2))/(4*d*f) + (((-3*
b*c*C + 8*b*B*d + 3*a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(
5/2))/(6*d*f) + (((48*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(3*b*c*C - 8*b
*B*d - 3*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(8*b
*f) + (((24*b*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3 - (3*(-(b*c) + a*d)*(48*
b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(3*b*c*C - 8*b*B*d - 3*a*C*d)))/8)*S
qrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]/(b*f) + ((24*(-(b^4*Sqrt
[-b^2]*d^2*(a^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - b^2*(c^2*C + 2
*B*c*d - C*d^2 - A*(c^2 - d^2)) + 2*a*b*(2*c*(A - C)*d + B*(c^2 - d^2)))
- b^5*d^2*(2*a*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) - a^2*(2*c*(A -
C)*d + B*(c^2 - d^2)) + b^2*(2*c*(A - C)*d + B*(c^2 - d^2))))*ArcTanh[(Sqr
t[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]
*Sqrt[c + d*Tan[e + f*x]])]/(b^2*Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b
^2]*d)/b]) - (24*b^2*d^2*(Sqrt[-b^2]*(a^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^
2 - d^2)) - b^2*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^2)) + 2*a*b*(2*c*(A
- C)*d + B*(c^2 - d^2))) - b*(2*a*b*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 - d^
2)) - a^2*(2*c*(A - C)*d + B*(c^2 - d^2)) + b^2*(2*c*(A - C)*d + B*(c^2 -
d^2))))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqr
t[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c
+ (Sqrt[-b^2]*d)/b]) + (3*Sqrt[b]*Sqrt[c - (a*d)/b]*Sqrt[(c/(c - (a*d)...
```

Rubi [A] (verified)

Time = 9.59 (sec) , antiderivative size = 711, normalized size of antiderivative = 1.04, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.327$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\frac{\int -\frac{1}{2}\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}((3bcC-3adC-8bBd)\tan^2(e+fx)-8(Ab-Cb+aB)d\tan(e+fx))^{4d}}{4df} \xrightarrow{27} \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{5/2}}{4df} -$$

$$\frac{\int \sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}((3bcC-3adC-8bBd)\tan^2(e+fx)-8(Ab-Cb+aB)d\tan(e+fx))^{8d}}{8d} \xrightarrow{3042} \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{5/2}}{4df} -$$

$$\frac{\int \sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}((3bcC-3adC-8bBd)\tan(e+fx)^2-8(Ab-Cb+aB)d\tan(e+fx))^{8d}}{8d} \xrightarrow{4130} \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{5/2}}{4df} -$$

$$\frac{\int -\frac{(c+d\tan(e+fx))^{3/2}(c(3cC-8Bd)b^2-2ad(3cC+20Bd)b+3a^2(16A-15C)d^2+(48b(Ab-Cb+aB)d^2+(bc-ad)(3bcC-3adC-8bBd))\tan^2(e+fx)+48(Ba^2+2bAb))}{2\sqrt{a+b\tan(e+fx)}}}{3d}}{8d} \xrightarrow{27} \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{5/2}}{4df} -$$

$$\frac{(-3aCd-8bBd+3bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3df} \int \frac{(c+d\tan(e+fx))^{3/2}(c(3cC-8Bd)b^2-2ad(3cC+20Bd)b+3a^2(16A-15C)d^2+(48b(Ab-Cb+aB)d^2+(bc-ad)(3bcC-3adC-8bBd))\tan^2(e+fx)+48(Ba^2+2bAb))}{2\sqrt{a+b\tan(e+fx)}}}{8d} \xrightarrow{3042} \frac{C(a+b\tan(e+fx))^{3/2}(c+d\tan(e+fx))^{5/2}}{4df} -$$

$$\frac{(-3aCd-8bBd+3bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3df} \int \frac{(c+d\tan(e+fx))^{3/2}(c(3cC-8Bd)b^2-2ad(3cC+20Bd)b+3a^2(16A-15C)d^2+(48b(Ab-Cb+aB)d^2+(bc-ad)(3bcC-3adC-8bBd))\tan^2(e+fx)+48(Ba^2+2bAb))}{2\sqrt{a+b\tan(e+fx)}}}{8d} \xrightarrow{4130}$$

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} - \frac{\int -\frac{3\sqrt{c+d \tan(e+fx)}(-c(3Cc^2-8Bdc-16(A-C)d^2)b^3+ad(9Cc^2+64Bdc+48(A-C)d^2))}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}}{3df} - \frac{(-3aCd-8bBd+3bcC)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3df}$$

↓ 27

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} - \frac{(-3aCd-8bBd+3bcC)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3df} - \frac{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}(48bd^2(aB+Ab-bC)+(bc-ad)(-3aCd-8bBd+3bcC))}{2bf}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} - \frac{(-3aCd-8bBd+3bcC)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3df} - \frac{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}(48bd^2(aB+Ab-bC)+(bc-ad)(-3aCd-8bBd+3bcC))}{2bf}$$

↓ 4130

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} - \frac{(-3aCd-8bBd+3bcC)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3df} - \frac{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}(48bd^2(aB+Ab-bC)+(bc-ad)(-3aCd-8bBd+3bcC))}{2bf}$$

↓ 27

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} - \frac{(-3aCd-8bBd+3bcC)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3df} - \frac{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}(48bd^2(aB+Ab-bC)+(bc-ad)(-3aCd-8bBd+3bcC))}{2bf}$$

↓ 3042

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} -$$

$$\frac{(-3aCd - 8bBd + 3bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{3df} - \frac{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}(48bd^2(aB + Ab - bC) + (bc - ad)(-3aCd - 8bBd + 3bcC))}{2bf}$$

4138

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} -$$

$$\frac{(-3aCd - 8bBd + 3bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{3df} - \frac{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}(48bd^2(aB + Ab - bC) + (bc - ad)(-3aCd - 8bBd + 3bcC))}{2bf}$$

2348

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} -$$

$$\frac{(3bcC - 3adC - 8bBd)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{3df} - \frac{(48b(Ab - Cb + aB)d^2 + (bc - ad)(3bcC - 3adC - 8bBd))\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{2bf}$$

2009

$$\frac{C(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}}{4df} -$$

$$\frac{(-3aCd - 8bBd + 3bcC)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{3df} - \frac{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}(48bd^2(aB + Ab - bC) + (bc - ad)(-3aCd - 8bBd + 3bcC))}{2bf}$$

input

```
Int[(a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f
*x] + C*Tan[e + f*x]^2),x]
```

output

$$\begin{aligned} & (C*(a + b*\text{Tan}[e + f*x])^{(3/2)}*(c + d*\text{Tan}[e + f*x])^{(5/2)})/(4*d*f) - (((3*b \\ & *c*C - 8*b*B*d - 3*a*C*d)*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]*(c + d*\text{Tan}[e + f*x])^{(5 \\ & /2)})/(3*d*f) - (((48*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(3*b*c*C - 8*b* \\ & B*d - 3*a*C*d))*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]*(c + d*\text{Tan}[e + f*x])^{(3/2)})/(2*b* \\ & f) - (3*(-1/2*(-128*(a - I*b)^{(3/2)}*b^2*(I*A + B - I*C)*(c - I*d)^{(3/2)}*d^ \\ & 2*\text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])]/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c + \\ & d*\text{Tan}[e + f*x]])) - 128*(a + I*b)^{(3/2)}*b^2*(B - I*(A - C))*(c + I*d)^{(3/ \\ & 2)}*d^2*\text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])]/(\text{Sqrt}[a + I*b]*\text{Sqr} \\ & t[c + d*\text{Tan}[e + f*x]])] + (2*(3*a^4*C*d^4 - 4*a^3*b*d^3*(3*c*C + 2*B*d) + \\ & 6*a^2*b^2*d^2*(3*c^2*C + 12*B*c*d + 8*(A - C)*d^2) - 12*a*b^3*d*(c^3*C - 6 \\ & *B*c^2*d - 24*c*(A - C)*d^2 + 16*B*d^3) + b^4*(3*c^4*C - 8*B*c^3*d + 48*c^ \\ & 2*(A - C)*d^2 - 192*B*c*d^3 - 128*(A - C)*d^4))*\text{ArcTanh}[(\text{Sqrt}[d]*\text{Sqrt}[a + \\ & b*\text{Tan}[e + f*x]])]/(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])]/(\text{Sqrt}[b]*\text{Sqrt}[d])/ \\ & (b*f) - ((64*b*(a^2*B - b^2*B + 2*a*b*(A - C))*d^3 + (b*c - a*d)*(48*b*(A*b \\ & + a*B - b*C)*d^2 + (b*c - a*d)*(3*b*c*C - 8*b*B*d - 3*a*C*d)))*\text{Sqrt}[a + b \\ & * \text{Tan}[e + f*x]]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]/(b*f)))/(4*b))/(6*d))/(8*d) \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_*)(G_x_) /; \text{FreeQ}[b, x]]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 2348

$$\begin{aligned} & \text{Int}[(P_x)*((c_) + (d_)*(x_))^{(m_)*((e_) + (f_)*(x_))^{(n_)*((a_) + (b_ \\ &)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(c + d*x)^m*(e + f*x)^ \\ & n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{PolyQ}[P \\ & x, x] \&\& (\text{IntegerQ}[p] \parallel (\text{IntegerQ}[2*p] \&\& \text{IntegerQ}[m] \&\& \text{ILtQ}[n, 0])) \&\& \\ & \text{!(IGtQ}[m, 0] \&\& \text{IGtQ}[n, 0]) \end{aligned}$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int (a + b \tan(fx + e))^{\frac{3}{2}} (c + d \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2) dx$$

input

```
int((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

output

```
int((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

Fricas [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^{\frac{3}{2}} (c + d \tan(e + fx))^{\frac{3}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**(3/2)*(c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**(3/2)*(c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan^2(fx + e) + B \tan(fx + e) + A) (b \tan(fx + e) + a)^{3/2} (c + d \tan(fx + e))^{3/2} dx$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)*(d*tan(f*x + e) + c)^(3/2), x)`

Giac [F]

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) (b \tan(fx + e) + a)^{3/2} (d \tan(fx + e) + c)^{3/2} dx$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)*(d*tan(f*x + e) + c)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `int((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `int((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))^{3/2} (c \\
& + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \right. \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) acd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) b^2 d \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) bc^2 \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) abd \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) ac^2 \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) b^2 c \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) a^2 d \\
& + 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) abc \\
& + \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) a^2 c
\end{aligned}$$

input

```
int((a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```


output

```
int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4,x)*b
*c*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*
**3,x)*a*c*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e
+ f*x)**3,x)*b**2*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a
)*tan(e + f*x)**3,x)*b*c**2 + 2*int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*tan(e + f*x)**2,x)*a*b*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(t
an(e + f*x)*b + a)*tan(e + f*x)**2,x)*a*c**2 + int(sqrt(tan(e + f*x)*d + c
)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2,x)*b**2*c + int(sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x),x)*a**2*d + 2*int(sqrt(tan
(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x),x)*a*b*c + int(sqrt
(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a),x)*a**2*c
```

3.136 $\int \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$

Optimal result	1533
Mathematica [A] (verified)	1534
Rubi [A] (verified)	1535
Maple [F(-1)]	1539
Fricas [F(-1)]	1540
Sympy [F]	1540
Maxima [F]	1541
Giac [F]	1541
Mupad [F(-1)]	1542
Reduce [F]	1542

Optimal result

Integrand size = 49, antiderivative size = 510

$$\int \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx =$$

$$\frac{\sqrt{a - ib}(iA + B - iC)(c - id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c - id}\sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib}\sqrt{c + d \tan(e + fx)}}\right)}{f}$$

$$+ \frac{\sqrt{a + ib}(iA - B - iC)(c + id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c + id}\sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib}\sqrt{c + d \tan(e + fx)}}\right)}{f}$$

$$+ \frac{(a^3Cd^3 - a^2bd^2(3cC + 2Bd) + ab^2d(3c^2C + 12Bcd + 8(A - C)d^2) - b^3(c^3C - 6Bc^2d - 24c(A - C)d^2)}{8b^{5/2}d^{3/2}f}$$

$$+ \frac{(8b(Ab + aB - bC)d^2 - (bc - ad)(bcC - 6bBd - aCd)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{8b^2df}$$

$$- \frac{(bcC - 6bBd - aCd) \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}{12bdf}$$

$$+ \frac{C \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{3df}$$

output

```

-(a-I*b)^(1/2)*(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f+(a+I*b)^(1/2)*(I*A-B-I*C)*(c+I*d)^(3/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f+1/8*(a^3*C*d^3-a^2*b*d^2*(2*B*d+3*C*c)+a*b^2*d*(3*c^2*C+12*B*c*d+8*(A-C)*d^2)-b^3*(c^3*C-6*B*c^2*d-24*c*(A-C)*d^2+16*B*d^3))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(5/2)/d^(3/2)/f+1/8*(8*b*(A*b+B*a-C*b)*d^2-(-a*d+b*c)*(-6*B*b*d-C*a*d+C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b^2/d/f-1/12*(-6*B*b*d-C*a*d+C*b*c)*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)/b/d/f+1/3*C*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)/d/f
    
```

Mathematica [A] (verified)

Time = 7.92 (sec) , antiderivative size = 867, normalized size of antiderivative = 1.70

$$\int \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \frac{C \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{3df}$$

$$+ \frac{(-bcC+6bBd+aCd)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{4bf} + \frac{3(8b(Ab+aB-bC)d^2-(bc-ad)(bcC-6bBd-aCd))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{4bf} +$$

input

```

Integrate[Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
    
```

output

```
(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2))/(3*d*f) + (((-b*c
*C) + 6*b*B*d + a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2)
)/(4*b*f) + ((3*(8*b*(A*b + a*B - b*C)*d^2 - (b*c - a*d)*(b*c*C - 6*b*B*d
- a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(4*b*f) + ((6
*b^2*d*(b*(2*a*A*c*d - 2*a*c*C*d + A*b*(c^2 - d^2) + a*B*(c^2 - d^2) - b*(
c^2*C + 2*B*c*d - C*d^2)) - Sqrt[-b^2]*(a*(c^2*C + 2*B*c*d - C*d^2 - A*(c^
2 - d^2)) + b*(2*c*(A - C)*d + B*(c^2 - d^2))))*ArcTanh[(Sqrt[-c + (Sqrt[-
b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan
[e + f*x]])]/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) - (6*b^2
*d*(b*(2*a*A*c*d - 2*a*c*C*d + A*b*(c^2 - d^2) + a*B*(c^2 - d^2) - b*(c^2*
C + 2*B*c*d - C*d^2)) + Sqrt[-b^2]*(a*(c^2*C + 2*B*c*d - C*d^2 - A*(c^2 -
d^2)) + b*(2*c*(A - C)*d + B*(c^2 - d^2))))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*
d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f
*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) + (3*Sqrt[b]*Sqr
t[c - (a*d)/b]*(a^3*C*d^3 - a^2*b*d^2*(3*c*C + 2*B*d) + a*b^2*d*(3*c^2*C +
12*B*c*d + 8*(A - C)*d^2) - b^3*(c^3*C - 6*B*c^2*d - 24*c*(A - C)*d^2 + 1
6*B*d^3))*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c - (a*
d)/b])]*Sqrt[(b*c + b*d*Tan[e + f*x])/(b*c - a*d)]/(4*Sqrt[d]*Sqrt[c + d*
Tan[e + f*x]])/(b^2*f))/(2*b))/(3*d)
```

Rubi [A] (verified)

Time = 5.99 (sec) , antiderivative size = 524, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \int \frac{(c+d \tan(e+fx))^{3/2} ((bcC-adC-6bBd) \tan^2(e+fx)-6(Ab-Cb+aB)d \tan(e+fx)+bcC-a(6A-5C)d)}{2\sqrt{a+b \tan(e+fx)}} dx + \\
 & \frac{3d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}} \\
 & \quad \downarrow 27 \\
 & \int \frac{(c+d \tan(e+fx))^{3/2} ((bcC-adC-6bBd) \tan^2(e+fx)-6(Ab-Cb+aB)d \tan(e+fx)+bcC-a(6A-5C)d)}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(c+d \tan(e+fx))^{3/2} ((bcC-adC-6bBd) \tan(e+fx)^2-6(Ab-Cb+aB)d \tan(e+fx)+bcC-a(6A-5C)d)}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \quad \downarrow 4130 \\
 & \int \frac{3\sqrt{c+d \tan(e+fx)}(c(cC+2Bd)b^2-2ad(4Ac-3Cc-3Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC)))}{2\sqrt{a+b \tan(e+fx)}} dx \\
 & \quad \downarrow 27 \\
 & \int \frac{\sqrt{c+d \tan(e+fx)}(c(cC+2Bd)b^2-2ad(4Ac-3Cc-3Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC)))}{4b\sqrt{a+b \tan(e+fx)}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\sqrt{c+d \tan(e+fx)}(c(cC+2Bd)b^2-2ad(4Ac-3Cc-3Bd)b-8d(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx)b+a^2Cd^2-(8b(Ab-Cb+aB)d^2-(bc-ad)(bcC-adC)))}{4b\sqrt{a+b \tan(e+fx)}} dx \\
 & \quad \downarrow 4130
 \end{aligned}$$

$$3 \left(\frac{C \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}}{3df} - \int \frac{-c(Cc^2 + 10Bdc + 8(A-C)d^2)b^3 - ad(13Cc^2 + 20Bdc - 8Cd^2 - 8A(2c^2 - d^2))b^2 + 16d(2aAc d - 2acCd + Ab(c^2 - d^2) + aB(c^2 - d^2) - b(Cc^2 + 2Bdc - Cd^2))}{2\sqrt{a + b \tan(e + fx)}} dx \right)$$

27

$$3 \left(\frac{C \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}}{3df} - \int \frac{-c(Cc^2 + 10Bdc + 8(A-C)d^2)b^3 + ad(16Ac^2 - 13Cc^2 - 20Bdc - 8Ad^2 + 8Cd^2)b^2 + 16d(2aAc d - 2acCd + Ab(c^2 - d^2) + aB(c^2 - d^2) - b(Cc^2 + 2Bdc - Cd^2))}{\sqrt{a + b \tan(e + fx)}} dx \right)$$

3042

$$3 \left(\frac{C \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}}{3df} - \int \frac{-c(Cc^2 + 10Bdc + 8(A-C)d^2)b^3 + ad(16Ac^2 - 13Cc^2 - 20Bdc - 8Ad^2 + 8Cd^2)b^2 + 16d(2aAc d - 2acCd + Ab(c^2 - d^2) + aB(c^2 - d^2) - b(Cc^2 + 2Bdc - Cd^2))}{\sqrt{a + b \tan(e + fx)}} dx \right)$$

4138

$$3 \left(\frac{C \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}}{3df} - \int \frac{-c(Cc^2 + 10Bdc + 8(A-C)d^2)b^3 + ad(16Ac^2 - 13Cc^2 - 20Bdc - 8Ad^2 + 8Cd^2)b^2 + 16d(2aAc d - 2acCd + Ab(c^2 - d^2) + aB(c^2 - d^2) - b(Cc^2 + 2Bdc - Cd^2))}{\sqrt{a + b \tan(e + fx)} \sqrt{c}} dx \right)$$

2348

$$\frac{(bcC - adC - 6bBd) \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}}{2bf} + 3 \left(\frac{C \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}}{3df} - \int \frac{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (8b(Ab - Cb + aB)d^2 - (bc - ad)(bcC - adC - 6bBd))}{bf} dx \right)$$

2009

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3df} - \frac{(-aCd-6bBd+bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} + 3 \left(-\frac{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(8bd^2(aB+Ab-bC)-(bc-ad)(-aCd-6bBd+bcC))}{bf} \right)$$

input

```
Int[Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]
] + C*Tan[e + f*x]^2),x]
```

output

```
(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2))/(3*d*f) - (((b*c*C
- 6*b*B*d - a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(
2*b*f) + (3*(-1/2*(-16*Sqrt[a - I*b]*b^2*(I*A + B - I*C)*(c - I*d)^(3/2)*d
*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c +
d*Tan[e + f*x]])] - 16*Sqrt[a + I*b]*b^2*(B - I*(A - C))*(c + I*d)^(3/2)*d
*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c +
d*Tan[e + f*x]])] + (2*(a^3*C*d^3 - a^2*b*d^2*(3*c*C + 2*B*d) + a*b^2*d*(3
*c^2*C + 12*B*c*d + 8*(A - C)*d^2) - b^3*(c^3*C - 6*B*c^2*d - 24*c*(A - C)
*d^2 + 16*B*d^3))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt
[c + d*Tan[e + f*x]])])/(Sqrt[b]*Sqrt[d]))/(b*f) - ((8*b*(A*b + a*B - b*C)
*d^2 - (b*c - a*d)*(b*c*C - 6*b*B*d - a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqr
t[c + d*Tan[e + f*x]]/(b*f)))/(4*b))/(6*d)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_
)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^
n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P
x, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) &&
!(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_
) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan^2(fx + e)} (c + d \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan^2(fx + e))^2 dx$$

input `int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

Fricas [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**(1/2)*(c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral(sqrt(a + b*tan(e + f*x))*(c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a} (d \tan(fx + e) + c)$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)*(d*tan(f*x + e) + c)^(3/2), x)`

Giac [F]

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a} (d \tan(fx + e) + c)$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)*(d*tan(f*x + e) + c)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (C \tan(e + fx)^2 + B \tan(e + fx) + A)$$

input `int((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `int((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\begin{aligned} & \int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) cd \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) bd \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) ad \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) bc \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) ac \end{aligned}$$

input `int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output

```
int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3,x)*c
*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2
,x)*b*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*
x)**2,x)*c**2 + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(
e + f*x),x)*a*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*ta
n(e + f*x),x)*b*c + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a),
x)*a*c
```

$$3.137 \quad \int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$$

Optimal result	1544
Mathematica [A] (verified)	1545
Rubi [A] (verified)	1546
Maple [F(-1)]	1549
Fricas [F(-1)]	1550
Sympy [F]	1550
Maxima [F]	1551
Giac [F]	1551
Mupad [F(-1)]	1551
Reduce [F]	1552

Optimal result

Integrand size = 49, antiderivative size = 382

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx =$$

$$\frac{(iA+B-ic)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a-ib}f}$$

$$- \frac{(B-i(A-C))(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a+ib}f}$$

$$+ \frac{(3a^2Cd^2 - 2abd(3cC + 2Bd) + b^2(3c^2C + 12Bcd + 8(A - C)d^2)) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{4b^{5/2}\sqrt{d}f}$$

$$+ \frac{(3bcC + 4bBd - 3aCd)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{4b^2f}$$

$$+ \frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(1/2)/f-(B-I*(A-C))*(c+I*d)^(3
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(1/2)/f+1/4*(3*a^2*C*d^2-2*a*b*d*(2*B*d+3*C*c)+b^2*(3
*c^2*C+12*B*c*d+8*(A-C)*d^2))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/
2)/(c+d*tan(f*x+e))^(1/2))/b^(5/2)/d^(1/2)/f+1/4*(4*B*b*d-3*C*a*d+3*C*b*c)
*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b^2/f+1/2*C*(a+b*tan(f*x+e)
)^(1/2)*(c+d*tan(f*x+e))^(3/2)/b/f

```

Mathematica [A] (verified)

Time = 6.55 (sec) , antiderivative size = 580, normalized size of antiderivative = 1.52

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \frac{4(-b(2c(A-C)d + B(c^2 - d^2)) + \sqrt{-b^2}(c^2 C + 2Bcd - Ad^2)) \operatorname{ArcTanh}\left[\frac{\sqrt{-c + (\sqrt{-b^2}d)/b} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right] - (4(b(2c(A-C)d + B(c^2 - d^2)) + \sqrt{-b^2}(c^2 C + 2Bcd - Ad^2)) \operatorname{ArcTanh}\left[\frac{\sqrt{c + (\sqrt{-b^2}d)/b} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right] + ((3b^2cC + 4b^2Bd - 3a^2C)d) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} / b + 2c \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2} + (\sqrt{c - (a/d)/b} (3a^2C d^2 - 2a^2b d (3c^2 C + 2Bd) + b^2 (3c^2 C + 12Bcd + 8(A-C)d^2)) \operatorname{ArcSinh}\left[\frac{\sqrt{d} \sqrt{a + b \tan(e + fx)}}{\sqrt{b} \sqrt{c - (a/d)/b}}\right] \sqrt{(b(c + d \tan(e + fx)) / (b^2 c - a^2 d)) / (b^{3/2} \sqrt{d} \sqrt{c + d \tan(e + fx)})}}{4b^2 f}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/Sqrt[a + b*Tan[e + f*x]],x]

```

output

```

((-4*(-(b*(2*c*(A - C)*d + B*(c^2 - d^2))) + Sqrt[-b^2]*(c^2*C + 2*B*c*d -
C*d^2 + A*(-c^2 + d^2)))*ArcTanh[(Sqrt[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*
Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[-a
+ Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) - (4*(b*(2*c*(A - C)*d + B*(c^
2 - d^2)) + Sqrt[-b^2]*(c^2*C + 2*B*c*d - C*d^2 + A*(-c^2 + d^2)))*ArcTanh
[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]
]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*
d)/b]) + ((3*b*c*C + 4*b*B*d - 3*a*C*d)*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c +
d*Tan[e + f*x]])/b + 2*C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/
2) + (Sqrt[c - (a*d)/b]*(3*a^2*C*d^2 - 2*a*b*d*(3*c^2*C + 2*B*d) + b^2*(3*c^
2*C + 12*B*c*d + 8*(A - C)*d^2))*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]
])/(Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)]
/(b^(3/2)*Sqrt[d]*Sqrt[c + d*Tan[e + f*x]]))/(4*b*f)

```

Rubi [A] (verified)

Time = 3.78 (sec) , antiderivative size = 393, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.224$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 25, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{\sqrt{a + b \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4130} \\
 & \frac{\int \frac{\sqrt{c+d \tan(e+fx)}((3bcC-3adC+4bBd) \tan^2(e+fx)+4b(Bc+(A-C)d) \tan(e+fx)+4Abc-C(bc+3ad))}{2\sqrt{a+b \tan(e+fx)}} dx}{\frac{2b}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} + \frac{2bf}{2bf}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{c+d \tan(e+fx)}((3bcC-3adC+4bBd) \tan^2(e+fx)+4b(Bc+(A-C)d) \tan(e+fx)+4Abc-bcC-3aCd)}{\sqrt{a+b \tan(e+fx)}} dx}{\frac{4b}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} + \frac{2bf}{2bf}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{c+d \tan(e+fx)}((3bcC-3adC+4bBd) \tan(e+fx)^2+4b(Bc+(A-C)d) \tan(e+fx)+4Abc-bcC-3aCd)}{\sqrt{a+b \tan(e+fx)}} dx}{\frac{4b}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} + \frac{2bf}{2bf}} \\
 & \quad \downarrow \text{4130}
 \end{aligned}$$

$$\int \frac{-8(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2-2c(4Abc-C(bc+3ad))b-(8d(Bc+(A-C)d)b^2+(bc-ad)(3bcC-3adC+4bBd)) \tan^2(e+fx)+(bc+ad)(3bcC-3adC-2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)})}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf}$$

4b

↓ 27

$$\int \frac{(-3aCd+4bBd+3bcC)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf} - \int \frac{8Ac^2b^2-c(5cC+4Bd)b^2+8(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2-2ad(3cC+2Bd)b+(8d(Bc+(A-C)d)b^2+(bc-ad)(3bcC-3adC+4bBd)) \tan^2(e+fx)+\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{2b}$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf}$$

4b

↓ 25

$$\int \frac{8Ac^2b^2-c(5cC+4Bd)b^2+8(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2-2ad(3cC+2Bd)b+3a^2Cd^2+(8d(Bc+(A-C)d)b^2+(bc-ad)(3bcC-3adC+4bBd)) \tan^2(e+fx)+\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{2b}$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf}$$

4b

↓ 3042

$$\int \frac{8Ac^2b^2-c(5cC+4Bd)b^2+8(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2-2ad(3cC+2Bd)b+3a^2Cd^2+(8d(Bc+(A-C)d)b^2+(bc-ad)(3bcC-3adC+4bBd)) \tan^2(e+fx)+\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{2b}$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf}$$

4b

↓ 4138

$$\int \frac{8Ac^2b^2-c(5cC+4Bd)b^2+8(2c(A-C)d+B(c^2-d^2)) \tan(e+fx)b^2-2ad(3cC+2Bd)b+3a^2Cd^2+(8d(Bc+(A-C)d)b^2+(bc-ad)(3bcC-3adC+4bBd)) \tan^2(e+fx)+\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(\tan^2(e+fx)+1)}{2bf}$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf}$$

4b

↓ 2348

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} + \frac{(-3aCd+4bBd+3bcC)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{bf} + \frac{\int\left(\frac{8Ad^2b^2-8Cd^2b^2+3c^2Cb^2+12Bcdb^2-4aBd^2b-6acCdb+3a^2Cd^2}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} + \frac{-8Bc^2b^2+8Bd^2b^2}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}\right)dx}{\sqrt{b}\sqrt{d}}$$

↓ 2009

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} + \frac{(-3aCd+4bBd+3bcC)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{bf} + \frac{2(3a^2Cd^2-2abd(2Bd+3cC)+b^2(8d^2(A-C)+12Bcd+3c^2C))\operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b\tan(e+fx)}}{\sqrt{b}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{b}\sqrt{d}}$$

4b

input `Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[a + b*Tan[e + f*x]],x]`

output `(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(2*b*f) + (((-8*b^2*(I*A + B - I*C)*(c - I*d)^(3/2)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b] - (8*b^2*(B - I*(A - C))*(c + I*d)^(3/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] + (2*(3*a^2*C*d^2 - 2*a*b*d*(3*c*C + 2*B*d) + b^2*(3*c^2*C + 12*B*c*d + 8*(A - C)*d^2))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/((Sqrt[b]*Sqrt[d]))/(2*b*f) + ((3*b*c*C + 4*b*B*d - 3*a*C*d)*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f))/(4*b)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_
)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^
n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P
x, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) &&
!(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4130

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{\sqrt{a + b \tan(fx + e)}} dx$$

input `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x)`

output `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(1/2),x)`

output `Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(a + b*tan(e + f*x)), x)`

Maxima [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) (d \tan(fx + e) + c)^{3/2}}{\sqrt{b \tan(fx + e) + a}} dx$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(3/2)/sqrt(b*tan(f*x + e) + a), x)`

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) (d \tan(fx + e) + c)^{3/2}}{\sqrt{b \tan(fx + e) + a}} dx$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(3/2)/sqrt(b*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Hanged}$$

input `int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(1/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)}}{\tan(fx + e)} dx \right) \tan(fx + e)$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) bd$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) c^2$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) ad$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) bc$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a}}{\tan(fx + e) b + a} dx \right) ac$$

input `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x)`

output `int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)*b + a),x)*c*d + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*b*d + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*c**2 + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*a*d + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*b*c + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)*b + a),x)*a*c`

3.138
$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$$

Optimal result	1553
Mathematica [B] (verified)	1554
Rubi [A] (verified)	1555
Maple [F(-1)]	1560
Fricas [F(-1)]	1560
Sympy [F]	1560
Maxima [F(-1)]	1561
Giac [F]	1561
Mupad [F(-1)]	1562
Reduce [F]	1562

Optimal result

Integrand size = 49, antiderivative size = 382

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx =$$

$$\frac{(iA+B-iC)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{3/2} f}$$

$$- \frac{(B-i(A-C))(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{3/2} f}$$

$$+ \frac{\sqrt{d}(3bcC+2bBd-3aCd) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{b^{5/2} f}$$

$$+ \frac{(2Ab^2-2abB+3a^2C+b^2C) d \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}}{b^2 (a^2+b^2) f}$$

$$- \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{b(a^2+b^2) f \sqrt{a+b \tan(e+fx)}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(3/2)/f-(B-I*(A-C))*(c+I*d)^(3
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(3/2)/f+d^(1/2)*(2*B*b*d-3*C*a*d+3*C*b*c)*arctanh(d^(
1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(5/2)/f+(2*A
*b^2-2*B*a*b+3*C*a^2+C*b^2)*d*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2
)/b^2/(a^2+b^2)/f-2*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(3/2)/b/(a^2+b^2
)/f/(a+b*tan(f*x+e))^(1/2)

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1664 vs. $2(382) = 764$.

Time = 6.80 (sec) , antiderivative size = 1664, normalized size of antiderivative = 4.36

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```

Integrate[(((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^(3/2),x]

```

output

```
(C*(c + d*Tan[e + f*x])^(3/2))/(b*f*Sqrt[a + b*Tan[e + f*x]]) + ((-2*b*(I*
A + B - I*C)*(-c + I*d)^(3/2)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f
*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((-a + I*b)^(3/2)*f) + (
2*b*(I*A - B - I*C)*(c + I*d)^(3/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[
e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((a + I*b)^(3/2)*f)
- (2*b*(A + I*B - C)*(I*c - d)*Sqrt[c + d*Tan[e + f*x]])/((a + I*b)*f*Sqrt
[a + b*Tan[e + f*x]]) + (2*b*(A - I*B - C)*(I*c + d)*Sqrt[c + d*Tan[e + f*
x]])/((a - I*b)*f*Sqrt[a + b*Tan[e + f*x]]) + (6*c*C*Sqrt[c + d*Tan[e + f*
x]]*(1 + (b*d*(a + b*Tan[e + f*x])))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a
*b*d)/(b*c - a*d))))^(3/2)*(1 - (Sqrt[b]*Sqrt[d]*ArcSinh[(Sqrt[b]*Sqrt[d]*
Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - (a*b
*d)/(b*c - a*d)])]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)
/(b*c - a*d) - (a*b*d)/(b*c - a*d)]*Sqrt[1 + (b*d*(a + b*Tan[e + f*x]))]/((
b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))))))/(Sqrt[b/((b^2*
c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)]*f*Sqrt[a + b*Tan[e + f*x]]*Sqrt[(b*
(c + d*Tan[e + f*x]))/(b*c - a*d)]*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c
- a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))))) + (4*B*d*Sqrt[c + d*
Tan[e + f*x]]*(1 + (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c -
a*d) - (a*b*d)/(b*c - a*d))))^(3/2)*(1 - (Sqrt[b]*Sqrt[d]*ArcSinh[(Sqrt[b]
]*Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c ...
```

Rubi [A] (verified)

Time = 4.25 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.204$, Rules used = {3042, 4128, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$\frac{2 \int \frac{\sqrt{c+d \tan(e+fx)}((3Ca^2-2bBa+2Ab^2+b^2C) d \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-3ad)+Ab(ac+3bd))}{2\sqrt{a+b \tan(e+fx)}}}{b(a^2+b^2)} \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 27

$$\frac{\int \frac{\sqrt{c+d \tan(e+fx)}((3Ca^2-2bBa+2Ab^2+b^2C) d \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-3ad)+Ab(ac+3bd))}{\sqrt{a+b \tan(e+fx)}}}{b(a^2+b^2)} \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 3042

$$\frac{\int \frac{\sqrt{c+d \tan(e+fx)}((3Ca^2-2bBa+2Ab^2+b^2C) d \tan(e+fx)^2-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-3ad)+Ab(ac+3bd))}{\sqrt{a+b \tan(e+fx)}}}{b(a^2+b^2)} \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 4130

$$\frac{\int \frac{-2(2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2-2c((bB-aC)(bc-3ad)+Ab(ac+3bd))b-(a^2+b^2)d(3bcC-3adC+2bBd)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}}{b} \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 27

$$\frac{d(3a^2C-2abB+2Ab^2+b^2C)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf} \frac{\int \frac{-2(2aAc d-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)}{b}}{b(a^2+b^2)} \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 3042

$$\frac{d(3a^2C - 2abB + 2Ab^2 + b^2C) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{bf} - \int \frac{-2(2aAc d - 2acCd - Ab(c^2 - d^2) + aB(c^2 - d^2) + b(Cc^2 + 2Bdc - Cd^2)) \tan(e + fx)}{b(a^2 + b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 4138

$$\frac{d(3a^2C - 2abB + 2Ab^2 + b^2C) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{bf} - \int \frac{-2(2aAc d - 2acCd - Ab(c^2 - d^2) + aB(c^2 - d^2) + b(Cc^2 + 2Bdc - Cd^2)) \tan(e + fx)}{b(a^2 + b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 2348

$$- \frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}} +$$

$$\frac{d(3a^2C - 2abB + 2Ab^2 + b^2C) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{bf} - \int \left(\frac{(a^2 + b^2) d(-3bcC + 3adC - 2bBd)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} + \frac{-2Ac^2b^3 + 2Ad^2b^3 - 2Cd^2b^3 + 2c^2Cb^3 + 4Bcd^2}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} \right)$$

↓ 2009

$$- \frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}} +$$

$$\frac{d(3a^2C - 2abB + 2Ab^2 + b^2C) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{bf} - \frac{2\sqrt{d}(a^2 + b^2)(-3aCd + 2bBd + 3bcC) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a + b \tan(e + fx)}}{\sqrt{b}\sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{b}} + \frac{2b^2(-bC + d)}{b(a^2 + b^2)}$$

$b(a^2 + b^2)$

input

```
Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(3/2),x]
```

output

$$\begin{aligned} & (-2*(A*b^2 - a*(b*B - a*C))*(c + d*\text{Tan}[e + f*x])^{(3/2)})/(b*(a^2 + b^2)*f*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]) + (-1/2*((2*(I*a - b)*b^2*(A - I*B - C)*(c - I*d) \\ & ^{(3/2)*\text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])}]/\text{Sqrt}[a - I*b] - (2*(a - I*b)*b^2*(I*A - B - I*C)* \\ & (c + I*d)^{(3/2)*\text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])}]/\text{Sqrt}[a + I*b] - (2*(a^2 + b^2)*\text{Sqrt}[d]*(\\ & 3*b*c*C + 2*b*B*d - 3*a*C*d)*\text{ArcTanh}[(\text{Sqrt}[d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])}]/\text{Sqrt}[b])/(b*f) + ((2*A*b^2 - 2*a*b*B + \\ & 3*a^2*C + b^2*C)*d*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/(b*f) \\ &))/(b*(a^2 + b^2)) \end{aligned}$$
Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] \text{ /; FreeQ}[b, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 2348

$$\begin{aligned} & \text{Int}[(P_x)*((c_) + (d_)*(x_))^{(m_)}*((e_) + (f_)*(x_))^{(n_)}*((a_) + (b_ \\ &)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P_x, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{IntegerQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0]) \end{aligned}$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4128

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]

```

rule 4130

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

rule 4138

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]

```

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan^2(fx + e))^2}{(a + b \tan(fx + e))^{\frac{3}{2}}} dx$$

input `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x)`

output `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \int \frac{(c + d \tan(e + fx))^{\frac{3}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(3/2),x)`

output

```
Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)*
*2)/(a + b*tan(e + f*x))**(3/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(3/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(d \tan(fx + e) + c)^{3/2}}{(b \tan(fx + e) + a)^{3/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(3/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(3/
2)/(b*tan(f*x + e) + a)^(3/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(3/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x)
```

output

```
(2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*d**2 + 2*sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b*c*d + int((sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*b**2 + 2*ta
n(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b*c*d**2*f - int((sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*b**2
+ 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*b**2*c**2*d*f + int((sqrt(tan
(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**
2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*c*d**2*f - int((sqrt(tan(e + f
*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*b**2
+ 2*tan(e + f*x)*a*b + a**2),x)*a*b*c**2*d*f + int((sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*b**2 + 2*ta
n(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**2*d**2*f + int((sqrt(tan(e + f
*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*b**2
+ 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b*c**2*d*f - int((sqrt(tan
(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**
2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*b**3*c*d*f - int((sqrt
(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*
x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*b**2*c**3*f + int(
(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e
+ f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*b*d**2*f + int((sq...
```


3.139
$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$$

Optimal result	1564
Mathematica [C] (verified)	1565
Rubi [A] (verified)	1566
Maple [F(-1)]	1570
Fricas [F(-1)]	1570
Sympy [F]	1570
Maxima [F(-1)]	1571
Giac [F]	1571
Mupad [F(-1)]	1572
Reduce [F]	1572

Optimal result

Integrand size = 49, antiderivative size = 402

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx =$$

$$\frac{(iA+B-iC)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{5/2} f}$$

$$- \frac{(B-i(A-C))(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{5/2} f}$$

$$+ \frac{2Cd^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{b^{5/2} f}$$

$$- \frac{2(a^4Cd+b^4(Bc+Ad)+2ab^3(Ac-cC-Bd)-a^2b^2(Bc+(A-3C)d))\sqrt{c+d \tan(e+fx)}}{b^2(a^2+b^2)^2 f \sqrt{a+b \tan(e+fx)}}$$

$$- \frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{3b(a^2+b^2) f (a+b \tan(e+fx))^{3/2}}$$

output

$$\begin{aligned}
& -(I*A+B-I*C)*(c-I*d)^{(3/2)}*\operatorname{arctanh}((c-I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/(a \\
& -I*b)^{(1/2)}/(c+d*\tan(f*x+e))^{(1/2)})/(a-I*b)^{(5/2)}/f-(B-I*(A-C))*(c+I*d)^{(3 \\
& /2)}*\operatorname{arctanh}((c+I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/(a+I*b)^{(1/2)}/(c+d*\tan(f* \\
& x+e))^{(1/2)})/(a+I*b)^{(5/2)}/f+2*C*d^{(3/2)}*\operatorname{arctanh}(d^{(1/2)}*(a+b*\tan(f*x+e))^{(\\
& 1/2)}/b^{(1/2)}/(c+d*\tan(f*x+e))^{(1/2)})/b^{(5/2)}/f-2*(a^4*C*d+b^4*(A*d+B*c)+2 \\
& *a*b^3*(A*c-B*d-C*c)-a^2*b^2*(B*c+(A-3*C)*d))*(c+d*\tan(f*x+e))^{(1/2)}/b^2/(\\
& a^2+b^2)^2/f/(a+b*\tan(f*x+e))^{(1/2)}-2/3*(A*b^2-a*(B*b-C*a))*(c+d*\tan(f*x+e \\
&))^{(3/2)}/b/(a^2+b^2)/f/(a+b*\tan(f*x+e))^{(3/2)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.47 (sec) , antiderivative size = 519, normalized size of antiderivative = 1.29

$$\begin{aligned}
& \int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx = \frac{(B+i(A-C))(c+d \tan(e+fx))}{3(a-ib)f(a+b \tan(e+fx))^{3/2}} \\
& - \frac{(iA-B-iC)(c+d \tan(e+fx))^{3/2}}{3(a+ib)f(a+b \tan(e+fx))^{3/2}} \\
& - \frac{2C(bc-ad) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, -\frac{d(a+b \tan(e+fx))}{bc-ad}\right) \sqrt{c+d \tan(e+fx)}}{3b^2 f(a+b \tan(e+fx))^{3/2} \sqrt{\frac{b(c+d \tan(e+fx))}{bc-ad}}} \\
& + \frac{(A-iB-C)(ic+d) \left(\frac{\sqrt{-c+id} \operatorname{arctanh}\left(\frac{\sqrt{-c+id} \sqrt{a+b \tan(e+fx)}}{\sqrt{-a+ib} \sqrt{c+d \tan(e+fx)}}\right)}{(-a+ib)^{3/2}} + \frac{\sqrt{c+d \tan(e+fx)}}{(a-ib) \sqrt{a+b \tan(e+fx)}} \right)}{(a-ib)f} \\
& + \frac{(A+iB-C)(ic-d) \left(\frac{\sqrt{c+id} \operatorname{arctanh}\left(\frac{\sqrt{c+id} \sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib} \sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{3/2}} - \frac{\sqrt{c+d \tan(e+fx)}}{(a+ib) \sqrt{a+b \tan(e+fx)}} \right)}{(a+ib)f}
\end{aligned}$$

input

```
Integrate[(((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^(5/2),x]
```

output

```

((B + I*(A - C))*(c + d*Tan[e + f*x])^(3/2))/(3*(a - I*b)*f*(a + b*Tan[e +
f*x])^(3/2)) - ((I*A - B - I*C)*(c + d*Tan[e + f*x])^(3/2))/(3*(a + I*b)*
f*(a + b*Tan[e + f*x])^(3/2)) - (2*C*(b*c - a*d)*Hypergeometric2F1[-3/2, -
3/2, -1/2, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d))]*Sqrt[c + d*Tan[e + f*x
]])/(3*b^2*f*(a + b*Tan[e + f*x])^(3/2)*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c
- a*d)]) + ((A - I*B - C)*(I*c + d)*((Sqrt[-c + I*d]*ArcTanh[(Sqrt[-c + I
*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/
(-a + I*b)^(3/2) + Sqrt[c + d*Tan[e + f*x]]/((a - I*b)*Sqrt[a + b*Tan[e +
f*x]])))/((a - I*b)*f) + ((A + I*B - C)*(I*c - d)*((Sqrt[c + I*d]*ArcTanh[
(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e +
f*x]])])/((a + I*b)^(3/2) - Sqrt[c + d*Tan[e + f*x]]/((a + I*b)*Sqrt[a + b
*Tan[e + f*x]])))/((a + I*b)*f)

```

Rubi [A] (verified)

Time = 4.67 (sec) , antiderivative size = 457, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.204$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$\frac{2 \int \frac{3\sqrt{c+d \tan(e+fx)}((a^2+b^2)Cd \tan^2(e+fx) - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd))}{2(a+b \tan(e+fx))^{3/2}} dx}{3b(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{3bf(a^2 + b^2)(a + b \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)}((a^2+b^2)Cd \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-ad)+Ab(ac+bd))}{(a+b \tan(e+fx))^{3/2}} dx$$

$$\frac{b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}} \\ \frac{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)}((a^2+b^2)Cd \tan(e+fx)^2-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-ad)+Ab(ac+bd))}{(a+b \tan(e+fx))^{3/2}} dx$$

$$\frac{b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}} \\ \frac{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 4128

$$2 \int \frac{-((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+bd)((bB-aC)(bc-ad)+Ab(ac+bd))b+(a^2+b^2)^2 Cd}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{-((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+bd)((bB-aC)(bc-ad)+Ab(ac+bd))b+(a^2+b^2)^2 Cd^2}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{-((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+bd)((bB-aC)(bc-ad)+Ab(ac+bd))b+(a^2+b^2)^2 Cd^2}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 4138

$$\int \frac{-\left(\left(ac+bd\right)\left(A-C\right)\left(bc-ad\right)-B\left(ac+bd\right)\right)+\left(bc-ad\right)\left(bBc+b\left(A-C\right)d+a\left(Ac-Cc-Bd\right)\right) \tan \left(e+f x\right) b^2+\left(ac+bd\right)\left(\left(bB-aC\right)\left(bc-ad\right)+A b\left(ac+bd\right)\right) b+\left(a^2+b^2\right)^2 C d^2}{\sqrt{a+b \tan \left(e+f x\right)} \sqrt{c+d \tan \left(e+f x\right)}\left(\tan ^2\left(e+f x\right)+1\right)} \frac{d f\left(a^2+b^2\right)}{b f\left(a^2+b^2\right)}$$

$$\frac{2\left(A b^2-a\left(b B-a C\right)\right)\left(c+d \tan \left(e+f x\right)\right)^{3 / 2}}{3 b f\left(a^2+b^2\right)\left(a+b \tan \left(e+f x\right)\right)^{3 / 2}}$$

2348

$$\int \left(\frac{\left(a^2+b^2\right)^2 C d^2}{\sqrt{a+b \tan \left(e+f x\right)} \sqrt{c+d \tan \left(e+f x\right)}}+\frac{B c^2 b^4-B d^2 b^4+2 A c d b^4-2 c C d b^4+2 a A c^2 b^3-2 a A d^2 b^3+2 a C d^2 b^3-2 a c^2 C b^3-4 a B c d b^3-a^2 B c^2 b^2+a^2 B d^2 b^2-2 a^2 A c d b^2}{2\left(a^2+b^2\right) \sqrt{a+b \tan \left(e+f x\right)} \sqrt{c+d \tan \left(e+f x\right)}}\right) \frac{d f\left(a^2+b^2\right)}{b f\left(a^2+b^2\right)}$$

$$\frac{2\left(A b^2-a\left(b B-a C\right)\right)\left(c+d \tan \left(e+f x\right)\right)^{3 / 2}}{3 b\left(a^2+b^2\right) f\left(a+b \tan \left(e+f x\right)\right)^{3 / 2}}$$

2009

$$\frac{2\left(A b^2-a\left(b B-a C\right)\right)\left(c+d \tan \left(e+f x\right)\right)^{3 / 2}}{3 b f\left(a^2+b^2\right)\left(a+b \tan \left(e+f x\right)\right)^{3 / 2}}+$$

$$\frac{2 \sqrt{c+d \tan \left(e+f x\right)}\left(a^4 C d-a^2 b^2\left(d\left(A-3 C\right)+B c\right)+2 a b^3\left(A c-B d-c C\right)+b^4\left(A d+B c\right)\right)}{b f\left(a^2+b^2\right) \sqrt{a+b \tan \left(e+f x\right)}}+\frac{2 C d^{3 / 2}\left(a^2+b^2\right)^2 \operatorname{arctanh}\left(\frac{\sqrt{d} \sqrt{a+b \tan \left(e+f x\right)}}{\sqrt{b} \sqrt{c+d \tan \left(e+f x\right)}}\right)}{\sqrt{b}}-\frac{b^2\left(a^2+b^2\right)}{b\left(a^2+b^2\right)}$$

input

```
Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(5/2),x]
```

output

```
(-2*(A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(3/2))/(3*b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^(3/2)) + (((-(((a + I*b)^2*b^2*(I*A + B - I*C)*(c - I*d)^(3/2)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b]) - ((a - I*b)^2*b^2*(B - I*(A - C))*(c + I*d)^(3/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] + (2*(a^2 + b^2)^2*C*d^(3/2)*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[b])/(b*(a^2 + b^2)*f) - (2*(a^4*C*d + b^4*(B*c + A*d) + 2*a*b^3*(A*c - c*C - B*d) - a^2*b^2*(B*c + (A - 3*C)*d))*Sqrt[c + d*Tan[e + f*x]])/(b*(a^2 + b^2)*f*Sqrt[a + b*Tan[e + f*x]])/(b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 2348 $\text{Int}[(Px_)*((c_) + (d_)*(x_))^{(m_)*((e_) + (f_)*(x_))^{(n_)*((a_) + (b_)*(x_)^2)^{(p_)}}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4128 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]^{(m_)*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]^{(n_)*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_)] + (C_.)*\tan[(e_.) + (f_.)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[(A*d^2 + c*(c*C - B*d))*(a + b*\tan[e + f*x])^m*((c + d*\tan[e + f*x])^{(n + 1)}/(d*f*(n + 1)*(c^2 + d^2))), x] - \text{Simp}[1/(d*(n + 1)*(c^2 + d^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m - 1)}*(c + d*\tan[e + f*x])^{(n + 1)}*\text{Simp}[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*\tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*\tan[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[n, -1]$
- rule 4138 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]^{(m_)*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]^{(n_)*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_)] + (C_.)*\tan[(e_.) + (f_.)*(x_)]^2)}, x_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\tan[e + f*x], x]\}, \text{Simp}[ff/f \ \text{Subst}[\text{Int}[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, \tan[e + f*x]/ff], x]] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan^2(fx + e))^2}{(a + b \tan(fx + e))^{\frac{5}{2}}} dx$$

input `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x)`

output `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{(c + d \tan(e + fx))^{\frac{3}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(5/2),x)`

output

```
Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)*
*2)/(a + b*tan(e + f*x))**(5/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(5/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(d \tan(fx + e) + c)^{3/2}}{(b \tan(fx + e) + a)^{5/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(5/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(3/
2)/(b*tan(f*x + e) + a)^(5/2), x)
```


Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(5/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x)
```

output

```
(6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*b*d**3
+ 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*c**2
*d**2 + 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b
**2*c*d**2 - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f
*x)*b*c**3*d + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**2*d*
*3 + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b*c*d**2 + 2*sq
rt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*c**3*d - 4*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b**2*c**2*d - 2*sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a)*b*c**4 + 3*int((sqrt(tan(e + f*x)*d + c)*sqrt
(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**3*b**3 + 3*tan(e + f*
x)**2*a*b**2 + 3*tan(e + f*x)*a**2*b + a**3),x)*tan(e + f*x)**2*a**2*b**2*
c*d**3*f - 6*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e
+ f*x)**3)/(tan(e + f*x)**3*b**3 + 3*tan(e + f*x)**2*a*b**2 + 3*tan(e + f*
x)*a**2*b + a**3),x)*tan(e + f*x)**2*a*b**3*c**2*d**2*f + 3*int((sqrt(tan(
e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**3
*b**3 + 3*tan(e + f*x)**2*a*b**2 + 3*tan(e + f*x)*a**2*b + a**3),x)*tan(e
+ f*x)**2*b**4*c**3*d*f + 6*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x
)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**3*b**3 + 3*tan(e + f*x)**2*a*b**2
+ 3*tan(e + f*x)*a**2*b + a**3),x)*tan(e + f*x)*a**3*b*c*d**3*f - 12*int(
(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(ta...
```

3.140
$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$$

Optimal result	1574
Mathematica [B] (verified)	1575
Rubi [A] (verified)	1576
Maple [F(-1)]	1582
Fricas [F(-1)]	1582
Sympy [F]	1582
Maxima [F(-2)]	1583
Giac [F]	1583
Mupad [F(-1)]	1584
Reduce [F]	1584

Optimal result

Integrand size = 49, antiderivative size = 586

$$\int \frac{(c+d \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx =$$

$$\frac{(iA+B-iC)(c-id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{7/2} f}$$

$$-\frac{(B-i(A-C))(c+id)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{7/2} f}$$

$$-\frac{2(2a^3bBd+3a^4Cd+b^4(5Bc+3Ad)+2ab^3(5Ac-5cC-4Bd)-a^2b^2(5Bc+7Ad-13Cd))\sqrt{c+d \tan(e+fx)}}{15b^2(a^2+b^2)^2 f(a+b \tan(e+fx))^{3/2}}$$

$$-\frac{2(2a^5bBd^2+3a^6Cd^2+a^4b^2d(10Bc+(8A+C)d)+a^2b^4(45Ac^2-45c^2C-90Bcd-49Ad^2+58Cd^2)-15b^2(a^2+b^2)^2 f(a+b \tan(e+fx))^{3/2}}{15b^2(a^2+b^2)^2 f(a+b \tan(e+fx))^{3/2}}$$

$$-\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{5b(a^2+b^2) f(a+b \tan(e+fx))^{5/2}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(3/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(7/2)/f-(B-I*(A-C))*(c+I*d)^(3
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(7/2)/f-2/15*(2*a^3*b*B*d+3*a^4*C*d+b^4*(3*A*d+5*B*c)
+2*a*b^3*(5*A*c-4*B*d-5*C*c)-a^2*b^2*(7*A*d+5*B*c-13*C*d))*(c+d*tan(f*x+e)
)^(1/2)/b^2/(a^2+b^2)^2/f/(a+b*tan(f*x+e))^(3/2)-2/15*(2*a^5*b*B*d^2+3*a^6
*C*d^2+a^4*b^2*d*(10*B*c+(8*A+C)*d)+a^2*b^4*(45*A*c^2-49*A*d^2-90*B*c*d-45
*C*c^2+58*C*d^2)-a^3*b^3*(50*c*(A-C)*d+B*(15*c^2-39*d^2))+a*b^5*(70*c*(A-C)
*d+B*(45*c^2-23*d^2))+b^6*(5*c*(4*B*d+3*C*c)-3*A*(5*c^2-d^2)))*(c+d*tan(f
*x+e))^(1/2)/b^2/(a^2+b^2)^3/(-a*d+b*c)/f/(a+b*tan(f*x+e))^(1/2)-2/5*(A*b^
2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(3/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^(5/2)

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 3134 vs. $2(586) = 1172$.

Time = 8.26 (sec) , antiderivative size = 3134, normalized size of antiderivative = 5.35

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Result too large to show}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^(7/2),x]

```

output

```

-((C*(c + d*Tan[e + f*x])^(3/2))/(b*f*(a + b*Tan[e + f*x])^(5/2))) - (-1/4
*((3*b*c*C - 2*b*B*d - 3*a*C*d)*Sqrt[c + d*Tan[e + f*x]])/(b*f*(a + b*Tan[
e + f*x])^(5/2)) - ((-2*((b^2*(8*A*b^2*c^2 + 3*a^2*C*d^2 - 2*a*b*d*(3*c*C
- B*d) - 5*b^2*c*(c*C + 2*B*d)))/4 - a*(-1/4*(a*(8*b^2*d*(B*c + (A - C)*d)
+ (b*c - a*d)*(3*b*c*C - 2*b*B*d - 3*a*C*d))) + 2*b^3*(2*c*(A - C)*d + B*
(c^2 - d^2))))*Sqrt[c + d*Tan[e + f*x]]/(5*(a^2 + b^2)*(b*c - a*d)*f*(a +
b*Tan[e + f*x])^(5/2)) - (2*((-2*(b^2*((2*b^2*d - (5*a*(b*c - a*d))/2)*(
8*A*b^2*c^2 + 3*a^2*C*d^2 - 2*a*b*d*(3*c*C - B*d) - 5*b^2*c*(c*C + 2*B*d)
)/4 + ((-5*b*c)/2 + (a*d)/2)*(-1/4*(a*(8*b^2*d*(B*c + (A - C)*d) + (b*c -
a*d)*(3*b*c*C - 2*b*B*d - 3*a*C*d))) + 2*b^3*(2*c*(A - C)*d + B*(c^2 - d^2
)))) - a*((5*b*(b*c - a*d)*((b*(8*A*b^2*c^2 + 3*a^2*C*d^2 - 2*a*b*d*(3*c*C
- B*d) - 5*b^2*c*(c*C + 2*B*d)))/4 - (b*(8*b^2*d*(B*c + (A - C)*d) + (b*c
- a*d)*(3*b*c*C - 2*b*B*d - 3*a*C*d)))/4 - 2*a*b^2*(2*c*(A - C)*d + B*(c^
2 - d^2))))/2 - 2*a*d*((b^2*(8*A*b^2*c^2 + 3*a^2*C*d^2 - 2*a*b*d*(3*c*C -
B*d) - 5*b^2*c*(c*C + 2*B*d)))/4 - a*(-1/4*(a*(8*b^2*d*(B*c + (A - C)*d) +
(b*c - a*d)*(3*b*c*C - 2*b*B*d - 3*a*C*d))) + 2*b^3*(2*c*(A - C)*d + B*(c
^2 - d^2)))))*Sqrt[c + d*Tan[e + f*x]]/(3*(a^2 + b^2)*(b*c - a*d)*f*(a +
b*Tan[e + f*x])^(3/2)) - (2*((-15*b^2*(b*c - a*d)^2*((3*a^2*A*b*c^2 - A*
b^3*c^2 - a^3*B*c^2 + 3*a*b^2*B*c^2 - 3*a^2*b*c^2*C + b^3*c^2*C - 2*a^3*A*
c*d + 6*a*A*b^2*c*d - 6*a^2*b*B*c*d + 2*b^3*B*c*d + 2*a^3*c*C*d - 6*a*b...

```

Rubi [A] (verified)

Time = 7.70 (sec) , antiderivative size = 678, normalized size of antiderivative = 1.16, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{7/2}} dx$$

↓ 4128

$$2 \int \frac{\sqrt{c+d \tan(e+fx)}(-((-3Ca^2-2bBa+2Ab^2-5b^2C)d \tan^2(e+fx))-5b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(5bc-3ad)+Ab(5ac+3bd))}{2(a+b \tan(e+fx))^{5/2}}$$

$$\frac{5b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}$$

$$\frac{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)}(-((-3Ca^2-2bBa+2Ab^2-5b^2C)d \tan^2(e+fx))-5b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(5bc-3ad)+Ab(5ac+3bd))}{(a+b \tan(e+fx))^{5/2}}$$

$$\frac{5b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}$$

$$\frac{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)}(-((-3Ca^2-2bBa+2Ab^2-5b^2C)d \tan^2(e+fx))-5b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(5bc-3ad)+Ab(5ac+3bd))}{(a+b \tan(e+fx))^{5/2}}$$

$$\frac{5b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}$$

$$\frac{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 4128

$$2 \int \frac{-15((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(3ac+bd)((bB-aC)(5bc-3ad)+Ab(5ac+3bd))b+d(3Cda^4)}{2(a+b \tan(e+fx))^{5/2}}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 27

$$\int \frac{-15((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(3ac+bd)((bB-aC)(5bc-3ad)+Ab(5ac+3bd))b+d(3Cda^4)}{(a+b \tan(e+fx))^{5/2}}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{3/2}}{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 3042

$$\int \frac{-15((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(3ac+bd)((bB-aC)(5bc-3ad)+Ab(5ac+3bd))b+d(3Cda^4)}{(a+b \tan(e+fx))^5 \sqrt{a^2+b^2}}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 4132

$$- \frac{2 \int \frac{15(b^2(bc-ad)((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^3-3b(2c(A-C)d+B(c^2-d^2))a^2-3b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))a+b^3(2c(A-C)d+B(c^2-d^2)))}{(a^2+b^2)^{5/2}}}{\sqrt{a+b \tan(e+fx)}}}{(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 27

$$- \frac{15 \int \frac{b^2(bc-ad)((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^3-3b(2c(A-C)d+B(c^2-d^2))a^2-3b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))a+b^3(2c(A-C)d+B(c^2-d^2)))}{(a^2+b^2)^{5/2}}}{\sqrt{a+b \tan(e+fx)}}}{(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 3042

$$- \frac{15 \int \frac{b^2(bc-ad)((Cc^2+2Bdc-Cd^2-A(c^2-d^2))a^3-3b(2c(A-C)d+B(c^2-d^2))a^2-3b^2(Cc^2+2Bdc-Cd^2-A(c^2-d^2))a+b^3(2c(A-C)d+B(c^2-d^2)))}{(a^2+b^2)^{5/2}}}{\sqrt{a+b \tan(e+fx)}}}{(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 4099

$$- \frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}} +$$

$$- \frac{2\sqrt{c+d \tan(e+fx)}(3a^4Cd+2a^3bBd-a^2b^2(7Ad+5Bc-13Cd)+2ab^3(5Ac-4Bd-5cC)+b^4(3Ad+5Bc))}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}} + \frac{2\sqrt{c+d \tan(e+fx)}(3a^6Cd^2+2a^5bBd)}{(a^2+b^2)}$$

↓ 3042

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c+d \tan(e+fx)}(3a^6Cd^2+2a^5bBd-a^2b^2(7Ad+5Bc-13Cd)+2ab^3(5Ac-4Bd-5cC)+b^4(3Ad+5Bc))}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 4098

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c+d \tan(e+fx)}(3a^6Cd^2+2a^5bBd-a^2b^2(7Ad+5Bc-13Cd)+2ab^3(5Ac-4Bd-5cC)+b^4(3Ad+5Bc))}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 104

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c+d \tan(e+fx)}(3a^6Cd^2+2a^5bBd-a^2b^2(7Ad+5Bc-13Cd)+2ab^3(5Ac-4Bd-5cC)+b^4(3Ad+5Bc))}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 221

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{3/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}} + \frac{2\sqrt{c+d \tan(e+fx)}(3a^6Cd^2+2a^5bBd-a^2b^2(7Ad+5Bc-13Cd)+2ab^3(5Ac-4Bd-5cC)+b^4(3Ad+5Bc))}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

input

```
Int[((c + d*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(7/2),x]
```


output

```
(-2*(A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(3/2))/(5*b*(a^2 + b^2)*f
*(a + b*Tan[e + f*x])^(5/2)) + ((-2*(2*a^3*b*B*d + 3*a^4*C*d + b^4*(5*B*c
+ 3*A*d) + 2*a*b^3*(5*A*c - 5*c*C - 4*B*d) - a^2*b^2*(5*B*c + 7*A*d - 13*C
*d))*Sqrt[c + d*Tan[e + f*x]])/(3*b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^(3/
2)) + ((-15*((I*(a + I*b)^3*b^2*(A - I*B - C)*(c - I*d)^(3/2)*(b*c - a*d)*
ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d
*Tan[e + f*x]])])/(Sqrt[a - I*b]*f) - (I*(a - I*b)^3*b^2*(A + I*B - C)*(c
+ I*d)^(3/2)*(b*c - a*d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/
(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*f)))/((a^2 + b^2
)*(b*c - a*d)) - (2*(2*a^5*b*B*d^2 + 3*a^6*C*d^2 + a^4*b^2*d*(10*B*c + (8*
A + C)*d) + a^2*b^4*(45*A*c^2 - 45*c^2*C - 90*B*c*d - 49*A*d^2 + 58*C*d^2)
- a^3*b^3*(50*c*(A - C)*d + B*(15*c^2 - 39*d^2)) + a*b^5*(70*c*(A - C)*d
+ B*(45*c^2 - 23*d^2)) + b^6*(5*c*(3*c*C + 4*B*d) - 3*A*(5*c^2 - d^2))*Sq
rt[c + d*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e + f*x]
])/((3*b*(a^2 + b^2)))/(5*b*(a^2 + b^2))
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 104

```
Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4098

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

rule 4099

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
(!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(a + b \tan(fx + e))^{\frac{7}{2}}} dx$$

input `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x)`

output `int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \int \frac{(c + d \tan(e + fx))^{\frac{3}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx$$

input `integrate((c+d*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(7/2),x)`

output

```
Integral((c + d*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)*
*2)/(a + b*tan(e + f*x))**(7/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(7/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(((2*b*d+2*a*c)^2>0)', see `assum
e?` for mo
```

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) * (d \tan(fx + e) + c)^{3/2}}{(b \tan(fx + e) + a)^{7/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(7/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(3/
2)/(b*tan(f*x + e) + a)^(7/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(7/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x)
```

output

```
(6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*a**2*
c*d**4 + 20*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)
**2*a*b**2*d**4 - 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan
(e + f*x)**2*a*b*c**2*d**3 + 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)
*b + a)*tan(e + f*x)**2*b**3*c*d**3 + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(
e + f*x)*b + a)*tan(e + f*x)**2*b**2*c**3*d**2 - 16*sqrt(tan(e + f*x)*d +
c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b**2*c*d**4 + 50*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b*d**4 + 12*sqrt(ta
n(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*c**2*d**3 + 2
0*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*b**2*c*
d**3 - 24*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a
*b*c**3*d**2 - 40*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e
+ f*x)*a*b*c*d**4 - 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*ta
n(e + f*x)*b**3*c**2*d**2 + 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*
b + a)*tan(e + f*x)*b**2*c**4*d + 8*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*tan(e + f*x)*b**2*c**2*d**3 + 30*sqrt(tan(e + f*x)*d + c)*sqrt
(tan(e + f*x)*b + a)*a**3*d**4 + 20*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*a**2*b*c*d**3 + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b
+ a)*a**2*c**3*d**2 - 30*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a
)*a**2*c*d**4 - 30*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a...
```

3.141 $\int \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$

Optimal result	1586
Mathematica [A] (verified)	1587
Rubi [A] (verified)	1588
Maple [F(-1)]	1593
Fricas [F(-1)]	1594
Sympy [F]	1594
Maxima [F]	1595
Giac [F]	1595
Mupad [F(-1)]	1596
Reduce [F]	1596

Optimal result

Integrand size = 49, antiderivative size = 695

$$\int \sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx =$$

$$\frac{\sqrt{a - ib}(iA + B - iC)(c - id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c - id}\sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib}\sqrt{c + d \tan(e + fx)}}\right)}{f} - \frac{\sqrt{a + ib}(B - i(A - C))(c + id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c + id}\sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib}\sqrt{c + d \tan(e + fx)}}\right)}{f}$$

$$\frac{(5a^4Cd^4 - 4a^3bd^3(5cC + 2Bd) + 2a^2b^2d^2(15c^2C + 20Bcd + 8(A - C)d^2) - 4ab^3d(5c^3C + 30Bc^2d + 40bc^2C + 48b^2d^2(Abc + aBc - bcC + aAd - bBd - aCd) + (bc - ad)(48b(Ab + aB - bC)d^2 - 5(bc - ad)(bcC - 8bBd - aCd))\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2} - (bcC - 8bBd - aCd)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}}{64b^3df + 96b^2df - 24bdf} + \frac{C\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{7/2}}{4df}$$

output

```

-(a-I*b)^(1/2)*(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-(a+I*b)^(1/2)*(B-I*(A-C))*(c+I*d)^(5/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/f-1/64*(5*a^4*C*d^4-4*a^3*b*d^3*(2*B*d+5*C*c)+2*a^2*b^2*d^2*(15*c^2*C+20*B*c*d+8*(A-C)*d^2)-4*a*b^3*d*(5*c^3*C+30*B*c^2*d+40*c*(A-C)*d^2-16*B*d^3)+b^4*(5*c^4*C-40*B*c^3*d-240*c^2*(A-C)*d^2+320*B*c*d^3+128*(A-C)*d^4)*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(7/2)/d^(3/2)/f+1/64*(64*b^2*d^2*(A*a*d+A*b*c+B*a*c-B*b*d-C*a*d-C*b*c)+(-a*d+b*c)*(48*b*(A*b+B*a-C*b)*d^2-5*(-a*d+b*c)*(-8*B*b*d-C*a*d+C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b^3/d/f+1/96*(48*b*(A*b+B*a-C*b)*d^2-5*(-a*d+b*c)*(-8*B*b*d-C*a*d+C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)/b^2/d/f-1/24*(-8*B*b*d-C*a*d+C*b*c)*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)/b/d/f+1/4*C*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(7/2)/d/f

```

Mathematica [A] (verified)

Time = 8.35 (sec) , antiderivative size = 1261, normalized size of antiderivative = 1.81

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Too large to display}$$

input

```

Integrate[Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]

```


output

```
(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(7/2))/(4*d*f) + (((-b*c
*C) + 8*b*B*d + a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2)
)/(6*b*f) + (((48*b*(A*b + a*B - b*C)*d^2 - 5*(b*c - a*d)*(b*c*C - 8*b*B*d
- a*C*d))*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(8*b*f) +
(((24*b^2*d^2*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d) - (3*(-b*c)
+ a*d)*(48*b*(A*b + a*B - b*C)*d^2 - 5*(b*c - a*d)*(b*c*C - 8*b*B*d - a*C
*d)))/8)*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]/(b*f) + ((-24*
b^3*d*(Sqrt[-b^2]*(b*(A - C)*d*(3*c^2 - d^2) + b*B*(c^3 - 3*c*d^2) - a*(A*
c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3)) - b*(A*(b*c^3 +
3*a*c^2*d - 3*b*c*d^2 - a*d^3) - b*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3)
+ a*(B*c^3 - 3*c^2*C*d - 3*B*c*d^2 + C*d^3)))*ArcTanh[(Sqrt[-c + (Sqrt[-b
^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan[
e + f*x]])]/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) - (24*b^3
*d*(Sqrt[-b^2]*(b*(A - C)*d*(3*c^2 - d^2) + b*B*(c^3 - 3*c*d^2) - a*(A*c^3
- c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3)) + b*(A*(b*c^3 + 3*a
*c^2*d - 3*b*c*d^2 - a*d^3) - b*(c^3*C + 3*B*c^2*d - 3*c*C*d^2 - B*d^3) +
a*(B*c^3 - 3*c^2*C*d - 3*B*c*d^2 + C*d^3)))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*
d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f
*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) - (3*Sqrt[b]*Sqr
t[c - (a*d)/b]*Sqrt[(c/(c - (a*d)/b) - (a*d)/(b*(c - (a*d)/b)))^(-1)]*S...
```

Rubi [A] (verified)

Time = 9.39 (sec) , antiderivative size = 717, normalized size of antiderivative = 1.03, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.327$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\begin{aligned}
 & \int \frac{(c+d \tan(e+fx))^{5/2} ((bcC-adC-8bBd) \tan^2(e+fx)-8(Ab-Cb+aB)d \tan(e+fx)+bcC-a(8A-7C)d)}{2\sqrt{a+b \tan(e+fx)}} dx \\
 & \frac{4d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{7/2}} + \\
 & \quad \downarrow 27 \\
 & \int \frac{(c+d \tan(e+fx))^{5/2} ((bcC-adC-8bBd) \tan^2(e+fx)-8(Ab-Cb+aB)d \tan(e+fx)+bcC-a(8A-7C)d)}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \frac{8d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{7/2}} - \\
 & \quad \downarrow 3042 \\
 & \int \frac{(c+d \tan(e+fx))^{5/2} ((bcC-adC-8bBd) \tan(e+fx)^2-8(Ab-Cb+aB)d \tan(e+fx)+bcC-a(8A-7C)d)}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \frac{8d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{7/2}} - \\
 & \quad \downarrow 4130 \\
 & \int \frac{(c+d \tan(e+fx))^{3/2} (c(5cC+8Bd)b^2-2ad(24Ac-19Cc-20Bd)b-48d(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx)b+5a^2Cd^2-(48b(Ab-Cb+aB)d^2-5(bc-ad))}{2\sqrt{a+b \tan(e+fx)}} dx \\
 & \frac{8d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{7/2}} - \\
 & \quad \downarrow 27 \\
 & \int \frac{(c+d \tan(e+fx))^{3/2} (c(5cC+8Bd)b^2-2ad(24Ac-19Cc-20Bd)b-48d(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx)b+5a^2Cd^2-(48b(Ab-Cb+aB)d^2-5(bc-ad))}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \frac{8d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{7/2}} - \\
 & \quad \downarrow 3042 \\
 & \int \frac{(c+d \tan(e+fx))^{3/2} (c(5cC+8Bd)b^2-2ad(24Ac-19Cc-20Bd)b-48d(Abc+aBc-bCc+aAd-bBd-aCd) \tan(e+fx)b+5a^2Cd^2-(48b(Ab-Cb+aB)d^2-5(bc-ad))}{\sqrt{a+b \tan(e+fx)}} dx \\
 & \frac{8d}{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{7/2}} - \\
 & \quad \downarrow 4130
 \end{aligned}$$

$$\int \frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\sqrt{c+d\tan(e+fx)}(-c(5Cc^2+24Bdc+16(A-C)d^2)b^3+ad(64Ac^2-49Cc^2-96Bdc-48Ad^2+48Cd^2)b^2+64d(2aAc d-2acCd+Ab(c^2-d^2)+aB(c^2-d^2))-b(Cc^3+3ad^2c-3ad^2c-3ad^2c-3ad^2c))}{4df}$$

↓ 27

$$\int \frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\sqrt{c+d\tan(e+fx)}(-c(5Cc^2+24Bdc+16(A-C)d^2)b^3+ad(64Ac^2-49Cc^2-96Bdc-48Ad^2+48Cd^2)b^2+64d(2aAc d-2acCd+Ab(c^2-d^2)+aB(c^2-d^2))-b(Cc^3+3ad^2c-3ad^2c-3ad^2c-3ad^2c))}{4df}$$

↓ 3042

$$\int \frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\sqrt{c+d\tan(e+fx)}(-c(5Cc^2+24Bdc+16(A-C)d^2)b^3+ad(64Ac^2-49Cc^2-96Bdc-48Ad^2+48Cd^2)b^2+64d(2aAc d-2acCd+Ab(c^2-d^2)+aB(c^2-d^2))-b(Cc^3+3ad^2c-3ad^2c-3ad^2c-3ad^2c))}{4df}$$

↓ 4130

$$\int \frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\left(\int \frac{c(5Cc^3+88Bdc^2+144(A-C)d^2c-64Bd^3)b^4+4ad(27Cc^3+66Bdc^2-56Cd^2c-16Bd^3-8A(4c^3-7cd^2))b^3-128d(A(bc^3+3adc^2-3bd^2c-ad^3))-b(Cc^3+3ad^2c-3ad^2c-3ad^2c-3ad^2c))}{4df} - \frac{c(5Cc^3+88Bdc^2+144(A-C)d^2c-64Bd^3)}{bf}\right)}{4df}$$

↓ 27

$$\int \frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\left(\int \frac{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(64b^2d^2(ad+aBc-aCd+Abc-bBd-bcC)+(bc-ad)(48bd^2(aB+Ab-bC)-5(bc-ad)(-aCd-8bBd+bcC))}{bf} - \frac{c(5Cc^3+88Bdc^2+144(A-C)d^2c-64Bd^3)}{bf}\right)}{4df}$$

↓ 3042

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\left(\frac{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(64b^2d^2(ad+aBc-aCd+Abc-bBd-bcC)+(bc-ad)(48bd^2(aB+Ab-bC)-5(bc-ad)(-aCd-8bBd+bcC)))}{bf} - \int \frac{c(5C^3+...}{...}\right)}{...}$$

↓ 4138

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{3\left(\frac{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(64b^2d^2(ad+aBc-aCd+Abc-bBd-bcC)+(bc-ad)(48bd^2(aB+Ab-bC)-5(bc-ad)(-aCd-8bBd+bcC)))}{bf} - \int \frac{c(5C^3+...}{...}\right)}{...}$$

↓ 2348

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{(bcC-adC-8bBd)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3bf} + \frac{(48b(Ab-Cb+aB)d^2-5(bc-ad)(bcC-adC-8bBd))\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf}$$

↓ 2009

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{7/2}}{4df} - \frac{(-aCd-8bBd+bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3bf} + \frac{\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}(48bd^2(aB+Ab-bC)-5(bc-ad)(-aCd-8bBd+bcC))}{2bf}$$

input `Int[Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]`

output

```
(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(7/2))/(4*d*f) - (((b*c*C
- 8*b*B*d - a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2))/(
3*b*f) + (-1/2*((48*b*(A*b + a*B - b*C)*d^2 - 5*(b*c - a*d)*(b*c*C - 8*b*B
*d - a*C*d))*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(b*f) -
(3*(-1/2*(128*Sqrt[a - I*b]*b^3*(B + I*(A - C))*(c - I*d)^(5/2)*d*ArcTanh[
(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e +
f*x]])] - 128*Sqrt[a + I*b]*b^3*(I*A - B - I*C)*(c + I*d)^(5/2)*d*ArcTanh
[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e
+ f*x]])] + (2*(5*a^4*C*d^4 - 4*a^3*b*d^3*(5*c*C + 2*B*d) + 2*a^2*b^2*d^2*
(15*c^2*C + 20*B*c*d + 8*(A - C)*d^2) - 4*a*b^3*d*(5*c^3*C + 30*B*c^2*d +
40*c*(A - C)*d^2 - 16*B*d^3) + b^4*(5*c^4*C - 40*B*c^3*d - 240*c^2*(A - C)
*d^2 + 320*B*c*d^3 + 128*(A - C)*d^4))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e +
f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[b]*Sqrt[d]))/(b*f) + ((
64*b^2*d^2*(A*b*c + a*B*c - b*c*C + a*A*d - b*B*d - a*C*d) + (b*c - a*d)*(
48*b*(A*b + a*B - b*C)*d^2 - 5*(b*c - a*d)*(b*c*C - 8*b*B*d - a*C*d))*Sqr
t[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f)))/(4*b))/(6*b))/(8*d
)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_
.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^
n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P
x, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) &&
!(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4130

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

rule 4138

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]

```

Maple [F(-1)]

Timed out.

hanged

input

```
int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

output

```
int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**(1/2)*(c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral(sqrt(a + b*tan(e + f*x))*(c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a} (d \tan(fx + e) + c)$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)*(d*tan(f*x + e) + c)^(5/2), x)`

Giac [F]

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a} (d \tan(fx + e) + c)$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)*(d*tan(f*x + e) + c)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx = \int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (C \tan(e + fx)^2 + B \tan(e + fx) + A)$$

input `int((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

output `int((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\begin{aligned} & \int \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^4 dx \right) c d^2 \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) b d^2 \\ &+ 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3 dx \right) c^2 d \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) a d^2 \\ &+ 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) b c d \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2 dx \right) c^3 \\ &+ 2 \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) a c d \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e) dx \right) b c^2 \\ &+ \left(\int \sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} dx \right) a c^2 \end{aligned}$$

input

```
int((a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*
x+e)^2),x)
```

output

```
int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4,x)*c
*d**2 + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)
**3,x)*b*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*ta
n(e + f*x)**3,x)*c**2*d + int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b
+ a)*tan(e + f*x)**2,x)*a*d**2 + 2*int(sqrt(tan(e + f*x)*d + c)*sqrt(tan(
e + f*x)*b + a)*tan(e + f*x)**2,x)*b*c*d + int(sqrt(tan(e + f*x)*d + c)*sq
rt(tan(e + f*x)*b + a)*tan(e + f*x)**2,x)*c**3 + 2*int(sqrt(tan(e + f*x)*d
+ c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x),x)*a*c*d + int(sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x),x)*b*c**2 + int(sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a),x)*a*c**2
```

$$3.142 \quad \int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx$$

Optimal result	1598
Mathematica [A] (verified)	1599
Rubi [A] (verified)	1600
Maple [F(-1)]	1605
Fricas [F(-1)]	1605
Sympy [F]	1605
Maxima [F(-1)]	1606
Giac [F]	1606
Mupad [F(-1)]	1607
Reduce [F]	1607

Optimal result

Integrand size = 49, antiderivative size = 507

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx =$$

$$-\frac{(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a-ib}f}$$

$$+\frac{(iA-B-iC)(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a+ib}f}$$

$$-\frac{(5a^3Cd^3-3a^2bd^2(5cC+2Bd)+ab^2d(15c^2C+20Bcd+8(A-C)d^2)-b^3(5c^3C+30Bc^2d+40c(A-C)d^2+3B^2d^2))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{8b^{7/2}\sqrt{d}f}$$

$$+\frac{(8b^2d(Bc+(A-C)d)+(bc-ad)(5bcC+6bBd-5aCd))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{8b^3f}$$

$$+\frac{(5bcC+6bBd-5aCd)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{12b^2f}$$

$$+\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3bf}$$

output

$$\begin{aligned}
& -(I*A+B-I*C)*(c-I*d)^{(5/2)}*\operatorname{arctanh}((c-I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/(a \\
& -I*b)^{(1/2)}/(c+d*\tan(f*x+e))^{(1/2)})/(a-I*b)^{(1/2)}/f+(I*A-B-I*C)*(c+I*d)^{(5 \\
& /2)}*\operatorname{arctanh}((c+I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/(a+I*b)^{(1/2)}/(c+d*\tan(f* \\
& x+e))^{(1/2)})/(a+I*b)^{(1/2)}/f-1/8*(5*a^3*C*d^3-3*a^2*b*d^2*(2*B*d+5*C*c)+a* \\
& b^2*d*(15*c^2*C+20*B*c*d+8*(A-C)*d^2)-b^3*(5*c^3*C+30*B*c^2*d+40*c*(A-C)*d \\
& ^2-16*B*d^3))*\operatorname{arctanh}(d^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/b^{(1/2)}/(c+d*\tan(f*x+ \\
& e))^{(1/2)})/b^{(7/2)}/d^{(1/2)}/f+1/8*(8*b^2*d*(B*c+(A-C)*d)+(-a*d+b*c)*(6*B*b* \\
& d-5*C*a*d+5*C*b*c))*(a+b*\tan(f*x+e))^{(1/2)}*(c+d*\tan(f*x+e))^{(1/2)}/b^3/f+1/ \\
& 12*(6*B*b*d-5*C*a*d+5*C*b*c)*(a+b*\tan(f*x+e))^{(1/2)}*(c+d*\tan(f*x+e))^{(3/2)} \\
& /b^2/f+1/3*C*(a+b*\tan(f*x+e))^{(1/2)}*(c+d*\tan(f*x+e))^{(5/2)}/b/f
\end{aligned}$$
Mathematica [A] (verified)

Time = 7.81 (sec) , antiderivative size = 780, normalized size of antiderivative = 1.54

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{a+b \tan(e+fx)}} dx = \frac{C \sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))}{3bf}$$

$$+ \frac{(5bcC+6bBd-5aCd) \sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{4bf} + \frac{3(8b^2d(Bc+(A-C)d)+(bc-ad)(5bcC+6bBd-5aCd) \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)})}{4bf}$$

input

```
Integrate[(((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[a + b*Tan[e + f*x]]),x]
```

output

```
(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2))/(3*b*f) + (((5*b*c
*C + 6*b*B*d - 5*a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2
))/ (4*b*f) + ((3*(8*b^2*d*(B*c + (A - C)*d) + (b*c - a*d)*(5*b*c*C + 6*b*B
*d - 5*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(4*b*f)
+ ((6*b^3*(b*(A - C)*d*(3*c^2 - d^2) + b*B*(c^3 - 3*c*d^2) + Sqrt[-b^2]*(A
*c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3))*ArcTanh[(Sqrt[-
c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqr
t[c + d*Tan[e + f*x]])]/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b
]) - (6*b^3*(b*(A - C)*d*(3*c^2 - d^2) + b*B*(c^3 - 3*c*d^2) - Sqrt[-b^2]*
(A*c^3 - c^3*C - 3*B*c^2*d - 3*A*c*d^2 + 3*c*C*d^2 + B*d^3))*ArcTanh[(Sqrt
[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqr
t[c + d*Tan[e + f*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b])
- (3*Sqrt[b]*Sqrt[c - (a*d)/b]*(5*a^3*C*d^3 - 3*a^2*b*d^2*(5*c*C + 2*B*d)
+ a*b^2*d*(15*c^2*C + 20*B*c*d + 8*(A - C)*d^2) - b^3*(5*c^3*C + 30*B*c^2
*d + 40*c*(A - C)*d^2 - 16*B*d^3))*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x
]])/(Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*c + b*d*Tan[e + f*x])/(b*c - a*d)
)]/(4*Sqrt[d]*Sqrt[c + d*Tan[e + f*x]])/(b^2*f)/(2*b)/(3*b)
```

Rubi [A] (verified)

Time = 5.87 (sec) , antiderivative size = 524, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

↓ 4130

$$\int \frac{(c + d \tan(e + fx))^{3/2} ((5bcC - 5adC + 6bBd) \tan^2(e + fx) + 6b(Bc + (A - C)d) \tan(e + fx) + 6Abc - C(bc + 5ad))}{2\sqrt{a + b \tan(e + fx)}} dx + \frac{3b}{3bf} \sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}$$

$$3 \left(\frac{\int \frac{16((A-C)d(3c^2-d^2)+B(c^3-3cd^2)) \tan(e+fx)b^3+2c(8Ac^2b^2-c(3cC+2Bd)b^2-2ad(5cC+3Bd)b+5a^2Cd^2)b+(16d(2c(A-C)d+B(c^2-d^2)))b^3+(bc-ad)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{\dots} \right)$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3bf}$$

↓ 27

$$3 \left(\frac{\int \frac{16((A-C)d(3c^2-d^2)+B(c^3-3cd^2)) \tan(e+fx)b^3+2c(8Ac^2b^2-c(3cC+2Bd)b^2-2ad(5cC+3Bd)b+5a^2Cd^2)b+(16d(2c(A-C)d+B(c^2-d^2)))b^3+(bc-ad)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{\dots} \right)$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3bf}$$

↓ 3042

$$3 \left(\frac{\int \frac{16((A-C)d(3c^2-d^2)+B(c^3-3cd^2)) \tan(e+fx)b^3+2c(8Ac^2b^2-c(3cC+2Bd)b^2-2ad(5cC+3Bd)b+5a^2Cd^2)b+(16d(2c(A-C)d+B(c^2-d^2)))b^3+(bc-ad)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{\dots} \right)$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3bf}$$

↓ 4138

$$3 \left(\frac{\int \frac{16((A-C)d(3c^2-d^2)+B(c^3-3cd^2)) \tan(e+fx)b^3+2c(8Ac^2b^2-c(3cC+2Bd)b^2-2ad(5cC+3Bd)b+5a^2Cd^2)b+(16d(2c(A-C)d+B(c^2-d^2)))b^3+(bc-ad)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{\dots} \right)$$

$$\frac{C\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}}{3bf}$$

↓ 2348

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3bf} + \frac{(-5aCd+6bBd+5bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} + \frac{3\left(\frac{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}((bc-ad)(-5aCd+6bBd+5bcC)+8b^2d(A-C)+}{bf}\right)}{1}$$

↓ 2009

$$\frac{C\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{5/2}}{3bf} + \frac{(-5aCd+6bBd+5bcC)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} + \frac{3\left(\frac{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}((bc-ad)(-5aCd+6bBd+5bcC)+8b^2d(A-C)+}{bf}\right)}{1}$$

input `Int(((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[a + b*Tan[e + f*x]],x)`

output `(C*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2))/(3*b*f) + (((5*b*c*C + 6*b*B*d - 5*a*C*d)*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(2*b*f) + (3*(((-16*b^3*(I*A + B - I*C)*(c - I*d)^(5/2)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b] - (16*b^3*(B - I*(A - C))*(c + I*d)^(5/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] - (2*(5*a^3*C*d^3 - 3*a^2*b*d^2*(5*c*C + 2*B*d) + a*b^2*d*(15*c^2*C + 20*B*c*d + 8*(A - C)*d^2) - b^3*(5*c^3*C + 30*B*c^2*d + 40*c*(A - C)*d^2 - 16*B*d^3))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[b]*Sqrt[d]))/(2*b*f) + ((8*b^2*d*(B*c + (A - C)*d) + (b*c - a*d)*(5*b*c*C + 6*b*B*d - 5*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]/(b*f)))/(4*b))/(6*b)`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2348 `Int[(P_x)*((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4130 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4138 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan^2(fx + e))^2}{\sqrt{a + b \tan(fx + e)}} dx$$

input `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x)`

output `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{\frac{5}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{\frac{5}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \int \frac{(c + d \tan(e + fx))^{\frac{5}{2}} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx$$

input `integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(1/2),x)`

output

```
Integral((c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)*
*2)/sqrt(a + b*tan(e + f*x)), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(1/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e) + A}{\sqrt{b \tan(fx + e) + a}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(
f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(5/
2)/sqrt(b*tan(f*x + e) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(1/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\begin{aligned} \int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{a + b \tan(e + fx)}} dx &= \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)}}{\tan(fx + e)} dx \right) \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3}{\tan(fx + e) b + a} dx \right) b d^2 \\ &+ 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^3}{\tan(fx + e) b + a} dx \right) c^2 d \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) a d^2 \\ &+ 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) b c d \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{\tan(fx + e) b + a} dx \right) c^3 \\ &+ 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) a c d \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)}{\tan(fx + e) b + a} dx \right) b c^2 \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a}}{\tan(fx + e) b + a} dx \right) a c^2 \end{aligned}$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(1/2),x)
```

output

```
int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(t
an(e + f*x)*b + a),x)*c*d**2 + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)*b + a),x)*b*d**2 + 2*int((sqrt(
tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x
)*b + a),x)*c**2*d + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a
)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x)*a*d**2 + 2*int((sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*b + a),x
)*b*c*d + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f
*x)**2)/(tan(e + f*x)*b + a),x)*c**3 + 2*int((sqrt(tan(e + f*x)*d + c)*sqr
t(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)*b + a),x)*a*c*d + int((s
qrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*
x)*b + a),x)*b*c**2 + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b +
a))/(tan(e + f*x)*b + a),x)*a*c**2
```

3.143
$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx$$

Optimal result	1609
Mathematica [B] (verified)	1610
Rubi [A] (verified)	1611
Maple [F(-1)]	1616
Fricas [F(-1)]	1617
Sympy [F]	1617
Maxima [F(-1)]	1617
Giac [F]	1618
Mupad [F(-1)]	1618
Reduce [F]	1618

Optimal result

Integrand size = 49, antiderivative size = 535

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{3/2}} dx =$$

$$-\frac{(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{3/2} f}$$

$$-\frac{(B-i(A-C))(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{3/2} f}$$

$$+\frac{\sqrt{d}(15a^2Cd^2-6abd(5cC+2Bd)+b^2(15c^2C+20Bcd+8(A-C)d^2)) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{4b^{7/2} f}$$

$$-\frac{d(15a^3Cd-8Ab^2(bc-ad)-3a^2b(5cC+4Bd)-b^3(7cC+4Bd)+ab^2(8Bc+7Cd)) \sqrt{a+b \tan(e+fx)}}{4b^3(a^2+b^2) f}$$

$$+\frac{(4Ab^2-4abB+5a^2C+b^2C) d \sqrt{a+b \tan(e+fx)} (c+d \tan(e+fx))^{3/2}}{2b^2(a^2+b^2) f}$$

$$-\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{b(a^2+b^2) f \sqrt{a+b \tan(e+fx)}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(3/2)/f-(B-I*(A-C))*(c+I*d)^(5
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(3/2)/f+1/4*d^(1/2)*(15*a^2*C*d^2-6*a*b*d*(2*B*d+5*C*
c)+b^2*(15*c^2*C+20*B*c*d+8*(A-C)*d^2))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(
1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(7/2)/f-1/4*d*(15*a^3*C*d-8*A*b^2*(
-a*d+b*c)-3*a^2*b*(4*B*d+5*C*c)-b^3*(4*B*d+7*C*c)+a*b^2*(8*B*c+7*C*d))*(a+
b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/b^3/(a^2+b^2)/f+1/2*(4*A*b^2-4*
B*a*b+5*C*a^2+C*b^2)*d*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(3/2)/b^2/(
a^2+b^2)/f-2*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(5/2)/b/(a^2+b^2)/f/(a+b
*tan(f*x+e))^(1/2)

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1774 vs. $2(535) = 1070$.

Time = 7.61 (sec) , antiderivative size = 1774, normalized size of antiderivative = 3.32

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^(3/2),x]

```

output

```
(C*(c + d*Tan[e + f*x])^(5/2))/(2*b*f*Sqrt[a + b*Tan[e + f*x]]) + (((5*b*c
*C + 4*b*B*d - 5*a*C*d)*(c + d*Tan[e + f*x])^(3/2))/(2*b*f*Sqrt[a + b*Tan[
e + f*x]]) + ((8*b^2*(I*A + B - I*C)*(-c + I*d)^(5/2)*ArcTanh[(Sqrt[-c + I
*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/
((-a + I*b)^(3/2)*f) - (8*b^2*(B - I*(A - C))*(c + I*d)^(5/2)*ArcTanh[(Sqr
t[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x
]])])/((a + I*b)^(3/2)*f) + (8*b^2*(I*A + B - I*C)*(c - I*d)^2*Sqrt[c + d*
Tan[e + f*x]])/((a - I*b)*f*Sqrt[a + b*Tan[e + f*x]]) + (8*b^2*(A + I*B -
C)*(c + I*d)^2*Sqrt[c + d*Tan[e + f*x]])/((I*a - b)*f*Sqrt[a + b*Tan[e + f
*x]]) + (30*a^2*C*d^2*Sqrt[c + d*Tan[e + f*x]]*(1 + (b*d*(a + b*Tan[e + f*
x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))))^(3/2)*(1 -
(Sqrt[b]*Sqrt[d]*ArcSinh[(Sqrt[b]*Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt
[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)])]*Sqrt[a + b*T
an[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a
*d)]*Sqrt[1 + (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d)
- (a*b*d)/(b*c - a*d))])))/((b*Sqrt[b/((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c
- a*d))]*f*Sqrt[a + b*Tan[e + f*x]]*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c -
a*d)]*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) -
(a*b*d)/(b*c - a*d))))) - (12*a*d*(5*c*C + 2*B*d)*Sqrt[c + d*Tan[e + f*x]
]*(1 + (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (...
```

Rubi [A] (verified)

Time = 7.25 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4128, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$2 \int \frac{(c+d \tan(e+fx))^{3/2} ((5Ca^2-4bBa+4Ab^2+b^2C)d \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-5ad)+Ab(ac+5bd))}{2\sqrt{a+b \tan(e+fx)} b(a^2+b^2)}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 27

$$\int \frac{(c+d \tan(e+fx))^{3/2} ((5Ca^2-4bBa+4Ab^2+b^2C)d \tan^2(e+fx)-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-5ad)+Ab(ac+5bd))}{\sqrt{a+b \tan(e+fx)} b(a^2+b^2)}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^{3/2} ((5Ca^2-4bBa+4Ab^2+b^2C)d \tan^2(e+fx)^2-b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(bc-5ad)+Ab(ac+5bd))}{\sqrt{a+b \tan(e+fx)} b(a^2+b^2)}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 4130

$$\int -\frac{\sqrt{c+d \tan(e+fx)}(-4(2aAcd-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2-4c((bB-aC)(bc-5ad)+Ab(ac+5bd))b+d(15Cda^3-3b^2d^2))}{2\sqrt{a+b \tan(e+fx)} 2b}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 27

$$\frac{d(5a^2C-4abB+4Ab^2+b^2C)\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}}{2bf} - \int \frac{\sqrt{c+d \tan(e+fx)}(-4(2aAcd-2acCd-Ab(c^2-d^2)+aB(c^2-d^2)+b(Cc^2+2Bdc-Cd^2)) \tan(e+fx)b^2-4c((bB-aC)(bc-5ad)+Ab(ac+5bd))b+d(15Cda^3-3b^2d^2))}{2b}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

↓ 3042

$$\frac{d(5a^2C-4abB+4Ab^2+b^2C)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \int \frac{\sqrt{c+d\tan(e+fx)}(-4(2aAcd-2acCd-Ab(c^2-d^2))+aB(c^2-d^2)+b(Cc^2+2Bd))}{bf}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 4130

$$\frac{d(5a^2C-4abB+4Ab^2+b^2C)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \int \frac{15Cd^3a^4-6bd^2(5cC+2Bd)a^3+b^2d(15C^2+20Bdc+(8A+7C)d^2)a^2-2b^3(4C^2+2Bd)}{bf}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 27

$$\frac{d(5a^2C-4abB+4Ab^2+b^2C)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \frac{d\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(15a^3Cd-3a^2b(4Bd+5cC)-8Ab^2(bc-ad)+4a^2b^2)}{bf}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 3042

$$\frac{d(5a^2C-4abB+4Ab^2+b^2C)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \frac{d\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(15a^3Cd-3a^2b(4Bd+5cC)-8Ab^2(bc-ad)+4a^2b^2)}{bf}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 4138

$$\frac{d(5a^2C-4abB+4Ab^2+b^2C)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \frac{d\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(15a^3Cd-3a^2b(4Bd+5cC)-8Ab^2(bc-ad)+4a^2b^2)}{bf}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{bf (a^2 + b^2) \sqrt{a + b \tan(e + fx)}}$$

↓ 2348

$$\frac{(5Ca^2 - 4bBa + 4Ab^2 + b^2C)d\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \frac{d(15Cda^3 - 3b(5cC + 4Bd)a^2 + b^2(8Bc + 7Cd)a - 8Ab^2(bc - ad) - b^3(7cC + 4Bd))\sqrt{a+b\tan(e+fx)}}{bf}$$

$$\frac{2(Ab^2 - a(bB - aC))(c + d\tan(e + fx))^{5/2}}{b(a^2 + b^2)f\sqrt{a + b\tan(e + fx)}}$$

↓ 2009

$$- \frac{2(Ab^2 - a(bB - aC))(c + d\tan(e + fx))^{5/2}}{bf(a^2 + b^2)\sqrt{a + b\tan(e + fx)}} +$$

$$\frac{d(5a^2C - 4abB + 4Ab^2 + b^2C)\sqrt{a+b\tan(e+fx)}(c+d\tan(e+fx))^{3/2}}{2bf} - \frac{d\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}(15a^3Cd - 3a^2b(4Bd + 5cC) - 8Ab^2(bc - ad) + ab^3(7cC + 4Bd))}{bf}$$

input

```
Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(3/2),x]
```

output

```
(-2*(A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(5/2))/(b*(a^2 + b^2)*f*Sqrt[a + b*Tan[e + f*x]]) + (((4*A*b^2 - 4*a*b*B + 5*a^2*C + b^2*C)*d*Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2))/(2*b*f) - (-1/2*((-8*(a + I*b)*b^3*(I*A + B - I*C)*(c - I*d)^(5/2)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b] + (8*b^3*(I*a + b)*(A + I*B - C)*(c + I*d)^(5/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] + (2*(a^2 + b^2)*Sqrt[d]*(15*a^2*C*d^2 - 6*a*b*d*(5*c*C + 2*B*d) + b^2*(15*c^2*C + 20*B*c*d + 8*(A - C)*d^2))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[b])/(b*f) + (d*(15*a^3*C*d - 8*A*b^2*(b*c - a*d) - 3*a^2*b*(5*c*C + 4*B*d) - b^3*(7*c*C + 4*B*d) + a*b^2*(8*B*c + 7*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f))/(4*b)/(b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 2348 $\text{Int}[(Px_)*((c_) + (d_)*(x_))^{(m_)*((e_) + (f_)*(x_))^{(n_)*((a_) + (b_)*(x_)^2)^{(p_)}}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4128 $\text{Int}[((a_) + (b_)*\tan[(e_) + (f_)*(x_)])^{(m_)*((c_) + (d_)*\tan[(e_) + (f_)*(x_)])^{(n_)*((A_) + (B_)*\tan[(e_) + (f_)*(x_)] + (C_)*\tan[(e_) + (f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[(A*d^2 + c*(c*C - B*d))*(a + b*\tan[e + f*x])^m*((c + d*\tan[e + f*x])^{(n + 1)}/(d*f*(n + 1)*(c^2 + d^2))), x] - \text{Simp}[1/(d*(n + 1)*(c^2 + d^2)) \ \text{Int}[(a + b*\tan[e + f*x])^{(m - 1)*(c + d*\tan[e + f*x])^{(n + 1)*\text{Simp}[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*\tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*\tan[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[n, -1]$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(a + b \tan(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(3/2),x)
```

output

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(3/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(3/2),x)
```

output

```
Integral((c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x))**(3/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e)}{(b \tan(fx + e))^{3/2}}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(5/2)/(b*tan(f*x + e) + a)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input `int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(3/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2),x)`

output

```
(4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*c*d**2 + 2*sqrt(tan
(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b*c**2*d + int((sqrt(tan(e + f*x
)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(tan(e + f*x)**2*b**2 +
2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b*c*d**3*f - int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(tan(e + f*x)**2*
b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*b**2*c**2*d**2*f + int((
sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(tan(e
+ f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*c*d**3*f - int((sqrt(t
an(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(tan(e + f*x)
**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a*b*c**2*d**2*f + int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*
b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**2*d**3*f + 2*int((s
qrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e +
f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b*c**2*d**2*f
- int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)
/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*b**3*c
*d**2*f - 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)**3)/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*
x)*b**2*c**3*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*
tan(e + f*x)**3)/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*...
```


3.144
$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx$$

Optimal result	1620
Mathematica [C] (verified)	1621
Rubi [A] (verified)	1622
Maple [F(-1)]	1627
Fricas [F(-1)]	1628
Sympy [F]	1628
Maxima [F(-1)]	1628
Giac [F]	1629
Mupad [F(-1)]	1629
Reduce [F]	1629

Optimal result

Integrand size = 49, antiderivative size = 545

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx =$$

$$-\frac{(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{5/2} f}$$

$$-\frac{(B-i(A-C))(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{5/2} f}$$

$$+\frac{d^{3/2}(5bcC+2bBd-5aCd) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{b^{7/2} f}$$

$$-\frac{d(2a^3bBd-5a^4Cd-2ab^3(2Ac-2cC-3Bd)+2a^2b^2(Bc-5Cd)-b^4(2Bc+(4A+C)d)) \sqrt{a+b \tan(e+fx)}}{b^3(a^2+b^2)^2 f}$$

$$+\frac{2(2a^3bBd-5a^4Cd-b^4(3Bc+5Ad)-2ab^3(3Ac-3cC-4Bd)+a^2b^2(3Bc+(A-11C)d))(c+d \tan(e+fx))}{3b^2(a^2+b^2)^2 f \sqrt{a+b \tan(e+fx)}}$$

$$-\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{3b(a^2+b^2) f(a+b \tan(e+fx))^{3/2}}$$

output

$$\begin{aligned}
& -(I*A+B-I*C)*(c-I*d)^{(5/2)}*\operatorname{arctanh}((c-I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/(a \\
& -I*b)^{(1/2)}/(c+d*\tan(f*x+e))^{(1/2)})/(a-I*b)^{(5/2)}/f-(B-I*(A-C))*(c+I*d)^{(5 \\
& /2)}*\operatorname{arctanh}((c+I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/(a+I*b)^{(1/2)}/(c+d*\tan(f* \\
& x+e))^{(1/2)})/(a+I*b)^{(5/2)}/f+d^{(3/2)}*(2*B*b*d-5*C*a*d+5*C*b*c)*\operatorname{arctanh}(d^{(\\
& 1/2)}*(a+b*\tan(f*x+e))^{(1/2)}/b^{(1/2)}/(c+d*\tan(f*x+e))^{(1/2)})/b^{(7/2)}/f-d*(2 \\
& *a^3*b*B*d-5*a^4*C*d-2*a*b^3*(2*A*c-3*B*d-2*C*c)+2*a^2*b^2*(B*c-5*C*d)-b^4 \\
& *(2*B*c+(4*A+C)*d))*(a+b*\tan(f*x+e))^{(1/2)}*(c+d*\tan(f*x+e))^{(1/2)}/b^3/(a^2 \\
& +b^2)^2/f+2/3*(2*a^3*b*B*d-5*a^4*C*d-b^4*(5*A*d+3*B*c)-2*a*b^3*(3*A*c-4*B* \\
& d-3*C*c)+a^2*b^2*(3*B*c+(A-11*C)*d))*(c+d*\tan(f*x+e))^{(3/2)}/b^2/(a^2+b^2)^ \\
& 2/f/(a+b*\tan(f*x+e))^{(1/2)}-2/3*(A*b^2-a*(B*b-C*a))*(c+d*\tan(f*x+e))^{(5/2)}/ \\
& b/(a^2+b^2)/f/(a+b*\tan(f*x+e))^{(3/2)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.99 (sec) , antiderivative size = 802, normalized size of antiderivative = 1.47

$$\begin{aligned}
& \int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{5/2}} dx = \frac{C(c+d \tan(e+fx))^{5/2}}{bf(a+b \tan(e+fx))^{3/2}} \\
& - \frac{2b(A-iB-C)(c-id)(c+d \tan(e+fx))^{3/2}}{3(ia+b)f(a+b \tan(e+fx))^{3/2}} + \frac{2b(A+iB-C)(c+id)(c+d \tan(e+fx))^{3/2}}{3(ia-b)f(a+b \tan(e+fx))^{3/2}} - \frac{10cC(bc-ad) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, -\frac{b \tan(e+fx)}{a+b \tan(e+fx)}\right)}{3bf(a+b \tan(e+fx))^{3/2} \sqrt{b}}
\end{aligned}$$

input

$$\text{Integrate}[\frac{(c+d*\text{Tan}[e+f*x])^{(5/2)}*(A+B*\text{Tan}[e+f*x]+C*\text{Tan}[e+f*x]^2)}{(a+b*\text{Tan}[e+f*x])^{(5/2)}},x]$$

output

```
(C*(c + d*Tan[e + f*x])^(5/2))/(b*f*(a + b*Tan[e + f*x])^(3/2)) + ((-2*b*(
A - I*B - C)*(c - I*d)*(c + d*Tan[e + f*x])^(3/2))/(3*(I*a + b)*f*(a + b*T
an[e + f*x])^(3/2)) + (2*b*(A + I*B - C)*(c + I*d)*(c + d*Tan[e + f*x])^(3
/2))/(3*(I*a - b)*f*(a + b*Tan[e + f*x])^(3/2)) - (10*c*C*(b*c - a*d)*Hype
rgeometric2F1[-3/2, -3/2, -1/2, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d))]*S
qrt[c + d*Tan[e + f*x]])/(3*b*f*(a + b*Tan[e + f*x])^(3/2)*Sqrt[(b*(c + d*
Tan[e + f*x]))/(b*c - a*d)]) - (4*B*d*(b*c - a*d)*Hypergeometric2F1[-3/2,
-3/2, -1/2, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d))]*Sqrt[c + d*Tan[e + f
x]])/(3*b*f*(a + b*Tan[e + f*x])^(3/2)*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c
- a*d)]) + (10*a*C*d*(b*c - a*d)*Hypergeometric2F1[-3/2, -3/2, -1/2, -((d*
(a + b*Tan[e + f*x]))/(b*c - a*d))]*Sqrt[c + d*Tan[e + f*x]])/(3*b^2*f*(a
+ b*Tan[e + f*x])^(3/2)*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)]) + (2*b
*(I*A + B - I*C)*(c - I*d)^2*((Sqrt[-c + I*d]*ArcTanh[(Sqrt[-c + I*d]*Sqrt
[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(-a + I*
b)^(3/2) + Sqrt[c + d*Tan[e + f*x]]/((a - I*b)*Sqrt[a + b*Tan[e + f*x]]))
/((a - I*b)*f) - (2*b*(A + I*B - C)*(c + I*d)^2*((Sqrt[c + I*d]*ArcTanh[(S
qrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f
*x]])])/((a + I*b)^(3/2) - Sqrt[c + d*Tan[e + f*x]]/((a + I*b)*Sqrt[a + b*T
an[e + f*x]])))/((I*a - b)*f))/(2*b)
```

Rubi [A] (verified)

Time = 8.33 (sec) , antiderivative size = 603, normalized size of antiderivative = 1.11, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$2 \int \frac{(c+d \tan(e+fx))^{3/2} ((5Ca^2-2bBa+2Ab^2+3b^2C)d \tan^2(e+fx)-3b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(3bc-5ad)+Ab(3ac+3bd))}{2(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}$$

$$\frac{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{(c+d \tan(e+fx))^{3/2} ((5Ca^2-2bBa+2Ab^2+3b^2C)d \tan^2(e+fx)-3b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(3bc-5ad)+Ab(3ac+3bd))}{(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}$$

$$\frac{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^{3/2} ((5Ca^2-2bBa+2Ab^2+3b^2C)d \tan^2(e+fx)-3b((A-C)(bc-ad)-B(ac+bd)) \tan(e+fx)+(bB-aC)(3bc-5ad)+Ab(3ac+3bd))}{(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3b(a^2+b^2)}{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}$$

$$\frac{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 4128

$$2 \int \frac{\sqrt{c+d \tan(e+fx)} (-3((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+3bd)((bB-aC)(3bc-5ad)+Ab(3ac+3bd))}{(a+b \tan(e+fx))^{3/2}}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)} (-3((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+3bd)((bB-aC)(3bc-5ad)+Ab(3ac+3bd))}{(a+b \tan(e+fx))^{3/2}}$$

$$\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{3bf(a^2+b^2)(a+b \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-3((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(ac+3bd)((bB-aC)(3bc-5ad)+Ab(3ac+ \right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3bf (a^2 + b^2) (a + b \tan(e + fx))^{3/2}}$$

↓ 4130

$$\int - \frac{3 \left(5Cd^3a^5 - bd^2(5cC+2Bd)a^4 + 10b^2Cd^3a^3 - 2b^3(Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2c + 8Cd^2c + 3Bd^3)a^2 - b^4(4Ad(3c^2 - d^2) - Cd(12c^2 + d^2) + 4B(c^3 - 3cd^2))a - (a^2 + \right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3bf (a^2 + b^2) (a + b \tan(e + fx))^{3/2}}$$

↓ 27

$$3 \int \frac{5Cd^3a^5 - bd^2(5cC+2Bd)a^4 + 10b^2Cd^3a^3 - 2b^3(Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2c + 8Cd^2c + 3Bd^3)a^2 - b^4(4Ad(3c^2 - d^2) - Cd(12c^2 + d^2) + 4B(c^3 - 3cd^2))a - (a^2 + \right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3bf (a^2 + b^2) (a + b \tan(e + fx))^{3/2}}$$

↓ 3042

$$3 \int \frac{5Cd^3a^5 - bd^2(5cC+2Bd)a^4 + 10b^2Cd^3a^3 - 2b^3(Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2c + 8Cd^2c + 3Bd^3)a^2 - b^4(4Ad(3c^2 - d^2) - Cd(12c^2 + d^2) + 4B(c^3 - 3cd^2))a - (a^2 + \right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3bf (a^2 + b^2) (a + b \tan(e + fx))^{3/2}}$$

↓ 4138

$$3 \int \frac{5Cd^3a^5 - bd^2(5cC+2Bd)a^4 + 10b^2Cd^3a^3 - 2b^3(Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2c + 8Cd^2c + 3Bd^3)a^2 - b^4(4Ad(3c^2 - d^2) - Cd(12c^2 + d^2) + 4B(c^3 - 3cd^2))a - (a^2 + \right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3bf (a^2 + b^2) (a + b \tan(e + fx))^{3/2}}$$

↓ 2348

$$\frac{2(-5Cda^4+2bBda^3+b^2(3Bc+(A-11C)d)a^2-2b^3(3Ac-3Cc-4Bd)a-b^4(3Bc+5Ad))(c+d \tan(e+fx))^{3/2}}{b(a^2+b^2)f\sqrt{a+b \tan(e+fx)}} + \frac{-3d\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{b(a^2+b^2)f\sqrt{a+b \tan(e+fx)}}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3b (a^2 + b^2) f (a + b \tan(e + fx))^{3/2}}$$

↓ 2009

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{3bf (a^2 + b^2) (a + b \tan(e + fx))^{3/2}} +$$

$$\frac{2(c+d \tan(e+fx))^{3/2}(-5a^4Cd+2a^3bBd+a^2b^2(d(A-11C)+3Bc)-2ab^3(3Ac-4Bd-3Cc)-b^4(5Ad+3Bc))}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}} + \frac{-3d\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{bf(a^2+b^2)\sqrt{a+b \tan(e+fx)}}$$

input

```
Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(5/2),x]
```

output

```
(-2*(A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(5/2))/(3*b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^(3/2)) + ((2*(2*a^3*b*B*d - 5*a^4*C*d - b^4*(3*B*c + 5*A*d) - 2*a*b^3*(3*A*c - 3*c*C - 4*B*d) + a^2*b^2*(3*B*c + (A - 11*C)*d))*(c + d*Tan[e + f*x])^(3/2))/(b*(a^2 + b^2)*f*Sqrt[a + b*Tan[e + f*x]]) + ((-3*((2*(a + I*b)^2*b^3*(B + I*(A - C))*(c - I*d)^(5/2)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b] - (2*(a - I*b)^2*b^3*(I*A - B - I*C)*(c + I*d)^(5/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] - (2*(a^2 + b^2)^2*d^(3/2)*(5*b*c*C + 2*b*B*d - 5*a*C*d)*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[b]))/(2*b*f) - (3*d*(2*a^3*b*B*d - 5*a^4*C*d - 2*a*b^3*(2*A*c - 2*c*C - 3*B*d) + 2*a^2*b^2*(B*c - 5*C*d) - b^4*(2*B*c + (4*A + C)*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(b*f))/(b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2348 `Int[(Px)*((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4128 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4130

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

rule 4138

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]

```

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(a + b \tan(fx + e))^{\frac{5}{2}}} dx$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(5/2),x)
```

output

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(5/2),x)
```


Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(5/2),x)
```

output

```
Integral((c + d*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(a + b*tan(e + f*x))**(5/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e)}{(b \tan(fx + e))^{5/2}}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(5/2)/(b*tan(f*x + e) + a)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input `int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(5/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{5/2}} dx = \int \frac{(d \tan(fx + e) + c)^{5/2} (A + B \tan(fx + e))}{(\tan(fx + e))^{5/2}}$$

input `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x)`

output `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2),x)`

3.145
$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx$$

Optimal result	1631
Mathematica [C] (verified)	1632
Rubi [A] (verified)	1633
Maple [F(-1)]	1638
Fricas [F(-1)]	1638
Sympy [F(-1)]	1638
Maxima [F(-1)]	1639
Giac [F]	1639
Mupad [F(-1)]	1639
Reduce [F]	1640

Optimal result

Integrand size = 49, antiderivative size = 590

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{7/2}} dx =$$

$$\frac{(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{7/2} f}$$

$$- \frac{(B-i(A-C))(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{7/2} f}$$

$$+ \frac{2Cd^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{b^{7/2} f}$$

$$- \frac{2(a^6Cd^2+3a^4b^2Cd^2-3a^2b^4(c^2C+2Bcd-2Cd^2-A(c^2-d^2))+b^6(c(cC+2Bd)-A(c^2-d^2))-a^3b^3(a^2+b^2)^3 f \sqrt{a+b \tan(e+fx)})}{3b^2(a^2+b^2)^2 f(a+b \tan(e+fx))^{3/2}}$$

$$- \frac{2(a^4Cd+b^4(Bc+Ad)+2ab^3(Ac-cC-Bd)-a^2b^2(Bc+(A-3C)d))(c+d \tan(e+fx))^{3/2}}{5b(a^2+b^2) f(a+b \tan(e+fx))^{5/2}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(7/2)/f-(B-I*(A-C))*(c+I*d)^(5
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(7/2)/f+2*C*d^(5/2)*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(
1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(7/2)/f-2*(a^6*C*d^2+3*a^4*b^2*C*d
^2-3*a^2*b^4*(c^2*C+2*B*c*d-2*C*d^2-A*(c^2-d^2))+b^6*(c*(2*B*d+C*c)-A*(c^2
-d^2))-a^3*b^3*(2*c*(A-C)*d+B*(c^2-d^2))+3*a*b^5*(2*c*(A-C)*d+B*(c^2-d^2))
)*(c+d*tan(f*x+e))^(1/2)/b^3/(a^2+b^2)^3/f/(a+b*tan(f*x+e))^(1/2)-2/3*(a^4
*C*d+b^4*(A*d+B*c)+2*a*b^3*(A*c-B*d-C*c)-a^2*b^2*(B*c+(A-3*C)*d))*(c+d*tan
(f*x+e))^(3/2)/b^2/(a^2+b^2)^2/f/(a+b*tan(f*x+e))^(3/2)-2/5*(A*b^2-a*(B*b-
C*a))*(c+d*tan(f*x+e))^(5/2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^(5/2)
    
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.71 (sec) , antiderivative size = 641, normalized size of antiderivative = 1.09

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \frac{(B + i(A - C))(c + d \tan(e + fx))}{5(a - ib)f(a + b \tan(e + fx))^{5/2}}$$

$$- \frac{(iA - B - iC)(c + d \tan(e + fx))^{5/2}}{5(a + ib)f(a + b \tan(e + fx))^{5/2}}$$

$$- \frac{2C(bc - ad)^2 \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, -\frac{5}{2}, -\frac{3}{2}, -\frac{d(a + b \tan(e + fx))}{bc - ad}\right) \sqrt{c + d \tan(e + fx)}}{5b^3 f(a + b \tan(e + fx))^{5/2} \sqrt{\frac{b(c + d \tan(e + fx))}{bc - ad}}}$$

$$+ \frac{(A - iB - C)(ic + d) \left(\frac{(c + d \tan(e + fx))^{3/2}}{(a - ib)(a + b \tan(e + fx))^{3/2}} + \frac{3(c - id) \left(\frac{\sqrt{-c + id} \operatorname{arctanh}\left(\frac{\sqrt{-c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{(-a + ib)^{3/2}} + \frac{\sqrt{c + d \tan(e + fx)}}{(a - ib) \sqrt{a + b \tan(e + fx)}}\right)}{a - ib}}{3(a - ib)f} \right)}{3(a + ib)f}$$

$$+ \frac{(A + iB - C)(ic - d) \left(\frac{(c + d \tan(e + fx))^{3/2}}{(a + ib)(a + b \tan(e + fx))^{3/2}} - \frac{3(c + id) \left(\frac{\sqrt{c + id} \operatorname{arctanh}\left(\frac{\sqrt{c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{(a + ib)^{3/2}} - \frac{\sqrt{c + d \tan(e + fx)}}{(a + ib) \sqrt{a + b \tan(e + fx)}}\right)}{a + ib}}{3(a + ib)f} \right)}{3(a + ib)f}$$

input

```
Integrate[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(7/2),x]
```

output

```
((B + I*(A - C))*(c + d*Tan[e + f*x])^(5/2))/(5*(a - I*b)*f*(a + b*Tan[e + f*x])^(5/2)) - ((I*A - B - I*C)*(c + d*Tan[e + f*x])^(5/2))/(5*(a + I*b)*f*(a + b*Tan[e + f*x])^(5/2)) - (2*C*(b*c - a*d)^2*Hypergeometric2F1[-5/2, -5/2, -3/2, -(d*(a + b*Tan[e + f*x]))/(b*c - a*d)]*Sqrt[c + d*Tan[e + f*x]])/(5*b^3*f*(a + b*Tan[e + f*x])^(5/2)*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)]) + ((A - I*B - C)*(I*c + d)*((c + d*Tan[e + f*x])^(3/2)/((a - I*b)*(a + b*Tan[e + f*x])^(3/2)) + (3*(c - I*d)*((Sqrt[-c + I*d]*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(-a + I*b)^(3/2) + Sqrt[c + d*Tan[e + f*x]]/((a - I*b)*Sqrt[a + b*Tan[e + f*x]])))/(a - I*b))/((3*(a - I*b)*f) - ((A + I*B - C)*(I*c - d)*((c + d*Tan[e + f*x])^(3/2)/((a + I*b)*(a + b*Tan[e + f*x])^(3/2)) - (3*(c + I*d)*((Sqrt[c + I*d]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((a + I*b)^(3/2) - Sqrt[c + d*Tan[e + f*x]]/((a + I*b)*Sqrt[a + b*Tan[e + f*x]])))/(a + I*b))/((3*(a + I*b)*f)
```

Rubi [A] (verified)

Time = 8.51 (sec) , antiderivative size = 658, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4128, 27, 3042, 4128, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{7/2}} dx$$

↓ 4128

$$2 \int \frac{5(c+d \tan(e+fx))^{3/2} ((a^2+b^2)Cd \tan^2(e+fx) - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd))}{2(a+b \tan(e+fx))^{5/2}} dx$$

$$\frac{5b(a^2+b^2)}{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}} \\ \frac{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 27

$$\int \frac{(c+d \tan(e+fx))^{3/2} ((a^2+b^2)Cd \tan^2(e+fx) - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd))}{(a+b \tan(e+fx))^{5/2}} dx$$

$$\frac{b(a^2+b^2)}{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}} \\ \frac{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 3042

$$\int \frac{(c+d \tan(e+fx))^{3/2} ((a^2+b^2)Cd \tan(e+fx)^2 - b((A-C)(bc-ad) - B(ac+bd)) \tan(e+fx) + (bB-aC)(bc-ad) + Ab(ac+bd))}{(a+b \tan(e+fx))^{5/2}} dx$$

$$\frac{b(a^2+b^2)}{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}} \\ \frac{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 4128

$$2 \int \frac{\sqrt{c+d \tan(e+fx)} \left(-((ac+bd)((A-C)(bc-ad) - B(ac+bd)) + (bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd)) \tan(e+fx)b^2) + (ac+bd)((bB-aC)(bc-ad) + Ab(ac+bd)) \right)}{2(a+b \tan(e+fx))^{3/2} 3b(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-((ac+bd)((A-C)(bc-ad) - B(ac+bd)) + (bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd)) \tan(e+fx)b^2) + (ac+bd)((bB-aC)(bc-ad) + Ab(ac+bd)) \right)}{(a+b \tan(e+fx))^{3/2} b(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5bf(a^2+b^2)(a+b \tan(e+fx))^{5/2}}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd)) \tan(e+fx)b^2) + (ac+bd)((bB-aC)(bc-ad)+Ab(ac+bd)) \right)}{(a+b \tan(e+fx))^{3/2} b(a^2+b^2)}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5bf (a^2 + b^2) (a + b \tan(e + fx))^{5/2}}$$

↓ 4128

$$2 \int \frac{Cd^3 a^6 + 3b^2 Cd^3 a^4 - b^3 (Cc^3 + 3Bdc^2 - 3Cd^2 c - Bd^3 - A(c^3 - 3cd^2)) a^3 + 3b^4 (Bc^3 + 3Adc^2 - 3Cdc^2 - 3Bd^2 c - Ad^3 + 2Cd^3) a^2 - 3b^5 (Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2 c + 3Cd^3)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5b (a^2 + b^2) f(a + b \tan(e + fx))^{5/2}}$$

↓ 27

$$\int \frac{Cd^3 a^6 + 3b^2 Cd^3 a^4 - b^3 (Cc^3 + 3Bdc^2 - 3Cd^2 c - Bd^3 - A(c^3 - 3cd^2)) a^3 + 3b^4 (Bc^3 + 3Adc^2 - 3Cdc^2 - 3Bd^2 c - Ad^3 + 2Cd^3) a^2 - 3b^5 (Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2 c + 3Cd^3)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5b (a^2 + b^2) f(a + b \tan(e + fx))^{5/2}}$$

↓ 3042

$$\int \frac{Cd^3 a^6 + 3b^2 Cd^3 a^4 - b^3 (Cc^3 + 3Bdc^2 - 3Cd^2 c - Bd^3 - A(c^3 - 3cd^2)) a^3 + 3b^4 (Bc^3 + 3Adc^2 - 3Cdc^2 - 3Bd^2 c - Ad^3 + 2Cd^3) a^2 - 3b^5 (Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2 c + 3Cd^3)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5b (a^2 + b^2) f(a + b \tan(e + fx))^{5/2}}$$

↓ 4138

$$\int \frac{Cd^3 a^6 + 3b^2 Cd^3 a^4 - b^3 (Cc^3 + 3Bdc^2 - 3Cd^2 c - Bd^3 - A(c^3 - 3cd^2)) a^3 + 3b^4 (Bc^3 + 3Adc^2 - 3Cdc^2 - 3Bd^2 c - Ad^3 + 2Cd^3) a^2 - 3b^5 (Ac^3 - Cc^3 - 3Bdc^2 - 3Ad^2 c + 3Cd^3)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5b (a^2 + b^2) f(a + b \tan(e + fx))^{5/2}}$$

↓ 2348

$$\int \left(\frac{(a^2+b^2)^3 C d^3}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} + \frac{-Ac^3 b^6 - Bd^3 b^6 + 3Ac d^2 b^6 - 3c C d^2 b^6 + c^3 C b^6 + 3B c^2 d b^6 + 3a B c^3 b^5 - 3a A d^3 b^5 + 3a C d^3 b^5 - 9a B c d^2 b^5 + 9a A c^2 d b^5 - 9a c^2 C d^2 b^5}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} \right) dx$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5b(a^2 + b^2) f(a + b \tan(e + fx))^{5/2}}$$

↓ 2009

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{5bf(a^2 + b^2) (a + b \tan(e + fx))^{5/2}} +$$

$$\frac{-2(c+d \tan(e+fx))^{3/2} (a^4 C d - a^2 b^2 (d(A-3C) + Bc) + 2ab^3 (Ac - Bd - cC) + b^4 (Ad + Bc))}{3bf(a^2 + b^2) (a + b \tan(e + fx))^{3/2}} + \frac{-2\sqrt{c+d \tan(e+fx)} (a^6 C d^2 + 3a^4 b^2 C d^2 - a^3 b^3 (2cd(A-3C) + b^2 (Ac - Bd - cC) + b^4 (Ad + Bc)))}{3bf(a^2 + b^2) (a + b \tan(e + fx))^{3/2}}$$

input

```
Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(7/2),x]
```

output

```
(-2*(A*b^2 - a*(b*B - a*C))*(c + d*Tan[e + f*x])^(5/2))/(5*b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^(5/2)) + ((-2*(a^4*C*d + b^4*(B*c + A*d) + 2*a*b^3*(A*c - c*C - B*d) - a^2*b^2*(B*c + (A - 3*C)*d))*(c + d*Tan[e + f*x])^(3/2))/(3*b*(a^2 + b^2)*f*(a + b*Tan[e + f*x])^(3/2)) + (((-((a + I*b)^3*b^3*(I*A + B - I*C)*(c - I*d)^(5/2)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a - I*b]) - (b^3*(I*a + b)^3*(A + I*B - C)*(c + I*d)^(5/2)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[a + I*b] + (2*(a^2 + b^2)^3*C*d^(5/2)*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[b])/(b*(a^2 + b^2)*f) - (2*(a^6*C*d^2 + 3*a^4*b^2*C*d^2 - 3*a^2*b^4*(c^2*C + 2*B*c*d - 2*C*d^2 - A*(c^2 - d^2)) + b^6*(c*(c*C + 2*B*d) - A*(c^2 - d^2)) - a^3*b^3*(2*c*(A - C)*d + B*(c^2 - d^2)) + 3*a*b^5*(2*c*(A - C)*d + B*(c^2 - d^2)))*Sqrt[c + d*Tan[e + f*x]])/(b*(a^2 + b^2)*f*Sqrt[a + b*Tan[e + f*x]])/(b*(a^2 + b^2)))/(b*(a^2 + b^2))
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 2348 $\text{Int}[(Px_)*((c_) + (d_)*(x_))^{(m_)*((e_) + (f_)*(x_))^{(n_)*((a_) + (b_)*(x_)^2)^{(p_)}}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4128 $\text{Int}[(a_*) + (b_*)\text{tan}[(e_*) + (f_)*(x_)]^{(m_)*((c_*) + (d_*)\text{tan}[(e_*) + (f_)*(x_)]^{(n_)*((A_*) + (B_*)\text{tan}[(e_*) + (f_)*(x_)] + (C_*)\text{tan}[(e_*) + (f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[(A*d^2 + c*(c*C - B*d))*(a + b*\text{Tan}[e + f*x])^m*((c + d*\text{Tan}[e + f*x])^{(n + 1)}/(d*f*(n + 1)*(c^2 + d^2))), x] - \text{Simp}[1/(d*(n + 1)*(c^2 + d^2)) \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m - 1)}*(c + d*\text{Tan}[e + f*x])^{(n + 1)}*\text{Simp}[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*\text{Tan}[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[n, -1]$
- rule 4138 $\text{Int}[(a_*) + (b_*)\text{tan}[(e_*) + (f_)*(x_)]^{(m_)*((c_*) + (d_*)\text{tan}[(e_*) + (f_)*(x_)]^{(n_)*((A_*) + (B_*)\text{tan}[(e_*) + (f_)*(x_)] + (C_*)\text{tan}[(e_*) + (f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Simp}[ff/f \ \text{Subst}[\text{Int}[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, \text{Tan}[e + f*x]/ff], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(a + b \tan(fx + e))^{\frac{7}{2}}} dx$$

input `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x)`

output `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(7/2),x)`

output `Timed out`

Maxima [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)}{(b \tan(fx + e) + a)^{7/2}}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(7/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(5/2)/(b*tan(f*x + e) + a)^(7/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(7/2),x)
```

output

\text{Hanged}

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{7/2}} dx = \text{too large to display}$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e)
)^(7/2),x)
```

output

```
(10*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*a**2
*b*d**5 + 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x
)**2*a**2*c**2*d**4 + 20*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)
*tan(e + f*x)**2*a*b**2*c*d**4 - 28*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*tan(e + f*x)**2*a*b*c**3*d**3 + 18*sqrt(tan(e + f*x)*d + c)*sq
rt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b**3*c**2*d**3 - 16*sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b**3*d**5 + 16*sqrt(tan
(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b**2*c**4*d**2 -
32*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b**2
*c**2*d**4 + 10*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)*a**3*d**5 + 90*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(
e + f*x)*a**2*b*c*d**4 + 14*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b +
a)*tan(e + f*x)*a**2*c**3*d**3 - 10*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +
f*x)*b + a)*tan(e + f*x)*a*b**2*c**2*d**3 - 40*sqrt(tan(e + f*x)*d + c)*s
qrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*b**2*d**5 - 36*sqrt(tan(e + f*x)*d
+ c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*b*c**4*d**2 - 80*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*b*c**2*d**4 + 6*sqrt(
tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**3*c**3*d**2 +
8*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**3*c*d
**4 + 22*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)...
```

3.146
$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{9/2}} dx$$

Optimal result	1641
Mathematica [C] (warning: unable to verify)	1642
Rubi [F]	1643
Maple [F(-1)]	1651
Fricas [F(-1)]	1651
Sympy [F(-1)]	1651
Maxima [F(-2)]	1652
Giac [F]	1652
Mupad [F(-1)]	1653
Reduce [F]	1653

Optimal result

Integrand size = 49, antiderivative size = 946

$$\int \frac{(c+d \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(a+b \tan(e+fx))^{9/2}} dx =$$

$$\frac{(iA+B-iC)(c-id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a-ib)^{9/2} f}$$

$$-\frac{(B-i(A-C))(c+id)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a+ib)^{9/2} f}$$

$$-\frac{2(6a^5bBd^2+15a^6Cd^2+a^4b^2d(14Bc+8Ad+37Cd)+3a^2b^4(35Ac^2-35c^2C-70Bcd-39Ad^2+54Cd^2+2(6a^7bBd^3+15a^8Cd^3+2a^6b^2d^2(7Bc+4Ad+26Cd)-2ab^7(210Ac^3-210c^3C-525Bc^2d-406Acd^2-2(2a^3bBd+5a^4Cd+b^4(7Bc+5Ad)+2ab^3(7Ac-7cC-6Bd)-a^2b^2(7Bc+9Ad-19Cd))(c+d \tan(e+fx)))^{5/2}}{35b^2(a^2+b^2)^2 f(a+b \tan(e+fx))^{5/2}}$$

$$-\frac{2(Ab^2-a(bB-aC))(c+d \tan(e+fx))^{5/2}}{7b(a^2+b^2) f(a+b \tan(e+fx))^{7/2}}$$

output

```

-(I*A+B-I*C)*(c-I*d)^(5/2)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(9/2)/f-(B-I*(A-C))*(c+I*d)^(5
/2)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(a+I*b)^(9/2)/f-2/105*(6*a^5*b*B*d^2+15*a^6*C*d^2+a^4*b^2*d*(
8*A*d+14*B*c+37*C*d)+3*a^2*b^4*(35*A*c^2-39*A*d^2-70*B*c*d-35*C*c^2+54*C*d
^2)-a^3*b^3*(98*c*(A-C)*d+B*(35*c^2-75*d^2))+a*b^5*(182*c*(A-C)*d+B*(105*c
^2-71*d^2))+b^6*(7*c*(8*B*d+5*C*c)-5*A*(7*c^2-3*d^2)))*(c+d*tan(f*x+e))^(1
/2)/b^3/(a^2+b^2)^3/f/(a+b*tan(f*x+e))^(3/2)-2/105*(6*a^7*b*B*d^3+15*a^8*C
*d^3+2*a^6*b^2*d^2*(4*A*d+7*B*c+26*C*d)-2*a*b^7*(210*A*c^3-406*A*c*d^2-525
*B*c^2*d+88*B*d^3-210*C*c^3+406*C*c*d^2)-a^4*b^4*(525*A*c^2*d-311*A*d^3+10
5*B*c^3-749*B*c*d^2-525*C*c^2*d+221*C*d^3)+2*a^2*b^6*(875*A*c^2*d-261*A*d^
3+315*B*c^3-812*B*c*d^2-875*C*c^2*d+291*C*d^3)+2*a^5*b^3*d*(56*c*(A-C)*d+B
*(35*c^2-12*d^2))-b^8*(5*d*(49*A*c^2-3*A*d^2-49*C*c^2)+7*B*(15*c^3-23*c*d^
2))-2*a^3*b^5*(210*c^3*C+700*B*c^2*d-798*C*c*d^2-317*B*d^3-42*A*(5*c^3-19*
c*d^2)))*(c+d*tan(f*x+e))^(1/2)/b^3/(a^2+b^2)^4/(-a*d+b*c)/f/(a+b*tan(f*x+
e))^(1/2)-2/35*(2*a^3*b*B*d+5*a^4*C*d+b^4*(5*A*d+7*B*c)+2*a*b^3*(7*A*c-6*B
*d-7*C*c)-a^2*b^2*(9*A*d+7*B*c-19*C*d))*(c+d*tan(f*x+e))^(3/2)/b^2/(a^2+b^
2)^2/f/(a+b*tan(f*x+e))^(5/2)-2/7*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(5/
2)/b/(a^2+b^2)/f/(a+b*tan(f*x+e))^(7/2)

```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 53.15 (sec) , antiderivative size = 2723641, normalized size of antiderivative = 2879.11

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \text{Result too large to show}$$

input

```

Integrate[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(a + b*Tan[e + f*x])^(9/2),x]

```

output

Result too large to show

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(a + b \tan(e + fx))^{9/2}} dx$$

↓ 4128

$$2 \int \frac{(c + d \tan(e + fx))^{3/2} (-((-5Ca^2 - 2bBa + 2Ab^2 - 7b^2C)d \tan^2(e + fx)) - 7b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(7bc - 5ad) + Ab^2)}{2(a + b \tan(e + fx))^{7/2}} dx$$

$$\frac{7b(a^2 + b^2)}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}} \frac{2(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}}$$

↓ 27

$$\int \frac{(c + d \tan(e + fx))^{3/2} (-((-5Ca^2 - 2bBa + 2Ab^2 - 7b^2C)d \tan^2(e + fx)) - 7b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(7bc - 5ad) + Ab^2)}{(a + b \tan(e + fx))^{7/2}} dx$$

$$\frac{7b(a^2 + b^2)}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}} \frac{2(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}}$$

↓ 3042

$$\int \frac{(c + d \tan(e + fx))^{3/2} (-((-5Ca^2 - 2bBa + 2Ab^2 - 7b^2C)d \tan^2(e + fx)^2) - 7b((A - C)(bc - ad) - B(ac + bd)) \tan(e + fx) + (bB - aC)(7bc - 5ad) + Ab^2)}{(a + b \tan(e + fx))^{7/2}} dx$$

$$\frac{7b(a^2 + b^2)}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}} \frac{2(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}}$$

↓ 4128

$$2 \int \frac{\sqrt{c + d \tan(e + fx)} (-35((ac + bd)((A - C)(bc - ad) - B(ac + bd)) + (bc - ad)(bBc + b(A - C)d + a(Ac - Cc - Bd))) \tan(e + fx) b^2 + (5ac + 3bd)((bB - aC)(7bc - 5ad) + Ab^2)}{2(a + b \tan(e + fx))^{7/2}} dx$$

$$\frac{2(Ab^2 - a(bB - aC))(c + d \tan(e + fx))^{5/2}}{7bf(a^2 + b^2)(a + b \tan(e + fx))^{7/2}}$$

↓ 27

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-35((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(5ac+3bd)((bB-aC)(7bc-5ad)+Ab(7a$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7bf (a^2 + b^2) (a + b \tan(e + fx))^{7/2}}$$

↓ 3042

$$\int \frac{\sqrt{c+d \tan(e+fx)} \left(-35((ac+bd)((A-C)(bc-ad)-B(ac+bd))+(bc-ad)(bBc+b(A-C)d+a(Ac-Cc-Bd))) \tan(e+fx)b^2+(5ac+3bd)((bB-aC)(7bc-5ad)+Ab(7a$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7bf (a^2 + b^2) (a + b \tan(e + fx))^{7/2}}$$

↓ 4128

$$2 \int \frac{15Cd^3 a^6 + 6bBd^3 a^5 + b^2 d^2 (14Bc + 8Ad + 37Cd) a^4 - b^3 (105Cc^3 + 245Bdc^2 - 203Cd^2 c - 75Bd^3 - 7A(15c^3 - 29cd^2)) a^3 + 3b^4 (35B(3c^3 - 5cd^2) + d(245Ac^2 - 245Cc^2$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b (a^2 + b^2) f (a + b \tan(e + fx))^{7/2}}$$

↓ 27

$$\int \frac{15Cd^3 a^6 + 6bBd^3 a^5 + b^2 d^2 (14Bc + 8Ad + 37Cd) a^4 - b^3 (105Cc^3 + 245Bdc^2 - 203Cd^2 c - 75Bd^3 - 7A(15c^3 - 29cd^2)) a^3 + 3b^4 (35B(3c^3 - 5cd^2) + d(245Ac^2 - 245Cc^2$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b (a^2 + b^2) f (a + b \tan(e + fx))^{7/2}}$$

↓ 3042

$$\int \frac{15Cd^3 a^6 + 6bBd^3 a^5 + b^2 d^2 (14Bc + 8Ad + 37Cd) a^4 - b^3 (105Cc^3 + 245Bdc^2 - 203Cd^2 c - 75Bd^3 - 7A(15c^3 - 29cd^2)) a^3 + 3b^4 (35B(3c^3 - 5cd^2) + d(245Ac^2 - 245Cc^2$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b (a^2 + b^2) f (a + b \tan(e + fx))^{7/2}}$$

↓ 4132

$$\frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-3111d^3))}{7b(a^2+b^2)f(a+b \tan(e+fx))^{7/2}}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 27

$$\frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-3111d^3))}{7b(a^2+b^2)f(a+b \tan(e+fx))^{7/2}}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int \frac{b^3(bc-ad)((Cc^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2-b^3(bc-ad))}{7b(a^2+b^2)f(a+b \tan(e+fx))^{7/2}}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$\frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-3111d^3))}{7b(a^2+b^2)f(a+b \tan(e+fx))^{7/2}}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$105 \int - \frac{b^3(bc-ad)\left((C^3+3Bdc^2-3Cd^2c-Bd^3-A(c^3-3cd^2))a^4-4b((A-C)d(3c^2-d^2)+B(c^3-3cd^2))a^3+6b^2(Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)a^2\right)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

↓ 25

$$- \frac{2\sqrt{c+d \tan(e+fx)}(15Cd^3a^8+6bBd^3a^7+2b^2d^2(7Bc+4Ad+26Cd)a^6+2b^3d(56c(A-C)d+B(35c^2-12d^2))a^5-b^4(105Bc^3+525Adc^2-525Cdc^2-749Bd^2c-311\dots)}{\dots}$$

$$\frac{2(Ab^2 - a(bB - aC)) (c + d \tan(e + fx))^{5/2}}{7b(a^2 + b^2) f(a + b \tan(e + fx))^{7/2}}$$

input `Int[((c + d*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(a + b*Tan[e + f*x])^(9/2),x]`

output \$Aborted

Defintions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \&\& \text{!Ma}$
 $\text{tchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_) /; \text{FreeQ}[\text{b}, \text{x}]$

rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinear}$
 $\text{Q}[\text{u}, \text{x}]$

rule 4128 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(x_)]^{(m)}*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) +$
 $(\text{f}_.)*(x_)]^{(n)}*((\text{A}_.) + (\text{B}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(x_)] + (\text{C}_.)*\text{tan}[(\text{e}_.)$
 $+ (\text{f}_.)*(x_)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{A}*d^2 + c*(c*C - B*d))*(a + b*\text{Tan}[e +$
 $f*x])^{(m)}*((c + d*\text{Tan}[e + f*x])^{(n+1)}/(d*f*(n+1)*(c^2 + d^2))), \text{x}] - \text{Sim}$
 $\text{p}[1/(d*(n+1)*(c^2 + d^2)) \quad \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}*(c + d*\text{Tan}[e$
 $+ f*x])^{(n+1)}*\text{Simp}[\text{A}*d*(b*d*m - a*c*(n+1)) + (c*C - B*d)*(b*c*m + a*d*$
 $(n+1)) - d*(n+1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*\text{Tan}[e + f*x] - b$
 $*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n+1)))*\text{Tan}[e + f*x]^2, \text{x}],$
 $\text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}\}, \text{x}] \&\& \text{NeQ}[\text{b}*c - \text{a}*d, 0] \&\& \text{NeQ}$
 $[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{GtQ}[\text{m}, 0] \&\& \text{LtQ}[\text{n}, -1]$

rule 4132 $\text{Int}[(\text{a}_.) + (\text{b}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(x_)]^{(m)}*((\text{c}_.) + (\text{d}_.)*\text{tan}[(\text{e}_.) +$
 $(\text{f}_.)*(x_)]^{(n)}*((\text{A}_.) + (\text{B}_.)*\text{tan}[(\text{e}_.) + (\text{f}_.)*(x_)] + (\text{C}_.)*\text{tan}[(\text{e}_.)$
 $+ (\text{f}_.)*(x_)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{A}*b^2 - a*(b*B - a*C))*(a + b*\text{Tan}[e +$
 $f*x])^{(m+1)}*((c + d*\text{Tan}[e + f*x])^{(n+1)}/(f*(m+1)*(b*c - a*d)*(a^2 +$
 $b^2))), \text{x}] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \quad \text{Int}[(a + b*\text{Tan}[e +$
 $f*x])^{(m+1)}*(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[\text{A}*(a*(b*c - a*d)*(m+1) - b^2*d*$
 $(m+n+2)) + (b*B - a*C)*(b*c*(m+1) + a*d*(n+1)) - (m+1)*(b*c - a*d$
 $)*(A*b - a*B - b*C)*\text{Tan}[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m+n+2)*\text{Ta}$
 $n[e + f*x]^2, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{B}, \text{C}, \text{n}\}, \text{x}] \&\& \text{NeQ}$
 $[\text{b}*c - \text{a}*d, 0] \&\& \text{NeQ}[\text{a}^2 + \text{b}^2, 0] \&\& \text{NeQ}[\text{c}^2 + \text{d}^2, 0] \&\& \text{LtQ}[\text{m}, -1] \&\&$
 $!(\text{ILtQ}[\text{n}, -1] \&\& (!\text{IntegerQ}[\text{m}] || (\text{EqQ}[\text{c}, 0] \&\& \text{NeQ}[\text{a}, 0])))$

Maple [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan^2(fx + e))^2}{(a + b \tan(fx + e))^{\frac{9}{2}}} dx$$

input `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x)`

output `int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \text{Timed out}$$

input `integrate((c+d*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(9/2),x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \text{Exception raised: ValueError}$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x, algorithm="maxima")
```

output

```
Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(((2*b*d+2*a*c)^2>0)', see `assume?` for mo
```

Giac [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) * (d \tan(fx + e) + c)^{5/2}}{(b \tan(fx + e) + a)^{9/2}} dx$$

input

```
integrate((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^(5/2)/(b*tan(f*x + e) + a)^(9/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \text{Hanged}$$

input

```
int(((c + d*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(a + b*tan(e + f*x))^(9/2),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{(c + d \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(a + b \tan(e + fx))^{9/2}} dx = \int \frac{(d \tan(fx + e) + c)^{5/2} (A + B \tan(fx + e))}{(\tan(fx + e) + \frac{a}{d})^{9/2}} dx$$

input

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x)
```

output

```
int((c+d*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(9/2),x)
```

$$3.147 \quad \int \frac{(a+b \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1654
Mathematica [A] (verified)	1655
Rubi [A] (verified)	1656
Maple [F(-1)]	1660
Fricas [F(-1)]	1661
Sympy [F]	1661
Maxima [F(-1)]	1661
Giac [F]	1662
Mupad [F(-1)]	1662
Reduce [F]	1663

Optimal result

Integrand size = 49, antiderivative size = 507

$$\int \frac{(a+b \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx =$$

$$\frac{(a-ib)^{5/2}(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c-id}f}$$

$$+ \frac{(a+ib)^{5/2}(iA-B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c+id}f}$$

$$+ \frac{(5a^3Cd^3-15a^2bd^2(cC-2Bd)+5ab^2d(3c^2C-4Bcd+8(A-C)d^2)-b^3(5c^3C-6Bc^2d+8c(A-C)d)}{8\sqrt{bd}^{7/2}f}$$

$$+ \frac{(8b(Ab+aB-bC)d^2+(bc-ad)(5bcC-6bBd-5aCd))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{8d^3f}$$

$$- \frac{(5bcC-6bBd-5aCd)(a+b \tan(e+fx))^{3/2}\sqrt{c+d \tan(e+fx)}}{12d^2f}$$

$$+ \frac{C(a+b \tan(e+fx))^{5/2}\sqrt{c+d \tan(e+fx)}}{3df}$$

output

```

-(a-I*b)^(5/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(1/2)/f+(a+I*b)^(5/2)*(I*A-B-I
*C)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(c+I*d)^(1/2)/f+1/8*(5*a^3*C*d^3-15*a^2*b*d^2*(-2*B*d+C*c)+5*
a*b^2*d*(3*c^2*C-4*B*c*d+8*(A-C)*d^2)-b^3*(5*c^3*C-6*B*c^2*d+8*c*(A-C)*d^2
+16*B*d^3))*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e)
)^(1/2))/b^(1/2)/d^(7/2)/f+1/8*(8*b*(A*b+B*a-C*b)*d^2+(-a*d+b*c)*(-6*B*b*d
-5*C*a*d+5*C*b*c))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/d^3/f-1/1
2*(-6*B*b*d-5*C*a*d+5*C*b*c)*(a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)
/d^2/f+1/3*C*(a+b*tan(f*x+e))^(5/2)*(c+d*tan(f*x+e))^(1/2)/d/f

```

Mathematica [A] (verified)

Time = 7.66 (sec) , antiderivative size = 785, normalized size of antiderivative = 1.55

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df}$$

$$+ \frac{(-5bcC + 6bBd + 5aCd)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{4df} + \frac{3(8b(Ab + aB - bC)d^2 + (bc - ad)(5bcC - 6bBd - 5aCd)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{4df}$$

input

```

Integrate[((a + b*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/Sqrt[c + d*Tan[e + f*x]],x]

```

output

```
(C*(a + b*Tan[e + f*x])^(5/2)*Sqrt[c + d*Tan[e + f*x]]/(3*d*f) + (((-5*b*
c*C + 6*b*B*d + 5*a*C*d)*(a + b*Tan[e + f*x])^(3/2)*Sqrt[c + d*Tan[e + f*x
]])/(4*d*f) + ((3*(8*b*(A*b + a*B - b*C)*d^2 + (b*c - a*d)*(5*b*c*C - 6*b*
B*d - 5*a*C*d))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]/(4*d*f)
+ ((-6*(Sqrt[-b^2]*(3*a^2*b*B - b^3*B - a^3*(A - C) + 3*a*b^2*(A - C)) -
b*(a^3*B - 3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C)))*d^3*ArcTanh[(Sqrt[-
c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqr
t[c + d*Tan[e + f*x]])]/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b
]) - (6*(Sqrt[-b^2]*(3*a^2*b*B - b^3*B - a^3*(A - C) + 3*a*b^2*(A - C)) +
b*(a^3*B - 3*a*b^2*B + 3*a^2*b*(A - C) - b^3*(A - C)))*d^3*ArcTanh[(Sqrt[c
+ (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[
c + d*Tan[e + f*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) +
(3*Sqrt[b]*Sqrt[c - (a*d)/b]*(5*a^3*C*d^3 - 15*a^2*b*d^2*(c*C - 2*B*d) +
5*a*b^2*d*(3*c^2*C - 4*B*c*d + 8*(A - C)*d^2) - b^3*(5*c^3*C - 6*B*c^2*d +
8*c*(A - C)*d^2 + 16*B*d^3))*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(
Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*c + b*d*Tan[e + f*x])/(b*c - a*d)]/(4
*Sqrt[d]*Sqrt[c + d*Tan[e + f*x]])/(b*d*f))/(2*d))/(3*d)
```

Rubi [A] (verified)

Time = 5.93 (sec) , antiderivative size = 524, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{\sqrt{c + d \tan(e + fx)}} dx$$

↓ 4130

$$\int -\frac{(a + b \tan(e + fx))^{3/2} ((5bcC - 5adC - 6bBd) \tan^2(e + fx) - 6(Ab - Cb + aB)d \tan(e + fx) + 5bcC - a(6A - C)d)}{2\sqrt{c + d \tan(e + fx)}} dx + \frac{3d}{3df} \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df}$$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\
 & \frac{\int \frac{(a + b \tan(e + fx))^{3/2} ((5bcC - 5adC - 6bBd) \tan^2(e + fx) - 6(Ab - Cb + aB)d \tan(e + fx) + 5bcC - a(6A - C)d)}{\sqrt{c + d \tan(e + fx)}} dx}{6d} \\
 & \downarrow 3042 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\
 & \frac{\int \frac{(a + b \tan(e + fx))^{3/2} ((5bcC - 5adC - 6bBd) \tan(e + fx)^2 - 6(Ab - Cb + aB)d \tan(e + fx) + 5bcC - a(6A - C)d)}{\sqrt{c + d \tan(e + fx)}} dx}{6d} \\
 & \downarrow 4130 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\
 & \frac{\int \frac{3\sqrt{a + b \tan(e + fx)} (c(5cC - 6Bd)b^2 - 2ad(5cC + Bd)b + a^2(8A - 3C)d^2 + (8b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 6bBd)) \tan^2(e + fx) + 8(Ba^2 + 2b(A - C)a - a^2))}{2\sqrt{c + d \tan(e + fx)}}}{2d}}{6d} \\
 & \downarrow 27 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\
 & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - 3 \int \frac{\sqrt{a + b \tan(e + fx)} (c(5cC - 6Bd)b^2 - 2ad(5cC + Bd)b + a^2(8A - 3C)d^2 + (8b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 6bBd)))}{\sqrt{c + d \tan(e + fx)}}}{6d} \\
 & \downarrow 3042 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\
 & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - 3 \int \frac{\sqrt{a + b \tan(e + fx)} (c(5cC - 6Bd)b^2 - 2ad(5cC + Bd)b + a^2(8A - 3C)d^2 + (8b(Ab - Cb + aB)d^2 + (bc - ad)(5bcC - 5adC - 6bBd)))}{\sqrt{c + d \tan(e + fx)}}}{6d} \\
 & \downarrow 4130 \\
 & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\
 & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - 3 \left(\int \frac{-c(5C^2 - 6Bdc + 8(A - C)d^2)b^3 + ad(15C^2 - 20Bdc - 8(A - C)d^2)b^2 - 3a^2d^2(5cC - 6Bd)}{\sqrt{c + d \tan(e + fx)}}}{6d} \right)
 \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \left(\frac{\int \frac{-c(5Cc^2 - 6Bdc + 8(A-C)d^2)b^3 + ad(15Cc^2 - 20Bdc - 8(A-C)d^2)b^2 - 3a^2d^2(5cC}{\dots}}{3} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \left(\frac{\int \frac{-c(5Cc^2 - 6Bdc + 8(A-C)d^2)b^3 + ad(15Cc^2 - 20Bdc - 8(A-C)d^2)b^2 - 3a^2d^2(5cC}{\dots}}{3} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4138 \\ & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \left(\frac{\int \frac{-c(5Cc^2 - 6Bdc + 8(A-C)d^2)b^3 + ad(15Cc^2 - 20Bdc - 8(A-C)d^2)b^2 - 3a^2d^2(5cC}{\dots}}{3} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2348 \\ & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \left(\frac{\int \frac{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (8bd^2(aB + Ab - bC) + (bc - ad)(-5aCd - 6bBd + 5bcC)}{df}}{3} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2009 \\ & \frac{C(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}}{3df} - \\ & \frac{(-5aCd - 6bBd + 5bcC)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \left(\frac{\int \frac{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (8bd^2(aB + Ab - bC) + (bc - ad)(-5aCd - 6bBd + 5bcC)}{df}}{3} \right) \end{aligned}$$

input $\text{Int}[(a + b \cdot \tan[e + f \cdot x])^{5/2} \cdot (A + B \cdot \tan[e + f \cdot x] + C \cdot \tan[e + f \cdot x]^2) / \sqrt{c + d \cdot \tan[e + f \cdot x]}, x]$

output $(C \cdot (a + b \cdot \tan[e + f \cdot x])^{5/2} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]}) / (3 \cdot d \cdot f) - ((5 \cdot b \cdot c \cdot C - 6 \cdot b \cdot B \cdot d - 5 \cdot a \cdot C \cdot d) \cdot (a + b \cdot \tan[e + f \cdot x])^{3/2} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]}) / (2 \cdot d \cdot f) - (3 \cdot (((-16 \cdot (a - I \cdot b)^{5/2} \cdot (I \cdot A + B - I \cdot C) \cdot d^3 \cdot \text{ArcTanh}[\sqrt{c - I \cdot d} \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]}]) / (\sqrt{a - I \cdot b} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]})) / \sqrt{c - I \cdot d} - (16 \cdot (a + I \cdot b)^{5/2} \cdot (B - I \cdot (A - C)) \cdot d^3 \cdot \text{ArcTanh}[\sqrt{c + I \cdot d} \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]}]) / (\sqrt{a + I \cdot b} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]})) / \sqrt{c + I \cdot d} + (2 \cdot (5 \cdot a^3 \cdot C \cdot d^3 - 15 \cdot a^2 \cdot b \cdot d^2 \cdot (c \cdot C - 2 \cdot B \cdot d) + 5 \cdot a \cdot b^2 \cdot d \cdot (3 \cdot c^2 \cdot C - 4 \cdot B \cdot c \cdot d + 8 \cdot (A - C) \cdot d^2) - b^3 \cdot (5 \cdot c^3 \cdot C - 6 \cdot B \cdot c^2 \cdot d + 8 \cdot c \cdot (A - C) \cdot d^2 + 16 \cdot B \cdot d^3)) \cdot \text{ArcTanh}[\sqrt{d} \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]}] / (\sqrt{b} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]})) / (\sqrt{b} \cdot \sqrt{d})) / (2 \cdot d \cdot f) + ((8 \cdot b \cdot (A \cdot b + a \cdot B - b \cdot C) \cdot d^2 + (b \cdot c - a \cdot d) \cdot (5 \cdot b \cdot c \cdot C - 6 \cdot b \cdot B \cdot d - 5 \cdot a \cdot C \cdot d)) \cdot \sqrt{a + b \cdot \tan[e + f \cdot x]} \cdot \sqrt{c + d \cdot \tan[e + f \cdot x]}) / (d \cdot f)) / (4 \cdot d) / (6 \cdot d)$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)(Gx_)] /; \text{FreeQ}[b, x]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2348 $\text{Int}[(Px_)((c_)+(d_)(x_))^{(m_)}((e_)+(f_)(x_))^{(n_)}((a_)+(b_)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[Px \cdot (c + d \cdot x)^m \cdot (e + f \cdot x)^n \cdot (a + b \cdot x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2 \cdot p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{\sqrt{c + d \tan(fx + e)}} dx$$

input

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(1/2),x)
```

output

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(1/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((a+b*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((a + b*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e)}{\sqrt{d \tan(fx + e) + c}}$$

input `integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(5/2)/sqrt(d*tan(f*x + e) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(a + b \tan(e + fx))^{5/2} (C \tan(e + fx) + B)}{\sqrt{c + d \tan(e + fx)}}$$

input `int(((a + b*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2),x)`

output `int(((a + b*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2), x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)}}{d \tan(fx + e) + c} \right. \\
+ 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)^3}{d \tan(fx + e) + c} dx \right) abc \\
+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)^3}{d \tan(fx + e) + c} dx \right) b^3 \\
+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) a^2 c \\
+ 3 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) a b^2 \\
+ 3 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)}{d \tan(fx + e) + c} dx \right) a^2 b \\
+ \left. \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a}{d \tan(fx + e) + c} dx \right) a^3 \right.$$

input

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(1/2),x)
```

output

```
int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(t
an(e + f*x)*d + c),x)*b**2*c + 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e
+ f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)*d + c),x)*a*b*c + int((sqrt(t
an(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)
*d + c),x)*b**3 + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*t
an(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a**2*c + 3*int((sqrt(tan(e + f*x)*
d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*
a*b**2 + 3*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f
*x))/(tan(e + f*x)*d + c),x)*a**2*b + int((sqrt(tan(e + f*x)*d + c)*sqrt(t
an(e + f*x)*b + a))/(tan(e + f*x)*d + c),x)*a**3
```

$$3.148 \quad \int \frac{(a+b \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1664
Mathematica [A] (verified)	1665
Rubi [A] (verified)	1666
Maple [F(-1)]	1669
Fricas [F(-1)]	1670
Sympy [F]	1670
Maxima [F]	1670
Giac [F]	1671
Mupad [F(-1)]	1671
Reduce [F]	1672

Optimal result

Integrand size = 49, antiderivative size = 381

$$\int \frac{(a+b \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx =$$

$$\frac{(a-ib)^{3/2}(iA+B-iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c-id}f}$$

$$- \frac{(a+ib)^{3/2}(B-i(A-C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c+id}f}$$

$$+ \frac{(3a^2Cd^2-6abd(cC-2Bd)+b^2(3c^2C-4Bcd+8(A-C)d^2)) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{4\sqrt{bd}^{5/2}f}$$

$$- \frac{(3bcC-4bBd-3aCd)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{4d^2f}$$

$$+ \frac{C(a+b \tan(e+fx))^{3/2}\sqrt{c+d \tan(e+fx)}}{2df}$$

output

$$\begin{aligned}
& -(a-I*b)^{(3/2)}*(I*A+B-I*C)*\operatorname{arctanh}((c-I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)})/(a \\
& -I*b)^{(1/2)}/(c+d*\tan(f*x+e))^{(1/2)})/(c-I*d)^{(1/2)}/f-(a+I*b)^{(3/2)}*(B-I*(A- \\
& C))*\operatorname{arctanh}((c+I*d)^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)})/(a+I*b)^{(1/2)}/(c+d*\tan(f* \\
& x+e))^{(1/2)})/(c+I*d)^{(1/2)}/f+1/4*(3*a^2*C*d^2-6*a*b*d*(-2*B*d+C*c)+b^2*(3* \\
& c^2*C-4*B*c*d+8*(A-C)*d^2))*\operatorname{arctanh}(d^{(1/2)}*(a+b*\tan(f*x+e))^{(1/2)})/b^{(1/2)} \\
& /((c+d*\tan(f*x+e))^{(1/2)})/b^{(1/2)}/d^{(5/2)}/f-1/4*(-4*B*b*d-3*C*a*d+3*C*b*c)* \\
& (a+b*\tan(f*x+e))^{(1/2)}*(c+d*\tan(f*x+e))^{(1/2)}/d^2/f+1/2*C*(a+b*\tan(f*x+e)) \\
& ^{(3/2)}*(c+d*\tan(f*x+e))^{(1/2)}/d/f
\end{aligned}$$

Mathematica [A] (verified)

Time = 4.64 (sec) , antiderivative size = 582, normalized size of antiderivative = 1.53

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \frac{4(b(a^2 B - b^2 B + 2ab(A - C)) - \sqrt{-b^2}(2abB + b^2(A - C))) \sqrt{c + d \tan(e + fx)}}{b \sqrt{-b^2}}$$

input

```
Integrate[((a + b*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]
```

output

$$\begin{aligned}
& ((4*(b*(a^2*B - b^2*B + 2*a*b*(A - C)) - \operatorname{Sqrt}[-b^2]*(2*a*b*B + b^2*(A - C) \\
& + a^2*(-A + C)))*d^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[-c + (\operatorname{Sqrt}[-b^2]*d)/b]*\operatorname{Sqrt}[a + b*\tan[e \\
& + f*x]])/(\operatorname{Sqrt}[-a + \operatorname{Sqrt}[-b^2]]*\operatorname{Sqrt}[c + d*\tan[e + f*x]])]/(b*\operatorname{Sqrt}[-a + \\
& \operatorname{Sqrt}[-b^2]]*\operatorname{Sqrt}[-c + (\operatorname{Sqrt}[-b^2]*d)/b]) - (4*(b*(a^2*B - b^2*B + 2*a*b*(A \\
& - C)) + \operatorname{Sqrt}[-b^2]*(2*a*b*B + b^2*(A - C) + a^2*(-A + C)))*d^2*\operatorname{ArcTanh}[(\\
& \operatorname{Sqrt}[c + (\operatorname{Sqrt}[-b^2]*d)/b]*\operatorname{Sqrt}[a + b*\tan[e + f*x]])/(\operatorname{Sqrt}[a + \operatorname{Sqrt}[-b^2]] \\
& *\operatorname{Sqrt}[c + d*\tan[e + f*x]])]/(b*\operatorname{Sqrt}[a + \operatorname{Sqrt}[-b^2]]*\operatorname{Sqrt}[c + (\operatorname{Sqrt}[-b^2]* \\
& d)/b]) + (-3*b*c*C + 4*b*B*d + 3*a*C*d)*\operatorname{Sqrt}[a + b*\tan[e + f*x]]*\operatorname{Sqrt}[c + \\
& d*\tan[e + f*x]] + 2*C*d*(a + b*\tan[e + f*x])^{(3/2)}*\operatorname{Sqrt}[c + d*\tan[e + f*x]] \\
&] + (\operatorname{Sqrt}[c - (a*d)/b]*(3*a^2*C*d^2 + 6*a*b*d*(-(c*C) + 2*B*d) + b^2*(3*c^2 \\
& *C - 4*B*c*d + 8*(A - C)*d^2))*\operatorname{ArcSinh}[(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a + b*\tan[e + f*x]]) \\
& /(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[c - (a*d)/b])]*\operatorname{Sqrt}[(b*(c + d*\tan[e + f*x]))/(b*c - a*d)]/ \\
& (\operatorname{Sqrt}[b]*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + d*\tan[e + f*x]])/(4*d^2*f)
\end{aligned}$$

Rubi [A] (verified)

Time = 3.56 (sec) , antiderivative size = 392, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.204$, Rules used = {3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{\sqrt{c + d \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4130} \\
 & \frac{\int -\frac{\sqrt{a+b \tan(e+fx)}((3bcC-3adC-4bBd) \tan^2(e+fx)-4(Ab-Cb+aB)d \tan(e+fx)+3bcC-a(4A-C)d)}{2\sqrt{c+d \tan(e+fx)}} dx}{\frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df}} + \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{a+b \tan(e+fx)}((3bcC-3adC-4bBd) \tan^2(e+fx)-4(Ab-Cb+aB)d \tan(e+fx)+3bcC-a(4A-C)d)}{\sqrt{c+d \tan(e+fx)}} dx}{\frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df}} - \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{a+b \tan(e+fx)}((3bcC-3adC-4bBd) \tan(e+fx)^2-4(Ab-Cb+aB)d \tan(e+fx)+3bcC-a(4A-C)d)}{\sqrt{c+d \tan(e+fx)}} dx}{\frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df}} - \\
 & \quad \downarrow \text{4130} \\
 & \frac{\int -\frac{c(3cC-4Bd)b^2-2ad(3cC+2Bd)b+a^2(8A-5C)d^2+(8b(Ab-Cb+aB)d^2+(bc-ad)(3bcC-3adC-4bBd)) \tan^2(e+fx)+8(Ba^2+2b(A-C)a-b^2B)d^2 \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{4d}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \\ & \frac{(-3aCd - 4bBd + 3bcC) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{df} - \frac{\int \frac{c(3cC - 4Bd)b^2 - 2ad(3cC + 2Bd)b + a^2(8A - 5C)d^2 + (8b(Ab - Cb + aB)d^2 + (bc - ad)(3bcC - 2ad^2)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{2d}}{4d} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \\ & \frac{(-3aCd - 4bBd + 3bcC) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{df} - \frac{\int \frac{c(3cC - 4Bd)b^2 - 2ad(3cC + 2Bd)b + a^2(8A - 5C)d^2 + (8b(Ab - Cb + aB)d^2 + (bc - ad)(3bcC - 2ad^2)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{2d}}{4d} \end{aligned}$$

$$\begin{aligned} & \downarrow 4138 \\ & \frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \\ & \frac{(-3aCd - 4bBd + 3bcC) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{df} - \frac{\int \frac{c(3cC - 4Bd)b^2 - 2ad(3cC + 2Bd)b + a^2(8A - 5C)d^2 + (8b(Ab - Cb + aB)d^2 + (bc - ad)(3bcC - 2ad^2)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{2df}}{4d} \end{aligned}$$

$$\begin{aligned} & \downarrow 2348 \\ & \frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \\ & \frac{(-3aCd - 4bBd + 3bcC) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{df} - \frac{\int \left(\frac{8Ad^2b^2 - 8Cd^2b^2 + 3c^2Cb^2 - 4Bcdb^2 + 12aBd^2b - 6acCdb + 3a^2Cd^2}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} + \frac{-16aAbd^2 - 8a^2Bd^2}{\sqrt{b} \sqrt{d}} \right) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{4d} \end{aligned}$$

$$\begin{aligned} & \downarrow 2009 \\ & \frac{C(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}{2df} - \\ & \frac{(-3aCd - 4bBd + 3bcC) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{df} - \frac{2(3a^2Cd^2 - 6abd(cC - 2Bd) + b^2(8d^2(A - C) - 4Bcd + 3c^2C)) \operatorname{arctanh}\left(\frac{\sqrt{d} \sqrt{a + b \tan(e + fx)}}{\sqrt{b} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{b} \sqrt{d}} \end{aligned}$$

4d

input

```
Int[((a + b*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]
```


output

```
(C*(a + b*Tan[e + f*x])^(3/2)*Sqrt[c + d*Tan[e + f*x]]/(2*d*f) - (-1/2*((-8*(a - I*b)^(3/2)*(I*A + B - I*C)*d^2*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c - I*d] - (8*(a + I*b)^(3/2)*(B - I*(A - C))*d^2*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c + I*d] + (2*(3*a^2*C*d^2 - 6*a*b*d*(c*C - 2*B*d) + b^2*(3*c^2*C - 4*B*c*d + 8*(A - C)*d^2))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/((Sqrt[b]*Sqrt[d]))/(d*f) + ((3*b*c*C - 4*b*B*d - 3*a*C*d)*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(d*f))/(4*d)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2348

```
Int[(P_x)*((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{\sqrt{c + d \tan(fx + e)}} dx$$

input

```
int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(1/2),x)
```

output

```
int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(1/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((a+b*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((a + b*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) (a + b \tan(fx + e))^{3/2}}{\sqrt{d \tan(fx + e) + c}}$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)/sqrt(d*tan(f*x + e) + c), x)
```

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A) (a + b \tan(fx + e))^{3/2}}{\sqrt{d \tan(fx + e) + c}} dx$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)/sqrt(d*tan(f*x + e) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \int \frac{(a + b \tan(e + fx))^{3/2} (C \tan(e + fx) + B + A \tan(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

input

```
int(((a + b*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2),x)
```

output

```
int(((a + b*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2), x)
```

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)}}{d \tan(fx + e) + c} dx \right) ac$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) ac$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) b^2$$

$$+ 2 \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a \tan(fx + e)}{d \tan(fx + e) + c} dx \right) ab$$

$$+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e)} b + a}{d \tan(fx + e) + c} dx \right) a^2$$

input

```
int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(1/2),x)
```

output

```
int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(t
an(e + f*x)*d + c),x)*b*c + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)
)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*a*c + int((sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*d + c)
,x)*b**2 + 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e
+ f*x))/(tan(e + f*x)*d + c),x)*a*b + int((sqrt(tan(e + f*x)*d + c)*sqrt(t
an(e + f*x)*b + a))/(tan(e + f*x)*d + c),x)*a**2
```

3.149
$$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1673
Mathematica [A] (verified)	1674
Rubi [A] (verified)	1674
Maple [F(-1)]	1677
Fricas [B] (verification not implemented)	1678
Sympy [F]	1678
Maxima [F]	1679
Giac [F]	1679
Mupad [F(-1)]	1680
Reduce [F]	1680

Optimal result

Integrand size = 49, antiderivative size = 290

$$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx$$

$$= -\frac{\sqrt{a-ib}(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c-id}f}$$

$$+ \frac{\sqrt{a+ib}(iA-B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c+id}f}$$

$$- \frac{(bcC-2bBd-aCd)\operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{bd}^{3/2}f}$$

$$+ \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df}$$

output

```

-(a-I*b)^(1/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(1/2)/f+(a+I*b)^(1/2)*(I*A-B-I
*C)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(c+I*d)^(1/2)/f-(-2*B*b*d-C*a*d+C*b*c)*arctanh(d^(1/2)*(a+b*t
an(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/b^(1/2)/d^(3/2)/f+C*(a+b*
tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/d/f
    
```

Mathematica [A] (verified)

Time = 4.41 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.55

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{C \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{df}$$

$$\frac{(-b(Ab + aB - bC) + \sqrt{-b^2}(bB + a(-A + C))) \operatorname{darctanh}\left(\frac{\sqrt{-c + \frac{\sqrt{-b^2}d}{b}} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{-a + \sqrt{-b^2}} \sqrt{-c + \frac{\sqrt{-b^2}d}{b}}} + \frac{(b(Ab + aB - bC) + \sqrt{-b^2}(bB + a(-A + C))) d}{\sqrt{a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}} + \frac{bdf}{bdf}$$

input

```
Integrate[(Sqrt[a + b*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]
```

output

```
(C*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(d*f) - (((-b*(A*b + a*B - b*C)) + Sqrt[-b^2]*(b*B + a*(-A + C)))*d*ArcTanh[(Sqrt[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) + ((b*(A*b + a*B - b*C) + Sqrt[-b^2]*(b*B + a*(-A + C)))*d*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) + (Sqrt[b]*Sqrt[c - (a*d)/b]*(b*c*C - 2*b*B*d - a*C*d)*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)])/(Sqrt[d]*Sqrt[c + d*Tan[e + f*x]])/(b*d*f)
```

Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{\sqrt{c+d \tan(e+fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan(e+fx)^2)}{\sqrt{c+d \tan(e+fx)}} dx \\
 & \quad \downarrow \text{4130} \\
 & \int \frac{(bcC-adC-2bBd) \tan^2(e+fx)-2(Ab-Cb+aB)d \tan(e+fx)+bcC-2aAd+aCd}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx + \\
 & \quad \frac{d}{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \frac{df}{df} \\
 & \quad \downarrow \text{27} \\
 & \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df} - \\
 & \int \frac{(bcC-adC-2bBd) \tan^2(e+fx)-2(Ab-Cb+aB)d \tan(e+fx)+bcC-a(2A-C)d}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx \\
 & \quad \frac{2d}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df} - \\
 & \int \frac{(bcC-adC-2bBd) \tan(e+fx)^2-2(Ab-Cb+aB)d \tan(e+fx)+bcC-a(2A-C)d}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx \\
 & \quad \frac{2d}{2d} \\
 & \quad \downarrow \text{4138} \\
 & \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df} - \\
 & \int \frac{(bcC-adC-2bBd) \tan^2(e+fx)-2(Ab-Cb+aB)d \tan(e+fx)+bcC-a(2A-C)d}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(\tan^2(e+fx)+1)}} dx \\
 & \quad \frac{2df}{2df} \\
 & \quad \downarrow \text{2348} \\
 & \frac{C\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df} - \\
 & \int \left(\frac{bcC-adC-2bBd}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} + \frac{2Abd+2aBd-2bCd+i(-2aAd+2bBd+2aCd)}{2(i-\tan(e+fx))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} + \frac{-2Abd-2aBd+2bCd+i(-2aAd+2bBd+2aCd)}{2(\tan(e+fx)+i)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \right) dx \\
 & \quad \frac{2df}{2df} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{C\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{df} - \frac{2d\sqrt{a-ib}(B+i(A-C))\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b\tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{c-id}} - \frac{2d\sqrt{a+ib}(iA-B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b\tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{c+id}} + \frac{2(-aCd-2bBd+bc)}{2df}$$

input

```
Int[(Sqrt[a + b*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/Sqrt[c + d*Tan[e + f*x]],x]
```

output

```
-1/2*((2*Sqrt[a - I*b]*(B + I*(A - C))*d*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c - I*d] - (2*Sqrt[a + I*b]*(I*A - B - I*C))*d*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c + I*d] + (2*(b*c*C - 2*b*B*d - a*C*d)*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/((Sqrt[b]*Sqrt[d]))/(d*f) + (C*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(d*f)
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2348

```
Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan (fx + e)} (A + B \tan (fx + e) + C \tan (fx + e)^2)}{\sqrt{c + d \tan (fx + e)}} dx$$

input

```
int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```
int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38950 vs. $2(226) = 452$.

Time = 127.24 (sec) , antiderivative size = 77916, normalized size of antiderivative = 268.68

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Too large to include`

Sympy [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((a+b*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral(sqrt(a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/sqrt(c + d*tan(e + f*x)), x)`

Maxima [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a}}{\sqrt{d \tan(fx + e) + c}} dx$$

input

```
integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)/sqrt(d*tan(f*x + e) + c), x)
```

Giac [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e) + A) \sqrt{b \tan(fx + e) + a}}{\sqrt{d \tan(fx + e) + c}} dx$$

input

```
integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)/sqrt(d*tan(f*x + e) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx = \text{Hanged}$$

input `int(((a + b*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(1/2),x)`

output `\text{Hanged}`

Reduce [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{\sqrt{c + d \tan(e + fx)}} dx \\ &= \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) c \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a} \tan(fx + e)}{d \tan(fx + e) + c} dx \right) b \\ &+ \left(\int \frac{\sqrt{d \tan(fx + e) + c} \sqrt{\tan(fx + e) b + a}}{d \tan(fx + e) + c} dx \right) a \end{aligned}$$

input `int(((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(1/2),x)`

output `int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*c + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)*d + c),x)*b + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)*d + c),x)*a`

3.150 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx$

Optimal result	1681
Mathematica [A] (verified)	1682
Rubi [A] (verified)	1682
Maple [F(-1)]	1684
Fricas [B] (verification not implemented)	1685
Sympy [F]	1685
Maxima [F]	1686
Giac [F]	1686
Mupad [F(-1)]	1687
Reduce [F]	1687

Optimal result

Integrand size = 49, antiderivative size = 237

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx$$

$$= -\frac{(B + i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c - id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a - ib} \sqrt{c - id} f}$$

$$- \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a + ib} \sqrt{c + id} f} + \frac{2C \operatorname{arctanh}\left(\frac{\sqrt{d} \sqrt{a + b \tan(e + fx)}}{\sqrt{b} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{b} \sqrt{d} f}$$

output

```

-(B+I*(A-C))*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c
+d*tan(f*x+e))^(1/2))/(a-I*b)^(1/2)/(c-I*d)^(1/2)/f-(B-I*(A-C))*arctanh((c
+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(
a+I*b)^(1/2)/(c+I*d)^(1/2)/f+2*C*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(
1/2)/(c+d*tan(f*x+e))^(1/2))/b^(1/2)/d^(1/2)/f
    
```

Mathematica [A] (verified)

Time = 1.45 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.53

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx$$

$$= \frac{(bB + \sqrt{-b^2}(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{-c + \frac{\sqrt{-b^2}d}{b}} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{-a + \sqrt{-b^2}} \sqrt{-c + \frac{\sqrt{-b^2}d}{b}}} - \frac{(bB + \sqrt{-b^2}(-A + C)) \operatorname{arctanh}\left(\frac{\sqrt{c + \frac{\sqrt{-b^2}d}{b}} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + \sqrt{-b^2}} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a + \sqrt{-b^2}} \sqrt{c + \frac{\sqrt{-b^2}d}{b}}} + \dots$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]),x]
```

output

```
((b*B + Sqrt[-b^2]*(A - C))*ArcTanh[(Sqrt[-c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[-a + Sqrt[-b^2]]*Sqrt[-c + (Sqrt[-b^2]*d)/b]) - (b*B + Sqrt[-b^2]*(-A + C))*ArcTanh[(Sqrt[c + (Sqrt[-b^2]*d)/b]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + d*Tan[e + f*x]])]/(Sqrt[a + Sqrt[-b^2]]*Sqrt[c + (Sqrt[-b^2]*d)/b]) + (2*Sqrt[b]*C*Sqrt[c - (a*d)/b]*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c - (a*d)/b])]*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)]/(Sqrt[d]*Sqrt[c + d*Tan[e + f*x]])]/(b*f)
```

Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.082$, Rules used = {3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx \\
& \quad \downarrow \text{4138} \\
& \int \frac{C \tan^2(e + fx) + B \tan(e + fx) + A}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)} (\tan^2(e + fx) + 1)} d \tan(e + fx) \\
& \quad \downarrow \text{2348} \\
& \int \left(\frac{i(A - C) - B}{2(i - \tan(e + fx)) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} + \frac{C}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} + \frac{B + i(A - C)}{2(\tan(e + fx) + i) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} \right) dx \\
& \quad \downarrow \text{2009} \\
& \frac{(B + i(A - C)) \operatorname{arctanh} \left(\frac{\sqrt{c - id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib} \sqrt{c + d \tan(e + fx)}} \right) - (B - i(A - C)) \operatorname{arctanh} \left(\frac{\sqrt{c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib} \sqrt{c + d \tan(e + fx)}} \right) + 2C \operatorname{arctanh} \left(\frac{\sqrt{d} \sqrt{a + b \tan(e + fx)}}{\sqrt{b} \sqrt{c + d \tan(e + fx)}} \right)}{f \sqrt{a - ib} \sqrt{c - id} \sqrt{a + ib} \sqrt{c + id} \sqrt{b} \sqrt{d}}
\end{aligned}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]),x]
```

output

```
(-(((B + I*(A - C))*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a - I*b]*Sqrt[c - I*d])) - ((B - I*(A - C))*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]) + (2*C*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[b]*Sqrt[d]))/f
```


Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2348 `Int[(Px_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4138 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan(fx + e) + C \tan(fx + e)^2}{\sqrt{a + b \tan(fx + e)} \sqrt{c + d \tan(fx + e)}} dx$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2),x)`

output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48154 vs. $2(180) = 360$.

Time = 128.69 (sec) , antiderivative size = 96324, normalized size of antiderivative = 406.43

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output Too large to include

Sympy [F]

$$\begin{aligned} & \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx \\ &= \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(1/2)/(c+d*tan(f*x+e))**(1/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(sqrt(a + b*tan(e + f*x))*sqrt(c + d*tan(e + f*x))), x)`

Maxima [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{\sqrt{b \tan(fx + e) + a} \sqrt{d \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/(sqrt(b*tan(f*x + e) + a)*sqrt(d*tan(f*x + e) + c)), x)`

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx$$

$$= \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{\sqrt{b \tan(fx + e) + a} \sqrt{d \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/(sqrt(b*tan(f*x + e) + a)*sqrt(d*tan(f*x + e) + c)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(1/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```

(2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a - 2*int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*
a*b*d**2 + tan(e + f*x)**2*b**2*c*d + tan(e + f*x)*a**2*d**2 + 2*tan(e + f
*x)*a*b*c*d + tan(e + f*x)*b**2*c**2 + a**2*c*d + a*b*c**2),x)*a**2*b*d**2
*f - 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)
**3)/(tan(e + f*x)**2*a*b*d**2 + tan(e + f*x)**2*b**2*c*d + tan(e + f*x)*a
**2*d**2 + 2*tan(e + f*x)*a*b*c*d + tan(e + f*x)*b**2*c**2 + a**2*c*d + a*
b*c**2),x)*a*b**2*c*d*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*
b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*a*b*d**2 + tan(e + f*x)**2*b**2*c
*d + tan(e + f*x)*a**2*d**2 + 2*tan(e + f*x)*a*b*c*d + tan(e + f*x)*b**2*c
**2 + a**2*c*d + a*b*c**2),x)*a**3*d**2*f - 2*int((sqrt(tan(e + f*x)*d + c
)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*a*b*d**2 + ta
n(e + f*x)**2*b**2*c*d + tan(e + f*x)*a**2*d**2 + 2*tan(e + f*x)*a*b*c*d +
tan(e + f*x)*b**2*c**2 + a**2*c*d + a*b*c**2),x)*a**2*b*c*d*f - int((sqrt
(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*
x)**2*a*b*d**2 + tan(e + f*x)**2*b**2*c*d + tan(e + f*x)*a**2*d**2 + 2*tan
(e + f*x)*a*b*c*d + tan(e + f*x)*b**2*c**2 + a**2*c*d + a*b*c**2),x)*a*b**
2*c**2*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)**2)/(tan(e + f*x)**2*b*d + tan(e + f*x)*a*d + tan(e + f*x)*b*c + a*c)
,x)*a*c*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*ta...

```

3.151
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1689
Mathematica [A] (verified)	1690
Rubi [A] (verified)	1690
Maple [F(-1)]	1694
Fricas [F(-1)]	1694
Sympy [F]	1694
Maxima [F]	1695
Giac [F]	1695
Mupad [F(-1)]	1696
Reduce [F]	1696

Optimal result

Integrand size = 49, antiderivative size = 251

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a - ib)^{3/2} \sqrt{c - idf}}$$

$$\frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a + ib)^{3/2} \sqrt{c + idf}}$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{(a^2 + b^2) (bc - ad) f \sqrt{a + b \tan(e + fx)}}$$

output

```
-(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(3/2)/(c-I*d)^(1/2)/f-(B-I*(A-C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a+I*b)^(3/2)/(c+I*d)^(1/2)/f-2*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(1/2)/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 1.78 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.05

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx = \frac{(a+ib)(iA+B-iC) \operatorname{arctanh}\left(\frac{\sqrt{-c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{-a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{-a+ib}\sqrt{-c+id}} + \frac{(ia+b)(A+iB)}{(a+ib)(A+iB)}$$

```
input Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(3/2)*Sqrt[c + d*Tan[e + f*x]]),x]
```

```
output (((a + I*b)*(I*A + B - I*C)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + I*b]*Sqrt[-c + I*d]) + ((I*a + b)*(A + I*B - C)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]) + (2*(A*b^2 + a*(-(b*B) + a*C))*Sqrt[c + d*Tan[e + f*x]])/((- (b*c) + a*d)*Sqrt[a + b*Tan[e + f*x]])/((a^2 + b^2)*f)
```

Rubi [A] (verified)

Time = 2.32 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.21, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.184$, Rules used = {3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx$$

↓ 4132

$$\begin{aligned}
 & \frac{2 \int -\frac{(bB+a(A-C))(bc-ad)-(Ab-Cb-aB)(bc-ad)\tan(e+fx)}{2\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx}{(a^2+b^2)(bc-ad)} \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{(bB+a(A-C))(bc-ad)-(Ab-Cb-aB)(bc-ad)\tan(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx}{(a^2+b^2)(bc-ad)} \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{(bB+a(A-C))(bc-ad)-(Ab-Cb-aB)(bc-ad)\tan(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx}{(a^2+b^2)(bc-ad)} \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} \\
 & \quad \downarrow 4099 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} + \\
 & \frac{\frac{1}{2}(a-ib)(A+iB-C)(bc-ad) \int \frac{1-i\tan(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx + \frac{1}{2}(a+ib)(A-iB-C)(bc-ad) \int \frac{i}{\sqrt{a+b\tan(e+fx)}} dx}{(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 3042 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} + \\
 & \frac{\frac{1}{2}(a-ib)(A+iB-C)(bc-ad) \int \frac{1-i\tan(e+fx)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx + \frac{1}{2}(a+ib)(A-iB-C)(bc-ad) \int \frac{i}{\sqrt{a+b\tan(e+fx)}} dx}{(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 4098 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} + \\
 & \frac{(a+ib)(A-iB-C)(bc-ad) \int \frac{1}{(1-i\tan(e+fx))\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} d\tan(e+fx)}{2f} + \frac{(a-ib)(A+iB-C)(bc-ad) \int \frac{1}{(i\tan(e+fx)+1)\sqrt{a+b\tan(e+fx)}} d\tan(e+fx)}{2f} \\
 & \quad \downarrow 104 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} + \\
 & \frac{(a+ib)(A-iB-C)(bc-ad) \int \frac{1}{(1-i\tan(e+fx))\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} d\tan(e+fx)}{2f} + \frac{(a-ib)(A+iB-C)(bc-ad) \int \frac{1}{(i\tan(e+fx)+1)\sqrt{a+b\tan(e+fx)}} d\tan(e+fx)}{2f} \\
 & \quad \downarrow 104
 \end{aligned}$$

rule 221 $\text{Int}[(a_.) + (b_.) \cdot (x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 4098 $\text{Int}[(a_.) + (b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(m_.)} \cdot ((A_.) + (B_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[A^2/f \text{ Subst}[\text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n / (A - B \cdot x)], x], x, \text{Tan}[e + f \cdot x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A^2 + B^2, 0]$

rule 4099 $\text{Int}[(a_.) + (b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(m_.)} \cdot ((A_.) + (B_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(A + I \cdot B)/2 \text{ Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^m \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot (1 - I \cdot \text{Tan}[e + f \cdot x]), x], x] + \text{Simp}[(A - I \cdot B)/2 \text{ Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^m \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot (1 + I \cdot \text{Tan}[e + f \cdot x]), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A^2 + B^2, 0]$

rule 4132 $\text{Int}[(a_.) + (b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(m_.)} \cdot ((c_.) + (d_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(n_.)} \cdot ((A_.) + (B_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)] + (C_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)]^2), x_Symbol] \rightarrow \text{Simp}[(A \cdot b^2 - a \cdot (b \cdot B - a \cdot C)) \cdot (a + b \cdot \text{Tan}[e + f \cdot x])^{(m+1)} \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^{(n+1)} / (f \cdot (m+1) \cdot (b \cdot c - a \cdot d) \cdot (a^2 + b^2)), x] + \text{Simp}[1 / ((m+1) \cdot (b \cdot c - a \cdot d) \cdot (a^2 + b^2)) \text{ Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^{(m+1)} \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot \text{Simp}[A \cdot (a \cdot (b \cdot c - a \cdot d) \cdot (m+1) - b^2 \cdot d \cdot (m+n+2)) + (b \cdot B - a \cdot C) \cdot (b \cdot c \cdot (m+1) + a \cdot d \cdot (n+1)) - (m+1) \cdot (b \cdot c - a \cdot d) \cdot (A \cdot b - a \cdot B - b \cdot C) \cdot \text{Tan}[e + f \cdot x] - d \cdot (A \cdot b^2 - a \cdot (b \cdot B - a \cdot C)) \cdot (m+n+2) \cdot \text{Tan}[e + f \cdot x]^2, x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, C, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ !(\text{ILtQ}[n, -1] \ \&\& \ (!\text{IntegerQ}[m] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{NeQ}[a, 0])))$

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan(fx + e) + C \tan(fx + e)^2}{(a + b \tan(fx + e))^{\frac{3}{2}} \sqrt{c + d \tan(fx + e)}} dx$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2),x)`

output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{3}{2}} \sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{3}{2}} \sqrt{c + d \tan(e + fx)}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{3}{2}} \sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(3/2)/(c+d*tan(f*x+e))**(1/2),x)`

output

```
Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**(3/2)*sqrt(c + d*tan(e + f*x))), x)
```

Maxima [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx = \int \frac{C \tan(fx + e)^2 + B \tan(fx + e) + A}{(b \tan(fx + e) + a)^{3/2} \sqrt{d \tan(fx + e) + c}} dx$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/((b*tan(f*x + e) + a)^(3/2)*sqrt(d*tan(f*x + e) + c)), x)
```

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx = \int \frac{C \tan(fx + e)^2 + B \tan(fx + e) + A}{(b \tan(fx + e) + a)^{3/2} \sqrt{d \tan(fx + e) + c}} dx$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/((b*tan(f*x + e) + a)^(3/2)*sqrt(d*tan(f*x + e) + c)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(1/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```

(2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*c + int((sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b**2*
d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d
+ 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*a*b**2*d*f - int((sqrt(t
an(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3
*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*
a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*tan(e + f*x)*b**3*c*f + int((sq
rt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x
)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f
*x)*a**2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a**2*b*d*f - int((sqrt(tan(
e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b*
*2*d + 2*tan(e + f*x)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**
2*d + 2*tan(e + f*x)*a*b*c + a**2*c),x)*a*b**2*c*f + int((sqrt(tan(e + f*x
)*d + c)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x
)**2*a*b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)
*a*b*c + a**2*c),x)*tan(e + f*x)*a**2*b*d*f - int((sqrt(tan(e + f*x)*d + c
)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*
b*d + tan(e + f*x)**2*b**2*c + tan(e + f*x)*a**2*d + 2*tan(e + f*x)*a*b*c
+ a**2*c),x)*tan(e + f*x)*a*b**2*c*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(
tan(e + f*x)*b + a))/(tan(e + f*x)**3*b**2*d + 2*tan(e + f*x)**2*a*b*d ...

```

3.152
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{5/2} \sqrt{c+d \tan(e+fx)}} dx$$

Optimal result	1698
Mathematica [A] (verified)	1699
Rubi [A] (verified)	1699
Maple [F(-1)]	1704
Fricas [F(-1)]	1704
Sympy [F]	1704
Maxima [F(-1)]	1705
Giac [F]	1705
Mupad [F(-1)]	1706
Reduce [F]	1706

Optimal result

Integrand size = 49, antiderivative size = 375

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a - ib)^{5/2} \sqrt{c - id} f}$$

$$- \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a + ib)^{5/2} \sqrt{c + id} f}$$

$$- \frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^{3/2}}$$

$$- \frac{2(5a^3bBd - 2a^4Cd + b^4(3Bc - 2Ad) + ab^3(6Ac - 6cC - Bd) - a^2b^2(3Bc + 8Ad - 4Cd)) \sqrt{c + d \tan(e + fx)}}{3(a^2 + b^2)^2 (bc - ad)^2 f \sqrt{a + b \tan(e + fx)}}$$

output

```
-(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(5/2)/(c-I*d)^(1/2)/f-(B-I*(A-C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a+I*b)^(5/2)/(c+I*d)^(1/2)/f-2/3*(A*b^2-a*(B*b-C*a))*(c+d*tan(f*x+e))^(1/2)/(a^2+b^2)/(-a*d+b*c)/f/(a+b*tan(f*x+e))^(3/2)-2/3*(5*a^3*b*B*d-2*a^4*C*d+b^4*(-2*A*d+3*B*c)+a*b^3*(6*A*c-B*d-6*C*c)-a^2*b^2*(8*A*d+3*B*c-4*C*d))*(c+d*tan(f*x+e))^(1/2)/(a^2+b^2)^2/(-a*d+b*c)^2/f/(a+b*tan(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 4.09 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.03

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx = \frac{3(a+ib)^2(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{-c+id}\sqrt{a+b\tan(e+fx)}}{\sqrt{-a+ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{-a+ib}\sqrt{-c+id}} + \frac{3i(a-ib)^2}{\dots}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(5/2)*Sqrt[c + d*Tan[e + f*x]]),x]
```

output

```
((3*(a + I*b)^2*(I*A + B - I*C)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + I*b]*Sqrt[-c + I*d]) + ((3*I)*(a - I*b)^2*(A + I*B - C)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]) + (2*(a^2 + b^2)*(A*b^2 + a*(-(b*B) + a*C))*Sqrt[c + d*Tan[e + f*x]])/((-b*c) + a*d)*(a + b*Tan[e + f*x])^(3/2) + (2*(-5*a^3*b*B*d + 2*a^4*C*d + b^4*(-3*B*c + 2*A*d) + a*b^3*(-6*A*c + 6*c*C + B*d) + a^2*b^2*(3*B*c + 8*A*d - 4*C*d))*Sqrt[c + d*Tan[e + f*x]])/((b*c - a*d)^2*Sqrt[a + b*Tan[e + f*x]])/(3*(a^2 + b^2)^2*f)
```

Rubi [A] (verified)Time = 3.91 (sec) , antiderivative size = 458, normalized size of antiderivative = 1.22, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx$$

↓ 4132

$$2 \int \frac{2Ab^2 + 2(Ab^2 - a(bB - aC))d \tan^2(e + fx) - 3aA(bc - ad) - (bB - aC)(3bc - ad) + 3(Ab - Cb - aB)(bc - ad) \tan(e + fx)}{2(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{3(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \\ \frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{2Ab^2 + 2(Ab^2 - a(bB - aC))d \tan^2(e + fx) - 3aA(bc - ad) - (bB - aC)(3bc - ad) + 3(Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{3(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \\ \frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{2Ab^2 + 2(Ab^2 - a(bB - aC))d \tan(e + fx)^2 - 3aA(bc - ad) - (bB - aC)(3bc - ad) + 3(Ab - Cb - aB)(bc - ad) \tan(e + fx)}{(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} dx$$

$$\frac{3(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \\ \frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 4132

$$\frac{2\sqrt{c + d \tan(e + fx)}(-2a^4Cd + 5a^3bBd - a^2b^2(8Ad + 3Bc - 4Cd) + ab^3(6Ac - Bd - 6cC) + b^4(3Bc - 2Ad))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}} - \frac{2 \int \frac{3((A - C)a^2 + 2bBa - b^2(A - C))(bc - ad)}{2\sqrt{a + b \tan(e + fx)}} dx}{(a^2 + b^2)(bc - ad)}$$

$$\frac{3(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \\ \frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 27

$$\frac{2\sqrt{c + d \tan(e + fx)}(-2a^4Cd + 5a^3bBd - a^2b^2(8Ad + 3Bc - 4Cd) + ab^3(6Ac - Bd - 6cC) + b^4(3Bc - 2Ad))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}} - \frac{3 \int \frac{((A - C)a^2 + 2bBa - b^2(A - C))(bc - ad)}{\sqrt{a + b \tan(e + fx)}} dx}{(a^2 + b^2)(bc - ad)}$$

$$\frac{3(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}} \\ \frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}}$$

↓ 3042

$$\begin{aligned}
 & \frac{2\sqrt{c+d\tan(e+fx)}(-2a^4Cd+5a^3bBd-a^2b^2(8Ad+3Bc-4Cd)+ab^3(6Ac-Bd-6cC)+b^4(3Bc-2Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} - 3 \int \frac{((A-C)a^2+2bBa-b^2(A-C))(bc-a}{\sqrt{a+b\tan(e+fx)}} \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{3f(a^2+b^2)(bc-ad)(a+b\tan(e+fx))^{3/2}} \quad 3(a^2+b^2)(bc-ad) \\
 & \quad \downarrow 4099 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{3f(a^2+b^2)(bc-ad)(a+b\tan(e+fx))^{3/2}} \\
 & \frac{2\sqrt{c+d\tan(e+fx)}(-2a^4Cd+5a^3bBd-a^2b^2(8Ad+3Bc-4Cd)+ab^3(6Ac-Bd-6cC)+b^4(3Bc-2Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} - 3 \left(\frac{1}{2}(a-ib)^2(A+iB-C)(bc-ad)^2 \int \frac{1}{\sqrt{a+b\tan(e+fx)}} \right) \\
 & \quad 3(a^2+b^2)(bc-ad) \\
 & \quad \downarrow 3042 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{3f(a^2+b^2)(bc-ad)(a+b\tan(e+fx))^{3/2}} \\
 & \frac{2\sqrt{c+d\tan(e+fx)}(-2a^4Cd+5a^3bBd-a^2b^2(8Ad+3Bc-4Cd)+ab^3(6Ac-Bd-6cC)+b^4(3Bc-2Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} - 3 \left(\frac{1}{2}(a-ib)^2(A+iB-C)(bc-ad)^2 \int \frac{1}{\sqrt{a+b\tan(e+fx)}} \right) \\
 & \quad 3(a^2+b^2)(bc-ad) \\
 & \quad \downarrow 4098 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{3f(a^2+b^2)(bc-ad)(a+b\tan(e+fx))^{3/2}} \\
 & \frac{2\sqrt{c+d\tan(e+fx)}(-2a^4Cd+5a^3bBd-a^2b^2(8Ad+3Bc-4Cd)+ab^3(6Ac-Bd-6cC)+b^4(3Bc-2Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} - 3 \left(\frac{(a+ib)^2(A-iB-C)(bc-ad)^2 \int \frac{1}{(1-i\tan(e+fx))\sqrt{a+b\tan(e+fx)}} \right) \\
 & \quad 3(a^2+b^2)(bc-ad) \\
 & \quad \downarrow 104 \\
 & \frac{2(Ab^2-a(bB-aC))\sqrt{c+d\tan(e+fx)}}{3f(a^2+b^2)(bc-ad)(a+b\tan(e+fx))^{3/2}} \\
 & \frac{2\sqrt{c+d\tan(e+fx)}(-2a^4Cd+5a^3bBd-a^2b^2(8Ad+3Bc-4Cd)+ab^3(6Ac-Bd-6cC)+b^4(3Bc-2Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}} - 3 \left(\frac{(a-ib)^2(A+iB-C)(bc-ad)^2 \int \frac{1}{(1-i\tan(e+fx))\sqrt{a+b\tan(e+fx)}} \right) \\
 & \quad 3(a^2+b^2)(bc-ad) \\
 & \quad \downarrow 221
 \end{aligned}$$

$$\frac{2(Ab^2 - a(bB - aC)) \sqrt{c + d \tan(e + fx)}}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2}} - \frac{2\sqrt{c+d \tan(e+fx)}(-2a^4Cd+5a^3bBd-a^2b^2(8Ad+3Bc-4Cd)+ab^3(6Ac-Bd-6cC)+b^4(3Bc-2Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}} - \frac{3 \left(\frac{i(a-ib)^2(A+ib-C)(bc-ad)^2 \arctan\left(\frac{f\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}{f\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{f\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}} \right)}{3(a^2 + b^2)(bc - ad)}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(5/2)*Sqrt[c + d*Tan[e + f*x]]),x]`

output `(-2*(A*b^2 - a*(b*B - a*C))*Sqrt[c + d*Tan[e + f*x]]/(3*(a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])^(3/2)) - ((-3*((-I)*(a + I*b)^2*(A - I*B - C)*(b*c - a*d)^2*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a - I*b]*Sqrt[c - I*d]*f) + (I*(a - I*b)^2*(A + I*B - C)*(b*c - a*d)^2*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]*f)))/((a^2 + b^2)*(b*c - a*d)) + (2*(5*a^3*b*B*d - 2*a^4*C*d + b^4*(3*B*c - 2*A*d) + a*b^3*(6*A*c - 6*c*C - B*d) - a^2*b^2*(3*B*c + 8*A*d - 4*C*d))*Sqrt[c + d*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e + f*x]]))/(3*(a^2 + b^2)*(b*c - a*d))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 221 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan(fx + e) + C \tan^2(fx + e)}{(a + b \tan(fx + e))^{\frac{5}{2}} \sqrt{c + d \tan(fx + e)}} dx$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(1/2),x)`

output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(1/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{5}{2}} \sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{5}{2}} \sqrt{c + d \tan(e + fx)}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{5}{2}} \sqrt{c + d \tan(e + fx)}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(5/2)/(c+d*tan(f*x+e))**(1/2),x)`

output

```
Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**(5/2)*sqrt(c + d*tan(e + f*x))), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx = \text{Timed out}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx = \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{(b \tan(fx + e) + a)^{5/2} \sqrt{d \tan(fx + e) + c}} dx$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(1/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/((b*tan(f*x + e) + a)^(5/2)*sqrt(d*tan(f*x + e) + c)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(5/2)*(c + d*tan(e + f*x))^(1/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} \sqrt{c + d \tan(e + fx)}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(1/2),x)
```

output

```
(4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b*c*d +
6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*c*d - 2*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b*c**2 + 3*int((sqrt(tan(e + f*x)*d
+ c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**3*d + 3*ta
n(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2*a**2*b
*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f*x)*a**
2*b*c + a**3*c),x)*tan(e + f*x)**2*a**2*b**3*d**2*f - 6*int((sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**3*d
+ 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)**2
*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e + f
*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*a*b**4*c*d*f + 3*int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**
3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*x)
**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(e
+ f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)**2*b**5*c**2*f + 6*int((sqrt(tan
(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b
**3*d + 3*tan(e + f*x)**3*a*b**2*d + tan(e + f*x)**3*b**3*c + 3*tan(e + f*
x)**2*a**2*b*d + 3*tan(e + f*x)**2*a*b**2*c + tan(e + f*x)*a**3*d + 3*tan(
e + f*x)*a**2*b*c + a**3*c),x)*tan(e + f*x)*a**3*b**2*d**2*f - 12*int((sqr
t(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f...
```


3.153
$$\int \frac{(a+b \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1708
Mathematica [B] (verified)	1709
Rubi [A] (verified)	1710
Maple [F(-1)]	1715
Fricas [F(-1)]	1716
Sympy [F]	1716
Maxima [F(-1)]	1716
Giac [F]	1717
Mupad [F(-1)]	1717
Reduce [F]	1717

Optimal result

Integrand size = 49, antiderivative size = 528

$$\int \frac{(a+b \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx =$$

$$-\frac{(a-ib)^{5/2}(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c-id)^{3/2}f}$$

$$-\frac{(a+ib)^{5/2}(B-i(A-C))\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c+id)^{3/2}f}$$

$$+\frac{\sqrt{b}(15a^2Cd^2-10abd(3cC-2Bd)+b^2(15c^2C-12Bcd+8(A-C)d^2))\operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{4d^{7/2}f}$$

$$-\frac{2(c^2C-Bcd+Ad^2)(a+b \tan(e+fx))^{5/2}}{d(c^2+d^2)f\sqrt{c+d \tan(e+fx)}}$$

$$-\frac{b(3(bc-ad)(5c^2C-4Bcd+(4A+C)d^2)-4d^2((A-C)(bc-ad)+B(ac+bd)))\sqrt{a+b \tan(e+fx)}}{4d^3(c^2+d^2)f}$$

$$+\frac{b(5c^2C-4Bcd+(4A+C)d^2)(a+b \tan(e+fx))^{3/2}\sqrt{c+d \tan(e+fx)}}{2d^2(c^2+d^2)f}$$

output

```

-(a-I*b)^(5/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(3/2)/f-(a+I*b)^(5/2)*(B-I*(A-
C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(c+I*d)^(3/2)/f+1/4*b^(1/2)*(15*a^2*C*d^2-10*a*b*d*(-2*B*d+3*
C*c)+b^2*(15*c^2*C-12*B*c*d+8*(A-C)*d^2))*arctanh(d^(1/2)*(a+b*tan(f*x+e))
^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/d^(7/2)/f-2*(A*d^2-B*c*d+C*c^2)*(a+
b*tan(f*x+e))^(5/2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)-1/4*b*(3*(-a*d+b*
c)*(5*c^2*C-4*B*c*d+(4*A+C)*d^2)-4*d^2*((A-C)*(-a*d+b*c)+B*(a*c+b*d)))*(a+
b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e))^(1/2)/d^3/(c^2+d^2)/f+1/2*b*(5*c^2*C-
4*B*c*d+(4*A+C)*d^2)*(a+b*tan(f*x+e))^(3/2)*(c+d*tan(f*x+e))^(1/2)/d^2/(c^
2+d^2)/f

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2245 vs. $2(528) = 1056$.

Time = 8.19 (sec) , antiderivative size = 2245, normalized size of antiderivative = 4.25

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Result too large to show}$$

input

```

Integrate[((a + b*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(c + d*Tan[e + f*x])^(3/2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^(5/2))/(2*d*f*Sqrt[c + d*Tan[e + f*x]]) + (((-5*b*
c*C + 4*b*B*d + 5*a*C*d)*(a + b*Tan[e + f*x])^(3/2))/(2*d*f*Sqrt[c + d*Tan
[e + f*x]]) + ((8*(-a + I*b)^(5/2)*(I*A + B - I*C)*d^2*ArcTanh[(Sqrt[-c +
I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])
/((-c + I*d)^(3/2)*f) - (8*(a + I*b)^(5/2)*(B - I*(A - C))*d^2*ArcTanh[(Sq
rt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*
x]])])/((c + I*d)^(3/2)*f) + (8*(a - I*b)^2*(I*A + B - I*C)*d^2*Sqrt[a + b
*Tan[e + f*x]])/((c - I*d)*f*Sqrt[c + d*Tan[e + f*x]]) + (8*(a + I*b)^2*(B
- I*(A - C))*d^2*Sqrt[a + b*Tan[e + f*x]])/((c + I*d)*f*Sqrt[c + d*Tan[e
+ f*x]]) + (30*a^2*C*(b*c - a*d)*(b/((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c -
a*d)))^(3/2)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))^2*Sqrt[(b*(c + d*
Tan[e + f*x]))/(b*c - a*d)]*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*
((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))))*(-((b*d*(a + b*Tan[e + f*x])
)/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))*(-1 - (b*d*(a
+ b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)
)))) - (Sqrt[b]*Sqrt[d]*ArcSinh[(Sqrt[b]*Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])
/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)])]*Sqrt[a
+ b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - (a*b*d)/(b
*c - a*d)]*Sqrt[1 + (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c
- a*d) - (a*b*d)/(b*c - a*d))])))/(b^2*f*Sqrt[a + b*Tan[e + f*x]]*Sqrt...
```

Rubi [A] (verified)

Time = 7.16 (sec) , antiderivative size = 555, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4128, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$2 \int \frac{(a+b \tan(e+fx))^{3/2} (b(5C^2-4Bdc+(4A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+5bd)+(5bc-ad)(cC-Bd))}{2\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\int \frac{(a+b \tan(e+fx))^{3/2} (b(5C^2-4Bdc+(4A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+5bd)+(5bc-ad)(cC-Bd))}{\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^{3/2} (b(5C^2-4Bdc+(4A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+5bd)+(5bc-ad)(cC-Bd))}{\sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4130

$$\int -\frac{\sqrt{a+b \tan(e+fx)} (-4((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d)) \tan(e+fx)d^2-4a(Ad(ac+5bd)+(5bc-ad)(cC-Bd))+b(3(bc-ad)(5C^2-4Bdc+(4A+C)d^2))}{2\sqrt{c+d \tan(e+fx)} 2d}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{b(d^2(4A+C)-4Bcd+5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \int \frac{\sqrt{a+b \tan(e+fx)} (-4((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d))}{2\sqrt{c+d \tan(e+fx)} 2d}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{b(d^2(4A+C)-4Bcd+5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{\int \frac{\sqrt{a+b \tan(e+fx)}(-4((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d))}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4130

$$\frac{b(d^2(4A+C)-4Bcd+5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{\int \frac{c(15Cc^3-12Bdc^2+(8A+7C)d^2c-4Bd^3)b^3-2ad(15Cc^3-10Bdc^2+3(4A+C)d)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\frac{b(d^2(4A+C)-4Bcd+5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{b\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}(3(bc-ad)(d^2(4A+C)-4Bcd+5c^2C)-4d^2((A+C)d^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d)))}{df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\frac{b(d^2(4A+C)-4Bcd+5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{b\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}(3(bc-ad)(d^2(4A+C)-4Bcd+5c^2C)-4d^2((A+C)d^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d)))}{df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 4138

$$\frac{b(d^2(4A+C)-4Bcd+5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{b\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}(3(bc-ad)(d^2(4A+C)-4Bcd+5c^2C)-4d^2((A+C)d^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d)))}{df}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

↓ 2348

$$\frac{b(5C^2 - 4Bdc + (4A+C)d^2)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{b(3(bc-ad)(5C^2 - 4Bdc + (4A+C)d^2) - 4d^2((A-C)(bc-ad) + B(ac+bd))) \sqrt{a+b \tan(e+fx)}}{df}$$

$$\frac{2(Cc^2 - Bdc + Ad^2)(a+b \tan(e+fx))^{5/2}}{d(c^2 + d^2) f \sqrt{c+d \tan(e+fx)}}$$

↓ 2009

$$- \frac{2(Ad^2 - Bcd + c^2C)(a+b \tan(e+fx))^{5/2}}{df(c^2 + d^2) \sqrt{c+d \tan(e+fx)}} +$$

$$\frac{b(d^2(4A+C) - 4Bcd + 5c^2C)(a+b \tan(e+fx))^{3/2} \sqrt{c+d \tan(e+fx)}}{2df} - \frac{b\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)} (3(bc-ad)(d^2(4A+C) - 4Bcd + 5c^2C) - 4d^2((A-C)(bc-ad) + B(ac+bd)))}{df}$$

input `Int(((a + b*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x)`

output `(-2*(c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^(5/2))/(d*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]]) + ((b*(5*c^2*C - 4*B*c*d + (4*A + C)*d^2)*(a + b*Tan[e + f*x])^(3/2)*Sqrt[c + d*Tan[e + f*x]])/(2*d*f) - (-1/2*((-8*(a - I*b)^(5/2)*(I*A + B - I*C)*(c + I*d)*d^3*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c - I*d] + (8*(a + I*b)^(5/2)*(A + I*B - C)*d^3*(I*c + d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c + I*d] + (2*Sqrt[b]*(c^2 + d^2)*(15*a^2*C*d^2 - 10*a*b*d*(3*c*C - 2*B*d) + b^2*(15*c^2*C - 12*B*c*d + 8*(A - C)*d^2))*ArcTanh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[d])/(d*f) + (b*(3*(b*c - a*d)*(5*c^2*C - 4*B*c*d + (4*A + C)*d^2) - 4*d^2*((A - C)*(b*c - a*d) + B*(a*c + b*d)))*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]])/(d*f))/(4*d))/(d*(c^2 + d^2))`

Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2348 `Int[(Px_)*((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4128 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(c + d \tan(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(3/2),x)
```

output

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(3/2),x)
```


Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

input

```
integrate((a+b*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)
```

output

```
Integral((a + b*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(3/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e)}{(d \tan(fx + e))^{3/2}}$$

input `integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(5/2)/(d*tan(f*x + e) + c)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))^{5/2} (C \tan(e + fx) + B)}{(c + d \tan(e + fx))^{3/2}}$$

input `int(((a + b*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2),x)`

output `int(((a + b*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2), x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input `int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)`

output

```
( - 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**2*b**2 + int((s
qrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)/(tan(e +
f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a*b**2*c*d**2*f
- int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**4)
/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*b**3*c
**2*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f
*x)**4)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*b**2*c**2*
d*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*
**4)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*b**3*c**3*f + 2*
int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(t
an(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a**2*b*c*
d**2*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*
x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*
a*b**3*d**2*f - 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*t
an(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(
e + f*x)*a*b**2*c**2*d*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)
*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2
),x)*tan(e + f*x)*b**4*c*d*f + 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e
+ f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d
+ c**2),x)*a**2*b*c**2*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e +...
```

3.154
$$\int \frac{(a+b \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1719
Mathematica [B] (verified)	1720
Rubi [A] (verified)	1721
Maple [F(-1)]	1726
Fricas [F(-1)]	1726
Sympy [F]	1726
Maxima [F(-1)]	1727
Giac [F]	1727
Mupad [F(-1)]	1728
Reduce [F]	1728

Optimal result

Integrand size = 49, antiderivative size = 380

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(a - ib)^{3/2} (iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c - id)^{3/2} f}$$

$$- \frac{(a + ib)^{3/2} (B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c + id)^{3/2} f}$$

$$- \frac{\sqrt{b}(3bcC - 2bBd - 3aCd) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{d^{5/2} f}$$

$$- \frac{2(c^2C - Bcd + Ad^2) (a + b \tan(e + fx))^{3/2}}{d(c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

$$+ \frac{b(3c^2C - 2Bcd + (2A + C)d^2) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{d^2 (c^2 + d^2) f}$$

output

```

-(a-I*b)^(3/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(3/2)/f-(a+I*b)^(3/2)*(B-I*(A
-C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(c+I*d)^(3/2)/f-b^(1/2)*(-2*B*b*d-3*C*a*d+3*C*b*c)*arctanh(d^(
1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/d^(5/2)/f-2*(
A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(3/2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(
1/2)+b*(3*c^2*C-2*B*c*d+(2*A+C)*d^2)*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f*x+e
))^1/2)/d^2/(c^2+d^2)/f

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2141 vs. $2(380) = 760$.

Time = 7.02 (sec) , antiderivative size = 2141, normalized size of antiderivative = 5.63

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Result too large to show}$$

input

```

Integrate[((a + b*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(c + d*Tan[e + f*x])^(3/2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^(3/2))/(d*f*Sqrt[c + d*Tan[e + f*x]]) + ((-2*(-a +
I*b)^(3/2)*(B + I*(A - C))*d*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f
*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((-c + I*d)^(3/2)*f) - (
2*(a + I*b)^(3/2)*(B - I*(A - C))*d*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[
e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((c + I*d)^(3/2)*f)
+ (2*(I*a + b)*(A - I*B - C)*d*Sqrt[a + b*Tan[e + f*x]])/((c - I*d)*f*Sqrt
[c + d*Tan[e + f*x]]) - (2*(I*a - b)*(A + I*B - C)*d*Sqrt[a + b*Tan[e + f*
x]])/((c + I*d)*f*Sqrt[c + d*Tan[e + f*x]]) - (6*c*C*(b*c - a*d)*(b/((b^2*
c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))^(3/2)*((b^2*c)/(b*c - a*d) - (a*b*d
)/(b*c - a*d))^2*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)]*(-1 - (b*d*(a
+ b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)
)))*(-((b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b
*d)/(b*c - a*d))*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b
*c - a*d) - (a*b*d)/(b*c - a*d)))))) - (Sqrt[b]*Sqrt[d]*ArcSinh[(Sqrt[b]*S
qrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d)
- (a*b*d)/(b*c - a*d)])]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[
(b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)]*Sqrt[1 + (b*d*(a + b*Tan[e + f*
x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))])))/(b*d^2*
f*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]*Sqrt[1 + (b*d*(a + b*T
an[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))]...
```

Rubi [A] (verified)

Time = 4.25 (sec) , antiderivative size = 417, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.204$, Rules used = {3042, 4128, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$\frac{2 \int \frac{\sqrt{a+b \tan(e+fx)}(b(3Cc^2-2Bdc+(2A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{2\sqrt{c+d \tan(e+fx)}}}{d(c^2+d^2)} \frac{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}}$$

↓ 27

$$\frac{\int \frac{\sqrt{a+b \tan(e+fx)}(b(3Cc^2-2Bdc+(2A+C)d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{\sqrt{c+d \tan(e+fx)}}}{d(c^2+d^2)} \frac{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}}$$

↓ 3042

$$\frac{\int \frac{\sqrt{a+b \tan(e+fx)}(b(3Cc^2-2Bdc+(2A+C)d^2) \tan^2(e+fx)^2+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+3bd)+(3bc-ad)(cC-Bd))}{\sqrt{c+d \tan(e+fx)}}}{d(c^2+d^2)} \frac{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}}$$

↓ 4130

$$\frac{\int -\frac{2((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d)) \tan(e+fx)d^2-2a(Ad(ac+3bd)+(3bc-ad)(cC-Bd))d+b(3bcC-3adC-2bBd)(c^2+d^2) \tan^2(e+fx)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}}{d}}{d(c^2+d^2)} \frac{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}}$$

↓ 27

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df} - \frac{\int \frac{-2((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d)) \tan(e+fx)d^2-2a(Ad(ac+3bd)+(3bc-ad)(cC-Bd))d+b(3bcC-3adC-2bBd)(c^2+d^2) \tan^2(e+fx)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}}{d}}{d(c^2+d^2)}$$

↓ 3042

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{df} - \frac{\int \frac{-2((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d))\tan(e+fx)d^2-2a(Ad^2-2Bcd+c^2C)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx}{d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

4138

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{df} - \frac{\int \frac{-2((Bc-(A-C)d)a^2+2b(Ac-Cc+Bd)a-b^2(Bc-(A-C)d))\tan(e+fx)d^2-2a(Ad^2-2Bcd+c^2C)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} dx}{d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}}$$

2348

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} +$$

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{df} - \frac{\int \left(\frac{b(3bcC-3adC-2bBd)(c^2+d^2)}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} + \frac{2Ab^2d^3-2a^2Ad^3+4abBd^3+2a^2Cd^3-2b^2C^2d^3}{\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} \right) dx}{d(c^2+d^2)}$$

2009

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{df (c^2 + d^2) \sqrt{c + d \tan(e + fx)}} +$$

$$\frac{b(d^2(2A+C)-2Bcd+3c^2C)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}}{df} - \frac{2d^2(a-ib)^{3/2}(-d+ic)(A-ib-C)\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b\tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{c-id}} - \frac{2d^2(a+ib)^{3/2}(-d-ic)(A+ib-C)\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b\tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d\tan(e+fx)}}\right)}{\sqrt{c+id}}$$

$d(c^2 + d^2)$

input

```
Int[((a + b*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x]
```


output

$$\begin{aligned} & (-2*(c^2*C - B*c*d + A*d^2)*(a + b*\text{Tan}[e + f*x])^{(3/2)})/(d*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]) + (-1/2*((2*(a - I*b)^{(3/2)}*(A - I*B - C)*(I*c - d)*d^2*\text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/(\text{Sqrt}[c - I*d] - (2*(a + I*b)^{(3/2)}*(I*A - B - I*C)*(c - I*d)*d^2*\text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/(\text{Sqrt}[c + I*d] + (2*\text{Sqrt}[b]*(3*b*c*C - 2*b*B*d - 3*a*C*d)*(c^2 + d^2)*\text{ArcTanh}[(\text{Sqrt}[d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/(\text{Sqrt}[d])/(d*f) + (b*(3*c^2*C - 2*B*c*d + (2*A + C)*d^2)*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])/(d*f))/(d*(c^2 + d^2)) \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] \text{ /; FreeQ}[b, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 2348

$$\begin{aligned} & \text{Int}[(P_x)*((c_) + (d_)*(x_))^{(m_)}*((e_) + (f_)*(x_))^{(n_)}*((a_) + (b_)* \\ & (x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P_x, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0]) \end{aligned}$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4128

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]

```

rule 4130

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))

```

rule 4138

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]

```

Maple [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan^2(fx + e)^2)}{(c + d \tan(fx + e))^{\frac{3}{2}}} dx$$

input `int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)`

output `int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))^{\frac{3}{2}} (A + B \tan(e + fx))}{(c + d \tan(e + fx))^{3/2}}$$

input `integrate((a+b*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)`

output

```
Integral((a + b*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)*
*2)/(c + d*tan(e + f*x))**(3/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(
f*x+e))^(3/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^{3/2}}{(d \tan(fx + e) + c)^{3/2}} dx$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(
f*x+e))^(3/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/
2)/(d*tan(f*x + e) + c)^(3/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(a + b \tan(e + fx))^{3/2} (C \tan(e + fx) + \dots)}{(c + d \tan(e + fx))^{3/2}}$$

input `int(((a + b*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2),x)`

output `int(((a + b*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2), x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(\tan(fx + e) b + a)^{3/2} (A + B \tan(fx + e))}{(d \tan(fx + e) + c)}$$

input `int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)`

output `int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)`

3.155
$$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1729
Mathematica [A] (verified)	1730
Rubi [A] (verified)	1730
Maple [F(-1)]	1733
Fricas [F(-1)]	1734
Sympy [F]	1734
Maxima [F(-1)]	1734
Giac [F]	1735
Mupad [F(-1)]	1735
Reduce [F]	1735

Optimal result

Integrand size = 49, antiderivative size = 299

$$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{3/2}} dx =$$

$$-\frac{\sqrt{a-ib}(iA+B-iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c-id)^{3/2}f}$$

$$-\frac{\sqrt{a+ib}(B-i(A-C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c+id)^{3/2}f}$$

$$+\frac{2\sqrt{b}C \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{d^{3/2}f} - \frac{2(c^2C - Bcd + Ad^2) \sqrt{a+b \tan(e+fx)}}{d(c^2 + d^2) f \sqrt{c+d \tan(e+fx)}}$$

output

```
-(a-I*b)^(1/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(3/2)/f-(a+I*b)^(1/2)*(B-I*(A-C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c+I*d)^(3/2)/f+2*b^(1/2)*C*arctanh(d^(1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/d^(3/2)/f-2*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(1/2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 3.50 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.35

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \frac{\sqrt{-a+ib}(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{-c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{-a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(-c+id)^{3/2}}$$

input `Integrate[(Sqrt[a + b*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(3/2),x]`

output `((Sqrt[-a + I*b]*(I*A + B - I*C)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(-c + I*d)^(3/2) + (I*Sqrt[a + I*b]*(A + I*B - C)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/((c + I*d)^(3/2) + ((B + I*(A - C))*Sqrt[a + b*Tan[e + f*x]])/((c - I*d)*Sqrt[c + d*Tan[e + f*x]]) + ((-I)*A + B + I*C)*Sqrt[a + b*Tan[e + f*x]])/((c + I*d)*Sqrt[c + d*Tan[e + f*x]]) + (2*C*(-(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]]) + Sqrt[b*c - a*d])*ArcSinh[(Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/Sqrt[b*c - a*d])*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d))]/(d^(3/2)*Sqrt[c + d*Tan[e + f*x]])/f`

Rubi [A] (verified)

Time = 2.46 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4128, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{3/2}} dx$$

↓ 4128

$$\begin{aligned}
 & \frac{2 \int \frac{bC(c^2+d^2) \tan^2(e+fx) + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+bd) + (bc-ad)(cC-Bd)}{2\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{d(c^2+d^2)}{2(Ad^2 - Bcd + c^2C) \sqrt{a+b \tan(e+fx)}}} \\
 & \qquad \qquad \qquad \frac{d(c^2+d^2)}{df(c^2+d^2) \sqrt{c+d \tan(e+fx)}} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{\int \frac{bC(c^2+d^2) \tan^2(e+fx) + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+bd) + (bc-ad)(cC-Bd)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{d(c^2+d^2)}{2(Ad^2 - Bcd + c^2C) \sqrt{a+b \tan(e+fx)}}} \\
 & \qquad \qquad \qquad \frac{d(c^2+d^2)}{df(c^2+d^2) \sqrt{c+d \tan(e+fx)}} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{\int \frac{bC(c^2+d^2) \tan(e+fx)^2 + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+bd) + (bc-ad)(cC-Bd)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx}{\frac{d(c^2+d^2)}{2(Ad^2 - Bcd + c^2C) \sqrt{a+b \tan(e+fx)}}} \\
 & \qquad \qquad \qquad \frac{d(c^2+d^2)}{df(c^2+d^2) \sqrt{c+d \tan(e+fx)}} \\
 & \qquad \qquad \qquad \downarrow 4138 \\
 & \frac{\int \frac{bC(c^2+d^2) \tan^2(e+fx) + d((A-C)(bc-ad) + B(ac+bd)) \tan(e+fx) + Ad(ac+bd) + (bc-ad)(cC-Bd)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)} (\tan^2(e+fx)+1)}}{df(c^2+d^2)} d \tan(e+fx) \\
 & \qquad \qquad \qquad \frac{d(c^2+d^2)}{2(Ad^2 - Bcd + c^2C) \sqrt{a+b \tan(e+fx)}} \\
 & \qquad \qquad \qquad \frac{d(c^2+d^2)}{df(c^2+d^2) \sqrt{c+d \tan(e+fx)}} \\
 & \qquad \qquad \qquad \downarrow 2348 \\
 & \frac{\int \left(\frac{bC(c^2+d^2)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} + \frac{aAd^2 - bBd^2 - aCd^2 - Abcd - aBcd + bcCd + i(Abd^2 + aBd^2 - bCd^2 + aAc d - bBcd - aCd)}{2(i - \tan(e+fx)) \sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} + \frac{-aAd^2 + \dots}{df(c^2+d^2)} \right)}{\frac{d(c^2+d^2)}{2(Ad^2 - Bcd + c^2C) \sqrt{a+b \tan(e+fx)}}} + \\
 & \qquad \qquad \qquad \frac{d(c^2+d^2)}{df(c^2+d^2) \sqrt{c+d \tan(e+fx)}} \\
 & \qquad \qquad \qquad \downarrow 2009 \\
 & \frac{d\sqrt{a-ib}(c+id)(iA+B-iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c-id}} + \frac{d\sqrt{a+ib}(d+ic)(A+iB-C) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{c+id}} + \frac{2\sqrt{b}C}{df(c^2+d^2)}
 \end{aligned}$$

input $\text{Int}[(\text{Sqrt}[a + b*\text{Tan}[e + f*x]]*(A + B*\text{Tan}[e + f*x] + C*\text{Tan}[e + f*x]^2))/(c + d*\text{Tan}[e + f*x])^{(3/2)},x]$

output $(-((\text{Sqrt}[a - I*b]*(I*A + B - I*C)*(c + I*d)*d*\text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/ \text{Sqrt}[c - I*d]) + (\text{Sqrt}[a + I*b]*(A + I*B - C)*d*(I*c + d)*\text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/ \text{Sqrt}[c + I*d] + (2*\text{Sqrt}[b]*C*(c^2 + d^2)*\text{ArcTanh}[(\text{Sqrt}[d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])])/ \text{Sqrt}[d])/(d*(c^2 + d^2)*f) - (2*(c^2*C - B*c*d + A*d^2)*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(d*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])$

Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 2348 $\text{Int}[(P_x)*((c_) + (d_)*(x_))^{(m_)}*((e_) + (f_)*(x_))^{(n_)}*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P_x, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan (fx + e)} (A + B \tan (fx + e) + C \tan (fx + e)^2)}{(c + d \tan (fx + e))^{\frac{3}{2}}} dx$$

input

```
int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(3/2),x)
```

output

```
int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(3/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((a+b*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral(sqrt(a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(3/2), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output Timed out

Giac [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e))}{(d \tan(fx + e))^{3/2}}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)/(d*tan(f*x + e) + c)^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \int \frac{\sqrt{a + b \tan(e + fx)}(C \tan(e + fx) + B)}{(c + d \tan(e + fx))^{3/2}}$$

input `int(((a + b*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2),x)`

output `int(((a + b*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(3/2), x)`

Reduce [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(3/2),x)`

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b**2 + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*a*c*d**2*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*tan(e + f*x)*b*c**2*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*a*c**2*d*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2)/(tan(e + f*x)**2*d**2 + 2*tan(e + f*x)*c*d + c**2),x)*b*c**3*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*tan(e + f*x)*a**2*b*d**2*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*tan(e + f*x)*a*b**2*c*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*a**2*b*c*d*f - int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + ...
```

3.156
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1737
Mathematica [A] (verified)	1738
Rubi [A] (verified)	1738
Maple [F(-1)]	1742
Fricas [F(-1)]	1742
Sympy [F]	1742
Maxima [F]	1743
Giac [F]	1743
Mupad [F(-1)]	1744
Reduce [F]	1744

Optimal result

Integrand size = 49, antiderivative size = 251

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(B + i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c - id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a - ib}(c - id)^{3/2} f}$$

$$+ \frac{(iA - B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a + ib}(c + id)^{3/2} f}$$

$$+ \frac{2(c^2 C - Bcd + Ad^2) \sqrt{a + b \tan(e + fx)}}{(bc - ad)(c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

output

```

-(B+I*(A-C))*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c
+d*tan(f*x+e))^(1/2))/(a-I*b)^(1/2)/(c-I*d)^(3/2)/f+(I*A-B-I*C)*arctanh((c
+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(
a+I*b)^(1/2)/(c+I*d)^(3/2)/f+2*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(1/2)/
(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)
    
```

Mathematica [A] (verified)

Time = 2.17 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.10

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(bc - ad) \left(\frac{(iA+B-iC)(c+id) \operatorname{arctanh}\left(\frac{\sqrt{-c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{-a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{-a+ib}\sqrt{-c+id}} + \frac{(A+iB-C)(ic+d) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a+ib}\sqrt{c+id}} \right) + 2(c$$

$$(-bc + ad)(c^2 + d^2) f$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

```
-(((b*c - a*d)*(((I*A + B - I*C)*(c + I*d)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + I*b]*Sqrt[-c + I*d]) + ((A + I*B - C)*(I*c + d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d])) + (2*(c^2*C - B*c*d + A*d^2)*Sqrt[a + b*Tan[e + f*x]])/Sqrt[c + d*Tan[e + f*x]]/((-b*c) + a*d)*(c^2 + d^2)*f))
```

Rubi [A] (verified)Time = 2.36 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.20, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.184$, Rules used = {3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx$$

$$\downarrow \text{4132}$$

$$\begin{aligned}
 & \frac{2 \int \frac{(bc-ad)(Ac-Cc+Bd)+(bc-ad)(Bc-(A-C)d) \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{(c^2+d^2)(bc-ad)} + \\
 & \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{(bc-ad)(Ac-Cc+Bd)+(bc-ad)(Bc-(A-C)d) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{(c^2+d^2)(bc-ad)} + \\
 & \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{(bc-ad)(Ac-Cc+Bd)+(bc-ad)(Bc-(A-C)d) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{(c^2+d^2)(bc-ad)} + \\
 & \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} \\
 & \quad \downarrow 4099 \\
 & \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} + \\
 & \frac{\frac{1}{2}(c-id)(A+iB-C)(bc-ad) \int \frac{1-i \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(c+id)(A-iB-C)(bc-ad) \int \frac{i}{\sqrt{a+b \tan(e+fx)}} dx}{(c^2+d^2)(bc-ad)} \\
 & \quad \downarrow 3042 \\
 & \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} + \\
 & \frac{\frac{1}{2}(c-id)(A+iB-C)(bc-ad) \int \frac{1-i \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx + \frac{1}{2}(c+id)(A-iB-C)(bc-ad) \int \frac{i}{\sqrt{a+b \tan(e+fx)}} dx}{(c^2+d^2)(bc-ad)} \\
 & \quad \downarrow 4098 \\
 & \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} + \\
 & \frac{(c+id)(A-iB-C)(bc-ad) \int \frac{1}{(1-i \tan(e+fx))\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} d \tan(e+fx)}{2f} + \frac{(c-id)(A+iB-C)(bc-ad) \int \frac{1}{(i \tan(e+fx)+1)\sqrt{a+b \tan(e+fx)}} d \tan(e+fx)}{2f} \\
 & \quad \downarrow 104 \\
 & \frac{\dots}{(c^2+d^2)(bc-ad)}
 \end{aligned}$$

$$\frac{\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}} + \frac{(c-id)(A+iB-C)(bc-ad) \int \frac{1}{-ia+b+\frac{(ic-d)(a+b \tan(e+fx))}{c+d \tan(e+fx)}} d \frac{\sqrt{a+b \tan(e+fx)}}{\sqrt{c+d \tan(e+fx)}}}{f} + \frac{(c+id)(A-iB-C)(bc-ad) \int \frac{1}{ia+b-\frac{(ic+d)(a+b \tan(e+fx))}{c+d \tan(e+fx)}} d \frac{\sqrt{a+b \tan(e+fx)}}{\sqrt{c+d \tan(e+fx)}}}{f}}{(c^2 + d^2)(bc - ad)}$$

↓ 221

$$\frac{\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{f(c^2 + d^2)(bc - ad) \sqrt{c + d \tan(e + fx)}} + \frac{i(c-id)(A+iB-C)(bc-ad) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{f\sqrt{a+ib}\sqrt{c+id}} - \frac{i(c+id)(A-iB-C)(bc-ad) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{f\sqrt{a-ib}\sqrt{c-id}}}{(c^2 + d^2)(bc - ad)}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

```
(((-I)*(A - I*B - C)*(c + I*d)*(b*c - a*d)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a - I*b]*Sqrt[c - I*d]*f) + (I*(A + I*B - C)*(c - I*d)*(b*c - a*d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]*f))/((b*c - a*d)*(c^2 + d^2)) + (2*(c^2*C - B*c*d + A*d^2)*Sqrt[a + b*Tan[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 104

```
Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

rule 221 $\text{Int}[(a_.) + (b_.) \cdot (x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 4098 $\text{Int}[(a_.) + (b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(m_.)} \cdot ((A_.) + (B_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[A^2/f \ \text{Subst}[\text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n / (A - B \cdot x)], x], x, \text{Tan}[e + f \cdot x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, m, n\}, x \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A^2 + B^2, 0]$

rule 4099 $\text{Int}[(a_.) + (b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(m_.)} \cdot ((A_.) + (B_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(A + I \cdot B)/2 \ \text{Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^m \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot (1 - I \cdot \text{Tan}[e + f \cdot x]), x], x] + \text{Simp}[(A - I \cdot B)/2 \ \text{Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^m \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot (1 + I \cdot \text{Tan}[e + f \cdot x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, m, n\}, x \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A^2 + B^2, 0]$

rule 4132 $\text{Int}[(a_.) + (b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(m_.)} \cdot ((c_.) + (d_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)])^{(n_.)} \cdot ((A_.) + (B_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)] + (C_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)]^2), x_Symbol] \rightarrow \text{Simp}[(A \cdot b^2 - a \cdot (b \cdot B - a \cdot C)) \cdot (a + b \cdot \text{Tan}[e + f \cdot x])^{(m+1)} \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^{(n+1)} / (f \cdot (m+1) \cdot (b \cdot c - a \cdot d) \cdot (a^2 + b^2)), x] + \text{Simp}[1 / ((m+1) \cdot (b \cdot c - a \cdot d) \cdot (a^2 + b^2)) \ \text{Int}[(a + b \cdot \text{Tan}[e + f \cdot x])^{(m+1)} \cdot (c + d \cdot \text{Tan}[e + f \cdot x])^n \cdot \text{Simp}[A \cdot (a \cdot (b \cdot c - a \cdot d) \cdot (m+1) - b^2 \cdot d \cdot (m+n+2)) + (b \cdot B - a \cdot C) \cdot (b \cdot c \cdot (m+1) + a \cdot d \cdot (n+1)) - (m+1) \cdot (b \cdot c - a \cdot d) \cdot (A \cdot b - a \cdot B - b \cdot C) \cdot \text{Tan}[e + f \cdot x] - d \cdot (A \cdot b^2 - a \cdot (b \cdot B - a \cdot C)) \cdot (m+n+2) \cdot \text{Tan}[e + f \cdot x]^2, x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ !(\text{ILtQ}[n, -1] \ \&\& \ (!\text{IntegerQ}[m] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{NeQ}[a, 0])))$

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan(fx + e) + C \tan(fx + e)^2}{\sqrt{a + b \tan(fx + e)} (c + d \tan(fx + e))^{\frac{3}{2}}} dx$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2),x)`

output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(1/2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(sqrt(a + b*tan(e + f*x))*
(c + d*tan(e + f*x))**(3/2)), x)`

Maxima [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx = \int \frac{C \tan(fx + e)^2 + B \tan(fx + e) + A}{\sqrt{b \tan(fx + e) + a}(d \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/(sqrt(b*tan(f*x + e) + a)
)*(d*tan(f*x + e) + c)^(3/2)), x)`

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx = \int \frac{C \tan(fx + e)^2 + B \tan(fx + e) + A}{\sqrt{b \tan(fx + e) + a}(d \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/(sqrt(b*tan(f*x + e) + a)
)*(d*tan(f*x + e) + c)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(3/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*c + int((sqrt(tan(
e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**3*b*
d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a
*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*tan(e + f*x)*a*b*d**2*f - int((sqr
t(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)
**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e +
f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*tan(e + f*x)*b**2*c*d*f + in
t((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e
+ f*x)**3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*ta
n(e + f*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*a*b*c*d*f - int((sqrt(
tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**
3*b*d**2 + tan(e + f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f
*x)*a*c*d + tan(e + f*x)*b*c**2 + a*c**2),x)*b**2*c**2*f + int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)**3*b*d**2 + tan(e +
f*x)**2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e +
f*x)*b*c**2 + a*c**2),x)*tan(e + f*x)*a**2*d**2*f - int((sqrt(tan(e + f*x)
*d + c)*sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**
2*a*d**2 + 2*tan(e + f*x)**2*b*c*d + 2*tan(e + f*x)*a*c*d + tan(e + f*x)*b
*c**2 + a*c**2),x)*tan(e + f*x)*a*b*c*d*f - int((sqrt(tan(e + f*x)*d + c)*
sqrt(tan(e + f*x)*b + a))/(tan(e + f*x)**3*b*d**2 + tan(e + f*x)**2*a*d...
```

3.157 $\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}} dx$

Optimal result	1746
Mathematica [A] (verified)	1747
Rubi [A] (verified)	1747
Maple [F(-1)]	1752
Fricas [F(-1)]	1752
Sympy [F]	1753
Maxima [F(-1)]	1753
Giac [F]	1753
Mupad [F(-1)]	1754
Reduce [F]	1754

Optimal result

Integrand size = 49, antiderivative size = 383

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a - ib)^{3/2}(c - id)^{3/2} f}$$

$$- \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a + ib)^{3/2}(c + id)^{3/2} f}$$

$$- \frac{2(Ab^2 - a(bB - aC))}{(a^2 + b^2)(bc - ad) f \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}$$

$$- \frac{2d(b^2c(cC - Bd) - abB(c^2 + d^2) + a^2(2c^2C - Bcd + Cd^2) + A(a^2d^2 + b^2(c^2 + 2d^2))) \sqrt{a + b \tan(e + fx)}}{(a^2 + b^2)(bc - ad)^2 (c^2 + d^2) f \sqrt{c + d \tan(e + fx)}}$$

output

```

-(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c
+d*tan(f*x+e))^(1/2))/(a-I*b)^(3/2)/(c-I*d)^(3/2)/f-(B-I*(A-C))*arctanh((c
+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(
a+I*b)^(3/2)/(c+I*d)^(3/2)/f-2*(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/
(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2)-2*d*(b^2*c*(-B*d+C*c)-a*b*B*
(c^2+d^2)+a^2*(-B*c*d+2*C*c^2+C*d^2)+A*(a^2*d^2+b^2*(c^2+2*d^2)))*(a+b*tan
(f*x+e))^(1/2)/(a^2+b^2)/(-a*d+b*c)^2/(c^2+d^2)/f/(c+d*tan(f*x+e))^(1/2)
    
```

Mathematica [A] (verified)

Time = 6.52 (sec) , antiderivative size = 484, normalized size of antiderivative = 1.26

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{2(Ab^2 - a(bB - aC))}{(a^2 + b^2)(bc - ad)f \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}} + \frac{(bc - ad)^2 \left(\frac{(a+ib)(iA+B-iC)(c+id) \operatorname{arctanh}\left(\frac{\sqrt{-c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{-a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{-a+ib}\sqrt{-c+id}} + \frac{(ia+b)(A+iB-C)(c-id) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{\sqrt{a+ib}\sqrt{c+id}} \right)}{2(-bc+ad)(c^2+d^2)f}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

```
(-2*(A*b^2 - a*(b*B - a*C)))/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]) - (2*((b*c - a*d)^2*((a + I*b)*(I*A + B - I*C)*(c + I*d)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + I*b]*Sqrt[-c + I*d]) + ((I*a + b)*(A + I*B - C)*(c - I*d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]))/(2*(-(b*c) + a*d)*(c^2 + d^2)*f) - (2*(-(c*(-(c*(A*b^2 - a*(b*B - a*C))*d) + ((A*b - a*B - b*C)*d*(b*c - a*d))/2)) + (d^2*(2*A*b^2*d - a*A*(b*c - a*d) - (b*B - a*C)*(b*c + a*d)))/2)*Sqrt[a + b*Tan[e + f*x]]/((-b*c) + a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])))/((a^2 + b^2)*(b*c - a*d))
```

Rubi [A] (verified)

Time = 4.13 (sec) , antiderivative size = 466, normalized size of antiderivative = 1.22, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx$$

↓ 4132

$$2 \int \frac{(A+C)da^2 - b(Ac - Cc + Bd)a + 2(Ab^2 - a(bB - aC))d \tan^2(e+fx) - b^2(Bc - 2Ad) + (Ab - Cb - aB)(bc - ad) \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))} \frac{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\int \frac{(A+C)da^2 - b(Ac - Cc + Bd)a + 2(Ab^2 - a(bB - aC))d \tan^2(e+fx) - b^2(Bc - 2Ad) + (Ab - Cb - aB)(bc - ad) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))} \frac{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}$$

↓ 3042

$$\int \frac{(A+C)da^2 - b(Ac - Cc + Bd)a + 2(Ab^2 - a(bB - aC))d \tan(e+fx)^2 - b^2(Bc - 2Ad) + (Ab - Cb - aB)(bc - ad) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))} \frac{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}$$

↓ 4132

$$2 \int \frac{(bBc - b(A - C)d + a(Ac - Cc + Bd))(bc - ad)^2 + (aBc + bCc - bBd + aCd - A(bc + ad)) \tan(e+fx)(bc - ad)^2}{2\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} dx + \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2 + a^2(-Bcd))}{(c^2 + d^2)(bc - ad) f(c^2 + d^2)}$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))} \frac{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}{f(a^2 + b^2)(bc - ad) \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}$$

↓ 27

$$\begin{aligned}
 & \frac{2d\sqrt{a+b\tan(e+fx)}(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \frac{\int \frac{(bBc-b(A-C)d+a(Ac-Cc+Bd))(bc-ad)}{\sqrt{a+b\tan(e+fx)}}}{(a^2+b^2)(bc-ad)} \\
 & \frac{2(Ab^2-a(bB-aC))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{2d\sqrt{a+b\tan(e+fx)}(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \frac{\int \frac{(bBc-b(A-C)d+a(Ac-Cc+Bd))(bc-ad)}{\sqrt{a+b\tan(e+fx)}}}{(a^2+b^2)(bc-ad)} \\
 & \frac{2(Ab^2-a(bB-aC))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} \\
 & \quad \downarrow 4099 \\
 & \frac{2(Ab^2-a(bB-aC))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} - \frac{2d\sqrt{a+b\tan(e+fx)}(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \frac{\frac{1}{2}(a-ib)(c-id)(A+iB-C)(bc-ad)^2 \int \frac{1}{\sqrt{a+b\tan(e+fx)}}}{(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 3042 \\
 & \frac{2(Ab^2-a(bB-aC))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} - \frac{2d\sqrt{a+b\tan(e+fx)}(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \frac{\frac{1}{2}(a-ib)(c-id)(A+iB-C)(bc-ad)^2 \int \frac{1}{\sqrt{a+b\tan(e+fx)}}}{(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 4098 \\
 & \frac{2(Ab^2-a(bB-aC))}{f(a^2+b^2)(bc-ad)\sqrt{a+b\tan(e+fx)}\sqrt{c+d\tan(e+fx)}} - \frac{2d\sqrt{a+b\tan(e+fx)}(a^2Ad^2+a^2(-Bcd+2c^2C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d\tan(e+fx)}} - \frac{\frac{(a+ib)(c+id)(A-iB-C)(bc-ad)^2 \int \frac{1}{(1-i\tan(e+fx))}}{(1-i\tan(e+fx))}}{(a^2+b^2)(bc-ad)} \\
 & \quad \downarrow 104
 \end{aligned}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}} - \frac{(a-ib)(c-id)(A+iB-C)(bc-ad)^2 f \frac{-ia+b+}{f}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2+a^2(-Bcd+2c^2 C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2 c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{(a^2 + b^2)(bc - ad)}{f\sqrt{a+ib}\sqrt{c+id}}$$

↓ 221

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}} - \frac{i(a-ib)(c-id)(A+iB-C)(bc-ad)^2 \arctan\frac{f}{f\sqrt{a+ib}\sqrt{c+id}}}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2+a^2(-Bcd+2c^2 C+Cd^2)-abB(c^2+d^2)+Ab^2(c^2+2d^2)+b^2 c(cC-Bd))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{(a^2 + b^2)(bc - ad)}{f\sqrt{a+ib}\sqrt{c+id}}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

```
(-2*(A*b^2 - a*(b*B - a*C)))/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]) - (-((((-I)*(a + I*b)*(A - I*B - C)*(c + I*d)*(b*c - a*d)^2*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a - I*b]*Sqrt[c - I*d]*f) + (I*(a - I*b)*(A + I*B - C)*(c - I*d)*(b*c - a*d)^2*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]*f))/((b*c - a*d)*(c^2 + d^2))) + (2*d*(a^2*A*d^2 + b^2*c*(c*C - B*d) - a*b*B*(c^2 + d^2) + A*b^2*(c^2 + 2*d^2) + a^2*(2*c^2*C - B*c*d + C*d^2))*Sqrt[a + b*Tan[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d))
```

Definitions of rubi rules used

- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`
- rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan(fx + e) + C \tan^2(fx + e)}{(a + b \tan(fx + e))^{\frac{3}{2}} (c + d \tan(fx + e))^{\frac{3}{2}}} dx$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e)
)^(3/2),x)
```

output

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e)
)^(3/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(
f*x+e))^(3/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{3}{2}} (c + d \tan(e + fx))^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e)**(3/2)/(c+d*tan(f*x+e)**(3/2)),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**(3/2)*(c + d*tan(e + f*x))**(3/2)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx = \int \frac{C \tan(fx + e)^2 + B \tan(fx + e) + A}{(b \tan(fx + e) + a)^{\frac{3}{2}} (d \tan(fx + e) + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/((b*tan(f*x + e) + a)^(3/2)*(d*tan(f*x + e) + c)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(3/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b*c*d
- 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*c*d - 2*sqrt(tan(
e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b*c**2 + int((sqrt(tan(e + f*x)*d
+ c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**2*d**2 +
2*tan(e + f*x)**3*a*b*d**2 + 2*tan(e + f*x)**3*b**2*c*d + tan(e + f*x)**2*
a**2*d**2 + 4*tan(e + f*x)**2*a*b*c*d + tan(e + f*x)**2*b**2*c**2 + 2*tan(
e + f*x)*a**2*c*d + 2*tan(e + f*x)*a*b*c**2 + a**2*c**2),x)*tan(e + f*x)**
2*a**2*b**2*d**3*f - 2*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b +
a)*tan(e + f*x))/(tan(e + f*x)**4*b**2*d**2 + 2*tan(e + f*x)**3*a*b*d**2
+ 2*tan(e + f*x)**3*b**2*c*d + tan(e + f*x)**2*a**2*d**2 + 4*tan(e + f*x)*
2*a*b*c*d + tan(e + f*x)**2*b**2*c**2 + 2*tan(e + f*x)*a**2*c*d + 2*tan(e
+ f*x)*a*b*c**2 + a**2*c**2),x)*tan(e + f*x)**2*a*b**3*c*d**2*f + int((sq
rt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)
)**4*b**2*d**2 + 2*tan(e + f*x)**3*a*b*d**2 + 2*tan(e + f*x)**3*b**2*c*d +
tan(e + f*x)**2*a**2*d**2 + 4*tan(e + f*x)**2*a*b*c*d + tan(e + f*x)**2*b
**2*c**2 + 2*tan(e + f*x)*a**2*c*d + 2*tan(e + f*x)*a*b*c**2 + a**2*c**2),
x)*tan(e + f*x)**2*b**4*c**2*d*f + int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(
e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b**2*d**2 + 2*tan(e + f*x)*
3*a*b*d**2 + 2*tan(e + f*x)**3*b**2*c*d + tan(e + f*x)**2*a**2*d**2 + 4*t
an(e + f*x)**2*a*b*c*d + tan(e + f*x)**2*b**2*c**2 + 2*tan(e + f*x)*a**...
```


3.158
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{5/2}(c+d \tan(e+fx))^{3/2}} dx$$

Optimal result	1756
Mathematica [A] (verified)	1757
Rubi [A] (verified)	1758
Maple [F(-1)]	1763
Fricas [F(-1)]	1764
Sympy [F]	1764
Maxima [F(-1)]	1765
Giac [F]	1765
Mupad [F(-1)]	1765
Reduce [F]	1766

Optimal result

Integrand size = 49, antiderivative size = 598

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2}(c + d \tan(e + fx))^{3/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a - ib)^{5/2}(c - id)^{3/2} f}$$

$$- \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a + ib)^{5/2}(c + id)^{3/2} f}$$

$$- \frac{2(Ab^2 - a(bB - aC))}{3(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}$$

$$- \frac{2(7a^3bBd - 4a^4Cd + b^4(3Bc - 4Ad) + ab^3(6Ac - 6cC + Bd) - a^2b^2(3Bc + 2(5A - C)d))}{3(a^2 + b^2)^2(bc - ad)^2 f \sqrt{a + b \tan(e + fx)} \sqrt{c + d \tan(e + fx)}}$$

$$- \frac{2d(8a^3bBd(c^2 + d^2) + 2ab^3(3Ac - 3cC + Bd)(c^2 + d^2) - a^4d(8c^2C - 3Bcd + (3A + 5C)d^2) - a^2b^2(3Bc + 2(5A - C)d))}{3(a^2 + b^2)^2(bc - ad)^2}$$

output

```

-(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c
+d*tan(f*x+e))^(1/2))/(a-I*b)^(5/2)/(c-I*d)^(3/2)/f-(B-I*(A-C))*arctanh((c
+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(
a+I*b)^(5/2)/(c+I*d)^(3/2)/f-2/3*(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/
f/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(1/2)-2/3*(7*a^3*b*B*d-4*a^4*C*d
+b^4*(-4*A*d+3*B*c)+a*b^3*(6*A*c+B*d-6*C*c)-a^2*b^2*(3*B*c+2*(5*A-C)*d))/(
a^2+b^2)^2/(-a*d+b*c)^2/f/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(1/2)-2/
3*d*(8*a^3*b*B*d*(c^2+d^2)+2*a*b^3*(3*A*c+B*d-3*C*c)*(c^2+d^2)-a^4*d*(8*c^
2*C-3*B*c*d+(3*A+5*C)*d^2)-a^2*b^2*(11*A*c^2*d+17*A*d^3+3*B*c^3-3*B*c*d^2+
5*C*c^2*d-C*d^3)-b^4*(d*(5*A*c^2+8*A*d^2+3*C*c^2)-3*B*(c^3+2*c*d^2)))*(a+b
*tan(f*x+e))^(1/2)/(a^2+b^2)^2/(-a*d+b*c)^3/(c^2+d^2)/f/(c+d*tan(f*x+e))^(
1/2)
    
```

Mathematica [A] (verified)

Time = 6.70 (sec) , antiderivative size = 902, normalized size of antiderivative = 1.51

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \frac{2(Ab^2 - a(bB - aC))}{3(a^2 + b^2)(bc - ad)f(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}}$$

$$2 \left(\frac{2(-a(-2a(Ab^2 - a(bB - aC))d + \frac{3}{2}b(Ab - aB - bC)(bc - ad) + \frac{1}{2}b^2(4Ab^2d - 3aA(bc - ad) - (bB - aC)(3bc + ad)))}{(a^2 + b^2)(bc - ad)f\sqrt{a + b \tan(e + fx)}\sqrt{c + d \tan(e + fx)}} \right) - \frac{3(bc - ad)^3 \left(\frac{(a + ib)}{2} \right)}{2}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(5
/2)*(c + d*Tan[e + f*x])^(3/2)),x]
    
```

output

```

(-2*(A*b^2 - a*(b*B - a*C)))/(3*(a^2 + b^2)*(b*c - a*d)*f*(a + b*Tan[e + f*x])^(3/2)*Sqrt[c + d*Tan[e + f*x]]) - (2*((-2*(-a*(-2*a*(A*b^2 - a*(b*B - a*C)))*d + (3*b*(A*b - a*B - b*C)*(b*c - a*d))/2)) + (b^2*(4*A*b^2*d - 3*a*A*(b*c - a*d) - (b*B - a*C)*(3*b*c + a*d)))/2)/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]) - (2*((-3*(b*c - a*d)^3*((a + I*b)^2*(I*A + B - I*C)*(c + I*d)*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + I*b]*Sqrt[-c + I*d]) + ((a - I*b)^2*(A + I*B - C)*(I*c + d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d])))/(4*(-(b*c) + a*d)*(c^2 + d^2)*f) - (2*(d^2*((-1/2*(b*c) - (a*d)/2)*(-2*a*(A*b^2 - a*(b*B - a*C))*d + (3*b*(A*b - a*B - b*C)*(b*c - a*d))/2) + ((b^2*d - (a*(b*c - a*d))/2)*(4*A*b^2*d - 3*a*A*(b*c - a*d) - (b*B - a*C)*(3*b*c + a*d)))/2) - c*((d*(b*c - a*d)*(-2*b*(A*b^2 - a*(b*B - a*C))*d - (3*a*(A*b - a*B - b*C)*(b*c - a*d))/2 + (b*(4*A*b^2*d - 3*a*A*(b*c - a*d) - (b*B - a*C)*(3*b*c + a*d)))/2))/2 - c*d*(-(a*(-2*a*(A*b^2 - a*(b*B - a*C))*d + (3*b*(A*b - a*B - b*C)*(b*c - a*d))/2)) + (b^2*(4*A*b^2*d - 3*a*A*(b*c - a*d) - (b*B - a*C)*(3*b*c + a*d)))/2)))*Sqrt[a + b*Tan[e + f*x]]/((- (b*c) + a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])/((a^2 + b^2)*(b*c - a*d)))/(3*(a^2 + b^2)*(b*c - a*d))

```

Rubi [A] (verified)

Time = 7.36 (sec) , antiderivative size = 704, normalized size of antiderivative = 1.18, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx$$

↓ 4132

$$\frac{2 \int \frac{4Adb^2+4(Ab^2-a(bB-aC))d \tan^2(e+fx)-3aA(bc-ad)-(bB-aC)(3bc+ad)+3(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{2(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(a^2+b^2)(bc-ad)}{2(Ab^2-a(bB-aC))}} \downarrow 27$$

$$\frac{\int \frac{4Adb^2+4(Ab^2-a(bB-aC))d \tan^2(e+fx)-3aA(bc-ad)-(bB-aC)(3bc+ad)+3(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(a^2+b^2)(bc-ad)}{2(Ab^2-a(bB-aC))}} \downarrow 3042$$

$$\frac{\int \frac{4Adb^2+4(Ab^2-a(bB-aC))d \tan(e+fx)^2-3aA(bc-ad)-(bB-aC)(3bc+ad)+3(Ab-Cb-aB)(bc-ad) \tan(e+fx)}{(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(a^2+b^2)(bc-ad)}{2(Ab^2-a(bB-aC))}} \downarrow 4132$$

$$\frac{2(-4a^4Cd+7a^3bBd-a^2b^2(10Ad+3Bc-2Cd)+ab^3(6Ac+Bd-6cC)+b^4(3Bc-4Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \downarrow 27$$

$$\frac{2(-4a^4Cd+7a^3bBd-a^2b^2(10Ad+3Bc-2Cd)+ab^3(6Ac+Bd-6cC)+b^4(3Bc-4Ad))}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \downarrow 3042$$

↓ 3042

$$\frac{2(Ab^2 - a(bB - aC))}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{2(-4a^4Cd+7a^3bBd-a^2b^2(10Ad+3Bc-2Cd)+ab^3(6Ac+Bd-6cC)+b^4(3Bc-4Ad))} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}$$

↓ 4098

$$\frac{2(Ab^2 - a(bB - aC))}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{2(-4a^4Cd+7a^3bBd-a^2b^2(10Ad+3Bc-2Cd)+ab^3(6Ac+Bd-6cC)+b^4(3Bc-4Ad))} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}$$

↓ 104

$$\frac{2(Ab^2 - a(bB - aC))}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{2(-4a^4Cd+7a^3bBd-a^2b^2(10Ad+3Bc-2Cd)+ab^3(6Ac+Bd-6cC)+b^4(3Bc-4Ad))} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}$$

↓ 221

$$\frac{2(Ab^2 - a(bB - aC))}{3f(a^2 + b^2)(bc - ad)(a + b \tan(e + fx))^{3/2} \sqrt{c + d \tan(e + fx)}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{2(-4a^4Cd+7a^3bBd-a^2b^2(10Ad+3Bc-2Cd)+ab^3(6Ac+Bd-6cC)+b^4(3Bc-4Ad))} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^4(-d)(d^2(3A+5C)-3Bcd+8c^2C)+8}{f(a^2+b^2)(bc-ad)\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}$$

input

```
Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(5/2)*(c + d*Tan[e + f*x])^(3/2)),x]
```

output

$$\begin{aligned} & (-2*(A*b^2 - a*(b*B - a*C)))/(3*(a^2 + b^2)*(b*c - a*d)*f*(a + b*\tan[e + f*x])^{3/2}*\sqrt{c + d*\tan[e + f*x]}) - ((2*(7*a^3*b*B*d - 4*a^4*C*d + b^4*(3*B*c - 4*A*d) + a*b^3*(6*A*c - 6*c*C + B*d) - a^2*b^2*(3*B*c + 10*A*d - 2*C*d)))/((a^2 + b^2)*(b*c - a*d)*f*\sqrt{a + b*\tan[e + f*x]})*\sqrt{c + d*\tan[e + f*x]}) - ((3*((-I)*(a + I*b)^2*(A - I*B - C)*(c + I*d)*(b*c - a*d)^3*\operatorname{ArcTanh}[(\sqrt{c - I*d})*\sqrt{a + b*\tan[e + f*x]})]/(\sqrt{a - I*b})*\sqrt{c + d*\tan[e + f*x]})/(\sqrt{a - I*b})*\sqrt{c - I*d}*f) + (I*(a - I*b)^2*(A + I*B - C)*(c - I*d)*(b*c - a*d)^3*\operatorname{ArcTanh}[(\sqrt{c + I*d})*\sqrt{a + b*\tan[e + f*x]})]/(\sqrt{a + I*b})*\sqrt{c + d*\tan[e + f*x]})/(\sqrt{a + I*b})*\sqrt{c + I*d}*f)))/((b*c - a*d)*(c^2 + d^2)) - (2*d*(8*a^3*b*B*d*(c^2 + d^2) + 2*a*b^3*(3*A*c - 3*c*C + B*d)*(c^2 + d^2) - a^4*d*(8*c^2*C - 3*B*c*d + (3*A + 5*C)*d^2) - a^2*b^2*(3*B*c^3 + 11*A*c^2*d + 5*c^2*C*d - 3*B*c*d^2 + 17*A*d^3 - C*d^3) - b^4*(d*(5*A*c^2 + 3*c^2*C + 8*A*d^2) - 3*B*(c^3 + 2*c*d^2)))*\sqrt{a + b*\tan[e + f*x]})/((b*c - a*d)*(c^2 + d^2)*f*\sqrt{c + d*\tan[e + f*x]})/((a^2 + b^2)*(b*c - a*d)))/(3*(a^2 + b^2)*(b*c - a*d)) \end{aligned}$$
Definitions of rubi rules used

rule 27

$$\operatorname{Int}[(a_*)*(F_x_), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[F_x, (b_)*(G_x_)] /; \operatorname{FreeQ}[b, x]$$

rule 104

$$\operatorname{Int}[(((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)})/((e_.) + (f_.)*(x_)), x_] \rightarrow \operatorname{With}\{q = \operatorname{Denominator}[m]\}, \operatorname{Simp}[q \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$$

rule 221

$$\operatorname{Int}[((a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x\} \ \&\& \ \operatorname{NegQ}[a/b]$$

rule 3042

$$\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4098

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

rule 4099

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan (fx + e) + C \tan (fx + e)^2}{(a + b \tan (fx + e))^{\frac{5}{2}} (c + d \tan (fx + e))^{\frac{3}{2}}} dx$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e)
)^(3/2),x)
```


output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(3/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(5/2)/(c+d*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**(5/2)*(c + d*tan(e + f*x))**(3/2)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{(b \tan(fx + e) + a)^{5/2} (d \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/((b*tan(f*x + e) + a)^(5/2)*(d*tan(f*x + e) + c)^(3/2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \text{Hanged}$$

input `int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(5/2)*(c + d*tan(e + f*x))^(3/2)),x)`

output `\text{Hanged}`

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{5/2} (c + d \tan(e + fx))^{3/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(5/2)/(c+d*tan(f*x+e))^(3/2),x)
```

output

```
( - 16*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b
**2*c*d**2 - 24*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)*a*b*c*d**2 - 8*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(
e + f*x)*b**2*c**2*d - 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)
*a**2*c*d**2 - 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b*c*
*2*d + 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b**2*c**3 + 3*i
nt((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e
+ f*x)**5*b**3*d**2 + 3*tan(e + f*x)**4*a*b**2*d**2 + 2*tan(e + f*x)**4*b
**3*c*d + 3*tan(e + f*x)**3*a**2*b*d**2 + 6*tan(e + f*x)**3*a*b**2*c*d + t
an(e + f*x)**3*b**3*c**2 + tan(e + f*x)**2*a**3*d**2 + 6*tan(e + f*x)**2*a
**2*b*c*d + 3*tan(e + f*x)**2*a*b**2*c**2 + 2*tan(e + f*x)*a**3*c*d + 3*ta
n(e + f*x)*a**2*b*c**2 + a**3*c**2),x)*tan(e + f*x)**3*a**3*b**3*d**4*f -
9*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(ta
n(e + f*x)**5*b**3*d**2 + 3*tan(e + f*x)**4*a*b**2*d**2 + 2*tan(e + f*x)**
4*b**3*c*d + 3*tan(e + f*x)**3*a**2*b*d**2 + 6*tan(e + f*x)**3*a*b**2*c*d
+ tan(e + f*x)**3*b**3*c**2 + tan(e + f*x)**2*a**3*d**2 + 6*tan(e + f*x)**
2*a**2*b*c*d + 3*tan(e + f*x)**2*a*b**2*c**2 + 2*tan(e + f*x)*a**3*c*d + 3
*tan(e + f*x)*a**2*b*c**2 + a**3*c**2),x)*tan(e + f*x)**3*a**2*b**4*c*d**3
*f + 9*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)
)/(tan(e + f*x)**5*b**3*d**2 + 3*tan(e + f*x)**4*a*b**2*d**2 + 2*tan(e ...
```

3.159
$$\int \frac{(a+b \tan(e+fx))^{5/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1767
Mathematica [B] (verified)	1768
Rubi [A] (verified)	1769
Maple [F(-1)]	1774
Fricas [F(-1)]	1775
Sympy [F]	1775
Maxima [F(-1)]	1775
Giac [F]	1776
Mupad [F(-1)]	1776
Reduce [F]	1776

Optimal result

Integrand size = 49, antiderivative size = 549

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$-\frac{(a - ib)^{5/2} (iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c - id)^{5/2} f}$$

$$-\frac{(a + ib)^{5/2} (B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c + id)^{5/2} f}$$

$$-\frac{b^{3/2} (5bcC - 2bBd - 5aCd) \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{d^{7/2} f}$$

$$-\frac{2(c^2C - Bcd + Ad^2) (a + b \tan(e + fx))^{5/2}}{3d(c^2 + d^2) f(c + d \tan(e + fx))^{3/2}}$$

$$-\frac{2(b(5c^4C - 2Bc^3d - c^2(A - 11C)d^2 - 8Bcd^3 + 5Ad^4) + 3ad^2(2c(A - C)d - B(c^2 - d^2))) (a + b \tan(e + fx))^{5/2}}{3d^2(c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

$$+\frac{b(b(5c^4C - 2Bc^3d + 10c^2Cd^2 - 6Bcd^3 + (4A + C)d^4) + 2ad^2(2c(A - C)d - B(c^2 - d^2))) \sqrt{a + b \tan(e + fx)}}{d^3(c^2 + d^2)^2 f}$$

output

```

-(a-I*b)^(5/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a
-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(5/2)/f-(a+I*b)^(5/2)*(B-I*(A
-C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*
x+e))^(1/2))/(c+I*d)^(5/2)/f-b^(3/2)*(-2*B*b*d-5*C*a*d+5*C*b*c)*arctanh(d^(
1/2)*(a+b*tan(f*x+e))^(1/2)/b^(1/2)/(c+d*tan(f*x+e))^(1/2))/d^(7/2)/f-2/3
*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(5/2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))
^(3/2)-2/3*(b*(5*c^4*C-2*B*c^3*d-c^2*(A-11*C)*d^2-8*B*c*d^3+5*A*d^4)+3*a*d
^2*(2*c*(A-C)*d-B*(c^2-d^2)))*(a+b*tan(f*x+e))^(3/2)/d^2/(c^2+d^2)^2/f/(c+
d*tan(f*x+e))^(1/2)+b*(b*(5*c^4*C-2*B*c^3*d+10*C*c^2*d^2-6*B*c*d^3+(4*A+C)
*d^4)+2*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*(a+b*tan(f*x+e))^(1/2)*(c+d*tan(f
*x+e))^(1/2)/d^3/(c^2+d^2)^2/f

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2650 vs. $2(549) = 1098$.

Time = 8.36 (sec) , antiderivative size = 2650, normalized size of antiderivative = 4.83

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Result too large to show}$$

input

```

Integrate[((a + b*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]
^2))/(c + d*Tan[e + f*x])^(5/2),x]

```

output

```
(C*(a + b*Tan[e + f*x])^(5/2))/(d*f*(c + d*Tan[e + f*x])^(3/2)) + ((2*(I*a
+ b)*(A - I*B - C)*d*(a + b*Tan[e + f*x])^(3/2))/(3*(c - I*d)*f*(c + d*Ta
n[e + f*x])^(3/2)) - (2*(I*a - b)*(A + I*B - C)*d*(a + b*Tan[e + f*x])^(3/
2))/(3*(c + I*d)*f*(c + d*Tan[e + f*x])^(3/2)) + (2*(a - I*b)^2*(I*A + B -
I*C)*d*((Sqrt[-a + I*b]*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])]
/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])))/(-c + I*d)^(3/2) + Sqrt[a + b
*Tan[e + f*x]]/((c - I*d)*Sqrt[c + d*Tan[e + f*x]]))/((c - I*d)*f) + (2*(
a + I*b)^2*(I*A - B - I*C)*d*((Sqrt[a + I*b]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a
+ b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])]/(c + I*d)^(
3/2) - Sqrt[a + b*Tan[e + f*x]]/((c + I*d)*Sqrt[c + d*Tan[e + f*x]]))/((c
+ I*d)*f) + (10*c*C*(b*c - a*d)*(b/((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c -
a*d)))^(5/2)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))^3*Sqrt[(b*(c + d*
Tan[e + f*x]))/(b*c - a*d)]*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*
((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))))^2*((b^2*d^2*(a + b*Tan[e + f
*x])^2)/(3*(b*c - a*d)^2*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))^2*(-1
- (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/
(b*c - a*d))))^2 - (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c
- a*d) - (a*b*d)/(b*c - a*d))*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c - a*d
)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))) - (Sqrt[b]*Sqrt[d]*ArcSin
h[(Sqrt[b]*Sqrt[d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2...
```

Rubi [A] (verified)

Time = 8.15 (sec) , antiderivative size = 605, normalized size of antiderivative = 1.10, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4130, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$2 \int \frac{(a+b \tan(e+fx))^{3/2} (b(5C^2-2Bdc+(2A+3C)d^2) \tan^2(e+fx)+3d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(3ac+5bd)+(5bc-3ad)(cC-2Ad))}{2(c+d \tan(e+fx))^{3/2}} \\ \frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{5/2}} \\ \frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{(a+b \tan(e+fx))^{3/2} (b(5C^2-2Bdc+(2A+3C)d^2) \tan^2(e+fx)+3d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(3ac+5bd)+(5bc-3ad)(cC-2Ad))}{(c+d \tan(e+fx))^{3/2}} \\ \frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{5/2}} \\ \frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^{3/2} (b(5C^2-2Bdc+(2A+3C)d^2) \tan^2(e+fx)+3d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(3ac+5bd)+(5bc-3ad)(cC-2Ad))}{(c+d \tan(e+fx))^{3/2}} \\ \frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{5/2}} \\ \frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 4128

$$2 \int \frac{\sqrt{a+b \tan(e+fx)} (3((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+3bd)(Ad(3ac+5bd)+(5bc-3ad)(cC-2Ad))}{(c+d \tan(e+fx))^{3/2}}$$

$$\frac{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{5/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{\sqrt{a+b \tan(e+fx)} (3((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+3bd)(Ad(3ac+5bd)+(5bc-3ad)(cC-2Ad))}{(c+d \tan(e+fx))^{3/2}}$$

$$\frac{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{5/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{\sqrt{a+b \tan(e+fx)} \left(3((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+3bd)(Ad(3ac+5bd)+(5bc-3ad)(cC-Ad)) \right)}{\dots}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 4130

$$\int \frac{3 \left(-c(5Cc^4 - 2Bdc^3 + 10Cd^2c^2 - 6Bd^3c + (4A+C)d^4) b^3 - (5bcC - 5adC - 2bBd)(c^2 + d^2)^2 \tan^2(e+fx)b^2 + ad(5Cc^4 - 2(3A-8C)d^2c^2 - 12Bd^3c + (6A-C)d^4) b^2 + 6 \right)}{\dots}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{-c(5Cc^4 - 2Bdc^3 + 10Cd^2c^2 - 6Bd^3c + (4A+C)d^4) b^3 - (5bcC - 5adC - 2bBd)(c^2 + d^2)^2 \tan^2(e+fx)b^2 + ad(5Cc^4 - 2(3A-8C)d^2c^2 - 12Bd^3c + (6A-C)d^4) b^2 + 6}{\dots}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{-c(5Cc^4 - 2Bdc^3 + 10Cd^2c^2 - 6Bd^3c + (4A+C)d^4) b^3 - (5bcC - 5adC - 2bBd)(c^2 + d^2)^2 \tan^2(e+fx)b^2 + ad(5Cc^4 - 2(3A-8C)d^2c^2 - 12Bd^3c + (6A-C)d^4) b^2 + 6}{\dots}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 4138

$$\int \frac{-c(5Cc^4 - 2Bdc^3 + 10Cd^2c^2 - 6Bd^3c + (4A+C)d^4) b^3 - (5bcC - 5adC - 2bBd)(c^2 + d^2)^2 \tan^2(e+fx)b^2 + ad(5Cc^4 - 2(3A-8C)d^2c^2 - 12Bd^3c + (6A-C)d^4) b^2 + 6}{\dots}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 2348

$$\frac{3b\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(2a(2c(A-C)d-B(c^2-d^2))d^2+b(5Cc^4-2Bdc^3+10Cd^2c^2-6Bd^3c+(4A+C)d^4))}{df} + \frac{3 \int \left(-\frac{b^2(5bcC-5adC-2bBd)(c^2+d^2)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} \right)}{df}$$

$$\frac{2(Cc^2 - Bdc + Ad^2) (a + b \tan(e + fx))^{5/2}}{3d(c^2 + d^2) f(c + d \tan(e + fx))^{3/2}}$$

↓ 2009

$$-\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{5/2}}{3df(c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$-\frac{2(a+b \tan(e+fx))^{3/2}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-11C)+5Ad^4-2Bc^3d-8Bcd^3+5c^4C))}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{3b\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}{df}$$

input

```
Int[((a + b*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2),x]
```

output

```
(-2*(c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^(5/2))/(3*d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) + ((-2*(b*(5*c^4*C - 2*B*c^3*d - c^2*(A - 11*C))*d^2 - 8*B*c*d^3 + 5*A*d^4) + 3*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*(a + b*Tan[e + f*x])^(3/2)/(d*(c^2 + d^2)*f*sqrt[c + d*Tan[e + f*x]]) + ((3*((-2*(a - I*b)^(5/2)*(I*A + B - I*C))*(c + I*d)^2*d^3*ArcTanh[(sqrt[c - I*d]*sqrt[a + b*Tan[e + f*x]])/(sqrt[a - I*b]*sqrt[c + d*Tan[e + f*x]])])/sqrt[c - I*d] - (2*(a + I*b)^(5/2)*(B - I*(A - C))*(c - I*d)^2*d^3*ArcTanh[(sqrt[c + I*d]*sqrt[a + b*Tan[e + f*x]])/(sqrt[a + I*b]*sqrt[c + d*Tan[e + f*x]])])/sqrt[c + I*d] - (2*b^(3/2)*(5*b*c*C - 2*b*B*d - 5*a*C*d)*(c^2 + d^2)^2*ArcTanh[(sqrt[d]*sqrt[a + b*Tan[e + f*x]])/(sqrt[b]*sqrt[c + d*Tan[e + f*x]])])/sqrt[d]))/(2*d*f) + (3*b*(b*(5*c^4*C - 2*B*c^3*d + 10*c^2*C*d^2 - 6*B*c*d^3 + (4*A + C)*d^4) + 2*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*sqrt[a + b*Tan[e + f*x]]*sqrt[c + d*Tan[e + f*x]])/(d*f))/(d*(c^2 + d^2))/(3*d*(c^2 + d^2))
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 2348 $\text{Int}[(Px_)*((c_) + (d_)*(x_))^{(m_)*((e_) + (f_)*(x_))^{(n_)*((a_) + (b_)*(x_)^2)^{(p_)}}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0])$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4128 $\text{Int}[(a_*) + (b_*)\tan[(e_*) + (f_)*(x_)]^{(m_)*((c_*) + (d_*)\tan[(e_*) + (f_)*(x_)]^{(n_)*((A_*) + (B_*)\tan[(e_*) + (f_)*(x_)] + (C_*)\tan[(e_*) + (f_)*(x_)]^2)}, x_Symbol] \rightarrow \text{Simp}[(A*d^2 + c*(c*C - B*d))*(a + b*\text{Tan}[e + f*x])^m*((c + d*\text{Tan}[e + f*x])^{(n + 1)}/(d*f*(n + 1)*(c^2 + d^2))), x] - \text{Simp}[1/(d*(n + 1)*(c^2 + d^2)) \ \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m - 1)}*(c + d*\text{Tan}[e + f*x])^{(n + 1)}*\text{Simp}[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*\text{Tan}[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[n, -1]$

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(fx + e))^{\frac{5}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(c + d \tan(fx + e))^{\frac{5}{2}}} dx$$

input

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

output

```
int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c + d \tan(e + fx))^{5/2}}$$

input

```
integrate((a+b*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(5/2),x)
```

output

```
Integral((a + b*tan(e + f*x))**(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(5/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^{5/2}}{(d \tan(fx + e) + c)^{5/2}}$$

input `integrate((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(5/2)/(d*tan(f*x + e) + c)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))^{5/2} (C \tan(e + fx) + B + A)}{(c + d \tan(e + fx))^{5/2}}$$

input `int(((a + b*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2),x)`

output `int(((a + b*tan(e + f*x))^(5/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2), x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^{5/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(\tan(fx + e) b + a)^{5/2} (A + B \tan(fx + e) + C \tan^2(fx + e))}{(d \tan(fx + e) + c)^{5/2}}$$

input `int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)`

output `int((a+b*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)`

3.160
$$\int \frac{(a+b \tan(e+fx))^{3/2} (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1778
Mathematica [B] (verified)	1779
Rubi [A] (verified)	1780
Maple [F(-1)]	1784
Fricas [F(-1)]	1785
Sympy [F]	1785
Maxima [F(-1)]	1785
Giac [F]	1786
Mupad [F(-1)]	1786
Reduce [F]	1786

Optimal result

Integrand size = 49, antiderivative size = 407

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(a - ib)^{3/2} (iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c - id)^{5/2} f}$$

$$- \frac{(a + ib)^{3/2} (B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c + id)^{5/2} f}$$

$$+ \frac{2b^{3/2} C \operatorname{arctanh}\left(\frac{\sqrt{d}\sqrt{a+b \tan(e+fx)}}{\sqrt{b}\sqrt{c+d \tan(e+fx)}}\right)}{d^{5/2} f} - \frac{2(c^2 C - Bcd + Ad^2) (a + b \tan(e + fx))^{3/2}}{3d (c^2 + d^2) f (c + d \tan(e + fx))^{3/2}}$$

$$- \frac{2(b(c^4 C - c^2(A - 3C)d^2 - 2Bcd^3 + Ad^4) + ad^2(2c(A - C)d - B(c^2 - d^2))) \sqrt{a + b \tan(e + fx)}}{d^2 (c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

output

$$\begin{aligned}
 & - (a - I * b)^{(3/2)} * (I * A + B - I * C) * \operatorname{arctanh}((c - I * d)^{(1/2)} * (a + b * \tan(f * x + e))^{(1/2)} / (a - I * b)^{(1/2)} / (c + d * \tan(f * x + e))^{(1/2)}) / (c - I * d)^{(5/2)} / f - (a + I * b)^{(3/2)} * (B - I * (A - C)) * \operatorname{arctanh}((c + I * d)^{(1/2)} * (a + b * \tan(f * x + e))^{(1/2)} / (a + I * b)^{(1/2)} / (c + d * \tan(f * x + e))^{(1/2)}) / (c + I * d)^{(5/2)} / f + 2 * b^{(3/2)} * C * \operatorname{arctanh}(d^{(1/2)} * (a + b * \tan(f * x + e))^{(1/2)} / b^{(1/2)} / (c + d * \tan(f * x + e))^{(1/2)}) / d^{(5/2)} / f - 2/3 * (A * d^2 - B * c * d + C * c^2) * (a + b * \tan(f * x + e))^{(3/2)} / d / (c^2 + d^2) / f / (c + d * \tan(f * x + e))^{(3/2)} - 2 * (b * (c^4 * C - c^2 * (A - 3 * C)) * d^2 - 2 * B * c * d^3 + A * d^4) + a * d^2 * (2 * c * (A - C) * d - B * (c^2 - d^2))) * (a + b * \tan(f * x + e))^{(1/2)} / d^2 / (c^2 + d^2)^2 / f / (c + d * \tan(f * x + e))^{(1/2)}
 \end{aligned}$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1135 vs. 2(407) = 814.

Time = 6.71 (sec) , antiderivative size = 1135, normalized size of antiderivative = 2.79

$$\begin{aligned}
 & \int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \frac{(B + i(A - C))(a + b \tan(e + fx))}{3(c - id)f(c + d \tan(e + fx))^{3/2}} \\
 & - \frac{(iA - B - iC)(a + b \tan(e + fx))^{3/2}}{3(c + id)f(c + d \tan(e + fx))^{3/2}} \\
 & + \frac{(ia + b)(A - iB - C) \left(\frac{\sqrt{-a + ib} \operatorname{arctanh}\left(\frac{\sqrt{-c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{-a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{(-c + id)^{3/2}} + \frac{\sqrt{a + b \tan(e + fx)}}{(c - id) \sqrt{c + d \tan(e + fx)}} \right)}{(c - id)f} \\
 & + \frac{(ia - b)(A + iB - C) \left(\frac{\sqrt{a + ib} \operatorname{arctanh}\left(\frac{\sqrt{c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{(c + id)^{3/2}} - \frac{\sqrt{a + b \tan(e + fx)}}{(c + id) \sqrt{c + d \tan(e + fx)}} \right)}{(c + id)f} \\
 & - \frac{2C(bc - ad) \left(\frac{b}{\frac{b^2c}{bc - ad} - \frac{abd}{bc - ad}} \right)^{5/2} \left(\frac{b^2c}{bc - ad} - \frac{abd}{bc - ad} \right)^3 \sqrt{\frac{b(c + d \tan(e + fx))}{bc - ad}} \left(-1 - \frac{bd(a + b \tan(e + fx))}{(bc - ad) \left(\frac{b^2c}{bc - ad} - \frac{abd}{bc - ad} \right)} \right)^2}{3(bc - ad)^2} \\
 & \qquad \qquad \qquad b^2 d^3 f \sqrt{a + b \tan(e + fx)}
 \end{aligned}$$

input

```
Integrate[((a + b*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2),x]
```


output

```

((B + I*(A - C))*(a + b*Tan[e + f*x])^(3/2))/(3*(c - I*d)*f*(c + d*Tan[e +
f*x])^(3/2)) - ((I*A - B - I*C)*(a + b*Tan[e + f*x])^(3/2))/(3*(c + I*d)*
f*(c + d*Tan[e + f*x])^(3/2)) + ((I*a + b)*(A - I*B - C)*((Sqrt[-a + I*b]*
ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c +
d*Tan[e + f*x]])])/(-c + I*d)^(3/2) + Sqrt[a + b*Tan[e + f*x]]/((c - I*d)
*Sqrt[c + d*Tan[e + f*x]])))/((c - I*d)*f) + ((I*a - b)*(A + I*B - C)*((Sqr
rt[a + I*b]*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b
]*Sqrt[c + d*Tan[e + f*x]])])/((c + I*d)^(3/2) - Sqrt[a + b*Tan[e + f*x]]/(
(c + I*d)*Sqrt[c + d*Tan[e + f*x]])))/((c + I*d)*f) - (2*C*(b*c - a*d)*(b/
((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)))^(5/2)*((b^2*c)/(b*c - a*d) -
(a*b*d)/(b*c - a*d))^3*Sqrt[(b*(c + d*Tan[e + f*x]))/(b*c - a*d)]*(-1 - (b
*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c
- a*d))))^2*((b^2*d^2*(a + b*Tan[e + f*x])^2)/(3*(b*c - a*d)^2*((b^2*c)/(b
*c - a*d) - (a*b*d)/(b*c - a*d)))^2*(-1 - (b*d*(a + b*Tan[e + f*x]))/((b*c
- a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))))^2) - (b*d*(a + b*Tan[
e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d))*(-1 -
(b*d*(a + b*Tan[e + f*x]))/((b*c - a*d)*((b^2*c)/(b*c - a*d) - (a*b*d)/(b*c
- a*d)))))) - (Sqrt[b]*Sqrt[d]*ArcSinh[(Sqrt[b]*Sqrt[d]*Sqrt[a + b*Tan[e
+ f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - (a*b*d)/(b*c - a*d)]]
]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[b*c - a*d]*Sqrt[(b^2*c)/(b*c - a*d) - ...

```

Rubi [A] (verified)

Time = 4.69 (sec) , antiderivative size = 462, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.204$, Rules used = {3042, 4128, 27, 3042, 4128, 27, 3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$2 \int \frac{3\sqrt{a+b \tan(e+fx)}(bC(c^2+d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+bd)+(bc-ad)(cC-Bd))}{2(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{\sqrt{a+b \tan(e+fx)}(bC(c^2+d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+bd)+(bc-ad)(cC-Bd))}{(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{\sqrt{a+b \tan(e+fx)}(bC(c^2+d^2) \tan^2(e+fx)+d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(ac+bd)+(bc-ad)(cC-Bd))}{(c+d \tan(e+fx))^{3/2}} dx$$

$$\frac{d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 4128

$$2 \int \frac{((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+bd)(Ad(ac+bd)+(bc-ad)(cC-Bd))d+b^2C(c^2+d^2)^2 \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 27

$$\int \frac{((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+bd)(Ad(ac+bd)+(bc-ad)(cC-Bd))d+b^2C(c^2+d^2)^2 \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx$$

$$\frac{d(c^2+d^2)}{2(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{3/2}}$$

$$\frac{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 3042

$$\int \frac{((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+bd)(Ad(ac+bd)+(bc-ad)(cC-Bd))d+b^2C(c^2+d^2)^2 \tan(e+fx)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)} d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

4138

$$\int \frac{((ac+bd)((A-C)(bc-ad)+B(ac+bd))-(bc-ad)(bBc-b(A-C)d-a(Ac-Cc+Bd))) \tan(e+fx)d^2+(ac+bd)(Ad(ac+bd)+(bc-ad)(cC-Bd))d+b^2C(c^2+d^2)^2 \tan(e+fx)}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)} (d \tan(e+fx) + c)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

2348

$$\int \left(\frac{b^2C(c^2+d^2)^2}{\sqrt{a+b \tan(e+fx)} \sqrt{c+d \tan(e+fx)}} + \frac{2aAbd^4+a^2Bd^4-b^2Bd^4-2abCd^4-2Ab^2cd^3+2a^2Acd^3-4abBcd^3-2a^2cCd^3+2b^2cCd^3-2aAbc^2d^2-a^2Bc^2d^2+b^2Bc^2d^2}{2(i-\tan(e+fx))} \right)$$

$$\frac{2(Cc^2 - Bdc + Ad^2) (a + b \tan(e + fx))^{3/2}}{3d (c^2 + d^2) f (c + d \tan(e + fx))^{3/2}}$$

2009

$$\frac{2(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{3/2}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$-\frac{2\sqrt{a+b \tan(e+fx)}(ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(A-3C)+Ad^4-2Bcd^3+c^4C))}{df(c^2+d^2)\sqrt{c+d \tan(e+fx)}} + \frac{d^2(a-ib)^{3/2}(c+id)^2(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c-id}}{\sqrt{a-id}}\right)}{\sqrt{c-id}}$$

$d(c^2 + d^2)$

input

```
Int[((a + b*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2),x]
```

output

$$\begin{aligned} & (-2*(c^2*C - B*c*d + A*d^2)*(a + b*\text{Tan}[e + f*x])^{(3/2)})/(3*d*(c^2 + d^2)*f \\ & *(c + d*\text{Tan}[e + f*x])^{(3/2)}) + (((-(a - I*b)^{(3/2)}*(I*A + B - I*C)*(c + I \\ & *d)^2*d^2*\text{ArcTanh}[\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[a - I*b]* \\ & \text{Sqrt}[c + d*\text{Tan}[e + f*x]])))/\text{Sqrt}[c - I*d] - ((a + I*b)^{(3/2)}*(B - I*(A - \\ & C))*(c - I*d)^2*d^2*\text{ArcTanh}[\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt} \\ & [a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])))/\text{Sqrt}[c + I*d] + (2*b^{(3/2)}*C*(c^2 + \\ & d^2)^2*\text{ArcTanh}[\text{Sqrt}[d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(\text{Sqrt}[b]*\text{Sqrt}[c + d*\text{Tan}[\\ & e + f*x]])))/\text{Sqrt}[d]/(d*(c^2 + d^2)*f) - (2*(b*(c^4*C - c^2*(A - 3*C)*d^2 \\ & - 2*B*c*d^3 + A*d^4) + a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*\text{Sqrt}[a + b* \\ & \text{Tan}[e + f*x]]/(d*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]))/(d*(c^2 + d^2)) \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)(F_x_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ /; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x_)] \text{ /; FreeQ}[b, x]$$

rule 2009

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 2348

$$\begin{aligned} & \text{Int}[(P_x)*((c_) + (d_)*(x_))^{(m_)}*((e_) + (f_)*(x_))^{(n_)}*((a_) + (b_ \\ &)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_x*(c + d*x)^m*(e + f*x)^ \\ & n*(a + b*x^2)^p, x], x] \text{ /; FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P \\ & x, x] \ \&\& \ (\text{IntegerQ}[p] \ || \ (\text{IntegerQ}[2*p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n, 0])) \ \&\& \\ & \ !(\text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0]) \end{aligned}$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4128

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(fx + e))^{\frac{3}{2}} (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(c + d \tan(fx + e))^{\frac{5}{2}}} dx$$

input

```
int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

output

```
int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c + d \tan(e + fx))^{5/2}}$$

input

```
integrate((a+b*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**(5/2),x)
```

output

```
Integral((a + b*tan(e + f*x))**(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**(5/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan(fx + e))^2 + B \tan(fx + e)}{(d \tan(fx + e))^{5/2}}$$

input `integrate((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^(3/2)/(d*tan(f*x + e) + c)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(a + b \tan(e + fx))^{3/2} (C \tan(e + fx) + B)}{(c + d \tan(e + fx))^{5/2}}$$

input `int(((a + b*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2),x)`

output `int(((a + b*tan(e + f*x))^(3/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2), x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^{3/2} (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input `int((a+b*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)`

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*
b*c*d + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a
*b**3*d + 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)
*a*b**2*c**2 + 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)*b**4*c - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**3*c*
d - 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**2*b**2*d + 2*sq
rt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**2*b*c**2 + 14*sqrt(tan(
e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b**3*c + 3*int((sqrt(tan(e + f*
x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3
+ 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)
**2*a**2*b*c*d**4*f - 6*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b
+ a)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3
*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)**2*a*b**2*c**2*d**3*f + 3*int
((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(
e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3
),x)*tan(e + f*x)**2*b**3*c**3*d**2*f + 6*int((sqrt(tan(e + f*x)*d + c)*sq
rt(tan(e + f*x)*b + a)*tan(e + f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e +
f*x)**2*c*d**2 + 3*tan(e + f*x)*c**2*d + c**3),x)*tan(e + f*x)*a**2*b*c**2
*d**3*f - 12*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e
+ f*x)**3)/(tan(e + f*x)**3*d**3 + 3*tan(e + f*x)**2*c*d**2 + 3*tan(e + ...
```


3.161
$$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1788
Mathematica [A] (verified)	1789
Rubi [A] (verified)	1789
Maple [F(-1)]	1794
Fricas [F(-1)]	1794
Sympy [F]	1795
Maxima [F(-2)]	1795
Giac [F]	1795
Mupad [F(-1)]	1796
Reduce [F]	1796

Optimal result

Integrand size = 49, antiderivative size = 373

$$\int \frac{\sqrt{a+b \tan(e+fx)}(A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^{5/2}} dx =$$

$$-\frac{\sqrt{a-ib}(iA+B-iC)\operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c-id)^{5/2}f}$$

$$-\frac{\sqrt{a+ib}(B-i(A-C))\operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(c+id)^{5/2}f}$$

$$-\frac{2(c^2C-Bcd+Ad^2)\sqrt{a+b \tan(e+fx)}}{3d(c^2+d^2)f(c+d \tan(e+fx))^{3/2}}$$

$$+\frac{2(b(c^4C+2Bc^3d-c^2(5A-7C)d^2-4Bcd^3+Ad^4)+3ad^2(2c(A-C)d-B(c^2-d^2)))\sqrt{a+b \tan(e+fx)}}{3d(bc-ad)(c^2+d^2)^2f\sqrt{c+d \tan(e+fx)}}$$

output

```
-(a-I*b)^(1/2)*(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c-I*d)^(5/2)/f-(a+I*b)^(1/2)*(B-I*(A-C))*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(c+I*d)^(5/2)/f-2/3*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(1/2)/d/(c^2+d^2)/f/(c+d*tan(f*x+e))^(3/2)+2/3*(b*(c^4*C+2*B*c^3*d-c^2*(5*A-7*C)*d^2-4*B*c*d^3+A*d^4)+3*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*(a+b*tan(f*x+e))^(1/2)/d/(-a*d+b*c)/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 6.07 (sec) , antiderivative size = 434, normalized size of antiderivative = 1.16

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \frac{-\frac{3C\sqrt{a+b \tan(e+fx)}}{(c+d \tan(e+fx))^{3/2}} + \frac{(c^2C+2Bcd+(-2A+3C))}{(c^2+d^2)(c+d \tan(e+fx))}}{(c+d \tan(e+fx))^{3/2}}$$

input

```
Integrate[(Sqrt[a + b*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2
))/(c + d*Tan[e + f*x])^(5/2),x]
```

output

```
((-3*C*Sqrt[a + b*Tan[e + f*x]])/(c + d*Tan[e + f*x])^(3/2) + ((c^2*C + 2*
B*c*d + (-2*A + 3*C)*d^2)*Sqrt[a + b*Tan[e + f*x]]/((c^2 + d^2)*(c + d*Ta
n[e + f*x])^(3/2)) + (-3*d*(b*c - a*d)*((Sqrt[-a + I*b]*(I*A + B - I*C)*(c
+ I*d)^2*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b
]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[-c + I*d] + (Sqrt[a + I*b]*((-I)*A + B
+ I*C)*(c - I*d)^2*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[
a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/Sqrt[c + I*d]) + (2*(b*(c^4*C + 2*B*c
^3*d + c^2*(-5*A + 7*C)*d^2 - 4*B*c*d^3 + A*d^4) + 3*a*d^2*(2*c*(A - C)*d
+ B*(-c^2 + d^2)))*Sqrt[a + b*Tan[e + f*x]]/Sqrt[c + d*Tan[e + f*x]]/((b
*c - a*d)*(c^2 + d^2)^2))/(3*d*f)
```

Rubi [A] (verified)

Time = 4.15 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.19, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$, Rules used = {3042, 4128, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx)^2)}{(c + d \tan(e + fx))^{5/2}} dx$$

↓ 4128

$$\frac{2 \int \frac{b(Cc^2+2Bdc-(2A-3C)d^2) \tan^2(e+fx)+3d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(3ac+bd)+(bc-3ad)(cC-Bd)}{2\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}} = \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 27

$$\frac{\int \frac{b(Cc^2+2Bdc-(2A-3C)d^2) \tan^2(e+fx)+3d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(3ac+bd)+(bc-3ad)(cC-Bd)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}} = \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 3042

$$\frac{\int \frac{b(Cc^2+2Bdc-(2A-3C)d^2) \tan(e+fx)^2+3d((A-C)(bc-ad)+B(ac+bd)) \tan(e+fx)+Ad(3ac+bd)+(bc-3ad)(cC-Bd)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}} = \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 4132

$$\frac{2 \int -\frac{3(d(bc-ad)(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2)))+d(bc-ad)(2aAcd-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2))}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}}}{(c^2+d^2)(bc-ad)} dx}{\frac{3d(c^2+d^2)}{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}} = \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 27

$$\frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \int \frac{d(bc-ad)(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))-b(2c(A-C)d-B(c^2-d^2)))+d(bc-ad)(2aAcd-2acCd-Ab(c^2-d^2)-aB(c^2-d^2)+b(Cc^2-2Bdc-Cd^2))}{(c^2+d^2)(bc-ad)} dx}{3d(c^2+d^2)}}{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}} = \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3df(c^2+d^2)(c+d \tan(e+fx))^{3/2}}$$

↓ 3042

$$\frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \int \frac{d(bc-ad)(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}}}{3d(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}}$$

↓ 4099

$$-\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \left(-\frac{1}{2}d(a+ib)(c-id)^2(A+iB-C)(bc-ad) \right)}{3d(c^2+d^2)}$$

↓ 3042

$$-\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \left(-\frac{1}{2}d(a+ib)(c-id)^2(A+iB-C)(bc-ad) \right)}{3d(c^2+d^2)}$$

↓ 4098

$$-\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \left(-\frac{d(a+ib)(c-id)^2(A+iB-C)(bc-ad)}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} \right)}{3d(c^2+d^2)}$$

↓ 104

$$-\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \left(-\frac{d(a+ib)(c-id)^2(A+iB-C)(bc-ad)}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} \right)}{3d(c^2+d^2)}$$

↓ 221

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3df (c^2 + d^2) (c + d \tan(e + fx))^{3/2}} + \frac{2\sqrt{a+b \tan(e+fx)}(3ad^2(2cd(A-C)-B(c^2-d^2))+b(-c^2d^2(5A-7C)+Ad^4+2Bc^3d-4Bcd^3+c^4C))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{\left(\frac{id\sqrt{a-ib}(c+id)^2(A-ib-C)(bc-ad)\arctan\left(\frac{c+id}{f\sqrt{c-d}}\right)}{f\sqrt{c-d}} \right)}{3d(c^2 + d^2)}$$

input `Int[(Sqrt[a + b*Tan[e + f*x]]*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^(5/2),x]`

output `(-2*(c^2*C - B*c*d + A*d^2)*Sqrt[a + b*Tan[e + f*x]])/(3*d*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) + ((-3*((I*Sqrt[a - I*b]*(A - I*B - C)*(c + I*d)^2*d*(b*c - a*d)*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[c - I*d]*f) - (I*Sqrt[a + I*b]*(A + I*B - C)*(c - I*d)^2*d*(b*c - a*d)*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[c + I*d]*f)))/((b*c - a*d)*(c^2 + d^2)) + (2*(b*(c^4*C + 2*B*c^3*d - c^2*(5*A - 7*C)*d^2 - 4*B*c*d^3 + A*d^4) + 3*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Sqrt[a + b*Tan[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])/(3*d*(c^2 + d^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 221 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132

```

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```

Maple [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan (fx + e)} (A + B \tan (fx + e) + C \tan (fx + e)^2)}{(c + d \tan (fx + e))^{\frac{5}{2}}} dx$$

input

```
int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

output

```
int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan (e + fx)} (A + B \tan (e + fx) + C \tan^2 (e + fx))}{(c + d \tan (e + fx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input

```
integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(
f*x+e))^(5/2),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((a+b*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)**(5/2),x)`

output `Integral(sqrt(a + b*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x)**(5/2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(((2*b*d+2*a*c)^2>0)', see `assume?` for more)`

Giac [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{(C \tan^2(fx + e) + B \tan(fx + e) + A) \sqrt{a + b \tan(fx + e)}}{(d \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*sqrt(b*tan(f*x + e) + a)/(d*tan(f*x + e) + c)^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \int \frac{\sqrt{a + b \tan(e + fx)}(C \tan(e + fx) + \dots)}{(c + d \tan(e + fx))^{5/2}}$$

input `int(((a + b*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2),x)`

output `int(((a + b*tan(e + f*x))^(1/2)*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^(5/2), x)`

Reduce [F]

$$\int \frac{\sqrt{a + b \tan(e + fx)}(A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input `int((a+b*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^(5/2),x)`

output

```
( - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*a*b*c
*d + 4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**3
*d + 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b**2
*c**2 - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a**2*c*d - 2*s
qrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b**2*d + 2*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b*c**2 + 6*sqrt(tan(e + f*x)*d + c
)*sqrt(tan(e + f*x)*b + a)*b**3*c + 6*int((sqrt(tan(e + f*x)*d + c)*sqrt(t
an(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3
*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3*tan(
e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3 + a*c
**3),x)*tan(e + f*x)**2*a**3*b*d**4*f - 12*int((sqrt(tan(e + f*x)*d + c)*s
qrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b*d**3 + tan(e + f*
x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2 + 3
*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3
+ a*c**3),x)*tan(e + f*x)**2*a**2*b**2*c*d**3*f - 3*int((sqrt(tan(e + f*x)
*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b*d**3 + t
an(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c
*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x
)*b*c**3 + a*c**3),x)*tan(e + f*x)**2*a**2*b*c*d**4*f + 6*int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b...
```

3.162
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1798
Mathematica [A] (verified)	1799
Rubi [A] (verified)	1799
Maple [F(-1)]	1804
Fricas [F(-1)]	1804
Sympy [F]	1804
Maxima [F(-1)]	1805
Giac [F]	1805
Mupad [F(-1)]	1806
Reduce [F]	1806

Optimal result

Integrand size = 49, antiderivative size = 379

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(B + i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c - id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a - ib} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a - ib}(c - id)^{5/2} f}$$

$$+ \frac{(iA - B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c + id} \sqrt{a + b \tan(e + fx)}}{\sqrt{a + ib} \sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{a + ib}(c + id)^{5/2} f}$$

$$+ \frac{2(c^2 C - Bcd + Ad^2) \sqrt{a + b \tan(e + fx)}}{3(bc - ad)(c^2 + d^2) f (c + d \tan(e + fx))^{3/2}}$$

$$+ \frac{2(b(2c^4 C - 5Bc^3 d + 4c^2(2A - C)d^2 + Bcd^3 + 2Ad^4) - 3ad^2(2c(A - C)d - B(c^2 - d^2))) \sqrt{a + b \tan(e + fx)}}{3(bc - ad)^2 (c^2 + d^2)^2 f \sqrt{c + d \tan(e + fx)}}$$

output

```
-(B+I*(A-C))*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a-I*b)^(1/2)/(c-I*d)^(5/2)/f+(I*A-B-I*C)*arctanh((c+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(a+I*b)^(1/2)/(c+I*d)^(5/2)/f+2/3*(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(1/2)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))^(3/2)+2/3*(b*(2*c^4*C-5*B*c^3*d+4*c^2*(2*A-C)*d^2+B*c*d^3+2*A*d^4)-3*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2)))*(a+b*tan(f*x+e))^(1/2)/(-a*d+b*c)^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)
```

Mathematica [A] (verified)

Time = 3.78 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.06

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx = \frac{3(bc - ad)^2 \left(\frac{(iA + B - iC)(c + id)^2 \operatorname{arctanh}\left(\frac{\sqrt{-c + id}\sqrt{a + b \tan(e + fx)}}{\sqrt{-a + ib}\sqrt{c + d \tan(e + fx)}}\right)}{\sqrt{-a + ib}\sqrt{-c + id}} \right)}{}$$

input

```
Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2)),x]
```

output

```
(3*(b*c - a*d)^2*((I*A + B - I*C)*(c + I*d)^2*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a + I*b]*Sqrt[-c + I*d]) + (I*(A + I*B - C)*(c - I*d)^2*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]) + (2*(b*c - a*d)*(c^2 + d^2)*(c^2*C - B*c*d + A*d^2)*Sqrt[a + b*Tan[e + f*x]])/(c + d*Tan[e + f*x])^(3/2) + (2*(b*(2*c^4*C - 5*B*c^3*d + 4*c^2*(2*A - C)*d^2 + B*c*d^3 + 2*A*d^4) + 3*a*d^2*(2*c*(-A + C)*d + B*(c^2 - d^2)))*Sqrt[a + b*Tan[e + f*x]])/Sqrt[c + d*Tan[e + f*x]])/(3*(b*c - a*d)^2*(c^2 + d^2)^2*f)
```

Rubi [A] (verified)

Time = 4.04 (sec) , antiderivative size = 461, normalized size of antiderivative = 1.22, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.245$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx$$

↓ 4132

$$\frac{2 \int \frac{2Abd^2+2b(Cc^2-Bdc+Ad^2) \tan^2(e+fx)+3Ac(bc-ad)-(bc-3ad)(cC-Bd)+3(bc-ad)(Bc-(A-C)d) \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(c^2+d^2)(bc-ad)}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}}}$$

↓ 27

$$\frac{\int \frac{2Abd^2+2b(Cc^2-Bdc+Ad^2) \tan^2(e+fx)+3Ac(bc-ad)-(bc-3ad)(cC-Bd)+3(bc-ad)(Bc-(A-C)d) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(c^2+d^2)(bc-ad)}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}}}$$

↓ 3042

$$\frac{\int \frac{2Abd^2+2b(Cc^2-Bdc+Ad^2) \tan(e+fx)^2+3Ac(bc-ad)-(bc-3ad)(cC-Bd)+3(bc-ad)(Bc-(A-C)d) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(c^2+d^2)(bc-ad)}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}}}$$

↓ 4132

$$\frac{2 \int -\frac{3((Cc^2-2Bdc-Cd^2-A(c^2-d^2))(bc-ad)^2+(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)(bc-ad)^2)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}} dx}{\frac{3(c^2+d^2)(bc-ad)}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}}}$$

↓ 27

$$\frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \int \frac{(Cc^2-2Bdc-Cd^2-A(c^2-d^2))(bc-ad)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx))^{3/2}} dx}{\frac{3(c^2+d^2)(bc-ad)}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2(Ad^2-Bcd+c^2C) \sqrt{a+b \tan(e+fx)}}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}}}$$

↓ 3042

$$\frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3 \int \frac{(Cc^2-2Bdc-Cd^2-A(c^2-d^2))(bc-ad)}{\sqrt{a+b \tan(e+fx)}}}{(c^2+d^2)}$$

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} + \frac{3(c^2 + d^2)(bc - ad)}{3(c^2 + d^2)(bc - ad)}$$

4099

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3\left(-\frac{1}{2}(c-id)^2(A+iB-C)(bc-ad)^2 \int \sqrt{\dots}\right)}{3(c^2 + d^2)(bc - ad)}$$

3042

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3\left(-\frac{1}{2}(c-id)^2(A+iB-C)(bc-ad)^2 \int \sqrt{\dots}\right)}{3(c^2 + d^2)(bc - ad)}$$

4098

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3\left(\frac{(c+id)^2(A-iB-C)(bc-ad)^2 \int \frac{\dots}{(1-i \tan \dots)}}{(1-i \tan \dots)}\right)}{3(c^2 + d^2)(bc - ad)}$$

104

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} +$$

$$\frac{2\sqrt{a+b \tan(e+fx)}(b(4c^2d^2(2A-C)+2Ad^4-5Bc^3d+Bcd^3+2c^4C)-3ad^2(2cd(A-C)-B(c^2-d^2)))}{f(c^2+d^2)(bc-ad)\sqrt{c+d \tan(e+fx)}} - \frac{3\left(\frac{(c-id)^2(A+iB-C)(bc-ad)^2 \int \frac{\dots}{-ia+b \dots}}{-ia+b \dots}\right)}{3(c^2 + d^2)(bc - ad)}$$

221

$$\frac{2(Ad^2 - Bcd + c^2C) \sqrt{a + b \tan(e + fx)}}{3f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^{3/2}} + \frac{2\sqrt{a + b \tan(e + fx)}(b(4c^2d^2(2A - C) + 2Ad^4 - 5Bc^3d + Bcd^3 + 2c^4C) - 3ad^2(2cd(A - C) - B(c^2 - d^2)))}{f(c^2 + d^2)(bc - ad)\sqrt{c + d \tan(e + fx)}} - \frac{3 \left(\frac{i(c+id)^2(A-iB-C)(bc-ad)^2 \arctanh\left(\frac{f\sqrt{a-ib}\sqrt{c-id}}{f\sqrt{a-ib}\sqrt{c-id}}\right)}{f\sqrt{a-ib}\sqrt{c-id}} \right)}{3(c^2 + d^2)(bc - ad)}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])^(5/2)),x]`

output `(2*(c^2*C - B*c*d + A*d^2)*Sqrt[a + b*Tan[e + f*x]])/(3*(b*c - a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) + ((-3*((I*(A - I*B - C)*(c + I*d)^2*(b*c - a*d)^2*ArcTanh[(Sqrt[c - I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a - I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a - I*b]*Sqrt[c - I*d]*f) - (I*(A + I*B - C)*(c - I*d)^2*(b*c - a*d)^2*ArcTanh[(Sqrt[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[a + I*b]*Sqrt[c + I*d]*f)))/((b*c - a*d)*(c^2 + d^2)) + (2*(b*(2*c^4*C - 5*B*c^3*d + 4*c^2*(2*A - C)*d^2 + B*c*d^3 + 2*A*d^4) - 3*a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)))*Sqrt[a + b*Tan[e + f*x]])/((b*c - a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e + f*x]])/(3*(b*c - a*d)*(c^2 + d^2))`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 221 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan(fx + e) + C \tan(fx + e)^2}{\sqrt{a + b \tan(fx + e)} (c + d \tan(fx + e))^{\frac{5}{2}}} dx$$

input `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(5/2),x)`

output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(5/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)} (c + d \tan(e + fx))^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(1/2)/(c+d*tan(f*x+e))**(5/2),x)`

output

```
Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(sqrt(a + b*tan(e + f*x)
)*(c + d*tan(e + f*x))**(5/2)), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(
f*x+e))^(5/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx = \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{\sqrt{b \tan(fx + e) + a}(d \tan(fx + e) + c)^{5/2}} dx$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(
f*x+e))^(5/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/(sqrt(b*tan(f*x + e) + a
)*(d*tan(f*x + e) + c)^(5/2)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(1/2)*(c + d*tan(e + f*x))^(5/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(5/2),x)
```

output

```
(4*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)*b*c*d -
2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*c*d + 6*sqrt(tan(e +
f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b*c**2 + 3*int((sqrt(tan(e + f*x)*d
+ c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b*d**3 + tan(
e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d*
*2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b
*c**3 + a*c**3),x)*tan(e + f*x)**2*a**2*b*d**4*f - 6*int((sqrt(tan(e + f*x)
)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b*d**3 +
tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*
c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(e + f*
x)*b*c**3 + a*c**3),x)*tan(e + f*x)**2*a*b**2*c*d**3*f + 3*int((sqrt(tan(e
+ f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)**4*b*d
**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)
**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d + tan(
e + f*x)*b*c**3 + a*c**3),x)*tan(e + f*x)**2*b**3*c**2*d**2*f + 6*int((sqr
t(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e + f*x)
**4*b*d**3 + tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2 + 3*tan(e
+ f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d
+ tan(e + f*x)*b*c**3 + a*c**3),x)*tan(e + f*x)*a**2*b*c*d**3*f - 12*int(
(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e...
```

3.163
$$\int \frac{A+B \tan(e+fx)+C \tan^2(e+fx)}{(a+b \tan(e+fx))^{3/2}(c+d \tan(e+fx))^{5/2}} dx$$

Optimal result	1808
Mathematica [A] (verified)	1809
Rubi [A] (verified)	1810
Maple [F(-1)]	1815
Fricas [F(-1)]	1816
Sympy [F]	1816
Maxima [F(-1)]	1817
Giac [F]	1817
Mupad [F(-1)]	1817
Reduce [F]	1818

Optimal result

Integrand size = 49, antiderivative size = 651

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2}(c + d \tan(e + fx))^{5/2}} dx =$$

$$\frac{(iA + B - iC) \operatorname{arctanh}\left(\frac{\sqrt{c-id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a-ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a - ib)^{3/2}(c - id)^{5/2} f}$$

$$- \frac{(B - i(A - C)) \operatorname{arctanh}\left(\frac{\sqrt{c+id}\sqrt{a+b \tan(e+fx)}}{\sqrt{a+ib}\sqrt{c+d \tan(e+fx)}}\right)}{(a + ib)^{3/2}(c + id)^{5/2} f}$$

$$- \frac{2(Ab^2 - a(bB - aC))}{(a^2 + b^2)(bc - ad)f\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

$$- \frac{2d(b^2c(cC - Bd) - 3abB(c^2 + d^2) + a^2(4c^2C - Bcd + 3Cd^2) + A(a^2d^2 + b^2(3c^2 + 4d^2)))\sqrt{a + b \tan(e + fx)}}{3(a^2 + b^2)(bc - ad)^2(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}}$$

$$- \frac{2d(b^3c(5c^3C - 8Bc^2d - cCd^2 - 2Bd^3) + a^2b(8c^4C - 8Bc^3d + 5c^2Cd^2 - 2Bcd^3 + 3Cd^4) + 3a^3d^2(2cCd^2 + 3(a^2 + b^2)cd))\sqrt{a + b \tan(e + fx)}}{3(a^2 + b^2)(bc - ad)^2(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}}$$

output

```

-(I*A+B-I*C)*arctanh((c-I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a-I*b)^(1/2)/(c
+d*tan(f*x+e))^(1/2))/(a-I*b)^(3/2)/(c-I*d)^(5/2)/f-(B-I*(A-C))*arctanh((c
+I*d)^(1/2)*(a+b*tan(f*x+e))^(1/2)/(a+I*b)^(1/2)/(c+d*tan(f*x+e))^(1/2))/(
a+I*b)^(3/2)/(c+I*d)^(5/2)/f-2*(A*b^2-a*(B*b-C*a))/(a^2+b^2)/(-a*d+b*c)/f/
(a+b*tan(f*x+e))^(1/2)/(c+d*tan(f*x+e))^(3/2)-2/3*d*(b^2*c*(-B*d+C*c)-3*a*
b*B*(c^2+d^2)+a^2*(-B*c*d+4*C*c^2+3*C*d^2)+A*(a^2*d^2+b^2*(3*c^2+4*d^2)))*
(a+b*tan(f*x+e))^(1/2)/(a^2+b^2)/(-a*d+b*c)^2/(c^2+d^2)/f/(c+d*tan(f*x+e))
^(3/2)-2/3*d*(b^3*c*(-8*B*c^2*d-2*B*d^3+5*C*c^3-C*c*d^2)+a^2*b*(-8*B*c^3*d
-2*B*c*d^3+8*C*c^4+5*C*c^2*d^2+3*C*d^4)+3*a^3*d^2*(2*C*c*d+B*(c^2-d^2))+3*
a*b^2*(2*c*C*d^3-B*(c^4+c^2*d^2+2*d^4))-A*(6*a^3*c*d^3+6*a*b^2*c*d^3-a^2*b
*d^2*(11*c^2+5*d^2)-b^3*(3*c^4+17*c^2*d^2+8*d^4)))*(a+b*tan(f*x+e))^(1/2)/
(a^2+b^2)/(-a*d+b*c)^3/(c^2+d^2)^2/f/(c+d*tan(f*x+e))^(1/2)
    
```

Mathematica [A] (verified)

Time = 6.72 (sec) , antiderivative size = 903, normalized size of antiderivative = 1.39

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \frac{2(Ab^2 - a(bB - aC))}{(a^2 + b^2)(bc - ad)f\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

$$2 \left(-\frac{2(-c(-2c(Ab^2 - a(bB - aC))d + \frac{1}{2}(Ab - aB - bC)d(bc - ad)) + \frac{1}{2}d^2(4Ab^2d - aA(bc - ad) - (bB - aC)(bc + 3ad)))\sqrt{a + b \tan(e + fx)}}{3(-bc + ad)(c^2 + d^2)f(c + d \tan(e + fx))^{3/2}} \right) - \frac{3C}{2} \frac{1}{(c + d \tan(e + fx))^{3/2}}$$

input

```

Integrate[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(3
/2)*(c + d*Tan[e + f*x])^(5/2)),x]
    
```

output

```

(-2*(A*b^2 - a*(b*B - a*C)))/((a^2 + b^2)*(b*c - a*d)*f*Sqrt[a + b*Tan[e +
f*x]]*(c + d*Tan[e + f*x])^(3/2)) - (2*((-2*(-c*(-2*c*(A*b^2 - a*(b*B -
a*C))*d + ((A*b - a*B - b*C)*d*(b*c - a*d))/2)) + (d^2*(4*A*b^2*d - a*A*(b
*c - a*d) - (b*B - a*C)*(b*c + 3*a*d)))/2)*Sqrt[a + b*Tan[e + f*x]]/(3*(-
(b*c) + a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^(3/2)) - (2*((3*(b*c - a*d
)^3*((a + I*b)*(I*A + B - I*C)*(c + I*d)^2*ArcTanh[(Sqrt[-c + I*d]*Sqrt[a
+ b*Tan[e + f*x]])/(Sqrt[-a + I*b]*Sqrt[c + d*Tan[e + f*x]])])/(Sqrt[-a +
I*b]*Sqrt[-c + I*d]) + ((I*a + b)*(A + I*B - C)*(c - I*d)^2*ArcTanh[(Sqrt
[c + I*d]*Sqrt[a + b*Tan[e + f*x]])/(Sqrt[a + I*b]*Sqrt[c + d*Tan[e + f*x]
]])/(Sqrt[a + I*b]*Sqrt[c + I*d])))/(4*(-(b*c) + a*d)*(c^2 + d^2)*f) - (2
*(d^2*((b*c)/2 - (3*a*d)/2)*(-2*c*(A*b^2 - a*(b*B - a*C))*d + ((A*b - a*B
- b*C)*d*(b*c - a*d))/2) + ((b*d^2 - (3*c*(-(b*c) + a*d))/2)*(4*A*b^2*d -
a*A*(b*c - a*d) - (b*B - a*C)*(b*c + 3*a*d)))/2) - c*((3*d*(-(b*c) + a*d)
*(-2*(A*b^2 - a*(b*B - a*C))*d^2 - (c*(A*b - a*B - b*C)*(b*c - a*d))/2 + (
d*(4*A*b^2*d - a*A*(b*c - a*d) - (b*B - a*C)*(b*c + 3*a*d)))/2))/2 - b*c*(
(-c*(-2*c*(A*b^2 - a*(b*B - a*C))*d + ((A*b - a*B - b*C)*d*(b*c - a*d))/2)
) + (d^2*(4*A*b^2*d - a*A*(b*c - a*d) - (b*B - a*C)*(b*c + 3*a*d)))/2))*S
qrt[a + b*Tan[e + f*x]]/((- (b*c) + a*d)*(c^2 + d^2)*f*Sqrt[c + d*Tan[e +
f*x]])))/(3*(-(b*c) + a*d)*(c^2 + d^2)))/((a^2 + b^2)*(b*c - a*d))

```

Rubi [A] (verified)

Time = 7.49 (sec) , antiderivative size = 751, normalized size of antiderivative = 1.15, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx) + C \tan(e + fx)^2}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx$$

↓ 4132

$$2 \int \frac{4Adb^2 + 4(Ab^2 - a(bB - aC))d \tan^2(e + fx) - aA(bc - ad) - (bB - aC)(bc + 3ad) + (Ab - Cb - aB)(bc - ad) \tan(e + fx)}{2\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))}$$

$$f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}$$

↓ 27

$$\int \frac{4Adb^2 + 4(Ab^2 - a(bB - aC))d \tan^2(e + fx) - aA(bc - ad) - (bB - aC)(bc + 3ad) + (Ab - Cb - aB)(bc - ad) \tan(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))}$$

$$f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}$$

↓ 3042

$$\int \frac{4Adb^2 + 4(Ab^2 - a(bB - aC))d \tan(e + fx)^2 - aA(bc - ad) - (bB - aC)(bc + 3ad) + (Ab - Cb - aB)(bc - ad) \tan(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2}} dx$$

$$\frac{(a^2 + b^2)(bc - ad)}{2(Ab^2 - a(bB - aC))}$$

$$f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}$$

↓ 4132

$$2 \int \frac{-3(aBc + bCc - bBd + aCd - A(bc + ad)) \tan(e + fx)(bc - ad)^2 + 2bd(Ad^2a^2 + (4Cc^2 - Bdc + 3Cd^2)a^2 - 3bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(3c^2 + 4d^2)) \tan^2(e + fx)}{2\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2} \cdot 3(c^2 + d^2)(bc - ad)}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

↓ 27

$$\int \frac{-3(aBc + bCc - bBd + aCd - A(bc + ad)) \tan(e + fx)(bc - ad)^2 + 2bd(Ad^2a^2 + (4Cc^2 - Bdc + 3Cd^2)a^2 - 3bB(c^2 + d^2)a + b^2c(cC - Bd) + Ab^2(3c^2 + 4d^2)) \tan^2(e + fx)}{\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{5/2} \cdot 3(c^2 + d^2)(bc - ad)}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

↓ 3042

$$\int \frac{-3(aBc+bCc-bBd+aCd-A(bc+ad)) \tan(e+fx)(bc-ad)^2+2bd(Ad^2a^2+(4Cc^2-Bdc+3Cd^2)a^2-3bB(c^2+d^2)a+b^2c(cC-Bd)+Ab^2(3c^2+4d^2)) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}(c+d \tan(e+fx)) \frac{3(c^2+d^2)(bc-ad)}{3(c^2+d^2)(bc-ad)}}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

4132

$$2 \int \frac{3\left(\left(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))+b(2c(A-C)d-B(c^2-d^2))\right)(bc-ad)^3+(2aAcd-2acCd+Ab(c^2-d^2)-aB(c^2-d^2)-b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{2\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(c^2+d^2)(bc-ad)} \tan(e+fx)}{(c^2+d^2)(bc-ad)}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

27

$$3 \int \frac{\left(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))+b(2c(A-C)d-B(c^2-d^2))\right)(bc-ad)^3+(2aAcd-2acCd+Ab(c^2-d^2)-aB(c^2-d^2)-b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(c^2+d^2)(bc-ad)} \tan(e+fx)}{(c^2+d^2)(bc-ad)}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

3042

$$3 \int \frac{\left(a(Cc^2-2Bdc-Cd^2-A(c^2-d^2))+b(2c(A-C)d-B(c^2-d^2))\right)(bc-ad)^3+(2aAcd-2acCd+Ab(c^2-d^2)-aB(c^2-d^2)-b(Cc^2-2Bdc-Cd^2)) \tan(e+fx)}{\sqrt{a+b \tan(e+fx)}\sqrt{c+d \tan(e+fx)}(c^2+d^2)(bc-ad)} \tan(e+fx)}{(c^2+d^2)(bc-ad)}$$

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

4099

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}}$$

$$\frac{2d\sqrt{a+b \tan(e+fx)}(a^2Ad^2+a^2(-Bcd+4c^2C+3Cd^2)-3abB(c^2+d^2)+Ab^2(3c^2+4d^2)+b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2d\sqrt{a+b \tan(e+fx)}(3a^3d^2(B(c^2-d^2)+$$

3042

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2 + a^2(-Bcd+4c^2 C+3Cd^2) - 3abB(c^2+d^2) + Ab^2(3c^2+4d^2) + b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2d\sqrt{a+b \tan(e+fx)}(3a^3 d^2(B(c^2-d^2)) + \dots)}{\dots}$$

↓ 4098

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2 + a^2(-Bcd+4c^2 C+3Cd^2) - 3abB(c^2+d^2) + Ab^2(3c^2+4d^2) + b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2d\sqrt{a+b \tan(e+fx)}(3a^3 d^2(B(c^2-d^2)) + \dots)}{\dots}$$

↓ 104

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2 + a^2(-Bcd+4c^2 C+3Cd^2) - 3abB(c^2+d^2) + Ab^2(3c^2+4d^2) + b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2d\sqrt{a+b \tan(e+fx)}(3a^3 d^2(B(c^2-d^2)) + \dots)}{\dots}$$

↓ 221

$$\frac{2(Ab^2 - a(bB - aC))}{f(a^2 + b^2)(bc - ad)\sqrt{a + b \tan(e + fx)}(c + d \tan(e + fx))^{3/2}} - \frac{2d\sqrt{a+b \tan(e+fx)}(a^2 Ad^2 + a^2(-Bcd+4c^2 C+3Cd^2) - 3abB(c^2+d^2) + Ab^2(3c^2+4d^2) + b^2c(cC-Bd))}{3f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^{3/2}} + \frac{2d\sqrt{a+b \tan(e+fx)}(3a^3 d^2(B(c^2-d^2)) + \dots)}{\dots}$$

input `Int[(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2)/((a + b*Tan[e + f*x])^(3/2)*(c + d*Tan[e + f*x])^(5/2)),x]`

output

$$\begin{aligned} & (-2*(A*b^2 - a*(b*B - a*C)))/((a^2 + b^2)*(b*c - a*d)*f*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]*(c + d*\text{Tan}[e + f*x])^{(3/2)}) - ((2*d*(a^2*A*d^2 + b^2*c*(c*C - B*d) \\ & - 3*a*b*B*(c^2 + d^2) + A*b^2*(3*c^2 + 4*d^2) + a^2*(4*c^2*C - B*c*d + 3*C*d^2))*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])/(3*(b*c - a*d)*(c^2 + d^2)*f*(c + d*\text{Tan}[e \\ & + f*x])^{(3/2)}) + ((3*((I*(a + I*b)*(A - I*B - C)*(c + I*d)^2*(b*c - a*d)^3 \\ & * \text{ArcTanh}[(\text{Sqrt}[c - I*d]*\text{Sqrt}[a + b*\text{Tan}[e + f*x]])]/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c + \\ & d*\text{Tan}[e + f*x]])))/(\text{Sqrt}[a - I*b]*\text{Sqrt}[c - I*d]*f) - (I*(a - I*b)*(A + I*B \\ & - C)*(c - I*d)^2*(b*c - a*d)^3 * \text{ArcTanh}[(\text{Sqrt}[c + I*d]*\text{Sqrt}[a + b*\text{Tan}[e + \\ & f*x]])]/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + d*\text{Tan}[e + f*x]])))/(\text{Sqrt}[a + I*b]*\text{Sqrt}[c + \\ & I*d]*f)))/((b*c - a*d)*(c^2 + d^2)) + (2*d*(b^3*c*(5*c^3*C - 8*B*c^2*d - \\ & c*C*d^2 - 2*B*d^3) + a^2*b*(8*c^4*C - 8*B*c^3*d + 5*c^2*C*d^2 - 2*B*c*d^3 \\ & + 3*C*d^4) + 3*a^3*d^2*(2*c*C*d + B*(c^2 - d^2)) + 3*a*b^2*(2*c*C*d^3 - B*(c^4 \\ & + c^2*d^2 + 2*d^4)) - A*(6*a^3*c*d^3 + 6*a*b^2*c*d^3 - a^2*b*d^2*(11*c^2 \\ & + 5*d^2) - b^3*(3*c^4 + 17*c^2*d^2 + 8*d^4)))*\text{Sqrt}[a + b*\text{Tan}[e + f*x]]) \\ &)/((b*c - a*d)*(c^2 + d^2)*f*\text{Sqrt}[c + d*\text{Tan}[e + f*x]]))/(3*(b*c - a*d)*(c^2 \\ & + d^2)))/((a^2 + b^2)*(b*c - a*d)) \end{aligned}$$

Defintions of rubi rules used

rule 27

$$\text{Int}[(a_*)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)*(G_x) \text{ ; FreeQ}[b, x]]$$

rule 104

$$\text{Int}[(((a_.) + (b_.)*(x_))^{(m_)*((c_.) + (d_.)*(x_))^{(n_))}/((e_.) + (f_.)*(x_)), x_] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Simp}[q \quad \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$$

rule 221

$$\text{Int}[((a_) + (b_.)*(x_)^2)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$$

rule 3042

$$\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

rule 4098

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

rule 4099

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

rule 4132

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F(-1)]

Timed out.

$$\int \frac{A + B \tan (fx + e) + C \tan (fx + e)^2}{(a + b \tan (fx + e))^{\frac{3}{2}} (c + d \tan (fx + e))^{\frac{5}{2}}} dx$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e)
)^(5/2),x)
```

output `int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(5/2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{\frac{3}{2}} (c + d \tan(e + fx))^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(a+b*tan(f*x+e))**(3/2)/(c+d*tan(f*x+e))**(5/2),x)`

output `Integral((A + B*tan(e + f*x) + C*tan(e + f*x)**2)/((a + b*tan(e + f*x))**(3/2)*(c + d*tan(e + f*x))**(5/2)), x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

output

Timed out

Giac [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \int \frac{C \tan^2(fx + e) + B \tan(fx + e) + A}{(b \tan(fx + e) + a)^{3/2} (d \tan(fx + e) + c)^{5/2}} dx$$

input

```
integrate((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(5/2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)/((b*tan(f*x + e) + a)^(3/2)*(d*tan(f*x + e) + c)^(5/2)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \text{Hanged}$$

input

```
int((A + B*tan(e + f*x) + C*tan(e + f*x)^2)/((a + b*tan(e + f*x))^(3/2)*(c + d*tan(e + f*x))^(5/2)),x)
```

output

\text{Hanged}

Reduce [F]

$$\int \frac{A + B \tan(e + fx) + C \tan^2(e + fx)}{(a + b \tan(e + fx))^{3/2} (c + d \tan(e + fx))^{5/2}} dx = \text{too large to display}$$

input

```
int((A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(a+b*tan(f*x+e))^(3/2)/(c+d*tan(f*x+e))^(5/2),x)
```

output

```
(16*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)**2*b**2
*c*d**2 + 8*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x)
*a*b*c*d**2 + 24*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e +
f*x)*b**2*c**2*d - 2*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a
**2*c*d**2 + 12*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*a*b*c**2*
d + 6*sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*b**2*c**3 + 3*int(
(sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e +
f*x)**5*b**2*d**3 + 2*tan(e + f*x)**4*a*b*d**3 + 3*tan(e + f*x)**4*b**2*c
*d**2 + tan(e + f*x)**3*a**2*d**3 + 6*tan(e + f*x)**3*a*b*c*d**2 + 3*tan(e
+ f*x)**3*b**2*c**2*d + 3*tan(e + f*x)**2*a**2*c*d**2 + 6*tan(e + f*x)**2*
a*b*c**2*d + tan(e + f*x)**2*b**2*c**3 + 3*tan(e + f*x)*a**2*c**2*d + 2*ta
n(e + f*x)*a*b*c**3 + a**2*c**3),x)*tan(e + f*x)**3*a**3*b**2*d**5*f - 9*i
nt((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(tan(e
+ f*x)**5*b**2*d**3 + 2*tan(e + f*x)**4*a*b*d**3 + 3*tan(e + f*x)**4*b**2
*c*d**2 + tan(e + f*x)**3*a**2*d**3 + 6*tan(e + f*x)**3*a*b*c*d**2 + 3*tan
(e + f*x)**3*b**2*c**2*d + 3*tan(e + f*x)**2*a**2*c*d**2 + 6*tan(e + f*x)*
**2*a*b*c**2*d + tan(e + f*x)**2*b**2*c**3 + 3*tan(e + f*x)*a**2*c**2*d + 2
*tan(e + f*x)*a*b*c**3 + a**2*c**3),x)*tan(e + f*x)**3*a**2*b**3*c*d**4*f
+ 9*int((sqrt(tan(e + f*x)*d + c)*sqrt(tan(e + f*x)*b + a)*tan(e + f*x))/(
tan(e + f*x)**5*b**2*d**3 + 2*tan(e + f*x)**4*a*b*d**3 + 3*tan(e + f*x)...
```

3.164 $\int (a+b \tan(e+fx))^m (c+d \tan(e+fx))^n (A + B \tan(e+fx) + C \tan^2(e+fx)) dx =$

Optimal result	1819
Mathematica [F]	1820
Rubi [A] (verified)	1820
Maple [F]	1822
Fricas [F]	1822
Sympy [F(-2)]	1823
Maxima [F]	1823
Giac [F]	1823
Mupad [F(-1)]	1824
Reduce [F]	1824

Optimal result

Integrand size = 45, antiderivative size = 376

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx =$$

$$\frac{(B + i(A - C)) \operatorname{AppellF1}\left(1 + m, -n, 1, 2 + m, -\frac{d(a + b \tan(e + fx))}{bc - ad}, \frac{a + b \tan(e + fx)}{a - ib}\right) (a + b \tan(e + fx))^{1+m}}{2(a - ib)f(1 + m)}$$

$$\frac{(A + iB - C) \operatorname{AppellF1}\left(1 + m, -n, 1, 2 + m, -\frac{d(a + b \tan(e + fx))}{bc - ad}, \frac{a + b \tan(e + fx)}{a + ib}\right) (a + b \tan(e + fx))^{1+m}}{2(ia - b)f(1 + m)}$$

$$+ \frac{C \operatorname{Hypergeometric2F1}\left(1 + m, -n, 2 + m, -\frac{d(a + b \tan(e + fx))}{bc - ad}\right) (a + b \tan(e + fx))^{1+m} (c + d \tan(e + fx))}{bf(1 + m)}$$

output

```
-1/2*(B+I*(A-C))*AppellF1(1+m,1,-n,2+m,(a+b*tan(f*x+e))/(a-I*b),-d*(a+b*tan(f*x+e))/(-a*d+b*c))*(a+b*tan(f*x+e))^(1+m)*(c+d*tan(f*x+e))^n/(a-I*b)/f/(1+m)/((b*(c+d*tan(f*x+e))/(-a*d+b*c))^n)-1/2*(A+I*B-C)*AppellF1(1+m,1,-n,2+m,(a+b*tan(f*x+e))/(a+I*b),-d*(a+b*tan(f*x+e))/(-a*d+b*c))*(a+b*tan(f*x+e))^(1+m)*(c+d*tan(f*x+e))^n/(I*a-b)/f/(1+m)/((b*(c+d*tan(f*x+e))/(-a*d+b*c))^n)+C*hypergeom([-n,1+m],[2+m],-d*(a+b*tan(f*x+e))/(-a*d+b*c))*(a+b*tan(f*x+e))^(1+m)*(c+d*tan(f*x+e))^n/b/f/(1+m)/((b*(c+d*tan(f*x+e))/(-a*d+b*c))^n)
```


Mathematica [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input

```
Integrate[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(A + B*Tan[e + f*x]
] + C*Tan[e + f*x]^2), x]
```

output

```
Integrate[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(A + B*Tan[e + f*x]
] + C*Tan[e + f*x]^2), x]
```

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 371, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4138, 2348, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) (c + d \tan(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)^2) (c + d \tan(e + fx))^n dx$$

$$\downarrow \text{4138}$$

$$\int \frac{(a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (C \tan^2(e + fx) + B \tan(e + fx) + A)}{\tan^2(e + fx) + 1} d \tan(e + fx)$$

$$\downarrow \text{2348}$$

$$\int \frac{C(c + d \tan(e + fx))^n (a + b \tan(e + fx))^m + \frac{(i(A - C) - B)(c + d \tan(e + fx))^n (a + b \tan(e + fx))^m}{2(i - \tan(e + fx))} + \frac{(B + i(A - C))(c + d \tan(e + fx))^n (a + b \tan(e + fx))^m}{2(\tan(e + fx) - i)}}{f} dx$$

↓ 2009

$$\frac{(B+i(A-C))(a+b\tan(e+fx))^{m+1}(c+d\tan(e+fx))^n \left(\frac{b(c+d\tan(e+fx))}{bc-ad}\right)^{-n} \operatorname{AppellF1}\left(m+1, -n, 1, m+2, -\frac{d(a+b\tan(e+fx))}{bc-ad}, \frac{a+b\tan(e+fx)}{a-ib}\right)}{2(m+1)(a-ib)}$$

input `Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]`

output `(-1/2*((B + I*(A - C))*AppellF1[1 + m, -n, 1, 2 + m, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d)), (a + b*Tan[e + f*x])/(a - I*b)]*(a + b*Tan[e + f*x])^(1 + m)*(c + d*Tan[e + f*x])^n)/((a - I*b)*(1 + m)*((b*(c + d*Tan[e + f*x]))/(b*c - a*d))^n) - ((A + I*B - C)*AppellF1[1 + m, -n, 1, 2 + m, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d)), (a + b*Tan[e + f*x])/(a + I*b)]*(a + b*Tan[e + f*x])^(1 + m)*(c + d*Tan[e + f*x])^n)/(2*(I*a - b)*(1 + m)*((b*(c + d*Tan[e + f*x]))/(b*c - a*d))^n) + (C*Hypergeometric2F1[1 + m, -n, 2 + m, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d))]*(a + b*Tan[e + f*x])^(1 + m)*(c + d*Tan[e + f*x])^n)/(b*(1 + m)*((b*(c + d*Tan[e + f*x]))/(b*c - a*d))^n)/f`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2348 `Int[(Px_)*((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(c + d*x)^m*(e + f*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && (IntegerQ[p] || (IntegerQ[2*p] && IntegerQ[m] && ILtQ[n, 0])) && !(IGtQ[m, 0] && IGtQ[n, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4138

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

Maple [F]

$$\int (a + b \tan (fx + e))^m (c + d \tan (fx + e))^n (A + B \tan (fx + e) + C \tan (fx + e)^2) dx$$

input

```
int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^n*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x)
```

output

```
int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^n*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x)
```

Fricas [F]

$$\int (a + b \tan (e + fx))^m (c + d \tan (e + fx))^n (A + B \tan (e + fx) + C \tan^2 (e + fx)) dx$$

$$= \int (C \tan (fx + e)^2 + B \tan (fx + e) + A) (b \tan (fx + e) + a)^m (d \tan (fx + e) + c)^n dx$$

input

```
integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^n*(A+B*tan(f*x+e)+C*tan(f*x+
e)^2),x, algorithm="fricas")
```

output

```
integral((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m*(d
*tan(f*x + e) + c)^n, x)
```

Sympy [F(-2)]

Exception generated.

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Exception raised: HeuristicGCDFailed

input

```
integrate((a+b*tan(f*x+e))**m*(c+d*tan(f*x+e))**n*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)
```

output

```
Exception raised: HeuristicGCDFailed >> no luck
```

Maxima [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m (d \tan(fx + e) + c)^n dx$$

input

```
integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^n*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m*(d*tan(f*x + e) + c)^n, x)
```

Giac [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m (d \tan(fx + e) + c)^n dx$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^n*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m*(d*tan(f*x + e) + c)^n, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (C \tan(e + fx)^2 + B \tan(e + fx) + A) dx$$

input `int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))^n*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))^n*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^n (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \left(\int (d \tan(fx + e) + c)^n (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) c$$

$$+ \left(\int (d \tan(fx + e) + c)^n (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) b$$

$$+ \left(\int (d \tan(fx + e) + c)^n (\tan(fx + e) b + a)^m dx \right) a$$

input `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^n*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)`

output

```
int((tan(e + f*x)*d + c)**n*(tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*c +  
int((tan(e + f*x)*d + c)**n*(tan(e + f*x)*b + a)**m*tan(e + f*x),x)*b + i  
nt((tan(e + f*x)*d + c)**n*(tan(e + f*x)*b + a)**m,x)*a
```

3.165 $\int (a+b \tan(e+fx))^m (c+d \tan(e+fx))^3 (A + B \tan(e+fx) + C \tan^2(e+fx)) dx$

Optimal result	1826
Mathematica [B] (verified)	1827
Rubi [A] (verified)	1828
Maple [F]	1834
Fricas [F]	1834
Sympy [F]	1834
Maxima [F(-1)]	1835
Giac [F]	1835
Mupad [F(-1)]	1836
Reduce [F]	1837

Optimal result

Integrand size = 45, antiderivative size = 560

$$\begin{aligned}
 & \int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
 = & \frac{(bc(2 + m)(b^2d(Bc + (A - C)d)(3 + m)(4 + m) - 2(bc - ad)(3aCd - b(3cC + Bd(4 + m)))) + d(b^3(2 + m)(3 + m)(4 + m) - 2(bc - ad)(3aCd - b(3cC + Bd(4 + m))))}{2(ia + b)f(1 + m)} \\
 & + \frac{(A - iB - C)(c - id)^3 \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a - ib}\right) (a + b \tan(e + fx))^{1+m}}{2(ia + b)f(1 + m)} \\
 & - \frac{(A + iB - C)(c + id)^3 \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a + ib}\right) (a + b \tan(e + fx))^{1+m}}{2(ia - b)f(1 + m)} \\
 & + \frac{d(b^2d(Bc + (A - C)d)(3 + m)(4 + m) - 2(bc - ad)(3aCd - b(3cC + Bd(4 + m)))) \tan(e + fx)(a + b \tan(e + fx))^{1+m}}{b^3f(2 + m)(3 + m)(4 + m)} \\
 & - \frac{(3aCd - b(3cC + Bd(4 + m)))(a + b \tan(e + fx))^{1+m}(c + d \tan(e + fx))^2}{b^2f(3 + m)(4 + m)} \\
 & + \frac{C(a + b \tan(e + fx))^{1+m}(c + d \tan(e + fx))^3}{bf(4 + m)}
 \end{aligned}$$

output

```
(b*c*(2+m)*(b^2*d*(B*c+(A-C)*d)*(3+m)*(4+m)-2*(-a*d+b*c)*(3*C*a*d-b*(3*C*c+B*d*(4+m))))+d*(b^3*(2*c*(A-C)*d+B*(c^2-d^2))*(2+m)*(3+m)*(4+m)-a*(b^2*d*(B*c+(A-C)*d)*(3+m)*(4+m)-2*(-a*d+b*c)*(3*C*a*d-b*(3*C*c+B*d*(4+m)))))*(a+b*tan(f*x+e))^(1+m)/b^4/f/(1+m)/(2+m)/(3+m)/(4+m)+1/2*(A-I*B-C)*(c-I*d)^3*hypergeom([1, 1+m], [2+m], (a+b*tan(f*x+e))/(a-I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a+b)/f/(1+m)-1/2*(A+I*B-C)*(c+I*d)^3*hypergeom([1, 1+m], [2+m], (a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a-b)/f/(1+m)+d*(b^2*d*(B*c+(A-C)*d)*(3+m)*(4+m)-2*(-a*d+b*c)*(3*C*a*d-b*(3*C*c+B*d*(4+m))))*tan(f*x+e)*(a+b*tan(f*x+e))^(1+m)/b^3/f/(2+m)/(3+m)/(4+m)-(3*C*a*d-b*(3*C*c+B*d*(4+m)))*(a+b*tan(f*x+e))^(1+m)*(c+d*tan(f*x+e))^2/b^2/f/(3+m)/(4+m)+C*(a+b*tan(f*x+e))^(1+m)*(c+d*tan(f*x+e))^3/b/f/(4+m)
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1390 vs. $2(560) = 1120$.

Time = 6.34 (sec) , antiderivative size = 1390, normalized size of antiderivative = 2.48

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Too large to display

input

```
Integrate[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```


output

```
(C*(a + b*Tan[e + f*x])^(1 + m)*(c + d*Tan[e + f*x])^3)/(b*f*(4 + m)) + ((
(3*b*c*C - 3*a*C*d + b*B*d*(4 + m))*(a + b*Tan[e + f*x])^(1 + m)*(c + d*Ta
n[e + f*x])^2)/(b*f*(3 + m)) + ((d*(b^2*d*(B*c + (A - C)*d)*(3 + m)*(4 + m)
) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m)))*Tan[e + f*x]*(a + b
*Tan[e + f*x])^(1 + m))/(b*f*(2 + m)) - (((-(b*c*(2 + m)*(b^2*d*(B*c + (A
- C)*d)*(3 + m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m)
))) + d*(-(b^3*(2*c*(A - C)*d + B*(c^2 - d^2))*(2 + m)*(3 + m)*(4 + m)) +
a*(b^2*d*(B*c + (A - C)*d)*(3 + m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*
C*d + b*B*d*(4 + m))))*(a + b*Tan[e + f*x])^(1 + m))/(b*f*(1 + m)) + ((I/
2)*(a*d*(b^2*d*(B*c + (A - C)*d)*(3 + m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C
- 3*a*C*d + b*B*d*(4 + m))) + b*c*(2 + m)*(b^2*d*(B*c + (A - C)*d)*(3 + m)
*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m))) - b*c*(2 + m)
)*(-(2*a*d + b*c*(1 + m))*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m))) + b*c*(3 +
m)*(A*b*c*(4 + m) - C*(3*a*d + b*c*(1 + m)))) - d*(-(b^3*(2*c*(A - C)*d +
B*(c^2 - d^2))*(2 + m)*(3 + m)*(4 + m)) + a*(b^2*d*(B*c + (A - C)*d)*(3 +
m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m)))) - I*b*(2
+ m)*(b^2*c*(2*c*(A - C)*d + B*(c^2 - d^2))*(3 + m)*(4 + m) - d*(b^2*d*(B
*c + (A - C)*d)*(3 + m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d
*(4 + m))) + d*(-(2*a*d + b*c*(1 + m))*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m)
)) + b*c*(3 + m)*(A*b*c*(4 + m) - C*(3*a*d + b*c*(1 + m)))))*Hypergeom...
```

Rubi [A] (verified)

Time = 5.84 (sec) , antiderivative size = 580, normalized size of antiderivative = 1.04, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.356$, Rules used = {3042, 4130, 3042, 4130, 25, 3042, 4120, 25, 3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))^3 (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (c + d \tan(e + fx))^3 (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4130$$

$$\frac{\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 ((3bcC - 3adC + bBd(m + 4)) \tan^2(e + fx) + b(Bc + (A - C)d)(m + 4))}{b(m + 4)}$$

$$\frac{C(c + d \tan(e + fx))^3 (a + b \tan(e + fx))^{m+1}}{bf(m + 4)}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 ((3bcC - 3adC + bBd(m + 4)) \tan(e + fx)^2 + b(Bc + (A - C)d)(m + 4))}{b(m + 4)}$$

$$\frac{C(c + d \tan(e + fx))^3 (a + b \tan(e + fx))^{m+1}}{bf(m + 4)}$$

↓ 4130

$$\frac{\int -(a + b \tan(e + fx))^m (c + d \tan(e + fx)) - ((2c(A - C)d + B(c^2 - d^2))(m + 3)(m + 4) \tan(e + fx)b^2) - c(m + 3)(Abc(m + 4) - C(3ad + bc(m + 1)))b}{b(m + 3)}$$

$$\frac{C(c + d \tan(e + fx))^3 (a + b \tan(e + fx))^{m+1}}{bf(m + 4)}$$

↓ 25

$$\frac{(c + d \tan(e + fx))^2 (-3aCd + bBd(m + 4) + 3bcC)(a + b \tan(e + fx))^{m+1}}{bf(m + 3)} - \frac{\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) - ((2c(A - C)d + B(c^2 - d^2))(m + 3)(m + 4) \tan(e + fx)b^2) - c(m + 3)(Abc(m + 4) - C(3ad + bc(m + 1)))b}{b(m + 3)}$$

$$\frac{C(c + d \tan(e + fx))^3 (a + b \tan(e + fx))^{m+1}}{bf(m + 4)}$$

↓ 3042

$$\frac{(c + d \tan(e + fx))^2 (-3aCd + bBd(m + 4) + 3bcC)(a + b \tan(e + fx))^{m+1}}{bf(m + 3)} - \frac{\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) - ((2c(A - C)d + B(c^2 - d^2))(m + 3)(m + 4) \tan(e + fx)b^2) - c(m + 3)(Abc(m + 4) - C(3ad + bc(m + 1)))b}{b(m + 3)}$$

$$\frac{C(c + d \tan(e + fx))^3 (a + b \tan(e + fx))^{m+1}}{bf(m + 4)}$$

↓ 4120

$$\frac{(c + d \tan(e + fx))^2 (-3aCd + bBd(m + 4) + 3bcC)(a + b \tan(e + fx))^{m+1}}{bf(m + 3)} - \frac{\int -(a + b \tan(e + fx))^m - (((A - C)d(3c^2 - d^2) + B(c^3 - 3cd^2))(m + 2)(m + 3))}{b(m + 3)}$$

$$\frac{C(c + d \tan(e + fx))^3 (a + b \tan(e + fx))^{m+1}}{bf(m + 4)}$$

↓ 25

$$\frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{f(a+b \tan(e+fx))^m(-((A-C)d(3c^2-d^2)+B(c^3-3cd^2))(m+2)(m^2+7m+10))}{bf(m+3)}$$

$$\frac{C(c+d \tan(e+fx))^3(a+b \tan(e+fx))^{m+1}}{bf(m+4)}$$

↓ 3042

$$\frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{f(a+b \tan(e+fx))^m(-((A-C)d(3c^2-d^2)+B(c^3-3cd^2))(m+2)(m^2+7m+10))}{bf(m+3)}$$

$$\frac{C(c+d \tan(e+fx))^3(a+b \tan(e+fx))^{m+1}}{bf(m+4)}$$

↓ 4113

$$\frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{f(a+b \tan(e+fx))^m(-((Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)(m+2)(m+3)))}{bf(m+3)}$$

$$\frac{C(c+d \tan(e+fx))^3(a+b \tan(e+fx))^{m+1}}{bf(m+4)}$$

↓ 3042

$$\frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{f(a+b \tan(e+fx))^m(-((Ac^3-Cc^3-3Bdc^2-3Ad^2c+3Cd^2c+Bd^3)(m+2)(m+3)))}{bf(m+3)}$$

$$\frac{C(c+d \tan(e+fx))^3(a+b \tan(e+fx))^{m+1}}{bf(m+4)}$$

↓ 4022

$$\frac{C(c+d \tan(e+fx))^3(a+b \tan(e+fx))^{m+1}}{bf(m+4)} +$$

$$\frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{d \tan(e+fx)(a+b \tan(e+fx))^{m+1}(2(bc-ad)(-3aCd+bBd(m+4)+3bcC)+b^2)}{bf(m+2)}$$

↓ 3042

$$\frac{C(c + d \tan(e + fx))^3(a + b \tan(e + fx))^{m+1}}{bf(m + 4)} + \frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{d \tan(e+fx)(a+b \tan(e+fx))^{m+1}(2(bc-ad)(-3aCd+bBd(m+4)+3bcC)+b^2)}{bf(m+2)}$$

↓ 4020

$$\frac{C(c + d \tan(e + fx))^3(a + b \tan(e + fx))^{m+1}}{bf(m + 4)} + \frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{d \tan(e+fx)(a+b \tan(e+fx))^{m+1}(2(bc-ad)(-3aCd+bBd(m+4)+3bcC)+b^2)}{bf(m+2)}$$

↓ 25

$$\frac{C(c + d \tan(e + fx))^3(a + b \tan(e + fx))^{m+1}}{bf(m + 4)} + \frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{d \tan(e+fx)(a+b \tan(e+fx))^{m+1}(2(bc-ad)(-3aCd+bBd(m+4)+3bcC)+b^2)}{bf(m+2)}$$

↓ 78

$$\frac{C(c + d \tan(e + fx))^3(a + b \tan(e + fx))^{m+1}}{bf(m + 4)} + \frac{(c+d \tan(e+fx))^2(-3aCd+bBd(m+4)+3bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+3)} - \frac{d \tan(e+fx)(a+b \tan(e+fx))^{m+1}(2(bc-ad)(-3aCd+bBd(m+4)+3bcC)+b^2)}{bf(m+2)}$$

input

```
Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^3*(A + B*Tan[e + f*x] + C*
Tan[e + f*x]^2),x]
```

output

$$\begin{aligned} & (C*(a + b*\text{Tan}[e + f*x])^{(1 + m)}*(c + d*\text{Tan}[e + f*x])^3)/(b*f*(4 + m)) + ((\\ & (3*b*c*C - 3*a*C*d + b*B*d*(4 + m))*(a + b*\text{Tan}[e + f*x])^{(1 + m)}*(c + d*\text{Tan} \\ & n[e + f*x])^2)/(b*f*(3 + m)) - (-((d*(b^2*d*(B*c + (A - C)*d)*(3 + m)*(4 + \\ & m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d*(4 + m)))*\text{Tan}[e + f*x]*(a + \\ & b*\text{Tan}[e + f*x])^{(1 + m)})/(b*f*(2 + m))) + (-(((b*c*(2 + m)*(b^2*d*(B*c + \\ & (A - C)*d)*(3 + m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C*d + b*B*d*(4 + \\ & m))) + d*(b^3*(2*c*(A - C)*d + B*(c^2 - d^2))*(2 + m)*(3 + m)*(4 + m) - a \\ & *(b^2*d*(B*c + (A - C)*d)*(3 + m)*(4 + m) + 2*(b*c - a*d)*(3*b*c*C - 3*a*C \\ & *d + b*B*d*(4 + m))))*(a + b*\text{Tan}[e + f*x])^{(1 + m)})/(b*f*(1 + m))) + ((I/ \\ & 2)*b^3*(A - I*B - C)*(c - I*d)^3*(2 + m)*(3 + m)*(4 + m)*\text{Hypergeometric2F1} \\ & [1, 1 + m, 2 + m, (a + b*\text{Tan}[e + f*x])/(a - I*b)]*(a + b*\text{Tan}[e + f*x])^{(1 \\ & + m)})/((a - I*b)*f*(1 + m)) - ((I/2)*b^3*(A + I*B - C)*(c + I*d)^3*(2 + m) \\ & *(3 + m)*(4 + m)*\text{Hypergeometric2F1}[1, 1 + m, 2 + m, (a + b*\text{Tan}[e + f*x])/(\\ & a + I*b)]*(a + b*\text{Tan}[e + f*x])^{(1 + m)})/((a + I*b)*f*(1 + m)))/(b*(2 + m) \\ &)/(b*(3 + m)))/(b*(4 + m)) \end{aligned}$$
Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), x_Symbol] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$$

rule 78

$$\begin{aligned} & \text{Int}[((a_) + (b_)*(x_))^{(m_)}*((c_) + (d_)*(x_))^{(n_)}, x_Symbol] \text{ :> } \text{Simp}[(b \\ & *c - a*d)^n*((a + b*x)^{(m + 1)})/(b^{(n + 1)}*(m + 1))*\text{Hypergeometric2F1}[-n, m \\ & + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] \text{ /; } \text{FreeQ}\{a, b, c, d, m\}, x] \\ & \&\& \text{!IntegerQ}[m] \&\& \text{IntegerQ}[n] \end{aligned}$$

rule 3042

$$\text{Int}[u_, x_Symbol] \text{ :> } \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ /; } \text{FunctionOfTrigOfLinear} \\ \text{Q}[u, x]$$

rule 4020

$$\begin{aligned} & \text{Int}(((a_.) + (b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^{(m_)}*((c_) + (d_.)*\text{tan}[(e_.) + \\ & (f_.)*(x_)]), x_Symbol] \text{ :> } \text{Simp}[c*(d/f) \quad \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + \\ & c*x), x], x, d*\text{Tan}[e + f*x], x] \text{ /; } \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[\\ & b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0] \end{aligned}$$

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F]

$$\int (a + b \tan(fx + e))^m (c + d \tan(fx + e))^3 (A + B \tan(fx + e) + C \tan(fx + e)^2) dx$$

input `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)`

output `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)`

Fricas [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) (d \tan(fx + e) + c)^3 (b \tan(fx + e) + a)^m dx$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `integral((C*d^3*tan(f*x + e)^5 + (3*C*c*d^2 + B*d^3)*tan(f*x + e)^4 + A*c^3 + (3*C*c^2*d + 3*B*c*d^2 + A*d^3)*tan(f*x + e)^3 + (C*c^3 + 3*B*c^2*d + 3*A*c*d^2)*tan(f*x + e)^2 + (B*c^3 + 3*A*c^2*d)*tan(f*x + e))*(b*tan(f*x + e) + a)^m, x)`

Sympy [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**m*(c+d*tan(f*x+e))**3*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**m*(c + d*tan(e + f*x))**3*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

= Timed out

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A)(d \tan(fx + e) + c)^3 (b \tan(fx + e) + a)^m dx$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^3*(b*tan(f*x + e) + a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + f x))^m (c + d \tan(e + f x))^3 (A + B \tan(e + f x) + C \tan^2(e + f x)) dx$$

$$= \int (a + b \tan(e + f x))^m (c + d \tan(e + f x))^3 (C \tan(e + f x)^2 + B \tan(e + f x) + A) dx$$

input

```
int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*
tan(e + f*x)^2),x)
```

output

```
int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))^3*(A + B*tan(e + f*x) + C*
tan(e + f*x)^2), x)
```

Reduce [F]

$$\begin{aligned}
& \int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^3 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\
&= \left(\int (\tan(fx + e) b + a)^m dx \right) a c^3 + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^5 dx \right) c d^3 \\
&\quad + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^4 dx \right) b d^3 \\
&\quad + 3 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^4 dx \right) c^2 d^2 \\
&\quad + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^3 dx \right) a d^3 \\
&\quad + 3 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^3 dx \right) b c d^2 \\
&\quad + 3 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^3 dx \right) c^3 d \\
&\quad + 3 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) a c d^2 \\
&\quad + 3 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) b c^2 d \\
&\quad + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) c^4 \\
&\quad + 3 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) a c^2 d \\
&\quad + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) b c^3
\end{aligned}$$

input `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^3*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)`

output

```
int((tan(e + f*x)*b + a)**m,x)*a*c**3 + int((tan(e + f*x)*b + a)**m*tan(e
+ f*x)**5,x)*c*d**3 + int((tan(e + f*x)*b + a)**m*tan(e + f*x)**4,x)*b*d**
3 + 3*int((tan(e + f*x)*b + a)**m*tan(e + f*x)**4,x)*c**2*d**2 + int((tan(
e + f*x)*b + a)**m*tan(e + f*x)**3,x)*a*d**3 + 3*int((tan(e + f*x)*b + a)*
**m*tan(e + f*x)**3,x)*b*c*d**2 + 3*int((tan(e + f*x)*b + a)**m*tan(e + f*x
)**3,x)*c**3*d + 3*int((tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*a*c*d**2
+ 3*int((tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*b*c**2*d + int((tan(e
+ f*x)*b + a)**m*tan(e + f*x)**2,x)*c**4 + 3*int((tan(e + f*x)*b + a)**m*t
an(e + f*x),x)*a*c**2*d + int((tan(e + f*x)*b + a)**m*tan(e + f*x),x)*b*c*
*3
```

3.166 $\int (a+b \tan(e+fx))^m (c+d \tan(e+fx))^2 (A + B \tan(e+fx) + C \tan^2(e+fx)) dx$

Optimal result	1839
Mathematica [A] (verified)	1840
Rubi [A] (verified)	1840
Maple [F]	1845
Fricas [F]	1845
Sympy [F]	1845
Maxima [F]	1846
Giac [F]	1846
Mupad [F(-1)]	1847
Reduce [F]	1847

Optimal result

Integrand size = 45, antiderivative size = 363

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(2a^2Cd^2 - abd(2cC + Bd)(3 + m) + b^2(2 + m)(2c^2C + 2Bcd(3 + m) + (A - C)d^2(3 + m)))(a + b \tan(e + fx))^{1+m}}{b^3 f(1 + m)(2 + m)(3 + m)}$$

$$+ \frac{(A - iB - C)(c - id)^2 \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a - ib}\right) (a + b \tan(e + fx))^{1+m}}{2(ia + b)f(1 + m)}$$

$$+ \frac{(iA - B - iC)(c + id)^2 \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a + ib}\right) (a + b \tan(e + fx))^{1+m}}{2(a + ib)f(1 + m)}$$

$$- \frac{d(2aCd - b(2cC + Bd(3 + m))) \tan(e + fx)(a + b \tan(e + fx))^{1+m}}{b^2 f(2 + m)(3 + m)}$$

$$+ \frac{C(a + b \tan(e + fx))^{1+m} (c + d \tan(e + fx))^2}{bf(3 + m)}$$

output

```
(2*a^2*C*d^2-a*b*d*(B*d+2*C*c)*(3+m)+b^2*(2+m)*(2*c^2*C+2*B*c*d*(3+m)+(A-C)*d^2*(3+m))*(a+b*tan(f*x+e))^(1+m)/b^3/f/(1+m)/(2+m)/(3+m)+1/2*(A-I*B-C)*(c-I*d)^2*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a-I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a+b)/f/(1+m)+1/2*(I*A-B-I*C)*(c+I*d)^2*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))^(1+m)/(a+I*b)/f/(1+m)-d*(2*C*a*d-b*(2*C*c+B*d*(3+m)))*tan(f*x+e)*(a+b*tan(f*x+e))^(1+m)/b^2/f/(2+m)/(3+m)+C*(a+b*tan(f*x+e))^(1+m)*(c+d*tan(f*x+e))^2/b/f/(3+m)
```

Mathematica [A] (verified)

Time = 5.03 (sec) , antiderivative size = 308, normalized size of antiderivative = 0.85

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(a + b \tan(e + fx))^{1+m} \left(\frac{2bc(2+m)(2bcC - 2aCd + bBd(3+m)) + 2d(b^2(Bc + (A-C)d)(2+m)(3+m) - a(2bcC - 2aCd + bBd(3+m)))}{(a - I*b)^2} \right)}{(a - I*b)^2}$$

input

```
Integrate[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]
```

output

```
((a + b*Tan[e + f*x])^(1 + m)*((2*b*c*(2 + m)*(2*b*c*C - 2*a*C*d + b*B*d*(3 + m)) + 2*d*(b^2*(B*c + (A - C)*d)*(2 + m)*(3 + m) - a*(2*b*c*C - 2*a*C*d + b*B*d*(3 + m))) - (I*b^3*(A - I*B - C)*(c - I*d)^2*(6 + 5*m + m^2)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)])/(a - I*b) + (I*b^3*(A + I*B - C)*(c + I*d)^2*(6 + 5*m + m^2)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)])/(a + I*b))/(b^2*(1 + m)*(2 + m)) + (2*d*(2*b*c*C - 2*a*C*d + b*B*d*(3 + m))*Tan[e + f*x])/(b*(2 + m)) + 2*C*(c + d*Tan[e + f*x])^2)/(2*b*f*(3 + m))
```

Rubi [A] (verified)

Time = 3.20 (sec) , antiderivative size = 385, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4130, 3042, 4120, 3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))^2 (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 3042

$$\int (c + d \tan(e + fx))^2 (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

↓ 4130

$$\frac{\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) ((2bcC - 2adC + bBd(m + 3)) \tan^2(e + fx) + b(Bc + (A - C)d)(m + 3))}{bf(m + 3)} \\ \frac{C(c + d \tan(e + fx))^2 (a + b \tan(e + fx))^{m+1}}{bf(m + 3)}$$

↓ 3042

$$\frac{\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) ((2bcC - 2adC + bBd(m + 3)) \tan(e + fx)^2 + b(Bc + (A - C)d)(m + 3))}{bf(m + 3)} \\ \frac{C(c + d \tan(e + fx))^2 (a + b \tan(e + fx))^{m+1}}{bf(m + 3)}$$

↓ 4120

$$\frac{d \tan(e + fx) (-2aCd + bBd(m + 3) + 2bcC) (a + b \tan(e + fx))^{m+1}}{bf(m + 2)} - \frac{\int (a + b \tan(e + fx))^m (-((2c(A - C)d + B(c^2 - d^2))(m + 2)(m + 3) \tan(e + fx)b^2))}{bf(m + 2)}$$

$$\frac{C(c + d \tan(e + fx))^2 (a + b \tan(e + fx))^{m+1}}{bf(m + 3)}$$

↓ 3042

$$\frac{d \tan(e + fx) (-2aCd + bBd(m + 3) + 2bcC) (a + b \tan(e + fx))^{m+1}}{bf(m + 2)} - \frac{\int (a + b \tan(e + fx))^m (-((2c(A - C)d + B(c^2 - d^2))(m + 2)(m + 3) \tan(e + fx)b^2))}{bf(m + 2)}$$

$$\frac{C(c + d \tan(e + fx))^2 (a + b \tan(e + fx))^{m+1}}{bf(m + 3)}$$

↓ 4113

$$\frac{d \tan(e + fx) (-2aCd + bBd(m + 3) + 2bcC) (a + b \tan(e + fx))^{m+1}}{bf(m + 2)} - \frac{\int (a + b \tan(e + fx))^m (-((Ac^2 - Cc^2 - 2Bdc - Ad^2 + Cd^2)(m + 2)(m + 3)b^2) - (2c(A - C)d + B(c^2 - d^2))(m + 2)(m + 3) \tan(e + fx)b^2))}{bf(m + 2)}$$

$$\frac{C(c + d \tan(e + fx))^2 (a + b \tan(e + fx))^{m+1}}{bf(m + 3)}$$

↓ 3042

$$\frac{d \tan(e+fx)(-2aCd+bBd(m+3)+2bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{f(a+b \tan(e+fx))^m(-((Ac^2-Cc^2-2Bdc-Ad^2+Cd^2)(m+2)(m+3)b^2)-(2$$

$$\frac{C(c+d \tan(e+fx))^2(a+b \tan(e+fx))^{m+1}}{bf(m+3)}$$

↓ 4022

$$\frac{C(c+d \tan(e+fx))^2(a+b \tan(e+fx))^{m+1}}{bf(m+3)} +$$

$$\frac{d \tan(e+fx)(-2aCd+bBd(m+3)+2bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{-\frac{1}{2}b^2(m+2)(m+3)(c+id)^2(A+iB-C) \int(1-i \tan(e+fx))(a+b \tan(e+fx))^m$$

↓ 3042

$$\frac{C(c+d \tan(e+fx))^2(a+b \tan(e+fx))^{m+1}}{bf(m+3)} +$$

$$\frac{d \tan(e+fx)(-2aCd+bBd(m+3)+2bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{-\frac{1}{2}b^2(m+2)(m+3)(c+id)^2(A+iB-C) \int(1-i \tan(e+fx))(a+b \tan(e+fx))^m$$

↓ 4020

$$\frac{C(c+d \tan(e+fx))^2(a+b \tan(e+fx))^{m+1}}{bf(m+3)} +$$

$$\frac{d \tan(e+fx)(-2aCd+bBd(m+3)+2bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{ib^2(m+2)(m+3)(c-id)^2(A-iB-C) \int \frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e+fx))}{2f} + \frac{ib^2(m+2)}$$

↓ 25

$$\frac{C(c+d \tan(e+fx))^2(a+b \tan(e+fx))^{m+1}}{bf(m+3)} +$$

$$\frac{d \tan(e+fx)(-2aCd+bBd(m+3)+2bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{ib^2(m+2)(m+3)(c-id)^2(A-iB-C) \int \frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e+fx))}{2f} - \frac{ib^2(m+2)}$$

↓ 78

$$\frac{C(c+d \tan(e+fx))^2(a+b \tan(e+fx))^{m+1}}{bf(m+3)} +$$

$$\frac{d \tan(e+fx)(-2aCd+bBd(m+3)+2bcC)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{(a+b \tan(e+fx))^{m+1}(2a^2Cd^2-abd(m+3)(Bd+2cC)+b^2(m+2)(d^2(m+3)(A-C)+$$

input `Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^2*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2),x]`

output `(C*(a + b*Tan[e + f*x])^(1 + m)*(c + d*Tan[e + f*x])^2)/(b*f*(3 + m)) + ((d*(2*b*c*C - 2*a*C*d + b*B*d*(3 + m))*Tan[e + f*x]*(a + b*Tan[e + f*x])^(1 + m))/(b*f*(2 + m)) - (((2*a^2*C*d^2 - a*b*d*(2*c*C + B*d)*(3 + m) + b^2*(2 + m)*(2*c^2*C + 2*B*c*d*(3 + m) + (A - C)*d^2*(3 + m)))*(a + b*Tan[e + f*x])^(1 + m))/(b*f*(1 + m))) + ((I/2)*b^2*(A - I*B - C)*(c - I*d)^2*(2 + m)*(3 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a - I*b)*f*(1 + m)) - ((I/2)*b^2*(A + I*B - C)*(c + I*d)^2*(2 + m)*(3 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a + I*b)*f*(1 + m)))/(b*(2 + m))/(b*(3 + m))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

rule 4120

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)]^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]
```

rule 4130

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e.
) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Maple [F]

$$\int (a + b \tan (fx + e))^m (c + d \tan (fx + e))^2 (A + B \tan (fx + e) + C \tan (fx + e)^2) dx$$

input `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)`

output `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2), x)`

Fricas [F]

$$\begin{aligned} & \int (a + b \tan (e + fx))^m (c + d \tan (e + fx))^2 (A + B \tan (e + fx) + C \tan^2 (e + fx)) dx \\ &= \int (C \tan (fx + e)^2 + B \tan (fx + e) + A) (d \tan (fx + e) + c)^2 (b \tan (fx + e) + a)^m dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `integral((C*d^2*tan(f*x + e)^4 + (2*C*c*d + B*d^2)*tan(f*x + e)^3 + A*c^2 + (C*c^2 + 2*B*c*d + A*d^2)*tan(f*x + e)^2 + (B*c^2 + 2*A*c*d)*tan(f*x + e))* (b*tan(f*x + e) + a)^m, x)`

Sympy [F]

$$\begin{aligned} & \int (a + b \tan (e + fx))^m (c + d \tan (e + fx))^2 (A + B \tan (e + fx) + C \tan^2 (e + fx)) dx \\ &= \int (a + b \tan (e + fx))^m (c + d \tan (e + fx))^2 (A + B \tan (e + fx) \\ & \hspace{15em} + C \tan^2 (e + fx)) dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))**m*(c+d*tan(f*x+e))**2*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**m*(c + d*tan(e + f*x))**2*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A)(d \tan(fx + e) + c)^2 (b \tan(fx + e) + a)^m dx$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^2*(b*tan(f*x + e) + a)^m, x)`

Giac [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A)(d \tan(fx + e) + c)^2 (b \tan(fx + e) + a)^m dx$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)^2*(b*tan(f*x + e) + a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (C \tan(e + fx)^2 + B \tan(e + fx) + A) dx$$

input

```
int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*
tan(e + f*x)^2),x)
```

output

```
int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))^2*(A + B*tan(e + f*x) + C*
tan(e + f*x)^2), x)
```

Reduce [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx))^2 (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \left(\int (\tan(fx + e) b + a)^m dx \right) a c^2 + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^4 dx \right) c d^2$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^3 dx \right) b d^2$$

$$+ 2 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^3 dx \right) c^2 d$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) a d^2$$

$$+ 2 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) b c d$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) c^3$$

$$+ 2 \left(\int (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) a c d$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) b c^2$$

input

```
int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))^2*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),
x)
```

output

```
int((tan(e + f*x)*b + a)**m,x)*a*c**2 + int((tan(e + f*x)*b + a)**m*tan(e
+ f*x)**4,x)*c*d**2 + int((tan(e + f*x)*b + a)**m*tan(e + f*x)**3,x)*b*d**
2 + 2*int((tan(e + f*x)*b + a)**m*tan(e + f*x)**3,x)*c**2*d + int((tan(e +
f*x)*b + a)**m*tan(e + f*x)**2,x)*a*d**2 + 2*int((tan(e + f*x)*b + a)**m*
tan(e + f*x)**2,x)*b*c*d + int((tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*
c**3 + 2*int((tan(e + f*x)*b + a)**m*tan(e + f*x),x)*a*c*d + int((tan(e +
f*x)*b + a)**m*tan(e + f*x),x)*b*c**2
```

3.167 $\int (a+b \tan(e+fx))^m (c+d \tan(e+fx)) (A+B \tan(e$

Optimal result	1849
Mathematica [A] (verified)	1850
Rubi [A] (verified)	1850
Maple [F]	1854
Fricas [F]	1854
Sympy [F]	1854
Maxima [F]	1855
Giac [F]	1855
Mupad [F(-1)]	1856
Reduce [F]	1856

Optimal result

Integrand size = 43, antiderivative size = 247

$$\int (a+b \tan(e+fx))^m (c+d \tan(e+fx)) (A+B \tan(e+fx) + C \tan^2(e+fx)) dx$$

$$= -\frac{(aCd - b(cC + Bd)(2+m))(a+b \tan(e+fx))^{1+m}}{b^2 f(1+m)(2+m)}$$

$$+ \frac{(A - iB - C)(c - id) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right) (a+b \tan(e+fx))^{1+m}}{2(ia+b)f(1+m)}$$

$$- \frac{(A + iB - C)(c + id) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a+ib}\right) (a+b \tan(e+fx))^{1+m}}{2(ia-b)f(1+m)}$$

$$+ \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{1+m}}{bf(2+m)}$$

output

```
-(C*a*d-b*(B*d+C*c)*(2+m))*(a+b*tan(f*x+e))^(1+m)/b^2/f/(1+m)/(2+m)+1/2*(A
-I*B-C)*(c-I*d)*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a-I*b))*(a+b*ta
n(f*x+e))^(1+m)/(I*a+b)/f/(1+m)-1/2*(A+I*B-C)*(c+I*d)*hypergeom([1, 1+m],[
2+m],(a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a-b)/f/(1+m)+C*d*
tan(f*x+e)*(a+b*tan(f*x+e))^(1+m)/b/f/(2+m)
```

Mathematica [A] (verified)

Time = 2.10 (sec) , antiderivative size = 202, normalized size of antiderivative = 0.82

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{(a + b \tan(e + fx))^{1+m} \left(\frac{-2aCd + 2b(cC + Bd)(2+m)}{b(1+m)} - \frac{ib(A - iB - C)(c - id)(2+m) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right)}{(a-ib)(1+m)} \right)}{2bf(2+m)}$$

input

```
Integrate[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x]
+ C*Tan[e + f*x]^2),x]
```

output

```
((a + b*Tan[e + f*x])^(1 + m)*((-2*a*C*d + 2*b*(c*C + B*d)*(2 + m))/(b*(1
+ m)) - (I*b*(A - I*B - C)*(c - I*d)*(2 + m)*Hypergeometric2F1[1, 1 + m, 2
+ m, (a + b*Tan[e + f*x])/(a - I*b)])/((a - I*b)*(1 + m)) + (I*b*(A + I*B
- C)*(c + I*d)*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e +
f*x])/(a + I*b)])/((a + I*b)*(1 + m)) + 2*C*d*Tan[e + f*x]))/(2*b*f*(2 + m
))
```

Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {3042, 4120, 3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + d \tan(e + fx))(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (c + d \tan(e + fx))(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4120$$

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \int (a+b \tan(e+fx))^m ((aCd - b(cC + Bd)(m+2)) \tan^2(e+fx) - b(Bc + (A-C)d)(m+2) \tan(e+fx) + a)}{b(m+2)}$$

↓ 3042

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \int (a+b \tan(e+fx))^m ((aCd - b(cC + Bd)(m+2)) \tan(e+fx)^2 - b(Bc + (A-C)d)(m+2) \tan(e+fx) + a)}{b(m+2)}$$

↓ 4113

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \int (a+b \tan(e+fx))^m (-b(Ac - Cc - Bd)(m+2) - b(Bc + (A-C)d) \tan(e+fx)(m+2)) dx + \frac{(aCd - b(m+2)(A-C)d)}{b(m+2)}}{b(m+2)}$$

↓ 3042

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \int (a+b \tan(e+fx))^m (-b(Ac - Cc - Bd)(m+2) - b(Bc + (A-C)d) \tan(e+fx)(m+2)) dx + \frac{(aCd - b(m+2)(A-C)d)}{b(m+2)}}{b(m+2)}$$

↓ 4022

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{1}{2}b(m+2)(c+id)(A+iB-C) \int (1-i \tan(e+fx))(a+b \tan(e+fx))^m dx - \frac{1}{2}b(m+2)(c-id)(A-iB-C)}{b(m+2)}$$

↓ 3042

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{1}{2}b(m+2)(c+id)(A+iB-C) \int (1-i \tan(e+fx))(a+b \tan(e+fx))^m dx - \frac{1}{2}b(m+2)(c-id)(A-iB-C)}{b(m+2)}$$

↓ 4020

$$\frac{\frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \frac{ib(m+2)(c-id)(A-iB-C)}{2f} \int -\frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e+fx)) + \frac{ib(m+2)(c+id)(A+iB-C)}{2f} \int -\frac{(a+b \tan(e+fx))^m}{i \tan(e+fx)+1} d(-i \tan(e+fx)) + (a)}{b(m+2)}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \\ & \frac{ib(m+2)(c-id)(A-iB-C) \int \frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e+fx))}{2f} - \frac{ib(m+2)(c+id)(A+iB-C) \int \frac{(a+b \tan(e+fx))^m}{i \tan(e+fx)+1} d(-i \tan(e+fx))}{2f} + \frac{(aCd-b)}{b(m+2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 78 \\ & \frac{Cd \tan(e+fx)(a+b \tan(e+fx))^{m+1}}{bf(m+2)} - \\ & \frac{ib(m+2)(c-id)(A-iB-C)(a+b \tan(e+fx))^{m+1} \operatorname{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a-ib}\right)}{2f(m+1)(a-ib)} - \frac{ib(m+2)(c+id)(A+iB-C)(a+b \tan(e+fx))^{m+1}}{b(m+2)} \end{aligned}$$

input

```
Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
(C*d*Tan[e + f*x]*(a + b*Tan[e + f*x])^(1 + m))/(b*f*(2 + m)) - (((a*C*d - b*(c*C + B*d)*(2 + m))*(a + b*Tan[e + f*x])^(1 + m))/(b*f*(1 + m)) + ((I/2)*b*(A - I*B - C)*(c - I*d)*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a - I*b)*f*(1 + m)) - ((I/2)*b*(A + I*B - C)*(c + I*d)*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a + I*b)*f*(1 + m)))/(b*(2 + m))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 78

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

Maple [F]

$$\int (a + b \tan (fx + e))^m (c + d \tan (fx + e)) (A + B \tan (fx + e) + C \tan (fx + e)^2) dx$$

input `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

Fricas [F]

$$\begin{aligned} & \int (a + b \tan (e + fx))^m (c + d \tan (e + fx)) (A + B \tan (e + fx) + C \tan^2 (e + fx)) dx \\ & = \int (C \tan (fx + e)^2 + B \tan (fx + e) + A) (d \tan (fx + e) + c) (b \tan (fx + e) + a)^m dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")`

output `integral((C*d*tan(f*x + e)^3 + (C*c + B*d)*tan(f*x + e)^2 + A*c + (B*c + A*d)*tan(f*x + e))*(b*tan(f*x + e) + a)^m, x)`

Sympy [F]

$$\begin{aligned} & \int (a + b \tan (e + fx))^m (c + d \tan (e + fx)) (A + B \tan (e + fx) + C \tan^2 (e + fx)) dx \\ & = \int (a + b \tan (e + fx))^m (c + d \tan (e + fx)) (A + B \tan (e + fx) + C \tan^2 (e + fx)) dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))**m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output

```
Integral((a + b*tan(e + f*x))**m*(c + d*tan(e + f*x))*(A + B*tan(e + f*x)
+ C*tan(e + f*x)**2), x)
```

Maxima [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) (d \tan(fx + e) + c) (b \tan(fx + e) + a)^m dx$$

input

```
integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x, algorithm="maxima")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)*(b*
tan(f*x + e) + a)^m, x)
```

Giac [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan(fx + e)^2 + B \tan(fx + e) + A) (d \tan(fx + e) + c) (b \tan(fx + e) + a)^m dx$$

input

```
integrate((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)
^2),x, algorithm="giac")
```

output

```
integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)*(b*
tan(f*x + e) + a)^m, x)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) (C \tan(e + fx)^2 + B \tan(e + fx) + A) dx$$

input `int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `int((a + b*tan(e + f*x))^m*(c + d*tan(e + f*x))*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\int (a + b \tan(e + fx))^m (c + d \tan(e + fx)) (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \left(\int (\tan(fx + e) b + a)^m dx \right) ac + \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^3 dx \right) cd$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) bd$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e)^2 dx \right) c^2$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) ad$$

$$+ \left(\int (\tan(fx + e) b + a)^m \tan(fx + e) dx \right) bc$$

input `int((a+b*tan(f*x+e))^m*(c+d*tan(f*x+e))*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output

```
int((tan(e + f*x)*b + a)**m,x)*a*c + int((tan(e + f*x)*b + a)**m*tan(e + f
*x)**3,x)*c*d + int((tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*b*d + int((
tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*c**2 + int((tan(e + f*x)*b + a)*
*m*tan(e + f*x),x)*a*d + int((tan(e + f*x)*b + a)**m*tan(e + f*x),x)*b*c
```

3.168 $\int (a+b \tan(e+fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$

Optimal result	1858
Mathematica [A] (verified)	1859
Rubi [A] (verified)	1859
Maple [F]	1862
Fricas [F]	1862
Sympy [F]	1863
Maxima [F]	1863
Giac [F]	1863
Mupad [F(-1)]	1864
Reduce [F]	1864

Optimal result

Integrand size = 33, antiderivative size = 178

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{C(a + b \tan(e + fx))^{1+m}}{bf(1 + m)}$$

$$+ \frac{(A - iB - C) \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a - ib}\right) (a + b \tan(e + fx))^{1+m}}{2(ia + b)f(1 + m)}$$

$$+ \frac{(iA - B - iC) \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a + ib}\right) (a + b \tan(e + fx))^{1+m}}{2(a + ib)f(1 + m)}$$

output

```
C*(a+b*tan(f*x+e))^(1+m)/b/f/(1+m)+1/2*(A-I*B-C)*hypergeom([1, 1+m], [2+m],
(a+b*tan(f*x+e))/(a-I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a+b)/f/(1+m)+1/2*(I*A-
B-I*C)*hypergeom([1, 1+m], [2+m], (a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))
^(1+m)/(a+I*b)/f/(1+m)
```

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.76

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \frac{\left(\frac{2C}{b} - \frac{i(A-iB-C) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right)}{a-ib} \right) + \frac{i(A+iB-C) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a+ib}\right)}{a+ib}}{2f(1+m)}$$

input

```
Integrate[(a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
((2*C)/b - (I*(A - I*B - C)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)])/(a - I*b) + (I*(A + I*B - C)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)])/(a + I*b))*(a + b*Tan[e + f*x])^(1 + m))/(2*f*(1 + m))
```

Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan(e + fx)^2) dx$$

$$\downarrow 4113$$

$$\int (a + b \tan(e + fx))^m (A - C + B \tan(e + fx)) dx + \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)}$$

$$\downarrow 3042$$

$$\begin{aligned}
& \int (a + b \tan(e + fx))^m (A - C + B \tan(e + fx)) dx + \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)} \\
& \quad \downarrow 4022 \\
& \frac{1}{2}(A + iB - C) \int (1 - i \tan(e + fx))(a + b \tan(e + fx))^m dx + \frac{1}{2}(A - iB - C) \int (i \tan(e + \\
& \quad fx) + 1)(a + b \tan(e + fx))^m dx + \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)} \\
& \quad \downarrow 3042 \\
& \frac{1}{2}(A + iB - C) \int (1 - i \tan(e + fx))(a + b \tan(e + fx))^m dx + \frac{1}{2}(A - iB - C) \int (i \tan(e + \\
& \quad fx) + 1)(a + b \tan(e + fx))^m dx + \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)} \\
& \quad \downarrow 4020 \\
& \frac{i(A - iB - C) \int -\frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e + fx))}{2f} - \\
& \frac{i(A + iB - C) \int -\frac{(a+b \tan(e+fx))^m}{i \tan(e+fx)+1} d(-i \tan(e + fx))}{2f} + \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)} \\
& \quad \downarrow 25 \\
& -\frac{i(A - iB - C) \int \frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e + fx))}{2f} + \\
& \frac{i(A + iB - C) \int \frac{(a+b \tan(e+fx))^m}{i \tan(e+fx)+1} d(-i \tan(e + fx))}{2f} + \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)} \\
& \quad \downarrow 78 \\
& \frac{i(A - iB - C)(a + b \tan(e + fx))^{m+1} \operatorname{Hypergeometric2F1}\left(1, m + 1, m + 2, \frac{a+b \tan(e+fx)}{a-ib}\right)}{2f(m+1)(a-ib)} + \\
& \frac{i(A + iB - C)(a + b \tan(e + fx))^{m+1} \operatorname{Hypergeometric2F1}\left(1, m + 1, m + 2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)} + \\
& \frac{C(a + b \tan(e + fx))^{m+1}}{bf(m+1)}
\end{aligned}$$

input

```
Int[(a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2), x]
```

output

```
(C*(a + b*Tan[e + f*x])^(1 + m))/(b*f*(1 + m)) - ((I/2)*(A - I*B - C)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a - I*b)*f*(1 + m)) + ((I/2)*(A + I*B - C)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a + I*b)*f*(1 + m))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4020

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4113

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

Maple [F]

$$\int (a + b \tan(fx + e))^m (A + B \tan(fx + e) + C \tan^2(fx + e))^2 dx$$

input

```
int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

output

```
int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)
```

Fricas [F]

$$\begin{aligned} & \int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m dx \end{aligned}$$

input

```
integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="fricas")
```

output

```
integral((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m, x)
```

Sympy [F]

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

input `integrate((a+b*tan(f*x+e))**m*(A+B*tan(f*x+e)+C*tan(f*x+e)**2),x)`

output `Integral((a + b*tan(e + f*x))**m*(A + B*tan(e + f*x) + C*tan(e + f*x)**2), x)`

Maxima [F]

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m, x)`

Giac [F]

$$\int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx$$

$$= \int (C \tan^2(fx + e) + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \int (a + b \tan(e + fx))^m (C \tan^2(e + fx) + B \tan(e + fx) + A) dx \end{aligned}$$

input `int((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2),x)`

output `int((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2), x)`

Reduce [F]

$$\begin{aligned} & \int (a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx)) dx \\ &= \left(\int (\tan(fx + e)b + a)^m dx \right) a + \left(\int (\tan(fx + e)b + a)^m \tan(fx + e)^2 dx \right) c \\ & \quad + \left(\int (\tan(fx + e)b + a)^m \tan(fx + e) dx \right) b \end{aligned}$$

input `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2),x)`

output `int((tan(e + f*x)*b + a)**m,x)*a + int((tan(e + f*x)*b + a)**m*tan(e + f*x)**2,x)*c + int((tan(e + f*x)*b + a)**m*tan(e + f*x),x)*b`

3.169
$$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx$$

Optimal result	1865
Mathematica [A] (verified)	1866
Rubi [A] (verified)	1866
Maple [F]	1870
Fricas [F]	1870
Sympy [F]	1871
Maxima [F]	1871
Giac [F]	1872
Mupad [F(-1)]	1872
Reduce [F]	1873

Optimal result

Integrand size = 45, antiderivative size = 258

$$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{c+d \tan(e+fx)} dx =$$

$$\frac{(iA+B-iC) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right) (a+b \tan(e+fx))^{1+m}}{2(a-ib)(c-id)f(1+m)}$$

$$- \frac{(A+iB-C) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a+ib}\right) (a+b \tan(e+fx))^{1+m}}{2(ia-b)(c+id)f(1+m)}$$

$$+ \frac{(c^2C-Bcd+Ad^2) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, -\frac{d(a+b \tan(e+fx))}{bc-ad}\right) (a+b \tan(e+fx))^{1+m}}{(bc-ad)(c^2+d^2)f(1+m)}$$

output

```
-1/2*(I*A+B-I*C)*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a-I*b))*(a+b*tan(f*x+e))^(1+m)/(a-I*b)/(c-I*d)/f/(1+m)-1/2*(A+I*B-C)*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a-b)/(c+I*d)/f/(1+m)+(A*d^2-B*c*d+C*c^2)*hypergeom([1, 1+m],[2+m],-d*(a+b*tan(f*x+e))/(-a*d+b*c))*(a+b*tan(f*x+e))^(1+m)/(-a*d+b*c)/(c^2+d^2)/f/(1+m)
```

Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.79

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \frac{\left(\frac{(A - iB - C)(-ic + d) \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a - ib}\right)}{a - ib} + \frac{(A + iB - C)(ic + d) \operatorname{Hypergeometric2F1}\left(1, 1 + m, 2 + m, \frac{a + b \tan(e + fx)}{a + ib}\right)}{a + ib} \right)}{2(c^2 + d^2) f(1 + m)}$$

input

```
Integrate[((a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x]),x]
```

output

```
((((A - I*B - C)*((-I)*c + d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)])/(a - I*b) + ((A + I*B - C)*(I*c + d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)])/(a + I*b) + (2*(c^2 * C - B*c*d + A*d^2)*Hypergeometric2F1[1, 1 + m, 2 + m, (d*(a + b*Tan[e + f*x]))/(-b*c + a*d)]/(b*c - a*d))*(a + b*Tan[e + f*x])^(1 + m))/(2*(c^2 + d^2)*f*(1 + m))
```

Rubi [A] (verified)

Time = 1.80 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4136, 3042, 4022, 3042, 4020, 25, 78, 4117, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan(e + fx)^2)}{c + d \tan(e + fx)} dx$$

↓ 4136

$$\begin{aligned}
 & \frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan^2(e+fx)+1)}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \\
 & \frac{\int (a + b \tan(e + fx))^m (Ac - Cc + Bd + (Bc - (A - C)d) \tan(e + fx)) dx}{c^2 + d^2} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{\int (a + b \tan(e + fx))^m (Ac - Cc + Bd + (Bc - (A - C)d) \tan(e + fx)) dx}{c^2 + d^2} + \\
 & \frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan(e+fx)^2+1)}{c+d \tan(e+fx)} dx}{c^2 + d^2} \\
 & \qquad \qquad \qquad \downarrow \text{4022} \\
 & \frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan(e+fx)^2+1)}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \\
 & \frac{\frac{1}{2}(c - id)(A + iB - C) \int (1 - i \tan(e + fx))(a + b \tan(e + fx))^m dx + \frac{1}{2}(c + id)(A - iB - C) \int (i \tan(e + fx) + 1) dx}{c^2 + d^2}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan(e+fx)^2+1)}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \\
 & \frac{\frac{1}{2}(c - id)(A + iB - C) \int (1 - i \tan(e + fx))(a + b \tan(e + fx))^m dx + \frac{1}{2}(c + id)(A - iB - C) \int (i \tan(e + fx) + 1) dx}{c^2 + d^2}} \\
 & \qquad \qquad \qquad \downarrow \text{4020} \\
 & \frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan(e+fx)^2+1)}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \\
 & \frac{\frac{i(c+id)(A-iB-C) \int -\frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e+fx))}{2f} - \frac{i(c-id)(A+iB-C) \int -\frac{(a+b \tan(e+fx))^m}{i \tan(e+fx)+1} d(-i \tan(e+fx))}{2f}}{c^2 + d^2}} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & \frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan(e+fx)^2+1)}{c+d \tan(e+fx)} dx}{c^2 + d^2} + \\
 & \frac{\frac{i(c-id)(A+iB-C) \int \frac{(a+b \tan(e+fx))^m}{i \tan(e+fx)+1} d(-i \tan(e+fx))}{2f} - \frac{i(c+id)(A-iB-C) \int \frac{(a+b \tan(e+fx))^m}{1-i \tan(e+fx)} d(i \tan(e+fx))}{2f}}{c^2 + d^2}} \\
 & \qquad \qquad \qquad \downarrow \text{78}
 \end{aligned}$$

$$\frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m (\tan(e+fx)^2+1) dx}{c+d \tan(e+fx)}}{c^2 + d^2} + \frac{i(c-id)(A+iB-C)(a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)} - \frac{i(c+id)(A-iB-C)(a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)}}{c^2 + d^2}$$

↓ 4117

$$\frac{(Ad^2 - Bcd + c^2C) \int \frac{(a+b \tan(e+fx))^m d \tan(e+fx)}{c+d \tan(e+fx)}}{f(c^2 + d^2)} + \frac{i(c-id)(A+iB-C)(a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)} - \frac{i(c+id)(A-iB-C)(a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)}}{c^2 + d^2}$$

↓ 78

$$\frac{(Ad^2 - Bcd + c^2C) (a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, -\frac{d(a+b \tan(e+fx))}{bc-ad}\right)}{f(m+1)(c^2 + d^2)(bc - ad)} + \frac{i(c-id)(A+iB-C)(a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)} - \frac{i(c+id)(A-iB-C)(a+b \tan(e+fx))^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{a+b \tan(e+fx)}{a+ib}\right)}{2f(m+1)(a+ib)}}{c^2 + d^2}$$

input

```
Int[((a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]),x]
```

output

```
((c^2*C - B*c*d + A*d^2)*Hypergeometric2F1[1, 1 + m, 2 + m, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d))]*(a + b*Tan[e + f*x])^(1 + m))/((b*c - a*d)*(c^2 + d^2)*f*(1 + m)) + (((-1/2*I)*(A - I*B - C)*(c + I*d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a - I*b)*f*(1 + m)) + ((I/2)*(A + I*B - C)*(c - I*d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a + I*b)*f*(1 + m)))/(c^2 + d^2)
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 78 $\text{Int}[(\text{a}_) + (\text{b}_) * (\text{x}_)^{(\text{m}_)} * ((\text{c}_) + (\text{d}_) * (\text{x}_)^{(\text{n}_)}), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{b} * \text{c} - \text{a} * \text{d})^{\text{n}} * ((\text{a} + \text{b} * \text{x})^{(\text{m} + 1)} / (\text{b}^{(\text{n} + 1)} * (\text{m} + 1))) * \text{Hypergeometric2F1}[-\text{n}, \text{m} + 1, \text{m} + 2, (-\text{d}) * ((\text{a} + \text{b} * \text{x}) / (\text{b} * \text{c} - \text{a} * \text{d}))], \text{x}] /;$ $\text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{m}\}, \text{x}]$ && $\text{!IntegerQ}[\text{m}]$ && $\text{IntegerQ}[\text{n}]$
- rule 3042 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /;$ $\text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$
- rule 4020 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{(\text{m}_)} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]), \text{x_Symbol}] \rightarrow \text{Simp}[\text{c} * (\text{d} / \text{f}) \quad \text{Subst}[\text{Int}[(\text{a} + (\text{b} / \text{d}) * \text{x})^{\text{m}} / (\text{d}^2 + \text{c} * \text{x}), \text{x}], \text{x}, \text{d} * \text{Tan}[\text{e} + \text{f} * \text{x}], \text{x}] /;$ $\text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}]$ && $\text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0]$ && $\text{NeQ}[\text{a}^2 + \text{b}^2, 0]$ && $\text{EqQ}[\text{c}^2 + \text{d}^2, 0]$
- rule 4022 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{(\text{m}_)} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{I} * \text{d}) / 2 \quad \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 - \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] + \text{Simp}[(\text{c} - \text{I} * \text{d}) / 2 \quad \text{Int}[(\text{a} + \text{b} * \text{Tan}[\text{e} + \text{f} * \text{x}])^{\text{m}} * (1 + \text{I} * \text{Tan}[\text{e} + \text{f} * \text{x}]), \text{x}], \text{x}] /;$ $\text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}]$ && $\text{NeQ}[\text{b} * \text{c} - \text{a} * \text{d}, 0]$ && $\text{NeQ}[\text{a}^2 + \text{b}^2, 0]$ && $\text{NeQ}[\text{c}^2 + \text{d}^2, 0]$ && $\text{!IntegerQ}[\text{m}]$
- rule 4117 $\text{Int}[(\text{a}_.) + (\text{b}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^{(\text{m}_)} * ((\text{c}_.) + (\text{d}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)])^{(\text{n}_)} * ((\text{A}_) + (\text{C}_.) * \tan[(\text{e}_.) + (\text{f}_.) * (\text{x}_)]^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{A} / \text{f} \quad \text{Subst}[\text{Int}[(\text{a} + \text{b} * \text{x})^{\text{m}} * (\text{c} + \text{d} * \text{x})^{\text{n}}, \text{x}], \text{x}, \text{Tan}[\text{e} + \text{f} * \text{x}], \text{x}] /;$ $\text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{A}, \text{C}, \text{m}, \text{n}\}, \text{x}]$ && $\text{EqQ}[\text{A}, \text{C}]$

rule 4136

```
Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

Maple [F]

$$\int \frac{(a + b \tan(fx + e))^m (A + B \tan(fx + e) + C \tan(fx + e)^2)}{c + d \tan(fx + e)} dx$$

input

```
int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)
```

output

```
int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)
```

Fricas [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{d \tan(fx + e) + c} dx$$

input

```
integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+
e)),x, algorithm="fricas")
```

output

```
integral((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d
*tan(f*x + e) + c), x)
```

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

input `integrate((a+b*tan(f*x+e))**m*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e)),x)`

output `Integral((a + b*tan(e + f*x))**m*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x)), x)`

Maxima [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{d \tan(fx + e) + c} dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d*tan(f*x + e) + c), x)`

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{d \tan(fx + e) + c} dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d*tan(f*x + e) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \int \frac{(a + b \tan(e + fx))^m (C \tan(e + fx)^2 + B \tan(e + fx) + A)}{c + d \tan(e + fx)} dx$$

input `int(((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x)),x)`

output `int(((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x)), x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{c + d \tan(e + fx)} dx$$

$$= \left(\int \frac{(\tan(fx + e)b + a)^m}{d \tan(fx + e) + c} dx \right) a + \left(\int \frac{(\tan(fx + e)b + a)^m \tan(fx + e)^2}{d \tan(fx + e) + c} dx \right) c$$

$$+ \left(\int \frac{(\tan(fx + e)b + a)^m \tan(fx + e)}{d \tan(fx + e) + c} dx \right) b$$

input `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e)),x)`

output `int((tan(e + f*x)*b + a)**m/(tan(e + f*x)*d + c),x)*a + int(((tan(e + f*x)*b + a)**m*tan(e + f*x)**2)/(tan(e + f*x)*d + c),x)*c + int(((tan(e + f*x)*b + a)**m*tan(e + f*x))/(tan(e + f*x)*d + c),x)*b`

3.170
$$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$$

Optimal result	1874
Mathematica [A] (verified)	1875
Rubi [A] (verified)	1875
Maple [F]	1880
Fricas [F]	1880
Sympy [F(-2)]	1880
Maxima [F]	1881
Giac [F]	1881
Mupad [F(-1)]	1882
Reduce [F]	1882

Optimal result

Integrand size = 45, antiderivative size = 403

$$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^2} dx$$

$$= \frac{(A-iB-C) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right) (a+b \tan(e+fx))^{1+m}}{2(ia+b)(c-id)^2 f(1+m)}$$

$$+ \frac{(iA-B-iC) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a+ib}\right) (a+b \tan(e+fx))^{1+m}}{2(a+ib)(c+id)^2 f(1+m)}$$

$$- \frac{(ad^2(2c(A-C)d-B(c^2-d^2))-b(Ad^2(c^2(2-m)-d^2m)-Bcd(c^2(1-m)-d^2(1+m)))-c^2C(c^2+d^2))}{(bc-ad)^2 (c^2+d^2)}$$

$$+ \frac{(c^2C-Bcd+Ad^2)(a+b \tan(e+fx))^{1+m}}{(bc-ad)(c^2+d^2)f(c+d \tan(e+fx))}$$

output

```
1/2*(A-I*B-C)*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a-I*b))*(a+b*tan(f*x+e))^(1+m)/(I*a+b)/(c-I*d)^2/f/(1+m)+1/2*(I*A-B-I*C)*hypergeom([1, 1+m],[2+m],(a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))^(1+m)/(a+I*b)/(c+I*d)^2/f/(1+m)-(a*d^2*(2*c*(A-C)*d-B*(c^2-d^2))-b*(A*d^2*(c^2*(2-m)-d^2*m)-B*c*d*(c^2*(1-m)-d^2*(1+m))-c^2*C*(c^2*m+d^2*(2+m)))*hypergeom([1, 1+m],[2+m],-d*(a+b*tan(f*x+e))/(-a*d+b*c))*(a+b*tan(f*x+e))^(1+m)/(-a*d+b*c)^2/(c^2+d^2)^2/f/(1+m)+(A*d^2-B*c*d+C*c^2)*(a+b*tan(f*x+e))^(1+m)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))
```

Mathematica [A] (verified)

Time = 6.02 (sec) , antiderivative size = 360, normalized size of antiderivative = 0.89

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \frac{(a + b \tan(e + fx))^{1+m} \left(-\frac{i \left(\frac{(A-iB-C)(c+id)^2(-bc+ad) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right)}{a-ib} + \frac{(A+iB-C)(c-id)^2(bc-ad) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a+ib}\right)}{a+ib} \right)}{(c^2+d^2)(1+m)} \right)}{(c^2+d^2)(1+m)}$$

input

```
Integrate[((a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/
/(c + d*Tan[e + f*x])^2,x]
```

output

```
((a + b*Tan[e + f*x])^(1 + m)*((( -I)*((A - I*B - C)*(c + I*d)^2*(-(b*c) +
a*d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)])/
(a - I*b) + ((A + I*B - C)*(c - I*d)^2*(b*c - a*d)*Hypergeometric2F1[1, 1
+ m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)])/(a + I*b)))/((c^2 + d^2)*(1 +
m)) + (2*(a*d^2*(2*c*(A - C)*d + B*(-c^2 + d^2)) + b*(A*c^2*d^2*(-2 + m)
+ c^4*C*m + A*d^4*m + c^2*C*d^2*(2 + m) - B*(c^3*d*(-1 + m) + c*d^3*(1 + m
))))*Hypergeometric2F1[1, 1 + m, 2 + m, (d*(a + b*Tan[e + f*x])/(-(b*c) +
a*d)])/((b*c - a*d)*(c^2 + d^2)*(1 + m)) - (2*(c^2*C - B*c*d + A*d^2))/(c
+ d*Tan[e + f*x]))/(2*(-(b*c) + a*d)*(c^2 + d^2)*f)
```

Rubi [A] (verified)

Time = 3.58 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4132, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 78, 4117, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^2} dx$$

↓ 4132

$$\frac{\int \frac{(a+b \tan(e+fx))^m (-b(Cc^2-Bdc+Ad^2)m \tan^2(e+fx)+(bc-ad)(Bc-(A-C)d) \tan(e+fx)+(cC-Bd)(ad-bc(m+1))-A(acd-b(c^2-d^2m)))}{c+d \tan(e+fx)} dx}{(c^2+d^2)(bc-ad)} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))}$$

↓ 3042

$$\frac{\int \frac{(a+b \tan(e+fx))^m (-b(Cc^2-Bdc+Ad^2)m \tan(e+fx)^2+(bc-ad)(Bc-(A-C)d) \tan(e+fx)+(cC-Bd)(ad-bc(m+1))-A(acd-b(c^2-d^2m)))}{c+d \tan(e+fx)} dx}{(c^2+d^2)(bc-ad)} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))}$$

↓ 4136

$$\frac{\int \frac{-(a+b \tan(e+fx))^m ((bc-ad)(Cc^2-2Bdc-Cd^2-A(c^2-d^2))+(bc-ad)(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)) dx}{c^2+d^2}}{(c^2+d^2)(bc-ad)} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))}$$

↓ 25

$$\frac{\int \frac{-(a+b \tan(e+fx))^m ((bc-ad)(Cc^2-2Bdc-Cd^2-A(c^2-d^2))+(bc-ad)(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)) dx}{c^2+d^2}}{(c^2+d^2)(bc-ad)} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))}$$

↓ 3042

$$\frac{\int \frac{-(a+b \tan(e+fx))^m ((bc-ad)(Cc^2-2Bdc-Cd^2-A(c^2-d^2))+(bc-ad)(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)) dx}{c^2+d^2}}{(c^2+d^2)(bc-ad)} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))}$$

↓ 4022

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} + \frac{(ad^2(2cd(A-C) - B(c^2 - d^2)) - b(Ac^2d^2(2-m) - Ad^4m - B(c^3d(1-m) - cd^3(m+1)) + c^4(-C)m - c^2Cd^2(m+2)))}{c^2 + d^2} \int \frac{(a + b \tan(e + fx))^m (\tan(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 3042

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} + \frac{(ad^2(2cd(A-C) - B(c^2 - d^2)) - b(Ac^2d^2(2-m) - Ad^4m - B(c^3d(1-m) - cd^3(m+1)) + c^4(-C)m - c^2Cd^2(m+2)))}{c^2 + d^2} \int \frac{(a + b \tan(e + fx))^m (\tan(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 4020

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} + \frac{(ad^2(2cd(A-C) - B(c^2 - d^2)) - b(Ac^2d^2(2-m) - Ad^4m - B(c^3d(1-m) - cd^3(m+1)) + c^4(-C)m - c^2Cd^2(m+2)))}{c^2 + d^2} \int \frac{(a + b \tan(e + fx))^m (\tan(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 25

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} + \frac{(ad^2(2cd(A-C) - B(c^2 - d^2)) - b(Ac^2d^2(2-m) - Ad^4m - B(c^3d(1-m) - cd^3(m+1)) + c^4(-C)m - c^2Cd^2(m+2)))}{c^2 + d^2} \int \frac{(a + b \tan(e + fx))^m (\tan(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 78

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} + \frac{(ad^2(2cd(A-C) - B(c^2 - d^2)) - b(Ac^2d^2(2-m) - Ad^4m - B(c^3d(1-m) - cd^3(m+1)) + c^4(-C)m - c^2Cd^2(m+2)))}{c^2 + d^2} \int \frac{(a + b \tan(e + fx))^m (\tan(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 4117

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} + \frac{(ad^2(2cd(A-C) - B(c^2 - d^2)) - b(Ac^2d^2(2-m) - Ad^4m - B(c^3d(1-m) - cd^3(m+1)) + c^4(-C)m - c^2Cd^2(m+2)))}{f(c^2 + d^2)} \int \frac{(a + b \tan(e + fx))^m (\tan(e + fx))}{c + d \tan(e + fx)} dx$$

↓ 78

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))} +$$

$$\frac{(a + b \tan(e + fx))^{m+1}(ad^2(2cd(A - C) - B(c^2 - d^2)) - b(Ad^2d^2(2 - m) - Ad^4m - B(c^3d(1 - m) - cd^3(m + 1)) + c^4(-C)m - c^2Cd^2(m + 2)))}{f(m + 1)(c^2 + d^2)(bc - ad)} \text{Hypergeometric2F1}$$

input

```
Int[((a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x])^2,x]
```

output

```
((c^2*C - B*c*d + A*d^2)*(a + b*Tan[e + f*x])^(1 + m))/((b*c - a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])) + (-(((a*d^2*(2*c*(A - C)*d - B*(c^2 - d^2)) - b*(A*c^2*d^2*(2 - m) - c^4*C*m - A*d^4*m - c^2*C*d^2*(2 + m) - B*(c^3*d*(1 - m) - c*d^3*(1 + m))))*Hypergeometric2F1[1, 1 + m, 2 + m, -((d*(a + b*Tan[e + f*x]))/(b*c - a*d))]*(a + b*Tan[e + f*x])^(1 + m))/((b*c - a*d)*(c^2 + d^2)*f*(1 + m)) - (((I/2)*(A - I*B - C)*(c + I*d)^2*(b*c - a*d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a - I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a - I*b)*f*(1 + m)) - ((I/2)*(A + I*B - C)*(c - I*d)^2*(b*c - a*d)*Hypergeometric2F1[1, 1 + m, 2 + m, (a + b*Tan[e + f*x])/(a + I*b)]*(a + b*Tan[e + f*x])^(1 + m))/((a + I*b)*f*(1 + m)))/(c^2 + d^2))/((b*c - a*d)*(c^2 + d^2))
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 78

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4020 $\text{Int}[\left((a_{.}) + (b_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)], x_{\text{Symbol}}] \rightarrow \text{Simp}[c(d/f) \text{Subst}[\text{Int}[(a + (b/d)x]^m/(d^2 + cx), x], x, d\tan[e + fx]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[\left((a_{.}) + (b_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)], x_{\text{Symbol}}] \rightarrow \text{Simp}[(c + I*d)/2 \text{Int}[(a + b\tan[e + fx])^m(1 - I\tan[e + fx]), x], x] + \text{Simp}[(c - I*d)/2 \text{Int}[(a + b\tan[e + fx])^m(1 + I\tan[e + fx]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!IntegerQ}[m]$

rule 4117 $\text{Int}[\left((a_{.}) + (b_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(n_{.})}\left((A_{.}) + (C_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[A/f \text{Subst}[\text{Int}[(a + b*x)^m(c + d*x)^n, x], x, \tan[e + fx]], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, C, m, n\}, x\} \&\& \text{EqQ}[A, C]$

rule 4132 $\text{Int}[\left((a_{.}) + (b_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(m_{.})}\left((c_{.}) + (d_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(n_{.})}\left((A_{.}) + (B_{.})\tan[e_{.}] + (f_{.})(x_{.})\right) + (C_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(A*b^2 - a*(b*B - a*C))*(a + b\tan[e + fx])^{(m+1)}*((c + d\tan[e + fx])^{(n+1)}/(f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{Int}[(a + b\tan[e + fx])^{(m+1)}*(c + d\tan[e + fx])^n \text{Simp}[A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) + (b*B - a*C)*(b*c*(m+1) + a*d*(n+1)) - (m+1)*(b*c - a*d)*(A*b - a*B - b*C)*\tan[e + fx] - d*(A*b^2 - a*(b*B - a*C))*(m+n+2)*\tan[e + fx]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{LtQ}[m, -1] \&\& \text{!(ILtQ}[n, -1] \&\& (\text{!IntegerQ}[m] || (\text{EqQ}[c, 0] \&\& \text{NeQ}[a, 0])))$

rule 4136 $\text{Int}[\left(\left((c_{.}) + (d_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^{(n_{.})}\left((A_{.}) + (B_{.})\tan[e_{.}] + (f_{.})(x_{.})\right) + (C_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)^2\right)/\left((a_{.}) + (b_{.})\tan[e_{.}] + (f_{.})(x_{.})\right)], x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(a^2 + b^2) \text{Int}[(c + d\tan[e + fx])^n \text{Simp}[b*B + a*(A - C) + (a*B - b*(A - C))*\tan[e + fx], x], x], x] + \text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{Int}[(c + d\tan[e + fx])^n((1 + \tan[e + fx]^2)/(a + b\tan[e + fx])), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!GtQ}[n, 0] \&\& \text{!LeQ}[n, -1]$

Maple [F]

$$\int \frac{(a + b \tan(fx + e))^m (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(c + d \tan(fx + e))^2} dx$$

input `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2, x)`

output `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2, x)`

Fricas [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{(d \tan(fx + e) + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="fricas")`

output `integral((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d^2*tan(f*x + e)^2 + 2*c*d*tan(f*x + e) + c^2), x)`

Sympy [F(-2)]

Exception generated.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

= Exception raised: HeuristicGCDFailed

input `integrate((a+b*tan(f*x+e))**m*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**2,x)`

output Exception raised: HeuristicGCDFailed >> no luck

Maxima [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{(d \tan(fx + e) + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="maxima")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d*tan(f*x + e) + c)^2, x)`

Giac [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{(d \tan(fx + e) + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d*tan(f*x + e) + c)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx$$

$$= \int \frac{(a + b \tan(e + fx))^m (C \tan^2(e + fx) + B \tan(e + fx) + A)}{(c + d \tan(e + fx))^2} dx$$

input `int(((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^2,x)`

output `int(((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^2, x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^2} dx = \text{too large to display}$$

input `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^2,x)`

output

```

((tan(e + f*x)*b + a)**m*a**2*d - (tan(e + f*x)*b + a)**m*a*b*c*m - (tan(e
+ f*x)*b + a)**m*b**2*d*m + (tan(e + f*x)*b + a)**m*b**2*d - int((tan(e +
f*x)*b + a)**m/(tan(e + f*x)**3*a*b*d**3 - tan(e + f*x)**3*b**2*c*d**2*m
+ tan(e + f*x)**2*a**2*d**3 - tan(e + f*x)**2*a*b*c*d**2*m + 2*tan(e + f*x)
)**2*a*b*c*d**2 - 2*tan(e + f*x)**2*b**2*c**2*d*m + 2*tan(e + f*x)*a**2*c*
d**2 - 2*tan(e + f*x)*a*b*c**2*d*m + tan(e + f*x)*a*b*c**2*d - tan(e + f*x)
)*b**2*c**3*m + a**2*c**2*d - a*b*c**3*m),x)*tan(e + f*x)*a**2*b**2*d**4*f
*m + int((tan(e + f*x)*b + a)**m/(tan(e + f*x)**3*a*b*d**3 - tan(e + f*x)*
**3*b**2*c*d**2*m + tan(e + f*x)**2*a**2*d**3 - tan(e + f*x)**2*a*b*c*d**2*
m + 2*tan(e + f*x)**2*a*b*c*d**2 - 2*tan(e + f*x)**2*b**2*c**2*d*m + 2*tan
(e + f*x)*a**2*c*d**2 - 2*tan(e + f*x)*a*b*c**2*d*m + tan(e + f*x)*a*b*c**
2*d - tan(e + f*x)*b**2*c**3*m + a**2*c**2*d - a*b*c**3*m),x)*tan(e + f*x)
*a**2*b**2*d**4*f + 2*int((tan(e + f*x)*b + a)**m/(tan(e + f*x)**3*a*b*d**
3 - tan(e + f*x)**3*b**2*c*d**2*m + tan(e + f*x)**2*a**2*d**3 - tan(e + f*
x)**2*a*b*c*d**2*m + 2*tan(e + f*x)**2*a*b*c*d**2 - 2*tan(e + f*x)**2*b**2
*c**2*d*m + 2*tan(e + f*x)*a**2*c*d**2 - 2*tan(e + f*x)*a*b*c**2*d*m + tan
(e + f*x)*a*b*c**2*d - tan(e + f*x)*b**2*c**3*m + a**2*c**2*d - a*b*c**3*m
),x)*tan(e + f*x)*a*b**3*c*d**3*f*m**2 - 2*int((tan(e + f*x)*b + a)**m/(ta
n(e + f*x)**3*a*b*d**3 - tan(e + f*x)**3*b**2*c*d**2*m + tan(e + f*x)**2*a
**2*d**3 - tan(e + f*x)**2*a*b*c*d**2*m + 2*tan(e + f*x)**2*a*b*c*d**2 ...

```


3.171
$$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$$

Optimal result	1884
Mathematica [B] (verified)	1885
Rubi [F]	1886
Maple [F]	1893
Fricas [F]	1894
Sympy [F]	1894
Maxima [F(-1)]	1895
Giac [F]	1895
Mupad [F(-1)]	1895
Reduce [F]	1896

Optimal result

Integrand size = 45, antiderivative size = 702

$$\int \frac{(a+b \tan(e+fx))^m (A+B \tan(e+fx)+C \tan^2(e+fx))}{(c+d \tan(e+fx))^3} dx$$

$$= \frac{(A-iB-C) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a-ib}\right) (a+b \tan(e+fx))^{1+m}}{2(ia+b)(c-id)^3 f(1+m)}$$

$$+ \frac{(A+iB-C) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, \frac{a+b \tan(e+fx)}{a+ib}\right) (a+b \tan(e+fx))^{1+m}}{2(a+ib)(ic-d)^3 f(1+m)}$$

$$+ \frac{(2a^2 d^3((A-C)d(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(B(6c^2d^2-c^4(2-m)-d^4m)+2c(A-C)d(c^2+d^2))+2ad^2(2c(A-C)d-B(c^2-d^2))-b(c^4C(1-m)+Ad^4(1-m)-Bc^3d(3-m)+Bcd^3(1+m)+c^2d^2))}{2(bc-ad)^2(c^2+d^2)^2 f(c+d \tan(e+fx))}$$

output

```

1/2*(A-I*B-C)*hypergeom([1, 1+m], [2+m], (a+b*tan(f*x+e))/(a-I*b))*(a+b*tan(
f*x+e))^(1+m)/(I*a+b)/(c-I*d)^3/f/(1+m)+1/2*(A+I*B-C)*hypergeom([1, 1+m], [
2+m], (a+b*tan(f*x+e))/(a+I*b))*(a+b*tan(f*x+e))^(1+m)/(a+I*b)/(I*c-d)^3/f/
(1+m)+1/2*(2*a^2*d^3*((A-C)*d*(3*c^2-d^2)-B*(c^3-3*c*d^2))-2*a*b*d^2*(B*(6
*c^2*d^2-c^4*(2-m)-d^4*m)+2*c*(A-C)*d*(c^2*(3-m)-d^2*(1+m)))-b^2*(A*d^2*(d
^4*(1-m)*m+2*c^2*d^2*(-m^2+3*m+1)-c^4*(m^2-5*m+6))+B*c*d*(d^4*m*(1+m)-2*c^
2*d^2*(-m^2+m+3)+c^4*(m^2-3*m+2))+c^2*C*(c^4*(1-m)*m+2*c^2*d^2*(-m^2-m+3)-
d^4*(m^2+3*m+2))))*hypergeom([1, 1+m], [2+m], -d*(a+b*tan(f*x+e))/(-a*d+b*c)
)*(a+b*tan(f*x+e))^(1+m)/(-a*d+b*c)^3/(c^2+d^2)^3/f/(1+m)+1/2*(A*d^2-B*c*d
+C*c^2)*(a+b*tan(f*x+e))^(1+m)/(-a*d+b*c)/(c^2+d^2)/f/(c+d*tan(f*x+e))^2-1
/2*(2*a*d^2*(2*c*(A-C)*d-B*(c^2-d^2))-b*(c^4*C*(1-m)+A*d^4*(1-m)-B*c^3*d*(
3-m)+B*c*d^3*(1+m)+c^2*d^2*(A*(5-m)-C*(3+m))))*(a+b*tan(f*x+e))^(1+m)/(-a*
d+b*c)^2/(c^2+d^2)^2/f/(c+d*tan(f*x+e))

```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2238 vs. $2(702) = 1404$.

Time = 6.20 (sec) , antiderivative size = 2238, normalized size of antiderivative = 3.19

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

= Result too large to show

input

```

Integrate[((a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))
/(c + d*Tan[e + f*x])^3,x]

```

output

```
-1/2*((A*d^2 - c*(-(c*C) + B*d))*(a + b*Tan[e + f*x])^(1 + m))/((-b*c) +
a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])^2 - ((((-(c*(2*d*(b*c - a*d)*(B*
c - (A - C)*d) - b*c*(c^2*C - B*c*d + A*d^2)*(1 - m))) + d^2*(A*(2*c*(b*c
- a*d) + b*d^2*(1 - m)) + (c*C - B*d)*(2*a*d - b*c*(1 + m))))*(a + b*Tan[e
+ f*x])^(1 + m))/((-b*c) + a*d)*(c^2 + d^2)*f*(c + d*Tan[e + f*x])) - (
-((((-(c*d*(-(b*c) + a*d)*(-2*c*(b*c - a*d)*(B*c - (A - C)*d) - b*d*(c^2*C
- B*c*d + A*d^2)*(1 - m) + d*(A*(2*c*(b*c - a*d) + b*d^2*(1 - m)) + (c*C -
B*d)*(2*a*d - b*c*(1 + m)))) - b*c^2*m*(-(c*(2*d*(b*c - a*d)*(B*c - (A -
C)*d) - b*c*(c^2*C - B*c*d + A*d^2)*(1 - m)) + d^2*(A*(2*c*(b*c - a*d) +
b*d^2*(1 - m)) + (c*C - B*d)*(2*a*d - b*c*(1 + m)))) + d^2*((2*d*(b*c - a
*d)*(B*c - (A - C)*d) - b*c*(c^2*C - B*c*d + A*d^2)*(1 - m))*(-(a*d) + b*c
*(1 + m)) + (-c*(-(b*c) + a*d)) - b*d^2*m*(A*(2*c*(b*c - a*d) + b*d^2*(1
- m)) + (c*C - B*d)*(2*a*d - b*c*(1 + m)))))*Hypergeometric2F1[1, 1 + m,
2 + m, (d*(a + b*Tan[e + f*x]))/((-b*c) + a*d)]*(a + b*Tan[e + f*x])^(1 +
m))/((-b*c) + a*d)*(c^2 + d^2)*f*(1 + m)) + (((I/2)*(d*(-(b*c) + a*d)*(-
2*c*(b*c - a*d)*(B*c - (A - C)*d) - b*d*(c^2*C - B*c*d + A*d^2)*(1 - m) +
d*(A*(2*c*(b*c - a*d) + b*d^2*(1 - m)) + (c*C - B*d)*(2*a*d - b*c*(1 + m)
)) + c*((2*d*(b*c - a*d)*(B*c - (A - C)*d) - b*c*(c^2*C - B*c*d + A*d^2)*(
1 - m))*(-(a*d) + b*c*(1 + m)) + (-c*(-(b*c) + a*d)) - b*d^2*m*(A*(2*c*(
b*c - a*d) + b*d^2*(1 - m)) + (c*C - B*d)*(2*a*d - b*c*(1 + m))) + b*m*...
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan(e + fx)^2)}{(c + d \tan(e + fx))^3} dx$$

↓ 4132

$$\frac{\int \frac{(a + b \tan(e + fx))^m (b(Cc^2 - Bdc + Ad^2)(1 - m) \tan^2(e + fx) + 2(bc - ad)(Bc - (A - C)d) \tan(e + fx) + A(b(1 - m)d^2 + 2c(bc - ad)) + (cC - Bd)(2ad - (c^2 + d^2)(bc - ad)))}{(c + d \tan(e + fx))^2}}{2(c^2 + d^2)(bc - ad)} \frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^m (b(Cc^2-Bdc+Ad^2)(1-m) \tan(e+fx)^2+2(bc-ad)(Bc-(A-C)d) \tan(e+fx)+A(b(1-m)d^2+2c(bc-ad))+(cC-Bd)(2ad-c+d \tan(e+fx))^2)}{2(c^2+d^2)(bc-ad)} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2}$$

↓ 4132

$$\int \frac{(a+b \tan(e+fx))^m (2(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)(bc-ad)^2-bm(2ad^2(2c(A-C)d-B(c^2-d^2))-b(C(1-m)c^4-Bd(3-m)c^3+d^2(A(5-m)-C(m+3))))}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2}$$

↓ 25

$$\int \frac{(a+b \tan(e+fx))^m (2(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)(bc-ad)^2-bm(2ad^2(2c(A-C)d-B(c^2-d^2))-b(C(1-m)c^4-Bd(3-m)c^3+d^2(A(5-m)-C(m+3))))}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2}$$

↓ 3042

$$\int \frac{(a+b \tan(e+fx))^m (2(2c(A-C)d-B(c^2-d^2)) \tan(e+fx)(bc-ad)^2-bm(2ad^2(2c(A-C)d-B(c^2-d^2))-b(C(1-m)c^4-Bd(3-m)c^3+d^2(A(5-m)-C(m+3))))}{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2}$$

↓ 4136

$$\int \frac{-2(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Cc^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} \frac{(Ad^2-Bcd+c^2C)(a+b \tan(e+fx))^{m+1}}{2f(c^2+d^2)(bc-ad)(c+d \tan(e+fx))^2}$$

↓ 27

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2} \tan(e+fx) dx$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3) - (bc-ad)^2 ((A-C)d(3c^2-d^2) - B(c^3-3cd^2))) \tan(e+fx) dx}{c^2+d^2} - (2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2} \tan(e+fx) dx$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2} \tan(e+fx) dx$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3) - (bc-ad)^2 ((A-C)d(3c^2-d^2) - B(c^3-3cd^2))) \tan(e+fx) dx}{c^2+d^2} - (2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2} \tan(e+fx) dx$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2} \tan(e+fx) dx$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Ce^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2 ((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^4m^2-4m^3+2m^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^4m^2-4m^3+2m^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Ce^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2 ((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^4m^2-4m^3+2m^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^4m^2-4m^3+2m^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Ce^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2 ((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^4m^2-4m^3+2m^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3) - (bc-ad)^2 ((A-C)d(3c^2-d^2) - B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3) - (bc-ad)^2 ((A-C)d(3c^2-d^2) - B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C)(a + b \tan(e + fx))^{m+1}}{2f(c^2 + d^2)(bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Ce^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2 ((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2)))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2)))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Ce^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2 ((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2)))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2)))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2 \int -(a+b \tan(e+fx))^m ((bc-ad)^2 (Ac^3-Ce^3+3Bdc^2-3Ad^2c+3Cd^2c-Bd^3)-(bc-ad)^2 ((A-C)d(3c^2-d^2)-B(c^3-3cd^2)) \tan(e+fx) dx}{c^2+d^2} - \frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2)))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{2f - (a + b \tan(e + fx))^m ((bc - ad)^2 (Ac^3 - Cc^3 + 3Bdc^2 - 3Ad^2c + 3Cd^2c - Bd^3) - (bc - ad)^2 ((A - C)d(3c^2 - d^2) - B(c^3 - 3cd^2)) \tan(e + fx) dx}{c^2 + d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

↓ 25

$$\frac{(2a^2d^3(d(A-C)(3c^2-d^2)-B(c^3-3cd^2))-2abd^2(2cd(A-C)(c^2(3-m)-d^2(m+1))+B(-(c^4(2-m))+6c^2d^2-d^4m))-b^2(Ad^2(-(c^4(m^2-5m+6))+2c^2d^2)))}{c^2+d^2}$$

$$\frac{(Ad^2 - Bcd + c^2C) (a + b \tan(e + fx))^{m+1}}{2f (c^2 + d^2) (bc - ad)(c + d \tan(e + fx))^2}$$

```
input Int[((a + b*Tan[e + f*x])^m*(A + B*Tan[e + f*x] + C*Tan[e + f*x]^2))/(c + d*Tan[e + f*x]^3,x]
```

```
output $Aborted
```

Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])], x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

Maple [F]

$$\int \frac{(a + b \tan(fx + e))^m (A + B \tan(fx + e) + C \tan(fx + e)^2)}{(c + d \tan(fx + e))^3} dx$$

input `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3, x)`

output `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3, x)`

Fricas [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

$$= \int \frac{(C \tan(fx + e)^2 + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{(d \tan(fx + e) + c)^3} dx$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="fricas")`

output `integral((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d^3*tan(f*x + e)^3 + 3*c*d^2*tan(f*x + e)^2 + 3*c^2*d*tan(f*x + e) + c^3), x)`

Sympy [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

$$= \int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx$$

input `integrate((a+b*tan(f*x+e))**m*(A+B*tan(f*x+e)+C*tan(f*x+e)**2)/(c+d*tan(f*x+e))**3,x)`

output `Integral((a + b*tan(e + f*x))**m*(A + B*tan(e + f*x) + C*tan(e + f*x)**2)/(c + d*tan(e + f*x))**3, x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{Timed out}$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="maxima")`

output `Timed out`

Giac [F]

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx \\ &= \int \frac{(C \tan^2(fx + e) + B \tan(fx + e) + A)(b \tan(fx + e) + a)^m}{(d \tan(fx + e) + c)^3} dx \end{aligned}$$

input `integrate((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((C*tan(f*x + e)^2 + B*tan(f*x + e) + A)*(b*tan(f*x + e) + a)^m/(d*tan(f*x + e) + c)^3, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx \\ &= \int \frac{(a + b \tan(e + fx))^m (C \tan^2(e + fx) + B \tan(e + fx) + A)}{(c + d \tan(e + fx))^3} dx \end{aligned}$$

input `int(((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^3,x)`

output `int(((a + b*tan(e + f*x))^m*(A + B*tan(e + f*x) + C*tan(e + f*x)^2))/(c + d*tan(e + f*x))^3, x)`

Reduce [F]

$$\int \frac{(a + b \tan(e + fx))^m (A + B \tan(e + fx) + C \tan^2(e + fx))}{(c + d \tan(e + fx))^3} dx = \text{too large to display}$$

input `int((a+b*tan(f*x+e))^m*(A+B*tan(f*x+e)+C*tan(f*x+e)^2)/(c+d*tan(f*x+e))^3,x)`

output `(2*(tan(e + f*x)*b + a)**m*a**2*d - (tan(e + f*x)*b + a)**m*a*b*c*m - (tan(e + f*x)*b + a)**m*b**2*d*m + 2*(tan(e + f*x)*b + a)**m*b**2*d + 2*int((tan(e + f*x)*b + a)**m/(tan(e + f*x)**4*b*d**3*m - 2*tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3*m - 2*tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2*m - 6*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2*m - 6*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d*m - 6*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d*m - 6*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3*m - 2*tan(e + f*x)*b*c**3 + a*c**3*m - 2*a*c**3),x)*tan(e + f*x)**2*a**3*d**4*f*m**2 - 4*int((tan(e + f*x)*b + a)**m/(tan(e + f*x)**4*b*d**3*m - 2*tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3*m - 2*tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2*m - 6*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2*m - 6*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d*m - 6*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d*m - 6*tan(e + f*x)*a*c**2*d + tan(e + f*x)*b*c**3*m - 2*tan(e + f*x)*b*c**3 + a*c**3*m - 2*a*c**3),x)*tan(e + f*x)**2*a**3*d**4*f*m - int((tan(e + f*x)*b + a)**m/(tan(e + f*x)**4*b*d**3*m - 2*tan(e + f*x)**4*b*d**3 + tan(e + f*x)**3*a*d**3*m - 2*tan(e + f*x)**3*a*d**3 + 3*tan(e + f*x)**3*b*c*d**2*m - 6*tan(e + f*x)**3*b*c*d**2 + 3*tan(e + f*x)**2*a*c*d**2*m - 6*tan(e + f*x)**2*a*c*d**2 + 3*tan(e + f*x)**2*b*c**2*d*m - 6*tan(e + f*x)**2*b*c**2*d + 3*tan(e + f*x)*a*c**2*d*m - 6*tan(e + f*x)*a*c**2*d + tan(e + f...`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 1897
4.2 Links to plain text integration problems used in this report for each CAS . 1915

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```



```
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
If[AppellFunctionQ[Head[expn]],
  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
If[Head[expn]===RootSum,
  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
If[Head[expn]===Integrate || Head[expn]===Int,
  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]
```

```
SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]
```

```
HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]
```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```



```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)=",type(result))
    print("type(optimal)=",type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file