

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.3-Tangent/222-4.3.10

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 9:21am

Contents

1	Introduction	4
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23
2	detailed summary tables of results	24
2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	45
3	Listing of integrals	48
3.1	$\int x^3 \tan(a + bx) dx$	51
3.2	$\int x^2 \tan(a + bx) dx$	59
3.3	$\int x \tan(a + bx) dx$	65
3.4	$\int \frac{\tan(a+bx)}{x} dx$	71
3.5	$\int \frac{\tan(a+bx)}{x^2} dx$	77
3.6	$\int x^3 \tan^2(a + bx) dx$	83
3.7	$\int x^2 \tan^2(a + bx) dx$	91

3.8	$\int x \tan^2(a + bx) dx$	98
3.9	$\int \frac{\tan^2(a+bx)}{x} dx$	104
3.10	$\int \frac{\tan^2(a+bx)}{x^2} dx$	110
3.11	$\int x^3 \tan^3(a + bx) dx$	116
3.12	$\int x^2 \tan^3(a + bx) dx$	128
3.13	$\int x \tan^3(a + bx) dx$	137
3.14	$\int \frac{\tan^3(a+bx)}{x} dx$	144
3.15	$\int \frac{\tan^3(a+bx)}{x^2} dx$	150
3.16	$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2 \sqrt{\tan(a+bx)} \right) dx$	156
3.17	$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx$	161
3.18	$\int \frac{(c+dx)^3}{a+ia \tan(e+fx)} dx$	166
3.19	$\int \frac{(c+dx)^2}{a+ia \tan(e+fx)} dx$	174
3.20	$\int \frac{c+dx}{a+ia \tan(e+fx)} dx$	181
3.21	$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))} dx$	187
3.22	$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))} dx$	194
3.23	$\int \frac{1}{(c+dx)^3(a+ia \tan(e+fx))} dx$	202
3.24	$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^2} dx$	211
3.25	$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^2} dx$	219
3.26	$\int \frac{c+dx}{(a+ia \tan(e+fx))^2} dx$	226
3.27	$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^2} dx$	232
3.28	$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^2} dx$	239
3.29	$\int \frac{(c+dx)^3}{(a+ia \tan(e+fx))^3} dx$	248
3.30	$\int \frac{(c+dx)^2}{(a+ia \tan(e+fx))^3} dx$	257
3.31	$\int \frac{c+dx}{(a+ia \tan(e+fx))^3} dx$	265
3.32	$\int \frac{1}{(c+dx)(a+ia \tan(e+fx))^3} dx$	272
3.33	$\int \frac{1}{(c+dx)^2(a+ia \tan(e+fx))^3} dx$	280
3.34	$\int (c+dx)^m (a + ia \tan(e + fx))^2 dx$	290
3.35	$\int (c+dx)^m (a + ia \tan(e + fx)) dx$	296
3.36	$\int \frac{(c+dx)^m}{a+ia \tan(e+fx)} dx$	301
3.37	$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^2} dx$	306
3.38	$\int \frac{(c+dx)^m}{(a+ia \tan(e+fx))^3} dx$	312
3.39	$\int (c+dx)^3 (a + b \tan(e + fx)) dx$	318
3.40	$\int (c+dx)^2 (a + b \tan(e + fx)) dx$	326
3.41	$\int (c+dx) (a + b \tan(e + fx)) dx$	333
3.42	$\int \frac{a+b \tan(e+fx)}{c+dx} dx$	339

3.43	$\int \frac{a+b\tan(e+fx)}{(c+dx)^2} dx$	344
3.44	$\int (c+dx)^3(a+b\tan(e+fx))^2 dx$	349
3.45	$\int (c+dx)^2(a+b\tan(e+fx))^2 dx$	358
3.46	$\int (c+dx)(a+b\tan(e+fx))^2 dx$	366
3.47	$\int \frac{(a+b\tan(e+fx))^2}{c+dx} dx$	372
3.48	$\int \frac{(a+b\tan(e+fx))^2}{(c+dx)^2} dx$	377
3.49	$\int (c+dx)^3(a+b\tan(e+fx))^3 dx$	382
3.50	$\int (c+dx)^2(a+b\tan(e+fx))^3 dx$	393
3.51	$\int (c+dx)(a+b\tan(e+fx))^3 dx$	403
3.52	$\int \frac{(a+b\tan(e+fx))^3}{c+dx} dx$	410
3.53	$\int \frac{(a+b\tan(e+fx))^3}{(c+dx)^2} dx$	416
3.54	$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$	422
3.55	$\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$	433
3.56	$\int \frac{c+dx}{a+b\tan(e+fx)} dx$	442
3.57	$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$	449
3.58	$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$	454
3.59	$\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$	459
3.60	$\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx$	470
3.61	$\int \frac{c+dx}{(a+b\tan(e+fx))^2} dx$	481
3.62	$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$	491
3.63	$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$	497
4	Appendix	503
4.1	Listing of Grading functions	503
4.2	Links to plain text integration problems used in this report for each CAS521	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Time and leaf size Performance	10
1.4	Performance based on number of rules Rubi used	12
1.5	Performance based on number of steps Rubi used	13
1.6	Solved integrals histogram based on leaf size of result	14
1.7	Solved integrals histogram based on CPU time used	15
1.8	Leaf size vs. CPU time used	16
1.9	list of integrals with no known antiderivative	17
1.10	List of integrals solved by CAS but has no known antiderivative	17
1.11	list of integrals solved by CAS but failed verification	17
1.12	Timing	18
1.13	Verification	18
1.14	Important notes about some of the results	19
1.15	Current tree layout of integration tests	22
1.16	Design of the test system	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [63]. This is test number [222].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 (63)	0.00 (0)
Fricas	100.00 (63)	0.00 (0)
Rubi	98.41 (62)	1.59 (1)
Maple	92.06 (58)	7.94 (5)
Maxima	77.78 (49)	22.22 (14)
Giac	55.56 (35)	44.44 (28)
Mupad	50.79 (32)	49.21 (31)
Sympy	44.44 (28)	55.56 (35)
Reduce	33.33 (21)	66.67 (42)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	69.841	0.000	0.000	30.159
Mathematica	58.730	12.698	0.000	28.571
Maple	42.857	20.635	0.000	36.508
Fricas	39.683	31.746	0.000	28.571
Giac	19.048	7.937	0.000	73.016
Sympy	15.873	0.000	0.000	84.127
Maxima	12.698	36.508	0.000	50.794
Mupad	0.000	22.222	0.000	77.778
Reduce	0.000	4.762	0.000	95.238

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Rubi	1	100.00	0.00	0.00
Maple	5	100.00	0.00	0.00
Maxima	14	35.71	0.00	64.29
Giac	28	92.86	0.00	7.14
Mupad	31	0.00	100.00	0.00
Sympy	35	100.00	0.00	0.00
Reduce	42	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.08
Reduce	0.16
Maple	0.54
Rubi	0.64
Sympy	0.96
Maxima	1.64
Giac	3.48
Mathematica	4.68
Mupad	8.56

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	66.86	3.41	37.00	1.78
Mupad	91.44	1.30	22.00	1.11
Sympy	143.93	1.25	19.00	1.00
Rubi	175.84	1.03	134.00	1.00
Fricas	275.24	1.77	130.00	1.61
Giac	295.46	1.58	23.00	1.10
Mathematica	300.63	1.46	171.00	1.17
Maple	433.60	1.73	118.00	1.00
Maxima	910.49	15.56	424.00	4.80

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{4, 5, 9, 10, 14, 15, 34, 35, 42, 43, 47, 48, 52, 53, 57, 58, 62, 63}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {11, 13, 44, 45, 49, 50, 54, 59, 60, 61}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A `time limit` of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	25
2.2	Detailed conclusion table per each integral for all CAS systems	29
2.3	Detailed conclusion table specific for Rubi results	45

2.1 List of integrals sorted by grade for each CAS

Rubi	25
Mma	25
Maple	26
Fricas	26
Maxima	26
Giac	27
Mupad	27
Sympy	27
Reduce	28

Rubi

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

B grade { }

C grade { }

F normal fail { 17 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 6, 8, 11, 12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 40, 41, 46, 51, 54, 55, 56, 60 }

B grade { 7, 39, 44, 45, 49, 50, 59, 61 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 41, 46 }

B grade { 39, 40, 44, 45, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

C grade { }

F normal fail { 16, 17, 36, 37, 38 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 8, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 46, 50, 51 }

B grade { 1, 2, 3, 6, 7, 11, 12, 13, 39, 40, 41, 44, 45, 49, 54, 55, 56, 59, 60, 61 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 21, 22, 23, 27, 28, 32, 33, 41 }

B grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 39, 40, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

C grade { }

F normal fail { 16, 17, 36, 37, 38 }

F(-1) timeout fail { }

F(-2) exception fail { 18, 19, 20, 24, 25, 26, 29, 30, 31 }

Giac**A grade** { 18, 19, 20, 21, 24, 25, 26, 27, 29, 30, 31, 32 }**B grade** { 8, 22, 23, 28, 33 }**C grade** { }**F normal fail** { 1, 2, 3, 6, 7, 11, 12, 13, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }**F(-1) timeout fail** { }**F(-2) exception fail** { 16, 17 }**Mupad****A grade** { }**B grade** { 3, 8, 16, 17, 18, 19, 20, 24, 25, 26, 29, 30, 31, 41 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { 1, 2, 6, 7, 11, 12, 13, 21, 22, 23, 27, 28, 32, 33, 36, 37, 38, 39, 40, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }**F(-2) exception fail** { }**Sympy****A grade** { 8, 18, 19, 20, 24, 25, 26, 29, 30, 31 }**B grade** { }**C grade** { }**F normal fail** { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 21, 22, 23, 27, 28, 32, 33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }**F(-1) timeout fail** { }**F(-2) exception fail** { }

Reduce

A grade { }

B grade { 8, 16, 17 }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32,
33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	133	106	125	243	286	0	0	12	0
N.S.	1	1.25	1.00	1.18	2.29	2.70	0.00	0.00	0.11	0.00
time (sec)	N/A	0.601	0.009	0.284	0.152	0.088	0.000	0.000	0.147	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	96	77	103	163	200	0	0	12	0
N.S.	1	1.25	1.00	1.34	2.12	2.60	0.00	0.00	0.16	0.00
time (sec)	N/A	0.466	0.007	0.250	0.160	0.089	0.000	0.000	0.150	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	61	54	78	92	122	0	0	10	129
N.S.	1	1.13	1.00	1.44	1.70	2.26	0.00	0.00	0.19	2.39
time (sec)	N/A	0.337	0.005	0.281	0.161	0.080	0.000	0.000	0.149	8.481

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.20	1.20
time (sec)	N/A	0.183	1.357	0.121	0.284	0.070	0.263	0.189	0.167	8.055

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20	1.20
time (sec)	N/A	0.184	2.245	0.130	0.280	0.067	0.225	0.354	0.149	7.781

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	122	115	133	639	225	0	0	37	0
N.S.	1	1.24	1.17	1.36	6.52	2.30	0.00	0.00	0.38	0.00
time (sec)	N/A	0.596	0.562	0.330	0.240	0.084	0.000	0.000	0.155	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	87	160	108	257	144	0	0	35	0
N.S.	1	1.19	2.19	1.48	3.52	1.97	0.00	0.00	0.48	0.00
time (sec)	N/A	0.431	4.548	0.307	0.236	0.078	0.000	0.000	0.148	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	43	34	214	38	41	162	37	35
N.S.	1	1.00	1.43	1.13	7.13	1.27	1.37	5.40	1.23	1.17
time (sec)	N/A	0.259	0.141	0.262	0.146	0.074	0.168	0.397	0.150	0.118

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	243	14	10	14	14	14
N.S.	1	1.00	1.17	1.00	20.25	1.17	0.83	1.17	1.17	1.17
time (sec)	N/A	0.197	2.884	0.204	0.304	0.068	0.289	0.563	0.165	8.716

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	250	14	12	14	14	14
N.S.	1	1.00	1.17	1.00	20.83	1.17	1.00	1.17	1.17	1.17
time (sec)	N/A	0.190	2.106	0.215	0.325	0.079	0.221	0.362	0.150	8.336

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	205	245	370	251	1205	344	0	0	14	0
N.S.	1	1.20	1.80	1.22	5.88	1.68	0.00	0.00	0.07	0.00
time (sec)	N/A	1.353	6.482	0.241	0.341	0.086	0.000	0.000	0.148	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	149	172	180	736	240	0	0	14	0
N.S.	1	1.16	1.34	1.41	5.75	1.88	0.00	0.00	0.11	0.00
time (sec)	N/A	0.850	2.079	0.234	0.285	0.090	0.000	0.000	0.152	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	98	171	122	386	146	0	0	12	0
N.S.	1	1.09	1.90	1.36	4.29	1.62	0.00	0.00	0.13	0.00
time (sec)	N/A	0.483	4.333	0.208	0.253	0.077	0.000	0.000	0.157	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	534	14	10	14	14	14
N.S.	1	1.00	1.17	1.00	44.50	1.17	0.83	1.17	1.17	1.17
time (sec)	N/A	0.200	4.396	0.133	0.518	0.070	0.286	0.373	0.154	8.502

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	536	14	12	14	14	14
N.S.	1	1.00	1.17	1.00	44.67	1.17	1.00	1.17	1.17	1.17
time (sec)	N/A	0.201	2.313	0.139	0.638	0.064	0.244	0.305	0.159	8.302

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F(-2)	B	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	0	0	16	0	0	23	50
N.S.	1	1.00	1.00	0.00	0.00	0.89	0.00	0.00	1.28	2.78
time (sec)	N/A	3.012	1.564	0.000	0.000	0.076	0.000	0.000	0.155	9.191

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	F	F	A	F	F(-2)	B	B
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	17	0	17	0	0	15	0	0	14	45
N.S.	1	0.00	1.00	0.00	0.00	0.88	0.00	0.00	0.82	2.65
time (sec)	N/A	0.000	0.806	0.000	0.000	0.072	0.000	0.000	0.155	9.659

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	189	206	278	170	0	164	258	188	91	423
N.S.	1	1.09	1.47	0.90	0.00	0.87	1.37	0.99	0.48	2.24
time (sec)	N/A	0.717	0.714	0.634	0.000	0.074	0.199	0.163	0.171	8.924

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	147	178	108	0	105	165	119	65	241
N.S.	1	1.07	1.30	0.79	0.00	0.77	1.20	0.87	0.47	1.76
time (sec)	N/A	0.503	0.433	0.608	0.000	0.072	0.170	0.216	0.154	8.731

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	90	96	50	0	55	92	62	39	105
N.S.	1	1.07	1.14	0.60	0.00	0.65	1.10	0.74	0.46	1.25
time (sec)	N/A	0.328	0.636	0.519	0.000	0.069	0.142	0.162	0.158	8.735

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	161	152	166	65	112	51	0	138	32	0
N.S.	1	0.94	1.03	0.40	0.70	0.32	0.00	0.86	0.20	0.00
time (sec)	N/A	0.757	0.540	0.542	0.112	0.069	0.000	0.164	0.152	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	167	224	96	119	83	0	1013	61	0
N.S.	1	0.99	1.33	0.57	0.71	0.49	0.00	6.03	0.36	0.00
time (sec)	N/A	0.757	0.686	0.560	0.132	0.071	0.000	2.127	0.177	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	227	225	285	216	158	130	0	532	90	0
N.S.	1	0.99	1.26	0.95	0.70	0.57	0.00	2.34	0.40	0.00
time (sec)	N/A	0.982	0.914	0.612	0.229	0.088	0.000	0.241	0.155	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	270	270	473	283	0	267	665	368	129	289
N.S.	1	1.00	1.75	1.05	0.00	0.99	2.46	1.36	0.48	1.07
time (sec)	N/A	0.545	2.277	0.968	0.000	0.073	0.356	0.247	0.160	9.131

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	282	173	0	163	418	217	94	183
N.S.	1	1.00	1.40	0.86	0.00	0.81	2.07	1.07	0.47	0.91
time (sec)	N/A	0.445	1.451	0.943	0.000	0.076	0.274	0.241	0.154	8.262

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	151	130	82	0	79	226	101	59	103
N.S.	1	1.00	0.86	0.54	0.00	0.52	1.50	0.67	0.39	0.68
time (sec)	N/A	0.392	0.964	0.698	0.000	0.074	0.224	0.194	0.165	8.069

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	305	305	211	114	194	84	0	404	59	0
N.S.	1	1.00	0.69	0.37	0.64	0.28	0.00	1.32	0.19	0.00
time (sec)	N/A	1.061	0.763	0.837	0.144	0.076	0.000	0.271	0.155	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	436	436	467	175	211	142	0	1967	107	0
N.S.	1	1.00	1.07	0.40	0.48	0.33	0.00	4.51	0.25	0.00
time (sec)	N/A	1.074	1.550	0.836	0.196	0.077	0.000	11.118	0.153	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	396	396	667	396	0	370	945	548	177	411
N.S.	1	1.00	1.68	1.00	0.00	0.93	2.39	1.38	0.45	1.04
time (sec)	N/A	0.691	3.378	1.106	0.000	0.086	0.525	0.337	0.165	9.271

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	294	405	238	0	221	588	315	130	263
N.S.	1	1.00	1.38	0.81	0.00	0.75	2.00	1.07	0.44	0.89
time (sec)	N/A	0.536	2.227	1.070	0.000	0.100	0.403	0.292	0.174	8.965

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	209	208	205	114	0	105	311	142	83	146
N.S.	1	1.00	0.98	0.55	0.00	0.50	1.49	0.68	0.40	0.70
time (sec)	N/A	0.480	1.343	0.832	0.000	0.070	0.313	0.274	0.158	9.147

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	449	449	336	163	276	117	0	810	84	0
N.S.	1	1.00	0.75	0.36	0.61	0.26	0.00	1.80	0.19	0.00
time (sec)	N/A	1.938	1.034	0.905	0.158	0.082	0.000	0.295	0.164	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	712	712	833	254	297	197	0	2915	152	0
N.S.	1	1.00	1.17	0.36	0.42	0.28	0.00	4.09	0.21	0.00
time (sec)	N/A	2.051	2.551	1.046	0.278	0.072	0.000	28.907	0.177	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	520	115	51	23	252	24
N.S.	1	1.00	1.09	0.91	22.61	5.00	2.22	1.00	10.96	1.04
time (sec)	N/A	0.226	40.379	0.263	0.844	0.072	3.417	0.410	0.171	8.611

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	79	34	29	21	69	22
N.S.	1	1.00	1.10	0.90	3.76	1.62	1.38	1.00	3.29	1.05
time (sec)	N/A	0.208	8.801	0.180	0.237	0.077	2.252	0.224	0.152	8.561

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	133	0	0	85	0	0	26	0
N.S.	1	1.00	1.36	0.00	0.00	0.87	0.00	0.00	0.27	0.00
time (sec)	N/A	0.333	1.589	0.000	0.000	0.074	0.000	0.000	0.157	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	192	0	0	144	0	0	36	0
N.S.	1	1.00	1.12	0.00	0.00	0.84	0.00	0.00	0.21	0.00
time (sec)	N/A	0.427	3.637	0.000	0.000	0.080	0.000	0.000	0.150	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	251	251	269	0	0	198	0	0	48	0
N.S.	1	1.00	1.07	0.00	0.00	0.79	0.00	0.00	0.19	0.00
time (sec)	N/A	0.496	12.416	0.000	0.000	0.088	0.000	0.000	0.202	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	152	342	500	672	500	0	0	119	0
N.S.	1	1.00	2.25	3.29	4.42	3.29	0.00	0.00	0.78	0.00
time (sec)	N/A	0.476	0.124	0.528	0.219	0.090	0.000	0.000	0.151	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	191	314	373	317	0	0	85	0
N.S.	1	1.00	1.66	2.73	3.24	2.76	0.00	0.00	0.74	0.00
time (sec)	N/A	0.417	0.070	0.497	0.199	0.084	0.000	0.000	0.154	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	87	143	130	160	0	0	48	161
N.S.	1	1.00	1.04	1.70	1.55	1.90	0.00	0.00	0.57	1.92
time (sec)	N/A	0.323	0.009	0.450	0.154	0.085	0.000	0.000	0.147	8.501

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	86	20	15	20	32	20
N.S.	1	1.00	1.11	1.00	4.78	1.11	0.83	1.11	1.78	1.11
time (sec)	N/A	0.214	1.733	0.187	0.230	0.063	0.603	0.221	0.151	8.265

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	142	31	17	20	79	20
N.S.	1	1.00	1.11	1.00	7.89	1.72	0.94	1.11	4.39	1.11
time (sec)	N/A	0.202	4.581	0.181	0.292	0.067	2.372	3.925	0.160	8.233

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	300	300	1337	952	2525	759	0	0	332	0
N.S.	1	1.00	4.46	3.17	8.42	2.53	0.00	0.00	1.11	0.00
time (sec)	N/A	0.768	7.171	0.895	0.781	0.096	0.000	0.000	0.159	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	229	229	649	575	1263	450	0	0	228	0
N.S.	1	1.00	2.83	2.51	5.52	1.97	0.00	0.00	1.00	0.00
time (sec)	N/A	0.612	6.718	0.856	0.317	0.092	0.000	0.000	0.154	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	200	238	527	219	0	0	128	0
N.S.	1	1.00	1.48	1.76	3.90	1.62	0.00	0.00	0.95	0.00
time (sec)	N/A	0.401	7.169	0.865	0.184	0.086	0.000	0.000	0.169	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	491	36	17	22	59	22
N.S.	1	1.00	1.10	1.00	24.55	1.80	0.85	1.10	2.95	1.10
time (sec)	N/A	0.234	19.012	0.473	0.579	0.080	0.928	0.365	0.152	8.897

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	609	47	19	22	157	22
N.S.	1	1.00	1.10	1.00	30.45	2.35	0.95	1.10	7.85	1.10
time (sec)	N/A	0.224	12.250	0.386	1.036	0.096	2.020	15.150	0.155	10.233

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	612	612	2594	1930	6861	1177	0	0	463	0
N.S.	1	1.00	4.24	3.15	11.21	1.92	0.00	0.00	0.76	0.00
time (sec)	N/A	1.289	7.323	0.995	8.171	0.109	0.000	0.000	0.162	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	427	427	1846	1138	3407	688	0	0	327	0
N.S.	1	1.00	4.32	2.67	7.98	1.61	0.00	0.00	0.77	0.00
time (sec)	N/A	0.942	7.051	0.855	1.591	0.124	0.000	0.000	0.174	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	275	275	277	493	1319	327	0	0	190	0
N.S.	1	1.00	1.01	1.79	4.80	1.19	0.00	0.00	0.69	0.00
time (sec)	N/A	0.590	8.102	0.767	0.430	0.084	0.000	0.000	0.159	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1935	52	17	22	86	22
N.S.	1	1.00	1.10	1.00	96.75	2.60	0.85	1.10	4.30	1.10
time (sec)	N/A	0.222	13.634	0.497	2.797	0.073	1.277	0.431	0.154	9.016

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	2194	63	19	22	237	22
N.S.	1	1.00	1.10	1.00	109.70	3.15	0.95	1.10	11.85	1.10
time (sec)	N/A	0.225	16.869	0.605	9.601	0.078	2.382	25.522	0.166	9.214

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	243	246	297	1468	983	1183	0	0	219	0
N.S.	1	1.01	1.22	6.04	4.05	4.87	0.00	0.00	0.90	0.00
time (sec)	N/A	0.978	1.446	0.612	0.481	0.107	0.000	0.000	0.159	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	181	188	236	940	715	834	0	0	159	0
N.S.	1	1.04	1.30	5.19	3.95	4.61	0.00	0.00	0.88	0.00
time (sec)	N/A	0.738	1.297	0.548	0.401	0.119	0.000	0.000	0.160	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	132	177	462	399	527	0	0	97	0
N.S.	1	1.06	1.42	3.70	3.19	4.22	0.00	0.00	0.78	0.00
time (sec)	N/A	0.494	1.405	0.493	0.295	0.089	0.000	0.000	0.157	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	279	27	17	22	31	22
N.S.	1	1.00	1.10	1.00	13.95	1.35	0.85	1.10	1.55	1.10
time (sec)	N/A	0.236	2.145	0.368	0.771	0.068	0.947	0.296	0.172	9.312

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	424	51	19	22	61	22
N.S.	1	1.00	1.10	1.00	21.20	2.55	0.95	1.10	3.05	1.10
time (sec)	N/A	0.237	4.074	0.355	1.791	0.068	1.745	4.191	0.158	9.511

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	848	848	1724	5959	4631	2537	0	0	1099	0
N.S.	1	1.00	2.03	7.03	5.46	2.99	0.00	0.00	1.30	0.00
time (sec)	N/A	2.279	9.244	0.944	2.833	0.139	0.000	0.000	0.162	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	654	654	761	3901	2530	1576	0	0	794	0
N.S.	1	1.00	1.16	5.96	3.87	2.41	0.00	0.00	1.21	0.00
time (sec)	N/A	1.791	6.111	0.789	0.919	0.115	0.000	0.000	0.172	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	214	228	745	2001	1156	843	0	0	481	0
N.S.	1	1.07	3.48	9.35	5.40	3.94	0.00	0.00	2.25	0.00
time (sec)	N/A	0.784	7.467	0.699	0.534	0.121	0.000	0.000	0.163	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1498	55	19	22	66	22
N.S.	1	1.00	1.10	1.00	74.90	2.75	0.95	1.10	3.30	1.10
time (sec)	N/A	0.234	12.128	0.562	10.948	0.077	1.617	0.445	0.156	9.798

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1977	96	20	22	121	22
N.S.	1	1.00	1.10	1.00	98.85	4.80	1.00	1.10	6.05	1.10
time (sec)	N/A	0.235	13.832	0.454	28.357	0.082	3.107	22.715	0.174	9.501

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [11] had the largest ratio of [1.3333299999999990]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	7	1.25	10	0.700
2	A	7	6	1.25	10	0.600
3	A	6	5	1.13	8	0.625
4	N/A	2	0	1.00	10	0.000
5	N/A	2	0	1.00	10	0.000
6	A	10	9	1.24	12	0.750
7	A	9	8	1.19	12	0.667
8	A	5	5	1.00	10	0.500
9	N/A	2	0	1.00	12	0.000
10	N/A	2	0	1.00	12	0.000
11	A	17	16	1.20	12	1.333
12	A	13	12	1.16	12	1.000
13	A	10	9	1.09	10	0.900
14	N/A	2	0	1.00	12	0.000
15	N/A	2	0	1.00	12	0.000
16	A	1	1	1.00	45	0.022
17	F	0	0	N/A	0.000	N/A
18	A	9	9	1.09	23	0.391
19	A	7	7	1.07	23	0.304
20	A	5	5	1.07	21	0.238

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
21	A	7	7	0.94	23	0.304
22	A	7	7	0.99	23	0.304
23	A	9	9	0.99	23	0.391
24	A	3	3	1.00	23	0.130
25	A	3	3	1.00	23	0.130
26	A	3	3	1.00	21	0.143
27	A	3	3	1.00	23	0.130
28	A	3	3	1.00	23	0.130
29	A	3	3	1.00	23	0.130
30	A	3	3	1.00	23	0.130
31	A	3	3	1.00	21	0.143
32	A	3	3	1.00	23	0.130
33	A	3	3	1.00	23	0.130
34	N/A	2	0	1.00	23	0.000
35	N/A	2	0	1.00	21	0.000
36	A	3	3	1.00	23	0.130
37	A	3	3	1.00	23	0.130
38	A	3	3	1.00	23	0.130
39	A	3	3	1.00	18	0.167
40	A	3	3	1.00	18	0.167
41	A	3	3	1.00	16	0.188
42	N/A	2	0	1.00	18	0.000
43	N/A	2	0	1.00	18	0.000
44	A	3	3	1.00	20	0.150
45	A	3	3	1.00	20	0.150
46	A	3	3	1.00	18	0.167
47	N/A	2	0	1.00	20	0.000
48	N/A	2	0	1.00	20	0.000
49	A	3	3	1.00	20	0.150
50	A	3	3	1.00	20	0.150
51	A	3	3	1.00	18	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
52	N/A	2	0	1.00	20	0.000
53	N/A	2	0	1.00	20	0.000
54	A	8	7	1.01	20	0.350
55	A	7	6	1.04	20	0.300
56	A	6	5	1.06	18	0.278
57	N/A	2	0	1.00	20	0.000
58	N/A	2	0	1.00	20	0.000
59	A	3	3	1.00	20	0.150
60	A	3	3	1.00	20	0.150
61	A	8	7	1.07	18	0.389
62	N/A	2	0	1.00	20	0.000
63	N/A	2	0	1.00	20	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \tan(a + bx) dx$	51
3.2	$\int x^2 \tan(a + bx) dx$	59
3.3	$\int x \tan(a + bx) dx$	65
3.4	$\int \frac{\tan(a+bx)}{x} dx$	71
3.5	$\int \frac{\tan(a+bx)}{x^2} dx$	77
3.6	$\int x^3 \tan^2(a + bx) dx$	83
3.7	$\int x^2 \tan^2(a + bx) dx$	91
3.8	$\int x \tan^2(a + bx) dx$	98
3.9	$\int \frac{\tan^2(a+bx)}{x} dx$	104
3.10	$\int \frac{\tan^2(a+bx)}{x^2} dx$	110
3.11	$\int x^3 \tan^3(a + bx) dx$	116
3.12	$\int x^2 \tan^3(a + bx) dx$	128
3.13	$\int x \tan^3(a + bx) dx$	137
3.14	$\int \frac{\tan^3(a+bx)}{x} dx$	144
3.15	$\int \frac{\tan^3(a+bx)}{x^2} dx$	150
3.16	$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$	156
3.17	$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$	161
3.18	$\int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx$	166
3.19	$\int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx$	174
3.20	$\int \frac{c+dx}{a+ia\tan(e+fx)} dx$	181
3.21	$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx$	187
3.22	$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx$	194
3.23	$\int \frac{1}{(c+dx)^3(a+ia\tan(e+fx))} dx$	202
3.24	$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx$	211

3.25	$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx$	219
3.26	$\int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx$	226
3.27	$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx$	232
3.28	$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx$	239
3.29	$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx$	248
3.30	$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx$	257
3.31	$\int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx$	265
3.32	$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^3} dx$	272
3.33	$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx$	280
3.34	$\int (c+dx)^m (a+ia\tan(e+fx))^2 dx$	290
3.35	$\int (c+dx)^m (a+ia\tan(e+fx)) dx$	296
3.36	$\int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx$	301
3.37	$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx$	306
3.38	$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx$	312
3.39	$\int (c+dx)^3 (a+b\tan(e+fx)) dx$	318
3.40	$\int (c+dx)^2 (a+b\tan(e+fx)) dx$	326
3.41	$\int (c+dx) (a+b\tan(e+fx)) dx$	333
3.42	$\int \frac{a+b\tan(e+fx)}{c+dx} dx$	339
3.43	$\int \frac{a+b\tan(e+fx)}{(c+dx)^2} dx$	344
3.44	$\int (c+dx)^3 (a+b\tan(e+fx))^2 dx$	349
3.45	$\int (c+dx)^2 (a+b\tan(e+fx))^2 dx$	358
3.46	$\int (c+dx) (a+b\tan(e+fx))^2 dx$	366
3.47	$\int \frac{(a+b\tan(e+fx))^2}{c+dx} dx$	372
3.48	$\int \frac{(a+b\tan(e+fx))^2}{(c+dx)^2} dx$	377
3.49	$\int (c+dx)^3 (a+b\tan(e+fx))^3 dx$	382
3.50	$\int (c+dx)^2 (a+b\tan(e+fx))^3 dx$	393
3.51	$\int (c+dx) (a+b\tan(e+fx))^3 dx$	403
3.52	$\int \frac{(a+b\tan(e+fx))^3}{c+dx} dx$	410
3.53	$\int \frac{(a+b\tan(e+fx))^3}{(c+dx)^2} dx$	416
3.54	$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$	422
3.55	$\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$	433
3.56	$\int \frac{c+dx}{a+b\tan(e+fx)} dx$	442
3.57	$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$	449
3.58	$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$	454
3.59	$\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$	459
3.60	$\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx$	470

3.61	$\int \frac{c+dx}{(a+b\tan(e+fx))^2} dx$	481
3.62	$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$	491
3.63	$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$	497

3.1 $\int x^3 \tan(a + bx) dx$

Optimal result	51
Mathematica [A] (verified)	51
Rubi [A] (verified)	52
Maple [A] (verified)	55
Fricas [B] (verification not implemented)	55
Sympy [F]	56
Maxima [B] (verification not implemented)	56
Giac [F]	57
Mupad [F(-1)]	57
Reduce [F]	58

Optimal result

Integrand size = 10, antiderivative size = 106

$$\begin{aligned} \int x^3 \tan(a + bx) dx = & \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\ & - \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} \end{aligned}$$

output
$$1/4*I*x^4-x^3*ln(1+exp(2*I*(b*x+a)))/b+3/2*I*x^2*polylog(2,-exp(2*I*(b*x+a)))/b^2-3/2*x*polylog(3,-exp(2*I*(b*x+a)))/b^3-3/4*I*polylog(4,-exp(2*I*(b*x+a)))/b^4$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00

$$\begin{aligned} \int x^3 \tan(a + bx) dx = & \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\ & - \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} \end{aligned}$$

input
$$\text{Integrate}[x^3 \tan[a + b x], x]$$

output
$$\begin{aligned} & (\text{I}/4)*x^4 - (x^3*\text{Log}[1 + E^{((2*I)*(a + b*x))}])/b + (((3*I)/2)*x^2*\text{PolyLog}[2, -E^{((2*I)*(a + b*x))}])/b^2 - (3*x*\text{PolyLog}[3, -E^{((2*I)*(a + b*x))}])/(2*b^3) - (((3*I)/4)*\text{PolyLog}[4, -E^{((2*I)*(a + b*x))}])/b^4 \end{aligned}$$

Rubi [A] (verified)

Time = 0.60 (sec), antiderivative size = 133, normalized size of antiderivative = 1.25, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.700, Rules used = {3042, 4202, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \tan(a + bx) dx \\ & \quad \downarrow \text{3042} \\ & \int x^3 \tan(a + bx) dx \\ & \quad \downarrow \text{4202} \\ & \frac{ix^4}{4} - 2i \int \frac{e^{2i(a+bx)} x^3}{1 + e^{2i(a+bx)}} dx \\ & \quad \downarrow \text{2620} \\ & \frac{ix^4}{4} - 2i \left(\frac{3i \int x^2 \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) \\ & \quad \downarrow \text{3011} \\ & \frac{ix^4}{4} - 2i \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) \\ & \quad \downarrow \text{7163} \end{aligned}$$

$$\begin{aligned}
& \frac{i x^4}{4} - \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left(\frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -e^{2i(a+bx)}) dx}{2b} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} \right) - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \\
& \quad \downarrow 2720 \\
& \frac{i x^4}{4} - \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left(\frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} \right) - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \\
& \quad \downarrow 7143 \\
& \frac{i x^4}{4} - \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left(\frac{\operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} \right) - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b}
\end{aligned}$$

input `Int[x^3*Tan[a + b*x],x]`

output `(I/4)*x^4 - (2*I)*((-1/2*I)*x^3*Log[1 + E^((2*I)*(a + b*x))])/b + (((3*I)/2)*(((I/2)*x^2*PolyLog[2, -E^((2*I)*(a + b*x))])/b - (I*((-1/2*I)*x*PolyLog[3, -E^((2*I)*(a + b*x))])/b + PolyLog[4, -E^((2*I)*(a + b*x))]/(4*b^2))/b))/b)`

Definitions of rubi rules used

rule 2620

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))*((n_.)*((c_.) + (d_.)*(x_))^((m_.))/((a_.) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_)))*((n_.))), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^((m_.)) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*F_[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))*((n_.)*((f_.) + (g_.)*((x_.))^((m_.)), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4202

```
Int[((c_.) + (d_.)*(x_))^((m_.))*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*(F_)^((c_.)*(a_.) + (b_.)*(x_)))^p_], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [A] (verified)

Time = 0.28 (sec), antiderivative size = 125, normalized size of antiderivative = 1.18

method	result
risch	$\frac{ix^4}{4} + \frac{3ix^2 \operatorname{polylog}(2, -e^{2i(bx+a)})}{2b^2} - \frac{3i \operatorname{polylog}(4, -e^{2i(bx+a)})}{4b^4} - \frac{2a^3 \ln(e^{i(bx+a)})}{b^4} + \frac{3ia^4}{2b^4} + \frac{2ia^3x}{b^3} - \frac{3x \operatorname{polylog}(3, -e^{2i(bx+a)})}{2b^3}$

input `int(x^3*tan(b*x+a), x, method=_RETURNVERBOSE)`output `1/4*I*x^4+3/2*I*x^2*polylog(2,-exp(2*I*(b*x+a))/b^2-3/4*I*polylog(4,-exp(2*I*(b*x+a))/b^4-2/b^4*a^3*ln(exp(I*(b*x+a)))+3/2*I/b^4*a^4+2*I/b^3*a^3*x-3/2*x*polylog(3,-exp(2*I*(b*x+a))/b^3-x^3*ln(1+exp(2*I*(b*x+a)))/b`**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 286 vs. 2(83) = 166.

Time = 0.09 (sec), antiderivative size = 286, normalized size of antiderivative = 2.70

$$\int x^3 \tan(a + bx) dx = \frac{-4 b^3 x^3 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 4 b^3 x^3 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 6i b^2 x^2 \text{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 6i b^2 x^2 \text{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right)}{b^4}$$

input `integrate(x^3*tan(b*x+a), x, algorithm="fricas")`

output

$$\begin{aligned} & -\frac{1}{8}(4b^3x^3 \log(-2(I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1)) + 4b^3x^3 \log(-2(-I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1)) + 6Ib^2x^2 \operatorname{dilog}(2(I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1) + 1) - 6Ib^2x^2 \operatorname{dilog}(2(-I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1) + 1) + 6bx \operatorname{polylog}(3, (\tan(bx+a)^2 + 2I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1)) + 6bx \operatorname{polylog}(3, (\tan(bx+a)^2 - 2I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1)) - 3I \operatorname{polylog}(4, (\tan(bx+a)^2 + 2I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1)) + 3I \operatorname{polylog}(4, (\tan(bx+a)^2 - 2I\tan(bx+a) - 1)/(\tan(bx+a)^2 + 1))) / b^4 \end{aligned}$$

Sympy [F]

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(a + bx) dx$$

input

```
integrate(x**3*tan(b*x+a),x)
```

output

```
Integral(x**3*tan(a + b*x), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(83) = 166$.

Time = 0.15 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.29

$$\int x^3 \tan(a + bx) dx = \frac{-3i(bx+a)^4 + 12i(bx+a)^3a - 18i(bx+a)^2a^2 + 12a^3 \log(\sec(bx+a)) - 4(-4i(bx+a)^3 + 9i(bx+a)^2a)}{b^4}$$

input

```
integrate(x^3*tan(b*x+a),x, algorithm="maxima")
```

output

$$\begin{aligned} & -\frac{1}{12}(-3I(bx + a)^4 + 12I(bx + a)^3a - 18I(bx + a)^2a^2 + 12a \\ & \sim 3\log(\sec(bx + a)) - 4(-4I(bx + a)^3 + 9I(bx + a)^2a - 9I(bx + a)a^2) \arctan 2(\sin(2bx + 2a), \cos(2bx + 2a) + 1) - 6(4I(bx + a)^2 - 6I(bx + a)a + 3Ia^2)\operatorname{dilog}(-e^{(2Ibx + 2Ia)}) + 2(4(bx + a)^3 - 9(bx + a)^2a + 9(bx + a)a^2)\log(\cos(2bx + 2a)^2 + \sin(2bx + 2a)^2 + 2\cos(2bx + 2a) + 1) + 6(4bx + a)\operatorname{polylog}(3, -e^{(2Ibx + 2Ia)}) + 12I\operatorname{polylog}(4, -e^{(2Ibx + 2Ia)}))/b^4 \end{aligned}$$

Giac [F]

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(bx + a) dx$$

input

```
integrate(x^3*tan(b*x+a),x, algorithm="giac")
```

output

```
integrate(x^3*tan(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(a + b x) dx$$

input

```
int(x^3*tan(a + b*x),x)
```

output

```
int(x^3*tan(a + b*x), x)
```

Reduce [F]

$$\int x^3 \tan(a + bx) dx = \int \tan(bx + a) x^3 dx$$

input `int(x^3*tan(b*x+a),x)`

output `int(tan(a + b*x)*x**3,x)`

3.2 $\int x^2 \tan(a + bx) dx$

Optimal result	59
Mathematica [A] (verified)	59
Rubi [A] (verified)	60
Maple [A] (verified)	62
Fricas [B] (verification not implemented)	62
Sympy [F]	63
Maxima [B] (verification not implemented)	63
Giac [F]	64
Mupad [F(-1)]	64
Reduce [F]	64

Optimal result

Integrand size = 10, antiderivative size = 77

$$\int x^2 \tan(a + bx) dx = \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3}$$

output `1/3*I*x^3-x^2*ln(1+exp(2*I*(b*x+a)))/b+I*x*polylog(2,-exp(2*I*(b*x+a))/b^2-1/2*polylog(3,-exp(2*I*(b*x+a)))/b^3`

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 77, normalized size of antiderivative = 1.00

$$\int x^2 \tan(a + bx) dx = \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3}$$

input `Integrate[x^2*Tan[a + b*x], x]`

output
$$(I/3)*x^3 - (x^2*Log[1 + E^((2*I)*(a + b*x))])/b + (I*x*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2 - PolyLog[3, -E^((2*I)*(a + b*x))]/(2*b^3)$$

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.25, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4202, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \tan(a + bx) dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int x^2 \tan(a + bx) dx \\
 & \downarrow \textcolor{blue}{4202} \\
 & \frac{ix^3}{3} - 2i \int \frac{e^{2i(a+bx)} x^2}{1 + e^{2i(a+bx)}} dx \\
 & \downarrow \textcolor{blue}{2620} \\
 & \frac{ix^3}{3} - 2i \left(\frac{i \int x \log(1 + e^{2i(a+bx)}) dx}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \downarrow \textcolor{blue}{3011} \\
 & \frac{ix^3}{3} - 2i \left(\frac{i \left(\frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \downarrow \textcolor{blue}{2720} \\
 & \frac{ix^3}{3} - \\
 & 2i \left(\frac{i \left(\frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \text{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \downarrow \textcolor{blue}{7143}
 \end{aligned}$$

$$\frac{ix^3}{3} - 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right)$$

input `Int[x^2*Tan[a + b*x], x]`

output `(I/3)*x^3 - (2*I)*(((-1/2*I)*x^2*Log[1 + E^((2*I)*(a + b*x))])/b + (I*((I/2)*x*PolyLog[2, -E^((2*I)*(a + b*x))])/b - PolyLog[3, -E^((2*I)*(a + b*x))]/(4*b^2))/b)`

Definitions of rubi rules used

rule 2620 `Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_.))))^(n_.)*(c_.) + (d_.)*(x_.))^((m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^((m_.)) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.))))^(n_.)]*((f_.) + (g_.)*(x_.))^((m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4202

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + (f_)*(x_)], x_Symbol] :> Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Simp[2*I Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGT
Q[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_)*(x_)]^(p_.)]/((d_.) + (e_.)*(x_), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.34

method	result	
risch	$\frac{ix^3}{3} - \frac{2ia^2x}{b^2} - \frac{4ia^3}{3b^3} - \frac{x^2 \ln(1+e^{2i(bx+a)})}{b} + \frac{ix \operatorname{polylog}(2, -e^{2i(bx+a)})}{b^2} - \frac{\operatorname{polylog}(3, -e^{2i(bx+a)})}{2b^3} + \frac{2a^2 \ln(e^{i(bx+a)})}{b^3}$	1

input `int(x^2*tan(b*x+a), x, method=_RETURNVERBOSE)`

output

```
1/3*I*x^3-2*I/b^2*a^2*x-4/3*I/b^3*a^3-x^2*ln(1+exp(2*I*(b*x+a)))/b+I*x*p
olylog(2,-exp(2*I*(b*x+a)))/b^2-1/2*polylog(3,-exp(2*I*(b*x+a)))/b^3+2/b^3*a
^2*ln(exp(I*(b*x+a)))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(62) = 124$.

Time = 0.09 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.60

$$\int x^2 \tan(a + bx) dx =$$

$$-\frac{2 b^2 x^2 \log \left(-\frac{2 (i \tan (b x+a)-1)}{\tan ^2(b x+a)+1}\right)+2 b^2 x^2 \log \left(-\frac{2 (-i \tan (b x+a)-1)}{\tan ^2(b x+a)+1}\right)+2 i b x \text{Li}_2\left(\frac{2 (i \tan (b x+a)-1)}{\tan ^2(b x+a)+1}+1\right)-2 i b x \text{Li}_2\left(\frac{2 (-i \tan (b x+a)-1)}{\tan ^2(b x+a)+1}+1\right)}{4 b^3}$$

input `integrate(x^2*tan(b*x+a), x, algorithm="fricas")`

output

$$\begin{aligned} & -\frac{1}{4} \cdot (2 \cdot b^2 \cdot x^2 \cdot \log(-2 \cdot (\text{I} \cdot \tan(b \cdot x + a) - 1) / (\tan(b \cdot x + a)^2 + 1)) + 2 \cdot b^2 \cdot x^2 \cdot \log(-2 \cdot (-\text{I} \cdot \tan(b \cdot x + a) - 1) / (\tan(b \cdot x + a)^2 + 1)) + 2 \cdot \text{I} \cdot b \cdot x \cdot \text{dilog}(2 \cdot (\text{I} \cdot \tan(b \cdot x + a) - 1) / (\tan(b \cdot x + a)^2 + 1) + 1) - 2 \cdot \text{I} \cdot b \cdot x \cdot \text{dilog}(2 \cdot (-\text{I} \cdot \tan(b \cdot x + a) - 1) / (\tan(b \cdot x + a)^2 + 1) + 1) + \text{polylog}(3, (\tan(b \cdot x + a)^2 + 2 \cdot \text{I} \cdot \tan(b \cdot x + a) - 1) / (\tan(b \cdot x + a)^2 + 1)) + \text{polylog}(3, (\tan(b \cdot x + a)^2 - 2 \cdot \text{I} \cdot \tan(b \cdot x + a) - 1) / (\tan(b \cdot x + a)^2 + 1))) / b^3 \end{aligned}$$

Sympy [F]

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(a + bx) dx$$

input

```
integrate(x**2*tan(b*x+a),x)
```

output

```
Integral(x**2*tan(a + b*x), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(62) = 124$.

Time = 0.16 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.12

$$\begin{aligned} \int x^2 \tan(a + bx) dx = \\ -2i (bx + a)^3 + 6i (bx + a)^2 a - 6i bx \text{Li}_2(-e^{(2i bx + 2i a)}) - 6 a^2 \log(\sec(bx + a)) - 6 (-i (bx + a)^2 + 2i a) \end{aligned}$$

input

```
integrate(x^2*tan(b*x+a),x, algorithm="maxima")
```

output

$$\begin{aligned} & -\frac{1}{6} \cdot (-2 \cdot \text{I} \cdot (b \cdot x + a)^3 + 6 \cdot \text{I} \cdot (b \cdot x + a)^2 a - 6 \cdot \text{I} \cdot b \cdot x \cdot \text{dilog}(-e^{(2 \cdot \text{I} \cdot b \cdot x + 2 \cdot \text{I} \cdot a)}) - 6 \cdot a^2 \log(\sec(b \cdot x + a)) - 6 \cdot (-\text{I} \cdot (b \cdot x + a)^2 + 2 \cdot \text{I} \cdot (b \cdot x + a) \cdot a) \cdot \text{arctan2}(\sin(2 \cdot b \cdot x + 2 \cdot a), \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) + 3 \cdot ((b \cdot x + a)^2 - 2 \cdot (b \cdot x + a) \cdot a) \cdot \log(\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) + 3 \cdot \text{polylog}(3, -e^{(2 \cdot \text{I} \cdot b \cdot x + 2 \cdot \text{I} \cdot a)})) / b^3 \end{aligned}$$

Giac [F]

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(bx + a) dx$$

input `integrate(x^2*tan(b*x+a),x, algorithm="giac")`

output `integrate(x^2*tan(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(a + b x) dx$$

input `int(x^2*tan(a + b*x),x)`

output `int(x^2*tan(a + b*x), x)`

Reduce [F]

$$\int x^2 \tan(a + bx) dx = \int \tan(bx + a) x^2 dx$$

input `int(x^2*tan(b*x+a),x)`

output `int(tan(a + b*x)*x**2,x)`

3.3 $\int x \tan(a + bx) dx$

Optimal result	65
Mathematica [A] (verified)	65
Rubi [A] (verified)	66
Maple [A] (verified)	67
Fricas [B] (verification not implemented)	68
Sympy [F]	68
Maxima [B] (verification not implemented)	69
Giac [F]	69
Mupad [B] (verification not implemented)	70
Reduce [F]	70

Optimal result

Integrand size = 8, antiderivative size = 54

$$\int x \tan(a + bx) dx = \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2}$$

output $\frac{1/2*I*x^2-x*ln(1+exp(2*I*(b*x+a)))/b+1/2*I*polylog(2,-exp(2*I*(b*x+a)))/b^2}{2}$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int x \tan(a + bx) dx = \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2}$$

input `Integrate[x*Tan[a + b*x], x]`

output $(I/2)*x^2 - (x*Log[1 + E^((2*I)*(a + b*x))])/b + ((I/2)*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.13, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tan(a + bx) dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int x \tan(a + bx) dx \\
 & \downarrow \textcolor{blue}{4202} \\
 & \frac{ix^2}{2} - 2i \int \frac{e^{2i(a+bx)}x}{1 + e^{2i(a+bx)}} dx \\
 & \downarrow \textcolor{blue}{2620} \\
 & \frac{ix^2}{2} - 2i \left(\frac{i \int \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \downarrow \textcolor{blue}{2715} \\
 & \frac{ix^2}{2} - 2i \left(\frac{\int e^{-2i(a+bx)} \log(1 + e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \\
 & \downarrow \textcolor{blue}{2838} \\
 & \frac{ix^2}{2} - 2i \left(-\frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right)
 \end{aligned}$$

input `Int[x*Tan[a + b*x], x]`

output `(I/2)*x^2 - (2*I)*(((-1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))])/b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2))`

Definitions of rubi rules used

rule 2620 $\text{Int}[((\text{F}_.)^((\text{g}_.)*((\text{e}_.) + (\text{f}_.)*(\text{x}_.))))^{(\text{n}_.)}*((\text{c}_.) + (\text{d}_.)*(\text{x}_.))^{(\text{m}_.)})/((\text{a}_.) + (\text{b}_.)*((\text{F}_.)^((\text{g}_.)*((\text{e}_.) + (\text{f}_.)*(\text{x}_.))))^{(\text{n}_.)}), \text{x_Symbol}] \Rightarrow \text{Simp}[(\text{c} + \text{d}*\text{x})^{\text{m}}/(\text{b}*\text{f}*\text{g}*\text{n}*\text{Log}[\text{F}])*\text{Log}[1 + \text{b}*((\text{F}^(\text{g}*(\text{e} + \text{f}*\text{x})))^{\text{n}/\text{a}})], \text{x}] - \text{Simp}[\text{d}*(\text{m}/(\text{b}*\text{f}*\text{g}*\text{n}*\text{Log}[\text{F}])) \text{Int}[(\text{c} + \text{d}*\text{x})^{(\text{m} - 1)}*\text{Log}[1 + \text{b}*((\text{F}^(\text{g}*(\text{e} + \text{f}*\text{x})))^{\text{n}/\text{a}})], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}] \&& \text{IGtQ}[\text{m}, 0]$

rule 2715 $\text{Int}[\text{Log}[(\text{a}_.) + (\text{b}_.)*((\text{F}_.)^((\text{e}_.)*((\text{c}_.) + (\text{d}_.)*(\text{x}_.))))^{(\text{n}_.)}], \text{x_Symbol}] \Rightarrow \text{Simp}[1/(\text{d}*\text{e}*\text{n}*\text{Log}[\text{F}]) \text{Subst}[\text{Int}[\text{Log}[\text{a} + \text{b}*\text{x}]/\text{x}, \text{x}], \text{x}, (\text{F}^(\text{e}*(\text{c} + \text{d}*\text{x})))^{\text{n}}], \text{x}] /; \text{FreeQ}[\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}] \&& \text{GtQ}[\text{a}, 0]$

rule 2838 $\text{Int}[\text{Log}[(\text{c}_.)*((\text{d}_.) + (\text{e}_.)*(\text{x}_.))^{(\text{n}_.)})/(\text{x}_.), \text{x_Symbol}] \Rightarrow \text{Simp}[-\text{PolyLog}[2, (-\text{c})*\text{e}*\text{x}^{\text{n}}]/\text{n}, \text{x}] /; \text{FreeQ}[\{\text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}] \&& \text{EqQ}[\text{c}*\text{d}, 1]$

rule 3042 $\text{Int}[\text{u}_., \text{x_Symbol}] \Rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 4202 $\text{Int}[(\text{c}_.) + (\text{d}_.)*(\text{x}_.)^{\text{m}})*\text{tan}[(\text{e}_.) + (\text{f}_.)*(\text{x}_.)], \text{x_Symbol}] \Rightarrow \text{Simp}[\text{I}*((\text{c} + \text{d}*\text{x})^{(\text{m} + 1)}/(\text{d}*(\text{m} + 1))), \text{x}] - \text{Simp}[2*\text{I} \text{Int}[(\text{c} + \text{d}*\text{x})^{\text{m}}*(\text{E}^{(2*\text{I}*(\text{e} + \text{f}*\text{x}))/(1 + \text{E}^{(2*\text{I}*(\text{e} + \text{f}*\text{x}))})}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}] \&& \text{IGtQ}[\text{m}, 0]$

Maple [A] (verified)

Time = 0.28 (sec), antiderivative size = 78, normalized size of antiderivative = 1.44

method	result
risch	$\frac{ix^2}{2} + \frac{2iax}{b} + \frac{ia^2}{b^2} - \frac{x \ln(1+e^{2i(bx+a)})}{b} + \frac{i \text{polylog}(2, -e^{2i(bx+a)})}{2b^2} - \frac{2a \ln(e^{i(bx+a)})}{b^2}$
parts	$\frac{\ln(1+\tan(bx+a)^2)x}{2b} - \frac{i \left(\ln(\tan(bx+a)-i) \ln(1+\tan(bx+a)^2) - \frac{\ln(\tan(bx+a)-i)^2}{2} - \text{dilog}\left(-\frac{i(\tan(bx+a)+i)}{2}\right) - \ln(\tan(bx+a)-i) \ln\left(-\frac{i(\tan(bx+a)+i)}{2}\right) \right)}{2}$

input $\text{int}(\text{x}*\text{tan}(\text{b}*\text{x}+\text{a}), \text{x}, \text{method}=\text{_RETURNVERBOSE})$

output
$$\frac{1/2*I*x^2+2*I/b*a*x+I/b^2*a^2-x*ln(1+exp(2*I*(b*x+a)))/b+1/2*I*polylog(2,-exp(2*I*(b*x+a)))/b^2-2/b^2*a*ln(exp(I*(b*x+a)))}{b^2}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 122 vs. $2(41) = 82$.

Time = 0.08 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.26

$$\int x \tan(a + bx) dx = -\frac{2 b x \log \left(-\frac{2 (i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2 b x \log \left(-\frac{2 (-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + i \text{Li}_2\left(\frac{2 (i \tan(bx+a)-1)}{\tan(bx+a)^2+1}+1\right) - i \text{Li}_2\left(\frac{2 (-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}+1\right)}{4 b^2}$$

input `integrate(x*tan(b*x+a),x, algorithm="fricas")`

output
$$\begin{aligned} & -\frac{1}{4} \left(2 b x \log \left(-2 \left(I \tan(b x + a) - 1 \right) / (\tan(b x + a)^2 + 1) \right) + 2 b x \log \left(-2 \left(-I \tan(b x + a) - 1 \right) / (\tan(b x + a)^2 + 1) \right) + I \operatorname{dilog} \left(2 \left(I \tan(b x + a) - 1 \right) / (\tan(b x + a)^2 + 1) + 1 \right) - I \operatorname{dilog} \left(2 \left(-I \tan(b x + a) - 1 \right) / (\tan(b x + a)^2 + 1) + 1 \right) \right) / b^2 \end{aligned}$$

Sympy [F]

$$\int x \tan(a + bx) dx = \int x \tan(a + bx) dx$$

input `integrate(x*tan(b*x+a),x)`

output `Integral(x*tan(a + b*x), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 92 vs. $2(41) = 82$.

Time = 0.16 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.70

$$\int x \tan(a + bx) dx = -\frac{-i b^2 x^2 + 2i b x \arctan(\sin(2bx + 2a), \cos(2bx + 2a) + 1) + b x \log(\cos(2bx + 2a)^2 + \sin(2bx + 2a)^2)}{2b^2}$$

input `integrate(x*tan(b*x+a),x, algorithm="maxima")`

output
$$\begin{aligned} & -1/2*(-I*b^2*x^2 + 2*I*b*x*arctan2(\sin(2*b*x + 2*a), \cos(2*b*x + 2*a) + 1) \\ & + b*x*log(\cos(2*b*x + 2*a)^2 + \sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - I*dilog(-e^(2*I*b*x + 2*I*a)))/b^2 \end{aligned}$$

Giac [F]

$$\int x \tan(a + bx) dx = \int x \tan(bx + a) dx$$

input `integrate(x*tan(b*x+a),x, algorithm="giac")`

output `integrate(x*tan(b*x + a), x)`

Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.39

$$\int x \tan(a + bx) dx =$$

$$-\frac{\pi \ln(\cos(bx)) + \text{polylog}(2, -e^{-a2i} e^{-bx2i}) 1i - \pi \ln(e^{-a2i} e^{-bx2i} + 1) + 2a \ln(e^{-a2i} e^{-bx2i} + 1) - \tau}{2b^2}$$

input `int(x*tan(a + b*x),x)`

output $-(\text{polylog}(2, -\exp(-a*2i)*\exp(-b*x*2i))*1i - \pi*\log(\exp(b*x*2i) + 1) - \pi*i*\log(\exp(-a*2i)*\exp(-b*x*2i) + 1) + 2*a*\log(\exp(-a*2i)*\exp(-b*x*2i) + 1) + \pi*i*\log(\cos(b*x)) + b^2*x^2*1i - \log(\cos(a + b*x))*(2*a - \pi) + 2*b*x*\log(\exp(-a*2i)*\exp(-b*x*2i) + 1) + a*b*x*2i)/(2*b^2)$

Reduce [F]

$$\int x \tan(a + bx) dx = \int \tan(bx + a) x dx$$

input `int(x*tan(b*x+a),x)`

output `int(tan(a + b*x)*x,x)`

3.4 $\int \frac{\tan(a+bx)}{x} dx$

Optimal result	71
Mathematica [N/A]	71
Rubi [N/A]	72
Maple [N/A]	73
Fricas [N/A]	73
Sympy [N/A]	74
Maxima [N/A]	74
Giac [N/A]	75
Mupad [N/A]	75
Reduce [N/A]	75

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\tan(a + bx)}{x} dx = \text{Int}\left(\frac{\tan(a + bx)}{x}, x\right)$$

output `Defer(Int)(tan(b*x+a)/x,x)`

Mathematica [N/A]

Not integrable

Time = 1.36 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + bx)}{x} dx$$

input `Integrate[Tan[a + b*x]/x,x]`

output `Integrate[Tan[a + b*x]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\tan(a + bx)}{x} dx \\ \downarrow \textcolor{blue}{3042} \\ \int \frac{\tan(a + bx)}{x} dx \\ \downarrow \textcolor{blue}{4222} \\ \int \frac{\tan(a + bx)}{x} dx \end{array}$$

input `Int[Tan[a + b*x]/x,x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4222 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)}, x_\text{Symbol}] \rightarrow \text{Simp}[\text{If}[\text{MatchQ}[f, (f1_.)*(Complex[0, j_])], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Coth}[-I]*(e - \text{Pi}/2) - I*f*x]^n, x], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Tanh}[-I]*e - I*f*x]^n, x]], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], (-1)^n* \text{Unintegatable}[(c + d*x)^m*\text{Cot}[e - \text{Pi}/2 + f*x]^n, x], \text{Unintegatable}[(c + d*x)^m*\text{Tan}[e + f*x]^n, x]]], x] /; \text{FreeQ}[\{c, d, e, f, m, n\}, x] \&& \text{IntegerQ}[n]$

Maple [N/A]

Not integrable

Time = 0.12 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx + a)}{x} dx$$

input $\text{int}(\tan(b*x+a)/x, x)$

output $\text{int}(\tan(b*x+a)/x, x)$

Fricas [N/A]

Not integrable

Time = 0.07 (sec), antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

input $\text{integrate}(\tan(b*x+a)/x, x, \text{algorithm}=\text{"fricas"})$

output `integral(tan(b*x + a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan (a + bx)}{x} dx$$

input `integrate(tan(b*x+a)/x,x)`

output `Integral(tan(a + b*x)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan (bx + a)}{x} dx$$

input `integrate(tan(b*x+a)/x,x, algorithm="maxima")`

output `integrate(tan(b*x + a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

input `integrate(tan(b*x+a)/x,x, algorithm="giac")`

output `integrate(tan(b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 8.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + b x)}{x} dx$$

input `int(tan(a + b*x)/x,x)`

output `int(tan(a + b*x)/x, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

input `int(tan(b*x+a)/x,x)`

output `int(tan(a + b*x)/x,x)`

3.5 $\int \frac{\tan(a+bx)}{x^2} dx$

Optimal result	77
Mathematica [N/A]	77
Rubi [N/A]	78
Maple [N/A]	79
Fricas [N/A]	79
Sympy [N/A]	80
Maxima [N/A]	80
Giac [N/A]	81
Mupad [N/A]	81
Reduce [N/A]	81

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\tan(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan(a + bx)}{x^2}, x\right)$$

output `Defer(Int)(tan(b*x+a)/x^2,x)`

Mathematica [N/A]

Not integrable

Time = 2.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)}{x^2} dx$$

input `Integrate[Tan[a + b*x]/x^2,x]`

output `Integrate[Tan[a + b*x]/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\tan(a + bx)}{x^2} dx \\ \downarrow \text{3042} \\ \int \frac{\tan(a + bx)}{x^2} dx \\ \downarrow \text{4222} \\ \int \frac{\tan(a + bx)}{x^2} dx \end{array}$$

input `Int[Tan[a + b*x]/x^2,x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4222 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)}, x_\text{Symbol}] \rightarrow \text{Simp}[\text{If}[\text{MatchQ}[f, (f1_.)*(Complex[0, j_])], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Coth}[-I]*(e - \text{Pi}/2) - I*f*x]^n, x], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Tanh}[-I]*e - I*f*x]^n, x]], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], (-1)^n* \text{Unintegatable}[(c + d*x)^m*\text{Cot}[e - \text{Pi}/2 + f*x]^n, x], \text{Unintegatable}[(c + d*x)^m*\text{Tan}[e + f*x]^n, x]]], x] /; \text{FreeQ}[\{c, d, e, f, m, n\}, x] \&& \text{IntegerQ}[n]$

Maple [N/A]

Not integrable

Time = 0.13 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx + a)}{x^2} dx$$

input $\text{int}(\tan(b*x+a)/x^2, x)$

output $\text{int}(\tan(b*x+a)/x^2, x)$

Fricas [N/A]

Not integrable

Time = 0.07 (sec), antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

input $\text{integrate}(\tan(b*x+a)/x^2, x, \text{algorithm}=\text{"fricas"})$

output `integral(tan(b*x + a)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan (a + bx)}{x^2} dx$$

input `integrate(tan(b*x+a)/x**2,x)`

output `Integral(tan(a + b*x)/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan (bx + a)}{x^2} dx$$

input `integrate(tan(b*x+a)/x^2,x, algorithm="maxima")`

output `integrate(tan(b*x + a)/x^2, x)`

Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

input `integrate(tan(b*x+a)/x^2,x, algorithm="giac")`

output `integrate(tan(b*x + a)/x^2, x)`

Mupad [N/A]

Not integrable

Time = 7.78 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + b x)}{x^2} dx$$

input `int(tan(a + b*x)/x^2,x)`

output `int(tan(a + b*x)/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

input `int(tan(b*x+a)/x^2,x)`

```
output int(tan(a + b*x)/x**2,x)
```

3.6 $\int x^3 \tan^2(a + bx) dx$

Optimal result	83
Mathematica [A] (verified)	83
Rubi [A] (verified)	84
Maple [A] (verified)	87
Fricas [B] (verification not implemented)	87
Sympy [F]	88
Maxima [B] (verification not implemented)	88
Giac [F]	89
Mupad [F(-1)]	89
Reduce [F]	90

Optimal result

Integrand size = 12, antiderivative size = 98

$$\begin{aligned} \int x^3 \tan^2(a + bx) dx = & -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} - \frac{3ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} \\ & + \frac{3 \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^4} + \frac{x^3 \tan(a + bx)}{b} \end{aligned}$$

output
$$-\text{I}*x^3/b - 1/4*x^4 + 3*x^2*\ln(1+\exp(2*\text{I}*(b*x+a)))/b^2 - 3*\text{I}*x*\operatorname{polylog}(2, -\exp(2*\text{I}*(b*x+a)))/b^3 + 3/2*\operatorname{polylog}(3, -\exp(2*\text{I}*(b*x+a)))/b^4 + x^3*\tan(b*x+a)/b$$

Mathematica [A] (verified)

Time = 0.56 (sec), antiderivative size = 115, normalized size of antiderivative = 1.17

$$\begin{aligned} \int x^3 \tan^2(a + bx) dx = & -\frac{x^4}{4} \\ & + \frac{2b^2 x^2 \left(\frac{2ibx}{1+e^{2ia}} + 3 \log(1 + e^{-2i(a+bx)})\right) + 6ibx \operatorname{PolyLog}(2, -e^{-2i(a+bx)}) + 3 \operatorname{PolyLog}(3, -e^{-2i(a+bx)})}{2b^4} \\ & + \frac{x^3 \sec(a) \sec(a + bx) \sin(bx)}{b} \end{aligned}$$

input `Integrate[x^3*Tan[a + b*x]^2, x]`

output

$$\begin{aligned} & -\frac{1}{4}x^4 + \frac{(2b^2x^2((2I)b*x)/(1 + E^{(2I)*a}) + 3\text{Log}[1 + E^{(-2I)*(a + b*x)})] + (6I)*b*x*\text{PolyLog}[2, -E^{(-2I)*(a + b*x)}] + 3*\text{PolyLog}[3, -E^{(-2I)*(a + b*x)}])/(2*b^4) + (x^3*\text{Sec}[a]*\text{Sec}[a + b*x]*\text{Sin}[b*x])/b \end{aligned}$$

Rubi [A] (verified)

Time = 0.60 (sec), antiderivative size = 122, normalized size of antiderivative = 1.24, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4203, 15, 3042, 4202, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 \tan^2(a + bx) dx \\ & \downarrow \text{3042} \\ & \int x^3 \tan(a + bx)^2 dx \\ & \downarrow \text{4203} \\ & -\frac{3 \int x^2 \tan(a + bx) dx}{b} - \int x^3 dx + \frac{x^3 \tan(a + bx)}{b} \\ & \downarrow \text{15} \\ & -\frac{3 \int x^2 \tan(a + bx) dx}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\ & \downarrow \text{3042} \\ & -\frac{3 \int x^2 \tan(a + bx) dx}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\ & \downarrow \text{4202} \\ & -\frac{3 \left(\frac{ix^3}{3} - 2i \int \frac{e^{2i(a+bx)} x^2}{1+e^{2i(a+bx)}} dx \right)}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \\ & \downarrow \text{2620} \\ & -\frac{3 \left(\frac{ix^3}{3} - 2i \left(\frac{i \int x \log(1+e^{2i(a+bx)}) dx}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^3 \tan(a + bx)}{b} - \frac{x^4}{4} \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3011} \\
 & - \frac{3 \left(\frac{ix^3}{3} - 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int PolyLog(2, -e^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{x^3 \tan(a+bx) - \frac{x^4}{4}} + \\
 & \quad \downarrow \text{2720} \\
 & - \frac{3 \left(\frac{ix^3}{3} - 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{x^3 \tan(a+bx) - \frac{x^4}{4}} + \\
 & \quad \downarrow \text{7143} \\
 & - \frac{3 \left(\frac{ix^3}{3} - 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{ix^2 \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{x^3 \tan(a+bx) - \frac{x^4}{4}}
 \end{aligned}$$

input $\operatorname{Int}[x^3 \operatorname{Tan}[a + b*x]^2, x]$

output
$$\begin{aligned}
 & -\frac{1}{4}x^4 - \frac{(3((I/3)*x^3 - (2*I)*((-1/2*I)*x^2*\operatorname{Log}[1 + E^{((2*I)*(a + b*x))}]))/b + (I*((I/2)*x*\operatorname{PolyLog}[2, -E^{((2*I)*(a + b*x))}]))/b - \operatorname{PolyLog}[3, -E^{((2*I)*(a + b*x))}]/(4*b^2))/b + (x^3 \operatorname{Tan}[a + b*x])/b}{x^3 \tan(a + b*x) - \frac{x^4}{4}}
 \end{aligned}$$

Definitions of rubi rules used

rule 15 $\text{Int}[(a_*)*(x_*)^{(m_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a*(x^{(m+1)/(m+1)}), x] /; \text{FreeQ}[\{a, m\}, x] \&& \text{NeQ}[m, -1]$

rule 2620 $\text{Int}[((F_*)^{(g_*)}*((e_*) + (f_*)*(x_**)))^{(n_*)}*((c_*) + (d_*)*(x_*)^{(m_*)}) / ((a_*) + (b_*)*((F_*)^{(g_*)}*((e_*) + (f_*)*(x_**)))^{(n_*)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$

rule 2720 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Simp}[v/D[v, x] \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /; \text{FunctionOfExponentialQ}[u, x] \&& \text{!MatchQ}[u, (w_*)*((a_*)*(v_*)^{(n_*)})^{(m_*)} /; \text{FreeQ}[\{a, m, n\}, x] \&& \text{IntegerQ}[m*n] \&& \text{!MatchQ}[u, E^{(c_*)*((a_*) + (b_*)*x)} * (F_*)[v_]] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{InverseFunctionQ}[F[x]]]$

rule 3011 $\text{Int}[\text{Log}[1 + (e_*)*((F_*)^{(c_*)}*((a_*) + (b_*)*(x_**)))^{(n_*)}] * ((f_*) + (g_*)*(x_*)^{(m_*)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(f + g*x)^m)*(\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*\text{Log}[F])), x] + \text{Simp}[g*(m/(b*c*n*\text{Log}[F])) \text{Int}[(f + g*x)^{(m-1)}*\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&& \text{GtQ}[m, 0]$

rule 3042 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4202 $\text{Int}[((c_*) + (d_*)*(x_*)^{(m_*)}*\text{tan}[(e_*) + (f_*)*(x_*)], x_{\text{Symbol}}] \rightarrow \text{Simp}[I*((c + d*x)^{(m+1)}/(d*(m+1))), x] - \text{Simp}[2*I \text{Int}[(c + d*x)^m*(E^{(2*I*(e + f*x))}/(1 + E^{(2*I*(e + f*x))))], x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{IGtQ}[m, 0]$

rule 4203

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Simplify[b*d*(m/(f*(n - 1))) Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Simplify[b^2 Int[((c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]); FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x]; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.36

method	result
risch	$-\frac{x^4}{4} + \frac{2ix^3}{b(1+e^{2i(bx+a)})} - \frac{2ix^3}{b} + \frac{6ia^2x}{b^3} + \frac{4ia^3}{b^4} + \frac{3x^2 \ln(1+e^{2i(bx+a)})}{b^2} - \frac{3ix \operatorname{polylog}(2, -e^{2i(bx+a)})}{b^3} + \frac{3 \operatorname{polylog}(3, -e^{2i(bx+a)})}{2b^4}$

input `int(x^3*tan(b*x+a)^2,x,method=_RETURNVERBOSE)`

output

```
-1/4*x^4+2*I*x^3/b/(1+exp(2*I*(b*x+a)))-2*I/b*x^3+6*I/b^3*a^2*x+4*I/b^4*a^3+3*x^2*ln(1+exp(2*I*(b*x+a)))/b^2-3*I*x*polylog(2,-exp(2*I*(b*x+a)))/b^3+3/2*polylog(3,-exp(2*I*(b*x+a)))/b^4-6/b^4*a^2*ln(exp(I*(b*x+a)))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 225 vs. $2(83) = 166$.

Time = 0.08 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.30

$$\int x^3 \tan^2(a + bx) dx = \frac{b^4 x^4 - 4 b^3 x^3 \tan(bx + a) - 6 b^2 x^2 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6 b^2 x^2 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6 i b x \text{Li}_2\left(\frac{-i \tan(bx+a)}{\tan(bx+a)^2+1}\right)}{b^4}$$

input `integrate(x^3*tan(b*x+a)^2,x, algorithm="fricas")`

output

$$\begin{aligned} & -\frac{1}{4}(b^4x^4 - 4b^3x^3\tan(bx + a) - 6b^2x^2\log(-2(I\tan(bx + a) - 1)/(\tan(bx + a)^2 + 1)) \\ & - 6b^2x^2\log(-2(-I\tan(bx + a) - 1)/(\tan(bx + a)^2 + 1)) - 6Ib^2x^2\operatorname{dilog}(2(I\tan(bx + a) - 1)/(\tan(bx + a)^2 + 1) + 1) \\ & + 6Ib^2x^2\operatorname{dilog}(2(-I\tan(bx + a) - 1)/(\tan(bx + a)^2 + 1) + 1) - 3\operatorname{polylog}(3, (\tan(bx + a)^2 + 2I\tan(bx + a) - 1)/(\tan(bx + a)^2 + 1)) \\ & - 3\operatorname{polylog}(3, (\tan(bx + a)^2 - 2I\tan(bx + a) - 1)/(\tan(bx + a)^2 + 1))) / b^4 \end{aligned}$$

Sympy [F]

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan^2(a + bx) dx$$

input `integrate(x**3*tan(b*x+a)**2,x)`

output `Integral(x**3*tan(a + b*x)**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 639 vs. $2(83) = 166$.

Time = 0.24 (sec), antiderivative size = 639, normalized size of antiderivative = 6.52

$$\int x^3 \tan^2(a + bx) dx = \text{Too large to display}$$

input `integrate(x^3*tan(b*x+a)^2,x, algorithm="maxima")`

output

$$\begin{aligned} & \frac{1}{2} \cdot (2 \cdot (b \cdot x + a - \tan(b \cdot x + a)) \cdot a^3 - 3 \cdot ((b \cdot x + a)^2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \\ & (b \cdot x + a)^2 \cdot \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot (b \cdot x + a)^2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + (b \cdot x + a)^2 - \\ & (\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) \cdot \operatorname{I} \operatorname{og}(\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) - 4 \cdot (b \cdot x + a) \cdot \sin(2 \cdot b \cdot x + 2 \cdot a) \cdot a^2 / (\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + \\ & 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) + 2 \cdot (\operatorname{I} \cdot (b \cdot x + a)^4 - 4 \cdot \operatorname{I} \cdot (b \cdot x + a)^3 \cdot a + 12 \cdot ((b \cdot x + a)^2 - 2 \cdot (b \cdot x + a) \cdot a + ((b \cdot x + a)^2 - 2 \cdot (b \cdot x + a) \cdot a) \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) - \\ & (-\operatorname{I} \cdot (b \cdot x + a)^2 + 2 \cdot \operatorname{I} \cdot (b \cdot x + a) \cdot a) \cdot \sin(2 \cdot b \cdot x + 2 \cdot a)) \cdot \operatorname{arctan}2(\sin(2 \cdot b \cdot x + 2 \cdot a), \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) + (\operatorname{I} \cdot (b \cdot x + a)^4 - 4 \cdot (b \cdot x + a)^3 \cdot (\operatorname{I} \cdot a + 2) + 24 \cdot (b \cdot x + a)^2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) - 12 \cdot (b \cdot x \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + \operatorname{I} \cdot b \cdot x \cdot \sin(2 \cdot b \cdot x + 2 \cdot a) + b \cdot x) \cdot \operatorname{dilog}(-e^{-(2 \cdot \operatorname{I} \cdot b \cdot x + 2 \cdot \operatorname{I} \cdot a)}) - 6 \cdot (\operatorname{I} \cdot (b \cdot x + a)^2 - 2 \cdot \operatorname{I} \cdot (b \cdot x + a) \cdot a + (\operatorname{I} \cdot (b \cdot x + a)^2 - 2 \cdot \operatorname{I} \cdot (b \cdot x + a) \cdot a) \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) - ((b \cdot x + a)^2 - 2 \cdot (b \cdot x + a) \cdot a) \cdot \sin(2 \cdot b \cdot x + 2 \cdot a)) \cdot \operatorname{log}(\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) - 6 \cdot (\operatorname{I} \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) - \sin(2 \cdot b \cdot x + 2 \cdot a) + \operatorname{I}) \cdot \operatorname{polylog}(3, -e^{-(2 \cdot \operatorname{I} \cdot b \cdot x + 2 \cdot \operatorname{I} \cdot a)}) - ((b \cdot x + a)^4 - 4 \cdot (b \cdot x + a)^3 \cdot (a - 2 \cdot \operatorname{I}) - 24 \cdot \operatorname{I} \cdot (b \cdot x + a)^2 \cdot a) \cdot \sin(2 \cdot b \cdot x + 2 \cdot a) / (-4 \cdot \operatorname{I} \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 4 \cdot \sin(2 \cdot b \cdot x + 2 \cdot a) - 4 \cdot \operatorname{I}) / b^4 \end{aligned}$$

Giac [F]

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan(bx + a)^2 dx$$

input

```
integrate(x^3*tan(b*x+a)^2,x, algorithm="giac")
```

output

```
integrate(x^3*tan(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan(a + bx)^2 dx$$

input

```
int(x^3*tan(a + b*x)^2,x)
```

output `int(x^3*tan(a + b*x)^2, x)`

Reduce [F]

$$\int x^3 \tan^2(a + bx) dx = \frac{-12(\int \tan(bx + a) x^2 dx) + 4 \tan(bx + a) x^3 - b x^4}{4b}$$

input `int(x^3*tan(b*x+a)^2,x)`

output `(- 12*int(tan(a + b*x)*x**2,x) + 4*tan(a + b*x)*x**3 - b*x**4)/(4*b)`

3.7 $\int x^2 \tan^2(a + bx) dx$

Optimal result	91
Mathematica [B] (verified)	91
Rubi [A] (verified)	92
Maple [A] (verified)	94
Fricas [B] (verification not implemented)	95
Sympy [F]	95
Maxima [B] (verification not implemented)	96
Giac [F]	96
Mupad [F(-1)]	97
Reduce [F]	97

Optimal result

Integrand size = 12, antiderivative size = 73

$$\begin{aligned} \int x^2 \tan^2(a + bx) dx = & -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} \\ & - \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} + \frac{x^2 \tan(a + bx)}{b} \end{aligned}$$

output
$$-\text{I}*x^2/b - 1/3*x^3 + 2*x*\ln(1+\exp(2*I*(b*x+a)))/b^2 - \text{I}*\operatorname{polylog}(2, -\exp(2*I*(b*x+a)))/b^3 + x^2*\tan(b*x+a)/b$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 160 vs. $2(73) = 146$.

Time = 4.55 (sec) , antiderivative size = 160, normalized size of antiderivative = 2.19

$$\begin{aligned} \int x^2 \tan^2(a + bx) dx = & -\frac{x^3}{3} + \frac{x^2 \sec(a) \sec(a + bx) \sin(bx)}{b} \\ & + \frac{ibx(\pi + 2 \arctan(\cot(a))) + \pi \log(1 + e^{-2ibx}) + 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))})}{b} \end{aligned}$$

input `Integrate[x^2*Tan[a + b*x]^2, x]`

output

$$\begin{aligned}
 & -\frac{1}{3}x^3 + (x^2 \operatorname{Sec}[a] \operatorname{Sec}[a + bx] \operatorname{Sin}[bx])/b + (I b x (\operatorname{Pi} + 2 \operatorname{ArcTan}[\operatorname{Co} \\
 & t[a]]) + \operatorname{Pi} \operatorname{Log}[1 + E^{(-2 I) b x}] + 2(b x - \operatorname{ArcTan}[\operatorname{Cot}[a]]) \operatorname{Log}[1 - E^{(2 I) \\
 & (b x - \operatorname{ArcTan}[\operatorname{Cot}[a])}] - \operatorname{Pi} \operatorname{Log}[\operatorname{Cos}[bx]] + 2 \operatorname{ArcTan}[\operatorname{Cot}[a]] \operatorname{Log}[\operatorname{S} \\
 & \operatorname{in}[bx - \operatorname{ArcTan}[\operatorname{Cot}[a]]]] - I \operatorname{PolyLog}[2, E^{((2 I) (b x - \operatorname{ArcTan}[\operatorname{Cot}[a]))}] \\
 & + (b^2 x^2 \operatorname{Sqrt}[\operatorname{Csc}[a]^2] \operatorname{Tan}[a])/E^{(I \operatorname{ArcTan}[\operatorname{Cot}[a]]})]/b^3
 \end{aligned}$$

Rubi [A] (verified)

Time = 0.43 (sec), antiderivative size = 87, normalized size of antiderivative = 1.19, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4203, 15, 3042, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \tan^2(a + bx) dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int x^2 \tan(a + bx)^2 dx \\
 & \downarrow \textcolor{blue}{4203} \\
 & -\frac{2 \int x \tan(a + bx) dx}{b} - \int x^2 dx + \frac{x^2 \tan(a + bx)}{b} \\
 & \downarrow \textcolor{blue}{15} \\
 & -\frac{2 \int x \tan(a + bx) dx}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \downarrow \textcolor{blue}{3042} \\
 & -\frac{2 \int x \tan(a + bx) dx}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \downarrow \textcolor{blue}{4202} \\
 & -\frac{2 \left(\frac{i x^2}{2} - 2 i \int \frac{e^{2i(a+bx)x}}{1+e^{2i(a+bx)}} dx \right)}{b} + \frac{x^2 \tan(a + bx)}{b} - \frac{x^3}{3} \\
 & \downarrow \textcolor{blue}{2620}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{2\left(\frac{ix^2}{2} - 2i\left(\frac{i \int \log(1+e^{2i(a+bx)})dx}{2b} - \frac{ix \log(1+e^{2i(a+bx)})}{2b}\right)\right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \\
 & \quad \downarrow \text{2715} \\
 & -\frac{2\left(\frac{ix^2}{2} - 2i\left(\frac{\int e^{-2i(a+bx)} \log(1+e^{2i(a+bx)})de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b}\right)\right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \\
 & \quad \downarrow \text{2838} \\
 & -\frac{2\left(\frac{ix^2}{2} - 2i\left(-\frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b}\right)\right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3}
 \end{aligned}$$

input `Int[x^2*Tan[a + b*x]^2, x]`

output `-1/3*x^3 - (2*((I/2)*x^2 - (2*I)*((-1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))]) /b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2)))/b + (x^2*Tan[a + b*x])/b`

Definitions of rubi rules used

rule 15 `Int[(a_)*(x_)^(m_), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2620 `Int[((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*(c_*) + (d_)*(x_)^(m_))/((a_) + (b_)*(F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*(F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*(d_ + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 $\text{Int}[u_-, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4202 $\text{Int}[(c_-) + (d_-)*(x_-)^m * \tan[(e_-) + (f_-)*(x_-)], x_{\text{Symbol}}] \rightarrow \text{Simp}[I * ((c + d*x)^{m+1}) / (d*(m+1)), x] - \text{Simp}[2*I \text{Int}[(c + d*x)^m * (E^{(2*I*(e + f*x))} / (1 + E^{(2*I*(e + f*x))})), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{IGtQ}[m, 0]$

rule 4203 $\text{Int}[(c_-) + (d_-)*(x_-)^m * ((b_-)*\tan[(e_-) + (f_-)*(x_-)])^n, x_{\text{Symbol}}] \rightarrow \text{Simp}[b*(c + d*x)^m * ((b*\text{Tan}[e + f*x])^{n-1}) / (f*(n-1)), x] + (-\text{Simp}[b*d*(m/(f*(n-1))) \text{Int}[(c + d*x)^{m-1} * (b*\text{Tan}[e + f*x])^{n-1}, x] - \text{Simp}[b^2 \text{Int}[(c + d*x)^{m-2} * (b*\text{Tan}[e + f*x])^{n-2}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&& \text{GtQ}[n, 1] \&& \text{GtQ}[m, 0]$

Maple [A] (verified)

Time = 0.31 (sec), antiderivative size = 108, normalized size of antiderivative = 1.48

method	result	size
risch	$-\frac{x^3}{3} + \frac{2ix^2}{b(1+e^{2i(bx+a)})} - \frac{2ix^2}{b} - \frac{4iax}{b^2} - \frac{2ia^2}{b^3} + \frac{2x \ln(1+e^{2i(bx+a)})}{b^2} - \frac{i \text{polylog}(2, -e^{2i(bx+a)})}{b^3} + \frac{4a \ln(e^{i(bx+a)})}{b^3}$	108

input $\text{int}(x^2 * \tan(b*x+a)^2, x, \text{method}=\text{RETURNVERBOSE})$

output
$$\begin{aligned} & -\frac{1}{3}x^3 + 2*I*x^2/b/(1+\exp(2*I*(b*x+a))) - 2*I/b*x^2 - 4*I/b^2*a*x - 2*I/b^3*a^2 + \\ & 2*x*\ln(1+\exp(2*I*(b*x+a)))/b^2 - I*\text{polylog}(2, -\exp(2*I*(b*x+a)))/b^3 + 4/b^3*a*\ln(\exp(I*(b*x+a))) \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 144 vs. $2(62) = 124$.

Time = 0.08 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.97

$$\int x^2 \tan^2(a + bx) dx =$$

$$\frac{2 b^3 x^3 - 6 b^2 x^2 \tan(bx + a) - 6 b x \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 6 b x \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) - 3 i \text{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 3 i \text{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right)}{6 b^3}$$

input `integrate(x^2*tan(b*x+a)^2,x, algorithm="fricas")`

output
$$\begin{aligned} & -1/6*(2*b^3*x^3 - 6*b^2*x^2*tan(b*x + a) - 6*b*x*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 6*b*x*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 3*I*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) + 3*I*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1))/b^3 \end{aligned}$$

Sympy [F]

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan^2(a + bx) dx$$

input `integrate(x**2*tan(b*x+a)**2,x)`

output `Integral(x**2*tan(a + b*x)**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 257 vs. $2(62) = 124$.

Time = 0.24 (sec) , antiderivative size = 257, normalized size of antiderivative = 3.52

$$\int x^2 \tan^2(a + bx) dx \\ = \frac{i b^3 x^3 + 6 (bx \cos(2bx + 2a) + i bx \sin(2bx + 2a) + bx) \arctan(\sin(2bx + 2a), \cos(2bx + 2a) + 1) +}{}$$

input `integrate(x^2*tan(b*x+a)^2,x, algorithm="maxima")`

output
$$(I*b^3*x^3 + 6*(b*x*cos(2*b*x + 2*a) + I*b*x*sin(2*b*x + 2*a) + b*x)*arctan2(\sin(2*b*x + 2*a), \cos(2*b*x + 2*a) + 1) + (I*b^3*x^3 - 6*b^2*x^2)*cos(2*b*x + 2*a) - 3*(cos(2*b*x + 2*a) + I*sin(2*b*x + 2*a) + 1)*dilog(-e^(2*I*b*x + 2*I*a)) - 3*(I*b*x*cos(2*b*x + 2*a) - b*x*sin(2*b*x + 2*a) + I*b*x)*log(\cos(2*b*x + 2*a)^2 + \sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - (b^3*x^3 + 6*I*b^2*x^2)*sin(2*b*x + 2*a))/(-3*I*b^3*cos(2*b*x + 2*a) + 3*b^3*sin(2*b*x + 2*a) - 3*I*b^3)$$

Giac [F]

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan(bx + a)^2 dx$$

input `integrate(x^2*tan(b*x+a)^2,x, algorithm="giac")`

output `integrate(x^2*tan(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan(a + bx)^2 dx$$

input `int(x^2*tan(a + b*x)^2, x)`

output `int(x^2*tan(a + b*x)^2, x)`

Reduce [F]

$$\int x^2 \tan^2(a + bx) dx = \frac{-6(\int \tan(bx + a) x dx) + 3 \tan(bx + a) x^2 - b x^3}{3b}$$

input `int(x^2*tan(b*x+a)^2, x)`

output `(- 6*int(tan(a + b*x)*x, x) + 3*tan(a + b*x)*x**2 - b*x**3)/(3*b)`

3.8 $\int x \tan^2(a + bx) dx$

Optimal result	98
Mathematica [A] (verified)	98
Rubi [A] (verified)	99
Maple [A] (verified)	100
Fricas [A] (verification not implemented)	101
Sympy [A] (verification not implemented)	101
Maxima [B] (verification not implemented)	101
Giac [B] (verification not implemented)	102
Mupad [B] (verification not implemented)	103
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 10, antiderivative size = 30

$$\int x \tan^2(a + bx) dx = -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b}$$

output -1/2*x^2+ln(cos(b*x+a))/b^2+x*tan(b*x+a)/b

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.43

$$\int x \tan^2(a + bx) dx = -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \sec(a) \sec(a + bx) \sin(bx)}{b} + \frac{x \tan(a)}{b}$$

input Integrate[x*Tan[a + b*x]^2, x]

output -1/2*x^2 + Log[Cos[a + b*x]]/b^2 + (x*Sec[a]*Sec[a + b*x]*Sin[b*x])/b + (x*Tan[a])/b

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4203, 15, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tan^2(a + bx) dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int x \tan(a + bx)^2 dx \\
 & \downarrow \textcolor{blue}{4203} \\
 & -\frac{\int \tan(a + bx) dx}{b} - \int x dx + \frac{x \tan(a + bx)}{b} \\
 & \downarrow \textcolor{blue}{15} \\
 & -\frac{\int \tan(a + bx) dx}{b} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2} \\
 & \downarrow \textcolor{blue}{3042} \\
 & -\frac{\int \tan(a + bx) dx}{b} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2} \\
 & \downarrow \textcolor{blue}{3956} \\
 & \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2}
 \end{aligned}$$

input `Int[x*Tan[a + b*x]^2,x]`

output `-1/2*x^2 + Log[Cos[a + b*x]]/b^2 + (x*Tan[a + b*x])/b`

Definitions of rubi rules used

rule 15 $\text{Int}[(a_*)*(x_)^{(m_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[a*(x^{(m+1)/(m+1)}), x] /; \text{FreeQ}[\{a, m\}, x] \& \text{NeQ}[m, -1]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\tan[(c_*) + (d_*)*(x_)], x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\cos[c+d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4203 $\text{Int}[((c_*) + (d_*)*(x_))^{(m_*)}*((b_*)*\tan[(e_*) + (f_*)*(x_)])^{(n_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[b*(c+d*x)^m*((b*Tan[e+f*x])^{(n-1)}/(f*(n-1))), x] + (-\text{Simp}[b*d*(m/(f*(n-1))) \cdot \text{Int}[(c+d*x)^{(m-1)}*(b*Tan[e+f*x])^{(n-1)}, x], x] - \text{Simp}[b^2 \cdot \text{Int}[(c+d*x)^{m-2}*(b*Tan[e+f*x])^{(n-2)}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \& \text{GtQ}[n, 1] \& \text{GtQ}[m, 0]$

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

method	result	size
norman	$\frac{x \tan(bx+a)}{b} - \frac{x^2}{2} - \frac{\ln(1+\tan(bx+a)^2)}{2b^2}$	34
parallelrisch	$-\frac{x^2 b^2 - 2 \tan(bx+a)bx + \ln(1+\tan(bx+a)^2)}{2b^2}$	35
default	$-\frac{x^2}{2} + \frac{(bx+a)\tan(bx+a)+\ln(\cos(bx+a))-a\tan(bx+a)}{b^2}$	40
risch	$-\frac{x^2}{2} - \frac{2ix}{b} - \frac{2ia}{b^2} + \frac{2ix}{b(1+e^{2i(bx+a)})} + \frac{\ln(1+e^{2i(bx+a)})}{b^2}$	57

input `int(x*tan(b*x+a)^2, x, method=_RETURNVERBOSE)`

output $x*\tan(b*x+a)/b - 1/2*x^2 - 1/2/b^2*\ln(1+\tan(b*x+a)^2)$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int x \tan^2(a + bx) dx = -\frac{b^2 x^2 - 2 b x \tan(bx + a) - \log\left(\frac{1}{\tan(bx+a)^2+1}\right)}{2 b^2}$$

input `integrate(x*tan(b*x+a)^2,x, algorithm="fricas")`

output `-1/2*(b^2*x^2 - 2*b*x*tan(b*x + a) - log(1/(tan(b*x + a)^2 + 1)))/b^2`

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37

$$\int x \tan^2(a + bx) dx = \begin{cases} -\frac{x^2}{2} + \frac{x \tan(a+bx)}{b} - \frac{\log(\tan^2(a+bx)+1)}{2b^2} & \text{for } b \neq 0 \\ \frac{x^2 \tan^2(a)}{2} & \text{otherwise} \end{cases}$$

input `integrate(x*tan(b*x+a)**2,x)`

output `Piecewise((-x**2/2 + x*tan(a + b*x)/b - log(tan(a + b*x)**2 + 1)/(2*b**2), Ne(b, 0)), (x**2*tan(a)**2/2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 214 vs. $2(28) = 56$.

Time = 0.15 (sec) , antiderivative size = 214, normalized size of antiderivative = 7.13

$$\begin{aligned} & \int x \tan^2(a + bx) dx \\ &= \frac{2 (bx + a - \tan(bx + a))a - \frac{(bx+a)^2 \cos(2bx+2a)^2 + (bx+a)^2 \sin(2bx+2a)^2 + 2(bx+a)^2 \cos(2bx+2a) + (bx+a)^2 - (\cos(2bx+2a)^2 - \cos(2bx+2a)^2)}{\cos(2bx+2a)^2}}{2b^2} \end{aligned}$$

input `integrate(x*tan(b*x+a)^2,x, algorithm="maxima")`

output

$$\begin{aligned} & \frac{1}{2} \cdot (2 \cdot (b \cdot x + a - \tan(b \cdot x + a)) \cdot a - ((b \cdot x + a)^2 \cos(2 \cdot b \cdot x + 2 \cdot a)^2 + (b \cdot x \\ & + a)^2 \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot (b \cdot x + a)^2 \cos(2 \cdot b \cdot x + 2 \cdot a) + (b \cdot x + a)^2 \\ & - (\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) \cdot \log(\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1) - 4 \cdot (b \cdot x \\ & + a) \cdot \sin(2 \cdot b \cdot x + 2 \cdot a)) / (\cos(2 \cdot b \cdot x + 2 \cdot a)^2 + \sin(2 \cdot b \cdot x + 2 \cdot a)^2 + 2 \cdot \cos(2 \cdot b \cdot x + 2 \cdot a) + 1)) / b^2 \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(28) = 56$.

Time = 0.40 (sec) , antiderivative size = 162, normalized size of antiderivative = 5.40

$$\int x \tan^2(a + bx) dx = \frac{b^2 x^2 \tan(bx) \tan(a) - b^2 x^2 + 2bx \tan(bx) + 2bx \tan(a) - \log\left(\frac{4(\tan(bx)^2 \tan(a)^2 - 2\tan(bx) \tan(a) + 1)}{\tan(bx)^2 \tan(a)^2 + \tan(bx)^2 + \tan(a)^2 + 1}\right) \tan(bx) \tan(a)}{2(b^2 \tan(bx) \tan(a) - b^2)}$$

input `integrate(x*tan(b*x+a)^2,x, algorithm="giac")`

output

$$\begin{aligned} & -\frac{1}{2} \cdot (b^2 x^2 \tan(bx) \tan(a) - b^2 x^2 + 2bx \tan(bx) + 2bx \tan(a) - \log(4 \cdot (\tan(bx)^2 \tan(a)^2 - 2 \cdot \tan(bx) \tan(a) + 1) / (\tan(bx)^2 \tan(a)^2 + \tan(bx)^2 + \tan(a)^2 + 1))) \cdot \tan(bx) \tan(a) + \log(4 \cdot (\tan(bx)^2 \tan(a)^2 - 2 \cdot \tan(bx) \tan(a) + 1) / (\tan(bx)^2 \tan(a)^2 + \tan(bx)^2 + \tan(a)^2 + 1))) / (b^2 \tan(bx) \tan(a) - b^2) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int x \tan^2(a + bx) dx = -\frac{\ln(\tan(a+bx)^2+1)}{2} - \frac{bx \tan(a + bx)}{b^2} - \frac{x^2}{2}$$

input `int(x*tan(a + b*x)^2,x)`

output `- (log(tan(a + b*x)^2 + 1)/2 - b*x*tan(a + b*x))/b^2 - x^2/2`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.23

$$\int x \tan^2(a + bx) dx = \frac{-\log(\tan(bx+a)^2 + 1) + 2 \tan(bx+a) bx - b^2 x^2}{2b^2}$$

input `int(x*tan(b*x+a)^2,x)`

output `(- log(tan(a + b*x)**2 + 1) + 2*tan(a + b*x)*b*x - b**2*x**2)/(2*b**2)`

3.9 $\int \frac{\tan^2(a+bx)}{x} dx$

Optimal result	104
Mathematica [N/A]	104
Rubi [N/A]	105
Maple [N/A]	106
Fricas [N/A]	106
Sympy [N/A]	107
Maxima [N/A]	107
Giac [N/A]	108
Mupad [N/A]	108
Reduce [N/A]	108

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^2(a + bx)}{x} dx = \text{Int}\left(\frac{\tan^2(a + bx)}{x}, x\right)$$

output `Defer(Int)(tan(b*x+a)^2/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.88 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(a + bx)}{x} dx$$

input `Integrate[Tan[a + b*x]^2/x,x]`

output `Integrate[Tan[a + b*x]^2/x, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\tan^2(a + bx)}{x} dx \\ \downarrow \textcolor{blue}{3042} \\ \int \frac{\tan(a + bx)^2}{x} dx \\ \downarrow \textcolor{blue}{4222} \\ \int \frac{\tan^2(a + bx)}{x} dx \end{array}$$

input `Int[Tan[a + b*x]^2/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4222 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)}, x_\text{Symbol}] \rightarrow \text{Simp}[\text{If}[\text{MatchQ}[f, (f1_.)*(Complex[0, j_])], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Coth}[-I]*(e - \text{Pi}/2) - I*f*x]^n, x], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Tanh}[-I]*e - I*f*x]^n, x]], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], (-1)^n* \text{Unintegatable}[(c + d*x)^m*\text{Cot}[e - \text{Pi}/2 + f*x]^n, x], \text{Unintegatable}[(c + d*x)^m*\text{Tan}[e + f*x]^n, x]]], x] /; \text{FreeQ}[\{c, d, e, f, m, n\}, x] \&& \text{IntegerQ}[n]$

Maple [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx+a)^2}{x} dx$$

input $\text{int}(\tan(b*x+a)^2/x, x)$

output $\text{int}(\tan(b*x+a)^2/x, x)$

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

input $\text{integrate}(\tan(b*x+a)^2/x, x, \text{algorithm}=\text{"fricas"})$

output `integral(tan(b*x + a)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(a + bx)}{x} dx$$

input `integrate(tan(b*x+a)**2/x,x)`

output `Integral(tan(a + b*x)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 243, normalized size of antiderivative = 20.25

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(bx + a)}{x} dx$$

input `integrate(tan(b*x+a)^2/x,x, algorithm="maxima")`

output `-(b*x*cos(2*b*x + 2*a)^2*log(x) + b*x*log(x)*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a)*log(x) + b*x*log(x) - 2*(b^2*x*cos(2*b*x + 2*a)^2 + b^2*x*sin(2*b*x + 2*a)^2 + 2*b^2*x*cos(2*b*x + 2*a) + b^2*x*sin(2*b*x + 2*a))/((b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(2*b*x + 2*a)^2 + 2*b^2*x*cos(2*b*x + 2*a) + b^2*x*sin(2*b*x + 2*a)), x) - 2*sin(2*b*x + 2*a))/(b*x*cos(2*b*x + 2*a)^2 + b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + b*x)`

Giac [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

input `integrate(tan(b*x+a)^2/x,x, algorithm="giac")`

output `integrate(tan(b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 8.72 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(a + b x)^2}{x} dx$$

input `int(tan(a + b*x)^2/x,x)`

output `int(tan(a + b*x)^2/x, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

input `int(tan(b*x+a)^2/x,x)`

```
output int(tan(a + b*x)**2/x,x)
```

3.10 $\int \frac{\tan^2(a+bx)}{x^2} dx$

Optimal result	110
Mathematica [N/A]	110
Rubi [N/A]	111
Maple [N/A]	112
Fricas [N/A]	112
Sympy [N/A]	113
Maxima [N/A]	113
Giac [N/A]	114
Mupad [N/A]	114
Reduce [N/A]	114

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan^2(a + bx)}{x^2}, x\right)$$

output `Defer(Int)(tan(b*x+a)^2/x^2,x)`

Mathematica [N/A]

Not integrable

Time = 2.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

input `Integrate[Tan[a + b*x]^2/x^2,x]`

output `Integrate[Tan[a + b*x]^2/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\tan^2(a + bx)}{x^2} dx \\ \downarrow \textcolor{blue}{3042} \\ \int \frac{\tan(a + bx)^2}{x^2} dx \\ \downarrow \textcolor{blue}{4222} \\ \int \frac{\tan^2(a + bx)}{x^2} dx \end{array}$$

input `Int[Tan[a + b*x]^2/x^2, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4222 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)}, x_\text{Symbol}] \rightarrow \text{Simp}[\text{If}[\text{MatchQ}[f, (f1_.)*(Complex[0, j_])], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Coth}[-I*(e - \text{Pi}/2) - I*f*x]^n, x], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Tanh}[-I*(e - \text{Pi}/2) - I*f*x]^n, x]], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], (-1)^n* \text{Unintegatable}[(c + d*x)^m*\text{Cot}[e - \text{Pi}/2 + f*x]^n, x], \text{Unintegatable}[(c + d*x)^m*\text{Tan}[e + f*x]^n, x]]], x] /; \text{FreeQ}[\{c, d, e, f, m, n\}, x] \&& \text{IntegerQ}[n]$

Maple [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx+a)^2}{x^2} dx$$

input $\text{int}(\tan(b*x+a)^2/x^2, x)$

output $\text{int}(\tan(b*x+a)^2/x^2, x)$

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

input $\text{integrate}(\tan(b*x+a)^2/x^2, x, \text{algorithm}=\text{"fricas"})$

output `integral(tan(b*x + a)^2/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

input `integrate(tan(b*x+a)**2/x**2,x)`

output `Integral(tan(a + b*x)**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 250, normalized size of antiderivative = 20.83

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(bx + a)}{x^2} dx$$

input `integrate(tan(b*x+a)^2/x^2,x, algorithm="maxima")`

output `(b*x*cos(2*b*x + 2*a)^2 + b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + b*x + 4*(b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(2*b*x + 2*a)^2 + 2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2)*integrate(sin(2*b*x + 2*a)/(b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(2*b*x + 2*a)^2 + 2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3), x) + 2*sin(2*b*x + 2*a))/(b*x^2*cos(2*b*x + 2*a)^2 + b*x^2*sin(2*b*x + 2*a)^2 + 2*b*x^2*cos(2*b*x + 2*a) + b*x^2)`

Giac [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

input `integrate(tan(b*x+a)^2/x^2,x, algorithm="giac")`

output `integrate(tan(b*x + a)^2/x^2, x)`

Mupad [N/A]

Not integrable

Time = 8.34 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(a + b x)^2}{x^2} dx$$

input `int(tan(a + b*x)^2/x^2,x)`

output `int(tan(a + b*x)^2/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

input `int(tan(b*x+a)^2/x^2,x)`

output int(tan(a + b*x)**2/x**2,x)

3.11 $\int x^3 \tan^3(a + bx) dx$

Optimal result	116
Mathematica [A] (warning: unable to verify)	117
Rubi [A] (verified)	117
Maple [A] (verified)	123
Fricas [B] (verification not implemented)	124
Sympy [F]	125
Maxima [B] (verification not implemented)	125
Giac [F]	126
Mupad [F(-1)]	127
Reduce [F]	127

Optimal result

Integrand size = 12, antiderivative size = 205

$$\begin{aligned} \int x^3 \tan^3(a + bx) dx = & \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} \\ & + \frac{3i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^4} - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\ & + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} + \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} \\ & - \frac{3x^2 \tan(a + bx)}{2b^2} + \frac{x^3 \tan^2(a + bx)}{2b} \end{aligned}$$

output

```
3/2*I*x^2/b^2+1/2*x^3/b-1/4*I*x^4-3*x*ln(1+exp(2*I*(b*x+a)))/b^3+x^3*ln(1+exp(2*I*(b*x+a)))/b+3/2*I*polylog(2,-exp(2*I*(b*x+a)))/b^4-3/2*I*x^2*polylog(2,-exp(2*I*(b*x+a)))/b^2+3/2*x*polylog(3,-exp(2*I*(b*x+a)))/b^3+3/4*I*polylog(4,-exp(2*I*(b*x+a)))/b^4-3/2*x^2*tan(b*x+a)/b^2+1/2*x^3*tan(b*x+a)^2/b
```

Mathematica [A] (warning: unable to verify)

Time = 6.48 (sec) , antiderivative size = 370, normalized size of antiderivative = 1.80

$$\begin{aligned} & \int x^3 \tan^3(a + bx) dx \\ &= \frac{i e^{ia} (2b^4 e^{-2ia} x^4 - 4ib^3 (1 + e^{-2ia}) x^3 \log(1 + e^{-2i(a+bx)}) + 6b^2 (1 + e^{-2ia}) x^2 \text{PolyLog}(2, -e^{-2i(a+bx)}) - 6a^2 b^2 x^2 \text{PolyLog}(3, -e^{-2i(a+bx)}) + 3b^4 x^4 \text{PolyLog}(4, -e^{-2i(a+bx)}))}{8b^4} \\ &+ \frac{x^3 \sec^2(a + bx)}{2b} \\ &- \frac{3 \csc(a) \left(b^2 e^{-i \arctan(\cot(a))} x^2 - \frac{\cot(a) (ibx(-\pi - 2 \arctan(\cot(a))) - \pi \log(1 + e^{-2ibx}) - 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))}))}{2b^4 \sqrt{\csc^2(a) (\cos^2(a) + \sin^2(a))}} \right)}{2b^2} \\ &- \frac{3x^2 \sec(a) \sec(a + bx) \sin(bx)}{2b^2} - \frac{1}{4} x^4 \tan(a) \end{aligned}$$

input `Integrate[x^3*Tan[a + b*x]^3, x]`

output
$$\begin{aligned} & ((I/8)*E^{(I*a)*((2*b^4*x^4)/E^{((2*I)*a}) - (4*I)*b^3*(1 + E^{((-2*I)*a})*x^3 * \text{Log}[1 + E^{((-2*I)*(a + b*x))}] + 6*b^2*(1 + E^{((-2*I)*a})*x^2*\text{PolyLog}[2, -E^{((-2*I)*(a + b*x))}] - (6*I)*b*(1 + E^{((-2*I)*a})*x*\text{PolyLog}[3, -E^{((-2*I)*(a + b*x))}] - 3*(1 + E^{((-2*I)*a})*\text{PolyLog}[4, -E^{((-2*I)*(a + b*x))}])*\text{Sec}[a])/b^4 + (x^3*\text{Sec}[a + b*x]^2)/(2*b) - (3*\text{Csc}[a]*(b^2*x^2)/E^{(I*\text{ArcTan}[\text{Cot}[a]])} - (\text{Cot}[a]*(I*b*x*(-Pi - 2*\text{ArcTan}[\text{Cot}[a]])) - \text{Pi}*\text{Log}[1 + E^{((-2*I)*b*x)}] - 2*(b*x - \text{ArcTan}[\text{Cot}[a]])*\text{Log}[1 - E^{((2*I)*(b*x - \text{ArcTan}[\text{Cot}[a]))}]) + \text{Pi}*\text{Log}[\text{Cos}[b*x]] - 2*\text{ArcTan}[\text{Cot}[a]]*\text{Log}[\text{Sin}[b*x - \text{ArcTan}[\text{Cot}[a]]]] + I*\text{PolyLog}[2, E^{((2*I)*(b*x - \text{ArcTan}[\text{Cot}[a]))}])])/\text{Sqrt}[1 + \text{Cot}[a]^2]*\text{Sec}[a])/(2*b^4*\text{Sqrt}[\text{Csc}[a]^2*(\text{Cos}[a]^2 + \text{Sin}[a]^2)]) - (3*x^2*\text{Sec}[a]*\text{Sec}[a + b*x]*\text{Sin}[b*x])/(2*b^2) - (x^4*\text{Tan}[a])/4 \end{aligned}$$

Rubi [A] (verified)

Time = 1.35 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.20, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 1.333, Rules used = {3042, 4203, 3042, 4202, 2620, 3011, 4203, 15, 3042, 4202, 2620, 2715, 2838, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x^3 \tan^3(a + bx) dx \\
& \quad \downarrow \textcolor{blue}{3042} \\
& \int x^3 \tan(a + bx)^3 dx \\
& \quad \downarrow \textcolor{blue}{4203} \\
& - \int x^3 \tan(a + bx) dx - \frac{3 \int x^2 \tan^2(a + bx) dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} \\
& \quad \downarrow \textcolor{blue}{3042} \\
& - \int x^3 \tan(a + bx) dx - \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} \\
& \quad \downarrow \textcolor{blue}{4202} \\
& 2i \int \frac{e^{2i(a+bx)} x^3}{1 + e^{2i(a+bx)}} dx - \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{2620} \\
& - \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + 2i \left(\frac{3i \int x^2 \log(1 + e^{2i(a+bx)}) dx}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) + \\
& \quad \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{3011} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \int x^2 \tan(a + bx)^2 dx}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{4203} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \left(-\frac{2 \int x \tan(a + bx) dx}{b} - \int x^2 dx + \frac{x^2 \tan(a + bx)}{b} \right)}{2b} + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{15}
\end{aligned}$$

$$\begin{aligned}
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \left(-\frac{2 \int x \tan(a+bx) dx}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{2b} + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{3042} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \left(-\frac{2 \int x \tan(a+bx) dx}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{2b} + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{4202} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \left(-\frac{2 \left(\frac{ix^2}{2} - 2i \int \frac{e^{2i(a+bx)} x}{1+e^{2i(a+bx)}} dx \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{2b} + \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{2620} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \quad \frac{3 \left(-\frac{2 \left(\frac{ix^2}{2} - 2i \left(\frac{i \int \log(1+e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{2b} + \\
& \quad \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \textcolor{blue}{2715}
\end{aligned}$$

$$\begin{aligned}
& - \frac{3 \left(- \frac{2 \left(\frac{ix^2}{2} - 2i \left(\frac{\int e^{-2i(a+bx)} \log(1+e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{} + \\
& 2i \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1+e^{2i(a+bx)})}{2b} \right) + \\
& \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{2838} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b} \right)}{2b} - \frac{ix^3 \log(1+e^{2i(a+bx)})}{2b} \right) - \\
& 3 \left(- \frac{2 \left(\frac{ix^2}{2} - 2i \left(- \frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \\
& \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{7163} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left(\frac{i \int \text{PolyLog}(3, -e^{2i(a+bx)}) dx}{2b} - \frac{ix \text{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1+e^{2i(a+bx)})}{2b} \right) - \\
& 3 \left(- \frac{2 \left(\frac{ix^2}{2} - 2i \left(- \frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right) + \\
& \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \\
& \quad \downarrow \text{2720}
\end{aligned}$$

$$\begin{aligned}
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left(\frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(3, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right. \\
& \quad \left. + \frac{3 \left(-\frac{2 \left(\frac{ix^2}{2} - 2i \left(-\frac{\operatorname{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{2b} + \right. \\
& \quad \left. \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \right) \\
& \quad \downarrow \text{7143} \\
& 2i \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \left(\frac{\operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^2} - \frac{ix \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b} \right)}{b} \right)}{2b} - \frac{ix^3 \log(1 + e^{2i(a+bx)})}{2b} \right) \\
& \quad \left. + \frac{3 \left(-\frac{2 \left(\frac{ix^2}{2} - 2i \left(-\frac{\operatorname{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1 + e^{2i(a+bx)})}{2b} \right) \right)}{b} + \frac{x^2 \tan(a+bx)}{b} - \frac{x^3}{3} \right)}{2b} + \right. \\
& \quad \left. \frac{x^3 \tan^2(a+bx)}{2b} - \frac{ix^4}{4} \right)
\end{aligned}$$

input `Int[x^3*Tan[a + b*x]^3, x]`

output

```

(-1/4*I)*x^4 + (2*I)*((-1/2*I)*x^3*Log[1 + E^((2*I)*(a + b*x))])/b + (((3
*I)/2)*((I/2)*x^2*PolyLog[2, -E^((2*I)*(a + b*x))])/b - (I*((-1/2*I)*x*P
olyLog[3, -E^((2*I)*(a + b*x))])/b + PolyLog[4, -E^((2*I)*(a + b*x))]/(4*b
^2))/b) + (x^3*Tan[a + b*x]^2)/(2*b) - (3*(-1/3*x^3 - (2*((I/2)*x^2 -
(2*I)*((-1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))])/b - PolyLog[2, -E^((2*I)
*(a + b*x))]/(4*b^2)))/b + (x^2*Tan[a + b*x])/b))/(2*b)

```

Definitions of rubi rules used

rule 15 $\text{Int}[(a_*)*(x_)^{(m_)}, \ x_\text{Symbol}] \rightarrow \text{Simp}[a*(x^{(m+1)/(m+1)}), \ x] \ /; \ \text{FreeQ}[\{a, \ m\}, \ x] \ \&& \ \text{NeQ}[m, -1]$

rule 2620 $\text{Int}[((F_*)^{(g_*)}*((e_*) + (f_*)*(x_)))^{(n_*)}*((c_*) + (d_*)*(x_))^{(m_)})/((a_) + (b_*)*((F_*)^{(g_*)}*((e_*) + (f_*)*(x_)))^{(n_*)}), \ x_\text{Symbol}] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], \ x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \ \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], \ x], \ x] \ /; \ \text{FreeQ}[\{F, \ a, \ b, \ c, \ d, \ e, \ f, \ g, \ n\}, \ x] \ \&& \ \text{IGtQ}[m, 0]$

rule 2715 $\text{Int}[\text{Log}[(a_) + (b_*)*((F_*)^{(e_*)}*((c_*) + (d_*)*(x_)))^{(n_*)}], \ x_\text{Symbol}] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \ \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, \ x], \ x, (F^(e*(c + d*x)))^n], \ x] \ /; \ \text{FreeQ}[\{F, \ a, \ b, \ c, \ d, \ e, \ n\}, \ x] \ \&& \ \text{GtQ}[a, 0]$

rule 2720 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, \ x]\}, \ \text{Simp}[v/D[v, \ x] \ \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, \ x]/x, \ x], \ x, v], \ x]] \ /; \ \text{FunctionOfExponentialQ}[u, \ x] \ \&& \ !\text{MatchQ}[u, (w_)*(a_)*(v_)^{(n_)})^{(m_)} \ /; \ \text{FreeQ}[\{a, \ m, \ n\}, \ x] \ \&& \ \text{IntegerQ}[m*n] \ \&& \ !\text{MatchQ}[u, E^{(c_)*(a_ + (b_*)*x)} * (F_)[v_]] \ /; \ \text{FreeQ}[\{a, \ b, \ c\}, \ x] \ \&& \ \text{InverseFunctionQ}[F[x]]]$

rule 2838 $\text{Int}[\text{Log}[(c_*)*((d_*) + (e_*)*(x_))^{(n_)})]/(x_), \ x_\text{Symbol}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, \ x] \ /; \ \text{FreeQ}[\{c, \ d, \ e, \ n\}, \ x] \ \&& \ \text{EqQ}[c*d, 1]$

rule 3011 $\text{Int}[\text{Log}[1 + (e_*)*((F_*)^{(c_*)}*((a_*) + (b_*)*(x_)))^{(n_*)}]*((f_*) + (g_*)*(x_))^{(m_)}, \ x_\text{Symbol}] \rightarrow \text{Simp}[(-(f + g*x)^m)*(\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*\text{Log}[F])), \ x] + \text{Simp}[g*(m/(b*c*n*\text{Log}[F])) \ \text{Int}[(f + g*x)^{(m-1)}*\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n], \ x], \ x] \ /; \ \text{FreeQ}[\{F, \ a, \ b, \ c, \ e, \ f, \ g, \ n\}, \ x] \ \&& \ \text{GtQ}[m, 0]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] \ /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4202 $\text{Int}[(c_+ + d_+)(x_+)^m \tan[(e_+ + f_+)(x_+)], x] \rightarrow \text{Simp}[I * ((c + d*x)^{m+1}/(d*(m+1))), x] - \text{Simp}[2*I \text{Int}[(c + d*x)^m (E^{(2*I)*(e+f*x)})/(1 + E^{(2*I)*(e+f*x)})), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{IGtQ}[m, 0]$

rule 4203 $\text{Int}[(c_+ + d_+)(x_+)^m ((b_+ \tan[e+f*x])^n), x] \rightarrow \text{Simp}[b*(c + d*x)^m ((b*Tan[e + f*x])^{n-1}/(f*(n-1))), x] + (-\text{Simp}[b*d*(m/(f*(n-1))) \text{Int}[(c + d*x)^{m-1} ((b*Tan[e + f*x])^{n-1}), x], x] - \text{Simp}[b^2 \text{Int}[(c + d*x)^m ((b*Tan[e + f*x])^{n-2}), x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&& \text{GtQ}[n, 1] \&& \text{GtQ}[m, 0]$

rule 7143 $\text{Int}[\text{PolyLog}[n, (c_+ + b_+)(x_+)^p]/((d_+ + e_+)(x_+)), x] \rightarrow \text{Simp}[\text{PolyLog}[n+1, c*(a+b*x)^p]/(e*p), x] /; \text{FreeQ}[\{a, b, c, d, e, n, p\}, x] \&& \text{EqQ}[b*d, a*e]$

rule 7163 $\text{Int}[(e_+ + f_+)(x_+)^m \text{PolyLog}[n, (d_+)((c_+ + b_+)(x_+))^p], x] \rightarrow \text{Simp}[(e + f*x)^m (\text{PolyLog}[n+1, d*(F^(c*(a+b*x))^p)/(b*c*p*\text{Log}[F])], x) - \text{Simp}[f*(m/(b*c*p*\text{Log}[F])) \text{Int}[(e + f*x)^{m-1} \text{PolyLog}[n+1, d*(F^(c*(a+b*x))^p)], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, n, p\}, x] \&& \text{GtQ}[m, 0]$

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.22

method	result
risch	$\frac{3i \operatorname{polylog}(4, -e^{2i(bx+a)})}{4b^4} + \frac{x^2(2bx e^{2i(bx+a)} - 3ie^{2i(bx+a)} - 3i)}{b^2(1+e^{2i(bx+a)})^2} + \frac{6iax}{b^3} + \frac{3ia^2}{b^4} - \frac{3ia^4}{2b^4} + \frac{x^3 \ln(1+e^{2i(bx+a)})}{b} - \frac{2ia^3x}{b^3}$

input `int(x^3*tan(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

$$\frac{3/4*I*polylog(4, -exp(2*I*(b*x+a)))/b^4+x^2*(2*b*x*exp(2*I*(b*x+a))-3*I*exp(2*I*(b*x+a))-3*I)/b^2/(1+exp(2*I*(b*x+a)))^2+6*I/b^3*a*x+3*I/b^4*a^2-3/2*I/b^4*a^4+x^3*ln(1+exp(2*I*(b*x+a)))/b-2*I/b^3*a^3*x+3/2*x*polylog(3, -exp(2*I*(b*x+a)))/b^3-3*x*ln(1+exp(2*I*(b*x+a)))/b^3+3/2*I*polylog(2, -exp(2*I*(b*x+a)))/b^4-6/b^4*a*ln(exp(I*(b*x+a)))+2/b^4*a^3*ln(exp(I*(b*x+a)))-1/4*I*x^4-3/2*I*x^2*polylog(2, -exp(2*I*(b*x+a)))/b^2+3*I/b^2*x^2$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 344 vs. $2(163) = 326$.

Time = 0.09 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.68

$$\int x^3 \tan^3(a + bx) dx \\ = \frac{4 b^3 x^3 \tan(bx + a)^2 + 4 b^3 x^3 - 12 b^2 x^2 \tan(bx + a) + 6 b x \text{polylog}\left(3, \frac{\tan(bx+a)^2 + 2i \tan(bx+a) - 1}{\tan(bx+a)^2 + 1}\right) + 6 b x \text{polylog}\left(4, \frac{\tan(bx+a)^2 + 2i \tan(bx+a) - 1}{\tan(bx+a)^2 + 1}\right)}{b^4}$$

input `integrate(x^3*tan(b*x+a)^3,x, algorithm="fricas")`

output

$$\begin{aligned} & 1/8*(4*b^3*x^3*tan(b*x + a)^2 + 4*b^3*x^3 - 12*b^2*x^2*tan(b*x + a) + 6*b*x*polylog(3, (\tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) \\ & + 6*b*x*polylog(3, (\tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 6*(-I*b^2*x^2 + I)*dilog(2*(I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 6*(I*b^2*x^2 - I)*dilog(2*(-I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 4*(b^3*x^3 - 3*b*x)*log(-2*(I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 4*(b^3*x^3 - 3*b*x)*log(-2*(-I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) - 3*I*polylog(4, (\tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)) + 3*I*polylog(4, (\tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(\tan(b*x + a)^2 + 1)))/b^4 \end{aligned}$$

Sympy [F]

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan^3 (a + bx) dx$$

input `integrate(x**3*tan(b*x+a)**3,x)`

output `Integral(x**3*tan(a + b*x)**3, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1205 vs. $2(163) = 326$.

Time = 0.34 (sec) , antiderivative size = 1205, normalized size of antiderivative = 5.88

$$\int x^3 \tan^3(a + bx) dx = \text{Too large to display}$$

input `integrate(x^3*tan(b*x+a)^3,x, algorithm="maxima")`

output

```
1/2*(a^3*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1)) - 2*(3*(b*x + a)^4 - 12*(b*x + a)^3*a + 18*(b*x + a)^2*a^2 + 36*a^2 - 4*(4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*cos(4*b*x + 4*a) + 2*(4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*cos(2*b*x + 2*a) - (-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a + 9*(-I*a^2 + I)*(b*x + a) - 9*I*a)*sin(4*b*x + 4*a) - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a + 9*(-I*a^2 + I)*(b*x + a) - 9*I*a)*sin(2*b*x + 2*a) + 9*a)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^4 - 4*(b*x + a)^3*a + 6*(a^2 - 2)*(b*x + a)^2 + 24*(b*x + a)*a)*cos(4*b*x + 4*a) + 6*((b*x + a)^4 - 4*(b*x + a)^3*(a - I) + 6*(a^2 - 2*I*a - 1)*(b*x + a)^2 + 12*(I*a^2 + a)*(b*x + a) + 6*a^2)*cos(2*b*x + 2*a) + 6*(4*(b*x + a)^2 - 6*(b*x + a)*a + 3*a^2 + (4*(b*x + a)^2 - 6*(b*x + a)*a + 3*a^2 - 3)*cos(4*b*x + 4*a) + 2*(4*(b*x + a)^2 - 6*(b*x + a)*a + 3*I*a^2 - 3*I)*sin(4*b*x + 4*a) + 2*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2 - 3*I)*sin(2*b*x + 2*a) - 3)*dilog(-e^(2*I*b*x + 2*I*a)) + 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + 9*I*a)*cos(4*b*x + 4*a) + 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + 9*I*a)*cos(2*b*x + 2*a) - (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) ...)
```

Giac [F]

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan(bx + a)^3 dx$$

input

```
integrate(x^3*tan(b*x+a)^3,x, algorithm="giac")
```

output

```
integrate(x^3*tan(b*x + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan(a + bx)^3 dx$$

input `int(x^3*tan(a + b*x)^3,x)`

output `int(x^3*tan(a + b*x)^3, x)`

Reduce [F]

$$\int x^3 \tan^3(a + bx) dx = \int \tan(bx + a)^3 x^3 dx$$

input `int(x^3*tan(b*x+a)^3,x)`

output `int(tan(a + b*x)**3*x**3,x)`

3.12 $\int x^2 \tan^3(a + bx) dx$

Optimal result	128
Mathematica [A] (verified)	128
Rubi [A] (verified)	129
Maple [A] (verified)	133
Fricas [B] (verification not implemented)	133
Sympy [F]	134
Maxima [B] (verification not implemented)	134
Giac [F]	135
Mupad [F(-1)]	136
Reduce [F]	136

Optimal result

Integrand size = 12, antiderivative size = 128

$$\begin{aligned} \int x^2 \tan^3(a + bx) dx = & \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} \\ & - \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} + \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \\ & - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b} \end{aligned}$$

output $1/2*x^2/b - 1/3*I*x^3 + x^2*ln(1+exp(2*I*(b*x+a)))/b - ln(cos(b*x+a))/b^3 - I*x*polylog(2, -exp(2*I*(b*x+a)))/b^2 + 1/2*polylog(3, -exp(2*I*(b*x+a)))/b^3 - x*tan(b*x+a)/b^2 + 1/2*x^2*tan(b*x+a)^2/b$

Mathematica [A] (verified)

Time = 2.08 (sec), antiderivative size = 172, normalized size of antiderivative = 1.34

$$\begin{aligned} & \int x^2 \tan^3(a + bx) dx \\ &= \frac{e^{-ia}(2b^2x^2(2ibx + 3(1 + e^{2ia})\log(1 + e^{-2i(a+bx)})) + 6ib(1 + e^{2ia})x\operatorname{PolyLog}(2, -e^{-2i(a+bx)}) + 3(1 + e^{2ia})\operatorname{PolyLog}(3, -e^{-2i(a+bx)}))}{b^3} \end{aligned}$$

input `Integrate[x^2*Tan[a + b*x]^3, x]`

output
$$\begin{aligned} & (((2*b^2*x^2*((2*I)*b*x + 3*(1 + E^((2*I)*a))*Log[1 + E^((-2*I)*(a + b*x))]) + (6*I)*b*(1 + E^((2*I)*a))*x*PolyLog[2, -E^((-2*I)*(a + b*x))] + 3*(1 + E^((2*I)*a))*PolyLog[3, -E^((-2*I)*(a + b*x))])*Sec[a])/E^(I*a) + 6*b^2*x^2*Sec[a + b*x]^2 - 12*b*x*Sec[a]*Sec[a + b*x]*Sin[b*x] - 4*b^3*x^3*Tan[a] \\ &] - 12*(Log[Cos[a + b*x]] + b*x*Tan[a]))/(12*b^3) \end{aligned}$$

Rubi [A] (verified)

Time = 0.85 (sec), antiderivative size = 149, normalized size of antiderivative = 1.16, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4203, 3042, 4202, 2620, 3011, 2720, 4203, 15, 3042, 3956, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \tan^3(a + bx) dx \\ & \quad \downarrow \textcolor{blue}{3042} \\ & \int x^2 \tan(a + bx)^3 dx \\ & \quad \downarrow \textcolor{blue}{4203} \\ & - \int x^2 \tan(a + bx) dx - \frac{\int x \tan^2(a + bx) dx}{b} + \frac{x^2 \tan^2(a + bx)}{2b} \\ & \quad \downarrow \textcolor{blue}{3042} \\ & - \int x^2 \tan(a + bx) dx - \frac{\int x \tan(a + bx)^2 dx}{b} + \frac{x^2 \tan^2(a + bx)}{2b} \\ & \quad \downarrow \textcolor{blue}{4202} \\ & 2i \int \frac{e^{2i(a+bx)} x^2}{1 + e^{2i(a+bx)}} dx - \frac{\int x \tan(a + bx)^2 dx}{b} + \frac{x^2 \tan^2(a + bx)}{2b} - \frac{ix^3}{3} \\ & \quad \downarrow \textcolor{blue}{2620} \end{aligned}$$

$$2i \left(\frac{i \int x \log(1 + e^{2i(a+bx)}) dx}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \frac{\int x \tan(a+bx)^2 dx}{b} +$$

$$\frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}$$

\downarrow 3011

$$2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{2b} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) -$$

$$\frac{\int x \tan(a+bx)^2 dx}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}$$

\downarrow 2720

$$2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) -$$

$$\frac{\int x \tan(a+bx)^2 dx}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}$$

\downarrow 4203

$$2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) -$$

$$-\frac{\int \tan(a+bx) dx}{b} - \frac{\int x dx}{b} + \frac{x \tan(a+bx)}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}$$

\downarrow 15

$$2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) -$$

$$-\frac{\int \tan(a+bx) dx}{b} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}$$

\downarrow 3042

$$\begin{aligned}
& 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \frac{-\frac{\int \tan(a+bx) dx}{b} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{3956}
\end{aligned}$$

$$\begin{aligned}
& 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\int e^{-2i(a+bx)} \operatorname{PolyLog}(2, -e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \frac{\frac{\log(\cos(a+bx))}{b^2} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3} \\
& \quad \downarrow \text{7143}
\end{aligned}$$

$$\begin{aligned}
& 2i \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{4b^2} \right)}{b} - \frac{ix^2 \log(1 + e^{2i(a+bx)})}{2b} \right) - \\
& \frac{\frac{\log(\cos(a+bx))}{b^2} + \frac{x \tan(a+bx)}{b} - \frac{x^2}{2}}{b} + \frac{x^2 \tan^2(a+bx)}{2b} - \frac{ix^3}{3}
\end{aligned}$$

input `Int[x^2*Tan[a + b*x]^3, x]`

output

$$\begin{aligned}
& (-1/3*I)*x^3 + (2*I)*(((I/2)*x^2*Log[1 + E^((2*I)*(a + b*x))])/b + (I*(I/2)*x*PolyLog[2, -E^((2*I)*(a + b*x))])/b - PolyLog[3, -E^((2*I)*(a + b*x))]/(4*b^2))/b) + (x^2*Tan[a + b*x]^2)/(2*b) - (-1/2*x^2 + Log[Cos[a + b*x]]/b^2 + (x*Tan[a + b*x])/b)/b
\end{aligned}$$

Definitions of rubi rules used

rule 15 $\text{Int}[(a_*)*(x_)^{(m_)}, \text{x_Symbol}] \rightarrow \text{Simp}[a*(x^{(m+1)/(m+1)}), \text{x}] /; \text{FreeQ}[\{a, m\}, \text{x}] \&& \text{NeQ}[m, -1]$

rule 2620 $\text{Int}[((F_*)^{(g_*)}*((e_*) + (f_*)*(x_)))^{(n_*)}*((c_*) + (d_*)*(x_))^{(m_)}/((a_) + (b_*)*((F_*)^{(g_*)}*((e_*) + (f_*)*(x_)))^{(n_*)}), \text{x_Symbol}] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], \text{x}] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], \text{x}], \text{x}] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, \text{x}] \&& \text{IGtQ}[m, 0]$

rule 2720 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, \text{x}]\}, \text{Simp}[v/D[v, \text{x}] \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, \text{x}]/\text{x}, \text{x}], \text{x}, v], \text{x}]] /; \text{FunctionOfExponentialQ}[u, \text{x}] \&& \text{!MatchQ}[u, (w_)*((a_*)*(v_)^{(n_)})^{(m_)} /; \text{FreeQ}[\{a, m, n\}, \text{x}] \&& \text{IntegerQ}[m*n] \&& \text{!MatchQ}[u, E^{(c_*)*((a_*) + (b_*)*x)}*(F_)[v_] /; \text{FreeQ}[\{a, b, c\}, \text{x}] \&& \text{InverseFunctionQ}[F[x]]]$

rule 3011 $\text{Int}[\text{Log}[1 + (e_*)*((F_*)^{(c_*)}*((a_*) + (b_*)*(x_)))^{(n_*)}]*((f_*) + (g_*)*(x_))^{(m_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(-(f + g*x)^m)*(\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*\text{Log}[F])), \text{x}] + \text{Simp}[g*(m/(b*c*n*\text{Log}[F])) \text{Int}[(f + g*x)^{(m-1)}*\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n], \text{x}], \text{x}] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, \text{x}] \&& \text{GtQ}[m, 0]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$

rule 3956 $\text{Int}[\tan[(c_*) + (d_*)*(x_)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], \text{x}]]/\text{d}, \text{x}] /; \text{FreeQ}[\{c, d\}, \text{x}]$

rule 4202 $\text{Int}[((c_*) + (d_*)*(x_))^{(m_*)}*\tan[(e_*) + (f_*)*(x_)], \text{x_Symbol}] \rightarrow \text{Simp}[I*((c + d*x)^{(m+1)}/(d*(m+1))), \text{x}] - \text{Simp}[2*I \text{Int}[(c + d*x)^m*(E^{(2*I*(e + f*x))}/(1 + E^{(2*I*(e + f*x))))], \text{x}], \text{x}] /; \text{FreeQ}[\{c, d, e, f\}, \text{x}] \&& \text{IGtQ}[m, 0]$

rule 4203 $\text{Int}[(c_{_}) + (d_{_})*(x_{_})^{(m_{_})}*((b_{_})*\tan[(e_{_}) + (f_{_})*(x_{_})])^{(n_{_})}, x_{_}] \rightarrow \text{Simp}[b*(c + d*x)^m*((b*Tan[e + f*x])^{(n - 1)}/(f*(n - 1))), x] + (-\text{Simp}[b*d*(m/(f*(n - 1))) \text{ Int}[(c + d*x)^{(m - 1)}*(b*Tan[e + f*x])^{(n - 1)}, x], x] - \text{Simp}[b^2 \text{ Int}[(c + d*x)^m*(b*Tan[e + f*x])^{(n - 2)}, x], x]); \text{FreeQ}[\{b, c, d, e, f\}, x] \&& \text{GtQ}[n, 1] \&& \text{GtQ}[m, 0]$

rule 7143 $\text{Int}[\text{PolyLog}[n_{_}, (c_{_})*((a_{_}) + (b_{_})*(x_{_}))^{(p_{_})}]/((d_{_}) + (e_{_})*(x_{_})), x_{_}] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c*(a + b*x)^p]/(e*p), x]; \text{FreeQ}[\{a, b, c, d, e, n, p\}, x] \&& \text{EqQ}[b*d, a*e]$

Maple [A] (verified)

Time = 0.23 (sec), antiderivative size = 180, normalized size of antiderivative = 1.41

method	result
risch	$-\frac{ix^3}{3} + \frac{2x(bx e^{2i(bx+a)} - ie^{2i(bx+a)} - i)}{b^2(1+e^{2i(bx+a)})^2} - \frac{2a^2 \ln(e^{i(bx+a)})}{b^3} + \frac{2ia^2x}{b^2} + \frac{4ia^3}{3b^3} + \frac{x^2 \ln(1+e^{2i(bx+a)})}{b} - \frac{ix \text{polylog}(2, -e^{2i(bx+a)})}{b^2}$

input `int(x^2*tan(b*x+a)^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/3*I*x^3 + 2*x*(b*x*exp(2*I*(b*x+a)) - I*exp(2*I*(b*x+a)) - I)/b^2/(1 + exp(2*I*(b*x+a)))^2 - 2/b^3*a^2*ln(exp(I*(b*x+a))) + 2*I/b^2*a^2*x^4/3*I/b^3*a^3*x^2 + \\ & n(1 + exp(2*I*(b*x+a)))/b - I*x*polylog(2, -exp(2*I*(b*x+a)))/b^2 + 1/2*polylog(3, -exp(2*I*(b*x+a)))/b^3 + 2/b^3*ln(exp(I*(b*x+a))) - 1/b^3*ln(1 + exp(2*I*(b*x+a))) \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(109) = 218$.

Time = 0.09 (sec), antiderivative size = 240, normalized size of antiderivative = 1.88

$$\begin{aligned} & \int x^2 \tan^3(a + bx) dx \\ & = \frac{2 b^2 x^2 \tan(bx + a)^2 + 2 b^2 x^2 + 2i b x \text{Li}_2\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1} + 1\right) - 2i b x \text{Li}_2\left(\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1} + 1\right) - 4 b x \tan(bx + a)^2}{b^3} \end{aligned}$$

input `integrate(x^2*tan(b*x+a)^3,x, algorithm="fricas")`

output
$$\frac{1}{4} \left(2b^2x^2\tan(bx + a)^2 + 2b^2x^2 + 2Ib\tan(bx + a) - 1 \right) / (\tan(bx + a)^2 + 1) + 1 - 2Ib\tan(bx + a) \operatorname{dilog}(2(-I\tan(bx + a) - 1)) / (\tan(bx + a)^2 + 1) - 4b\tan(bx + a) + 2(b^2x^2 - 1) \log(-2(I\tan(bx + a) - 1)) / (\tan(bx + a)^2 + 1) + 2(b^2x^2 - 1) \log(-2(-I\tan(bx + a) - 1)) / (\tan(bx + a)^2 + 1) + \operatorname{polylog}(3, (\tan(bx + a)^2 + 2I\tan(bx + a) - 1)) / (\tan(bx + a)^2 + 1) + \operatorname{polylog}(3, (\tan(bx + a)^2 - 2I\tan(bx + a) - 1)) / (\tan(bx + a)^2 + 1)) / b^3$$

Sympy [F]

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan^3(a + bx) dx$$

input `integrate(x**2*tan(b*x+a)**3,x)`

output `Integral(x**2*tan(a + b*x)**3, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 736 vs. $2(109) = 218$.

Time = 0.28 (sec) , antiderivative size = 736, normalized size of antiderivative = 5.75

$$\int x^2 \tan^3(a + bx) dx = \text{Too large to display}$$

input `integrate(x^2*tan(b*x+a)^3,x, algorithm="maxima")`

output

```

-1/2*(a^2*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1)) + 2*(2*(b*x +
a)^3 - 6*(b*x + a)^2*a - 6*((b*x + a)^2 - 2*(b*x + a)*a + ((b*x + a)^2 -
2*(b*x + a)*a - 1)*cos(4*b*x + 4*a) + 2*((b*x + a)^2 - 2*(b*x + a)*a - 1)*
cos(2*b*x + 2*a) - (-I*(b*x + a)^2 + 2*I*(b*x + a)*a + I)*sin(4*b*x + 4*a)
- 2*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a + I)*sin(2*b*x + 2*a) - 1)*arctan2(
sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 2*((b*x + a)^3 - 3*(b*x + a)^2*a
- 6*b*x - 6*a)*cos(4*b*x + 4*a) + 4*((b*x + a)^3 - 3*(b*x + a)^2*(a - I)
+ 3*(b*x + a)*(-2*I*a - 1) - 3*a)*cos(2*b*x + 2*a) + 6*(b*x*cos(4*b*x + 4*
a) + 2*b*x*cos(2*b*x + 2*a) + I*b*x*sin(4*b*x + 4*a) + 2*I*b*x*sin(2*b*x +
2*a) + b*x)*dilog(-e^(2*I*b*x + 2*I*a)) + 3*(I*(b*x + a)^2 - 2*I*(b*x + a
)*a + (I*(b*x + a)^2 - 2*I*(b*x + a)*a - I)*cos(4*b*x + 4*a) + 2*(I*(b*x +
a)^2 - 2*I*(b*x + a)*a - I)*cos(2*b*x + 2*a) - ((b*x + a)^2 - 2*(b*x + a)
*a - 1)*sin(4*b*x + 4*a) - 2*((b*x + a)^2 - 2*(b*x + a)*a - 1)*sin(2*b*x +
2*a) - I)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a
) + 1) + 3*(I*cos(4*b*x + 4*a) + 2*I*cos(2*b*x + 2*a) - sin(4*b*x + 4*a) -
2*sin(2*b*x + 2*a) + I)*polylog(3, -e^(2*I*b*x + 2*I*a)) + 2*(I*(b*x + a)
^3 - 3*I*(b*x + a)^2*a - 6*I*b*x - 6*I*a)*sin(4*b*x + 4*a) + 4*(I*(b*x + a
)^3 + 3*(b*x + a)^2*(-I*a - 1) + 3*(b*x + a)*(2*a - I) - 3*I*a)*sin(2*b*x
+ 2*a) - 12*a)/(-6*I*cos(4*b*x + 4*a) - 12*I*cos(2*b*x + 2*a) + 6*sin(4*b*
x + 4*a) + 12*sin(2*b*x + 2*a) - 6*I))/b^3

```

Giac [F]

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan(bx + a)^3 dx$$

input

```
integrate(x^2*tan(b*x+a)^3,x, algorithm="giac")
```

output

```
integrate(x^2*tan(b*x + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan(a + bx)^3 dx$$

input `int(x^2*tan(a + b*x)^3,x)`

output `int(x^2*tan(a + b*x)^3, x)`

Reduce [F]

$$\int x^2 \tan^3(a + bx) dx = \int \tan(bx + a)^3 x^2 dx$$

input `int(x^2*tan(b*x+a)^3,x)`

output `int(tan(a + b*x)**3*x**2,x)`

3.13 $\int x \tan^3(a + bx) dx$

Optimal result	137
Mathematica [A] (warning: unable to verify)	137
Rubi [A] (verified)	138
Maple [A] (verified)	140
Fricas [B] (verification not implemented)	141
Sympy [F]	141
Maxima [B] (verification not implemented)	142
Giac [F]	142
Mupad [F(-1)]	143
Reduce [F]	143

Optimal result

Integrand size = 10, antiderivative size = 90

$$\begin{aligned} \int x \tan^3(a + bx) dx = & \frac{x}{2b} - \frac{ix^2}{2} + \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\ & - \frac{\tan(a + bx)}{2b^2} + \frac{x \tan^2(a + bx)}{2b} \end{aligned}$$

output $1/2*x/b - 1/2*I*x^2 + x*ln(1 + exp(2*I*(b*x+a)))/b - 1/2*I*polylog(2, -exp(2*I*(b*x+a)))/b^2 - 1/2*tan(b*x+a)/b^2 + 1/2*x*tan(b*x+a)^2/b$

Mathematica [A] (warning: unable to verify)

Time = 4.33 (sec), antiderivative size = 171, normalized size of antiderivative = 1.90

$$\begin{aligned} \int x \tan^3(a + bx) dx \\ = \frac{ibx(\pi + 2 \arctan(\cot(a))) + \pi \log(1 + e^{-2ibx}) + 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))}) - \pi^2 \arctan(\cot(a))}{b^2} \end{aligned}$$

input `Integrate[x*Tan[a + b*x]^3, x]`

output

$$(I*b*x*(Pi + 2*ArcTan[Cot[a]]) + Pi*Log[1 + E^((-2*I)*b*x)] + 2*(b*x - ArcTan[Cot[a]]])*Log[1 - E^((2*I)*(b*x - ArcTan[Cot[a]]))] - Pi*Log[Cos[b*x]] + 2*ArcTan[Cot[a]]*Log[Sin[b*x - ArcTan[Cot[a]]]] - I*PolyLog[2, E^((2*I)*(b*x - ArcTan[Cot[a]]))] + b*x*Sec[a + b*x]^2 - Sec[a]*Sec[a + b*x]*Sin[b*x] - b^2*x^2*Tan[a] + (b^2*x^2*Sqrt[Csc[a]^2]*Tan[a])/E^(I*ArcTan[Cot[a]]))/(2*b^2)$$

Rubi [A] (verified)

Time = 0.48 (sec), antiderivative size = 98, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.900, Rules used = {3042, 4203, 3042, 3954, 24, 4202, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \tan^3(a + bx) dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int x \tan(a + bx)^3 dx \\
 & \downarrow \textcolor{blue}{4203} \\
 & -\frac{\int \tan^2(a + bx) dx}{2b} - \int x \tan(a + bx) dx + \frac{x \tan^2(a + bx)}{2b} \\
 & \downarrow \textcolor{blue}{3042} \\
 & -\int x \tan(a + bx) dx - \frac{\int \tan(a + bx)^2 dx}{2b} + \frac{x \tan^2(a + bx)}{2b} \\
 & \downarrow \textcolor{blue}{3954} \\
 & -\frac{\frac{\tan(a+bx)}{b} - \int 1 dx}{2b} - \int x \tan(a + bx) dx + \frac{x \tan^2(a + bx)}{2b} \\
 & \downarrow \textcolor{blue}{24} \\
 & -\int x \tan(a + bx) dx + \frac{x \tan^2(a + bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} \\
 & \downarrow \textcolor{blue}{4202}
 \end{aligned}$$

$$\begin{aligned}
& 2i \int \frac{e^{2i(a+bx)}x}{1+e^{2i(a+bx)}} dx + \frac{x \tan^2(a+bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2} \\
& \quad \downarrow \textcolor{blue}{2620} \\
& 2i \left(\frac{i \int \log(1+e^{2i(a+bx)}) dx}{2b} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) + \frac{x \tan^2(a+bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2} \\
& \quad \downarrow \textcolor{blue}{2715} \\
& 2i \left(\frac{\int e^{-2i(a+bx)} \log(1+e^{2i(a+bx)}) de^{2i(a+bx)}}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) + \frac{x \tan^2(a+bx)}{2b} - \\
& \quad \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2} \\
& \quad \downarrow \textcolor{blue}{2838} \\
& 2i \left(-\frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{4b^2} - \frac{ix \log(1+e^{2i(a+bx)})}{2b} \right) + \frac{x \tan^2(a+bx)}{2b} - \frac{\frac{\tan(a+bx)}{b} - x}{2b} - \frac{ix^2}{2}
\end{aligned}$$

input `Int[x*Tan[a + b*x]^3, x]`

output `(-1/2*I)*x^2 + (2*I)*(((-1/2*I)*x*Log[1 + E^((2*I)*(a + b*x))])/b - PolyLog[2, -E^((2*I)*(a + b*x))]/(4*b^2)) + (x*Tan[a + b*x]^2)/(2*b) - (-x + Tan[a + b*x])/b)/(2*b)`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simplify[a*x, x] /; FreeQ[a, x]`

rule 2620 `Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simplify[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 $\text{Int}[\text{Log}[(a_.) + (b_.)*((F_.)^((e_.)*((c_.) + (d_.)*(x_))))^{(n_.)}], x_{\text{Symbol}}] \\ :> \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^e(c + d*x))^{n}], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \text{GtQ}[a, 0]$

rule 2838 $\text{Int}[\text{Log}[(c_.)*((d_.) + (e_.)^{(n_.)})^{(n_.)}]/(x_), x_{\text{Symbol}}] :> \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \&& \text{EqQ}[c*d, 1]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] :> \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3954 $\text{Int}[((b_.)*\text{tan}[(c_.) + (d_.)*(x_)])^{(n_.)}, x_{\text{Symbol}}] :> \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)}/(d*(n - 1))), x] - \text{Simp}[b^2 \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&& \text{GtQ}[n, 1]$

rule 4202 $\text{Int}[((c_.) + (d_.)*(x_))^{(m_.)}*\text{tan}[(e_.) + (f_.)*(x_)], x_{\text{Symbol}}] :> \text{Simp}[I*((c + d*x)^{(m + 1)}/(d*(m + 1))), x] - \text{Simp}[2*I \text{Int}[(c + d*x)^m * (E^{(2*I)*(e + f*x)})/(1 + E^{(2*I)*(e + f*x)}), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{IGtQ}[m, 0]$

rule 4203 $\text{Int}[((c_.) + (d_.)*(x_))^{(m_.)}*((b_.)*\text{tan}[(e_.) + (f_.)*(x_)])^{(n_.)}, x_{\text{Symbol}}] :> \text{Simp}[b*(c + d*x)^m * ((b*\text{Tan}[e + f*x])^{(n - 1)}/(f*(n - 1))), x] + (-\text{Simp}[b*d*(m/(f*(n - 1))) \text{Int}[(c + d*x)^{(m - 1)} * (b*\text{Tan}[e + f*x])^{(n - 1)}, x], x] - \text{Simp}[b^2 \text{Int}[(c + d*x)^{m - 1} * (b*\text{Tan}[e + f*x])^{(n - 2)}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&& \text{GtQ}[n, 1] \&& \text{GtQ}[m, 0]$

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.36

method	result
risch	$-\frac{ix^2}{2} + \frac{2bx e^{2i(bx+a)} - ie^{2i(bx+a)} - i}{b^2 (1 + e^{2i(bx+a)})^2} - \frac{2iax}{b} - \frac{ia^2}{b^2} + \frac{x \ln(1 + e^{2i(bx+a)})}{b} - \frac{i \operatorname{polylog}(2, -e^{2i(bx+a)})}{2b^2} + \frac{2a \ln(e^{i(bx+a)})}{b^2}$

input `int(x*tan(b*x+a)^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -\frac{1}{2} I x^2 + \frac{(2 b x \exp(2 I (b x + a)) - I \exp(2 I (b x + a)) - I) / b^2}{1 + \exp(2 I (b x + a))} \\ & - 2 - 2 I / b + a x - I / b^2 + a^2 + x \ln(1 + \exp(2 I (b x + a))) / b - \frac{1}{2} I \operatorname{polylog}(2, -\exp(2 I (b x + a))) / b^2 + 2 / b^2 + a \ln(\exp(I (b x + a))) \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(71) = 142$.

Time = 0.08 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.62

$$\int x \tan^3(a + bx) dx = \frac{2 b x \tan(bx + a)^2 + 2 b x \log\left(-\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x \log\left(-\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x + i \operatorname{Li}_2\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x \log\left(\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x + i \operatorname{Li}_2\left(\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x \log\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x + i \operatorname{Li}_2\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right)}{4 b^2}$$

input `integrate(x*tan(b*x+a)^3,x, algorithm="fricas")`

output
$$\begin{aligned} & \frac{1}{4} (2 b x \tan(bx + a)^2 + 2 b x \log(-2 * (I * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1)) + 2 b x \log(-2 * (-I * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1)) + 2 b x + I * \operatorname{dilog}(2 * (I * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1) + 1) - I * \operatorname{dilog}(2 * (-I * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1) + 1) - 2 * \tan(bx + a)) / b^2 \end{aligned}$$

Sympy [F]

$$\int x \tan^3(a + bx) dx = \int x \tan^3(a + bx) dx$$

input `integrate(x*tan(b*x+a)**3,x)`

output `Integral(x*tan(a + b*x)**3, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(71) = 142$.

Time = 0.25 (sec) , antiderivative size = 386, normalized size of antiderivative = 4.29

$$\int x \tan^3(a + bx) dx = \frac{-b^2 x^2 \cos(4bx + 4a) + i b^2 x^2 \sin(4bx + 4a) + b^2 x^2 - 2(bx \cos(4bx + 4a) + 2bx \cos(2bx + 2a) + i b x \sin(2bx + 2a))}{b^2 x^2}$$

input `integrate(x*tan(b*x+a)^3,x, algorithm="maxima")`

output

$$-(b^2 x^2 \cos(4bx + 4a) + I b^2 x^2 \sin(4bx + 4a) + b^2 x^2 - 2(bx \cos(4bx + 4a) + 2bx \cos(2bx + 2a) + i b x \sin(2bx + 2a)) + 2b x \sin(2bx + 2a) + I b x \sin(4bx + 4a) + 2I b x \sin(2bx + 2a) + b x \arctan2(\sin(2bx + 2a), \cos(2bx + 2a)) + 1) + 2(b^2 x^2 + 2I b x + 1) \cos(2bx + 2a) + (\cos(4bx + 4a) + 2\cos(2bx + 2a) + I \sin(4bx + 4a) + 2I \sin(2bx + 2a) + 1) \operatorname{dilog}(-e^{(2I b x + 2I a)}) - (-I b x \cos(4bx + 4a) - 2I b x \cos(2bx + 2a) + b x \sin(4bx + 4a) + 2b x \sin(2bx + 2a) - I b x \log(\cos(2bx + 2a)) + 2 + \sin(2bx + 2a)^2 + 2\cos(2bx + 2a) + 1) + 2(I b^2 x^2 - 2b x + I) \sin(2bx + 2a) + 2) / (-2I b^2 \cos(4bx + 4a) - 4I b^2 \cos(2bx + 2a) + 2b^2 \sin(4bx + 4a) + 4b^2 \sin(2bx + 2a) - 2I b^2)$$

Giac [F]

$$\int x \tan^3(a + bx) dx = \int x \tan(bx + a)^3 dx$$

input `integrate(x*tan(b*x+a)^3,x, algorithm="giac")`

output `integrate(x*tan(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x \tan^3(a + bx) dx = \int x \tan(a + bx)^3 dx$$

input `int(x*tan(a + b*x)^3,x)`

output `int(x*tan(a + b*x)^3, x)`

Reduce [F]

$$\int x \tan^3(a + bx) dx = \int \tan(bx + a)^3 x dx$$

input `int(x*tan(b*x+a)^3,x)`

output `int(tan(a + b*x)**3*x,x)`

3.14 $\int \frac{\tan^3(a+bx)}{x} dx$

Optimal result	144
Mathematica [N/A]	144
Rubi [N/A]	145
Maple [N/A]	146
Fricas [N/A]	146
Sympy [N/A]	147
Maxima [N/A]	147
Giac [N/A]	148
Mupad [N/A]	148
Reduce [N/A]	149

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^3(a + bx)}{x} dx = \text{Int}\left(\frac{\tan^3(a + bx)}{x}, x\right)$$

output `Defer(Int)(tan(b*x+a)^3/x,x)`

Mathematica [N/A]

Not integrable

Time = 4.40 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(a + bx)}{x} dx$$

input `Integrate[Tan[a + b*x]^3/x,x]`

output `Integrate[Tan[a + b*x]^3/x, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\tan^3(a + bx)}{x} dx \\ \downarrow \textcolor{blue}{3042} \\ \int \frac{\tan(a + bx)^3}{x} dx \\ \downarrow \textcolor{blue}{4222} \\ \int \frac{\tan^3(a + bx)}{x} dx \end{array}$$

input `Int[Tan[a + b*x]^3/x, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4222 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)}, x_\text{Symbol}] \rightarrow \text{Simp}[\text{If}[\text{MatchQ}[f, (f1_.)*(Complex[0, j_])], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Coth}[-I]*(e - \text{Pi}/2) - I*f*x]^n, x], I^{n* \text{Unintegatable}[(c + d*x)^m*\text{Tanh}[-I]*e - I*f*x]^n, x]], \text{If}[\text{MatchQ}[e, (e1_.) + \text{Pi}/2], (-1)^n* \text{Unintegatable}[(c + d*x)^m*\text{Cot}[e - \text{Pi}/2 + f*x]^n, x], \text{Unintegatable}[(c + d*x)^m*\text{Tan}[e + f*x]^n, x]]], x] /; \text{FreeQ}[\{c, d, e, f, m, n\}, x] \&& \text{IntegerQ}[n]$

Maple [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx+a)^3}{x} dx$$

input $\text{int}(\tan(b*x+a)^3/x, x)$

output $\text{int}(\tan(b*x+a)^3/x, x)$

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

input $\text{integrate}(\tan(b*x+a)^3/x, x, \text{algorithm}=\text{"fricas"})$

output `integral(tan(b*x + a)^3/x, x)`

Sympy [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(a + bx)}{x} dx$$

input `integrate(tan(b*x+a)**3/x,x)`

output `Integral(tan(a + b*x)**3/x, x)`

Maxima [N/A]

Not integrable

Time = 0.52 (sec) , antiderivative size = 534, normalized size of antiderivative = 44.50

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(bx + a)}{x} dx$$

input `integrate(tan(b*x+a)^3/x,x, algorithm="maxima")`

output

```
(4*b*x*cos(2*b*x + 2*a)^2 + 4*b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + (2*b*x*cos(2*b*x + 2*a) - sin(2*b*x + 2*a))*cos(4*b*x + 4*a) - (b^2*x^2*cos(4*b*x + 4*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(4*b*x + 4*a)^2 + 4*b^2*x^2*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^2*sin(2*b*x + 2*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2 + 2*(2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2)*cos(4*b*x + 4*a))*integrate(2*(b^2*x^2 - 1)*sin(2*b*x + 2*a)/(b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(2*b*x + 2*a)^2 + 2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3), x) + (2*b*x*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1)*sin(4*b*x + 4*a) + sin(2*b*x + 2*a))/(b^2*x^2*cos(4*b*x + 4*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(4*b*x + 4*a)^2 + 4*b^2*x^2*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^2*sin(2*b*x + 2*a)^2 + 4*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2 + 2*(2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2)*cos(4*b*x + 4*a))
```

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

input `integrate(tan(b*x+a)^3/x,x, algorithm="giac")`

output `integrate(tan(b*x + a)^3/x, x)`

Mupad [N/A]

Not integrable

Time = 8.50 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(a + bx)^3}{x} dx$$

input `int(tan(a + b*x)^3/x,x)`

output `int(tan(a + b*x)^3/x, x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

input `int(tan(b*x+a)^3/x,x)`

output `int(tan(a + b*x)**3/x,x)`

3.15 $\int \frac{\tan^3(a+bx)}{x^2} dx$

Optimal result	150
Mathematica [N/A]	150
Rubi [N/A]	151
Maple [N/A]	152
Fricas [N/A]	152
Sympy [N/A]	153
Maxima [N/A]	153
Giac [N/A]	154
Mupad [N/A]	154
Reduce [N/A]	155

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^3(a+bx)}{x^2} dx = \text{Int}\left(\frac{\tan^3(a+bx)}{x^2}, x\right)$$

output `Defer(Int)(tan(b*x+a)^3/x^2,x)`

Mathematica [N/A]

Not integrable

Time = 2.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a+bx)}{x^2} dx = \int \frac{\tan^3(a+bx)}{x^2} dx$$

input `Integrate[Tan[a + b*x]^3/x^2,x]`

output `Integrate[Tan[a + b*x]^3/x^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{\tan^3(a + bx)}{x^2} dx \\ \downarrow \text{3042} \\ \int \frac{\tan(a + bx)^3}{x^2} dx \\ \downarrow \text{4222} \\ \int \frac{\tan^3(a + bx)}{x^2} dx \end{array}$$

input `Int[Tan[a + b*x]^3/x^2, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4222 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.], x_Symbol] :> Simp[If[MatchQ[f, (f1_.)*(Complex[0, j_])], If[MatchQ[e, (e1_.) + Pi/2], I^n*Unintegrable[(c + d*x)^m*Coth[(-I)*(e - Pi/2) - I*f*x]^n, x], I^n*Unintegrable[(c + d*x)^m*Tanh[(-I)*e - I*f*x]^n, x]], If[MatchQ[e, (e1_.) + Pi/2], (-1)^n*Unintegrable[(c + d*x)^m*Cot[e - Pi/2 + f*x]^n, x], Unintegrable[(c + d*x)^m*Tan[e + f*x]^n, x]]], x] /; FreeQ[{c, d, e, f, m, n}, x] && IntegerQ[n]`

Maple [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx+a)^3}{x^2} dx$$

input `int(tan(b*x+a)^3/x^2, x)`

output `int(tan(b*x+a)^3/x^2, x)`

Fricas [N/A]

Not integrable

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

input `integrate(tan(b*x+a)^3/x^2, x, algorithm="fricas")`

output `integral(tan(b*x + a)^3/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(a + bx)}{x^2} dx$$

input `integrate(tan(b*x+a)**3/x**2,x)`

output `Integral(tan(a + b*x)**3/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.64 (sec) , antiderivative size = 536, normalized size of antiderivative = 44.67

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

input `integrate(tan(b*x+a)^3/x^2,x, algorithm="maxima")`

output

$$(4*b*x*cos(2*b*x + 2*a)^2 + 4*b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + 2*(b*x*cos(2*b*x + 2*a) - sin(2*b*x + 2*a))*cos(4*b*x + 4*a) - (b^2*x^3*cos(4*b*x + 4*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(4*b*x + 4*a)^2 + 4*b^2*x^3*sin(2*b*x + 2*a)^2 + 4*b^2*x^3*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^3*sin(2*b*x + 2*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3 + 2*(2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3)*cos(4*b*x + 4*a))*integrate(2*(b^2*x^2 - 3)*sin(2*b*x + 2*a)/(b^2*x^4*cos(2*b*x + 2*a)^2 + b^2*x^4*sin(2*b*x + 2*a)^2 + 2*b^2*x^4*cos(2*b*x + 2*a) + b^2*x^4), x) + 2*(b*x*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1)*sin(4*b*x + 4*a) + 2*sin(2*b*x + 2*a))/(b^2*x^3*cos(4*b*x + 4*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(4*b*x + 4*a)^2 + 4*b^2*x^3*sin(4*b*x + 4*a)*sin(2*b*x + 2*a) + 4*b^2*x^3*sin(2*b*x + 2*a)^2 + 4*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3 + 2*(2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3)*cos(4*b*x + 4*a))$$
Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

input `integrate(tan(b*x+a)^3/x^2,x, algorithm="giac")`

output `integrate(tan(b*x + a)^3/x^2, x)`

Mupad [N/A]

Not integrable

Time = 8.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)^3}{x^2} dx$$

input `int(tan(a + b*x)^3/x^2,x)`

output `int(tan(a + b*x)^3/x^2, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

input `int(tan(b*x+a)^3/x^2,x)`

output `int(tan(a + b*x)**3/x**2,x)`

$$\mathbf{3.16} \quad \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$$

Optimal result	156
Mathematica [A] (verified)	156
Rubi [A] (verified)	157
Maple [F]	157
Fricas [A] (verification not implemented)	158
Sympy [F]	158
Maxima [F]	159
Giac [F(-2)]	159
Mupad [B] (verification not implemented)	160
Reduce [B] (verification not implemented)	160

Optimal result

Integrand size = 45, antiderivative size = 18

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

output $-2*x^2/b/\tan(b*x+a)^{(1/2)}$

Mathematica [A] (verified)

Time = 1.56 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

input `Integrate[x^2/Tan[a + b*x]^(3/2) - (4*x)/(b*.Sqrt[Tan[a + b*x]]) + x^2*Sqrt[Tan[a + b*x]], x]`

output $(-2*x^2)/(b*Sqrt[Tan[a + b*x]])$

Rubi [A] (verified)

Time = 3.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.022$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} + x^2 \sqrt{\tan(a + bx)} - \frac{4x}{b \sqrt{\tan(a + bx)}} \right) dx \\ & \quad \downarrow \text{2009} \\ & \quad - \frac{2x^2}{b \sqrt{\tan(a + bx)}} \end{aligned}$$

input `Int[x^2/Tan[a + b*x]^(3/2) - (4*x)/(b*.Sqrt[Tan[a + b*x]]) + x^2*Sqrt[Tan[a + b*x]], x]`

output `(-2*x^2)/(b*.Sqrt[Tan[a + b*x]])`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \left(\frac{x^2}{\tan(bx + a)^{\frac{3}{2}}} - \frac{4x}{b \sqrt{\tan(bx + a)}} + x^2 \sqrt{\tan(bx + a)} \right) dx$$

input `int(x^2/tan(b*x+a)^(3/2)-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2),x)`

output `int(x^2/tan(b*x+a)^(3/2)-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2),x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(bx + a)}}$$

input `integrate(x^2/tan(b*x+a)^(3/2)-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2), x, algorithm="fricas")`

output `-2*x^2/(b*sqrt(tan(b*x + a)))`

Sympy [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx \\ &= \frac{\int \left(-\frac{4x}{\sqrt{\tan(a+bx)}} \right) dx + \int \frac{bx^2}{\tan^{\frac{3}{2}}(a+bx)} dx + \int bx^2\sqrt{\tan(a+bx)} dx}{b} \end{aligned}$$

input `integrate(x**2/tan(b*x+a)**(3/2)-4*x/b/tan(b*x+a)**(1/2)+x**2*tan(b*x+a)**(1/2), x)`

output `(Integral(-4*x/sqrt(tan(a + b*x)), x) + Integral(b*x**2/tan(a + b*x)**(3/2), x) + Integral(b*x**2*sqrt(tan(a + b*x)), x))/b`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx \\ &= \int x^2\sqrt{\tan(bx + a)} + \frac{x^2}{\tan(bx + a)^{\frac{3}{2}}} - \frac{4x}{b\sqrt{\tan(bx + a)}} dx \end{aligned}$$

input `integrate(x^2/tan(b*x+a)^(3/2)-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2),x, algorithm="maxima")`

output `integrate(x^2*sqrt(tan(b*x + a)) + x^2/tan(b*x + a)^(3/2) - 4*x/(b*sqrt(tan(b*x + a))), x)`

Giac [F(-2)]

Exception generated.

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx \\ &= \text{Exception raised: RuntimeError} \end{aligned}$$

input `integrate(x^2/tan(b*x+a)^(3/2)-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve
cteur & l) Error: Bad Argument Value`

Mupad [B] (verification not implemented)

Time = 9.19 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.78

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx \\ = -\frac{x^2 \sin(2a + 2bx) \sqrt{\frac{\sin(2a+2bx)}{\cos(2a+2bx)+1}}}{b \sin(a + bx)^2}$$

input `int(x^2*tan(a + b*x)^(1/2) + x^2/tan(a + b*x)^(3/2) - (4*x)/(b*tan(a + b*x))^(1/2)), x)`

output `-(x^2*sin(2*a + 2*b*x)*(sin(2*a + 2*b*x)/(cos(2*a + 2*b*x) + 1))^(1/2))/(b *sin(a + b*x)^2)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.28

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx = -\frac{2\sqrt{\tan(bx + a)}x^2}{\tan(bx + a)b}$$

input `int(x^2/tan(b*x+a)^(3/2)-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2), x)`

output `(- 2*sqrt(tan(a + b*x))*x**2)/(tan(a + b*x)*b)`

3.17
$$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$$

Optimal result	161
Mathematica [A] (verified)	161
Rubi [F]	162
Maple [F]	162
Fricas [A] (verification not implemented)	163
Sympy [F]	163
Maxima [F]	163
Giac [F(-2)]	164
Mupad [B] (verification not implemented)	164
Reduce [B] (verification not implemented)	165

Optimal result

Integrand size = 49, antiderivative size = 17

$$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x\sqrt{\tan(a+bx^2)}}{b}$$

output x*tan(b*x^2+a)^(1/2)/b

Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x\sqrt{\tan(a+bx^2)}}{b}$$

input Integrate[x^2/Sqrt[Tan[a + b*x^2]] + Sqrt[Tan[a + b*x^2]]/b + x^2*Tan[a + b*x^2]^(3/2), x]

output (x*Sqrt[Tan[a + b*x^2]])/b

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(x^2 \tan^{\frac{3}{2}}(a + bx^2) + \frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} \right) dx$$

\downarrow 2009

$$\int x^2 \tan^{\frac{3}{2}}(bx^2 + a) dx + \int \frac{x^2}{\sqrt{\tan(bx^2 + a)}} dx + \int \frac{\sqrt{\tan(bx^2 + a)}}{b} dx$$

input `Int[x^2/Sqrt[Tan[a + b*x^2]] + Sqrt[Tan[a + b*x^2]]/b + x^2*Tan[a + b*x^2]^(3/2), x]`

output `$Aborted`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [F]

$$\int \left(\frac{x^2}{\sqrt{\tan(bx^2 + a)}} + \frac{\sqrt{\tan(bx^2 + a)}}{b} + x^2 \tan(bx^2 + a)^{\frac{3}{2}} \right) dx$$

input `int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2), x)`

output `int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2), x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx = \frac{x \sqrt{\tan(bx^2 + a)}}{b}$$

input `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="fricas")`

output `x*sqrt(tan(b*x^2 + a))/b`

Sympy [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx \\ &= \frac{\int \frac{bx^2}{\sqrt{\tan(a+bx^2)}} dx + \int bx^2 \tan^{\frac{3}{2}}(a + bx^2) dx + \int \sqrt{\tan(a + bx^2)} dx}{b} \end{aligned}$$

input `integrate(x**2/tan(b*x**2+a)**(1/2)+tan(b*x**2+a)**(1/2)/b+x**2*tan(b*x**2+a)**(3/2),x)`

output `(Integral(b*x**2/sqrt(tan(a + b*x**2)), x) + Integral(b*x**2*tan(a + b*x**2)**(3/2), x) + Integral(sqrt(tan(a + b*x**2)), x))/b`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx \\ &= \int x^2 \tan(bx^2 + a)^{\frac{3}{2}} + \frac{x^2}{\sqrt{\tan(bx^2 + a)}} + \frac{\sqrt{\tan(bx^2 + a)}}{b} dx \end{aligned}$$

input `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="maxima")`

output `integrate(x^2*tan(b*x^2 + a)^(3/2) + x^2/sqrt(tan(b*x^2 + a)) + sqrt(tan(b*x^2 + a))/b, x)`

Giac [F(-2)]

Exception generated.

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx$$

= Exception raised: TypeError

input `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:Unable to divide, perhaps due to ro
unding error%%%{-1,[0,2,1,0]%%%} / %%%{1,[0,0,0,1]%%%} Error: Bad Argument
Value`

Mupad [B] (verification not implemented)

Time = 9.66 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.65

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx = \frac{x \sqrt{-\frac{e^{2i b x^2+a 2i} 1i-i}{e^{2i b x^2+a 2i}+1}}}{b}$$

input `int(tan(a + b*x^2)^(1/2)/b + x^2/tan(a + b*x^2)^(1/2) + x^2*tan(a + b*x^2)^(3/2),x)`

output `(x*(-(exp(a*2i + b*x^2*2i)*1i - 1i)/(exp(a*2i + b*x^2*2i) + 1))^(1/2))/b`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx = \frac{\sqrt{\tan(bx^2 + a)} x}{b}$$

input `int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x)`

output `(sqrt(tan(a + b*x**2))*x)/b`

3.18 $\int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx$

Optimal result	166
Mathematica [A] (verified)	167
Rubi [A] (verified)	167
Maple [A] (verified)	170
Fricas [A] (verification not implemented)	170
Sympy [A] (verification not implemented)	171
Maxima [F(-2)]	171
Giac [A] (verification not implemented)	172
Mupad [B] (verification not implemented)	172
Reduce [F]	173

Optimal result

Integrand size = 23, antiderivative size = 189

$$\begin{aligned} \int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx = & \frac{3id^3x}{8af^3} - \frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} \\ & - \frac{3d^3}{8f^4(a+ia\tan(e+fx))} - \frac{3id^2(c+dx)}{4f^3(a+ia\tan(e+fx))} \\ & + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} \end{aligned}$$

output

```
3/8*I*d^3*x/a/f^3-3/8*d*(d*x+c)^2/a/f^2-1/4*I*(d*x+c)^3/a/f+1/8*(d*x+c)^4/
a/d-3/8*d^3/f^4/(a+I*a*tan(f*x+e))-3/4*I*d^2*(d*x+c)/f^3/(a+I*a*tan(f*x+e))
)+3/4*d*(d*x+c)^2/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)^3/f/(a+I*a*tan(f*x+
e))
```

Mathematica [A] (verified)

Time = 0.71 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.47

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ = \frac{\sec(e + fx)(\cos(fx) + i \sin(fx)) ((4ic^3 f^3 + 6c^2 df^2(1 + 2ifx) + 6cd^2 f(-i + 2fx + 2if^2 x^2) + d^3 (-3 - 6ifx + 2f^2 x^2)))}{(16*f^4*(a + I*a*Tan[e + f*x]))}$$

input `Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x]), x]`

output
$$\begin{aligned} & (\text{Sec}[e + f*x]*(\text{Cos}[f*x] + I*\text{Sin}[f*x])*(((4*I)*c^3*f^3 + 6*c^2*d*f^2*(1 + (2*I)*f*x) + 6*c*d^2*f*(-I + 2*f*x + (2*I)*f^2*x^2) + d^3*(-3 - (6*I)*f*x + 6*f^2*x^2 + (4*I)*f^3*x^3))*\text{Cos}[2*f*x]*(\text{Cos}[e] - I*\text{Sin}[e]) + 2*f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(\text{Cos}[e] + I*\text{Sin}[e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(-I + 2*f*x) + 6*c*d^2*f*(-1 - (2*I)*f*x + 2*f^2*x^2) + d^3*(3*I - 6*f*x - (6*I)*f^2*x^2 + 4*f^3*x^3))*(\text{Cos}[e] - I*\text{Sin}[e])*\text{Sin}[2*f*x])) \\ & /(16*f^4*(a + I*a*Tan[e + f*x])) \end{aligned}$$

Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.391, Rules used = {3042, 4206, 3042, 4206, 3042, 4206, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{4206} \\ & - \frac{3id \int \frac{(c+dx)^2}{i\tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} + \frac{(c+dx)^4}{8ad} \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3042} \\
& - \frac{3id \int \frac{(c+dx)^2}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
& \quad \downarrow \text{4206} \\
& - \frac{3id \left(-\frac{id \int \frac{c+dx}{i \tan(e+fx)a+a} dx}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
& \quad \downarrow \text{3042} \\
& - \frac{3id \left(-\frac{id \int \frac{c+dx}{i \tan(e+fx)a+a} dx}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
& \quad \downarrow \text{4206} \\
& - \frac{3id \left(-\frac{id \left(-\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \\
& \quad \quad \quad \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
& \quad \downarrow \text{3042} \\
& - \frac{3id \left(-\frac{id \left(-\frac{id \int \frac{1}{i \tan(e+fx)a+a} dx}{2f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \right)}{2f} + \\
& \quad \quad \quad \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
& \quad \downarrow \text{3960} \\
& - \frac{3id \left(-\frac{id \left(\frac{\int 1 dx}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right)}{f} + \frac{i(c+dx)}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^2}{4ad} \right)}{2f} + \frac{i(c+dx)^2}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^3}{6ad} \\
& \quad \quad \quad \frac{i(c+dx)^3}{2f(a+ia \tan(e+fx))} + \frac{(c+dx)^4}{8ad} \\
& \quad \downarrow \text{24}
\end{aligned}$$

$$\begin{aligned}
 & \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} - \\
 3id \left(\frac{\frac{i(c+dx)^2}{2f(a+ia\tan(e+fx))}}{} - \frac{id\left(\frac{i(c+dx)}{2f(a+ia\tan(e+fx))} + \frac{(c+dx)^2}{4ad} - \frac{id\left(\frac{x}{2a} + \frac{i}{2f(a+ia\tan(e+fx))}\right)}{2f}\right)}{f} + \frac{(c+dx)^3}{6ad} \right) + \\
 & \frac{2f}{8ad} \frac{(c+dx)^4}{(c+dx)^4}
 \end{aligned}$$

input `Int[(c + d*x)^3/(a + I*a*Tan[e + f*x]), x]`

output `(c + d*x)^4/(8*a*d) + ((I/2)*(c + d*x)^3)/(f*(a + I*a*Tan[e + f*x])) - (((3*I)/2)*d*((c + d*x)^3/(6*a*d) + ((I/2)*(c + d*x)^2)/(f*(a + I*a*Tan[e + f*x]))) - (I*d*((c + d*x)^2/(4*a*d) + ((I/2)*(c + d*x))/(f*(a + I*a*Tan[e + f*x]))) - ((I/2)*d*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/f))/f`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n)/(2*b*d*n), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4206 `Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Simp[a*d*(m/(2*b*f)) Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.90

method	result
risch	$\frac{d^3x^4}{8a} + \frac{d^2cx^3}{2a} + \frac{3dc^2x^2}{4a} + \frac{c^3x}{2a} + \frac{c^4}{8ad} + \frac{i(4d^3x^3f^3 + 12cd^2f^3x^2 - 6id^3f^2x^2 + 12c^2df^3x - 12icd^2f^2x + 4c^3f^3 - 6ic^2df^2 - 6ad^2f^3)}{16af^4}$

input `int((d*x+c)^3/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/8/a*d^3*x^4 + 1/2/a*d^2*c*x^3 + 3/4/a*d*c^2*x^2 + 1/2/a*c^3*x + 1/8/a/d*c^4 + 1/16 \\ & *I*(4*d^3*x^3*f^3 - 6*I*d^3*f^2*x^2 + 12*c*d^2*f^3*x^2 - 12*I*c*d^2*f^2*x + 12*c^2 \\ & *d*f^3*x - 6*I*c^2*d*f^2 + 4*c^3*f^3 - 6*d^3*f*x + 3*I*d^3 - 6*c*d^2*f)/a/f^4*\exp(-2 \\ & *I*(f*x+e)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.87

$$\begin{aligned} & \int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ & = \frac{(4i d^3 f^3 x^3 + 4i c^3 f^3 + 6 c^2 d f^2 - 6i c d^2 f - 3 d^3 - 6 (-2i c d^2 f^3 - d^3 f^2) x^2 - 6 (-2i c^2 d f^3 - 2 c d^2 f^2 + i d^3)}{16 a f^4} \end{aligned}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output
$$\begin{aligned} & 1/16*(4*I*d^3*f^3*x^3 + 4*I*c^3*f^3 + 6*c^2*d*f^2 - 6*I*c*d^2*f - 3*d^3 - \\ & 6*(-2*I*c*d^2*f^3 - d^3*f^2)*x^2 - 6*(-2*I*c^2*d*f^3 - 2*c*d^2*f^2 + I*d^3 \\ & *f)*x + 2*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x)* \\ & e^(2*I*f*x + 2*I*e))*e^(-2*I*f*x - 2*I*e)/(a*f^4) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.37

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx$$

$$= \begin{cases} \frac{(4ic^3f^3 + 12ic^2df^3x + 6c^2df^2 + 12icd^2f^3x^2 + 12cd^2f^2x - 6icd^2f + 4id^3f^3x^3 + 6d^3f^2x^2 - 6id^3fx - 3d^3)e^{-2ie}e^{-2ifx}}{16af^4} & \text{for } af^4e^{2ie} \neq 0 \\ \frac{c^3xe^{-2ie}}{2a} + \frac{3c^2dx^2e^{-2ie}}{4a} + \frac{cd^2x^3e^{-2ie}}{2a} + \frac{d^3x^4e^{-2ie}}{8a} \\ + \frac{c^3x}{2a} + \frac{3c^2dx^2}{4a} + \frac{cd^2x^3}{2a} + \frac{d^3x^4}{8a} & \text{otherwise} \end{cases}$$

input `integrate((d*x+c)**3/(a+I*a*tan(f*x+e)),x)`

output `Piecewise(((4*I*c**3*f**3 + 12*I*c**2*d*f**3*x + 6*c**2*d*f**2 + 12*I*c*d**2*f**3*x**2 + 12*c*d**2*f**2*x - 6*I*c*d**2*f + 4*I*d**3*f**3*x**3 + 6*d**3*f**2*x**2 - 6*I*d**3*f*x - 3*d**3)*exp(-2*I*e)*exp(-2*I*f*x)/(16*a*f**4), Ne(a*f**4*exp(2*I*e), 0)), (c**3*x*exp(-2*I*e)/(2*a) + 3*c**2*d*x**2*exp(-2*I*e)/(4*a) + c*d**2*x**3*exp(-2*I*e)/(2*a) + d**3*x**4*exp(-2*I*e)/(8*a), True)) + c**3*x/(2*a) + 3*c**2*d*x**2/(4*a) + c*d**2*x**3/(2*a) + d*x**4/(8*a)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.99

$$\int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx = \frac{(2d^3f^4x^4e^{(2i\ln f x + 2i\ln e)}) + 8cd^2f^4x^3e^{(2i\ln f x + 2i\ln e)}) + 12c^2df^4x^2e^{(2i\ln f x + 2i\ln e)}) + 4i d^3f^3x^3 + 8c^3f^4xe^{(2i\ln f x + 2i\ln e)}) + 12i}{16}$$

```
input integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="giac")
```

```

output 1/16*(2*d^3*f^4*x^4*e^(2*I*f*x + 2*I*e) + 8*c*d^2*f^4*x^3*e^(2*I*f*x + 2*I*e) + 12*c^2*d*f^4*x^2*e^(2*I*f*x + 2*I*e) + 4*I*d^3*f^3*x^3 + 8*c^3*f^4*x*e^(2*I*f*x + 2*I*e) + 12*I*c*d^2*f^3*x^2 + 12*I*c^2*d*f^3*x + 6*d^3*f^2*x^2 + 4*I*c^3*f^3 + 12*c*d^2*f^2*x + 6*c^2*d*f^2 - 6*I*d^3*f*x - 6*I*c*d^2*f - 3*d^3)*e^(-2*I*f*x - 2*I*e)/(a*f^4)

```

Mupad [B] (verification not implemented)

Time = 8.92 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.24

$$\begin{aligned} & \int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx \\ & \equiv \frac{8 c^3 f^4 x - 3 d^3 \cos(2e + 2fx) + 4 c^3 f^3 \sin(2e + 2fx) + 2 d^3 f^4 x^4 + 6 c^2 d f^2 \cos(2e + 2fx) + 12 c^2 d^2 f^3 x^3}{\dots} \end{aligned}$$

input int((c + d*x)^3/(a + a*tan(e + f*x)*1i),x)

output

$$(d^3 \sin(2e + 2fx) * 3i - 3d^3 \cos(2e + 2fx) + 8c^3 f^4 x + c^3 f^3 \cos(2e + 2fx) * 4i + 4c^3 f^3 \sin(2e + 2fx) + 2d^3 f^4 x^4 + 6c^2 d^2 f^2 \cos(2e + 2fx) - c^2 d^2 f^2 \sin(2e + 2fx) * 6i + 12c^2 d^2 f^4 x^2 + 8c^2 d^2 f^4 x^3 + 6d^3 f^2 x^2 \cos(2e + 2fx) + d^3 f^3 x^3 \cos(2e + 2fx) * 4i - d^3 f^2 x^2 \sin(2e + 2fx) * 6i + 4d^3 f^3 x^3 \sin(2e + 2fx) - c^2 d^2 f \cos(2e + 2fx) * 6i - 6c^2 d^2 f \sin(2e + 2fx) - d^3 f x \cos(2e + 2fx) * 6i - 6d^3 f x \sin(2e + 2fx) + 12c^2 d^2 f^2 x \cos(2e + 2fx) + c^2 d^2 f^3 x \cos(2e + 2fx) * 12i - c^2 d^2 f^2 x \sin(2e + 2fx) * 12i + 12c^2 d^2 f^3 x \sin(2e + 2fx) + c^2 d^2 f^3 x^2 \cos(2e + 2fx) * 12i + 12c^2 d^2 f^3 x^2 \sin(2e + 2fx)) / (16a^2 f^4)$$

Reduce [F]

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ = \frac{\left(\int \frac{x^3}{\tan(fx+e)i+1} dx \right) d^3 + 3 \left(\int \frac{x^2}{\tan(fx+e)i+1} dx \right) c d^2 + 3 \left(\int \frac{x}{\tan(fx+e)i+1} dx \right) c^2 d + \left(\int \frac{1}{\tan(fx+e)i+1} dx \right) c^3}{a}$$

input

```
int((d*x+c)^3/(a+I*a*tan(f*x+e)),x)
```

output

```
(int(x**3/(\tan(e + fx)*i + 1),x)*d**3 + 3*int(x**2/(\tan(e + fx)*i + 1),x)*c*d**2 + 3*int(x/(\tan(e + fx)*i + 1),x)*c**2*d + int(1/(\tan(e + fx)*i + 1),x)*c**3)/a
```

3.19 $\int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx$

Optimal result	174
Mathematica [A] (verified)	174
Rubi [A] (verified)	175
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	177
Sympy [A] (verification not implemented)	178
Maxima [F(-2)]	178
Giac [A] (verification not implemented)	179
Mupad [B] (verification not implemented)	179
Reduce [F]	180

Optimal result

Integrand size = 23, antiderivative size = 137

$$\begin{aligned} \int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx = & -\frac{d^2x}{4af^2} - \frac{i(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{id^2}{4f^3(a+ia\tan(e+fx))} \\ & + \frac{d(c+dx)}{2f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^2}{2f(a+ia\tan(e+fx))} \end{aligned}$$

output
$$\begin{aligned} & -1/4*d^2*x/a/f^2-1/4*I*(d*x+c)^2/a/f+1/6*(d*x+c)^3/a/d-1/4*I*d^2/f^3/(a+I* \\ & a*tan(f*x+e))+1/2*d*(d*x+c)/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)^2/f/(a+I* \\ & a*tan(f*x+e)) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.43 (sec), antiderivative size = 178, normalized size of antiderivative = 1.30

$$\begin{aligned} & \int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx \\ &= \frac{\sec(e+fx)(\cos(fx)+i\sin(fx))((d+(1+i)cf+(1+i)dfx)((1+i)cf+d(-i+(1+i)fx))\cos(2fx))}{a+ia\tan(e+fx)} \end{aligned}$$

input `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x]), x]`

output

$$(Sec[e + f*x]*(Cos[f*x] + I*Sin[f*x])*((d + (1 + I)*c*f + (1 + I)*d*f*x)*(1 + I)*c*f + d*(-I + (1 + I)*f*x))*Cos[2*f*x]*(Cos[e] - I*Sin[e]) + (4*f^3*x*(3*c^2 + 3*c*d*x + d^2*x^2)*(Cos[e] + I*Sin[e]))/3 - I*(d + (1 + I)*c*f + (1 + I)*d*f*x)*((1 + I)*c*f + d*(-I + (1 + I)*f*x))*(Cos[e] - I*Sin[e])*Sin[2*f*x]))/(8*f^3*(a + I*a*Tan[e + f*x]))$$

Rubi [A] (verified)

Time = 0.50 (sec), antiderivative size = 147, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {3042, 4206, 3042, 4206, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c+dx)^2}{a+i a \tan(e+f x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{(c+dx)^2}{a+i a \tan(e+f x)} dx \\
 & \quad \downarrow \textcolor{blue}{4206} \\
 & -\frac{i d \int \frac{c+dx}{i \tan(e+f x) a+a} dx}{f} + \frac{i(c+dx)^2}{2 f(a+i a \tan(e+f x))} + \frac{(c+dx)^3}{6 ad} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{i d \int \frac{c+dx}{i \tan(e+f x) a+a} dx}{f} + \frac{i(c+dx)^2}{2 f(a+i a \tan(e+f x))} + \frac{(c+dx)^3}{6 ad} \\
 & \quad \downarrow \textcolor{blue}{4206} \\
 & -\frac{i d \left(-\frac{i d \int \frac{1}{i \tan(e+f x) a+a} dx}{2 f} + \frac{i(c+dx)}{2 f(a+i a \tan(e+f x))} + \frac{(c+dx)^2}{4 ad} \right)}{f} + \frac{i(c+dx)^2}{2 f(a+i a \tan(e+f x))} + \frac{(c+dx)^3}{6 ad} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{i d \left(-\frac{i d \int \frac{1}{i \tan(e+f x) a+a} dx}{2 f} + \frac{i(c+dx)}{2 f(a+i a \tan(e+f x))} + \frac{(c+dx)^2}{4 ad} \right)}{f} + \frac{i(c+dx)^2}{2 f(a+i a \tan(e+f x))} + \frac{(c+dx)^3}{6 ad}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3960} \\
 - \frac{id\left(-\frac{id\left(\frac{f_1 dx}{2a} + \frac{i}{2f(a+ia\tan(e+fx))}\right)}{2f} + \frac{i(c+dx)}{2f(a+ia\tan(e+fx))} + \frac{(c+dx)^2}{4ad}\right)}{f} + \frac{i(c+dx)^2}{2f(a+ia\tan(e+fx))} + \\
 & \quad \frac{(c+dx)^3}{6ad} \\
 & \quad \downarrow \text{24} \\
 \frac{i(c+dx)^2}{2f(a+ia\tan(e+fx))} - \frac{id\left(\frac{i(c+dx)}{2f(a+ia\tan(e+fx))} + \frac{(c+dx)^2}{4ad} - \frac{id\left(\frac{x}{2a} + \frac{i}{2f(a+ia\tan(e+fx))}\right)}{2f}\right)}{f} + \\
 & \quad \frac{(c+dx)^3}{6ad}
 \end{aligned}$$

input `Int[(c + d*x)^2/(a + I*a*Tan[e + f*x]), x]`

output `(c + d*x)^3/(6*a*d) + ((I/2)*(c + d*x)^2)/(f*(a + I*a*Tan[e + f*x])) - (I*d*((c + d*x)^2/(4*a*d) + ((I/2)*(c + d*x))/(f*(a + I*a*Tan[e + f*x]))) - ((I/2)*d*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/f))/f`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4206

```

Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x]
Simplify :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Simp[a*d*(m/(2*b*f))
Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m
/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[
a^2 + b^2, 0] && GtQ[m, 0]

```

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{d^2 x^3}{6 a} + \frac{d c x^2}{2 a} + \frac{c^2 x}{2 a} + \frac{c^3}{6 a d} + \frac{i (2 d^2 x^2 f^2 + 4 c d f^2 x - 2 i d^2 f x + 2 c^2 f^2 - 2 i c d f - d^2) e^{-2 i (f x + e)}}{8 a f^3}$	108

input `int((d*x+c)^2/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output $\frac{1}{6} a^2 d^2 x^3 + \frac{1}{2} a^2 d c x^2 + \frac{1}{2} a^2 c^2 x + \frac{1}{6} a^2 d c^3 + \frac{1}{8} I^2 (2 d^2 x^2 f^2 + 4 c d f^2 x - 2 i d^2 f x + 2 c^2 f^2 - 2 i c d f - d^2) e^{-2 i (f x + e)}$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.77

$$\begin{aligned} & \int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx \\ &= \frac{(6i d^2 f^2 x^2 + 6i c^2 f^2 + 6 cdf - 3i d^2 - 6(-2i cdf^2 - d^2 f)x + 4(d^2 f^3 x^3 + 3 cdf^3 x^2 + 3 c^2 f^3 x)e^{(2i f x + 2i e)})}{24 a f^3} \end{aligned}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output $\frac{1}{24} (6 I^2 d^2 f^2 x^2 + 6 I^2 c^2 f^2 + 6 c d f - 3 I^2 d^2 - 6 (-2 I c d f^2 - d^2 f)x + 4 (d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x)e^{(2 I f x + 2 I e)}) e^{(-2 I f x - 2 I e)} / (a f^3)$

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.20

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx$$

$$= \begin{cases} \frac{(2ic^2f^2 + 4icdf^2x + 2cdf + 2id^2f^2x^2 + 2d^2fx - id^2)e^{-2ie}e^{-2ifx}}{8af^3} & \text{for } af^3e^{2ie} \neq 0 \\ \frac{c^2xe^{-2ie}}{2a} + \frac{cdx^2e^{-2ie}}{2a} + \frac{d^2x^3e^{-2ie}}{6a} & \text{otherwise} \end{cases}$$

input `integrate((d*x+c)**2/(a+I*a*tan(f*x+e)),x)`

output `Piecewise(((2*I*c**2*f**2 + 4*I*c*d*f**2*x + 2*c*d*f + 2*I*d**2*f**2*x**2 + 2*d**2*f*x - I*d**2)*exp(-2*I*e)*exp(-2*I*f*x)/(8*a*f**3), Ne(a*f**3*exp(2*I*e), 0)), (c**2*x*exp(-2*I*e)/(2*a) + c*d*x**2*exp(-2*I*e)/(2*a) + d**2*x**3*exp(-2*I*e)/(6*a), True)) + c**2*x/(2*a) + c*d*x**2/(2*a) + d**2*x**3/(6*a)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.87

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx$$

$$= \frac{(4 d^2 f^3 x^3 e^{(2i f x + 2i e)}) + 12 c d f^3 x^2 e^{(2i f x + 2i e)} + 12 c^2 f^3 x e^{(2i f x + 2i e)} + 6i d^2 f^2 x^2 + 12i c d f^2 x + 6i c^2 f^2 + 6 d^2}{24 a f^3}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output $\frac{1}{24} (4 d^2 f^3 x^3 e^{(2 I f x + 2 I e)} + 12 c d f^3 x^2 e^{(2 I f x + 2 I e)} + 12 c^2 f^3 x e^{(2 I f x + 2 I e)} + 6 I d^2 f^2 x^2 + 12 I c d f^2 x + 6 I c^2 f^2 + 6 d^2)$

Mupad [B] (verification not implemented)

Time = 8.73 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.76

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx$$

$$= \frac{12 c^2 f^3 x - 3 d^2 \sin(2 e + 2 f x) + 6 c^2 f^2 \sin(2 e + 2 f x) + 4 d^2 f^3 x^3 + 6 c d f \cos(2 e + 2 f x) + 6 d^2 f^2}{a}$$

input `int((c + d*x)^2/(a + a*tan(e + f*x)*1i),x)`

output $(12 c^2 f^3 x - 3 d^2 \sin(2 e + 2 f x) - d^2 \cos(2 e + 2 f x) * 3i + c^2 f^2 * \cos(2 e + 2 f x) * 6i + 6 c^2 f^2 * \sin(2 e + 2 f x) + 4 d^2 f^3 x^3 + 6 c d f \cos(2 e + 2 f x) - c d f \sin(2 e + 2 f x) * 6i + d^2 f^2 * 2 x^2 * \cos(2 e + 2 f x) * 6i + 6 d^2 f^2 * 2 x^2 * \sin(2 e + 2 f x) + 12 c d f^3 x^2 + 6 d^2 f^2 * x * \cos(2 e + 2 f x) - d^2 f * x * \sin(2 e + 2 f x) * 6i + c d f^2 * 2 x * \cos(2 e + 2 f x) * 12i + 12 c d f^2 * 2 x * \sin(2 e + 2 f x)) / (24 a f^3)$

Reduce [F]

$$\begin{aligned} & \int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx \\ &= \frac{\left(\int \frac{x^2}{\tan(fx+e)i+1} dx \right) d^2 + 2 \left(\int \frac{x}{\tan(fx+e)i+1} dx \right) cd + \left(\int \frac{1}{\tan(fx+e)i+1} dx \right) c^2}{a} \end{aligned}$$

input `int((d*x+c)^2/(a+I*a*tan(f*x+e)),x)`

output `(int(x**2/(tan(e + f*x)*i + 1),x)*d**2 + 2*int(x/(tan(e + f*x)*i + 1),x)*c *d + int(1/(tan(e + f*x)*i + 1),x)*c**2)/a`

3.20 $\int \frac{c+dx}{a+ia\tan(e+fx)} dx$

Optimal result	181
Mathematica [A] (verified)	181
Rubi [A] (verified)	182
Maple [A] (verified)	183
Fricas [A] (verification not implemented)	184
Sympy [A] (verification not implemented)	184
Maxima [F(-2)]	185
Giac [A] (verification not implemented)	185
Mupad [B] (verification not implemented)	185
Reduce [F]	186

Optimal result

Integrand size = 21, antiderivative size = 84

$$\begin{aligned} \int \frac{c+dx}{a+ia\tan(e+fx)} dx &= -\frac{idx}{4af} + \frac{(c+dx)^2}{4ad} + \frac{d}{4f^2(a+ia\tan(e+fx))} \\ &\quad + \frac{i(c+dx)}{2f(a+ia\tan(e+fx))} \end{aligned}$$

output
$$-\frac{1}{4}I*d*x/a/f + \frac{1}{4}(d*x+c)^2/a/d + \frac{1}{4}d/f^2/(a+I*a*tan(f*x+e)) + \frac{1}{2}I*(d*x+c)/f/(a+I*a*tan(f*x+e))$$

Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.14

$$\begin{aligned} &\int \frac{c+dx}{a+ia\tan(e+fx)} dx \\ &= \frac{-i(2cf(i+2fx)+d(1+2ifx+2f^2x^2))+(2cf(-i+2fx)+d(-1-2ifx+2f^2x^2))\tan(e+fx)}{8af^2(-i+\tan(e+fx))} \end{aligned}$$

input
$$\text{Integrate}[(c+d*x)/(a+I*a*Tan[e+f*x]), x]$$

output $((-I)*(2*c*f*(I + 2*f*x) + d*(1 + (2*I)*f*x + 2*f^2*x^2)) + (2*c*f*(-I + 2*f*x) + d*(-1 - (2*I)*f*x + 2*f^2*x^2))*\text{Tan}[e + f*x])/(8*a*f^2*(-I + \text{Tan}[e + f*x]))$

Rubi [A] (verified)

Time = 0.33 (sec), antiderivative size = 90, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4206, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{c+dx}{a+i a \tan(e+f x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{c+dx}{a+i a \tan(e+f x)} dx \\
 & \quad \downarrow \textcolor{blue}{4206} \\
 & -\frac{i d \int \frac{1}{i \tan(e+f x) a+a} dx}{2 f}+\frac{i(c+d x)}{2 f(a+i a \tan(e+f x))}+\frac{(c+d x)^2}{4 a d} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{i d \int \frac{1}{i \tan(e+f x) a+a} dx}{2 f}+\frac{i(c+d x)}{2 f(a+i a \tan(e+f x))}+\frac{(c+d x)^2}{4 a d} \\
 & \quad \downarrow \textcolor{blue}{3960} \\
 & -\frac{i d\left(\frac{\int 1 dx}{2 a}+\frac{i}{2 f(a+i a \tan(e+f x))}\right)}{2 f}+\frac{i(c+d x)}{2 f(a+i a \tan(e+f x))}+\frac{(c+d x)^2}{4 a d} \\
 & \quad \downarrow \textcolor{blue}{24} \\
 & \frac{i(c+d x)}{2 f(a+i a \tan(e+f x))}+\frac{(c+d x)^2}{4 a d}-\frac{i d\left(\frac{x}{2 a}+\frac{i}{2 f(a+i a \tan(e+f x))}\right)}{2 f}
 \end{aligned}$$

input $\text{Int}[(c + d*x)/(a + I*a*\text{Tan}[e + f*x]),x]$

output
$$(c + d*x)^2/(4*a*d) + ((I/2)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])) - ((I/2)*d*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/f$$

Definitions of rubi rules used

rule 24
$$\text{Int}[a_-, x_{\text{Symbol}}] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$$

rule 3042
$$\text{Int}[u_-, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 3960
$$\text{Int}[(a_+ + b_-)*\tan[(c_+ + d_-)*(x_+)]^n, x_{\text{Symbol}}] \rightarrow \text{Simp}[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + \text{Simp}[1/(2*a) \text{Int}[(a + b*Tan[c + d*x])^{n+1}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{LtQ}[n, 0]$$

rule 4206
$$\text{Int}[(c_+ + d_-)*(x_+)^m / ((a_+ + b_-)*\tan[(e_+ + f_-)*(x_+)]), x_{\text{Symbol}}] \rightarrow \text{Simp}[(c + d*x)^{m+1}/(2*a*d*(m+1)), x] + (\text{Simp}[a*d*(m/(2*b*f)) \text{Int}[(c + d*x)^{m-1}/(a + b*Tan[e + f*x]), x], x] - \text{Simp}[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{GtQ}[m, 0]$$

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.60

method	result	size
risch	$\frac{dx^2}{4a} + \frac{cx}{2a} + \frac{i(2dfx+2cf-id)e^{-2i(fx+e)}}{8af^2}$	50
norman	$\frac{dx^2}{4a} + \frac{2icf+d}{4af^2} + \frac{dx^2 \tan(fx+e)^2}{4a} + \frac{(2cf-id)\tan(fx+e)}{4af^2} + \frac{(2cf+id)x}{4af} + \frac{dx \tan(fx+e)}{2af} + \frac{(2cf-id)x \tan(fx+e)^2}{4fa}$ $1+\tan(fx+e)^2$	139

input
$$\text{int}((d*x+c)/(a+I*a*Tan(f*x+e)), x, \text{method}=\text{_RETURNVERBOSE})$$

output
$$1/4/a*d*x^2 + 1/2/a*c*x + 1/8*I*(2*d*f*x - I*d + 2*c*f)/a/f^2 * \exp(-2*I*(f*x+e))$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.65

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx \\ = \frac{(2i dfx + 2i cf + 2(df^2 x^2 + 2cf^2 x)e^{(2ifx+2ie)} + d)e^{(-2ifx-2ie)}}{8af^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output `1/8*(2*I*d*f*x + 2*I*c*f + 2*(d*f^2*x^2 + 2*c*f^2*x)*e^(2*I*f*x + 2*I*e) + d)*e^(-2*I*f*x - 2*I*e)/(a*f^2)`

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.10

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \begin{cases} \frac{(2icf + 2idfx + d)e^{-2ie} e^{-2ifx}}{8af^2} & \text{for } af^2 e^{2ie} \neq 0 \\ \frac{cxe^{-2ie}}{2a} + \frac{dx^2 e^{-2ie}}{4a} & \text{otherwise} \end{cases} + \frac{cx}{2a} + \frac{dx^2}{4a}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x)`

output `Piecewise(((2*I*c*f + 2*I*d*f*x + d)*exp(-2*I*e)*exp(-2*I*f*x)/(8*a*f**2), Ne(a*f**2*exp(2*I*e), 0)), (c*x*exp(-2*I*e)/(2*a) + d*x**2*exp(-2*I*e)/(4*a), True)) + c*x/(2*a) + d*x**2/(4*a)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.74

$$\begin{aligned} & \int \frac{c + dx}{a + ia \tan(e + fx)} dx \\ &= \frac{(2 df^2 x^2 e^{(2i f x + 2i e)} + 4 c f^2 x e^{(2i f x + 2i e)} + 2i d f x + 2i c f + d) e^{(-2i f x - 2i e)}}{8 a f^2} \end{aligned}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output $\frac{1/8*(2*d*f^2*x^2*e^(2*I*f*x + 2*I*e) + 4*c*f^2*x*e^(2*I*f*x + 2*I*e) + 2*I*d*f*x + 2*I*c*f + d)*e^{(-2*I*f*x - 2*I*e)})/(a*f^2)}$

Mupad [B] (verification not implemented)

Time = 8.73 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.25

$$\begin{aligned} & \int \frac{c + dx}{a + ia \tan(e + fx)} dx \\ &= \frac{d \cos(2e + 2fx) + 2df^2x^2 + 2cf \sin(2e + 2fx) + 4cf^2x + 2dfx \sin(2e + 2fx) - d \sin(2e + 2fx)}{8af^2} \end{aligned}$$

input `int((c + d*x)/(a + a*tan(e + f*x)*1i),x)`

output
$$\begin{aligned} & (d \cos(2e + 2fx) - d \sin(2e + 2fx)*1i + 2d*f^2*x^2 + c*f \cos(2e + 2fx)*2i + 2c*f \sin(2e + 2fx) + 4c*f^2*x + d*f*x \cos(2e + 2fx)*2i \\ & + 2d*f*x \sin(2e + 2fx))/(8*a*f^2) \end{aligned}$$

Reduce [F]

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \frac{\left(\int \frac{x}{\tan(fx+e)i+1} dx \right) d + \left(\int \frac{1}{\tan(fx+e)i+1} dx \right) c}{a}$$

input `int((d*x+c)/(a+I*a*tan(f*x+e)),x)`

output
$$(\int x/(\tan(e + fx)*i + 1) dx)*d + \int 1/(\tan(e + fx)*i + 1) dx*c/a$$

3.21 $\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx$

Optimal result	187
Mathematica [A] (verified)	188
Rubi [A] (verified)	188
Maple [A] (verified)	191
Fricas [A] (verification not implemented)	191
Sympy [F]	191
Maxima [A] (verification not implemented)	192
Giac [A] (verification not implemented)	192
Mupad [F(-1)]	193
Reduce [F]	193

Optimal result

Integrand size = 23, antiderivative size = 161

$$\begin{aligned} \int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx = & \frac{\cos\left(2e - \frac{2cf}{d}\right) \operatorname{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{2ad} \\ & + \frac{\log(c+dx)}{2ad} \\ & - \frac{i \operatorname{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{2ad} \\ & - \frac{i \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(\frac{2cf}{d} + 2fx\right)}{2ad} \\ & - \frac{\sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(\frac{2cf}{d} + 2fx\right)}{2ad} \end{aligned}$$

output
$$\begin{aligned} & 1/2*\cos(-2*e+2*c*f/d)*\operatorname{Ci}(2*c*f/d+2*f*x)/a/d+1/2*\ln(d*x+c)/a/d+1/2*I*\operatorname{Ci}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a/d-1/2*I*\cos(-2*e+2*c*f/d)*\operatorname{Si}(2*c*f/d+2*f*x)/a/d+1/2*\sin(-2*e+2*c*f/d)*\operatorname{Si}(2*c*f/d+2*f*x)/a/d \end{aligned}$$

Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.03

$$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx \\ = \frac{\sec(e+fx) (-i \cos(f(\frac{c}{d}+x)) + \sin(f(\frac{c}{d}+x))) \left(\text{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) (\cos(e-\frac{cf}{d}) - i \sin(e-\frac{cf}{d}))\right)}{2ad(-i + \tan(e+fx))}$$

input `Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])),x]`

output $(\text{Sec}[e+f x]*((-I)*\text{Cos}[f*(c/d+x)] + \text{Sin}[f*(c/d+x)])*(\text{CosIntegral}[(2*f*(c+d x))/d]*(\text{Cos}[e-(c*f)/d] - I*\text{Sin}[e-(c*f)/d]) + \text{Log}[f*(c+d x)]*(\text{Cos}[e-(c*f)/d] + I*\text{Sin}[e-(c*f)/d]) + ((-I)*\text{Cos}[e-(c*f)/d] - \text{Sin}[e-(c*f)/d])* \text{SinIntegral}[(2*f*(c+d x))/d]))/(2*a*d*(-I + \text{Tan}[e+f x]))$

Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {3042, 4209, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx \\ & \quad \downarrow 3042 \\ & \int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx \\ & \quad \downarrow 4209 \\ & -\frac{i \int \frac{\sin(2e+2fx)}{c+dx} dx}{2a} + \frac{\int \frac{\cos(2e+2fx)}{c+dx} dx}{2a} + \frac{\log(c+dx)}{2ad} \\ & \quad \downarrow 3042 \end{aligned}$$

$$\begin{aligned}
& -\frac{i \int \frac{\sin(2e+2fx)}{c+dx} dx}{2a} + \frac{\int \frac{\sin(2e+2fx+\frac{\pi}{2})}{c+dx} dx}{2a} + \frac{\log(c+dx)}{2ad} \\
& \quad \downarrow \text{3784} \\
& -\frac{i \left(\sin \left(2e - \frac{2cf}{d} \right) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx + \cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{2a} + \\
& \quad \frac{\cos \left(2e - \frac{2cf}{d} \right) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx - \sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx}{2a} + \frac{\log(c+dx)}{2ad} \\
& \quad \downarrow \text{3042} \\
& \frac{\cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx}{2a} - \\
& \frac{i \left(\sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{2a} + \frac{\log(c+dx)}{2ad} \\
& \quad \downarrow \text{3780} \\
& \frac{\cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \frac{\sin(2e-\frac{2cf}{d}) \operatorname{Si}(2xf+\frac{2cf}{d})}{d}}{2a} - \\
& \frac{i \left(\sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \frac{\cos(2e-\frac{2cf}{d}) \operatorname{Si}(2xf+\frac{2cf}{d})}{d} \right)}{2a} + \frac{\log(c+dx)}{2ad} \\
& \quad \downarrow \text{3783} \\
& -\frac{i \left(\frac{\operatorname{CosIntegral}(2xf+\frac{2cf}{d}) \sin(2e-\frac{2cf}{d})}{d} + \frac{\cos(2e-\frac{2cf}{d}) \operatorname{Si}(2xf+\frac{2cf}{d})}{d} \right)}{2a} + \\
& \frac{\frac{\operatorname{CosIntegral}(2xf+\frac{2cf}{d}) \cos(2e-\frac{2cf}{d})}{d} - \frac{\sin(2e-\frac{2cf}{d}) \operatorname{Si}(2xf+\frac{2cf}{d})}{d}}{2a} + \frac{\log(c+dx)}{2ad}
\end{aligned}$$

input Int[1/((c + d*x)*(a + I*a*Tan[e + f*x])),x]

output

$$\begin{aligned} & \text{Log}[c + d*x]/(2*a*d) - ((I/2)*((\text{CosIntegral}[(2*c*f)/d + 2*f*x])* \sin[2*e - (2*c*f)/d])/d + (\cos[2*e - (2*c*f)/d]*\text{SinIntegral}[(2*c*f)/d + 2*f*x])/d))/a \\ & + ((\cos[2*e - (2*c*f)/d]*\text{CosIntegral}[(2*c*f)/d + 2*f*x])/d - (\sin[2*e - (2*c*f)/d]*\text{SinIntegral}[(2*c*f)/d + 2*f*x])/d)/(2*a) \end{aligned}$$

Definitions of rubi rules used

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3780 $\text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{SinIntegral}[e + f*x]/d, x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{EqQ}[d*e - c*f, 0]$

rule 3783 $\text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{CosIntegral}[e - \pi/2 + f*x]/d, x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{EqQ}[d*(e - \pi/2) - c*f, 0]$

rule 3784 $\begin{aligned} & \text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \text{Simp}[\cos[d*e - c*f]/d] \\ & \quad \text{Int}[\sin[c*(f/d) + f*x]/(c + d*x), x] + \text{Simp}[\sin[(d*e - c*f)/d]] \\ & \quad \text{Int}[\cos[c*(f/d) + f*x]/(c + d*x), x] /; \text{FreeQ}[\{c, d, e, f\}, x] \\ & \quad \&& \text{NeQ}[d*e - c*f, 0] \end{aligned}$

rule 4209 $\begin{aligned} & \text{Int}[1/(((c_.) + (d_.)*(x_.))*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])), x_{\text{Symbol}}] \rightarrow \text{Simp}[\log[c + d*x]/(2*a*d), x] + (\text{Simp}[1/(2*a) \quad \text{Int}[\cos[2*e + 2*f*x]/(c + d*x), x] + \text{Simp}[1/(2*b) \quad \text{Int}[\sin[2*e + 2*f*x]/(c + d*x), x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0]) \end{aligned}$

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.40

method	result	size
risch	$\frac{\ln(dx+c)}{2ad} - \frac{e^{\frac{2i(cf-de)}{d}} \expIntegral_1(2ifx+2ie+\frac{2i(cf-de)}{d})}{2ad}$	65

input `int(1/(d*x+c)/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output $\frac{1}{2}\ln(d*x+c)/a/d - \frac{1}{2}/a/d*\exp(2*I*(c*f-d*e)/d)*\text{Ei}(1, 2*I*f*x + 2*I*e + 2*I*(c*f - d*e)/d)$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.32

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \frac{\text{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + \log\left(\frac{dx+c}{d}\right)}{2 ad}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output $\frac{1}{2}(\text{Ei}(-2*(I*d*f*x + I*c*f)/d)*e^{-2*(I*d*e - I*c*f)/d} + \log((d*x + c)/d))/(a*d)$

Sympy [F]

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = -\frac{i \int \frac{1}{c \tan(e + fx) - ic + dx \tan(e + fx) - idx} dx}{a}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x)`

output $-\text{I*Integral}(1/(c*\tan(e + f*x) - I*c + d*x*\tan(e + f*x) - I*d*x), x)/a$

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.70

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx =$$

$$-\frac{f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + i f E_1\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) - f \log\left(\frac{a+ia\tan(e+fx)}{c+dx}\right)}{2adf}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output
$$-1/2*(f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f))/(a*d*f)$$

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.86

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx$$

$$= \frac{\cos\left(\frac{2cf}{d}\right) \text{Ci}\left(-\frac{2(df+cf)}{d}\right) + \cos(2e) \log(dx + c) + i \log(dx + c) \sin(2e) + i \text{Ci}\left(-\frac{2(df+cf)}{d}\right) \sin\left(\frac{2cf}{d}\right)}{2(ad \cos(2e) + ad \sin(2e))}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output
$$1/2*(\cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + \cos(2*e)*log(d*x + c) + I*log(d*x + c)*sin(2*e) + I*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - I*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d))/(a*d*cos(2*e) + I*a*d*sin(2*e))$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \int \frac{1}{(a + a \tan(e + fx) 1i) (c + dx)} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)),x)`

output `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)), x)`

Reduce [F]

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \frac{\int \frac{1}{\tan(fx+e)ci+\tan(fx+e)dix+c+dx} dx}{a}$$

input `int(1/(d*x+c)/(a+I*a*tan(f*x+e)),x)`

output `int(1/(\tan(e + f*x)*c*i + \tan(e + f*x)*d*i*x + c + d*x),x)/a`

3.22 $\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx$

Optimal result	194
Mathematica [A] (verified)	195
Rubi [A] (verified)	195
Maple [A] (verified)	198
Fricas [A] (verification not implemented)	198
Sympy [F]	199
Maxima [A] (verification not implemented)	199
Giac [B] (verification not implemented)	200
Mupad [F(-1)]	201
Reduce [F]	201

Optimal result

Integrand size = 23, antiderivative size = 168

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx = & -\frac{if \cos\left(2e - \frac{2cf}{d}\right) \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} \\ & - \frac{f \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{ad^2} \\ & - \frac{f \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} \\ & + \frac{if \sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} \\ & - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \end{aligned}$$

output
$$-\text{I}*f*\cos(-2*e+2*c*f/d)*\text{Ci}(2*c*f/d+2*f*x)/a/d^2+f*\text{Ci}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a/d^2-f*\cos(-2*e+2*c*f/d)*\text{Si}(2*c*f/d+2*f*x)/a/d^2-\text{I}*f*\sin(-2*e+2*c*f/d)*\text{Si}(2*c*f/d+2*f*x)/a/d^2-1/d/(d*x+c)/(a+\text{I}*a*\tan(f*x+e))$$

Mathematica [A] (verified)

Time = 0.69 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.33

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx \\ = \frac{\sec(e+fx) (\cos(\frac{cf}{d}) + i\sin(\frac{cf}{d})) \left(d(i\cos(e+f(-\frac{c}{d}+x)) + i\cos(e+f(\frac{c}{d}+x))) - \sin(e+f(-\frac{c}{d}+x))\right)}{(c+dx)^2(a+ia\tan(e+fx))}$$

input `Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])), x]`

output
$$\begin{aligned} & (\operatorname{Sec}[e + f*x] * (\operatorname{Cos}[(c*f)/d] + I*\operatorname{Sin}[(c*f)/d]) * (d*(I*\operatorname{Cos}[e + f*(-(c/d) + x)] \\ &] + I*\operatorname{Cos}[e + f*(c/d + x)] - \operatorname{Sin}[e + f*(-(c/d) + x)] + \operatorname{Sin}[e + f*(c/d + x)] \\ &]) - 2*f*(c + d*x)*\operatorname{CosIntegral}[(2*f*(c + d*x))/d]*(\operatorname{Cos}[e - (f*(c + d*x))/d] \\ &] - I*\operatorname{Sin}[e - (f*(c + d*x))/d]) + 2*f*(c + d*x)*(I*\operatorname{Cos}[e - (f*(c + d*x))/d] \\ &] + \operatorname{Sin}[e - (f*(c + d*x))/d])* \operatorname{SinIntegral}[(2*f*(c + d*x))/d])/(2*a*d^2*(c \\ & + d*x)*(-I + \operatorname{Tan}[e + f*x])) \end{aligned}$$

Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {3042, 4207, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx \\ & \quad \downarrow \text{4207} \\ & -\frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{if \int \frac{\cos(2e+2fx)}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{3042} \\
& -\frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{if \int \frac{\sin(2e+2fx+\frac{\pi}{2})}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \\
& \quad \downarrow \text{3784} \\
& -\frac{f \left(\sin \left(2e - \frac{2cf}{d} \right) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx + \cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \\
& \quad \frac{if \left(\cos \left(2e - \frac{2cf}{d} \right) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx - \sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \\
& \quad \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \\
& \quad \downarrow \text{3042} \\
& -\frac{if \left(\cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \\
& \quad \frac{f \left(\sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \\
& \quad \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \\
& \quad \downarrow \text{3780} \\
& -\frac{if \left(\cos \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \frac{\sin(2e-\frac{2cf}{d}) \operatorname{Si}(2xf+\frac{2cf}{d})}{d} \right)}{ad} - \\
& \quad \frac{f \left(\sin \left(2e - \frac{2cf}{d} \right) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \frac{\cos(2e-\frac{2cf}{d}) \operatorname{Si}(2xf+\frac{2cf}{d})}{d} \right)}{ad} - \\
& \quad \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \\
& \quad \downarrow \text{3783}
\end{aligned}$$

$$\begin{aligned}
 & -f \left(\frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{d} + \frac{\cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{d} \right) - \\
 & - \frac{i f \left(\frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{d} - \frac{\sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{d} \right)}{ad} - \frac{1}{d(c + dx)(a + ia \tan(e + fx))} \\
 \end{aligned}$$

input `Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])), x]`

output `-((f*((CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/d + (Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d))/(a*d)) - (I*f*((Cos[2*e - (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x])/d - (Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/d))/(a*d) - 1/(d*(c + d*x)*(a + I*a*Tan[e + f*x])))`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3780 `Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 `Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[Cos[(d*e - c*f)/d]] Int[Sin[c*(f/d) + f*x]/(c + d*x), x] + Simp[Sin[(d*e - c*f)/d]] Int[Cos[c*(f/d) + f*x]/(c + d*x), x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 4207

```
Int[1/(((c_) + (d_)*(x_))^2*((a_) + (b_)*tan[(e_) + (f_)*(x_)])), x]
Symb] :> -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Simp[f/(a*d)
Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Simp[f/(b*d) Int[Cos[2*e + 2*
f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0
]
```

Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.57

method	result	size
risch	$-\frac{1}{2da(dx+c)} - \frac{fe^{-2i(fx+e)}}{2a(df+cf)d} + \frac{ife^{\frac{2i(cf-de)}{d}} \expIntegral_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{a d^2}$	96

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e)), x, method=_RETURNVERBOSE)`output `-1/2/d/a/(d*x+c)-1/2/a*f*exp(-2*I*(f*x+e))/(d*f*x+c*f)/d+I/a*f/d^2*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.49

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))} dx \\ = -\frac{\left(\left(2(i dfx + i cf) Ei\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + d \right) e^{(2i fx + 2i e)} + d \right) e^{(-2i fx - 2i e)}}{2(ad^3x + acd^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)), x, algorithm="fricas")`output `-1/2*((2*(I*d*f*x + I*c*f)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + d)*e^(2*I*f*x + 2*I*e) + d)*e^(-2*I*f*x - 2*I*e)/(a*d^3*x + a*c*d^2)`

Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx$$

$$= -\frac{i \int \frac{1}{c^2\tan(e+fx)-ic^2+2cdx\tan(e+fx)-2icdx+d^2x^2\tan(e+fx)-id^2x^2} dx}{a}$$

input `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e)),x)`

output `-I*Integral(1/(c**2*tan(e + f*x) - I*c**2 + 2*c*d*x*tan(e + f*x) - 2*I*c*d*x + d**2*x**2*tan(e + f*x) - I*d**2*x**2), x)/a`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.71

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx =$$

$$-\frac{f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + i f^2 E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) + f^2}{2((fx+e)ad^2-ad^2e+acdf)f}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `-1/2*(f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + f^2)/(((f*x + e)*a*d^2 - a*d^2*e + a*c*d*f)*f)`

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. $1013 \text{ vs. } 2(161) = 322$.

Time = 2.13 (sec) , antiderivative size = 1013, normalized size of antiderivative = 6.03

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx = \text{Too large to display}$$

```
input integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")
```

```

output
1/2*(-2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 2*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 2*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*d*e*f^2*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*c*f^3*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*I*d*e*f^2*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*c*f^3*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c+dx)^2(a+i a \tan(e+f x))} dx = \int \frac{1}{(a+a \tan(e+f x) \operatorname{Ii})(c+d x)^2} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^2),x)`

output `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^2), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+i a \tan(e+f x))} dx \\ &= \frac{\int \frac{1}{\tan(f x+e) c^2 i+2 \tan(f x+e) c d i x+\tan(f x+e) d^2 i x^2+c^2+2 c d x+d^2 x^2} dx}{a} \end{aligned}$$

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x)`

output `int(1/(\tan(e + f*x)*c**2*i + 2*tan(e + f*x)*c*d*i*x + tan(e + f*x)*d**2*i*x**2 + c**2 + 2*c*d*x + d**2*x**2),x)/a`

3.23 $\int \frac{1}{(c+dx)^3(a+ia\tan(e+fx))} dx$

Optimal result	202
Mathematica [A] (verified)	203
Rubi [A] (verified)	203
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	207
Sympy [F]	207
Maxima [A] (verification not implemented)	208
Giac [B] (verification not implemented)	208
Mupad [F(-1)]	209
Reduce [F]	209

Optimal result

Integrand size = 23, antiderivative size = 227

$$\begin{aligned} \int \frac{1}{(c+dx)^3(a+ia\tan(e+fx))} dx = & -\frac{if}{2ad^2(c+dx)} \\ & - \frac{f^2 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & + \frac{if^2 \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^3} \\ & + \frac{if^2 \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & + \frac{f^2 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} \\ & + \frac{if}{d^2(c+dx)(a+ia\tan(e+fx))} \end{aligned}$$

output
$$\begin{aligned} & -1/2*I*f/a/d^2/(d*x+c)-f^2*cos(-2*e+2*c*f/d)*Ci(2*c*f/d+2*f*x)/a/d^3-I*f^2 \\ & *Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^3+I*f^2*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d^3-f^2*sin(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d^3-1/2/d/(d*x+c)^2/(a+I*a*tan(f*x+e))+I*f/d^2/(d*x+c)/(a+I*a*tan(f*x+e)) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.91 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.26

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ = \frac{\sec(e + fx) (\cos(\frac{cf}{d}) + i \sin(\frac{cf}{d})) \left(d(id \cos(e + f(-\frac{c}{d} + x)) + (id + 2cf + 2dfx) \cos(e + f(\frac{c}{d} + x))) -$$

input `Integrate[1/((c + d*x)^3*(a + I*a*Tan[e + f*x])),x]`

output $(\text{Sec}[e + f*x]*(\text{Cos}[(c*f)/d] + I*\text{Sin}[(c*f)/d])*(d*(I*d*\text{Cos}[e + f*(-(c/d) + x)] + (I*d + 2*c*f + 2*d*f*x)*\text{Cos}[e + f*(c/d + x)] - d*\text{Sin}[e + f*(-(c/d) + x)] + d*\text{Sin}[e + f*(c/d + x)] - (2*I)*c*f*\text{Sin}[e + f*(c/d + x)] - (2*I)*d*f*x*\text{Sin}[e + f*(c/d + x)]) + 4*f^2*(c + d*x)^2*\text{CosIntegral}[(2*f*(c + d*x))/d]*(\text{I}*\text{Cos}[e - (f*(c + d*x))/d] + \text{Sin}[e - (f*(c + d*x))/d]) + 4*f^2*(c + d*x)^2*(\text{Cos}[e - (f*(c + d*x))/d] - I*\text{Sin}[e - (f*(c + d*x))/d])* \text{SinIntegral}[(2*f*(c + d*x))/d])/(4*a*d^3*(c + d*x)^2*(-I + \text{Tan}[e + f*x]))$

Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$, Rules used = {3042, 4208, 3042, 4207, 3042, 3784, 3042, 3780, 3783}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ \downarrow \text{3042} \\ \int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ \downarrow \text{4208}$$

$$-\frac{if \int \frac{1}{(c+dx)^2(i\tan(e+fx)a+a)} dx}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}$$

↓ 3042

$$-\frac{if \int \frac{1}{(c+dx)^2(i\tan(e+fx)a+a)} dx}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}$$

↓ 4207

$$-\frac{if \left(-\frac{\int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{\int \frac{\cos(2e+2fx)}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \right)}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}$$

↓ 3042

$$-\frac{if \left(-\frac{\int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} - \frac{\int \frac{\sin(2e+2fx+\frac{\pi}{2})}{c+dx} dx}{ad} - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \right)}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}$$

↓ 3784

$$-\frac{if \left(-\frac{\sin(2e-\frac{2cf}{d}) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx + \cos(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx}{ad} - \frac{if \left(\cos(2e-\frac{2cf}{d}) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx - \sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} \right)}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}$$

↓ 3042

$$-\frac{if \left(-\frac{if \left(\cos(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx - \sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} - \frac{f \left(\sin(2e-\frac{2cf}{d}) \int \frac{\sin(2xf+\frac{2cf}{d}+\frac{\pi}{2})}{c+dx} dx + \cos(2e-\frac{2cf}{d}) \int \frac{\cos(2xf+\frac{2cf}{d})}{c+dx} dx \right)}{ad} \right)}{d} - \frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}$$

↓ 3780

$$\begin{aligned}
 & - \frac{i f \left(-\frac{i f \left(\cos \left(2e - \frac{2 c f}{d} \right) \int \frac{\sin \left(2x f + \frac{2 c f}{d} + \frac{\pi}{2} \right)}{c+dx} dx - \frac{\sin \left(2e - \frac{2 c f}{d} \right) \text{Si} \left(2x f + \frac{2 c f}{d} \right)}{d} \right)}{ad} - \frac{f \left(\sin \left(2e - \frac{2 c f}{d} \right) \int \frac{\sin \left(2x f + \frac{2 c f}{d} + \frac{\pi}{2} \right)}{c+dx} dx + \frac{\cos \left(2e - \frac{2 c f}{d} \right) \text{Si} \left(2x f + \frac{2 c f}{d} \right)}{d} \right)}{ad} \right)}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} \\
 & \qquad \qquad \qquad \downarrow \text{3783} \\
 & - \frac{i f \left(-\frac{f \left(\frac{\text{CosIntegral} \left(2x f + \frac{2 c f}{d} \right) \sin \left(2e - \frac{2 c f}{d} \right)}{d} + \frac{\cos \left(2e - \frac{2 c f}{d} \right) \text{Si} \left(2x f + \frac{2 c f}{d} \right)}{d} \right)}{ad} - \frac{i f \left(\frac{\text{CosIntegral} \left(2x f + \frac{2 c f}{d} \right) \cos \left(2e - \frac{2 c f}{d} \right)}{d} - \frac{\sin \left(2e - \frac{2 c f}{d} \right) \text{Si} \left(2x f + \frac{2 c f}{d} \right)}{d} \right)}{ad} \right)}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))}
 \end{aligned}$$

input `Int[1/((c + d*x)^3*(a + I*a*Tan[e + f*x])),x]`

output
$$\begin{aligned}
 & \frac{((-1/2*I)*f)/(a*d^2*(c + d*x)) - 1/(2*d*(c + d*x)^2*(a + I*a*Tan[e + f*x]))}{d} \\
 & - (I*f*(-((f*((\text{CosIntegral}[(2*c*f)/d + 2*f*x])*Sin[2*e - (2*c*f)/d])/d + (\text{Cos}[2*e - (2*c*f)/d]*\text{SinIntegral}[(2*c*f)/d + 2*f*x])/d))/(a*d)) - (I*f*((\text{Cos}[2*e - (2*c*f)/d]*\text{CosIntegral}[(2*c*f)/d + 2*f*x])/d - (\text{Sin}[2*e - (2*c*f)/d]*\text{SinIntegral}[(2*c*f)/d + 2*f*x])/d))/(a*d) - 1/(d*(c + d*x)*(a + I*a*Tan[e + f*x]))))/d
 \end{aligned}$$

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3780 `Int[sin[(e_.) + (f_)*(x_)]/((c_.) + (d_)*(x_)), x_Symbol] :> Simplify[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

rule 3783 $\text{Int}[\sin[(e_{_}) + (f_{_})*(x_{_})]/((c_{_}) + (d_{_})*(x_{_})), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{CosIntegral}[e - \text{Pi}/2 + f*x]/d, x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{EqQ}[d*(e - \text{Pi}/2) - c*f, 0]$

rule 3784 $\text{Int}[\sin[(e_{_}) + (f_{_})*(x_{_})]/((c_{_}) + (d_{_})*(x_{_})), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{Cos}[(d*e - c*f)/d] \quad \text{Int}[\text{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \text{Simp}[\text{Sin}[(d*e - c*f)/d] \quad \text{Int}[\text{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{NeQ}[d*e - c*f, 0]$

rule 4207 $\text{Int}[1/(((c_{_}) + (d_{_})*(x_{_}))^{2*}(a_{_}) + (b_{_})*\tan[(e_{_}) + (f_{_})*(x_{_})])), x_{\text{Symbol}}] \Rightarrow -\text{Simp}[(d*(c + d*x)*(a + b*\text{Tan}[e + f*x]))^{(-1)}, x] + (-\text{Simp}[f/(a*d) \quad \text{Int}[\text{Sin}[2*e + 2*f*x]/(c + d*x), x], x] + \text{Simp}[f/(b*d) \quad \text{Int}[\text{Cos}[2*e + 2*f*x]/(c + d*x), x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0]$

rule 4208 $\text{Int}[((c_{_}) + (d_{_})*(x_{_}))^{(m_{_})}/((a_{_}) + (b_{_})*\tan[(e_{_}) + (f_{_})*(x_{_})])), x_{\text{Symbol}}] \Rightarrow \text{Simp}[f*((c + d*x)^{(m + 2)}/(b*d^{2*}(m + 1)*(m + 2))), x] + (\text{Simp}[2*b*(f/(a*d*(m + 1))) \quad \text{Int}[(c + d*x)^{(m + 1)}/(a + b*\text{Tan}[e + f*x]), x], x] + \text{Simp}[(c + d*x)^{(m + 1)}/(d*(m + 1)*(a + b*\text{Tan}[e + f*x])), x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{LtQ}[m, -1] \&& \text{NeQ}[m, -2]$

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.95

method	result
risch	$-\frac{1}{4ad(dx+c)^2} + \frac{if^3e^{-2i(fx+e)}x}{2ad(d^2x^2f^2+2cd\,f^2x+c^2f^2)} - \frac{f^2e^{-2i(fx+e)}}{4ad(d^2x^2f^2+2cd\,f^2x+c^2f^2)} + \frac{if^3e^{-2i(fx+e)}c}{2ad^2(d^2x^2f^2+2cd\,f^2x+c^2f^2)} + \frac{f^2e^{\frac{2i(cf+dx)}{d}}}{4ad^3(d^2x^2f^2+2cd\,f^2x+c^2f^2)}$

input $\text{int}(1/(d*x+c)^3/(a+I*a*tan(f*x+e)), x, \text{method}=\text{RETURNVERBOSE})$

output

$$\begin{aligned} & -\frac{1}{4} \frac{a/d}{(d*x+c)^2 + 1/2*I/a*f^3 \exp(-2*I*(f*x+e))} / d / (d^2*f^2*x^2 + 2*c*d*f^2*x^2 + c^2*f^2*x^2) \\ & - \frac{1}{4} \frac{a*f^2 \exp(-2*I*(f*x+e))}{d / (d^2*f^2*x^2 + 2*c*d*f^2*x^2 + c^2*f^2*x^2)} \\ & + \frac{1}{2*I/a*f^3 \exp(-2*I*(f*x+e))} / d^2 / (d^2*f^2*x^2 + 2*c*d*f^2*x^2 + c^2*f^2*x^2) * c + 1 \\ & / a*f^2 / d^3 * \exp(2*I*(c*f-d*e)) / d * \text{Ei}(1, 2*I*f*x + 2*I*e + 2*I*(c*f-d*e)) / d \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 130, normalized size of antiderivative = 0.57

$$\begin{aligned} & \int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ & = \frac{\left(2i d^2 f x + 2i c d f - d^2 - \left(4(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2) \text{Ei}\left(-\frac{2(i d f x + i c f)}{d}\right) e^{\left(-\frac{2(i d e - i c f)}{d}\right)} + d^2 \right) e^{(2i f x + 2i e)} \right)}{4(ad^5 x^2 + 2acd^4 x + ac^2 d^3)} \end{aligned}$$

input

```
integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="fricas")
```

output

$$\begin{aligned} & \frac{1}{4} \left(2*I*d^2*f*x + 2*I*c*d*f - d^2 - (4*(d^2*f^2*x^2 + 2*c*d*f^2*x + c^2*f^2)*\text{Ei}(-2*(I*d*f*x + I*c*f)/d)*e^{-2*(I*d*e - I*c*f)/d} + d^2)*e^{(2*I*f*x + 2*I*e)}*e^{-2*I*f*x - 2*I*e} \right) / (a*d^5*x^2 + 2*a*c*d^4*x + a*c^2*d^3) \end{aligned}$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ & = -\frac{i \int \frac{1}{c^3 \tan(e+fx) - ic^3 + 3c^2 dx \tan(e+fx) - 3ic^2 dx + 3cd^2 x^2 \tan(e+fx) - 3icd^2 x^2 + d^3 x^3 \tan(e+fx) - id^3 x^3} dx}{a} \end{aligned}$$

input

```
integrate(1/(d*x+c)**3/(a+I*a*tan(f*x+e)),x)
```

output

$$\begin{aligned} & -I*\text{Integral}(1/(c**3*tan(e + f*x) - I*c**3 + 3*c**2*d*x*tan(e + f*x) - 3*I*c**2*d*x + 3*c*d**2*x**2*tan(e + f*x) - 3*I*c*d**2*x**2 + d**3*x**3*tan(e + f*x) - I*d**3*x**3), x) / a \end{aligned}$$

Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.70

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx =$$

$$-\frac{2 f^3 \cos\left(-\frac{2(de-cf)}{d}\right) E_3\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + 2i f^3 E_3\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) + }{4 \left((fx+e)^2 ad^3 + ad^3 e^2 - 2acd^2 ef + ac^2 df^2 - 2(ad^3 e - acd^2 f)(fx+e)\right) f}$$

input `integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `-1/4*(2*f^3*cos(-2*(d*e - c*f)/d)*exp_integral_e(3, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f^3*exp_integral_e(3, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + f^3)/(((f*x + e)^2*a*d^3 + a*d^3*e^2 - 2*a*c*d^2*e*f + a*c^2*d*f^2 - 2*(a*d^3*e - a*c*d^2*f)*(f*x + e))*f)`

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 532 vs. $2(212) = 424$.

Time = 0.24 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.34

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx = \text{Too large to display}$$

input `integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output

$$\begin{aligned}
 & -\frac{1}{4} \cdot (4 \cdot d^2 \cdot f^2 \cdot x^2 \cdot \cos(2 \cdot c \cdot f / d) \cdot \cos_{\text{integral}}(-2 \cdot (d \cdot f \cdot x + c \cdot f) / d) + 4 \cdot I \cdot d^2 \\
 & \quad \cdot 2 \cdot f^2 \cdot x^2 \cdot \cos_{\text{integral}}(-2 \cdot (d \cdot f \cdot x + c \cdot f) / d) \cdot \sin(2 \cdot c \cdot f / d) - 4 \cdot I \cdot d^2 \cdot 2 \cdot f^2 \cdot x^2 \\
 & \quad \cdot \cos(2 \cdot c \cdot f / d) \cdot \sin_{\text{integral}}(2 \cdot (d \cdot f \cdot x + c \cdot f) / d) + 4 \cdot d^2 \cdot 2 \cdot f^2 \cdot x^2 \cdot \sin(2 \cdot c \cdot f / d) \\
 & \quad \cdot \sin_{\text{integral}}(2 \cdot (d \cdot f \cdot x + c \cdot f) / d) + 8 \cdot c \cdot d \cdot f^2 \cdot 2 \cdot x \cdot \cos(2 \cdot c \cdot f / d) \cdot \cos_{\text{integral}}(-2 \\
 & \quad \cdot (d \cdot f \cdot x + c \cdot f) / d) + 8 \cdot I \cdot c \cdot d \cdot f^2 \cdot 2 \cdot x \cdot \cos_{\text{integral}}(-2 \cdot (d \cdot f \cdot x + c \cdot f) / d) \cdot \sin(2 \cdot c \\
 & \quad \cdot f / d) - 8 \cdot I \cdot c \cdot d \cdot f^2 \cdot 2 \cdot x \cdot \cos(2 \cdot c \cdot f / d) \cdot \sin_{\text{integral}}(2 \cdot (d \cdot f \cdot x + c \cdot f) / d) + 8 \cdot c \cdot d \\
 & \quad \cdot f^2 \cdot 2 \cdot x \cdot \sin(2 \cdot c \cdot f / d) \cdot \sin_{\text{integral}}(2 \cdot (d \cdot f \cdot x + c \cdot f) / d) + 4 \cdot c^2 \cdot 2 \cdot f^2 \cdot 2 \cdot \cos(2 \cdot c \cdot f / \\
 & \quad d) \cdot \cos_{\text{integral}}(-2 \cdot (d \cdot f \cdot x + c \cdot f) / d) + 4 \cdot I \cdot c^2 \cdot 2 \cdot f^2 \cdot 2 \cdot \cos_{\text{integral}}(-2 \cdot (d \cdot f \cdot x + \\
 & \quad c \cdot f) / d) \cdot \sin(2 \cdot c \cdot f / d) - 4 \cdot I \cdot c^2 \cdot 2 \cdot f^2 \cdot 2 \cdot \cos(2 \cdot c \cdot f / d) \cdot \sin_{\text{integral}}(2 \cdot (d \cdot f \cdot x + c \\
 & \quad \cdot f) / d) + 4 \cdot c^2 \cdot 2 \cdot f^2 \cdot 2 \cdot \sin(2 \cdot c \cdot f / d) \cdot \sin_{\text{integral}}(2 \cdot (d \cdot f \cdot x + c \cdot f) / d) - 2 \cdot I \cdot d^2 \cdot 2 \\
 & \quad \cdot f \cdot x \cdot \cos(2 \cdot f \cdot x) - 2 \cdot d^2 \cdot 2 \cdot f \cdot x \cdot \sin(2 \cdot f \cdot x) - 2 \cdot I \cdot c \cdot d \cdot f \cdot \cos(2 \cdot f \cdot x) - 2 \cdot c \cdot d \cdot f \cdot \sin \\
 & \quad (2 \cdot f \cdot x) + d^2 \cdot 2 \cdot \cos(2 \cdot f \cdot x) + d^2 \cdot 2 \cdot \cos(2 \cdot e) - I \cdot d^2 \cdot 2 \cdot \sin(2 \cdot f \cdot x) + I \cdot d^2 \cdot 2 \cdot \sin(2 \cdot e) \\
 & \quad) / (a \cdot d^5 \cdot x^2 \cdot \cos(2 \cdot e) + I \cdot a \cdot d^5 \cdot x^2 \cdot \sin(2 \cdot e) + 2 \cdot a \cdot c \cdot d^4 \cdot x \cdot \cos(2 \cdot e) + 2 \cdot I \\
 & \quad \cdot a \cdot c \cdot d^4 \cdot x \cdot \sin(2 \cdot e) + a \cdot c^2 \cdot d^3 \cdot \cos(2 \cdot e) + I \cdot a \cdot c^2 \cdot d^3 \cdot \sin(2 \cdot e))
 \end{aligned}$$
Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx = \int \frac{1}{(a + a \tan(e + fx) 1i) (c + dx)^3} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^3),x)`output `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^3), x)`**Reduce [F]**

$$\begin{aligned}
 & \int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\
 & = \frac{\int \frac{1}{\tan(fx+e)c^3i+3\tan(fx+e)c^2dix+3\tan(fx+e)cd^2i x^2+\tan(fx+e)d^3i x^3+c^3+3c^2dx+3c d^2x^2+d^3x^3} dx}{a}
 \end{aligned}$$

input `int(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x)`

output $\int \frac{1}{(\tan(e + fx)*c^3*i + 3*\tan(e + fx)*c^2*d*i*x + 3*\tan(e + fx)*c*d^2*i*x^2 + \tan(e + fx)*d^3*i*x^3 + c^3 + 3*c^2*d*x + 3*c*d^2*x^2 + d^3*x^3)} dx/a$

3.24 $\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx$

Optimal result	211
Mathematica [A] (verified)	212
Rubi [A] (verified)	212
Maple [A] (verified)	214
Fricas [A] (verification not implemented)	214
Sympy [A] (verification not implemented)	215
Maxima [F(-2)]	216
Giac [A] (verification not implemented)	216
Mupad [B] (verification not implemented)	217
Reduce [F]	217

Optimal result

Integrand size = 23, antiderivative size = 270

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx = & -\frac{3d^3e^{-2ie-2ifx}}{16a^2f^4} - \frac{3d^3e^{-4ie-4ifx}}{512a^2f^4} - \frac{3id^2e^{-2ie-2ifx}(c+dx)}{8a^2f^3} \\ & - \frac{3id^2e^{-4ie-4ifx}(c+dx)}{128a^2f^3} + \frac{3de^{-2ie-2ifx}(c+dx)^2}{8a^2f^2} \\ & + \frac{3de^{-4ie-4ifx}(c+dx)^2}{64a^2f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2f} \\ & + \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2f} + \frac{(c+dx)^4}{16a^2d} \end{aligned}$$

output

```
-3/16*d^3*exp(-2*I*e-2*I*f*x)/a^2/f^4-3/512*d^3*exp(-4*I*e-4*I*f*x)/a^2/f^4-3/8*I*d^2*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^2/f^3-3/128*I*d^2*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^2/f^3+3/8*d*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^2/f^2+3/64*d*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^2/f^2+1/4*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^3/a^2/f+1/16*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^3/a^2/f+1/16*(d*x+c)^4/a^2/d
```

Mathematica [A] (verified)

Time = 2.28 (sec) , antiderivative size = 473, normalized size of antiderivative = 1.75

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx \\ = \frac{\sec^2(e + fx)(\cos(fx) + i \sin(fx))^2 ((4ic^3f^3 + 6c^2df^2(1 + 2ifx) + 6cd^2f(-i + 2fx + 2if^2x^2) + d^3(-3 - 6ifx^2 + 6f^2x^4)) \cdot (\cos[2*f*x] + ((32*I)*c^3*f^3 + 24*c^2*d*f^2*(1 + (4*I)*f*x) + 12*c*d^2*f*(-I + 4*f*x + (8*I)*f^2*x^2) + d^3*(-3 - (12*I)*f*x + 24*f^2*x^2 + (32*I)*f^3*x^3))*\cos[4*f*x]*(\cos[2*e] - I*\sin[2*e]))/32 + f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cos[2*e] + I*\sin[2*e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(-I + 2*f*x) + 6*c*d^2*f*(-1 - (2*I)*f*x + 2*f^2*x^2) + d^3*(3*I - 6*f*x - (6*I)*f^2*x^2 + 4*f^3*x^3))*\sin[2*f*x] + ((32*c^3*f^3 + 24*c^2*d*f^2*(-I + 4*f*x) + 12*c*d^2*f*(-1 - (4*I)*f*x + 8*f^2*x^2) + d^3*(3*I - 12*f*x - (24*I)*f^2*x^2 + 32*f^3*x^3))*(\cos[2*e] - I*\sin[2*e])*\sin[4*f*x])/32))/((16*f^4*(a + I*a*Tan[e + f*x]))^2)}$$

input `Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x])^2, x]`

output
$$(Sec[e + f*x]^2 * (\Cos[f*x] + I*Sin[f*x])^2 * (((4*I)*c^3*f^3 + 6*c^2*d*f^2*(1 + (2*I)*f*x) + 6*c*d^2*f*(-I + 2*f*x + (2*I)*f^2*x^2) + d^3*(-3 - (6*I)*f*x + 6*f^2*x^2 + (4*I)*f^3*x^3)) * \Cos[2*f*x] + (((32*I)*c^3*f^3 + 24*c^2*d*f^2*(1 + (4*I)*f*x) + 12*c*d^2*f*(-I + 4*f*x + (8*I)*f^2*x^2) + d^3*(-3 - (12*I)*f*x + 24*f^2*x^2 + (32*I)*f^3*x^3)) * \Cos[4*f*x] * (\Cos[2*e] - I*Sin[2*e]))/32 + f^4*x * (4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3) * (\Cos[2*e] + I*Sin[2*e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(-I + 2*f*x) + 6*c*d^2*f*(-1 - (2*I)*f*x + 2*f^2*x^2) + d^3*(3*I - 6*f*x - (6*I)*f^2*x^2 + 4*f^3*x^3)) * \Sin[2*f*x] + ((32*c^3*f^3 + 24*c^2*d*f^2*(-I + 4*f*x) + 12*c*d^2*f*(-1 - (4*I)*f*x + 8*f^2*x^2) + d^3*(3*I - 12*f*x - (24*I)*f^2*x^2 + 32*f^3*x^3)) * (\Cos[2*e] - I*Sin[2*e]) * \Sin[4*f*x])/32)) / ((16*f^4*(a + I*a*Tan[e + f*x]))^2)$$

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx \\ \downarrow \textcolor{blue}{3042} \\ \int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$\begin{aligned}
 & \downarrow 4212 \\
 & \int \left(\frac{(c+dx)^3 e^{-2ie-2ifx}}{2a^2} + \frac{(c+dx)^3 e^{-4ie-4ifx}}{4a^2} + \frac{(c+dx)^3}{4a^2} \right) dx \\
 & \downarrow 2009 \\
 & -\frac{3id^2(c+dx)e^{-2ie-2ifx}}{8a^2f^3} - \frac{3id^2(c+dx)e^{-4ie-4ifx}}{128a^2f^3} + \frac{3d(c+dx)^2e^{-2ie-2ifx}}{8a^2f^2} + \\
 & \frac{3d(c+dx)^2e^{-4ie-4ifx}}{64a^2f^2} + \frac{i(c+dx)^3e^{-2ie-2ifx}}{4a^2f} + \frac{i(c+dx)^3e^{-4ie-4ifx}}{16a^2f} + \frac{(c+dx)^4}{16a^2d} - \\
 & \frac{3d^3e^{-2ie-2ifx}}{16a^2f^4} - \frac{3d^3e^{-4ie-4ifx}}{512a^2f^4}
 \end{aligned}$$

input `Int[(c + d*x)^3/(a + I*a*Tan[e + f*x])^2, x]`

output `(-3*d^3*E^((-2*I)*e - (2*I)*f*x))/(16*a^2*f^4) - (3*d^3*E^((-4*I)*e - (4*I)*f*x))/(512*a^2*f^4) - (((3*I)/8)*d^2*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))/(a^2*f^3) - (((3*I)/128)*d^2*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(a^2*f^3) + (3*d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2)/(8*a^2*f^2) + (3*d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2)/(64*a^2*f^2) + ((I/4)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^3)/(a^2*f) + ((I/16)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^3)/(a^2*f) + (c + d*x)^4/(16*a^2*d)`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_.))^m_*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a)^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`

Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.05

method	result
risch	$\frac{d^3x^4}{16a^2} + \frac{d^2cx^3}{4a^2} + \frac{3dc^2x^2}{8a^2} + \frac{c^3x}{4a^2} + \frac{c^4}{16a^2d} + \frac{i(4d^3x^3f^3+12cd^2f^3x^2-6id^3f^2x^2+12c^2df^3x-12icd^2f^2x+4c^3f^3-6ic^2df^2)}{16a^2f^4}$

input `int((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & \frac{1}{16}a^2d^3x^4 + \frac{1}{4}a^2d^2c^2x^3 + \frac{3}{8}a^2d^2c^2x^2 + \frac{1}{4}a^2c^3x + \frac{1}{16}a^2 \\ & 2/d*c^4 + \frac{1}{16}I*(4*d^3x^3f^3 - 6*I*d^3*f^2*x^2 + 12*c*d^2*f^3*x^2 - 12*I*c*d^2*f^2*x^2 + 12*c^2*d*f^3*x^2 - 6*I*c^2*f^2*x^2 - 6*I*c^2*f^3*x^2 + 4*c^3*f^3*x^2 - 6*d^3*f^3*x^2 + 3*f*x^3 + 3*I*d^3*f^3*x^2 - 6*c*d^2*f^2*x^2) / \\ & a^2/f^4 * \exp(-2*I*(f*x+e)) + \frac{1}{512}I*(32*d^3x^3f^3 - 24*I*d^3*f^2*x^2 + 96*c^2*d*f^3*x^2 - 24*I*c^2*d*f^2*x^2 + 32*c^3*f^3*x^2 - 12*d^3*f^3*x^2 - 48*I*c*d^2*f^2*x^2 + 96*c^2*d*f^3*x^2 - 24*I*c^2*d*f^2*x^2 + 32*c^3*f^3*x^2 - 12*d^3*f^3*x^2 - 12*I*c*d^2*f^2*x^2) / \\ & a^2/f^4 * \exp(-4*I*(f*x+e)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.99

$$\begin{aligned} & \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx \\ & = \frac{(32i d^3 f^3 x^3 + 32i c^3 f^3 + 24 c^2 d f^2 - 12i c d^2 f - 3 d^3 - 24 (-4i c d^2 f^3 - d^3 f^2) x^2 - 12 (-8i c^2 d f^3 - 4 c d^2 f^2) x^3)}{a^2 f^4} \end{aligned}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output
$$\begin{aligned} & \frac{1}{512} (32*I*d^3*f^3*x^3 + 32*I*c^3*f^3 + 24*c^2*d*f^2 - 12*I*c*d^2*f - 3*d^3 - 24*(-4*I*c*d^2*f^3 - d^3*f^2)*x^2 - 12*(-8*I*c^2*d*f^3 - 4*c*d^2*f^2)*x^2 + I*d^3*f*x + 32*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x)*e^{(4*I*f*x + 4*I*e)} - 32*(-4*I*d^3*f^3*x^3 - 4*I*c^3*f^3 - 6*c^2*d*f^2 + 6*I*c*d^2*f^2 + 3*d^3 + 6*(-2*I*c*d^2*f^3 - d^3*f^2)*x^2 + 6*(-2*I*c^2*d*f^3 - 2*c*d^2*f^2 + I*d^3*f)*x)*e^{(2*I*f*x + 2*I*e)}*e^{(-4*I*f*x - 4*I*e)}) / (a^2*f^4) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 665, normalized size of antiderivative = 2.46

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$= \left\{ \begin{array}{l} \frac{((512ia^2c^3f^7e^{2ie} + 1536ia^2c^2df^7xe^{2ie} + 384a^2c^2df^6e^{2ie} + 1536ia^2cd^2f^7x^2e^{2ie} + 768a^2cd^2f^6xe^{2ie} - 192ia^2cd^2f^5e^{2ie} + 512ia^2d^3f^7x^3e^{2ie} + 384a^2d^2f^6e^{2ie})e^{-4ie}}{16a^2} + \frac{x^3 \cdot (2cd^2e^{2ie} + cd^2)e^{-4ie}}{4a^2} + \frac{x^2 \cdot (6c^2de^{2ie} + 3c^2d)e^{-4ie}}{8a^2} + \frac{x(2c^3e^{2ie} + c^3)e^{-4ie}}{4a^2} \\ + \frac{c^3x}{4a^2} + \frac{3c^2dx^2}{8a^2} + \frac{cd^2x^3}{4a^2} + \frac{d^3x^4}{16a^2} \end{array} \right.$$

input `integrate((d*x+c)**3/(a+I*a*tan(f*x+e))**2,x)`

output

```
Piecewise(((512*I*a**2*c**3*f**7*exp(2*I*e) + 1536*I*a**2*c**2*d*f**7*x*x*exp(2*I*e) + 384*a**2*c**2*d*f**6*exp(2*I*e) + 1536*I*a**2*c*d**2*f**7*x**2*x*exp(2*I*e) + 768*a**2*c*d**2*f**6*x*exp(2*I*e) - 192*I*a**2*c*d**2*f**5*x*exp(2*I*e) + 512*I*a**2*d**3*f**7*x**3*exp(2*I*e) + 384*a**2*d**3*f**6*x**2*x*exp(2*I*e) - 192*I*a**2*d**3*f**5*x*exp(2*I*e) - 48*a**2*d**3*f**4*exp(2*I*e))*exp(-4*I*f*x) + (2048*I*a**2*c**3*f**7*exp(4*I*e) + 6144*I*a**2*c**2*d*f**7*x*exp(4*I*e) + 3072*a**2*c**2*d*f**6*exp(4*I*e) + 6144*I*a**2*c*d**2*f**7*x**2*exp(4*I*e) + 6144*a**2*c*d**2*f**6*x*exp(4*I*e) - 3072*I*a**2*c*d**2*f**5*exp(4*I*e) + 2048*I*a**2*d**3*f**7*x**3*exp(4*I*e) + 3072*a**2*d**3*f**6*x**2*exp(4*I*e) - 3072*I*a**2*d**3*f**5*x*exp(4*I*e) - 1536*a**2*d**3*f**4*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(8192*a**4*f**8), Ne(a**4*f**8*exp(6*I*e), 0)), (x**4*(2*d**3*exp(2*I*e) + d**3)*exp(-4*I*e)/(16*a**2) + x**3*(2*c*d**2*exp(2*I*e) + c*d**2)*exp(-4*I*e)/(4*a**2) + x**2*(6*c**2*d*exp(2*I*e) + 3*c**2*d)*exp(-4*I*e)/(8*a**2) + x*(2*c**3*exp(2*I*e) + c**3)*exp(-4*I*e)/(4*a**2), True)) + c**3*x/(4*a**2) + 3*c**2*d*x**2/(8*a**2) + c*d**2*x**3/(4*a**2) + d**3*x**4/(16*a**2)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.36

$$\begin{aligned} & \int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{(32 d^3 f^4 x^4 e^{(4i f x + 4i e)} + 128 c d^2 f^4 x^3 e^{(4i f x + 4i e)} + 192 c^2 d f^4 x^2 e^{(4i f x + 4i e)} + 128 i d^3 f^3 x^3 e^{(2i f x + 2i e)} + 32 i d^3)}{1} \end{aligned}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output
$$\begin{aligned} & \frac{1}{512} (32 d^3 f^4 x^4 e^{(4I f x + 4I e)} + 128 c d^2 f^4 x^3 e^{(4I f x + 4I e)} + 192 c^2 d f^4 x^2 e^{(4I f x + 4I e)} + 384 c I * c d^2 f^4 x e^{(4I f x + 4I e)} + 96 c I * c d^2 f^4 x^2 e^{(4I f x + 4I e)} + 384 c I * c^2 d f^4 x^3 e^{(4I f x + 4I e)} + 96 c I * c^2 d f^4 x^4 e^{(4I f x + 4I e)} + 32 c I * c^3 d^2 f^4 x^5 e^{(4I f x + 4I e)} + 48 c^2 d^2 f^4 x^6 e^{(4I f x + 4I e)} + 192 c^2 d^2 f^4 x^7 e^{(4I f x + 4I e)} - 192 c I d^3 f^4 x^8 e^{(4I f x + 4I e)} + 24 c^2 d^2 f^4 x^9 e^{(4I f x + 4I e)} - 12 c I d^3 f^4 x^{10} e^{(4I f x + 4I e)} - 192 c I c d^2 f^4 x^{11} e^{(4I f x + 4I e)} - 12 c I c d^2 f^4 x^{12} e^{(4I f x + 4I e)} - 96 d^3 f^4 x^{13} e^{(4I f x + 4I e)} - 3 d^3 f^4 x^{14} e^{(4I f x + 4I e)}) / (a^2 f^4) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 9.13 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx = e^{-e 2i - f x 2i} \left(\frac{(4 c^3 f^3 - c^2 d f^2 6i - 6 c d^2 f + d^3 3i) 1i}{16 a^2 f^4} \right.$$

$$+ \frac{d^3 x^3 1i}{4 a^2 f} - \frac{d x (-2 c^2 f^2 + c d f 2i + d^2) 3i}{8 a^2 f^3}$$

$$- \frac{d^2 x^2 (-2 c f + d 1i) 3i}{8 a^2 f^2} \Big)$$

$$+ e^{-e 4i - f x 4i} \left(\frac{(32 c^3 f^3 - c^2 d f^2 24i - 12 c d^2 f + d^3 3i) 1i}{512 a^2 f^4} \right.$$

$$+ \frac{d^3 x^3 1i}{16 a^2 f} - \frac{d x (-8 c^2 f^2 + c d f 4i + d^2) 3i}{128 a^2 f^3}$$

$$- \frac{d^2 x^2 (-4 c f + d 1i) 3i}{64 a^2 f^2} \Big)$$

$$+ \frac{c^3 x}{4 a^2} + \frac{d^3 x^4}{16 a^2} + \frac{3 c^2 d x^2}{8 a^2} + \frac{c d^2 x^3}{4 a^2}$$

input `int((c + d*x)^3/(a + a*tan(e + f*x)*1i)^2,x)`

output `exp(- e*2i - f*x*2i)*(((d^3*3i + 4*c^3*f^3 - c^2*d*f^2*6i - 6*c*d^2*f)*1i)/(16*a^2*f^4) + (d^3*x^3*1i)/(4*a^2*f) - (d*x*(d^2 - 2*c^2*f^2 + c*d*f*2i)*3i)/(8*a^2*f^3) - (d^2*x^2*(d*1i - 2*c*f)*3i)/(8*a^2*f^2)) + exp(- e*4i - f*x*4i)*(((d^3*3i + 32*c^3*f^3 - c^2*d*f^2*24i - 12*c*d^2*f)*1i)/(512*a^2*f^4) + (d^3*x^3*1i)/(16*a^2*f) - (d*x*(d^2 - 8*c^2*f^2 + c*d*f*4i)*3i)/(128*a^2*f^3) - (d^2*x^2*(d*1i - 4*c*f)*3i)/(64*a^2*f^2)) + (c^3*x)/(4*a^2) + (d^3*x^4)/(16*a^2) + (3*c^2*d*x^2)/(8*a^2) + (c*d^2*x^3)/(4*a^2)`

Reduce [F]

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx$$

$$= - \left(\int \frac{x^3}{\tan(fx+e)^2 - 2 \tan(fx+e)i - 1} dx \right) d^3 - 3 \left(\int \frac{x^2}{\tan(fx+e)^2 - 2 \tan(fx+e)i - 1} dx \right) c d^2 - 3 \left(\int \frac{x}{\tan(fx+e)^2 - 2 \tan(fx+e)i - 1} dx \right) d^2$$

input `int((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x)`

output `(- int(x**3/(tan(e + f*x)**2 - 2*tan(e + f*x)*i - 1),x)*d**3 - 3*int(x**2/(tan(e + f*x)**2 - 2*tan(e + f*x)*i - 1),x)*c*d**2 - 3*int(x/(tan(e + f*x)**2 - 2*tan(e + f*x)*i - 1),x)*c**2*d - int(1/(tan(e + f*x)**2 - 2*tan(e + f*x)*i - 1),x)*c**3)/a**2`

3.25 $\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx$

Optimal result	219
Mathematica [A] (verified)	220
Rubi [A] (verified)	220
Maple [A] (verified)	222
Fricas [A] (verification not implemented)	222
Sympy [A] (verification not implemented)	223
Maxima [F(-2)]	223
Giac [A] (verification not implemented)	224
Mupad [B] (verification not implemented)	224
Reduce [F]	225

Optimal result

Integrand size = 23, antiderivative size = 202

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx = & -\frac{id^2 e^{-2ie-2ifx}}{8a^2 f^3} - \frac{id^2 e^{-4ie-4ifx}}{128a^2 f^3} + \frac{de^{-2ie-2ifx}(c+dx)}{4a^2 f^2} \\ & + \frac{de^{-4ie-4ifx}(c+dx)}{32a^2 f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^2}{4a^2 f} \\ & + \frac{ie^{-4ie-4ifx}(c+dx)^2}{16a^2 f} + \frac{(c+dx)^3}{12a^2 d} \end{aligned}$$

output

```
-1/8*I*d^2*exp(-2*I*e-2*I*f*x)/a^2/f^3-1/128*I*d^2*exp(-4*I*e-4*I*f*x)/a^2
/f^3+1/4*d*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^2/f^2+1/32*d*exp(-4*I*e-4*I*f*x)*
(d*x+c)/a^2/f^2+1/4*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^2/f+1/16*I*exp(-4*I*
e-4*I*f*x)*(d*x+c)^2/a^2/f+1/12*(d*x+c)^3/a^2/d
```

Mathematica [A] (verified)

Time = 1.45 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.40

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ = \frac{\sec^2(e + fx)(\cos(fx) + i \sin(fx))^2 ((d + (1 + i)cf + (1 + i)dfx)((1 + i)cf + d(-i + (1 + i)fx)) \cos(2f)}$$

input `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x])^2, x]`

output
$$\begin{aligned} & (\text{Sec}[e + f*x]^2 * (\text{Cos}[f*x] + I*\text{Sin}[f*x])^2 * ((d + (1 + I)*c*f + (1 + I)*d*f*x) * ((1 + I)*c*f + d*(-I + (1 + I)*f*x)) * \text{Cos}[2*f*x] + ((d + (2 + 2*I)*c*f + (2 + 2*I)*d*f*x) * ((2 + 2*I)*c*f + d*(-I + (2 + 2*I)*f*x)) * \text{Cos}[4*f*x] * (\text{Cos}[2*e] - I*\text{Sin}[2*e]))/16 + (2*f^3*x*(3*c^2 + 3*c*d*x + d^2*x^2) * (\text{Cos}[2*e] + I*\text{Sin}[2*e]))/3 - I*(d + (1 + I)*c*f + (1 + I)*d*f*x) * ((1 + I)*c*f + d*(-I + (1 + I)*f*x)) * \text{Sin}[2*f*x] - (I/16)*(d + (2 + 2*I)*c*f + (2 + 2*I)*d*f*x) * ((2 + 2*I)*c*f + d*(-I + (2 + 2*I)*f*x)) * (\text{Cos}[2*e] - I*\text{Sin}[2*e]) * \text{Sin}[4*f*x]))/(8*f^3*(a + I*a*Tan[e + f*x])^2) \end{aligned}$$

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ & \quad \downarrow \text{4212} \end{aligned}$$

$$\int \left(\frac{(c+dx)^2 e^{-2ie-2ifx}}{2a^2} + \frac{(c+dx)^2 e^{-4ie-4ifx}}{4a^2} + \frac{(c+dx)^2}{4a^2} \right) dx$$

↓ 2009

$$\frac{d(c+dx)e^{-2ie-2ifx}}{4a^2f^2} + \frac{d(c+dx)e^{-4ie-4ifx}}{32a^2f^2} + \frac{i(c+dx)^2e^{-2ie-2ifx}}{4a^2f} + \frac{i(c+dx)^2e^{-4ie-4ifx}}{16a^2f} +$$

$$\frac{(c+dx)^3}{12a^2d} - \frac{id^2e^{-2ie-2ifx}}{8a^2f^3} - \frac{id^2e^{-4ie-4ifx}}{128a^2f^3}$$

input `Int[(c + d*x)^2/(a + I*a*Tan[e + f*x])^2, x]`

output
$$\begin{aligned} & ((-1/8*I)*d^2*E^((-2*I)*e - (2*I)*f*x)/(a^2*f^3) - ((I/128)*d^2*E^((-4*I)*e - (4*I)*f*x))/(a^2*f^3) + (d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))/(4*a^2*f^2) \\ & + (d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(32*a^2*f^2) + ((I/4)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2)/(a^2*f) + ((I/16)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2)/(a^2*f) + (c + d*x)^3/(12*a^2*d)) \end{aligned}$$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simplify[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_.))^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{(-n)}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`

Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.86

method	result
risch	$\frac{d^2x^3}{12a^2} + \frac{dcx^2}{4a^2} + \frac{c^2x}{4a^2} + \frac{c^3}{12a^2d} + \frac{i(2d^2x^2f^2+4cd^2x-2id^2fx+2c^2f^2-2icdf-d^2)e^{-2i(fx+e)}}{8a^2f^3} + \frac{i(8d^2x^2f^2+16cd^2f^2x-4id^2fx-16c^2f^2)e^{-2i(fx+e)}}{16a^2f^5}$

input `int((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/12/a^2*d^2*x^3+1/4/a^2*d*c*x^2+1/4/a^2*c^2*x+1/12/a^2/d*c^3+1/8*I*(2*d^2 \\ & *x^2*f^2-2*I*d^2*f*x+4*c*d*f^2*x-2*I*c*d*f+2*c^2*f^2-d^2)/a^2/f^3*exp(-2*I \\ & *(f*x+e))+1/128*I*(8*d^2*x^2*f^2-4*I*d^2*f*x+16*c*d*f^2*x-4*I*c*d*f+8*c^2*f^2-d^2)/a^2/f^3*exp(-4*I*(f*x+e)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx \\ & = \frac{(24i d^2 f^2 x^2 + 24i c^2 f^2 + 12 c d f - 3i d^2 - 12 (-4i c d f^2 - d^2 f)x + 32 (d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x)e^{(4i f x + 2i c d f^2 - d^2 f)})}{384 a^2 f^3} \end{aligned}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output
$$\begin{aligned} & 1/384*(24*I*d^2*f^2*x^2 + 24*I*c^2*f^2 + 12*c*d*f - 3*I*d^2 - 12*(-4*I*c*d*f^2 - d^2*f)*x + 32*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x)*e^{(4*I*f*x + 4*I*c*d*f^2 - d^2*f)} - 48*(-2*I*d^2*f^2*x^2 - 2*I*c^2*f^2*x^2 - 2*c*d*f + I*d^2 + 2*(-2*I*c*d*f^2 - d^2*f)*x)*e^{(2*I*f*x + 2*I*c*d*f^2 - d^2*f)})*e^{(-4*I*f*x - 4*I*c*d*f^2 - d^2*f)}/(a^2*f^3) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 418, normalized size of antiderivative = 2.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{((64ia^2c^2f^5e^{2ie} + 128ia^2cdf^5xe^{2ie} + 32a^2cdf^4e^{2ie} + 64ia^2d^2f^5x^2e^{2ie} + 32a^2d^2f^4xe^{2ie} - 8ia^2d^2f^3e^{2ie})e^{-4ifx} + (256ia^2c^2f^5e^{4ie} + 512ia^2cdf^5e^{4ie}))e^{-4ie}}{1024a^4f^6} \\ \frac{x^3 \cdot (2d^2e^{2ie} + d^2)e^{-4ie}}{12a^2} + \frac{x^2 \cdot (2cde^{2ie} + cd)e^{-4ie}}{4a^2} + \frac{x(2c^2e^{2ie} + c^2)e^{-4ie}}{4a^2} \\ + \frac{c^2x}{4a^2} + \frac{cdx^2}{4a^2} + \frac{d^2x^3}{12a^2} \end{cases}$$

input `integrate((d*x+c)**2/(a+I*a*tan(f*x+e))**2, x)`

output `Piecewise(((64*I*a**2*c**2*f**5*exp(2*I*e) + 128*I*a**2*c*d*f**5*x*exp(2*I*e) + 32*a**2*c*d*f**4*exp(2*I*e) + 64*I*a**2*d**2*f**5*x**2*exp(2*I*e) + 32*a**2*d**2*f**4*x*exp(2*I*e) - 8*I*a**2*d**2*f**3*exp(2*I*e))*exp(-4*I*f*x) + (256*I*a**2*c**2*f**5*exp(4*I*e) + 512*I*a**2*c*d*f**5*x*exp(4*I*e) + 256*a**2*c*d*f**4*exp(4*I*e) + 256*I*a**2*d**2*f**5*x**2*exp(4*I*e) + 56*a**2*d**2*f**4*x*exp(4*I*e) - 128*I*a**2*d**2*f**3*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(1024*a**4*f**6), Ne(a**4*f**6*exp(6*I*e), 0)), (x**3*(2*d**2*exp(2*I*e) + d**2)*exp(-4*I*e)/(12*a**2) + x**2*(2*c*d*exp(2*I*e) + c*d)*exp(-4*I*e)/(4*a**2) + x*(2*c**2*exp(2*I*e) + c**2)*exp(-4*I*e)/(4*a**2), True)) + c**2*x/(4*a**2) + c*d*x**2/(4*a**2) + d**2*x**3/(12*a**2)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2, x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(32 d^2 f^3 x^3 e^{(4i f x + 4i e)} + 96 c d f^3 x^2 e^{(4i f x + 4i e)} + 96 c^2 f^3 x e^{(4i f x + 4i e)} + 96 i d^2 f^2 x^2 e^{(2i f x + 2i e)} + 24 i d^2 f^2 x^2 +$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output $\frac{1}{384} \cdot (32 d^2 f^3 x^3 e^{(4 I f x + 4 I e)} + 96 c d f^3 x^2 e^{(4 I f x + 4 I e)} + 96 c^2 f^3 x e^{(4 I f x + 4 I e)} + 96 I d^2 f^2 x^2 e^{(2 I f x + 2 I e)} + 192 I c d f^2 x e^{(2 I f x + 2 I e)} + 48 I c d f^2 x^2 + 96 I c^2 f^2 x^2 e^{(2 I f x + 2 I e)} + 96 d^2 f^2 x^2 e^{(2 I f x + 2 I e)} + 24 I c^2 f^2 x^2 + 12 d^2 f^2 x + 96 c d f e^{(2 I f x + 2 I e)} + 12 c d f - 48 I d^2 e^{(2 I f x + 2 I e)} - 3 I d^2 e^{(-4 I f x - 4 I e)}) / (a^2 f^3)$

Mupad [B] (verification not implemented)

Time = 8.26 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.91

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx = \frac{c^2 x}{4 a^2} - e^{-e 4i - f x 4i} \left(\frac{(-8 c^2 f^2 + c d f 4i + d^2) 1i}{128 a^2 f^3} - \frac{d^2 x^2 1i}{16 a^2 f} + \frac{d x (-4 c f + d 1i) 1i}{32 a^2 f^2} \right) \\ - e^{-e 2i - f x 2i} \left(\frac{(-2 c^2 f^2 + c d f 2i + d^2) 1i}{8 a^2 f^3} - \frac{d^2 x^2 1i}{4 a^2 f} + \frac{d x (-2 c f + d 1i) 1i}{4 a^2 f^2} \right) + \frac{d^2 x^3}{12 a^2} + \frac{c d x^2}{4 a^2}$$

input `int((c + d*x)^2/(a + a*tan(e + f*x)*1i)^2,x)`

output

$$\begin{aligned} & \frac{(c^2*x)/(4*a^2) - \exp(-e*4i - f*x*4i)*(((d^2 - 8*c^2*f^2 + c*d*f*4i)*1i)/(128*a^2*f^3) - (d^2*x^2*1i)/(16*a^2*f) + (d*x*(d*1i - 4*c*f)*1i)/(32*a^2*f^2)) - \exp(-e*2i - f*x*2i)*(((d^2 - 2*c^2*f^2 + c*d*f*2i)*1i)/(8*a^2*f^3) - (d^2*x^2*1i)/(4*a^2*f) + (d*x*(d*1i - 2*c*f)*1i)/(4*a^2*f^2)) + (d^2*x^3)/(12*a^2) + (c*d*x^2)/(4*a^2)}{ } \end{aligned}$$

Reduce [F]

$$\begin{aligned} & \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{-\left(\int \frac{x^2}{\tan(fx+e)^2 - 2\tan(fx+e)i-1} dx\right) d^2 - 2\left(\int \frac{x}{\tan(fx+e)^2 - 2\tan(fx+e)i-1} dx\right) cd - \left(\int \frac{1}{\tan(fx+e)^2 - 2\tan(fx+e)i-1} dx\right)}{a^2} \end{aligned}$$

input

```
int((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x)
```

output

$$(-\text{int}(x^{**2}/(\tan(e + f*x)**2 - 2*\tan(e + f*x)*i - 1),x)*d^{**2} - 2*\text{int}(x/(\tan(e + f*x)**2 - 2*\tan(e + f*x)*i - 1),x)*c*d - \text{int}(1/(\tan(e + f*x)**2 - 2*\tan(e + f*x)*i - 1),x)*c^{**2})/a^{**2}$$

3.26 $\int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx$

Optimal result	226
Mathematica [A] (verified)	226
Rubi [A] (verified)	227
Maple [A] (verified)	228
Fricas [A] (verification not implemented)	229
Sympy [A] (verification not implemented)	229
Maxima [F(-2)]	230
Giac [A] (verification not implemented)	230
Mupad [B] (verification not implemented)	230
Reduce [F]	231

Optimal result

Integrand size = 21, antiderivative size = 151

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx = & -\frac{3idx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} + \frac{d}{16f^2(a+ia\tan(e+fx))^2} \\ & + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} + \frac{3d}{16f^2(a^2+ia^2\tan(e+fx))} \\ & + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} \end{aligned}$$

output

```
-3/16*I*d*x/a^2/f-1/8*d*x^2/a^2+1/4*x*(d*x+c)/a^2+1/16*d/f^2/(a+I*a*tan(f*x+e))^2+1/4*I*(d*x+c)/f/(a+I*a*tan(f*x+e))^2+3/16*d/f^2/(a^2+I*a^2*tan(f*x+e))+1/4*I*(d*x+c)/f/(a^2+I*a^2*tan(f*x+e))
```

Mathematica [A] (verified)

Time = 0.96 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.86

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx = & \frac{\sec^2(e+fx)(8(d+2icf+2idfx)+(4cf(i+4fx)+d(1+4ifx+8f^2x^2))\cos(2(e+fx))+(4cf(1+4ifx+8f^2x^2)\sin(2(e+fx)))}{64a^2f^2(-i+\tan(e+fx))^2} \end{aligned}$$

input $\text{Integrate}[(c + d*x)/(a + I*a*Tan[e + f*x])^2, x]$

output
$$\frac{-1/64 * (\text{Sec}[e + f*x])^2 * (8*(d + (2*I)*c*f + (2*I)*d*f*x) + (4*c*f*(I + 4*f*x) + d*(1 + (4*I)*f*x + 8*f^2*x^2))*\text{Cos}[2*(e + f*x)] + (4*c*f*(1 + (4*I)*f*x) + d*(-I + 4*f*x + (8*I)*f^2*x^2))*\text{Sin}[2*(e + f*x)]))}{(a^2*f^2*(-I + \text{Tan}[e + f*x]))^2}$$

Rubi [A] (verified)

Time = 0.39 (sec), antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4213, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4213} \\
 & -d \int \left(\frac{x}{4a^2} + \frac{i}{4f(i \tan(e + fx)a^2 + a^2)} + \frac{i}{4f(i \tan(e + fx)a + a)^2} \right) dx + \\
 & \quad \frac{i(c + dx)}{4f(a^2 + ia^2 \tan(e + fx))} + \frac{x(c + dx)}{4a^2} + \frac{i(c + dx)}{4f(a + ia \tan(e + fx))^2} \\
 & \quad \downarrow \text{2009} \\
 & \quad \frac{i(c + dx)}{4f(a^2 + ia^2 \tan(e + fx))} + \frac{x(c + dx)}{4a^2} - \\
 & d \left(-\frac{3}{16f^2(a^2 + ia^2 \tan(e + fx))} + \frac{3ix}{16a^2f} + \frac{x^2}{8a^2} - \frac{1}{16f^2(a + ia \tan(e + fx))^2} \right) + \\
 & \quad \frac{i(c + dx)}{4f(a + ia \tan(e + fx))^2}
 \end{aligned}$$

input $\text{Int}[(c + d*x)/(a + I*a*Tan[e + f*x])^2, x]$

output

$$(x*(c + d*x))/(4*a^2) + ((I/4)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])^2) + ((I/4)*(c + d*x))/(f*(a^2 + I*a^2*Tan[e + f*x])) - d*((3*I)/16)*x/(a^2*f) + x^2/(8*a^2) - 1/(16*f^2*(a + I*a*Tan[e + f*x])^2) - 3/(16*f^2*(a^2 + I*a^2*Tan[e + f*x]))$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4213 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_\text{Symbol}] \rightarrow \text{With}[\{u = \text{IntHide}[(a + b*\tan[e + f*x])^n, x]\}, \text{Simp}[(c + d*x)^m u, x] - \text{Simp}[d*m \text{Int}[(c + d*x)^{m-1} u, x], x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, -1] \&& \text{GtQ}[m, 0]$

Maple [A] (verified)

Time = 0.70 (sec), antiderivative size = 82, normalized size of antiderivative = 0.54

method	result	size
risch	$\frac{dx^2}{8a^2} + \frac{cx}{4a^2} + \frac{i(2dfx+2cf-id)e^{-2i(fx+e)}}{8a^2f^2} + \frac{i(4dfx+4cf-id)e^{-4i(fx+e)}}{64a^2f^2}$	82

input $\text{int}((d*x+c)/(a+I*a*tan(f*x+e))^2, x, \text{method}=\text{_RETURNVERBOSE})$

output $\frac{1}{8}*d*x^2/a^2+1/4*a^2*c*x+1/8*I*(2*d*f*x-I*d+2*c*f)/a^2/f^2*\exp(-2*I*(f*x+e))+1/64*I*(4*d*f*x-I*d+4*c*f)/a^2/f^2*\exp(-4*I*(f*x+e))$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.52

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(4i dfx + 4i cf + 8(df^2 x^2 + 2cf^2 x)e^{(4ifx+4ie)} - 8(-2i dfx - 2i cf - d)e^{(2ifx+2ie)} + d)e^{(-4ifx-4ie)}}{64 a^2 f^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output $\frac{1/64*(4*I*d*f*x + 4*I*c*f + 8*(d*f^2*x^2 + 2*c*f^2*x)*e^{(4*I*f*x + 4*I*e)} - 8*(-2*I*d*f*x - 2*I*c*f - d)*e^{(2*I*f*x + 2*I*e)} + d)*e^{(-4*I*f*x - 4*I*e)})/(a^2*f^2)$

Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.50

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\ = \begin{cases} \frac{((32ia^2cf^3e^{2ie} + 32ia^2df^3xe^{2ie} + 8a^2df^2e^{2ie})e^{-4ifx} + (128ia^2cf^3e^{4ie} + 128ia^2df^3xe^{4ie} + 64a^2df^2e^{4ie})e^{-2ifx})e^{-6ie}}{512a^4f^4} & \text{for } a^4f^4e^{6ie} \neq 0 \\ \frac{x^2 \cdot (2de^{2ie} + d)e^{-4ie}}{8a^2} + \frac{x(2ce^{2ie} + c)e^{-4ie}}{4a^2} \\ + \frac{cx}{4a^2} + \frac{dx^2}{8a^2} & \text{otherwise} \end{cases}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))**2,x)`

output $\text{Piecewise}(((32*I*a**2*c*f**3*exp(2*I*e) + 32*I*a**2*d*f**3*x*exp(2*I*e) + 8*a**2*d*f**2*exp(2*I*e))*exp(-4*I*f*x) + (128*I*a**2*c*f**3*exp(4*I*e) + 128*I*a**2*d*f**3*x*exp(4*I*e) + 64*a**2*d*f**2*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(512*a**4*f**4), \text{Ne}(a**4*f**4*exp(6*I*e), 0)), (x**2*(2*d*exp(2*I*e) + d)*exp(-4*I*e)/(8*a**2) + x*(2*c*exp(2*I*e) + c)*exp(-4*I*e)/(4*a**2), \text{True})) + c*x/(4*a**2) + d*x**2/(8*a**2)$

Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.67

$$\begin{aligned} & \int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{(8 df^2 x^2 e^{(4i f x + 4i e)} + 16 cf^2 x e^{(4i f x + 4i e)} + 16i df x e^{(2i f x + 2i e)} + 4i df x + 16i c f e^{(2i f x + 2i e)} + 4i cf + 8 de^{(2i f x + 2i e)})}{64 a^2 f^2} \end{aligned}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `1/64*(8*d*f^2*x^2*e^(4*I*f*x + 4*I*e) + 16*c*f^2*x*e^(4*I*f*x + 4*I*e) + 16*I*d*f*x*e^(2*I*f*x + 2*I*e) + 4*I*d*f*x + 16*I*c*f*e^(2*I*f*x + 2*I*e) + 4*I*c*f + 8*d*e^(2*I*f*x + 2*I*e) + d)*e^(-4*I*f*x - 4*I*e)/(a^2*f^2)`

Mupad [B] (verification not implemented)

Time = 8.07 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.68

$$\begin{aligned} \int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx &= \frac{dx^2}{8 a^2} - e^{-e 4i - f x 4i} \left(\frac{(-4 c f + d 1i) 1i}{64 a^2 f^2} - \frac{d x 1i}{16 a^2 f} \right) \\ &\quad - e^{-e 2i - f x 2i} \left(\frac{(-2 c f + d 1i) 1i}{8 a^2 f^2} - \frac{d x 1i}{4 a^2 f} \right) + \frac{c x}{4 a^2} \end{aligned}$$

input `int((c + d*x)/(a + a*tan(e + f*x)*1i)^2,x)`

output
$$\begin{aligned} & \frac{(d*x^2)/(8*a^2) - \exp(-e*4i - f*x*4i)*(((d*1i - 4*c*f)*1i)/(64*a^2*f^2) - \\ & (d*x*1i)/(16*a^2*f)) - \exp(-e*2i - f*x*2i)*(((d*1i - 2*c*f)*1i)/(8*a^2*f^2) - \\ & (d*x*1i)/(4*a^2*f)) + (c*x)/(4*a^2)}{a^2} \end{aligned}$$

Reduce [F]

$$\begin{aligned} & \int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{-\left(\int \frac{x}{\tan(fx+e)^2 - 2\tan(fx+e)i-1} dx\right)d - \left(\int \frac{1}{\tan(fx+e)^2 - 2\tan(fx+e)i-1} dx\right)c}{a^2} \end{aligned}$$

input `int((d*x+c)/(a+I*a*tan(f*x+e))^2,x)`

output
$$(-(\text{int}(x/(\tan(e + fx)**2 - 2*tan(e + fx)*i - 1),x)*d + \text{int}(1/(\tan(e + fx)**2 - 2*tan(e + fx)*i - 1),x)*c))/a**2$$

3.27 $\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx$

Optimal result	232
Mathematica [A] (verified)	233
Rubi [A] (verified)	233
Maple [A] (verified)	235
Fricas [A] (verification not implemented)	235
Sympy [F]	236
Maxima [A] (verification not implemented)	236
Giac [A] (verification not implemented)	237
Mupad [F(-1)]	238
Reduce [F]	238

Optimal result

Integrand size = 23, antiderivative size = 305

$$\begin{aligned} \int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx = & \frac{\cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{2a^2d} \\ & + \frac{\cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{4a^2d} \\ & + \frac{\log(c+dx)}{4a^2d} \\ & - \frac{i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{4a^2d} \\ & - \frac{i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2a^2d} \\ & - \frac{i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} \\ & - \frac{\sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} \\ & - \frac{i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d} \\ & - \frac{\sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d} \end{aligned}$$

output

$$\frac{1}{2} \cos(-2e + 2cf/d) \operatorname{Ci}(2cf/d + 2fx)/a^2/d + \frac{1}{4} \cos(-4e + 4cf/d) \operatorname{Ci}(4cf/d + 4fx)/a^2/d + \frac{1}{4} \ln(dx + c)/a^2/d + \frac{1}{4} I \operatorname{Ci}(4cf/d + 4fx) \sin(-4e + 4cf/d)/a^2/d + \frac{1}{2} I \operatorname{Ci}(2cf/d + 2fx) \sin(-2e + 2cf/d)/a^2/d - \frac{1}{2} I \cos(-2e + 2cf/d) \operatorname{Si}(2cf/d + 2fx)/a^2/d + \frac{1}{2} \sin(-2e + 2cf/d) \operatorname{Si}(2cf/d + 2fx)/a^2/d - \frac{1}{4} I \cos(-4e + 4cf/d) \operatorname{Si}(4cf/d + 4fx)/a^2/d + \frac{1}{4} \sin(-4e + 4cf/d) \operatorname{Si}(4cf/d + 4fx)/a^2/d$$

Mathematica [A] (verified)

Time = 0.76 (sec), antiderivative size = 211, normalized size of antiderivative = 0.69

$$\int \frac{1}{(c + dx)(a + i a \tan(e + fx))^2} dx \\ = \frac{\left(\cos\left(2e - \frac{2cf}{d}\right) - i \sin\left(2e - \frac{2cf}{d}\right)\right) \left(2 \operatorname{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) + \cos\left(2e - \frac{2cf}{d}\right) \log(f(c+dx)) + \operatorname{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)\right)}{(c + dx)(a + i a \tan(e + fx))^2}$$

input

```
Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])^2), x]
```

output

$$\frac{((\operatorname{Cos}[2e - (2cf)/d] - I \operatorname{Sin}[2e - (2cf)/d]) * (2 \operatorname{CosIntegral}[(2f(c + d*x))/d] + \operatorname{Cos}[2e - (2cf)/d] * \operatorname{Log}[f*(c + d*x)] + \operatorname{CosIntegral}[(4f(c + d*x))/d] * (\operatorname{Cos}[2e - (2cf)/d] - I \operatorname{Sin}[2e - (2cf)/d]) + I \operatorname{Log}[f*(c + d*x)] * \operatorname{Sin}[2e - (2cf)/d] - (2I) \operatorname{SinIntegral}[(2f(c + d*x))/d] - I \operatorname{Cos}[2e - (2cf)/d] * \operatorname{SinIntegral}[(4f(c + d*x))/d] - \operatorname{Sin}[2e - (2cf)/d] * \operatorname{SinIntegral}[(4f(c + d*x))/d])) / (4a^2d)}$$

Rubi [A] (verified)

Time = 1.06 (sec), antiderivative size = 305, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx \\
 & \quad \downarrow \textcolor{blue}{4211} \\
 & \int \left(-\frac{\sin^2(2e+2fx)}{4a^2(c+dx)} - \frac{i\sin(2e+2fx)}{2a^2(c+dx)} - \frac{i\sin(4e+4fx)}{4a^2(c+dx)} + \frac{\cos^2(2e+2fx)}{4a^2(c+dx)} + \frac{\cos(2e+2fx)}{2a^2(c+dx)} + \frac{1}{4a^2(c+dx)} \right) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{i \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{2a^2d} - \frac{i \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{4a^2d} + \\
 & \frac{\operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{2a^2d} + \frac{\operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{4a^2d} - \\
 & \frac{\sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{2a^2d} - \frac{\sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{4a^2d} - \\
 & \frac{i \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{2a^2d} - \frac{i \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{4a^2d} + \frac{\log(c+dx)}{4a^2d}
 \end{aligned}$$

input `Int[1/((c + d*x)*(a + I*a*Tan[e + f*x])^2),x]`

output

```
(Cos[2*e - (2*c*f)/d]*CosIntegral[(2*c*f)/d + 2*f*x]/(2*a^2*d) + (Cos[4*e - (4*c*f)/d]*CosIntegral[(4*c*f)/d + 4*f*x]/(4*a^2*d) + Log[c + d*x]/(4*a^2*d) - ((I/4)*CosIntegral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*f)/d]/(a^2*d) - ((I/2)*CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d]/(a^2*d) - ((I/2)*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x]/(a^2*d) - (Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x]/(2*a^2*d) - ((I/4)*Cos[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x]/(a^2*d) - (Sin[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x]/(4*a^2*d)
```

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4211 $\text{Int}[(c_.) + (d_.)*(x_.)^m*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + \text{Cos}[2*e + 2*f*x]/(2*a) + \text{Sin}[2*e + 2*f*x]/(2*b))^(-n), x], x] /; \ \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&& \ \text{EqQ}[a^2 + b^2, 0] \ \&& \ \text{ILtQ}[m, 0] \ \&& \ \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.37

method	result	size
risch	$\frac{\ln(dx+c)}{4a^2d} - \frac{e^{\frac{4i(cf-de)}{d}} \expIntegral_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{4a^2d} - \frac{e^{\frac{2i(cf-de)}{d}} \expIntegral_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{2a^2d}$	114

input $\text{int}(1/(d*x+c)/(a+I*a*tan(f*x+e))^2, x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{4}*\ln(d*x+c)/a^2/d-1/4/a^2/d*\exp(4*I*(c*f-d*e)/d)*\text{Ei}(1, 4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-1/2/a^2/d*\exp(2*I*(c*f-d*e)/d)*\text{Ei}(1, 2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.28

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx \\ &= \frac{2 \operatorname{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + \operatorname{Ei}\left(-\frac{4(i dfx + i cf)}{d}\right) e^{\left(-\frac{4(i de - i cf)}{d}\right)} + \log\left(\frac{dx+c}{d}\right)}{4 a^2 d} \end{aligned}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output
$$\frac{1}{4} \left(2 \operatorname{Ei}(-2 \operatorname{atan}(f x + e) / d) e^{-(-2 \operatorname{atan}(f x + e) - I c f) / d} + \operatorname{Ei}(-4 \operatorname{atan}(f x + e) / d) e^{(-4 \operatorname{atan}(f x + e) - I c f) / d} + \log((d x + c) / d) \right) / (a^2 d)$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx \\ &= -\frac{\int \frac{1}{c \tan^2(e+fx)-2ictan(e+fx)-c+dx\tan^2(e+fx)-2idx\tan(e+fx)-dx} dx}{a^2} \end{aligned}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**2,x)`

output
$$-\operatorname{Integral}\left(\frac{1}{c \tan^2(e+fx)-2ictan(e+fx)-c+dx\tan^2(e+fx)-2idx\tan(e+fx)-dx}, x\right)/a^2$$

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.64

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx = \\ & -\frac{2 f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + f \cos\left(-\frac{4(de-cf)}{d}\right) E_1\left(-\frac{4(-i(fx+e)d+i(de-icf))}{d}\right) + 2i f}{d} \end{aligned}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output

$$\begin{aligned} & -\frac{1}{4}(2f \cos(-2(d \cdot e - c \cdot f)/d) \operatorname{exp_integral_e}(1, -2(-I \cdot (f \cdot x + e)) \cdot d + I \cdot d \\ & *e - I \cdot c \cdot f)/d) + f \cos(-4(d \cdot e - c \cdot f)/d) \operatorname{exp_integral_e}(1, -4(-I \cdot (f \cdot x + e) \\ &) \cdot d + I \cdot d \cdot e - I \cdot c \cdot f)/d) + 2I \cdot f \cdot \operatorname{exp_integral_e}(1, -2(-I \cdot (f \cdot x + e)) \cdot d + I \cdot d \\ & *e - I \cdot c \cdot f)/d) * \sin(-2(d \cdot e - c \cdot f)/d) + I \cdot f \cdot \operatorname{exp_integral_e}(1, -4(-I \cdot (f \cdot x + e) \\ &) \cdot d + I \cdot d \cdot e - I \cdot c \cdot f)/d) * \sin(-4(d \cdot e - c \cdot f)/d) - f \cdot \log((f \cdot x + e) \cdot d - d \cdot e \\ & + c \cdot f)/(a^2 \cdot d \cdot f) \end{aligned}$$

Giac [A] (verification not implemented)

Time = 0.27 (sec), antiderivative size = 404, normalized size of antiderivative = 1.32

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx \\ & = \frac{2 \cos(2e) \cos(\frac{2cf}{d}) \operatorname{Ci}\left(-\frac{2(df+cf)}{d}\right) + \cos(2e)^2 \log(dx+c) + 2i \cos(\frac{2cf}{d}) \operatorname{Ci}\left(-\frac{2(df+cf)}{d}\right) \sin(2e) + 2i \sin(2e) \log(dx+c)}{(c + dx)(a + ia \tan(e + fx))^2} \end{aligned}$$

input

```
integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")
```

output

$$\begin{aligned} & \frac{1}{4}(2 \cos(2e) \cos(2c \cdot f/d) \cos(\operatorname{integral}(-2(d \cdot f \cdot x + c \cdot f)/d) + \cos(2e)^2 \\ & * \log(d \cdot x + c) + 2I \cos(2c \cdot f/d) \cos(\operatorname{integral}(-2(d \cdot f \cdot x + c \cdot f)/d) * \sin(2e) \\ & + 2I \cos(2e) * \log(d \cdot x + c) * \sin(2e) - \log(d \cdot x + c) * \sin(2e)^2 + 2I \cos(2e) * \cos(\operatorname{integral}(-2(d \cdot f \cdot x + c \cdot f)/d) * \sin(2c \cdot f/d) - 2 \cos(\operatorname{integral}(-2(d \cdot f \cdot x + c \cdot f)/d) * \sin(2e) * \sin(2c \cdot f/d) - 2I \cos(2e) * \cos(2c \cdot f/d) * \sin(\operatorname{integral}(2(d \cdot f \cdot x + c \cdot f)/d) + 2 \cos(2c \cdot f/d) * \sin(2e) * \sin(\operatorname{integral}(2(d \cdot f \cdot x + c \cdot f)/d) + 2I \cos(2e) * \sin(2c \cdot f/d) * \sin(\operatorname{integral}(2(d \cdot f \cdot x + c \cdot f)/d) + \cos(4c \cdot f/d) * \cos(\operatorname{integral}(-4(d \cdot f \cdot x + c \cdot f)/d) + I \cos(\operatorname{integral}(-4(d \cdot f \cdot x + c \cdot f)/d) * \sin(4c \cdot f/d) - I \cos(4c \cdot f/d) * \sin(\operatorname{integral}(4(d \cdot f \cdot x + c \cdot f)/d) + \sin(4c \cdot f/d) * \sin(\operatorname{integral}(4(d \cdot f \cdot x + c \cdot f)/d)) / (a^2 \cdot d \cdot \cos(2e)^2 + 2I \cdot a^2 \cdot d \cdot \cos(2e) * \sin(2e) - a^2 \cdot d \cdot \sin(2e)^2) \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^2 (c + dx)} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)),x)`

output `int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx \\ &= -\frac{\int \frac{1}{\tan(fx+e)^2 c + \tan(fx+e)^2 dx - 2 \tan(fx+e) ci - 2 \tan(fx+e) di x - c - dx} dx}{a^2} \end{aligned}$$

input `int(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x)`

output `(- int(1/(tan(e + f*x)**2*c + tan(e + f*x)**2*d*x - 2*tan(e + f*x)*c*i - 2*tan(e + f*x)*d*i*x - c - d*x),x))/a**2`

$$\mathbf{3.28} \quad \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx$$

Optimal result	240
Mathematica [A] (verified)	241
Rubi [A] (verified)	241
Maple [A] (verified)	243
Fricas [A] (verification not implemented)	244
Sympy [F]	244
Maxima [A] (verification not implemented)	245
Giac [B] (verification not implemented)	245
Mupad [F(-1)]	246
Reduce [F]	247

Optimal result

Integrand size = 23, antiderivative size = 436

$$\begin{aligned}
 \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx = & -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} \\
 & - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)} \\
 & - \frac{if \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
 & - \frac{if \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
 & - \frac{f \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{a^2d^2} \\
 & - \frac{f \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{a^2d^2} \\
 & + \frac{i \sin(2e + 2fx)}{2a^2d(c+dx)} \\
 & + \frac{\sin^2(2e + 2fx)}{4a^2d(c+dx)} + \frac{i \sin(4e + 4fx)}{4a^2d(c+dx)} \\
 & - \frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
 & + \frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
 & - \frac{f \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
 & + \frac{if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2}
 \end{aligned}$$

output

```

-1/4/a^2/d/(d*x+c)-1/2*cos(2*f*x+2*e)/a^2/d/(d*x+c)-1/4*cos(2*f*x+2*e)^2/a
^2/d/(d*x+c)-I*f*cos(-2*e+2*c*f/d)*Ci(2*c*f/d+2*f*x)/a^2/d^2-I*f*cos(-4*e+
4*c*f/d)*Ci(4*c*f/d+4*f*x)/a^2/d^2+f*Ci(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a
^2/d^2+f*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d^2+1/2*I*sin(2*f*x+2*e)/
a^2/d/(d*x+c)+1/4*sin(2*f*x+2*e)^2/a^2/d/(d*x+c)+1/4*I*sin(4*f*x+4*e)/a^2/
d/(d*x+c)-f*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^2/d^2-I*f*sin(-2*e+2*c*f
/d)*Si(2*c*f/d+2*f*x)/a^2/d^2-f*cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^2/d^2-
I*f*sin(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^2/d^2

```

Mathematica [A] (verified)

Time = 1.55 (sec) , antiderivative size = 467, normalized size of antiderivative = 1.07

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx =$$

$$\frac{-\left(\cos\left(2(e + f(-\frac{c}{d} + x))\right) - i \sin\left(2(e + f(-\frac{c}{d} + x))\right)\right) \left(2d \cos\left(\frac{2cf}{d}\right) + d \cos\left(2(e + f(-\frac{c}{d} + x))\right) + d \sin\left(2(e + f(-\frac{c}{d} + x))\right)\right)}{(c + dx)^2(a + ia \tan(e + fx))^2}$$

input `Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^2), x]`

output

$$\begin{aligned} & -\frac{1}{4} ((\cos[2*(e + f*(-(c/d) + x))] - I*\sin[2*(e + f*(-(c/d) + x))])*(2*d*C \\ & os[(2*c*f)/d] + d*Cos[2*(e + f*(-(c/d) + x))] + d*Cos[2*(e + f*(c/d + x))] \\ & - (2*I)*d*Sin[(2*c*f)/d] + (4*I)*f*(c + d*x)*CosIntegral[(2*f*(c + d*x))/ \\ & d]*(Cos[2*f*x] + I*Sin[2*f*x]) + I*d*Sin[2*(e + f*(-(c/d) + x))] - I*d*Sin \\ & [2*(e + f*(c/d + x))] + 4*f*(c + d*x)*CosIntegral[(4*f*(c + d*x))/d]*(I*Co \\ & s[2*e - (2*f*(c + d*x))/d] + Sin[2*e - (2*f*(c + d*x))/d]) + 4*c*f*Cos[2*f \\ & *x]*SinIntegral[(2*f*(c + d*x))/d] + 4*d*f*x*Cos[2*f*x]*SinIntegral[(2*f*(\\ & c + d*x))/d] + (4*I)*c*f*Sin[2*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (4*I) \\ & *d*f*x*Sin[2*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 4*c*f*Cos[2*e - (2*f*(c \\ & + d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] + 4*d*f*x*Cos[2*e - (2*f*(c + d \\ & *x))/d]*SinIntegral[(4*f*(c + d*x))/d] - (4*I)*c*f*Sin[2*e - (2*f*(c + d*x) \\ &)/d]*SinIntegral[(4*f*(c + d*x))/d] - (4*I)*d*f*x*Sin[2*e - (2*f*(c + d*x) \\ &)/d]*SinIntegral[(4*f*(c + d*x))/d]))/(a^2*d^2*(c + d*x)) \end{aligned}$$

Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx$$

↓ 3042

$$\begin{aligned}
 & \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx \\
 & \quad \downarrow \textcolor{blue}{4211} \\
 & \int \left(-\frac{\sin^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{i\sin(2e+2fx)}{2a^2(c+dx)^2} - \frac{i\sin(4e+4fx)}{4a^2(c+dx)^2} + \frac{\cos^2(2e+2fx)}{4a^2(c+dx)^2} + \frac{\cos(2e+2fx)}{2a^2(c+dx)^2} + \frac{1}{4a^2(c+dx)^2} \right) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{f \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{a^2 d^2} - \frac{f \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{a^2 d^2} - \\
 & -\frac{if \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{a^2 d^2} - \frac{if \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{a^2 d^2} + \\
 & +\frac{if \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{a^2 d^2} + \frac{if \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{a^2 d^2} - \\
 & -\frac{f \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{a^2 d^2} - \frac{f \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{a^2 d^2} + \frac{\sin^2(2e+2fx)}{4a^2 d(c+dx)} + \\
 & +\frac{i \sin(2e+2fx)}{2a^2 d(c+dx)} + \frac{i \sin(4e+4fx)}{4a^2 d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2 d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2 d(c+dx)} - \frac{1}{4a^2 d(c+dx)}
 \end{aligned}$$

input `Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^2),x]`

output

$$\begin{aligned}
 & -1/4 * 1/(a^2 * d * (c + d*x)) - \operatorname{Cos}[2*e + 2*f*x]/(2*a^2 * d * (c + d*x)) - \operatorname{Cos}[2*e \\
 & + 2*f*x]^2/(4*a^2 * d * (c + d*x)) - (I*f * \operatorname{Cos}[2*e - (2*c*f)/d] * \operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/(a^2 * d^2) - \\
 & - (I*f * \operatorname{Cos}[4*e - (4*c*f)/d] * \operatorname{CosIntegral}[(4*c*f)/d + 4*f*x])/(a^2 * d^2) - (f * \operatorname{CosIntegral}[(4*c*f)/d + 4*f*x] * \operatorname{Sin}[4*e - (4*c*f)/d])/(a^2 * d^2) - \\
 & - (f * \operatorname{CosIntegral}[(2*c*f)/d + 2*f*x] * \operatorname{Sin}[2*e - (2*c*f)/d])/(a^2 * d^2) + ((I/2) * \operatorname{Sin}[2*e + 2*f*x])/(a^2 * d * (c + d*x)) + \\
 & + \operatorname{Sin}[2*e + 2*f*x]^2/(4*a^2 * d * (c + d*x)) + ((I/4) * \operatorname{Sin}[4*e + 4*f*x])/(a^2 * d * (c + d*x)) - (f * \operatorname{C} \\
 & \operatorname{os}[2*e - (2*c*f)/d] * \operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a^2 * d^2) + (I*f * \operatorname{Sin}[2 \\
 & *e - (2*c*f)/d] * \operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a^2 * d^2) - (f * \operatorname{Cos}[4*e - (4*c*f)/d] * \operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/(a^2 * d^2) + \\
 & + (I*f * \operatorname{Sin}[4*e - (4*c*f)/d] * \operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/(a^2 * d^2)
 \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4211 $\text{Int}[(c_.) + (d_.)*(x_.)^m*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + \text{Cos}[2*e + 2*f*x]/(2*a) + \text{Sin}[2*e + 2*f*x]/(2*b))^(-n), x], x] /; \ \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&& \ \text{EqQ}[a^2 + b^2, 0] \ \&& \ \text{ILtQ}[m, 0] \ \&& \ \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.40

method	result
risch	$-\frac{1}{4a^2d(dx+c)} - \frac{fe^{-4i(fx+e)}}{4a^2(df+cf)d} + \frac{ife^{\frac{4i(cf-de)}{d}} \expIntegral_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{a^2d^2} - \frac{fe^{-2i(fx+e)}}{2a^2(df+cf)d} + \frac{ife^{\frac{2i(cf-de)}{d}} \expIntegral_1\left(4ifx+4ie+\frac{2i(cf-de)}{d}\right)}{a^2d^2}$

input $\text{int}(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2, x, \text{method}=\text{RETURNVERBOSE})$

output
$$\begin{aligned} & -1/4/a^2/d/(d*x+c) - 1/4/a^2/f*\exp(-4*I*(f*x+e))/(d*f*x+c*f)/d + I/a^2*f/d^2*\exp(4*I*(c*f-d*e)/d)*\text{Ei}(1, 4*I*f*x+4*I*e+4*I*(c*f-d*e)/d) - 1/2/a^2*f*\exp(-2*I*(f*x+e))/(d*f*x+c*f)/d + I/a^2*f/d^2*\exp(2*I*(c*f-d*e)/d)*\text{Ei}(1, 2*I*f*x+2*I*e+2*I*(c*f-d*e)/d) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.33

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx =$$

$$-\frac{\left(\left(4(i dfx + i cf) \text{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + 4(i dfx + i cf) \text{Ei}\left(-\frac{4(i dfx + i cf)}{d}\right) e^{\left(-\frac{4(i de - i cf)}{d}\right)} + a^2 d^3 x + a^2 c d^2 \right) \right)}{4(a^2 d^3 x + a^2 c d^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output
$$\begin{aligned} & -1/4 * ((4*(I*d*f*x + I*c*f)*\text{Ei}(-2*(I*d*f*x + I*c*f)/d)*e^{-(-2*(I*d*e - I*c*f)/d)} + 4*(I*d*f*x + I*c*f)*\text{Ei}(-4*(I*d*f*x + I*c*f)/d)*e^{-(-4*(I*d*e - I*c*f)/d)} + d)*e^{(4*I*f*x + 4*I*e)} + 2*d*e^{(2*I*f*x + 2*I*e)} + d)*e^{-(-4*I*f*x - 4*I*e)/(a^2*d^3*x + a^2*c*d^2)}) \end{aligned}$$

Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx =$$

$$-\frac{\int \frac{1}{c^2 \tan^2(e+fx) - 2ic^2 \tan(e+fx) - c^2 + 2cdx \tan^2(e+fx) - 4icdx \tan(e+fx) - 2cdx + d^2 x^2 \tan^2(e+fx) - 2id^2 x^2 \tan(e+fx) - d^2 x^2} dx}{a^2}$$

input `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e))**2,x)`

output
$$\begin{aligned} & -\text{Integral}(1/(c**2*tan(e + f*x)**2 - 2*I*c**2*tan(e + f*x) - c**2 + 2*c*d*x*tan(e + f*x)**2 - 4*I*c*d*x*tan(e + f*x) - 2*c*d*x + d**2*x**2*tan(e + f*x)**2 - 2*I*d**2*x**2*tan(e + f*x) - d**2*x**2), x)/a**2 \end{aligned}$$

Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.48

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx =$$

$$\frac{-2 f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + f^2 \cos\left(-\frac{4(de-cf)}{d}\right) E_2\left(-\frac{4(-i(fx+e)d+i(de-icf))}{d}\right) + 2i}{4((fx+e)a^2d^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output

$$\begin{aligned} & -1/4*(2*f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f^2*cos(-4*(d*e - c*f)/d)*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + I*f^2*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + f^2)/(((f*x + e)*a^2*d^2 - a^2*d^2*e + a^2*c*d*f)*f) \end{aligned}$$
Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1967 vs. $2(410) = 820$.

Time = 11.12 (sec) , antiderivative size = 1967, normalized size of antiderivative = 4.51

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output

```
1/4*(-4*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*I*d*e*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*I*c*f^3*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 4*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 4*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-4*(d*e - c*f)/d) - 4*d*e*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-4*(d*e - c*f)/d) + 4*c*f^3*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) ...
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + i a \tan(e + fx))^2} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^2 (c + dx)^2} dx$$

input

```
int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)^2),x)
```

output

```
int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)^2), x)
```

Reduce [F]

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx =$$

$$-\frac{\int \frac{1}{\tan(fx+e)^2 c^2 + 2 \tan(fx+e)^2 c dx + \tan(fx+e)^2 d^2 x^2 - 2 \tan(fx+e) c^2 i - 4 \tan(fx+e) c d i x - 2 \tan(fx+e) d^2 i x^2 - c^2 - 2 c d x - d^2 x^2} dx}{a^2}$$

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x)`

output `(- int(1/(tan(e + f*x)**2*c**2 + 2*tan(e + f*x)**2*c*d*x + tan(e + f*x)**2*d**2*x**2 - 2*tan(e + f*x)*c**2*i - 4*tan(e + f*x)*c*d*i*x - 2*tan(e + f*x)*d**2*i*x**2 - c**2 - 2*c*d*x - d**2*x**2),x))/a**2`

$$3.29 \quad \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx$$

Optimal result	248
Mathematica [A] (verified)	249
Rubi [A] (verified)	250
Maple [A] (verified)	251
Fricas [A] (verification not implemented)	252
Sympy [A] (verification not implemented)	253
Maxima [F(-2)]	253
Giac [A] (verification not implemented)	254
Mupad [B] (verification not implemented)	255
Reduce [F]	256

Optimal result

Integrand size = 23, antiderivative size = 396

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx = & -\frac{9d^3e^{-2ie-2ifx}}{64a^3f^4} - \frac{9d^3e^{-4ie-4ifx}}{1024a^3f^4} - \frac{d^3e^{-6ie-6ifx}}{1728a^3f^4} \\ & - \frac{9id^2e^{-2ie-2ifx}(c+dx)}{32a^3f^3} - \frac{9id^2e^{-4ie-4ifx}(c+dx)}{256a^3f^3} \\ & - \frac{id^2e^{-6ie-6ifx}(c+dx)}{288a^3f^3} + \frac{9de^{-2ie-2ifx}(c+dx)^2}{32a^3f^2} \\ & + \frac{9de^{-4ie-4ifx}(c+dx)^2}{128a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)^2}{96a^3f^2} \\ & + \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} \\ & + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d} \end{aligned}$$

output

$$\begin{aligned} & -\frac{9}{64}d^3 \exp(-2*I*e-2*I*f*x)/a^3/f^4 - \frac{9}{1024}d^3 \exp(-4*I*e-4*I*f*x)/a^3/f \\ & ^4 - \frac{1}{1728}d^3 \exp(-6*I*e-6*I*f*x)/a^3/f^4 - \frac{9}{32}d^2 \exp(-2*I*e-2*I*f*x)*(d*x+c)/a^3/f^3 \\ & - \frac{9}{256}d^2 \exp(-4*I*e-4*I*f*x)*(d*x+c)/a^3/f^3 + \frac{1}{32}d \exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^3/f^2 \\ & - \frac{9}{128}d \exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^3/f^2 + \frac{1}{16}d \exp(-2*I*e-2*I*f*x)*(d*x+c)^3/a^3/f^1 \\ & - \frac{9}{48}d \exp(-6*I*e-6*I*f*x)*(d*x+c)^3/a^3/f^1 + \frac{1}{32}d \exp(-6*I*e-6*I*f*x)*(d*x+c)^3/a^3/f^1 \\ & - \frac{9}{32}d \exp(-4*I*e-4*I*f*x)*(d*x+c)^3/a^3/f^1 + \frac{1}{48}d \exp(-6*I*e-6*I*f*x)*(d*x+c)^3/a^3/f^1 \end{aligned}$$

Mathematica [A] (verified)

Time = 3.38 (sec) , antiderivative size = 667, normalized size of antiderivative = 1.68

$$\begin{aligned} & \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx \\ & = \frac{i \sec^3(e+fx) (243(32ic^3f^3 + 8c^2df^2(5 + 12ifx) + 4cd^2f(-9i + 20fx + 24if^2x^2) + d^3(-17 - 36ifx + 12if^2x^2)))}{\dots} \end{aligned}$$

input

```
Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x])^3, x]
```

output

$$\begin{aligned} & ((I/27648)*\text{Sec}[e + f*x]^3*(243*((32*I)*c^3*f^3 + 8*c^2*d*f^2*(5 + (12*I)*f*x) + 4*c*d^2*f*(-9*I + 20*f*x + (24*I)*f^2*x^2) + d^3*(-17 - (36*I)*f*x + 40*f^2*x^2 + (32*I)*f^3*x^3))*\text{Cos}[e + f*x] + 16*(36*c^3*f^3*(I + 6*f*x) + 18*c^2*d*f^2*(1 + (6*I)*f*x + 18*f^2*x^2) + 6*c*d^2*f*(-I + 6*f*x + (18*I)*f^2*x^2 + 36*f^3*x^3) + d^3*(-1 - (6*I)*f*x + 18*f^2*x^2 + (36*I)*f^3*x^3 + 54*f^4*x^4))*\text{Cos}[3*(e + f*x)] - (3645*I)*d^3*\text{Sin}[e + f*x] + 6804*c*d^2*f*\text{Sin}[e + f*x] + (5832*I)*c^2*d*f^2*\text{Sin}[e + f*x] - 2592*c^3*f^3*\text{Sin}[e + f*x] + 6804*d^3*f*x*\text{Sin}[e + f*x] + (11664*I)*c*d^2*f^2*x*\text{Sin}[e + f*x] - 7776*c^2*d*f^3*x*\text{Sin}[e + f*x] + (5832*I)*d^3*f^2*x^2*\text{Sin}[e + f*x] - 2592*d^3*f^3*x^3*\text{Sin}[e + f*x] + (16*I)*d^3*\text{Sin}[3*(e + f*x)] - 96*c*d^2*f*\text{Sin}[3*(e + f*x)] - (288*I)*c^2*d*f^2*\text{Sin}[3*(e + f*x)] + 576*c^3*f^3*\text{Sin}[3*(e + f*x)] - 96*d^3*f*x*\text{Sin}[3*(e + f*x)] - (576*I)*c*d^2*f^2*x*\text{Sin}[3*(e + f*x)] + 1728*c^2*d*f^3*x*\text{Sin}[3*(e + f*x)] + (3456*I)*c^3*f^4*x*\text{Sin}[3*(e + f*x)] - (288*I)*d^3*f^2*x^2*\text{Sin}[3*(e + f*x)] + 1728*c^2*d*f^4*x^2*\text{Sin}[3*(e + f*x)] + (5184*I)*c^2*d*f^4*x^2*\text{Sin}[3*(e + f*x)] + 576*d^3*f^3*x^3*\text{Sin}[3*(e + f*x)] + (3456*I)*c*d^2*f^4*x^3*\text{Sin}[3*(e + f*x)] + (864*I)*d^3*f^4*x^4*\text{Sin}[3*(e + f*x)]))/((a^3*f^4*(-I + \text{Tan}[e + f*x]))^3) \end{aligned}$$

Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx \\
 & \quad \downarrow \text{4212} \\
 & \int \left(\frac{3(c+dx)^3 e^{-2ie-2ifx}}{8a^3} + \frac{3(c+dx)^3 e^{-4ie-4ifx}}{8a^3} + \frac{(c+dx)^3 e^{-6ie-6ifx}}{8a^3} + \frac{(c+dx)^3}{8a^3} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -\frac{9id^2(c+dx)e^{-2ie-2ifx}}{32a^3f^3} - \frac{9id^2(c+dx)e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2(c+dx)e^{-6ie-6ifx}}{288a^3f^3} + \\
 & \frac{9d(c+dx)^2e^{-2ie-2ifx}}{32a^3f^2} + \frac{9d(c+dx)^2e^{-4ie-4ifx}}{128a^3f^2} + \frac{d(c+dx)^2e^{-6ie-6ifx}}{96a^3f^2} + \\
 & \frac{3i(c+dx)^3e^{-2ie-2ifx}}{16a^3f} + \frac{3i(c+dx)^3e^{-4ie-4ifx}}{32a^3f} + \frac{i(c+dx)^3e^{-6ie-6ifx}}{48a^3f} + \frac{(c+dx)^4}{32a^3d} - \\
 & \frac{9d^3e^{-2ie-2ifx}}{64a^3f^4} - \frac{9d^3e^{-4ie-4ifx}}{1024a^3f^4} - \frac{d^3e^{-6ie-6ifx}}{1728a^3f^4}
 \end{aligned}$$

input `Int[(c + d*x)^3/(a + I*a*Tan[e + f*x])^3,x]`

output

$$\begin{aligned} & \frac{(-9d^3E^{\left((-2I)\right)}e - (2I)f^2x)/(64a^3f^4) - (9d^3E^{\left((-4I)\right)}e - (4I)f^2x)/(1024a^3f^4) - (d^3E^{\left((-6I)\right)}e - (6I)f^2x)/(1728a^3f^4) - ((9I)/32)d^2E^{\left((-2I)\right)}e - (2I)f^2x)(c + d^2x)/(a^3f^3) - (((9I)/256)d^2E^{\left((-4I)\right)}e - (4I)f^2x)(c + d^2x)/(a^3f^3) - ((I/288)d^2E^{\left((-6I)\right)}e - (6I)f^2x)(c + d^2x)/(a^3f^3) + (9d^2E^{\left((-2I)\right)}e - (2I)f^2x)(c + d^2x)^2/(32a^3f^2) + (9d^2E^{\left((-4I)\right)}e - (4I)f^2x)(c + d^2x)^2/(128a^3f^2) + (d^2E^{\left((-6I)\right)}e - (6I)f^2x)(c + d^2x)^2/(96a^3f^2) + ((3I)/16)E^{\left((-2I)\right)}e - (2I)f^2x)(c + d^2x)^3/(a^3f) + ((3I)/32)E^{\left((-4I)\right)}e - (4I)f^2x)(c + d^2x)^3/(a^3f) + ((I/48)E^{\left((-6I)\right)}e - (6I)f^2x)(c + d^2x)^3/(a^3f) + (c + d^2x)^4/(32a^3d) \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4212 $\text{Int}[(c_.) + (d_.)(x_.)^m_*(a_.) + (b_.)*\tan[(e_.) + (f_.)(x_.)]^n_, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d^2x)^m, (1/(2a) + E^{(2(a/b)*(e + f)x)})/(2a)^{-n}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 1.11 (sec), antiderivative size = 396, normalized size of antiderivative = 1.00

method	result
risch	$\frac{d^3x^4}{32a^3} + \frac{d^2cx^3}{8a^3} + \frac{3d^2c^2x^2}{16a^3} + \frac{c^3x}{8a^3} + \frac{c^4}{32a^3d} + \frac{3i(4d^3x^3f^3 + 12cd^2f^3x^2 - 6id^3f^2x^2 + 12c^2df^3x - 12cd^2f^2x + 4c^3f^3 - 6ic^2df^2)}{64a^3f^4}$

input $\text{int}((d^2x^3 + c)^3/(a + I*a*\tan(f*x + e))^3, x, \text{method} = \text{RETURNVERBOSE})$

output

$$\begin{aligned} & \frac{1}{32} a^3 d^3 x^4 + \frac{1}{8} a^3 d^2 c x^3 + \frac{3}{16} a^3 d c^2 x^2 + \frac{1}{8} a^3 c^3 x + \frac{1}{32} a \\ & ^3 d c^4 + \frac{3}{64} I^*(4 d^3 x^3 f^3 - 6 I d^3 f^2 x^2 + 12 c d^2 f^3 x^2 - 12 I c d^2 f \\ & * f^2 x + 12 c^2 d f^3 x - 6 I c^2 d f^2 + 4 c^3 f^3 - 6 d^3 f x + 3 I d^3 - 6 c d^2 f) \\ & / a^3 / f^4 \exp(-2 I^*(f x + e)) + \frac{3}{1024} I^*(32 d^3 x^3 f^3 - 24 I d^3 f^2 x^2 + 96 c \\ & d^2 f^3 x^2 - 48 I c d^2 f^2 x + 96 c^2 d f^3 x - 24 I c^2 d f^2 + 32 c^3 f^3 - 12 d \\ & ^3 f x + 3 I d^3 - 12 c d^2 f) / a^3 / f^4 \exp(-4 I^*(f x + e)) + \frac{1}{1728} I^*(36 d^3 x^3 f^3 \\ & - 18 I d^3 f^2 x^2 + 108 c d^2 f^3 x^2 - 36 I c d^2 f^2 x + 108 c^2 d f^3 x - 18 \\ & * I c^2 d f^2 + 36 c^3 f^3 - 6 d^3 f x + I d^3 - 6 c d^2 f) / a^3 / f^4 \exp(-6 I^*(f x + e)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 370, normalized size of antiderivative = 0.93

$$\begin{aligned} & \int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx \\ & = \frac{(576 i d^3 f^3 x^3 + 576 i c^3 f^3 + 288 c^2 d f^2 - 96 i c d^2 f - 16 d^3 - 288 (-6 i c d^2 f^3 - d^3 f^2) x^2 - 96 (-18 i c^2 d f^3 - 6 i c^3 d^2 f^2) x)}{1} \end{aligned}$$

input

```
integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")
```

output

$$\begin{aligned} & \frac{1}{27648} (576 I d^3 f^3 x^3 + 576 I c^3 f^3 + 288 c^2 d f^2 - 96 I c d^2 f \\ & - 16 d^3 - 288 (-6 I c d^2 f^3 - d^3 f^2) x^2 - 96 (-18 I c^2 d f^3 - 6 c \\ & d^2 f^2 + I d^3 f) x + 864 (d^3 f^4 x^4 + 4 c d^2 f^4 x^3 + 6 c^2 d f^4 x^2 \\ & + 4 c^3 f^4 x) e^{(6 I f x + 6 I e)} - 1296 (-4 I d^3 f^3 x^3 - 4 I c^3 f^3 \\ & x^3 - 6 c^2 d f^2 + 6 I c d^2 f^2 + 3 d^3 + 6 (-2 I c d^2 f^3 - d^3 f^2) x^2 + \\ & 6 (-2 I c^2 d f^3 - 2 c d^2 f^2 + I d^3 f) x) e^{(4 I f x + 4 I e)} - 81 (- \\ & 32 I d^3 f^3 x^3 - 32 I c^3 f^3 - 24 c^2 d f^2 + 12 I c d^2 f^2 + 3 d^3 + 24 \\ & (-4 I c d^2 f^3 - d^3 f^2) x^2 + 12 (-8 I c^2 d f^3 - 4 c d^2 f^2 + I d^3 \\ & f) x) e^{(2 I f x + 2 I e)} e^{(-6 I f x - 6 I e)} / (a^3 f^4) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 945, normalized size of antiderivative = 2.39

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = \text{Too large to display}$$

input `integrate((d*x+c)**3/(a+I*a*tan(f*x+e))**3, x)`

output

```
Piecewise(((2359296*I*a**6*c**3*f**11*exp(6*I*e) + 7077888*I*a**6*c**2*d*f**11*x*exp(6*I*e) + 1179648*a**6*c**2*d*f**10*exp(6*I*e) + 7077888*I*a**6*c*d**2*f**11*x**2*exp(6*I*e) + 2359296*a**6*c*d**2*f**10*x*exp(6*I*e) - 393216*I*a**6*c*d**2*f**9*exp(6*I*e) + 2359296*I*a**6*d**3*f**11*x**3*exp(6*I*e) + 1179648*a**6*d**3*f**10*x**2*exp(6*I*e) - 393216*I*a**6*d**3*f**9*x*exp(6*I*e) - 65536*a**6*d**3*f**8*exp(6*I*e))*exp(-6*I*f*x) + (10616832*I*a**6*c**3*f**11*exp(8*I*e) + 31850496*I*a**6*c**2*d*f**11*x*exp(8*I*e) + 7962624*a**6*c**2*d*f**10*exp(8*I*e) + 31850496*I*a**6*c*d**2*f**11*x**2*exp(8*I*e) + 15925248*a**6*c*d**2*f**10*x*exp(8*I*e) - 3981312*I*a**6*c*d**2*f**9*exp(8*I*e) + 10616832*I*a**6*d**3*f**11*x**3*exp(8*I*e) + 7962624*a**6*d**3*f**10*x**2*exp(8*I*e) - 3981312*I*a**6*d**3*f**9*x*exp(8*I*e) - 995328*a**6*d**3*f**8*exp(8*I*e))*exp(-4*I*f*x) + (21233664*I*a**6*c**3*f**11*exp(10*I*e) + 63700992*I*a**6*c**2*d*f**11*x*exp(10*I*e) + 31850496*a**6*c**2*d*f**10*exp(10*I*e) + 63700992*I*a**6*c*d**2*f**11*x**2*exp(10*I*e) + 63700992*a**6*c*d**2*f**10*x*exp(10*I*e) - 31850496*I*a**6*c*d**2*f**9*exp(10*I*e) + 21233664*I*a**6*d**3*f**11*x**3*exp(10*I*e) + 31850496*a**6*d**3*f**10*x**2*exp(10*I*e) - 31850496*I*a**6*d**3*f**9*x*exp(10*I*e) - 15925248*a**6*d**3*f**8*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(113246208*a**9*f**12), Ne(a**9*f**12*exp(12*I*e), 0)), (x**4*(3*d**3*exp(4*I*e) + 3*d**3*exp(2*I*e) + d**3)*exp(-6*I*e)/(32*a**3) + x**3*(3*c*d**2*exp(4*I...)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3, x, algorithm="maxima")`

output

```
Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 548, normalized size of antiderivative = 1.38

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(864 d^3 f^4 x^4 e^{(6i f x + 6i e)} + 3456 c d^2 f^4 x^3 e^{(6i f x + 6i e)} + 5184 c^2 d f^4 x^2 e^{(6i f x + 6i e)} + 5184 i d^3 f^3 x^3 e^{(4i f x + 4i e)} + 2592 c^2 d^2 f^3 x^2 e^{(4i f x + 4i e)} + 15552 c d^2 f^3 x e^{(4i f x + 4i e)} + 7776 c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} + 1944 c^2 d^2 f^2 x e^{(4i f x + 4i e)} + 576 c^3 d^2 f x^3 e^{(4i f x + 4i e)} + 3888 c^3 d^2 f x^2 e^{(4i f x + 4i e)} + 1296 c^3 d^2 f x e^{(4i f x + 4i e)} + 3456 c^4 f x^4 e^{(4i f x + 4i e)} + 10368 c^4 f^2 x^3 e^{(4i f x + 4i e)} + 20736 c^4 f^3 x^2 e^{(4i f x + 4i e)} + 15552 c^4 f^4 x e^{(4i f x + 4i e)} + 7776 c^5 f^3 x^3 e^{(4i f x + 4i e)} + 23328 c^5 f^4 x^2 e^{(4i f x + 4i e)} + 23328 c^5 f^5 x e^{(4i f x + 4i e)} + 576 c^6 f^4 x^2 e^{(4i f x + 4i e)} + 1152 c^6 f^5 x e^{(4i f x + 4i e)} + 1152 c^7 f^5 x e^{(4i f x + 4i e)} + 23328 c^8 f^6 x e^{(4i f x + 4i e)} + 15552 c^9 f^7 x e^{(4i f x + 4i e)} + 7776 c^{10} f^8 x e^{(4i f x + 4i e)} + 2592 c^{11} f^9 x e^{(4i f x + 4i e)} + 5184 c^{12} f^{10} x e^{(4i f x + 4i e)} + 864 c^{13} f^{11} x e^{(4i f x + 4i e)} + 3456 c^{14} f^{12} x e^{(4i f x + 4i e)} + 15552 c^{15} f^{13} x e^{(4i f x + 4i e)} + 576 c^{16} f^{14} x e^{(4i f x + 4i e)} + 1944 c^{17} f^{15} x e^{(4i f x + 4i e)} + 5184 i c^{18} f^{16} x e^{(4i f x + 4i e)} + 1296 i c^{19} f^{17} x e^{(4i f x + 4i e)} + 3888 i c^{20} f^{18} x e^{(4i f x + 4i e)} + 10368 i c^{21} f^{19} x e^{(4i f x + 4i e)} + 23328 i c^{22} f^{20} x e^{(4i f x + 4i e)} + 46656 i c^{23} f^{21} x e^{(4i f x + 4i e)} + 86400 i c^{24} f^{22} x e^{(4i f x + 4i e)} + 155520 i c^{25} f^{23} x e^{(4i f x + 4i e)} + 233280 i c^{26} f^{24} x e^{(4i f x + 4i e)} + 233280 i c^{27} f^{25} x e^{(4i f x + 4i e)} + 155520 i c^{28} f^{26} x e^{(4i f x + 4i e)} + 86400 i c^{29} f^{27} x e^{(4i f x + 4i e)} + 46656 i c^{30} f^{28} x e^{(4i f x + 4i e)} + 15552 i c^{31} f^{29} x e^{(4i f x + 4i e)} + 3456 i c^{32} f^{30} x e^{(4i f x + 4i e)} + 864 i c^{33} f^{31} x e^{(4i f x + 4i e)} + 23328 i c^{34} f^{32} x e^{(4i f x + 4i e)} + 46656 i c^{35} f^{33} x e^{(4i f x + 4i e)} + 86400 i c^{36} f^{34} x e^{(4i f x + 4i e)} + 155520 i c^{37} f^{35} x e^{(4i f x + 4i e)} + 233280 i c^{38} f^{36} x e^{(4i f x + 4i e)} + 233280 i c^{39} f^{37} x e^{(4i f x + 4i e)} + 155520 i c^{40} f^{38} x e^{(4i f x + 4i e)} + 86400 i c^{41} f^{39} x e^{(4i f x + 4i e)} + 46656 i c^{42} f^{40} x e^{(4i f x + 4i e)} + 15552 i c^{43} f^{41} x e^{(4i f x + 4i e)} + 3456 i c^{44} f^{42} x e^{(4i f x + 4i e)} + 864 i c^{45} f^{43} x e^{(4i f x + 4i e)} + 23328 i c^{46} f^{44} x e^{(4i f x + 4i e)} + 46656 i c^{47} f^{45} x e^{(4i f x + 4i e)} + 86400 i c^{48} f^{46} x e^{(4i f x + 4i e)} + 155520 i c^{49} f^{47} x e^{(4i f x + 4i e)} + 233280 i c^{50} f^{48} x e^{(4i f x + 4i e)} + 233280 i c^{51} f^{49} x e^{(4i f x + 4i e)} + 155520 i c^{52} f^{50} x e^{(4i f x + 4i e)} + 86400 i c^{53} f^{51} x e^{(4i f x + 4i e)} + 46656 i c^{54} f^{52} x e^{(4i f x + 4i e)} + 15552 i c^{55} f^{53} x e^{(4i f x + 4i e)} + 3456 i c^{56} f^{54} x e^{(4i f x + 4i e)} + 864 i c^{57} f^{55} x e^{(4i f x + 4i e)} + 23328 i c^{58} f^{56} x e^{(4i f x + 4i e)} + 46656 i c^{59} f^{57} x e^{(4i f x + 4i e)} + 86400 i c^{60} f^{58} x e^{(4i f x + 4i e)} + 155520 i c^{61} f^{59} x e^{(4i f x + 4i e)} + 233280 i c^{62} f^{60} x e^{(4i f x + 4i e)} + 233280 i c^{63} f^{61} x e^{(4i f x + 4i e)} + 155520 i c^{64} f^{62} x e^{(4i f x + 4i e)} + 86400 i c^{65} f^{63} x e^{(4i f x + 4i e)} + 46656 i c^{66} f^{64} x e^{(4i f x + 4i e)} + 15552 i c^{67} f^{65} x e^{(4i f x + 4i e)} + 3456 i c^{68} f^{66} x e^{(4i f x + 4i e)} + 864 i c^{69} f^{67} x e^{(4i f x + 4i e)} + 23328 i c^{70} f^{68} x e^{(4i f x + 4i e)} + 46656 i c^{71} f^{69} x e^{(4i f x + 4i e)} + 86400 i c^{72} f^{70} x e^{(4i f x + 4i e)} + 155520 i c^{73} f^{71} x e^{(4i f x + 4i e)} + 233280 i c^{74} f^{72} x e^{(4i f x + 4i e)} + 233280 i c^{75} f^{73} x e^{(4i f x + 4i e)} + 155520 i c^{76} f^{74} x e^{(4i f x + 4i e)} + 86400 i c^{77} f^{75} x e^{(4i f x + 4i e)} + 46656 i c^{78} f^{76} x e^{(4i f x + 4i e)} + 15552 i c^{79} f^{77} x e^{(4i f x + 4i e)} + 3456 i c^{80} f^{78} x e^{(4i f x + 4i e)} + 864 i c^{81} f^{79} x e^{(4i f x + 4i e)} + 23328 i c^{82} f^{80} x e^{(4i f x + 4i e)} + 46656 i c^{83} f^{81} x e^{(4i f x + 4i e)} + 86400 i c^{84} f^{82} x e^{(4i f x + 4i e)} + 155520 i c^{85} f^{83} x e^{(4i f x + 4i e)} + 233280 i c^{86} f^{84} x e^{(4i f x + 4i e)} + 233280 i c^{87} f^{85} x e^{(4i f x + 4i e)} + 155520 i c^{88} f^{86} x e^{(4i f x + 4i e)} + 86400 i c^{89} f^{87} x e^{(4i f x + 4i e)} + 46656 i c^{90} f^{88} x e^{(4i f x + 4i e)} + 15552 i c^{91} f^{89} x e^{(4i f x + 4i e)} + 3456 i c^{92} f^{90} x e^{(4i f x + 4i e)} + 864 i c^{93} f^{91} x e^{(4i f x + 4i e)} + 23328 i c^{94} f^{92} x e^{(4i f x + 4i e)} + 46656 i c^{95} f^{93} x e^{(4i f x + 4i e)} + 86400 i c^{96} f^{94} x e^{(4i f x + 4i e)} + 155520 i c^{97} f^{95} x e^{(4i f x + 4i e)} + 233280 i c^{98} f^{96} x e^{(4i f x + 4i e)} + 233280 i c^{99} f^{97} x e^{(4i f x + 4i e)} + 155520 i c^{100} f^{98} x e^{(4i f x + 4i e)} + 86400 i c^{101} f^{99} x e^{(4i f x + 4i e)} + 46656 i c^{102} f^{100} x e^{(4i f x + 4i e)} + 15552 i c^{103} f^{101} x e^{(4i f x + 4i e)} + 3456 i c^{104} f^{102} x e^{(4i f x + 4i e)} + 864 i c^{105} f^{103} x e^{(4i f x + 4i e)} + 23328 i c^{106} f^{104} x e^{(4i f x + 4i e)} + 46656 i c^{107} f^{105} x e^{(4i f x + 4i e)} + 86400 i c^{108} f^{106} x e^{(4i f x + 4i e)} + 155520 i c^{109} f^{107} x e^{(4i f x + 4i e)} + 233280 i c^{110} f^{108} x e^{(4i f x + 4i e)} + 233280 i c^{111} f^{109} x e^{(4i f x + 4i e)} + 155520 i c^{112} f^{110} x e^{(4i f x + 4i e)} + 86400 i c^{113} f^{111} x e^{(4i f x + 4i e)} + 46656 i c^{114} f^{112} x e^{(4i f x + 4i e)} + 15552 i c^{115} f^{113} x e^{(4i f x + 4i e)} + 3456 i c^{116} f^{114} x e^{(4i f x + 4i e)} + 864 i c^{117} f^{115} x e^{(4i f x + 4i e)} + 23328 i c^{118} f^{116} x e^{(4i f x + 4i e)} + 46656 i c^{119} f^{117} x e^{(4i f x + 4i e)} + 86400 i c^{120} f^{118} x e^{(4i f x + 4i e)} + 155520 i c^{121} f^{119} x e^{(4i f x + 4i e)} + 233280 i c^{122} f^{120} x e^{(4i f x + 4i e)} + 233280 i c^{123} f^{121} x e^{(4i f x + 4i e)} + 155520 i c^{124} f^{122} x e^{(4i f x + 4i e)} + 86400 i c^{125} f^{123} x e^{(4i f x + 4i e)} + 46656 i c^{126} f^{124} x e^{(4i f x + 4i e)} + 15552 i c^{127} f^{125} x e^{(4i f x + 4i e)} + 3456 i c^{128} f^{126} x e^{(4i f x + 4i e)} + 864 i c^{129} f^{127} x e^{(4i f x + 4i e)} + 23328 i c^{130} f^{128} x e^{(4i f x + 4i e)} + 46656 i c^{131} f^{129} x e^{(4i f x + 4i e)} + 86400 i c^{132} f^{130} x e^{(4i f x + 4i e)} + 155520 i c^{133} f^{131} x e^{(4i f x + 4i e)} + 233280 i c^{134} f^{132} x e^{(4i f x + 4i e)} + 233280 i c^{135} f^{133} x e^{(4i f x + 4i e)} + 155520 i c^{136} f^{134} x e^{(4i f x + 4i e)} + 86400 i c^{137} f^{135} x e^{(4i f x + 4i e)} + 46656 i c^{138} f^{136} x e^{(4i f x + 4i e)} + 15552 i c^{139} f^{137} x e^{(4i f x + 4i e)} + 3456 i c^{140} f^{138} x e^{(4i f x + 4i e)} + 864 i c^{141} f^{139} x e^{(4i f x + 4i e)} + 23328 i c^{142} f^{140} x e^{(4i f x + 4i e)} + 46656 i c^{143} f^{141} x e^{(4i f x + 4i e)} + 86400 i c^{144} f^{142} x e^{(4i f x + 4i e)} + 155520 i c^{145} f^{143} x e^{(4i f x + 4i e)} + 233280 i c^{146} f^{144} x e^{(4i f x + 4i e)} + 233280 i c^{147} f^{145} x e^{(4i f x + 4i e)} + 155520 i c^{148} f^{146} x e^{(4i f x + 4i e)} + 86400 i c^{149} f^{147} x e^{(4i f x + 4i e)} + 46656 i c^{150} f^{148} x e^{(4i f x + 4i e)} + 15552 i c^{151} f^{149} x e^{(4i f x + 4i e)} + 3456 i c^{152} f^{150} x e^{(4i f x + 4i e)} + 864 i c^{153} f^{151} x e^{(4i f x + 4i e)} + 23328 i c^{154} f^{152} x e^{(4i f x + 4i e)} + 46656 i c^{155} f^{153} x e^{(4i f x + 4i e)} + 86400 i c^{156} f^{154} x e^{(4i f x + 4i e)} + 155520 i c^{157} f^{155} x e^{(4i f x + 4i e)} + 233280 i c^{158} f^{156} x e^{(4i f x + 4i e)} + 233280 i c^{159} f^{157} x e^{(4i f x + 4i e)} + 155520 i c^{160} f^{158} x e^{(4i f x + 4i e)} + 86400 i c^{161} f^{159} x e^{(4i f x + 4i e)} + 46656 i c^{162} f^{160} x e^{(4i f x + 4i e)} + 15552 i c^{163} f^{161} x e^{(4i f x + 4i e)} + 3456 i c^{164} f^{162} x e^{(4i f x + 4i e)} + 864 i c^{165} f^{163} x e^{(4i f x + 4i e)} + 23328 i c^{166} f^{164} x e^{(4i f x + 4i e)} + 46656 i c^{167} f^{165} x e^{(4i f x + 4i e)} + 86400 i c^{168} f^{166} x e^{(4i f x + 4i e)} + 155520 i c^{169} f^{167} x e^{(4i f x + 4i e)} + 233280 i c^{170} f^{168} x e^{(4i f x + 4i e)} + 233280 i c^{171} f^{169} x e^{(4i f x + 4i e)} + 155520 i c^{172} f^{170} x e^{(4i f x + 4i e)} + 86400 i c^{173} f^{171} x e^{(4i f x + 4i e)} + 46656 i c^{174} f^{172} x e^{(4i f x + 4i e)} + 15552 i c^{175} f^{173} x e^{(4i f x + 4i e)} + 3456 i c^{176} f^{174} x e^{(4i f x + 4i e)} + 864 i c^{177} f^{175} x e^{(4i f x + 4i e)} + 23328 i c^{178} f^{176} x e^{(4i f x + 4i e)} + 46656 i c^{179} f^{177} x e^{(4i f x + 4i e)} + 86400 i c^{180} f^{178} x e^{(4i f x + 4i e)} + 155520 i c^{181} f^{179} x e^{(4i f x + 4i e)} + 233280 i c^{182} f^{180} x e^{(4i f x + 4i e)} + 233280 i c^{183} f^{181} x e^{(4i f x + 4i e)} + 155520 i c^{184} f^{182} x e^{(4i f x + 4i e)} + 86400 i c^{185} f^{183} x e^{(4i f x + 4i e)} + 46656 i c^{186} f^{184} x e^{(4i f x + 4i e)} + 15552 i c^{187} f^{185} x e^{(4i f x + 4i e)} + 3456 i c^{188} f^{186} x e^{(4i f x + 4i e)} + 864 i c^{189} f^{187} x e^{(4i f x + 4i e)} + 23328 i c^{190} f^{188} x e^{(4i f x + 4i e)} + 46656 i c^{191} f^{189} x e^{(4i f x + 4i e)} + 86400 i c^{192} f^{190} x e^{(4i f x + 4i e)} + 155520 i c^{193} f^{191} x e^{(4i f x + 4i e)} + 233280 i c^{194} f^{192} x e^{(4i f x + 4i e)} + 233280 i c^{195} f^{193} x e^{(4i f x + 4i e)} + 155520 i c^{196} f^{194} x e^{(4i f x + 4i e)} + 86400 i c^{197} f^{195} x e^{(4i f x + 4i e)} + 46656 i c^{198} f^{196} x e^{(4i f x + 4i e)} + 15552 i c^{199} f^{197} x e^{(4i f x + 4i e)} + 3456 i c^{200} f^{198} x e^{(4i f x + 4i e)} + 864 i c^{201} f^{199} x e^{(4i f x + 4i e)} + 23328 i c^{202} f^{200} x e^{(4i f x + 4i e)} + 46656 i c^{203} f^{201} x e^{(4i f x + 4i e)} + 86400 i c^{204} f^{202} x e^{(4i f x + 4i e)} + 155520 i c^{205} f^{203} x e^{(4i f x + 4i e)} + 233280 i c^{206} f^{204} x e^{(4i f x + 4i e)} + 233280 i c^{207} f^{205} x e^{(4i f x + 4i e)} + 155520 i c^{208} f^{206} x e^{(4i f x + 4i e)} + 86400 i c^{209} f^{207} x e^{(4i f x + 4i e)} + 46656 i c^{210} f^{208} x e^{(4i f x + 4i e)} + 15552 i c^{211} f^{209} x e^{(4i f x + 4i e)} + 3456 i c^{212} f^{210} x e^{(4i f x + 4i e)} + 864 i c^{213} f^{211} x e^{(4i f x + 4i e)} + 23328 i c^{214} f^{212} x e^{(4i f x + 4i e)} + 46656 i c^{215} f^{213} x e^{(4i f x + 4i e)} + 86400 i c^{216} f^{214} x e^{(4i f x + 4i e)} + 155520 i c^{217} f^{215} x e^{(4i f x + 4i e)} + 233280 i c^{218} f^{216} x e^{(4i f x + 4i e)} + 233280 i c^{219} f^{217} x e^{(4i f x + 4i e)} + 155520 i c^{220} f^{218} x e^{(4i f x + 4i e)} + 86400 i c^{221} f^{219} x e^{(4i f x + 4i e)} + 46656 i c^{222} f^{220} x e^{(4i f x + 4i e)} + 15552 i c^{223} f^{221} x e^{(4i f x + 4i e)} + 3456 i c^{224} f^{222} x e^{(4i f x + 4i e)} + 864 i c^{225} f^{223} x e^{(4i f x + 4i e)} + 23328 i c^{226} f^{224} x e^{(4i f x + 4i e)} + 46656 i c^{227} f^{225} x e^{(4i f x + 4i e)} + 86400 i c^{228} f^{226} x e^{(4i f x + 4i e)} + 155520 i c^{229} f^{227} x e^{(4i f x + 4i e)} + 233280 i c^{230} f^{228} x e^{(4i f x + 4i e)} + 233280 i c^{231} f^{229} x e^{(4i f x + 4i e)} + 155520 i c^{232} f^{230} x e^{(4i f x + 4i e)} + 86400 i c^{233} f^{231} x e^{(4i f x + 4i e)} + 46656 i c^{234} f^{232} x e^{(4i f x + 4i e)} + 15552 i c^{235} f^{233} x e^{(4i f x + 4i e)} + 3456 i c^{236} f^{234} x e^{(4i f x + 4i e)} + 864 i c^{237} f^{235} x e^{(4i f x + 4i e)} + 23328 i c^{238} f^{236} x e^{(4i f x + 4i e)} + 46656 i c^{239} f^{237} x e^{(4i f x + 4i e)} + 86400 i c^{240} f^{238} x e^{(4i f x + 4i e)} + 155520 i c^{241} f^{239} x e^{(4i f x + 4i e)} + 233280 i c^{242} f^{240} x e^{(4i f x + 4i e)} + 233280 i c^{243} f^{241} x e^{(4i f x + 4i e)} + 155520 i c^{244} f^{242} x e^{(4i f x + 4i e)} + 86400 i c^{245} f^{243} x e^{(4i f x + 4i e)} + 46656 i c^{246} f^{244} x e^{(4i f x + 4i e)} + 15552 i c^{247} f^{245} x e^{(4i f x + 4i e)} + 3456 i c^{248} f^{246} x e^{(4i f x + 4i e)} + 864 i c^{249} f^{247} x e^{(4i f x + 4i e)} + 23328 i c^{250} f^{248} x e^{(4i f x + 4i e)} + 46656 i c^{251} f^{249} x e^{(4i f x + 4i e)} + 86400 i c^{252} f^{250} x e^{(4i f x + 4i e)} + 155520 i c^{253} f^{251} x e^{(4i f x + 4i e)} + 233280 i c^{254} f^{252} x e^{(4i f x + 4i e)} + 233280 i c^{255} f^{253} x e^{(4i f x + 4i e)} + 155520 i c^{256} f^{254} x e^{(4i f x + 4i e)} + 86400 i c^{257} f^{255} x e^{(4i f x + 4i e)} + 46656 i c^{258} f^{256} x e^{(4i f x + 4i e)} + 15552 i c^{259} f^{257} x e^{(4i f x + 4i e)} + 3456 i c^{260} f^{258} x e^{(4i f x + 4i e)} + 864 i c^{261} f^{259} x e^{(4i f x + 4i e)} + 23328 i c^{262} f^{260} x e^{(4i f x + 4i e)} + 46656 i c^{263} f^{261} x e^{(4i f x + 4i e)} + 86400 i c^{264} f^{262} x e^{(4i f x + 4i e)} + 155520 i c^{265} f^{263} x e^{(4i f x + 4i e)} + 233280 i c^{266} f^{264} x e^{(4i f x + 4i e)} + 233280 i c^{267} f^{265} x e^{(4i f x + 4i e)} + 155520 i c^{268} f^{266} x e^{(4i f x + 4i e)} + 86400 i c^{269} f^{267} x e^{(4i f x + 4i e)} + 46656 i c^{270} f^{268} x e^{(4i f x + 4i e)} + 15552 i c^{271} f^{269} x e^{(4i f x + 4i e)} + 3456 i c^{272} f^{270} x e^{(4i f x + 4i e)} + 864 i c^{273} f^{271} x e^{(4i f x + 4i e)} + 23328 i c^{274} f^{272} x e^{(4i f x + 4i e)} + 46656 i c^{275} f^{273} x e^{(4i f x + 4i e)} + 86400 i c^{276} f^{274} x e^{(4i f x + 4i e)} + 155520 i c^{277} f^{275} x e^{(4i f x + 4i e)} + 233280 i c^{278} f^{276} x e^{(4i f x + 4i e)} + 233280 i c^{279} f^{277} x e^{(4i f x + 4i e)} + 155520 i c^{280} f^{278} x e^{(4i f x + 4i e)} + 86400 i c^{281} f^{279} x e^{(4i f x + 4i e)} + 46656 i c^{282} f^{280} x e^{(4i f x + 4i e)} + 15552 i c^{283} f^{281} x e^{(4i f x + 4i e)} + 3456 i c^{284} f^{282} x e^{(4i f x + 4i e)} + 864 i c^{285} f^{283} x e^{(4i f x + 4i e)} + 23328 i c^{286} f^{284} x e^{(4i f x + 4i e)} + 46656 i c^{287} f^{285} x e^{(4i f x + 4i e)} + 86400 i c^{288} f^{286} x e^{(4i f x + 4i e)} + 155520 i c^{289} f^{287} x e^{(4i f x + 4i e)} + 233280 i c^{290} f^{288} x e^{(4i f x + 4i e)} + 233280 i c^{291} f^{289} x e^{(4i f x + 4i e)} + 155520 i c^{292} f^{290} x e^{(4i f x + 4i e)} + 86400 i c^{293} f^{291} x e^{(4i f x + 4i e)} + 46656 i c^{294} f^{292} x e^{(4i f x + 4i e)} + 15552 i c^{295} f^{293} x e^{(4i f x + 4i e)} + 3456 i c^{296} f^{294} x e^{(4i f x + 4i e)} + 864 i c^{297} f^{295} x e^{(4i f x + 4i e)} + 23328 i c^{298} f^{296} x e^{(4i f x + 4i e)} + 46656 i c^{299} f^{297} x e^{(4i f x + 4i e)} + 86400 i c^{300} f^{298} x e^{(4i f x + 4i e)} + 155520 i c^{301} f^{299} x e^{(4i f x + 4i e)} + 233280 i c^{302} f^{300} x e^{(4i f x + 4i e)} + 233280 i c^{303} f^{301} x e^{(4i f x + 4i e)} + 155520 i c^{304} f^{302} x e^{(4i f x + 4i e)} + 86400 i c^{305} f^{303} x e^{(4i f x + 4i e)} + 46656 i c^{306} f^{304} x e^{(4i f x + 4i e)} + 15552 i c^{307} f^{305} x e^{(4i f x + 4i e)} + 3456 i c^{308} f^{306} x e^{(4i f x + 4i e)} + 864 i c^{309} f^{307} x e^{(4i f x + 4i e)} + 23328 i c^{310} f^{308} x e^{(4i f x + 4i e)} + 46656 i c^{311} f^{309} x e^{(4i f x + 4i e)} + 86400 i c^{312} f^{310} x e^{(4i f x + 4i e)} + 155520 i c^{313} f^{311} x e^{(4i f x + 4i e)} + 233280 i c^{314} f^{312} x e^{(4i f x + 4i e)} + 233280 i c^{315} f^{313} x e^{(4i f x + 4i e)} + 155520 i c^{316} f^{314} x e^{(4i f x + 4i e)} + 86400 i c^{317} f^{315} x e^{(4i f x + 4i e)} + 46656 i c^{318} f^{316} x e^{(4i f x + 4i e)} + 15552 i c^{319} f^{317} x e^{(4i f x + 4i e)} + 3456 i c^{320} f^{318} x e^{(4i f x + 4i e)} + 864 i c^{321$$

Mupad [B] (verification not implemented)

Time = 9.27 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.04

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = e^{-e 2i - f x 2i} \left(\frac{(12 c^3 f^3 - c^2 d f^2 18i - 18 c d^2 f + d^3 9i) 1i}{64 a^3 f^4} \right.$$

$$+ \frac{d^3 x^3 3i}{16 a^3 f} - \frac{d x (-2 c^2 f^2 + c d f 2i + d^2) 9i}{32 a^3 f^3}$$

$$- \frac{d^2 x^2 (-2 c f + d 1i) 9i}{32 a^3 f^2} \Big)$$

$$+ e^{-e 4i - f x 4i} \left(\frac{(96 c^3 f^3 - c^2 d f^2 72i - 36 c d^2 f + d^3 9i) 1i}{1024 a^3 f^4} \right.$$

$$+ \frac{d^3 x^3 3i}{32 a^3 f} - \frac{d x (-8 c^2 f^2 + c d f 4i + d^2) 9i}{256 a^3 f^3}$$

$$- \frac{d^2 x^2 (-4 c f + d 1i) 9i}{128 a^3 f^2} \Big)$$

$$+ e^{-e 6i - f x 6i} \left(\frac{(36 c^3 f^3 - c^2 d f^2 18i - 6 c d^2 f + d^3 1i) 1i}{1728 a^3 f^4} \right.$$

$$+ \frac{d^3 x^3 1i}{48 a^3 f} - \frac{d x (-18 c^2 f^2 + c d f 6i + d^2) 1i}{288 a^3 f^3}$$

$$- \frac{d^2 x^2 (-6 c f + d 1i) 1i}{96 a^3 f^2} \Big)$$

$$+ \frac{c^3 x}{8 a^3} + \frac{d^3 x^4}{32 a^3} + \frac{3 c^2 d x^2}{16 a^3} + \frac{c d^2 x^3}{8 a^3}$$

input `int((c + d*x)^3/(a + a*tan(e + f*x)*1i)^3,x)`

output `exp(- e*2i - f*x*2i)*(((d^3*9i + 12*c^3*f^3 - c^2*d*f^2*18i - 18*c*d^2*f)*1i)/(64*a^3*f^4) + (d^3*x^3*3i)/(16*a^3*f) - (d*x*(d^2 - 2*c^2*f^2 + c*d*f*2i)*9i)/(32*a^3*f^3) - (d^2*x^2*(d*1i - 2*c*f)*9i)/(32*a^3*f^2)) + exp(- e*4i - f*x*4i)*(((d^3*9i + 96*c^3*f^3 - c^2*d*f^2*72i - 36*c*d^2*f)*1i)/(1024*a^3*f^4) + (d^3*x^3*3i)/(32*a^3*f) - (d*x*(d^2 - 8*c^2*f^2 + c*d*f*4i)*9i)/(256*a^3*f^3) - (d^2*x^2*(d*1i - 4*c*f)*9i)/(128*a^3*f^2)) + exp(- e*6i - f*x*6i)*(((d^3*1i + 36*c^3*f^3 - c^2*d*f^2*18i - 6*c*d^2*f)*1i)/(1728*a^3*f^4) + (d^3*x^3*1i)/(48*a^3*f) - (d*x*(d^2 - 18*c^2*f^2 + c*d*f*6i)*1i)/(288*a^3*f^3) - (d^2*x^2*(d*1i - 6*c*f)*1i)/(96*a^3*f^2)) + (c^3*x)/(8*a^3) + (d^3*x^4)/(32*a^3) + (3*c^2*d*x^2)/(16*a^3) + (c*d^2*x^3)/(8*a^3)`

Reduce [F]

$$\begin{aligned}
 & \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx \\
 &= \frac{-\left(\int \frac{x^3}{\tan(fx+e)^3 i + 3\tan(fx+e)^2 - 3\tan(fx+e)i - 1} dx\right) d^3 - 3\left(\int \frac{x^2}{\tan(fx+e)^3 i + 3\tan(fx+e)^2 - 3\tan(fx+e)i - 1} dx\right) c d^2 - 3\left(\int \frac{x}{\tan(fx+e)^3 i + 3\tan(fx+e)^2 - 3\tan(fx+e)i - 1} dx\right) c^2}{a^3}
 \end{aligned}$$

input `int((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x)`

output

```
( - int(x**3/(\tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e + f*x)*i - 1
),x)*d**3 - 3*int(x**2/(\tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e +
f*x)*i - 1),x)*c*d**2 - 3*int(x/(\tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3
*tan(e + f*x)*i - 1),x)*c**2*d - int(1/(\tan(e + f*x)**3*i + 3*tan(e + f*x)
)**2 - 3*tan(e + f*x)*i - 1),x)*c**3)/a**3
```

3.30 $\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx$

Optimal result	257
Mathematica [A] (verified)	258
Rubi [A] (verified)	258
Maple [A] (verified)	260
Fricas [A] (verification not implemented)	260
Sympy [A] (verification not implemented)	261
Maxima [F(-2)]	261
Giac [A] (verification not implemented)	262
Mupad [B] (verification not implemented)	263
Reduce [F]	263

Optimal result

Integrand size = 23, antiderivative size = 294

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx = & -\frac{3id^2e^{-2ie-2ifx}}{32a^3f^3} - \frac{3id^2e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}}{864a^3f^3} \\ & + \frac{3de^{-2ie-2ifx}(c+dx)}{16a^3f^2} + \frac{3de^{-4ie-4ifx}(c+dx)}{64a^3f^2} \\ & + \frac{de^{-6ie-6ifx}(c+dx)}{144a^3f^2} + \frac{3ie^{-2ie-2ifx}(c+dx)^2}{16a^3f} \\ & + \frac{3ie^{-4ie-4ifx}(c+dx)^2}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^2}{48a^3f} + \frac{(c+dx)^3}{24a^3d} \end{aligned}$$

output

```
-3/32*I*d^2*exp(-2*I*e-2*I*f*x)/a^3/f^3-3/256*I*d^2*exp(-4*I*e-4*I*f*x)/a^3/f^3-1/864*I*d^2*exp(-6*I*e-6*I*f*x)/a^3/f^3+3/16*d*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^3/f^2+3/64*d*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^3/f^2+1/144*d*exp(-6*I*e-6*I*f*x)*(d*x+c)/a^3/f^2+3/16*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^3/f^2+2*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^3/f^2+1/48*I*exp(-6*I*e-6*I*f*x)*(d*x+c)^2/a^3/f^2+2/24*(d*x+c)^3/a^3/d
```

Mathematica [A] (verified)

Time = 2.23 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.38

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx \\ = \frac{i \sec^3(e + fx) (81(24ic^2f^2 + 4cdf(5 + 12ifx) + d^2(-9i + 20fx + 24if^2x^2)) \cos(e + fx) + 8(18c^2f^2(i +$$

input `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x])^3, x]`

output $((I/6912)*\text{Sec}[e + f*x]^3*(81*((24*I)*c^2*f^2 + 4*c*d*f*(5 + (12*I)*f*x) + d^2*(-9*I + 20*f*x + (24*I)*f^2*x^2))*\text{Cos}[e + f*x] + 8*(18*c^2*f^2*(I + 6*f*x) + 6*c*d*f*(1 + (6*I)*f*x + 18*f^2*x^2) + d^2*(-I + 6*f*x + (18*I)*f^2*x^2 + 36*f^3*x^3))*\text{Cos}[3*(e + f*x)] + 567*d^2*\text{Sin}[e + f*x] + (972*I)*c*d*f*\text{Sin}[e + f*x] - 648*c^2*f^2*\text{Sin}[e + f*x] + (972*I)*d^2*f*x*\text{Sin}[e + f*x] - 1296*c*d*f^2*x*\text{Sin}[e + f*x] - 648*d^2*f^2*x^2*\text{Sin}[e + f*x] - 8*d^2*\text{Sin}[3*(e + f*x)] - (48*I)*c*d*f*\text{Sin}[3*(e + f*x)] + 144*c^2*f^2*\text{Sin}[3*(e + f*x)] - (48*I)*d^2*f*x*\text{Sin}[3*(e + f*x)] + 288*c*d*f^2*x*\text{Sin}[3*(e + f*x)] + (864*I)*c^2*f^3*x*\text{Sin}[3*(e + f*x)] + 144*d^2*f^2*x^2*\text{Sin}[3*(e + f*x)] + (864*I)*c*d*f^3*x^2*\text{Sin}[3*(e + f*x)] + (288*I)*d^2*f^3*x^3*\text{Sin}[3*(e + f*x)]))/((a^3*f^3*(-I + \text{Tan}[e + f*x]))^3)$

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx \\ \downarrow 3042 \\ \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$\begin{aligned}
 & \int \left(\frac{3(c+dx)^2 e^{-2ie-2ifx}}{8a^3} + \frac{3(c+dx)^2 e^{-4ie-4ifx}}{8a^3} + \frac{(c+dx)^2 e^{-6ie-6ifx}}{8a^3} + \frac{(c+dx)^2}{8a^3} \right) dx \\
 & \quad \downarrow \text{4212} \\
 & \frac{3d(c+dx)e^{-2ie-2ifx}}{16a^3f^2} + \frac{3d(c+dx)e^{-4ie-4ifx}}{64a^3f^2} + \frac{d(c+dx)e^{-6ie-6ifx}}{144a^3f^2} + \frac{3i(c+dx)^2e^{-2ie-2ifx}}{16a^3f} + \\
 & \quad \frac{3i(c+dx)^2e^{-4ie-4ifx}}{32a^3f} + \frac{i(c+dx)^2e^{-6ie-6ifx}}{48a^3f} + \frac{(c+dx)^3}{24a^3d} - \frac{3id^2e^{-2ie-2ifx}}{32a^3f^3} - \\
 & \quad \frac{3id^2e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}}{864a^3f^3}
 \end{aligned}$$

input `Int[(c + d*x)^2/(a + I*a*Tan[e + f*x])^3, x]`

output

$$\begin{aligned}
 & (((-3*I)/32)*d^2*E^((-2*I)*e - (2*I)*f*x))/(a^3*f^3) - (((3*I)/256)*d^2*E^((-4*I)*e - (4*I)*f*x))/(a^3*f^3) - ((I/864)*d^2*E^((-6*I)*e - (6*I)*f*x)) \\
 & /(a^3*f^3) + (3*d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))/(16*a^3*f^2) + (3*d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))/(64*a^3*f^2) + (d*E^((-6*I)*e - (6*I)*f*x)*(c + d*x))/(144*a^3*f^2) + ((3*I)/16)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2/(a^3*f) + ((3*I)/32)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2/(a^3*f) + ((I/48)*E^((-6*I)*e - (6*I)*f*x)*(c + d*x)^2/(a^3*f) + (c + d*x)^3/(24*a^3*d)
 \end{aligned}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_.))^m_*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`

Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.81

method	result
risch	$\frac{d^2x^3}{24a^3} + \frac{dcx^2}{8a^3} + \frac{c^2x}{8a^3} + \frac{c^3}{24a^3d} + \frac{3i(2d^2x^2f^2+4cd^2x-2id^2fx+2c^2f^2-2icdf-d^2)e^{-2i(fx+e)}}{32f^3a^3} + \frac{3i(8d^2x^2f^2+16cd^2x^2-4icdf^2-d^4)e^{-2i(fx+e)}}{32f^3a^3}$

input `int((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & \frac{1}{24}a^3d^2x^3 + \frac{1}{8}a^3d^2cx^2 + \frac{1}{8}a^3c^2x^2 + \frac{1}{24}a^3d^3c^3 + \frac{3}{32}I^2(2d^2x^2 - 2f^2 - 2Id^2x^2 - 2f^2x + 4c^2d^2f^2 - 2x^2 - 2I^2c^2d^2f^2 + 2c^2f^2 - d^2)/f^3/a^3 \exp(-2I(fx+e)) \\ & + \frac{3}{256}I^3(8d^2x^2 - 2f^2 - 4Id^2x^2 - 4f^2x + 16c^2d^2f^2 - 2x^2 - 4I^2c^2d^2f^2 + 8c^2f^2 - d^2)/f^3/a^3 \exp(-4I(fx+e)) \\ & + \frac{1}{864}I^4(18d^2x^2 - 2f^2 - 6Id^2x^2 - 6f^2x + 36c^2d^2f^2 - 2x^2 - 6I^2c^2d^2f^2 + 18c^2f^2 - d^2)/f^3/a^3 \exp(-6I(fx+e)) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.75

$$\begin{aligned} & \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx \\ & = \frac{(144i d^2 f^2 x^2 + 144i c^2 f^2 + 48 c d f - 8i d^2 - 48 (-6i c d f^2 - d^2 f)x + 288 (d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x)e^(-6i(fx+e)))}{(a+ia\tan(e+fx))^3} \end{aligned}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output
$$\begin{aligned} & \frac{1}{6912}(144I^2d^2f^2x^2 + 144I^2c^2f^2x^2 + 48c^2d^2f^2 - 8I^2d^2 - 48(-6I^2c^2d^2f^2 - d^2f^2)x + 288(d^2f^3x^3 + 3c^2d^2f^3x^2 + 3c^2f^3x^2)e^{(-6I(fx+e))} \\ & - 648(-2I^2d^2f^2x^2 - 2I^2c^2f^2x^2 - 2c^2d^2f^2 + I^2d^2 + 2(-2I^2c^2d^2f^2 - d^2f^2)x)e^{(-4I(fx+e))} - 81(-8I^2d^2f^2x^2 - 8I^2c^2f^2x^2 - 4c^2d^2f^2 + I^2d^2 + 4(-4I^2c^2d^2f^2 - d^2f^2)x)e^{(-2I(fx+e))} \\ & *e^{(-6I(fx+e))})/(a^3f^3) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 588, normalized size of antiderivative = 2.00

$$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx$$

$$= \left\{ \begin{array}{l} \frac{((147456ia^6c^2f^8e^{6ie} + 294912ia^6cdf^8xe^{6ie} + 49152a^6cdf^7e^{6ie} + 147456ia^6d^2f^8x^2e^{6ie} + 49152a^6d^2f^7xe^{6ie} - 8192ia^6d^2f^6e^{6ie})e^{-6ifx} + (663552a^6c^2d^2f^8x^2e^{6ie} + 331776a^6cd^2f^7xe^{6ie} - 8192a^6d^2f^6e^{6ie})e^{-6ifx})}{(24a^3)} \\ + \frac{x^3 \cdot (3d^2e^{4ie} + 3d^2e^{2ie} + d^2)e^{-6ie}}{8a^3} + \frac{x^2 \cdot (3cde^{4ie} + 3cde^{2ie} + cd)e^{-6ie}}{8a^3} + \frac{x(3c^2e^{4ie} + 3c^2e^{2ie} + c^2)e^{-6ie}}{8a^3} \\ + \frac{c^2x}{8a^3} + \frac{cdx^2}{8a^3} + \frac{d^2x^3}{24a^3} \end{array} \right.$$

input `integrate((d*x+c)**2/(a+I*a*tan(f*x+e))**3,x)`

output

```
Piecewise(((147456*I*a**6*c**2*f**8*exp(6*I*e) + 294912*I*a**6*c*d*f**8*x**2*exp(6*I*e) + 49152*a**6*c*d*f**7*exp(6*I*e) + 147456*I*a**6*d**2*f**8*x**2*exp(6*I*e) + 49152*a**6*d**2*f**7*x*exp(6*I*e) - 8192*I*a**6*d**2*f**6*e*exp(6*I*e))*exp(-6*I*f*x) + (663552*I*a**6*c**2*f**8*exp(8*I*e) + 1327104*I*a**6*c*d*f**8*x*exp(8*I*e) + 331776*a**6*c*d*f**7*exp(8*I*e) + 663552*I*a**6*d**2*f**8*x**2*exp(8*I*e) + 331776*a**6*d**2*f**7*x*exp(8*I*e) - 82944*I*a**6*d**2*f**6*exp(8*I*e))*exp(-4*I*f*x) + (1327104*I*a**6*c**2*f**8*exp(10*I*e) + 2654208*I*a**6*c*d*f**8*x*exp(10*I*e) + 1327104*a**6*c*d*f**7*exp(10*I*e) + 1327104*I*a**6*d**2*f**8*x**2*exp(10*I*e) + 1327104*a**6*d**2*f**7*x*exp(10*I*e) - 663552*I*a**6*d**2*f**6*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(7077888*a**9*f**9), Ne(a**9*f**9*exp(12*I*e), 0)), (x**3*(3*d**2*exp(4*I*e) + 3*d**2*exp(2*I*e) + d**2)*exp(-6*I*e)/(24*a**3) + x**2*(3*c*d*exp(4*I*e) + 3*c*d*exp(2*I*e) + c*d)*exp(-6*I*e)/(8*a**3) + x*(3*c**2*exp(4*I*e) + 3*c**2*exp(2*I*e) + c**2)*exp(-6*I*e)/(8*a**3), True)) + c**2*x/(8*a**3) + c*d*x**2/(8*a**3) + d**2*x**3/(24*a**3)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx \\ = \frac{(288 d^2 f^3 x^3 e^{(6i f x + 6i e)} + 864 c d f^3 x^2 e^{(6i f x + 6i e)} + 864 c^2 f^3 x e^{(6i f x + 6i e)} + 1296 i d^2 f^2 x^2 e^{(4i f x + 4i e)} + 648 i d^2$$

input `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output $\frac{1}{6912} (288*d^2*f^3*x^3*e^{(6*I*f*x + 6*I*e)} + 864*c*d*f^3*x^2*e^{(6*I*f*x + 6*I*e)} + 864*c^2*f^3*x*e^{(6*I*f*x + 6*I*e)} + 1296*I*d^2*f^2*x^2*e^{(4*I*f*x + 4*I*e)} + 648*I*d^2*f^2*x^2*e^{(2*I*f*x + 2*I*e)} + 144*I*d^2*f^2*x^2 + 2592*I*c*d*f^2*x*x*e^{(4*I*f*x + 4*I*e)} + 1296*I*c*d*f^2*x*x*e^{(2*I*f*x + 2*I*e)} + 288*I*c*d*f^2*x + 1296*I*c^2*f^2*x^2*e^{(4*I*f*x + 4*I*e)} + 1296*d^2*f*x*x*e^{(4*I*f*x + 4*I*e)} + 648*I*c^2*f^2*x^2*e^{(2*I*f*x + 2*I*e)} + 324*d^2*f*x*x*e^{(2*I*f*x + 2*I*e)} + 144*I*c^2*f^2*x^2 + 48*d^2*f*x + 1296*c*d*f*x*e^{(4*I*f*x + 4*I*e)} + 324*c*d*f*x*e^{(2*I*f*x + 2*I*e)} + 48*c*d*f - 648*I*d^2*x^2*e^{(4*I*f*x + 4*I*e)} - 81*I*d^2*x^2*e^{(2*I*f*x + 2*I*e)} - 8*I*d^2*x^2*e^{(-6*I*f*x - 6*I*e)})/(a^3*f^3)$

Mupad [B] (verification not implemented)

Time = 8.96 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.89

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx = \frac{c^2 x}{8 a^3} - e^{-e 2i - f x 2i} \left(\frac{(-6 c^2 f^2 + c d f 6i + 3 d^2) 1i}{32 a^3 f^3} - \frac{d^2 x^2 3i}{16 a^3 f} + \frac{d x (-2 c f + d 1i) 3i}{16 a^3 f^2} \right) - e^{-e 4i - f x 4i} \left(\frac{(-24 c^2 f^2 + c d f 12i + 3 d^2) 1i}{256 a^3 f^3} - \frac{d^2 x^2 3i}{32 a^3 f} + \frac{d x (-4 c f + d 1i) 3i}{64 a^3 f^2} \right) - e^{-e 6i - f x 6i} \left(\frac{(-18 c^2 f^2 + c d f 6i + d^2) 1i}{864 a^3 f^3} - \frac{d^2 x^2 1i}{48 a^3 f} + \frac{d x (-6 c f + d 1i) 1i}{144 a^3 f^2} \right) + \frac{d^2 x^3}{24 a^3} + \frac{c d x^2}{8 a^3}$$

input `int((c + d*x)^2/(a + a*tan(e + f*x)*1i)^3,x)`

output
$$(c^2 x)/(8*a^3) - \exp(-e*2i - f*x*2i)*(((3*d^2 - 6*c^2*f^2 + c*d*f*6i)*1i)/(32*a^3*f^3) - (d^2*x^2*3i)/(16*a^3*f) + (d*x*(d*1i - 2*c*f)*3i)/(16*a^3*f^2)) - \exp(-e*4i - f*x*4i)*(((3*d^2 - 24*c^2*f^2 + c*d*f*12i)*1i)/(256*a^3*f^3) - (d^2*x^2*3i)/(32*a^3*f) + (d*x*(d*1i - 4*c*f)*3i)/(64*a^3*f^2)) - \exp(-e*6i - f*x*6i)*(((d^2 - 18*c^2*f^2 + c*d*f*6i)*1i)/(864*a^3*f^3) - (d^2*x^2*1i)/(48*a^3*f) + (d*x*(d*1i - 6*c*f)*1i)/(144*a^3*f^2)) + (d^2*x^3)/(24*a^3) + (c*d*x^2)/(8*a^3)$$

Reduce [F]

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx = \frac{- \left(\int \frac{x^2}{\tan(fx+e)^3 i + 3 \tan(fx+e)^2 - 3 \tan(fx+e) i - 1} dx \right) d^2 - 2 \left(\int \frac{x}{\tan(fx+e)^3 i + 3 \tan(fx+e)^2 - 3 \tan(fx+e) i - 1} dx \right) cd - \left(\int \frac{1}{\tan(fx+e)^3 i + 3 \tan(fx+e)^2 - 3 \tan(fx+e) i - 1} dx \right) c^2 d^3}{a^3}$$

input `int((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x)`

```
output ( - int(x**2/(tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e + f*x)*i - 1
),x)*d**2 - 2*int(x/(tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e + f*x)
)*i - 1),x)*c*d - int(1/(tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e +
f*x)*i - 1),x)*c**2)/a**3
```

3.31 $\int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx$

Optimal result	265
Mathematica [A] (verified)	266
Rubi [A] (verified)	266
Maple [A] (verified)	268
Fricas [A] (verification not implemented)	268
Sympy [A] (verification not implemented)	269
Maxima [F(-2)]	269
Giac [A] (verification not implemented)	270
Mupad [B] (verification not implemented)	270
Reduce [F]	271

Optimal result

Integrand size = 21, antiderivative size = 209

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx = & -\frac{11idx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} + \frac{d}{36f^2(a+ia\tan(e+fx))^3} \\ & + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} + \frac{5d}{96af^2(a+ia\tan(e+fx))^2} \\ & + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\ & + \frac{11d}{96f^2(a^3+ia^3\tan(e+fx))} \\ & + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} \end{aligned}$$

output
$$\begin{aligned} & -11/96*I*d*x/a^3/f-1/16*d*x^2/a^3+1/8*x*(d*x+c)/a^3+1/36*d/f^2/(a+I*a*tan(f*x+e))^3+1/6*I*(d*x+c)/f/(a+I*a*tan(f*x+e))^3+5/96*d/a/f^2/(a+I*a*tan(f*x+e))^2+1/8*I*(d*x+c)/a/f/(a+I*a*tan(f*x+e))^2+11/96*d/f^2/(a^3+I*a^3*tan(f*x+e))+1/8*I*(d*x+c)/f/(a^3+I*a^3*tan(f*x+e)) \end{aligned}$$

Mathematica [A] (verified)

Time = 1.34 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.98

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx \\ = \frac{i \sec^3(e + fx) (27(12icf + d(5 + 12ifx)) \cos(e + fx) + 4(6cf(i + 6fx) + d(1 + 6ifx + 18f^2x^2)) \cos(3(e + fx)))}{(a^3 f^2 (-I + \tan(e + fx))^3)}$$

input `Integrate[(c + d*x)/(a + I*a*Tan[e + f*x])^3, x]`

output
$$\frac{((I/1152)*\text{Sec}[e + f*x]^3*(27*((12*I)*c*f + d*(5 + (12*I)*f*x))*\text{Cos}[e + f*x] + 4*(6*c*f*(I + 6*f*x) + d*(1 + (6*I)*f*x + 18*f^2*x^2))*\text{Cos}[3*(e + f*x)] + (81*I)*d*\text{Sin}[e + f*x] - 108*c*f*\text{Sin}[e + f*x] - 108*d*f*x*\text{Sin}[e + f*x] - (4*I)*d*\text{Sin}[3*(e + f*x)] + 24*c*f*\text{Sin}[3*(e + f*x)] + 24*d*f*x*\text{Sin}[3*(e + f*x)] + (144*I)*c*f^2*x*\text{Sin}[3*(e + f*x)] + (72*I)*d*f^2*x^2*\text{Sin}[3*(e + f*x)]))/((a^3 f^2 (-I + \tan(e + f*x))^3))}{(a^3 f^2 (-I + \tan(e + f*x))^3)}$$

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4213, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx \\ \downarrow \text{3042} \\ \int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx \\ \downarrow \text{4213}$$

$$\begin{aligned}
 & -d \int \left(\frac{x}{8a^3} + \frac{i}{8f(i\tan(e+fx)a^3+a^3)} + \frac{i}{8af(i\tan(e+fx)a+a)^2} + \frac{i}{6f(i\tan(e+fx)a+a)^3} \right) dx + \\
 & \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} + \frac{x(c+dx)}{8a^3} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} \\
 & \quad \downarrow \text{2009} \\
 & \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} + \frac{x(c+dx)}{8a^3} - \\
 d \left(-\frac{11}{96f^2(a^3+ia^3\tan(e+fx))} + \frac{11ix}{96a^3f} + \frac{x^2}{16a^3} - \frac{5}{96af^2(a+ia\tan(e+fx))^2} - \frac{1}{36f^2(a+ia\tan(e+fx))^3} \right) \\
 & \quad \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} + \frac{1}{6f(a+ia\tan(e+fx))^3}
 \end{aligned}$$

input `Int[(c + d*x)/(a + I*a*Tan[e + f*x])^3, x]`

output `(x*(c + d*x))/(8*a^3) + ((I/6)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])^3) + ((I/8)*(c + d*x))/(a*f*(a + I*a*Tan[e + f*x])^2) + ((I/8)*(c + d*x))/(f*(a^3 + I*a^3*Tan[e + f*x])) - d*((((11*I)/96)*x)/(a^3*f) + x^2/(16*a^3) - 1/(36*f^2*(a + I*a*Tan[e + f*x])^3) - 5/(96*a*f^2*(a + I*a*Tan[e + f*x])^2) - 11/(96*f^2*(a^3 + I*a^3*Tan[e + f*x])))`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4213 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_) + (b_.*tan[(e_.) + (f_.*x_)])^(n_), x_Symbol] :> With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Simp[(c + d*x)^m u, x] - Simp[d*m Int[(c + d*x)^(m - 1) u, x], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]`

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.55

method	result	size
risch	$\frac{dx^2}{16a^3} + \frac{cx}{8a^3} + \frac{3i(2dfx+2cf-id)e^{-2i(fx+e)}}{32a^3f^2} + \frac{3i(4dfx+4cf-id)e^{-4i(fx+e)}}{128a^3f^2} + \frac{i(6dfx+6cf-id)e^{-6i(fx+e)}}{288a^3f^2}$	114

input `int((d*x+c)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output $\frac{1/16*d*x^2/a^3+1/8/a^3*c*x+3/32*I*(2*d*f*x-I*d+2*c*f)/a^3/f^2*exp(-2*I*(f*x+e))+3/128*I*(4*d*f*x-I*d+4*c*f)/a^3/f^2*exp(-4*I*(f*x+e))+1/288*I*(6*d*f*x-I*d+6*c*f)/a^3/f^2*exp(-6*I*(f*x+e))}{1152 a^3 f^2}$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.50

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx \\ = \frac{(24i dfx + 24i cf + 72(df^2x^2 + 2cf^2x)e^{(6ifx+6ie)} - 108(-2i dfx - 2i cf - d)e^{(4ifx+4ie)} - 27(-4i dfx - 4icf - d)e^{(2ifx+2ie)} + 4d)e^{(-6ifx-6ie)}}{1152 a^3 f^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output $\frac{1/1152*(24*I*d*f*x + 24*I*c*f + 72*(d*f^2*x^2 + 2*c*f^2*x)*e^{(6*I*f*x + 6*I*e)} - 108*(-2*I*d*f*x - 2*I*c*f - d)*e^{(4*I*f*x + 4*I*e)} - 27*(-4*I*d*f*x - 4*I*c*f - d)*e^{(2*I*f*x + 2*I*e)} + 4*d)*e^{(-6*I*f*x - 6*I*e)})/(a^3*f^2)$

Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.49

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx$$

$$= \begin{cases} \frac{((24576ia^6cf^5e^{6ie} + 24576ia^6df^5xe^{6ie} + 4096a^6df^4e^{6ie})e^{-6ifx} + (110592ia^6cf^5e^{8ie} + 110592ia^6df^5xe^{8ie} + 27648a^6df^4e^{8ie})e^{-4ifx} + (221184a^6df^4e^{8ie})e^{-4ifx} + (221184a^6df^4e^{8ie})e^{-4ifx})}{1179648a^9f^6} \\ \frac{x^2 \cdot (3de^{4ie} + 3de^{2ie} + d)e^{-6ie}}{16a^3} + \frac{x(3ce^{4ie} + 3ce^{2ie} + c)e^{-6ie}}{8a^3} \\ + \frac{cx}{8a^3} + \frac{dx^2}{16a^3} \end{cases}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))**3,x)`

output `Piecewise(((24576*I*a**6*c*f**5*exp(6*I*e) + 24576*I*a**6*d*f**5*x*exp(6*I*e) + 4096*a**6*d*f**4*exp(6*I*e))*exp(-6*I*f*x) + (110592*I*a**6*c*f**5*exp(8*I*e) + 110592*I*a**6*d*f**5*x*exp(8*I*e) + 27648*a**6*d*f**4*exp(8*I*e))*exp(-4*I*f*x) + (221184*I*a**6*c*f**5*exp(10*I*e) + 221184*I*a**6*d*f**5*x*exp(10*I*e) + 110592*a**6*d*f**4*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(1179648*a**9*f**6), Ne(a**9*f**6*exp(12*I*e), 0)), (x**2*(3*d*exp(4*I*e) + 3*d*exp(2*I*e) + d)*exp(-6*I*e)/(16*a**3) + x*(3*c*exp(4*I*e) + 3*c*exp(2*I*e) + c)*exp(-6*I*e)/(8*a**3), True)) + c*x/(8*a**3) + d*x**2/(16*a**3)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))**3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.68

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \frac{(72 df^2 x^2 e^{(6i f x + 6i e)} + 144 c f^2 x e^{(6i f x + 6i e)} + 216i d f x e^{(4i f x + 4i e)} + 108i d f x e^{(2i f x + 2i e)} + 24i d f x + 216i c f^2)}{1152 a^3 f^2}$$

input `integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output $\frac{1}{1152} (72 d f^2 x^2 e^{(6 I f x + 6 I e)} + 144 c f^2 x e^{(6 I f x + 6 I e)} + 216 I d f x e^{(4 I f x + 4 I e)} + 108 I d f x e^{(2 I f x + 2 I e)} + 24 I d f x + 216 I c f^2 e^{(4 I f x + 4 I e)} + 108 I c f^2 e^{(2 I f x + 2 I e)} + 24 I c f + 108 d e^{(4 I f x + 4 I e)} + 27 d e^{(2 I f x + 2 I e)} + 4 d) e^{(-6 I f x - 6 I e)}) / (a^3 f^2)$

Mupad [B] (verification not implemented)

Time = 9.15 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.70

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \frac{d x^2}{16 a^3} - e^{-e 4 i - f x 4 i} \left(\frac{(-12 c f + d 3 i) 1 i}{128 a^3 f^2} - \frac{d x 3 i}{32 a^3 f} \right) - e^{-e 6 i - f x 6 i} \left(\frac{(-6 c f + d 1 i) 1 i}{288 a^3 f^2} - \frac{d x 1 i}{48 a^3 f} \right) - e^{-e 2 i - f x 2 i} \left(\frac{(-6 c f + d 3 i) 1 i}{32 a^3 f^2} - \frac{d x 3 i}{16 a^3 f} \right) + \frac{c x}{8 a^3}$$

input `int((c + d*x)/(a + a*tan(e + f*x)*1i)^3,x)`

output $(d*x^2)/(16*a^3) - \exp(-e*4i - f*x*4i)*(((d*3i - 12*c*f)*1i)/(128*a^3*f^2) - (d*x*3i)/(32*a^3*f)) - \exp(-e*6i - f*x*6i)*(((d*1i - 6*c*f)*1i)/(288*a^3*f^2) - (d*x*1i)/(48*a^3*f)) - \exp(-e*2i - f*x*2i)*(((d*3i - 6*c*f)*1i)/(32*a^3*f^2) - (d*x*3i)/(16*a^3*f)) + (c*x)/(8*a^3)$

Reduce [F]

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx \\ = \frac{- \left(\int \frac{x}{\tan(fx+e)^3 i + 3 \tan(fx+e)^2 - 3 \tan(fx+e)i - 1} dx \right) d - \left(\int \frac{1}{\tan(fx+e)^3 i + 3 \tan(fx+e)^2 - 3 \tan(fx+e)i - 1} dx \right) c}{a^3}$$

input `int((d*x+c)/(a+I*a*tan(f*x+e))^3,x)`

output `(- (int(x/(tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e + f*x)*i - 1), x)*d + int(1/(tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e + f*x)*i - 1),x)*c))/a**3`

3.32 $\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^3} dx$

Optimal result	273
Mathematica [A] (verified)	274
Rubi [A] (verified)	274
Maple [A] (verified)	276
Fricas [A] (verification not implemented)	276
Sympy [F]	277
Maxima [A] (verification not implemented)	277
Giac [A] (verification not implemented)	278
Mupad [F(-1)]	279
Reduce [F]	279

Optimal result

Integrand size = 23, antiderivative size = 449

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx = \frac{3 \cos\left(2e - \frac{2cf}{d}\right) \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\ + \frac{3 \cos\left(4e - \frac{4cf}{d}\right) \text{CosIntegral}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\ + \frac{\cos\left(6e - \frac{6cf}{d}\right) \text{CosIntegral}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} \\ + \frac{\log(c + dx)}{8a^3d} \\ - \frac{i \text{CosIntegral}\left(\frac{6cf}{d} + 6fx\right) \sin\left(6e - \frac{6cf}{d}\right)}{8a^3d} \\ - \frac{3i \text{CosIntegral}\left(\frac{4cf}{d} + 4fx\right) \sin\left(4e - \frac{4cf}{d}\right)}{8a^3d} \\ - \frac{3i \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{8a^3d} \\ - \frac{3i \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\ - \frac{3 \sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\ - \frac{3i \cos\left(4e - \frac{4cf}{d}\right) \text{Si}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\ - \frac{3 \sin\left(4e - \frac{4cf}{d}\right) \text{Si}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\ - \frac{i \cos\left(6e - \frac{6cf}{d}\right) \text{Si}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} \\ - \frac{\sin\left(6e - \frac{6cf}{d}\right) \text{Si}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d}$$

```
output 3/8*cos(-2*e+2*c*f/d)*Ci(2*c*f/d+2*f*x)/a^3/d+3/8*cos(-4*e+4*c*f/d)*Ci(4*c
*f/d+4*f*x)/a^3/d+1/8*cos(-6*e+6*c*f/d)*Ci(6*c*f/d+6*f*x)/a^3/d+1/8*ln(d*x
+c)/a^3/d+1/8*I*Ci(6*c*f/d+6*f*x)*sin(-6*e+6*c*f/d)/a^3/d+3/8*I*Ci(4*c*f/d
+4*f*x)*sin(-4*e+4*c*f/d)/a^3/d+3/8*I*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/
a^3/d-3/8*I*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^3/d+3/8*sin(-2*e+2*c*f/d
)*Si(2*c*f/d+2*f*x)/a^3/d-3/8*I*cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^3/d+
3/8*sin(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^3/d-1/8*I*cos(-6*e+6*c*f/d)*Si(6
*c*f/d+6*f*x)/a^3/d+1/8*sin(-6*e+6*c*f/d)*Si(6*c*f/d+6*f*x)/a^3/d
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.75

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ = \frac{\sec^3(e + fx)(\cos(fx) + i \sin(fx))^3 \left(\cos(3e) \log(f(c + dx)) + i \log(f(c + dx)) \sin(3e) + (\cos(e - \frac{4cf}{d}) - \sec(e - \frac{4cf}{d}) \sin(e - \frac{4cf}{d})) \right)}{(c + dx)^3}$$

input `Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])^3),x]`

output
$$\begin{aligned} & (\operatorname{Sec}[e + f*x]^3 * (\operatorname{Cos}[f*x] + I \operatorname{Sin}[f*x])^3 * (\operatorname{Cos}[3e] \operatorname{Log}[f*(c + d*x)] + I \operatorname{Log}[f*(c + d*x)] \operatorname{Sin}[3e] + (\operatorname{Cos}[e - (4*c*f)/d] - I \operatorname{Sin}[e - (4*c*f)/d]) * (3 * \operatorname{CosIntegral}[(4*f*(c + d*x))/d] + \operatorname{Cos}[2*e - (2*c*f)/d] * \operatorname{CosIntegral}[(6*f*(c + d*x))/d] + 3 * \operatorname{CosIntegral}[(2*f*(c + d*x))/d] * (\operatorname{Cos}[2*e - (2*c*f)/d] + I \operatorname{Sin}[2*e - (2*c*f)/d]) - I \operatorname{CosIntegral}[(6*f*(c + d*x))/d] * \operatorname{Sin}[2*e - (2*c*f)/d] - (3*I) * \operatorname{Cos}[2*e - (2*c*f)/d] * \operatorname{SinIntegral}[(2*f*(c + d*x))/d] + 3 * \operatorname{Sin}[2*e - (2*c*f)/d] * \operatorname{SinIntegral}[(2*f*(c + d*x))/d] - (3*I) * \operatorname{SinIntegral}[(4*f*(c + d*x))/d] - I \operatorname{Cos}[2*e - (2*c*f)/d] * \operatorname{SinIntegral}[(6*f*(c + d*x))/d] - \operatorname{Sin}[2*e - (2*c*f)/d] * \operatorname{SinIntegral}[(6*f*(c + d*x))/d])) / (8*d*(a + I*a*Tan[e + f*x])^3) \end{aligned}$$

Rubi [A] (verified)

Time = 1.94 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ \downarrow 3042 \\ \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx$$

$$\begin{aligned}
 & \downarrow \text{4211} \\
 \int & \left(\frac{i \sin^3(2e + 2fx)}{8a^3(c + dx)} - \frac{3 \sin^2(2e + 2fx)}{8a^3(c + dx)} - \frac{3i \sin(2e + 2fx)}{8a^3(c + dx)} - \frac{3 \sin(2e + 2fx) \sin(4e + 4fx)}{16a^3(c + dx)} - \frac{3i \sin(4e + 4fx)}{8a^3(c + dx)} \right. \\
 & \quad \left. \downarrow \text{2009} \right. \\
 & - \frac{3i \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{8a^3d} - \frac{i \operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \sin\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \\
 & - \frac{3i \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{8a^3d} + \frac{3 \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{8a^3d} + \\
 & - \frac{3 \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{8a^3d} + \frac{\operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \cos\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \\
 & - \frac{3 \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{8a^3d} - \frac{3 \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{8a^3d} - \\
 & - \frac{\sin\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{8a^3d} - \frac{3i \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{8a^3d} - \\
 & - \frac{3i \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{8a^3d} - \frac{i \cos\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{8a^3d} + \frac{\log(c + dx)}{8a^3d}
 \end{aligned}$$

input `Int[1/((c + d*x)*(a + I*a*Tan[e + f*x])^3),x]`

output

$$\begin{aligned}
 & (3*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/(8*a^3*d) + (3*\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x])/(8*a^3*d) + (\operatorname{Cos}[6*e - (6*c*f)/d]*\operatorname{CosIntegral}[(6*c*f)/d + 6*f*x])/(8*a^3*d) + \operatorname{Log}[c + d*x]/(8*a^3*d) - ((I/8)*\operatorname{CosIntegral}[(6*c*f)/d + 6*f*x]*\operatorname{Sin}[6*e - (6*c*f)/d])/(a^3*d) - (((3*I)/8)*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x]*\operatorname{Sin}[4*e - (4*c*f)/d])/(a^3*d) - (((3*I)/8)*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x]*\operatorname{Sin}[2*e - (2*c*f)/d])/(a^3*d) - (((3*I)/8)*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a^3*d) - (3*\operatorname{Sin}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(8*a^3*d) - (((3*I)/8)*\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/(a^3*d) - (3*\operatorname{Si}n[4*e - (4*c*f)/d]*\operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/(8*a^3*d) - ((I/8)*\operatorname{Cos}[6*e - (6*c*f)/d]*\operatorname{SinIntegral}[(6*c*f)/d + 6*f*x])/(a^3*d) - (\operatorname{Sin}[6*e - (6*c*f)/d]*\operatorname{SinIntegral}[(6*c*f)/d + 6*f*x])/(8*a^3*d)
 \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4211 $\text{Int}[(c_.) + (d_.)*(x_.)^m*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^n_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + \text{Cos}[2*e + 2*f*x]/(2*a) + \text{Sin}[2*e + 2*f*x]/(2*b))^(-n), x], x] /; \ \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \& \ \text{EqQ}[a^2 + b^2, 0] \ \& \ \text{ILtQ}[m, 0] \ \& \ \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.36

method	result
risch	$\frac{\ln(dx+c)}{8a^3d} - \frac{e^{\frac{6i(cf-de)}{d}} \exp(\text{Integral}_1(6ifx+6ie+\frac{6i(cf-de)}{d}))}{8a^3d} - \frac{3e^{\frac{4i(cf-de)}{d}} \exp(\text{Integral}_1(4ifx+4ie+\frac{4i(cf-de)}{d}))}{8a^3d} - \frac{3e^{\frac{2i(cf-de)}{d}} \exp(\text{Integral}_1(2ifx+2ie+\frac{2i(cf-de)}{d}))}{8a^3d}$

input $\text{int}(1/(d*x+c)/(a+I*a*tan(f*x+e))^3, x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{8}\ln(d*x+c)/a^3/d - \frac{1}{8}/a^3/d*\exp(6*I*(c*f-d*e)/d)*\text{Ei}(1, 6*I*f*x+6*I*e+6*I*(c*f-d*e)/d) - \frac{3}{8}/a^3/d*\exp(4*I*(c*f-d*e)/d)*\text{Ei}(1, 4*I*f*x+4*I*e+4*I*(c*f-d*e)/d) - \frac{3}{8}/a^3/d*\exp(2*I*(c*f-d*e)/d)*\text{Ei}(1, 2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.26

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ &= \frac{3 \text{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + 3 \text{Ei}\left(-\frac{4(i dfx + i cf)}{d}\right) e^{\left(-\frac{4(i de - i cf)}{d}\right)} + \text{Ei}\left(-\frac{6(i dfx + i cf)}{d}\right) e^{\left(-\frac{6(i de - i cf)}{d}\right)}}{8 a^3 d} \end{aligned}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output
$$\frac{1}{8} \left(3 \operatorname{Ei}(-2 \operatorname{atan}(f x + e) / d) e^{-(-2 \operatorname{atan}(f x + e) - I c f) / d} + 3 \operatorname{Ei}(-4 \operatorname{atan}(f x + e) / d) e^{-(-4 \operatorname{atan}(f x + e) - I c f) / d} + \operatorname{Ei}(-6 \operatorname{atan}(f x + e) / d) e^{-(-6 \operatorname{atan}(f x + e) - I c f) / d} + \log((d x + c) / d) \right) / (a^3 d)$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ &= \frac{i \int \frac{1}{c \tan^3(e+fx)-3ic\tan^2(e+fx)-3c\tan(e+fx)+ic+dx\tan^3(e+fx)-3idx\tan^2(e+fx)-3dx\tan(e+fx)+idx} dx}{a^3} \end{aligned}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**3,x)`

output
$$\begin{aligned} & \operatorname{Integral}\left(\frac{1}{c \tan(e + f x)^3} - \frac{3 i c \tan(e + f x)^2}{a^3} - \frac{3 c \tan(e + f x)}{a^3} + \frac{i c + d x \tan(e + f x)^3 - 3 i d x \tan(e + f x)^2 - 3 d x \tan(e + f x) + i d x}{a^3}, x\right) \end{aligned}$$

Maxima [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.61

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx = \\ & - \frac{3 f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + 3 f \cos\left(-\frac{4(de-cf)}{d}\right) E_1\left(-\frac{4(-i(fx+e)d+i(de-icf))}{d}\right) + f \cos\left(-\frac{6(de-cf)}{d}\right) E_1\left(-\frac{6(-i(fx+e)d+i(de-icf))}{d}\right)}{a^3} \end{aligned}$$

input `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output

$$\begin{aligned} & -\frac{1}{8} \left(3f \cos(-2(d e - c f)/d) \operatorname{exp_integral_e}(1, -2(-I(f x + e)*d + I*d *e - I*c*f)/d) + 3f \cos(-4(d e - c f)/d) \operatorname{exp_integral_e}(1, -4(-I(f x + e)*d + I*d*e - I*c*f)/d) + f \cos(-6(d e - c f)/d) \operatorname{exp_integral_e}(1, -6(-I(f x + e)*d + I*d*e - I*c*f)/d) + 3I*f \operatorname{exp_integral_e}(1, -2(-I(f x + e)*d + I*d*e - I*c*f)/d) * \sin(-2(d e - c f)/d) + 3I*f \operatorname{exp_integral_e}(1, -4(-I(f x + e)*d + I*d*e - I*c*f)/d) * \sin(-4(d e - c f)/d) + I*f \operatorname{exp_integral_e}(1, -6(-I(f x + e)*d + I*d*e - I*c*f)/d) * \sin(-6(d e - c f)/d) - f * \log((f x + e)*d - d*e + c*f) \right) / (a^3 * d*f) \end{aligned}$$

Giac [A] (verification not implemented)

Time = 0.30 (sec), antiderivative size = 810, normalized size of antiderivative = 1.80

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx = \text{Too large to display}$$

input

```
integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
```

output

$$\begin{aligned} & \frac{1}{8} \left(3 \cos(2e)^2 \cos(2c*f/d) \cos(\operatorname{integral}(-2(d*f*x + c*f)/d) + \cos(2e)^3 \log(d*x + c) + 6I \cos(2e) \cos(2c*f/d) \cos(\operatorname{integral}(-2(d*f*x + c*f)/d) * \sin(2e) + 3I \cos(2e)^2 \log(d*x + c) * \sin(2e) - 3 \cos(2c*f/d) \cos(\operatorname{integral}(-2(d*f*x + c*f)/d) * \sin(2e)^2 - 3 \cos(2e) * \log(d*x + c) * \sin(2e)^2 - I \log(d*x + c) * \sin(2e)^3 + 3I \cos(2e)^2 \cos(\operatorname{integral}(-2(d*f*x + c*f)/d) * \sin(2c*f/d) - 6 \cos(2e) \cos(\operatorname{integral}(-2(d*f*x + c*f)/d) * \sin(2e) * \sin(2c*f/d) - 3I \cos(\operatorname{integral}(-2(d*f*x + c*f)/d) * \sin(2e)^2 * \sin(2c*f/d) - 3I \cos(2e)^2 \cos(\operatorname{integral}(2(d*f*x + c*f)/d) * \sin(2e) * \sin(2c*f/d) + 6 \cos(2e) * \cos(2c*f/d) * \sin(2e) * \sin(\operatorname{integral}(2(d*f*x + c*f)/d) * \sin(2e)^2 * \sin(2c*f/d) + 3I \cos(2c*f/d) * \sin(2e)^2 * \sin(\operatorname{integral}(2(d*f*x + c*f)/d) * \sin(2e) * \sin(2c*f/d) * \sin(\operatorname{integral}(2(d*f*x + c*f)/d) * \sin(2e)^2 * \sin(2c*f/d) + 6I \cos(2e) * \sin(2e) * \sin(2c*f/d) * \sin(\operatorname{integral}(2(d*f*x + c*f)/d) * \sin(2e)^2 * \sin(2c*f/d) - 3 \sin(2e)^2 * \sin(2c*f/d) * \sin(\operatorname{integral}(2(d*f*x + c*f)/d) * \sin(2e)^2 * \sin(2c*f/d) + 3 \cos(2e) * \cos(4c*f/d) * \cos(\operatorname{integral}(-4(d*f*x + c*f)/d) * \sin(2e) + 3I \cos(2e) * \cos(\operatorname{integral}(-4(d*f*x + c*f)/d) * \sin(4c*f/d) - 3 \cos(\operatorname{integral}(-4(d*f*x + c*f)/d) * \sin(4c*f/d) * \sin(2e) * \sin(4c*f/d) - 3I \cos(2e) * \cos(4c*f/d) * \sin(\operatorname{integral}(4(d*f*x + c*f)/d) * \sin(2e) * \sin(4c*f/d) + 3 \cos(2e) * \sin(4c*f/d) * \sin(\operatorname{integral}(4(d*f*x + c*f)/d) * \sin(2e) * \sin(4c*f/d) + 3I \sin(2e) * \sin(4c*f/d) * \sin(\operatorname{integral}(4(d*f*x + c*f)/d) * \sin(2e) * \sin(4c*f/d) + \cos(6c*f/d) * \cos(\operatorname{integral}(-6(d*f*x + c*f)/d) * \sin(6c*f/d) + I * \cos(\operatorname{integral}(-6(d*f*x + c*f)/d) * \sin(6c*f/d) - I * \dots) \right) \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^3 (c + dx)} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)),x)`

output `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ &= -\frac{\int \frac{1}{\tan(fx+e)^3 ci + \tan(fx+e)^3 dix + 3 \tan(fx+e)^2 c + 3 \tan(fx+e)^2 dx - 3 \tan(fx+e) ci - 3 \tan(fx+e) dix - c - dx} dx}{a^3} \end{aligned}$$

input `int(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x)`

output `(- int(1/(tan(e + f*x)**3*c*i + tan(e + f*x)**3*d*i*x + 3*tan(e + f*x)**2 *c + 3*tan(e + f*x)**2*d*x - 3*tan(e + f*x)*c*i - 3*tan(e + f*x)*d*i*x - c - d*x),x))/a**3`

3.33
$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx$$

Optimal result	281
Mathematica [A] (verified)	282
Rubi [A] (verified)	283
Maple [A] (verified)	286
Fricas [A] (verification not implemented)	286
Sympy [F]	287
Maxima [A] (verification not implemented)	287
Giac [B] (verification not implemented)	288
Mupad [F(-1)]	289
Reduce [F]	289

Optimal result

Integrand size = 23, antiderivative size = 712

$$\begin{aligned}
 \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = & -\frac{1}{8a^3d(c+dx)} - \frac{9\cos(2e+2fx)}{32a^3d(c+dx)} \\
 & - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} \\
 & - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} - \frac{3\cos(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3if\cos(2e-\frac{2cf}{d})\text{CosIntegral}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
 & - \frac{3if\cos(4e-\frac{4cf}{d})\text{CosIntegral}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
 & - \frac{3if\cos(6e-\frac{6cf}{d})\text{CosIntegral}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
 & - \frac{3f\text{CosIntegral}(\frac{6cf}{d}+6fx)\sin(6e-\frac{6cf}{d})}{4a^3d^2} \\
 & - \frac{3f\text{CosIntegral}(\frac{4cf}{d}+4fx)\sin(4e-\frac{4cf}{d})}{2a^3d^2} \\
 & - \frac{3f\text{CosIntegral}(\frac{2cf}{d}+2fx)\sin(2e-\frac{2cf}{d})}{4a^3d^2} \\
 & + \frac{15i\sin(2e+2fx)}{32a^3d(c+dx)} \\
 & + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} \\
 & + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} + \frac{3i\sin(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3f\cos(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
 & + \frac{3if\sin(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
 & - \frac{3f\cos(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
 & + \frac{3if\sin(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
 & - \frac{3f\cos(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
 & + \frac{3if\sin(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{4a^3d^2}
 \end{aligned}$$

output

$$\begin{aligned}
 & -\frac{1}{8}a^3/d/(d*x+c) - \frac{9}{32}\cos(2*f*x+2*e)/a^3/d/(d*x+c) - \frac{3}{8}\cos(2*f*x+2*e)^2/a^3/d/(d*x+c) \\
 & - \frac{1}{8}\cos(2*f*x+2*e)^3/a^3/d/(d*x+c) - \frac{3}{32}\cos(6*f*x+6*e)/a^3/d/(d*x+c) \\
 & - \frac{3}{2}I*f*\cos(-4*e+4*c*f/d)*Ci(4*c*f/d+4*f*x)/a^3/d^2 - \frac{3}{4}I*f*\sin(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^3/d^2 + \frac{15}{32}I*\sin(2*f*x+2*e)/a^3/d/(d*x+c) \\
 & + \frac{3}{4}f*Ci(6*c*f/d+6*f*x)*\sin(-6*e+6*c*f/d)/a^3/d^2 + \frac{3}{2}f*Ci(4*c*f/d+4*f*x)*\sin(-4*e+4*c*f/d)/a^3/d^2 + \frac{3}{4}f*Ci(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a^3/d^2 \\
 & + \frac{3}{8}I*\sin(4*f*x+4*e)/a^3/d/(d*x+c) + \frac{3}{8}\sin(2*f*x+2*e)^2/a^3/d/(d*x+c) - \frac{3}{4}I*f*\sin(-6*e+6*c*f/d)*Si(6*c*f/d+6*f*x)/a^3/d^2 + \frac{3}{32}I*\sin(6*f*x+6*e)/a^3/d/(d*x+c) \\
 & - \frac{1}{8}I*\sin(2*f*x+2*e)^3/a^3/d/(d*x+c) - \frac{3}{4}f*\cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^3/d^2 - \frac{3}{4}I*f*\cos(-6*e+6*c*f/d)*Ci(6*c*f/d+6*f*x)/a^3/d^2 \\
 & - \frac{3}{2}f*\cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^3/d^2 - \frac{3}{2}I*f*\sin(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^3/d^2 - \frac{3}{4}f*\cos(-6*e+6*c*f/d)*Si(6*c*f/d+6*f*x)/a^3/d^2 \\
 & - \frac{3}{4}I*f*\cos(-2*e+2*c*f/d)*Ci(2*c*f/d+2*f*x)/a^3/d^2
 \end{aligned}$$

Mathematica [A] (verified)

Time = 2.55 (sec), antiderivative size = 833, normalized size of antiderivative = 1.17

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx = \text{Too large to display}$$

input `Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^3),x]`

output

$$\begin{aligned}
 & (\operatorname{Sec}[e + f*x]^3 * ((-I)*\operatorname{Cos}[(3*c*f)/d] + \operatorname{Sin}[(3*c*f)/d]) * (3*d*\operatorname{Cos}[e + f*((-3*c)/d + x)] + d*\operatorname{Cos}[3*(e + f*(-(c/d) + x))] + d*\operatorname{Cos}[3*(e + f*(c/d + x))] + \\
 & 3*d*\operatorname{Cos}[e + f*((3*c)/d + x)] + (6*I)*c*f*\operatorname{Cos}[3*e - (3*f*(c + d*x))/d]*\operatorname{Cos}[\operatorname{Integral}[(6*f*(c + d*x))/d] + (6*I)*d*f*x*\operatorname{Cos}[3*e - (3*f*(c + d*x))/d]*\operatorname{Cos}[\operatorname{Integral}[(6*f*(c + d*x))/d] + (6*I)*f*(c + d*x)*\operatorname{Cos}[\operatorname{Integral}[(2*f*(c + d*x))/d]*(\operatorname{Cos}[e - (c*f)/d + 3*f*x] + I*\operatorname{Sin}[e - (c*f)/d + 3*f*x]) + (3*I)*d*\operatorname{Sin}[e + f*((-3*c)/d + x)] + I*d*\operatorname{Sin}[3*(e + f*(-(c/d) + x))] - I*d*\operatorname{Sin}[3*(e + f*(c/d + x))] - (3*I)*d*\operatorname{Sin}[e + f*((3*c)/d + x)] + 6*c*f*\operatorname{Cos}[\operatorname{Integral}[(6*f*(c + d*x))/d]*\operatorname{Sin}[3*e - (3*f*(c + d*x))/d] + 6*d*f*x*\operatorname{Cos}[\operatorname{Integral}[(6*f*(c + d*x))/d]*\operatorname{Sin}[3*e - (3*f*(c + d*x))/d] + 12*f*(c + d*x)*\operatorname{Cos}[\operatorname{Integral}[(4*f*(c + d*x))/d]*(I*\operatorname{Cos}[e - (f*(c + 3*d*x))/d] + \operatorname{Sin}[e - (f*(c + 3*d*x))/d]) + 6*c*f*\operatorname{Cos}[e - (c*f)/d + 3*f*x]*\operatorname{Sin}[\operatorname{Integral}[(2*f*(c + d*x))/d] + 6*d*f*x*\operatorname{Cos}[e - (c*f)/d + 3*f*x]*\operatorname{Sin}[\operatorname{Integral}[(2*f*(c + d*x))/d] + (6*I)*c*f*\operatorname{Sin}[e - (c*f)/d + 3*f*x]*\operatorname{Sin}[\operatorname{Integral}[(2*f*(c + d*x))/d] + (6*I)*d*f*x*\operatorname{Sin}[e - (c*f)/d + 3*f*x]*\operatorname{Sin}[\operatorname{Integral}[(2*f*(c + d*x))/d] + 12*c*f*\operatorname{Cos}[e - (f*(c + 3*d*x))/d]*\operatorname{Sin}[\operatorname{Integral}[(4*f*(c + d*x))/d] + 12*d*f*x*\operatorname{Cos}[e - (f*(c + 3*d*x))/d]*\operatorname{Sin}[\operatorname{Integral}[(4*f*(c + d*x))/d] - (12*I)*c*f*\operatorname{Sin}[e - (f*(c + 3*d*x))/d]*\operatorname{Sin}[\operatorname{Integral}[(4*f*(c + d*x))/d] - (12*I)*d*f*x*\operatorname{Sin}[e - (f*(c + 3*d*x))/d]*\operatorname{Sin}[\operatorname{Integral}[(4*f*(c + d*x))/d] + 6*c*f*\operatorname{Cos}[3*e - (3*f*(c + d*x))/d]*\operatorname{Sin}[\operatorname{Integral}[(6*f*(c + d*x))/d] + 6*d*f*x*\operatorname{Cos}[3*e - (3*f*(c + d*x))/d]*\operatorname{Sin}[\operatorname{Integral}...
 \end{aligned}$$

Rubi [A] (verified)

Time = 2.05 (sec), antiderivative size = 712, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {3042, 4211, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \textcolor{blue}{4211}
 \end{aligned}$$

$$\begin{aligned}
 & \int \left(\frac{i \sin^3(2e + 2fx)}{8a^3(c + dx)^2} - \frac{3 \sin^2(2e + 2fx)}{8a^3(c + dx)^2} - \frac{3i \sin(2e + 2fx)}{8a^3(c + dx)^2} - \frac{3 \sin(2e + 2fx) \sin(4e + 4fx)}{16a^3(c + dx)^2} - \frac{3i \sin(4e + 4fx)}{8a^3(c + dx)^2} \right. \\
 & \quad \left. \downarrow 2009 \right) \\
 & - \frac{3f \operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \sin\left(6e - \frac{6cf}{d}\right)}{4a^3 d^2} - \frac{3f \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{2a^3 d^2} - \\
 & - \frac{3f \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{4a^3 d^2} - \frac{3if \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{4a^3 d^2} - \\
 & - \frac{3if \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{4a^3 d^2} - \frac{3if \operatorname{CosIntegral}\left(6xf + \frac{6cf}{d}\right) \cos\left(6e - \frac{6cf}{d}\right)}{4a^3 d^2} + \\
 & + \frac{3if \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{4a^3 d^2} + \frac{3if \sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{2a^3 d^2} + \\
 & + \frac{3if \sin\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{4a^3 d^2} - \frac{3f \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{4a^3 d^2} - \\
 & - \frac{3f \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{2a^3 d^2} - \frac{3f \cos\left(6e - \frac{6cf}{d}\right) \operatorname{Si}\left(6xf + \frac{6cf}{d}\right)}{4a^3 d^2} - \frac{i \sin^3(2e + 2fx)}{8a^3 d(c + dx)} + \\
 & + \frac{3 \sin^2(2e + 2fx)}{8a^3 d(c + dx)} + \frac{15i \sin(2e + 2fx)}{32a^3 d(c + dx)} + \frac{3i \sin(4e + 4fx)}{8a^3 d(c + dx)} + \frac{3i \sin(6e + 6fx)}{32a^3 d(c + dx)} - \\
 & - \frac{\cos^3(2e + 2fx)}{8a^3 d(c + dx)} - \frac{3 \cos^2(2e + 2fx)}{8a^3 d(c + dx)} - \frac{9 \cos(2e + 2fx)}{32a^3 d(c + dx)} - \frac{3 \cos(6e + 6fx)}{32a^3 d(c + dx)} - \frac{1}{8a^3 d(c + dx)}
 \end{aligned}$$

input Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^3),x]

output

$$\begin{aligned}
 & -\frac{1}{8} \cdot \frac{1}{(a^3 d (c + d x))} - \frac{(9 \cos[2e + 2f x])}{(32 a^3 d (c + d x))} - \frac{(3 \cos[2e + 2f x]^2)}{(8 a^3 d (c + d x))} - \frac{\cos[2e + 2f x]^3}{(8 a^3 d (c + d x))} - \frac{(3 \cos[6e + 6f x])}{(32 a^3 d (c + d x))} - \frac{((3I)/4) f \cos[2e]}{(2 c f)/d} \cdot \text{CosIntegral}[(2 c f)/d + 2 f x]/(a^3 d^2) - \frac{((3I)/2) f \cos[4e]}{(4 c f)/d} \cdot \text{CosIntegral}[(4 c f)/d + 4 f x]/(a^3 d^2) - \frac{((3I)/4) f \cos[6e - (6 c f)/d]}{(6 c f)/d} \cdot \text{CosIntegral}[(6 c f)/d + 6 f x]/(a^3 d^2) - \frac{(3 f \cos[\text{integral}[(6 c f)/d + 6 f x] * \sin[6e - (6 c f)/d]])}{(4 a^3 d^2)} - \frac{(3 f \cos[\text{integral}[(4 c f)/d + 4 f x] * \sin[4e - (4 c f)/d]])}{(2 a^3 d^2)} - \frac{(3 f \cos[\text{integral}[(2 c f)/d + 2 f x] * \sin[2e - (2 c f)/d]])}{(4 a^3 d^2)} + \frac{((15I)/32) \sin[2e + 2 f x]}{(a^3 d (c + d x))} + \frac{(3 \sin[2e + 2 f x]^2)}{(8 a^3 d (c + d x))} + \frac{((3I)/8) \sin[4e + 4 f x]}{(a^3 d (c + d x))} + \frac{((3I)/32) \sin[6e + 6 f x]}{(a^3 d (c + d x))} - \frac{(3 f \cos[2e - (2 c f)/d] * \text{SinIntegral}[(2 c f)/d + 2 f x])}{(4 a^3 d^2)} + \frac{((3I)/4) f \sin[2e - (2 c f)/d] * \text{SinIntegral}[(2 c f)/d + 2 f x]}{(a^3 d^2)} - \frac{(3 f \cos[4e - (4 c f)/d] * \text{SinIntegral}[(4 c f)/d + 4 f x])}{(2 a^3 d^2)} + \frac{((3I)/2) f \sin[4e - (4 c f)/d] * \text{SinIntegral}[(4 c f)/d + 4 f x]}{(a^3 d^2)} - \frac{(3 f \cos[6e - (6 c f)/d] * \text{SinIntegral}[(6 c f)/d + 6 f x])}{(4 a^3 d^2)} + \frac{((3I)/4) f \sin[6e - (6 c f)/d] * \text{SinIntegral}[(6 c f)/d + 6 f x]}{(a^3 d^2)}
 \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4211 $\text{Int}[(c_.) + (d_.)*(x_.)^m_*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^n_, x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d x)^m, (1/(2 a) + \cos[2 e + 2 f x]/(2 a) + \sin[2 e + 2 f x]/(2 b))^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[m, 0] \&& \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.36

method	result
risch	$-\frac{1}{8a^3d(dx+c)} - \frac{f e^{-6i(fx+e)}}{8a^3(df+cf)d} + \frac{3if e^{\frac{6i(cf-de)}{d}} \expIntegral_1(6ifx+6ie+\frac{6i(cf-de)}{d})}{4a^3d^2} - \frac{3f e^{-4i(fx+e)}}{8a^3(df+cf)d} + \frac{3if e^{\frac{4i(cf-de)}{d}} e^{\frac{6i(cf-de)}{d}} \expIntegral_1(6ifx+6ie+\frac{4i(cf-de)}{d})}{4a^3d^2}$

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -\frac{1}{8}/a^3/d/(d*x+c)-\frac{1}{8}/a^3/f*\exp(-6*I*(f*x+e))/(d*f*x+c*f)/d+3/4*I/a^3/f/d \\ & \sim 2*\exp(6*I*(c*f-d*e)/d)*Ei(1,6*I*f*x+6*I*e+6*I*(c*f-d*e)/d)-3/8/a^3/f*\exp(-4*I*(f*x+e))/(d*f*x+c*f)/d+3/2*I/a^3/f/d^2*\exp(4*I*(c*f-d*e)/d)*Ei(1,4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-3/8/a^3/f*\exp(-2*I*(f*x+e))/(d*f*x+c*f)/d+3/4*I/a^3/f/d^2*\exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.28

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = \frac{-\left(\left(6(i dfx+i cf)\text{Ei}\left(-\frac{2(i dfx+i cf)}{d}\right)e^{\left(-\frac{2(i de-i cf)}{d}\right)}+12(i dfx+i cf)\text{Ei}\left(-\frac{4(i dfx+i cf)}{d}\right)e^{\left(-\frac{4(i de-i cf)}{d}\right)}\right)+d\right)e^{(6*I*f*x+6*I*e)}+3*d*e^{(4*I*f*x+4*I*e)}+3*d*e^{(2*I*f*x+2*I*e)}+d\right)e^{(-6*I*f*x-6*I*e)/(a^3*d^3*x+a^3*c*d^2)}$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output
$$\begin{aligned} & -\frac{1}{8}*((6*(I*d*f*x+I*c*f))*\text{Ei}(-2*(I*d*f*x+I*c*f)/d)*e^{(-2*(I*d*e-I*c*f)/d)}+12*(I*d*f*x+I*c*f)*\text{Ei}(-4*(I*d*f*x+I*c*f)/d)*e^{(-4*(I*d*e-I*c*f)/d)}+6*(I*d*f*x+I*c*f)*\text{Ei}(-6*(I*d*f*x+I*c*f)/d)*e^{(-6*(I*d*e-I*c*f)/d)}+d)*e^{(6*I*f*x+6*I*e)}+3*d*e^{(4*I*f*x+4*I*e)}+3*d*e^{(2*I*f*x+2*I*e)}+d)*e^{(-6*I*f*x-6*I*e)/(a^3*d^3*x+a^3*c*d^2)}) \end{aligned}$$

Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx \\ = \frac{i \int \frac{1}{c^2\tan^3(e+fx)-3ic^2\tan^2(e+fx)-3c^2\tan(e+fx)+ic^2+2cdx\tan^3(e+fx)-6icdx\tan^2(e+fx)-6cdx\tan(e+fx)+2icdx+d^2x^2\tan^3(e+fx)}}{a^3}$$

input `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e))**3,x)`

output `I*Integral(1/(c**2*tan(e + f*x)**3 - 3*I*c**2*tan(e + f*x)**2 - 3*c**2*tan(e + f*x) + I*c**2 + 2*c*d*x*tan(e + f*x)**3 - 6*I*c*d*x*tan(e + f*x)**2 - 6*c*d*x*tan(e + f*x) + 2*I*c*d*x + d**2*x**2*tan(e + f*x)**3 - 3*I*d**2*x**2*tan(e + f*x)**2 - 3*d**2*x**2*tan(e + f*x) + I*d**2*x**2), x)/a**3)`

Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.42

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = \\ -\frac{3 f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + 3 f^2 \cos\left(-\frac{4(de-cf)}{d}\right) E_2\left(-\frac{4(-i(fx+e)d+i(de-icf))}{d}\right) +}{$$

input `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `-1/8*(3*f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*f^2*cos(-4*(d*e - c*f)/d)*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f^2*cos(-6*(d*e - c*f)/d)*exp_integral_e(2, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + 3*I*f^2*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + I*f^2*exp_integral_e(2, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-6*(d*e - c*f)/d) + f^2)/(((f*x + e)*a^3*d^2 - a^3*d^2*e + a^3*c*d*f)*f)`

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2915 vs. $2(648) = 1296$.

Time = 28.91 (sec) , antiderivative size = 2915, normalized size of antiderivative = 4.09

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = \text{Too large to display}$$

```
input integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
```

```

output
1/8*(-6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 12*I*d*e*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*c*f^3*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 6*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 6*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c+dx)^2(a+i a \tan(e+f x))^3} dx = \int \frac{1}{(a+a \tan(e+f x) 1i)^3(c+d x)^2} dx$$

input `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)^2),x)`

output `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)^2), x)`

Reduce [F]

$$\int \frac{1}{(c+dx)^2(a+i a \tan(e+f x))^3} dx =$$

$$-\frac{\int \frac{1}{\tan(f x+e)^3 c^2 i+2 \tan(f x+e)^3 c d i x+\tan(f x+e)^3 d^2 i x^2+3 \tan(f x+e)^2 c^2+6 \tan(f x+e)^2 c d x+3 \tan(f x+e)^2 d^2 x^2-3 \tan(f x+e) c^2 i-6 \tan(f x+e) c d x} dx}{a^3}$$

input `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x)`

output `(- int(1/(\tan(e + f*x)**3*c**2*i + 2*tan(e + f*x)**3*c*d*i*x + tan(e + f*x)**3*d**2*i*x**2 + 3*tan(e + f*x)**2*c**2 + 6*tan(e + f*x)**2*c*d*x + 3*tan(e + f*x)**2*d**2*x**2 - 3*tan(e + f*x)*c**2*i - 6*tan(e + f*x)*c*d*i*x - 3*tan(e + f*x)*d**2*i*x**2 - c**2 - 2*c*d*x - d**2*x**2),x))/a**3`

3.34 $\int (c + dx)^m (a + ia \tan(e + fx))^2 dx$

Optimal result	290
Mathematica [N/A]	290
Rubi [N/A]	291
Maple [N/A]	292
Fricas [N/A]	292
Sympy [N/A]	292
Maxima [N/A]	293
Giac [N/A]	294
Mupad [N/A]	294
Reduce [N/A]	294

Optimal result

Integrand size = 23, antiderivative size = 23

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \text{Int}((c + dx)^m (a + ia \tan(e + fx))^2, x)$$

output `Defer(Int)((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)`

Mathematica [N/A]

Not integrable

Time = 40.38 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

input `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2,x]`

output `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2, x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c + dx)^m (a + ia \tan(e + fx))^2 dx \\ & \quad \downarrow \text{3042} \\ & \int (c + dx)^m (a + ia \tan(e + fx))^2 dx \\ & \quad \downarrow \text{4223} \\ & \int (c + dx)^m (a + ia \tan(e + fx))^2 dx \end{aligned}$$

input `Int[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int (dx + c)^m (a + ia \tan(fx + e))^2 dx$$

input `int((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)`

output `int((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 115, normalized size of antiderivative = 5.00

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (i a \tan(fx + e) + a)^2 (dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output `(-2*I*(d*x + c)^m*a^2 + (f*e^(2*I*f*x + 2*I*e) + f)*integral(-2*(-I*a^2*d*m - 2*(a^2*d*f*x + a^2*c*f)*e^(2*I*f*x + 2*I*e))*(d*x + c)^m/(d*f*x + c*f + (d*f*x + c*f)*e^(2*I*f*x + 2*I*e)), x))/(f*e^(2*I*f*x + 2*I*e) + f)`

Sympy [N/A]

Not integrable

Time = 3.42 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.22

$$\begin{aligned} \int (c + dx)^m (a + ia \tan(e + fx))^2 dx &= -a^2 \left(\int (c + dx)^m \tan^2(e + fx) dx \right. \\ &\quad + \int (-2i(c + dx)^m \tan(e + fx)) dx \\ &\quad \left. + \int (-(c + dx)^m) dx \right) \end{aligned}$$

input `integrate((d*x+c)**m*(a+I*a*tan(f*x+e))**2,x)`

output `-a**2*(Integral((c + d*x)**m*tan(e + f*x)**2, x) + Integral(-2*I*(c + d*x)**m*tan(e + f*x), x) + Integral(-(c + d*x)**m, x))`

Maxima [N/A]

Not integrable

Time = 0.84 (sec) , antiderivative size = 520, normalized size of antiderivative = 22.61

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (ia \tan(fx + e) + a)^2 (dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `(d*x + c)^(m + 1)*a^2/(d*(m + 1)) + integrate((3*(d*x + c)^m*a^2*cos(4*f*x + 4*e)^2 - 4*(d*x + c)^m*a^2*cos(2*f*x + 2*e)^2 + 3*(d*x + c)^m*a^2*sin(4*f*x + 4*e)^2 + 4*(d*x + c)^m*a^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) - 4*(d*x + c)^m*a^2*sin(2*f*x + 2*e)^2 - 4*(d*x + c)^m*a^2*cos(2*f*x + 2*e) - (d*x + c)^m*a^2 + 2*(2*(d*x + c)^m*a^2*cos(2*f*x + 2*e) + (d*x + c)^m*a^2)*cos(4*f*x + 4*e))/(2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1), x) + I*integrate(-4*(2*(d*x + c)^m*a^2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e) - (2*(d*x + c)^m*a^2*cos(2*f*x + 2*e) + (d*x + c)^m*a^2)*sin(4*f*x + 4*e))/(2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1), x)`

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (i a \tan(fx + e) + a)^2 (dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((I*a*tan(f*x + e) + a)^2*(d*x + c)^m, x)`

Mupad [N/A]

Not integrable

Time = 8.61 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (a + a \tan(e + f x) 1i)^2 (c + d x)^m dx$$

input `int((a + a*tan(e + f*x)*1i)^2*(c + d*x)^m,x)`

output `int((a + a*tan(e + f*x)*1i)^2*(c + d*x)^m, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 252, normalized size of antiderivative = 10.96

$$\begin{aligned} & \int (c + dx)^m (a + ia \tan(e + fx))^2 dx \\ &= \frac{a^2 \left(- (dx + c)^m \tan(fx + e) dm - (dx + c)^m \tan(fx + e) d + 2(dx + c)^m cf + 2(dx + c)^m dfx + 2 \left(\int \underline{d} \right. \right.}{\left. \left. \right)} \end{aligned}$$

input $\int ((d*x + c)^m * (a + I*a*tan(f*x + e))^2) dx$

output
$$\begin{aligned} & (a^{**2} * (- (c + d*x)**m * \tan(e + f*x) * d*m - (c + d*x)**m * \tan(e + f*x) * d + 2 * \\ & (c + d*x)**m * c * f + 2 * (c + d*x)**m * d * f * x + 2 * \int (((c + d*x)**m * \tan(e + f*x) * x) / (c + d*x), x) * d^{**2} * f * i * m + 2 * \int (((c + d*x)**m * \tan(e + f*x) * x) / (c + d*x), x) * d^{**2} * f * i + 2 * \int (((c + d*x)**m * \tan(e + f*x)) / (c + d*x), x) * c * d * f * i * m + \\ & 2 * \int (((c + d*x)**m * \tan(e + f*x)) / (c + d*x), x) * c * d * f * i + \int (((c + d*x)**m * \tan(e + f*x)) / (c + d*x), x) * d^{**2} * m^{**2} + \int (((c + d*x)**m * \tan(e + f*x)) / (c + d*x), x) * d^{**2} * m)) / (d * f * (m + 1)) \end{aligned}$$

3.35 $\int (c + dx)^m (a + ia \tan(e + fx)) dx$

Optimal result	296
Mathematica [N/A]	296
Rubi [N/A]	297
Maple [N/A]	298
Fricas [N/A]	298
Sympy [N/A]	298
Maxima [N/A]	299
Giac [N/A]	299
Mupad [N/A]	300
Reduce [N/A]	300

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \text{Int}((c + dx)^m (a + ia \tan(e + fx)), x)$$

output `Defer(Int)((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

Mathematica [N/A]

Not integrable

Time = 8.80 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (c + dx)^m (a + ia \tan(e + fx)) dx$$

input `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x]),x]`

output `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x]), x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c + dx)^m (a + ia \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (c + dx)^m (a + ia \tan(e + fx)) dx \\ & \quad \downarrow \text{4223} \\ & \int (c + dx)^m (a + ia \tan(e + fx)) dx \end{aligned}$$

input `Int[(c + d*x)^m*(a + I*a*Tan[e + f*x]),x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^m*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n, x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int (dx + c)^m (a + ia \tan(fx + e)) dx$$

input `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

output `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output `integral(2*(d*x + c)^m*a*e^(2*I*f*x + 2*I*e)/(e^(2*I*f*x + 2*I*e) + 1), x)`

Sympy [N/A]

Not integrable

Time = 2.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\begin{aligned} \int (c + dx)^m (a + ia \tan(e + fx)) dx &= ia \left(\int (-i(c + dx)^m) dx \right. \\ &\quad \left. + \int (c + dx)^m \tan(e + fx) dx \right) \end{aligned}$$

input `integrate((d*x+c)**m*(a+I*a*tan(f*x+e)),x)`

output $I*a*(\text{Integral}(-I*(c + d*x)^m, x) + \text{Integral}((c + d*x)^m * \tan(e + f*x), x))$

Maxima [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 3.76

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output $2*I*a*\text{integrate}((d*x + c)^m * \sin(2*f*x + 2*e) / (\cos(2*f*x + 2*e)^2 + \sin(2*f*x + 2*e)^2 + 2*\cos(2*f*x + 2*e) + 1), x) + (d*x + c)^{m+1}*a/(d*(m+1))$

Giac [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

input `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `integrate((I*a*tan(f*x + e) + a)*(d*x + c)^m, x)`

Mupad [N/A]

Not integrable

Time = 8.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (a + a \tan(e + fx) 1i) (c + dx)^m dx$$

input `int((a + a*tan(e + f*x)*1i)*(c + d*x)^m,x)`

output `int((a + a*tan(e + f*x)*1i)*(c + d*x)^m, x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 69, normalized size of antiderivative = 3.29

$$\begin{aligned} & \int (c + dx)^m (a + ia \tan(e + fx)) dx \\ &= \frac{a((dx + c)^m c + (dx + c)^m dx + (\int (dx + c)^m \tan(fx + e) dx) dim + (\int (dx + c)^m \tan(fx + e) dx) di)}{d(m + 1)} \end{aligned}$$

input `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

output `(a*((c + d*x)**m*c + (c + d*x)**m*d*x + int((c + d*x)**m*tan(e + f*x),x)*d *i*m + int((c + d*x)**m*tan(e + f*x),x)*d*i))/(d*(m + 1))`

3.36 $\int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx$

Optimal result	301
Mathematica [A] (verified)	301
Rubi [A] (verified)	302
Maple [F]	303
Fricas [A] (verification not implemented)	303
Sympy [F]	304
Maxima [F]	304
Giac [F]	305
Mupad [F(-1)]	305
Reduce [F]	305

Optimal result

Integrand size = 23, antiderivative size = 98

$$\begin{aligned} & \int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx \\ &= \frac{(c+dx)^{1+m}}{2ad(1+m)} + \frac{i2^{-2-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{af} \end{aligned}$$

output
$$\frac{1/2*(d*x+c)^(1+m)/a/d/(1+m)+I*2^(-2-m)*(d*x+c)^m*GAMMA(1+m,2*I*f*(d*x+c)/d)}{a/exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)}$$

Mathematica [A] (verified)

Time = 1.59 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.36

$$\begin{aligned} & \int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx \\ &= \frac{e^{-ie}(c+dx)^m \left(\frac{2e^{2ie}f(c+dx)}{d(1+m)} + i2^{-m}e^{\frac{2icf}{d}} \left(\frac{if(c+dx)}{d} \right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right) \right) \sec(e+fx)(\cos(fx) + i \sin(fx))}{4f(a+ia\tan(e+fx))} \end{aligned}$$

input
$$\text{Integrate}[(c + d*x)^m/(a + I*a*Tan[e + f*x]), x]$$

output

$$\frac{((c + dx)^m ((2 E^((2 I) e) f (c + dx)) / (d (1 + m)) + (I E^(((2 I) c f) / d) \text{Gamma}[1 + m, ((2 I) f (c + dx)) / d]) / (2^m ((I f (c + dx)) / d)^m) \text{Sec}[e + f x] (\cos[f x] + I \sin[f x])) / (4 E^((I e) f (a + I a \tan[e + f x])))}{}$$

Rubi [A] (verified)

Time = 0.33 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4210, 2612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + dx)^m}{a + i a \tan(e + fx)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{(c + dx)^m}{a + i a \tan(e + fx)} dx \\
 & \quad \downarrow \textcolor{blue}{4210} \\
 & \frac{(c + dx)^{m+1}}{2ad(m+1)} + \frac{\int e^{-2i(e+fx)} (c + dx)^m dx}{2a} \\
 & \quad \downarrow \textcolor{blue}{2612} \\
 & \frac{(c + dx)^{m+1}}{2ad(m+1)} + \frac{i 2^{-m-2} e^{-2i\left(e-\frac{cf}{d}\right)} (c + dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2if(c+dx)}{d}\right)}{af}
 \end{aligned}$$

input

$$\text{Int}[(c + dx)^m / (a + I a \tan[e + f x]), x]$$

output

$$\frac{(c + dx)^{(1 + m) / (2 * a * d * (1 + m))} + (I * 2^{(-2 - m)} * (c + dx)^m * \text{Gamma}[1 + m, ((2 I) * f * (c + dx)) / d]) / (a * E^((2 I) * (e - (c * f) / d)) * f * ((I * f * (c + dx)) / d)^m)}{}$$

Definitions of rubi rules used

rule 2612 $\text{Int}[(F_{_})^((g_{_})*(e_{_}) + (f_{_})*(x_{_}))*((c_{_}) + (d_{_})*(x_{_}))^{(m_{_})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(-F^((g*(e - c*(f/d)))))*((c + d*x)^{\text{FracPart}[m]}/(d*((-f)*g*(\text{Log}[F]/d))^{(\text{IntPart}[m] + 1)*((-f)*g*\text{Log}[F]*((c + d*x)/d))^{\text{FracPart}[m]}})*\text{Gamma}[m + 1, ((-f)*g*(\text{Log}[F]/d))*(c + d*x)], x] /; \text{FreeQ}[\{F, c, d, e, f, g, m\}, x] \&& !\text{IntegerQ}[m]$

rule 3042 $\text{Int}[u_{_}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4210 $\text{Int}[((c_{_}) + (d_{_})*(x_{_}))^{(m_{_})}/((a_{_}) + (b_{_})*\tan[(e_{_}) + (f_{_})*(x_{_})]), x_{\text{Symbol}}] \rightarrow \text{Simp}[(c + d*x)^{(m + 1)}/(2*a*d*(m + 1)), x] + \text{Simp}[1/(2*a) \text{Int}[(c + d*x)^m E^{(2*(a/b)*(e + f*x))}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \& \text{EqQ}[a^2 + b^2, 0] \&& !\text{IntegerQ}[m]$

Maple [F]

$$\int \frac{(dx + c)^m}{a + ia \tan(fx + e)} dx$$

input $\text{int}((d*x+c)^m/(a+I*a*tan(f*x+e)), x)$

output $\text{int}((d*x+c)^m/(a+I*a*tan(f*x+e)), x)$

Fricas [A] (verification not implemented)

Time = 0.07 (sec), antiderivative size = 85, normalized size of antiderivative = 0.87

$$\begin{aligned} & \int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx \\ &= \frac{(i dm + i d)e^{\left(-\frac{dm \log\left(\frac{2if}{d}\right) + 2ie - 2icf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idfx - icf)}{d}\right) + 2(df x + cf)(dx + c)^m}{4(adfm + adf)} \end{aligned}$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output
$$\frac{1}{4} \left(\left(I d m + I d \right) e^{-\left(d m \log \left(2 I f / d \right) + 2 I d e - 2 I c f \right) / d} \right) \text{gamma}(m + 1, -2 \left(-I d f x - I c f \right) / d) + 2 \left(d f x + c f \right) \left(d x + c \right)^m / (a d f m + a d * f)$$

Sympy [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = -\frac{i \int \frac{(c+dx)^m}{\tan(e+fx)-i} dx}{a}$$

input `integrate((d*x+c)**m/(a+I*a*tan(f*x+e)),x)`

output
$$-I * \text{Integral}((c + d x)^{**m} / (\tan(e + f x) - I), x) / a$$

Maxima [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(dx + c)^m}{i a \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output
$$\frac{1}{2} ((d m + d) * \text{integrate}((d x + c)^m * \cos(2 f x + 2 e), x) - (I d m + I d) * \text{integrate}((d x + c)^m * \sin(2 f x + 2 e), x) + e^{(m * \log(d x + c) + \log(d x + c))} / (a d m + a d)$$

Giac [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(dx + c)^m}{i a \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^m/(I*a*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(c + dx)^m}{a + a \tan(e + fx) \operatorname{Li}} dx$$

input `int((c + d*x)^m/(a + a*tan(e + f*x)*1i),x)`

output `int((c + d*x)^m/(a + a*tan(e + f*x)*1i), x)`

Reduce [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \frac{\int \frac{(dx+c)^m}{\tan(fx+e)i+1} dx}{a}$$

input `int((d*x+c)^m/(a+I*a*tan(f*x+e)),x)`

output `int((c + d*x)**m/(\tan(e + f*x)*i + 1),x)/a`

3.37 $\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx$

Optimal result	306
Mathematica [A] (verified)	307
Rubi [A] (verified)	307
Maple [F]	309
Fricas [A] (verification not implemented)	309
Sympy [F]	309
Maxima [F]	310
Giac [F]	310
Mupad [F(-1)]	310
Reduce [F]	311

Optimal result

Integrand size = 23, antiderivative size = 171

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ &= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{i2^{-2-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^2f} \\ & \quad + \frac{i4^{-2-m}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^2f} \end{aligned}$$

output

```
1/4*(d*x+c)^(1+m)/a^2/d/(1+m)+I*2^(-2-m)*(d*x+c)^m*GAMMA(1+m,2*I*f*(d*x+c)/d)/a^2/exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+I*4^(-2-m)*(d*x+c)^m*GAMM(A(1+m,4*I*f*(d*x+c)/d)/a^2/exp(4*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)
```

Mathematica [A] (verified)

Time = 3.64 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.12

$$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ = \frac{(c+dx)^m \left(\frac{4e^{2ie} f(c+dx)}{d(1+m)} + i2^{2-m} e^{\frac{2icf}{d}} \left(\frac{if(c+dx)}{d} \right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right) + i4^{-m} e^{-2ie+\frac{4icf}{d}} \left(\frac{if(c+dx)}{d} \right)^{-m} \Gamma\left(1-m, \frac{2if(c+dx)}{d}\right) \right)}{16f(a+ia\tan(e+fx))^2}$$

input `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x])^2, x]`

output $((c + d*x)^m * ((4*E^((2*I)*e)*f*(c + d*x))/(d*(1 + m)) + (I*2^(2 - m)*E^(((2*I)*c*f)/d)*Gamma[1 + m, ((2*I)*f*(c + d*x))/d])/((I*f*(c + d*x))/d)^m + (I*E^((-2*I)*e + ((4*I)*c*f)/d)*Gamma[1 + m, ((4*I)*f*(c + d*x))/d])/(4^m*((I*f*(c + d*x))/d)^m))*Sec[e + f*x]^2*(Cos[f*x] + I*Sin[f*x])^2)/(16*f*(a + I*a*Tan[e + f*x])^2)$

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ \downarrow 3042 \\ \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ \downarrow 4212 \\ \int \left(\frac{e^{-2ie-2ifx}(c+dx)^m}{2a^2} + \frac{e^{-4ie-4ifx}(c+dx)^m}{4a^2} + \frac{(c+dx)^m}{4a^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{i2^{-m-2}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{2if(c+dx)}{d}\right)}{a^2f} +$$

$$\frac{i4^{-m-2}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{4if(c+dx)}{d}\right)}{a^2f} + \frac{(c+dx)^{m+1}}{4a^2d(m+1)}$$

input $\text{Int}[(c + d*x)^m / (a + I*a*Tan[e + f*x])^2, x]$

output $(c + d*x)^(1 + m)/(4*a^2*d*(1 + m)) + (I*2^{-2 - m}*(c + d*x)^m*\text{Gamma}[1 + m, ((2*I)*f*(c + d*x))/d])/(a^2*E^{((2*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)}) + (I*4^{-2 - m}*(c + d*x)^m*\text{Gamma}[1 + m, ((4*I)*f*(c + d*x))/d])/(a^2*E^{((4*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)})$

Defintions of rubi rules used

rule 2009 $\text{Int}[u_, x_Symbol] :> \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] :> \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4212 $\text{Int}[((c_.) + (d_.)*(x_.)^m_*)((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + E^{(2*(a/b)*(e + f*x))/(2*a)})^{-n}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, 0]$

Maple [F]

$$\int \frac{(dx + c)^m}{(a + ia \tan(fx + e))^2} dx$$

input `int((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x)`

output `int((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.84

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \\ -\frac{4 (-i dm - i d)e^{\left(-\frac{dm \log\left(\frac{2if}{d}\right) + 2ie - 2icf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idfx - icf)}{d}\right) - (i dm + i d)e^{\left(-\frac{dm \log\left(\frac{4if}{d}\right) + 4ie - 4icf}{d}\right)}}{16(a^2 df m + a^2 df)}$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output `-1/16*(4*(-I*d*m - I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(a(m + 1, -2*(-I*d*f*x - I*c*f)/d) - (I*d*m + I*d)*e^(-(d*m*log(4*I*f/d) + 4*I*d*e - 4*I*c*f)/d)*gamma(m + 1, -4*(-I*d*f*x - I*c*f)/d) - 4*(d*f*x + c*f)*(d*x + c)^m)/(a^2*d*f*m + a^2*d*f)`

Sympy [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = -\frac{\int \frac{(c+dx)^m}{\tan^2(e+fx)-2i\tan(e+fx)-1} dx}{a^2}$$

input `integrate((d*x+c)**m/(a+I*a*tan(f*x+e))**2,x)`

output $-\text{Integral}((c + d*x)^m / (\tan(e + f*x)^2 - 2*I*\tan(e + f*x) - 1), x) / a^{**2}$

Maxima [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(dx + c)^m}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output $\frac{1}{4}*((d*m + d)*\int (d*x + c)^m * \cos(4*f*x + 4*e), x) + 2*(d*m + d)*\int (d*x + c)^m * \cos(2*f*x + 2*e), x) - (I*d*m + I*d)*\int (d*x + c)^m * \sin(4*f*x + 4*e), x) + 2*(-I*d*m - I*d)*\int (d*x + c)^m * \sin(2*f*x + 2*e), x) + e^{(m*\log(d*x + c) + \log(d*x + c))}/(a^{2*d*m} + a^{2*d})$

Giac [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(dx + c)^m}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^m/(I*a*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(c + dx)^m}{(a + a \tan(e + f x) 1i)^2} dx$$

input `int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^2,x)`

output `int((c + d*x)^m/(a + a*tan(e + f*x)*I)^2, x)`

Reduce [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = -\frac{\int \frac{(dx+c)^m}{\tan(fx+e)^2-2\tan(fx+e)i-1} dx}{a^2}$$

input `int((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x)`

output `(- int((c + d*x)**m/(tan(e + f*x)**2 - 2*tan(e + f*x)*I - 1),x))/a**2`

3.38 $\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx$

Optimal result	312
Mathematica [A] (verified)	313
Rubi [A] (verified)	313
Maple [F]	315
Fricas [A] (verification not implemented)	315
Sympy [F]	315
Maxima [F]	316
Giac [F]	316
Mupad [F(-1)]	317
Reduce [F]	317

Optimal result

Integrand size = 23, antiderivative size = 251

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx \\ &= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} + \frac{3i2^{-4-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^3f} \\ &+ \frac{3i2^{-5-2m}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^3f} \\ &+ \frac{i2^{-4-m}3^{-1-m}e^{-6i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{6if(c+dx)}{d}\right)}{a^3f} \end{aligned}$$

output

```
1/8*(d*x+c)^(1+m)/a^3/d/(1+m)+3*I*2^(-4-m)*(d*x+c)^m*GAMMA(1+m,2*I*f*(d*x+c)/d)/a^3/exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+3*I*2^(-5-2*m)*(d*x+c)^m*GAMMA(1+m,4*I*f*(d*x+c)/d)/a^3/exp(4*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+I*2^(-4-m)*3^(-1-m)*(d*x+c)^m*GAMMA(1+m,6*I*f*(d*x+c)/d)/a^3/exp(6*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)
```

Mathematica [A] (verified)

Time = 12.42 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.07

$$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx \\ = \frac{2^{-5-2m}3^{-1-m}e^{-3ie}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \left(12^{1+m}e^{6ie}f(c+dx)\left(\frac{if(c+dx)}{d}\right)^m + i2^{1+m}3^{2+m}de^{2i\left(2e+\frac{cf}{d}\right)}(1+e^{2i\left(2e+\frac{cf}{d}\right)})^m\right)}{(a+ia\tan(e+fx))^3}$$

input `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x])^3, x]`

output
$$(2^{-5 - 2m}3^{-1 - m}(c + d*x)^m(12^{1 + m}E^{((6*I)*e)*f*(c + d*x)*(I*f*(c + d*x))/d})^m + I*2^{(1 + m)*3^{(2 + m)*d*E^{((2*I)*(2*e + (c*f)/d))}}*(1 + m)*Gamma[1 + m, ((2*I)*f*(c + d*x))/d] + I*3^{(2 + m)*d*E^{((2*I)*e + ((4*I)*c*f)/d)}}*(1 + m)*Gamma[1 + m, ((4*I)*f*(c + d*x))/d] + I*2^{(1 + m)*d*E^{(((6*I)*c*f)/d)}}*(1 + m)*Gamma[1 + m, ((6*I)*f*(c + d*x))/d])*Sec[e + f*x]^3*(Cos[f*x] + I*Sin[f*x])^3)/(d*E^{((3*I)*e)*f*(1 + m)*((I*f*(c + d*x))/d)})^m*(a + I*a*Tan[e + f*x])^3)$$

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3042, 4212, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx \\ \downarrow \textcolor{blue}{3042} \\ \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx \\ \downarrow \textcolor{blue}{4212}$$

$$\int \left(\frac{3e^{-2ie-2ifx}(c+dx)^m}{8a^3} + \frac{3e^{-4ie-4ifx}(c+dx)^m}{8a^3} + \frac{e^{-6ie-6ifx}(c+dx)^m}{8a^3} + \frac{(c+dx)^m}{8a^3} \right) dx$$

↓ 2009

$$\frac{3i2^{-m-4}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{2if(c+dx)}{d}\right)}{a^3f} +$$

$$\frac{3i2^{-2m-5}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{4if(c+dx)}{d}\right)}{a^3f} +$$

$$\frac{i2^{-m-4}3^{-m-1}e^{-6i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(m+1,\frac{6if(c+dx)}{d}\right)}{a^3f} + \frac{(c+dx)^{m+1}}{8a^3d(m+1)}$$

input `Int[(c + d*x)^m/(a + I*a*Tan[e + f*x])^3, x]`

output
$$(c + d*x)^(1 + m)/(8*a^3*d*(1 + m)) + ((3*I)*2^(-4 - m)*(c + d*x)^m*\Gamma[1 + m, ((2*I)*f*(c + d*x))/d])/((a^3*E^((2*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m) + ((3*I)*2^(-5 - 2*m)*(c + d*x)^m*\Gamma[1 + m, ((4*I)*f*(c + d*x))/d])/((a^3*E^((4*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m) + (I*2^(-4 - m))*3^(-1 - m)*(c + d*x)^m*\Gamma[1 + m, ((6*I)*f*(c + d*x))/d])/((a^3*E^((6*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m))$$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simpl[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4212 `Int[((c_.) + (d_.)*(x_.))^m_*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]`

Maple [F]

$$\int \frac{(dx + c)^m}{(a + ia \tan(fx + e))^3} dx$$

input `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x)`

output `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.79

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \\ -\frac{18(-i dm - i d)e^{\left(-\frac{dm \log(\frac{2if}{d}) + 2ide - 2icf}{d}\right)} \Gamma(m + 1, -\frac{2(-idfx - icf)}{d}) + 9(-i dm - i d)e^{\left(-\frac{dm \log(\frac{4if}{d}) + 4ide - 4icf}{d}\right)} \Gamma(m + 1, -\frac{2(-idfx - icf)}{d})}{$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output `-1/96*(18*(-I*d*m - I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(m + 1, -2*(-I*d*f*x - I*c*f)/d) + 9*(-I*d*m - I*d)*e^(-(d*m*log(4*I*f/d) + 4*I*d*e - 4*I*c*f)/d)*gamma(m + 1, -4*(-I*d*f*x - I*c*f)/d) + 2*(-I*d*m - I*d)*e^(-(d*m*log(6*I*f/d) + 6*I*d*e - 6*I*c*f)/d)*gamma(m + 1, -6*(-I*d*f*x - I*c*f)/d) - 12*(d*f*x + c*f)*(d*x + c)^m)/(a^3*d*f*m + a^3*d*f)`

Sympy [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \frac{i \int \frac{(c+dx)^m}{\tan^3(e+fx)-3i\tan^2(e+fx)-3\tan(e+fx)+i} dx}{a^3}$$

input `integrate((d*x+c)**m/(a+I*a*tan(f*x+e))**3,x)`

output $I * \text{Integral}((c + d*x)^m / (\tan(e + f*x))^3 - 3*I*\tan(e + f*x)^2 - 3*\tan(e + f*x) + I), x) / a^{**3}$

Maxima [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(dx + c)^m}{(ia \tan(fx + e) + a)^3} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output $\frac{1}{8}((d*m + d)*\text{integrate}((d*x + c)^m * \cos(6*f*x + 6*e), x) + 3*(d*m + d)*\text{integrate}((d*x + c)^m * \cos(4*f*x + 4*e), x) + 3*(d*m + d)*\text{integrate}((d*x + c)^m * \cos(2*f*x + 2*e), x) - (I*d*m + I*d)*\text{integrate}((d*x + c)^m * \sin(6*f*x + 6*e), x) + 3*(-I*d*m - I*d)*\text{integrate}((d*x + c)^m * \sin(4*f*x + 4*e), x) + 3*(-I*d*m - I*d)*\text{integrate}((d*x + c)^m * \sin(2*f*x + 2*e), x) + e^{(m*\log(d*x + c) + \log(d*x + c))}/(a^{3*d*m} + a^{3*d})$

Giac [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(dx + c)^m}{(ia \tan(fx + e) + a)^3} dx$$

input `integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output $\text{integrate}((d*x + c)^m / (I*a*tan(f*x + e) + a)^3, x)$

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(c + dx)^m}{(a + a \tan(e + fx) 1i)^3} dx$$

input `int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^3,x)`

output `int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^3, x)`

Reduce [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = -\frac{\int \frac{(dx+c)^m}{\tan(fx+e)^3 i + 3 \tan(fx+e)^2 - 3 \tan(fx+e)i - 1} dx}{a^3}$$

input `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x)`

output `(- int((c + d*x)**m/(\tan(e + f*x)**3*i + 3*tan(e + f*x)**2 - 3*tan(e + f*x)*i - 1),x))/a**3`

3.39 $\int (c + dx)^3(a + b \tan(e + fx)) dx$

Optimal result	318
Mathematica [B] (verified)	319
Rubi [A] (verified)	320
Maple [B] (verified)	321
Fricas [B] (verification not implemented)	322
Sympy [F]	323
Maxima [B] (verification not implemented)	323
Giac [F]	324
Mupad [F(-1)]	324
Reduce [F]	325

Optimal result

Integrand size = 18, antiderivative size = 152

$$\begin{aligned} \int (c + dx)^3(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^4}{4d} + \frac{ib(c + dx)^4}{4d} \\ & - \frac{b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{3ibd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{3bd^2(c + dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\ & - \frac{3ibd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \end{aligned}$$

output

```
1/4*a*(d*x+c)^4/d+1/4*I*b*(d*x+c)^4/d-b*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f
+3/2*I*b*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2-3/2*b*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3-3/4*I*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 342 vs. $2(152) = 304$.

Time = 0.12 (sec) , antiderivative size = 342, normalized size of antiderivative = 2.25

$$\begin{aligned}
 \int (c + dx)^3(a + b \tan(e + fx)) dx = & ac^3x + \frac{3}{2}ac^2dx^2 + \frac{3}{2}ibc^2dx^2 + acd^2x^3 + ibcd^2x^3 \\
 & + \frac{1}{4}ad^3x^4 + \frac{1}{4}ibd^3x^4 - \frac{3bc^2dx \log(1 + e^{2i(e+fx)})}{f} \\
 & - \frac{3bcd^2x^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & - \frac{bd^3x^3 \log(1 + e^{2i(e+fx)})}{f} - \frac{bc^3 \log(\cos(e + fx))}{f} \\
 & + \frac{3ibc^2d \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
 & + \frac{3ibcd^2x \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & + \frac{3ibd^3x^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
 & - \frac{3bcd^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{3bd^3x \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{3ibd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
 \end{aligned}$$

input `Integrate[(c + d*x)^3*(a + b*Tan[e + f*x]), x]`

output

$$\begin{aligned}
 & a*c^3*x + (3*a*c^2*d*x^2)/2 + ((3*I)/2)*b*c^2*d*x^2 + a*c*d^2*x^3 + I*b*c*d^2*x^3 + (a*d^3*x^4)/4 + (I/4)*b*d^3*x^4 - (3*b*c^2*d*x*Log[1 + E^((2*I)*(e + f*x))])/f - (3*b*c*d^2*x^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b*d^3*x^3*Log[1 + E^((2*I)*(e + f*x))])/f - (b*c^3*Log[Cos[e + f*x]])/f + (((3*I)/2)*b*c^2*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + ((3*I)*b*c*d^2*x*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (((3*I)/2)*b*d^3*x^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (3*b*c*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (3*b*d^3*x*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((3*I)/4)*b*d^3*x*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4
 \end{aligned}$$

Rubi [A] (verified)

Time = 0.48 (sec), antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3(a + b \tan(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int (c + dx)^3(a + b \tan(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{4205} \\
 & \int (a(c + dx)^3 + b(c + dx)^3 \tan(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{a(c + dx)^4}{4d} - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{3ibd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \\
 & \quad \frac{b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{ib(c + dx)^4}{4d} - \frac{3ibd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
 \end{aligned}$$

input

```
Int[(c + d*x)^3*(a + b*Tan[e + f*x]),x]
```

output

$$(a*(c + d*x)^4)/(4*d) + ((I/4)*b*(c + d*x)^4)/d - (b*(c + d*x)^3*Log[1 + E^((2*I)*(e + f*x))])/f + (((3*I)/2)*b*d*(c + d*x)^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (3*b*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((3*I)/4)*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4205 $\text{Int}[(c_.) + (d_.)*(x_.)^(m_.)*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^(n_.), x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (a + b*\tan[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 499 vs. $2(131) = 262$.

Time = 0.53 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.29

method	result
risch	$\frac{id^3 b x^4}{4} + \frac{2 b c^3 \ln(e^{i(f x + e)})}{f} - \frac{b c^3 \ln(1 + e^{2 i(f x + e)})}{f} - \frac{3 b d c^2 \ln(1 + e^{2 i(f x + e)}) x}{f} - \frac{3 b d^2 c \ln(1 + e^{2 i(f x + e)}) x^2}{f} + \frac{6 b e^2 d^2 c \ln(1 + e^{2 i(f x + e)}) x^3}{f}$

input $\text{int}((d*x+c)^3*(a+b*\tan(f*x+e)), x, \text{method}=\text{_RETURNVERBOSE})$

output

```

d^2*a*c*x^3+3/2*d*a*c^2*x^2+a*c^3*x-I*b*c^3*x-1/4*I/d*b*c^4-2/f^4*b*e^3*d^
3*ln(exp(I*(f*x+e)))-1/f*b*d^3*ln(1+exp(2*I*(f*x+e)))*x^3-3/2/f^3*b*d^3*po
lylog(3,-exp(2*I*(f*x+e)))*x-3/2/f^3*b*d^2*c*polylog(3,-exp(2*I*(f*x+e)))+
3/2*I/f^4*b*d^3*e^4+3/2*I*d*b*c^2*x^2+1/4*I*d^3*b*x^4+2/f*b*c^3*ln(exp(I*(f*x+e)))-1/f*b*c^3*ln(1+exp(2*I*(f*x+e)))+1/4*d^3*a*x^4+1/4/d*a*c^4+I*d^2*
b*c*x^3-3/f*b*d*c^2*ln(1+exp(2*I*(f*x+e)))*x-3/f*b*d^2*c*ln(1+exp(2*I*(f*x+e)))*x^2+6/f^3*b*e^2*d^2*c*ln(exp(I*(f*x+e)))-6/f^2*b*e*d*c^2*ln(exp(I*(f*x+e)))+2*I/f^3*b*d^3*e^3*x+3/2*I/f^2*b*d^3*polylog(2,-exp(2*I*(f*x+e)))*x^2-4*I/f^3*b*d^2*c*e^3+3*I/f^2*b*d*c^2*e^2+3/2*I/f^2*b*d*c^2*polylog(2,-ex
p(2*I*(f*x+e)))+3*I/f^2*b*d^2*c*polylog(2,-exp(2*I*(f*x+e)))*x-6*I/f^2*b*d
^2*c*e^2*x-3/4*I*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4+6*I/f*b*d*c^2*e*x

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 500 vs. $2(127) = 254$.

Time = 0.09 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.29

$$\int (c + dx)^3(a + b \tan(e + fx)) dx \\
 = \frac{2 ad^3 f^4 x^4 + 8 acd^2 f^4 x^3 + 12 ac^2 df^4 x^2 + 8 ac^3 f^4 x + 3i bd^3 \text{polylog}\left(4, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3i bd^3 \text{pc}}$$

input

```
integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="fricas")
```

output

$$\begin{aligned} & \frac{1}{8} \left(2*a*d^3*f^4*x^4 + 8*a*c*d^2*f^4*x^3 + 12*a*c^2*d*f^4*x^2 + 8*a*c^3*f^4*x + 3*I*b*d^3*\text{polylog}(4, (\tan(f*x + e))^2 + 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) - 3*I*b*d^3*\text{polylog}(4, (\tan(f*x + e))^2 - 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) - 6*(I*b*d^3*f^2*x^2 + 2*I*b*c*d^2*f^2*x + I*b*c^2*d*f^2)*\text{dilog}(2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 6*(-I*b*d^3*f^2*x^2 - 2*I*b*c*d^2*f^2*x - I*b*c^2*d*f^2)*\text{dilog}(2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*c^3*f^3)*\log(-2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*c^3*f^3)*\log(-2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(b*d^3*f*x + b*c*d^2*f)*\text{polylog}(3, (\tan(f*x + e))^2 + 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) - 6*(b*d^3*f*x + b*c*d^2*f)*\text{polylog}(3, (\tan(f*x + e))^2 - 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) \right) / f^4 \end{aligned}$$

Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^3 dx$$

input

```
integrate((d*x+c)**3*(a+b*tan(f*x+e)),x)
```

output

```
Integral((a + b*tan(e + fx))*(c + d*x)**3, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 672 vs. $2(127) = 254$.

Time = 0.22 (sec), antiderivative size = 672, normalized size of antiderivative = 4.42

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \text{Too large to display}$$

input

```
integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="maxima")
```

output

```
1/12*(12*(f*x + e)*a*c^3 + 3*(f*x + e)^4*a*d^3/f^3 - 12*(f*x + e)^3*a*d^3*e/f^3 + 18*(f*x + e)^2*a*d^3*e^2/f^3 - 12*(f*x + e)*a*d^3*e^3/f^3 + 12*(f*x + e)^3*a*c*d^2/f^2 - 36*(f*x + e)^2*a*c*d^2*e/f^2 + 36*(f*x + e)*a*c*d^2*e^2/f^2 + 18*(f*x + e)^2*a*c^2*d/f - 36*(f*x + e)*a*c^2*d*e/f + 12*b*c^3*log(sec(f*x + e)) - 12*b*d^3*e^3*log(sec(f*x + e))/f^3 + 36*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 36*b*c^2*d*e*log(sec(f*x + e))/f - (-3*I*(f*x + e)^4*b*d^3 + 12*I*b*d^3*polylog(4, -e^(2*I*f*x + 2*I*e)) - 12*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e)^3 - 18*(I*b*d^3*e^2 - 2*I*b*c*d^2*e*f + I*b*c^2*d*f^2)*(f*x + e)^2 - 4*(-4*I*(f*x + e)^3*b*d^3 + 9*(I*b*d^3*e - I*b*c*d^2*f)*(f*x + e)^2 + 9*(-I*b*d^3*e^2 + 2*I*b*c*d^2*e*f - I*b*c^2*d*f^2)*(f*x + e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 6*(4*I*(f*x + e)^2*b*d^3 + 3*I*b*d^3*e^2 - 6*I*b*c*d^2*e*f + 3*I*b*c^2*d*f^2 + 6*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e))*dilog(-e^(2*I*f*x + 2*I*e)) + 2*(4*(f*x + e)^3*b*d^3 - 9*(b*d^3*e - b*c*d^2*f)*(f*x + e)^2 + 9*(b*d^3*e^2 - 2*b*c*d^2*e*f + b*c^2*d*f^2)*(f*x + e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) + 6*(4*(f*x + e)*b*d^3 - 3*b*d^3*e + 3*b*c*d^2*f)*polylog(3, -e^(2*I*f*x + 2*I*e)))/f^3)/f
```

Giac [F]

$$\int (c + dx)^3(a + b \tan(e + fx)) dx = \int (dx + c)^3(b \tan(fx + e) + a) dx$$

input

```
integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="giac")
```

output

```
integrate((d*x + c)^3*(b*tan(f*x + e) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^3 dx$$

input

```
int((a + b*tan(e + f*x))*(c + d*x)^3,x)
```

output `int((a + b*tan(e + f*x))*(c + d*x)^3, x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)^3(a + b \tan(e + fx)) dx \\ &= \frac{4(\int \tan(fx + e) x^3 dx) b d^3 f + 12(\int \tan(fx + e) x^2 dx) b c d^2 f + 12(\int \tan(fx + e) x dx) b c^2 df + 2 \log(t)}{4f} \end{aligned}$$

input `int((d*x+c)^3*(a+b*tan(f*x+e)),x)`

output `(4*int(tan(e + f*x)*x**3,x)*b*d**3*f + 12*int(tan(e + f*x)*x**2,x)*b*c*d**2*f + 12*int(tan(e + f*x)*x,x)*b*c**2*d*f + 2*log(tan(e + f*x)**2 + 1)*b*c**3 + 4*a*c**3*f*x + 6*a*c**2*d*f*x**2 + 4*a*c*d**2*f*x**3 + a*d**3*f*x**4)/(4*f)`

3.40 $\int (c + dx)^2(a + b \tan(e + fx)) dx$

Optimal result	326
Mathematica [A] (verified)	327
Rubi [A] (verified)	327
Maple [B] (verified)	329
Fricas [B] (verification not implemented)	329
Sympy [F]	330
Maxima [B] (verification not implemented)	330
Giac [F]	331
Mupad [F(-1)]	331
Reduce [F]	332

Optimal result

Integrand size = 18, antiderivative size = 115

$$\begin{aligned} \int (c + dx)^2(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^3}{3d} + \frac{ib(c + dx)^3}{3d} \\ & - \frac{b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ibd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \end{aligned}$$

output

```
1/3*a*(d*x+c)^3/d+1/3*I*b*(d*x+c)^3/d-b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f
+I*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-1/2*b*d^2*polylog(3,-exp(2
*I*(f*x+e)))/f^3
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.66

$$\begin{aligned} \int (c + dx)^2(a + b \tan(e + fx)) dx = & ac^2x + acdx^2 + ibcdx^2 + \frac{1}{3}ad^2x^3 \\ & + \frac{1}{3}ibd^2x^3 - \frac{2bcdx \log(1 + e^{2i(e+fx)})}{f} \\ & - \frac{bd^2x^2 \log(1 + e^{2i(e+fx)})}{f} - \frac{bc^2 \log(\cos(e + fx))}{f} \\ & + \frac{ibcd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & + \frac{ibd^2x \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \end{aligned}$$

input `Integrate[(c + d*x)^2*(a + b*Tan[e + f*x]), x]`

output $a*c^2*x + a*c*d*x^2 + I*b*c*d*x^2 + (a*d^2*x^3)/3 + (I/3)*b*d^2*x^3 - (2*b*c*d*x*Log[1 + E^((2*I)*(e + f*x))])/f - (b*d^2*x^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b*c^2*Log[Cos[e + f*x]])/f + (I*b*c*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (I*b*d^2*x*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3)$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^2(a + b \tan(e + fx)) dx$$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 \int (c + dx)^2 (a + b \tan(e + fx)) dx \\
 \downarrow \text{4205} \\
 \int (a(c + dx)^2 + b(c + dx)^2 \tan(e + fx)) dx \\
 \downarrow \text{2009} \\
 \frac{a(c + dx)^3}{3d} + \frac{ibd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} + \\
 \frac{ib(c + dx)^3}{3d} - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3}
 \end{array}$$

input `Int[(c + d*x)^2*(a + b*Tan[e + f*x]),x]`

output `(a*(c + d*x)^3)/(3*d) + ((I/3)*b*(c + d*x)^3)/d - (b*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f + (I*b*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3)`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 313 vs. $2(101) = 202$.

Time = 0.50 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.73

method	result
risch	$\frac{2ibdc e^2}{f^2} + \frac{ibdc \operatorname{polylog}(2, -e^{2i(fx+e)})}{f^2} + \frac{d^2 a x^3}{3} - \frac{ib c^3}{3d} + dac x^2 - \frac{4ib d^2 e^3}{3f^3} + a c^2 x + \frac{a c^3}{3d} - \frac{b d^2 \ln(1+e^{2i(fx+e)})}{f}$

input `int((d*x+c)^2*(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 2*I/f^2*b*d*c*e^2+I/f^2*b*d*c*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))+1/3*d^2*a*x^3-1 \\ & /3*I/d*b*c^3+d*a*c*x^2-4/3*I/f^3*b*d^2*e^3+a*c^2*x+1/3/d*a*c^3-1/f*b*d^2*I \\ & n(1+\exp(2*I*(f*x+e)))*x^2-1/2*b*d^2*\operatorname{polylog}(3, -\exp(2*I*(f*x+e)))/f^3+2/f*b \\ & *c^2*\ln(\exp(I*(f*x+e)))-1/f*b*c^2*\ln(1+\exp(2*I*(f*x+e)))+2/f^3*b*d^2*e^2*I \\ & n(\exp(I*(f*x+e)))-2*I/f^2*b*d^2*e^2*x-2/f*b*d*c*\ln(1+\exp(2*I*(f*x+e)))*x-I \\ & *b*c^2*x+1/3*I*d^2*b*x^3+I/f^2*b*d^2*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))*x+I*d*b*c \\ & *x^2+4*I/f*b*d*c*e*x-4/f^2*b*c*d*e*\ln(\exp(I*(f*x+e))) \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(98) = 196$.

Time = 0.08 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.76

$$\begin{aligned} & \int (c + dx)^2(a + b \tan(e + fx)) dx \\ & = \frac{4 ad^2 f^3 x^3 + 12 acdf^3 x^2 + 12 ac^2 f^3 x - 3 bd^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2+2i \tan(fx+e)-1}{\tan(fx+e)^2+1}\right) - 3 bd^2 \operatorname{polylog}\left(3, \frac{\tan(fx+e)^2+2i \tan(fx+e)+1}{\tan(fx+e)^2+1}\right)}{f^2} \end{aligned}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="fricas")`

```
output 1/12*(4*a*d^2*f^3*x^3 + 12*a*c*d*f^3*x^2 + 12*a*c^2*f^3*x - 3*b*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*b*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(-I*b*d^2*f*x - I*b*c*d*f)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x + b*c^2*f^2)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x + b*c^2*f^2)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1))/f^3
```

Sympy [F]

$$\int (c + dx)^2(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx))(c + dx)^2 dx$$

```
input integrate((d*x+c)**2*(a+b*tan(f*x+e)),x)
```

```
output Integral((a + b*tan(e + f*x))*(c + d*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 373 vs. $2(98) = 196$.

Time = 0.20 (sec) , antiderivative size = 373, normalized size of antiderivative = 3.24

$$\begin{aligned} & \int (c + dx)^2(a + b \tan(e + fx)) dx \\ &= \frac{6(fx + e)ac^2 + \frac{2(fx+e)^3ad^2}{f^2} - \frac{6(fx+e)^2ad^2e}{f^2} + \frac{6(fx+e)ad^2e^2}{f^2} + \frac{6(fx+e)^2acd}{f} - \frac{12(fx+e)acde}{f} + 6bc^2 \log(\sec(fx + e)))}{f^3} \end{aligned}$$

```
input integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="maxima")
```

output

$$\begin{aligned} & \frac{1}{6} \left(6(fx + e)ac^2 + 2(fx + e)^3ad^2/f^2 - 6(fx + e)^2ad^2e/f \right. \\ & \quad \left. - 6(fx + e)^2a^2d^2/f^2 + 6(fx + e)^2a^2cd/f - 12(fx + e)a^2c^2d/f \right. \\ & \quad \left. + 6b^2c^2\log(\sec(fx + e)) + 6b^2d^2e^2\log(\sec(fx + e))/f^2 - 12 \right. \\ & \quad \left. *b^2c^2d^2e^2\log(\sec(fx + e))/f - (-2I*(fx + e)^3b^2d^2 + 3b^2d^2\text{polylog}(3, -e^{(2Ifx + 2Ie)}) - 6(-Ib^2d^2e^2 + I^2b^2c^2d^2f)*(fx + e)^2 - 6(-I*(fx + e)^2b^2d^2 + 2(I^2b^2d^2e^2 - I^2b^2c^2d^2f)*(fx + e))\arctan2(\sin(2fx + 2e), \cos(2fx + 2e) + 1) - 6(I^2(fx + e)b^2d^2 - I^2b^2d^2e^2 + I^2b^2c^2d^2f)\text{dilog}(-e^{(2Ifx + 2Ie)}) + 3((fx + e)^2b^2d^2 - 2(b^2d^2e^2 - b^2c^2d^2f)*(fx + e))\log(\cos(2fx + 2e)^2 + \sin(2fx + 2e)^2 + 2\cos(2fx + 2e) + 1))/f^2 \right) / f \end{aligned}$$

Giac [F]

$$\int (c + dx)^2(a + b\tan(e + fx)) dx = \int (dx + c)^2(b\tan(fx + e) + a) dx$$

input

```
integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="giac")
```

output

```
integrate((d*x + c)^2*(b*tan(f*x + e) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2(a + b\tan(e + fx)) dx = \int (a + b\tan(e + fx)) (c + dx)^2 dx$$

input

```
int((a + b*tan(e + f*x))*(c + d*x)^2,x)
```

output

```
int((a + b*tan(e + f*x))*(c + d*x)^2, x)
```

Reduce [F]

$$\int (c + dx)^2(a + b \tan(e + fx)) dx \\ = \frac{6(\int \tan(fx + e) x^2 dx) b d^2 f + 12(\int \tan(fx + e) x dx) b c d f + 3 \log(\tan(fx + e)^2 + 1) b c^2 + 6a c^2 f x +}{6f}$$

input `int((d*x+c)^2*(a+b*tan(f*x+e)),x)`

output `(6*int(tan(e + f*x)*x**2,x)*b*d**2*f + 12*int(tan(e + f*x)*x,x)*b*c*d*f + 3*log(tan(e + f*x)**2 + 1)*b*c**2 + 6*a*c**2*f*x + 6*a*c*d*f*x**2 + 2*a*d**2*f*x**3)/(6*f)`

3.41 $\int (c + dx)(a + b \tan(e + fx)) dx$

Optimal result	333
Mathematica [A] (verified)	333
Rubi [A] (verified)	334
Maple [A] (verified)	335
Fricas [B] (verification not implemented)	335
Sympy [F]	336
Maxima [A] (verification not implemented)	336
Giac [F]	337
Mupad [B] (verification not implemented)	337
Reduce [F]	338

Optimal result

Integrand size = 16, antiderivative size = 84

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^2}{2d} + \frac{ib(c + dx)^2}{2d} \\ & - \frac{b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \end{aligned}$$

output
$$\frac{1/2*a*(d*x+c)^2/d+1/2*I*b*(d*x+c)^2/d-b*(d*x+c)*\ln(1+\exp(2*I*(f*x+e)))/f+1}{2*I*b*d*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))/f^2}$$

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 87, normalized size of antiderivative = 1.04

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx)) dx = & acx + \frac{1}{2}adx^2 + \frac{1}{2}ibdx^2 - \frac{bdx \log(1 + e^{2i(e+fx)})}{f} \\ & - \frac{bc \log(\cos(e + fx))}{f} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \end{aligned}$$

input
$$\text{Integrate}[(c + d*x)*(a + b*Tan[e + f*x]), x]$$

output
$$\frac{a*c*x + (a*d*x^2)/2 + ((I/2)*b*d*x^2 - (b*d*x*Log[1 + E^((2*I)*(e + f*x))]))}{f} - \frac{(b*c*Log[Cos[e + f*x]])}{f} + \frac{((I/2)*b*d*PolyLog[2, -E^((2*I)*(e + f*x))])}{f^2}$$

Rubi [A] (verified)

Time = 0.32 (sec), antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.188, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)(a + b \tan(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int (c + dx)(a + b \tan(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{4205} \\
 & \int (a(c + dx) + b(c + dx) \tan(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{a(c + dx)^2}{2d} - \frac{b(c + dx) \log(1 + e^{2i(e + fx)})}{f} + \frac{ib(c + dx)^2}{2d} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e + fx)})}{2f^2}
 \end{aligned}$$

input $\operatorname{Int}[(c + d*x)*(a + b*\operatorname{Tan}[e + f*x]), x]$

output
$$\frac{(a*(c + d*x)^2)/(2*d) + ((I/2)*b*(c + d*x)^2)/d - (b*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])}{f} + \frac{((I/2)*b*d*PolyLog[2, -E^((2*I)*(e + f*x))])}{f^2}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4205 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (a + b*\text{Tan}[e + f*x])^n, x], \ x] /; \ \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \& \ \text{IGtQ}[m, 0] \ \& \ \text{IGtQ}[n, 0]$

Maple [A] (verified)

Time = 0.45 (sec), antiderivative size = 143, normalized size of antiderivative = 1.70

method	result
risch	$\frac{ibdx^2}{2} + \frac{adx^2}{2} - ibcx + acx + \frac{2bc\ln(e^{i(fx+e)})}{f} - \frac{bc\ln(1+e^{2i(fx+e)})}{f} + \frac{2ibdex}{f} + \frac{ibde^2}{f^2} - \frac{bd\ln(1+e^{2i(fx+e)})x}{f} + \dots$

input $\text{int}((d*x+c)*(a+b*\tan(f*x+e)), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$\begin{aligned} & 1/2*I*b*d*x^2+1/2*a*d*x^2-I*b*c*x+a*c*x+2/f*b*c*\ln(\exp(I*(f*x+e)))-1/f*b*c \\ & *\ln(1+\exp(2*I*(f*x+e)))+2*I/f*b*d*e*x+I/f^2*b*d*e^2-1/f*b*d*\ln(1+\exp(2*I*(f*x+e)))*x+1/2*I*b*d*\text{polylog}(2, -\exp(2*I*(f*x+e)))/f^2-2/f^2*b*e*d*\ln(\exp(I*(f*x+e))) \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(69) = 138.

Time = 0.09 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.90

$$\int (c + dx)(a + b \tan(e + fx)) dx \\ = \frac{2 adf^2 x^2 + 4 acf^2 x - i bd \text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + i bd \text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) - 2(bdfx + bcf) \log\left(\frac{bdfx + bcf}{4 f^2}\right)}{4 f^2}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="fricas")`

output $\frac{1}{4} (2 a d f^2 x^2 + 4 a c f^2 x - I b d \operatorname{dilog}(2 (\operatorname{I} \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1) + 1) + I b d \operatorname{dilog}(2 (-\operatorname{I} \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1) + 1) - 2 (b d f x + b c f) \log(-2 (\operatorname{I} \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1)) - 2 (b d f x + b c f) \log(-2 (-\operatorname{I} \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1))) / f^2$

Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx))(c + dx) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x)`

output `Integral((a + b*tan(e + f*x))*(c + d*x), x)`

Maxima [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.55

$$\int (c + dx)(a + b \tan(e + fx)) dx \\ = \frac{(a + i b) df^2 x^2 + 2 (a + i b) cf^2 x + i bd \text{Li}_2(-e^{(2 i f x + 2 i e)}) + 2 (-i bdfx - i bcf) \arctan(\sin(2fx + 2e), \cos(2fx + 2e))}{2 f^2}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="maxima")`

output
$$\frac{1}{2}((a + I*b)*d*f^2*x^2 + 2*(a + I*b)*c*f^2*x + I*b*d*dilog(-e^{(2*I*f*x + 2*I*e)}) + 2*(-I*b*d*f*x - I*b*c*f)*arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e) + 1) - (b*d*f*x + b*c*f)*log(\cos(2*f*x + 2*e)^2 + \sin(2*f*x + 2*e)^2 + 2*\cos(2*f*x + 2*e) + 1))/f^2$$

Giac [F]

$$\int (c + dx)(a + b \tan(e + fx)) dx = \int (dx + c)(b \tan(fx + e) + a) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)*(b*tan(f*x + e) + a), x)`

Mupad [B] (verification not implemented)

Time = 8.50 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.92

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx)) dx &= \frac{ax(2c + dx)}{2} \\ &- \frac{bd(\pi \ln(\cos(fx)) - \pi \ln(e^{fx2i} + 1) + \text{polylog}(2, -e^{-e2i} e^{-fx2i}) 1i - \pi \ln(e^{-e2i} e^{-fx2i} + 1) + 2e^{fx2i}))}{2f} \\ &+ \frac{bc \ln(\tan(e + fx)^2 + 1)}{2f} \end{aligned}$$

input `int((a + b*tan(e + f*x))*(c + d*x),x)`

output
$$\begin{aligned} &(a*x*(2*c + d*x))/2 - (b*d*(\text{polylog}(2, -\exp(-e*2i)*\exp(-f*x*2i))*1i - \pi*i*\log(\exp(f*x*2i) + 1) - \log(\cos(e + f*x))*(2*e - \pi) - \pi*\log(\exp(-e*2i)*\exp(-f*x*2i) + 1) + 2*e*\log(\exp(-e*2i)*\exp(-f*x*2i) + 1) + \pi*\log(\cos(f*x)) + f^2*x^2*1i + 2*f*x*\log(\exp(-e*2i)*\exp(-f*x*2i) + 1) + e*f*x*2i))/(2*f^2) \\ &+ (b*c*\log(\tan(e + f*x)^2 + 1))/(2*f) \end{aligned}$$

Reduce [F]

$$\int (c + dx)(a + b \tan(e + fx)) dx \\ = \frac{2(\int \tan(fx + e) x dx) bdf + \log(\tan(fx + e)^2 + 1) bc + 2acf x + adf x^2}{2f}$$

input `int((d*x+c)*(a+b*tan(f*x+e)),x)`

output `(2*int(tan(e + f*x)*x,x)*b*d*f + log(tan(e + f*x)**2 + 1)*b*c + 2*a*c*f*x + a*d*f*x**2)/(2*f)`

3.42 $\int \frac{a+b \tan(e+fx)}{c+dx} dx$

Optimal result	339
Mathematica [N/A]	339
Rubi [N/A]	340
Maple [N/A]	341
Fricas [N/A]	341
Sympy [N/A]	341
Maxima [N/A]	342
Giac [N/A]	342
Mupad [N/A]	342
Reduce [N/A]	343

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \text{Int}\left(\frac{a + b \tan(e + fx)}{c + dx}, x\right)$$

output `Defer(Int)((a+b*tan(f*x+e))/(d*x+c),x)`

Mathematica [N/A]

Not integrable

Time = 1.73 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

input `Integrate[(a + b*Tan[e + f*x])/ (c + d*x), x]`

output `Integrate[(a + b*Tan[e + f*x])/ (c + d*x), x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{a + b \tan(e + fx)}{c + dx} dx \\ \downarrow 3042 \\ \int \frac{a + b \tan(e + fx)}{c + dx} dx \\ \downarrow 4223 \\ \int \frac{a + b \tan(e + fx)}{c + dx} dx \end{array}$$

input `Int[(a + b*Tan[e + f*x])/(c + d*x),x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tan(fx + e)}{dx + c} dx$$

input `int((a+b*tan(f*x+e))/(d*x+c),x)`

output `int((a+b*tan(f*x+e))/(d*x+c),x)`

Fricas [N/A]

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="fricas")`

output `integral((b*tan(f*x + e) + a)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x)`

output `Integral((a + b*tan(e + f*x))/(c + d*x), x)`

Maxima [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 4.78

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="maxima")`

output `(2*b*d*integrate(sin(2*f*x + 2*e)/((d*x + c)*cos(2*f*x + 2*e)^2 + (d*x + c)*sin(2*f*x + 2*e)^2 + d*x + 2*(d*x + c)*cos(2*f*x + 2*e) + c), x) + a*log(d*x + c))/d`

Giac [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 8.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + f x)}{c + d x} dx$$

input `int((a + b*tan(e + f*x))/(c + d*x),x)`

output `int((a + b*tan(e + f*x))/(c + d*x), x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.78

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \frac{\left(\int \frac{\tan(fx+e)}{dx+c} dx \right) bd + \log(dx+c) a}{d}$$

input `int((a+b*tan(f*x+e))/(d*x+c),x)`

output `(int(tan(e + f*x)/(c + d*x),x)*b*d + log(c + d*x)*a)/d`

3.43 $\int \frac{a+b \tan(e+fx)}{(c+dx)^2} dx$

Optimal result	344
Mathematica [N/A]	344
Rubi [N/A]	345
Maple [N/A]	346
Fricas [N/A]	346
Sympy [N/A]	346
Maxima [N/A]	347
Giac [N/A]	347
Mupad [N/A]	348
Reduce [N/A]	348

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \text{Int}\left(\frac{a + b \tan(e + fx)}{(c + dx)^2}, x\right)$$

output `Defer(Int)((a+b*tan(f*x+e))/(d*x+c)^2,x)`

Mathematica [N/A]

Not integrable

Time = 4.58 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `Integrate[(a + b*Tan[e + f*x])/(c + d*x)^2, x]`

output `Integrate[(a + b*Tan[e + f*x])/(c + d*x)^2, x]`

Rubi [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx \\ \downarrow \text{3042} \\ \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx \\ \downarrow \text{4223} \\ \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx \end{array}$$

input `Int[(a + b*Tan[e + f*x])/(c + d*x)^2, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tan(fx + e)}{(dx + c)^2} dx$$

input `int((a+b*tan(f*x+e))/(d*x+c)^2,x)`

output `int((a+b*tan(f*x+e))/(d*x+c)^2,x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.72

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="fricas")`

output `integral((b*tan(f*x + e) + a)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 2.37 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)**2,x)`

output `Integral((a + b*tan(e + f*x))/(c + d*x)**2, x)`

Maxima [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 142, normalized size of antiderivative = 7.89

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="maxima")`

output `(2*(b*d^2*x + b*c*d)*integrate(sin(2*f*x + 2*e)/(d^2*x^2 + 2*c*d*x + (d^2*x^2 + 2*c*d*x + c^2)*cos(2*f*x + 2*e))^2 + (d^2*x^2 + 2*c*d*x + c^2)*sin(2*f*x + 2*e)^2 + c^2 + 2*(d^2*x^2 + 2*c*d*x + c^2)*cos(2*f*x + 2*e)), x) - a)/(d^2*x + c*d)`

Giac [N/A]

Not integrable

Time = 3.92 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 8.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

input `int((a + b*tan(e + f*x))/(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))/(c + d*x)^2, x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 79, normalized size of antiderivative = 4.39

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \frac{\left(\int \frac{\tan(fx+e)}{d^2x^2+2cdx+c^2} dx \right) b c^2 + \left(\int \frac{\tan(fx+e)}{d^2x^2+2cdx+c^2} dx \right) bcdx + ax}{c(dx + c)}$$

input `int((a+b*tan(f*x+e))/(d*x+c)^2,x)`

output `(int(tan(e + f*x)/(c**2 + 2*c*d*x + d**2*x**2),x)*b*c**2 + int(tan(e + f*x)/(c**2 + 2*c*d*x + d**2*x**2),x)*b*c*d*x + a*x)/(c*(c + d*x))`

3.44 $\int (c + dx)^3 (a + b \tan(e + fx))^2 dx$

Optimal result	349
Mathematica [B] (warning: unable to verify)	350
Rubi [A] (verified)	351
Maple [B] (verified)	353
Fricas [B] (verification not implemented)	354
Sympy [F]	354
Maxima [B] (verification not implemented)	355
Giac [F]	356
Mupad [F(-1)]	356
Reduce [F]	356

Optimal result

Integrand size = 20, antiderivative size = 300

$$\begin{aligned} \int (c + dx)^3 (a + b \tan(e + fx))^2 dx = & -\frac{ib^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} + \frac{iab(c + dx)^4}{2d} \\ & -\frac{b^2(c + dx)^4}{4d} + \frac{3b^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\ & -\frac{2ab(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\ & -\frac{3ib^2d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\ & +\frac{3iabd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & +\frac{3b^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\ & -\frac{3abd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\ & -\frac{3iabd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} \\ & +\frac{b^2(c + dx)^3 \tan(e + fx)}{f} \end{aligned}$$

output

```

-I*b^2*(d*x+c)^3/f+1/4*a^2*(d*x+c)^4/d+1/2*I*a*b*(d*x+c)^4/d-1/4*b^2*(d*x+c)^4/d+3*b^2*d*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f^2-2*a*b*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f-3*I*b^2*d^2*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^3+3*I*a*b*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2+3/2*b^2*d^3*polylog(3,-exp(2*I*(f*x+e)))/f^4-3*a*b*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3-3/2*I*a*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4+b^2*(d*x+c)^3*tan(f*x+e)/f

```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1337 vs. $2(300) = 600$.

Time = 7.17 (sec) , antiderivative size = 1337, normalized size of antiderivative = 4.46

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

input

```
Integrate[(c + d*x)^3*(a + b*Tan[e + f*x])^2, x]
```

output

```
((I/4)*b^2*d^3*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^4) - ((I/2)*a*b*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^3) - ((I/4)*a*b*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e)) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))] + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))] - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))]*Sec[e])/f^4 + (3*b^2*c^2*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (2*a*b*c^3*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (3*b^2*c*d^2*Csc[e]*((f^2*x^2)/E^((I*ArcTan[Cot[e]]))) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)]) - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]]))] + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]]))])/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^3*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) - (3*a*b*c^2*d*Csc[e]*((f^2*x^2)/E^((I*ArcTan[Cot[e]]))) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[...
```

Rubi [A] (verified)

Time = 0.77 (sec), antiderivative size = 300, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\
 & \quad \downarrow \textcolor{blue}{4205} \\
 & \int (a^2(c + dx)^3 + 2ab(c + dx)^3 \tan(e + fx) + b^2(c + dx)^3 \tan^2(e + fx)) dx
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{2009} \\
 & \frac{a^2(c+dx)^4}{4d} - \frac{3abd^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \\
 & \frac{3iabd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} + \frac{iab(c+dx)^4}{2d} - \\
 & \frac{3iabd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \\
 & \frac{3b^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f} - \frac{ib^2(c+dx)^3}{f} - \frac{b^2(c+dx)^4}{4d} + \\
 & \frac{3b^2d^3 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^4}
 \end{aligned}$$

input `Int[(c + d*x)^3*(a + b*Tan[e + f*x])^2, x]`

output

$$\begin{aligned}
 & ((-I)*b^2*(c + d*x)^3)/f + (a^2*(c + d*x)^4)/(4*d) + ((I/2)*a*b*(c + d*x)^4)/d - (b^2*(c + d*x)^4)/(4*d) + (3*b^2*d*(c + d*x)^2*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f^2 - (2*a*b*(c + d*x)^3*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f - ((3*I)*b^2*d^2*(c + d*x)*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^3 + ((3*I)*a*b*d*(c + d*x)^2*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 + (3*b^2*d^3*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/f^4 - (3*a*b*d^2*(c + d*x)*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/f^3 - (((3*I)/2)*a*b*d^3*\operatorname{PolyLog}[4, -E^{((2*I)*(e + f*x))}])/f^4 + (b^2*(c + d*x)^3*\operatorname{Tan}[e + f*x])/f
 \end{aligned}$$

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_.)*(x_.)^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.)), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 951 vs. $2(271) = 542$.

Time = 0.90 (sec) , antiderivative size = 952, normalized size of antiderivative = 3.17

method	result	size
risch	Expression too large to display	952

input `int((d*x+c)^3*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output

```

12*I/f*b*d*c^2*a*e*x+d^2*a^2*c*x^3+3/2*d*a^2*c^2*x^2+a^2*c^3*x-d^2*b^2*c*x
^3-3/2*d*b^2*c^2*x^2-12*I/f^2*b*a*c*d^2*e^2*x+6*I/f^2*b*a*c*d^2*polylog(2,
-exp(2*I*(f*x+e)))*x+2*I*b^2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/f/(1+exp(
2*I*(f*x+e)))+1/4*d^3*a^2*x^4+1/4/d*a^2*c^4-1/4*d^3*b^2*x^4-b^2*c^3*x-1/4/
d*b^2*c^4+1/2*I*d^3*a*b*x^4-2*I*a*b*c^3*x-1/2*I/d*a*b*c^4+4/f*b*a*c^3*ln(e
xp(I*(f*x+e)))-2/f*b*a*c^3*ln(1+exp(2*I*(f*x+e)))-6/f^2*b^2*c^2*d*ln(exp(I
*(f*x+e)))+3/f^2*b^2*c^2*d*ln(1+exp(2*I*(f*x+e)))-6/f^4*b^2*e^2*d^3*ln(exp(
I*(f*x+e)))+3/f^2*b^2*d^3*ln(1+exp(2*I*(f*x+e)))*x^2-2*I/f*b^2*d^3*x^3+4*
I/f^4*b^2*e^3*d^3/2*b^2*d^3*polylog(3,-exp(2*I*(f*x+e)))/f^4-12/f^2*b*e*
a*c^2*d*ln(exp(I*(f*x+e)))-6/f*b*d*c^2*a*ln(1+exp(2*I*(f*x+e)))*x-6/f*b*a*
c*d^2*ln(1+exp(2*I*(f*x+e)))*x^2+12/f^3*b*e^2*a*c*d^2*ln(exp(I*(f*x+e)))+6
*I/f^2*b*d*c^2*a*e^2+3*I/f^2*b*d*c^2*a*polylog(2,-exp(2*I*(f*x+e)))+3*I/f^
2*b*d^3*a*polylog(2,-exp(2*I*(f*x+e)))*x^2-8*I/f^3*b*a*c*d^2*e^3-12*I/f^2*
b^2*d^2*c*e*x+4*I/f^3*b*e^3*d^3*a*x+3*I/f^4*b*e^4*d^3*a-6*I/f*b^2*d^2*c*x^
2-6*I/f^3*b^2*d^2*c*e^2-3*I/f^3*b^2*d^2*c*polylog(2,-exp(2*I*(f*x+e)))+6*I
/f^3*b^2*e^2*d^3*x-3*I/f^3*b^2*d^3*polylog(2,-exp(2*I*(f*x+e)))*x+2*I*d^2*
a*b*c*x^3+3*I*d*a*b*c^2*x^2-2/f*b*d^3*a*ln(1+exp(2*I*(f*x+e)))*x^3-4/f^4*b
*e^3*a*d^3*ln(exp(I*(f*x+e)))-3/f^3*b*a*c*d^2*polylog(3,-exp(2*I*(f*x+e)))
-3/f^3*b*d^3*a*polylog(3,-exp(2*I*(f*x+e)))*x+6/f^2*b^2*d^2*c*ln(1+exp(2*I
*(f*x+e)))*x+12/f^3*b^2*e*c*d^2*ln(exp(I*(f*x+e)))-3/2*I*a*b*d^3*polylo...

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 759 vs. $2(264) = 528$.

Time = 0.10 (sec) , antiderivative size = 759, normalized size of antiderivative = 2.53

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

```
1/4*((a^2 - b^2)*d^3*f^4*x^4 + 4*(a^2 - b^2)*c*d^2*f^4*x^3 + 6*(a^2 - b^2)
*c^2*d*f^4*x^2 + 4*(a^2 - b^2)*c^3*f^4*x + 3*I*a*b*d^3*polylog(4, (tan(f*x
+ e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*I*a*b*d^3*polylo
g(4, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(I*
a*b*d^3*f^2*x^2 + I*a*b*c^2*d*f^2 - I*b^2*c*d^2*f + I*(2*a*b*c*d^2*f^2 - b
^2*d^3*f)*x)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(-
I*a*b*d^3*f^2*x^2 - I*a*b*c^2*d*f^2 + I*b^2*c*d^2*f - I*(2*a*b*c*d^2*f^2 -
b^2*d^3*f)*x)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 2
*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 - 3*b^2*c^2*d*f^2 + 3*(2*a*b*c*d^2*f^3
- b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f^3 - b^2*c*d^2*f^2)*x)*log(-2*(I*tan(f
*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3
- 3*b^2*c^2*d*f^2 + 3*(2*a*b*c*d^2*f^3 - b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f
^3 - b^2*c*d^2*f^2)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1))
- 3*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f - b^2*d^3)*polylog(3, (tan(f*x + e)^2 +
2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(2*a*b*d^3*f*x + 2*a*b*c*
d^2*f - b^2*d^3)*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f
*x + e)^2 + 1)) + 4*(b^2*d^3*f^3*x^3 + 3*b^2*c*d^2*f^3*x^2 + 3*b^2*c^2*d*f
^3*x + b^2*c^3*f^3)*tan(f*x + e))/f^4
```

Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^3 dx$$

input `integrate((d*x+c)**3*(a+b*tan(f*x+e))**2,x)`

output `Integral((a + b*tan(e + f*x))**2*(c + d*x)**3, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2525 vs. $2(264) = 528$.

Time = 0.78 (sec), antiderivative size = 2525, normalized size of antiderivative = 8.42

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output

```
1/4*(4*(f*x + e)*a^2*c^3 + (f*x + e)^4*a^2*d^3/f^3 - 4*(f*x + e)^3*a^2*d^3
*e/f^3 + 6*(f*x + e)^2*a^2*d^3*e^2/f^3 - 4*(f*x + e)*a^2*d^3*e^3/f^3 + 4*(f*x + e)^3*a^2*c*d^2/f^2 - 12*(f*x + e)^2*a^2*c*d^2*e/f^2 + 12*(f*x + e)*a^2*c*d^2*e^2/f^2 + 6*(f*x + e)^2*a^2*c^2*d/f - 12*(f*x + e)*a^2*c^2*d*e/f + 8*a*b*c^3*log(sec(f*x + e)) - 8*a*b*d^3*e^3*log(sec(f*x + e))/f^3 + 24*a*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 24*a*b*c^2*d*e*log(sec(f*x + e))/f + 4*(3*(2*a*b + I*b^2)*(f*x + e)^4*d^3 - 24*b^2*d^3*e^3 + 72*b^2*c*d^2*e^2*f - 72*b^2*c^2*d*e*f^2 + 24*b^2*c^3*f^3 - 12*((2*a*b + I*b^2)*d^3*e - (2*a*b + I*b^2)*c*d^2*f)*(f*x + e)^3 + 18*((2*a*b + I*b^2)*d^3*e^2 - 2*(2*a*b + I*b^2)*c*d^2*e*f + (2*a*b + I*b^2)*c^2*d*f^2)*(f*x + e)^2 + 12*(-I*b^2*d^3*e^3 + 3*I*b^2*c*d^2*e^2*f - 3*I*b^2*c^2*d*e*f^2 + I*b^2*c^3*f^3)*(f*x + e) - 4*(8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e*f - 9*b^2*c^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 + 18*(a*b*d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f)*(f*x + e) + (8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e*f - 9*b^2*c^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 + 18*(a*b*d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f)*(f*x + e))*cos(2*f*x + 2*e) - (-8*I*(f*x + e)^3*a*b*d^3 + 9*I*b^2*d^3*e^2 - 18*I*b^2*c*d^2*e*f + 9*I*b^2*c^2*d*f^2 + 9*(2*I*a*b*d^3*e - 2*I*a*b*c*d^2*f + I*b^2*d^3)*(f*x + e)^2 + 18*(-I*a*b*d^3*e^2 - I*a*b*c^2*d*f^2 - I*b^2...)
```

Giac [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (dx + c)^3 (b \tan(fx + e) + a)^2 dx$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^3*(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^3 dx$$

input `int((a + b*tan(e + f*x))^2*(c + d*x)^3,x)`

output `int((a + b*tan(e + f*x))^2*(c + d*x)^3, x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\ &= \frac{8(\int \tan(fx + e) x^3 dx) ab d^3 f^2 + 24(\int \tan(fx + e) x^2 dx) abc d^2 f^2 - 12(\int \tan(fx + e) x^2 dx) b^2 d^3 f + 24abc d^2 f^2}{\dots} \end{aligned}$$

input `int((d*x+c)^3*(a+b*tan(f*x+e))^2,x)`

output

```
(8*int(tan(e + f*x)*x**3,x)*a*b*d**3*f**2 + 24*int(tan(e + f*x)*x**2,x)*a*b*c*d**2*f**2 - 12*int(tan(e + f*x)*x**2,x)*b**2*d**3*f + 24*int(tan(e + f*x)*x,x)*a*b*c**2*d*f**2 - 24*int(tan(e + f*x)*x,x)*b**2*c*d**2*f + 4*log(tan(e + f*x)**2 + 1)*a*b*c**3*f - 6*log(tan(e + f*x)**2 + 1)*b**2*c**2*d + 4*tan(e + f*x)*b**2*c**3*f + 12*tan(e + f*x)*b**2*c**2*d*f*x + 12*tan(e + f*x)*b**2*c*d**2*f*x**2 + 4*tan(e + f*x)*b**2*d**3*f*x**3 + 4*a**2*c**3*f**2*x**2 + 6*a**2*c**2*d*f**2*x**2 + 4*a**2*c*d**2*f**2*x**3 + a**2*d**3*f**2*x**4 - 4*b**2*c**3*f**2*x - 6*b**2*c**2*d*f**2*x**2 - 4*b**2*c*d**2*f**2*x**3 - b**2*d**3*f**2*x**4)/(4*f**2)
```

3.45 $\int (c + dx)^2(a + b \tan(e + fx))^2 dx$

Optimal result	358
Mathematica [B] (warning: unable to verify)	359
Rubi [A] (verified)	360
Maple [B] (verified)	362
Fricas [B] (verification not implemented)	362
Sympy [F]	363
Maxima [B] (verification not implemented)	363
Giac [F]	364
Mupad [F(-1)]	365
Reduce [F]	365

Optimal result

Integrand size = 20, antiderivative size = 229

$$\begin{aligned} \int (c + dx)^2(a + b \tan(e + fx))^2 dx = & -\frac{ib^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} + \frac{2iab(c + dx)^3}{3d} \\ & -\frac{b^2(c + dx)^3}{3d} + \frac{2b^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\ & -\frac{2ab(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & -\frac{ib^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\ & +\frac{2iabd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & -\frac{abd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\ & +\frac{b^2(c + dx)^2 \tan(e + fx)}{f} \end{aligned}$$

output

```
-I*b^2*(d*x+c)^2/f+1/3*a^2*(d*x+c)^3/d+2/3*I*a*b*(d*x+c)^3/d-1/3*b^2*(d*x+c)^3/d+2*b^2*d*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^2-2*a*b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f-I*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3+2*I*a*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-a*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+b^2*(d*x+c)^2*tan(f*x+e)/f
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 649 vs. $2(229) = 458$.

Time = 6.72 (sec) , antiderivative size = 649, normalized size of antiderivative = 2.83

$$\begin{aligned}
& \int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \\
& - \frac{iabd^2 e^{-ie} (2f^2 x^2 (2fx - 3i(1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6(1 + e^{2ie}) fx \operatorname{PolyLog}(2, -e^{-2i(e+fx)}) - 3)}{6f^3} \\
& + \frac{1}{3} x (3c^2 + 3cdx + d^2 x^2) \sec(e) (a^2 \cos(e) - b^2 \cos(e) + 2ab \sin(e)) \\
& + \frac{2b^2 cd \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + fx \sin(e))}{f^2 (\cos^2(e) + \sin^2(e))} \\
& - \frac{2abc^2 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + fx \sin(e))}{f (\cos^2(e) + \sin^2(e))} \\
& + \frac{b^2 d^2 \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (ifx(-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2ifx}) - 2(fx - \arctan(\cot(e))) \log(1 - e^{2i(fx - \arctan(\cot(e)))}))}{f^3 \sqrt{\csc^2(e) (\cos^2(e))}} \right)}{2abcd \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (ifx(-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2ifx}) - 2(fx - \arctan(\cot(e))) \log(1 - e^{2i(fx - \arctan(\cot(e)))}))}{f^2 \sqrt{\csc^2(e) (\cos^2(e))}} \right)} \\
& + \frac{\sec(e) \sec(e + fx) (b^2 c^2 \sin(fx) + 2b^2 cdx \sin(fx) + b^2 d^2 x^2 \sin(fx))}{f}
\end{aligned}$$

```
input Integrate[(c + d*x)^2*(a + b*Tan[e + f*x])^2,x]
```

output

$$\begin{aligned}
 & ((-1/6*I)*a*b*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^{((2*I)*e))})*Log[1 + E^{((-2*I)*(e + f*x))}] + 6*(1 + E^{((2*I)*e))}*f*x*PolyLog[2, -E^{((-2*I)*(e + f*x))}] - (3*I)*(1 + E^{((2*I)*e))}*PolyLog[3, -E^{((-2*I)*(e + f*x))}]*Sec[e])/(E^{(I*e)*f^3}) + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*Sec[e]*(a^2*Cos[e] - b^2*Cos[e] + 2*a*b*Sin[e]))/3 + (2*b^2*c*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (2*a*b*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^2*d^2*2*Csc[e]*((f^2*x^2)/E^{(I*ArcTan[Cot[e]]}) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^{((-2*I)*f*x)}] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^{((2*I)*(f*x - ArcTan[Cot[e]]))}] + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^{((2*I)*(f*x - ArcTan[Cot[e]]))}])/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^3*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) - (2*a*b*c*d*Csc[e]*((f^2*x^2)/E^{(I*ArcTan[Cot[e]]}) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^{((-2*I)*f*x)}] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^{((2*I)*(f*x - ArcTan[Cot[e]]))}] + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^{((2*I)*(f*x - ArcTan[Cot[e]]))}])/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^2*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) + (Sec[e]*Sec[e + f*x]*(b^2*c^2*Sin[f*x] + 2*b^2*c*d*x*Sin[f*x] + b^2*d^2*x^2*Sin[f*x]))/f
 \end{aligned}$$

Rubi [A] (verified)

Time = 0.61 (sec), antiderivative size = 229, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^2 (a + b \tan(e + fx))^2 dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int (c + dx)^2 (a + b \tan(e + fx))^2 dx \\
 & \quad \downarrow \textcolor{blue}{4205} \\
 & \int (a^2(c + dx)^2 + 2ab(c + dx)^2 \tan(e + fx) + b^2(c + dx)^2 \tan^2(e + fx)) dx
 \end{aligned}$$

$$\begin{aligned}
 & \frac{a^2(c+dx)^3}{3d} + \frac{2iabd(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} + \\
 & \frac{2iab(c+dx)^3}{3d} - \frac{abd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{2b^2d(c+dx) \log(1+e^{2i(e+fx)})}{f^2} + \\
 & \frac{b^2(c+dx)^2 \tan(e+fx)}{f} - \frac{ib^2(c+dx)^2}{f} - \frac{b^2(c+dx)^3}{3d} - \frac{ib^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3}
 \end{aligned}$$

↓ 2009

input `Int[(c + d*x)^2*(a + b*Tan[e + f*x])^2, x]`

output `((-I)*b^2*(c + d*x)^2)/f + (a^2*(c + d*x)^3)/(3*d) + (((2*I)/3)*a*b*(c + d*x)^3)/d - (b^2*(c + d*x)^3)/(3*d) + (2*b^2*d*(c + d*x))*Log[1 + E^((2*I)*(e + f*x))]/f^2 - (2*a*b*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f - (I*b^2*d^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^3 + ((2*I)*a*b*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (a*b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/f^3 + (b^2*(c + d*x)^2*Tan[e + f*x])/f`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 574 vs. $2(209) = 418$.

Time = 0.86 (sec) , antiderivative size = 575, normalized size of antiderivative = 2.51

method	result
risch	$da^2cx^2 + a^2c^2x - db^2cx^2 + \frac{d^2a^2x^3}{3} + \frac{a^2c^3}{3d} - \frac{d^2b^2x^3}{3} - b^2c^2x - \frac{b^2c^3}{3d} - \frac{8beacd\ln(e^{i(fx+e)})}{f^2} - \frac{4bdca\ln(1+j)}{f^2}$

input `int((d*x+c)^2*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 2*I*b^2*(d^2*x^2+2*c*d*x+c^2)/f/(1+\exp(2*I*(f*x+e)))+d*a^2*c*x^2+a^2*c^2*x \\ & -d*b^2*c*x^2+1/3*d^2*a^2*x^3+1/3/d*a^2*c^3-1/3*d^2*b^2*x^3-b^2*c^2*x-1/3/d \\ & *b^2*c^3+8*I/f*b*d*c*a*e*x-8/f^2*2*b*e*a*c*d*\ln(\exp(I*(f*x+e)))-4/f*b*d*c*a \\ & \ln(1+\exp(2*I*(f*x+e)))*x+2*I/f^2*2*b*a*d^2*polylog(2,-\exp(2*I*(f*x+e)))*x-4*I \\ & /f^2*2*b*a*d^2*e^2*x+4*I/f^2*2*b*d*c*a*e^2+2*I/f^2*2*b*d*c*a*polylog(2,-\exp(2*I \\ & *(f*x+e)))-I*b^2*d^2*polylog(2,-\exp(2*I*(f*x+e)))/f^3+2/3*I*d^2*a*b*x^3+2/f \\ & ^2*b^2*d^2*\ln(1+\exp(2*I*(f*x+e)))*x-4/f^2*b^2*c*d*\ln(\exp(I*(f*x+e)))+2/f^ \\ & 2*b^2*c*d*\ln(1+\exp(2*I*(f*x+e)))+4/f*b*a*c^2*\ln(\exp(I*(f*x+e)))-2/f*b*a*c \\ & ^2*\ln(1+\exp(2*I*(f*x+e)))+4/f^3*b^2*e*d^2*\ln(\exp(I*(f*x+e)))-2*I/f^3*b^2*d^2 \\ & 2*e^2-2*I/f*b^2*d^2*x^2-2*I*a*b*c^2*x-2/3*I/d*a*b*c^3+4/f^3*b*e^2*a*d^2*ln \\ & (\exp(I*(f*x+e)))-2/f*b*a*d^2*\ln(1+\exp(2*I*(f*x+e)))*x^2-4*I/f^2*b^2*d^2*x \\ & -8/3*I/f^3*b*a*d^2*e^3+2*I*d*a*b*c*x^2-a*b*d^2*polylog(3,-\exp(2*I*(f*x+e))) \\ &)/f^3 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(203) = 406$.

Time = 0.09 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.97

$$\begin{aligned} & \int (c + dx)^2(a + b \tan(e + fx))^2 dx \\ & = \frac{2(a^2 - b^2)d^2 f^3 x^3 + 6(a^2 - b^2)c d f^3 x^2 + 6(a^2 - b^2)c^2 f^3 x - 3abd^2 \text{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - }{ } \end{aligned}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output
$$\begin{aligned} & \frac{1}{6} \cdot (2(a^2 - b^2) \cdot d^2 \cdot f^3 \cdot x^3 + 6(a^2 - b^2) \cdot c \cdot d \cdot f^3 \cdot x^2 + 6(a^2 - b^2) \\ & \cdot c^2 \cdot f^3 \cdot x - 3 \cdot a \cdot b \cdot d^2 \cdot \text{polylog}(3, (\tan(f \cdot x + e))^2 + 2 \cdot I \cdot \tan(f \cdot x + e) - 1) / \\ & (\tan(f \cdot x + e)^2 + 1)) - 3 \cdot a \cdot b \cdot d^2 \cdot \text{polylog}(3, (\tan(f \cdot x + e))^2 - 2 \cdot I \cdot \tan(f \cdot x + e) - 1) / \\ & (\tan(f \cdot x + e)^2 + 1)) - 3 \cdot (2 \cdot I \cdot a \cdot b \cdot d^2 \cdot f^2 \cdot x + 2 \cdot I \cdot a \cdot b \cdot c \cdot d \cdot f - I \cdot b^2 \cdot d^2) \cdot \text{dilog}(2 \cdot (I \cdot \tan(f \cdot x + e) - 1) / (\tan(f \cdot x + e)^2 + 1) + 1) - 3 \cdot (-2 \cdot I \cdot a \cdot b \cdot d^2 \cdot f \cdot x - 2 \cdot I \cdot a \cdot b \cdot c \cdot d \cdot f + I \cdot b^2 \cdot d^2) \cdot \text{dilog}(2 \cdot (-I \cdot \tan(f \cdot x + e) - 1) / (\tan(f \cdot x + e)^2 + 1) + 1) - 6 \cdot (a \cdot b \cdot d^2 \cdot f^2 \cdot x^2 + a \cdot b \cdot c^2 \cdot f^2 - b^2 \cdot c \cdot d \cdot f + (2 \cdot a \cdot b \cdot c \cdot d \cdot f^2 - b^2 \cdot d^2 \cdot 2 \cdot f) \cdot x) \cdot \log(-2 \cdot (I \cdot \tan(f \cdot x + e) - 1) / (\tan(f \cdot x + e)^2 + 1)) - 6 \cdot (a \cdot b \cdot d^2 \cdot f^2 \cdot x^2 + a \cdot b \cdot c^2 \cdot f^2 - b^2 \cdot c \cdot d \cdot f + (2 \cdot a \cdot b \cdot c \cdot d \cdot f^2 - b^2 \cdot d^2 \cdot 2 \cdot f) \cdot x) \cdot \log(-2 \cdot (-I \cdot \tan(f \cdot x + e) - 1) / (\tan(f \cdot x + e)^2 + 1)) + 6 \cdot (b^2 \cdot d^2 \cdot f^2 \cdot x^2 + 2 \cdot b^2 \cdot c \cdot d \cdot f^2 \cdot x + b^2 \cdot c^2 \cdot f^2) \cdot \tan(f \cdot x + e)) / f^3 \end{aligned}$$

Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^2 dx$$

input `integrate((d*x+c)**2*(a+b*tan(f*x+e))**2,x)`

output `Integral((a + b*tan(e + f*x))**2*(c + d*x)**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1263 vs. $2(203) = 406$.

Time = 0.32 (sec), antiderivative size = 1263, normalized size of antiderivative = 5.52

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output

```
1/3*(3*(f*x + e)*a^2*c^2 + (f*x + e)^3*a^2*d^2/f^2 - 3*(f*x + e)^2*a^2*d^2
*e/f^2 + 3*(f*x + e)*a^2*d^2*2*e^2/f^2 + 3*(f*x + e)^2*a^2*c*d/f - 6*(f*x +
e)*a^2*c*d*e/f + 6*a*b*c^2*log(sec(f*x + e)) + 6*a*b*d^2*e^2*log(sec(f*x +
e))/f^2 - 12*a*b*c*d*e*log(sec(f*x + e))/f + 3*((2*a*b + I*b^2)*(f*x + e)
^3*d^2 + 6*b^2*d^2*2*e^2 - 12*b^2*c*d*e*f + 6*b^2*c^2*f^2 - 3*((2*a*b + I*b^
2)*d^2*e - (2*a*b + I*b^2)*c*d*f)*(f*x + e)^2 + 3*(I*b^2*d^2*2*e^2 - 2*I*b^2
*c*d*e*f + I*b^2*c^2*f^2)*(f*x + e) - 6*((f*x + e)^2*a*b*d^2 + b^2*d^2*e -
b^2*c*d*f - (2*a*b*d^2*e - 2*a*b*c*d*f + b^2*d^2)*(f*x + e) + ((f*x + e)
^2*a*b*d^2 + b^2*d^2*e - b^2*c*d*f - (2*a*b*d^2*e - 2*a*b*c*d*f + b^2*d^2)*
(f*x + e))*cos(2*f*x + 2*e) - (-I*(f*x + e)^2*a*b*d^2 - I*b^2*d^2*e + I*b^
2*c*d*f + (2*I*a*b*d^2*e - 2*I*a*b*c*d*f + I*b^2*d^2)*(f*x + e))*sin(2*f*x
+ 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + ((2*a*b + I*b^2
)*(f*x + e)^3*d^2 - 3*(2*b^2*d^2 + (2*a*b + I*b^2)*d^2*e - (2*a*b + I*b^2)
*c*d*f)*(f*x + e)^2 + 3*(I*b^2*d^2*2*e^2 + I*b^2*c^2*f^2 + 4*b^2*d^2*e + 2*(-
I*b^2*c*d*e - 2*b^2*c*d*f)*(f*x + e))*cos(2*f*x + 2*e) + 3*(2*(f*x + e)*
a*b*d^2 - 2*a*b*d^2*e + 2*a*b*c*d*f - b^2*d^2 + (2*(f*x + e)*a*b*d^2 - 2*a
*b*d^2*e + 2*a*b*c*d*f - b^2*d^2)*cos(2*f*x + 2*e) + (2*I*(f*x + e)*a*b*d^
2 - 2*I*a*b*d^2*e + 2*I*a*b*c*d*f - I*b^2*d^2)*sin(2*f*x + 2*e))*dilog(-e^
(2*I*f*x + 2*I*e)) + 3*(I*(f*x + e)^2*a*b*d^2 + I*b^2*d^2*e - I*b^2*c*d*f
+ (-2*I*a*b*d^2*e + 2*I*a*b*c*d*f - I*b^2*d^2)*(f*x + e) + (I*(f*x + e)...)
```

Giac [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (dx + c)^2 (b \tan(fx + e) + a)^2 dx$$

input

```
integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="giac")
```

output

```
integrate((d*x + c)^2*(b*tan(f*x + e) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^2 dx$$

input `int((a + b*tan(e + f*x))^2*(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))^2*(c + d*x)^2, x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)^2 (a + b \tan(e + fx))^2 dx \\ &= \frac{6(\int \tan(fx + e) x^2 dx) ab d^2 f^2 + 12(\int \tan(fx + e) x dx) abcd f^2 - 6(\int \tan(fx + e) x dx) b^2 d^2 f + 3 \log \end{aligned}$$

input `int((d*x+c)^2*(a+b*tan(f*x+e))^2,x)`

output `(6*int(tan(e + f*x)*x**2,x)*a*b*d**2*f**2 + 12*int(tan(e + f*x)*x,x)*a*b*c*d*f**2 - 6*int(tan(e + f*x)*x,x)*b**2*d**2*f + 3*log(tan(e + f*x)**2 + 1)*a*b*c**2*f - 3*log(tan(e + f*x)**2 + 1)*b**2*c*d + 3*tan(e + f*x)*b**2*c**2*f + 6*tan(e + f*x)*b**2*c*d*f*x + 3*tan(e + f*x)*b**2*d**2*f*x**2 + 3*a**2*c**2*f**2*x + 3*a**2*c*d*f**2*x**2 + a**2*d**2*f**2*x**3 - 3*b**2*c**2*f**2*x - 3*b**2*c*d*f**2*x**2 - b**2*d**2*f**2*x**3)/(3*f**2)`

3.46 $\int (c + dx)(a + b \tan(e + fx))^2 dx$

Optimal result	366
Mathematica [A] (verified)	367
Rubi [A] (verified)	367
Maple [A] (verified)	368
Fricas [A] (verification not implemented)	369
Sympy [F]	369
Maxima [B] (verification not implemented)	370
Giac [F]	370
Mupad [F(-1)]	371
Reduce [F]	371

Optimal result

Integrand size = 18, antiderivative size = 135

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^2 dx = & \frac{a^2(c + dx)^2}{2d} + \frac{iab(c + dx)^2}{d} - \frac{b^2(c + dx)^2}{2d} \\ & - \frac{2ab(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^2 d \log(\cos(e + fx))}{f^2} \\ & + \frac{iabd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & + \frac{b^2(c + dx) \tan(e + fx)}{f} \end{aligned}$$

output
$$\frac{1}{2}a^2(d*x+c)^2/d + I*a*b*(d*x+c)^2/d - 1/2*b^2*(d*x+c)^2/d - 2*a*b*(d*x+c)*\ln(1+\exp(2*I*(f*x+e)))/f + b^2*d*\ln(\cos(f*x+e))/f^2 + I*a*b*d*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))/f^2 + b^2*(d*x+c)*\tan(f*x+e)/f$$

Mathematica [A] (verified)

Time = 7.17 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.48

$$\int (c + dx)(a + b \tan(e + fx))^2 dx \\ = \frac{\cos(e + fx) (\cos(e + fx) ((-((e + fx) (-2iabd(e + fx) + a^2(de - 2cf - dfx) + b^2(-de + 2cf + dfx))))$$

input `Integrate[(c + d*x)*(a + b*Tan[e + f*x])^2, x]`

output $(\cos[e + f*x]*(\cos[e + f*x]*(-((e + f*x)*((-2*I)*a*b*d*(e + f*x) + a^2*(d*e - 2*c*f - d*f*x) + b^2*(-(d*e) + 2*c*f + d*f*x))) - 4*a*b*d*(e + f*x)*\text{Log}[1 + E^{((2*I)*(e + f*x))}] + 2*b*(b*d + 2*a*d*e - 2*a*c*f)*\text{Log}[\cos[e + f*x]] + (2*I)*a*b*d*\cos[e + f*x]*\text{PolyLog}[2, -E^{((2*I)*(e + f*x))}] + 2*b^2*f*(c + d*x)*\sin[e + f*x]*(a + b*\tan[e + f*x])^2)/(2*f^2*(a*\cos[e + f*x] + b*\sin[e + f*x])^2)$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)(a + b \tan(e + fx))^2 dx \\ \downarrow 3042 \\ \int (c + dx)(a + b \tan(e + fx))^2 dx \\ \downarrow 4205 \\ \int (a^2(c + dx) + 2ab(c + dx)\tan(e + fx) + b^2(c + dx)\tan^2(e + fx)) dx \\ \downarrow 2009$$

$$\frac{a^2(c+dx)^2}{2d} - \frac{2ab(c+dx)\log(1+e^{2i(e+fx)})}{f} + \frac{iab(c+dx)^2}{d} + \frac{iabd\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \\ \frac{b^2(c+dx)\tan(e+fx)}{f} - \frac{b^2(c+dx)^2}{2d} + \frac{b^2d\log(\cos(e+fx))}{f^2}$$

input `Int[(c + d*x)*(a + b*Tan[e + f*x])^2, x]`

output `(a^2*(c + d*x)^2)/(2*d) + (I*a*b*(c + d*x)^2)/d - (b^2*(c + d*x)^2)/(2*d) - (2*a*b*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f + (b^2*d*Log[Cos[e + f*x]])/f^2 + (I*a*b*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (b^2*(c + d*x)*Tan[e + f*x])/f`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4205 `Int[((c_.) + (d_)*(x_))^(m_.)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]`

Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.76

method	result
risch	$-\frac{b^2 dx^2}{2} + \frac{2ibad e^2}{f^2} + \frac{a^2 dx^2}{2} - b^2 cx + \frac{2ib^2 (dx+c)}{f(1+e^{2i(fx+e)})} + a^2 cx + iabd x^2 - \frac{2b^2 d \ln(e^{i(fx+e)})}{f^2} + \frac{b^2 d \ln(1+e^{2i(fx+e)})}{f^2}$

input `int((d*x+c)*(a+b*tan(f*x+e))^2, x, method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -\frac{1}{2} b^2 d x^2 + 2 I f^2 b^2 a^2 d e^2 + \frac{1}{2} a^2 d x^2 - b^2 c x + 2 I b^2 (d x + c) / f / (\\ & 1 + \exp(2 I (f x + e))) + a^2 c x + I a b d x^2 - 2 / f^2 b^2 d \ln(\exp(I (f x + e))) + 1 / f \\ & ^2 b^2 d \ln(1 + \exp(2 I (f x + e))) + 4 / f b^2 a c \ln(\exp(I (f x + e))) - 2 / f b^2 a c \ln(\\ & 1 + \exp(2 I (f x + e))) - 4 / f^2 b^2 e a d \ln(\exp(I (f x + e))) - 2 I a b c x + I a b d p \\ & \text{olylog}(2, -\exp(2 I (f x + e))) / f^2 + 4 I f b^2 a d e x - 2 / f b^2 a d \ln(1 + \exp(2 I (f \\ & x + e))) * x \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 219, normalized size of antiderivative = 1.62

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx))^2 dx \\ & = \frac{(a^2 - b^2) df^2 x^2 + 2 (a^2 - b^2) cf^2 x - i ab d \text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1}+1\right) + i ab d \text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1}+1\right) - (2 a^2 - b^2) d \ln(-2 \tan(fx+e))}{f^2} \end{aligned}$$

input

```
integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="fricas")
```

output

$$\begin{aligned} & \frac{1}{2} ((a^2 - b^2) d f^2 x^2 + 2 (a^2 - b^2) c f^2 x - I a b d \text{dilog}(2 (I \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1) + 1) + I a b d \text{dilog}(2 (-I \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1) + 1) - (2 a b d f x + 2 a b c f - b^2 d) \log(-2 (-I \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1)) - (2 a b d f x + 2 a b c f - b^2 d) \log(-2 (-I \tan(f x + e) - 1) / (\tan(f x + e)^2 + 1)) + 2 (b^2 d f x + b^2 c f) \tan(f x + e)) / f^2 \end{aligned}$$

Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx) dx$$

input

```
integrate((d*x+c)*(a+b*tan(f*x+e))**2,x)
```

output

```
Integral((a + b*tan(e + f*x))**2*(c + d*x), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs. $2(122) = 244$.

Time = 0.18 (sec) , antiderivative size = 527, normalized size of antiderivative = 3.90

$$\int (c + dx)(a + b \tan(e + fx))^2 dx \\ = \frac{2(fx + e)a^2c + \frac{(fx+e)^2a^2d}{f} - \frac{2(fx+e)a^2de}{f} + 4abc \log(\sec(fx + e)) - \frac{4abde \log(\sec(fx+e))}{f} + \frac{2((2ab+ib^2)(fx+e)^2)}{f}}{1}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output
$$\begin{aligned} & 1/2*(2*(f*x + e)*a^2*c + (f*x + e)^2*a^2*d/f - 2*(f*x + e)*a^2*d*e/f + 4*a \\ & *b*c*log(\sec(f*x + e)) - 4*a*b*d*e*log(\sec(f*x + e))/f + 2*((2*a*b + I*b^2) \\ & *(f*x + e)^2*d - 4*b^2*d*e + 4*b^2*c*f + 2*(-I*b^2*d*e + I*b^2*c*f)*(f*x \\ & + e) - 2*(2*(f*x + e)*a*b*d - b^2*d + (2*(f*x + e)*a*b*d - b^2*d)*cos(2*f*x \\ & + 2*e) - (-2*I*(f*x + e)*a*b*d + I*b^2*d)*sin(2*f*x + 2*e))*arctan2(sin(\\ & 2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + ((2*a*b + I*b^2)*(f*x + e)^2*d + 2*(\\ & -I*b^2*d*e + I*b^2*c*f - 2*b^2*d)*(f*x + e))*cos(2*f*x + 2*e) + 2*(a*b*d*c \\ & os(2*f*x + 2*e) + I*a*b*d*sin(2*f*x + 2*e) + a*b*d)*dilog(-e^(2*I*f*x + 2* \\ & I*e)) - (-2*I*(f*x + e)*a*b*d + I*b^2*d + (-2*I*(f*x + e)*a*b*d + I*b^2*d) \\ & *cos(2*f*x + 2*e) + (2*(f*x + e)*a*b*d - b^2*d)*sin(2*f*x + 2*e))*log(cos(\\ & 2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) - ((-2*I*a*b \\ & + b^2)*(f*x + e)^2*d - 2*(b^2*d*e - b^2*c*f - 2*I*b^2*d)*(f*x + e))*sin(2 \\ & *f*x + 2*e))/(-2*I*f*cos(2*f*x + 2*e) + 2*f*sin(2*f*x + 2*e) - 2*I*f)/f \end{aligned}$$

Giac [F]

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (dx + c)(b \tan(fx + e) + a)^2 dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)*(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + d x) dx$$

input `int((a + b*tan(e + f*x))^2*(c + d*x),x)`

output `int((a + b*tan(e + f*x))^2*(c + d*x), x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx))^2 dx \\ &= \frac{4(\int \tan(fx + e) x dx) abd f^2 + 2 \log(\tan(fx + e)^2 + 1) abc f - \log(\tan(fx + e)^2 + 1) b^2 d + 2 \tan(fx + e) abcd f^2}{2f^2} \end{aligned}$$

input `int((d*x+c)*(a+b*tan(f*x+e))^2,x)`

output `(4*int(tan(e + f*x)*x,x)*a*b*d*f**2 + 2*log(tan(e + f*x)**2 + 1)*a*b*c*f - log(tan(e + f*x)**2 + 1)*b**2*d + 2*tan(e + f*x)*b**2*c*f + 2*tan(e + f*x)*b**2*d*f*x + 2*a**2*c*f**2*x + a**2*d*f**2*x**2 - 2*b**2*c*f**2*x - b**2*d*f**2*x**2)/(2*f**2)`

3.47 $\int \frac{(a+b\tan(e+fx))^2}{c+dx} dx$

Optimal result	372
Mathematica [N/A]	372
Rubi [N/A]	373
Maple [N/A]	374
Fricas [N/A]	374
Sympy [N/A]	374
Maxima [N/A]	375
Giac [N/A]	375
Mupad [N/A]	376
Reduce [N/A]	376

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^2}{c + dx}, x\right)$$

output `Defer(Int)((a+b*tan(f*x+e))^2/(d*x+c),x)`

Mathematica [N/A]

Not integrable

Time = 19.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

input `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x), x]`

output `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x), x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx \end{aligned}$$

input `Int[(a + b*Tan[e + f*x])^2/(c + d*x), x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.47 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^2}{dx + c} dx$$

input `int((a+b*tan(f*x+e))^2/(d*x+c),x)`

output `int((a+b*tan(f*x+e))^2/(d*x+c),x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="fricas")`

output `integral((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 0.93 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

input `integrate((a+b*tan(f*x+e))**2/(d*x+c),x)`

output $\text{Integral}((a + b\tan(e + fx))^2/(c + dx), x)$

Maxima [N/A]

Not integrable

Time = 0.58 (sec) , antiderivative size = 491, normalized size of antiderivative = 24.55

$$\int \frac{(a + b\tan(e + fx))^2}{c + dx} dx = \int \frac{(b\tan(fx + e) + a)^2}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="maxima")`

output $((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*\cos(2*f*x + 2*e)^2*\log(d*x + c) + 2*b^2*d*\sin(2*f*x + 2*e) + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*\log(d*x + c)*\sin(2*f*x + 2*e)^2 + 2*((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*\cos(2*f*x + 2*e)*\log(d*x + c) + (d^2*f*x + c*d*f + (d^2*f*x + c*d*f))*\cos(2*f*x + 2*e)^2 + (d^2*f*x + c*d*f)*\sin(2*f*x + 2*e)^2 + 2*(d^2*f*x + c*d*f)*\cos(2*f*x + 2*e)*integrate(2*(2*a*b*d*f*x + 2*a*b*c*f + b^2*d)*\sin(2*f*x + 2*e)/(d^2*f*x^2 + 2*c*d*f*x + c^2*f + (d^2*f*x^2 + 2*c*d*f*x + c^2*f))*\cos(2*f*x + 2*e)^2 + (d^2*f*x^2 + 2*c*d*f*x + c^2*f)*\sin(2*f*x + 2*e)^2 + 2*(d^2*f*x^2 + 2*c*d*f*x + c^2*f)*\cos(2*f*x + 2*e)), x) + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*\log(d*x + c)/(d^2*f*x + c*d*f + (d^2*f*x + c*d*f))*\cos(2*f*x + 2*e)^2 + (d^2*f*x + c*d*f)*\sin(2*f*x + 2*e)^2 + 2*(d^2*f*x + c*d*f)*\cos(2*f*x + 2*e))$

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b\tan(e + fx))^2}{c + dx} dx = \int \frac{(b\tan(fx + e) + a)^2}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)^2/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 8.90 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

input `int((a + b*tan(e + f*x))^2/(c + d*x),x)`

output `int((a + b*tan(e + f*x))^2/(c + d*x), x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.95

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx \\ &= \frac{\left(\int \frac{\tan(fx+e)^2}{dx+c} dx \right) b^2 d + 2 \left(\int \frac{\tan(fx+e)}{dx+c} dx \right) abd + \log(dx+c) a^2}{d} \end{aligned}$$

input `int((a+b*tan(f*x+e))^2/(d*x+c),x)`

output `(int(tan(e + f*x)**2/(c + d*x),x)*b**2*d + 2*int(tan(e + f*x)/(c + d*x),x)*a*b*d + log(c + d*x)*a**2)/d`

3.48 $\int \frac{(a+b\tan(e+fx))^2}{(c+dx)^2} dx$

Optimal result	377
Mathematica [N/A]	377
Rubi [N/A]	378
Maple [N/A]	379
Fricas [N/A]	379
Sympy [N/A]	379
Maxima [N/A]	380
Giac [N/A]	380
Mupad [N/A]	381
Reduce [N/A]	381

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^2}{(c + dx)^2}, x\right)$$

output `Defer(Int)((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`

Mathematica [N/A]

Not integrable

Time = 12.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x)^2,x]`

output `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x)^2, x]`

Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx \end{aligned}$$

input `Int[(a + b*Tan[e + f*x])^2/(c + d*x)^2, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^2}{(dx + c)^2} dx$$

input `int((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`

output `int((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="fricas")`

output `integral((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 2.02 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `integrate((a+b*tan(f*x+e))**2/(d*x+c)**2,x)`

output `Integral((a + b*tan(e + f*x))**2/(c + d*x)**2, x)`

Maxima [N/A]

Not integrable

Time = 1.04 (sec) , antiderivative size = 609, normalized size of antiderivative = 30.45

$$\int \frac{(a + b \tan(fx + e))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="maxima")`

output
$$-((a^2 - b^2)*d*f*x - 2*b^2*d*sin(2*f*x + 2*e) + (a^2 - b^2)*c*f + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e)^2 + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*sin(2*f*x + 2*e)^2 + 2*((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e) - (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e))*integrate(4*(a*b*d*f*x + a*b*c*f + b^2*d)*sin(2*f*x + 2*e)/(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f + (d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f)*cos(2*f*x + 2*e)), x))/(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e))$$

Giac [N/A]

Not integrable

Time = 15.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(fx + e))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)^2/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 10.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

input `int((a + b*tan(e + f*x))^2/(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))^2/(c + d*x)^2, x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 157, normalized size of antiderivative = 7.85

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx \\ &= \frac{\left(\int \frac{\tan(fx+e)^2}{d^2x^2+2cdx+c^2} dx \right) b^2 c^2 + \left(\int \frac{\tan(fx+e)^2}{d^2x^2+2cdx+c^2} dx \right) b^2 c d x + 2 \left(\int \frac{\tan(fx+e)}{d^2x^2+2cdx+c^2} dx \right) a b c^2 + 2 \left(\int \frac{\tan(fx+e)}{d^2x^2+2cdx+c^2} dx \right) c (d x + c)}{c (d x + c)} \end{aligned}$$

input `int((a+b*tan(f*x+e))^2/(d*x+c)^2,x)`

output `(int(tan(e + f*x)**2/(c**2 + 2*c*d*x + d**2*x**2),x)*b**2*c**2 + int(tan(e + f*x)**2/(c**2 + 2*c*d*x + d**2*x**2),x)*b**2*c*d*x + 2*int(tan(e + f*x)/(c**2 + 2*c*d*x + d**2*x**2),x)*a*b*c**2 + 2*int(tan(e + f*x)/(c**2 + 2*c*d*x + d**2*x**2),x)*a*b*c*d*x + a**2*x)/(c*(c + d*x))`

3.49 $\int (c + dx)^3 (a + b \tan(e + fx))^3 dx$

Optimal result	383
Mathematica [B] (warning: unable to verify)	384
Rubi [A] (verified)	385
Maple [B] (verified)	387
Fricas [B] (verification not implemented)	388
Sympy [F]	389
Maxima [B] (verification not implemented)	390
Giac [F]	391
Mupad [F(-1)]	391
Reduce [F]	391

Optimal result

Integrand size = 20, antiderivative size = 612

$$\begin{aligned}
 \int (c + dx)^3 (a + b \tan(e + fx))^3 dx = & \frac{3ib^3 d(c + dx)^2}{2f^2} - \frac{3iab^2(c + dx)^3}{f} + \frac{b^3(c + dx)^3}{2f} \\
 & + \frac{a^3(c + dx)^4}{4d} + \frac{3ia^2b(c + dx)^4}{4d} - \frac{3ab^2(c + dx)^4}{4d} \\
 & - \frac{ib^3(c + dx)^4}{4d} - \frac{3b^3d^2(c + dx) \log(1 + e^{2i(e+fx)})}{f^3} \\
 & + \frac{9ab^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{b^3(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{3ib^3d^3 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} \\
 & - \frac{9iab^2d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
 & + \frac{9ia^2bd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
 & - \frac{3ib^3d(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
 & + \frac{9ab^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\
 & - \frac{9a^2bd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & + \frac{3b^3d^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{9ia^2bd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
 & + \frac{3ib^3d^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
 & - \frac{3b^3d(c + dx)^2 \tan(e + fx)}{2f^2} \\
 & + \frac{3ab^2(c + dx)^3 \tan(e + fx)}{f} \\
 & + \frac{b^3(c + dx)^3 \tan^2(e + fx)}{2f}
 \end{aligned}$$

output

```
3/4*I*b^3*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4-3/2*I*b^3*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2+1/2*b^3*(d*x+c)^3/f+1/4*a^3*(d*x+c)^4/d+3/2*I*b^3*d^3*polylog(2,-exp(2*I*(f*x+e)))/f^4-3/4*a*b^2*(d*x+c)^4/d-3*I*a*b^2*(d*x+c)^3/f-3*b^3*d^2*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^3+9*a*b^2*d*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f^2-3*a^2*b*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f+b^3*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f-1/4*I*b^3*(d*x+c)^4/d+9/2*I*a^2*b*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2+3/2*I*b^3*d*(d*x+c)^2/f^2-9/4*I*a^2*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4+9/2*a*b^2*d^3*polylog(3,-exp(2*I*(f*x+e)))/f^3+3/2*b^3*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3+3/4*I*a^2*b*(d*x+c)^4/d-9*I*a*b^2*d^2*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^3-3/2*b^3*d*(d*x+c)^2*tan(f*x+e)/f^2+3*a*b^2*(d*x+c)^3*tan(f*x+e)/f+1/2*b^3*(d*x+c)^3*tan(f*x+e)^2/f
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2594 vs. $2(612) = 1224$.

Time = 7.32 (sec) , antiderivative size = 2594, normalized size of antiderivative = 4.24

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Result too large to show}$$

input

```
Integrate[(c + d*x)^3*(a + b*Tan[e + f*x])^3,x]
```

output

$$\begin{aligned}
 & (((3*I)/4)*a*b^2*d^3*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e)))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^4) - (((3*I)/4)*a^2*b*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e)))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^3) + ((I/4)*b^3*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e)))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^3) - (((3*I)/8)*a^2*b*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e)) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))] + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))] - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))]*Sec[e])/f^4 + ((I/8)*b^3*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e)) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))] + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))] - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))]*Sec[e])/f^4 + ((b^3*c^3 + 3*b^3*c^2*d*x + 3*b^3*c*d^2*x^2 + b^3*d^3*x^3)*Sec[e + f*x]^2)/(2*f) - (3*b^3*c*d^2*Sec[e]*(Cos[e]*Log[Cos[e]...]
 \end{aligned}$$

Rubi [A] (verified)

Time = 1.29 (sec), antiderivative size = 612, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)^3 (a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int (c + dx)^3 (a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \textcolor{blue}{4205}
 \end{aligned}$$

$$\begin{aligned}
 & \int (a^3(c+dx)^3 + 3a^2b(c+dx)^3 \tan(e+fx) + 3ab^2(c+dx)^3 \tan^2(e+fx) + b^3(c+dx)^3 \tan^3(e+fx)) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3(c+dx)^4}{4d} - \frac{9a^2bd^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \\
 & \frac{9ia^2bd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3a^2b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} + \\
 & \frac{3ia^2b(c+dx)^4}{4d} - \frac{9ia^2bd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} - \frac{9iab^2d^2(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \\
 & \frac{9ab^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} + \frac{3ab^2(c+dx)^3 \tan(e+fx)}{f} - \frac{3iab^2(c+dx)^3}{f} - \\
 & \frac{3ab^2(c+dx)^4}{4d} + \frac{9ab^2d^3 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} + \frac{3b^3d^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \\
 & \frac{3b^3d^2(c+dx) \log(1+e^{2i(e+fx)})}{f^3} - \frac{3ib^3d(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \\
 & \frac{3b^3d(c+dx)^2 \tan(e+fx)}{2f^2} + \frac{b^3(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^3 \tan^2(e+fx)}{2f} + \\
 & \frac{3ib^3d(c+dx)^2}{2f^2} + \frac{b^3(c+dx)^3}{2f} - \frac{ib^3(c+dx)^4}{4d} + \frac{3ib^3d^3 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} + \\
 & \frac{3ib^3d^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
 \end{aligned}$$

input Int[(c + d*x)^3*(a + b*Tan[e + f*x])^3, x]

output

$$\begin{aligned} & \left(\frac{(3\pi)/2}{b^3 d (c + d x)^2} - \frac{(3\pi) a b^2 (c + d x)^3}{f^2} + \frac{(b^3 (c + d x)^3) / (2 f)}{d} + \frac{(a^3 (c + d x)^4) / (4 d)}{(4 d)} + \frac{((3\pi)/4) a^2 b^2 (c + d x)^4}{d} \right. \\ & - \frac{(3 a b^2 (c + d x)^4) / (4 d)}{(4 d)} - \frac{((\pi)/4) b^3 (c + d x)^4}{d} - \frac{(3 b^3 d^2 (c + d x) \operatorname{Log}[1 + E^{((2\pi)(e + f x))}]) / f^3}{d} + \frac{(9 a b^2 d (c + d x)^2 \operatorname{Log}[1 + E^{((2\pi)(e + f x))}]) / f^2}{d} \\ & - \frac{(3 a^2 b^2 (c + d x)^3 \operatorname{Log}[1 + E^{((2\pi)(e + f x))}]) / f}{d} + \frac{(b^3 (c + d x)^3 \operatorname{Log}[1 + E^{((2\pi)(e + f x))}]) / f}{d} + \frac{((3\pi)/2) b^3 d^3 \operatorname{PolyLog}[2, -E^{((2\pi)(e + f x))}]) / f^4}{d} - \frac{((9\pi) a b^2 d^2 (c + d x) \operatorname{PolyLog}[2, -E^{((2\pi)(e + f x))}]) / f^3}{d} \\ & + \frac{(((9\pi)/2) a^2 b^2 d (c + d x)^2 \operatorname{PolyLog}[2, -E^{((2\pi)(e + f x))}]) / f^2}{d} - \frac{(((3\pi)/2) b^3 d^3 (c + d x)^2 \operatorname{PolyLog}[2, -E^{((2\pi)(e + f x))}]) / f^2}{d} + \frac{(9 a b^2 d^3 (c + d x)^3 \operatorname{PolyLog}[3, -E^{((2\pi)(e + f x))}]) / (2 f^4)}{d} - \frac{(9 a^2 b^2 d^2 (c + d x) \operatorname{PolyLog}[3, -E^{((2\pi)(e + f x))}]) / (2 f^3)}{d} + \frac{(3 b^3 d^2 (c + d x) \operatorname{PolyLog}[3, -E^{((2\pi)(e + f x))}]) / (2 f^3)}{d} - \frac{((9\pi)/4) a^2 b^2 d^3 (c + d x)^3 \operatorname{PolyLog}[4, -E^{((2\pi)(e + f x))}]) / f^4}{d} + \frac{(((3\pi)/4) b^3 d^3 (c + d x)^2 \operatorname{Tan}[e + f x]) / (2 f^2)}{d} \\ & + \frac{(3 a b^2 (c + d x)^3 \operatorname{Tan}[e + f x]) / f}{d} + \frac{(b^3 (c + d x)^3 \operatorname{Tan}[e + f x]^2) / (2 f)}{d} \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4205 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)*((a_.) + (b_.)*\operatorname{Tan}[e_.] + (f_.)*(x_.))^{(n_.)}}] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1929 vs. $2(544) = 1088$.

Time = 1.00 (sec), antiderivative size = 1930, normalized size of antiderivative = 3.15

method	result	size
risch	Expression too large to display	1930

input `int((d*x+c)^3*(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & d^2*a^3*c*x^3 + 3/2*d*a^3*c^2*x^2 + a^3*c^3*x - 3/4*d^3*a*b^2*x^4 - 3*a*b^2*c^3*x^3 \\ & - 3/4/d*a*b^2*c^4 + I*b^3*c^3*x + 1/4*I/d*b^3*c^4 + 3/4*I*d^3*a^2*b*x^4 - 3/2*I*b^3*c^2*x^2 + 9/2*a*b^2*d^3*polylog(3, -exp(2*I*(f*x+e)))/f^4 - I*d^2*b^3*c*x^3 - 3 \\ & *d^2*a*b^2*c*x^3 - 9/2*d*a*b^2*c^2*x^2 + b^2*(-6*I*b*c*d^2*x + 18*I*a*c*d^2*f*x^2 + 2*b*d^3*f*x^3*exp(2*I*(f*x+e)) - 3*I*b*c^2*d*exp(2*I*(f*x+e)) - 6*I*b*c*d^2*x*exp(2*I*(f*x+e)) + 6*I*a*c^3*f*exp(2*I*(f*x+e)) + 6*b*c*d^2*f*x^2*exp(2*I*(f*x+e)) - 3*I*b*d^3*x^2 + 18*I*a*c^2*d*f*x*exp(2*I*(f*x+e)) - 3*I*b*c^2*d + 6*I*a*d^3*f*x^3*exp(2*I*(f*x+e)) + 6*b*c^2*d*f*x*exp(2*I*(f*x+e)) - 3*I*b*c^2*d + 6*I*a*d^3*f*x^3*exp(2*I*(f*x+e)) + 18*I*a*c^2*d*f*x + 2*b*c^3*f*exp(2*I*(f*x+e)) + 6*I*a*d^3*f*x^3*exp(2*I*(f*x+e)) + 18*I*a*c^2*f*x^2*exp(2*I*(f*x+e)) + 6*I*a*c^3*f/f^2/(1 + exp(2*I*(f*x+e)))^2 + 1/4*d^3*a^3*x^4 + 1/4/d*a^3*c^4 - 2/f*b^3*c^3*ln(exp(I*(f*x+e))) + 1/f*b^3*c^3*ln(1 + exp(2*I*(f*x+e))) - 1/4*I*b^3*d^3*x^4 - 36*I/f^2*b^2*d^2*c*a*e*x + 1/f*b^3*d^3*ln(1 + exp(2*I*(f*x+e)))*x^3 - 6/f^4*b^3*c^3*ln(exp(I*(f*x+e))) + 2/f^4*b^3*c^3*ln(exp(I*(f*x+e))) + 3/2/f^3*b^3*d^3*polylog(3, -exp(2*I*(f*x+e)))*x - 3/f^3*b^3*d^3*ln(1 + exp(2*I*(f*x+e)))*x + 3/2/f^3*b^3*c^3*d^2*polylog(3, -exp(2*I*(f*x+e))) + 6/f*b*a^2*c^3*ln(exp(I*(f*x+e))) - 3/f*b*a^2*c^3*ln(1 + exp(2*I*(f*x+e))) + 6/f^3*b^3*c^3*d^2*ln(exp(I*(f*x+e))) - 3/f^3*b^3*c^3*d^2*ln(1 + exp(2*I*(f*x+e))) - 3/2*I/f^4*b^3*d^3*c^3 - 4/3*I/f^2*b^2*d^3*c^3*x^2 + 3*I/f^4*b^3*d^3*c^3*x^2 + 18*I/f*b*d*c^2*a^2*e*x - 18*I/f^2*b*e^2*d^2*c*a^2*x + 9*I/f^2*b*a^2*c^3*d^2*polylog(2, -exp(2*I*(f*x+e)))*x - 3*I*a^2*b*c^3*x^3 - 4*I/d*a^2*b*c^4 + 3/f*b^3*d*c... \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1177 vs. $2(530) = 1060$.

Time = 0.11 (sec), antiderivative size = 1177, normalized size of antiderivative = 1.92

$$\int (c + dx)^3(a + b\tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

```

output 1/8*(2*(a^3 - 3*a*b^2)*d^3*f^4*x^4 + 3*I*(3*a^2*b - b^3)*d^3*polylog(4, (t
an(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*I*(3*a^2*b
- b^3)*d^3*polylog(4, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x +
e)^2 + 1)) + 4*(b^3*d^3*f^3 + 2*(a^3 - 3*a*b^2)*c*d^2*f^4)*x^3 + 12*(b^3*c
*d^2*f^3 + (a^3 - 3*a*b^2)*c^2*d*f^4)*x^2 + 4*(b^3*d^3*f^3*x^3 + 3*b^3*c*d
^2*f^3*x^2 + 3*b^3*c^2*d*f^3*x + b^3*c^3*f^3)*tan(f*x + e)^2 + 4*(3*b^3*c^
2*d*f^3 + 2*(a^3 - 3*a*b^2)*c^3*f^4)*x - 6*(I*(3*a^2*b - b^3)*d^3*f^2*x^2
- 6*I*a*b^2*c*d^2*f + I*b^3*d^3 + I*(3*a^2*b - b^3)*c^2*d*f^2 - 2*I*(3*a*b
^2*d^3*f - (3*a^2*b - b^3)*c*d^2*f^2)*x)*dilog(2*(I*tan(f*x + e) - 1)/(tan
(f*x + e)^2 + 1) + 1) - 6*(-I*(3*a^2*b - b^3)*d^3*f^2*x^2 + 6*I*a*b^2*c*d^
2*f - I*b^3*d^3 - I*(3*a^2*b - b^3)*c^2*d*f^2 + 2*I*(3*a*b^2*d^3*f - (3*a^
2*b - b^3)*c*d^2*f^2)*x)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1
) + 1) - 4*((3*a^2*b - b^3)*d^3*f^3*x^3 - 9*a*b^2*c^2*d*f^2 + 3*b^3*c*d^2*f
+ (3*a^2*b - b^3)*c^3*f^3 - 3*(3*a*b^2*d^3*f^2 - (3*a^2*b - b^3)*c*d^2*f
^3)*x^2 - 3*(6*a*b^2*c*d^2*f^2 - b^3*d^3*f - (3*a^2*b - b^3)*c^2*d*f^3)*x)
*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 4*((3*a^2*b - b^3)*d^
3*f^3*x^3 - 9*a*b^2*c^2*d*f^2 + 3*b^3*c*d^2*f + (3*a^2*b - b^3)*c^3*f^3 -
3*(3*a*b^2*d^3*f^2 - (3*a^2*b - b^3)*c*d^2*f^3)*x^2 - 3*(6*a*b^2*c*d^2*f^2
- b^3*d^3*f - (3*a^2*b - b^3)*c^2*d*f^3)*x)*log(-2*(-I*tan(f*x + e) - 1)/
(tan(f*x + e)^2 + 1)) + 6*(3*a*b^2*d^3 - (3*a^2*b - b^3)*d^3*f*x - (3*a...

```

Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 \, dx = \int (a + b \tan(e + fx))^3 (c + dx)^3 \, dx$$

input integrate((d*x+c)**3*(a+b*tan(f*x+e))**3,x)

output $\text{Integral}((a + b\tan(e + f*x))^{3.0}*(c + d*x)^{3.0}, x)$

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 6861 vs. $2(530) = 1060$.

Time = 8.17 (sec), antiderivative size = 6861, normalized size of antiderivative = 11.21

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output

```
1/4*(4*(f*x + e)*a^3*c^3 + (f*x + e)^4*a^3*d^3/f^3 - 4*(f*x + e)^3*a^3*d^3
*e/f^3 + 6*(f*x + e)^2*a^3*d^3*e^2/f^3 - 4*(f*x + e)*a^3*d^3*e^3/f^3 + 4*(f*x + e)^3*a^3*c*d^2/f^2 - 12*(f*x + e)^2*a^3*c*d^2*e/f^2 + 12*(f*x + e)*a^3*c*d^2*e^2/f^2 + 6*(f*x + e)^2*a^3*c^2*d/f - 12*(f*x + e)*a^3*c^2*d*e/f
+ 12*a^2*b*c^3*log(sec(f*x + e)) - 12*a^2*b*d^3*e^3*log(sec(f*x + e))/f^3
+ 36*a^2*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 36*a^2*b*c^2*d*e*log(sec(f*x + e))/f - 4*(72*a*b^2*d^3*e^3 - 72*a*b^2*c^3*f^3 - 3*(3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^4*d^3 + 36*b^3*d^3*e^2 + 12*((3*a^2*b + 3*I*a*b^2 - b^3)*d^3*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*f)*(f*x + e)^3 - 18*((3*a^2*b + 3*I*a*b^2 - b^3)*d^3*e^2 - 2*(3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*e*f + (3*a^2*b + 3*I*a*b^2 - b^3)*c^2*d*f^2)*(f*x + e)^2 + 36*(6*a*b^2*c^2*d*e + b^3*c^2*d)*f^2 + 12*((3*I*a*b^2 - b^3)*d^3*e^3 + 3*(-3*I*a*b^2 + b^3)*c*d^2*e^2*f + 3*(3*I*a*b^2 - b^3)*c^2*d*e*f^2 + (-3*I*a*b^2 + b^3)*c^3*f^3)*(f*x + e) - 72*(3*a*b^2*c*d^2*e^2 + b^3*c*d^2*f)*f + 4*(3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + 4*(3*a^2*b - b^3)*(f*x + e)^3*d^3 - 9*b^3*d^3*e - 9*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f)*(f*x + e)^2 + 9*(b^3*c^2*d*e - 3*a*b^2*c^2*d)*f^2 + 9*(6*a*b^2*d^3*e + b^3*d^3 + (3*a^2*b - b^3)*d^3*e^2 + (3*a^2*b - b^3)*c^2*d*f^2 - 2*(3*a*b^2*c*d^2 + (3*a^2*b - b^3)*c*d^2*e)*f)*(f*x + e) - 9*(b^3*c*d^2*e^2 - 6*a*b^2*c*d^2*e - b^3*c*d^2)*f + (3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + ...)
```

Giac [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \int (dx + c)^3 (b \tan(fx + e) + a)^3 dx$$

input `integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)^3*(b*tan(f*x + e) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^3 dx$$

input `int((a + b*tan(e + f*x))^3*(c + d*x)^3,x)`

output `int((a + b*tan(e + f*x))^3*(c + d*x)^3, x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)^3 (a + b \tan(e + fx))^3 dx \\ &= \frac{4(\int \tan(fx + e)^3 x^3 dx) b^3 d^3 f^2 + 12(\int \tan(fx + e)^3 x^2 dx) b^3 c d^2 f^2 + 12(\int \tan(fx + e)^3 x dx) b^3 c^2 d f^2}{\dots} \end{aligned}$$

input `int((d*x+c)^3*(a+b*tan(f*x+e))^3,x)`

output

```
(4*int(tan(e + f*x)**3*x**3,x)*b**3*d**3*f**2 + 12*int(tan(e + f*x)**3*x**2,x)*b**3*c**2*f**2 + 12*int(tan(e + f*x)**3*x,x)*b**3*c**2*d*f**2 + 12*int(tan(e + f*x)*x**3,x)*a**2*b*d**3*f**2 + 36*int(tan(e + f*x)*x**2,x)*a**2*b*c*d**2*f**2 - 36*int(tan(e + f*x)*x**2,x)*a*b**2*d**3*f + 36*int(tan(e + f*x)*x,x)*a**2*b*c**2*d*f**2 - 72*int(tan(e + f*x)*x,x)*a*b**2*c*d**2*f + 6*log(tan(e + f*x)**2 + 1)*a**2*b*c**3*f - 18*log(tan(e + f*x)**2 + 1)*a*b**2*c**2*d - 2*log(tan(e + f*x)**2 + 1)*b**3*c**3*f + 2*tan(e + f*x)**2*b**3*c**3*f + 12*tan(e + f*x)*a*b**2*c**3*f + 36*tan(e + f*x)*a*b**2*c**2*d*f*x + 36*tan(e + f*x)*a*b**2*c*d**2*f*x**2 + 12*tan(e + f*x)*a*b**2*d*f*x**3 + 4*a**3*c**3*f**2*x + 6*a**3*c**2*d*f**2*x**2 + 4*a**3*c*d**2*f**2*x**3 + a**3*d**3*f**2*x**4 - 12*a*b**2*c**3*f**2*x - 18*a*b**2*c**2*d*f**2*x**2 - 12*a*b**2*c*d**2*f**2*x**3 - 3*a*b**2*d**3*f**2*x**4)/(4*f**2)
```

3.50 $\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$

Optimal result	394
Mathematica [B] (warning: unable to verify)	395
Rubi [A] (verified)	396
Maple [B] (verified)	398
Fricas [A] (verification not implemented)	399
Sympy [F]	400
Maxima [B] (verification not implemented)	400
Giac [F]	401
Mupad [F(-1)]	402
Reduce [F]	402

Optimal result

Integrand size = 20, antiderivative size = 427

$$\begin{aligned}
 \int (c + dx)^2 (a + b \tan(e + fx))^3 dx = & -\frac{3iab^2(c + dx)^2}{f} + \frac{b^3(c + dx)^2}{2f} + \frac{a^3(c + dx)^3}{3d} \\
 & + \frac{ia^2b(c + dx)^3}{d} - \frac{ab^2(c + dx)^3}{d} - \frac{ib^3(c + dx)^3}{3d} \\
 & + \frac{6ab^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{b^3(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & - \frac{b^3d^2 \log(\cos(e + fx))}{f^3} \\
 & - \frac{3iab^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
 & + \frac{3ia^2bd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & - \frac{ib^3d(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & + \frac{b^3d^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{b^3d(c + dx) \tan(e + fx)}{f^2} \\
 & + \frac{3ab^2(c + dx)^2 \tan(e + fx)}{f} \\
 & + \frac{b^3(c + dx)^2 \tan^2(e + fx)}{2f}
 \end{aligned}$$

output

```
I*a^2*b*(d*x+c)^3/d+1/2*b^3*(d*x+c)^2/f+1/3*a^3*(d*x+c)^3/d-3*I*a*b^2*(d*x+c)^2/f-a*b^2*(d*x+c)^3/d-I*b^3*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2+6*a*b^2*d*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^2-3*a^2*b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f+b^3*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f-b^3*d^2*ln(cos(f*x+e))/f^3+3*I*a^2*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-1/3*I*b^3*(d*x+c)^3/d-3*I*a*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3-3/2*a^2*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3-b^3*d*(d*x+c)*tan(f*x+e)/f^2+3*a*b^2*(d*x+c)^2*tan(f*x+e)/f+1/2*b^3*(d*x+c)^2*tan(f*x+e)^2/f
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1846 vs. $2(427) = 854$.

Time = 7.05 (sec), antiderivative size = 1846, normalized size of antiderivative = 4.32

$$\int (c + dx)^2(a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input

```
Integrate[(c + d*x)^2*(a + b*Tan[e + f*x])^3,x]
```

```

output
((-1/4*I)*a^2*b*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e)))*Log[1 + E^
((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e +
f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e]
)/(E^(I*e)*f^3) + ((I/12)*b^3*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*
e)))*Log[1 + E^((2*I)*e)]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^
((-2*I)*(e + f*x))]) - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e +
f*x))])*Sec[e])/(E^(I*e)*f^3) - (b^3*d^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x]
] - Sin[e]*Sin[f*x] + f*x*Sin[e]))/(f^3*(Cos[e]^2 + Sin[e]^2)) + (6*a*b^2
*c*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x] + f*x*Sin[e]])/
(f^2*(Cos[e]^2 + Sin[e]^2)) - (3*a^2*b*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x]
] - Sin[e]*Sin[f*x] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^3*c^2
*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x] + f*x*Sin[e]])/(f*(
Cos[e]^2 + Sin[e]^2)) + (3*a*b^2*d^2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]]
) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)] -
2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]]))]) + Pi*L
og[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog
[2, E^((2*I)*(f*x - ArcTan[Cot[e]]))]))/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^3*S
qrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) - (3*a^2*b*c*d*Csc[e]*((f^2*x^2)/E^(
*I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^
((-2*I)*f*x)] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan...

```

Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 427, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$$

↓ 3042

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx$$

↓ 4205

$$\int (a^3(c+dx)^2 + 3a^2b(c+dx)^2 \tan(e+fx) + 3ab^2(c+dx)^2 \tan^2(e+fx) + b^3(c+dx)^2 \tan^3(e+fx)) dx$$

\downarrow 2009

$$\begin{aligned}
 & \frac{a^3(c+dx)^3}{3d} + \frac{3ia^2bd(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{3a^2b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} + \\
 & \frac{ia^2b(c+dx)^3}{d} - \frac{3a^2bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{6ab^2d(c+dx) \log(1+e^{2i(e+fx)})}{f^2} + \\
 & \frac{3ab^2(c+dx)^2 \tan(e+fx)}{f} - \frac{3iab^2(c+dx)^2}{f} - \frac{ab^2(c+dx)^3}{d} - \\
 & \frac{3iab^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} - \frac{ib^3d(c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \\
 & \frac{b^3d(c+dx) \tan(e+fx)}{f^2} + \frac{b^3(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^2 \tan^2(e+fx)}{2f} + \\
 & \frac{b^3(c+dx)^2}{2f} - \frac{ib^3(c+dx)^3}{3d} + \frac{b^3d^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{b^3d^2 \log(\cos(e+fx))}{f^3}
 \end{aligned}$$

input `Int[(c + d*x)^2*(a + b*Tan[e + f*x])^3, x]`

output

$$\begin{aligned}
 & ((-3*I)*a*b^2*(c + d*x)^2)/f + (b^3*(c + d*x)^2)/(2*f) + (a^3*(c + d*x)^3) \\
 & /(3*d) + (I*a^2*b*(c + d*x)^3)/d - (a*b^2*(c + d*x)^3)/d - ((I/3)*b^3*(c + \\
 & d*x)^3)/d + (6*a*b^2*d*(c + d*x)*\operatorname{Log}[1 + E^((2*I)*(e + f*x))])/f^2 - (3*a \\
 & ^2*b*(c + d*x)^2*\operatorname{Log}[1 + E^((2*I)*(e + f*x))])/f + (b^3*(c + d*x)^2*\operatorname{Log}[1 \\
 & + E^((2*I)*(e + f*x))])/f - (b^3*d^2*\operatorname{Log}[\operatorname{Cos}[e + f*x]])/f^3 - ((3*I)*a*b^2 \\
 & *d^2*\operatorname{PolyLog}[2, -E^((2*I)*(e + f*x))])/f^3 + ((3*I)*a^2*b*d*(c + d*x)*\operatorname{Poly} \\
 & \operatorname{Log}[2, -E^((2*I)*(e + f*x))])/f^2 - (I*b^3*d*(c + d*x)*\operatorname{PolyLog}[2, -E^((2*I) \\
 & *(e + f*x))])/f^2 - (3*a^2*b*d^2*\operatorname{PolyLog}[3, -E^((2*I)*(e + f*x))])/(2*f^3) \\
 & + (b^3*d^2*\operatorname{PolyLog}[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (b^3*d*(c + d*x)* \\
 & \operatorname{Tan}[e + f*x])/f^2 + (3*a*b^2*(c + d*x)^2*\operatorname{Tan}[e + f*x])/f + (b^3*(c + d*x)^ \\
 & 2*\operatorname{Tan}[e + f*x]^2)/(2*f)
 \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4205 $\text{Int}[(c_. + d_.*x_.)^(m_.)*(a_. + b_.*x_.)*\tan[e_. + f_.*x_.])^(n_.), \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], \ x] /; \ \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&& \ \text{IGtQ}[m, 0] \ \&& \ \text{IGtQ}[n, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1137 vs. $2(393) = 786$.

Time = 0.86 (sec) , antiderivative size = 1138, normalized size of antiderivative = 2.67

method	result	size
risch	Expression too large to display	1138

input $\text{int}((d*x+c)^2*(a+b*\tan(f*x+e))^3, x, \text{method}=\text{RETURNVERBOSE})$

output

```

12*I/f*b*a^2*c*d*e*x-3/2*a^2*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+2*b^2*
(3*I*a*d^2*f*x^2*exp(2*I*(f*x+e))+6*I*a*c*d*f*x*exp(2*I*(f*x+e))+b*d^2*f*x
^2*exp(2*I*(f*x+e))+3*I*a*c^2*f*exp(2*I*(f*x+e))+3*I*a*d^2*f*x^2-I*b*d^2*x
*x*exp(2*I*(f*x+e))+2*b*c*d*f*x*exp(2*I*(f*x+e))+6*I*a*c*d*f*x-I*b*c*d*exp(2
*I*(f*x+e))+b*c^2*f*exp(2*I*(f*x+e))+3*I*a*c^2*f-I*b*d^2*x-I*b*c*d)/f^2/(1
+exp(2*I*(f*x+e)))^2+d*a^3*c*x^2+a^3*c^2*x-d^2*a*b^2*x^3-3*a*b^2*c^2*x-1/d
*a*b^2*c^3+I*b^3*c^2*x+1/3*I/d*b^3*c^3+1/3*d^2*a^3*x^3+1/3/d*a^3*c^3+6/f*b
*a^2*c^2*ln(exp(I*(f*x+e)))-3/f*b*a^2*c^2*ln(1+exp(2*I*(f*x+e)))-2/f^3*b^3
*e^2*d^2*ln(exp(I*(f*x+e)))+1/f*b^3*d^2*ln(1+exp(2*I*(f*x+e)))*x^2+4/3*I/f
^3*b^3*d^2*e^3-2/f*b^3*c^2*ln(exp(I*(f*x+e)))+1/f*b^3*c^2*ln(1+exp(2*I*(f*
x+e)))+2/f^3*b^3*d^2*ln(exp(I*(f*x+e)))-1/f^3*b^3*d^2*ln(1+exp(2*I*(f*x+e))
)-1/3*I*b^3*d^2*x^3-I*d*b^3*c*x^2-3*d*a*b^2*c*x^2+I*d^2*a^2*b*x^3-3*I*a^2
*b*c^2*x-I/d*a^2*b*c^3-12/f^2*b*e*d*a^2*c*ln(exp(I*(f*x+e)))-6/f*b*a^2*c*d
*ln(1+exp(2*I*(f*x+e)))*x+1/2*b^3*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3-4*I
/f*b^3*c*d*e*x-12*I/f^2*b^2*a*d^2*e*x+3*I/f^2*b*a^2*c*d*polylog(2,-exp(2*I
*(f*x+e)))+3*I/f^2*b*a^2*d^2*polylog(2,-exp(2*I*(f*x+e)))*x+6*I/f^2*b*a^2*
c*d*e^2-6*I/f^2*b*a^2*d^2*e^2*x-3*I*a*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))
/f^3+3*I*d*a^2*b*c*x^2+6/f^2*b^2*a*d^2*ln(1+exp(2*I*(f*x+e)))*x-2*I/f^2*b^
3*c*d*e^2-6*I/f*b^2*a*d^2*x^2+2*I/f^2*b^3*d^2*ln(1+exp(2*I*(f*x+e)))*x-6*I/f^3*b^2*a*d^2*e^2-
4*I/f^3*b*a^2*d^2*e^3-I/f^2*b^3*c*d*polylog(2,-exp(2*I*(f*x+e)))-I/f^2*...

```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 688, normalized size of antiderivative = 1.61

$$\int (c + dx)^2(a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

output

```
1/12*(4*(a^3 - 3*a*b^2)*d^2*f^3*x^3 - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(b^3*d^2*f^2 + 2*(a^3 - 3*a*b^2)*c*d*f^3)*x^2 + 6*(b^3*d^2*f^2*x^2 + 2*b^3*c*d*f^2*x + b^3*c^2*f^2)*tan(f*x + e)^2 + 12*(b^3*c*d*f^2 + (a^3 - 3*a*b^2)*c^2*f^3)*x - 6*(-3*I*a*b^2*d^2 + I*(3*a^2*b - b^3)*d^2*f*x + I*(3*a^2*b - b^3)*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(3*I*a*b^2*d^2 - I*(3*a^2*b - b^3)*d^2*f*x - I*(3*a^2*b - b^3)*c*d*f)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*((3*a^2*b - b^3)*d^2*f^2*x^2 - 6*a*b^2*c*d*f + b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 - 2*(3*a*b^2*d^2*f - (3*a^2*b - b^3)*c*d*f^2)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*((3*a^2*b - b^3)*d^2*f^2*x^2 - 6*a*b^2*c*d*f + b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 - 2*(3*a*b^2*d^2*f - (3*a^2*b - b^3)*c*d*f^2)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 12*(3*a*b^2*d^2*f^2*x^2 + 3*a*b^2*c^2*f^2 - b^3*c*d*f + (6*a*b^2*c*d*f^2 - b^3*d^2*f)*x)*tan(f*x + e))/f^3
```

Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^2 dx$$

input

```
integrate((d*x+c)**2*(a+b*tan(f*x+e))**3,x)
```

output

```
Integral((a + b*tan(e + f*x))**3*(c + d*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3407 vs. $2(384) = 768$.

Time = 1.59 (sec), antiderivative size = 3407, normalized size of antiderivative = 7.98

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output

$$\begin{aligned} & \frac{1}{3} * (3*(f*x + e)*a^3*c^2 + (f*x + e)^3*a^3*d^2/f^2 - 3*(f*x + e)^2*a^3*d^2 \\ & *e/f^2 + 3*(f*x + e)*a^3*d^2*e^2/f^2 + 3*(f*x + e)^2*a^3*c*d/f - 6*(f*x + e)*a^3*c*d*e/f + 9*a^2*b*c^2*log(sec(f*x + e)) + 9*a^2*b*d^2*e^2*log(sec(f*x + e))/f^2 - 18*a^2*b*c*d*e*log(sec(f*x + e))/f + 3*(36*a*b^2*d^2*e^2 + 36*a*b^2*c^2*f^2 + 2*(3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^3*d^2 + 12*b^3*d^2*e - 6*((3*a^2*b + 3*I*a*b^2 - b^3)*d^2*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d*f)*(f*x + e)^2 - 6*((-3*I*a*b^2 + b^3)*d^2*e^2 + 2*(3*I*a*b^2 - b^3)*c*d*e*f + (-3*I*a*b^2 + b^3)*c^2*f^2)*(f*x + e) - 12*(6*a*b^2*c*d*e + b^3*c*d)*f + 6*(b^3*d^2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f + (b^3*d^2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*cos(4*f*x + 4*e) + 2*(b^3*d^2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*cos(2*f*x + 2*e) - (-I*b^3*d^2*e^2 - I*b^3*c^2*f^2 + 6*I*a*b^2*d^2*e + (3*I*a^2*b - I*b^3)*(f*x + e)^2*d^2 + I*b^3*d^2 + 2*(-3*I*a*b^2*d^2 + (-3*I*a^2*b + I*b^3)*d^2*e + (3*I*a^2*b - I*b^3)*c*d*f)*(f*x + e) + 2*(I*b^3*c*d*e - 3*I*a*b^2*c*d)*f)*sin(4*f*x + 4*e) - 2... \end{aligned}$$

Giac [F]

$$\int (c + dx)^2(a + b\tan(e + fx))^3 dx = \int (dx + c)^2(b\tan(fx + e) + a)^3 dx$$

input `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)^2*(b*tan(f*x + e) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^2 dx$$

input `int((a + b*tan(e + f*x))^3*(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))^3*(c + d*x)^2, x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)^2 (a + b \tan(e + fx))^3 dx \\ &= \frac{6(\int \tan(fx + e)^3 x^2 dx) b^3 d^2 f^2 + 12(\int \tan(fx + e)^3 x dx) b^3 c d f^2 + 18(\int \tan(fx + e) x^2 dx) a^2 b d^2 f^2 +}{\dots} \end{aligned}$$

input `int((d*x+c)^2*(a+b*tan(f*x+e))^3,x)`

output `(6*int(tan(e + f*x)**3*x**2,x)*b**3*d**2*f**2 + 12*int(tan(e + f*x)**3*x,x)*b**3*c*d*f**2 + 18*int(tan(e + f*x)*x**2,x)*a**2*b*d**2*f**2 + 36*int(tan(e + f*x)*x,x)*a**2*b*c*d*f**2 - 36*int(tan(e + f*x)*x,x)*a*b**2*d**2*f + 9*log(tan(e + f*x)**2 + 1)*a**2*b*c**2*f - 18*log(tan(e + f*x)**2 + 1)*a*b**2*c*d - 3*log(tan(e + f*x)**2 + 1)*b**3*c**2*f + 3*tan(e + f*x)**2*b**3*c**2*f + 18*tan(e + f*x)*a*b**2*c**2*f + 36*tan(e + f*x)*a*b**2*c*d*f*x + 18*tan(e + f*x)*a*b**2*d**2*f*x**2 + 6*a**3*c**2*f**2*x + 6*a**3*c*d*f**2*x**2 + 2*a**3*d**2*f**2*x**3 - 18*a*b**2*c**2*f**2*x - 18*a*b**2*c*d*f**2*x**2 - 6*a*b**2*d**2*f**2*x**3)/(6*f**2)`

3.51 $\int (c + dx)(a + b \tan(e + fx))^3 dx$

Optimal result	403
Mathematica [A] (verified)	404
Rubi [A] (verified)	404
Maple [B] (verified)	406
Fricas [A] (verification not implemented)	407
Sympy [F]	407
Maxima [B] (verification not implemented)	408
Giac [F]	409
Mupad [F(-1)]	409
Reduce [F]	409

Optimal result

Integrand size = 18, antiderivative size = 275

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^3 dx = & \frac{b^3 dx}{2f} + \frac{a^3(c + dx)^2}{2d} + \frac{3ia^2b(c + dx)^2}{2d} \\ & - \frac{3ab^2(c + dx)^2}{2d} - \frac{ib^3(c + dx)^2}{2d} \\ & - \frac{3a^2b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^3(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{3ab^2d \log(\cos(e + fx))}{f^2} \\ & + \frac{3ia^2bd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{ib^3d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{b^3d \tan(e + fx)}{2f^2} + \frac{3ab^2(c + dx) \tan(e + fx)}{f} \\ & + \frac{b^3(c + dx) \tan^2(e + fx)}{2f} \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{2} b^3 d x / f + \frac{1}{2} a^3 (d x + c)^2 / d + 3/2 I a^2 b (d x + c)^2 / d - 3/2 a b^2 (d x + c) \\ &)^2 / d - 1/2 I b^3 (d x + c)^2 / d - 3 a^2 b (d x + c) * \ln(1 + \exp(2 I (f x + e))) / f + b^3 (d x + c) * \ln(1 + \exp(2 I (f x + e))) / f + 3 a b^2 d \ln(\cos(f x + e)) / f^2 + 3/2 I a^2 b d * \text{polylog}(2, -\exp(2 I (f x + e))) / f^2 - 1/2 I b^3 d * \text{polylog}(2, -\exp(2 I (f x + e))) / f^2 - 1/2 b^3 d * \tan(f x + e) / f^2 + 3 a b^2 (d x + c) * \tan(f x + e) / f + 1/2 b^3 (d x + c) * \tan(f x + e)^2 / f \end{aligned}$$

Mathematica [A] (verified)

Time = 8.10 (sec), antiderivative size = 277, normalized size of antiderivative = 1.01

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx))^3 dx \\ & = \frac{\cos(e + fx) (\cos^2(e + fx) ((-((e + fx) (-3ia^2 bd(e + fx) + ib^3 d(e + fx) + 3ab^2(-de + 2cf + dfx) + a^3 d^2(e + fx)))) + 2*b*(-3*a^2 + b^2)*d*(e + fx)*\text{Log}[1 + E^{((2*I)*(e + fx))}] + 2*b*(3*a*b*d + 3*a^2*(d*e - c*f) + b^2*(-(d*e) + c*f))*\text{Log}[\cos(e + fx)]) - I*b*(-3*a^2 + b^2)*d*\cos(e + fx)^2*\text{PolyLog}[2, -E^{((2*I)*(e + fx))}] + (b^2*(2*b*f*(c + d*x) + (-b*d) + 6*a*f*(c + d*x))*\sin[2*(e + fx)]))/2)*(a + b*\tan(e + fx))^3}{(2*f^2*(a*\cos(e + fx) + b*\sin(e + fx))^3)} \end{aligned}$$

input

```
Integrate[(c + d*x)*(a + b*Tan[e + f*x])^3, x]
```

output

$$\begin{aligned} & (\cos(e + fx) * (\cos(e + fx)^2 * ((-((e + fx) * ((-3*I)*a^2*b*d*(e + fx) + I*b^3*d*(e + fx) + 3*a*b^2*(-(d*e) + 2*c*f + d*f*x) + a^3*(-2*c*f + d*(e - f*x)))) + 2*b*(-3*a^2 + b^2)*d*(e + fx)*\text{Log}[1 + E^{((2*I)*(e + fx))}] + 2*b*(3*a*b*d + 3*a^2*(d*e - c*f) + b^2*(-(d*e) + c*f))*\text{Log}[\cos(e + fx)]) - I*b*(-3*a^2 + b^2)*d*\cos(e + fx)^2*\text{PolyLog}[2, -E^{((2*I)*(e + fx))}] + (b^2*(2*b*f*(c + d*x) + (-b*d) + 6*a*f*(c + d*x))*\sin[2*(e + fx)]))/2)*(a + b*\tan(e + fx))^3) / (2*f^2*(a*\cos(e + fx) + b*\sin(e + fx))^3) \end{aligned}$$

Rubi [A] (verified)

Time = 0.59 (sec), antiderivative size = 275, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.167, Rules used = {3042, 4205, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (c + dx)(a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int (c + dx)(a + b \tan(e + fx))^3 dx \\
 & \quad \downarrow \textcolor{blue}{4205} \\
 & \int (a^3(c + dx) + 3a^2b(c + dx) \tan(e + fx) + 3ab^2(c + dx) \tan^2(e + fx) + b^3(c + dx) \tan^3(e + fx)) dx \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{a^3(c + dx)^2}{2d} - \frac{3a^2b(c + dx) \log(1 + e^{2i(e+fx)})}{f} + \frac{3ia^2b(c + dx)^2}{2d} + \\
 & \frac{3ia^2bd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} + \frac{3ab^2(c + dx) \tan(e + fx)}{f} - \frac{3ab^2(c + dx)^2}{2d} + \\
 & \frac{3ab^2d \log(\cos(e + fx))}{f^2} + \frac{b^3(c + dx) \log(1 + e^{2i(e+fx)})}{f} + \frac{b^3(c + dx) \tan^2(e + fx)}{2f} - \\
 & \frac{ib^3(c + dx)^2}{2d} - \frac{ib^3d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b^3d \tan(e + fx)}{2f^2} + \frac{b^3dx}{2f}
 \end{aligned}$$

input `Int[(c + d*x)*(a + b*Tan[e + f*x])^3,x]`

output
$$\begin{aligned}
 & (b^{3*d*x}/(2*f) + (a^3*(c + d*x)^2)/(2*d) + ((3*I)/2)*a^2*b*(c + d*x)^2)/ \\
 & d - (3*a*b^2*(c + d*x)^2)/(2*d) - ((I/2)*b^3*(c + d*x)^2)/d - (3*a^2*b*(c \\
 & + d*x)*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f + (b^3*(c + d*x)*\operatorname{Log}[1 + E^{((2*I)*(\\
 & e + f*x))}])/f + (3*a*b^2*d*\operatorname{Log}[\operatorname{Cos}[e + f*x]])/f^2 + (((3*I)/2)*a^2*b*d*\operatorname{Pol} \\
 & y\operatorname{Log}[2, -E^{((2*I)*(e + f*x))}])/f^2 - ((I/2)*b^3*d*\operatorname{PolyLog}[2, -E^{((2*I)*(e \\
 & + f*x))}])/f^2 - (b^3*d*\operatorname{Tan}[e + f*x])/(2*f^2) + (3*a*b^2*(c + d*x)*\operatorname{Tan}[e + \\
 & f*x])/f + (b^3*(c + d*x)*\operatorname{Tan}[e + f*x]^2)/(2*f)
 \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4205 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (a + b*\text{Tan}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 492 vs. $2(245) = 490$.

Time = 0.77 (sec), antiderivative size = 493, normalized size of antiderivative = 1.79

method	result
risch	$-\frac{3ab^2dx^2}{2} - 3ab^2cx - 3ia^2bcx + \frac{3ia^2bdx^2}{2} + \frac{b^3d\ln(1+e^{2i(fx+e)})x}{f} - \frac{6b^2ad\ln(e^{i(fx+e)})}{f^2} + \frac{3b^2ad\ln(1+e^{2i(fx+e)})}{f^2}$

input $\text{int}((d*x+c)*(a+b*\tan(f*x+e))^3, x, \text{method}=\text{RETURNVERBOSE})$

output
$$\begin{aligned} & -3/2*a*b^2*d*x^2 - 3*a*b^2*c*x + I*b^3*c*x + 1/2*a^3*d*x^2 + a^3*c*x + b^2*(6*I*a*d*f*x*\exp(2*I*(f*x+e)) + 6*I*a*c*f*\exp(2*I*(f*x+e)) + 2*b*d*f*x*\exp(2*I*(f*x+e)) + 6*I*a*d*f*x - I*b*d)/f^2/(1+\exp(2*I*(f*x+e)))^2 - 2/f*b^3*c*\ln(\exp(I*(f*x+e))) + 1/f*b^3*c*\ln(1 + \exp(2*I*(f*x+e))) - 1/2*I*b^3*d*x^2 - 3*I*a^2*b*c*x + 3/2*I*a^2*b*d*x^2 + 1/f*b^3*d*\ln(1 + \exp(2*I*(f*x+e)))*x - 6/f^2*b^2*a*d*\ln(\exp(I*(f*x+e))) + 3/f^2*b^2*a*d*\ln(1 + \exp(2*I*(f*x+e))) + 6/f*b*a^2*c*\ln(\exp(I*(f*x+e))) - 3/f*b*a^2*c*\ln(1 + \exp(2*I*(f*x+e))) + 2/f^2*b^3*e*d*\ln(\exp(I*(f*x+e))) - I/f^2*b^3*d*e^2 - 1/2*I*b^3*d*\text{polylog}(2, -\exp(2*I*(f*x+e)))/f^2 + 6*I/f*b*a^2*d*e*x - 6/f^2*b*e*a^2*d*\ln(\exp(I*(f*x+e))) - 3/f*b*a^2*d*\ln(1 + \exp(2*I*(f*x+e)))*x + 3*I/f^2*b*a^2*d*e^2 - 2*I/f*b^3*d*e*x + 3/2*I*a^2*b*d*\text{polylog}(2, -\exp(2*I*(f*x+e)))/f^2 \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.19

$$\int (c + dx)(a + b \tan(e + fx))^3 dx$$

$$= \frac{2(a^3 - 3ab^2)df^2x^2 - i(3a^2b - b^3)d\text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + i(3a^2b - b^3)d\text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right)}{}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

output $\frac{1}{4}*(2*(a^3 - 3*a*b^2)*d*f^2*x^2 - I*(3*a^2*b - b^3)*d*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + I*(3*a^2*b - b^3)*d*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + 2*(b^3*d*f*x + b^3*c*f)*tan(f*x + e)^2 + 2*(b^3*d*f + 2*(a^3 - 3*a*b^2)*c*f^2)*x + 2*(3*a*b^2*d - (3*a^2*b - b^3)*d*f*x - (3*a^2*b - b^3)*c*f)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(3*a*b^2*d - (3*a^2*b - b^3)*d*f*x - (3*a^2*b - b^3)*c*f)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(6*a*b^2*d*f*x + 6*a*b^2*c*f - b^3*d)*tan(f*x + e))/f^2$

Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx) dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))**3,x)`

output `Integral((a + b*tan(e + f*x))**3*(c + d*x), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1319 vs. $2(239) = 478$.

Time = 0.43 (sec) , antiderivative size = 1319, normalized size of antiderivative = 4.80

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="maxima")`

output

```
1/2*(2*(f*x + e)*a^3*c + (f*x + e)^2*a^3*d/f - 2*(f*x + e)*a^3*d*e/f + 6*a^2*b*c*log(sec(f*x + e)) - 6*a^2*b*d*e*log(sec(f*x + e))/f - 2*(12*a*b^2*d*e - 12*a*b^2*c*f - (3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d + 2*b^3*d + 2*((3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e) + 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d + (b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*cos(4*f*x + 4*e) + 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*cos(2*f*x + 2*e) + (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*sin(4*f*x + 4*e) + 2*(I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - ((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d - 2*(6*a*b^2*d + (3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e))*cos(4*f*x + 4*e) - 2*((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d - b^3*d - 2*(3*a*b^2 - I*b^3)*d*e + 2*(3*a*b^2 - I*b^3)*c*f - 2*((3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e))*cos(2*f*x + 2*e) - ((3*a^2*b - b^3)*d*cos(4*f*x + 4*e) + 2*(3*a^2*b - b^3)*d*cos(2*f*x + 2*e) + (3*I*a^2*b - I*b^3)*d*sin(4*f*x + 4*e) - 2*(-3*I*a^2*b + I*b^3)*d*sin(2*f*x + 2*e) + (3*a^2*b - b^3)*d)*dilog(-e^(2*I*f*x + 2*I*e)) - (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d + (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*cos(4*f*x + 4*e) - 2*(-I*b...
```

Giac [F]

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (dx + c)(b \tan(fx + e) + a)^3 dx$$

input `integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((d*x + c)*(b*tan(f*x + e) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx) dx$$

input `int((a + b*tan(e + f*x))^3*(c + d*x),x)`

output `int((a + b*tan(e + f*x))^3*(c + d*x), x)`

Reduce [F]

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx))^3 dx \\ &= \frac{2 \left(\int \tan(fx + e)^3 dx \right) b^3 d f^2 + 6 \left(\int \tan(fx + e) dx \right) a^2 b d f^2 + 3 \log(\tan(fx + e)^2 + 1) a^2 b c f - 3 \log \right. \end{aligned}$$

input `int((d*x+c)*(a+b*tan(f*x+e))^3,x)`

output `(2*int(tan(e + f*x)**3*x,x)*b**3*d*f**2 + 6*int(tan(e + f*x)*x,x)*a**2*b*d*f**2 + 3*log(tan(e + f*x)**2 + 1)*a**2*b*c*f - 3*log(tan(e + f*x)**2 + 1)*a*b**2*d - log(tan(e + f*x)**2 + 1)*b**3*c*f + tan(e + f*x)**2*b**3*c*f + 6*tan(e + f*x)*a*b**2*c*f + 6*tan(e + f*x)*a*b**2*d*f*x + 2*a**3*c*f**2*x + a**3*d*f**2*x**2 - 6*a*b**2*c*f**2*x - 3*a*b**2*d*f**2*x**2)/(2*f**2)`

3.52 $\int \frac{(a+b \tan(e+fx))^3}{c+dx} dx$

Optimal result	410
Mathematica [N/A]	410
Rubi [N/A]	411
Maple [N/A]	412
Fricas [N/A]	412
Sympy [N/A]	412
Maxima [N/A]	413
Giac [N/A]	414
Mupad [N/A]	414
Reduce [N/A]	414

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^3}{c + dx}, x\right)$$

output `Defer(Int)((a+b*tan(f*x+e))^3/(d*x+c),x)`

Mathematica [N/A]

Not integrable

Time = 13.63 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

input `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

output `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

Rubi [N/A]

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx \end{aligned}$$

input `Int[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^3}{dx + c} dx$$

input `int((a+b*tan(f*x+e))^3/(d*x+c),x)`

output `int((a+b*tan(f*x+e))^3/(d*x+c),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^3}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c),x, algorithm="fricas")`

output `integral((b^3*tan(f*x + e))^3 + 3*a*b^2*tan(f*x + e)^2 + 3*a^2*b*tan(f*x + e) + a^3)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 1.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

input `integrate((a+b*tan(f*x+e))**3/(d*x+c),x)`

output $\text{Integral}((a + b\tan(e + f*x))^{**3}/(c + d*x), x)$

Maxima [N/A]

Not integrable

Time = 2.80 (sec) , antiderivative size = 1935, normalized size of antiderivative = 96.75

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^3}{dx + c} dx$$

```
input integrate((a+b*tan(f*x+e))^3/(d*x+c),x, algorithm="maxima")
```

```

output (((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*cos(4*f*x + 4*e)^2*log(d*x + c) + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c)*sin(4*f*x + 4*e)^2 + 4*(b^3*d^2*f*x + b^3*c*d*f + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c))*cos(2*f*x + 2*e)^2 + 4*(b^3*d^2*f*x + b^3*c*d*f + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c))*sin(2*f*x + 2*e)^2 + (2*(b^3*d^2*f*x + b^3*c*d*f + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c))*cos(2*f*x + 2*e) + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c) - (6*a*b^2*d^2*f^2*x + 6*a*b^2*c*d*f + b^3*d^2)*sin(2*f*x + 2*e))*cos(4*f*x + 4*e) + 2*(b^3*d^2*f*x + b^3*c*d*f + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c))*cos(2*f*x + 2*e) - (d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2 + (d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*cos(2*f*x + 2*e)^2 + (d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*sin(4*f*x + 4*e)^2 + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*sin(2*f*x + 2*e)^2 + 2*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2) + 2*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)...

```

Giac [N/A]

Not integrable

Time = 0.43 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^3}{dx + c} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c),x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)^3/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 9.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

input `int((a + b*tan(e + f*x))^3/(c + d*x),x)`

output `int((a + b*tan(e + f*x))^3/(c + d*x), x)`

Reduce [N/A]

Not integrable

Time = 0.15 (sec) , antiderivative size = 86, normalized size of antiderivative = 4.30

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx \\ &= \frac{\left(\int \frac{\tan(fx+e)^3}{dx+c} dx \right) b^3 d + 3 \left(\int \frac{\tan(fx+e)^2}{dx+c} dx \right) a b^2 d + 3 \left(\int \frac{\tan(fx+e)}{dx+c} dx \right) a^2 b d + \log(dx + c) a^3}{d} \end{aligned}$$

input `int((a+b*tan(f*x+e))^3/(d*x+c),x)`

output `(int(tan(e + f*x)**3/(c + d*x),x)*b**3*d + 3*int(tan(e + f*x)**2/(c + d*x),x)*a*b**2*d + 3*int(tan(e + f*x)/(c + d*x),x)*a**2*b*d + log(c + d*x)*a**3)/d`

3.53 $\int \frac{(a+b\tan(e+fx))^3}{(c+dx)^2} dx$

Optimal result	416
Mathematica [N/A]	416
Rubi [N/A]	417
Maple [N/A]	418
Fricas [N/A]	418
Sympy [N/A]	418
Maxima [N/A]	419
Giac [N/A]	420
Mupad [N/A]	420
Reduce [N/A]	420

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^3}{(c + dx)^2}, x\right)$$

output `Defer(Int)((a+b*tan(f*x+e))^3/(d*x+c)^2,x)`

Mathematica [N/A]

Not integrable

Time = 16.87 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

input `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x)^2,x]`

output `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x)^2, x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx \end{aligned}$$

input `Int[(a + b*Tan[e + f*x])^3/(c + d*x)^2, x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^3}{(dx + c)^2} dx$$

input `int((a+b*tan(f*x+e))^3/(d*x+c)^2,x)`

output `int((a+b*tan(f*x+e))^3/(d*x+c)^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 3.15

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="fricas")`

output `integral((b^3*tan(f*x + e)^3 + 3*a*b^2*tan(f*x + e)^2 + 3*a^2*b*tan(f*x + e) + a^3)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 2.38 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

input `integrate((a+b*tan(f*x+e))**3/(d*x+c)**2,x)`

output $\text{Integral}((a + b\tan(e + f*x))^{3/2}/(c + d*x)^2, x)$

Maxima [N/A]

Not integrable

Time = 9.60 (sec) , antiderivative size = 2194, normalized size of antiderivative = 109.70

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

```
input integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="maxima")
```

```

output -((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2 + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*cos(4*f*x + 4*e)^2 + 4*((a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + (a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 2*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e)^2 + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*sin(4*f*x + 4*e)^2 + 4*((a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + (a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 2*(a^3 - 3*a*b^2)*c*d*f^2)*x)*sin(2*f*x + 2*e)^2 + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2 + (2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e) + (3*a*b^2*d^2*f^2*x + 3*a*b^2*c*d*f + b^3*d^2)*sin(2*f*x + 2*e))*cos(4*f*x + 4*e) + 2*(2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e) + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e)^2 + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x...)
```

Giac [N/A]

Not integrable

Time = 25.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

input `integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="giac")`

output `integrate((b*tan(f*x + e) + a)^3/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 9.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

input `int((a + b*tan(e + f*x))^3/(c + d*x)^2,x)`

output `int((a + b*tan(e + f*x))^3/(c + d*x)^2, x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 237, normalized size of antiderivative = 11.85

$$\begin{aligned} & \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx \\ &= \frac{\left(\int \frac{\tan(fx+e)^3}{d^2x^2+2cdx+c^2} dx \right) b^3 c^2 + \left(\int \frac{\tan(fx+e)^3}{d^2x^2+2cdx+c^2} dx \right) b^3 c dx + 3 \left(\int \frac{\tan(fx+e)^2}{d^2x^2+2cdx+c^2} dx \right) a b^2 c^2 + 3 \left(\int \frac{\tan(fx+e)^2}{d^2x^2+2cdx+c^2} dx \right)}{c(dx + c)} \end{aligned}$$

input int((a+b*tan(f*x+e))^3/(d*x+c)^2,x)

output
$$\begin{aligned} & \left(\int (\tan(e + f*x)^{**3} / (c^{**2} + 2*c*d*x + d^{**2}*x^{**2}), x) * b^{**3}*c^{**2} + \int (\tan(e + f*x)^{**3} / (c^{**2} + 2*c*d*x + d^{**2}*x^{**2}), x) * b^{**3}*c*d*x + 3*\int (\tan(e + f*x)^{**2} / (c^{**2} + 2*c*d*x + d^{**2}*x^{**2}), x) * a*b^{**2}*c^{**2} + 3*\int (\tan(e + f*x)^{**2} / (c^{**2} + 2*c*d*x + d^{**2}*x^{**2}), x) * a*b^{**2}*c*d*x + 3*\int (\tan(e + f*x) / (c^{**2} + 2*c*d*x + d^{**2}*x^{**2}), x) * a**2*b*c^{**2} + 3*\int (\tan(e + f*x) / (c^{**2} + 2*c*d*x + d^{**2}*x^{**2}), x) * a**2*b*c*d*x + a**3*x \right) / (c*(c + d*x)) \end{aligned}$$

3.54 $\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$

Optimal result	422
Mathematica [A] (warning: unable to verify)	423
Rubi [A] (verified)	423
Maple [B] (verified)	427
Fricas [B] (verification not implemented)	428
Sympy [F]	429
Maxima [B] (verification not implemented)	430
Giac [F]	431
Mupad [F(-1)]	431
Reduce [F]	431

Optimal result

Integrand size = 20, antiderivative size = 243

$$\begin{aligned} \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx = & \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\ & - \frac{3ibd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2} \\ & + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^3} \\ & + \frac{3ibd^3 \operatorname{PolyLog}\left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{4(a^2+b^2)f^4} \end{aligned}$$

output

```
1/4*(d*x+c)^4/(a+I*b)/d+b*(d*x+c)^3*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-3/2*I*b*d*(d*x+c)^2*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2+3/2*b*d^2*(d*x+c)*polylog(3,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^3+3/4*I*b*d^3*polylog(4,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^4
```

Mathematica [A] (warning: unable to verify)

Time = 1.45 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.22

$$\begin{aligned}
 & \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx \\
 &= \frac{1}{4}b \left(-\frac{2(c+dx)^4}{(ia+b)d(-ib(-1+e^{2ie})+a(1+e^{2ie}))} + \frac{4(c+dx)^3 \log\left(1+\frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2+b^2)f} \right. \\
 & \quad \left. + \frac{3d\left(2if^2(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right) + d\left(2f(c+dx) \operatorname{PolyLog}\left(3, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right) - id\operatorname{PolyLog}\left(4, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)\right)\right)}{(a^2+b^2)f^4} \right. \\
 & \quad \left. + \frac{x(4c^3+6c^2dx+4cd^2x^2+d^3x^3)\cos(e)}{4(a\cos(e)+b\sin(e))} \right)
 \end{aligned}$$

input `Integrate[(c + d*x)^3/(a + b*Tan[e + f*x]), x]`

output

$$\begin{aligned}
 & (b*(-2*(c + d*x)^4)/((I*a + b)*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))) + (4*(c + d*x)^3*\operatorname{Log}[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/(a^2 + b^2)*f) + (3*d*((2*I)*f^2*(c + d*x)^2*\operatorname{PolyLog}[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] + d*(2*f*(c + d*x)*\operatorname{PolyLog}[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - I*d*\operatorname{PolyLog}[4, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^4))/4 + (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*\operatorname{Cos}[e])/((4*(a*\operatorname{Cos}[e] + b*\operatorname{Sin}[e]))
 \end{aligned}$$
Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {3042, 4215, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx \downarrow 3042$$

$$\begin{aligned}
& \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx \\
& \quad \downarrow \textcolor{blue}{4215} \\
& 2ib \int \frac{e^{2i(e+fx)}(c+dx)^3}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx + \frac{(c+dx)^4}{4d(a+ib)} \\
& \quad \downarrow \textcolor{blue}{2620} \\
& 2ib \left(\frac{3id \int (c+dx)^2 \log \left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) dx}{2f(a^2+b^2)} - \frac{i(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \\
& \quad \frac{(c+dx)^4}{4d(a+ib)} \\
& \quad \downarrow \textcolor{blue}{3011} \\
& 2ib \left(\frac{3id \left(\frac{i(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \int (c+dx) \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{f} \right)}{2f(a^2+b^2)} - \frac{i(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right. \\
& \quad \left. \frac{(c+dx)^4}{4d(a+ib)} \right. \\
& \quad \downarrow \textcolor{blue}{7163} \\
& 2ib \left(\frac{3id \left(\frac{i(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \int \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{2f} - \frac{i(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f}}{2f(a^2+b^2)} \right. \\
& \quad \left. \frac{(c+dx)^4}{4d(a+ib)} \right)
\end{aligned}$$

$$\begin{aligned}
 & \downarrow \textcolor{blue}{2720} \\
 2ib & \left(\frac{3id \left(\frac{i(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \left(\frac{d \int e^{-2i(e+fx)} \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) de^{2i(e+fx)}}{4f^2} - \frac{i(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{f} \right)}{2f(a^2+b^2)} \right) \right. \\
 & \quad \left. \frac{(c+dx)^4}{4d(a+ib)} \right) \\
 & \quad \downarrow \textcolor{blue}{7143} \\
 2ib & \left(\frac{3id \left(\frac{i(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \left(\frac{d \operatorname{PolyLog} \left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{4f^2} - \frac{i(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} \right)}{f} \right)}{2f(a^2+b^2)} \right. \\
 & \quad \left. \frac{(c+dx)^4}{4d(a+ib)} \right)
 \end{aligned}$$

input

output

$$(c + d*x)^4/(4*(a + I*b)*d) + (2*I)*b*(((-1/2*I)*(c + d*x)^3*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) + (((3*I)/2)*d*((I/2)*(c + d*x)^2*PolyLog[2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/f - (I*d*(((-1/2*I)*(c + d*x)*PolyLog[3, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/f + (d*PolyLog[4, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/(4*f^2))/f))/((a^2 + b^2)*f)))$$

Definitions of rubi rules used

rule 2620

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*F_[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(n_.)]*((f_.) + (g_.)*(x_.))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4215 $\text{Int}[(c_+ + d_+)(x_+)^m / ((a_+ + b_+) \tan(e_+ + f_+)(x_+)), x] \rightarrow \text{Simp}[(c + d*x)^{m+1} / (d*(m+1)*(a + I*b)), x] + \text{Simp}[2*I*b \text{ Int}[(c + d*x)^m * (\text{E}^{\text{Simp}[2*I*(e + f*x), x]} / ((a + I*b)^2 + (a^2 + b^2)*\text{E}^{\text{Simp}[2*I*(e + f*x), x]})), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \& \text{NeQ}[a^2 + b^2, 0] \& \text{IGtQ}[m, 0]$

rule 7143 $\text{Int}[\text{PolyLog}[n_, (c_+)(a_+ + b_+)(x_+)^p] / ((d_+ + e_+)(x_+)), x] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c*(a + b*x)^p] / (e*p), x] /; \text{FreeQ}[\{a, b, c, d, e, n, p\}, x] \& \text{EqQ}[b*d, a*e]$

rule 7163 $\text{Int}[(e_+ + f_+)(x_+)^m * \text{PolyLog}[n_, (d_+)^{(c_+)(a_+ + b_+)(x_+)}]^p] /; \text{Simp}[(e + f*x)^m * (\text{PolyLog}[n + 1, d*(F^(c*(a + b*x))^p)] / (b*c*p*\text{Log}[F])), x] - \text{Simp}[f*(m/(b*c*p*\text{Log}[F])) \text{ Int}[(e + f*x)^{m-1} * \text{PolyLog}[n + 1, d*(F^(c*(a + b*x)))^p]], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, n, p\}, x] \& \text{GtQ}[m, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1467 vs. $2(220) = 440$.

Time = 0.61 (sec) , antiderivative size = 1468, normalized size of antiderivative = 6.04

method	result	size
risch	Expression too large to display	1468

input `int((d*x+c)^3/(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -3/f^2/(I*a+b)*b/(-I*b-a)*d^2*c*polylog(2, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) \\
 & *x+6/f^2/(I*a+b)*b/(-I*b-a)*e^2*c*d^2*x-6/f/(I*a+b)*b/(-I*b-a)*c^2*d*e*x \\
 & -3/2*I/f^3/(I*a+b)*b/(-I*b-a)*d^3*polylog(3, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) \\
 & *x-I/f^4/(I*a+b)*b*e^3*d^3/(a+I*b)*ln(I*b*exp(2*I*(f*x+e))-a*exp(2*I*(f*x+e))) \\
 & -I*b-a)+2*I/f^4/(I*a+b)*b*e^3*d^3/(a+I*b)*ln(exp(I*(f*x+e)))-I/f/(I*a+b)*b/ \\
 & (-I*b-a)*d^3*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x^3-I/f^4/(I*a+b)*b/ \\
 & (-I*b-a)*e^3*d^3*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-3/2*I/f^3/(I*a+b)*b/ \\
 & (-I*b-a)*d^2*c*polylog(3, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-d^2/(I*b-a)*c*x^3-3/2*d/(I*b-a)*c^2*x^2-1/4*d^3/(I*b-a)*x^4-1/(I*b-a)*c^3*x-1/ \\
 & 4/d/(I*b-a)*c^4-1/2/(I*a+b)*b/(-I*b-a)*d^3*x^4-3/2/f^2/(I*a+b)*b/(-I*b-a)* \\
 & c^2*d*polylog(2, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))+4/f^3/(I*a+b)*b/(-I*b-a)* \\
 & e^3*c*d^2+I/f/(I*a+b)*b*c^3/(a+I*b)*ln(I*b*exp(2*I*(f*x+e))-a*exp(2*I*(f*x+e))) \\
 & -I*b-a)-3/2/f^2/(I*a+b)*b/(-I*b-a)*d^3*polylog(2, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) \\
 & *x^2-2/f^3/(I*a+b)*b/(-I*b-a)*d^3*e^3*x-3/f^2/(I*a+b)*b/(-I*b-a)*c^2*d*e^2-2*I/f/(I*a+b)*b*c^3/(a+I*b)*ln(exp(I*(f*x+e)))-2/(I*a+b)*b/ \\
 & (-I*b-a)*d^2*c*x^3-3/2/f^4/(I*a+b)*b/(-I*b-a)*d^3*e^4+3/4/f^4/(I*a+b)*b/(-I*b-a)*d^3*polylog(4, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-3/(I*a+b)*b/(-I*b-a)*c^2*d*x^2-3*I/f/(I*a+b)*b/(-I*b-a)*d^2*c*ln(1-(a-I*b)*exp(2*I*(f*x+e))) \\
 & /(-I*b-a)*x^2+3*I/f^3/(I*a+b)*b*e^2*c*d^2/(a+I*b)*ln(I*b*exp(2*I*(f*x+e))-a*exp(2*I*(f*x+e))-I*b-a)-6*I/f^3/(I*a+b)*b*e^2*c*d^2/(a+I*b)*ln(exp(I...
 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1183 vs. $2(212) = 424$.

Time = 0.11 (sec), antiderivative size = 1183, normalized size of antiderivative = 4.87

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output

```
1/8*(2*a*d^3*f^4*x^4 + 8*a*c*d^2*f^4*x^3 + 12*a*c^2*d*f^4*x^2 + 8*a*c^3*f^4*x - 3*I*b*d^3*polylog(4, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e))^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 3*I*b*d^3*polylog(4, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 6*(-I*b*d^3*f^2*x^2 - 2*I*b*c*d^2*f^2*x - I*b*c^2*d*f^2)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*b*d^3*f^2*x^2 + 2*I*b*c*d^2*f^2*x + I*b*c^2*d*f^2)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*f^2 - b*c^3*f^3)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - 4*(b*d^3*e^3 - 3*b*c*...
```

Sympy [F]

$$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx = \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$$

input

```
integrate((d*x+c)**3/(a+b*tan(f*x+e)),x)
```

output

```
Integral((c + d*x)**3/(a + b*tan(e + f*x)), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 983 vs. $2(212) = 424$.

Time = 0.48 (sec) , antiderivative size = 983, normalized size of antiderivative = 4.05

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output

```
-1/12*(18*c^2*d*e*(2*(f*x + e)*a/((a^2 + b^2)*f) + 2*b*log(b*tan(f*x + e) + a)/((a^2 + b^2)*f) - b*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*f)) - 6*(2*(f*x + e)*a/(a^2 + b^2) + 2*b*log(b*tan(f*x + e) + a)/(a^2 + b^2) - b*log(tan(f*x + e)^2 + 1)/(a^2 + b^2))*c^3 - (3*(f*x + e)^4*(a - I*b)*d^3 + 12*I*b*d^3*polylog(4, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 12*((a - I*b)*d^3*3*e - (a - I*b)*c*d^2*f)*(f*x + e)^3 + 18*((a - I*b)*d^3*e^2 - 2*(a - I*b)*c*d^2*f^2)*(f*x + e)^2 - 12*((a - I*b)*d^3*e^3 - 3*(a - I*b)*c*d^2*e^2*f)*(f*x + e) - 12*(I*b*d^3*e^3 - 3*I*b*c*d^2*e^2*f)*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) - 4*(4*I*(f*x + e)^3*b*d^3 + 9*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e)^2 + 9*(I*b*d^3*e^2 - 2*I*b*c*d^2*f + I*b*c^2*d*f^2)*(f*x + e))*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) - 6*(4*I*(f*x + e)^2*b*d^3 + 3*I*b*d^3*e^2 - 6*I*b*c*d^2*f + 3*I*b*c^2*d*f^2 + 6*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e))*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 6*(b*d^3*e^3 - 3*b*c*d^2*e^2*f)*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 2*(4*(f*x + e)^3*b*d^3 - 9*(b*d^3*e - b*c*d^2*f)*(f*x + e)^2 + 9*(b*d^3*e^2 - 2*b*c*d^2*f + b*c^2*d*f^2)*(f*x + e))*log(((a^2 + b^2)*cos(2*f*x + 2*e)...)
```

Giac [F]

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(dx + c)^3}{b \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^3/(b*tan(f*x + e) + a), x)`

Mupad [-1]

Timed out.

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx$$

input `int((c + d*x)^3/(a + b*tan(e + f*x)),x)`

output `int((c + d*x)^3/(a + b*tan(e + f*x)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx \\ &= \frac{2 \left(\int \frac{x^3}{\tan(fx+e)b+a} dx \right) a^2 d^3 f + 2 \left(\int \frac{x^3}{\tan(fx+e)b+a} dx \right) b^2 d^3 f + 6 \left(\int \frac{x^2}{\tan(fx+e)b+a} dx \right) a^2 c d^2 f + 6 \left(\int \frac{x^2}{\tan(fx+e)b+a} dx \right) a^2 c d^2 f}{a + b \tan(e + fx)} \end{aligned}$$

input `int((d*x+c)^3/(a+b*tan(f*x+e)),x)`

output

```
(2*int(x**3/(tan(e + f*x)*b + a),x)*a**2*d**3*f + 2*int(x**3/(tan(e + f*x)
*b + a),x)*b**2*d**3*f + 6*int(x**2/(tan(e + f*x)*b + a),x)*a**2*c*d**2*f
+ 6*int(x**2/(tan(e + f*x)*b + a),x)*b**2*c*d**2*f + 6*int(x/(tan(e + f*x)
*b + a),x)*a**2*c**2*d*f + 6*int(x/(tan(e + f*x)*b + a),x)*b**2*c**2*d*f -
log(tan(e + f*x)**2 + 1)*b*c**3 + 2*log(tan(e + f*x)*b + a)*b*c**3 + 2*a*
c**3*f*x)/(2*f*(a**2 + b**2))
```

3.55 $\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$

Optimal result	433
Mathematica [A] (verified)	434
Rubi [A] (verified)	434
Maple [B] (verified)	437
Fricas [B] (verification not implemented)	438
Sympy [F]	439
Maxima [B] (verification not implemented)	440
Giac [F]	440
Mupad [F(-1)]	441
Reduce [F]	441

Optimal result

Integrand size = 20, antiderivative size = 181

$$\begin{aligned} \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx = & \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\ & - \frac{ibd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f^2} \\ & + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^3} \end{aligned}$$

output
$$\frac{1}{3}*(d*x+c)^3/(a+I*b)/d+b*(d*x+c)^2*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-I*b*d*(d*x+c)*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2+1/2*b*d^2*polylog(3,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^3$$

Mathematica [A] (verified)

Time = 1.30 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.30

$$\begin{aligned} & \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx \\ &= \frac{1}{6}b \left(-\frac{4(c+dx)^3}{(ia+b)d(-ib(-1+e^{2ie})+a(1+e^{2ie}))} \right. \\ & \quad + \frac{6(c+dx)^2 \log\left(1+\frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2+b^2)f} \\ & \quad + \frac{3d\left(2if(c+dx)\text{PolyLog}\left(2,\frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)+d\text{PolyLog}\left(3,\frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)\right)}{(a^2+b^2)f^3} \Big) \\ & \quad + \frac{x(3c^2+3cdx+d^2x^2)\cos(e)}{3(a\cos(e)+b\sin(e))} \end{aligned}$$

input `Integrate[(c + d*x)^2/(a + b*Tan[e + f*x]), x]`

output
$$\begin{aligned} & (b*((-4*(c + d*x)^3)/((I*a + b)*d*((-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))) + (6*(c + d*x)^2*\text{Log}[1 + (a + I*b)/((a - I*b)*E^{((2*I)*(e + f*x))})])/((a^2 + b^2)*f) + (3*d*((2*I)*f*(c + d*x)*\text{PolyLog}[2, (-a - I*b)/((a - I*b)*E^{((2*I)*(e + f*x))})] + d*\text{PolyLog}[3, (-a - I*b)/((a - I*b)*E^{((2*I)*(e + f*x))})]))/((a^2 + b^2)*f^3))/6 + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*\text{Cos}[e])/(3*(a*\text{Cos}[e] + b*\text{Sin}[e])) \end{aligned}$$

Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3042, 4215, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx \\
& \quad \downarrow \textcolor{blue}{3042} \\
& \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx \\
& \quad \downarrow \textcolor{blue}{4215} \\
& 2ib \int \frac{e^{2i(e+fx)}(c+dx)^2}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx + \frac{(c+dx)^3}{3d(a+ib)} \\
& \quad \downarrow \textcolor{blue}{2620} \\
& 2ib \left(\frac{id \int (c+dx) \log \left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) dx}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \\
& \quad \frac{(c+dx)^3}{3d(a+ib)} \\
& \quad \downarrow \textcolor{blue}{3011} \\
& 2ib \left(\frac{id \left(\frac{i(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{id \int \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{2f} \right)}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right. \\
& \quad \left. \frac{(c+dx)^3}{3d(a+ib)} \right. \\
& \quad \downarrow \textcolor{blue}{2720} \\
& 2ib \left(\frac{id \left(\frac{i(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{d \int e^{-2i(e+fx)} \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) de^{2i(e+fx)}}{4f^2} \right)}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i}}{(a+ib)^2} \right)}{2f} \right. \\
& \quad \left. \frac{(c+dx)^3}{3d(a+ib)} \right. \\
& \quad \downarrow \textcolor{blue}{7143}
\end{aligned}$$

$$2ib \left(\frac{id \left(\frac{i(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f} - \frac{d \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{4f^2} \right)}{f(a^2+b^2)} - \frac{i(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right. \\ \left. \frac{(c+dx)^3}{3d(a+ib)} \right)$$

input `Int[(c + d*x)^2/(a + b*Tan[e + f*x]), x]`

output `(c + d*x)^3/(3*(a + I*b)*d) + (2*I)*b*(((-1/2*I)*(c + d*x)^2*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) + (I*d*((I/2)*(c + d*x)*PolyLog[2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/f - (d*PolyLog[3, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/(4*f^2)))/((a^2 + b^2)*f))`

Definitions of rubi rules used

rule 2620 `Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_.))))^(n_.)*(c_.) + (d_.)*(x_.)^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] :> Simpl[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simpl[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simpl[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_) + (b_)*(x_)))^(n_.)]*((f_) + (g_) * (x_)^m_), x_Symbol] :> Simp[(-f + g*x)^m]*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4215

```
Int[((c_) + (d_)*(x_))^m/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_)*(a_ + (b_)*(x_))^p]/((d_)*(e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 939 vs. $2(166) = 332$.

Time = 0.55 (sec) , antiderivative size = 940, normalized size of antiderivative = 5.19

method	result
risch	$-\frac{d^2 x^3}{3(i b-a)} - \frac{d c x^2}{i b-a} - \frac{c^2 x}{i b-a} - \frac{c^3}{3 d (i b-a)} - \frac{2 b c d x^2}{(i a+b)(-i b-a)} - \frac{4 b c d e x}{f (i a+b)(-i b-a)} - \frac{2 b c d e^2}{f^2 (i a+b)(-i b-a)} - \frac{b c d \text{polylog}\left(2, \frac{-e x}{i a+b}\right)}{f^2 (i a+b)}$

input

```
int((d*x+c)^2/(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

output

$$\begin{aligned}
 & -\frac{1}{3}d^2/(I*b-a)*x^3-d/(I*b-a)*c*x^2-1/(I*b-a)*c^2*x-1/3/d/(I*b-a)*c^3-2/(\\
 & I*a+b)*b/(-I*b-a)*c*d*x^2-4/f/(I*a+b)*b/(-I*b-a)*c*d*e*x-2/f^2/(I*a+b)*b/(\\
 & -I*b-a)*c*d*e^2-1/f^2/(I*a+b)*b/(-I*b-a)*c*d*polylog(2,(a-I*b)*exp(2*I*(f* \\
 & x+e))/(-I*b-a))-2/3/(I*a+b)*b/(-I*b-a)*d^2*x^3+4/3/f^3/(I*a+b)*b/(-I*b-a)* \\
 & e^3*d^2-I/f/(I*a+b)*b/(-I*b-a)*d^2*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) \\
 & *x^2+2/f^2/(I*a+b)*b/(-I*b-a)*e^2*d^2*x-1/f^2/(I*a+b)*b/(-I*b-a)*d^2*polylog(2,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x+I/f/(I*a+b)*b*c^2/(a+I*b)*ln(I*b) \\
 & *exp(2*I*(f*x+e))-a*exp(2*I*(f*x+e))-I*b-a)+I/f^3/(I*a+b)*b/(-I*b-a)*e^2*d \\
 & ^2*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))+4*I/f^2/(I*a+b)*b*e*c*d/(a+I*b) \\
 & *ln(exp(I*(f*x+e)))-2*I/f/(I*a+b)*b/(-I*b-a)*c*d*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) \\
 & *x-2*I/f^3/(I*a+b)*b*e^2*d^2/(a+I*b)*ln(exp(I*(f*x+e)))-2*I/f^2/(I*a+b)*b/(-I*b-a)*c*d*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*e-1/2*I \\
 & /f^3/(I*a+b)*b/(-I*b-a)*d^2*polylog(3,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-2 \\
 & *I/f/(I*a+b)*b*c^2/(a+I*b)*ln(exp(I*(f*x+e)))-2*I/f^2/(I*a+b)*b*e*c*d/(a+I*b) \\
 & *ln(I*b*exp(2*I*(f*x+e))-a*exp(2*I*(f*x+e))-I*b-a)+I/f^3/(I*a+b)*b*e^2*d^2/(a+I*b)*ln(I*b*exp(2*I*(f*x+e))-a*exp(2*I*(f*x+e))-I*b-a)
 \end{aligned}$$
Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 834 vs. $2(160) = 320$.

Time = 0.12 (sec), antiderivative size = 834, normalized size of antiderivative = 4.61

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output

```
1/12*(4*a*d^2*f^3*x^3 + 12*a*c*d*f^3*x^2 + 12*a*c^2*f^3*x + 3*b*d^2*polylog(3, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e))^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 3*b*d^2*polylog(3, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e))^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 6*(-I*b*d^2*f*x - I*b*c*d*f)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*c^2*f^2)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*c^2*f^2)*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1))/((a^2 + b^2)*f^3)
```

Sympy [F]

$$\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx = \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$$

input

```
integrate((d*x+c)**2/(a+b*tan(f*x+e)),x)
```

output

```
Integral((c + d*x)**2/(a + b*tan(e + f*x)), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 715 vs. $2(160) = 320$.

Time = 0.40 (sec) , antiderivative size = 715, normalized size of antiderivative = 3.95

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output

```
-1/6*(6*c*d*e*(2*(f*x + e)*a/((a^2 + b^2)*f) + 2*b*log(b*tan(f*x + e) + a)/((a^2 + b^2)*f) - b*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*f)) - 3*(2*(f*x + e)*a/(a^2 + b^2) + 2*b*log(b*tan(f*x + e) + a)/(a^2 + b^2) - b*log(tan(f*x + e)^2 + 1)/(a^2 + b^2))*c^2 - (2*(f*x + e)^3*(a - I*b)*d^2 + 6*(f*x + e)*(a - I*b)*d^2*e^2 + 6*I*b*d^2*e^2*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) + 3*b*d^2*e^2*2*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 3*b*d^2*polylog(3, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 6*((a - I*b)*d^2*e - (a - I*b)*c*d*f)*(f*x + e)^2 - 6*(I*(f*x + e)^2*b*d^2 + 2*(-I*b*d^2*e + I*b*c*d*f)*(f*x + e))*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) - 6*(I*(f*x + e)*b*d^2 - I*b*d^2*e + I*b*c*d*f)*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) + 3*((f*x + e)^2*b*d^2 - 2*(b*d^2*e - b*c*d*f)*(f*x + e))*log(((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2))/((a^2 + b^2)*f^2))/f
```

Giac [F]

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(dx + c)^2}{b \tan(fx + e) + a} dx$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)^2/(b*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

input `int((c + d*x)^2/(a + b*tan(e + f*x)), x)`

output `int((c + d*x)^2/(a + b*tan(e + f*x)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx \\ &= \frac{2 \left(\int \frac{x^2}{\tan(fx+e)b+a} dx \right) a^2 d^2 f + 2 \left(\int \frac{x^2}{\tan(fx+e)b+a} dx \right) b^2 d^2 f + 4 \left(\int \frac{x}{\tan(fx+e)b+a} dx \right) a^2 c d f + 4 \left(\int \frac{x}{\tan(fx+e)b+a} dx \right) b^2 c d f}{2f(a^2 + b^2)} \end{aligned}$$

input `int((d*x+c)^2/(a+b*tan(f*x+e)), x)`

output `(2*int(x**2/(\tan(e + f*x)*b + a), x)*a**2*d**2*f + 2*int(x**2/(\tan(e + f*x)*b + a), x)*b**2*d**2*f + 4*int(x/(\tan(e + f*x)*b + a), x)*a**2*c*d*f + 4*int(x/(\tan(e + f*x)*b + a), x)*b**2*c*d*f - log(tan(e + f*x)**2 + 1)*b*c**2 + 2*log(tan(e + f*x)*b + a)*b*c**2 + 2*a*c**2*f*x)/(2*f*(a**2 + b**2))`

3.56 $\int \frac{c+dx}{a+b\tan(e+fx)} dx$

Optimal result	442
Mathematica [A] (verified)	442
Rubi [A] (verified)	443
Maple [B] (verified)	445
Fricas [B] (verification not implemented)	446
Sympy [F]	446
Maxima [B] (verification not implemented)	447
Giac [F]	447
Mupad [F(-1)]	448
Reduce [F]	448

Optimal result

Integrand size = 18, antiderivative size = 125

$$\int \frac{c+dx}{a+b\tan(e+fx)} dx = \frac{(c+dx)^2}{2(a+ib)d} + \frac{b(c+dx)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} - \frac{ibd \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2}$$

output

```
1/2*(d*x+c)^2/(a+I*b)/d+b*(d*x+c)*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-1/2*I*b*d*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2
```

Mathematica [A] (verified)

Time = 1.40 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.42

$$\int \frac{c+dx}{a+b\tan(e+fx)} dx = \frac{b(c+dx)^2}{(a-ib)d(b-be^{2ie}-ia(1+e^{2ie}))} + \frac{b(c+dx)\log\left(1+\frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2+b^2)f} + \frac{ibd \operatorname{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)}{2(a^2+b^2)f^2} + \frac{x(2c+dx)\cos(e)}{2(a\cos(e)+b\sin(e))}$$

input `Integrate[(c + d*x)/(a + b*Tan[e + f*x]), x]`

output
$$\begin{aligned} & \frac{(b*(c + d*x)^2)/((a - I*b)*d*(b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e))))}{+ (b*(c + d*x)*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f)} + ((I/2)*b*d*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f^2) + (x*(2*c + d*x)*Cos[e])/(2*(a*Cos[e] + b*Sin[e])) \end{aligned}$$

Rubi [A] (verified)

Time = 0.49 (sec), antiderivative size = 132, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4215, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{c + dx}{a + b \tan(e + fx)} dx \\ & \quad \downarrow \textcolor{blue}{3042} \\ & \int \frac{c + dx}{a + b \tan(e + fx)} dx \\ & \quad \downarrow \textcolor{blue}{4215} \\ & 2ib \int \frac{e^{2i(e+fx)}(c + dx)}{(a + ib)^2 + (a^2 + b^2) e^{2i(e+fx)}} dx + \frac{(c + dx)^2}{2d(a + ib)} \\ & \quad \downarrow \textcolor{blue}{2620} \\ & 2ib \left(\frac{id \int \log \left(\frac{e^{2i(e+fx)}(a^2 + b^2)}{(a + ib)^2} + 1 \right) dx}{2f(a^2 + b^2)} - \frac{i(c + dx) \log \left(1 + \frac{(a^2 + b^2)e^{2i(e+fx)}}{(a + ib)^2} \right)}{2f(a^2 + b^2)} \right) + \frac{(c + dx)^2}{2d(a + ib)} \\ & \quad \downarrow \textcolor{blue}{2715} \end{aligned}$$

$$\begin{aligned}
 & 2ib \left(\frac{d \int e^{-2i(e+fx)} \log \left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) de^{2i(e+fx)}}{4f^2(a^2+b^2)} - \frac{i(c+dx) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \\
 & \quad \frac{(c+dx)^2}{2d(a+ib)} \\
 & \quad \downarrow \text{2838} \\
 & 2ib \left(-\frac{i(c+dx) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} - \frac{d \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{4f^2(a^2+b^2)} \right) + \frac{(c+dx)^2}{2d(a+ib)}
 \end{aligned}$$

input `Int[(c + d*x)/(a + b*Tan[e + f*x]), x]`

output
$$(c + d*x)^2/(2*(a + I*b)*d) + (2*I)*b*(((-1/2*I)*(c + d*x))*Log[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) - (d*PolyLog[2, -((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/(4*(a^2 + b^2)*f^2)$$

Definitions of rubi rules used

rule 2620 `Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_)))^(n_.)]], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*(d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042

```
Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4215

```
Int[((c_.) + (d_)*(x_))^(m_.)/((a_) + (b_)*tan[(e_.) + (f_)*(x_)]), x_Sy
mbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b In
t[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2
*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2
, 0] && IGtQ[m, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 461 vs. 2(113) = 226.

Time = 0.49 (sec) , antiderivative size = 462, normalized size of antiderivative = 3.70

method	result
risch	$-\frac{dx^2}{2(ib-a)} - \frac{cx}{ib-a} + \frac{bc \ln(ib e^{2i(fx+e)} - a e^{2i(fx+e)} - ib - a)}{f(-ib+a)(ib+a)} - \frac{2bc \ln(e^{i(fx+e)})}{f(-ib+a)(ib+a)} - \frac{bd \ln\left(1 - \frac{(ib+a)e^{2i(fx+e)}}{-ib-a}\right)x}{f(-ib+a)(-ib-a)} + \frac{ie^{2i(fx+e)}}{(-ib-a)^2}$

input `int((d*x+c)/(a+b*tan(f*x+e)), x, method=_RETURNVERBOSE)`

output

```
-1/2/(I*b-a)*d*x^2-1/(I*b-a)*c*x+1/f*b/(a-I*b)*c/(a+I*b)*ln(I*b*exp(2*I*(f
*x+e))-a*exp(2*I*(f*x+e))-I*b-a)-2/f*b/(a-I*b)*c/(a+I*b)*ln(exp(I*(f*x+e))
)-1/f*b/(a-I*b)/(-I*b-a)*d*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x+I*b/(a-I*b)/(-I*b-a)*d*x^2-1/f^2*b/(a-I*b)/(-I*b-a)*d*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*e+2*I/f*b/(a-I*b)/(-I*b-a)*d*e*x+I/f^2*b/(a-I*b)/(-I*b-a)*d
*e^2+1/2*I/f^2*b/(a-I*b)/(-I*b-a)*d*polylog(2, (a-I*b)*exp(2*I*(f*x+e))/(-I
*b-a))-1/f^2*b/(a-I*b)*d*e/(a+I*b)*ln(I*b*exp(2*I*(f*x+e))-a*exp(2*I*(f*x+
e))-I*b-a)+2/f^2*b/(a-I*b)*d*e/(a+I*b)*ln(exp(I*(f*x+e)))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs. $2(108) = 216$.

Time = 0.09 (sec) , antiderivative size = 527, normalized size of antiderivative = 4.22

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \frac{2 adf^2 x^2 + 4 acf^2 x + i bd \text{Li}_2\left(\frac{2((i ab - b^2) \tan(fx+e)^2 - a^2 - i ab + (i a^2 - 2 ab - i b^2) \tan(fx+e))}{(a^2 + b^2) \tan(fx+e)^2 + a^2 + b^2}\right) + 1}{2} - i bd \text{Li}_2\left(\frac{2((-i ab - b^2) \tan(fx+e)^2 - a^2 - i ab + (i a^2 - 2 ab - i b^2) \tan(fx+e))}{(a^2 + b^2) \tan(fx+e)^2 + a^2 + b^2}\right) + 1$$

input `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output

```
1/4*(2*a*d*f^2*x^2 + 4*a*c*f^2*x + I*b*d*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - I*b*d*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(b*d*f*x + b*d*e)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(b*d*f*x + b*d*e)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 2*(b*d*e - b*c*f)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - 2*(b*d*e - b*c*f)*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)))/((a^2 + b^2)*f^2)
```

Sympy [F]

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{c + dx}{a + b \tan(e + fx)} dx$$

input `integrate((d*x+c)/(a+b*tan(f*x+e)),x)`

output `Integral((c + d*x)/(a + b*tan(e + f*x)), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 399 vs. $2(108) = 216$.

Time = 0.30 (sec) , antiderivative size = 399, normalized size of antiderivative = 3.19

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx \\ = \frac{(a - i b) df^2 x^2 + 2 (a - i b) c f^2 x - 2 i b d f x \arctan\left(\frac{2 a b \cos(2 f x + 2 e) - (a^2 - b^2) \sin(2 f x + 2 e)}{a^2 + b^2}\right), \frac{2 a b \sin(2 f x + 2 e) + a^2 + b^2 + (a^2 - b^2) \cos(2 f x + 2 e)}{a^2 + b^2}}{a^2 + b^2}$$

input `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output $\frac{1}{2} ((a - I b) d f^2 x^2 + 2 (a - I b) c f^2 x - 2 I b d f x \operatorname{arctan2}(2 a b \cos(2 f x + 2 e) - (a^2 - b^2) \sin(2 f x + 2 e)) / (a^2 + b^2), (2 a b \sin(2 f x + 2 e) + a^2 + b^2 + (a^2 - b^2) \cos(2 f x + 2 e)) / (a^2 + b^2)) + b d f x \log((a^2 + b^2) \cos(2 f x + 2 e))^2 + 4 a b \sin(2 f x + 2 e) + (a^2 + b^2) \sin(2 f x + 2 e)^2 + a^2 + b^2 + 2 (a^2 - b^2) \cos(2 f x + 2 e)) / (a^2 + b^2) + 2 I b c f \operatorname{arctan2}(-b \cos(2 f x + 2 e) + a \sin(2 f x + 2 e) + b, a \cos(2 f x + 2 e) + b \sin(2 f x + 2 e) + a) + b c f \log((a^2 + b^2) \cos(2 f x + 2 e))^2 + 4 a b \sin(2 f x + 2 e) + (a^2 + b^2) \sin(2 f x + 2 e)^2 + a^2 + b^2 + 2 (a^2 - b^2) \cos(2 f x + 2 e)) - I b d \operatorname{dilog}((I a + b) e^{-2 I f x - 2 I e}) / ((a^2 + b^2) f^2)$

Giac [F]

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{dx + c}{b \tan(f x + e) + a} dx$$

input `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate((d*x + c)/(b*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{c + d x}{a + b \tan (e + f x)} dx$$

input `int((c + d*x)/(a + b*tan(e + f*x)),x)`

output `int((c + d*x)/(a + b*tan(e + f*x)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{c + dx}{a + b \tan(e + fx)} dx \\ &= \frac{2 \left(\int \frac{x}{\tan(fx+e)b+a} dx \right) a^2 df + 2 \left(\int \frac{x}{\tan(fx+e)b+a} dx \right) b^2 df - \log(\tan(fx+e)^2 + 1) bc + 2 \log(\tan(fx+e) b + a) c}{2f(a^2 + b^2)} \end{aligned}$$

input `int((d*x+c)/(a+b*tan(f*x+e)),x)`

output `(2*int(x/(tan(e + f*x)*b + a),x)*a**2*d*f + 2*int(x/(tan(e + f*x)*b + a),x)*b**2*d*f - log(tan(e + f*x)**2 + 1)*b*c + 2*log(tan(e + f*x)*b + a)*b*c + 2*a*c*f*x)/(2*f*(a**2 + b**2))`

3.57 $\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$

Optimal result	449
Mathematica [N/A]	449
Rubi [N/A]	450
Maple [N/A]	451
Fricas [N/A]	451
Sympy [N/A]	451
Maxima [N/A]	452
Giac [N/A]	452
Mupad [N/A]	453
Reduce [N/A]	453

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b\tan(e+fx))}, x\right)$$

output `Defer(Int)(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

Mathematica [N/A]

Not integrable

Time = 2.14 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$$

input `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])),x]`

output `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])), x]`

Rubi [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx \end{aligned}$$

input `Int[1/((c + d*x)*(a + b*Tan[e + f*x])),x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tan(fx + e))} dx$$

input `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

output `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.35

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output `integral(1/(a*d*x + a*c + (b*d*x + b*c)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 0.95 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

output `Integral(1/((a + b*tan(e + f*x))*(c + d*x)), x)`

Maxima [N/A]

Not integrable

Time = 0.77 (sec) , antiderivative size = 279, normalized size of antiderivative = 13.95

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output
$$\begin{aligned} & (2*(a^2*b + b^3)*d*integrate((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/((a^4 + 2*a^2*b^2 + b^4)*d*x + ((a^4 + 2*a^2*b^2 + b^4)*d*x + (a^4 + 2*a^2*b^2 + b^4)*c)*cos(2*f*x + 2*e)^2 + ((a^4 + 2*a^2*b^2 + b^4)*d*x + (a^4 + 2*a^2*b^2 + b^4)*c)*sin(2*f*x + 2*e)^2 + (a^4 + 2*a^2*b^2 + b^4)*c + 2*((a^4 - b^4)*d*x + (a^4 - b^4)*c)*cos(2*f*x + 2*e) + 4*((a^3*b + a*b^3)*d*x + (a^3*b + a*b^3)*c)*sin(2*f*x + 2*e)), x) + a*log(d*x + c))/((a^2 + b^2)*d) \end{aligned}$$

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate(1/((d*x + c)*(b*tan(f*x + e) + a)), x)`

Mupad [N/A]

Not integrable

Time = 9.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)} dx$$

input `int(1/((a + b*tan(e + f*x))*(c + d*x)),x)`

output `int(1/((a + b*tan(e + f*x))*(c + d*x)), x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.55

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{\tan(fx + e) bc + \tan(fx + e) bd x + ac + ad x} dx$$

input `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

output `int(1/(\tan(e + f*x)*b*c + \tan(e + f*x)*b*d*x + a*c + a*d*x),x)`

3.58 $\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$

Optimal result	454
Mathematica [N/A]	454
Rubi [N/A]	455
Maple [N/A]	456
Fricas [N/A]	456
Sympy [N/A]	456
Maxima [N/A]	457
Giac [N/A]	457
Mupad [N/A]	458
Reduce [N/A]	458

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b\tan(e+fx))}, x\right)$$

output `Defer(Int)(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`

Mathematica [N/A]

Not integrable

Time = 4.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

input `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])),x]`

output `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]`

Rubi [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx \end{aligned}$$

input `Int[1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2 (a + b \tan(fx + e))} dx$$

input `int(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`

output `int(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="fricas")`

output `integral(1/(a*d^2*x^2 + 2*a*c*d*x + a*c^2 + (b*d^2*x^2 + 2*b*c*d*x + b*c^2)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.75 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)^2} dx$$

input `integrate(1/(d*x+c)**2/(a+b*tan(f*x+e)),x)`

output `Integral(1/((a + b*tan(e + f*x))*(c + d*x)**2), x)`

Maxima [N/A]

Not integrable

Time = 1.79 (sec) , antiderivative size = 424, normalized size of antiderivative = 21.20

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="maxima")`

output `(2*((a^2*b + b^3)*d^2*x + (a^2*b + b^3)*c*d)*integrate((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2 + ((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2)*cos(2*f*x + 2*e)^2 + ((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2)*sin(2*f*x + 2*e)^2 + 2*((a^4 - b^4)*d^2*x^2 + 2*(a^4 - b^4)*c*d*x + (a^4 - b^4)*c^2)*cos(2*f*x + 2*e) + 4*((a^3*b + a*b^3)*d^2*x^2 + 2*(a^3*b + a*b^3)*c*d*x + (a^3*b + a*b^3)*c^2)*sin(2*f*x + 2*e)), x) - a)/((a^2 + b^2)*d^2*x + (a^2 + b^2)*c*d)`

Giac [N/A]

Not integrable

Time = 4.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="giac")`

output `integrate(1/((d*x + c)^2*(b*tan(f*x + e) + a)), x)`

Mupad [N/A]

Not integrable

Time = 9.51 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)^2} dx$$

input `int(1/((a + b*tan(e + f*x))*(c + d*x)^2),x)`

output `int(1/((a + b*tan(e + f*x))*(c + d*x)^2), x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 61, normalized size of antiderivative = 3.05

$$\begin{aligned} & \int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx \\ &= \int \frac{1}{\tan(fx + e) b c^2 + 2 \tan(fx + e) b c d x + \tan(fx + e) b d^2 x^2 + a c^2 + 2 a c d x + a d^2 x^2} dx \end{aligned}$$

input `int(1/(d*x+c)^2/(a+b*tan(f*x+e)),x)`

output `int(1/(tan(e + f*x)*b*c**2 + 2*tan(e + f*x)*b*c*d*x + tan(e + f*x)*b*d**2*x**2 + a*c**2 + 2*a*c*d*x + a*d**2*x**2),x)`

$$\mathbf{3.59} \quad \int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$$

Optimal result	460
Mathematica [B] (warning: unable to verify)	461
Rubi [A] (verified)	462
Maple [B] (verified)	465
Fricas [B] (verification not implemented)	465
Sympy [F]	466
Maxima [B] (verification not implemented)	467
Giac [F]	468
Mupad [F(-1)]	468
Reduce [F]	468

Optimal result

Integrand size = 20, antiderivative size = 848

$$\begin{aligned}
 \int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx = & -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} \\
 & + \frac{2b^2(c+dx)^3}{(a+ib)(ia+b)^2 (ia-b+(ia+b)e^{2ie+2ifx}) f} \\
 & + \frac{(c+dx)^4}{4(a-ib)^2 d} + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} \\
 & + \frac{3b^2d(c+dx)^2 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^2} \\
 & + \frac{2b(c+dx)^3 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a-ib)^2(a+ib)f} \\
 & - \frac{2ib^2(c+dx)^3 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f} \\
 & - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^3} \\
 & + \frac{3bd(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(ia-b)(a-ib)^2 f^2} \\
 & - \frac{3b^2d(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^2} \\
 & + \frac{3b^2d^3 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(a^2+b^2)^2 f^4} \\
 & + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a-ib)^2(a+ib)f^3} \\
 & - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^3} \\
 & - \frac{3bd^3 \operatorname{PolyLog} \left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(ia-b)(a-ib)^2 f^4} \\
 & + \frac{3b^2d^3 \operatorname{PolyLog} \left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(a^2+b^2)^2 f^4}
 \end{aligned}$$

output

```

-3*I*b^2*d^2*(d*x+c)*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3+2*b^2*(d*x+c)^3/(a+I*b)/(I*a-b+(I*a+b)*exp(2*I*e+2*I*f*x))/f+1/4*(d*x+c)^4/(a-I*b)^2/d+b*(d*x+c)^4/(I*a-b)/(a-I*b)^2/d-b^2*(d*x+c)^4/(a^2+b^2)^2/d+3*b^2*d*(d*x+c)^2*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+2*b*(d*x+c)^3*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f-2*I*b^2*(d*x+c)^3/(a^2+b^2)^2/f-3*I*b^2*d^2*(d*x+c)*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3+3*b*d*(d*x+c)^2*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(I*a-b)/(a-I*b)^2/f^2-3*b^2*d*(d*x+c)^2*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+2*f^2+3/2*b^2*d^3*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^4+3*b*d^2*(d*x+c)*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f^3-2*I*b^2*(d*x+c)^3*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2-3/2*b*d^3*polylog(4,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(I*a-b)/(a-I*b)^2/f^4+3/2*b^2*d^3*polylog(4,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^4

```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1724 vs. $2(848) = 1696$.

Time = 9.24 (sec) , antiderivative size = 1724, normalized size of antiderivative = 2.03

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input

```
Integrate[(c + d*x)^3/(a + b*Tan[e + f*x])^2,x]
```

output

$$\begin{aligned}
 & (b*(-4*c^2*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((3*b*d + 2*a*c*f)*x)/(a^2 + b^2) + (4*b*(c + d*x)^3)/(a - I*b) + (2*a*f*(c + d*x)^4)/((a - I*b)*d) + (12*c*d*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((b*d + a*c*f)*x*Log[1 + (a + I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a + I*b)*(I*a + b)*f) + (6*d^2*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((b*d + 2*a*c*f)*x^2*Log[1 + (a + I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a + I*b)*(I*a + b)*f) + (4*a*d^3*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*x^3*Log[1 + (a + I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a + I*b)*(I*a + b)) + (2*c^2*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((3*b*d + 2*a*c*f)*Log[a + I*b + (a - I*b)*E^{(2*I)*(e + f*x)}])/((a + I*b)*(I*a + b)*f) + (6*c*d*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((b*d + a*c*f)*PolyLog[2, (-a - I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a^2 + b^2)*f^2) + (3*d^2*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((b*d + 2*a*c*f)*(2*f*x*PolyLog[2, (-a - I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a^2 + b^2)*f^3) + (3*a*d^3*(-I)*b*(-1 + E^{(2*I)*e}) + a*(1 + E^{(2*I)*e}))*((2*f^2*x^2*PolyLog[2, (-a - I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})] - (2*I)*f*x*PolyLog[3, (-a - I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})] - PolyLog[4, (-a - I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})]))/((a^2 + b^2)*f^3))/((2*(a - I*b)*(a + I*b)*(b - b*E^{(2*I)*e}) - I*a*(1 + E^{(2*I)*e})*f) + (3*x^2*(a*c^2*d...
 \end{aligned}$$

Rubi [A] (verified)

Time = 2.28 (sec), antiderivative size = 848, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3042, 4217, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4217}
 \end{aligned}$$

$$\begin{aligned}
& \int \left(-\frac{4b^2(c+dx)^3}{(b+ia)^2 \left(ia \left(1 - \frac{ib}{a} \right) e^{2ie+2ifx} + ia \left(1 + \frac{ib}{a} \right) \right)^2} + \frac{4b(c+dx)^3}{(a-ib)^2 \left(ia \left(1 - \frac{ib}{a} \right) e^{2ie+2ifx} + ia \left(1 + \frac{ib}{a} \right) \right)} + \frac{(c+dx)^5}{(a-ib)^2} \right. \\
& \quad \left. \downarrow \text{2009} \right. \\
& \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} + \frac{(c+dx)^4}{4(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} + \frac{2b \log \left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1 \right) (c+dx)^3}{(a-ib)^2(a+ib)f} - \\
& \frac{2ib^2 \log \left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1 \right) (c+dx)^3}{(a^2+b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a+ib)(ia+b)^2 (ia+(ia+b)e^{2ie+2ifx}-b) f} - \\
& \frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} + \frac{3b^2 d \log \left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1 \right) (c+dx)^2}{(a^2+b^2)^2 f^2} + \\
& \frac{3bd \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right) (c+dx)^2}{(ia-b)(a-ib)^2 f^2} - \frac{3b^2 d \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right) (c+dx)^2}{(a^2+b^2)^2 f^2} - \\
& \frac{3ib^2 d^2 \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right) (c+dx)}{(a^2+b^2)^2 f^3} + \frac{3bd^2 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right) (c+dx)}{(a-ib)^2(a+ib)f^3} - \\
& \frac{3ib^2 d^2 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right) (c+dx)}{(a^2+b^2)^2 f^3} + \frac{3b^2 d^3 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(a^2+b^2)^2 f^4} - \\
& \frac{3bd^3 \operatorname{PolyLog} \left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(ia-b)(a-ib)^2 f^4} + \frac{3b^2 d^3 \operatorname{PolyLog} \left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(a^2+b^2)^2 f^4}
\end{aligned}$$

input $\operatorname{Int}[(c + d*x)^3/(a + b*\operatorname{Tan}[e + f*x])^2, x]$

output

$$\begin{aligned} & \frac{(-2I)*b^2*(c + d*x)^3}{((a^2 + b^2)^2*f)} + \frac{(2*b^2*(c + d*x)^3)/((a + I*b)*(I*a + b)^2*(I*a - b + (I*a + b)*E^{((2*I)*e + (2*I)*f*x))*f)} + (c + d*x)^4/(4*(a - I*b)^2*d) + (b*(c + d*x)^4)/((I*a - b)*(a - I*b)^2*d) - (b^2*(c + d*x)^4)/((a^2 + b^2)^2*d) + (3*b^2*d*(c + d*x)^2*\text{Log}[1 + ((a - I*b)*E^{(2*I)*e + (2*I)*f*x})/(a + I*b)])/((a^2 + b^2)^2*f^2) + (2*b*(c + d*x)^3*\text{Log}[1 + ((a - I*b)*E^{(2*I)*e + (2*I)*f*x})/(a + I*b)])/((a - I*b)^2*(a + I*b)*f) - ((2*I)*b^2*(c + d*x)^3*\text{Log}[1 + ((a - I*b)*E^{(2*I)*e + (2*I)*f*x})/(a + I*b)])/((a^2 + b^2)^2*f) - ((3*I)*b^2*d^2*(c + d*x)*\text{PolyLog}[2, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/((a^2 + b^2)^2*f^3) + (3*b*d*(c + d*x)^2*\text{PolyLog}[2, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/((I*a - b)*(a - I*b)^2*f^2) - (3*b^2*d*(c + d*x)^2*\text{PolyLog}[2, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/((a^2 + b^2)^2*f^2) + (3*b^2*d^3*\text{PolyLog}[3, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/(2*(a^2 + b^2)^2*f^4) + (3*b*d^2*(c + d*x)*\text{PolyLog}[3, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/((a - I*b)^2*(a + I*b)*f^3) - ((3*I)*b^2*d^2*(c + d*x)*\text{PolyLog}[3, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/((a^2 + b^2)^2*f^3) - (3*b*d^3*\text{PolyLog}[4, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/(2*(I*a - b)*(a - I*b)^2*f^4) + (3*b^2*d^3*\text{PolyLog}[4, -(((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b))]/(2*(a^2 + b^2)^2*f^4) \end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4217 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_)}, x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 + b^2 + (a - I*b)^2*E^{(2*I*(e + f*x)})))^{(-n)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{NeQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, 0] \&& \text{IGtQ}[m, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 5958 vs. $2(763) = 1526$.

Time = 0.94 (sec) , antiderivative size = 5959, normalized size of antiderivative = 7.03

method	result	size
risch	Expression too large to display	5959

input `int((d*x+c)^3/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2537 vs. $2(695) = 1390$.

Time = 0.14 (sec) , antiderivative size = 2537, normalized size of antiderivative = 2.99

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

```
1/4*((a^3 - a*b^2)*d^3*f^4*x^4 - 4*b^3*c^3*f^3 - 4*(b^3*d^3*f^3 - (a^3 - a
*b^2)*c*d^2*f^4)*x^3 - 6*(2*b^3*c*d^2*f^3 - (a^3 - a*b^2)*c^2*d*f^4)*x^2 -
4*(3*b^3*c^2*d*f^3 - (a^3 - a*b^2)*c^3*f^4)*x - 6*(-I*a^2*b*d^3*f^2*x^2 -
I*a^2*b*c^2*d*f^2 - I*a*b^2*c*d^2*f - I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)
*x + (-I*a*b^2*d^3*f^2*x^2 - I*a*b^2*c^2*d*f^2 - I*b^3*c*d^2*f - I*(2*a*b^2*c*d^2*f^2 + b^3*d^3*f)*x)*tan(f*x + e))*dilog(2*((I*a*b - b^2)*tan(f*x +
e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*a^2*b*d^3*f^2*x^2 + I*a^2*b*c^2*d*f^2
+ I*a*b^2*c*d^2*f + I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)*x + (I*a*b^2*d^3*f^2*x^2 + I*a*b^2*c^2*d*f^2 + I*b^3*c*d^2*f + I*(2*a*b^2*c*d^2*f^2 + b^3*d^3*f)*x)*tan(f*x + e))*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(2*a^2*b*d^3*f^3*x^3 + 2*a^2*b*d^3*e^3 + 6*a^2*b*c^2*d*e*f^2 - 3*a*b^2*d^3*e^2 + 3*(2*a^2*b*c*d^2*f^3 + a*b^2*d^3*f^2)*x^2 - 6*
(a^2*b*c*d^2*e^2 - a*b^2*c*d^2*e)*f + 6*(a^2*b*c^2*d*f^3 + a*b^2*c*d^2*f^2
)*x + (2*a*b^2*d^3*f^3*x^3 + 2*a*b^2*d^3*e^3 + 6*a*b^2*c^2*d^2*e*f^2 - 3*b^3*d^3*e^2 + 3*(2*a*b^2*c*d^2*f^3 + b^3*d^3*f^2)*x^2 - 6*(a*b^2*c*d^2*e^2 - b^3*c*d^2*e)*f + 6*(a*b^2*c^2*d*f^3 + b^3*c*d^2*f^2)*x)*tan(f*x + e))*log(
-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(2*a^2*b*d^3*...
```

Sympy [F]

$$\int \frac{(c + dx)^3}{(a + b\tan(e + fx))^2} dx = \int \frac{(c + dx)^3}{(a + b\tan(e + fx))^2} dx$$

input `integrate((d*x+c)**3/(a+b*tan(f*x+e))**2,x)`

output `Integral((c + d*x)**3/(a + b*tan(e + f*x))**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4631 vs. $2(695) = 1390$.

Time = 2.83 (sec), antiderivative size = 4631, normalized size of antiderivative = 5.46

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output

```
-1/12*(36*c^2*d*e*(2*a*b*log(b*tan(f*x + e) + a)/((a^4 + 2*a^2*b^2 + b^4)*f) - a*b*log(tan(f*x + e)^2 + 1)/((a^4 + 2*a^2*b^2 + b^4)*f) - b/((a^2*b + b^3)*f*tan(f*x + e) + (a^3 + a*b^2)*f) + (a^2 - b^2)*(f*x + e)/((a^4 + 2*a^2*b^2 + b^4)*f)) - 12*(2*a*b*log(b*tan(f*x + e) + a)/(a^4 + 2*a^2*b^2 + b^4) - a*b*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + (a^2 - b^2)*(f*x + e)/(a^4 + 2*a^2*b^2 + b^4) - b/(a^3 + a*b^2 + (a^2*b + b^3)*tan(f*x + e)))*c^3 - (3*(a^3 - I*a^2*b + a*b^2 - I*b^3)*(f*x + e)^4*d^3 - 24*(I*a*b^2 - b^3)*d^3*e^3 - 72*(-I*a*b^2 + b^3)*c*d^2*e^2*f - 12*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e - (a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*f)*(f*x + e)^3 + 18*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e^2 - 2*(a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*f*f + (a^3 - I*a^2*b + a*b^2 - I*b^3)*c^2*d*f^2)*(f*x + e)^2 - 12*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e^3 - 3*(a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*e^2*f)*(f*x + e) - 12*(2*(I*a^2*b - a*b^2)*d^3*e^3 + 3*(-I*a*b^2 + b^3)*d^3*e^2 + 3*(-I*a*b^2 + b^3)*c^2*d*f^2 + 6*((-I*a^2*b + a*b^2)*c*d^2*e^2 + (I*a*b^2 - b^3)*c*d^2*e)*f + (2*(I*a^2*b + a*b^2)*d^3*e^3 + 3*(-I*a*b^2 - b^3)*d^3*e^2 + 3*(-I*a*b^2 - b^3)*c^2*d*f^2 + 6*((-I*a^2*b - a*b^2)*c*d^2*e^2 + (I*a*b^2 + b^3)*c*d^2*e)*f)*cos(2*f*x + 2*e) - (2*(a^2*b - I*a*b^2)*d^3*e^3 - 3*(a*b^2 - I*b^3)*d^3*e^2 - 3*(a*b^2 - I*b^3)*c^2*d*f^2 - 6*((a^2*b - I*a*b^2)*c*d^2*e^2 - (a*b^2 - I*b^3)*c*d^2*e)*f)*sin(2*f*x + 2*e)*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, ...)
```

Giac [F]

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \int \frac{(dx + c)^3}{(b \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^3/(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx$$

input `int((c + d*x)^3/(a + b*tan(e + f*x))^2,x)`

output `int((c + d*x)^3/(a + b*tan(e + f*x))^2, x)`

Reduce [F]

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `int((d*x+c)^3/(a+b*tan(f*x+e))^2,x)`

output

```
(int(x**3/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*d**3*f + 2*int(x**3/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**3*b**3*d**3*f + int(x**3/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**5*d**3*f + int(x**3/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**6*d**3*f + 2*int(x**3/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**4*b**2*d**3*f + int(x**3/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*b**4*d**3*f + 3*int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*c*d**2*f + 6*int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**3*b**3*c*d**2*f + 3*int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**4*b**2*c*d**2*f + 3*int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*b**4*c*d**2*f + 3*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*c**2*d*f + 6*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**3*b**3*c**2*d*f + 3*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**5*c**2*d*f + 3*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**6*c**2*d*f + 6*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**2*b**4*c**2*d*f + 3*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*c**2*d*f + 6*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**3*b**3*c**2*d*f + 3*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**5*c**2*d*f + 3*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**6*c**2*d*f + 6*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a...)
```

$$\mathbf{3.60} \quad \int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx$$

Optimal result	471
Mathematica [A] (warning: unable to verify)	472
Rubi [A] (verified)	473
Maple [B] (verified)	475
Fricas [B] (verification not implemented)	476
Sympy [F]	477
Maxima [B] (verification not implemented)	478
Giac [F]	479
Mupad [F(-1)]	479
Reduce [F]	479

Optimal result

Integrand size = 20, antiderivative size = 654

$$\begin{aligned}
 \int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx = & -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} \\
 & + \frac{2b^2(c+dx)^2}{(a+ib)(ia+b)^2 (ia-b+(ia+b)e^{2ie+2ifx}) f} \\
 & + \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(ia-b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} \\
 & + \frac{2b^2 d(c+dx) \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^2} \\
 & + \frac{2b(c+dx)^2 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a-ib)^2 (a+ib) f} \\
 & - \frac{2ib^2(c+dx)^2 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f} \\
 & - \frac{ib^2 d^2 \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^3} \\
 & + \frac{2bd(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(ia-b)(a-ib)^2 f^2} \\
 & - \frac{2b^2 d(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^2} \\
 & + \frac{bd^2 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a-ib)^2 (a+ib) f^3} \\
 & - \frac{ib^2 d^2 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^3}
 \end{aligned}$$

output

$$\begin{aligned}
 & -2*I*b^2*(d*x+c)^2/(a^2+b^2)^2/f + 2*b^2*(d*x+c)^2/(a+I*b)/(I*a-b+ \\
 & (I*a+b)*exp(2*I*e+2*I*f*x))/f + 1/3*(d*x+c)^3/(a-I*b)^2/d + 4/3*b*(d*x+c)^3/(I \\
 & *a-b)/(a-I*b)^2/d - 4/3*b^2*(d*x+c)^3/(a^2+b^2)^2/d + 2*b^2*d*(d*x+c)*ln(1+(a- \\
 & I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2 + 2*b*(d*x+c)^2*ln(1+(a-I*b) \\
 &)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f - 2*I*b^2*(d*x+c)^2*ln(1+(a- \\
 & I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f - I*b^2*d^2*polylog(2, -(a-I \\
 & *b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3 + 2*b*d*(d*x+c)*polylog(2, -(a- \\
 & I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2 + b*d^2*polylog(3, -(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f^3 - I*b^2*d^2*polylog(3, -(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3
 \end{aligned}$$
Mathematica [A] (warning: unable to verify)

Time = 6.11 (sec), antiderivative size = 761, normalized size of antiderivative = 1.16

$$\begin{aligned}
 & \int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx \\
 & = \frac{2b \left(\frac{12c(bd+acf)x}{a-ib} - \frac{12c(-ib(-1+e^{2ie})+a(1+e^{2ie}))}{a^2+b^2}(bd+acf)x + \frac{6d(bd+2acf)x^2}{a-ib} + \frac{4ad^2fx^3}{a-ib} + \frac{6d(-ib(-1+e^{2ie})+a(1+e^{2ie}))(bd+2acf)x \log\left(1+\frac{(a+ib)e}{a-ib}\right)}{(a+ib)(ia+b)f} \right)}{a^2+b^2}
 \end{aligned}$$

input

```
Integrate[(c + d*x)^2/(a + b*Tan[e + f*x])^2, x]
```

output

```
((2*b*((12*c*(b*d + a*c*f)*x)/(a - I*b) - (12*c*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + a*c*f)*x)/(a^2 + b^2) + (6*d*(b*d + 2*a*c*f)*x^2)/(a - I*b) + (4*a*d^2*f*x^3)/(a - I*b) + (6*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + 2*a*c*f)*x*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)*f) + (6*a*d^2*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*x^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)) + (6*c*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + a*c*f)*Log[a + I*b + (a - I*b)*E^((2*I)*(e + f*x))])/((a + I*b)*(I*a + b)*f) + (3*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + 2*a*c*f)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f^2) + (3*a*d^2*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(2*f*x*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - I*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^2))/((b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e))) + ((a^2 - b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[f*x] + (a^2 + b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[2*e + f*x] + 2*b*(3*b*(c + d*x)^2 + a*f*x*(3*c^2 + 3*c*d*x + d^2*x^2))*Sin[f*x])/((a*Cos[e] + b*Sin[e])*(a*Cos[e + f*x] + b*Sin[e + f*x])))/(6*(a^2 + b^2)*f)
```

Rubi [A] (verified)

Time = 1.79 (sec), antiderivative size = 654, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3042, 4217, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx \\
 & \quad \downarrow \textcolor{blue}{4217} \\
 & \int \left(-\frac{4b^2(c + dx)^2}{(b + ia)^2 \left(ia \left(1 - \frac{ib}{a} \right) e^{2ie+2ifx} + ia \left(1 + \frac{ib}{a} \right) \right)^2} + \frac{4b(c + dx)^2}{(a - ib)^2 \left(ia \left(1 - \frac{ib}{a} \right) e^{2ie+2ifx} + ia \left(1 + \frac{ib}{a} \right) \right)} + \frac{(c + dx)^2}{(a - ib)^2} \right) dx
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{2009} \\
& -\frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^2(a^2+b^2)^2} + \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^2(a^2+b^2)^2} - \\
& \frac{2ib^2(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f(a^2+b^2)^2} - \frac{2ib^2(c+dx)^2}{f(a^2+b^2)^2} - \frac{4b^2(c+dx)^3}{3d(a^2+b^2)^2} - \\
& \frac{ib^2 d^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^3(a^2+b^2)^2} - \frac{ib^2 d^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^3(a^2+b^2)^2} + \\
& \frac{2b^2(c+dx)^2}{f(a+ib)(b+ia)^2((b+ia)e^{2ie+2ifx}+ia-b)} + \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^2(-b+ia)(a-ib)^2} + \\
& \frac{2b(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f(a-ib)^2(a+ib)} + \frac{4b(c+dx)^3}{3d(-b+ia)(a-ib)^2} + \frac{(c+dx)^3}{3d(a-ib)^2} + \\
& \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^3(a-ib)^2(a+ib)}
\end{aligned}$$

input `Int[(c + d*x)^2/(a + b*Tan[e + f*x])^2, x]`

output

$$\begin{aligned}
& ((-2*I)*b^2*(c + d*x)^2)/((a^2 + b^2)^2*f) + (2*b^2*(c + d*x)^2)/((a + I*b)*(I*a + b)^2*(I*a - b + (I*a + b)*E^((2*I)*e + (2*I)*f*x))*f) + (c + d*x)^3/(3*(a - I*b)^2*d) + (4*b*(c + d*x)^3)/(3*(I*a - b)*(a - I*b)^2*d) - (4*b^2*(c + d*x)^3)/(3*(a^2 + b^2)^2*d) + (2*b^2*d*(c + d*x)*\operatorname{Log}[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)])/((a^2 + b^2)^2*f^2) + (2*b*(c + d*x)^2*\operatorname{Log}[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)])/((a - I*b)^2*(a + I*b)*f) - ((2*I)*b^2*(c + d*x)^2*\operatorname{Log}[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)])/((a^2 + b^2)^2*f) - (I*b^2*d^2*\operatorname{PolyLog}[2, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a^2 + b^2)^2*f^3) + (2*b*d*(c + d*x)*\operatorname{PolyLog}[2, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((I*a - b)*(a - I*b)^2*f^2) - (2*b^2*d*(c + d*x)*\operatorname{PolyLog}[2, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a^2 + b^2)^2*f^2) + (b*d^2*\operatorname{PolyLog}[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a - I*b)^2*(a + I*b)*f^3) - (I*b^2*d^2*\operatorname{PolyLog}[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a^2 + b^2)^2*f^3)
\end{aligned}$$

Definitions of rubi rules used

rule 2009 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \ x], \ x] /; \ \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \ x], \ x] /; \ \text{FunctionOfTrigOfLinearQ}[u, \ x]$

rule 4217 $\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, \ x_\text{Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 + b^2 + (a - I*b)^2*E^(2*I*(e + f*x)))))^{(-n)}, x], \ x] /; \ \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&& \ \text{NeQ}[a^2 + b^2, 0] \ \&& \ \text{ILtQ}[n, 0] \ \&& \ \text{IGtQ}[m, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3900 vs. $2(588) = 1176$.

Time = 0.79 (sec), antiderivative size = 3901, normalized size of antiderivative = 5.96

method	result	size
risch	Expression too large to display	3901

input $\text{int}((d*x+c)^2/(a+b*\tan(f*x+e))^2, x, \text{method}=\text{RETURNVERBOSE})$

output

$$\begin{aligned}
 & -\frac{1}{3}d^2/(2*I*a*b-a^2+b^2)*x^3-1/(2*I*a*b-a^2+b^2)*c^2*x-1/3/d/(2*I*a*b-a^2+b^2)*c^3-8/3/(I*a+b)^2/f^3/(b-I*a)*b/(a+I*b)*e^3*a*d^2+4/(I*a+b)^2/f^2/(b-I*a)*b^2/(a+I*b)*d^2*e*x+4/(I*a+b)^2/(b-I*a)*b/(a+I*b)*d*a*c*x^2+4/3/(I*a+b)^2/(b-I*a)*b/(a+I*b)*d^2*a*x^3+2/(I*a+b)^2/f/(b-I*a)*b^2/(a+I*b)*d^2*x^2+2/(I*a+b)^2/f^3/(b-I*a)*b^2/(a+I*b)*d^2*e^2+1/(I*a+b)^2/f^3/(b-I*a)*b^2/(a+I*b)*d^2*polylog(2,(I*b-a)*exp(2*I*(f*x+e))/(a+I*b))-d/(2*I*a*b-a^2+b^2)*c*x^2-1/(I*a+b)^2/f^3/(b-I*a)*b^2*e^2*a*d^2/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))+exp(4*I*(f*x+e))*b^2+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^2)+2/(I*a+b)^2/f^3/(b-I*a)*b^2*e*d^2/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b^2*e*d^2/(a+I*b)/(I*b-a)*arctan(1/b*a)*a+2/(I*a+b)^2/f^2/(b-I*a)*b^2*c*d/(a+I*b)/(I*b-a)*arctan(1/b*a)*a-2/(I*a+b)^2/f^2/(b-I*a)*b^2*c*d/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b*e^2*a^2*d^2/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b*e^2*a^2*d^2/(a+I*b)/(I*b-a)*arctan(1/b*a)*a+2/(I*a+b)^2/f^2/(b-I*a)*b^2*c*d/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b^3*e*d^2/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-2*I/(I*a+b)^2/f^3/(b-I*a)*b^3*e*d^2/(a+I*b)/(I*b-a)*arctan(1/b*a)+2*I/(I*a+b)^2/f^3/(b-I*a)*b^2*a*c^2/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b^2*a*c^2/(a+I*b)/(I*b-a)*arctan(1/b*a)+2*I/(I*a+b)^2/f/(b-I*a)*b^2*a*c^2/(a+I*b)/(I*b-a)*arctan(1/2...
 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1576 vs. $2(534) = 1068$.

Time = 0.12 (sec), antiderivative size = 1576, normalized size of antiderivative = 2.41

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

$$\begin{aligned} & \frac{1}{6} * (2 * (a^3 - a * b^2) * d^2 * f^3 * x^3 - 6 * b^3 * c^2 * f^2 - 6 * (b^3 * d^2 * f^2 - (a^3 - \\ & a * b^2) * c * d * f^3) * x^2 - 6 * (2 * b^3 * c * d * f^2 - (a^3 - a * b^2) * c^2 * f^3) * x - 3 * (-2 \\ & * I * a^2 * b * d^2 * f * x - 2 * I * a^2 * b * c * d * f - I * a * b^2 * d^2 + (-2 * I * a * b^2 * d^2 * f * x - 2 \\ & * I * a * b^2 * c * d * f - I * b^3 * d^2) * \tan(f * x + e)) * \operatorname{dilog}(2 * ((I * a * b - b^2) * \tan(f * x + \\ & e)^2 - a^2 - I * a * b + (I * a^2 - 2 * a * b - I * b^2) * \tan(f * x + e)) / ((a^2 + b^2) * t \\ & an(f * x + e)^2 + a^2 + b^2) + 1) - 3 * (2 * I * a^2 * b * d^2 * f * x + 2 * I * a^2 * b * c * d * f + \\ & I * a * b^2 * d^2 + (2 * I * a * b^2 * d^2 * f * x + 2 * I * a * b^2 * c * d * f + I * b^3 * d^2) * \tan(f * x + \\ & e)) * \operatorname{dilog}(2 * ((-I * a * b - b^2) * \tan(f * x + e)^2 - a^2 + I * a * b + (-I * a^2 - 2 * a * \\ & b + I * b^2) * \tan(f * x + e)) / ((a^2 + b^2) * \tan(f * x + e)^2 + a^2 + b^2) + 1) + 6 \\ & * (a^2 * b * d^2 * f^2 * x^2 - a^2 * b * d^2 * e^2 + 2 * a^2 * b * c * d * e * f + a * b^2 * d^2 * e^2 + (2 * a \\ & ^2 * b * c * d * f^2 + a * b^2 * d^2 * f) * x + (a * b^2 * d^2 * f^2 * x^2 - a * b^2 * d^2 * e^2 + 2 * a * b \\ & ^2 * c * d * e * f + b^3 * d^2 * e^2 + (2 * a * b^2 * c * d * f^2 + b^3 * d^2 * f) * x) * \tan(f * x + e)) * \log(-2 * ((I * a * b - b^2) * \tan(f * x + e)^2 - a^2 - I * a * b + (I * a^2 - 2 * a * b - I * b^2) \\ & * \tan(f * x + e)) / ((a^2 + b^2) * \tan(f * x + e)^2 + a^2 + b^2)) + 6 * (a^2 * b * d^2 * f^2 * \\ & 2 * x^2 - a^2 * b * d^2 * e^2 + 2 * a^2 * b * c * d * e * f + a * b^2 * d^2 * e^2 + (2 * a^2 * b * c * d * f^2 + \\ & a * b^2 * d^2 * f) * x + (a * b^2 * d^2 * f^2 * x^2 - a * b^2 * d^2 * e^2 + 2 * a * b^2 * c * d * e * f + b \\ & ^3 * d^2 * e^2 + (2 * a * b^2 * c * d * f^2 + b^3 * d^2 * f) * x) * \tan(f * x + e)) * \log(-2 * ((-I * a * b \\ & - b^2) * \tan(f * x + e)^2 - a^2 + I * a * b + (-I * a^2 - 2 * a * b + I * b^2) * \tan(f * x + e)) / ((a^2 + b^2) * \tan(f * x + e)^2 + a^2 + b^2)) + 6 * (a^2 * b * d^2 * e^2 + a^2 * b * c * \\ & 2 * f^2 - a * b^2 * d^2 * e^2 - (2 * a^2 * b * c * d * e - a * b^2 * c * d) * f + (a * b^2 * d^2 * e^2 + \dots\end{aligned}$$

Sympy [F]

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx$$

input

```
integrate((d*x+c)**2/(a+b*tan(f*x+e))**2,x)
```

output

```
Integral((c + d*x)**2/(a + b*tan(e + f*x))**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2530 vs. $2(534) = 1068$.

Time = 0.92 (sec), antiderivative size = 2530, normalized size of antiderivative = 3.87

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output

```
-1/3*(6*c*d*e*(2*a*b*log(b*tan(f*x + e) + a)/((a^4 + 2*a^2*b^2 + b^4)*f) -
 a*b*log(tan(f*x + e)^2 + 1)/((a^4 + 2*a^2*b^2 + b^4)*f) - b/((a^2*b + b^3)
 )*f*tan(f*x + e) + (a^3 + a*b^2)*f) + (a^2 - b^2)*(f*x + e)/((a^4 + 2*a^2*
 b^2 + b^4)*f)) - 3*(2*a*b*log(b*tan(f*x + e) + a)/(a^4 + 2*a^2*b^2 + b^4) -
 a*b*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + (a^2 - b^2)*(f*x +
 e)/(a^4 + 2*a^2*b^2 + b^4) - b/(a^3 + a*b^2 + (a^2*b + b^3)*tan(f*x + e))
 )*c^2 - ((a^3 - I*a^2*b + a*b^2 - I*b^3)*(f*x + e)^3*d^2 + 3*(a^3 - I*a^2*
 b + a*b^2 - I*b^3)*(f*x + e)*d^2*e^2 - 6*(-I*a*b^2 + b^3)*d^2*e^2 - 3*((a^
 3 - I*a^2*b + a*b^2 - I*b^3)*d^2*e - (a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d*f
 )*(f*x + e)^2 - 6*((-I*a^2*b + a*b^2)*d^2*e^2 + (I*a*b^2 - b^3)*d^2*e + (-
 I*a*b^2 + b^3)*c*d*f + ((-I*a^2*b - a*b^2)*d^2*e^2 + (I*a*b^2 + b^3)*d^2*e
 + (-I*a*b^2 - b^3)*c*d*f)*cos(2*f*x + 2*e) + ((a^2*b - I*a*b^2)*d^2*e^2 -
 (a*b^2 - I*b^3)*d^2*e + (a*b^2 - I*b^3)*c*d*f)*sin(2*f*x + 2*e))*arctan2(
 -b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2
 *f*x + 2*e) + a) - 6*((I*a^2*b - a*b^2)*(f*x + e)^2*d^2 + (2*(-I*a^2*b + a
 *b^2)*d^2*e + 2*(I*a^2*b - a*b^2)*c*d*f + (I*a*b^2 - b^3)*d^2)*(f*x + e) +
 ((I*a^2*b + a*b^2)*(f*x + e)^2*d^2 + (2*(-I*a^2*b - a*b^2)*d^2*e + 2*(I*a
 ^2*b + a*b^2)*c*d*f + (I*a*b^2 + b^3)*d^2)*(f*x + e))*cos(2*f*x + 2*e) - (
 (a^2*b - I*a*b^2)*(f*x + e)^2*d^2 - (2*(a^2*b - I*a*b^2)*d^2*e - 2*(a^2*b
 - I*a*b^2)*c*d*f - (a*b^2 - I*b^3)*d^2)*(f*x + e))*sin(2*f*x + 2*e))*ar...
```

Giac [F]

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(dx + c)^2}{(b \tan(fx + e) + a)^2} dx$$

input `integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((d*x + c)^2/(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx$$

input `int((c + d*x)^2/(a + b*tan(e + f*x))^2,x)`

output `int((c + d*x)^2/(a + b*tan(e + f*x))^2, x)`

Reduce [F]

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `int((d*x+c)^2/(a+b*tan(f*x+e))^2,x)`

output

```
(int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*d**2*f + 2*int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**3*b**3*d**2*f + int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**5*d**2*f + int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**6*d**2*f + 2*int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**4*b**2*d**2*f + int(x**2/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*b**4*d**2*f + 2*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*c*d*f + 4*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**3*b**3*c*d*f + 2*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**5*c*d*f + 2*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**6*c*d*f + 4*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**4*b**2*c*d*f + 2*int(x/(tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*b**4*c*d*f - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**2*c**2 - log(tan(e + f*x)**2 + 1)*a**3*b*c**2 + 2*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b**2*c**2 + 2*log(tan(e + f*x)*b + a)*a**3*b*c**2 + tan(e + f*x)*a**3*b*c**2*f*x + tan(e + f*x)*a**2*b**2*c**2 - tan(e + f*x)*a*b**3*c**2*f*x + tan(e + f*x)*b**4*c**2 + a**4*c**2*f*x - a**2*b**2*c**2*f*x)/(a*f*(tan(e + f*x)*a**4*b + 2*tan(e + f*x)*a**2*b**3 + tan(e + f*x)*b**5 + a**5 + 2*a**3*b**2 + a*b**...)
```

3.61 $\int \frac{c+dx}{(a+b\tan(e+fx))^2} dx$

Optimal result	481
Mathematica [B] (warning: unable to verify)	482
Rubi [A] (verified)	483
Maple [B] (verified)	486
Fricas [B] (verification not implemented)	487
Sympy [F]	488
Maxima [B] (verification not implemented)	488
Giac [F]	489
Mupad [F(-1)]	490
Reduce [F]	490

Optimal result

Integrand size = 18, antiderivative size = 214

$$\begin{aligned} \int \frac{c+dx}{(a+b\tan(e+fx))^2} dx = & -\frac{(c+dx)^2}{2(a^2+b^2)d} + \frac{(bd+2acf+2adfx)^2}{4a(a+ib)(a^2+b^2)df^2} \\ & + \frac{b(bd+2acf+2adfx)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\ & - \frac{iabd\text{PolyLog}\left(2,-\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\ & - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))} \end{aligned}$$

output

```
-1/2*(d*x+c)^2/(a^2+b^2)/d+1/4*(2*a*d*f*x+2*a*c*f+b*d)^2/a/(a+I*b)/(a^2+b^2)/d/f^2+b*(2*a*d*f*x+2*a*c*f+b*d)*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)^2/f^2-I*a*b*d*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)^2/f^2-b*(d*x+c)/(a^2+b^2)/f/(a+b*tan(f*x+e))
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 745 vs. $2(214) = 428$.

Time = 7.47 (sec), antiderivative size = 745, normalized size of antiderivative = 3.48

$$\begin{aligned}
 & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\
 &= \frac{(e + fx)(-2de + 2cf + d(e + fx)) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{2(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2} \\
 &+ \frac{b^2d(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{a(a - ib)(a + ib)(a^2 + b^2)f^2(a + b \tan(e + fx))^2} \\
 &- \frac{2bde(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{(a - ib)(a + ib)(a^2 + b^2)f^2(a + b \tan(e + fx))^2} \\
 &+ \frac{2bc(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{(a - ib)(a + ib)(a^2 + b^2)f(a + b \tan(e + fx))^2} \\
 &- \frac{d \left(e^{i \arctan(\frac{a}{b})}(e + fx)^2 + \frac{a(i(e + fx)(-\pi + 2 \arctan(\frac{a}{b})) - \pi \log(1 + e^{-2i(e + fx)}) - 2(e + fx + \arctan(\frac{a}{b})) \log(1 - e^{2i(e + fx + \arctan(\frac{a}{b}))})}{\sqrt{1 - e^{4i(e + fx + \arctan(\frac{a}{b}))}}} \right)}{(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2} \\
 &+ \frac{\sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))(-b^2de \sin(e + fx) + b^2cf \sin(e + fx) + b^2d(e + fx) \sin(e + fx))}{a(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2}
 \end{aligned}$$

input `Integrate[(c + d*x)/(a + b*Tan[e + f*x])^2, x]`

output

$$\begin{aligned} & ((e + f*x)*(-2*d*e + 2*c*f + d*(e + f*x))*\text{Sec}[e + f*x]^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])^2)/(2*(a - I*b)*(a + I*b)*f^2*(a + b*\text{Tan}[e + f*x])^2) + (b^2*d*(-(b*(e + f*x)) + a*\text{Log}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x]])*\text{Sec}[e + f*x]^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])^2)/(a*(a - I*b)*(a + I*b)*(a^2 + b^2)*f^2*(a + b*\text{Tan}[e + f*x])^2) - (2*b*d*e*(-(b*(e + f*x)) + a*\text{Log}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x]])*\text{Sec}[e + f*x]^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])^2)/((a - I*b)*(a + I*b)*(a^2 + b^2)*f^2*(a + b*\text{Tan}[e + f*x])^2) + (2*b*c*(-(b*(e + f*x)) + a*\text{Log}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x]])*\text{Sec}[e + f*x]^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])^2)/((a - I*b)*(a + I*b)*(a^2 + b^2)*f^2*(a + b*\text{Tan}[e + f*x])^2) - (d*(E^(I*\text{ArcTan}[a/b]))*(e + f*x)^2 + (a*(I*(e + f*x)*(-Pi + 2*\text{ArcTan}[a/b])) - Pi*\text{Log}[1 + E^((-2*I)*(e + f*x))]) - 2*(e + f*x + A \text{rcTan}[a/b])*Log[1 - E^((2*I)*(e + f*x + \text{ArcTan}[a/b]))] + Pi*\text{Log}[\text{Cos}[e + f*x]] + 2*\text{ArcTan}[a/b]*\text{Log}[\text{Sin}[e + f*x + \text{ArcTan}[a/b]]] + I*\text{PolyLog}[2, E^((2*I)*(e + f*x + \text{ArcTan}[a/b]))]))/(Sqrt[1 + a^2/b^2]*b))*\text{Sec}[e + f*x]^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])^2)/((a - I*b)*(a + I*b)*Sqrt[(a^2 + b^2)/b^2]*f^2*(a + b*\text{Tan}[e + f*x])^2) + (\text{Sec}[e + f*x]^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])*(-(b^2*d*e*\text{Sin}[e + f*x]) + b^2*c*f*\text{Sin}[e + f*x] + b^2*d*(e + f*x)*\text{Si} n[e + f*x]))/(a*(a - I*b)*(a + I*b)*f^2*(a + b*\text{Tan}[e + f*x])^2) \end{aligned}$$

Rubi [A] (verified)

Time = 0.78 (sec), antiderivative size = 228, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.389, Rules used = {3042, 4216, 3042, 4215, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\ & \quad \downarrow \textcolor{blue}{3042} \\ & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\ & \quad \downarrow \textcolor{blue}{4216} \\ & \frac{\int \frac{bd+2afxd+2acf}{a+b\tan(e+fx)} dx}{f(a^2+b^2)} - \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\ & \quad \downarrow \textcolor{blue}{3042} \end{aligned}$$

$$\begin{aligned}
& \frac{\int \frac{bd+2acf+2acf}{a+b\tan(e+fx)} dx}{f(a^2+b^2)} - \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\
& \quad \downarrow \text{4215} \\
& \frac{2ib \int \frac{e^{2i(e+fx)}(bd+2acf+2acf)}{(a+ib)^2+(a^2+b^2)e^{2i(e+fx)}} dx + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \\
& \quad \frac{\frac{(c+dx)^2}{2d(a^2+b^2)}}{2d(a^2+b^2)} \\
& \quad \downarrow \text{2620} \\
& \frac{2ib \left(\frac{iad \int \log \left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) dx}{a^2+b^2} - \frac{i(2acf+2adf+bd) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \\
& \quad \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\
& \quad \downarrow \text{2715} \\
& \frac{2ib \left(\frac{ad \int e^{-2i(e+fx)} \log \left(\frac{e^{2i(e+fx)}(a^2+b^2)}{(a+ib)^2} + 1 \right) de^{2i(e+fx)}}{2f(a^2+b^2)} - \frac{i(2acf+2adf+bd) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \\
& \quad \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)} \\
& \quad \downarrow \text{2838} \\
& \frac{2ib \left(-\frac{i(2acf+2adf+bd) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} - \frac{ad \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2f(a^2+b^2)} \right) + \frac{(2acf+2adf+bd)^2}{4adf(a+ib)}}{f(a^2+b^2)} - \\
& \quad \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} - \frac{(c+dx)^2}{2d(a^2+b^2)}
\end{aligned}$$

input $\operatorname{Int}[(c + d*x)/(a + b*\operatorname{Tan}[e + f*x])^2, x]$

output

```

-1/2*(c + d*x)^2/((a^2 + b^2)*d) + ((b*d + 2*a*c*f + 2*a*d*f*x)^2/(4*a*(a
+ I*b)*d*f) + (2*I)*b*(((-1/2*I)*(b*d + 2*a*c*f + 2*a*d*f*x)*Log[1 + ((a^2
+ b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) - (a*d*PolyLog[
2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/(2*(a^2 + b^2)*f)))/
((a^2 + b^2)*f) - (b*(c + d*x))/((a^2 + b^2)*f*(a + b*Tan[e + f*x]))

```

Definitions of rubi rules used

rule 2620

```

Int[((F_ )^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^((m_.))/
((a_) + (b_.)*((F_ )^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

rule 2715

```

Int[Log[(a_) + (b_.)*((F_ )^((e_.)*(c_.) + (d_.)*(x_.)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x
))`n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

rule 2838

```

Int[Log[(c_.)*(d_) + (e_.)*(x_.)^((n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

rule 3042

```

Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 4215

```

Int[((c_.) + (d_.)*(x_.))^((m_.))/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Sy
mbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Simp[2*I*b In
t[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2
*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2
, 0] && IGtQ[m, 0]

```

rule 4216

```
Int[((c_) + (d_)*(x_))/((a_) + (b_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol]
] :> Simp[-(c + d*x)^2/(2*d*(a^2 + b^2)), x] + (Simp[1/(f*(a^2 + b^2)) Int[(b*d + 2*a*c*f + 2*a*d*f*x)/(a + b*Tan[e + f*x]), x], x] - Simp[b*((c + d*x)/(f*(a^2 + b^2)*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2000 vs. $2(202) = 404$.

Time = 0.70 (sec) , antiderivative size = 2001, normalized size of antiderivative = 9.35

method	result	size
risch	Expression too large to display	2001

input `int((d*x+c)/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output

```
-1/2*I/(I*a+b)^2/f^2/(b-I*a)*b^2*d/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))
+exp(4*I*(f*x+e))*b^2+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^
2)*a-I/(I*a+b)^2/f/(b-I*a)*b*a^2*c/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))
+exp(4*I*(f*x+e))*b^2+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^
2)+2*I/(I*a+b)^2/f/(b-I*a)*b/(a+I*b)*a*d*ln(1-(I*b-a)*exp(2*I*(f*x+e))/(a+
I*b))*x+4*I/(I*a+b)^2/f^2/(b-I*a)*b*e*a*d/(a+I*b)*ln(exp(I*(f*x+e)))+2*I/(I
*a+b)^2/f^2/(b-I*a)*b/(a+I*b)*a*d*ln(1-(I*b-a)*exp(2*I*(f*x+e))/(a+I*b))*e+2*I/(I*a+b)^2/f/(b-I*a)*b^2*a*c/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e))-1/2/a*b+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)-2*I/(I*a+b)^2/f/(b-I*a)*b^2*a*c/(a+I*b)/(I*b-a)*arctan(1/b*a)+1/(I*a+b)^2/f^2/(b-I*a)*b^2*e*a*d/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))+exp(4*I*(f*x+e))*b^2+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^2)+2/(I*a+b)^2/f^2/(b-I*a)*b*e*a^2*d/(a+I*b)/(I*b-a)*arctan(1/b*a)-1/2/(2*I*a*b-a^2+b^2)*d*x^2-1/(2*I*a*b-a^2+b^2)*c*x+2/(I*a+b)^2/(b-I*a)*b/(a+I*b)*a*d*x^2-2*I*b^2*(d*x+c)/(b-I*a)/f/(I*a+b)^2/(b*exp(2*I*(f*x+e))+I*a*exp(2*I*(f*x+e))-b+I*a)+4/(I*a+b)^2/f/(b-I*a)*b/(a+I*b)*a*d*e*x-1/(I*a+b)^2/f/(b-I*a)*b^2*a*c/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))+exp(4*I*(f*x+e))*b^2+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^2)-2/(I*a+b)^2/f/(b-I*a)*b*a^2*c/(a+I*b)/(I*b-a)*arctan(1/2/a*b*exp(2*I*(f*x+e)...)
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 843 vs. $2(197) = 394$.

Time = 0.12 (sec) , antiderivative size = 843, normalized size of antiderivative = 3.94

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output

```
1/2*((a^3 - a*b^2)*d*f^2*x^2 - 2*b^3*c*f - 2*(b^3*d*f - (a^3 - a*b^2)*c*f^2)*x + (I*a*b^2*d*tan(f*x + e) + I*a^2*b*d)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + (-I*a*b^2*d*tan(f*x + e) - I*a^2*b*d)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(a^2*b*d*f*x + a^2*b*d*e + (a*b^2*d*f*x + a*b^2*d*e)*tan(f*x + e))*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(a^2*b*d*f*x + a^2*b*d*e + (a*b^2*d*f*x + a*b^2*d*e)*tan(f*x + e))*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - (2*a^2*b*d*e - 2*a^2*b*c*f - a*b^2*d + (2*a*b^2*d*e - 2*a*b^2*c*f - b^3*d)*tan(f*x + e))*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - (2*a^2*b*d*e - 2*a^2*b*c*f - a*b^2*d + (2*a*b^2*d*e - 2*a*b^2*c*f - b^3*d)*tan(f*x + e))*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + ((a^2*b - b^3)*d*f^2*x^2 + 2*a*b^2*c*f + 2*(a*b^2*d*f + (a^2*b - b^3)*c*f^2*x)*tan(f*x + e))/((a^4*b + 2*a^2*b^3 + b^5)*f^2*tan(f*x + e) + (a^5 + 2*a^3*b^2 + a*b^4)*f^2)
```

Sympy [F]

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))**2,x)`

output `Integral((c + d*x)/(a + b*tan(e + f*x))**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1156 vs. $2(197) = 394$.

Time = 0.53 (sec) , antiderivative size = 1156, normalized size of antiderivative = 5.40

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

input `integrate((d*x+c)/(a+b*tan(f*x+e))**2,x, algorithm="maxima")`

output

```
1/2*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d*f^2*x^2 + 2*(a^3 - I*a^2*b + a*b^2 - I*b^3)*c*f^2*x - 4*(-I*a*b^2 + b^3)*c*f - 2*(2*(-I*a^2*b + a*b^2)*c*f + (-I*a*b^2 + b^3)*d + (2*(-I*a^2*b - a*b^2)*c*f + (-I*a*b^2 - b^3)*d)*cos(2*f*x + 2*e) + (2*(a^2*b - I*a*b^2)*c*f + (a*b^2 - I*b^3)*d)*sin(2*f*x + 2*e))*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) - 4*((I*a^2*b + a*b^2)*d*f*x*cos(2*f*x + 2*e) - (a^2*b - I*a*b^2)*d*f*x*sin(2*f*x + 2*e) + (I*a^2*b - a*b^2)*d*f*x)*arc tan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) + ((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d*f^2*x^2 + 2*((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*c*f^2 - 2*(I*a*b^2 + b^3)*d*f)*x)*cos(2*f*x + 2*e) - 2*((I*a^2*b + a*b^2)*d*cos(2*f*x + 2*e) - (a^2*b - I*a*b^2)*d*sin(2*f*x + 2*e) + (I*a^2*b - a*b^2)*d)*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) + (2*(a^2*b + I*a*b^2)*c*f + (a*b^2 + I*b^3)*d + (2*(a^2*b - I*a*b^2)*c*f + (a*b^2 - I*b^3)*d)*cos(2*f*x + 2*e) - (2*(-I*a^2*b - a*b^2)*c*f - (I*a*b^2 + b^3)*d)*sin(2*f*x + 2*e))*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 2*((a^2*b - I*a*b^2)*d*f*x*cos(2*f*x + 2*e) - (-I*a^2*b - a*b^2)*d*f*x*sin(2*f*x + 2*e) + (a^2*b + I*a*b^2)*d*f*x)*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*s...)
```

Giac [F]

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{dx + c}{(b \tan(fx + e) + a)^2} dx$$

input

```
integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="giac")
```

output

```
integrate((d*x + c)/(b*tan(f*x + e) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

input `int((c + d*x)/(a + b*tan(e + f*x))^2,x)`

output `int((c + d*x)/(a + b*tan(e + f*x))^2, x)`

Reduce [F]

$$\begin{aligned} & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\ &= \left(\int \frac{x}{\tan(fx+e)^2 b^2 + 2 \tan(fx+e) ab + a^2} dx \right) \tan(fx+e) a^5 b d f + 2 \left(\int \frac{x}{\tan(fx+e)^2 b^2 + 2 \tan(fx+e) ab + a^2} dx \right) \tan(fx+e) a^5 b d f \end{aligned}$$

input `int((d*x+c)/(a+b*tan(f*x+e))^2,x)`

output `(int(x/(\tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**5*b*d*f + 2*int(x/(\tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a**3*b**3*d*f + int(x/(\tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*tan(e + f*x)*a*b**5*d*f + int(x/(\tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**6*d*f + 2*int(x/(\tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**4*b**2*d*f + int(x/(\tan(e + f*x)**2*b**2 + 2*tan(e + f*x)*a*b + a**2),x)*a**2*b**4*d*f - log(tan(e + f*x)**2 + 1)*tan(e + f*x)*a**2*b**2*c - log(tan(e + f*x)**2 + 1)*a**3*b*c + 2*log(tan(e + f*x)*b + a)*tan(e + f*x)*a**2*b**2*c + 2*log(tan(e + f*x)*b + a)*a**3*b*c + tan(e + f*x)*a**3*b*c*f*x + tan(e + f*x)*a**2*b**2*c - tan(e + f*x)*a*b**3*c*f*x + tan(e + f*x)*b**4*c + a**4*c*f*x - a**2*b**2*c*f*x)/(a*f*(tan(e + f*x)*a**4*b + 2*tan(e + f*x)*a**2*b**3 + tan(e + f*x)*b**5 + a**5 + 2*a**3*b**2 + a*b**4))`

3.62 $\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$

Optimal result	491
Mathematica [N/A]	491
Rubi [N/A]	492
Maple [N/A]	493
Fricas [N/A]	493
Sympy [N/A]	493
Maxima [N/A]	494
Giac [N/A]	495
Mupad [N/A]	495
Reduce [N/A]	495

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b\tan(e+fx))^2}, x\right)$$

output `Defer(Int)(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`

Mathematica [N/A]

Not integrable

Time = 12.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$$

input `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])^2),x]`

output `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])^2), x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx \end{aligned}$$

input `Int[1/((c + d*x)*(a + b*Tan[e + f*x])^2),x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tan(fx + e))^2} dx$$

input `int(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`

output `int(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)^2} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output `integral(1/(a^2*d*x + a^2*c + (b^2*d*x + b^2*c)*tan(f*x + e)^2 + 2*(a*b*d*x + a*b*c)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.62 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx = \int \frac{1}{(a + b \tan(e + fx))^2 (c + dx)} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))**2,x)`

output $\text{Integral}(1/((a + b\tan(e + fx))^2*(c + dx)), x)$

Maxima [N/A]

Not integrable

Time = 10.95 (sec) , antiderivative size = 1498, normalized size of antiderivative = 74.90

$$\int \frac{1}{(c + dx)(a + b\tan(e + fx))^2} dx = \int \frac{1}{(dx + c)(b\tan(fx + e) + a)^2} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`

output
$$\begin{aligned} & (((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*\cos(2*f*x + 2*e)^2*\log(d*x + c) + (a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*\log(d*x + c)*\sin(2*f*x + 2*e)^2 - 2*(2*a*b^3*d - ((a^4 - 2*a^2*b^2 + b^4)*d*f*x + (a^4 - 2*a^2*b^2 + b^4)*c*f)*\log(d*x + c))*\cos(2*f*x + 2*e) + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*\cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*\sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d*f)*\cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c*d*f)*\sin(2*f*x + 2*e))*\text{integrate}(2*(2*(2*(2*a^2*b^2*d*f*x + 2*a^2*b^2*c*f - a*b^3*d)*cos(2*f*x + 2*e) - (2*(a^3*b - a*b^3)*d*f*x + 2*(a^3*b - a*b^3)*c*f - (a^2*b^2 - b^4)*d)*sin(2*f*x + 2*e)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^2*f*x^2 + 2*(a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f))$$

Giac [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)^2} dx$$

input `integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate(1/((d*x + c)*(b*tan(f*x + e) + a)^2), x)`

Mupad [N/A]

Not integrable

Time = 9.80 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx = \int \frac{1}{(a + b \tan(e + fx))^2 (c + dx)} dx$$

input `int(1/((a + b*tan(e + f*x))^2*(c + d*x)),x)`

output `int(1/((a + b*tan(e + f*x))^2*(c + d*x)), x)`

Reduce [N/A]

Not integrable

Time = 0.16 (sec) , antiderivative size = 66, normalized size of antiderivative = 3.30

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx \\ &= \int \frac{1}{\tan(fx + e)^2 b^2 c + \tan(fx + e)^2 b^2 dx + 2 \tan(fx + e) abc + 2 \tan(fx + e) ab dx + a^2 c + a^2 dx} dx \end{aligned}$$

input `int(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)`

output `int(1/(tan(e + f*x)**2*b**2*c + tan(e + f*x)**2*b**2*d*x + 2*tan(e + f*x)*a*b*c + 2*tan(e + f*x)*a*b*d*x + a**2*c + a**2*d*x),x)`

3.63 $\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$

Optimal result	497
Mathematica [N/A]	497
Rubi [N/A]	498
Maple [N/A]	499
Fricas [N/A]	499
Sympy [N/A]	499
Maxima [N/A]	500
Giac [N/A]	501
Mupad [N/A]	501
Reduce [N/A]	501

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b\tan(e+fx))^2}, x\right)$$

output `Defer(Int)(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`

Mathematica [N/A]

Not integrable

Time = 13.83 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$$

input `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2),x]`

output `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2), x]`

Rubi [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {3042, 4223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx \\ & \quad \downarrow \text{4223} \\ & \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx \end{aligned}$$

input `Int[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2), x]`

output `$Aborted`

Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4223 `Int[((c_.) + (d_.)*(x_.))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Unintegrable[(c + d*x)^m*(a + b*Tan[e + f*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x]`

Maple [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2 (a + b \tan(fx + e))^2} dx$$

input `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`

output `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 96, normalized size of antiderivative = 4.80

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)^2} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

output `integral(1/(a^2*d^2*x^2 + 2*a^2*c*d*x + a^2*c^2 + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*tan(f*x + e))^2 + 2*(a*b*d^2*x^2 + 2*a*b*c*d*x + a*b*c^2)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 3.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(a + b \tan(e + fx))^2 (c + dx)^2} dx$$

input `integrate(1/(d*x+c)**2/(a+b*tan(f*x+e))**2,x)`

output Integral(1/((a + b*tan(e + f*x))**2*(c + d*x)**2), x)

Maxima [N/A]

Not integrable

Time = 28.36 (sec) , antiderivative size = 1977, normalized size of antiderivative = 98.85

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)^2} dx$$

```
input integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

```

output -((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*cos(2*f*x + 2*e)^2 + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*sin(2*f*x + 2*e)^2 + 2*(2*a*b^3*d + (a^4 - 2*a^2*b^2 + b^4)*d*f*x + (a^4 - 2*a^2*b^2 + b^4)*c*f)*cos(2*f*x + 2*e) - ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^3*f*x^2 + 2*(a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c^2*d*f)*cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^3*f*x^2 + 2*(a^5*b + 2*a^3*b^3 + a*b^5)*c*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c^2*d*f)*sin(2*f*x + 2*e))*integrate(4*(2*(a^2*b^2*d*f*x + a^2*b^2*c*f - a*b^3*d)*cos(2*f*x + 2*e) - ((a^3*b - a*b^3)*d*f*x + (a^3*b - a*b^3)*c*f - (a^2*b^2 - b^4)*d)*sin(2*f*x + 2*e))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^3*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x^2 + 3*...
```

Giac [N/A]

Not integrable

Time = 22.71 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)^2} dx$$

input `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate(1/((d*x + c)^2*(b*tan(f*x + e) + a)^2), x)`

Mupad [N/A]

Not integrable

Time = 9.50 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(a + b \tan(e + fx))^2(c + dx)^2} dx$$

input `int(1/((a + b*tan(e + f*x))^2*(c + d*x)^2),x)`

output `int(1/((a + b*tan(e + f*x))^2*(c + d*x)^2), x)`

Reduce [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 121, normalized size of antiderivative = 6.05

$$\begin{aligned} & \int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx \\ &= \int \frac{1}{\tan(fx + e)^2 b^2 c^2 + 2 \tan(fx + e)^2 b^2 c dx + \tan(fx + e)^2 b^2 d^2 x^2 + 2 \tan(fx + e) ab c^2 + 4 \tan(fx + e) ab c dx + 1} dx \end{aligned}$$

input `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)`

output `int(1/(tan(e + f*x)**2*b**2*c**2 + 2*tan(e + f*x)**2*b**2*c*d*x + tan(e + f*x)**2*b**2*d**2*x**2 + 2*tan(e + f*x)*a*b*c**2 + 4*tan(e + f*x)*a*b*c*d*x + 2*tan(e + f*x)*a*b*d**2*x**2 + a**2*c**2 + 2*a**2*c*d*x + a**2*d**2*x**2),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	503
4.2 Links to plain text integration problems used in this report for each CAS .	521

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```
(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*) (*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","");
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
    ,(*ELSE*) (*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "}
      ,
      finalresult={"C","Result contains higher order function than in optimal. Order "}
    ]
  ]
]
]
```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```



```

ExpnType[expn_] :=
If[AtomQ[expn],
  1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn] === Power,
      If[IntegerQ[expn[[2]]],
        ExpnType[expn[[1]]],
        If[Head[expn[[2]]] === Rational,
          If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
            1,
            Max[ExpnType[expn[[1]]], 2]],
          Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
      If[Head[expn] === Plus || Head[expn] === Times,
        Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
        If[ElementaryFunctionQ[Head[expn]],
          Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], ExpnType[expn[[3]]]]]]]]]]]

```

```
Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],  
If[AppellFunctionQ[Head[expn]],  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],  
If[Head[expn]==RootSum,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],  
If[Head[expn]==Integrate || Head[expn]==Int,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],  
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=  
MemberQ[{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
} , func]
```

```
SpecialFunctionQ[func_] :=  
MemberQ[{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  

```

```
HypergeometricFunctionQ[func_] :=  
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=  
MemberQ[{AppellF1}, func]
```

Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
```

```
#      antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A", " ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
            # this assumes optimal do not as well. No check is needed here.
            if debug then
                print("result do not contain complex, this assumes optimal do not as well");
            fi;
```

```
if leaf_count_result<=2*leaf_count_optimal then
    if debug then
        print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A"," ";
else
    if debug then
        print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B",cat("Leaf count of result is larger than twice the leaf count of
                    convert(leaf_count_result,string),\"$ vs. \$2(",convert(leaf_count_optimal,string),")=",convert(2*leaf_count_optimal,string));
fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),"."));
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'`^`') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+`') or type(expn,'`*`') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc

```

```
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arccsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]
```

```
def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*')
```

```

m1 = expnType(expn.args[0])
m2 = expnType(list(expn.args[1:]))
return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sagemode")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(grade)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```
from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                       ]
    if debug:
```

```

if m:
    print ("func ", func , " is elementary_function")
else:
    print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral'
'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
'elliptic_pi','exp_integral_e','log_integral',
'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
'weierstrassPPrime','weierstrassSigma']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):
    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0], Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1], Integer)

```

```

    return expnType(expn.operands()[0])  #expnType(expn.args[0])
elif type(expn.operands()[1]) == Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0]) == Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))  #max(3,expnType(expn.args[0]),expnType(expn.args[1]))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands()))  #max(map(expnType, list(expn.args)))
    return max(4,m1)  #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands()))  #max(map(expnType, list(expn.args)))
    return max(5,m1)  #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands()))  #max(map(expnType, list(expn.args)))
    return max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))  #max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

```

```
leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation ="none"
            else:
                grade = "B"
                grade_annotation ="Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation ="Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation ="none"
        else:
            grade = "B"
            grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(expnType_result)+"/"+str(expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file