

# Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.4-Cotangent/224-4.4.0

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 9:07pm

# Contents

<b>1</b>	<b>Introduction</b>	<b>4</b>
1.1	Listing of CAS systems tested . . . . .	5
1.2	Results . . . . .	6
1.3	Time and leaf size Performance . . . . .	10
1.4	Performance based on number of rules Rubi used . . . . .	12
1.5	Performance based on number of steps Rubi used . . . . .	13
1.6	Solved integrals histogram based on leaf size of result . . . . .	14
1.7	Solved integrals histogram based on CPU time used . . . . .	15
1.8	Leaf size vs. CPU time used . . . . .	16
1.9	list of integrals with no known antiderivative . . . . .	17
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	17
1.11	list of integrals solved by CAS but failed verification . . . . .	17
1.12	Timing . . . . .	18
1.13	Verification . . . . .	18
1.14	Important notes about some of the results . . . . .	19
1.15	Current tree layout of integration tests . . . . .	22
1.16	Design of the test system . . . . .	23
<b>2</b>	<b>detailed summary tables of results</b>	<b>24</b>
2.1	List of integrals sorted by grade for each CAS . . . . .	25
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	29
2.3	Detailed conclusion table specific for Rubi results . . . . .	43
<b>3</b>	<b>Listing of integrals</b>	<b>45</b>
3.1	$\int \cot(a + bx) dx$ . . . . .	47
3.2	$\int \cot^2(a + bx) dx$ . . . . .	52
3.3	$\int \cot^3(a + bx) dx$ . . . . .	57
3.4	$\int \cot^4(a + bx) dx$ . . . . .	63
3.5	$\int \cot^5(a + bx) dx$ . . . . .	69
3.6	$\int \cot^6(a + bx) dx$ . . . . .	76
3.7	$\int \cot^7(a + bx) dx$ . . . . .	82

3.8	$\int \cot^8(a + bx) dx$	90
3.9	$\int (c \cot(a + bx))^{7/2} dx$	97
3.10	$\int (c \cot(a + bx))^{5/2} dx$	108
3.11	$\int (c \cot(a + bx))^{3/2} dx$	118
3.12	$\int \sqrt{c \cot(a + bx)} dx$	128
3.13	$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx$	136
3.14	$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx$	145
3.15	$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx$	155
3.16	$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx$	165
3.17	$\int (c \cot(a + bx))^{4/3} dx$	175
3.18	$\int (c \cot(a + bx))^{2/3} dx$	185
3.19	$\int \sqrt[3]{c \cot(a + bx)} dx$	195
3.20	$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx$	204
3.21	$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx$	213
3.22	$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx$	223
3.23	$\int \cot^n(a + bx) dx$	234
3.24	$\int (b \cot(c + dx))^n dx$	239
3.25	$\int (a \cot^2(x))^{3/2} dx$	244
3.26	$\int \sqrt{a \cot^2(x)} dx$	250
3.27	$\int \frac{1}{\sqrt{a \cot^2(x)}} dx$	255
3.28	$\int \frac{1}{(a \cot^2(x))^{3/2}} dx$	260
3.29	$\int (a \cot^3(x))^{3/2} dx$	266
3.30	$\int \sqrt{a \cot^3(x)} dx$	276
3.31	$\int \frac{1}{\sqrt{a \cot^3(x)}} dx$	286
3.32	$\int \frac{1}{(a \cot^3(x))^{3/2}} dx$	295
3.33	$\int (a \cot^4(x))^{3/2} dx$	305
3.34	$\int \sqrt{a \cot^4(x)} dx$	311
3.35	$\int \frac{1}{\sqrt{a \cot^4(x)}} dx$	316
3.36	$\int \frac{1}{(a \cot^4(x))^{3/2}} dx$	321
3.37	$\int (b \cot^p(c + dx))^n dx$	327
3.38	$\int (a(b \cot(c + dx))^p)^n dx$	332
3.39	$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx$	337
3.40	$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx$	342
3.41	$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx$	347
3.42	$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx$	352
3.43	$\int (d \cot(e + fx))^n \csc^6(e + fx) dx$	358

---

3.44	$\int (d \cot(e + fx))^n \csc^4(e + fx) dx$	364
3.45	$\int (d \cot(e + fx))^n \csc^2(e + fx) dx$	370
3.46	$\int (d \cot(e + fx))^n \sin^2(e + fx) dx$	375
3.47	$\int (d \cot(e + fx))^n \sin^4(e + fx) dx$	380
3.48	$\int (d \cot(e + fx))^n \csc^3(e + fx) dx$	385
3.49	$\int (d \cot(e + fx))^n \csc(e + fx) dx$	390
3.50	$\int (d \cot(e + fx))^n \sin(e + fx) dx$	395
3.51	$\int (d \cot(e + fx))^n \sin^3(e + fx) dx$	400
3.52	$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx$	405
<b>4</b>	<b>Appendix</b>	<b>410</b>
4.1	Listing of Grading functions	410
4.2	Links to plain text integration problems used in this report for each CAS428	

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	5
1.2	Results . . . . .	6
1.3	Time and leaf size Performance . . . . .	10
1.4	Performance based on number of rules Rubi used . . . . .	12
1.5	Performance based on number of steps Rubi used . . . . .	13
1.6	Solved integrals histogram based on leaf size of result . . . . .	14
1.7	Solved integrals histogram based on CPU time used . . . . .	15
1.8	Leaf size vs. CPU time used . . . . .	16
1.9	list of integrals with no known antiderivative . . . . .	17
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	17
1.11	list of integrals solved by CAS but failed verification . . . . .	17
1.12	Timing . . . . .	18
1.13	Verification . . . . .	18
1.14	Important notes about some of the results . . . . .	19
1.15	Current tree layout of integration tests . . . . .	22
1.16	Design of the test system . . . . .	23

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 52 ]. This is test number [ 224 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 52 )	0.00 ( 0 )
Mathematica	100.00 ( 52 )	0.00 ( 0 )
Maple	71.15 ( 37 )	28.85 ( 15 )
Fricas	71.15 ( 37 )	28.85 ( 15 )
Maxima	71.15 ( 37 )	28.85 ( 15 )
Mupad	50.00 ( 26 )	50.00 ( 26 )
Giac	32.69 ( 17 )	67.31 ( 35 )
Reduce	30.77 ( 16 )	69.23 ( 36 )
Sympy	15.38 ( 8 )	84.62 ( 44 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.

Table 1.2: Description of grading applied to integration result


Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.

System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	71.154	0.000	28.846	0.000
Maple	71.154	0.000	0.000	28.846
Maxima	71.154	0.000	0.000	28.846
Giac	25.000	7.692	0.000	67.308
Fricas	13.462	57.692	0.000	28.846
Sympy	9.615	5.769	0.000	84.615
Mupad	0.000	50.000	0.000	50.000
Reduce	0.000	30.769	0.000	69.231


Table 1.3: Antiderivative Grade distribution of each CAS



The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	15	100.00	0.00	0.00
Maple	15	100.00	0.00	0.00
Maxima	15	100.00	0.00	0.00
Mupad	26	0.00	100.00	0.00
Giac	35	94.29	0.00	5.71
Reduce	36	100.00	0.00	0.00
Sympy	44	95.45	4.55	0.00

Table 1.4: Failure statistics for each CAS

### 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.09
Maxima	0.10
Reduce	0.16
Sympy	0.20
Giac	0.22
Rubi	0.34
Mathematica	0.40
Maple	1.03
Mupad	9.37

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Reduce	43.69	1.33	36.00	1.14
Sympy	45.25	1.42	45.00	1.26
Giac	56.53	1.32	46.00	1.11
Maxima	93.62	0.90	84.00	0.84
Rubi	94.52	0.99	72.00	1.00
Maple	98.92	1.00	90.00	0.97
Mathematica	107.87	1.28	67.00	0.94
Mupad	112.04	1.40	77.50	1.01
Fricas	240.05	2.30	168.00	2.17

Table 1.6: Leaf size performance for each CAS

# 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

# 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used



## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22}

Mathematica {39, 48, 50, 51, 52}

Maple {}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Reduce** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

## Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

### Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

# 1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
January 13, 2024  
Design note



# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	25
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	29
2.3	Detailed conclusion table specific for Rubi results . . . . .	43

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	25
Mma . . . . .	25
Maple . . . . .	26
Fricas . . . . .	26
Maxima . . . . .	26
Giac . . . . .	27
Mupad . . . . .	27
Sympy . . . . .	27
Reduce . . . . .	28

### Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

**B grade** { }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### Mma

**A grade** { 1, 3, 5, 7, 9, 10, 11, 12, 13, 14, 15, 16, 19, 20, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 35, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 49 }

**B grade** { }

**C grade** { 2, 4, 6, 8, 17, 18, 21, 22, 33, 34, 39, 48, 50, 51, 52 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maple

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 43, 44, 45 }

**B grade** { }

**C grade** { }

**F normal fail** { 23, 24, 37, 38, 39, 40, 41, 42, 46, 47, 48, 49, 50, 51, 52 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Fricas

**A grade** { 1, 3, 25, 33, 43, 44, 45 }

**B grade** { 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 26, 27, 28, 29, 30, 31, 32, 34, 35, 36 }

**C grade** { }

**F normal fail** { 23, 24, 37, 38, 39, 40, 41, 42, 46, 47, 48, 49, 50, 51, 52 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maxima

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 43, 44, 45 }

**B grade** { }

**C grade** { }

**F normal fail** { 23, 24, 37, 38, 39, 40, 41, 42, 46, 47, 48, 49, 50, 51, 52 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Giac

**A grade** { 1, 3, 5, 7, 19, 20, 25, 26, 27, 28, 33, 34, 45 }

**B grade** { 2, 4, 6, 8 }

**C grade** { }

**F normal fail** { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 21, 22, 23, 24, 29, 30, 31, 32, 37, 38, 39, 40, 41, 42, 43, 44, 46, 47, 48, 49, 50, 51, 52 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { 35, 36 }

## Mupad

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 27, 43, 44, 45 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 23, 24, 25, 26, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 46, 47, 48, 49, 50, 51, 52 }

**F(-2) exception fail** { }

## Sympy

**A grade** { 2, 4, 6, 7, 8 }

**B grade** { 1, 3, 5 }

**C grade** { }

**F normal fail** { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 52 }

**F(-1) timedout fail** { 43, 51 }

**F(-2) exception fail** { }

## Reduce

**A grade** { }

**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 25, 26, 27, 28, 33, 34, 35, 36 }

**C grade** { }

**F normal fail** { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 29, 30, 31, 32, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	13	11	17	11	19	29	12	31	24
N.S.	1	1.18	1.00	1.55	1.00	1.73	2.64	1.09	2.82	2.18
time (sec)	N/A	0.169	0.004	0.082	0.041	0.076	0.092	0.143	0.146	10.171

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	29	18	18	39	17	35	17	15
N.S.	1	1.00	1.93	1.20	1.20	2.60	1.13	2.33	1.13	1.00
time (sec)	N/A	0.172	0.008	0.085	0.109	0.069	0.070	0.213	0.163	0.066

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	30	28	29	23	47	53	27	67	78
N.S.	1	1.07	1.00	1.04	0.82	1.68	1.89	0.96	2.39	2.79
time (sec)	N/A	0.233	0.008	0.145	0.035	0.079	0.235	0.190	0.162	12.322

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	33	29	34	84	27	62	28	23
N.S.	1	1.00	1.22	1.07	1.26	3.11	1.00	2.30	1.04	0.85
time (sec)	N/A	0.234	0.010	0.115	0.117	0.076	0.091	0.159	0.169	0.070

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	44	39	39	38	83	66	38	79	182
N.S.	1	1.05	0.93	0.93	0.90	1.98	1.57	0.90	1.88	4.33
time (sec)	N/A	0.330	0.015	0.164	0.029	0.083	0.322	0.172	0.148	13.761

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	33	39	44	123	39	91	38	36
N.S.	1	1.00	0.73	0.87	0.98	2.73	0.87	2.02	0.84	0.80
time (sec)	N/A	0.303	0.022	0.128	0.111	0.114	0.117	0.146	0.158	9.521

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	60	58	49	48	126	80	49	89	340
N.S.	1	1.03	1.00	0.84	0.83	2.17	1.38	0.84	1.53	5.86
time (sec)	N/A	0.403	0.012	0.187	0.032	0.097	0.471	0.147	0.177	15.646

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	33	49	54	168	51	116	48	43
N.S.	1	1.00	0.58	0.86	0.95	2.95	0.89	2.04	0.84	0.75
time (sec)	N/A	0.376	0.009	0.158	0.122	0.087	0.179	0.159	0.146	9.073

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	177	227	175	169	197	426	0	0	58	91
N.S.	1	1.28	0.99	0.95	1.11	2.41	0.00	0.00	0.33	0.51
time (sec)	N/A	0.573	0.307	0.176	0.114	0.099	0.000	0.000	0.150	9.398

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	197	101	154	185	380	0	0	24	74
N.S.	1	1.25	0.64	0.98	1.18	2.42	0.00	0.00	0.15	0.47
time (sec)	N/A	0.477	0.214	0.145	0.132	0.118	0.000	0.000	0.153	9.030

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	154	201	161	149	179	316	0	0	38	75
N.S.	1	1.31	1.05	0.97	1.16	2.05	0.00	0.00	0.25	0.49
time (sec)	N/A	0.479	0.144	0.137	0.114	0.094	0.000	0.000	0.173	8.681



Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	136	174	71	136	165	286	0	0	12	50
N.S.	1	1.28	0.52	1.00	1.21	2.10	0.00	0.00	0.09	0.37
time (sec)	N/A	0.420	0.067	0.191	0.125	0.110	0.000	0.000	0.157	8.566

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	180	131	138	165	268	0	0	24	57
N.S.	1	1.31	0.96	1.01	1.20	1.96	0.00	0.00	0.18	0.42
time (sec)	N/A	0.395	0.076	0.168	0.121	0.082	0.000	0.000	0.158	9.654

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	197	82	157	187	377	0	0	24	76
N.S.	1	1.25	0.52	1.00	1.19	2.40	0.00	0.00	0.15	0.48
time (sec)	N/A	0.456	0.086	0.141	0.121	0.110	0.000	0.000	0.194	9.907

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	158	205	86	157	188	380	0	0	24	77
N.S.	1	1.30	0.54	0.99	1.19	2.41	0.00	0.00	0.15	0.49
time (sec)	N/A	0.475	0.177	0.140	0.128	0.097	0.000	0.000	0.164	9.081

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	178	225	97	171	205	464	0	0	24	91
N.S.	1	1.26	0.54	0.96	1.15	2.61	0.00	0.00	0.13	0.51
time (sec)	N/A	0.574	0.234	0.141	0.120	0.088	0.000	0.000	0.163	9.344

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	242	217	205	214	196	431	0	0	31	246
N.S.	1	0.90	0.85	0.88	0.81	1.78	0.00	0.00	0.13	1.02
time (sec)	N/A	0.505	0.236	0.201	0.121	0.101	0.000	0.000	0.187	10.290

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	225	196	185	191	182	416	0	0	14	260
N.S.	1	0.87	0.82	0.85	0.81	1.85	0.00	0.00	0.06	1.16
time (sec)	N/A	0.434	0.154	0.240	0.120	0.082	0.000	0.000	0.202	10.530

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	112	106	108	102	211	0	145	14	134
N.S.	1	0.85	0.81	0.82	0.78	1.61	0.00	1.11	0.11	1.02
time (sec)	N/A	0.313	0.214	0.127	0.117	0.084	0.000	0.856	0.159	9.237

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	112	98	108	103	639	0	145	14	128
N.S.	1	0.85	0.75	0.82	0.79	4.88	0.00	1.11	0.11	0.98
time (sec)	N/A	0.319	0.100	0.119	0.119	0.091	0.000	0.356	0.154	9.980

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	225	196	189	200	182	384	0	0	14	231
N.S.	1	0.87	0.84	0.89	0.81	1.71	0.00	0.00	0.06	1.03
time (sec)	N/A	0.453	0.151	0.223	0.113	0.084	0.000	0.000	0.155	10.988

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	244	219	254	212	204	660	0	0	14	277
N.S.	1	0.90	1.04	0.87	0.84	2.70	0.00	0.00	0.06	1.14
time (sec)	N/A	0.497	0.174	0.191	0.116	0.103	0.000	0.000	0.188	9.558

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	46	0	0	0	0	0	10	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.22	0.00
time (sec)	N/A	0.201	0.097	0.000	0.000	0.000	0.000	0.000	0.148	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	52	0	0	0	0	0	14	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.216	0.093	0.000	0.000	0.000	0.000	0.000	0.153	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	27	27	29	30	52	0	31	41	0
N.S.	1	0.75	0.75	0.81	0.83	1.44	0.00	0.86	1.14	0.00
time (sec)	N/A	0.284	0.017	0.145	0.113	0.088	0.000	0.110	0.178	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	22	20	43	0	20	20	0
N.S.	1	1.00	1.00	1.38	1.25	2.69	0.00	1.25	1.25	0.00
time (sec)	N/A	0.229	0.007	0.104	0.120	0.074	0.000	0.115	0.179	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	26	12	45	0	19	34	25
N.S.	1	1.00	1.00	1.53	0.71	2.65	0.00	1.12	2.00	1.47
time (sec)	N/A	0.223	0.009	0.102	0.113	0.080	0.000	0.132	0.153	8.776

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	28	30	36	22	74	0	46	92	0
N.S.	1	0.72	0.77	0.92	0.56	1.90	0.00	1.18	2.36	0.00
time (sec)	N/A	0.271	0.027	0.100	0.115	0.074	0.000	0.136	0.175	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	149	80	189	113	411	0	0	14	0
N.S.	1	0.96	0.52	1.22	0.73	2.65	0.00	0.00	0.09	0.00
time (sec)	N/A	0.531	0.194	0.187	0.121	0.080	0.000	0.000	0.156	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	136	122	166	94	387	0	0	21	0
N.S.	1	1.03	0.92	1.26	0.71	2.93	0.00	0.00	0.16	0.00
time (sec)	N/A	0.485	0.090	0.154	0.121	0.081	0.000	0.000	0.159	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	136	60	164	94	382	0	0	16	0
N.S.	1	1.02	0.45	1.23	0.71	2.87	0.00	0.00	0.12	0.00
time (sec)	N/A	0.481	0.063	0.152	0.137	0.093	0.000	0.000	0.158	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	166	151	71	184	109	495	0	0	16	0
N.S.	1	0.91	0.43	1.11	0.66	2.98	0.00	0.00	0.10	0.00
time (sec)	N/A	0.532	0.090	0.136	0.124	0.089	0.000	0.000	0.151	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	40	30	40	37	110	0	57	25	0
N.S.	1	0.57	0.43	0.57	0.53	1.57	0.00	0.81	0.36	0.00
time (sec)	N/A	0.338	0.016	0.145	0.111	0.078	0.000	0.118	0.149	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	<b>F</b>	A	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	23	28	27	16	61	0	21	8	0
N.S.	1	0.72	0.88	0.84	0.50	1.91	0.00	0.66	0.25	0.00
time (sec)	N/A	0.234	0.012	0.110	0.110	0.073	0.000	0.116	0.152	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F(-2)</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	21	23	26	13	80	0	0	17	0
N.S.	1	0.68	0.74	0.84	0.42	2.58	0.00	0.00	0.55	0.00
time (sec)	N/A	0.225	0.018	0.108	0.112	0.089	0.000	0.000	0.154	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	<b>F(-2)</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	40	41	42	29	142	0	0	65	0
N.S.	1	0.52	0.53	0.55	0.38	1.84	0.00	0.00	0.84	0.00
time (sec)	N/A	0.345	0.046	0.108	0.121	0.078	0.000	0.000	0.187	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	60	0	0	0	0	0	16	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.310	0.082	0.000	0.000	0.000	0.000	0.000	0.160	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	62	0	0	0	0	0	21	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.34	0.00
time (sec)	N/A	0.330	0.075	0.000	0.000	0.000	0.000	0.000	0.160	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	289	0	0	0	0	0	26	0
N.S.	1	1.00	3.32	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.373	1.805	0.000	0.000	0.000	0.000	0.000	0.168	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	83	0	0	0	0	0	26	0
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.384	0.597	0.000	0.000	0.000	0.000	0.000	0.161	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	65	0	0	0	0	0	19	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.279	0.097	0.000	0.000	0.000	0.000	0.000	0.189	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	83	0	0	0	0	0	26	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.492	0.507	0.000	0.000	0.000	0.000	0.000	0.158	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	71	73	90	84	145	0	0	23	123
N.S.	1	0.93	0.96	1.18	1.11	1.91	0.00	0.00	0.30	1.62
time (sec)	N/A	0.267	0.412	29.432	0.045	0.111	0.000	0.000	0.154	10.175



Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	49	45	60	56	87	0	0	23	84
N.S.	1	0.96	0.88	1.18	1.10	1.71	0.00	0.00	0.45	1.65
time (sec)	N/A	0.254	0.281	3.280	0.037	0.081	0.000	0.000	0.154	10.291

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	26	26	25	41	0	47	23	73
N.S.	1	1.00	1.04	1.04	1.00	1.64	0.00	1.88	0.92	2.92
time (sec)	N/A	0.219	0.014	0.454	0.027	0.080	0.000	0.388	0.176	9.444

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	59	0	0	0	0	0	23	0
N.S.	1	1.00	1.16	0.00	0.00	0.00	0.00	0.00	0.45	0.00
time (sec)	N/A	0.237	0.316	0.000	0.000	0.000	0.000	0.000	0.160	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	59	0	0	0	0	0	23	0
N.S.	1	1.00	1.16	0.00	0.00	0.00	0.00	0.00	0.45	0.00
time (sec)	N/A	0.231	0.352	0.000	0.000	0.000	0.000	0.000	0.162	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	741	0	0	0	0	0	23	0
N.S.	1	1.00	9.38	0.00	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.242	7.808	0.000	0.000	0.000	0.000	0.000	0.154	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	69	0	0	0	0	0	21	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.27	0.00
time (sec)	N/A	0.225	0.224	0.000	0.000	0.000	0.000	0.000	0.165	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	264	0	0	0	0	0	21	0
N.S.	1	1.00	3.62	0.00	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.227	0.894	0.000	0.000	0.000	0.000	0.000	0.187	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	477	0	0	0	0	0	23	0
N.S.	1	1.00	6.04	0.00	0.00	0.00	0.00	0.00	0.29	0.00
time (sec)	N/A	0.237	1.981	0.000	0.000	0.000	0.000	0.000	0.157	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	306	0	0	0	0	0	26	0
N.S.	1	1.00	3.69	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.237	1.937	0.000	0.000	0.000	0.000	0.000	0.155	0.000

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [7] had the largest ratio of [1.8750000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.18	6	0.500
2	A	3	3	1.00	8	0.375
3	A	7	7	1.07	8	0.875
4	A	5	5	1.00	8	0.625
5	A	11	11	1.05	8	1.375
6	A	7	7	1.00	8	0.875
7	A	15	15	1.03	8	1.875
8	A	9	9	1.00	8	1.125
9	A	16	15	1.28	12	1.250
10	A	14	13	1.25	12	1.083
11	A	14	13	1.31	12	1.083
12	A	12	11	1.28	12	0.917
13	A	12	11	1.31	12	0.917
14	A	14	13	1.25	12	1.083
15	A	14	13	1.30	12	1.083
16	A	16	15	1.26	12	1.250
17	A	14	13	0.90	12	1.083
18	A	12	11	0.87	12	0.917
19	A	12	11	0.85	12	0.917
20	A	12	11	0.85	12	0.917
21	A	12	11	0.87	12	0.917

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	14	13	0.90	12	1.083
23	A	4	3	1.00	8	0.375
24	A	4	3	1.00	10	0.300
25	A	9	9	0.75	10	0.900
26	A	5	5	1.00	10	0.500
27	A	4	4	1.00	10	0.400
28	A	6	6	0.72	10	0.600
29	A	18	17	0.96	10	1.700
30	A	16	15	1.03	10	1.500
31	A	16	15	1.02	10	1.500
32	A	18	17	0.91	10	1.700
33	A	9	9	0.57	10	0.900
34	A	5	5	0.72	10	0.500
35	A	5	5	0.68	10	0.500
36	A	9	9	0.52	10	0.900
37	A	6	5	1.00	12	0.417
38	A	6	5	1.00	14	0.357
39	A	4	4	1.00	21	0.190
40	A	4	4	1.00	21	0.190
41	A	5	4	1.00	21	0.190
42	A	6	6	1.00	21	0.286
43	A	5	4	0.93	19	0.211
44	A	5	4	0.96	19	0.211
45	A	4	3	1.00	19	0.158
46	A	4	3	1.00	19	0.158
47	A	4	3	1.00	19	0.158
48	A	2	2	1.00	19	0.105
49	A	2	2	1.00	17	0.118
50	A	2	2	1.00	17	0.118
51	A	2	2	1.00	19	0.105
52	A	2	2	1.00	21	0.095

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int \cot(a + bx) dx$ . . . . .	47
3.2	$\int \cot^2(a + bx) dx$ . . . . .	52
3.3	$\int \cot^3(a + bx) dx$ . . . . .	57
3.4	$\int \cot^4(a + bx) dx$ . . . . .	63
3.5	$\int \cot^5(a + bx) dx$ . . . . .	69
3.6	$\int \cot^6(a + bx) dx$ . . . . .	76
3.7	$\int \cot^7(a + bx) dx$ . . . . .	82
3.8	$\int \cot^8(a + bx) dx$ . . . . .	90
3.9	$\int (c \cot(a + bx))^{7/2} dx$ . . . . .	97
3.10	$\int (c \cot(a + bx))^{5/2} dx$ . . . . .	108
3.11	$\int (c \cot(a + bx))^{3/2} dx$ . . . . .	118
3.12	$\int \sqrt{c \cot(a + bx)} dx$ . . . . .	128
3.13	$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx$ . . . . .	136
3.14	$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx$ . . . . .	145
3.15	$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx$ . . . . .	155
3.16	$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx$ . . . . .	165
3.17	$\int (c \cot(a + bx))^{4/3} dx$ . . . . .	175
3.18	$\int (c \cot(a + bx))^{2/3} dx$ . . . . .	185
3.19	$\int \sqrt[3]{c \cot(a + bx)} dx$ . . . . .	195
3.20	$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx$ . . . . .	204
3.21	$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx$ . . . . .	213
3.22	$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx$ . . . . .	223
3.23	$\int \cot^n(a + bx) dx$ . . . . .	234
3.24	$\int (b \cot(c + dx))^n dx$ . . . . .	239
3.25	$\int (a \cot^2(x))^{3/2} dx$ . . . . .	244

3.26	$\int \sqrt{a \cot^2(x)} dx$	250
3.27	$\int \frac{1}{\sqrt{a \cot^2(x)}} dx$	255
3.28	$\int \frac{1}{(a \cot^2(x))^{3/2}} dx$	260
3.29	$\int (a \cot^3(x))^{3/2} dx$	266
3.30	$\int \sqrt{a \cot^3(x)} dx$	276
3.31	$\int \frac{1}{\sqrt{a \cot^3(x)}} dx$	286
3.32	$\int \frac{1}{(a \cot^3(x))^{3/2}} dx$	295
3.33	$\int (a \cot^4(x))^{3/2} dx$	305
3.34	$\int \sqrt{a \cot^4(x)} dx$	311
3.35	$\int \frac{1}{\sqrt{a \cot^4(x)}} dx$	316
3.36	$\int \frac{1}{(a \cot^4(x))^{3/2}} dx$	321
3.37	$\int (b \cot^p(c + dx))^n dx$	327
3.38	$\int (a(b \cot(c + dx))^p)^n dx$	332
3.39	$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx$	337
3.40	$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx$	342
3.41	$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx$	347
3.42	$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx$	352
3.43	$\int (d \cot(e + fx))^n \csc^6(e + fx) dx$	358
3.44	$\int (d \cot(e + fx))^n \csc^4(e + fx) dx$	364
3.45	$\int (d \cot(e + fx))^n \csc^2(e + fx) dx$	370
3.46	$\int (d \cot(e + fx))^n \sin^2(e + fx) dx$	375
3.47	$\int (d \cot(e + fx))^n \sin^4(e + fx) dx$	380
3.48	$\int (d \cot(e + fx))^n \csc^3(e + fx) dx$	385
3.49	$\int (d \cot(e + fx))^n \csc(e + fx) dx$	390
3.50	$\int (d \cot(e + fx))^n \sin(e + fx) dx$	395
3.51	$\int (d \cot(e + fx))^n \sin^3(e + fx) dx$	400
3.52	$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx$	405

### 3.1 $\int \cot(a + bx) dx$

Optimal result . . . . .	47
Mathematica [A] (verified) . . . . .	47
Rubi [A] (verified) . . . . .	48
Maple [A] (verified) . . . . .	49
Fricas [A] (verification not implemented) . . . . .	49
Sympy [B] (verification not implemented) . . . . .	50
Maxima [A] (verification not implemented) . . . . .	50
Giac [A] (verification not implemented) . . . . .	50
Mupad [B] (verification not implemented) . . . . .	51
Reduce [B] (verification not implemented) . . . . .	51

#### Optimal result

Integrand size = 6, antiderivative size = 11

$$\int \cot(a + bx) dx = \frac{\log(\sin(a + bx))}{b}$$

output `ln(sin(b*x+a))/b`

#### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cot(a + bx) dx = \frac{\log(\sin(a + bx))}{b}$$

input `Integrate[Cot[a + b*x],x]`

output `Log[Sin[a + b*x]]/b`



**Rubi [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.18, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \cot(a + bx) dx \\
 \downarrow 3042 \\
 \int -\tan\left(a + bx + \frac{\pi}{2}\right) dx \\
 \downarrow 25 \\
 -\int \tan\left(\frac{1}{2}(2a + \pi) + bx\right) dx \\
 \downarrow 3956 \\
 \frac{\log(-\sin(a + bx))}{b}
 \end{array}$$

input `Int[Cot[a + b*x],x]`

output `Log[-Sin[a + b*x]]/b`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

**Maple [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

method	result	size
derivativedivides	$-\frac{\ln(\cot(bx+a)^2+1)}{2b}$	17
default	$-\frac{\ln(\cot(bx+a)^2+1)}{2b}$	17
parallelrisc	$\frac{\ln(\tan(bx+a))+\ln\left(\frac{1}{\sqrt{\sec(bx+a)^2}}\right)}{b}$	24
norman	$\frac{\ln(\tan(bx+a))}{b} - \frac{\ln(1+\tan(bx+a)^2)}{2b}$	29
risc	$-ix - \frac{2ia}{b} + \frac{\ln(e^{2i(bx+a)}-1)}{b}$	29

input

```
int(cot(b*x+a),x,method=_RETURNVERBOSE)
```

output

```
-1/2/b*ln(cot(b*x+a)^2+1)
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.73

$$\int \cot(a + bx) dx = \frac{\log\left(-\frac{1}{2} \cos(2bx + 2a) + \frac{1}{2}\right)}{2b}$$

input

```
integrate(cot(b*x+a),x, algorithm="fricas")
```

output

```
1/2*log(-1/2*cos(2*b*x + 2*a) + 1/2)/b
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 29 vs.  $2(8) = 16$ .

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.64

$$\int \cot(a + bx) dx = \begin{cases} -\frac{\log(\tan^2(a+bx)+1)}{2b} + \frac{\log(\tan(a+bx))}{b} & \text{for } b \neq 0 \\ x \cot(a) & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a), x)`

output `Piecewise((-log(tan(a + b*x)**2 + 1)/(2*b) + log(tan(a + b*x))/b, Ne(b, 0)), (x*cot(a), True))`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \cot(a + bx) dx = \frac{\log(\sin(bx + a))}{b}$$

input `integrate(cot(b*x+a), x, algorithm="maxima")`

output `log(sin(b*x + a))/b`

**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int \cot(a + bx) dx = \frac{\log(|\sin(bx + a)|)}{b}$$

input `integrate(cot(b*x+a), x, algorithm="giac")`

output `log(abs(sin(b*x + a)))/b`

**Mupad [B] (verification not implemented)**

Time = 10.17 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.18

$$\int \cot(a + bx) dx = -x \operatorname{li} + \frac{\ln(e^{a+2i} e^{bx+2i} - 1)}{b}$$

input `int(cot(a + b*x),x)`

output `log(exp(a*2i)*exp(b*x*2i) - 1)/b - x*1i`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.82

$$\int \cot(a + bx) dx = \frac{-\log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + 1\right) + \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$$

input `int(cot(b*x+a),x)`

output `( - log(tan((a + b*x)/2)**2 + 1) + log(tan((a + b*x)/2)))/b`

## 3.2 $\int \cot^2(a + bx) dx$

Optimal result . . . . .	52
Mathematica [C] (verified) . . . . .	52
Rubi [A] (verified) . . . . .	53
Maple [A] (verified) . . . . .	54
Fricas [B] (verification not implemented) . . . . .	54
Sympy [A] (verification not implemented) . . . . .	55
Maxima [A] (verification not implemented) . . . . .	55
Giac [B] (verification not implemented) . . . . .	55
Mupad [B] (verification not implemented) . . . . .	56
Reduce [B] (verification not implemented) . . . . .	56

### Optimal result

Integrand size = 8, antiderivative size = 15

$$\int \cot^2(a + bx) dx = -x - \frac{\cot(a + bx)}{b}$$

output `-x-cot(b*x+a)/b`

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.93

$$\int \cot^2(a + bx) dx = -\frac{\cot(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(a + bx)\right)}{b}$$

input `Integrate[Cot[a + b*x]^2,x]`

output `-((Cot[a + b*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[a + b*x]^2])/b)`

**Rubi [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^2(a + bx) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan\left(a + bx + \frac{\pi}{2}\right)^2 dx \\ & \quad \downarrow \text{3954} \\ & - \int 1 dx - \frac{\cot(a + bx)}{b} \\ & \quad \downarrow \text{24} \\ & -\frac{\cot(a + bx)}{b} - x \end{aligned}$$

input `Int[Cot[a + b*x]^2,x]`

output `-x - Cot[a + b*x]/b`

**Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

**Maple [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

method	result	size
parallelrisch	$-\frac{bx - \cot(bx+a)}{b}$	18
risch	$-x - \frac{2i}{b(e^{2i(bx+a)} - 1)}$	24
norman	$-\frac{\frac{1}{b} - x \tan(bx+a)}{\tan(bx+a)}$	25
derivativedivides	$-\frac{\cot(bx+a) + \frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))}{b}$	26
default	$-\frac{\cot(bx+a) + \frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))}{b}$	26

input

```
int(cot(b*x+a)^2, x, method=_RETURNVERBOSE)
```

output

```
1/b*(-b*x-cot(b*x+a))
```

**Fricas [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 39 vs.  $2(15) = 30$ .

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.60

$$\int \cot^2(a + bx) dx = -\frac{bx \sin(2bx + 2a) + \cos(2bx + 2a) + 1}{b \sin(2bx + 2a)}$$

input

```
integrate(cot(b*x+a)^2, x, algorithm="fricas")
```

output

```
-(b*x*sin(2*b*x + 2*a) + cos(2*b*x + 2*a) + 1)/(b*sin(2*b*x + 2*a))
```

**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cot^2(a + bx) dx = \begin{cases} -x - \frac{\cot(a+bx)}{b} & \text{for } b \neq 0 \\ x \cot^2(a) & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a)**2,x)`

output `Piecewise((-x - cot(a + b*x)/b, Ne(b, 0)), (x*cot(a)**2, True))`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \cot^2(a + bx) dx = -\frac{bx + a + \frac{1}{\tan(bx+a)}}{b}$$

input `integrate(cot(b*x+a)^2,x, algorithm="maxima")`

output `-(b*x + a + 1/tan(b*x + a))/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(15) = 30.

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.33

$$\int \cot^2(a + bx) dx = -\frac{2bx + 2a + \frac{1}{\tan(\frac{1}{2}bx + \frac{1}{2}a)} - \tan(\frac{1}{2}bx + \frac{1}{2}a)}{2b}$$

input `integrate(cot(b*x+a)^2,x, algorithm="giac")`

output `-1/2*(2*b*x + 2*a + 1/tan(1/2*b*x + 1/2*a) - tan(1/2*b*x + 1/2*a))/b`



**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cot^2(a + bx) dx = -x - \frac{\cot(a + bx)}{b}$$

input `int(cot(a + b*x)^2,x)`output `- x - cot(a + b*x)/b`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \cot^2(a + bx) dx = \frac{-\cot(bx + a) - bx}{b}$$

input `int(cot(b*x+a)^2,x)`output `( - (cot(a + b*x) + b*x))/b`

### 3.3 $\int \cot^3(a + bx) dx$

Optimal result . . . . .	57
Mathematica [A] (verified) . . . . .	57
Rubi [A] (verified) . . . . .	58
Maple [A] (verified) . . . . .	59
Fricas [A] (verification not implemented) . . . . .	60
Sympy [B] (verification not implemented) . . . . .	60
Maxima [A] (verification not implemented) . . . . .	61
Giac [A] (verification not implemented) . . . . .	61
Mupad [B] (verification not implemented) . . . . .	61
Reduce [B] (verification not implemented) . . . . .	62

#### Optimal result

Integrand size = 8, antiderivative size = 28

$$\int \cot^3(a + bx) dx = -\frac{\cot^2(a + bx)}{2b} - \frac{\log(\sin(a + bx))}{b}$$

output `-1/2*cot(b*x+a)^2/b-ln(sin(b*x+a))/b`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \cot^3(a + bx) dx = -\frac{\csc^2(a + bx)}{2b} - \frac{\log(\sin(a + bx))}{b}$$

input `Integrate[Cot[a + b*x]^3,x]`

output `-1/2*Csc[a + b*x]^2/b - Log[Sin[a + b*x]]/b`

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 25, 3954, 25, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(a + bx + \frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(\frac{1}{2}(2a + \pi) + bx\right)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & \int -\cot(a + bx) dx - \frac{\cot^2(a + bx)}{2b} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot(a + bx) dx - \frac{\cot^2(a + bx)}{2b} \\
 & \quad \downarrow \text{3042} \\
 & -\int -\tan\left(a + bx + \frac{\pi}{2}\right) dx - \frac{\cot^2(a + bx)}{2b} \\
 & \quad \downarrow \text{25} \\
 & \int \tan\left(\frac{1}{2}(2a + \pi) + bx\right) dx - \frac{\cot^2(a + bx)}{2b} \\
 & \quad \downarrow \text{3956} \\
 & -\frac{\cot^2(a + bx)}{2b} - \frac{\log(-\sin(a + bx))}{b}
 \end{aligned}$$

input

Int[Cot[a + b\*x]^3,x]

output  $-1/2*\text{Cot}[a + b*x]^2/b - \text{Log}[-\text{Sin}[a + b*x]]/b$

### Defintions of rubi rules used

rule 25  $\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$

rule 3042  $\text{Int}[\text{u}_, \text{x\_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] \text{ ; FunctionOfTrigOfLinear} \text{ Q}[\text{u}, \text{x}]$

rule 3954  $\text{Int}[(\text{b}_.*\text{tan}[(\text{c}_.) + (\text{d}_.)*(\text{x}_)])^{(\text{n}_)}, \text{x\_Symbol}] \rightarrow \text{Simp}[\text{b}*((\text{b}*\text{Tan}[\text{c} + \text{d} * \text{x}])^{(\text{n} - 1)}/(\text{d}*(\text{n} - 1))), \text{x}] - \text{Simp}[\text{b}^2 \text{ Int}[(\text{b}*\text{Tan}[\text{c} + \text{d}*\text{x}])^{(\text{n} - 2)}, \text{x}], \text{x}] \text{ ; FreeQ}\{\text{b}, \text{c}, \text{d}\}, \text{x}\} \&\& \text{GtQ}[\text{n}, 1]$

rule 3956  $\text{Int}[\text{tan}[(\text{c}_.) + (\text{d}_.)*(\text{x}_)], \text{x\_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[\text{c} + \text{d} * \text{x}], \text{x}]]/\text{d}, \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}\}, \text{x}\}$

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$\frac{-\frac{\cot(bx+a)^2}{2} + \frac{\ln(\cot(bx+a)^2+1)}{2}}{b}$	29
default	$\frac{-\frac{\cot(bx+a)^2}{2} + \frac{\ln(\cot(bx+a)^2+1)}{2}}{b}$	29
parallelrisch	$\frac{-\cot(bx+a)^2 - 2\ln(\tan(bx+a)) + \ln(\sec(bx+a)^2)}{2b}$	35
norman	$-\frac{1}{2b \tan(bx+a)^2} - \frac{\ln(\tan(bx+a))}{b} + \frac{\ln(1+\tan(bx+a)^2)}{2b}$	43
risch	$ix + \frac{2ia}{b} + \frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}-1)^2} - \frac{\ln(e^{2i(bx+a)}-1)}{b}$	57

input  $\text{int}(\cot(b*x+a)^3, \text{x}, \text{method}=\_RETURNVERBOSE)$

output `1/b*(-1/2*cot(b*x+a)^2+1/2*ln(cot(b*x+a)^2+1))`

### Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.68

$$\int \cot^3(a + bx) dx = -\frac{(\cos(2bx + 2a) - 1) \log\left(-\frac{1}{2} \cos(2bx + 2a) + \frac{1}{2}\right) - 2}{2(b \cos(2bx + 2a) - b)}$$

input `integrate(cot(b*x+a)^3,x, algorithm="fricas")`

output `-1/2*((cos(2*b*x + 2*a) - 1)*log(-1/2*cos(2*b*x + 2*a) + 1/2) - 2)/(b*cos(2*b*x + 2*a) - b)`

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(22) = 44$ .

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.89

$$\int \cot^3(a + bx) dx = \begin{cases} \tilde{\infty}x & \text{for } a = 0 \wedge b = 0 \\ x \cot^3(a) & \text{for } b = 0 \\ \tilde{\infty}x & \text{for } a = -bx \\ \frac{\log(\tan^2(a+bx)+1)}{2b} - \frac{\log(\tan(a+bx))}{b} - \frac{1}{2b \tan^2(a+bx)} & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a)**3,x)`

output `Piecewise((zoo*x, Eq(a, 0) & Eq(b, 0)), (x*cot(a)**3, Eq(b, 0)), (zoo*x, Eq(a, -b*x)), (log(tan(a + b*x)**2 + 1)/(2*b) - log(tan(a + b*x))/b - 1/(2*b*tan(a + b*x)**2), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.82

$$\int \cot^3(a + bx) dx = -\frac{\frac{1}{\sin(bx+a)^2} + \log(\sin(bx+a)^2)}{2b}$$

input `integrate(cot(b*x+a)^3,x, algorithm="maxima")`output `-1/2*(1/sin(b*x + a)^2 + log(sin(b*x + a)^2))/b`**Giac [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \cot^3(a + bx) dx = -\frac{\log(|\sin(bx+a)|)}{b} - \frac{1}{2b \sin(bx+a)^2}$$

input `integrate(cot(b*x+a)^3,x, algorithm="giac")`output `-log(abs(sin(b*x + a)))/b - 1/2/(b*sin(b*x + a)^2)`**Mupad [B] (verification not implemented)**

Time = 12.32 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.79

$$\int \cot^3(a + bx) dx = x \operatorname{li} - \frac{\ln(e^{a+2i} e^{bx+2i} - 1)}{b} + \frac{2}{b(e^{a+2i+bx+2i} - 1)} + \frac{2}{b(1 + e^{a+4i+bx+4i} - 2e^{a+2i+bx+2i})}$$

input `int(cot(a + b*x)^3,x)`output `x*li - log(exp(a*2i)*exp(b*x*2i) - 1)/b + 2/(b*(exp(a*2i + b*x*2i) - 1)) + 2/(b*(exp(a*4i + b*x*4i) - 2*exp(a*2i + b*x*2i) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.39

$$\int \cot^3(a + bx) dx$$

$$= \frac{4 \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + 1\right) \sin(bx + a)^2 - 4 \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \sin(bx + a)^2 + \sin(bx + a)^2 - 2}{4 \sin(bx + a)^2 b}$$

input `int(cot(b*x+a)^3,x)`output `(4*log(tan((a + b*x)/2)**2 + 1)*sin(a + b*x)**2 - 4*log(tan((a + b*x)/2))*sin(a + b*x)**2 + sin(a + b*x)**2 - 2)/(4*sin(a + b*x)**2*b)`

### 3.4 $\int \cot^4(a + bx) dx$

Optimal result	63
Mathematica [C] (verified)	63
Rubi [A] (verified)	64
Maple [A] (verified)	65
Fricas [B] (verification not implemented)	66
Sympy [A] (verification not implemented)	66
Maxima [A] (verification not implemented)	67
Giac [B] (verification not implemented)	67
Mupad [B] (verification not implemented)	68
Reduce [B] (verification not implemented)	68

#### Optimal result

Integrand size = 8, antiderivative size = 27

$$\int \cot^4(a + bx) dx = x + \frac{\cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b}$$

output `x+cot(b*x+a)/b-1/3*cot(b*x+a)^3/b`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.22

$$\int \cot^4(a + bx) dx = -\frac{\cot^3(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(a + bx)\right)}{3b}$$

input `Integrate[Cot[a + b*x]^4,x]`

output `-1/3*(Cot[a + b*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[a + b*x]^2])/b`



**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(a + bx + \frac{\pi}{2}\right)^4 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int \cot^2(a + bx) dx - \frac{\cot^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & - \int \tan\left(a + bx + \frac{\pi}{2}\right)^2 dx - \frac{\cot^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3954} \\
 & \int 1 dx - \frac{\cot^3(a + bx)}{3b} + \frac{\cot(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & - \frac{\cot^3(a + bx)}{3b} + \frac{\cot(a + bx)}{b} + x
 \end{aligned}$$

input

Int[Cot[a + b\*x]^4,x]

output

x + Cot[a + b\*x]/b - Cot[a + b\*x]^3/(3\*b)

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

### Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

method	result	size
parallelrisc	$\frac{-\cot(bx+a)^3 + 3bx + 3\cot(bx+a)}{3b}$	29
derivativedivides	$\frac{-\frac{\cot(bx+a)^3}{3} + \cot(bx+a) - \frac{\pi}{2} + \operatorname{arccot}(\cot(bx+a))}{b}$	32
default	$\frac{-\frac{\cot(bx+a)^3}{3} + \cot(bx+a) - \frac{\pi}{2} + \operatorname{arccot}(\cot(bx+a))}{b}$	32
norman	$\frac{\frac{\tan(bx+a)^2}{b} + x \tan(bx+a)^3 - \frac{1}{3b}}{\tan(bx+a)^3}$	38
risc	$x + \frac{4i(3e^{4i(bx+a)} - 3e^{2i(bx+a)} + 2)}{3b(e^{2i(bx+a)} - 1)^3}$	46

input `int(cot(b*x+a)^4, x, method=_RETURNVERBOSE)`

output `1/3*(-cot(b*x+a)^3+3*b*x+3*cot(b*x+a))/b`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(25) = 50.

Time = 0.08 (sec) , antiderivative size = 84, normalized size of antiderivative = 3.11

$$\int \cot^4(a + bx) dx = \frac{4 \cos(2bx + 2a)^2 + 3(bx \cos(2bx + 2a) - bx) \sin(2bx + 2a) + 2 \cos(2bx + 2a) - 2}{3(b \cos(2bx + 2a) - b) \sin(2bx + 2a)}$$

input `integrate(cot(b*x+a)^4,x, algorithm="fricas")`

output `1/3*(4*cos(2*b*x + 2*a)^2 + 3*(b*x*cos(2*b*x + 2*a) - b*x)*sin(2*b*x + 2*a) + 2*cos(2*b*x + 2*a) - 2)/((b*cos(2*b*x + 2*a) - b)*sin(2*b*x + 2*a))`

**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \cot^4(a + bx) dx = \begin{cases} x - \frac{\cot^3(a+bx)}{3b} + \frac{\cot(a+bx)}{b} & \text{for } b \neq 0 \\ x \cot^4(a) & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a)**4,x)`

output `Piecewise((x - cot(a + b*x)**3/(3*b) + cot(a + b*x)/b, Ne(b, 0)), (x*cot(a)**4, True))`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int \cot^4(a + bx) dx = \frac{3bx + 3a + \frac{3 \tan(bx+a)^2 - 1}{\tan(bx+a)^3}}{3b}$$

input `integrate(cot(b*x+a)^4,x, algorithm="maxima")`

output `1/3*(3*b*x + 3*a + (3*tan(b*x + a)^2 - 1)/tan(b*x + a)^3)/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(25) = 50.

Time = 0.16 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.30

$$\int \cot^4(a + bx) dx$$

$$= \frac{\tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3 + 24bx + 24a + \frac{15 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - 1}{\tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3} - 15 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)}{24b}$$

input `integrate(cot(b*x+a)^4,x, algorithm="giac")`

output `1/24*(tan(1/2*b*x + 1/2*a)^3 + 24*b*x + 24*a + (15*tan(1/2*b*x + 1/2*a)^2 - 1)/tan(1/2*b*x + 1/2*a)^3 - 15*tan(1/2*b*x + 1/2*a))/b`

**Mupad [B] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \cot^4(a + bx) dx = x + \frac{\cot(a + bx) - \frac{\cot(a+bx)^3}{3}}{b}$$

input `int(cot(a + b*x)^4,x)`

output `x + (cot(a + b*x) - cot(a + b*x)^3/3)/b`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.04

$$\int \cot^4(a + bx) dx = \frac{-\cot(bx + a)^3 + 3\cot(bx + a) + 3bx}{3b}$$

input `int(cot(b*x+a)^4,x)`

output `( - cot(a + b*x)**3 + 3*cot(a + b*x) + 3*b*x)/(3*b)`

### 3.5 $\int \cot^5(a + bx) dx$

Optimal result . . . . .	69
Mathematica [A] (verified) . . . . .	69
Rubi [A] (verified) . . . . .	70
Maple [A] (verified) . . . . .	72
Fricas [B] (verification not implemented) . . . . .	72
Sympy [B] (verification not implemented) . . . . .	73
Maxima [A] (verification not implemented) . . . . .	73
Giac [A] (verification not implemented) . . . . .	74
Mupad [B] (verification not implemented) . . . . .	74
Reduce [B] (verification not implemented) . . . . .	75

#### Optimal result

Integrand size = 8, antiderivative size = 42

$$\int \cot^5(a + bx) dx = \frac{\cot^2(a + bx)}{2b} - \frac{\cot^4(a + bx)}{4b} + \frac{\log(\sin(a + bx))}{b}$$

output

```
1/2*cot(b*x+a)^2/b-1/4*cot(b*x+a)^4/b+ln(sin(b*x+a))/b
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.93

$$\int \cot^5(a + bx) dx = \frac{\csc^2(a + bx)}{b} - \frac{\csc^4(a + bx)}{4b} + \frac{\log(\sin(a + bx))}{b}$$

input

```
Integrate[Cot[a + b*x]^5,x]
```

output

```
Csc[a + b*x]^2/b - Csc[a + b*x]^4/(4*b) + Log[Sin[a + b*x]]/b
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.375$ , Rules used = {3042, 25, 3954, 25, 3042, 25, 3954, 25, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^5(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(a + bx + \frac{\pi}{2}\right)^5 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(\frac{1}{2}(2a + \pi) + bx\right)^5 dx \\
 & \quad \downarrow \text{3954} \\
 & \int -\cot^3(a + bx) dx - \frac{\cot^4(a + bx)}{4b} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot^3(a + bx) dx - \frac{\cot^4(a + bx)}{4b} \\
 & \quad \downarrow \text{3042} \\
 & -\int -\tan\left(a + bx + \frac{\pi}{2}\right)^3 dx - \frac{\cot^4(a + bx)}{4b} \\
 & \quad \downarrow \text{25} \\
 & \int \tan\left(\frac{1}{2}(2a + \pi) + bx\right)^3 dx - \frac{\cot^4(a + bx)}{4b} \\
 & \quad \downarrow \text{3954} \\
 & -\int -\cot(a + bx) dx - \frac{\cot^4(a + bx)}{4b} + \frac{\cot^2(a + bx)}{2b} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
& \int \cot(a + bx) dx - \frac{\cot^4(a + bx)}{4b} + \frac{\cot^2(a + bx)}{2b} \\
& \quad \downarrow \text{3042} \\
& \int -\tan\left(a + bx + \frac{\pi}{2}\right) dx - \frac{\cot^4(a + bx)}{4b} + \frac{\cot^2(a + bx)}{2b} \\
& \quad \downarrow \text{25} \\
& -\int \tan\left(\frac{1}{2}(2a + \pi) + bx\right) dx - \frac{\cot^4(a + bx)}{4b} + \frac{\cot^2(a + bx)}{2b} \\
& \quad \downarrow \text{3956} \\
& -\frac{\cot^4(a + bx)}{4b} + \frac{\cot^2(a + bx)}{2b} + \frac{\log(-\sin(a + bx))}{b}
\end{aligned}$$

input `Int[Cot[a + b*x]^5,x]`

output `Cot[a + b*x]^2/(2*b) - Cot[a + b*x]^4/(4*b) + Log[-Sin[a + b*x]]/b`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`



**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$\frac{-\frac{\cot(bx+a)^4}{4} + \frac{\cot(bx+a)^2}{2} - \frac{\ln(\cot(bx+a)^2+1)}{2}}{b}$	39
default	$\frac{-\frac{\cot(bx+a)^4}{4} + \frac{\cot(bx+a)^2}{2} - \frac{\ln(\cot(bx+a)^2+1)}{2}}{b}$	39
parallelrisch	$\frac{-\cot(bx+a)^4 + 2\cot(bx+a)^2 + 4\ln(\tan(bx+a)) - 2\ln(\sec(bx+a)^2)}{4b}$	47
norman	$\frac{-\frac{1}{4b} + \frac{\tan(bx+a)^2}{2b}}{\tan(bx+a)^4} + \frac{\ln(\tan(bx+a))}{b} - \frac{\ln(1+\tan(bx+a)^2)}{2b}$	57
risch	$-ix - \frac{2ia}{b} - \frac{4(e^{6i(bx+a)} - e^{4i(bx+a)} + e^{2i(bx+a)})}{b(e^{2i(bx+a)} - 1)^4} + \frac{\ln(e^{2i(bx+a)} - 1)}{b}$	77

input `int(cot(b*x+a)^5,x,method=_RETURNVERBOSE)`output `1/b*(-1/4*cot(b*x+a)^4+1/2*cot(b*x+a)^2-1/2*ln(cot(b*x+a)^2+1))`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. 2(38) = 76.

Time = 0.08 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.98

$$\int \cot^5(a + bx) dx$$

$$= \frac{(\cos(2bx + 2a)^2 - 2\cos(2bx + 2a) + 1) \log\left(-\frac{1}{2}\cos(2bx + 2a) + \frac{1}{2}\right) - 4\cos(2bx + 2a) + 2}{2(b\cos(2bx + 2a)^2 - 2b\cos(2bx + 2a) + b)}$$

input `integrate(cot(b*x+a)^5,x, algorithm="fricas")`output `1/2*((cos(2*b*x + 2*a)^2 - 2*cos(2*b*x + 2*a) + 1)*log(-1/2*cos(2*b*x + 2*a) + 1/2) - 4*cos(2*b*x + 2*a) + 2)/(b*cos(2*b*x + 2*a)^2 - 2*b*cos(2*b*x + 2*a) + b)`

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 66 vs.  $2(32) = 64$ .

Time = 0.32 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.57

$$\int \cot^5(a + bx) dx = \begin{cases} \tilde{\infty}x & \text{for } a = 0 \wedge b = 0 \\ x \cot^5(a) & \text{for } b = 0 \\ \tilde{\infty}x & \text{for } a = -bx \\ -\frac{\log(\tan^2(a+bx)+1)}{2b} + \frac{\log(\tan(a+bx))}{b} + \frac{1}{2b \tan^2(a+bx)} - \frac{1}{4b \tan^4(a+bx)} & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a)**5,x)`

output `Piecewise((zoo*x, Eq(a, 0) & Eq(b, 0)), (x*cot(a)**5, Eq(b, 0)), (zoo*x, Eq(a, -b*x)), (-log(tan(a + b*x)**2 + 1)/(2*b) + log(tan(a + b*x))/b + 1/(2*b*tan(a + b*x)**2) - 1/(4*b*tan(a + b*x)**4), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.90

$$\int \cot^5(a + bx) dx = \frac{\frac{4 \sin(bx+a)^2 - 1}{\sin(bx+a)^4} + 2 \log(\sin(bx + a)^2)}{4b}$$

input `integrate(cot(b*x+a)^5,x, algorithm="maxima")`

output `1/4*((4*sin(b*x + a)^2 - 1)/sin(b*x + a)^4 + 2*log(sin(b*x + a)^2))/b`

**Giac [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.90

$$\int \cot^5(a + bx) dx = \frac{\log(|\sin(bx + a)|)}{b} + \frac{4 \sin(bx + a)^2 - 1}{4 b \sin(bx + a)^4}$$

input `integrate(cot(b*x+a)^5,x, algorithm="giac")`

output `log(abs(sin(b*x + a)))/b + 1/4*(4*sin(b*x + a)^2 - 1)/(b*sin(b*x + a)^4)`

**Mupad [B] (verification not implemented)**

Time = 13.76 (sec) , antiderivative size = 182, normalized size of antiderivative = 4.33

$$\begin{aligned} \int \cot^5(a + bx) dx = & -x \operatorname{li} + \frac{\ln(e^{a+2i} e^{bx+2i} - 1)}{b} - \frac{4}{b(e^{a+2i+bx+2i} - 1)} \\ & - \frac{b(1 + e^{a+4i+bx+4i} - 2e^{a+2i+bx+2i})}{8} \\ & - \frac{b(3e^{a+2i+bx+2i} - 3e^{a+4i+bx+4i} + e^{a+6i+bx+6i} - 1)}{4} \\ & - \frac{b(1 + 6e^{a+4i+bx+4i} - 4e^{a+6i+bx+6i} + e^{a+8i+bx+8i} - 4e^{a+2i+bx+2i})}{4} \end{aligned}$$

input `int(cot(a + b*x)^5,x)`

output `log(exp(a*2i)*exp(b*x*2i) - 1)/b - x*1i - 4/(b*(exp(a*2i + b*x*2i) - 1)) - 8/(b*(exp(a*4i + b*x*4i) - 2*exp(a*2i + b*x*2i) + 1)) - 8/(b*(3*exp(a*2i + b*x*2i) - 3*exp(a*4i + b*x*4i) + exp(a*6i + b*x*6i) - 1)) - 4/(b*(6*exp(a*4i + b*x*4i) - 4*exp(a*2i + b*x*2i) - 4*exp(a*6i + b*x*6i) + exp(a*8i + b*x*8i) + 1))`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.88

$$\int \cot^5(a + bx) dx$$

$$= \frac{-32 \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + 1\right) \sin(bx + a)^4 + 32 \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \sin(bx + a)^4 - 13 \sin(bx + a)^4 + 32 \sin(bx + a)^2 - 8}{32 \sin(bx + a)^4 b}$$

input `int(cot(b*x+a)^5,x)`output `( - 32*log(tan((a + b*x)/2)**2 + 1)*sin(a + b*x)**4 + 32*log(tan((a + b*x)/2))*sin(a + b*x)**4 - 13*sin(a + b*x)**4 + 32*sin(a + b*x)**2 - 8)/(32*sin(a + b*x)**4*b)`

### 3.6 $\int \cot^6(a + bx) dx$

Optimal result . . . . .	76
Mathematica [C] (verified) . . . . .	76
Rubi [A] (verified) . . . . .	77
Maple [A] (verified) . . . . .	78
Fricas [B] (verification not implemented) . . . . .	79
Sympy [A] (verification not implemented) . . . . .	79
Maxima [A] (verification not implemented) . . . . .	80
Giac [B] (verification not implemented) . . . . .	80
Mupad [B] (verification not implemented) . . . . .	81
Reduce [B] (verification not implemented) . . . . .	81

#### Optimal result

Integrand size = 8, antiderivative size = 45

$$\int \cot^6(a + bx) dx = -x - \frac{\cot(a + bx)}{b} + \frac{\cot^3(a + bx)}{3b} - \frac{\cot^5(a + bx)}{5b}$$

output `-x-cot(b*x+a)/b+1/3*cot(b*x+a)^3/b-1/5*cot(b*x+a)^5/b`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \cot^6(a + bx) dx = -\frac{\cot^5(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{2}, 1, -\frac{3}{2}, -\tan^2(a + bx)\right)}{5b}$$

input `Integrate[Cot[a + b*x]^6,x]`

output `-1/5*(Cot[a + b*x]^5*Hypergeometric2F1[-5/2, 1, -3/2, -Tan[a + b*x]^2])/b`

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$ , Rules used = {3042, 3954, 3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^6(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(a + bx + \frac{\pi}{2}\right)^6 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int \cot^4(a + bx) dx - \frac{\cot^5(a + bx)}{5b} \\
 & \quad \downarrow \text{3042} \\
 & - \int \tan\left(a + bx + \frac{\pi}{2}\right)^4 dx - \frac{\cot^5(a + bx)}{5b} \\
 & \quad \downarrow \text{3954} \\
 & \int \cot^2(a + bx) dx - \frac{\cot^5(a + bx)}{5b} + \frac{\cot^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(a + bx + \frac{\pi}{2}\right)^2 dx - \frac{\cot^5(a + bx)}{5b} + \frac{\cot^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3954} \\
 & - \int 1 dx - \frac{\cot^5(a + bx)}{5b} + \frac{\cot^3(a + bx)}{3b} - \frac{\cot(a + bx)}{b} \\
 & \quad \downarrow \text{24} \\
 & - \frac{\cot^5(a + bx)}{5b} + \frac{\cot^3(a + bx)}{3b} - \frac{\cot(a + bx)}{b} - x
 \end{aligned}$$

input

Int[Cot[a + b\*x]^6,x]

output  $-x - \cot[a + b*x]/b + \cot[a + b*x]^3/(3*b) - \cot[a + b*x]^5/(5*b)$

### Defintions of rubi rules used

rule 24  $\text{Int}[a_, x\_Symbol] \text{ :> Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

rule 3042  $\text{Int}[u_, x\_Symbol] \text{ :> Int}[\text{DeactivateTrig}[u, x], x] \text{ /; FunctionOfTrigOfLinearQ}[u, x]$

rule 3954  $\text{Int}[(b\_)*\tan[(c\_)+(d\_)*(x\_)]^{(n\_)}, x\_Symbol] \text{ :> Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)})/(d*(n - 1)), x] - \text{Simp}[b^2 \text{ Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] \text{ /; FreeQ}\{b, c, d\}, x] \ \&\& \text{GtQ}[n, 1]$

### Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

method	result	size
parallelsch	$\frac{-3 \cot(bx+a)^5 + 5 \cot(bx+a)^3 - 15bx - 15 \cot(bx+a)}{15b}$	39
derivativedivides	$\frac{-\frac{\cot(bx+a)^5}{5} + \frac{\cot(bx+a)^3}{3} - \cot(bx+a) + \frac{\pi}{2} - \text{arccot}(\cot(bx+a))}{b}$	46
default	$\frac{-\frac{\cot(bx+a)^5}{5} + \frac{\cot(bx+a)^3}{3} - \cot(bx+a) + \frac{\pi}{2} - \text{arccot}(\cot(bx+a))}{b}$	46
norman	$\frac{-\frac{1}{5b} + \frac{\tan(bx+a)^2}{3b} - \frac{\tan(bx+a)^4}{b} - x \tan(bx+a)^5}{\tan(bx+a)^5}$	53
risch	$-x - \frac{2i(45 e^{8i(bx+a)} - 90 e^{6i(bx+a)} + 140 e^{4i(bx+a)} - 70 e^{2i(bx+a)} + 23)}{15b(e^{2i(bx+a)} - 1)^5}$	70

input  $\text{int}(\cot(b*x+a)^6, x, \text{method}=\_RETURNVERBOSE)$

output  $1/15*(-3*\cot(b*x+a)^5+5*\cot(b*x+a)^3-15*b*x-15*\cot(b*x+a))/b$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 123 vs.  $2(41) = 82$ .

Time = 0.11 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.73

$$\int \cot^6(a + bx) dx = \frac{23 \cos(2bx + 2a)^3 - \cos(2bx + 2a)^2 + 15 (bx \cos(2bx + 2a)^2 - 2bx \cos(2bx + 2a) + bx) \sin(2bx + 2a)}{15 (b \cos(2bx + 2a)^2 - 2b \cos(2bx + 2a) + b) \sin(2bx + 2a)}$$

input `integrate(cot(b*x+a)^6,x, algorithm="fricas")`

output `-1/15*(23*cos(2*b*x + 2*a)^3 - cos(2*b*x + 2*a)^2 + 15*(b*x*cos(2*b*x + 2*a)^2 - 2*b*x*cos(2*b*x + 2*a) + b*x)*sin(2*b*x + 2*a) - 11*cos(2*b*x + 2*a) + 13)/((b*cos(2*b*x + 2*a)^2 - 2*b*cos(2*b*x + 2*a) + b)*sin(2*b*x + 2*a))`

**Sympy [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \cot^6(a + bx) dx = \begin{cases} -x - \frac{\cot^5(a+bx)}{5b} + \frac{\cot^3(a+bx)}{3b} - \frac{\cot(a+bx)}{b} & \text{for } b \neq 0 \\ x \cot^6(a) & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a)**6,x)`

output `Piecewise((-x - cot(a + b*x)**5/(5*b) + cot(a + b*x)**3/(3*b) - cot(a + b*x)/b, Ne(b, 0)), (x*cot(a)**6, True))`



**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

$$\int \cot^6(a + bx) dx = -\frac{15bx + 15a + \frac{15 \tan(bx+a)^4 - 5 \tan(bx+a)^2 + 3}{\tan(bx+a)^5}}{15b}$$

input `integrate(cot(b*x+a)^6,x, algorithm="maxima")`

output `-1/15*(15*b*x + 15*a + (15*tan(b*x + a)^4 - 5*tan(b*x + a)^2 + 3)/tan(b*x + a)^5)/b`

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 91 vs.  $2(41) = 82$ .

Time = 0.15 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.02

$$\int \cot^6(a + bx) dx = \frac{3 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^5 - 35 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3 - 480bx - 480a - \frac{330 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 - 35 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 + 3}{\tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^5} + 330 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)}{480b}$$

input `integrate(cot(b*x+a)^6,x, algorithm="giac")`

output `1/480*(3*tan(1/2*b*x + 1/2*a)^5 - 35*tan(1/2*b*x + 1/2*a)^3 - 480*b*x - 480*a - (330*tan(1/2*b*x + 1/2*a)^4 - 35*tan(1/2*b*x + 1/2*a)^2 + 3)/tan(1/2*b*x + 1/2*a)^5 + 330*tan(1/2*b*x + 1/2*a))/b`

**Mupad [B] (verification not implemented)**

Time = 9.52 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int \cot^6(a + bx) dx = -x - \frac{\cot(a+bx)^5}{5} - \frac{\cot(a+bx)^3}{3} + \cot(a + bx)$$

input `int(cot(a + b*x)^6,x)`output `- x - (cot(a + b*x) - cot(a + b*x)^3/3 + cot(a + b*x)^5/5)/b`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.84

$$\int \cot^6(a + bx) dx = \frac{-3 \cot (bx + a)^5 + 5 \cot (bx + a)^3 - 15 \cot (bx + a) - 15bx}{15b}$$

input `int(cot(b*x+a)^6,x)`output `( - 3*cot(a + b*x)**5 + 5*cot(a + b*x)**3 - 15*cot(a + b*x) - 15*b*x)/(15*b)`

### 3.7 $\int \cot^7(a + bx) dx$

Optimal result . . . . .	82
Mathematica [A] (verified) . . . . .	82
Rubi [A] (verified) . . . . .	83
Maple [A] (verified) . . . . .	85
Fricas [B] (verification not implemented) . . . . .	86
Sympy [A] (verification not implemented) . . . . .	86
Maxima [A] (verification not implemented) . . . . .	87
Giac [A] (verification not implemented) . . . . .	87
Mupad [B] (verification not implemented) . . . . .	88
Reduce [B] (verification not implemented) . . . . .	88

#### Optimal result

Integrand size = 8, antiderivative size = 58

$$\int \cot^7(a + bx) dx = -\frac{\cot^2(a + bx)}{2b} + \frac{\cot^4(a + bx)}{4b} - \frac{\cot^6(a + bx)}{6b} - \frac{\log(\sin(a + bx))}{b}$$

output `-1/2*cot(b*x+a)^2/b+1/4*cot(b*x+a)^4/b-1/6*cot(b*x+a)^6/b-ln(sin(b*x+a))/b`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \cot^7(a + bx) dx = -\frac{3 \csc^2(a + bx)}{2b} + \frac{3 \csc^4(a + bx)}{4b} - \frac{\csc^6(a + bx)}{6b} - \frac{\log(\sin(a + bx))}{b}$$

input `Integrate[Cot[a + b*x]^7,x]`

output `(-3*Csc[a + b*x]^2)/(2*b) + (3*Csc[a + b*x]^4)/(4*b) - Csc[a + b*x]^6/(6*b) - Log[Sin[a + b*x]]/b`

**Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.875$ , Rules used = {3042, 25, 3954, 25, 3042, 25, 3954, 25, 3042, 25, 3954, 25, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^7(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\tan\left(a + bx + \frac{\pi}{2}\right)^7 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \tan\left(\frac{1}{2}(2a + \pi) + bx\right)^7 dx \\
 & \quad \downarrow \text{3954} \\
 & \int -\cot^5(a + bx) dx - \frac{\cot^6(a + bx)}{6b} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot^5(a + bx) dx - \frac{\cot^6(a + bx)}{6b} \\
 & \quad \downarrow \text{3042} \\
 & -\int -\tan\left(a + bx + \frac{\pi}{2}\right)^5 dx - \frac{\cot^6(a + bx)}{6b} \\
 & \quad \downarrow \text{25} \\
 & \int \tan\left(\frac{1}{2}(2a + \pi) + bx\right)^5 dx - \frac{\cot^6(a + bx)}{6b} \\
 & \quad \downarrow \text{3954} \\
 & -\int -\cot^3(a + bx) dx - \frac{\cot^6(a + bx)}{6b} + \frac{\cot^4(a + bx)}{4b} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
& \int \cot^3(a+bx)dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} \\
& \quad \downarrow \text{3042} \\
& \int -\tan\left(a+bx+\frac{\pi}{2}\right)^3 dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} \\
& \quad \downarrow \text{25} \\
& -\int \tan\left(\frac{1}{2}(2a+\pi)+bx\right)^3 dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} \\
& \quad \downarrow \text{3954} \\
& \int -\cot(a+bx)dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} - \frac{\cot^2(a+bx)}{2b} \\
& \quad \downarrow \text{25} \\
& -\int \cot(a+bx)dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} - \frac{\cot^2(a+bx)}{2b} \\
& \quad \downarrow \text{3042} \\
& -\int -\tan\left(a+bx+\frac{\pi}{2}\right) dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} - \frac{\cot^2(a+bx)}{2b} \\
& \quad \downarrow \text{25} \\
& \int \tan\left(\frac{1}{2}(2a+\pi)+bx\right) dx - \frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} - \frac{\cot^2(a+bx)}{2b} \\
& \quad \downarrow \text{3956} \\
& -\frac{\cot^6(a+bx)}{6b} + \frac{\cot^4(a+bx)}{4b} - \frac{\cot^2(a+bx)}{2b} - \frac{\log(-\sin(a+bx))}{b}
\end{aligned}$$

input

Int[Cot[a + b\*x]^7,x]

output

$$-\frac{1}{2}\cot[a + b*x]^2/b + \cot[a + b*x]^4/(4*b) - \cot[a + b*x]^6/(6*b) - \text{Log}[-\text{Sin}[a + b*x]]/b$$

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d *x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d *x], x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{-\frac{\cot(bx+a)^6}{6} + \frac{\cot(bx+a)^4}{4} - \frac{\cot(bx+a)^2}{2} + \frac{\ln(\cot(bx+a)^2+1)}{2}}{b}$	49
default	$\frac{-\frac{\cot(bx+a)^6}{6} + \frac{\cot(bx+a)^4}{4} - \frac{\cot(bx+a)^2}{2} + \frac{\ln(\cot(bx+a)^2+1)}{2}}{b}$	49
parallelrisc	$\frac{-2 \cot(bx+a)^6 + 3 \cot(bx+a)^4 - 6 \cot(bx+a)^2 - 12 \ln(\tan(bx+a)) + 6 \ln(\sec(bx+a)^2)}{12b}$	57
norman	$\frac{-\frac{1}{6b} + \frac{\tan(bx+a)^2}{4b} - \frac{\tan(bx+a)^4}{2b}}{\tan(bx+a)^6} - \frac{\ln(\tan(bx+a))}{b} + \frac{\ln(1+\tan(bx+a)^2)}{2b}$	71
risc	$ix + \frac{2ia}{b} + \frac{6e^{10i(bx+a)} - 12e^{8i(bx+a)} + \frac{68e^{6i(bx+a)}}{3} - 12e^{4i(bx+a)} + 6e^{2i(bx+a)}}{b(e^{2i(bx+a)} - 1)^6} - \frac{\ln(e^{2i(bx+a)} - 1)}{b}$	104

input `int(cot(b*x+a)^7, x, method=_RETURNVERBOSE)`

output `1/b*(-1/6*cot(b*x+a)^6+1/4*cot(b*x+a)^4-1/2*cot(b*x+a)^2+1/2*ln(cot(b*x+a)^2+1))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 126 vs.  $2(52) = 104$ .

Time = 0.10 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.17

$$\int \cot^7(a + bx) dx$$

$$= \frac{18 \cos(2bx + 2a)^2 - 3(\cos(2bx + 2a)^3 - 3 \cos(2bx + 2a)^2 + 3 \cos(2bx + 2a) - 1) \log\left(-\frac{1}{2} \cos(2bx + 2a) + \frac{1}{2}\right) - 18 \cos(2bx + 2a) + 8}{6(b \cos(2bx + 2a))^3 - 3b \cos(2bx + 2a)^2 + 3b \cos(2bx + 2a) - b}$$

input `integrate(cot(b*x+a)^7,x, algorithm="fricas")`

output `1/6*(18*cos(2*b*x + 2*a)^2 - 3*(cos(2*b*x + 2*a)^3 - 3*cos(2*b*x + 2*a)^2 + 3*cos(2*b*x + 2*a) - 1)*log(-1/2*cos(2*b*x + 2*a) + 1/2) - 18*cos(2*b*x + 2*a) + 8)/(b*cos(2*b*x + 2*a)^3 - 3*b*cos(2*b*x + 2*a)^2 + 3*b*cos(2*b*x + 2*a) - b)`

**Sympy [A] (verification not implemented)**

Time = 0.47 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.38

$$\int \cot^7(a + bx) dx$$

$$= \begin{cases} \tilde{\infty}x & \text{for } a = 0 \wedge b = 0 \\ x \cot^7(a) & \text{for } b = 0 \\ \tilde{\infty}x & \text{for } a = -bx \\ \frac{\log(\tan^2(a+bx)+1)}{2b} - \frac{\log(\tan(a+bx))}{b} - \frac{1}{2b \tan^2(a+bx)} + \frac{1}{4b \tan^4(a+bx)} - \frac{1}{6b \tan^6(a+bx)} & \text{otherwise} \end{cases}$$

input `integrate(cot(b*x+a)**7,x)`

output `Piecewise((zoo*x, Eq(a, 0) & Eq(b, 0)), (x*cot(a)**7, Eq(b, 0)), (zoo*x, Eq(a, -b*x)), (log(tan(a + b*x)**2 + 1)/(2*b) - log(tan(a + b*x))/b - 1/(2*b*tan(a + b*x)**2) + 1/(4*b*tan(a + b*x)**4) - 1/(6*b*tan(a + b*x)**6), True))`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.83

$$\int \cot^7(a + bx) dx = -\frac{\frac{18 \sin^4(bx+a) - 9 \sin^2(bx+a) + 2}{\sin^6(bx+a)} + 6 \log(\sin(bx+a)^2)}{12b}$$

input `integrate(cot(b*x+a)^7,x, algorithm="maxima")`

output `-1/12*((18*sin(b*x + a)^4 - 9*sin(b*x + a)^2 + 2)/sin(b*x + a)^6 + 6*log(sin(b*x + a)^2))/b`

**Giac [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.84

$$\int \cot^7(a + bx) dx = -\frac{\log(|\sin(bx+a)|)}{b} - \frac{18 \sin^4(bx+a) - 9 \sin^2(bx+a) + 2}{12b \sin^6(bx+a)}$$

input `integrate(cot(b*x+a)^7,x, algorithm="giac")`

output `-log(abs(sin(b*x + a)))/b - 1/12*(18*sin(b*x + a)^4 - 9*sin(b*x + a)^2 + 2)/(b*sin(b*x + a)^6)`



**Mupad [B] (verification not implemented)**

Time = 15.65 (sec) , antiderivative size = 340, normalized size of antiderivative = 5.86

$$\int \cot^7(a + bx) dx = x \operatorname{li} - \frac{\ln(e^{a+2i} e^{b+2i} - 1)}{b} + \frac{32}{b(5e^{a+2i+b+2i} - 10e^{a+4i+b+4i} + 10e^{a+6i+b+6i} - 5e^{a+8i+b+8i} + e^{a+10i+b+10i} - 1)} + \frac{32}{3b(1 + 15e^{a+4i+b+4i} - 20e^{a+6i+b+6i} + 15e^{a+8i+b+8i} - 6e^{a+10i+b+10i} + e^{a+12i+b+12i} - 6e^{a+2i+b+2i})} + \frac{6}{b(e^{a+2i+b+2i} - 1)} + \frac{18}{b(1 + e^{a+4i+b+4i} - 2e^{a+2i+b+2i})} + \frac{104}{3b(3e^{a+2i+b+2i} - 3e^{a+4i+b+4i} + e^{a+6i+b+6i} - 1)} + \frac{44}{b(1 + 6e^{a+4i+b+4i} - 4e^{a+6i+b+6i} + e^{a+8i+b+8i} - 4e^{a+2i+b+2i})}$$

input `int(cot(a + b*x)^7,x)`

output

```
x*i - log(exp(a*2i)*exp(b*x*2i) - 1)/b + 32/(b*(5*exp(a*2i + b*x*2i) - 10*exp(a*4i + b*x*4i) + 10*exp(a*6i + b*x*6i) - 5*exp(a*8i + b*x*8i) + exp(a*10i + b*x*10i) - 1)) + 32/(3*b*(15*exp(a*4i + b*x*4i) - 6*exp(a*2i + b*x*2i) - 20*exp(a*6i + b*x*6i) + 15*exp(a*8i + b*x*8i) - 6*exp(a*10i + b*x*10i) + exp(a*12i + b*x*12i) + 1)) + 6/(b*(exp(a*2i + b*x*2i) - 1)) + 18/(b*(exp(a*4i + b*x*4i) - 2*exp(a*2i + b*x*2i) + 1)) + 104/(3*b*(3*exp(a*2i + b*x*2i) - 3*exp(a*4i + b*x*4i) + exp(a*6i + b*x*6i) - 1)) + 44/(b*(6*exp(a*4i + b*x*4i) - 4*exp(a*2i + b*x*2i) - 4*exp(a*6i + b*x*6i) + exp(a*8i + b*x*8i) + 1))
```

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.53

$$\int \cot^7(a + bx) dx = \frac{48 \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2 + 1\right) \sin(bx + a)^6 - 48 \log\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \sin(bx + a)^6 + 25 \sin(bx + a)^6 - 72 \sin(bx + a)^6}{48 \sin(bx + a)^6 b}$$

input `int(cot(b*x+a)^7,x)`

output `(48*log(tan((a + b*x)/2)**2 + 1)*sin(a + b*x)**6 - 48*log(tan((a + b*x)/2)  
)*sin(a + b*x)**6 + 25*sin(a + b*x)**6 - 72*sin(a + b*x)**4 + 36*sin(a + b  
*x)**2 - 8)/(48*sin(a + b*x)**6*b)`

### 3.8 $\int \cot^8(a + bx) dx$

Optimal result . . . . .	90
Mathematica [C] (verified) . . . . .	90
Rubi [A] (verified) . . . . .	91
Maple [A] (verified) . . . . .	93
Fricas [B] (verification not implemented) . . . . .	93
Sympy [A] (verification not implemented) . . . . .	94
Maxima [A] (verification not implemented) . . . . .	94
Giac [B] (verification not implemented) . . . . .	95
Mupad [B] (verification not implemented) . . . . .	95
Reduce [B] (verification not implemented) . . . . .	96

#### Optimal result

Integrand size = 8, antiderivative size = 57

$$\int \cot^8(a + bx) dx = x + \frac{\cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} + \frac{\cot^5(a + bx)}{5b} - \frac{\cot^7(a + bx)}{7b}$$

output `x+cot(b*x+a)/b-1/3*cot(b*x+a)^3/b+1/5*cot(b*x+a)^5/b-1/7*cot(b*x+a)^7/b`

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.58

$$\int \cot^8(a + bx) dx = -\frac{\cot^7(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{2}, 1, -\frac{5}{2}, -\tan^2(a + bx)\right)}{7b}$$

input `Integrate[Cot[a + b*x]^8,x]`

output `-1/7*(Cot[a + b*x]^7*Hypergeometric2F1[-7/2, 1, -5/2, -Tan[a + b*x]^2])/b`

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$ , Rules used = {3042, 3954, 3042, 3954, 3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^8(a + bx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(a + bx + \frac{\pi}{2}\right)^8 dx \\
 & \quad \downarrow \text{3954} \\
 & - \int \cot^6(a + bx) dx - \frac{\cot^7(a + bx)}{7b} \\
 & \quad \downarrow \text{3042} \\
 & - \int \tan\left(a + bx + \frac{\pi}{2}\right)^6 dx - \frac{\cot^7(a + bx)}{7b} \\
 & \quad \downarrow \text{3954} \\
 & \int \cot^4(a + bx) dx - \frac{\cot^7(a + bx)}{7b} + \frac{\cot^5(a + bx)}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(a + bx + \frac{\pi}{2}\right)^4 dx - \frac{\cot^7(a + bx)}{7b} + \frac{\cot^5(a + bx)}{5b} \\
 & \quad \downarrow \text{3954} \\
 & - \int \cot^2(a + bx) dx - \frac{\cot^7(a + bx)}{7b} + \frac{\cot^5(a + bx)}{5b} - \frac{\cot^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3042} \\
 & - \int \tan\left(a + bx + \frac{\pi}{2}\right)^2 dx - \frac{\cot^7(a + bx)}{7b} + \frac{\cot^5(a + bx)}{5b} - \frac{\cot^3(a + bx)}{3b} \\
 & \quad \downarrow \text{3954}
 \end{aligned}$$

$$\int 1 dx - \frac{\cot^7(a+bx)}{7b} + \frac{\cot^5(a+bx)}{5b} - \frac{\cot^3(a+bx)}{3b} + \frac{\cot(a+bx)}{b}$$

↓ 24

$$-\frac{\cot^7(a+bx)}{7b} + \frac{\cot^5(a+bx)}{5b} - \frac{\cot^3(a+bx)}{3b} + \frac{\cot(a+bx)}{b} + x$$

input `Int[Cot[a + b*x]^8,x]`

output `x + Cot[a + b*x]/b - Cot[a + b*x]^3/(3*b) + Cot[a + b*x]^5/(5*b) - Cot[a + b*x]^7/(7*b)`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :=> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] :=> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :=> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

**Maple [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.86

method	result	size
parallelrisc	$\frac{-15 \cot(bx+a)^7 + 21 \cot(bx+a)^5 - 35 \cot(bx+a)^3 + 105bx + 105 \cot(bx+a)}{105b}$	49
derivativedivides	$\frac{-\frac{\cot(bx+a)^7}{7} + \frac{\cot(bx+a)^5}{5} - \frac{\cot(bx+a)^3}{3} + \cot(bx+a) - \frac{\pi}{2} + \operatorname{arccot}(\cot(bx+a))}{b}$	52
default	$\frac{-\frac{\cot(bx+a)^7}{7} + \frac{\cot(bx+a)^5}{5} - \frac{\cot(bx+a)^3}{3} + \cot(bx+a) - \frac{\pi}{2} + \operatorname{arccot}(\cot(bx+a))}{b}$	52
norman	$\frac{\frac{\tan(bx+a)^6}{b} + x \tan(bx+a)^7 - \frac{1}{7b} + \frac{\tan(bx+a)^2}{5b} - \frac{\tan(bx+a)^4}{3b}}{\tan(bx+a)^7}$	64
risc	$x + \frac{8i(105 e^{12i(bx+a)} - 315 e^{10i(bx+a)} + 770 e^{8i(bx+a)} - 770 e^{6i(bx+a)} + 609 e^{4i(bx+a)} - 203 e^{2i(bx+a)} + 44)}{105b(e^{2i(bx+a)} - 1)^7}$	90

input `int (cot (b*x+a)^8,x,method=_RETURNVERBOSE)`output `1/105*(-15*cot (b*x+a)^7+21*cot (b*x+a)^5-35*cot (b*x+a)^3+105*b*x+105*cot (b*x+a))/b`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(51) = 102.

Time = 0.09 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.95

$$\int \cot^8(a + bx) dx$$

$$= \frac{176 \cos(2bx + 2a)^4 - 108 \cos(2bx + 2a)^3 + 20 \cos(2bx + 2a)^2 + 105 (bx \cos(2bx + 2a)^3 - 3bx \cos(2bx + 2a))}{105 (b \cos(2bx + 2a)^3 - 3b \cos(2bx + 2a)^2 + 3b \cos(2bx + 2a))}$$

input `integrate(cot (b*x+a)^8,x, algorithm="fricas")`

output

```
1/105*(176*cos(2*b*x + 2*a)^4 - 108*cos(2*b*x + 2*a)^3 + 20*cos(2*b*x + 2*
a)^2 + 105*(b*x*cos(2*b*x + 2*a)^3 - 3*b*x*cos(2*b*x + 2*a)^2 + 3*b*x*cos(
2*b*x + 2*a) - b*x)*sin(2*b*x + 2*a) + 228*cos(2*b*x + 2*a) - 76)/((b*cos(
2*b*x + 2*a)^3 - 3*b*cos(2*b*x + 2*a)^2 + 3*b*cos(2*b*x + 2*a) - b)*sin(2*
b*x + 2*a))
```

**Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.89

$$\int \cot^8(a + bx) dx = \begin{cases} x - \frac{\cot^7(a+bx)}{7b} + \frac{\cot^5(a+bx)}{5b} - \frac{\cot^3(a+bx)}{3b} + \frac{\cot(a+bx)}{b} & \text{for } b \neq 0 \\ x \cot^8(a) & \text{otherwise} \end{cases}$$

input

```
integrate(cot(b*x+a)**8,x)
```

output

```
Piecewise((x - cot(a + b*x)**7/(7*b) + cot(a + b*x)**5/(5*b) - cot(a + b*x)
)**3/(3*b) + cot(a + b*x)/b, Ne(b, 0)), (x*cot(a)**8, True))
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.95

$$\int \cot^8(a + bx) dx = \frac{105 bx + 105 a + \frac{105 \tan(bx+a)^6 - 35 \tan(bx+a)^4 + 21 \tan(bx+a)^2 - 15}{\tan(bx+a)^7}}{105 b}$$

input

```
integrate(cot(b*x+a)^8,x, algorithm="maxima")
```

output

```
1/105*(105*b*x + 105*a + (105*tan(b*x + a)^6 - 35*tan(b*x + a)^4 + 21*tan(
b*x + a)^2 - 15)/tan(b*x + a)^7)/b
```

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 116 vs.  $2(51) = 102$ .

Time = 0.16 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.04

$$\int \cot^8(a + bx) dx$$

$$= \frac{15 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^7 - 189 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^5 + 1295 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3 + 13440bx + 13440a + \frac{9765 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^6 - 1295 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + 189 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - 15}{13440b}$$

input `integrate(cot(b*x+a)^8,x, algorithm="giac")`

output

```
1/13440*(15*tan(1/2*b*x + 1/2*a)^7 - 189*tan(1/2*b*x + 1/2*a)^5 + 1295*tan
(1/2*b*x + 1/2*a)^3 + 13440*b*x + 13440*a + (9765*tan(1/2*b*x + 1/2*a)^6 -
1295*tan(1/2*b*x + 1/2*a)^4 + 189*tan(1/2*b*x + 1/2*a)^2 - 15)/tan(1/2*b*
x + 1/2*a)^7 - 9765*tan(1/2*b*x + 1/2*a))/b
```

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.75

$$\int \cot^8(a + bx) dx = x + \frac{-\frac{\cot(a+bx)^7}{7} + \frac{\cot(a+bx)^5}{5} - \frac{\cot(a+bx)^3}{3} + \cot(a+bx)}{b}$$

input `int(cot(a + b*x)^8,x)`

output

```
x + (cot(a + b*x) - cot(a + b*x)^3/3 + cot(a + b*x)^5/5 - cot(a + b*x)^7/7
)/b
```



**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.84

$$\int \cot^8(a + bx) dx$$
$$= \frac{-15 \cot (bx + a)^7 + 21 \cot (bx + a)^5 - 35 \cot (bx + a)^3 + 105 \cot (bx + a) + 105bx}{105b}$$

input `int(cot(b*x+a)^8,x)`

output `( - 15*cot(a + b*x)**7 + 21*cot(a + b*x)**5 - 35*cot(a + b*x)**3 + 105*cot(a + b*x) + 105*b*x)/(105*b)`

### 3.9 $\int (c \cot(a + bx))^{7/2} dx$

Optimal result	97
Mathematica [A] (verified)	98
Rubi [A] (warning: unable to verify)	98
Maple [A] (verified)	104
Fricas [B] (verification not implemented)	104
Sympy [F]	105
Maxima [A] (verification not implemented)	105
Giac [F]	106
Mupad [B] (verification not implemented)	106
Reduce [F]	107

#### Optimal result

Integrand size = 12, antiderivative size = 177

$$\int (c \cot(a + bx))^{7/2} dx = \frac{c^{7/2} \arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} - \frac{c^{7/2} \arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} - \frac{c^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}b} + \frac{2c^3 \sqrt{c \cot(a + bx)}}{b} - \frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

output

```
1/2*c^(7/2)*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b-1/2*c
^(7/2)*arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b-1/2*c^(7/2)
)*arctanh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a)))*2^(1/
2)/b+2*c^3*(c*cot(b*x+a))^(1/2)/b-2/5*c*(c*cot(b*x+a))^(5/2)/b
```

**Mathematica [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.99

$$\int (c \cot(a + bx))^{7/2} dx = \frac{(c \cot(a + bx))^{7/2} \left( -\frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\cot(a+bx)}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{\arctan\left(\frac{1 + \sqrt{2}\sqrt{\cot(a+bx)}}{\sqrt{2}}\right)}{\sqrt{2}} - 2\sqrt{\cot(a + bx)} + \frac{2}{5} \cot^{5/2}(a + bx) - \dots \right)}{b \cot^{7/2}(a + bx)}$$

input

```
Integrate[(c*Cot[a + b*x])^(7/2),x]
```

output

```
-(((c*Cot[a + b*x])^(7/2)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[a + b*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[a + b*x]]]/Sqrt[2] - 2*Sqrt[Cot[a + b*x]] + (2*Cot[a + b*x]^(5/2))/5 - Log[1 - Sqrt[2]*Sqrt[Cot[a + b*x]] + Cot[a + b*x]]/(2*Sqrt[2]) + Log[1 + Sqrt[2]*Sqrt[Cot[a + b*x]] + Cot[a + b*x]]/(2*Sqrt[2])))/(b*Cot[a + b*x]^(7/2)))
```

**Rubi [A] (warning: unable to verify)**

Time = 0.57 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.28, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$ , Rules used = {3042, 3954, 3042, 3954, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c \cot(a + bx))^{7/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \left( -c \tan\left(a + bx + \frac{\pi}{2}\right) \right)^{7/2} dx \\ & \quad \downarrow \text{3954} \\ & -c^2 \int (c \cot(a + bx))^{3/2} dx - \frac{2c(c \cot(a + bx))^{5/2}}{5b} \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& -c^2 \int \left( -c \tan \left( a + bx + \frac{\pi}{2} \right) \right)^{3/2} dx - \frac{2c(c \cot(a + bx))^{5/2}}{5b} \\
& \downarrow 3954 \\
& -c^2 \left( c^2 \left( - \int \frac{1}{\sqrt{c \cot(a + bx)}} dx \right) - \frac{2c\sqrt{c \cot(a + bx)}}{b} \right) - \frac{2c(c \cot(a + bx))^{5/2}}{5b} \\
& \downarrow 3042 \\
& -c^2 \left( c^2 \left( - \int \frac{1}{\sqrt{-c \tan \left( a + bx + \frac{\pi}{2} \right)}} dx \right) - \frac{2c\sqrt{c \cot(a + bx)}}{b} \right) - \frac{2c(c \cot(a + bx))^{5/2}}{5b} \\
& \downarrow 3957 \\
& -c^2 \left( \frac{c^3 \int \frac{1}{\sqrt{c \cot(a + bx)(\cot^2(a + bx)c^2 + c^2)}} d(c \cot(a + bx))}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \right) - \\
& \quad \frac{2c(c \cot(a + bx))^{5/2}}{5b} \\
& \downarrow 266 \\
& -c^2 \left( \frac{2c^3 \int \frac{1}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \right) - \frac{2c(c \cot(a + bx))^{5/2}}{5b} \\
& \downarrow 755 \\
& -c^2 \left( \frac{2c^3 \left( \frac{\int \frac{c - c^2 \cot^2(a + bx)}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{2c} + \frac{\int \frac{c^2 \cot^2(a + bx) + c}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{2c} \right)}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \right) - \\
& \quad \frac{2c(c \cot(a + bx))^{5/2}}{5b} \\
& \downarrow 1476
\end{aligned}$$

$$-c^2 \left( \frac{2c^3 \left( \frac{\frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)} \right) + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx) + c^2} d\sqrt{c \cot(a+bx)}}{2c}}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

1082

$$-c^2 \left( \frac{2c^3 \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} d(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} d(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} \right) + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx) + c^2} d\sqrt{c \cot(a+bx)}}{2c}}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

217

$$-c^2 \left( \frac{2c^3 \left( \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx) + c^2} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1) - \arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

1479

$$-c^2 \left( \frac{2c^3 \left( \frac{\int \frac{\sqrt{2}\sqrt{c} - 2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{2}(\sqrt{c} + \sqrt{2}\sqrt{c \cot(a+bx)})}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} \right) + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}}}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

25

$$-c^2 \left( \frac{2c^3 \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right)}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

↓ 27

$$-c^2 \left( \frac{2c^3 \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right)}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

↓ 1103

$$-c^2 \left( \frac{2c^3 \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c}\cot(a+bx))}{\sqrt{2}\sqrt{c}} + \frac{\log(\sqrt{2}c^{3/2}\cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(-\sqrt{2}c^{3/2}\cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} \right)}{b} \right)$$

$$\frac{2c(c \cot(a + bx))^{5/2}}{5b}$$

input `Int[(c*Cot[a + b*x])^(7/2), x]`

output

$$\begin{aligned} & (-2*c*(c*\cot[a + b*x])^{5/2})/(5*b) - c^2*((-2*c*\sqrt{c*\cot[a + b*x]})/b + \\ & (2*c^3*((-\arctan[1 - \sqrt{2}*\sqrt{c}*\cot[a + b*x]]/(\sqrt{2}*\sqrt{c})) + \\ & \arctan[1 + \sqrt{2}*\sqrt{c}*\cot[a + b*x]]/(\sqrt{2}*\sqrt{c}))/2*c) + (-1/2* \\ & \log[c - \sqrt{2}*c^{3/2}*\cot[a + b*x] + c^2*\cot[a + b*x]^2/(\sqrt{2}*\sqrt{c} \\ & )] + \log[c + \sqrt{2}*c^{3/2}*\cot[a + b*x] + c^2*\cot[a + b*x]^2/(2*\sqrt{2} \\ & *\sqrt{c}))/2*c))/b \end{aligned}$$

### Definitions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x\_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$$

rule 217

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{-1} * \arctan[\text{Rt}[-\text{b}, 2]*(x/\text{Rt}[-\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{LtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$$

rule 266

$$\text{Int}[(\text{c}_)*(x_)^m * (\text{a}_) + (\text{b}_)*(x_)^2)^p, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(\text{a} + \text{b}*x^{2*k}/\text{c}^2))^p, \text{x}], \text{x}, (\text{c}*x)^{1/k}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \ \&\& \ \text{FractionQ}[\text{m}] \ \&\& \ \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$$

rule 755

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_)^4)^{-1}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(\text{r} - \text{s}*x^2)/(\text{a} + \text{b}*x^4)], \text{x}], \text{x}] + \text{Simp}[1/(2*r) \quad \text{Int}[(\text{r} + \text{s}*x^2)/(\text{a} + \text{b}*x^4)], \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ (\text{GtQ}[\text{a}/\text{b}, 0] \ || \ (\text{PosQ}[\text{a}/\text{b}] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{a}]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{b}]])$$

rule 1082

$$\text{Int}[(\text{a}_) + (\text{b}_)*(x_) + (\text{c}_)*(x_)^2)^{-1}, \text{x\_Symbol}] \rightarrow \text{With}[\{\text{q} = 1 - 4*\text{Simplify}[\text{a}*(\text{c}/\text{b}^2)]\}, \text{Simp}[-2/\text{b} \quad \text{Subst}[\text{Int}[1/(\text{q} - x^2)], \text{x}], \text{x}, 1 + 2*c*(x/\text{b})], \text{x}] \text{ ; RationalQ}[\text{q}] \ \&\& \ (\text{EqQ}[\text{q}^2, 1] \ || \ \text{!RationalQ}[\text{b}^2 - 4*\text{a}*c]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$$

rule 1103  $\text{Int}[\frac{(d_.) + (e_.)x}{(a_.) + (b_.)x + (c_.)x^2}, x\_Symbol] \rightarrow \text{Simp}[d \cdot \text{Log}[\text{RemoveContent}[a + b x + c x^2, x]]/b, x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 c d - b^2 e, 0]$

rule 1476  $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x\_Symbol] \rightarrow \text{With}\{q = \text{Rt}[2(d/e), 2]\}, \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e + q x + x^2, x], x], x] + \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e - q x + x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c d^2 - a e^2, 0] \ \&\& \ \text{PosQ}[d e]$

rule 1479  $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x\_Symbol] \rightarrow \text{With}\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2c q) \text{Int}[(q - 2x)/\text{Simp}[d/e + q x - x^2, x], x], x] + \text{Simp}[e/(2c q) \text{Int}[(q + 2x)/\text{Simp}[d/e - q x - x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c d^2 - a e^2, 0] \ \&\& \ \text{NegQ}[d e]$

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3954  $\text{Int}[(b_.) \tan[(c_.) + (d_.)x]^n, x\_Symbol] \rightarrow \text{Simp}[b \cdot ((b \cdot \tan[c + d x])^{n-1} / (d \cdot (n-1))), x] - \text{Simp}[b^2 \text{Int}[(b \cdot \tan[c + d x])^{n-2}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1]$

rule 3957  $\text{Int}[(b_.) \tan[(c_.) + (d_.)x]^n, x\_Symbol] \rightarrow \text{Simp}[b/d \text{Subst}[\text{Int}[x^n / (b^2 + x^2), x], x, b \cdot \tan[c + d x]], x] /; \text{FreeQ}\{b, c, d, n\}, x] \ \&\& \ \text{!IntegerQ}[n]$



**Maple [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.95

method	result
derivativedivides	$2c \left( \frac{(c \cot(bx+a))^{\frac{5}{2}}}{5} - c^2 \sqrt{c \cot(bx+a)} + \frac{c^2 (c^2)^{\frac{1}{4}} \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right) \right)}{8} \right)$
default	$2c \left( \frac{(c \cot(bx+a))^{\frac{5}{2}}}{5} - c^2 \sqrt{c \cot(bx+a)} + \frac{c^2 (c^2)^{\frac{1}{4}} \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right) \right)}{8} \right)$

input `int((c*cot(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

output 
$$-2/b*c*(1/5*(c*\cot(b*x+a))^{5/2}-c^2*(c*\cot(b*x+a))^{1/2}+1/8*c^2*(c^2)^{1/4}*2^{1/2}*(\ln((c*\cot(b*x+a)+(c^2)^{1/4}*(c*\cot(b*x+a))^{1/2}*2^{1/2}+(c^2)^{1/2}))/((c*\cot(b*x+a)-(c^2)^{1/4}*(c*\cot(b*x+a))^{1/2}*2^{1/2}+(c^2)^{1/2}))+2*\arctan(2^{1/2}/(c^2)^{1/4}*(c*\cot(b*x+a))^{1/2}+1)-2*\arctan(-2^{1/2}/((c^2)^{1/4}*(c*\cot(b*x+a))^{1/2}+1)))$$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 426 vs. 2(140) = 280.

Time = 0.10 (sec) , antiderivative size = 426, normalized size of antiderivative = 2.41

$$\int (c \cot(a + bx))^{7/2} dx =$$

$$10 \sqrt{2} (c^3 \cos(2bx + 2a) - c^3) \sqrt{c} \arctan \left( \frac{\sqrt{2} \sqrt{c} \sqrt{\frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)}} + c}{c} \right) + 10 \sqrt{2} (c^3 \cos(2bx + 2a) - c^3) \sqrt{c} \arctan \left( \frac{\sqrt{2} \sqrt{c} \sqrt{\frac{c \cos(2bx + 2a) - c}{\sin(2bx + 2a)}} + c}{c} \right)$$

input `integrate((c*cot(b*x+a))^(7/2),x, algorithm="fricas")`

output

```
-1/20*(10*sqrt(2)*(c^3*cos(2*b*x + 2*a) - c^3)*sqrt(c)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) + c)/c) + 10*sqrt(2)*(c^3*cos(2*b*x + 2*a) - c^3)*sqrt(c)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) - c)/c) + 5*sqrt(2)*(c^3*cos(2*b*x + 2*a) - c^3)*sqrt(c)*log((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) + c*cos(2*b*x + 2*a) + c*sin(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) - 5*sqrt(2)*(c^3*cos(2*b*x + 2*a) - c^3)*sqrt(c)*log(-(sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) - c*cos(2*b*x + 2*a) - c*sin(2*b*x + 2*a) - c)/sin(2*b*x + 2*a)) - 16*(3*c^3*cos(2*b*x + 2*a) - 2*c^3)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)))/(b*cos(2*b*x + 2*a) - b)
```

**Sympy [F]**

$$\int (c \cot(a + bx))^{7/2} dx = \int (c \cot(a + bx))^{\frac{7}{2}} dx$$

input

```
integrate((c*cot(b*x+a))**(7/2),x)
```

output

```
Integral((c*cot(a + b*x))**(7/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.11

$$\int (c \cot(a + bx))^{7/2} dx =$$

$$\left( 10 \sqrt{2} c^{\frac{5}{2}} \arctan \left( \frac{\sqrt{2} \left( \sqrt{2} \sqrt{c} + 2 \sqrt{\frac{c}{\tan(bx+a)}} \right)}{2 \sqrt{c}} \right) \right) + 10 \sqrt{2} c^{\frac{5}{2}} \arctan \left( -\frac{\sqrt{2} \left( \sqrt{2} \sqrt{c} - 2 \sqrt{\frac{c}{\tan(bx+a)}} \right)}{2 \sqrt{c}} \right) + 5 \sqrt{2} c^{\frac{5}{2}} \log \left( \dots \right)$$

input

```
integrate((c*cot(b*x+a))^(7/2),x, algorithm="maxima")
```

output

```
-1/20*(10*sqrt(2)*c^(5/2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(b*x + a)))/sqrt(c)) + 10*sqrt(2)*c^(5/2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c)) + 5*sqrt(2)*c^(5/2)*log(sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a)) - 5*sqrt(2)*c^(5/2)*log(-sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a)) - 40*c^2*sqrt(c/tan(b*x + a)) + 8*(c/tan(b*x + a))^(5/2))*c/b
```

**Giac [F]**

$$\int (c \cot(a + bx))^{7/2} dx = \int (c \cot(bx + a))^{7/2} dx$$

input

```
integrate((c*cot(b*x+a))^(7/2),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(7/2), x)
```

**Mupad [B] (verification not implemented)**

Time = 9.40 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.51

$$\int (c \cot(a + bx))^{7/2} dx = \frac{2c^3 \sqrt{c \cot(a + bx)}}{b} - \frac{2c (c \cot(a + bx))^{5/2}}{5b} + \frac{(-1)^{1/4} c^{7/2} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right) \operatorname{li}}{b} + \frac{(-1)^{1/4} c^{7/2} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)} \operatorname{li}}{\sqrt{c}}\right)}{b}$$

input

```
int((c*cot(a + b*x))^(7/2),x)
```

output

```
(2*c^3*(c*cot(a + b*x))^(1/2))/b - (2*c*(c*cot(a + b*x))^(5/2))/(5*b) + ((-1)^(1/4)*c^(7/2)*atan(((1/4)*(-1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2))*li)/b + ((-1)^(1/4)*c^(7/2)*atan(((1/4)*(-1)^(1/4)*(c*cot(a + b*x))^(1/2)*li)/c^(1/2)))/b
```

**Reduce [F]**

$$\int (c \cot(a + bx))^{7/2} dx = \frac{\sqrt{c} c^3 \left( -2\sqrt{\cot(bx + a)} \cot(bx + a)^2 + 10\sqrt{\cot(bx + a)} + 5 \left( \int \frac{\sqrt{\cot(bx+a)}}{\cot(bx+a)} dx \right) b \right)}{5b}$$

input `int((c*cot(b*x+a))^(7/2),x)`

output `(sqrt(c)*c**3*( - 2*sqrt(cot(a + b*x))*cot(a + b*x)**2 + 10*sqrt(cot(a + b*x)) + 5*int(sqrt(cot(a + b*x))/cot(a + b*x),x)*b))/(5*b)`

### 3.10 $\int (c \cot(a + bx))^{5/2} dx$

Optimal result	108
Mathematica [A] (verified)	109
Rubi [A] (warning: unable to verify)	109
Maple [A] (verified)	113
Fricas [B] (verification not implemented)	114
Sympy [F]	115
Maxima [A] (verification not implemented)	115
Giac [F]	116
Mupad [B] (verification not implemented)	116
Reduce [F]	117

#### Optimal result

Integrand size = 12, antiderivative size = 157

$$\int (c \cot(a + bx))^{5/2} dx = -\frac{c^{5/2} \arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} + \frac{c^{5/2} \arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} - \frac{c^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

output

```
-1/2*c^(5/2)*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b+1/2*c^(5/2)*arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b-1/2*c^(5/2)*arctanh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a)))*2^(1/2)/b-2/3*c*(c*cot(b*x+a))^(3/2)/b
```

**Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.64

$$\int (c \cot(a + bx))^{5/2} dx = \frac{c(c \cot(a + bx))^{3/2} \left( -3 \arctan \left( \sqrt[4]{-\cot^2(a + bx)} \right) \sqrt[4]{-\cot(a + bx)} + 3 \operatorname{arctanh} \left( \sqrt[4]{-\cot^2(a + bx)} \right) \sqrt[4]{-\cot(a + bx)} \right)}{3b \cot^{7/4}(a + bx)}$$

input

```
Integrate[(c*Cot[a + b*x])^(5/2),x]
```

output

```
-1/3*(c*(c*Cot[a + b*x])^(3/2)*(-3*ArcTan[(-Cot[a + b*x]^2)^(1/4)]*(-Cot[a + b*x])^(1/4) + 3*ArcTanh[(-Cot[a + b*x]^2)^(1/4)]*(-Cot[a + b*x])^(1/4) + 2*Cot[a + b*x]^(7/4)))/(b*Cot[a + b*x]^(7/4))
```

**Rubi [A] (warning: unable to verify)**

Time = 0.48 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.25, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$ , Rules used = {3042, 3954, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c \cot(a + bx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \left( -c \tan \left( a + bx + \frac{\pi}{2} \right) \right)^{5/2} dx \\ & \quad \downarrow \text{3954} \\ & c^2 \left( - \int \sqrt{c \cot(a + bx)} dx \right) - \frac{2c(c \cot(a + bx))^{3/2}}{3b} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$c^2 \left( - \int \sqrt{-c \tan \left( a + bx + \frac{\pi}{2} \right)} dx \right) - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 3957

$$\frac{c^3 \int \frac{\sqrt{c \cot(a+bx)}}{\cot^2(a+bx)c^2+c^2} d(c \cot(a + bx))}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 266

$$\frac{2c^3 \int \frac{c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)}}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 826

$$\frac{2c^3 \left( \frac{1}{2} \int \frac{c^2 \cot^2(a+bx)+c}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)} - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)} \right)}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 1476

$$\frac{2c^3 \left( \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a + bx)} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a + bx)} \right)}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 1082

$$\frac{2c^3 \left( \frac{1}{2} \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx)-1} d(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx)-1} d(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)} \right)}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 217

$$\frac{2c^3 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)} \right)}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

↓ 1479

$$2c^3 \left( \frac{1}{2} \left( \frac{\int -\frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int -\frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \arctan\left(\frac{\sqrt{2}\sqrt{c}\cot(a+bx)+1}{\sqrt{2}\sqrt{c}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{c}\cot(a+bx)}{\sqrt{2}\sqrt{c}}\right) \right) \right)$$

$$\frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

25

$$2c^3 \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \arctan\left(\frac{\sqrt{2}\sqrt{c}\cot(a+bx)+1}{\sqrt{2}\sqrt{c}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{c}\cot(a+bx)}{\sqrt{2}\sqrt{c}}\right) \right) \right)$$

$$\frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

27

$$2c^3 \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{c}} \right) + \frac{1}{2} \left( \arctan\left(\frac{\sqrt{2}\sqrt{c}\cot(a+bx)+1}{\sqrt{2}\sqrt{c}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{c}\cot(a+bx)}{\sqrt{2}\sqrt{c}}\right) \right) \right)$$

$$\frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

1103

$$2c^3 \left( \frac{1}{2} \left( \frac{\arctan\left(\frac{\sqrt{2}\sqrt{c}\cot(a+bx)+1}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}\sqrt{c}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{c}\cot(a+bx)}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\log\left(\frac{-\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c}{2\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}\sqrt{c}} - \frac{\log\left(\frac{\sqrt{2}c^3 \cot(a+bx)+c^2 \cot^2(a+bx)+c}{2\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}\sqrt{c}} \right) \right)$$

$$\frac{2c(c \cot(a + bx))^{3/2}}{3b}$$

input `Int[(c*Cot[a + b*x])^(5/2),x]`

output `(-2*c*(c*Cot[a + b*x])^(3/2))/(3*b) + (2*c^3*((-(ArcTan[1 - Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c])) + ArcTan[1 + Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c]))/2 + (Log[c - Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]) - Log[c + Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]))/2)/b`



## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.98

method	result
derivativedivides	$2c \left( \frac{(c \cot(bx+a))^{\frac{3}{2}}}{3} - \frac{c^2 \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right)}{8 (c^2)^{\frac{1}{4}}} \right)$
default	$2c \left( \frac{(c \cot(bx+a))^{\frac{3}{2}}}{3} - \frac{c^2 \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right)}{8 (c^2)^{\frac{1}{4}}} \right)$

```
input int((c*cot(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/b*c*(1/3*(c*cot(b*x+a))^(3/2)-1/8*c^2/(c^2)^(1/4)*2^(1/2)*(ln((c*cot(b*x+a)-(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))/(c*cot(b*x+a)+(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))+2*arctan(2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)-2*arctan(-2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 380 vs. 2(122) = 244.

Time = 0.12 (sec) , antiderivative size = 380, normalized size of antiderivative = 2.42

$$\int (c \cot(a + bx))^{5/2} dx = \frac{6 \sqrt{2} c^{\frac{5}{2}} \arctan \left( \frac{\sqrt{2} \sqrt{c} \sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}} + c}{c} \right) \sin(2bx+2a) + 6 \sqrt{2} c^{\frac{5}{2}} \arctan \left( \frac{\sqrt{2} \sqrt{c} \sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}}}{c} \right)}{b}$$

```
input integrate((c*cot(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
1/12*(6*sqrt(2)*c^(5/2)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) + c)/c)*sin(2*b*x + 2*a) + 6*sqrt(2)*c^(5/2)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) - c)/c)*sin(2*b*x + 2*a) - 3*sqrt(2)*c^(5/2)*log((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) + c*cos(2*b*x + 2*a) + c*sin(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) + 3*sqrt(2)*c^(5/2)*log(-(sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) - c*cos(2*b*x + 2*a) - c*sin(2*b*x + 2*a) - c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) - 8*(c^2*cos(2*b*x + 2*a) + c^2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)))/(b*sin(2*b*x + 2*a))
```

### Sympy [F]

$$\int (c \cot(a + bx))^{5/2} dx = \int (c \cot(a + bx))^{\frac{5}{2}} dx$$

input

```
integrate((c*cot(b*x+a))**(5/2), x)
```

output

```
Integral((c*cot(a + b*x))**(5/2), x)
```

### Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.18

$$\int (c \cot(a + bx))^{5/2} dx = \frac{3c^2 \left( \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}\sqrt{c} + 2\sqrt{\tan\frac{c}{bx+a}})}{2\sqrt{c}}\right)}{\sqrt{c}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}\sqrt{c} - 2\sqrt{\tan\frac{c}{bx+a}})}{2\sqrt{c}}\right)}{\sqrt{c}} \right) - \sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\tan\frac{c}{bx+a}}\right)}{12b}$$

input

```
integrate((c*cot(b*x+a))^(5/2), x, algorithm="maxima")
```

output

```
1/12*(3*c^2*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(
b*x + a)))/sqrt(c))/sqrt(c) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(
c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c))/sqrt(c) - sqrt(2)*log(sqrt(2)*sqrt(c)
)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c) + sqrt(2)*log(-sqrt(2)
)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c) - 8*(c/tan(b
*x + a))^(3/2))*c/b
```

**Giac [F]**

$$\int (c \cot(a + bx))^{5/2} dx = \int (c \cot(bx + a))^{5/2} dx$$

input

```
integrate((c*cot(b*x+a))^(5/2),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(5/2), x)
```

**Mupad [B] (verification not implemented)**

Time = 9.03 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.47

$$\int (c \cot(a + bx))^{5/2} dx = \frac{(-1)^{1/4} c^{5/2} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right)}{b} - \frac{2c(c \cot(a + bx))^{3/2}}{3b} - \frac{(-1)^{1/4} c^{5/2} \operatorname{atanh}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right)}{b}$$

input

```
int((c*cot(a + b*x))^(5/2),x)
```

output

```
((-1)^(1/4)*c^(5/2)*atan((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2)))/b -
(2*c*(c*cot(a + b*x))^(3/2))/(3*b) - ((-1)^(1/4)*c^(5/2)*atanh((( -1)^(1/4)
)*(c*cot(a + b*x))^(1/2))/c^(1/2)))/b
```

**Reduce [F]**

$$\int (c \cot(a + bx))^{5/2} dx = \sqrt{c} \left( \int \sqrt{\cot(bx + a)} \cot(bx + a)^2 dx \right) c^2$$

input `int((c*cot(b*x+a))^(5/2),x)`

output `sqrt(c)*int(sqrt(cot(a + b*x))*cot(a + b*x)**2,x)*c**2`

### 3.11 $\int (c \cot(a + bx))^{3/2} dx$

Optimal result . . . . .	118
Mathematica [A] (verified) . . . . .	119
Rubi [A] (warning: unable to verify) . . . . .	119
Maple [A] (verified) . . . . .	124
Fricas [B] (verification not implemented) . . . . .	124
Sympy [F] . . . . .	125
Maxima [A] (verification not implemented) . . . . .	125
Giac [F] . . . . .	126
Mupad [B] (verification not implemented) . . . . .	126
Reduce [F] . . . . .	127

#### Optimal result

Integrand size = 12, antiderivative size = 154

$$\int (c \cot(a + bx))^{3/2} dx = -\frac{c^{3/2} \arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} + \frac{c^{3/2} \arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} + \frac{c^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}b} - \frac{2c\sqrt{c \cot(a + bx)}}{b}$$

output

```
-1/2*c^(3/2)*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b+1/2*c^(3/2)*arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b+1/2*c^(3/2)*arctanh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a)))*2^(1/2)/b-2*c*(c*cot(b*x+a))^(1/2)/b
```

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.05

$$\int (c \cot(a + bx))^{3/2} dx = \frac{(c \cot(a + bx))^{3/2} \left( \frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\cot(a+bx)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1 + \sqrt{2}\sqrt{\cot(a+bx)}}{\sqrt{2}}\right)}{\sqrt{2}} + 2\sqrt{\cot(a + bx)} + \frac{\log\left(\frac{1 - \sqrt{2}\sqrt{\cot(a+bx)} + \cot(a + bx)}{2\sqrt{2}}\right)}{2\sqrt{2}} \right)}{b \cot^{\frac{3}{2}}(a + bx)}$$

input `Integrate[(c*Cot[a + b*x])^(3/2),x]`

output `-(((c*Cot[a + b*x])^(3/2)*(ArcTan[1 - Sqrt[2]*Sqrt[Cot[a + b*x]]]/Sqrt[2] - ArcTan[1 + Sqrt[2]*Sqrt[Cot[a + b*x]]]/Sqrt[2] + 2*Sqrt[Cot[a + b*x]] + Log[1 - Sqrt[2]*Sqrt[Cot[a + b*x]] + Cot[a + b*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[a + b*x]] + Cot[a + b*x]]/(2*Sqrt[2])))/(b*Cot[a + b*x]^(3/2)))`

**Rubi [A] (warning: unable to verify)**

Time = 0.48 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.31, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$ , Rules used = {3042, 3954, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (c \cot(a + bx))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \left(-c \tan\left(a + bx + \frac{\pi}{2}\right)\right)^{3/2} dx \\ & \quad \downarrow \text{3954} \\ & c^2 \left(-\int \frac{1}{\sqrt{c \cot(a + bx)}} dx\right) - \frac{2c\sqrt{c \cot(a + bx)}}{b} \end{aligned}$$



$$\begin{aligned}
 & \downarrow 3042 \\
 & c^2 \left( - \int \frac{1}{\sqrt{-c \tan(a + bx + \frac{\pi}{2})}} dx \right) - \frac{2c\sqrt{c \cot(a + bx)}}{b} \\
 & \downarrow 3957 \\
 & \frac{c^3 \int \frac{1}{\sqrt{c \cot(a+bx)(\cot^2(a+bx)c^2+c^2)}} d(c \cot(a + bx))}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \\
 & \downarrow 266 \\
 & \frac{2c^3 \int \frac{1}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)}}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \\
 & \downarrow 755 \\
 & \frac{2c^3 \left( \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\int \frac{c^2 \cot^2(a+bx)+c}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)}{b} - \frac{2c\sqrt{c \cot(a + bx)}}{b} \\
 & \downarrow 1476 \\
 & \frac{2c^3 \left( \frac{\frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2c} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)}{b} \\
 & \frac{2c\sqrt{c \cot(a + bx)}}{b} \\
 & \downarrow 1082 \\
 & \frac{2c^3 \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} d(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}}}{2c} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} d(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}}}{2c} + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)}{b} \\
 & \frac{2c\sqrt{c \cot(a + bx)}}{b} \\
 & \downarrow 217 \\
 & \frac{2c^3 \left( \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)}{b} \\
 & \frac{2c\sqrt{c \cot(a + bx)}}{b}
 \end{aligned}$$

↓ 1479

$$2c^3 \left( \frac{\int -\frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} - \frac{\int -\frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right)$$

---


$$\frac{2c\sqrt{c}\cot(a+bx)}{b}$$

↓ 25

$$2c^3 \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)-1)}{\sqrt{2}\sqrt{c}} \right)$$

---


$$\frac{2c\sqrt{c}\cot(a+bx)}{b}$$

↓ 27

$$2c^3 \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)-1)}{\sqrt{2}\sqrt{c}} \right)$$

---


$$\frac{2c\sqrt{c}\cot(a+bx)}{b}$$

↓ 1103

$$2c^3 \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c}\cot(a+bx))}{\sqrt{2}\sqrt{c}} + \frac{\log(\sqrt{2}c^{3/2}\cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(-\sqrt{2}c^{3/2}\cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} \right)$$

---


$$\frac{2c\sqrt{c}\cot(a+bx)}{b}$$

input

Int[(c\*Cot[a + b\*x])^(3/2),x]

output

$$\frac{(-2*c*\sqrt{c*\cot[a + b*x]})/b + (2*c^3*((-\operatorname{ArcTan}[1 - \sqrt{2}*\sqrt{c}*\cot[a + b*x]]/(\sqrt{2}*\sqrt{c})) + \operatorname{ArcTan}[1 + \sqrt{2}*\sqrt{c}*\cot[a + b*x]]/(\sqrt{2}*\sqrt{c}))/2*c) + (-1/2*\log[c - \sqrt{2}*c^{3/2}*\cot[a + b*x] + c^2*\cot[a + b*x]^2]/(\sqrt{2}*\sqrt{c}) + \log[c + \sqrt{2}*c^{3/2}*\cot[a + b*x] + c^2*\cot[a + b*x]^2]/(2*\sqrt{2}*\sqrt{c}))/2*c)/b$$

### Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$$

rule 27

$$\operatorname{Int}[(a_*)*(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!MatchQ}[F_x, (b_*)*(G_x)] /; \operatorname{FreeQ}[b, x]$$

rule 217

$$\operatorname{Int}[(a_*) + (b_*)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[-b, 2])^{-1})*\operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{LtQ}[a, 0] \operatorname{||} \operatorname{LtQ}[b, 0])$$

rule 266

$$\operatorname{Int}[(c_*)*(x_)^m*((a_*) + (b_*)*(x_)^2)^p], x\_Symbol] \rightarrow \operatorname{With}[\{k = \operatorname{Denominator}[m]\}, \operatorname{Simp}[k/c \operatorname{Subst}[\operatorname{Int}[x^{k*(m+1)-1}*(a + b*(x^{2*k}/c^2))^p, x], x, (c*x)^{1/k}], x]] /; \operatorname{FreeQ}[\{a, b, c, p\}, x] \&\& \operatorname{FractionQ}[m] \&\& \operatorname{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 755

$$\operatorname{Int}[(a_*) + (b_*)*(x_)^4)^{-1}, x\_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numerator}[\operatorname{Rt}[a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[a/b, 2]]\}, \operatorname{Simp}[1/(2*r) \operatorname{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \operatorname{Simp}[1/(2*r) \operatorname{Int}[(r + s*x^2)/(a + b*x^4), x], x]] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& (\operatorname{GtQ}[a/b, 0] \operatorname{||} (\operatorname{PosQ}[a/b] \&\& \operatorname{AtomQ}[\operatorname{SplitProduct}[\operatorname{SumBaseQ}, a]] \&\& \operatorname{AtomQ}[\operatorname{SplitProduct}[\operatorname{SumBaseQ}, b]]))$$

rule 1082

$$\operatorname{Int}[(a_*) + (b_*)*(x_) + (c_*)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{With}[\{q = 1 - 4*\operatorname{Simplify}[a*(c/b^2)]\}, \operatorname{Simp}[-2/b \operatorname{Subst}[\operatorname{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \operatorname{RationalQ}[q] \&\& (\operatorname{EqQ}[q^2, 1] \operatorname{||} \operatorname{!RationalQ}[b^2 - 4*a*c])] /; \operatorname{FreeQ}[\{a, b, c\}, x]$$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S  
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,  
e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[  
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[  
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]  
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[  
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],  
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F  
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d  
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]  
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int  
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&  
!IntegerQ[n]`

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.97

method	result
derivativdivides	$2c \left( \frac{(c^2)^{\frac{1}{4}} \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)} + 1}{(c^2)^{\frac{1}{4}}} \right) - 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right)}{\sqrt{c \cot(bx+a)}} \right) - \frac{\dots}{b}$
default	$2c \left( \frac{(c^2)^{\frac{1}{4}} \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)} + 1}{(c^2)^{\frac{1}{4}}} \right) - 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right)}{\sqrt{c \cot(bx+a)}} \right) - \frac{\dots}{b}$

```
input int((c*cot(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2/b*c*((c*cot(b*x+a))^(1/2)-1/8*(c^2)^(1/4)*2^(1/2)*(ln((c*cot(b*x+a)+(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))/(c*cot(b*x+a)-(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))+2*arctan(2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1))-2*arctan(-2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 316 vs. 2(122) = 244.

Time = 0.09 (sec) , antiderivative size = 316, normalized size of antiderivative = 2.05

$$\int (c \cot(a + bx))^{3/2} dx = \frac{2\sqrt{2}c^{\frac{3}{2}} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}}+c}{c}\right) + 2\sqrt{2}c^{\frac{3}{2}} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}}-c}{c}\right) + \sqrt{2}c^{\frac{3}{2}} \log\left(\frac{c \cot(a + bx) + \sqrt{c \cot(a + bx)}}{c \cot(a + bx) - \sqrt{c \cot(a + bx)}}\right)}{b}$$

```
input integrate((c*cot(b*x+a))^(3/2),x, algorithm="fricas")
```

output

```
1/4*(2*sqrt(2)*c^(3/2)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) +
c)/sin(2*b*x + 2*a)) + c)/c) + 2*sqrt(2)*c^(3/2)*arctan((sqrt(2)*sqrt(c)*s
qrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) - c)/c) + sqrt(2)*c^(3/2)*l
og((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*
b*x + 2*a) + c*cos(2*b*x + 2*a) + c*sin(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)
) - sqrt(2)*c^(3/2)*log(-(sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/si
n(2*b*x + 2*a))*sin(2*b*x + 2*a) - c*cos(2*b*x + 2*a) - c*sin(2*b*x + 2*a)
- c)/sin(2*b*x + 2*a)) - 8*c*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*
a)))/b
```

**Sympy [F]**

$$\int (c \cot(a + bx))^{3/2} dx = \int (c \cot(a + bx))^{\frac{3}{2}} dx$$

input

```
integrate((c*cot(b*x+a))**(3/2),x)
```

output

```
Integral((c*cot(a + b*x))**(3/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.16

$$\int (c \cot(a + bx))^{3/2} dx = \frac{\left( 2 \sqrt{2} \sqrt{c} \arctan \left( \frac{\sqrt{2} (\sqrt{2} \sqrt{c} + 2 \sqrt{\frac{c}{\tan(bx+a)}})}{2 \sqrt{c}} \right) \right) + 2 \sqrt{2} \sqrt{c} \arctan \left( -\frac{\sqrt{2} (\sqrt{2} \sqrt{c} - 2 \sqrt{\frac{c}{\tan(bx+a)}})}{2 \sqrt{c}} \right) + \sqrt{2} \sqrt{c} \ln \left( \frac{\sqrt{2} \sqrt{c} + 2 \sqrt{\frac{c}{\tan(bx+a)}}}{\sqrt{2} \sqrt{c} - 2 \sqrt{\frac{c}{\tan(bx+a)}}} \right)}{2 \sqrt{2} \sqrt{c}}$$

input

```
integrate((c*cot(b*x+a))^(3/2),x, algorithm="maxima")
```

output

```
1/4*(2*sqrt(2)*sqrt(c)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(
b*x + a)))/sqrt(c)) + 2*sqrt(2)*sqrt(c)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(
c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c)) + sqrt(2)*sqrt(c)*log(sqrt(2)*sqrt(c)
)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a)) - sqrt(2)*sqrt(c)*log(-sqrt(2)
)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a)) - 8*sqrt(c/tan(b*x +
a))*c/b
```

**Giac [F]**

$$\int (c \cot(a + bx))^{3/2} dx = \int (c \cot(bx + a))^{\frac{3}{2}} dx$$

input

```
integrate((c*cot(b*x+a))^(3/2),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(3/2), x)
```

**Mupad [B] (verification not implemented)**

Time = 8.68 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.49

$$\int (c \cot(a + bx))^{3/2} dx = -\frac{2c \sqrt{c \cot(a + bx)}}{b} - \frac{(-1)^{1/4} c^{3/2} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right) \operatorname{li}}{b} - \frac{(-1)^{1/4} c^{3/2} \operatorname{atanh}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right) \operatorname{li}}{b}$$

input

```
int((c*cot(a + b*x))^(3/2),x)
```

output

```
- (2*c*(c*cot(a + b*x))^(1/2))/b - ((-1)^(1/4)*c^(3/2)*atan(((-1)^(1/4)*(c
*cot(a + b*x))^(1/2))/c^(1/2))*1i)/b - ((-1)^(1/4)*c^(3/2)*atanh(((-1)^(1/
4)*(c*cot(a + b*x))^(1/2))/c^(1/2))*1i)/b
```

**Reduce [F]**

$$\int (c \cot(a + bx))^{3/2} dx = \frac{\sqrt{c} c \left( -2\sqrt{\cot(bx + a)} - \left( \int \frac{\sqrt{\cot(bx+a)}}{\cot(bx+a)} dx \right) b \right)}{b}$$

input `int((c*cot(b*x+a))^(3/2),x)`

output `(sqrt(c)*c*( - 2*sqrt(cot(a + b*x)) - int(sqrt(cot(a + b*x))/cot(a + b*x), x)*b))/b`



### 3.12 $\int \sqrt{c \cot(a + bx)} dx$

Optimal result . . . . .	128
Mathematica [A] (verified) . . . . .	128
Rubi [A] (warning: unable to verify) . . . . .	129
Maple [A] (verified) . . . . .	132
Fricas [B] (verification not implemented) . . . . .	133
Sympy [F] . . . . .	133
Maxima [A] (verification not implemented) . . . . .	134
Giac [F] . . . . .	134
Mupad [B] (verification not implemented) . . . . .	135
Reduce [F] . . . . .	135

#### Optimal result

Integrand size = 12, antiderivative size = 136

$$\int \sqrt{c \cot(a + bx)} dx = \frac{\sqrt{c} \arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} - \frac{\sqrt{c} \arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b} + \frac{\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}b}$$

output

```
1/2*c^(1/2)*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b-1/2*c^(1/2)*arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b+1/2*c^(1/2)*arctanh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a)))*2^(1/2)/b
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.52

$$\int \sqrt{c \cot(a + bx)} dx = \frac{\left(-\arctan\left(\sqrt[4]{-\cot^2(a + bx)}\right) + \operatorname{arctanh}\left(\sqrt[4]{-\cot^2(a + bx)}\right)\right) \sqrt[4]{-\cot(a + bx)} \sqrt{c \cot(a + bx)}}{b \cot^{\frac{3}{4}}(a + bx)}$$

input `Integrate[Sqrt[c*Cot[a + b*x]],x]`

output  $((-\text{ArcTan}[(-\text{Cot}[a + b*x]^2)^{(1/4)}] + \text{ArcTanh}[(-\text{Cot}[a + b*x]^2)^{(1/4)}]) * (-\text{Cot}[a + b*x])^{(1/4)} * \text{Sqrt}[c * \text{Cot}[a + b*x]] / (b * \text{Cot}[a + b*x]^{(3/4)})$

### Rubi [A] (warning: unable to verify)

Time = 0.42 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.28, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{c \cot(a + bx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-c \tan\left(a + bx + \frac{\pi}{2}\right)} dx \\
 & \quad \downarrow \text{3957} \\
 & \frac{c \int \frac{\sqrt{c \cot(a+bx)}}{\cot^2(a+bx)c^2+c^2} d(c \cot(a + bx))}{b} \\
 & \quad \downarrow \text{266} \\
 & \frac{2c \int \frac{c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{826} \\
 & \frac{2c \left( \frac{1}{2} \int \frac{c^2 \cot^2(a+bx)+c}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)} - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a + bx)} \right)}{b} \\
 & \quad \downarrow \text{1476} \\
 & \frac{2c \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a + bx)} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a + bx)} \right) \right)}{b}
 \end{aligned}$$

↓ 1082

$$2c \left( \frac{\int \frac{-c^2 \cot^2(a+bx)-1}{\sqrt{2}\sqrt{c}} d(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} - \frac{\int \frac{-c^2 \cot^2(a+bx)-1}{\sqrt{2}\sqrt{c}} d(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}$$


---

$b$

↓ 217

$$2c \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} \right)$$


---

$b$

↓ 1479

$$2c \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c} \cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c} \cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1) - \arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx)) \right) \right)$$


---

$b$

↓ 25

$$2c \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c} \cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c} \cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1) - \arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx)) \right) \right)$$


---

$b$

↓ 27

$$2c \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c} \cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c} \cot(a+bx)}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{c}} \right) + \frac{1}{2} \left( \arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1) - \arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx)) \right) \right)$$


---

$b$

↓ 1103

$$2c \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\log(-\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} \right) \right)$$


---

$b$

input `Int[Sqrt[c*Cot[a + b*x]], x]`

output 
$$\frac{(-2*c*((-ArcTan[1 - Sqrt[2]*Sqrt[c]*Cot[a + b*x])/(Sqrt[2]*Sqrt[c])) + ArcTan[1 + Sqrt[2]*Sqrt[c]*Cot[a + b*x])/(Sqrt[2]*Sqrt[c]))/2 + (Log[c - Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]) - Log[c + Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]))/2)/b$$

### Defintions of rubi rules used

rule 25 
$$\text{Int}[-(F_x), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$$

rule 27 
$$\text{Int}[(a_*)(F_x), x\_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$$

rule 217 
$$\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 266 
$$\text{Int}[(c_*)(x_)^m * ((a_*) + (b_*)(x_)^2)^p], x\_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{k*(m+1)-1}*(a + b*(x^{2*k}/c^2))^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 826 
$$\text{Int}[(x_)^2/((a_*) + (b_*)(x_)^4), x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \text{ Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \text{ Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 1082 
$$\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{c\sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2+\sqrt{c^2}}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2+\sqrt{c^2}}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( -\frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right) \right)}{4b(c^2)^{\frac{1}{4}}}$
default	$\frac{c\sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2+\sqrt{c^2}}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2+\sqrt{c^2}}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( -\frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right) \right)}{4b(c^2)^{\frac{1}{4}}}$

input `int((c*cot(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output

```
-1/4/b*c/(c^2)^(1/4)*2^(1/2)*(ln((c*cot(b*x+a)-(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2))/(c*cot(b*x+a)+(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))+2*arctan(2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)-2*arctan(-2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 286 vs.  $2(106) = 212$ .

Time = 0.11 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.10

$$\int \sqrt{c \cot(a + bx)} dx =$$

$$2\sqrt{2}\sqrt{c} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}}+c}{c}\right) + 2\sqrt{2}\sqrt{c} \arctan\left(\frac{\sqrt{2}\sqrt{c}\sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}}-c}{c}\right) - \sqrt{2}\sqrt{c} \log\left(\frac{\sqrt{2}\sqrt{c}}{\dots}\right)$$

input

```
integrate((c*cot(b*x+a))^(1/2),x, algorithm="fricas")
```

output

```
-1/4*(2*sqrt(2)*sqrt(c)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) + c)/c) + 2*sqrt(2)*sqrt(c)*arctan((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) - c)/c) - sqrt(2)*sqrt(c)*log((sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) + c*cos(2*b*x + 2*a) + c*sin(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)) + sqrt(2)*sqrt(c)*log(-(sqrt(2)*sqrt(c)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a) - c*cos(2*b*x + 2*a) - c*sin(2*b*x + 2*a) - c)/sin(2*b*x + 2*a)))/b
```

### Sympy [F]

$$\int \sqrt{c \cot(a + bx)} dx = \int \sqrt{c \cot(a + bx)} dx$$

input

```
integrate((c*cot(b*x+a))**(1/2),x)
```

output `Integral(sqrt(c*cot(a + b*x)), x)`

### Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.21

$$\int \sqrt{c \cot(a + bx)} dx =$$

$$\frac{c \left( \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}\sqrt{c} + 2\sqrt{\frac{c}{\tan(bx+a)}})}{2\sqrt{c}}\right)}{\sqrt{c}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}\sqrt{c} - 2\sqrt{\frac{c}{\tan(bx+a)}})}{2\sqrt{c}}\right)}{\sqrt{c}} - \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\frac{c}{\tan(bx+a)}} + c + \frac{c}{\tan(bx+a)}\right)}{\sqrt{c}} \right)}{4b}$$

input `integrate((c*cot(b*x+a))^(1/2),x, algorithm="maxima")`

output `-1/4*c*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(b*x + a)))/sqrt(c))/sqrt(c) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c))/sqrt(c) - sqrt(2)*log(sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c) + sqrt(2)*log(-sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c))/b`

### Giac [F]

$$\int \sqrt{c \cot(a + bx)} dx = \int \sqrt{c \cot(bx + a)} dx$$

input `integrate((c*cot(b*x+a))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(c*cot(b*x + a)), x)`

**Mupad [B] (verification not implemented)**

Time = 8.57 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.37

$$\int \sqrt{c \cot(a + bx)} dx$$

$$= -\frac{(-1)^{1/4} \sqrt{c} \left( \operatorname{atan} \left( \frac{(-1)^{1/4} \sqrt{c \cot(a+bx)}}{\sqrt{c}} \right) - \operatorname{atanh} \left( \frac{(-1)^{1/4} \sqrt{c \cot(a+bx)}}{\sqrt{c}} \right) \right)}{b}$$

input `int((c*cot(a + b*x))^(1/2),x)`output `-((-1)^(1/4)*c^(1/2)*(atan(((1/4)*(-1)*c*cot(a + b*x))^(1/2))/c^(1/2)) - atanh(((1/4)*(-1)*c*cot(a + b*x))^(1/2))/c^(1/2)))/b`**Reduce [F]**

$$\int \sqrt{c \cot(a + bx)} dx = \sqrt{c} \left( \int \sqrt{\cot(bx + a)} dx \right)$$

input `int((c*cot(b*x+a))^(1/2),x)`output `sqrt(c)*int(sqrt(cot(a + b*x)),x)`



### 3.13 $\int \frac{1}{\sqrt{c \cot(a+bx)}} dx$

Optimal result	136
Mathematica [A] (verified)	136
Rubi [A] (warning: unable to verify)	137
Maple [A] (verified)	141
Fricas [B] (verification not implemented)	141
Sympy [F]	142
Maxima [A] (verification not implemented)	142
Giac [F]	143
Mupad [B] (verification not implemented)	143
Reduce [F]	144

#### Optimal result

Integrand size = 12, antiderivative size = 137

$$\int \frac{1}{\sqrt{c \cot(a+bx)}} dx = \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b\sqrt{c}} - \frac{\arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}b\sqrt{c}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}b\sqrt{c}}$$

output

```
1/2*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(1/2)-1/2*arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(1/2)-1/2*arctanh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a))*2^(1/2)/b/c^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{c \cot(a+bx)}} dx = \frac{\sqrt{\cot(a+bx)} \left( 2 \arctan\left(1 - \sqrt{2}\sqrt{\cot(a+bx)}\right) - 2 \arctan\left(1 + \sqrt{2}\sqrt{\cot(a+bx)}\right) + \log\left(1 - \sqrt{2}\sqrt{\cot(a+bx)}\right) \right)}{2\sqrt{2}b\sqrt{c \cot(a+bx)}}$$

input `Integrate[1/Sqrt[c*Cot[a + b*x]],x]`

output `(Sqrt[Cot[a + b*x]]*(2*ArcTan[1 - Sqrt[2]*Sqrt[Cot[a + b*x]]] - 2*ArcTan[1 + Sqrt[2]*Sqrt[Cot[a + b*x]]) + Log[1 - Sqrt[2]*Sqrt[Cot[a + b*x]] + Cot[a + b*x]] - Log[1 + Sqrt[2]*Sqrt[Cot[a + b*x]] + Cot[a + b*x]])/(2*Sqrt[2]*b*Sqrt[c*Cot[a + b*x]])`

### Rubi [A] (warning: unable to verify)

Time = 0.40 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.31, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{c \cot(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-c \tan(a + bx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{3957} \\
 & \frac{c \int \frac{1}{\sqrt{c \cot(a + bx)} (\cot^2(a + bx) c^2 + c^2)} d(c \cot(a + bx))}{b} \\
 & \quad \downarrow \text{266} \\
 & \frac{2c \int \frac{1}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{755} \\
 & \frac{2c \left( \frac{\int \frac{c - c^2 \cot^2(a + bx)}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{2c} + \frac{\int \frac{c^2 \cot^2(a + bx) + c}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{2c} \right)}{b} \\
 & \quad \downarrow \text{1476}
 \end{aligned}$$

$$2c \left( \frac{\frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)} + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx) + c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)$$

$b$

↓ 1082

$$2c \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} \frac{d(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}}}{2c} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} \frac{d(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}}}{2c} + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx) + c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)$$

$b$

↓ 217

$$2c \left( \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx) + c^2} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)$$

$b$

↓ 1479

$$2c \left( \frac{\int \frac{\sqrt{2}\sqrt{c} - 2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{2}(\sqrt{c} + \sqrt{2}\sqrt{c \cot(a+bx)})}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} \right)$$

$b$

↓ 25

$$2c \left( \frac{\int \frac{\sqrt{2}\sqrt{c} - 2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{2}(\sqrt{c} + \sqrt{2}\sqrt{c \cot(a+bx)})}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)$$

$b$

↓ 27

$$2c \left( \frac{\int \frac{\sqrt{2}\sqrt{c} - 2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{c} + \sqrt{2}\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)$$

$b$

↓ 1103

$$\frac{2c \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c}\cot(a+bx))}{\sqrt{2}\sqrt{c}} + \frac{\log(\sqrt{2}c^{3/2}\cot(a+bx)+c^2\cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(-\sqrt{2}c^{3/2}\cot(a+bx)+c^2\cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} \right)}{b}$$

input `Int[1/Sqrt[c*Cot[a + b*x]],x]`

output `(-2*c*((-(ArcTan[1 - Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c])) + ArcTan[1 + Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c]))/(2*c) + (-1/2*Log[c - Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(Sqrt[2]*Sqrt[c]) + Log[c + Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]))/(2*c))/b`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755  $\text{Int}[(a + (b \cdot x^4)^{-1}), x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2 \cdot r) \int (r - s \cdot x^2)/(a + b \cdot x^4), x], x] + \text{Simp}[1/(2 \cdot r) \int (r + s \cdot x^2)/(a + b \cdot x^4), x], x] /; \text{FreeQ}\{a, b\}, x] \&\& (\text{GtQ}[a/b, 0] \parallel (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 1082  $\text{Int}[(a + (b \cdot x) + (c \cdot x^2)^{-1}), x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\int 1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4 \cdot a \cdot c]) /; \text{FreeQ}\{a, b, c\}, x]$

rule 1103  $\text{Int}[(d + (e \cdot x))/(a + (b \cdot x) + (c \cdot x^2)), x\_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1476  $\text{Int}[(d + (e \cdot x^2))/(a + (c \cdot x^4)), x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \int 1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \int 1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \&\& \text{PosQ}[d \cdot e]$

rule 1479  $\text{Int}[(d + (e \cdot x^2))/(a + (c \cdot x^4)), x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \int (q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \int (q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \&\& \text{NegQ}[d \cdot e]$

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3957  $\text{Int}[(b \cdot \tan[(c \cdot x) + (d \cdot x)])^n], x\_Symbol] \rightarrow \text{Simp}[b/d \text{ Subst}[\int x^n/(b^2 + x^2), x], x, b \cdot \text{Tan}[c + d \cdot x]], x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[n]$

### Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.01

method	result
derivativedivides	$\frac{(c^2)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{c\cot(bx+a)+(c^2)^{\frac{1}{4}}\sqrt{c\cot(bx+a)}\sqrt{2+\sqrt{c^2}}}{c\cot(bx+a)-(c^2)^{\frac{1}{4}}\sqrt{c\cot(bx+a)}\sqrt{2+\sqrt{c^2}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{c\cot(bx+a)}}{(c^2)^{\frac{1}{4}}}+1\right)-2\arctan\left(-\frac{\sqrt{2}\sqrt{c\cot(bx+a)}}{(c^2)^{\frac{1}{4}}}\right)}{4bc}$
default	$\frac{(c^2)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{c\cot(bx+a)+(c^2)^{\frac{1}{4}}\sqrt{c\cot(bx+a)}\sqrt{2+\sqrt{c^2}}}{c\cot(bx+a)-(c^2)^{\frac{1}{4}}\sqrt{c\cot(bx+a)}\sqrt{2+\sqrt{c^2}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{c\cot(bx+a)}}{(c^2)^{\frac{1}{4}}}+1\right)-2\arctan\left(-\frac{\sqrt{2}\sqrt{c\cot(bx+a)}}{(c^2)^{\frac{1}{4}}}\right)}{4bc}$

input `int(1/(c*cot(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/4/b/c*(c^2)^{(1/4)}*2^{(1/2)}*(\ln((c*\cot(b*x+a)+(c^2)^{(1/4)}*(c*\cot(b*x+a))^{(1/2)}*2^{(1/2)}+(c^2)^{(1/2)})/(c*\cot(b*x+a)-(c^2)^{(1/4)}*(c*\cot(b*x+a))^{(1/2)}*2^{(1/2)}+(c^2)^{(1/2)}))+2*\arctan(2^{(1/2)}/(c^2)^{(1/4)}*(c*\cot(b*x+a))^{(1/2)}+1)-2*\arctan(-2^{(1/2)}/(c^2)^{(1/4)}*(c*\cot(b*x+a))^{(1/2)}+1))$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. 2(106) = 212.

Time = 0.08 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.96

$$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx =$$

$$\frac{2\sqrt{2}\arctan\left(\frac{\sqrt{2}\sqrt{\frac{c\cos(2bx+2a)+c}{\sin(2bx+2a)}}}{\sqrt{c}}+1\right)}{\sqrt{c}} + \frac{2\sqrt{2}\arctan\left(\frac{\sqrt{2}\sqrt{\frac{c\cos(2bx+2a)+c}{\sin(2bx+2a)}}}{\sqrt{c}}-1\right)}{\sqrt{c}} + \frac{\sqrt{2}\log\left(\frac{\sqrt{2}\sqrt{\frac{c\cos(2bx+2a)+c}{\sin(2bx+2a)}}\sin(2bx+2a)}{\sqrt{c}}+\cos(2bx+2a)\right)}{\sqrt{c}}$$

$4b$

input `integrate(1/(c*cot(b*x+a))^(1/2),x, algorithm="fricas")`

output

```
-1/4*(2*sqrt(2)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))/sqrt(c) + 1)/sqrt(c) + 2*sqrt(2)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))/sqrt(c) - 1)/sqrt(c) + sqrt(2)*log((sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) + cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/sin(2*b*x + 2*a))/sqrt(c) - sqrt(2)*log(-(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) - cos(2*b*x + 2*a) - sin(2*b*x + 2*a) - 1)/sin(2*b*x + 2*a))/sqrt(c))/b
```

### Sympy [F]

$$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx = \int \frac{1}{\sqrt{c \cot(a + bx)}} dx$$

input

```
integrate(1/(c*cot(b*x+a))**(1/2), x)
```

output

```
Integral(1/sqrt(c*cot(a + b*x)), x)
```

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.20

$$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx =$$

$$c \left( \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}\sqrt{c} + 2\sqrt{\tan(bx+a)})}{2\sqrt{c}}\right)}{c^{\frac{3}{2}}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}\sqrt{c} - 2\sqrt{\tan(bx+a)})}{2\sqrt{c}}\right)}{c^{\frac{3}{2}}} + \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\frac{c}{\tan(bx+a)} + c} + \frac{c}{\tan(bx+a)}\right)}{c^{\frac{3}{2}}} \right)$$


---

4 b

input

```
integrate(1/(c*cot(b*x+a))^(1/2), x, algorithm="maxima")
```

output

```
-1/4*c*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(b*x + a)))/sqrt(c))/c^(3/2) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c))/c^(3/2) + sqrt(2)*log(sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/c^(3/2) - sqrt(2)*log(-sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/c^(3/2))/b
```

**Giac [F]**

$$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx = \int \frac{1}{\sqrt{c \cot(bx + a)}} dx$$

input

```
integrate(1/(c*cot(b*x+a))^(1/2),x, algorithm="giac")
```

output

```
integrate(1/sqrt(c*cot(b*x + a)), x)
```

**Mupad [B] (verification not implemented)**

Time = 9.65 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.42

$$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx = \frac{(-1)^{1/4} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a+bx)}}{\sqrt{c}}\right) \operatorname{li}}{b \sqrt{c}} + \frac{(-1)^{1/4} \operatorname{atanh}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a+bx)}}{\sqrt{c}}\right) \operatorname{li}}{b \sqrt{c}}$$

input

```
int(1/(c*cot(a + b*x))^(1/2),x)
```

output

```
((-1)^(1/4)*atan((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2))*li)/(b*c^(1/2)) + ((-1)^(1/4)*atanh((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2))*li)/(b*c^(1/2))
```



**Reduce [F]**

$$\int \frac{1}{\sqrt{c \cot(a + bx)}} dx = \frac{\sqrt{c} \left( \int \frac{\sqrt{\cot(bx+a)}}{\cot(bx+a)} dx \right)}{c}$$

input `int(1/(c*cot(b*x+a))^(1/2),x)`

output `(sqrt(c)*int(sqrt(cot(a + b*x))/cot(a + b*x),x))/c`

### 3.14 $\int \frac{1}{(c \cot(a+bx))^{3/2}} dx$

Optimal result	145
Mathematica [A] (verified)	145
Rubi [A] (warning: unable to verify)	146
Maple [A] (verified)	150
Fricas [B] (verification not implemented)	151
Sympy [F]	152
Maxima [A] (verification not implemented)	152
Giac [F]	153
Mupad [B] (verification not implemented)	153
Reduce [F]	154

#### Optimal result

Integrand size = 12, antiderivative size = 157

$$\int \frac{1}{(c \cot(a+bx))^{3/2}} dx = -\frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}bc^{3/2}} + \frac{\arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}bc^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c+\sqrt{c \cot(a+bx)}}}\right)}{\sqrt{2}bc^{3/2}} + \frac{2}{bc\sqrt{c \cot(a+bx)}}$$

output

```
-1/2*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(3/2)+1/2*
arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(3/2)-1/2*arcta
nh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a)))*2^(1/2)/b/c^
(3/2)+2/b/c/(c*cot(b*x+a))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.52

$$\int \frac{1}{(c \cot(a+bx))^{3/2}} dx = \frac{2 + \arctan\left(\sqrt[4]{-\cot^2(a+bx)}\right) \sqrt[4]{-\cot^2(a+bx)} - \operatorname{arctanh}\left(\sqrt[4]{-\cot^2(a+bx)}\right)}{bc\sqrt{c \cot(a+bx)}}$$

input `Integrate[(c*Cot[a + b*x])^(-3/2),x]`

output  $(2 + \text{ArcTan}[(-\text{Cot}[a + b*x]^2)^{(1/4)}]*(-\text{Cot}[a + b*x]^2)^{(1/4)} - \text{ArcTanh}[(-\text{Cot}[a + b*x]^2)^{(1/4)}]*(-\text{Cot}[a + b*x]^2)^{(1/4)})/(b*c*\text{Sqrt}[c*\text{Cot}[a + b*x]])$

### Rubi [A] (warning: unable to verify)

Time = 0.46 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.25, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$ , Rules used = {3042, 3955, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \cot(a + bx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{3955} \\
 & \frac{2}{bc\sqrt{c \cot(a + bx)}} - \frac{\int \sqrt{c \cot(a + bx)} dx}{c^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{bc\sqrt{c \cot(a + bx)}} - \frac{\int \sqrt{-c \tan(a + bx + \frac{\pi}{2})} dx}{c^2} \\
 & \quad \downarrow \text{3957} \\
 & \frac{\int \frac{\sqrt{c \cot(a + bx)}}{\cot^2(a + bx)c^2 + c^2} d(c \cot(a + bx))}{bc} + \frac{2}{bc\sqrt{c \cot(a + bx)}} \\
 & \quad \downarrow \text{266} \\
 & \frac{2 \int \frac{c^2 \cot^2(a + bx)}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{bc} + \frac{2}{bc\sqrt{c \cot(a + bx)}}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 826 \\
& \frac{2\left(\frac{1}{2} \int \frac{c^2 \cot^2(a+bx)+c}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}\right)}{\frac{bc}{2}} + \\
& \frac{bc}{\sqrt{c \cot(a+bx)}} \\
& \downarrow 1476 \\
& \frac{2\left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}\right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}\right)}{\frac{2}{bc}} - \\
& \frac{bc}{\sqrt{c \cot(a+bx)}} \\
& \downarrow 1082 \\
& \frac{2\left(\frac{1}{2} \left(\frac{\int \frac{1}{-c^2 \cot^2(a+bx)-1} d(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx)-1} d(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}}\right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}\right)}{\frac{2}{bc}} - \\
& \frac{bc}{\sqrt{c \cot(a+bx)}} \\
& \downarrow 217 \\
& \frac{2\left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}}\right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}\right)}{\frac{bc}{2}} + \\
& \frac{bc}{\sqrt{c \cot(a+bx)}} \\
& \downarrow 1479 \\
& \frac{2\left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c} \cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c} \cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}}\right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}}\right)\right)}{\frac{2}{bc}} + \\
& \frac{bc}{\sqrt{c \cot(a+bx)}} \\
& \downarrow 25
\end{aligned}$$

$$\begin{aligned}
 & 2 \left( \frac{1}{2} \left( - \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c \cot(a+bx)})}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c \cot(a+bx)})}{\sqrt{2}\sqrt{c}} \right) \right) \\
 & \qquad \qquad \qquad \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & 2 \left( \frac{1}{2} \left( - \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c \cot(a+bx)})}{\sqrt{2}\sqrt{c}} \right) \right) \\
 & \qquad \qquad \qquad \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \qquad \qquad \qquad \downarrow 1103 \\
 & 2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c \cot(a+bx)+1})}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c \cot(a+bx)})}{\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\log(-\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} \right) \right) \\
 & \qquad \qquad \qquad \frac{2}{bc\sqrt{c \cot(a+bx)}}
 \end{aligned}$$

input `Int[(c*Cot[a + b*x])^(-3/2),x]`

output `2/(b*c*Sqrt[c*Cot[a + b*x]]) + (2*((-(ArcTan[1 - Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c])) + ArcTan[1 + Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c])))/2 + (Log[c - Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]) - Log[c + Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]))/2)/(b*c)`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.00

method	result
derivativedivides	$2c \left( -\frac{1}{c^2 \sqrt{c \cot(bx+a)}} - \frac{\sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right)}{8c^2 (c^2)^{\frac{1}{4}}} \right)$
default	$2c \left( -\frac{1}{c^2 \sqrt{c \cot(bx+a)}} - \frac{\sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} \right)}{8c^2 (c^2)^{\frac{1}{4}}} \right)$

```
input int(1/(c*cot(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -2/b*c*(-1/c^2/(c*cot(b*x+a))^(1/2)-1/8/c^2/(c^2)^(1/4)*2^(1/2)*(ln((c*cot
(b*x+a)-(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2))/(c*cot(b*x+a)
)+(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))+2*arctan(2^(1/2)/
(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)-2*arctan(-2^(1/2)/(c^2)^(1/4)*(c*cot(b
*x+a))^(1/2)+1)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 377 vs. 2(124) = 248.

Time = 0.11 (sec) , antiderivative size = 377, normalized size of antiderivative = 2.40

$$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx = \frac{2\sqrt{2}(c \cos(2bx + 2a) + c) \arctan \left( \frac{\sqrt{2} \sqrt{\frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)}}}{\sqrt{c}} + 1 \right)}{\sqrt{c}} + \frac{2\sqrt{2}(c \cos(2bx + 2a) + c) \arctan \left( \frac{\sqrt{2} \sqrt{\frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)}}}{\sqrt{c}} \right)}{\sqrt{c}}$$

```
input integrate(1/(c*cot(b*x+a))^(3/2),x, algorithm="fricas")
```



output

```
1/4*(2*sqrt(2)*(c*cos(2*b*x + 2*a) + c)*arctan(sqrt(2)*sqrt((c*cos(2*b*x +
2*a) + c)/sin(2*b*x + 2*a))/sqrt(c) + 1)/sqrt(c) + 2*sqrt(2)*(c*cos(2*b*x
+ 2*a) + c)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)
)/sqrt(c) - 1)/sqrt(c) - sqrt(2)*(c*cos(2*b*x + 2*a) + c)*log((sqrt(2)*sqr
t((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) + co
s(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/sin(2*b*x + 2*a))/sqrt(c) + sqrt(2)
*(c*cos(2*b*x + 2*a) + c)*log(-(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(
2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) - cos(2*b*x + 2*a) - sin(2*b*x + 2*
a) - 1)/sin(2*b*x + 2*a))/sqrt(c) + 8*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*
b*x + 2*a))*sin(2*b*x + 2*a))/(b*c^2*cos(2*b*x + 2*a) + b*c^2)
```

### Sympy [F]

$$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx = \int \frac{1}{(c \cot(a + bx))^{\frac{3}{2}}} dx$$

input

```
integrate(1/(c*cot(b*x+a))**(3/2), x)
```

output

```
Integral((c*cot(a + b*x))**(-3/2), x)
```

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.19

$$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx = \frac{c \left( \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}\sqrt{c}+2\sqrt{\tan(bx+a)})}{2\sqrt{c}}\right)}{\sqrt{c}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}\sqrt{c}-2\sqrt{\tan(bx+a)})}{2\sqrt{c}}\right)}{\sqrt{c}} \right) - \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\tan(bx+a)}\right)}{c^2}}{4b}$$

input

```
integrate(1/(c*cot(b*x+a))^(3/2), x, algorithm="maxima")
```

output

```
1/4*c*((2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(b*x +
a)))/sqrt(c))/sqrt(c) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(c) -
2*sqrt(c/tan(b*x + a)))/sqrt(c))/sqrt(c) - sqrt(2)*log(sqrt(2)*sqrt(c)*sq
r t(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c) + sqrt(2)*log(-sqrt(2)*sq
r t(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c))/c^2 + 8/(c^2*sqrt
(c/tan(b*x + a)))/b
```

**Giac [F]**

$$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx = \int \frac{1}{(c \cot(bx + a))^{\frac{3}{2}}} dx$$

input

```
integrate(1/(c*cot(b*x+a))^(3/2),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(-3/2), x)
```

**Mupad [B] (verification not implemented)**

Time = 9.91 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.48

$$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx = \frac{2}{bc \sqrt{c \cot(a + bx)}} + \frac{(-1)^{1/4} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right)}{bc^{3/2}} - \frac{(-1)^{1/4} \operatorname{atanh}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right)}{bc^{3/2}}$$

input

```
int(1/(c*cot(a + b*x))^(3/2),x)
```

output

```
2/(b*c*(c*cot(a + b*x))^(1/2)) + ((-1)^(1/4)*atan((( -1)^(1/4)*(c*cot(a + b
*x))^(1/2))/c^(1/2)))/(b*c^(3/2)) - ((-1)^(1/4)*atanh((( -1)^(1/4)*(c*cot(a
+ b*x))^(1/2))/c^(1/2)))/(b*c^(3/2))
```

**Reduce [F]**

$$\int \frac{1}{(c \cot(a + bx))^{3/2}} dx = \frac{\sqrt{c} \left( \int \frac{\sqrt{\cot(bx+a)}}{\cot(bx+a)^2} dx \right)}{c^2}$$

input `int(1/(c*cot(b*x+a))^(3/2),x)`

output `(sqrt(c)*int(sqrt(cot(a + b*x))/cot(a + b*x)**2,x))/c**2`

### 3.15 $\int \frac{1}{(c \cot(a+bx))^{5/2}} dx$

Optimal result	155
Mathematica [A] (verified)	155
Rubi [A] (warning: unable to verify)	156
Maple [A] (verified)	160
Fricas [B] (verification not implemented)	161
Sympy [F]	162
Maxima [A] (verification not implemented)	162
Giac [F]	163
Mupad [B] (verification not implemented)	163
Reduce [F]	164

#### Optimal result

Integrand size = 12, antiderivative size = 158

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = -\frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}bc^{5/2}} + \frac{\arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}bc^{5/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}bc^{5/2}} + \frac{2}{3bc(c \cot(a + bx))^{3/2}}$$

output

```
-1/2*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(5/2)+1/2*
arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(5/2)+1/2*arcta
nh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a)))*2^(1/2)/b/c^
(5/2)+2/3/b/c/(c*cot(b*x+a))^(3/2)
```

#### Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.54

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \frac{-2 + 3 \arctan\left(\sqrt[4]{-\cot^2(a + bx)}\right) (-\cot^2(a + bx))^{3/4} + 3 \operatorname{arctanh}\left(\sqrt[4]{-\cot^2(a + bx)}\right) (-\cot^2(a + bx))^{3/4}}{3bc(c \cot(a + bx))^{3/2}}$$

input `Integrate[(c*Cot[a + b*x])^(-5/2),x]`

output `-1/3*(-2 + 3*ArcTan[(-Cot[a + b*x]^2)^(1/4)]*(-Cot[a + b*x]^2)^(3/4) + 3*ArcTanh[(-Cot[a + b*x]^2)^(1/4)]*(-Cot[a + b*x]^2)^(3/4))/(b*c*(c*Cot[a + b*x])^(3/2))`

### Rubi [A] (warning: unable to verify)

Time = 0.47 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.30, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$ , Rules used = {3042, 3955, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \cot(a + bx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{3955} \\
 & \frac{2}{3bc(c \cot(a + bx))^{3/2}} - \frac{\int \frac{1}{\sqrt{c \cot(a + bx)}} dx}{c^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{3bc(c \cot(a + bx))^{3/2}} - \frac{\int \frac{1}{\sqrt{-c \tan(a + bx + \frac{\pi}{2})}} dx}{c^2} \\
 & \quad \downarrow \text{3957} \\
 & \frac{\int \frac{1}{\sqrt{c \cot(a + bx)}(\cot^2(a + bx)c^2 + c^2)} d(c \cot(a + bx))}{bc} + \frac{2}{3bc(c \cot(a + bx))^{3/2}} \\
 & \quad \downarrow \text{266} \\
 & \frac{2 \int \frac{1}{c^4 \cot^4(a + bx) + c^2} d\sqrt{c \cot(a + bx)}}{bc} + \frac{2}{3bc(c \cot(a + bx))^{3/2}}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 755 \\
 & \frac{2 \left( \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\int \frac{c^2 \cot^2(a+bx)+c}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)}{bc} + \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \downarrow 1476 \\
 & \frac{2 \left( \frac{\frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2c} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)}{bc} \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \downarrow 1082 \\
 & \frac{2 \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} d(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - 1} d(\sqrt{2}\sqrt{c} \cot(a+bx) + 1)}{\sqrt{2}\sqrt{c}} + \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} \right)}{bc} + \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \downarrow 217 \\
 & \frac{2 \left( \frac{\int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{2c} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1) - \arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)}{bc} + \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \downarrow 1479 \\
 & \frac{2 \left( \frac{\int -\frac{\sqrt{2}\sqrt{c} - 2\sqrt{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} - \frac{\int -\frac{\sqrt{2}(\sqrt{c} + \sqrt{2}\sqrt{c \cot(a+bx)})}{c^2 \cot^2(a+bx) + \sqrt{2}c^{3/2} \cot(a+bx) + c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx) + 1) - \arctan(1 - \sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right)}{bc} \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \downarrow 25
 \end{aligned}$$

$$\begin{aligned}
 & 2 \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(\dots)}{2c} \right) \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & 2 \left( \frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{2}\sqrt{c}} + \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)}}{2\sqrt{c}} + \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(\dots)}{2c} \right) \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}} \\
 & \quad \downarrow 1103 \\
 & 2 \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c}\cot(a+bx))}{\sqrt{2}\sqrt{c}} + \frac{\log(\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(-\sqrt{2}c^{3/2} \cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} \right) \\
 & \frac{2}{3bc(c \cot(a+bx))^{3/2}}
 \end{aligned}$$

input `Int[(c*Cot[a + b*x])^(-5/2),x]`

output `2/(3*b*c*(c*Cot[a + b*x])^(3/2)) + (2*((-(ArcTan[1 - Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c])) + ArcTan[1 + Sqrt[2]*Sqrt[c]*Cot[a + b*x]]/(Sqrt[2]*Sqrt[c])))/(2*c) + (-1/2*Log[c - Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(Sqrt[2]*Sqrt[c]) + Log[c + Sqrt[2]*c^(3/2)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/(2*Sqrt[2]*Sqrt[c]))/(b*c)`

## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`



rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.99

method	result
derivativedivides	$2c \left( \frac{1}{3c^2(c \cot(bx+a))^{\frac{3}{2}}} - \frac{(c^2)^{\frac{1}{4}} \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2} \right)}{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)} + 1}{(c^2)^{\frac{1}{4}}} \right)}{8c^4} \right) - \frac{b}{b}$
default	$2c \left( \frac{1}{3c^2(c \cot(bx+a))^{\frac{3}{2}}} - \frac{(c^2)^{\frac{1}{4}} \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2} \right)}{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)} + 1}{(c^2)^{\frac{1}{4}}} \right)}{8c^4} \right) - \frac{b}{b}$

```
input int(1/(c*cot(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -2/b*c*(-1/3/c^2/(c*cot(b*x+a))^(3/2)-1/8/c^4*(c^2)^(1/4)*2^(1/2)*(ln((c*cot(b*x+a)+(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))/(c*cot(b*x+a)-(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))+2*arctan(2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)-2*arctan(-2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 380 vs. 2(124) = 248.

Time = 0.10 (sec) , antiderivative size = 380, normalized size of antiderivative = 2.41

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \frac{6 \sqrt{2}(c \cos(2bx + 2a) + c) \arctan \left( \frac{\sqrt{2} \sqrt{\frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)}} + 1}{\sqrt{c}} \right)}{\sqrt{c}} + \frac{6 \sqrt{2}(c \cos(2bx + 2a) + c) \arctan \left( \frac{\sqrt{2} \sqrt{\frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)}}}{\sqrt{c}} \right)}{\sqrt{c}}$$

```
input integrate(1/(c*cot(b*x+a))^(5/2),x, algorithm="fricas")
```

output

```
1/12*(6*sqrt(2)*(c*cos(2*b*x + 2*a) + c)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))/sqrt(c) + 1)/sqrt(c) + 6*sqrt(2)*(c*cos(2*b*x + 2*a) + c)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)))/sqrt(c) - 1)/sqrt(c) + 3*sqrt(2)*(c*cos(2*b*x + 2*a) + c)*log((sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) + cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/sin(2*b*x + 2*a))/sqrt(c) - 3*sqrt(2)*(c*cos(2*b*x + 2*a) + c)*log(-(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) - cos(2*b*x + 2*a) - sin(2*b*x + 2*a) - 1)/sin(2*b*x + 2*a))/sqrt(c) - 8*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*(cos(2*b*x + 2*a) - 1)/(b*c^3*cos(2*b*x + 2*a) + b*c^3)
```

### Sympy [F]

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \int \frac{1}{(c \cot(a + bx))^{\frac{5}{2}}} dx$$

input

```
integrate(1/(c*cot(b*x+a))**(5/2), x)
```

output

```
Integral((c*cot(a + b*x))**(-5/2), x)
```

### Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.19

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \frac{c \left( 3 \left( \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}\sqrt{c} + 2\sqrt{\tan\frac{c}{bx+a}})}{2\sqrt{c}}\right)}{c^{\frac{3}{2}}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}\sqrt{c} - 2\sqrt{\tan\frac{c}{bx+a}})}{2\sqrt{c}}\right)}{c^{\frac{3}{2}}}\right) + \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\tan\frac{c}{bx+a}}\right)}{c^2} \right)}{c^2}$$

12b

input

```
integrate(1/(c*cot(b*x+a))^(5/2), x, algorithm="maxima")
```

output

```
1/12*c*(3*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(b*x + a)))/sqrt(c))/c^(3/2) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c))/c^(3/2) + sqrt(2)*log(sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/c^(3/2) - sqrt(2)*log(-sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/c^(3/2))/c^2 + 8/(c^2*(c/tan(b*x + a)^(3/2)))/b
```

**Giac [F]**

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \int \frac{1}{(c \cot(bx + a))^{\frac{5}{2}}} dx$$

input

```
integrate(1/(c*cot(b*x+a))^(5/2),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(-5/2), x)
```

**Mupad [B] (verification not implemented)**

Time = 9.08 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.49

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \frac{2}{3bc(c \cot(a + bx))^{3/2}} - \frac{(-1)^{1/4} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right) \operatorname{li}}{bc^{5/2}} - \frac{(-1)^{1/4} \operatorname{atanh}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a + bx)}}{\sqrt{c}}\right) \operatorname{li}}{bc^{5/2}}$$

input

```
int(1/(c*cot(a + b*x))^(5/2),x)
```

output

```
2/(3*b*c*(c*cot(a + b*x))^(3/2)) - ((-1)^(1/4)*atan((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2))*1i)/(b*c^(5/2)) - ((-1)^(1/4)*atanh((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2))*1i)/(b*c^(5/2))
```

**Reduce [F]**

$$\int \frac{1}{(c \cot(a + bx))^{5/2}} dx = \frac{\sqrt{c} \left( \int \frac{\sqrt{\cot(bx+a)}}{\cot(bx+a)^3} dx \right)}{c^3}$$

input `int(1/(c*cot(b*x+a))^(5/2),x)`

output `(sqrt(c)*int(sqrt(cot(a + b*x))/cot(a + b*x)**3,x))/c**3`

### 3.16 $\int \frac{1}{(c \cot(a+bx))^{7/2}} dx$

Optimal result	165
Mathematica [A] (verified)	165
Rubi [A] (warning: unable to verify)	166
Maple [A] (verified)	171
Fricas [B] (verification not implemented)	171
Sympy [F]	172
Maxima [A] (verification not implemented)	173
Giac [F]	173
Mupad [B] (verification not implemented)	174
Reduce [F]	174

#### Optimal result

Integrand size = 12, antiderivative size = 178

$$\int \frac{1}{(c \cot(a+bx))^{7/2}} dx = \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}bc^{7/2}} - \frac{\arctan\left(1 + \frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{\sqrt{2}bc^{7/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{c \cot(a+bx)}}{\sqrt{c} + \sqrt{c \cot(a+bx)}}\right)}{\sqrt{2}bc^{7/2}} + \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2}{bc^3\sqrt{c \cot(a+bx)}}$$

output

```
1/2*arctan(1-2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(7/2)-1/2*arctan(1+2^(1/2)*(c*cot(b*x+a))^(1/2)/c^(1/2))*2^(1/2)/b/c^(7/2)+1/2*arctanh(2^(1/2)*(c*cot(b*x+a))^(1/2)/(c^(1/2)+c^(1/2)*cot(b*x+a))*2^(1/2)/b/c^(7/2)+2/5/b/c/(c*cot(b*x+a))^(5/2)-2/b/c^3/(c*cot(b*x+a))^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.54

$$\int \frac{1}{(c \cot(a+bx))^{7/2}} dx = \frac{-5 \arctan\left(\sqrt[4]{-\cot^2(a+bx)}\right) \sqrt[4]{-\cot^2(a+bx)} + 5 \operatorname{arctanh}\left(\sqrt[4]{-\cot^2(a+bx)}\right)}{5bc^3\sqrt{c \cot(a+bx)}}$$

input `Integrate[(c*Cot[a + b*x])^(-7/2),x]`

output `(-5*ArcTan[(-Cot[a + b*x]^2)^(1/4)]*(-Cot[a + b*x]^2)^(1/4) + 5*ArcTanh[(-Cot[a + b*x]^2)^(1/4)]*(-Cot[a + b*x]^2)^(1/4) + 2*(-5 + Tan[a + b*x]^2))/(5*b*c^3*Sqrt[c*Cot[a + b*x]])`

### Rubi [A] (warning: unable to verify)

Time = 0.57 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.26, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$ , Rules used = {3042, 3955, 3042, 3955, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(c \cot(a + bx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{3955} \\
 & \frac{2}{5bc(c \cot(a + bx))^{5/2}} - \frac{\int \frac{1}{(c \cot(a + bx))^{3/2}} dx}{c^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2}{5bc(c \cot(a + bx))^{5/2}} - \frac{\int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{3/2}} dx}{c^2} \\
 & \quad \downarrow \text{3955} \\
 & \frac{2}{5bc(c \cot(a + bx))^{5/2}} - \frac{\frac{2}{bc\sqrt{c \cot(a + bx)}} - \frac{\int \sqrt{c \cot(a + bx)} dx}{c^2}}{c^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2}{bc\sqrt{c \cot(a+bx)}} - \frac{\int \sqrt{-c \tan(a+bx+\frac{\pi}{2})} dx}{c^2} \\
 & \quad \downarrow \text{3957} \\
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{\int \frac{\sqrt{c \cot(a+bx)}}{\cot^2(a+bx)c^2+c^2} d(c \cot(a+bx))}{bc} + \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \quad \downarrow \text{266} \\
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2 \int \frac{c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)}}{bc} + \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \quad \downarrow \text{826} \\
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2 \left( \frac{1}{2} \int \frac{c^2 \cot^2(a+bx)+c}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} \right)}{bc} + \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \quad \downarrow \text{1476} \\
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)} + \frac{1}{2} \int \frac{1}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2} \cot(a+bx)+c} d\sqrt{c \cot(a+bx)} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} \right)}{bc} + \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \quad \downarrow \text{1082} \\
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2 \left( \frac{1}{2} \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx)-1} d(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx)-1} d(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} \right)}{bc} + \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \quad \downarrow \text{217} \\
 & \frac{2}{5bc(c \cot(a+bx))^{5/2}} - \frac{2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c} \cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c} \cot(a+bx))}{\sqrt{2}\sqrt{c}} \right) - \frac{1}{2} \int \frac{c-c^2 \cot^2(a+bx)}{c^4 \cot^4(a+bx)+c^2} d\sqrt{c \cot(a+bx)} \right)}{bc} + \frac{2}{bc\sqrt{c \cot(a+bx)}} \\
 & \quad \downarrow \\
 & \frac{2}{c^2}
 \end{aligned}$$



$$\begin{array}{c} \downarrow 1479 \\ 2 \\ \frac{5bc(c \cot(a + bx))^{5/2}}{2} \\ \hline 2 \left( \frac{1}{2} \left( \frac{\int -\frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} + \frac{\int -\frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right) \right) \\ \hline bc \\ \hline c^2 \end{array}$$

$$\begin{array}{c} \downarrow 25 \\ 2 \\ \frac{5bc(c \cot(a + bx))^{5/2}}{2} \\ \hline 2 \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{2}(\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx))}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right) \right) \\ \hline bc \\ \hline c^2 \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ 2 \\ \frac{5bc(c \cot(a + bx))^{5/2}}{2} \\ \hline 2 \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}\sqrt{c}-2\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)-\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{2}\sqrt{c}} - \frac{\int \frac{\sqrt{c}+\sqrt{2}\sqrt{c}\cot(a+bx)}{c^2 \cot^2(a+bx)+\sqrt{2}c^{3/2}\cot(a+bx)+c} d\sqrt{c}\cot(a+bx)}{2\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} \right) \right) \\ \hline bc \\ \hline c^2 \end{array}$$

$$\begin{array}{c} \downarrow 1103 \\ 2 \\ \frac{5bc(c \cot(a + bx))^{5/2}}{2} \\ \hline 2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{c}\cot(a+bx)+1)}{\sqrt{2}\sqrt{c}} - \frac{\arctan(1-\sqrt{2}\sqrt{c}\cot(a+bx))}{\sqrt{2}\sqrt{c}} \right) + \frac{1}{2} \left( \frac{\log(-\sqrt{2}c^{3/2}\cot(a+bx)+c^2 \cot^2(a+bx)+c)}{2\sqrt{2}\sqrt{c}} - \frac{\log(\sqrt{2}c^{3/2}\cot(a+bx)+c^2 \cot^2(a+bx))}{2\sqrt{2}\sqrt{c}} \right) \right) \\ \hline bc \\ \hline c^2 \end{array}$$

input `Int[(c*Cot[a + b*x])^(-7/2),x]`

output

$$\frac{2}{5bc}(c \cot[a + bx])^{5/2} - \frac{2}{b^2c} \sqrt{c \cot[a + bx]} + \frac{2}{c} \left( -\operatorname{ArcTan}\left[\frac{1 - \sqrt{2} \sqrt{c \cot[a + bx]}}{\sqrt{2} \sqrt{c}}\right] + \operatorname{ArcTan}\left[\frac{1 + \sqrt{2} \sqrt{c \cot[a + bx]}}{\sqrt{2} \sqrt{c}}\right] \right) / 2 + \frac{\log[c - \sqrt{2} c^{3/2} \cot[a + bx] + c^2 \cot[a + bx]^2]}{2 \sqrt{2} \sqrt{c}} - \frac{\log[c + \sqrt{2} c^{3/2} \cot[a + bx] + c^2 \cot[a + bx]^2]}{2 \sqrt{2} \sqrt{c}} \Big/ (bc)^2$$

### Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$$

rule 27

$$\operatorname{Int}[(a_*)(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[F_x, (b_*)(G_x)] /; \operatorname{FreeQ}[b, x]$$

rule 217

$$\operatorname{Int}[(a_*) + (b_*)(x_*)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2] \operatorname{Rt}[-b, 2])^{-1} \operatorname{ArcTan}[\operatorname{Rt}[-b, 2] (x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b] \ \&\& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$$

rule 266

$$\operatorname{Int}[(c_*)(x_*)^m ((a_*) + (b_*)(x_*)^2)^p], x\_Symbol] \rightarrow \operatorname{With}[\{k = \operatorname{Denominator}[m]\}, \operatorname{Simp}[k/c \operatorname{Subst}[\operatorname{Int}[x^{k(m+1)-1} (a + b(x^{2k}/c^2))^p], x], (c*x)^{1/k}], x] /; \operatorname{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \operatorname{FractionQ}[m] \ \&\& \ \operatorname{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 826

$$\operatorname{Int}[(x_*)^2 / ((a_*) + (b_*)(x_*)^4), x\_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numerator}[\operatorname{Rt}[a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[a/b, 2]]\}, \operatorname{Simp}[1/(2*s) \operatorname{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \operatorname{Simp}[1/(2*s) \operatorname{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ (\operatorname{GtQ}[a/b, 0] \ || \ (\operatorname{PosQ}[a/b] \ \&\& \ \operatorname{AtomQ}[\operatorname{SplitProduct}[\operatorname{SumBaseQ}, a]] \ \&\& \ \operatorname{AtomQ}[\operatorname{SplitProduct}[\operatorname{SumBaseQ}, b]]))$$

rule 1082

$$\operatorname{Int}[(a_*) + (b_*)(x_*) + (c_*)(x_*)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{With}[\{q = 1 - 4*S \operatorname{implify}[a*(c/b^2)]\}, \operatorname{Simp}[-2/b \operatorname{Subst}[\operatorname{Int}[1/(q - x^2), x], x], 1 + 2*c*(x/b)], x] /; \operatorname{RationalQ}[q] \ \&\& \ (\operatorname{EqQ}[q^2, 1] \ || \ !\operatorname{RationalQ}[b^2 - 4*a*c]) /; \operatorname{FreeQ}[\{a, b, c\}, x]$$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.96

method	result
derivativedivides	$2c \frac{\left( \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( -\frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) \right)}{8c^4 (c^2)^{\frac{1}{4}}}$
default	$2c \frac{\left( \sqrt{2} \left( \ln \left( \frac{c \cot(bx+a) - (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}}{c \cot(bx+a) + (c^2)^{\frac{1}{4}} \sqrt{c \cot(bx+a)} \sqrt{2} + \sqrt{c^2}} \right) + 2 \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) - 2 \arctan \left( -\frac{\sqrt{2} \sqrt{c \cot(bx+a)}}{(c^2)^{\frac{1}{4}}} + 1 \right) \right)}{8c^4 (c^2)^{\frac{1}{4}}}$

```
input int(1/(c*cot(b*x+a))^(7/2),x,method=_RETURNVERBOSE)
```

```
output -2/b*c*(1/8/c^4/(c^2)^(1/4)*2^(1/2)*(ln((c*cot(b*x+a)-(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))/(c*cot(b*x+a)+(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)*2^(1/2)+(c^2)^(1/2)))+2*arctan(2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1)-2*arctan(-2^(1/2)/(c^2)^(1/4)*(c*cot(b*x+a))^(1/2)+1))-1/5/c^2/(c*cot(b*x+a))^(5/2)+1/c^4/(c*cot(b*x+a))^(1/2))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 464 vs. 2(142) = 284.

Time = 0.09 (sec) , antiderivative size = 464, normalized size of antiderivative = 2.61

$$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx =$$

$$16 \sqrt{\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}} (3 \cos(2bx+2a) + 2) \sin(2bx+2a) + \frac{10 \sqrt{2} (c \cos(2bx+2a)^2 + 2c \cos(2bx+2a) + c) \arctan \left( \frac{\sqrt{2} \sqrt{c \cot(a+bx)}}{(c^2)^{\frac{1}{4}}} + 1 \right)}{\sqrt{c}}$$

```
input integrate(1/(c*cot(b*x+a))^(7/2),x, algorithm="fricas")
```

output

```
-1/20*(16*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*(3*cos(2*b*x + 2*a) + 2)*sin(2*b*x + 2*a) + 10*sqrt(2)*(c*cos(2*b*x + 2*a)^2 + 2*c*cos(2*b*x + 2*a) + c)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))/sqrt(c) + 1)/sqrt(c) + 10*sqrt(2)*(c*cos(2*b*x + 2*a)^2 + 2*c*cos(2*b*x + 2*a) + c)*arctan(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a)))/sqrt(c) - 1)/sqrt(c) - 5*sqrt(2)*(c*cos(2*b*x + 2*a)^2 + 2*c*cos(2*b*x + 2*a) + c)*log((sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) + cos(2*b*x + 2*a) + sin(2*b*x + 2*a) + 1)/sin(2*b*x + 2*a))/sqrt(c) + 5*sqrt(2)*(c*cos(2*b*x + 2*a)^2 + 2*c*cos(2*b*x + 2*a) + c)*log(-(sqrt(2)*sqrt((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))*sin(2*b*x + 2*a)/sqrt(c) - cos(2*b*x + 2*a) - sin(2*b*x + 2*a) - 1)/sin(2*b*x + 2*a))/sqrt(c))/(b*c^4*cos(2*b*x + 2*a)^2 + 2*b*c^4*cos(2*b*x + 2*a) + b*c^4)
```

### Sympy [F]

$$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx = \int \frac{1}{(c \cot(a + bx))^{\frac{7}{2}}} dx$$

input

```
integrate(1/(c*cot(b*x+a))**(7/2), x)
```

output

```
Integral((c*cot(a + b*x))**(-7/2), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.15

$$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx =$$

$$\frac{5 \left( \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}\sqrt{c}+2\sqrt{\tan(bx+a)})}{2\sqrt{c}}\right)}{\sqrt{c}} + \frac{2\sqrt{2} \arctan\left(\frac{-\sqrt{2}(\sqrt{2}\sqrt{c}-2\sqrt{\tan(bx+a)})}{2\sqrt{c}}\right)}{\sqrt{c}} \right) - \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\frac{c}{\tan(bx+a)}+c+\frac{c}{\tan(bx+a)}}\right) + \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\frac{c}{\tan(bx+a)}-c+\frac{c}{\tan(bx+a)}}\right)}{c^4}}{c^4} + \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\frac{c}{\tan(bx+a)}+c+\frac{c}{\tan(bx+a)}}\right) + \frac{\sqrt{2} \log\left(\sqrt{2}\sqrt{c}\sqrt{\frac{c}{\tan(bx+a)}-c+\frac{c}{\tan(bx+a)}}\right)}{c^4}}{20b}$$

input `integrate(1/(c*cot(b*x+a))^(7/2),x, algorithm="maxima")`

output `-1/20*c*(5*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(c) + 2*sqrt(c/tan(b*x + a)))/sqrt(c))/sqrt(c) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(c) - 2*sqrt(c/tan(b*x + a)))/sqrt(c))/sqrt(c) - sqrt(2)*log(sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c) + sqrt(2)*log(-sqrt(2)*sqrt(c)*sqrt(c/tan(b*x + a)) + c + c/tan(b*x + a))/sqrt(c))/c^4 - 8*(c^2 - 5*c^2/tan(b*x + a)^2)/(c^4*(c/tan(b*x + a))^(5/2)))/b`

**Giac [F]**

$$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx = \int \frac{1}{(c \cot(bx + a))^{\frac{7}{2}}} dx$$

input `integrate(1/(c*cot(b*x+a))^(7/2),x, algorithm="giac")`

output `integrate((c*cot(b*x + a))^(-7/2), x)`

**Mupad [B] (verification not implemented)**

Time = 9.34 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.51

$$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx = \frac{\frac{2}{5c} - \frac{2 \cot(a+bx)^2}{c}}{b (c \cot(a + bx))^{5/2}} - \frac{(-1)^{1/4} \operatorname{atan}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{b c^{7/2}} + \frac{(-1)^{1/4} \operatorname{atanh}\left(\frac{(-1)^{1/4} \sqrt{c \cot(a+bx)}}{\sqrt{c}}\right)}{b c^{7/2}}$$

input `int(1/(c*cot(a + b*x))^(7/2),x)`output `(2/(5*c) - (2*cot(a + b*x)^2)/c)/(b*(c*cot(a + b*x))^(5/2)) - ((-1)^(1/4)*atan((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2)))/(b*c^(7/2)) + ((-1)^(1/4)*atanh((( -1)^(1/4)*(c*cot(a + b*x))^(1/2))/c^(1/2)))/(b*c^(7/2))`**Reduce [F]**

$$\int \frac{1}{(c \cot(a + bx))^{7/2}} dx = \frac{\sqrt{c} \left( \int \frac{\sqrt{\cot(bx+a)}}{\cot(bx+a)^4} dx \right)}{c^4}$$

input `int(1/(c*cot(b*x+a))^(7/2),x)`output `(sqrt(c)*int(sqrt(cot(a + b*x))/cot(a + b*x)**4,x))/c**4`

### 3.17 $\int (c \cot(a + bx))^{4/3} dx$

Optimal result	175
Mathematica [C] (verified)	176
Rubi [A] (warning: unable to verify)	176
Maple [A] (verified)	181
Fricas [B] (verification not implemented)	181
Sympy [F]	182
Maxima [A] (verification not implemented)	182
Giac [F]	183
Mupad [B] (verification not implemented)	183
Reduce [F]	184

#### Optimal result

Integrand size = 12, antiderivative size = 242

$$\int (c \cot(a + bx))^{4/3} dx = \frac{c^{4/3} \arctan\left(\frac{\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{b} - \frac{c^{4/3} \arctan\left(\sqrt{3} - \frac{2\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{2b} + \frac{c^{4/3} \arctan\left(\sqrt{3} + \frac{2\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{2b} - \frac{3c\sqrt[3]{c \cot(a + bx)}}{b} - \frac{\sqrt{3}c^{4/3} \log\left(c^{2/3} - \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a + bx)} + (c \cot(a + bx))^{2/3}\right)}{4b} + \frac{\sqrt{3}c^{4/3} \log\left(c^{2/3} + \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a + bx)} + (c \cot(a + bx))^{2/3}\right)}{4b}$$

output

```
c^(4/3)*arctan((c*cot(b*x+a))^(1/3)/c^(1/3))/b+1/2*c^(4/3)*arctan(-3^(1/2)
+2*(c*cot(b*x+a))^(1/3)/c^(1/3))/b+1/2*c^(4/3)*arctan(3^(1/2)+2*(c*cot(b*x
+a))^(1/3)/c^(1/3))/b-3*c*(c*cot(b*x+a))^(1/3)/b-1/4*3^(1/2)*c^(4/3)*ln(c^
(2/3)-3^(1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b+1/4*3^(
1/2)*c^(4/3)*ln(c^(2/3)+3^(1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a)
)^(2/3))/b
```



**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.85

$$\int (c \cot(a + bx))^{4/3} dx =$$

$$\frac{c^3 \sqrt[3]{c \cot(a + bx)} \left( 6 \sqrt[6]{\cot^2(a + bx)} - i \log \left( 1 - i \sqrt[6]{\cot^2(a + bx)} \right) + i \log \left( 1 + i \sqrt[6]{\cot^2(a + bx)} \right) - (-1) \right)}{-}$$

input `Integrate[(c*Cot[a + b*x])^(4/3),x]`

output

```
-1/2*(c*(c*Cot[a + b*x])^(1/3)*(6*(Cot[a + b*x]^2)^(1/6) - I*Log[1 - I*(Cot[a + b*x]^2)^(1/6)] + I*Log[1 + I*(Cot[a + b*x]^2)^(1/6)] - (-1)^(5/6)*Log[1 - (-1)^(1/6)*(Cot[a + b*x]^2)^(1/6)] + (-1)^(5/6)*Log[1 + (-1)^(1/6)*(Cot[a + b*x]^2)^(1/6)] - (-1)^(1/6)*Log[1 - (-1)^(5/6)*(Cot[a + b*x]^2)^(1/6)] + (-1)^(1/6)*Log[1 + (-1)^(5/6)*(Cot[a + b*x]^2)^(1/6)]))/(b*(Cot[a + b*x]^2)^(1/6))
```

**Rubi [A] (warning: unable to verify)**

Time = 0.51 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.90, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$ , Rules used = {3042, 3954, 3042, 3957, 266, 753, 27, 216, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c \cot(a + bx))^{4/3} dx$$

$$\downarrow \text{3042}$$

$$\int \left( -c \tan \left( a + bx + \frac{\pi}{2} \right) \right)^{4/3} dx$$

$$\downarrow \text{3954}$$

$$\begin{aligned}
 & c^2 \left( - \int \frac{1}{(c \cot(a + bx))^{2/3}} dx \right) - \frac{3c \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{3042} \\
 & c^2 \left( - \int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{2/3}} dx \right) - \frac{3c \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{3957} \\
 & \frac{c^3 \int \frac{1}{(c \cot(a + bx))^{2/3} (\cot^2(a + bx) c^2 + c^2)} d(c \cot(a + bx))}{b} - \frac{3c \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{266} \\
 & \frac{3c^3 \int \frac{1}{c^6 \cot^6(a + bx) + c^2} d \sqrt[3]{c \cot(a + bx)}}{b} - \frac{3c \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{753} \\
 & 3c^3 \left( \frac{\int \frac{1}{c^2 \cot^2(a + bx) + c^{2/3}} d \sqrt[3]{c \cot(a + bx)}}{3c^{4/3}} + \frac{\int \frac{2 \sqrt[3]{c - \sqrt{3}} \sqrt[3]{c \cot(a + bx)}}{2(c^2 \cot^2(a + bx) - \sqrt{3} c^{4/3} \cot(a + bx) + c^{2/3})} d \sqrt[3]{c \cot(a + bx)}}{3c^{5/3}} + \frac{\int \frac{2 \sqrt[3]{c + \sqrt{3}} \sqrt[3]{c \cot(a + bx)}}{2(c^2 \cot^2(a + bx) + \sqrt{3} c^{4/3} \cot(a + bx) + c^{2/3})} d \sqrt[3]{c \cot(a + bx)}}{3c^{5/3}} \right) \\
 & \quad \hrule \\
 & \frac{3c \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{27} \\
 & 3c^3 \left( \frac{\int \frac{1}{c^2 \cot^2(a + bx) + c^{2/3}} d \sqrt[3]{c \cot(a + bx)}}{3c^{4/3}} + \frac{\int \frac{2 \sqrt[3]{c - \sqrt{3}} \sqrt[3]{c \cot(a + bx)}}{c^2 \cot^2(a + bx) - \sqrt{3} c^{4/3} \cot(a + bx) + c^{2/3}} d \sqrt[3]{c \cot(a + bx)}}{6c^{5/3}} + \frac{\int \frac{2 \sqrt[3]{c + \sqrt{3}} \sqrt[3]{c \cot(a + bx)}}{c^2 \cot^2(a + bx) + \sqrt{3} c^{4/3} \cot(a + bx) + c^{2/3}} d \sqrt[3]{c \cot(a + bx)}}{6c^{5/3}} \right) \\
 & \quad \hrule \\
 & \frac{3c \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{216} \\
 & 3c^3 \left( \frac{\int \frac{2 \sqrt[3]{c - \sqrt{3}} \sqrt[3]{c \cot(a + bx)}}{c^2 \cot^2(a + bx) - \sqrt{3} c^{4/3} \cot(a + bx) + c^{2/3}} d \sqrt[3]{c \cot(a + bx)}}{6c^{5/3}} + \frac{\int \frac{2 \sqrt[3]{c + \sqrt{3}} \sqrt[3]{c \cot(a + bx)}}{c^2 \cot^2(a + bx) + \sqrt{3} c^{4/3} \cot(a + bx) + c^{2/3}} d \sqrt[3]{c \cot(a + bx)}}{6c^{5/3}} + \frac{\arctan(\dots)}{b} \right) \\
 & \quad \hrule \\
 & \frac{3c \sqrt[3]{c \cot(a + bx)}}{b}
 \end{aligned}$$

↓ 1142

$$3c^3 \left( \frac{\frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c} - 2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} \right)$$

$$\frac{3c^3 \sqrt[3]{c \cot(a+bx)}}{b}$$

↓ 25

$$3c^3 \left( \frac{\frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)} + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c} - 2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} \right) +$$

$$\frac{3c^3 \sqrt[3]{c \cot(a+bx)}}{b}$$

↓ 1082

$$3c^3 \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - \frac{1}{3}} d \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c} - 2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} \right) + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c} + 2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}$$

$$\frac{3c^3 \sqrt[3]{c \cot(a+bx)}}{b}$$

↓ 217

$$3c^3 \left( \frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c} - 2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \arctan \left( \sqrt{3} \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right) \right)}{6c^{5/3}} \right) + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c} + 2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}$$

$$\frac{3c^3 \sqrt[3]{c \cot(a+bx)}}{b}$$

↓ 1103

$$\frac{3c^3 \left( \frac{\arctan\left(\frac{c^{2/3} \cot(a+bx)}{3c^{5/3}}\right)}{3c^{5/3}} + \frac{-\arctan\left(\sqrt{3}\left(1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}}\right)\right) - \frac{1}{2}\sqrt{3} \log\left(-\sqrt{3}c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3}\right)}{6c^{5/3}} + \frac{\arctan\left(\sqrt{3}\left(1 + \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}}\right)\right) - \frac{1}{2}\sqrt{3} \log\left(-\sqrt{3}c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3}\right)}{6c^{5/3}}}{b} \right)}{3c^3 \sqrt[3]{c \cot(a+bx)}}$$

input `Int[(c*Cot[a + b*x])^(4/3), x]`

output `(-3*c*(c*Cot[a + b*x])^(1/3))/b + (3*c^3*(ArcTan[c^(2/3)*Cot[a + b*x]]/(3*c^(5/3)) + (-ArcTan[Sqrt[3]*(1 - (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]]) - (Sqrt[3]*Log[c^(2/3) - Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(5/3)) + (ArcTan[Sqrt[3]*(1 + (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]]) + (Sqrt[3]*Log[c^(2/3) + Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(5/3)))/b`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 753 `Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u, v}, Simp[u = Int[(r - s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; 2*(r^2/(a*n)) Int[1/(r^2 + s^2*x^2), x] + 2*(r/(a*n)) Sum[u, {k, 1, (n - 2)/4}], x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && PosQ[a/b]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

**Maple [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.88

method	result
derivativedivides	$-\frac{3c(c \cot(bx+a))^{\frac{1}{3}}}{b} - \frac{c\sqrt{3}(c^2)^{\frac{1}{6}} \ln\left((c \cot(bx+a))^{\frac{2}{3}} - \sqrt{3}(c^2)^{\frac{1}{6}}(c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}}\right)}{4b} + \frac{c(c^2)^{\frac{1}{6}} \arctan\left(\frac{2(c \cot(bx+a))^{\frac{1}{3}}}{(c^2)^{\frac{1}{6}} - \sqrt{3}(c \cot(bx+a))^{\frac{1}{3}}}\right)}{2b}$
default	$-\frac{3c(c \cot(bx+a))^{\frac{1}{3}}}{b} - \frac{c\sqrt{3}(c^2)^{\frac{1}{6}} \ln\left((c \cot(bx+a))^{\frac{2}{3}} - \sqrt{3}(c^2)^{\frac{1}{6}}(c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}}\right)}{4b} + \frac{c(c^2)^{\frac{1}{6}} \arctan\left(\frac{2(c \cot(bx+a))^{\frac{1}{3}}}{(c^2)^{\frac{1}{6}} - \sqrt{3}(c \cot(bx+a))^{\frac{1}{3}}}\right)}{2b}$

input

```
int((c*cot(b*x+a))^(4/3),x,method=_RETURNVERBOSE)
```

output

```
-3*c*(c*cot(b*x+a))^(1/3)/b-1/4/b*c*3^(1/2)*(c^2)^(1/6)*ln((c*cot(b*x+a))^(2/3)-3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/2/b*c*(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)-3^(1/2))+1/b*c*(c^2)^(1/6)*arctan((c*cot(b*x+a))^(1/3)/(c^2)^(1/6))+1/4/b*c*3^(1/2)*(c^2)^(1/6)*ln((c*cot(b*x+a))^(2/3)+3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/2/b*c*(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)+3^(1/2))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 431 vs. 2(184) = 368.

Time = 0.10 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.78

$$\int (c \cot(a + bx))^{4/3} dx = \frac{\left(-\frac{c^8}{b^6}\right)^{\frac{1}{6}} (\sqrt{-3b + b}) \log\left(c\left(\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}\right)^{\frac{1}{3}} + \frac{1}{2}\left(-\frac{c^8}{b^6}\right)^{\frac{1}{6}} (\sqrt{-3b + b})\right) - \left(-\frac{c^8}{b^6}\right)^{\frac{1}{6}} (\sqrt{-3b + b})}{4b}$$

input `integrate((c*cot(b*x+a))^(4/3),x, algorithm="fricas")`

output `1/4*((-c^8/b^6)^(1/6)*(sqrt(-3)*b + b)*log(c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) + 1/2*(-c^8/b^6)^(1/6)*(sqrt(-3)*b + b)) - (-c^8/b^6)^(1/6)*(sqrt(-3)*b + b)*log(c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) - 1/2*(-c^8/b^6)^(1/6)*(sqrt(-3)*b + b)) + (-c^8/b^6)^(1/6)*(sqrt(-3)*b - b)*log(c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) + 1/2*(-c^8/b^6)^(1/6)*(sqrt(-3)*b - b)) - (-c^8/b^6)^(1/6)*(sqrt(-3)*b - b)*log(c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) - 1/2*(-c^8/b^6)^(1/6)*(sqrt(-3)*b - b)) + 2*(-c^8/b^6)^(1/6)*b*log(c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) + (-c^8/b^6)^(1/6)*b) - 2*(-c^8/b^6)^(1/6)*b*log(c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) - (-c^8/b^6)^(1/6)*b) - 1/2*c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3))/b`

### Sympy [F]

$$\int (c \cot(a + bx))^{4/3} dx = \int (c \cot(a + bx))^{4/3} dx$$

input `integrate((c*cot(b*x+a))**(4/3),x)`

output `Integral((c*cot(a + b*x))**(4/3), x)`

### Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.81

$$\int (c \cot(a + bx))^{4/3} dx = \frac{\left( \sqrt{3}c^{1/3} \log \left( \sqrt{3}c^{1/3} \left( \frac{c}{\tan(bx+a)} \right)^{1/3} + c^{2/3} + \left( \frac{c}{\tan(bx+a)} \right)^{2/3} \right) - \sqrt{3}c^{1/3} \log \left( -\sqrt{3}c^{1/3} \left( \frac{c}{\tan(bx+a)} \right)^{1/3} + c^{2/3} \right) \right)}{3}$$

input `integrate((c*cot(b*x+a))^(4/3),x, algorithm="maxima")`

output

```
1/4*(sqrt(3)*c^(1/3)*log(sqrt(3)*c^(1/3)*(c/tan(b*x + a))^(1/3) + c^(2/3)
+ (c/tan(b*x + a))^(2/3)) - sqrt(3)*c^(1/3)*log(-sqrt(3)*c^(1/3)*(c/tan(b*
x + a))^(1/3) + c^(2/3) + (c/tan(b*x + a))^(2/3)) + 2*c^(1/3)*arctan((sqrt
(3)*c^(1/3) + 2*(c/tan(b*x + a))^(1/3))/c^(1/3)) + 2*c^(1/3)*arctan(-(sqrt
(3)*c^(1/3) - 2*(c/tan(b*x + a))^(1/3))/c^(1/3)) + 4*c^(1/3)*arctan((c/tan
(b*x + a))^(1/3)/c^(1/3)) - 12*(c/tan(b*x + a))^(1/3))*c/b
```

**Giac [F]**

$$\int (c \cot(a + bx))^{4/3} dx = \int (c \cot(bx + a))^{4/3} dx$$

input

```
integrate((c*cot(b*x+a))^(4/3),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(4/3), x)
```

**Mupad [B] (verification not implemented)**

Time = 10.29 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.02

$$\begin{aligned} \int (c \cot(a + bx))^{4/3} dx &= -\frac{3c(c \cot(a + bx))^{1/3}}{b} \\ &+ \frac{(-1)^{1/6} c^{4/3} \operatorname{atan}\left(\frac{(-1)^{5/6} (c \cot(a + bx))^{1/3} \operatorname{li}}{c^{1/3}}\right) \operatorname{li}}{b} \\ &- \frac{(-1)^{1/6} c^{4/3} \ln\left((-1)^{1/6} c^{1/3} - 2(c \cot(a + bx))^{1/3} + (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2b} \\ &- \frac{(-1)^{1/6} c^{4/3} \ln\left(2(c \cot(a + bx))^{1/3} + (-1)^{1/6} c^{1/3} - (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2b} \\ &+ \frac{(-1)^{1/6} c^{4/3} \ln\left(2(c \cot(a + bx))^{1/3} + (-1)^{1/6} c^{1/3} + (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b} \\ &+ \frac{(-1)^{1/6} c^{4/3} \ln\left(2(c \cot(a + bx))^{1/3} - (-1)^{1/6} c^{1/3} + (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(-\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b} \end{aligned}$$



input `int((c*cot(a + b*x))^(4/3),x)`

output `((-1)^(1/6)*c^(4/3)*atan((( -1)^(5/6)*(c*cot(a + b*x))^(1/3)*1i)/c^(1/3))*1i)/b - (3*c*(c*cot(a + b*x))^(1/3))/b - ((-1)^(1/6)*c^(4/3)*log((-1)^(1/6)*c^(1/3) - 2*(c*cot(a + b*x))^(1/3) + (-1)^(2/3)*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/2 + 1/2))/(2*b) - ((-1)^(1/6)*c^(4/3)*log(2*(c*cot(a + b*x))^(1/3) + (-1)^(1/6)*c^(1/3) - (-1)^(2/3)*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/2 - 1/2))/(2*b) + ((-1)^(1/6)*c^(4/3)*log(2*(c*cot(a + b*x))^(1/3) + (-1)^(1/6)*c^(1/3) + (-1)^(2/3)*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/4 + 1/4))/b + ((-1)^(1/6)*c^(4/3)*log(2*(c*cot(a + b*x))^(1/3) - (-1)^(1/6)*c^(1/3) + (-1)^(2/3)*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/4 - 1/4))/b`

### Reduce [F]

$$\int (c \cot(a + bx))^{4/3} dx = \frac{c^{4/3} \left( -3 \cot(bx + a)^{1/3} - \left( \int \frac{1}{\cot(bx+a)^{2/3}} dx \right) b \right)}{b}$$

input `int((c*cot(b*x+a))^(4/3),x)`

output `(c**(1/3)*c*( - 3*cot(a + b*x)**(1/3) - int(cot(a + b*x)**(1/3)/cot(a + b*x),x)*b))/b`

### 3.18 $\int (c \cot(a + bx))^{2/3} dx$

Optimal result	185
Mathematica [C] (verified)	186
Rubi [A] (warning: unable to verify)	186
Maple [A] (verified)	190
Fricas [B] (verification not implemented)	191
Sympy [F]	192
Maxima [A] (verification not implemented)	192
Giac [F]	193
Mupad [B] (verification not implemented)	193
Reduce [F]	194

#### Optimal result

Integrand size = 12, antiderivative size = 225

$$\int (c \cot(a + bx))^{2/3} dx = -\frac{c^{2/3} \arctan\left(\frac{\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{b} + \frac{c^{2/3} \arctan\left(\sqrt{3} - \frac{2\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{2b} - \frac{c^{2/3} \arctan\left(\sqrt{3} + \frac{2\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{2b} - \frac{\sqrt{3}c^{2/3} \log\left(c^{2/3} - \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a + bx)} + (c \cot(a + bx))^{2/3}\right)}{4b} + \frac{\sqrt{3}c^{2/3} \log\left(c^{2/3} + \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a + bx)} + (c \cot(a + bx))^{2/3}\right)}{4b}$$

output

```
-c^(2/3)*arctan((c*cot(b*x+a))^(1/3)/c^(1/3))/b-1/2*c^(2/3)*arctan(-3^(1/2)
)+2*(c*cot(b*x+a))^(1/3)/c^(1/3))/b-1/2*c^(2/3)*arctan(3^(1/2)+2*(c*cot(b*
x+a))^(1/3)/c^(1/3))/b-1/4*3^(1/2)*c^(2/3)*ln(c^(2/3)-3^(1/2)*c^(1/3)*(c*c
ot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b+1/4*3^(1/2)*c^(2/3)*ln(c^(2/3)+3^(
1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.82

$$\int (c \cot(a + bx))^{2/3} dx = \frac{(c \cot(a + bx))^{5/3} \left( -i \log \left( 1 - i \sqrt[6]{\cot^2(a + bx)} \right) + i \log \left( 1 + i \sqrt[6]{\cot^2(a + bx)} \right) + \sqrt[6]{-1} \left( - \right) \right)}{2 * b * c * (\cot[a + b * x]^2)^{(5/6)}}$$

input `Integrate[(c*Cot[a + b*x])^(2/3),x]`

output `((Cot[a + b*x])^(5/3)*((-I)*Log[1 - I*(Cot[a + b*x]^2)^(1/6)] + I*Log[1 + I*(Cot[a + b*x]^2)^(1/6)] + (-1)^(1/6)*(-Log[1 - (-1)^(1/6)*(Cot[a + b*x]^2)^(1/6)] + Log[1 + (-1)^(1/6)*(Cot[a + b*x]^2)^(1/6)] + (-1)^(2/3)*(-Log[1 - (-1)^(5/6)*(Cot[a + b*x]^2)^(1/6)] + Log[1 + (-1)^(5/6)*(Cot[a + b*x]^2)^(1/6)])))/(2*b*c*(Cot[a + b*x]^2)^(5/6))`

**Rubi [A] (warning: unable to verify)**

Time = 0.43 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.87, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3957, 266, 824, 27, 216, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (c \cot(a + bx))^{2/3} dx$$

$$\downarrow \text{3042}$$

$$\int \left( -c \tan \left( a + bx + \frac{\pi}{2} \right) \right)^{2/3} dx$$

$$\downarrow \text{3957}$$

$$\frac{c \int \frac{(c \cot(a+bx))^{2/3}}{\cot^2(a+bx)c^2+c^2} d(c \cot(a+bx))}{b}$$

↓ 266

$$\frac{3c \int \frac{c^4 \cot^4(a+bx)}{c^6 \cot^6(a+bx)+c^2} d^3 \sqrt{c \cot(a+bx)}}{b}$$

↓ 824

---


$$3c \left( \frac{1}{3} \int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} + \frac{\int -\frac{\sqrt[3]{c}-\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{2(c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3})} d^3 \sqrt{c \cot(a+bx)}}{3\sqrt[3]{c}} + \frac{\int -\frac{\sqrt[3]{c}+\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{2(c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3})} d^3 \sqrt{c \cot(a+bx)}}{3\sqrt[3]{c}} \right)$$

↓ 27

---


$$3c \left( \frac{1}{3} \int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{\int \frac{\sqrt[3]{c}-\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6\sqrt[3]{c}} - \frac{\int \frac{\sqrt[3]{c}+\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6\sqrt[3]{c}} \right)$$

↓ 216

---


$$3c \left( -\frac{\int \frac{\sqrt[3]{c}-\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6\sqrt[3]{c}} - \frac{\int \frac{\sqrt[3]{c}+\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6\sqrt[3]{c}} + \arctan \frac{\sqrt[3]{c}-\sqrt{3} \sqrt[3]{c \cot(a+bx)}}{\sqrt[3]{c}+\sqrt{3} \sqrt[3]{c \cot(a+bx)}} \right)$$

↓ 1142

---


$$3c \left( -\frac{\frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{1}{2} \sqrt{3} \int -\frac{\sqrt{3} \sqrt[3]{c}-2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6\sqrt[3]{c}} \right)$$

↓ 25

---


$$3c \left( -\frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c}-2 \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6\sqrt[3]{c}} \right)$$

$$\begin{aligned}
 & \downarrow 1082 \\
 & 3c \left( \frac{\frac{1}{2}\sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} dx \sqrt[3]{c \cot(a+bx)} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - \frac{1}{3}} d \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - \frac{1}{3}}}{\sqrt{3}}}{6 \sqrt[3]{c}} \right) \\
 & \downarrow 217 \\
 & 3c \left( \frac{\frac{1}{2}\sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} dx \sqrt[3]{c \cot(a+bx)} + \arctan \left( \sqrt{3} \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right) \right)}{6 \sqrt[3]{c}} - \frac{\frac{1}{2}\sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c+2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} dx \sqrt[3]{c \cot(a+bx)}}{6 \sqrt[3]{c}} \right) \\
 & \downarrow 1103 \\
 & 3c \left( \frac{\arctan(c^{2/3} \cot(a+bx))}{3 \sqrt[3]{c}} - \frac{\arctan \left( \sqrt{3} \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right) \right) - \frac{1}{2}\sqrt{3} \log \left( -\sqrt{3}c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3} \right)}{6 \sqrt[3]{c}} - \frac{\frac{1}{2}\sqrt{3} \log \left( \sqrt{3}c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3} \right)}{6 \sqrt[3]{c}} \right)
 \end{aligned}$$

input `Int[(c*Cot[a + b*x])^(2/3),x]`

output `(-3*c*(ArcTan[c^(2/3)*Cot[a + b*x]]/(3*c^(1/3)) - (ArcTan[Sqrt[3]*(1 - (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]]) - (Sqrt[3]*Log[c^(2/3) - Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(1/3)) - (-ArcTan[Sqrt[3]*(1 + (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]]) + (Sqrt[3]*Log[c^(2/3) + Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(1/3)))/b`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 216  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 217  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[(c_ \cdot)(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^2)^p, x\_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k(m+1)-1} \cdot (a + b \cdot x^{2k}/c^2)]^p, x], x, (c \cdot x)^{1/k}], x] /;$   $\text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 824  $\text{Int}[(x_ )^m / ((a_ + (b_ \cdot)(x_ )^n)), x\_Symbol] \rightarrow \text{Module}\{r = \text{Numerator}[\text{Rt}[a/b, n]], s = \text{Denominator}[\text{Rt}[a/b, n]], k, u\}, \text{Simp}[u = \text{Int}[(r \cdot \text{Cos}[(2k-1) \cdot m \cdot (\text{Pi}/n)] - s \cdot \text{Cos}[(2k-1) \cdot (m+1) \cdot (\text{Pi}/n)] \cdot x] / (r^2 - 2 \cdot r \cdot s \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x + s^2 \cdot x^2), x] + \text{Int}[(r \cdot \text{Cos}[(2k-1) \cdot m \cdot (\text{Pi}/n)] + s \cdot \text{Cos}[(2k-1) \cdot (m+1) \cdot (\text{Pi}/n)] \cdot x] / (r^2 + 2 \cdot r \cdot s \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x + s^2 \cdot x^2), x] ; 2 \cdot (-1)^{(m/2)} \cdot (r^{m+2} / (a \cdot n \cdot s^m)) \ \text{Int}[1/(r^2 + s^2 \cdot x^2), x] + 2 \cdot (r^{m+1} / (a \cdot n \cdot s^m)) \ \text{Sum}[u, \{k, 1, (n-2)/4\}], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{IGtQ}[(n-2)/4, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LtQ}[m, n-1] \ \&\& \ \text{PosQ}[a/b]$

rule 1082  $\text{Int}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot s \ \text{implify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$   $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$   $\text{FreeQ}\{a, b, c, x\}$

rule 1103  $\text{Int}[(d_ + (e_ \cdot)(x_ )) / ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2), x\_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142  $\text{Int}[(d_ + (e_ \cdot)(x_ )) / ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2), x\_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e) / (2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e / (2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x) / (a + b \cdot x + c \cdot x^2), x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\}$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.85

method	result
derivativedivides	$3c \left( \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln \left( (c \cot(bx+a))^{\frac{2}{3}} - \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}} \right)}{12c^2} + \frac{\arctan \left( \frac{2(c \cot(bx+a))^{\frac{1}{3}} - \sqrt{3}}{(c^2)^{\frac{1}{6}}} \right)}{6(c^2)^{\frac{1}{6}}} - \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln \left( (c \cot(bx+a))^{\frac{2}{3}} + \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}} \right)}{12c^2} + \frac{\arctan \left( \frac{2(c \cot(bx+a))^{\frac{1}{3}} + \sqrt{3}}{(c^2)^{\frac{1}{6}}} \right)}{6(c^2)^{\frac{1}{6}}} \right) \frac{1}{b}$
default	$3c \left( \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln \left( (c \cot(bx+a))^{\frac{2}{3}} - \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}} \right)}{12c^2} + \frac{\arctan \left( \frac{2(c \cot(bx+a))^{\frac{1}{3}} - \sqrt{3}}{(c^2)^{\frac{1}{6}}} \right)}{6(c^2)^{\frac{1}{6}}} - \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln \left( (c \cot(bx+a))^{\frac{2}{3}} + \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}} \right)}{12c^2} + \frac{\arctan \left( \frac{2(c \cot(bx+a))^{\frac{1}{3}} + \sqrt{3}}{(c^2)^{\frac{1}{6}}} \right)}{6(c^2)^{\frac{1}{6}}} \right) \frac{1}{b}$

input `int((c*cot(b*x+a))^(2/3),x,method=_RETURNVERBOSE)`

output `-3/b*c*(1/12/c^2*3^(1/2)*(c^2)^(5/6)*ln((c*cot(b*x+a))^(2/3)-3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/6/(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)-3^(1/2))-1/12/c^2*3^(1/2)*(c^2)^(5/6)*ln((c*cot(b*x+a))^(2/3)+3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/6/(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)+3^(1/2))+1/3/(c^2)^(1/6)*arctan((c*cot(b*x+a))^(1/3)/(c^2)^(1/6))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 416 vs.  $2(169) = 338$ .

Time = 0.08 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.85

$$\begin{aligned}
 & \int (c \cot(a \\
 & + bx))^{2/3} dx = \frac{1}{4} (\sqrt{-3} - 1) \left( -\frac{c^4}{b^6} \right)^{\frac{1}{6}} \log \left( c^3 \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right. \\
 & \left. + \frac{1}{2} (\sqrt{-3}b^5 + b^5) \left( -\frac{c^4}{b^6} \right)^{\frac{5}{6}} \right) \\
 & - \frac{1}{4} (\sqrt{-3} - 1) \left( -\frac{c^4}{b^6} \right)^{\frac{1}{6}} \log \left( c^3 \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right. \\
 & \left. - \frac{1}{2} (\sqrt{-3}b^5 + b^5) \left( -\frac{c^4}{b^6} \right)^{\frac{5}{6}} \right) \\
 & + \frac{1}{4} (\sqrt{-3} + 1) \left( -\frac{c^4}{b^6} \right)^{\frac{1}{6}} \log \left( c^3 \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right. \\
 & \left. + \frac{1}{2} (\sqrt{-3}b^5 - b^5) \left( -\frac{c^4}{b^6} \right)^{\frac{5}{6}} \right) \\
 & - \frac{1}{4} (\sqrt{-3} + 1) \left( -\frac{c^4}{b^6} \right)^{\frac{1}{6}} \log \left( c^3 \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right. \\
 & \left. - \frac{1}{2} (\sqrt{-3}b^5 - b^5) \left( -\frac{c^4}{b^6} \right)^{\frac{5}{6}} \right) \\
 & - \frac{1}{2} \left( -\frac{c^4}{b^6} \right)^{\frac{1}{6}} \log \left( b^5 \left( -\frac{c^4}{b^6} \right)^{\frac{5}{6}} + c^3 \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right) \\
 & + \frac{1}{2} \left( -\frac{c^4}{b^6} \right)^{\frac{1}{6}} \log \left( -b^5 \left( -\frac{c^4}{b^6} \right)^{\frac{5}{6}} + c^3 \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right)
 \end{aligned}$$

input `integrate((c*cot(b*x+a))^(2/3),x, algorithm="fricas")`



output

```
1/4*(sqrt(-3) - 1)*(-c^4/b^6)^(1/6)*log(c^3*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) + 1/2*(sqrt(-3)*b^5 + b^5)*(-c^4/b^6)^(5/6)) - 1/4*(sqrt(-3) - 1)*(-c^4/b^6)^(1/6)*log(c^3*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) - 1/2*(sqrt(-3)*b^5 + b^5)*(-c^4/b^6)^(5/6)) + 1/4*(sqrt(-3) + 1)*(-c^4/b^6)^(1/6)*log(c^3*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) + 1/2*(sqrt(-3)*b^5 - b^5)*(-c^4/b^6)^(5/6)) - 1/4*(sqrt(-3) + 1)*(-c^4/b^6)^(1/6)*log(c^3*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3) - 1/2*(sqrt(-3)*b^5 - b^5)*(-c^4/b^6)^(5/6)) - 1/2*(-c^4/b^6)^(1/6)*log(b^5*(-c^4/b^6)^(5/6) + c^3*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) + 1/2*(-c^4/b^6)^(1/6)*log(-b^5*(-c^4/b^6)^(5/6) + c^3*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3))
```

**Sympy [F]**

$$\int (c \cot(a + bx))^{2/3} dx = \int (c \cot(a + bx))^{2/3} dx$$

input

```
integrate((c*cot(b*x+a))**(2/3),x)
```

output

```
Integral((c*cot(a + b*x))**(2/3), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.81

$$\int (c \cot(a + bx))^{2/3} dx = \frac{\sqrt{3} \log\left(\sqrt{3}c^{1/3} \left(\frac{c}{\tan(bx+a)}\right)^{1/3} + c^{2/3} + \left(\frac{c}{\tan(bx+a)}\right)^{2/3}\right)}{c^{1/3}} - \frac{\sqrt{3} \log\left(-\sqrt{3}c^{1/3} \left(\frac{c}{\tan(bx+a)}\right)^{1/3} + c^{2/3} + \left(\frac{c}{\tan(bx+a)}\right)^{2/3}\right)}{c^{1/3}} - \frac{2 \arctan\left(\frac{c^{1/3} \left(\frac{c}{\tan(bx+a)}\right)^{1/3} + c^{2/3}}{\sqrt{3}c^{1/3} \left(\frac{c}{\tan(bx+a)}\right)^{1/3} + c^{2/3} + \left(\frac{c}{\tan(bx+a)}\right)^{2/3}}\right)}{4b}$$

input

```
integrate((c*cot(b*x+a))^(2/3),x, algorithm="maxima")
```

output

```
1/4*(sqrt(3)*log(sqrt(3)*c^(1/3)*(c/tan(b*x + a))^(1/3) + c^(2/3) + (c/tan
(b*x + a))^(2/3))/c^(1/3) - sqrt(3)*log(-sqrt(3)*c^(1/3)*(c/tan(b*x + a))^(
1/3) + c^(2/3) + (c/tan(b*x + a))^(2/3))/c^(1/3) - 2*arctan((sqrt(3)*c^(1
/3) + 2*(c/tan(b*x + a))^(1/3))/c^(1/3))/c^(1/3) - 2*arctan(-(sqrt(3)*c^(1
/3) - 2*(c/tan(b*x + a))^(1/3))/c^(1/3))/c^(1/3) - 4*arctan((c/tan(b*x + a
))^(1/3)/c^(1/3))/c^(1/3))*c/b
```

**Giac [F]**

$$\int (c \cot(a + bx))^{2/3} dx = \int (c \cot(bx + a))^{2/3} dx$$

input

```
integrate((c*cot(b*x+a))^(2/3),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(2/3), x)
```

**Mupad [B] (verification not implemented)**

Time = 10.53 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.16

$$\int (c \cot(a + bx))^{2/3} dx = -\frac{(-1)^{1/6} c^{2/3} \operatorname{atan}\left(\frac{(-1)^{2/3} (c \cot(a+bx))^{1/3}}{c^{1/3}}\right) \operatorname{li}}{b}$$

$$-\frac{(-1)^{1/6} c^{2/3} \ln\left(\frac{972 c^9}{b^3} - \frac{486 (-1)^{1/6} c^{26/3} (-1+\sqrt{3} \operatorname{li}) (c \cot(a+bx))^{1/3}}{b^3}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2b}$$

$$-\frac{(-1)^{1/6} c^{2/3} \ln\left(\frac{972 c^9}{b^3} - \frac{486 (-1)^{1/6} c^{26/3} (1+\sqrt{3} \operatorname{li}) (c \cot(a+bx))^{1/3}}{b^3}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2b}$$

$$+\frac{(-1)^{1/6} c^{2/3} \ln\left(\frac{972 c^9}{b^3} + \frac{486 (-1)^{1/6} c^{26/3} (-1+\sqrt{3} \operatorname{li}) (c \cot(a+bx))^{1/3}}{b^3}\right) \left(-\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b}$$

$$+\frac{(-1)^{1/6} c^{2/3} \ln\left(\frac{972 c^9}{b^3} + \frac{486 (-1)^{1/6} c^{26/3} (1+\sqrt{3} \operatorname{li}) (c \cot(a+bx))^{1/3}}{b^3}\right) \left(\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b}$$

input `int((c*cot(a + b*x))^(2/3),x)`

output 
$$\begin{aligned} &((-1)^{1/6}c^{2/3}\log((972c^9)/b^3 + (486(-1)^{1/6}c^{26/3})(3^{1/2})^* \\ &1i - 1)(c*\cot(a + b*x))^{1/3})/b^3*((3^{1/2})^*1i)/4 - 1/4)/b - ((-1)^{1/6} \\ &6)c^{2/3}\log((972c^9)/b^3 - (486(-1)^{1/6}c^{26/3})(3^{1/2})^*1i - 1)(c* \\ &c*\cot(a + b*x))^{1/3})/b^3*((3^{1/2})^*1i)/2 - 1/2)/(2*b) - ((-1)^{1/6}c^{2/3} \\ &(2/3)*\log((972c^9)/b^3 - (486(-1)^{1/6}c^{26/3})(3^{1/2})^*1i + 1)(c*\cot \\ &(a + b*x))^{1/3})/b^3*((3^{1/2})^*1i)/2 + 1/2)/(2*b) - ((-1)^{1/6}c^{2/3} \\ &*\operatorname{atan}(((1)^{2/3})(c*\cot(a + b*x))^{1/3})/c^{1/3})^*1i)/b + ((-1)^{1/6}c^{2/3} \\ &(2/3)*\log((972c^9)/b^3 + (486(-1)^{1/6}c^{26/3})(3^{1/2})^*1i + 1)(c*\cot \\ &a + b*x))^{1/3})/b^3*((3^{1/2})^*1i)/4 + 1/4)/b \end{aligned}$$

### Reduce [F]

$$\int (c \cot(a + bx))^{2/3} dx = c^{2/3} \left( \int \cot(bx + a)^{2/3} dx \right)$$

input `int((c*cot(b*x+a))^(2/3),x)`

output `c**(2/3)*int(cot(a + b*x)**(2/3),x)`

### 3.19 $\int \sqrt[3]{c \cot(a + bx)} dx$

Optimal result	195
Mathematica [A] (verified)	195
Rubi [A] (warning: unable to verify)	196
Maple [A] (verified)	199
Fricas [B] (verification not implemented)	200
Sympy [F]	201
Maxima [A] (verification not implemented)	201
Giac [A] (verification not implemented)	202
Mupad [B] (verification not implemented)	202
Reduce [F]	203

#### Optimal result

Integrand size = 12, antiderivative size = 131

$$\int \sqrt[3]{c \cot(a + bx)} dx = \frac{\sqrt{3} \sqrt[3]{c} \arctan\left(\frac{c^{2/3} - 2(c \cot(a + bx))^{2/3}}{\sqrt{3} c^{2/3}}\right)}{2b} + \frac{\sqrt[3]{c} \log(c^{2/3} + (c \cot(a + bx))^{2/3})}{2b} - \frac{\sqrt[3]{c} \log(c^{4/3} - c^{2/3}(c \cot(a + bx))^{2/3} + (c \cot(a + bx))^{4/3})}{4b}$$

output

```
1/2*3^(1/2)*c^(1/3)*arctan(1/3*(c^(2/3)-2*(c*cot(b*x+a))^(2/3))*3^(1/2)/c^(2/3))/b+1/2*c^(1/3)*ln(c^(2/3)+(c*cot(b*x+a))^(2/3))/b-1/4*c^(1/3)*ln(c^(4/3)-c^(2/3)*(c*cot(b*x+a))^(2/3)+(c*cot(b*x+a))^(4/3))/b
```

#### Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.81

$$\int \sqrt[3]{c \cot(a + bx)} dx = \frac{(c \cot(a + bx))^{4/3} \left( \log\left(1 + \sqrt[3]{\cot^2(a + bx)}\right) - \sqrt[3]{-1} \log\left(1 - \sqrt[3]{-1} \sqrt[3]{\cot^2(a + bx)}\right) + (-1)^{2/3} \log\left(1 - \sqrt[3]{-1} \sqrt[3]{\cot^2(a + bx)}\right) \right)}{2bc \cot^2(a + bx)^{2/3}}$$

input `Integrate[(c*Cot[a + b*x])^(1/3),x]`

output  $((c*\text{Cot}[a + b*x])^{4/3}*(\text{Log}[1 + (\text{Cot}[a + b*x]^2)^{1/3}] - (-1)^{1/3}*\text{Log}[1 - (-1)^{1/3}*(\text{Cot}[a + b*x]^2)^{1/3}] + (-1)^{2/3}*\text{Log}[1 + (-1)^{2/3}*(\text{Cot}[a + b*x]^2)^{1/3}]))/(2*b*c*(\text{Cot}[a + b*x]^2)^{2/3})$

### Rubi [A] (warning: unable to verify)

Time = 0.31 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.85, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3957, 266, 807, 821, 16, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[3]{c \cot(a + bx)} dx$$

$$\downarrow 3042$$

$$\int \sqrt[3]{-c \tan\left(a + bx + \frac{\pi}{2}\right)} dx$$

$$\downarrow 3957$$

$$c \int \frac{\sqrt[3]{c \cot(a + bx)}}{\cot^2(a + bx)c^2 + c^2} d(c \cot(a + bx))$$

$$\frac{c \int \sqrt[3]{c \cot(a + bx)}}{b}$$

$$\downarrow 266$$

$$3c \int \frac{c^3 \cot^3(a + bx)}{c^6 \cot^6(a + bx) + c^2} d \sqrt[3]{c \cot(a + bx)}$$

$$\frac{3c \int \frac{c^3 \cot^3(a + bx)}{c^6 \cot^6(a + bx) + c^2} d \sqrt[3]{c \cot(a + bx)}}{b}$$

$$\downarrow 807$$

$$3c \int \frac{c^2 \cot^2(a + bx)}{c^3 \cot^3(a + bx) + c^2} d(c^2 \cot^2(a + bx))$$

$$\frac{3c \int \frac{c^2 \cot^2(a + bx)}{c^3 \cot^3(a + bx) + c^2} d(c^2 \cot^2(a + bx))}{2b}$$

$$\downarrow 821$$

$$\begin{array}{c}
\frac{3c \left( \frac{\int \frac{c^2 \cot^2(a+bx)+c^{2/3}}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx)) - \frac{\int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d(c^2 \cot^2(a+bx))}{3c^{2/3}} \right)}{2b} \\
\downarrow 16 \\
\frac{3c \left( \frac{\int \frac{c^2 \cot^2(a+bx)+c^{2/3}}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx)) - \frac{\log(c^2 \cot^2(a+bx)+c^{2/3})}{3c^{2/3}} \right)}{2b} \\
\downarrow 1142 \\
\frac{3c \left( \frac{\frac{3}{2} c^{2/3} \int \frac{1}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx)) + \frac{1}{2} \int -\frac{c^{2/3}-2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{2/3}} - \frac{\log(c^2 \cot^2(a+bx)+c^{2/3})}{3c^{2/3}} \right)}{2b} \\
\downarrow 25 \\
\frac{3c \left( \frac{\frac{3}{2} c^{2/3} \int \frac{1}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx)) - \frac{1}{2} \int \frac{c^{2/3}-2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{2/3}} - \frac{\log(c^2 \cot^2(a+bx)+c^{2/3})}{3c^{2/3}} \right)}{2b} \\
\downarrow 1082 \\
\frac{3c \left( \frac{3 \int \frac{1}{2 \sqrt[3]{C \cot(a+bx)-4}} d(1-2 \sqrt[3]{C \cot(a+bx)}) - \frac{1}{2} \int \frac{c^{2/3}-2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{2/3}} - \frac{\log(c^2 \cot^2(a+bx)+c^{2/3})}{3c^{2/3}} \right)}{2b} \\
\downarrow 217 \\
\frac{3c \left( \frac{-\frac{1}{2} \int \frac{c^{2/3}-2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx)-c^{5/3} \cot(a+bx)+c^{4/3}} d(c^2 \cot^2(a+bx)) - \sqrt{3} \arctan\left(\frac{1-2 \sqrt[3]{C \cot(a+bx)}}{\sqrt{3}}\right)}{3c^{2/3}} - \frac{\log(c^2 \cot^2(a+bx)+c^{2/3})}{3c^{2/3}} \right)}{2b} \\
\downarrow 1103 \\
\frac{3c \left( \frac{\frac{1}{2} \log(-c^{5/3} \cot(a+bx)+c^2 \cot^2(a+bx)+c^{4/3}) - \sqrt{3} \arctan\left(\frac{1-2 \sqrt[3]{C \cot(a+bx)}}{\sqrt{3}}\right)}{3c^{2/3}} - \frac{\log(c^2 \cot^2(a+bx)+c^{2/3})}{3c^{2/3}} \right)}{2b}
\end{array}$$

input `Int[(c*Cot[a + b*x])^(1/3),x]`

output `(-3*c*(-1/3*Log[c^(2/3) + c^2*Cot[a + b*x]^2]/c^(2/3) + (-(Sqrt[3]*ArcTan[  
(1 - 2*c^(1/3)*Cot[a + b*x])/Sqrt[3]]) + Log[c^(4/3) - c^(5/3)*Cot[a + b*x]  
] + c^2*Cot[a + b*x]^2/2)/(3*c^(2/3)))/(2*b)`

### Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +  
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(  
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &  
& (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = De  
nominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))  
^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && I  
ntBinomialQ[a, b, c, 2, m, p, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m  
+ 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x,  
x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 821 `Int[(x_)/((a_) + (b_.)*(x_)^3), x_Symbol] := Simp[-(3*Rt[a, 3]*Rt[b, 3])^(  
-1) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]*Rt[b, 3])  
Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2  
*x^2), x], x] /; FreeQ[{a, b}, x]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

## Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.82



method	result
derivativeldivides	$3c \left[ \frac{\ln\left((c \cot(bx+a))^{\frac{2}{3}} + (c^2)^{\frac{1}{3}}\right)}{6(c^2)^{\frac{1}{3}}} + \frac{\ln\left((c \cot(bx+a))^{\frac{4}{3}} - (c \cot(bx+a))^{\frac{2}{3}} (c^2)^{\frac{1}{3}} + (c^2)^{\frac{2}{3}}\right)}{12(c^2)^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3} \left(\frac{2(c \cot(bx+a))}{(c^2)^{\frac{1}{3}}}\right)}{6(c^2)^{\frac{1}{3}}}\right)}{6(c^2)^{\frac{1}{3}}}$ <hr/> $b$
default	$3c \left[ \frac{\ln\left((c \cot(bx+a))^{\frac{2}{3}} + (c^2)^{\frac{1}{3}}\right)}{6(c^2)^{\frac{1}{3}}} + \frac{\ln\left((c \cot(bx+a))^{\frac{4}{3}} - (c \cot(bx+a))^{\frac{2}{3}} (c^2)^{\frac{1}{3}} + (c^2)^{\frac{2}{3}}\right)}{12(c^2)^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3} \left(\frac{2(c \cot(bx+a))}{(c^2)^{\frac{1}{3}}}\right)}{6(c^2)^{\frac{1}{3}}}\right)}{6(c^2)^{\frac{1}{3}}}$ <hr/> $b$

input

```
int((c*cot(b*x+a))^(1/3),x,method=_RETURNVERBOSE)
```

output

```
-3/b*c*(-1/6/(c^2)^(1/3)*ln((c*cot(b*x+a))^(2/3)+(c^2)^(1/3))+1/12/(c^2)^(1/3)*ln((c*cot(b*x+a))^(4/3)-(c*cot(b*x+a))^(2/3)*(c^2)^(1/3)+(c^2)^(2/3))+1/6*3^(1/2)/(c^2)^(1/3)*arctan(1/3*3^(1/2)*(2*(c*cot(b*x+a))^(2/3)/(c^2)^(1/3)-1)))
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. 2(100) = 200.

Time = 0.08 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.61

$$\int \sqrt[3]{c \cot(a + bx)} dx = \frac{2\sqrt{3}c^{\frac{1}{3}} \arctan\left(-\frac{\sqrt{3}c - 2\sqrt{3}c^{\frac{1}{3}}\left(\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}\right)^{\frac{2}{3}}}{3c}\right) - 2c^{\frac{1}{3}} \log\left(c^{\frac{2}{3}} + \left(\frac{c \cos(2bx+2a)+c}{\sin(2bx+2a)}\right)^{\frac{2}{3}}\right) + c^{\frac{1}{3}} \log\left(\frac{c^{\frac{4}{3}} \sin(2bx+2a)}{c^{\frac{4}{3}} \sin(2bx+2a)}\right)}{4b}$$

input `integrate((c*cot(b*x+a))^(1/3),x, algorithm="fricas")`

output `-1/4*(2*sqrt(3)*c^(1/3)*arctan(-1/3*(sqrt(3)*c - 2*sqrt(3)*c^(1/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3))/c) - 2*c^(1/3)*log(c^(2/3) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)) + c^(1/3)*log((c^(4/3)*sin(2*b*x + 2*a) - c^(2/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3))*sin(2*b*x + 2*a) + (c*cos(2*b*x + 2*a) + c)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3))/sin(2*b*x + 2*a))/b`

## Sympy [F]

$$\int \sqrt[3]{c \cot(a + bx)} dx = \int \sqrt[3]{c \cot(a + bx)} dx$$

input `integrate((c*cot(b*x+a))**(1/3),x)`

output `Integral((c*cot(a + b*x))**(1/3), x)`

## Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.78

$$\int \sqrt[3]{c \cot(a + bx)} dx = \frac{c \left( \frac{2\sqrt{3} \arctan \left( -\frac{\sqrt{3} \left( c^{\frac{2}{3}} - 2 \left( \frac{c}{\tan(bx+a)} \right)^{\frac{2}{3}} \right)}{3c^{\frac{2}{3}}} \right)}{c^{\frac{2}{3}}} + \frac{\log \left( c^{\frac{4}{3}} - c^{\frac{2}{3}} \left( \frac{c}{\tan(bx+a)} \right)^{\frac{2}{3}} + \left( \frac{c}{\tan(bx+a)} \right)^{\frac{4}{3}} \right)}{c^{\frac{2}{3}}} - \frac{2 \log \left( c^{\frac{2}{3}} + \left( \frac{c}{\tan(bx+a)} \right)^{\frac{2}{3}} \right)}{c^{\frac{2}{3}}} \right)}{4b}$$

input `integrate((c*cot(b*x+a))^(1/3),x, algorithm="maxima")`

output

```
-1/4*c*(2*sqrt(3)*arctan(-1/3*sqrt(3)*(c^(2/3) - 2*(c/tan(b*x + a))^(2/3))
/c^(2/3))/c^(2/3) + log(c^(4/3) - c^(2/3)*(c/tan(b*x + a))^(2/3) + (c/tan(
b*x + a))^(4/3))/c^(2/3) - 2*log(c^(2/3) + (c/tan(b*x + a))^(2/3))/c^(2/3)
)/b
```

**Giac [A] (verification not implemented)**

Time = 0.86 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.11

$$\int \sqrt[3]{c \cot(a + bx)} dx$$

$$c \left( \frac{2\sqrt{3}(-c)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left((-c)^{\frac{1}{3}} + 2\left(-c + \frac{c}{\cos(bx+a)^2}\right)^{\frac{1}{3}}\right)}{3(-c)^{\frac{1}{3}}}\right)}{c} + \frac{(-c)^{\frac{1}{3}} \log\left(\left(-c\right)^{\frac{2}{3}} + (-c)^{\frac{1}{3}}\left(-c + \frac{c}{\cos(bx+a)^2}\right)^{\frac{1}{3}} + \left(-c + \frac{c}{\cos(bx+a)^2}\right)^{\frac{2}{3}}\right)}{c} \right) - \dots$$


---


$$= \frac{\dots}{4b}$$

input

```
integrate((c*cot(b*x+a))^(1/3),x, algorithm="giac")
```

output

```
1/4*c*(2*sqrt(3)*(-c)^(1/3)*arctan(1/3*sqrt(3)*((-c)^(1/3) + 2*(-c + c/cos
(b*x + a)^2)^(1/3))/(-c)^(1/3))/c + (-c)^(1/3)*log((-c)^(2/3) + (-c)^(1/3)
*(-c + c/cos(b*x + a)^2)^(1/3) + (-c + c/cos(b*x + a)^2)^(2/3))/c - 2*(-c)
^(1/3)*log(abs(-(-c)^(1/3) + (-c + c/cos(b*x + a)^2)^(1/3)))/c)/b
```

**Mupad [B] (verification not implemented)**

Time = 9.24 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.02

$$\int \sqrt[3]{c \cot(a + bx)} dx = \frac{c^{1/3} \ln\left(81 c^{16/3} (c \cot(a + bx))^{2/3} + 81 c^6\right)}{2b}$$

$$- \frac{c^{1/3} \ln\left(\frac{81 c^6}{b^4} - \frac{81 c^{16/3} \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2}\right) (c \cot(a + bx))^{2/3}}{b^4}\right) \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2}\right)}{2b}$$

$$+ \frac{c^{1/3} \ln\left(\frac{81 c^6}{b^4} + \frac{162 c^{16/3} \left(-\frac{1}{4} + \frac{\sqrt{3} i i}{4}\right) (c \cot(a + bx))^{2/3}}{b^4}\right) \left(-\frac{1}{4} + \frac{\sqrt{3} i i}{4}\right)}{b}$$

input `int((c*cot(a + b*x))^(1/3),x)`

output `(c^(1/3)*log(81*c^(16/3)*(c*cot(a + b*x))^(2/3) + 81*c^6))/(2*b) - (c^(1/3)*log((81*c^6)/b^4 - (81*c^(16/3)*((3^(1/2)*1i)/2 + 1/2)*(c*cot(a + b*x))^(2/3))/b^4)*((3^(1/2)*1i)/2 + 1/2))/(2*b) + (c^(1/3)*log((81*c^6)/b^4 + (162*c^(16/3)*((3^(1/2)*1i)/4 - 1/4)*(c*cot(a + b*x))^(2/3))/b^4)*((3^(1/2)*1i)/4 - 1/4))/b`

### Reduce [F]

$$\int \sqrt[3]{c \cot(a + bx)} dx = c^{\frac{1}{3}} \left( \int \cot(bx + a)^{\frac{1}{3}} dx \right)$$

input `int((c*cot(b*x+a))^(1/3),x)`

output `c**(1/3)*int(cot(a + b*x)**(1/3),x)`

**3.20**  $\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx$

Optimal result . . . . .	204
Mathematica [A] (verified) . . . . .	204
Rubi [A] (warning: unable to verify) . . . . .	205
Maple [A] (verified) . . . . .	209
Fricas [B] (verification not implemented) . . . . .	209
Sympy [F] . . . . .	210
Maxima [A] (verification not implemented) . . . . .	211
Giac [A] (verification not implemented) . . . . .	211
Mupad [B] (verification not implemented) . . . . .	212
Reduce [F] . . . . .	212

**Optimal result**

Integrand size = 12, antiderivative size = 131

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx = \frac{\sqrt{3} \arctan\left(\frac{c^{2/3} - 2(c \cot(a + bx))^{2/3}}{\sqrt{3}c^{2/3}}\right)}{2b\sqrt[3]{c}} - \frac{\log(c^{2/3} + (c \cot(a + bx))^{2/3})}{2b\sqrt[3]{c}} + \frac{\log(c^{4/3} - c^{2/3}(c \cot(a + bx))^{2/3} + (c \cot(a + bx))^{4/3})}{4b\sqrt[3]{c}}$$

output `1/2*3^(1/2)*arctan(1/3*(c^(2/3)-2*(c*cot(b*x+a))^(2/3))*3^(1/2)/c^(2/3))/b  
/c^(1/3)-1/2*ln(c^(2/3)+(c*cot(b*x+a))^(2/3))/b/c^(1/3)+1/4*ln(c^(4/3)-c^(  
2/3)*(c*cot(b*x+a))^(2/3)+(c*cot(b*x+a))^(4/3))/b/c^(1/3)`

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx = \frac{\sqrt[3]{\cot(a + bx)} \left( -2\sqrt{3} \arctan\left(\frac{-1+2\cot^{\frac{2}{3}}(a+bx)}{\sqrt{3}}\right) - 2 \log\left(1 + \cot^{\frac{2}{3}}(a + bx)\right) + \log\left(1 - \cot^{\frac{2}{3}}(a + bx) + c\right) \right)}{4b\sqrt[3]{c \cot(a + bx)}}$$

input `Integrate[(c*Cot[a + b*x])^(-1/3),x]`

output `(Cot[a + b*x]^(1/3)*(-2*Sqrt[3]*ArcTan[(-1 + 2*Cot[a + b*x]^(2/3))/Sqrt[3]] - 2*Log[1 + Cot[a + b*x]^(2/3)] + Log[1 - Cot[a + b*x]^(2/3) + Cot[a + b*x]^(4/3)]))/(4*b*(c*Cot[a + b*x])^(1/3))`

### Rubi [A] (warning: unable to verify)

Time = 0.32 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.85, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3957, 266, 807, 750, 16, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt[3]{-c \tan\left(a + bx + \frac{\pi}{2}\right)}} dx \\
 & \quad \downarrow \text{3957} \\
 & - \frac{c \int \frac{1}{\sqrt[3]{c \cot(a + bx)} (\cot^2(a + bx) c^2 + c^2)} d(c \cot(a + bx))}{b} \\
 & \quad \downarrow \text{266} \\
 & - \frac{3c \int \frac{\sqrt[3]{c \cot(a + bx)}}{c^6 \cot^6(a + bx) + c^2} d \sqrt[3]{c \cot(a + bx)}}{b} \\
 & \quad \downarrow \text{807} \\
 & - \frac{3c \int \frac{1}{c^3 \cot^3(a + bx) + c^2} d(c^2 \cot^2(a + bx))}{2b} \\
 & \quad \downarrow \text{750}
 \end{aligned}$$

$$\begin{array}{c}
\frac{3c \left( \frac{\int \frac{1}{c^2 \cot^2(a+bx) + c^{2/3}} d(c^2 \cot^2(a+bx))}{3c^{4/3}} + \frac{\int \frac{2c^{2/3} - c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{4/3}} \right)}{2b} \\
\downarrow 16 \\
\frac{3c \left( \frac{\int \frac{2c^{2/3} - c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{4/3}} + \frac{\log(c^2 \cot^2(a+bx) + c^{2/3})}{3c^{4/3}} \right)}{2b} \\
\downarrow 1142 \\
\frac{3c \left( \frac{\frac{3}{2} c^{2/3} \int \frac{1}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx)) - \frac{1}{2} \int \frac{c^{2/3} - 2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{4/3}} + \frac{\log(c^2 \cot^2(a+bx) + c^{2/3})}{3c^{4/3}} \right)}{2b} \\
\downarrow 25 \\
\frac{3c \left( \frac{\frac{3}{2} c^{2/3} \int \frac{1}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx)) + \frac{1}{2} \int \frac{c^{2/3} - 2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx))}{3c^{4/3}} + \frac{\log(c^2 \cot^2(a+bx) + c^{2/3})}{3c^{4/3}} \right)}{2b} \\
\downarrow 1082 \\
\frac{3c \left( \frac{\frac{1}{2} \int \frac{c^{2/3} - 2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx)) + 3 \int \frac{1}{2 \sqrt[3]{C \cot(a+bx) - 4}} d(1 - 2 \sqrt[3]{C \cot(a+bx)})}{3c^{4/3}} + \frac{\log(c^2 \cot^2(a+bx) + c^{2/3})}{3c^{4/3}} \right)}{2b} \\
\downarrow 217 \\
\frac{3c \left( \frac{\frac{1}{2} \int \frac{c^{2/3} - 2c^2 \cot^2(a+bx)}{c^2 \cot^2(a+bx) - c^{5/3} \cot(a+bx) + c^{4/3}} d(c^2 \cot^2(a+bx)) - \sqrt{3} \arctan \left( \frac{1 - 2 \sqrt[3]{C \cot(a+bx)}}{\sqrt{3}} \right)}{3c^{4/3}} + \frac{\log(c^2 \cot^2(a+bx) + c^{2/3})}{3c^{4/3}} \right)}{2b} \\
\downarrow 1103
\end{array}$$

$$\frac{3c \left( \frac{-\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{c} \cot(a+bx)}{\sqrt{3}}\right) - \frac{1}{2} \log(-c^{5/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{4/3})}{3c^{4/3}} + \frac{\log(c^2 \cot^2(a+bx) + c^{2/3})}{3c^{4/3}} \right)}{2b}$$

input `Int[(c*Cot[a + b*x])^(-1/3),x]`

output `(-3*c*(Log[c^(2/3) + c^2*Cot[a + b*x]^2]/(3*c^(4/3)) + (-Sqrt[3]*ArcTan[(1 - 2*c^(1/3)*Cot[a + b*x])/Sqrt[3]]) - Log[c^(4/3) - c^(5/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2]/2)/(3*c^(4/3)))/(2*b)`

### Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 750 `Int[((a_) + (b_.)*(x_)^3)^(-1), x_Symbol] := Simp[1/(3*Rt[a, 3]^2) Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Simp[1/(3*Rt[a, 3]^2) Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]`



rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.82

method	result
derivativedivides	$3c \left( \frac{\ln\left(\frac{(c \cot(bx+a))^{\frac{2}{3}} + (c^2)^{\frac{1}{3}}}{6(c^2)^{\frac{2}{3}}}\right) - \ln\left(\frac{(c \cot(bx+a))^{\frac{4}{3}} - (c \cot(bx+a))^{\frac{2}{3}}(c^2)^{\frac{1}{3}} + (c^2)^{\frac{2}{3}}}{12(c^2)^{\frac{2}{3}}}\right)}{b} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3} \left(\frac{2(c \cot(bx+a))^{\frac{2}{3}}}{(c^2)^{\frac{1}{3}}}\right)}{6(c^2)^{\frac{2}{3}}}\right)}{6(c^2)^{\frac{2}{3}}}$
default	$3c \left( \frac{\ln\left(\frac{(c \cot(bx+a))^{\frac{2}{3}} + (c^2)^{\frac{1}{3}}}{6(c^2)^{\frac{2}{3}}}\right) - \ln\left(\frac{(c \cot(bx+a))^{\frac{4}{3}} - (c \cot(bx+a))^{\frac{2}{3}}(c^2)^{\frac{1}{3}} + (c^2)^{\frac{2}{3}}}{12(c^2)^{\frac{2}{3}}}\right)}{b} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3} \left(\frac{2(c \cot(bx+a))^{\frac{2}{3}}}{(c^2)^{\frac{1}{3}}}\right)}{6(c^2)^{\frac{2}{3}}}\right)}{6(c^2)^{\frac{2}{3}}}$

input `int(1/(c*cot(b*x+a))^(1/3),x,method=_RETURNVERBOSE)`

output `-3/b*c*(1/6/(c^2)^(2/3)*ln((c*cot(b*x+a))^(2/3)+(c^2)^(1/3))-1/12/(c^2)^(2/3)*ln((c*cot(b*x+a))^(4/3)-(c*cot(b*x+a))^(2/3)*(c^2)^(1/3)+(c^2)^(2/3))+1/6/(c^2)^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2*(c*cot(b*x+a))^(2/3)/(c^2)^(1/3)-1)))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 268 vs. 2(100) = 200.

Time = 0.09 (sec) , antiderivative size = 639, normalized size of antiderivative = 4.88

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx = \text{Too large to display}$$

input `integrate(1/(c*cot(b*x+a))^(1/3),x, algorithm="fricas")`

output `[1/4*(sqrt(3)*c*sqrt((-c)^(1/3)/c)*log(1/2*sqrt(3)*((-c)^(2/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)*(cos(2*b*x + 2*a) - 1) - 2*c*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)*sin(2*b*x + 2*a) + (c*cos(2*b*x + 2*a) - c)*(-c)^(1/3))*sqrt((-c)^(1/3)/c) - 3/2*(-c)^(1/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)*(cos(2*b*x + 2*a) - 1) + 3/2*c*cos(2*b*x + 2*a) + 1/2*c) - 2*(-c)^(2/3)*log((-c)^(2/3) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)) + (-c)^(2/3)*log(-((-c)^(1/3)*c*sin(2*b*x + 2*a) + (-c)^(2/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)*sin(2*b*x + 2*a) - (c*cos(2*b*x + 2*a) + c)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3))/sin(2*b*x + 2*a)))/(b*c), -1/4*(2*sqrt(3)*c*sqrt(-(-c)^(1/3)/c)*arctan(1/3*(sqrt(3)*(-c)^(1/3)*c*sqrt(-(-c)^(1/3)/c) + 2*sqrt(3)*(-c)^(2/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)*sqrt(-(-c)^(1/3)/c))/c) + 2*(-c)^(2/3)*log((-c)^(2/3) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)) - (-c)^(2/3)*log(-((-c)^(1/3)*c*sin(2*b*x + 2*a) + (-c)^(2/3)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)*sin(2*b*x + 2*a) - (c*cos(2*b*x + 2*a) + c)*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3))/sin(2*b*x + 2*a)))/(b*c)]`

## Sympy [F]

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx = \int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx$$

input `integrate(1/(c*cot(b*x+a))**(1/3),x)`

output `Integral((c*cot(a + b*x))**(-1/3), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx =$$

$$\frac{c \left( \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(c^{\frac{2}{3}} - 2\left(\frac{c}{\tan(bx+a)}\right)^{\frac{2}{3}}\right)}{3c^{\frac{2}{3}}}\right)}{c^{\frac{4}{3}}} - \frac{\log\left(c^{\frac{4}{3}} - c^{\frac{2}{3}}\left(\frac{c}{\tan(bx+a)}\right)^{\frac{2}{3}} + \left(\frac{c}{\tan(bx+a)}\right)^{\frac{4}{3}}\right)}{c^{\frac{4}{3}}} + \frac{2 \log\left(c^{\frac{2}{3}} + \left(\frac{c}{\tan(bx+a)}\right)^{\frac{2}{3}}\right)}{c^{\frac{4}{3}}}\right)}{4b}$$

input `integrate(1/(c*cot(b*x+a))^(1/3),x, algorithm="maxima")`output `-1/4*c*(2*sqrt(3)*arctan(-1/3*sqrt(3)*(c^(2/3) - 2*(c/tan(b*x + a))^(2/3))/c^(2/3))/c^(4/3) - log(c^(4/3) - c^(2/3)*(c/tan(b*x + a))^(2/3) + (c/tan(b*x + a))^(4/3))/c^(4/3) + 2*log(c^(2/3) + (c/tan(b*x + a))^(2/3))/c^(4/3))/b`**Giac [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.11

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx =$$

$$\frac{2\sqrt{3}(-c)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left((-c)^{\frac{1}{3}} + 2\left(-c + \frac{c}{\cos(bx+a)^2}\right)^{\frac{1}{3}}\right)}{3(-c)^{\frac{1}{3}}}\right)}{c} - \frac{(-c)^{\frac{2}{3}} \log\left((-c)^{\frac{2}{3}} + (-c)^{\frac{1}{3}}\left(-c + \frac{c}{\cos(bx+a)^2}\right)^{\frac{1}{3}} + \left(-c + \frac{c}{\cos(bx+a)^2}\right)^{\frac{2}{3}}\right)}{c} + \dots$$

$4b$

input `integrate(1/(c*cot(b*x+a))^(1/3),x, algorithm="giac")`

output

```
-1/4*(2*sqrt(3)*(-c)^(2/3)*arctan(1/3*sqrt(3)*((-c)^(1/3) + 2*(-c + c/cos(
b*x + a)^2)^(1/3))/(-c)^(1/3))/c - (-c)^(2/3)*log((-c)^(2/3) + (-c)^(1/3)*
(-c + c/cos(b*x + a)^2)^(1/3) + (-c + c/cos(b*x + a)^2)^(2/3))/c + 2*(-c)^(
2/3)*log(abs(-(-c)^(1/3) + (-c + c/cos(b*x + a)^2)^(1/3)))/c)/b
```

**Mupad [B] (verification not implemented)**

Time = 9.98 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.98

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx = -\frac{\ln\left((c \cot(a + bx))^{2/3} + c^{2/3}\right)}{2bc^{1/3}} - \frac{\ln\left(\frac{81c^{11/3}(-1+\sqrt{3}i)}{b^3} + \frac{162c^3(c \cot(a+bx))^{2/3}}{b^3}\right)(-1+\sqrt{3}i)}{4bc^{1/3}} + \frac{\ln\left(\frac{81c^{11/3}(1+\sqrt{3}i)}{b^3} - \frac{162c^3(c \cot(a+bx))^{2/3}}{b^3}\right)(1+\sqrt{3}i)}{4bc^{1/3}}$$

input

```
int(1/(c*cot(a + b*x))^(1/3),x)
```

output

```
(log((81*c^(11/3)*(3^(1/2)*1i + 1))/b^3 - (162*c^3*(c*cot(a + b*x))^(2/3))
/b^3)*(3^(1/2)*1i + 1))/(4*b*c^(1/3)) - (log((81*c^(11/3)*(3^(1/2)*1i - 1)
)/b^3 + (162*c^3*(c*cot(a + b*x))^(2/3))/b^3*(3^(1/2)*1i - 1))/(4*b*c^(1/
3)) - log((c*cot(a + b*x))^(2/3) + c^(2/3))/(2*b*c^(1/3))
```

**Reduce [F]**

$$\int \frac{1}{\sqrt[3]{c \cot(a + bx)}} dx = \frac{\int \frac{1}{\cot(bx+a)^{\frac{1}{3}}} dx}{c^{\frac{1}{3}}}$$

input

```
int(1/(c*cot(b*x+a))^(1/3),x)
```

output

```
int(1/cot(a + b*x)**(1/3),x)/c**(1/3)
```

### 3.21 $\int \frac{1}{(c \cot(a+bx))^{2/3}} dx$

Optimal result	213
Mathematica [C] (verified)	214
Rubi [A] (warning: unable to verify)	214
Maple [A] (verified)	218
Fricas [B] (verification not implemented)	219
Sympy [F]	220
Maxima [A] (verification not implemented)	220
Giac [F]	221
Mupad [B] (verification not implemented)	221
Reduce [F]	222

#### Optimal result

Integrand size = 12, antiderivative size = 225

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx = -\frac{\arctan\left(\frac{\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{bc^{2/3}} + \frac{\arctan\left(\sqrt{3} - \frac{2\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{2bc^{2/3}} - \frac{\arctan\left(\sqrt{3} + \frac{2\sqrt[3]{c \cot(a + bx)}}{\sqrt[3]{c}}\right)}{2bc^{2/3}} + \frac{\sqrt{3} \log\left(c^{2/3} - \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a + bx)} + (c \cot(a + bx))^{2/3}\right)}{4bc^{2/3}} - \frac{\sqrt{3} \log\left(c^{2/3} + \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a + bx)} + (c \cot(a + bx))^{2/3}\right)}{4bc^{2/3}}$$

output

```
-arctan((c*cot(b*x+a))^(1/3)/c^(1/3))/b/c^(2/3)-1/2*arctan(-3^(1/2)+2*(c*cot(b*x+a))^(1/3)/c^(1/3))/b/c^(2/3)-1/2*arctan(3^(1/2)+2*(c*cot(b*x+a))^(1/3)/c^(1/3))/b/c^(2/3)+1/4*3^(1/2)*ln(c^(2/3)-3^(1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b/c^(2/3)-1/4*3^(1/2)*ln(c^(2/3)+3^(1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b/c^(2/3)
```

**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.84

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx = \frac{\sqrt[3]{c \cot(a + bx)} \left( -i \log \left( 1 - i \sqrt[6]{\cot^2(a + bx)} \right) + i \log \left( 1 + i \sqrt[6]{\cot^2(a + bx)} \right) \right)}{b}$$

input

```
Integrate[(c*Cot[a + b*x])^(-2/3),x]
```

output

```
((c*Cot[a + b*x])^(1/3)*((-I)*Log[1 - I*(Cot[a + b*x]^2)^(1/6)] + I*Log[1 + I*(Cot[a + b*x]^2)^(1/6)]) + (-1)^(1/6)*(-((-1)^(2/3)*Log[1 - (-1)^(1/6)*(Cot[a + b*x]^2)^(1/6)]) + (-1)^(2/3)*Log[1 + (-1)^(1/6)*(Cot[a + b*x]^2)^(1/6)]) - Log[1 - (-1)^(5/6)*(Cot[a + b*x]^2)^(1/6)] + Log[1 + (-1)^(5/6)*(Cot[a + b*x]^2)^(1/6)]))/(2*b*c*(Cot[a + b*x]^2)^(1/6))
```

**Rubi [A] (warning: unable to verify)**

Time = 0.45 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.87, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$ , Rules used = {3042, 3957, 266, 753, 27, 216, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(c \cot(a + bx))^{2/3}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{2/3}} dx \\ & \quad \downarrow \text{3957} \\ & - \frac{c \int \frac{1}{(c \cot(a + bx))^{2/3} (\cot^2(a + bx) c^2 + c^2)} d(c \cot(a + bx))}{b} \end{aligned}$$

$$\begin{aligned} & \downarrow 266 \\ & \frac{3c \int \frac{1}{c^6 \cot^6(a+bx)+c^2} d^3 \sqrt{c \cot(a+bx)}}{b} \\ & \downarrow 753 \\ & \frac{3c \left( \int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} \right)}{3c^{4/3}} + \frac{\int \frac{2 \sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{2(c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3})} d^3 \sqrt{c \cot(a+bx)}}{3c^{5/3}} + \frac{\int \frac{2 \sqrt[3]{c+\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{2(c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3})} d^3 \sqrt{c \cot(a+bx)}}{3c^{5/3}}}{b} \\ & \downarrow 27 \\ & \frac{3c \left( \int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} \right)}{3c^{4/3}} + \frac{\int \frac{2 \sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} + \frac{\int \frac{2 \sqrt[3]{c+\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}}}{b} \\ & \downarrow 216 \\ & \frac{3c \left( \int \frac{2 \sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)} \right)}{6c^{5/3}} + \frac{\int \frac{2 \sqrt[3]{c+\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} + \frac{\arctan \left( \frac{\sqrt{3} \sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} \right)}{b} \\ & \downarrow 1142 \\ & \frac{3c \left( \frac{\frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} - \frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} \right)}{6c^{5/3}} \\ & \downarrow 25 \\ & \frac{3c \left( \frac{\frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} + \frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6c^{5/3}} \right)}{6c^{5/3}} \\ & \downarrow 1082 \end{aligned}$$



$$\begin{aligned}
 & \frac{3c \left( \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - \frac{1}{3}} d \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3} c^{4/3} \cot(a+bx) + c^{2/3}} d \sqrt[3]{c \cot(a+bx)} + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c+2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3} c^{4/3} \cot(a+bx) + c^{2/3}} d \sqrt[3]{c \cot(a+bx)} \right)}{b} \\
 & \quad \downarrow 217 \\
 & \frac{3c \left( \frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3} c^{4/3} \cot(a+bx) + c^{2/3}} d \sqrt[3]{c \cot(a+bx)} - \arctan \left( \sqrt{3} \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right) \right)}{6c^{5/3}} + \frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c+2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3} c^{4/3} \cot(a+bx) + c^{2/3}} d \sqrt[3]{c \cot(a+bx)} \right)}{b} \\
 & \quad \downarrow 1103 \\
 & \frac{3c \left( \frac{\arctan(c^{2/3} \cot(a+bx))}{3c^{5/3}} + \frac{-\arctan \left( \sqrt{3} \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right) \right) - \frac{1}{2} \sqrt{3} \log \left( -\sqrt{3} c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3} \right)}{6c^{5/3}} + \frac{\arctan \left( \sqrt{3} \left( 1 + \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right) \right)}{3c^{5/3}} \right)}{b}
 \end{aligned}$$

```
input Int[(c*Cot[a + b*x])^(-2/3), x]
```

```
output (-3*c*(ArcTan[c^(2/3)*Cot[a + b*x]]/(3*c^(5/3)) + (-ArcTan[Sqrt[3]*(1 - (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]]) - (Sqrt[3]*Log[c^(2/3) - Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(5/3)) + (ArcTan[Sqrt[3]*(1 + (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]]) + (Sqrt[3]*Log[c^(2/3) + Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(5/3)))/b
```

Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 216  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 217  $\text{Int}[(a_ + (b_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[(c_ \cdot)(x_ )^m \cdot (a_ + (b_ \cdot)(x_ )^2)^p, x\_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k(m+1)-1} \cdot (a + b \cdot x^{2k}/c^2)]^p, x], x, (c \cdot x)^{1/k}], x] /;$   $\text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 753  $\text{Int}[(a_ + (b_ \cdot)(x_ )^n)^{-1}, x\_Symbol] \rightarrow \text{Module}\{r = \text{Numerator}[\text{Rt}[a/b, n]], s = \text{Denominator}[\text{Rt}[a/b, n]], k, u, v\}, \text{Simp}[u = \text{Int}[(r - s \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x)/(r^2 - 2 \cdot r \cdot s \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x + s^2 \cdot x^2), x] + \text{Int}[(r + s \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x)/(r^2 + 2 \cdot r \cdot s \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x + s^2 \cdot x^2), x]; 2 \cdot (r^2/(a \cdot n)) \ \text{Int}[1/(r^2 + s^2 \cdot x^2), x] + 2 \cdot (r/(a \cdot n)) \ \text{Sum}[u, \{k, 1, (n-2)/4\}], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{IGtQ}[(n-2)/4, 0] \ \&\& \ \text{PosQ}[a/b]$

rule 1082  $\text{Int}[(a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$   $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$   $\text{FreeQ}\{a, b, c, x\}$

rule 1103  $\text{Int}[(d_ + (e_ \cdot)(x_ )) / ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2), x\_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142  $\text{Int}[(d_ + (e_ \cdot)(x_ )) / ((a_ + (b_ \cdot)(x_ ) + (c_ \cdot)(x_ )^2), x\_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\}$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.89

method	result
derivativedivides	$3c \frac{\left( \frac{\sqrt{3} (c^2)^{\frac{1}{6}} \ln \left( (c \cot(bx+a))^{\frac{2}{3}} + \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}} \right)}{12c^2} \right) + \frac{(c^2)^{\frac{1}{6}} \arctan \left( \frac{2(c \cot(bx+a))^{\frac{1}{3}} + \sqrt{3}}{(c^2)^{\frac{1}{6}}} \right)}{6c^2}}{\sqrt{3} (c^2)^{\frac{1}{6}}}$
default	$3c \frac{\left( \frac{\sqrt{3} (c^2)^{\frac{1}{6}} \ln \left( (c \cot(bx+a))^{\frac{2}{3}} + \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}} \right)}{12c^2} \right) + \frac{(c^2)^{\frac{1}{6}} \arctan \left( \frac{2(c \cot(bx+a))^{\frac{1}{3}} + \sqrt{3}}{(c^2)^{\frac{1}{6}}} \right)}{6c^2}}{\sqrt{3} (c^2)^{\frac{1}{6}}}$

input `int(1/(c*cot(b*x+a))^(2/3),x,method=_RETURNVERBOSE)`

output `-3/b*c*(1/12/c^2*3^(1/2)*(c^2)^(1/6)*ln((c*cot(b*x+a))^(2/3)+3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/6/c^2*(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)+3^(1/2))-1/12/c^2*3^(1/2)*(c^2)^(1/6)*ln((c*cot(b*x+a))^(2/3)-3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/6/c^2*(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)-3^(1/2))+1/3/c^2*(c^2)^(1/6)*arctan((c*cot(b*x+a))^(1/3)/(c^2)^(1/6))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 384 vs.  $2(169) = 338$ .

Time = 0.08 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.71

$$\begin{aligned}
& \int \frac{1}{(c \cot(a + bx))^{2/3}} dx = \\
& -\frac{1}{4} (\sqrt{-3} + 1) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \log \left( \frac{1}{2} (\sqrt{-3}bc + bc) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \right. \\
& \left. + \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right) \\
& + \frac{1}{4} (\sqrt{-3} + 1) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \log \left( -\frac{1}{2} (\sqrt{-3}bc + bc) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \right. \\
& \left. + \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right) \\
& - \frac{1}{4} (\sqrt{-3} - 1) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \log \left( \frac{1}{2} (\sqrt{-3}bc - bc) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \right. \\
& \left. + \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right) \\
& + \frac{1}{4} (\sqrt{-3} - 1) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \log \left( -\frac{1}{2} (\sqrt{-3}bc - bc) \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \right. \\
& \left. + \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right) \\
& - \frac{1}{2} \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \log \left( bc \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} + \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right) \\
& + \frac{1}{2} \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} \log \left( -bc \left(-\frac{1}{b^6 c^4}\right)^{\frac{1}{6}} + \left( \frac{c \cos(2bx + 2a) + c}{\sin(2bx + 2a)} \right)^{\frac{1}{3}} \right)
\end{aligned}$$

input `integrate(1/(c*cot(b*x+a))^(2/3),x, algorithm="fricas")`

output

```
-1/4*(sqrt(-3) + 1)*(-1/(b^6*c^4))^(1/6)*log(1/2*(sqrt(-3)*b*c + b*c)*(-1/
(b^6*c^4))^(1/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) + 1/
4*(sqrt(-3) + 1)*(-1/(b^6*c^4))^(1/6)*log(-1/2*(sqrt(-3)*b*c + b*c)*(-1/(b
^6*c^4))^(1/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) - 1/4*
(sqrt(-3) - 1)*(-1/(b^6*c^4))^(1/6)*log(1/2*(sqrt(-3)*b*c - b*c)*(-1/(b^6*
c^4))^(1/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) + 1/4*(sq
rt(-3) - 1)*(-1/(b^6*c^4))^(1/6)*log(-1/2*(sqrt(-3)*b*c - b*c)*(-1/(b^6*c^
4))^(1/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) - 1/2*(-1/(
b^6*c^4))^(1/6)*log(b*c*(-1/(b^6*c^4))^(1/6) + ((c*cos(2*b*x + 2*a) + c)/s
in(2*b*x + 2*a))^(1/3)) + 1/2*(-1/(b^6*c^4))^(1/6)*log(-b*c*(-1/(b^6*c^4))
^(1/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3))
```

**Sympy [F]**

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx = \int \frac{1}{(c \cot(a + bx))^{2/3}} dx$$

input

```
integrate(1/(c*cot(b*x+a))**(2/3), x)
```

output

```
Integral((c*cot(a + b*x))**(-2/3), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.81

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx =$$

$$c \left( \frac{\sqrt{3} \log \left( \sqrt{3} c^{1/3} \left( \frac{c}{\tan(bx+a)} \right)^{1/3} + c^{2/3} + \left( \frac{c}{\tan(bx+a)} \right)^{2/3} \right)}{c^{5/3}} - \frac{\sqrt{3} \log \left( -\sqrt{3} c^{1/3} \left( \frac{c}{\tan(bx+a)} \right)^{1/3} + c^{2/3} + \left( \frac{c}{\tan(bx+a)} \right)^{2/3} \right)}{c^{5/3}} + \frac{2 \arctan \left( \frac{\sqrt{3} c^{1/3} + 2 \left( \frac{c}{\tan(bx+a)} \right)^{1/3}}{c^{1/3}} \right)}{c^{5/3}} \right)$$

4b

input

```
integrate(1/(c*cot(b*x+a))^(2/3), x, algorithm="maxima")
```

output

```
-1/4*c*(sqrt(3)*log(sqrt(3)*c^(1/3)*(c/tan(b*x + a))^(1/3) + c^(2/3) + (c/
tan(b*x + a))^(2/3))/c^(5/3) - sqrt(3)*log(-sqrt(3)*c^(1/3)*(c/tan(b*x + a
))^(1/3) + c^(2/3) + (c/tan(b*x + a))^(2/3))/c^(5/3) + 2*arctan((sqrt(3)*c
^(1/3) + 2*(c/tan(b*x + a))^(1/3))/c^(1/3))/c^(5/3) + 2*arctan(-(sqrt(3)*c
^(1/3) - 2*(c/tan(b*x + a))^(1/3))/c^(1/3))/c^(5/3) + 4*arctan((c/tan(b*x
+ a))^(1/3)/c^(1/3))/c^(5/3))/b
```

**Giac [F]**

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx = \int \frac{1}{(c \cot(bx + a))^{2/3}} dx$$

input

```
integrate(1/(c*cot(b*x+a))^(2/3),x, algorithm="giac")
```

output

```
integrate((c*cot(b*x + a))^(-2/3), x)
```

**Mupad [B] (verification not implemented)**

Time = 10.99 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.03

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx = -\frac{(-1)^{1/6} \operatorname{atan}\left(\frac{(-1)^{5/6} (c \cot(a + bx))^{1/3} \operatorname{li}}{c^{1/3}}\right) \operatorname{li}}{b c^{2/3}}$$

$$-\frac{(-1)^{1/6} \ln\left(2 (c \cot(a + bx))^{1/3} + (-1)^{1/6} c^{1/3} + (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2 b c^{2/3}}$$

$$-\frac{(-1)^{1/6} \ln\left(2 (c \cot(a + bx))^{1/3} - (-1)^{1/6} c^{1/3} + (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2 b c^{2/3}}$$

$$+\frac{(-1)^{1/6} \ln\left((-1)^{1/6} c^{1/3} - 2 (c \cot(a + bx))^{1/3} + (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b c^{2/3}}$$

$$+\frac{(-1)^{1/6} \ln\left(2 (c \cot(a + bx))^{1/3} + (-1)^{1/6} c^{1/3} - (-1)^{2/3} \sqrt{3} c^{1/3}\right) \left(-\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b c^{2/3}}$$

input

```
int(1/(c*cot(a + b*x))^(2/3),x)
```

output

```
((-1)^(1/6)*log((-1)^(1/6)*c^(1/3) - 2*(c*cot(a + b*x))^(1/3) + (-1)^(2/3)
*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/4 + 1/4)/(b*c^(2/3)) - ((-1)^(1/6)*log(2*
(c*cot(a + b*x))^(1/3) + (-1)^(1/6)*c^(1/3) + (-1)^(2/3)*3^(1/2)*c^(1/3))*
((3^(1/2)*1i)/2 + 1/2))/(2*b*c^(2/3)) - ((-1)^(1/6)*log(2*(c*cot(a + b*x))
^(1/3) - (-1)^(1/6)*c^(1/3) + (-1)^(2/3)*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/2
- 1/2))/(2*b*c^(2/3)) - ((-1)^(1/6)*atan((( -1)^(5/6)*(c*cot(a + b*x))^(1/3)
)*1i)/c^(1/3))*1i)/(b*c^(2/3)) + ((-1)^(1/6)*log(2*(c*cot(a + b*x))^(1/3)
+ (-1)^(1/6)*c^(1/3) - (-1)^(2/3)*3^(1/2)*c^(1/3))*((3^(1/2)*1i)/4 - 1/4))
/(b*c^(2/3))
```

**Reduce [F]**

$$\int \frac{1}{(c \cot(a + bx))^{2/3}} dx = \frac{\int \frac{1}{\cot(bx+a)^{2/3}} dx}{c^{2/3}}$$

input

```
int(1/(c*cot(b*x+a))^(2/3),x)
```

output

```
int(1/cot(a + b*x)**(2/3),x)/c**(2/3)
```

### 3.22 $\int \frac{1}{(c \cot(a+bx))^{4/3}} dx$

Optimal result	223
Mathematica [C] (verified)	224
Rubi [A] (warning: unable to verify)	224
Maple [A] (verified)	229
Fricas [B] (verification not implemented)	230
Sympy [F]	230
Maxima [A] (verification not implemented)	231
Giac [F]	231
Mupad [B] (verification not implemented)	232
Reduce [F]	233

#### Optimal result

Integrand size = 12, antiderivative size = 244

$$\int \frac{1}{(c \cot(a+bx))^{4/3}} dx = \frac{\arctan\left(\frac{\sqrt[3]{c \cot(a+bx)}}{\sqrt[3]{c}}\right)}{bc^{4/3}} - \frac{\arctan\left(\sqrt{3} - \frac{2\sqrt[3]{c \cot(a+bx)}}{\sqrt[3]{c}}\right)}{2bc^{4/3}} + \frac{\arctan\left(\sqrt{3} + \frac{2\sqrt[3]{c \cot(a+bx)}}{\sqrt[3]{c}}\right)}{2bc^{4/3}} + \frac{3}{bc\sqrt[3]{c \cot(a+bx)}} + \frac{\sqrt{3} \log\left(c^{2/3} - \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a+bx)} + (c \cot(a+bx))^{2/3}\right)}{4bc^{4/3}} - \frac{\sqrt{3} \log\left(c^{2/3} + \sqrt{3}\sqrt[3]{c}\sqrt[3]{c \cot(a+bx)} + (c \cot(a+bx))^{2/3}\right)}{4bc^{4/3}}$$

output

```
arctan((c*cot(b*x+a))^(1/3)/c^(1/3))/b/c^(4/3)+1/2*arctan(-3^(1/2)+2*(c*cot(b*x+a))^(1/3)/c^(1/3))/b/c^(4/3)+1/2*arctan(3^(1/2)+2*(c*cot(b*x+a))^(1/3)/c^(1/3))/b/c^(4/3)+3/b/c/(c*cot(b*x+a))^(1/3)+1/4*3^(1/2)*ln(c^(2/3)-3^(1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b/c^(4/3)-1/4*3^(1/2)*ln(c^(2/3)+3^(1/2)*c^(1/3)*(c*cot(b*x+a))^(1/3)+(c*cot(b*x+a))^(2/3))/b/c^(4/3)
```



**Mathematica [C] (verified)**

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.04

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx = \frac{6 + i\sqrt[6]{\cot^2(a + bx)} \log\left(1 - i\sqrt[6]{\cot^2(a + bx)}\right) - i\sqrt[6]{\cot^2(a + bx)} \log\left(1 + i\sqrt[6]{\cot^2(a + bx)}\right)}{2bc(c \cot(a + bx))^{1/3}}$$

input `Integrate[(c*Cot[a + b*x])^(-4/3),x]`

output  $(6 + I*(\text{Cot}[a + b*x]^2)^{(1/6)}*\text{Log}[1 - I*(\text{Cot}[a + b*x]^2)^{(1/6})] - I*(\text{Cot}[a + b*x]^2)^{(1/6)}*\text{Log}[1 + I*(\text{Cot}[a + b*x]^2)^{(1/6})] + (-1)^{(1/6)}*(\text{Cot}[a + b*x]^2)^{(1/6)}*\text{Log}[1 - (-1)^{(1/6)}*(\text{Cot}[a + b*x]^2)^{(1/6})] - (-1)^{(1/6)}*(\text{Cot}[a + b*x]^2)^{(1/6)}*\text{Log}[1 + (-1)^{(1/6)}*(\text{Cot}[a + b*x]^2)^{(1/6})] + (-1)^{(5/6)}*(\text{Cot}[a + b*x]^2)^{(1/6)}*\text{Log}[1 - (-1)^{(5/6)}*(\text{Cot}[a + b*x]^2)^{(1/6})] - (-1)^{(5/6)}*(\text{Cot}[a + b*x]^2)^{(1/6)}*\text{Log}[1 + (-1)^{(5/6)}*(\text{Cot}[a + b*x]^2)^{(1/6})])/(2 * b * c * (c * \text{Cot}[a + b*x])^{(1/3)})$

**Rubi [A] (warning: unable to verify)**

Time = 0.50 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.90, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.083$ , Rules used = {3042, 3955, 3042, 3957, 266, 824, 27, 216, 1142, 25, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx$$

↓ 3042

$$\int \frac{1}{(-c \tan(a + bx + \frac{\pi}{2}))^{4/3}} dx$$

↓ 3955

$$\frac{3}{bc\sqrt[3]{c \cot(a+bx)}} - \frac{\int (c \cot(a+bx))^{2/3} dx}{c^2}$$

↓ 3042

$$\frac{3}{bc\sqrt[3]{c \cot(a+bx)}} - \frac{\int (-c \tan(a+bx + \frac{\pi}{2}))^{2/3} dx}{c^2}$$

↓ 3957

$$\frac{\int \frac{(c \cot(a+bx))^{2/3}}{\cot^2(a+bx)c^2+c^2} d(c \cot(a+bx))}{bc} + \frac{3}{bc\sqrt[3]{c \cot(a+bx)}}$$

↓ 266

$$\frac{3 \int \frac{c^4 \cot^4(a+bx)}{c^6 \cot^6(a+bx)+c^2} d\sqrt[3]{c \cot(a+bx)}}{bc} + \frac{3}{bc\sqrt[3]{c \cot(a+bx)}}$$

↓ 824

$$3 \left( \frac{1}{3} \int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d\sqrt[3]{c \cot(a+bx)} + \frac{\int -\frac{\sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{2(c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3})} d\sqrt[3]{c \cot(a+bx)}}{3\sqrt[3]{c}} + \frac{\int -\frac{\sqrt[3]{c+\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{2(c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3})} d\sqrt[3]{c \cot(a+bx)}}{3\sqrt[3]{c}} \right)$$

---


$$\frac{3}{bc\sqrt[3]{c \cot(a+bx)}}$$

↓ 27

$$3 \left( \frac{1}{3} \int \frac{1}{c^2 \cot^2(a+bx)+c^{2/3}} d\sqrt[3]{c \cot(a+bx)} - \frac{\int \frac{\sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)-\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d\sqrt[3]{c \cot(a+bx)}}{6\sqrt[3]{c}} - \frac{\int \frac{\sqrt[3]{c+\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx)+\sqrt{3}c^{4/3} \cot(a+bx)+c^{2/3}} d\sqrt[3]{c \cot(a+bx)}}{6\sqrt[3]{c}} \right)$$

---


$$\frac{3}{bc\sqrt[3]{c \cot(a+bx)}}$$

↓ 216

$$3 \left( -\frac{\int \frac{\sqrt[3]{c-\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6 \sqrt[3]{c}} - \frac{\int \frac{\sqrt[3]{c+\sqrt{3}} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6 \sqrt[3]{c}} + \arctan(c) \right)$$

---


$$\frac{3}{bc \sqrt[3]{c \cot(a+bx)}}$$

↓ 1142

$$3 \left( -\frac{-\frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{1}{2} \sqrt{3} \int -\frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6 \sqrt[3]{c}} \right)$$

---


$$\frac{3}{bc \sqrt[3]{c \cot(a+bx)}}$$

↓ 25

$$3 \left( -\frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{1}{2} \sqrt[3]{c} \int \frac{1}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)}}{6 \sqrt[3]{c}} \right)$$

---


$$\frac{3}{bc \sqrt[3]{c \cot(a+bx)}}$$

↓ 1082

$$3 \left( -\frac{\frac{1}{2} \sqrt{3} \int \frac{\sqrt{3} \sqrt[3]{c-2} \sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt{c \cot(a+bx)} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - \frac{1}{\sqrt{3}}} d \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right)}{\sqrt{3}}}{6 \sqrt[3]{c}} - \frac{\int \frac{1}{-c^2 \cot^2(a+bx) - \frac{1}{\sqrt{3}}} d \left( 1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}} \right)}{\sqrt{3}}}{bc} \right)$$

---


$$\frac{3}{bc \sqrt[3]{c \cot(a+bx)}}$$

↓ 217

$$3 \left( \frac{\frac{1}{2}\sqrt{3} \int \frac{\sqrt{3}\sqrt[3]{c-2}\sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) - \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}} d^3 \sqrt[3]{c \cot(a+bx)} + \arctan\left(\sqrt{3}\left(1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}}\right)\right)}{6\sqrt[3]{c}} - \frac{\frac{1}{2}\sqrt{3} \int \frac{\sqrt{3}\sqrt[3]{c+2}\sqrt[3]{c \cot(a+bx)}}{c^2 \cot^2(a+bx) + \sqrt{3}c^{4/3} \cot(a+bx) + c^{2/3}}}{6\sqrt[3]{c}} \right) \frac{bc}{3} \frac{3}{bc^3 \sqrt[3]{c \cot(a+bx)}} \downarrow 1103$$

$$3 \left( \frac{\arctan(c^{2/3} \cot(a+bx))}{3\sqrt[3]{c}} - \frac{\arctan\left(\sqrt{3}\left(1 - \frac{2c^{2/3} \cot(a+bx)}{\sqrt{3}}\right)\right) - \frac{1}{2}\sqrt{3} \log\left(-\sqrt{3}c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3}\right)}{6\sqrt[3]{c}} - \frac{\frac{1}{2}\sqrt{3} \log\left(\sqrt{3}c^{4/3} \cot(a+bx) + c^2 \cot^2(a+bx) + c^{2/3}\right)}{6\sqrt[3]{c}} \right) \frac{bc}{3} \frac{3}{bc^3 \sqrt[3]{c \cot(a+bx)}}$$

input `Int[(c*Cot[a + b*x])^(-4/3), x]`

output `3/(b*c*(c*Cot[a + b*x])^(1/3)) + (3*(ArcTan[c^(2/3)*Cot[a + b*x]]/(3*c^(1/3)) - (ArcTan[Sqrt[3]*(1 - (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]])/Sqrt[3]) - (Sqrt[3]*Log[c^(2/3) - Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(1/3)) - (-ArcTan[Sqrt[3]*(1 + (2*c^(2/3)*Cot[a + b*x])/Sqrt[3]])/Sqrt[3]) + (Sqrt[3]*Log[c^(2/3) + Sqrt[3]*c^(4/3)*Cot[a + b*x] + c^2*Cot[a + b*x]^2])/2)/(6*c^(1/3)))/(b*c)`

**Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 217  $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$   $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[(c_ \cdot x)^m \cdot (a_ + (b_ \cdot x)^2)^p, x\_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k(m+1)-1} \cdot (a + b \cdot x^{2k}/c^2)]^p, x], x, (c \cdot x)^{1/k}], x] /;$   $\text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 824  $\text{Int}[(x_)^m / ((a_ + (b_ \cdot x)^n)), x\_Symbol] \rightarrow \text{Module}[\{r = \text{Numerator}[\text{Rt}[a/b, n]], s = \text{Denominator}[\text{Rt}[a/b, n]], k, u\}, \text{Simp}[u = \text{Int}[(r \cdot \text{Cos}[(2k-1)m \cdot (\text{Pi}/n)] - s \cdot \text{Cos}[(2k-1)(m+1) \cdot (\text{Pi}/n)] \cdot x) / (r^2 - 2rs \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x + s^2 \cdot x^2), x] + \text{Int}[(r \cdot \text{Cos}[(2k-1)m \cdot (\text{Pi}/n)] + s \cdot \text{Cos}[(2k-1)(m+1) \cdot (\text{Pi}/n)] \cdot x) / (r^2 + 2rs \cdot \text{Cos}[(2k-1) \cdot (\text{Pi}/n)] \cdot x + s^2 \cdot x^2), x] ; 2 \cdot (-1)^{m/2} \cdot (r^{m+2} / (a \cdot n \cdot s^m)) \ \text{Int}[1 / (r^2 + s^2 \cdot x^2), x] + 2 \cdot (r^{m+1} / (a \cdot n \cdot s^m)) \ \text{Sum}[u, \{k, 1, (n-2)/4\}], x] /;$   $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[(n-2)/4, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{LtQ}[m, n-1] \ \&\& \ \text{PosQ}[a/b]$

rule 1082  $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$   $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$   $\text{FreeQ}[\{a, b, c\}, x]$

rule 1103  $\text{Int}[(d_ + (e_ \cdot x)) / ((a_ + (b_ \cdot x) + (c_ \cdot x)^2), x\_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$   $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142  $\text{Int}[(d_ + (e_ \cdot x)) / ((a_ + (b_ \cdot x) + (c_ \cdot x)^2), x\_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e) / (2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e / (2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x) / (a + b \cdot x + c \cdot x^2), x], x] /;$   $\text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

```
rule 3955 Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

```
rule 3957 Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

### Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.87

method	result
derivativedivides	$3c \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln\left(\frac{(c \cot(bx+a))^{\frac{2}{3}} - \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}}}{12c^2}\right) + \frac{\arctan\left(\frac{2(c \cot(bx+a))^{\frac{1}{3}} - \sqrt{3}}{(c^2)^{\frac{1}{6}}}\right)}{6(c^2)^{\frac{1}{6}}}}{c^2} - \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln\left(\dots\right)}{c^2}$
default	$3c \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln\left(\frac{(c \cot(bx+a))^{\frac{2}{3}} - \sqrt{3} (c^2)^{\frac{1}{6}} (c \cot(bx+a))^{\frac{1}{3}} + (c^2)^{\frac{1}{3}}}{12c^2}\right) + \frac{\arctan\left(\frac{2(c \cot(bx+a))^{\frac{1}{3}} - \sqrt{3}}{(c^2)^{\frac{1}{6}}}\right)}{6(c^2)^{\frac{1}{6}}}}{c^2} - \frac{\sqrt{3} (c^2)^{\frac{5}{6}} \ln\left(\dots\right)}{c^2}$

```
input int(1/(c*cot(b*x+a))^(4/3), x, method=_RETURNVERBOSE)
```

```
output -3/b*c*(-(1/12/c^2*3^(1/2)*(c^2)^(5/6)*ln((c*cot(b*x+a))^(2/3)-3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/6/(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)-3^(1/2)))-1/12/c^2*3^(1/2)*(c^2)^(5/6)*ln((c*cot(b*x+a))^(2/3)+3^(1/2)*(c^2)^(1/6)*(c*cot(b*x+a))^(1/3)+(c^2)^(1/3))+1/6/(c^2)^(1/6)*arctan(2*(c*cot(b*x+a))^(1/3)/(c^2)^(1/6)+3^(1/2))+1/3/(c^2)^(1/6)*arctan((c*cot(b*x+a))^(1/3)/(c^2)^(1/6)))/c^2-1/c^2/(c*cot(b*x+a))^(1/3))
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 660 vs.  $2(186) = 372$ .

Time = 0.10 (sec) , antiderivative size = 660, normalized size of antiderivative = 2.70

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx = \text{Too large to display}$$

input `integrate(1/(c*cot(b*x+a))^(4/3),x, algorithm="fricas")`

output

```
1/4*(2*(b*c^2*cos(2*b*x + 2*a) + b*c^2)*(-1/(b^6*c^8))^(1/6)*log(b^5*c^7*(
-1/(b^6*c^8))^(5/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) -
2*(b*c^2*cos(2*b*x + 2*a) + b*c^2)*(-1/(b^6*c^8))^(1/6)*log(-b^5*c^7*(-1/
(b^6*c^8))^(5/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) - (s
qrt(-3)*b*c^2 - b*c^2 + (sqrt(-3)*b*c^2 - b*c^2)*cos(2*b*x + 2*a))*(-1/(b^
6*c^8))^(1/6)*log(1/2*(sqrt(-3)*b^5*c^7 + b^5*c^7)*(-1/(b^6*c^8))^(5/6) +
((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) + (sqrt(-3)*b*c^2 - b*c
^2 + (sqrt(-3)*b*c^2 - b*c^2)*cos(2*b*x + 2*a))*(-1/(b^6*c^8))^(1/6)*log(-
1/2*(sqrt(-3)*b^5*c^7 + b^5*c^7)*(-1/(b^6*c^8))^(5/6) + ((c*cos(2*b*x + 2*
a) + c)/sin(2*b*x + 2*a))^(1/3)) - (sqrt(-3)*b*c^2 + b*c^2 + (sqrt(-3)*b*c
^2 + b*c^2)*cos(2*b*x + 2*a))*(-1/(b^6*c^8))^(1/6)*log(1/2*(sqrt(-3)*b^5*c
^7 - b^5*c^7)*(-1/(b^6*c^8))^(5/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x +
2*a))^(1/3)) + (sqrt(-3)*b*c^2 + b*c^2 + (sqrt(-3)*b*c^2 + b*c^2)*cos(2*b
*x + 2*a))*(-1/(b^6*c^8))^(1/6)*log(-1/2*(sqrt(-3)*b^5*c^7 - b^5*c^7)*(-1/
(b^6*c^8))^(5/6) + ((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(1/3)) + 12
*((c*cos(2*b*x + 2*a) + c)/sin(2*b*x + 2*a))^(2/3)*sin(2*b*x + 2*a)/(b*c^
2*cos(2*b*x + 2*a) + b*c^2)
```

**Sympy [F]**

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx = \int \frac{1}{(c \cot(a + bx))^{4/3}} dx$$

input `integrate(1/(c*cot(b*x+a))**(4/3),x)`

output `Integral((c*cot(a + b*x))**(-4/3), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.84

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx =$$

$$c \left( \frac{\sqrt{3} \log \left( \sqrt{3} c^{1/3} \left( \frac{c}{\tan(bx+a)} \right)^{1/3} + c^{2/3} + \left( \frac{c}{\tan(bx+a)} \right)^{2/3} \right)}{c^{1/3}} - \frac{\sqrt{3} \log \left( -\sqrt{3} c^{1/3} \left( \frac{c}{\tan(bx+a)} \right)^{1/3} + c^{2/3} + \left( \frac{c}{\tan(bx+a)} \right)^{2/3} \right)}{c^{1/3}} - \frac{2 \arctan \left( \frac{\sqrt{3} c^{1/3} + 2 \left( \frac{c}{\tan(bx+a)} \right)^{1/3}}{c^{1/3}} \right)}{c^2} \right)$$

4b

input `integrate(1/(c*cot(b*x+a))^(4/3),x, algorithm="maxima")`output `-1/4*c*((sqrt(3)*log(sqrt(3)*c^(1/3)*(c/tan(b*x + a))^(1/3) + c^(2/3) + (c/tan(b*x + a))^(2/3))/c^(1/3) - sqrt(3)*log(-sqrt(3)*c^(1/3)*(c/tan(b*x + a))^(1/3) + c^(2/3) + (c/tan(b*x + a))^(2/3))/c^(1/3) - 2*arctan((sqrt(3)*c^(1/3) + 2*(c/tan(b*x + a))^(1/3))/c^(1/3))/c^(1/3) - 2*arctan(-(sqrt(3)*c^(1/3) - 2*(c/tan(b*x + a))^(1/3))/c^(1/3))/c^(1/3) - 4*arctan((c/tan(b*x + a))^(1/3)/c^(1/3))/c^(1/3))/c^2 - 12/(c^2*(c/tan(b*x + a))^(1/3)))/b`**Giac [F]**

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx = \int \frac{1}{(c \cot(bx + a))^{4/3}} dx$$

input `integrate(1/(c*cot(b*x+a))^(4/3),x, algorithm="giac")`output `integrate((c*cot(b*x + a))^(-4/3), x)`



**Mupad [B] (verification not implemented)**

Time = 9.56 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.14

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx = \frac{3}{bc (c \cot(a + bx))^{1/3}} + \frac{(-1)^{1/6} \operatorname{atan}\left(\frac{(-1)^{2/3} (c \cot(a + bx))^{1/3}}{c^{1/3}}\right) \operatorname{li}}{bc^{4/3}}$$

$$- \frac{(-1)^{1/6} \ln\left(972 b^6 c^{12} + 972 (-1)^{1/6} b^6 c^{35/3} \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right) (c \cot(a + bx))^{1/3}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2 b c^{4/3}}$$

$$- \frac{(-1)^{1/6} \ln\left(972 b^6 c^{12} + 972 (-1)^{1/6} b^6 c^{35/3} \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right) (c \cot(a + bx))^{1/3}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2 b c^{4/3}}$$

$$+ \frac{(-1)^{1/6} \ln\left(972 b^6 c^{12} - 1944 (-1)^{1/6} b^6 c^{35/3} \left(-\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right) (c \cot(a + bx))^{1/3}\right) \left(-\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b c^{4/3}}$$

$$+ \frac{(-1)^{1/6} \ln\left(972 b^6 c^{12} - 1944 (-1)^{1/6} b^6 c^{35/3} \left(\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right) (c \cot(a + bx))^{1/3}\right) \left(\frac{1}{4} + \frac{\sqrt{3} \operatorname{li}}{4}\right)}{b c^{4/3}}$$

input `int(1/(c*cot(a + b*x))^(4/3),x)`output

```
3/(b*c*(c*cot(a + b*x))^(1/3)) + ((-1)^(1/6)*atan((( -1)^(2/3)*(c*cot(a + b
*x))^(1/3))/c^(1/3))*1i)/(b*c^(4/3)) - ((-1)^(1/6)*log(972*b^6*c^12 + 972*
(-1)^(1/6)*b^6*c^(35/3)*((3^(1/2)*1i)/2 - 1/2)*(c*cot(a + b*x))^(1/3))*((3
^(1/2)*1i)/2 - 1/2))/(2*b*c^(4/3)) - ((-1)^(1/6)*log(972*b^6*c^12 + 972*(-
1)^(1/6)*b^6*c^(35/3)*((3^(1/2)*1i)/2 + 1/2)*(c*cot(a + b*x))^(1/3))*((3^(
1/2)*1i)/2 + 1/2))/(2*b*c^(4/3)) + ((-1)^(1/6)*log(972*b^6*c^12 - 1944*(-1
)^(1/6)*b^6*c^(35/3)*((3^(1/2)*1i)/4 - 1/4)*(c*cot(a + b*x))^(1/3))*((3^(1
/2)*1i)/4 - 1/4))/(b*c^(4/3)) + ((-1)^(1/6)*log(972*b^6*c^12 - 1944*(-1)^(
1/6)*b^6*c^(35/3)*((3^(1/2)*1i)/4 + 1/4)*(c*cot(a + b*x))^(1/3))*((3^(1/2
*1i)/4 + 1/4))/(b*c^(4/3))
```

**Reduce [F]**

$$\int \frac{1}{(c \cot(a + bx))^{4/3}} dx = \frac{\int \frac{1}{\cot(bx+a)^{4/3}} dx}{c^{4/3}}$$

input `int(1/(c*cot(b*x+a))^(4/3),x)`

output `int(1/(cot(a + b*x)**(1/3)*cot(a + b*x)),x)/(c**(1/3)*c)`

### 3.23 $\int \cot^n(a + bx) dx$

Optimal result	234
Mathematica [A] (verified)	234
Rubi [A] (verified)	235
Maple [F]	236
Fricas [F]	236
Sympy [F]	237
Maxima [F]	237
Giac [F]	237
Mupad [F(-1)]	238
Reduce [F]	238

#### Optimal result

Integrand size = 8, antiderivative size = 46

$$\int \cot^n(a + bx) dx = -\frac{\cot^{1+n}(a + bx) \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, -\cot^2(a + bx)\right)}{b(1+n)}$$

output

```
-cot(b*x+a)^(1+n)*hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], -cot(b*x+a)^2)/b/(1+n)
```

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \cot^n(a + bx) dx = -\frac{\cot^{1+n}(a + bx) \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, -\cot^2(a + bx)\right)}{b(1+n)}$$

input

```
Integrate[Cot[a + b*x]^n,x]
```

output

```
-((Cot[a + b*x]^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, -Cot[a + b*x]^2])/(b*(1 + n)))
```

**Rubi [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^n(a + bx) dx$$

$$\downarrow 3042$$

$$\int \left(-\tan\left(a + bx + \frac{\pi}{2}\right)\right)^n dx$$

$$\downarrow 3957$$

$$-\frac{\int \frac{\cot^n(a+bx)}{\cot^2(a+bx)+1} d \cot(a+bx)}{b}$$

$$\downarrow 278$$

$$-\frac{\cot^{n+1}(a+bx) \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, -\cot^2(a+bx)\right)}{b(n+1)}$$

input `Int[Cot[a + b*x]^n,x]`

output `-((Cot[a + b*x]^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, -Cot[a + b*x]^2])/(b*(1 + n)))`

**Defintions of rubi rules used**

rule 278

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

### Maple [F]

$$\int \cot (bx + a)^n dx$$

input `int(cot(b*x+a)^n,x)`

output `int(cot(b*x+a)^n,x)`

### Fricas [F]

$$\int \cot^n(a + bx) dx = \int \cot (bx + a)^n dx$$

input `integrate(cot(b*x+a)^n,x, algorithm="fricas")`

output `integral(cot(b*x + a)^n, x)`

**Sympy [F]**

$$\int \cot^n(a + bx) dx = \int \cot^n(a + bx) dx$$

input `integrate(cot(b*x+a)**n,x)`

output `Integral(cot(a + b*x)**n, x)`

**Maxima [F]**

$$\int \cot^n(a + bx) dx = \int \cot(bx + a)^n dx$$

input `integrate(cot(b*x+a)^n,x, algorithm="maxima")`

output `integrate(cot(b*x + a)^n, x)`

**Giac [F]**

$$\int \cot^n(a + bx) dx = \int \cot(bx + a)^n dx$$

input `integrate(cot(b*x+a)^n,x, algorithm="giac")`

output `integrate(cot(b*x + a)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int \cot^n(a + bx) dx = \int \cot(a + bx)^n dx$$

input `int(cot(a + b*x)^n, x)`output `int(cot(a + b*x)^n, x)`**Reduce [F]**

$$\int \cot^n(a + bx) dx = \int \cot(bx + a)^n dx$$

input `int(cot(b*x+a)^n, x)`output `int(cot(a + b*x)**n, x)`

### 3.24 $\int (b \cot(c + dx))^n dx$

Optimal result	239
Mathematica [A] (verified)	239
Rubi [A] (verified)	240
Maple [F]	241
Fricas [F]	241
Sympy [F]	242
Maxima [F]	242
Giac [F]	242
Mupad [F(-1)]	243
Reduce [F]	243

#### Optimal result

Integrand size = 10, antiderivative size = 51

$$\int (b \cot(c + dx))^n dx = -\frac{(b \cot(c + dx))^{1+n} \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, -\cot^2(c + dx)\right)}{bd(1+n)}$$

output

`-(b*cot(d*x+c))^(1+n)*hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], -cot(d*x+c)^2)/b/d/(1+n)`

#### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.02

$$\int (b \cot(c + dx))^n dx = -\frac{\cot(c + dx)(b \cot(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, \frac{1+n}{2}, \frac{3+n}{2}, -\cot^2(c + dx)\right)}{d(1+n)}$$

input

`Integrate[(b*Cot[c + d*x])^n,x]`



output  $-\left(\cot[c + dx] \cdot (b \cot[c + dx])^n \operatorname{Hypergeometric2F1}\left[1, (1 + n)/2, (3 + n)/2, -\cot[c + dx]^2\right]\right) / (d(1 + n))$

### Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (b \cot(c + dx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \left(-b \tan\left(c + dx + \frac{\pi}{2}\right)\right)^n dx \\ & \quad \downarrow \text{3957} \\ & - \frac{b \int \frac{(b \cot(c + dx))^n}{\cot^2(c + dx) b^2 + b^2} d(b \cot(c + dx))}{d} \\ & \quad \downarrow \text{278} \\ & - \frac{(b \cot(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, \frac{n+1}{2}, \frac{n+3}{2}, -\cot^2(c + dx)\right)}{bd(n+1)} \end{aligned}$$

input  $\operatorname{Int}[(b \cot[c + dx])^n, x]$

output  $-\left(\left(b \cot[c + dx]\right)^{(1 + n)} \operatorname{Hypergeometric2F1}\left[1, (1 + n)/2, (3 + n)/2, -\cot[c + dx]^2\right]\right) / (b \cdot d \cdot (1 + n))$

**Defintions of rubi rules used**

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

**Maple [F]**

$$\int (b \cot(dx + c))^n dx$$

input `int((b*cot(d*x+c))^n,x)`

output `int((b*cot(d*x+c))^n,x)`

**Fricas [F]**

$$\int (b \cot(c + dx))^n dx = \int (b \cot(dx + c))^n dx$$

input `integrate((b*cot(d*x+c))^n,x, algorithm="fricas")`

output `integral((b*cot(d*x + c))^n, x)`

**Sympy [F]**

$$\int (b \cot(c + dx))^n dx = \int (b \cot(c + dx))^n dx$$

input `integrate((b*cot(d*x+c))**n,x)`

output `Integral((b*cot(c + d*x))**n, x)`

**Maxima [F]**

$$\int (b \cot(c + dx))^n dx = \int (b \cot(dx + c))^n dx$$

input `integrate((b*cot(d*x+c))^n,x, algorithm="maxima")`

output `integrate((b*cot(d*x + c))^n, x)`

**Giac [F]**

$$\int (b \cot(c + dx))^n dx = \int (b \cot(dx + c))^n dx$$

input `integrate((b*cot(d*x+c))^n,x, algorithm="giac")`

output `integrate((b*cot(d*x + c))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \cot(c + dx))^n dx = \int (b \cot(c + dx))^n dx$$

input `int((b*cot(c + d*x))^n,x)`output `int((b*cot(c + d*x))^n, x)`**Reduce [F]**

$$\int (b \cot(c + dx))^n dx = b^n \left( \int \cot(dx + c)^n dx \right)$$

input `int((b*cot(d*x+c))^n,x)`output `b**n*int(cot(c + d*x)**n,x)`

## 3.25 $\int (a \cot^2(x))^{3/2} dx$

Optimal result . . . . .	244
Mathematica [A] (verified) . . . . .	244
Rubi [A] (verified) . . . . .	245
Maple [A] (verified) . . . . .	247
Fricas [A] (verification not implemented) . . . . .	247
Sympy [F] . . . . .	248
Maxima [A] (verification not implemented) . . . . .	248
Giac [A] (verification not implemented) . . . . .	248
Mupad [F(-1)] . . . . .	249
Reduce [B] (verification not implemented) . . . . .	249

### Optimal result

Integrand size = 10, antiderivative size = 36

$$\int (a \cot^2(x))^{3/2} dx = -\frac{1}{2}a \cot(x) \sqrt{a \cot^2(x)} - a \sqrt{a \cot^2(x)} \log(\sin(x)) \tan(x)$$

output

```
-1/2*a*cot(x)*(a*cot(x)^2)^(1/2)-a*(a*cot(x)^2)^(1/2)*ln(sin(x))*tan(x)
```

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75

$$\int (a \cot^2(x))^{3/2} dx = -\frac{1}{2}a \sqrt{a \cot^2(x)} (\csc^2(x) + 2 \log(\sin(x))) \tan(x)$$

input

```
Integrate[(a*Cot[x]^2)^(3/2),x]
```

output

```
-1/2*(a*Sqrt[a*Cot[x]^2]*(Csc[x]^2 + 2*Log[Sin[x]])*Tan[x])
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.75, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {3042, 4141, 3042, 25, 3954, 25, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cot^2(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \tan \left( x + \frac{\pi}{2} \right)^2 \right)^{3/2} dx \\
 & \quad \downarrow \text{4141} \\
 & a \tan(x) \sqrt{a \cot^2(x)} \int \cot^3(x) dx \\
 & \quad \downarrow \text{3042} \\
 & a \tan(x) \sqrt{a \cot^2(x)} \int -\tan \left( x + \frac{\pi}{2} \right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & -a \tan(x) \sqrt{a \cot^2(x)} \int \tan \left( x + \frac{\pi}{2} \right)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & -a \tan(x) \sqrt{a \cot^2(x)} \left( \frac{\cot^2(x)}{2} - \int -\cot(x) dx \right) \\
 & \quad \downarrow \text{25} \\
 & -a \tan(x) \sqrt{a \cot^2(x)} \left( \int \cot(x) dx + \frac{\cot^2(x)}{2} \right) \\
 & \quad \downarrow \text{3042} \\
 & -a \tan(x) \sqrt{a \cot^2(x)} \left( \int -\tan \left( x + \frac{\pi}{2} \right) dx + \frac{\cot^2(x)}{2} \right) \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
 & -a \tan(x) \sqrt{a \cot^2(x)} \left( \frac{\cot^2(x)}{2} - \int \tan\left(x + \frac{\pi}{2}\right) dx \right) \\
 & \quad \downarrow \text{3956} \\
 & -a \tan(x) \sqrt{a \cot^2(x)} \left( \frac{\cot^2(x)}{2} + \log(\sin(x)) \right)
 \end{aligned}$$

input `Int[(a*Cot[x]^2)^(3/2),x]`

output `-(a*Sqrt[a*Cot[x]^2]*(Cot[x]^2/2 + Log[Sin[x]])*Tan[x])`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{(a \cot(x)^2)^{\frac{3}{2}} (-\cot(x)^2 + \ln(\cot(x)^2 + 1))}{2 \cot(x)^3}$	29
default	$\frac{(a \cot(x)^2)^{\frac{3}{2}} (-\cot(x)^2 + \ln(\cot(x)^2 + 1))}{2 \cot(x)^3}$	29
risch	$a \sqrt{-\frac{\alpha (e^{2ix} + 1)^2}{(e^{2ix} - 1)^2}} (ie^{4ix} \ln(e^{2ix} - 1) + e^{4ix} x - 2ie^{2ix} \ln(e^{2ix} - 1) - 2ie^{2ix} - 2e^{2ix} x + i \ln(e^{2ix} - 1) + x)$ $(e^{2ix} + 1)(e^{2ix} - 1)$	112

input `int((a*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `1/2*(a*cot(x)^2)^(3/2)*(-cot(x)^2+ln(cot(x)^2+1))/cot(x)^3`

**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.44

$$\int (a \cot^2(x))^{3/2} dx = \frac{((a \cos(2x) - a) \log(-\frac{1}{2} \cos(2x) + \frac{1}{2}) - 2a) \sqrt{-\frac{a \cos(2x) + a}{\cos(2x) - 1}}}{2 \sin(2x)}$$

input `integrate((a*cot(x)^2)^(3/2),x, algorithm="fricas")`

output `1/2*((a*cos(2*x) - a)*log(-1/2*cos(2*x) + 1/2) - 2*a)*sqrt(-(a*cos(2*x) + a)/(cos(2*x) - 1))/sin(2*x)`



**Sympy [F]**

$$\int (a \cot^2(x))^{3/2} dx = \int (a \cot^2(x))^{\frac{3}{2}} dx$$

input `integrate((a*cot(x)**2)**(3/2), x)`

output `Integral((a*cot(x)**2)**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.83

$$\int (a \cot^2(x))^{3/2} dx = \frac{1}{2} a^{\frac{3}{2}} \log(\tan(x)^2 + 1) - a^{\frac{3}{2}} \log(\tan(x)) - \frac{a^{\frac{3}{2}}}{2 \tan(x)^2}$$

input `integrate((a*cot(x)^2)^(3/2), x, algorithm="maxima")`

output `1/2*a^(3/2)*log(tan(x)^2 + 1) - a^(3/2)*log(tan(x)) - 1/2*a^(3/2)/tan(x)^2`

**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int (a \cot^2(x))^{3/2} dx = \frac{1}{2} a^{\frac{3}{2}} \left( \frac{1}{\cos(x)^2 - 1} - \log(-\cos(x)^2 + 1) \right) \operatorname{sgn}(\cos(x)) \operatorname{sgn}(\sin(x))$$

input `integrate((a*cot(x)^2)^(3/2), x, algorithm="giac")`

output `1/2*a^(3/2)*(1/(cos(x)^2 - 1) - log(-cos(x)^2 + 1))*sgn(cos(x))*sgn(sin(x))`

**Mupad [F(-1)]**

Timed out.

$$\int (a \cot^2(x))^{3/2} dx = \int (a \cot(x)^2)^{3/2} dx$$

input `int((a*cot(x)^2)^(3/2), x)`output `int((a*cot(x)^2)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.14

$$\int (a \cot^2(x))^{3/2} dx = \frac{\sqrt{a} a \left( 4 \log \left( \tan \left( \frac{x}{2} \right)^2 + 1 \right) \sin(x)^2 - 4 \log \left( \tan \left( \frac{x}{2} \right) \right) \sin(x)^2 + \sin(x)^2 - 2 \right)}{4 \sin(x)^2}$$

input `int((a*cot(x)^2)^(3/2), x)`output `(sqrt(a)*a*(4*log(tan(x/2)**2 + 1)*sin(x)**2 - 4*log(tan(x/2))*sin(x)**2 + sin(x)**2 - 2))/(4*sin(x)**2)`

### 3.26 $\int \sqrt{a \cot^2(x)} dx$

Optimal result	250
Mathematica [A] (verified)	250
Rubi [A] (verified)	251
Maple [A] (verified)	252
Fricas [B] (verification not implemented)	253
Sympy [F]	253
Maxima [A] (verification not implemented)	253
Giac [A] (verification not implemented)	254
Mupad [F(-1)]	254
Reduce [B] (verification not implemented)	254

#### Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \sqrt{a \cot^2(x)} dx = \sqrt{a \cot^2(x)} \log(\sin(x)) \tan(x)$$

output

```
(a*cot(x)^2)^(1/2)*ln(sin(x))*tan(x)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \sqrt{a \cot^2(x)} dx = \sqrt{a \cot^2(x)} \log(\sin(x)) \tan(x)$$

input

```
Integrate[Sqrt[a*Cot[x]^2],x]
```

output

```
Sqrt[a*Cot[x]^2]*Log[Sin[x]]*Tan[x]
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4141, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \cot^2(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \tan\left(x + \frac{\pi}{2}\right)^2} dx \\
 & \quad \downarrow \text{4141} \\
 & \tan(x) \sqrt{a \cot^2(x)} \int \cot(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \tan(x) \sqrt{a \cot^2(x)} \int -\tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{25} \\
 & \tan(x) \left(-\sqrt{a \cot^2(x)}\right) \int \tan\left(x + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3956} \\
 & \tan(x) \sqrt{a \cot^2(x)} \log(\sin(x))
 \end{aligned}$$

input

 $\text{Int}[\text{Sqrt}[a \cdot \text{Cot}[x]^2], x]$ 

output

 $\text{Sqrt}[a \cdot \text{Cot}[x]^2] \cdot \text{Log}[\text{Sin}[x]] \cdot \text{Tan}[x]$

## Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)^(n_)^(p_)], x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

method	result	size
derivativedivides	$-\frac{\sqrt{a \cot(x)^2} \ln(\cot(x)^2 + 1)}{2 \cot(x)}$	22
default	$-\frac{\sqrt{a \cot(x)^2} \ln(\cot(x)^2 + 1)}{2 \cot(x)}$	22
risch	$-\frac{\sqrt{-\frac{a(e^{2ix}+1)^2}{(e^{2ix}-1)^2}}(e^{2ix}-1)x}{e^{2ix}+1} - i\frac{\sqrt{-\frac{a(e^{2ix}+1)^2}{(e^{2ix}-1)^2}}(e^{2ix}-1)\ln(e^{2ix}-1)}{e^{2ix}+1}$	94

input `int((a*cot(x)^2)^(1/2), x, method=_RETURNVERBOSE)`

output `-1/2*(a*cot(x)^2)^(1/2)/cot(x)*ln(cot(x)^2+1)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 43 vs.  $2(14) = 28$ .

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.69

$$\int \sqrt{a \cot^2(x)} dx = \frac{\sqrt{-\frac{a \cos(2x)+a}{\cos(2x)-1}} \log\left(-\frac{1}{2} \cos(2x) + \frac{1}{2}\right) \sin(2x)}{2(\cos(2x) + 1)}$$

input `integrate((a*cot(x)^2)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(-(a*cos(2*x) + a)/(cos(2*x) - 1))*log(-1/2*cos(2*x) + 1/2)*sin(2*x)/(cos(2*x) + 1)`

**Sympy [F]**

$$\int \sqrt{a \cot^2(x)} dx = \int \sqrt{a \cot^2(x)} dx$$

input `integrate((a*cot(x)**2)**(1/2),x)`

output `Integral(sqrt(a*cot(x)**2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \sqrt{a \cot^2(x)} dx = -\frac{1}{2} \sqrt{a} \log(\tan(x)^2 + 1) + \sqrt{a} \log(\tan(x))$$

input `integrate((a*cot(x)^2)^(1/2),x, algorithm="maxima")`

output `-1/2*sqrt(a)*log(tan(x)^2 + 1) + sqrt(a)*log(tan(x))`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \sqrt{a \cot^2(x)} dx = \frac{1}{2} \sqrt{a} \log(-\cos(x)^2 + 1) \operatorname{sgn}(\cos(x)) \operatorname{sgn}(\sin(x))$$

input `integrate((a*cot(x)^2)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(a)*log(-cos(x)^2 + 1)*sgn(cos(x))*sgn(sin(x))`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \cot^2(x)} dx = \int \sqrt{a \cot(x)^2} dx$$

input `int((a*cot(x)^2)^(1/2),x)`

output `int((a*cot(x)^2)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \sqrt{a \cot^2(x)} dx = \sqrt{a} \left( -\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + \log\left(\tan\left(\frac{x}{2}\right)\right) \right)$$

input `int((a*cot(x)^2)^(1/2),x)`

output `sqrt(a)*(-log(tan(x/2)**2 + 1) + log(tan(x/2)))`

### 3.27 $\int \frac{1}{\sqrt{a \cot^2(x)}} dx$

Optimal result	255
Mathematica [A] (verified)	255
Rubi [A] (verified)	256
Maple [A] (verified)	257
Fricas [B] (verification not implemented)	258
Sympy [F]	258
Maxima [A] (verification not implemented)	258
Giac [A] (verification not implemented)	259
Mupad [B] (verification not implemented)	259
Reduce [B] (verification not implemented)	259

#### Optimal result

Integrand size = 10, antiderivative size = 17

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = -\frac{\cot(x) \log(\cos(x))}{\sqrt{a \cot^2(x)}}$$

output `-cot(x)*ln(cos(x))/(a*cot(x)^2)^(1/2)`

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = -\frac{\cot(x) \log(\cos(x))}{\sqrt{a \cot^2(x)}}$$

input `Integrate[1/Sqrt[a*Cot[x]^2],x]`

output `-((Cot[x]*Log[Cos[x]])/Sqrt[a*Cot[x]^2])`



**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {3042, 4141, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \cot^2(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a \tan(x + \frac{\pi}{2})^2}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\cot(x) \int \tan(x) dx}{\sqrt{a \cot^2(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot(x) \int \tan(x) dx}{\sqrt{a \cot^2(x)}} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\cot(x) \log(\cos(x))}{\sqrt{a \cot^2(x)}}
 \end{aligned}$$

input `Int [1/Sqrt [a*Cot [x] ^2] , x]`

output `-((Cot [x]*Log [Cos [x]])/Sqrt [a*Cot [x] ^2])`

## Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.53

method	result	size
derivativedivides	$\frac{\cot(x) \left( \ln(\cot(x)^2 + 1) - 2 \ln(\cot(x)) \right)}{2\sqrt{a \cot(x)^2}}$	26
default	$\frac{\cot(x) \left( \ln(\cot(x)^2 + 1) - 2 \ln(\cot(x)) \right)}{2\sqrt{a \cot(x)^2}}$	26
risch	$-\frac{(e^{2ix} + 1)x}{\sqrt{-\frac{a(e^{2ix} + 1)^2}{(e^{2ix} - 1)^2} (e^{2ix} - 1)}} - \frac{i(e^{2ix} + 1) \ln(e^{2ix} + 1)}{\sqrt{-\frac{a(e^{2ix} + 1)^2}{(e^{2ix} - 1)^2} (e^{2ix} - 1)}}$	94

input `int(1/(a*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*cot(x)*(ln(cot(x)^2+1)-2*ln(cot(x)))/(a*cot(x)^2)^(1/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(15) = 30.

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.65

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = -\frac{\sqrt{-\frac{a \cos(2x)+a}{\cos(2x)-1}} \log\left(\frac{1}{2} \cos(2x) + \frac{1}{2}\right) \sin(2x)}{2(a \cos(2x) + a)}$$

input `integrate(1/(a*cot(x)^2)^(1/2),x, algorithm="fricas")`

output `-1/2*sqrt(-(a*cos(2*x) + a)/(cos(2*x) - 1))*log(1/2*cos(2*x) + 1/2)*sin(2*x)/(a*cos(2*x) + a)`

**Sympy [F]**

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = \int \frac{1}{\sqrt{a \cot^2(x)}} dx$$

input `integrate(1/(a*cot(x)**2)**(1/2),x)`

output `Integral(1/sqrt(a*cot(x)**2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = \frac{\log(\tan(x)^2 + 1)}{2\sqrt{a}}$$

input `integrate(1/(a*cot(x)^2)^(1/2),x, algorithm="maxima")`

output `1/2*log(tan(x)^2 + 1)/sqrt(a)`

**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = -\frac{\log(|\cos(x)|)}{\sqrt{a} \operatorname{sgn}(\cos(x)) \operatorname{sgn}(\sin(x))}$$

input `integrate(1/(a*cot(x)^2)^(1/2),x, algorithm="giac")`output `-log(abs(cos(x)))/(sqrt(a)*sgn(cos(x))*sgn(sin(x)))`**Mupad [B] (verification not implemented)**

Time = 8.78 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.47

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = -\frac{\operatorname{atan}\left(\frac{\sqrt{-a} \cot(x)}{\sqrt{a} \sqrt{\cot(x)^2}}\right)}{\sqrt{-a}}$$

input `int(1/(a*cot(x)^2)^(1/2),x)`output `-atan(((a)^(1/2)*cot(x))/(a^(1/2)*(cot(x)^2)^(1/2)))/(a)^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{a \cot^2(x)}} dx = \frac{\sqrt{a} \left( \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) - \log\left(\tan\left(\frac{x}{2}\right) - 1\right) - \log\left(\tan\left(\frac{x}{2}\right) + 1\right) \right)}{a}$$

input `int(1/(a*cot(x)^2)^(1/2),x)`output `(sqrt(a)*(log(tan(x/2)**2 + 1) - log(tan(x/2) - 1) - log(tan(x/2) + 1)))/a`

$$3.28 \quad \int \frac{1}{(a \cot^2(x))^{3/2}} dx$$

Optimal result	260
Mathematica [A] (verified)	260
Rubi [A] (verified)	261
Maple [A] (verified)	262
Fricas [B] (verification not implemented)	263
Sympy [F]	263
Maxima [A] (verification not implemented)	264
Giac [A] (verification not implemented)	264
Mupad [F(-1)]	264
Reduce [B] (verification not implemented)	265

### Optimal result

Integrand size = 10, antiderivative size = 39

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \frac{\cot(x) \log(\cos(x))}{a \sqrt{a \cot^2(x)}} + \frac{\tan(x)}{2a \sqrt{a \cot^2(x)}}$$

output `cot(x)*ln(cos(x))/a/(a*cot(x)^2)^(1/2)+1/2*tan(x)/a/(a*cot(x)^2)^(1/2)`

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.77

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \frac{2 \cot(x) \log(\cos(x)) + \csc(x) \sec(x)}{2a \sqrt{a \cot^2(x)}}$$

input `Integrate[(a*Cot[x]^2)^(-3/2),x]`

output `(2*Cot[x]*Log[Cos[x]] + Csc[x]*Sec[x])/(2*a*Sqrt[a*Cot[x]^2])`

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \cot^2(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\left(a \tan\left(x + \frac{\pi}{2}\right)\right)^{3/2}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\cot(x) \int \tan^3(x) dx}{a \sqrt{a \cot^2(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot(x) \int \tan(x)^3 dx}{a \sqrt{a \cot^2(x)}} \\
 & \quad \downarrow \text{3954} \\
 & \frac{\cot(x) \left(\frac{\tan^2(x)}{2} - \int \tan(x) dx\right)}{a \sqrt{a \cot^2(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot(x) \left(\frac{\tan^2(x)}{2} - \int \tan(x) dx\right)}{a \sqrt{a \cot^2(x)}} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\cot(x) \left(\frac{\tan^2(x)}{2} + \log(\cos(x))\right)}{a \sqrt{a \cot^2(x)}}
 \end{aligned}$$

input

```
Int[(a*Cot[x]^2)^(-3/2),x]
```

output  $(\cot(x) \cdot (\log(\cos(x)) + \tan(x)^{2/2})) / (a \sqrt{a \cot(x)^2})$

**Defintions of rubi rules used**

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 3954  $\text{Int}(((b \cdot) \tan[(c \cdot) + (d \cdot)(x \cdot)])^{(n \cdot)}, x\_Symbol) \rightarrow \text{Simp}[b \cdot ((b \cdot \tan[c + d \cdot x])^{(n - 1)} / (d \cdot (n - 1))), x] - \text{Simp}[b^2 \text{ Int}[(b \cdot \tan[c + d \cdot x])^{(n - 2)}, x], x] \text{ ; FreeQ}\{b, c, d\}, x \ \&\& \text{GtQ}[n, 1]$

rule 3956  $\text{Int}[\tan[(c \cdot) + (d \cdot)(x \cdot)], x\_Symbol] \rightarrow \text{Simp}[-\log[\text{RemoveContent}[\cos[c + d \cdot x], x]] / d, x] \text{ ; FreeQ}\{c, d\}, x]$

rule 4141  $\text{Int}[(u \cdot) \cdot ((b \cdot) \tan[(e \cdot) + (f \cdot)(x \cdot)])^{(n \cdot)})^{(p \cdot)}, x\_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\tan[e + f \cdot x], x]\}, \text{Simp}[(b \cdot ff^n)^{\text{IntPart}[p]} \cdot ((b \cdot \tan[e + f \cdot x])^{(n \cdot \text{FracPart}[p])} / (\tan[e + f \cdot x] / ff)^{(n \cdot \text{FracPart}[p])}) \text{ Int}[\text{ActivateTrig}[u] \cdot (\tan[e + f \cdot x] / ff)^{(n \cdot p)}, x], x]\} \text{ ; FreeQ}\{b, e, f, n, p\}, x \ \&\& \text{!IntegerQ}[p] \ \&\& \text{IntegerQ}[n] \ \&\& (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d \cdot) \cdot (\text{trig\_})[e + f \cdot x])^{(m \cdot)}] / \text{ ; FreeQ}\{d, m\}, x \ \&\& \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\})]$

**Maple [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$-\frac{\cot(x) \left( \ln(\cot(x)^2 + 1) \cot(x)^2 - 2 \ln(\cot(x)) \cot(x)^2 - 1 \right)}{2 \left( a \cot(x)^2 \right)^{\frac{3}{2}}}$	36
default	$-\frac{\cot(x) \left( \ln(\cot(x)^2 + 1) \cot(x)^2 - 2 \ln(\cot(x)) \cot(x)^2 - 1 \right)}{2 \left( a \cot(x)^2 \right)^{\frac{3}{2}}}$	36
risch	$\frac{i e^{4ix} \ln(e^{2ix} + 1) + e^{4ix} x + 2 i e^{2ix} \ln(e^{2ix} + 1) + 2 i e^{2ix} + 2 e^{2ix} x + i \ln(e^{2ix} + 1) + x}{a(e^{2ix} + 1)(e^{2ix} - 1) \sqrt{-\frac{a(e^{2ix} + 1)^2}{(e^{2ix} - 1)^2}}}$	114

input `int(1/(a*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

output `-1/2*cot(x)*(ln(cot(x)^2+1)*cot(x)^2-2*ln(cot(x))*cot(x)^2-1)/(a*cot(x)^2)^(3/2)`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 74 vs.  $2(33) = 66$ .

Time = 0.07 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.90

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \frac{((\cos(2x) + 1) \log\left(\frac{1}{2} \cos(2x) + \frac{1}{2}\right) \sin(2x) + 2 \sin(2x)) \sqrt{-\frac{a \cos(2x) + a}{\cos(2x) - 1}}}{2(a^2 \cos(2x))^2 + 2a^2 \cos(2x) + a^2}$$

input `integrate(1/(a*cot(x)^2)^(3/2),x, algorithm="fricas")`

output `1/2*((cos(2*x) + 1)*log(1/2*cos(2*x) + 1/2)*sin(2*x) + 2*sin(2*x))*sqrt(-(a*cos(2*x) + a)/(cos(2*x) - 1))/(a^2*cos(2*x)^2 + 2*a^2*cos(2*x) + a^2)`

### Sympy [F]

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \int \frac{1}{(a \cot^2(x))^{\frac{3}{2}}} dx$$

input `integrate(1/(a*cot(x)**2)**(3/2),x)`

output `Integral((a*cot(x)**2)**(-3/2), x)`



**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.56

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \frac{\tan(x)^2}{2 a^{3/2}} - \frac{\log(\tan(x)^2 + 1)}{2 a^{3/2}}$$

input `integrate(1/(a*cot(x)^2)^(3/2),x, algorithm="maxima")`output `1/2*tan(x)^2/a^(3/2) - 1/2*log(tan(x)^2 + 1)/a^(3/2)`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.18

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = -\frac{\frac{\operatorname{sgn}(\sin(x))}{\sqrt{a}} - \frac{2\sqrt{a}\log(|\cos(x)|) + \frac{\sqrt{a}}{\cos(x)^2}}{a\operatorname{sgn}(\cos(x))\operatorname{sgn}(\sin(x))}}{2a}$$

input `integrate(1/(a*cot(x)^2)^(3/2),x, algorithm="giac")`output `-1/2*(sgn(sin(x))/sqrt(a) - (2*sqrt(a)*log(abs(cos(x))) + sqrt(a)/cos(x)^2)/(a*sgn(cos(x))*sgn(sin(x))))/a`**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \int \frac{1}{(a \cot(x)^2)^{3/2}} dx$$

input `int(1/(a*cot(x)^2)^(3/2),x)`output `int(1/(a*cot(x)^2)^(3/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.36

$$\int \frac{1}{(a \cot^2(x))^{3/2}} dx = \frac{\sqrt{a} \left( -2 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^2 + 2 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + 2 \log\left(\tan\left(\frac{x}{2}\right) - 1\right) \sin(x)^2 - 2 \log\left(\tan\left(\frac{x}{2}\right) - 1\right) + 2 \log\left(\tan\left(\frac{x}{2}\right) + 1\right) \sin(x)^2 - 2 \log\left(\tan\left(\frac{x}{2}\right) + 1\right) - \sin(x)^2 \right)}{2a^2 (\sin(x)^2 - 1)}$$

input `int(1/(a*cot(x)^2)^(3/2),x)`output `(sqrt(a)*(-2*log(tan(x/2)**2 + 1)*sin(x)**2 + 2*log(tan(x/2)**2 + 1) + 2*log(tan(x/2) - 1)*sin(x)**2 - 2*log(tan(x/2) - 1) + 2*log(tan(x/2) + 1)*sin(x)**2 - 2*log(tan(x/2) + 1) - sin(x)**2))/(2*a**2*(sin(x)**2 - 1))`

### 3.29 $\int (a \cot^3(x))^{3/2} dx$

Optimal result	266
Mathematica [A] (verified)	267
Rubi [A] (verified)	267
Maple [A] (verified)	272
Fricas [B] (verification not implemented)	273
Sympy [F]	273
Maxima [A] (verification not implemented)	274
Giac [F]	274
Mupad [F(-1)]	274
Reduce [F]	275

#### Optimal result

Integrand size = 10, antiderivative size = 155

$$\int (a \cot^3(x))^{3/2} dx = \frac{2}{3}a\sqrt{a \cot^3(x)} + \frac{a \arctan\left(1 - \sqrt{2}\sqrt{\cot(x)}\right) \sqrt{a \cot^3(x)}}{\sqrt{2} \cot^{\frac{3}{2}}(x)} - \frac{a \arctan\left(1 + \sqrt{2}\sqrt{\cot(x)}\right) \sqrt{a \cot^3(x)}}{\sqrt{2} \cot^{\frac{3}{2}}(x)} + \frac{a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\cot(x)}}{1+\cot(x)}\right) \sqrt{a \cot^3(x)}}{\sqrt{2} \cot^{\frac{3}{2}}(x)} - \frac{2}{7}a \cot^2(x) \sqrt{a \cot^3(x)}$$

output

```
2/3*a*(a*cot(x)^3)^(1/2)-1/2*a*arctan(-1+2^(1/2)*cot(x)^(1/2))*(a*cot(x)^3)^(1/2)*2^(1/2)/cot(x)^(3/2)-1/2*a*arctan(1+2^(1/2)*cot(x)^(1/2))*(a*cot(x)^3)^(1/2)*2^(1/2)/cot(x)^(3/2)+1/2*a*arctanh(2^(1/2)*cot(x)^(1/2)/(1+cot(x)))*(a*cot(x)^3)^(1/2)*2^(1/2)/cot(x)^(3/2)-2/7*a*cot(x)^2*(a*cot(x)^3)^(1/2)
```

### Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.52

$$\int (a \cot^3(x))^{3/2} dx = \frac{a\sqrt{a \cot^3(x)} \left( -21 \arctan \left( \sqrt[4]{-\cot^2(x)} \right) \sqrt[4]{-\cot(x)} + 21 \operatorname{arctanh} \left( \sqrt[4]{-\cot^2(x)} \right) \right)}{21 \cot^{7/4}(x)}$$

input `Integrate[(a*Cot[x]^3)^(3/2),x]`

output `(a*Sqrt[a*Cot[x]^3]*(-21*ArcTan[(-Cot[x]^2)^(1/4)]*(-Cot[x])^(1/4) + 21*ArcTanh[(-Cot[x]^2)^(1/4)]*(-Cot[x])^(1/4) + 2*Cot[x]^(7/4)*(7 - 3*Cot[x]^2)))/(21*Cot[x]^(7/4))`

### Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.96, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.700$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3954, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a \cot^3(x))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \int \left( -a \tan \left( x + \frac{\pi}{2} \right)^3 \right)^{3/2} dx \\ & \quad \downarrow \text{4141} \\ & \frac{a\sqrt{a \cot^3(x)} \int \cot^{9/2}(x) dx}{\cot^{3/2}(x)} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
& \frac{a\sqrt{a\cot^3(x)} \int (-\tan(x + \frac{\pi}{2}))^{9/2} dx}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{3954} \\
& \frac{a\sqrt{a\cot^3(x)} \left( -\int \cot^{\frac{5}{2}}(x) dx - \frac{2}{7} \cot^{\frac{7}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{3042} \\
& \frac{a\sqrt{a\cot^3(x)} \left( -\int (-\tan(x + \frac{\pi}{2}))^{5/2} dx - \frac{2}{7} \cot^{\frac{7}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{3954} \\
& \frac{a\sqrt{a\cot^3(x)} \left( \int \sqrt{\cot(x)} dx - \frac{2}{7} \cot^{\frac{7}{2}}(x) + \frac{2}{3} \cot^{\frac{3}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{3042} \\
& \frac{a\sqrt{a\cot^3(x)} \left( \int \sqrt{-\tan(x + \frac{\pi}{2})} dx - \frac{2}{7} \cot^{\frac{7}{2}}(x) + \frac{2}{3} \cot^{\frac{3}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{3957} \\
& \frac{a\sqrt{a\cot^3(x)} \left( -\int \frac{\sqrt{\cot(x)}}{\cot^2(x)+1} d\cot(x) - \frac{2}{7} \cot^{\frac{7}{2}}(x) + \frac{2}{3} \cot^{\frac{3}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{266} \\
& \frac{a\sqrt{a\cot^3(x)} \left( -2 \int \frac{\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} - \frac{2}{7} \cot^{\frac{7}{2}}(x) + \frac{2}{3} \cot^{\frac{3}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{826} \\
& \frac{a\sqrt{a\cot^3(x)} \left( -2 \left( \frac{1}{2} \int \frac{\cot(x)+1}{\cot^2(x)+1} d\sqrt{\cot(x)} - \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) - \frac{2}{7} \cot^{\frac{7}{2}}(x) + \frac{2}{3} \cot^{\frac{3}{2}}(x) \right)}{\cot^{\frac{3}{2}}(x)} \\
& \quad \downarrow \text{1476} \\
& \frac{a\sqrt{a\cot^3(x)} \left( -2 \left( \frac{1}{2} \int \frac{1}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} + \frac{1}{2} \int \frac{1}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) - \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)}
\end{aligned}$$

↓ 1082

$$\frac{a\sqrt{a\cot^3(x)}\left(-2\left(\frac{1}{2}\left(\frac{\int\frac{1}{\cot(x)-1}d(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}}-\frac{\int\frac{1}{\cot(x)-1}d(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}}\right)-\frac{1}{2}\int\frac{1-\cot(x)}{\cot^2(x)+1}d\sqrt{\cot(x)}\right)-\frac{2}{7}\cot^{\frac{7}{2}}(x)\right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 217

$$\frac{a\sqrt{a\cot^3(x)}\left(-2\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}}\right)-\frac{1}{2}\int\frac{1-\cot(x)}{\cot^2(x)+1}d\sqrt{\cot(x)}\right)-\frac{2}{7}\cot^{\frac{7}{2}}(x)+\frac{2}{3}\cot^{\frac{3}{2}}(x)\right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 1479

$$\frac{a\sqrt{a\cot^3(x)}\left(-2\left(\frac{1}{2}\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}}d\sqrt{\cot(x)}}{2\sqrt{2}}+\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)+1})}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}}d\sqrt{\cot(x)}}{2\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}}\right)\right)-\frac{2}{7}\cot^{\frac{7}{2}}(x)+\frac{2}{3}\cot^{\frac{3}{2}}(x)\right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 25

$$\frac{a\sqrt{a\cot^3(x)}\left(-2\left(\frac{1}{2}\left(-\frac{\int\frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}}d\sqrt{\cot(x)}}{2\sqrt{2}}-\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)+1})}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}}d\sqrt{\cot(x)}}{2\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}}\right)\right)-\frac{2}{7}\cot^{\frac{7}{2}}(x)+\frac{2}{3}\cot^{\frac{3}{2}}(x)\right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 27

$$\frac{a\sqrt{a\cot^3(x)}\left(-2\left(\frac{1}{2}\left(-\frac{\int\frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}}d\sqrt{\cot(x)}}{2\sqrt{2}}-\frac{1}{2}\int\frac{\sqrt{2}\sqrt{\cot(x)+1}}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}}d\sqrt{\cot(x)}\right)+\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}}\right)\right)-\frac{2}{7}\cot^{\frac{7}{2}}(x)+\frac{2}{3}\cot^{\frac{3}{2}}(x)\right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 1103

$$\frac{a\sqrt{a\cot^3(x)}\left(-2\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}}-\frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}}\right)+\frac{1}{2}\left(\frac{\log(\cot(x)-\sqrt{2}\sqrt{\cot(x)+1})}{2\sqrt{2}}-\frac{\log(\cot(x)+\sqrt{2}\sqrt{\cot(x)+1})}{2\sqrt{2}}\right)\right)-\frac{2}{7}\cot^{\frac{7}{2}}(x)+\frac{2}{3}\cot^{\frac{3}{2}}(x)\right)}{\cot^{\frac{3}{2}}(x)}$$

input `Int[(a*Cot[x]^3)^(3/2),x]`

output `(a*Sqrt[a*Cot[x]^3]*((2*Cot[x]^(3/2))/3 - (2*Cot[x]^(7/2))/7 - 2*((-ArcTan[1 - Sqrt[2]*Sqrt[Cot[x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]/(2*Sqrt[2]))/2))/Cot[x]^(3/2)`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`



rule 4141

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Ta
n[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /
; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

**Maple [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{(a \cot(x)^3)^{\frac{3}{2}} \left( 24(a \cot(x))^{\frac{7}{2}} (a^2)^{\frac{1}{4}} - 56a^2(a \cot(x))^{\frac{3}{2}} (a^2)^{\frac{1}{4}} + 21a^4\sqrt{2} \ln \left( -\frac{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2-a \cot(x)-\sqrt{a^2}}}{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2+\sqrt{a^2}}} \right) \right)}{84 \cot(x)^3 (a \cot(x))^{\frac{3}{2}} a^2 (a^2)^{\frac{1}{4}}}$
default	$\frac{(a \cot(x)^3)^{\frac{3}{2}} \left( 24(a \cot(x))^{\frac{7}{2}} (a^2)^{\frac{1}{4}} - 56a^2(a \cot(x))^{\frac{3}{2}} (a^2)^{\frac{1}{4}} + 21a^4\sqrt{2} \ln \left( -\frac{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2-a \cot(x)-\sqrt{a^2}}}{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2+\sqrt{a^2}}} \right) \right)}{84 \cot(x)^3 (a \cot(x))^{\frac{3}{2}} a^2 (a^2)^{\frac{1}{4}}}$

input

```
int((a*cot(x)^3)^(3/2),x,method=_RETURNVERBOSE)
```

output

```
-1/84*(a*cot(x)^3)^(3/2)*(24*(a*cot(x))^(7/2)*(a^2)^(1/4)-56*a^2*(a*cot(x)
)^(3/2)*(a^2)^(1/4)+21*a^4*2^(1/2)*ln(-((a^2)^(1/4)*(a*cot(x))^(1/2)*2^(1/
2)-a*cot(x)-(a^2)^(1/2))/(a*cot(x)+(a^2)^(1/4)*(a*cot(x))^(1/2)*2^(1/2)+(a
^2)^(1/2)))+42*a^4*2^(1/2)*arctan((2^(1/2)*(a*cot(x))^(1/2)+(a^2)^(1/4))/(
a^2)^(1/4))+42*a^4*2^(1/2)*arctan((2^(1/2)*(a*cot(x))^(1/2)-(a^2)^(1/4))/(
a^2)^(1/4)))/cot(x)^3/(a*cot(x))^(3/2)/a^2/(a^2)^(1/4)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 411 vs.  $2(118) = 236$ .

Time = 0.08 (sec) , antiderivative size = 411, normalized size of antiderivative = 2.65

$$\int (a \cot^3(x))^{3/2} dx =$$

$$42 \sqrt{2}(a \cos(2x) - a)\sqrt{a} \arctan\left(\frac{\sqrt{2}\sqrt{a}\sqrt{-\frac{a \cos(2x)^2 + 2a \cos(2x) + a}{(\cos(2x) - 1)\sin(2x)}} \sin(2x) + a \cos(2x) + a}{a \cos(2x) + a}\right) + 42 \sqrt{2}(a \cos(2x) - a)$$

input `integrate((a*cot(x)^3)^(3/2),x, algorithm="fricas")`

output `-1/84*(42*sqrt(2)*(a*cos(2*x) - a)*sqrt(a)*arctan((sqrt(2)*sqrt(a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*sin(2*x) + a*cos(2*x) + a)/(a*cos(2*x) + a)) + 42*sqrt(2)*(a*cos(2*x) - a)*sqrt(a)*arctan((sqrt(2)*sqrt(a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*sin(2*x) - a*cos(2*x) - a)/(a*cos(2*x) + a)) + 21*sqrt(2)*(a*cos(2*x) - a)*sqrt(a)*log((sqrt(2)*sqrt(a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*(cos(2*x) - 1) + a*cos(2*x) + a*sin(2*x) + a)/sin(2*x)) - 21*sqrt(2)*(a*cos(2*x) - a)*sqrt(a)*log(-(sqrt(2)*sqrt(a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*(cos(2*x) - 1) - a*cos(2*x) - a*sin(2*x) - a)/sin(2*x)) - 16*(5*a*cos(2*x) - 2*a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))/(cos(2*x) - 1)`

**Sympy [F]**

$$\int (a \cot^3(x))^{3/2} dx = \int (a \cot^3(x))^{\frac{3}{2}} dx$$

input `integrate((a*cot(x)**3)**(3/2),x)`

output `Integral((a*cot(x)**3)**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.73

$$\int (a \cot^3(x))^{3/2} dx = \frac{1}{4} \left( 2\sqrt{2}\sqrt{a} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(x)})\right) + 2\sqrt{2}\sqrt{a} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(x)})\right) \right) + \frac{2a^{3/2}}{3\tan(x)^{3/2}} - \frac{2a^{3/2}}{7\tan(x)^{7/2}}$$

input `integrate((a*cot(x)^3)^(3/2),x, algorithm="maxima")`

output `1/4*(2*sqrt(2)*sqrt(a)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(x)))) + 2*sqrt(2)*sqrt(a)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(x)))) + sqrt(2)*sqrt(a)*log(sqrt(2)*sqrt(tan(x)) + tan(x) + 1) - sqrt(2)*sqrt(a)*log(-sqrt(2)*sqrt(tan(x)) + tan(x) + 1))*a + 2/3*a^(3/2)/tan(x)^(3/2) - 2/7*a^(3/2)/tan(x)^(7/2)`

**Giac [F]**

$$\int (a \cot^3(x))^{3/2} dx = \int (a \cot(x)^3)^{3/2} dx$$

input `integrate((a*cot(x)^3)^(3/2),x, algorithm="giac")`

output `integrate((a*cot(x)^3)^(3/2), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a \cot^3(x))^{3/2} dx = \int (a \cot(x)^3)^{3/2} dx$$

input `int((a*cot(x)^3)^(3/2),x)`

output `int((a*cot(x)^3)^(3/2), x)`

**Reduce [F]**

$$\int (a \cot^3(x))^{3/2} dx = \sqrt{a} \left( \int \sqrt{\cot(x)} \cot(x)^4 dx \right) a$$

input `int((a*cot(x)^3)^(3/2), x)`

output `sqrt(a)*int(sqrt(cot(x))*cot(x)**4, x)*a`

### 3.30 $\int \sqrt{a \cot^3(x)} dx$

Optimal result	276
Mathematica [A] (verified)	277
Rubi [A] (verified)	277
Maple [A] (verified)	282
Fricas [B] (verification not implemented)	282
Sympy [F]	283
Maxima [A] (verification not implemented)	283
Giac [F]	284
Mupad [F(-1)]	284
Reduce [F]	284

#### Optimal result

Integrand size = 10, antiderivative size = 132

$$\int \sqrt{a \cot^3(x)} dx = -\frac{\arctan\left(1 - \sqrt{2}\sqrt{\cot(x)}\right) \sqrt{a \cot^3(x)}}{\sqrt{2} \cot^{\frac{3}{2}}(x)} + \frac{\arctan\left(1 + \sqrt{2}\sqrt{\cot(x)}\right) \sqrt{a \cot^3(x)}}{\sqrt{2} \cot^{\frac{3}{2}}(x)} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\cot(x)}}{1+\cot(x)}\right) \sqrt{a \cot^3(x)}}{\sqrt{2} \cot^{\frac{3}{2}}(x)} - 2\sqrt{a \cot^3(x)} \tan(x)$$

output

```
1/2*arctan(-1+2^(1/2)*cot(x)^(1/2))*(a*cot(x)^3)^(1/2)*2^(1/2)/cot(x)^(3/2)
)+1/2*arctan(1+2^(1/2)*cot(x)^(1/2))*(a*cot(x)^3)^(1/2)*2^(1/2)/cot(x)^(3/2)
)+1/2*arctanh(2^(1/2)*cot(x)^(1/2)/(1+cot(x)))*(a*cot(x)^3)^(1/2)*2^(1/2)
/cot(x)^(3/2)-2*(a*cot(x)^3)^(1/2)*tan(x)
```

### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.92

$$\int \sqrt{a \cot^3(x)} dx = \frac{\sqrt{a \cot^3(x)} \left( 2\sqrt{2} \arctan \left( 1 - \sqrt{2}\sqrt{\cot(x)} \right) - 2\sqrt{2} \arctan \left( 1 + \sqrt{2}\sqrt{\cot(x)} \right) + 8\sqrt{\cot(x)} + \sqrt{2} \log \left( \dots \right) \right)}{4 \cot^{\frac{3}{2}}(x)}$$

input `Integrate[Sqrt[a*Cot[x]^3],x]`

output `-1/4*(Sqrt[a*Cot[x]^3]*(2*Sqrt[2]*ArcTan[1 - Sqrt[2]*Sqrt[Cot[x]]] - 2*Sqrt[2]*ArcTan[1 + Sqrt[2]*Sqrt[Cot[x]]] + 8*Sqrt[Cot[x]] + Sqrt[2]*Log[1 - Sqrt[2]*Sqrt[Cot[x]] + Cot[x]] - Sqrt[2]*Log[1 + Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]))/Cot[x]^(3/2)`

### Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.03, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a \cot^3(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{-a \tan \left( x + \frac{\pi}{2} \right)^3} dx \\ & \quad \downarrow \text{4141} \\ & \frac{\sqrt{a \cot^3(x)} \int \cot^{\frac{3}{2}}(x) dx}{\cot^{\frac{3}{2}}(x)} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{3042} \\ & \frac{\sqrt{a \cot^3(x)} \int (-\tan(x + \frac{\pi}{2}))^{3/2} dx}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{3954} \\ & \frac{\sqrt{a \cot^3(x)} \left( -\int \frac{1}{\sqrt{\cot(x)}} dx - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{3042} \\ & \frac{\sqrt{a \cot^3(x)} \left( -\int \frac{1}{\sqrt{-\tan(x + \frac{\pi}{2})}} dx - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{3957} \\ & \frac{\sqrt{a \cot^3(x)} \left( \int \frac{1}{\sqrt{\cot(x)(\cot^2(x)+1)}} d \cot(x) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{266} \\ & \frac{\sqrt{a \cot^3(x)} \left( 2 \int \frac{1}{\cot^2(x)+1} d\sqrt{\cot(x)} - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{755} \\ & \frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \int \frac{\cot(x)+1}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{1476} \\ & \frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} + \frac{1}{2} \int \frac{1}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) \right)}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{1082} \\ & \frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \left( \frac{\int \frac{1}{-\cot(x)-1} d(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(x)-1} d(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} \right) \right) \right) - 2\sqrt{\cot(x)}}{\cot^{\frac{3}{2}}(x)} \\ & \downarrow \text{217} \end{aligned}$$

$$\frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \int \frac{1 - \cot(x)}{\cot^2(x) + 1} d\sqrt{\cot(x)} + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 1479

$$\frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \left( - \frac{\int - \frac{\sqrt{2} - 2\sqrt{\cot(x)}}{\cot(x) - \sqrt{2}\sqrt{\cot(x)} + 1} d\sqrt{\cot(x)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)} + 1)}{\cot(x) + \sqrt{2}\sqrt{\cot(x)} + 1} d\sqrt{\cot(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)} + 1)}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\cot(x)}) \right) \right) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 25

$$\frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(x)}}{\cot(x) - \sqrt{2}\sqrt{\cot(x)} + 1} d\sqrt{\cot(x)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)} + 1)}{\cot(x) + \sqrt{2}\sqrt{\cot(x)} + 1} d\sqrt{\cot(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)} + 1)}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\cot(x)}) \right) \right) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 27

$$\frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(x)}}{\cot(x) - \sqrt{2}\sqrt{\cot(x)} + 1} d\sqrt{\cot(x)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(x)} + 1}{\cot(x) + \sqrt{2}\sqrt{\cot(x)} + 1} d\sqrt{\cot(x)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)} + 1)}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\cot(x)}) \right) \right) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)}$$

↓ 1103

$$\frac{\sqrt{a \cot^3(x)} \left( 2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\cot(x) + \sqrt{2}\sqrt{\cot(x)} + 1)}{2\sqrt{2}} - \frac{\log(\cot(x) - \sqrt{2}\sqrt{\cot(x)})}{2\sqrt{2}} \right) \right) - 2\sqrt{\cot(x)} \right)}{\cot^{\frac{3}{2}}(x)}$$

input

Int [Sqrt [a\*Cot [x] ^3] , x]

output

(Sqrt [a\*Cot [x] ^3] \* (-2\*Sqrt [Cot [x]] + 2\*((- (ArcTan [1 - Sqrt [2]\*Sqrt [Cot [x]]] / Sqrt [2]) + ArcTan [1 + Sqrt [2]\*Sqrt [Cot [x]]] / Sqrt [2])) / 2 + (-1/2\*Log [1 - Sqrt [2]\*Sqrt [Cot [x]] + Cot [x]] / Sqrt [2] + Log [1 + Sqrt [2]\*Sqrt [Cot [x]] + Cot [x]] / (2\*Sqrt [2])) / 2)) / Cot [x] ^ (3/2)



## Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

### Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.26

method	result
derivativedivides	$\frac{\sqrt{a \cot(x)^3} \left( -(a^2)^{\frac{1}{4}} \sqrt{2} \ln \left( -\frac{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} + \sqrt{a^2}}{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} - a \cot(x) - \sqrt{a^2}} \right) - 2(a^2)^{\frac{1}{4}} \sqrt{2} \arctan \left( \frac{\sqrt{2} \sqrt{a \cot(x)} + (a^2)^{\frac{1}{4}}}{(a^2)^{\frac{1}{4}}} \right) \right)}{4 \cot(x) \sqrt{a \cot(x)}}$
default	$\frac{\sqrt{a \cot(x)^3} \left( -(a^2)^{\frac{1}{4}} \sqrt{2} \ln \left( -\frac{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} + \sqrt{a^2}}{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} - a \cot(x) - \sqrt{a^2}} \right) - 2(a^2)^{\frac{1}{4}} \sqrt{2} \arctan \left( \frac{\sqrt{2} \sqrt{a \cot(x)} + (a^2)^{\frac{1}{4}}}{(a^2)^{\frac{1}{4}}} \right) \right)}{4 \cot(x) \sqrt{a \cot(x)}}$

input `int((a*cot(x)^3)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/4*(a*\cot(x)^3)^{(1/2)}*(-(a^2)^{(1/4)}*2^{(1/2)}*\ln(-(a*\cot(x)+(a^2)^{(1/4)}*(a*\cot(x))^{(1/2)}*2^{(1/2)+(a^2)^{(1/2)})/((a^2)^{(1/4)}*(a*\cot(x))^{(1/2)}*2^{(1/2)-a*\cot(x)-(a^2)^{(1/2)}))-2*(a^2)^{(1/4)}*2^{(1/2)}*\arctan((2^{(1/2)}*(a*\cot(x))^{(1/2)+(a^2)^{(1/4)})/(a^2)^{(1/4)}-2*(a^2)^{(1/4)}*2^{(1/2)}*\arctan((2^{(1/2)}*(a*\cot(x))^{(1/2)-(a^2)^{(1/4)})/(a^2)^{(1/4)}+8*(a*\cot(x))^{(1/2)})/\cot(x)/(a*\cot(x))^{(1/2)}$$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 387 vs. 2(101) = 202.

Time = 0.08 (sec) , antiderivative size = 387, normalized size of antiderivative = 2.93

$$\int \sqrt{a \cot^3(x)} dx$$

$$= \frac{2 \sqrt{2} \sqrt{a} (\cos(2x) + 1) \arctan \left( \frac{\sqrt{2} \sqrt{a} \sqrt{-\frac{a \cos(2x)^2 + 2a \cos(2x) + a}{(\cos(2x) - 1) \sin(2x)}} \sin(2x) + a \cos(2x) + a}{a \cos(2x) + a} \right) + 2 \sqrt{2} \sqrt{a} (\cos(2x) + 1) \arctan \left( \frac{\sqrt{2} \sqrt{a} \sqrt{-\frac{a \cos(2x)^2 + 2a \cos(2x) + a}{(\cos(2x) - 1) \sin(2x)}} \sin(2x) + a \cos(2x) + a}{a \cos(2x) + a} \right)}{4 \cot(x) \sqrt{a \cot(x)}}$$

input `integrate((a*cot(x)^3)^(1/2),x, algorithm="fricas")`

output

```
1/4*(2*sqrt(2)*sqrt(a)*(cos(2*x) + 1)*arctan((sqrt(2)*sqrt(a)*sqrt(-(a*cos
(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x))))*sin(2*x) + a*cos(2*
x) + a)/(a*cos(2*x) + a)) + 2*sqrt(2)*sqrt(a)*(cos(2*x) + 1)*arctan((sqrt(
2)*sqrt(a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x
))))*sin(2*x) - a*cos(2*x) - a)/(a*cos(2*x) + a)) - sqrt(2)*sqrt(a)*(cos(2*
x) + 1)*log((sqrt(2)*sqrt(a)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos
(2*x) - 1)*sin(2*x))))*(cos(2*x) - 1) + a*cos(2*x) + a*sin(2*x) + a)/sin(2*
x)) + sqrt(2)*sqrt(a)*(cos(2*x) + 1)*log(-(sqrt(2)*sqrt(a)*sqrt(-(a*cos(2*
x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x))))*(cos(2*x) - 1) - a*cos
(2*x) - a*sin(2*x) - a)/sin(2*x)) - 8*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) +
a)/((cos(2*x) - 1)*sin(2*x)))*sin(2*x))/(cos(2*x) + 1)
```

**Sympy [F]**

$$\int \sqrt{a \cot^3(x)} dx = \int \sqrt{a \cot^3(x)} dx$$

input

```
integrate((a*cot(x)**3)**(1/2), x)
```

output

```
Integral(sqrt(a*cot(x)**3), x)
```

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.71

$$\int \sqrt{a \cot^3(x)} dx =$$

$$-\frac{1}{4} \left( 2\sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (\sqrt{2} + 2\sqrt{\tan(x)}) \right) \right) + 2\sqrt{2} \arctan \left( -\frac{1}{2} \sqrt{2} (\sqrt{2} - 2\sqrt{\tan(x)}) \right) - \sqrt{2} \log \left( \frac{2\sqrt{a}}{\sqrt{\tan(x)}} \right)$$

input

```
integrate((a*cot(x)^3)^(1/2), x, algorithm="maxima")
```

output

```
-1/4*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(x)))) + 2*sqrt(2)
*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(x)))) - sqrt(2)*log(sqrt(2)*sq
r t(tan(x)) + tan(x) + 1) + sqrt(2)*log(-sqrt(2)*sqrt(tan(x)) + tan(x) + 1))
*sqrt(a) - 2*sqrt(a)/sqrt(tan(x))
```

**Giac [F]**

$$\int \sqrt{a \cot^3(x)} dx = \int \sqrt{a \cot(x)^3} dx$$

input

```
integrate((a*cot(x)^3)^(1/2),x, algorithm="giac")
```

output

```
integrate(sqrt(a*cot(x)^3), x)
```

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \cot^3(x)} dx = \int \sqrt{a \cot(x)^3} dx$$

input

```
int((a*cot(x)^3)^(1/2),x)
```

output

```
int((a*cot(x)^3)^(1/2), x)
```

**Reduce [F]**

$$\int \sqrt{a \cot^3(x)} dx = \sqrt{a} \left( -2\sqrt{\cot(x)} - \left( \int \frac{\sqrt{\cot(x)}}{\cot(x)} dx \right) \right)$$

input

```
int((a*cot(x)^3)^(1/2),x)
```

output `sqrt(a)*( - 2*sqrt(cot(x)) - int(sqrt(cot(x))/cot(x),x))`

### 3.31 $\int \frac{1}{\sqrt{a \cot^3(x)}} dx$

Optimal result	286
Mathematica [A] (verified)	286
Rubi [A] (verified)	287
Maple [A] (verified)	291
Fricas [B] (verification not implemented)	292
Sympy [F]	293
Maxima [A] (verification not implemented)	293
Giac [F]	294
Mupad [F(-1)]	294
Reduce [F]	294

#### Optimal result

Integrand size = 10, antiderivative size = 133

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \frac{2 \cot(x)}{\sqrt{a \cot^3(x)}} - \frac{\arctan\left(1 - \sqrt{2}\sqrt{\cot(x)}\right) \cot^{\frac{3}{2}}(x)}{\sqrt{2}\sqrt{a \cot^3(x)}} + \frac{\arctan\left(1 + \sqrt{2}\sqrt{\cot(x)}\right) \cot^{\frac{3}{2}}(x)}{\sqrt{2}\sqrt{a \cot^3(x)}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\cot(x)}}{1+\cot(x)}\right) \cot^{\frac{3}{2}}(x)}{\sqrt{2}\sqrt{a \cot^3(x)}}$$

output

```
2*cot(x)/(a*cot(x)^3)^(1/2)+1/2*arctan(-1+2^(1/2)*cot(x)^(1/2))*cot(x)^(3/2)*2^(1/2)/(a*cot(x)^3)^(1/2)+1/2*arctan(1+2^(1/2)*cot(x)^(1/2))*cot(x)^(3/2)*2^(1/2)/(a*cot(x)^3)^(1/2)-1/2*arctanh(2^(1/2)*cot(x)^(1/2)/(1+cot(x)))*cot(x)^(3/2)*2^(1/2)/(a*cot(x)^3)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.45

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \frac{\cot(x) \left( 2 + \arctan\left(\sqrt[4]{-\cot^2(x)}\right) \sqrt[4]{-\cot^2(x)} - \operatorname{arctanh}\left(\sqrt[4]{-\cot^2(x)}\right) \sqrt[4]{-\cot^2(x)} \right)}{\sqrt{a \cot^3(x)}}$$

input `Integrate[1/Sqrt[a*Cot[x]^3],x]`

output `(Cot[x]*(2 + ArcTan[(-Cot[x]^2)^(1/4)]*(-Cot[x]^2)^(1/4) - ArcTanh[(-Cot[x]^2)^(1/4)]*(-Cot[x]^2)^(1/4)))/Sqrt[a*Cot[x]^3]`

### Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$ , Rules used = {3042, 4141, 3042, 3955, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \cot^3(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{-a \tan(x + \frac{\pi}{2})^3}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\cot^{\frac{3}{2}}(x) \int \frac{1}{\cot^{\frac{3}{2}}(x)} dx}{\sqrt{a \cot^3(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot^{\frac{3}{2}}(x) \int \frac{1}{(-\tan(x + \frac{\pi}{2}))^{3/2}} dx}{\sqrt{a \cot^3(x)}} \\
 & \quad \downarrow \text{3955} \\
 & \frac{\cot^{\frac{3}{2}}(x) \left( \frac{2}{\sqrt{\cot(x)}} - \int \sqrt{\cot(x)} dx \right)}{\sqrt{a \cot^3(x)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$



$$\begin{aligned}
& \frac{\cot^{\frac{3}{2}}(x) \left( \frac{2}{\sqrt{\cot(x)}} - \int \sqrt{-\tan\left(x + \frac{\pi}{2}\right)} dx \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{3957} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( \int \frac{\sqrt{\cot(x)}}{\cot^2(x)+1} d\cot(x) + \frac{2}{\sqrt{\cot(x)}} \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{266} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( 2 \int \frac{\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{2}{\sqrt{\cot(x)}} \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{826} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \int \frac{\cot(x)+1}{\cot^2(x)+1} d\sqrt{\cot(x)} - \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) + \frac{2}{\sqrt{\cot(x)}} \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{1476} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} + \frac{1}{2} \int \frac{1}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) - \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{1082} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( \frac{\int \frac{1}{-\cot(x)-1} d(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(x)-1} d(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) + \frac{2}{\sqrt{\cot(x)}} \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{217} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) + \frac{2}{\sqrt{\cot(x)}} \right)}{\sqrt{a \cot^3(x)}} \\
& \quad \downarrow \text{1479} \\
& \frac{\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)+1})}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) \right)}{\sqrt{a \cot^3(x)}}
\end{aligned}$$

$$\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( - \int \frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)+1})}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} \right) \right) \right) \frac{1}{\sqrt{a \cot^3(x)}}$$

$$\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( - \int \frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(x)+1}}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} \right) \right) \right) \frac{1}{\sqrt{a \cot^3(x)}}$$

$$\cot^{\frac{3}{2}}(x) \left( 2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\log(\cot(x)-\sqrt{2}\sqrt{\cot(x)+1})}{2\sqrt{2}} - \frac{\log(\cot(x)+\sqrt{2}\sqrt{\cot(x)+1})}{2\sqrt{2}} \right) \right) \right) \frac{1}{\sqrt{a \cot^3(x)}}$$

input

```
Int[1/Sqrt[a*Cot[x]^3], x]
```

output

```
(Cot[x]^(3/2)*(2/Sqrt[Cot[x]] + 2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]/(2*Sqrt[2]))/2))/Sqrt[a*Cot[x]^3]
```

### Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 217  $\text{Int}[\{(a\_)+(b\_)*(x\_)^2\}^{-1}, x\_Symbol] \rightarrow \text{Simp}[\{-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1}\}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$   $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 266  $\text{Int}[\{(c\_)*(x\_)\}^{m\_}*\{(a\_)+(b\_)*(x\_)^2\}^{p\_}, x\_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(a+b*(x^{2*k}/c^2))^p, x], x, (c*x)^{1/k}], x]] /;$   $\text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 826  $\text{Int}[(x\_)^2/\{(a\_)+(b\_)*(x\_)^4\}, x\_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \ \text{Int}[(r+s*x^2)/(a+b*x^4), x], x] - \text{Simp}[1/(2*s) \ \text{Int}[(r-s*x^2)/(a+b*x^4), x], x]] /;$   $\text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 1082  $\text{Int}[\{(a\_)+(b\_)*(x\_)+(c\_)*(x\_)^2\}^{-1}, x\_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q-x^2), x], x, 1+2*c*(x/b)], x] /;$   $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4*a*c]) /;$   $\text{FreeQ}[\{a, b, c\}, x]$

rule 1103  $\text{Int}[\{(d\_)+(e\_)*(x\_)/\{(a\_)+(b\_)*(x\_)+(c\_)*(x\_)^2\}, x\_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a+b*x+c*x^2, x]]/b), x] /;$   $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1476  $\text{Int}[\{(d\_)+(e\_)*(x\_)^2\}/\{(a\_)+(c\_)*(x\_)^4\}, x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \ \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \ \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /;$   $\text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[d*e]$

rule 1479  $\text{Int}[\{(d\_)+(e\_)*(x\_)^2\}/\{(a\_)+(c\_)*(x\_)^4\}, x\_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \ \text{Int}[(q-2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \ \text{Int}[(q+2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /;$   $\text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

### Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.23

method	result
derivativedivides	$\frac{\cot(x) \left( \sqrt{2} \sqrt{a \cot(x)} \ln \left( -\frac{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} - a \cot(x) - \sqrt{a^2}}{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} + \sqrt{a^2}} \right) + 2\sqrt{2} \sqrt{a \cot(x)} \arctan \left( \frac{\sqrt{2} \sqrt{a \cot(x)} + (a^2)^{\frac{1}{4}}}{(a^2)^{\frac{1}{4}}} \right) + 2\sqrt{2} \sqrt{a \cot(x)} \right)}{4\sqrt{a \cot(x)^3 (a^2)^{\frac{1}{4}}}}$
default	$\frac{\cot(x) \left( \sqrt{2} \sqrt{a \cot(x)} \ln \left( -\frac{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} - a \cot(x) - \sqrt{a^2}}{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} + \sqrt{a^2}} \right) + 2\sqrt{2} \sqrt{a \cot(x)} \arctan \left( \frac{\sqrt{2} \sqrt{a \cot(x)} + (a^2)^{\frac{1}{4}}}{(a^2)^{\frac{1}{4}}} \right) + 2\sqrt{2} \sqrt{a \cot(x)} \right)}{4\sqrt{a \cot(x)^3 (a^2)^{\frac{1}{4}}}}$

input `int(1/(a*cot(x)^3)^(1/2), x, method=_RETURNVERBOSE)`

output

```
1/4*cot(x)*(2^(1/2)*(a*cot(x))^(1/2)*ln(-((a^2)^(1/4)*(a*cot(x))^(1/2)*2^(1/2)-a*cot(x)-(a^2)^(1/2))/(a*cot(x)+(a^2)^(1/4)*(a*cot(x))^(1/2)*2^(1/2)+(a^2)^(1/2)))+2*2^(1/2)*(a*cot(x))^(1/2)*arctan((2^(1/2)*(a*cot(x))^(1/2)+(a^2)^(1/4))/(a^2)^(1/4))+2*2^(1/2)*(a*cot(x))^(1/2)*arctan((2^(1/2)*(a*cot(x))^(1/2)-(a^2)^(1/4))/(a^2)^(1/4))+8*(a^2)^(1/4)/(a*cot(x)^3)^(1/2)/(a^2)^(1/4)
```

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 382 vs.  $2(101) = 202$ .

Time = 0.09 (sec) , antiderivative size = 382, normalized size of antiderivative = 2.87

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx$$

$$= \frac{2\sqrt{2}(a \cos(2x)+a) \arctan\left(\frac{\sqrt{2}\sqrt{-\frac{a \cos(2x)^2+2a \cos(2x)+a}{(\cos(2x)-1)\sin(2x)}} \sin(2x)}{\frac{\sqrt{a}}{\cos(2x)+1}} + \cos(2x)+1\right)}{\sqrt{a}} + \frac{2\sqrt{2}(a \cos(2x)+a) \arctan\left(\frac{\sqrt{2}\sqrt{-\frac{a \cos(2x)^2+2a \cos(2x)+a}{(\cos(2x)-1)\sin(2x)}}}{\frac{\sqrt{a}}{\cos(2x)+1}}\right)}{\sqrt{a}}$$

input

```
integrate(1/(a*cot(x)^3)^(1/2),x, algorithm="fricas")
```

output

```
1/4*(2*sqrt(2)*(a*cos(2*x) + a)*arctan((sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x))))*sin(2*x)/sqrt(a) + cos(2*x) + 1)/(cos(2*x) + 1)/sqrt(a) + 2*sqrt(2)*(a*cos(2*x) + a)*arctan((sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x))))*sin(2*x)/sqrt(a) - cos(2*x) - 1)/(cos(2*x) + 1)/sqrt(a) + sqrt(2)*(a*cos(2*x) + a)*log((sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x))))*(cos(2*x) - 1)/sqrt(a) + cos(2*x) + sin(2*x) + 1)/sin(2*x))/sqrt(a) - sqrt(2)*(a*cos(2*x) + a)*log(-(sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x))))*(cos(2*x) - 1)/sqrt(a) - cos(2*x) - sin(2*x) - 1)/sin(2*x))/sqrt(a) - 8*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*(cos(2*x) - 1)/(a*cos(2*x) + a)
```

**Sympy [F]**

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \int \frac{1}{\sqrt{a \cot^3(x)}} dx$$

input `integrate(1/(a*cot(x)**3)**(1/2),x)`

output `Integral(1/sqrt(a*cot(x)**3), x)`

**Maxima [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \frac{2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(x)}\right)\right) + 2\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(x)}\right)\right) + \sqrt{2} \log\left(\sqrt{2}\sqrt{\tan(x)} + \tan(x) + 1\right) - \sqrt{2} \log\left(-\sqrt{2}\sqrt{\tan(x)} + \tan(x) + 1\right)}{4\sqrt{a}} + \frac{2\sqrt{\tan(x)}}{\sqrt{a}}$$

input `integrate(1/(a*cot(x)^3)^(1/2),x, algorithm="maxima")`

output `-1/4*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(x)))) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(x)))) + sqrt(2)*log(sqrt(2)*sqrt(tan(x)) + tan(x) + 1) - sqrt(2)*log(-sqrt(2)*sqrt(tan(x)) + tan(x) + 1))/sqrt(a) + 2*sqrt(tan(x))/sqrt(a)`

**Giac [F]**

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \int \frac{1}{\sqrt{a \cot(x)^3}} dx$$

input `integrate(1/(a*cot(x)^3)^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(a*cot(x)^3), x)`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \int \frac{1}{\sqrt{a \cot(x)^3}} dx$$

input `int(1/(a*cot(x)^3)^(1/2),x)`

output `int(1/(a*cot(x)^3)^(1/2), x)`

**Reduce [F]**

$$\int \frac{1}{\sqrt{a \cot^3(x)}} dx = \frac{\sqrt{a} \left( \int \frac{\sqrt{\cot(x)}}{\cot(x)^2} dx \right)}{a}$$

input `int(1/(a*cot(x)^3)^(1/2),x)`

output `(sqrt(a)*int(sqrt(cot(x))/cot(x)**2,x))/a`

### 3.32 $\int \frac{1}{(a \cot^3(x))^{3/2}} dx$

Optimal result	295
Mathematica [A] (verified)	296
Rubi [A] (verified)	296
Maple [A] (verified)	301
Fricas [B] (verification not implemented)	302
Sympy [F]	302
Maxima [A] (verification not implemented)	303
Giac [F]	303
Mupad [F(-1)]	304
Reduce [F]	304

#### Optimal result

Integrand size = 10, antiderivative size = 166

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = -\frac{2}{3a\sqrt{a \cot^3(x)}} + \frac{\arctan\left(1 - \sqrt{2}\sqrt{\cot(x)}\right) \cot^{\frac{3}{2}}(x)}{\sqrt{2}a\sqrt{a \cot^3(x)}} - \frac{\arctan\left(1 + \sqrt{2}\sqrt{\cot(x)}\right) \cot^{\frac{3}{2}}(x)}{\sqrt{2}a\sqrt{a \cot^3(x)}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\cot(x)}}{1+\cot(x)}\right) \cot^{\frac{3}{2}}(x)}{\sqrt{2}a\sqrt{a \cot^3(x)}} + \frac{2 \tan^2(x)}{7a\sqrt{a \cot^3(x)}}$$

output

```
-2/3/a/(a*cot(x)^3)^(1/2)-1/2*arctan(-1+2^(1/2)*cot(x)^(1/2))*cot(x)^(3/2)
*2^(1/2)/a/(a*cot(x)^3)^(1/2)-1/2*arctan(1+2^(1/2)*cot(x)^(1/2))*cot(x)^(3
/2)*2^(1/2)/a/(a*cot(x)^3)^(1/2)-1/2*arctanh(2^(1/2)*cot(x)^(1/2)/(1+cot(x
)))*cot(x)^(3/2)*2^(1/2)/a/(a*cot(x)^3)^(1/2)+2/7*tan(x)^2/a/(a*cot(x)^3)^(
1/2)
```



**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.43

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \frac{-14 + 21 \arctan\left(\sqrt[4]{-\cot^2(x)}\right) (-\cot^2(x))^{3/4} + 21 \operatorname{arctanh}\left(\sqrt[4]{-\cot^2(x)}\right) (-\cot^2(x))^{3/4}}{21a\sqrt{a \cot^3(x)}}$$

input `Integrate[(a*Cot[x]^3)^(-3/2),x]`

output `(-14 + 21*ArcTan[(-Cot[x]^2)^(1/4)]*(-Cot[x]^2)^(3/4) + 21*ArcTanh[(-Cot[x]^2)^(1/4)]*(-Cot[x]^2)^(3/4) + 6*Tan[x]^2)/(21*a*Sqrt[a*Cot[x]^3])`

**Rubi [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.91, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.700$ , Rules used = {3042, 4141, 3042, 3955, 3042, 3955, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{1}{(a \cot^3(x))^{3/2}} dx \\ \downarrow 3042 \\ \int \frac{1}{\left(-a \tan\left(x + \frac{\pi}{2}\right)\right)^{3/2}} dx \\ \downarrow 4141 \\ \frac{\cot^{\frac{3}{2}}(x) \int \frac{1}{\cot^{\frac{3}{2}}(x)} dx}{a\sqrt{a \cot^3(x)}} \\ \downarrow 3042 \end{array}$$

$$\frac{\cot^{\frac{3}{2}}(x) \int \frac{1}{(-\tan(x+\frac{\pi}{2}))^{9/2}} dx}{a\sqrt{a \cot^3(x)}}$$

↓ 3955

$$\frac{\cot^{\frac{3}{2}}(x) \left( \frac{2}{7 \cot^{\frac{7}{2}}(x)} - \int \frac{1}{\cot^{\frac{5}{2}}(x)} dx \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 3042

$$\frac{\cot^{\frac{3}{2}}(x) \left( \frac{2}{7 \cot^{\frac{7}{2}}(x)} - \int \frac{1}{(-\tan(x+\frac{\pi}{2}))^{5/2}} dx \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 3955

$$\frac{\cot^{\frac{3}{2}}(x) \left( \int \frac{1}{\sqrt{\cot(x)}} dx - \frac{2}{3 \cot^{\frac{3}{2}}(x)} + \frac{2}{7 \cot^{\frac{7}{2}}(x)} \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 3042

$$\frac{\cot^{\frac{3}{2}}(x) \left( \int \frac{1}{\sqrt{-\tan(x+\frac{\pi}{2})}} dx - \frac{2}{3 \cot^{\frac{3}{2}}(x)} + \frac{2}{7 \cot^{\frac{7}{2}}(x)} \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 3957

$$\frac{\cot^{\frac{3}{2}}(x) \left( - \int \frac{1}{\sqrt{\cot(x)(\cot^2(x)+1)}} d \cot(x) - \frac{2}{3 \cot^{\frac{3}{2}}(x)} + \frac{2}{7 \cot^{\frac{7}{2}}(x)} \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 266

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \int \frac{1}{\cot^2(x)+1} d\sqrt{\cot(x)} - \frac{2}{3 \cot^{\frac{3}{2}}(x)} + \frac{2}{7 \cot^{\frac{7}{2}}(x)} \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 755

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \int \frac{\cot(x)+1}{\cot^2(x)+1} d\sqrt{\cot(x)} \right) - \frac{2}{3 \cot^{\frac{3}{2}}(x)} + \frac{2}{7 \cot^{\frac{7}{2}}(x)} \right)}{a\sqrt{a \cot^3(x)}}$$

↓ 1476

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \left( \frac{1}{2} \int \frac{1}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} + \frac{1}{2} \int \frac{1}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) \right) \right)}{a\sqrt{a\cot^3(x)}}$$

↓ 1082

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \left( \frac{\int \frac{1}{\cot(x)-1} d(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} - \frac{\int \frac{1}{\cot(x)-1} d(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} \right) \right) \right) - \frac{2}{3\cot^{\frac{3}{2}}(x)}}{a\sqrt{a\cot^3(x)}}$$

↓ 217

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \int \frac{1-\cot(x)}{\cot^2(x)+1} d\sqrt{\cot(x)} + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) \right) - \frac{2}{3\cot^{\frac{3}{2}}(x)} + \frac{2}{7\cot^{\frac{7}{2}}(x)}}{a\sqrt{a\cot^3(x)}}$$

↓ 1479

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \left( -\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)+1})}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) \right)}{a\sqrt{a\cot^3(x)}}$$

↓ 25

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(x)+1})}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) \right)}{a\sqrt{a\cot^3(x)}}$$

↓ 27

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \left( \frac{\int \frac{\sqrt{2}-2\sqrt{\cot(x)}}{\cot(x)-\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(x)+1}}{\cot(x)+\sqrt{2}\sqrt{\cot(x)+1}} d\sqrt{\cot(x)} \right) + \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) \right)}{a\sqrt{a\cot^3(x)}}$$

↓ 1103

$$\frac{\cot^{\frac{3}{2}}(x) \left( -2 \left( \frac{1}{2} \left( \frac{\arctan(\sqrt{2}\sqrt{\cot(x)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(x)})}{\sqrt{2}} \right) \right) + \frac{1}{2} \left( \frac{\log(\cot(x)+\sqrt{2}\sqrt{\cot(x)+1})}{2\sqrt{2}} - \frac{\log(\cot(x)-\sqrt{2}\sqrt{\cot(x)})}{2\sqrt{2}} \right) \right)}{a\sqrt{a\cot^3(x)}}$$

input `Int[(a*Cot[x]^3)^(-3/2),x]`

output `(Cot[x]^(3/2)*(2/(7*Cot[x]^(7/2)) - 2/(3*Cot[x]^(3/2)) - 2*((-ArcTan[1 - Sqrt[2]*Sqrt[Cot[x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[x]] + Cot[x]]/(2*Sqrt[2]))/2))/(a*Sqrt[a*Cot[x]^3])`

### Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4141

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])) Int[ActivateTrig[u]*(Ta
n[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /
; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{\cot(x) \left( 21(a^2)^{\frac{1}{4}} \sqrt{2} (a \cot(x))^{\frac{7}{2}} \ln \left( -\frac{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} + \sqrt{a^2}}{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} - a \cot(x) - \sqrt{a^2}} \right) + 42(a^2)^{\frac{1}{4}} \sqrt{2} (a \cot(x))^{\frac{7}{2}} \arctan \left( \frac{\sqrt{2} \sqrt{a \cot(x)}}{a \cot(x) - \sqrt{a^2}} \right) \right)}{84a^4 (a \cot(x)^3)}$
default	$\frac{\cot(x) \left( 21(a^2)^{\frac{1}{4}} \sqrt{2} (a \cot(x))^{\frac{7}{2}} \ln \left( -\frac{a \cot(x) + (a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} + \sqrt{a^2}}{(a^2)^{\frac{1}{4}} \sqrt{a \cot(x)} \sqrt{2} - a \cot(x) - \sqrt{a^2}} \right) + 42(a^2)^{\frac{1}{4}} \sqrt{2} (a \cot(x))^{\frac{7}{2}} \arctan \left( \frac{\sqrt{2} \sqrt{a \cot(x)}}{a \cot(x) - \sqrt{a^2}} \right) \right)}{84a^4 (a \cot(x)^3)}$

input

```
int(1/(a*cot(x)^3)^(3/2), x, method=_RETURNVERBOSE)
```

output

```
-1/84*cot(x)/a^4*(21*(a^2)^(1/4)*2^(1/2)*(a*cot(x))^(7/2)*ln(-(a*cot(x)+(a
^2)^(1/4)*(a*cot(x))^(1/2)*2^(1/2)+(a^2)^(1/2)))/((a^2)^(1/4)*(a*cot(x))^(1
/2)*2^(1/2)-a*cot(x)-(a^2)^(1/2)))+42*(a^2)^(1/4)*2^(1/2)*(a*cot(x))^(7/2)
*arctan((2^(1/2)*(a*cot(x))^(1/2)+(a^2)^(1/4))/(a^2)^(1/4))-42*(a^2)^(1/4)
*2^(1/2)*(a*cot(x))^(7/2)*arctan((-2^(1/2)*(a*cot(x))^(1/2)+(a^2)^(1/4))/(
a^2)^(1/4))+56*a^4*cot(x)^2-24*a^4)/(a*cot(x)^3)^(3/2)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 495 vs.  $2(128) = 256$ .

Time = 0.09 (sec) , antiderivative size = 495, normalized size of antiderivative = 2.98

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \text{Too large to display}$$

input `integrate(1/(a*cot(x)^3)^(3/2),x, algorithm="fricas")`

output

```
1/84*(16*(5*cos(2*x)^2 - 3*cos(2*x) - 2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x)
) + a)/((cos(2*x) - 1)*sin(2*x))*sin(2*x) - 42*sqrt(2)*(a*cos(2*x)^3 + 3*
a*cos(2*x)^2 + 3*a*cos(2*x) + a)*arctan((sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a
*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*sin(2*x)/sqrt(a) + cos(2*x) + 1)
/(cos(2*x) + 1))/sqrt(a) - 42*sqrt(2)*(a*cos(2*x)^3 + 3*a*cos(2*x)^2 + 3*a
*cos(2*x) + a)*arctan((sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((c
os(2*x) - 1)*sin(2*x)))*sin(2*x)/sqrt(a) - cos(2*x) - 1)/(cos(2*x) + 1))/s
qrt(a) + 21*sqrt(2)*(a*cos(2*x)^3 + 3*a*cos(2*x)^2 + 3*a*cos(2*x) + a)*log
((sqrt(2)*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)
)))*(cos(2*x) - 1)/sqrt(a) + cos(2*x) + sin(2*x) + 1)/sin(2*x))/sqrt(a) - 2
1*sqrt(2)*(a*cos(2*x)^3 + 3*a*cos(2*x)^2 + 3*a*cos(2*x) + a)*log(-(sqrt(2)
*sqrt(-(a*cos(2*x)^2 + 2*a*cos(2*x) + a)/((cos(2*x) - 1)*sin(2*x)))*(cos(2
*x) - 1)/sqrt(a) - cos(2*x) - sin(2*x) - 1)/sin(2*x))/sqrt(a))/(a^2*cos(2*
x)^3 + 3*a^2*cos(2*x)^2 + 3*a^2*cos(2*x) + a^2)
```

**Sympy [F]**

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \int \frac{1}{(a \cot^3(x))^{\frac{3}{2}}} dx$$

input `integrate(1/(a*cot(x)**3)**(3/2),x)`

output `Integral((a*cot(x)**3)**(-3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.66

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \frac{2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(x)}\right)\right) + 2\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(x)}\right)\right)}{4a} + \frac{2\left(3\sqrt{a}\tan(x)^{7/2} - 7\sqrt{a}\tan(x)^{3/2}\right)}{21a^2}$$

input `integrate(1/(a*cot(x)^3)^(3/2),x, algorithm="maxima")`output `1/4*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(x)))) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(x)))) - sqrt(2)*log(sqrt(2)*sqrt(tan(x)) + tan(x) + 1) + sqrt(2)*log(-sqrt(2)*sqrt(tan(x)) + tan(x) + 1))/a^(3/2) + 2/21*(3*sqrt(a)*tan(x)^(7/2) - 7*sqrt(a)*tan(x)^(3/2))/a^2`**Giac [F]**

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \int \frac{1}{(a \cot(x)^3)^{3/2}} dx$$

input `integrate(1/(a*cot(x)^3)^(3/2),x, algorithm="giac")`output `integrate((a*cot(x)^3)^(-3/2), x)`



**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \int \frac{1}{(a \cot(x)^3)^{3/2}} dx$$

input `int(1/(a*cot(x)^3)^(3/2), x)`output `int(1/(a*cot(x)^3)^(3/2), x)`**Reduce [F]**

$$\int \frac{1}{(a \cot^3(x))^{3/2}} dx = \frac{\sqrt{a} \left( \int \frac{\sqrt{\cot(x)}}{\cot(x)^5} dx \right)}{a^2}$$

input `int(1/(a*cot(x)^3)^(3/2), x)`output `(sqrt(a)*int(sqrt(cot(x))/cot(x)**5, x))/a**2`

### 3.33 $\int (a \cot^4(x))^{3/2} dx$

Optimal result . . . . .	305
Mathematica [C] (verified) . . . . .	305
Rubi [A] (verified) . . . . .	306
Maple [A] (verified) . . . . .	308
Fricas [A] (verification not implemented) . . . . .	308
Sympy [F] . . . . .	309
Maxima [A] (verification not implemented) . . . . .	309
Giac [A] (verification not implemented) . . . . .	309
Mupad [F(-1)] . . . . .	310
Reduce [B] (verification not implemented) . . . . .	310

#### Optimal result

Integrand size = 10, antiderivative size = 70

$$\int (a \cot^4(x))^{3/2} dx = \frac{1}{3}a \cot(x) \sqrt{a \cot^4(x)} - \frac{1}{5}a \cot^3(x) \sqrt{a \cot^4(x)} - a \sqrt{a \cot^4(x)} \tan(x) - ax \sqrt{a \cot^4(x)} \tan^2(x)$$

output

```
1/3*a*cot(x)*(a*cot(x)^4)^(1/2)-1/5*a*cot(x)^3*(a*cot(x)^4)^(1/2)-a*(a*cot(x)^4)^(1/2)*tan(x)-a*x*(a*cot(x)^4)^(1/2)*tan(x)^2
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.43

$$\int (a \cot^4(x))^{3/2} dx = -\frac{1}{5}(a \cot^4(x))^{3/2} \text{Hypergeometric2F1}\left(-\frac{5}{2}, 1, -\frac{3}{2}, -\tan^2(x)\right) \tan(x)$$

input

```
Integrate[(a*Cot[x]^4)^(3/2),x]
```

output

```
-1/5*((a*Cot[x]^4)^(3/2)*Hypergeometric2F1[-5/2, 1, -3/2, -Tan[x]^2]*Tan[x
1])
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.57, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cot^4(x))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \tan \left( x + \frac{\pi}{2} \right)^4 \right)^{3/2} dx \\
 & \quad \downarrow \text{4141} \\
 & a \tan^2(x) \sqrt{a \cot^4(x)} \int \cot^6(x) dx \\
 & \quad \downarrow \text{3042} \\
 & a \tan^2(x) \sqrt{a \cot^4(x)} \int \tan \left( x + \frac{\pi}{2} \right)^6 dx \\
 & \quad \downarrow \text{3954} \\
 & a \tan^2(x) \sqrt{a \cot^4(x)} \left( - \int \cot^4(x) dx - \frac{1}{5} \cot^5(x) \right) \\
 & \quad \downarrow \text{3042} \\
 & a \tan^2(x) \sqrt{a \cot^4(x)} \left( - \int \tan \left( x + \frac{\pi}{2} \right)^4 dx - \frac{1}{5} \cot^5(x) \right) \\
 & \quad \downarrow \text{3954} \\
 & a \tan^2(x) \sqrt{a \cot^4(x)} \left( \int \cot^2(x) dx - \frac{1}{5} \cot^5(x) + \frac{\cot^3(x)}{3} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
 & a \tan^2(x) \sqrt{a \cot^4(x)} \left( \int \tan \left( x + \frac{\pi}{2} \right)^2 dx - \frac{1}{5} \cot^5(x) + \frac{\cot^3(x)}{3} \right) \\
 & \quad \downarrow \text{3954} \\
 & a \tan^2(x) \sqrt{a \cot^4(x)} \left( - \int 1 dx - \frac{1}{5} \cot^5(x) + \frac{\cot^3(x)}{3} - \cot(x) \right) \\
 & \quad \downarrow \text{24} \\
 & a \tan^2(x) \left( -x - \frac{1}{5} \cot^5(x) + \frac{\cot^3(x)}{3} - \cot(x) \right) \sqrt{a \cot^4(x)}
 \end{aligned}$$

input `Int[(a*Cot[x]^4)^(3/2),x]`

output `a*Sqrt[a*Cot[x]^4]*(-x - Cot[x] + Cot[x]^3/3 - Cot[x]^5/5)*Tan[x]^2`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

**Maple [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.57

method	result	size
derivativedivides	$\frac{(a \cot(x)^4)^{\frac{3}{2}} \left( -3 \cot(x)^5 + 5 \cot(x)^3 + \frac{15\pi}{2} - 15 \operatorname{arccot}(\cot(x)) - 15 \cot(x) \right)}{15 \cot(x)^6}$	40
default	$\frac{(a \cot(x)^4)^{\frac{3}{2}} \left( -3 \cot(x)^5 + 5 \cot(x)^3 + \frac{15\pi}{2} - 15 \operatorname{arccot}(\cot(x)) - 15 \cot(x) \right)}{15 \cot(x)^6}$	40
risch	$\frac{a(e^{2ix}-1)^2 \sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}} x}{(e^{2ix}+1)^2} + \frac{2ia \sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}} (45 e^{8ix} - 90 e^{6ix} + 140 e^{4ix} - 70 e^{2ix} + 23)}{15(e^{2ix}+1)^2 (e^{2ix}-1)^3}$	119

input `int((a*cot(x)^4)^(3/2),x,method=_RETURNVERBOSE)`output `1/15*(a*cot(x)^4)^(3/2)*(-3*cot(x)^5+5*cot(x)^3+15/2*Pi-15*arccot(cot(x))-15*cot(x))/cot(x)^6`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.57

$$\int (a \cot^4(x))^{3/2} dx = \frac{(23 a \cos(2x))^3 - a \cos(2x)^2 - 11 a \cos(2x) + 15 (ax \cos(2x))^2 - 2 ax \cos(2x) + a}{15 (\cos(2x)^2 - 1) \sin(2x)}$$

input `integrate((a*cot(x)^4)^(3/2),x, algorithm="fricas")`output `1/15*(23*a*cos(2*x)^3 - a*cos(2*x)^2 - 11*a*cos(2*x) + 15*(a*x*cos(2*x))^2 - 2*a*x*cos(2*x) + a*x)*sin(2*x) + 13*a)*sqrt((a*cos(2*x)^2 + 2*a*cos(2*x) + a)/(cos(2*x)^2 - 2*cos(2*x) + 1))/((cos(2*x)^2 - 1)*sin(2*x))`

**Sympy [F]**

$$\int (a \cot^4(x))^{3/2} dx = \int (a \cot^4(x))^{\frac{3}{2}} dx$$

input `integrate((a*cot(x)**4)**(3/2),x)`

output `Integral((a*cot(x)**4)**(3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.53

$$\int (a \cot^4(x))^{3/2} dx = -a^{\frac{3}{2}}x - \frac{15 a^{\frac{3}{2}} \tan(x)^4 - 5 a^{\frac{3}{2}} \tan(x)^2 + 3 a^{\frac{3}{2}}}{15 \tan(x)^5}$$

input `integrate((a*cot(x)^4)^(3/2),x, algorithm="maxima")`

output `-a^(3/2)*x - 1/15*(15*a^(3/2)*tan(x)^4 - 5*a^(3/2)*tan(x)^2 + 3*a^(3/2))/tan(x)^5`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.81

$$\int (a \cot^4(x))^{3/2} dx = \frac{1}{480} \left( 3 \tan\left(\frac{1}{2}x\right)^5 - 35 \tan\left(\frac{1}{2}x\right)^3 - 480x - \frac{330 \tan\left(\frac{1}{2}x\right)^4 - 35 \tan\left(\frac{1}{2}x\right)^2 + 3}{\tan\left(\frac{1}{2}x\right)^5} \right)$$

input `integrate((a*cot(x)^4)^(3/2),x, algorithm="giac")`

output `1/480*(3*tan(1/2*x)^5 - 35*tan(1/2*x)^3 - 480*x - (330*tan(1/2*x)^4 - 35*tan(1/2*x)^2 + 3)/tan(1/2*x)^5 + 330*tan(1/2*x))*a^(3/2)`

**Mupad [F(-1)]**

Timed out.

$$\int (a \cot^4(x))^{3/2} dx = \int (a \cot(x)^4)^{3/2} dx$$

input `int((a*cot(x)^4)^(3/2),x)`output `int((a*cot(x)^4)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.36

$$\int (a \cot^4(x))^{3/2} dx = \frac{\sqrt{a} a (-3 \cot(x)^5 + 5 \cot(x)^3 - 15 \cot(x) - 15x)}{15}$$

input `int((a*cot(x)^4)^(3/2),x)`output `(sqrt(a)*a*( - 3*cot(x)**5 + 5*cot(x)**3 - 15*cot(x) - 15*x))/15`

### 3.34 $\int \sqrt{a \cot^4(x)} dx$

Optimal result	311
Mathematica [C] (verified)	311
Rubi [A] (verified)	312
Maple [A] (verified)	313
Fricas [B] (verification not implemented)	314
Sympy [F]	314
Maxima [A] (verification not implemented)	314
Giac [A] (verification not implemented)	315
Mupad [F(-1)]	315
Reduce [B] (verification not implemented)	315

#### Optimal result

Integrand size = 10, antiderivative size = 32

$$\int \sqrt{a \cot^4(x)} dx = -\sqrt{a \cot^4(x)} \tan(x) - x \sqrt{a \cot^4(x)} \tan^2(x)$$

output

```
-(a*cot(x)^4)^(1/2)*tan(x)-x*(a*cot(x)^4)^(1/2)*tan(x)^2
```

#### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \sqrt{a \cot^4(x)} dx = -\sqrt{a \cot^4(x)} \operatorname{Hypergeometric2F1} \left( -\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(x) \right) \tan(x)$$

input

```
Integrate[Sqrt[a*Cot[x]^4],x]
```

output

```
-(Sqrt[a*Cot[x]^4]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[x]^2]*Tan[x])
```



**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.72, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4141, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a \cot^4(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a \tan\left(x + \frac{\pi}{2}\right)^4} dx \\
 & \quad \downarrow \text{4141} \\
 & \tan^2(x) \sqrt{a \cot^4(x)} \int \cot^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \tan^2(x) \sqrt{a \cot^4(x)} \int \tan\left(x + \frac{\pi}{2}\right)^2 dx \\
 & \quad \downarrow \text{3954} \\
 & \tan^2(x) \sqrt{a \cot^4(x)} \left(-\int 1 dx - \cot(x)\right) \\
 & \quad \downarrow \text{24} \\
 & \tan^2(x) (-x - \cot(x)) \sqrt{a \cot^4(x)}
 \end{aligned}$$

input

```
Int[Sqrt[a*Cot[x]^4],x]
```

output

```
(-x - Cot[x])*Sqrt[a*Cot[x]^4]*Tan[x]^2
```

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\sqrt{a \cot(x)^4} \left(-\cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x))\right)}{\cot(x)^2}$	27
default	$\frac{\sqrt{a \cot(x)^4} \left(-\cot(x) + \frac{\pi}{2} - \operatorname{arccot}(\cot(x))\right)}{\cot(x)^2}$	27
risch	$\sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}} (e^{2ix}-1)^2 x + \frac{2i \sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}} (e^{2ix}-1)}{(e^{2ix}+1)^2}$	85

input `int((a*cot(x)^4)^(1/2), x, method=_RETURNVERBOSE)`

output `(a*cot(x)^4)^(1/2)/cot(x)^2*(-cot(x)+1/2*Pi-arccot(cot(x)))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 61 vs.  $2(28) = 56$ .

Time = 0.07 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.91

$$\int \sqrt{a \cot^4(x)} dx = \frac{(x \cos(2x) - x - \sin(2x)) \sqrt{\frac{a \cos(2x)^2 + 2a \cos(2x) + a}{\cos(2x)^2 - 2 \cos(2x) + 1}}}{\cos(2x) + 1}$$

input `integrate((a*cot(x)^4)^(1/2),x, algorithm="fricas")`

output `(x*cos(2*x) - x - sin(2*x))*sqrt((a*cos(2*x)^2 + 2*a*cos(2*x) + a)/(cos(2*x)^2 - 2*cos(2*x) + 1))/(cos(2*x) + 1)`

**Sympy [F]**

$$\int \sqrt{a \cot^4(x)} dx = \int \sqrt{a \cot^4(x)} dx$$

input `integrate((a*cot(x)**4)**(1/2),x)`

output `Integral(sqrt(a*cot(x)**4), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.50

$$\int \sqrt{a \cot^4(x)} dx = -\sqrt{ax} - \frac{\sqrt{a}}{\tan(x)}$$

input `integrate((a*cot(x)^4)^(1/2),x, algorithm="maxima")`

output `-sqrt(a)*x - sqrt(a)/tan(x)`

**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.66

$$\int \sqrt{a \cot^4(x)} dx = -\frac{1}{2} \sqrt{a} \left( 2x + \frac{1}{\tan\left(\frac{1}{2}x\right)} - \tan\left(\frac{1}{2}x\right) \right)$$

input `integrate((a*cot(x)^4)^(1/2),x, algorithm="giac")`

output `-1/2*sqrt(a)*(2*x + 1/tan(1/2*x) - tan(1/2*x))`

**Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a \cot^4(x)} dx = \int \sqrt{a \cot(x)^4} dx$$

input `int((a*cot(x)^4)^(1/2),x)`

output `int((a*cot(x)^4)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.25

$$\int \sqrt{a \cot^4(x)} dx = -\sqrt{a} (\cot(x) + x)$$

input `int((a*cot(x)^4)^(1/2),x)`

output `- sqrt(a)*(cot(x) + x)`

### 3.35 $\int \frac{1}{\sqrt{a \cot^4(x)}} dx$

Optimal result	316
Mathematica [A] (verified)	316
Rubi [A] (verified)	317
Maple [A] (verified)	318
Fricas [B] (verification not implemented)	319
Sympy [F]	319
Maxima [A] (verification not implemented)	319
Giac [F(-2)]	320
Mupad [F(-1)]	320
Reduce [B] (verification not implemented)	320

#### Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \frac{\cot(x)}{\sqrt{a \cot^4(x)}} - \frac{x \cot^2(x)}{\sqrt{a \cot^4(x)}}$$

output

```
cot(x)/(a*cot(x)^4)^(1/2)-x*cot(x)^2/(a*cot(x)^4)^(1/2)
```

#### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \frac{\cot(x) - \arctan(\tan(x)) \cot^2(x)}{\sqrt{a \cot^4(x)}}$$

input

```
Integrate[1/Sqrt[a*Cot[x]^4],x]
```

output

```
(Cot[x] - ArcTan[Tan[x]]*Cot[x]^2)/Sqrt[a*Cot[x]^4]
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {3042, 4141, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{a \cot^4(x)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\sqrt{a \tan(x + \frac{\pi}{2})^4}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\cot^2(x) \int \tan^2(x) dx}{\sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot^2(x) \int \tan(x)^2 dx}{\sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3954} \\
 & \frac{\cot^2(x)(\tan(x) - \int 1 dx)}{\sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{24} \\
 & \frac{(\tan(x) - x) \cot^2(x)}{\sqrt{a \cot^4(x)}}
 \end{aligned}$$

input `Int [1/Sqrt [a*Cot [x]^4] ,x]`

output `(Cot [x]^2*(-x + Tan [x]))/Sqrt [a*Cot [x]^4]`

## Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

## Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$\frac{\cot(x)\left(\left(\frac{\pi}{2}-\operatorname{arccot}(\cot(x))\right)\cot(x)+1\right)}{\sqrt{a\cot(x)^4}}$	26
default	$\frac{\cot(x)\left(\left(\frac{\pi}{2}-\operatorname{arccot}(\cot(x))\right)\cot(x)+1\right)}{\sqrt{a\cot(x)^4}}$	26
risch	$\frac{(e^{2ix}+1)^2x}{\sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}(e^{2ix}-1)^2}} - \frac{2i(e^{2ix}+1)}{\sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}(e^{2ix}-1)^2}}$	85

input `int(1/(a*cot(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

output `cot(x)*((1/2*Pi-arccot(cot(x)))*cot(x)+1)/(a*cot(x)^4)^(1/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 80 vs.  $2(27) = 54$ .

Time = 0.09 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.58

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \frac{(x \cos(2x))^2 - (\cos(2x) - 1) \sin(2x) - x \sqrt{\frac{a \cos(2x)^2 + 2a \cos(2x) + a}{\cos(2x)^2 - 2 \cos(2x) + 1}}}{a \cos(2x)^2 + 2a \cos(2x) + a}$$

input `integrate(1/(a*cot(x)^4)^(1/2),x, algorithm="fricas")`

output `(x*cos(2*x)^2 - (cos(2*x) - 1)*sin(2*x) - x)*sqrt((a*cos(2*x)^2 + 2*a*cos(2*x) + a)/(cos(2*x)^2 - 2*cos(2*x) + 1))/(a*cos(2*x)^2 + 2*a*cos(2*x) + a)`

**Sympy [F]**

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \int \frac{1}{\sqrt{a \cot^4(x)}} dx$$

input `integrate(1/(a*cot(x)**4)**(1/2),x)`

output `Integral(1/sqrt(a*cot(x)**4), x)`

**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.42

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = -\frac{x}{\sqrt{a}} + \frac{\tan(x)}{\sqrt{a}}$$

input `integrate(1/(a*cot(x)^4)^(1/2),x, algorithm="maxima")`

output `-x/sqrt(a) + tan(x)/sqrt(a)`



**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a*cot(x)^4)^(1/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \int \frac{1}{\sqrt{a \cot(x)^4}} dx$$

input `int(1/(a*cot(x)^4)^(1/2),x)`

output `int(1/(a*cot(x)^4)^(1/2), x)`

**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.55

$$\int \frac{1}{\sqrt{a \cot^4(x)}} dx = \frac{\sqrt{a}(-\cot(x)x + 1)}{\cot(x)a}$$

input `int(1/(a*cot(x)^4)^(1/2),x)`

output `(sqrt(a)*(-cot(x)*x + 1))/(cot(x)*a)`

$$3.36 \quad \int \frac{1}{(a \cot^4(x))^{3/2}} dx$$

Optimal result	321
Mathematica [A] (verified)	321
Rubi [A] (verified)	322
Maple [A] (verified)	324
Fricas [B] (verification not implemented)	324
Sympy [F]	325
Maxima [A] (verification not implemented)	325
Giac [F(-2)]	325
Mupad [F(-1)]	326
Reduce [B] (verification not implemented)	326

### Optimal result

Integrand size = 10, antiderivative size = 77

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \frac{\cot(x)}{a\sqrt{a \cot^4(x)}} - \frac{x \cot^2(x)}{a\sqrt{a \cot^4(x)}} - \frac{\tan(x)}{3a\sqrt{a \cot^4(x)}} + \frac{\tan^3(x)}{5a\sqrt{a \cot^4(x)}}$$

output

```
cot(x)/a/(a*cot(x)^4)^(1/2)-x*cot(x)^2/a/(a*cot(x)^4)^(1/2)-1/3*tan(x)/a/(a*cot(x)^4)^(1/2)+1/5*tan(x)^3/a/(a*cot(x)^4)^(1/2)
```

### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.53

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \frac{15 \cot(x) - 15 \arctan(\tan(x)) \cot^2(x) - 5 \tan(x) + 3 \tan^3(x)}{15a\sqrt{a \cot^4(x)}}$$

input

```
Integrate[(a*Cot[x]^4)^(-3/2),x]
```

output

```
(15*Cot[x] - 15*ArcTan[Tan[x]]*Cot[x]^2 - 5*Tan[x] + 3*Tan[x]^3)/(15*a*Sqrt[a*Cot[x]^4])
```

**Rubi [A] (verified)**

Time = 0.34 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.52, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {3042, 4141, 3042, 3954, 3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a \cot^4(x))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\left(a \tan\left(x + \frac{\pi}{2}\right)\right)^{3/2}} dx \\
 & \quad \downarrow \text{4141} \\
 & \frac{\cot^2(x) \int \tan^6(x) dx}{a \sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot^2(x) \int \tan(x)^6 dx}{a \sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3954} \\
 & \frac{\cot^2(x) \left(\frac{\tan^5(x)}{5} - \int \tan^4(x) dx\right)}{a \sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\cot^2(x) \left(\frac{\tan^5(x)}{5} - \int \tan(x)^4 dx\right)}{a \sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3954} \\
 & \frac{\cot^2(x) \left(\int \tan^2(x) dx + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3}\right)}{a \sqrt{a \cot^4(x)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{\cot^2(x) \left( \int \tan(x)^2 dx + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} \right)}{a\sqrt{a \cot^4(x)}}$$

$$\downarrow \text{3954}$$

$$\frac{\cot^2(x) \left( -\int 1 dx + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} + \tan(x) \right)}{a\sqrt{a \cot^4(x)}}$$

$$\downarrow \text{24}$$

$$\frac{\left( -x + \frac{\tan^5(x)}{5} - \frac{\tan^3(x)}{3} + \tan(x) \right) \cot^2(x)}{a\sqrt{a \cot^4(x)}}$$

input `Int[(a*Cot[x]^4)^(-3/2),x]`

output `(Cot[x]^2*(-x + Tan[x] - Tan[x]^3/3 + Tan[x]^5/5))/(a*Sqrt[a*Cot[x]^4])`

### Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 4141 `Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])] Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]`

**Maple [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$\frac{\cot(x) \left( 15 \left( \frac{\pi}{2} - \operatorname{arccot}(\cot(x)) \right) \cot(x)^5 + 15 \cot(x)^4 - 5 \cot(x)^2 + 3 \right)}{15 \left( a \cot(x)^4 \right)^{\frac{3}{2}}}$	42
default	$\frac{\cot(x) \left( 15 \left( \frac{\pi}{2} - \operatorname{arccot}(\cot(x)) \right) \cot(x)^5 + 15 \cot(x)^4 - 5 \cot(x)^2 + 3 \right)}{15 \left( a \cot(x)^4 \right)^{\frac{3}{2}}}$	42
risch	$\frac{(e^{2ix}+1)^2 x}{a(e^{2ix}-1)^2 \sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}}} - \frac{2i(45 e^{8ix}+90 e^{6ix}+140 e^{4ix}+70 e^{2ix}+23)}{15a(e^{2ix}+1)^3 (e^{2ix}-1)^2 \sqrt{\frac{a(e^{2ix}+1)^4}{(e^{2ix}-1)^4}}}$	123

input `int(1/(a*cot(x)^4)^(3/2),x,method=_RETURNVERBOSE)`

output `1/15*cot(x)*(15*(1/2*Pi-arccot(cot(x)))*cot(x)^5+15*cot(x)^4-5*cot(x)^2+3)/(a*cot(x)^4)^(3/2)`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. 2(65) = 130.

Time = 0.08 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.84

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \frac{(15 x \cos(2x))^4 + 30 x \cos(2x)^3 - 30 x \cos(2x) - (23 \cos(2x))^3 + \cos(2x)^2 - 11 \cos(2x)}{15 (a^2 \cos(2x))^4 + 4 a^2 \cos(2x)^3 + 6 a^2 \cos(2x)^2 + \dots}$$

input `integrate(1/(a*cot(x)^4)^(3/2),x, algorithm="fricas")`

output `1/15*(15*x*cos(2*x)^4 + 30*x*cos(2*x)^3 - 30*x*cos(2*x) - (23*cos(2*x))^3 + cos(2*x)^2 - 11*cos(2*x) - 13)*sin(2*x) - 15*x)*sqrt((a*cos(2*x)^2 + 2*a*cos(2*x) + a)/(cos(2*x)^2 - 2*cos(2*x) + 1))/(a^2*cos(2*x)^4 + 4*a^2*cos(2*x)^3 + 6*a^2*cos(2*x)^2 + 4*a^2*cos(2*x) + a^2)`

**Sympy [F]**

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \int \frac{1}{(a \cot^4(x))^{\frac{3}{2}}} dx$$

input `integrate(1/(a*cot(x)**4)**(3/2),x)`

output `Integral((a*cot(x)**4)**(-3/2), x)`

**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.38

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \frac{3 \tan(x)^5 - 5 \tan(x)^3 + 15 \tan(x)}{15 a^{\frac{3}{2}}} - \frac{x}{a^{\frac{3}{2}}}$$

input `integrate(1/(a*cot(x)^4)^(3/2),x, algorithm="maxima")`

output `1/15*(3*tan(x)^5 - 5*tan(x)^3 + 15*tan(x))/a^(3/2) - x/a^(3/2)`

**Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a*cot(x)^4)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \int \frac{1}{(a \cot(x)^4)^{3/2}} dx$$

input `int(1/(a*cot(x)^4)^(3/2),x)`output `int(1/(a*cot(x)^4)^(3/2), x)`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.84

$$\int \frac{1}{(a \cot^4(x))^{3/2}} dx = \frac{\sqrt{a} (-15 \cos(x) \sin(x)^4 x + 30 \cos(x) \sin(x)^2 x - 15 \cos(x) x + 23 \sin(x)^5 - 35 \sin(x)^3)}{15 \cos(x) a^2 (\sin(x)^4 - 2 \sin(x)^2 + 1)}$$

input `int(1/(a*cot(x)^4)^(3/2),x)`output `(sqrt(a)*(-15*cos(x)*sin(x)**4*x + 30*cos(x)*sin(x)**2*x - 15*cos(x)*x + 23*sin(x)**5 - 35*sin(x)**3 + 15*sin(x)))/(15*cos(x)*a**2*(sin(x)**4 - 2*sin(x)**2 + 1))`

### 3.37 $\int (b \cot^p(c + dx))^n dx$

Optimal result	327
Mathematica [A] (verified)	327
Rubi [A] (verified)	328
Maple [F]	329
Fricas [F]	330
Sympy [F]	330
Maxima [F]	330
Giac [F]	331
Mupad [F(-1)]	331
Reduce [F]	331

#### Optimal result

Integrand size = 12, antiderivative size = 60

$$\int (b \cot^p(c + dx))^n dx = \frac{\cot(c + dx) (b \cot^p(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), -\cot^2(c + dx)\right)}{d(1 + np)}$$

output

```
-cot(d*x+c)*(b*cot(d*x+c)^p)^n*hypergeom([1, 1/2*n*p+1/2],[1/2*n*p+3/2],-cot(d*x+c)^2)/d/(n*p+1)
```

#### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00

$$\int (b \cot^p(c + dx))^n dx = \frac{\cot(c + dx) (b \cot^p(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), -\cot^2(c + dx)\right)}{d(1 + np)}$$

input

```
Integrate[(b*Cot[c + d*x]^p)^n,x]
```



output

$$-\left(\cot[c + dx] \cdot (b \cot[c + dx]^p)^n \operatorname{Hypergeometric2F1}\left[1, \frac{(1 + np)}{2}, \frac{(3 + np)}{2}, -\cot[c + dx]^2\right] / (d(1 + np))\right)$$
**Rubi [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {3042, 4142, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (b \cot^p(c + dx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \left(b \left(-\tan\left(c + dx + \frac{\pi}{2}\right)\right)^p\right)^n dx \\ & \quad \downarrow \text{4142} \\ & \cot^{-np}(c + dx) (b \cot^p(c + dx))^n \int \cot^{np}(c + dx) dx \\ & \quad \downarrow \text{3042} \\ & \cot^{-np}(c + dx) (b \cot^p(c + dx))^n \int \left(-\tan\left(c + dx + \frac{\pi}{2}\right)\right)^{np} dx \\ & \quad \downarrow \text{3957} \\ & \frac{\cot^{-np}(c + dx) (b \cot^p(c + dx))^n \int \frac{\cot^{np}(c + dx)}{\cot^2(c + dx) + 1} d \cot(c + dx)}{d} \\ & \quad \downarrow \text{278} \\ & \frac{\cot(c + dx) (b \cot^p(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(np + 1), \frac{1}{2}(np + 3), -\cot^2(c + dx)\right)}{d(np + 1)} \end{aligned}$$

input

$$\operatorname{Int}[(b \cot[c + dx]^p)^n, x]$$

output  $-\left(\cot[c + dx] \cdot (b \cot[c + dx]^p)^n \operatorname{Hypergeometric2F1}\left[1, (1 + np)/2, (3 + np)/2, -\cot[c + dx]^2\right]\right) / (d(1 + np))$

### Defintions of rubi rules used

rule 278  $\operatorname{Int}[(c \cdot x)^m \cdot (a + (b \cdot x^2)^p), x\_Symbol] \rightarrow \operatorname{Simp}[a^p \cdot (c \cdot x)^{m+1} / (c \cdot (m+1)) \cdot \operatorname{Hypergeometric2F1}[-p, (m+1)/2, (m+1)/2 + 1, (-b) \cdot (x^2/a)], x] /;$   $\operatorname{FreeQ}\{a, b, c, m, p\}, x \ \&\& \ !\operatorname{IGtQ}[p, 0] \ \&\& \ (\operatorname{ILtQ}[p, 0] \ || \ \operatorname{GtQ}[a, 0])$

rule 3042  $\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /;$   $\operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3957  $\operatorname{Int}[(b \cdot \tan[c] + (d \cdot x))^n], x\_Symbol] \rightarrow \operatorname{Simp}[b/d \ \operatorname{Subst}[\operatorname{Int}[x^n / (b^2 + x^2), x], x, b \cdot \tan[c + dx]], x] /;$   $\operatorname{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\operatorname{IntegerQ}[n]$

rule 4142  $\operatorname{Int}[(u \cdot (b \cdot (c \cdot \tan[e] + (f \cdot x))^n))^p], x\_Symbol] \rightarrow \operatorname{Simp}[b^{\operatorname{IntPart}[p]} \cdot (b \cdot (c \cdot \tan[e + f \cdot x])^n)^{\operatorname{FracPart}[p]} / (c \cdot \tan[e + f \cdot x])^{n \cdot \operatorname{FracPart}[p]}] \ \operatorname{Int}[\operatorname{ActivateTrig}[u] \cdot (c \cdot \tan[e + f \cdot x])^{n \cdot p}, x], x] /;$   $\operatorname{FreeQ}\{b, c, e, f, n, p\}, x \ \&\& \ !\operatorname{IntegerQ}[p] \ \&\& \ !\operatorname{IntegerQ}[n] \ \&\& \ (\operatorname{EqQ}[u, 1] \ || \ \operatorname{MatchQ}[u, ((d \cdot (\operatorname{trig}_)[e + f \cdot x])^m)]) /;$   $\operatorname{FreeQ}\{d, m\}, x \ \&\& \ \operatorname{MemberQ}\{\sin, \cos, \tan, \cot, \sec, \csc\}, \operatorname{trig}]$

### Maple [F]

$$\int (b \cot(dx + c)^p)^n dx$$

input  $\operatorname{int}((b \cdot \cot(dx + c)^p)^n, x)$

output  $\operatorname{int}((b \cdot \cot(dx + c)^p)^n, x)$

**Fricas [F]**

$$\int (b \cot^p(c + dx))^n dx = \int (b \cot(dx + c)^p)^n dx$$

input `integrate((b*cot(d*x+c)^p)^n,x, algorithm="fricas")`

output `integral((b*cot(d*x + c)^p)^n, x)`

**Sympy [F]**

$$\int (b \cot^p(c + dx))^n dx = \int (b \cot^p(c + dx))^n dx$$

input `integrate((b*cot(d*x+c)**p)**n,x)`

output `Integral((b*cot(c + d*x)**p)**n, x)`

**Maxima [F]**

$$\int (b \cot^p(c + dx))^n dx = \int (b \cot(dx + c)^p)^n dx$$

input `integrate((b*cot(d*x+c)^p)^n,x, algorithm="maxima")`

output `integrate((b*cot(d*x + c)^p)^n, x)`

**Giac [F]**

$$\int (b \cot^p(c + dx))^n dx = \int (b \cot(dx + c)^p)^n dx$$

input `integrate((b*cot(d*x+c)^p)^n,x, algorithm="giac")`

output `integrate((b*cot(d*x + c)^p)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \cot^p(c + dx))^n dx = \int (b \cot(c + dx)^p)^n dx$$

input `int((b*cot(c + d*x)^p)^n,x)`

output `int((b*cot(c + d*x)^p)^n, x)`

**Reduce [F]**

$$\int (b \cot^p(c + dx))^n dx = b^n \left( \int \cot(dx + c)^{np} dx \right)$$

input `int((b*cot(d*x+c)^p)^n,x)`

output `b**n*int(cot(c + d*x)**(n*p),x)`

### 3.38 $\int (a(b \cot(c + dx))^p)^n dx$

Optimal result	332
Mathematica [A] (verified)	332
Rubi [A] (verified)	333
Maple [F]	334
Fricas [F]	335
Sympy [F]	335
Maxima [F]	335
Giac [F]	336
Mupad [F(-1)]	336
Reduce [F]	336

#### Optimal result

Integrand size = 14, antiderivative size = 62

$$\int (a(b \cot(c + dx))^p)^n dx = \frac{\cot(c + dx) (a(b \cot(c + dx))^p)^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), -\cot^2(c + dx)\right)}{d(1 + np)}$$

output

```
-cot(d*x+c)*(a*(b*cot(d*x+c))^p)^n*hypergeom([1, 1/2*n*p+1/2], [1/2*n*p+3/2], -cot(d*x+c)^2)/d/(n*p+1)
```

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00

$$\int (a(b \cot(c + dx))^p)^n dx = \frac{\cot(c + dx) (a(b \cot(c + dx))^p)^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), -\cot^2(c + dx)\right)}{d(1 + np)}$$

input

```
Integrate[(a*(b*Cot[c + d*x])^p)^n,x]
```

output

$$-\left(\cot[c + dx] \cdot (a \cdot (b \cdot \cot[c + dx])^p)^n \cdot \text{Hypergeometric2F1}\left[1, \frac{(1 + np)}{2}, \frac{(3 + np)}{2}, -\cot[c + dx]^2\right] / (d \cdot (1 + np))\right)$$
**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$ , Rules used = {3042, 4142, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a(b \cot(c + dx))^p)^n dx \\ & \quad \downarrow \text{3042} \\ & \int \left(a\left(-b \tan\left(c + dx + \frac{\pi}{2}\right)\right)^p\right)^n dx \\ & \quad \downarrow \text{4142} \\ & (b \cot(c + dx))^{-np} (a(b \cot(c + dx))^p)^n \int (b \cot(c + dx))^{np} dx \\ & \quad \downarrow \text{3042} \\ & (b \cot(c + dx))^{-np} (a(b \cot(c + dx))^p)^n \int \left(-b \tan\left(c + dx + \frac{\pi}{2}\right)\right)^{np} dx \\ & \quad \downarrow \text{3957} \\ & \frac{b(b \cot(c + dx))^{-np} (a(b \cot(c + dx))^p)^n \int \frac{(b \cot(c + dx))^{np}}{\cot^2(c + dx) b^2 + b^2} d(b \cot(c + dx))}{d} \\ & \quad \downarrow \text{278} \\ & \frac{\cot(c + dx) \text{Hypergeometric2F1}\left(1, \frac{1}{2}(np + 1), \frac{1}{2}(np + 3), -\cot^2(c + dx)\right) (a(b \cot(c + dx))^p)^n}{d(np + 1)} \end{aligned}$$

input

$$\text{Int}[(a \cdot (b \cdot \cot[c + dx])^p)^n, x]$$

output  $-\left(\cot[c + dx] \cdot (a \cdot (b \cdot \cot[c + dx])^p)^n \cdot \text{Hypergeometric2F1}\left[1, (1 + np)/2, (3 + np)/2, -\cot[c + dx]^2\right]\right) / (d \cdot (1 + np))$

### Defintions of rubi rules used

rule 278  $\text{Int}[(c \cdot x)^m \cdot (a + (b \cdot x^2)^p), x\_Symbol] \rightarrow \text{Simp}[a^p \cdot (c \cdot x)^{m+1} / (c \cdot (m+1)) \cdot \text{Hypergeometric2F1}[-p, (m+1)/2, (m+1)/2 + 1, (-b) \cdot (x^2/a)], x] /;$   $\text{FreeQ}\{a, b, c, m, p\}, x \} \&\& \text{!IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] \mid \mid \text{GtQ}[a, 0])$

rule 3042  $\text{Int}[u, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$   $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3957  $\text{Int}[(b \cdot \tan[c] + d \cdot x)^n, x\_Symbol] \rightarrow \text{Simp}[b/d \text{ Subst}[\text{Int}[x^n / (b^2 + x^2), x], x, b \cdot \tan[c + dx]], x] /;$   $\text{FreeQ}\{b, c, d, n\}, x \} \&\& \text{!IntegerQ}[n]$

rule 4142  $\text{Int}[(u \cdot (b \cdot (c \cdot \tan[e] + f \cdot x))^n)^p, x\_Symbol] \rightarrow \text{Simp}[b^{\text{IntPart}[p]} \cdot (b \cdot (c \cdot \tan[e + f \cdot x])^n)^{\text{FracPart}[p]} / (c \cdot \tan[e + f \cdot x])^{n \cdot \text{FracPart}[p]}] \cdot \text{Int}[\text{ActivateTrig}[u] \cdot (c \cdot \tan[e + f \cdot x])^{n \cdot p}, x], x] /;$   $\text{FreeQ}\{b, c, e, f, n, p\}, x \} \&\& \text{!IntegerQ}[p] \&\& \text{!IntegerQ}[n] \&\& (\text{EqQ}[u, 1] \mid \mid \text{MatchQ}[u, ((d \cdot (\text{trig}_1)[e + f \cdot x])^m]) /;$   $\text{FreeQ}\{d, m\}, x \} \&\& \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}_1\}$

### Maple [F]

$$\int (a(b \cot(dx + c))^p)^n dx$$

input  $\text{int}((a \cdot (b \cdot \cot(dx + c))^p)^n, x)$

output  $\text{int}((a \cdot (b \cdot \cot(dx + c))^p)^n, x)$

**Fricas [F]**

$$\int (a(b \cot(c + dx))^p)^n dx = \int ((b \cot(dx + c))^p a)^n dx$$

input `integrate((a*(b*cot(d*x+c))^p)^n,x, algorithm="fricas")`

output `integral(((b*cot(d*x + c))^p*a)^n, x)`

**Sympy [F]**

$$\int (a(b \cot(c + dx))^p)^n dx = \int (a(b \cot(c + dx))^p)^n dx$$

input `integrate((a*(b*cot(d*x+c))**p)**n,x)`

output `Integral((a*(b*cot(c + d*x))**p)**n, x)`

**Maxima [F]**

$$\int (a(b \cot(c + dx))^p)^n dx = \int ((b \cot(dx + c))^p a)^n dx$$

input `integrate((a*(b*cot(d*x+c))^p)^n,x, algorithm="maxima")`

output `integrate(((b*cot(d*x + c))^p*a)^n, x)`



**Giac [F]**

$$\int (a(b \cot(c + dx))^p)^n dx = \int ((b \cot(dx + c))^p a)^n dx$$

input `integrate((a*(b*cot(d*x+c))^p)^n,x, algorithm="giac")`

output `integrate(((b*cot(d*x + c))^p*a)^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a(b \cot(c + dx))^p)^n dx = \int (a(b \cot(c + dx))^p)^n dx$$

input `int((a*(b*cot(c + d*x))^p)^n,x)`

output `int((a*(b*cot(c + d*x))^p)^n, x)`

**Reduce [F]**

$$\int (a(b \cot(c + dx))^p)^n dx = b^{np} a^n \left( \int \cot(dx + c)^{np} dx \right)$$

input `int((a*(b*cot(d*x+c))^p)^n,x)`

output `b**(n*p)*a**n*int(cot(c + d*x)**(n*p),x)`

### 3.39 $\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx$

Optimal result	337
Mathematica [C] (warning: unable to verify)	337
Rubi [A] (verified)	338
Maple [F]	339
Fricas [F]	340
Sympy [F]	340
Maxima [F]	340
Giac [F]	341
Mupad [F(-1)]	341
Reduce [F]	341

#### Optimal result

Integrand size = 21, antiderivative size = 87

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \frac{(b \cot(e + fx))^{1+n} \text{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{2}(1 - m + n), \frac{3+n}{2}, \cos^2(e + fx)\right) (a \sin(e + fx))^m \sin^2(e + fx)}{bf(1+n)}$$

output

```
-(b*cot(f*x+e))^(1+n)*hypergeom([1/2+1/2*n, 1/2-1/2*m+1/2*n], [3/2+1/2*n], cos(f*x+e)^2)*(a*sin(f*x+e))^m*(sin(f*x+e)^2)^(1/2-1/2*m+1/2*n)/b/f/(1+n)
```

#### Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.81 (sec) , antiderivative size = 289, normalized size of antiderivative = 3.32

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \frac{(3+m)}{f(1+m-n)} \left( (3+m-n) \text{AppellF1}\left(\frac{1}{2}(1+m-n), -n, 1+m, \frac{1}{2}(3+m-n), \tan^2\left(\frac{1}{2}(e+fx)\right)\right), -\tan\left(\frac{1}{2}(e+fx)\right) \right)$$

input

```
Integrate[(b*Cot[e + f*x])^n*(a*Sin[e + f*x])^m,x]
```

output

```
((3 + m - n)*AppellF1[(1 + m - n)/2, -n, 1 + m, (3 + m - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(b*Cot[e + f*x])^n*Sin[e + f*x]*(a*Sin[e + f*x])^m)/(f*(1 + m - n)*((3 + m - n)*AppellF1[(1 + m - n)/2, -n, 1 + m, (3 + m - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*(n*AppellF1[(3 + m - n)/2, 1 - n, 1 + m, (5 + m - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + (1 + m)*AppellF1[(3 + m - n)/2, -n, 2 + m, (5 + m - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))
```

**Rubi [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {3042, 3083, 3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a \sin(e + fx))^m (b \cot(e + fx))^n dx$$

$$\downarrow 3042$$

$$\int \left( a \cos \left( e + fx - \frac{\pi}{2} \right) \right)^m \left( -b \tan \left( e + fx - \frac{\pi}{2} \right) \right)^n dx$$

$$\downarrow 3083$$

$$(a \sin(e + fx))^m \left( \frac{\csc(e + fx)}{a} \right)^m \int (b \cot(e + fx))^n \left( \frac{\csc(e + fx)}{a} \right)^{-m} dx$$

$$\downarrow 3042$$

$$(a \sin(e + fx))^m \left( \frac{\csc(e + fx)}{a} \right)^m \int \left( \frac{\sec \left( e + fx - \frac{\pi}{2} \right)}{a} \right)^{-m} \left( -b \tan \left( e + fx - \frac{\pi}{2} \right) \right)^n dx$$

$$\downarrow 3097$$

$$\frac{(a \sin(e + fx))^m (b \cot(e + fx))^{n+1} \sin^2(e + fx)^{\frac{1}{2}(-m+n+1)} \text{Hypergeometric2F1} \left( \frac{n+1}{2}, \frac{1}{2}(-m+n+1), \frac{n+3}{2}, \cos^2(e + fx) \right)}{bf(n+1)}$$

input

```
Int[(b*Cot[e + f*x])^n*(a*Sin[e + f*x])^m,x]
```

output  $-\left(\left(b \cot [e+f x]\right)^{(1+n)} \operatorname{Hypergeometric2F1}\left[\frac{(1+n)}{2}, \frac{(1-m+n)}{2}, \frac{(3+n)}{2}, \cos [e+f x]^2\right] \left(a \sin [e+f x]\right)^m \left(\sin [e+f x]^2\right)^{\frac{(1-m+n)}{2}}\right) / \left(b f(1+n)\right)$

### Defintions of rubi rules used

rule 3042  $\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] / ; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3083  $\operatorname{Int}[(\cos [(e_{.})+(f_{.})(x_{.})] \cdot (a_{.}))^{(m_{.})} \cdot ((b_{.}) \cdot \tan [(e_{.})+(f_{.})(x_{.})])^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(a \cdot \cos [e+f x])^{\operatorname{FracPart}[m]} \cdot (\sec [e+f x] / a)^{\operatorname{FracPart}[m]} \operatorname{Int}[(b \cdot \tan [e+f x])^n / (\sec [e+f x] / a)^m, x], x] / ; \operatorname{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n]$

rule 3097  $\operatorname{Int}[(a_{.}) \cdot \sec [(e_{.})+(f_{.})(x_{.})])^{(m_{.})} \cdot ((b_{.}) \cdot \tan [(e_{.})+(f_{.})(x_{.})])^{(n_{.})}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(a \cdot \sec [e+f x])^m \cdot (b \cdot \tan [e+f x])^{(n+1)} \cdot (\cos [e+f x]^2)^{\frac{(m+n+1)}{2}} / (b f(n+1))] \cdot \operatorname{Hypergeometric2F1}\left[\frac{(n+1)}{2}, \frac{(m+n+1)}{2}, \frac{(n+3)}{2}, \sin [e+f x]^2\right], x] / ; \operatorname{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& !\operatorname{IntegerQ}[(n-1) / 2] \&\& !\operatorname{IntegerQ}[m / 2]$

### Maple **[F]**

$$\int (b \cot (f x+e))^n (a \sin (f x+e))^m dx$$

input  $\operatorname{int}((b \cdot \cot (f \cdot x+e))^n \cdot (a \cdot \sin (f \cdot x+e))^m, x)$

output  $\operatorname{int}((b \cdot \cot (f \cdot x+e))^n \cdot (a \cdot \sin (f \cdot x+e))^m, x)$

**Fricas [F]**

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \int (b \cot(fx + e))^n (a \sin(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="fricas")`

output `integral((b*cot(f*x + e))^n*(a*sin(f*x + e))^m, x)`

**Sympy [F]**

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \int (a \sin(e + fx))^m (b \cot(e + fx))^n dx$$

input `integrate((b*cot(f*x+e))**n*(a*sin(f*x+e))**m,x)`

output `Integral((a*sin(e + f*x))**m*(b*cot(e + f*x))**n, x)`

**Maxima [F]**

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \int (b \cot(fx + e))^n (a \sin(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="maxima")`

output `integrate((b*cot(f*x + e))^n*(a*sin(f*x + e))^m, x)`

**Giac [F]**

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \int (b \cot(fx + e))^n (a \sin(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="giac")`

output `integrate((b*cot(f*x + e))^n*(a*sin(f*x + e))^m, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = \int (b \cot(e + fx))^n (a \sin(e + fx))^m dx$$

input `int((b*cot(e + f*x))^n*(a*sin(e + f*x))^m,x)`

output `int((b*cot(e + f*x))^n*(a*sin(e + f*x))^m, x)`

**Reduce [F]**

$$\int (b \cot(e + fx))^n (a \sin(e + fx))^m dx = b^n a^m \left( \int \sin(fx + e)^m \cot(fx + e)^n dx \right)$$

input `int((b*cot(f*x+e))^n*(a*sin(f*x+e))^m,x)`

output `b**n*a**m*int(sin(e + f*x)**m*cot(e + f*x)**n,x)`

### 3.40 $\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx$

Optimal result	342
Mathematica [A] (verified)	342
Rubi [A] (verified)	343
Maple [F]	344
Fricas [F]	345
Sympy [F]	345
Maxima [F]	345
Giac [F]	346
Mupad [F(-1)]	346
Reduce [F]	346

#### Optimal result

Integrand size = 21, antiderivative size = 84

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \frac{(a \cos(e + fx))^m (b \cot(e + fx))^{1+n} \operatorname{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{2}(1+m+n), \frac{1}{2}(3+m+n), \cos^2(e + fx)\right)}{bf(1+m+n)}$$

output

```
-(a*cos(f*x+e))^m*(b*cot(f*x+e))^(1+n)*hypergeom([1/2+1/2*n, 1/2+1/2*m+1/2*n], [3/2+1/2*m+1/2*n], cos(f*x+e)^2)*(sin(f*x+e)^2)^(1/2+1/2*n)/b/f/(1+m+n)
```

#### Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.99

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \frac{b(a \cos(e + fx))^m (b \cot(e + fx))^{-1+n} \operatorname{Hypergeometric2F1}\left(\frac{2+m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, -\tan^2(e + fx)\right) \sec^2(e + fx)}{f(-1+n)}$$

input

```
Integrate[(a*Cos[e + f*x])^m*(b*Cot[e + f*x])^n,x]
```

output

```

-((b*(a*cos[e + f*x])^m*(b*cot[e + f*x])^(-1 + n)*Hypergeometric2F1[(2 + m
)/2, (1 - n)/2, (3 - n)/2, -Tan[e + f*x]^2]*(Sec[e + f*x]^2)^(m/2))/(f*(-1
+ n))

```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {3042, 3082, 3042, 3056}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cos(e + fx))^m (b \cot(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sin \left( e + fx + \frac{\pi}{2} \right) \right)^m \left( -b \tan \left( e + fx + \frac{\pi}{2} \right) \right)^n dx \\
 & \quad \downarrow \text{3082} \\
 & \frac{a(-\sin(e + fx))^{n+1} (a \cos(e + fx))^{-n-1} (b \cot(e + fx))^{n+1} \int (a \cos(e + fx))^{m+n} (-\sin(e + fx))^{-n} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(-\sin(e + fx))^{n+1} (a \cos(e + fx))^{-n-1} (b \cot(e + fx))^{n+1} \int (a \cos(e + fx))^{m+n} (-\sin(e + fx))^{-n} dx}{b} \\
 & \quad \downarrow \text{3056} \\
 & \frac{\sin^2(e + fx)^{\frac{n+1}{2}} (a \cos(e + fx))^m (b \cot(e + fx))^{n+1} \text{Hypergeometric2F1} \left( \frac{n+1}{2}, \frac{1}{2}(m+n+1), \frac{1}{2}(m+n+3), \cot^2(e + fx) \right)}{bf(m+n+1)}
 \end{aligned}$$

input

```

Int[(a*cos[e + f*x])^m*(b*cot[e + f*x])^n,x]

```



output  $-\left(\left(a \cos [e+f x]\right)^m\left(b \cot [e+f x]\right)^{(1+n)} \operatorname{Hypergeometric2F1}\left[\frac{(1+n)}{2}, \frac{(1+m+n)}{2}, \frac{(3+m+n)}{2}, \cos [e+f x]^2\right] \frac{\left(\sin [e+f x]^2\right)^{\frac{(1+n)}{2}}}{\left(b f(1+m+n)\right)}\right)$

### Definitions of rubi rules used

rule 3042  $\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] / ; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3056  $\operatorname{Int}[(\cos [(e.)+(f.)*(x_*)]*(a.))^{(m)}*((b.)*\sin [(e.)+(f.)*(x_*)])^{(n)}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-b^{(2*\operatorname{IntPart}[(n-1)/2]+1)}*(b*\sin [e+f x])^{(2*\operatorname{FracPart}[(n-1)/2]}*((a*\cos [e+f x])^{(m+1)} / (a*f*(m+1)*(\sin [e+f x]^2)^{\operatorname{FracPart}[(n-1)/2]})))*\operatorname{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \cos [e+f x]^2], x] / ; \operatorname{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \operatorname{SimplerQ}[n, m]$

rule 3082  $\operatorname{Int}[(a.*\sin [(e.)+(f.)*(x_*)])^{(m)}*((b.)*\tan [(e.)+(f.)*(x_*)])^{(n)}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[a*\cos [e+f x]^{(n+1)}*((b*\tan [e+f x])^{(n+1)} / (b*(a*\sin [e+f x])^{(n+1)})) \operatorname{Int}[(a*\sin [e+f x])^{(m+n)} / \cos [e+f x]^n, x], x] / ; \operatorname{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& !\operatorname{IntegerQ}[n]$

### Maple **[F]**

$$\int (a \cos (f x + e))^m (b \cot (f x + e))^n dx$$

input  $\operatorname{int}((a*\cos(f*x+e))^m*(b*\cot(f*x+e))^n,x)$

output  $\operatorname{int}((a*\cos(f*x+e))^m*(b*\cot(f*x+e))^n,x)$

**Fricas [F]**

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cos(fx + e))^m (b \cot(fx + e))^n dx$$

input `integrate((a*cos(f*x+e))^m*(b*cot(f*x+e))^n,x, algorithm="fricas")`

output `integral((a*cos(f*x + e))^m*(b*cot(f*x + e))^n, x)`

**Sympy [F]**

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cos(e + fx))^m (b \cot(e + fx))^n dx$$

input `integrate((a*cos(f*x+e))**m*(b*cot(f*x+e))**n,x)`

output `Integral((a*cos(e + f*x))**m*(b*cot(e + f*x))**n, x)`

**Maxima [F]**

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cos(fx + e))^m (b \cot(fx + e))^n dx$$

input `integrate((a*cos(f*x+e))^m*(b*cot(f*x+e))^n,x, algorithm="maxima")`

output `integrate((a*cos(f*x + e))^m*(b*cot(f*x + e))^n, x)`

**Giac [F]**

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cos(fx + e))^m (b \cot(fx + e))^n dx$$

input `integrate((a*cos(f*x+e))^m*(b*cot(f*x+e))^n,x, algorithm="giac")`

output `integrate((a*cos(f*x + e))^m*(b*cot(f*x + e))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cos(e + fx))^m (b \cot(e + fx))^n dx$$

input `int((a*cos(e + f*x))^m*(b*cot(e + f*x))^n,x)`

output `int((a*cos(e + f*x))^m*(b*cot(e + f*x))^n, x)`

**Reduce [F]**

$$\int (a \cos(e + fx))^m (b \cot(e + fx))^n dx = b^n a^m \left( \int \cot(fx + e)^n \cos(fx + e)^m dx \right)$$

input `int((a*cos(f*x+e))^m*(b*cot(f*x+e))^n,x)`

output `b**n*a**m*int(cot(e + f*x)**n*cos(e + f*x)**m,x)`

### 3.41 $\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx$

Optimal result	347
Mathematica [A] (verified)	347
Rubi [A] (verified)	348
Maple [F]	349
Fricas [F]	350
Sympy [F]	350
Maxima [F]	350
Giac [F]	351
Mupad [F(-1)]	351
Reduce [F]	351

#### Optimal result

Integrand size = 21, antiderivative size = 64

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \frac{(a \cot(e + fx))^{1+m} (b \cot(e + fx))^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + m + n), \frac{1}{2}(3 + m + n), -\cot^2(e + fx)\right)}{af(1 + m + n)}$$

output

```
-(a*cot(f*x+e))^(1+m)*(b*cot(f*x+e))^n*hypergeom([1, 1/2+1/2*m+1/2*n], [3/2+1/2*m+1/2*n], -cot(f*x+e)^2)/a/f/(1+m+n)
```

#### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.02

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \frac{\cot(e + fx) (a \cot(e + fx))^m (b \cot(e + fx))^n \operatorname{Hypergeometric2F1}\left(1, \frac{1}{2}(1 + m + n), \frac{1}{2}(3 + m + n), -\cot^2(e + fx)\right)}{f(1 + m + n)}$$

input

```
Integrate[(a*Cot[e + f*x])^m*(b*Cot[e + f*x])^n,x]
```

output

```
-((Cot[e + f*x]*(a*Cot[e + f*x])^m*(b*Cot[e + f*x])^n*Hypergeometric2F1[1,
(1 + m + n)/2, (3 + m + n)/2, -Cot[e + f*x]^2])/(f*(1 + m + n)))
```

**Rubi [A] (verified)**

Time = 0.28 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {2034, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \cot(e + fx))^m (b \cot(e + fx))^n dx \\
 & \quad \downarrow \text{2034} \\
 & (a \cot(e + fx))^{-n} (b \cot(e + fx))^n \int (a \cot(e + fx))^{m+n} dx \\
 & \quad \downarrow \text{3042} \\
 & (a \cot(e + fx))^{-n} (b \cot(e + fx))^n \int \left(-a \tan\left(e + fx + \frac{\pi}{2}\right)\right)^{m+n} dx \\
 & \quad \downarrow \text{3957} \\
 & \frac{a(a \cot(e + fx))^{-n} (b \cot(e + fx))^n \int \frac{(a \cot(e + fx))^{m+n}}{\cot^2(e + fx) a^2 + a^2} d(a \cot(e + fx))}{f} \\
 & \quad \downarrow \text{278} \\
 & \frac{(a \cot(e + fx))^{m+1} (b \cot(e + fx))^n \text{Hypergeometric2F1}\left(1, \frac{1}{2}(m + n + 1), \frac{1}{2}(m + n + 3), -\cot^2(e + fx)\right)}{af(m + n + 1)}
 \end{aligned}$$

input

```
Int[(a*Cot[e + f*x])^m*(b*Cot[e + f*x])^n,x]
```

output

```
-(((a*Cot[e + f*x])^(1 + m)*(b*Cot[e + f*x])^n*Hypergeometric2F1[1, (1 + m
+ n)/2, (3 + m + n)/2, -Cot[e + f*x]^2])/(a*f*(1 + m + n)))
```

## Definitions of rubi rules used

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2034 `Int[(F*x_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Simp[b^IntPart[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])) Int[(a*v)^(m + n)*Fx, x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

Maple **[F]**

$$\int (a \cot (fx + e))^m (b \cot (fx + e))^n dx$$

input `int((a*cot(f*x+e))^m*(b*cot(f*x+e))^n,x)`

output `int((a*cot(f*x+e))^m*(b*cot(f*x+e))^n,x)`

**Fricas [F]**

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cot(fx + e))^m (b \cot(fx + e))^n dx$$

input `integrate((a*cot(f*x+e))^m*(b*cot(f*x+e))^n,x, algorithm="fricas")`

output `integral((a*cot(f*x + e))^m*(b*cot(f*x + e))^n, x)`

**Sympy [F]**

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cot(e + fx))^m (b \cot(e + fx))^n dx$$

input `integrate((a*cot(f*x+e))**m*(b*cot(f*x+e))**n,x)`

output `Integral((a*cot(e + f*x))**m*(b*cot(e + f*x))**n, x)`

**Maxima [F]**

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cot(fx + e))^m (b \cot(fx + e))^n dx$$

input `integrate((a*cot(f*x+e))^m*(b*cot(f*x+e))^n,x, algorithm="maxima")`

output `integrate((a*cot(f*x + e))^m*(b*cot(f*x + e))^n, x)`

**Giac [F]**

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cot(fx + e))^m (b \cot(fx + e))^n dx$$

input `integrate((a*cot(f*x+e))^m*(b*cot(f*x+e))^n,x, algorithm="giac")`

output `integrate((a*cot(f*x + e))^m*(b*cot(f*x + e))^n, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = \int (a \cot(e + fx))^m (b \cot(e + fx))^n dx$$

input `int((a*cot(e + f*x))^m*(b*cot(e + f*x))^n,x)`

output `int((a*cot(e + f*x))^m*(b*cot(e + f*x))^n, x)`

**Reduce [F]**

$$\int (a \cot(e + fx))^m (b \cot(e + fx))^n dx = b^n a^m \left( \int \cot(fx + e)^{m+n} dx \right)$$

input `int((a*cot(f*x+e))^m*(b*cot(f*x+e))^n,x)`

output `b**n*a**m*int(cot(e + f*x)**(m + n),x)`



### 3.42 $\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx$

Optimal result	352
Mathematica [A] (verified)	352
Rubi [A] (verified)	353
Maple [F]	355
Fricas [F]	355
Sympy [F]	355
Maxima [F]	356
Giac [F]	356
Mupad [F(-1)]	356
Reduce [F]	357

#### Optimal result

Integrand size = 21, antiderivative size = 90

$$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx = \frac{(b \cot(e + fx))^{1+n} \text{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{2}(1-m+n), \frac{1}{2}(3-m+n), \cos^2(e + fx)\right) (a \sec(e + fx))^m}{bf(1-m+n)}$$

output

```
-(b*cot(f*x+e))^(1+n)*hypergeom([1/2+1/2*n, 1/2-1/2*m+1/2*n], [3/2-1/2*m+1/2*n], cos(f*x+e)^2)*(a*sec(f*x+e))^m*(sin(f*x+e)^2)^(1/2+1/2*n)/b/f/(1-m+n)
```

#### Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.92

$$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx = \frac{b(b \cot(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(1 - \frac{m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, -\tan^2(e + fx)\right) (a \sec(e + fx))^m \sec^2(e + fx)}{f(-1+n)}$$

input

```
Integrate[(b*Cot[e + f*x])^n*(a*Sec[e + f*x])^m,x]
```

output

```

-((b*(b*Cot[e + f*x])^(-1 + n)*Hypergeometric2F1[1 - m/2, (1 - n)/2, (3 -
n)/2, -Tan[e + f*x]^2]*(a*Sec[e + f*x])^m)/(f*(-1 + n)*(Sec[e + f*x]^2)^(m
/2)))

```

**Rubi [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {3042, 3098, 3042, 3082, 3042, 3056}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \sec(e + fx))^m (b \cot(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \csc\left(e + fx + \frac{\pi}{2}\right) \right)^m \left( -b \tan\left(e + fx + \frac{\pi}{2}\right) \right)^n dx \\
 & \quad \downarrow \text{3098} \\
 & \left( \frac{\cos(e + fx)}{a} \right)^m (a \sec(e + fx))^m \int \left( \frac{\cos(e + fx)}{a} \right)^{-m} (b \cot(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \left( \frac{\cos(e + fx)}{a} \right)^m (a \sec(e + fx))^m \int \left( \frac{\sin\left(e + fx + \frac{\pi}{2}\right)}{a} \right)^{-m} \left( -b \tan\left(e + fx + \frac{\pi}{2}\right) \right)^n dx \\
 & \quad \downarrow \text{3082} \\
 & \frac{(-\sin(e + fx))^{n+1} (a \sec(e + fx))^m (b \cot(e + fx))^{n+1} \left( \frac{\cos(e+fx)}{a} \right)^{m-n-1} \int \left( \frac{\cos(e+fx)}{a} \right)^{n-m} (-\sin(e + fx))^{-n} dx}{ab} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-\sin(e + fx))^{n+1} (a \sec(e + fx))^m (b \cot(e + fx))^{n+1} \left( \frac{\cos(e+fx)}{a} \right)^{m-n-1} \int \left( \frac{\cos(e+fx)}{a} \right)^{n-m} (-\sin(e + fx))^{-n} dx}{ab} \\
 & \quad \downarrow \text{3056}
 \end{aligned}$$

$$\frac{\sin^2(e + fx)^{\frac{n+1}{2}} (a \sec(e + fx))^m (b \cot(e + fx))^{n+1} \text{Hypergeometric2F1}\left(\frac{n+1}{2}, \frac{1}{2}(-m + n + 1), \frac{1}{2}(-m + n + 3)\right)}{bf(-m + n + 1)}$$

input `Int[(b*Cot[e + f*x])^n*(a*Sec[e + f*x])^m,x]`

output `-(((b*Cot[e + f*x])^(1 + n)*Hypergeometric2F1[(1 + n)/2, (1 - m + n)/2, (3 - m + n)/2, Cos[e + f*x]^2]*(a*Sec[e + f*x])^m*(Sin[e + f*x]^2)^((1 + n)/2))/(b*f*(1 - m + n))`

### Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3056 `Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^m_*((b_.)*sin[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] := Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sine[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]`

rule 3082 `Int[((a_.)*sin[(e_.) + (f_.)*(x_.)])^m_*((b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] := Simp[a*Cos[e + f*x]^(n + 1)*((b*Tan[e + f*x])^(n + 1)/(b*(a*Sine[e + f*x])^(n + 1))) Int[(a*Sine[e + f*x])^(m + n)/Cos[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n]`

rule 3098 `Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m_*((b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] := Simp[(a*Csc[e + f*x])^FracPart[m]*(Sin[e + f*x]/a)^FracPart[m] Int[(b*Tan[e + f*x])^n/(Sin[e + f*x]/a)^m, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]`

**Maple [F]**

$$\int (b \cot (fx + e))^n (a \sec (fx + e))^m dx$$

input `int((b*cot(f*x+e))^n*(a*sec(f*x+e))^m,x)`

output `int((b*cot(f*x+e))^n*(a*sec(f*x+e))^m,x)`

**Fricas [F]**

$$\int (b \cot (e + fx))^n (a \sec (e + fx))^m dx = \int (b \cot (fx + e))^n (a \sec (fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="fricas")`

output `integral((b*cot(f*x + e))^n*(a*sec(f*x + e))^m, x)`

**Sympy [F]**

$$\int (b \cot (e + fx))^n (a \sec (e + fx))^m dx = \int (a \sec (e + fx))^m (b \cot (e + fx))^n dx$$

input `integrate((b*cot(f*x+e))**n*(a*sec(f*x+e))**m,x)`

output `Integral((a*sec(e + f*x))**m*(b*cot(e + f*x))**n, x)`

**Maxima [F]**

$$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx = \int (b \cot(fx + e))^n (a \sec(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="maxima")`

output `integrate((b*cot(f*x + e))^n*(a*sec(f*x + e))^m, x)`

**Giac [F]**

$$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx = \int (b \cot(fx + e))^n (a \sec(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="giac")`

output `integrate((b*cot(f*x + e))^n*(a*sec(f*x + e))^m, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx = \int (b \cot(e + fx))^n \left( \frac{a}{\cos(e + fx)} \right)^m dx$$

input `int((b*cot(e + f*x))^n*(a/cos(e + f*x))^m,x)`

output `int((b*cot(e + f*x))^n*(a/cos(e + f*x))^m, x)`

**Reduce [F]**

$$\int (b \cot(e + fx))^n (a \sec(e + fx))^m dx = b^n a^m \left( \int \sec(fx + e)^m \cot(fx + e)^n dx \right)$$

input `int((b*cot(f*x+e))^n*(a*sec(f*x+e))^m,x)`

output `b**n*a**m*int(sec(e + f*x)**m*cot(e + f*x)**n,x)`

### 3.43 $\int (d \cot(e + fx))^n \csc^6(e + fx) dx$

Optimal result	358
Mathematica [A] (verified)	358
Rubi [A] (verified)	359
Maple [A] (verified)	360
Fricas [A] (verification not implemented)	361
Sympy [F(-1)]	361
Maxima [A] (verification not implemented)	361
Giac [F]	362
Mupad [B] (verification not implemented)	362
Reduce [F]	363

#### Optimal result

Integrand size = 19, antiderivative size = 76

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = -\frac{(d \cot(e + fx))^{1+n}}{df(1+n)} - \frac{2(d \cot(e + fx))^{3+n}}{d^3 f(3+n)} - \frac{(d \cot(e + fx))^{5+n}}{d^5 f(5+n)}$$

output

```
-(d*cot(f*x+e))^(1+n)/d/f/(1+n)-2*(d*cot(f*x+e))^(3+n)/d^3/f/(3+n)-(d*cot(f*x+e))^(5+n)/d^5/f/(5+n)
```

#### Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = \frac{(8 + 6n + n^2 - 2(3 + n) \cos(2(e + fx)) + \cos(4(e + fx))) \cot(e + fx) (d \cot(e + fx))^n \csc^4(e + fx)}{f(1+n)(3+n)(5+n)}$$

input

```
Integrate[(d*Cot[e + f*x])^n*Csc[e + f*x]^6,x]
```

output

$$-\left(\left(8 + 6n + n^2 - 2(3 + n)\cos[2(e + fx)] + \cos[4(e + fx)]\right)\cot[e + fx] \cdot (d \cot[e + fx])^n \operatorname{Csc}[e + fx]^4 / (f(1 + n)(3 + n)(5 + n))\right)$$

**Rubi [A] (verified)**

Time = 0.27 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3087, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{csc}^6(e + fx)(d \cot(e + fx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \sec\left(e + fx - \frac{\pi}{2}\right)^6 \left(-d \tan\left(e + fx - \frac{\pi}{2}\right)\right)^n dx \\ & \quad \downarrow \text{3087} \\ & \frac{\int (d \cot(e + fx))^n (\cot^2(e + fx) + 1)^2 d(-\cot(e + fx))}{f} \\ & \quad \downarrow \text{244} \\ & \frac{\int \left( (d \cot(e + fx))^n + \frac{2(d \cot(e + fx))^{n+2}}{d^2} + \frac{(d \cot(e + fx))^{n+4}}{d^4} \right) d(-\cot(e + fx))}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{-\frac{(d \cot(e + fx))^{n+5}}{d^5(n+5)} - \frac{2(d \cot(e + fx))^{n+3}}{d^3(n+3)} - \frac{(d \cot(e + fx))^{n+1}}{d(n+1)}}{f} \end{aligned}$$

input

$$\operatorname{Int}[(d \cot[e + fx])^n \operatorname{Csc}[e + fx]^6, x]$$

output

$$\left(-\frac{(d \cot[e + fx])^{1+n}}{d(1+n)} - \frac{2(d \cot[e + fx])^{3+n}}{d^3(3+n)} - \frac{(d \cot[e + fx])^{5+n}}{d^5(5+n)}\right) / f$$



## Definitions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

## Maple [A] (verified)

Time = 29.43 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.18

method	result	size
derivativedivides	$-\frac{\cot(fx+e)e^{n \ln(d \cot(fx+e))}}{f(1+n)} - \frac{2 \cot(fx+e)^3 e^{n \ln(d \cot(fx+e))}}{f(3+n)} - \frac{\cot(fx+e)^5 e^{n \ln(d \cot(fx+e))}}{f(5+n)}$	90
default	$-\frac{\cot(fx+e)e^{n \ln(d \cot(fx+e))}}{f(1+n)} - \frac{2 \cot(fx+e)^3 e^{n \ln(d \cot(fx+e))}}{f(3+n)} - \frac{\cot(fx+e)^5 e^{n \ln(d \cot(fx+e))}}{f(5+n)}$	90
risch	Expression too large to display	10532

input `int((d*cot(f*x+e))^n*csc(f*x+e)^6,x,method=_RETURNVERBOSE)`

output `-1/f/(1+n)*cot(f*x+e)*exp(n*ln(d*cot(f*x+e)))-2/f/(3+n)*cot(f*x+e)^3*exp(n*ln(d*cot(f*x+e)))-1/f/(5+n)*cot(f*x+e)^5*exp(n*ln(d*cot(f*x+e)))`

**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.91

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = \frac{(8 \cos(fx + e))^5 - 4(n + 5) \cos(fx + e)^3 + (n^2 + 8n + 15) \cos(fx + e) \left(\frac{d}{s}\right)}{((fn^3 + 9fn^2 + 23fn + 15f) \cos(fx + e)^4 + fn^3 + 9fn^2 - 2(fn^3 + 9fn^2 + 23fn + 15f) \cos(fx + e)^2 + 23fn + 15f) \sin(fx + e)}$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^6,x, algorithm="fricas")`

output `-(8*cos(f*x + e)^5 - 4*(n + 5)*cos(f*x + e)^3 + (n^2 + 8*n + 15)*cos(f*x + e))*(d*cos(f*x + e)/sin(f*x + e))^n/(((f*n^3 + 9*f*n^2 + 23*f*n + 15*f)*cos(f*x + e)^4 + f*n^3 + 9*f*n^2 - 2*(f*n^3 + 9*f*n^2 + 23*f*n + 15*f)*cos(f*x + e)^2 + 23*f*n + 15*f)*sin(f*x + e))`

**Sympy [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = \text{Timed out}$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)**6,x)`

output `Timed out`

**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.11

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = -\frac{\left(\frac{d}{\tan(fx+e)}\right)^{n+1}}{d(n+1)} + \frac{2d^n \tan(fx+e)^{-n}}{(n+3) \tan(fx+e)^3} + \frac{d^n \tan(fx+e)^{-n}}{(n+5) \tan(fx+e)^5} f$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^6,x, algorithm="maxima")`

output

```
-((d/tan(f*x + e))^(n + 1)/(d*(n + 1)) + 2*d^n*tan(f*x + e)^(-n)/((n + 3)*
tan(f*x + e)^3) + d^n*tan(f*x + e)^(-n)/((n + 5)*tan(f*x + e)^5))/f
```

**Giac [F]**

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e)^6 dx$$

input

```
integrate((d*cot(f*x+e))^n*csc(f*x+e)^6,x, algorithm="giac")
```

output

```
integrate((d*cot(f*x + e))^n*csc(f*x + e)^6, x)
```

**Mupad [B] (verification not implemented)**

Time = 10.17 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.62

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx =$$

$$\frac{\left(\frac{d \cos(e+fx)}{2 \cos\left(\frac{e}{2}+\frac{fx}{2}\right) \sin\left(\frac{e}{2}+\frac{fx}{2}\right)}\right)^n \left(5 \cos(e + fx) - \frac{5 \cos(3e+3fx)}{2} + \frac{\cos(5e+5fx)}{2} + 5n \cos(e + fx) - n \cos(3e + 3fx)\right)}{f \sin(e + fx)^5 (n + 1) (n + 3) (n + 5)}$$

input

```
int((d*cot(e + f*x))^n/sin(e + f*x)^6,x)
```

output

```
-(((d*cos(e + f*x))/(2*cos(e/2 + (f*x)/2)*sin(e/2 + (f*x)/2)))^n*(5*cos(e
+ f*x) - (5*cos(3*e + 3*f*x))/2 + cos(5*e + 5*f*x)/2 + 5*n*cos(e + f*x) -
n*cos(3*e + 3*f*x) + n^2*cos(e + f*x)))/(f*sin(e + f*x)^5*(n + 1)*(n + 3)*
(n + 5))
```

**Reduce [F]**

$$\int (d \cot(e + fx))^n \csc^6(e + fx) dx = d^n \left( \int \cot(fx + e)^n \csc(fx + e)^6 dx \right)$$

input `int((d*cot(f*x+e))^n*csc(f*x+e)^6,x)`

output `d**n*int(cot(e + f*x)**n*csc(e + f*x)**6,x)`

### 3.44 $\int (d \cot(e + fx))^n \csc^4(e + fx) dx$

Optimal result	364
Mathematica [A] (verified)	364
Rubi [A] (verified)	365
Maple [A] (verified)	366
Fricas [A] (verification not implemented)	367
Sympy [F]	367
Maxima [A] (verification not implemented)	367
Giac [F]	368
Mupad [B] (verification not implemented)	368
Reduce [F]	369

#### Optimal result

Integrand size = 19, antiderivative size = 51

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx = -\frac{(d \cot(e + fx))^{1+n}}{df(1+n)} - \frac{(d \cot(e + fx))^{3+n}}{d^3 f(3+n)}$$

output

$$-(d*\cot(f*x+e))^{(1+n)}/d/f/(1+n)-(d*\cot(f*x+e))^{(3+n)}/d^3/f/(3+n)$$

#### Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.88

$$\begin{aligned} & \int (d \cot(e + fx))^n \csc^4(e + fx) dx \\ &= -\frac{\cot(e + fx)(d \cot(e + fx))^n (2 + (1 + n) \csc^2(e + fx))}{f(1 + n)(3 + n)} \end{aligned}$$

input

$$\text{Integrate}[(d*\text{Cot}[e + f*x])^n*\text{Csc}[e + f*x]^4,x]$$

output

$$-((\text{Cot}[e + f*x]*(d*\text{Cot}[e + f*x])^n*(2 + (1 + n)*\text{Csc}[e + f*x]^2))/(f*(1 + n)*(3 + n)))$$

**Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {3042, 3087, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \csc^4(e + fx)(d \cot(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sec\left(e + fx - \frac{\pi}{2}\right)^4 \left(-d \tan\left(e + fx - \frac{\pi}{2}\right)\right)^n dx \\
 & \quad \downarrow \text{3087} \\
 & \frac{\int (d \cot(e + fx))^n (\cot^2(e + fx) + 1) d(-\cot(e + fx))}{f} \\
 & \quad \downarrow \text{244} \\
 & \frac{\int \left( (d \cot(e + fx))^n + \frac{(d \cot(e + fx))^{n+2}}{d^2} \right) d(-\cot(e + fx))}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{-\frac{(d \cot(e + fx))^{n+3}}{d^3(n+3)} - \frac{(d \cot(e + fx))^{n+1}}{d(n+1)}}{f}
 \end{aligned}$$

input

```
Int[(d*Cot[e + f*x])^n*Csc[e + f*x]^4,x]
```

output

```
((-(d*Cot[e + f*x])^(1 + n)/(d*(1 + n))) - (d*Cot[e + f*x])^(3 + n)/(d^3*(3 + n)))/f
```

## Definitions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

## Maple [A] (verified)

Time = 3.28 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.18

method	result	size
derivativedivides	$-\frac{\cot(fx+e)e^{n \ln(d \cot(fx+e))}}{f(1+n)} - \frac{\cot(fx+e)^3 e^{n \ln(d \cot(fx+e))}}{f(3+n)}$	60
default	$-\frac{\cot(fx+e)e^{n \ln(d \cot(fx+e))}}{f(1+n)} - \frac{\cot(fx+e)^3 e^{n \ln(d \cot(fx+e))}}{f(3+n)}$	60
risch	Expression too large to display	5257

input `int((d*cot(f*x+e))^n*csc(f*x+e)^4,x,method=_RETURNVERBOSE)`

output `-1/f/(1+n)*cot(f*x+e)*exp(n*ln(d*cot(f*x+e)))-1/f/(3+n)*cot(f*x+e)^3*exp(n*ln(d*cot(f*x+e)))`

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.71

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx$$

$$= \frac{(2 \cos(fx + e))^3 - (n + 3) \cos(fx + e) \left(\frac{d \cos(fx + e)}{\sin(fx + e)}\right)^n}{(fn^2 - (fn^2 + 4fn + 3f) \cos(fx + e)^2 + 4fn + 3f) \sin(fx + e)}$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^4,x, algorithm="fricas")`output `(2*cos(f*x + e)^3 - (n + 3)*cos(f*x + e))*(d*cos(f*x + e)/sin(f*x + e))^n/  
((f*n^2 - (f*n^2 + 4*f*n + 3*f)*cos(f*x + e)^2 + 4*f*n + 3*f)*sin(f*x + e)  
)`**Sympy [F]**

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx = \int (d \cot(e + fx))^n \csc^4(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*csc(f*x+e)**4,x)`output `Integral((d*cot(e + f*x))**n*csc(e + f*x)**4, x)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.10

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx = -\frac{\left(\frac{d}{\tan(fx + e)}\right)^{n+1}}{d(n+1)} + \frac{d^n \tan(fx + e)^{-n}}{(n+3) \tan(fx + e)^3} \frac{1}{f}$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^4,x, algorithm="maxima")`



output

```
-((d/tan(f*x + e))^(n + 1)/(d*(n + 1)) + d^n*tan(f*x + e)^(-n)/((n + 3)*tan(f*x + e)^3))/f
```

**Giac [F]**

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e)^4 dx$$

input

```
integrate((d*cot(f*x+e))^n*csc(f*x+e)^4,x, algorithm="giac")
```

output

```
integrate((d*cot(f*x + e))^n*csc(f*x + e)^4, x)
```

**Mupad [B] (verification not implemented)**

Time = 10.29 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.65

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx$$

$$= -\frac{\left(\frac{3 \cos(e+fx)}{2} - \frac{\cos(3e+3fx)}{2} + n \cos(e + fx)\right) \left(\frac{d \cos(e+fx)}{2 \cos\left(\frac{e}{2} + \frac{fx}{2}\right) \sin\left(\frac{e}{2} + \frac{fx}{2}\right)}\right)^n}{f \sin(e + fx)^3 (n + 1) (n + 3)}$$

input

```
int((d*cot(e + f*x))^n/sin(e + f*x)^4,x)
```

output

```
-(((3*cos(e + f*x))/2 - cos(3*e + 3*f*x)/2 + n*cos(e + f*x))*((d*cos(e + f*x))/(2*cos(e/2 + (f*x)/2)*sin(e/2 + (f*x)/2)))^n)/(f*sin(e + f*x)^3*(n + 1)*(n + 3))
```

**Reduce [F]**

$$\int (d \cot(e + fx))^n \csc^4(e + fx) dx = d^n \left( \int \cot(fx + e)^n \csc(fx + e)^4 dx \right)$$

input `int((d*cot(f*x+e))^n*csc(f*x+e)^4,x)`

output `d**n*int(cot(e + f*x)**n*csc(e + f*x)**4,x)`

### 3.45 $\int (d \cot(e + fx))^n \csc^2(e + fx) dx$

Optimal result	370
Mathematica [A] (verified)	370
Rubi [A] (verified)	371
Maple [A] (verified)	372
Fricas [A] (verification not implemented)	372
Sympy [F]	373
Maxima [A] (verification not implemented)	373
Giac [A] (verification not implemented)	373
Mupad [B] (verification not implemented)	374
Reduce [F]	374

#### Optimal result

Integrand size = 19, antiderivative size = 25

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = -\frac{(d \cot(e + fx))^{1+n}}{df(1+n)}$$

output

```
-(d*cot(f*x+e))^(1+n)/d/f/(1+n)
```

#### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = -\frac{\cot(e + fx)(d \cot(e + fx))^n}{f(1+n)}$$

input

```
Integrate[(d*Cot[e + f*x])^n*Csc[e + f*x]^2,x]
```

output

```
-((Cot[e + f*x]*(d*Cot[e + f*x])^n)/(f*(1 + n)))
```

**Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3087, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \csc^2(e + fx)(d \cot(e + fx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \sec\left(e + fx - \frac{\pi}{2}\right)^2 \left(-d \tan\left(e + fx - \frac{\pi}{2}\right)\right)^n dx$$

$$\downarrow \text{3087}$$

$$\frac{\int (d \cot(e + fx))^n d(-\cot(e + fx))}{f}$$

$$\downarrow \text{17}$$

$$-\frac{(d \cot(e + fx))^{n+1}}{df(n+1)}$$

input `Int[(d*Cot[e + f*x])^n*Csc[e + f*x]^2,x]`

output `-((d*Cot[e + f*x])^(1 + n)/(d*f*(1 + n)))`

**Defintions of rubi rules used**

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol]
:> Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x]
;/; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])
```

**Maple [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$-\frac{(d \cot(fx+e))^{1+n}}{df(1+n)}$	26
default	$-\frac{(d \cot(fx+e))^{1+n}}{df(1+n)}$	26
risch	Expression too large to display	1742

input

```
int((d*cot(f*x+e))^n*csc(f*x+e)^2,x,method=_RETURNVERBOSE)
```

output

```
-(d*cot(f*x+e))^(1+n)/d/f/(1+n)
```

**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = -\frac{\left(\frac{d \cos(fx+e)}{\sin(fx+e)}\right)^n \cos(fx + e)}{(fn + f) \sin(fx + e)}$$

input

```
integrate((d*cot(f*x+e))^n*csc(f*x+e)^2,x, algorithm="fricas")
```

output

```
-(d*cos(f*x + e)/sin(f*x + e))^n*cos(f*x + e)/((f*n + f)*sin(f*x + e))
```

**Sympy [F]**

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = \int (d \cot(e + fx))^n \csc^2(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*csc(f*x+e)**2,x)`

output `Integral((d*cot(e + f*x))**n*csc(e + f*x)**2, x)`

**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = -\frac{(d \cot(fx + e))^{n+1}}{df(n+1)}$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^2,x, algorithm="maxima")`

output `-(d*cot(f*x + e))^(n + 1)/(d*f*(n + 1))`

**Giac [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.88

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = -\frac{\left(-\frac{d \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - d}{2 \tan(\frac{1}{2} fx + \frac{1}{2} e)}\right)^{n+1}}{df(n+1)}$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^2,x, algorithm="giac")`

output `-(-1/2*(d*tan(1/2*f*x + 1/2*e)^2 - d)/tan(1/2*f*x + 1/2*e))^(n + 1)/(d*f*(n + 1))`

**Mupad [B] (verification not implemented)**

Time = 9.44 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.92

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = \frac{\left( \frac{\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2}{2n+2} - \frac{1}{2n+2} \right) \left( -\frac{d\left(\tan\left(\frac{e}{2} + \frac{fx}{2}\right)^2 - 1\right)}{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)} \right)^n}{f \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}$$

input `int((d*cot(e + f*x))^n/sin(e + f*x)^2,x)`output `((tan(e/2 + (f*x)/2)^2/(2*n + 2) - 1/(2*n + 2))*(-(d*(tan(e/2 + (f*x)/2)^2 - 1))/(2*tan(e/2 + (f*x)/2)))^n)/(f*tan(e/2 + (f*x)/2))`**Reduce [F]**

$$\int (d \cot(e + fx))^n \csc^2(e + fx) dx = d^n \left( \int \cot(fx + e)^n \csc(fx + e)^2 dx \right)$$

input `int((d*cot(f*x+e))^n*csc(f*x+e)^2,x)`output `d**n*int(cot(e + f*x)**n*csc(e + f*x)**2,x)`

### 3.46 $\int (d \cot(e + fx))^n \sin^2(e + fx) dx$

Optimal result	375
Mathematica [A] (verified)	375
Rubi [A] (verified)	376
Maple [F]	377
Fricas [F]	377
Sympy [F]	378
Maxima [F]	378
Giac [F]	378
Mupad [F(-1)]	379
Reduce [F]	379

#### Optimal result

Integrand size = 19, antiderivative size = 51

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx$$

$$= -\frac{(d \cot(e + fx))^{1+n} \operatorname{Hypergeometric2F1}\left(2, \frac{1+n}{2}, \frac{3+n}{2}, -\cot^2(e + fx)\right)}{df(1+n)}$$

output

`-(d*cot(f*x+e))^(1+n)*hypergeom([2, 1/2+1/2*n], [3/2+1/2*n], -cot(f*x+e)^2)/d/f/(1+n)`

#### Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.16

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx$$

$$= \frac{(d \cot(e + fx))^n \operatorname{Hypergeometric2F1}\left(2, \frac{3}{2} - \frac{n}{2}, \frac{5}{2} - \frac{n}{2}, -\tan^2(e + fx)\right) \tan^3(e + fx)}{f(3-n)}$$

input

`Integrate[(d*Cot[e + f*x])^n*Sin[e + f*x]^2,x]`



output  $((d*\cot[e + f*x])^n*\text{Hypergeometric2F1}[2, 3/2 - n/2, 5/2 - n/2, -\tan[e + f*x]^2]*\tan[e + f*x]^3)/(f*(3 - n))$

### Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3087, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^2(e + fx)(d \cot(e + fx))^n dx \\ & \quad \downarrow 3042 \\ & \int \frac{(-d \tan(e + fx - \frac{\pi}{2}))^n}{\sec(e + fx - \frac{\pi}{2})^2} dx \\ & \quad \downarrow 3087 \\ & \int \frac{(d \cot(e + fx))^n d(-\cot(e + fx))}{(\cot^2(e + fx) + 1)^2} \\ & \quad \quad \quad \downarrow 278 \\ & -\frac{(d \cot(e + fx))^{n+1} \text{Hypergeometric2F1}(2, \frac{n+1}{2}, \frac{n+3}{2}, -\cot^2(e + fx))}{df(n+1)} \end{aligned}$$

input  $\text{Int}[(d*\cot[e + f*x])^n*\sin[e + f*x]^2,x]$

output  $-(((d*\cot[e + f*x])^{(1 + n)}*\text{Hypergeometric2F1}[2, (1 + n)/2, (3 + n)/2, -\cot[e + f*x]^2])/(d*f*(1 + n)))$

### Defintions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

### Maple [F]

$$\int (d \cot (fx + e))^n \sin (fx + e)^2 dx$$

input `int((d*cot(f*x+e))^n*sin(f*x+e)^2,x)`

output `int((d*cot(f*x+e))^n*sin(f*x+e)^2,x)`

### Fricas [F]

$$\int (d \cot (e + fx))^n \sin ^2 (e + fx) dx = \int (d \cot (fx + e))^n \sin (fx + e)^2 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^2,x, algorithm="fricas")`

output `integral(-(cos(f*x + e)^2 - 1)*(d*cot(f*x + e))^n, x)`

**Sympy [F]**

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx = \int (d \cot(e + fx))^n \sin^2(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*sin(f*x+e)**2,x)`

output `Integral((d*cot(e + f*x))**n*sin(e + f*x)**2, x)`

**Maxima [F]**

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e)^2 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^2,x, algorithm="maxima")`

output `integrate((d*cot(f*x + e))^n*sin(f*x + e)^2, x)`

**Giac [F]**

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e)^2 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^2,x, algorithm="giac")`

output `integrate((d*cot(f*x + e))^n*sin(f*x + e)^2, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx = \int \sin(e + fx)^2 (d \cot(e + fx))^n dx$$

input `int(sin(e + f*x)^2*(d*cot(e + f*x))^n,x)`

output `int(sin(e + f*x)^2*(d*cot(e + f*x))^n, x)`

**Reduce [F]**

$$\int (d \cot(e + fx))^n \sin^2(e + fx) dx = d^n \left( \int \cot(fx + e)^n \sin(fx + e)^2 dx \right)$$

input `int((d*cot(f*x+e))^n*sin(f*x+e)^2,x)`

output `d**n*int(cot(e + f*x)**n*sin(e + f*x)**2,x)`

### 3.47 $\int (d \cot(e + fx))^n \sin^4(e + fx) dx$

Optimal result	380
Mathematica [A] (verified)	380
Rubi [A] (verified)	381
Maple [F]	382
Fricas [F]	382
Sympy [F]	383
Maxima [F]	383
Giac [F]	383
Mupad [F(-1)]	384
Reduce [F]	384

#### Optimal result

Integrand size = 19, antiderivative size = 51

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx$$

$$= -\frac{(d \cot(e + fx))^{1+n} \operatorname{Hypergeometric2F1}\left(3, \frac{1+n}{2}, \frac{3+n}{2}, -\cot^2(e + fx)\right)}{df(1+n)}$$

output

$-(d*\cot(f*x+e))^{(1+n)}*\operatorname{hypergeom}([3, 1/2+1/2*n], [3/2+1/2*n], -\cot(f*x+e)^2)/d/f/(1+n)$

#### Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.16

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx$$

$$= \frac{(d \cot(e + fx))^n \operatorname{Hypergeometric2F1}\left(3, \frac{5}{2} - \frac{n}{2}, \frac{7}{2} - \frac{n}{2}, -\tan^2(e + fx)\right) \tan^5(e + fx)}{f(5-n)}$$

input

$\operatorname{Integrate}[(d*\operatorname{Cot}[e + f*x])^n*\operatorname{Sin}[e + f*x]^4,x]$

output  $((d*\cot[e + f*x])^n*\text{Hypergeometric2F1}[3, 5/2 - n/2, 7/2 - n/2, -\tan[e + f*x]^2*\tan[e + f*x]^5)/(f*(5 - n))$

### Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$ , Rules used = {3042, 3087, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sin^4(e + fx)(d \cot(e + fx))^n dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(-d \tan(e + fx - \frac{\pi}{2}))^n}{\sec(e + fx - \frac{\pi}{2})^4} dx \\ & \quad \downarrow \text{3087} \\ & \int \frac{(d \cot(e + fx))^n d(-\cot(e + fx))}{(\cot^2(e + fx) + 1)^3} \\ & \quad \quad \quad \downarrow \text{278} \\ & \frac{(d \cot(e + fx))^{n+1} \text{Hypergeometric2F1}\left(3, \frac{n+1}{2}, \frac{n+3}{2}, -\cot^2(e + fx)\right)}{df(n+1)} \end{aligned}$$

input  $\text{Int}[(d*\cot[e + f*x])^n*\sin[e + f*x]^4,x]$

output  $-(((d*\cot[e + f*x])^{(1 + n)}*\text{Hypergeometric2F1}[3, (1 + n)/2, (3 + n)/2, -\cot[e + f*x]^2])/(d*f*(1 + n)))$

**Defintions of rubi rules used**

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3087 `Int[sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[1/f Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

**Maple [F]**

$$\int (d \cot (fx + e))^n \sin (fx + e)^4 dx$$

input `int((d*cot(f*x+e))^n*sin(f*x+e)^4,x)`

output `int((d*cot(f*x+e))^n*sin(f*x+e)^4,x)`

**Fricas [F]**

$$\int (d \cot (e + fx))^n \sin^4 (e + fx) dx = \int (d \cot (fx + e))^n \sin (fx + e)^4 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^4,x, algorithm="fricas")`

output `integral((cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*(d*cot(f*x + e))^n, x)`

**Sympy [F]**

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx = \int (d \cot(e + fx))^n \sin^4(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*sin(f*x+e)**4,x)`

output `Integral((d*cot(e + f*x))**n*sin(e + f*x)**4, x)`

**Maxima [F]**

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e)^4 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^4,x, algorithm="maxima")`

output `integrate((d*cot(f*x + e))^n*sin(f*x + e)^4, x)`

**Giac [F]**

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e)^4 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^4,x, algorithm="giac")`

output `integrate((d*cot(f*x + e))^n*sin(f*x + e)^4, x)`



**Mupad [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx = \int \sin(e + fx)^4 (d \cot(e + fx))^n dx$$

input `int(sin(e + f*x)^4*(d*cot(e + f*x))^n,x)`output `int(sin(e + f*x)^4*(d*cot(e + f*x))^n, x)`**Reduce [F]**

$$\int (d \cot(e + fx))^n \sin^4(e + fx) dx = d^n \left( \int \cot(fx + e)^n \sin(fx + e)^4 dx \right)$$

input `int((d*cot(f*x+e))^n*sin(f*x+e)^4,x)`output `d**n*int(cot(e + f*x)**n*sin(e + f*x)**4,x)`

### 3.48 $\int (d \cot(e + fx))^n \csc^3(e + fx) dx$

Optimal result	385
Mathematica [C] (warning: unable to verify)	385
Rubi [A] (verified)	386
Maple [F]	387
Fricas [F]	388
Sympy [F]	388
Maxima [F]	388
Giac [F]	389
Mupad [F(-1)]	389
Reduce [F]	389

#### Optimal result

Integrand size = 19, antiderivative size = 79

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \frac{(d \cot(e + fx))^{1+n} \csc^3(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{4+n}{2}, \frac{3+n}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)^{\frac{4+n}{2}}}{df(1+n)}$$

output

```
-(d*cot(f*x+e))^(1+n)*csc(f*x+e)^3*hypergeom([2+1/2*n, 1/2+1/2*n],[3/2+1/2*n],cos(f*x+e)^2)*(sin(f*x+e)^2)^(2+1/2*n)/d/f/(1+n)
```

#### Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 7.81 (sec) , antiderivative size = 741, normalized size of antiderivative = 9.38

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \text{Too large to display}$$

input

```
Integrate[(d*Cot[e + f*x])^n*Csc[e + f*x]^3,x]
```

output

```

((d*Cot[e + f*x])^n*(-1/2*(Cot[(e + f*x)/2]^2*Hypergeometric2F1[-1 - n/2,
-n, -1/2*n, Tan[(e + f*x)/2]^2])/((2 + n)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2
)^n) + (2*Hypergeometric2F1[1 - n/2, -n, 2 - n/2, Tan[(e + f*x)/2]^2]*Tan[
(e + f*x)/2]^2)/((8 - 4*n)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n) - ((-4 + n
)*AppellF1[1 - n/2, -n, 1, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^
2]*Sin[(e + f*x)/2]^2)/((-2 + n)*((-4 + n)*AppellF1[1 - n/2, -n, 1, 2 - n/
2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*(n*AppellF1[2 - n/2, 1 - n
, 1, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[2 - n/2,
-n, 2, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2
]^2)) + ((-4 + n)*Cos[(e + f*x)/2]^2*(-2 + n)*Hypergeometric2F1[-n, -1/2*
n, 1 - n/2, Tan[(e + f*x)/2]^2] - n*AppellF1[1 - n/2, -n, 1, 2 - n/2, Tan[
(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Tan[(e + f*x)/2]^2))/((-2 + n)*n*(4*(
Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n - n*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n
+ (-4 + n)*AppellF1[1 - n/2, -n, 1, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e
+ f*x)/2]^2]*Tan[(e + f*x)/2]^2 + 2*n*AppellF1[2 - n/2, 1 - n, 1, 3 - n/2,
Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Tan[(e + f*x)/2]^4 + 2*AppellF1[
2 - n/2, -n, 2, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Tan[(e +
f*x)/2]^4))))/(2*f)

```

## Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \csc^3(e + fx)(d \cot(e + fx))^n dx$$

↓ 3042

$$\int \sec\left(e + fx - \frac{\pi}{2}\right)^3 \left(-d \tan\left(e + fx - \frac{\pi}{2}\right)\right)^n dx$$

↓ 3097

$$\frac{\csc^3(e + fx) \sin^2(e + fx)^{\frac{n+4}{2}} (d \cot(e + fx))^{n+1} \text{Hypergeometric2F1}\left(\frac{n+1}{2}, \frac{n+4}{2}, \frac{n+3}{2}, \cos^2(e + fx)\right)}{df(n+1)}$$

input `Int[(d*Cot[e + f*x])^n*Csc[e + f*x]^3,x]`

output `-(((d*Cot[e + f*x])^(1 + n)*Csc[e + f*x]^3*Hypergeometric2F1[(1 + n)/2, (4 + n)/2, (3 + n)/2, Cos[e + f*x]^2]*(Sin[e + f*x]^2)^((4 + n)/2))/(d*f*(1 + n))`

### Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3097 `Int[((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n + 1)*((Cos[e + f*x]^2)^((m + n + 1)/2)/(b*f*(n + 1)))*Hypergeometric2F1[(n + 1)/2, (m + n + 1)/2, (n + 3)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[(n - 1)/2] && !IntegerQ[m/2]`

### Maple [F]

$$\int (d \cot (fx + e))^n \csc (fx + e)^3 dx$$

input `int((d*cot(f*x+e))^n*csc(f*x+e)^3,x)`

output `int((d*cot(f*x+e))^n*csc(f*x+e)^3,x)`

**Fricas [F]**

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e)^3 dx$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^3,x, algorithm="fricas")`

output `integral((d*cot(f*x + e))^n*csc(f*x + e)^3, x)`

**Sympy [F]**

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \int (d \cot(e + fx))^n \csc^3(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*csc(f*x+e)**3,x)`

output `Integral((d*cot(e + f*x))**n*csc(e + f*x)**3, x)`

**Maxima [F]**

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e)^3 dx$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^3,x, algorithm="maxima")`

output `integrate((d*cot(f*x + e))^n*csc(f*x + e)^3, x)`

**Giac [F]**

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e)^3 dx$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e)^3,x, algorithm="giac")`

output `integrate((d*cot(f*x + e))^n*csc(f*x + e)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = \int \frac{(d \cot(e + fx))^n}{\sin(e + fx)^3} dx$$

input `int((d*cot(e + f*x))^n/sin(e + f*x)^3,x)`

output `int((d*cot(e + f*x))^n/sin(e + f*x)^3, x)`

**Reduce [F]**

$$\int (d \cot(e + fx))^n \csc^3(e + fx) dx = d^n \left( \int \cot(fx + e)^n \csc(fx + e)^3 dx \right)$$

input `int((d*cot(f*x+e))^n*csc(f*x+e)^3,x)`

output `d**n*int(cot(e + f*x)**n*csc(e + f*x)**3,x)`

### 3.49 $\int (d \cot(e + fx))^n \csc(e + fx) dx$

Optimal result	390
Mathematica [A] (verified)	390
Rubi [A] (verified)	391
Maple [F]	392
Fricas [F]	392
Sympy [F]	393
Maxima [F]	393
Giac [F]	393
Mupad [F(-1)]	394
Reduce [F]	394

#### Optimal result

Integrand size = 17, antiderivative size = 77

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = \frac{(d \cot(e + fx))^{1+n} \csc(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{2+n}{2}, \frac{3+n}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)^{\frac{2+n}{2}}}{df(1+n)}$$

output

```
-(d*cot(f*x+e))^(1+n)*csc(f*x+e)*hypergeom([1+1/2*n, 1/2+1/2*n], [3/2+1/2*n], cos(f*x+e)^2)*(sin(f*x+e)^2)^(1+1/2*n)/d/f/(1+n)
```

#### Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.90

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = \frac{(d \cot(e + fx))^n \operatorname{Hypergeometric2F1}\left(-n, -\frac{n}{2}, 1 - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right)\right) (\cos(e + fx) \sec^2\left(\frac{1}{2}(e + fx)\right))}{fn}$$

input

```
Integrate[(d*Cot[e + f*x])^n*Csc[e + f*x], x]
```

output 
$$-\left(\left(\frac{d \cot(e + f x)}{dx}\right)^n \operatorname{Hypergeometric2F1}\left[-n, -\frac{1}{2}n, 1 - \frac{n}{2}, \tan\left(\frac{e + f x}{2}\right)^2\right]\right) / \left(f^n \cos\left(\frac{e + f x}{2}\right) \sec\left(\frac{e + f x}{2}\right)^{2n}\right)$$

### Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \csc(e + f x) (d \cot(e + f x))^n dx$$

$$\downarrow \text{3042}$$

$$\int \sec\left(e + f x - \frac{\pi}{2}\right) \left(-d \tan\left(e + f x - \frac{\pi}{2}\right)\right)^n dx$$

$$\downarrow \text{3097}$$

$$\frac{\csc(e + f x) \sin^2(e + f x)^{\frac{n+2}{2}} (d \cot(e + f x))^{n+1} \operatorname{Hypergeometric2F1}\left(\frac{n+1}{2}, \frac{n+2}{2}, \frac{n+3}{2}, \cos^2(e + f x)\right)}{df(n+1)}$$

input 
$$\operatorname{Int}\left[\left(\frac{d \cot(e + f x)}{dx}\right)^n \operatorname{Csc}[e + f x], x\right]$$

output 
$$-\left(\left(\frac{d \cot(e + f x)}{dx}\right)^{(1+n)} \operatorname{Csc}[e + f x] \operatorname{Hypergeometric2F1}\left[\frac{(1+n)}{2}, \frac{(2+n)}{2}, \frac{(3+n)}{2}, \cos[e + f x]^2\right] \left(\sin[e + f x]^2\right)^{\frac{(2+n)}{2}}\right) / \left(d f (1+n)\right)$$



## Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3097 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n + 1)*((Cos[e + f*x]^2)^(m + n + 1)/2)/(b*f*(n + 1))*Hypergeometric2F1[(n + 1)/2, (m + n + 1)/2, (n + 3)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[(n - 1)/2] && !IntegerQ[m/2]`

## Maple [F]

$$\int (d \cot (fx + e))^n \csc (fx + e) dx$$

input `int((d*cot(f*x+e))^n*csc(f*x+e),x)`

output `int((d*cot(f*x+e))^n*csc(f*x+e),x)`

## Fricas [F]

$$\int (d \cot (e + fx))^n \csc (e + fx) dx = \int (d \cot (fx + e))^n \csc (fx + e) dx$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e),x, algorithm="fricas")`

output `integral((d*cot(f*x + e))^n*csc(f*x + e), x)`

**Sympy [F]**

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = \int (d \cot(e + fx))^n \csc(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*csc(f*x+e),x)`

output `Integral((d*cot(e + f*x))**n*csc(e + f*x), x)`

**Maxima [F]**

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e) dx$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e),x, algorithm="maxima")`

output `integrate((d*cot(f*x + e))^n*csc(f*x + e), x)`

**Giac [F]**

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = \int (d \cot(fx + e))^n \csc(fx + e) dx$$

input `integrate((d*cot(f*x+e))^n*csc(f*x+e),x, algorithm="giac")`

output `integrate((d*cot(f*x + e))^n*csc(f*x + e), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = \int \frac{(d \cot(e + fx))^n}{\sin(e + fx)} dx$$

input `int((d*cot(e + f*x))^n/sin(e + f*x),x)`output `int((d*cot(e + f*x))^n/sin(e + f*x), x)`**Reduce [F]**

$$\int (d \cot(e + fx))^n \csc(e + fx) dx = d^n \left( \int \cot(fx + e)^n \csc(fx + e) dx \right)$$

input `int((d*cot(f*x+e))^n*csc(f*x+e),x)`output `d**n*int(cot(e + f*x)**n*csc(e + f*x),x)`

### 3.50 $\int (d \cot(e + fx))^n \sin(e + fx) dx$

Optimal result	395
Mathematica [C] (warning: unable to verify)	395
Rubi [A] (verified)	396
Maple [F]	397
Fricas [F]	397
Sympy [F]	398
Maxima [F]	398
Giac [F]	398
Mupad [F(-1)]	399
Reduce [F]	399

#### Optimal result

Integrand size = 17, antiderivative size = 73

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = \frac{(d \cot(e + fx))^{1+n} \operatorname{Hypergeometric2F1}\left(\frac{n}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(e + fx)\right) \sin(e + fx) \sin^2(e + fx)^{n/2}}{df(1+n)}$$

output

```
-(d*cot(f*x+e))^(1+n)*hypergeom([1/2*n, 1/2+1/2*n],[3/2+1/2*n],cos(f*x+e)^2)*sin(f*x+e)*(sin(f*x+e)^2)^(1/2*n)/d/f/(1+n)
```

#### Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 0.89 (sec) , antiderivative size = 264, normalized size of antiderivative = 3.62

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = \frac{8(-4 + n) \operatorname{AppellF1}\left(1 - \frac{n}{2}, -n, 2, 2 - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) \cos^2\left(\frac{1}{2}(e + fx)\right)}{f(-2 + n) (2(-4 + n) \operatorname{AppellF1}\left(1 - \frac{n}{2}, -n, 2, 2 - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) \cos^2\left(\frac{1}{2}(e + fx)\right))}$$

input

```
Integrate[(d*Cot[e + f*x])^n*Sin[e + f*x],x]
```

output

```
(-8*(-4 + n)*AppellF1[1 - n/2, -n, 2, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^4*(d*Cot[e + f*x])^n*Sin[(e + f*x)/2]^2)/(f*(-2 + n)*(2*(-4 + n)*AppellF1[1 - n/2, -n, 2, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 - 2*(n*AppellF1[2 - n/2, 1 - n, 2, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*AppellF1[2 - n/2, -n, 3, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*(-1 + Cos[e + f*x])))
```

**Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$ , Rules used = {3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin(e + fx)(d \cot(e + fx))^n dx$$

↓ 3042

$$\int \frac{(-d \tan(e + fx - \frac{\pi}{2}))^n}{\sec(e + fx - \frac{\pi}{2})} dx$$

↓ 3097

$$\frac{\sin(e + fx) \sin^2(e + fx)^{n/2} (d \cot(e + fx))^{n+1} \text{Hypergeometric2F1}\left(\frac{n}{2}, \frac{n+1}{2}, \frac{n+3}{2}, \cos^2(e + fx)\right)}{df(n+1)}$$

input

```
Int[(d*Cot[e + f*x])^n*Sin[e + f*x],x]
```

output

```
-(((d*Cot[e + f*x])^(1 + n)*Hypergeometric2F1[n/2, (1 + n)/2, (3 + n)/2, Cos[e + f*x]^2]*Sin[e + f*x]*(Sin[e + f*x]^2)^(n/2))/(d*f*(1 + n)))
```

## Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3097 `Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n + 1)*((Cos[e + f*x]^2)^(m + n + 1)/2)/(b*f*(n + 1))*Hypergeometric2F1[(n + 1)/2, (m + n + 1)/2, (n + 3)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[(n - 1)/2] && !IntegerQ[m/2]`

## Maple [F]

$$\int (d \cot (fx + e))^n \sin (fx + e) dx$$

input `int((d*cot(f*x+e))^n*sin(f*x+e),x)`

output `int((d*cot(f*x+e))^n*sin(f*x+e),x)`

## Fricas [F]

$$\int (d \cot (e + fx))^n \sin (e + fx) dx = \int (d \cot (fx + e))^n \sin (fx + e) dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e),x, algorithm="fricas")`

output `integral((d*cot(f*x + e))^n*sin(f*x + e), x)`

**Sympy [F]**

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = \int (d \cot(e + fx))^n \sin(e + fx) dx$$

input `integrate((d*cot(f*x+e))**n*sin(f*x+e),x)`

output `Integral((d*cot(e + f*x))**n*sin(e + f*x), x)`

**Maxima [F]**

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e) dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e),x, algorithm="maxima")`

output `integrate((d*cot(f*x + e))^n*sin(f*x + e), x)`

**Giac [F]**

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e) dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e),x, algorithm="giac")`

output `integrate((d*cot(f*x + e))^n*sin(f*x + e), x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = \int \sin(e + fx) (d \cot(e + fx))^n dx$$

input `int(sin(e + f*x)*(d*cot(e + f*x))^n,x)`output `int(sin(e + f*x)*(d*cot(e + f*x))^n, x)`**Reduce [F]**

$$\int (d \cot(e + fx))^n \sin(e + fx) dx = d^n \left( \int \cot(fx + e)^n \sin(fx + e) dx \right)$$

input `int((d*cot(f*x+e))^n*sin(f*x+e),x)`output `d**n*int(cot(e + f*x)**n*sin(e + f*x),x)`



### 3.51 $\int (d \cot(e + fx))^n \sin^3(e + fx) dx$

Optimal result	400
Mathematica [C] (warning: unable to verify)	400
Rubi [A] (verified)	401
Maple [F]	402
Fricas [F]	402
Sympy [F(-1)]	403
Maxima [F]	403
Giac [F]	403
Mupad [F(-1)]	404
Reduce [F]	404

#### Optimal result

Integrand size = 19, antiderivative size = 79

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = \frac{(d \cot(e + fx))^{1+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}(-2 + n), \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(e + fx)\right) \sin^3(e + fx) \sin^2(e + fx)}{df(1 + n)}$$

output

$$-(d*\cot(f*x+e))^{(1+n)}*hypergeom([-1+1/2*n, 1/2+1/2*n], [3/2+1/2*n], \cos(f*x+e)^2)*\sin(f*x+e)^3*(\sin(f*x+e)^2)^{(-1+1/2*n)}/d/f/(1+n)$$

#### Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.98 (sec) , antiderivative size = 477, normalized size of antiderivative = 6.04

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = \frac{f(-2 + n) (2(-4 + n) \operatorname{AppellF1}\left(1 - \frac{n}{2}, -n, 3, 2 - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) \cos^2\left(\frac{1}{2}(e + fx)\right) \sin^3(e + fx)}{df(1 + n)}$$

input

$$\operatorname{Integrate}[(d*\cot[e + f*x])^n*\sin[e + f*x]^3,x]$$

output

```
(-4*(-4 + n)*(AppellF1[1 - n/2, -n, 3, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - AppellF1[1 - n/2, -n, 4, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Cos[(e + f*x)/2]^3*(d*Cot[e + f*x])^n*Sin[(e + f*x)/2]*Sin[e + f*x]^3)/(f*(-2 + n)*(2*(-4 + n)*AppellF1[1 - n/2, -n, 3, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 - 2*(-4 + n)*AppellF1[1 - n/2, -n, 4, 2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 - 2*(n*AppellF1[2 - n/2, 1 - n, 3, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - n*AppellF1[2 - n/2, 1 - n, 4, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 3*AppellF1[2 - n/2, -n, 4, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 4*AppellF1[2 - n/2, -n, 5, 3 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]))*(-1 + Cos[e + f*x]))
```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sin^3(e + fx)(d \cot(e + fx))^n dx$$

↓ 3042

$$\int \frac{(-d \tan(e + fx - \frac{\pi}{2}))^n}{\sec(e + fx - \frac{\pi}{2})^3} dx$$

↓ 3097

$$\frac{\sin^3(e + fx) \sin^2(e + fx)^{\frac{n-2}{2}} (d \cot(e + fx))^{n+1} \text{Hypergeometric2F1}\left(\frac{n-2}{2}, \frac{n+1}{2}, \frac{n+3}{2}, \cos^2(e + fx)\right)}{df(n+1)}$$

input

```
Int[(d*Cot[e + f*x])^n*Sin[e + f*x]^3,x]
```

output  $-\left(\left(d \cot [e+f x]\right)^{(1+n)} \operatorname{Hypergeometric2F1}\left[\frac{-2+n}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos [e+f x]^2\right] \sin [e+f x]^3 \left(\sin [e+f x]^2\right)^{\frac{-2+n}{2}}\right) / \left(d f (1+n)\right)$

### Defintions of rubi rules used

rule 3042  $\operatorname{Int}[u, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] / ; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3097  $\operatorname{Int}\left[\left(\left(a_{\cdot}\right) \sec \left[\left(e_{\cdot}\right)+\left(f_{\cdot}\right)\left(x_{\cdot}\right)\right]\right)^{\left(m_{\cdot}\right)} \left(\left(b_{\cdot}\right) \tan \left[\left(e_{\cdot}\right)+\left(f_{\cdot}\right)\left(x_{\cdot}\right)\right]\right)^{\left(n_{\cdot}\right)}\right], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}\left[\left(a \sec [e+f x]\right)^m \left(b \tan [e+f x]\right)^{n+1} \left(\cos [e+f x]^2\right)^{\frac{m+n+1}{2}} / \left(b f (n+1)\right) \operatorname{Hypergeometric2F1}\left[\frac{n+1}{2}, \frac{m+n+1}{2}, \frac{n+3}{2}, \sin [e+f x]^2\right], x\right] / ; \operatorname{FreeQ}\{a, b, e, f, m, n, x\} \&\& !\operatorname{IntegerQ}\left[\frac{n-1}{2}\right] \&\& !\operatorname{IntegerQ}\left[\frac{m}{2}\right]$

### Maple [F]

$$\int (d \cot (f x+e))^n \sin (f x+e)^3 dx$$

input  $\operatorname{int}\left(\left(d \cot (f x+e)\right)^n \sin (f x+e)^3, x\right)$

output  $\operatorname{int}\left(\left(d \cot (f x+e)\right)^n \sin (f x+e)^3, x\right)$

### Fricas [F]

$$\int (d \cot (e+f x))^n \sin ^3(e+f x) dx = \int (d \cot (f x+e))^n \sin (f x+e)^3 dx$$

input  $\operatorname{integrate}\left(\left(d \cot (f x+e)\right)^n \sin (f x+e)^3, x, \text{algorithm}=\text{"fricas"}\right)$

output  $\operatorname{integral}\left(-\left(\cos (f x+e)\right)^2-1\right) \cdot\left(d \cot (f x+e)\right)^n \sin (f x+e), x$

**Sympy [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = \text{Timed out}$$

input `integrate((d*cot(f*x+e))**n*sin(f*x+e)**3,x)`output `Timed out`**Maxima [F]**

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e)^3 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^3,x, algorithm="maxima")`output `integrate((d*cot(f*x + e))^n*sin(f*x + e)^3, x)`**Giac [F]**

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = \int (d \cot(fx + e))^n \sin(fx + e)^3 dx$$

input `integrate((d*cot(f*x+e))^n*sin(f*x+e)^3,x, algorithm="giac")`output `integrate((d*cot(f*x + e))^n*sin(f*x + e)^3, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = \int \sin(e + fx)^3 (d \cot(e + fx))^n dx$$

input `int(sin(e + f*x)^3*(d*cot(e + f*x))^n,x)`

output `int(sin(e + f*x)^3*(d*cot(e + f*x))^n, x)`

**Reduce [F]**

$$\int (d \cot(e + fx))^n \sin^3(e + fx) dx = d^n \left( \int \cot(fx + e)^n \sin(fx + e)^3 dx \right)$$

input `int((d*cot(f*x+e))^n*sin(f*x+e)^3,x)`

output `d**n*int(cot(e + f*x)**n*sin(e + f*x)**3,x)`

### 3.52 $\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx$

Optimal result	405
Mathematica [C] (warning: unable to verify)	405
Rubi [A] (verified)	406
Maple [F]	407
Fricas [F]	407
Sympy [F]	408
Maxima [F]	408
Giac [F]	408
Mupad [F(-1)]	409
Reduce [F]	409

#### Optimal result

Integrand size = 21, antiderivative size = 83

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = \frac{(b \cot(e + fx))^{1+n} (a \csc(e + fx))^m \operatorname{Hypergeometric2F1}\left(\frac{1+n}{2}, \frac{1}{2}(1+m+n), \frac{3+n}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)}{bf(1+n)}$$

output

```
-(b*cot(f*x+e))^(1+n)*(a*csc(f*x+e))^m*hypergeom([1/2+1/2*n, 1/2+1/2*m+1/2*n], [3/2+1/2*n], cos(f*x+e)^2)*(sin(f*x+e)^2)^(1/2+1/2*m+1/2*n)/b/f/(1+n)
```

#### Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.94 (sec) , antiderivative size = 306, normalized size of antiderivative = 3.69

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = \frac{f(-1+m+n) ((-3+m+n) \operatorname{AppellF1}\left(\frac{1}{2}(1-m-n), -n, 1-m, \frac{1}{2}(3-m-n), \tan^2\left(\frac{1}{2}(e+fx)\right)\right) \tan^2\left(\frac{1}{2}(e+fx)\right)}{f(-1+m+n)}$$

input

```
Integrate[(b*Cot[e + f*x])^n*(a*Csc[e + f*x])^m,x]
```

output

```

-((a*(-3 + m + n)*AppellF1[(1 - m - n)/2, -n, 1 - m, (3 - m - n)/2, Tan[(e
+ f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(b*Cot[e + f*x])^n*(a*Csc[e + f*x])^(-1
+ m))/(f*(-1 + m + n)*((-3 + m + n)*AppellF1[(1 - m - n)/2, -n, 1 - m, (3
- m - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*(n*AppellF1[(3 -
m - n)/2, 1 - n, 1 - m, (5 - m - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)
/2]^2] - (-1 + m)*AppellF1[(3 - m - n)/2, -n, 2 - m, (5 - m - n)/2, Tan[(e
+ f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))

```

**Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {3042, 3097}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a \csc(e + fx))^m (b \cot(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left( a \sec\left(e + fx - \frac{\pi}{2}\right) \right)^m \left( -b \tan\left(e + fx - \frac{\pi}{2}\right) \right)^n dx \\
 & \quad \downarrow \text{3097} \\
 & \frac{(a \csc(e + fx))^m (b \cot(e + fx))^{n+1} \sin^2(e + fx)^{\frac{1}{2}(m+n+1)} \text{Hypergeometric2F1}\left(\frac{n+1}{2}, \frac{1}{2}(m+n+1), \frac{n+3}{2}, \cos^2(e + fx)\right)}{bf(n+1)}
 \end{aligned}$$

input

```
Int[(b*Cot[e + f*x])^n*(a*Csc[e + f*x])^m,x]
```

output

```

-(((b*Cot[e + f*x])^(1 + n)*(a*Csc[e + f*x])^m*Hypergeometric2F1[(1 + n)/2
, (1 + m + n)/2, (3 + n)/2, Cos[e + f*x]^2]*(Sin[e + f*x]^2)^((1 + m + n)/
2))/(b*f*(1 + n))

```

## Definitions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3097 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n + 1)*((Cos[e + f*x]^2)^(m + n + 1)/2)/(b*f*(n + 1))*Hypergeometric2F1[(n + 1)/2, (m + n + 1)/2, (n + 3)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[(n - 1)/2] && !IntegerQ[m/2]`

## Maple [F]

$$\int (b \cot (fx + e))^n (a \csc (fx + e))^m dx$$

input `int((b*cot(f*x+e))^n*(a*csc(f*x+e))^m,x)`

output `int((b*cot(f*x+e))^n*(a*csc(f*x+e))^m,x)`

## Fricas [F]

$$\int (b \cot (e + fx))^n (a \csc (e + fx))^m dx = \int (b \cot (fx + e))^n (a \csc (fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*csc(f*x+e))^m,x, algorithm="fricas")`

output `integral((b*cot(f*x + e))^n*(a*csc(f*x + e))^m, x)`



**Sympy [F]**

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = \int (a \csc(e + fx))^m (b \cot(e + fx))^n dx$$

input `integrate((b*cot(f*x+e))**n*(a*csc(f*x+e))**m,x)`

output `Integral((a*csc(e + f*x))**m*(b*cot(e + f*x))**n, x)`

**Maxima [F]**

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = \int (b \cot(fx + e))^n (a \csc(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*csc(f*x+e))^m,x, algorithm="maxima")`

output `integrate((b*cot(f*x + e))^n*(a*csc(f*x + e))^m, x)`

**Giac [F]**

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = \int (b \cot(fx + e))^n (a \csc(fx + e))^m dx$$

input `integrate((b*cot(f*x+e))^n*(a*csc(f*x+e))^m,x, algorithm="giac")`

output `integrate((b*cot(f*x + e))^n*(a*csc(f*x + e))^m, x)`

**Mupad [F(-1)]**

Timed out.

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = \int (b \cot(e + fx))^n \left( \frac{a}{\sin(e + fx)} \right)^m dx$$

input `int((b*cot(e + f*x))^n*(a/sin(e + f*x))^m,x)`output `int((b*cot(e + f*x))^n*(a/sin(e + f*x))^m, x)`**Reduce [F]**

$$\int (b \cot(e + fx))^n (a \csc(e + fx))^m dx = b^n a^m \left( \int \csc(fx + e)^m \cot(fx + e)^n dx \right)$$

input `int((b*cot(f*x+e))^n*(a*csc(f*x+e))^m,x)`output `b**n*a**m*int(csc(e + f*x)**m*cot(e + f*x)**n,x)`

# CHAPTER 4

## APPENDIX

4.1	Listing of Grading functions . . . . .	410
4.2	Links to plain text integration problems used in this report for each CAS .	428

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "
  ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],  
If [AppellFunctionQ [Head [expn]],  
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],  
If [Head [expn] == RootSum,  
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],  
If [Head [expn] == Integrate || Head [expn] == Int,  
Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],  
9]]]]]]]]]]
```

```
ElementaryFunctionQ [func_] :=  
MemberQ [{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
}, func]
```

```
SpecialFunctionQ [func_] :=  
MemberQ [{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,  
Gamma, LogGamma, PolyGamma,  
Zeta, PolyLog, ProductLog,  
EllipticF, EllipticE, EllipticPi  
}, func]
```

```
HypergeometricFunctionQ [func_] :=  
MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ [func_] :=  
MemberQ [{AppellF1}, func]
```

## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```



```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
            fi;
        fi;
    else #ExpnType(result) > ExpnType(optimal)
        if debug then
            print("ExpnType(result) > ExpnType(optimal)");
        fi;
        return "C",cat("Result contains higher order function than in optimal. Order ",
                        convert(ExpnType_result,string)," vs. order ",
                        convert(ExpnType_optimal,string),".");
    fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9
end proc

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

## Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```



```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

## 4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file