

Computer Algebra Independent Integration Tests

Summer 2024

4-Trig-functions/4.6-Cosecant/246-4.6.1.4

Nasser M. Abbasi

May 18, 2024 Compiled on May 18, 2024 at 1:17am

Contents

1	Introduction	3
1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22
2	detailed summary tables of results	23
2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	34
3	Listing of integrals	36
3.1	$\int \frac{\tan^4(x)}{a+a \csc(x)} dx$	38
3.2	$\int \frac{\tan^3(x)}{a+a \csc(x)} dx$	46
3.3	$\int \frac{\tan^2(x)}{a+a \csc(x)} dx$	52
3.4	$\int \frac{\tan(x)}{a+a \csc(x)} dx$	58
3.5	$\int \frac{\cot(x)}{a+a \csc(x)} dx$	64
3.6	$\int \frac{\cot^2(x)}{a+a \csc(x)} dx$	69

3.7	$\int \frac{\cot^3(x)}{a+a \csc(x)} dx$	74
3.8	$\int \frac{\cot^4(x)}{a+a \csc(x)} dx$	79
3.9	$\int \frac{\cot^5(x)}{a+a \csc(x)} dx$	86
3.10	$\int \frac{\cot^6(x)}{a+a \csc(x)} dx$	92
3.11	$\int \frac{\cot^7(x)}{a+a \csc(x)} dx$	99
3.12	$\int \frac{\tan^5(x)}{a+b \csc(x)} dx$	105
3.13	$\int \frac{\tan^3(x)}{a+b \csc(x)} dx$	113
3.14	$\int \frac{\tan(x)}{a+b \csc(x)} dx$	120
3.15	$\int \frac{\cot(x)}{a+b \csc(x)} dx$	126
3.16	$\int \frac{\cot^3(x)}{a+b \csc(x)} dx$	131
3.17	$\int \frac{\cot^5(x)}{a+b \csc(x)} dx$	137
3.18	$\int \frac{\cot^7(x)}{a+b \csc(x)} dx$	144
3.19	$\int \frac{\tan^4(x)}{a+b \csc(x)} dx$	151
3.20	$\int \frac{\tan^2(x)}{a+b \csc(x)} dx$	163
3.21	$\int \frac{\cot^2(x)}{a+b \csc(x)} dx$	172
3.22	$\int \frac{\cot^4(x)}{a+b \csc(x)} dx$	181
3.23	$\int \frac{\cot^6(x)}{a+b \csc(x)} dx$	190
4	Appendix	199
4.1	Listing of Grading functions	199
4.2	Links to plain text integration problems used in this report for each CAS217	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	Performance based on number of rules Rubi used	11
1.5	Performance based on number of steps Rubi used	12
1.6	Solved integrals histogram based on leaf size of result	13
1.7	Solved integrals histogram based on CPU time used	14
1.8	Leaf size vs. CPU time used	15
1.9	list of integrals with no known antiderivative	16
1.10	List of integrals solved by CAS but has no known antiderivative	16
1.11	list of integrals solved by CAS but failed verification	16
1.12	Timing	17
1.13	Verification	17
1.14	Important notes about some of the results	18
1.15	Current tree layout of integration tests	21
1.16	Design of the test system	22

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [23]. This is test number [246].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (23)	0.00 (0)
Mathematica	100.00 (23)	0.00 (0)
Maple	100.00 (23)	0.00 (0)
Fricas	100.00 (23)	0.00 (0)
Mupad	100.00 (23)	0.00 (0)
Giac	100.00 (23)	0.00 (0)
Reduce	100.00 (23)	0.00 (0)
Maxima	78.26 (18)	21.74 (5)
Sympy	0.00 (0)	100.00 (23)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Maple	95.652	0.000	4.348	0.000
Mathematica	86.957	13.043	0.000	0.000
Giac	86.957	13.043	0.000	0.000
Fricas	73.913	26.087	0.000	0.000
Maxima	60.870	17.391	0.000	21.739
Mupad	0.000	100.000	0.000	0.000
Reduce	0.000	100.000	0.000	0.000
Sympy	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	0	0.00	0.00	0.00
Mupad	0	0.00	0.00	0.00
Giac	0	0.00	0.00	0.00
Reduce	0	0.00	0.00	0.00
Maxima	5	0.00	0.00	100.00
Sympy	23	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.05
Fricas	0.12
Giac	0.15
Reduce	0.18
Mathematica	0.20
Maple	0.39
Rubi	0.39
Mupad	15.87
Sympy	-nan(ind)

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	75.91	1.06	58.00	1.11
Maxima	79.56	1.45	54.00	1.12
Maple	79.70	1.16	60.00	1.19
Mathematica	89.26	1.30	71.00	1.04
Giac	91.52	1.23	65.00	1.23
Fricas	166.35	1.92	70.00	1.27
Reduce	192.74	2.25	92.00	1.93
Mupad	689.09	6.59	113.00	2.33
Sympy	-nan(ind)	-nan(ind)	nan	nan

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A `time limit` of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests




Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	24
2.2	Detailed conclusion table per each integral for all CAS systems	28
2.3	Detailed conclusion table specific for Rubi results	34

2.1 List of integrals sorted by grade for each CAS

Rubi	24
Mma	24
Maple	25
Fricas	25
Maxima	25
Giac	26
Mupad	26
Sympy	26
Reduce	27

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

B grade { 1, 8, 10 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple**A grade** { 1, 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }**B grade** { }**C grade** { 3 }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Fricas****A grade** { 1, 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 19, 20, 21 }**B grade** { 8, 10, 17, 18, 22, 23 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Maxima****A grade** { 2, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18 }**B grade** { 1, 3, 8, 10 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { 19, 20, 21, 22, 23 }

Giac**A grade** { 1, 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23 }**B grade** { 8, 10, 22 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Mupad****A grade** { }**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Sympy****A grade** { }**B grade** { }**C grade** { }**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }**F(-1) timeout fail** { }**F(-2) exception fail** { }

Reduce**A grade** { }**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	64	111	84	194	58	0	80	181	83
N.S.	1	1.16	2.02	1.53	3.53	1.05	0.00	1.45	3.29	1.51
time (sec)	N/A	0.456	0.196	0.207	0.113	0.076	0.000	0.132	0.164	15.272

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	57	50	44	58	71	0	52	181	134
N.S.	1	0.84	0.74	0.65	0.85	1.04	0.00	0.76	2.66	1.97
time (sec)	N/A	0.260	0.109	0.167	0.033	0.087	0.000	0.140	0.169	15.423

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	43	62	50	94	38	0	49	61	51
N.S.	1	1.10	1.59	1.28	2.41	0.97	0.00	1.26	1.56	1.31
time (sec)	N/A	0.353	0.150	0.125	0.104	0.075	0.000	0.165	0.165	15.243

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	35	30	28	31	35	0	34	77	62
N.S.	1	0.88	0.75	0.70	0.78	0.88	0.00	0.85	1.92	1.55
time (sec)	N/A	0.235	0.042	0.113	0.030	0.088	0.000	0.135	0.172	15.222

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	17	9	9	0	9	25	25
N.S.	1	1.00	1.00	1.89	1.00	1.00	0.00	1.00	2.78	2.78
time (sec)	N/A	0.182	0.008	0.107	0.024	0.072	0.000	0.132	0.160	15.011

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	30	18	30	25	0	17	13	45
N.S.	1	1.00	2.00	1.20	2.00	1.67	0.00	1.13	0.87	3.00
time (sec)	N/A	0.209	0.110	0.138	0.107	0.081	0.000	0.159	0.164	14.951

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	14	11	13	18	19	0	14	31	36
N.S.	1	0.88	0.69	0.81	1.12	1.19	0.00	0.88	1.94	2.25
time (sec)	N/A	0.220	0.018	0.210	0.023	0.082	0.000	0.112	0.160	14.922

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	36	90	51	86	64	0	65	38	85
N.S.	1	1.16	2.90	1.65	2.77	2.06	0.00	2.10	1.23	2.74
time (sec)	N/A	0.285	0.103	0.315	0.107	0.091	0.000	0.137	0.162	15.047

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	26	26	25	29	45	0	29	53	80
N.S.	1	0.72	0.72	0.69	0.81	1.25	0.00	0.81	1.47	2.22
time (sec)	N/A	0.229	0.022	0.456	0.025	0.078	0.000	0.139	0.167	15.186

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	58	163	83	134	104	0	109	54	123
N.S.	1	1.18	3.33	1.69	2.73	2.12	0.00	2.22	1.10	2.51
time (sec)	N/A	0.380	0.112	0.632	0.118	0.088	0.000	0.138	0.173	15.241

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	44	39	37	42	70	0	41	65	113
N.S.	1	0.76	0.67	0.64	0.72	1.21	0.00	0.71	1.12	1.95
time (sec)	N/A	0.254	0.038	1.018	0.037	0.087	0.000	0.165	0.170	15.575

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	178	210	172	160	251	260	0	252	1150	445
N.S.	1	1.18	0.97	0.90	1.41	1.46	0.00	1.42	6.46	2.50
time (sec)	N/A	0.514	0.582	0.412	0.036	0.219	0.000	0.178	0.186	17.288

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	139	115	101	120	144	0	139	497	172
N.S.	1	1.23	1.02	0.89	1.06	1.27	0.00	1.23	4.40	1.52
time (sec)	N/A	0.389	0.363	0.215	0.033	0.130	0.000	0.167	0.201	16.002

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	80	56	60	50	58	0	53	110	80
N.S.	1	1.21	0.85	0.91	0.76	0.88	0.00	0.80	1.67	1.21
time (sec)	N/A	0.302	0.042	0.133	0.031	0.093	0.000	0.112	0.185	15.140

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	22	11	21	11	11	0	12	34	55
N.S.	1	1.16	0.58	1.11	0.58	0.58	0.00	0.63	1.79	2.89
time (sec)	N/A	0.195	0.009	0.108	0.024	0.096	0.000	0.148	0.196	15.590

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	42	39	43	42	47	0	44	92	75
N.S.	1	1.11	1.03	1.13	1.11	1.24	0.00	1.16	2.42	1.97
time (sec)	N/A	0.247	0.039	0.271	0.035	0.092	0.000	0.151	0.200	15.504

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	77	85	86	84	151	0	90	194	157
N.S.	1	1.07	1.18	1.19	1.17	2.10	0.00	1.25	2.69	2.18
time (sec)	N/A	0.282	0.099	0.673	0.027	0.104	0.000	0.179	0.209	15.928

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	125	132	149	149	327	0	155	311	284
N.S.	1	1.02	1.08	1.22	1.22	2.68	0.00	1.27	2.55	2.33
time (sec)	N/A	0.329	0.210	1.413	0.028	0.127	0.000	0.151	0.184	15.496

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	177	194	190	0	483	0	216	632	4468
N.S.	1	1.11	1.22	1.19	0.00	3.04	0.00	1.36	3.97	28.10
time (sec)	N/A	1.170	0.899	0.289	0.000	0.113	0.000	0.148	0.183	21.907

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	107	87	106	0	296	0	103	162	2341
N.S.	1	1.27	1.04	1.26	0.00	3.52	0.00	1.23	1.93	27.87
time (sec)	N/A	0.625	0.272	0.143	0.000	0.109	0.000	0.170	0.181	16.932

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	73	71	75	0	206	0	80	53	697
N.S.	1	1.20	1.16	1.23	0.00	3.38	0.00	1.31	0.87	11.43
time (sec)	N/A	0.601	0.071	0.223	0.000	0.112	0.000	0.162	0.182	15.177

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	107	158	141	0	453	0	163	152	2163
N.S.	1	1.18	1.74	1.55	0.00	4.98	0.00	1.79	1.67	23.77
time (sec)	N/A	0.671	0.394	0.512	0.000	0.193	0.000	0.194	0.200	16.014

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	186	186	312	251	0	852	0	299	267	4075
N.S.	1	1.00	1.68	1.35	0.00	4.58	0.00	1.61	1.44	21.91
time (sec)	N/A	0.536	0.748	1.023	0.000	0.379	0.000	0.193	0.189	16.890

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [19] had the largest ratio of [1.6153800000000004]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	12	12	1.16	13	0.923
2	A	6	5	0.84	13	0.385
3	A	9	9	1.10	13	0.692
4	A	6	5	0.88	11	0.455
5	A	4	3	1.00	11	0.273
6	A	3	3	1.00	13	0.231
7	A	6	5	0.88	13	0.385
8	A	6	6	1.16	13	0.462
9	A	6	5	0.72	13	0.385
10	A	8	8	1.18	13	0.615
11	A	6	5	0.76	13	0.385
12	A	5	4	1.18	13	0.308
13	A	5	4	1.23	13	0.308
14	A	5	4	1.21	11	0.364
15	A	6	5	1.16	11	0.455
16	A	5	4	1.11	13	0.308
17	A	5	4	1.07	13	0.308
18	A	5	4	1.02	13	0.308
19	A	22	21	1.11	13	1.615
20	A	15	14	1.27	13	1.077
21	A	15	14	1.20	13	1.077

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	13	12	1.18	13	0.923
23	A	5	5	1.00	13	0.385

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{\tan^4(x)}{a+a \csc(x)} dx$	38
3.2	$\int \frac{\tan^3(x)}{a+a \csc(x)} dx$	46
3.3	$\int \frac{\tan^2(x)}{a+a \csc(x)} dx$	52
3.4	$\int \frac{\tan(x)}{a+a \csc(x)} dx$	58
3.5	$\int \frac{\cot(x)}{a+a \csc(x)} dx$	64
3.6	$\int \frac{\cot^2(x)}{a+a \csc(x)} dx$	69
3.7	$\int \frac{\cot^3(x)}{a+a \csc(x)} dx$	74
3.8	$\int \frac{\cot^4(x)}{a+a \csc(x)} dx$	79
3.9	$\int \frac{\cot^5(x)}{a+a \csc(x)} dx$	86
3.10	$\int \frac{\cot^6(x)}{a+a \csc(x)} dx$	92
3.11	$\int \frac{\cot^7(x)}{a+a \csc(x)} dx$	99
3.12	$\int \frac{\tan^5(x)}{a+b \csc(x)} dx$	105
3.13	$\int \frac{\tan^3(x)}{a+b \csc(x)} dx$	113
3.14	$\int \frac{\tan(x)}{a+b \csc(x)} dx$	120
3.15	$\int \frac{\cot(x)}{a+b \csc(x)} dx$	126
3.16	$\int \frac{\cot^3(x)}{a+b \csc(x)} dx$	131
3.17	$\int \frac{\cot^5(x)}{a+b \csc(x)} dx$	137
3.18	$\int \frac{\cot^7(x)}{a+b \csc(x)} dx$	144
3.19	$\int \frac{\tan^4(x)}{a+b \csc(x)} dx$	151
3.20	$\int \frac{\tan^2(x)}{a+b \csc(x)} dx$	163
3.21	$\int \frac{\cot^2(x)}{a+b \csc(x)} dx$	172
3.22	$\int \frac{\cot^4(x)}{a+b \csc(x)} dx$	181

3.1 $\int \frac{\tan^4(x)}{a+a \csc(x)} dx$

Optimal result	38
Mathematica [B] (verified)	38
Rubi [A] (verified)	39
Maple [A] (verified)	41
Fricas [A] (verification not implemented)	42
Sympy [F]	42
Maxima [B] (verification not implemented)	43
Giac [A] (verification not implemented)	43
Mupad [B] (verification not implemented)	44
Reduce [B] (verification not implemented)	44

Optimal result

Integrand size = 13, antiderivative size = 55

$$\begin{aligned} \int \frac{\tan^4(x)}{a + a \csc(x)} dx &= \frac{x}{a} - \frac{(15 - 8 \csc(x)) \tan(x)}{15a} \\ &\quad + \frac{(5 - 4 \csc(x)) \tan^3(x)}{15a} - \frac{(1 - \csc(x)) \tan^5(x)}{5a} \end{aligned}$$

output $x/a-1/15*(15-8*csc(x))*tan(x)/a+1/15*(5-4*csc(x))*tan(x)^3/a-1/5*(1-csc(x))*tan(x)^5/a$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 111 vs. $2(55) = 110$.

Time = 0.20 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.02

$$\begin{aligned} \int \frac{\tan^4(x)}{a + a \csc(x)} dx \\ = \frac{200 + 6(-89 + 120x) \cos(x) + 128 \cos(2x) - 178 \cos(3x) + 240x \cos(3x) + 184 \cos(4x) - 64 \sin(x) - 176x \sin(x)}{960a (\cos(\frac{x}{2}) - \sin(\frac{x}{2}))^3 (\cos(\frac{x}{2}) + \sin(\frac{x}{2}))} \end{aligned}$$

input `Integrate[Tan[x]^4/(a + a*Csc[x]), x]`

output

$$(200 + 6*(-89 + 120*x)*\cos[x] + 128*\cos[2*x] - 178*\cos[3*x] + 240*x*\cos[3*x] + 184*\cos[4*x] - 64*\sin[x] - 178*\sin[2*x] + 240*x*\sin[2*x] - 128*\sin[3*x] - 89*\sin[4*x] + 120*x*\sin[4*x])/((960*a*(\cos[x/2] - \sin[x/2]))^3*(\cos[x/2] + \sin[x/2])^5)$$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.16, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.923, Rules used = {3042, 4376, 25, 3042, 4370, 25, 3042, 4370, 25, 3042, 4370, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^4(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cot(x)^4(a \csc(x) + a)} dx \\
 & \quad \downarrow \textcolor{blue}{4376} \\
 & \frac{\int -((a - a \csc(x)) \tan^6(x)) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & -\frac{\int (a - a \csc(x)) \tan^6(x) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{\int \frac{a - a \csc(x)}{\cot(x)^6} dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{4370} \\
 & -\frac{\frac{1}{5} \int -((5a - 4a \csc(x)) \tan^4(x)) dx + \frac{1}{5} \tan^5(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & -\frac{\frac{1}{5} \tan^5(x)(a - a \csc(x)) - \frac{1}{5} \int (5a - 4a \csc(x)) \tan^4(x) dx}{a^2}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{\frac{1}{5} \tan^5(x)(a - a \csc(x)) - \frac{1}{5} \int \frac{5a - 4a \csc(x)}{\cot(x)^4} dx}{a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{5} \left(-\frac{1}{3} \int ((15a - 8a \csc(x)) \tan^2(x)) dx - \frac{1}{3} \tan^3(x)(5a - 4a \csc(x)) \right) + \frac{1}{5} \tan^5(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\frac{1}{5} \left(\frac{1}{3} \int (15a - 8a \csc(x)) \tan^2(x) dx - \frac{1}{3} \tan^3(x)(5a - 4a \csc(x)) \right) + \frac{1}{5} \tan^5(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{1}{5} \left(\frac{1}{3} \int \frac{15a - 8a \csc(x)}{\cot(x)^2} dx - \frac{1}{3} \tan^3(x)(5a - 4a \csc(x)) \right) + \frac{1}{5} \tan^5(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \text{4370} \\
 & \frac{-\frac{1}{5} \left(\frac{1}{3} (-15adx + \tan(x)(15a - 8a \csc(x))) - \frac{1}{3} \tan^3(x)(5a - 4a \csc(x)) \right) + \frac{1}{5} \tan^5(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \text{24} \\
 & \frac{-\frac{1}{5} \tan^5(x)(a - a \csc(x)) + \frac{1}{5} \left(\frac{1}{3} (\tan(x)(15a - 8a \csc(x)) - 15ax) - \frac{1}{3} \tan^3(x)(5a - 4a \csc(x)) \right)}{a^2}
 \end{aligned}$$

input `Int[Tan[x]^4/(a + a*Csc[x]), x]`

output `-(((a - a*Csc[x])*Tan[x]^5)/5 + (-1/3*((5*a - 4*a*Csc[x])*Tan[x]^3) + (-1/5*a*x + (15*a - 8*a*Csc[x])*Tan[x])/3)/5)/a^2)`

Definitions of rubi rules used

rule 24 $\text{Int}[a_-, x_{\text{Symbol}}] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

rule 25 $\text{Int}[-(F_x_-, x_{\text{Symbol}}) \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 3042 $\text{Int}[u_-, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4370 $\text{Int}[(\cot[(c_-. + (d_-.)*(x_-)*(e_-))^m]*(\csc[(c_-. + (d_-.)*(x_-))*b_-.] + (a_-.)), x_{\text{Symbol}}) \rightarrow \text{Simp}[-(e*\text{Cot}[c + d*x])^{(m + 1)}*((a + b*\text{Csc}[c + d*x])/((d*e*(m + 1))), x] - \text{Simp}[1/(e^2*(m + 1)) \quad \text{Int}[(e*\text{Cot}[c + d*x])^{(m + 2)}*(a*(m + 1) + b*(m + 2)*\text{Csc}[c + d*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{LtQ}[m, -1]$

rule 4376 $\text{Int}[(\cot[(c_-. + (d_-.)*(x_-)*(e_-))^m]*(\csc[(c_-. + (d_-.)*(x_-))*b_-.] + (a_-.))^{(n)}, x_{\text{Symbol}}) \rightarrow \text{Simp}[a^{(2*n)}/e^{(2*n)} \quad \text{Int}[(e*\text{Cot}[c + d*x])^{(m + 2*n)/(-a + b*\text{Csc}[c + d*x])^n}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{EqQ}[a^{(2 - b^2), 0] \&& \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 0.21 (sec), antiderivative size = 84, normalized size of antiderivative = 1.53

method	result
default	$\frac{2 \arctan(\tan(\frac{x}{2})) + \frac{2}{5(\tan(\frac{x}{2})+1)^5} - \frac{1}{(\tan(\frac{x}{2})+1)^4} + \frac{1}{(\tan(\frac{x}{2})+1)^2} + \frac{11}{8(\tan(\frac{x}{2})+1)} - \frac{1}{6(\tan(\frac{x}{2})-1)^3} - \frac{1}{4(\tan(\frac{x}{2})-1)^2} + \frac{5}{8(\tan(\frac{x}{2})-1)}}{a}$
risch	$\frac{x + \frac{\frac{62 i e^{2 i x}}{15} + \frac{146 e^{3 i x}}{15} + \frac{62 e^{i x}}{15} + \frac{10 i e^{4 i x}}{3} + \frac{26 e^{5 i x}}{3} + \frac{46 i}{15} - 2 i e^{6 i x} + 2 e^{7 i x}}{(e^{i x} + i)^5 (e^{i x} - i)^3 a}}$

input $\text{int}(\tan(x)^4/(a+a*\csc(x)), x, \text{method}=\text{RETURNVERBOSE})$

output
$$\frac{64/a*(1/32*arctan(tan(1/2*x))+1/160/(tan(1/2*x)+1)^5-1/64/(tan(1/2*x)+1)^4+1/64/(tan(1/2*x)+1)^2+11/512/(tan(1/2*x)+1)-1/384/(tan(1/2*x)-1)^3-1/256/(tan(1/2*x)-1)^2+5/512/(tan(1/2*x)-1))}{}$$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.05

$$\int \frac{\tan^4(x)}{a + a \csc(x)} dx = \frac{15 x \cos(x)^3 + 23 \cos(x)^4 - 19 \cos(x)^2 + (15 x \cos(x)^3 - 8 \cos(x)^2 + 1) \sin(x) + 4}{15 (a \cos(x)^3 \sin(x) + a \cos(x)^3)}$$

input `integrate(tan(x)^4/(a+a*csc(x)),x, algorithm="fricas")`

output
$$\frac{1/15*(15*x*cos(x)^3 + 23*cos(x)^4 - 19*cos(x)^2 + (15*x*cos(x)^3 - 8*cos(x)^2 + 1)*sin(x) + 4)/(a*cos(x)^3*sin(x) + a*cos(x)^3)}{}$$

Sympy [F]

$$\int \frac{\tan^4(x)}{a + a \csc(x)} dx = \frac{\int \frac{\tan^4(x)}{\csc(x)+1} dx}{a}$$

input `integrate(tan(x)**4/(a+a*csc(x)),x)`

output `Integral(tan(x)**4/(csc(x) + 1), x)/a`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. $2(47) = 94$.

Time = 0.11 (sec) , antiderivative size = 194, normalized size of antiderivative = 3.53

$$\begin{aligned} & \int \frac{\tan^4(x)}{a + a \csc(x)} dx \\ &= \frac{2 \left(\frac{\sin(x)}{\cos(x)+1} - \frac{46 \sin(x)^2}{(\cos(x)+1)^2} - \frac{13 \sin(x)^3}{(\cos(x)+1)^3} + \frac{100 \sin(x)^4}{(\cos(x)+1)^4} + \frac{35 \sin(x)^5}{(\cos(x)+1)^5} - \frac{30 \sin(x)^6}{(\cos(x)+1)^6} - \frac{15 \sin(x)^7}{(\cos(x)+1)^7} + 8 \right)}{15 \left(a + \frac{2 a \sin(x)}{\cos(x)+1} - \frac{2 a \sin(x)^2}{(\cos(x)+1)^2} - \frac{6 a \sin(x)^3}{(\cos(x)+1)^3} + \frac{6 a \sin(x)^5}{(\cos(x)+1)^5} + \frac{2 a \sin(x)^6}{(\cos(x)+1)^6} - \frac{2 a \sin(x)^7}{(\cos(x)+1)^7} - \frac{a \sin(x)^8}{(\cos(x)+1)^8} \right)} \\ &+ \frac{2 \arctan \left(\frac{\sin(x)}{\cos(x)+1} \right)}{a} \end{aligned}$$

input `integrate(tan(x)^4/(a+a*csc(x)),x, algorithm="maxima")`

output
$$\begin{aligned} & \frac{2/15 * (\sin(x)/(\cos(x) + 1) - 46*\sin(x)^2/(\cos(x) + 1)^2 - 13*\sin(x)^3/(\cos(x) + 1)^3 + 100*\sin(x)^4/(\cos(x) + 1)^4 + 35*\sin(x)^5/(\cos(x) + 1)^5 - 30*\sin(x)^6/(\cos(x) + 1)^6 - 15*\sin(x)^7/(\cos(x) + 1)^7 + 8)/(a + 2*a*\sin(x)/(\cos(x) + 1) - 2*a*\sin(x)^2/(\cos(x) + 1)^2 - 6*a*\sin(x)^3/(\cos(x) + 1)^3 + 6*a*\sin(x)^5/(\cos(x) + 1)^5 + 2*a*\sin(x)^6/(\cos(x) + 1)^6 - 2*a*\sin(x)^7/(\cos(x) + 1)^7 - a*\sin(x)^8/(\cos(x) + 1)^8) + 2*\arctan(\sin(x)/(\cos(x) + 1))/a \end{aligned}$$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.45

$$\begin{aligned} & \int \frac{\tan^4(x)}{a + a \csc(x)} dx \\ &= \frac{x}{a} + \frac{15 \tan \left(\frac{1}{2} x \right)^2 - 36 \tan \left(\frac{1}{2} x \right) + 17}{24 a \left(\tan \left(\frac{1}{2} x \right) - 1 \right)^3} \\ &+ \frac{55 \tan \left(\frac{1}{2} x \right)^4 + 260 \tan \left(\frac{1}{2} x \right)^3 + 450 \tan \left(\frac{1}{2} x \right)^2 + 300 \tan \left(\frac{1}{2} x \right) + 71}{40 a \left(\tan \left(\frac{1}{2} x \right) + 1 \right)^5} \end{aligned}$$

input `integrate(tan(x)^4/(a+a*csc(x)),x, algorithm="giac")`

output

$$\frac{x/a + 1/24*(15*tan(1/2*x)^2 - 36*tan(1/2*x) + 17)/(a*(tan(1/2*x) - 1)^3) + 1/40*(55*tan(1/2*x)^4 + 260*tan(1/2*x)^3 + 450*tan(1/2*x)^2 + 300*tan(1/2*x) + 71)/(a*(tan(1/2*x) + 1)^5)}{}$$

Mupad [B] (verification not implemented)

Time = 15.27 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.51

$$\int \frac{\tan^4(x)}{a + a \csc(x)} dx = \frac{x}{a} - \frac{-2 \tan\left(\frac{x}{2}\right)^7 - 4 \tan\left(\frac{x}{2}\right)^6 + \frac{14 \tan\left(\frac{x}{2}\right)^5}{3} + \frac{40 \tan\left(\frac{x}{2}\right)^4}{3} - \frac{26 \tan\left(\frac{x}{2}\right)^3}{15} - \frac{92 \tan\left(\frac{x}{2}\right)^2}{15} + \frac{2 \tan\left(\frac{x}{2}\right)}{15} + \frac{16}{15}}{a \left(\tan\left(\frac{x}{2}\right) - 1\right)^3 \left(\tan\left(\frac{x}{2}\right) + 1\right)^5}$$

input

```
int(tan(x)^4/(a + a/sin(x)),x)
```

output

$$\frac{x/a - ((2*tan(x/2))/15 - (92*tan(x/2)^2)/15 - (26*tan(x/2)^3)/15 + (40*tan(x/2)^4)/3 + (14*tan(x/2)^5)/3 - 4*tan(x/2)^6 - 2*tan(x/2)^7 + 16/15)/(a*(tan(x/2) - 1)^3*(tan(x/2) + 1)^5)}{}$$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 181, normalized size of antiderivative = 3.29

$$\int \frac{\tan^4(x)}{a + a \csc(x)} dx = \frac{5 \cos(x) \sin(x)^3 \tan(x)^3 - 15 \cos(x) \sin(x)^3 \tan(x) + 15 \cos(x) \sin(x)^3 x + 8 \cos(x) \sin(x)^3 + 5 \cos(x) \sin(x)^3 \tan(x)^3}{a}$$

input

```
int(tan(x)^4/(a+a*csc(x)),x)
```

output

```
(5*cos(x)*sin(x)**3*tan(x)**3 - 15*cos(x)*sin(x)**3*tan(x) + 15*cos(x)*sin(x)**3*x + 8*cos(x)*sin(x)**3 + 5*cos(x)*sin(x)**2*tan(x)**3 - 15*cos(x)*sin(x)**2*tan(x)**2*x + 15*cos(x)*sin(x)**2*x + 8*cos(x)*sin(x)**2 - 5*cos(x)*sin(x)*tan(x)**3 + 15*cos(x)*sin(x)*tan(x) - 15*cos(x)*sin(x)*x - 8*cos(x)*sin(x) - 5*cos(x)*tan(x)**3 + 15*cos(x)*tan(x) - 15*cos(x)*x - 8*cos(x) - 3*sin(x)**4 + 12*sin(x)**3 + 12*sin(x)**2 - 8*sin(x) - 8)/(15*cos(x)*a*(sin(x)**3 + sin(x)**2 - sin(x) - 1))
```

3.2 $\int \frac{\tan^3(x)}{a+a \csc(x)} dx$

Optimal result	46
Mathematica [A] (verified)	46
Rubi [A] (verified)	47
Maple [A] (verified)	48
Fricas [A] (verification not implemented)	49
Sympy [F]	49
Maxima [A] (verification not implemented)	50
Giac [A] (verification not implemented)	50
Mupad [B] (verification not implemented)	51
Reduce [B] (verification not implemented)	51

Optimal result

Integrand size = 13, antiderivative size = 68

$$\begin{aligned} \int \frac{\tan^3(x)}{a + a \csc(x)} dx = & \frac{5 \log(1 - \sin(x))}{16a} + \frac{11 \log(1 + \sin(x))}{16a} \\ & + \frac{1}{8a(1 - \sin(x))} - \frac{1}{8a(1 + \sin(x))^2} + \frac{3}{4a(1 + \sin(x))} \end{aligned}$$

output
$$\frac{5/16 \ln(1-\sin(x))/a + 11/16 \ln(1+\sin(x))/a + 1/8/a/(1-\sin(x)) - 1/8/a/(1+\sin(x))}{\sim 2+3/4/a/(1+\sin(x))}$$

Mathematica [A] (verified)

Time = 0.11 (sec), antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx = \frac{5 \log(1 - \sin(x)) + 11 \log(1 + \sin(x)) + \frac{2(-6 - 3 \sin(x) + 5 \sin^2(x))}{(-1 + \sin(x))(1 + \sin(x))^2}}{16a}$$

input `Integrate[Tan[x]^3/(a + a*Csc[x]),x]`

output
$$\frac{(5 \operatorname{Log}[1 - \operatorname{Sin}[x]] + 11 \operatorname{Log}[1 + \operatorname{Sin}[x]] + (2 (-6 - 3 \operatorname{Sin}[x] + 5 \operatorname{Sin}[x]^2))}{((-1 + \operatorname{Sin}[x]) (1 + \operatorname{Sin}[x])^2)/(16 a)}$$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.84, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 4367, 27, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cot(x)^3(a \csc(x) + a)} dx \\
 & \quad \downarrow \textcolor{blue}{4367} \\
 & a^4 \int \frac{\sin^4(x)}{a^5(1 - \sin(x))^2(\sin(x) + 1)^3} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{27} \\
 & \frac{\int \frac{\sin^4(x)}{(1 - \sin(x))^2(\sin(x) + 1)^3} d\sin(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{99} \\
 & \frac{\int \left(\frac{11}{16(\sin(x) + 1)} - \frac{3}{4(\sin(x) + 1)^2} + \frac{1}{4(\sin(x) + 1)^3} + \frac{5}{16(\sin(x) - 1)} + \frac{1}{8(\sin(x) - 1)^2} \right) d\sin(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{\frac{1}{8(1 - \sin(x))} + \frac{3}{4(\sin(x) + 1)} - \frac{1}{8(\sin(x) + 1)^2} + \frac{5}{16} \log(1 - \sin(x)) + \frac{11}{16} \log(\sin(x) + 1)}{a}
 \end{aligned}$$

input `Int[Tan[x]^3/(a + a*Csc[x]),x]`

output `((5*Log[1 - Sin[x]])/16 + (11*Log[1 + Sin[x]])/16 + 1/(8*(1 - Sin[x])) - 1/(8*(1 + Sin[x])^2) + 3/(4*(1 + Sin[x]))) / a`

Definitions of rubi rules used

rule 27 $\text{Int}[(a_*)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& !\text{MatchQ}[F_x, (b_*)*(G_x_) /; \text{FreeQ}[b, x]]$

rule 99 $\text{Int}[(a_*) + (b_*)*(x_*)^m * ((c_*) + (d_*)*(x_*)^n * ((e_*) + (f_*)*(x_*)^p), x_*) \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, p\}, x] \&& \text{IntegersQ}[m, n] \&& (\text{IntegerQ}[p] \mid (\text{GtQ}[m, 0] \&& \text{GeQ}[n, -1]))]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4367 $\text{Int}[\cot[(c_*) + (d_*)*(x_*)]^m * (\csc[(c_*) + (d_*)*(x_*)]*((b_*) + (a_*)^n), x_Symbol) \rightarrow \text{Simp}[1/(a^{(m-n-1)*b^n*d}) \text{ Subst}[\text{Int}[(a - b*x)^{(m-1)/2} * ((a + b*x)^{(m-1)/2+n}/x^{(m+n)}), x], x, \text{Sin}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{IntegerQ}[(m-1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{-\frac{1}{8(1+\sin(x))^2} + \frac{3}{4(1+\sin(x))} + \frac{\frac{11 \ln(1+\sin(x))}{16}}{a} - \frac{1}{8(\sin(x)-1)} + \frac{5 \ln(\sin(x)-1)}{16}}{a}$	44
risch	$-\frac{ix}{a} + \frac{i(6ie^{2ix}+5e^{ix}-6ie^{4ix}+14e^{3ix}+5e^{5ix})}{4(e^{ix}-i)^2(e^{ix}+i)^4a} + \frac{5\ln(e^{ix}-i)}{8a} + \frac{11\ln(e^{ix}+i)}{8a}$	101

input $\text{int}(\tan(x)^3/(a+a*\csc(x)), x, \text{method}=\text{_RETURNVERBOSE})$

output $\frac{1}{a} \left(-\frac{1}{8} \right) \left(\frac{1}{1 + \sin(x)} \right)^2 + \frac{3}{4} \left(\frac{1}{1 + \sin(x)} \right) + \frac{11}{16} \ln(1 + \sin(x)) - \frac{1}{8} \left(\frac{1}{\sin(x) - 1} \right) + \frac{5}{16} \ln(\sin(x) - 1)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 71, normalized size of antiderivative = 1.04

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx \\ = \frac{10 \cos(x)^2 + 11 (\cos(x)^2 \sin(x) + \cos(x)^2) \log(\sin(x) + 1) + 5 (\cos(x)^2 \sin(x) + \cos(x)^2) \log(-\sin(x) + 1)}{16 (a \cos(x)^2 \sin(x) + a \cos(x)^2)}$$

input `integrate(tan(x)^3/(a+a*csc(x)),x, algorithm="fricas")`

output $\frac{1}{16} (10 \cos(x)^2 + 11 (\cos(x)^2 \sin(x) + \cos(x)^2) \log(\sin(x) + 1) + 5 (\cos(x)^2 \sin(x) + \cos(x)^2) \log(-\sin(x) + 1) + 6 \sin(x) + 2) / (a \cos(x)^2 \sin(x) + a \cos(x)^2)$

Sympy [F]

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx = \frac{\int \frac{\tan^3(x)}{\csc(x)+1} dx}{a}$$

input `integrate(tan(x)**3/(a+a*csc(x)),x)`

output `Integral(tan(x)**3/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.85

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx = \frac{5 \sin(x)^2 - 3 \sin(x) - 6}{8(a \sin(x)^3 + a \sin(x)^2 - a \sin(x) - a)} + \frac{11 \log(\sin(x) + 1)}{16 a} + \frac{5 \log(\sin(x) - 1)}{16 a}$$

input `integrate(tan(x)^3/(a+a*csc(x)),x, algorithm="maxima")`

output `1/8*(5*sin(x)^2 - 3*sin(x) - 6)/(a*sin(x)^3 + a*sin(x)^2 - a*sin(x) - a) + 11/16*log(sin(x) + 1)/a + 5/16*log(sin(x) - 1)/a`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.76

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx = \frac{11 \log(\sin(x) + 1)}{16 a} + \frac{5 \log(-\sin(x) + 1)}{16 a} + \frac{5 \sin(x)^2 - 3 \sin(x) - 6}{8 a (\sin(x) + 1)^2 (\sin(x) - 1)}$$

input `integrate(tan(x)^3/(a+a*csc(x)),x, algorithm="giac")`

output `11/16*log(sin(x) + 1)/a + 5/16*log(-sin(x) + 1)/a + 1/8*(5*sin(x)^2 - 3*sin(x) - 6)/(a*(sin(x) + 1)^2*(sin(x) - 1))`

Mupad [B] (verification not implemented)

Time = 15.42 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.97

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx = \frac{5 \ln(\tan(\frac{x}{2}) - 1)}{8a} + \frac{11 \ln(\tan(\frac{x}{2}) + 1)}{8a} - \frac{\ln(\tan(\frac{x}{2})^2 + 1)}{a} \\ + \frac{-\frac{3 \tan(\frac{x}{2})^5}{4} + \frac{\tan(\frac{x}{2})^4}{2} + \frac{9 \tan(\frac{x}{2})^3}{2} + \frac{\tan(\frac{x}{2})^2}{2} - \frac{3 \tan(\frac{x}{2})}{4}}{a \tan(\frac{x}{2})^6 + 2a \tan(\frac{x}{2})^5 - a \tan(\frac{x}{2})^4 - 4a \tan(\frac{x}{2})^3 - a \tan(\frac{x}{2})^2 + 2a \tan(\frac{x}{2}) + a}$$

input `int(tan(x)^3/(a + a/sin(x)),x)`

output
$$\frac{(5 \log(\tan(x/2) - 1))}{(8*a)} + \frac{(11 \log(\tan(x/2) + 1))}{(8*a)} - \log(\tan(x/2)^2 + 1)/a + (\tan(x/2)^2/2 - (3 \tan(x/2))/4 + (9 \tan(x/2)^3)/2 + \tan(x/2)^4/2 - (3 \tan(x/2)^5)/4)/(a + 2*a*\tan(x/2) - a*\tan(x/2)^2 - 4*a*\tan(x/2)^3 - a*\tan(x/2)^4 + 2*a*\tan(x/2)^5 + a*\tan(x/2)^6)$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 181, normalized size of antiderivative = 2.66

$$\int \frac{\tan^3(x)}{a + a \csc(x)} dx = -\frac{-8 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^3 - 8 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^2 + 8 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x) + 8 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a}$$

input int(tan(x)^3/(a+a*csc(x)),x)

```

output (- 8*log(tan(x/2)**2 + 1)*sin(x)**3 - 8*log(tan(x/2)**2 + 1)*sin(x)**2 +
8*log(tan(x/2)**2 + 1)*sin(x) + 8*log(tan(x/2)**2 + 1) + 5*log(tan(x/2) -
1)*sin(x)**3 + 5*log(tan(x/2) - 1)*sin(x)**2 - 5*log(tan(x/2) - 1)*sin(x) -
5*log(tan(x/2) - 1) + 11*log(tan(x/2) + 1)*sin(x)**3 + 11*log(tan(x/2) +
1)*sin(x)**2 - 11*log(tan(x/2) + 1)*sin(x) - 11*log(tan(x/2) + 1) - 3*sin(
x)**3 + 2*sin(x)**2 - 3)/(8*a*(sin(x)**3 + sin(x)**2 - sin(x) - 1))

```

3.3 $\int \frac{\tan^2(x)}{a+a \csc(x)} dx$

Optimal result	52
Mathematica [A] (verified)	52
Rubi [A] (verified)	53
Maple [C] (verified)	55
Fricas [A] (verification not implemented)	55
Sympy [F]	56
Maxima [B] (verification not implemented)	56
Giac [A] (verification not implemented)	56
Mupad [B] (verification not implemented)	57
Reduce [B] (verification not implemented)	57

Optimal result

Integrand size = 13, antiderivative size = 39

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = -\frac{x}{a} + \frac{(3 - 2 \csc(x)) \tan(x)}{3a} - \frac{(1 - \csc(x)) \tan^3(x)}{3a}$$

output
$$-\frac{x}{a} + \frac{(3 - 2 \csc(x)) \tan(x)}{3a} - \frac{(1 - \csc(x)) \tan^3(x)}{3a}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.59

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = -\frac{4 \cos(2x) - 2 \sin(x) + (-5 + 6x) \cos(x)(1 + \sin(x))}{6a (\cos(\frac{x}{2}) - \sin(\frac{x}{2})) (\cos(\frac{x}{2}) + \sin(\frac{x}{2}))^3}$$

input `Integrate[Tan[x]^2/(a + a*Csc[x]), x]`

output
$$-\frac{1}{6}(4 \cos(2x) - 2 \sin(x) + (-5 + 6x) \cos(x)(1 + \sin(x))) / (a (\cos(\frac{x}{2}) - \sin(\frac{x}{2})) (\cos(\frac{x}{2}) + \sin(\frac{x}{2}))^3)$$

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {3042, 4376, 25, 3042, 4370, 25, 3042, 4370, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cot(x)^2(a \csc(x) + a)} dx \\
 & \quad \downarrow \textcolor{blue}{4376} \\
 & \frac{\int -((a - a \csc(x)) \tan^4(x)) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & -\frac{\int (a - a \csc(x)) \tan^4(x) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{\int \frac{a-a \csc(x)}{\cot(x)^4} dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{4370} \\
 & -\frac{\frac{1}{3} \int -((3a - 2a \csc(x)) \tan^2(x)) dx + \frac{1}{3} \tan^3(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & -\frac{\frac{1}{3} \tan^3(x)(a - a \csc(x)) - \frac{1}{3} \int (3a - 2a \csc(x)) \tan^2(x) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{\frac{1}{3} \tan^3(x)(a - a \csc(x)) - \frac{1}{3} \int \frac{3a-2a \csc(x)}{\cot(x)^2} dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{4370}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{\frac{1}{3}(-\int -3adx - (\tan(x)(3a - 2a \csc(x)))) + \frac{1}{3}\tan^3(x)(a - a \csc(x))}{a^2} \\
 & \quad \downarrow \text{24} \\
 & -\frac{\frac{1}{3}\tan^3(x)(a - a \csc(x)) + \frac{1}{3}(3ax - \tan(x)(3a - 2a \csc(x)))}{a^2}
 \end{aligned}$$

input `Int[Tan[x]^2/(a + a*Csc[x]), x]`

output `-(((a - a*Csc[x])*Tan[x]^3)/3 + (3*a*x - (3*a - 2*a*Csc[x])*Tan[x])/3)/a^2`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4370 `Int[(cot[(c_.) + (d_)*(x_)]*(e_.))^m_*(csc[(c_.) + (d_)*(x_)]*(b_.) + (a_)), x_Symbol] :> Simp[(-(e*Cot[c + d*x])^(m + 1))*(a + b*Csc[c + d*x])/((d*e*(m + 1))), x] - Simp[1/(e^2*(m + 1)) Int[(e*Cot[c + d*x])^(m + 2)*(a*(m + 1) + b*(m + 2)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e}, x] && LtQ[m, -1]`

rule 4376 `Int[(cot[(c_.) + (d_)*(x_)]*(e_.))^m_*(csc[(c_.) + (d_)*(x_)]*(b_.) + (a_))^(n_), x_Symbol] :> Simp[a^(2*n)/e^(2*n) Int[(e*Cot[c + d*x])^(m + 2*n)/(-a + b*Csc[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[a^2 - b^2, 0] && ILtQ[n, 0]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.28

method	result	size
risch	$-\frac{x}{a} - \frac{2(5e^{ix} + 4i + 3e^{3ix})}{3(e^{ix} + i)^3(e^{ix} - i)a}$	50
default	$\frac{-2\arctan(\tan(\frac{x}{2})) + \frac{2}{3(\tan(\frac{x}{2})+1)^3} - \frac{1}{(\tan(\frac{x}{2})+1)^2} - \frac{3}{2(\tan(\frac{x}{2})+1)} - \frac{1}{2(\tan(\frac{x}{2})-1)}}{a}$	54

input `int(tan(x)^2/(a+a*csc(x)),x,method=_RETURNVERBOSE)`

output `-x/a-2/3*(5*exp(I*x)+4*I+3*exp(3*I*x))/(exp(I*x)+I)^3/(exp(I*x)-I)/a`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.97

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = -\frac{3 x \cos(x) + 4 \cos(x)^2 + (3 x \cos(x) - 1) \sin(x) - 2}{3 (a \cos(x) \sin(x) + a \cos(x))}$$

input `integrate(tan(x)^2/(a+a*csc(x)),x, algorithm="fricas")`

output `-1/3*(3*x*cos(x) + 4*cos(x)^2 + (3*x*cos(x) - 1)*sin(x) - 2)/(a*cos(x)*sin(x) + a*cos(x))`

Sympy [F]

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = \frac{\int \frac{\tan^2(x)}{\csc(x)+1} dx}{a}$$

input `integrate(tan(x)**2/(a+a*csc(x)),x)`

output `Integral(tan(x)**2/(csc(x) + 1), x)/a`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(33) = 66$.

Time = 0.10 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.41

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = -\frac{2 \left(\frac{\sin(x)}{\cos(x)+1} - \frac{6 \sin(x)^2}{(\cos(x)+1)^2} - \frac{3 \sin(x)^3}{(\cos(x)+1)^3} + 2 \right)}{3 \left(a + \frac{2 a \sin(x)}{\cos(x)+1} - \frac{2 a \sin(x)^3}{(\cos(x)+1)^3} - \frac{a \sin(x)^4}{(\cos(x)+1)^4} \right)} - \frac{2 \arctan \left(\frac{\sin(x)}{\cos(x)+1} \right)}{a}$$

input `integrate(tan(x)^2/(a+a*csc(x)),x, algorithm="maxima")`

output `-2/3*(sin(x)/(cos(x) + 1) - 6*sin(x)^2/(cos(x) + 1)^2 - 3*sin(x)^3/(cos(x) + 1)^3 + 2)/(a + 2*a*sin(x)/(cos(x) + 1) - 2*a*sin(x)^3/(cos(x) + 1)^3 - a*sin(x)^4/(cos(x) + 1)^4) - 2*arctan(sin(x)/(cos(x) + 1))/a`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.26

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = -\frac{x}{a} - \frac{1}{2 a (\tan(\frac{1}{2} x) - 1)} - \frac{9 \tan(\frac{1}{2} x)^2 + 24 \tan(\frac{1}{2} x) + 11}{6 a (\tan(\frac{1}{2} x) + 1)^3}$$

input `integrate(tan(x)^2/(a+a*csc(x)),x, algorithm="giac")`

output
$$\frac{-x/a - 1/2/(a*(\tan(1/2*x) - 1)) - 1/6*(9*\tan(1/2*x)^2 + 24*\tan(1/2*x) + 11)}{(a*(\tan(1/2*x) + 1)^3)}$$

Mupad [B] (verification not implemented)

Time = 15.24 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.31

$$\int \frac{\tan^2(x)}{a + a \csc(x)} dx = \frac{-2 \tan(\frac{x}{2})^3 - 4 \tan(\frac{x}{2})^2 + \frac{2 \tan(\frac{x}{2})}{3} + \frac{4}{3}}{a (\tan(\frac{x}{2}) - 1) (\tan(\frac{x}{2}) + 1)^3} - \frac{x}{a}$$

input `int(tan(x)^2/(a + a/sin(x)),x)`

output
$$\frac{((2*\tan(x/2))/3 - 4*\tan(x/2)^2 - 2*\tan(x/2)^3 + 4/3)/(a*(\tan(x/2) - 1)*(ta n(x/2) + 1)^3) - x/a}{3}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.56

$$\begin{aligned} & \int \frac{\tan^2(x)}{a + a \csc(x)} dx \\ &= \frac{3 \cos(x) \sin(x) \tan(x) - 3 \cos(x) \sin(x) x - 2 \cos(x) \sin(x) + 3 \cos(x) \tan(x) - 3 \cos(x) x - 2 \cos(x)}{3 \cos(x) a (\sin(x) + 1)} \end{aligned}$$

input `int(tan(x)^2/(a+a*csc(x)),x)`

output
$$\frac{(3*\cos(x)*\sin(x)*\tan(x) - 3*\cos(x)*\sin(x)*x - 2*\cos(x)*\sin(x) + 3*\cos(x)*ta n(x) - 3*\cos(x)*x - 2*\cos(x) + \sin(x)**2 - 2*\sin(x) - 2)/(3*\cos(x)*a*(\sin(x) + 1))}{3}$$

3.4 $\int \frac{\tan(x)}{a+a \csc(x)} dx$

Optimal result	58
Mathematica [A] (verified)	58
Rubi [A] (verified)	59
Maple [A] (verified)	60
Fricas [A] (verification not implemented)	61
Sympy [F]	61
Maxima [A] (verification not implemented)	61
Giac [A] (verification not implemented)	62
Mupad [B] (verification not implemented)	62
Reduce [B] (verification not implemented)	62

Optimal result

Integrand size = 11, antiderivative size = 40

$$\int \frac{\tan(x)}{a + a \csc(x)} dx = -\frac{\log(1 - \sin(x))}{4a} - \frac{3 \log(1 + \sin(x))}{4a} - \frac{1}{2a(1 + \sin(x))}$$

output -1/4*ln(1-sin(x))/a-3/4*ln(1+sin(x))/a-1/2/a/(1+sin(x))

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.75

$$\int \frac{\tan(x)}{a + a \csc(x)} dx = -\frac{\log(1 - \sin(x)) + 3 \log(1 + \sin(x)) + \frac{2}{1+\sin(x)}}{4a}$$

input Integrate[Tan[x]/(a + a*Csc[x]),x]

output -1/4*(Log[1 - Sin[x]] + 3*Log[1 + Sin[x]] + 2/(1 + Sin[x]))/a

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4367, 27, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cot(x)(a \csc(x) + a)} dx \\
 & \quad \downarrow \textcolor{blue}{4367} \\
 & a^2 \int \frac{\sin^2(x)}{a^3(1 - \sin(x))(\sin(x) + 1)^2} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{27} \\
 & \int \frac{\sin^2(x)}{(1 - \sin(x))(\sin(x) + 1)^2} d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{a} \\
 & \int \left(-\frac{3}{4(\sin(x) + 1)} + \frac{1}{2(\sin(x) + 1)^2} - \frac{1}{4(\sin(x) - 1)} \right) d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{99} \\
 & \int \left(-\frac{1}{2(\sin(x) + 1)} - \frac{1}{4} \log(1 - \sin(x)) - \frac{3}{4} \log(\sin(x) + 1) \right) d\sin(x) \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{1}{2(\sin(x) + 1)} - \frac{1}{4} \log(1 - \sin(x)) - \frac{3}{4} \log(\sin(x) + 1)
 \end{aligned}$$

input `Int[Tan[x]/(a + a*Csc[x]),x]`

output `(-1/4*Log[1 - Sin[x]] - (3*Log[1 + Sin[x]])/4 - 1/(2*(1 + Sin[x])))/a`

Definitions of rubi rules used

rule 27 $\text{Int}[(a_*)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[F_x, (b_*)*(G_x_) /; \text{FreeQ}[b, x]]$

rule 99 $\text{Int}[(a_*) + (b_*)*(x_*)^m * ((c_*) + (d_*)*(x_*)^n * ((e_*) + (f_*)*(x_*)^p), x_*) \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, p\}, x] \&& \text{IntegersQ}[m, n] \&& (\text{IntegerQ}[p] \mid (\text{GtQ}[m, 0] \&& \text{GeQ}[n, -1]))]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4367 $\text{Int}[\cot[(c_*) + (d_*)*(x_*)]^m * (\csc[(c_*) + (d_*)*(x_*)]*((b_*) + (a_*))^n, x_Symbol] \rightarrow \text{Simp}[1/(a^{(m-n-1)*b^n*d}) \text{ Subst}[\text{Int}[(a - b*x)^{(m-1)/2} * ((a + b*x)^{(m-1)/2+n}/x^{(m+n)}), x], x, \text{Sin}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{IntegerQ}[(m-1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{-\frac{1}{2(1+\sin(x))}-\frac{3 \ln(1+\sin(x))}{4}-\frac{\ln(\sin(x)-1)}{4}}{a}$	28
risch	$\frac{ix}{a}-\frac{ie^{ix}}{(e^{ix}+i)^2 a}-\frac{\ln(e^{ix}-i)}{2 a}-\frac{3 \ln(e^{ix}+i)}{2 a}$	58

input `int(tan(x)/(a+a*csc(x)),x,method=_RETURNVERBOSE)`

output `1/a*(-1/2/(1+sin(x))-3/4*ln(1+sin(x))-1/4*ln(sin(x)-1))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88

$$\int \frac{\tan(x)}{a + a \csc(x)} dx \\ = -\frac{3 (\sin(x) + 1) \log(\sin(x) + 1) + (\sin(x) + 1) \log(-\sin(x) + 1) + 2}{4 (a \sin(x) + a)}$$

input `integrate(tan(x)/(a+a*csc(x)),x, algorithm="fricas")`

output `-1/4*(3*(sin(x) + 1)*log(sin(x) + 1) + (sin(x) + 1)*log(-sin(x) + 1) + 2)/(a*sin(x) + a)`

Sympy [F]

$$\int \frac{\tan(x)}{a + a \csc(x)} dx = \frac{\int \frac{\tan(x)}{\csc(x)+1} dx}{a}$$

input `integrate(tan(x)/(a+a*csc(x)),x)`

output `Integral(tan(x)/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.78

$$\int \frac{\tan(x)}{a + a \csc(x)} dx = -\frac{3 \log(\sin(x) + 1)}{4 a} - \frac{\log(\sin(x) - 1)}{4 a} - \frac{1}{2 (a \sin(x) + a)}$$

input `integrate(tan(x)/(a+a*csc(x)),x, algorithm="maxima")`

output `-3/4*log(sin(x) + 1)/a - 1/4*log(sin(x) - 1)/a - 1/2/(a*sin(x) + a)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int \frac{\tan(x)}{a + a \csc(x)} dx = -\frac{3 \log(\sin(x) + 1)}{4a} - \frac{\log(-\sin(x) + 1)}{4a} - \frac{1}{2a(\sin(x) + 1)}$$

input `integrate(tan(x)/(a+a*csc(x)),x, algorithm="giac")`

output `-3/4*log(sin(x) + 1)/a - 1/4*log(-sin(x) + 1)/a - 1/2/(a*(sin(x) + 1))`

Mupad [B] (verification not implemented)

Time = 15.22 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.55

$$\begin{aligned} \int \frac{\tan(x)}{a + a \csc(x)} dx &= \frac{\tan(\frac{x}{2})}{a \tan^2(\frac{x}{2}) + 2a \tan(\frac{x}{2}) + a} - \frac{3 \ln(\tan(\frac{x}{2}) + 1)}{2a} \\ &\quad - \frac{\ln(\tan(\frac{x}{2}) - 1)}{2a} + \frac{\ln(\tan(\frac{x}{2})^2 + 1)}{a} \end{aligned}$$

input `int(tan(x)/(a + a/sin(x)),x)`

output `tan(x/2)/(a + 2*a*tan(x/2) + a*tan(x/2)^2) - (3*log(tan(x/2) + 1))/(2*a) - log(tan(x/2) - 1)/(2*a) + log(tan(x/2)^2 + 1)/a`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.92

$$\begin{aligned} \int \frac{\tan(x)}{a + a \csc(x)} dx \\ = \frac{2 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x) + 2 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) - \log\left(\tan\left(\frac{x}{2}\right) - 1\right) \sin(x) - \log\left(\tan\left(\frac{x}{2}\right) - 1\right) - 3 \log\left(\sin(x) + 1\right)}{2a(\sin(x) + 1)} \end{aligned}$$

input `int(tan(x)/(a+a*csc(x)),x)`

output
$$\frac{(2\log(\tan(x/2)^{**2} + 1)*\sin(x) + 2*\log(\tan(x/2)^{**2} + 1) - \log(\tan(x/2) - 1)*\sin(x) - \log(\tan(x/2) - 1) - 3*\log(\tan(x/2) + 1)*\sin(x) - 3*\log(\tan(x/2) + 1) - 1)/(2*a*(\sin(x) + 1))}{}$$

3.5 $\int \frac{\cot(x)}{a+a \csc(x)} dx$

Optimal result	64
Mathematica [A] (verified)	64
Rubi [A] (verified)	65
Maple [A] (verified)	66
Fricas [A] (verification not implemented)	66
Sympy [F]	67
Maxima [A] (verification not implemented)	67
Giac [A] (verification not implemented)	67
Mupad [B] (verification not implemented)	68
Reduce [B] (verification not implemented)	68

Optimal result

Integrand size = 11, antiderivative size = 9

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{\log(1 + \sin(x))}{a}$$

output `ln(1+sin(x))/a`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{\log(1 + \sin(x))}{a}$$

input `Integrate[Cot[x]/(a + a*Csc[x]),x]`

output `Log[1 + Sin[x]]/a`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4367, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{4367} \\
 & \int \frac{1}{a \sin(x) + a} d \sin(x) \\
 & \quad \downarrow \textcolor{blue}{16} \\
 & \frac{\log(\sin(x) + 1)}{a}
 \end{aligned}$$

input `Int[Cot[x]/(a + a*Csc[x]),x]`

output `Log[1 + Sin[x]]/a`

Definitions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4367 $\text{Int}[\cot[(c_{_}) + (d_{_})*(x_{_})]^{(m_{_})}*(\csc[(c_{_}) + (d_{_})*(x_{_})]*(b_{_}) + (a_{_}))^{(n_{_})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(a^{(m - n - 1)}*b^{n*d}) \text{Subst}[\text{Int}[(a - b*x)^{(m - 1)/2}*(a + b*x)^{((m - 1)/2 + n)}/x^{(m + n)}, x], x, \text{Sin}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \& \text{IntegerQ}[(m - 1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Maple [A] (verified)

Time = 0.11 (sec), antiderivative size = 17, normalized size of antiderivative = 1.89

method	result	size
derivativedivides	$-\frac{-\ln(\csc(x)+1)+\ln(\csc(x))}{a}$	17
default	$-\frac{-\ln(\csc(x)+1)+\ln(\csc(x))}{a}$	17
risch	$-\frac{ix}{a} + \frac{2\ln(e^{ix}+i)}{a}$	23

input `int(cot(x)/(a+a*csc(x)),x,method=_RETURNVERBOSE)`

output `-1/a*(-ln(csc(x)+1)+ln(csc(x)))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec), antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{\log(\sin(x) + 1)}{a}$$

input `integrate(cot(x)/(a+a*csc(x)),x, algorithm="fricas")`

output `log(sin(x) + 1)/a`

Sympy [F]

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot(x)}{\csc(x)+1} dx}{a}$$

input `integrate(cot(x)/(a+a*csc(x)),x)`

output `Integral(cot(x)/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{\log(\sin(x) + 1)}{a}$$

input `integrate(cot(x)/(a+a*csc(x)),x, algorithm="maxima")`

output `log(sin(x) + 1)/a`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{\log(\sin(x) + 1)}{a}$$

input `integrate(cot(x)/(a+a*csc(x)),x, algorithm="giac")`

output `log(sin(x) + 1)/a`

Mupad [B] (verification not implemented)

Time = 15.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.78

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{2 \ln(\tan(\frac{x}{2}) + 1) - \ln(\tan(\frac{x}{2})^2 + 1)}{a}$$

input `int(cot(x)/(a + a/sin(x)),x)`

output `(2*log(tan(x/2) + 1) - log(tan(x/2)^2 + 1))/a`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.78

$$\int \frac{\cot(x)}{a + a \csc(x)} dx = \frac{-\log(\tan(\frac{x}{2})^2 + 1) + 2 \log(\tan(\frac{x}{2}) + 1)}{a}$$

input `int(cot(x)/(a+a*csc(x)),x)`

output `(- log(tan(x/2)**2 + 1) + 2*log(tan(x/2) + 1))/a`

3.6 $\int \frac{\cot^2(x)}{a+a \csc(x)} dx$

Optimal result	69
Mathematica [A] (verified)	69
Rubi [A] (verified)	70
Maple [A] (verified)	71
Fricas [A] (verification not implemented)	71
Sympy [F]	72
Maxima [A] (verification not implemented)	72
Giac [A] (verification not implemented)	72
Mupad [B] (verification not implemented)	73
Reduce [B] (verification not implemented)	73

Optimal result

Integrand size = 13, antiderivative size = 15

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = -\frac{x}{a} - \frac{\operatorname{arctanh}(\cos(x))}{a}$$

output $-\frac{x}{a} - \frac{\operatorname{arctanh}(\cos(x))}{a}$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.00

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = -\frac{x}{a} - \frac{\log(\cos(\frac{x}{2}))}{a} + \frac{\log(\sin(\frac{x}{2}))}{a}$$

input `Integrate[Cot[x]^2/(a + a*Csc[x]),x]`

output $-\frac{x}{a} - \frac{\log(\cos(\frac{x}{2}))}{a} + \frac{\log(\sin(\frac{x}{2}))}{a}$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4376, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^2(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^2}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{4376} \\
 & \frac{\int (a \csc(x) - a) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{-a \operatorname{arctanh}(\cos(x)) - ax}{a^2}
 \end{aligned}$$

input `Int[Cot[x]^2/(a + a*Csc[x]),x]`

output `(-(a*x) - a*ArcTanh[Cos[x]])/a^2`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4376

```
Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^(m_)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] :> Simp[a^(2*n)/e^(2*n) Int[(e*Cot[c + d*x])^(m + 2*n)/(-a + b*Csc[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[a^2 - b^2, 0] && ILtQ[n, 0]
```

Maple [A] (verified)

Time = 0.14 (sec), antiderivative size = 18, normalized size of antiderivative = 1.20

method	result	size
default	$\frac{\ln(\tan(\frac{x}{2})) - 2 \arctan(\tan(\frac{x}{2}))}{a}$	18
risch	$-\frac{x}{a} - \frac{\ln(e^{ix} + 1)}{a} + \frac{\ln(e^{ix} - 1)}{a}$	33

input `int(cot(x)^2/(a+a*csc(x)),x,method=_RETURNVERBOSE)`

output `1/a*(ln(tan(1/2*x))-2*arctan(tan(1/2*x)))`

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 25, normalized size of antiderivative = 1.67

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = -\frac{2x + \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)}{2a}$$

input `integrate(cot(x)^2/(a+a*csc(x)),x, algorithm="fricas")`

output `-1/2*(2*x + log(1/2*cos(x) + 1/2) - log(-1/2*cos(x) + 1/2))/a`

Sympy [F]

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot^2(x)}{\csc(x)+1} dx}{a}$$

input `integrate(cot(x)**2/(a+a*csc(x)),x)`

output `Integral(cot(x)**2/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.00

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = -\frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} + \frac{\log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

input `integrate(cot(x)^2/(a+a*csc(x)),x, algorithm="maxima")`

output `-2*arctan(sin(x)/(cos(x) + 1))/a + log(sin(x)/(cos(x) + 1))/a`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = -\frac{x}{a} + \frac{\log\left(|\tan\left(\frac{1}{2}x\right)|\right)}{a}$$

input `integrate(cot(x)^2/(a+a*csc(x)),x, algorithm="giac")`

output `-x/a + log(abs(tan(1/2*x)))/a`

Mupad [B] (verification not implemented)

Time = 14.95 (sec) , antiderivative size = 45, normalized size of antiderivative = 3.00

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = \frac{2 \operatorname{atan}\left(\frac{4}{4 \tan\left(\frac{x}{2}\right) + 4} - \frac{4 \tan\left(\frac{x}{2}\right)}{4 \tan\left(\frac{x}{2}\right) + 4}\right)}{a} + \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{a}$$

input `int(cot(x)^2/(a + a/sin(x)),x)`

output `(2*atan(4/(4*tan(x/2) + 4) - (4*tan(x/2))/(4*tan(x/2) + 4)))/a + log(tan(x/2))/a`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cot^2(x)}{a + a \csc(x)} dx = \frac{\log\left(\tan\left(\frac{x}{2}\right)\right) - x}{a}$$

input `int(cot(x)^2/(a+a*csc(x)),x)`

output `(log(tan(x/2)) - x)/a`

3.7 $\int \frac{\cot^3(x)}{a+a \csc(x)} dx$

Optimal result	74
Mathematica [A] (verified)	74
Rubi [A] (verified)	75
Maple [A] (verified)	76
Fricas [A] (verification not implemented)	77
Sympy [F]	77
Maxima [A] (verification not implemented)	77
Giac [A] (verification not implemented)	78
Mupad [B] (verification not implemented)	78
Reduce [B] (verification not implemented)	78

Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = -\frac{\csc(x)}{a} - \frac{\log(\sin(x))}{a}$$

output $-\csc(x)/a - \ln(\sin(x))/a$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = -\frac{\csc(x) + \log(\sin(x))}{a}$$

input `Integrate[Cot[x]^3/(a + a*Csc[x]), x]`

output $-((\text{Csc}[x] + \text{Log}[\text{Sin}[x]])/a)$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 4367, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^3}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{4367} \\
 & \frac{\int a \csc^2(x)(1 - \sin(x))d\sin(x)}{a^2} \\
 & \quad \downarrow \textcolor{blue}{27} \\
 & \frac{\int \csc^2(x)(1 - \sin(x))d\sin(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{49} \\
 & \frac{\int (\csc^2(x) - \csc(x)) d\sin(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{-\csc(x) - \log(\sin(x))}{a}
 \end{aligned}$$

input `Int[Cot[x]^3/(a + a*Csc[x]),x]`

output `(-Csc[x] - Log[Sin[x]])/a`

Definitions of rubi rules used

rule 27 $\text{Int}[(a_*)*(F_x_), x_{\text{Symbol}}] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[F_x, (b_*)*(G_x_) /; \text{FreeQ}[b, x]]$

rule 49 $\text{Int}[(a_*) + (b_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}), x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}[a, b, c, d, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[m + n + 2, 0]$

rule 2009 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4367 $\text{Int}[\cot[(c_*) + (d_*)*(x_*)^{(m_*)}*(\csc[(c_*) + (d_*)*(x_*)]*(b_*) + (a_*))^{(n_*)}], x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(a^{(m - n - 1)}*b^n*d) \text{ Subst}[\text{Int}[(a - b*x)^{((m - 1)/2)*((a + b*x)^{(m - 1)/2} + n)/x^{(m + n)}}, x], x, \text{Sin}[c + d*x]], x] /; \text{FreeQ}[a, b, c, d, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Maple [A] (verified)

Time = 0.21 (sec), antiderivative size = 13, normalized size of antiderivative = 0.81

method	result	size
derivativeDivides	$\frac{-\csc(x)+\ln(\csc(x))}{a}$	13
default	$\frac{-\csc(x)+\ln(\csc(x))}{a}$	13
risch	$\frac{ix}{a} - \frac{2ie^{ix}}{a(e^{2ix}-1)} - \frac{\ln(e^{2ix}-1)}{a}$	42

input $\text{int}(\cot(x)^3/(a+a*\csc(x)), x, \text{method}=\text{_RETURNVERBOSE})$

output $1/a*(-\csc(x)+\ln(\csc(x)))$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = -\frac{\log\left(\frac{1}{2} \sin(x)\right) \sin(x) + 1}{a \sin(x)}$$

input `integrate(cot(x)^3/(a+a*csc(x)),x, algorithm="fricas")`

output `-(log(1/2*sin(x))*sin(x) + 1)/(a*sin(x))`

Sympy [F]

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot^3(x)}{\csc(x)+1} dx}{a}$$

input `integrate(cot(x)**3/(a+a*csc(x)),x)`

output `Integral(cot(x)**3/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = -\frac{\log(\sin(x))}{a} - \frac{1}{a \sin(x)}$$

input `integrate(cot(x)^3/(a+a*csc(x)),x, algorithm="maxima")`

output `-log(sin(x))/a - 1/(a*sin(x))`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = -\frac{\frac{1}{\sin(x)} + \log(|\sin(x)|)}{a}$$

input `integrate(cot(x)^3/(a+a*csc(x)),x, algorithm="giac")`

output `-(1/sin(x) + log(abs(sin(x))))/a`

Mupad [B] (verification not implemented)

Time = 14.92 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.25

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = -\frac{\frac{\tan(\frac{x}{2})}{2} - \ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + \ln\left(\tan\left(\frac{x}{2}\right)\right) + \frac{1}{2\tan(\frac{x}{2})}}{a}$$

input `int(cot(x)^3/(a + a/sin(x)),x)`

output `-(tan(x/2)/2 - log(tan(x/2)^2 + 1) + log(tan(x/2)) + 1/(2*tan(x/2)))/a`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.94

$$\int \frac{\cot^3(x)}{a + a \csc(x)} dx = \frac{\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x) - \log\left(\tan\left(\frac{x}{2}\right)\right) \sin(x) - 1}{\sin(x) a}$$

input `int(cot(x)^3/(a+a*csc(x)),x)`

output `(log(tan(x/2)**2 + 1)*sin(x) - log(tan(x/2))*sin(x) - 1)/(sin(x)*a)`

3.8 $\int \frac{\cot^4(x)}{a+a \csc(x)} dx$

Optimal result	79
Mathematica [B] (verified)	79
Rubi [A] (verified)	80
Maple [A] (verified)	82
Fricas [B] (verification not implemented)	82
Sympy [F]	83
Maxima [B] (verification not implemented)	83
Giac [B] (verification not implemented)	84
Mupad [B] (verification not implemented)	84
Reduce [B] (verification not implemented)	85

Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\cot^4(x)}{a + a \csc(x)} dx = \frac{x}{a} + \frac{\operatorname{arctanh}(\cos(x))}{2a} + \frac{\cot(x)(2 - \csc(x))}{2a}$$

output `x/a+1/2*arctanh(cos(x))/a+1/2*cot(x)*(2-csc(x))/a`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 90 vs. $2(31) = 62$.

Time = 0.10 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.90

$$\begin{aligned} \int \frac{\cot^4(x)}{a + a \csc(x)} dx = & \frac{x}{a} + \frac{\cot\left(\frac{x}{2}\right)}{2a} - \frac{\csc^2\left(\frac{x}{2}\right)}{8a} + \frac{\log\left(\cos\left(\frac{x}{2}\right)\right)}{2a} \\ & - \frac{\log\left(\sin\left(\frac{x}{2}\right)\right)}{2a} + \frac{\sec^2\left(\frac{x}{2}\right)}{8a} - \frac{\tan\left(\frac{x}{2}\right)}{2a} \end{aligned}$$

input `Integrate[Cot[x]^4/(a + a*Csc[x]),x]`

output $\frac{x/a + \cot[x/2]/(2*a) - \csc[x/2]^2/(8*a) + \log[\cos[x/2]]/(2*a) - \log[\sin[x/2]]/(2*a) + \sec[x/2]^2/(8*a) - \tan[x/2]/(2*a)}{2}$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 36, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3042, 4376, 25, 3042, 4369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^4(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^4}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{4376} \\
 & \frac{\int -\cot^2(x)(a - a \csc(x)) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & -\frac{\int \cot^2(x)(a - a \csc(x)) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & -\frac{\int \cot(x)^2(a - a \csc(x)) dx}{a^2} \\
 & \quad \downarrow \textcolor{blue}{4369} \\
 & -\frac{\frac{1}{2} \int (2a - a \csc(x)) dx - \frac{1}{2} \cot(x)(2a - a \csc(x))}{a^2} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & -\frac{\frac{1}{2}(-a \operatorname{arctanh}(\cos(x)) - 2ax) - \frac{1}{2} \cot(x)(2a - a \csc(x))}{a^2}
 \end{aligned}$$

input $\text{Int}[\cot[x]^4/(a + a*\csc[x]), x]$

output $-(((-2*a*x - a*\text{ArcTanh}[\cos[x]])/2 - (\cot[x]*(2*a - a*\csc[x]))/2)/a^2)$

Definitions of rubi rules used

rule 25 $\text{Int}[-(F[x], x), x] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F[x], x], x]$

rule 2009 $\text{Int}[u, x], x \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u, x], x \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4369 $\text{Int}[(\cot[(c_.) + (d_.)*(x_.)]*(e_.)])^{(m_*)}(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_)), x], x \rightarrow \text{Simp}[(-e)*(e*\cot[c + d*x])^{(m - 1)}((a*m + b*(m - 1))*\csc[c + d*x])/((d*m*(m - 1))), x] - \text{Simp}[e^{2/m} \quad \text{Int}[(e*\cot[c + d*x])^{(m - 2)}*(a*m + b*(m - 1))*\csc[c + d*x]], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{GtQ}[m, 1]$

rule 4376 $\text{Int}[(\cot[(c_.) + (d_.)*(x_.)]*(e_.)])^{(m_*)}(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_))^{(n_*)}, x], x \rightarrow \text{Simp}[a^{(2*n)}/e^{(2*n)} \quad \text{Int}[(e*\cot[c + d*x])^{(m + 2*n)}/(-a + b*\csc[c + d*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{EqQ}[a^{2 - b^2}, 0] \&& \text{ILtQ}[n, 0]$

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.65

method	result	size
default	$\frac{\frac{\tan(\frac{x}{2})^2}{2} - 2\tan(\frac{x}{2}) - \frac{1}{2\tan(\frac{x}{2})^2} + \frac{2}{\tan(\frac{x}{2})} - 2\ln(\tan(\frac{x}{2})) + 8\arctan(\tan(\frac{x}{2}))}{4a}$	51
risch	$\frac{x}{a} + \frac{2ie^{2ix} + e^{3ix} - 2i + e^{ix}}{a(e^{2ix} - 1)^2} - \frac{\ln(e^{ix} - 1)}{2a} + \frac{\ln(e^{ix} + 1)}{2a}$	67

input `int(cot(x)^4/(a+a*csc(x)),x,method=_RETURNVERBOSE)`

output `1/4/a*(1/2*tan(1/2*x)^2-2*tan(1/2*x)-1/2/tan(1/2*x)^2+2/tan(1/2*x)-2*ln(tan(1/2*x))+8*arctan(tan(1/2*x)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 64 vs. $2(25) = 50$.

Time = 0.09 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.06

$$\int \frac{\cot^4(x)}{a + a \csc(x)} dx \\ = \frac{4 x \cos(x)^2 + (\cos(x)^2 - 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x)^2 - 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 4 \cos(x) \sin(x)}{4 (a \cos(x)^2 - a)}$$

input `integrate(cot(x)^4/(a+a*csc(x)),x, algorithm="fricas")`

output `1/4*(4*x*cos(x)^2 + (cos(x)^2 - 1)*log(1/2*cos(x) + 1/2) - (cos(x)^2 - 1)*log(-1/2*cos(x) + 1/2) - 4*cos(x)*sin(x) - 4*x + 2*cos(x))/(a*cos(x)^2 - a)`

Sympy [F]

$$\int \frac{\cot^4(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot^4(x)}{\csc(x)+1} dx}{a}$$

input `integrate(cot(x)**4/(a+a*csc(x)),x)`

output `Integral(cot(x)**4/(csc(x) + 1), x)/a`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 86 vs. $2(25) = 50$.

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.77

$$\begin{aligned} \int \frac{\cot^4(x)}{a + a \csc(x)} dx &= -\frac{\frac{4 \sin(x)}{\cos(x)+1} - \frac{\sin(x)^2}{(\cos(x)+1)^2}}{8 a} + \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} \\ &\quad - \frac{\log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{2 a} + \frac{\left(\frac{4 \sin(x)}{\cos(x)+1} - 1\right)(\cos(x) + 1)^2}{8 a \sin(x)^2} \end{aligned}$$

input `integrate(cot(x)^4/(a+a*csc(x)),x, algorithm="maxima")`

output `-1/8*(4*sin(x)/(cos(x) + 1) - sin(x)^2/(cos(x) + 1)^2)/a + 2*arctan(sin(x)/(cos(x) + 1))/a - 1/2*log(sin(x)/(cos(x) + 1))/a + 1/8*(4*sin(x)/(cos(x) + 1) - 1)*(cos(x) + 1)^2/(a*sin(x)^2)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(25) = 50$.

Time = 0.14 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.10

$$\int \frac{\cot^4(x)}{a + a \csc(x)} dx = \frac{x}{a} - \frac{\log(|\tan(\frac{1}{2}x)|)}{2a} + \frac{a \tan(\frac{1}{2}x)^2 - 4a \tan(\frac{1}{2}x)}{8a^2} \\ + \frac{6 \tan(\frac{1}{2}x)^2 + 4 \tan(\frac{1}{2}x) - 1}{8a \tan(\frac{1}{2}x)^2}$$

input `integrate(cot(x)^4/(a+a*csc(x)),x, algorithm="giac")`

output `x/a - 1/2*log(abs(tan(1/2*x)))/a + 1/8*(a*tan(1/2*x)^2 - 4*a*tan(1/2*x))/a^2 + 1/8*(6*tan(1/2*x)^2 + 4*tan(1/2*x) - 1)/(a*tan(1/2*x)^2)`

Mupad [B] (verification not implemented)

Time = 15.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.74

$$\int \frac{\cot^4(x)}{a + a \csc(x)} dx = \frac{\tan(\frac{x}{2})^2}{8a} - \frac{2 \operatorname{atan}\left(\frac{4}{4 \tan(\frac{x}{2})+2}-\frac{2 \tan(\frac{x}{2})}{4 \tan(\frac{x}{2})+2}\right)}{a} \\ - \frac{\tan(\frac{x}{2})}{2a} - \frac{\ln(\tan(\frac{x}{2}))}{2a} + \frac{2 \tan(\frac{x}{2})-\frac{1}{2}}{4a \tan(\frac{x}{2})^2}$$

input `int(cot(x)^4/(a + a/sin(x)),x)`

output `tan(x/2)^2/(8*a) - (2*atan(4/(4*tan(x/2) + 2) - (2*tan(x/2))/(4*tan(x/2) + 2)))/a - tan(x/2)/(2*a) - log(tan(x/2))/(2*a) + (2*tan(x/2) - 1/2)/(4*a*tan(x/2)^2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

$$\int \frac{\cot^4(x)}{a + a \csc(x)} dx = \frac{2 \cos(x) \sin(x) - \cos(x) - \log(\tan(\frac{x}{2})) \sin(x)^2 + 2 \sin(x)^2 x}{2 \sin(x)^2 a}$$

input `int(cot(x)^4/(a+a*csc(x)),x)`

output `(2*cos(x)*sin(x) - cos(x) - log(tan(x/2))*sin(x)**2 + 2*sin(x)**2*x)/(2*sin(x)**2*a)`

3.9 $\int \frac{\cot^5(x)}{a+a \csc(x)} dx$

Optimal result	86
Mathematica [A] (verified)	86
Rubi [A] (verified)	87
Maple [A] (verified)	88
Fricas [A] (verification not implemented)	89
Sympy [F]	89
Maxima [A] (verification not implemented)	89
Giac [A] (verification not implemented)	90
Mupad [B] (verification not implemented)	90
Reduce [B] (verification not implemented)	91

Optimal result

Integrand size = 13, antiderivative size = 36

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx = \frac{\csc(x)}{a} + \frac{\csc^2(x)}{2a} - \frac{\csc^3(x)}{3a} + \frac{\log(\sin(x))}{a}$$

output `csc(x)/a+1/2*csc(x)^2/a-1/3*csc(x)^3/a+ln(sin(x))/a`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx = \frac{\csc(x) + \frac{\csc^2(x)}{2} - \frac{\csc^3(x)}{3} + \log(\sin(x))}{a}$$

input `Integrate[Cot[x]^5/(a + a*Csc[x]),x]`

output `(Csc[x] + Csc[x]^2/2 - Csc[x]^3/3 + Log[Sin[x]])/a`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 4367, 27, 84, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^5(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^5}{a \csc(x) + a} dx \\
 & \quad \downarrow \textcolor{blue}{4367} \\
 & \frac{\int a^3 \csc^4(x)(1 - \sin(x))^2(\sin(x) + 1)d\sin(x)}{a^4} \\
 & \quad \downarrow \textcolor{blue}{27} \\
 & \frac{\int \csc^4(x)(1 - \sin(x))^2(\sin(x) + 1)d\sin(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{84} \\
 & \frac{\int (\csc^4(x) - \csc^3(x) - \csc^2(x) + \csc(x)) d\sin(x)}{a} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{-\frac{1}{3} \csc^3(x) + \frac{\csc^2(x)}{2} + \csc(x) + \log(\sin(x))}{a}
 \end{aligned}$$

input `Int[Cot[x]^5/(a + a*Csc[x]),x]`

output `(Csc[x] + Csc[x]^2/2 - Csc[x]^3/3 + Log[Sin[x]])/a`

Definitions of rubi rules used

rule 27 $\text{Int}[(a_*)(F_x_), x_{\text{Symbol}}] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[F_x, (b_*)(G_x_) /; \text{FreeQ}[b, x]]$

rule 84 $\text{Int}[(d_*)(x_*)^{(n_*)}((a_*) + (b_*)(x_*)*((e_*) + (f_*)(x_*)^{(p_*)}), x] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, d, e, f, n\}, x] \&& \text{IGtQ}[p, 0] \&& \text{EqQ}[b*e + a*f, 0] \&& \text{!(ILtQ}[n + p + 2, 0] \&& \text{GtQ}[n + 2*p, 0])$

rule 2009 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4367 $\text{Int}[\cot[(c_*) + (d_*)(x_*)]^{(m_*)}(\csc[(c_*) + (d_*)(x_*)]*(b_*) + (a_*)^{(n_*)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(a^{(m - n - 1)}*b^n*d) \text{ Subst}[\text{Int}[(a - b*x)^{((m - 1)/2)*((m - 1)/2 + n)}/x^{(m + n)}, x], x, \text{Sin}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Maple [A] (verified)

Time = 0.46 (sec), antiderivative size = 25, normalized size of antiderivative = 0.69

method	result	size
derivativeDivides	$\frac{\csc(x) - \frac{\csc(x)^3}{3} + \frac{\csc(x)^2}{2}}{a} - \ln(\csc(x))$	25
default	$\frac{\csc(x) - \frac{\csc(x)^3}{3} + \frac{\csc(x)^2}{2}}{a} - \ln(\csc(x))$	25
risch	$-\frac{ix}{a} + \frac{2i(3ie^{4ix} + 3e^{5ix} - 3ie^{2ix} - 2e^{3ix} + 3e^{ix})}{3a(e^{2ix} - 1)^3} + \frac{\ln(e^{2ix} - 1)}{a}$	74

input `int(cot(x)^5/(a+a*csc(x)),x,method=_RETURNVERBOSE)`

output $\frac{1}{a} \left(\csc(x) - \frac{1}{3} \csc(x)^3 + \frac{1}{2} \csc(x)^2 - \ln(\csc(x)) \right)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec), antiderivative size = 45, normalized size of antiderivative = 1.25

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx = \frac{6 (\cos(x)^2 - 1) \log\left(\frac{1}{2} \sin(x)\right) \sin(x) + 6 \cos(x)^2 - 3 \sin(x) - 4}{6 (a \cos(x)^2 - a) \sin(x)}$$

input `integrate(cot(x)^5/(a+a*csc(x)),x, algorithm="fricas")`

output $\frac{1}{6} \left(6 (\cos(x)^2 - 1) \log(1/2 \sin(x)) \sin(x) + 6 \cos(x)^2 - 3 \sin(x) - 4 \right) / ((a \cos(x)^2 - a) \sin(x))$

Sympy [F]

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot^5(x)}{\csc(x)+1} dx}{a}$$

input `integrate(cot(x)**5/(a+a*csc(x)),x)`

output `Integral(cot(x)**5/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.03 (sec), antiderivative size = 29, normalized size of antiderivative = 0.81

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx = \frac{\log(\sin(x))}{a} + \frac{6 \sin(x)^2 + 3 \sin(x) - 2}{6 a \sin(x)^3}$$

input `integrate(cot(x)^5/(a+a*csc(x)),x, algorithm="maxima")`

output $\log(\sin(x))/a + 1/6*(6*\sin(x)^2 + 3*\sin(x) - 2)/(a*\sin(x)^3)$

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.81

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx = \frac{\frac{6 \sin(x)^2 + 3 \sin(x) - 2}{\sin(x)^3} + 6 \log(|\sin(x)|)}{6 a}$$

input `integrate(cot(x)^5/(a+a*csc(x)),x, algorithm="giac")`

output $1/6*((6*\sin(x)^2 + 3*\sin(x) - 2)/\sin(x)^3 + 6*\log(\text{abs}(\sin(x))))/a$

Mupad [B] (verification not implemented)

Time = 15.19 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.22

$$\begin{aligned} \int \frac{\cot^5(x)}{a + a \csc(x)} dx = & \frac{3 \tan(\frac{x}{2})}{8 a} - \frac{\ln(\tan(\frac{x}{2})^2 + 1)}{a} + \frac{\tan(\frac{x}{2})^2}{8 a} - \frac{\tan(\frac{x}{2})^3}{24 a} \\ & + \frac{\ln(\tan(\frac{x}{2}))}{a} + \frac{3 \tan(\frac{x}{2})^2 + \tan(\frac{x}{2}) - \frac{1}{3}}{8 a \tan(\frac{x}{2})^3} \end{aligned}$$

input `int(cot(x)^5/(a + a/sin(x)),x)`

output $(3*\tan(x/2))/(8*a) - \log(\tan(x/2)^2 + 1)/a + \tan(x/2)^2/(8*a) - \tan(x/2)^3/(24*a) + \log(\tan(x/2))/a + (\tan(x/2) + 3*\tan(x/2)^2 - 1/3)/(8*a*\tan(x/2)^3)$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.47

$$\int \frac{\cot^5(x)}{a + a \csc(x)} dx \\ = \frac{-12 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^3 + 12 \log\left(\tan\left(\frac{x}{2}\right)\right) \sin(x)^3 - 3 \sin(x)^3 + 12 \sin(x)^2 + 6 \sin(x) - 4}{12 \sin(x)^3 a}$$

input `int(cot(x)^5/(a+a*csc(x)),x)`

output `(- 12*log(tan(x/2)**2 + 1)*sin(x)**3 + 12*log(tan(x/2))*sin(x)**3 - 3*sin(x)**3 + 12*sin(x)**2 + 6*sin(x) - 4)/(12*sin(x)**3*a)`

$$3.10 \quad \int \frac{\cot^6(x)}{a+a \csc(x)} dx$$

Optimal result	92
Mathematica [B] (verified)	92
Rubi [A] (verified)	93
Maple [A] (verified)	95
Fricas [B] (verification not implemented)	95
Sympy [F]	96
Maxima [B] (verification not implemented)	96
Giac [B] (verification not implemented)	97
Mupad [B] (verification not implemented)	98
Reduce [B] (verification not implemented)	98

Optimal result

Integrand size = 13, antiderivative size = 49

$$\begin{aligned} \int \frac{\cot^6(x)}{a + a \csc(x)} dx &= -\frac{x}{a} - \frac{3 \operatorname{arctanh}(\cos(x))}{8a} \\ &\quad + \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a} \end{aligned}$$

output
$$-\frac{x}{a} - \frac{3}{8} \operatorname{arctanh}(\cos(x)) \cdot \frac{1}{a+1} + \frac{1}{12} \cot(x)^3 \cdot (4 - 3 \csc(x)) \cdot \frac{1}{a-1} - \frac{8}{a} \cot(x) \cdot (8 - 3 \csc(x)) \cdot \frac{1}{a}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 163 vs. $2(49) = 98$.

Time = 0.11 (sec) , antiderivative size = 163, normalized size of antiderivative = 3.33

$$\int \frac{\cot^6(x)}{a + a \csc(x)} dx = -\frac{x}{a} - \frac{2 \cot\left(\frac{x}{2}\right)}{3a} + \frac{5 \csc^2\left(\frac{x}{2}\right)}{32a} + \frac{\cot\left(\frac{x}{2}\right) \csc^2\left(\frac{x}{2}\right)}{24a} - \frac{\csc^4\left(\frac{x}{2}\right)}{64a} - \frac{3 \log\left(\cos\left(\frac{x}{2}\right)\right)}{8a} + \frac{3 \log\left(\sin\left(\frac{x}{2}\right)\right)}{8a} - \frac{5 \sec^2\left(\frac{x}{2}\right)}{32a} + \frac{\sec^4\left(\frac{x}{2}\right)}{64a} + \frac{2 \tan\left(\frac{x}{2}\right)}{3a} - \frac{\sec^2\left(\frac{x}{2}\right) \tan\left(\frac{x}{2}\right)}{24a}$$

input `Integrate[Cot[x]^6/(a + a*Csc[x]),x]`

output
$$\begin{aligned} & -(x/a) - (2*\text{Cot}[x/2])/(3*a) + (5*\text{Csc}[x/2]^2)/(32*a) + (\text{Cot}[x/2]*\text{Csc}[x/2]^2) \\ & /(24*a) - \text{Csc}[x/2]^4/(64*a) - (3*\text{Log}[\text{Cos}[x/2]])/(8*a) + (3*\text{Log}[\text{Sin}[x/2]]) \\ & /(8*a) - (5*\text{Sec}[x/2]^2)/(32*a) + \text{Sec}[x/2]^4/(64*a) + (2*\text{Tan}[x/2])/(3*a) - \\ & (\text{Sec}[x/2]^2*\text{Tan}[x/2])/(24*a) \end{aligned}$$

Rubi [A] (verified)

Time = 0.38 (sec), antiderivative size = 58, normalized size of antiderivative = 1.18, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$, Rules used = {3042, 4376, 25, 3042, 4369, 3042, 4369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^6(x)}{a \csc(x) + a} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cot(x)^6}{a \csc(x) + a} dx \\ & \quad \downarrow \text{4376} \\ & \frac{\int -\cot^4(x)(a - a \csc(x)) dx}{a^2} \\ & \quad \downarrow \text{25} \\ & -\frac{\int \cot^4(x)(a - a \csc(x)) dx}{a^2} \\ & \quad \downarrow \text{3042} \\ & -\frac{\int \cot(x)^4(a - a \csc(x)) dx}{a^2} \\ & \quad \downarrow \text{4369} \\ & -\frac{-\frac{1}{4} \int \cot^2(x)(4a - 3a \csc(x)) dx - \frac{1}{12} \cot^3(x)(4a - 3a \csc(x))}{a^2} \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & -\frac{-\frac{1}{4} \int \cot(x)^2 (4a - 3a \csc(x)) dx - \frac{1}{12} \cot^3(x) (4a - 3a \csc(x))}{a^2} \\
 & \quad \downarrow \text{4369} \\
 & -\frac{\frac{1}{4} \left(\frac{1}{2} \int (8a - 3a \csc(x)) dx + \frac{1}{2} \cot(x) (8a - 3a \csc(x)) \right) - \frac{1}{12} \cot^3(x) (4a - 3a \csc(x))}{a^2} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\frac{1}{4} \left(\frac{1}{2} (3a \operatorname{arctanh}(\cos(x)) + 8ax) + \frac{1}{2} \cot(x) (8a - 3a \csc(x)) \right) - \frac{1}{12} \cot^3(x) (4a - 3a \csc(x))}{a^2}
 \end{aligned}$$

input `Int[Cot[x]^6/(a + a*Csc[x]), x]`

output `-((-1/12*(Cot[x]^3*(4*a - 3*a*Csc[x])) + ((8*a*x + 3*a*ArcTanh[Cos[x]])/2 + (Cot[x]*(8*a - 3*a*Csc[x]))/2)/4)/a^2)`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4369 `Int[(cot[(c_.) + (d_.)*(x_.)]*(e_.))^m_]*(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.), x_Symbol] :> Simp[(-e)*(e*Cot[c + d*x])^(m - 1)*((a*m + b*(m - 1)*Csc[c + d*x])/(d*m*(m - 1))), x] - Simp[e^(2/m) Int[(e*Cot[c + d*x])^(m - 2)*((a*m + b*(m - 1)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e}, x] && GtQ[m, 1]`

rule 4376

```
Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^(m_)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] :> Simp[a^(2*n)/e^(2*n) Int[(e*Cot[c + d*x])^(m + 2*n)/(-a + b*Csc[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[a^2 - b^2, 0] && ILtQ[n, 0]
```

Maple [A] (verified)

Time = 0.63 (sec), antiderivative size = 83, normalized size of antiderivative = 1.69

method	result	
default	$\frac{\tan(\frac{x}{2})^4 - \frac{2 \tan(\frac{x}{2})^3}{3} - 2 \tan(\frac{x}{2})^2 + 10 \tan(\frac{x}{2}) - \frac{1}{4 \tan(\frac{x}{2})^4} + \frac{2}{3 \tan(\frac{x}{2})^3} + \frac{2}{\tan(\frac{x}{2})^2} - \frac{10}{\tan(\frac{x}{2})} + 6 \ln(\tan(\frac{x}{2})) - 32 \arctan(\tan(\frac{x}{2}))}{16a}$	8
risch	$-\frac{x}{a} - \frac{48ie^{6ix}+15e^{7ix}-96ie^{4ix}+9e^{5ix}+80ie^{2ix}+9e^{3ix}-32i+15e^{ix}}{12a(e^{2ix}-1)^4} + \frac{3\ln(e^{ix}-1)}{8a} - \frac{3\ln(e^{ix}+1)}{8a}$	1

input `int(cot(x)^6/(a+a*csc(x)),x,method=_RETURNVERBOSE)`output
$$\frac{1/16/a*(1/4*tan(1/2*x)^4-2/3*tan(1/2*x)^3-2*tan(1/2*x)^2+10*tan(1/2*x)-1/4/tan(1/2*x)^4+2/3/tan(1/2*x)^3+2/tan(1/2*x)^2-10/tan(1/2*x)+6*ln(tan(1/2*x))) - 32*arctan(tan(1/2*x)))$$
Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 104 vs. $2(43) = 86$.

Time = 0.09 (sec), antiderivative size = 104, normalized size of antiderivative = 2.12

$$\int \frac{\cot^6(x)}{a + a \csc(x)} dx = \frac{-48 x \cos(x)^4 - 96 x \cos(x)^2 + 30 \cos(x)^3 + 9 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(\frac{1}{2} \cos(x) + \frac{1}{2}) - 9 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(-\frac{1}{2} \cos(x) + \frac{1}{2})}{48 (a \cos(x)^4 - 2 a \cos(x)^2 + a)}$$

input `integrate(cot(x)^6/(a+a*csc(x)),x, algorithm="fricas")`

output

$$-1/48*(48*x*cos(x)^4 - 96*x*cos(x)^2 + 30*cos(x)^3 + 9*(cos(x)^4 - 2*cos(x)^2 + 1)*log(1/2*cos(x) + 1/2) - 9*(cos(x)^4 - 2*cos(x)^2 + 1)*log(-1/2*cos(x) + 1/2) - 16*(4*cos(x)^3 - 3*cos(x))*sin(x) + 48*x - 18*cos(x))/(a*cos(x)^4 - 2*a*cos(x)^2 + a)$$

Sympy [F]

$$\int \frac{\cot^6(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot^6(x)}{\csc(x)+1} dx}{a}$$

input

```
integrate(cot(x)**6/(a+a*csc(x)),x)
```

output

```
Integral(cot(x)**6/(csc(x) + 1), x)/a
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. $2(43) = 86$.

Time = 0.12 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.73

$$\begin{aligned} \int \frac{\cot^6(x)}{a + a \csc(x)} dx &= \frac{\frac{120 \sin(x)}{\cos(x)+1} - \frac{24 \sin(x)^2}{(\cos(x)+1)^2} - \frac{8 \sin(x)^3}{(\cos(x)+1)^3} + \frac{3 \sin(x)^4}{(\cos(x)+1)^4}}{192 a} \\ &\quad - \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} + \frac{3 \log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{8 a} \\ &\quad + \frac{\left(\frac{8 \sin(x)}{\cos(x)+1} + \frac{24 \sin(x)^2}{(\cos(x)+1)^2} - \frac{120 \sin(x)^3}{(\cos(x)+1)^3} - 3\right)(\cos(x) + 1)^4}{192 a \sin(x)^4} \end{aligned}$$

input

```
integrate(cot(x)**6/(a+a*csc(x)),x, algorithm="maxima")
```

output

$$\frac{1}{192} \left(\frac{120 \sin(x)}{\cos(x) + 1} - \frac{24 \sin(x)^2}{\cos(x) + 1} - \frac{8 \sin(x)^3}{\cos(x) + 1} - \frac{3 \sin(x)^4}{\cos(x) + 1} + \frac{3 \sin(x)^4}{(\cos(x) + 1)^4} \right) + \frac{2 \arctan(\sin(x)/(\cos(x) + 1))}{a} + \frac{3/8 \log(\sin(x)/(\cos(x) + 1))}{a} + \frac{1}{192} \left(\frac{8 \sin(x)}{\cos(x) + 1} + \frac{24 \sin(x)^2}{\cos(x) + 1} - \frac{120 \sin(x)^3}{\cos(x) + 1} - \frac{3 (\cos(x) + 1)^4}{a \sin(x)^4} \right)$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 109 vs. $2(43) = 86$.

Time = 0.14 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.22

$$\begin{aligned} & \int \frac{\cot^6(x)}{a + a \csc(x)} dx \\ &= -\frac{x}{a} + \frac{3 \log(|\tan(\frac{1}{2}x)|)}{8a} \\ &+ \frac{3 a^3 \tan(\frac{1}{2}x)^4 - 8 a^3 \tan(\frac{1}{2}x)^3 - 24 a^3 \tan(\frac{1}{2}x)^2 + 120 a^3 \tan(\frac{1}{2}x)}{192 a^4} \\ &- \frac{150 \tan(\frac{1}{2}x)^4 + 120 \tan(\frac{1}{2}x)^3 - 24 \tan(\frac{1}{2}x)^2 - 8 \tan(\frac{1}{2}x) + 3}{192 a \tan(\frac{1}{2}x)^4} \end{aligned}$$

input

```
integrate(cot(x)^6/(a+a*csc(x)),x, algorithm="giac")
```

output

$$\begin{aligned} & -\frac{x}{a} + \frac{3/8 \log(\tan(1/2*x))}{a} + \frac{1}{192} \left(3 a^3 \tan(1/2*x)^4 - 8 a^3 \tan(1/2*x)^3 - 24 a^3 \tan(1/2*x)^2 + 120 a^3 \tan(1/2*x) \right) / a^4 - \frac{1}{192} \left(150 \tan(1/2*x)^4 + 120 \tan(1/2*x)^3 - 24 \tan(1/2*x)^2 - 8 \tan(1/2*x) + 3 \right) / (a \tan(1/2*x)^4) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 15.24 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.51

$$\int \frac{\cot^6(x)}{a + a \csc(x)} dx = \frac{5 \tan(\frac{x}{2})}{8 a} + \frac{2 \operatorname{atan}\left(\frac{4}{4 \tan(\frac{x}{2}) + 3/2} - \frac{3 \tan(\frac{x}{2})}{2(4 \tan(\frac{x}{2}) + 3/2)}\right)}{a} \\ - \frac{\tan(\frac{x}{2})^2}{8 a} - \frac{\tan(\frac{x}{2})^3}{24 a} + \frac{\tan(\frac{x}{2})^4}{64 a} + \frac{3 \ln(\tan(\frac{x}{2}))}{8 a} \\ + \frac{-10 \tan(\frac{x}{2})^3 + 2 \tan(\frac{x}{2})^2 + \frac{2 \tan(\frac{x}{2})}{3} - \frac{1}{4}}{16 a \tan(\frac{x}{2})^4}$$

input `int(cot(x)^6/(a + a/sin(x)),x)`

output `(5*tan(x/2))/(8*a) + (2*atan(4/(4*tan(x/2) + 3/2) - (3*tan(x/2))/(2*(4*tan(x/2) + 3/2))))/a - tan(x/2)^2/(8*a) - tan(x/2)^3/(24*a) + tan(x/2)^4/(64*a) + (3*log(tan(x/2)))/(8*a) + ((2*tan(x/2))/3 + 2*tan(x/2)^2 - 10*tan(x/2)^3 - 1/4)/(16*a*tan(x/2)^4)`

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.10

$$\int \frac{\cot^6(x)}{a + a \csc(x)} dx \\ = \frac{-32 \cos(x) \sin(x)^3 + 15 \cos(x) \sin(x)^2 + 8 \cos(x) \sin(x) - 6 \cos(x) + 9 \log(\tan(\frac{x}{2})) \sin(x)^4 - 24 \sin(x)^4 a}{24 \sin(x)^4 a}$$

input `int(cot(x)^6/(a+a*csc(x)),x)`

output `(- 32*cos(x)*sin(x)**3 + 15*cos(x)*sin(x)**2 + 8*cos(x)*sin(x) - 6*cos(x) + 9*log(tan(x/2))*sin(x)**4 - 24*sin(x)**4*x)/(24*sin(x)**4*a)`

3.11 $\int \frac{\cot^7(x)}{a+a \csc(x)} dx$

Optimal result	99
Mathematica [A] (verified)	99
Rubi [A] (verified)	100
Maple [A] (verified)	101
Fricas [A] (verification not implemented)	102
Sympy [F]	102
Maxima [A] (verification not implemented)	103
Giac [A] (verification not implemented)	103
Mupad [B] (verification not implemented)	103
Reduce [B] (verification not implemented)	104

Optimal result

Integrand size = 13, antiderivative size = 58

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx = -\frac{\csc(x)}{a} - \frac{\csc^2(x)}{a} + \frac{2 \csc^3(x)}{3a} + \frac{\csc^4(x)}{4a} - \frac{\csc^5(x)}{5a} - \frac{\log(\sin(x))}{a}$$

output
$$-\csc(x)/a - \csc(x)^2/a + 2/3*\csc(x)^3/a + 1/4*\csc(x)^4/a - 1/5*\csc(x)^5/a - \ln(\sin(x))/a$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.67

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx = -\frac{\csc(x) + \csc^2(x) - \frac{2 \csc^3(x)}{3} - \frac{\csc^4(x)}{4} + \frac{\csc^5(x)}{5} + \log(\sin(x))}{a}$$

input `Integrate[Cot[x]^7/(a + a*Csc[x]),x]`

output
$$-((Csc[x] + Csc[x]^2 - (2*Csc[x]^3)/3 - Csc[x]^4/4 + Csc[x]^5/5 + Log[Sin[x]])/a)$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.76, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 4367, 27, 99, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^7(x)}{a \csc(x) + a} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(x)^7}{a \csc(x) + a} dx \\
 & \quad \downarrow \text{4367} \\
 & \frac{\int a^5 \csc^6(x)(1 - \sin(x))^3(\sin(x) + 1)^2 d\sin(x)}{a^6} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \csc^6(x)(1 - \sin(x))^3(\sin(x) + 1)^2 d\sin(x)}{a} \\
 & \quad \downarrow \text{99} \\
 & \frac{\int (\csc^6(x) - \csc^5(x) - 2\csc^4(x) + 2\csc^3(x) + \csc^2(x) - \csc(x)) d\sin(x)}{a} \\
 & \quad \downarrow \text{2009} \\
 & \frac{-\frac{1}{5}\csc^5(x) + \frac{\csc^4(x)}{4} + \frac{2\csc^3(x)}{3} - \csc^2(x) - \csc(x) - \log(\sin(x))}{a}
 \end{aligned}$$

input `Int[Cot[x]^7/(a + a*Csc[x]),x]`

output `(-Csc[x] - Csc[x]^2 + (2*Csc[x]^3)/3 + Csc[x]^4/4 - Csc[x]^5/5 - Log[Sin[x]])/a`

Definitions of rubi rules used

rule 27 $\text{Int}[(a_*)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[F_x, (b_*)*(G_x_) /; \text{FreeQ}[b, x]]$

rule 99 $\text{Int}[(a_*) + (b_*)*(x_*)^m * ((c_*) + (d_*)*(x_*)^n * ((e_*) + (f_*)*(x_*)^p), x_*) \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, p\}, x] \&& \text{IntegersQ}[m, n] \&& (\text{IntegerQ}[p] \mid (\text{GtQ}[m, 0] \&& \text{GeQ}[n, -1]))]$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4367 $\text{Int}[\cot[(c_*) + (d_*)*(x_*)]^m * (\csc[(c_*) + (d_*)*(x_*)]*b_*) + (a_*)^n, x_Symbol] \rightarrow \text{Simp}[1/(a^{(m-n-1)*b^n*d}) \text{ Subst}[\text{Int}[(a - b*x)^{(m-1)/2} * ((a + b*x)^{(m-1)/2+n}/x^{(m+n)}), x], x, \text{Sin}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{IntegerQ}[(m-1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.64

method	result	size
derivativedivides	$\frac{-\csc(x)^5 + \frac{\csc(x)^4}{4} + \frac{2\csc(x)^3}{3} - \csc(x)^2 - \csc(x) + \ln(\csc(x))}{a}$	37
default	$\frac{-\csc(x)^5 + \frac{\csc(x)^4}{4} + \frac{2\csc(x)^3}{3} - \csc(x)^2 - \csc(x) + \ln(\csc(x))}{a}$	37
risch	$\frac{ix}{a} - \frac{2i(30ie^{8ix} + 15e^{9ix} - 60ie^{6ix} - 20e^{7ix} + 60ie^{4ix} + 58e^{5ix} - 30ie^{2ix} - 20e^{3ix} + 15e^{ix})}{15a(e^{2ix}-1)^5} - \frac{\ln(e^{2ix}-1)}{a}$	105

input $\text{int}(\cot(x)^7/(a+a*\csc(x)), x, \text{method}=\text{RETURNVERBOSE})$

output $\frac{1}{a} \left(-\frac{1}{5} \csc(x)^5 + \frac{1}{4} \csc(x)^4 + \frac{2}{3} \csc(x)^3 - \csc(x)^2 - \csc(x) + \ln(\csc(x)) \right)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec), antiderivative size = 70, normalized size of antiderivative = 1.21

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx =$$

$$-\frac{60 \cos(x)^4 + 60 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(\frac{1}{2} \sin(x)) \sin(x) - 80 \cos(x)^2 - 15 (4 \cos(x)^2 - 3) \sin(x)}{60 (a \cos(x)^4 - 2 a \cos(x)^2 + a) \sin(x)}$$

input `integrate(cot(x)^7/(a+a*csc(x)),x, algorithm="fricas")`

output
$$\begin{aligned} & -\frac{1}{60} (60 \cos(x)^4 + 60 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log(\frac{1}{2} \sin(x)) \sin(x) \\ & - 80 \cos(x)^2 - 15 (4 \cos(x)^2 - 3) \sin(x) + 32) / ((a \cos(x)^4 - 2 a \cos(x)^2 + a) \sin(x)) \end{aligned}$$

Sympy [F]

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx = \frac{\int \frac{\cot^7(x)}{\csc(x)+1} dx}{a}$$

input `integrate(cot(x)**7/(a+a*csc(x)),x)`

output `Integral(cot(x)**7/(csc(x) + 1), x)/a`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.72

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx = -\frac{\log(\sin(x))}{a} - \frac{60 \sin(x)^4 + 60 \sin(x)^3 - 40 \sin(x)^2 - 15 \sin(x) + 12}{60 a \sin(x)^5}$$

input `integrate(cot(x)^7/(a+a*csc(x)),x, algorithm="maxima")`

output `-log(sin(x))/a - 1/60*(60*sin(x)^4 + 60*sin(x)^3 - 40*sin(x)^2 - 15*sin(x) + 12)/(a*sin(x)^5)`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.71

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx = -\frac{\frac{60 \sin(x)^4 + 60 \sin(x)^3 - 40 \sin(x)^2 - 15 \sin(x) + 12}{\sin(x)^5} + 60 \log(|\sin(x)|)}{60 a}$$

input `integrate(cot(x)^7/(a+a*csc(x)),x, algorithm="giac")`

output `-1/60*((60*sin(x)^4 + 60*sin(x)^3 - 40*sin(x)^2 - 15*sin(x) + 12)/sin(x)^5 + 60*log(abs(sin(x))))/a`

Mupad [B] (verification not implemented)

Time = 15.57 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.95

$$\int \frac{\cot^7(x)}{a + a \csc(x)} dx = -\frac{180 \tan(\frac{x}{2})^3 - 960 \tan(\frac{x}{2})^5 \ln\left(\tan(\frac{x}{2})^2 + 1\right) - 50 \tan(\frac{x}{2})^2 - 15 \tan(\frac{x}{2}) + 300 \tan(\frac{x}{2})^4 + 300 \tan(\frac{x}{2})^6}{960 a \tan(\frac{x}{2})^5}$$

input `int(cot(x)^7/(a + a/sin(x)),x)`

output
$$\begin{aligned} & -(180\tan(x/2)^3 - 960\tan(x/2)^5\log(\tan(x/2)^2 + 1) - 50\tan(x/2)^2 - 15 \\ & * \tan(x/2) + 300\tan(x/2)^4 + 300\tan(x/2)^6 + 180\tan(x/2)^7 - 50\tan(x/2)^8 - 15\tan(x/2)^9 + 6\tan(x/2)^{10} + 960\tan(x/2)^5\log(\tan(x/2)) + 6)/(96 \\ & 0*a\tan(x/2)^5) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.12

$$\begin{aligned} & \int \frac{\cot^7(x)}{a + a \csc(x)} dx \\ & = \frac{480 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^5 - 480 \log(\tan(\frac{x}{2})) \sin(x)^5 + 195 \sin(x)^5 - 480 \sin(x)^4 - 480 \sin(x)^3 +}{480 \sin(x)^5 a} \end{aligned}$$

input `int(cot(x)^7/(a+a*csc(x)),x)`

output
$$\begin{aligned} & (480*\log(\tan(x/2)^2 + 1)*\sin(x)^5 - 480*\log(\tan(x/2))*\sin(x)^5 + 195*\sin(x)^5 - 480*\sin(x)^4 - 480*\sin(x)^3 + 320*\sin(x)^2 + 120*\sin(x) - 96) \\ & /(480*\sin(x)^5*a) \end{aligned}$$

3.12 $\int \frac{\tan^5(x)}{a+b \csc(x)} dx$

Optimal result	105
Mathematica [A] (verified)	106
Rubi [A] (verified)	106
Maple [A] (verified)	108
Fricas [A] (verification not implemented)	108
Sympy [F]	109
Maxima [A] (verification not implemented)	109
Giac [A] (verification not implemented)	110
Mupad [B] (verification not implemented)	110
Reduce [B] (verification not implemented)	111

Optimal result

Integrand size = 13, antiderivative size = 178

$$\begin{aligned} \int \frac{\tan^5(x)}{a+b \csc(x)} dx = & \frac{1}{16(a+b)(1-\csc(x))^2} + \frac{5a+7b}{16(a+b)^2(1-\csc(x))} \\ & + \frac{1}{16(a-b)(1+\csc(x))^2} + \frac{5a-7b}{16(a-b)^2(1+\csc(x))} \\ & - \frac{(8a^2+21ab+15b^2)\log(1-\csc(x))}{16(a+b)^3} \\ & - \frac{(8a^2-21ab+15b^2)\log(1+\csc(x))}{16(a-b)^3} \\ & + \frac{b^6\log(a+b \csc(x))}{a(a^2-b^2)^3} - \frac{\log(\sin(x))}{a} \end{aligned}$$

output

```
1/16/(a+b)/(1-csc(x))^2+1/16*(5*a+7*b)/(a+b)^2/(1-csc(x))+1/16/(a-b)/(1+csc(x))^2+1/16*(5*a-7*b)/(a-b)^2/(1+csc(x))-1/16*(8*a^2+21*a*b+15*b^2)*ln(1-csc(x))/(a+b)^3-1/16*(8*a^2-21*a*b+15*b^2)*ln(1+csc(x))/(a-b)^3+b^6*ln(a+b*csc(x))/a/(a^2-b^2)^3-ln(sin(x))/a
```

Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.97

$$\int \frac{\tan^5(x)}{a + b \csc(x)} dx \\ = \frac{\csc(x)(b + a \sin(x)) \left(-\frac{(8a^2+21ab+15b^2) \log(1-\sin(x))}{(a+b)^3} - \frac{(8a^2-21ab+15b^2) \log(1+\sin(x))}{(a-b)^3} + \frac{16b^6 \log(b+a \sin(x))}{a(a-b)^3(a+b)^3} + \frac{1}{(a+b)(-1+a \sin(x))^3} \right)}{16(a + b \csc(x))}$$

input `Integrate[Tan[x]^5/(a + b*Csc[x]), x]`

output $(\csc(x)*(b + a*\sin(x))*(-(8*a^2 + 21*a*b + 15*b^2)*\log[1 - \sin(x])/(a + b)^3 - (8*a^2 - 21*a*b + 15*b^2)*\log[1 + \sin(x)]/(a - b)^3 + (16*b^6*\log[b + a*\sin(x])/(a*(a - b)^3*(a + b)^3) + 1/((a + b)*(-1 + \sin(x))^2) + (7*a + 9*b)/((a + b)^2*(-1 + \sin(x))) + 1/((a - b)*(1 + \sin(x))^2) + (-7*a + 9*b)/((a - b)^2*(1 + \sin(x)))))/(16*(a + b*\csc(x)))$

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.18, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 4373, 615, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^5(x)}{a + b \csc(x)} dx \\ & \downarrow 3042 \\ & \int \frac{1}{\cot(x)^5(a + b \csc(x))} dx \\ & \downarrow 4373 \\ & b^6 \int \frac{\sin(x)}{b(a + b \csc(x))(b^2 - b^2 \csc^2(x))^3} d(b \csc(x)) \\ & \downarrow 615 \end{aligned}$$

$$b^6 \int \left(\frac{7b - 5a}{16(a-b)^2 b^5 (\csc(x)b + b)^2} + \frac{\sin(x)}{ab^7} + \frac{8a^2 + 21ba + 15b^2}{16b^6(a+b)^3(b - b \csc(x))} + \frac{1}{a(a-b)^3(a+b)^3(a+b \csc(x))} + \frac{1}{16b^6(a-b)^3(a+b \csc(x))^2} \right) dx$$

\downarrow 2009

$$b^6 \left(\frac{\log(a + b \csc(x))}{a (a^2 - b^2)^3} - \frac{(8a^2 + 21ab + 15b^2) \log(b - b \csc(x))}{16b^6(a+b)^3} - \frac{(8a^2 - 21ab + 15b^2) \log(b \csc(x) + b)}{16b^6(a-b)^3} + \frac{\log(b \csc(x) + b)}{ab^5} \right)$$

input `Int[Tan[x]^5/(a + b*Csc[x]), x]`

output $b^{12} \left(\frac{1}{16b^4(a+b)^2} + \frac{5a+7b}{16b^5(a+b)^2} + \frac{1}{16b^4(a-b)^2} + \frac{5a-7b}{16b^5(a-b)^2} + \frac{\operatorname{Log}[b \csc(x)]}{a^2 b^6} - \frac{(8a^2 + 21ab + 15b^2) \operatorname{Log}[b - b \csc(x)]}{16b^6(a+b)^3} + \frac{\operatorname{Log}[a + b \csc(x)]}{a^2 (a^2 - b^2)^3} - \frac{(8a^2 - 21ab + 15b^2) \operatorname{Log}[b + b \csc(x)]}{16b^6(a-b)^3} - \frac{(8a^2 - 21ab + 15b^2) \operatorname{Log}[b - b \csc(x)]}{16b^6(a-b)^3} \right)$

Definitions of rubi rules used

rule 615 `Int[((e_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x]; FreeQ[{a, b, c, d, e, m, n}, x] && ILtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4373 `Int[cot[(c_)*(x_)]^(m_)*csc[(c_)*(x_)]*(b_)*(a_))^(n_), x_Symbol] :> Simp[-(-1)^((m - 1)/2)/(d*b^(m - 1)) Subst[Int[(b^2 - x^2)^((m - 1)/2)*((a + x)^n/x), x], x, b*Csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]`

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.90

method	result
default	$\frac{1}{2(8a-8b)(1+\sin(x))^2} - \frac{7a-9b}{16(a-b)^2(1+\sin(x))} + \frac{(-8a^2+21ba-15b^2)\ln(1+\sin(x))}{16(a-b)^3} + \frac{b^6\ln(a\sin(x)+b)}{(a+b)^3(a-b)^3a} + \frac{1}{2(8a+8b)(\sin(x)-1)^2}$
risch	$-\frac{21ixa}{8(a^3-3a^2b+3ab^2-b^3)} + \frac{ixa^2}{a^3+3a^2b+3ab^2+b^3} + \frac{21ixa}{8(a^3+3a^2b+3ab^2+b^3)} - \frac{2ixb^6}{a(a^6-3a^4b^2+3a^2b^4-b^6)} + \frac{15ixb^2}{8(a^3-3a^2b+3ab^2-b^3)}$

input `int(tan(x)^5/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/2/(8*a-8*b)/(1+\sin(x))^2 - 1/16*(7*a-9*b)/(a-b)^2/(1+\sin(x)) + 1/16/(a-b)^3* \\ & (-8*a^2+21*a*b-15*b^2)*\ln(1+\sin(x))+b^6/(a+b)^3/(a-b)^3/a*\ln(a*\sin(x)+b)+1 \\ & /2/(8*a+8*b)/(sin(x)-1)^2 - 1/16*(-7*a-9*b)/(a+b)^2/(sin(x)-1) + 1/16/(a+b)^3* \\ & (-8*a^2-21*a*b-15*b^2)*\ln(sin(x)-1) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.46

$$\begin{aligned} & \int \frac{\tan^5(x)}{a+b\csc(x)} dx \\ & = \frac{16 b^6 \cos(x)^4 \log(a \sin(x) + b) + 4 a^6 - 8 a^4 b^2 + 4 a^2 b^4 - (8 a^6 + 3 a^5 b - 24 a^4 b^2 - 10 a^3 b^3 + 24 a^2 b^4 + 1)}{1} \end{aligned}$$

input `integrate(tan(x)^5/(a+b*csc(x)),x, algorithm="fricas")`

output
$$\begin{aligned} & 1/16*(16*b^6*cos(x)^4*log(a*sin(x) + b) + 4*a^6 - 8*a^4*b^2 + 4*a^2*b^4 - \\ & (8*a^6 + 3*a^5*b - 24*a^4*b^2 - 10*a^3*b^3 + 24*a^2*b^4 + 15*a*b^5)*cos(x) \\ & ^4*log(sin(x) + 1) - (8*a^6 - 3*a^5*b - 24*a^4*b^2 + 10*a^3*b^3 + 24*a^2*b^4 - \\ & 15*a*b^5)*cos(x)^4*log(-sin(x) + 1) - 8*(2*a^6 - 5*a^4*b^2 + 3*a^2*b^4)*cos(x)^2 - \\ & 2*(2*a^5*b - 4*a^3*b^3 + 2*a*b^5 - (5*a^5*b - 14*a^3*b^3 + 9 \\ & *a*b^5)*cos(x)^2)*sin(x))/((a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6)*cos(x)^4) \end{aligned}$$

Sympy [F]

$$\int \frac{\tan^5(x)}{a + b \csc(x)} dx = \int \frac{\tan^5(x)}{a + b \csc(x)} dx$$

input `integrate(tan(x)**5/(a+b*csc(x)),x)`

output `Integral(tan(x)**5/(a + b*csc(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.41

$$\begin{aligned} \int \frac{\tan^5(x)}{a + b \csc(x)} dx &= \frac{b^6 \log(a \sin(x) + b)}{a^7 - 3a^5b^2 + 3a^3b^4 - ab^6} \\ &- \frac{(8a^2 - 21ab + 15b^2) \log(\sin(x) + 1)}{16(a^3 - 3a^2b + 3ab^2 - b^3)} - \frac{(8a^2 + 21ab + 15b^2) \log(\sin(x) - 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} \\ &- \frac{(5a^2b - 9b^3) \sin(x)^3 + 6a^3 - 10ab^2 - 4(2a^3 - 3ab^2) \sin(x)^2 - (3a^2b - 7b^3) \sin(x)}{8((a^4 - 2a^2b^2 + b^4) \sin(x)^4 + a^4 - 2a^2b^2 + b^4 - 2(a^4 - 2a^2b^2 + b^4) \sin(x)^2)} \end{aligned}$$

input `integrate(tan(x)^5/(a+b*csc(x)),x, algorithm="maxima")`

output `b^6*log(a*sin(x) + b)/(a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6) - 1/16*(8*a^2 - 21*a*b + 15*b^2)*log(sin(x) + 1)/(a^3 - 3*a^2*b + 3*a*b^2 - b^3) - 1/16*(8*a^2 + 21*a*b + 15*b^2)*log(sin(x) - 1)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 1/8*((5*a^2*b - 9*b^3)*sin(x)^3 + 6*a^3 - 10*a*b^2 - 4*(2*a^3 - 3*a*b^2)*sin(x)^2 - (3*a^2*b - 7*b^3)*sin(x))/((a^4 - 2*a^2*b^2 + b^4)*sin(x)^4 + a^4 - 2*a^2*b^2 + b^4 - 2*(a^4 - 2*a^2*b^2 + b^4)*sin(x)^2)`

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.42

$$\int \frac{\tan^5(x)}{a + b \csc(x)} dx = \frac{b^6 \log(|a \sin(x) + b|)}{a^7 - 3a^5b^2 + 3a^3b^4 - ab^6} - \frac{(8a^2 - 21ab + 15b^2) \log(\sin(x) + 1)}{16(a^3 - 3a^2b + 3ab^2 - b^3)} - \frac{(8a^2 + 21ab + 15b^2) \log(-\sin(x) + 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} - \frac{6a^5 - 16a^3b^2 + 10ab^4 + (5a^4b - 14a^2b^3 + 9b^5) \sin(x)^3 - 4(2a^5 - 5a^3b^2 + 3ab^4) \sin(x)^2 - (3a^4b - 10a^2b^3 + 7b^5) \sin(x)}{8(a+b)^3(a-b)^3(\sin(x)+1)^2(\sin(x)-1)^2}$$

input `integrate(tan(x)^5/(a+b*csc(x)),x, algorithm="giac")`

output $b^{6} \log(\left|a \sin(x) + b\right|)/(a^7 - 3a^5b^2 + 3a^3b^4 - ab^6) - 1/16*(8*a^2 - 21*a*b + 15*b^2)*\log(\sin(x) + 1)/(a^3 - 3a^2*b + 3*a*b^2 - b^3) - 1/16*(8*a^2 + 21*a*b + 15*b^2)*\log(-\sin(x) + 1)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 1/8*(6*a^5 - 16*a^3*b^2 + 10*a*b^4 + (5*a^4*b - 14*a^2*b^3 + 9*b^5)*\sin(x)^3 - 4*(2*a^5 - 5*a^3*b^2 + 3*a*b^4)*\sin(x)^2 - (3*a^4*b - 10*a^2*b^3 + 7*b^5)*\sin(x))/((a + b)^3*(a - b)^3*(\sin(x) + 1)^2*(\sin(x) - 1)^2)$

Mupad [B] (verification not implemented)

Time = 17.29 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.50

$$\int \frac{\tan^5(x)}{a + b \csc(x)} dx = \frac{\frac{2 \tan(\frac{x}{2})^2 (2 a b^2 - a^3)}{a^4 - 2 a^2 b^2 + b^4} - \frac{4 \tan(\frac{x}{2})^4 (3 a b^2 - 2 a^3)}{a^4 - 2 a^2 b^2 + b^4} + \frac{2 \tan(\frac{x}{2})^6 (2 a b^2 - a^3)}{a^4 - 2 a^2 b^2 + b^4} + \frac{\tan(\frac{x}{2})^7 (3 a^2 b - 7 b^3)}{4 (a^4 - 2 a^2 b^2 + b^4)} - \frac{\tan(\frac{x}{2})^3 (11 a^2 b - 15 b^3)}{4 (a^4 - 2 a^2 b^2 + b^4)} - \frac{\tan(\frac{x}{2})^8 - 4 \tan(\frac{x}{2})^6 + 6 \tan(\frac{x}{2})^4 - 4 \tan(\frac{x}{2})^2 + 1}{\tan(\frac{x}{2})^8 - 4 \tan(\frac{x}{2})^6 + 6 \tan(\frac{x}{2})^4 - 4 \tan(\frac{x}{2})^2 + 1} - \ln\left(\tan\left(\frac{x}{2}\right) - 1\right) \left(\frac{5 b}{8 (a + b)^2} + \frac{1}{a + b} + \frac{b^2}{4 (a + b)^3}\right) - \ln\left(\tan\left(\frac{x}{2}\right) + 1\right) \left(\frac{b^2}{4 (a - b)^3} - \frac{5 b}{8 (a - b)^2} + \frac{1}{a - b}\right) + \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{b^6 \ln\left(b \tan\left(\frac{x}{2}\right)^2 + 2 a \tan\left(\frac{x}{2}\right) + b\right)}{-a^7 + 3 a^5 b^2 - 3 a^3 b^4 + a b^6}}$$

input `int(tan(x)^5/(a + b/sin(x)),x)`

output

$$\begin{aligned} & ((2*\tan(x/2)^2*(2*a*b^2 - a^3))/(a^4 + b^4 - 2*a^2*b^2) - (4*\tan(x/2)^4*(3*a*b^2 - 2*a^3))/(a^4 + b^4 - 2*a^2*b^2) + (2*\tan(x/2)^6*(2*a*b^2 - a^3))/(a^4 + b^4 - 2*a^2*b^2) + (\tan(x/2)^7*(3*a^2*b - 7*b^3))/(4*(a^4 + b^4 - 2*a^2*b^2)) - (\tan(x/2)^3*(11*a^2*b - 15*b^3))/(4*(a^4 + b^4 - 2*a^2*b^2)) - (\tan(x/2)^5*(11*a^2*b - 15*b^3))/(4*(a^4 + b^4 - 2*a^2*b^2)) + (b*\tan(x/2)*(3*a^2 - 7*b^2))/(4*(a^4 + b^4 - 2*a^2*b^2)))/(6*\tan(x/2)^4 - 4*\tan(x/2)^2 - 4*\tan(x/2)^6 + \tan(x/2)^8 + 1) - \log(\tan(x/2) - 1)*((5*b)/(8*(a + b)^2) + 1/(a + b) + b^2/(4*(a + b)^3)) - \log(\tan(x/2) + 1)*(b^2/(4*(a - b)^3) - (5*b)/(8*(a - b)^2) + 1/(a - b)) + \log(\tan(x/2)^2 + 1)/a - (b^6*\log(b + 2*a*\tan(x/2) + b*\tan(x/2)^2))/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.19 (sec), antiderivative size = 1150, normalized size of antiderivative = 6.46

$$\int \frac{\tan^5(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input

int(tan(x)^5/(a+b*csc(x)),x)

output

```
(8*log(tan(x/2)**2 + 1)*sin(x)**4*a**6 - 24*log(tan(x/2)**2 + 1)*sin(x)**4*a**4*b**2 + 24*log(tan(x/2)**2 + 1)*sin(x)**4*a**2*b**4 - 8*log(tan(x/2)*  
*2 + 1)*sin(x)**4*b**6 - 16*log(tan(x/2)**2 + 1)*sin(x)**2*a**6 + 48*log(t  
an(x/2)**2 + 1)*sin(x)**2*a**4*b**2 - 48*log(tan(x/2)**2 + 1)*sin(x)**2*a*  
*2*b**4 + 16*log(tan(x/2)**2 + 1)*sin(x)**2*b**6 + 8*log(tan(x/2)**2 + 1)*  
a**6 - 24*log(tan(x/2)**2 + 1)*a**4*b**2 + 24*log(tan(x/2)**2 + 1)*a**2*b*  
*4 - 8*log(tan(x/2)**2 + 1)*b**6 - 8*log(tan(x/2) - 1)*sin(x)**4*a**6 + 3*  
log(tan(x/2) - 1)*sin(x)**4*a**5*b + 24*log(tan(x/2) - 1)*sin(x)**4*a**4*b  
**2 - 10*log(tan(x/2) - 1)*sin(x)**4*a**3*b**3 - 24*log(tan(x/2) - 1)*sin(  
x)**4*a**2*b**4 + 15*log(tan(x/2) - 1)*sin(x)**4*a*b**5 + 16*log(tan(x/2)  
- 1)*sin(x)**2*a**6 - 6*log(tan(x/2) - 1)*sin(x)**2*a**5*b - 48*log(tan(x/  
2) - 1)*sin(x)**2*a**4*b**2 + 20*log(tan(x/2) - 1)*sin(x)**2*a**3*b**3 + 4  
8*log(tan(x/2) - 1)*sin(x)**2*a**2*b**4 - 30*log(tan(x/2) - 1)*sin(x)**2*a  
*b**5 - 8*log(tan(x/2) - 1)*a**6 + 3*log(tan(x/2) - 1)*a**5*b + 24*log(tan  
(x/2) - 1)*a**4*b**2 - 10*log(tan(x/2) - 1)*a**3*b**3 - 24*log(tan(x/2) -  
1)*a**2*b**4 + 15*log(tan(x/2) - 1)*a*b**5 - 8*log(tan(x/2) + 1)*sin(x)**4  
*a**6 - 3*log(tan(x/2) + 1)*sin(x)**4*a**5*b + 24*log(tan(x/2) + 1)*sin(x)  
**4*a**4*b**2 + 10*log(tan(x/2) + 1)*sin(x)**4*a**3*b**3 - 24*log(tan(x/2)  
+ 1)*sin(x)**4*a**2*b**4 - 15*log(tan(x/2) + 1)*sin(x)**4*a*b**5 + 16*log  
(tan(x/2) + 1)*sin(x)**2*a**6 + 6*log(tan(x/2) + 1)*sin(x)**2*a**5*b - ...
```

3.13 $\int \frac{\tan^3(x)}{a+b \csc(x)} dx$

Optimal result	113
Mathematica [A] (verified)	113
Rubi [A] (verified)	114
Maple [A] (verified)	115
Fricas [A] (verification not implemented)	116
Sympy [F]	116
Maxima [A] (verification not implemented)	117
Giac [A] (verification not implemented)	117
Mupad [B] (verification not implemented)	118
Reduce [B] (verification not implemented)	118

Optimal result

Integrand size = 13, antiderivative size = 113

$$\begin{aligned} \int \frac{\tan^3(x)}{a + b \csc(x)} dx = & -\frac{1}{4(a+b)(1-\csc(x))} - \frac{1}{4(a-b)(1+\csc(x))} \\ & + \frac{(2a+3b)\log(1-\csc(x))}{4(a+b)^2} + \frac{(2a-3b)\log(1+\csc(x))}{4(a-b)^2} \\ & + \frac{b^4 \log(a+b \csc(x))}{a (a^2-b^2)^2} + \frac{\log(\sin(x))}{a} \end{aligned}$$

output
$$-1/4/(a+b)/(1-\csc(x))-1/4/(a-b)/(1+\csc(x))+1/4*(2*a+3*b)*\ln(1-\csc(x))/(a+b)^2+1/4*(2*a-3*b)*\ln(1+\csc(x))/(a-b)^2+b^4*\ln(a+b*csc(x))/a/(a^2-b^2)^2+ln(\sin(x))/a$$

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.02

$$\begin{aligned} & \int \frac{\tan^3(x)}{a + b \csc(x)} dx \\ &= \frac{\csc(x)(b + a \sin(x)) \left(\frac{(2a+3b)\log(1-\sin(x))}{(a+b)^2} + \frac{(2a-3b)\log(1+\sin(x))}{(a-b)^2} + \frac{4b^4 \log(b+a \sin(x))}{a(a-b)^2(a+b)^2} - \frac{1}{(a+b)(-1+\sin(x))} + \frac{1}{(a-b)(1+\sin(x))} \right)}{4(a + b \csc(x))} \end{aligned}$$

input $\text{Integrate}[\tan[x]^3/(a + b*\csc[x]), x]$

output $(\csc[x]*(b + a*\sin[x])*((2*a + 3*b)*\log[1 - \sin[x]])/(a + b)^2 + ((2*a - 3*b)*\log[1 + \sin[x]])/(a - b)^2 + (4*b^4*\log[b + a*\sin[x]])/(a*(a - b)^2*(a + b)^2) - 1/((a + b)*(-1 + \sin[x])) + 1/((a - b)*(1 + \sin[x]))))/(4*(a + b*\csc[x]))$

Rubi [A] (verified)

Time = 0.39 (sec), antiderivative size = 139, normalized size of antiderivative = 1.23, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 4373, 615, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\cot(x)^3(a + b \csc(x))} dx \\
 & \quad \downarrow \text{4373} \\
 & -b^4 \int \frac{\sin(x)}{b(a + b \csc(x))(b^2 - b^2 \csc^2(x))^2} d(b \csc(x)) \\
 & \quad \downarrow \text{615} \\
 & -b^4 \int \left(\frac{3b - 2a}{4(a - b)^2 b^4 (\csc(x) b + b)} + \frac{\sin(x)}{ab^5} + \frac{2a + 3b}{4b^4 (a + b)^2 (b - b \csc(x))} - \frac{1}{a(a - b)^2 (a + b)^2 (a + b \csc(x))} + \frac{1}{4b^3 (a + b)^3} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & -b^4 \left(-\frac{\log(a + b \csc(x))}{a (a^2 - b^2)^2} + \frac{\log(b \csc(x))}{ab^4} - \frac{(2a + 3b) \log(b - b \csc(x))}{4b^4 (a + b)^2} - \frac{(2a - 3b) \log(b \csc(x) + b)}{4b^4 (a - b)^2} + \frac{1}{4b^3 (a + b)^3} \right)
 \end{aligned}$$

input $\text{Int}[\tan[x]^3/(a + b*\csc[x]), x]$

output

$$-(b^4*(1/(4*b^3*(a+b)*(b-b*Csc[x])) + 1/(4*(a-b)*b^3*(b+b*Csc[x]))) + Log[b*Csc[x]]/(a*b^4) - ((2*a+3*b)*Log[b-b*Csc[x]])/(4*b^4*(a+b)^2) - Log[a+b*Csc[x]]/(a*(a^2-b^2)^2) - ((2*a-3*b)*Log[b+b*Csc[x]])/(4*(a-b)^2*b^4)))$$

Definitions of rubi rules used

rule 615 $\text{Int}[(e_*)(x_())^{(m_())}((c_()) + (d_*)(x_())^{(n_())}((a_()) + (b_())*(x_())^2)^{(p_())}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(e*x)^m*(c+d*x)^n*(a+b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \&& \text{ILtQ}[p, 0]$

rule 2009 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4373 $\text{Int}[\cot[(c_()) + (d_*)(x_())^{(m_())}(\csc[(c_()) + (d_*)(x_())]*(b_()) + (a_())^{(n_())}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-(-1)^{(m-1)/2}/(d*b^{(m-1)}) \text{Subst}[\text{Int}[(b^2-x^2)^{((m-1)/2)}*((a+x)^n/x), x], x, b*Csc[c+d*x]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&& \text{IntegerQ}[(m-1)/2] \&& \text{NeQ}[a^2-b^2, 0]$

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.89

method	result
default	$\frac{1}{(4a-4b)(1+\sin(x))} + \frac{(2a-3b)\ln(1+\sin(x))}{4(a-b)^2} + \frac{b^4\ln(a\sin(x)+b)}{(a+b)^2(a-b)^2a} - \frac{1}{(4a+4b)(\sin(x)-1)} + \frac{(2a+3b)\ln(\sin(x)-1)}{4(a+b)^2}$
risch	$\frac{ix}{a} - \frac{ixa}{a^2-2ba+b^2} + \frac{3ixb}{2(a^2-2ba+b^2)} - \frac{ixa}{a^2+2ba+b^2} - \frac{3ixb}{2(a^2+2ba+b^2)} - \frac{2ixb^4}{a(a^4-2a^2b^2+b^4)} + \frac{i(-2ia e^{2ix}+b e^{3ix}-b e^{ix})}{(e^{2ix}+1)^2(a^2-b^2)}$

input $\text{int}(\tan(x)^3/(a+b*csc(x)), x, \text{method}=\text{_RETURNVERBOSE})$

output
$$\frac{1/(4*a-4*b)/(1+\sin(x))+1/4*(2*a-3*b)/(a-b)^2*\ln(1+\sin(x))+b^4/(a+b)^2/(a-b)^2/a*\ln(a*\sin(x)+b)-1/(4*a+4*b)/(sin(x)-1)+1/4*(2*a+3*b)/(a+b)^2*\ln(\sin(x)-1)}$$

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.27

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx \\ = \frac{4 b^4 \cos(x)^2 \log(a \sin(x) + b) + 2 a^4 - 2 a^2 b^2 + (2 a^4 + a^3 b - 4 a^2 b^2 - 3 a b^3) \cos(x)^2 \log(\sin(x) + 1) + (4 a^5 - 2 a^3 b^2 + a b^4) \cos(x)^2}{4 (a^5 - 2 a^3 b^2 + a b^4) \cos(x)}$$

input `integrate(tan(x)^3/(a+b*csc(x)),x, algorithm="fricas")`

output
$$\frac{1/4*(4*b^4*cos(x)^2*log(a*sin(x) + b) + 2*a^4 - 2*a^2*b^2 + (2*a^4 + a^3*b - 4*a^2*b^2 - 3*a*b^3)*cos(x)^2*log(sin(x) + 1) + (2*a^4 - a^3*b - 4*a^2*b^2 + 3*a*b^3)*cos(x)^2*log(-sin(x) + 1) - 2*(a^3*b - a*b^3)*sin(x))/((a^5 - 2*a^3*b^2 + a*b^4)*cos(x)^2)}$$

Sympy [F]

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx = \int \frac{\tan^3(x)}{a + b \csc(x)} dx$$

input `integrate(tan(x)**3/(a+b*csc(x)),x)`

output `Integral(tan(x)**3/(a + b*csc(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.06

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx = \frac{b^4 \log(a \sin(x) + b)}{a^5 - 2a^3b^2 + ab^4} + \frac{(2a - 3b) \log(\sin(x) + 1)}{4(a^2 - 2ab + b^2)} \\ + \frac{(2a + 3b) \log(\sin(x) - 1)}{4(a^2 + 2ab + b^2)} + \frac{b \sin(x) - a}{2((a^2 - b^2) \sin(x)^2 - a^2 + b^2)}$$

input `integrate(tan(x)^3/(a+b*csc(x)),x, algorithm="maxima")`

output $b^{4 \log(a \sin(x) + b) / (a^5 - 2a^3b^2 + ab^4)} + 1/4*(2*a - 3*b)*\log(\sin(x) + 1) / (a^2 - 2*a*b + b^2) + 1/4*(2*a + 3*b)*\log(\sin(x) - 1) / (a^2 + 2*a*b + b^2) + 1/2*(b*\sin(x) - a) / ((a^2 - b^2)*\sin(x)^2 - a^2 + b^2)$

Giac [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.23

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx = \frac{b^4 \log(|a \sin(x) + b|)}{a^5 - 2a^3b^2 + ab^4} + \frac{(2a - 3b) \log(\sin(x) + 1)}{4(a^2 - 2ab + b^2)} \\ + \frac{(2a + 3b) \log(-\sin(x) + 1)}{4(a^2 + 2ab + b^2)} \\ - \frac{a^3 - ab^2 - (a^2b - b^3) \sin(x)}{2(a + b)^2(a - b)^2(\sin(x) + 1)(\sin(x) - 1)}$$

input `integrate(tan(x)^3/(a+b*csc(x)),x, algorithm="giac")`

output $b^{4 \log(\text{abs}(a \sin(x) + b)) / (a^5 - 2a^3b^2 + ab^4)} + 1/4*(2*a - 3*b)*\log(\sin(x) + 1) / (a^2 - 2*a*b + b^2) + 1/4*(2*a + 3*b)*\log(-\sin(x) + 1) / (a^2 + 2*a*b + b^2) - 1/2*(a^3 - a*b^2 - (a^2b - b^3)*\sin(x)) / ((a + b)^2(a - b)^2*(\sin(x) + 1)*(\sin(x) - 1))$

Mupad [B] (verification not implemented)

Time = 16.00 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.52

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx = \frac{\ln(\tan(\frac{x}{2}) + 1) (2a - 3b)}{2(a - b)^2} - \frac{\frac{b \tan(\frac{x}{2})^3}{a^2 - b^2} - \frac{2a \tan(\frac{x}{2})^2}{a^2 - b^2} + \frac{b \tan(\frac{x}{2})}{a^2 - b^2}}{\tan(\frac{x}{2})^4 - 2 \tan(\frac{x}{2})^2 + 1}$$

$$- \frac{\ln(\tan(\frac{x}{2})^2 + 1)}{a} + \frac{\ln(\tan(\frac{x}{2}) - 1) (2a + 3b)}{2(a + b)^2}$$

$$+ \frac{b^4 \ln(b \tan(\frac{x}{2})^2 + 2a \tan(\frac{x}{2}) + b)}{a (a^2 - b^2)^2}$$

input `int(tan(x)^3/(a + b/sin(x)),x)`

output `(log(tan(x/2) + 1)*(2*a - 3*b))/(2*(a - b)^2) - ((b*tan(x/2)^3)/(a^2 - b^2) - (2*a*tan(x/2)^2)/(a^2 - b^2) + (b*tan(x/2))/(a^2 - b^2))/((tan(x/2)^4 - 2*tan(x/2)^2 + 1) - log(tan(x/2)^2 + 1)/a + (log(tan(x/2) - 1)*(2*a + 3*b))/(2*(a + b)^2) + (b^4*log(b + 2*a*tan(x/2) + b*tan(x/2)^2))/(a*(a^2 - b^2)^2))`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 497, normalized size of antiderivative = 4.40

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `int(tan(x)^3/(a+b*csc(x)),x)`

output

```
( - 2*log(tan(x/2)**2 + 1)*sin(x)**2*a**4 + 4*log(tan(x/2)**2 + 1)*sin(x)*  
*2*a**2*b**2 - 2*log(tan(x/2)**2 + 1)*sin(x)**2*b**4 + 2*log(tan(x/2)**2 +  
1)*a**4 - 4*log(tan(x/2)**2 + 1)*a**2*b**2 + 2*log(tan(x/2)**2 + 1)*b**4  
+ 2*log(tan(x/2) - 1)*sin(x)**2*a**4 - log(tan(x/2) - 1)*sin(x)**2*a**3*b  
- 4*log(tan(x/2) - 1)*sin(x)**2*a**2*b**2 + 3*log(tan(x/2) - 1)*sin(x)**2*  
a*b**3 - 2*log(tan(x/2) - 1)*a**4 + log(tan(x/2) - 1)*a**3*b + 4*log(tan(x/  
2) - 1)*a**2*b**2 - 3*log(tan(x/2) - 1)*a*b**3 + 2*log(tan(x/2) + 1)*sin(  
x)**2*a**4 + log(tan(x/2) + 1)*sin(x)**2*a**3*b - 4*log(tan(x/2) + 1)*sin(  
x)**2*a**2*b**2 - 3*log(tan(x/2) + 1)*sin(x)**2*a*b**3 - 2*log(tan(x/2) +  
1)*a**4 - log(tan(x/2) + 1)*a**3*b + 4*log(tan(x/2) + 1)*a**2*b**2 + 3*log(  
(tan(x/2) + 1)*a*b**3 + 2*log(tan(x/2)**2*b + 2*tan(x/2)*a + b)*sin(x)**2*  
b**4 - 2*log(tan(x/2)**2*b + 2*tan(x/2)*a + b)*b**4 + sin(x)**2*a**4 - sin(  
(x)**2*a**2*b**2 + sin(x)*a**3*b - sin(x)*a*b**3 - 2*a**4 + 2*a**2*b**2)/(  
2*a*(sin(x)**2*a**4 - 2*sin(x)**2*a**2*b**2 + sin(x)**2*b**4 - a**4 + 2*a*  
*2*b**2 - b**4))
```

3.14 $\int \frac{\tan(x)}{a+b \csc(x)} dx$

Optimal result	120
Mathematica [A] (verified)	120
Rubi [A] (verified)	121
Maple [A] (verified)	122
Fricas [A] (verification not implemented)	123
Sympy [F]	123
Maxima [A] (verification not implemented)	123
Giac [A] (verification not implemented)	124
Mupad [B] (verification not implemented)	124
Reduce [B] (verification not implemented)	124

Optimal result

Integrand size = 11, antiderivative size = 66

$$\begin{aligned} \int \frac{\tan(x)}{a + b \csc(x)} dx = & -\frac{\log(1 - \csc(x))}{2(a + b)} - \frac{\log(1 + \csc(x))}{2(a - b)} \\ & + \frac{b^2 \log(a + b \csc(x))}{a(a^2 - b^2)} - \frac{\log(\sin(x))}{a} \end{aligned}$$

output
$$-\frac{1}{2} \ln(1 - \csc(x)) / (a + b) - \ln(1 + \csc(x)) / (2a - 2b) + b^2 \ln(a + b \csc(x)) / a / (a^2 - b^2) - \ln(\sin(x)) / a$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.85

$$\begin{aligned} \int \frac{\tan(x)}{a + b \csc(x)} dx \\ = -\frac{a(a - b) \log(1 - \sin(x)) + a(a + b) \log(1 + \sin(x)) - 2b^2 \log(b + a \sin(x))}{2a(a - b)(a + b)} \end{aligned}$$

input `Integrate[Tan[x]/(a + b*Csc[x]),x]`

output
$$\frac{-\frac{1}{2}a(a-b)\ln(1-\sin(x)) + a(a+b)\ln(1+\sin(x)) - 2b^2\ln(b+a\sin(x))}{a(a-b)(a+b)}$$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 4373, 615, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(x)}{a+b \csc(x)} dx \\
 & \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cot(x)(a+b \csc(x))} dx \\
 & \downarrow \textcolor{blue}{4373} \\
 & b^2 \int \frac{\sin(x)}{b(a+b \csc(x))(b^2-b^2 \csc^2(x))} d(b \csc(x)) \\
 & \downarrow \textcolor{blue}{615} \\
 & b^2 \int \left(\frac{\sin(x)}{ab^3} + \frac{1}{2b^2(a+b)(b-b \csc(x))} + \frac{1}{a(a-b)(a+b)(a+b \csc(x))} - \frac{1}{2(a-b)b^2(\csc(x)b+b)} \right) d(b \csc(x)) \\
 & \downarrow \textcolor{blue}{2009} \\
 & b^2 \left(\frac{\log(a+b \csc(x))}{a(a^2-b^2)} + \frac{\log(b \csc(x))}{ab^2} - \frac{\log(b-b \csc(x))}{2b^2(a+b)} - \frac{\log(b \csc(x)+b)}{2b^2(a-b)} \right)
 \end{aligned}$$

input
$$\text{Int}[\tan[x]/(a+b \csc[x]), x]$$

output
$$\frac{b^2 (\ln(b \csc(x))/a^2 - \ln(b - b \csc(x))/(2b^2(a+b)) + \ln(a+b \csc(x))/(a(a^2-b^2)) - \ln(b + b \csc(x))/(2(a-b)b^2))}{a^2 b^2}$$

Definitions of rubi rules used

rule 615 $\text{Int}[(e_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*))^{(n_*)}*((a_*) + (b_*)*(x_*)^2)^{(p_*)}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x]; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \& \text{ILtQ}[p, 0]$

rule 2009 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_*, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4373 $\text{Int}[\cot[(c_*) + (d_*)*(x_*)]^{(m_*)}*(\csc[(c_*) + (d_*)*(x_*)]*(b_*) + (a_*)^{(n_*)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[-(-1)^{((m - 1)/2)} / (d*b^{(m - 1)}) \text{Subst}[\text{Int}[(b^2 - x^2)^{((m - 1)/2)}*((a + x)^n/x), x], x, b*\text{Csc}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \& \text{IntegerQ}[(m - 1)/2] \& \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.91

method	result	size
default	$-\frac{\ln(1+\sin(x))}{2a-2b} + \frac{b^2 \ln(a \sin(x)+b)}{(a+b)(a-b)a} - \frac{\ln(\sin(x)-1)}{2a+2b}$	60
risch	$-\frac{ix}{a} + \frac{ix}{a+b} + \frac{ix}{a-b} - \frac{2ix b^2}{(a^2-b^2)a} - \frac{\ln(e^{ix}-i)}{a+b} - \frac{\ln(e^{ix}+i)}{a-b} + \frac{b^2 \ln(e^{2ix}-1+\frac{2ib e^{ix}}{a})}{(a^2-b^2)a}$	122

input $\text{int}(1/(a+b*\csc(x))*\tan(x), x, \text{method}=\text{RETURNVERBOSE})$

output $-\ln(1+\sin(x))/(2*a-2*b)+b^2/(a+b)/(a-b)/a*\ln(a*\sin(x)+b)-1/(2*a+2*b)*\ln(\sin(x)-1)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.88

$$\int \frac{\tan(x)}{a + b \csc(x)} dx = \frac{2 b^2 \log(a \sin(x) + b) - (a^2 + ab) \log(\sin(x) + 1) - (a^2 - ab) \log(-\sin(x) + 1)}{2(a^3 - ab^2)}$$

input `integrate(tan(x)/(a+b*csc(x)),x, algorithm="fricas")`

output $\frac{1/2*(2*b^2*log(a*sin(x) + b) - (a^2 + a*b)*log(sin(x) + 1) - (a^2 - a*b)*log(-sin(x) + 1))}{(a^3 - a*b^2)}$

Sympy [F]

$$\int \frac{\tan(x)}{a + b \csc(x)} dx = \int \frac{\tan(x)}{a + b \csc(x)} dx$$

input `integrate(tan(x)/(a+b*csc(x)),x)`

output `Integral(tan(x)/(a + b*csc(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.76

$$\int \frac{\tan(x)}{a + b \csc(x)} dx = \frac{b^2 \log(a \sin(x) + b)}{a^3 - ab^2} - \frac{\log(\sin(x) + 1)}{2(a - b)} - \frac{\log(\sin(x) - 1)}{2(a + b)}$$

input `integrate(tan(x)/(a+b*csc(x)),x, algorithm="maxima")`

output $b^2*\log(a*\sin(x) + b)/(a^3 - a*b^2) - 1/2*\log(\sin(x) + 1)/(a - b) - 1/2*\log(\sin(x) - 1)/(a + b)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.80

$$\int \frac{\tan(x)}{a + b \csc(x)} dx = \frac{b^2 \log(|a \sin(x) + b|)}{a^3 - ab^2} - \frac{\log(\sin(x) + 1)}{2(a - b)} - \frac{\log(-\sin(x) + 1)}{2(a + b)}$$

input `integrate(tan(x)/(a+b*csc(x)),x, algorithm="giac")`

output `b^2*log(abs(a*sin(x) + b))/(a^3 - a*b^2) - 1/2*log(sin(x) + 1)/(a - b) - 1/2*log(-sin(x) + 1)/(a + b)`

Mupad [B] (verification not implemented)

Time = 15.14 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.21

$$\begin{aligned} \int \frac{\tan(x)}{a + b \csc(x)} dx &= \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{\ln\left(\tan\left(\frac{x}{2}\right) + 1\right)}{a - b} - \frac{\ln\left(\tan\left(\frac{x}{2}\right) - 1\right)}{a + b} \\ &\quad + \frac{b^2 \ln\left(b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b\right)}{a (a^2 - b^2)} \end{aligned}$$

input `int(tan(x)/(a + b/sin(x)),x)`

output `log(tan(x/2)^2 + 1)/a - log(tan(x/2) + 1)/(a - b) - log(tan(x/2) - 1)/(a + b) + (b^2*log(b + 2*a*tan(x/2) + b*tan(x/2)^2))/(a*(a^2 - b^2))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.67

$$\begin{aligned} \int \frac{\tan(x)}{a + b \csc(x)} dx \\ = \frac{\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) a^2 - \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) b^2 - \log\left(\tan\left(\frac{x}{2}\right) - 1\right) a^2 + \log\left(\tan\left(\frac{x}{2}\right) - 1\right) ab - \log\left(\tan\left(\frac{x}{2}\right) + 1\right) b^2}{a (a^2 - b^2)} \end{aligned}$$

input `int(tan(x)/(a+b*csc(x)),x)`

output
$$\frac{(\log(\tan(x/2)^2 + 1)*a^2 - \log(\tan(x/2)^2 + 1)*b^2 - \log(\tan(x/2) - 1)*a^2 + \log(\tan(x/2) - 1)*a*b - \log(\tan(x/2) + 1)*a^2 - \log(\tan(x/2) + 1)*a*b + \log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*b^2)/(a*(a^2 - b^2))$$

3.15 $\int \frac{\cot(x)}{a+b \csc(x)} dx$

Optimal result	126
Mathematica [A] (verified)	126
Rubi [A] (verified)	127
Maple [A] (verified)	128
Fricas [A] (verification not implemented)	129
Sympy [F]	129
Maxima [A] (verification not implemented)	129
Giac [A] (verification not implemented)	130
Mupad [B] (verification not implemented)	130
Reduce [B] (verification not implemented)	130

Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{\log(a + b \csc(x))}{a} + \frac{\log(\sin(x))}{a}$$

output `ln(a+b*csc(x))/a+ln(sin(x))/a`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{\log(b + a \sin(x))}{a}$$

input `Integrate[Cot[x]/(a + b*Csc[x]),x]`

output `Log[b + a*Sin[x]]/a`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4373, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{4373} \\
 & - \int \frac{\sin(x)}{b(a + b \csc(x))} d(b \csc(x)) \\
 & \quad \downarrow \textcolor{blue}{47} \\
 & \frac{\int \frac{1}{a+b \csc(x)} d(b \csc(x))}{a} - \frac{\int \frac{\sin(x)}{b} d(b \csc(x))}{a} \\
 & \quad \downarrow \textcolor{blue}{14} \\
 & \frac{\int \frac{1}{a+b \csc(x)} d(b \csc(x))}{a} - \frac{\log(b \csc(x))}{a} \\
 & \quad \downarrow \textcolor{blue}{16} \\
 & \frac{\log(a + b \csc(x))}{a} - \frac{\log(b \csc(x))}{a}
 \end{aligned}$$

input `Int[Cot[x]/(a + b*Csc[x]),x]`

output `-(Log[b*Csc[x]]/a) + Log[a + b*Csc[x]]/a`

Definitions of rubi rules used

rule 14 $\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] /; \text{FreeQ}[a, x]$

rule 16 $\text{Int}[(c_)/((a_) + (b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 47 $\text{Int}[1/(((a_) + (b_)*(x_))*(c_*) + (d_*)*(x_)), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{Int}[1/(a + b*x), x] - \text{Simp}[d/(b*c - a*d) \text{Int}[1/(c + d*x), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4373 $\text{Int}[\cot[(c_*) + (d_*)*(x_)]^{(m_)}*(\csc[(c_*) + (d_*)*(x_)]*(b_*) + (a_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[-(-1)^{((m - 1)/2)} / (d*b^{(m - 1)}) \text{Subst}[\text{Int}[(b^2 - x^2)^{((m - 1)/2)}*((a + x)^{n/x}), x], x, b*Csc[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

method	result	size
derivativedivides	$\frac{\ln(a+b\csc(x))}{a} - \frac{\ln(\csc(x))}{a}$	21
default	$\frac{\ln(a+b\csc(x))}{a} - \frac{\ln(\csc(x))}{a}$	21
risch	$-\frac{ix}{a} + \frac{\ln(e^{2ix}-1+\frac{2ib e^{ix}}{a})}{a}$	33

input $\text{int}(1/(a+b*\csc(x))*\cot(x), x, \text{method}=\text{RETURNVERBOSE})$

output $\ln(a+b\csc(x))/a - 1/a*\ln(\csc(x))$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{\log(a \sin(x) + b)}{a}$$

input `integrate(cot(x)/(a+b*csc(x)),x, algorithm="fricas")`

output `log(a*sin(x) + b)/a`

Sympy [F]

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \int \frac{\cot(x)}{a + b \csc(x)} dx$$

input `integrate(cot(x)/(a+b*csc(x)),x)`

output `Integral(cot(x)/(a + b*csc(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.58

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{\log(a \sin(x) + b)}{a}$$

input `integrate(cot(x)/(a+b*csc(x)),x, algorithm="maxima")`

output `log(a*sin(x) + b)/a`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.63

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{\log(|a \sin(x) + b|)}{a}$$

input `integrate(cot(x)/(a+b*csc(x)),x, algorithm="giac")`

output `log(abs(a*sin(x) + b))/a`

Mupad [B] (verification not implemented)

Time = 15.59 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.89

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{2 \operatorname{atanh}\left(\frac{a \left(2 b^3 \sin(x) + \frac{5 a b^2}{2} - a^3 - \frac{a b^2 \cos(2x)}{2}\right)}{(-a^2 + \sin(x) a b + 2 b^2)^2}\right)}{a}$$

input `int(cot(x)/(a + b/sin(x)),x)`

output `(2*atanh((a*(2*b^3*sin(x) + (5*a*b^2)/2 - a^3 - (a*b^2*cos(2*x))/2))/(2*b^2 - a^2 + a*b*sin(x))^2))/a`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.79

$$\int \frac{\cot(x)}{a + b \csc(x)} dx = \frac{-\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right)}{a}$$

input `int(cot(x)/(a+b*csc(x)),x)`

output `(- log(tan(x/2)**2 + 1) + log(tan(x/2)**2*b + 2*tan(x/2)*a + b))/a`

3.16 $\int \frac{\cot^3(x)}{a+b \csc(x)} dx$

Optimal result	131
Mathematica [A] (verified)	131
Rubi [A] (verified)	132
Maple [A] (verified)	133
Fricas [A] (verification not implemented)	134
Sympy [F]	134
Maxima [A] (verification not implemented)	134
Giac [A] (verification not implemented)	135
Mupad [B] (verification not implemented)	135
Reduce [B] (verification not implemented)	135

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx = -\frac{\csc(x)}{b} - \frac{\left(1 - \frac{a^2}{b^2}\right) \log(a + b \csc(x))}{a} - \frac{\log(\sin(x))}{a}$$

output
$$-\csc(x)/b - (1-a^2/b^2)*\ln(a+b*\csc(x))/a - \ln(\sin(x))/a$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx = \frac{-ab \csc(x) - a^2 \log(\sin(x)) + (a^2 - b^2) \log(b + a \sin(x))}{ab^2}$$

input `Integrate[Cot[x]^3/(a + b*Csc[x]), x]`

output
$$(-(a*b*Csc[x]) - a^2*Log[Sin[x]] + (a^2 - b^2)*Log[b + a*Sin[x]])/(a*b^2)$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 4373, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^3}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{4373} \\
 & \frac{\int \frac{(b^2 - b^2 \csc^2(x)) \sin(x)}{b(a+b \csc(x))} d(b \csc(x))}{b^2} \\
 & \quad \downarrow \textcolor{blue}{522} \\
 & \frac{\int \left(\frac{a^2 - b^2}{a(a+b \csc(x))} + \frac{b \sin(x)}{a} - 1 \right) d(b \csc(x))}{b^2} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{(a^2 - b^2) \log(a+b \csc(x))}{a} + \frac{b^2 \log(b \csc(x))}{a} - b \csc(x)
 \end{aligned}$$

input `Int[Cot[x]^3/(a + b*Csc[x]),x]`

output
$$-\frac{b \csc (x)}{b^2}+\frac{\log (b \csc (x))}{a}+\frac{\left(a^2-b^2\right) \log (a+b \csc (x))}{a^2}$$

Definitions of rubi rules used

rule 522 $\text{Int}[(e_*)(x_*)^{(m_*)}((c_*) + (d_*)(x_*))^{(n_*)}((a_*) + (b_*)(x_*)^2)^{(p_*)}], \text{x_Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p], \text{x}], \text{x} /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, \text{x}] \&& \text{IGtQ}[p, 0]$

rule 2009 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \text{x}], \text{x}] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$

rule 4373 $\text{Int}[\cot[(c_*) + (d_*)(x_*)]^{(m_*)}(\csc[(c_*) + (d_*)(x_*)]*(b_*) + (a_*)^{(n_*)}], \text{x_Symbol}] \rightarrow \text{Simp}[-(-1)^{((m - 1)/2)} / (d*b^{(m - 1)}) \text{Subst}[\text{Int}[(b^2 - x^2)^{((m - 1)/2)}*((a + x)^n/x), \text{x}], \text{x}, b*\text{Csc}[c + d*x]], \text{x}] /; \text{FreeQ}[\{a, b, c, d, n\}, \text{x}] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

method	result	size
default	$-\frac{1}{b \sin(x)} - \frac{a \ln(\sin(x))}{b^2} + \frac{(a^2 - b^2) \ln(a \sin(x) + b)}{b^2 a}$	43
risch	$\frac{i x}{a} - \frac{2 i e^{i x}}{b(e^{2 i x} - 1)} - \frac{a \ln(e^{2 i x} - 1)}{b^2} + \frac{a \ln\left(e^{2 i x} - 1 + \frac{2 i b e^{i x}}{a}\right)}{b^2} - \frac{\ln\left(e^{2 i x} - 1 + \frac{2 i b e^{i x}}{a}\right)}{a}$	93

input `int(cot(x)^3/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output $-1/b/\sin(x) - a/b^2*\ln(\sin(x)) + (a^2 - b^2)/b^2/a*\ln(a*\sin(x) + b)$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.24

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx \\ = -\frac{a^2 \log\left(-\frac{1}{2} \sin(x)\right) \sin(x) - (a^2 - b^2) \log(a \sin(x) + b) \sin(x) + ab}{ab^2 \sin(x)}$$

input `integrate(cot(x)^3/(a+b*csc(x)),x, algorithm="fricas")`

output `-(a^2*log(-1/2*sin(x))*sin(x) - (a^2 - b^2)*log(a*sin(x) + b)*sin(x) + a*b)/(a*b^2*sin(x))`

Sympy [F]

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx = \int \frac{\cot^3(x)}{a + b \csc(x)} dx$$

input `integrate(cot(x)**3/(a+b*csc(x)),x)`

output `Integral(cot(x)**3/(a + b*csc(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx = -\frac{a \log(\sin(x))}{b^2} + \frac{(a^2 - b^2) \log(a \sin(x) + b)}{ab^2} - \frac{1}{b \sin(x)}$$

input `integrate(cot(x)^3/(a+b*csc(x)),x, algorithm="maxima")`

output `-a*log(sin(x))/b^2 + (a^2 - b^2)*log(a*sin(x) + b)/(a*b^2) - 1/(b*sin(x))`

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx = -\frac{a \log(|\sin(x)|)}{b^2} + \frac{(a^2 - b^2) \log(|a \sin(x) + b|)}{ab^2} - \frac{1}{b \sin(x)}$$

input `integrate(cot(x)^3/(a+b*csc(x)),x, algorithm="giac")`

output `-a*log(abs(sin(x)))/b^2 + (a^2 - b^2)*log(abs(a*sin(x) + b))/(a*b^2) - 1/(b*sin(x))`

Mupad [B] (verification not implemented)

Time = 15.50 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.97

$$\begin{aligned} \int \frac{\cot^3(x)}{a + b \csc(x)} dx &= \ln \left(b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b \right) \left(\frac{a}{b^2} - \frac{1}{a} \right) - \frac{\tan\left(\frac{x}{2}\right)}{2b} \\ &\quad + \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{1}{2b \tan\left(\frac{x}{2}\right)} - \frac{a \ln\left(\tan\left(\frac{x}{2}\right)\right)}{b^2} \end{aligned}$$

input `int(cot(x)^3/(a + b/sin(x)),x)`

output `log(b + 2*a*tan(x/2) + b*tan(x/2)^2)*(a/b^2 - 1/a) - tan(x/2)/(2*b) + log(tan(x/2)^2 + 1)/a - 1/(2*b*tan(x/2)) - (a*log(tan(x/2)))/b^2`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.42

$$\begin{aligned} \int \frac{\cot^3(x)}{a + b \csc(x)} dx \\ = \frac{\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x) b^2 + \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right) \sin(x) a^2 - \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right) \sin(x) a b^2}{\sin(x) a b^2} \end{aligned}$$

input `int(cot(x)^3/(a+b*csc(x)),x)`

output
$$\frac{(\log(\tan(x/2)^2 + 1)*\sin(x)*b^2 + \log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*\sin(x)*a^2 - \log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*\sin(x)*b^2 - \log(\tan(x/2))*\sin(x)*a^2 - a*b)/(\sin(x)*a*b^2)}$$

3.17 $\int \frac{\cot^5(x)}{a+b \csc(x)} dx$

Optimal result	137
Mathematica [A] (verified)	137
Rubi [A] (verified)	138
Maple [A] (verified)	139
Fricas [B] (verification not implemented)	140
Sympy [F]	140
Maxima [A] (verification not implemented)	141
Giac [A] (verification not implemented)	141
Mupad [B] (verification not implemented)	142
Reduce [B] (verification not implemented)	142

Optimal result

Integrand size = 13, antiderivative size = 72

$$\begin{aligned} \int \frac{\cot^5(x)}{a + b \csc(x)} dx = & -\frac{(a^2 - 2b^2) \csc(x)}{b^3} + \frac{a \csc^2(x)}{2b^2} - \frac{\csc^3(x)}{3b} \\ & + \frac{(a^2 - b^2)^2 \log(a + b \csc(x))}{ab^4} + \frac{\log(\sin(x))}{a} \end{aligned}$$

output $-\frac{(a^2 - 2b^2) \csc(x)}{b^3} + \frac{a \csc^2(x)}{2b^2} - \frac{\csc^3(x)}{3b} + \frac{(a^2 - b^2)^2 \log(a + b \csc(x))}{ab^4} + \frac{\log(\sin(x))}{a}$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.18

$$\begin{aligned} \int \frac{\cot^5(x)}{a + b \csc(x)} dx \\ = \frac{-6ab(a^2 - 2b^2) \csc(x) + 3a^2b^2 \csc^2(x) - 2ab^3 \csc^3(x) - 6a^2(a^2 - 2b^2) \log(\sin(x)) + 6(a^2 - b^2)^2 \log(b + a \csc(x))}{6ab^4} \end{aligned}$$

input `Integrate[Cot[x]^5/(a + b*Csc[x]),x]`

output

$$\frac{(-6*a*b*(a^2 - 2*b^2)*Csc[x] + 3*a^2*b^2*Csc[x]^2 - 2*a*b^3*Csc[x]^3 - 6*a^2*(a^2 - 2*b^2)*Log[Sin[x]] + 6*(a^2 - b^2)^2*Log[b + a*Sin[x]])/(6*a*b^4)}$$

Rubi [A] (verified)

Time = 0.28 (sec), antiderivative size = 77, normalized size of antiderivative = 1.07, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.308, Rules used = {3042, 4373, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^5(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^5}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{4373} \\
 & - \frac{\int \frac{(b^2 - b^2 \csc^2(x))^2 \sin(x)}{b(a+b \csc(x))} d(b \csc(x))}{b^4} \\
 & \quad \downarrow \textcolor{blue}{522} \\
 & - \frac{\int \left(\frac{\sin(x)b^3}{a} + \csc^2(x)b^2 - a \csc(x)b + a^2 \left(1 - \frac{2b^2}{a^2} \right) - \frac{(a^2 - b^2)^2}{a(a+b \csc(x))} \right) d(b \csc(x))}{b^4} \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & - \frac{b(a^2 - 2b^2) \csc(x) - \frac{(a^2 - b^2)^2 \log(a+b \csc(x))}{a} + \frac{b^4 \log(b \csc(x))}{a} - \frac{1}{2}ab^2 \csc^2(x) + \frac{1}{3}b^3 \csc^3(x)}{b^4}
 \end{aligned}$$

input

$$\text{Int}[\cot[x]^5/(a + b*\csc[x]), x]$$

output

$$-\frac{((b*(a^2 - 2*b^2)*Csc[x] - (a*b^2*Csc[x]^2)/2 + (b^3*Csc[x]^3)/3 + (b^4*Log[b*Csc[x]])/a - ((a^2 - b^2)^2*Log[a + b*Csc[x]])/a)/b^4}$$

Definitions of rubi rules used

rule 522 $\text{Int}[(e_*)(x_*)^{(m_*)}((c_*) + (d_*)(x_*))^{(n_*)}((a_*) + (b_*)(x_*)^2)^{(p_*)}], \text{x_Symbol}] \rightarrow \text{Int}[\text{ExpandIntegrand}[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p], \text{x}], \text{x} /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, \text{x}] \&& \text{IGtQ}[p, 0]$

rule 2009 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Simp}[\text{IntSum}[u, \text{x}], \text{x}] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[u, \text{x}]$

rule 4373 $\text{Int}[\cot[(c_*) + (d_*)(x_*)]^{(m_*)}(\csc[(c_*) + (d_*)(x_*)]*(b_*) + (a_*)^{(n_*)}], \text{x_Symbol}] \rightarrow \text{Simp}[-(-1)^{((m - 1)/2)} / (d*b^{(m - 1)}) \text{Subst}[\text{Int}[(b^2 - x^2)^{((m - 1)/2)}*((a + x)^n/x), \text{x}], \text{x}, b*\text{Csc}[c + d*x]], \text{x}] /; \text{FreeQ}[\{a, b, c, d, n\}, \text{x}] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{NeQ}[a^2 - b^2, 0]$

Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.19

method	result
default	$-\frac{1}{3b \sin(x)^3} - \frac{a^2 - 2b^2}{b^3 \sin(x)} + \frac{a}{2b^2 \sin(x)^2} - \frac{(a^2 - 2b^2)a \ln(\sin(x))}{b^4} + \frac{(a^4 - 2a^2b^2 + b^4) \ln(a \sin(x) + b)}{b^4 a}$
risch	$-\frac{ix}{a} - \frac{2i(-3iab e^{4ix} + 3a^2 e^{5ix} - 6b^2 e^{5ix} + 3iab e^{2ix} - 6a^2 e^{3ix} + 8b^2 e^{3ix} + 3e^{ix} a^2 - 6e^{ix} b^2)}{3b^3 (e^{2ix} - 1)^3} - \frac{a^3 \ln(e^{2ix} - 1)}{b^4} + \frac{2a \ln(e^{2ix} - 1)}{b^2} +$

input $\text{int}(\cot(x)^5 / (a + b * \csc(x)), \text{x}, \text{method} = \text{RETURNVERBOSE})$

output $-1/3/b/\sin(x)^3 - (a^2 - 2*b^2)/b^3/\sin(x) + 1/2*a/b^2/\sin(x)^2 - (a^2 - 2*b^2)/b^4*a*\ln(\sin(x)) + (a^4 - 2*a^2*b^2 + b^4)/b^4/a*\ln(a*\sin(x) + b)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(68) = 136$.

Time = 0.10 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.10

$$\int \frac{\cot^5(x)}{a + b \csc(x)} dx = -\frac{3 a^2 b^2 \sin(x) - 6 a^3 b + 10 a b^3 + 6 (a^3 b - 2 a b^3) \cos(x)^2 + 6 (a^4 - 2 a^2 b^2 + b^4 - (a^4 - 2 a^2 b^2 + b^4) \cos(x)^2)}{6 (ab^4 \cos(x)^2 - a^2 b^2)}$$

input `integrate(cot(x)^5/(a+b*csc(x)),x, algorithm="fricas")`

output
$$\begin{aligned} & -1/6*(3*a^2*b^2*sin(x) - 6*a^3*b + 10*a*b^3 + 6*(a^3*b - 2*a*b^3)*cos(x)^2 \\ & + 6*(a^4 - 2*a^2*b^2 + b^4 - (a^4 - 2*a^2*b^2 + b^4)*cos(x)^2)*log(a*sin(x) + b)*sin(x) - 6*(a^4 - 2*a^2*b^2 - (a^4 - 2*a^2*b^2)*cos(x)^2)*log(1/2*sin(x))*sin(x))/((a*b^4*cos(x)^2 - a*b^4)*sin(x)) \end{aligned}$$

Sympy [F]

$$\int \frac{\cot^5(x)}{a + b \csc(x)} dx = \int \frac{\cot^5(x)}{a + b \csc(x)} dx$$

input `integrate(cot(x)**5/(a+b*csc(x)),x)`

output `Integral(cot(x)**5/(a + b*csc(x)), x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.17

$$\int \frac{\cot^5(x)}{a + b \csc(x)} dx = -\frac{(a^3 - 2ab^2) \log(\sin(x))}{b^4} + \frac{(a^4 - 2a^2b^2 + b^4) \log(a \sin(x) + b)}{ab^4}$$

$$+ \frac{3ab \sin(x) - 6(a^2 - 2b^2) \sin(x)^2 - 2b^2}{6b^3 \sin(x)^3}$$

input `integrate(cot(x)^5/(a+b*csc(x)),x, algorithm="maxima")`

output `-(a^3 - 2*a*b^2)*log(sin(x))/b^4 + (a^4 - 2*a^2*b^2 + b^4)*log(a*sin(x) + b)/(a*b^4) + 1/6*(3*a*b*sin(x) - 6*(a^2 - 2*b^2)*sin(x)^2 - 2*b^2)/(b^3*sin(x)^3)`

Giac [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.25

$$\int \frac{\cot^5(x)}{a + b \csc(x)} dx = -\frac{(a^3 - 2ab^2) \log(|\sin(x)|)}{b^4} + \frac{(a^4 - 2a^2b^2 + b^4) \log(|a \sin(x) + b|)}{ab^4}$$

$$+ \frac{3ab^2 \sin(x) - 2b^3 - 6(a^2b - 2b^3) \sin(x)^2}{6b^4 \sin(x)^3}$$

input `integrate(cot(x)^5/(a+b*csc(x)),x, algorithm="giac")`

output `-(a^3 - 2*a*b^2)*log(abs(sin(x)))/b^4 + (a^4 - 2*a^2*b^2 + b^4)*log(abs(a*sin(x) + b))/(a*b^4) + 1/6*(3*a*b^2*sin(x) - 2*b^3 - 6*(a^2*b - 2*b^3)*sin(x)^2)/(b^4*sin(x)^3)`

Mupad [B] (verification not implemented)

Time = 15.93 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.18

$$\begin{aligned} \int \frac{\cot^5(x)}{a + b \csc(x)} dx &= \tan\left(\frac{x}{2}\right) \left(\frac{7}{8b} - \frac{a^2}{2b^3} \right) - \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{\tan\left(\frac{x}{2}\right)^3}{24b} \\ &\quad + \frac{a \tan\left(\frac{x}{2}\right)^2}{8b^2} - \frac{\tan\left(\frac{x}{2}\right)^2 (4a^2 - 7b^2) + \frac{b^2}{3} - ab \tan\left(\frac{x}{2}\right)}{8b^3 \tan\left(\frac{x}{2}\right)^3} \\ &\quad + \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right) (2ab^2 - a^3)}{b^4} \\ &\quad + \frac{\ln\left(b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b\right) (a^2 - b^2)^2}{ab^4} \end{aligned}$$

input `int(cot(x)^5/(a + b/sin(x)),x)`

output $\tan(x/2)*(7/(8*b) - a^2/(2*b^3)) - \log(\tan(x/2)^2 + 1)/a - \tan(x/2)^3/(24*b) + (a*\tan(x/2)^2)/(8*b^2) - (\tan(x/2)^2*(4*a^2 - 7*b^2) + b^2/3 - a*b*\tan(x/2))/(8*b^3*tan(x/2)^3) + (\log(\tan(x/2))*(2*a*b^2 - a^3))/b^4 + (\log(b + 2*a*tan(x/2) + b*tan(x/2)^2)*(a^2 - b^2)^2)/(a*b^4)$

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 194, normalized size of antiderivative = 2.69

$$\begin{aligned} \int \frac{\cot^5(x)}{a + b \csc(x)} dx \\ = \frac{-12 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^3 b^4 + 12 \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right) \sin(x)^3 a^4 - 24 \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right) \sin(x)^3 a^3 b^3}{a^5 b^5} \end{aligned}$$

input `int(cot(x)^5/(a+b*csc(x)),x)`

```
output ( - 12*log(tan(x/2)**2 + 1)*sin(x)**3*b**4 + 12*log(tan(x/2)**2*b + 2*tan(x/2)*a + b)*sin(x)**3*a**4 - 24*log(tan(x/2)**2*b + 2*tan(x/2)*a + b)*sin(x)**3*a**2*b**2 + 12*log(tan(x/2)**2*b + 2*tan(x/2)*a + b)*sin(x)**3*b**4 - 12*log(tan(x/2))*sin(x)**3*a**4 + 24*log(tan(x/2))*sin(x)**3*a**2*b**2 - 3*sin(x)**3*a**2*b**2 - 12*sin(x)**2*a**3*b + 24*sin(x)**2*a*b**3 + 6*sin(x)*a**2*b**2 - 4*a*b**3)/(12*sin(x)**3*a*b**4)
```

3.18 $\int \frac{\cot^7(x)}{a+b \csc(x)} dx$

Optimal result	144
Mathematica [A] (verified)	144
Rubi [A] (verified)	145
Maple [A] (verified)	146
Fricas [B] (verification not implemented)	147
Sympy [F]	147
Maxima [A] (verification not implemented)	148
Giac [A] (verification not implemented)	148
Mupad [B] (verification not implemented)	149
Reduce [B] (verification not implemented)	149

Optimal result

Integrand size = 13, antiderivative size = 122

$$\begin{aligned} \int \frac{\cot^7(x)}{a + b \csc(x)} dx = & -\frac{(a^4 - 3a^2b^2 + 3b^4) \csc(x)}{b^5} + \frac{a(a^2 - 3b^2) \csc^2(x)}{2b^4} \\ & - \frac{(a^2 - 3b^2) \csc^3(x)}{3b^3} + \frac{a \csc^4(x)}{4b^2} - \frac{\csc^5(x)}{5b} \\ & + \frac{(a^2 - b^2)^3 \log(a + b \csc(x))}{ab^6} - \frac{\log(\sin(x))}{a} \end{aligned}$$

output

```
-(a^4-3*a^2*b^2+3*b^4)*csc(x)/b^5+1/2*a*(a^2-3*b^2)*csc(x)^2/b^4-1/3*(a^2-3*b^2)*csc(x)^3/b^3+1/4*a*csc(x)^4/b^2-1/5*csc(x)^5/b+(a^2-b^2)^3*ln(a+b*csc(x))/a/b^6-1n(sin(x))/a
```

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.08

$$\begin{aligned} \int \frac{\cot^7(x)}{a + b \csc(x)} dx \\ = \frac{-60b(a^4 - 3a^2b^2 + 3b^4) \csc(x) + 30ab^2(a^2 - 3b^2) \csc^2(x) - 20b^3(a^2 - 3b^2) \csc^3(x) + 15ab^4 \csc^4(x) - 12b^6 \csc^5(x)}{60b^6} \end{aligned}$$

input $\text{Integrate}[\cot[x]^7/(a + b \csc[x]), x]$

output
$$\begin{aligned} & (-60*b*(a^4 - 3*a^2*b^2 + 3*b^4)*\csc[x] + 30*a*b^2*(a^2 - 3*b^2)*\csc[x]^2 \\ & - 20*b^3*(a^2 - 3*b^2)*\csc[x]^3 + 15*a*b^4*\csc[x]^4 - 12*b^5*\csc[x]^5 - 60 \\ & *a*(a^4 - 3*a^2*b^2 + 3*b^4)*\log[\sin[x]] + (60*(a^2 - b^2)^3*3*\log[b + a*\sin \\ & [x]])/a)/(60*b^6) \end{aligned}$$

Rubi [A] (verified)

Time = 0.33 (sec), antiderivative size = 125, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 4373, 522, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^7(x)}{a + b \csc(x)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cot(x)^7}{a + b \csc(x)} dx \\ & \quad \downarrow \text{4373} \\ & \frac{\int \frac{(b^2 - b^2 \csc^2(x))^3 \sin(x)}{b(a+b \csc(x))} d(b \csc(x))}{b^6} \\ & \quad \downarrow \text{522} \\ & \frac{\int \left(\frac{\sin(x)b^5}{a} - \csc^4(x)b^4 + a \csc^3(x)b^3 - (a^2 - 3b^2) \csc^2(x)b^2 + a(a^2 - 3b^2) \csc(x)b - a^4 \left(\frac{3(b^2 - a^2)b^2}{a^4} + 1 \right) + \frac{(a^2 - 3b^2)^3}{a(a+b \csc(x))} \right) dx}{b^6} \\ & \quad \downarrow \text{2009} \\ & \frac{\frac{1}{2}ab^2(a^2 - 3b^2)\csc^2(x) + \frac{(a^2 - b^2)^3 \log(a+b \csc(x))}{a} - \frac{1}{3}b^3(a^2 - 3b^2)\csc^3(x) - b(a^4 - 3a^2b^2 + 3b^4)\csc(x) + \frac{b^6 \log(b \csc(x))}{a}}{b^6} \end{aligned}$$

input $\text{Int}[\cot[x]^7/(a + b*\csc[x]), x]$

output
$$\begin{aligned} & \left(-\frac{(b^4(a^4 - 3a^2b^2 + 3b^4)\csc[x])}{2} + \frac{(a^2b^2(a^2 - 3b^2)\csc[x]^2)}{2} \right. \\ & \left. - \frac{(b^3(a^2 - 3b^2)\csc[x]^3)}{3} + \frac{(a^4b^4\csc[x]^4)}{4} - \frac{(b^5\csc[x]^5)}{5} + \right. \\ & \left. \frac{(b^6\log[b*\csc[x]])}{a} + \frac{((a^2 - b^2)^3\log[a + b*\csc[x]])}{a} \right) / b^6 \end{aligned}$$

Definitions of rubi rules used

rule 522
$$\text{Int}[(e_*(x_*)^{(m_*)}((c_*) + (d_*)*(x_*)^{(n_*)}((a_*) + (b_*)*(x_*)^2)^{(p_*)}), x_{\text{Symbol}}] \Rightarrow \text{Int}[\text{ExpandIntegrand}[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x] \&& \text{IGtQ}[p, 0]$$

rule 2009
$$\text{Int}[u_, x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 3042
$$\text{Int}[u_, x_{\text{Symbol}}] \Rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$$

rule 4373
$$\text{Int}[\cot[(c_*) + (d_*)*(x_*)^{(m_*)}(\csc[(c_*) + (d_*)*(x_*)]*(b_*) + (a_*)^{(n_*)}), x_{\text{Symbol}}] \Rightarrow \text{Simp}[-(-1)^{(m - 1)/2}/(d*b^{(m - 1)}) \text{Subst}[\text{Int}[(b^2 - x^2)^{(m - 1)/2}*((a + x)^n/x), x], x, b*\csc[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{NeQ}[a^2 - b^2, 0]$$

Maple [A] (verified)

Time = 1.41 (sec), antiderivative size = 149, normalized size of antiderivative = 1.22

method	result
default	$-\frac{1}{5b\sin(x)^5} - \frac{a^2 - 3b^2}{3b^3\sin(x)^3} - \frac{a^4 - 3a^2b^2 + 3b^4}{b^5\sin(x)} + \frac{a}{4b^2\sin(x)^4} + \frac{(a^2 - 3b^2)a}{2b^4\sin(x)^2} - \frac{(a^4 - 3a^2b^2 + 3b^4)a\ln(\sin(x))}{b^6} + \frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\ln(\sin(x))}{5b^5\sin(x)^5}$
risch	$\frac{ix}{a} - \frac{2i(-45ia^3b^3e^{2ix} + 45ia^3be^{6ix} + 15a^4e^{9ix} - 45a^2b^2e^{9ix} + 45b^4e^{9ix} - 15ia^3be^{8ix} - 45ia^3be^{4ix} - 60a^4e^{7ix} + 160a^2b^2e^{7ix} - 120b^4e^{7ix})}{a^7}$

input
$$\text{int}(\cot(x)^7/(a+b*\csc(x)), x, \text{method}=\text{_RETURNVERBOSE})$$

output

$$\begin{aligned} & -\frac{1}{5} \frac{b}{\sin(x)^5} - \frac{1}{3} \frac{(a^2 - 3b^2)}{b^3 \sin(x)^3} - \frac{(a^4 - 3a^2 b^2 + 3b^4)}{b^5 \sin(x)} \\ & + \frac{1}{4} \frac{a}{b^2 \sin(x)^4} + \frac{1}{2} \frac{(a^2 - 3b^2)}{b^4 a \sin(x)^2} - \frac{(a^4 - 3a^2 b^2 + 3b^4)}{b^6 a \ln(\sin(x))} \\ & + \frac{(a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6)}{b^6 a \ln(a \sin(x) + b)} \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 327 vs. $2(114) = 228$.

Time = 0.13 (sec), antiderivative size = 327, normalized size of antiderivative = 2.68

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx =$$

$$\frac{60 a^5 b - 160 a^3 b^3 + 132 a b^5 + 60 (a^5 b - 3 a^3 b^3 + 3 a b^5) \cos(x)^4 - 20 (6 a^5 b - 17 a^3 b^3 + 15 a b^5) \cos(x)^2}{a^6 - 3 a^4 b^2 + 3 a^2 b^4 - b^6 + (a^6 - 3 a^4 b^2 + 3 a^2 b^4 - b^6) \cos(x)^4 - 2 (a^6 - 3 a^4 b^2 + 3 a^2 b^4 - b^6) \cos(x)^2} \log(a \sin(x) + b) \sin(x)$$

input

```
integrate(cot(x)^7/(a+b*csc(x)),x, algorithm="fricas")
```

output

$$\begin{aligned} & -\frac{1}{60} (60 a^5 b - 160 a^3 b^3 + 132 a b^5 + 60 (a^5 b - 3 a^3 b^3 + 3 a b^5) \cos(x)^4 - 20 (6 a^5 b - 17 a^3 b^3 + 15 a b^5) \cos(x)^2 - 60 (a^6 - 3 a^4 b^2 + 3 a^2 b^4 - b^6) \cos(x)^4 - 2 (a^6 - 3 a^4 b^2 + 3 a^2 b^4 - b^6) \cos(x)^2) \log(a \sin(x) + b) \sin(x) \\ & + 60 (a^6 - 3 a^4 b^2 + 3 a^2 b^4 + (a^6 - 3 a^4 b^2 + 3 a^2 b^4) \cos(x)^4 - 2 (a^6 - 3 a^4 b^2 + 3 a^2 b^4) \cos(x)^2) \log(-1/2 \sin(x)) \sin(x) - 15 (2 a^4 b^2 - 5 a^2 b^4 - 2 (a^4 b^2 - 3 a^2 b^4) \cos(x)^2) \sin(x) / ((a b^6 \cos(x)^4 - 2 a^2 b^6 \cos(x)^2 + a^2 b^6) \sin(x)) \end{aligned}$$

Sympy [F]

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx = \int \frac{\cot^7(x)}{a + b \csc(x)} dx$$

input

```
integrate(cot(x)**7/(a+b*csc(x)),x)
```

output

```
Integral(cot(x)**7/(a + b*csc(x)), x)
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.22

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx$$

$$= -\frac{(a^5 - 3a^3b^2 + 3ab^4) \log(\sin(x))}{b^6} + \frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \log(a \sin(x) + b)}{ab^6}$$

$$+ \frac{15ab^3 \sin(x) - 60(a^4 - 3a^2b^2 + 3b^4) \sin(x)^4 - 12b^4 + 30(a^3b - 3ab^3) \sin(x)^3 - 20(a^2b^2 - 3b^4) \sin(x)^2}{60b^5 \sin(x)^5}$$

input `integrate(cot(x)^7/(a+b*csc(x)),x, algorithm="maxima")`

output $-(a^5 - 3a^3b^2 + 3ab^4) \log(\sin(x))/b^6 + (a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \log(a \sin(x) + b)/(a*b^6) + 1/60*(15*a*b^3*\sin(x) - 60*(a^4 - 3a^2b^2 + 3b^4)*\sin(x)^4 - 12*b^4 + 30*(a^3b - 3ab^3)*\sin(x)^3 - 20*(a^2b^2 - 3b^4)*\sin(x)^2)/(b^5*\sin(x)^5)$

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.27

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx$$

$$= -\frac{(a^5 - 3a^3b^2 + 3ab^4) \log(|\sin(x)|)}{b^6} + \frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \log(|a \sin(x) + b|)}{ab^6}$$

$$+ \frac{15ab^4 \sin(x) - 12b^5 - 60(a^4b - 3a^2b^3 + 3b^5) \sin(x)^4 + 30(a^3b^2 - 3ab^4) \sin(x)^3 - 20(a^2b^3 - 3b^5) \sin(x)^2}{60b^6 \sin(x)^5}$$

input `integrate(cot(x)^7/(a+b*csc(x)),x, algorithm="giac")`

output $-(a^5 - 3a^3b^2 + 3ab^4) \log(\text{abs}(\sin(x)))/b^6 + (a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \log(\text{abs}(a \sin(x) + b))/(a*b^6) + 1/60*(15*a*b^4*\sin(x) - 12*b^5 - 60*(a^4b - 3a^2b^3 + 3b^5)*\sin(x)^4 + 30*(a^3b^2 - 3ab^4)*\sin(x)^3 - 20*(a^2b^3 - 3b^5)*\sin(x)^2)/(b^6*\sin(x)^5)$

Mupad [B] (verification not implemented)

Time = 15.50 (sec) , antiderivative size = 284, normalized size of antiderivative = 2.33

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx = \tan\left(\frac{x}{2}\right)^3 \left(\frac{3}{32b} - \frac{a^2}{24b^3} \right) - \frac{19 \tan\left(\frac{x}{2}\right)}{16b} \\ + \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{\tan\left(\frac{x}{2}\right)^5}{160b} - \tan\left(\frac{x}{2}\right)^2 \left(\frac{a}{32b^2} + \frac{a\left(\frac{9}{32b} - \frac{a^2}{8b^3}\right)}{b} \right) \\ + \frac{11a^2 \tan\left(\frac{x}{2}\right)}{8b^3} + \frac{a \tan\left(\frac{x}{2}\right)^4}{64b^2} - \frac{a^4 \tan\left(\frac{x}{2}\right)}{2b^5} - \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right) (a^5 - 3a^3b^2 + 3ab^4)}{b^6} \\ - \frac{\cot\left(\frac{x}{2}\right)^5 \left(\tan\left(\frac{x}{2}\right)^3 (10ab^3 - 4a^3b) - \tan\left(\frac{x}{2}\right)^2 \left(3b^4 - \frac{4a^2b^2}{3} \right) + \tan\left(\frac{x}{2}\right)^4 (16a^4 - 44a^2b^2 + 38b^4) + \frac{b^4}{5} \right)}{32b^5} \\ + \frac{\ln\left(b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b\right) (a^2 - b^2)^3}{ab^6}$$

input `int(cot(x)^7/(a + b/sin(x)),x)`

output $\tan(x/2)^3*(3/(32*b) - a^2/(24*b^3)) - (19*tan(x/2))/(16*b) + \log(\tan(x/2)^2 + 1)/a - \tan(x/2)^5/(160*b) - \tan(x/2)^2*(a/(32*b^2) + (a*(9/(32*b) - a^2/(8*b^3))/b) + (11*a^2*tan(x/2))/(8*b^3) + (a*tan(x/2)^4)/(64*b^2) - (a^4*tan(x/2))/(2*b^5) - (\log(\tan(x/2))*(3*a*b^4 + a^5 - 3*a^3*b^2))/b^6 - (\cot(x/2)^5*(\tan(x/2)^3*(10*a*b^3 - 4*a^3*b) - \tan(x/2)^2*(3*b^4 - (4*a^2*b^2)/3) + \tan(x/2)^4*(16*a^4 + 38*b^4 - 44*a^2*b^2) + b^4/5 - (a*b^3*tan(x/2))/2))/(32*b^5) + (\log(b + 2*a*tan(x/2) + b*tan(x/2)^2)*(a^2 - b^2)^3)/(a*b^6)$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 311, normalized size of antiderivative = 2.55

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx \\ = \frac{480 \log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) \sin(x)^5 b^6 + 480 \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right) \sin(x)^5 a^6 - 1440 \log\left(\tan\left(\frac{x}{2}\right)^2 b + 2 \tan\left(\frac{x}{2}\right) a + b\right) \sin(x)^5 a^5 b^5}{a^6 b^6}$$

input `int(cot(x)^7/(a+b*csc(x)),x)`

output
$$\begin{aligned} & (480*\log(\tan(x/2)^2 + 1)*\sin(x)^5*b^6 + 480*\log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*\sin(x)^5*a^6 - 1440*\log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*\sin(x)^5*a^4*b^2 + 1440*\log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*\sin(x)^5*a^2*b^4 - 480*\log(\tan(x/2)^2*b + 2*\tan(x/2)*a + b)*\sin(x)^5*b^6 - 480*\log(\tan(x/2))*\sin(x)^5*a^6 + 1440*\log(\tan(x/2))*\sin(x)^5*a^4*b^2 - 1440*\log(\tan(x/2))*\sin(x)^5*a^2*b^4 - 120*\sin(x)^5*a^4*b^2 + 315*\sin(x)^5*a^2*b^4 - 480*\sin(x)^4*a^5*b + 1440*\sin(x)^4*a^3*b^3 - 1440*\sin(x)^4*a*b^5 + 240*\sin(x)^3*a^4*b^2 - 720*\sin(x)^3*a^2*b^4 - 160*\sin(x)^2*a^3*b^3 + 480*\sin(x)^2*a*b^5 + 120*\sin(x)*a^2*b^4 - 96*a*b^5)/(480*\sin(x)^5*a*b^6) \end{aligned}$$

3.19 $\int \frac{\tan^4(x)}{a+b \csc(x)} dx$

Optimal result	151
Mathematica [A] (verified)	151
Rubi [A] (verified)	152
Maple [A] (verified)	158
Fricas [A] (verification not implemented)	159
Sympy [F]	159
Maxima [F(-2)]	160
Giac [A] (verification not implemented)	160
Mupad [B] (verification not implemented)	161
Reduce [B] (verification not implemented)	161

Optimal result

Integrand size = 13, antiderivative size = 159

$$\begin{aligned} \int \frac{\tan^4(x)}{a+b \csc(x)} dx = & \frac{x}{a} + \frac{2b^5 \operatorname{arctanh}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a(a^2-b^2)^{5/2}} - \frac{b^3 \sec(x)}{(a^2-b^2)^2} + \frac{b \sec(x)}{a^2-b^2} \\ & - \frac{b \sec^3(x)}{3(a^2-b^2)} + \frac{ab^2 \tan(x)}{(a^2-b^2)^2} - \frac{a \tan(x)}{a^2-b^2} + \frac{a \tan^3(x)}{3(a^2-b^2)} \end{aligned}$$

output $x/a+2*b^5*arctanh((a+b*tan(1/2*x))/(a^2-b^2)^(1/2))/a/(a^2-b^2)^(5/2)-b^3*\sec(x)/(a^2-b^2)^2+b*\sec(x)/(a^2-b^2)-b*\sec(x)^3/(3*a^2-3*b^2)+a*b^2*tan(x)/(a^2-b^2)^2-a*tan(x)/(a^2-b^2)+a*tan(x)^3/(3*a^2-3*b^2)$

Mathematica [A] (verified)

Time = 0.90 (sec), antiderivative size = 194, normalized size of antiderivative = 1.22

$$\begin{aligned} & \int \frac{\tan^4(x)}{a+b \csc(x)} dx \\ & \csc(x)(b+a \sin(x)) \left(-\frac{24 b^5 \operatorname{arctan}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{-a^2+b^2}}\right)}{(-a^2+b^2)^{5/2}} + \frac{\sec^3(x) (2 a^3 b - 8 a b^3 + 9 (a^2-b^2)^2 x \cos(x) + 6 a b (a^2-2 b^2) \cos(2 x) + 3 a^4 x \cos(3 x))}{(a-b)^2 (a+b)^2} \right) \\ & = \frac{12 a (a+b \csc(x))}{12 a (a+b \csc(x))} \end{aligned}$$

input $\text{Integrate}[\tan[x]^4/(a + b \csc[x]), x]$

output
$$\frac{(\csc[x] * (b + a \sin[x]) * ((-24*b^5 * \text{ArcTan}[(a + b \tan[x/2]) / \sqrt{-a^2 + b^2}] / (-a^2 + b^2)^{5/2} + (\sec[x]^3 * (2*a^3 * b - 8*a*b^3 + 9*(a^2 - b^2)^2 * x * \cos[x] + 6*a*b*(a^2 - 2*b^2) * \cos[2*x] + 3*a^4 * x * \cos[3*x] - 6*a^2 * b^2 * x * \cos[3*x] + 3*b^4 * x * \cos[3*x] + 3*a^2 * b^2 * \sin[x] - 4*a^4 * \sin[3*x] + 7*a^2 * b^2 * \sin[3*x])) / ((a - b)^2 * (a + b)^2))) / (12*a*(a + b \csc[x]))}{}$$

Rubi [A] (verified)

Time = 1.17 (sec), antiderivative size = 177, normalized size of antiderivative = 1.11, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 1.615$, Rules used = {3042, 4386, 3042, 3381, 3042, 3086, 2009, 3381, 3042, 3086, 24, 3214, 3042, 3139, 1083, 219, 3954, 24, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^4(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1}{\cot(x)^4(a + b \csc(x))} dx \\
 & \quad \downarrow \text{4386} \\
 & \int \frac{\sin(x) \tan^4(x)}{a \sin(x) + b} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sin(x)^5}{\cos(x)^4(a \sin(x) + b)} dx \\
 & \quad \downarrow \text{3381} \\
 & \frac{a \int \tan^4(x) dx}{a^2 - b^2} + \frac{b^2 \int \frac{\sin(x) \tan^2(x)}{b + a \sin(x)} dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan^3(x) dx}{a^2 - b^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \int \frac{\sin(x)^3}{\cos(x)^2(b+a \sin(x))} dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan(x)^3 dx}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{3086} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} - \frac{b \int (\sec^2(x) - 1) d \sec(x)}{a^2 - b^2} + \frac{b^2 \int \frac{\sin(x)^3}{\cos(x)^2(b+a \sin(x))} dx}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{2009} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \int \frac{\sin(x)^3}{\cos(x)^2(b+a \sin(x))} dx}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{3381} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan^2(x) dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan(x) dx}{a^2 - b^2} \right)}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{3042} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan(x) dx}{a^2 - b^2} \right)}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{3086} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \int 1 d \sec(x)}{a^2 - b^2} \right)}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{24} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \right)}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{3214} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \left(\frac{x}{a} - \frac{b \int \frac{1}{b+a \sin(x)} dx}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \right)}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \textcolor{blue}{3042}
\end{aligned}$$

$$\begin{aligned}
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \left(\frac{x}{a} - \frac{b \int \frac{1}{b+a \sin(x)} dx}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \right)}{a^2 - b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \text{3139} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \left(\frac{x}{a} - \frac{2b \int \frac{1}{b \tan^2(\frac{x}{2}) + 2a \tan(\frac{x}{2}) + b} d \tan(\frac{x}{2})}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \right)}{a^2 - b^2} - \\
& \quad \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \text{1083} \\
& \frac{a \int \tan(x)^4 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \left(\frac{4b \int \frac{1}{4(a^2 - b^2) - (2a + 2b \tan(\frac{x}{2}))^2} d(2a + 2b \tan(\frac{x}{2}))}{a} + \frac{x}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \right)}{a^2 - b^2} - \\
& \quad \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \text{219} \\
& \frac{b^2 \left(\frac{b^2 \arctanh \left(\frac{2a + 2b \tan(\frac{x}{2})}{2\sqrt{a^2 - b^2}} \right)}{a\sqrt{a^2 - b^2}} + \frac{x}{a} \right)}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \frac{\frac{a \int \tan(x)^2 dx}{a^2 - b^2} + \frac{a^2 - b^2}{a^2 - b^2} \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2}}{a^2 - b^2} + \frac{a \int \tan(x)^4 dx}{a^2 - b^2} - \\
& \quad \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2 - b^2} \\
& \quad \downarrow \text{3954}
\end{aligned}$$

$$\begin{aligned}
& \frac{b^2 \left(\frac{a(\tan(x) - \int 1 dx)}{a^2 - b^2} + \frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a+2b \tan \left(\frac{x}{2} \right)}{2\sqrt{a^2-b^2}} \right) + \frac{x}{a}}{a\sqrt{a^2-b^2}} \right)}{a^2-b^2} - \frac{b \sec(x)}{a^2-b^2} \right)}{a^2-b^2} + \\
& \frac{a \left(\frac{\tan^3(x)}{3} - \int \tan^2(x) dx \right)}{a^2-b^2} - \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2-b^2} \\
& \quad \downarrow 24 \\
& \frac{b^2 \left(\frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a+2b \tan \left(\frac{x}{2} \right)}{2\sqrt{a^2-b^2}} \right) + \frac{x}{a}}{a\sqrt{a^2-b^2}} \right)}{a^2-b^2} + \frac{a(\tan(x)-x)}{a^2-b^2} - \frac{b \sec(x)}{a^2-b^2} \right)}{a^2-b^2} - \\
& \frac{a \left(\frac{\tan^3(x)}{3} - \int \tan^2(x) dx \right)}{a^2-b^2} + \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2-b^2} \\
& \quad \downarrow 3042 \\
& \frac{b^2 \left(\frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a+2b \tan \left(\frac{x}{2} \right)}{2\sqrt{a^2-b^2}} \right) + \frac{x}{a}}{a\sqrt{a^2-b^2}} \right)}{a^2-b^2} + \frac{a(\tan(x)-x)}{a^2-b^2} - \frac{b \sec(x)}{a^2-b^2} \right)}{a^2-b^2} - \\
& \frac{a \left(\frac{\tan^3(x)}{3} - \int \tan(x)^2 dx \right)}{a^2-b^2} + \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2-b^2} \\
& \quad \downarrow 3954
\end{aligned}$$

$$\begin{aligned}
 & \frac{a \left(\int 1 dx + \frac{\tan^3(x)}{3} - \tan(x) \right)}{a^2 - b^2} + \frac{b^2 \left(\frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a+2b \tan(\frac{x}{2})}{2\sqrt{a^2-b^2}} \right) + \frac{x}{a}}{a\sqrt{a^2-b^2}} \right)}{a^2-b^2} + \frac{a(\tan(x)-x)}{a^2-b^2} - \frac{b \sec(x)}{a^2-b^2} \right)}{a^2-b^2} - \\
 & \quad \downarrow 24 \\
 & \frac{b^2 \left(\frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a+2b \tan(\frac{x}{2})}{2\sqrt{a^2-b^2}} \right) + \frac{x}{a}}{a\sqrt{a^2-b^2}} \right)}{a^2-b^2} + \frac{a(\tan(x)-x)}{a^2-b^2} - \frac{b \sec(x)}{a^2-b^2} \right)}{a^2-b^2} + \frac{a \left(x + \frac{\tan^3(x)}{3} - \tan(x) \right)}{a^2-b^2} - \\
 & \quad \frac{b \left(\frac{\sec^3(x)}{3} - \sec(x) \right)}{a^2-b^2}
 \end{aligned}$$

input `Int[Tan[x]^4/(a + b*Csc[x]), x]`

output `-((b*(-Sec[x] + Sec[x]^3/3))/(a^2 - b^2)) + (a*(x - Tan[x] + Tan[x]^3/3))/ (a^2 - b^2) + (b^2*((b^2*(x/a + (2*b*ArcTanh[(2*a + 2*b*Tan[x/2])/(2*.Sqrt[a^2 - b^2]]])/ (a*.Sqrt[a^2 - b^2]))/(a^2 - b^2) - (b*Sec[x])/(a^2 - b^2) + (a*(-x + Tan[x]))/(a^2 - b^2)))/(a^2 - b^2)`

Definitions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))* ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1083 $\text{Int}[(a_ + b_)*(x_ + c_)*x_2^{-1}, x] \rightarrow \text{Simp}[-2 \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 2009 $\text{Int}[u, x] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u, x] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3086 $\text{Int}[(a_*\sec[e_* + f_*)*(x_])^{(m_*)}((b_*)\tan[e_* + f_*)*(x_])^{(n_*)}, x] \rightarrow \text{Simp}[a/f \text{Subst}[\text{Int}[(a*x)^{(m-1)}(-1 + x^2)^{((n-1)/2)}, x], x, \text{Sec}[e + f*x], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \&& \text{IntegerQ}[(n-1)/2] \&& !(\text{IntegerQ}[m/2] \&& \text{LtQ}[0, m, n+1])]$

rule 3139 $\text{Int}[(a_ + b_)\sin[(c_ + d_)*x_]]^{(-1)}, x] \rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Simp}[2*(e/d) \text{Subst}[\text{Int}[1/(a + 2*b*e*x + a*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[a^2 - b^2, 0]]$

rule 3214 $\text{Int}[(a_* + b_*)\sin[(e_* + f_*)*(x_)]/((c_ + d_*)\sin[(e_* + f_*)*(x_)]), x] \rightarrow \text{Simp}[b*(x/d), x] - \text{Simp}[(b*c - a*d)/d \text{Int}[1/(c + d*\text{Sin}[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{NeQ}[b*c - a*d, 0]]$

rule 3381 $\text{Int}[((\cos[e_* + f_*)*(g_*)*(g_*)^{(p_*)}((d_*)\sin[e_* + f_*)*(x_])^{(n_*)})/((a_ + b_*)\sin[(e_* + f_*)*(x_)]), x] \rightarrow \text{Simp}[a*(d^2/(a^2 - b^2)) \text{Int}[(g*\text{Cos}[e + f*x])^p*(d*\text{Sin}[e + f*x])^{(n-2)}, x], x] + (-\text{Simp}[b*(d/(a^2 - b^2)) \text{Int}[(g*\text{Cos}[e + f*x])^p*(d*\text{Sin}[e + f*x])^{(n-1)}, x], x] - \text{Simp}[a^2*(d^2/(g^2*(a^2 - b^2))) \text{Int}[(g*\text{Cos}[e + f*x])^{(p+2)}*(d*\text{Sin}[e + f*x])^{(n-2)/(a + b*\text{Sin}[e + f*x])), x], x]) /; \text{FreeQ}[\{a, b, d, e, f, g\}, x] \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegersQ}[2*n, 2*p] \&& \text{LtQ}[p, -1] \&& \text{GtQ}[n, 1]]$

rule 3954 $\text{Int}[(b_*)\tan(c_*) + (d_*)(x_*)]^{(n_)}, x_{\text{Symbol}} \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)}/(d*(n - 1))), x] - \text{Simp}[b^2 \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&& \text{GtQ}[n, 1]$

rule 4386 $\text{Int}[\cot(c_*) + (d_*)(x_*)]^{(m_)}\text{csc}(c_*) + (d_*)(x_*)*(b_*) + (a_*)^{(n_)}, x_{\text{Symbol}} \rightarrow \text{Int}[\cos[c + d*x]^m * ((b + a*\sin[c + d*x])^n/\sin[c + d*x]^{(m + n)}), x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n] \&& \text{IntegerQ}[m] \&& (\text{IntegerQ}[m/2] \mid\mid \text{LeQ}[m, 1])$

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.19

method	result
default	$\frac{2 \arctan(\tan(\frac{x}{2}))}{a} - \frac{64}{3(\tan(\frac{x}{2})+1)^3(64a-64b)} + \frac{32}{(64a-64b)(\tan(\frac{x}{2})+1)^2} - \frac{-2a+3b}{2(a-b)^2(\tan(\frac{x}{2})+1)} - \frac{2b^5 \arctan\left(\frac{2b \tan(\frac{x}{2})}{2\sqrt{-a^2+b^2}}\right)}{(a-b)^2(a+b)^2 a \sqrt{-a^2+b^2}}$
risch	$\frac{x}{a} - \frac{2(6ia^3e^{4ix}-9ia\,b^2e^{4ix}-3a^2be^{5ix}+6b^3e^{5ix}+6ia^3e^{2ix}-12ia\,b^2e^{2ix}-2a^2be^{3ix}+8b^3e^{3ix}+4ia^3-7ia\,b^2-3a^2be^{ix}+6e^{ix}b^3)}{3(-a^2+b^2)^2(e^{2ix}+1)^3} +$

input `int(tan(x)^4/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 2/a*\arctan(\tan(1/2*x))-64/3/(\tan(1/2*x)+1)^3/(64*a-64*b)+32/(64*a-64*b)/(\tan(1/2*x)+1)^2-1/2/(a-b)^2*(-2*a+3*b)/(\tan(1/2*x)+1)-2/(a-b)^2/(a+b)^2*b^5 \\ & /a/(-a^2+b^2)^(1/2)*\arctan(1/2*(2*b*\tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2))-64/3 \\ & /(\tan(1/2*x)-1)^3/(64*a+64*b)-32/(64*a+64*b)/(\tan(1/2*x)-1)^2-1/2/(a+b)^2*(-2*a-3*b)/(\tan(1/2*x)-1) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 483, normalized size of antiderivative = 3.04

$$\int \frac{\tan^4(x)}{a + b \csc(x)} dx$$

$$= \left[\frac{3 \sqrt{a^2 - b^2} b^5 \cos(x)^3 \log\left(\frac{(a^2 - b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 + 2(b \cos(x) \sin(x) + a \cos(x))\sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2ab \sin(x) - a^2 - b^2}\right) - 2a^5 b + 4a^3 b^3 - \dots}{\dots} \right]$$

input `integrate(tan(x)^4/(a+b*csc(x)),x, algorithm="fricas")`

output

$$\begin{aligned} & [1/6*(3*sqrt(a^2 - b^2)*b^5*cos(x)^3*log(((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*s\\ & in(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*c\\ & os(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) - 2*a^5*b + 4*a^3*b^3 - 2*a*b^5 + 6*(\\ & a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*x*cos(x)^3 + 6*(a^5*b - 3*a^3*b^3 + 2*a\\ & *b^5)*cos(x)^2 + 2*(a^6 - 2*a^4*b^2 + a^2*b^4 - (4*a^6 - 11*a^4*b^2 + 7*a^\\ & 2*b^4)*cos(x)^2)*sin(x))/((a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6)*cos(x)^3), \\ & 1/3*(3*sqrt(-a^2 + b^2)*b^5*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2\\ & - b^2)*cos(x)))*cos(x)^3 - a^5*b + 2*a^3*b^3 - a*b^5 + 3*(a^6 - 3*a^4*b^2\\ & + 3*a^2*b^4 - b^6)*x*cos(x)^3 + 3*(a^5*b - 3*a^3*b^3 + 2*a*b^5)*cos(x)^2\\ & + (a^6 - 2*a^4*b^2 + a^2*b^4 - (4*a^6 - 11*a^4*b^2 + 7*a^2*b^4)*cos(x)^2)*\\ & sin(x))/((a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6)*cos(x)^3)] \end{aligned}$$

Sympy [F]

$$\int \frac{\tan^4(x)}{a + b \csc(x)} dx = \int \frac{\tan^4(x)}{a + b \csc(x)} dx$$

input `integrate(tan(x)**4/(a+b*csc(x)),x)`

output `Integral(tan(x)**4/(a + b*csc(x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^4(x)}{a + b \csc(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(tan(x)^4/(a+b*csc(x)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more de

Giac [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.36

$$\begin{aligned} \int \frac{\tan^4(x)}{a + b \csc(x)} dx = & -\frac{2 \left(\pi \lfloor \frac{x}{2\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan(\frac{1}{2}x) + a}{\sqrt{-a^2 + b^2}} \right) \right) b^5}{(a^5 - 2a^3b^2 + ab^4)\sqrt{-a^2 + b^2}} + \frac{x}{a} \\ & + \frac{2 \left(3a^3 \tan(\frac{1}{2}x)^5 - 6ab^2 \tan(\frac{1}{2}x)^5 + 3b^3 \tan(\frac{1}{2}x)^4 - 10a^3 \tan(\frac{1}{2}x)^3 + 16ab^2 \tan(\frac{1}{2}x)^3 + 6a^2b \tan(\frac{1}{2}x)^2 - 3(a^4 - 2a^2b^2 + b^4)(\tan(\frac{1}{2}x)^2 - 1) \right)}{3(a^4 - 2a^2b^2 + b^4)} \end{aligned}$$

input `integrate(tan(x)^4/(a+b*csc(x)),x, algorithm="giac")`

output
$$\begin{aligned} & -2*(\pi*\operatorname{floor}(1/2*x/\pi + 1/2)*\operatorname{sgn}(b) + \arctan((b*\tan(1/2*x) + a)/\sqrt{-a^2 + b^2}))/((a^5 - 2a^3b^2 + ab^4)\sqrt{-a^2 + b^2}) + x/a + 2/3*(3*a^3*\tan(1/2*x)^5 - 6*a*b^2*\tan(1/2*x)^5 + 3*b^3*\tan(1/2*x)^4 - 10*a^3*\tan(1/2*x)^3 + 16*a*b^2*\tan(1/2*x)^3 + 6*a^2*b*\tan(1/2*x)^2 - 12*b^3*\tan(1/2*x)^2 + 3*a^3*\tan(1/2*x) - 6*a*b^2*\tan(1/2*x) - 2*a^2*b + 5*b^3)/((a^4 - 2a^2b^2 + b^4)*(\tan(1/2*x)^2 - 1)^3) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 21.91 (sec) , antiderivative size = 4468, normalized size of antiderivative = 28.10

$$\int \frac{\tan^4(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `int(tan(x)^4/(a + b/sin(x)),x)`

output

$$\begin{aligned} & ((4*\tan(x/2)^2*(a^2*b - 2*b^3))/(a^4 + b^4 - 2*a^2*b^2) - (2*(2*a^2*b - 5*b^3))/(3*(a^4 + b^4 - 2*a^2*b^2)) + (2*b^3*tan(x/2)^4)/(a^4 + b^4 - 2*a^2*b^2) \\ & - (4*a*tan(x/2)^3*(5*a^2 - 8*b^2))/(3*(a^4 + b^4 - 2*a^2*b^2)) + (2*a*tan(x/2)*(a^2 - 2*b^2))/(a^4 + b^4 - 2*a^2*b^2) + (2*a*tan(x/2)^5*(a^2 - 2*b^2))/(a^4 + b^4 - 2*a^2*b^2))/(3*tan(x/2)^2 - 3*tan(x/2)^4 + tan(x/2)^6 - 1) + (2*atan((320*a*b^21*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) + (64*a^21*b*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) - (2560*a^3*b^19*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) + (9280*a^5*b^17*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) - (20160*a^7*b^15*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) + (29184*a^9*b^13*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7...)) \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 632, normalized size of antiderivative = 3.97

$$\int \frac{\tan^4(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `int(tan(x)^4/(a+b*csc(x)),x)`

output

```
(6*sqrt( - a**2 + b**2)*atan((tan(x/2)*b + a)/sqrt( - a**2 + b**2))*cos(x)
 *sin(x)**2*b**5 - 6*sqrt( - a**2 + b**2)*atan((tan(x/2)*b + a)/sqrt( - a**
 2 + b**2))*cos(x)*b**5 + cos(x)*sin(x)**2*tan(x)**3*a**6 - 3*cos(x)*sin(x)
 **2*tan(x)**3*a**4*b**2 + 3*cos(x)*sin(x)**2*tan(x)**3*a**2*b**4 - cos(x)*
 sin(x)**2*tan(x)**3*b**6 - 3*cos(x)*sin(x)**2*tan(x)*a**6 + 9*cos(x)*sin(x)
 )**2*tan(x)*a**4*b**2 - 9*cos(x)*sin(x)**2*tan(x)*a**2*b**4 + 3*cos(x)*sin
 (x)**2*tan(x)*b**6 + 3*cos(x)*sin(x)**2*a**6*x + 2*cos(x)*sin(x)**2*a**5*b
 - 9*cos(x)*sin(x)**2*a**4*b**2*x - 5*cos(x)*sin(x)**2*a**3*b**3 + 9*cos(x)
 )*sin(x)**2*a**2*b**4*x + 3*cos(x)*sin(x)**2*a*b**5 - 3*cos(x)*sin(x)**2*b
 **6*x - cos(x)*tan(x)**3*a**6 + 3*cos(x)*tan(x)**3*a**4*b**2 - 3*cos(x)*ta
 n(x)**3*a**2*b**4 + cos(x)*tan(x)**3*b**6 + 3*cos(x)*tan(x)*a**6 - 9*cos(x)
 )*tan(x)*a**4*b**2 + 9*cos(x)*tan(x)*a**2*b**4 - 3*cos(x)*tan(x)*b**6 - 3*
 cos(x)*a**6*x - 2*cos(x)*a**5*b + 9*cos(x)*a**4*b**2*x + 5*cos(x)*a**3*b**
 3 - 9*cos(x)*a**2*b**4*x - 3*cos(x)*a*b**5 + 3*cos(x)*b**6*x - sin(x)**3*a
 **4*b**2 + 5*sin(x)**3*a**2*b**4 - 4*sin(x)**3*b**6 + 3*sin(x)**2*a**5*b -
 9*sin(x)**2*a**3*b**3 + 6*sin(x)**2*a*b**5 - 3*sin(x)*a**2*b**4 + 3*sin(x)
 )*b**6 - 2*a**5*b + 7*a**3*b**3 - 5*a*b**5)/(3*cos(x)*a*(sin(x)**2*a**6 -
 3*sin(x)**2*a**4*b**2 + 3*sin(x)**2*a**2*b**4 - sin(x)**2*b**6 - a**6 + 3*
 a**4*b**2 - 3*a**2*b**4 + b**6))
```

3.20 $\int \frac{\tan^2(x)}{a+b \csc(x)} dx$

Optimal result	163
Mathematica [A] (verified)	163
Rubi [A] (verified)	164
Maple [A] (verified)	168
Fricas [A] (verification not implemented)	168
Sympy [F]	169
Maxima [F(-2)]	169
Giac [A] (verification not implemented)	169
Mupad [B] (verification not implemented)	170
Reduce [B] (verification not implemented)	171

Optimal result

Integrand size = 13, antiderivative size = 84

$$\int \frac{\tan^2(x)}{a + b \csc(x)} dx = -\frac{x}{a} + \frac{2b^3 \operatorname{arctanh}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{a(a^2-b^2)^{3/2}} - \frac{b \sec(x)}{a^2-b^2} + \frac{a \tan(x)}{a^2-b^2}$$

output
$$-\frac{x}{a} + \frac{2b^3 \operatorname{arctanh}\left(\frac{a+b \tan(1/2*x)}{\sqrt{a^2-b^2}}\right)}{a(a^2-b^2)^{3/2}} - \frac{b \sec(x)}{a^2-b^2} + \frac{a \tan(x)}{a^2-b^2}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.04

$$\begin{aligned} & \int \frac{\tan^2(x)}{a + b \csc(x)} dx \\ &= -\frac{-2b^3 \operatorname{arctan}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{-a^2+b^2}}\right) + \sqrt{-a^2+b^2}(-a^2x + b^2x - ab \sec(x) + a^2 \tan(x))}{a(-a^2+b^2)^{3/2}} \end{aligned}$$

input
$$\text{Integrate}[\tan[x]^2/(a + b*\csc[x]), x]$$

output

$$-\left(\left(-2 b^3 \operatorname{ArcTan}\left[\left(a+b \tan \left[\frac{x}{2}\right]\right) / \sqrt{-a^2+b^2}\right]+\sqrt{-a^2+b^2} * \left(-\left(a^2 x\right)+b^2 x-a b \sec [x]+a^2 \tan [x]\right)\right) /\left(a \left(-a^2+b^2\right)^{(3/2)}\right)$$

Rubi [A] (verified)

Time = 0.62 (sec), antiderivative size = 107, normalized size of antiderivative = 1.27, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.077$, Rules used = {3042, 4386, 3042, 3381, 3042, 3086, 24, 3214, 3042, 3139, 1083, 219, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{1}{\cot(x)^2(a + b \csc(x))} dx \\
 & \quad \downarrow \textcolor{blue}{4386} \\
 & \int \frac{\sin(x) \tan^2(x)}{a \sin(x) + b} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\sin(x)^3}{\cos(x)^2(a \sin(x) + b)} dx \\
 & \quad \downarrow \textcolor{blue}{3381} \\
 & \frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan^2(x) dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan(x) dx}{a^2 - b^2} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan(x) dx}{a^2 - b^2} \\
 & \quad \downarrow \textcolor{blue}{3086} \\
 & \frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \int 1 d \sec(x)}{a^2 - b^2}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow \text{24} \\
& \frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{3214} \\
& \frac{b^2 \left(\frac{x}{a} - \frac{b \int \frac{1}{b+a \sin(x)} dx}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{b^2 \left(\frac{x}{a} - \frac{b \int \frac{1}{b+a \sin(x)} dx}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{3139} \\
& \frac{b^2 \left(\frac{x}{a} - \frac{2b \int \frac{1}{b \tan^2(\frac{x}{2}) + 2a \tan(\frac{x}{2}) + b} d \tan(\frac{x}{2})}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{1083} \\
& \frac{b^2 \left(\frac{4b \int \frac{1}{4(a^2 - b^2) - (2a + 2b \tan(\frac{x}{2}))^2} d(2a + 2b \tan(\frac{x}{2}))}{a} + \frac{x}{a} \right)}{a^2 - b^2} + \frac{a \int \tan(x)^2 dx}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{219} \\
& \frac{a \int \tan(x)^2 dx}{a^2 - b^2} + \frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a + 2b \tan(\frac{x}{2})}{2\sqrt{a^2 - b^2}} \right)}{a\sqrt{a^2 - b^2}} + \frac{x}{a} \right)}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{3954} \\
& \frac{a(\tan(x) - \int 1 dx)}{a^2 - b^2} + \frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a + 2b \tan(\frac{x}{2})}{2\sqrt{a^2 - b^2}} \right)}{a\sqrt{a^2 - b^2}} + \frac{x}{a} \right)}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} \\
& \quad \downarrow \text{24}
\end{aligned}$$

$$\frac{b^2 \left(\frac{2b \operatorname{arctanh} \left(\frac{2a+2b \tan \left(\frac{x}{2} \right)}{2\sqrt{a^2-b^2}} \right)}{a\sqrt{a^2-b^2}} + \frac{x}{a} \right)}{a^2-b^2} + \frac{a(\tan(x)-x)}{a^2-b^2} - \frac{b \sec(x)}{a^2-b^2}$$

input `Int[Tan[x]^2/(a + b*Csc[x]), x]`

output $\frac{(b^2*(x/a + (2*b*\operatorname{ArcTanh}[(2*a + 2*b*Tan[x/2])/(2*\operatorname{Sqrt}[a^2 - b^2])])/(a*\operatorname{Sqr}t[a^2 - b^2]))/(a^2 - b^2) - (b*\operatorname{Sec}[x])/(a^2 - b^2) + (a*(-x + \operatorname{Tan}[x]))/(a^2 - b^2)}$

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

rule 3139 $\text{Int}[(a_.) + (b_.) \cdot \sin[(c_.) + (d_.) \cdot (x_.)]]^{(-1)}, x_{\text{Symbol}}] \Rightarrow \text{With}\{\text{e} = \text{FreeFactors}[\text{Tan}[(c + d \cdot x)/2], x]\}, \text{Simp}[2 \cdot (\text{e}/d) \cdot \text{Subst}[\text{Int}[1/(a + 2 \cdot b \cdot e \cdot x + a \cdot e^2 \cdot x^2), x], x, \text{Tan}[(c + d \cdot x)/2]/\text{e}], x]] /; \text{FreeQ}\{\{a, b, c, d\}, x\} \&& \text{NeQ}[a^2 - b^2, 0]$

rule 3214 $\text{Int}[(a_.) + (b_.) \cdot \sin[(e_.) + (f_.) \cdot (x_.)]] / ((c_.) + (d_.) \cdot \sin[(e_.) + (f_.) \cdot (x_.)]), x_{\text{Symbol}}] \Rightarrow \text{Simp}[b \cdot (x/d), x] - \text{Simp}[(b \cdot c - a \cdot d)/d \cdot \text{Int}[1/(c + d \cdot \text{Sin}[e + f \cdot x]), x], x] /; \text{FreeQ}\{\{a, b, c, d, e, f\}, x\} \&& \text{NeQ}[b \cdot c - a \cdot d, 0]$

rule 3381 $\text{Int}[(\cos[(e_.) + (f_.) \cdot (x_.)] \cdot (g_.)^{(p)}) \cdot ((d_.) \cdot \sin[(e_.) + (f_.) \cdot (x_.)])^{(n)}) / ((a_.) + (b_.) \cdot \sin[(e_.) + (f_.) \cdot (x_.)]), x_{\text{Symbol}}] \Rightarrow \text{Simp}[a \cdot (d^2/(a^2 - b^2)) \cdot \text{Int}[(g \cdot \text{Cos}[e + f \cdot x])^{p \cdot (d \cdot \text{Sin}[e + f \cdot x])^{(n - 2)}}, x], x] + (-\text{Simp}[b \cdot (d/(a^2 - b^2)) \cdot \text{Int}[(g \cdot \text{Cos}[e + f \cdot x])^{p \cdot (d \cdot \text{Sin}[e + f \cdot x])^{(n - 1)}}, x], x] - \text{Simp}[a^2 \cdot (d^2/(g^2 \cdot (a^2 - b^2))) \cdot \text{Int}[(g \cdot \text{Cos}[e + f \cdot x])^{(p + 2) \cdot (d \cdot \text{Sin}[e + f \cdot x])^{(n - 2)} / (a + b \cdot \text{Sin}[e + f \cdot x])}, x], x]) /; \text{FreeQ}\{\{a, b, d, e, f, g\}, x\} \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegersQ}[2 \cdot n, 2 \cdot p] \&& \text{LtQ}[p, -1] \&& \text{GtQ}[n, 1]$

rule 3954 $\text{Int}[(b_.) \cdot \tan[(c_.) + (d_.) \cdot (x_.)]]^{(n)}, x_{\text{Symbol}}] \Rightarrow \text{Simp}[b \cdot ((b \cdot \text{Tan}[c + d \cdot x])^{(n - 1)} / (d \cdot (n - 1))), x] - \text{Simp}[b^2 \cdot \text{Int}[(b \cdot \text{Tan}[c + d \cdot x])^{(n - 2)}, x], x] /; \text{FreeQ}\{\{b, c, d\}, x\} \&& \text{GtQ}[n, 1]$

rule 4386 $\text{Int}[\cot[(c_.) + (d_.) \cdot (x_.)]^{(m_.)} \cdot (\csc[(c_.) + (d_.) \cdot (x_.)] \cdot (b_.) + (a_.)^{(n)}), x_{\text{Symbol}}] \Rightarrow \text{Int}[\text{Cos}[c + d \cdot x]^m \cdot ((b + a \cdot \text{Sin}[c + d \cdot x])^n / \text{Sin}[c + d \cdot x]^{(m + n)}), x] /; \text{FreeQ}\{\{a, b, c, d\}, x\} \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n] \&& \text{IntegerQ}[m] \&& (\text{IntegerQ}[m/2] \text{ || } \text{LeQ}[m, 1])$

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.26

method	result	size
default	$-\frac{2 \arctan(\tan(\frac{x}{2}))}{a} - \frac{16}{(16a+16b)(\tan(\frac{x}{2})-1)} - \frac{2b^3 \arctan\left(\frac{2b \tan(\frac{x}{2})+2a}{2\sqrt{-a^2+b^2}}\right)}{(a-b)(a+b)a\sqrt{-a^2+b^2}} - \frac{16}{(16a-16b)(\tan(\frac{x}{2})+1)}$	106
risch	$-\frac{x}{a} + \frac{-2ia+2b e^{ix}}{(e^{2ix}+1)(-a^2+b^2)} - \frac{b^3 \ln\left(\frac{e^{ix}+ib\sqrt{a^2-b^2}-a^2+b^2}{a\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}(a+b)(a-b)a} + \frac{b^3 \ln\left(\frac{e^{ix}+ib\sqrt{a^2-b^2}+a^2-b^2}{a\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}(a+b)(a-b)a}$	195

input `int(tan(x)^2/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -2/a*\arctan(\tan(1/2*x))-16/(16*a+16*b)/(\tan(1/2*x)-1)-2/(a-b)/(a+b)*b^3/a/ \\ & (-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*b*tan(1/2*x)+2*a)/(-a^2+b^2)^{(1/2)})-16/(16* \\ & a-16*b)/(\tan(1/2*x)+1) \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 296, normalized size of antiderivative = 3.52

$$\begin{aligned} & \int \frac{\tan^2(x)}{a + b \csc(x)} dx \\ &= \left[-\frac{\sqrt{a^2 - b^2} b^3 \cos(x) \log \left(-\frac{(a^2 - 2 b^2) \cos(x)^2 + 2 a b \sin(x) + a^2 + b^2 - 2 (b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2 a b \sin(x) - a^2 - b^2} \right) + 2 a^3 b - 2 a b^3 + \right. \\ & \quad \left. 2 (a^5 - 2 a^3 b^2 + a b^4) \cos(x)}{2 (a^5 - 2 a^3 b^2 + a b^4)} \right] \end{aligned}$$

input `integrate(tan(x)^2/(a+b*csc(x)),x, algorithm="fricas")`

output
$$\begin{aligned} & [-1/2*(\sqrt(a^2 - b^2)*b^3*cos(x)*log(-((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 - 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) + 2*a^3*b - 2*a*b^3 + 2*(a^4 - 2*a^2*b^2 + b^4)*x*cos(x) - 2*(a^4 - a^2*b^2)*sin(x))/((a^5 - 2*a^3*b^2 + a*b^4)*cos(x)), (\sqrt(-a^2 + b^2)*b^3*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a))/((a^2 - b^2)*cos(x)))*cos(x) - a^3*b + a*b^3 - (a^4 - 2*a^2*b^2 + b^4)*x*cos(x) + (a^4 - a^2*b^2)*sin(x))/((a^5 - 2*a^3*b^2 + a*b^4)*cos(x))] \end{aligned}$$

Sympy [F]

$$\int \frac{\tan^2(x)}{a + b \csc(x)} dx = \int \frac{\tan^2(x)}{a + b \csc(x)} dx$$

input `integrate(tan(x)**2/(a+b*csc(x)),x)`

output `Integral(tan(x)**2/(a + b*csc(x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^2(x)}{a + b \csc(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(tan(x)^2/(a+b*csc(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details)`

Giac [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.23

$$\begin{aligned} \int \frac{\tan^2(x)}{a + b \csc(x)} dx = & -\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan \left(\frac{b \tan(\frac{1}{2}x) + a}{\sqrt{-a^2 + b^2}} \right)\right) b^3}{(a^3 - ab^2) \sqrt{-a^2 + b^2}} \\ & - \frac{x}{a} - \frac{2 (a \tan(\frac{1}{2}x) - b)}{(a^2 - b^2) (\tan(\frac{1}{2}x)^2 - 1)} \end{aligned}$$

input `integrate(tan(x)^2/(a+b*csc(x)),x, algorithm="giac")`

output

$$\frac{-2*(\pi*\text{floor}(1/2*x/\pi + 1/2)*\text{sgn}(b) + \arctan((b*\tan(1/2*x) + a)/\sqrt{-a^2 + b^2}))*b^3/((a^3 - a*b^2)*\sqrt{-a^2 + b^2}) - x/a - 2*(a*\tan(1/2*x) - b)/((a^2 - b^2)*(\tan(1/2*x)^2 - 1))}{}$$

Mupad [B] (verification not implemented)

Time = 16.93 (sec), antiderivative size = 2341, normalized size of antiderivative = 27.87

$$\int \frac{\tan^2(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input

```
int(tan(x)^2/(a + b/sin(x)),x)
```

output

$$\begin{aligned} & \frac{((2*b)/(a^2 - b^2) - (2*a*tan(x/2))/(a^2 - b^2))/(\tan(x/2)^2 - 1) + (2*a*tan((64*a^11*b*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) - (192*a*b^11*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) + (768*a^3*b^9*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) - (1216*a^5*b^7*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) + (960*a^7*b^5*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) - (384*a^9*b^3*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3))/a + (b^3*atan(((b^3*((a + b)^3*(a - b)^3)^(1/2)*(tan(x/2)*(64*a^12*b + 64*b^13 - 320*a^2*b^11 + 736*a^4*b^9 - 992*a^6*b^7 + 800*a^8*b^5 - 352*a^10*b^3) - 32*a*b^12 + 160*a^3*b^10 - 320*a^5*b^8 + 320*a^7*b^6 - 160*a^9*b^4 + 32*a^11*b^2 + (b^3*((a + b)^3*(a - b)^3)^(1/2)*(tan(x/2)*(64*a^2*b^12 - 256*a^4*b^10 + 384*a^6*b^8 - 256*a^8*b^6 + 64*a^10*b^4) - 32*a^13*b + 64*a^3*b^11 - 288*a^5*b^9 + 512*a^7*b^7 - 448*a^9*b^5 + 192*a^11*b^3 + (b^3*((a + b)^3*(a - b)^3)^(1/2)*(tan(x/2)*(96*a^14*b + 64*a^2*b^13 - 416*a^4*b^11 + 1120*a^6*b^9 - 1600*a^8*b^7 + 1280*a^10*b^5 - 544*a^12*b^3) - 32*a^3*b^12 + 160*a^5*b^10 - 320*a^7*b^8 + 320*a^9*b^6 - 160*a^11*b^4 + 32*a^13*b^2)))/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)))*1i)/(a*b^6 - \dots} \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.93

$$\int \frac{\tan^2(x)}{a + b \csc(x)} dx \\ = \frac{2\sqrt{-a^2 + b^2} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})b+a}{\sqrt{-a^2+b^2}}\right) \cos(x)b^3 + \cos(x)\tan(x)a^4 - 2\cos(x)\tan(x)a^2b^2 + \cos(x)\tan(x)b^4 - \cos(x)a(a^4 - 2a^2b^2 + b^4)}{\cos(x)a(a^4 - 2a^2b^2 + b^4)}$$

input `int(tan(x)^2/(a+b*csc(x)),x)`

output `(2*sqrt(-a**2 + b**2)*atan((tan(x/2)*b + a)/sqrt(-a**2 + b**2))*cos(x)*b**3 + cos(x)*tan(x)*a**4 - 2*cos(x)*tan(x)*a**2*b**2 + cos(x)*tan(x)*b**4 - cos(x)*a**4*x + cos(x)*a**3*b + 2*cos(x)*a**2*b**2*x - cos(x)*a*b**3 - cos(x)*b**4*x + sin(x)*a**2*b**2 - sin(x)*b**4 - a**3*b + a*b**3)/(cos(x)*a*(a**4 - 2*a**2*b**2 + b**4))`

3.21 $\int \frac{\cot^2(x)}{a+b \csc(x)} dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [A] (verified)	176
Fricas [A] (verification not implemented)	177
Sympy [F]	177
Maxima [F(-2)]	178
Giac [A] (verification not implemented)	178
Mupad [B] (verification not implemented)	179
Reduce [B] (verification not implemented)	179

Optimal result

Integrand size = 13, antiderivative size = 61

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx = -\frac{x}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} + \frac{2\sqrt{a^2 - b^2} \operatorname{arctanh}\left(\frac{a+b\tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{ab}$$

output
$$-\frac{x}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} + \frac{2\sqrt{a^2 - b^2} \operatorname{arctanh}\left(\frac{a+b\tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{ab}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.16

$$\begin{aligned} & \int \frac{\cot^2(x)}{a + b \csc(x)} dx \\ &= \frac{-bx + 2\sqrt{-a^2 + b^2} \operatorname{arctan}\left(\frac{a+b\tan(\frac{x}{2})}{\sqrt{-a^2+b^2}}\right) - a \log(\cos(\frac{x}{2})) + a \log(\sin(\frac{x}{2}))}{ab} \end{aligned}$$

input `Integrate[Cot[x]^2/(a + b*Csc[x]), x]`

output
$$\frac{(-b*x) + 2*\text{Sqrt}[-a^2 + b^2]*\text{ArcTan}[(a + b*\text{Tan}[x/2])/\text{Sqrt}[-a^2 + b^2]] - a * \text{Log}[\text{Cos}[x/2]] + a*\text{Log}[\text{Sin}[x/2]])/(a*b)}$$

Rubi [A] (verified)

Time = 0.60 (sec), antiderivative size = 73, normalized size of antiderivative = 1.20, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.077$, Rules used = {3042, 4382, 3042, 4539, 25, 3042, 4257, 4407, 3042, 4318, 3042, 3139, 1083, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^2(x)}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cot(x)^2}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{4382} \\
 & \int \frac{\csc^2(x) - 1}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\csc(x)^2 - 1}{a + b \csc(x)} dx \\
 & \quad \downarrow \textcolor{blue}{4539} \\
 & \frac{\int -\frac{b+a \csc(x)}{a+b \csc(x)} dx}{b} + \frac{\int \csc(x) dx}{b} \\
 & \quad \downarrow \textcolor{blue}{25} \\
 & \frac{\int \csc(x) dx}{b} - \frac{\int \frac{b+a \csc(x)}{a+b \csc(x)} dx}{b} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{\int \csc(x) dx}{b} - \frac{\int \frac{b+a \csc(x)}{a+b \csc(x)} dx}{b}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 4257 \\
 & -\frac{\int \frac{b+a \csc(x)}{a+b \csc(x)} dx}{b} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 4407 \\
 & -\frac{\frac{(a^2-b^2) \int \frac{\csc(x)}{a+b \csc(x)} dx}{a}}{b} + \frac{bx}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 3042 \\
 & -\frac{\frac{(a^2-b^2) \int \frac{\csc(x)}{a+b \csc(x)} dx}{a}}{b} + \frac{bx}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 4318 \\
 & -\frac{\frac{(a^2-b^2) \int \frac{1}{\frac{a \sin(x)}{b}+1} dx}{ab}}{b} + \frac{bx}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 3042 \\
 & -\frac{\frac{(a^2-b^2) \int \frac{1}{\frac{a \sin(x)}{b}+1} dx}{ab}}{b} + \frac{bx}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 3139 \\
 & -\frac{\frac{2(a^2-b^2) \int \frac{1}{\tan^2\left(\frac{x}{2}\right)+\frac{2a \tan\left(\frac{x}{2}\right)}{b}+1} d \tan\left(\frac{x}{2}\right)}{ab}}{b} + \frac{bx}{a} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 1083 \\
 & -\frac{\frac{bx}{a} - \frac{4(a^2-b^2) \int \frac{1}{-\left(\frac{2a}{b}+2 \tan\left(\frac{x}{2}\right)\right)^2-4\left(1-\frac{a^2}{b^2}\right)} d\left(\frac{2a}{b}+2 \tan\left(\frac{x}{2}\right)\right)}{ab}}{b} - \frac{\operatorname{arctanh}(\cos(x))}{b} \\
 & \quad \downarrow 219 \\
 & -\frac{\frac{bx}{a} - \frac{2 \sqrt{a^2-b^2} \operatorname{arctanh}\left(\frac{b\left(\frac{2a}{b}+2 \tan\left(\frac{x}{2}\right)\right)}{2 \sqrt{a^2-b^2}}\right)}{a}}{b} - \frac{\operatorname{arctanh}(\cos(x))}{b}
 \end{aligned}$$

input

output $-\left(\operatorname{ArcTanh}[\cos[x]]/b\right) - \frac{(b \cos[x])/a - (2 \sqrt{a^2 - b^2}) \operatorname{ArcTanh}\left[\frac{b((2 a)/b + 2 \tan[x/2])}{2 \sqrt{a^2 - b^2}}\right]}{2 \sqrt{a^2 - b^2}}/a/b$

Definitions of rubi rules used

rule 25 $\operatorname{Int}[-(F x_), x \text{Symbol}] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$

rule 219 $\operatorname{Int}[(a_+ + b_-)(x_-)^2, x \text{Symbol}] \rightarrow \operatorname{Simp}[(1/(Rt[a, 2]*Rt[-b, 2])) * \operatorname{ArcTanh}[Rt[-b, 2]*x/Rt[a, 2]], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&& \operatorname{NegQ}[a/b] \&& (\operatorname{GtQ}[a, 0] \text{||} \operatorname{LtQ}[b, 0])$

rule 1083 $\operatorname{Int}[(a_+ + b_-)(x_-) + (c_-)(x_-)^2, x \text{Symbol}] \rightarrow \operatorname{Simp}[-2 \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[b^2 - 4 a c - x^2, x], x], x, b + 2 c x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x]$

rule 3042 $\operatorname{Int}[u_, x \text{Symbol}] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3139 $\operatorname{Int}[(a_+ + b_-)\sin[(c_-) + (d_-)(x_-)], x \text{Symbol}] \rightarrow \operatorname{With}[\{e = \operatorname{FreeFactors}[\operatorname{Tan}[(c + d x)/2], x]\}, \operatorname{Simp}[2*(e/d) \operatorname{Subst}[\operatorname{Int}[1/(a + 2 b e x + a *e^2 x^2), x], x, \operatorname{Tan}[(c + d x)/2]/e], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&& \operatorname{NeQ}[a^2 - b^2, 0]$

rule 4257 $\operatorname{Int}[\csc[(c_-) + (d_-)(x_-)], x \text{Symbol}] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\cos[c + d x]/d, x] /; \operatorname{FreeQ}[\{c, d\}, x]$

rule 4318 $\operatorname{Int}[\csc[(e_-) + (f_-)(x_-)]/(\csc[(e_-) + (f_-)(x_-)]*(b_-) + (a_)), x \text{Symbol}] \rightarrow \operatorname{Simp}[1/b \operatorname{Int}[1/(1 + (a/b)*\operatorname{Sin}[e + f x]), x], x] /; \operatorname{FreeQ}[\{a, b, e, f\}, x] \&& \operatorname{NeQ}[a^2 - b^2, 0]$

rule 4382

```
Int[cot[(c_.) + (d_.)*(x_)]^2*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_),  
x_Symbol] :> Int[(-1 + Csc[c + d*x]^2)*(a + b*Csc[c + d*x])^n, x] /; FreeQ[  
{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0]
```

rule 4407

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/csc[(e_.) + (f_.)*(x_)]*(b_.) +  
(a_), x_Symbol] :> Simp[c*(x/a), x] - Simp[(b*c - a*d)/a Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

rule 4539

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))/csc[(e_.) + (f_.)*(x_)]*(b_.)  
+ (a_), x_Symbol] :> Simp[C/b Int[Csc[e + f*x], x], x] + Simp[1/b Int[(A*b - a*C*Csc[e + f*x])/a + b*Csc[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, C}, x]
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.23

method	result	size
default	$\frac{\ln(\tan(\frac{x}{2}))}{b} - \frac{2\arctan(\tan(\frac{x}{2}))}{a} + \frac{(-2a^2+2b^2)\arctan\left(\frac{2b\tan(\frac{x}{2})+2a}{2\sqrt{-a^2+b^2}}\right)}{ba\sqrt{-a^2+b^2}}$	75
risch	$-\frac{x}{a} - \frac{\sqrt{a^2-b^2}\ln\left(e^{ix}+\frac{ib-\sqrt{a^2-b^2}}{a}\right)}{ba} + \frac{\sqrt{a^2-b^2}\ln\left(e^{ix}+\frac{ib+\sqrt{a^2-b^2}}{a}\right)}{ba} - \frac{\ln(e^{ix}+1)}{b} + \frac{\ln(e^{ix}-1)}{b}$	126

input `int(cot(x)^2/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output

```
1/b*ln(tan(1/2*x))-2/a*arctan(tan(1/2*x))+(-2*a^2+2*b^2)/b/a/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 206, normalized size of antiderivative = 3.38

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx$$

$$= \left[-\frac{2bx + a \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - a \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - \sqrt{a^2 - b^2} \log\left(\frac{(a^2 - 2b^2) \cos(x)^2 + 2ab \sin(x) + a^2 + b^2 + 2}{a^2 \cos(x)^2 - 2ab \sin(x)}\right)}{2ab} \right.$$

$$\left. - \frac{2bx + a \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - a \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 2\sqrt{-a^2 + b^2} \arctan\left(-\frac{\sqrt{-a^2 + b^2}(b \sin(x) + a)}{(a^2 - b^2) \cos(x)}\right)}{2ab} \right]$$

input `integrate(cot(x)^2/(a+b*csc(x)),x, algorithm="fricas")`

output `[-1/2*(2*b*x + a*log(1/2*cos(x) + 1/2) - a*log(-1/2*cos(x) + 1/2) - sqrt(a^2 - b^2)*log((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2))/(a*b), -1/2*(2*b*x + a*log(1/2*cos(x) + 1/2) - a*log(-1/2*cos(x) + 1/2) - 2*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2 - b^2)*cos(x))))/(a*b)]`

Sympy [F]

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx = \int \frac{\cot^2(x)}{a + b \csc(x)} dx$$

input `integrate(cot(x)**2/(a+b*csc(x)),x)`

output `Integral(cot(x)**2/(a + b*csc(x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(cot(x)^2/(a+b*csc(x)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details)

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.31

$$\begin{aligned} \int \frac{\cot^2(x)}{a + b \csc(x)} dx = & -\frac{x}{a} + \frac{\log(|\tan(\frac{1}{2}x)|)}{b} \\ & - \frac{2 \left(\pi \lfloor \frac{x}{2\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(b) + \arctan\left(\frac{b \tan(\frac{1}{2}x) + a}{\sqrt{-a^2 + b^2}}\right) \right) (a^2 - b^2)}{\sqrt{-a^2 + b^2} ab} \end{aligned}$$

input `integrate(cot(x)^2/(a+b*csc(x)),x, algorithm="giac")`

output `-x/a + log(abs(tan(1/2*x)))/b - 2*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))*(a^2 - b^2)/(sqrt(-a^2 + b^2)*a*b)`

Mupad [B] (verification not implemented)

Time = 15.18 (sec) , antiderivative size = 697, normalized size of antiderivative = 11.43

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `int(cot(x)^2/(a + b/sin(x)),x)`

output
$$\begin{aligned} & (2*\operatorname{atan}((64*b^3)/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2)) \\ & + (64*a^3*tan(x/2))/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2)) - (64*a^2*b)/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2)) - (64*a*b^2*tan(x/2))/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2))) / a + \log(\tan(x/2)) / b - (2*\operatorname{atanh}((512*a^4*(a^2 - b^2)^(1/2)) / (256*a*b^4 - 64*b^5*tan(x/2) + 512*a^5 - 768*a^3*b^2 + 832*a^2*b^3*tan(x/2) + (1024*a^6*tan(x/2)) / b - 1792*a^4*b*tan(x/2)) - (512*a^2*(a^2 - b^2)^(1/2)) / (256*a*b^2 - 64*b^3*tan(x/2) - 768*a^3 + (512*a^5) / b^2 - (1792*a^4*tan(x/2)) / b + (1024*a^6*tan(x/2)) / b^3 + 832*a^2*b*tan(x/2)) + (64*b^2*(a^2 - b^2)^(1/2)) / (256*a*b^2 - 64*b^3*tan(x/2) - 768*a^3 + (512*a^5) / b^2 - (1792*a^4*tan(x/2)) / b + (1024*a^6*tan(x/2)) / b^3 + 832*a^2*b*tan(x/2)) - (1280*a^3*tan(x/2)*(a^2 - b^2)^(1/2)) / (256*a*b^3 - 64*b^4*tan(x/2) - 1792*a^4*tan(x/2) - 768*a^3*b + (512*a^5) / b + 832*a^2*b^2*tan(x/2) + (1024*a^6*tan(x/2)) / b^2 + (1024*a^5*tan(x/2)*(a^2 - b^2)^(1/2)) / (1024*a^6*tan(x/2) - 64*b^6*tan(x/2) + 256*a*b^5 + 512*a^5*b - 768*a^3*b^3 + 832*a^2*b^4*tan(x/2) - 1792*a^4*b^2*tan(x/2) + (320*a*b*tan(x/2)*(a^2 - b^2)^(1/2)) / (256*a*b^2 - 64*b^3*tan(x/2) - 768*a^3 + (512*a^5) / b^2 - (1792*a^4*tan(x/2)) / b + (1024*a^6*tan(x/2)) / b^3 + 832*a^2*b*tan(x/2))) * (a^2 - b^2)^(1/2)) / (a*b) \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx = \frac{2\sqrt{-a^2 + b^2} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})b+a}{\sqrt{-a^2+b^2}}\right) + \log(\tan(\frac{x}{2}))a - bx}{ab}$$

input `int(cot(x)^2/(a+b*csc(x)),x)`

output
$$\frac{(2\sqrt{-a^2 + b^2}) \operatorname{atan}(\tan(x/2) \cdot b + a) / \sqrt{-a^2 + b^2}}{\tan(x/2) \cdot a - b \cdot x} + \log(\tan(x/2) \cdot a - b \cdot x) / (a \cdot b)$$

3.22 $\int \frac{\cot^4(x)}{a+b \csc(x)} dx$

Optimal result	181
Mathematica [A] (verified)	181
Rubi [A] (verified)	182
Maple [A] (verified)	185
Fricas [B] (verification not implemented)	186
Sympy [F]	186
Maxima [F(-2)]	187
Giac [B] (verification not implemented)	187
Mupad [B] (verification not implemented)	188
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 13, antiderivative size = 91

$$\begin{aligned} \int \frac{\cot^4(x)}{a + b \csc(x)} dx &= \frac{x}{a} - \frac{(2a^2 - 3b^2) \operatorname{arctanh}(\cos(x))}{2b^3} \\ &\quad + \frac{2(a^2 - b^2)^{3/2} \operatorname{arctanh}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{ab^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \end{aligned}$$

output $x/a - 1/2*(2*a^2 - 3*b^2)*\operatorname{arctanh}(\cos(x))/b^3 + 2*(a^2 - b^2)^{3/2}*\operatorname{arctanh}((a+b*tan(1/2*x))/(a^2 - b^2)^{1/2})/a/b^3 + a*cot(x)/b^2 - 1/2*cot(x)*csc(x)/b$

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.74

$$\begin{aligned} \int \frac{\cot^4(x)}{a + b \csc(x)} dx \\ = \frac{8b^3x - 16(-a^2 + b^2)^{3/2} \operatorname{arctan}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{-a^2+b^2}}\right) + 4a^2b \cot(\frac{x}{2}) - ab^2 \csc^2(\frac{x}{2}) - 8a^3 \log(\cos(\frac{x}{2})) + 12ab^2 \log(\cos(\frac{x}{2}))}{8ab^3} \end{aligned}$$

input `Integrate[Cot[x]^4/(a + b*Csc[x]), x]`

output

$$(8*b^3*x - 16*(-a^2 + b^2)^(3/2)*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]] + 4*a^2*b*Cot[x/2] - a*b^2*Csc[x/2]^2 - 8*a^3*Log[Cos[x/2]] + 12*a*b^2*Log[Cos[x/2]] + 8*a^3*Log[Sin[x/2]] - 12*a*b^2*Log[Sin[x/2]] + a*b^2*Sec[x/2]^2 - 4*a^2*b*Tan[x/2])/(8*a*b^3)$$

Rubi [A] (verified)

Time = 0.67 (sec), antiderivative size = 107, normalized size of antiderivative = 1.18, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.923, Rules used = {3042, 4386, 3042, 3372, 25, 3042, 3536, 3042, 3139, 1083, 219, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^4(x)}{a + b \csc(x)} dx \\ & \quad \downarrow \textcolor{blue}{3042} \\ & \int \frac{\cot(x)^4}{a + b \csc(x)} dx \\ & \quad \downarrow \textcolor{blue}{4386} \\ & \int \frac{\cos(x) \cot^3(x)}{a \sin(x) + b} dx \\ & \quad \downarrow \textcolor{blue}{3042} \\ & \int \frac{\cos(x)^4}{\sin(x)^3(a \sin(x) + b)} dx \\ & \quad \downarrow \textcolor{blue}{3372} \\ & - \frac{\int -\frac{\csc(x)(2a^2+b \sin(x)a-3b^2+2b^2 \sin^2(x))}{b+a \sin(x)} dx}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\ & \quad \downarrow \textcolor{blue}{25} \\ & \frac{\int \frac{\csc(x)(2a^2+b \sin(x)a-3b^2+2b^2 \sin^2(x))}{b+a \sin(x)} dx}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\ & \quad \downarrow \textcolor{blue}{3042} \end{aligned}$$

$$\begin{aligned}
 & \frac{\int \frac{2a^2+b \sin(x)a-3b^2+2b^2 \sin(x)^2}{\sin(x)(b+a \sin(x))} dx}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\
 & \quad \downarrow \textcolor{blue}{3536} \\
 & \frac{-\frac{2(a^2-b^2)^2 \int \frac{1}{b+a \sin(x)} dx}{ab} + \frac{(2a^2-3b^2) \int \csc(x) dx}{b} + \frac{2b^2x}{a}}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \frac{-\frac{2(a^2-b^2)^2 \int \frac{1}{b+a \sin(x)} dx}{ab} + \frac{(2a^2-3b^2) \int \csc(x) dx}{b} + \frac{2b^2x}{a}}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\
 & \quad \downarrow \textcolor{blue}{3139} \\
 & \frac{-\frac{4(a^2-b^2)^2 \int \frac{1}{b \tan^2(\frac{x}{2})+2a \tan(\frac{x}{2})+b} d \tan(\frac{x}{2})}{ab} + \frac{(2a^2-3b^2) \int \csc(x) dx}{b} + \frac{2b^2x}{a}}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\
 & \quad \downarrow \textcolor{blue}{1083} \\
 & \frac{\frac{8(a^2-b^2)^2 \int \frac{1}{4(a^2-b^2)-(2a+2b \tan(\frac{x}{2}))^2} d(2a+2b \tan(\frac{x}{2}))}{ab} + \frac{(2a^2-3b^2) \int \csc(x) dx}{b} + \frac{2b^2x}{a}}{2b^2} + \frac{a \cot(x)}{b^2} - \\
 & \quad \frac{\frac{\cot(x) \csc(x)}{2b}}{2b} \\
 & \quad \downarrow \textcolor{blue}{219} \\
 & \frac{\frac{(2a^2-3b^2) \int \csc(x) dx}{b} + \frac{4(a^2-b^2)^{3/2} \operatorname{arctanh}\left(\frac{2a+2b \tan(\frac{x}{2})}{2\sqrt{a^2-b^2}}\right)}{ab} + \frac{2b^2x}{a}}{2b^2} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} \\
 & \quad \downarrow \textcolor{blue}{4257} \\
 & \frac{-\frac{(2a^2-3b^2) \operatorname{arctanh}(\cos(x))}{b} + \frac{4(a^2-b^2)^{3/2} \operatorname{arctanh}\left(\frac{2a+2b \tan(\frac{x}{2})}{2\sqrt{a^2-b^2}}\right)}{ab}}{2b^2} + \frac{2b^2x}{a} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b}
 \end{aligned}$$

input `Int [Cot[x]^4/(a + b*Csc[x]),x]`

output `((2*b^2*x)/a - ((2*a^2 - 3*b^2)*ArcTanh[Cos[x]])/b + (4*(a^2 - b^2)^(3/2)*ArcTanh[(2*a + 2*b*Tan[x/2])/((2*Sqrt[a^2 - b^2])])/(a*b))/(2*b^2) + (a*Cot[x])/b^2 - (Cot[x]*Csc[x])/(2*b)`

Definitions of rubi rules used

rule 25 $\text{Int}[-(\text{Fx}__), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$

rule 219 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__)^2)^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[-\text{b}, 2]))*\text{ArcTanh}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \&& \text{NegQ}[\text{a}/\text{b}] \&& (\text{GtQ}[\text{a}, 0] \mid\mid \text{LtQ}[\text{b}, 0])$

rule 1083 $\text{Int}[(\text{a}__) + (\text{b}__)*(\text{x}__) + (\text{c}__)*(\text{x}__)^2)^{(-1)}, \text{x_Symbol}] \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[\text{b}^2 - 4*\text{a}*\text{c} - \text{x}^2, \text{x}], \text{x}], \text{x}, \text{b} + 2*\text{c}*\text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$

rule 3042 $\text{Int}[\text{u}__, \text{x_Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[\text{u}, \text{x}], \text{x}] /; \text{FunctionOfTrigOfLinearQ}[\text{u}, \text{x}]$

rule 3139 $\text{Int}[(\text{a}__) + (\text{b}__)*\text{sin}[(\text{c}__) + (\text{d}__)*(\text{x}__)])^{(-1)}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{e} = \text{FreeFactors}[\text{Tan}[(\text{c} + \text{d}*\text{x})/2], \text{x}]\}, \text{Simp}[2*(\text{e}/\text{d}) \quad \text{Subst}[\text{Int}[1/(\text{a} + 2*\text{b}*\text{e}*\text{x} + \text{a}*\text{e}^2*\text{x}^2), \text{x}], \text{x}, \text{Tan}[(\text{c} + \text{d}*\text{x})/2]/\text{e}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}] \&& \text{NeQ}[\text{a}^2 - \text{b}^2, 0]$

rule 3372 $\text{Int}[\cos[(\text{e}__) + (\text{f}__)*(\text{x}__)]^{4*((\text{d}__)*\text{sin}[(\text{e}__) + (\text{f}__)*(\text{x}__)])^{(\text{n}__)}}*(\text{a}__) + (\text{b}__)*\text{sin}[(\text{e}__) + (\text{f}__)*(\text{x}__)])^{(\text{m}__)}, \text{x_Symbol}] \rightarrow \text{Simp}[\text{Cos}[\text{e} + \text{f}*\text{x}]*(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*\text{x}])^{(\text{m} + 1)}*((\text{d}*\text{Sin}[\text{e} + \text{f}*\text{x}])^{(\text{n} + 1)}/(\text{a}*\text{d}*\text{f}*(\text{n} + 1))), \text{x}] + (-\text{Sin}[\text{b}*(\text{m} + \text{n} + 2)*\text{Cos}[\text{e} + \text{f}*\text{x}]*(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*\text{x}])^{(\text{m} + 1)}*((\text{d}*\text{Sin}[\text{e} + \text{f}*\text{x}])^{(\text{n} + 2)}/(\text{a}^2*\text{d}^2*\text{f}*(\text{n} + 1)*(\text{n} + 2))), \text{x}] - \text{Simp}[1/(\text{a}^2*\text{d}^2*(\text{n} + 1)*(\text{n} + 2)) \quad \text{Int}[(\text{a} + \text{b}*\text{Sin}[\text{e} + \text{f}*\text{x}])^{\text{m}}*(\text{d}*\text{Sin}[\text{e} + \text{f}*\text{x}])^{(\text{n} + 2)}*\text{Simp}[\text{a}^{2*\text{n}}*(\text{n} + 2) - \text{b}^{2*(\text{m} + \text{n} + 2)}*(\text{m} + \text{n} + 3) + \text{a}*\text{b}*\text{m}*\text{Sin}[\text{e} + \text{f}*\text{x}] - (\text{a}^{2*(\text{n} + 1)}*(\text{n} + 2) - \text{b}^{2*(\text{m} + \text{n} + 2)}*(\text{m} + \text{n} + 4))*\text{Sin}[\text{e} + \text{f}*\text{x}]^2, \text{x}], \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{d}, \text{e}, \text{f}, \text{m}\}, \text{x}] \&& \text{NeQ}[\text{a}^2 - \text{b}^2, 0] \&& (\text{IGtQ}[\text{m}, 0] \mid\mid \text{IntegersQ}[2*\text{m}, 2*\text{n}]) \&& \text{!m} < -1 \&& \text{LtQ}[\text{n}, -1] \&& (\text{LtQ}[\text{n}, -2] \mid\mid \text{EqQ}[\text{m} + \text{n} + 4, 0])$

rule 3536 $\text{Int}[(A_{\cdot}) + (B_{\cdot}) \sin[(e_{\cdot}) + (f_{\cdot})x_{\cdot}] + (C_{\cdot}) \sin[(e_{\cdot}) + (f_{\cdot})x_{\cdot}]^2 / (((a_{\cdot}) + (b_{\cdot}) \sin[(e_{\cdot}) + (f_{\cdot})x_{\cdot}]) * ((c_{\cdot}) + (d_{\cdot}) \sin[(e_{\cdot}) + (f_{\cdot})x_{\cdot}]))], x_{\text{Symbol}} :> \text{Simp}[C*(x/(b*d)), x] + (\text{Simp}[(A*b^2 - a*b*B + a^2*C) / (b*(b*c - a*d)) \text{Int}[1/(a + b*\text{Sin}[e + f*x]), x], x] - \text{Simp}[(c^2*C - B*c*d + A*d^2) / (d*(b*c - a*d)) \text{Int}[1/(c + d*\text{Sin}[e + f*x]), x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, C\}, x] \&& \text{NeQ}[b*c - a*d, 0] \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{NeQ}[c^2 - d^2, 0]$

rule 4257 $\text{Int}[\csc[(c_{\cdot}) + (d_{\cdot})x_{\cdot}], x_{\text{Symbol}} :> \text{Simp}[-\text{ArcTanh}[\cos[c + d*x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4386 $\text{Int}[\cot[(c_{\cdot}) + (d_{\cdot})x_{\cdot}]^m * (\csc[(c_{\cdot}) + (d_{\cdot})x_{\cdot}] * (b_{\cdot}) + (a_{\cdot}))^{n-m}, x_{\text{Symbol}} :> \text{Int}[\cos[c + d*x]^m * ((b + a*\text{Sin}[c + d*x])^n / \sin[c + d*x]^{m+n}), x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n] \&& \text{IntegerQ}[m] \&& (\text{IntegerQ}[m/2] \text{||} \text{LeQ}[m, 1])]$

Maple [A] (verified)

Time = 0.51 (sec), antiderivative size = 141, normalized size of antiderivative = 1.55

method	result
default	$-\frac{b \tan(\frac{x}{2})^2 + 2a \tan(\frac{x}{2})}{4b^2} - \frac{1}{8b \tan(\frac{x}{2})^2} + \frac{(4a^2 - 6b^2) \ln(\tan(\frac{x}{2}))}{4b^3} + \frac{a}{2b^2 \tan(\frac{x}{2})} + \frac{2 \arctan(\tan(\frac{x}{2}))}{a} + \frac{(-8a^4 + 16a^2b^2 - 4a^2b^4)}{4b^4}$
risch	$\frac{x}{a} + \frac{2ia e^{2ix} + b e^{3ix} - 2ia + b e^{ix}}{(e^{2ix} - 1)^2 b^2} + \frac{\ln(e^{ix} - 1) a^2}{b^3} - \frac{3 \ln(e^{ix} - 1)}{2b} - \frac{\ln(e^{ix} + 1) a^2}{b^3} + \frac{3 \ln(e^{ix} + 1)}{2b} + \frac{\sqrt{a^2 - b^2} a \ln\left(e^{ix} + \frac{ib + \sqrt{a^2 - b^2}}{a}\right)}{b^3}$

input `int(cot(x)^4/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/4/b^2*(-1/2*b*\tan(1/2*x)^2+2*a*\tan(1/2*x))-1/8/b/\tan(1/2*x)^2+1/4/b^3*(\\ & 4*a^2-6*b^2)*\ln(\tan(1/2*x))+1/2*a/b^2/\tan(1/2*x)+2/a*\arctan(\tan(1/2*x))+1/ \\ & 4*(-8*a^4+16*a^2*b^2-8*b^4)/a/b^3/(-a^2+b^2)^(1/2)*\arctan(1/2*(2*b*\tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 203 vs. $2(81) = 162$.

Time = 0.19 (sec) , antiderivative size = 453, normalized size of antiderivative = 4.98

$$\int \frac{\cot^4(x)}{a + b \csc(x)} dx \\ = \left[\frac{4 b^3 x \cos(x)^2 - 4 a^2 b \cos(x) \sin(x) - 4 b^3 x + 2 a b^2 \cos(x) - 2 ((a^2 - b^2) \cos(x)^2 - a^2 + b^2) \sqrt{a^2 - b^2} \log(-((a^2 - b^2) \cos(x)^2 - a^2 + b^2))}{\dots} \right]$$

input `integrate(cot(x)^4/(a+b*csc(x)),x, algorithm="fricas")`

output $[1/4*(4*b^3*x*cos(x)^2 - 4*a^2*b*cos(x)*sin(x) - 4*b^3*x + 2*a*b^2*cos(x) - 2*((a^2 - b^2)*cos(x)^2 - a^2 + b^2)*sqrt(a^2 - b^2)*log(-((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 - 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) + (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(1/2*cos(x) + 1/2) - (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(-1/2*cos(x) + 1/2))/(a*b^3*cos(x)^2 - a*b^3), 1/4*(4*b^3*x*cos(x)^2 - 4*a^2*b*cos(x)*sin(x) - 4*b^3*x + 2*a*b^2*cos(x) + 4*((a^2 - b^2)*cos(x)^2 - a^2 + b^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2 - b^2)*cos(x))) + (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(1/2*cos(x) + 1/2) - (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(-1/2*cos(x) + 1/2))/(a*b^3*cos(x)^2 - a*b^3)]$

Sympy [F]

$$\int \frac{\cot^4(x)}{a + b \csc(x)} dx = \int \frac{\cot^4(x)}{a + b \csc(x)} dx$$

input `integrate(cot(x)**4/(a+b*csc(x)),x)`

output `Integral(cot(x)**4/(a + b*csc(x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^4(x)}{a + b \csc(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(cot(x)^4/(a+b*csc(x)),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more de

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(81) = 162$.

Time = 0.19 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.79

$$\begin{aligned} \int \frac{\cot^4(x)}{a + b \csc(x)} dx = & \frac{x}{a} + \frac{b \tan\left(\frac{1}{2}x\right)^2 - 4a \tan\left(\frac{1}{2}x\right)}{8b^2} + \frac{(2a^2 - 3b^2) \log(|\tan(\frac{1}{2}x)|)}{2b^3} \\ & - \frac{2(a^4 - 2a^2b^2 + b^4) \left(\pi \lfloor \frac{x}{2\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(b) + \arctan\left(\frac{b \tan(\frac{1}{2}x) + a}{\sqrt{-a^2 + b^2}}\right) \right)}{\sqrt{-a^2 + b^2} ab^3} \\ & - \frac{12a^2 \tan\left(\frac{1}{2}x\right)^2 - 18b^2 \tan\left(\frac{1}{2}x\right)^2 - 4ab \tan\left(\frac{1}{2}x\right) + b^2}{8b^3 \tan\left(\frac{1}{2}x\right)^2} \end{aligned}$$

input `integrate(cot(x)^4/(a+b*csc(x)),x, algorithm="giac")`

output $x/a + 1/8*(b*\tan(1/2*x)^2 - 4*a*\tan(1/2*x))/b^2 + 1/2*(2*a^2 - 3*b^2)*\log(\operatorname{abs}(\tan(1/2*x)))/b^3 - 2*(a^4 - 2*a^2*b^2 + b^4)*(\pi*\operatorname{floor}(1/2*x/\pi + 1/2)*\operatorname{sgn}(b) + \arctan((b*\tan(1/2*x) + a)/\sqrt{-a^2 + b^2})) / (\sqrt{-a^2 + b^2} * a * b^3) - 1/8*(12*a^2*\tan(1/2*x)^2 - 18*b^2*\tan(1/2*x)^2 - 4*a*b*\tan(1/2*x) + b^2) / (b^3*\tan(1/2*x)^2)$

Mupad [B] (verification not implemented)

Time = 16.01 (sec) , antiderivative size = 2163, normalized size of antiderivative = 23.77

$$\int \frac{\cot^4(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `int(cot(x)^4/(a + b/sin(x)),x)`

output

```
(a^3*log(sin(x/2)/cos(x/2)))/(a*b^3 - a*b^3*cos(2*x)) - (2*b^3*atan((2*b^3*cos(x/2) + 2*a^3*sin(x/2) - 3*a*b^2*sin(x/2))/(2*b^3*sin(x/2) - 2*a^3*cos(x/2) + 3*a*b^2*cos(x/2)))/(a*b^3 - a*b^3*cos(2*x)) - (atan((32*a^6*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) - 14*b^6*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) - 14*b^12*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) + 19*a*b^5*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) + 16*a^5*b*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) + 13*a*b^11*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) - 36*a^3*b^3*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) - 24*a^3*b^9*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) + 8*a^5*b^7*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) + 2*a^7*b^5*cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) - 82*a^4*b^2*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(3/2) + 72*a^2*b^10*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) - 106*a^4*b^8*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) + 56*a^6*b^6*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) - 11*a^8*b^4*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2) + 2*a^10*b^2*sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^(1/2))/(a^15*sin(x/2)*32i + a^2*b^13*cos(x/2)*27i - a^4*b^11*cos(x/2)*165i + a^6*b^9*cos(x/2)*390i - a^8*b^7*cos(x/2)*469i + a^10*b^5*cos(x/2)*309i - a^12*b^3*cos(x/2)*108i + a^3*b^12*sin(x/2)*108i - a^5...
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.67

$$\begin{aligned} & \int \frac{\cot^4(x)}{a + b \csc(x)} dx \\ &= \frac{4\sqrt{-a^2 + b^2} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})b+a}{\sqrt{-a^2+b^2}}\right) \sin(x)^2 a^2 - 4\sqrt{-a^2 + b^2} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})b+a}{\sqrt{-a^2+b^2}}\right) \sin(x)^2 b^2 + 2 \cos(x) \sin(x) a^2 b^2}{2 \sin(x)^2 a b^3} \end{aligned}$$

input `int(cot(x)^4/(a+b*csc(x)),x)`

output
$$\begin{aligned} & (4*\sqrt{-a^2 + b^2}*\text{atan}((\tan(x/2)*b + a)/\sqrt{-a^2 + b^2})*\sin(x) \\ & \quad - 4*\sqrt{-a^2 + b^2}*\text{atan}((\tan(x/2)*b + a)/\sqrt{-a^2 + b^2}) \\ & \quad *(\sin(x)^2*b^2 + 2*\cos(x)*\sin(x)*a^2*b - \cos(x)*a*b^2 + 2*\log(\tan(x/2)) * \sin(x)^2*a^3 - 3*\log(\tan(x/2))*\sin(x)^2*a*b^2 + 2*\sin(x)^2*b^3*x) \\ & \quad)/(2*\sin(x)^2*a*b^3) \end{aligned}$$

3.23 $\int \frac{\cot^6(x)}{a+b \csc(x)} dx$

Optimal result	190
Mathematica [A] (verified)	191
Rubi [A] (verified)	191
Maple [A] (verified)	193
Fricas [B] (verification not implemented)	194
Sympy [F]	195
Maxima [F(-2)]	195
Giac [A] (verification not implemented)	196
Mupad [B] (verification not implemented)	196
Reduce [B] (verification not implemented)	197

Optimal result

Integrand size = 13, antiderivative size = 186

$$\begin{aligned} \int \frac{\cot^6(x)}{a + b \csc(x)} dx = & -\frac{x}{a} - \frac{3 \operatorname{arctanh}(\cos(x))}{8b} - \frac{(a^2 - 3b^2) \operatorname{arctanh}(\cos(x))}{2b^3} \\ & - \frac{(a^4 - 3a^2b^2 + 3b^4) \operatorname{arctanh}(\cos(x))}{b^5} \\ & + \frac{2(a^2 - b^2)^{5/2} \operatorname{arctanh}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{ab^5} + \frac{a \cot(x)}{b^2} \\ & + \frac{a(a^2 - 3b^2) \cot(x)}{b^4} + \frac{a \cot^3(x)}{3b^2} - \frac{3 \cot(x) \csc(x)}{8b} \\ & - \frac{(a^2 - 3b^2) \cot(x) \csc(x)}{2b^3} - \frac{\cot(x) \csc^3(x)}{4b} \end{aligned}$$

output

```
-x/a-3/8*arctanh(cos(x))/b-1/2*(a^2-3*b^2)*arctanh(cos(x))/b^3-(a^4-3*a^2*b^2+3*b^4)*arctanh(cos(x))/b^5+2*(a^2-b^2)^(5/2)*arctanh((a+b*tan(1/2*x))/(a^2-b^2)^(1/2))/a/b^5+a*cot(x)/b^2+a*(a^2-3*b^2)*cot(x)/b^4+1/3*a*cot(x)^3/b^2-3/8*cot(x)*csc(x)/b-1/2*(a^2-3*b^2)*cot(x)*csc(x)/b^3-1/4*cot(x)*csc(x)^3/b
```

Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 312, normalized size of antiderivative = 1.68

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx \\ = \frac{-192b^5x + 384(-a^2 + b^2)^{5/2} \arctan\left(\frac{a+b\tan(\frac{x}{2})}{\sqrt{-a^2+b^2}}\right) + 32a^2b(3a^2 - 7b^2)\cot\left(\frac{x}{2}\right) - 24a^3b^2\csc^2\left(\frac{x}{2}\right) + 54ab^4\csc(x)}{}$$

input `Integrate[Cot[x]^6/(a + b*Csc[x]), x]`

output
$$(-192*b^5*x + 384*(-a^2 + b^2)^{(5/2)}*\text{ArcTan}[(a + b*\text{Tan}[x/2])/\text{Sqrt}[-a^2 + b^2]] + 32*a^2*b*(3*a^2 - 7*b^2)*\text{Cot}[x/2] - 24*a^3*b^2*\text{Csc}[x/2]^2 + 54*a*b^4*\text{Csc}[x/2]^2 - 3*a*b^4*\text{Csc}[x/2]^4 - 192*a^5*\text{Log}[\text{Cos}[x/2]] + 480*a^3*b^2*\text{Log}[\text{Cos}[x/2]] - 360*a*b^4*\text{Log}[\text{Cos}[x/2]] + 192*a^5*\text{Log}[\text{Sin}[x/2]] - 480*a^3*b^2*\text{Log}[\text{Sin}[x/2]] + 360*a*b^4*\text{Log}[\text{Sin}[x/2]] + 24*a^3*b^2*\text{Sec}[x/2]^2 - 54*a*b^4*\text{Sec}[x/2]^2 + 3*a*b^4*\text{Sec}[x/2]^4 - 64*a^2*b^3*\text{Csc}[x]^3*\text{Sin}[x/2]^4 + 4*a^2*b^3*\text{Csc}[x/2]^4*\text{Sin}[x] - 96*a^4*b*\text{Tan}[x/2] + 224*a^2*b^3*\text{Tan}[x/2])/(192*a*b^5)$$

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 4386, 3042, 3376, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx \\ \downarrow 3042 \\ \int \frac{\cot(x)^6}{a + b \csc(x)} dx \\ \downarrow 4386$$

$$\begin{aligned}
 & \int \frac{\cos(x) \cot^5(x)}{a \sin(x) + b} dx \\
 & \quad \downarrow \textcolor{blue}{3042} \\
 & \int \frac{\cos(x)^6}{\sin(x)^5(a \sin(x) + b)} dx \\
 & \quad \downarrow \textcolor{blue}{3376} \\
 & \int \left(\frac{(3ab^2 - a^3) \csc^2(x)}{b^4} - \frac{(a^2 - b^2)^3}{ab^5(a \sin(x) + b)} + \frac{(a^2 - 3b^2) \csc^3(x)}{b^3} + \frac{(a^4 - 3a^2b^2 + 3b^4) \csc(x)}{b^5} - \frac{a \csc^4(x)}{b^2} - \frac{1}{a} \right. \\
 & \quad \downarrow \textcolor{blue}{2009} \\
 & \frac{2(a^2 - b^2)^{5/2} \operatorname{arctanh}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{ab^5} - \frac{(a^2 - 3b^2) \operatorname{arctanh}(\cos(x))}{2b^3} + \frac{a(a^2 - 3b^2) \cot(x)}{b^4} - \\
 & \frac{(a^2 - 3b^2) \cot(x) \csc(x)}{2b^3} - \frac{(a^4 - 3a^2b^2 + 3b^4) \operatorname{arctanh}(\cos(x))}{b^5} + \frac{a \cot^3(x)}{3b^2} + \frac{a \cot(x)}{b^2} - \frac{x}{a} - \\
 & \left. \frac{3 \operatorname{arctanh}(\cos(x))}{8b} - \frac{\cot(x) \csc^3(x)}{4b} - \frac{3 \cot(x) \csc(x)}{8b} \right)
 \end{aligned}$$

input `Int[Cot[x]^6/(a + b*Csc[x]),x]`

output `-(x/a) - (3*ArcTanh[Cos[x]])/(8*b) - ((a^2 - 3*b^2)*ArcTanh[Cos[x]])/(2*b^3) - ((a^4 - 3*a^2*b^2 + 3*b^4)*ArcTanh[Cos[x]])/b^5 + (2*(a^2 - b^2)^(5/2)*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a*b^5) + (a*Cot[x])/b^2 + (a*(a^2 - 3*b^2)*Cot[x])/b^4 + (a*Cot[x]^3)/(3*b^2) - (3*Cot[x]*Csc[x])/(8*b) - ((a^2 - 3*b^2)*Cot[x]*Csc[x])/(2*b^3) - (Cot[x]*Csc[x]^3)/(4*b)`

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3376 $\text{Int}[\cos[(e_.) + (f_.)*(x_)]^{(p_.)}*((d_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)}*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_.)}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandTrig}[(d*\sin[e + f*x])^n*(a + b*\sin[e + f*x])^m*(1 - \sin[e + f*x]^2)^{(p/2)}, x], x] /; \text{FreeQ}[\{a, b, d, e, f\}, x] \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegersQ}[m, 2*n, p/2] \&& (\text{LtQ}[m, -1] \text{ || } (\text{EqQ}[m, -1] \&& \text{GtQ}[p, 0]))$

rule 4386 $\text{Int}[\cot[(c_.) + (d_.)*(x_)]^{(m_.)}*(\csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^{(n_.)}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{Cos}[c + d*x]^m*((b + a*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^{(m + n)}), x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n] \&& \text{IntegerQ}[m] \&& (\text{IntegerQ}[m/2] \text{ || } \text{LeQ}[m, 1])$

Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.35

method	result
default	$-\frac{\tan(\frac{x}{2})^4 b^3}{4} + \frac{2 \tan(\frac{x}{2})^3 a b^2}{3} - 2 \tan(\frac{x}{2})^2 a^2 b + 4 \tan(\frac{x}{2})^2 b^3 + 8 \tan(\frac{x}{2}) a^3 - 18 \tan(\frac{x}{2}) a b^2 - \frac{1}{64 b \tan(\frac{x}{2})^4} - \frac{4 a^2 - 8 b^2}{32 b^3 \tan(\frac{x}{2})^2} + \dots$
risch	$-\frac{x}{a} - \frac{72 i a b^2 e^{6ix} - 168 i a b^2 e^{4ix} - 12 a^2 b e^{7ix} + 27 b^3 e^{7ix} + 72 i a^3 e^{4ix} - 56 i a b^2 + 12 a^2 b e^{5ix} - 3 b^3 e^{5ix} - 72 i a^3 e^{2ix} + 24 i a^3 + 12 a^2 b e^{3ix} - 12 b^4 (e^{2ix} - 1)^4}{12 b^4 (e^{2ix} - 1)^4}$

input `int(cot(x)^6/(a+b*csc(x)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/16/b^4*(-1/4*tan(1/2*x)^4*b^3+2/3*tan(1/2*x)^3*a*b^2-2*tan(1/2*x)^2*a^2 \\ & *b+4*tan(1/2*x)^2*b^3+8*tan(1/2*x)*a^3-18*tan(1/2*x)*a*b^2)-1/64/b/tan(1/2*x)^4-1/32*(4*a^2-8*b^2)/b^3/tan(1/2*x)^2+1/16/b^5*(16*a^4-40*a^2*b^2+30*b^4)*ln(tan(1/2*x))+1/24*a/b^2/tan(1/2*x)^3+1/8*a*(4*a^2-9*b^2)/b^4/tan(1/2*x)-2/a*arctan(tan(1/2*x))+1/16*(-32*a^6+96*a^4*b^2-96*a^2*b^4+32*b^6)/b^5 \\ & /a/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*tan(1/2*x)+2*a)/(-a^2+b^2)^(1/2)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 403 vs. $2(168) = 336$.

Time = 0.38 (sec) , antiderivative size = 852, normalized size of antiderivative = 4.58

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `integrate(cot(x)^6/(a+b*csc(x)),x, algorithm="fricas")`

output

```
[-1/48*(48*b^5*x*cos(x)^4 - 96*b^5*x*cos(x)^2 + 48*b^5*x - 6*(4*a^3*b^2 - 9*a*b^4)*cos(x)^3 - 24*((a^4 - 2*a^2*b^2 + b^4)*cos(x)^4 + a^4 - 2*a^2*b^2 + b^4 - 2*(a^4 - 2*a^2*b^2 + b^4)*cos(x)^2)*sqrt(a^2 - b^2)*log((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) + 6*(4*a^3*b^2 - 7*a*b^4)*cos(x) + 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(1/2*cos(x) + 1/2) - 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(-1/2*cos(x) + 1/2) + 16*((3*a^4*b - 7*a^2*b^3)*cos(x)^3 - 3*(a^4*b - 2*a^2*b^3)*cos(x)^2*sin(x))/(a*b^5*cos(x)^4 - 2*a*b^5*cos(x)^2 + a*b^5), -1/48*(48*b^5*x*cos(x)^4 - 96*b^5*x*cos(x)^2 + 48*b^5*x - 6*(4*a^3*b^2 - 9*a*b^4)*cos(x)^3 - 48*((a^4 - 2*a^2*b^2 + b^4)*cos(x)^4 + a^4 - 2*a^2*b^2 + b^4 - 2*(a^4 - 2*a^2*b^2 + b^4)*cos(x)^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2 - b^2)*cos(x))) + 6*(4*a^3*b^2 - 7*a*b^4)*cos(x) + 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(1/2*cos(x) + 1/2) - 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(-1/2*cos(x) + 1/2) + 16*((3*a^4*b - 7*a^2*b^3)*cos(x)^3 - 3*(a^4*b - 2*a^2*b^3)*cos(x))*sin(x)...]
```

Sympy [F]

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx = \int \frac{\cot^6(x)}{a + b \csc(x)} dx$$

input `integrate(cot(x)**6/(a+b*csc(x)),x)`

output `Integral(cot(x)**6/(a + b*csc(x)), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx = \text{Exception raised: ValueError}$$

input `integrate(cot(x)^6/(a+b*csc(x)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more de`

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.61

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx = -\frac{x}{a} + \frac{3 b^3 \tan\left(\frac{1}{2}x\right)^4 - 8 a b^2 \tan\left(\frac{1}{2}x\right)^3 + 24 a^2 b \tan\left(\frac{1}{2}x\right)^2 - 48 b^3 \tan\left(\frac{1}{2}x\right)^2 - 96 a^3 \tan\left(\frac{1}{2}x\right) + 216 a b^2 \tan\left(\frac{1}{2}x\right)}{192 b^4} + \frac{(8 a^4 - 20 a^2 b^2 + 15 b^4) \log\left(|\tan\left(\frac{1}{2}x\right)|\right)}{8 b^5} - \frac{2 (a^6 - 3 a^4 b^2 + 3 a^2 b^4 - b^6) \left(\pi \lfloor \frac{x}{2\pi} + \frac{1}{2} \rfloor \operatorname{sgn}(b) + \arctan\left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}}\right)\right)}{\sqrt{-a^2 + b^2} ab^5} - \frac{400 a^4 \tan\left(\frac{1}{2}x\right)^4 - 1000 a^2 b^2 \tan\left(\frac{1}{2}x\right)^4 + 750 b^4 \tan\left(\frac{1}{2}x\right)^4 - 96 a^3 b \tan\left(\frac{1}{2}x\right)^3 + 216 a b^3 \tan\left(\frac{1}{2}x\right)^3 + 192 b^5 \tan\left(\frac{1}{2}x\right)^4}{192 b^5 \tan\left(\frac{1}{2}x\right)^4}$$

input `integrate(cot(x)^6/(a+b*csc(x)),x, algorithm="giac")`

output

```
-x/a + 1/192*(3*b^3*tan(1/2*x)^4 - 8*a*b^2*tan(1/2*x)^3 + 24*a^2*b*tan(1/2*x)^2 - 48*b^3*tan(1/2*x)^2 - 96*a^3*tan(1/2*x) + 216*a*b^2*tan(1/2*x))/b^4 + 1/8*(8*a^4 - 20*a^2*b^2 + 15*b^4)*log(abs(tan(1/2*x)))/b^5 - 2*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))/(sqrt(-a^2 + b^2)*a*b^5) - 1/192*(400*a^4*tan(1/2*x)^4 - 1000*a^2*b^2*tan(1/2*x)^4 + 750*b^4*tan(1/2*x)^4 - 96*a^3*b*tan(1/2*x)^3 + 216*a*b^3*tan(1/2*x)^3 + 24*a^2*b^2*tan(1/2*x)^2 - 48*b^4*tan(1/2*x)^2 - 8*a*b^3*tan(1/2*x) + 3*b^4)/(b^5*tan(1/2*x)^4)
```

Mupad [B] (verification not implemented)

Time = 16.89 (sec) , antiderivative size = 4075, normalized size of antiderivative = 21.91

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx = \text{Too large to display}$$

input `int(cot(x)^6/(a + b/sin(x)),x)`

output

$$\begin{aligned}
 & \tan(x/2) * (a/(8*b^2) + (2*a*(1/(2*b) - a^2/(4*b^3)))/b) - \tan(x/2)^2 * (1/(4*b) - a^2/(8*b^3)) + \tan(x/2)^4/(64*b) - (2*atan(((4*(24*a*b^16 + 58*a^3*b^14 - 345*a^5*b^12 + 543*a^7*b^10 - 440*a^9*b^8 + 184*a^11*b^6 - 32*a^13*b^4))/b^12 + (((4*(24*a^3*b^16 - 32*a^5*b^14))/b^12 - (4*tan(x/2)*(16*a^2*b^17 - 136*a^4*b^15 + 128*a^6*b^13))/b^12)*1i)/a - (4*(93*a^3*b^15 - 240*a^5*b^13 + 208*a^7*b^11 - 64*a^9*b^9))/b^12 + (4*tan(x/2)*(62*a^2*b^16 - 335*a^4*b^14 + 604*a^6*b^12 - 456*a^8*b^10 + 128*a^10*b^8))/b^12)*1i)/a - (4*tan(x/2)*(16*b^17 - 180*a^2*b^15 + 252*a^4*b^13 - 148*a^6*b^11 + 100*a^8*b^9 - 40*a^10*b^7 + 8*a^12*b^5))/b^12)*1i)/a - (4*(53*a*b^15 - 120*a^3*b^13 + 48*a^5*b^11 + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^11*b^5))/b^12 + (4*tan(x/2)*(8*a^16 + 62*b^16 - 410*a^2*b^14 + 929*a^4*b^12 - 1096*a^6*b^10 + 873*a^8*b^8 - 550*a^10*b^6 + 255*a^12*b^4 - 68*a^14*b^2))/b^12)/a - ((4*(53*a*b^15 - 120*a^3*b^13 + 48*a^5*b^11 + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^11*b^5))/b^12 + ((4*(24*a*b^16 + 58*a^3*b^14 - 345*a^5*b^12 + 543*a^7*b^10 - 440*a^9*b^8 + 184*a^11*b^6 - 32*a^13*b^4))/b^12 + (((4*(93*a^3*b^15 - 240*a^5*b^13 + 208*a^7*b^11 - 64*a^9*b^9))/b^12 + (((4*(24*a^3*b^16 - 32*a^5*b^14))/b^12 - (4*tan(x/2)*(16*a^2*b^17 - 136*a^4*b^15 + 128*a^6*b^13))/b^12)*1i)/a - (4*tan(x/2)*(62*a^2*b^16 - 335*a^4*b^14 + 604*a^6*b^12 - 456*a^8*b^10 + 128*a^10*b^8))/b^12)*1i)/a - (4*tan(x/2)*(16*b^17 - 180*a^2*b^15 + 252*a^4*b^13 - 148*a^6*b^11 + 100*a^8*b^9 - 40*a^10*b^7 + 8*a^12*b^5))/b^12...
 \end{aligned}$$

Reduce [B] (verification not implemented)

Time = 0.19 (sec), antiderivative size = 267, normalized size of antiderivative = 1.44

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx = \frac{48\sqrt{-a^2 + b^2} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})b+a}{\sqrt{-a^2+b^2}}\right) \sin(x)^4 a^4 - 96\sqrt{-a^2 + b^2} \operatorname{atan}\left(\frac{\tan(\frac{x}{2})b+a}{\sqrt{-a^2+b^2}}\right) \sin(x)^4 a^2 b^2 + 48\sqrt{-a^2 + b^2} a^4 \sin(x)^4 b^4}{a^4 b^4}$$

input

```
int(cot(x)^6/(a+b*csc(x)),x)
```

```
output (48*sqrt( - a**2 + b**2)*atan((tan(x/2)*b + a)/sqrt( - a**2 + b**2))*sin(x)**4*a**4 - 96*sqrt( - a**2 + b**2)*atan((tan(x/2)*b + a)/sqrt( - a**2 + b**2))*sin(x)**4*a**2*b**2 + 48*sqrt( - a**2 + b**2)*atan((tan(x/2)*b + a)/sqrt( - a**2 + b**2))*sin(x)**4*b**4 + 24*cos(x)*sin(x)**3*a**4*b - 56*cos(x)*sin(x)**3*a**2*b**3 - 12*cos(x)*sin(x)**2*a**3*b**2 + 27*cos(x)*sin(x)**2*a*b**4 + 8*cos(x)*sin(x)*a**2*b**3 - 6*cos(x)*a*b**4 + 24*log(tan(x/2))*sin(x)**4*a**5 - 60*log(tan(x/2))*sin(x)**4*a**3*b**2 + 45*log(tan(x/2))*sin(x)**4*a*b**4 - 24*sin(x)**4*b**5*x)/(24*sin(x)**4*a*b**5)
```

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	199
4.2 Links to plain text integration problems used in this report for each CAS .	217

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```
(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*) (*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","");
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "}
  ,
]
```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```



```

ExpnType[expn_] :=
If[AtomQ[expn],
  1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn] === Power,
      If[IntegerQ[expn[[2]]],
        ExpnType[expn[[1]]],
        If[Head[expn[[2]]] === Rational,
          If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
            1,
            Max[ExpnType[expn[[1]]], 2]],
          Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
      If[Head[expn] === Plus || Head[expn] === Times,
        Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
        If[ElementaryFunctionQ[Head[expn]],
          Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3],
              Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], ExpnType[expn[[3]]]]]]]]]]]

```

```
Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],  
If[AppellFunctionQ[Head[expn]],  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],  
If[Head[expn]==RootSum,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],  
If[Head[expn]==Integrate || Head[expn]==Int,  
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],  
9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=  
MemberQ[{  
Exp, Log,  
Sin, Cos, Tan, Cot, Sec, Csc,  
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  
Sinh, Cosh, Tanh, Coth, Sech, Csch,  
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  
} , func]
```

```
SpecialFunctionQ[func_] :=  
MemberQ[{  
Erf, Erfc, Erfi,  
FresnelS, FresnelC,  
ExpIntegralE, ExpIntegralEi, LogIntegral,  

```

```
HypergeometricFunctionQ[func_] :=  
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=  
MemberQ[{AppellF1}, func]
```

Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issue
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
```

```
#      antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
            # this assumes optimal do not as well. No check is needed here.
            if debug then
                print("result do not contain complex, this assumes optimal do not as well");
            fi;
```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                           convert(leaf_count_result,string),\"$ vs. \$2(", 
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_co
                           fi;
            fi;
        else #ExpnType(result) > ExpnType(optimal)
            if debug then
                print("ExpnType(result) > ExpnType(optimal)");
            fi;
            return "C",cat("Result contains higher order function than in optimal. Order ",
                           convert(ExpnType_result,string)," vs. order ",
                           convert(ExpnType_optimal,string),"."));
        fi;

    end proc:

    #

    # is_contains_complex(result)
    # takes expressions and returns true if it contains "I" else false
    #
    #Nasser 032417
    is_contains_complex:= proc(expression)
        return (has(expression,I));
    end proc:

    # The following summarizes the type number assigned an expression
    # based on the functions it involves
    # 1 = rational function
    # 2 = algebraic function
    # 3 = elementary function
    # 4 = special function
    # 5 = hypergeometric function

```

```
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'`^`') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then
            if type(op(1,expn),'rational') then
                1
            else
                max(2,ExpnType(op(1,expn)))
            end if
        else
            max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
        end if
    elif type(expn,'`+`') or type(expn,'`*`') then
        max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif ElementaryFunctionQ(op(0,expn)) then
        max(3,ExpnType(op(1,expn)))
    elif SpecialFunctionQ(op(0,expn)) then
        max(4,apply(max,map(ExpnType,[op(expn)])))
    elif HypergeometricFunctionQ(op(0,expn)) then
        max(5,apply(max,map(ExpnType,[op(expn)])))
    elif AppellFunctionQ(op(0,expn)) then
        max(6,apply(max,map(ExpnType,[op(expn)])))
    elif op(0,expn)='int' then
        max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
```

```
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arccsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]
```

```
def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*')
```

```

m1 = expnType(expn.args[0])
m2 = expnType(list(expn.args[1:]))
return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sagemath")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(grade)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(grade)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```
from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                       ]
    if debug:
```

```

if m:
    print ("func ", func , " is elementary_function")
else:
    print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral'
'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
'elliptic_pi','exp_integral_e','log_integral',
'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
'weierstrassPPrime','weierstrassSigma']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):
    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0], Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1], Integer)

```

```

    return expnType(expn.operands()[0])  #expnType(expn.args[0])
elif type(expn.operands()[1]) == Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0]) == Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))  #max(3,expnType(expn.args[0]),expnType(expn.args[1]))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(4,m1)  #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(5,m1)  #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))      #max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

```

```
leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation ="none"
            else:
                grade = "B"
                grade_annotation ="Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation ="Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation ="none"
        else:
            grade = "B"
            grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(expnType_result)+"/"+str(expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file