

Computer Algebra Independent Integration Tests

Summer 2024

5-Inverse-trig-functions/5.3-Inverse-tangent/276-5.3

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 9:09pm

Contents

1	Introduction	7
1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26
2	detailed summary tables of results	27
2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	72
3	Listing of integrals	78
3.1	$\int x^3 \arctan(a + bx^4) dx$	84
3.2	$\int x^{-1+n} \arctan(a + bx^n) dx$	90
3.3	$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	96
3.4	$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	104
3.5	$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	111

3.6	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx$	117
3.7	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$	126
3.8	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$	131
3.9	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$	137
3.10	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$	144
3.11	$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	152
3.12	$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	159
3.13	$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	165
3.14	$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	171
3.15	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$	176
3.16	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$	182
3.17	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$	189
3.18	$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	196
3.19	$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	203
3.20	$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	210
3.21	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$	216
3.22	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$	222
3.23	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$	228
3.24	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$	235
3.25	$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	243
3.26	$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	251
3.27	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$	259
3.28	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$	266

3.29	$\int \frac{\arctan\left(\frac{\sqrt{-cx}}{\sqrt{d+cx^2}}\right)}{x^{9/2}} dx$	274
3.30	$\int \frac{\arctan(1+x+x^2)}{x^2} dx$	283
3.31	$\int \frac{(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^n}{1-c^2x^2} dx$	289
3.32	$\int \frac{(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^3}{1-c^2x^2} dx$	294
3.33	$\int \frac{(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^2}{1-c^2x^2} dx$	303
3.34	$\int \frac{a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1-c^2x^2} dx$	312
3.35	$\int \frac{1}{(1-c^2x^2)(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))} dx$	318
3.36	$\int \frac{1}{(1-c^2x^2)(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right))^2} dx$	323
3.37	$\int x^m \arctan(\tan(a+bx)) dx$	329
3.38	$\int x^2 \arctan(\tan(a+bx)) dx$	334
3.39	$\int x \arctan(\tan(a+bx)) dx$	339
3.40	$\int \arctan(\tan(a+bx)) dx$	344
3.41	$\int \frac{\arctan(\tan(a+bx))}{x} dx$	349
3.42	$\int x^m \arctan(\cot(a+bx)) dx$	354
3.43	$\int x^2 \arctan(\cot(a+bx)) dx$	360
3.44	$\int x \arctan(\cot(a+bx)) dx$	365
3.45	$\int \arctan(\cot(a+bx)) dx$	370
3.46	$\int \frac{\arctan(\cot(a+bx))}{x} dx$	375
3.47	$\int \arctan(\tan(a+bx)) dx$	380
3.48	$\int x^2 \arctan(c+d \tan(a+bx)) dx$	385
3.49	$\int x \arctan(c+d \tan(a+bx)) dx$	397
3.50	$\int \arctan(c+d \tan(a+bx)) dx$	406
3.51	$\int \frac{\arctan(c+d \tan(a+bx))}{x} dx$	414
3.52	$\int x^2 \arctan(c+(1+ic) \tan(a+bx)) dx$	419
3.53	$\int x \arctan(c+(1+ic) \tan(a+bx)) dx$	427
3.54	$\int \arctan(c+(1+ic) \tan(a+bx)) dx$	435
3.55	$\int \frac{\arctan(c+(1+ic) \tan(a+bx))}{x} dx$	442
3.56	$\int x^2 \arctan(c+(-1+ic) \tan(a+bx)) dx$	447
3.57	$\int x \arctan(c+(-1+ic) \tan(a+bx)) dx$	456
3.58	$\int \arctan(c+(-1+ic) \tan(a+bx)) dx$	464
3.59	$\int \frac{\arctan(c+(-1+ic) \tan(a+bx))}{x} dx$	472
3.60	$\int \arctan(\cot(a+bx)) dx$	477
3.61	$\int x^2 \arctan(c+d \cot(a+bx)) dx$	482
3.62	$\int x \arctan(c+d \cot(a+bx)) dx$	494
3.63	$\int \arctan(c+d \cot(a+bx)) dx$	503

3.64	$\int \frac{\arctan(c+d\cot(a+bx))}{x} dx$	512
3.65	$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx$	517
3.66	$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx$	526
3.67	$\int \arctan(c + (1 - ic) \cot(a + bx)) dx$	534
3.68	$\int \frac{\arctan(c+(1-ic)\cot(a+bx))}{x} dx$	542
3.69	$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx$	547
3.70	$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx$	555
3.71	$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx$	563
3.72	$\int \frac{\arctan(c+(-1-ic)\cot(a+bx))}{x} dx$	570
3.73	$\int \arctan(\sinh(x)) dx$	575
3.74	$\int x \arctan(\sinh(x)) dx$	581
3.75	$\int x^2 \arctan(\sinh(x)) dx$	588
3.76	$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx$	595
3.77	$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx$	606
3.78	$\int (e + fx) \arctan(\tanh(a + bx)) dx$	615
3.79	$\int \arctan(\tanh(a + bx)) dx$	623
3.80	$\int \frac{\arctan(\tanh(a+bx))}{e+fx} dx$	629
3.81	$\int x^2 \arctan(c + d \tanh(a + bx)) dx$	634
3.82	$\int x \arctan(c + d \tanh(a + bx)) dx$	644
3.83	$\int \arctan(c + d \tanh(a + bx)) dx$	653
3.84	$\int \frac{\arctan(c+d\tanh(a+bx))}{x} dx$	661
3.85	$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx$	666
3.86	$\int x \arctan(c + (i + c) \tanh(a + bx)) dx$	675
3.87	$\int \arctan(c + (i + c) \tanh(a + bx)) dx$	683
3.88	$\int \frac{\arctan(c+(i+c)\tanh(a+bx))}{x} dx$	690
3.89	$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx$	695
3.90	$\int x \arctan(c - (i - c) \tanh(a + bx)) dx$	703
3.91	$\int \arctan(c - (i - c) \tanh(a + bx)) dx$	711
3.92	$\int \frac{\arctan(c-(i-c)\tanh(a+bx))}{x} dx$	718
3.93	$\int (e + fx)^3 \arctan(\coth(a + bx)) dx$	723
3.94	$\int (e + fx)^2 \arctan(\coth(a + bx)) dx$	734
3.95	$\int (e + fx) \arctan(\coth(a + bx)) dx$	743
3.96	$\int \arctan(\coth(a + bx)) dx$	751
3.97	$\int \frac{\arctan(\coth(a+bx))}{e+fx} dx$	757
3.98	$\int x^2 \arctan(c + d \coth(a + bx)) dx$	762
3.99	$\int x \arctan(c + d \coth(a + bx)) dx$	772
3.100	$\int \arctan(c + d \coth(a + bx)) dx$	781
3.101	$\int \frac{\arctan(c+d\coth(a+bx))}{x} dx$	789
3.102	$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx$	794

3.103	$\int x \arctan(c + (i + c) \coth(a + bx)) dx$	803
3.104	$\int \arctan(c + (i + c) \coth(a + bx)) dx$	811
3.105	$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx$	818
3.106	$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx$	823
3.107	$\int x \arctan(c - (i - c) \coth(a + bx)) dx$	832
3.108	$\int \arctan(c - (i - c) \coth(a + bx)) dx$	840
3.109	$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx$	847
3.110	$\int \arctan(e^x) dx$	852
3.111	$\int x \arctan(e^x) dx$	857
3.112	$\int x^2 \arctan(e^x) dx$	862
3.113	$\int \arctan(e^{a+bx}) dx$	868
3.114	$\int x \arctan(e^{a+bx}) dx$	874
3.115	$\int x^2 \arctan(e^{a+bx}) dx$	880
3.116	$\int \arctan(a + bf^{c+dx}) dx$	887
3.117	$\int x \arctan(a + bf^{c+dx}) dx$	894
3.118	$\int x^2 \arctan(a + bf^{c+dx}) dx$	901
3.119	$\int e^{-x} \arctan(e^x) dx$	909
3.120	$\int \frac{\arctan(x)}{(-1+x)^3} dx$	915
3.121	$\int \frac{\arctan(1+2x)}{(4+3x)^3} dx$	921
3.122	$\int \arctan(\sqrt{1+x}) dx$	929
3.123	$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx$	935
3.124	$\int \frac{1}{(a+ax^2)(b-2b\arctan(x))} dx$	940
3.125	$\int \frac{x+x^3+(1+x)^2 \arctan(x)}{(1+x)^2(1+x^2)} dx$	945
3.126	$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx$	950
3.127	$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx$	957
3.128	$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx$	964
3.129	$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx$	970
3.130	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} dx$	976
3.131	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^2} dx$	982
3.132	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^3} dx$	989
3.133	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^4} dx$	995
3.134	$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$	1002
3.135	$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$	1008
3.136	$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$	1013

3.137 $\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx \dots\dots\dots 1018$

3.138 $\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx \dots\dots\dots 1024$

3.139 $\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^m} dx \dots\dots\dots 1029$

3.140 $\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2 x^2}}\right)}{\sqrt{a + bx^2}} dx \dots\dots\dots 1034$

3.141 $\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2 x^2}}\right)^2}{\sqrt{a + bx^2}} dx \dots\dots\dots 1040$

3.142 $\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2 x^2}}\right)}{\sqrt{a + bx^2}} dx \dots\dots\dots 1045$

3.143 $\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2 x^2}}\right)} dx \dots\dots\dots 1050$

3.144 $\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2 x^2}}\right)^2} dx \dots\dots\dots 1056$

3.145 $\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2 x^2}}\right)^3} dx \dots\dots\dots 1061$

3.146 $\int \frac{\arctan(c(a + bx)) \log(d(a + bx))}{a + bx} dx \dots\dots\dots 1066$

3.147 $\int e^{c(a + bx)} \arctan(\sinh(ac + bcx)) dx \dots\dots\dots 1073$

3.148 $\int e^{c(a + bx)} \arctan(\cosh(ac + bcx)) dx \dots\dots\dots 1079$

3.149 $\int e^{c(a + bx)} \arctan(\tanh(ac + bcx)) dx \dots\dots\dots 1087$

3.150 $\int e^{c(a + bx)} \arctan(\coth(ac + bcx)) dx \dots\dots\dots 1097$

3.151 $\int e^{c(a + bx)} \arctan(\operatorname{sech}(ac + bcx)) dx \dots\dots\dots 1107$

3.152 $\int e^{c(a + bx)} \arctan(\operatorname{csch}(ac + bcx)) dx \dots\dots\dots 1115$

3.153 $\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx \dots\dots\dots 1122$

4 Appendix 1128

4.1 Listing of Grading functions 1128

4.2 Links to plain text integration problems used in this report for each CAS 1146

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [153]. This is test number [276].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (153)	0.00 (0)
Mathematica	98.69 (151)	1.31 (2)
Fricas	93.46 (143)	6.54 (10)
Maple	87.58 (134)	12.42 (19)
Maxima	56.21 (86)	43.79 (67)
Reduce	47.06 (72)	52.94 (81)
Giac	40.52 (62)	59.48 (91)
Mupad	35.95 (55)	64.05 (98)
Sympy	33.33 (51)	66.67 (102)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	88.235	0.000	0.654	11.111
Mathematica	69.935	5.229	12.418	12.418
Fricas	51.634	30.719	0.000	17.647
Maxima	36.601	12.418	0.000	50.980
Maple	29.412	20.261	26.797	23.529
Giac	24.837	4.575	0.000	70.588
Sympy	18.954	6.536	4.575	69.935
Mupad	0.000	24.837	0.000	75.163
Reduce	0.000	35.948	0.000	64.052

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	2	100.00	0.00	0.00
Fricas	10	100.00	0.00	0.00
Maple	19	100.00	0.00	0.00
Maxima	67	79.10	2.99	17.91
Reduce	81	100.00	0.00	0.00
Sympy	102	46.08	29.41	24.51
Giac	91	95.60	4.40	0.00
Mupad	98	0.00	100.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.12
Reduce	0.20
Maxima	0.39
Rubi	0.47
Mupad	0.66
Mathematica	0.95
Giac	2.89
Maple	4.00
Sympy	8.22

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	44.58	1.11	25.00	1.05
Reduce	64.79	1.21	48.00	1.13
Giac	71.89	1.29	37.50	0.95
Sympy	91.73	1.67	61.00	1.01
Maxima	111.91	1.50	70.00	1.00
Rubi	124.65	1.07	89.00	1.00
Mathematica	135.46	1.29	79.00	0.95
Fricas	240.76	1.71	83.00	1.29
Maple	926.89	5.40	163.00	1.84

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{31, 35, 36, 51, 55, 59, 64, 68, 72, 80, 84, 88, 92, 97, 101, 105, 109}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {1, 2}

Mathematica {50, 54, 58, 63, 67, 71, 81, 82, 83, 98, 99}

Maple {48, 49, 52, 53, 56, 57, 61, 62, 65, 66, 69, 70, 74, 75, 76, 77, 78, 81, 82, 85, 86, 89, 90, 93, 94, 95, 98, 99, 102, 103, 106, 107, 146, 147, 148, 149, 150, 151, 152, 153}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)), output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals.

These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```


For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	72

2.1 List of integrals sorted by grade for each CAS

Rubi	28
Mma	29
Maple	29
Fricas	30
Maxima	30
Giac	31
Mupad	31
Sympy	32
Reduce	32

Rubi

A grade { 1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 32, 33, 34, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 56, 57, 58, 60, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 76, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 96, 98, 99, 100, 102, 103, 104, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153 }

B grade { }

C grade { 6 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 30, 34, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 56, 57, 60, 61, 62, 65, 66, 69, 70, 73, 74, 75, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 94, 95, 96, 98, 99, 100, 102, 103, 104, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 152 }

B grade { 50, 54, 58, 63, 67, 71, 76, 93 }

C grade { 15, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 121, 148, 149, 150, 151, 153 }

F normal fail { 32, 33 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 8, 9, 10, 15, 16, 17, 30, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 60, 73, 79, 96, 111, 112, 116, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 132, 133, 134, 135, 136, 137, 138, 139 }

B grade { 3, 4, 5, 7, 11, 12, 13, 14, 32, 33, 34, 50, 54, 58, 63, 67, 71, 83, 87, 91, 100, 104, 108, 110, 113, 114, 115, 117, 118, 130, 131 }

C grade { 2, 48, 49, 52, 53, 56, 57, 61, 62, 65, 66, 69, 70, 74, 75, 76, 77, 78, 81, 82, 85, 86, 89, 90, 93, 94, 95, 98, 99, 102, 103, 106, 107, 146, 147, 148, 149, 150, 151, 152, 153 }

F normal fail { 6, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 140, 141, 142, 143, 144, 145 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 60, 65, 66, 67, 69, 70, 71, 74, 75, 111, 112, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 131, 132, 133, 137, 138, 139, 140, 143, 144, 145, 147, 149, 150 }

B grade { 48, 49, 50, 52, 53, 54, 56, 57, 58, 61, 62, 63, 73, 76, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 96, 98, 99, 100, 102, 103, 104, 106, 107, 108, 110, 113, 114, 134, 148, 151, 152, 153 }

C grade { }

F normal fail { 6, 32, 33, 34, 130, 135, 136, 141, 142, 146 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 7, 8, 9, 10, 11, 12, 13, 30, 37, 40, 41, 42, 43, 44, 45, 46, 47, 60, 85, 86, 87, 89, 90, 91, 102, 103, 104, 106, 107, 108, 116, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 138, 139, 147, 148, 149, 150, 151, 152 }

B grade { 14, 38, 39, 50, 52, 53, 54, 56, 57, 58, 63, 65, 66, 67, 69, 70, 71, 110, 113 }

C grade { }

F normal fail { 3, 4, 5, 6, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 33, 34, 48, 49, 61, 62, 73, 74, 75, 76, 77, 78, 79, 81, 82, 83, 93, 94, 95, 96, 98, 99, 100, 111, 112, 114, 115, 117, 118, 135, 136, 137, 153 }

F(-1) timedout fail { 51, 64 }

F(-2) exception fail { 55, 59, 68, 72, 134, 140, 141, 142, 143, 144, 145, 146 }

Giac

A grade { 1, 2, 3, 4, 5, 11, 12, 13, 14, 15, 16, 17, 30, 40, 42, 43, 44, 45, 46, 47, 60, 119, 120, 122, 123, 124, 126, 127, 128, 129, 131, 132, 133, 147, 148, 149, 151, 152 }

B grade { 7, 8, 9, 10, 121, 125, 150 }

C grade { }

F normal fail { 6, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 33, 34, 37, 38, 39, 41, 48, 49, 50, 52, 53, 54, 56, 57, 58, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 95, 96, 98, 99, 100, 102, 103, 104, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 130, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 153 }

F(-1) timedout fail { 76, 77, 93, 94 }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 14, 30, 38, 39, 40, 43, 44, 45, 47, 60, 110, 113, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 131, 132, 133, 134, 137, 138, 139, 147, 148, 149, 150, 151, 152 }

C grade { }

F normal fail { }

F(-1) timedout fail { 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 33, 34, 37, 41, 42, 46, 48, 49, 50, 52, 53, 54, 56, 57, 58, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 76, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 96, 98, 99, 100, 102, 103, 104, 106, 107, 108, 111, 112, 114, 115, 116, 117, 118, 130, 135, 136, 140, 141, 142, 143, 144, 145, 146, 153 }

F(-2) exception fail { }

Sympy

A grade { 1, 3, 4, 5, 7, 8, 11, 12, 13, 14, 15, 16, 17, 30, 38, 39, 41, 43, 44, 45, 46, 60, 119, 122, 123, 124, 127, 128, 129 }

B grade { 9, 10, 37, 40, 42, 47, 120, 121, 125, 131 }

C grade { 19, 20, 21, 22, 26, 27, 28 }

F normal fail { 6, 32, 33, 34, 49, 50, 73, 74, 75, 76, 77, 78, 79, 82, 83, 93, 94, 95, 96, 100, 110, 111, 112, 113, 114, 115, 116, 117, 118, 130, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 148, 149, 150, 152 }

F(-1) timeout fail { 18, 23, 24, 25, 29, 48, 55, 59, 61, 62, 63, 64, 68, 72, 81, 84, 88, 92, 97, 98, 99, 101, 105, 109, 126, 132, 133, 147, 151, 153 }

F(-2) exception fail { 2, 52, 53, 54, 56, 57, 58, 65, 66, 67, 69, 70, 71, 85, 86, 87, 89, 90, 91, 102, 103, 104, 106, 107, 108 }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 30, 37, 40, 42, 45, 47, 60, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 147, 148, 149, 150, 151, 152 }

C grade { }

F normal fail { 6, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 32, 33, 34, 38, 39, 41, 43, 44, 46, 48, 49, 50, 52, 53, 54, 56, 57, 58, 61, 62, 63, 65, 66, 67, 69, 70, 71, 73, 74, 75, 76, 77, 78, 79, 81, 82, 83, 85, 86, 87, 89, 90, 91, 93, 94, 95, 96, 98, 99, 100, 102, 103, 104, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 130, 146, 153 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	34	37	37	37	45	60	37	223	230
N.S.	1	0.81	0.88	0.88	0.88	1.07	1.43	0.88	5.31	5.48
time (sec)	N/A	0.268	0.012	0.566	0.029	0.108	0.634	0.128	0.215	0.786

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	F(-2)	A	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	36	40	140	40	58	0	40	58	58
N.S.	1	0.80	0.89	3.11	0.89	1.29	0.00	0.89	1.29	1.29
time (sec)	N/A	0.285	0.025	7.032	0.028	0.123	0.000	0.130	0.180	0.649

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	144	149	86	260	0	76	138	94	159	0
N.S.	1	1.03	0.60	1.81	0.00	0.53	0.96	0.65	1.10	0.00
time (sec)	N/A	0.306	0.051	0.514	0.000	0.101	0.988	0.219	0.192	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	119	74	212	0	65	107	80	137	0
N.S.	1	1.03	0.64	1.83	0.00	0.56	0.92	0.69	1.18	0.00
time (sec)	N/A	0.262	0.036	0.023	0.000	0.103	0.432	0.218	0.191	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	89	59	164	0	49	73	63	138	0
N.S.	1	1.01	0.67	1.86	0.00	0.56	0.83	0.72	1.57	0.00
time (sec)	N/A	0.242	0.028	0.023	0.000	0.104	0.150	0.189	0.183	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	288	189	171	0	0	0	0	0	22	0
N.S.	1	0.66	0.59	0.00	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	0.754	1.651	0.000	0.000	0.000	0.000	0.000	0.189	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	54	122	58	44	53	104	69	0
N.S.	1	1.00	0.95	2.14	1.02	0.77	0.93	1.82	1.21	0.00
time (sec)	N/A	0.223	0.025	0.026	0.092	0.119	1.855	0.230	0.197	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	84	67	69	68	58	83	198	98	0
N.S.	1	0.99	0.79	0.81	0.80	0.68	0.98	2.33	1.15	0.00
time (sec)	N/A	0.232	0.032	0.030	0.050	0.114	2.273	0.238	0.180	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	114	78	117	109	68	352	275	120	0
N.S.	1	1.01	0.69	1.04	0.96	0.60	3.12	2.43	1.06	0.00
time (sec)	N/A	0.260	0.040	0.028	0.056	0.125	2.905	0.244	0.188	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	141	144	89	165	132	80	575	351	142	0
N.S.	1	1.02	0.63	1.17	0.94	0.57	4.08	2.49	1.01	0.00
time (sec)	N/A	0.284	0.044	0.030	0.042	0.146	3.912	0.245	0.191	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	116	83	231	167	79	136	137	132	0
N.S.	1	0.94	0.67	1.86	1.35	0.64	1.10	1.10	1.06	0.00
time (sec)	N/A	0.284	0.067	0.030	0.041	0.157	1.645	0.199	0.199	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	97	72	183	139	68	105	101	110	0
N.S.	1	0.98	0.73	1.85	1.40	0.69	1.06	1.02	1.11	0.00
time (sec)	N/A	0.260	0.058	0.024	0.048	0.157	0.705	0.205	0.190	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	76	60	135	111	56	75	64	88	0
N.S.	1	1.03	0.81	1.82	1.50	0.76	1.01	0.86	1.19	0.00
time (sec)	N/A	0.243	0.052	0.023	0.053	0.169	0.335	0.191	0.187	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	47	43	86	77	41	41	40	64	35
N.S.	1	1.09	1.00	2.00	1.79	0.95	0.95	0.93	1.49	0.81
time (sec)	N/A	0.203	0.012	0.017	0.048	0.095	0.135	0.161	0.181	0.637

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	86	90	0	148	66	53	113	0
N.S.	1	1.00	1.46	1.53	0.00	2.51	1.12	0.90	1.92	0.00
time (sec)	N/A	0.229	0.040	0.030	0.000	0.121	2.075	0.200	0.192	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	85	101	130	0	198	82	89	137	0
N.S.	1	0.93	1.11	1.43	0.00	2.18	0.90	0.98	1.51	0.00
time (sec)	N/A	0.240	0.060	0.027	0.000	0.137	4.028	0.198	0.199	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	119	115	114	178	0	228	148	116	162	0
N.S.	1	0.97	0.96	1.50	0.00	1.92	1.24	0.97	1.36	0.00
time (sec)	N/A	0.251	0.075	0.029	0.000	0.143	6.659	0.192	0.211	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	211	218	170	0	0	96	0	0	24	0
N.S.	1	1.03	0.81	0.00	0.00	0.45	0.00	0.00	0.11	0.00
time (sec)	N/A	0.321	11.391	0.000	0.000	0.146	0.000	0.000	0.234	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	181	186	158	0	0	85	75	0	24	0
N.S.	1	1.03	0.87	0.00	0.00	0.47	0.41	0.00	0.13	0.00
time (sec)	N/A	0.306	0.296	0.000	0.000	0.135	41.386	0.000	0.211	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	153	154	147	0	0	71	75	0	21	0
N.S.	1	1.01	0.96	0.00	0.00	0.46	0.49	0.00	0.14	0.00
time (sec)	N/A	0.278	0.202	0.000	0.000	0.129	9.094	0.000	0.192	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	115	0	0	52	71	0	24	0
N.S.	1	1.00	0.94	0.00	0.00	0.43	0.58	0.00	0.20	0.00
time (sec)	N/A	0.252	0.096	0.000	0.000	0.110	8.789	0.000	0.190	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	156	154	150	0	0	73	78	0	24	0
N.S.	1	0.99	0.96	0.00	0.00	0.47	0.50	0.00	0.15	0.00
time (sec)	N/A	0.279	0.191	0.000	0.000	0.121	43.998	0.000	0.212	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	186	186	162	0	0	86	0	0	24	0
N.S.	1	1.00	0.87	0.00	0.00	0.46	0.00	0.00	0.13	0.00
time (sec)	N/A	0.309	0.222	0.000	0.000	0.109	0.000	0.000	0.207	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	216	218	171	0	0	97	0	0	24	0
N.S.	1	1.01	0.79	0.00	0.00	0.45	0.00	0.00	0.11	0.00
time (sec)	N/A	0.328	0.402	0.000	0.000	0.122	0.000	0.000	0.231	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	326	321	139	0	0	94	0	0	24	0
N.S.	1	0.98	0.43	0.00	0.00	0.29	0.00	0.00	0.07	0.00
time (sec)	N/A	0.442	10.112	0.000	0.000	0.138	0.000	0.000	0.221	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	296	289	119	0	0	79	75	0	22	0
N.S.	1	0.98	0.40	0.00	0.00	0.27	0.25	0.00	0.07	0.00
time (sec)	N/A	0.391	0.076	0.000	0.000	0.127	10.341	0.000	0.195	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	260	253	89	0	0	55	71	0	24	0
N.S.	1	0.97	0.34	0.00	0.00	0.21	0.27	0.00	0.09	0.00
time (sec)	N/A	0.350	0.072	0.000	0.000	0.118	9.590	0.000	0.190	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	298	285	121	0	0	80	78	0	24	0
N.S.	1	0.96	0.41	0.00	0.00	0.27	0.26	0.00	0.08	0.00
time (sec)	N/A	0.399	0.091	0.000	0.000	0.122	10.401	0.000	0.201	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	331	317	137	0	0	94	0	0	24	0
N.S.	1	0.96	0.41	0.00	0.00	0.28	0.00	0.00	0.07	0.00
time (sec)	N/A	0.426	0.082	0.000	0.000	0.110	0.000	0.000	0.221	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	50	43	42	47	41	43	55	42
N.S.	1	1.00	1.00	0.86	0.84	0.94	0.82	0.86	1.10	0.84
time (sec)	N/A	0.290	0.013	1.006	0.117	0.093	0.288	0.127	0.182	0.735

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	42	36	39	38	34	38	39	39
N.S.	1	1.00	1.05	0.90	0.98	0.95	0.85	0.95	0.98	0.98
time (sec)	N/A	0.247	0.130	1.234	0.872	0.125	4.157	0.347	0.621	0.850

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	431	429	0	1664	0	0	0	0	150	0
N.S.	1	1.00	0.00	3.86	0.00	0.00	0.00	0.00	0.35	0.00
time (sec)	N/A	1.105	0.000	1.542	0.000	0.000	0.000	0.000	0.209	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	283	285	0	916	0	0	0	0	108	0
N.S.	1	1.01	0.00	3.24	0.00	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	0.802	0.000	0.121	0.000	0.000	0.000	0.000	0.201	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	93	93	368	0	0	0	0	64	0
N.S.	1	0.95	0.95	3.76	0.00	0.00	0.00	0.00	0.65	0.00
time (sec)	N/A	0.324	0.028	0.109	0.000	0.000	0.000	0.000	0.190	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	42	36	39	48	61	38	63	39
N.S.	1	1.00	1.05	0.90	0.98	1.20	1.52	0.95	1.58	0.98
time (sec)	N/A	0.241	0.196	0.651	0.338	0.089	3.309	0.204	0.190	0.596

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	42	36	163	91	126	38	123	39
N.S.	1	1.00	1.05	0.90	4.08	2.28	3.15	0.95	3.08	0.98
time (sec)	N/A	0.249	0.893	0.612	0.429	0.085	7.168	0.322	0.213	1.415

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	34	41	38	33	158	0	38	0
N.S.	1	1.00	0.92	1.11	1.03	0.89	4.27	0.00	1.03	0.00
time (sec)	N/A	0.187	0.045	0.440	0.049	0.093	0.811	0.000	0.182	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	20	81	13	32	0	13	19
N.S.	1	1.00	0.87	0.87	3.52	0.57	1.39	0.00	0.57	0.83
time (sec)	N/A	0.167	0.013	0.091	0.039	0.077	0.156	0.000	0.196	0.139

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	20	57	13	32	0	11	19
N.S.	1	1.00	0.87	0.87	2.48	0.57	1.39	0.00	0.48	0.83
time (sec)	N/A	0.177	0.011	0.041	0.030	0.078	0.097	0.000	0.188	0.071

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	15	12	10	42	26	14	16
N.S.	1	1.00	1.12	0.94	0.75	0.62	2.62	1.62	0.88	1.00
time (sec)	N/A	0.153	0.005	0.097	0.032	0.074	0.076	0.121	0.188	0.045

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	19	23	42	8	34	0	13	0
N.S.	1	1.00	0.90	1.10	2.00	0.38	1.62	0.00	0.62	0.00
time (sec)	N/A	0.190	0.011	0.071	0.113	0.102	0.388	0.000	0.174	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	31	56	40	42	160	62	47	0
N.S.	1	1.00	0.86	1.56	1.11	1.17	4.44	1.72	1.31	0.00
time (sec)	N/A	0.181	0.035	0.523	0.027	0.087	2.049	0.134	0.184	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	26	17	17	26	19	22	25
N.S.	1	1.00	0.87	1.13	0.74	0.74	1.13	0.83	0.96	1.09
time (sec)	N/A	0.163	0.013	0.278	0.051	0.088	0.130	0.129	0.192	0.502

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	20	26	17	17	26	19	20	25
N.S.	1	1.00	0.87	1.13	0.74	0.74	1.13	0.83	0.87	1.09
time (sec)	N/A	0.182	0.012	0.077	0.026	0.079	0.093	0.129	0.182	0.080

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	22	15	15	24	15	21	21
N.S.	1	1.00	1.12	1.38	0.94	0.94	1.50	0.94	1.31	1.31
time (sec)	N/A	0.159	0.006	0.091	0.044	0.070	0.103	0.127	0.186	0.450

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	27	14	14	27	15	21	0
N.S.	1	1.00	1.00	1.42	0.74	0.74	1.42	0.79	1.11	0.00
time (sec)	N/A	0.182	0.013	0.093	0.033	0.084	1.615	0.126	0.185	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	15	12	10	42	26	14	16
N.S.	1	1.00	1.12	0.94	0.75	0.62	2.62	1.62	0.88	1.00
time (sec)	N/A	0.158	0.000	0.000	0.028	0.092	0.089	0.129	0.180	0.002

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	403	519	371	8039	0	1965	0	0	17	0
N.S.	1	1.29	0.92	19.95	0.00	4.88	0.00	0.00	0.04	0.00
time (sec)	N/A	1.567	2.878	49.339	0.000	0.183	0.000	0.000	0.182	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	305	396	281	7647	0	1545	0	0	15	0
N.S.	1	1.30	0.92	25.07	0.00	5.07	0.00	0.00	0.05	0.00
time (sec)	N/A	1.148	2.576	3.917	0.000	0.164	0.000	0.000	0.198	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	277	549	1001	433	1101	0	0	13	0
N.S.	1	1.40	2.77	5.06	2.19	5.56	0.00	0.00	0.07	0.00
time (sec)	N/A	0.707	5.102	2.898	0.205	0.167	0.000	0.000	0.190	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	0	17	14	17	17	17
N.S.	1	1.00	1.13	1.00	0.00	1.13	0.93	1.13	1.13	1.13
time (sec)	N/A	0.312	1.858	0.441	0.000	0.091	71.748	0.776	0.200	0.577

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	154	198	140	1487	310	322	0	0	24	0
N.S.	1	1.29	0.91	9.66	2.01	2.09	0.00	0.00	0.16	0.00
time (sec)	N/A	0.837	0.259	4.920	0.058	0.127	0.000	0.000	0.188	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	123	155	110	1452	218	271	0	0	22	0
N.S.	1	1.26	0.89	11.80	1.77	2.20	0.00	0.00	0.18	0.00
time (sec)	N/A	0.660	0.156	1.335	0.059	0.103	0.000	0.000	0.184	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	107	967	563	448	202	0	0	20	0
N.S.	1	1.26	11.38	6.62	5.27	2.38	0.00	0.00	0.24	0.00
time (sec)	N/A	0.449	8.552	1.550	0.170	0.102	0.000	0.000	0.182	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	0	37	0	21	24	22
N.S.	1	1.00	1.10	0.90	0.00	1.76	0.00	1.00	1.14	1.05
time (sec)	N/A	0.312	0.443	0.552	0.000	0.090	0.000	0.692	0.194	0.703

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	155	199	137	1488	310	322	0	0	26	0
N.S.	1	1.28	0.88	9.60	2.00	2.08	0.00	0.00	0.17	0.00
time (sec)	N/A	0.804	0.269	5.005	0.070	0.110	0.000	0.000	0.189	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	124	156	111	1453	218	271	0	0	24	0
N.S.	1	1.26	0.90	11.72	1.76	2.19	0.00	0.00	0.19	0.00
time (sec)	N/A	0.623	0.176	1.498	0.055	0.098	0.000	0.000	0.184	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	108	847	594	448	200	0	0	22	0
N.S.	1	1.26	9.85	6.91	5.21	2.33	0.00	0.00	0.26	0.00
time (sec)	N/A	0.454	6.252	1.592	0.198	0.091	0.000	0.000	0.191	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	0	37	0	21	26	22
N.S.	1	1.00	1.10	0.90	0.00	1.76	0.00	1.00	1.24	1.05
time (sec)	N/A	0.320	0.734	0.451	0.000	0.098	0.000	0.857	0.190	0.854

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	22	15	15	24	15	21	21
N.S.	1	1.00	1.12	1.38	0.94	0.94	1.50	0.94	1.31	1.31
time (sec)	N/A	0.156	0.001	0.017	0.038	0.069	0.080	0.120	0.186	0.002

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	399	515	360	7869	0	1589	0	0	17	0
N.S.	1	1.29	0.90	19.72	0.00	3.98	0.00	0.00	0.04	0.00
time (sec)	N/A	1.555	2.655	60.347	0.000	0.271	0.000	0.000	0.218	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	303	394	275	7501	0	1289	0	0	15	0
N.S.	1	1.30	0.91	24.76	0.00	4.25	0.00	0.00	0.05	0.00
time (sec)	N/A	1.133	2.107	4.211	0.000	0.236	0.000	0.000	0.208	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	198	277	702	1145	532	965	0	0	13	0
N.S.	1	1.40	3.55	5.78	2.69	4.87	0.00	0.00	0.07	0.00
time (sec)	N/A	0.689	9.774	4.129	0.212	0.220	0.000	0.000	0.194	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-1)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	0	17	0	17	17	17
N.S.	1	1.00	1.13	1.00	0.00	1.13	0.00	1.13	1.13	1.13
time (sec)	N/A	0.298	1.692	0.493	0.000	0.092	0.000	2.654	0.194	0.631

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	154	198	136	1487	309	166	0	0	30	0
N.S.	1	1.29	0.88	9.66	2.01	1.08	0.00	0.00	0.19	0.00
time (sec)	N/A	0.811	0.240	5.201	0.083	0.094	0.000	0.000	0.224	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	123	155	110	1452	217	144	0	0	28	0
N.S.	1	1.26	0.89	11.80	1.76	1.17	0.00	0.00	0.23	0.00
time (sec)	N/A	0.675	0.160	1.483	0.065	0.099	0.000	0.000	0.231	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	F(-2)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	107	929	584	458	112	0	0	26	0
N.S.	1	1.26	10.93	6.87	5.39	1.32	0.00	0.00	0.31	0.00
time (sec)	N/A	0.463	9.465	1.727	0.145	0.093	0.000	0.000	0.225	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	23	0	37	0	25	30	23
N.S.	1	1.00	1.10	1.10	0.00	1.76	0.00	1.19	1.43	1.10
time (sec)	N/A	0.302	0.473	0.865	0.000	0.086	0.000	1.540	0.194	0.883

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	155	199	140	1488	311	166	0	0	28	0
N.S.	1	1.28	0.90	9.60	2.01	1.07	0.00	0.00	0.18	0.00
time (sec)	N/A	0.848	0.250	4.981	0.058	0.098	0.000	0.000	0.225	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	B	A	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	124	156	110	1453	219	144	0	0	26	0
N.S.	1	1.26	0.89	11.72	1.77	1.16	0.00	0.00	0.21	0.00
time (sec)	N/A	0.644	0.163	1.536	0.067	0.092	0.000	0.000	0.222	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	B	B	A	F(-2)	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	108	872	625	458	112	0	0	24	0
N.S.	1	1.26	10.14	7.27	5.33	1.30	0.00	0.00	0.28	0.00
time (sec)	N/A	0.448	5.750	1.597	0.155	0.083	0.000	0.000	0.216	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	23	23	0	37	0	25	28	23
N.S.	1	1.00	1.10	1.10	0.00	1.76	0.00	1.19	1.33	1.10
time (sec)	N/A	0.304	0.481	0.825	0.000	0.086	0.000	1.750	0.200	0.944

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	56	52	0	58	0	0	5	0
N.S.	1	1.00	1.44	1.33	0.00	1.49	0.00	0.00	0.13	0.00
time (sec)	N/A	0.302	0.038	2.021	0.000	0.116	0.000	0.000	0.196	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	74	77	93	632	0	93	0	0	7	0
N.S.	1	1.04	1.26	8.54	0.00	1.26	0.00	0.00	0.09	0.00
time (sec)	N/A	0.434	0.035	1.305	0.000	0.108	0.000	0.000	0.174	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	108	113	121	658	0	125	0	0	9	0
N.S.	1	1.05	1.12	6.09	0.00	1.16	0.00	0.00	0.08	0.00
time (sec)	N/A	0.576	0.064	1.911	0.000	0.095	0.000	0.000	0.183	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	F	B	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	299	327	600	3570	0	1460	0	0	67	0
N.S.	1	1.09	2.01	11.94	0.00	4.88	0.00	0.00	0.22	0.00
time (sec)	N/A	1.150	0.671	38.757	0.000	0.206	0.000	0.000	0.203	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	229	251	375	2668	0	1002	0	0	46	0
N.S.	1	1.10	1.64	11.65	0.00	4.38	0.00	0.00	0.20	0.00
time (sec)	N/A	0.869	0.349	26.380	0.000	0.164	0.000	0.000	0.194	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	159	175	237	1776	0	600	0	0	25	0
N.S.	1	1.10	1.49	11.17	0.00	3.77	0.00	0.00	0.16	0.00
time (sec)	N/A	0.588	0.194	2.996	0.000	0.134	0.000	0.000	0.190	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	80	91	107	0	334	0	0	9	0
N.S.	1	1.08	1.23	1.45	0.00	4.51	0.00	0.00	0.12	0.00
time (sec)	N/A	0.350	0.020	1.533	0.000	0.147	0.000	0.000	0.180	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	3	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	0.20	1.13	1.13
time (sec)	N/A	0.213	10.976	0.768	1.617	0.094	3.796	87.325	0.190	0.508

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	355	441	438	6917	0	1289	0	0	17	0
N.S.	1	1.24	1.23	19.48	0.00	3.63	0.00	0.00	0.05	0.00
time (sec)	N/A	1.448	0.853	50.624	0.000	0.180	0.000	0.000	0.191	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	267	343	330	6567	0	1067	0	0	15	0
N.S.	1	1.28	1.24	24.60	0.00	4.00	0.00	0.00	0.06	0.00
time (sec)	N/A	1.104	0.726	3.451	0.000	0.170	0.000	0.000	0.202	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	237	288	352	0	825	0	0	13	0
N.S.	1	1.36	1.66	2.02	0.00	4.74	0.00	0.00	0.07	0.00
time (sec)	N/A	0.664	2.217	1.431	0.000	0.250	0.000	0.000	0.199	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	0	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.00	1.13	1.13	1.13
time (sec)	N/A	0.295	5.830	0.135	1.049	0.091	0.000	0.354	0.188	0.610

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	142	167	134	1406	129	293	0	0	25	0
N.S.	1	1.18	0.94	9.90	0.91	2.06	0.00	0.00	0.18	0.00
time (sec)	N/A	0.902	0.145	1.135	1.372	0.137	0.000	0.000	0.191	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	113	132	103	1370	106	247	0	0	23	0
N.S.	1	1.17	0.91	12.12	0.94	2.19	0.00	0.00	0.20	0.00
time (sec)	N/A	0.668	0.074	0.542	1.463	0.126	0.000	0.000	0.204	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	92	71	545	80	187	0	0	21	0
N.S.	1	1.16	0.90	6.90	1.01	2.37	0.00	0.00	0.27	0.00
time (sec)	N/A	0.434	1.441	0.536	1.526	0.112	0.000	0.000	0.207	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	17	72	37	0	19	25	20
N.S.	1	1.00	1.11	0.89	3.79	1.95	0.00	1.00	1.32	1.05
time (sec)	N/A	0.294	2.933	0.189	0.583	0.111	0.000	0.173	0.197	0.943

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	145	170	134	1417	129	293	0	0	26	0
N.S.	1	1.17	0.92	9.77	0.89	2.02	0.00	0.00	0.18	0.00
time (sec)	N/A	0.797	0.151	1.291	1.423	0.109	0.000	0.000	0.205	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	116	136	103	1381	107	247	0	0	24	0
N.S.	1	1.17	0.89	11.91	0.92	2.13	0.00	0.00	0.21	0.00
time (sec)	N/A	0.646	0.084	0.680	1.362	0.105	0.000	0.000	0.201	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	96	71	516	80	187	0	0	22	0
N.S.	1	1.17	0.87	6.29	0.98	2.28	0.00	0.00	0.27	0.00
time (sec)	N/A	0.452	1.472	0.698	1.368	0.101	0.000	0.000	0.217	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	77	37	0	19	26	20
N.S.	1	1.00	1.09	0.91	3.50	1.68	0.00	0.86	1.18	0.91
time (sec)	N/A	0.292	2.955	0.260	0.513	0.077	0.000	0.174	0.205	0.943

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	C	F	B	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	299	327	600	3570	0	1460	0	0	67	0
N.S.	1	1.09	2.01	11.94	0.00	4.88	0.00	0.00	0.22	0.00
time (sec)	N/A	1.182	0.267	7.812	0.000	0.189	0.000	0.000	0.211	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	229	251	375	2668	0	1002	0	0	46	0
N.S.	1	1.10	1.64	11.65	0.00	4.38	0.00	0.00	0.20	0.00
time (sec)	N/A	0.856	0.162	4.629	0.000	0.154	0.000	0.000	0.234	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	159	175	237	1777	0	600	0	0	25	0
N.S.	1	1.10	1.49	11.18	0.00	3.77	0.00	0.00	0.16	0.00
time (sec)	N/A	0.589	0.105	0.605	0.000	0.159	0.000	0.000	0.199	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	79	91	106	0	334	0	0	9	0
N.S.	1	1.08	1.25	1.45	0.00	4.58	0.00	0.00	0.12	0.00
time (sec)	N/A	0.378	0.019	0.293	0.000	0.141	0.000	0.000	0.194	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	0	3	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.00	0.20	1.13	1.13
time (sec)	N/A	0.244	0.657	0.092	1.958	0.082	0.000	73.997	0.209	0.547

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	351	437	442	6845	0	1269	0	0	17	0
N.S.	1	1.25	1.26	19.50	0.00	3.62	0.00	0.00	0.05	0.00
time (sec)	N/A	1.506	0.872	16.229	0.000	0.188	0.000	0.000	0.198	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	265	339	334	6495	0	1051	0	0	15	0
N.S.	1	1.28	1.26	24.51	0.00	3.97	0.00	0.00	0.06	0.00
time (sec)	N/A	1.175	0.634	2.719	0.000	0.251	0.000	0.000	0.198	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	237	287	352	0	813	0	0	13	0
N.S.	1	1.36	1.65	2.02	0.00	4.67	0.00	0.00	0.07	0.00
time (sec)	N/A	0.689	1.919	1.568	0.000	0.221	0.000	0.000	0.189	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	0	17	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.00	1.13	1.13	1.13
time (sec)	N/A	0.288	6.085	0.140	0.994	0.091	0.000	0.371	0.196	0.638

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	142	167	134	1405	129	293	0	0	25	0
N.S.	1	1.18	0.94	9.89	0.91	2.06	0.00	0.00	0.18	0.00
time (sec)	N/A	0.810	0.144	1.115	1.721	0.103	0.000	0.000	0.205	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	113	133	103	1369	106	247	0	0	23	0
N.S.	1	1.18	0.91	12.12	0.94	2.19	0.00	0.00	0.20	0.00
time (sec)	N/A	0.622	0.069	0.580	1.367	0.095	0.000	0.000	0.195	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	92	71	545	80	187	0	0	21	0
N.S.	1	1.16	0.90	6.90	1.01	2.37	0.00	0.00	0.27	0.00
time (sec)	N/A	0.438	1.289	0.622	1.340	0.116	0.000	0.000	0.196	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	17	77	37	0	19	25	20
N.S.	1	1.00	1.11	0.89	4.05	1.95	0.00	1.00	1.32	1.05
time (sec)	N/A	0.289	3.079	0.233	0.555	0.090	0.000	0.183	0.200	0.973

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	145	170	134	1418	129	293	0	0	26	0
N.S.	1	1.17	0.92	9.78	0.89	2.02	0.00	0.00	0.18	0.00
time (sec)	N/A	0.806	0.150	1.280	1.513	0.120	0.000	0.000	0.199	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	116	135	103	1382	107	247	0	0	24	0
N.S.	1	1.16	0.89	11.91	0.92	2.13	0.00	0.00	0.21	0.00
time (sec)	N/A	0.651	0.083	0.803	1.632	0.135	0.000	0.000	0.212	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	B	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	96	71	516	80	187	0	0	22	0
N.S.	1	1.17	0.87	6.29	0.98	2.28	0.00	0.00	0.27	0.00
time (sec)	N/A	0.442	1.266	0.724	1.699	0.106	0.000	0.000	0.190	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	72	37	0	19	26	20
N.S.	1	1.00	1.09	0.91	3.27	1.68	0.00	0.86	1.18	0.91
time (sec)	N/A	0.283	3.102	0.299	0.535	0.091	0.000	0.186	0.195	0.938

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	59	53	34	40	0	0	6	21
N.S.	1	1.00	1.90	1.71	1.10	1.29	0.00	0.00	0.19	0.68
time (sec)	N/A	0.230	0.012	0.081	0.116	0.113	0.000	0.000	0.188	0.637

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	57	50	44	0	65	0	0	8	0
N.S.	1	0.90	0.79	0.70	0.00	1.03	0.00	0.00	0.13	0.00
time (sec)	N/A	0.336	0.008	0.083	0.000	0.106	0.000	0.000	0.193	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	93	80	70	0	87	0	0	10	0
N.S.	1	1.02	0.88	0.77	0.00	0.96	0.00	0.00	0.11	0.00
time (sec)	N/A	0.474	0.012	0.109	0.000	0.095	0.000	0.000	0.185	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	B	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	43	83	95	63	103	0	0	10	37
N.S.	1	0.96	1.84	2.11	1.40	2.29	0.00	0.00	0.22	0.82
time (sec)	N/A	0.243	0.146	0.086	0.135	0.101	0.000	0.000	0.186	0.688

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	87	71	349	0	151	0	0	12	0
N.S.	1	0.96	0.78	3.84	0.00	1.66	0.00	0.00	0.13	0.00
time (sec)	N/A	0.428	0.010	0.128	0.000	0.104	0.000	0.000	0.186	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	141	115	407	0	187	0	0	14	0
N.S.	1	1.06	0.86	3.06	0.00	1.41	0.00	0.00	0.11	0.00
time (sec)	N/A	0.586	0.015	0.178	0.000	0.106	0.000	0.000	0.182	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	196	177	167	162	189	212	0	0	14	0
N.S.	1	0.90	0.85	0.83	0.96	1.08	0.00	0.00	0.07	0.00
time (sec)	N/A	0.558	0.278	0.582	0.188	0.108	0.000	0.000	0.182	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	232	238	236	672	0	304	0	0	16	0
N.S.	1	1.03	1.02	2.90	0.00	1.31	0.00	0.00	0.07	0.00
time (sec)	N/A	0.753	0.060	0.685	0.000	0.118	0.000	0.000	0.183	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	302	321	299	758	0	378	0	0	18	0
N.S.	1	1.06	0.99	2.51	0.00	1.25	0.00	0.00	0.06	0.00
time (sec)	N/A	1.013	0.032	1.111	0.000	0.124	0.000	0.000	0.197	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	33	25	23	19	28	19	20	33	20
N.S.	1	1.32	1.00	0.92	0.76	1.12	0.76	0.80	1.32	0.80
time (sec)	N/A	0.264	0.012	0.045	0.034	0.102	1.259	0.129	0.187	0.099

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	53	35	32	31	50	153	32	71	31
N.S.	1	1.18	0.78	0.71	0.69	1.11	3.40	0.71	1.58	0.69
time (sec)	N/A	0.239	0.024	0.112	0.125	0.084	0.300	0.122	0.187	0.130

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	74	81	54	54	77	223	657	119	46
N.S.	1	1.16	1.27	0.84	0.84	1.20	3.48	10.27	1.86	0.72
time (sec)	N/A	0.315	0.031	0.128	0.113	0.100	0.319	0.237	0.185	0.664

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	22	25	24	18	26	24	21	24
N.S.	1	1.00	0.73	0.83	0.80	0.60	0.87	0.80	0.70	0.80
time (sec)	N/A	0.194	0.017	0.050	0.114	0.091	0.106	0.121	0.187	0.086

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	5	5	5	5	5	5
N.S.	1	1.00	1.00	1.20	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.188	0.042	0.031	0.036	0.087	0.115	0.126	0.190	0.250

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	16	15	14	16	15	15
N.S.	1	1.00	1.00	0.82	0.94	0.88	0.82	0.94	0.88	0.88
time (sec)	N/A	0.209	0.094	0.056	0.046	0.078	0.247	0.123	0.192	0.142

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	16	26	53	104	35	16
N.S.	1	1.00	1.00	0.94	0.89	1.44	2.94	5.78	1.94	0.89
time (sec)	N/A	0.343	0.021	0.214	0.231	0.097	0.306	0.130	0.199	0.137

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	74	58	45	44	40	0	44	51	72
N.S.	1	1.09	0.85	0.66	0.65	0.59	0.00	0.65	0.75	1.06
time (sec)	N/A	0.265	0.040	0.129	0.157	0.092	0.000	0.133	0.217	1.722

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	65	53	40	39	35	46	39	44	65
N.S.	1	1.10	0.90	0.68	0.66	0.59	0.78	0.66	0.75	1.10
time (sec)	N/A	0.253	0.022	0.031	0.159	0.103	85.402	0.129	0.201	0.942

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	56	48	35	34	28	39	34	37	58
N.S.	1	1.12	0.96	0.70	0.68	0.56	0.78	0.68	0.74	1.16
time (sec)	N/A	0.246	0.022	0.029	0.161	0.089	21.453	0.131	0.196	0.845

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	41	31	28	26	22	29	27	27	40
N.S.	1	1.11	0.84	0.76	0.70	0.59	0.78	0.73	0.73	1.08
time (sec)	N/A	0.222	0.070	0.015	0.202	0.098	7.747	0.127	0.200	0.921

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	84	374	43	0	0	0	16	0
N.S.	1	1.00	2.00	8.90	1.02	0.00	0.00	0.00	0.38	0.00
time (sec)	N/A	0.315	0.127	0.257	0.128	0.000	0.000	0.000	0.211	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	44	40	57	29	28	537	28	33	44
N.S.	1	1.07	0.98	1.39	0.71	0.68	13.10	0.68	0.80	1.07
time (sec)	N/A	0.237	0.021	0.035	0.153	0.089	43.712	0.132	0.199	1.397

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	56	48	35	34	35	0	34	38	49
N.S.	1	1.12	0.96	0.70	0.68	0.70	0.00	0.68	0.76	0.98
time (sec)	N/A	0.246	0.021	0.040	0.171	0.100	0.000	0.139	0.201	1.280

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	65	51	40	39	40	0	39	49	56
N.S.	1	1.10	0.86	0.68	0.66	0.68	0.00	0.66	0.83	0.95
time (sec)	N/A	0.255	0.028	0.042	0.149	0.099	0.000	0.128	0.210	0.869

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	59	0	126	0	0	52	57
N.S.	1	1.00	1.00	0.94	0.00	2.00	0.00	0.00	0.83	0.90
time (sec)	N/A	0.377	0.041	0.471	0.000	0.132	0.000	0.000	0.192	0.695

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	59	57	0	0	0	0	31	0
N.S.	1	1.00	1.00	0.97	0.00	0.00	0.00	0.00	0.53	0.00
time (sec)	N/A	0.365	0.023	0.088	0.000	0.000	0.000	0.000	0.188	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	59	57	0	0	0	0	31	0
N.S.	1	1.00	1.00	0.97	0.00	0.00	0.00	0.00	0.53	0.00
time (sec)	N/A	0.319	0.016	0.082	0.000	0.000	0.000	0.000	0.181	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	55	55	0	83	0	0	29	49
N.S.	1	1.00	1.00	1.00	0.00	1.51	0.00	0.00	0.53	0.89
time (sec)	N/A	0.366	0.116	0.254	0.000	0.077	0.000	0.000	0.195	0.556

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	57	57	29	82	0	0	31	51
N.S.	1	1.00	1.00	1.00	0.51	1.44	0.00	0.00	0.54	0.89
time (sec)	N/A	0.347	0.018	0.085	0.246	0.087	0.000	0.000	0.194	0.544

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	59	57	29	82	0	0	31	51
N.S.	1	1.00	1.00	0.97	0.49	1.39	0.00	0.00	0.53	0.86
time (sec)	N/A	0.355	0.018	0.086	0.331	0.086	0.000	0.000	0.191	0.534

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	66	0	0	132	0	0	104	0
N.S.	1	1.00	0.92	0.00	0.00	1.83	0.00	0.00	1.44	0.00
time (sec)	N/A	0.420	0.101	0.000	0.000	0.112	0.000	0.000	0.188	0.000

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	62	0	0	0	0	0	53	0
N.S.	1	1.00	0.91	0.00	0.00	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	0.398	0.059	0.000	0.000	0.000	0.000	0.000	0.192	0.000

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	62	0	0	0	0	0	53	0
N.S.	1	1.00	0.91	0.00	0.00	0.00	0.00	0.00	0.78	0.00
time (sec)	N/A	0.358	0.028	0.000	0.000	0.000	0.000	0.000	0.185	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	58	0	0	83	0	0	53	0
N.S.	1	1.00	0.91	0.00	0.00	1.30	0.00	0.00	0.83	0.00
time (sec)	N/A	0.387	0.048	0.000	0.000	0.099	0.000	0.000	0.183	0.000

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	60	0	0	82	0	0	52	0
N.S.	1	1.00	0.91	0.00	0.00	1.24	0.00	0.00	0.79	0.00
time (sec)	N/A	0.365	0.046	0.000	0.000	0.092	0.000	0.000	0.200	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F(-2)	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	62	0	0	83	0	0	53	0
N.S.	1	1.00	0.91	0.00	0.00	1.22	0.00	0.00	0.78	0.00
time (sec)	N/A	0.396	0.040	0.000	0.000	0.090	0.000	0.000	0.201	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F(-2)	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	101	89	79	1087	0	0	0	0	28	0
N.S.	1	0.88	0.78	10.76	0.00	0.00	0.00	0.00	0.28	0.00
time (sec)	N/A	0.679	0.088	6.987	0.000	0.000	0.000	0.000	0.194	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	F(-1)	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	48	46	61	1299	48	75	0	66	65	66
N.S.	1	0.96	1.27	27.06	1.00	1.56	0.00	1.38	1.35	1.38
time (sec)	N/A	0.351	0.055	0.893	0.133	0.096	0.000	0.125	0.198	0.671

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	F	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	103	104	146	1371	131	221	0	155	178	133
N.S.	1	1.01	1.42	13.31	1.27	2.15	0.00	1.50	1.73	1.29
time (sec)	N/A	0.534	0.075	4.483	0.145	0.105	0.000	0.133	0.247	0.348

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	F	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	134	175	89	1355	167	210	0	236	169	164
N.S.	1	1.31	0.66	10.11	1.25	1.57	0.00	1.76	1.26	1.22
time (sec)	N/A	0.511	0.057	0.909	0.143	0.100	0.000	0.129	0.202	2.028

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	A	F	B	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	135	175	89	1355	167	210	0	239	169	164
N.S.	1	1.30	0.66	10.04	1.24	1.56	0.00	1.77	1.25	1.21
time (sec)	N/A	0.519	0.057	0.835	0.175	0.126	0.000	0.127	0.199	1.134

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	A	B	F(-1)	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	103	104	145	838	169	276	0	155	173	135
N.S.	1	1.01	1.41	8.14	1.64	2.68	0.00	1.50	1.68	1.31
time (sec)	N/A	0.469	0.076	4.096	0.143	0.117	0.000	0.137	0.256	0.808

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	F	A	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	47	44	57	885	47	131	0	65	63	67
N.S.	1	0.94	1.21	18.83	1.00	2.79	0.00	1.38	1.34	1.43
time (sec)	N/A	0.331	0.050	0.540	0.135	0.102	0.000	0.126	0.205	0.677

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	116	547	0	250	0	0	64	0
N.S.	1	1.00	0.71	3.36	0.00	1.53	0.00	0.00	0.39	0.00
time (sec)	N/A	0.810	0.182	159.477	0.000	0.133	0.000	0.000	0.192	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [73] had the largest ratio of [1.66667000000000010]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	5	4	0.81	12	0.333
2	A	5	4	0.80	14	0.286
3	A	7	6	1.03	25	0.240
4	A	6	5	1.03	25	0.200
5	A	5	4	1.01	23	0.174
6	C	12	11	0.66	25	0.440
7	A	2	2	1.00	25	0.080
8	A	3	3	0.99	25	0.120
9	A	4	4	1.01	25	0.160
10	A	5	5	1.02	25	0.200
11	A	5	4	0.94	25	0.160
12	A	5	4	0.98	25	0.160
13	A	5	4	1.03	25	0.160
14	A	2	2	1.09	21	0.095
15	A	5	4	1.00	25	0.160
16	A	6	5	0.93	25	0.200
17	A	7	6	0.97	25	0.240
18	A	7	6	1.03	27	0.222
19	A	6	5	1.03	27	0.185
20	A	5	4	1.01	27	0.148
21	A	4	3	1.00	27	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	5	4	0.99	27	0.148
23	A	6	5	1.00	27	0.185
24	A	7	6	1.01	27	0.222
25	A	9	8	0.98	27	0.296
26	A	8	7	0.98	27	0.259
27	A	7	6	0.97	27	0.222
28	A	8	7	0.96	27	0.259
29	A	9	8	0.96	27	0.296
30	A	3	3	1.00	11	0.273
31	N/A	1	0	1.00	40	0.000
32	A	7	6	1.00	40	0.150
33	A	6	5	1.01	40	0.125
34	A	4	3	0.95	38	0.079
35	N/A	1	0	1.00	40	0.000
36	N/A	1	0	1.00	40	0.000
37	A	2	2	1.00	11	0.182
38	A	2	2	1.00	11	0.182
39	A	2	2	1.00	9	0.222
40	A	3	2	1.00	7	0.286
41	A	2	2	1.00	11	0.182
42	A	2	2	1.00	11	0.182
43	A	2	2	1.00	11	0.182
44	A	2	2	1.00	9	0.222
45	A	3	2	1.00	7	0.286
46	A	2	2	1.00	11	0.182
47	A	3	2	1.00	7	0.286
48	A	7	6	1.29	15	0.400
49	A	6	5	1.30	13	0.385
50	A	5	4	1.40	11	0.364
51	N/A	1	0	1.00	15	0.000
52	A	8	7	1.29	21	0.333
53	A	7	6	1.26	19	0.316

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	6	5	1.26	17	0.294
55	N/A	1	0	1.00	21	0.000
56	A	9	8	1.28	21	0.381
57	A	8	7	1.26	19	0.368
58	A	7	6	1.26	17	0.353
59	N/A	1	0	1.00	21	0.000
60	A	3	2	1.00	7	0.286
61	A	7	6	1.29	15	0.400
62	A	6	5	1.30	13	0.385
63	A	5	4	1.40	11	0.364
64	N/A	1	0	1.00	15	0.000
65	A	9	8	1.29	21	0.381
66	A	8	7	1.26	19	0.368
67	A	7	6	1.26	17	0.353
68	N/A	1	0	1.00	21	0.000
69	A	8	7	1.28	21	0.333
70	A	7	6	1.26	19	0.316
71	A	6	5	1.26	17	0.294
72	N/A	1	0	1.00	21	0.000
73	A	6	5	1.00	3	1.667
74	A	7	6	1.04	5	1.200
75	A	8	7	1.05	7	1.000
76	A	9	8	1.09	15	0.533
77	A	8	7	1.10	15	0.467
78	A	7	6	1.10	13	0.462
79	A	6	5	1.08	7	0.714
80	N/A	1	0	1.00	15	0.000
81	A	7	6	1.24	15	0.400
82	A	6	5	1.28	13	0.385
83	A	5	4	1.36	11	0.364
84	N/A	1	0	1.00	15	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
85	A	9	8	1.18	19	0.421
86	A	8	7	1.17	17	0.412
87	A	7	6	1.16	15	0.400
88	N/A	1	0	1.00	19	0.000
89	A	8	7	1.17	22	0.318
90	A	7	6	1.17	20	0.300
91	A	6	5	1.17	18	0.278
92	N/A	1	0	1.00	22	0.000
93	A	9	8	1.09	15	0.533
94	A	8	7	1.10	15	0.467
95	A	7	6	1.10	13	0.462
96	A	6	5	1.08	7	0.714
97	N/A	1	0	1.00	15	0.000
98	A	7	6	1.25	15	0.400
99	A	6	5	1.28	13	0.385
100	A	5	4	1.36	11	0.364
101	N/A	1	0	1.00	15	0.000
102	A	9	8	1.18	19	0.421
103	A	8	7	1.18	17	0.412
104	A	7	6	1.16	15	0.400
105	N/A	1	0	1.00	19	0.000
106	A	8	7	1.17	22	0.318
107	A	7	6	1.16	20	0.300
108	A	6	5	1.17	18	0.278
109	N/A	1	0	1.00	22	0.000
110	A	4	3	1.00	4	0.750
111	A	5	4	0.90	6	0.667
112	A	6	5	1.02	8	0.625
113	A	4	3	0.96	8	0.375
114	A	5	4	0.96	10	0.400
115	A	6	5	1.06	12	0.417
116	A	9	8	0.90	12	0.667

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
117	A	6	5	1.03	14	0.357
118	A	7	6	1.06	16	0.375
119	A	7	6	1.32	10	0.600
120	A	4	4	1.18	8	0.500
121	A	6	6	1.16	14	0.429
122	A	6	5	1.00	8	0.625
123	A	1	1	1.00	14	0.071
124	A	1	1	1.00	19	0.053
125	A	2	2	1.00	26	0.077
126	A	11	10	1.09	21	0.476
127	A	10	9	1.10	21	0.429
128	A	9	8	1.12	19	0.421
129	A	8	7	1.11	18	0.389
130	A	7	6	1.00	21	0.286
131	A	8	7	1.07	21	0.333
132	A	9	8	1.12	21	0.381
133	A	10	9	1.10	21	0.429
134	A	2	2	1.00	39	0.051
135	A	2	2	1.00	39	0.051
136	A	2	2	1.00	37	0.054
137	A	2	2	1.00	39	0.051
138	A	2	2	1.00	39	0.051
139	A	2	2	1.00	39	0.051
140	A	2	2	1.00	40	0.050
141	A	2	2	1.00	40	0.050
142	A	2	2	1.00	38	0.053
143	A	2	2	1.00	40	0.050
144	A	2	2	1.00	40	0.050
145	A	2	2	1.00	40	0.050
146	A	6	5	0.88	24	0.208
147	A	6	5	0.96	20	0.250
148	A	8	7	1.01	20	0.350

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
149	A	13	12	1.31	20	0.600
150	A	13	12	1.30	20	0.600
151	A	9	8	1.01	20	0.400
152	A	7	6	0.94	20	0.300
153	A	2	2	1.00	24	0.083

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \arctan(a + bx^4) dx$	84
3.2	$\int x^{-1+n} \arctan(a + bx^n) dx$	90
3.3	$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	96
3.4	$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	104
3.5	$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	111
3.6	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx$	117
3.7	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$	126
3.8	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$	131
3.9	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$	137
3.10	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$	144
3.11	$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	152
3.12	$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	159
3.13	$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	165
3.14	$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	171
3.15	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$	176
3.16	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$	182
3.17	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$	189

3.18	$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	196
3.19	$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	203
3.20	$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	210
3.21	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$	216
3.22	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$	222
3.23	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$	228
3.24	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$	235
3.25	$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	243
3.26	$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$	251
3.27	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$	259
3.28	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$	266
3.29	$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx$	274
3.30	$\int \frac{\arctan(1+x+x^2)}{x^2} dx$	283
3.31	$\int \frac{(a+b \arctan(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^n}{1-c^2x^2} dx$	289
3.32	$\int \frac{(a+b \arctan(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^3}{1-c^2x^2} dx$	294
3.33	$\int \frac{(a+b \arctan(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^2}{1-c^2x^2} dx$	303
3.34	$\int \frac{a+b \arctan(\frac{\sqrt{1-cx}}{\sqrt{1+cx}})}{1-c^2x^2} dx$	312
3.35	$\int \frac{1}{(1-c^2x^2)(a+b \arctan(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))} dx$	318
3.36	$\int \frac{1}{(1-c^2x^2)(a+b \arctan(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}))^2} dx$	323
3.37	$\int x^m \arctan(\tan(a+bx)) dx$	329
3.38	$\int x^2 \arctan(\tan(a+bx)) dx$	334
3.39	$\int x \arctan(\tan(a+bx)) dx$	339
3.40	$\int \arctan(\tan(a+bx)) dx$	344
3.41	$\int \frac{\arctan(\tan(a+bx))}{x} dx$	349
3.42	$\int x^m \arctan(\cot(a+bx)) dx$	354
3.43	$\int x^2 \arctan(\cot(a+bx)) dx$	360
3.44	$\int x \arctan(\cot(a+bx)) dx$	365
3.45	$\int \arctan(\cot(a+bx)) dx$	370

3.46	$\int \frac{\arctan(\cot(a+bx))}{x} dx$	375
3.47	$\int \arctan(\tan(a+bx)) dx$	380
3.48	$\int x^2 \arctan(c+d \tan(a+bx)) dx$	385
3.49	$\int x \arctan(c+d \tan(a+bx)) dx$	397
3.50	$\int \arctan(c+d \tan(a+bx)) dx$	406
3.51	$\int \frac{\arctan(c+d \tan(a+bx))}{x} dx$	414
3.52	$\int x^2 \arctan(c+(1+ic) \tan(a+bx)) dx$	419
3.53	$\int x \arctan(c+(1+ic) \tan(a+bx)) dx$	427
3.54	$\int \arctan(c+(1+ic) \tan(a+bx)) dx$	435
3.55	$\int \frac{\arctan(c+(1+ic) \tan(a+bx))}{x} dx$	442
3.56	$\int x^2 \arctan(c+(-1+ic) \tan(a+bx)) dx$	447
3.57	$\int x \arctan(c+(-1+ic) \tan(a+bx)) dx$	456
3.58	$\int \arctan(c+(-1+ic) \tan(a+bx)) dx$	464
3.59	$\int \frac{\arctan(c+(-1+ic) \tan(a+bx))}{x} dx$	472
3.60	$\int \arctan(\cot(a+bx)) dx$	477
3.61	$\int x^2 \arctan(c+d \cot(a+bx)) dx$	482
3.62	$\int x \arctan(c+d \cot(a+bx)) dx$	494
3.63	$\int \arctan(c+d \cot(a+bx)) dx$	503
3.64	$\int \frac{\arctan(c+d \cot(a+bx))}{x} dx$	512
3.65	$\int x^2 \arctan(c+(1-ic) \cot(a+bx)) dx$	517
3.66	$\int x \arctan(c+(1-ic) \cot(a+bx)) dx$	526
3.67	$\int \arctan(c+(1-ic) \cot(a+bx)) dx$	534
3.68	$\int \frac{\arctan(c+(1-ic) \cot(a+bx))}{x} dx$	542
3.69	$\int x^2 \arctan(c+(-1-ic) \cot(a+bx)) dx$	547
3.70	$\int x \arctan(c+(-1-ic) \cot(a+bx)) dx$	555
3.71	$\int \arctan(c+(-1-ic) \cot(a+bx)) dx$	563
3.72	$\int \frac{\arctan(c+(-1-ic) \cot(a+bx))}{x} dx$	570
3.73	$\int \arctan(\sinh(x)) dx$	575
3.74	$\int x \arctan(\sinh(x)) dx$	581
3.75	$\int x^2 \arctan(\sinh(x)) dx$	588
3.76	$\int (e+fx)^3 \arctan(\tanh(a+bx)) dx$	595
3.77	$\int (e+fx)^2 \arctan(\tanh(a+bx)) dx$	606
3.78	$\int (e+fx) \arctan(\tanh(a+bx)) dx$	615
3.79	$\int \arctan(\tanh(a+bx)) dx$	623
3.80	$\int \frac{\arctan(\tanh(a+bx))}{e+fx} dx$	629
3.81	$\int x^2 \arctan(c+d \tanh(a+bx)) dx$	634
3.82	$\int x \arctan(c+d \tanh(a+bx)) dx$	644
3.83	$\int \arctan(c+d \tanh(a+bx)) dx$	653
3.84	$\int \frac{\arctan(c+d \tanh(a+bx))}{x} dx$	661

3.85	$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx$	666
3.86	$\int x \arctan(c + (i + c) \tanh(a + bx)) dx$	675
3.87	$\int \arctan(c + (i + c) \tanh(a + bx)) dx$	683
3.88	$\int \frac{\arctan(c+(i+c) \tanh(a+bx))}{x} dx$	690
3.89	$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx$	695
3.90	$\int x \arctan(c - (i - c) \tanh(a + bx)) dx$	703
3.91	$\int \arctan(c - (i - c) \tanh(a + bx)) dx$	711
3.92	$\int \frac{\arctan(c-(i-c) \tanh(a+bx))}{x} dx$	718
3.93	$\int (e + fx)^3 \arctan(\coth(a + bx)) dx$	723
3.94	$\int (e + fx)^2 \arctan(\coth(a + bx)) dx$	734
3.95	$\int (e + fx) \arctan(\coth(a + bx)) dx$	743
3.96	$\int \arctan(\coth(a + bx)) dx$	751
3.97	$\int \frac{\arctan(\coth(a+bx))}{e+fx} dx$	757
3.98	$\int x^2 \arctan(c + d \coth(a + bx)) dx$	762
3.99	$\int x \arctan(c + d \coth(a + bx)) dx$	772
3.100	$\int \arctan(c + d \coth(a + bx)) dx$	781
3.101	$\int \frac{\arctan(c+d \coth(a+bx))}{x} dx$	789
3.102	$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx$	794
3.103	$\int x \arctan(c + (i + c) \coth(a + bx)) dx$	803
3.104	$\int \arctan(c + (i + c) \coth(a + bx)) dx$	811
3.105	$\int \frac{\arctan(c+(i+c) \coth(a+bx))}{x} dx$	818
3.106	$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx$	823
3.107	$\int x \arctan(c - (i - c) \coth(a + bx)) dx$	832
3.108	$\int \arctan(c - (i - c) \coth(a + bx)) dx$	840
3.109	$\int \frac{\arctan(c-(i-c) \coth(a+bx))}{x} dx$	847
3.110	$\int \arctan(e^x) dx$	852
3.111	$\int x \arctan(e^x) dx$	857
3.112	$\int x^2 \arctan(e^x) dx$	862
3.113	$\int \arctan(e^{a+bx}) dx$	868
3.114	$\int x \arctan(e^{a+bx}) dx$	874
3.115	$\int x^2 \arctan(e^{a+bx}) dx$	880
3.116	$\int \arctan(a + bf^{c+dx}) dx$	887
3.117	$\int x \arctan(a + bf^{c+dx}) dx$	894
3.118	$\int x^2 \arctan(a + bf^{c+dx}) dx$	901
3.119	$\int e^{-x} \arctan(e^x) dx$	909
3.120	$\int \frac{\arctan(x)}{(-1+x)^3} dx$	915
3.121	$\int \frac{\arctan(1+2x)}{(4+3x)^3} dx$	921
3.122	$\int \arctan(\sqrt{1+x}) dx$	929

3.123	$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx$	935
3.124	$\int \frac{1}{(a+ax^2)(b-2b\arctan(x))} dx$	940
3.125	$\int \frac{x+x^3+(1+x)^2\arctan(x)}{(1+x)^2(1+x^2)} dx$	945
3.126	$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx$	950
3.127	$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx$	957
3.128	$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx$	964
3.129	$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx$	970
3.130	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} dx$	976
3.131	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^2} dx$	982
3.132	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^3} dx$	989
3.133	$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^4} dx$	995
3.134	$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$	1002
3.135	$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$	1008
3.136	$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$	1013
3.137	$\int \frac{1}{\sqrt{d-\frac{c^2dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)} dx$	1018
3.138	$\int \frac{1}{\sqrt{d-\frac{c^2dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2} dx$	1024
3.139	$\int \frac{1}{\sqrt{d-\frac{c^2dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^3} dx$	1029
3.140	$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx$	1034
3.141	$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx$	1040
3.142	$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx$	1045
3.143	$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)} dx$	1050
3.144	$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2} dx$	1056

3.145	$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^3} dx$ 1061
3.146	$\int \frac{\arctan(c(a+bx)) \log(d(a+bx))}{a+bx} dx$ 1066
3.147	$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx$ 1073
3.148	$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx$ 1079
3.149	$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$ 1087
3.150	$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx$ 1097
3.151	$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx$ 1107
3.152	$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx$ 1115
3.153	$\int \frac{(a+b \arctan(cx^n))(d+e \log(fx^m))}{x} dx$ 1122

3.1 $\int x^3 \arctan(a + bx^4) dx$

Optimal result	84
Mathematica [A] (verified)	84
Rubi [A] (warning: unable to verify)	85
Maple [A] (verified)	86
Fricas [A] (verification not implemented)	87
Sympy [A] (verification not implemented)	87
Maxima [A] (verification not implemented)	88
Giac [A] (verification not implemented)	88
Mupad [B] (verification not implemented)	88
Reduce [B] (verification not implemented)	89

Optimal result

Integrand size = 12, antiderivative size = 42

$$\int x^3 \arctan(a + bx^4) dx = \frac{(a + bx^4) \arctan(a + bx^4)}{4b} - \frac{\log(1 + (a + bx^4)^2)}{8b}$$

output

```
1/4*(b*x^4+a)*arctan(b*x^4+a)/b-1/8*ln(1+(b*x^4+a)^2)/b
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x^3 \arctan(a + bx^4) dx = -\frac{-2(a + bx^4) \arctan(a + bx^4) + \log(1 + (a + bx^4)^2)}{8b}$$

input

```
Integrate[x^3*ArcTan[a + b*x^4],x]
```

output

```
-1/8*(-2*(a + b*x^4)*ArcTan[a + b*x^4] + Log[1 + (a + b*x^4)^2])/b
```

Rubi [A] (warning: unable to verify)

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {7266, 5562, 5345, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x^3 \arctan(a + bx^4) dx \\
 \downarrow 7266 \\
 \frac{1}{4} \int \arctan(bx^4 + a) dx^4 \\
 \downarrow 5562 \\
 \frac{\int \arctan(bx^4 + a) d(bx^4 + a)}{4b} \\
 \downarrow 5345 \\
 \frac{(a + bx^4) \arctan(a + bx^4) - \int \frac{bx^4 + a}{x^8 + 1} d(bx^4 + a)}{4b} \\
 \downarrow 240 \\
 \frac{(a + bx^4) \arctan(a + bx^4) - \frac{1}{2} \log(x^8 + 1)}{4b}
 \end{array}$$

input `Int[x^3*ArcTan[a + b*x^4],x]`

output `((a + b*x^4)*ArcTan[a + b*x^4] - Log[1 + x^8]/2)/(4*b)`

Defintions of rubi rules used

rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 5345 `Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_), x_Symbol] := Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Simp[b*c*n*p Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

rule 5562 `Int[((a_) + ArcTan[(c_) + (d_)*(x_)])*(b_)^(p_), x_Symbol] := Simp[1/d Subst[Int[(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]`

rule 7266 `Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

method	result
derivativdivides	$\frac{(bx^4+a) \arctan(bx^4+a) - \frac{\ln(1+(bx^4+a)^2)}{2}}{4b}$
default	$\frac{(bx^4+a) \arctan(bx^4+a) - \frac{\ln(1+(bx^4+a)^2)}{2}}{4b}$
parallelrisch	$-\frac{-2 \arctan(bx^4+a)x^4b^2 - 2a \arctan(bx^4+a)b + \ln(b^2x^8 + 2abx^4 + a^2 + 1)b}{8b^2}$
parts	$\frac{x^4 \arctan(bx^4+a)}{4} - b \left(\frac{\ln(b^2x^8 + 2abx^4 + a^2 + 1)}{8b^2} - \frac{a \arctan\left(\frac{2b^2x^4 + 2ab}{4b^2}\right)}{4b^2} \right)$
risch	$-\frac{ix^4 \ln(1+i(bx^4+a))}{8} + \frac{ix^4 \ln(1-i(bx^4+a))}{8} + \frac{a \arctan\left(\frac{x^4b}{a^2+1} + \frac{a^2bx^4}{a^2+1} + \frac{a^3}{a^2+1} + \frac{a}{a^2+1}\right)}{4b} - \frac{a \arctan(a)}{4b} - \frac{\ln(1+a^2)}{4b}$

input `int(x^3*arctan(b*x^4+a),x,method=_RETURNVERBOSE)`

output `1/4/b*((b*x^4+a)*arctan(b*x^4+a)-1/2*ln(1+(b*x^4+a)^2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.07

$$\int x^3 \arctan(a + bx^4) dx = \frac{2(bx^4 + a) \arctan(bx^4 + a) - \log(b^2x^8 + 2abx^4 + a^2 + 1)}{8b}$$

input `integrate(x^3*arctan(b*x^4+a),x, algorithm="fricas")`

output `1/8*(2*(b*x^4 + a)*arctan(b*x^4 + a) - log(b^2*x^8 + 2*a*b*x^4 + a^2 + 1)) /b`

Sympy [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.43

$$\int x^3 \arctan(a + bx^4) dx = \begin{cases} \frac{a \operatorname{atan}(a+bx^4)}{4b} + \frac{x^4 \operatorname{atan}(a+bx^4)}{4} - \frac{\log(a^2+2abx^4+b^2x^8+1)}{8b} & \text{for } b \neq 0 \\ \frac{x^4 \operatorname{atan}(a)}{4} & \text{otherwise} \end{cases}$$

input `integrate(x**3*atan(b*x**4+a),x)`

output `Piecewise((a*atan(a + b*x**4)/(4*b) + x**4*atan(a + b*x**4)/4 - log(a**2 + 2*a*b*x**4 + b**2*x**8 + 1)/(8*b), Ne(b, 0)), (x**4*atan(a)/4, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x^3 \arctan(a + bx^4) dx = \frac{2(bx^4 + a) \arctan(bx^4 + a) - \log((bx^4 + a)^2 + 1)}{8b}$$

input `integrate(x^3*arctan(b*x^4+a),x, algorithm="maxima")`output `1/8*(2*(b*x^4 + a)*arctan(b*x^4 + a) - log((b*x^4 + a)^2 + 1))/b`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.88

$$\int x^3 \arctan(a + bx^4) dx = \frac{2(bx^4 + a) \arctan(bx^4 + a) - \log((bx^4 + a)^2 + 1)}{8b}$$

input `integrate(x^3*arctan(b*x^4+a),x, algorithm="giac")`output `1/8*(2*(b*x^4 + a)*arctan(b*x^4 + a) - log((b*x^4 + a)^2 + 1))/b`**Mupad [B] (verification not implemented)**

Time = 0.79 (sec) , antiderivative size = 230, normalized size of antiderivative = 5.48

$$\int x^3 \arctan(a + bx^4) dx = \frac{x^4 \operatorname{atan}(bx^4 + a)}{4} - \frac{\ln(a^2 + 2abx^4 + b^2x^8 + 1)}{8b} + \frac{a \operatorname{atan}\left(\frac{a}{a^6+3a^4+3a^2+1} + \frac{3a^3}{a^6+3a^4+3a^2+1} + \frac{3a^5}{a^6+3a^4+3a^2+1} + \frac{a^7}{a^6+3a^4+3a^2+1} + \frac{bx^4}{a^6+3a^4+3a^2+1} + \frac{3a^2bx^4}{a^6+3a^4+3a^2+1}\right)}{4b}$$

input `int(x^3*atan(a + b*x^4),x)`

output

```
(x^4*atan(a + b*x^4))/4 - log(a^2 + b^2*x^8 + 2*a*b*x^4 + 1)/(8*b) + (a*atan(a/(3*a^2 + 3*a^4 + a^6 + 1) + (3*a^3)/(3*a^2 + 3*a^4 + a^6 + 1) + (3*a^5)/(3*a^2 + 3*a^4 + a^6 + 1) + a^7/(3*a^2 + 3*a^4 + a^6 + 1) + (b*x^4)/(3*a^2 + 3*a^4 + a^6 + 1) + (3*a^2*b*x^4)/(3*a^2 + 3*a^4 + a^6 + 1) + (3*a^4*b*x^4)/(3*a^2 + 3*a^4 + a^6 + 1) + (a^6*b*x^4)/(3*a^2 + 3*a^4 + a^6 + 1)))/(4*b)
```

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 223, normalized size of antiderivative = 5.31

$$\int x^3 \arctan(a + bx^4) dx$$

$$= \frac{2atan(bx^4 + a)a + 2atan(bx^4 + a)bx^4 - \log\left((a^2 + 1)^{\frac{1}{4}} - b^{\frac{1}{4}}\sqrt{2(a^2 + 1)^{\frac{1}{4}} - \sqrt{\sqrt{a^2 + 1} - a}\sqrt{2}x + \sqrt{a^2 + 1}}\right)}{8b}$$

input

```
int(x^3*atan(b*x^4+a),x)
```

output

```
(2*atan(a + b*x**4)*a + 2*atan(a + b*x**4)*b*x**4 - log((a**2 + 1)**(1/4) - b**(1/4)*sqrt(2*(a**2 + 1)**(1/4) - sqrt(sqrt(a**2 + 1) - a)*sqrt(2))*x + sqrt(b)*x**2) - log((a**2 + 1)**(1/4) - b**(1/4)*sqrt(2*(a**2 + 1)**(1/4) + sqrt(sqrt(a**2 + 1) - a)*sqrt(2))*x + sqrt(b)*x**2) - log((a**2 + 1)**(1/4) + b**(1/4)*sqrt(2*(a**2 + 1)**(1/4) - sqrt(sqrt(a**2 + 1) - a)*sqrt(2))*x + sqrt(b)*x**2) - log((a**2 + 1)**(1/4) + b**(1/4)*sqrt(2*(a**2 + 1)**(1/4) + sqrt(sqrt(a**2 + 1) - a)*sqrt(2))*x + sqrt(b)*x**2))/(8*b)
```

3.2 $\int x^{-1+n} \arctan(a + bx^n) dx$

Optimal result	90
Mathematica [A] (verified)	90
Rubi [A] (warning: unable to verify)	91
Maple [C] (verified)	92
Fricas [A] (verification not implemented)	93
Sympy [F(-2)]	93
Maxima [A] (verification not implemented)	93
Giac [A] (verification not implemented)	94
Mupad [B] (verification not implemented)	94
Reduce [B] (verification not implemented)	94

Optimal result

Integrand size = 14, antiderivative size = 45

$$\int x^{-1+n} \arctan(a + bx^n) dx = \frac{(a + bx^n) \arctan(a + bx^n)}{bn} - \frac{\log(1 + (a + bx^n)^2)}{2bn}$$

output

```
(a+b*x^n)*arctan(a+b*x^n)/b/n-1/2*ln(1+(a+b*x^n)^2)/b/n
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.89

$$\int x^{-1+n} \arctan(a + bx^n) dx = -\frac{-2(a + bx^n) \arctan(a + bx^n) + \log(1 + (a + bx^n)^2)}{2bn}$$

input

```
Integrate[x^(-1 + n)*ArcTan[a + b*x^n], x]
```

output

```
-1/2*(-2*(a + b*x^n)*ArcTan[a + b*x^n] + Log[1 + (a + b*x^n)^2])/(b*n)
```

Rubi [A] (warning: unable to verify)

Time = 0.28 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {7266, 5562, 5345, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int x^{n-1} \arctan(a + bx^n) dx \\
 \downarrow 7266 \\
 \frac{\int \arctan(bx^n + a) dx^n}{n} \\
 \downarrow 5562 \\
 \frac{\int \arctan(bx^n + a) d(bx^n + a)}{bn} \\
 \downarrow 5345 \\
 \frac{(a + bx^n) \arctan(a + bx^n) - \int \frac{bx^n + a}{x^{2n+1}} d(bx^n + a)}{bn} \\
 \downarrow 240 \\
 \frac{(a + bx^n) \arctan(a + bx^n) - \frac{1}{2} \log(x^{2n} + 1)}{bn}
 \end{array}$$

input `Int[x^(-1 + n)*ArcTan[a + b*x^n], x]`

output `((a + b*x^n)*ArcTan[a + b*x^n] - Log[1 + x^(2*n)])/2)/(b*n)`

Defintions of rubi rules used

rule 240 $\text{Int}[(x_)/((a_)+(b_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^2, x]]/(2*b), x] /; \text{FreeQ}\{a, b\}, x]$

rule 5345 $\text{Int}(((a_)+\text{ArcTan}[(c_)*(x_)]*(b_))^p, x_Symbol) \rightarrow \text{Simp}[x*(a + b*\text{ArcTan}[c*x^n])^p, x] - \text{Simp}[b*c*n*p \text{Int}[x^n*((a + b*\text{ArcTan}[c*x^n])^{p-1})/(1 + c^2*x^{2*n})], x], x] /; \text{FreeQ}\{a, b, c, n\}, x \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ (\text{EqQ}[n, 1] \ || \ \text{EqQ}[p, 1])]$

rule 5562 $\text{Int}(((a_)+\text{ArcTan}[(c_)+(d_)*(x_)]*(b_))^p, x_Symbol) \rightarrow \text{Simp}[1/d \text{Subst}[\text{Int}[(a + b*\text{ArcTan}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{IGtQ}[p, 0]$

rule 7266 $\text{Int}[(u_)*(x_)^m, x_Symbol] \rightarrow \text{Simp}[1/(m + 1) \text{Subst}[\text{Int}[\text{SubstFor}[x^{(m+1)}, u, x], x, x^{(m+1)}], x], x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ \text{FunctionOfQ}[x^{(m+1)}, u, x]$

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 7.03 (sec) , antiderivative size = 140, normalized size of antiderivative = 3.11

method	result
risch	$-\frac{ix^n \ln(1+i(a+bx^n))}{2n} + \frac{ix^n \ln(1-i(a+bx^n))}{2n} - \frac{\ln\left(\frac{a+i}{b}+x^n\right)}{2nb} - \frac{\ln\left(x^n-\frac{i-a}{b}\right)}{2nb} + \frac{i \ln\left(\frac{a+i}{b}+x^n\right)a}{2nb} - \frac{i \ln\left(x^n-\frac{i-a}{b}\right)a}{2nb}$

input `int(x^(-1+n)*arctan(a+b*x^n),x,method=_RETURNVERBOSE)`

output
$$-1/2*I/n*x^n*\ln(1+I*(a+b*x^n))+1/2*I/n*x^n*\ln(1-I*(a+b*x^n))-1/2/n/b*\ln((a+I)/b+x^n)-1/2/n/b*\ln(x^n-(I-a)/b)+1/2*I/n/b*\ln((a+I)/b+x^n)*a-1/2*I/n/b*n(x^n-(I-a)/b)*a$$

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int x^{-1+n} \arctan(a + bx^n) dx = \frac{2bx^n \arctan(bx^n + a) + 2a \arctan(bx^n + a) - \log(b^2x^{2n} + 2abx^n + a^2 + 1)}{2bn}$$

input `integrate(x^(-1+n)*arctan(a+b*x^n),x, algorithm="fricas")`

output `1/2*(2*b*x^n*arctan(b*x^n + a) + 2*a*arctan(b*x^n + a) - log(b^2*x^(2*n) + 2*a*b*x^n + a^2 + 1))/(b*n)`

Sympy [F(-2)]

Exception generated.

$$\int x^{-1+n} \arctan(a + bx^n) dx = \text{Exception raised: HeuristicGCDFailed}$$

input `integrate(x**(-1+n)*atan(a+b*x**n),x)`

output `Exception raised: HeuristicGCDFailed >> no luck`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.89

$$\int x^{-1+n} \arctan(a + bx^n) dx = \frac{2(bx^n + a) \arctan(bx^n + a) - \log((bx^n + a)^2 + 1)}{2bn}$$

input `integrate(x^(-1+n)*arctan(a+b*x^n),x, algorithm="maxima")`

output `1/2*(2*(b*x^n + a)*arctan(b*x^n + a) - log((b*x^n + a)^2 + 1))/(b*n)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.89

$$\int x^{-1+n} \arctan(a + bx^n) dx = \frac{2(bx^n + a) \arctan(bx^n + a) - \log((bx^n + a)^2 + 1)}{2bn}$$

input `integrate(x^(-1+n)*arctan(a+b*x^n),x, algorithm="giac")`

output `1/2*(2*(b*x^n + a)*arctan(b*x^n + a) - log((b*x^n + a)^2 + 1))/(b*n)`

Mupad [B] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int x^{-1+n} \arctan(a + bx^n) dx = \frac{x^n \operatorname{atan}(a + bx^n)}{n} - \frac{\ln(a^2 + b^2 x^{2n} + 2abx^n + 1) - 2a \operatorname{atan}(a + bx^n)}{2bn}$$

input `int(x^(n - 1)*atan(a + b*x^n),x)`

output `(x^n*atan(a + b*x^n))/n - (log(a^2 + b^2*x^(2*n) + 2*a*b*x^n + 1) - 2*a*atan(a + b*x^n))/(2*b*n)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int x^{-1+n} \arctan(a + bx^n) dx = \frac{2x^n \operatorname{atan}(x^n b + a) b + 2 \operatorname{atan}(x^n b + a) a - \log(x^{2n} b^2 + 2x^n ab + a^2 + 1)}{2bn}$$

input `int(x^(-1+n)*atan(a+b*x^n),x)`

output
$$\frac{(2x^{2n}\operatorname{atan}(x^n b + a)b + 2\operatorname{atan}(x^n b + a)a - \log(x^{2n}b^2 + 2x^n a b + a^2 + 1))}{(2b^n)}$$

3.3 $\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	96
Mathematica [A] (verified)	96
Rubi [A] (verified)	97
Maple [B] (verified)	100
Fricas [A] (verification not implemented)	101
Sympy [A] (verification not implemented)	101
Maxima [F]	102
Giac [A] (verification not implemented)	102
Mupad [F(-1)]	103
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 25, antiderivative size = 144

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{5d^2x\sqrt{d+ex^2}}{96(-e)^{5/2}} + \frac{5dx^3\sqrt{d+ex^2}}{144(-e)^{3/2}} + \frac{x^5\sqrt{d+ex^2}}{36\sqrt{-e}} + \frac{1}{6}x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + \frac{5d^3\sqrt{-e}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{96e^{7/2}}$$

output

$$\frac{5}{96}d^2x\sqrt{d+ex^2}/(-e)^{5/2} + \frac{5}{144}d^3x^3\sqrt{d+ex^2}/(-e)^{3/2} + \frac{1}{36}x^5\sqrt{d+ex^2}/(-e)^{1/2} + \frac{1}{6}x^6\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + \frac{5}{96}d^3(-e)^{1/2}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)/e^{7/2}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.60

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{-ex}\sqrt{d+ex^2}(-15d^2 + 10dex^2 - 8e^2x^4) + 3(5d^3 + 16e^3x^6) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{288e^3}$$

input `Integrate[x^5*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(Sqrt[-e]*x*Sqrt[d + e*x^2]*(-15*d^2 + 10*d*e*x^2 - 8*e^2*x^4) + 3*(5*d^3 + 16*e^3*x^6)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(288*e^3)`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {5674, 262, 262, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow \text{5674} \\
 & \frac{1}{6}x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{-e} \int \frac{x^6}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{6}x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{-e} \left(\frac{x^5\sqrt{d+ex^2}}{6e} - \frac{5d \int \frac{x^4}{\sqrt{ex^2+d}} dx}{6e} \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{6}x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{-e} \left(\frac{x^5\sqrt{d+ex^2}}{6e} - \frac{5d \left(\frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \int \frac{x^2}{\sqrt{ex^2+d}} dx}{4e} \right)}{6e} \right) \\
 & \quad \downarrow \text{262}
 \end{aligned}$$

$$\frac{1}{6} \sqrt{-e} \left(\frac{\frac{1}{6} x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{x^5 \sqrt{d+ex^2}}{6e} - \frac{5d \left(\frac{x^3 \sqrt{d+ex^2}}{4e} - \frac{3d \left(\frac{x \sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{\sqrt{ex^2+d}} dx}{2e} \right)}{4e} \right)}{6e}}{6e} \right)$$

224

$$\frac{1}{6} \sqrt{-e} \left(\frac{\frac{1}{6} x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{x^5 \sqrt{d+ex^2}}{6e} - \frac{5d \left(\frac{x^3 \sqrt{d+ex^2}}{4e} - \frac{3d \left(\frac{x \sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{1 - \frac{ex^2}{ex^2+d}} d \frac{x}{\sqrt{ex^2+d}} \right)}{4e} \right)}{6e}}{6e} \right)$$

219

$$\frac{1}{6} \sqrt{-e} \left(\frac{\frac{1}{6} x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{x^5 \sqrt{d+ex^2}}{6e} - \frac{5d \left(\frac{x^3 \sqrt{d+ex^2}}{4e} - \frac{3d \left(\frac{x \sqrt{d+ex^2}}{2e} - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2e^{3/2}} \right)}{4e} \right)}{6e}}{6e} \right)$$

input `Int[x^5*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output

```
(x^6*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/6 - (Sqrt[-e]*((x^5*Sqrt[d + e*
x^2]))/(6*e) - (5*d*((x^3*Sqrt[d + e*x^2]))/(4*e) - (3*d*((x*Sqrt[d + e*x^2]
)/(2*e) - (d*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(2*e^(3/2))))/(4*e)))/(
6*e))/6
```

Defintions of rubi rules used

rule 219

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 224

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

rule 262

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/
(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

rule 5674

```
Int[ArcTan[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_))^(m_), x_S
ymbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 259 vs. 2(110) = 220.

Time = 0.51 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.81

method	result
default	$\frac{x^6 \arctan\left(\frac{\sqrt{-e} x}{\sqrt{e x^2 + d}}\right)}{6} + \frac{\sqrt{-e} e \frac{x^7 \sqrt{e x^2 + d}}{8e} - \left(\frac{7d \frac{x^5 \sqrt{e x^2 + d}}{6e} - \left(\frac{5d \left(\frac{x^3 \sqrt{e x^2 + d}}{4e} - \frac{3d \left(\frac{x \sqrt{e x^2 + d}}{2e} - \frac{d \ln(\sqrt{e} x + \sqrt{e x^2 + d})}{2e^{\frac{3}{2}}}\right)}{4e} \right)}{6e} \right)}{8e} \right)}{6d}$
parts	$\frac{x^6 \arctan\left(\frac{\sqrt{-e} x}{\sqrt{e x^2 + d}}\right)}{6} + \frac{\sqrt{-e} e \frac{x^7 \sqrt{e x^2 + d}}{8e} - \left(\frac{7d \frac{x^5 \sqrt{e x^2 + d}}{6e} - \left(\frac{5d \left(\frac{x^3 \sqrt{e x^2 + d}}{4e} - \frac{3d \left(\frac{x \sqrt{e x^2 + d}}{2e} - \frac{d \ln(\sqrt{e} x + \sqrt{e x^2 + d})}{2e^{\frac{3}{2}}}\right)}{4e} \right)}{6e} \right)}{8e} \right)}{6d}$

input

```
int(x^5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
1/6*x^6*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/6*(-e)^(1/2)*e/d*(1/8*x^7/e
*(e*x^2+d)^(1/2)-7/8*d/e*(1/6*x^5/e*(e*x^2+d)^(1/2)-5/6*d/e*(1/4*x^3/e*(e*
x^2+d)^(1/2)-3/4*d/e*(1/2*x/e*(e*x^2+d)^(1/2)-1/2*d/e^(3/2)*ln(e^(1/2)*x+(
e*x^2+d)^(1/2)))))-1/6*(-e)^(1/2)/d*(1/8*x^5*(e*x^2+d)^(3/2)/e-5/8*d/e*(1
/6*x^3*(e*x^2+d)^(3/2)/e-1/2*d/e*(1/4*x*(e*x^2+d)^(3/2)/e-1/4*d/e*(1/2*x*(
e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.53

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= -\frac{(8e^2x^5 - 10dex^3 + 15d^2x)\sqrt{ex^2+d}\sqrt{-e} - 3(16e^3x^6 + 5d^3)\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{288e^3}$$

input

```
integrate(x^5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")
```

output

```
-1/288*((8*e^2*x^5 - 10*d*e*x^3 + 15*d^2*x)*sqrt(e*x^2 + d)*sqrt(-e) - 3*(
16*e^3*x^6 + 5*d^3)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/e^3
```

Sympy [A] (verification not implemented)

Time = 0.99 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.96

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \begin{cases} \frac{5d^3 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{96e^3} - \frac{5d^2x\sqrt{-e}\sqrt{d+ex^2}}{96e^3} + \frac{5dx^3\sqrt{-e}\sqrt{d+ex^2}}{144e^2} + \frac{x^6 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{6} - \frac{x^5\sqrt{-e}\sqrt{d+ex^2}}{36e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input

```
integrate(x**5*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)
```

output

```
Piecewise((5*d**3*atan(x*sqrt(-e)/sqrt(d + e*x**2))/(96*e**3) - 5*d**2*x*sqrt(-e)*sqrt(d + e*x**2)/(96*e**3) + 5*d*x**3*sqrt(-e)*sqrt(d + e*x**2)/(144*e**2) + x**6*atan(x*sqrt(-e)/sqrt(d + e*x**2))/6 - x**5*sqrt(-e)*sqrt(d + e*x**2)/(36*e), Ne(e, 0)), (0, True))
```

Maxima [F]

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input

```
integrate(x^5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
1/6*x^6*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - d*sqrt(-e)*integrate(-1/6*sqrt(e*x^2 + d)*x^6/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)
```

Giac [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.65

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{6} x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \frac{1}{288} \sqrt{-e^2x^2 - de} \left(2x^2 \left(\frac{4x^2}{e} - \frac{5d}{e^2}\right) + \frac{15d^2}{e^3}\right) x - \frac{5d^3 \arcsin\left(\frac{ex}{\sqrt{-de}}\right) \operatorname{sgn}(e)}{96e^2|e|}$$

input

```
integrate(x^5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
1/6*x^6*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/288*sqrt(-e^2*x^2 - d*e)*(2*x^2*(4*x^2/e - 5*d/e^2) + 15*d^2/e^3)*x - 5/96*d^3*arcsin(e*x/sqrt(-d*e))*sgn(e)/(e^2*abs(e))
```

Mupad [F(-1)]

Timed out.

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^5 \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^5*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2),x)`

output `int(x^5*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.10

$$\int x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{15 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+ex^2}}\right) d^3 + 48 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+ex^2}}\right) e^3 x^6 - 15\sqrt{e}\sqrt{ex^2+d} d^2 i x + 10\sqrt{e}\sqrt{ex^2-d}}{288e^3}$$

input `int(x^5*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2),x)`

output `(15*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d**3 + 48*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e**3*x**6 - 15*sqrt(e)*sqrt(d + e*x**2)*d**2*i*x + 10*sqrt(e)*sqrt(d + e*x**2)*d*e*i*x**3 - 8*sqrt(e)*sqrt(d + e*x**2)*e**2*i*x**5)/(288*e**3)`

3.4 $\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [B] (verified)	107
Fricas [A] (verification not implemented)	107
Sympy [A] (verification not implemented)	108
Maxima [F]	108
Giac [A] (verification not implemented)	109
Mupad [F(-1)]	109
Reduce [B] (verification not implemented)	110

Optimal result

Integrand size = 25, antiderivative size = 116

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{3dx\sqrt{d+ex^2}}{32(-e)^{3/2}} + \frac{x^3\sqrt{d+ex^2}}{16\sqrt{-e}} + \frac{1}{4}x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{3d^2\sqrt{-e}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{32e^{5/2}}$$

output

```
3/32*d*x*(e*x^2+d)^(1/2)/(-e)^(3/2)+1/16*x^3*(e*x^2+d)^(1/2)/(-e)^(1/2)+1/4*x^4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))-3/32*d^2*(-e)^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/e^(5/2)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.64

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{-ex}(3d-2ex^2)\sqrt{d+ex^2} + (-3d^2+8e^2x^4)\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{32e^2}$$

input `Integrate[x^3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(Sqrt[-e]*x*(3*d - 2*e*x^2)*Sqrt[d + e*x^2] + (-3*d^2 + 8*e^2*x^4)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(32*e^2)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5674, 262, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow \text{5674} \\
 & \frac{1}{4}x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{-e} \int \frac{x^4}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4}x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{-e} \left(\frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \int \frac{x^2}{\sqrt{ex^2+d}} dx}{4e} \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{1}{4}x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{-e} \left(\frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left(\frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{\sqrt{ex^2+d}} dx}{2e} \right)}{4e} \right) \\
 & \quad \downarrow \text{224} \\
 & \frac{1}{4}x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{-e} \left(\frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d \left(\frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{1-\frac{ex^2}{d}} d\frac{x}{\sqrt{ex^2+d}} dx}{2e} \right)}{4e} \right)
 \end{aligned}$$

$$\frac{1}{4}x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{4}\sqrt{-e} \left(\frac{x^3\sqrt{d+ex^2}}{4e} - \frac{3d\left(\frac{x\sqrt{d+ex^2}}{2e} - \frac{d\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2e^{3/2}}\right)}{4e} \right)$$

input `Int[x^3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(x^4*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/4 - (Sqrt[-e]*((x^3*Sqrt[d + e*x^2])/(4*e) - (3*d*((x*Sqrt[d + e*x^2])/(2*e) - (d*ArcTanh[(Sqrt[e]*x)/Sqrt[d + e*x^2]])/(2*e^(3/2)))))/(4*e))/4`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. 2(88) = 176.

Time = 0.02 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.83

method	result
default	$\frac{x^4 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{4} + \frac{\sqrt{-e}e \left(\frac{x^5 \sqrt{ex^2+d}}{6e} - \frac{5d \left(\frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left(\frac{x\sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}}\right)}{4e} \right)}{6e} \right)}{4d} - \frac{\sqrt{-e} \left(\frac{x^3(e x^2+d)}{6e} \right)}{4d}$
parts	$\frac{x^4 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{4} + \frac{\sqrt{-e}e \left(\frac{x^5 \sqrt{ex^2+d}}{6e} - \frac{5d \left(\frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left(\frac{x\sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}}\right)}{4e} \right)}{6e} \right)}{4d} - \frac{\sqrt{-e} \left(\frac{x^3(e x^2+d)}{6e} \right)}{4d}$

input

```
int(x^3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
1/4*x^4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/4*(-e)^(1/2)*e/d*(1/6*x^5/e
*(e*x^2+d)^(1/2)-5/6*d/e*(1/4*x^3/e*(e*x^2+d)^(1/2)-3/4*d/e*(1/2*x/e*(e*x^
2+d)^(1/2)-1/2*d/e^(3/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))-1/4*(-e)^(1/2)/d
*(1/6*x^3*(e*x^2+d)^(3/2)/e-1/2*d/e*(1/4*x*(e*x^2+d)^(3/2)/e-1/4*d/e*(1/2*
x*(e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.56

$$\int x^3 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right) dx$$

$$= -\frac{(2ex^3 - 3dx)\sqrt{ex^2+d}\sqrt{-e} - (8e^2x^4 - 3d^2) \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{32e^2}$$

input `integrate(x^3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `-1/32*((2*e*x^3 - 3*d*x)*sqrt(e*x^2 + d)*sqrt(-e) - (8*e^2*x^4 - 3*d^2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/e^2`

Sympy [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.92

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \begin{cases} -\frac{3d^2 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{32e^2} + \frac{3dx\sqrt{-e}\sqrt{d+ex^2}}{32e^2} + \frac{x^4 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{4} - \frac{x^3\sqrt{-e}\sqrt{d+ex^2}}{16e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x**3*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Piecewise((-3*d**2*atan(x*sqrt(-e)/sqrt(d + e*x**2))/(32*e**2) + 3*d*x*sqrt(-e)*sqrt(d + e*x**2)/(32*e**2) + x**4*atan(x*sqrt(-e)/sqrt(d + e*x**2))/4 - x**3*sqrt(-e)*sqrt(d + e*x**2)/(16*e), Ne(e, 0)), (0, True))`

Maxima [F]

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `1/4*x^4*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - d*sqrt(-e)*integrate(-1/4*sqrt(e*x^2 + d)*x^4/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

Giac [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.69

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{4} x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \frac{1}{32} \sqrt{-e^2x^2-dex} \left(\frac{2x^2}{e} - \frac{3d}{e^2}\right) + \frac{3d^2 \arcsin\left(\frac{ex}{\sqrt{-de}}\right) \operatorname{sgn}(e)}{32e|e|}$$

input `integrate(x^3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `1/4*x^4*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/32*sqrt(-e^2*x^2 - d*e)*x*(2*x^2/e - 3*d/e^2) + 3/32*d^2*arcsin(e*x/sqrt(-d*e))*sgn(e)/(e*abs(e))`

Mupad [F(-1)]

Timed out.

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^3 \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^3*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2),x)`

output `int(x^3*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.18

$$\int x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{-3\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}+d+ex^2}\right) d^2 + 8\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}+d+ex^2}\right) e^2 x^4 + 3\sqrt{e}\sqrt{ex^2+d} dix - 2\sqrt{e}\sqrt{ex^2+d}}{32e^2}$$

input `int(x^3*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output

```
( - 3*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d**2 + 8*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e**2*x**4 + 3*sqrt(e)*sqrt(d + e*x**2)*d*i*x - 2*sqrt(e)*sqrt(d + e*x**2)*e*i*x**3)/(32*e**2)
```

3.5 $\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	111
Mathematica [A] (verified)	111
Rubi [A] (verified)	112
Maple [B] (verified)	113
Fricas [A] (verification not implemented)	114
Sympy [A] (verification not implemented)	114
Maxima [F]	115
Giac [A] (verification not implemented)	115
Mupad [F(-1)]	116
Reduce [B] (verification not implemented)	116

Optimal result

Integrand size = 23, antiderivative size = 88

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{x\sqrt{d+ex^2}}{4\sqrt{-e}} + \frac{1}{2}x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + \frac{d\sqrt{-e}\operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{4e^{3/2}}$$

output

```
1/4*x*(e*x^2+d)^(1/2)/(-e)^(1/2)+1/2*x^2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/4*d*(-e)^(1/2)*arctanh(e^(1/2)*x/(e*x^2+d)^(1/2))/e^(3/2)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.67

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{-\sqrt{-ex}\sqrt{d+ex^2} + (d+2ex^2) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{4e}$$

input

```
Integrate[x*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]
```


output

$$\frac{-(\text{Sqrt}[-e]*x*\text{Sqrt}[d + e*x^2]) + (d + 2*e*x^2)*\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]]}{4*e}$$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {5674, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{5674} \\ & \frac{1}{2}x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{-e} \int \frac{x^2}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \\ & \frac{1}{2}x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{-e} \left(\frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{\sqrt{ex^2+d}} dx}{2e} \right) \\ & \quad \downarrow \text{224} \\ & \frac{1}{2}x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{-e} \left(\frac{x\sqrt{d+ex^2}}{2e} - \frac{d \int \frac{1}{1-\frac{ex^2}{ex^2+d}} d\frac{x}{\sqrt{ex^2+d}}}{2e} \right) \\ & \quad \downarrow \text{219} \\ & \frac{1}{2}x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{2}\sqrt{-e} \left(\frac{x\sqrt{d+ex^2}}{2e} - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{ex}}{\sqrt{d+ex^2}}\right)}{2e^{3/2}} \right) \end{aligned}$$

input

$$\text{Int}[x*\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]], x]$$

output

$$\frac{(x^2*\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]])/2 - (\text{Sqrt}[-e]*((x*\text{Sqrt}[d + e*x^2])/2)))/(2*e) - (d*\text{ArcTanh}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d + e*x^2]])/(2*e^{(3/2)})/2$$

Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 224 Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

```
rule 262 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)
^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/
(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b
, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c
, 2, m, p, x]
```

```
rule 5674 Int[ArcTan[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_))^(m_), x_S
ymbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(66) = 132.

Time = 0.02 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.86

method	result
default	$\frac{x^2 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{2} + \frac{\sqrt{-e}e \left(\frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left(\frac{x\sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{2d} - \frac{\sqrt{-e} \left(\frac{x(ex^2+d)^{\frac{3}{2}}}{4e} - \frac{d \left(\frac{x\sqrt{ex^2+d}}{2} + \dots \right)}{2d} \right)}{2d}$
parts	$\frac{x^2 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{2} + \frac{\sqrt{-e}e \left(\frac{x^3 \sqrt{ex^2+d}}{4e} - \frac{3d \left(\frac{x\sqrt{ex^2+d}}{2e} - \frac{d \ln(\sqrt{e}x + \sqrt{ex^2+d})}{2e^{\frac{3}{2}}} \right)}{4e} \right)}{2d} - \frac{\sqrt{-e} \left(\frac{x(ex^2+d)^{\frac{3}{2}}}{4e} - \frac{d \left(\frac{x\sqrt{ex^2+d}}{2} + \dots \right)}{2d} \right)}{2d}$

input `int(x*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output `1/2*x^2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/2*(-e)^(1/2)*e/d*(1/4*x^3/e*(e*x^2+d)^(1/2)-3/4*d/e*(1/2*x/e*(e*x^2+d)^(1/2)-1/2*d/e^(3/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2))))-1/2*(-e)^(1/2)/d*(1/4*x*(e*x^2+d)^(3/2)/e-1/4*d/e*(1/2*x*(e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.56

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = -\frac{\sqrt{ex^2+d}\sqrt{-ex} - (2ex^2+d) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{4e}$$

input `integrate(x*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `-1/4*(sqrt(e*x^2 + d)*sqrt(-e)*x - (2*e*x^2 + d)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/e`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.83

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \begin{cases} \frac{d \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{4e} + \frac{x^2 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{2} - \frac{x\sqrt{-e}\sqrt{d+ex^2}}{4e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output

```
Piecewise((d*atan(x*sqrt(-e)/sqrt(d + e*x**2))/(4*e) + x**2*atan(x*sqrt(-e)
)/sqrt(d + e*x**2))/2 - x*sqrt(-e)*sqrt(d + e*x**2)/(4*e), Ne(e, 0)), (0,
True))
```

Maxima [F]

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input

```
integrate(x*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
1/2*x^2*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - d*sqrt(-e)*integrate(-1/2*s
qrt(e*x^2 + d)*x^2/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)
```

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.72

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{2} x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \frac{d \arcsin\left(\frac{ex}{\sqrt{-de}}\right) \operatorname{sgn}(e)}{4|e|} - \frac{\sqrt{-e^2x^2-dex}}{4e}$$

input

```
integrate(x*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
1/2*x^2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/4*d*arcsin(e*x/sqrt(-d*e))*
sgn(e)/abs(e) - 1/4*sqrt(-e^2*x^2 - d*e)*x/e
```

Mupad [F(-1)]

Timed out.

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2),x)`

output `int(x*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.57

$$\int x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{2 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+d+ex^2}}\right) d + 2 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+dx+ex^2}}{\sqrt{e}\sqrt{ex^2+dx+d+ex^2}}\right) ex^2 - \sqrt{e}\sqrt{ex^2+d}ix - \log\left(\frac{\sqrt{ex^2+d}+\sqrt{ex}}{\sqrt{d}}\right) di}{4e}$$

input `int(x*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2),x)`

output `(2*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d + 2*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e*x**2 - sqrt(e)*sqrt(d + e*x**2)*i*x - log((sqrt(d + e*x**2) + sqrt(e)*x)/sqrt(d))*d*i)/(4*e)`

3.6
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx$$

Optimal result	117
Mathematica [A] (verified)	118
Rubi [C] (verified)	118
Maple [F]	123
Fricas [F]	123
Sympy [F]	123
Maxima [F]	124
Giac [F]	124
Mupad [F(-1)]	124
Reduce [F]	125

Optimal result

Integrand size = 25, antiderivative size = 288

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx = -\frac{\sqrt{d}\sqrt{-e}\sqrt{1+\frac{ex^2}{d}}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)^2}{2\sqrt{e}\sqrt{d+ex^2}} + \frac{\sqrt{d}\sqrt{-e}\sqrt{1+\frac{ex^2}{d}}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)}{\sqrt{e}\sqrt{d+ex^2}} - \frac{\sqrt{d}\sqrt{-e}\sqrt{1+\frac{ex^2}{d}}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log(x)}{\sqrt{e}\sqrt{d+ex^2}} + \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)\log(x) + \frac{\sqrt{d}\sqrt{-e}\sqrt{1+\frac{ex^2}{d}}\operatorname{PolyLog}\left(2,e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)}{2\sqrt{e}\sqrt{d+ex^2}}$$

output

$$\begin{aligned}
& -1/2*d^{(1/2)}*(-e)^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{arcsinh}(e^{(1/2)}*x/d^{(1/2)})^2/e^{(1/2)}/(e*x^2+d)^{(1/2)}+d^{(1/2)}*(-e)^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{arcsinh}(e^{(1/2)}*x/d^{(1/2)})*\ln(1-(e^{(1/2)}*x/d^{(1/2)}+(1+e*x^2/d)^{(1/2)})^2)/e^{(1/2)}/(e*x^2+d)^{(1/2)}-d^{(1/2)}*(-e)^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{arcsinh}(e^{(1/2)}*x/d^{(1/2)})*\ln(x)/e^{(1/2)}/(e*x^2+d)^{(1/2)}+\arctan((-e)^{(1/2)}*x/(e*x^2+d)^{(1/2)})*\ln(x)+1/2*d^{(1/2)}*(-e)^{(1/2)}*(1+e*x^2/d)^{(1/2)}*\operatorname{polylog}(2,(e^{(1/2)}*x/d^{(1/2)}+(1+e*x^2/d)^{(1/2)})^2)/e^{(1/2)}/(e*x^2+d)^{(1/2)}
\end{aligned}$$
Mathematica [A] (verified)

Time = 1.65 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.59

$$\begin{aligned}
& \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) \log(x) \\
& + \frac{\sqrt{-e}\sqrt{1+\frac{ex^2}{d}}\left(\operatorname{arcsinh}\left(\sqrt{\frac{e}{d}}x\right)^2 + 2\operatorname{arcsinh}\left(\sqrt{\frac{e}{d}}x\right) \log\left(1 - e^{-2\operatorname{arcsinh}\left(\sqrt{\frac{e}{d}}x\right)}\right) - 2\log(x) \log\left(\sqrt{\frac{e}{d}}x + \sqrt{d+ex^2}\right)\right)}{2\sqrt{\frac{e}{d}}\sqrt{d+ex^2}}
\end{aligned}$$

input

`Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x,x]`

output

$$\begin{aligned}
& \operatorname{ArcTan}\left[\frac{\operatorname{Sqrt}[-e]*x}{\operatorname{Sqrt}[d + e*x^2]}\right]*\operatorname{Log}[x] + \left(\operatorname{Sqrt}[-e]*\operatorname{Sqrt}\left[1 + \frac{e*x^2}{d}\right]/d\right)*\left(\operatorname{ArcSinh}\left[\operatorname{Sqrt}[e/d]*x\right]^2 + 2*\operatorname{ArcSinh}\left[\operatorname{Sqrt}[e/d]*x\right]*\operatorname{Log}\left[1 - E^{-2*\operatorname{ArcSinh}\left[\operatorname{Sqrt}[e/d]*x\right]}\right] - 2*\operatorname{Log}[x]*\operatorname{Log}\left[\operatorname{Sqrt}[e/d]*x + \operatorname{Sqrt}\left[1 + \frac{e*x^2}{d}\right]\right] - \operatorname{PolyLog}\left[2, E^{-2*\operatorname{ArcSinh}\left[\operatorname{Sqrt}[e/d]*x\right]}\right]\right)\right)/(2*\operatorname{Sqrt}[e/d]*\operatorname{Sqrt}[d + e*x^2])
\end{aligned}$$
Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.66, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {5672, 2764, 2762, 6190, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx \\
& \quad \downarrow \text{5672} \\
& \log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \sqrt{-e} \int \frac{\log(x)}{\sqrt{ex^2+d}} dx \\
& \quad \downarrow \text{2764} \\
& \log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \int \frac{\log(x)}{\sqrt{\frac{ex^2}{d}+1}} dx}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{2762} \\
& \log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} - \sqrt{d} \int \frac{\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} dx \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{6190} \\
& \frac{\log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}\sqrt{\frac{ex^2}{d}+1}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{ex}} d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{3042} \\
& \log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} - \frac{\sqrt{d} \int -i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) \tan\left(i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{26} \\
& \log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d} \int \operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) \tan\left(i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) + \frac{\pi}{2}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} \right)}{\sqrt{d+ex^2}} \\
& \quad \downarrow \text{4199}
\end{aligned}$$

$$\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(2i\int \frac{e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} \operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) - \frac{1}{2}i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)^2\right)}{1-e}}{\sqrt{e}} \right)$$

$\sqrt{d+ex^2}$

↓ 25

$$\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\int \frac{e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} \operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) - \frac{1}{2}i\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)^2\right)}{1-e}}{\sqrt{e}} \right)$$

$\sqrt{d+ex^2}$

↓ 2620

$$\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\left(\frac{1}{2}\int \log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right) d\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) - \frac{1}{2}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) \log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)\right)}{\sqrt{e}} \right)$$

$\sqrt{d+ex^2}$

↓ 2715

$$\sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\left(\frac{1}{4}\int e^{-2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} \log\left(1-e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right) de^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)} - \frac{1}{2}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\right)}{\sqrt{e}} \right)$$

$\sqrt{d+ex^2}$

↓ 2838

$$\frac{\log(x) \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \sqrt{-e}\sqrt{\frac{ex^2}{d}+1} \left(\frac{\sqrt{d}\log(x)\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{i\sqrt{d}\left(-2i\left(-\frac{1}{4}\operatorname{PolyLog}\left(2, e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)\right) - \frac{1}{2}\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)\log\left(1 - e^{2\operatorname{arcsinh}\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}\right)\right)}{\sqrt{e}} \right)}{\sqrt{d+ex^2}}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x,x]`

output `ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]*Log[x] - (Sqrt[-e]*Sqrt[1 + (e*x^2)/d] * ((Sqrt[d]*ArcSinh[(Sqrt[e]*x)/Sqrt[d]]*Log[x])/Sqrt[e] + (I*Sqrt[d]*((-1/2*I)*ArcSinh[(Sqrt[e]*x)/Sqrt[d]]^2 - (2*I)*(-1/2*(ArcSinh[(Sqrt[e]*x)/Sqrt[d]]*Log[1 - E^(2*ArcSinh[(Sqrt[e]*x)/Sqrt[d]])) - PolyLog[2, E^(2*ArcSinh[(Sqrt[e]*x)/Sqrt[d]]])/4)))/Sqrt[e]))/Sqrt[d + e*x^2]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2762 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[e, 2]*(x/Sqrt[d])]*((a + b*Log[c*x^n])/Rt[e, 2]), x] - Simp[b*(n/Rt[e, 2]) Int[ArcSinh[Rt[e, 2]*(x/Sqrt[d])]/x, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && GtQ[d, 0] && PosQ[e]`

rule 2764 `Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[Sqrt[1 + (e/d)*x^2]/Sqrt[d + e*x^2] Int[(a + b*Log[c*x^n])/Sqrt[1 + (e/d)*x^2], x], x] /; FreeQ[{a, b, c, d, e, n}, x] && !GtQ[d, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 5672 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]/(x_), x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])*Log[x], x] - Simp[c Int[Log[x]/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c}, x] && EqQ[b + c^2, 0]`

rule 6190 `Int[((a_.) + ArcSinh[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Coth[-a/b + x/b], x], x, a + b*ArcSinh[c*x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x,x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x,x)`

Fricas [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x,x, algorithm="fricas")`

output `integral(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x, x)`

Sympy [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{x} dx$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x,x)`

output `Integral(atan(x*sqrt(-e)/sqrt(d + e*x**2))/x, x)`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x,x, algorithm="maxima")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x, x)`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x,x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x,x`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x, x`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x,x)`

output `int(atan((sqrt(e)*i*x)/sqrt(d + e*x**2))/x,x)`

3.7 $\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$

Optimal result	126
Mathematica [A] (verified)	126
Rubi [A] (verified)	127
Maple [B] (verified)	128
Fricas [A] (verification not implemented)	128
Sympy [A] (verification not implemented)	129
Maxima [A] (verification not implemented)	129
Giac [B] (verification not implemented)	129
Mupad [F(-1)]	130
Reduce [B] (verification not implemented)	130

Optimal result

Integrand size = 25, antiderivative size = 57

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{2dx} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{2x^2}$$

output

```
-1/2*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x-1/2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.95

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\sqrt{-ex}\sqrt{d+ex^2} + d \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{2dx^2}$$

input

```
Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^3,x]
```

output

```
-1/2*(Sqrt[-e]*x*Sqrt[d + e*x^2] + d*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(d*x^2)
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {5674, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx$$

↓ 5674

$$\frac{1}{2}\sqrt{-e} \int \frac{1}{x^2\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{2x^2}$$

↓ 242

$$-\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{2x^2} - \frac{\sqrt{-e}\sqrt{d+ex^2}}{2dx}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^3,x]`

output `-1/2*(Sqrt[-e]*Sqrt[d + e*x^2])/(d*x) - ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(2*x^2)`

Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] :> Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. 2(45) = 90.

Time = 0.03 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.14

method	result	size
default	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{e x^2+d}}\right)}{2x^2} - \frac{\sqrt{-e}\sqrt{e}\ln(\sqrt{e}x+\sqrt{e x^2+d})}{2d} + \frac{\sqrt{-e}\left(-\frac{(e x^2+d)^{\frac{3}{2}}}{dx} + \frac{2e\left(\frac{x\sqrt{e x^2+d}}{2} + \frac{d\ln(\sqrt{e}x+\sqrt{e x^2+d})}{2\sqrt{e}}\right)}{d}\right)}{2d}$	122
parts	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{e x^2+d}}\right)}{2x^2} - \frac{\sqrt{-e}\sqrt{e}\ln(\sqrt{e}x+\sqrt{e x^2+d})}{2d} + \frac{\sqrt{-e}\left(-\frac{(e x^2+d)^{\frac{3}{2}}}{dx} + \frac{2e\left(\frac{x\sqrt{e x^2+d}}{2} + \frac{d\ln(\sqrt{e}x+\sqrt{e x^2+d})}{2\sqrt{e}}\right)}{d}\right)}{2d}$	122

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x,method=_RETURNVERBOSE)`

output `-1/2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2-1/2*(-e)^(1/2)*e^(1/2)/d*ln(e^(1/2)*x+(e*x^2+d)^(1/2))+1/2*(-e)^(1/2)/d*(-1/d/x*(e*x^2+d)^(3/2)+2*e/d*(1/2*x*(e*x^2+d)^(1/2)+1/2*d/e^(1/2)*ln(e^(1/2)*x+(e*x^2+d)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.77

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\sqrt{ex^2+d}\sqrt{-e}x + d\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{2dx^2}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x, algorithm="fricas")`

output `-1/2*(sqrt(e*x^2 + d)*sqrt(-e)*x + d*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/(d*x^2)`

Sympy [A] (verification not implemented)

Time = 1.86 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.93

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{2x^2} - \frac{\sqrt{e}\sqrt{-e}\sqrt{\frac{d}{ex^2} + 1}}{2d}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**3,x)`

output `-atan(x*sqrt(-e)/sqrt(d + e*x**2))/(2*x**2) - sqrt(e)*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(2*d)`

Maxima [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.02

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = -\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{2x^2} - \frac{\sqrt{-e}ex^2 + d\sqrt{-e}}{2\sqrt{ex^2+dx}}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x, algorithm="maxima")`

output `-1/2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^2 - 1/2*(sqrt(-e)*e*x^2 + d*sqrt(-e))/(sqrt(e*x^2 + d)*d*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs. 2(45) = 90.

Time = 0.23 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.82

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \frac{e^4 x}{4(\sqrt{-dee} + \sqrt{-e^2 x^2 - de}|e|)d|e|} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{2x^2} - \frac{\sqrt{-dee} + \sqrt{-e^2 x^2 - de}|e|}{4dx|e|}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x, algorithm="giac")`

output `1/4*e^4*x/((sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))*d*abs(e)) - 1/2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^2 - 1/4*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))/(d*x*abs(e))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^3} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^3,x)`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^3, x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.21

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^3} dx = \frac{-\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d - \sqrt{e}\sqrt{ex^2+d}ix}{2dx^2}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^3,x)`

output `(- (atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d + sqrt(e)*sqrt(d + e*x**2)*i*x)/(2*d*x**2)`

$$3.8 \quad \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$$

Optimal result	131
Mathematica [A] (verified)	131
Rubi [A] (verified)	132
Maple [A] (verified)	133
Fricas [A] (verification not implemented)	134
Sympy [A] (verification not implemented)	134
Maxima [A] (verification not implemented)	134
Giac [B] (verification not implemented)	135
Mupad [F(-1)]	135
Reduce [B] (verification not implemented)	136

Optimal result

Integrand size = 25, antiderivative size = 85

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{12dx^3} - \frac{(-e)^{3/2}\sqrt{d+ex^2}}{6d^2x} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

output

```
-1/12*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^3-1/6*(-e)^(3/2)*(e*x^2+d)^(1/2)/d^2/
x-1/4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^4
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.79

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \frac{\sqrt{-ex}\sqrt{d+ex^2}(-d+2ex^2) - 3d^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{12d^2x^4}$$

input

```
Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^5,x]
```

output

```
(Sqrt[-e]*x*Sqrt[d + e*x^2]*(-d + 2*e*x^2) - 3*d^2*ArcTan[(Sqrt[-e]*x)/Sqr
t[d + e*x^2]])/(12*d^2*x^4)
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.99, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5674, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$$

↓ 5674

$$\frac{1}{4}\sqrt{-e} \int \frac{1}{x^4\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

↓ 245

$$\frac{1}{4}\sqrt{-e} \left(-\frac{2e \int \frac{1}{x^2\sqrt{ex^2+d}} dx}{3d} - \frac{\sqrt{d+ex^2}}{3dx^3} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

↓ 242

$$\frac{1}{4}\sqrt{-e} \left(\frac{2e\sqrt{d+ex^2}}{3d^2x} - \frac{\sqrt{d+ex^2}}{3dx^3} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{4x^4}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^5,x]`

output `(Sqrt[-e]*(-1/3*Sqrt[d + e*x^2]/(d*x^3) + (2*e*Sqrt[d + e*x^2])/(3*d^2*x)))/4 - ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(4*x^4)`

Definitions of rubi rules used

rule 242 $\text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^2)^p, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot (a + b \cdot x^2)^{p+1} / (a \cdot c \cdot (m+1)), x] /;$ $\text{FreeQ}\{a, b, c, m, p\}, x]$ && $\text{EqQ}[m + 2 \cdot p + 3, 0]$ && $\text{NeQ}[m, -1]$

rule 245 $\text{Int}[x^m \cdot (a + b \cdot x^2)^p, x_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot (a + b \cdot x^2)^{p+1} / (a \cdot (m+1)), x] - \text{Simp}[b \cdot (m + 2 \cdot (p + 1) + 1) / (a \cdot (m + 1)) \cdot \text{Int}[x^{m+2} \cdot (a + b \cdot x^2)^p, x], x] /;$ $\text{FreeQ}\{a, b, m, p\}, x]$ && $\text{ILtQ}[\text{Simplify}[(m + 1) / 2 + p + 1], 0]$ && $\text{NeQ}[m, -1]$

rule 5674 $\text{Int}[\text{ArcTan}[(c \cdot x) / \text{Sqrt}[a + b \cdot x^2]] \cdot (d \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[(d \cdot x)^{m+1} \cdot \text{ArcTan}[c \cdot x / \text{Sqrt}[a + b \cdot x^2]] / (d \cdot (m + 1)), x] - \text{Simp}[c / (d \cdot (m + 1)) \cdot \text{Int}[(d \cdot x)^{m+1} / \text{Sqrt}[a + b \cdot x^2], x], x] /;$ $\text{FreeQ}\{a, b, c, d, m\}, x]$ && $\text{EqQ}[b + c^2, 0]$ && $\text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{4x^4} + \frac{\sqrt{-e}e\sqrt{ex^2+d}}{4d^2x} - \frac{\sqrt{-e}(ex^2+d)^{\frac{3}{2}}}{12d^2x^3}$	69
parts	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{4x^4} + \frac{\sqrt{-e}e\sqrt{ex^2+d}}{4d^2x} - \frac{\sqrt{-e}(ex^2+d)^{\frac{3}{2}}}{12d^2x^3}$	69

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x,method=_RETURNVERBOSE)`

output
$$-1/4 \cdot \arctan((-e)^{1/2} \cdot x / (e \cdot x^2 + d)^{1/2}) / x^4 + 1/4 \cdot (-e)^{1/2} \cdot e / d^2 \cdot (e \cdot x^2 + d)^{1/2} / x - 1/12 \cdot (-e)^{1/2} / d^2 \cdot x^3 \cdot (e \cdot x^2 + d)^{3/2}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.68

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = -\frac{3d^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - (2ex^3 - dx)\sqrt{ex^2+d}\sqrt{-e}}{12d^2x^4}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x, algorithm="fricas")`output `-1/12*(3*d^2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - (2*e*x^3 - d*x)*sqrt(e*x^2 + d)*sqrt(-e))/(d^2*x^4)`**Sympy [A] (verification not implemented)**

Time = 2.27 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.98

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = -\frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{4x^4} - \frac{\sqrt{e}\sqrt{-e}\sqrt{\frac{d}{ex^2} + 1}}{12dx^2} + \frac{e^{\frac{3}{2}}\sqrt{-e}\sqrt{\frac{d}{ex^2} + 1}}{6d^2}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**5,x)`output `-atan(x*sqrt(-e)/sqrt(d + e*x**2))/(4*x**4) - sqrt(e)*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(12*d*x**2) + e**(3/2)*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(6*d**2)`**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.80

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \frac{\sqrt{ex^2+d}\sqrt{-e}}{4d^2x} - \frac{(ex^2+d)^{\frac{3}{2}}\sqrt{-e}}{12d^2x^3} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{4x^4}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x, algorithm="maxima")`

output

```
1/4*sqrt(e*x^2 + d)*sqrt(-e)*e/(d^2*x) - 1/12*(e*x^2 + d)^(3/2)*sqrt(-e)/(d^2*x^3) - 1/4*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^4
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. $2(67) = 134$.

Time = 0.24 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.33

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = -\frac{\left(e^3 + \frac{9(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^2}{ex^2}\right)e^6x^3}{96(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^3d^2|e|} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{4x^4} + \frac{\frac{9(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)d^4e^3}{x} + \frac{(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^3d^4}{ex^3}}{96d^6e^2|e|}$$

input

```
integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x, algorithm="giac")
```

output

```
-1/96*(e^3 + 9*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^2/(e*x^2))*e^6*x^3/((sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^3*d^2*abs(e)) - 1/4*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^4 + 1/96*(9*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))*d^4*e^3/x + (sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^3*d^4/(e*x^3))/(d^6*e^2*abs(e))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^5} dx$$

input

```
int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^5,x
```

output

```
int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^5, x
```


Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.15

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^5} dx$$

$$= \frac{-3\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d^2 - \sqrt{e}\sqrt{ex^2+d}dix + 2\sqrt{e}\sqrt{ex^2+d}ei x^3 - 2e^2i x^4}{12d^2x^4}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^5,x)`output `(- 3*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d**2 - sqrt(e)*sqrt(d + e*x**2)*d*i*x + 2*sqrt(e)*sqrt(d + e*x**2)*e*i*x**3 - 2*e**2*i*x**4)/(12*d**2*x**4)`

3.9
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

Optimal result	137
Mathematica [A] (verified)	137
Rubi [A] (verified)	138
Maple [A] (verified)	139
Fricas [A] (verification not implemented)	140
Sympy [B] (verification not implemented)	141
Maxima [A] (verification not implemented)	142
Giac [B] (verification not implemented)	142
Mupad [F(-1)]	143
Reduce [B] (verification not implemented)	143

Optimal result

Integrand size = 25, antiderivative size = 113

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{30dx^5} - \frac{2(-e)^{3/2}\sqrt{d+ex^2}}{45d^2x^3} - \frac{4(-e)^{5/2}\sqrt{d+ex^2}}{45d^3x} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

output
$$-1/30*(-e)^{(1/2)}*(e*x^2+d)^{(1/2)}/d/x^5-2/45*(-e)^{(3/2)}*(e*x^2+d)^{(1/2)}/d^2/x^3-4/45*(-e)^{(5/2)}*(e*x^2+d)^{(1/2)}/d^3/x-1/6*\arctan((-e)^{(1/2)}*x/(e*x^2+d)^{(1/2)})/x^6$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.69

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \frac{\sqrt{-ex}\sqrt{d+ex^2}(-3d^2+4dex^2-8e^2x^4)-15d^3\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{90d^3x^6}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^7,x]`

output `(Sqrt[-e]*x*Sqrt[d + e*x^2]*(-3*d^2 + 4*d*e*x^2 - 8*e^2*x^4) - 15*d^3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(90*d^3*x^6)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5674, 245, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

$$\downarrow 5674$$

$$\frac{1}{6}\sqrt{-e} \int \frac{1}{x^6\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

$$\downarrow 245$$

$$\frac{1}{6}\sqrt{-e} \left(-\frac{4e \int \frac{1}{x^4\sqrt{ex^2+d}} dx}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

$$\downarrow 245$$

$$\frac{1}{6}\sqrt{-e} \left(-\frac{4e \left(-\frac{2e \int \frac{1}{x^2\sqrt{ex^2+d}} dx}{3d} - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

$$\downarrow 242$$

$$\frac{1}{6}\sqrt{-e} \left(-\frac{4e \left(\frac{2e\sqrt{d+ex^2}}{3d^2x} - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^7,x]`

output `(Sqrt[-e]*(-1/5*Sqrt[d + e*x^2]/(d*x^5) - (4*e*(-1/3*Sqrt[d + e*x^2]/(d*x^3) + (2*e*Sqrt[d + e*x^2])/(3*d^2*x)))/(5*d))/6 - ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(6*x^6)`

Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 245 `Int[(x_)^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^2)^(p + 1)/(a*(m + 1))), x] - Simp[b*((m + 2*(p + 1) + 1)/(a*(m + 1)) Int[x^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]`

rule 5674 `Int[ArcTan[((c_.)*(x_)/Sqrt[(a_.) + (b_.)*(x_)^2])*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1)), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.04

method	result	size
default	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{6x^6} - \frac{\sqrt{-e}e\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{6d} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2+d)^{\frac{3}{2}}}{15d^2x^3}\right)}{6d}$	117
parts	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{6x^6} - \frac{\sqrt{-e}e\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{6d} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2+d)^{\frac{3}{2}}}{15d^2x^3}\right)}{6d}$	117

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x,method=_RETURNVERBOSE)`

output `-1/6*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^6-1/6*(-e)^(1/2)*e/d*(-1/3*(e*x^2+d)^(1/2)/d/x^3+2/3*e/d^2*(e*x^2+d)^(1/2)/x)+1/6*(-e)^(1/2)/d*(-1/5/d/x^5*(e*x^2+d)^(3/2)+2/15*e/d^2/x^3*(e*x^2+d)^(3/2))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.60

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

$$= -\frac{15d^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) + (8e^2x^5 - 4dex^3 + 3d^2x)\sqrt{ex^2+d}\sqrt{-e}}{90d^3x^6}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x, algorithm="fricas")`

output `-1/90*(15*d^3*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + (8*e^2*x^5 - 4*d*e*x^3 + 3*d^2*x)*sqrt(e*x^2 + d)*sqrt(-e))/(d^3*x^6)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 352 vs. $2(102) = 204$.

Time = 2.91 (sec) , antiderivative size = 352, normalized size of antiderivative = 3.12

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = -\frac{d^4 e^{\frac{9}{2}} \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{30d^5 e^4 x^4 + 60d^4 e^5 x^6 + 30d^3 e^6 x^8} - \frac{d^3 e^{\frac{11}{2}} x^2 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{45d^5 e^4 x^4 + 90d^4 e^5 x^6 + 45d^3 e^6 x^8} - \frac{d^2 e^{\frac{13}{2}} x^4 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{30d^5 e^4 x^4 + 60d^4 e^5 x^6 + 30d^3 e^6 x^8} - \frac{2de^{\frac{15}{2}} x^6 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{15d^5 e^4 x^4 + 30d^4 e^5 x^6 + 15d^3 e^6 x^8} - \frac{4e^{\frac{17}{2}} x^8 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{45d^5 e^4 x^4 + 90d^4 e^5 x^6 + 45d^3 e^6 x^8} - \frac{\arctan\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{6x^6}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**7,x)`

output `-d**4*e**(9/2)*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(30*d**5*e**4*x**4 + 60*d**4*e**5*x**6 + 30*d**3*e**6*x**8) - d**3*e**(11/2)*x**2*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(45*d**5*e**4*x**4 + 90*d**4*e**5*x**6 + 45*d**3*e**6*x**8) - d**2*e**(13/2)*x**4*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(30*d**5*e**4*x**4 + 60*d**4*e**5*x**6 + 30*d**3*e**6*x**8) - 2*d*e**(15/2)*x**6*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(15*d**5*e**4*x**4 + 30*d**4*e**5*x**6 + 15*d**3*e**6*x**8) - 4*e**(17/2)*x**8*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(45*d**5*e**4*x**4 + 90*d**4*e**5*x**6 + 45*d**3*e**6*x**8) - atan(x*sqrt(-e)/sqrt(d + e*x**2))/(6*x**6)`

Maxima [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.96

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = -\frac{(2e^2x^4 + dex^2 - d^2)\sqrt{-e}}{18\sqrt{ex^2+d}d^3x^3} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{6x^6} + \frac{(2e^2x^4 - dex^2 - 3d^2)\sqrt{ex^2+d}\sqrt{-e}}{90d^3x^5}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x, algorithm="maxima")`

output `-1/18*(2*e^2*x^4 + d*e*x^2 - d^2)*sqrt(-e)*e/(sqrt(e*x^2 + d)*d^3*x^3) - 1/6*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^6 + 1/90*(2*e^2*x^4 - d*e*x^2 - 3*d^2)*sqrt(e*x^2 + d)*sqrt(-e)/(d^3*x^5)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. 2(89) = 178.

Time = 0.24 (sec) , antiderivative size = 275, normalized size of antiderivative = 2.43

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \frac{\left(3e^4 + \frac{25(\sqrt{-dee} + \sqrt{-e^2x^2 - de|e|})^2}{x^2} + \frac{150(\sqrt{-dee} + \sqrt{-e^2x^2 - de|e|})^4}{e^4x^4}\right)e^{10}x^5}{2880(\sqrt{-dee} + \sqrt{-e^2x^2 - de|e|})^5d^3|e|} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{6x^6} - \frac{\frac{150(\sqrt{-dee} + \sqrt{-e^2x^2 - de|e|})d^{12}e^6}{x} + \frac{25(\sqrt{-dee} + \sqrt{-e^2x^2 - de|e|})^3d^{12}e^2}{x^3} + \frac{3(\sqrt{-dee} + \sqrt{-e^2x^2 - de|e|})^5d^{12}}{e^2x^5}}{2880d^{15}e^4|e|}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x, algorithm="giac")`

output

```
1/2880*(3*e^4 + 25*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^2/x^2 + 15
0*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^4/(e^4*x^4))*e^10*x^5/((sqr
t(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^5*d^3*abs(e)) - 1/6*arctan(sqrt(-
e)*x/sqrt(e*x^2 + d))/x^6 - 1/2880*(150*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*
e)*abs(e))*d^12*e^6/x + 25*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^3*
d^12*e^2/x^3 + 3*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^5*d^12/(e^2*
x^5))/(d^15*e^4*abs(e))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^7} dx$$

input

```
int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^7,x)
```

output

```
int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^7, x)
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.06

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^7} dx$$

$$= \frac{-15 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d^3 - 3\sqrt{e}\sqrt{ex^2+d}d^2ix + 4\sqrt{e}\sqrt{ex^2+d}deix^3 - 8\sqrt{e}\sqrt{ex^2+d}e^2ix^5 + 8e^3ix^6}{90d^3x^6}$$

input

```
int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^7,x)
```

output

```
( - 15*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e
x**2)*x + d + e*x**2))*d**3 - 3*sqrt(e)*sqrt(d + e*x**2)*d**2*i*x + 4*sqrt
(e)*sqrt(d + e*x**2)*d*e*i*x**3 - 8*sqrt(e)*sqrt(d + e*x**2)*e**2*i*x**5 +
8*e**3*i*x**6)/(90*d**3*x**6)
```


3.10
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$$

Optimal result	144
Mathematica [A] (verified)	144
Rubi [A] (verified)	145
Maple [A] (verified)	147
Fricas [A] (verification not implemented)	147
Sympy [B] (verification not implemented)	148
Maxima [A] (verification not implemented)	149
Giac [B] (verification not implemented)	150
Mupad [F(-1)]	151
Reduce [B] (verification not implemented)	151

Optimal result

Integrand size = 25, antiderivative size = 141

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{56dx^7} - \frac{3(-e)^{3/2}\sqrt{d+ex^2}}{140d^2x^5} - \frac{(-e)^{5/2}\sqrt{d+ex^2}}{35d^3x^3} - \frac{2(-e)^{7/2}\sqrt{d+ex^2}}{35d^4x} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

output `-1/56*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^7-3/140*(-e)^(3/2)*(e*x^2+d)^(1/2)/d^2/x^5-1/35*(-e)^(5/2)*(e*x^2+d)^(1/2)/d^3/x^3-2/35*(-e)^(7/2)*(e*x^2+d)^(1/2)/d^4/x-1/8*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^8`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.63

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \frac{\sqrt{-ex}\sqrt{d+ex^2}(-5d^3 + 6d^2ex^2 - 8de^2x^4 + 16e^3x^6) - 35d^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{280d^4x^8}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^9,x]`

output `(Sqrt[-e]*x*Sqrt[d + e*x^2]*(-5*d^3 + 6*d^2*e*x^2 - 8*d*e^2*x^4 + 16*e^3*x^6) - 35*d^4*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(280*d^4*x^8)`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5674, 245, 245, 245, 242}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$$

$$\downarrow 5674$$

$$\frac{1}{8}\sqrt{-e} \int \frac{1}{x^8\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

$$\downarrow 245$$

$$\frac{1}{8}\sqrt{-e} \left(-\frac{6e \int \frac{1}{x^6\sqrt{ex^2+d}} dx}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

$$\downarrow 245$$

$$\frac{1}{8}\sqrt{-e} \left(-\frac{6e \left(-\frac{4e \int \frac{1}{x^4\sqrt{ex^2+d}} dx}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right)}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

$$\downarrow 245$$

$$\frac{1}{8}\sqrt{-e} \left(-\frac{6e \left(-\frac{4e \left(-\frac{2e \int \frac{1}{x^2 \sqrt{ex^2+d}} dx - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right)}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{8x^8} \right)$$

↓ 242

$$\frac{1}{8}\sqrt{-e} \left(-\frac{6e \left(-\frac{4e \left(\frac{2e\sqrt{d+ex^2}}{3d^2x} - \frac{\sqrt{d+ex^2}}{3dx^3} \right)}{5d} - \frac{\sqrt{d+ex^2}}{5dx^5} \right)}{7d} - \frac{\sqrt{d+ex^2}}{7dx^7} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{8x^8} \right)$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^9,x]`

output `(Sqrt[-e]*(-1/7*Sqrt[d + e*x^2]/(d*x^7) - (6*e*(-1/5*Sqrt[d + e*x^2]/(d*x^5) - (4*e*(-1/3*Sqrt[d + e*x^2]/(d*x^3) + (2*e*Sqrt[d + e*x^2]/(3*d^2*x)))/(5*d)))/(7*d)))/8 - ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(8*x^8)`

Defintions of rubi rules used

rule 242 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, p}, x] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 245 `Int[(x_)^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^2)^(p + 1)/(a*(m + 1))), x] - Simp[b*((m + 2*(p + 1) + 1)/(a*(m + 1)) Int[x^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, m, p}, x] && ILtQ[Simplify[(m + 1)/2 + p + 1], 0] && NeQ[m, -1]`

rule 5674

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_S
ymbol] :> Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.17

method	result
default	$-\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{8x^8} - \frac{\sqrt{-e}e\left(-\frac{\sqrt{ex^2+d}}{5dx^5} - \frac{4e\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{5d}\right)}{8d} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{7dx^7} - \frac{4e\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2)}{15d^2}\right)}{7d}\right)}{8d}$
parts	$-\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{8x^8} - \frac{\sqrt{-e}e\left(-\frac{\sqrt{ex^2+d}}{5dx^5} - \frac{4e\left(-\frac{\sqrt{ex^2+d}}{3dx^3} + \frac{2e\sqrt{ex^2+d}}{3d^2x}\right)}{5d}\right)}{8d} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{7dx^7} - \frac{4e\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{5dx^5} + \frac{2e(ex^2)}{15d^2}\right)}{7d}\right)}{8d}$

input

```
int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x,method=_RETURNVERBOSE)
```

output

```
-1/8*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^8-1/8*(-e)^(1/2)*e/d*(-1/5*(e*
x^2+d)^(1/2)/d/x^5-4/5*e/d*(-1/3*(e*x^2+d)^(1/2)/d/x^3+2/3*e/d^2*(e*x^2+d)
^(1/2)/x))+1/8*(-e)^(1/2)/d*(-1/7/d/x^7*(e*x^2+d)^(3/2)-4/7*e/d*(-1/5/d/x^
5*(e*x^2+d)^(3/2)+2/15*e/d^2/x^3*(e*x^2+d)^(3/2)))
```

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.57

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$$

$$= -\frac{35d^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - (16e^3x^7 - 8de^2x^5 + 6d^2ex^3 - 5d^3x)\sqrt{ex^2+d}\sqrt{-e}}{280d^4x^8}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x, algorithm="fricas")`

output
$$-1/280*(35*d^4*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - (16*e^3*x^7 - 8*d*e^2*x^5 + 6*d^2*e*x^3 - 5*d^3*x)*sqrt(e*x^2 + d)*sqrt(-e))/(d^4*x^8)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 575 vs. $2(128) = 256$.

Time = 3.91 (sec) , antiderivative size = 575, normalized size of antiderivative = 4.08

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = -\frac{5d^6 e^{\frac{19}{2}} \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{280d^7 e^9 x^6 + 840d^6 e^{10} x^8 + 840d^5 e^{11} x^{10} + 280d^4 e^{12} x^{12}}$$

$$- \frac{9d^5 e^{\frac{21}{2}} x^2 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{280d^7 e^9 x^6 + 840d^6 e^{10} x^8 + 840d^5 e^{11} x^{10} + 280d^4 e^{12} x^{12}}$$

$$- \frac{5d^4 e^{\frac{23}{2}} x^4 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{280d^7 e^9 x^6 + 840d^6 e^{10} x^8 + 840d^5 e^{11} x^{10} + 280d^4 e^{12} x^{12}}$$

$$+ \frac{5d^3 e^{\frac{25}{2}} x^6 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{280d^7 e^9 x^6 + 840d^6 e^{10} x^8 + 840d^5 e^{11} x^{10} + 280d^4 e^{12} x^{12}}$$

$$+ \frac{15d^2 e^{\frac{27}{2}} x^8 \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{140d^7 e^9 x^6 + 420d^6 e^{10} x^8 + 420d^5 e^{11} x^{10} + 140d^4 e^{12} x^{12}}$$

$$+ \frac{5de^{\frac{29}{2}} x^{10} \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{35d^7 e^9 x^6 + 105d^6 e^{10} x^8 + 105d^5 e^{11} x^{10} + 35d^4 e^{12} x^{12}}$$

$$+ \frac{2e^{\frac{31}{2}} x^{12} \sqrt{-e} \sqrt{\frac{d}{ex^2} + 1}}{35d^7 e^9 x^6 + 105d^6 e^{10} x^8 + 105d^5 e^{11} x^{10} + 35d^4 e^{12} x^{12}}$$

$$- \frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{8x^8}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**9,x)`

output

```
-5*d**6*e**(19/2)*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(280*d**7*e**9*x**6 + 840*
d**6*e**10*x**8 + 840*d**5*e**11*x**10 + 280*d**4*e**12*x**12) - 9*d**5*e*
*(21/2)*x**2*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(280*d**7*e**9*x**6 + 840*d**6*
e**10*x**8 + 840*d**5*e**11*x**10 + 280*d**4*e**12*x**12) - 5*d**4*e**(23/
2)*x**4*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(280*d**7*e**9*x**6 + 840*d**6*e**10
*x**8 + 840*d**5*e**11*x**10 + 280*d**4*e**12*x**12) + 5*d**3*e**(25/2)*x*
*6*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(280*d**7*e**9*x**6 + 840*d**6*e**10*x**8
+ 840*d**5*e**11*x**10 + 280*d**4*e**12*x**12) + 15*d**2*e**(27/2)*x**8*s
qrt(-e)*sqrt(d/(e*x**2) + 1)/(140*d**7*e**9*x**6 + 420*d**6*e**10*x**8 + 4
20*d**5*e**11*x**10 + 140*d**4*e**12*x**12) + 5*d*e**(29/2)*x**10*sqrt(-e)
*sqrt(d/(e*x**2) + 1)/(35*d**7*e**9*x**6 + 105*d**6*e**10*x**8 + 105*d**5*
e**11*x**10 + 35*d**4*e**12*x**12) + 2*e**(31/2)*x**12*sqrt(-e)*sqrt(d/(e*
x**2) + 1)/(35*d**7*e**9*x**6 + 105*d**6*e**10*x**8 + 105*d**5*e**11*x**10
+ 35*d**4*e**12*x**12) - atan(x*sqrt(-e)/sqrt(d + e*x**2))/(8*x**8)
```

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.94

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \frac{(8e^3x^6 + 4de^2x^4 - d^2ex^2 + 3d^3)\sqrt{-e}}{120\sqrt{ex^2+d}d^4x^5} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{8x^8} - \frac{(8e^3x^6 - 4de^2x^4 + 3d^2ex^2 + 15d^3)\sqrt{ex^2+d}\sqrt{-e}}{840d^4x^7}$$

input

```
integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x, algorithm="maxima")
```

output

```
1/120*(8*e^3*x^6 + 4*d*e^2*x^4 - d^2*e*x^2 + 3*d^3)*sqrt(-e)*e/(sqrt(e*x^2
+ d)*d^4*x^5) - 1/8*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^8 - 1/840*(8*e^3
*x^6 - 4*d*e^2*x^4 + 3*d^2*e*x^2 + 15*d^3)*sqrt(e*x^2 + d)*sqrt(-e)/(d^4*x
^7)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 351 vs. $2(111) = 222$.

Time = 0.24 (sec) , antiderivative size = 351, normalized size of antiderivative = 2.49

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx =$$

$$\frac{\left(5e^5 + \frac{49(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^2}{x^2}e + \frac{245(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^4}{e^3x^4} + \frac{1225(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^6}{e^7x^6}\right)e^{14}x^7}{35840(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^7d^4|e|}$$

$$- \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{8x^8}$$

$$+ \frac{1225(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)d^{24}e^9}{x} + \frac{245(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^3d^{24}e^5}{x^3} + \frac{49(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^5d^{24}e}{x^5} + \frac{5(\sqrt{-dee} + \sqrt{-e^2x^2 - de}|e|)^7d^{24}}{35840d^{28}e^6|e|}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x, algorithm="giac")`

output

```
-1/35840*(5*e^5 + 49*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^2*e/x^2
+ 245*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^4/(e^3*x^4) + 1225*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^6/(e^7*x^6))*e^14*x^7/((sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^7*d^4*abs(e)) - 1/8*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^8 + 1/35840*(1225*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))*d^24*e^9/x + 245*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^3*d^24*e^5/x^3 + 49*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^5*d^24*e/x^5 + 5*(sqrt(-d*e)*e + sqrt(-e^2*x^2 - d*e)*abs(e))^7*d^24/(e^3*x^7))/(d^28*e^6*abs(e))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^9} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^9,x`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^9, x`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.01

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^9} dx$$

$$= \frac{-35 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d^4 - 5\sqrt{e}\sqrt{ex^2+d}d^3ix + 6\sqrt{e}\sqrt{ex^2+d}d^2eix^3 - 8\sqrt{e}\sqrt{ex^2+d}de^2ix^5}{280d^4x^8}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^9,x`

output `(- 35*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d**4 - 5*sqrt(e)*sqrt(d + e*x**2)*d**3*i*x + 6*sqrt(e)*sqrt(d + e*x**2)*d**2*e*i*x**3 - 8*sqrt(e)*sqrt(d + e*x**2)*d*e**2*i*x**5 + 16*sqrt(e)*sqrt(d + e*x**2)*e**3*i*x**7 - 16*e**4*i*x**8)/(280*d**4*x**8)`

3.11 $\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	152
Mathematica [A] (verified)	152
Rubi [A] (verified)	153
Maple [B] (verified)	154
Fricas [A] (verification not implemented)	155
Sympy [A] (verification not implemented)	156
Maxima [A] (verification not implemented)	156
Giac [A] (verification not implemented)	157
Mupad [F(-1)]	157
Reduce [B] (verification not implemented)	158

Optimal result

Integrand size = 25, antiderivative size = 124

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{d^3\sqrt{d+ex^2}}{7(-e)^{7/2}} - \frac{d^2(d+ex^2)^{3/2}}{7(-e)^{7/2}} + \frac{3d(d+ex^2)^{5/2}}{35(-e)^{7/2}} - \frac{(d+ex^2)^{7/2}}{49(-e)^{7/2}} + \frac{1}{7}x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

output

```
1/7*d^3*(e*x^2+d)^(1/2)/(-e)^(7/2)-1/7*d^2*(e*x^2+d)^(3/2)/(-e)^(7/2)+3/35
*d*(e*x^2+d)^(5/2)/(-e)^(7/2)-1/49*(e*x^2+d)^(7/2)/(-e)^(7/2)+1/7*x^7*arct
an((-e)^(1/2)*x/(e*x^2+d)^(1/2))
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.67

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{d+ex^2}(16d^3 - 8d^2ex^2 + 6de^2x^4 - 5e^3x^6)}{245(-e)^{7/2}} + \frac{1}{7}x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

input

```
Integrate[x^6*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]
```

output

$$\frac{(\sqrt{d + ex^2} * (16d^3 - 8d^2 * ex^2 + 6d * e^2 * x^4 - 5e^3 * x^6)) / (245 * (-e)^{(7/2)}) + (x^7 * \text{ArcTan}[(\sqrt{-e} * x) / \sqrt{d + ex^2}])}{7}$$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.94, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5674, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{5674} \\ & \frac{1}{7} x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{7} \sqrt{-e} \int \frac{x^7}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{7} x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{14} \sqrt{-e} \int \frac{x^6}{\sqrt{ex^2+d}} dx^2 \\ & \quad \downarrow \text{53} \\ & \frac{1}{7} x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \\ & \frac{1}{14} \sqrt{-e} \int \left(-\frac{d^3}{e^3 \sqrt{ex^2+d}} + \frac{3\sqrt{ex^2+d} d^2}{e^3} - \frac{3(ex^2+d)^{3/2} d}{e^3} + \frac{(ex^2+d)^{5/2}}{e^3} \right) dx^2 \\ & \quad \downarrow \text{2009} \\ & \frac{1}{7} x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \\ & \frac{1}{14} \sqrt{-e} \left(-\frac{2d^3 \sqrt{d+ex^2}}{e^4} + \frac{2d^2 (d+ex^2)^{3/2}}{e^4} + \frac{2(d+ex^2)^{7/2}}{7e^4} - \frac{6d(d+ex^2)^{5/2}}{5e^4} \right) \end{aligned}$$

input

$$\text{Int}[x^6 * \text{ArcTan}[(\sqrt{-e} * x) / \sqrt{d + ex^2}], x]$$

output

$$-1/14*(\text{Sqrt}[-e]*((-2*d^3*\text{Sqrt}[d + e*x^2])/e^4 + (2*d^2*(d + e*x^2)^{(3/2)})/e^4 - (6*d*(d + e*x^2)^{(5/2)})/(5*e^4) + (2*(d + e*x^2)^{(7/2)})/(7*e^4))) + (x^7*\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]])/7$$
Defintions of rubi rules used

rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

rule 243

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 5674

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_S
ymbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. $2(94) = 188$.

Time = 0.03 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.86

method	result
default	$\frac{x^7 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{7} + \frac{\sqrt{-e}e \left(\frac{x^8 \sqrt{ex^2+d}}{9e} - \frac{8d \left(\frac{x^6 \sqrt{ex^2+d}}{7e} - \frac{6d \left(\frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{7e} \right)}{9e} \right)}{7d}$
parts	$\frac{x^7 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{7} + \frac{\sqrt{-e}e \left(\frac{x^8 \sqrt{ex^2+d}}{9e} - \frac{8d \left(\frac{x^6 \sqrt{ex^2+d}}{7e} - \frac{6d \left(\frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{7e} \right)}{9e} \right)}{7d}$

```
input int(x^6*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)
```

```
output 1/7*x^7*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/7*(-e)^(1/2)*e/d*(1/9*x^8/e
*(e*x^2+d)^(1/2)-8/9*d/e*(1/7*x^6/e*(e*x^2+d)^(1/2)-6/7*d/e*(1/5*x^4/e*(e*
x^2+d)^(1/2)-4/5*d/e*(1/3*x^2/e*(e*x^2+d)^(1/2)-2/3*d/e^2*(e*x^2+d)^(1/2)
)))-1/7*(-e)^(1/2)/d*(1/9*x^6*(e*x^2+d)^(3/2)/e-2/3*d/e*(1/7*x^4*(e*x^2+d)
^(3/2)/e-4/7*d/e*(1/5*x^2*(e*x^2+d)^(3/2)/e-2/15*d/e^2*(e*x^2+d)^(3/2)))
```

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.64

$$\int x^6 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right) dx = \frac{35 e^4 x^7 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) - (5 e^3 x^6 - 6 d e^2 x^4 + 8 d^2 e x^2 - 16 d^3) \sqrt{ex^2+d} \sqrt{-e}}{245 e^4}$$

input `integrate(x^6*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output $\frac{1}{245}*(35*e^4*x^7*arctan(\sqrt{-e}*x/\sqrt{e*x^2 + d}) - (5*e^3*x^6 - 6*d*e^2*x^4 + 8*d^2*e*x^2 - 16*d^3)*\sqrt{e*x^2 + d}*\sqrt{-e})/e^4$

Sympy [A] (verification not implemented)

Time = 1.65 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.10

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \begin{cases} \frac{16d^3\sqrt{-e}\sqrt{d+ex^2}}{245e^4} - \frac{8d^2x^2\sqrt{-e}\sqrt{d+ex^2}}{245e^3} + \frac{6dx^4\sqrt{-e}\sqrt{d+ex^2}}{245e^2} + \frac{x^7 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{7} - \frac{x^6\sqrt{-e}\sqrt{d+ex^2}}{49e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x**6*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Piecewise((16*d**3*sqrt(-e)*sqrt(d + e*x**2)/(245*e**4) - 8*d**2*x**2*sqrt(-e)*sqrt(d + e*x**2)/(245*e**3) + 6*d*x**4*sqrt(-e)*sqrt(d + e*x**2)/(245*e**2) + x**7*atan(x*sqrt(-e)/sqrt(d + e*x**2))/7 - x**6*sqrt(-e)*sqrt(d + e*x**2)/(49*e), Ne(e, 0)), (0, True))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.35

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{7} x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)$$

$$- \frac{\left(35(ex^2+d)^{\frac{9}{2}} - 135(ex^2+d)^{\frac{7}{2}}d + 189(ex^2+d)^{\frac{5}{2}}d^2 - 105(ex^2+d)^{\frac{3}{2}}d^3\right)\sqrt{-e}}{2205de^4}$$

$$+ \frac{\left(35(ex^2+d)^{\frac{9}{2}} - 180(ex^2+d)^{\frac{7}{2}}d + 378(ex^2+d)^{\frac{5}{2}}d^2 - 420(ex^2+d)^{\frac{3}{2}}d^3 + 315\sqrt{ex^2+dd^4}\right)\sqrt{-e}}{2205de^4}$$

input `integrate(x^6*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output

```
1/7*x^7*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/2205*(35*(e*x^2 + d)^(9/2)
- 135*(e*x^2 + d)^(7/2)*d + 189*(e*x^2 + d)^(5/2)*d^2 - 105*(e*x^2 + d)^(3
/2)*d^3)*sqrt(-e)/(d*e^4) + 1/2205*(35*(e*x^2 + d)^(9/2) - 180*(e*x^2 + d)
^(7/2)*d + 378*(e*x^2 + d)^(5/2)*d^2 - 420*(e*x^2 + d)^(3/2)*d^3 + 315*sqr
t(e*x^2 + d)*d^4)*sqrt(-e)/(d*e^4)
```

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.10

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{7} x^7 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) + \frac{\sqrt{-e^2x^2-d}ed^3}{7e^4} + \frac{35(-e^2x^2-de)^{\frac{3}{2}}d^2e^2 + 21(e^2x^2+de)^2\sqrt{-e^2x^2-d}ede - 5(e^2x^2+de)^3\sqrt{-e^2x^2-de}}{245e^7}$$

input

```
integrate(x^6*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
1/7*x^7*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 1/7*sqrt(-e^2*x^2 - d*e)*d^3/
e^4 + 1/245*(35*(-e^2*x^2 - d*e)^(3/2)*d^2*e^2 + 21*(e^2*x^2 + d*e)^2*sqrt
(-e^2*x^2 - d*e)*d*e - 5*(e^2*x^2 + d*e)^3*sqrt(-e^2*x^2 - d*e))/e^7
```

Mupad [F(-1)]

Timed out.

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^6 \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input

```
int(x^6*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)),x)
```

output

```
int(x^6*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)), x)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.06

$$\int x^6 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{35 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) e^4 x^7 + 16\sqrt{e}\sqrt{ex^2+d}d^3i - 8\sqrt{e}\sqrt{ex^2+d}d^2ei x^2 + 6\sqrt{e}\sqrt{ex^2+d}de^2i x^4}{245e^4}$$

input

```
int(x^6*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)
```

output

```
(35*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e**4*x**7 + 16*sqrt(e)*sqrt(d + e*x**2)*d**3*i - 8*sqrt(e)*sqrt(d + e*x**2)*d**2*e*i*x**2 + 6*sqrt(e)*sqrt(d + e*x**2)*d*e**2*i*x**4 - 5*sqrt(e)*sqrt(d + e*x**2)*e**3*i*x**6)/(245*e**4)
```

3.12 $\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	159
Mathematica [A] (verified)	159
Rubi [A] (verified)	160
Maple [B] (verified)	161
Fricas [A] (verification not implemented)	162
Sympy [A] (verification not implemented)	162
Maxima [A] (verification not implemented)	163
Giac [A] (verification not implemented)	163
Mupad [F(-1)]	164
Reduce [B] (verification not implemented)	164

Optimal result

Integrand size = 25, antiderivative size = 99

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{d^2\sqrt{d+ex^2}}{5(-e)^{5/2}} - \frac{2d(d+ex^2)^{3/2}}{15(-e)^{5/2}} + \frac{(d+ex^2)^{5/2}}{25(-e)^{5/2}} + \frac{1}{5}x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

output `1/5*d^2*(e*x^2+d)^(1/2)/(-e)^(5/2)-2/15*d*(e*x^2+d)^(3/2)/(-e)^(5/2)+1/25*(e*x^2+d)^(5/2)/(-e)^(5/2)+1/5*x^5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.73

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{d+ex^2}(8d^2 - 4dex^2 + 3e^2x^4)}{75(-e)^{5/2}} + \frac{1}{5}x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

input `Integrate[x^4*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output

```
(Sqrt[d + e*x^2]*(8*d^2 - 4*d*e*x^2 + 3*e^2*x^4))/(75*(-e)^(5/2)) + (x^5*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/5
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5674, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow \text{5674} \\
 & \frac{1}{5}x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{5}\sqrt{-e} \int \frac{x^5}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{5}x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{10}\sqrt{-e} \int \frac{x^4}{\sqrt{ex^2+d}} dx^2 \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{5}x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{10}\sqrt{-e} \int \left(\frac{d^2}{e^2\sqrt{ex^2+d}} - \frac{2\sqrt{ex^2+d}d}{e^2} + \frac{(ex^2+d)^{3/2}}{e^2} \right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{5}x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{10}\sqrt{-e} \left(\frac{2d^2\sqrt{d+ex^2}}{e^3} + \frac{2(d+ex^2)^{5/2}}{5e^3} - \frac{4d(d+ex^2)^{3/2}}{3e^3} \right)
 \end{aligned}$$

input

```
Int[x^4*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
-1/10*(Sqrt[-e]*((2*d^2*Sqrt[d + e*x^2])/e^3 - (4*d*(d + e*x^2)^(3/2))/(3*e^3) + (2*(d + e*x^2)^(5/2))/(5*e^3))) + (x^5*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/5
```

Defintions of rubi rules used

```
rule 53 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5674 Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_S
ymbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(75) = 150.

Time = 0.02 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.85

method	result
default	$\frac{x^5 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{5} + \frac{\sqrt{-e}e \left(\frac{x^6 \sqrt{ex^2+d}}{7e} - \frac{6d \left(\frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{7e} \right)}{5d} - \frac{\sqrt{-e} \left(\frac{x^4 (ex^2+d)^{\frac{3}{2}}}{7e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{5d}$
parts	$\frac{x^5 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{5} + \frac{\sqrt{-e}e \left(\frac{x^6 \sqrt{ex^2+d}}{7e} - \frac{6d \left(\frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{7e} \right)}{5d} - \frac{\sqrt{-e} \left(\frac{x^4 (ex^2+d)^{\frac{3}{2}}}{7e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{5d}$

input `int(x^4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output `1/5*x^5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/5*(-e)^(1/2)*e/d*(1/7*x^6/e*(e*x^2+d)^(1/2)-6/7*d/e*(1/5*x^4/e*(e*x^2+d)^(1/2)-4/5*d/e*(1/3*x^2/e*(e*x^2+d)^(1/2)-2/3*d/e^2*(e*x^2+d)^(1/2))))-1/5*(-e)^(1/2)/d*(1/7*x^4*(e*x^2+d)^(3/2)/e-4/7*d/e*(1/5*x^2*(e*x^2+d)^(3/2)/e-2/15*d/e^2*(e*x^2+d)^(3/2))`

Fricas [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.69

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{15e^3x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - (3e^2x^4 - 4dex^2 + 8d^2)\sqrt{ex^2+d}\sqrt{-e}}{75e^3}$$

input `integrate(x^4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `1/75*(15*e^3*x^5*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - (3*e^2*x^4 - 4*d*e*x^2 + 8*d^2)*sqrt(e*x^2 + d)*sqrt(-e))/e^3`

Sympy [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.06

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \begin{cases} -\frac{8d^2\sqrt{-e}\sqrt{d+ex^2}}{75e^3} + \frac{4dx^2\sqrt{-e}\sqrt{d+ex^2}}{75e^2} + \frac{x^5 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{5} - \frac{x^4\sqrt{-e}\sqrt{d+ex^2}}{25e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(x**4*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output

```
Piecewise((-8*d**2*sqrt(-e)*sqrt(d + e*x**2)/(75*e**3) + 4*d*x**2*sqrt(-e)*sqrt(d + e*x**2)/(75*e**2) + x**5*atan(x*sqrt(-e)/sqrt(d + e*x**2))/5 - x**4*sqrt(-e)*sqrt(d + e*x**2)/(25*e), Ne(e, 0)), (0, True))
```

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.40

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{1}{5} x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)$$

$$- \frac{\left(15(ex^2+d)^{\frac{7}{2}} - 42(ex^2+d)^{\frac{5}{2}}d + 35(ex^2+d)^{\frac{3}{2}}d^2\right)\sqrt{-e}}{525de^3}$$

$$+ \frac{\left(5(ex^2+d)^{\frac{7}{2}} - 21(ex^2+d)^{\frac{5}{2}}d + 35(ex^2+d)^{\frac{3}{2}}d^2 - 35\sqrt{ex^2+dd^3}\right)\sqrt{-e}}{175de^3}$$

input

```
integrate(x^4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
1/5*x^5*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/525*(15*(e*x^2 + d)^(7/2) - 42*(e*x^2 + d)^(5/2)*d + 35*(e*x^2 + d)^(3/2)*d^2)*sqrt(-e)/(d*e^3) + 1/175*(5*(e*x^2 + d)^(7/2) - 21*(e*x^2 + d)^(5/2)*d + 35*(e*x^2 + d)^(3/2)*d^2 - 35*sqrt(e*x^2 + d)*d^3)*sqrt(-e)/(d*e^3)
```

Giac [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.02

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{5} x^5 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \frac{\sqrt{-e^2x^2 - ded^2}}{5e^3}$$

$$- \frac{10(-e^2x^2 - de)^{\frac{3}{2}}de + 3(e^2x^2 + de)^2\sqrt{-e^2x^2 - de}}{75e^5}$$

input

```
integrate(x^4*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")
```

output

```
1/5*x^5*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/5*sqrt(-e^2*x^2 - d*e)*d^2/
e^3 - 1/75*(10*(-e^2*x^2 - d*e)^(3/2)*d*e + 3*(e^2*x^2 + d*e)^2*sqrt(-e^2*
x^2 - d*e))/e^5
```

Mupad [F(-1)]

Timed out.

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^4 \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input

```
int(x^4*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2),x)
```

output

```
int(x^4*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2), x)
```

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.11

$$\int x^4 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{15 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) e^3 x^5 - 8\sqrt{e}\sqrt{ex^2+d}d^2i + 4\sqrt{e}\sqrt{ex^2+d}dei x^2 - 3\sqrt{e}\sqrt{ex^2+d}e^2i x^4}{75e^3}$$

input

```
int(x^4*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2),x)
```

output

```
(15*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**
2)*x + d + e*x**2))*e**3*x**5 - 8*sqrt(e)*sqrt(d + e*x**2)*d**2*i + 4*sqrt
(e)*sqrt(d + e*x**2)*d*e*i*x**2 - 3*sqrt(e)*sqrt(d + e*x**2)*e**2*i*x**4)/
(75*e**3)
```

3.13 $\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	165
Mathematica [A] (verified)	165
Rubi [A] (verified)	166
Maple [B] (verified)	167
Fricas [A] (verification not implemented)	168
Sympy [A] (verification not implemented)	168
Maxima [A] (verification not implemented)	169
Giac [A] (verification not implemented)	169
Mupad [F(-1)]	170
Reduce [B] (verification not implemented)	170

Optimal result

Integrand size = 25, antiderivative size = 74

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{d\sqrt{d+ex^2}}{3(-e)^{3/2}} - \frac{(d+ex^2)^{3/2}}{9(-e)^{3/2}} + \frac{1}{3}x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

output

```
1/3*d*(e*x^2+d)^(1/2)/(-e)^(3/2)-1/9*(e*x^2+d)^(3/2)/(-e)^(3/2)+1/3*x^3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.81

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{9} \left(\frac{(2d-ex^2)\sqrt{d+ex^2}}{(-e)^{3/2}} + 3x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) \right)$$

input

```
Integrate[x^2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
((2*d - e*x^2)*Sqrt[d + e*x^2])/(-e)^(3/2) + 3*x^3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/9
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5674, 243, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\
 & \quad \downarrow \text{5674} \\
 & \frac{1}{3}x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{3}\sqrt{-e} \int \frac{x^3}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{3}x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{-e} \int \frac{x^2}{\sqrt{ex^2+d}} dx^2 \\
 & \quad \downarrow \text{53} \\
 & \frac{1}{3}x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{-e} \int \left(\frac{\sqrt{ex^2+d}}{e} - \frac{d}{e\sqrt{ex^2+d}}\right) dx^2 \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{3}x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{1}{6}\sqrt{-e} \left(\frac{2(d+ex^2)^{3/2}}{3e^2} - \frac{2d\sqrt{d+ex^2}}{e^2}\right)
 \end{aligned}$$

input `Int[x^2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `-1/6*(Sqrt[-e]*((-2*d*Sqrt[d + e*x^2])/e^2 + (2*(d + e*x^2)^(3/2))/(3*e^2)) + (x^3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/3`

Defintions of rubi rules used

```
rule 53 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5674 Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^(m_.), x_S
ymbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(56) = 112.

Time = 0.02 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.82

method	result	size
default	$\frac{x^3 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{3} + \frac{\sqrt{-e}e \left(\frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{3d} - \frac{\sqrt{-e} \left(\frac{x^2 (ex^2+d)^{\frac{3}{2}}}{5e} - \frac{2d (ex^2+d)^{\frac{3}{2}}}{15e^2} \right)}{3d}$	135
parts	$\frac{x^3 \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{3} + \frac{\sqrt{-e}e \left(\frac{x^4 \sqrt{ex^2+d}}{5e} - \frac{4d \left(\frac{x^2 \sqrt{ex^2+d}}{3e} - \frac{2d \sqrt{ex^2+d}}{3e^2} \right)}{5e} \right)}{3d} - \frac{\sqrt{-e} \left(\frac{x^2 (ex^2+d)^{\frac{3}{2}}}{5e} - \frac{2d (ex^2+d)^{\frac{3}{2}}}{15e^2} \right)}{3d}$	135

```
input int(x^2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)
```


output

```
1/3*x^3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+1/3*(-e)^(1/2)*e/d*(1/5*x^4/e
*(e*x^2+d)^(1/2)-4/5*d/e*(1/3*x^2/e*(e*x^2+d)^(1/2)-2/3*d/e^2*(e*x^2+d)^(1
/2)))-1/3*(-e)^(1/2)/d*(1/5*x^2*(e*x^2+d)^(3/2)/e-2/15*d/e^2*(e*x^2+d)^(3/
2))
```

Fricas [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.76

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{3e^2x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \sqrt{ex^2+d}(ex^2-2d)\sqrt{-e}}{9e^2}$$

input

```
integrate(x^2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")
```

output

```
1/9*(3*e^2*x^3*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - sqrt(e*x^2 + d)*(e*x^2
- 2*d)*sqrt(-e))/e^2
```

Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.01

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \begin{cases} \frac{2d\sqrt{-e}\sqrt{d+ex^2}}{9e^2} + \frac{x^3 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{3} - \frac{x^2\sqrt{-e}\sqrt{d+ex^2}}{9e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input

```
integrate(x**2*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)
```

output

```
Piecewise((2*d*sqrt(-e)*sqrt(d + e*x**2)/(9*e**2) + x**3*atan(x*sqrt(-e)/s
qrt(d + e*x**2))/3 - x**2*sqrt(-e)*sqrt(d + e*x**2)/(9*e), Ne(e, 0)), (0,
True))
```

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.50

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{1}{3} x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \frac{\left(3(ex^2+d)^{\frac{5}{2}} - 5(ex^2+d)^{\frac{3}{2}}d\right)\sqrt{-e}}{45de^2}$$

$$+ \frac{\left(3(ex^2+d)^{\frac{5}{2}} - 10(ex^2+d)^{\frac{3}{2}}d + 15\sqrt{ex^2+dd^2}\right)\sqrt{-e}}{45de^2}$$

input `integrate(x^2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`output `1/3*x^3*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/45*(3*(e*x^2 + d)^(5/2) - 5*(e*x^2 + d)^(3/2)*d)*sqrt(-e)/(d*e^2) + 1/45*(3*(e*x^2 + d)^(5/2) - 10*(e*x^2 + d)^(3/2)*d + 15*sqrt(e*x^2 + d)*d^2)*sqrt(-e)/(d*e^2)`**Giac [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.86

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{1}{3} x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)$$

$$+ \frac{\sqrt{-e^2x^2 - ded}}{3e^2} + \frac{(-e^2x^2 - de)^{\frac{3}{2}}}{9e^3}$$

input `integrate(x^2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`output `1/3*x^3*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 1/3*sqrt(-e^2*x^2 - d*e)*d/e^2 + 1/9*(-e^2*x^2 - d*e)^(3/2)/e^3`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^2 \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^2*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2),x)`

output `int(x^2*atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.19

$$\int x^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{3 \operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) e^2 x^3 + 2\sqrt{e}\sqrt{ex^2+d}di - \sqrt{e}\sqrt{ex^2+d}ei x^2}{9e^2}$$

input `int(x^2*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2),x)`

output `(3*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e**2*x**3 + 2*sqrt(e)*sqrt(d + e*x**2)*d*i - sqrt(e)*sqrt(d + e*x**2)*e*i*x**2)/(9*e**2)`

3.14 $\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	171
Mathematica [A] (verified)	171
Rubi [A] (verified)	172
Maple [B] (verified)	173
Fricas [A] (verification not implemented)	173
Sympy [A] (verification not implemented)	174
Maxima [B] (verification not implemented)	174
Giac [A] (verification not implemented)	175
Mupad [B] (verification not implemented)	175
Reduce [B] (verification not implemented)	175

Optimal result

Integrand size = 21, antiderivative size = 43

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{d+ex^2}}{\sqrt{-e}} + x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

output $(e*x^2+d)^{(1/2)} / (-e)^{(1/2)} + x*\arctan((-e)^{(1/2)}*x / (e*x^2+d)^{(1/2)})$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{d+ex^2}}{\sqrt{-e}} + x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]], x]`

output $\text{Sqrt}[d + e*x^2] / \text{Sqrt}[-e] + x*\text{ArcTan}[(\text{Sqrt}[-e]*x) / \text{Sqrt}[d + e*x^2]]$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5670, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$\downarrow \text{5670}$$

$$x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \sqrt{-e} \int \frac{x}{\sqrt{ex^2+d}} dx$$

$$\downarrow \text{241}$$

$$x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{-e}\sqrt{d+ex^2}}{e}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `-((Sqrt[-e]*Sqrt[d + e*x^2])/e) + x*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 5670 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]], x_Symbol] := Simp[x*ArcTan[(c*x)/Sqrt[a + b*x^2]], x] - Simp[c Int[x/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c}, x] && EqQ[b + c^2, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(35) = 70.

Time = 0.02 (sec) , antiderivative size = 86, normalized size of antiderivative = 2.00

method	result	size
default	$x \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) + \frac{\sqrt{-e}e\left(\frac{x^2\sqrt{ex^2+d}}{3e} - \frac{2d\sqrt{ex^2+d}}{3e^2}\right)}{d} - \frac{\sqrt{-e}(ex^2+d)^{\frac{3}{2}}}{3de}$	86
parts	$x \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) + \frac{\sqrt{-e}e\left(\frac{x^2\sqrt{ex^2+d}}{3e} - \frac{2d\sqrt{ex^2+d}}{3e^2}\right)}{d} - \frac{\sqrt{-e}(ex^2+d)^{\frac{3}{2}}}{3de}$	86

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x,method=_RETURNVERBOSE)`

output `x*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+(-e)^(1/2)*e/d*(1/3*x^2/e*(e*x^2+d)^(1/2)-2/3*d/e^2*(e*x^2+d)^(1/2))-1/3*(-e)^(1/2)/d*(e*x^2+d)^(3/2)/e`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{ex \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \sqrt{ex^2+d}\sqrt{-e}}{e}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `(e*x*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - sqrt(e*x^2 + d)*sqrt(-e))/e`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \begin{cases} x \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt{-e}\sqrt{d+ex^2}}{e} & \text{for } e \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `Piecewise((x*atan(x*sqrt(-e)/sqrt(d + e*x**2)) - sqrt(-e)*sqrt(d + e*x**2)/e, Ne(e, 0)), (0, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(35) = 70.

Time = 0.05 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.79

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{(ex^2 + d)^{\frac{3}{2}}\sqrt{-e}}{3de} + \frac{\left((ex^2 + d)^{\frac{3}{2}} - 3\sqrt{ex^2 + d}\right)\sqrt{-e}}{3de}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `x*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 1/3*(e*x^2 + d)^(3/2)*sqrt(-e)/(d*e) + 1/3*((e*x^2 + d)^(3/2) - 3*sqrt(e*x^2 + d)*d)*sqrt(-e)/(d*e)`

Giac [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = x \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - \frac{\sqrt{-e^2x^2-de}}{e}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `x*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - sqrt(-e^2*x^2 - d*e)/e`

Mupad [B] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.81

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\sqrt{ex^2+d}}{\sqrt{-e}} + x \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `(d + e*x^2)^(1/2)/(-e)^(1/2) + x*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.49

$$\int \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) ex - \sqrt{e}\sqrt{ex^2+d}i}{e}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `(atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*e*x - sqrt(e)*sqrt(d + e*x**2)*i)/e`

3.15
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

Optimal result	176
Mathematica [C] (verified)	176
Rubi [A] (verified)	177
Maple [A] (verified)	178
Fricas [A] (verification not implemented)	179
Sympy [A] (verification not implemented)	179
Maxima [F]	180
Giac [A] (verification not implemented)	180
Mupad [F(-1)]	180
Reduce [B] (verification not implemented)	181

Optimal result

Integrand size = 25, antiderivative size = 59

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = -\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} - \frac{\sqrt{-e}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{\sqrt{d}}$$

output

`-arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x-(-e)^(1/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2))/d^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.46

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = -\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} + \frac{i\sqrt{e}\log\left(\frac{2i\sqrt{d}}{\sqrt{ex}} - \frac{2\sqrt{-e}\sqrt{d+ex^2}}{ex}\right)}{\sqrt{d}}$$

input

`Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^2,x]`

output

$$-(\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]]/x) + (I*\text{Sqrt}[e]*\text{Log}[(2*I)*\text{Sqrt}[d]) / (\text{Sqrt}[e]*x) - (2*\text{Sqrt}[-e]*\text{Sqrt}[d + e*x^2])/(e*x))/\text{Sqrt}[d]$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5674, 243, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

$$\downarrow 5674$$

$$\sqrt{-e} \int \frac{1}{x\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x}$$

$$\downarrow 243$$

$$\frac{1}{2}\sqrt{-e} \int \frac{1}{x^2\sqrt{ex^2+d}} dx^2 - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x}$$

$$\downarrow 73$$

$$\frac{\sqrt{-e} \int \frac{1}{\frac{x^4-d}{e}-\frac{d}{e}} d\sqrt{ex^2+d}}{e} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x}$$

$$\downarrow 221$$

$$-\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x} - \frac{\sqrt{-e}\text{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{\sqrt{d}}$$

input

$$\text{Int}[\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]]/x^2,x]$$

output

$$-(\text{ArcTan}[(\text{Sqrt}[-e]*x)/\text{Sqrt}[d + e*x^2]]/x) - (\text{Sqrt}[-e]*\text{ArcTanh}[\text{Sqrt}[d + e*x^2]/\text{Sqrt}[d]])/\text{Sqrt}[d]$$

Definitions of rubi rules used

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 243

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
ntegerQ[(m - 1)/2]
```

rule 5674

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_S
ymbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x
] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ
[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.53

method	result	size
default	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{x} - \frac{\sqrt{-e}\sqrt{ex^2+d}}{d} + \frac{\sqrt{-e}\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{d}$	90
parts	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{x} - \frac{\sqrt{-e}\sqrt{ex^2+d}}{d} + \frac{\sqrt{-e}\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{d}$	90

input

```
int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x,method=_RETURNVERBOSE)
```

output

```
-arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x-(-e)^(1/2)/d*(e*x^2+d)^(1/2)+(-e)^(
1/2)/d*((e*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(e*x^2+d)^(1/2))/x))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 148, normalized size of antiderivative = 2.51

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

$$= \left[\frac{x\sqrt{-\frac{e}{d}} \log\left(-\frac{e^2x^2+2\sqrt{ex^2+d}\sqrt{-e}\sqrt{-\frac{e}{d}}+2de}{x^2}\right) - 2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{2x}, \right.$$

$$\left. - \frac{x\sqrt{\frac{e}{d}} \arctan\left(\frac{\sqrt{ex^2+d}\sqrt{-e}\sqrt{\frac{e}{d}}}{e^2x^2+de}\right) + \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x} \right]$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x, algorithm="fricas")`

output `[1/2*(x*sqrt(-e/d)*log(-(e^2*x^2 + 2*sqrt(e*x^2 + d)*d*sqrt(-e)*sqrt(-e/d) + 2*d*e)/x^2) - 2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/x, -(x*sqrt(e/d)*arctan(sqrt(e*x^2 + d)*d*sqrt(-e)*sqrt(e/d)/(e^2*x^2 + d*e)) + arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/x]`

Sympy [A] (verification not implemented)

Time = 2.08 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.12

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \frac{\sqrt{-e} \left(\begin{cases} \frac{2 \operatorname{atan}\left(\frac{\sqrt{d+ex^2}}{\sqrt{-d}}\right)}{\sqrt{-d}} & \text{for } e \neq 0 \\ \frac{\log(x^2)}{\sqrt{d}} & \text{otherwise} \end{cases} \right)}{2} - \frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{x}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**2,x)`

output `sqrt(-e)*Piecewise((2*atan(sqrt(d + e*x**2)/sqrt(-d))/sqrt(-d), Ne(e, 0)), (log(x**2)/sqrt(d), True))/2 - atan(x*sqrt(-e)/sqrt(d + e*x**2))/x`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x, algorithm="maxima")`

output `(d*sqrt(-e)*x*integrate(-sqrt(e*x^2 + d)/(e^2*x^5 + d*e*x^3 - (e*x^3 + d*x)*(e*x^2 + d)), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)))/x`

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = -\frac{e \arctan\left(\frac{\sqrt{-e^2x^2-de}}{\sqrt{de}}\right)}{\sqrt{de}} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x, algorithm="giac")`

output `-e*arctan(sqrt(-e^2*x^2 - d*e)/sqrt(d*e))/sqrt(d*e) - arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^2,x)`

output `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^2, x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.92

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^2} dx$$

$$= \frac{-\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d + \sqrt{e}\sqrt{d}\log\left(\frac{\sqrt{ex^2+d}-\sqrt{d}+\sqrt{ex}}{\sqrt{d}}\right) ix - \sqrt{e}\sqrt{d}\log\left(\frac{\sqrt{ex^2+d}+\sqrt{d}+\sqrt{ex}}{\sqrt{d}}\right) ix}{dx}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^2,x)`output `(- atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d + sqrt(e)*sqrt(d)*log((sqrt(d + e*x**2) - sqrt(d) + sqrt(e)*x)/sqrt(d))*i*x - sqrt(e)*sqrt(d)*log((sqrt(d + e*x**2) + sqrt(d) + sqrt(e)*x)/sqrt(d))*i*x)/(d*x)`

3.16
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

Optimal result	182
Mathematica [A] (verified)	182
Rubi [A] (verified)	183
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	185
Sympy [A] (verification not implemented)	186
Maxima [F]	186
Giac [A] (verification not implemented)	187
Mupad [F(-1)]	187
Reduce [B] (verification not implemented)	187

Optimal result

Integrand size = 25, antiderivative size = 91

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{6dx^2} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3} - \frac{(-e)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{6d^{3/2}}$$

output

$$-1/6*(-e)^{(1/2)}*(e*x^2+d)^{(1/2)}/d/x^2-1/3*\arctan((-e)^{(1/2)}*x/(e*x^2+d)^{(1/2)})/x^3-1/6*(-e)^{(3/2)}*\operatorname{arctanh}((e*x^2+d)^{(1/2)}/d^{(1/2)})/d^{(3/2)}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.11

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{6dx^2} + \frac{e^{3/2}\arctan\left(\frac{\sqrt{d}\sqrt{-e}}{\sqrt{e}\sqrt{d+ex^2}}\right)}{6d^{3/2}} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

input

`Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^4,x]`

output

```
-1/6*(Sqrt[-e]*Sqrt[d + e*x^2])/(d*x^2) + (e^(3/2)*ArcTan[(Sqrt[d]*Sqrt[-e])/(Sqrt[e]*Sqrt[d + e*x^2])])/(6*d^(3/2)) - ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(3*x^3)
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5674, 243, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

↓ 5674

$$\frac{1}{3}\sqrt{-e} \int \frac{1}{x^3\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

↓ 243

$$\frac{1}{6}\sqrt{-e} \int \frac{1}{x^4\sqrt{ex^2+d}} dx^2 - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

↓ 52

$$\frac{1}{6}\sqrt{-e} \left(-\frac{e \int \frac{1}{x^2\sqrt{ex^2+d}} dx^2}{2d} - \frac{\sqrt{d+ex^2}}{dx^2} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

↓ 73

$$\frac{1}{6}\sqrt{-e} \left(-\frac{\int \frac{1}{x^4-d} d\sqrt{ex^2+d}}{d} - \frac{\sqrt{d+ex^2}}{dx^2} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

↓ 221

$$\frac{1}{6}\sqrt{-e} \left(\frac{\text{earctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{d^{3/2}} - \frac{\sqrt{d+ex^2}}{dx^2} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^3}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^4,x]`

output `-1/3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^3 + (Sqrt[-e]*(-(Sqrt[d + e*x^2]/(d*x^2)) + (e*ArcTanh[Sqrt[d + e*x^2]/Sqrt[d]])/d^(3/2)))/6`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.43

method	result	size
default	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{3x^3} + \frac{\sqrt{-e}e \ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)}{3d^{\frac{3}{2}}} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{2dx^2} + \frac{e\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{2d}\right)}{3d}$	130
parts	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{3x^3} + \frac{\sqrt{-e}e \ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)}{3d^{\frac{3}{2}}} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{2dx^2} + \frac{e\left(\sqrt{ex^2+d}-\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)\right)}{2d}\right)}{3d}$	130

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^3+1/3*(-e)^(1/2)*e/d^(3/2)*ln((2*d+2*d^(1/2)*(e*x^2+d)^(1/2))/x)+1/3*(-e)^(1/2)/d*(-1/2/d/x^2*(e*x^2+d)^(3/2)+1/2*e/d*((e*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(e*x^2+d)^(1/2))/x)))`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.18

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{x^4} dx = \left[\frac{ex^3 \sqrt{-\frac{e}{d}} \log\left(-\frac{e^2x^2-2\sqrt{ex^2+dd}\sqrt{-e}\sqrt{-\frac{e}{d}}+2de}{x^2}\right) - 2\sqrt{ex^2+d}\sqrt{-e}x - 4d \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{12 dx^3}, \frac{ex^3 \sqrt{\frac{e}{d}} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{12 dx^3} \right]$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x, algorithm="fricas")`

output

```
[1/12*(e*x^3*sqrt(-e/d)*log(-(e^2*x^2 - 2*sqrt(e*x^2 + d)*d*sqrt(-e)*sqrt(-e/d) + 2*d*e)/x^2) - 2*sqrt(e*x^2 + d)*sqrt(-e)*x - 4*d*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/(d*x^3), 1/6*(e*x^3*sqrt(e/d)*arctan(sqrt(e*x^2 + d)*d*sqrt(-e)*sqrt(e/d)/(e^2*x^2 + d*e)) - sqrt(e*x^2 + d)*sqrt(-e)*x - 2*d*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/(d*x^3)]
```

Sympy [A] (verification not implemented)

Time = 4.03 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.90

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = -\frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{3x^3} - \frac{\sqrt{e}\sqrt{-e}\sqrt{\frac{d}{ex^2}+1}}{6dx} + \frac{e\sqrt{-e}\operatorname{asinh}\left(\frac{\sqrt{d}}{\sqrt{ex}}\right)}{6d^{\frac{3}{2}}}$$

input

```
integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**4,x)
```

output

```
-atan(x*sqrt(-e)/sqrt(d + e*x**2))/(3*x**3) - sqrt(e)*sqrt(-e)*sqrt(d/(e*x**2) + 1)/(6*d*x) + e*sqrt(-e)*asinh(sqrt(d)/(sqrt(e)*x))/(6*d**(3/2))
```

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input

```
integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x, algorithm="maxima")
```

output

```
1/3*(3*d*sqrt(-e)*x^3*integrate(-1/3*sqrt(e*x^2 + d)/(e^2*x^7 + d*e*x^5 - (e*x^5 + d*x^3)*(e*x^2 + d)), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)))/x^3
```

Giac [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.98

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

$$= \frac{1}{6} e^3 \left(\frac{\arctan\left(\frac{\sqrt{-e^2x^2-de}}{\sqrt{de}}\right)}{\sqrt{dede}} - \frac{\sqrt{-e^2x^2-de}}{de^3x^2} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{3x^3}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x, algorithm="giac")`output `1/6*e^3*(arctan(sqrt(-e^2*x^2 - d*e)/sqrt(d*e))/(sqrt(d*e)*d*e) - sqrt(-e^2*x^2 - d*e)/(d*e^3*x^2)) - 1/3*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^3`**Mupad [F(-1)]**

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^4,x)`output `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^4, x)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.51

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^4} dx$$

$$= \frac{-2\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d^2 - \sqrt{e}\sqrt{ex^2+d}dix - \sqrt{e}\sqrt{d}\log\left(\frac{\sqrt{ex^2+d}-\sqrt{d}+\sqrt{ex}}{\sqrt{d}}\right) ei x^3 + \sqrt{e}\sqrt{d}\log\left(\frac{\sqrt{ex^2+d}+\sqrt{d}+\sqrt{ex}}{\sqrt{d}}\right) ei x^3}{6d^2x^3}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^4,x)`

output `(- 2*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d**2 - sqrt(e)*sqrt(d + e*x**2)*d*i*x - sqrt(e)*sqrt(d)*log((sqrt(d + e*x**2) - sqrt(d) + sqrt(e)*x)/sqrt(d))*e*i*x**3 + sqrt(e)*sqrt(d)*log((sqrt(d + e*x**2) + sqrt(d) + sqrt(e)*x)/sqrt(d))*e*i*x**3)/(6*d**2*x**3)`

3.17
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [A] (verified)	192
Fricas [A] (verification not implemented)	193
Sympy [A] (verification not implemented)	193
Maxima [F]	194
Giac [A] (verification not implemented)	194
Mupad [F(-1)]	195
Reduce [B] (verification not implemented)	195

Optimal result

Integrand size = 25, antiderivative size = 119

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = -\frac{\sqrt{-e}\sqrt{d+ex^2}}{20dx^4} - \frac{3(-e)^{3/2}\sqrt{d+ex^2}}{40d^2x^2} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5} - \frac{3(-e)^{5/2}\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{40d^{5/2}}$$

output `-1/20*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^4-3/40*(-e)^(3/2)*(e*x^2+d)^(1/2)/d^2/x^2-1/5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^5-3/40*(-e)^(5/2)*arctanh((e*x^2+d)^(1/2)/d^(1/2))/d^(5/2)`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.96

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \sqrt{-e} \left(-\frac{1}{20dx^4} + \frac{3e}{40d^2x^2} \right) \sqrt{d+ex^2} - \frac{3e^{5/2} \arctan\left(\frac{\sqrt{d}\sqrt{-e}}{\sqrt{e}\sqrt{d+ex^2}}\right)}{40d^{5/2}} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^6,x]`

output `Sqrt[-e]*(-1/20*1/(d*x^4) + (3*e)/(40*d^2*x^2))*Sqrt[d + e*x^2] - (3*e^(5/2))*ArcTan[(Sqrt[d]*Sqrt[-e])/(Sqrt[e]*Sqrt[d + e*x^2])]/(40*d^(5/2)) - ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(5*x^5)`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$, Rules used = {5674, 243, 52, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx \\
 & \quad \downarrow \text{5674} \\
 & \frac{1}{5}\sqrt{-e} \int \frac{1}{x^5\sqrt{ex^2+d}} dx - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{10}\sqrt{-e} \int \frac{1}{x^6\sqrt{ex^2+d}} dx^2 - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{10}\sqrt{-e} \left(-\frac{3e \int \frac{1}{x^4\sqrt{ex^2+d}} dx^2}{4d} - \frac{\sqrt{d+ex^2}}{2dx^4} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{10}\sqrt{-e} \left(-\frac{3e \left(-\frac{e \int \frac{1}{x^2\sqrt{ex^2+d}} dx^2}{2d} - \frac{\sqrt{d+ex^2}}{dx^2} \right)}{4d} - \frac{\sqrt{d+ex^2}}{2dx^4} \right) - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{1}{10}\sqrt{-e}\left(-\frac{3e\left(-\frac{\int\frac{1}{x^4-\frac{d}{e}}d\sqrt{ex^2+d}}{d}-\frac{\sqrt{d+ex^2}}{dx^2}\right)}{4d}-\frac{\sqrt{d+ex^2}}{2dx^4}\right)-\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5}$$

↓ 221

$$\frac{1}{10}\sqrt{-e}\left(-\frac{3e\left(\frac{e\operatorname{arctanh}\left(\frac{\sqrt{d+ex^2}}{\sqrt{d}}\right)}{d^{3/2}}-\frac{\sqrt{d+ex^2}}{dx^2}\right)}{4d}-\frac{\sqrt{d+ex^2}}{2dx^4}\right)-\frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^5}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^6,x]`

output `-1/5*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^5 + (Sqrt[-e]*(-1/2*Sqrt[d + e*x^2]/(d*x^4) - (3*e*(-(Sqrt[d + e*x^2]/(d*x^2)) + (e*ArcTanh[Sqrt[d + e*x^2]/Sqrt[d]])/d^(3/2)))/(4*d))/10`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 243 $\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[1/2 \text{ Subst}[\text{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p\}, x] \&\& \text{IntegerQ}[(m - 1)/2]$

rule 5674 $\text{Int}[\text{ArcTan}[(c_.)*(x_)]/\text{Sqrt}[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m + 1)}*(\text{ArcTan}[c*x]/\text{Sqrt}[a + b*x^2])/(d*(m + 1)), x] - \text{Simp}[c/(d*(m + 1)) \text{ Int}[(d*x)^{(m + 1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \&\& \text{EqQ}[b + c^2, 0] \&\& \text{NeQ}[m, -1]$

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.50

method	result
default	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{5x^5} - \frac{\sqrt{-e}e\left(-\frac{\sqrt{ex^2+d}}{2x^2d} + \frac{e\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)}{2d^{\frac{3}{2}}}\right)}{5d} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{4dx^4} - \frac{e\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{2dx^2} + \frac{e(\sqrt{ex^2+d}-\sqrt{-e})}{4d}\right)}{5d}\right)}{5d}$
parts	$-\frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{5x^5} - \frac{\sqrt{-e}e\left(-\frac{\sqrt{ex^2+d}}{2x^2d} + \frac{e\ln\left(\frac{2d+2\sqrt{d}\sqrt{ex^2+d}}{x}\right)}{2d^{\frac{3}{2}}}\right)}{5d} + \frac{\sqrt{-e}\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{4dx^4} - \frac{e\left(-\frac{(ex^2+d)^{\frac{3}{2}}}{2dx^2} + \frac{e(\sqrt{ex^2+d}-\sqrt{-e})}{4d}\right)}{5d}\right)}{5d}$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x,method=_RETURNVERBOSE)`

output
$$-1/5*\arctan((-e)^{(1/2)}*x/(e*x^2+d)^{(1/2)})/x^5-1/5*(-e)^{(1/2)}*e/d*(-1/2*(e*x^2+d)^{(1/2)}/x^2/d+1/2*e/d^{(3/2)}*\ln((2*d+2*d^{(1/2)}*(e*x^2+d)^{(1/2)})/x))+1/5*(-e)^{(1/2)}/d*(-1/4/d/x^4*(e*x^2+d)^{(3/2)}-1/4*e/d*(-1/2/d/x^2*(e*x^2+d)^{(3/2)}+1/2*e/d*((e*x^2+d)^{(1/2)}-d^{(1/2)}*\ln((2*d+2*d^{(1/2)}*(e*x^2+d)^{(1/2)})/x))))$$

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.92

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$$

$$= \left[\frac{3e^2x^5\sqrt{-\frac{e}{d}}\log\left(-\frac{e^2x^2+2\sqrt{ex^2+d}\sqrt{-e}\sqrt{-\frac{e}{d}}+2de}{x^2}\right) - 16d^2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) + 2(3ex^3 - 2dx)\sqrt{ex^2+d}\sqrt{-e}}{80d^2x^5} \right. \\ \left. - \frac{3e^2x^5\sqrt{\frac{e}{d}}\arctan\left(\frac{\sqrt{ex^2+d}\sqrt{-e}\sqrt{\frac{e}{d}}}{e^2x^2+de}\right) + 8d^2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - (3ex^3 - 2dx)\sqrt{ex^2+d}\sqrt{-e}}{40d^2x^5} \right]$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x, algorithm="fricas")`output `[1/80*(3*e^2*x^5*sqrt(-e/d)*log(-(e^2*x^2 + 2*sqrt(e*x^2 + d)*d*sqrt(-e)*sqrt(-e/d) + 2*d*e)/x^2) - 16*d^2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 2*(3*e*x^3 - 2*d*x)*sqrt(e*x^2 + d)*sqrt(-e))/(d^2*x^5), -1/40*(3*e^2*x^5*sqrt(e/d)*arctan(sqrt(e*x^2 + d)*d*sqrt(-e)*sqrt(e/d)/(e^2*x^2 + d*e)) + 8*d^2*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - (3*e*x^3 - 2*d*x)*sqrt(e*x^2 + d)*sqrt(-e))/(d^2*x^5)]`**Sympy [A] (verification not implemented)**

Time = 6.66 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.24

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = -\frac{\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{5x^5} - \frac{\sqrt{-e}}{20\sqrt{ex^5}\sqrt{\frac{d}{ex^2}+1}} + \frac{\sqrt{e}\sqrt{-e}}{40dx^3\sqrt{\frac{d}{ex^2}+1}} \\ + \frac{3e^{\frac{3}{2}}\sqrt{-e}}{40d^2x\sqrt{\frac{d}{ex^2}+1}} - \frac{3e^2\sqrt{-e}\operatorname{asinh}\left(\frac{\sqrt{d}}{\sqrt{ex}}\right)}{40d^{\frac{5}{2}}}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**6,x)`

output

```
-atan(x*sqrt(-e)/sqrt(d + e*x**2))/(5*x**5) - sqrt(-e)/(20*sqrt(e)*x**5*sqrt(d/(e*x**2) + 1)) + sqrt(e)*sqrt(-e)/(40*d*x**3*sqrt(d/(e*x**2) + 1)) + 3*e**(3/2)*sqrt(-e)/(40*d**2*x*sqrt(d/(e*x**2) + 1)) - 3*e**2*sqrt(-e)*asinh(sqrt(d)/(sqrt(e)*x))/(40*d**(5/2))
```

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input

```
integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x, algorithm="maxima")
```

output

```
1/5*(5*d*sqrt(-e)*x^5*integrate(-1/5*sqrt(e*x^2 + d)/(e^2*x^9 + d*e*x^7 - (e*x^7 + d*x^5)*(e*x^2 + d)), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)))/x^5
```

Giac [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.97

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = -\frac{3e^4 \arctan\left(\frac{\sqrt{-e^2x^2-de}}{\sqrt{de}}\right)}{\sqrt{ded^2}} + \frac{5\sqrt{-e^2x^2-de}de^5 + 3(-e^2x^2-de)^{\frac{3}{2}}e^4}{40e} - \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{5x^5}$$

input

```
integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x, algorithm="giac")
```

output

```
-1/40*(3*e^4*arctan(sqrt(-e^2*x^2 - d*e)/sqrt(d*e))/(sqrt(d*e)*d^2) + (5*sqrt(-e^2*x^2 - d*e)*d*e^5 + 3*(-e^2*x^2 - d*e)^(3/2)*e^4)/(d^2*e^4*x^4))/e - 1/5*arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^5
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^6,x`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^6, x`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.36

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^6} dx$$

$$= \frac{-8\operatorname{atan}\left(\frac{\sqrt{e}\sqrt{ex^2+d}ix+ei x^2}{\sqrt{e}\sqrt{ex^2+d}x+d+ex^2}\right) d^3 - 2\sqrt{e}\sqrt{ex^2+d}d^2ix + 3\sqrt{e}\sqrt{ex^2+d}dei x^3 + 3\sqrt{e}\sqrt{d}\log\left(\frac{\sqrt{ex^2+d}-\sqrt{d}}{\sqrt{d}}\right)}{40d^3x^5}$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^6,x`

output `(- 8*atan((sqrt(e)*sqrt(d + e*x**2)*i*x + e*i*x**2)/(sqrt(e)*sqrt(d + e*x**2)*x + d + e*x**2))*d**3 - 2*sqrt(e)*sqrt(d + e*x**2)*d**2*i*x + 3*sqrt(e)*sqrt(d + e*x**2)*d*e*i*x**3 + 3*sqrt(e)*sqrt(d)*log((sqrt(d + e*x**2) - sqrt(d) + sqrt(e)*x)/sqrt(d))*e**2*i*x**5 - 3*sqrt(e)*sqrt(d)*log((sqrt(d + e*x**2) + sqrt(d) + sqrt(e)*x)/sqrt(d))*e**2*i*x**5)/(40*d**3*x**5)`

3.18 $\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	196
Mathematica [C] (verified)	197
Rubi [A] (verified)	197
Maple [F]	200
Fricas [A] (verification not implemented)	200
Sympy [F(-1)]	201
Maxima [F]	201
Giac [F]	201
Mupad [F(-1)]	202
Reduce [F]	202

Optimal result

Integrand size = 27, antiderivative size = 211

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{60d^2\sqrt{x}\sqrt{d+ex^2}}{847(-e)^{5/2}} + \frac{36dx^{5/2}\sqrt{d+ex^2}}{847(-e)^{3/2}} + \frac{4x^{9/2}\sqrt{d+ex^2}}{121\sqrt{-e}}$$

$$+ \frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + \frac{30d^{11/4}\sqrt{-e}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{847e^{13/4}\sqrt{d+ex^2}}$$

output

```
60/847*d^2*x^(1/2)*(e*x^2+d)^(1/2)/(-e)^(5/2)+36/847*d*x^(5/2)*(e*x^2+d)^(1/2)/(-e)^(3/2)+4/121*x^(9/2)*(e*x^2+d)^(1/2)/(-e)^(1/2)+2/11*x^(11/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+30/847*d^(11/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(13/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 11.39 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.81

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{4\sqrt{x}\sqrt{d+ex^2}(15d^2 - 9dex^2 + 7e^2x^4)}{847(-e)^{5/2}} + \frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{60id^3\sqrt{1+\frac{d}{ex^2}}x \operatorname{EllipticF}\left(\operatorname{iarcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{847\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}(-e)^{5/2}\sqrt{d+ex^2}}$$

input `Integrate[x^(9/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(4*Sqrt[x]*Sqrt[d + e*x^2]*(15*d^2 - 9*d*e*x^2 + 7*e^2*x^4))/(847*(-e)^(5/2)) + (2*x^(11/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/11 - (((60*I)/847)*d^3*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(Sqrt[(I*Sqrt[d])/Sqrt[e]]*(-e)^(5/2)*Sqrt[d + e*x^2])`

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5674, 262, 262, 262, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

↓ 5674

$$\frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{-e} \int \frac{x^{11/2}}{\sqrt{ex^2+d}} dx$$

↓ 262

$$\begin{aligned}
 & \frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{-e} \left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d \int \frac{x^{7/2}}{\sqrt{ex^2+d}} dx}{11e} \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{-e} \left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \int \frac{x^{3/2}}{\sqrt{ex^2+d}} dx}{7e} \right)}{11e} \right) \\
 & \quad \downarrow \text{262} \\
 & \frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{-e} \left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx}{3e} \right)}{7e} \right)}{11e} \right) \\
 & \quad \downarrow \text{266} \\
 & \frac{2}{11}x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{11}\sqrt{-e} \left(\frac{2x^{9/2}\sqrt{d+ex^2}}{11e} - \frac{9d \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{2d \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3e} \right)}{7e} \right)}{11e} \right) \\
 & \quad \downarrow \text{761}
 \end{aligned}$$

$$\frac{2}{11} \sqrt{-e} \left(\frac{2x^{9/2} \sqrt{d+ex^2}}{11e} - \frac{\frac{2}{11} x^{11/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - 9d \left(\frac{2x^{5/2} \sqrt{d+ex^2}}{7e} - \frac{5d \left(\frac{2\sqrt{x} \sqrt{d+ex^2}}{3e} - \frac{d^{3/4} (\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d} + \sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)\right)}{3e^{5/4} \sqrt{d+ex^2}} \right)}{7e} \right)}{11e} \right)$$

```
input Int [x^(9/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]], x]
```

```
output (2*x^(11/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/11 - (2*Sqrt[-e]*((2*x^(9/2)*Sqrt[d + e*x^2])/(11*e) - (9*d*((2*x^(5/2)*Sqrt[d + e*x^2])/(7*e) - (5*d*((2*Sqrt[x]*Sqrt[d + e*x^2])/(3*e) - (d^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(3*e^(5/4)*Sqrt[d + e*x^2])))/(7*e)))/(11*e)))/11
```

Defintions of rubi rules used

```
rule 262 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

```
rule 266 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]
```


rule 761

```
Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

rule 5674

```
Int[ArcTan[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [F]

$$\int x^{\frac{9}{2}} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) dx$$

input

```
int(x^(9/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)
```

output

```
int(x^(9/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)
```

Fricas [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.45

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right) dx = \frac{2\left(77e^4x^{\frac{11}{2}} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) + 30d^3\sqrt{-e}\sqrt{e}\text{weierstrassPInverse}\left(-\frac{4d}{e}, 0\right)\right)}{847e^4}$$

input

```
integrate(x^(9/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")
```

output

```
2/847*(77*e^4*x^(11/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 30*d^3*sqrt(-e)*sqrt(e)*weierstrassPInverse(-4*d/e, 0, x) - 2*(7*e^3*x^4 - 9*d*e^2*x^2 + 15*d^2*e)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(x))/e^4
```

Sympy [F(-1)]

Timed out.

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input `integrate(x**(9/2)*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output Timed out

Maxima [F]

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(9/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `2/11*x^(11/2)*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - 2*d*sqrt(-e)*integrate(-1/11*x*e^(1/2*log(e*x^2 + d) + 9/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

Giac [F]

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(9/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(9/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{9/2} \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(9/2)*atan((-e)^(1/2)*x/(d + e*x^2)^(1/2)),x)`

output `int(x^(9/2)*atan((-e)^(1/2)*x/(d + e*x^2)^(1/2)), x)`

Reduce [F]

$$\int x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right) x^4 dx$$

input `int(x^(9/2)*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2))*x**4,x)`

3.19 $\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	203
Mathematica [C] (verified)	204
Rubi [A] (verified)	204
Maple [F]	206
Fricas [A] (verification not implemented)	207
Sympy [C] (verification not implemented)	207
Maxima [F]	208
Giac [F]	208
Mupad [F(-1)]	208
Reduce [F]	209

Optimal result

Integrand size = 27, antiderivative size = 181

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{20d\sqrt{x}\sqrt{d+ex^2}}{147(-e)^{3/2}} + \frac{4x^{5/2}\sqrt{d+ex^2}}{49\sqrt{-e}}$$

$$+ \frac{2}{7}x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{10d^{7/4}\sqrt{-e}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{147e^{9/4}\sqrt{d+ex^2}}$$

output

```
20/147*d*x^(1/2)*(e*x^2+d)^(1/2)/(-e)^(3/2)+4/49*x^(5/2)*(e*x^2+d)^(1/2)/(-e)^(1/2)+2/7*x^(7/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))-10/147*d^(7/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(9/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.87

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2}{147} \sqrt{x} \left(\frac{2(5d-3ex^2)\sqrt{d+ex^2}}{(-e)^{3/2}} \right. \\ \left. + 21x^3 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) \right) \\ - \frac{20id^2 \sqrt{1+\frac{d}{ex^2}} x \operatorname{EllipticF}\left(i \operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{147 \sqrt{\frac{i\sqrt{d}}{\sqrt{e}}} (-e)^{3/2} \sqrt{d+ex^2}}$$

input `Integrate[x^(5/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(2*Sqrt[x]*((2*(5*d - 3*e*x^2)*Sqrt[d + e*x^2])/(-e)^(3/2) + 21*x^3*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]))/147 - (((20*I)/147)*d^2*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/Sqrt[(I*Sqrt[d])/Sqrt[e]]*(-e)^(3/2)*Sqrt[d + e*x^2])`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$, Rules used = {5674, 262, 262, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\ \downarrow \text{5674}$$

$$\begin{aligned}
 & \frac{2}{7}x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{-e} \int \frac{x^{7/2}}{\sqrt{ex^2+d}} dx \\
 & \quad \downarrow 262 \\
 & \frac{2}{7}x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{-e} \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \int \frac{x^{3/2}}{\sqrt{ex^2+d}} dx}{7e} \right) \\
 & \quad \downarrow 262 \\
 & \frac{2}{7}x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{-e} \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx}{3e} \right)}{7e} \right) \\
 & \quad \downarrow 266 \\
 & \frac{2}{7}x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{-e} \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{2d \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3e} \right)}{7e} \right) \\
 & \quad \downarrow 761 \\
 & \frac{2}{7}x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{7}\sqrt{-e} \left(\frac{2x^{5/2}\sqrt{d+ex^2}}{7e} - \frac{5d \left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d^{3/4}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{3e^{5/4}\sqrt{d+ex^2}} \right)}{7e} \right)
 \end{aligned}$$

input `Int [x^(5/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]], x]`

output `(2*x^(7/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/7 - (2*Sqrt[-e]*((2*x^(5/2)*Sqrt[d + e*x^2])/(7*e) - (5*d*((2*Sqrt[x]*Sqrt[d + e*x^2])/(3*e) - (d^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*e^(5/4)*Sqrt[d + e*x^2])))/(7*e)))/7`

Definitions of rubi rules used

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1))) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int x^{\frac{5}{2}} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(5/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(5/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.47

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2\left(21e^3x^{7/2}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - 10d^2\sqrt{-e}\sqrt{e}\operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) - 2(3e^2x^2 - 5d)e\sqrt{ex^2+d}\sqrt{-e}\sqrt{x}\right)}{147e^3}$$

input `integrate(x^(5/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `2/147*(21*e^3*x^(7/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 10*d^2*sqrt(-e)*sqrt(e)*weierstrassPInverse(-4*d/e, 0, x) - 2*(3*e^2*x^2 - 5*d*e)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(x))/e^3`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 41.39 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.41

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{7/2} \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{7} - \frac{x^{9/2}\sqrt{-e}\Gamma\left(\frac{9}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{9}{4} \middle| \frac{ex^2e^{i\pi}}{d}\right)}{7\sqrt{d}\Gamma\left(\frac{13}{4}\right)}$$

input `integrate(x**(5/2)*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `2*x**(7/2)*atan(x*sqrt(-e)/sqrt(d + e*x**2))/7 - x**(9/2)*sqrt(-e)*gamma(9/4)*hyper((1/2, 9/4), (13/4,), e*x**2*exp_polar(I*pi)/d)/(7*sqrt(d)*gamma(13/4))`

Maxima [F]

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(5/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `2/7*x^(7/2)*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - 2*d*sqrt(-e)*integrate(-1/7*x*e^(1/2*log(e*x^2 + d) + 5/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

Giac [F]

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(5/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(5/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{5/2} \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(5/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(5/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

Reduce [F]

$$\int x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right) x^2 dx$$

input `int(x^(5/2)*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2))*x**2,x)`

3.20 $\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	210
Mathematica [C] (verified)	211
Rubi [A] (verified)	211
Maple [F]	213
Fricas [A] (verification not implemented)	213
Sympy [C] (verification not implemented)	214
Maxima [F]	214
Giac [F]	215
Mupad [F(-1)]	215
Reduce [F]	215

Optimal result

Integrand size = 27, antiderivative size = 153

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$$

$$= \frac{4\sqrt{x}\sqrt{d+ex^2}}{9\sqrt{-e}} + \frac{2}{3}x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)$$

$$+ \frac{2d^{3/4}\sqrt{-e}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{9e^{5/4}\sqrt{d+ex^2}}$$

output

```
4/9*x^(1/2)*(e*x^2+d)^(1/2)/(-e)^(1/2)+2/3*x^(3/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+2/9*d^(3/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(5/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.96

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{4\sqrt{x}\sqrt{d+ex^2}}{9\sqrt{-e}} + \frac{2}{3}x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{4id\sqrt{1+\frac{d}{ex^2}}x \operatorname{EllipticF}\left(\operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{9\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}\sqrt{-e}\sqrt{d+ex^2}}$$

input `Integrate[Sqrt[x]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(4*Sqrt[x]*Sqrt[d + e*x^2])/(9*Sqrt[-e]) + (2*x^(3/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/3 - (((4*I)/9)*d*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[-e]*Sqrt[d + e*x^2])`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$, Rules used = {5674, 262, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{5674} \\ & \frac{2}{3}x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{3}\sqrt{-e} \int \frac{x^{3/2}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \end{aligned}$$

$$\begin{aligned}
& \frac{2}{3}x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{3}\sqrt{-e}\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{d \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx}{3e}\right) \\
& \quad \downarrow \text{266} \\
& \frac{2}{3}x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{3}\sqrt{-e}\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{2d \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3e}\right) \\
& \quad \downarrow \text{761} \\
& \frac{2}{3}\sqrt{-e}\left(\frac{2\sqrt{x}\sqrt{d+ex^2}}{3e} - \frac{\frac{2}{3}x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - d^{3/4}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d} + \sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{3e^{5/4}\sqrt{d+ex^2}}\right)
\end{aligned}$$

input `Int[Sqrt[x]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(2*x^(3/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/3 - (2*Sqrt[-e]*((2*Sqrt[x]*Sqrt[d + e*x^2])/(3*e) - (d^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*e^(5/4)*Sqrt[d + e*x^2])))`

Defintions of rubi rules used

rule 262 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 5674 `Int[ArcTan[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2 + d}}\right) dx$$

input `int(x^(1/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(1/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.46

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{d + ex^2}}\right) dx$$

$$= \frac{2\left(3e^2x^{\frac{3}{2}} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) + 2d\sqrt{-e}\sqrt{e}\text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) - 2\sqrt{ex^2+d}\sqrt{-e}e\sqrt{x}\right)}{9e^2}$$

input `integrate(x^(1/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `2/9*(3*e^2*x^(3/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 2*d*sqrt(-e)*sqrt(e)*weierstrassPInverse(-4*d/e, 0, x) - 2*sqrt(e*x^2 + d)*sqrt(-e)*e*sqrt(x))/e^2`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 9.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.49

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{\frac{3}{2}} \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{3} - \frac{x^{\frac{5}{2}} \sqrt{-e} \Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{ex^2 e^{i\pi}}{d}\right)}{3\sqrt{d} \Gamma\left(\frac{9}{4}\right)}$$

input `integrate(x**(1/2)*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `2*x**(3/2)*atan(x*sqrt(-e)/sqrt(d + e*x**2))/3 - x**(5/2)*sqrt(-e)*gamma(5/4)*hyper((1/2, 5/4), (9/4,), e*x**2*exp_polar(I*pi)/d)/(3*sqrt(d)*gamma(9/4))`

Maxima [F]

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(1/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `2/3*x^(3/2)*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - 2*d*sqrt(-e)*integrate(-1/3*x*e^(1/2*log(e*x^2 + d) + 1/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

Giac [F]

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(1/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(1/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(1/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

Reduce [F]

$$\int \sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(1/2)*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)),x)`

3.21
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$$

Optimal result	216
Mathematica [C] (verified)	217
Rubi [A] (verified)	217
Maple [F]	219
Fricas [A] (verification not implemented)	219
Sympy [C] (verification not implemented)	219
Maxima [F]	220
Giac [F]	220
Mupad [F(-1)]	221
Reduce [F]	221

Optimal result

Integrand size = 27, antiderivative size = 122

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} + \frac{2\sqrt{-e}\left(\sqrt{d} + \sqrt{ex}\right) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{d}\sqrt[4]{e}\sqrt{d+ex^2}}$$

output

```
-2*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2)+2*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(1/4)/e^(1/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.94

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} + \frac{4i\sqrt{-e}\sqrt{1+\frac{d}{ex^2}}x \operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}\sqrt{d+ex^2}}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(3/2), x]`

output `(-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/Sqrt[x] + ((4*I)*Sqrt[-e]*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[d + e*x^2]`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {5674, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx$$

$$\downarrow \text{5674}$$

$$2\sqrt{-e} \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}}$$

$$\downarrow \text{266}$$

$$4\sqrt{-e} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x} - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}}$$

↓ 761

$$\frac{2\sqrt{-e}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{d}\sqrt[4]{e}\sqrt{d+ex^2}} - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(3/2),x]`

output `(-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/Sqrt[x] + (2*Sqrt[-e]*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(d^(1/4)*e^(1/4)*Sqrt[d + e*x^2])`

Defintions of rubi rules used

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{3}{2}}} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.43

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \frac{2\left(2\sqrt{-e}\sqrt{ex}\text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) - e\sqrt{x}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)\right)}{ex}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x, algorithm="fricas")`

output `2*(2*sqrt(-e)*sqrt(e)*x*weierstrassPInverse(-4*d/e, 0, x) - e*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/(e*x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 8.79 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.58

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = -\frac{2\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} + \frac{\sqrt{x}\sqrt{-e}\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{ex^2 e^{i\pi}}{d}\right)}{\sqrt{d}\Gamma\left(\frac{5}{4}\right)}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(3/2),x)`

output `-2*atan(x*sqrt(-e)/sqrt(d + e*x**2))/sqrt(x) + sqrt(x)*sqrt(-e)*gamma(1/4)
*hyper((1/4, 1/2), (5/4,), e*x**2*exp_polar(I*pi)/d)/(sqrt(d)*gamma(5/4))`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{3}{2}}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x, algorithm="maxima")`

output `2*(d*sqrt(-e)*sqrt(x)*integrate(-sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(3/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 3/2*log(x))), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d))/sqrt(x)`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{3}{2}}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^(3/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{3/2}} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2)/x^(3/2),x)`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2)/x^(3/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{3/2}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x^2} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x**2,x)`

3.22
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$$

Optimal result	222
Mathematica [C] (verified)	223
Rubi [A] (verified)	223
Maple [F]	225
Fricas [A] (verification not implemented)	225
Sympy [C] (verification not implemented)	226
Maxima [F]	226
Giac [F]	227
Mupad [F(-1)]	227
Reduce [F]	227

Optimal result

Integrand size = 27, antiderivative size = 156

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = -\frac{4\sqrt{-e}\sqrt{d+ex^2}}{15dx^{3/2}} - \frac{2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} - \frac{2\sqrt{-e}e^{3/4}\left(\sqrt{d} + \sqrt{ex}\right)\sqrt{\frac{d+ex^2}{\left(\sqrt{d}+\sqrt{ex}\right)^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{15d^{5/4}\sqrt{d+ex^2}}$$

output

```
-4/15*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^(3/2)-2/5*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2)-2/15*(-e)^(1/2)*e^(3/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(5/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.96

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = -\frac{2\left(2\sqrt{-ex}\sqrt{d+ex^2} + 3d \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)\right)}{15dx^{5/2}} + \frac{4i(-e)^{3/2}\sqrt{1+\frac{d}{ex^2}}x \operatorname{EllipticF}\left(\operatorname{iarcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{15d\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}\sqrt{d+ex^2}}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(7/2),x]`

output `(-2*(2*Sqrt[-e]*x*Sqrt[d + e*x^2] + 3*d*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]))/(15*d*x^(5/2)) + (((4*I)/15)*(-e)^(3/2)*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(d*Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[d + e*x^2])`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$, Rules used = {5674, 264, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx$$

↓ 5674

$$\frac{2}{5}\sqrt{-e} \int \frac{1}{x^{5/2}\sqrt{ex^2+d}} dx - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}}$$

$$\begin{aligned}
 & \downarrow 264 \\
 & \frac{2}{5}\sqrt{-e} \left(-\frac{e \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} \\
 & \downarrow 266 \\
 & \frac{2}{5}\sqrt{-e} \left(-\frac{2e \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} \\
 & \downarrow 761 \\
 & \frac{2}{5}\sqrt{-e} \left(-\frac{e^{3/4}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{3d^{5/4}\sqrt{d+ex^2}} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right) - \\
 & \qquad \qquad \qquad \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}}
 \end{aligned}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(7/2),x]`

output `(-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(5*x^(5/2)) + (2*Sqrt[-e]*((-2*Sqrt[d + e*x^2])/(3*d*x^(3/2)) - (e^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2]))/(3*d^(5/4)*Sqrt[d + e*x^2]))/5`

Defintions of rubi rules used

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761

```
Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

rule 5674

```
Int[ArcTan[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input

```
int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x)
```

output

```
int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.47

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \frac{2\left(2\sqrt{-e}\sqrt{ex^3}\text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) + 2\sqrt{ex^2+d}\sqrt{-ex}^{\frac{3}{2}} + 3d\sqrt{x}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)\right)}{15dx^3}$$

input

```
integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x, algorithm="fricas")
```

output

```
-2/15*(2*sqrt(-e)*sqrt(e)*x^3*weierstrassPInverse(-4*d/e, 0, x) + 2*sqrt(e*x^2 + d)*sqrt(-e)*x^(3/2) + 3*d*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/(d*x^3)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 44.00 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.50

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = -\frac{2 \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{5x^{5/2}} + \frac{\sqrt{-e}\Gamma\left(-\frac{3}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{3}{4}, \frac{1}{2} \\ \frac{1}{4} \end{matrix} \middle| \frac{ex^2 e^{i\pi}}{d} \right)}{5\sqrt{d}x^{3/2}\Gamma\left(\frac{1}{4}\right)}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(7/2), x)`

output `-2*atan(x*sqrt(-e)/sqrt(d + e*x**2))/(5*x**(5/2)) + sqrt(-e)*gamma(-3/4)*hyper((-3/4, 1/2), (1/4,), e*x**2*exp_polar(I*pi)/d)/(5*sqrt(d)*x**(3/2)*gamma(1/4))`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2), x, algorithm="maxima")`

output `2/5*(5*d*sqrt(-e)*x^(5/2)*integrate(-1/5*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(7/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 7/2*log(x))), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d))/x^(5/2)`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^(7/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{7/2}} dx$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(7/2),x)`

output `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(7/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{7/2}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x^4} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x**4,x)`

3.23
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$$

Optimal result	228
Mathematica [C] (verified)	229
Rubi [A] (verified)	229
Maple [F]	231
Fricas [A] (verification not implemented)	232
Sympy [F(-1)]	232
Maxima [F]	232
Giac [F]	233
Mupad [F(-1)]	233
Reduce [F]	234

Optimal result

Integrand size = 27, antiderivative size = 186

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = -\frac{4\sqrt{-e}\sqrt{d+ex^2}}{63dx^{7/2}} - \frac{20(-e)^{3/2}\sqrt{d+ex^2}}{189d^2x^{3/2}} - \frac{2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} + \frac{10\sqrt{-e}e^{7/4}\left(\sqrt{d} + \sqrt{ex}\right)\sqrt{\frac{d+ex^2}{\left(\sqrt{d}+\sqrt{ex}\right)^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{189d^{9/4}\sqrt{d+ex^2}}$$

output

```
-4/63*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^(7/2)-20/189*(-e)^(3/2)*(e*x^2+d)^(1/2)/d^2/x^(3/2)-2/9*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2)+10/189*(-e)^(1/2)*e^(7/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(9/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.87

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \frac{4\sqrt{-ex}\sqrt{d+ex^2}(-3d+5ex^2) - 42d^2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{189d^2x^{9/2}} + \frac{20i(-e)^{5/2}\sqrt{1+\frac{d}{ex^2}}x \operatorname{EllipticF}\left(i\operatorname{arcsinh}\left(\frac{\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}}{\sqrt{x}}\right), -1\right)}{189d^2\sqrt{\frac{i\sqrt{d}}{\sqrt{e}}}\sqrt{d+ex^2}}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(11/2),x]`

output `(4*Sqrt[-e]*x*Sqrt[d + e*x^2]*(-3*d + 5*e*x^2) - 42*d^2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(189*d^2*x^(9/2)) + (((20*I)/189)*(-e)^(5/2)*Sqrt[1 + d/(e*x^2)]*x*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(d^2*Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[d + e*x^2])`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$, Rules used = {5674, 264, 264, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx$$

↓ 5674

$$\frac{2}{9}\sqrt{-e} \int \frac{1}{x^{9/2}\sqrt{ex^2+d}} dx - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}}$$

$$\begin{aligned}
& \downarrow 264 \\
& \frac{2}{9}\sqrt{-e} \left(-\frac{5e \int \frac{1}{x^{5/2}\sqrt{ex^2+d}} dx}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} \\
& \downarrow 264 \\
& \frac{2}{9}\sqrt{-e} \left(-\frac{5e \left(-\frac{e \int \frac{1}{\sqrt{x}\sqrt{ex^2+d}} dx}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} \\
& \downarrow 266 \\
& \frac{2}{9}\sqrt{-e} \left(-\frac{5e \left(-\frac{2e \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}} \\
& \downarrow 761 \\
& \frac{2}{9}\sqrt{-e} \left(-\frac{5e \left(-\frac{e^{3/4}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{3d^{5/4}\sqrt{d+ex^2}} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{9x^{9/2}}
\end{aligned}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(11/2),x]`

output `(-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(9*x^(9/2)) + (2*Sqrt[-e]*((-2*Sqrt[d + e*x^2])/(7*d*x^(7/2)) - (5*e*((-2*Sqrt[d + e*x^2])/(3*d*x^(3/2)) - (e^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(3*d^(5/4)*Sqrt[d + e*x^2])))/(7*d)))/9`

Definitions of rubi rules used

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{11}{2}}} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.46

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \frac{2\left(10\sqrt{-e}e^{\frac{3}{2}}x^5\text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) - 21d^2\sqrt{x}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) + 2\right)}{189d^2x^5}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x, algorithm="fricas")`

output `2/189*(10*sqrt(-e)*e^(3/2)*x^5*weierstrassPInverse(-4*d/e, 0, x) - 21*d^2*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 2*(5*e*x^3 - 3*d*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(x))/(d^2*x^5)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \text{Timed out}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(11/2),x)`

output `Timed out`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{11}{2}}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x, algorithm="maxima")`

output $2/9*(9*d*\sqrt{-e}*x^{(9/2)}*\text{integrate}(-1/9*\sqrt{e*x^2 + d}*x/((e^2*x^4 + d*e*x^2)*x^{(11/2)} - (e*x^2 + d)*e^{(\log(e*x^2 + d) + 11/2*\log(x))}), x) - \text{arctan}2(\sqrt{-e}*x, \sqrt{e*x^2 + d}))/x^{(9/2)}$

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{11/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^(11/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\text{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{11/2}} dx$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(11/2),x)`

output `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(11/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{11/2}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x^6} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(11/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x**6,x)`

3.24
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$$

Optimal result	235
Mathematica [C] (verified)	236
Rubi [A] (verified)	236
Maple [F]	239
Fricas [A] (verification not implemented)	240
Sympy [F(-1)]	240
Maxima [F]	241
Giac [F]	241
Mupad [F(-1)]	241
Reduce [F]	242

Optimal result

Integrand size = 27, antiderivative size = 216

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = -\frac{4\sqrt{-e}\sqrt{d+ex^2}}{143dx^{11/2}} - \frac{36(-e)^{3/2}\sqrt{d+ex^2}}{1001d^2x^{7/2}} - \frac{60(-e)^{5/2}\sqrt{d+ex^2}}{1001d^3x^{3/2}} - \frac{2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}} - \frac{30\sqrt{-e}e^{11/4}\left(\sqrt{d} + \sqrt{ex}\right)\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{1001d^{13/4}\sqrt{d+ex^2}}$$

output

```
-4/143*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^(11/2)-36/1001*(-e)^(3/2)*(e*x^2+d)^(1/2)/d^2/x^(7/2)-60/1001*(-e)^(5/2)*(e*x^2+d)^(1/2)/d^3/x^(3/2)-2/13*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(13/2)-30/1001*(-e)^(1/2)*e^(11/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(13/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.79

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \frac{2 \left(-\frac{2\sqrt{-e}\sqrt{d+ex^2}(7d^2x-9dex^3+15e^2x^5)}{d^3} - 77 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + \frac{30i(-e)^{7/2}\sqrt{1+\frac{d}{ex^2}}x^{15/2} \operatorname{EllipticF}\left[\operatorname{ArcSinh}\left[\frac{\sqrt{d}}{\sqrt{e}}\right]/\sqrt{x}\right], -1\right)}{d^3\sqrt{d+ex^2}} \right)}{1001x^{13/2}}$$

input

```
Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(15/2), x]
```

output

```
(2*((-2*Sqrt[-e]*Sqrt[d + e*x^2]*(7*d^2*x - 9*d*e*x^3 + 15*e^2*x^5))/d^3 - 77*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] + ((30*I)*(-e)^(7/2)*Sqrt[1 + d/(e*x^2)]*x^(15/2)*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[d])/Sqrt[e]]/Sqrt[x]], -1])/(d^3*Sqrt[(I*Sqrt[d])/Sqrt[e]]*Sqrt[d + e*x^2])))/(1001*x^(13/2))
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5674, 264, 264, 264, 266, 761}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx$$

↓ 5674

$$\frac{2}{13}\sqrt{-e} \int \frac{1}{x^{13/2}\sqrt{ex^2+d}} dx - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}}$$

↓ 264

$$\begin{aligned}
 & \frac{2}{13} \sqrt{-e} \left(-\frac{9e \int \frac{1}{x^{9/2} \sqrt{ex^2+d}} dx}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \frac{2 \arctan \left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}} \right)}{13x^{13/2}} \\
 & \quad \downarrow 264 \\
 & \frac{2}{13} \sqrt{-e} \left(-\frac{9e \left(-\frac{5e \int \frac{1}{x^{5/2} \sqrt{ex^2+d}} dx}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \frac{2 \arctan \left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}} \right)}{13x^{13/2}} \\
 & \quad \downarrow 264 \\
 & \frac{2}{13} \sqrt{-e} \left(-\frac{9e \left(\frac{5e \left(-\frac{e \int \frac{1}{\sqrt{x} \sqrt{ex^2+d}} dx}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \\
 & \quad \frac{2 \arctan \left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}} \right)}{13x^{13/2}} \\
 & \quad \downarrow 266 \\
 & \frac{2}{13} \sqrt{-e} \left(-\frac{9e \left(-\frac{5e \left(-\frac{2e \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{3d} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \\
 & \quad \frac{2 \arctan \left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}} \right)}{13x^{13/2}} \\
 & \quad \downarrow 761
 \end{aligned}$$

$$\left(\frac{\frac{2}{13}\sqrt{-e}}{\frac{11d}{13x^{13/2}}} - \frac{9e \left(\frac{5e \left(\frac{e^{3/4}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{3d^{5/4}\sqrt{d+ex^2}} - \frac{2\sqrt{d+ex^2}}{3dx^{3/2}} \right)}{7d} - \frac{2\sqrt{d+ex^2}}{7dx^{7/2}} \right)}{11d} - \frac{2\sqrt{d+ex^2}}{11dx^{11/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{13x^{13/2}}$$

```
input Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(15/2),x]
```

```
output (-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(13*x^(13/2)) + (2*Sqrt[-e]*((-2*Sqrt[d + e*x^2])/(11*d*x^(11/2)) - (9*e*((-2*Sqrt[d + e*x^2])/(7*d*x^(7/2))) - (5*e*((-2*Sqrt[d + e*x^2])/(3*d*x^(3/2))) - (e^(3/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(3*d^(5/4)*Sqrt[d + e*x^2])))/(7*d)))/(11*d))/13
```

Definitions of rubi rules used

rule 264 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*(m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{15}{2}}} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.45

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx =$$

$$\frac{2\left(30\sqrt{-e}e^{\frac{5}{2}}x^7\text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right) + 77d^3\sqrt{x}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) + 2(15e^2x^5 - 9dex^3 + 7d^2x)\right)}{1001d^3x^7}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x, algorithm="fricas")`

output `-2/1001*(30*sqrt(-e)*e^(5/2)*x^7*weierstrassPInverse(-4*d/e, 0, x) + 77*d^3*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 2*(15*e^2*x^5 - 9*d*e*x^3 + 7*d^2*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(x))/(d^3*x^7)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \text{Timed out}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(15/2),x)`

output `Timed out`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x, algorithm="maxima")`

output `2/13*(13*d*sqrt(-e)*x^(13/2)*integrate(-1/13*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(15/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 15/2*log(x))), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d))/x^(13/2)`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^(15/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{15/2}} dx$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(15/2),x)`

output `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(15/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{15/2}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x^8} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(15/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x**8,x)`

3.25 $\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	243
Mathematica [C] (verified)	244
Rubi [A] (verified)	244
Maple [F]	248
Fricas [A] (verification not implemented)	248
Sympy [F(-1)]	249
Maxima [F]	249
Giac [F]	250
Mupad [F(-1)]	250
Reduce [F]	250

Optimal result

Integrand size = 27, antiderivative size = 326

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{28dx^{3/2}\sqrt{d+ex^2}}{405(-e)^{3/2}} + \frac{4x^{7/2}\sqrt{d+ex^2}}{81\sqrt{-e}} - \frac{28d^2\sqrt{-e}\sqrt{x}\sqrt{d+ex^2}}{135e^{5/2}(\sqrt{d}+\sqrt{ex})} + \frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + \frac{28d^{9/4}\sqrt{-e}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{135e^{11/4}\sqrt{d+ex^2}} - \frac{14d^{9/4}\sqrt{-e}}{135e^{11/4}\sqrt{d+ex^2}}$$

output

```
28/405*d*x^(3/2)*(e*x^2+d)^(1/2)/(-e)^(3/2)+4/81*x^(7/2)*(e*x^2+d)^(1/2)/(-e)^(1/2)-28/135*d^2*(-e)^(1/2)*x^(1/2)*(e*x^2+d)^(1/2)/e^(5/2)/(d^(1/2)+e^(1/2)*x)+2/9*x^(9/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+28/135*d^(9/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x))^2^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/e^(11/4)/(e*x^2+d)^(1/2)-14/135*d^(9/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x))^2^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(11/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.11 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.43

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{3/2}\left(2\sqrt{-e}(7d^2 + 2dex^2 - 5e^2x^4) + 45e^2x^3\sqrt{d+ex^2}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)\right)}{405e^2\sqrt{d+ex^2}}$$

input

```
Integrate[x^(7/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
(2*x^(3/2)*(2*Sqrt[-e]*(7*d^2 + 2*d*e*x^2 - 5*e^2*x^4) + 45*e^2*x^3*Sqrt[d + e*x^2]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] - 14*d^2*Sqrt[-e]*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)])/(405*e^2*Sqrt[d + e*x^2])
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 321, normalized size of antiderivative = 0.98, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {5674, 262, 262, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{5674} \\ & \frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{-e} \int \frac{x^{9/2}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \\ & \frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{-e} \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \int \frac{x^{5/2}}{\sqrt{ex^2+d}} dx}{9e} \right) \\ & \quad \downarrow \text{262} \end{aligned}$$

$$\frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{-e} \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{3d \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{5e} \right)}{9e} \right)$$

↓ 266

$$\frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{9}\sqrt{-e} \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x}}{5e} \right)}{9e} \right)$$

↓ 834

$$\frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right)}{9e} \right)$$

↓ 27

$$\frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right)}{9e} \right)$$

↓ 761

$$\frac{2}{9}\sqrt{-e} \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}}) \sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \int \frac{\sqrt{d-\sqrt{ex}}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right)}{9e} \right)$$

1510

$$\frac{2}{9}\sqrt{-e} \left(\frac{2x^{7/2}\sqrt{d+ex^2}}{9e} - \frac{7d \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\frac{2}{9}x^{9/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}}) \sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}})}{\sqrt{e}} \right)}{5e} \right)}{9e} \right)$$

input `Int[x^(7/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]`

output `(2*x^(9/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/9 - (2*Sqrt[-e]*((2*x^(7/2)*Sqrt[d + e*x^2])/(9*e) - (7*d*((2*x^(3/2)*Sqrt[d + e*x^2])/(5*e) - (6*d*(-((-(Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d + Sqrt[e]*x]^2)*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2]])/(e^(1/4)*Sqrt[d + e*x^2]))/Sqrt[e]) + (d^(1/4)*(Sqrt[d + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d + Sqrt[e]*x]^2)*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2]])/(2*e^(3/4)*Sqrt[d + e*x^2])))/(5*e)))/(9*e))/9`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 834 `Int[(x_)^2/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[1/q Int[1/Sqrt[a + b*x^4], x], x] - Simp[1/q Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1510 `Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

rule 5674 `Int[ArcTan[((c_)*(x_))/Sqrt[(a_) + (b_)*(x_)^2]]*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(7/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(7/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.29

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2 \left(45 e^3 x^{\frac{9}{2}} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + 42 d^2 \sqrt{-e} \sqrt{e} \text{weierstrassZeta}\left(-\frac{4d}{e}, 0, \text{weierstrassZeta}\right) \right)}{405 e}$$

input `integrate(x^(7/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output

```
2/405*(45*e^3*x^(9/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 42*d^2*sqrt(-e)
*sqrt(e)*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)) - 2
*(5*e^2*x^3 - 7*d*e*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(x))/e^3
```

Sympy [F(-1)]

Timed out.

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \text{Timed out}$$

input

```
integrate(x**(7/2)*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)
```

output

Timed out

Maxima [F]

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input

```
integrate(x^(7/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")
```

output

```
2/9*x^(9/2)*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - 2*d*sqrt(-e)*integrate(
-1/9*x*e^(1/2*log(e*x^2 + d) + 7/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)
)^2), x)
```

Giac [F]

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(7/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(7/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{7/2} \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(7/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(7/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

Reduce [F]

$$\int x^{7/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right) x^3 dx$$

input `int(x^(7/2)*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2))*x**3,x)`

3.26 $\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx$

Optimal result	251
Mathematica [C] (verified)	252
Rubi [A] (verified)	252
Maple [F]	255
Fricas [A] (verification not implemented)	256
Sympy [C] (verification not implemented)	256
Maxima [F]	257
Giac [F]	257
Mupad [F(-1)]	257
Reduce [F]	258

Optimal result

Integrand size = 27, antiderivative size = 296

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{4x^{3/2}\sqrt{d+ex^2}}{25\sqrt{-e}} + \frac{12d\sqrt{-e}\sqrt{x}\sqrt{d+ex^2}}{25e^{3/2}(\sqrt{d}+\sqrt{ex})}$$

$$+ \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{12d^{5/4}\sqrt{-e}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{25e^{7/4}\sqrt{d+ex^2}} + \frac{6d^{5/4}\sqrt{-e}}{25e^{7/4}\sqrt{d+ex^2}}$$

output

```
4/25*x^(3/2)*(e*x^2+d)^(1/2)/(-e)^(1/2)+12/25*d*(-e)^(1/2)*x^(1/2)*(e*x^2+d)^(1/2)/e^(3/2)/(d^(1/2)+e^(1/2)*x)+2/5*x^(5/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))-12/25*d^(5/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/e^(7/4)/(e*x^2+d)^(1/2)+6/25*d^(5/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(7/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.08 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.40

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{3/2}\left(-2\sqrt{-e}(d+ex^2) + 5ex\sqrt{d+ex^2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + 2d\sqrt{-e}\sqrt{1+\frac{ex^2}{d}}\right)}{25e\sqrt{d+ex^2}}$$

input

```
Integrate[x^(3/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]],x]
```

output

```
(2*x^(3/2)*(-2*Sqrt[-e]*(d + e*x^2) + 5*e*x*Sqrt[d + e*x^2]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] + 2*d*Sqrt[-e]*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)])/(25*e*Sqrt[d + e*x^2])
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$, Rules used = {5674, 262, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx \\ & \quad \downarrow \text{5674} \\ & \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{-e} \int \frac{x^{5/2}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{262} \\ & \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{-e} \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{3d \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{5e} \right) \\ & \quad \downarrow \text{266} \end{aligned}$$

$$\begin{aligned}
& \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{-e} \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x}}{5e} \right) \\
& \quad \downarrow \text{834} \\
& \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \\
& \frac{2}{5}\sqrt{-e} \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right) \\
& \quad \downarrow \text{27} \\
& \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{2}{5}\sqrt{-e} \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right) \\
& \quad \downarrow \text{761} \\
& \frac{2}{5}x^{5/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \\
& \frac{2}{5}\sqrt{-e} \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{5e} \right) \\
& \quad \downarrow \text{1510}
\end{aligned}$$

$$\frac{2}{5}\sqrt{-e} \left(\frac{2x^{3/2}\sqrt{d+ex^2}}{5e} - \frac{6d \left(\frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} \right) - \frac{\sqrt[4]{d}(\sqrt{d+\sqrt{ex}})\sqrt{\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}} E\left(\frac{d+ex^2}{(\sqrt{d+\sqrt{ex}})^2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}}}{5e} \right)$$

```
input Int[x^(3/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]], x]
```

```
output (2*x^(5/2)*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/5 - (2*Sqrt[-e]*((2*x^(3/2)*Sqrt[d + e*x^2])/(5*e) - (6*d*(-((-((Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)^2)*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(e^(1/4)*Sqrt[d + e*x^2])/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)^2)*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(2*e^(3/4)*Sqrt[d + e*x^2])))/(5*e))/5
```

Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 262 Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 761 `Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 834 `Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[1/q Int[1/Sqrt[a + b*x^4], x], x] - Simp[1/q Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]`

rule 1510 `Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]`

rule 5674 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*(ArcTan[(c*x)/Sqrt[a + b*x^2]]/(d*(m + 1))), x] - Simp[c/(d*(m + 1)) Int[(d*x)^(m + 1)/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

Maple [F]

$$\int x^{\frac{3}{2}} \arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(3/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(x^(3/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.27

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2\left(5e^2x^{5/2}\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) - 2\sqrt{ex^2+d}\sqrt{-ex}^{3/2} - 6d\sqrt{-e}\sqrt{e}\text{weierstrass}\right)}{25e^2}$$

input `integrate(x^(3/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="fricas")`

output `2/25*(5*e^2*x^(5/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) - 2*sqrt(e*x^2 + d)*sqrt(-e)*e*x^(3/2) - 6*d*sqrt(-e)*sqrt(e)*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)))/e^2`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 10.34 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.25

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \frac{2x^{5/2} \operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{5} - \frac{x^{7/2}\sqrt{-e}\Gamma\left(\frac{7}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{7}{4} \middle| \frac{ex^2e^{i\pi}}{d}\right)}{5\sqrt{d}\Gamma\left(\frac{11}{4}\right)}$$

input `integrate(x**(3/2)*atan((-e)**(1/2)*x/(e*x**2+d)**(1/2)),x)`

output `2*x**(5/2)*atan(x*sqrt(-e)/sqrt(d + e*x**2))/5 - x**(7/2)*sqrt(-e)*gamma(7/4)*hyper((1/2, 7/4), (11/4,), e*x**2*exp_polar(I*pi)/d)/(5*sqrt(d)*gamma(11/4))`

Maxima [F]

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(3/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="maxima")`

output `2/5*x^(5/2)*arctan2(sqrt(-e)*x, sqrt(e*x^2 + d)) - 2*d*sqrt(-e)*integrate(-1/5*x*e^(1/2*log(e*x^2 + d) + 3/2*log(x))/(e^2*x^4 + d*e*x^2 - (e*x^2 + d)^2), x)`

Giac [F]

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `integrate(x^(3/2)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x, algorithm="giac")`

output `integrate(x^(3/2)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int x^{3/2} \operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right) dx$$

input `int(x^(3/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)),x)`

output `int(x^(3/2)*atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2)), x)`

Reduce [F]

$$\int x^{3/2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) dx = \int \sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right) x dx$$

input `int(x^(3/2)*atan((-e)^(1/2)*x/(e*x^2+d)^(1/2)),x)`

output `int(sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2))*x,x)`

3.27
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$$

Optimal result	259
Mathematica [C] (verified)	260
Rubi [A] (verified)	260
Maple [F]	263
Fricas [A] (verification not implemented)	263
Sympy [C] (verification not implemented)	263
Maxima [F]	264
Giac [F]	264
Mupad [F(-1)]	265
Reduce [F]	265

Optimal result

Integrand size = 27, antiderivative size = 260

$$\begin{aligned} & \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx \\ &= -\frac{4\sqrt{-e}\sqrt{x}\sqrt{d+ex^2}}{\sqrt{e}(\sqrt{d}+\sqrt{ex})} + 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) \\ & \quad + \frac{4\sqrt[4]{d}\sqrt{-e}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{e^{3/4}\sqrt{d+ex^2}} \\ & \quad - \frac{2\sqrt[4]{d}\sqrt{-e}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{e^{3/4}\sqrt{d+ex^2}} \end{aligned}$$

output

```
-4*(-e)^(1/2)*x^(1/2)*(e*x^2+d)^(1/2)/e^(1/2)/(d^(1/2)+e^(1/2)*x)+2*x^(1/2)
)*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))+4*d^(1/4)*(-e)^(1/2)*(d^(1/2)+e^(1/2)
)*x*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*EllipticE(sin(2*arctan(e^(1/4)
)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/e^(3/4)/(e*x^2+d)^(1/2)-2*d^(1/4)*(-e)^(
1/2)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*InverseJa
cobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/e^(3/4)/(e*x^2+d)^(1
/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.07 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.34

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \frac{4\sqrt{-ex}^{3/2} \sqrt{1 + \frac{ex^2}{d}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{ex^2}{d}\right)}{3\sqrt{d+ex^2}}$$

input `Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/Sqrt[x], x]`

output `2*Sqrt[x]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] - (4*Sqrt[-e]*x^(3/2)*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)]/(3*Sqrt[d + e*x^2]))`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 253, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5674, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx \\ & \quad \downarrow \text{5674} \\ & 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - 2\sqrt{-e} \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx \\ & \quad \downarrow \text{266} \\ & 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - 4\sqrt{-e} \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x} \end{aligned}$$

$$\begin{aligned}
& \downarrow 834 \\
& 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - 4\sqrt{-e} \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right) \\
& \downarrow 27 \\
& 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - 4\sqrt{-e} \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right) \\
& \downarrow 761 \\
& 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \\
& 4\sqrt{-e} \left(\frac{\sqrt[4]{d}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right) \\
& \downarrow 1510 \\
& 2\sqrt{x} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - \\
& 4\sqrt{-e} \left(\frac{\sqrt[4]{d}(\sqrt{d} + \sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}} \right)
\end{aligned}$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/Sqrt[x], x]`

output `2*Sqrt[x]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] - 4*Sqrt[-e]*(-((-(Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(e^(1/4)*Sqrt[d + e*x^2]))/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2])/(2*e^(3/4)*Sqrt[d + e*x^2]))`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 266 $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)}(a + b*(x^{(2*k)/c^2)})^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 761 $\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^4]/(a*(1 + q^2*x^2)^2))/(2*q*\text{Sqrt}[a + b*x^4))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 834 $\text{Int}[(x_)^2/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 2]\}, \text{Simp}[1/q \text{ Int}[1/\text{Sqrt}[a + b*x^4], x], x] - \text{Simp}[1/q \text{ Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^4], x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 1510 $\text{Int}[((d_*) + (e_*)(x_)^2)/\text{Sqrt}[(a_*) + (c_*)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2)^2))/(q*\text{Sqrt}[a + c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$
- rule 5674 $\text{Int}[\text{ArcTan}[(c_*)(x_)/\text{Sqrt}[(a_*) + (b_*)(x_)^2]]*((d_*)(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTan}[(c*x)/\text{Sqrt}[a + b*x^2]]/(d*(m+1))), x] - \text{Simp}[c/(d*(m+1)) \text{ Int}[(d*x)^{(m+1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b + c^2, 0] \ \&\& \ \text{NeQ}[m, -1]$

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.21

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx$$

$$= \frac{2\left(e\sqrt{x}\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right) + 2\sqrt{-e}\sqrt{e}\operatorname{weierstrassZeta}\left(-\frac{4d}{e}, 0, \operatorname{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)\right)\right)}{e}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x, algorithm="fricas")`

output `2*(e*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 2*sqrt(-e)*sqrt(e)*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)))/e`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 9.59 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.27

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = 2\sqrt{x}\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right) - \frac{x^{\frac{3}{2}}\sqrt{-e}\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{ex^2e^{i\pi}}{d}\right)}{\sqrt{d}\Gamma\left(\frac{7}{4}\right)}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(1/2),x)`

output `2*sqrt(x)*atan(x*sqrt(-e)/sqrt(d + e*x**2)) - x**(3/2)*sqrt(-e)*gamma(3/4)
*hyper((1/2, 3/4), (7/4,), e*x**2*exp_polar(I*pi)/d)/(sqrt(d)*gamma(7/4))`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x, algorithm="maxima")`

output `-2*d*sqrt(-e)*integrate(sqrt(e*x^2 + d)*x/((e*x^2 + d)*e^(log(e*x^2 + d) +
1/2*log(x)) - (e^2*x^4 + d*e*x^2)*sqrt(x)), x) + 2*sqrt(x)*arctan2(sqrt(-
e)*x, sqrt(e*x^2 + d))`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/sqrt(x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{\sqrt{x}} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2)/x^(1/2),x)`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2)/x^(1/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{\sqrt{x}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(1/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x,x)`

3.28
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx$$

Optimal result	266
Mathematica [C] (verified)	267
Rubi [A] (verified)	267
Maple [F]	271
Fricas [A] (verification not implemented)	271
Sympy [C] (verification not implemented)	271
Maxima [F]	272
Giac [F]	272
Mupad [F(-1)]	273
Reduce [F]	273

Optimal result

Integrand size = 27, antiderivative size = 298

$$\begin{aligned} \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = & -\frac{4\sqrt{-e}\sqrt{d+ex^2}}{3d\sqrt{x}} \\ & + \frac{4\sqrt{-e^2}\sqrt{x}\sqrt{d+ex^2}}{3d(\sqrt{d}+\sqrt{ex})} - \frac{2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\ & - \frac{4\sqrt{-e}\sqrt[4]{e}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{3d^{3/4}\sqrt{d+ex^2}} \\ & + \frac{2\sqrt{-e}\sqrt[4]{e}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{3d^{3/4}\sqrt{d+ex^2}} \end{aligned}$$

output

```
-4/3*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^(1/2)+4/3*(-e^2)^(1/2)*x^(1/2)*(e*x^2+d)^(1/2)/d/(d^(1/2)+e^(1/2)*x)-2/3*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(3/2)-4/3*(-e)^(1/2)*e^(1/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x))^2)^(1/2)*EllipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/d^(3/4)/(e*x^2+d)^(1/2)+2/3*(-e)^(1/2)*e^(1/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x))^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2*2^(1/2))/d^(3/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.09 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.41

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx =$$

$$\frac{2\left(6\sqrt{-ex}(d+ex^2) + 3d\sqrt{d+ex^2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) + 2(-e)^{3/2}x^3\sqrt{1+\frac{ex^2}{d}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}\right)\right)}{9dx^{3/2}\sqrt{d+ex^2}}$$

input

```
Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(5/2), x]
```

output

```
(-2*(6*Sqrt[-e]*x*(d + e*x^2) + 3*d*Sqrt[d + e*x^2]*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] + 2*(-e)^(3/2)*x^3*Sqrt[1 + (e*x^2)/d]*Hypergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)]))/(9*d*x^(3/2)*Sqrt[d + e*x^2])
```

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 285, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$, Rules used = {5674, 264, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx \\
& \quad \downarrow \text{5674} \\
& \frac{2}{3}\sqrt{-e} \int \frac{1}{x^{3/2}\sqrt{ex^2+d}} dx - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow \text{264} \\
& \frac{2}{3}\sqrt{-e} \left(\frac{e \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow \text{266} \\
& \frac{2}{3}\sqrt{-e} \left(\frac{2e \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x}}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow \text{834} \\
& \frac{2}{3}\sqrt{-e} \left(\frac{2e \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{2}{3}\sqrt{-e} \left(\frac{2e \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \\
& \quad \downarrow \text{761}
\end{aligned}$$

$$\frac{2}{3}\sqrt{-e} \left(\frac{2e \left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}}$$

↓ 1510

$$\frac{2}{3}\sqrt{-e} \left(\frac{2e \left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}} \right)}{d} - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{3x^{3/2}} \right)$$

input `Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(5/2),x]`

output `(-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]])/(3*x^(3/2)) + (2*Sqrt[-e]*((-2*Sqrt[d + e*x^2]))/(d*Sqrt[x]) + (2*e*(-((-(Sqrt[x]*Sqrt[d + e*x^2]))/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2]))/(e^(1/4)*Sqrt[d + e*x^2])/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2]))/(2*e^(3/4)*Sqrt[d + e*x^2]))/d)/3`

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 264 $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^{2*(m+1)})) \text{ Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 266 $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(2*k)}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 761 $\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^4]/(a*(1 + q^2*x^2)^2)]/(2*q*\text{Sqrt}[a + b*x^4))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x]] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 834 $\text{Int}[(x_)^2/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b/a, 2]\}, \text{Simp}[1/q \text{ Int}[1/\text{Sqrt}[a + b*x^4], x], x] - \text{Simp}[1/q \text{ Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^4], x], x]] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 1510 $\text{Int}[((d_*) + (e_*)(x_)^2)/\text{Sqrt}[(a_*) + (c_*)(x_)^4], x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*\text{Sqrt}[a + c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$
- rule 5674 $\text{Int}[\text{ArcTan}[(c_*)(x_)/\text{Sqrt}[(a_*) + (b_*)(x_)^2]]*((d_*)(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTan}[(c*x)/\text{Sqrt}[a + b*x^2]]/(d*(m+1))), x] - \text{Simp}[c/(d*(m+1)) \text{ Int}[(d*x)^{(m+1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b + c^2, 0] \ \&\& \ \text{NeQ}[m, -1]$

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{ex^2+d}}\right)}{x^{\frac{5}{2}}} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x)`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.27

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \frac{2\left(2\sqrt{-e}\sqrt{ex^2}\text{weierstrassZeta}\left(-\frac{4d}{e}, 0, \text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)\right) + 2\sqrt{ex^2+d}\sqrt{-e}x^{\frac{3}{2}} + d\sqrt{x}\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)\right)}{3dx^2}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x, algorithm="fricas")`

output `-2/3*(2*sqrt(-e)*sqrt(e)*x^2*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)) + 2*sqrt(e*x^2 + d)*sqrt(-e)*x^(3/2) + d*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)))/(d*x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 10.40 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.26

$$\int \frac{\arctan\left(\frac{\sqrt{-e}x}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = -\frac{2\operatorname{atan}\left(\frac{x\sqrt{-e}}{\sqrt{d+ex^2}}\right)}{3x^{\frac{3}{2}}} + \frac{\sqrt{-e}\Gamma\left(-\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{4}, \frac{1}{2} \\ \frac{3}{4} \end{matrix} \middle| \frac{ex^2e^{i\pi}}{d} \right)}{3\sqrt{d}\sqrt{x}\Gamma\left(\frac{3}{4}\right)}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(5/2),x)`

output `-2*atan(x*sqrt(-e)/sqrt(d + e*x**2))/(3*x**(3/2)) + sqrt(-e)*gamma(-1/4)*hyper((-1/4, 1/2), (3/4,), e*x**2*exp_polar(I*pi)/d)/(3*sqrt(d)*sqrt(x)*gamma(3/4))`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{5/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x, algorithm="maxima")`

output `2/3*(3*d*sqrt(-e)*x^(3/2)*integrate(-1/3*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(5/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 5/2*log(x))), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d))/x^(3/2)`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{5/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^(5/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{5/2}} dx$$

input `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2)/x^(5/2),x)`

output `int(atan((-e)^(1/2)*x)/(d + e*x^2)^(1/2)/x^(5/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{5/2}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x^3} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(5/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x**3,x)`

3.29
$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx$$

Optimal result	274
Mathematica [C] (verified)	275
Rubi [A] (verified)	275
Maple [F]	280
Fricas [A] (verification not implemented)	280
Sympy [F(-1)]	280
Maxima [F]	281
Giac [F]	281
Mupad [F(-1)]	281
Reduce [F]	282

Optimal result

Integrand size = 27, antiderivative size = 331

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = -\frac{4\sqrt{-e}\sqrt{d+ex^2}}{35dx^{5/2}} - \frac{12(-e)^{3/2}\sqrt{d+ex^2}}{35d^2\sqrt{x}}$$

$$- \frac{12\sqrt{-e}e^{3/2}\sqrt{x}\sqrt{d+ex^2}}{35d^2(\sqrt{d}+\sqrt{ex})} - \frac{2\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}}$$

$$+ \frac{12\sqrt{-e}e^{5/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}E\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right)\middle|\frac{1}{2}\right)}{35d^{7/4}\sqrt{d+ex^2}}$$

$$- \frac{6\sqrt{-e}e^{5/4}(\sqrt{d}+\sqrt{ex})\sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}}\text{EllipticF}\left(2\arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right),\frac{1}{2}\right)}{35d^{7/4}\sqrt{d+ex^2}}$$

output

```
-4/35*(-e)^(1/2)*(e*x^2+d)^(1/2)/d/x^(5/2)-12/35*(-e)^(3/2)*(e*x^2+d)^(1/2)
)/d^2/x^(1/2)-12/35*(-e)^(1/2)*e^(3/2)*x^(1/2)*(e*x^2+d)^(1/2)/d^2/(d^(1/2)
)+e^(1/2)*x)-2/7*arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(7/2)+12/35*(-e)^(
1/2)*e^(5/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)+e^(1/2)*x)^2)^(1/2)*E
llipticE(sin(2*arctan(e^(1/4)*x^(1/2)/d^(1/4))),1/2*2^(1/2))/d^(7/4)/(e*x^
2+d)^(1/2)-6/35*(-e)^(1/2)*e^(5/4)*(d^(1/2)+e^(1/2)*x)*((e*x^2+d)/(d^(1/2)
+e^(1/2)*x)^2)^(1/2)*InverseJacobiAM(2*arctan(e^(1/4)*x^(1/2)/d^(1/4)),1/2
*2^(1/2))/d^(7/4)/(e*x^2+d)^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.08 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.41

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \frac{4\sqrt{-ex}(-d^2 + 2dex^2 + 3e^2x^4) - 10d^2\sqrt{d+ex^2} \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right) - 4(-e)^{5/2}x^5\sqrt{d+ex^2}}{35d^2x^{7/2}\sqrt{d+ex^2}}$$

input

```
Integrate[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(9/2),x]
```

output

```
(4*Sqrt[-e]*x*(-d^2 + 2*d*e*x^2 + 3*e^2*x^4) - 10*d^2*Sqrt[d + e*x^2]*ArcT
an[(Sqrt[-e]*x)/Sqrt[d + e*x^2]] - 4*(-e)^(5/2)*x^5*Sqrt[1 + (e*x^2)/d]*Hy
pergeometric2F1[1/2, 3/4, 7/4, -((e*x^2)/d)]/(35*d^2*x^(7/2)*Sqrt[d + e*x
^2])
```

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {5674, 264, 264, 266, 834, 27, 761, 1510}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx \\
 & \quad \downarrow \text{5674} \\
 & \frac{2}{7}\sqrt{-e} \int \frac{1}{x^{7/2}\sqrt{ex^2+d}} dx - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
 & \quad \downarrow \text{264} \\
 & \frac{2}{7}\sqrt{-e} \left(-\frac{3e \int \frac{1}{x^{3/2}\sqrt{ex^2+d}} dx}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
 & \quad \downarrow \text{264} \\
 & \frac{2}{7}\sqrt{-e} \left(-\frac{3e \left(\frac{e \int \frac{\sqrt{x}}{\sqrt{ex^2+d}} dx}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
 & \quad \downarrow \text{266} \\
 & \frac{2}{7}\sqrt{-e} \left(-\frac{3e \left(\frac{2e \int \frac{x}{\sqrt{ex^2+d}} d\sqrt{x}}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
 & \quad \downarrow \text{834} \\
 & \frac{2}{7}\sqrt{-e} \left(-\frac{3e \left(\frac{2e \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\sqrt{d} \int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{d}\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \\
 & \quad \frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\frac{2}{7}\sqrt{-e} \left(\frac{3e \left(\frac{2e \left(\frac{\sqrt{d} \int \frac{1}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{d} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2 \arctan \left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}} \right)}{7x^{7/2}}$$

↓ 761

$$\frac{2}{7}\sqrt{-e} \left(\frac{3e \left(\frac{2e \left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF} \left(2 \arctan \left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt{d}} \right), \frac{1}{2} \right) - \frac{\int \frac{\sqrt{d}-\sqrt{ex}}{\sqrt{ex^2+d}} d\sqrt{x}}{\sqrt{e}} \right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{2\sqrt{d+ex^2}}{d\sqrt{x}} \right)}{5d} - \frac{2\sqrt{d+ex^2}}{5dx^{5/2}} \right) - \frac{2 \arctan \left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}} \right)}{7x^{7/2}}$$

↓ 1510

$$\left(\frac{2e}{3e} \left(\frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{2e^{3/4}\sqrt{d+ex^2}} - \frac{\sqrt[4]{d}(\sqrt{d}+\sqrt{ex}) \sqrt{\frac{d+ex^2}{(\sqrt{d}+\sqrt{ex})^2}} E\left(2 \arctan\left(\frac{\sqrt[4]{e}\sqrt{x}}{\sqrt[4]{d}}\right), \frac{1}{2}\right)}{\sqrt[4]{e}\sqrt{d+ex^2}\sqrt{e}} \right) - \frac{2}{7}\sqrt{-e} \right) \frac{d}{5d}$$

$$\frac{2 \arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{7x^{7/2}}$$

input

```
Int[ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/x^(9/2),x]
```

output

```
(-2*ArcTan[(Sqrt[-e]*x)/Sqrt[d + e*x^2]]/(7*x^(7/2)) + (2*Sqrt[-e]*((-2*Sqrt[d + e*x^2])/(5*d*x^(5/2)) - (3*e*((-2*Sqrt[d + e*x^2])/(d*Sqrt[x]) + (2*e*(-((-(Sqrt[x]*Sqrt[d + e*x^2])/(Sqrt[d] + Sqrt[e]*x)) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticE[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(e^(1/4)*Sqrt[d + e*x^2])/Sqrt[e]) + (d^(1/4)*(Sqrt[d] + Sqrt[e]*x)*Sqrt[(d + e*x^2)/(Sqrt[d] + Sqrt[e]*x)^2]*EllipticF[2*ArcTan[(e^(1/4)*Sqrt[x])/d^(1/4)], 1/2)]/(2*e^(3/4)*Sqrt[d + e*x^2])))/d))/(5*d))/7
```

Definitions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 264 $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m+1)}((a + b*x^2)^{(p+1})/(a*c*(m+1))), x] - \text{Simp}[b*((m+2*p+3)/(a*c^{2*(m+1)})) \text{ Int}[(c*x)^{(m+2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 266 $\text{Int}[((c_*)(x_))^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{ Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(2*k)}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 761 $\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^4]/(a*(1 + q^2*x^2)^2)]/(2*q*\text{Sqrt}[a + b*x^4))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x]] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 834 $\text{Int}[(x_)^2/\text{Sqrt}[(a_*) + (b_*)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 2]\}, \text{Simp}[1/q \text{ Int}[1/\text{Sqrt}[a + b*x^4], x], x] - \text{Simp}[1/q \text{ Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^4], x], x]] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$
- rule 1510 $\text{Int}[((d_*) + (e_*)(x_)^2)/\text{Sqrt}[(a_*) + (c_*)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*\text{Sqrt}[a + c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$
- rule 5674 $\text{Int}[\text{ArcTan}[(c_*)(x_)/\text{Sqrt}[(a_*) + (b_*)(x_)^2]]*((d_*)(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*(\text{ArcTan}[(c*x)/\text{Sqrt}[a + b*x^2]]/(d*(m+1))), x] - \text{Simp}[c/(d*(m+1)) \text{ Int}[(d*x)^{(m+1)}/\text{Sqrt}[a + b*x^2], x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b + c^2, 0] \ \&\& \ \text{NeQ}[m, -1]$

Maple [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{\frac{9}{2}}} dx$$

input `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x)`

output `int(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.28

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \frac{2\left(6\sqrt{-e}e^{\frac{3}{2}}x^4\text{weierstrassZeta}\left(-\frac{4d}{e}, 0, \text{weierstrassPInverse}\left(-\frac{4d}{e}, 0, x\right)\right) - 5d^2\sqrt{x}\right)}{35d^2x^4}$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x, algorithm="fricas")`

output `2/35*(6*sqrt(-e)*e^(3/2)*x^4*weierstrassZeta(-4*d/e, 0, weierstrassPInverse(-4*d/e, 0, x)) - 5*d^2*sqrt(x)*arctan(sqrt(-e)*x/sqrt(e*x^2 + d)) + 2*(3*e*x^3 - d*x)*sqrt(e*x^2 + d)*sqrt(-e)*sqrt(x))/(d^2*x^4)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \text{Timed out}$$

input `integrate(atan((-e)**(1/2)*x/(e*x**2+d)**(1/2))/x**(9/2),x)`

output `Timed out`

Maxima [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{9/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x, algorithm="maxima")`

output `2/7*(7*d*sqrt(-e)*x^(7/2)*integrate(-1/7*sqrt(e*x^2 + d)*x/((e^2*x^4 + d*e*x^2)*x^(9/2) - (e*x^2 + d)*e^(log(e*x^2 + d) + 9/2*log(x))), x) - arctan2(sqrt(-e)*x, sqrt(e*x^2 + d))/x^(7/2)`

Giac [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{9/2}} dx$$

input `integrate(arctan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x, algorithm="giac")`

output `integrate(arctan(sqrt(-e)*x/sqrt(e*x^2 + d))/x^(9/2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\operatorname{atan}\left(\frac{\sqrt{-ex}}{\sqrt{ex^2+d}}\right)}{x^{9/2}} dx$$

input `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(9/2),x)`

output `int(atan(((e)^(1/2)*x)/(d + e*x^2)^(1/2))/x^(9/2), x)`

Reduce [F]

$$\int \frac{\arctan\left(\frac{\sqrt{-ex}}{\sqrt{d+ex^2}}\right)}{x^{9/2}} dx = \int \frac{\sqrt{x} \operatorname{atan}\left(\frac{\sqrt{e}ix}{\sqrt{ex^2+d}}\right)}{x^5} dx$$

input `int(atan((-e)^(1/2)*x/(e*x^2+d)^(1/2))/x^(9/2),x)`

output `int((sqrt(x)*atan((sqrt(e)*i*x)/sqrt(d + e*x**2)))/x**5,x)`

3.30 $\int \frac{\arctan(1+x+x^2)}{x^2} dx$

Optimal result	283
Mathematica [A] (verified)	283
Rubi [A] (verified)	284
Maple [A] (verified)	285
Fricas [A] (verification not implemented)	286
Sympy [A] (verification not implemented)	286
Maxima [A] (verification not implemented)	287
Giac [A] (verification not implemented)	287
Mupad [B] (verification not implemented)	288
Reduce [B] (verification not implemented)	288

Optimal result

Integrand size = 11, antiderivative size = 50

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = \frac{1}{2} \arctan(1+x) - \frac{\arctan(1+x+x^2)}{x} + \frac{\log(x)}{2} - \frac{1}{2} \log(1+x^2) + \frac{1}{4} \log(2+2x+x^2)$$

output

```
1/2*arctan(1+x)-arctan(x^2+x+1)/x+1/2*ln(x)-1/2*ln(x^2+1)+1/4*ln(x^2+2*x+2)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = \frac{1}{2} \arctan(1+x) - \frac{\arctan(1+x+x^2)}{x} + \frac{\log(x)}{2} - \frac{1}{2} \log(1+x^2) + \frac{1}{4} \log(2+2x+x^2)$$

input

```
Integrate[ArcTan[1 + x + x^2]/x^2,x]
```

output

```
ArcTan[1 + x]/2 - ArcTan[1 + x + x^2]/x + Log[x]/2 - Log[1 + x^2]/2 + Log[
2 + 2*x + x^2]/4
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {5728, 2462, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(x^2 + x + 1)}{x^2} dx$$

$$\downarrow 5728$$

$$\int \frac{2x + 1}{x(x^4 + 2x^3 + 3x^2 + 2x + 2)} dx - \frac{\arctan(x^2 + x + 1)}{x}$$

$$\downarrow 2462$$

$$\int \left(-\frac{x}{x^2 + 1} + \frac{x + 2}{2(x^2 + 2x + 2)} + \frac{1}{2x} \right) dx - \frac{\arctan(x^2 + x + 1)}{x}$$

$$\downarrow 2009$$

$$-\frac{\arctan(x^2 + x + 1)}{x} + \frac{1}{2} \arctan(x + 1) - \frac{1}{2} \log(x^2 + 1) + \frac{1}{4} \log(x^2 + 2x + 2) + \frac{\log(x)}{2}$$

input

```
Int[ArcTan[1 + x + x^2]/x^2,x]
```

output

```
ArcTan[1 + x]/2 - ArcTan[1 + x + x^2]/x + Log[x]/2 - Log[1 + x^2]/2 + Log[
2 + 2*x + x^2]/4
```

Definitions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2462 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{Qx = Factor[Px]}, Int[ExpandIntegr
and[u*Qx^p, x], x] /; !SumQ[NonfreeFactors[Qx, x]] /; PolyQ[Px, x] && GtQ
[Expon[Px, x], 2] && !BinomialQ[Px, x] && !TrinomialQ[Px, x] && ILtQ[p, 0
] && RationalFunctionQ[u, x]`

rule 5728 `Int[((a_.) + ArcTan[u]*(b_.))*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Sim
p[(c + d*x)^(m + 1)*((a + b*ArcTan[u])/(d*(m + 1))), x] - Simp[b/(d*(m + 1
) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(1 + u^2)), x], x]
/; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] &
& !FunctionOfQ[(c + d*x)^(m + 1), u, x] && FalseQ[PowerVariableExpn[u, m +
1, x]]`

Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.86

method	result
default	$\frac{\arctan(x+1)}{2} - \frac{\arctan(x^2+x+1)}{x} + \frac{\ln(x)}{2} - \frac{\ln(x^2+1)}{2} + \frac{\ln(x^2+2x+2)}{4}$
parts	$\frac{\arctan(x+1)}{2} - \frac{\arctan(x^2+x+1)}{x} + \frac{\ln(x)}{2} - \frac{\ln(x^2+1)}{2} + \frac{\ln(x^2+2x+2)}{4}$
risch	$\frac{i \ln(1+i(x^2+x+1))}{2x} - \frac{i(2i \ln(x)x+i \ln(1-i+x)x+i \ln(1+i+x)x-2i \ln(x^2+1)x+\ln(1-i+x)x-\ln(1+i+x)x+2 \ln(1-i(x^2+x+1)))}{4x}$
parallelrisch	$\frac{4i \ln(x-i)x-4i \ln(x+i)x-7i \ln(1-i+x)x+7i \ln(1+i+x)x+6 \ln(x)x-6 \ln(x-i)x-6 \ln(x+i)x+3 \ln(1-i+x)x+3 \ln(1+i+x)x}{12x}$

input `int(arctan(x^2+x+1)/x^2,x,method=_RETURNVERBOSE)`

output `1/2*arctan(x+1)-arctan(x^2+x+1)/x+1/2*ln(x)-1/2*ln(x^2+1)+1/4*ln(x^2+2*x+2)`
)

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.94

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = \frac{2x \arctan(x+1) + x \log(x^2+2x+2) - 2x \log(x^2+1) + 2x \log(x) - 4 \arctan(x^2+x+1)}{4x}$$

input `integrate(arctan(x^2+x+1)/x^2,x, algorithm="fricas")`

output `1/4*(2*x*arctan(x + 1) + x*log(x^2 + 2*x + 2) - 2*x*log(x^2 + 1) + 2*x*log(x) - 4*arctan(x^2 + x + 1))/x`

Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.82

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = \frac{\log(x)}{2} - \frac{\log(x^2+1)}{2} + \frac{\log(x^2+2x+2)}{4} + \frac{\operatorname{atan}(x+1)}{2} - \frac{\operatorname{atan}(x^2+x+1)}{x}$$

input `integrate(atan(x**2+x+1)/x**2,x)`

output `log(x)/2 - log(x**2 + 1)/2 + log(x**2 + 2*x + 2)/4 + atan(x + 1)/2 - atan(x**2 + x + 1)/x`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.84

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = -\frac{\arctan(x^2+x+1)}{x} + \frac{1}{2} \arctan(x+1) + \frac{1}{4} \log(x^2+2x+2) - \frac{1}{2} \log(x^2+1) + \frac{1}{2} \log(x)$$

input `integrate(arctan(x^2+x+1)/x^2,x, algorithm="maxima")`output `-arctan(x^2 + x + 1)/x + 1/2*arctan(x + 1) + 1/4*log(x^2 + 2*x + 2) - 1/2*log(x^2 + 1) + 1/2*log(x)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.86

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = -\frac{\arctan(x^2+x+1)}{x} + \frac{1}{2} \arctan(x+1) + \frac{1}{4} \log(x^2+2x+2) - \frac{1}{2} \log(x^2+1) + \frac{1}{2} \log(|x|)$$

input `integrate(arctan(x^2+x+1)/x^2,x, algorithm="giac")`output `-arctan(x^2 + x + 1)/x + 1/2*arctan(x + 1) + 1/4*log(x^2 + 2*x + 2) - 1/2*log(x^2 + 1) + 1/2*log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.84

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = \frac{\operatorname{atan}(x+1)}{2} + \frac{\ln(x^2+2x+2)}{4} - \frac{\ln(x^2+1)}{2} + \frac{\ln(x)}{2} - \frac{\operatorname{atan}(x^2+x+1)}{x}$$

input `int(atan(x + x^2 + 1)/x^2,x)`output `atan(x + 1)/2 + log(2*x + x^2 + 2)/4 - log(x^2 + 1)/2 + log(x)/2 - atan(x + x^2 + 1)/x`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(1+x+x^2)}{x^2} dx = \frac{-2\operatorname{atan}(x^2+x+1)x - 4\operatorname{atan}(x^2+x+1) + 2\operatorname{atan}(x)x + \log(x^2+2x+2)x - 2\log(x^2+1)x + 2\log(x)}{4x}$$

input `int(atan(x^2+x+1)/x^2,x)`output `(- 2*atan(x**2 + x + 1)*x - 4*atan(x**2 + x + 1) + 2*atan(x)*x + log(x**2 + 2*x + 2)*x - 2*log(x**2 + 1)*x + 2*log(x)*x)/(4*x)`

$$3.31 \quad \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx$$

Optimal result	289
Mathematica [N/A]	289
Rubi [N/A]	290
Maple [N/A]	290
Fricas [N/A]	291
Sympy [N/A]	291
Maxima [N/A]	292
Giac [N/A]	292
Mupad [N/A]	293
Reduce [N/A]	293

Optimal result

Integrand size = 40, antiderivative size = 40

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx = \text{Int}\left(\frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2}, x\right)$$

output `Defer(Int)((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x)`

Mathematica [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx = \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1-c^2x^2} dx$$

input `Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^n/(1 - c^2*x^2),x]`

output `Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^n/(1 - c^2*x^2), x]`

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^n}{1 - c^2x^2} dx$$

↓ 7234

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^n}{1 - c^2x^2} dx$$

input

```
Int[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^n/(1 - c^2*x^2), x]
```

output

```
$Aborted
```

Maple [N/A]

Not integrable

Time = 1.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)\right)^n}{-c^2x^2 + 1} dx$$

input

```
int((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1), x)
```

output

```
int((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1), x)
```

Fricas [N/A]

Not integrable

Time = 0.13 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x, alg
orithm="fricas")`

output `integral(-(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^n/(c^2*x^2 - 1), x)`

Sympy [N/A]

Not integrable

Time = 4.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = -\int \frac{\left(a + b \operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*atan((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**n/(-c**2*x**2+1),x)`

output `-Integral((a + b*atan(sqrt(-c*x + 1)/sqrt(c*x + 1)))**n/(c**2*x**2 - 1), x
)`

Maxima [N/A]

Not integrable

Time = 0.87 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x, algorithm="maxima")`

output `-integrate((b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^n/(c^2*x^2 - 1), x)`

Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^n}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x, algorithm="giac")`

output `integrate(-(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^n/(c^2*x^2 - 1), x)`

Mupad [N/A]

Not integrable

Time = 0.85 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = - \int \frac{\left(a + b \operatorname{atan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^n}{c^2x^2 - 1} dx$$

input `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^n/(c^2*x^2 - 1),x)`

output `-int((a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^n/(c^2*x^2 - 1), x)`

Reduce [N/A]

Not integrable

Time = 0.62 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^n}{1 - c^2x^2} dx = - \left(\int \frac{\left(\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) b + a\right)^n}{c^2x^2 - 1} dx \right)$$

input `int((a+b*atan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^n/(-c^2*x^2+1),x)`

output `- int((atan(sqrt(- c*x + 1)/sqrt(c*x + 1))*b + a)**n/(c**2*x**2 - 1),x)`

$$3.32 \quad \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1-c^2x^2} dx$$

Optimal result	294
Mathematica [F]	295
Rubi [A] (verified)	295
Maple [B] (verified)	298
Fricas [F]	299
Sympy [F]	300
Maxima [F]	300
Giac [F]	301
Mupad [F(-1)]	301
Reduce [F]	301

Optimal result

Integrand size = 40, antiderivative size = 431

$$\begin{aligned} & \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1-c^2x^2} dx \\ &= -\frac{2\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3 \operatorname{arctanh}\left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad + \frac{3ib\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad - \frac{3ib\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2 \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad + \frac{3b^2\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad - \frac{3b^2\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \\ & \quad - \frac{3ib^3 \operatorname{PolyLog}\left(4, 1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{4c} + \frac{3ib^3 \operatorname{PolyLog}\left(4, -1 + \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{4c} \end{aligned}$$

output

```
2*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3*arctanh(-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+3/2*I*b*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*polylog(2,1-2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-3/2*I*b*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*polylog(2,-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+3/2*b^2*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*polylog(3,1-2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-3/2*b^2*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*polylog(3,-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-3/4*I*b^3*polylog(4,1-2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+3/4*I*b^3*polylog(4,-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c
```

Mathematica [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx$$

input

```
Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3/(1 - c^2*x^2), x]
```

output

```
Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3/(1 - c^2*x^2), x]
```

Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 429, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {7232, 5357, 5523, 5529, 5533, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^3}{1 - c^2x^2} dx$$

↓ 7232

$$\int \frac{\sqrt{cx+1} \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}}$$

c
↓ 5357

$$\frac{2 \operatorname{arctanh} \left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \int \frac{\left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^2 \operatorname{arctanh} \left(1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1} \right)}{\frac{1-cx}{cx+1} + 1} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}}}{c}$$

↓ 5523

$$\frac{2 \operatorname{arctanh} \left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left(\frac{1}{2} \int \frac{\left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^2 \log \left(2 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1} \right)}{\frac{1-cx}{cx+1} + 1} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}} - \frac{1}{2} \right)}{c}$$

↓ 5529

$$\frac{2 \operatorname{arctanh} \left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog} \left(2, 1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1} \right) \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right) \right)}{c}$$

↓ 5533

$$\frac{2 \operatorname{arctanh} \left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog} \left(2, 1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1} \right) \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right) \right)}{c}$$

↓ 7164

$$\frac{2 \operatorname{arctanh} \left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}} \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)^3 - 6b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog} \left(2, 1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1} \right) \right) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right) \right)}{c}$$

input

```
Int[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3/(1 - c^2*x^2),x]
```

output

```

-((2*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^3*ArcTanh[1 - 2/(1 + (I*S
qrt[1 - c*x])/Sqrt[1 + c*x]]) - 6*b*((I/2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sq
rt[1 + c*x]])^2*PolyLog[2, 1 - 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) -
I*b*((I/2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[3, 1 - 2/(1
+ (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) + (b*PolyLog[4, 1 - 2/(1 + (I*Sqrt[1
- c*x])/Sqrt[1 + c*x]))/4))/2 + ((-1/2*I)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqr
t[1 + c*x]])^2*PolyLog[2, -1 + 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) +
I*b*((I/2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[3, -1 + 2/(
1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) + (b*PolyLog[4, -1 + 2/(1 + (I*Sqrt[
1 - c*x])/Sqrt[1 + c*x]))/4))/2))/c)

```

Defintions of rubi rules used

rule 5357

```

Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_/(x_), x_Symbol] := Simp[2*(a +
b*ArcTan[c*x])^p*ArcTanh[1 - 2/(1 + I*c*x)], x] - Simp[2*b*c*p Int[(a + b
*ArcTan[c*x])^(p - 1)*(ArcTanh[1 - 2/(1 + I*c*x)]/(1 + c^2*x^2)), x], x] /;
FreeQ[{a, b, c}, x] && IGtQ[p, 1]

```

rule 5523

```

Int[(ArcTanh[u_]*((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_)/((d_) + (e_.)*(x
_)^2), x_Symbol] := Simp[1/2 Int[Log[1 + u]*((a + b*ArcTan[c*x])^p/(d + e
*x^2)), x], x] - Simp[1/2 Int[Log[1 - u]*((a + b*ArcTan[c*x])^p/(d + e*x^
2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] &&
EqQ[u^2 - (1 - 2*(I/(I - c*x)))^2, 0]

```

rule 5529

```

Int[(Log[u_]*((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_)/((d_) + (e_.)*(x_)^2
), x_Symbol] := Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)
), x] + Simp[b*p*(I/2 Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/
(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c
^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]

```

rule 5533

```

Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^p_*PolyLog[k_, u_]/((d_) + (e_.
)*(x_)^2), x_Symbol] := Simp[I*(a + b*ArcTan[c*x])^p*(PolyLog[k + 1, u]/(2*
c*d)), x] - Simp[b*p*(I/2 Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[k + 1
, u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, k}, x] && IGtQ[p, 0] &&
EqQ[e, c^2*d] && EqQ[u^2 - (1 - 2*(I/(I - c*x)))^2, 0]

```

rule 7164 `Int[(u_)*PolyLog[n_, v_], x_Symbol] :=> With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]`

rule 7232 `Int[((a_.) + (b_.)*(F_)[((c_.)*Sqrt[(d_.) + (e_.)*(x_)])/Sqrt[(f_.) + (g_.)*(x_)])^(n_.)/((A_.) + (C_.)*(x_)^2), x_Symbol] :=> Simp[2*e*(g/(C*(e*f - d*g))) Subst[Int[(a + b*F[c*x])^n/x, x], x, Sqrt[d + e*x]/Sqrt[f + g*x]], x] /; FreeQ[{a, b, c, d, e, f, g, A, C, F}, x] && EqQ[C*d*f - A*e*g, 0] && EqQ[e*f + d*g, 0] && IGtQ[n, 0]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1663 vs. $2(360) = 720$.

Time = 1.54 (sec) , antiderivative size = 1664, normalized size of antiderivative = 3.86

method	result	size
default	Expression too large to display	1664
parts	Expression too large to display	1664

input `int((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x,method=_RETURNVERBOSE)`

output

```

-1/2*a^3/c*ln(c*x-1)+1/2*a^3/c*ln(c*x+1)-b^3*(1/c*arctan((-c*x+1)^(1/2)/(c
*x+1)^(1/2))^3*ln(1+(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1
)^(1/2))-3*I/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))^2*polylog(2,-(1+I*(-c*
x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))+6/c*arctan((-c*x+1)^(
1/2)/(c*x+1)^(1/2))*polylog(3,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+
1)/(c*x+1)+1)^(1/2))+6*I/c*polylog(4,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/
((-c*x+1)/(c*x+1)+1)^(1/2))-1/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))^3*ln((
1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1)+1)+3/2*I/c*arctan
((-c*x+1)^(1/2)/(c*x+1)^(1/2))^2*polylog(2,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1
/2))^2/((-c*x+1)/(c*x+1)+1))-3/2/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*po
lylog(3,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1))-3/4*I/
c*polylog(4,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1))+1/
c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))^3*ln(1-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(
1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-3*I/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/
2))^2*polylog(2,(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1
/2))+6/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(3,(1+I*(-c*x+1)^(1/2
)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))+6*I/c*polylog(4,(1+I*(-c*x+1)
^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-3*a*b^2*(1/c*arctan((-c
*x+1)^(1/2)/(c*x+1)^(1/2))^2*ln(1+(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*
x+1)/(c*x+1)+1)^(1/2))-2*I/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polyl...

```

Fricas [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^3}{c^2x^2 - 1} dx$$

input

```

integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x, alg
orithm="fricas")

```

output

```

integral(-(b^3*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))^3 + 3*a*b^2*arctan(sqrt
(-c*x + 1)/sqrt(c*x + 1))^2 + 3*a^2*b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)
) + a^3)/(c^2*x^2 - 1), x)

```

Sympy [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = - \int \frac{a^3}{c^2x^2 - 1} dx - \int \frac{b^3 \operatorname{atan}^3\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

$$- \int \frac{3ab^2 \operatorname{atan}^2\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

$$- \int \frac{3a^2b \operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

input `integrate((a+b*atan((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**3/(-c**2*x**2+1),x)`

output `-Integral(a**3/(c**2*x**2 - 1), x) - Integral(b**3*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))**3/(c**2*x**2 - 1), x) - Integral(3*a*b**2*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1), x) - Integral(3*a**2*b*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)`

Maxima [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int - \frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^3}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x, algorithm="maxima")`

output `1/2*a^3*(log(c*x + 1)/c - log(c*x - 1)/c) + 1/64*(4*(b^3*log(c*x + 1) - b^3*log(-c*x + 1))*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1))^3 - 3*(b^3*log(2)^2*log(c*x + 1) - b^3*log(2)^2*log(-c*x + 1))*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1)) - 64*c*integrate(1/128*(112*b^3*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1))^3 + 384*a*b^2*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1))^2 - 3*(b^3*log(2)^2*log(c*x + 1) - b^3*log(2)^2*log(-c*x + 1) - 4*(b^3*log(c*x + 1) - b^3*log(-c*x + 1))*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1))^2)*sqrt(c*x + 1)*sqrt(-c*x + 1) + 12*(b^3*log(2)^2 + 32*a^2*b)*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1)))/(c^2*x^2 - 1), x)/c`

Giac [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^3}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x, algorithm="giac")`

output `integrate(-(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^3/(c^2*x^2 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx = \int -\frac{\left(a + b \operatorname{atan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^3}{c^2x^2 - 1} dx$$

input `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^3/(c^2*x^2 - 1),x)`

output `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^3/(c^2*x^2 - 1), x)`

Reduce [F]

$$\frac{\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^3}{1 - c^2x^2} dx - 6\left(\int \frac{\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2-1} dx\right) a^2bc - 2\left(\int \frac{\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^3}{c^2x^2-1} dx\right) b^3c - 6\left(\int \frac{\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2}{c^2x^2-1} dx\right) a b^2c - \log(c^2x - c)}{2c}$$

input `int((a+b*atan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^3/(-c^2*x^2+1),x)`

output

```
( - 6*int(atan(sqrt( - c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1),x)*a**2*b*c
- 2*int(atan(sqrt( - c*x + 1)/sqrt(c*x + 1))**3/(c**2*x**2 - 1),x)*b**3*c
- 6*int(atan(sqrt( - c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1),x)*a*b**2
*c - log(c**2*x - c)*a**3 + log(c**2*x + c)*a**3)/(2*c)
```

3.33
$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1-c^2x^2} dx$$

Optimal result	303
Mathematica [F]	304
Rubi [A] (verified)	304
Maple [B] (verified)	307
Fricas [F]	308
Sympy [F]	309
Maxima [F]	309
Giac [F]	310
Mupad [F(-1)]	310
Reduce [F]	310

Optimal result

Integrand size = 40, antiderivative size = 283

$$\begin{aligned} & \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1-c^2x^2} dx \\ &= -\frac{2\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2 \operatorname{arctanh}\left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad + \frac{ib\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad - \frac{ib\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right) \operatorname{PolyLog}\left(2, -1 + \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{c} \\ & \quad + \frac{b^2 \operatorname{PolyLog}\left(3, 1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} - \frac{b^2 \operatorname{PolyLog}\left(3, -1 + \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}}\right)}{2c} \end{aligned}$$

output

```
2*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2*arctanh(-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+I*b*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,1-2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-I*b*(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c+1/2*b^2*polylog(3,1-2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c-1/2*b^2*polylog(3,-1+2/(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2)))/c
```

Mathematica [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx$$

input

```
Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2/(1 - c^2*x^2), x]
```

output

```
Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2/(1 - c^2*x^2), x]
```

Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {7232, 5357, 5523, 5529, 7164}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2}{1 - c^2x^2} dx$$

↓ 7232

$$\int \frac{\sqrt{cx+1} \left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2}{\sqrt{1-cx}} d\frac{\sqrt{1-cx}}{\sqrt{cx+1}}$$

c

↓ 5357

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \int \frac{\left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right) \operatorname{arctanh}\left(1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1}\right) d\frac{\sqrt{1-cx}}{\sqrt{cx+1}}}{\frac{1-cx}{cx+1} + 1}}{c}$$

↓ 5523

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \left(\frac{1}{2} \int \frac{\left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right) \log\left(2 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1}\right) d\frac{\sqrt{1-cx}}{\sqrt{cx+1}} - \frac{1}{2}\right)}{c}}$$

↓ 5529

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1}\right)\right) \left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)\right)}{c}$$

↓ 7164

$$\frac{2\operatorname{arctanh}\left(1 - \frac{2}{1 + \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}}\right) \left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2 - 4b \left(\frac{1}{2} \left(\frac{1}{2} i \operatorname{PolyLog}\left(2, 1 - \frac{2}{\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1}\right)\right) \left(a + b \operatorname{arctan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)\right)}{c}$$

input `Int[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2/(1 - c^2*x^2), x]`

output `-((2*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2*ArcTanh[1 - 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) - 4*b*((I/2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[2, 1 - 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) + (b*PolyLog[3, 1 - 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x])])/4)/2 + ((-1/2*I)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])*PolyLog[2, -1 + 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]) - (b*PolyLog[3, -1 + 2/(1 + (I*Sqrt[1 - c*x])/Sqrt[1 + c*x])])/4)/2))/c`

Definitions of rubi rules used

rule 5357 $\text{Int}[(a + \text{ArcTan}[c \cdot x])^p \cdot \text{ArcTanh}[1 - 2/(1 + I \cdot c \cdot x)], x] - \text{Simp}[2 \cdot b \cdot c \cdot p \cdot \text{Int}[(a + b \cdot \text{ArcTan}[c \cdot x])^{p-1} \cdot \text{ArcTanh}[1 - 2/(1 + I \cdot c \cdot x)]/(1 + c^2 \cdot x^2)], x] /;$
 $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{IGtQ}[p, 1]$

rule 5523 $\text{Int}[(\text{ArcTanh}[u] \cdot (a + \text{ArcTan}[c \cdot x])^p) / ((d + e \cdot x)^2), x] :> \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 + u] \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^p / (d + e \cdot x^2)], x] - \text{Simp}[1/2 \cdot \text{Int}[\text{Log}[1 - u] \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^p / (d + e \cdot x^2)], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[e, c^2 \cdot d] \ \&\& \ \text{EqQ}[u^2 - (1 - 2 \cdot (I / (I - c \cdot x)))^2, 0]$

rule 5529 $\text{Int}[(\text{Log}[u] \cdot (a + \text{ArcTan}[c \cdot x])^p) / ((d + e \cdot x)^2), x] :> \text{Simp}[(-I) \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^p \cdot (\text{PolyLog}[2, 1 - u] / (2 \cdot c \cdot d)), x] + \text{Simp}[b \cdot p \cdot (I/2) \cdot \text{Int}[(a + b \cdot \text{ArcTan}[c \cdot x])^{p-1} \cdot (\text{PolyLog}[2, 1 - u] / (d + e \cdot x^2))], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{EqQ}[e, c^2 \cdot d] \ \&\& \ \text{EqQ}[(1 - u)^2 - (1 - 2 \cdot (I / (I - c \cdot x)))^2, 0]$

rule 7164 $\text{Int}[u \cdot \text{PolyLog}[n, v], x] :> \text{With}\{w = \text{DerivativeDivides}[v, u \cdot v, x]\}, \text{Simp}[w \cdot \text{PolyLog}[n + 1, v], x] /;$
 $! \text{FalseQ}[w] /;$
 $\text{FreeQ}[n, x]$

rule 7232 $\text{Int}[(a + b \cdot (F)[(c \cdot \text{Sqrt}[d + e \cdot x]) / \text{Sqrt}[f + g \cdot x]) \cdot (x)]^n / ((A + C \cdot x)^2), x] :> \text{Simp}[2 \cdot e \cdot (g / (C \cdot (e \cdot f - d \cdot g))) \cdot \text{Subst}[\text{Int}[(a + b \cdot F[c \cdot x])^n / x], x, \text{Sqrt}[d + e \cdot x] / \text{Sqrt}[f + g \cdot x]], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, f, g, A, C, F, x\} \ \&\& \ \text{EqQ}[C \cdot d \cdot f - A \cdot e \cdot g, 0] \ \&\& \ \text{EqQ}[e \cdot f + d \cdot g, 0] \ \&\& \ \text{IGtQ}[n, 0]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 915 vs. $2(240) = 480$.

Time = 0.12 (sec) , antiderivative size = 916, normalized size of antiderivative = 3.24

method	result
default	$-\frac{a^2 \ln(cx-1)}{2c} + \frac{a^2 \ln(cx+1)}{2c} - b^2 \left(\frac{\arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2 \ln\left(1 + \frac{1 + i\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} - \frac{2i \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \operatorname{polylog}\left(2, -\frac{1 + i\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} \right)$
parts	$-\frac{a^2 \ln(cx-1)}{2c} + \frac{a^2 \ln(cx+1)}{2c} - b^2 \left(\frac{\arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2 \ln\left(1 + \frac{1 + i\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} - \frac{2i \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \operatorname{polylog}\left(2, -\frac{1 + i\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} \right)$

input

```
int((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x,method=_RE
TURNVERBOSE)
```

output

```

-1/2*a^2/c*ln(c*x-1)+1/2*a^2/c*ln(c*x+1)-b^2*(1/c*arctan((-c*x+1)^(1/2)/(c
*x+1)^(1/2))^2*ln(1+(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1
)^(1/2))-2*I/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-(1+I*(-c*x+
1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))+2/c*polylog(3,-(1+I*(-
c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-1/c*arctan((-c*x+1
)^(1/2)/(c*x+1)^(1/2))^2*ln((1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)
/(c*x+1)+1)+1)+I/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,-(1+I*(-
c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1))-1/2/c*polylog(3,-(1+I*
(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1))+1/c*arctan((-c*x+1)^(
1/2)/(c*x+1)^(1/2))^2*ln(1-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(
c*x+1)+1)^(1/2))-2*I/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*polylog(2,(1+I
*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))+2/c*polylog(3,(
1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-2*a*b*(1/c*
arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1+(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2
))/((-c*x+1)/(c*x+1)+1)^(1/2))-I/c*polylog(2,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(
1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-1/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)
)*ln((1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1)+1)+1/2*I/c*
polylog(2,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1))+1/c*
arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2
))/((-c*x+1)/(c*x+1)+1)^(1/2))-I/c*polylog(2,(1+I*(-c*x+1)^(1/2)/(c*x+1...

```

Fricas [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^2}{c^2x^2 - 1} dx$$

input

```

integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x, alg
orithm="fricas")

```

output

```

integral(-(b^2*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))^2 + 2*a*b*arctan(sqrt(
-c*x + 1)/sqrt(c*x + 1)) + a^2)/(c^2*x^2 - 1), x)

```

Sympy [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = - \int \frac{a^2}{c^2x^2 - 1} dx - \int \frac{b^2 \operatorname{atan}^2\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx - \int \frac{2ab \operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

input

```
integrate((a+b*atan((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**2/(-c**2*x**2+1),x)
```

output

```
-Integral(a**2/(c**2*x**2 - 1), x) - Integral(b**2*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1), x) - Integral(2*a*b*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)
```

Maxima [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int - \frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^2}{c^2x^2 - 1} dx$$

input

```
integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x, algorithm="maxima")
```

output

```
1/2*a^2*(log(c*x + 1)/c - log(c*x - 1)/c) - 1/32*(b^2*log(2)^2*log(c*x + 1) - b^2*log(2)^2*log(-c*x + 1) - 4*(b^2*log(c*x + 1) - b^2*log(-c*x + 1))*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1))^2 - (b^2*(log(c*x + 1)/c - log(c*x - 1)/c)*log(2)^2 - 64*b^2*integrate(1/16*sqrt(c*x + 1)*sqrt(-c*x + 1)*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))*log(c*x + 1)/(c^2*x^2 - 1), x) + 64*b^2*integrate(1/16*sqrt(c*x + 1)*sqrt(-c*x + 1)*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))*log(-c*x + 1)/(c^2*x^2 - 1), x) - 384*b^2*integrate(1/16*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))^2/(c^2*x^2 - 1), x) - 1024*a*b*integrate(1/16*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c^2*x^2 - 1), x))*c)/c
```

Giac [F]

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int -\frac{\left(b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a\right)^2}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x, algorithm="giac")`

output `integrate(-(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^2/(c^2*x^2 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx = \int -\frac{\left(a + b \operatorname{atan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)\right)^2}{c^2x^2 - 1} dx$$

input `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2/(c^2*x^2 - 1),x)`

output `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2/(c^2*x^2 - 1), x)`

Reduce [F]

$$\frac{\int \frac{\left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}{1 - c^2x^2} dx - 4 \left(\int \frac{\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2-1} dx \right) abc - 2 \left(\int \frac{\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)^2}{c^2x^2-1} dx \right) b^2c - \log(c^2x - c) a^2 + \log(c^2x + c) a^2}{2c}$$

input `int((a+b*atan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2/(-c^2*x^2+1),x)`

output

```
( - 4*int(atan(sqrt( - c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1),x)*a*b*c -  
2*int(atan(sqrt( - c*x + 1)/sqrt(c*x + 1))**2/(c**2*x**2 - 1),x)*b**2*c -  
log(c**2*x - c)*a**2 + log(c**2*x + c)*a**2)/(2*c)
```


3.34
$$\int \frac{a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1-c^2x^2} dx$$

Optimal result	312
Mathematica [A] (verified)	312
Rubi [A] (verified)	313
Maple [B] (verified)	314
Fricas [F]	315
Sympy [F]	315
Maxima [F]	316
Giac [F]	316
Mupad [F(-1)]	317
Reduce [F]	317

Optimal result

Integrand size = 38, antiderivative size = 98

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = -\frac{a \log\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{c} - \frac{ib \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{2c} + \frac{ib \operatorname{PolyLog}\left(2, \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{2c}$$

output

```
-a*ln((-c*x+1)^(1/2)/(c*x+1)^(1/2))/c-1/2*I*b*polylog(2,-I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/c+1/2*I*b*polylog(2,I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/c
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.95

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = -\frac{a \log\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right) + \frac{1}{2}ib \operatorname{PolyLog}\left(2, -\frac{i\sqrt{1-cx}}{\sqrt{1+cx}}\right) - \frac{1}{2}ib \operatorname{PolyLog}\left(2, \frac{i\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{c}$$

input `Integrate[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])/(1 - c^2*x^2),x]`

output `-((a*Log[Sqrt[1 - c*x]/Sqrt[1 + c*x]] + (I/2)*b*PolyLog[2, ((-I)*Sqrt[1 - c*x])/Sqrt[1 + c*x]] - (I/2)*b*PolyLog[2, (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]])/c)`

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.95, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.079$, Rules used = {7232, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{1 - c^2x^2} dx \\
 & \quad \downarrow \text{7232} \\
 & \int \frac{\sqrt{cx+1} \left(a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right) \right)}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}} \\
 & \quad \downarrow \text{5355} \\
 & \frac{\frac{1}{2}ib \int \frac{\sqrt{cx+1} \log\left(1 - \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}} - \frac{1}{2}ib \int \frac{\sqrt{cx+1} \log\left(\frac{i\sqrt{1-cx}}{\sqrt{cx+1}} + 1\right)}{\sqrt{1-cx}} d \frac{\sqrt{1-cx}}{\sqrt{cx+1}} + a \log\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{c} \\
 & \quad \downarrow \text{2838} \\
 & \frac{a \log\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right) + \frac{1}{2}ib \text{PolyLog}\left(2, -\frac{i\sqrt{1-cx}}{\sqrt{cx+1}}\right) - \frac{1}{2}ib \text{PolyLog}\left(2, \frac{i\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{c}
 \end{aligned}$$

input `Int[(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])/(1 - c^2*x^2),x]`

output

```

-((a*Log[Sqrt[1 - c*x]/Sqrt[1 + c*x]] + (I/2)*b*PolyLog[2, ((-I)*Sqrt[1 -
c*x])/Sqrt[1 + c*x]] - (I/2)*b*PolyLog[2, (I*Sqrt[1 - c*x])/Sqrt[1 + c*x]]
)/c)
    
```

Defintions of rubi rules used

rule 2838

```

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
    
```

rule 5355

```

Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] := Simp[a*Log[x], x]
+ (Simp[I*(b/2) Int[Log[1 - I*c*x]/x, x], x] - Simp[I*(b/2) Int[Log[1
+ I*c*x]/x, x], x]) /; FreeQ[{a, b, c}, x]
    
```

rule 7232

```

Int[((a_.) + (b_.)*(F_)[((c_.)*Sqrt[(d_.) + (e_.)*(x_)])/Sqrt[(f_.) + (g_.)
*(x_)])^(n_.))/(A_.) + (C_.)*(x_)^2, x_Symbol] := Simp[2*e*(g/(C*(e*f - d
*g))) Subst[Int[(a + b*F[c*x])^n/x, x], x, Sqrt[d + e*x]/Sqrt[f + g*x]],
x] /; FreeQ[{a, b, c, d, e, f, g, A, C, F}, x] && EqQ[C*d*f - A*e*g, 0] &&
EqQ[e*f + d*g, 0] && IGtQ[n, 0]
    
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 367 vs. 2(78) = 156.

Time = 0.11 (sec) , antiderivative size = 368, normalized size of antiderivative = 3.76

method	result
default	$-\frac{a \ln(cx-1)}{2c} + \frac{a \ln(cx+1)}{2c} - b \left(\frac{\arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \ln\left(1 + \frac{1 + \frac{i\sqrt{-cx+1}}{\sqrt{cx+1}}}{\sqrt{\frac{-cx+1}{cx+1} + 1}}\right)}{c} - \frac{i \operatorname{polylog}\left(2, -\frac{1 + \frac{i\sqrt{-cx+1}}{\sqrt{cx+1}}}{\sqrt{\frac{-cx+1}{cx+1} + 1}}\right)}{c} - \frac{\arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} \right)$
parts	$-\frac{a \ln(cx-1)}{2c} + \frac{a \ln(cx+1)}{2c} - b \left(\frac{\arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) \ln\left(1 + \frac{1 + \frac{i\sqrt{-cx+1}}{\sqrt{cx+1}}}{\sqrt{\frac{-cx+1}{cx+1} + 1}}\right)}{c} - \frac{i \operatorname{polylog}\left(2, -\frac{1 + \frac{i\sqrt{-cx+1}}{\sqrt{cx+1}}}{\sqrt{\frac{-cx+1}{cx+1} + 1}}\right)}{c} - \frac{\arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c} \right)$

input `int((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x,method=_RETURNERVERBOSE)`

output `-1/2*a/c*ln(c*x-1)+1/2*a/c*ln(c*x+1)-b*(1/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1+(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-I/c*polylog(2,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-1/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln((1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))^2/((-c*x+1)/(c*x+1)+1)+1)+1/2*I/c*polylog(2,-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)+1/c*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))*ln(1-(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2))-I/c*polylog(2,(1+I*(-c*x+1)^(1/2)/(c*x+1)^(1/2))/((-c*x+1)/(c*x+1)+1)^(1/2)))`

Fricas [F]

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \int -\frac{b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x,algorithm="fricas")`

output `integral(-(b*arctan(sqrt(-c*x + 1))/sqrt(c*x + 1)) + a)/(c^2*x^2 - 1), x)`

Sympy [F]

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = -\int \frac{a}{c^2x^2 - 1} dx - \int \frac{b \operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

input `integrate((a+b*atan((-c*x+1)**(1/2)/(c*x+1)**(1/2)))/(-c**2*x**2+1),x)`

output `-Integral(a/(c**2*x**2 - 1), x) - Integral(b*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)`

Maxima [F]

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \int -\frac{b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x, algorith="maxima")`

output `1/2*a*(log(c*x + 1)/c - log(c*x - 1)/c) + 1/2*((log(c*x + 1) - log(-c*x + 1))*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1)) - 2*c*integrate(1/2*(e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1))*log(c*x + 1) - e^(1/2*log(c*x + 1) + 1/2*log(-c*x + 1))*log(-c*x + 1))/((c^2*x^2 - 1)*(c*x + 1) - (c^2*x^2 - 1)*(c*x - 1)), x))*b/c`

Giac [F]

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \int -\frac{b \arctan\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right) + a}{c^2x^2 - 1} dx$$

input `integrate((a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1),x, algorith="giac")`

output `integrate(-(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)/(c^2*x^2 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx = \int -\frac{a + b \operatorname{atan}\left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}}\right)}{c^2x^2 - 1} dx$$

input `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))/(c^2*x^2 - 1), x)`

output `int(-(a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))/(c^2*x^2 - 1), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{a + b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)}{1 - c^2x^2} dx \\ & - 2 \left(\int \frac{\operatorname{atan}\left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}}\right)}{c^2x^2-1} dx \right) bc - \log(c^2x - c)a + \log(c^2x + c)a \\ & = \frac{\hspace{10em}}{2c} \end{aligned}$$

input `int((a+b*atan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))/(-c^2*x^2+1), x)`

output `(- 2*int(atan(sqrt(-c*x + 1)/sqrt(c*x + 1))/(c**2*x**2 - 1), x)*b*c - lo
g(c**2*x - c)*a + log(c**2*x + c)*a)/(2*c)`

$$3.35 \quad \int \frac{1}{(1-c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx$$

Optimal result	318
Mathematica [N/A]	318
Rubi [N/A]	319
Maple [N/A]	319
Fricas [N/A]	320
Sympy [N/A]	320
Maxima [N/A]	321
Giac [N/A]	321
Mupad [N/A]	322
Reduce [N/A]	322

Optimal result

Integrand size = 40, antiderivative size = 40

$$\int \frac{1}{(1-c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \text{Int} \left(\frac{1}{(1-c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)}, x \right)$$

output

```
Defer(Int)(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)
```

Mathematica [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{1}{(1-c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \int \frac{1}{(1-c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx$$

input

```
Integrate[1/((1 - c^2*x^2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])),x]
```

output

```
Integrate[1/((1 - c^2*x^2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])), x
]
```

Rubi [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)} dx$$

↓ 7234

$$\int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{cx+1}} \right) \right)} dx$$

input

```
Int[1/((1 - c^2*x^2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])),x]
```

output

```
$Aborted
```

Maple [N/A]

Not integrable

Time = 0.65 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \frac{1}{(-c^2x^2 + 1) \left(a + b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) \right)} dx$$

input

```
int(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)
```


output `int(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.20

$$\int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \int -\frac{1}{(c^2x^2 - 1) \left(b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x, algorithm="fricas")`

output `integral(-1/(a*c^2*x^2 + (b*c^2*x^2 - b)*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) - a), x)`

Sympy [N/A]

Not integrable

Time = 3.31 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.52

$$\begin{aligned} & \int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx \\ &= - \int \frac{1}{ac^2x^2 - a + bc^2x^2 \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) - b \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)} dx \end{aligned}$$

input `integrate(1/(-c**2*x**2+1)/(a+b*atan((-c*x+1)**(1/2)/(c*x+1)**(1/2))),x)`

output `-Integral(1/(a*c**2*x**2 - a + b*c**2*x**2*atan(sqrt(-c*x + 1)/sqrt(c*x + 1)) - b*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))), x)`

Maxima [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \int -\frac{1}{(c^2x^2 - 1) \left(b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x, algorithm="maxima")`

output `-integrate(1/((c^2*x^2 - 1)*(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)), x)`

Giac [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)} dx = \int -\frac{1}{(c^2x^2 - 1) \left(b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x, algorithm="giac")`

output `integrate(-1/((c^2*x^2 - 1)*(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)), x)`

Mupad [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)} dx = - \int \frac{1}{\left(a + b \operatorname{atan} \left(\frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right) (c^2 x^2 - 1)} dx$$

input `int(-1/((a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))*(c^2*x^2 - 1)),x)`

output `-int(1/((a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))*(c^2*x^2 - 1)), x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.58

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)} dx$$

$$= - \left(\int \frac{1}{\operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) b c^2 x^2 - \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) b + a c^2 x^2 - a} dx \right)$$

input `int(1/(-c^2*x^2+1)/(a+b*atan((-c*x+1)^(1/2)/(c*x+1)^(1/2))),x)`

output `- int(1/(atan(sqrt(-c*x + 1)/sqrt(c*x + 1))*b*c**2*x**2 - atan(sqrt(-c*x + 1)/sqrt(c*x + 1))*b + a*c**2*x**2 - a),x)`

$$3.36 \quad \int \frac{1}{(1-c^2x^2) \left(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2} dx$$

Optimal result	323
Mathematica [N/A]	323
Rubi [N/A]	324
Maple [N/A]	325
Fricas [N/A]	325
Sympy [N/A]	326
Maxima [N/A]	326
Giac [N/A]	327
Mupad [N/A]	327
Reduce [N/A]	328

Optimal result

Integrand size = 40, antiderivative size = 40

$$\int \frac{1}{(1-c^2x^2) \left(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2} dx$$

$$= \text{Int} \left(\frac{1}{(1-c^2x^2) \left(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2}, x \right)$$

output `Defer(Int)(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

Mathematica [N/A]

Not integrable

Time = 0.89 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05

$$\int \frac{1}{(1-c^2x^2) \left(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2} dx = \int \frac{1}{(1-c^2x^2) \left(a+b \arctan\left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}}\right)\right)^2} dx$$

input `Integrate[1/((1 - c^2*x^2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2), x]`

output `Integrate[1/((1 - c^2*x^2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2), x]`

Rubi [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right)^2} dx$$

↓ 7234

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right)^2} dx$$

input `Int[1/((1 - c^2*x^2)*(a + b*ArcTan[Sqrt[1 - c*x]/Sqrt[1 + c*x]])^2),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.61 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \frac{1}{(-c^2x^2 + 1) \left(a + b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) \right)^2} dx$$

input `int(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

output `int(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 91, normalized size of antiderivative = 2.28

$$\int \frac{1}{(1 - c^2x^2) \left(a + b \arctan \left(\frac{\sqrt{1-cx}}{\sqrt{1+cx}} \right) \right)^2} dx = \int -\frac{1}{(c^2x^2 - 1) \left(b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)^2} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x, algorithm="fricas")`

output `integral(-1/(a^2*c^2*x^2 + (b^2*c^2*x^2 - b^2)*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))^2 - a^2 + 2*(a*b*c^2*x^2 - a*b)*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1))), x)`

Sympy [N/A]

Not integrable

Time = 7.17 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.15

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx =$$

$$- \int \frac{1}{a^2 c^2 x^2 - a^2 + 2abc^2 x^2 \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) - 2ab \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + b^2 c^2 x^2 \operatorname{atan}^2 \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) - b^2 \operatorname{atan}^2 \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)} dx$$

input `integrate(1/(-c**2*x**2+1)/(a+b*atan((-c*x+1)**(1/2)/(c*x+1)**(1/2)))**2,x)`

output `-Integral(1/(a**2*c**2*x**2 - a**2 + 2*a*b*c**2*x**2*atan(sqrt(-c*x + 1)/sqrt(c*x + 1)) - 2*a*b*atan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + b**2*c**2*x**2*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))**2 - b**2*atan(sqrt(-c*x + 1)/sqrt(c*x + 1))**2), x)`

Maxima [N/A]

Not integrable

Time = 0.43 (sec) , antiderivative size = 163, normalized size of antiderivative = 4.08

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx = \int - \frac{1}{(c^2 x^2 - 1) \left(b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)^2} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x, algorithm="maxima")`

output `2*(2*(b^2*c^2*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1)) + a*b*c^2)*sqrt(c*x + 1)*sqrt(-c*x + 1)*integrate(1/2*x/((a*b*c^2*x^2 - a*b + (b^2*c^2*x^2 - b^2)*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1)))*sqrt(c*x + 1)*sqrt(-c*x + 1)), x) + 1)/((b^2*c*arctan2(sqrt(-c*x + 1), sqrt(c*x + 1)) + a*b*c)*sqrt(c*x + 1)*sqrt(-c*x + 1))`

Giac [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.95

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx = \int -\frac{1}{(c^2 x^2 - 1) \left(b \arctan \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) + a \right)^2} dx$$

input `integrate(1/(-c^2*x^2+1)/(a+b*arctan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x, algorithm="giac")`

output `integrate(-1/((c^2*x^2 - 1)*(b*arctan(sqrt(-c*x + 1)/sqrt(c*x + 1)) + a)^2), x)`

Mupad [N/A]

Not integrable

Time = 1.42 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.98

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx = - \int \frac{1}{\left(a + b \operatorname{atan} \left(\frac{\sqrt{1 - cx}}{\sqrt{cx + 1}} \right) \right)^2 (c^2 x^2 - 1)} dx$$

input `int(-1/((a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2*(c^2*x^2 - 1)),x)`

output `-int(1/((a + b*atan((1 - c*x)^(1/2)/(c*x + 1)^(1/2)))^2*(c^2*x^2 - 1)), x)`

Reduce [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 123, normalized size of antiderivative = 3.08

$$\int \frac{1}{(1 - c^2 x^2) \left(a + b \arctan \left(\frac{\sqrt{1 - cx}}{\sqrt{1 + cx}} \right) \right)^2} dx =$$

$$- \left(\int \frac{1}{\operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)^2 b^2 c^2 x^2 - \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right)^2 b^2 + 2 \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) ab c^2 x^2 - 2 \operatorname{atan} \left(\frac{\sqrt{-cx+1}}{\sqrt{cx+1}} \right) ab + a^2} dx \right)$$

input

```
int(1/(-c^2*x^2+1)/(a+b*atan((-c*x+1)^(1/2)/(c*x+1)^(1/2)))^2,x)
```

output

```
- int(1/(atan(sqrt(-c*x+1)/sqrt(c*x+1))**2*b**2*c**2*x**2 - atan(sqrt(-c*x+1)/sqrt(c*x+1))**2*b**2 + 2*atan(sqrt(-c*x+1)/sqrt(c*x+1))*a*b*c**2*x**2 - 2*atan(sqrt(-c*x+1)/sqrt(c*x+1))*a*b + a**2*c**2*x**2 - a**2),x)
```

3.37 $\int x^m \arctan(\tan(a + bx)) dx$

Optimal result	329
Mathematica [A] (verified)	329
Rubi [A] (verified)	330
Maple [A] (verified)	331
Fricas [A] (verification not implemented)	331
Sympy [B] (verification not implemented)	331
Maxima [A] (verification not implemented)	332
Giac [F]	333
Mupad [F(-1)]	333
Reduce [B] (verification not implemented)	333

Optimal result

Integrand size = 11, antiderivative size = 37

$$\int x^m \arctan(\tan(a + bx)) dx = -\frac{bx^{2+m}}{2 + 3m + m^2} + \frac{x^{1+m} \arctan(\tan(a + bx))}{1 + m}$$

output `-b*x^(2+m)/(m^2+3*m+2)+x^(1+m)*arctan(tan(b*x+a))/(1+m)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int x^m \arctan(\tan(a + bx)) dx = x^m \left(\frac{bx^2}{2 + m} + \frac{x(-bx + \arctan(\tan(a + bx)))}{1 + m} \right)$$

input `Integrate[x^m*ArcTan[Tan[a + b*x]],x]`

output `x^m*((b*x^2)/(2 + m) + (x*(-(b*x) + ArcTan[Tan[a + b*x]])))/(1 + m)`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \arctan(\tan(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{x^{m+1} \arctan(\tan(a + bx))}{m + 1} - \frac{b \int x^{m+1} dx}{m + 1}$$

$$\downarrow \text{15}$$

$$\frac{x^{m+1} \arctan(\tan(a + bx))}{m + 1} - \frac{bx^{m+2}}{(m + 1)(m + 2)}$$

input `Int[x^m*ArcTan[Tan[a + b*x]],x]`

output `-((b*x^(2 + m))/((1 + m)*(2 + m))) + (x^(1 + m)*ArcTan[Tan[a + b*x]])/(1 + m)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1))) Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11

method	result
default	$\frac{b x^2 e^{m \ln(x)}}{2+m} + \frac{(\arctan(\tan(bx+a))-bx)x e^{m \ln(x)}}{1+m}$
parallelrisc	$-\frac{b x^m x^2 - 2 \arctan(\tan(bx+a)) x x^m - x x^m \arctan(\tan(bx+a)) m}{(1+m)(2+m)}$
risc	$-\frac{i x x^m \ln(e^{i(bx+a)})}{1+m} - x \left(4bx + 2\pi \operatorname{csgn}(ie^{2i(bx+a)})^3 + 2\pi \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)^3 + \pi m \operatorname{csgn}(ie^{2i(bx+a)})^3 + \pi m \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)^3 \right)$

input `int(x^m*arctan(tan(b*x+a)),x,method=_RETURNVERBOSE)`output `b/(2+m)*x^2*exp(m*ln(x))+(arctan(tan(b*x+a))-b*x)/(1+m)*x*exp(m*ln(x))`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int x^m \arctan(\tan(a + bx)) dx = \frac{((bm + b)x^2 + (am + 2a)x)x^m}{m^2 + 3m + 2}$$

input `integrate(x^m*arctan(tan(b*x+a)),x, algorithm="fricas")`output `((b*m + b)*x^2 + (a*m + 2*a)*x)*x^m/(m^2 + 3*m + 2)`**Sympy [B] (verification not implemented)**Leaf count of result is larger than twice the leaf count of optimal. 158 vs. $2(31) = 62$.

Time = 0.81 (sec) , antiderivative size = 158, normalized size of antiderivative = 4.27

$$\int x^m \arctan(\tan(a + bx)) dx$$

$$= \begin{cases} b \log(x) - \frac{\operatorname{atan}(\tan(a+bx)) + \pi \left\lfloor \frac{a+bx-\frac{\pi}{2}}{\pi} \right\rfloor}{x} & \text{for } m = -2 \\ -bx \log(x) + bx + \left(\operatorname{atan}(\tan(a + bx)) + 2\pi \left\lfloor \frac{a+bx-\frac{\pi}{2}}{\pi} \right\rfloor \right) \log(x) & \text{for } m = -1 \\ -\frac{bx^2 x^m}{m^2+3m+2} + \frac{m x x^m \left(\operatorname{atan}(\tan(a+bx)) + \pi \left\lfloor \frac{a+bx-\frac{\pi}{2}}{\pi} \right\rfloor \right)}{m^2+3m+2} + \frac{2 x x^m \left(\operatorname{atan}(\tan(a+bx)) + \pi \left\lfloor \frac{a+bx-\frac{\pi}{2}}{\pi} \right\rfloor \right)}{m^2+3m+2} & \text{otherwise} \end{cases}$$

input `integrate(x**m*atan(tan(b*x+a)),x)`

output `Piecewise((b*log(x) - (atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi)) /x, Eq(m, -2)), (-b*x*log(x) + b*x + (atan(tan(a + b*x)) + 2*pi*floor((a + b*x - pi/2)/pi))*log(x), Eq(m, -1)), (-b*x**2*x**m/(m**2 + 3*m + 2) + m*x*x**m*(atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))/(m**2 + 3*m + 2) + 2*x*x**m*(atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))/(m**2 + 3*m + 2), True))`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int x^m \arctan(\tan(a + bx)) dx = -\frac{bx^2 x^m}{(m+2)(m+1)} + \frac{x^{m+1} \arctan(\tan(bx + a))}{m+1}$$

input `integrate(x^m*arctan(tan(b*x+a)),x, algorithm="maxima")`

output `-b*x^2*x^m/((m + 2)*(m + 1)) + x^(m + 1)*arctan(tan(b*x + a))/(m + 1)`

Giac [F]

$$\int x^m \arctan(\tan(a + bx)) dx = \int x^m \arctan(\tan(bx + a)) dx$$

input `integrate(x^m*arctan(tan(b*x+a)),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int x^m \arctan(\tan(a + bx)) dx = \int x^m \operatorname{atan}(\tan(a + bx)) dx$$

input `int(x^m*atan(tan(a + b*x)),x)`

output `int(x^m*atan(tan(a + b*x)), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int x^m \arctan(\tan(a + bx)) dx = \frac{x^m x (\operatorname{atan}(\tan(bx + a)) m + 2 \operatorname{atan}(\tan(bx + a)) - bx)}{m^2 + 3m + 2}$$

input `int(x^m*atan(tan(b*x+a)),x)`

output `(x**m*x*(atan(tan(a + b*x))*m + 2*atan(tan(a + b*x)) - b*x)/(m**2 + 3*m + 2)`

3.38 $\int x^2 \arctan(\tan(a + bx)) dx$

Optimal result	334
Mathematica [A] (verified)	334
Rubi [A] (verified)	335
Maple [A] (verified)	336
Fricas [A] (verification not implemented)	336
Sympy [A] (verification not implemented)	337
Maxima [B] (verification not implemented)	337
Giac [F]	338
Mupad [B] (verification not implemented)	338
Reduce [F]	338

Optimal result

Integrand size = 11, antiderivative size = 23

$$\int x^2 \arctan(\tan(a + bx)) dx = -\frac{bx^4}{12} + \frac{1}{3}x^3 \arctan(\tan(a + bx))$$

output

```
-1/12*b*x^4+1/3*x^3*arctan(tan(b*x+a))
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^2 \arctan(\tan(a + bx)) dx = -\frac{1}{12}x^3(bx - 4 \arctan(\tan(a + bx)))$$

input

```
Integrate[x^2*ArcTan[Tan[a + b*x]],x]
```

output

```
-1/12*(x^3*(b*x - 4*ArcTan[Tan[a + b*x]]))
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(\tan(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{1}{3}x^3 \arctan(\tan(a + bx)) - \frac{b \int x^3 dx}{3}$$

$$\downarrow \text{15}$$

$$\frac{1}{3}x^3 \arctan(\tan(a + bx)) - \frac{bx^4}{12}$$

input `Int[x^2*ArcTan[Tan[a + b*x]],x]`

output `-1/12*(b*x^4) + (x^3*ArcTan[Tan[a + b*x]])/3`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n])`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result
default	$-\frac{bx^4}{12} + \frac{x^3 \arctan(\tan(bx+a))}{3}$
parallelrisc	$-\frac{bx^4}{12} + \frac{x^3 \arctan(\tan(bx+a))}{3}$
parts	$-\frac{bx^4}{12} + \frac{x^3 \arctan(\tan(bx+a))}{3}$
risc	$-\frac{ix^3 \ln(e^{i(bx+a)})}{3} - \frac{\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}(ie^{2i(bx+a)}) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)}{12} + \frac{\pi x^3 \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)}{12}$

input `int(x^2*arctan(tan(b*x+a)),x,method=_RETURNVERBOSE)`output `-1/12*b*x^4+1/3*x^3*arctan(tan(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int x^2 \arctan(\tan(a + bx)) dx = \frac{1}{4} bx^4 + \frac{1}{3} ax^3$$

input `integrate(x^2*arctan(tan(b*x+a)),x, algorithm="fricas")`output `1/4*b*x^4 + 1/3*a*x^3`

Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int x^2 \arctan(\tan(a + bx)) dx = -\frac{bx^4}{12} + \frac{x^3 \left(\operatorname{atan}(\tan(a + bx)) + \pi \left\lfloor \frac{a + bx - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{3}$$

input `integrate(x**2*atan(tan(b*x+a)),x)`

output `-b*x**4/12 + x**3*(atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))/3`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 81 vs. 2(19) = 38.

Time = 0.04 (sec) , antiderivative size = 81, normalized size of antiderivative = 3.52

$$\int x^2 \arctan(\tan(a + bx)) dx$$

$$= \frac{4 \left((bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2 \right) \arctan(\tan(bx+a))}{b^2} - \frac{(bx+a)^4 - 4(bx+a)^3 a + 6(bx+a)^2 a^2}{b^2}$$

$$12b$$

input `integrate(x^2*arctan(tan(b*x+a)),x, algorithm="maxima")`

output `1/12*(4*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctan(tan(b*x + a))/b^2 - ((b*x + a)^4 - 4*(b*x + a)^3*a + 6*(b*x + a)^2*a^2)/b^2)/b`

Giac [F]

$$\int x^2 \arctan(\tan(a + bx)) dx = \int x^2 \arctan(\tan(bx + a)) dx$$

input `integrate(x^2*arctan(tan(b*x+a)),x, algorithm="giac")`

output `sage0*x`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \arctan(\tan(a + bx)) dx = \frac{x^3 \operatorname{atan}(\tan(a + bx))}{3} - \frac{bx^4}{12}$$

input `int(x^2*atan(tan(a + b*x)),x)`

output `(x^3*atan(tan(a + b*x)))/3 - (b*x^4)/12`

Reduce [F]

$$\int x^2 \arctan(\tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a)) x^2 dx$$

input `int(x^2*atan(tan(b*x+a)),x)`

output `int(atan(tan(a + b*x))*x**2,x)`

3.39 $\int x \arctan(\tan(a + bx)) dx$

Optimal result	339
Mathematica [A] (verified)	339
Rubi [A] (verified)	340
Maple [A] (verified)	341
Fricas [A] (verification not implemented)	341
Sympy [A] (verification not implemented)	342
Maxima [B] (verification not implemented)	342
Giac [F]	342
Mupad [B] (verification not implemented)	343
Reduce [F]	343

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x \arctan(\tan(a + bx)) dx = -\frac{bx^3}{6} + \frac{1}{2}x^2 \arctan(\tan(a + bx))$$

output `-1/6*b*x^3+1/2*x^2*arctan(tan(b*x+a))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x \arctan(\tan(a + bx)) dx = -\frac{1}{6}x^2(bx - 3 \arctan(\tan(a + bx)))$$

input `Integrate[x*ArcTan[Tan[a + b*x]],x]`

output `-1/6*(x^2*(b*x - 3*ArcTan[Tan[a + b*x]]))`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5694, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(\tan(a + bx)) dx$$

$$\downarrow 5694$$

$$\frac{1}{2}x^2 \arctan(\tan(a + bx)) - \frac{1}{2}ib \int -ix^2 dx$$

$$\downarrow 15$$

$$\frac{1}{2}x^2 \arctan(\tan(a + bx)) - \frac{bx^3}{6}$$

input `Int[x*ArcTan[Tan[a + b*x]],x]`

output `-1/6*(b*x^3) + (x^2*ArcTan[Tan[a + b*x]])/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 5694 `Int[ArcTan[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tan[a + b*x]]/(f*(m + 1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result
default	$-\frac{x^3 b}{6} + \frac{x^2 \arctan(\tan(bx+a))}{2}$
parallelrisc	$-\frac{x^3 b}{6} + \frac{x^2 \arctan(\tan(bx+a))}{2}$
parts	$-\frac{x^3 b}{6} + \frac{x^2 \arctan(\tan(bx+a))}{2}$
risc	$-\frac{ix^2 \ln(e^{i(bx+a)})}{2} - \frac{\pi x^2 \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}(ie^{2i(bx+a)}) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)}{8} + \frac{\pi x^2 \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)}{8}$

input `int(x*arctan(tan(b*x+a)),x,method=_RETURNVERBOSE)`output `-1/6*x^3*b+1/2*x^2*arctan(tan(b*x+a))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int x \arctan(\tan(a + bx)) dx = \frac{1}{3} bx^3 + \frac{1}{2} ax^2$$

input `integrate(x*arctan(tan(b*x+a)),x, algorithm="fricas")`output `1/3*b*x^3 + 1/2*a*x^2`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int x \arctan(\tan(a + bx)) dx = -\frac{bx^3}{6} + \frac{x^2 \left(\operatorname{atan}(\tan(a + bx)) + \pi \left\lfloor \frac{a + bx - \frac{\pi}{2}}{\pi} \right\rfloor \right)}{2}$$

input `integrate(x*atan(tan(b*x+a)),x)`

output `-b*x**3/6 + x**2*(atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. 2(19) = 38.

Time = 0.03 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.48

$$\int x \arctan(\tan(a + bx)) dx = \frac{3 \left((bx+a)^2 - 2(bx+a)a \right) \arctan(\tan(bx+a)) - \frac{(bx+a)^3 - 3(bx+a)^2 a}{b}}{6b}$$

input `integrate(x*arctan(tan(b*x+a)),x, algorithm="maxima")`

output `1/6*(3*((b*x + a)^2 - 2*(b*x + a)*a)*arctan(tan(b*x + a))/b - ((b*x + a)^3 - 3*(b*x + a)^2*a)/b)/b`

Giac [F]

$$\int x \arctan(\tan(a + bx)) dx = \int x \arctan(\tan(bx + a)) dx$$

input `integrate(x*arctan(tan(b*x+a)),x, algorithm="giac")`

output `sage0*x`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \arctan(\tan(a + bx)) dx = \frac{x^2 \operatorname{atan}(\tan(a + bx))}{2} - \frac{bx^3}{6}$$

input `int(x*atan(tan(a + b*x)),x)`

output `(x^2*atan(tan(a + b*x)))/2 - (b*x^3)/6`

Reduce [F]

$$\int x \arctan(\tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a)) x dx$$

input `int(x*atan(tan(b*x+a)),x)`

output `int(atan(tan(a + b*x))*x,x)`

3.40 $\int \arctan(\tan(a + bx)) dx$

Optimal result	344
Mathematica [A] (verified)	344
Rubi [A] (verified)	345
Maple [A] (verified)	346
Fricas [A] (verification not implemented)	346
Sympy [B] (verification not implemented)	347
Maxima [A] (verification not implemented)	347
Giac [A] (verification not implemented)	347
Mupad [B] (verification not implemented)	348
Reduce [B] (verification not implemented)	348

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \arctan(\tan(a + bx)) dx = \frac{\arctan(\tan(a + bx))^2}{2b}$$

output `1/2*arctan(tan(b*x+a))^2/b`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \arctan(\tan(a + bx)) dx = -\frac{bx^2}{2} + x \arctan(\tan(a + bx))$$

input `Integrate[ArcTan[Tan[a + b*x]],x]`

output `-1/2*(b*x^2) + x*ArcTan[Tan[a + b*x]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(\tan(a + bx)) dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \arctan(\tan(a + bx)) d \arctan(\tan(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{\arctan(\tan(a + bx))^2}{2b}$$

input `Int[ArcTan[Tan[a + b*x]],x]`

output `ArcTan[Tan[a + b*x]]^2/(2*b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\arctan(\tan(bx+a))^2}{2b}$
default	$\frac{\arctan(\tan(bx+a))^2}{2b}$
parallelrisch	$-\frac{bx^2}{2} + x \arctan(\tan(bx+a))$
parts	$-\frac{bx^2}{2} + x \arctan(\tan(bx+a))$
risch	$-ix \ln(e^{i(bx+a)}) - \frac{\pi x \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}(ie^{2i(bx+a)}) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)}{4} + \frac{\pi x \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right)}{4}$

input `int(arctan(tan(b*x+a)),x,method=_RETURNVERBOSE)`output `1/2*arctan(tan(b*x+a))^2/b`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \arctan(\tan(a + bx)) dx = \frac{1}{2} bx^2 + ax$$

input `integrate(arctan(tan(b*x+a)),x, algorithm="fricas")`output `1/2*b*x^2 + a*x`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(12) = 24$.

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.62

$$\int \arctan(\tan(a + bx)) dx = \begin{cases} \frac{\left(\operatorname{atan}(\tan(a + bx)) + \pi \left\lfloor \frac{a + bx - \frac{\pi}{2}}{\pi} \right\rfloor\right)^2}{2b} & \text{for } b \neq 0 \\ x \left(\operatorname{atan}(\tan(a)) + \pi \left\lfloor \frac{a - \frac{\pi}{2}}{\pi} \right\rfloor\right) & \text{otherwise} \end{cases}$$

input `integrate(atan(tan(b*x+a)),x)`

output `Piecewise(((atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))**2/(2*b), Ne(b, 0)), (x*(atan(tan(a)) + pi*floor((a - pi/2)/pi)), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \arctan(\tan(a + bx)) dx = \frac{(bx + a)^2}{2b}$$

input `integrate(arctan(tan(b*x+a)),x, algorithm="maxima")`

output `1/2*(b*x + a)^2/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int \arctan(\tan(a + bx)) dx = \frac{1}{2} bx^2 - \pi x \left[\frac{bx + a}{\pi} + \frac{1}{2} \right] + ax$$

input `integrate(arctan(tan(b*x+a)),x, algorithm="giac")`

output `1/2*b*x^2 - pi*x*floor((b*x + a)/pi + 1/2) + a*x`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \arctan(\tan(a + bx)) dx = x \operatorname{atan}(\tan(a + bx)) - \frac{bx^2}{2}$$

input `int(atan(tan(a + b*x)),x)`

output `x*atan(tan(a + b*x)) - (b*x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \arctan(\tan(a + bx)) dx = \frac{\operatorname{atan}(\tan(bx + a))^2}{2b}$$

input `int(atan(tan(b*x+a)),x)`

output `atan(tan(a + b*x))**2/(2*b)`

3.41 $\int \frac{\arctan(\tan(a+bx))}{x} dx$

Optimal result	349
Mathematica [A] (verified)	349
Rubi [A] (verified)	350
Maple [A] (verified)	351
Fricas [A] (verification not implemented)	351
Sympy [A] (verification not implemented)	351
Maxima [A] (verification not implemented)	352
Giac [F]	352
Mupad [F(-1)]	353
Reduce [F]	353

Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = bx - (bx - \arctan(\tan(a + bx))) \log(x)$$

output `b*x-(b*x-arctan(tan(b*x+a)))*ln(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = bx + (-bx + \arctan(\tan(a + bx))) \log(x)$$

input `Integrate[ArcTan[Tan[a + b*x]]/x,x]`

output `b*x + -(b*x) + ArcTan[Tan[a + b*x]]*Log[x]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(\tan(a + bx))}{x} dx$$

$$\downarrow \text{2589}$$

$$bx - (bx - \arctan(\tan(a + bx))) \int \frac{1}{x} dx$$

$$\downarrow \text{14}$$

$$bx - \log(x)(bx - \arctan(\tan(a + bx)))$$

input `Int[ArcTan[Tan[a + b*x]]/x,x]`

output `b*x - (b*x - ArcTan[Tan[a + b*x]])*Log[x]`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

method	result
default	$\ln(x) \arctan(\tan(bx + a)) - b(\ln(x)x - x)$
parts	$\ln(x) \arctan(\tan(bx + a)) - b(\ln(x)x - x)$
risch	$-i \ln(x) \ln(e^{i(bx+a)}) - \ln(x)bx + bx - \frac{\pi \left(\operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}(ie^{2i(bx+a)}) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right) - \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right) \right)}{2}$

input `int(arctan(tan(b*x+a))/x,x,method=_RETURNVERBOSE)`output `ln(x)*arctan(tan(b*x+a))-b*(ln(x)*x-x)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.38

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = bx + a \log(x)$$

input `integrate(arctan(tan(b*x+a))/x,x, algorithm="fricas")`output `b*x + a*log(x)`**Sympy [A] (verification not implemented)**

Time = 0.39 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = -bx \log(x) + bx + \left(\operatorname{atan}(\tan(a + bx)) + \pi \left\lfloor \frac{a + bx - \frac{\pi}{2}}{\pi} \right\rfloor \right) \log(x)$$

input `integrate(atan(tan(b*x+a))/x,x)`

output `-b*x*log(x) + b*x + (atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))*log(x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.00

$$\int \frac{\arctan(\tan(a + bx))}{x} dx$$

$$= \frac{b \arctan(\tan(bx + a)) \log(bx) + (bx - (bx + a)) \log(bx) + a \log(bx) + a}{b}$$

input `integrate(arctan(tan(b*x+a))/x,x, algorithm="maxima")`

output `(b*arctan(tan(b*x + a))*log(b*x) + (b*x - (b*x + a)*log(b*x) + a*log(b*x) + a)*b)/b`

Giac [F]

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = \int \frac{\arctan(\tan(bx + a))}{x} dx$$

input `integrate(arctan(tan(b*x+a))/x,x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\tan(a + bx))}{x} dx$$

input `int(atan(tan(a + b*x))/x,x)`output `int(atan(tan(a + b*x))/x, x)`**Reduce [F]**

$$\int \frac{\arctan(\tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\tan(bx + a))}{x} dx$$

input `int(atan(tan(b*x+a))/x,x)`output `int(atan(tan(a + b*x))/x,x)`

3.42 $\int x^m \arctan(\cot(a + bx)) dx$

Optimal result	354
Mathematica [A] (verified)	354
Rubi [A] (verified)	355
Maple [A] (verified)	356
Fricas [A] (verification not implemented)	356
Sympy [B] (verification not implemented)	357
Maxima [A] (verification not implemented)	357
Giac [A] (verification not implemented)	358
Mupad [F(-1)]	358
Reduce [B] (verification not implemented)	358

Optimal result

Integrand size = 11, antiderivative size = 36

$$\int x^m \arctan(\cot(a + bx)) dx = \frac{bx^{2+m}}{2 + 3m + m^2} + \frac{x^{1+m} \arctan(\cot(a + bx))}{1 + m}$$

output

```
b*x^(2+m)/(m^2+3*m+2)+x^(1+m)*(1/2*Pi-arccot(cot(b*x+a)))/(1+m)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.86

$$\int x^m \arctan(\cot(a + bx)) dx = \frac{x^{1+m}(bx + (2 + m) \arctan(\cot(a + bx)))}{(1 + m)(2 + m)}$$

input

```
Integrate[x^m*ArcTan[Cot[a + b*x]],x]
```

output

```
(x^(1 + m)*(b*x + (2 + m)*ArcTan[Cot[a + b*x]]))/((1 + m)*(2 + m))
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^m \arctan(\cot(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{b \int x^{m+1} dx}{m+1} + \frac{x^{m+1} \arctan(\cot(a + bx))}{m+1}$$

$$\downarrow \text{15}$$

$$\frac{x^{m+1} \arctan(\cot(a + bx))}{m+1} + \frac{bx^{m+2}}{(m+1)(m+2)}$$

input `Int[x^m*ArcTan[Cot[a + b*x]],x]`

output `(b*x^(2 + m))/((1 + m)*(2 + m)) + (x^(1 + m)*ArcTan[Cot[a + b*x]])/(1 + m)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.56

method	result
default	$\frac{\pi x^{1+m}}{2m+2} - \frac{b x^2 e^{m \ln(x)}}{2+m} - \frac{(\operatorname{arccot}(\cot(bx+a)) - bx) x e^{m \ln(x)}}{1+m}$
parts	$\frac{\pi x^{1+m}}{2m+2} - \frac{b x^2 e^{m \ln(x)}}{2+m} - \frac{(\operatorname{arccot}(\cot(bx+a)) - bx) x e^{m \ln(x)}}{1+m}$
parallelrisch	$\frac{2b x^m x^2 + 2\pi x x^m - 4 \operatorname{arccot}(\cot(bx+a)) x x^m + \pi x x^m m - 2x x^m \operatorname{arccot}(\cot(bx+a)) m}{2(1+m)(2+m)}$
risch	$\frac{ix x^m \ln(e^{i(bx+a)})}{1+m} + \frac{x \left(4\pi + 4bx + 2\pi m + 2\pi \operatorname{csgn}(ie^{2i(bx+a)})^3 + \pi m \operatorname{csgn}(ie^{2i(bx+a)})^3 + 2\pi \operatorname{csgn}(ie^{i(bx+a)})^2 \operatorname{csgn}(ie^{2i(bx+a)}) \right)}{2(1+m)(2+m)}$

input `int(x^m*(1/2*Pi-arccot(cot(b*x+a))),x,method=_RETURNVERBOSE)`

output `1/2*Pi*x^(1+m)/(1+m)-b/(2+m)*x^2*exp(m*ln(x))-(arccot(cot(b*x+a))-b*x)/(1+m)*x*exp(m*ln(x))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.17

$$\int x^m \arctan(\cot(a + bx)) dx = -\frac{(2(bm + b)x^2 - (\pi(m + 2) - 2am - 4a)x)x^m}{2(m^2 + 3m + 2)}$$

input `integrate(x^m*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="fricas")`

output `-1/2*(2*(b*m + b)*x^2 - (pi*(m + 2) - 2*a*m - 4*a)*x)*x^m/(m^2 + 3*m + 2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(34) = 68$.

Time = 2.05 (sec) , antiderivative size = 160, normalized size of antiderivative = 4.44

$$\int x^m \arctan(\cot(a + bx)) dx$$

$$= \begin{cases} -b \log(x) + \frac{\operatorname{acot}(\cot(a+bx))}{x} - \frac{\pi}{2x} & \text{for } m = -2 \\ bx \log(x) - bx - \log(x) \operatorname{acot}(\cot(a + bx)) + \frac{\pi \log(x)}{2} & \text{for } m = -1 \\ \frac{2bx^2x^m}{2m^2+6m+4} - \frac{2mxx^m \operatorname{acot}(\cot(a+bx))}{2m^2+6m+4} + \frac{\pi mxx^m}{2m^2+6m+4} - \frac{4xx^m \operatorname{acot}(\cot(a+bx))}{2m^2+6m+4} + \frac{2\pi xx^m}{2m^2+6m+4} & \text{otherwise} \end{cases}$$

input `integrate(x**m*(1/2*pi-acot(cot(b*x+a))),x)`

output `Piecewise((-b*log(x) + acot(cot(a + b*x))/x - pi/(2*x), Eq(m, -2)), (b*x*log(x) - b*x - log(x)*acot(cot(a + b*x)) + pi*log(x)/2, Eq(m, -1)), (2*b*x**2*x**m/(2*m**2 + 6*m + 4) - 2*m*x*x**m*acot(cot(a + b*x))/(2*m**2 + 6*m + 4) + pi*m*x*x**m/(2*m**2 + 6*m + 4) - 4*x*x**m*acot(cot(a + b*x))/(2*m**2 + 6*m + 4) + 2*pi*x*x**m/(2*m**2 + 6*m + 4), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.11

$$\int x^m \arctan(\cot(a + bx)) dx = -\frac{bx^{m+2}}{m+2} + \frac{\pi x^{m+1}}{2(m+1)} - \frac{ax^{m+1}}{m+1}$$

input `integrate(x^m*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="maxima")`

output `-b*x^(m + 2)/(m + 2) + 1/2*pi*x^(m + 1)/(m + 1) - a*x^(m + 1)/(m + 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.72

$$\int x^m \arctan(\cot(a + bx)) dx$$

$$= -\frac{2bm x^2 x^m - \pi m x x^m + 2am x x^m + 2bx^2 x^m - 2\pi x x^m + 4ax x^m}{2(m^2 + 3m + 2)}$$

input `integrate(x^m*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="giac")`output `-1/2*(2*b*m*x^2*x^m - pi*m*x*x^m + 2*a*m*x*x^m + 2*b*x^2*x^m - 2*pi*x*x^m + 4*a*x*x^m)/(m^2 + 3*m + 2)`**Mupad [F(-1)]**

Timed out.

$$\int x^m \arctan(\cot(a + bx)) dx = \int x^m \left(\frac{\Pi}{2} - \operatorname{acot}(\cot(a + bx)) \right) dx$$

input `int(x^m*(Pi/2 - acot(cot(a + b*x))),x)`output `int(x^m*(Pi/2 - acot(cot(a + b*x))), x)`**Reduce [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.31

$$\int x^m \arctan(\cot(a + bx)) dx$$

$$= \frac{x^m x (-2 \operatorname{acot}(\cot(bx + a)) m - 4 \operatorname{acot}(\cot(bx + a)) + 2bx + m\pi + 2\pi)}{2m^2 + 6m + 4}$$

input `int(x^m*(1/2*Pi-acot(cot(b*x+a))),x)`

output
$$\frac{(x^m \cdot (-2 \operatorname{acot}(\cot(a + b \cdot x)) \cdot m - 4 \operatorname{acot}(\cot(a + b \cdot x)) + 2 \cdot b \cdot x + m \cdot \pi + 2 \cdot \pi))}{2 \cdot (m^2 + 3 \cdot m + 2)}$$

3.43 $\int x^2 \arctan(\cot(a + bx)) dx$

Optimal result	360
Mathematica [A] (verified)	360
Rubi [A] (verified)	361
Maple [A] (verified)	362
Fricas [A] (verification not implemented)	362
Sympy [A] (verification not implemented)	363
Maxima [A] (verification not implemented)	363
Giac [A] (verification not implemented)	363
Mupad [B] (verification not implemented)	364
Reduce [F]	364

Optimal result

Integrand size = 11, antiderivative size = 23

$$\int x^2 \arctan(\cot(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \arctan(\cot(a + bx))$$

output `1/12*b*x^4+1/3*x^3*(1/2*Pi-arccot(cot(b*x+a)))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x^2 \arctan(\cot(a + bx)) dx = \frac{1}{12}x^3(bx + 4 \arctan(\cot(a + bx)))$$

input `Integrate[x^2*ArcTan[Cot[a + b*x]],x]`

output `(x^3*(b*x + 4*ArcTan[Cot[a + b*x]]))/12`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2599, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(\cot(a + bx)) dx$$

$$\downarrow \text{2599}$$

$$\frac{b \int x^3 dx}{3} + \frac{1}{3} x^3 \arctan(\cot(a + bx))$$

$$\downarrow \text{15}$$

$$\frac{1}{3} x^3 \arctan(\cot(a + bx)) + \frac{bx^4}{12}$$

input `Int[x^2*ArcTan[Cot[a + b*x]],x]`

output `(b*x^4)/12 + (x^3*ArcTan[Cot[a + b*x]])/3`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2599 `Int[(u_)^(m_)*(v_)^(n_.), x_Symbol] := With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[u^(m + 1)*(v^n/(a*(m + 1))), x] - Simp[b*(n/(a*(m + 1)))] Int[u^(m + 1)*v^(n - 1), x], x] /; NeQ[b*u - a*v, 0] /; FreeQ[{m, n}, x] && PiecewiseLinearQ[u, v, x] && NeQ[m, -1] && ((LtQ[m, -1] && GtQ[n, 0]) && !(ILtQ[m + n, -2] && (FractionQ[m] || GeQ[2*n + m + 1, 0]))) || (IGtQ[n, 0] && IGtQ[m, 0] && LeQ[n, m]) || (IGtQ[n, 0] && !IntegerQ[m]) || (ILtQ[m, 0] && !IntegerQ[n]))`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

method	result
parallelrisch	$-\frac{x^3 \operatorname{arccot}(\cot(bx+a))}{3} + \frac{\pi x^3}{6} + \frac{bx^4}{12}$
default	$\frac{\pi x^3}{6} - \frac{x^3 \operatorname{arccot}(\cot(bx+a))}{3} - \frac{-(bx+a)^4 + a(bx+a)^3 - \frac{3a^2(bx+a)^2}{2} + a^3(bx+a)}{3b^3}$
parts	$\frac{\pi x^3}{6} - \frac{x^3 \operatorname{arccot}(\cot(bx+a))}{3} - \frac{-(bx+a)^4 + a(bx+a)^3 - \frac{3a^2(bx+a)^2}{2} + a^3(bx+a)}{3b^3}$
risch	$\frac{ix^3 \ln(e^{i(bx+a)})}{3} + \frac{\pi x^3 \operatorname{csgn}(ie^{i(bx+a)})^2 \operatorname{csgn}(ie^{2i(bx+a)})}{12} - \frac{\pi x^3 \operatorname{csgn}(ie^{i(bx+a)}) \operatorname{csgn}(ie^{2i(bx+a)})^2}{6} + \frac{\pi x^3 \operatorname{csgn}(ie^{2i(bx+a)})}{12}$

input `int(x^2*(1/2*Pi-arccot(cot(b*x+a))),x,method=_RETURNVERBOSE)`

output `-1/3*x^3*arccot(cot(b*x+a))+1/6*Pi*x^3+1/12*b*x^4`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int x^2 \arctan(\cot(a + bx)) dx = -\frac{1}{4}bx^4 + \frac{1}{6}(\pi - 2a)x^3$$

input `integrate(x^2*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="fricas")`

output `-1/4*b*x^4 + 1/6*(pi - 2*a)*x^3`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int x^2 \arctan(\cot(a + bx)) dx = \frac{bx^4}{12} - \frac{x^3 \operatorname{acot}(\cot(a + bx))}{3} + \frac{\pi x^3}{6}$$

input `integrate(x**2*(1/2*pi-acot(cot(b*x+a))),x)`output `b*x**4/12 - x**3*acot(cot(a + b*x))/3 + pi*x**3/6`**Maxima [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int x^2 \arctan(\cot(a + bx)) dx = -\frac{1}{4} bx^4 + \frac{1}{6} (\pi - 2a)x^3$$

input `integrate(x^2*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="maxima")`output `-1/4*b*x^4 + 1/6*(pi - 2*a)*x^3`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x^2 \arctan(\cot(a + bx)) dx = -\frac{1}{4} bx^4 + \frac{1}{6} \pi x^3 - \frac{1}{3} ax^3$$

input `integrate(x^2*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="giac")`output `-1/4*b*x^4 + 1/6*pi*x^3 - 1/3*a*x^3`

Mupad [B] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int x^2 \arctan(\cot(a + bx)) dx = \frac{\pi x^3}{6} + \frac{bx^4}{12} - \frac{x^3 \operatorname{acot}(\cot(a + bx))}{3}$$

input `int(x^2*(Pi/2 - acot(cot(a + b*x))),x)`output `(Pi*x^3)/6 + (b*x^4)/12 - (x^3*acot(cot(a + b*x)))/3`**Reduce [F]**

$$\int x^2 \arctan(\cot(a + bx)) dx = -\left(\int \operatorname{acot}(\cot(bx + a)) x^2 dx\right) + \frac{\pi x^3}{6}$$

input `int(x^2*(1/2*Pi-acot(cot(b*x+a))),x)`output `(- 6*int(acot(cot(a + b*x))*x**2,x) + pi*x**3)/6`

3.44 $\int x \arctan(\cot(a + bx)) dx$

Optimal result	365
Mathematica [A] (verified)	365
Rubi [A] (verified)	366
Maple [A] (verified)	367
Fricas [A] (verification not implemented)	367
Sympy [A] (verification not implemented)	368
Maxima [A] (verification not implemented)	368
Giac [A] (verification not implemented)	368
Mupad [B] (verification not implemented)	369
Reduce [F]	369

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int x \arctan(\cot(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \arctan(\cot(a + bx))$$

output `1/6*b*x^3+1/2*x^2*(1/2*Pi-arccot(cot(b*x+a)))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int x \arctan(\cot(a + bx)) dx = \frac{1}{6}x^2(bx + 3 \arctan(\cot(a + bx)))$$

input `Integrate[x*ArcTan[Cot[a + b*x]],x]`

output `(x^2*(b*x + 3*ArcTan[Cot[a + b*x]]))/6`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {5696, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(\cot(a + bx)) dx$$

$$\downarrow \text{5696}$$

$$\frac{1}{2}x^2 \arctan(\cot(a + bx)) - \frac{1}{2}ib \int ix^2 dx$$

$$\downarrow \text{15}$$

$$\frac{1}{2}x^2 \arctan(\cot(a + bx)) + \frac{bx^3}{6}$$

input `Int[x*ArcTan[Cot[a + b*x]],x]`

output `(b*x^3)/6 + (x^2*ArcTan[Cot[a + b*x]])/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 5696 `Int[ArcTan[(c_.) + Cot[(a_.) + (b_.)*(x_)]]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Cot[a + b*x]]/(f*(m + 1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - I*d)^2, -1]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

method	result
parallelrisch	$\frac{x^3 b}{6} - \frac{x^2 \operatorname{arccot}(\cot(bx+a))}{2} + \frac{\pi x^2}{4}$
default	$\frac{\pi x^2}{4} - \frac{x^2 \operatorname{arccot}(\cot(bx+a))}{2} - \frac{-(bx+a)^3 + a(bx+a)^2 - a^2(bx+a)}{2b^2}$
parts	$\frac{\pi x^2}{4} - \frac{x^2 \operatorname{arccot}(\cot(bx+a))}{2} - \frac{-(bx+a)^3 + a(bx+a)^2 - a^2(bx+a)}{2b^2}$
risch	$\frac{ix^2 \ln(e^{i(bx+a)})}{2} + \frac{\pi x^2 \operatorname{csgn}(ie^{i(bx+a)})^2 \operatorname{csgn}(ie^{2i(bx+a)})}{8} - \frac{\pi x^2 \operatorname{csgn}(ie^{i(bx+a)}) \operatorname{csgn}(ie^{2i(bx+a)})^2}{4} + \frac{\pi x^2 \operatorname{csgn}(ie^{2i(bx+a)})}{8}$

input `int(x*(1/2*Pi-arccot(cot(b*x+a))),x,method=_RETURNVERBOSE)`

output `1/6*x^3*b-1/2*x^2*arccot(cot(b*x+a))+1/4*Pi*x^2`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int x \arctan(\cot(a + bx)) dx = -\frac{1}{3} bx^3 + \frac{1}{4} (\pi - 2a)x^2$$

input `integrate(x*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="fricas")`

output `-1/3*b*x^3 + 1/4*(pi - 2*a)*x^2`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int x \arctan(\cot(a + bx)) dx = \frac{bx^3}{6} - \frac{x^2 \operatorname{acot}(\cot(a + bx))}{2} + \frac{\pi x^2}{4}$$

input `integrate(x*(1/2*pi-acot(cot(b*x+a))),x)`output `b*x**3/6 - x**2*acot(cot(a + b*x))/2 + pi*x**2/4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int x \arctan(\cot(a + bx)) dx = -\frac{1}{3}bx^3 + \frac{1}{4}(\pi - 2a)x^2$$

input `integrate(x*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="maxima")`output `-1/3*b*x^3 + 1/4*(pi - 2*a)*x^2`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int x \arctan(\cot(a + bx)) dx = -\frac{1}{3}bx^3 + \frac{1}{4}\pi x^2 - \frac{1}{2}ax^2$$

input `integrate(x*(1/2*pi-arccot(cot(b*x+a))),x, algorithm="giac")`output `-1/3*b*x^3 + 1/4*pi*x^2 - 1/2*a*x^2`

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int x \arctan(\cot(a + bx)) dx = \frac{\pi x^2}{4} + \frac{bx^3}{6} - \frac{x^2 \operatorname{acot}(\cot(a + bx))}{2}$$

input `int(x*(Pi/2 - acot(cot(a + b*x))),x)`output `(Pi*x^2)/4 + (b*x^3)/6 - (x^2*acot(cot(a + b*x)))/2`**Reduce [F]**

$$\int x \arctan(\cot(a + bx)) dx = -\left(\int \operatorname{acot}(\cot(bx + a)) x dx\right) + \frac{\pi x^2}{4}$$

input `int(x*(1/2*Pi-acot(cot(b*x+a))),x)`output `(- 4*int(acot(cot(a + b*x))*x,x) + pi*x**2)/4`

3.45 $\int \arctan(\cot(a + bx)) dx$

Optimal result	370
Mathematica [A] (verified)	370
Rubi [A] (verified)	371
Maple [A] (verified)	372
Fricas [A] (verification not implemented)	372
Sympy [A] (verification not implemented)	373
Maxima [A] (verification not implemented)	373
Giac [A] (verification not implemented)	373
Mupad [B] (verification not implemented)	374
Reduce [B] (verification not implemented)	374

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \arctan(\cot(a + bx)) dx = -\frac{\arctan(\cot(a + bx))^2}{2b}$$

output

```
-1/2*(1/2*Pi-arccot(cot(b*x+a)))^2/b
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \arctan(\cot(a + bx)) dx = \frac{bx^2}{2} + x \arctan(\cot(a + bx))$$

input

```
Integrate[ArcTan[Cot[a + b*x]],x]
```

output

```
(b*x^2)/2 + x*ArcTan[Cot[a + b*x]]
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(\cot(a + bx)) dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \arctan(\cot(a + bx)) d \arctan(\cot(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{\arctan(\cot(a + bx))^2}{2b}$$

input `Int[ArcTan[Cot[a + b*x]],x]`

output `-1/2*ArcTan[Cot[a + b*x]]^2/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

method	result
parallelsch	$\frac{bx^2}{2} - x \operatorname{arccot}(\cot(bx+a)) + \frac{\pi x}{2}$
derivativedivides	$\frac{-\pi(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))) - \operatorname{arccot}(\cot(bx+a))^2}{2b}$
default	$\frac{\pi x}{2} - \frac{-(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))) \operatorname{arccot}(\cot(bx+a)) - \frac{(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a)))^2}{2}}{b}$
parts	$\frac{\pi x}{2} - \frac{-(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))) \operatorname{arccot}(\cot(bx+a)) - \frac{(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a)))^2}{2}}{b}$
risch	$\frac{\pi x}{2} + ix \ln(e^{i(bx+a)}) + \frac{\pi x \operatorname{csgn}(ie^{i(bx+a)})^2 \operatorname{csgn}(ie^{2i(bx+a)})}{4} - \frac{\pi x \operatorname{csgn}(ie^{i(bx+a)}) \operatorname{csgn}(ie^{2i(bx+a)})^2}{2} + \pi$

input `int(1/2*Pi-arccot(cot(b*x+a)),x,method=_RETURNVERBOSE)`output `1/2*b*x^2-x*arccot(cot(b*x+a))+1/2*Pi*x`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \arctan(\cot(a+bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}(\pi - 2a)x$$

input `integrate(1/2*pi-arccot(cot(b*x+a)),x, algorithm="fricas")`output `-1/2*b*x^2 + 1/2*(pi - 2*a)*x`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

$$\int \arctan(\cot(a + bx)) dx = \frac{\pi x}{2} - \begin{cases} \frac{\operatorname{acot}^2(\cot(a+bx))}{2b} & \text{for } b \neq 0 \\ x \operatorname{acot}(\cot(a)) & \text{otherwise} \end{cases}$$

input `integrate(1/2*pi-acot(cot(b*x+a)),x)`output `pi*x/2 - Piecewise((acot(cot(a + b*x))**2/(2*b), Ne(b, 0)), (x*acot(cot(a)), True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \arctan(\cot(a + bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}\pi x - ax$$

input `integrate(1/2*pi-arccot(cot(b*x+a)),x, algorithm="maxima")`output `-1/2*b*x^2 + 1/2*pi*x - a*x`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \arctan(\cot(a + bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}\pi x - ax$$

input `integrate(1/2*pi-arccot(cot(b*x+a)),x, algorithm="giac")`output `-1/2*b*x^2 + 1/2*pi*x - a*x`

Mupad [B] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \arctan(\cot(a + bx)) dx = \frac{\Pi x}{2} - x \operatorname{acot}(\cot(a + bx)) + \frac{b x^2}{2}$$

input `int(Pi/2 - acot(cot(a + b*x)),x)`output `(Pi*x)/2 - x*acot(cot(a + b*x)) + (b*x^2)/2`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \arctan(\cot(a + bx)) dx = \frac{-\operatorname{acot}(\cot(bx + a))^2 + b\pi x}{2b}$$

input `int(1/2*Pi-acot(cot(b*x+a)),x)`output `(- acot(cot(a + b*x))**2 + b*pi*x)/(2*b)`

3.46 $\int \frac{\arctan(\cot(a+bx))}{x} dx$

Optimal result	375
Mathematica [A] (verified)	375
Rubi [A] (verified)	376
Maple [A] (verified)	377
Fricas [A] (verification not implemented)	377
Sympy [A] (verification not implemented)	377
Maxima [A] (verification not implemented)	378
Giac [A] (verification not implemented)	378
Mupad [F(-1)]	378
Reduce [F]	379

Optimal result

Integrand size = 11, antiderivative size = 19

$$\int \frac{\arctan(\cot(a+bx))}{x} dx = -bx + (bx + \arctan(\cot(a+bx))) \log(x)$$

output

```
-b*x+(b*x+1/2*Pi-arc cot(cot(b*x+a)))*ln(x)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(\cot(a+bx))}{x} dx = -bx + (bx + \arctan(\cot(a+bx))) \log(x)$$

input

```
Integrate[ArcTan[Cot[a + b*x]]/x,x]
```

output

```
-(b*x) + (b*x + ArcTan[Cot[a + b*x]])*Log[x]
```


Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2589, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(\cot(a + bx))}{x} dx$$

$$\downarrow \text{2589}$$

$$(\arctan(\cot(a + bx)) + bx) \int \frac{1}{x} dx - bx$$

$$\downarrow \text{14}$$

$$\log(x)(\arctan(\cot(a + bx)) + bx) - bx$$

input `Int[ArcTan[Cot[a + b*x]]/x,x]`

output `-(b*x) + (b*x + ArcTan[Cot[a + b*x]])*Log[x]`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 2589 `Int[(v_)/(u_), x_Symbol] :> With[{a = Simplify[D[u, x]], b = Simplify[D[v, x]]}, Simp[b*(x/a), x] - Simp[(b*u - a*v)/a Int[1/u, x], x] /; NeQ[b*u - a*v, 0]] /; PiecewiseLinearQ[u, v, x]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

method	result
default	$\frac{\pi \ln(x)}{2} - bx - (\operatorname{arccot}(\cot(bx + a)) - bx) \ln(x)$
parts	$\frac{\pi \ln(x)}{2} - bx - (\operatorname{arccot}(\cot(bx + a)) - bx) \ln(x)$
risch	$i \ln(x) \ln(e^{i(bx+a)}) + \ln(x) bx - bx + \frac{\pi \left(\operatorname{csgn}(ie^{i(bx+a)})^2 \operatorname{csgn}(ie^{2i(bx+a)}) - 2 \operatorname{csgn}(ie^{i(bx+a)}) \operatorname{csgn}(ie^{2i(bx+a)})^2 + \dots \right)}{2}$

input `int((1/2*Pi-arccot(cot(b*x+a)))/x,x,method=_RETURNVERBOSE)`

output `1/2*Pi*ln(x)-b*x-(arccot(cot(b*x+a))-b*x)*ln(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{\arctan(\cot(a + bx))}{x} dx = -bx + \frac{1}{2} (\pi - 2a) \log(x)$$

input `integrate((1/2*pi-arccot(cot(b*x+a)))/x,x, algorithm="fricas")`

output `-b*x + 1/2*(pi - 2*a)*log(x)`

Sympy [A] (verification not implemented)

Time = 1.62 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

$$\int \frac{\arctan(\cot(a + bx))}{x} dx = bx \log(x) - bx - \log(x) \operatorname{acot}(\cot(a + bx)) + \frac{\pi \log(x)}{2}$$

input `integrate((1/2*pi-acot(cot(b*x+a)))/x,x)`

output `b*x*log(x) - b*x - log(x)*acot(cot(a + b*x)) + pi*log(x)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.74

$$\int \frac{\arctan(\cot(a + bx))}{x} dx = -bx + \frac{1}{2} (\pi - 2a) \log(x)$$

input `integrate((1/2*pi-arccot(cot(b*x+a)))/x,x, algorithm="maxima")`

output `-b*x + 1/2*(pi - 2*a)*log(x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{\arctan(\cot(a + bx))}{x} dx = -bx + \frac{1}{2} (\pi - 2a) \log(|x|)$$

input `integrate((1/2*pi-arccot(cot(b*x+a)))/x,x, algorithm="giac")`

output `-b*x + 1/2*(pi - 2*a)*log(abs(x))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(\cot(a + bx))}{x} dx = \int \frac{\frac{\pi}{2} - \operatorname{acot}(\cot(a + bx))}{x} dx$$

input `int((Pi/2 - acot(cot(a + b*x)))/x,x)`

output `int((Pi/2 - acot(cot(a + b*x)))/x, x)`

Reduce [F]

$$\int \frac{\arctan(\cot(a + bx))}{x} dx = - \left(\int \frac{\operatorname{acot}(\cot(bx + a))}{x} dx \right) + \frac{\log(x) \pi}{2}$$

input `int((1/2*Pi-acot(cot(b*x+a)))/x,x)`

output `(- 2*int(acot(cot(a + b*x))/x,x) + log(x)*pi)/2`

3.47 $\int \arctan(\tan(a + bx)) dx$

Optimal result	380
Mathematica [A] (verified)	380
Rubi [A] (verified)	381
Maple [A] (verified)	382
Fricas [A] (verification not implemented)	382
Sympy [B] (verification not implemented)	383
Maxima [A] (verification not implemented)	383
Giac [A] (verification not implemented)	383
Mupad [B] (verification not implemented)	384
Reduce [B] (verification not implemented)	384

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \arctan(\tan(a + bx)) dx = \frac{\arctan(\tan(a + bx))^2}{2b}$$

output

```
1/2*arctan(tan(b*x+a))^2/b
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \arctan(\tan(a + bx)) dx = -\frac{bx^2}{2} + x \arctan(\tan(a + bx))$$

input

```
Integrate[ArcTan[Tan[a + b*x]],x]
```

output

```
-1/2*(b*x^2) + x*ArcTan[Tan[a + b*x]]
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(\tan(a + bx)) dx$$

$$\downarrow 2588$$

$$\frac{\int \arctan(\tan(a + bx)) d \arctan(\tan(a + bx))}{b}$$

$$\downarrow 15$$

$$\frac{\arctan(\tan(a + bx))^2}{2b}$$

input `Int[ArcTan[Tan[a + b*x]],x]`

output `ArcTan[Tan[a + b*x]]^2/(2*b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result
derivativedivides	$\frac{\arctan(\tan(bx+a))^2}{2b}$
default	$\frac{\arctan(\tan(bx+a))^2}{2b}$
parallelrisch	$-\frac{bx^2}{2} + x \arctan(\tan(bx+a))$
parts	$-\frac{bx^2}{2} + x \arctan(\tan(bx+a))$
risch	$-ix \ln(e^{i(bx+a)}) - \frac{\pi x \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right) \operatorname{csgn}(ie^{2i(bx+a)}) \operatorname{csgn}\left(\frac{ie^{2i(bx+a)}}{e^{2i(bx+a)}+1}\right)}{4} + \frac{\pi x \operatorname{csgn}\left(\frac{i}{e^{2i(bx+a)}+1}\right)}{4}$

input `int(arctan(tan(b*x+a)),x,method=_RETURNVERBOSE)`output `1/2*arctan(tan(b*x+a))^2/b`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

$$\int \arctan(\tan(a + bx)) dx = \frac{1}{2} bx^2 + ax$$

input `integrate(arctan(tan(b*x+a)),x, algorithm="fricas")`output `1/2*b*x^2 + a*x`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(12) = 24$.

Time = 0.09 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.62

$$\int \arctan(\tan(a + bx)) dx = \begin{cases} \frac{\left(\operatorname{atan}(\tan(a + bx)) + \pi \left\lfloor \frac{a + bx - \frac{\pi}{2}}{\pi} \right\rfloor\right)^2}{2b} & \text{for } b \neq 0 \\ x \left(\operatorname{atan}(\tan(a)) + \pi \left\lfloor \frac{a - \frac{\pi}{2}}{\pi} \right\rfloor\right) & \text{otherwise} \end{cases}$$

input `integrate(atan(tan(b*x+a)),x)`

output `Piecewise(((atan(tan(a + b*x)) + pi*floor((a + b*x - pi/2)/pi))**2/(2*b), Ne(b, 0)), (x*(atan(tan(a)) + pi*floor((a - pi/2)/pi)), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.75

$$\int \arctan(\tan(a + bx)) dx = \frac{(bx + a)^2}{2b}$$

input `integrate(arctan(tan(b*x+a)),x, algorithm="maxima")`

output `1/2*(b*x + a)^2/b`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.62

$$\int \arctan(\tan(a + bx)) dx = \frac{1}{2} bx^2 - \pi x \left[\frac{bx + a}{\pi} + \frac{1}{2} \right] + ax$$

input `integrate(arctan(tan(b*x+a)),x, algorithm="giac")`

output `1/2*b*x^2 - pi*x*floor((b*x + a)/pi + 1/2) + a*x`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \arctan(\tan(a + bx)) dx = x \operatorname{atan}(\tan(a + bx)) - \frac{bx^2}{2}$$

input `int(atan(tan(a + b*x)),x)`

output `x*atan(tan(a + b*x)) - (b*x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \arctan(\tan(a + bx)) dx = \frac{\operatorname{atan}(\tan(bx + a))^2}{2b}$$

input `int(atan(tan(b*x+a)),x)`

output `atan(tan(a + b*x))**2/(2*b)`

3.48 $\int x^2 \arctan(c + d \tan(a + bx)) dx$

Optimal result	385
Mathematica [A] (verified)	386
Rubi [A] (verified)	387
Maple [C] (warning: unable to verify)	393
Fricas [B] (verification not implemented)	393
Sympy [F(-1)]	394
Maxima [F]	395
Giac [F]	395
Mupad [F(-1)]	396
Reduce [F]	396

Optimal result

Integrand size = 15, antiderivative size = 403

$$\begin{aligned}
 \int x^2 \arctan(c + d \tan(a + bx)) dx &= \frac{1}{3} x^3 \arctan(c + d \tan(a + bx)) \\
 &+ \frac{1}{6} i x^3 \log \left(1 + \frac{(1 + ic + d)e^{2ia+2ibx}}{1 + ic - d} \right) \\
 &- \frac{1}{6} i x^3 \log \left(1 + \frac{(c + i(1 - d))e^{2ia+2ibx}}{c + i(1 + d)} \right) \\
 &+ \frac{x^2 \operatorname{PolyLog} \left(2, -\frac{(1+ic+d)e^{2ia+2ibx}}{1+ic-d} \right)}{4b} \\
 &- \frac{x^2 \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(1+d)} \right)}{4b} \\
 &+ \frac{ix \operatorname{PolyLog} \left(3, -\frac{(1+ic+d)e^{2ia+2ibx}}{1+ic-d} \right)}{4b^2} \\
 &- \frac{ix \operatorname{PolyLog} \left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(1+d)} \right)}{4b^2} \\
 &- \frac{\operatorname{PolyLog} \left(4, -\frac{(1+ic+d)e^{2ia+2ibx}}{1+ic-d} \right)}{8b^3} \\
 &+ \frac{\operatorname{PolyLog} \left(4, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(1+d)} \right)}{8b^3}
 \end{aligned}$$

output

```
1/3*x^3*arctan(c+d*tan(b*x+a))+1/6*I*x^3*ln(1+(1+I*c+d)*exp(2*I*a+2*I*b*x)
/(1+I*c-d))-1/6*I*x^3*ln(1+(c+I*(1-d))*exp(2*I*a+2*I*b*x)/(c+I*(1+d)))+1/4
*x^2*polylog(2,-(1+I*c+d)*exp(2*I*a+2*I*b*x)/(1+I*c-d))/b-1/4*x^2*polylog(
2,-(c+I*(1-d))*exp(2*I*a+2*I*b*x)/(c+I*(1+d)))/b+1/4*I*x*polylog(3,-(1+I*c
+d)*exp(2*I*a+2*I*b*x)/(1+I*c-d))/b^2-1/4*I*x*polylog(3,-(c+I*(1-d))*exp(2
*I*a+2*I*b*x)/(c+I*(1+d)))/b^2-1/8*polylog(4,-(1+I*c+d)*exp(2*I*a+2*I*b*x)
/(1+I*c-d))/b^3+1/8*polylog(4,-(c+I*(1-d))*exp(2*I*a+2*I*b*x)/(c+I*(1+d))
)/b^3
```

Mathematica [A] (verified)

Time = 2.88 (sec) , antiderivative size = 371, normalized size of antiderivative = 0.92

$$\int x^2 \arctan(c + d \tan(a + bx)) dx$$

$$= \frac{8b^3x^3 \arctan(c + d \tan(a + bx)) + 4ib^3x^3 \log\left(1 + \frac{(c+i(-1+d))e^{-2i(a+bx)}}{c-i(1+d)}\right) - 4ib^3x^3 \log\left(1 + \frac{(c+i(1+d))e^{-2i(a+bx)}}{i+c-id}\right)}{1}$$

input

```
Integrate[x^2*ArcTan[c + d*Tan[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTan[c + d*Tan[a + b*x]] + (4*I)*b^3*x^3*Log[1 + (c + I*(-1 +
d))/((c - I*(1 + d))*E^((2*I)*(a + b*x)))] - (4*I)*b^3*x^3*Log[1 + (c + I
*(1 + d))/((I + c - I*d)*E^((2*I)*(a + b*x)))] + 6*b^2*x^2*PolyLog[2, (-c
- I*(1 + d))/((c - I*(-1 + d))*E^((2*I)*(a + b*x)))] - 6*b^2*x^2*PolyLog[2
, (I - c - I*d)/((c - I*(1 + d))*E^((2*I)*(a + b*x)))] - (6*I)*b*x*PolyLog
[3, (-c - I*(1 + d))/((I + c - I*d)*E^((2*I)*(a + b*x)))] + (6*I)*b*x*Poly
Log[3, (I - c - I*d)/((c - I*(1 + d))*E^((2*I)*(a + b*x)))] - 3*PolyLog[4,
(-c - I*(1 + d))/((I + c - I*d)*E^((2*I)*(a + b*x)))] + 3*PolyLog[4, (I -
c - I*d)/((c - I*(1 + d))*E^((2*I)*(a + b*x)))])/(24*b^3)
```

Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 519, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5698, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(d \tan(a + bx) + c) dx \\
 & \quad \downarrow \text{5698} \\
 & \frac{1}{3}b(-ic - d + 1) \int \frac{e^{2ia+2ibx} x^3}{-ic + (-ic - d + 1)e^{2ia+2ibx} + d + 1} dx - \frac{1}{3}b(ic + d + 1) \\
 & 1) \int \frac{e^{2ia+2ibx} x^3}{ic + (ic + d + 1)e^{2ia+2ibx} - d + 1} dx + \frac{1}{3}x^3 \arctan(d \tan(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & -\frac{1}{3}b(ic + d + 1) \left(\frac{3 \int x^2 \log \left(\frac{e^{2ia+2ibx}(ic+d+1)}{ic-d+1} + 1 \right) dx}{2b(c - i(d + 1))} - \frac{x^3 \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c - i(d + 1))} \right) + \\
 & \frac{1}{3}b(-ic - d + 1) \left(\frac{x^3 \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c + i(1 - d))} - \frac{3 \int x^2 \log \left(\frac{e^{2ia+2ibx}(c+i(1-d))}{c+i(d+1)} + 1 \right) dx}{2b(c + i(1 - d))} \right) + \\
 & \quad \frac{1}{3}x^3 \arctan(d \tan(a + bx) + c) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$\begin{aligned}
 & 1) \left(\frac{3 \left(\frac{ix^2 \operatorname{PolyLog} \left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b} - \frac{i \int x \operatorname{PolyLog} \left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right) dx}{b} \right)}{2b(c-i(d+1))} - \frac{x^3 \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c-i(d+1))} \right) + \\
 & 1) \left(\frac{x^3 \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c+i(1-d))} - \frac{3 \left(\frac{ix^2 \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b} - \frac{i \int x \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right) dx}{b} \right)}{2b(c+i(1-d))} \right) \\
 & \frac{1}{3} x^3 \arctan(d \tan(a + bx) + c)
 \end{aligned}$$

↓ 7163

$$\begin{aligned}
 & 1) \left(\frac{3 \left(\frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} - \frac{-\frac{1}{3}b(ic+d+1) + i \left(\frac{i \int \operatorname{PolyLog}\left(3, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right) dx}{2b} - \frac{ix \operatorname{PolyLog}\left(3, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} \right)}{b}}{2b(c-i(d+1))} - \frac{x^3 \log\left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b(c-i(d+1))} \right)}{2b(c-i(d+1))} \\
 & 1) \left(\frac{x^3 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c+i(1-d))} - \frac{3 \left(\frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} - \frac{\frac{1}{3}b(-ic-d+1) + i \left(\frac{i \int \operatorname{PolyLog}\left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right) dx}{2b} - \frac{ix \operatorname{PolyLog}\left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} \right)}{b}}{2b(c+i(1-d))} - \frac{x^3 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c+i(1-d))} \right)}{2b(c+i(1-d))} \\
 & \frac{1}{3}x^3 \arctan(d \tan(a + bx) + c)
 \end{aligned}$$

↓ 2720

$$1) \left(\frac{3 \left(\frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} - \frac{-\frac{1}{3}b(ic+d+1) \int e^{-2ia-2ibx} \operatorname{PolyLog}\left(3, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right) de^{2ia+2ibx}}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b}}{b} \right)}{2b(c-i(d+1))}$$

$$1) \left(\frac{x^3 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c+i(1-d))} - \frac{\frac{1}{3}b(-ic-d+1) \left(\frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} - \frac{i \left(\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right) de^{2ia+2ibx}}{4b^2} \right)}{b} \right)}{2b(c+i(1-d))}$$

$$\frac{1}{3}x^3 \arctan(d \tan(a+bx) + c)$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{3}x^3 \arctan(d \tan(a + bx) + c) - \frac{1}{3}b(ic + d + \\
 & 1) \left(\frac{3 \left(\frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} - i \left(\frac{\operatorname{PolyLog}\left(4, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} \right)}{b} \right)}{2b(c - i(d + 1))} - \frac{x^3 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c + i(1 - d))} \right) \\
 & 1) \left(\frac{x^3 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c + i(1 - d))} - \frac{\frac{1}{3}b(-ic - d + \left(\frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} - i \left(\frac{\operatorname{PolyLog}\left(4, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} \right)}{b} \right)}{2b(c + i(1 - d))} \right)
 \end{aligned}$$

input `Int[x^2*ArcTan[c + d*Tan[a + b*x]],x]`

output `(x^3*ArcTan[c + d*Tan[a + b*x]])/3 - (b*(1 + I*c + d)*(-1/2*(x^3*Log[1 + (1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d)))/(b*(c - I*(1 + d))) + (3*(((I/2)*x^2*PolyLog[2, -(((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d)))]/b - (I*(((1/2)*x*PolyLog[3, -(((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d)))]/b + PolyLog[4, -(((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d))]/(4*b^2)))/b)/(2*b*(c - I*(1 + d)))))/3 + (b*(1 - I*c - d)*((x^3*Log[1 + ((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/(2*b*(c + I*(1 - d))) - (3*(((I/2)*x^2*PolyLog[2, -(((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/b - (I*(((1/2)*x*PolyLog[3, -(((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/b + PolyLog[4, -(((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/(4*b^2)))/b)/(2*b*(c + I*(1 - d)))))/3`

Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5698

```
Int[ArcTan[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tan[a + b*x]]/(f*(m +
1))), x] + (Simp[b*((1 - I*c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(
2*I*a + 2*I*b*x))/(1 - I*c + d + (1 - I*c - d)*E^(2*I*a + 2*I*b*x)), x], x]
- Simp[b*((1 + I*c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*I*a + 2
*I*b*x))/(1 + I*c - d + (1 + I*c + d)*E^(2*I*a + 2*I*b*x)), x], x]) /; Free
Q[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c + I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 49.34 (sec) , antiderivative size = 8039, normalized size of antiderivative = 19.95

method	result	size
risch	Expression too large to display	8039

input

```
int(x^2*arctan(c+d*tan(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1965 vs. $2(290) = 580$.

Time = 0.18 (sec) , antiderivative size = 1965, normalized size of antiderivative = 4.88

$$\int x^2 \arctan(c + d \tan(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(x^2*arctan(c+d*tan(b*x+a)),x, algorithm="fricas")
```

output

```

1/48*(16*b^3*x^3*arctan(d*tan(b*x + a) + c) + 6*b^2*x^2*dilog(2*((I*c*d -
d^2 + d)*tan(b*x + a)^2 - c^2 - I*c*d + (I*c^2 - 2*c*d - I*d^2 + I)*tan(b*
x + a) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*d +
1) + 1) - 6*b^2*x^2*dilog(2*((I*c*d - d^2 - d)*tan(b*x + a)^2 - c^2 - I*c*
d + (I*c^2 - 2*c*d - I*d^2 + I)*tan(b*x + a) - d - 1)/((c^2 + d^2 + 2*d +
1)*tan(b*x + a)^2 + c^2 + d^2 + 2*d + 1) + 1) + 6*b^2*x^2*dilog(2*((-I*c*d
- d^2 + d)*tan(b*x + a)^2 - c^2 + I*c*d + (-I*c^2 - 2*c*d + I*d^2 - I)*ta
n(b*x + a) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*
d + 1) + 1) - 6*b^2*x^2*dilog(2*((-I*c*d - d^2 - d)*tan(b*x + a)^2 - c^2 +
I*c*d + (-I*c^2 - 2*c*d + I*d^2 - I)*tan(b*x + a) - d - 1)/((c^2 + d^2 +
2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*d + 1) + 1) + 4*I*a^3*log(((I*c*d
+ d^2 + d)*tan(b*x + a)^2 - c^2 + I*c*d + (I*c^2 + I*d^2 + 2*I*d + I)*tan(
b*x + a) - d - 1)/(tan(b*x + a)^2 + 1)) - 4*I*a^3*log(((I*c*d + d^2 - d)*t
an(b*x + a)^2 - c^2 + I*c*d + (I*c^2 + I*d^2 - 2*I*d + I)*tan(b*x + a) +
d - 1)/(tan(b*x + a)^2 + 1)) + 4*I*a^3*log(((I*c*d - d^2 + d)*tan(b*x + a)^
2 + c^2 + I*c*d + (I*c^2 + I*d^2 - 2*I*d + I)*tan(b*x + a) - d + 1)/(tan(b
*x + a)^2 + 1)) - 4*I*a^3*log(((I*c*d - d^2 - d)*tan(b*x + a)^2 + c^2 + I*
c*d + (I*c^2 + I*d^2 + 2*I*d + I)*tan(b*x + a) + d + 1)/(tan(b*x + a)^2 +
1)) + 6*I*b*x*polylog(3, ((c^2 + 2*I*c*d - d^2 + 1)*tan(b*x + a)^2 - c^2 -
2*I*c*d + d^2 - 2*(-I*c^2 + 2*c*d + I*d^2 - I)*tan(b*x + a) - 1)/((c^2...

```

Sympy [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \tan(a + bx)) dx = \text{Timed out}$$

input

```
integrate(x**2*atan(c+d*tan(b*x+a)),x)
```

output

Timed out

Maxima [F]

$$\int x^2 \arctan(c + d \tan(a + bx)) dx = \int x^2 \arctan(d \tan(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*tan(b*x+a)),x, algorithm="maxima")`

output

```
1/6*x^3*arctan2(c*cos(2*b*x + 2*a) + (d + 1)*sin(2*b*x + 2*a) + c, (d + 1)
*cos(2*b*x + 2*a) - c*sin(2*b*x + 2*a) - d + 1) + 1/6*x^3*arctan2(c*cos(2*
b*x + 2*a) + (d - 1)*sin(2*b*x + 2*a) + c, -(d - 1)*cos(2*b*x + 2*a) + c*s
in(2*b*x + 2*a) + d + 1) + 4*b*d*integrate(-1/3*(2*(c^2 + d^2 + 1)*x^3*cos
(2*b*x + 2*a)^2 + 2*c*d*x^3*sin(2*b*x + 2*a) + 2*(c^2 + d^2 + 1)*x^3*sin(2
*b*x + 2*a)^2 + (c^2 - d^2 + 1)*x^3*cos(2*b*x + 2*a) - (2*c*d*x^3*sin(2*b*
x + 2*a) - (c^2 - d^2 + 1)*x^3*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d
*x^3*cos(2*b*x + 2*a) + (c^2 - d^2 + 1)*x^3*sin(2*b*x + 2*a))*sin(4*b*x +
4*a))/(c^4 + d^4 + 2*(c^2 - 1)*d^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2
+ 1)*cos(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*cos(
2*b*x + 2*a)^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2 + 1)*sin(4*b*x + 4*a
)^2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*sin(2*b*x + 2*a)^2 + 2*c
^2 + 2*(c^4 + d^4 - 2*(3*c^2 + 1)*d^2 + 2*c^2 + 2*(c^4 - d^4 + 2*c^2 + 1)*
cos(2*b*x + 2*a) - 4*(c*d^3 + (c^3 + c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x
+ 4*a) + 4*(c^4 - d^4 + 2*c^2 + 1)*cos(2*b*x + 2*a) - 4*(2*c*d^3 - 2*(c^3
+ c)*d - 2*(c*d^3 + (c^3 + c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 + 2*c^2 +
1)*sin(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 + c)*d)*sin(2*b*x
+ 2*a) + 1), x)
```

Giac [F]

$$\int x^2 \arctan(c + d \tan(a + bx)) dx = \int x^2 \arctan(d \tan(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan(d*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \tan(a + bx)) dx = \int x^2 \operatorname{atan}(c + d \tan(a + bx)) dx$$

input `int(x^2*atan(c + d*tan(a + b*x)),x)`output `int(x^2*atan(c + d*tan(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \arctan(c + d \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a)d + c) x^2 dx$$

input `int(x^2*atan(c+d*tan(b*x+a)),x)`output `int(atan(tan(a + b*x)*d + c)*x**2,x)`

3.49 $\int x \arctan(c + d \tan(a + bx)) dx$

Optimal result	397
Mathematica [A] (verified)	398
Rubi [A] (verified)	398
Maple [C] (warning: unable to verify)	402
Fricas [B] (verification not implemented)	402
Sympy [F]	403
Maxima [F]	404
Giac [F]	404
Mupad [F(-1)]	405
Reduce [F]	405

Optimal result

Integrand size = 13, antiderivative size = 305

$$\begin{aligned}
 \int x \arctan(c + d \tan(a + bx)) dx = & \frac{1}{2}x^2 \arctan(c + d \tan(a + bx)) \\
 & + \frac{1}{4}ix^2 \log \left(1 + \frac{(1 + ic + d)e^{2ia+2ibx}}{1 + ic - d} \right) \\
 & - \frac{1}{4}ix^2 \log \left(1 + \frac{(c + i(1 - d))e^{2ia+2ibx}}{c + i(1 + d)} \right) \\
 & + \frac{x \operatorname{PolyLog} \left(2, -\frac{(1+ic+d)e^{2ia+2ibx}}{1+ic-d} \right)}{4b} \\
 & - \frac{x \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(1+d)} \right)}{4b} \\
 & + \frac{i \operatorname{PolyLog} \left(3, -\frac{(1+ic+d)e^{2ia+2ibx}}{1+ic-d} \right)}{8b^2} \\
 & - \frac{i \operatorname{PolyLog} \left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(1+d)} \right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & 1/2*x^2*\arctan(c+d*\tan(b*x+a))+1/4*I*x^2*\ln(1+(1+I*c+d)*\exp(2*I*a+2*I*b*x) \\ & /((1+I*c-d))-1/4*I*x^2*\ln(1+(c+I*(1-d))*\exp(2*I*a+2*I*b*x)/(c+I*(1+d)))+1/4 \\ & *x*\text{polylog}(2,-(1+I*c+d)*\exp(2*I*a+2*I*b*x)/(1+I*c-d))/b-1/4*x*\text{polylog}(2,-(\\ & c+I*(1-d))*\exp(2*I*a+2*I*b*x)/(c+I*(1+d)))/b+1/8*I*\text{polylog}(3,-(1+I*c+d)*\exp \\ & (2*I*a+2*I*b*x)/(1+I*c-d))/b^2-1/8*I*\text{polylog}(3,-(c+I*(1-d))*\exp(2*I*a+2*I \\ & *b*x)/(c+I*(1+d)))/b^2 \end{aligned}$$
Mathematica [A] (verified)

Time = 2.58 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.92

$$\int x \arctan(c + d \tan(a + bx)) dx$$

$$= \frac{4b^2x^2 \arctan(c + d \tan(a + bx)) + 2ib^2x^2 \log\left(1 + \frac{(c+i(-1+d))e^{-2i(a+bx)}}{c-i(1+d)}\right) - 2ib^2x^2 \log\left(1 + \frac{(c+i(1+d))e^{-2i(a+bx)}}{i+c-id}\right)}{1}$$

input

Integrate[x*ArcTan[c + d*Tan[a + b*x]],x]

output

$$\begin{aligned} & (4*b^2*x^2*ArcTan[c + d*Tan[a + b*x]] + (2*I)*b^2*x^2*Log[1 + (c + I*(-1 + \\ & d))/((c - I*(1 + d))*E^((2*I)*(a + b*x)))] - (2*I)*b^2*x^2*Log[1 + (c + I \\ & *(1 + d))/((I + c - I*d)*E^((2*I)*(a + b*x)))] + 2*b*x*PolyLog[2, (-c - I* \\ & (1 + d))/((c - I*(-1 + d))*E^((2*I)*(a + b*x)))] - 2*b*x*PolyLog[2, (I - c \\ & - I*d)/((c - I*(1 + d))*E^((2*I)*(a + b*x)))] - I*PolyLog[3, (-c - I*(1 + \\ & d))/((I + c - I*d)*E^((2*I)*(a + b*x)))] + I*PolyLog[3, (I - c - I*d)/((c \\ & - I*(1 + d))*E^((2*I)*(a + b*x)))])/(8*b^2) \end{aligned}$$
Rubi [A] (verified)Time = 1.15 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.30, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {5698, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x \arctan(d \tan(a + bx) + c) dx \\
& \quad \downarrow \text{5698} \\
& \frac{1}{2}b(-ic - d + 1) \int \frac{e^{2ia+2ibx} x^2}{-ic + (-ic - d + 1)e^{2ia+2ibx} + d + 1} dx - \frac{1}{2}b(ic + d + \\
& 1) \int \frac{e^{2ia+2ibx} x^2}{ic + (ic + d + 1)e^{2ia+2ibx} - d + 1} dx + \frac{1}{2}x^2 \arctan(d \tan(a + bx) + c) \\
& \quad \downarrow \text{2620} \\
& -\frac{1}{2}b(ic + d + 1) \left(\frac{\int x \log \left(\frac{e^{2ia+2ibx}(ic+d+1)}{ic-d+1} + 1 \right) dx}{b(c - i(d + 1))} - \frac{x^2 \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c - i(d + 1))} \right) + \\
& \frac{1}{2}b(-ic - d + 1) \left(\frac{x^2 \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c + i(1 - d))} - \frac{\int x \log \left(\frac{e^{2ia+2ibx}(c+i(1-d))}{c+i(d+1)} + 1 \right) dx}{b(c + i(1 - d))} \right) + \\
& \quad \frac{1}{2}x^2 \arctan(d \tan(a + bx) + c) \\
& \quad \downarrow \text{3011} \\
& -\frac{1}{2}b(ic + d + \\
& 1) \left(\frac{\frac{ix \operatorname{PolyLog} \left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right) dx}{2b}}{b(c - i(d + 1))} - \frac{x^2 \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c - i(d + 1))} \right) + \\
& \quad \frac{1}{2}b(-ic - d + \\
& 1) \left(\frac{x^2 \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c + i(1 - d))} - \frac{\frac{ix \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right) dx}{2b}}{b(c + i(1 - d))} \right) + \\
& \quad \frac{1}{2}x^2 \arctan(d \tan(a + bx) + c) \\
& \quad \downarrow \text{2720}
\end{aligned}$$

$$\begin{aligned}
& \frac{-\frac{1}{2}b(ic+d+)}{2} \\
1) & \left(\frac{\frac{ix \operatorname{PolyLog}\left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right) de^{2ia+2ibx}}{4b^2}}{b(c-i(d+1))} - \frac{x^2 \log\left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b(c-i(d+1))} \right. \\
& \left. \frac{\frac{1}{2}b(-ic-d+)}{2} \right. \\
1) & \left(\frac{x^2 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c+i(1-d))} - \frac{ix \operatorname{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right) de^{2ia+2ibx}}{4b^2}}{b(c+i(1-d))} \right) \\
& \frac{1}{2}x^2 \arctan(d \tan(a+bx) + c)
\end{aligned}$$

↓ 7143

$$\begin{aligned}
& \frac{1}{2}x^2 \arctan(d \tan(a+bx) + c) - \frac{1}{2}b(ic+d+)}{2} \\
1) & \left(\frac{\frac{ix \operatorname{PolyLog}\left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{4b^2}}{b(c-i(d+1))} - \frac{x^2 \log\left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1}\right)}{2b(c-i(d+1))} \right) + \\
& \frac{1}{2}b(-ic-d+)}{2} \\
1) & \left(\frac{x^2 \log\left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b(c+i(1-d))} - \frac{ix \operatorname{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)}\right)}{4b^2}}{b(c+i(1-d))} \right)
\end{aligned}$$

input

```
Int[x*ArcTan[c + d*Tan[a + b*x]],x]
```

output

```
(x^2*ArcTan[c + d*Tan[a + b*x]])/2 - (b*(1 + I*c + d)*(-1/2*(x^2*Log[1 + (1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d)))/(b*(c - I*(1 + d))) + (((I/2)*x*PolyLog[2, -(((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d)))]/b - PolyLog[3, -(((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x)]/(1 + I*c - d))]/(4*b^2)))/(b*(c - I*(1 + d))))/2 + (b*(1 - I*c - d)*((x^2*Log[1 + ((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/(2*b*(c + I*(1 - d))) - (((I/2)*x*PolyLog[2, -(((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/b - PolyLog[3, -(((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 + d)))]/(4*b^2)))/(b*(c + I*(1 - d)))))/2
```

Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5698

```
Int[ArcTan[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tan[a + b*x]]/(f*(m +
1))), x] + (Simp[b*((1 - I*c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(
2*I*a + 2*I*b*x))/(1 - I*c + d + (1 - I*c - d)*E^(2*I*a + 2*I*b*x)), x], x]
- Simp[b*((1 + I*c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*I*a + 2
*I*b*x))/(1 + I*c - d + (1 + I*c + d)*E^(2*I*a + 2*I*b*x)), x], x]) /; Free
Q[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c + I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.92 (sec) , antiderivative size = 7647, normalized size of antiderivative = 25.07

method	result	size
risch	Expression too large to display	7647

input `int(x*arctan(c+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1545 vs. $2(217) = 434$.

Time = 0.16 (sec) , antiderivative size = 1545, normalized size of antiderivative = 5.07

$$\int x \arctan(c + d \tan(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctan(c+d*tan(b*x+a)),x, algorithm="fricas")`

output

```

1/16*(8*b^2*x^2*arctan(d*tan(b*x + a) + c) + 2*b*x*dilog(2*((I*c*d - d^2 +
d)*tan(b*x + a)^2 - c^2 - I*c*d + (I*c^2 - 2*c*d - I*d^2 + I)*tan(b*x + a
) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*d + 1) +
1) - 2*b*x*dilog(2*((I*c*d - d^2 - d)*tan(b*x + a)^2 - c^2 - I*c*d + (I*c^
2 - 2*c*d - I*d^2 + I)*tan(b*x + a) - d - 1)/((c^2 + d^2 + 2*d + 1)*tan(b*
x + a)^2 + c^2 + d^2 + 2*d + 1) + 1) + 2*b*x*dilog(2*((-I*c*d - d^2 + d)*t
an(b*x + a)^2 - c^2 + I*c*d + (-I*c^2 - 2*c*d + I*d^2 - I)*tan(b*x + a) +
d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*d + 1) + 1) -
2*b*x*dilog(2*((-I*c*d - d^2 - d)*tan(b*x + a)^2 - c^2 + I*c*d + (-I*c^2
- 2*c*d + I*d^2 - I)*tan(b*x + a) - d - 1)/((c^2 + d^2 + 2*d + 1)*tan(b*x
+ a)^2 + c^2 + d^2 + 2*d + 1) + 1) - 2*I*a^2*log(((I*c*d + d^2 + d)*tan(b*
x + a)^2 - c^2 + I*c*d + (I*c^2 + I*d^2 + 2*I*d + I)*tan(b*x + a) - d - 1)
/(tan(b*x + a)^2 + 1)) + 2*I*a^2*log(((I*c*d + d^2 - d)*tan(b*x + a)^2 - c
^2 + I*c*d + (I*c^2 + I*d^2 - 2*I*d + I)*tan(b*x + a) + d - 1)/(tan(b*x +
a)^2 + 1)) - 2*I*a^2*log(((I*c*d - d^2 + d)*tan(b*x + a)^2 + c^2 + I*c*d +
(I*c^2 + I*d^2 - 2*I*d + I)*tan(b*x + a) - d + 1)/(tan(b*x + a)^2 + 1)) +
2*I*a^2*log(((I*c*d - d^2 - d)*tan(b*x + a)^2 + c^2 + I*c*d + (I*c^2 + I*
d^2 + 2*I*d + I)*tan(b*x + a) + d + 1)/(tan(b*x + a)^2 + 1)) - 2*(I*b^2*x^
2 - I*a^2)*log(-2*((I*c*d - d^2 + d)*tan(b*x + a)^2 - c^2 - I*c*d + (I*c^2
- 2*c*d - I*d^2 + I)*tan(b*x + a) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(...

```

Sympy [F]

$$\int x \arctan(c + d \tan(a + bx)) dx = \int x \operatorname{atan}(c + d \tan(a + bx)) dx$$

input

```
integrate(x*atan(c+d*tan(b*x+a)),x)
```

output

```
Integral(x*atan(c + d*tan(a + b*x)), x)
```

Maxima [F]

$$\int x \arctan(c + d \tan(a + bx)) dx = \int x \arctan(d \tan(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*tan(b*x+a)),x, algorithm="maxima")`

output

```
1/4*x^2*arctan2(c*cos(2*b*x + 2*a) + (d + 1)*sin(2*b*x + 2*a) + c, (d + 1)
*cos(2*b*x + 2*a) - c*sin(2*b*x + 2*a) - d + 1) + 1/4*x^2*arctan2(c*cos(2*
b*x + 2*a) + (d - 1)*sin(2*b*x + 2*a) + c, -(d - 1)*cos(2*b*x + 2*a) + c*s
in(2*b*x + 2*a) + d + 1) + 2*b*d*integrate(-(2*(c^2 + d^2 + 1)*x^2*cos(2*b
*x + 2*a)^2 + 2*c*d*x^2*sin(2*b*x + 2*a) + 2*(c^2 + d^2 + 1)*x^2*sin(2*b*x
+ 2*a)^2 + (c^2 - d^2 + 1)*x^2*cos(2*b*x + 2*a) - (2*c*d*x^2*sin(2*b*x +
2*a) - (c^2 - d^2 + 1)*x^2*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d*x^2
*cos(2*b*x + 2*a) + (c^2 - d^2 + 1)*x^2*sin(2*b*x + 2*a))*sin(4*b*x + 4*a)
)/(c^4 + d^4 + 2*(c^2 - 1)*d^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2 + 1)
*cos(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*cos(2*b*
x + 2*a)^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2 + 1)*sin(4*b*x + 4*a)^2
+ 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*sin(2*b*x + 2*a)^2 + 2*c^2 +
2*(c^4 + d^4 - 2*(3*c^2 + 1)*d^2 + 2*c^2 + 2*(c^4 - d^4 + 2*c^2 + 1)*cos(
2*b*x + 2*a) - 4*(c*d^3 + (c^3 + c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x + 4
*a) + 4*(c^4 - d^4 + 2*c^2 + 1)*cos(2*b*x + 2*a) - 4*(2*c*d^3 - 2*(c^3 + c
)*d - 2*(c*d^3 + (c^3 + c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 + 2*c^2 + 1)*s
in(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 + c)*d)*sin(2*b*x + 2*
a) + 1), x)
```

Giac [F]

$$\int x \arctan(c + d \tan(a + bx)) dx = \int x \arctan(d \tan(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan(d*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + d \tan(a + bx)) dx = \int x \operatorname{atan}(c + d \tan(a + bx)) dx$$

input `int(x*atan(c + d*tan(a + b*x)),x)`output `int(x*atan(c + d*tan(a + b*x)), x)`**Reduce [F]**

$$\int x \arctan(c + d \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a)d + c) x dx$$

input `int(x*atan(c+d*tan(b*x+a)),x)`output `int(atan(tan(a + b*x)*d + c)*x,x)`

3.50 $\int \arctan(c + d \tan(a + bx)) dx$

Optimal result	406
Mathematica [B] (warning: unable to verify)	407
Rubi [A] (verified)	407
Maple [B] (verified)	410
Fricas [B] (verification not implemented)	411
Sympy [F]	412
Maxima [B] (verification not implemented)	412
Giac [F]	413
Mupad [F(-1)]	413
Reduce [F]	413

Optimal result

Integrand size = 11, antiderivative size = 198

$$\int \arctan(c + d \tan(a + bx)) dx = x \arctan(c + d \tan(a + bx)) + \frac{1}{2}ix \log\left(1 + \frac{(1 + ic + d)e^{2ia+2ibx}}{1 + ic - d}\right) - \frac{1}{2}ix \log\left(1 + \frac{(c + i(1 - d))e^{2ia+2ibx}}{c + i(1 + d)}\right) + \frac{\text{PolyLog}\left(2, -\frac{(1+ic+d)e^{2ia+2ibx}}{1+ic-d}\right)}{4b} - \frac{\text{PolyLog}\left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(1+d)}\right)}{4b}$$

output

```
x*arctan(c+d*tan(b*x+a))+1/2*I*x*ln(1+(1+I*c+d)*exp(2*I*a+2*I*b*x)/(1+I*c-d))-1/2*I*x*ln(1+(c+I*(1-d))*exp(2*I*a+2*I*b*x)/(c+I*(1+d)))+1/4*polylog(2,-(1+I*c+d)*exp(2*I*a+2*I*b*x)/(1+I*c-d))/b-1/4*polylog(2,-(c+I*(1-d))*exp(2*I*a+2*I*b*x)/(c+I*(1+d)))/b
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 549 vs. $2(198) = 396$.

Time = 5.10 (sec) , antiderivative size = 549, normalized size of antiderivative = 2.77

$$\int \arctan(c + d \tan(a + bx)) dx = x \arctan(c + d \tan(a + bx))$$

$$x \left(4a\sqrt{-d^2} \arctan(c + d \tan(a + bx)) - id \log(1 - i \tan(a + bx)) \log\left(\frac{-cd + \sqrt{-d^2} - d^2 \tan(a + bx)}{-cd + id^2 + \sqrt{-d^2}}\right) + id \log\left(\frac{-cd + \sqrt{-d^2} + d^2 \tan(a + bx)}{-cd + id^2 + \sqrt{-d^2}}\right) \right)$$

input `Integrate[ArcTan[c + d*Tan[a + b*x]],x]`

output

```
x*ArcTan[c + d*Tan[a + b*x]] - (x*(4*a*Sqrt[-d^2]*ArcTan[c + d*Tan[a + b*x]] - I*d*Log[1 - I*Tan[a + b*x]]*Log[(-(c*d) + Sqrt[-d^2] - d^2*Tan[a + b*x])/(-(c*d) + I*d^2 + Sqrt[-d^2])] + I*d*Log[1 + I*Tan[a + b*x]]*Log[(c*d - Sqrt[-d^2] + d^2*Tan[a + b*x])/(c*d + I*d^2 - Sqrt[-d^2])] + I*d*Log[1 - I*Tan[a + b*x]]*Log[(c*d + Sqrt[-d^2] + d^2*Tan[a + b*x])/(c*d - I*d^2 + Sqrt[-d^2])] - I*d*Log[1 + I*Tan[a + b*x]]*Log[(c*d + Sqrt[-d^2] + d^2*Tan[a + b*x])/(c*d + I*d^2 + Sqrt[-d^2])] - I*d*PolyLog[2, (d^2*(1 - I*Tan[a + b*x]))/(I*c*d + d^2 - I*Sqrt[-d^2])] + I*d*PolyLog[2, (d^2*(1 - I*Tan[a + b*x]))/(I*c*d + d^2 + I*Sqrt[-d^2])] + I*d*PolyLog[2, (d^2*(1 + I*Tan[a + b*x]))/((-I)*c*d + d^2 + I*Sqrt[-d^2])] - I*d*PolyLog[2, (d^2*(1 + I*Tan[a + b*x]))/(d^2 - I*(c*d + Sqrt[-d^2]))])/(2*Sqrt[-d^2]*(2*a - I*Log[1 - I*Tan[a + b*x]] + I*Log[1 + I*Tan[a + b*x]]))
```

Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.40, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {5690, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(d \tan(a + bx) + c) dx$$

↓ 5690

$$b(-ic - d + 1) \int \frac{e^{2ia+2ibx} x}{-ic + (-ic - d + 1)e^{2ia+2ibx} + d + 1} dx - b(ic + d + 1) \int \frac{e^{2ia+2ibx}}{ic + (ic + d + 1)e^{2ia+2ibx} - d + 1} dx + x \arctan(d \tan(a + bx) + c)$$

↓ 2620

$$-b(ic + d + 1) \left(\frac{\int \log \left(\frac{e^{2ia+2ibx}(ic+d+1)}{ic-d+1} + 1 \right) dx}{2b(c - i(d+1))} - \frac{x \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c - i(d+1))} \right) + b(-ic - d + 1) \left(\frac{x \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c + i(1-d))} - \frac{\int \log \left(\frac{e^{2ia+2ibx}(c+i(1-d))}{c+i(d+1)} + 1 \right) dx}{2b(c + i(1-d))} \right) + x \arctan(d \tan(a + bx) + c)$$

↓ 2715

$$1) \left(-\frac{i \int e^{-2ia-2ibx} \log \left(\frac{e^{2ia+2ibx}(ic+d+1)}{ic-d+1} + 1 \right) de^{2ia+2ibx}}{4b^2(c - i(d+1))} - \frac{x \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c - i(d+1))} \right) + 1) \left(\frac{i \int e^{-2ia-2ibx} \log \left(\frac{e^{2ia+2ibx}(c+i(1-d))}{c+i(d+1)} + 1 \right) de^{2ia+2ibx}}{4b^2(c + i(1-d))} + \frac{x \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c + i(1-d))} \right) + x \arctan(d \tan(a + bx) + c)$$

↓ 2838

$$1) \left(\frac{x \arctan(d \tan(a + bx) + c) - b(ic + d + 1) \operatorname{PolyLog} \left(2, -\frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{4b^2(c - i(d+1))} - \frac{x \log \left(1 + \frac{(ic+d+1)e^{2ia+2ibx}}{ic-d+1} \right)}{2b(c - i(d+1))} \right) + b(-ic - d + 1) \left(\frac{x \log \left(1 + \frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{2b(c + i(1-d))} - \frac{i \operatorname{PolyLog} \left(2, -\frac{(c+i(1-d))e^{2ia+2ibx}}{c+i(d+1)} \right)}{4b^2(c + i(1-d))} \right)$$

input

Int[ArcTan[c + d*Tan[a + b*x]], x]

output

```
x*ArcTan[c + d*Tan[a + b*x]] - b*(1 + I*c + d)*(-1/2*(x*Log[1 + ((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c - d)]/(b*(c - I*(1 + d))) + ((I/4)*PolyLog[2, -(((1 + I*c + d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c - d)])/(b^2*(c - I*(1 + d)))) + b*(1 - I*c - d)*((x*Log[1 + ((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 + d))])/(2*b*(c + I*(1 - d))) - ((I/4)*PolyLog[2, -(((c + I*(1 - d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 + d))])/(b^2*(c + I*(1 - d))))
```

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 5690

```
Int[ArcTan[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]], x_Symbol] := Simp[x*ArcTan[c + d*Tan[a + b*x]], x] + (Simp[b*(1 - I*c - d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 - I*c + d + (1 - I*c - d)*E^(2*I*a + 2*I*b*x))], x], x] - Simp[b*(1 + I*c + d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 + I*c - d + (1 + I*c + d)*E^(2*I*a + 2*I*b*x))], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[(c + I*d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1000 vs. $2(168) = 336$.

Time = 2.90 (sec) , antiderivative size = 1001, normalized size of antiderivative = 5.06

method	result	size
derivativeldivides	Expression too large to display	1001
default	Expression too large to display	1001
risch	Expression too large to display	4973

input `int(arctan(c+d*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\frac{1}{b/d} \left(d \arctan(\tan(b*x+a)) \arctan(c+d*\tan(b*x+a)) - d^2 \left(\frac{1}{2} \frac{I}{d} \arctan\left(-\frac{c+d*\tan(b*x+a)}{d+c/d}\right) \ln\left(1 - (-I-I*d+c) \left(1 + I \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2\right)\right) \right)^2 / \left(\left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + 1 \right) / (-I*d+I-c) - \frac{1}{2} \frac{I}{d} \arctan\left(-\frac{c+d*\tan(b*x+a)}{d+c/d}\right) \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + \frac{1}{4} \frac{I}{d} \operatorname{polylog}\left(2, (-I-I*d+c) \left(1 + I \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2\right)\right) / \left(\left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + 1 \right) / (-I*d+I-c) + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right) \arctan\left(-\frac{c+d*\tan(b*x+a)}{d+c/d}\right) + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 / \left(\left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + 1 \right) / (-I*d-I-c) \right) \arctan\left(-\frac{c+d*\tan(b*x+a)}{d+c/d}\right) - \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right) \arctan\left(-\frac{c+d*\tan(b*x+a)}{d+c/d}\right) + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + \frac{1}{4} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right) \operatorname{polylog}\left(2, (-I-I*d+c) \left(1 + I \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2\right)\right) / \left(\left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + 1 \right) / (-I*d-I-c) + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + \frac{1}{2} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right) \arctan\left(-\frac{c+d*\tan(b*x+a)}{d+c/d}\right) + \frac{1}{4} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right) \operatorname{polylog}\left(2, (-I-I*d+c) \left(1 + I \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2\right)\right) / \left(\left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + 1 \right) / (-I*d-I-c) + \frac{1}{4} \frac{I}{d} \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right) \operatorname{polylog}\left(2, (-I-I*d+c) \left(1 + I \left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2\right)\right) / \left(\left(\frac{c+d*\tan(b*x+a)}{d-c/d}\right)^2 + 1 \right) / (-I*d-I-c) \right) \right) \right)$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1101 vs. $2(141) = 282$.

Time = 0.17 (sec) , antiderivative size = 1101, normalized size of antiderivative = 5.56

$$\int \arctan(c + d \tan(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctan(c+d*tan(b*x+a)),x, algorithm="fricas")`

output

```
1/8*(8*b*x*arctan(d*tan(b*x + a) + c) - 2*(I*b*x + I*a)*log(-2*((I*c*d - d
^2 + d)*tan(b*x + a)^2 - c^2 - I*c*d + (I*c^2 - 2*c*d - I*d^2 + I)*tan(b*x
+ a) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 - 2*d + 1
)) - 2*(-I*b*x - I*a)*log(-2*((I*c*d - d^2 - d)*tan(b*x + a)^2 - c^2 - I*c
*d + (I*c^2 - 2*c*d - I*d^2 + I)*tan(b*x + a) - d - 1)/((c^2 + d^2 + 2*d +
1)*tan(b*x + a)^2 + c^2 + d^2 + 2*d + 1)) - 2*(-I*b*x - I*a)*log(-2*((-I*
c*d - d^2 + d)*tan(b*x + a)^2 - c^2 + I*c*d + (-I*c^2 - 2*c*d + I*d^2 - I)
*tan(b*x + a) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 -
2*d + 1)) - 2*(I*b*x + I*a)*log(-2*((-I*c*d - d^2 - d)*tan(b*x + a)^2 - c
^2 + I*c*d + (-I*c^2 - 2*c*d + I*d^2 - I)*tan(b*x + a) - d - 1)/((c^2 + d
^2 + 2*d + 1)*tan(b*x + a)^2 + c^2 + d^2 + 2*d + 1)) + 2*I*a*log(((I*c*d +
d^2 + d)*tan(b*x + a)^2 - c^2 + I*c*d + (I*c^2 + I*d^2 + 2*I*d + I)*tan(b*
x + a) - d - 1)/(tan(b*x + a)^2 + 1)) - 2*I*a*log(((I*c*d + d^2 - d)*tan(b
*x + a)^2 - c^2 + I*c*d + (I*c^2 + I*d^2 - 2*I*d + I)*tan(b*x + a) + d - 1
)/(tan(b*x + a)^2 + 1)) + 2*I*a*log(((I*c*d - d^2 + d)*tan(b*x + a)^2 + c^
2 + I*c*d + (I*c^2 + I*d^2 - 2*I*d + I)*tan(b*x + a) - d + 1)/(tan(b*x + a
)^2 + 1)) - 2*I*a*log(((I*c*d - d^2 - d)*tan(b*x + a)^2 + c^2 + I*c*d + (I
*c^2 + I*d^2 + 2*I*d + I)*tan(b*x + a) + d + 1)/(tan(b*x + a)^2 + 1)) + di
log(2*((I*c*d - d^2 + d)*tan(b*x + a)^2 - c^2 - I*c*d + (I*c^2 - 2*c*d - I
*d^2 + I)*tan(b*x + a) + d - 1)/((c^2 + d^2 - 2*d + 1)*tan(b*x + a)^2 + ...
```

Sympy [F]

$$\int \arctan(c + d \tan(a + bx)) dx = \int \operatorname{atan}(c + d \tan(a + bx)) dx$$

input `integrate(atan(c+d*tan(b*x+a)),x)`

output `Integral(atan(c + d*tan(a + b*x)), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 433 vs. $2(141) = 282$.

Time = 0.21 (sec) , antiderivative size = 433, normalized size of antiderivative = 2.19

$$\int \arctan(c + d \tan(a + bx)) dx$$

$$= d \left(\frac{8(bx+a) \arctan\left(\frac{d^2 \tan(bx+a)+cd}{d}\right)}{d} - \frac{4(bx+a) \arctan\left(\frac{cd+(d^2+d)\tan(bx+a)}{c^2+d^2+2d+1}, \frac{cd \tan(bx+a)+c^2+d+1}{c^2+d^2+2d+1}\right) - 4(bx+a) \arctan\left(\frac{cd+(d^2-d)\tan(bx+a)}{c^2+d^2-2d+1}\right)}{d} \right)$$

input `integrate(arctan(c+d*tan(b*x+a)),x, algorithm="maxima")`

output `1/8*(d*(8*(b*x + a)*arctan((d^2*tan(b*x + a) + c*d)/d)/d - (4*(b*x + a)*arctan2((c*d + (d^2 + d)*tan(b*x + a))/(c^2 + d^2 + 2*d + 1), (c*d*tan(b*x + a) + c^2 + d + 1)/(c^2 + d^2 + 2*d + 1)) - 4*(b*x + a)*arctan2((c*d + (d^2 - d)*tan(b*x + a))/(c^2 + d^2 - 2*d + 1), (c*d*tan(b*x + a) + c^2 - d + 1)/(c^2 + d^2 - 2*d + 1)) + log(tan(b*x + a)^2 + 1)*log((d^2*tan(b*x + a)^2 + 2*c*d*tan(b*x + a) + c^2 + 1)/(c^2 + d^2 + 2*d + 1)) - log(tan(b*x + a)^2 + 1)*log((d^2*tan(b*x + a)^2 + 2*c*d*tan(b*x + a) + c^2 + 1)/(c^2 + d^2 - 2*d + 1)) + 2*dilog(-(I*d*tan(b*x + a) - d)/(I*c + d + 1)) - 2*dilog(-(I*d*tan(b*x + a) - d)/(I*c + d - 1)) + 2*dilog((I*d*tan(b*x + a) + d)/(-I*c + d + 1)) - 2*dilog((I*d*tan(b*x + a) + d)/(-I*c + d - 1)))/d + 8*(b*x + a)*arctan(d*tan(b*x + a) + c) - 8*(b*x + a)*arctan((d^2*tan(b*x + a) + c*d)/d))/b`

Giac [F]

$$\int \arctan(c + d \tan(a + bx)) dx = \int \arctan(d \tan(bx + a) + c) dx$$

input `integrate(arctan(c+d*tan(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(d*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + d \tan(a + bx)) dx = \int \operatorname{atan}(c + d \tan(a + bx)) dx$$

input `int(atan(c + d*tan(a + b*x)),x)`

output `int(atan(c + d*tan(a + b*x)), x)`

Reduce [F]

$$\int \arctan(c + d \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) d + c) dx$$

input `int(atan(c+d*tan(b*x+a)),x)`

output `int(atan(tan(a + b*x)*d + c),x)`

3.51 $\int \frac{\arctan(c+d \tan(a+bx))}{x} dx$

Optimal result	414
Mathematica [N/A]	414
Rubi [N/A]	415
Maple [N/A]	415
Fricas [N/A]	416
Sympy [N/A]	416
Maxima [F(-1)]	416
Giac [N/A]	417
Mupad [N/A]	417
Reduce [N/A]	417

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + d \tan(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+d*tan(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 1.86 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \int \frac{\arctan(c + d \tan(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + d*Tan[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + d*Tan[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(d \tan(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\arctan(d \tan(a + bx) + c)}{x} dx$$

input `Int[ArcTan[c + d*Tan[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + d \tan(bx + a))}{x} dx$$

input `int(arctan(c+d*tan(b*x+a))/x,x)`

output `int(arctan(c+d*tan(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \int \frac{\arctan(d \tan(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*tan(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctan(d*tan(b*x + a) + c)/x, x)`

Sympy [N/A]

Not integrable

Time = 71.75 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + d \tan(a + bx))}{x} dx$$

input `integrate(atan(c+d*tan(b*x+a))/x,x)`

output `Integral(atan(c + d*tan(a + b*x))/x, x)`

Maxima [F(-1)]

Timed out.

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(arctan(c+d*tan(b*x+a))/x,x, algorithm="maxima")`

output `Timed out`

Giac [N/A]

Not integrable

Time = 0.78 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \int \frac{\arctan(d \tan(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*tan(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctan(d*tan(b*x + a) + c)/x, x)`

Mupad [N/A]

Not integrable

Time = 0.58 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + d \tan(a + bx))}{x} dx$$

input `int(atan(c + d*tan(a + b*x))/x,x)`

output `int(atan(c + d*tan(a + b*x))/x, x)`

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\tan(bx + a) d + c)}{x} dx$$

input `int(atan(c+d*tan(b*x+a))/x,x)`

output `int(atan(tan(a + b*x)*d + c)/x,x)`

3.52 $\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx$

Optimal result	419
Mathematica [A] (verified)	419
Rubi [A] (verified)	420
Maple [C] (warning: unable to verify)	423
Fricas [B] (verification not implemented)	424
Sympy [F(-2)]	424
Maxima [B] (verification not implemented)	425
Giac [F]	425
Mupad [F(-1)]	426
Reduce [F]	426

Optimal result

Integrand size = 21, antiderivative size = 154

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx$$

$$= -\frac{bx^4}{12} + \frac{1}{3}x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{6}ix^3 \log(1 - ice^{2ia+2ibx})$$

$$- \frac{x^2 \text{PolyLog}(2, ice^{2ia+2ibx})}{4b} - \frac{ix \text{PolyLog}(3, ice^{2ia+2ibx})}{4b^2} + \frac{\text{PolyLog}(4, ice^{2ia+2ibx})}{8b^3}$$

output

```
-1/12*b*x^4+1/3*x^3*arctan(c+(1+I*c)*tan(b*x+a))-1/6*I*x^3*ln(1-I*c*exp(2*I*a+2*I*b*x))-1/4*x^2*polylog(2,I*c*exp(2*I*a+2*I*b*x))/b-1/4*I*x*polylog(3,I*c*exp(2*I*a+2*I*b*x))/b^2+1/8*polylog(4,I*c*exp(2*I*a+2*I*b*x))/b^3
```

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.91

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx = \frac{1}{3}x^3 \arctan(c + (1 + ic) \tan(a + bx))$$

$$- \frac{4ib^3x^3 \log\left(1 + \frac{ie^{-2i(a+bx)}}{c}\right) - 6b^2x^2 \text{PolyLog}\left(2, -\frac{ie^{-2i(a+bx)}}{c}\right) + 6ibx \text{PolyLog}\left(3, -\frac{ie^{-2i(a+bx)}}{c}\right) + 3 \text{PolyLog}\left(4, -\frac{ie^{-2i(a+bx)}}{c}\right)}{24b^3}$$

input `Integrate[x^2*ArcTan[c + (1 + I*c)*Tan[a + b*x]],x]`

output `(x^3*ArcTan[c + (1 + I*c)*Tan[a + b*x]])/3 - ((4*I)*b^3*x^3*Log[1 + I/(c*E^((2*I)*(a + b*x)))] - 6*b^2*x^2*PolyLog[2, (-I)/(c*E^((2*I)*(a + b*x)))] + (6*I)*b*x*PolyLog[3, (-I)/(c*E^((2*I)*(a + b*x)))] + 3*PolyLog[4, (-I)/(c*E^((2*I)*(a + b*x)))])/(24*b^3)`

Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.29, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5694, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx \\
 & \quad \downarrow \text{5694} \\
 & \frac{1}{3} x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{3} ib \int \frac{x^3}{e^{2ia+2ibx} c + i} dx \\
 & \quad \downarrow \text{2615} \\
 & \frac{1}{3} x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{3} ib \left(ic \int \frac{e^{2ia+2ibx} x^3}{e^{2ia+2ibx} c + i} dx - \frac{ix^4}{4} \right) \\
 & \quad \downarrow \text{2620} \\
 & \frac{1}{3} x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \\
 & \frac{1}{3} ib \left(ic \left(\frac{3i \int x^2 \log(1 - ice^{2ia+2ibx}) dx}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^4}{4} \right) \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{3} x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \\
 & \frac{1}{3} ib \left(ic \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \int x \text{PolyLog}(2, ice^{2ia+2ibx}) dx}{b} \right)}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^4}{4} \right)
 \end{aligned}$$

$$\begin{aligned} & \downarrow 7163 \\ & \frac{1}{3}x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \\ & \frac{1}{3}ib \left(ic \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int \operatorname{PolyLog}(3, ice^{2ia+2ibx}) dx}{2b} - \frac{ix \operatorname{PolyLog}(3, ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2720 \\ & \frac{1}{3}x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \\ & \frac{1}{3}ib \left(ic \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(3, ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} - \frac{ix \operatorname{PolyLog}(3, ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} - ix^3 \log \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 7143 \\ & \frac{1}{3}x^3 \arctan(c + (1 + ic) \tan(a + bx)) - \\ & \frac{1}{3}ib \left(ic \left(\frac{3i \left(\frac{ix^2 \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\operatorname{PolyLog}(4, ice^{2ia+2ibx})}{4b^2} - \frac{ix \operatorname{PolyLog}(3, ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) \right) \end{aligned}$$

```
input Int[x^2*ArcTan[c + (1 + I*c)*Tan[a + b*x]], x]
```

```
output (x^3*ArcTan[c + (1 + I*c)*Tan[a + b*x]])/3 - (I/3)*b*((-1/4*I)*x^4 + I*c*((-1/2*I)*x^3*Log[1 - I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) + (((3*I)/2)*(((I/2)*x^2*PolyLog[2, I*c*E^((2*I)*a + (2*I)*b*x)]/b - (I*(((1/2*I)*x*PolyLog[3, I*c*E^((2*I)*a + (2*I)*b*x)]/b + PolyLog[4, I*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2))))/b))/(b*c)))
```

Definitions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 5694

```
Int[ArcTan[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tan[a + b*x]]/(f*(m + 1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.92 (sec) , antiderivative size = 1487, normalized size of antiderivative = 9.66

method	result	size
risch	Expression too large to display	1487

input

```
int(x^2*arctan(c+(1+I*c)*tan(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*I/b^3*a^3*ln(1-I*exp(I*(b*x+a))*(-I*c)^(1/2))-1/3*I*x^3*ln(exp(I*(b*x
+a)))+1/2*I/b^2*ln(1-I*exp(2*I*(b*x+a))*c)*x*a^2+1/6*I/b^3*a^3*ln(exp(2*I*
(b*x+a))*c+I)-1/6*I*x^3*ln(1-I*exp(2*I*(b*x+a))*c)-1/2*I/b^3*a^3*ln(1+I*ex
p(I*(b*x+a))*(-I*c)^(1/2))-1/4*x^2*polylog(2,I*exp(2*I*(b*x+a))*c)/b+1/4/b
^3*polylog(2,I*exp(2*I*(b*x+a))*c)*a^2+1/3*I/b^3*ln(1-I*exp(2*I*(b*x+a))*c
)*a^3+1/8*polylog(4,I*exp(2*I*(b*x+a))*c)/b^3-1/4*I*x*polylog(3,I*exp(2*I*
(b*x+a))*c)/b^2-1/2*I/b^2*a^2*ln(1+I*exp(I*(b*x+a))*(-I*c)^(1/2))*x-1/2*I/
b^2*a^2*ln(1-I*exp(I*(b*x+a))*(-I*c)^(1/2))*x+1/6*I*x^3*ln(exp(2*I*(b*x+a)
)*c+I)+1/12*I*(-2*I*Pi-2*ln(-I+c)+I*Pi*csgn(I*exp(2*I*(b*x+a))))*csgn(I*(-I
+c)/(exp(2*I*(b*x+a))+1))*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a)
)+1))-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn(
I*exp(2*I*(b*x+a)))^3+I*Pi*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a)
)+1))^3-I*Pi*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1))^3-I*Pi*csgn
(exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn(I*(-I+c)/(exp(2
*I*(b*x+a))+1))^3-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a)
))*c+I))*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1))+I*Pi*csgn(I*ex
p(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))+1))*csgn(exp(2*I*(b*x+a))*(-I+c)/
(exp(2*I*(b*x+a))+1))-2*I*Pi*csgn(I*exp(I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a)
))^2-I*Pi*csgn(I*exp(2*I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I
*(b*x+a))+1))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(-I+c)/(exp(2*...
```


Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(107) = 214$.

Time = 0.13 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.09

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx = \frac{b^4 x^4 - 2i b^3 x^3 \log\left(-\frac{ce^{(2i bx + 2i a) + i} e^{(-2i bx - 2i a)}}{c - i}\right) + 6 b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{4i} ce^{(i bx + i a)}\right) + 6 b^2 x^2 \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{4i} ce^{(i bx + i a)}\right)}{1}$$

input `integrate(x^2*arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="fricas")`

output `-1/12*(b^4*x^4 - 2*I*b^3*x^3*log(-(c*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/(c - I)) + 6*b^2*x^2*dilog(1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + 6*b^2*x^2*dilog(-1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) - a^4 - 2*I*a^3*log(1/2*(2*c*e^(I*b*x + I*a) + I*sqrt(4*I*c))/c) - 2*I*a^3*log(1/2*(2*c*e^(I*b*x + I*a) - I*sqrt(4*I*c))/c) + 12*I*b*x*polylog(3, 1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + 12*I*b*x*polylog(3, -1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + 2*(I*b^3*x^3 + I*a^3)*log(1/2*sqrt(4*I*c)*e^(I*b*x + I*a) + 1) + 2*(I*b^3*x^3 + I*a^3)*log(-1/2*sqrt(4*I*c)*e^(I*b*x + I*a) + 1) - 12*polylog(4, 1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) - 12*polylog(4, -1/2*sqrt(4*I*c)*e^(I*b*x + I*a)))/b^3`

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x**2*atan(c+(1+I*c)*tan(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) + 1 of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 310 vs. $2(107) = 214$.

Time = 0.06 (sec) , antiderivative size = 310, normalized size of antiderivative = 2.01

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx$$

$$= \frac{4 \left((bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2 \right) \arctan((ic+1) \tan(bx+a) + c)}{b^2} - \frac{(-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(4i(bx+a)^3 - 9i(bx+a)^2 a + 3i a^2)) \operatorname{arctan}_2(c \cos(2bx+2a), c \sin(2bx+2a) + 1) - 3(4i(bx+a)^2 - 6i(bx+a)a + 3i a^2) \operatorname{dilog}(ic e^{(2i b x + 2i a)}) + (4(bx+a)^3 - 9(bx+a)^2 a + 9(bx+a)a^2) \log(c^2 \cos(2bx+2a)^2 + c^2 \sin(2bx+2a)^2 + 2c \sin(2bx+2a) + 1) + 3(4bx+a) \operatorname{polylog}(3, ic e^{(2i b x + 2i a)}) + 6i \operatorname{polylog}(4, ic e^{(2i b x + 2i a)})}{b^2} (Ic + 1) / (b^2 (c - I)) / b$$

input `integrate(x^2*arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="maxima")`

output `1/12*(4*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctan((I*c + 1)*tan(b*x + a) + c)/b^2 - (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*x + a)^2*a^2 - 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*I*(b*x + a)*a^2)*arctan2(c*cos(2*b*x + 2*a), c*sin(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog(I*c*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 + 2*c*sin(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog(3, I*c*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, I*c*e^(2*I*b*x + 2*I*a))*(I*c + 1)/(b^2*(c - I))/b`

Giac [F]

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx = \int x^2 \arctan((ic + 1) \tan(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan((I*c + 1)*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx = \int x^2 \operatorname{atan}(c + \tan(a + bx) (1 + c \operatorname{li})) dx$$

input `int(x^2*atan(c + tan(a + b*x)*(c*1i + 1)),x)`

output `int(x^2*atan(c + tan(a + b*x)*(c*1i + 1)), x)`

Reduce [F]

$$\int x^2 \arctan(c + (1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) ci + \tan(bx + a) + c) x^2 dx$$

input `int(x^2*atan(c+(1+I*c)*tan(b*x+a)),x)`

output `int(atan(tan(a + b*x)*c*i + tan(a + b*x) + c)*x**2,x)`

3.53 $\int x \arctan(c + (1 + ic) \tan(a + bx)) dx$

Optimal result	427
Mathematica [A] (verified)	428
Rubi [A] (verified)	428
Maple [C] (warning: unable to verify)	431
Fricas [B] (verification not implemented)	432
Sympy [F(-2)]	432
Maxima [B] (verification not implemented)	433
Giac [F]	433
Mupad [F(-1)]	434
Reduce [F]	434

Optimal result

Integrand size = 19, antiderivative size = 123

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = -\frac{bx^3}{6} + \frac{1}{2}x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{4}ix^2 \log(1 - ice^{2ia+2ibx}) - \frac{x \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{4b} - \frac{i \operatorname{PolyLog}(3, ice^{2ia+2ibx})}{8b^2}$$

output

```
-1/6*b*x^3+1/2*x^2*arctan(c+(1+I*c)*tan(b*x+a))-1/4*I*x^2*ln(1-I*c*exp(2*I*a+2*I*b*x))-1/4*x*polylog(2,I*c*exp(2*I*a+2*I*b*x))/b-1/8*I*polylog(3,I*c*exp(2*I*a+2*I*b*x))/b^2
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.89

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = \frac{1}{2} x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{i \left(2b^2 x^2 \log \left(1 + \frac{ie^{-2i(a+bx)}}{c} \right) + 2ibx \operatorname{PolyLog} \left(2, -\frac{ie^{-2i(a+bx)}}{c} \right) + \operatorname{PolyLog} \left(3, -\frac{ie^{-2i(a+bx)}}{c} \right) \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c + (1 + I*c)*Tan[a + b*x]], x]
```

output

```
(x^2*ArcTan[c + (1 + I*c)*Tan[a + b*x]])/2 - ((I/8)*(2*b^2*x^2*Log[1 + I/(c*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, (-I)/(c*E^((2*I)*(a + b*x)))] + PolyLog[3, (-I)/(c*E^((2*I)*(a + b*x)))]))/b^2
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.26, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {5694, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arctan(c + (1 + ic) \tan(a + bx)) dx \\ & \quad \downarrow \text{5694} \\ & \frac{1}{2} x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2} ib \int \frac{x^2}{e^{2ia+2ibx} c + i} dx \\ & \quad \downarrow \text{2615} \\ & \frac{1}{2} x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2} ib \left(ic \int \frac{e^{2ia+2ibx} x^2}{e^{2ia+2ibx} c + i} dx - \frac{ix^3}{3} \right) \\ & \quad \downarrow \text{2620} \end{aligned}$$

$$\frac{1}{2}x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2}ib \left(ic \left(\frac{i \int x \log(1 - ice^{2ia+2ibx}) dx}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

↓ 3011

$$\frac{1}{2}x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2}ib \left(ic \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \int \operatorname{PolyLog}(2, ice^{2ia+2ibx}) dx}{2b} \right)}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

↓ 2720

$$\frac{1}{2}x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2}ib \left(ic \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} \right)}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

↓ 7143

$$\frac{1}{2}x^2 \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2}ib \left(ic \left(\frac{i \left(\frac{ix \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{\operatorname{PolyLog}(3, ice^{2ia+2ibx})}{4b^2} \right)}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

input `Int[x*ArcTan[c + (1 + I*c)*Tan[a + b*x]],x]`

output `(x^2*ArcTan[c + (1 + I*c)*Tan[a + b*x]])/2 - (I/2)*b*((-1/3*I)*x^3 + I*c*((-1/2*I)*x^2*Log[1 - I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) + I*(((I/2)*x*PolyLog[2, I*c*E^((2*I)*a + (2*I)*b*x)]/b - PolyLog[3, I*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/(b*c))`

Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5694 `Int[ArcTan[(c_.) + (d_.)*Tan[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tan[a + b*x]]/(f*(m + 1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c + I*d)^2, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.34 (sec) , antiderivative size = 1452, normalized size of antiderivative = 11.80

method	result	size
risch	Expression too large to display	1452

input `int(x*arctan(c+(1+I*c)*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-1/4*I/b^2*ln(1-I*exp(2*I*(b*x+a))*c)*a^2-1/6*x^3*b-1/8*I/b^2*polylog(3,I*
exp(2*I*(b*x+a))*c)+1/2*I/b*a*ln(1-I*exp(I*(b*x+a))*(-I*c)^(1/2))*x+1/8*I*
x^2*(-2*I*Pi-2*ln(-I+c)+I*Pi*csgn(I*exp(2*I*(b*x+a)))*csgn(I*(-I+c)/(exp(2
*I*(b*x+a))+1))*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))+1))-I*Pi*
csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn(I*exp(2*I*
(b*x+a)))^3+I*Pi*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))+1))^3-I*
Pi*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1))^3-I*Pi*csgn(exp(2*I*(
b*x+a))*(-I+c)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn(I*(-I+c)/(exp(2*I*(b*x+a)
)+1))^3-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))*c+I))*c
sgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1))+I*Pi*csgn(I*exp(2*I*(b*
x+a))*(-I+c)/(exp(2*I*(b*x+a))+1))*csgn(exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(
b*x+a))+1))-2*I*Pi*csgn(I*exp(I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a)))^2-I*Pi*
csgn(I*exp(2*I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))+
1))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(-I+c)/(exp(2*I*(b*x+a))+1)
)^2-I*Pi*csgn(I*(-I+c))*csgn(I*(-I+c)/(exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I*
exp(I*(b*x+a)))^2*csgn(I*exp(2*I*(b*x+a)))+I*Pi*csgn(I*(exp(2*I*(b*x+a))*c
+I)/(exp(2*I*(b*x+a))+1))*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))+1)
)^2+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2
*I*(b*x+a))+1))^2+I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I))*csgn(I*(exp(2*I*(b*x
+a))*c+I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*(-I+c)/(exp(2*I*(b*x+a))+...
```


Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 271 vs. $2(85) = 170$.

Time = 0.10 (sec) , antiderivative size = 271, normalized size of antiderivative = 2.20

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = \frac{2b^3x^3 - 3ib^2x^2 \log\left(-\frac{ce^{(2ibx+2ia)+i}e^{(-2ibx-2ia)}}{c-i}\right) + 2a^3 + 6bx\text{Li}_2\left(\frac{1}{2}\sqrt{4i}ce^{(ibx+ia)}\right) + 6bx\text{Li}_2\left(-\frac{1}{2}\sqrt{4i}\right)}{b^2}$$

input `integrate(x*arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="fricas")`

output `-1/12*(2*b^3*x^3 - 3*I*b^2*x^2*log(-(c*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/(c - I)) + 2*a^3 + 6*b*x*dilog(1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + 6*b*x*dilog(-1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + 3*I*a^2*log(1/2*(2*c*e^(I*b*x + I*a) + I*sqrt(4*I*c))/c) + 3*I*a^2*log(1/2*(2*c*e^(I*b*x + I*a) - I*sqrt(4*I*c))/c) + 3*(I*b^2*x^2 - I*a^2)*log(1/2*sqrt(4*I*c)*e^(I*b*x + I*a) + 1) + 3*(I*b^2*x^2 - I*a^2)*log(-1/2*sqrt(4*I*c)*e^(I*b*x + I*a) + 1) + 6*I*polylog(3, 1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + 6*I*polylog(3, -1/2*sqrt(4*I*c)*e^(I*b*x + I*a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x*atan(c+(1+I*c)*tan(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) + 1 o f type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(85) = 170$.

Time = 0.06 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.77

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx$$

$$= \frac{6 \left((bx+a)^2 - 2(bx+a)a \right) \arctan((ic+1) \tan(bx+a)+c)}{b} - \frac{(-4i(bx+a)^3 + 12i(bx+a)^2a - 6i b x \operatorname{Li}_2(i c e^{(2i bx+2i a)}) - 6(i(bx+a)^2 - 2i(bx+a)a))}{b}$$

input `integrate(x*arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="maxima")`

output `1/12*(6*((b*x + a)^2 - 2*(b*x + a)*a)*arctan((I*c + 1)*tan(b*x + a) + c)/b - (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog(I*c*e^(2*I*b*x + 2*I*a)) - 6*(I*(b*x + a)^2 - 2*I*(b*x + a)*a)*arctan2(c*cos(2*b*x + 2*a), c*sin(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 + 2*c*sin(2*b*x + 2*a) + 1) + 3*polylog(3, I*c*e^(2*I*b*x + 2*I*a)))*(I*c + 1)/(b*(c - I)))/b`

Giac [F]

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = \int x \arctan((ic + 1) \tan(bx + a) + c) dx$$

input `integrate(x*arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan((I*c + 1)*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = \int x \operatorname{atan}(c + \tan(a + bx) (1 + c li)) dx$$

input `int(x*atan(c + tan(a + b*x)*(c*1i + 1)),x)`

output `int(x*atan(c + tan(a + b*x)*(c*1i + 1)), x)`

Reduce [F]

$$\int x \arctan(c + (1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) ci + \tan(bx + a) + c) x dx$$

input `int(x*atan(c+(1+I*c)*tan(b*x+a)),x)`

output `int(atan(tan(a + b*x)*c*i + tan(a + b*x) + c)*x,x)`

3.54 $\int \arctan(c + (1 + ic) \tan(a + bx)) dx$

Optimal result	435
Mathematica [B] (warning: unable to verify)	435
Rubi [A] (verified)	436
Maple [B] (verified)	438
Fricas [B] (verification not implemented)	439
Sympy [F(-2)]	440
Maxima [B] (verification not implemented)	440
Giac [F]	441
Mupad [F(-1)]	441
Reduce [F]	441

Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = -\frac{bx^2}{2} + x \arctan(c + (1 + ic) \tan(a + bx)) - \frac{1}{2}ix \log(1 - ice^{2ia+2ibx}) - \frac{\text{PolyLog}(2, ice^{2ia+2ibx})}{4b}$$

output

```
-1/2*b*x^2+x*arctan(c+(1+I*c)*tan(b*x+a))-1/2*I*x*ln(1-I*c*exp(2*I*a+2*I*b*x))-1/4*polylog(2,I*c*exp(2*I*a+2*I*b*x))/b
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 967 vs. 2(85) = 170.

Time = 8.55 (sec) , antiderivative size = 967, normalized size of antiderivative = 11.38

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = x \arctan(c + (1 + ic) \tan(a + bx)) + \frac{((i + c) \cos(a + bx) + (1 + ic) \sin(a + bx)) \left(2bx - i \log \left(1 - \frac{\sec(bx)((-i+c) \cos(a)+i(i+c) \sin(a))(\cos(a+bx)-i \sin(a+bx))}{2c} \right) \right)}{((i + c) \cos(a + bx) + (1 + ic) \sin(a + bx))}$$

input `Integrate[ArcTan[c + (1 + I*c)*Tan[a + b*x]],x]`

output

```
x*ArcTan[c + (1 + I*c)*Tan[a + b*x]] + (I*x*((2*I)*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] - Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x])])/(2*c)]*Log[1 - I*Tan[b*x]] + Log[(Sec[b*x]*((1 - I*c)*Cos[a + b*x] + (-I + c)*Sin[a + b*x])])/(2*Cos[a] - (2*I)*Sin[a])]*Log[1 + I*Tan[b*x]] - PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] - PolyLog[2, (Sec[b*x]*((-I + c)*Cos[a] + I*(I + c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)] + PolyLog[2, (Sec[b*x]*((1 + I*c)*Cos[a] - (I + c)*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2]*Sec[a + b*x]^2*(Cos[b*x] + I*Sin[b*x])*(I*Cos[b*x] + Sin[b*x])*((1 - I*c)*Cos[a + b*x] + (-I + c)*Sin[a + b*x])/(((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x])*(2*b*x - I*Log[1 - (Sec[b*x]*((-I + c)*Cos[a] + I*(I + c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)] - I*Log[1 + (Sec[b*x]*((-1 - I*c)*Cos[a] + (I + c)*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2] - (I*(-I + c)*Cos[a + b*x]*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]])))/((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x]) + ((I + c)*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]])*Sin[a + b*x])/((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x]) - (2*I)*b*x*Tan[b*x] + Log[1 - (Sec[b*x]*((-I + c)*Cos[a] + I*(I + c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)]*Tan[b*x] - Log[1 + (Sec[b*x]*((-1 - I*c)*Cos[a] + (I + c)*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x]))/2]*Tan[b*x] - Log[1 - I*Tan[b*x]]*Tan[b*x] + Cos[a]^2*Log[1 + I*Tan[b*x]]*Tan[b*x] + Log[1 + I*Tan[b*x]]...
```

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.26, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {5686, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx$$

$$\downarrow 5686$$

$$x \arctan(c + (1 + ic) \tan(a + bx)) - ib \int \frac{x}{e^{2ia+2ibx} c + i} dx$$

$$\begin{aligned}
& \downarrow \text{2615} \\
& x \arctan(c + (1 + ic) \tan(a + bx)) - ib \left(ic \int \frac{e^{2ia+2ibx} x}{e^{2ia+2ibx} c + i} dx - \frac{ix^2}{2} \right) \\
& \downarrow \text{2620} \\
& x \arctan(c + (1 + ic) \tan(a + bx)) - \\
& ib \left(ic \left(\frac{i \int \log(1 - ice^{2ia+2ibx}) dx}{2bc} - \frac{ix \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right) \\
& \downarrow \text{2715} \\
& x \arctan(c + (1 + ic) \tan(a + bx)) - \\
& ib \left(ic \left(\frac{\int e^{-2ia-2ibx} \log(1 - ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2c} - \frac{ix \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right) \\
& \downarrow \text{2838} \\
& x \arctan(c + (1 + ic) \tan(a + bx)) - \\
& ib \left(ic \left(-\frac{\text{PolyLog}(2, ice^{2ia+2ibx})}{4b^2c} - \frac{ix \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right)
\end{aligned}$$

input `Int[ArcTan[c + (1 + I*c)*Tan[a + b*x]], x]`

output `x*ArcTan[c + (1 + I*c)*Tan[a + b*x]] - I*b*((-1/2*I)*x^2 + I*c((((-1/2*I)*x*Log[1 - I*c*E^((2*I)*a + (2*I)*b*x)])/(b*c) - PolyLog[2, I*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2*c)))`

Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

```
rule 2620 Int[((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 5686 Int[ArcTan[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]], x_Symbol] := Simp[x*ArcT
an[c + d*Tan[a + b*x]], x] - Simp[I*b Int[x/(c + I*d + c*E^(2*I*a + 2*I*b
*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c + I*d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 562 vs. 2(69) = 138.

Time = 1.55 (sec) , antiderivative size = 563, normalized size of antiderivative = 6.62

method	result
derivativedivides	$\frac{\arctan(c+(ic+1)\tan(bx+a))\ln(-c+(ic+1)\tan(bx+a)+i)c^2}{2i-2c} - \frac{2i\arctan(c+(ic+1)\tan(bx+a))\ln(-c+(ic+1)\tan(bx+a)+i)c}{2i-2c} - \arctan(c+(ic+1)\tan(bx+a))\ln(-c+(ic+1)\tan(bx+a)+i)c^2$
default	$\frac{\arctan(c+(ic+1)\tan(bx+a))\ln(-c+(ic+1)\tan(bx+a)+i)c^2}{2i-2c} - \frac{2i\arctan(c+(ic+1)\tan(bx+a))\ln(-c+(ic+1)\tan(bx+a)+i)c}{2i-2c} - \arctan(c+(ic+1)\tan(bx+a))\ln(-c+(ic+1)\tan(bx+a)+i)c^2$
risch	Expression too large to display

```
input int(arctan(c+(1+I*c)*tan(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/b/(1+I*c)*(arctan(c+(1+I*c)*tan(b*x+a))/(2*I-2*c)*ln(-c+(1+I*c)*tan(b*x+
a)+I)*c^2-2*I*arctan(c+(1+I*c)*tan(b*x+a))/(2*I-2*c)*ln(-c+(1+I*c)*tan(b*x
+a)+I)*c-arctan(c+(1+I*c)*tan(b*x+a))/(2*I-2*c)*ln(-c+(1+I*c)*tan(b*x+a)+I
)-arctan(c+(1+I*c)*tan(b*x+a))/(2*I-2*c)*ln(-I+c+(1+I*c)*tan(b*x+a))*c^2+2
*I*arctan(c+(1+I*c)*tan(b*x+a))/(2*I-2*c)*ln(-I+c+(1+I*c)*tan(b*x+a))*c+ar
ctan(c+(1+I*c)*tan(b*x+a))/(2*I-2*c)*ln(-I+c+(1+I*c)*tan(b*x+a))-(1+I*c)^2
*(1/2/(I-c)*(-1/4*I*ln(-I+c+(1+I*c)*tan(b*x+a))^2+1/2*I*(dilog(-1/2*I*(c+(
1+I*c)*tan(b*x+a)+I))+ln(-I+c+(1+I*c)*tan(b*x+a))*ln(-1/2*I*(c+(1+I*c)*tan
(b*x+a)+I))))-1/2/(I-c)*(1/2*I*(dilog(1/2*(c+(1+I*c)*tan(b*x+a)+I)/c)+ln(-
c+(1+I*c)*tan(b*x+a)+I)*ln(1/2*(c+(1+I*c)*tan(b*x+a)+I)/c))-1/2*I*(dilog((
-I+c+(1+I*c)*tan(b*x+a))/(-2*I+2*c))+ln(-c+(1+I*c)*tan(b*x+a)+I)*ln((-I+c+
(1+I*c)*tan(b*x+a))/(-2*I+2*c))))))

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 202 vs. $2(60) = 120$.

Time = 0.10 (sec) , antiderivative size = 202, normalized size of antiderivative = 2.38

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = \frac{b^2 x^2 - i b x \log\left(-\frac{(ce^{2i bx + 2i a}) + i}{c - i} e^{(-2i bx - 2i a)}\right) - a^2 - (-i b x - i a) \log\left(\frac{1}{2} \sqrt{4i} ce^{(i bx + i a)} + 1\right) - (-i b x - i a) \log\left(\frac{1}{2} \sqrt{4i} ce^{(i bx + i a)} - 1\right)}{b}$$

input

```
integrate(arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="fricas")
```

output

```

-1/2*(b^2*x^2 - I*b*x*log(-(c*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)
)/(c - I)) - a^2 - (-I*b*x - I*a)*log(1/2*sqrt(4*I*c)*e^(I*b*x + I*a) + 1)
- (-I*b*x - I*a)*log(-1/2*sqrt(4*I*c)*e^(I*b*x + I*a) + 1) - I*a*log(1/2*
(2*c*e^(I*b*x + I*a) + I*sqrt(4*I*c))/c) - I*a*log(1/2*(2*c*e^(I*b*x + I*a)
) - I*sqrt(4*I*c))/c) + dilog(1/2*sqrt(4*I*c)*e^(I*b*x + I*a)) + dilog(-1/
2*sqrt(4*I*c)*e^(I*b*x + I*a))/b

```


Sympy [F(-2)]

Exception generated.

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(atan(c+(1+I*c)*tan(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) + 1 o
f type <class 'sympy.core.add.Add'> to QQ_I[b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(60) = 120$.

Time = 0.17 (sec) , antiderivative size = 448, normalized size of antiderivative = 5.27

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="maxima")`

output `-1/8*((I*c + 1)*(4*I*(b*x + a)*log(-2*(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - 2*c + I)/(2*I*c^2 - 2*(c^2 - 2*I*c - 1)*tan(b*x + a) + 2*I))/(I*c + 1) - I*(4*(b*x + a)*(log(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - 2*c + I) - log(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - I)) + I*log(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - 2*c + I)^2 - 2*I*log(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - I)*log(-1/2*(c - I)*tan(b*x + a) + 1/2*I*c + 1/2) + 2*I*log(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - I)*log(-1/2*((I*c + 1)*tan(b*x + a) + c + I)/c + 1) - 2*I*log(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - 2*c + I)*log(-1/2*I*tan(b*x + a) + 1/2) - 2*I*dilog(1/2*(c - I)*tan(b*x + a) - 1/2*I*c + 1/2) + 2*I*dilog(1/2*((I*c + 1)*tan(b*x + a) + c + I)/c) - 2*I*dilog(1/2*I*tan(b*x + a) + 1/2))/(I*c + 1)) - 8*(b*x + a)*arctan((I*c + 1)*tan(b*x + a) + c) + 4*(-I*b*x - I*a)*log(-2*(-I*c^2 + (c^2 - 2*I*c - 1)*tan(b*x + a) - 2*c + I)/(2*I*c^2 - 2*(c^2 - 2*I*c - 1)*tan(b*x + a) + 2*I)))/b`

Giac [F]

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = \int \arctan((ic + 1) \tan(bx + a) + c) dx$$

input `integrate(arctan(c+(1+I*c)*tan(b*x+a)),x, algorithm="giac")`

output `integrate(arctan((I*c + 1)*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(c + \tan(a + bx) (1 + ci)) dx$$

input `int(atan(c + tan(a + b*x)*(c*1i + 1)),x)`

output `int(atan(c + tan(a + b*x)*(c*1i + 1)), x)`

Reduce [F]

$$\int \arctan(c + (1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) ci + \tan(bx + a) + c) dx$$

input `int(atan(c+(1+I*c)*tan(b*x+a)),x)`

output `int(atan(tan(a + b*x)*c*i + tan(a + b*x) + c),x)`

3.55 $\int \frac{\arctan(c+(1+ic)\tan(a+bx))}{x} dx$

Optimal result	442
Mathematica [N/A]	442
Rubi [N/A]	443
Maple [N/A]	443
Fricas [N/A]	444
Sympy [F(-1)]	444
Maxima [F(-2)]	444
Giac [N/A]	445
Mupad [N/A]	445
Reduce [N/A]	446

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\arctan(c + (1 + ic)\tan(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + (1 + ic)\tan(a + bx))}{x}, x\right)$$

output

```
Defer(Int)(arctan(c+(1+I*c)*tan(b*x+a))/x,x)
```

Mathematica [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(c + (1 + ic)\tan(a + bx))}{x} dx = \int \frac{\arctan(c + (1 + ic)\tan(a + bx))}{x} dx$$

input

```
Integrate[ArcTan[c + (1 + I*c)*Tan[a + b*x]]/x,x]
```

output

```
Integrate[ArcTan[c + (1 + I*c)*Tan[a + b*x]]/x, x]
```

Rubi [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx$$

input `Int[ArcTan[c + (1 + I*c)*Tan[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\arctan(c + (ic + 1) \tan(bx + a))}{x} dx$$

input `int(arctan(c+(1+I*c)*tan(b*x+a))/x,x)`

output `int(arctan(c+(1+I*c)*tan(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx = \int \frac{\arctan((ic + 1) \tan(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+(1+I*c)*tan(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c*e^(2*I*b*x + 2*I*a) + I)*e^(-2*I*b*x - 2*I*a)/(c - I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+(1+I*c)*tan(b*x+a))/x,x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx = \text{Exception raised: ValueError}$$

input `integrate(arctan(c+(1+I*c)*tan(b*x+a))/x,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(c-1>0)', see `assume?` for more
details)Is
```

Giac [N/A]

Not integrable

Time = 0.69 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx = \int \frac{\arctan((ic + 1) \tan(bx + a) + c)}{x} dx$$

input

```
integrate(arctan(c+(1+I*c)*tan(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctan((I*c + 1)*tan(b*x + a) + c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.70 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + \tan(a + bx) (1 + c li))}{x} dx$$

input

```
int(atan(c + tan(a + b*x)*(c*1i + 1))/x,x)
```

output

```
int(atan(c + tan(a + b*x)*(c*1i + 1))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

$$\int \frac{\arctan(c + (1 + ic) \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\tan(bx + a) ci + \tan(bx + a) + c)}{x} dx$$

input `int(atan(c+(1+I*c)*tan(b*x+a))/x,x)`output `int(atan(tan(a + b*x)*c*i + tan(a + b*x) + c)/x,x)`

3.56 $\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx$

Optimal result	447
Mathematica [A] (verified)	448
Rubi [A] (verified)	448
Maple [C] (warning: unable to verify)	452
Fricas [B] (verification not implemented)	453
Sympy [F(-2)]	453
Maxima [B] (verification not implemented)	454
Giac [F]	454
Mupad [F(-1)]	455
Reduce [F]	455

Optimal result

Integrand size = 21, antiderivative size = 155

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx)) + \frac{1}{6}ix^3 \log(1 + ice^{2ia+2ibx}) + \frac{x^2 \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{4b} + \frac{ix \operatorname{PolyLog}(3, -ice^{2ia+2ibx})}{4b^2} - \frac{\operatorname{PolyLog}(4, -ice^{2ia+2ibx})}{8b^3}$$

output

```
1/12*b*x^4+1/3*x^3*arctan(c-(1-I*c)*tan(b*x+a))+1/6*I*x^3*ln(1+I*c*exp(2*I*a+2*I*b*x))+1/4*x^2*polylog(2,-I*c*exp(2*I*a+2*I*b*x))/b+1/4*I*x*polylog(3,-I*c*exp(2*I*a+2*I*b*x))/b^2-1/8*polylog(4,-I*c*exp(2*I*a+2*I*b*x))/b^3
```


Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.88

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx = \frac{1}{24} \left(8x^3 \arctan(c + i(i + c) \tan(a + bx)) \right. \\ \left. + 4ix^3 \log \left(1 - \frac{ie^{-2i(a+bx)}}{c} \right) \right. \\ \left. - \frac{6x^2 \operatorname{PolyLog} \left(2, \frac{ie^{-2i(a+bx)}}{c} \right)}{b} \right. \\ \left. + \frac{6ix \operatorname{PolyLog} \left(3, \frac{ie^{-2i(a+bx)}}{c} \right)}{b^2} \right. \\ \left. + \frac{3 \operatorname{PolyLog} \left(4, \frac{ie^{-2i(a+bx)}}{c} \right)}{b^3} \right)$$

input

```
Integrate[x^2*ArcTan[c + (-1 + I*c)*Tan[a + b*x]],x]
```

output

```
(8*x^3*ArcTan[c + I*(I + c)*Tan[a + b*x]] + (4*I)*x^3*Log[1 - I/(c*E^((2*I)*(a + b*x)))] - (6*x^2*PolyLog[2, I/(c*E^((2*I)*(a + b*x)))]/b + ((6*I)*x*PolyLog[3, I/(c*E^((2*I)*(a + b*x)))]/b^2 + (3*PolyLog[4, I/(c*E^((2*I)*(a + b*x)))]/b^3))/24
```

Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.28, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {5694, 25, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

↓ 5694

$$\frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx)) - \frac{1}{3}ib \int -\frac{x^3}{i - ce^{2ia+2ibx}} dx$$

↓ 25

$$\frac{1}{3}ib \int \frac{x^3}{i - ce^{2ia+2ibx}} dx + \frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 2615

$$\frac{1}{3}ib \left(-ic \int \frac{e^{2ia+2ibx} x^3}{i - ce^{2ia+2ibx}} dx - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 2620

$$\frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \int x^2 \log(ie^{2ia+2ibx}c + 1) dx}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 3011

$$\frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \int x \text{PolyLog}(2, -ice^{2ia+2ibx}) dx}{b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 7163

$$\frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{i \int \text{PolyLog}(3, -ice^{2ia+2ibx}) dx}{2b} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} \right) + \frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx)) \right)$$

↓ 2720

$$\frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int e^{-2ia-2ibx} \text{PolyLog}(3, -ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{b} \right)}{2bc} \right)}{2bc} \right) \right)$$

$$\frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 7143

$$\frac{1}{3}x^3 \arctan(c - (1 - ic) \tan(a + bx)) + \frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\text{PolyLog}(4, -ice^{2ia+2ibx})}{4b^2} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{2b} \right)}{2bc} \right)}{2bc} \right) \right)$$

```
input Int[x^2*ArcTan[c + (-1 + I*c)*Tan[a + b*x]],x]
```

```
output (x^3*ArcTan[c - (1 - I*c)*Tan[a + b*x]]/3 + (I/3)*b*((-1/4*I)*x^4 - I*c*((I/2)*x^3*Log[1 + I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) - (((3*I)/2)*((I/2)*x^2*PolyLog[2, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/b - (I*((-1/2*I)*x*PolyLog[3, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/b + PolyLog[4, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/b)/(b*c))
```

Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 2615 Int[((c._) + (d._)*(x._))^(m._)/((a._) + (b._)*((F._)^((g._)*((e._) + (f._)*(x._))))^(n._), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5694

```
Int[ArcTan[(c_) + (d_)*Tan[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tan[a + b*x]]/(f*(m +
1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c + I*d + c*E^(2
*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && E
qQ[(c + I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[(((e_) + (f_)*(x_)^(m_))*PolyLog[n_, (d_)*((F_)^((c_)*((a_) + (b_
)*(x_))))^(p_)]), x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.00 (sec) , antiderivative size = 1488, normalized size of antiderivative = 9.60

method	result	size
risch	Expression too large to display	1488

input `int(x^2*arctan(c+(-1+I*c)*tan(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/6*I/b^3*a^3*\ln(-\exp(2*I*(b*x+a))*c+I)+1/12*I*(-2*I*Pi+2*\ln(I+c)-I*Pi*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))*csgn((\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))*csgn(\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))+I*Pi*csgn(I/(\exp(2*I*(b*x+a))+1)*(I+c))*csgn(I*\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I/(\exp(2*I*(b*x+a))+1))*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*(\exp(2*I*(b*x+a))*c-I))*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I*(I+c))*csgn(I/(\exp(2*I*(b*x+a))+1)*(I+c))^2+I*Pi*csgn(I*\exp(2*I*(b*x+a)))*csgn(I*\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*\exp(2*I*(b*x+a)))*csgn(I/(\exp(2*I*(b*x+a))+1)*(I+c))*csgn(I*\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))-I*Pi*csgn(I/(\exp(2*I*(b*x+a))+1))*csgn(I*(I+c))*csgn(I/(\exp(2*I*(b*x+a))+1)*(I+c))+I*Pi*csgn(I/(\exp(2*I*(b*x+a))+1))*csgn(I*(\exp(2*I*(b*x+a))*c-I))*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))-I*Pi*csgn(\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))^3-I*Pi*csgn(I/(\exp(2*I*(b*x+a))+1)*(I+c))^3+I*Pi*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))^3-I*Pi*csgn(I*\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))^3+I*Pi*csgn((\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I*\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))*csgn(\exp(2*I*(b*x+a))*(I+c)/(\exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I/(\exp(2*I*(b*x+a))+1))*csgn(I/(\exp(2*I*(b*x+a))+1)*(I+c))^2+I*Pi*csgn(e...
 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(108) = 216$.

Time = 0.11 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.08

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

$$= \frac{b^4 x^4 + 2i b^3 x^3 \log\left(-\frac{(c+i)e^{(2i bx+2i a)}}{ce^{(2i bx+2i a)}-i}\right) + 6 b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4i} ce^{(i bx+i a)}\right) + 6 b^2 x^2 \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{-4i} ce^{(i bx+i a)}\right) - a^4 - 2I a^3 \log\left(\frac{1}{2}(2c e^{(I b x + I a)} + I \sqrt{-4I c})/c\right) - 2I a^3 \log\left(\frac{1}{2}(2c e^{(I b x + I a)} - I \sqrt{-4I c})/c\right) + 12I b x \operatorname{polylog}(3, \frac{1}{2} \sqrt{-4I c} e^{(I b x + I a)}) + 12I b x \operatorname{polylog}(3, -\frac{1}{2} \sqrt{-4I c} e^{(I b x + I a)}) - 2(-I b^3 x^3 - I a^3) \log\left(\frac{1}{2} \sqrt{-4I c} e^{(I b x + I a)} + 1\right) - 2(-I b^3 x^3 - I a^3) \log\left(-\frac{1}{2} \sqrt{-4I c} e^{(I b x + I a)} + 1\right) - 12 \operatorname{polylog}(4, \frac{1}{2} \sqrt{-4I c} e^{(I b x + I a)}) - 12 \operatorname{polylog}(4, -\frac{1}{2} \sqrt{-4I c} e^{(I b x + I a)})}{b^3}$$

input `integrate(x^2*arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="fricas")`

output `1/12*(b^4*x^4 + 2*I*b^3*x^3*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x + 2*I*a) - I)) + 6*b^2*x^2*dilog(1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) + 6*b^2*x^2*dilog(-1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) - a^4 - 2*I*a^3*log(1/2*(2*c*e^(I*b*x + I*a) + I*sqrt(-4*I*c))/c) - 2*I*a^3*log(1/2*(2*c*e^(I*b*x + I*a) - I*sqrt(-4*I*c))/c) + 12*I*b*x*polylog(3, 1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) + 12*I*b*x*polylog(3, -1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) - 2*(-I*b^3*x^3 - I*a^3)*log(1/2*sqrt(-4*I*c)*e^(I*b*x + I*a) + 1) - 2*(-I*b^3*x^3 - I*a^3)*log(-1/2*sqrt(-4*I*c)*e^(I*b*x + I*a) + 1) - 12*polylog(4, 1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) - 12*polylog(4, -1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)))/b^3`

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x**2*atan(c+(-1+I*c)*tan(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) + 1 of f type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 310 vs. $2(108) = 216$.

Time = 0.07 (sec) , antiderivative size = 310, normalized size of antiderivative = 2.00

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

$$= \frac{4((bx+a)^3 - 3(bx+a)^2a + 3(bx+a)a^2) \arctan((ic-1) \tan(bx+a) + c)}{b^2} + \frac{(-3i(bx+a)^4 + 12i(bx+a)^3a - 18i(bx+a)^2a^2 - 2(-4i(bx+a)^3 + 9i(bx+a)^2a - 9i(bx+a)a^2) \arctan2(c \cos(2bx+2a), -c \sin(2bx+2a) + 1) - 3(4I(bx+a)^2 - 6I(bx+a)a + 3Ia^2) \operatorname{dilog}(-Ic e^{(2Ibx+2Ia)}) + (4(bx+a)^3 - 9(bx+a)^2a + 9(bx+a)a^2) \log(c^2 \cos(2bx+2a)^2 + c^2 \sin(2bx+2a)^2 - 2c \sin(2bx+2a) + 1) + 3(4bx+a) \operatorname{polylog}(3, -Ic e^{(2Ibx+2Ia)}) + 6I \operatorname{polylog}(4, -Ic e^{(2Ibx+2Ia)})) (Ic - 1) / (b^2(c + I))}{b}$$

input `integrate(x^2*arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="maxima")`

output `1/12*(4*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctan((I*c - 1)*tan(b*x + a) + c)/b^2 + (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*x + a)^2*a^2 - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a - 9*I*(b*x + a)*a^2)*arctan2(c*cos(2*b*x + 2*a), -c*sin(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog(-I*c*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 - 2*c*sin(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog(3, -I*c*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, -I*c*e^(2*I*b*x + 2*I*a)))*(I*c - 1)/(b^2*(c + I))/b`

Giac [F]

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int x^2 \arctan((ic - 1) \tan(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan((I*c - 1)*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int x^2 \operatorname{atan}(c + \tan(a + bx) (-1 + ci)) dx$$

input `int(x^2*atan(c + tan(a + b*x)*(c*1i - 1)),x)`output `int(x^2*atan(c + tan(a + b*x)*(c*1i - 1)), x)`**Reduce [F]**

$$\int x^2 \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) ci - \tan(bx + a) + c) x^2 dx$$

input `int(x^2*atan(c+(-1+I*c)*tan(b*x+a)),x)`output `int(atan(tan(a + b*x)*c*i - tan(a + b*x) + c)*x**2,x)`

3.57 $\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx$

Optimal result	456
Mathematica [A] (verified)	457
Rubi [A] (verified)	457
Maple [C] (warning: unable to verify)	460
Fricas [B] (verification not implemented)	461
Sympy [F(-2)]	461
Maxima [B] (verification not implemented)	462
Giac [F]	462
Mupad [F(-1)]	463
Reduce [F]	463

Optimal result

Integrand size = 19, antiderivative size = 124

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \arctan(c - (1 - ic) \tan(a + bx))$$

$$+ \frac{1}{4}ix^2 \log(1 + ice^{2ia+2ibx})$$

$$+ \frac{x \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{4b}$$

$$+ \frac{i \operatorname{PolyLog}(3, -ice^{2ia+2ibx})}{8b^2}$$

output

```
1/6*b*x^3+1/2*x^2*arctan(c-(1-I*c)*tan(b*x+a))+1/4*I*x^2*ln(1+I*c*exp(2*I*
a+2*I*b*x))+1/4*x*polylog(2,-I*c*exp(2*I*a+2*I*b*x))/b+1/8*I*polylog(3,-I*
c*exp(2*I*a+2*I*b*x))/b^2
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.90

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx = \frac{1}{2} x^2 \arctan(c + i(i + c) \tan(a + bx)) + \frac{i \left(2b^2 x^2 \log \left(1 - \frac{ie^{-2i(a+bx)}}{c} \right) + 2ibx \operatorname{PolyLog} \left(2, \frac{ie^{-2i(a+bx)}}{c} \right) + \operatorname{PolyLog} \left(3, \frac{ie^{-2i(a+bx)}}{c} \right) \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c + (-1 + I*c)*Tan[a + b*x]],x]
```

output

```
(x^2*ArcTan[c + I*(I + c)*Tan[a + b*x]])/2 + ((I/8)*(2*b^2*x^2*Log[1 - I/(c*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, I/(c*E^((2*I)*(a + b*x)))] + PolyLog[3, I/(c*E^((2*I)*(a + b*x)))]))/b^2
```

Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.26, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {5694, 25, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arctan(c + (-1 + ic) \tan(a + bx)) dx \\ & \quad \downarrow \text{5694} \\ & \frac{1}{2} x^2 \arctan(c - (1 - ic) \tan(a + bx)) - \frac{1}{2} ib \int -\frac{x^2}{i - ce^{2ia+2ibx}} dx \\ & \quad \downarrow \text{25} \\ & \frac{1}{2} ib \int \frac{x^2}{i - ce^{2ia+2ibx}} dx + \frac{1}{2} x^2 \arctan(c - (1 - ic) \tan(a + bx)) \\ & \quad \downarrow \text{2615} \\ & \frac{1}{2} ib \left(-ic \int \frac{e^{2ia+2ibx} x^2}{i - ce^{2ia+2ibx}} dx - \frac{ix^3}{3} \right) + \frac{1}{2} x^2 \arctan(c - (1 - ic) \tan(a + bx)) \end{aligned}$$

$$\frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \int x \log(ie^{2ia+2ibx}c + 1) dx}{bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 2620

$$\frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \left(\frac{ix \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -ice^{2ia+2ibx}) dx}{2b} \right)}{bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 2720

$$\frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \left(\frac{ix \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, -ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} \right)}{bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c - (1 - ic) \tan(a + bx))$$

↓ 7143

$$\frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \left(\frac{ix \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{\operatorname{PolyLog}(3, -ice^{2ia+2ibx})}{4b^2} \right)}{bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c - (1 - ic) \tan(a + bx))$$

input `Int[x*ArcTan[c + (-1 + I*c)*Tan[a + b*x]],x]`

output

```
(x^2*ArcTan[c - (1 - I*c)*Tan[a + b*x]])/2 + (I/2)*b*((-1/3*I)*x^3 - I*c*((I/2)*x^2*Log[1 + I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) - (I*(((I/2)*x*PolyLog[2, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/b - PolyLog[3, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/(b*c)))
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$
- rule 2615 $\text{Int}[((c_.) + (d_.)*(x_))^m / ((a_.) + (b_.)*((F_)^{(g_.)*((e_.) + (f_.)*(x_))})^n)), x_Symbol] \rightarrow \text{Simp}[(c + d*x)^{m+1} / (a*d*(m+1)), x] - \text{Simp}[b/a \quad \text{Int}[(c + d*x)^m * ((F^{(g*(e + f*x)))^n} / (a + b*(F^{(g*(e + f*x)))^n}))], x], x] /;$ $\text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$
- rule 2620 $\text{Int}[(((F_)^{(g_.)*((e_.) + (f_.)*(x_))})^n) * ((c_.) + (d_.)*(x_))^m / ((a_.) + (b_.)*((F_)^{(g_.)*((e_.) + (f_.)*(x_))})^n)), x_Symbol] \rightarrow \text{Simp}[(c + d*x)^m / (b*f*g*n*\text{Log}[F])] * \text{Log}[1 + b*((F^{(g*(e + f*x)))^n} / a)], x] - \text{Simp}[d*(m / (b*f*g*n*\text{Log}[F])) \quad \text{Int}[(c + d*x)^{m-1} * \text{Log}[1 + b*((F^{(g*(e + f*x)))^n} / a)]], x], x] /;$ $\text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$
- rule 2720 $\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Simp}[v/D[v, x] \quad \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /;$ $\text{FunctionOfExponentialQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (w_)*((a_)*(v_)^n)^m] /;$ $\text{FreeQ}\{a, m, n\}, x\} \ \&\& \ \text{IntegerQ}[m*n] \ \&\& \ !\text{MatchQ}[u, E^{(c_)*((a_.) + (b_.)*x)} * (F_)[v_]] /;$ $\text{FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{InverseFunctionQ}[F[x]]]$
- rule 3011 $\text{Int}[\text{Log}[1 + (e_)*((F_)^{(c_)*((a_.) + (b_.)*(x_))})^n)] * ((f_.) + (g_.)*(x_))^m, x_Symbol] \rightarrow \text{Simp}[(-f + g*x)^m * (\text{PolyLog}[2, (-e)*(F^{(c*(a + b*x)))^n} / (b*c*n*\text{Log}[F])]), x] + \text{Simp}[g*(m / (b*c*n*\text{Log}[F])) \quad \text{Int}[(f + g*x)^{m-1} * \text{PolyLog}[2, (-e)*(F^{(c*(a + b*x)))^n}], x], x] /;$ $\text{FreeQ}\{F, a, b, c, e, f, g, n\}, x\} \ \&\& \ \text{GtQ}[m, 0]$
- rule 5694 $\text{Int}[\text{ArcTan}[(c_.) + (d_.)*\text{Tan}[(a_.) + (b_.)*(x_)]] * ((e_.) + (f_.)*(x_))^m, x_Symbol] \rightarrow \text{Simp}[(e + f*x)^{m+1} * (\text{ArcTan}[c + d*\text{Tan}[a + b*x]] / (f*(m + 1))), x] - \text{Simp}[I*(b / (f*(m + 1))) \quad \text{Int}[(e + f*x)^{m+1} / (c + I*d + c*E^{(2*I*a + 2*I*b*x)}), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{EqQ}[(c + I*d)^2, -1]$

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.50 (sec) , antiderivative size = 1453, normalized size of antiderivative = 11.72

method	result	size
risch	Expression too large to display	1453

input

```
int(x*arctan(c+(-1+I*c)*tan(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*I/b*a*ln(1+I*exp(I*(b*x+a))*(I*c)^(1/2))*x-1/2*I/b^2*a^2*ln(1+I*exp(I*(b*x+a))*(I*c)^(1/2))+1/8*I*x^2*(-2*I*Pi+2*ln(I+c)-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))*csgn((exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))*csgn(exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1)*(I+c))*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c-I))*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I*(I+c))*csgn(I/(exp(2*I*(b*x+a))+1)*(I+c))^2+I*Pi*csgn(I*exp(2*I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))^2-I*Pi*csgn(I*exp(2*I*(b*x+a)))*csgn(I/(exp(2*I*(b*x+a))+1)*(I+c))*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(I+c))*csgn(I/(exp(2*I*(b*x+a))+1)*(I+c))+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I*(exp(2*I*(b*x+a))*c-I))*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))-I*Pi*csgn(exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))^3-I*Pi*csgn(I/(exp(2*I*(b*x+a))+1)*(I+c))^3+I*Pi*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))^3-I*Pi*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))^3+I*Pi*csgn((exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))*csgn(exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))+1))^2+I*Pi*csgn(I/(exp(2*I*(b*x+a))+1))*csgn(I/(exp(2*I*(b...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 271 vs. $2(86) = 172$.

Time = 0.10 (sec) , antiderivative size = 271, normalized size of antiderivative = 2.19

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

$$= \frac{2b^3x^3 + 3ib^2x^2 \log\left(-\frac{(c+i)e^{(2ibx+2ia)}}{ce^{(2ibx+2ia)}-i}\right) + 2a^3 + 6bx\text{Li}_2\left(\frac{1}{2}\sqrt{-4i}ce^{(ibx+ia)}\right) + 6bx\text{Li}_2\left(-\frac{1}{2}\sqrt{-4i}ce^{(ibx+ia)}\right)}{b^2}$$

input `integrate(x*arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*b^3*x^3 + 3*I*b^2*x^2*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x + 2*I*a) - I)) + 2*a^3 + 6*b*x*dilog(1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) + 6*b*x*dilog(-1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) + 3*I*a^2*log(1/2*(2*c*e^(I*b*x + I*a) + I*sqrt(-4*I*c))/c) + 3*I*a^2*log(1/2*(2*c*e^(I*b*x + I*a) - I*sqrt(-4*I*c))/c) - 3*(-I*b^2*x^2 + I*a^2)*log(1/2*sqrt(-4*I*c)*e^(I*b*x + I*a) + 1) - 3*(-I*b^2*x^2 + I*a^2)*log(-1/2*sqrt(-4*I*c)*e^(I*b*x + I*a) + 1) + 6*I*polylog(3, 1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)) + 6*I*polylog(3, -1/2*sqrt(-4*I*c)*e^(I*b*x + I*a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x*atan(c+(-1+I*c)*tan(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) + 1 o f type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(86) = 172$.

Time = 0.05 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.76

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

$$= \frac{6((bx+a)^2 - 2(bx+a)a) \arctan((ic-1) \tan(bx+a) + c)}{b} + \frac{(-4i(bx+a)^3 + 12i(bx+a)^2 a - 6i b x \operatorname{Li}_2(-i c e^{2i bx + 2i a})) - 6(-i(bx+a)^2 + 2i(bx+a)a)}{b}$$

input `integrate(x*arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="maxima")`

output `1/12*(6*((b*x + a)^2 - 2*(b*x + a)*a)*arctan((I*c - 1)*tan(b*x + a) + c)/b + (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog(-I*c*e^(2*I*b*x + 2*I*a)) - 6*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*arctan2(c*cos(2*b*x + 2*a), -c*sin(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 - 2*c*sin(2*b*x + 2*a) + 1) + 3*polylog(3, -I*c*e^(2*I*b*x + 2*I*a)))*(I*c - 1)/(b*(c + I))/b`

Giac [F]

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int x \arctan((ic - 1) \tan(bx + a) + c) dx$$

input `integrate(x*arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan((I*c - 1)*tan(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int x \operatorname{atan}(c + \tan(a + bx) (-1 + ci)) dx$$

input `int(x*atan(c + tan(a + b*x)*(c*1i - 1)),x)`

output `int(x*atan(c + tan(a + b*x)*(c*1i - 1)), x)`

Reduce [F]

$$\int x \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) ci - \tan(bx + a) + c) x dx$$

input `int(x*atan(c+(-1+I*c)*tan(b*x+a)),x)`

output `int(atan(tan(a + b*x)*c*i - tan(a + b*x) + c)*x,x)`

3.58 $\int \arctan(c + (-1 + ic) \tan(a + bx)) dx$

Optimal result	464
Mathematica [B] (warning: unable to verify)	464
Rubi [A] (verified)	465
Maple [B] (verified)	467
Fricas [B] (verification not implemented)	468
Sympy [F(-2)]	469
Maxima [B] (verification not implemented)	469
Giac [F]	470
Mupad [F(-1)]	470
Reduce [F]	471

Optimal result

Integrand size = 17, antiderivative size = 86

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = \frac{bx^2}{2} + x \arctan(c - (1 - ic) \tan(a + bx)) + \frac{1}{2}ix \log(1 + ice^{2ia+2ibx}) + \frac{\text{PolyLog}(2, -ice^{2ia+2ibx})}{4b}$$

output

```
1/2*b*x^2+x*arctan(c-(1-I*c)*tan(b*x+a))+1/2*I*x*ln(1+I*c*exp(2*I*a+2*I*b*x))+1/4*polylog(2,-I*c*exp(2*I*a+2*I*b*x))/b
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 847 vs. 2(86) = 172.

Time = 6.25 (sec) , antiderivative size = 847, normalized size of antiderivative = 9.85

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = x \arctan(c + i(i + c) \tan(a + bx)) + \frac{ix \left(-2ibx \log(2 \cos(bx)(\cos(bx) - i \sin(bx))) + \log \left(\frac{\sec(bx)(\cos(a+bx) - i \sin(a+bx))}{2c} \right) \right)}{((-i + c) \cos(a + bx) + i(i + c) \sin(a + bx)) \left(-2bx + i \log \left(1 - \frac{\sec(bx)((i+c) \cos(a)+(1+ic) \sin(a))(\cos(a+bx)-i \sin(a+bx))}{2c} \right) \right)}$$

input `Integrate[ArcTan[c + (-1 + I*c)*Tan[a + b*x]],x]`

output

```
x*ArcTan[c + I*(I + c)*Tan[a + b*x]] + (I*x*((-2*I)*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] + Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((-I + c)*Cos[a + b*x] + I*(I + c)*Sin[a + b*x]))/(2*c)]*Log[1 - I*Tan[b*x]] - Log[(Sec[b*x]*(Cos[a] + I*Sin[a])*((1 + I*c)*Cos[a + b*x] - (I + c)*Sin[a + b*x]))/2]*Log[1 + I*Tan[b*x]] + PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] + PolyLog[2, (Sec[b*x]*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)] - PolyLog[2, ((Cos[a] + I*Sin[a])*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(-I + Tan[b*x]))/2])*Sec[a + b*x]*(Cos[b*x] + I*Sin[b*x])*(I*Cos[b*x] + Sin[b*x]))/(((I + c)*Cos[a + b*x] + I*(I + c)*Sin[a + b*x])*(-2*b*x + I*Log[1 - (Sec[b*x]*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)] + (I*(I + c)*Cos[a + b*x]*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]]))/((-I + c)*Cos[a + b*x] + I*(I + c)*Sin[a + b*x]) + ((1 + I*c)*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]])*Sin[a + b*x])/((-I - I*c)*Cos[a + b*x] + (I + c)*Sin[a + b*x]) + (2*I)*b*x*Tan[b*x] - Log[1 - (Sec[b*x]*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)]*Tan[b*x] + Log[1 - I*Tan[b*x]]*Tan[b*x] - Log[1 + I*Tan[b*x]]*Tan[b*x] - (Log[(Sec[b*x]*(Cos[a] + I*Sin[a])*((1 + I*c)*Cos[a + b*x] - (I + c)*Sin[a + b*x]))/2]*Sec[b*x]^2)/(-I + Tan[b*x]) + (Log[1 - ((Cos[a] + I*Sin[a])*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(-I + Tan[b*x]))/2]*Sec[b*x]^2)/(-I + Tan[b*x]) + (Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((-I + c)*Cos...
```

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.26, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {5686, 25, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

$$\downarrow 5686$$

$$x \arctan(c - (1 - ic) \tan(a + bx)) - ib \int -\frac{x}{i - ce^{2ia+2ibx}} dx$$

$$\begin{aligned}
& \downarrow 25 \\
& ib \int \frac{x}{i - ce^{2ia+2ibx}} dx + x \arctan(c - (1 - ic) \tan(a + bx)) \\
& \downarrow 2615 \\
& ib \left(-ic \int \frac{e^{2ia+2ibx} x}{i - ce^{2ia+2ibx}} dx - \frac{ix^2}{2} \right) + x \arctan(c - (1 - ic) \tan(a + bx)) \\
& \downarrow 2620 \\
& ib \left(-ic \left(\frac{ix \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \int \log(ie^{2ia+2ibx} c + 1) dx}{2bc} \right) - \frac{ix^2}{2} \right) + x \arctan(c - (1 - ic) \tan(a + bx)) \\
& \downarrow 2715 \\
& ib \left(-ic \left(\frac{ix \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{\int e^{-2ia-2ibx} \log(ie^{2ia+2ibx} c + 1) de^{2ia+2ibx}}{4b^2 c} \right) - \frac{ix^2}{2} \right) + \\
& \quad x \arctan(c - (1 - ic) \tan(a + bx)) \\
& \downarrow 2838 \\
& \quad x \arctan(c - (1 - ic) \tan(a + bx)) + \\
& ib \left(-ic \left(\frac{\text{PolyLog}(2, -ice^{2ia+2ibx})}{4b^2 c} + \frac{ix \log(1 + ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right)
\end{aligned}$$

input `Int[ArcTan[c + (-1 + I*c)*Tan[a + b*x]],x]`

output `x*ArcTan[c - (1 - I*c)*Tan[a + b*x]] + I*b*((-1/2*I)*x^2 - I*c*((I/2)*x*Log[1 + I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) + PolyLog[2, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2*c))`

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2615 $\text{Int}[(((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.}))))^{\text{n}_.}], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m} + 1} / (\text{a} * \text{d} * (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b}/\text{a} \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m}} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}}) / (\text{a} + \text{b} * (\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}})], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2620 $\text{Int}[(((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.}], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m}} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}]) * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}}) / \text{a}], \text{x}] - \text{Simp}[\text{d} * (\text{m} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m} - 1} * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}}) / \text{a}], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2715 $\text{Int}[\text{Log}[(\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.}], \text{x_Symbol}] \rightarrow \text{Simp}[1 / (\text{d} * \text{e} * \text{n} * \text{Log}[\text{F}]) \quad \text{Subst}[\text{Int}[\text{Log}[\text{a} + \text{b} * \text{x}] / \text{x}, \text{x}], \text{x}, (\text{F}^{\text{e}}(\text{c} + \text{d} * \text{x}))^{\text{n}}], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}\} \&\& \text{GtQ}[\text{a}, 0]$
- rule 2838 $\text{Int}[\text{Log}[(\text{c}_.) * ((\text{d}_.) + (\text{e}_.) * (\text{x}_.)^{\text{n}_.})] / (\text{x}_.), \text{x_Symbol}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-\text{c}) * \text{e} * \text{x}^{\text{n}}] / \text{n}, \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}\} \&\& \text{EqQ}[\text{c} * \text{d}, 1]$
- rule 5686 $\text{Int}[\text{ArcTan}[(\text{c}_.) + (\text{d}_.) * \text{Tan}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)]], \text{x_Symbol}] \rightarrow \text{Simp}[\text{x} * \text{ArcTan}[\text{c} + \text{d} * \text{Tan}[\text{a} + \text{b} * \text{x}]], \text{x}] - \text{Simp}[\text{I} * \text{b} \quad \text{Int}[\text{x} / (\text{c} + \text{I} * \text{d} + \text{c} * \text{E}^{(2 * \text{I} * \text{a} + 2 * \text{I} * \text{b} * \text{x})}), \text{x}], \text{x}] \text{ ; FreeQ}\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}\} \&\& \text{EqQ}[(\text{c} + \text{I} * \text{d})^2, -1]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 593 vs. $2(70) = 140$.

Time = 1.59 (sec) , antiderivative size = 594, normalized size of antiderivative = 6.91

method	result
derivativdivides	$-\frac{\arctan(c+(ic-1)\tan(bx+a))\ln(c-(ic-1)\tan(bx+a)+i)c^2}{2i+2c} - \frac{2i\arctan(c+(ic-1)\tan(bx+a))\ln(c-(ic-1)\tan(bx+a)+i)c}{2i+2c} + \arctan(c+(ic-1)\tan(bx+a))\ln(c-(ic-1)\tan(bx+a)+i)$
default	$-\frac{\arctan(c+(ic-1)\tan(bx+a))\ln(c-(ic-1)\tan(bx+a)+i)c^2}{2i+2c} - \frac{2i\arctan(c+(ic-1)\tan(bx+a))\ln(c-(ic-1)\tan(bx+a)+i)c}{2i+2c} + \arctan(c+(ic-1)\tan(bx+a))\ln(c-(ic-1)\tan(bx+a)+i)$
risch	Expression too large to display

```
input int(arctan(c+(-1+I*c)*tan(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output 1/b/(-1+I*c)*(-arctan(c+(-1+I*c)*tan(b*x+a))/(2*I+2*c)*ln(c-(-1+I*c)*tan(b*x+a)+I)*c^2-2*I*arctan(c+(-1+I*c)*tan(b*x+a))/(2*I+2*c)*ln(c-(-1+I*c)*tan(b*x+a)+I)+arctan(c+(-1+I*c)*tan(b*x+a))/(2*I+2*c)*ln(c-(-1+I*c)*tan(b*x+a)+I)+arctan(c+(-1+I*c)*tan(b*x+a))/(2*I+2*c)*ln(I+c+(-1+I*c)*tan(b*x+a))*c^2+2*I*arctan(c+(-1+I*c)*tan(b*x+a))/(2*I+2*c)*ln(I+c+(-1+I*c)*tan(b*x+a))*c-arctan(c+(-1+I*c)*tan(b*x+a))/(2*I+2*c)*ln(I+c+(-1+I*c)*tan(b*x+a))+(-1+I*c)^2*(1/2/(I+c)*(1/4*I*ln(I+c+(-1+I*c)*tan(b*x+a))^2-1/2*I*((ln(I+c+(-1+I*c)*tan(b*x+a))-ln(-1/2*I*(I+c+(-1+I*c)*tan(b*x+a))))*ln(-1/2*I*(I-c-(-1+I*c)*tan(b*x+a)))-dilog(-1/2*I*(I+c+(-1+I*c)*tan(b*x+a)))))-1/2/(I+c)*(1/2*I*(dilog((-I-c-(-1+I*c)*tan(b*x+a))/(-2*I-2*c))+ln(c-(-1+I*c)*tan(b*x+a)+I)*ln((-I-c-(-1+I*c)*tan(b*x+a))/(-2*I-2*c)))-1/2*I*(dilog(-1/2*(I-c-(-1+I*c)*tan(b*x+a))/c)+ln(c-(-1+I*c)*tan(b*x+a)+I)*ln(-1/2*(I-c-(-1+I*c)*tan(b*x+a))/c))))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(61) = 122$.

Time = 0.09 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.33

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx$$

$$= \frac{b^2 x^2 + i b x \log\left(-\frac{(c+i)e^{(2i bx + 2i a)}}{ce^{(2i bx + 2i a)} - i}\right) - a^2 + (i b x + i a) \log\left(\frac{1}{2} \sqrt{-4i} ce^{(i bx + i a)} + 1\right) + (i b x + i a) \log\left(-\frac{1}{2} \sqrt{-4i} ce^{(i bx + i a)} - 1\right)}{b}$$

input `integrate(arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="fricas")`

output
$$\frac{1}{2}(b^2x^2 + Ibx \log(-c + I)e^{(2Ibx + 2Ia)}/(ce^{(2Ibx + 2Ia)} - I)) - a^2 + (Ibx + Ia) \log(1/2\sqrt{-4Ic}e^{(Ibx + Ia)} + 1) + (Ibx + Ia) \log(-1/2\sqrt{-4Ic}e^{(Ibx + Ia)} + 1) - Ia \log(1/2(2ce^{(Ibx + Ia)} + I\sqrt{-4Ic})/c) - Ia \log(1/2(2ce^{(Ibx + Ia)} - I\sqrt{-4Ic})/c) + \operatorname{dilog}(1/2\sqrt{-4Ic}e^{(Ibx + Ia)}) + \operatorname{dilog}(-1/2\sqrt{-4Ic}e^{(Ibx + Ia)})/b$$

Sympy [F(-2)]

Exception generated.

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(atan(c+(-1+I*c)*tan(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) + 1 of type <class 'sympy.core.add.Add'> to QQ_I[b, _t0, exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(61) = 122$.

Time = 0.20 (sec) , antiderivative size = 448, normalized size of antiderivative = 5.21

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = \text{Too large to display}$$

input `integrate(arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="maxima")`

output

```
-1/8*((I*c - 1)*(4*I*(b*x + a)*log(-2*(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) - I)/(2*I*c^2 - 2*(c^2 + 2*I*c - 1)*tan(b*x + a) - 4*c - 2*I))/(I*c - 1) + I*(4*(b*x + a)*(log(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) + 2*c + I) - log(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) - I)) + I*log(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) + 2*c + I)^2 - 2*I*log(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) - I)*log(1/2*(c + I)*tan(b*x + a) - 1/2*I*c + 1/2) + 2*I*log(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) - I)*log(-1/2*((I*c - 1)*tan(b*x + a) + c - I)/c + 1) - 2*I*log(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) + 2*c + I)*log(-1/2*I*tan(b*x + a) + 1/2) - 2*I*dilog(-1/2*(c + I)*tan(b*x + a) + 1/2*I*c + 1/2) + 2*I*dilog(1/2*((I*c - 1)*tan(b*x + a) + c - I)/c) - 2*I*dilog(1/2*I*tan(b*x + a) + 1/2))/(I*c - 1)) - 8*(b*x + a)*arctan((I*c - 1)*tan(b*x + a) + c) + 4*(-I*b*x - I*a)*log(-2*(-I*c^2 + (c^2 + 2*I*c - 1)*tan(b*x + a) - I)/(2*I*c^2 - 2*(c^2 + 2*I*c - 1)*tan(b*x + a) - 4*c - 2*I)))/b
```

Giac [F]

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int \arctan((ic - 1) \tan(bx + a) + c) dx$$

input

```
integrate(arctan(c+(-1+I*c)*tan(b*x+a)),x, algorithm="giac")
```

output

```
integrate(arctan((I*c - 1)*tan(b*x + a) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(c + \tan(a + bx) (-1 + c1i)) dx$$

input

```
int(atan(c + tan(a + b*x)*(c*1i - 1)),x)
```

output

```
int(atan(c + tan(a + b*x)*(c*1i - 1)), x)
```

Reduce [F]

$$\int \arctan(c + (-1 + ic) \tan(a + bx)) dx = \int \operatorname{atan}(\tan(bx + a) ci - \tan(bx + a) + c) dx$$

input `int(atan(c+(-1+I*c)*tan(b*x+a)),x)`

output `int(atan(tan(a + b*x)*c*i - tan(a + b*x) + c),x)`

3.59 $\int \frac{\arctan(c+(-1+ic)\tan(a+bx))}{x} dx$

Optimal result	472
Mathematica [N/A]	472
Rubi [N/A]	473
Maple [N/A]	473
Fricas [N/A]	474
Sympy [F(-1)]	474
Maxima [F(-2)]	474
Giac [N/A]	475
Mupad [N/A]	475
Reduce [N/A]	476

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\arctan(c + (-1 + ic)\tan(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + (-1 + ic)\tan(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+(-1+I*c)*tan(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 0.73 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(c + (-1 + ic)\tan(a + bx))}{x} dx = \int \frac{\arctan(c + (-1 + ic)\tan(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + (-1 + I*c)*Tan[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + (-1 + I*c)*Tan[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx$$

input `Int[ArcTan[c + (-1 + I*c)*Tan[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\arctan(c + (ic - 1) \tan(bx + a))}{x} dx$$

input `int(arctan(c+(-1+I*c)*tan(b*x+a))/x,x)`

output `int(arctan(c+(-1+I*c)*tan(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.10 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx = \int \frac{\arctan((ic - 1) \tan(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+(-1+I*c)*tan(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x + 2*I*a) - I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+(-1+I*c)*tan(b*x+a))/x,x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx = \text{Exception raised: ValueError}$$

input `integrate(arctan(c+(-1+I*c)*tan(b*x+a))/x,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(c-1>0)', see `assume?` for more
details)Is
```

Giac [N/A]

Not integrable

Time = 0.86 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx = \int \frac{\arctan((ic - 1) \tan(bx + a) + c)}{x} dx$$

input

```
integrate(arctan(c+(-1+I*c)*tan(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctan((I*c - 1)*tan(b*x + a) + c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.85 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + \tan(a + bx) (-1 + c1i))}{x} dx$$

input

```
int(atan(c + tan(a + b*x)*(c*1i - 1))/x,x)
```

output

```
int(atan(c + tan(a + b*x)*(c*1i - 1))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24

$$\int \frac{\arctan(c + (-1 + ic) \tan(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\tan(bx + a) ci - \tan(bx + a) + c)}{x} dx$$

input `int(atan(c+(-1+I*c)*tan(b*x+a))/x,x)`output `int(atan(tan(a + b*x)*c*i - tan(a + b*x) + c)/x,x)`

3.60 $\int \arctan(\cot(a + bx)) dx$

Optimal result	477
Mathematica [A] (verified)	477
Rubi [A] (verified)	478
Maple [A] (verified)	479
Fricas [A] (verification not implemented)	479
Sympy [A] (verification not implemented)	480
Maxima [A] (verification not implemented)	480
Giac [A] (verification not implemented)	480
Mupad [B] (verification not implemented)	481
Reduce [B] (verification not implemented)	481

Optimal result

Integrand size = 7, antiderivative size = 16

$$\int \arctan(\cot(a + bx)) dx = -\frac{\arctan(\cot(a + bx))^2}{2b}$$

output

```
-1/2*(1/2*Pi-arccot(cot(b*x+a)))^2/b
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \arctan(\cot(a + bx)) dx = \frac{bx^2}{2} + x \arctan(\cot(a + bx))$$

input

```
Integrate[ArcTan[Cot[a + b*x]],x]
```

output

```
(b*x^2)/2 + x*ArcTan[Cot[a + b*x]]
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2588, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(\cot(a + bx)) dx$$

$$\downarrow \text{2588}$$

$$\frac{\int \arctan(\cot(a + bx)) d \arctan(\cot(a + bx))}{b}$$

$$\downarrow \text{15}$$

$$\frac{\arctan(\cot(a + bx))^2}{2b}$$

input `Int[ArcTan[Cot[a + b*x]],x]`

output `-1/2*ArcTan[Cot[a + b*x]]^2/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2588 `Int[(u_)^(m_.), x_Symbol] := With[{c = Simplify[D[u, x]]}, Simp[1/c Subst[Int[x^m, x], x, u], x] /; FreeQ[m, x] && PiecewiseLinearQ[u, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

method	result
parallelsch	$\frac{bx^2}{2} - x \operatorname{arccot}(\cot(bx+a)) + \frac{\pi x}{2}$
derivativedivides	$\frac{-\pi(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))) - \operatorname{arccot}(\cot(bx+a))^2}{2b}$
default	$\frac{\pi x}{2} - \frac{-(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))) \operatorname{arccot}(\cot(bx+a)) - \frac{(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a)))^2}{2}}{b}$
parts	$\frac{\pi x}{2} - \frac{-(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a))) \operatorname{arccot}(\cot(bx+a)) - \frac{(\frac{\pi}{2} - \operatorname{arccot}(\cot(bx+a)))^2}{2}}{b}$
risch	$\frac{\pi x}{2} + ix \ln(e^{i(bx+a)}) + \frac{\pi x \operatorname{csgn}(ie^{i(bx+a)})^2 \operatorname{csgn}(ie^{2i(bx+a)})}{4} - \frac{\pi x \operatorname{csgn}(ie^{i(bx+a)}) \operatorname{csgn}(ie^{2i(bx+a)})^2}{2} + \pi$

input `int(1/2*Pi-arccot(cot(b*x+a)),x,method=_RETURNVERBOSE)`output `1/2*b*x^2-x*arccot(cot(b*x+a))+1/2*Pi*x`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \arctan(\cot(a+bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}(\pi - 2a)x$$

input `integrate(1/2*pi-arccot(cot(b*x+a)),x, algorithm="fricas")`output `-1/2*b*x^2 + 1/2*(pi - 2*a)*x`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

$$\int \arctan(\cot(a + bx)) dx = \frac{\pi x}{2} - \begin{cases} \frac{\operatorname{acot}^2(\cot(a+bx))}{2b} & \text{for } b \neq 0 \\ x \operatorname{acot}(\cot(a)) & \text{otherwise} \end{cases}$$

input `integrate(1/2*pi-acot(cot(b*x+a)),x)`output `pi*x/2 - Piecewise((acot(cot(a + b*x))**2/(2*b), Ne(b, 0)), (x*acot(cot(a)), True))`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \arctan(\cot(a + bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}\pi x - ax$$

input `integrate(1/2*pi-arccot(cot(b*x+a)),x, algorithm="maxima")`output `-1/2*b*x^2 + 1/2*pi*x - a*x`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \arctan(\cot(a + bx)) dx = -\frac{1}{2}bx^2 + \frac{1}{2}\pi x - ax$$

input `integrate(1/2*pi-arccot(cot(b*x+a)),x, algorithm="giac")`output `-1/2*b*x^2 + 1/2*pi*x - a*x`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \arctan(\cot(a + bx)) dx = \frac{\Pi x}{2} - x \operatorname{acot}(\cot(a + bx)) + \frac{b x^2}{2}$$

input `int(Pi/2 - acot(cot(a + b*x)),x)`output `(Pi*x)/2 - x*acot(cot(a + b*x)) + (b*x^2)/2`**Reduce [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \arctan(\cot(a + bx)) dx = \frac{-\operatorname{acot}(\cot(bx + a))^2 + b\pi x}{2b}$$

input `int(1/2*Pi-acot(cot(b*x+a)),x)`output `(- acot(cot(a + b*x))**2 + b*pi*x)/(2*b)`

3.61 $\int x^2 \arctan(c + d \cot(a + bx)) dx$

Optimal result	482
Mathematica [A] (verified)	483
Rubi [A] (verified)	484
Maple [C] (warning: unable to verify)	490
Fricas [B] (verification not implemented)	490
Sympy [F(-1)]	491
Maxima [F]	492
Giac [F]	492
Mupad [F(-1)]	493
Reduce [F]	493

Optimal result

Integrand size = 15, antiderivative size = 399

$$\begin{aligned}
 \int x^2 \arctan(c + d \cot(a + bx)) dx &= \frac{1}{3}x^3 \arctan(c + d \cot(a + bx)) \\
 &+ \frac{1}{6}ix^3 \log\left(1 - \frac{(1 + ic - d)e^{2ia+2ibx}}{1 + ic + d}\right) \\
 &- \frac{1}{6}ix^3 \log\left(1 - \frac{(c + i(1 + d))e^{2ia+2ibx}}{c + i(1 - d)}\right) \\
 &+ \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(1+ic-d)e^{2ia+2ibx}}{1+ic+d}\right)}{4b} \\
 &- \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(c+i(1+d))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b} \\
 &+ \frac{ix \operatorname{PolyLog}\left(3, \frac{(1+ic-d)e^{2ia+2ibx}}{1+ic+d}\right)}{4b^2} \\
 &- \frac{ix \operatorname{PolyLog}\left(3, \frac{(c+i(1+d))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b^2} \\
 &- \frac{\operatorname{PolyLog}\left(4, \frac{(1+ic-d)e^{2ia+2ibx}}{1+ic+d}\right)}{8b^3} \\
 &+ \frac{\operatorname{PolyLog}\left(4, \frac{(c+i(1+d))e^{2ia+2ibx}}{c+i(1-d)}\right)}{8b^3}
 \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{3}x^3 \arctan(c+d \cot(bx+a)) + \frac{1}{6}I x^3 \ln(1 - (1+Ic-d) \exp(2Ia+2Ibx)) \\ & / (1+Ic+d) - \frac{1}{6}I x^3 \ln(1 - (c+I(1+d)) \exp(2Ia+2Ibx)/(c+I(1-d))) + \frac{1}{4} \\ & * x^2 \operatorname{polylog}(2, (1+Ic-d) \exp(2Ia+2Ibx)/(1+Ic+d))/b - \frac{1}{4}x^2 \operatorname{polylog}(2, \\ & (c+I(1+d)) \exp(2Ia+2Ibx)/(c+I(1-d)))/b + \frac{1}{4}I x \operatorname{polylog}(3, (1+Ic-d) \\ & * \exp(2Ia+2Ibx)/(1+Ic+d))/b^2 - \frac{1}{4}I x \operatorname{polylog}(3, (c+I(1+d)) \exp(2Ia \\ & + 2Ibx)/(c+I(1-d)))/b^2 - \frac{1}{8} \operatorname{polylog}(4, (1+Ic-d) \exp(2Ia+2Ibx)/(1+I \\ & * c+d))/b^3 + \frac{1}{8} \operatorname{polylog}(4, (c+I(1+d)) \exp(2Ia+2Ibx)/(c+I(1-d)))/b^3 \end{aligned}$$
Mathematica [A] (verified)

Time = 2.65 (sec) , antiderivative size = 360, normalized size of antiderivative = 0.90

$$\int x^2 \arctan(c + d \cot(a + bx)) dx$$

$$= \frac{8b^3 x^3 \arctan(c + d \cot(a + bx)) + 4ib^3 x^3 \log\left(1 + \frac{(-c+i(1+d))e^{-2i(a+bx)}}{c+i(-1+d)}\right) - 4ib^3 x^3 \log\left(1 + \frac{(-c+i(-1+d))e^{-2i(a+bx)}}{c+i(1+d)}\right)}{24b^3}$$

input

`Integrate[x^2*ArcTan[c + d*Cot[a + b*x]],x]`

output

$$\begin{aligned} & (8b^3 x^3 \operatorname{ArcTan}[c + d \operatorname{Cot}[a + b x]] + (4I) b^3 x^3 \operatorname{Log}[1 + (-c + I(1 + \\ & d))/((c + I(-1 + d))E^{((2I)(a + b x))})] - (4I) b^3 x^3 \operatorname{Log}[1 + (-c + \\ & I(-1 + d))/((c + I(1 + d))E^{((2I)(a + b x))})] - 6b^2 x^2 \operatorname{PolyLog}[2, \\ & (c - I(1 + d))/((c + I(-1 + d))E^{((2I)(a + b x))})] + 6b^2 x^2 \operatorname{PolyLog}[2, \\ & (I + c - Id)/((c + I(1 + d))E^{((2I)(a + b x))})] + (6I) b x \operatorname{PolyLog}[3, \\ & (c - I(1 + d))/((c + I(-1 + d))E^{((2I)(a + b x))})] - (6I) b x \operatorname{PolyLog}[3, \\ & (I + c - Id)/((c + I(1 + d))E^{((2I)(a + b x))})] + 3 \operatorname{PolyLog}[4, \\ & (c - I(1 + d))/((c + I(-1 + d))E^{((2I)(a + b x))})] - 3 \operatorname{PolyLog}[4, \\ & (I + c - Id)/((c + I(1 + d))E^{((2I)(a + b x))})])]/(24b^3) \end{aligned}$$

Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 515, normalized size of antiderivative = 1.29, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5700, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(d \cot(a + bx) + c) dx \\
 & \quad \downarrow \text{5700} \\
 & \frac{1}{3}b(ic - d + 1) \int \frac{e^{2ia+2ibx} x^3}{ic - (ic - d + 1)e^{2ia+2ibx} + d + 1} dx - \frac{1}{3}b(-ic + d + \\
 & 1) \int \frac{e^{2ia+2ibx} x^3}{-ic - (-ic + d + 1)e^{2ia+2ibx} - d + 1} dx + \frac{1}{3}x^3 \arctan(d \cot(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & \frac{1}{3}b(ic - d + 1) \left(\frac{x^3 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c - i(1-d))} - \frac{3 \int x^2 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) dx}{2b(c - i(1-d))} \right) - \\
 & \frac{1}{3}b(-ic + d + 1) \left(\frac{3 \int x^2 \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) dx}{2b(c + i(d+1))} - \frac{x^3 \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c + i(d+1))} \right) + \\
 & \quad \frac{1}{3}x^3 \arctan(d \cot(a + bx) + c) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$\begin{aligned}
 & 1) \left(\frac{x^3 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c-i(1-d))} - \frac{\frac{\frac{1}{3}b(ic-d+1)}{3} \left(\frac{ix^2 \operatorname{PolyLog} \left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b} - \frac{i \int x \operatorname{PolyLog} \left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) dx}{b} \right)}{2b(c-i(1-d))} \right) - \\
 & 1) \left(\frac{\frac{\frac{1}{3}b(-ic+d+1)}{3} \left(\frac{ix^2 \operatorname{PolyLog} \left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b} - \frac{i \int x \operatorname{PolyLog} \left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) dx}{b} \right)}{2b(c+i(d+1))} - \frac{x^3 \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c+i(d+1))} \right) + \\
 & \frac{1}{3}x^3 \arctan(d \cot(a+bx) + c)
 \end{aligned}$$

↓ 7163

$$1) \left(\frac{x^3 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c-i(1-d))} - \frac{\frac{1}{3}b(ic-d+1) \left(\frac{ix^2 \operatorname{PolyLog} \left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b} - \frac{i \left(\frac{i \int \operatorname{PolyLog} \left(3, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) dx}{2b} - \frac{ix \operatorname{PolyLog} \left(3, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{b} \right)}{b} \right)}{2b(c-i(1-d))} \right)$$

$$1) \left(\frac{\frac{1}{3}b(-ic+d+1) \left(\frac{ix^2 \operatorname{PolyLog} \left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b} - \frac{i \left(\frac{i \int \operatorname{PolyLog} \left(3, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) dx}{2b} - \frac{ix \operatorname{PolyLog} \left(3, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b} \right)}{b} \right)}{2b(c+i(d+1))} - \frac{x^3 \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c+i(d+1))} \right)$$

$$\frac{1}{3}x^3 \arctan(d \cot(a + bx) + c)$$

↓ 2720

$$1) \left(\frac{x^3 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c-i(1-d))} - \frac{\frac{1}{3}b(ic-d+1) \left(\frac{ix^2 \operatorname{PolyLog} \left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b} - i \left(\frac{\int e^{-2ia-2ibx} \operatorname{PolyLog} \left(3, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) de^{2ia+2ibx}}{4b^2} \right) \right)}{2b(c-i(1-d))} \right)$$

$$1) \left(\frac{\frac{1}{3}b(-ic+d+1) \left(\frac{ix^2 \operatorname{PolyLog} \left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b} - i \left(\frac{\int e^{-2ia-2ibx} \operatorname{PolyLog} \left(3, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) de^{2ia+2ibx}}{4b^2} \right) \right) - ix \operatorname{PolyLog} \left(3, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c+i(d+1))} \right)$$

$$\frac{1}{3}x^3 \arctan(d \cot(a+bx) + c)$$

↓ 7143

$$\begin{aligned}
 & \frac{1}{3}x^3 \arctan(d \cot(a + bx) + c) + \frac{1}{3}b(ic - d + \\
 1) & \left(\frac{x^3 \log\left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{2b(c-i(1-d))} - \frac{3 \left(\frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{2b} - i \left(\frac{\operatorname{PolyLog}\left(4, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{b} \right)}{2b(c-i(1-d))} \right)}{2b(c-i(1-d))} \right) \\
 1) & \left(\frac{3 \left(\frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b} - i \left(\frac{\operatorname{PolyLog}\left(4, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b^2} - \frac{ix \operatorname{PolyLog}\left(3, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b} \right)}{b} \right)}{2b(c+i(d+1))} - \frac{\frac{1}{3}b(-ic+d+x^3 \log\left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right))}{2} \right)
 \end{aligned}$$

input `Int[x^2*ArcTan[c + d*Cot[a + b*x]],x]`

output `(x^3*ArcTan[c + d*Cot[a + b*x]])/3 + (b*(1 + I*c - d)*((x^3*Log[1 - ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)]/(2*b*(c - I*(1 - d))) - (3*(((I/2)*x^2*PolyLog[2, ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)])/b - (I*(((1/2)*x*PolyLog[3, ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)])/b + PolyLog[4, ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)]/(4*b^2))/b)/(2*b*(c - I*(1 - d)))))/3 - (b*(1 - I*c + d)*(-1/2*(x^3*Log[1 - ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d))])/(b*(c + I*(1 + d))) + (3*(((I/2)*x^2*PolyLog[2, ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d)))]/b - (I*(((1/2)*x*PolyLog[3, ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d)))]/b + PolyLog[4, ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d))]/(4*b^2))/b)/(2*b*(c + I*(1 + d)))))/3`

Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5700

```
Int[ArcTan[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Cot[a + b*x]]/(f*(m +
1))), x] + (Simp[b*((1 + I*c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(
2*I*a + 2*I*b*x)/(1 + I*c + d - (1 + I*c - d)*E^(2*I*a + 2*I*b*x))), x], x]
- Simp[b*((1 - I*c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*I*a + 2
*I*b*x)/(1 - I*c - d - (1 - I*c + d)*E^(2*I*a + 2*I*b*x))), x], x]) /; Free
Q[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c - I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_)^(m_.))*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 60.35 (sec) , antiderivative size = 7869, normalized size of antiderivative = 19.72

method	result	size
risch	Expression too large to display	7869

input

```
int(x^2*arctan(c+d*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1589 vs. $2(283) = 566$.

Time = 0.27 (sec) , antiderivative size = 1589, normalized size of antiderivative = 3.98

$$\int x^2 \arctan(c + d \cot(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(x^2*arctan(c+d*cot(b*x+a)),x, algorithm="fricas")
```

output

```

1/48*(16*b^3*x^3*arctan(d*cot(b*x + a) + c) + 6*b^2*x^2*dilog(-(c^2 + d^2
- (c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-I*c^2 + 2*c*d + I*d^2 - I
)*sin(2*b*x + 2*a) + 2*d + 1)/(c^2 + d^2 + 2*d + 1) + 1) + 6*b^2*x^2*dilog
(-(c^2 + d^2 - (c^2 - 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*c*d
- I*d^2 + I)*sin(2*b*x + 2*a) + 2*d + 1)/(c^2 + d^2 + 2*d + 1) + 1) - 6*b
^2*x^2*dilog(-(c^2 + d^2 - (c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-
I*c^2 + 2*c*d + I*d^2 - I)*sin(2*b*x + 2*a) - 2*d + 1)/(c^2 + d^2 - 2*d +
1) + 1) - 6*b^2*x^2*dilog(-(c^2 + d^2 - (c^2 - 2*I*c*d - d^2 + 1)*cos(2*b*
x + 2*a) + (I*c^2 + 2*c*d - I*d^2 + I)*sin(2*b*x + 2*a) - 2*d + 1)/(c^2 +
d^2 - 2*d + 1) + 1) - 4*I*a^3*log(1/2*c^2 + I*c*d - 1/2*d^2 - 1/2*(c^2 + d
^2 + 2*d + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*d + I)*sin(2*b*x
+ 2*a) + 1/2) + 4*I*a^3*log(1/2*c^2 + I*c*d - 1/2*d^2 - 1/2*(c^2 + d^2 -
2*d + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*d + I)*sin(2*b*x + 2
*a) + 1/2) + 4*I*a^3*log(-1/2*c^2 + I*c*d + 1/2*d^2 + 1/2*(c^2 + d^2 + 2*d
+ 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*d + I)*sin(2*b*x + 2*a) -
1/2) - 4*I*a^3*log(-1/2*c^2 + I*c*d + 1/2*d^2 + 1/2*(c^2 + d^2 - 2*d + 1)
*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*d + I)*sin(2*b*x + 2*a) - 1/2
) + 6*I*b*x*polylog(3, ((c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^
2 - 2*c*d - I*d^2 + I)*sin(2*b*x + 2*a))/(c^2 + d^2 + 2*d + 1)) - 6*I*b*x*
polylog(3, ((c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2 - 2*c*d...

```

Sympy [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \cot(a + bx)) dx = \text{Timed out}$$

input

```
integrate(x**2*atan(c+d*cot(b*x+a)),x)
```

output

Timed out

Maxima [F]

$$\int x^2 \arctan(c + d \cot(a + bx)) dx = \int x^2 \arctan(d \cot(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*cot(b*x+a)),x, algorithm="maxima")`

output

```
-1/6*x^3*arctan2(-c*cos(2*b*x + 2*a) + (d + 1)*sin(2*b*x + 2*a) + c, (d +
1)*cos(2*b*x + 2*a) + c*sin(2*b*x + 2*a) + d - 1) - 1/6*x^3*arctan2(-c*cos
(2*b*x + 2*a) + (d - 1)*sin(2*b*x + 2*a) + c, -(d - 1)*cos(2*b*x + 2*a) -
c*sin(2*b*x + 2*a) - d - 1) + 4*b*d*integrate(1/3*(2*(c^2 + d^2 + 1)*x^3*c
os(2*b*x + 2*a)^2 + 2*c*d*x^3*sin(2*b*x + 2*a) + 2*(c^2 + d^2 + 1)*x^3*sin
(2*b*x + 2*a)^2 - (c^2 - d^2 + 1)*x^3*cos(2*b*x + 2*a) - (2*c*d*x^3*sin(2*
b*x + 2*a) + (c^2 - d^2 + 1)*x^3*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c
*d*x^3*cos(2*b*x + 2*a) - (c^2 - d^2 + 1)*x^3*sin(2*b*x + 2*a))*sin(4*b*x
+ 4*a))/(c^4 + d^4 + 2*(c^2 - 1)*d^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^
2 + 1)*cos(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*co
s(2*b*x + 2*a)^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2 + 1)*sin(4*b*x + 4
*a)^2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*sin(2*b*x + 2*a)^2 + 2
*c^2 + 2*(c^4 + d^4 - 2*(3*c^2 + 1)*d^2 + 2*c^2 - 2*(c^4 - d^4 + 2*c^2 + 1
)*cos(2*b*x + 2*a) - 4*(c*d^3 + (c^3 + c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b
*x + 4*a) - 4*(c^4 - d^4 + 2*c^2 + 1)*cos(2*b*x + 2*a) + 4*(2*c*d^3 - 2*(c
^3 + c)*d + 2*(c*d^3 + (c^3 + c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 + 2*c^2
+ 1)*sin(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 + c)*d)*sin(2*b*
x + 2*a) + 1), x)
```

Giac [F]

$$\int x^2 \arctan(c + d \cot(a + bx)) dx = \int x^2 \arctan(d \cot(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan(d*cot(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \cot(a + bx)) dx = \int x^2 \operatorname{atan}(c + d \cot(a + bx)) dx$$

input `int(x^2*atan(c + d*cot(a + b*x)),x)`output `int(x^2*atan(c + d*cot(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \arctan(c + d \cot(a + bx)) dx = \int \operatorname{atan}(\cot(bx + a)d + c) x^2 dx$$

input `int(x^2*atan(c+d*cot(b*x+a)),x)`output `int(atan(cot(a + b*x)*d + c)*x**2,x)`

3.62 $\int x \arctan(c + d \cot(a + bx)) dx$

Optimal result	494
Mathematica [A] (verified)	495
Rubi [A] (verified)	495
Maple [C] (warning: unable to verify)	499
Fricas [B] (verification not implemented)	499
Sympy [F(-1)]	500
Maxima [F]	501
Giac [F]	501
Mupad [F(-1)]	502
Reduce [F]	502

Optimal result

Integrand size = 13, antiderivative size = 303

$$\begin{aligned}
 \int x \arctan(c + d \cot(a + bx)) dx &= \frac{1}{2}x^2 \arctan(c + d \cot(a + bx)) \\
 &+ \frac{1}{4}ix^2 \log\left(1 - \frac{(1 + ic - d)e^{2ia+2ibx}}{1 + ic + d}\right) \\
 &- \frac{1}{4}ix^2 \log\left(1 - \frac{(c + i(1 + d))e^{2ia+2ibx}}{c + i(1 - d)}\right) \\
 &+ \frac{x \operatorname{PolyLog}\left(2, \frac{(1+ic-d)e^{2ia+2ibx}}{1+ic+d}\right)}{4b} \\
 &- \frac{x \operatorname{PolyLog}\left(2, \frac{(c+i(1+d))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b} \\
 &+ \frac{i \operatorname{PolyLog}\left(3, \frac{(1+ic-d)e^{2ia+2ibx}}{1+ic+d}\right)}{8b^2} \\
 &- \frac{i \operatorname{PolyLog}\left(3, \frac{(c+i(1+d))e^{2ia+2ibx}}{c+i(1-d)}\right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & 1/2*x^2*\arctan(c+d*\cot(b*x+a))+1/4*I*x^2*\ln(1-(1+I*c-d)*\exp(2*I*a+2*I*b*x) \\ & /((1+I*c+d))-1/4*I*x^2*\ln(1-(c+I*(1+d))*\exp(2*I*a+2*I*b*x)/(c+I*(1-d)))+1/4 \\ & *x*\text{polylog}(2,(1+I*c-d)*\exp(2*I*a+2*I*b*x)/(1+I*c+d))/b-1/4*x*\text{polylog}(2,(c+ \\ & I*(1+d))*\exp(2*I*a+2*I*b*x)/(c+I*(1-d)))/b+1/8*I*\text{polylog}(3,(1+I*c-d)*\exp(2 \\ & *I*a+2*I*b*x)/(1+I*c+d))/b^2-1/8*I*\text{polylog}(3,(c+I*(1+d))*\exp(2*I*a+2*I*b*x) \\ &)/(c+I*(1-d)))/b^2 \end{aligned}$$
Mathematica [A] (verified)

Time = 2.11 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.91

$$\int x \arctan(c + d \cot(a + bx)) dx$$

$$= \frac{4b^2x^2 \arctan(c + d \cot(a + bx)) + 2ib^2x^2 \log\left(1 + \frac{(-c+i(1+d))e^{-2i(a+bx)}}{c+i(-1+d)}\right) - 2ib^2x^2 \log\left(1 + \frac{(-c+i(-1+d))e^{-2i(a+bx)}}{c+i(1+d)}\right)}{1}$$

input

`Integrate[x*ArcTan[c + d*Cot[a + b*x]],x]`

output

$$\begin{aligned} & (4*b^2*x^2*ArcTan[c + d*Cot[a + b*x]] + (2*I)*b^2*x^2*Log[1 + (-c + I*(1 + \\ & d))/((c + I*(-1 + d))*E^((2*I)*(a + b*x)))] - (2*I)*b^2*x^2*Log[1 + (-c + \\ & I*(-1 + d))/((c + I*(1 + d))*E^((2*I)*(a + b*x)))] - 2*b*x*PolyLog[2, (c \\ & - I*(1 + d))/((c + I*(-1 + d))*E^((2*I)*(a + b*x)))] + 2*b*x*PolyLog[2, (I \\ & + c - I*d)/((c + I*(1 + d))*E^((2*I)*(a + b*x)))] + I*PolyLog[3, (c - I*(\\ & 1 + d))/((c + I*(-1 + d))*E^((2*I)*(a + b*x)))] - I*PolyLog[3, (I + c - I* \\ & d)/((c + I*(1 + d))*E^((2*I)*(a + b*x)))])/(8*b^2) \end{aligned}$$
Rubi [A] (verified)Time = 1.13 (sec) , antiderivative size = 394, normalized size of antiderivative = 1.30, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {5700, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(d \cot(a + bx) + c) dx$$

↓ 5700

$$\frac{1}{2}b(ic - d + 1) \int \frac{e^{2ia+2ibx} x^2}{ic - (ic - d + 1)e^{2ia+2ibx} + d + 1} dx - \frac{1}{2}b(-ic + d + 1) \int \frac{e^{2ia+2ibx} x^2}{-ic - (-ic + d + 1)e^{2ia+2ibx} - d + 1} dx + \frac{1}{2}x^2 \arctan(d \cot(a + bx) + c)$$

↓ 2620

$$\frac{1}{2}b(ic - d + 1) \left(\frac{x^2 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c-i(1-d))} - \frac{\int x \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) dx}{b(c-i(1-d))} \right) - \frac{1}{2}b(-ic + d + 1) \left(\frac{\int x \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) dx}{b(c+i(d+1))} - \frac{x^2 \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c+i(d+1))} \right) + \frac{1}{2}x^2 \arctan(d \cot(a + bx) + c)$$

↓ 3011

$$\frac{1}{2}b(ic - d + 1) \left(\frac{x^2 \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c-i(1-d))} - \frac{ix \operatorname{PolyLog} \left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) dx}{2b} \right) - \frac{1}{2}b(-ic + d + 1) \left(\frac{ix \operatorname{PolyLog} \left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b} - \frac{i \int \operatorname{PolyLog} \left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) dx}{2b} - \frac{x^2 \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c+i(d+1))} \right) + \frac{1}{2}x^2 \arctan(d \cot(a + bx) + c)$$

↓ 2720

$$\begin{aligned}
 & \frac{1}{2}b(ic - d + \\
 1) & \left(\frac{x^2 \log\left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{2b(c-i(1-d))} - \frac{ix \operatorname{PolyLog}\left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right) de^{2ia+2ibx}}{4b^2} \right) \\
 & \frac{1}{2}b(-ic + d + \\
 1) & \left(\frac{ix \operatorname{PolyLog}\left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}\left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right) de^{2ia+2ibx}}{4b^2} - \frac{x^2 \log\left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b(c+i(d+1))} \right) \\
 & \frac{1}{2}x^2 \arctan(d \cot(a + bx) + c)
 \end{aligned}$$

7143

$$\begin{aligned}
 & \frac{1}{2}x^2 \arctan(d \cot(a + bx) + c) + \frac{1}{2}b(ic - d + \\
 1) & \left(\frac{x^2 \log\left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{2b(c-i(1-d))} - \frac{ix \operatorname{PolyLog}\left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{4b^2} \right) - \\
 & \frac{1}{2}b(-ic + d + \\
 1) & \left(\frac{ix \operatorname{PolyLog}\left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b} - \frac{\operatorname{PolyLog}\left(3, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b^2} - \frac{x^2 \log\left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b(c+i(d+1))} \right)
 \end{aligned}$$

```
input Int[x*ArcTan[c + d*Cot[a + b*x]],x]
```

```
output (x^2*ArcTan[c + d*Cot[a + b*x]])/2 + (b*(1 + I*c - d)*((x^2*Log[1 - ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)]/(2*b*(c - I*(1 - d))) - (((I/2)*x*PolyLog[2, ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)])/b - PolyLog[3, ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)]/(4*b^2))/(b*(c - I*(1 - d))))/2 - (b*(1 - I*c + d)*(-1/2*(x^2*Log[1 - ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d))]/(b*(c + I*(1 + d))) + (((I/2)*x*PolyLog[2, ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d))])/b - PolyLog[3, ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x))/(c + I*(1 - d))]/(4*b^2))/(b*(c + I*(1 + d))))/2
```

Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5700

```
Int[ArcTan[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Cot[a + b*x]]/(f*(m +
1))), x] + (Simp[b*((1 + I*c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(
2*I*a + 2*I*b*x)/(1 + I*c + d - (1 + I*c - d)*E^(2*I*a + 2*I*b*x))), x], x]
- Simp[b*((1 - I*c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*I*a + 2
*I*b*x)/(1 - I*c - d - (1 - I*c + d)*E^(2*I*a + 2*I*b*x))), x], x]) /; Free
Q[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c - I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.21 (sec) , antiderivative size = 7501, normalized size of antiderivative = 24.76

method	result	size
risch	Expression too large to display	7501

input `int(x*arctan(c+d*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1289 vs. $2(213) = 426$.

Time = 0.24 (sec) , antiderivative size = 1289, normalized size of antiderivative = 4.25

$$\int x \arctan(c + d \cot(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctan(c+d*cot(b*x+a)),x, algorithm="fricas")`

output

```

1/16*(8*b^2*x^2*arctan(d*cot(b*x + a) + c) + 2*b*x*dilog(-(c^2 + d^2 - (c^
2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-I*c^2 + 2*c*d + I*d^2 - I)*sin
(2*b*x + 2*a) + 2*d + 1)/(c^2 + d^2 + 2*d + 1) + 1) + 2*b*x*dilog(-(c^2 +
d^2 - (c^2 - 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*c*d - I*d^2
+ I)*sin(2*b*x + 2*a) + 2*d + 1)/(c^2 + d^2 + 2*d + 1) + 1) - 2*b*x*dilog(
-(c^2 + d^2 - (c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-I*c^2 + 2*c*d
+ I*d^2 - I)*sin(2*b*x + 2*a) - 2*d + 1)/(c^2 + d^2 - 2*d + 1) + 1) - 2*b
*x*dilog(-(c^2 + d^2 - (c^2 - 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2
+ 2*c*d - I*d^2 + I)*sin(2*b*x + 2*a) - 2*d + 1)/(c^2 + d^2 - 2*d + 1) +
1) + 2*I*a^2*log(1/2*c^2 + I*c*d - 1/2*d^2 - 1/2*(c^2 + d^2 + 2*d + 1)*cos
(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*d + I)*sin(2*b*x + 2*a) + 1/2) -
2*I*a^2*log(1/2*c^2 + I*c*d - 1/2*d^2 - 1/2*(c^2 + d^2 - 2*d + 1)*cos(2*b*
x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*d + I)*sin(2*b*x + 2*a) + 1/2) - 2*I*a
^2*log(-1/2*c^2 + I*c*d + 1/2*d^2 + 1/2*(c^2 + d^2 + 2*d + 1)*cos(2*b*x +
2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*d + I)*sin(2*b*x + 2*a) - 1/2) + 2*I*a^2*1
og(-1/2*c^2 + I*c*d + 1/2*d^2 + 1/2*(c^2 + d^2 - 2*d + 1)*cos(2*b*x + 2*a)
+ 1/2*(I*c^2 + I*d^2 - 2*I*d + I)*sin(2*b*x + 2*a) - 1/2) - 2*(-I*b^2*x^2
+ I*a^2)*log((c^2 + d^2 - (c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-
I*c^2 + 2*c*d + I*d^2 - I)*sin(2*b*x + 2*a) + 2*d + 1)/(c^2 + d^2 + 2*d +
1)) - 2*(I*b^2*x^2 - I*a^2)*log((c^2 + d^2 - (c^2 - 2*I*c*d - d^2 + 1)*...

```

Sympy [F(-1)]

Timed out.

$$\int x \arctan(c + d \cot(a + bx)) dx = \text{Timed out}$$

input

```
integrate(x*atan(c+d*cot(b*x+a)),x)
```

output

Timed out

Maxima [F]

$$\int x \arctan(c + d \cot(a + bx)) dx = \int x \arctan(d \cot(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*cot(b*x+a)),x, algorithm="maxima")`

output

```
-1/4*x^2*arctan2(-c*cos(2*b*x + 2*a) + (d + 1)*sin(2*b*x + 2*a) + c, (d +
1)*cos(2*b*x + 2*a) + c*sin(2*b*x + 2*a) + d - 1) - 1/4*x^2*arctan2(-c*cos
(2*b*x + 2*a) + (d - 1)*sin(2*b*x + 2*a) + c, -(d - 1)*cos(2*b*x + 2*a) -
c*sin(2*b*x + 2*a) - d - 1) + 2*b*d*integrate((2*(c^2 + d^2 + 1)*x^2*cos(2
*b*x + 2*a)^2 + 2*c*d*x^2*sin(2*b*x + 2*a) + 2*(c^2 + d^2 + 1)*x^2*sin(2*b
*x + 2*a)^2 - (c^2 - d^2 + 1)*x^2*cos(2*b*x + 2*a) - (2*c*d*x^2*sin(2*b*x
+ 2*a) + (c^2 - d^2 + 1)*x^2*cos(2*b*x + 2*a))*cos(4*b*x + 4*a) + (2*c*d*x
^2*cos(2*b*x + 2*a) - (c^2 - d^2 + 1)*x^2*sin(2*b*x + 2*a))*sin(4*b*x + 4*
a))/(c^4 + d^4 + 2*(c^2 - 1)*d^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2 +
1)*cos(4*b*x + 4*a)^2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*cos(2*
b*x + 2*a)^2 + (c^4 + d^4 + 2*(c^2 - 1)*d^2 + 2*c^2 + 1)*sin(4*b*x + 4*a)^
2 + 4*(c^4 + d^4 + 2*(c^2 + 1)*d^2 + 2*c^2 + 1)*sin(2*b*x + 2*a)^2 + 2*c^2
+ 2*(c^4 + d^4 - 2*(3*c^2 + 1)*d^2 + 2*c^2 - 2*(c^4 - d^4 + 2*c^2 + 1)*co
s(2*b*x + 2*a) - 4*(c*d^3 + (c^3 + c)*d)*sin(2*b*x + 2*a) + 1)*cos(4*b*x +
4*a) - 4*(c^4 - d^4 + 2*c^2 + 1)*cos(2*b*x + 2*a) + 4*(2*c*d^3 - 2*(c^3 +
c)*d + 2*(c*d^3 + (c^3 + c)*d)*cos(2*b*x + 2*a) - (c^4 - d^4 + 2*c^2 + 1)
*sin(2*b*x + 2*a))*sin(4*b*x + 4*a) + 8*(c*d^3 + (c^3 + c)*d)*sin(2*b*x +
2*a) + 1), x)
```

Giac [F]

$$\int x \arctan(c + d \cot(a + bx)) dx = \int x \arctan(d \cot(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan(d*cot(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + d \cot(a + bx)) dx = \int x \operatorname{atan}(c + d \cot(a + bx)) dx$$

input `int(x*atan(c + d*cot(a + b*x)),x)`output `int(x*atan(c + d*cot(a + b*x)), x)`**Reduce [F]**

$$\int x \arctan(c + d \cot(a + bx)) dx = \int \operatorname{atan}(\cot(bx + a) d + c) x dx$$

input `int(x*atan(c+d*cot(b*x+a)),x)`output `int(atan(cot(a + b*x)*d + c)*x,x)`

3.63 $\int \arctan(c + d \cot(a + bx)) dx$

Optimal result	503
Mathematica [B] (warning: unable to verify)	504
Rubi [A] (verified)	505
Maple [B] (verified)	507
Fricas [B] (verification not implemented)	508
Sympy [F(-1)]	509
Maxima [B] (verification not implemented)	510
Giac [F]	511
Mupad [F(-1)]	511
Reduce [F]	511

Optimal result

Integrand size = 11, antiderivative size = 198

$$\int \arctan(c + d \cot(a + bx)) dx = x \arctan(c + d \cot(a + bx)) + \frac{1}{2}ix \log\left(1 - \frac{(1 + ic - d)e^{2ia+2ibx}}{1 + ic + d}\right) - \frac{1}{2}ix \log\left(1 - \frac{(c + i(1 + d))e^{2ia+2ibx}}{c + i(1 - d)}\right) + \frac{\text{PolyLog}\left(2, \frac{(1+ic-d)e^{2ia+2ibx}}{1+ic+d}\right)}{4b} - \frac{\text{PolyLog}\left(2, \frac{(c+i(1+d))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b}$$

output

```
x*arctan(c+d*cot(b*x+a))+1/2*I*x*ln(1-(1+I*c-d)*exp(2*I*a+2*I*b*x)/(1+I*c+d))-1/2*I*x*ln(1-(c+I*(1+d))*exp(2*I*a+2*I*b*x)/(c+I*(1-d)))+1/4*polylog(2,(1+I*c-d)*exp(2*I*a+2*I*b*x)/(1+I*c+d))/b-1/4*polylog(2,(c+I*(1+d))*exp(2*I*a+2*I*b*x)/(c+I*(1-d)))/b
```


Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 702 vs. $2(198) = 396$.

Time = 9.77 (sec) , antiderivative size = 702, normalized size of antiderivative = 3.55

$$\int \arctan(c + d \cot(a + bx)) dx = x \left(\arctan(c + d \cot(a + bx)) \right. \\ \left. - \left(4a\sqrt{-d^2} \arctan\left(c + \frac{(1+c^2)\tan(a+bx)}{d}\right) + id \log(1 + i \tan(a + bx)) \log\left(\frac{-cd + \sqrt{-d^2} - (1+c^2)\tan(a+bx)}{-i - ic^2 - cd + \sqrt{-d^2}}\right) - id \right) \right)$$

input `Integrate[ArcTan[c + d*Cot[a + b*x]],x]`

output `x*(ArcTan[c + d*Cot[a + b*x]] - ((4*a*Sqrt[-d^2]*ArcTan[c + ((1 + c^2)*Tan[a + b*x])/d] + I*d*Log[1 + I*Tan[a + b*x]]*Log[(-(c*d) + Sqrt[-d^2] - (1 + c^2)*Tan[a + b*x])/(-I - I*c^2 - c*d + Sqrt[-d^2])] - I*d*Log[1 - I*Tan[a + b*x]]*Log[(-(c*d) + Sqrt[-d^2] - (1 + c^2)*Tan[a + b*x])/(I + I*c^2 - c*d + Sqrt[-d^2])] + I*d*Log[1 - I*Tan[a + b*x]]*Log[(c*d + Sqrt[-d^2] + (1 + c^2)*Tan[a + b*x])/(-I - I*c^2 + c*d + Sqrt[-d^2])] - I*d*Log[1 + I*Tan[a + b*x]]*Log[(c*d + Sqrt[-d^2] + (1 + c^2)*Tan[a + b*x])/(I + I*c^2 + c*d + Sqrt[-d^2])] - I*d*PolyLog[2, ((1 + c^2)*(1 - I*Tan[a + b*x])/(1 + c^2 + I*c*d - I*Sqrt[-d^2]))] + I*d*PolyLog[2, ((1 + c^2)*(1 - I*Tan[a + b*x])/(1 + c^2 + I*c*d + I*Sqrt[-d^2]))] - I*d*PolyLog[2, ((1 + c^2)*(1 + I*Tan[a + b*x])/(1 + c^2 - I*c*d - I*Sqrt[-d^2]))] + I*d*PolyLog[2, ((1 + c^2)*(1 + I*Tan[a + b*x])/(1 + c^2 - I*c*d + I*Sqrt[-d^2]))]*((-(c*d) + Sqrt[-d^2])*Cos[a + b*x] - (1 + c^2)*Sin[a + b*x])*((c*d + Sqrt[-d^2])*Cos[a + b*x] + (1 + c^2)*Sin[a + b*x]))/((1 + c^2)*Sqrt[-d^2]*(2*a - I*Log[1 - I*Tan[a + b*x]] + I*Log[1 + I*Tan[a + b*x]])*(1 + c^2 + d^2 + (-1 - c^2 + d^2)*Cos[2*(a + b*x)] + 2*c*d*Sin[2*(a + b*x)]))`

Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.40, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {5692, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(d \cot(a + bx) + c) dx \\
 & \quad \downarrow \text{5692} \\
 & b(ic - d + 1) \int \frac{e^{2ia+2ibx} x}{ic - (ic - d + 1)e^{2ia+2ibx} + d + 1} dx - b(-ic + d + \\
 & 1) \int \frac{e^{2ia+2ibx} x}{-ic - (-ic + d + 1)e^{2ia+2ibx} - d + 1} dx + x \arctan(d \cot(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & b(ic - d + 1) \left(\frac{x \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c - i(1-d))} - \frac{\int \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) dx}{2b(c - i(1-d))} \right) - b(-ic + d + \\
 & 1) \left(\frac{\int \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) dx}{2b(c + i(d+1))} - \frac{x \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c + i(d+1))} \right) + x \arctan(d \cot(a + \\
 & \quad bx) + c) \\
 & \quad \downarrow \text{2715} \\
 & 1) \left(\frac{b(ic - d +}{4b^2(c - i(1-d))} \int e^{-2ia-2ibx} \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right) de^{2ia+2ibx} + \frac{x \log \left(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1} \right)}{2b(c - i(1-d))} \right) - \\
 & 1) \left(\frac{b(-ic + d +}{4b^2(c + i(d+1))} \int e^{-2ia-2ibx} \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right) de^{2ia+2ibx} - \frac{x \log \left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)} \right)}{2b(c + i(d+1))} \right) + \\
 & \quad x \arctan(d \cot(a + bx) + c) \\
 & \quad \downarrow \text{2838}
 \end{aligned}$$

$$1) \left(\frac{x \arctan(d \cot(a + bx) + c) + b(ic - d + \frac{x \log(1 - \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}}{2b(c-i(1-d))})}{2b(c-i(1-d))} - \frac{i \operatorname{PolyLog}\left(2, \frac{(ic-d+1)e^{2ia+2ibx}}{ic+d+1}\right)}{4b^2(c-i(1-d))} \right) - b(-ic + d + 1) \left(\frac{i \operatorname{PolyLog}\left(2, \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{4b^2(c+i(d+1))} - \frac{x \log\left(1 - \frac{(c+i(d+1))e^{2ia+2ibx}}{c+i(1-d)}\right)}{2b(c+i(d+1))} \right)$$

input `Int[ArcTan[c + d*Cot[a + b*x]],x]`

output `x*ArcTan[c + d*Cot[a + b*x]] + b*(1 + I*c - d)*((x*Log[1 - ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)]/(2*b*(c - I*(1 - d)))) - ((I/4)*PolyLog[2, ((1 + I*c - d)*E^((2*I)*a + (2*I)*b*x))/(1 + I*c + d)]/(b^2*(c - I*(1 - d)))) - b*(1 - I*c + d)*(-1/2*(x*Log[1 - ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 - d)))]/(b*(c + I*(1 + d))) + ((I/4)*PolyLog[2, ((c + I*(1 + d))*E^((2*I)*a + (2*I)*b*x)]/(c + I*(1 - d)))]/(b^2*(c + I*(1 + d))))`

Defintions of rubi rules used

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5692

```
Int[ArcTan[(c_.) + Cot[(a_.) + (b_.)*(x_.)]*(d_.)], x_Symbol] := Simp[x*ArcTan[c + d*Cot[a + b*x]], x] + (Simp[b*(1 + I*c - d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 + I*c + d - (1 + I*c - d)*E^(2*I*a + 2*I*b*x))], x], x] - Simp[b*(1 - I*c + d) Int[x*(E^(2*I*a + 2*I*b*x))/(1 - I*c - d - (1 - I*c + d)*E^(2*I*a + 2*I*b*x))], x], x) /; FreeQ[{a, b, c, d}, x] && NeQ[(c + I*d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1144 vs. $2(168) = 336$.

Time = 4.13 (sec) , antiderivative size = 1145, normalized size of antiderivative = 5.78

method	result	size
derivativdivides	Expression too large to display	1145
default	Expression too large to display	1145
risch	Expression too large to display	4986

input

```
int(arctan(c+d*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/b/d*(-d*(1/2*Pi-arccot(cot(b*x+a)))*arctan(c+d*cot(b*x+a))+d^2*(-1/d*arc
tan(d*((c+d*cot(b*x+a))/d-c/d)+c)*arctan(-(c+d*cot(b*x+a))/d+c/d)-1/d^2*(-
1/2*I*d*arctan(d*((c+d*cot(b*x+a))/d-c/d)+c)*ln(1-(I*d+I+c)*(1+I*(d*((c+d*
cot(b*x+a))/d-c/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(-I*d+I-c))-
1/2*d*arctan(d*((c+d*cot(b*x+a))/d-c/d)+c)^2-1/4*d*polylog(2,(I*d+I+c)*(1+
I*(d*((c+d*cot(b*x+a))/d-c/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(-
I*d+I-c))+1/2*I*d^2*ln(1-(I-I*d+c)*(1+I*(d*((c+d*cot(b*x+a))/d-c/d)+c))^2
/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(I*d+I-c))*arctan(d*((c+d*cot(b*x+a)
)/d-c/d)+c)/(1+I*c+d)+1/2*I*d*ln(1-(I-I*d+c)*(1+I*(d*((c+d*cot(b*x+a))/d-c
/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(I*d+I-c))*arctan(d*((c+d*c
ot(b*x+a))/d-c/d)+c)/(1+I*c+d)+1/2*I*d/(-I-I*d+c)*ln(1-(I-I*d+c)*(1+I*(d*(
c+d*cot(b*x+a))/d-c/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(I*d+I-
c))*c*arctan(d*((c+d*cot(b*x+a))/d-c/d)+c)+1/2*d^2*arctan(d*((c+d*cot(b*x+
a))/d-c/d)+c)^2/(1+I*c+d)+1/4*d^2*polylog(2,(I-I*d+c)*(1+I*(d*((c+d*cot(b*
x+a))/d-c/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(I*d+I-c))/(1+I*c+
d)+1/2*d*arctan(d*((c+d*cot(b*x+a))/d-c/d)+c)^2/(1+I*c+d)+1/2*d/(-I-I*d+c)
*c*arctan(d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1/4*d*polylog(2,(I-I*d+c)*(1+I*(
d*((c+d*cot(b*x+a))/d-c/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(I*d
+I-c))/(1+I*c+d)+1/4*d/(-I-I*d+c)*polylog(2,(I-I*d+c)*(1+I*(d*((c+d*cot(b*
x+a))/d-c/d)+c))^2/((d*((c+d*cot(b*x+a))/d-c/d)+c)^2+1)/(I*d+I-c))*c)))

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 965 vs. $2(140) = 280$.

Time = 0.22 (sec) , antiderivative size = 965, normalized size of antiderivative = 4.87

$$\int \arctan(c + d \cot(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(arctan(c+d*cot(b*x+a)),x, algorithm="fricas")
```

output

```

1/8*(8*b*x*arctan(d*cot(b*x + a) + c) - 2*I*a*log(1/2*c^2 + I*c*d - 1/2*d^
2 - 1/2*(c^2 + d^2 + 2*d + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*
d + I)*sin(2*b*x + 2*a) + 1/2) + 2*I*a*log(1/2*c^2 + I*c*d - 1/2*d^2 - 1/2
*(c^2 + d^2 - 2*d + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*d + I)*
sin(2*b*x + 2*a) + 1/2) + 2*I*a*log(-1/2*c^2 + I*c*d + 1/2*d^2 + 1/2*(c^2
+ d^2 + 2*d + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 + 2*I*d + I)*sin(2*
b*x + 2*a) - 1/2) - 2*I*a*log(-1/2*c^2 + I*c*d + 1/2*d^2 + 1/2*(c^2 + d^2
- 2*d + 1)*cos(2*b*x + 2*a) + 1/2*(I*c^2 + I*d^2 - 2*I*d + I)*sin(2*b*x +
2*a) - 1/2) - 2*(-I*b*x - I*a)*log((c^2 + d^2 - (c^2 + 2*I*c*d - d^2 + 1)*
cos(2*b*x + 2*a) + (-I*c^2 + 2*c*d + I*d^2 - I)*sin(2*b*x + 2*a) + 2*d + 1
)/(c^2 + d^2 + 2*d + 1)) - 2*(I*b*x + I*a)*log((c^2 + d^2 - (c^2 - 2*I*c*d
- d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*c*d - I*d^2 + I)*sin(2*b*x + 2*a
) + 2*d + 1)/(c^2 + d^2 + 2*d + 1)) - 2*(I*b*x + I*a)*log((c^2 + d^2 - (c^
2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-I*c^2 + 2*c*d + I*d^2 - I)*sin
(2*b*x + 2*a) - 2*d + 1)/(c^2 + d^2 - 2*d + 1)) - 2*(-I*b*x - I*a)*log((c^
2 + d^2 - (c^2 - 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*c*d - I*
d^2 + I)*sin(2*b*x + 2*a) - 2*d + 1)/(c^2 + d^2 - 2*d + 1)) + dilog(-(c^2
+ d^2 - (c^2 + 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (-I*c^2 + 2*c*d + I*d
^2 - I)*sin(2*b*x + 2*a) + 2*d + 1)/(c^2 + d^2 + 2*d + 1) + 1) + dilog(-(c
^2 + d^2 - (c^2 - 2*I*c*d - d^2 + 1)*cos(2*b*x + 2*a) + (I*c^2 + 2*c*d ...

```

Sympy [F(-1)]

Timed out.

$$\int \arctan(c + d \cot(a + bx)) dx = \text{Timed out}$$

input

```
integrate(atan(c+d*cot(b*x+a)),x)
```

output

Timed out

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 532 vs. $2(140) = 280$.

Time = 0.21 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.69

$$\int \arctan(c + d \cot(a + bx)) dx =$$

$$d \left(\frac{8(bx+a) \arctan\left(\frac{cd+(c^2+1)\tan(bx+a)}{d}\right)}{d} - \frac{8(bx+a) \arctan\left(\frac{cd+(c^2+1)\tan(bx+a)}{d}\right) - 4 \arctan\left(\frac{cd+(c^2+1)\tan(bx+a)}{d}\right) \arctan\left(\frac{cd+(c^2+1)\tan(bx+a)}{d}\right)}{d} \right)$$

input `integrate(arctan(c+d*cot(b*x+a)),x, algorithm="maxima")`

output

```
-1/8*(d*(8*(b*x + a)*arctan((c*d + (c^2 + 1)*tan(b*x + a))/d)/d - (8*(b*x + a)*arctan((c*d + (c^2 + 1)*tan(b*x + a))/d) - 4*arctan((c*d + (c^2 + 1)*tan(b*x + a))/d)*arctan2((c*d + (c^2 + d + 1)*tan(b*x + a))/(c^2 + d^2 + 2*d + 1), -(c*d*tan(b*x + a) - c^2 - d - 1)/(c^2 + d^2 + 2*d + 1)) + 4*arctan((c*d + (c^2 + 1)*tan(b*x + a))/d)*arctan2(-(c*d + (c^2 - d + 1)*tan(b*x + a))/(c^2 + d^2 - 2*d + 1), -(c*d*tan(b*x + a) - c^2 + d - 1)/(c^2 + d^2 - 2*d + 1)) - (log(((c^2 + 1)*tan(b*x + a)^2 + c^2 + 1)/(c^2 + d^2 + 2*d + 1)) - log(((c^2 + 1)*tan(b*x + a)^2 + c^2 + 1)/(c^2 + d^2 - 2*d + 1)))*log((c^2 + 1)*d^2 + 2*(c^3 + c)*d*tan(b*x + a) + (c^4 + 2*c^2 + 1)*tan(b*x + a)^2) - 2*dilog(((I*c - 1)*tan(b*x + a) + I*d)/(c + I*d + I)) + 2*dilog(((I*c + 1)*tan(b*x + a) + I*d)/(c + I*d - I)) + 2*dilog(-(I*c - 1)*tan(b*x + a) + I*d)/(c - I*d + I) - 2*dilog(-(I*c + 1)*tan(b*x + a) + I*d)/(c - I*d - I))/d - 8*(b*x + a)*arctan(c + d/tan(b*x + a)) - 8*(b*x + a)*arctan((c*d + (c^2 + 1)*tan(b*x + a))/d))/b
```

Giac [F]

$$\int \arctan(c + d \cot(a + bx)) dx = \int \arctan(d \cot(bx + a) + c) dx$$

input `integrate(arctan(c+d*cot(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(d*cot(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + d \cot(a + bx)) dx = \int \operatorname{atan}(c + d \cot(a + bx)) dx$$

input `int(atan(c + d*cot(a + b*x)),x)`

output `int(atan(c + d*cot(a + b*x)), x)`

Reduce [F]

$$\int \arctan(c + d \cot(a + bx)) dx = \int \operatorname{atan}(\cot(bx + a) d + c) dx$$

input `int(atan(c+d*cot(b*x+a)),x)`

output `int(atan(cot(a + b*x)*d + c),x)`

3.64 $\int \frac{\arctan(c+d \cot(a+bx))}{x} dx$

Optimal result	512
Mathematica [N/A]	512
Rubi [N/A]	513
Maple [N/A]	513
Fricas [N/A]	514
Sympy [F(-1)]	514
Maxima [F(-1)]	514
Giac [N/A]	515
Mupad [N/A]	515
Reduce [N/A]	515

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + d \cot(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+d*cot(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 1.69 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \int \frac{\arctan(c + d \cot(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + d*Cot[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + d*Cot[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(d \cot(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\arctan(d \cot(a + bx) + c)}{x} dx$$

input `Int[ArcTan[c + d*Cot[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.49 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + d \cot(bx + a))}{x} dx$$

input `int(arctan(c+d*cot(b*x+a))/x,x)`

output `int(arctan(c+d*cot(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \int \frac{\arctan(d \cot(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*cot(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctan(d*cot(b*x + a) + c)/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+d*cot(b*x+a))/x,x)`

output `Timed out`

Maxima [F(-1)]

Timed out.

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(arctan(c+d*cot(b*x+a))/x,x, algorithm="maxima")`

output `Timed out`

Giac [N/A]

Not integrable

Time = 2.65 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \int \frac{\arctan(d \cot(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*cot(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctan(d*cot(b*x + a) + c)/x, x)`

Mupad [N/A]

Not integrable

Time = 0.63 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + d \cot(a + bx))}{x} dx$$

input `int(atan(c + d*cot(a + b*x))/x,x)`

output `int(atan(c + d*cot(a + b*x))/x, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \cot(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\cot(bx + a) d + c)}{x} dx$$

input `int(atan(c+d*cot(b*x+a))/x,x)`

output `int(atan(cot(a + b*x)*d + c)/x,x)`

3.65 $\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx$

Optimal result	517
Mathematica [A] (verified)	518
Rubi [A] (verified)	518
Maple [C] (warning: unable to verify)	522
Fricas [A] (verification not implemented)	523
Sympy [F(-2)]	523
Maxima [B] (verification not implemented)	524
Giac [F]	524
Mupad [F(-1)]	525
Reduce [F]	525

Optimal result

Integrand size = 21, antiderivative size = 154

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx = \frac{bx^4}{12} + \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx))$$

$$+ \frac{1}{6}ix^3 \log(1 - ice^{2ia+2ibx})$$

$$+ \frac{x^2 \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{4b}$$

$$+ \frac{ix \operatorname{PolyLog}(3, ice^{2ia+2ibx})}{4b^2}$$

$$- \frac{\operatorname{PolyLog}(4, ice^{2ia+2ibx})}{8b^3}$$

output

```
1/12*b*x^4-1/3*x^3*arctan(-c-(1-I*c)*cot(b*x+a))+1/6*I*x^3*ln(1-I*c*exp(2*
I*a+2*I*b*x))+1/4*x^2*polylog(2,I*c*exp(2*I*a+2*I*b*x))/b+1/4*I*x*polylog(
3,I*c*exp(2*I*a+2*I*b*x))/b^2-1/8*polylog(4,I*c*exp(2*I*a+2*I*b*x))/b^3
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.88

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx = \frac{1}{24} \left(8x^3 \arctan(c + (1 - ic) \cot(a + bx)) \right. \\ \left. + 4ix^3 \log \left(1 + \frac{ie^{-2i(a+bx)}}{c} \right) \right. \\ \left. - \frac{6x^2 \operatorname{PolyLog} \left(2, -\frac{ie^{-2i(a+bx)}}{c} \right)}{b} \right. \\ \left. + \frac{6ix \operatorname{PolyLog} \left(3, -\frac{ie^{-2i(a+bx)}}{c} \right)}{b^2} \right. \\ \left. + \frac{3 \operatorname{PolyLog} \left(4, -\frac{ie^{-2i(a+bx)}}{c} \right)}{b^3} \right)$$

input

```
Integrate[x^2*ArcTan[c + (1 - I*c)*Cot[a + b*x]],x]
```

output

```
(8*x^3*ArcTan[c + (1 - I*c)*Cot[a + b*x]] + (4*I)*x^3*Log[1 + I/(c*E^((2*I)*(a + b*x))))] - (6*x^2*PolyLog[2, (-I)/(c*E^((2*I)*(a + b*x))))]/b + ((6*I)*x*PolyLog[3, (-I)/(c*E^((2*I)*(a + b*x))))]/b^2 + (3*PolyLog[4, (-I)/(c*E^((2*I)*(a + b*x))))]/b^3)/24
```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.29, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {5696, 25, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx$$

↓ 5696

$$\begin{aligned}
 & \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx)) - \frac{1}{3}ib \int -\frac{x^3}{e^{2ia+2ibx}c+i} dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3}ib \int \frac{x^3}{e^{2ia+2ibx}c+i} dx + \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx)) \\
 & \quad \downarrow \text{2615} \\
 & \frac{1}{3}ib \left(ic \int \frac{e^{2ia+2ibx}x^3}{e^{2ia+2ibx}c+i} dx - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx)) \\
 & \quad \downarrow \text{2620} \\
 & \frac{1}{3}ib \left(ic \left(\frac{3i \int x^2 \log(1 - ice^{2ia+2ibx}) dx}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + \\
 & \quad \quad \quad (1 - ic) \cot(a + bx)) \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{3}ib \left(ic \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \int x \text{PolyLog}(2, ice^{2ia+2ibx}) dx}{b} \right)}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^4}{4} \right) + \\
 & \quad \quad \quad \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx)) \\
 & \quad \downarrow \text{7163} \\
 & \frac{1}{3}ib \left(ic \left(\frac{3i \left(\frac{ix^2 \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int \text{PolyLog}(3, ice^{2ia+2ibx}) dx}{2b} - \frac{ix \text{PolyLog}(3, ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right) \right) + \\
 & \quad \quad \quad \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx)) \\
 & \quad \downarrow \text{2720}
 \end{aligned}$$

$$\frac{1}{3}ib \left(ic \frac{\left(3i \left(\frac{ix^2 \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int e^{-2ia-2ibx} \text{PolyLog}(3, ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} - \frac{ix \text{PolyLog}(3, ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} - ix^3 \log \right)}{\right)} - \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx))$$

7143

$$\frac{1}{3}ib \left(ic \frac{\left(3i \left(\frac{ix^2 \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\text{PolyLog}(4, ice^{2ia+2ibx})}{4b^2} - \frac{ix \text{PolyLog}(3, ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} - \frac{ix^3 \log(1 - ice^{2ia+2ibx})}{2bc} \right)}{\right)} + \frac{1}{3}x^3 \arctan(c + (1 - ic) \cot(a + bx))$$

```
input Int[x^2*ArcTan[c + (1 - I*c)*Cot[a + b*x]], x]
```

```
output (x^3*ArcTan[c + (1 - I*c)*Cot[a + b*x]])/3 + (I/3)*b*((-1/4*I)*x^4 + I*c*((-1/2*I)*x^3*Log[1 - I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) + (((3*I)/2)*(((I/2)*x^2*PolyLog[2, I*c*E^((2*I)*a + (2*I)*b*x)]/b - (I*(((1/2*I)*x*PolyLog[3, I*c*E^((2*I)*a + (2*I)*b*x)]/b + PolyLog[4, I*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2))))/b))/(b*c)))
```

Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 2615 Int[(((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5696

```
Int[ArcTan[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Cot[a + b*x]]/(f*(m +
1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2
*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && E
qQ[(c - I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[(((e_) + (f_)*(x_)^(m_))*PolyLog[n_, (d_)*((F_)^((c_)*((a_) + (b_
)*(x_))))^(p_)]), x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.20 (sec) , antiderivative size = 1487, normalized size of antiderivative = 9.66

method	result	size
risch	Expression too large to display	1487

input `int(-x^2*arctan(-c-(1-I*c)*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
1/2*I/b^3*a^3*ln(1-I*exp(I*(b*x+a))*(-I*c)^(1/2))+1/3*I*x^3*ln(exp(I*(b*x+a)))
-1/2*I/b^2*ln(1-I*exp(2*I*(b*x+a))*c)*x*a^2-1/6*I/b^3*a^3*ln(exp(2*I*(b*x+a))*c+I)
+1/6*I*x^3*ln(1-I*exp(2*I*(b*x+a))*c)+1/2*I/b^3*a^3*ln(1+I*exp(I*(b*x+a))*(-I*c)^(1/2))
+1/4*x^2*polylog(2,I*exp(2*I*(b*x+a))*c)/b-1/4/b^3*polylog(2,I*exp(2*I*(b*x+a))*c)*a^2
-1/3*I/b^3*ln(1-I*exp(2*I*(b*x+a))*c)*a^3-1/8*polylog(4,I*exp(2*I*(b*x+a))*c)/b^3
-1/12*I*(2*I*Pi+I*Pi*csgn(I*exp(2*I*(b*x+a))))^3-2*I*Pi*csgn(I*exp(I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a)))^2
+I*Pi*csgn(I*exp(I*(b*x+a)))^2*csgn(I*exp(2*I*(b*x+a)))-2*ln(I+c)+I*Pi*csgn(I*exp(2*I*(b*x+a)))*(I+c)
/(exp(2*I*(b*x+a))-1)*csgn(exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I)
/(exp(2*I*(b*x+a))-1))*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))+I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I)
/(exp(2*I*(b*x+a))-1))*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))^2-I*Pi*csgn(I*exp(2*I*(b*x+a)))*(I+c)
/(exp(2*I*(b*x+a))-1))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))^2-I*Pi*csgn(I*(I+c))*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I*(exp(2*I*(b*x+a))*c+I))*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))+I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I*(I+c))*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))+I*Pi*csgn(I*exp(2*I*(b*x+a)))*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))-I*Pi*csgn((exp(2*...
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.08

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx$$

$$= \frac{2b^4x^4 + 4ib^3x^3 \log\left(-\frac{(c+i)e^{(2ibx+2ia)}}{ce^{(2ibx+2ia)+i}}\right) + 6b^2x^2 \operatorname{Li}_2\left(ice^{(2ibx+2ia)}\right) - 2a^4 - 4ia^3 \log\left(\frac{ce^{(2ibx+2ia)+i}}{c}\right) + 6ibx}{24b^3}$$

input `integrate(-x^2*arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="fricas")`

output `1/24*(2*b^4*x^4 + 4*I*b^3*x^3*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x + 2*I*a) + I)) + 6*b^2*x^2*dilog(I*c*e^(2*I*b*x + 2*I*a)) - 2*a^4 - 4*I*a^3*log((c*e^(2*I*b*x + 2*I*a) + I)/c) + 6*I*b*x*polylog(3, I*c*e^(2*I*b*x + 2*I*a)) - 4*(-I*b^3*x^3 - I*a^3)*log(-I*c*e^(2*I*b*x + 2*I*a) + 1) - 3*polylog(4, I*c*e^(2*I*b*x + 2*I*a)))/b^3`

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(-x**2*atan(-c-(1-I*c)*cot(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) - 1 of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 309 vs. $2(110) = 220$.

Time = 0.08 (sec) , antiderivative size = 309, normalized size of antiderivative = 2.01

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx$$

$$= \frac{4 \left((bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2 \right) \arctan((-ic+1) \cot(bx+a)+c)}{b^2} + \frac{(-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(4i(bx+a)^3 - 9i(bx+a)a^2)) \operatorname{arctan}^2(c \cos(2bx+2a), c \sin(2bx+2a) + 1) - 3(4I(bx+a)^2 - 6I(bx+a)a + 3Ia^2) \operatorname{dilog}(Ic e^{(2Ibx+2Ia)}) + (4(bx+a)^3 - 9(bx+a)^2 a + 9(bx+a)a^2) \log(c^2 \cos(2bx+2a)^2 + c^2 \sin(2bx+2a)^2 + 2c \sin(2bx+2a) + 1) + 3(4bx+a) \operatorname{polylog}(3, Ic e^{(2Ibx+2Ia)}) + 6I \operatorname{polylog}(4, Ic e^{(2Ibx+2Ia)})}{b^2} (Ic - 1) / (b^2 (c + I)) / b$$

input `integrate(-x^2*arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="maxima")`

output

```
1/12*(4*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctan((-I*c + 1)
)*cot(b*x + a) + c)/b^2 + (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b
*x + a)^2*a^2 - 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*I*(b*x + a)*a^2
)*arctan2(c*cos(2*b*x + 2*a), c*sin(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2
- 6*I*(b*x + a)*a + 3*I*a^2)*dilog(I*c*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a
)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*
sin(2*b*x + 2*a)^2 + 2*c*sin(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog(3,
I*c*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, I*c*e^(2*I*b*x + 2*I*a))*(I*c -
1)/(b^2*(c + I))/b
```

Giac [F]

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx = \int -x^2 \arctan(-(-ic + 1) \cot(bx + a) - c) dx$$

input `integrate(-x^2*arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="giac")`

output

```
integrate(-x^2*arctan(-(-I*c + 1)*cot(b*x + a) - c), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx = \int x^2 \operatorname{atan}(c - \cot(a + bx) (-1 + ci)) dx$$

input `int(x^2*atan(c - cot(a + b*x)*(c*1i - 1)),x)`

output `int(x^2*atan(c - cot(a + b*x)*(c*1i - 1)), x)`

Reduce [F]

$$\int x^2 \arctan(c + (1 - ic) \cot(a + bx)) dx = - \left(\int \operatorname{atan}(\cot(bx + a) ci - \cot(bx + a) - c) x^2 dx \right)$$

input `int(-x^2*atan(-c-(1-I*c)*cot(b*x+a)),x)`

output `- int(atan(cot(a + b*x)*c*i - cot(a + b*x) - c)*x**2,x)`

3.66 $\int x \arctan(c + (1 - ic) \cot(a + bx)) dx$

Optimal result	526
Mathematica [A] (verified)	527
Rubi [A] (verified)	527
Maple [C] (warning: unable to verify)	530
Fricas [A] (verification not implemented)	531
Sympy [F(-2)]	531
Maxima [B] (verification not implemented)	532
Giac [F]	532
Mupad [F(-1)]	533
Reduce [F]	533

Optimal result

Integrand size = 19, antiderivative size = 123

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx = \frac{bx^3}{6} + \frac{1}{2}x^2 \arctan(c + (1 - ic) \cot(a + bx)) + \frac{1}{4}ix^2 \log(1 - ice^{2ia+2ibx}) + \frac{x \operatorname{PolyLog}(2, ice^{2ia+2ibx})}{4b} + \frac{i \operatorname{PolyLog}(3, ice^{2ia+2ibx})}{8b^2}$$

output

```
1/6*b*x^3-1/2*x^2*arctan(-c-(1-I*c)*cot(b*x+a))+1/4*I*x^2*ln(1-I*c*exp(2*I*a+2*I*b*x))+1/4*x*polylog(2,I*c*exp(2*I*a+2*I*b*x))/b+1/8*I*polylog(3,I*c*exp(2*I*a+2*I*b*x))/b^2
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.89

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx = \frac{1}{2} x^2 \arctan(c + (1 - ic) \cot(a + bx)) + \frac{i \left(2b^2 x^2 \log \left(1 + \frac{ie^{-2i(a+bx)}}{c} \right) + 2ibx \operatorname{PolyLog} \left(2, -\frac{ie^{-2i(a+bx)}}{c} \right) + \operatorname{PolyLog} \left(3, -\frac{ie^{-2i(a+bx)}}{c} \right) \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c + (1 - I*c)*Cot[a + b*x]], x]
```

output

```
(x^2*ArcTan[c + (1 - I*c)*Cot[a + b*x]])/2 + ((I/8)*(2*b^2*x^2*Log[1 + I/(c*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, (-I)/(c*E^((2*I)*(a + b*x)))] + PolyLog[3, (-I)/(c*E^((2*I)*(a + b*x)))]))/b^2
```

Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.26, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {5696, 25, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arctan(c + (1 - ic) \cot(a + bx)) dx \\ & \quad \downarrow \text{5696} \\ & \frac{1}{2} x^2 \arctan(c + (1 - ic) \cot(a + bx)) - \frac{1}{2} ib \int -\frac{x^2}{e^{2ia+2ibx} c + i} dx \\ & \quad \downarrow \text{25} \\ & \frac{1}{2} ib \int \frac{x^2}{e^{2ia+2ibx} c + i} dx + \frac{1}{2} x^2 \arctan(c + (1 - ic) \cot(a + bx)) \\ & \quad \downarrow \text{2615} \\ & \frac{1}{2} ib \left(ic \int \frac{e^{2ia+2ibx} x^2}{e^{2ia+2ibx} c + i} dx - \frac{ix^3}{3} \right) + \frac{1}{2} x^2 \arctan(c + (1 - ic) \cot(a + bx)) \end{aligned}$$

$$\frac{1}{2}ib \left(ic \left(\frac{i \int x \log(1 - ice^{2ia+2ibx}) dx}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c + (1 - ic) \cot(a + bx))$$

↓ 2620

$$\frac{1}{2}ib \left(ic \left(\frac{i \left(\frac{ix \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{i \int \text{PolyLog}(2, ice^{2ia+2ibx}) dx}{2b} \right)}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c + (1 - ic) \cot(a + bx))$$

↓ 2720

$$\frac{1}{2}ib \left(ic \left(\frac{i \left(\frac{ix \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \text{PolyLog}(2, ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} \right)}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c + (1 - ic) \cot(a + bx))$$

↓ 7143

$$\frac{1}{2}ib \left(ic \left(\frac{i \left(\frac{ix \text{PolyLog}(2, ice^{2ia+2ibx})}{2b} - \frac{\text{PolyLog}(3, ice^{2ia+2ibx})}{4b^2} \right)}{bc} - \frac{ix^2 \log(1 - ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c + (1 - ic) \cot(a + bx))$$

input `Int[x*ArcTan[c + (1 - I*c)*Cot[a + b*x]],x]`

output

```
(x^2*ArcTan[c + (1 - I*c)*Cot[a + b*x]])/2 + (I/2)*b*((-1/3*I)*x^3 + I*c*((-1/2*I)*x^2*Log[1 - I*c*E^((2*I)*a + (2*I)*b*x)])/(b*c) + I*(((I/2)*x*PolyLog[2, I*c*E^((2*I)*a + (2*I)*b*x)]/b - PolyLog[3, I*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/(b*c))
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2615 $\text{Int}[\left(\left(\left(\text{c}_.\right) + \left(\text{d}_.\right) \cdot \left(\text{x}_.\right)^{\left(\text{m}_.\right)}\right) / \left(\left(\text{a}_.\right) + \left(\text{b}_.\right) \cdot \left(\text{F}_.\right)^{\left(\left(\text{g}_.\right) \cdot \left(\text{e}_.\right) + \left(\text{f}_.\right) \cdot \left(\text{x}_.\right)\right)\right)\right)^{\left(\text{n}_.\right)}, \text{x_Symbol}] \rightarrow \text{Simp}[\left(\text{c} + \text{d} \cdot \text{x}\right)^{\left(\text{m} + 1\right)} / \left(\text{a} \cdot \text{d} \cdot \left(\text{m} + 1\right)\right), \text{x}] - \text{Simp}\left[\frac{\text{b}}{\text{a}} \quad \text{Int}\left[\left(\text{c} + \text{d} \cdot \text{x}\right)^{\text{m}} \cdot \left(\text{F}^{\left(\text{g} \cdot \left(\text{e} + \text{f} \cdot \text{x}\right)\right)}\right)^{\text{n}} / \left(\text{a} + \text{b} \cdot \left(\text{F}^{\left(\text{g} \cdot \left(\text{e} + \text{f} \cdot \text{x}\right)\right)}\right)^{\text{n}}\right), \text{x}\right] \text{ ; FreeQ}\left\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\right\}, \text{x}\right] \&\& \text{IGtQ}\left[\text{m}, 0\right]$
- rule 2620 $\text{Int}[\left(\left(\left(\text{F}_.\right)^{\left(\left(\text{g}_.\right) \cdot \left(\text{e}_.\right) + \left(\text{f}_.\right) \cdot \left(\text{x}_.\right)\right)\right)\right)^{\left(\text{n}_.\right)} \cdot \left(\left(\text{c}_.\right) + \left(\text{d}_.\right) \cdot \left(\text{x}_.\right)^{\left(\text{m}_.\right)}\right) / \left(\left(\text{a}_.\right) + \left(\text{b}_.\right) \cdot \left(\text{F}_.\right)^{\left(\left(\text{g}_.\right) \cdot \left(\text{e}_.\right) + \left(\text{f}_.\right) \cdot \left(\text{x}_.\right)\right)\right)^{\left(\text{n}_.\right)}, \text{x_Symbol}] \rightarrow \text{Simp}\left[\left(\left(\text{c} + \text{d} \cdot \text{x}\right)^{\text{m}} / \left(\text{b} \cdot \text{f} \cdot \text{g} \cdot \text{n} \cdot \text{Log}\left[\text{F}\right]\right)\right) \cdot \text{Log}\left[1 + \text{b} \cdot \left(\text{F}^{\left(\text{g} \cdot \left(\text{e} + \text{f} \cdot \text{x}\right)\right)}\right)^{\text{n}} / \text{a}\right], \text{x}\right] - \text{Simp}\left[\text{d} \cdot \left(\text{m} / \left(\text{b} \cdot \text{f} \cdot \text{g} \cdot \text{n} \cdot \text{Log}\left[\text{F}\right]\right)\right) \quad \text{Int}\left[\left(\text{c} + \text{d} \cdot \text{x}\right)^{\left(\text{m} - 1\right)} \cdot \text{Log}\left[1 + \text{b} \cdot \left(\text{F}^{\left(\text{g} \cdot \left(\text{e} + \text{f} \cdot \text{x}\right)\right)}\right)^{\text{n}} / \text{a}\right], \text{x}\right], \text{x}\right] \text{ ; FreeQ}\left\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\right\}, \text{x}\right] \&\& \text{IGtQ}\left[\text{m}, 0\right]$
- rule 2720 $\text{Int}\left[\text{u}_., \text{x_Symbol}\right] \rightarrow \text{With}\left[\left\{\text{v} = \text{FunctionOfExponential}\left[\text{u}, \text{x}\right]\right\}, \text{Simp}\left[\text{v} / \text{D}\left[\text{v}, \text{x}\right] \quad \text{Subst}\left[\text{Int}\left[\text{FunctionOfExponentialFunction}\left[\text{u}, \text{x}\right] / \text{x}, \text{x}\right], \text{x}, \text{v}\right], \text{x}\right] \text{ ; FunctionOfExponentialQ}\left[\text{u}, \text{x}\right] \&\& \text{!MatchQ}\left[\text{u}, \left(\text{w}_.\right) \cdot \left(\left(\text{a}_.\right) \cdot \left(\text{v}_.\right)^{\left(\text{n}_.\right)}\right)^{\left(\text{m}_.\right)}\right] \text{ ; FreeQ}\left\{\text{a}, \text{m}, \text{n}\right\}, \text{x}\right] \&\& \text{IntegerQ}\left[\text{m} \cdot \text{n}\right] \&\& \text{!MatchQ}\left[\text{u}, \text{E}^{\left(\left(\text{c}_.\right) \cdot \left(\left(\text{a}_.\right) + \left(\text{b}_.\right) \cdot \text{x}\right)\right)} \cdot \left(\text{F}_.\right)\left[\text{v}_.\right]\right] \text{ ; FreeQ}\left\{\text{a}, \text{b}, \text{c}\right\}, \text{x}\right] \&\& \text{InverseFunctionQ}\left[\text{F}\left[\text{x}\right]\right]$
- rule 3011 $\text{Int}\left[\text{Log}\left[1 + \left(\text{e}_.\right) \cdot \left(\left(\text{F}_.\right)^{\left(\left(\text{c}_.\right) \cdot \left(\left(\text{a}_.\right) + \left(\text{b}_.\right) \cdot \left(\text{x}_.\right)\right)\right)\right)^{\left(\text{n}_.\right)}\right] \cdot \left(\left(\text{f}_.\right) + \left(\text{g}_.\right) \cdot \left(\text{x}_.\right)^{\left(\text{m}_.\right)}\right), \text{x_Symbol}] \rightarrow \text{Simp}\left[\left(-\left(\text{f} + \text{g} \cdot \text{x}\right)^{\text{m}} \cdot \left(\text{PolyLog}\left[2, \left(-\text{e}\right) \cdot \left(\text{F}^{\left(\text{c} \cdot \left(\text{a} + \text{b} \cdot \text{x}\right)\right)}\right)^{\text{n}}\right] / \left(\text{b} \cdot \text{c} \cdot \text{n} \cdot \text{Log}\left[\text{F}\right]\right)\right), \text{x}\right] + \text{Simp}\left[\text{g} \cdot \left(\text{m} / \left(\text{b} \cdot \text{c} \cdot \text{n} \cdot \text{Log}\left[\text{F}\right]\right)\right) \quad \text{Int}\left[\left(\text{f} + \text{g} \cdot \text{x}\right)^{\left(\text{m} - 1\right)} \cdot \text{PolyLog}\left[2, \left(-\text{e}\right) \cdot \left(\text{F}^{\left(\text{c} \cdot \left(\text{a} + \text{b} \cdot \text{x}\right)\right)}\right)^{\text{n}}\right], \text{x}\right], \text{x}\right] \text{ ; FreeQ}\left\{\text{F}, \text{a}, \text{b}, \text{c}, \text{e}, \text{f}, \text{g}, \text{n}\right\}, \text{x}\right] \&\& \text{GtQ}\left[\text{m}, 0\right]$
- rule 5696 $\text{Int}\left[\text{ArcTan}\left[\left(\text{c}_.\right) + \text{Cot}\left[\left(\text{a}_.\right) + \left(\text{b}_.\right) \cdot \left(\text{x}_.\right)\right] \cdot \left(\text{d}_.\right)\right] \cdot \left(\left(\text{e}_.\right) + \left(\text{f}_.\right) \cdot \left(\text{x}_.\right)^{\left(\text{m}_.\right)}\right), \text{x_Symbol}] \rightarrow \text{Simp}\left[\left(\text{e} + \text{f} \cdot \text{x}\right)^{\left(\text{m} + 1\right)} \cdot \left(\text{ArcTan}\left[\text{c} + \text{d} \cdot \text{Cot}\left[\text{a} + \text{b} \cdot \text{x}\right]\right] / \left(\text{f} \cdot \left(\text{m} + 1\right)\right)\right), \text{x}\right] - \text{Simp}\left[\text{I} \cdot \left(\text{b} / \left(\text{f} \cdot \left(\text{m} + 1\right)\right)\right) \quad \text{Int}\left[\left(\text{e} + \text{f} \cdot \text{x}\right)^{\left(\text{m} + 1\right)} / \left(\text{c} - \text{I} \cdot \text{d} - \text{c} \cdot \text{E}^{\left(2 \cdot \text{I} \cdot \text{a} + 2 \cdot \text{I} \cdot \text{b} \cdot \text{x}\right)}\right), \text{x}\right], \text{x}\right] \text{ ; FreeQ}\left\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\right\}, \text{x}\right] \&\& \text{IGtQ}\left[\text{m}, 0\right] \&\& \text{EqQ}\left[\left(\text{c} - \text{I} \cdot \text{d}\right)^2, -1\right]$

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.48 (sec) , antiderivative size = 1452, normalized size of antiderivative = 11.80

method	result	size
risch	Expression too large to display	1452

input

```
int(-x*arctan(-c-(1-I*c)*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/8*I*x^2*(2*I*Pi+I*Pi*csgn(I*exp(2*I*(b*x+a)))^3-2*I*Pi*csgn(I*exp(I*(b*x+a)))
*csgn(I*exp(2*I*(b*x+a)))^2+I*Pi*csgn(I*exp(I*(b*x+a)))^2*csgn(I*exp(2*I*(b*x+a)))
-2*ln(I+c)+I*Pi*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))*csgn(exp(2*I*(b*x+a))
*(I+c)/(exp(2*I*(b*x+a))-1))-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))*csgn(
(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))
+I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))^2-
I*Pi*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))*csgn(exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))^2-
I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I/(exp(2*I*(b*x+a))-1))
*(I+c)^2-I*Pi*csgn(I*(I+c))*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I*(exp(2*I*(b*x+a))*c+I))
*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))+I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I*(I+c))*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))
+I*Pi*csgn(I*exp(2*I*(b*x+a)))
*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))-I*Pi*csgn((exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))^2-
I*Pi*csgn(I*(exp(2*I*(b*x+a))*c+I)/(exp(2*I*(b*x+a))-1))^3+I*Pi*csgn(I*exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))^3+
I*Pi*csgn(exp(2*I*(b*x+a))*(I+c)/(exp(2*I*(b*x+a))-1))^3+I*Pi*csgn(I/(exp(2*I*(b*x+a))-1)*(I+c))^3-I*Pi*csgn(exp(2*I*(b*x+a))
*(I+c)/(exp(2*I*(b*x+a))-1))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))-1))^2-I*P...
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.17

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx$$

$$= \frac{4b^3x^3 + 6ib^2x^2 \log\left(-\frac{(c+i)e^{(2ibx+2ia)}}{ce^{(2ibx+2ia)}+i}\right) + 4a^3 + 6bx \operatorname{Li}_2\left(ice^{(2ibx+2ia)}\right) + 6ia^2 \log\left(\frac{ce^{(2ibx+2ia)}+i}{c}\right) - 6(-ib^2x^2 + I a^2) \log(-I c e^{(2ibx+2ia)} + 1) + 3I \operatorname{polylog}(3, I c e^{(2ibx+2ia)})}{24b^2}$$

input `integrate(-x*arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="fricas")`

output `1/24*(4*b^3*x^3 + 6*I*b^2*x^2*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x + 2*I*a) + I)) + 4*a^3 + 6*b*x*dilog(I*c*e^(2*I*b*x + 2*I*a)) + 6*I*a^2*log((c*e^(2*I*b*x + 2*I*a) + I)/c) - 6*(-I*b^2*x^2 + I*a^2)*log(-I*c*e^(2*I*b*x + 2*I*a) + 1) + 3*I*polylog(3, I*c*e^(2*I*b*x + 2*I*a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(-x*atan(-c-(1-I*c)*cot(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) - 1 of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 217 vs. $2(88) = 176$.

Time = 0.07 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.76

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx$$

$$= \frac{6((bx+a)^2 - 2(bx+a)a) \arctan((-ic+1) \cot(bx+a)+c)}{b} + \frac{(-4i(bx+a)^3 + 12i(bx+a)^2a - 6ibx \operatorname{Li}_2(ice^{(2ibx+2ia)}) - 6(i(bx+a)^2 - 2i(bx+a)a))}{b}$$

input `integrate(-x*arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="maxima")`

output `1/12*(6*((b*x + a)^2 - 2*(b*x + a)*a)*arctan((-I*c + 1)*cot(b*x + a) + c)/b + (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog(I*c*e^(2*I*b*x + 2*I*a)) - 6*(I*(b*x + a)^2 - 2*I*(b*x + a)*a)*arctan2(c*cos(2*b*x + 2*a), c*sin(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 + 2*c*sin(2*b*x + 2*a) + 1) + 3*polylog(3, I*c*e^(2*I*b*x + 2*I*a)))*(I*c - 1)/(b*(c + I))/b`

Giac [F]

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx = \int -x \arctan(-(-ic + 1) \cot(bx + a) - c) dx$$

input `integrate(-x*arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-x*arctan(-(-I*c + 1)*cot(b*x + a) - c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx = \int x \operatorname{atan}(c - \cot(a + bx) (-1 + ci)) dx$$

input `int(x*atan(c - cot(a + b*x)*(c*1i - 1)),x)`

output `int(x*atan(c - cot(a + b*x)*(c*1i - 1)), x)`

Reduce [F]

$$\int x \arctan(c + (1 - ic) \cot(a + bx)) dx = - \left(\int \operatorname{atan}(\cot(bx + a) ci - \cot(bx + a) - c) x dx \right)$$

input `int(-x*atan(-c-(1-I*c)*cot(b*x+a)),x)`

output `- int(atan(cot(a + b*x)*c*i - cot(a + b*x) - c)*x,x)`

3.67 $\int \arctan(c + (1 - ic) \cot(a + bx)) dx$

Optimal result	534
Mathematica [B] (warning: unable to verify)	534
Rubi [A] (verified)	535
Maple [B] (verified)	537
Fricas [A] (verification not implemented)	538
Sympy [F(-2)]	539
Maxima [B] (verification not implemented)	539
Giac [F]	540
Mupad [F(-1)]	540
Reduce [F]	541

Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = \frac{bx^2}{2} + x \arctan(c + (1 - ic) \cot(a + bx)) + \frac{1}{2}ix \log(1 - ice^{2ia+2ibx}) + \frac{\text{PolyLog}(2, ice^{2ia+2ibx})}{4b}$$

output `1/2*b*x^2-x*arctan(-c-(1-I*c)*cot(b*x+a))+1/2*I*x*ln(1-I*c*exp(2*I*a+2*I*b*x))+1/4*polylog(2,I*c*exp(2*I*a+2*I*b*x))/b`

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 929 vs. 2(85) = 170.

Time = 9.46 (sec) , antiderivative size = 929, normalized size of antiderivative = 10.93

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = x \arctan(c + (1 - ic) \cot(a + bx))$$

$$(i + \cot(a + bx))(1 + ic + (i + c) \cot(a + bx)) \left(2ibx + \log \left(1 - \frac{\sec(bx)((-i+c) \cos(a)+i(i+c) \sin(a))(\cos(a+bx)-2c}{2c} \right) \right)$$

input `Integrate[ArcTan[c + (1 - I*c)*Cot[a + b*x]],x]`

output `x*ArcTan[c + (1 - I*c)*Cot[a + b*x]] - (I*x*Csc[a + b*x]^2*(2*b*x*Log[2*Cos[b*x]*(Cos[b*x] - I*Sin[b*x])] + I*Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x])]/(2*c)]*Log[1 - I*Tan[b*x]] - I*Log[(Sec[b*x]*((1 - I*c)*Cos[a + b*x] + (-I + c)*Sin[a + b*x])]/(2*Cos[a] - (2*I)*Sin[a])]*Log[1 + I*Tan[b*x]] + I*PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x]] + I*PolyLog[2, (Sec[b*x]*((-I + c)*Cos[a] + I*(I + c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x])]/(2*c))] - I*PolyLog[2, (Sec[b*x]*((1 + I*c)*Cos[a] - (I + c)*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x])/2)]*(Cos[b*x] - I*Sin[b*x])*(Cos[b*x] + I*Sin[b*x])/((I + Cot[a + b*x])*(1 + I*c + (I + c)*Cot[a + b*x])*((2*I)*b*x + Log[1 - (Sec[b*x]*((-I + c)*Cos[a] + I*(I + c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x])]/(2*c))] + Log[1 + (Sec[b*x]*((-1 - I*c)*Cos[a] + (I + c)*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x])/2] + ((-I + c)*Cos[a + b*x]*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]]))/((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x]) + ((I + c)*(Log[1 - I*Tan[b*x]] - Log[1 + I*Tan[b*x]])*Sin[a + b*x])/((1 - I*c)*Cos[a + b*x] + (-I + c)*Sin[a + b*x]) + 2*b*x*Tan[b*x] + I*Log[1 - (Sec[b*x]*((-I + c)*Cos[a] + I*(I + c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x])]/(2*c)]*Tan[b*x] - I*Log[1 + (Sec[b*x]*((-1 - I*c)*Cos[a] + (I + c)*Sin[a])*(Cos[a + b*x] + I*Sin[a + b*x])/2]*Tan[b*x] - I*Log[1 - I*Tan[b*x]]*Tan[b*x] + I*Cos[a]^2*Log[1 + I*Tan[b*x]]*Tan[b*x] + I*Log[1 + I*Tan[b*x]]*Sin[a]^2*Tan[b*x] + (I*Log[(Sec[...`

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.26, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {5688, 25, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx$$

$$\downarrow 5688$$

$$x \arctan(c + (1 - ic) \cot(a + bx)) - ib \int -\frac{x}{e^{2ia+2ibx}c + i} dx$$

$$\begin{aligned}
& \downarrow 25 \\
& ib \int \frac{x}{e^{2ia+2ibx}c+i} dx + x \arctan(c + (1-ic) \cot(a+bx)) \\
& \downarrow 2615 \\
& ib \left(ic \int \frac{e^{2ia+2ibx}x}{e^{2ia+2ibx}c+i} dx - \frac{ix^2}{2} \right) + x \arctan(c + (1-ic) \cot(a+bx)) \\
& \downarrow 2620 \\
& ib \left(ic \left(\frac{i \int \log(1-ice^{2ia+2ibx}) dx}{2bc} - \frac{ix \log(1-ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right) + x \arctan(c + (1-ic) \cot(a+bx)) \\
& \downarrow 2715 \\
& ib \left(ic \left(\frac{\int e^{-2ia-2ibx} \log(1-ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2c} - \frac{ix \log(1-ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right) + \\
& \quad x \arctan(c + (1-ic) \cot(a+bx)) \\
& \downarrow 2838 \\
& \quad x \arctan(c + (1-ic) \cot(a+bx)) + \\
& ib \left(ic \left(-\frac{\text{PolyLog}(2, ice^{2ia+2ibx})}{4b^2c} - \frac{ix \log(1-ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right)
\end{aligned}$$

input `Int[ArcTan[c + (1 - I*c)*Cot[a + b*x]],x]`

output `x*ArcTan[c + (1 - I*c)*Cot[a + b*x]] + I*b*((-1/2*I)*x^2 + I*c((((-1/2*I)*x*Log[1 - I*c*E^((2*I)*a + (2*I)*b*x)])/(b*c) - PolyLog[2, I*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2*c)))`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2615 $\text{Int}[(((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.}))))^{\text{n}_.}), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m} + 1} / (\text{a} * \text{d} * (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b}/\text{a} \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m}} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}}) / (\text{a} + \text{b} * (\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}})), \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2620 $\text{Int}[(((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.}), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m}} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}]) * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}}) / \text{a}], \text{x}] - \text{Simp}[\text{d} * (\text{m} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m} - 1} * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}}) / \text{a}], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2715 $\text{Int}[\text{Log}[(\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.} * ((\text{e}_.) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.}), \text{x_Symbol}] \rightarrow \text{Simp}[1 / (\text{d} * \text{e} * \text{n} * \text{Log}[\text{F}]) \quad \text{Subst}[\text{Int}[\text{Log}[\text{a} + \text{b} * \text{x}] / \text{x}, \text{x}], \text{x}, (\text{F}^{\text{e}}(\text{c} + \text{d} * \text{x}))^{\text{n}}], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}\} \&\& \text{GtQ}[\text{a}, 0]$
- rule 2838 $\text{Int}[\text{Log}[(\text{c}_.) * ((\text{d}_.) + (\text{e}_.) * (\text{x}_.)^{\text{n}_.})] / (\text{x}_.), \text{x_Symbol}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-\text{c}) * \text{e} * \text{x}^{\text{n}}] / \text{n}, \text{x}] \text{ ; FreeQ}\{\text{c}, \text{d}, \text{e}, \text{n}\}, \text{x}\} \&\& \text{EqQ}[\text{c} * \text{d}, 1]$
- rule 5688 $\text{Int}[\text{ArcTan}[(\text{c}_.) + \text{Cot}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)] * (\text{d}_.)], \text{x_Symbol}] \rightarrow \text{Simp}[\text{x} * \text{ArcTan}[\text{c} + \text{d} * \text{Cot}[\text{a} + \text{b} * \text{x}]], \text{x}] - \text{Simp}[\text{I} * \text{b} \quad \text{Int}[\text{x} / (\text{c} - \text{I} * \text{d} - \text{c} * \text{E}^{(2 * \text{I} * \text{a} + 2 * \text{I} * \text{b} * \text{x})}), \text{x}], \text{x}] \text{ ; FreeQ}\{\text{a}, \text{b}, \text{c}, \text{d}\}, \text{x}\} \&\& \text{EqQ}[(\text{c} - \text{I} * \text{d})^2, -1]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 583 vs. $2(73) = 146$.

Time = 1.73 (sec) , antiderivative size = 584, normalized size of antiderivative = 6.87

method	result
derivativdivides	$-\frac{\arctan(-c+(ic-1)\cot(bx+a))\ln((ic-1)\cot(bx+a)+c+i)c^2}{2i+2c} - \frac{2i\arctan(-c+(ic-1)\cot(bx+a))\ln((ic-1)\cot(bx+a)+c+i)c}{2i+2c} + \dots$
default	$-\frac{\arctan(-c+(ic-1)\cot(bx+a))\ln((ic-1)\cot(bx+a)+c+i)c^2}{2i+2c} - \frac{2i\arctan(-c+(ic-1)\cot(bx+a))\ln((ic-1)\cot(bx+a)+c+i)c}{2i+2c} + \dots$
risch	Expression too large to display

```
input int(-arctan(-c-(1-I*c)*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

```
output -1/b/(-1+I*c)*(-arctan(-c+(-1+I*c)*cot(b*x+a))/(2*I+2*c)*ln((-1+I*c)*cot(b
*x+a)+c+I)*c^2-2*I*arctan(-c+(-1+I*c)*cot(b*x+a))/(2*I+2*c)*ln((-1+I*c)*co
t(b*x+a)+c+I)*c+arctan(-c+(-1+I*c)*cot(b*x+a))/(2*I+2*c)*ln((-1+I*c)*cot(b
*x+a)+c+I)+arctan(-c+(-1+I*c)*cot(b*x+a))/(2*I+2*c)*ln(-I+(-1+I*c)*cot(b*x
+a)-c)*c^2+2*I*arctan(-c+(-1+I*c)*cot(b*x+a))/(2*I+2*c)*ln(-I+(-1+I*c)*cot
(b*x+a)-c)*c-arctan(-c+(-1+I*c)*cot(b*x+a))/(2*I+2*c)*ln(-I+(-1+I*c)*cot(b
*x+a)-c)-(-1+I*c)^2*(-1/2/(I+c)*(1/2*I*(dilog(-1/2*I*((-1+I*c)*cot(b*x+a)-
c+I))+ln(-I+(-1+I*c)*cot(b*x+a)-c)*ln(-1/2*I*((-1+I*c)*cot(b*x+a)-c+I)))-1
/4*I*ln(-I+(-1+I*c)*cot(b*x+a)-c)^2)+1/2/(I+c)*(-1/2*I*(dilog((-I+(-1+I*c)
*cot(b*x+a)-c)/(-2*I-2*c))+ln((-1+I*c)*cot(b*x+a)+c+I)*ln((-I+(-1+I*c)*cot
(b*x+a)-c)/(-2*I-2*c)))+1/2*I*(dilog(-1/2*((-1+I*c)*cot(b*x+a)-c+I)/c)+ln(
(-1+I*c)*cot(b*x+a)+c+I)*ln(-1/2*((-1+I*c)*cot(b*x+a)-c+I)/c))))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.32

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx$$

$$= \frac{2b^2x^2 + 2ibx \log\left(-\frac{(c+i)e^{(2ibx+2ia)}}{ce^{(2ibx+2ia)}+i}\right) - 2a^2 - 2(-ibx - ia) \log(-ice^{(2ibx+2ia)} + 1) - 2ia \log\left(\frac{ce^{(2ibx+2ia)}}{c}\right)}{4b}$$

```
input integrate(-arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="fricas")
```

output

```
1/4*(2*b^2*x^2 + 2*I*b*x*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x +
2*I*a) + I)) - 2*a^2 - 2*(-I*b*x - I*a)*log(-I*c*e^(2*I*b*x + 2*I*a) + 1)
- 2*I*a*log((c*e^(2*I*b*x + 2*I*a) + I)/c) + dilog(I*c*e^(2*I*b*x + 2*I*a)
))/b
```

Sympy [F(-2)]

Exception generated.

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input

```
integrate(-atan(-c-(1-I*c)*cot(b*x+a)),x)
```

output

```
Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) - 1 o
f type <class 'sympy.core.add.Add'> to QQ_I[b,_t0,exp(I*a)]
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 458 vs. $2(64) = 128$.

Time = 0.15 (sec) , antiderivative size = 458, normalized size of antiderivative = 5.39

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = \text{Too large to display}$$

input

```
integrate(-arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="maxima")
```

output

```
-1/8*((I*c - 1)*(4*I*(b*x + a)*log(-2*(I*c^2 - (c^2 + 1)*tan(b*x + a) - 2*c - I)/(-2*I*c^2 + 2*(c^2 + 1)*tan(b*x + a) - 2*I))/(I*c - 1) - I*(4*(b*x + a)*(log(-I*c^2 + (c^2 + 1)*tan(b*x + a) + 2*c + I) - log(-I*c^2 + (c^2 + 1)*tan(b*x + a) - I)) - 2*I*log(-I*c^2 + (c^2 + 1)*tan(b*x + a) + 2*c + I)*log(-1/2*((I*c + 1)*tan(b*x + a) + c + I)/c + 1) + 2*I*log(-I*c^2 + (c^2 + 1)*tan(b*x + a) + 2*c + I)*log(tan(b*x + a) - I) - 2*I*log(1/2*(c - I)*tan(b*x + a) - 1/2*I*c + 1/2)*log(tan(b*x + a) - I) - I*log(tan(b*x + a) - I)^2 - 2*I*log(c^2 + 1)*log(I*tan(b*x + a) + 1) + 2*I*log(tan(b*x + a) - I)*log(-1/2*I*tan(b*x + a) + 1/2) + 2*I*log(c^2 + 1)*log(-I*tan(b*x + a) + 1) - 2*I*dilog(-1/2*(c - I)*tan(b*x + a) + 1/2*I*c + 1/2) - 2*I*dilog(1/2*((I*c + 1)*tan(b*x + a) + c + I)/c) + 2*I*dilog(1/2*I*tan(b*x + a) + 1/2))/(I*c - 1)) - 8*(b*x + a)*arctan(c + (-I*c + 1)/tan(b*x + a)) + 4*(-I*b*x - I*a)*log(-2*(I*c^2 - (c^2 + 1)*tan(b*x + a) - 2*c - I)/(-2*I*c^2 + 2*(c^2 + 1)*tan(b*x + a) - 2*I)))/b
```

Giac [F]

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = \int -\arctan(-(-ic + 1) \cot(bx + a) - c) dx$$

input

```
integrate(-arctan(-c-(1-I*c)*cot(b*x+a)),x, algorithm="giac")
```

output

```
integrate(-arctan(-(-I*c + 1)*cot(b*x + a) - c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = \int \operatorname{atan}(c - \cot(a + bx) (-1 + c1i)) dx$$

input

```
int(atan(c - cot(a + b*x)*(c*1i - 1)),x)
```

output

```
int(atan(c - cot(a + b*x)*(c*1i - 1)), x)
```

Reduce [F]

$$\int \arctan(c + (1 - ic) \cot(a + bx)) dx = - \left(\int \operatorname{atan}(\cot(bx + a) ci - \cot(bx + a) - c) dx \right)$$

input `int(-atan(-c-(1-I*c)*cot(b*x+a)),x)`

output `- int(atan(cot(a + b*x)*c*i - cot(a + b*x) - c),x)`

3.68 $\int \frac{\arctan(c+(1-ic)\cot(a+bx))}{x} dx$

Optimal result	542
Mathematica [N/A]	542
Rubi [N/A]	543
Maple [N/A]	543
Fricas [N/A]	544
Sympy [F(-1)]	544
Maxima [F(-2)]	544
Giac [N/A]	545
Mupad [N/A]	545
Reduce [N/A]	546

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\arctan(c + (1 - ic)\cot(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + (1 - ic)\cot(a + bx))}{x}, x\right)$$

output `Defer(Int)(-arctan(-c-(1-I*c)*cot(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 0.47 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(c + (1 - ic)\cot(a + bx))}{x} dx = \int \frac{\arctan(c + (1 - ic)\cot(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + (1 - I*c)*Cot[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + (1 - I*c)*Cot[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx$$

input `Int[ArcTan[c + (1 - I*c)*Cot[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.86 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int -\frac{\arctan(-c - (-ic + 1) \cot(bx + a))}{x} dx$$

input `int(-arctan(-c-(1-I*c)*cot(b*x+a))/x,x)`

output `int(-arctan(-c-(1-I*c)*cot(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx = \int -\frac{\arctan(-(-ic + 1) \cot(bx + a) - c)}{x} dx$$

input `integrate(-arctan(-c-(1-I*c)*cot(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c + I)*e^(2*I*b*x + 2*I*a)/(c*e^(2*I*b*x + 2*I*a) + I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(-atan(-c-(1-I*c)*cot(b*x+a))/x,x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx = \text{Exception raised: ValueError}$$

input `integrate(-arctan(-c-(1-I*c)*cot(b*x+a))/x,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(c-1>0)', see `assume?` for more
details)Is
```

Giac [N/A]

Not integrable

Time = 1.54 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx = \int -\frac{\arctan(-(-ic + 1) \cot(bx + a) - c)}{x} dx$$

input

```
integrate(-arctan(-c-(1-I*c)*cot(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(-arctan(-(-I*c + 1)*cot(b*x + a) - c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.88 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c - \cot(a + bx) (-1 + ci))}{x} dx$$

input

```
int(atan(c - cot(a + b*x)*(c*1i - 1))/x,x)
```

output

```
int(atan(c - cot(a + b*x)*(c*1i - 1))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.43

$$\int \frac{\arctan(c + (1 - ic) \cot(a + bx))}{x} dx$$

$$= - \left(\int \frac{\operatorname{atan}(\cot(bx + a) ci - \cot(bx + a) - c)}{x} dx \right)$$

input `int(-atan(-c-(1-I*c)*cot(b*x+a))/x,x)`output `- int(atan(cot(a + b*x)*c*i - cot(a + b*x) - c)/x,x)`

3.69 $\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx$

Optimal result	547
Mathematica [A] (verified)	548
Rubi [A] (verified)	548
Maple [C] (warning: unable to verify)	551
Fricas [A] (verification not implemented)	552
Sympy [F(-2)]	553
Maxima [B] (verification not implemented)	553
Giac [F]	554
Mupad [F(-1)]	554
Reduce [F]	554

Optimal result

Integrand size = 21, antiderivative size = 155

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx = -\frac{bx^4}{12} + \frac{1}{3}x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{6}ix^3 \log(1 + ice^{2ia+2ibx}) - \frac{x^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{4b} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{4b^2} + \frac{\text{PolyLog}(4, -ice^{2ia+2ibx})}{8b^3}$$

output

```
-1/12*b*x^4-1/3*x^3*arctan(-c+(1+I*c)*cot(b*x+a))-1/6*I*x^3*ln(1+I*c*exp(2
*I*a+2*I*b*x))-1/4*x^2*polylog(2,-I*c*exp(2*I*a+2*I*b*x))/b-1/4*I*x*polylo
g(3,-I*c*exp(2*I*a+2*I*b*x))/b^2+1/8*polylog(4,-I*c*exp(2*I*a+2*I*b*x))/b^
3
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.90

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx = \frac{1}{3} x^3 \arctan(c + (-1 - ic) \cot(a + bx)) - \frac{4ib^3 x^3 \log\left(1 - \frac{ie^{-2i(a+bx)}}{c}\right) - 6b^2 x^2 \operatorname{PolyLog}\left(2, \frac{ie^{-2i(a+bx)}}{c}\right) + 6ibx \operatorname{PolyLog}\left(3, \frac{ie^{-2i(a+bx)}}{c}\right) + 3 \operatorname{PolyLog}\left(4, \frac{ie^{-2i(a+bx)}}{c}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTan[c + (-1 - I*c)*Cot[a + b*x]], x]
```

output

```
(x^3*ArcTan[c + (-1 - I*c)*Cot[a + b*x]])/3 - ((4*I)*b^3*x^3*Log[1 - I/(c*E^((2*I)*(a + b*x)))] - 6*b^2*x^2*PolyLog[2, I/(c*E^((2*I)*(a + b*x)))] + (6*I)*b*x*PolyLog[3, I/(c*E^((2*I)*(a + b*x)))] + 3*PolyLog[4, I/(c*E^((2*I)*(a + b*x)))])/(24*b^3)
```

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.28, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5696, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx$$

$$\downarrow \text{5696}$$

$$\frac{1}{3} x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3} ib \int \frac{x^3}{i - ce^{2ia+2ibx}} dx$$

$$\downarrow \text{2615}$$

$$\frac{1}{3} x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3} ib \left(-ic \int \frac{e^{2ia+2ibx} x^3}{i - ce^{2ia+2ibx}} dx - \frac{ix^4}{4} \right)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \int x^2 \log(ie^{2ia+2ibx}c + 1) dx}{2bc} \right) - \frac{ix^4}{4} \right)$$

↓ 3011

$$\frac{1}{3}x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \int x \text{PolyLog}(2, -ice^{2ia+2ibx}) dx}{b} \right)}{2bc} \right) - \frac{ix^4}{4} \right)$$

↓ 7163

$$\frac{1}{3}x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int \text{PolyLog}(3, -ice^{2ia+2ibx}) dx}{2b} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} \right) \right)$$

↓ 2720

$$\frac{1}{3}x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\int e^{-2ia-2ibx} \text{PolyLog}(3, -ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{b} \right)}{b} \right)}{2bc} \right) \right)$$

↓ 7143

$$\frac{1}{3}x^3 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{3}ib \left(-ic \left(\frac{ix^3 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{3i \left(\frac{ix^2 \text{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \left(\frac{\text{PolyLog}(4, -ice^{2ia+2ibx})}{4b^2} - \frac{ix \text{PolyLog}(3, -ice^{2ia+2ibx})}{2b} \right)}{b} \right)}{2bc} \right) \right)$$

input `Int[x^2*ArcTan[c + (-1 - I*c)*Cot[a + b*x]],x]`

output `(x^3*ArcTan[c - (1 + I*c)*Cot[a + b*x]])/3 - (I/3)*b*((-1/4*I)*x^4 - I*c*((I/2)*x^3*Log[1 + I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) - (((3*I)/2)*((I/2)*x^2*PolyLog[2, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/b - (I*(((1/2*I)*x*PolyLog[3, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/b + PolyLog[4, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/b))/(b*c))`

Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_.))))^(n_.)]*((f_.) + (g_.)*(x_.))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5696

```
Int[ArcTan[(c_.) + Cot[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_)^(m_.
), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Cot[a + b*x]]/(f*(m +
1))), x] - Simp[I*(b/(f*(m + 1))) Int[(e + f*x)^(m + 1)/(c - I*d - c*E^(2
*I*a + 2*I*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && E
qq[(c - I*d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_)^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.98 (sec) , antiderivative size = 1488, normalized size of antiderivative = 9.60

method	result	size
risch	Expression too large to display	1488

input

```
int(-x^2*arctan(-c-(-1-I*c)*cot(b*x+a)),x,method=_RETURNVERBOSE)
```


output

```

1/6*I/b^3*a^3*ln(-exp(2*I*(b*x+a))*c+I)-1/3*I*x^3*ln(exp(I*(b*x+a)))-1/12*
I*(2*I*Pi+2*ln(-I+c)-I*Pi*csgn(I*exp(I*(b*x+a))))^2*csgn(I*exp(2*I*(b*x+a))
)+2*I*Pi*csgn(I*exp(I*(b*x+a)))*csgn(I*exp(2*I*(b*x+a)))^2-I*Pi*csgn(I*exp
(2*I*(b*x+a)))^3-I*Pi*csgn((exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))-1))^2+
I*Pi*csgn(exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))-1))^3-I*Pi*csgn(exp(2*
I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))-1))^2-I*Pi*csgn(I/(exp(2*I*(b*x+a))-1)
*(-I+c))*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I*(-I+c))-I*Pi*csgn(I/(exp(2*I*
(b*x+a))-1)*(-I+c))*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a))-1))*c
sgn(I*exp(2*I*(b*x+a)))+I*Pi*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a)
))-1))*csgn(I/(exp(2*I*(b*x+a))-1))*csgn(I*(exp(2*I*(b*x+a))*c-I))-I*Pi*cs
gn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))-1))*csgn((exp(2*I*(b*x+a))*c
-I)/(exp(2*I*(b*x+a))-1))^2-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c-I)/(exp(2*I*(b
*x+a))-1))^2*csgn(I/(exp(2*I*(b*x+a))-1))-I*Pi*csgn(I*(exp(2*I*(b*x+a))*c-
I)/(exp(2*I*(b*x+a))-1))^2*csgn(I*(exp(2*I*(b*x+a))*c-I))+I*Pi*csgn(I*(exp
(2*I*(b*x+a))*c-I)/(exp(2*I*(b*x+a))-1))*csgn((exp(2*I*(b*x+a))*c-I)/(exp(
2*I*(b*x+a))-1))+I*Pi*csgn(I/(exp(2*I*(b*x+a))-1)*(-I+c))^2*csgn(I/(exp(2*
I*(b*x+a))-1))+I*Pi*csgn(I/(exp(2*I*(b*x+a))-1)*(-I+c))^2*csgn(I*(-I+c))+I
*Pi*csgn(I/(exp(2*I*(b*x+a))-1)*(-I+c))*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(ex
p(2*I*(b*x+a))-1))^2+I*Pi*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(exp(2*I*(b*x+a)
-1))^2*csgn(I*exp(2*I*(b*x+a)))-I*Pi*csgn(I*exp(2*I*(b*x+a))*(-I+c)/(ex...

```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.07

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx =$$

$$\frac{2b^4x^4 - 4ib^3x^3 \log\left(-\frac{ce^{(2ibx+2ia)-i}e^{(-2ibx-2ia)}}{c-i}\right) + 6b^2x^2 \operatorname{Li}_2(-ice^{(2ibx+2ia)}) - 2a^4 - 4ia^3 \log\left(\frac{ce^{(2ibx+2ia)}}{c}\right)}{1}$$

input

```
integrate(-x^2*arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="fricas")
```

output

```

-1/24*(2*b^4*x^4 - 4*I*b^3*x^3*log(-(c*e^(2*I*b*x + 2*I*a) - I)*e^(-2*I*b*x
x - 2*I*a)/(c - I)) + 6*b^2*x^2*dilog(-I*c*e^(2*I*b*x + 2*I*a)) - 2*a^4 -
4*I*a^3*log((c*e^(2*I*b*x + 2*I*a) - I)/c) + 6*I*b*x*polylog(3, -I*c*e^(2*
I*b*x + 2*I*a)) + 4*(I*b^3*x^3 + I*a^3)*log(I*c*e^(2*I*b*x + 2*I*a) + 1) -
3*polylog(4, -I*c*e^(2*I*b*x + 2*I*a)))/b^3

```

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(-x**2*atan(-c-(-1-I*c)*cot(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) - 1 of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 311 vs. $2(109) = 218$.

Time = 0.06 (sec) , antiderivative size = 311, normalized size of antiderivative = 2.01

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx$$

$$= \frac{4 \left((bx+a)^3 - 3(bx+a)^2 a + 3(bx+a)a^2 \right) \arctan((-ic-1) \cot(bx+a)+c)}{b^2} - \frac{(-3i(bx+a)^4 + 12i(bx+a)^3 a - 18i(bx+a)^2 a^2 - 2(-4i(bx+a)^3 + 9i(bx+a)^2 a - 9i(bx+a)a^2) \arctan_2(c \cos(2bx+2a), -c \sin(2bx+2a) + 1) - 3(4I(bx+a)^2 - 6I(bx+a)a + 3Ia^2) \operatorname{dilog}(-Ic e^{(2Ibx+2Ia)}) + (4(bx+a)^3 - 9(bx+a)^2 a + 9(bx+a)a^2) \log(c^2 \cos(2bx+2a)^2 + c^2 \sin(2bx+2a)^2 - 2c \sin(2bx+2a) + 1) + 3(4bx+a) \operatorname{polylog}(3, -Ic e^{(2Ibx+2Ia)}) + 6I \operatorname{polylog}(4, -Ic e^{(2Ibx+2Ia)})}{(Ic+1)(b^2(c-I))} / b$$

input `integrate(-x^2*arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="maxima")`

output `1/12*(4*((b*x + a)^3 - 3*(b*x + a)^2*a + 3*(b*x + a)*a^2)*arctan((-I*c - 1)*cot(b*x + a) + c)/b^2 - (-3*I*(b*x + a)^4 + 12*I*(b*x + a)^3*a - 18*I*(b*x + a)^2*a^2 - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a - 9*I*(b*x + a)*a^2)*arctan_2(c*cos(2*b*x + 2*a), -c*sin(2*b*x + 2*a) + 1) - 3*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2)*dilog(-I*c*e^(2*I*b*x + 2*I*a)) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(b*x + a)*a^2)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 - 2*c*sin(2*b*x + 2*a) + 1) + 3*(4*b*x + a)*polylog(3, -I*c*e^(2*I*b*x + 2*I*a)) + 6*I*polylog(4, -I*c*e^(2*I*b*x + 2*I*a))* (I*c + 1)/(b^2*(c - I))/b`

Giac [F]

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx$$

$$= \int -x^2 \arctan(-(-ic - 1) \cot(bx + a) - c) dx$$

input `integrate(-x^2*arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-x^2*arctan(-(-I*c - 1)*cot(b*x + a) - c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx = \int x^2 \operatorname{atan}(c - \cot(a + bx) (1 + ci)) dx$$

input `int(x^2*atan(c - cot(a + b*x)*(c*1i + 1)),x)`

output `int(x^2*atan(c - cot(a + b*x)*(c*1i + 1)), x)`

Reduce [F]

$$\int x^2 \arctan(c + (-1 - ic) \cot(a + bx)) dx = - \left(\int \operatorname{atan}(\cot(bx + a) ci + \cot(bx + a) - c) x^2 dx \right)$$

input `int(-x^2*atan(-c-(-1-I*c)*cot(b*x+a)),x)`

output `- int(atan(cot(a + b*x)*c*i + cot(a + b*x) - c)*x**2,x)`

3.70 $\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx$

Optimal result	555
Mathematica [A] (verified)	556
Rubi [A] (verified)	556
Maple [C] (warning: unable to verify)	559
Fricas [A] (verification not implemented)	560
Sympy [F(-2)]	560
Maxima [B] (verification not implemented)	561
Giac [F]	561
Mupad [F(-1)]	562
Reduce [F]	562

Optimal result

Integrand size = 19, antiderivative size = 124

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = -\frac{bx^3}{6} + \frac{1}{2}x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{4}ix^2 \log(1 + ice^{2ia+2ibx}) - \frac{x \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{4b} - \frac{i \operatorname{PolyLog}(3, -ice^{2ia+2ibx})}{8b^2}$$

output

```
-1/6*b*x^3-1/2*x^2*arctan(-c+(1+I*c)*cot(b*x+a))-1/4*I*x^2*ln(1+I*c*exp(2*I*a+2*I*b*x))-1/4*x*polylog(2,-I*c*exp(2*I*a+2*I*b*x))/b-1/8*I*polylog(3,-I*c*exp(2*I*a+2*I*b*x))/b^2
```

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.89

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = \frac{1}{2} x^2 \arctan(c + (-1 - ic) \cot(a + bx)) - \frac{i \left(2b^2 x^2 \log \left(1 - \frac{ie^{-2i(a+bx)}}{c} \right) + 2ibx \operatorname{PolyLog} \left(2, \frac{ie^{-2i(a+bx)}}{c} \right) + \operatorname{PolyLog} \left(3, \frac{ie^{-2i(a+bx)}}{c} \right) \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c + (-1 - I*c)*Cot[a + b*x]],x]
```

output

```
(x^2*ArcTan[c + (-1 - I*c)*Cot[a + b*x]])/2 - ((I/8)*(2*b^2*x^2*Log[1 - I/(c*E^((2*I)*(a + b*x)))] + (2*I)*b*x*PolyLog[2, I/(c*E^((2*I)*(a + b*x)))] + PolyLog[3, I/(c*E^((2*I)*(a + b*x)))]))/b^2
```

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.26, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {5696, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x \arctan(c + (-1 - ic) \cot(a + bx)) dx \\ & \quad \downarrow \text{5696} \\ & \frac{1}{2} x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2} ib \int \frac{x^2}{i - ce^{2ia+2ibx}} dx \\ & \quad \downarrow \text{2615} \\ & \frac{1}{2} x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2} ib \left(-ic \int \frac{e^{2ia+2ibx} x^2}{i - ce^{2ia+2ibx}} dx - \frac{ix^3}{3} \right) \\ & \quad \downarrow \text{2620} \end{aligned}$$

$$\frac{1}{2}x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \int x \log(ie^{2ia+2ibx}c + 1) dx}{bc} \right) - \frac{ix^3}{3} \right)$$

↓ 3011

$$\frac{1}{2}x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \left(\frac{ix \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{i \int \operatorname{PolyLog}(2, -ice^{2ia+2ibx}) dx}{2b} \right)}{bc} \right) - \frac{ix^3}{3} \right)$$

↓ 2720

$$\frac{1}{2}x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \left(\frac{ix \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{\int e^{-2ia-2ibx} \operatorname{PolyLog}(2, -ice^{2ia+2ibx}) de^{2ia+2ibx}}{4b^2} \right)}{bc} \right) - \frac{ix^3}{3} \right)$$

↓ 7143

$$\frac{1}{2}x^2 \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2}ib \left(-ic \left(\frac{ix^2 \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \left(\frac{ix \operatorname{PolyLog}(2, -ice^{2ia+2ibx})}{2b} - \frac{\operatorname{PolyLog}(3, -ice^{2ia+2ibx})}{4b^2} \right)}{bc} \right) - \frac{ix^3}{3} \right)$$

input `Int[x*ArcTan[c + (-1 - I*c)*Cot[a + b*x]],x]`

output `(x^2*ArcTan[c - (1 + I*c)*Cot[a + b*x]]/2 - (I/2)*b*((-1/3*I)*x^3 - I*c*((I/2)*x^2*Log[1 + I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) - (I*((I/2)*x*PolyLog[2, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/b - PolyLog[3, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2)))/(b*c))`

Definitions of rubi rules used

rule 2615 $\text{Int}[\frac{(c + d x)^m}{(a + (b (F^{g(e + f x)}))^{n+1})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\frac{(c + d x)^{m+1}}{a d (m+1)}, x] - \text{Simp}[\frac{b}{a} \text{Int}[(c + d x)^m (F^{g(e + f x)})^n / (a + b (F^{g(e + f x)})^n)], x], x] /;$ $\text{FreeQ}\{F, a, b, c, d, e, f, g, n, x\} \ \&\& \ \text{IGtQ}[m, 0]$

rule 2620 $\text{Int}[\frac{(F^{g(e + f x)})^n (c + d x)^m}{(a + (b (F^{g(e + f x)}))^{n+1})}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\frac{(c + d x)^m (b f g n \text{Log}[F]) \text{Log}[1 + b (F^{g(e + f x)})^n / a]}{(c + d x)^m (b f g n \text{Log}[F])} \text{Int}[(c + d x)^{m-1} \text{Log}[1 + b (F^{g(e + f x)})^n / a]], x], x] /;$ $\text{FreeQ}\{F, a, b, c, d, e, f, g, n, x\} \ \&\& \ \text{IGtQ}[m, 0]$

rule 2720 $\text{Int}[u, x_{\text{Symbol}}] \rightarrow \text{With}\{v = \text{FunctionOfExponential}[u, x]\}, \text{Simp}[v/D[v, x] \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /;$ $\text{FunctionOfExponentialQ}[u, x] \ \&\& \ \text{!MatchQ}[u, (w) * (a) * (v)^n]^m /;$ $\text{FreeQ}\{a, m, n, x\} \ \&\& \ \text{IntegerQ}[m * n] \ \&\& \ \text{!MatchQ}[u, E^{(c) * (a) + (b) * x}] * (F)[v] /;$ $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{InverseFunctionQ}[F[x]]$

rule 3011 $\text{Int}[\text{Log}[1 + (e) * (F^{(c) * (a) + (b) * x})^n] * ((f) + (g) * (x))^m], x_{\text{Symbol}}] \rightarrow \text{Simp}[\frac{(-f + g x)^m * (\text{PolyLog}[2, (-e) * (F^{c(a + b x)})^n])}{(b c n \text{Log}[F])}, x] + \text{Simp}[g * (m / (b c n \text{Log}[F])) \text{Int}[(f + g x)^{m-1} * \text{PolyLog}[2, (-e) * (F^{c(a + b x)})^n]], x], x] /;$ $\text{FreeQ}\{F, a, b, c, e, f, g, n, x\} \ \&\& \ \text{GtQ}[m, 0]$

rule 5696 $\text{Int}[\text{ArcTan}[(c) + \text{Cot}[(a) + (b) * (x)] * (d)] * ((e) + (f) * (x))^m], x_{\text{Symbol}}] \rightarrow \text{Simp}[(e + f x)^{m+1} * (\text{ArcTan}[c + d \text{Cot}[a + b x]] / (f * (m + 1))), x] - \text{Simp}[I * (b / (f * (m + 1))) \text{Int}[(e + f x)^{m+1} / (c - I * d - c * E^{(2 * I * a + 2 * I * b * x)}), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{EqQ}[(c - I * d)^2, -1]$

rule 7143 $\text{Int}[\text{PolyLog}[n, (c) * (a) + (b) * (x)]^p / ((d) + (e) * (x)), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c * (a + b x)^p] / (e * p), x] /;$ $\text{FreeQ}\{a, b, c, d, e, n, p, x\} \ \&\& \ \text{EqQ}[b * d, a * e]$

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.54 (sec) , antiderivative size = 1453, normalized size of antiderivative = 11.72

method	result	size
risch	Expression too large to display	1453

input `int(-x*arctan(-c-(-1-I*c)*cot(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/8*I*x^2*(2*I*Pi+2*\ln(-I+c)-I*Pi*csgn(I*\exp(I*(b*x+a)))^2*csgn(I*\exp(2*I \\
 & *(b*x+a))) + 2*I*Pi*csgn(I*\exp(I*(b*x+a)))*csgn(I*\exp(2*I*(b*x+a)))^2 - I*Pi*c \\
 & sgn(I*\exp(2*I*(b*x+a)))^3 - I*Pi*csgn((\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a) \\
 &))-1))^2 + I*Pi*csgn(\exp(2*I*(b*x+a))*(-I+c)/(\exp(2*I*(b*x+a))-1))^3 - I*Pi*cs \\
 & gn(\exp(2*I*(b*x+a))*(-I+c)/(\exp(2*I*(b*x+a))-1))^2 - I*Pi*csgn(I/(\exp(2*I*(b \\
 & *x+a))-1)*(-I+c))*csgn(I/(\exp(2*I*(b*x+a))-1))*csgn(I*(-I+c)) - I*Pi*csgn(I/ \\
 & (\exp(2*I*(b*x+a))-1)*(-I+c))*csgn(I*\exp(2*I*(b*x+a))*(-I+c)/(\exp(2*I*(b*x+ \\
 & a))-1))*csgn(I*\exp(2*I*(b*x+a))) + I*Pi*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2 \\
 & *I*(b*x+a))-1))*csgn(I/(\exp(2*I*(b*x+a))-1))*csgn(I*(\exp(2*I*(b*x+a))*c-I) \\
 &) - I*Pi*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))-1))*csgn((\exp(2*I*(\\
 & b*x+a))*c-I)/(\exp(2*I*(b*x+a))-1))^2 - I*Pi*csgn(I*(\exp(2*I*(b*x+a))*c-I)/(e \\
 & xp(2*I*(b*x+a))-1))^2*csgn(I/(\exp(2*I*(b*x+a))-1)) - I*Pi*csgn(I*(\exp(2*I*(b \\
 & *x+a))*c-I)/(\exp(2*I*(b*x+a))-1))^2*csgn(I*(\exp(2*I*(b*x+a))*c-I)) + I*Pi*cs \\
 & gn(I*(\exp(2*I*(b*x+a))*c-I)/(\exp(2*I*(b*x+a))-1))*csgn((\exp(2*I*(b*x+a))*c \\
 & -I)/(\exp(2*I*(b*x+a))-1)) + I*Pi*csgn(I/(\exp(2*I*(b*x+a))-1)*(-I+c))^2*csgn(\\
 & I/(\exp(2*I*(b*x+a))-1)) + I*Pi*csgn(I/(\exp(2*I*(b*x+a))-1)*(-I+c))^2*csgn(I* \\
 & (-I+c)) + I*Pi*csgn(I/(\exp(2*I*(b*x+a))-1)*(-I+c))*csgn(I*\exp(2*I*(b*x+a))* \\
 & (-I+c)/(\exp(2*I*(b*x+a))-1))^2 + I*Pi*csgn(I*\exp(2*I*(b*x+a))*(-I+c)/(\exp(2*I \\
 & *(b*x+a))-1))^2*csgn(I*\exp(2*I*(b*x+a))) - I*Pi*csgn(I*\exp(2*I*(b*x+a))*(-I+ \\
 & c)/(\exp(2*I*(b*x+a))-1))*csgn(\exp(2*I*(b*x+a))*(-I+c)/(\exp(2*I*(b*x+a))\dots
 \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.16

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = \frac{4b^3x^3 - 6ib^2x^2 \log\left(-\frac{ce^{(2ibx+2ia)} - i}{c-i} e^{(-2ibx-2ia)}\right) + 4a^3 + 6bx \operatorname{Li}_2(-i ce^{(2ibx+2ia)}) + 6ia^2 \log\left(\frac{ce^{(2ibx+2ia)}}{c}\right)}{24b^2}$$

input `integrate(-x*arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="fricas")`

output `-1/24*(4*b^3*x^3 - 6*I*b^2*x^2*log(-(c*e^(2*I*b*x + 2*I*a) - I)*e^(-2*I*b*x - 2*I*a)/(c - I)) + 4*a^3 + 6*b*x*dilog(-I*c*e^(2*I*b*x + 2*I*a)) + 6*I*a^2*log((c*e^(2*I*b*x + 2*I*a) - I)/c) + 6*(I*b^2*x^2 - I*a^2)*log(I*c*e^(2*I*b*x + 2*I*a) + 1) + 3*I*polylog(3, -I*c*e^(2*I*b*x + 2*I*a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(-x*atan(-c-(-1-I*c)*cot(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) - 1 of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 219 vs. $2(87) = 174$.

Time = 0.07 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.77

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx$$

$$= \frac{6((bx+a)^2 - 2(bx+a)a) \arctan((-ic-1) \cot(bx+a)+c)}{b} - \frac{(-4i(bx+a)^3 + 12i(bx+a)^2a - 6ibx \operatorname{Li}_2(-ice^{(2ibx+2ia)}) - 6(-i(bx+a)^2 + 2i(bx+a)a) \log(c^2 \cos(2bx+2a) + c^2 \sin(2bx+2a)^2 - 2c \sin(2bx+2a) + 1) + 3p \operatorname{olylog}(3, -Ic * e^{(2I*bx+2I*a)})) * (Ic + 1) / (b * (c - I)))}{b}$$

input `integrate(-x*arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="maxima")`

output `1/12*(6*((b*x + a)^2 - 2*(b*x + a)*a)*arctan((-I*c - 1)*cot(b*x + a) + c)/b - (-4*I*(b*x + a)^3 + 12*I*(b*x + a)^2*a - 6*I*b*x*dilog(-I*c*e^(2*I*b*x + 2*I*a)) - 6*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*arctan2(c*cos(2*b*x + 2*a), -c*sin(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log(c^2*cos(2*b*x + 2*a)^2 + c^2*sin(2*b*x + 2*a)^2 - 2*c*sin(2*b*x + 2*a) + 1) + 3*polylog(3, -I*c*e^(2*I*b*x + 2*I*a)))*(I*c + 1)/(b*(c - I)))/b`

Giac [F]

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = \int -x \arctan(-(-ic - 1) \cot(bx + a) - c) dx$$

input `integrate(-x*arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-x*arctan(-(-I*c - 1)*cot(b*x + a) - c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = \int x \operatorname{atan}(c - \cot(a + bx) (1 + ci)) dx$$

input `int(x*atan(c - cot(a + b*x)*(c*1i + 1)),x)`

output `int(x*atan(c - cot(a + b*x)*(c*1i + 1)), x)`

Reduce [F]

$$\int x \arctan(c + (-1 - ic) \cot(a + bx)) dx = - \left(\int \operatorname{atan}(\cot(bx + a) ci + \cot(bx + a) - c) x dx \right)$$

input `int(-x*atan(-c-(-1-I*c)*cot(b*x+a)),x)`

output `- int(atan(cot(a + b*x)*c*i + cot(a + b*x) - c)*x,x)`

3.71 $\int \arctan(c + (-1 - ic) \cot(a + bx)) dx$

Optimal result	563
Mathematica [B] (warning: unable to verify)	563
Rubi [A] (verified)	564
Maple [B] (verified)	566
Fricas [A] (verification not implemented)	567
Sympy [F(-2)]	568
Maxima [B] (verification not implemented)	568
Giac [F]	569
Mupad [F(-1)]	569
Reduce [F]	569

Optimal result

Integrand size = 17, antiderivative size = 86

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = -\frac{bx^2}{2} + x \arctan(c - (1 + ic) \cot(a + bx)) - \frac{1}{2}ix \log(1 + ice^{2ia+2ibx}) - \frac{\text{PolyLog}(2, -ice^{2ia+2ibx})}{4b}$$

output

```
-1/2*b*x^2-x*arctan(-c+(1+I*c)*cot(b*x+a))-1/2*I*x*ln(1+I*c*exp(2*I*a+2*I*b*x))-1/4*polylog(2,-I*c*exp(2*I*a+2*I*b*x))/b
```

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 872 vs. 2(86) = 172.

Time = 5.75 (sec) , antiderivative size = 872, normalized size of antiderivative = 10.14

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = x \arctan(c + (-1 - ic) \cot(a + bx)) + \frac{ix \csc(a + bx) \left(2bx \log(2 \cos(bx)(\cos(bx) - i \sin(bx))) + i \log \left((i + \cot(a + bx))((-i + c) \cos(a + bx) + i(i + c) \sin(a + bx)) \right) \left(-2ibx - \log \left(1 - \frac{\sec(bx)((i+c) \cos(a) + (1+i))}{\dots} \right) \right)}{\dots}$$

input `Integrate[ArcTan[c + (-1 - I*c)*Cot[a + b*x]],x]`

output

```
x*ArcTan[c + (-1 - I*c)*Cot[a + b*x]] + (I*x*Csc[a + b*x]*(2*b*x*Log[2*Cos
[b*x]*(Cos[b*x] - I*Sin[b*x])] + I*Log[(Sec[b*x]*(Cos[a] - I*Sin[a])*((-I
+ c)*Cos[a + b*x] + I*(I + c)*Sin[a + b*x]))/(2*c)]*Log[1 - I*Tan[b*x]] -
I*Log[(Sec[b*x]*(Cos[a] + I*Sin[a])*((1 + I*c)*Cos[a + b*x] - (I + c)*Sin[
a + b*x]))/2]*Log[1 + I*Tan[b*x]] + I*PolyLog[2, -Cos[2*b*x] + I*Sin[2*b*x
]] + I*PolyLog[2, (Sec[b*x]*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(Cos[a + b
*x] - I*Sin[a + b*x]))/(2*c)] - I*PolyLog[2, ((Cos[a] + I*Sin[a])*((I + c)
*cos[a] + (1 + I*c)*Sin[a])*(-I + Tan[b*x]))/2])*(Cos[b*x] - I*Sin[b*x])*
(Cos[b*x] + I*Sin[b*x]))/((I + Cot[a + b*x])*((-I + c)*Cos[a + b*x] + I*(I
+ c)*Sin[a + b*x])*((-2*I)*b*x - Log[1 - (Sec[b*x]*((I + c)*Cos[a] + (1 +
I*c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]))/(2*c)] - (Log[1 - I*Tan[b*x]
]*((I + c)*Cos[a + b*x] + (1 + I*c)*Sin[a + b*x]))/((-I + c)*Cos[a + b*x]
+ I*(I + c)*Sin[a + b*x]) + (Log[1 + I*Tan[b*x]]*((I + c)*Cos[a + b*x] + (
1 + I*c)*Sin[a + b*x]))/((-I + c)*Cos[a + b*x] + I*(I + c)*Sin[a + b*x]) +
(Log[(Sec[b*x]*(Cos[a] + I*Sin[a])*((1 + I*c)*Cos[a + b*x] - (I + c)*Sin[
a + b*x]))/2]*Sec[b*x]^2)/(1 + I*Tan[b*x]) - 2*b*x*Tan[b*x] - I*Log[1 - (S
ec[b*x]*((I + c)*Cos[a] + (1 + I*c)*Sin[a])*(Cos[a + b*x] - I*Sin[a + b*x]
))/2]*Tan[b*x] + I*Log[1 - I*Tan[b*x]]*Tan[b*x] - I*Log[1 + I*Tan[b*x]
]*Tan[b*x] + (I*Log[1 - ((Cos[a] + I*Sin[a])*((I + c)*Cos[a] + (1 + I*c)*S
in[a])*(-I + Tan[b*x]))/2]*Sec[b*x]^2)/(-I + Tan[b*x]) + (I*Log[(Sec[b*...
```

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.26, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {5688, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx$$

$$\downarrow 5688$$

$$x \arctan(c - (1 + ic) \cot(a + bx)) - ib \int \frac{x}{i - ce^{2ia+2ibx}} dx$$

$$\begin{aligned}
& \downarrow 2615 \\
& x \arctan(c - (1 + ic) \cot(a + bx)) - ib \left(-ic \int \frac{e^{2ia+2ibx} x}{i - ce^{2ia+2ibx}} dx - \frac{ix^2}{2} \right) \\
& \downarrow 2620 \\
& ib \left(-ic \left(\frac{x \arctan(c - (1 + ic) \cot(a + bx)) - ix \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{i \int \log(ie^{2ia+2ibx} c + 1) dx}{2bc} \right) - \frac{ix^2}{2} \right) \\
& \downarrow 2715 \\
& ib \left(-ic \left(\frac{x \arctan(c - (1 + ic) \cot(a + bx)) - ix \log(1 + ice^{2ia+2ibx})}{2bc} - \frac{\int e^{-2ia-2ibx} \log(ie^{2ia+2ibx} c + 1) de^{2ia+2ibx}}{4b^2c} \right) - \frac{ix^2}{2} \right) \\
& \downarrow 2838 \\
& ib \left(-ic \left(\frac{x \arctan(c - (1 + ic) \cot(a + bx)) - \text{PolyLog}(2, -ice^{2ia+2ibx})}{4b^2c} + \frac{ix \log(1 + ice^{2ia+2ibx})}{2bc} \right) - \frac{ix^2}{2} \right)
\end{aligned}$$

input `Int[ArcTan[c + (-1 - I*c)*Cot[a + b*x]],x]`

output `x*ArcTan[c - (1 + I*c)*Cot[a + b*x]] - I*b*((-1/2*I)*x^2 - I*c*((I/2)*x*Log[1 + I*c*E^((2*I)*a + (2*I)*b*x)]/(b*c) + PolyLog[2, (-I)*c*E^((2*I)*a + (2*I)*b*x)]/(4*b^2*c)))`

Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

```
rule 2620 Int[((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 5688 Int[ArcTan[(c_) + Cot[(a_) + (b_)*(x_)]*(d_)], x_Symbol] := Simp[x*ArcT
an[c + d*Cot[a + b*x]], x] - Simp[I*b Int[x/(c - I*d - c*E^(2*I*a + 2*I*b
*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - I*d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 624 vs. 2(72) = 144.

Time = 1.60 (sec) , antiderivative size = 625, normalized size of antiderivative = 7.27

method	result
derivativdivides	$-\frac{\arctan(-c+(ic+1)\cot(bx+a))\ln(i+(ic+1)\cot(bx+a)-c)^2}{2i-2c} + \frac{2i\arctan(-c+(ic+1)\cot(bx+a))\ln(i+(ic+1)\cot(bx+a)-c)c}{2i-2c} + \arctan(-c+(ic+1)\cot(bx+a))\ln(i+(ic+1)\cot(bx+a)-c)$
default	$-\frac{\arctan(-c+(ic+1)\cot(bx+a))\ln(i+(ic+1)\cot(bx+a)-c)^2}{2i-2c} + \frac{2i\arctan(-c+(ic+1)\cot(bx+a))\ln(i+(ic+1)\cot(bx+a)-c)c}{2i-2c} + \arctan(-c+(ic+1)\cot(bx+a))\ln(i+(ic+1)\cot(bx+a)-c)$
risch	Expression too large to display

```
input int(-arctan(-c-(-1-I*c)*cot(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/b/(1+I*c)*(-arctan(-c+(1+I*c)*cot(b*x+a))/(2*I-2*c)*ln(I+(1+I*c)*cot(b*x+a)-c)*c^2+2*I*arctan(-c+(1+I*c)*cot(b*x+a))/(2*I-2*c)*ln(I+(1+I*c)*cot(b*x+a)-c)*c+arctan(-c+(1+I*c)*cot(b*x+a))/(2*I-2*c)*ln(I+(1+I*c)*cot(b*x+a)-c)+arctan(-c+(1+I*c)*cot(b*x+a))/(2*I-2*c)*ln(-(1+I*c)*cot(b*x+a)-c+I)*c^2-2*I*arctan(-c+(1+I*c)*cot(b*x+a))/(2*I-2*c)*ln(-(1+I*c)*cot(b*x+a)-c+I)*c-arctan(-c+(1+I*c)*cot(b*x+a))/(2*I-2*c)*ln(-(1+I*c)*cot(b*x+a)-c+I)+(1+I*c)^2*(-1/2/(I-c)*(1/4*I*ln(I+(1+I*c)*cot(b*x+a)-c)^2-1/2*I*((ln(I+(1+I*c)*cot(b*x+a)-c)-ln(-1/2*I*(I+(1+I*c)*cot(b*x+a)-c)))*ln(-1/2*I*(I-(1+I*c)*cot(b*x+a)+c))-dilog(-1/2*I*(I+(1+I*c)*cot(b*x+a)-c)))+1/2/(I-c)*(-1/2*I*(dilog(1/2*(I-(1+I*c)*cot(b*x+a)+c)/c)+ln(-(1+I*c)*cot(b*x+a)-c+I)*ln(1/2*(I-(1+I*c)*cot(b*x+a)+c)/c))+1/2*I*(dilog((-I-(1+I*c)*cot(b*x+a)+c)/(-2*I+2*c))+ln(-(1+I*c)*cot(b*x+a)-c+I)*ln((-I-(1+I*c)*cot(b*x+a)+c)/(-2*I+2*c))))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.30

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = \frac{2b^2x^2 - 2ibx \log\left(-\frac{(ce^{2ibx+2ia}-i)e^{-2ibx-2ia}}{c-i}\right) - 2a^2 + 2(ibx + ia) \log(ice^{2ibx+2ia} + 1) - 2ia \log\left(\frac{ce^{2ibx+2ia}-i}{c-i}\right)}{4b}$$

input

```
integrate(-arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="fricas")
```

output

```
-1/4*(2*b^2*x^2 - 2*I*b*x*log(-(c*e^(2*I*b*x + 2*I*a) - I)*e^(-2*I*b*x - 2*I*a)/(c - I)) - 2*a^2 + 2*(I*b*x + I*a)*log(I*c*e^(2*I*b*x + 2*I*a) + 1) - 2*I*a*log((c*e^(2*I*b*x + 2*I*a) - I)/c) + dilog(-I*c*e^(2*I*b*x + 2*I*a)))/b
```


Sympy [F(-2)]

Exception generated.

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(-atan(-c-(-1-I*c)*cot(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2*exp(2*I*a) - 1 of type <class 'sympy.core.add.Add'> to QQ_I[b,_t0,exp(I*a)]`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 458 vs. $2(63) = 126$.

Time = 0.15 (sec) , antiderivative size = 458, normalized size of antiderivative = 5.33

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = \text{Too large to display}$$

input `integrate(-arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="maxima")`

output `-1/8*((I*c + 1)*(4*I*(b*x + a)*log(-2*(I*c^2 - (c^2 + 1)*tan(b*x + a) + I)/(-2*I*c^2 + 2*(c^2 + 1)*tan(b*x + a) - 4*c + 2*I))/(I*c + 1) + I*(4*(b*x + a)*(log(-I*c^2 + (c^2 + 1)*tan(b*x + a) - 2*c + I) - log(-I*c^2 + (c^2 + 1)*tan(b*x + a) - I)) - 2*I*log(-I*c^2 + (c^2 + 1)*tan(b*x + a) - 2*c + I)*log(-1/2*((I*c - 1)*tan(b*x + a) + c - I)/c + 1) + 2*I*log(-I*c^2 + (c^2 + 1)*tan(b*x + a) - 2*c + I)*log(tan(b*x + a) - I) - 2*I*log(-1/2*(c + I)*tan(b*x + a) + 1/2*I*c + 1/2)*log(tan(b*x + a) - I) - I*log(tan(b*x + a) - I)^2 - 2*I*log(c^2 + 1)*log(I*tan(b*x + a) + 1) + 2*I*log(tan(b*x + a) - I)*log(-1/2*I*tan(b*x + a) + 1/2) + 2*I*log(c^2 + 1)*log(-I*tan(b*x + a) + 1) - 2*I*dilog(1/2*(c + I)*tan(b*x + a) - 1/2*I*c + 1/2) - 2*I*dilog(1/2*((I*c - 1)*tan(b*x + a) + c - I)/c) + 2*I*dilog(1/2*I*tan(b*x + a) + 1/2))/(I*c + 1) - 8*(b*x + a)*arctan(c + (-I*c - 1)/tan(b*x + a)) + 4*(-I*b*x - I*a)*log(-2*(I*c^2 - (c^2 + 1)*tan(b*x + a) + I)/(-2*I*c^2 + 2*(c^2 + 1)*tan(b*x + a) - 4*c + 2*I))/b`

Giac [F]

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = \int -\arctan(-(-ic - 1) \cot(bx + a) - c) dx$$

input `integrate(-arctan(-c-(-1-I*c)*cot(b*x+a)),x, algorithm="giac")`

output `integrate(-arctan(-(-I*c - 1)*cot(b*x + a) - c), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = \int \operatorname{atan}(c - \cot(a + bx) (1 + ci)) dx$$

input `int(atan(c - cot(a + b*x)*(c*1i + 1)),x)`

output `int(atan(c - cot(a + b*x)*(c*1i + 1)), x)`

Reduce [F]

$$\int \arctan(c + (-1 - ic) \cot(a + bx)) dx = - \left(\int \operatorname{atan}(\cot(bx + a) ci + \cot(bx + a) - c) dx \right)$$

input `int(-atan(-c-(-1-I*c)*cot(b*x+a)),x)`

output `- int(atan(cot(a + b*x)*c*i + cot(a + b*x) - c),x)`

3.72 $\int \frac{\arctan(c+(-1-ic)\cot(a+bx))}{x} dx$

Optimal result	570
Mathematica [N/A]	570
Rubi [N/A]	571
Maple [N/A]	571
Fricas [N/A]	572
Sympy [F(-1)]	572
Maxima [F(-2)]	572
Giac [N/A]	573
Mupad [N/A]	573
Reduce [N/A]	574

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\arctan(c + (-1 - ic)\cot(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + (-1 - ic)\cot(a + bx))}{x}, x\right)$$

output `Defer(Int)(-arctan(-c-(-1-I*c)*cot(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 0.48 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(c + (-1 - ic)\cot(a + bx))}{x} dx = \int \frac{\arctan(c + (-1 - ic)\cot(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + (-1 - I*c)*Cot[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + (-1 - I*c)*Cot[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx$$

input `Int[ArcTan[c + (-1 - I*c)*Cot[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.82 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int -\frac{\arctan(-c - (-ic - 1) \cot(bx + a))}{x} dx$$

input `int(-arctan(-c-(-1-I*c)*cot(b*x+a))/x,x)`

output `int(-arctan(-c-(-1-I*c)*cot(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.76

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx = \int -\frac{\arctan(-(-ic - 1) \cot(bx + a) - c)}{x} dx$$

input `integrate(-arctan(-c-(-1-I*c)*cot(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c*e^(2*I*b*x + 2*I*a) - I)*e^(-2*I*b*x - 2*I*a)/(c - I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(-atan(-c-(-1-I*c)*cot(b*x+a))/x,x)`

output `Timed out`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx = \text{Exception raised: ValueError}$$

input `integrate(-arctan(-c-(-1-I*c)*cot(b*x+a))/x,x, algorithm="maxima")`

output

```
Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(c-1>0)', see `assume?` for more
details)Is
```

Giac [N/A]

Not integrable

Time = 1.75 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.19

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx = \int -\frac{\arctan(-(-ic - 1) \cot(bx + a) - c)}{x} dx$$

input

```
integrate(-arctan(-c-(-1-I*c)*cot(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(-arctan(-(-I*c - 1)*cot(b*x + a) - c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c - \cot(a + bx) (1 + ci))}{x} dx$$

input

```
int(atan(c - cot(a + b*x)*(c*1i + 1))/x,x)
```

output

```
int(atan(c - cot(a + b*x)*(c*1i + 1))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.33

$$\int \frac{\arctan(c + (-1 - ic) \cot(a + bx))}{x} dx$$

$$= - \left(\int \frac{\operatorname{atan}(\cot(bx + a) ci + \cot(bx + a) - c)}{x} dx \right)$$

input `int(-atan(-c-(-1-I*c)*cot(b*x+a))/x,x)`output `- int(atan(cot(a + b*x)*c*i + cot(a + b*x) - c)/x,x)`

3.73 $\int \arctan(\sinh(x)) dx$

Optimal result	575
Mathematica [A] (verified)	575
Rubi [A] (verified)	576
Maple [A] (verified)	577
Fricas [B] (verification not implemented)	578
Sympy [F]	578
Maxima [F]	579
Giac [F]	579
Mupad [F(-1)]	579
Reduce [F]	580

Optimal result

Integrand size = 3, antiderivative size = 39

$$\int \arctan(\sinh(x)) dx = -2x \arctan(e^x) + x \arctan(\sinh(x)) \\ + i \operatorname{PolyLog}(2, -ie^x) - i \operatorname{PolyLog}(2, ie^x)$$

output

```
-2*x*arctan(exp(x))+x*arctan(sinh(x))+I*polylog(2,-I*exp(x))-I*polylog(2,I*exp(x))
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.44

$$\int \arctan(\sinh(x)) dx = x \arctan(\sinh(x)) - i(x(\log(1 - ie^x) - \log(1 + ie^x))) \\ - \operatorname{PolyLog}(2, -ie^x) + \operatorname{PolyLog}(2, ie^x)$$

input

```
Integrate[ArcTan[Sinh[x]],x]
```

output

```
x*ArcTan[Sinh[x]] - I*(x*(Log[1 - I*E^x] - Log[1 + I*E^x]) - PolyLog[2, (-I)*E^x] + PolyLog[2, I*E^x])
```


Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 1.667$, Rules used = {5726, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(\sinh(x)) dx \\
 & \quad \downarrow 5726 \\
 & x \arctan(\sinh(x)) - \int x \operatorname{sech}(x) dx \\
 & \quad \downarrow 3042 \\
 & x \arctan(\sinh(x)) - \int x \csc\left(ix + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow 4668 \\
 & i \int \log(1 - ie^x) dx - i \int \log(1 + ie^x) dx - 2x \arctan(e^x) + x \arctan(\sinh(x)) \\
 & \quad \downarrow 2715 \\
 & i \int e^{-x} \log(1 - ie^x) de^x - i \int e^{-x} \log(1 + ie^x) de^x - 2x \arctan(e^x) + x \arctan(\sinh(x)) \\
 & \quad \downarrow 2838 \\
 & -2x \arctan(e^x) + x \arctan(\sinh(x)) + i \operatorname{PolyLog}(2, -ie^x) - i \operatorname{PolyLog}(2, ie^x)
 \end{aligned}$$

input `Int[ArcTan[Sinh[x]], x]`

output `-2*x*ArcTan[E^x] + x*ArcTan[Sinh[x]] + I*PolyLog[2, (-I)*E^x] - I*PolyLog[2, I*E^x]`

Definitions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m], x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5726 `Int[ArcTan[u_], x_Symbol] :> Simp[x*ArcTan[u], x] - Int[SimplifyIntegrand[x
*(D[u, x]/(1 + u^2)), x], x] /; InverseFunctionFreeQ[u, x]`

Maple [A] (verified)

Time = 2.02 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.33

method	result	size
default	$x \arctan(\sinh(x)) - ix(\ln(1 - ie^x) - \ln(1 + ie^x)) + i \operatorname{dilog}(1 + ie^x) - i \operatorname{dilog}(1 - ie^x)$	52
parts	$x \arctan(\sinh(x)) - ix(\ln(1 - ie^x) - \ln(1 + ie^x)) + i \operatorname{dilog}(1 + ie^x) - i \operatorname{dilog}(1 - ie^x)$	52
risch	Expression too large to display	651

input `int(arctan(sinh(x)),x,method=_RETURNVERBOSE)`

output `x*arctan(sinh(x))-I*x*(ln(1-I*exp(x))-ln(1+I*exp(x)))+I*dilog(1+I*exp(x))-I*dilog(1-I*exp(x))`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(26) = 52$.

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.49

$$\int \arctan(\sinh(x)) dx = x \arctan(\sinh(x)) + i x \log(i \cosh(x) + i \sinh(x) + 1) - i x \log(-i \cosh(x) - i \sinh(x) + 1) - i \operatorname{Li}_2(i \cosh(x) + i \sinh(x)) + i \operatorname{Li}_2(-i \cosh(x) - i \sinh(x))$$

input `integrate(arctan(sinh(x)),x, algorithm="fricas")`

output `x*arctan(sinh(x)) + I*x*log(I*cosh(x) + I*sinh(x) + 1) - I*x*log(-I*cosh(x) - I*sinh(x) + 1) - I*dilog(I*cosh(x) + I*sinh(x)) + I*dilog(-I*cosh(x) - I*sinh(x))`

Sympy [F]

$$\int \arctan(\sinh(x)) dx = \int \operatorname{atan}(\sinh(x)) dx$$

input `integrate(atan(sinh(x)),x)`

output `Integral(atan(sinh(x)), x)`

Maxima [F]

$$\int \arctan(\sinh(x)) dx = \int \arctan(\sinh(x)) dx$$

input `integrate(arctan(sinh(x)),x, algorithm="maxima")`

output `x*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - 2*integrate(x*e^x/(e^(2*x) + 1), x)`

Giac [F]

$$\int \arctan(\sinh(x)) dx = \int \arctan(\sinh(x)) dx$$

input `integrate(arctan(sinh(x)),x, algorithm="giac")`

output `integrate(arctan(sinh(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(\sinh(x)) dx = \int \operatorname{atan}(\sinh(x)) dx$$

input `int(atan(sinh(x)),x)`

output `int(atan(sinh(x)), x)`

Reduce [F]

$$\int \arctan(\sinh(x)) dx = \int \operatorname{atan}(\sinh(x)) dx$$

input `int(atan(sinh(x)),x)`

output `int(atan(sinh(x)),x)`

3.74 $\int x \arctan(\sinh(x)) dx$

Optimal result	581
Mathematica [A] (verified)	581
Rubi [A] (verified)	582
Maple [C] (warning: unable to verify)	584
Fricas [A] (verification not implemented)	585
Sympy [F]	586
Maxima [F]	586
Giac [F]	586
Mupad [F(-1)]	587
Reduce [F]	587

Optimal result

Integrand size = 5, antiderivative size = 74

$$\int x \arctan(\sinh(x)) dx = -x^2 \arctan(e^x) + \frac{1}{2}x^2 \arctan(\sinh(x)) + ix \operatorname{PolyLog}(2, -ie^x) - ix \operatorname{PolyLog}(2, ie^x) - i \operatorname{PolyLog}(3, -ie^x) + i \operatorname{PolyLog}(3, ie^x)$$

output

```
-x^2*arctan(exp(x))+1/2*x^2*arctan(sinh(x))+I*x*polylog(2,-I*exp(x))-I*x*polylog(2,I*exp(x))-I*polylog(3,-I*exp(x))+I*polylog(3,I*exp(x))
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.26

$$\int x \arctan(\sinh(x)) dx = \frac{1}{2}x^2 \arctan(\sinh(x)) - \frac{1}{2}i(x^2 \log(1 - ie^x) - x^2 \log(1 + ie^x) - 2x \operatorname{PolyLog}(2, -ie^x) + 2x \operatorname{PolyLog}(2, ie^x) + 2 \operatorname{PolyLog}(3, -ie^x) - 2 \operatorname{PolyLog}(3, ie^x))$$

input

```
Integrate[x*ArcTan[Sinh[x]],x]
```

output

```
(x^2*ArcTan[Sinh[x]])/2 - (I/2)*(x^2*Log[1 - I*E^x] - x^2*Log[1 + I*E^x] -
2*x*PolyLog[2, (-I)*E^x] + 2*x*PolyLog[2, I*E^x] + 2*PolyLog[3, (-I)*E^x]
- 2*PolyLog[3, I*E^x])
```

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$, Rules used = {5728, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \arctan(\sinh(x)) dx \\
 & \quad \downarrow \text{5728} \\
 & \frac{1}{2} x^2 \arctan(\sinh(x)) - \frac{1}{2} \int x^2 \operatorname{sech}(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} x^2 \arctan(\sinh(x)) - \frac{1}{2} \int x^2 \csc\left(ix + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{4668} \\
 & \frac{1}{2} x^2 \arctan(\sinh(x)) + \frac{1}{2} \left(2i \int x \log(1 - ie^x) dx - 2i \int x \log(1 + ie^x) dx - 2x^2 \arctan(e^x) \right) \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{2} x^2 \arctan(\sinh(x)) + \\
 & \frac{1}{2} \left(-2i \left(\int \operatorname{PolyLog}(2, -ie^x) dx - x \operatorname{PolyLog}(2, -ie^x) \right) + 2i \left(\int \operatorname{PolyLog}(2, ie^x) dx - x \operatorname{PolyLog}(2, ie^x) \right) - 2x^2 \arctan(e^x) \right) \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2} x^2 \arctan(\sinh(x)) + \\
 & \frac{1}{2} \left(-2i \left(\int e^{-x} \operatorname{PolyLog}(2, -ie^x) de^x - x \operatorname{PolyLog}(2, -ie^x) \right) + 2i \left(\int e^{-x} \operatorname{PolyLog}(2, ie^x) de^x - x \operatorname{PolyLog}(2, ie^x) \right) - 2x^2 \arctan(e^x) \right) \\
 & \quad \downarrow \text{7143}
 \end{aligned}$$

$$\frac{1}{2}x^2 \arctan(\sinh(x)) + \frac{1}{2}(-2x^2 \arctan(e^x) - 2i(\text{PolyLog}(3, -ie^x) - x \text{PolyLog}(2, -ie^x)) + 2i(\text{PolyLog}(3, ie^x) - x \text{PolyLog}(2, ie^x)))$$

input `Int[x*ArcTan[Sinh[x]],x]`

output `(x^2*ArcTan[Sinh[x]])/2 + (-2*x^2*ArcTan[E^x] - (2*I)*(-(x*PolyLog[2, (-I)*E^x]) + PolyLog[3, (-I)*E^x]) + (2*I)*(-(x*PolyLog[2, I*E^x]) + PolyLog[3, I*E^x]))/2`

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^(((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5728

```
Int[((a_.) + ArcTan[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] :> Simp
p[(c + d*x)^(m + 1)*((a + b*ArcTan[u])/(d*(m + 1))), x] - Simp[b/(d*(m + 1)
) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(1 + u^2)), x], x]
/; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] &
& !FunctionOfQ[(c + d*x)^(m + 1), u, x] && FalseQ[PowerVariableExpn[u, m +
1, x]]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.30 (sec) , antiderivative size = 632, normalized size of antiderivative = 8.54

method	result	size
risch	Expression too large to display	632

input

```
int(x*arctan(sinh(x)),x,method=_RETURNVERBOSE)
```

output

```

1/2*I*x^2*ln(exp(x)+I)-1/2*I*x^2*ln(exp(x)-I)+1/2*I*x^2*ln(1+I*exp(x))+I*x
*polylog(2,-I*exp(x))-I*polylog(3,-I*exp(x))-1/8*Pi*(csgn(I*(exp(x)-I))^2*
csgn(I*(exp(x)-I)^2)-2*csgn(I*(exp(x)-I))*csgn(I*(exp(x)-I)^2)+csgn(I*(e
xp(x)-I)^2)^3+csgn(I*(exp(x)-I)^2)*csgn(I*exp(-x))*csgn(I*exp(-x)*(exp(x)-
I)^2)-csgn(I*(exp(x)-I)^2)*csgn(I*exp(-x)*(exp(x)-I)^2)^2-csgn(I*(exp(x)+I
))^2*csgn(I*(exp(x)+I)^2)+2*csgn(I*(exp(x)+I))*csgn(I*(exp(x)+I)^2)^2-csgn
(I*(exp(x)+I)^2)^3-csgn(I*(exp(x)+I)^2)*csgn(I*exp(-x))*csgn(I*exp(-x)*(ex
p(x)+I)^2)+csgn(I*(exp(x)+I)^2)*csgn(I*exp(-x)*(exp(x)+I)^2)^2-csgn(I*exp(-
x))*csgn(I*exp(-x)*(exp(x)-I)^2)^2+csgn(I*exp(-x))*csgn(I*exp(-x)*(exp(x)
+I)^2)^2-csgn(I*exp(-x)*(exp(x)+I)^2)*csgn(exp(-x)*(exp(x)+I)^2)+csgn(exp(-
x)*(exp(x)+I)^2)^2+csgn(I*exp(-x)*(exp(x)-I)^2)*csgn(exp(-x)*(exp(x)-I)^2
)+csgn(exp(-x)*(exp(x)-I)^2)^2+csgn(I*exp(-x)*(exp(x)-I)^2)^3-csgn(I*exp(-
x)*(exp(x)-I)^2)*csgn(exp(-x)*(exp(x)-I)^2)^2-csgn(I*exp(-x)*(exp(x)+I)^2
)^3+csgn(I*exp(-x)*(exp(x)+I)^2)*csgn(exp(-x)*(exp(x)+I)^2)^2-csgn(exp(-x)*
(exp(x)+I)^2)^3-csgn(exp(-x)*(exp(x)-I)^2)^3-2)*x^2-1/2*I*x^2*ln(1-I*exp(x
))-I*x*polylog(2,I*exp(x))+I*polylog(3,I*exp(x))

```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.26

$$\begin{aligned}
\int x \arctan(\sinh(x)) dx &= \frac{1}{2} x^2 \arctan(\sinh(x)) + \frac{1}{2} i x^2 \log(i \cosh(x) + i \sinh(x) + 1) \\
&\quad - \frac{1}{2} i x^2 \log(-i \cosh(x) - i \sinh(x) + 1) \\
&\quad - i x \operatorname{Li}_2(i \cosh(x) + i \sinh(x)) \\
&\quad + i x \operatorname{Li}_2(-i \cosh(x) - i \sinh(x)) \\
&\quad + i \operatorname{polylog}(3, i \cosh(x) + i \sinh(x)) \\
&\quad - i \operatorname{polylog}(3, -i \cosh(x) - i \sinh(x))
\end{aligned}$$

input

```
integrate(x*arctan(sinh(x)),x, algorithm="fricas")
```

output

```

1/2*x^2*arctan(sinh(x)) + 1/2*I*x^2*log(I*cosh(x) + I*sinh(x) + 1) - 1/2*I
*x^2*log(-I*cosh(x) - I*sinh(x) + 1) - I*x*dilog(I*cosh(x) + I*sinh(x)) +
I*x*dilog(-I*cosh(x) - I*sinh(x)) + I*polylog(3, I*cosh(x) + I*sinh(x)) -
I*polylog(3, -I*cosh(x) - I*sinh(x))

```

Sympy [F]

$$\int x \arctan(\sinh(x)) dx = \int x \operatorname{atan}(\sinh(x)) dx$$

input `integrate(x*atan(sinh(x)),x)`

output `Integral(x*atan(sinh(x)), x)`

Maxima [F]

$$\int x \arctan(\sinh(x)) dx = \int x \arctan(\sinh(x)) dx$$

input `integrate(x*arctan(sinh(x)),x, algorithm="maxima")`

output `1/2*x^2*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - integrate(x^2*e^x/(e^(2*x) + 1), x)`

Giac [F]

$$\int x \arctan(\sinh(x)) dx = \int x \arctan(\sinh(x)) dx$$

input `integrate(x*arctan(sinh(x)),x, algorithm="giac")`

output `integrate(x*arctan(sinh(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(\sinh(x)) dx = \int x \operatorname{atan}(\sinh(x)) dx$$

input `int(x*atan(sinh(x)),x)`output `int(x*atan(sinh(x)), x)`**Reduce [F]**

$$\int x \arctan(\sinh(x)) dx = \int \operatorname{atan}(\sinh(x)) x dx$$

input `int(x*atan(sinh(x)),x)`output `int(atan(sinh(x))*x,x)`

3.75 $\int x^2 \arctan(\sinh(x)) dx$

Optimal result	588
Mathematica [A] (verified)	588
Rubi [A] (verified)	589
Maple [C] (warning: unable to verify)	591
Fricas [A] (verification not implemented)	592
Sympy [F]	593
Maxima [F]	593
Giac [F]	594
Mupad [F(-1)]	594
Reduce [F]	594

Optimal result

Integrand size = 7, antiderivative size = 108

$$\int x^2 \arctan(\sinh(x)) dx = -\frac{2}{3}x^3 \arctan(e^x) + \frac{1}{3}x^3 \arctan(\sinh(x)) + ix^2 \operatorname{PolyLog}(2, -ie^x) - ix^2 \operatorname{PolyLog}(2, ie^x) - 2ix \operatorname{PolyLog}(3, -ie^x) + 2ix \operatorname{PolyLog}(3, ie^x) + 2i \operatorname{PolyLog}(4, -ie^x) - 2i \operatorname{PolyLog}(4, ie^x)$$

output

```
-2/3*x^3*arctan(exp(x))+1/3*x^3*arctan(sinh(x))+I*x^2*polylog(2,-I*exp(x))
-I*x^2*polylog(2,I*exp(x))-2*I*x*polylog(3,-I*exp(x))+2*I*x*polylog(3,I*exp(x))
+2*I*polylog(4,-I*exp(x))-2*I*polylog(4,I*exp(x))
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.12

$$\int x^2 \arctan(\sinh(x)) dx = \frac{1}{3}x^3 \arctan(\sinh(x)) - \frac{1}{3}i(x^3 \log(1 - ie^x) - x^3 \log(1 + ie^x) - 3x^2 \operatorname{PolyLog}(2, -ie^x) + 3x^2 \operatorname{PolyLog}(2, ie^x) + 6x \operatorname{PolyLog}(3, -ie^x) - 6x \operatorname{PolyLog}(3, ie^x) - 6 \operatorname{PolyLog}(4, -ie^x) + 6 \operatorname{PolyLog}(4, ie^x))$$

input `Integrate[x^2*ArcTan[Sinh[x]],x]`

output `(x^3*ArcTan[Sinh[x]])/3 - (I/3)*(x^3*Log[1 - I*E^x] - x^3*Log[1 + I*E^x] - 3*x^2*PolyLog[2, (-I)*E^x] + 3*x^2*PolyLog[2, I*E^x] + 6*x*PolyLog[3, (-I)*E^x] - 6*x*PolyLog[3, I*E^x] - 6*PolyLog[4, (-I)*E^x] + 6*PolyLog[4, I*E^x])`

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5728, 3042, 4668, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(\sinh(x)) dx \\
 & \quad \downarrow 5728 \\
 & \frac{1}{3} x^3 \arctan(\sinh(x)) - \frac{1}{3} \int x^3 \operatorname{sech}(x) dx \\
 & \quad \downarrow 3042 \\
 & \frac{1}{3} x^3 \arctan(\sinh(x)) - \frac{1}{3} \int x^3 \csc\left(ix + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow 4668 \\
 & \frac{1}{3} x^3 \arctan(\sinh(x)) + \frac{1}{3} \left(3i \int x^2 \log(1 - ie^x) dx - 3i \int x^2 \log(1 + ie^x) dx - 2x^3 \arctan(e^x) \right) \\
 & \quad \downarrow 3011 \\
 & \frac{1}{3} x^3 \arctan(\sinh(x)) + \\
 & \frac{1}{3} \left(-3i \left(2 \int x \operatorname{PolyLog}(2, -ie^x) dx - x^2 \operatorname{PolyLog}(2, -ie^x) \right) + 3i \left(2 \int x \operatorname{PolyLog}(2, ie^x) dx - x^2 \operatorname{PolyLog}(2, ie^x) \right) \right) \\
 & \quad \downarrow 7163
 \end{aligned}$$

$$\frac{1}{3} \left(-3i \left(2 \left(x \operatorname{PolyLog}(3, -ie^x) - \int \operatorname{PolyLog}(3, -ie^x) dx \right) - x^2 \operatorname{PolyLog}(2, -ie^x) \right) + 3i \left(2 \left(x \operatorname{PolyLog}(3, ie^x) \right. \right. \right.$$

↓ 2720

$$\left. \left. \left. \frac{1}{3} x^3 \arctan(\sinh(x)) + \right. \right. \right. \frac{1}{3} \left(-3i \left(2 \left(x \operatorname{PolyLog}(3, -ie^x) - \int e^{-x} \operatorname{PolyLog}(3, -ie^x) dx \right) - x^2 \operatorname{PolyLog}(2, -ie^x) \right) + 3i \left(2 \left(x \operatorname{PolyLog}(3, \right. \right. \right.$$

↓ 7143

$$\left. \left. \left. \frac{1}{3} x^3 \arctan(\sinh(x)) + \right. \right. \right. \frac{1}{3} \left(-2x^3 \arctan(e^x) - 3i \left(2 \left(x \operatorname{PolyLog}(3, -ie^x) - \operatorname{PolyLog}(4, -ie^x) \right) - x^2 \operatorname{PolyLog}(2, -ie^x) \right) + 3i \left(2 \left(x \operatorname{PolyLog}(3, \right. \right. \right.$$

input `Int[x^2*ArcTan[Sinh[x]],x]`

output `(x^3*ArcTan[Sinh[x]])/3 + (-2*x^3*ArcTan[E^x] - (3*I)*(-(x^2*PolyLog[2, (-I)*E^x]) + 2*(x*PolyLog[3, (-I)*E^x] - PolyLog[4, (-I)*E^x])) + (3*I)*(-(x^2*PolyLog[2, I*E^x]) + 2*(x*PolyLog[3, I*E^x] - PolyLog[4, I*E^x]))) / 3`

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x, x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5728 `Int[((a_.) + ArcTan[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcTan[u])/(d*(m + 1))), x] - Simp[b/(d*(m + 1)) Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(1 + u^2)), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && FalseQ[PowerVariableExpn[u, m + 1, x]]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*(F^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.91 (sec) , antiderivative size = 658, normalized size of antiderivative = 6.09

method	result	size
risch	Expression too large to display	658

input `int(x^2*arctan(sinh(x)),x,method=_RETURNVERBOSE)`

output

```

1/3*I*x^3*ln(exp(x)+I)-1/3*I*x^3*ln(exp(x)-I)+1/3*I*x^3*ln(1+I*exp(x))+I*x
^2*polylog(2,-I*exp(x))-2*I*x*polylog(3,-I*exp(x))+2*I*polylog(4,-I*exp(x)
)-1/12*Pi*(csgn(I*(exp(x)-I))^2*csgn(I*(exp(x)-I)^2)-2*csgn(I*(exp(x)-I))*
csgn(I*(exp(x)-I)^2)+csgn(I*(exp(x)-I)^2)^3+csgn(I*(exp(x)-I)^2)*csgn(I*
exp(-x))*csgn(I*exp(-x)*(exp(x)-I)^2)-csgn(I*(exp(x)-I)^2)*csgn(I*exp(-x)*
(exp(x)-I)^2)^2-csgn(I*(exp(x)+I))^2*csgn(I*(exp(x)+I)^2)+2*csgn(I*(exp(x)
+I))*csgn(I*(exp(x)+I)^2)^2-csgn(I*(exp(x)+I)^2)^3-csgn(I*(exp(x)+I)^2)*cs
gn(I*exp(-x))*csgn(I*exp(-x)*(exp(x)+I)^2)+csgn(I*(exp(x)+I)^2)*csgn(I*exp
(-x)*(exp(x)+I)^2)^2-csgn(I*exp(-x))*csgn(I*exp(-x)*(exp(x)-I)^2)^2+csgn(I
*exp(-x))*csgn(I*exp(-x)*(exp(x)+I)^2)^2-csgn(I*exp(-x)*(exp(x)+I)^2)*csgn
(exp(-x)*(exp(x)+I)^2)+csgn(exp(-x)*(exp(x)+I)^2)^2+csgn(I*exp(-x)*(exp(x)
-I)^2)*csgn(exp(-x)*(exp(x)-I)^2)+csgn(exp(-x)*(exp(x)-I)^2)^2+csgn(I*exp(
-x)*(exp(x)-I)^2)^3-csgn(I*exp(-x)*(exp(x)-I)^2)*csgn(exp(-x)*(exp(x)-I)^2
)^2-csgn(I*exp(-x)*(exp(x)+I)^2)^3+csgn(I*exp(-x)*(exp(x)+I)^2)*csgn(exp(-
x)*(exp(x)+I)^2)^2-csgn(exp(-x)*(exp(x)+I)^2)^3-csgn(exp(-x)*(exp(x)-I)^2
)^3-2)*x^3-1/3*I*x^3*ln(1-I*exp(x))-I*x^2*polylog(2,I*exp(x))+2*I*x*polylog
(3,I*exp(x))-2*I*polylog(4,I*exp(x))

```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.16

$$\begin{aligned}
 \int x^2 \arctan(\sinh(x)) dx &= \frac{1}{3} x^3 \arctan(\sinh(x)) + \frac{1}{3} i x^3 \log(i \cosh(x) + i \sinh(x) + 1) \\
 &\quad - \frac{1}{3} i x^3 \log(-i \cosh(x) - i \sinh(x) + 1) \\
 &\quad - i x^2 \operatorname{Li}_2(i \cosh(x) + i \sinh(x)) \\
 &\quad + i x^2 \operatorname{Li}_2(-i \cosh(x) - i \sinh(x)) \\
 &\quad + 2i x \operatorname{polylog}(3, i \cosh(x) + i \sinh(x)) \\
 &\quad - 2i x \operatorname{polylog}(3, -i \cosh(x) - i \sinh(x)) \\
 &\quad - 2i \operatorname{polylog}(4, i \cosh(x) + i \sinh(x)) \\
 &\quad + 2i \operatorname{polylog}(4, -i \cosh(x) - i \sinh(x))
 \end{aligned}$$

input `integrate(x^2*arctan(sinh(x)),x, algorithm="fricas")`

output

```
1/3*x^3*arctan(sinh(x)) + 1/3*I*x^3*log(I*cosh(x) + I*sinh(x) + 1) - 1/3*I
*x^3*log(-I*cosh(x) - I*sinh(x) + 1) - I*x^2*dilog(I*cosh(x) + I*sinh(x))
+ I*x^2*dilog(-I*cosh(x) - I*sinh(x)) + 2*I*x*polylog(3, I*cosh(x) + I*sin
h(x)) - 2*I*x*polylog(3, -I*cosh(x) - I*sinh(x)) - 2*I*polylog(4, I*cosh(x)
) + I*sinh(x)) + 2*I*polylog(4, -I*cosh(x) - I*sinh(x))
```

Sympy [F]

$$\int x^2 \arctan(\sinh(x)) dx = \int x^2 \operatorname{atan}(\sinh(x)) dx$$

input

```
integrate(x**2*atan(sinh(x)),x)
```

output

```
Integral(x**2*atan(sinh(x)), x)
```

Maxima [F]

$$\int x^2 \arctan(\sinh(x)) dx = \int x^2 \arctan(\sinh(x)) dx$$

input

```
integrate(x^2*arctan(sinh(x)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctan(1/2*(e^(2*x) - 1)*e^(-x)) - 2*integrate(1/3*x^3*e^x/(e^(2*x)
+ 1), x)
```

Giac [F]

$$\int x^2 \arctan(\sinh(x)) dx = \int x^2 \arctan(\sinh(x)) dx$$

input `integrate(x^2*arctan(sinh(x)),x, algorithm="giac")`

output `integrate(x^2*arctan(sinh(x)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(\sinh(x)) dx = \int x^2 \operatorname{atan}(\sinh(x)) dx$$

input `int(x^2*atan(sinh(x)),x)`

output `int(x^2*atan(sinh(x)), x)`

Reduce [F]

$$\int x^2 \arctan(\sinh(x)) dx = \int \operatorname{atan}(\sinh(x)) x^2 dx$$

input `int(x^2*atan(sinh(x)),x)`

output `int(atan(sinh(x))*x**2,x)`

3.76 $\int (e + fx)^3 \arctan(\tanh(a + bx)) dx$

Optimal result	595
Mathematica [B] (verified)	596
Rubi [A] (verified)	597
Maple [C] (warning: unable to verify)	601
Fricas [B] (verification not implemented)	602
Sympy [F]	603
Maxima [F]	604
Giac [F(-1)]	604
Mupad [F(-1)]	604
Reduce [F]	605

Optimal result

Integrand size = 15, antiderivative size = 299

$$\begin{aligned}
 \int (e + fx)^3 \arctan(\tanh(a + bx)) dx = & -\frac{(e + fx)^4 \arctan(e^{2a+2bx})}{4f} \\
 & + \frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} \\
 & + \frac{i(e + fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} \\
 & - \frac{i(e + fx)^3 \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b} \\
 & - \frac{3if(e + fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{8b^2} \\
 & + \frac{3if(e + fx)^2 \operatorname{PolyLog}(3, ie^{2a+2bx})}{8b^2} \\
 & + \frac{3if^2(e + fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{8b^3} \\
 & - \frac{3if^2(e + fx) \operatorname{PolyLog}(4, ie^{2a+2bx})}{8b^3} \\
 & - \frac{3if^3 \operatorname{PolyLog}(5, -ie^{2a+2bx})}{16b^4} \\
 & + \frac{3if^3 \operatorname{PolyLog}(5, ie^{2a+2bx})}{16b^4}
 \end{aligned}$$

output

```
-1/4*(f*x+e)^4*arctan(exp(2*b*x+2*a))/f+1/4*(f*x+e)^4*arctan(tanh(b*x+a))/
f+1/4*I*(f*x+e)^3*polylog(2,-I*exp(2*b*x+2*a))/b-1/4*I*(f*x+e)^3*polylog(2
,I*exp(2*b*x+2*a))/b-3/8*I*f*(f*x+e)^2*polylog(3,-I*exp(2*b*x+2*a))/b^2+3/
8*I*f*(f*x+e)^2*polylog(3,I*exp(2*b*x+2*a))/b^2+3/8*I*f^2*(f*x+e)*polylog(
4,-I*exp(2*b*x+2*a))/b^3-3/8*I*f^2*(f*x+e)*polylog(4,I*exp(2*b*x+2*a))/b^3
-3/16*I*f^3*polylog(5,-I*exp(2*b*x+2*a))/b^4+3/16*I*f^3*polylog(5,I*exp(2*
b*x+2*a))/b^4
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 600 vs. $2(299) = 598$.

Time = 0.67 (sec) , antiderivative size = 600, normalized size of antiderivative = 2.01

$$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx$$

$$= \frac{1}{4} x (4e^3 + 6e^2 fx + 4ef^2 x^2 + f^3 x^3) \arctan(\tanh(a + bx))$$

$$- \frac{i(8b^4 e^3 x \log(1 - ie^{2(a+bx)}) + 12b^4 e^2 f x^2 \log(1 - ie^{2(a+bx)}) + 8b^4 e f^2 x^3 \log(1 - ie^{2(a+bx)}) + 2b^4 f^3 x^4 \log(1 - ie^{2(a+bx)}))}{4}$$

input

```
Integrate[(e + f*x)^3*ArcTan[Tanh[a + b*x]],x]
```

output

```
(x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcTan[Tanh[a + b*x]])/4 -
((I/16)*(8*b^4*e^3*x*Log[1 - I*E^(2*(a + b*x))] + 12*b^4*e^2*f*x^2*Log[1 -
I*E^(2*(a + b*x))] + 8*b^4*e*f^2*x^3*Log[1 - I*E^(2*(a + b*x))] + 2*b^4*f
^3*x^4*Log[1 - I*E^(2*(a + b*x))] - 8*b^4*e^3*x*Log[1 + I*E^(2*(a + b*x))]
- 12*b^4*e^2*f*x^2*Log[1 + I*E^(2*(a + b*x))] - 8*b^4*e*f^2*x^3*Log[1 + I
*E^(2*(a + b*x))] - 2*b^4*f^3*x^4*Log[1 + I*E^(2*(a + b*x))] - 4*b^3*(e +
f*x)^3*PolyLog[2, (-I)*E^(2*(a + b*x))] + 4*b^3*(e + f*x)^3*PolyLog[2, I*E
^(2*(a + b*x))] + 6*b^2*e^2*f*PolyLog[3, (-I)*E^(2*(a + b*x))] + 12*b^2*e*
f^2*x*PolyLog[3, (-I)*E^(2*(a + b*x))] + 6*b^2*f^3*x^2*PolyLog[3, (-I)*E^(
2*(a + b*x))] - 6*b^2*e^2*f*PolyLog[3, I*E^(2*(a + b*x))] - 12*b^2*e*f^2*x
*PolyLog[3, I*E^(2*(a + b*x))] - 6*b^2*f^3*x^2*PolyLog[3, I*E^(2*(a + b*x)
)] - 6*b*e*f^2*PolyLog[4, (-I)*E^(2*(a + b*x))] - 6*b*f^3*x*PolyLog[4, (-I
)*E^(2*(a + b*x))] + 6*b*e*f^2*PolyLog[4, I*E^(2*(a + b*x))] + 6*b*f^3*x*P
olyLog[4, I*E^(2*(a + b*x))] + 3*f^3*PolyLog[5, (-I)*E^(2*(a + b*x))] - 3*
f^3*PolyLog[5, I*E^(2*(a + b*x))])/b^4
```

Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {5706, 3042, 4668, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)^3 \arctan(\tanh(a + bx)) dx \\
 & \quad \downarrow \text{5706} \\
 & \frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \frac{b \int (e + fx)^4 \operatorname{sech}(2a + 2bx) dx}{4f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \frac{b \int (e + fx)^4 \csc\left(2ia + 2ibx + \frac{\pi}{2}\right) dx}{4f} \\
 & \quad \downarrow \text{4668} \\
 & \frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \\
 & \frac{b \left(-\frac{2if \int (e+fx)^3 \log(1-ie^{2a+2bx}) dx}{b} + \frac{2if \int (e+fx)^3 \log(1+ie^{2a+2bx}) dx}{b} + \frac{(e+fx)^4 \arctan(e^{2a+2bx})}{b} \right)}{4f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \\
 & b \left(\frac{2if \left(\frac{3f \int (e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{2if \left(\frac{3f \int (e+fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx}) dx}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{b} \right) \\
 & \quad \downarrow \text{7163}
 \end{aligned}$$

$$\frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \int (e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx}) dx}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(1, -ie^{2a+2bx})}{2b} \right)}{b}$$

4f

↓ 7163

$$\frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(4, -ie^{2a+2bx}) dx}{2b} \right)}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(1, -ie^{2a+2bx})}{2b} \right)}{b}$$

↓ 2720

$$\frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{2b} - \frac{f \int e^{-2a-2bx} \operatorname{PolyLog}(4, -ie^{2a+2bx}) de^{2a+2bx}}{4b^2} \right)}{b} \right)}{2b} \right)}{b} - \frac{(e+fx)^3 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b}$$

7143

$$\frac{(e + fx)^4 \arctan(\tanh(a + bx))}{4f} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{2b} - \frac{f \operatorname{PolyLog}(5, -ie^{2a+2bx})}{4b^2} \right)}{b} \right)}{2b} \right)}{b} + \frac{(e+fx)^4 \arctan(e^{2a+2bx})}{b}$$

input `Int[(e + f*x)^3*ArcTan[Tanh[a + b*x]],x]`

output

```
((e + f*x)^4*ArcTan[Tanh[a + b*x]]/(4*f) - (b*(((e + f*x)^4*ArcTan[E^(2*a
+ 2*b*x)])/b + ((2*I)*f*(-1/2*((e + f*x)^3*PolyLog[2, (-I)*E^(2*a + 2*b*x
)])/b + (3*f*(((e + f*x)^2*PolyLog[3, (-I)*E^(2*a + 2*b*x)])/(2*b) - (f*((
e + f*x)*PolyLog[4, (-I)*E^(2*a + 2*b*x)])/(2*b) - (f*PolyLog[5, (-I)*E^(
2*a + 2*b*x)]/(4*b^2)))/b))/(2*b)))/b - ((2*I)*f*(-1/2*((e + f*x)^3*PolyL
og[2, I*E^(2*a + 2*b*x)])/b + (3*f*(((e + f*x)^2*PolyLog[3, I*E^(2*a + 2*b
*x)])/(2*b) - (f*((e + f*x)*PolyLog[4, I*E^(2*a + 2*b*x)])/(2*b) - (f*Pol
yLog[5, I*E^(2*a + 2*b*x)]/(4*b^2)))/b))/(2*b)))/b)/(4*f)
```

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4668

```
Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m, x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5706 `Int[ArcTan[Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Simp[(e + f*x)^(m + 1)*(ArcTan[Tanh[a + b*x]]/(f*(m + 1))), x] - Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a,
b, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 38.76 (sec) , antiderivative size = 3570, normalized size of antiderivative = 11.94

method	result	size
risch	Expression too large to display	3570

input `int((f*x+e)^3*arctan(tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

-3/16*I*f^3*polylog(5,-I*exp(2*b*x+2*a))/b^4+3/16*I*f^3*polylog(5,I*exp(2*
b*x+2*a))/b^4+1/16*Pi*(csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))*csgn(
(1+I)*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))-csgn((1+I)*(exp(2*b*x+2*a)+I)
/(exp(2*b*x+2*a)+1))^2-csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1))*csgn(
(1-I)*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1))-csgn((1-I)*(exp(2*b*x+2*a)-I)
/(exp(2*b*x+2*a)+1))^2-csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*(exp(2*b*x+2*a)-I
))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1))+csgn(I/(exp(2*b*x+2*a)+1)
)*csgn(I*(exp(2*b*x+2*a)+I))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))
+csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1))^
2-csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))
^2+csgn(I*(exp(2*b*x+2*a)-I))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1)
)^2-csgn(I*(exp(2*b*x+2*a)+I))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1
))^2-csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1))^3+csgn(I*(exp(2*b*x+2*a
)-I)/(exp(2*b*x+2*a)+1))*csgn((1-I)*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)+1))
^2+csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))^3-csgn(I*(exp(2*b*x+2*a)+
I)/(exp(2*b*x+2*a)+1))*csgn((1+I)*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))^2
+csgn((1+I)*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)+1))^3+csgn((1-I)*(exp(2*b*x
+2*a)-I)/(exp(2*b*x+2*a)+1))^3+1)*(f*x+e)^4/f-1/8*I*f^3*ln(exp(2*b*x+2*a)-
I)*x^4-1/2*I*ln(exp(2*b*x+2*a)-I)*x*e^3-1/8*I/f*ln(exp(2*b*x+2*a)-I)*e^4+3
/2*I*f/b*a*e^2*ln((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))*x+3/2*I*f/b*a*e^2...

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1460 vs. $2(236) = 472$.

Time = 0.21 (sec) , antiderivative size = 1460, normalized size of antiderivative = 4.88

$$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^3*arctan(tanh(b*x+a)),x, algorithm="fricas")
```

output

```

1/8*(24*I*f^3*polylog(5, 1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) +
24*I*f^3*polylog(5, -1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 24*I
*f^3*polylog(5, 1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 24*I*f^3
*polylog(5, -1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 2*(b^4*f^3*
x^4 + 4*b^4*e*f^2*x^3 + 6*b^4*e^2*f*x^2 + 4*b^4*e^3*x)*arctan(sinh(b*x + a
)/cosh(b*x + a)) - 4*(I*b^3*f^3*x^3 + 3*I*b^3*e*f^2*x^2 + 3*I*b^3*e^2*f*x
+ I*b^3*e^3)*dilog(1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 4*(I*b
^3*f^3*x^3 + 3*I*b^3*e*f^2*x^2 + 3*I*b^3*e^2*f*x + I*b^3*e^3)*dilog(-1/2*s
qrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 4*(-I*b^3*f^3*x^3 - 3*I*b^3*e*
f^2*x^2 - 3*I*b^3*e^2*f*x - I*b^3*e^3)*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a)
+ sinh(b*x + a))) - 4*(-I*b^3*f^3*x^3 - 3*I*b^3*e*f^2*x^2 - 3*I*b^3*e^2*f
*x - I*b^3*e^3)*dilog(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + (
-I*b^4*f^3*x^4 - 4*I*b^4*e*f^2*x^3 - 6*I*b^4*e^2*f*x^2 - 4*I*b^4*e^3*x - 4
*I*a*b^3*e^3 + 6*I*a^2*b^2*e^2*f - 4*I*a^3*b*e*f^2 + I*a^4*f^3)*log(1/2*sq
rt(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b^4*f^3*x^4 - 4*I*b^4*e
*f^2*x^3 - 6*I*b^4*e^2*f*x^2 - 4*I*b^4*e^3*x - 4*I*a*b^3*e^3 + 6*I*a^2*b^2
*e^2*f - 4*I*a^3*b*e*f^2 + I*a^4*f^3)*log(-1/2*sqrt(4*I)*(cosh(b*x + a) +
sinh(b*x + a)) + 1) + (I*b^4*f^3*x^4 + 4*I*b^4*e*f^2*x^3 + 6*I*b^4*e^2*f*x
^2 + 4*I*b^4*e^3*x + 4*I*a*b^3*e^3 - 6*I*a^2*b^2*e^2*f + 4*I*a^3*b*e*f^2 -
I*a^4*f^3)*log(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (...

```

Sympy [F]

$$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx = \int (e + fx)^3 \operatorname{atan}(\tanh(a + bx)) dx$$

input

```
integrate((f*x+e)**3*atan(tanh(b*x+a)), x)
```

output

```
Integral((e + f*x)**3*atan(tanh(a + b*x)), x)
```

Maxima [F]

$$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx = \int (fx + e)^3 \arctan(\tanh(bx + a)) dx$$

input `integrate((f*x+e)^3*arctan(tanh(b*x+a)),x, algorithm="maxima")`

output `1/4*(f^3*x^4 + 4*e*f^2*x^3 + 6*e^2*f*x^2 + 4*e^3*x)*arctan((e^(2*b*x + 2*a) - 1)/(e^(2*b*x + 2*a) + 1)) - integrate(1/2*(b*f^3*x^4*e^(2*a) + 4*b*e*f^2*x^3*e^(2*a) + 6*b*e^2*f*x^2*e^(2*a) + 4*b*e^3*x*e^(2*a))*e^(2*b*x)/(e^(4*b*x + 4*a) + 1), x)`

Giac [F(-1)]

Timed out.

$$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx = \text{Timed out}$$

input `integrate((f*x+e)^3*arctan(tanh(b*x+a)),x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^3 \arctan(\tanh(a + bx)) dx = \int \text{atan}(\tanh(a + bx)) (e + fx)^3 dx$$

input `int(atan(tanh(a + b*x))*(e + f*x)^3,x)`

output `int(atan(tanh(a + b*x))*(e + f*x)^3, x)`

Reduce [F]

$$\begin{aligned} \int (e + fx)^3 \arctan(\tanh(a + bx)) dx &= \left(\int \operatorname{atan}(\tanh(bx + a)) dx \right) e^3 \\ &+ \left(\int \operatorname{atan}(\tanh(bx + a)) x^3 dx \right) f^3 \\ &+ 3 \left(\int \operatorname{atan}(\tanh(bx + a)) x^2 dx \right) e f^2 \\ &+ 3 \left(\int \operatorname{atan}(\tanh(bx + a)) x dx \right) e^2 f \end{aligned}$$

input `int((f*x+e)^3*atan(tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)),x)*e**3 + int(atan(tanh(a + b*x))*x**3,x)*f**3 + 3
*int(atan(tanh(a + b*x))*x**2,x)*e*f**2 + 3*int(atan(tanh(a + b*x))*x,x)*e
**2*f`

3.77 $\int (e + fx)^2 \arctan(\tanh(a + bx)) dx$

Optimal result	606
Mathematica [A] (verified)	607
Rubi [A] (verified)	607
Maple [C] (warning: unable to verify)	611
Fricas [B] (verification not implemented)	612
Sympy [F]	613
Maxima [F]	613
Giac [F(-1)]	613
Mupad [F(-1)]	614
Reduce [F]	614

Optimal result

Integrand size = 15, antiderivative size = 229

$$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx = -\frac{(e + fx)^3 \arctan(e^{2a+2bx})}{3f} + \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} + \frac{i(e + fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} - \frac{i(e + fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b} - \frac{if(e + fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{4b^2} + \frac{if(e + fx) \operatorname{PolyLog}(3, ie^{2a+2bx})}{4b^2} + \frac{if^2 \operatorname{PolyLog}(4, -ie^{2a+2bx})}{8b^3} - \frac{if^2 \operatorname{PolyLog}(4, ie^{2a+2bx})}{8b^3}$$

$$\begin{aligned}
 & \int (e + fx)^2 \arctan(\tanh(a + bx)) dx \\
 & \quad \downarrow \text{5706} \\
 & \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \frac{b \int (e + fx)^3 \operatorname{sech}(2a + 2bx) dx}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \frac{b \int (e + fx)^3 \csc\left(2ia + 2ibx + \frac{\pi}{2}\right) dx}{3f} \\
 & \quad \downarrow \text{4668} \\
 & \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \\
 & \frac{b \left(-\frac{3if \int (e+fx)^2 \log(1-ie^{2a+2bx}) dx}{2b} + \frac{3if \int (e+fx)^2 \log(1+ie^{2a+2bx}) dx}{2b} + \frac{(e+fx)^3 \arctan(e^{2a+2bx})}{b} \right)}{3f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \\
 & b \left(\frac{3if \left(\frac{f \int (e+fx) \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \int (e+fx) \operatorname{PolyLog}(2, ie^{2a+2bx}) dx}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{2b} \right) \\
 & \quad \downarrow \text{7163} \\
 & \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \\
 & b \left(\frac{3if \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(3, -ie^{2a+2bx}) dx}{2b} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(3, ie^{2a+2bx}) dx}{2b} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{2b} \right) \\
 & \quad \downarrow \text{2720} \\
 & \frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} -
 \end{aligned}$$

$$\frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \frac{b \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \int e^{-2a-2bx} \operatorname{PolyLog}(3, -ie^{2a+2bx}) de^{2a+2bx}}{4b^2} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \left(\frac{(e+fx)}{2b} \right)}{3f} \right)}{3f}$$

3f

7143

$$\frac{(e + fx)^3 \arctan(\tanh(a + bx))}{3f} - \frac{b \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \operatorname{PolyLog}(4, -ie^{2a+2bx})}{4b^2} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \left(\frac{(e+fx)^3 \arctan(e^{2a+2bx})}{b} \right)}{3f} \right)}{3f}$$

3f

input `Int[(e + f*x)^2*ArcTan[Tanh[a + b*x]],x]`

output `((e + f*x)^3*ArcTan[Tanh[a + b*x]])/(3*f) - (b*(((e + f*x)^3*ArcTan[E^(2*a + 2*b*x)])/b + (((3*I)/2)*f*(-1/2*((e + f*x)^2*PolyLog[2, (-I)*E^(2*a + 2*b*x)])/b + (f*(((e + f*x)*PolyLog[3, (-I)*E^(2*a + 2*b*x)])/(2*b) - (f*PolyLog[4, (-I)*E^(2*a + 2*b*x)])/(4*b^2)))/b))/b - (((3*I)/2)*f*(-1/2*((e + f*x)^2*PolyLog[2, I*E^(2*a + 2*b*x)])/b + (f*(((e + f*x)*PolyLog[3, I*E^(2*a + 2*b*x)])/(2*b) - (f*PolyLog[4, I*E^(2*a + 2*b*x)])/(4*b^2)))/b))/b)/(3*f)`

Definitions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*(f_ + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5706 `Int[ArcTan[Tanh[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_))^(m_), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[Tanh[a + b*x]]/(f*(m + 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a, b, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 26.38 (sec) , antiderivative size = 2668, normalized size of antiderivative = 11.65

method	result	size
risch	Expression too large to display	2668

input

```
int((f*x+e)^2*arctan(tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/8*I*f^2*polylog(4,I*exp(2*b*x+2*a))/b^3+1/8*I*f^2*polylog(4,-I*exp(2*b*
x+2*a))/b^3-1/6*I/f*ln(exp(2*b*x+2*a)+I)*e^3-1/6*I*f^2*ln(1-I*exp(2*b*x+2*
a))*x^3-1/2*I/b*e^2*dilog(((I)^(1/2)-exp(b*x+a))/(I)^(1/2))-1/2*I/b*e^2*
dilog(((I)^(1/2)+exp(b*x+a))/(I)^(1/2))-1/2*I*e^2*ln(((I)^(1/2)-exp(b*x
+a))/(I)^(1/2))*x-1/2*I*e^2*ln(((I)^(1/2)+exp(b*x+a))/(I)^(1/2))*x+1/6*
I/f*e^3*ln(-exp(2*b*x+2*a)+I)+1/6*I*f^2*ln(1+I*exp(2*b*x+2*a))*x^3+1/2*I*e
^2*ln(1+exp(b*x+a)*(-1)^(3/4))*x+1/2*I*e^2*ln(1-exp(b*x+a)*(-1)^(3/4))*x+
1/2*I/b*e^2*dilog(1+exp(b*x+a)*(-1)^(3/4))+1/2*I/b*e^2*dilog(1-exp(b*x+a)*
(-1)^(3/4))+I*f/b^2*a^2*e*ln(((I)^(1/2)-exp(b*x+a))/(I)^(1/2))+I*f/b^2*a^
2*e*ln(((I)^(1/2)+exp(b*x+a))/(I)^(1/2))+I*f/b^2*a*e*dilog(((I)^(1/2)-e
xp(b*x+a))/(I)^(1/2))+I*f/b^2*a*e*dilog(((I)^(1/2)+exp(b*x+a))/(I)^(1/2
))-1/2*I*f/b^2*e*ln(1-I*exp(2*b*x+2*a))*a^2-1/2*I*f/b*e*polylog(2,I*exp(2*
b*x+2*a))*x-1/2*I*f/b^2*e*polylog(2,I*exp(2*b*x+2*a))*a-1/2*I*f^2*a^2/b^2*
ln(((I)^(1/2)-exp(b*x+a))/(I)^(1/2))*x-1/2*I*f^2*a^2/b^2*ln(((I)^(1/2)+
exp(b*x+a))/(I)^(1/2))*x+1/2*I*f^2/b^2*ln(1-I*exp(2*b*x+2*a))*a^2*x-1/2*I
*f/b^2*a^2*e*ln(exp(2*b*x+2*a)+I)-I*f/b^2*a^2*e*ln(1-exp(b*x+a)*(-1)^(3/4)
)-I*f/b^2*a*e*dilog(1+exp(b*x+a)*(-1)^(3/4))-I*f/b^2*a*e*dilog(1-exp(b*x+a)
)*(-1)^(3/4))-1/2*I*f^2/b^2*ln(1+I*exp(2*b*x+2*a))*a^2*x+1/2*I*f/b^2*a^2*
e*ln(-exp(2*b*x+2*a)+I)+1/2*I*f/b^2*e*ln(1+I*exp(2*b*x+2*a))*a^2+1/2*I*f/b*
e*polylog(2,-I*exp(2*b*x+2*a))*x+1/2*I*f/b^2*e*polylog(2,-I*exp(2*b*x+2...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1002 vs. $2(180) = 360$.

Time = 0.16 (sec) , antiderivative size = 1002, normalized size of antiderivative = 4.38

$$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)^2*arctan(tanh(b*x+a)),x, algorithm="fricas")`

output

```
1/6*(-6*I*f^2*polylog(4, 1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) -
6*I*f^2*polylog(4, -1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 6*I*f^2*po
lylog(4, 1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 6*I*f^2*po
lylog(4, -1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 2*(b^3*f^2*x^3
+ 3*b^3*e*f*x^2 + 3*b^3*e^2*x)*arctan(sinh(b*x + a)/cosh(b*x + a)) - 3*(I
*b^2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog(1/2*sqrt(4*I)*(cosh(b*x +
a) + sinh(b*x + a))) - 3*(I*b^2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog
(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 3*(-I*b^2*f^2*x^2 - 2*I
*b^2*e*f*x - I*b^2*e^2)*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a
))) - 3*(-I*b^2*f^2*x^2 - 2*I*b^2*e*f*x - I*b^2*e^2)*dilog(-1/2*sqrt(-4*I)
*(cosh(b*x + a) + sinh(b*x + a))) + (-I*b^3*f^2*x^3 - 3*I*b^3*e*f*x^2 - 3*
I*b^3*e^2*x - 3*I*a*b^2*e^2 + 3*I*a^2*b*e*f - I*a^3*f^2)*log(1/2*sqrt(4*I)
*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b^3*f^2*x^3 - 3*I*b^3*e*f*x^2
- 3*I*b^3*e^2*x - 3*I*a*b^2*e^2 + 3*I*a^2*b*e*f - I*a^3*f^2)*log(-1/2*sqrt
(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^3*f^2*x^3 + 3*I*b^3*e*f*
x^2 + 3*I*b^3*e^2*x + 3*I*a*b^2*e^2 - 3*I*a^2*b*e*f + I*a^3*f^2)*log(1/2*sq
rt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^3*f^2*x^3 + 3*I*b^3*
e*f*x^2 + 3*I*b^3*e^2*x + 3*I*a*b^2*e^2 - 3*I*a^2*b*e*f + I*a^3*f^2)*log(-
1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (3*I*a*b^2*e^2 - 3*I
*a^2*b*e*f + I*a^3*f^2)*log(I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x ...
```

Sympy [F]

$$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx = \int (e + fx)^2 \operatorname{atan}(\tanh(a + bx)) dx$$

input `integrate((f*x+e)**2*atan(tanh(b*x+a)),x)`

output `Integral((e + f*x)**2*atan(tanh(a + b*x)), x)`

Maxima [F]

$$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx = \int (fx + e)^2 \arctan(\tanh(bx + a)) dx$$

input `integrate((f*x+e)^2*arctan(tanh(b*x+a)),x, algorithm="maxima")`

output `1/3*(f^2*x^3 + 3*e*f*x^2 + 3*e^2*x)*arctan((e^(2*b*x + 2*a) - 1)/(e^(2*b*x + 2*a) + 1)) - integrate(2/3*(b*f^2*x^3*e^(2*a) + 3*b*e*f*x^2*e^(2*a) + 3*b*e^2*x*e^(2*a))*e^(2*b*x)/(e^(4*b*x + 4*a) + 1), x)`

Giac [F(-1)]

Timed out.

$$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx = \text{Timed out}$$

input `integrate((f*x+e)^2*arctan(tanh(b*x+a)),x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 \arctan(\tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(a + bx)) (e + fx)^2 dx$$

input `int(atan(tanh(a + b*x))*(e + f*x)^2,x)`output `int(atan(tanh(a + b*x))*(e + f*x)^2, x)`**Reduce [F]**

$$\begin{aligned} \int (e + fx)^2 \arctan(\tanh(a + bx)) dx &= \left(\int \operatorname{atan}(\tanh(bx + a)) dx \right) e^2 \\ &+ \left(\int \operatorname{atan}(\tanh(bx + a)) x^2 dx \right) f^2 \\ &+ 2 \left(\int \operatorname{atan}(\tanh(bx + a)) x dx \right) ef \end{aligned}$$

input `int((f*x+e)^2*atan(tanh(b*x+a)),x)`output `int(atan(tanh(a + b*x)),x)*e**2 + int(atan(tanh(a + b*x))*x**2,x)*f**2 + 2*int(atan(tanh(a + b*x))*x,x)*e*f`

3.78 $\int (e + fx) \arctan(\tanh(a + bx)) dx$

Optimal result	615
Mathematica [A] (verified)	616
Rubi [A] (verified)	616
Maple [C] (warning: unable to verify)	619
Fricas [B] (verification not implemented)	620
Sympy [F]	621
Maxima [F]	621
Giac [F]	621
Mupad [F(-1)]	622
Reduce [F]	622

Optimal result

Integrand size = 13, antiderivative size = 159

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = -\frac{(e + fx)^2 \arctan(e^{2a+2bx})}{2f} + \frac{(e + fx)^2 \arctan(\tanh(a + bx))}{2f} + \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} - \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b} - \frac{if \operatorname{PolyLog}(3, -ie^{2a+2bx})}{8b^2} + \frac{if \operatorname{PolyLog}(3, ie^{2a+2bx})}{8b^2}$$

output

```
-1/2*(f*x+e)^2*arctan(exp(2*b*x+2*a))/f+1/2*(f*x+e)^2*arctan(tanh(b*x+a))/f+1/4*I*(f*x+e)*polylog(2,-I*exp(2*b*x+2*a))/b-1/4*I*(f*x+e)*polylog(2,I*exp(2*b*x+2*a))/b-1/8*I*f*polylog(3,-I*exp(2*b*x+2*a))/b^2+1/8*I*f*polylog(3,I*exp(2*b*x+2*a))/b^2
```


Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.49

$$\int (e+fx) \arctan(\tanh(a+bx)) dx = ex \arctan(\tanh(a+bx)) + \frac{1}{2} fx^2 \arctan(\tanh(a+bx))$$

$$\frac{ie(2bx(\log(1 - ie^{2(a+bx)}) - \log(1 + ie^{2(a+bx)})) - \text{PolyLog}(2, -ie^{2(a+bx)}) + \text{PolyLog}(2, ie^{2(a+bx)}))}{8b^2}$$

$$\frac{if(2b^2x^2 \log(1 - ie^{2(a+bx)}) - 2b^2x^2 \log(1 + ie^{2(a+bx)}) - 2bx \text{PolyLog}(2, -ie^{2(a+bx)}) + 2bx \text{PolyLog}(2, ie^{2(a+bx)}))}{8b^2}$$

input

```
Integrate[(e + f*x)*ArcTan[Tanh[a + b*x]], x]
```

output

```
e*x*ArcTan[Tanh[a + b*x]] + (f*x^2*ArcTan[Tanh[a + b*x]])/2 - ((I/4)*e*(2*
b*x*(Log[1 - I*E^(2*(a + b*x))] - Log[1 + I*E^(2*(a + b*x))]) - PolyLog[2,
(-I)*E^(2*(a + b*x))] + PolyLog[2, I*E^(2*(a + b*x))])/b - ((I/8)*f*(2*b
^2*x^2*Log[1 - I*E^(2*(a + b*x))] - 2*b^2*x^2*Log[1 + I*E^(2*(a + b*x))] -
2*b*x*PolyLog[2, (-I)*E^(2*(a + b*x))] + 2*b*x*PolyLog[2, I*E^(2*(a + b*x
))] + PolyLog[3, (-I)*E^(2*(a + b*x))] - PolyLog[3, I*E^(2*(a + b*x))])/b
^2
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.10,
 number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules
 used = {5706, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e+fx) \arctan(\tanh(a+bx)) dx$$

$$\downarrow \text{5706}$$

$$\frac{(e+fx)^2 \arctan(\tanh(a+bx))}{2f} - \frac{b \int (e+fx)^2 \operatorname{sech}(2a+2bx) dx}{2f}$$

$$\downarrow \text{3042}$$

$$\frac{(e + fx)^2 \arctan(\tanh(a + bx))}{2f} - \frac{b \int (e + fx)^2 \csc(2ia + 2ibx + \frac{\pi}{2}) dx}{2f}$$

↓ 4668

$$\frac{(e + fx)^2 \arctan(\tanh(a + bx))}{2f} - \frac{b \left(-\frac{if \int (e+fx) \log(1-ie^{2a+2bx}) dx}{b} + \frac{if \int (e+fx) \log(1+ie^{2a+2bx}) dx}{b} + \frac{(e+fx)^2 \arctan(e^{2a+2bx})}{b} \right)}{2f}$$

↓ 3011

$$\frac{(e + fx)^2 \arctan(\tanh(a + bx))}{2f} - \frac{b \left(\frac{if \left(\frac{f \int \text{PolyLog}(2, -ie^{2a+2bx}) dx}{2b} - \frac{(e+fx) \text{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{if \left(\frac{f \int \text{PolyLog}(2, ie^{2a+2bx}) dx}{2b} - \frac{(e+fx) \text{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{b} \right) + \frac{(e+fx)^2 \arctan(e^{2a+2bx})}{b}}{2f}$$

↓ 2720

$$\frac{(e + fx)^2 \arctan(\tanh(a + bx))}{2f} - \frac{b \left(\frac{if \left(\frac{f \int e^{-2a-2bx} \text{PolyLog}(2, -ie^{2a+2bx}) de^{2a+2bx}}{4b^2} - \frac{(e+fx) \text{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{if \left(\frac{f \int e^{-2a-2bx} \text{PolyLog}(2, ie^{2a+2bx}) de^{2a+2bx}}{4b^2} - \frac{(e+fx) \text{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{b} \right)}{2f}$$

↓ 7143

$$\frac{(e + fx)^2 \arctan(\tanh(a + bx))}{2f} - \frac{b \left(\frac{(e+fx)^2 \arctan(e^{2a+2bx})}{b} + \frac{if \left(\frac{f \text{PolyLog}(3, -ie^{2a+2bx})}{4b^2} - \frac{(e+fx) \text{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{if \left(\frac{f \text{PolyLog}(3, ie^{2a+2bx})}{4b^2} - \frac{(e+fx) \text{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{b} \right)}{2f}$$

input

```
Int[(e + f*x)*ArcTan[Tanh[a + b*x]], x]
```

output

```
((e + f*x)^2*ArcTan[Tanh[a + b*x]])/(2*f) - (b*(((e + f*x)^2*ArcTan[E^(2*a + 2*b*x)]))/b + (I*f*(-1/2*((e + f*x)*PolyLog[2, (-I)*E^(2*a + 2*b*x)]))/b + (f*PolyLog[3, (-I)*E^(2*a + 2*b*x)])/(4*b^2)))/b - (I*f*(-1/2*((e + f*x)*PolyLog[2, I*E^(2*a + 2*b*x)]))/b + (f*PolyLog[3, I*E^(2*a + 2*b*x)])/(4*b^2))/b)/(2*f)
```

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4668

```
Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5706

```
Int[ArcTan[Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[Tanh[a + b*x]]/(f*(m + 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a, b, e, f}, x] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.00 (sec) , antiderivative size = 1776, normalized size of antiderivative = 11.17

method	result	size
risch	Expression too large to display	1776

input

```
int((f*x+e)*arctan(tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/8*I*f*polylog(3,-I*exp(2*b*x+2*a))/b^2+1/8*I*f*polylog(3,I*exp(2*b*x+2*a))/b^2-1/2*I*e*ln((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2)*x-1/2*I*e*ln((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2)*x-1/2*I*e/b*dilog((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2)-1/2*I*e/b*dilog((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2)-1/4*I*f*ln(1-I*exp(2*b*x+2*a))*x^2+1/4*I*f*ln(1+I*exp(2*b*x+2*a))*x^2+1/2*I*e*ln(1+exp(b*x+a))*(-1)^(3/4)*x+1/2*I*e*ln(1-exp(b*x+a))*(-1)^(3/4)*x+1/2*I*e/b*dilog(1+exp(b*x+a))*(-1)^(3/4)+1/2*I*e/b*dilog(1-exp(b*x+a))*(-1)^(3/4)-1/2*I*e/b*ln((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2)*a-1/2*I*e/b*ln((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2)*a+1/2*I*f/b^2*a^2*ln((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))+1/2*I*f/b^2*a^2*ln((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2))+1/2*I*f/b^2*a*dilog((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))+1/2*I*f/b^2*a*dilog((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2))-1/2*I*f/b^2*a^2*ln(1+exp(b*x+a))*(-1)^(3/4))-1/2*I*f/b^2*a^2*ln(1-exp(b*x+a))*(-1)^(3/4))-1/2*I*f/b^2*a*dilog(1+exp(b*x+a))*(-1)^(3/4))-1/2*I*f/b^2*a*dilog(1-exp(b*x+a))*(-1)^(3/4))-1/4*I*f/b^2*ln(1-I*exp(2*b*x+2*a))*a^2-1/4*I*f/b*polylog(2,I*exp(2*b*x+2*a))*x-1/4*I*f/b^2*polylog(2,I*exp(2*b*x+2*a))*a+1/2*I*e/b*ln(1+exp(b*x+a))*(-1)^(3/4))*a+1/2*I*e/b*ln(1-exp(b*x+a))*(-1)^(3/4))*a+1/4*I*f/b^2*ln(1+I*exp(2*b*x+2*a))*a^2+1/4*I*f/b*polylog(2,-I*exp(2*b*x+2*a))*x+1/4*I*f/b^2*polylog(2,-I*exp(2*b*x+2*a))*a-1/4*I/b^2*f*a^2*ln(exp(2*b*x+2*a)+I)+1/2*I/b*a*e*ln(exp(2*b*x+2*a)+I)-1/2*I*e/b*a*ln(-exp(2*b*x+2*a)+I)+1/4*I*f/b^2*a^2*ln(-exp(2*b*x+2*a))+...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 600 vs. $2(130) = 260$.

Time = 0.13 (sec) , antiderivative size = 600, normalized size of antiderivative = 3.77

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)*arctan(tanh(b*x+a)),x, algorithm="fricas")`

output

```
1/4*(2*(b^2*f*x^2 + 2*b^2*e*x)*arctan(sinh(b*x + a)/cosh(b*x + a)) - 2*(I*
b*f*x + I*b*e)*dilog(1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 2*(I
*b*f*x + I*b*e)*dilog(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 2*
(-I*b*f*x - I*b*e)*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) -
2*(-I*b*f*x - I*b*e)*dilog(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)
)) + (-I*b^2*f*x^2 - 2*I*b^2*e*x - 2*I*a*b*e + I*a^2*f)*log(1/2*sqrt(4*I)*
(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b^2*f*x^2 - 2*I*b^2*e*x - 2*I*a
*b*e + I*a^2*f)*log(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) +
(I*b^2*f*x^2 + 2*I*b^2*e*x + 2*I*a*b*e - I*a^2*f)*log(1/2*sqrt(-4*I)*(cosh
(b*x + a) + sinh(b*x + a)) + 1) + (I*b^2*f*x^2 + 2*I*b^2*e*x + 2*I*a*b*e -
I*a^2*f)*log(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (2*I*
a*b*e - I*a^2*f)*log(I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) + (2
*I*a*b*e - I*a^2*f)*log(-I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a))
+ (-2*I*a*b*e + I*a^2*f)*log(I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh(b*x +
a)) + (-2*I*a*b*e + I*a^2*f)*log(-I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh
(b*x + a)) + 2*I*f*polylog(3, 1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a)
)) + 2*I*f*polylog(3, -1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 2*
I*f*polylog(3, 1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 2*I*f*pol
ylog(3, -1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)))/b^2
```

Sympy [F]

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = \int (e + fx) \operatorname{atan}(\tanh(a + bx)) dx$$

input `integrate((f*x+e)*atan(tanh(b*x+a)),x)`

output `Integral((e + f*x)*atan(tanh(a + b*x)), x)`

Maxima [F]

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = \int (fx + e) \arctan(\tanh(bx + a)) dx$$

input `integrate((f*x+e)*arctan(tanh(b*x+a)),x, algorithm="maxima")`

output `1/2*(f*x^2 + 2*e*x)*arctan((e^(2*b*x + 2*a) - 1)/(e^(2*b*x + 2*a) + 1)) -
integrate((b*f*x^2*e^(2*a) + 2*b*e*x*e^(2*a))*e^(2*b*x)/(e^(4*b*x + 4*a) +
1), x)`

Giac [F]

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = \int (fx + e) \arctan(\tanh(bx + a)) dx$$

input `integrate((f*x+e)*arctan(tanh(b*x+a)),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(a + bx)) (e + fx) dx$$

input `int(atan(tanh(a + b*x))*(e + f*x),x)`output `int(atan(tanh(a + b*x))*(e + f*x), x)`**Reduce [F]**

$$\int (e + fx) \arctan(\tanh(a + bx)) dx = \left(\int \operatorname{atan}(\tanh(bx + a)) dx \right) e + \left(\int \operatorname{atan}(\tanh(bx + a)) x dx \right) f$$

input `int((f*x+e)*atan(tanh(b*x+a)),x)`output `int(atan(tanh(a + b*x)),x)*e + int(atan(tanh(a + b*x))*x,x)*f`

3.79 $\int \arctan(\tanh(a + bx)) dx$

Optimal result	623
Mathematica [A] (verified)	623
Rubi [A] (verified)	624
Maple [A] (verified)	626
Fricas [B] (verification not implemented)	626
Sympy [F]	627
Maxima [F]	627
Giac [F]	628
Mupad [F(-1)]	628
Reduce [F]	628

Optimal result

Integrand size = 7, antiderivative size = 74

$$\int \arctan(\tanh(a + bx)) dx = -x \arctan(e^{2a+2bx}) + x \arctan(\tanh(a + bx)) + \frac{i \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} - \frac{i \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b}$$

output

```
-x*arctan(exp(2*b*x+2*a))+x*arctan(tanh(b*x+a))+1/4*I*polylog(2,-I*exp(2*b*x+2*a))/b-1/4*I*polylog(2,I*exp(2*b*x+2*a))/b
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.23

$$\int \arctan(\tanh(a + bx)) dx = x \arctan(\tanh(a + bx)) - \frac{i(2bx(\log(1 - ie^{2(a+bx)}) - \log(1 + ie^{2(a+bx)})) - \operatorname{PolyLog}(2, -ie^{2(a+bx)}) + \operatorname{PolyLog}(2, ie^{2(a+bx)}))}{4b}$$

input

```
Integrate[ArcTan[Tanh[a + b*x]],x]
```


output

```
x*ArcTan[Tanh[a + b*x]] - ((I/4)*(2*b*x*(Log[1 - I*E^(2*(a + b*x))] - Log[
1 + I*E^(2*(a + b*x))]) - PolyLog[2, (-I)*E^(2*(a + b*x))] + PolyLog[2, I*
E^(2*(a + b*x))]))/b
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {5702, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(\tanh(a + bx)) dx \\
 & \quad \downarrow 5702 \\
 & x \arctan(\tanh(a + bx)) - b \int x \operatorname{sech}(2a + 2bx) dx \\
 & \quad \downarrow 3042 \\
 & x \arctan(\tanh(a + bx)) - b \int x \csc\left(2ia + 2ibx + \frac{\pi}{2}\right) dx \\
 & \quad \downarrow 4668 \\
 & x \arctan(\tanh(a + bx)) - \\
 & b \left(-\frac{i \int \log(1 - ie^{2a+2bx}) dx}{2b} + \frac{i \int \log(1 + ie^{2a+2bx}) dx}{2b} + \frac{x \arctan(e^{2a+2bx})}{b} \right) \\
 & \quad \downarrow 2715 \\
 & x \arctan(\tanh(a + bx)) - \\
 & b \left(-\frac{i \int e^{-2a-2bx} \log(1 - ie^{2a+2bx}) de^{2a+2bx}}{4b^2} + \frac{i \int e^{-2a-2bx} \log(1 + ie^{2a+2bx}) de^{2a+2bx}}{4b^2} + \frac{x \arctan(e^{2a+2bx})}{b} \right) \\
 & \quad \downarrow 2838 \\
 & x \arctan(\tanh(a + bx)) - \\
 & b \left(\frac{x \arctan(e^{2a+2bx})}{b} - \frac{i \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b^2} + \frac{i \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b^2} \right)
 \end{aligned}$$

input `Int[ArcTan[Tanh[a + b*x]],x]`

output `x*ArcTan[Tanh[a + b*x]] - b*((x*ArcTan[E^(2*a + 2*b*x)])/b - ((I/4)*PolyLog[2, (-I)*E^(2*a + 2*b*x)]/b^2 + ((I/4)*PolyLog[2, I*E^(2*a + 2*b*x)]/b^2)`

Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_)], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5702 `Int[ArcTan[Tanh[(a_) + (b_)*(x_)]], x_Symbol] :> Simp[x*ArcTan[Tanh[a + b*x]], x] - Simp[b Int[x*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a, b}, x]`

Maple [A] (verified)

Time = 1.53 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.45

method	result
parts	$x \arctan(\tanh(bx + a)) - \frac{-a \arctan(e^{2bx+2a}) + \frac{i(bx+a)(\ln(1-ie^{2bx+2a}) - \ln(1+ie^{2bx+2a}))}{2}}{b} - \frac{i \operatorname{dilog}(1+ie^{2bx+2a})}{4}$
derivativedivides	$\frac{\operatorname{arctanh}(\tanh(bx+a)) \arctan(\tanh(bx+a)) - \frac{i \operatorname{arctanh}(\tanh(bx+a)) \left(\ln\left(1 - \frac{i(\tanh(bx+a)+1)^2}{1-\tanh(bx+a)^2}\right) - \ln\left(1 + \frac{i(\tanh(bx+a)+1)^2}{1-\tanh(bx+a)^2}\right) \right)}{2}}{b}$
default	$\frac{\operatorname{arctanh}(\tanh(bx+a)) \arctan(\tanh(bx+a)) - \frac{i \operatorname{arctanh}(\tanh(bx+a)) \left(\ln\left(1 - \frac{i(\tanh(bx+a)+1)^2}{1-\tanh(bx+a)^2}\right) - \ln\left(1 + \frac{i(\tanh(bx+a)+1)^2}{1-\tanh(bx+a)^2}\right) \right)}{2}}{b}$
risch	Expression too large to display

```
input int(arctan(tanh(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output x*arctan(tanh(b*x+a))-1/b*(-a*arctan(exp(2*b*x+2*a))+1/2*I*(b*x+a)*(ln(1-I*exp(2*b*x+2*a))-ln(1+I*exp(2*b*x+2*a)))-1/4*I*dilog(1+I*exp(2*b*x+2*a))+1/4*I*dilog(1-I*exp(2*b*x+2*a)))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(57) = 114.

Time = 0.15 (sec) , antiderivative size = 334, normalized size of antiderivative = 4.51

$$\int \arctan(\tanh(a + bx)) dx$$

$$= \frac{2bx \arctan\left(\frac{\sinh(bx+a)}{\cosh(bx+a)}\right) + (-ibx - ia) \log\left(\frac{1}{2}\sqrt{4i}(\cosh(bx+a) + \sinh(bx+a)) + 1\right) + (-ibx - ia) \log\left(\frac{1}{2}\sqrt{4i}(\cosh(bx+a) - \sinh(bx+a)) + 1\right)}{b}$$

```
input integrate(arctan(tanh(b*x+a)), x, algorithm="fricas")
```

output

```

1/2*(2*b*x*arctan(sinh(b*x + a)/cosh(b*x + a)) + (-I*b*x - I*a)*log(1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b*x - I*a)*log(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b*x + I*a)*log(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b*x + I*a)*log(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + I*a*log(I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) + I*a*log(-I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) - I*a*log(I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) - I*a*log(-I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) - I*dilog(1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - I*dilog(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) + I*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + I*dilog(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))))/b

```

Sympy [F]

$$\int \arctan(\tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(a + bx)) dx$$

input

```
integrate(atan(tanh(b*x+a)),x)
```

output

```
Integral(atan(tanh(a + b*x)), x)
```

Maxima [F]

$$\int \arctan(\tanh(a + bx)) dx = \int \arctan(\tanh(bx + a)) dx$$

input

```
integrate(arctan(tanh(b*x+a)),x, algorithm="maxima")
```

output

```
x*arctan((e^(2*b*x + 2*a) - 1)/(e^(2*b*x + 2*a) + 1)) - 2*b*integrate(x*e^(2*b*x + 2*a)/(e^(4*b*x + 4*a) + 1), x)
```

Giac [F]

$$\int \arctan(\tanh(a + bx)) dx = \int \arctan(\tanh(bx + a)) dx$$

input `integrate(arctan(tanh(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(tanh(b*x + a)), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(\tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(a + bx)) dx$$

input `int(atan(tanh(a + b*x)),x)`

output `int(atan(tanh(a + b*x)), x)`

Reduce [F]

$$\int \arctan(\tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a)) dx$$

input `int(atan(tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)),x)`

3.80 $\int \frac{\arctan(\tanh(a+bx))}{e+fx} dx$

Optimal result	629
Mathematica [N/A]	629
Rubi [N/A]	630
Maple [N/A]	630
Fricas [N/A]	631
Sympy [N/A]	631
Maxima [N/A]	631
Giac [N/A]	632
Mupad [N/A]	632
Reduce [N/A]	633

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \text{Int}\left(\frac{\arctan(\tanh(a + bx))}{e + fx}, x\right)$$

output `Defer(Int)(arctan(tanh(b*x+a))/(f*x+e), x)`

Mathematica [N/A]

Not integrable

Time = 10.98 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\arctan(\tanh(a + bx))}{e + fx} dx$$

input `Integrate[ArcTan[Tanh[a + b*x]]/(e + f*x), x]`

output `Integrate[ArcTan[Tanh[a + b*x]]/(e + f*x), x]`

Rubi [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx$$

↓ 7299

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx$$

input `Int[ArcTan[Tanh[a + b*x]]/(e + f*x),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.77 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(\tanh(bx + a))}{fx + e} dx$$

input `int(arctan(tanh(b*x+a))/(f*x+e),x)`

output `int(arctan(tanh(b*x+a))/(f*x+e),x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\arctan(\tanh(bx + a))}{fx + e} dx$$

input `integrate(arctan(tanh(b*x+a))/(f*x+e),x, algorithm="fricas")`

output `integral(arctan(tanh(b*x + a))/(f*x + e), x)`

Sympy [N/A]

Not integrable

Time = 3.80 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\operatorname{atan}(\tanh(a + bx))}{e + fx} dx$$

input `integrate(atan(tanh(b*x+a))/(f*x+e),x)`

output `Integral(atan(tanh(a + b*x))/(e + f*x), x)`

Maxima [N/A]

Not integrable

Time = 1.62 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\arctan(\tanh(bx + a))}{fx + e} dx$$

input `integrate(arctan(tanh(b*x+a))/(f*x+e),x, algorithm="maxima")`

output `integrate(arctan(tanh(b*x + a))/(f*x + e), x)`

Giac [N/A]

Not integrable

Time = 87.33 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.20

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\arctan(\tanh(bx + a))}{fx + e} dx$$

input `integrate(arctan(tanh(b*x+a))/(f*x+e),x, algorithm="giac")`

output `sage0*x`

Mupad [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\operatorname{atan}(\tanh(a + bx))}{e + fx} dx$$

input `int(atan(tanh(a + b*x))/(e + f*x),x)`

output `int(atan(tanh(a + b*x))/(e + f*x), x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\tanh(a + bx))}{e + fx} dx = \int \frac{\operatorname{atan}(\tanh(bx + a))}{fx + e} dx$$

input `int(atan(tanh(b*x+a))/(f*x+e),x)`output `int(atan(tanh(a + b*x))/(e + f*x),x)`

3.81 $\int x^2 \arctan(c + d \tanh(a + bx)) dx$

Optimal result	634
Mathematica [A] (warning: unable to verify)	635
Rubi [A] (verified)	636
Maple [C] (warning: unable to verify)	640
Fricas [B] (verification not implemented)	641
Sympy [F(-1)]	642
Maxima [F]	642
Giac [F]	642
Mupad [F(-1)]	643
Reduce [F]	643

Optimal result

Integrand size = 15, antiderivative size = 355

$$\begin{aligned}
 \int x^2 \arctan(c + d \tanh(a + bx)) dx = & \frac{1}{3}x^3 \arctan(c + d \tanh(a + bx)) \\
 & + \frac{1}{6}ix^3 \log\left(1 + \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right) \\
 & - \frac{1}{6}ix^3 \log\left(1 + \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right) \\
 & + \frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(i-c-d)e^{2a+2bx}}{i-c+d}\right)}{4b} \\
 & - \frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{(i+c+d)e^{2a+2bx}}{i+c-d}\right)}{4b} \\
 & - \frac{ix \operatorname{PolyLog}\left(3, -\frac{(i-c-d)e^{2a+2bx}}{i-c+d}\right)}{4b^2} \\
 & + \frac{ix \operatorname{PolyLog}\left(3, -\frac{(i+c+d)e^{2a+2bx}}{i+c-d}\right)}{4b^2} \\
 & + \frac{i \operatorname{PolyLog}\left(4, -\frac{(i-c-d)e^{2a+2bx}}{i-c+d}\right)}{8b^3} \\
 & - \frac{i \operatorname{PolyLog}\left(4, -\frac{(i+c+d)e^{2a+2bx}}{i+c-d}\right)}{8b^3}
 \end{aligned}$$

output

```
1/3*x^3*arctan(c+d*tanh(b*x+a))+1/6*I*x^3*ln(1+(I-c-d)*exp(2*b*x+2*a)/(I-c
+d))-1/6*I*x^3*ln(1+(I+c+d)*exp(2*b*x+2*a)/(I+c-d))+1/4*I*x^2*polylog(2,-(
I-c-d)*exp(2*b*x+2*a)/(I-c+d))/b-1/4*I*x^2*polylog(2,-(I+c+d)*exp(2*b*x+2*
a)/(I+c-d))/b-1/4*I*x*polylog(3,-(I-c-d)*exp(2*b*x+2*a)/(I-c+d))/b^2+1/4*I
*x*polylog(3,-(I+c+d)*exp(2*b*x+2*a)/(I+c-d))/b^2+1/8*I*polylog(4,-(I-c-d)
*exp(2*b*x+2*a)/(I-c+d))/b^3-1/8*I*polylog(4,-(I+c+d)*exp(2*b*x+2*a)/(I+c-
d))/b^3
```

Mathematica [A] (warning: unable to verify)

Time = 0.85 (sec) , antiderivative size = 438, normalized size of antiderivative = 1.23

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \frac{1}{3} x^3 \arctan(c + d \tanh(a + bx))$$

$$- \frac{d \left(4b^3 x^3 \log \left(1 + \frac{2(1+(c+d)^2)e^{2(a+bx)}}{2+2c^2-2d^2-4\sqrt{-d^2}} \right) - 4b^3 x^3 \log \left(1 + \frac{(1+(c+d)^2)e^{2(a+bx)}}{1+c^2-d^2+2\sqrt{-d^2}} \right) + 6b^2 x^2 \text{PolyLog} \left(2, \frac{(1+c^2+2cd+2d^2)e^{2(a+bx)}}{-1-c^2+d^2} \right) \right)}{24b^3 \sqrt{-d^2}}$$

input

```
Integrate[x^2*ArcTan[c + d*Tanh[a + b*x]],x]
```

output

```
(x^3*ArcTan[c + d*Tanh[a + b*x]])/3 - (d*(4*b^3*x^3*Log[1 + (2*(1 + (c + d)
^2)*E^(2*(a + b*x)))/(2 + 2*c^2 - 2*d^2 - 4*Sqrt[-d^2])] - 4*b^3*x^3*Log[
1 + ((1 + (c + d)^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2])] + 6*
b^2*x^2*PolyLog[2, ((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(-1 - c^2 + d
^2 + 2*Sqrt[-d^2])] - 6*b^2*x^2*PolyLog[2, -(((1 + (c + d)^2)*E^(2*(a + b*
x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2]))] - 6*b*x*PolyLog[3, ((1 + c^2 + 2*c*d
+ d^2)*E^(2*(a + b*x)))/(-1 - c^2 + d^2 + 2*Sqrt[-d^2])] + 6*b*x*PolyLog[
3, -(((1 + (c + d)^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2]))] +
3*PolyLog[4, (-2*(1 + (c + d)^2)*E^(2*(a + b*x)))/(2 + 2*c^2 - 2*d^2 - 4*S
qrt[-d^2])] - 3*PolyLog[4, -(((1 + (c + d)^2)*E^(2*(a + b*x)))/(1 + c^2 -
d^2 + 2*Sqrt[-d^2]))])/(24*b^3*Sqrt[-d^2])
```

Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.24, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5722, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(d \tanh(a + bx) + c) dx \\
 & \quad \downarrow \text{5722} \\
 & -\frac{1}{3}b(1 + i(c + d)) \int \frac{e^{2a+2bx} x^3}{-c + (-c - d + i)e^{2a+2bx} + d + i} dx + \frac{1}{3}b(1 - i(c + \\
 & d)) \int \frac{e^{2a+2bx} x^3}{c + (c + d + i)e^{2a+2bx} - d + i} dx + \frac{1}{3}x^3 \arctan(d \tanh(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & -\frac{1}{3}b(1 + i(c + d)) \left(\frac{x^3 \log \left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} - \frac{3 \int x^2 \log \left(\frac{e^{2a+2bx}(-c-d+i)}{-c+d+i} + 1 \right) dx}{2b(-c-d+i)} \right) + \\
 & \frac{1}{3}b(1 - i(c + d)) \left(\frac{x^3 \log \left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} - \frac{3 \int x^2 \log \left(\frac{e^{2a+2bx}(c+d+i)}{c-d+i} + 1 \right) dx}{2b(c+d+i)} \right) + \\
 & \quad \frac{1}{3}x^3 \arctan(d \tanh(a + bx) + c) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$d)) \left(\frac{x^3 \log \left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} - \frac{\frac{-\frac{1}{3}b(1+i(c+\int x \operatorname{PolyLog} \left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) dx}{b} - \frac{x^2 \operatorname{PolyLog} \left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b} \right)}{3}}{2b(-c-d+i)} \right) +$$

$$d)) \left(\frac{x^3 \log \left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} - \frac{\frac{\frac{1}{3}b(1-i(c+\int x \operatorname{PolyLog} \left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) dx}{b} - \frac{x^2 \operatorname{PolyLog} \left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b} \right)}{3}}{2b(c+d+i)} \right) +$$

$$\frac{1}{3}x^3 \arctan(d \tanh(a+bx) + c)$$

↓ 7163

$$d)) \left(\frac{x^3 \log \left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} - \frac{\frac{-\frac{1}{3}b(1+i(c+\frac{x \operatorname{PolyLog} \left(3, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b} - \frac{\int \operatorname{PolyLog} \left(3, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) dx}{b} - \frac{x^2 \operatorname{PolyLog} \left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b} \right)}{3}}{2b(-c-d+i)} \right) +$$

$$d)) \left(\frac{x^3 \log \left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} - \frac{\frac{\frac{1}{3}b(1-i(c+\frac{x \operatorname{PolyLog} \left(3, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b} - \frac{\int \operatorname{PolyLog} \left(3, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) dx}{b} - \frac{x^2 \operatorname{PolyLog} \left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b} \right)}{3}}{2b(c+d+i)} \right) +$$

$$\frac{1}{3}x^3 \arctan(d \tanh(a+bx) + c)$$

↓ 2720

$$d)) \left(\frac{x^3 \log\left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} - \frac{\frac{-\frac{1}{3}b(1+i(c + \frac{x \operatorname{PolyLog}\left(3, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right) de^{2a+2bx}}{b}}{4b^2}}{2b(-c-d+i)} \right.$$

$$d)) \left(\frac{x^3 \log\left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} - \frac{\frac{\frac{1}{3}b(1-i(c + \frac{x \operatorname{PolyLog}\left(3, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right) de^{2a+2bx}}{b}}{4b^2}}{2b(c+d+i)} \right.$$

$$\frac{1}{3}x^3 \arctan(d \tanh(a + bx) + c)$$

↓ 7143

$$d)) \left(\frac{x^3 \log\left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} - \frac{\frac{\frac{1}{3}x^3 \arctan(d \tanh(a + bx) + c) - \frac{1}{3}b(1+i(c + \frac{x \operatorname{PolyLog}\left(3, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{4b^2}}{b}}{2b(-c-d+i)} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b}}{2b(-c-d+i)} \right.$$

$$d)) \left(\frac{x^3 \log\left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} - \frac{\frac{\frac{1}{3}b(1-i(c + \frac{x \operatorname{PolyLog}\left(3, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{4b^2}}{b}}{2b(c+d+i)} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b}}{2b(c+d+i)} \right.$$

input Int[x^2*ArcTan[c + d*Tanh[a + b*x]], x]

output

```
(x^3*ArcTan[c + d*Tanh[a + b*x]])/3 - (b*(1 + I*(c + d))*((x^3*Log[1 + ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)])/(2*b*(I - c - d)) - (3*(-1/2*(x^2*PolyLog[2, -(((I - c - d)*E^(2*a + 2*b*x))/(I - c + d))])/b + ((x*PolyLog[3, -(((I - c - d)*E^(2*a + 2*b*x))/(I - c + d))]/(2*b) - PolyLog[4, -(((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)]/(4*b^2))/b))/(2*b*(I - c - d))))/3 + (b*(1 - I*(c + d))*((x^3*Log[1 + ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)])/(2*b*(I + c + d)) - (3*(-1/2*(x^2*PolyLog[2, -(((I + c + d)*E^(2*a + 2*b*x))/(I + c - d))])/b + ((x*PolyLog[3, -(((I + c + d)*E^(2*a + 2*b*x))/(I + c - d))]/(2*b) - PolyLog[4, -(((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)]/(4*b^2))/b))/(2*b*(I + c + d)))))/3
```

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*(f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```


rule 5722

```
Int[ArcTan[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] + (Simp[I*b*((I - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(I - c + d + (I - c - d)*E^(2*a + 2*b*x))], x], x] - Simp[I*b*((I + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(I + c - d + (I + c + d)*E^(2*a + 2*b*x))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c - d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 50.62 (sec) , antiderivative size = 6917, normalized size of antiderivative = 19.48

method	result	size
risch	Expression too large to display	6917

input `int(x^2*arctan(c+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1289 vs. $2(263) = 526$.

Time = 0.18 (sec) , antiderivative size = 1289, normalized size of antiderivative = 3.63

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(x^2*arctan(c+d*tanh(b*x+a)),x, algorithm="fricas")`

output

```

1/6*(2*b^3*x^3*arctan((c*cosh(b*x + a) + d*sinh(b*x + a))/cosh(b*x + a)) +
3*I*b^2*x^2*dilog(sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*
(cosh(b*x + a) + sinh(b*x + a))) + 3*I*b^2*x^2*dilog(-sqrt(-(c^2 - d^2 + 2
*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 3*I*
b^2*x^2*dilog(sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh
(b*x + a) + sinh(b*x + a))) - 3*I*b^2*x^2*dilog(-sqrt(-(c^2 - d^2 - 2*I*d
+ 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - I*a^3*log
(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(
b*x + a) + 2*(c^2 - d^2 - 2*I*d + 1)*sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 -
2*c*d + d^2 + 1))) - I*a^3*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2
*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 - 2*I*d + 1)*sqrt(-(
c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) + I*a^3*log(2*(c^2 + 2*c*
d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(
c^2 - d^2 + 2*I*d + 1)*sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 +
1))) + I*a^3*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d
+ d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 + 2*I*d + 1)*sqrt(-(c^2 - d^2 - 2*
I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) - 6*I*b*x*polylog(3, sqrt(-(c^2 - d^2 +
2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 6*
I*b*x*polylog(3, -sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*
(cosh(b*x + a) + sinh(b*x + a))) + 6*I*b*x*polylog(3, sqrt(-(c^2 - d^2 - ...

```

Sympy [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \text{Timed out}$$

input `integrate(x**2*atan(c+d*tanh(b*x+a)),x)`

output `Timed out`

Maxima [F]

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \int x^2 \arctan(d \tanh(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*tanh(b*x+a)),x, algorithm="maxima")`

output `1/3*x^3*arctan(((c*e^(2*a) + d*e^(2*a))*e^(2*b*x) + c - d)/(e^(2*b*x + 2*a) + 1)) - 4*b*d*integrate(1/3*x^3*e^(2*b*x + 2*a)/(c^2 - 2*c*d + d^2 + (c^2*e^(4*a) + 2*c*d*e^(4*a) + d^2*e^(4*a) + e^(4*a))*e^(4*b*x) + 2*(c^2*e^(2*a) - d^2*e^(2*a) + e^(2*a))*e^(2*b*x) + 1), x)`

Giac [F]

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \int x^2 \arctan(d \tanh(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan(d*tanh(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \int x^2 \operatorname{atan}(c + d \tanh(a + bx)) dx$$

input `int(x^2*atan(c + d*tanh(a + b*x)),x)`output `int(x^2*atan(c + d*tanh(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \arctan(c + d \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a)d + c) x^2 dx$$

input `int(x^2*atan(c+d*tanh(b*x+a)),x)`output `int(atan(tanh(a + b*x)*d + c)*x**2,x)`

3.82 $\int x \arctan(c + d \tanh(a + bx)) dx$

Optimal result	644
Mathematica [A] (warning: unable to verify)	645
Rubi [A] (verified)	645
Maple [C] (warning: unable to verify)	649
Fricas [B] (verification not implemented)	649
Sympy [F]	650
Maxima [F]	651
Giac [F]	651
Mupad [F(-1)]	651
Reduce [F]	652

Optimal result

Integrand size = 13, antiderivative size = 267

$$\begin{aligned} \int x \arctan(c + d \tanh(a + bx)) dx = & \frac{1}{2} x^2 \arctan(c + d \tanh(a + bx)) \\ & + \frac{1}{4} i x^2 \log \left(1 + \frac{(i - c - d) e^{2a + 2bx}}{i - c + d} \right) \\ & - \frac{1}{4} i x^2 \log \left(1 + \frac{(i + c + d) e^{2a + 2bx}}{i + c - d} \right) \\ & + \frac{i x \operatorname{PolyLog} \left(2, -\frac{(i - c - d) e^{2a + 2bx}}{i - c + d} \right)}{4b} \\ & - \frac{i x \operatorname{PolyLog} \left(2, -\frac{(i + c + d) e^{2a + 2bx}}{i + c - d} \right)}{4b} \\ & - \frac{i \operatorname{PolyLog} \left(3, -\frac{(i - c - d) e^{2a + 2bx}}{i - c + d} \right)}{8b^2} \\ & + \frac{i \operatorname{PolyLog} \left(3, -\frac{(i + c + d) e^{2a + 2bx}}{i + c - d} \right)}{8b^2} \end{aligned}$$

output

$$\frac{1}{2}x^2 \arctan(c+d \tanh(bx+a)) + \frac{1}{4}I^*x^2 \ln(1+(I-c-d)\exp(2bx+2a)/(I-c+d)) - \frac{1}{4}I^*x^2 \ln(1+(I+c+d)\exp(2bx+2a)/(I+c-d)) + \frac{1}{4}I^*x \operatorname{polylog}(2, -(I-c-d)\exp(2bx+2a)/(I-c+d))/b - \frac{1}{4}I^*x \operatorname{polylog}(2, -(I+c+d)\exp(2bx+2a)/(I+c-d))/b - \frac{1}{8}I^* \operatorname{polylog}(3, -(I-c-d)\exp(2bx+2a)/(I-c+d))/b^2 + \frac{1}{8}I^* \operatorname{polylog}(3, -(I+c+d)\exp(2bx+2a)/(I+c-d))/b^2$$
Mathematica [A] (warning: unable to verify)

Time = 0.73 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.24

$$\int x \arctan(c + d \tanh(a + bx)) dx = \frac{1}{2}x^2 \arctan(c + d \tanh(a + bx)) - \frac{d \left(2b^2x^2 \log \left(1 + \frac{2(1+(c+d)^2)e^{2(a+bx)}}{2+2c^2-2d^2-4\sqrt{-d^2}} \right) - 2b^2x^2 \log \left(1 + \frac{(1+(c+d)^2)e^{2(a+bx)}}{1+c^2-d^2+2\sqrt{-d^2}} \right) + 2bx \operatorname{PolyLog} \left(2, \frac{(1+c^2+2cd+d^2)}{-1-c^2+d^2+} \right) \right)}{b^2}$$

input

Integrate[x*ArcTan[c + d*Tanh[a + b*x]],x]

output

$$\frac{(x^2 \operatorname{ArcTan}[c + d \operatorname{Tanh}[a + b x]])}{2} - \frac{(d(2b^2x^2 \operatorname{Log}[1 + (2(1 + (c + d)^2)E^{2(a + b x)})]/(2 + 2c^2 - 2d^2 - 4\sqrt{-d^2})] - 2b^2x^2 \operatorname{Log}[1 + ((1 + (c + d)^2)E^{2(a + b x)})]/(1 + c^2 - d^2 + 2\sqrt{-d^2})] + 2bx \operatorname{PolyLog}[2, ((1 + c^2 + 2cd + d^2)E^{2(a + b x)})/(-1 - c^2 + d^2 + 2\sqrt{-d^2})] - 2bx \operatorname{PolyLog}[2, -(((1 + (c + d)^2)E^{2(a + b x)})/(1 + c^2 - d^2 + 2\sqrt{-d^2})]) - \operatorname{PolyLog}[3, (-2(1 + (c + d)^2)E^{2(a + b x)})/(2 + 2c^2 - 2d^2 - 4\sqrt{-d^2})] + \operatorname{PolyLog}[3, -(((1 + (c + d)^2)E^{2(a + b x)})/(1 + c^2 - d^2 + 2\sqrt{-d^2})])])/(8b^2\sqrt{-d^2})$$
Rubi [A] (verified)Time = 1.10 (sec) , antiderivative size = 343, normalized size of antiderivative = 1.28, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {5722, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \arctan(d \tanh(a + bx) + c) dx \\
 & \quad \downarrow 5722 \\
 & -\frac{1}{2}b(1 + i(c + d)) \int \frac{e^{2a+2bx} x^2}{-c + (-c - d + i)e^{2a+2bx} + d + i} dx + \frac{1}{2}b(1 - i(c + \\
 & d)) \int \frac{e^{2a+2bx} x^2}{c + (c + d + i)e^{2a+2bx} - d + i} dx + \frac{1}{2}x^2 \arctan(d \tanh(a + bx) + c) \\
 & \quad \downarrow 2620 \\
 & -\frac{1}{2}b(1 + i(c + d)) \left(\frac{x^2 \log\left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} - \frac{\int x \log\left(\frac{e^{2a+2bx}(-c-d+i)}{-c+d+i} + 1\right) dx}{b(-c-d+i)} \right) + \\
 & \frac{1}{2}b(1 - i(c + d)) \left(\frac{x^2 \log\left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} - \frac{\int x \log\left(\frac{e^{2a+2bx}(c+d+i)}{c-d+i} + 1\right) dx}{b(c+d+i)} \right) + \\
 & \frac{1}{2}x^2 \arctan(d \tanh(a + bx) + c) \\
 & \quad \downarrow 3011 \\
 & d)) \left(\frac{x^2 \log\left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} - \frac{\frac{1}{2}b(1 + i(c + \int \text{PolyLog}\left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right) dx - \frac{x \text{PolyLog}\left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b}\right)}{b(-c-d+i)}}{b(-c-d+i)} \right) + \\
 & d)) \left(\frac{x^2 \log\left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} - \frac{\frac{1}{2}b(1 - i(c + \int \text{PolyLog}\left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right) dx - \frac{x \text{PolyLog}\left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b}\right)}{b(c+d+i)}}{b(c+d+i)} \right) + \\
 & \frac{1}{2}x^2 \arctan(d \tanh(a + bx) + c) \\
 & \quad \downarrow 2720
 \end{aligned}$$

$$\begin{aligned}
 & d)) \left(\frac{x^2 \log\left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b} \right) \\
 & d)) \left(\frac{x^2 \log\left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right) de^{2a+2bx}}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b} \right) + \\
 & \frac{1}{2}x^2 \arctan(d \tanh(a + bx) + c)
 \end{aligned}$$

↓ 7143

$$\begin{aligned}
 & d)) \left(\frac{x^2 \log\left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} - \frac{\operatorname{PolyLog}\left(3, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b} \right) + \\
 & d)) \left(\frac{x^2 \log\left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} - \frac{\operatorname{PolyLog}\left(3, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{4b^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b} \right)
 \end{aligned}$$

input `Int[x*ArcTan[c + d*Tanh[a + b*x]],x]`

output `(x^2*ArcTan[c + d*Tanh[a + b*x]])/2 - (b*(1 + I*(c + d))*((x^2*Log[1 + ((I - c - d)*E^(2*a + 2*b*x)]/(I - c + d)]/(2*b*(I - c - d)) - (-1/2*(x*PolyLog[2, -(((I - c - d)*E^(2*a + 2*b*x)]/(I - c + d)))]/b + PolyLog[3, -(((I - c - d)*E^(2*a + 2*b*x)]/(I - c + d))]/(4*b^2))/(b*(I - c - d))))/2 + (b*(1 - I*(c + d))*((x^2*Log[1 + ((I + c + d)*E^(2*a + 2*b*x)]/(I + c - d)])/(2*b*(I + c + d)) - (-1/2*(x*PolyLog[2, -(((I + c + d)*E^(2*a + 2*b*x)]/(I + c - d)))]/b + PolyLog[3, -(((I + c + d)*E^(2*a + 2*b*x)]/(I + c - d))]/(4*b^2))/(b*(I + c + d))))/2`

Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5722

```
Int[ArcTan[(c_) + (d_)*Tanh[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_)^(m_
)), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tanh[a + b*x]]/(f*(m
+ 1))), x] + (Simp[I*b*((I - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E
^(2*a + 2*b*x))/(I - c + d + (I - c - d)*E^(2*a + 2*b*x))], x], x] - Simp[I*
b*((I + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(I + c
- d + (I + c + d)*E^(2*a + 2*b*x))], x], x] /; FreeQ[{a, b, c, d, e, f},
x] && IGtQ[m, 0] && NeQ[(c - d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_)^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.45 (sec) , antiderivative size = 6567, normalized size of antiderivative = 24.60

method	result	size
risch	Expression too large to display	6567

input `int(x*arctan(c+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1067 vs. $2(197) = 394$.

Time = 0.17 (sec) , antiderivative size = 1067, normalized size of antiderivative = 4.00

$$\int x \arctan(c + d \tanh(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctan(c+d*tanh(b*x+a)),x, algorithm="fricas")`

output

```

1/4*(2*b^2*x^2*arctan((c*cosh(b*x + a) + d*sinh(b*x + a))/cosh(b*x + a)) +
  2*I*b*x*dilog(sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cos
h(b*x + a) + sinh(b*x + a))) + 2*I*b*x*dilog(-sqrt(-(c^2 - d^2 + 2*I*d + 1
)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 2*I*b*x*dilo
g(sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) +
sinh(b*x + a))) - 2*I*b*x*dilog(-sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*
d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) + I*a^2*log(2*(c^2 + 2*c*d
+ d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(c^
2 - d^2 - 2*I*d + 1)*sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1
))) + I*a^2*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d +
d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 - 2*I*d + 1)*sqrt(-(c^2 - d^2 + 2*I*
d + 1)/(c^2 - 2*c*d + d^2 + 1))) - I*a^2*log(2*(c^2 + 2*c*d + d^2 + 1)*cos
h(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(c^2 - d^2 + 2*I*
d + 1)*sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) - I*a^2*log
(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(
b*x + a) - 2*(c^2 - d^2 + 2*I*d + 1)*sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 -
2*c*d + d^2 + 1))) + (I*b^2*x^2 - I*a^2)*log(sqrt(-(c^2 - d^2 + 2*I*d + 1)
/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^2*x^
2 - I*a^2)*log(-sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(co
sh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b^2*x^2 + I*a^2)*log(sqrt(-(c^2...

```

Sympy [F]

$$\int x \arctan(c + d \tanh(a + bx)) dx = \int x \operatorname{atan}(c + d \tanh(a + bx)) dx$$

input

```
integrate(x*atan(c+d*tanh(b*x+a)),x)
```

output

```
Integral(x*atan(c + d*tanh(a + b*x)), x)
```

Maxima [F]

$$\int x \arctan(c + d \tanh(a + bx)) dx = \int x \arctan(d \tanh(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*tanh(b*x+a)),x, algorithm="maxima")`

output `1/2*x^2*arctan(((c*e^(2*a) + d*e^(2*a))*e^(2*b*x) + c - d)/(e^(2*b*x + 2*a) + 1)) - 2*b*d*integrate(x^2*e^(2*b*x + 2*a)/(c^2 - 2*c*d + d^2 + (c^2*e^(4*a) + 2*c*d*e^(4*a) + d^2*e^(4*a) + e^(4*a))*e^(4*b*x) + 2*(c^2*e^(2*a) - d^2*e^(2*a) + e^(2*a))*e^(2*b*x) + 1), x)`

Giac [F]

$$\int x \arctan(c + d \tanh(a + bx)) dx = \int x \arctan(d \tanh(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan(d*tanh(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + d \tanh(a + bx)) dx = \int x \operatorname{atan}(c + d \tanh(a + bx)) dx$$

input `int(x*atan(c + d*tanh(a + b*x)),x)`

output `int(x*atan(c + d*tanh(a + b*x)), x)`

Reduce [F]

$$\int x \arctan(c + d \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) d + c) x dx$$

input `int(x*atan(c+d*tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)*d + c)*x,x)`

3.83 $\int \arctan(c + d \tanh(a + bx)) dx$

Optimal result	653
Mathematica [A] (warning: unable to verify)	654
Rubi [A] (verified)	654
Maple [B] (verified)	656
Fricas [B] (verification not implemented)	657
Sympy [F]	658
Maxima [F]	659
Giac [F]	659
Mupad [F(-1)]	659
Reduce [F]	660

Optimal result

Integrand size = 11, antiderivative size = 174

$$\int \arctan(c + d \tanh(a + bx)) dx = x \arctan(c + d \tanh(a + bx)) + \frac{1}{2}ix \log\left(1 + \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right) - \frac{1}{2}ix \log\left(1 + \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right) + \frac{i \operatorname{PolyLog}\left(2, -\frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right)}{4b} - \frac{i \operatorname{PolyLog}\left(2, -\frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right)}{4b}$$

output

```
x*arctan(c+d*tanh(b*x+a))+1/2*I*x*ln(1+(I-c-d)*exp(2*b*x+2*a)/(I-c+d))-1/2
*I*x*ln(1+(I+c+d)*exp(2*b*x+2*a)/(I+c-d))+1/4*I*polylog(2,-(I-c-d)*exp(2*b
*x+2*a)/(I-c+d))/b-1/4*I*polylog(2,-(I+c+d)*exp(2*b*x+2*a)/(I+c-d))/b
```

Mathematica [A] (warning: unable to verify)

Time = 2.22 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.66

$$\int \arctan(c + d \tanh(a + bx)) dx = x \arctan(c + d \tanh(a + bx)) + \frac{4a\sqrt{-d^2} \arctan\left(\frac{1+c^2-d^2+(1+c^2+2cd+d^2)e^{2(a+bx)}}{2d}\right) - 2d(a+bx) \log\left(1 + \frac{2(1+(c+d)^2)e^{2(a+bx)}}{2+2c^2-2d^2-4\sqrt{-d^2}}\right) + 2d(a+bx) \log\left(1 + \frac{2(1+(c+d)^2)e^{2(a+bx)}}{2+2c^2-2d^2-4\sqrt{-d^2}}\right)}{4b\sqrt{-d^2}}$$

input

```
Integrate[ArcTan[c + d*Tanh[a + b*x]], x]
```

output

```
x*ArcTan[c + d*Tanh[a + b*x]] + (4*a*Sqrt[-d^2]*ArcTan[(1 + c^2 - d^2 + (1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(2*d)] - 2*d*(a + b*x)*Log[1 + (2*(1 + (c + d)^2)*E^(2*(a + b*x)))/(2 + 2*c^2 - 2*d^2 - 4*Sqrt[-d^2])] + 2*d*(a + b*x)*Log[1 + ((1 + (c + d)^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2])] + d*PolyLog[2, -(((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2]))] - d*PolyLog[2, ((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(-1 - c^2 + d^2 + 2*Sqrt[-d^2])]/(4*b*Sqrt[-d^2])
```

Rubi [A] (verified)Time = 0.66 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.36, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {5714, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(d \tanh(a + bx) + c) dx$$

$$\downarrow 5714$$

$$-b(1 + i(c + d)) \int \frac{e^{2a+2bx} x}{-c + (-c - d + i)e^{2a+2bx} + d + i} dx + b(1 - i(c + d)) \int \frac{e^{2a+2bx} x}{c + (c + d + i)e^{2a+2bx} - d + i} dx + x \arctan(d \tanh(a + bx) + c)$$

$$\downarrow 2620$$

$$-b(1+i(c+d)) \left(\frac{x \log \left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} - \frac{\int \log \left(\frac{e^{2a+2bx}(-c-d+i)}{-c+d+i} + 1 \right) dx}{2b(-c-d+i)} \right) + b(1-i(c+d)) \left(\frac{x \log \left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} - \frac{\int \log \left(\frac{e^{2a+2bx}(c+d+i)}{c-d+i} + 1 \right) dx}{2b(c+d+i)} \right) + x \arctan(d \tanh(a+bx) + c)$$

↓ 2715

$$d) \left(\frac{x \log \left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} - \frac{\int e^{-2a-2bx} \log \left(\frac{e^{2a+2bx}(-c-d+i)}{-c+d+i} + 1 \right) de^{2a+2bx}}{4b^2(-c-d+i)} \right) + b(1-i(c+d)) \left(\frac{x \log \left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} - \frac{\int e^{-2a-2bx} \log \left(\frac{e^{2a+2bx}(c+d+i)}{c-d+i} + 1 \right) de^{2a+2bx}}{4b^2(c+d+i)} \right) + x \arctan(d \tanh(a+bx) + c)$$

↓ 2838

$$d) \left(\frac{x \arctan(d \tanh(a+bx) + c) - b(1+i(c+d)) \operatorname{PolyLog} \left(2, -\frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{4b^2(-c-d+i)} + \frac{x \log \left(1 + \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) + b(1-i(c+d)) \left(\frac{\operatorname{PolyLog} \left(2, -\frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{4b^2(c+d+i)} + \frac{x \log \left(1 + \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right)$$

input `Int[ArcTan[c + d*Tanh[a + b*x]],x]`

output `x*ArcTan[c + d*Tanh[a + b*x]] - b*(1 + I*(c + d))*((x*Log[1 + ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)])/(2*b*(I - c - d)) + PolyLog[2, -(((I - c - d)*E^(2*a + 2*b*x))/(I - c + d))]/(4*b^2*(I - c - d))) + b*(1 - I*(c + d))*((x*Log[1 + ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)])/(2*b*(I + c + d)) + PolyLog[2, -(((I + c + d)*E^(2*a + 2*b*x))/(I + c - d))]/(4*b^2*(I + c + d)))`

Definitions of rubi rules used

rule 2620 $\text{Int}[\frac{((F_)^{((g_.) * (e_.) + (f_.) * (x_.))})^{(n_.)} * ((c_.) + (d_.) * (x_.))^{(m_.)}}{((a_.) + (b_.) * (F_)^{((g_.) * (e_.) + (f_.) * (x_.))})^{(n_.)}}, x_Symbol] \rightarrow \text{Simp}[\frac{(c + d*x)^m}{(b*f*g*n*\text{Log}[F])} * \text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x] - \text{Simp}[d*(m/(b*f*g*n*\text{Log}[F])) \text{Int}[(c + d*x)^{(m-1)} * \text{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x], x] /; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \&\& \text{IGtQ}[m, 0]$

rule 2715 $\text{Int}[\text{Log}[(a_) + (b_.) * (F_)^{((e_.) * (c_.) + (d_.) * (x_.))}]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n)], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \&\& \text{GtQ}[a, 0]$

rule 2838 $\text{Int}[\text{Log}[(c_.) * (d_) + (e_.) * (x_)^{(n_.)}] / (x_), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \&\& \text{EqQ}[c*d, 1]$

rule 5714 $\text{Int}[\text{ArcTan}[(c_.) + (d_.) * \text{Tanh}[(a_.) + (b_.) * (x_)]], x_Symbol] \rightarrow \text{Simp}[x * \text{ArcTan}[c + d * \text{Tanh}[a + b*x]], x] + (\text{Simp}[I*b*(I - c - d) \text{Int}[x*(E^{(2*a + 2*b*x)}) / (I - c + d + (I - c - d)*E^{(2*a + 2*b*x)})], x], x] - \text{Simp}[I*b*(I + c + d) \text{Int}[x*(E^{(2*a + 2*b*x)}) / (I + c - d + (I + c + d)*E^{(2*a + 2*b*x)})], x], x] /; \text{FreeQ}\{a, b, c, d\}, x\} \&\& \text{NeQ}[(c - d)^2, -1]$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 351 vs. $2(150) = 300$.

Time = 1.43 (sec) , antiderivative size = 352, normalized size of antiderivative = 2.02

method	result
derivativedivides	$\frac{\arctan(c+d \tanh(bx+a))d \ln(-d \tanh(bx+a)-d) - \arctan(c+d \tanh(bx+a))d \ln(-d \tanh(bx+a)+d)}{2} + \frac{d^2 \left(\frac{i \ln(-d \tanh(bx+a)+d) \ln(-d \tanh(bx+a)-d)}{2} \right)}{2}$
default	$\frac{\arctan(c+d \tanh(bx+a))d \ln(-d \tanh(bx+a)-d) - \arctan(c+d \tanh(bx+a))d \ln(-d \tanh(bx+a)+d)}{2} + \frac{d^2 \left(\frac{i \ln(-d \tanh(bx+a)+d) \ln(-d \tanh(bx+a)-d)}{2} \right)}{2}$
risch	Expression too large to display

```
input int(arctan(c+d*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

```
output 1/b/d*(1/2*arctan(c+d*tanh(b*x+a))*d*ln(-d*tanh(b*x+a)-d)-1/2*arctan(c+d*tanh(b*x+a))*d*ln(-d*tanh(b*x+a)+d)+1/2*d^2*(1/d*(1/2*I*ln(-d*tanh(b*x+a)+d))*ln((I+d*tanh(b*x+a)+c)/(I+c+d))-1/2*I*ln(-d*tanh(b*x+a)+d)*ln((I-d*tanh(b*x+a)-c)/(I-c-d))+1/2*I*dilog((I+d*tanh(b*x+a)+c)/(I+c+d))-1/2*I*dilog((I-d*tanh(b*x+a)-c)/(I-c-d)))-1/d*(1/2*I*ln(-d*tanh(b*x+a)-d)*ln((I+d*tanh(b*x+a)+c)/(I+c-d))-1/2*I*ln(-d*tanh(b*x+a)-d)*ln((I-d*tanh(b*x+a)-c)/(I-c+d)))+1/2*I*dilog((I+d*tanh(b*x+a)+c)/(I+c-d))-1/2*I*dilog((I-d*tanh(b*x+a)-c)/(I-c+d))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 825 vs. 2(128) = 256.

Time = 0.25 (sec) , antiderivative size = 825, normalized size of antiderivative = 4.74

$$\int \arctan(c + d \tanh(a + bx)) dx = \text{Too large to display}$$

```
input integrate(arctan(c+d*tanh(b*x+a)),x, algorithm="fricas")
```

output

```

1/2*(2*b*x*arctan((c*cosh(b*x + a) + d*sinh(b*x + a))/cosh(b*x + a)) - I*a
*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*s
inh(b*x + a) + 2*(c^2 - d^2 - 2*I*d + 1)*sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^
2 - 2*c*d + d^2 + 1))) - I*a*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) +
2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 - 2*I*d + 1)*sqrt(
-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) + I*a*log(2*(c^2 + 2*c*
d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(
c^2 - d^2 + 2*I*d + 1)*sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 +
1))) + I*a*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d +
d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 + 2*I*d + 1)*sqrt(-(c^2 - d^2 - 2*I*
d + 1)/(c^2 - 2*c*d + d^2 + 1))) + (I*b*x + I*a)*log(sqrt(-(c^2 - d^2 + 2*
I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (
I*b*x + I*a)*log(-sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*
(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b*x - I*a)*log(sqrt(-(c^2 - d^2
- 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1)
+ (-I*b*x - I*a)*log(-sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 +
1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + I*dilog(sqrt(-(c^2 - d^2 + 2*I*
d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) + I*dilog
(-sqrt(-(c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) +
sinh(b*x + a))) - I*dilog(sqrt(-(c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + ...

```

Sympy [F]

$$\int \arctan(c + d \tanh(a + bx)) dx = \int \operatorname{atan}(c + d \tanh(a + bx)) dx$$

input

```
integrate(atan(c+d*tanh(b*x+a)),x)
```

output

```
Integral(atan(c + d*tanh(a + b*x)), x)
```

Maxima [F]

$$\int \arctan(c + d \tanh(a + bx)) dx = \int \arctan(d \tanh(bx + a) + c) dx$$

input `integrate(arctan(c+d*tanh(b*x+a)),x, algorithm="maxima")`

output `-4*b*d*integrate(x*e^(2*b*x + 2*a)/(c^2 - 2*c*d + d^2 + (c^2*e^(4*a) + 2*c*d*e^(4*a) + d^2*e^(4*a) + e^(4*a))*e^(4*b*x) + 2*(c^2*e^(2*a) - d^2*e^(2*a) + e^(2*a))*e^(2*b*x) + 1), x) + x*arctan(((c*e^(2*a) + d*e^(2*a))*e^(2*b*x) + c - d)/(e^(2*b*x + 2*a) + 1))`

Giac [F]

$$\int \arctan(c + d \tanh(a + bx)) dx = \int \arctan(d \tanh(bx + a) + c) dx$$

input `integrate(arctan(c+d*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(d*tanh(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + d \tanh(a + bx)) dx = \int \operatorname{atan}(c + d \tanh(a + bx)) dx$$

input `int(atan(c + d*tanh(a + b*x)),x)`

output `int(atan(c + d*tanh(a + b*x)), x)`

Reduce [F]

$$\int \arctan(c + d \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) d + c) dx$$

input `int(atan(c+d*tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)*d + c),x)`

3.84 $\int \frac{\arctan(c+d \tanh(a+bx))}{x} dx$

Optimal result	661
Mathematica [N/A]	661
Rubi [N/A]	662
Maple [N/A]	662
Fricas [N/A]	663
Sympy [F(-1)]	663
Maxima [N/A]	663
Giac [N/A]	664
Mupad [N/A]	664
Reduce [N/A]	664

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + d \tanh(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+d*tanh(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 5.83 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \int \frac{\arctan(c + d \tanh(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + d*Tanh[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + d*Tanh[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(d \tanh(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\arctan(d \tanh(a + bx) + c)}{x} dx$$

input `Int[ArcTan[c + d*Tanh[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + d \tanh(bx + a))}{x} dx$$

input `int(arctan(c+d*tanh(b*x+a))/x,x)`

output `int(arctan(c+d*tanh(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \int \frac{\arctan(d \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*tanh(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctan(d*tanh(b*x + a) + c)/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+d*tanh(b*x+a))/x,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 1.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \int \frac{\arctan(d \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*tanh(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctan(d*tanh(b*x + a) + c)/x, x)`

Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \int \frac{\arctan(d \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*tanh(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctan(d*tanh(b*x + a) + c)/x, x)`

Mupad [N/A]

Not integrable

Time = 0.61 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + d \tanh(a + bx))}{x} dx$$

input `int(atan(c + d*tanh(a + b*x))/x,x)`

output `int(atan(c + d*tanh(a + b*x))/x, x)`

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\tanh(bx + a) d + c)}{x} dx$$

input `int(atan(c+d*tanh(b*x+a))/x,x)`

output `int(atan(tanh(a + b*x)*d + c)/x,x)`

3.85 $\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx$

Optimal result	666
Mathematica [A] (verified)	667
Rubi [A] (verified)	667
Maple [C] (warning: unable to verify)	670
Fricas [B] (verification not implemented)	671
Sympy [F(-2)]	672
Maxima [A] (verification not implemented)	672
Giac [F]	673
Mupad [F(-1)]	673
Reduce [F]	673

Optimal result

Integrand size = 19, antiderivative size = 142

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx = -\frac{1}{12}ibx^4 + \frac{1}{3}x^3 \arctan(c + (i + c) \tanh(a + bx))$$

$$+ \frac{1}{6}ix^3 \log(1 + ice^{2a+2bx})$$

$$+ \frac{ix^2 \operatorname{PolyLog}(2, -ice^{2a+2bx})}{4b}$$

$$- \frac{ix \operatorname{PolyLog}(3, -ice^{2a+2bx})}{4b^2}$$

$$+ \frac{i \operatorname{PolyLog}(4, -ice^{2a+2bx})}{8b^3}$$

output

```
-1/12*I*b*x^4+1/3*x^3*arctan(c+(I+c)*tanh(b*x+a))+1/6*I*x^3*ln(1+I*c*exp(2
*b*x+2*a))+1/4*I*x^2*polylog(2,-I*c*exp(2*b*x+2*a))/b-1/4*I*x*polylog(3,-I
*c*exp(2*b*x+2*a))/b^2+1/8*I*polylog(4,-I*c*exp(2*b*x+2*a))/b^3
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.94

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= \frac{8b^3 x^3 \arctan(c + (i + c) \tanh(a + bx)) + 4ib^3 x^3 \log\left(1 - \frac{ie^{-2(a+bx)}}{c}\right) - 6ib^2 x^2 \text{PolyLog}\left(2, \frac{ie^{-2(a+bx)}}{c}\right) - 6ibx \text{PolyLog}\left(3, \frac{ie^{-2(a+bx)}}{c}\right) - 6i \text{PolyLog}\left(4, \frac{ie^{-2(a+bx)}}{c}\right)}{24b^3}$$

input

```
Integrate[x^2*ArcTan[c + (I + c)*Tanh[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTan[c + (I + c)*Tanh[a + b*x]] + (4*I)*b^3*x^3*Log[1 - I/(c*E^(2*(a + b*x)))] - (6*I)*b^2*x^2*PolyLog[2, I/(c*E^(2*(a + b*x)))] - (6*I)*b*x*PolyLog[3, I/(c*E^(2*(a + b*x)))] - (3*I)*PolyLog[4, I/(c*E^(2*(a + b*x)))])/(24*b^3)
```

Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {5718, 25, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(c + (c + i) \tanh(a + bx)) dx$$

$$\downarrow \text{5718}$$

$$\frac{1}{3}x^3 \arctan(c + (c + i) \tanh(a + bx)) - \frac{1}{3}b \int -\frac{x^3}{i - ce^{2a+2bx}} dx$$

$$\downarrow \text{25}$$

$$\frac{1}{3}b \int \frac{x^3}{i - ce^{2a+2bx}} dx + \frac{1}{3}x^3 \arctan(c + (c + i) \tanh(a + bx))$$

$$\downarrow \text{2615}$$

$$\frac{1}{3}b \left(-ic \int \frac{e^{2a+2bx} x^3}{i - ce^{2a+2bx}} dx - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c+i) \tanh(a+bx))$$

↓ 2620

$$\frac{1}{3}b \left(-ic \left(\frac{3 \int x^2 \log(ie^{2a+2bx}c + 1) dx}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c+i) \tanh(a+bx))$$

↓ 3011

$$\frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\int x \text{PolyLog}(2, -ice^{2a+2bx}) dx}{b} - \frac{x^2 \text{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c+i) \tanh(a+bx))$$

↓ 7163

$$\frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\frac{x \text{PolyLog}(3, -ice^{2a+2bx})}{2b} - \frac{\int \text{PolyLog}(3, -ice^{2a+2bx}) dx}{b}}{2bc} - \frac{x^2 \text{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) \right) + \frac{1}{3}x^3 \arctan(c + (c+i) \tanh(a+bx))$$

↓ 2720

$$\frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\frac{x \text{PolyLog}(3, -ice^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \text{PolyLog}(3, -ice^{2a+2bx}) de^{2a+2bx}}{4b^2}}{b}}{2bc} - \frac{x^2 \text{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) \right) + \frac{1}{3}x^3 \arctan(c + (c+i) \tanh(a+bx))$$

↓ 7143

$$\frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\frac{x \text{PolyLog}(3, -ice^{2a+2bx})}{2b} - \frac{\text{PolyLog}(4, -ice^{2a+2bx})}{4b^2}}{b}}{2bc} - \frac{x^2 \text{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c+i) \tanh(a+bx))$$

input `Int[x^2*ArcTan[c + (I + c)*Tanh[a + b*x]],x]`

output `(x^3*ArcTan[c + (I + c)*Tanh[a + b*x]])/3 + (b*((-1/4*I)*x^4 - I*c*(-1/2*(x^3*Log[1 + I*c*E^(2*a + 2*b*x)])/(b*c) + (3*(-1/2*(x^2*PolyLog[2, (-I)*c*E^(2*a + 2*b*x)])/b + ((x*PolyLog[3, (-I)*c*E^(2*a + 2*b*x)])/(2*b) - PolyLog[4, (-I)*c*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b*c)))/3`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5718

```
Int[ArcTan[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_)^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_)^(m_.))*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.14 (sec) , antiderivative size = 1406, normalized size of antiderivative = 9.90

method	result	size
risch	Expression too large to display	1406

input

```
int(x^2*arctan(c+(1+c)*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

-1/12*I*b*c/(I+c)*x^4+1/12*Pi*(csgn(I*(2*exp(2*b*x+2*a)*c-2*I))*csgn(I*(2*
exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))^2-csgn(I*(2*exp(2*b*x+2*a)*c-2*I
))*csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))*csgn(I/(exp(2*b*x+2
*a)+1))-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(2*I*exp(2*
b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^2+csgn(I*(2*I*exp(2*b*x+2
*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(e
xp(2*b*x+2*a)+1))*csgn(I/(exp(2*b*x+2*a)+1))-csgn(I*(2*exp(2*b*x+2*a)*c-2*
I)/(exp(2*b*x+2*a)+1))^3+csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1
))^2*csgn(I/(exp(2*b*x+2*a)+1))+csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x
+2*a)+1))*csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))^2-csgn(I*(2*ex
p(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))*csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp
(2*b*x+2*a)+1))+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+
2*a)+1))^3-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+
1))^2*csgn(I/(exp(2*b*x+2*a)+1))-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*
a)*c)/(exp(2*b*x+2*a)+1))*csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(ex
p(2*b*x+2*a)+1))^2+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b
*x+2*a)+1))*csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1
))+csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^3-csgn
((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^2+csgn((2*exp
(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))^3-csgn((2*exp(2*b*x+2*a)*c-2*I)/...

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs. $2(105) = 210$.

Time = 0.14 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.06

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= \frac{-i b^4 x^4 + 2i b^3 x^3 \log\left(-\frac{(c+i)e^{(2bx+2a)}}{ce^{(2bx+2a)}-i}\right) + 6i b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4i} ce^{(bx+a)}\right) + 6i b^2 x^2 \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{-4i} ce^{(bx+a)}\right) + i}{}$$

input

```
integrate(x^2*arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="fricas")
```


output

```
1/12*(-I*b^4*x^4 + 2*I*b^3*x^3*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) - I)) + 6*I*b^2*x^2*dilog(1/2*sqrt(-4*I*c)*e^(b*x + a)) + 6*I*b^2*x^2*dilog(-1/2*sqrt(-4*I*c)*e^(b*x + a)) + I*a^4 - 2*I*a^3*log(1/2*(2*c*e^(b*x + a) + I*sqrt(-4*I*c))/c) - 2*I*a^3*log(1/2*(2*c*e^(b*x + a) - I*sqrt(-4*I*c))/c) - 12*I*b*x*polylog(3, 1/2*sqrt(-4*I*c)*e^(b*x + a)) - 12*I*b*x*polylog(3, -1/2*sqrt(-4*I*c)*e^(b*x + a)) - 2*(-I*b^3*x^3 - I*a^3)*log(1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) - 2*(-I*b^3*x^3 - I*a^3)*log(-1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) + 12*I*polylog(4, 1/2*sqrt(-4*I*c)*e^(b*x + a)) + 12*I*polylog(4, -1/2*sqrt(-4*I*c)*e^(b*x + a)))/b^3
```

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input

```
integrate(x**2*atan(c+(I+c)*tanh(b*x+a)),x)
```

output

```
Exception raised: CoercionFailed >> Cannot convert _t0**2 + exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]
```

Maxima [A] (verification not implemented)

Time = 1.37 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.91

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx = \frac{1}{3} x^3 \arctan((c + i) \tanh(bx + a) + c) + \frac{4}{9} \left(\frac{3x^4}{4ic - 4} - \frac{4b^3x^3 \log(i ce^{(2bx+2a)} + 1) + 6b^2x^2 \text{Li}_2(-i ce^{(2bx+2a)}) - 6bx \text{Li}_3(-i ce^{(2bx+2a)}) + 3 \text{Li}_4(-i ce^{(2bx+2a)})}{-2b^4(-ic + 1)} \right)$$

input

```
integrate(x^2*arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctan((c + I)*tanh(b*x + a) + c) + 4/9*(3*x^4/(4*I*c - 4) - (4*b^
3*x^3*log(I*c*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog(-I*c*e^(2*b*x + 2*a))
- 6*b*x*polylog(3, -I*c*e^(2*b*x + 2*a)) + 3*polylog(4, -I*c*e^(2*b*x + 2
*a)))/(b^4*(2*I*c - 2))*b*(c + I)
```

Giac [F]

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx = \int x^2 \arctan((c + i) \tanh(bx + a) + c) dx$$

input

```
integrate(x^2*arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="giac")
```

output

```
integrate(x^2*arctan((c + I)*tanh(b*x + a) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx = \int x^2 \operatorname{atan}(c + \tanh(a + bx) (c + 1i)) dx$$

input

```
int(x^2*atan(c + tanh(a + b*x)*(c + 1i)),x)
```

output

```
int(x^2*atan(c + tanh(a + b*x)*(c + 1i)), x)
```

Reduce [F]

$$\int x^2 \arctan(c + (i + c) \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) c + \tanh(bx + a) i + c) x^2 dx$$

input

```
int(x^2*atan(c+(I+c)*tanh(b*x+a)),x)
```

output `int(atan(tanh(a + b*x)*c + tanh(a + b*x)*i + c)*x**2,x)`

3.86 $\int x \arctan(c + (i + c) \tanh(a + bx)) dx$

Optimal result	675
Mathematica [A] (verified)	676
Rubi [A] (verified)	676
Maple [C] (warning: unable to verify)	679
Fricas [B] (verification not implemented)	680
Sympy [F(-2)]	680
Maxima [A] (verification not implemented)	681
Giac [F]	681
Mupad [F(-1)]	681
Reduce [F]	682

Optimal result

Integrand size = 17, antiderivative size = 113

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx = -\frac{1}{6}ibx^3 + \frac{1}{2}x^2 \arctan(c + (i + c) \tanh(a + bx)) + \frac{1}{4}ix^2 \log(1 + ice^{2a+2bx}) + \frac{ix \operatorname{PolyLog}(2, -ice^{2a+2bx})}{4b} - \frac{i \operatorname{PolyLog}(3, -ice^{2a+2bx})}{8b^2}$$

output

```
-1/6*I*b*x^3+1/2*x^2*arctan(c+(I+c)*tanh(b*x+a))+1/4*I*x^2*ln(1+I*c*exp(2*b*x+2*a))+1/4*I*x*polylog(2,-I*c*exp(2*b*x+2*a))/b-1/8*I*polylog(3,-I*c*exp(2*b*x+2*a))/b^2
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.91

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= \frac{2b^2x^2 \left(2 \arctan(c + (i + c) \tanh(a + bx)) + i \log \left(1 - \frac{ie^{-2(a+bx)}}{c} \right) \right) - 2ibx \operatorname{PolyLog} \left(2, \frac{ie^{-2(a+bx)}}{c} \right) - i \operatorname{PolyLog} \left(3, \frac{ie^{-2(a+bx)}}{c} \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c + (I + c)*Tanh[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTan[c + (I + c)*Tanh[a + b*x]] + I*Log[1 - I/(c*E^(2*(a + b*x))])) - (2*I)*b*x*PolyLog[2, I/(c*E^(2*(a + b*x)))] - I*PolyLog[3, I/(c*E^(2*(a + b*x)))]/(8*b^2)
```

Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.17, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {5718, 25, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(c + (c + i) \tanh(a + bx)) dx$$

$$\downarrow \text{5718}$$

$$\frac{1}{2}x^2 \arctan(c + (c + i) \tanh(a + bx)) - \frac{1}{2}b \int -\frac{x^2}{i - ce^{2a+2bx}} dx$$

$$\downarrow \text{25}$$

$$\frac{1}{2}b \int \frac{x^2}{i - ce^{2a+2bx}} dx + \frac{1}{2}x^2 \arctan(c + (c + i) \tanh(a + bx))$$

$$\downarrow \text{2615}$$

$$\frac{1}{2}b \left(-ic \int \frac{e^{2a+2bx} x^2}{i - ce^{2a+2bx}} dx - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c + (c + i) \tanh(a + bx))$$

$$\frac{1}{2}b \left(-ic \left(\frac{\int x \log(ice^{2a+2bx}c+1) dx}{bc} - \frac{x^2 \log(1+ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c+(c+i) \tanh(a+bx))$$

↓ 2620

$$\frac{1}{2}b \left(-ic \left(\frac{\int \text{PolyLog}(2,-ice^{2a+2bx}) dx}{2b} - \frac{x \text{PolyLog}(2,-ice^{2a+2bx})}{2b} - \frac{x^2 \log(1+ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c+(c+i) \tanh(a+bx))$$

↓ 3011

$$\frac{1}{2}b \left(-ic \left(\frac{\int e^{-2a-2bx} \text{PolyLog}(2,-ice^{2a+2bx}) de^{2a+2bx}}{4b^2} - \frac{x \text{PolyLog}(2,-ice^{2a+2bx})}{2b} - \frac{x^2 \log(1+ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c+(c+i) \tanh(a+bx))$$

↓ 2720

$$\frac{1}{2}b \left(-ic \left(\frac{\text{PolyLog}(3,-ice^{2a+2bx})}{4b^2} - \frac{x \text{PolyLog}(2,-ice^{2a+2bx})}{2b} - \frac{x^2 \log(1+ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c+(c+i) \tanh(a+bx))$$

↓ 7143

input `Int[x*ArcTan[c + (I + c)*Tanh[a + b*x]],x]`

output `(x^2*ArcTan[c + (I + c)*Tanh[a + b*x]])/2 + (b*((-1/3*I)*x^3 - I*c*(-1/2*(x^2*Log[1 + I*c*E^(2*a + 2*b*x)])/(b*c) + (-1/2*(x*PolyLog[2, (-I)*c*E^(2*a + 2*b*x)])/b + PolyLog[3, (-I)*c*E^(2*a + 2*b*x)]/(4*b^2))/(b*c)))/2`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \text{ :> Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2615 $\text{Int}[\text{((c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{(m}_.)}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{(g}_.) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{(n}_.)}))})^{\text{(n}_.)}), \text{x_Symbol}] \text{ :> Simp}[(\text{c} + \text{d} * \text{x})^{\text{(m} + 1)} / (\text{a} * \text{d} * (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b/a} \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m}} * ((\text{F}^{\text{(g}} * (\text{e} + \text{f} * \text{x}))^{\text{n}}) / (\text{a} + \text{b} * (\text{F}^{\text{(g}} * (\text{e} + \text{f} * \text{x}))^{\text{n}}))], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2620 $\text{Int}[\text{((F}_.)^{\text{(g}_.) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{(n}_.)})) * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{(m}_.)}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{(g}_.) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{(n}_.)}))})^{\text{(n}_.)}), \text{x_Symbol}] \text{ :> Simp}[(\text{c} + \text{d} * \text{x})^{\text{m}} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) * \text{Log}[1 + \text{b} * ((\text{F}^{\text{(g}} * (\text{e} + \text{f} * \text{x}))^{\text{n}}) / \text{a})], \text{x}] - \text{Simp}[\text{d} * (\text{m} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{(m} - 1)} * \text{Log}[1 + \text{b} * ((\text{F}^{\text{(g}} * (\text{e} + \text{f} * \text{x}))^{\text{n}}) / \text{a})], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2720 $\text{Int}[\text{u}_, \text{x_Symbol}] \text{ :> With}\{\text{v} = \text{FunctionOfExponential}[\text{u}, \text{x}]\}, \text{Simp}[\text{v}/\text{D}[\text{v}, \text{x}] \quad \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[\text{u}, \text{x}]/\text{x}, \text{x}], \text{x}, \text{v}], \text{x}] \text{ ; FunctionOfExponentialQ}[\text{u}, \text{x}] \&\& \text{!MatchQ}[\text{u}, (\text{w}_.) * ((\text{a}_.) * (\text{v}_.)^{\text{(n}_.)})^{\text{(m}_.)}] \text{ ; FreeQ}\{\text{a}, \text{m}, \text{n}\}, \text{x}\} \&\& \text{IntegerQ}[\text{m} * \text{n}] \&\& \text{!MatchQ}[\text{u}, \text{E}^{\text{(c}_.) * ((\text{a}_.) + (\text{b}_.) * \text{x}))} * (\text{F}_.)[\text{v}_.] \text{ ; FreeQ}\{\text{a}, \text{b}, \text{c}\}, \text{x}\} \&\& \text{InverseFunctionQ}[\text{F}[\text{x}]]]$
- rule 3011 $\text{Int}[\text{Log}[1 + (\text{e}_.) * ((\text{F}_.)^{\text{(c}_.) * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{\text{(n}_.)}))})^{\text{(n}_.)}] * ((\text{f}_.) + (\text{g}_.) * (\text{x}_.)^{\text{(m}_.)}), \text{x_Symbol}] \text{ :> Simp}[(\text{-(f} + \text{g} * \text{x})^{\text{m}} * (\text{PolyLog}[2, (\text{-e}) * (\text{F}^{\text{(c}} * (\text{a} + \text{b} * \text{x}))^{\text{n}}) / (\text{b} * \text{c} * \text{n} * \text{Log}[\text{F}]))], \text{x}) + \text{Simp}[\text{g} * (\text{m} / (\text{b} * \text{c} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{f} + \text{g} * \text{x})^{\text{(m} - 1)} * \text{PolyLog}[2, (\text{-e}) * (\text{F}^{\text{(c}} * (\text{a} + \text{b} * \text{x}))^{\text{n}})], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{GtQ}[\text{m}, 0]$
- rule 5718 $\text{Int}[\text{ArcTan}[(\text{c}_.) + (\text{d}_.) * \text{Tanh}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)]] * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{(m}_.)}), \text{x_Symbol}] \text{ :> Simp}[(\text{e} + \text{f} * \text{x})^{\text{(m} + 1)} * (\text{ArcTan}[\text{c} + \text{d} * \text{Tanh}[\text{a} + \text{b} * \text{x}]] / (\text{f} * (\text{m} + 1))), \text{x}] - \text{Simp}[\text{b} / (\text{f} * (\text{m} + 1)) \quad \text{Int}[(\text{e} + \text{f} * \text{x})^{\text{(m} + 1)} / (\text{c} - \text{d} + \text{c} * \text{E}^{\text{(2} * \text{a} + 2 * \text{b} * \text{x}))}], \text{x}], \text{x}] \text{ ; FreeQ}\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{EqQ}[(\text{c} - \text{d})^{\text{2}}, -1]$

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.54 (sec) , antiderivative size = 1370, normalized size of antiderivative = 12.12

method	result	size
risch	Expression too large to display	1370

input

```
int(x*arctan(c+(I+c)*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*I/b^2*ln(1+I*exp(b*x+a)*(I*c)^(1/2))*a^2+1/8*Pi*(csgn(I*(2*exp(2*b*x+
2*a)*c-2*I))*csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))^2-csgn(I*
(2*exp(2*b*x+2*a)*c-2*I))*csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+
1))*csgn(I/(exp(2*b*x+2*a)+1))-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)
*c))*csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^2+
csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(2*I*exp(2*b*x+2*a)
+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))*csgn(I/(exp(2*b*x+2*a)+1))-csgn(I
*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))^3+csgn(I*(2*exp(2*b*x+2*a)*c
-2*I)/(exp(2*b*x+2*a)+1))^2*csgn(I/(exp(2*b*x+2*a)+1))+csgn(I*(2*exp(2*b*x
+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))*csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+
2*a)+1))^2-csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))*csgn((2*exp
(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*
b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^3-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2
*a)*c)/(exp(2*b*x+2*a)+1))^2*csgn(I/(exp(2*b*x+2*a)+1))-csgn(I*(2*I*exp(2*
b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))*csgn((2*I*exp(2*b*x+2*a)+
2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^2+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp
(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))*csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*
a)*c)/(exp(2*b*x+2*a)+1))+csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(ex
p(2*b*x+2*a)+1))^3-csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x
+2*a)+1))^2+csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)+1))^3-csgn((2...
```


Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(83) = 166$.

Time = 0.13 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.19

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= \frac{-2i b^3 x^3 + 3i b^2 x^2 \log\left(-\frac{(c+i)e^{(2bx+2a)}}{ce^{(2bx+2a)}-i}\right) - 2i a^3 + 6i bx \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4i} ce^{(bx+a)}\right) + 6i bx \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{-4i} ce^{(bx+a)}\right)}{b^2}$$

input `integrate(x*arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="fricas")`

output `1/12*(-2*I*b^3*x^3 + 3*I*b^2*x^2*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) - I)) - 2*I*a^3 + 6*I*b*x*dilog(1/2*sqrt(-4*I*c)*e^(b*x + a)) + 6*I*b*x*dilog(-1/2*sqrt(-4*I*c)*e^(b*x + a)) + 3*I*a^2*log(1/2*(2*c*e^(b*x + a) + I*sqrt(-4*I*c))/c) + 3*I*a^2*log(1/2*(2*c*e^(b*x + a) - I*sqrt(-4*I*c))/c) - 3*(-I*b^2*x^2 + I*a^2)*log(1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) - 3*(-I*b^2*x^2 + I*a^2)*log(-1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) - 6*I*polylog(3, 1/2*sqrt(-4*I*c)*e^(b*x + a)) - 6*I*polylog(3, -1/2*sqrt(-4*I*c)*e^(b*x + a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x*atan(c+(I+c)*tanh(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 + exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.46 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.94

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= \left(\frac{2x^3}{3ic - 3} - \frac{2b^2x^2 \log(ice^{2bx+2a}) + 1 + 2bx \operatorname{Li}_2(-ice^{2bx+2a}) - \operatorname{Li}_3(-ice^{2bx+2a})}{-2b^3(-ic + 1)} \right) b(c + i)$$

$$+ \frac{1}{2} x^2 \arctan((c + i) \tanh(bx + a) + c)$$

input `integrate(x*arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="maxima")`output `(2*x^3/(3*I*c - 3) - (2*b^2*x^2*log(I*c*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog(-I*c*e^(2*b*x + 2*a)) - polylog(3, -I*c*e^(2*b*x + 2*a)))/(b^3*(2*I*c - 2)))*b*(c + I) + 1/2*x^2*arctan((c + I)*tanh(b*x + a) + c)`**Giac [F]**

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx = \int x \arctan((c + i) \tanh(bx + a) + c) dx$$

input `integrate(x*arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="giac")`output `integrate(x*arctan((c + I)*tanh(b*x + a) + c), x)`**Mupad [F(-1)]**

Timed out.

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx = \int x \operatorname{atan}(c + \tanh(a + bx) (c + 1i)) dx$$

input `int(x*atan(c + tanh(a + b*x)*(c + 1i)),x)`

output `int(x*atan(c + tanh(a + b*x)*(c + 1i)), x)`

Reduce [F]

$$\int x \arctan(c + (i + c) \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) c + \tanh(bx + a) i + c) x dx$$

input `int(x*atan(c+(I+c)*tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)*c + tanh(a + b*x)*i + c)*x,x)`

3.87 $\int \arctan(c + (i + c) \tanh(a + bx)) dx$

Optimal result	683
Mathematica [A] (verified)	683
Rubi [A] (verified)	684
Maple [B] (verified)	686
Fricas [B] (verification not implemented)	687
Sympy [F(-2)]	687
Maxima [A] (verification not implemented)	688
Giac [F]	688
Mupad [F(-1)]	688
Reduce [F]	689

Optimal result

Integrand size = 15, antiderivative size = 79

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx = -\frac{1}{2}ibx^2 + x \arctan(c + (i + c) \tanh(a + bx)) + \frac{1}{2}ix \log(1 + ice^{2a+2bx}) + \frac{i \operatorname{PolyLog}(2, -ice^{2a+2bx})}{4b}$$

output

```
-1/2*I*b*x^2+x*arctan(c+(I+c)*tanh(b*x+a))+1/2*I*x*ln(1+I*c*exp(2*b*x+2*a))
+1/4*I*polylog(2,-I*c*exp(2*b*x+2*a))/b
```

Mathematica [A] (verified)

Time = 1.44 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.90

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx = x \arctan(c + (i + c) \tanh(a + bx)) + \frac{i \left(2bx \log \left(1 - \frac{ie^{-2(a+bx)}}{c} \right) - \operatorname{PolyLog} \left(2, \frac{ie^{-2(a+bx)}}{c} \right) \right)}{4b}$$

input `Integrate[ArcTan[c + (I + c)*Tanh[a + b*x]], x]`

output `x*ArcTan[c + (I + c)*Tanh[a + b*x]] + ((I/4)*(2*b*x*Log[1 - I/(c*E^(2*(a + b*x)))] - PolyLog[2, I/(c*E^(2*(a + b*x))]))/b`

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5710, 25, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(c + (c + i) \tanh(a + bx)) dx \\
 & \quad \downarrow \text{5710} \\
 & x \arctan(c + (c + i) \tanh(a + bx)) - b \int -\frac{x}{i - ce^{2a+2bx}} dx \\
 & \quad \downarrow \text{25} \\
 & b \int \frac{x}{i - ce^{2a+2bx}} dx + x \arctan(c + (c + i) \tanh(a + bx)) \\
 & \quad \downarrow \text{2615} \\
 & b \left(-ic \int \frac{e^{2a+2bx} x}{i - ce^{2a+2bx}} dx - \frac{ix^2}{2} \right) + x \arctan(c + (c + i) \tanh(a + bx)) \\
 & \quad \downarrow \text{2620} \\
 & b \left(-ic \left(\frac{\int \log(i e^{2a+2bx} c + 1) dx}{2bc} - \frac{x \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right) + x \arctan(c + (c + i) \tanh(a + bx)) \\
 & \quad \downarrow \text{2715} \\
 & b \left(-ic \left(\frac{\int e^{-2a-2bx} \log(i e^{2a+2bx} c + 1) de^{2a+2bx}}{4b^2 c} - \frac{x \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right) + x \arctan(c + (c + i) \tanh(a + bx))
 \end{aligned}$$

$$\begin{array}{c} \downarrow 2838 \\ b \left(-ic \left(-\frac{x \arctan(c + (c + i) \tanh(a + bx)) + \text{PolyLog}(2, -ice^{2a+2bx})}{4b^2c} - \frac{x \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right) \end{array}$$

input `Int[ArcTan[c + (I + c)*Tanh[a + b*x]],x]`

output `x*ArcTan[c + (I + c)*Tanh[a + b*x]] + b*((-1/2*I)*x^2 - I*c*(-1/2*(x*Log[1 + I*c*E^(2*a + 2*b*x)])/(b*c) - PolyLog[2, (-I)*c*E^(2*a + 2*b*x)]/(4*b^2*c)))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2615 `Int[((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5710

```
Int[ArcTan[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]], x_Symbol] := Simp[x*ArcTan[c + d*Tanh[a + b*x]], x] - Simp[b Int[x/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 544 vs. 2(65) = 130.

Time = 0.54 (sec) , antiderivative size = 545, normalized size of antiderivative = 6.90

method	result
derivativedivides	$\frac{\arctan(c+(i+c)\tanh(bx+a))\ln(c-(i+c)\tanh(bx+a)+i)}{2i+2c} - \frac{2i\arctan(c+(i+c)\tanh(bx+a))\ln(c-(i+c)\tanh(bx+a)+i)c}{2i+2c} - \arctan(c+(i+c)\tanh(bx+a))\ln(c-(i+c)\tanh(bx+a)+i)$
default	$\frac{\arctan(c+(i+c)\tanh(bx+a))\ln(c-(i+c)\tanh(bx+a)+i)}{2i+2c} - \frac{2i\arctan(c+(i+c)\tanh(bx+a))\ln(c-(i+c)\tanh(bx+a)+i)c}{2i+2c} - \arctan(c+(i+c)\tanh(bx+a))\ln(c-(i+c)\tanh(bx+a)+i)$
risch	Expression too large to display

input

```
int(arctan(c+(I+c)*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/b/(I+c)*(arctan(c+(I+c)*tanh(b*x+a))/(2*I+2*c)*ln(c-(I+c)*tanh(b*x+a)+I)
-2*I*arctan(c+(I+c)*tanh(b*x+a))/(2*I+2*c)*ln(c-(I+c)*tanh(b*x+a)+I)*c-arc
tan(c+(I+c)*tanh(b*x+a))/(2*I+2*c)*ln(c-(I+c)*tanh(b*x+a)+I)*c^2-arctan(c+
(I+c)*tanh(b*x+a))/(2*I+2*c)*ln(I+c+(I+c)*tanh(b*x+a))+2*I*arctan(c+(I+c)*
tanh(b*x+a))/(2*I+2*c)*ln(I+c+(I+c)*tanh(b*x+a))*c+arctan(c+(I+c)*tanh(b*x
+a))/(2*I+2*c)*ln(I+c+(I+c)*tanh(b*x+a))*c^2-(I+c)^2*(1/2/(I+c)*(-1/2*I*((
ln(I+c+(I+c)*tanh(b*x+a))-ln(-1/2*I*(I+c+(I+c)*tanh(b*x+a))))*ln(-1/2*I*(I
-c-(I+c)*tanh(b*x+a)))-dilog(-1/2*I*(I+c+(I+c)*tanh(b*x+a))))+1/4*I*ln(I+c
+(I+c)*tanh(b*x+a))^2-1/2/(I+c)*(1/2*I*(dilog((-I-c-(I+c)*tanh(b*x+a))/(-
2*I-2*c))+ln(c-(I+c)*tanh(b*x+a)+I)*ln((-I-c-(I+c)*tanh(b*x+a))/(-2*I-2*c)
))-1/2*I*(dilog(-1/2*(I-c-(I+c)*tanh(b*x+a))/c)+ln(c-(I+c)*tanh(b*x+a)+I)*
ln(-1/2*(I-c-(I+c)*tanh(b*x+a))/c))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(58) = 116$.

Time = 0.11 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.37

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= \frac{-i b^2 x^2 + i b x \log\left(-\frac{(c+i)e^{(2bx+2a)}}{ce^{(2bx+2a)}-i}\right) + i a^2 + (i b x + i a) \log\left(\frac{1}{2} \sqrt{-4i} c e^{(bx+a)} + 1\right) + (i b x + i a) \log\left(-\frac{1}{2} \sqrt{-4i} c e^{(bx+a)} - 1\right)}{b}$$

input `integrate(arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="fricas")`

output `1/2*(-I*b^2*x^2 + I*b*x*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) - I)) + I*a^2 + (I*b*x + I*a)*log(1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) + (I*b*x + I*a)*log(-1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) - I*a*log(1/2*(2*c*e^(b*x + a) + I*sqrt(-4*I*c))/c) - I*a*log(1/2*(2*c*e^(b*x + a) - I*sqrt(-4*I*c))/c) + I*dilog(1/2*sqrt(-4*I*c)*e^(b*x + a)) + I*dilog(-1/2*sqrt(-4*I*c)*e^(b*x + a)))/b`

Sympy [F(-2)]

Exception generated.

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(atan(c+(I+c)*tanh(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 + exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[b, _t0, exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.53 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx$$

$$= 2b(c + i) \left(\frac{2x^2}{2ic - 2} - \frac{2bx \log(i ce^{(2bx+2a)} + 1) + \text{Li}_2(-i ce^{(2bx+2a)})}{-2b^2(-ic + 1)} \right)$$

$$+ x \arctan((c + i) \tanh(bx + a) + c)$$

input `integrate(arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="maxima")`output `2*b*(c + I)*(2*x^2/(2*I*c - 2) - (2*b*x*log(I*c*e^(2*b*x + 2*a) + 1) + dilog(-I*c*e^(2*b*x + 2*a)))/(b^2*(2*I*c - 2))) + x*arctan((c + I)*tanh(b*x + a) + c)`**Giac [F]**

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx = \int \arctan((c + i) \tanh(bx + a) + c) dx$$

input `integrate(arctan(c+(I+c)*tanh(b*x+a)),x, algorithm="giac")`output `integrate(arctan((c + I)*tanh(b*x + a) + c), x)`**Mupad [F(-1)]**

Timed out.

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx = \int \text{atan}(c + \tanh(a + bx) (c + 1i)) dx$$

input `int(atan(c + tanh(a + b*x)*(c + 1i)),x)`

output `int(atan(c + tanh(a + b*x)*(c + 1i)), x)`

Reduce [F]

$$\int \arctan(c + (i + c) \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) c + \tanh(bx + a) i + c) dx$$

input `int(atan(c+(I+c)*tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)*c + tanh(a + b*x)*i + c),x)`

3.88 $\int \frac{\arctan(c+(i+c) \tanh(a+bx))}{x} dx$

Optimal result	690
Mathematica [N/A]	690
Rubi [N/A]	691
Maple [N/A]	691
Fricas [N/A]	692
Sympy [F(-1)]	692
Maxima [N/A]	692
Giac [N/A]	693
Mupad [N/A]	693
Reduce [N/A]	694

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + (i + c) \tanh(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+(I+c)*tanh(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.93 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + (I + c)*Tanh[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + (I + c)*Tanh[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c + (c + i) \tanh(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c + (c + i) \tanh(a + bx))}{x} dx$$

input `Int[ArcTan[c + (I + c)*Tanh[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\arctan(c + (i + c) \tanh(bx + a))}{x} dx$$

input `int(arctan(c+(I+c)*tanh(b*x+a))/x,x)`

output `int(arctan(c+(I+c)*tanh(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.95

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \int \frac{\arctan((c + i) \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+(I+c)*tanh(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) - I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+(I+c)*tanh(b*x+a))/x,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.58 (sec) , antiderivative size = 72, normalized size of antiderivative = 3.79

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \int \frac{\arctan((c + i) \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+(I+c)*tanh(b*x+a))/x,x, algorithm="maxima")`

output

```
I*b*x - 1/4*(4*pi - 4*I*a - 2*arctan(c) - I*log(c^2 + 1))*log(x) + 1/2*integrate(arctan(c*e^(2*b*x + 2*a))/x, x) - 1/4*I*integrate(log(c^2*e^(4*b*x + 4*a) + 1)/x, x)
```

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \int \frac{\arctan((c + i) \tanh(bx + a) + c)}{x} dx$$

input

```
integrate(arctan(c+(I+c)*tanh(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctan((c + I)*tanh(b*x + a) + c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + \tanh(a + bx) (c + 1i))}{x} dx$$

input

```
int(atan(c + tanh(a + b*x)*(c + 1i))/x,x)
```

output

```
int(atan(c + tanh(a + b*x)*(c + 1i))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.32

$$\int \frac{\arctan(c + (i + c) \tanh(a + bx))}{x} dx$$
$$= \int \frac{\operatorname{atan}(\tanh(bx + a) c + \tanh(bx + a) i + c)}{x} dx$$

input `int(atan(c+(I+c)*tanh(b*x+a))/x,x)`output `int(atan(tanh(a + b*x)*c + tanh(a + b*x)*i + c)/x,x)`

3.89 $\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx$

Optimal result	695
Mathematica [A] (verified)	695
Rubi [A] (verified)	696
Maple [C] (warning: unable to verify)	699
Fricas [B] (verification not implemented)	700
Sympy [F(-2)]	700
Maxima [A] (verification not implemented)	701
Giac [F]	701
Mupad [F(-1)]	701
Reduce [F]	702

Optimal result

Integrand size = 22, antiderivative size = 145

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= \frac{1}{12} ibx^4 + \frac{1}{3} x^3 \arctan(c - (i - c) \tanh(a + bx)) - \frac{1}{6} ix^3 \log(1 - ice^{2a+2bx})$$

$$- \frac{ix^2 \operatorname{PolyLog}(2, ice^{2a+2bx})}{4b} + \frac{ix \operatorname{PolyLog}(3, ice^{2a+2bx})}{4b^2} - \frac{i \operatorname{PolyLog}(4, ice^{2a+2bx})}{8b^3}$$

output

```
1/12*I*b*x^4+1/3*x^3*arctan(c-(I-c)*tanh(b*x+a))-1/6*I*x^3*ln(1-I*c*exp(2*b*x+2*a))-1/4*I*x^2*polylog(2,I*c*exp(2*b*x+2*a))/b+1/4*I*x*polylog(3,I*c*exp(2*b*x+2*a))/b^2-1/8*I*polylog(4,I*c*exp(2*b*x+2*a))/b^3
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.92

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= \frac{8b^3 x^3 \arctan(c + (-i + c) \tanh(a + bx)) - 4ib^3 x^3 \log\left(1 + \frac{ie^{-2(a+bx)}}{c}\right) + 6ib^2 x^2 \operatorname{PolyLog}\left(2, -\frac{ie^{-2(a+bx)}}{c}\right)}{24b^3}$$

input `Integrate[x^2*ArcTan[c - (I - c)*Tanh[a + b*x]],x]`

output $(8*b^3*x^3*ArcTan[c + (-I + c)*Tanh[a + b*x]] - (4*I)*b^3*x^3*Log[1 + I/(c * E^{2*(a + b*x)})] + (6*I)*b^2*x^2*PolyLog[2, (-I)/(c * E^{2*(a + b*x)})] + (6*I)*b*x*PolyLog[3, (-I)/(c * E^{2*(a + b*x)})] + (3*I)*PolyLog[4, (-I)/(c * E^{2*(a + b*x)})])/(24*b^3)$

Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.17, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {5718, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(c - (-c + i) \tanh(a + bx)) dx$$

$$\downarrow 5718$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{3}b \int \frac{x^3}{e^{2a+2bx}c + i} dx$$

$$\downarrow 2615$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{3}b \left(ic \int \frac{e^{2a+2bx}x^3}{e^{2a+2bx}c + i} dx - \frac{ix^4}{4} \right)$$

$$\downarrow 2620$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \int x^2 \log(1 - ice^{2a+2bx}) dx}{2bc} \right) - \frac{ix^4}{4} \right)$$

$$\downarrow 3011$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{\int x \text{PolyLog}(2, ice^{2a+2bx}) dx}{b} - \frac{x^2 \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right)$$

$$\begin{aligned} & \downarrow 7163 \\ & \frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \\ & \frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{x \operatorname{PolyLog}(3, ice^{2a+2bx})}{2b} - \frac{\int \operatorname{PolyLog}(3, ice^{2a+2bx}) dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2720 \\ & \frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \\ & \frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{x \operatorname{PolyLog}(3, ice^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(3, ice^{2a+2bx}) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 7143 \\ & \frac{1}{3}x^3 \arctan(c - (-c + i) \tanh(a + bx)) - \\ & \frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{x \operatorname{PolyLog}(3, ice^{2a+2bx})}{2b} - \frac{\operatorname{PolyLog}(4, ice^{2a+2bx})}{4b^2} - \frac{x^2 \operatorname{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) \end{aligned}$$

input `Int[x^2*ArcTan[c - (I - c)*Tanh[a + b*x]],x]`

output `(x^3*ArcTan[c - (I - c)*Tanh[a + b*x]])/3 - (b*((-1/4*I)*x^4 + I*c*((x^3*Log[1 - I*c*E^(2*a + 2*b*x)])/(2*b*c) - (3*(-1/2*(x^2*PolyLog[2, I*c*E^(2*a + 2*b*x)]))/b + ((x*PolyLog[3, I*c*E^(2*a + 2*b*x)])/(2*b) - PolyLog[4, I*c*E^(2*a + 2*b*x)]/(4*b^2))/b))/(2*b*c)))/3`

Definitions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5718 `Int[ArcTan[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tanh[a + b*x]]/(f*(m + 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.29 (sec) , antiderivative size = 1417, normalized size of antiderivative = 9.77

method	result	size
risch	Expression too large to display	1417

input

```
int(x^2*arctan(c-(1-c)*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/6*I*x^3*ln(-2*exp(2*b*x+2*a)*c-2*I)-1/12*Pi*(csgn(I*(-2*I*exp(2*b*x+2*a)
+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))*csgn((-2*I*exp(2*b*x+2*a)+2*exp(2
*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))-csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a
)*c)/(exp(2*b*x+2*a)+1))^2-csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*(2*exp(2*b*x+
2*a)*c+2*I))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))+csgn(I/(e
xp(2*b*x+2*a)+1))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*
(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))+csgn(I/(exp(2
*b*x+2*a)+1))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))^2-csgn(I
/(exp(2*b*x+2*a)+1))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(
2*b*x+2*a)+1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))*csgn(
(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))-csgn((2*exp(2*b*x+2*a)*c+2*I)
/(exp(2*b*x+2*a)+1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I))*csgn(I*(2*exp(2*b*
x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))^2+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*
x+2*a)*c))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)
+1))^2+csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))^3-csgn(I*(2*exp
(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))*csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(
2*b*x+2*a)+1))^2-csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*
x+2*a)+1))^3+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*
a)+1))*csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^2
-csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^3-cs...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs. $2(105) = 210$.

Time = 0.11 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.02

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= \frac{i b^4 x^4 + 2i b^3 x^3 \log\left(-\frac{ce^{(2bx+2a)+i}e^{(-2bx-2a)}}{c-i}\right) - 6i b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{4i} ce^{(bx+a)}\right) - 6i b^2 x^2 \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{4i} ce^{(bx+a)}\right)}{b^3}$$

input `integrate(x^2*arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="fricas")`

output `1/12*(I*b^4*x^4 + 2*I*b^3*x^3*log(-(c*e^(2*b*x + 2*a) + I)*e^(-2*b*x - 2*a))/(c - I)) - 6*I*b^2*x^2*dilog(1/2*sqrt(4*I*c)*e^(b*x + a)) - 6*I*b^2*x^2*dilog(-1/2*sqrt(4*I*c)*e^(b*x + a)) - I*a^4 + 2*I*a^3*log(1/2*(2*c*e^(b*x + a) + I*sqrt(4*I*c))/c) + 2*I*a^3*log(1/2*(2*c*e^(b*x + a) - I*sqrt(4*I*c))/c) + 12*I*b*x*polylog(3, 1/2*sqrt(4*I*c)*e^(b*x + a)) + 12*I*b*x*polylog(3, -1/2*sqrt(4*I*c)*e^(b*x + a)) - 2*(I*b^3*x^3 + I*a^3)*log(1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - 2*(I*b^3*x^3 + I*a^3)*log(-1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - 12*I*polylog(4, 1/2*sqrt(4*I*c)*e^(b*x + a)) - 12*I*polylog(4, -1/2*sqrt(4*I*c)*e^(b*x + a)))/b^3`

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x**2*atan(c-(I-c)*tanh(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 + exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.42 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.89

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx = \frac{1}{3} x^3 \arctan((c - i) \tanh(bx + a) + c) - \frac{4}{9} \left(\frac{3x^4}{4ic + 4} - \frac{4b^3 x^3 \log(-ice^{(2bx+2a)} + 1) + 6b^2 x^2 \text{Li}_2(ice^{(2bx+2a)}) - 6bx \text{Li}_3(ice^{(2bx+2a)}) + 3 \text{Li}_4(ice^{(2bx+2a)})}{-2b^4(-ic - 1)} \right)$$

input `integrate(x^2*arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="maxima")`

output

```
1/3*x^3*arctan((c - I)*tanh(b*x + a) + c) - 4/9*(3*x^4/(4*I*c + 4) - (4*b^3*x^3*log(-I*c*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog(I*c*e^(2*b*x + 2*a)) - 6*b*x*polylog(3, I*c*e^(2*b*x + 2*a)) + 3*polylog(4, I*c*e^(2*b*x + 2*a)))/(b^4*(2*I*c + 2)))*b*(c - I)
```

Giac [F]

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx = \int x^2 \arctan((c - i) \tanh(bx + a) + c) dx$$

input `integrate(x^2*arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="giac")`

output

```
integrate(x^2*arctan((c - I)*tanh(b*x + a) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx = \int x^2 \text{atan}(c + \tanh(a + bx) (c - i)) dx$$

input `int(x^2*atan(c + tanh(a + b*x)*(c - 1i)),x)`

output `int(x^2*atan(c + tanh(a + b*x)*(c - 1i)), x)`

Reduce [F]

$$\int x^2 \arctan(c - (i - c) \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) c - \tanh(bx + a) i + c) x^2 dx$$

input `int(x^2*atan(c-(I-c)*tanh(b*x+a)), x)`

output `int(atan(tanh(a + b*x)*c - tanh(a + b*x)*i + c)*x**2,x)`

3.90 $\int x \arctan(c - (i - c) \tanh(a + bx)) dx$

Optimal result	703
Mathematica [A] (verified)	704
Rubi [A] (verified)	704
Maple [C] (warning: unable to verify)	707
Fricas [B] (verification not implemented)	708
Sympy [F(-2)]	708
Maxima [A] (verification not implemented)	709
Giac [F]	709
Mupad [F(-1)]	709
Reduce [F]	710

Optimal result

Integrand size = 20, antiderivative size = 116

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx = \frac{1}{6}ibx^3 + \frac{1}{2}x^2 \arctan(c - (i - c) \tanh(a + bx)) - \frac{1}{4}ix^2 \log(1 - ice^{2a+2bx}) - \frac{ix \operatorname{PolyLog}(2, ice^{2a+2bx})}{4b} + \frac{i \operatorname{PolyLog}(3, ice^{2a+2bx})}{8b^2}$$

output

```
1/6*I*b*x^3+1/2*x^2*arctan(c-(I-c)*tanh(b*x+a))-1/4*I*x^2*ln(1-I*c*exp(2*b*x+2*a))-1/4*I*x*polylog(2,I*c*exp(2*b*x+2*a))/b+1/8*I*polylog(3,I*c*exp(2*b*x+2*a))/b^2
```


Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.89

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= \frac{2b^2 x^2 \left(2 \arctan(c + (-i + c) \tanh(a + bx)) - i \log \left(1 + \frac{ie^{-2(a+bx)}}{c} \right) \right) + 2ibx \operatorname{PolyLog} \left(2, -\frac{ie^{-2(a+bx)}}{c} \right) + i}{8b^2}$$

input

```
Integrate[x*ArcTan[c - (I - c)*Tanh[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTan[c + (-I + c)*Tanh[a + b*x]] - I*Log[1 + I/(c*E^(2*(a + b*x))])) + (2*I)*b*x*PolyLog[2, (-I)/(c*E^(2*(a + b*x)))] + I*PolyLog[3, (-I)/(c*E^(2*(a + b*x)))]/(8*b^2)
```

Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.17, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5718, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(c - (-c + i) \tanh(a + bx)) dx$$

$$\downarrow \text{5718}$$

$$\frac{1}{2} x^2 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{2} b \int \frac{x^2}{e^{2a+2bx} c + i} dx$$

$$\downarrow \text{2615}$$

$$\frac{1}{2} x^2 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{2} b \left(ic \int \frac{e^{2a+2bx} x^2}{e^{2a+2bx} c + i} dx - \frac{ix^3}{3} \right)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int x \log(1 - ice^{2a+2bx}) dx}{bc} \right) - \frac{ix^3}{3} \right)$$

↓ 3011

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int \text{PolyLog}(2, ice^{2a+2bx}) dx}{2b} - \frac{x \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right) - \frac{ix^3}{3} \right)$$

↓ 2720

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int e^{-2a-2bx} \text{PolyLog}(2, ice^{2a+2bx}) de^{2a+2bx}}{4b^2} - \frac{x \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right) - \frac{ix^3}{3} \right)$$

↓ 7143

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \tanh(a + bx)) - \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\text{PolyLog}(3, ice^{2a+2bx})}{4b^2} - \frac{x \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right) - \frac{ix^3}{3} \right)$$

input `Int[x*ArcTan[c - (I - c)*Tanh[a + b*x]],x]`

output `(x^2*ArcTan[c - (I - c)*Tanh[a + b*x]])/2 - (b*((-1/3*I)*x^3 + I*c*((x^2*Log[1 - I*c*E^(2*a + 2*b*x)])/(2*b*c) - (-1/2*(x*PolyLog[2, I*c*E^(2*a + 2*b*x)])/b + PolyLog[3, I*c*E^(2*a + 2*b*x)]/(4*b^2))/(b*c)))/2`

Defintions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5718

```
Int[ArcTan[(c_) + (d_)*Tanh[(a_) + (b_)*(x_)]]*((e_) + (f_)*(x_))^(m_
.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Tanh[a + b*x]]/(f*(m
+ 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d + c*E^(2*a +
2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c -
d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.68 (sec) , antiderivative size = 1381, normalized size of antiderivative = 11.91

method	result	size
risch	Expression too large to display	1381

input `int(x*arctan(c-(I-c)*tanh(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-1/4*I*x^2*ln(1-I*c*exp(2*b*x+2*a))-1/8*Pi*(csgn(I*(-2*I*exp(2*b*x+2*a)+2*
exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))*csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*
x+2*a)*c)/(exp(2*b*x+2*a)+1))-csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c
)/(exp(2*b*x+2*a)+1))^2-csgn(I/(exp(2*b*x+2*a)+1))*csgn(I*(2*exp(2*b*x+2*a
)*c+2*I))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))+csgn(I/(exp(
2*b*x+2*a)+1))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(-2
*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))+csgn(I/(exp(2*b*
x+2*a)+1))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))^2-csgn(I/(e
xp(2*b*x+2*a)+1))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b
*x+2*a)+1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))*csgn((2*
exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))-csgn((2*exp(2*b*x+2*a)*c+2*I)/(e
xp(2*b*x+2*a)+1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I))*csgn(I*(2*exp(2*b*x+2
*a)*c+2*I)/(exp(2*b*x+2*a)+1))^2+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2
*a)*c))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1
))^2+csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))^3-csgn(I*(2*exp(2*
b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)+1))*csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b
*x+2*a)+1))^2-csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2
*a)+1))^3+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+
1))*csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^2-cs
gn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)+1))^3-csgn(...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(83) = 166$.

Time = 0.11 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.13

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= \frac{2i b^3 x^3 + 3i b^2 x^2 \log\left(-\frac{ce^{(2bx+2a)+i}e^{(-2bx-2a)}}{c-i}\right) + 2i a^3 - 6i bx \operatorname{Li}_2\left(\frac{1}{2} \sqrt{4i ce^{(bx+a)}}\right) - 6i bx \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{4i ce^{(bx+a)}}\right)}{b^2}$$

input `integrate(x*arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*I*b^3*x^3 + 3*I*b^2*x^2*log(-(c*e^(2*b*x + 2*a) + I)*e^(-2*b*x - 2*a)/(c - I)) + 2*I*a^3 - 6*I*b*x*dilog(1/2*sqrt(4*I*c)*e^(b*x + a)) - 6*I*b*x*dilog(-1/2*sqrt(4*I*c)*e^(b*x + a)) - 3*I*a^2*log(1/2*(2*c*e^(b*x + a) + I*sqrt(4*I*c))/c) - 3*I*a^2*log(1/2*(2*c*e^(b*x + a) - I*sqrt(4*I*c))/c) - 3*(I*b^2*x^2 - I*a^2)*log(1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - 3*(I*b^2*x^2 - I*a^2)*log(-1/2*sqrt(4*I*c)*e^(b*x + a) + 1) + 6*I*polylog(3, 1/2*sqrt(4*I*c)*e^(b*x + a)) + 6*I*polylog(3, -1/2*sqrt(4*I*c)*e^(b*x + a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x*atan(c-(I-c)*tanh(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 + exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.92

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx =$$

$$-\left(\frac{2x^3}{3ic + 3} - \frac{2b^2x^2 \log(-ice^{(2bx+2a)} + 1) + 2bx \operatorname{Li}_2(ice^{(2bx+2a)}) - \operatorname{Li}_3(ice^{(2bx+2a)})}{-2b^3(-ic - 1)} \right) b(c - i)$$

$$+ \frac{1}{2} x^2 \arctan((c - i) \tanh(bx + a) + c)$$

input `integrate(x*arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="maxima")`

output `-(2*x^3/(3*I*c + 3) - (2*b^2*x^2*log(-I*c*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog(I*c*e^(2*b*x + 2*a)) - polylog(3, I*c*e^(2*b*x + 2*a)))/(b^3*(2*I*c + 2)))*b*(c - I) + 1/2*x^2*arctan((c - I)*tanh(b*x + a) + c)`

Giac [F]

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx = \int x \arctan((c - i) \tanh(bx + a) + c) dx$$

input `integrate(x*arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan((c - I)*tanh(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx = \int x \operatorname{atan}(c + \tanh(a + bx) (c - i)) dx$$

input `int(x*atan(c + tanh(a + b*x)*(c - 1i)),x)`

output `int(x*atan(c + tanh(a + b*x)*(c - 1i)), x)`

Reduce [F]

$$\int x \arctan(c - (i - c) \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx+a) c - \tanh(bx+a) i + c) x dx$$

input `int(x*atan(c-(I-c)*tanh(b*x+a)), x)`

output `int(atan(tanh(a + b*x)*c - tanh(a + b*x)*i + c)*x, x)`

3.91 $\int \arctan(c - (i - c) \tanh(a + bx)) dx$

Optimal result	711
Mathematica [A] (verified)	711
Rubi [A] (verified)	712
Maple [B] (verified)	714
Fricas [B] (verification not implemented)	715
Sympy [F(-2)]	715
Maxima [A] (verification not implemented)	716
Giac [F]	716
Mupad [F(-1)]	716
Reduce [F]	717

Optimal result

Integrand size = 18, antiderivative size = 82

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx = \frac{1}{2}ibx^2 + x \arctan(c - (i - c) \tanh(a + bx)) - \frac{1}{2}ix \log(1 - ice^{2a+2bx}) - \frac{i \operatorname{PolyLog}(2, ice^{2a+2bx})}{4b}$$

output

```
1/2*I*b*x^2+x*arctan(c-(I-c)*tanh(b*x+a))-1/2*I*x*ln(1-I*c*exp(2*b*x+2*a))
-1/4*I*polylog(2,I*c*exp(2*b*x+2*a))/b
```

Mathematica [A] (verified)

Time = 1.47 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.87

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx = x \arctan(c + (-i + c) \tanh(a + bx)) - \frac{i \left(2bx \log \left(1 + \frac{ie^{-2(a+bx)}}{c} \right) - \operatorname{PolyLog} \left(2, -\frac{ie^{-2(a+bx)}}{c} \right) \right)}{4b}$$

input `Integrate[ArcTan[c - (I - c)*Tanh[a + b*x]], x]`

output `x*ArcTan[c + (-I + c)*Tanh[a + b*x]] - ((I/4)*(2*b*x*Log[1 + I/(c*E^(2*(a + b*x))]) - PolyLog[2, (-I)/(c*E^(2*(a + b*x))]))/b`

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.17, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {5710, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(c - (-c + i) \tanh(a + bx)) dx$$

$$\downarrow \text{5710}$$

$$x \arctan(c - (-c + i) \tanh(a + bx)) - b \int \frac{x}{e^{2a+2bx} c + i} dx$$

$$\downarrow \text{2615}$$

$$x \arctan(c - (-c + i) \tanh(a + bx)) - b \left(ic \int \frac{e^{2a+2bx} x}{e^{2a+2bx} c + i} dx - \frac{ix^2}{2} \right)$$

$$\downarrow \text{2620}$$

$$b \left(ic \left(\frac{x \arctan(c - (-c + i) \tanh(a + bx)) - \frac{x \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int \log(1 - ice^{2a+2bx}) dx}{2bc}}{2bc} \right) - \frac{ix^2}{2} \right)$$

$$\downarrow \text{2715}$$

$$b \left(ic \left(\frac{x \arctan(c - (-c + i) \tanh(a + bx)) - \frac{x \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int e^{-2a-2bx} \log(1 - ice^{2a+2bx}) de^{2a+2bx}}{4b^2 c}}{2bc} \right) - \frac{ix^2}{2} \right)$$

$$\downarrow \text{2838}$$

$$b \left(ic \left(\frac{x \arctan(c - (-c + i) \tanh(a + bx)) - \text{PolyLog}(2, ice^{2a+2bx})}{4b^2c} + \frac{x \log(1 - ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right)$$

input `Int[ArcTan[c - (I - c)*Tanh[a + b*x]],x]`

output `x*ArcTan[c - (I - c)*Tanh[a + b*x]] - b*((-1/2*I)*x^2 + I*c*((x*Log[1 - I*c*E^(2*a + 2*b*x)])/(2*b*c) + PolyLog[2, I*c*E^(2*a + 2*b*x)]/(4*b^2*c)))`

Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5710 `Int[ArcTan[(c_.) + (d_.)*Tanh[(a_.) + (b_.)*(x_)]], x_Symbol] := Simp[x*ArcTan[c + d*Tanh[a + b*x]], x] - Simp[b Int[x/(c - d + c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, -1]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 515 vs. $2(68) = 136$.

Time = 0.70 (sec) , antiderivative size = 516, normalized size of antiderivative = 6.29

method	result
derivativedivides	$-\frac{\arctan(c+\tanh(bx+a)(-i+c))\ln(\tanh(bx+a)(-i+c)-c+i)}{2i-2c} - \frac{2i\arctan(c+\tanh(bx+a)(-i+c))\ln(\tanh(bx+a)(-i+c)-c+i)c}{2i-2c} + \arctan(c+\tanh(bx+a)(-i+c))\ln(\tanh(bx+a)(-i+c)-c+i)$
default	$-\frac{\arctan(c+\tanh(bx+a)(-i+c))\ln(\tanh(bx+a)(-i+c)-c+i)}{2i-2c} - \frac{2i\arctan(c+\tanh(bx+a)(-i+c))\ln(\tanh(bx+a)(-i+c)-c+i)c}{2i-2c} + \arctan(c+\tanh(bx+a)(-i+c))\ln(\tanh(bx+a)(-i+c)-c+i)$
risch	Expression too large to display

```
input int(arctan(c-(I-c)*tanh(b*x+a)),x,method=_RETURNVERBOSE)
```

```
output 1/b/(-I+c)*(-arctan(c+tanh(b*x+a)*(-I+c))/(2*I-2*c)*ln(tanh(b*x+a)*(-I+c)-c+I)-2*I*arctan(c+tanh(b*x+a)*(-I+c))/(2*I-2*c)*ln(tanh(b*x+a)*(-I+c)-c+I)*c+arctan(c+tanh(b*x+a)*(-I+c))/(2*I-2*c)*ln(tanh(b*x+a)*(-I+c)-c+I)*c^2+arctan(c+tanh(b*x+a)*(-I+c))/(2*I-2*c)*ln(-I+tanh(b*x+a)*(-I+c)+c)+2*I*arctan(c+tanh(b*x+a)*(-I+c))/(2*I-2*c)*ln(-I+tanh(b*x+a)*(-I+c)+c)*c-arctan(c+tanh(b*x+a)*(-I+c))/(2*I-2*c)*ln(-I+tanh(b*x+a)*(-I+c)+c)*c^2+(I-c)^2*(1/2/(I-c)*(1/2*I*(dilog(-1/2*I*(tanh(b*x+a)*(-I+c)+c+I))+ln(-I+tanh(b*x+a)*(-I+c)+c)*ln(-1/2*I*(tanh(b*x+a)*(-I+c)+c+I)))-1/4*I*ln(-I+tanh(b*x+a)*(-I+c)+c)^2)-1/2/(I-c)*(-1/2*I*(dilog((-I+tanh(b*x+a)*(-I+c)+c)/(-2*I+2*c))+ln(tanh(b*x+a)*(-I+c)-c+I)*ln((-I+tanh(b*x+a)*(-I+c)+c)/(-2*I+2*c)))+1/2*I*(dilog(1/2*(tanh(b*x+a)*(-I+c)+c+I)/c)+ln(tanh(b*x+a)*(-I+c)-c+I)*ln(1/2*(tanh(b*x+a)*(-I+c)+c+I)/c))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(58) = 116$.

Time = 0.10 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.28

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= \frac{i b^2 x^2 + i b x \log\left(-\frac{(c e^{2bx+2a} + i) e^{-2bx-2a}}{c-i}\right) - i a^2 + (-i b x - i a) \log\left(\frac{1}{2} \sqrt{4i c e^{bx+a}} + 1\right) + (-i b x - i a)}{b}$$

input `integrate(arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="fricas")`

output `1/2*(I*b^2*x^2 + I*b*x*log(-(c*e^(2*b*x + 2*a) + I)*e^(-2*b*x - 2*a)/(c - I)) - I*a^2 + (-I*b*x - I*a)*log(1/2*sqrt(4*I*c)*e^(b*x + a) + 1) + (-I*b*x - I*a)*log(-1/2*sqrt(4*I*c)*e^(b*x + a) + 1) + I*a*log(1/2*(2*c*e^(b*x + a) + I*sqrt(4*I*c))/c) + I*a*log(1/2*(2*c*e^(b*x + a) - I*sqrt(4*I*c))/c) - I*dilog(1/2*sqrt(4*I*c)*e^(b*x + a)) - I*dilog(-1/2*sqrt(4*I*c)*e^(b*x + a)))/b`

Sympy [F(-2)]

Exception generated.

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(atan(c-(I-c)*tanh(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 + exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[b, _t0, exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.37 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.98

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx$$

$$= -2b(c - i) \left(\frac{2x^2}{2ic + 2} - \frac{2bx \log(-ice^{(2bx+2a)} + 1) + \text{Li}_2(ice^{(2bx+2a)})}{-2b^2(-ic - 1)} \right)$$

$$+ x \arctan((c - i) \tanh(bx + a) + c)$$

input `integrate(arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="maxima")`output `-2*b*(c - I)*(2*x^2/(2*I*c + 2) - (2*b*x*log(-I*c*e^(2*b*x + 2*a) + 1) + dilog(I*c*e^(2*b*x + 2*a)))/(b^2*(2*I*c + 2))) + x*arctan((c - I)*tanh(b*x + a) + c)`**Giac [F]**

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx = \int \arctan((c - i) \tanh(bx + a) + c) dx$$

input `integrate(arctan(c-(I-c)*tanh(b*x+a)),x, algorithm="giac")`output `integrate(arctan((c - I)*tanh(b*x + a) + c), x)`**Mupad [F(-1)]**

Timed out.

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx = \int \text{atan}(c + \tanh(a + bx) (c - i)) dx$$

input `int(atan(c + tanh(a + b*x)*(c - 1i)),x)`

output `int(atan(c + tanh(a + b*x)*(c - 1i)), x)`

Reduce [F]

$$\int \arctan(c - (i - c) \tanh(a + bx)) dx = \int \operatorname{atan}(\tanh(bx + a) c - \tanh(bx + a) i + c) dx$$

input `int(atan(c-(I-c)*tanh(b*x+a)),x)`

output `int(atan(tanh(a + b*x)*c - tanh(a + b*x)*i + c),x)`

3.92 $\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx$

Optimal result	718
Mathematica [N/A]	718
Rubi [N/A]	719
Maple [N/A]	719
Fricas [N/A]	720
Sympy [F(-1)]	720
Maxima [N/A]	720
Giac [N/A]	721
Mupad [N/A]	721
Reduce [N/A]	722

Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c - (i - c) \tanh(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c-(I-c)*tanh(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 2.96 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx$$

input `Integrate[ArcTan[c - (I - c)*Tanh[a + b*x]]/x,x]`

output `Integrate[ArcTan[c - (I - c)*Tanh[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c - (-c + i) \tanh(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c - (-c + i) \tanh(a + bx))}{x} dx$$

input `Int[ArcTan[c - (I - c)*Tanh[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\arctan(c - (i - c) \tanh(bx + a))}{x} dx$$

input `int(arctan(c-(I-c)*tanh(b*x+a))/x,x)`

output `int(arctan(c-(I-c)*tanh(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \int \frac{\arctan((c - i) \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c-(I-c)*tanh(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c*e^(2*b*x + 2*a) + I)*e^(-2*b*x - 2*a)/(c - I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c-(I-c)*tanh(b*x+a))/x,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 77, normalized size of antiderivative = 3.50

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \int \frac{\arctan((c - i) \tanh(bx + a) + c)}{x} dx$$

input `integrate(arctan(c-(I-c)*tanh(b*x+a))/x,x, algorithm="maxima")`

output

```
-I*b*x - 1/2*pi*log(x) - 1/4*(2*pi + 4*I*a - 2*arctan(c) + I*log(c^2 + 1))
*log(x) + 1/2*integrate(arctan(c*e^(2*b*x + 2*a))/x, x) + 1/4*I*integrate(
log(c^2*e^(4*b*x + 4*a) + 1)/x, x)
```

Giac [N/A]

Not integrable

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \int \frac{\arctan((c - i) \tanh(bx + a) + c)}{x} dx$$

input

```
integrate(arctan(c-(I-c)*tanh(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctan((c - I)*tanh(b*x + a) + c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + \tanh(a + bx) (c - i))}{x} dx$$

input

```
int(atan(c + tanh(a + b*x)*(c - 1i))/x,x)
```

output

```
int(atan(c + tanh(a + b*x)*(c - 1i))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{\arctan(c - (i - c) \tanh(a + bx))}{x} dx$$
$$= \int \frac{\operatorname{atan}(\tanh(bx + a) c - \tanh(bx + a) i + c)}{x} dx$$

input `int(atan(c-(I-c)*tanh(b*x+a))/x,x)`output `int(atan(tanh(a + b*x)*c - tanh(a + b*x)*i + c)/x,x)`

3.93 $\int (e + fx)^3 \arctan(\coth(a + bx)) dx$

Optimal result	723
Mathematica [B] (verified)	724
Rubi [A] (verified)	725
Maple [C] (warning: unable to verify)	729
Fricas [B] (verification not implemented)	730
Sympy [F]	731
Maxima [F]	732
Giac [F(-1)]	732
Mupad [F(-1)]	732
Reduce [F]	733

Optimal result

Integrand size = 15, antiderivative size = 299

$$\begin{aligned}
 \int (e + fx)^3 \arctan(\coth(a + bx)) dx = & \frac{(e + fx)^4 \arctan(e^{2a+2bx})}{4f} \\
 & + \frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} \\
 & - \frac{i(e + fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} \\
 & + \frac{i(e + fx)^3 \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b} \\
 & + \frac{3if(e + fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{8b^2} \\
 & - \frac{3if(e + fx)^2 \operatorname{PolyLog}(3, ie^{2a+2bx})}{8b^2} \\
 & - \frac{3if^2(e + fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{8b^3} \\
 & + \frac{3if^2(e + fx) \operatorname{PolyLog}(4, ie^{2a+2bx})}{8b^3} \\
 & + \frac{3if^3 \operatorname{PolyLog}(5, -ie^{2a+2bx})}{16b^4} \\
 & - \frac{3if^3 \operatorname{PolyLog}(5, ie^{2a+2bx})}{16b^4}
 \end{aligned}$$

output

```
1/4*(f*x+e)^4*arctan(exp(2*b*x+2*a))/f+1/4*(f*x+e)^4*arctan(coth(b*x+a))/f
-1/4*I*(f*x+e)^3*polylog(2,-I*exp(2*b*x+2*a))/b+1/4*I*(f*x+e)^3*polylog(2,
I*exp(2*b*x+2*a))/b+3/8*I*f*(f*x+e)^2*polylog(3,-I*exp(2*b*x+2*a))/b^2-3/8
*I*f*(f*x+e)^2*polylog(3,I*exp(2*b*x+2*a))/b^2-3/8*I*f^2*(f*x+e)*polylog(4
,-I*exp(2*b*x+2*a))/b^3+3/8*I*f^2*(f*x+e)*polylog(4,I*exp(2*b*x+2*a))/b^3+
3/16*I*f^3*polylog(5,-I*exp(2*b*x+2*a))/b^4-3/16*I*f^3*polylog(5,I*exp(2*b
*x+2*a))/b^4
```

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 600 vs. $2(299) = 598$.

Time = 0.27 (sec) , antiderivative size = 600, normalized size of antiderivative = 2.01

$$\int (e + fx)^3 \arctan(\coth(a + bx)) dx$$

$$= \frac{1}{4}x(4e^3 + 6e^2fx + 4ef^2x^2 + f^3x^3) \arctan(\coth(a + bx))$$

$$+ \frac{i(8b^4e^3x \log(1 - ie^{2(a+bx)}) + 12b^4e^2fx^2 \log(1 - ie^{2(a+bx)}) + 8b^4ef^2x^3 \log(1 - ie^{2(a+bx)}) + 2b^4f^3x^4 \log(1 - ie^{2(a+bx)}))}{4}$$

input

```
Integrate[(e + f*x)^3*ArcTan[Coth[a + b*x]],x]
```

output

```
(x*(4*e^3 + 6*e^2*f*x + 4*e*f^2*x^2 + f^3*x^3)*ArcTan[Coth[a + b*x]])/4 +
((I/16)*(8*b^4*e^3*x*Log[1 - I*E^(2*(a + b*x))] + 12*b^4*e^2*f*x^2*Log[1 -
I*E^(2*(a + b*x))] + 8*b^4*e*f^2*x^3*Log[1 - I*E^(2*(a + b*x))] + 2*b^4*f
^3*x^4*Log[1 - I*E^(2*(a + b*x))] - 8*b^4*e^3*x*Log[1 + I*E^(2*(a + b*x))]
- 12*b^4*e^2*f*x^2*Log[1 + I*E^(2*(a + b*x))] - 8*b^4*e*f^2*x^3*Log[1 + I
*E^(2*(a + b*x))] - 2*b^4*f^3*x^4*Log[1 + I*E^(2*(a + b*x))] - 4*b^3*(e +
f*x)^3*PolyLog[2, (-I)*E^(2*(a + b*x))] + 4*b^3*(e + f*x)^3*PolyLog[2, I*E
^(2*(a + b*x))] + 6*b^2*e^2*f*PolyLog[3, (-I)*E^(2*(a + b*x))] + 12*b^2*e*
f^2*x*PolyLog[3, (-I)*E^(2*(a + b*x))] + 6*b^2*f^3*x^2*PolyLog[3, (-I)*E^(
2*(a + b*x))] - 6*b^2*e^2*f*PolyLog[3, I*E^(2*(a + b*x))] - 12*b^2*e*f^2*x
*PolyLog[3, I*E^(2*(a + b*x))] - 6*b^2*f^3*x^2*PolyLog[3, I*E^(2*(a + b*x)
)] - 6*b*e*f^2*PolyLog[4, (-I)*E^(2*(a + b*x))] - 6*b*f^3*x*PolyLog[4, (-I
)*E^(2*(a + b*x))] + 6*b*e*f^2*PolyLog[4, I*E^(2*(a + b*x))] + 6*b*f^3*x*P
olyLog[4, I*E^(2*(a + b*x))] + 3*f^3*PolyLog[5, (-I)*E^(2*(a + b*x))] - 3*
f^3*PolyLog[5, I*E^(2*(a + b*x))])/b^4
```

Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {5708, 3042, 4668, 3011, 7163, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (e + fx)^3 \arctan(\coth(a + bx)) dx \\
 & \quad \downarrow \text{5708} \\
 & \frac{b \int (e + fx)^4 \operatorname{sech}(2a + 2bx) dx}{4f} + \frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \frac{b \int (e + fx)^4 \csc\left(2ia + 2ibx + \frac{\pi}{2}\right) dx}{4f} \\
 & \quad \downarrow \text{4668} \\
 & \frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \\
 & \frac{b \left(-\frac{2if \int (e+fx)^3 \log(1-ie^{2a+2bx}) dx}{b} + \frac{2if \int (e+fx)^3 \log(1+ie^{2a+2bx}) dx}{b} + \frac{(e+fx)^4 \arctan(e^{2a+2bx})}{b} \right)}{4f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \\
 & b \left(\frac{2if \left(\frac{3f \int (e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{2if \left(\frac{3f \int (e+fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx}) dx}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{b} \right) \\
 & \quad \downarrow \text{7163}
 \end{aligned}$$

$$\frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \int (e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx}) dx}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(1, -ie^{2a+2bx})}{2b} \right)}{b}$$

4f

↓ 7163

$$\frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(4, -ie^{2a+2bx}) dx}{2b} \right)}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{b} - \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{b} \right)}{2b} - \frac{(e+fx)^3 \operatorname{PolyLog}(1, -ie^{2a+2bx})}{2b} \right)}{b}$$

↓ 2720

$$\frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{2b} - \frac{f \int e^{-2a-2bx} \operatorname{PolyLog}(4, -ie^{2a+2bx}) de^{2a+2bx}}{4b^2} \right)}{b} \right)}{2b} \right)}{b} - \frac{(e+fx)^3 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b}$$

7143

$$\frac{(e + fx)^4 \arctan(\coth(a + bx))}{4f} + \frac{2if \left(\frac{3f \left(\frac{(e+fx)^2 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(4, -ie^{2a+2bx})}{2b} - \frac{f \operatorname{PolyLog}(5, -ie^{2a+2bx})}{4b^2} \right)}{b} \right)}{2b} \right)}{b} - \frac{(e+fx)^4 \arctan(e^{2a+2bx})}{b} + \frac{(e+fx)^3 \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b}$$

input `Int[(e + f*x)^3*ArcTan[Coth[a + b*x]],x]`

output

```
((e + f*x)^4*ArcTan[Coth[a + b*x]])/(4*f) + (b*(((e + f*x)^4*ArcTan[E^(2*a
+ 2*b*x)])/b + ((2*I)*f*(-1/2*((e + f*x)^3*PolyLog[2, (-I)*E^(2*a + 2*b*x
)])/b + (3*f*(((e + f*x)^2*PolyLog[3, (-I)*E^(2*a + 2*b*x)])/(2*b) - (f*((
e + f*x)*PolyLog[4, (-I)*E^(2*a + 2*b*x)])/(2*b) - (f*PolyLog[5, (-I)*E^(
2*a + 2*b*x)])/(4*b^2)))/b))/(2*b))/b - ((2*I)*f*(-1/2*((e + f*x)^3*PolyL
og[2, I*E^(2*a + 2*b*x)])/b + (3*f*(((e + f*x)^2*PolyLog[3, I*E^(2*a + 2*b
*x)])/(2*b) - (f*((e + f*x)*PolyLog[4, I*E^(2*a + 2*b*x)])/(2*b) - (f*Pol
yLog[5, I*E^(2*a + 2*b*x)])/(4*b^2)))/b))/(2*b))/b)/(4*f)
```

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4668

```
Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m, x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5708 `Int[ArcTan[Coth[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol]
:> Simp[(e + f*x)^(m + 1)*(ArcTan[Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/
(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a,
b, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 7.81 (sec) , antiderivative size = 3570, normalized size of antiderivative = 11.94

method	result	size
risch	Expression too large to display	3570

input `int((f*x+e)^3*arctan(coth(b*x+a)),x,method=_RETURNVERBOSE)`

output

```

3/16*I*f^3*polylog(5,-I*exp(2*b*x+2*a))/b^4+1/8*I*f^3*ln(exp(2*b*x+2*a)-I)
*x^4+1/2*I*ln(exp(2*b*x+2*a)-I)*x*e^3+1/8*I/f*ln(exp(2*b*x+2*a)-I)*e^4+1/2
*I*f^2*ln(exp(2*b*x+2*a)-I)*x^3*e+3/4*I*f*ln(exp(2*b*x+2*a)-I)*x^2*e^2-1/8
*I*(f*x+e)^4/f*ln(exp(2*b*x+2*a)+I)-3/2*I*f^2/b^2*e*ln(1-I*exp(2*b*x+2*a))
*a^2*x+3/2*I*f/b*e^2*ln(1-I*exp(2*b*x+2*a))*a*x+3/2*I*f^2/b^2*a^2*e*ln(((
-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))*x+3/2*I*f^2/b^2*a^2*e*ln(((I)^(1/2)+exp(
b*x+a))/(-I)^(1/2))*x-3/2*I*f/b*a*e^2*ln(((I)^(1/2)-exp(b*x+a))/(-I)^(1/2)
))*x-3/2*I*f/b*a*e^2*ln(((I)^(1/2)+exp(b*x+a))/(-I)^(1/2))*x+3/2*I*f^2/b^
3*a^3*e*ln(((I)^(1/2)-exp(b*x+a))/(-I)^(1/2))+3/2*I*f^2/b^3*a^3*e*ln(((I)
)^(1/2)+exp(b*x+a))/(-I)^(1/2))+3/2*I*f^2/b^3*a^2*e*dilog(((I)^(1/2)-exp(
b*x+a))/(-I)^(1/2))+3/2*I*f^2/b^3*a^2*e*dilog(((I)^(1/2)+exp(b*x+a))/(-I)
^(1/2))-3/2*I*f/b^2*a^2*e^2*ln(((I)^(1/2)-exp(b*x+a))/(-I)^(1/2))-3/2*I*f
/b^2*a^2*e^2*ln(((I)^(1/2)+exp(b*x+a))/(-I)^(1/2))-3/2*I*f/b^2*a*e^2*dilo
g(((I)^(1/2)-exp(b*x+a))/(-I)^(1/2))-3/2*I*f/b^2*a*e^2*dilog(((I)^(1/2)+
exp(b*x+a))/(-I)^(1/2))-1/2*I*f^2/b^3*a^3*e*ln(exp(2*b*x+2*a)+I)+3/4*I*f/b
^2*a^2*e^2*ln(exp(2*b*x+2*a)+I)-I*f^2/b^3*e*ln(1-I*exp(2*b*x+2*a))*a^3+3/4
*I*f^2/b*e*polylog(2,I*exp(2*b*x+2*a))*x^2-3/4*I*f^2/b^3*e*polylog(2,I*exp
(2*b*x+2*a))*a^2-3/4*I*f^2/b^2*e*polylog(3,I*exp(2*b*x+2*a))*x+3/4*I*f/b^2
*e^2*ln(1-I*exp(2*b*x+2*a))*a^2+3/4*I*f/b*e^2*polylog(2,I*exp(2*b*x+2*a))*
x+3/4*I*f/b^2*e^2*polylog(2,I*exp(2*b*x+2*a))*a+1/2*I*f^3/b^3*ln(1-I*ex...

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1460 vs. $2(236) = 472$.

Time = 0.19 (sec) , antiderivative size = 1460, normalized size of antiderivative = 4.88

$$\int (e + fx)^3 \arctan(\coth(a + bx)) dx = \text{Too large to display}$$

input

```
integrate((f*x+e)^3*arctan(coth(b*x+a)),x, algorithm="fricas")
```

output

```

1/8*(-24*I*f^3*polylog(5, 1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) -
24*I*f^3*polylog(5, -1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 24*
I*f^3*polylog(5, 1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 24*I*f^
3*polylog(5, -1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 2*(b^4*f^3
*x^4 + 4*b^4*e*f^2*x^3 + 6*b^4*e^2*f*x^2 + 4*b^4*e^3*x)*arctan(cosh(b*x +
a)/sinh(b*x + a)) - 4*(-I*b^3*f^3*x^3 - 3*I*b^3*e*f^2*x^2 - 3*I*b^3*e^2*f*x
- I*b^3*e^3)*dilog(1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 4*(-
I*b^3*f^3*x^3 - 3*I*b^3*e*f^2*x^2 - 3*I*b^3*e^2*f*x - I*b^3*e^3)*dilog(-1/
2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 4*(I*b^3*f^3*x^3 + 3*I*b^3*
e*f^2*x^2 + 3*I*b^3*e^2*f*x + I*b^3*e^3)*dilog(1/2*sqrt(-4*I)*(cosh(b*x +
a) + sinh(b*x + a))) - 4*(I*b^3*f^3*x^3 + 3*I*b^3*e*f^2*x^2 + 3*I*b^3*e^2*
f*x + I*b^3*e^3)*dilog(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) +
(I*b^4*f^3*x^4 + 4*I*b^4*e*f^2*x^3 + 6*I*b^4*e^2*f*x^2 + 4*I*b^4*e^3*x + 4
*I*a*b^3*e^3 - 6*I*a^2*b^2*e^2*f + 4*I*a^3*b*e*f^2 - I*a^4*f^3)*log(1/2*sq
rt(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^4*f^3*x^4 + 4*I*b^4*e*
f^2*x^3 + 6*I*b^4*e^2*f*x^2 + 4*I*b^4*e^3*x + 4*I*a*b^3*e^3 - 6*I*a^2*b^2*
e^2*f + 4*I*a^3*b*e*f^2 - I*a^4*f^3)*log(-1/2*sqrt(4*I)*(cosh(b*x + a) + s
inh(b*x + a)) + 1) + (-I*b^4*f^3*x^4 - 4*I*b^4*e*f^2*x^3 - 6*I*b^4*e^2*f*x
^2 - 4*I*b^4*e^3*x - 4*I*a*b^3*e^3 + 6*I*a^2*b^2*e^2*f - 4*I*a^3*b*e*f^2 +
I*a^4*f^3)*log(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (...

```

Sympy [F]

$$\int (e + fx)^3 \arctan(\coth(a + bx)) dx = \int (e + fx)^3 \operatorname{atan}(\coth(a + bx)) dx$$

input

```
integrate((f*x+e)**3*atan(coth(b*x+a)),x)
```

output

```
Integral((e + f*x)**3*atan(coth(a + b*x)), x)
```

Maxima [F]

$$\int (e + fx)^3 \arctan(\coth(a + bx)) dx = \int (fx + e)^3 \arctan(\coth(bx + a)) dx$$

input `integrate((f*x+e)^3*arctan(coth(b*x+a)),x, algorithm="maxima")`

output `1/4*(f^3*x^4 + 4*e*f^2*x^3 + 6*e^2*f*x^2 + 4*e^3*x)*arctan2(e^(2*b*x + 2*a) + 1, e^(2*b*x + 2*a) - 1) + integrate(1/2*(b*f^3*x^4*e^(2*a) + 4*b*e*f^2*x^3*e^(2*a) + 6*b*e^2*f*x^2*e^(2*a) + 4*b*e^3*x*e^(2*a))*e^(2*b*x)/(e^(4*b*x + 4*a) + 1), x)`

Giac [F(-1)]

Timed out.

$$\int (e + fx)^3 \arctan(\coth(a + bx)) dx = \text{Timed out}$$

input `integrate((f*x+e)^3*arctan(coth(b*x+a)),x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^3 \arctan(\coth(a + bx)) dx = \int \operatorname{atan}(\coth(a + bx)) (e + fx)^3 dx$$

input `int(atan(coth(a + b*x))*(e + f*x)^3,x)`

output `int(atan(coth(a + b*x))*(e + f*x)^3, x)`

Reduce [F]

$$\begin{aligned} \int (e + fx)^3 \arctan(\coth(a + bx)) dx &= \left(\int \operatorname{atan}(\coth(bx + a)) dx \right) e^3 \\ &+ \left(\int \operatorname{atan}(\coth(bx + a)) x^3 dx \right) f^3 \\ &+ 3 \left(\int \operatorname{atan}(\coth(bx + a)) x^2 dx \right) e f^2 \\ &+ 3 \left(\int \operatorname{atan}(\coth(bx + a)) x dx \right) e^2 f \end{aligned}$$

input `int((f*x+e)^3*atan(coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)),x)*e**3 + int(atan(coth(a + b*x))*x**3,x)*f**3 + 3
*int(atan(coth(a + b*x))*x**2,x)*e*f**2 + 3*int(atan(coth(a + b*x))*x,x)*e
**2*f`

3.94 $\int (e + fx)^2 \arctan(\coth(a + bx)) dx$

Optimal result	734
Mathematica [A] (verified)	735
Rubi [A] (verified)	735
Maple [C] (warning: unable to verify)	739
Fricas [B] (verification not implemented)	740
Sympy [F]	741
Maxima [F]	741
Giac [F(-1)]	741
Mupad [F(-1)]	742
Reduce [F]	742

Optimal result

Integrand size = 15, antiderivative size = 229

$$\int (e + fx)^2 \arctan(\coth(a + bx)) dx = \frac{(e + fx)^3 \arctan(e^{2a+2bx})}{3f} + \frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} - \frac{i(e + fx)^2 \text{PolyLog}(2, -ie^{2a+2bx})}{4b} + \frac{i(e + fx)^2 \text{PolyLog}(2, ie^{2a+2bx})}{4b} + \frac{if(e + fx) \text{PolyLog}(3, -ie^{2a+2bx})}{4b^2} - \frac{if(e + fx) \text{PolyLog}(3, ie^{2a+2bx})}{4b^2} - \frac{if^2 \text{PolyLog}(4, -ie^{2a+2bx})}{8b^3} + \frac{if^2 \text{PolyLog}(4, ie^{2a+2bx})}{8b^3}$$

$$\begin{aligned}
 & \int (e + fx)^2 \arctan(\coth(a + bx)) dx \\
 & \quad \downarrow \text{5708} \\
 & \frac{b \int (e + fx)^3 \operatorname{sech}(2a + 2bx) dx}{3f} + \frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} + \frac{b \int (e + fx)^3 \csc(2ia + 2ibx + \frac{\pi}{2}) dx}{3f} \\
 & \quad \downarrow \text{4668} \\
 & \frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} + \\
 & \frac{b \left(-\frac{3if \int (e+fx)^2 \log(1-ie^{2a+2bx}) dx}{2b} + \frac{3if \int (e+fx)^2 \log(1+ie^{2a+2bx}) dx}{2b} + \frac{(e+fx)^3 \arctan(e^{2a+2bx})}{b} \right)}{3f} \\
 & \quad \downarrow \text{3011} \\
 & \frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} + \\
 & b \left(\frac{3if \left(\frac{f \int (e+fx) \operatorname{PolyLog}(2, -ie^{2a+2bx}) dx}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \int (e+fx) \operatorname{PolyLog}(2, ie^{2a+2bx}) dx}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{2b} \right) \\
 & \quad \downarrow \text{7163} \\
 & \frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} + \\
 & b \left(\frac{3if \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(3, -ie^{2a+2bx}) dx}{2b} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, ie^{2a+2bx})}{2b} - \frac{f \int \operatorname{PolyLog}(3, ie^{2a+2bx}) dx}{2b} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, ie^{2a+2bx})}{2b} \right)}{2b} \right) \\
 & \quad \downarrow \text{2720}
 \end{aligned}$$

$$\frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} + \frac{3if \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \int e^{-2a-2bx} \operatorname{PolyLog}(3, -ie^{2a+2bx}) de^{2a+2bx}}{4b^2} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \left(\frac{(e+fx)}{2b} \right)}{2b} \right)}{3f}$$

↓ 7143

$$\frac{(e + fx)^3 \arctan(\coth(a + bx))}{3f} + \frac{3if \left(\frac{f \left(\frac{(e+fx) \operatorname{PolyLog}(3, -ie^{2a+2bx})}{2b} - \frac{f \operatorname{PolyLog}(4, -ie^{2a+2bx})}{4b^2} \right)}{b} - \frac{(e+fx)^2 \operatorname{PolyLog}(2, -ie^{2a+2bx})}{2b} \right)}{2b} - \frac{3if \left(\frac{f \left(\frac{(e+fx)^3 \arctan(e^{2a+2bx})}{b} \right)}{2b} \right)}{3f}$$

input `Int[(e + f*x)^2*ArcTan[Coth[a + b*x]],x]`

output `((e + f*x)^3*ArcTan[Coth[a + b*x]])/(3*f) + (b*(((e + f*x)^3*ArcTan[E^(2*a + 2*b*x)])/b + (((3*I)/2)*f*(-1/2*((e + f*x)^2*PolyLog[2, (-I)*E^(2*a + 2*b*x)])/b + (f*(((e + f*x)*PolyLog[3, (-I)*E^(2*a + 2*b*x)])/(2*b) - (f*PolyLog[4, (-I)*E^(2*a + 2*b*x)])/(4*b^2)))/b))/b - (((3*I)/2)*f*(-1/2*((e + f*x)^2*PolyLog[2, I*E^(2*a + 2*b*x)])/b + (f*(((e + f*x)*PolyLog[3, I*E^(2*a + 2*b*x)])/(2*b) - (f*PolyLog[4, I*E^(2*a + 2*b*x)])/(4*b^2)))/b))/b)/(3*f)`

Definitions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.))] * ((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)] * ((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5708 `Int[ArcTan[Coth[(a_.) + (b_.)*(x_)] * ((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a, b, e, f}, x] && IGtQ[m, 0]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 4.63 (sec) , antiderivative size = 2668, normalized size of antiderivative = 11.65

method	result	size
risch	Expression too large to display	2668

input

```
int((f*x+e)^2*arctan(coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/8*I*f^2*polylog(4,I*exp(2*b*x+2*a))/b^3-1/8*I*f^2*polylog(4,-I*exp(2*b*x
+2*a))/b^3+1/6*I*f^2*ln(1-I*exp(2*b*x+2*a))*x^3+1/6*I/f*e^3*ln(exp(2*b*x+2
*a)+I)+1/2*I/b*e^2*dilog(((I)^(1/2)-exp(b*x+a))/(I)^(1/2))+1/2*I/b*e^2*d
ilog(((I)^(1/2)+exp(b*x+a))/(I)^(1/2))+1/2*I*e^2*ln(((I)^(1/2)-exp(b*x+
a))/(I)^(1/2))*x+1/2*I*e^2*ln(((I)^(1/2)+exp(b*x+a))/(I)^(1/2))*x-1/6*I
/f*e^3*ln(-exp(2*b*x+2*a)+I)-1/6*I*f^2*ln(1+I*exp(2*b*x+2*a))*x^3-1/2*I/b*
e^2*dilog(1+exp(b*x+a)*(-1)^(3/4))-1/2*I/b*e^2*dilog(1-exp(b*x+a)*(-1)^(3/
4))-1/2*I*e^2*ln(1+exp(b*x+a)*(-1)^(3/4))*x-1/2*I*e^2*ln(1-exp(b*x+a)*(-1)
^(3/4))*x+1/12*Pi*(csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*(exp(2*b*x+2*a)-I))*c
sgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))-csgn(I/(exp(2*b*x+2*a)-1))*csg
gn(I*(exp(2*b*x+2*a)+I))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)-1))-csg
n(I/(exp(2*b*x+2*a)-1))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))^2+csg
gn(I/(exp(2*b*x+2*a)-1))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)-1))^2-c
sgn(I*(exp(2*b*x+2*a)-I))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))^2+
csgn(I*(exp(2*b*x+2*a)+I))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)-1))^2
+csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))*csgn((1+I)*(exp(2*b*x+2*a)-
I)/(exp(2*b*x+2*a)-1))-csgn((1+I)*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))^2
-csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)-1))*csgn((1-I)*(exp(2*b*x+2*a)+
I)/(exp(2*b*x+2*a)-1))-csgn((1-I)*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)-1))^2
+csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))^3-csgn(I*(exp(2*b*x+2*a)...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1002 vs. $2(180) = 360$.

Time = 0.15 (sec) , antiderivative size = 1002, normalized size of antiderivative = 4.38

$$\int (e + fx)^2 \arctan(\coth(a + bx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)^2*arctan(coth(b*x+a)),x, algorithm="fricas")`

output

```
1/6*(6*I*f^2*polylog(4, 1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 6
*I*f^2*polylog(4, -1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 6*I*f^
2*polylog(4, 1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 6*I*f^2*pol
ylog(4, -1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 2*(b^3*f^2*x^3
+ 3*b^3*e*f*x^2 + 3*b^3*e^2*x)*arctan(cosh(b*x + a)/sinh(b*x + a)) - 3*(-I
*b^2*f^2*x^2 - 2*I*b^2*e*f*x - I*b^2*e^2)*dilog(1/2*sqrt(4*I)*(cosh(b*x +
a) + sinh(b*x + a))) - 3*(-I*b^2*f^2*x^2 - 2*I*b^2*e*f*x - I*b^2*e^2)*dilo
g(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 3*(I*b^2*f^2*x^2 + 2*I
*b^2*e*f*x + I*b^2*e^2)*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a
))) - 3*(I*b^2*f^2*x^2 + 2*I*b^2*e*f*x + I*b^2*e^2)*dilog(-1/2*sqrt(-4*I)*
(cosh(b*x + a) + sinh(b*x + a))) + (I*b^3*f^2*x^3 + 3*I*b^3*e*f*x^2 + 3*I*
b^3*e^2*x + 3*I*a*b^2*e^2 - 3*I*a^2*b*e*f + I*a^3*f^2)*log(1/2*sqrt(4*I)*
(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^3*f^2*x^3 + 3*I*b^3*e*f*x^2 + 3
*I*b^3*e^2*x + 3*I*a*b^2*e^2 - 3*I*a^2*b*e*f + I*a^3*f^2)*log(-1/2*sqrt(4*
I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b^3*f^2*x^3 - 3*I*b^3*e*f*x^
2 - 3*I*b^3*e^2*x - 3*I*a*b^2*e^2 + 3*I*a^2*b*e*f - I*a^3*f^2)*log(1/2*sq
rt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b^3*f^2*x^3 - 3*I*b^3*
e*f*x^2 - 3*I*b^3*e^2*x - 3*I*a*b^2*e^2 + 3*I*a^2*b*e*f - I*a^3*f^2)*log(-1
/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-3*I*a*b^2*e^2 + 3*I
*a^2*b*e*f - I*a^3*f^2)*log(I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x ...
```

Sympy [F]

$$\int (e + fx)^2 \arctan(\coth(a + bx)) dx = \int (e + fx)^2 \operatorname{atan}(\coth(a + bx)) dx$$

input `integrate((f*x+e)**2*atan(coth(b*x+a)),x)`

output `Integral((e + f*x)**2*atan(coth(a + b*x)), x)`

Maxima [F]

$$\int (e + fx)^2 \arctan(\coth(a + bx)) dx = \int (fx + e)^2 \arctan(\coth(bx + a)) dx$$

input `integrate((f*x+e)^2*arctan(coth(b*x+a)),x, algorithm="maxima")`

output `1/3*(f^2*x^3 + 3*e*f*x^2 + 3*e^2*x)*arctan2(e^(2*b*x + 2*a) + 1, e^(2*b*x + 2*a) - 1) + integrate(2/3*(b*f^2*x^3*e^(2*a) + 3*b*e*f*x^2*e^(2*a) + 3*b*e^2*x*e^(2*a))*e^(2*b*x)/(e^(4*b*x + 4*a) + 1), x)`

Giac [F(-1)]

Timed out.

$$\int (e + fx)^2 \arctan(\coth(a + bx)) dx = \text{Timed out}$$

input `integrate((f*x+e)^2*arctan(coth(b*x+a)),x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int (e + fx)^2 \arctan(\coth(a + bx)) dx = \int \operatorname{atan}(\coth(a + bx)) (e + fx)^2 dx$$

input `int(atan(coth(a + b*x))*(e + f*x)^2,x)`output `int(atan(coth(a + b*x))*(e + f*x)^2, x)`**Reduce [F]**

$$\begin{aligned} \int (e + fx)^2 \arctan(\coth(a + bx)) dx &= \left(\int \operatorname{atan}(\coth(bx + a)) dx \right) e^2 \\ &+ \left(\int \operatorname{atan}(\coth(bx + a)) x^2 dx \right) f^2 \\ &+ 2 \left(\int \operatorname{atan}(\coth(bx + a)) x dx \right) ef \end{aligned}$$

input `int((f*x+e)^2*atan(coth(b*x+a)),x)`output `int(atan(coth(a + b*x)),x)*e**2 + int(atan(coth(a + b*x))*x**2,x)*f**2 + 2
*int(atan(coth(a + b*x))*x,x)*e*f`

3.95 $\int (e + fx) \arctan(\coth(a + bx)) dx$

Optimal result	743
Mathematica [A] (verified)	744
Rubi [A] (verified)	744
Maple [C] (warning: unable to verify)	747
Fricas [B] (verification not implemented)	748
Sympy [F]	749
Maxima [F]	749
Giac [F]	749
Mupad [F(-1)]	750
Reduce [F]	750

Optimal result

Integrand size = 13, antiderivative size = 159

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \frac{(e + fx)^2 \arctan(e^{2a+2bx})}{2f} + \frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f} - \frac{i(e + fx) \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} + \frac{i(e + fx) \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b} + \frac{if \operatorname{PolyLog}(3, -ie^{2a+2bx})}{8b^2} - \frac{if \operatorname{PolyLog}(3, ie^{2a+2bx})}{8b^2}$$

output

```
1/2*(f*x+e)^2*arctan(exp(2*b*x+2*a))/f+1/2*(f*x+e)^2*arctan(coth(b*x+a))/f
-1/4*I*(f*x+e)*polylog(2,-I*exp(2*b*x+2*a))/b+1/4*I*(f*x+e)*polylog(2,I*ex
p(2*b*x+2*a))/b+1/8*I*f*polylog(3,-I*exp(2*b*x+2*a))/b^2-1/8*I*f*polylog(3
,I*exp(2*b*x+2*a))/b^2
```


Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.49

$$\int (e + fx) \arctan(\coth(a + bx)) dx = ex \arctan(\coth(a + bx)) + \frac{1}{2} fx^2 \arctan(\coth(a + bx)) + \frac{ie(2bx(\log(1 - ie^{2(a+bx)}) - \log(1 + ie^{2(a+bx)})) - \text{PolyLog}(2, -ie^{2(a+bx)}) + \text{PolyLog}(2, ie^{2(a+bx)}))}{4b} + \frac{if(2b^2x^2 \log(1 - ie^{2(a+bx)}) - 2b^2x^2 \log(1 + ie^{2(a+bx)}) - 2bx \text{PolyLog}(2, -ie^{2(a+bx)}) + 2bx \text{PolyLog}(2, ie^{2(a+bx)}))}{8b^2}$$

input

```
Integrate[(e + f*x)*ArcTan[Coth[a + b*x]],x]
```

output

```
e*x*ArcTan[Coth[a + b*x]] + (f*x^2*ArcTan[Coth[a + b*x]])/2 + ((I/4)*e*(2*b*x*(Log[1 - I*E^(2*(a + b*x))] - Log[1 + I*E^(2*(a + b*x))]) - PolyLog[2, (-I)*E^(2*(a + b*x))] + PolyLog[2, I*E^(2*(a + b*x))])/b + ((I/8)*f*(2*b^2*x^2*Log[1 - I*E^(2*(a + b*x))] - 2*b^2*x^2*Log[1 + I*E^(2*(a + b*x))] - 2*b*x*PolyLog[2, (-I)*E^(2*(a + b*x))] + 2*b*x*PolyLog[2, I*E^(2*(a + b*x))]) + PolyLog[3, (-I)*E^(2*(a + b*x))] - PolyLog[3, I*E^(2*(a + b*x))])/b^2
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.10, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {5708, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (e + fx) \arctan(\coth(a + bx)) dx$$

$$\downarrow \text{5708}$$

$$\frac{b \int (e + fx)^2 \operatorname{sech}(2a + 2bx) dx}{2f} + \frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f}$$

$$\downarrow \text{3042}$$

$$\frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f} + \frac{b \int (e + fx)^2 \csc(2ia + 2ibx + \frac{\pi}{2}) dx}{2f}$$

↓ 4668

$$\frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f} + \frac{b \left(-\frac{if \int (e + fx) \log(1 - ie^{2a + 2bx}) dx}{b} + \frac{if \int (e + fx) \log(1 + ie^{2a + 2bx}) dx}{b} + \frac{(e + fx)^2 \arctan(e^{2a + 2bx})}{b} \right)}{2f}$$

↓ 3011

$$\frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f} + \frac{b \left(\frac{if \left(\frac{f \int \text{PolyLog}(2, -ie^{2a + 2bx}) dx}{2b} - \frac{(e + fx) \text{PolyLog}(2, -ie^{2a + 2bx})}{2b} \right)}{b} - \frac{if \left(\frac{f \int \text{PolyLog}(2, ie^{2a + 2bx}) dx}{2b} - \frac{(e + fx) \text{PolyLog}(2, ie^{2a + 2bx})}{2b} \right)}{b} \right) + \frac{(e + fx)^2 \arctan(e^{2a + 2bx})}{b}}{2f}$$

↓ 2720

$$\frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f} + \frac{b \left(\frac{if \left(\frac{f \int e^{-2a - 2bx} \text{PolyLog}(2, -ie^{2a + 2bx}) de^{2a + 2bx}}{4b^2} - \frac{(e + fx) \text{PolyLog}(2, -ie^{2a + 2bx})}{2b} \right)}{b} - \frac{if \left(\frac{f \int e^{-2a - 2bx} \text{PolyLog}(2, ie^{2a + 2bx}) de^{2a + 2bx}}{4b^2} - \frac{(e + fx) \text{PolyLog}(2, ie^{2a + 2bx})}{2b} \right)}{b} \right)}{2f}$$

↓ 7143

$$\frac{(e + fx)^2 \arctan(\coth(a + bx))}{2f} + \frac{b \left(\frac{(e + fx)^2 \arctan(e^{2a + 2bx})}{b} + \frac{if \left(\frac{f \text{PolyLog}(3, -ie^{2a + 2bx})}{4b^2} - \frac{(e + fx) \text{PolyLog}(2, -ie^{2a + 2bx})}{2b} \right)}{b} - \frac{if \left(\frac{f \text{PolyLog}(3, ie^{2a + 2bx})}{4b^2} - \frac{(e + fx) \text{PolyLog}(2, ie^{2a + 2bx})}{2b} \right)}{b} \right)}{2f}$$

input

```
Int[(e + f*x)*ArcTan[Coth[a + b*x]], x]
```

output

```
((e + f*x)^2*ArcTan[Coth[a + b*x]])/(2*f) + (b*(((e + f*x)^2*ArcTan[E^(2*a + 2*b*x)])/b + (I*f*(-1/2*((e + f*x)*PolyLog[2, (-I)*E^(2*a + 2*b*x)]))/b + (f*PolyLog[3, (-I)*E^(2*a + 2*b*x)])/(4*b^2)))/b - (I*f*(-1/2*((e + f*x)*PolyLog[2, I*E^(2*a + 2*b*x)]))/b + (f*PolyLog[3, I*E^(2*a + 2*b*x)])/(4*b^2))/b)/(2*f)
```

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 3042

```
Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4668

```
Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 5708

```
Int[ArcTan[Coth[(a_.) + (b_.)*(x_)]]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[Coth[a + b*x]]/(f*(m + 1))), x] + Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a, b, e, f}, x] && IGtQ[m, 0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.60 (sec) , antiderivative size = 1777, normalized size of antiderivative = 11.18

method	result	size
risch	Expression too large to display	1777

input

```
int((f*x+e)*arctan(coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/8*I*f*polylog(3,-I*exp(2*b*x+2*a))/b^2+1/4*I/b^2*f*a^2*ln(exp(2*b*x+2*a)
+I)-1/2*I/b*e*a*ln(exp(2*b*x+2*a)+I)-1/8*I*f*polylog(3,I*exp(2*b*x+2*a))/b
^2+1/2*I*ln(exp(2*b*x+2*a)-I)*e*x+1/4*I*ln(exp(2*b*x+2*a)-I)*f*x^2-1/2*I*f
/b^2*a^2*ln((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))-1/2*I*f/b^2*a^2*ln((-I)^(
1/2)+exp(b*x+a))/(-I)^(1/2))-1/2*I*f/b^2*a*dilog((-I)^(1/2)-exp(b*x+a))/(-
I)^(1/2))-1/2*I*f/b^2*a*dilog((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2))+1/2*I*e
*ln((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))*x+1/2*I*e*ln((-I)^(1/2)+exp(b*x+a)
))/(-I)^(1/2))*x+1/2*I*e/b*dilog((-I)^(1/2)-exp(b*x+a))/(-I)^(1/2))+1/2*I
*e/b*dilog((-I)^(1/2)+exp(b*x+a))/(-I)^(1/2))-1/2*I*f/b*ln(1+I*exp(2*b*x+
2*a))*a*x+1/2*I*f/b*a*ln(1+exp(b*x+a)*(-1)^(3/4))*x+1/2*I*(-1/2*f*x^2-e*x)
*ln(exp(2*b*x+2*a)+I)+1/4*Pi*(csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*(exp(2*b*x
+2*a)-I))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))-csgn(I/(exp(2*b*x+
2*a)-1))*csgn(I*(exp(2*b*x+2*a)+I))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2
*a)-1))-csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*
a)-1))^2+csgn(I/(exp(2*b*x+2*a)-1))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2
*a)-1))^2-csgn(I*(exp(2*b*x+2*a)-I))*csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+
2*a)-1))^2+csgn(I*(exp(2*b*x+2*a)+I))*csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x
+2*a)-1))^2+csgn(I*(exp(2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))*csgn((1+I)*(exp(
2*b*x+2*a)-I)/(exp(2*b*x+2*a)-1))-csgn((1+I)*(exp(2*b*x+2*a)-I)/(exp(2*b*x
+2*a)-1))^2-csgn(I*(exp(2*b*x+2*a)+I)/(exp(2*b*x+2*a)-1))*csgn((1-I)*(e...
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 600 vs. $2(130) = 260$.

Time = 0.16 (sec) , antiderivative size = 600, normalized size of antiderivative = 3.77

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \text{Too large to display}$$

input `integrate((f*x+e)*arctan(coth(b*x+a)),x, algorithm="fricas")`

output

```
1/4*(2*(b^2*f*x^2 + 2*b^2*e*x)*arctan(cosh(b*x + a)/sinh(b*x + a)) - 2*(-I
*b*f*x - I*b*e)*dilog(1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) - 2*(
-I*b*f*x - I*b*e)*dilog(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) -
2*(I*b*f*x + I*b*e)*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)))
- 2*(I*b*f*x + I*b*e)*dilog(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)
)) + (I*b^2*f*x^2 + 2*I*b^2*e*x + 2*I*a*b*e - I*a^2*f)*log(1/2*sqrt(4*I)*(
cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^2*f*x^2 + 2*I*b^2*e*x + 2*I*a*b
*e - I*a^2*f)*log(-1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-
I*b^2*f*x^2 - 2*I*b^2*e*x - 2*I*a*b*e + I*a^2*f)*log(1/2*sqrt(-4*I)*(cosh(
b*x + a) + sinh(b*x + a)) + 1) + (-I*b^2*f*x^2 - 2*I*b^2*e*x - 2*I*a*b*e +
I*a^2*f)*log(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-2*I
*a*b*e + I*a^2*f)*log(I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) + (
-2*I*a*b*e + I*a^2*f)*log(-I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)
) + (2*I*a*b*e - I*a^2*f)*log(I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh(b*x
+ a)) + (2*I*a*b*e - I*a^2*f)*log(-I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh
(b*x + a)) - 2*I*f*polylog(3, 1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a)
)) - 2*I*f*polylog(3, -1/2*sqrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 2*
I*f*polylog(3, 1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))) + 2*I*f*pol
ylog(3, -1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a)))/b^2
```

Sympy [F]

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \int (e + fx) \operatorname{atan}(\coth(a + bx)) dx$$

input `integrate((f*x+e)*atan(coth(b*x+a)),x)`

output `Integral((e + f*x)*atan(coth(a + b*x)), x)`

Maxima [F]

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \int (fx + e) \arctan(\coth(bx + a)) dx$$

input `integrate((f*x+e)*arctan(coth(b*x+a)),x, algorithm="maxima")`

output `1/2*(f*x^2 + 2*e*x)*arctan2(e^(2*b*x + 2*a) + 1, e^(2*b*x + 2*a) - 1) + integrate((b*f*x^2*e^(2*a) + 2*b*e*x*e^(2*a))*e^(2*b*x)/(e^(4*b*x + 4*a) + 1), x)`

Giac [F]

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \int (fx + e) \arctan(\coth(bx + a)) dx$$

input `integrate((f*x+e)*arctan(coth(b*x+a)),x, algorithm="giac")`

output `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \int \operatorname{atan}(\coth(a + bx)) (e + fx) dx$$

input `int(atan(coth(a + b*x))*(e + f*x),x)`output `int(atan(coth(a + b*x))*(e + f*x), x)`**Reduce [F]**

$$\int (e + fx) \arctan(\coth(a + bx)) dx = \left(\int \operatorname{atan}(\coth(bx + a)) dx \right) e + \left(\int \operatorname{atan}(\coth(bx + a)) x dx \right) f$$

input `int((f*x+e)*atan(coth(b*x+a)),x)`output `int(atan(coth(a + b*x)),x)*e + int(atan(coth(a + b*x))*x,x)*f`

3.96 $\int \arctan(\coth(a + bx)) dx$

Optimal result	751
Mathematica [A] (verified)	751
Rubi [A] (verified)	752
Maple [A] (verified)	754
Fricas [B] (verification not implemented)	754
Sympy [F]	755
Maxima [F]	755
Giac [F]	756
Mupad [F(-1)]	756
Reduce [F]	756

Optimal result

Integrand size = 7, antiderivative size = 73

$$\int \arctan(\coth(a + bx)) dx = x \arctan(e^{2a+2bx}) + x \arctan(\coth(a + bx)) - \frac{i \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b} + \frac{i \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b}$$

output `x*arctan(exp(2*b*x+2*a))+x*arctan(coth(b*x+a))-1/4*I*polylog(2,-I*exp(2*b*x+2*a))/b+1/4*I*polylog(2,I*exp(2*b*x+2*a))/b`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.25

$$\int \arctan(\coth(a + bx)) dx = x \arctan(\coth(a + bx)) + \frac{i(2bx(\log(1 - ie^{2(a+bx)}) - \log(1 + ie^{2(a+bx)})) - \operatorname{PolyLog}(2, -ie^{2(a+bx)}) + \operatorname{PolyLog}(2, ie^{2(a+bx)}))}{4b}$$

input `Integrate[ArcTan[Coth[a + b*x]],x]`

output

```
x*ArcTan[Coth[a + b*x]] + ((1/4)*(2*b*x*(Log[1 - I*E^(2*(a + b*x))] - Log[1 + I*E^(2*(a + b*x))]) - PolyLog[2, (-I)*E^(2*(a + b*x))] + PolyLog[2, I*E^(2*(a + b*x))])/b
```

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {5704, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(\coth(a + bx)) dx$$

$$\downarrow 5704$$

$$b \int x \operatorname{sech}(2a + 2bx) dx + x \arctan(\coth(a + bx))$$

$$\downarrow 3042$$

$$x \arctan(\coth(a + bx)) + b \int x \csc\left(2ia + 2ibx + \frac{\pi}{2}\right) dx$$

$$\downarrow 4668$$

$$x \arctan(\coth(a + bx)) + b \left(-\frac{i \int \log(1 - ie^{2a+2bx}) dx}{2b} + \frac{i \int \log(1 + ie^{2a+2bx}) dx}{2b} + \frac{x \arctan(e^{2a+2bx})}{b} \right)$$

$$\downarrow 2715$$

$$b \left(-\frac{i \int e^{-2a-2bx} \log(1 - ie^{2a+2bx}) de^{2a+2bx}}{4b^2} + \frac{i \int e^{-2a-2bx} \log(1 + ie^{2a+2bx}) de^{2a+2bx}}{4b^2} + \frac{x \arctan(e^{2a+2bx})}{b} \right)$$

$$\downarrow 2838$$

$$b \left(\frac{x \arctan(e^{2a+2bx})}{b} - \frac{i \operatorname{PolyLog}(2, -ie^{2a+2bx})}{4b^2} + \frac{i \operatorname{PolyLog}(2, ie^{2a+2bx})}{4b^2} \right)$$

input `Int[ArcTan[Coth[a + b*x]],x]`

output `x*ArcTan[Coth[a + b*x]] + b*((x*ArcTan[E^(2*a + 2*b*x)])/b - ((I/4)*PolyLog[2, (-I)*E^(2*a + 2*b*x)])/b^2 + ((I/4)*PolyLog[2, I*E^(2*a + 2*b*x)])/b^2)`

Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_)]], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 5704 `Int[ArcTan[Coth[(a_) + (b_)*(x_)]], x_Symbol] :> Simp[x*ArcTan[Coth[a + b*x]], x] + Simp[b Int[x*Sech[2*a + 2*b*x], x], x] /; FreeQ[{a, b}, x]`

Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.45

method	result
parts	$x \arctan(\coth(bx + a)) - \frac{i(bx+a)(\ln(1-ie^{2bx+2a}) - \ln(1+ie^{2bx+2a}))}{2} + \frac{i \operatorname{dilog}(1+ie^{2bx+2a})}{4} - \frac{i \operatorname{dilog}(1-ie^{2bx+2a})}{4}$
derivativedivides	$\frac{\operatorname{arctanh}(\coth(bx+a)) \arctan(\coth(bx+a)) - \frac{i \operatorname{arctanh}(\coth(bx+a)) \left(\ln\left(1 - \frac{i(\coth(bx+a)+1)^2}{1-\coth(bx+a)^2}\right) - \ln\left(1 + \frac{i(\coth(bx+a)+1)^2}{1-\coth(bx+a)^2}\right) \right)}{2}}{b}$
default	$\frac{\operatorname{arctanh}(\coth(bx+a)) \arctan(\coth(bx+a)) - \frac{i \operatorname{arctanh}(\coth(bx+a)) \left(\ln\left(1 - \frac{i(\coth(bx+a)+1)^2}{1-\coth(bx+a)^2}\right) - \ln\left(1 + \frac{i(\coth(bx+a)+1)^2}{1-\coth(bx+a)^2}\right) \right)}{2}}{b}$
risch	Expression too large to display

```
input int(arctan(coth(b*x+a)), x, method=_RETURNVERBOSE)
```

```
output x*arctan(coth(b*x+a))-1/b*(-1/2*I*(b*x+a)*(ln(1-I*exp(2*b*x+2*a))-ln(1+I*exp(2*b*x+2*a)))+1/4*I*dilog(1+I*exp(2*b*x+2*a))-1/4*I*dilog(1-I*exp(2*b*x+2*a))+a*arctan(exp(2*b*x+2*a)))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(56) = 112.

Time = 0.14 (sec) , antiderivative size = 334, normalized size of antiderivative = 4.58

$$\int \arctan(\coth(a + bx)) dx$$

$$= \frac{2bx \arctan\left(\frac{\cosh(bx+a)}{\sinh(bx+a)}\right) + (ibx + ia) \log\left(\frac{1}{2} \sqrt{4i}(\cosh(bx+a) + \sinh(bx+a)) + 1\right) + (ibx + ia) \log\left(\frac{1}{2} \sqrt{4i}(\cosh(bx+a) - \sinh(bx+a)) + 1\right)}{b}$$

```
input integrate(arctan(coth(b*x+a)), x, algorithm="fricas")
```

output

```

1/2*(2*b*x*arctan(cosh(b*x + a)/sinh(b*x + a)) + (I*b*x + I*a)*log(1/2*sqrt(
4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b*x + I*a)*log(-1/2*sqrt(
4*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b*x - I*a)*log(1/2*sqrt(-4
*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*b*x - I*a)*log(-1/2*sqrt(-4
*I)*(cosh(b*x + a) + sinh(b*x + a)) + 1) - I*a*log(I*sqrt(4*I) + 2*cosh(b*
x + a) + 2*sinh(b*x + a)) - I*a*log(-I*sqrt(4*I) + 2*cosh(b*x + a) + 2*sin
h(b*x + a)) + I*a*log(I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) +
I*a*log(-I*sqrt(-4*I) + 2*cosh(b*x + a) + 2*sinh(b*x + a)) + I*dilog(1/2*s
qrt(4*I)*(cosh(b*x + a) + sinh(b*x + a))) + I*dilog(-1/2*sqrt(4*I)*(cosh(b
*x + a) + sinh(b*x + a))) - I*dilog(1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b
*x + a))) - I*dilog(-1/2*sqrt(-4*I)*(cosh(b*x + a) + sinh(b*x + a))))/b

```

Sympy [F]

$$\int \arctan(\coth(a + bx)) dx = \int \operatorname{atan}(\coth(a + bx)) dx$$

input

```
integrate(atan(coth(b*x+a)),x)
```

output

```
Integral(atan(coth(a + b*x)), x)
```

Maxima [F]

$$\int \arctan(\coth(a + bx)) dx = \int \arctan(\coth(bx + a)) dx$$

input

```
integrate(arctan(coth(b*x+a)),x, algorithm="maxima")
```

output

```

x*arctan2(e^(2*b*x + 2*a) + 1, e^(2*b*x + 2*a) - 1) + 2*b*integrate(x*e^(2
*b*x + 2*a)/(e^(4*b*x + 4*a) + 1), x)

```

Giac [F]

$$\int \arctan(\coth(a + bx)) dx = \int \arctan(\coth(bx + a)) dx$$

input `integrate(arctan(coth(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(coth(b*x + a)), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(\coth(a + bx)) dx = \int \operatorname{atan}(\coth(a + bx)) dx$$

input `int(atan(coth(a + b*x)),x)`

output `int(atan(coth(a + b*x)), x)`

Reduce [F]

$$\int \arctan(\coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a)) dx$$

input `int(atan(coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)),x)`

3.97 $\int \frac{\arctan(\coth(a+bx))}{e+fx} dx$

Optimal result	757
Mathematica [N/A]	757
Rubi [N/A]	758
Maple [N/A]	758
Fricas [N/A]	759
Sympy [F(-1)]	759
Maxima [N/A]	759
Giac [N/A]	760
Mupad [N/A]	760
Reduce [N/A]	760

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \text{Int}\left(\frac{\arctan(\coth(a + bx))}{e + fx}, x\right)$$

output `Defer(Int)(arctan(coth(b*x+a))/(f*x+e), x)`

Mathematica [N/A]

Not integrable

Time = 0.66 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \int \frac{\arctan(\coth(a + bx))}{e + fx} dx$$

input `Integrate[ArcTan[Coth[a + b*x]]/(e + f*x), x]`

output `Integrate[ArcTan[Coth[a + b*x]]/(e + f*x), x]`

Rubi [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx$$

↓ 7299

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx$$

input `Int[ArcTan[Coth[a + b*x]]/(e + f*x),x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(\coth(bx + a))}{fx + e} dx$$

input `int(arctan(coth(b*x+a))/(f*x+e),x)`

output `int(arctan(coth(b*x+a))/(f*x+e),x)`

Fricas [N/A]

Not integrable

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \int \frac{\arctan(\coth(bx + a))}{fx + e} dx$$

input `integrate(arctan(coth(b*x+a))/(f*x+e),x, algorithm="fricas")`

output `integral(arctan(coth(b*x + a))/(f*x + e), x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \text{Timed out}$$

input `integrate(atan(coth(b*x+a))/(f*x+e),x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 1.96 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \int \frac{\arctan(\coth(bx + a))}{fx + e} dx$$

input `integrate(arctan(coth(b*x+a))/(f*x+e),x, algorithm="maxima")`

output `integrate(arctan(coth(b*x + a))/(f*x + e), x)`

Giac [N/A]

Not integrable

Time = 74.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.20

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \int \frac{\arctan(\coth(bx + a))}{fx + e} dx$$

input `integrate(arctan(coth(b*x+a))/(f*x+e),x, algorithm="giac")`

output `sage0*x`

Mupad [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \int \frac{\operatorname{atan}(\coth(a + bx))}{e + fx} dx$$

input `int(atan(coth(a + b*x))/(e + f*x),x)`

output `int(atan(coth(a + b*x))/(e + f*x), x)`

Reduce [N/A]

Not integrable

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(\coth(a + bx))}{e + fx} dx = \int \frac{\operatorname{atan}(\coth(bx + a))}{fx + e} dx$$

input `int(atan(coth(b*x+a))/(f*x+e),x)`

output `int(atan(coth(a + b*x))/(e + f*x),x)`

3.98 $\int x^2 \arctan(c + d \coth(a + bx)) dx$

Optimal result	762
Mathematica [A] (warning: unable to verify)	763
Rubi [A] (verified)	764
Maple [C] (warning: unable to verify)	768
Fricas [B] (verification not implemented)	769
Sympy [F(-1)]	770
Maxima [F]	770
Giac [F]	770
Mupad [F(-1)]	771
Reduce [F]	771

Optimal result

Integrand size = 15, antiderivative size = 351

$$\begin{aligned}
 \int x^2 \arctan(c + d \coth(a + bx)) dx = & \frac{1}{3}x^3 \arctan(c + d \coth(a + bx)) \\
 & + \frac{1}{6}ix^3 \log\left(1 - \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right) \\
 & - \frac{1}{6}ix^3 \log\left(1 - \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right) \\
 & + \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(i-c-d)e^{2a+2bx}}{i-c+d}\right)}{4b} \\
 & - \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{(i+c+d)e^{2a+2bx}}{i+c-d}\right)}{4b} \\
 & - \frac{ix \operatorname{PolyLog}\left(3, \frac{(i-c-d)e^{2a+2bx}}{i-c+d}\right)}{4b^2} \\
 & + \frac{ix \operatorname{PolyLog}\left(3, \frac{(i+c+d)e^{2a+2bx}}{i+c-d}\right)}{4b^2} \\
 & + \frac{i \operatorname{PolyLog}\left(4, \frac{(i-c-d)e^{2a+2bx}}{i-c+d}\right)}{8b^3} \\
 & - \frac{i \operatorname{PolyLog}\left(4, \frac{(i+c+d)e^{2a+2bx}}{i+c-d}\right)}{8b^3}
 \end{aligned}$$

output

$$\begin{aligned} & \frac{1}{3}x^3 \arctan(c+d\coth(bx+a)) + \frac{1}{6}I^3 x^3 \ln(1 - (I-c-d)\exp(2bx+2a)/(I-c+d)) \\ & - \frac{1}{6}I^3 x^3 \ln(1 - (I+c+d)\exp(2bx+2a)/(I+c-d)) + \frac{1}{4}I^2 x^2 \operatorname{polylog}(2, (I-c-d)\exp(2bx+2a)/(I-c-d)) \\ & /b - \frac{1}{4}I^2 x^2 \operatorname{polylog}(2, (I+c+d)\exp(2bx+2a)/(I+c-d)) /b - \frac{1}{4}I^2 x^2 \operatorname{polylog}(3, (I-c-d)\exp(2bx+2a)/(I-c-d)) /b^2 \\ & + \frac{1}{4}I^2 x^2 \operatorname{polylog}(3, (I+c+d)\exp(2bx+2a)/(I+c-d)) /b^2 + \frac{1}{8}I \operatorname{polylog}(4, (I-c-d)\exp(2bx+2a)/(I-c-d)) /b^3 \\ & - \frac{1}{8}I \operatorname{polylog}(4, (I+c+d)\exp(2bx+2a)/(I+c-d)) /b^3 \end{aligned}$$
Mathematica [A] (warning: unable to verify)

Time = 0.87 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.26

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \frac{1}{3}x^3 \arctan(c + d \coth(a + bx)) + \frac{d \left(4b^3 x^3 \log \left(1 - \frac{(1+(c+d)^2)e^{2(a+bx)}}{1+c^2-d^2+2\sqrt{-d^2}} \right) - 4b^3 x^3 \log \left(1 + \frac{(1+(c+d)^2)e^{2(a+bx)}}{-1-c^2+d^2+2\sqrt{-d^2}} \right) + 6b^2 x^2 \operatorname{PolyLog} \left(2, \frac{1+c^2+2cd+d^2}{1+c^2-d^2+2\sqrt{-d^2}} \right) \right)}{24b^3 \sqrt{-d^2}}$$

input

Integrate[x^2*ArcTan[c + d*Coth[a + b*x]],x]

output

$$\begin{aligned} & (x^3 \operatorname{ArcTan}[c + d \operatorname{Coth}[a + b x]])/3 + (d*(4b^3 x^3 \operatorname{Log}[1 - ((1 + (c + d)^2)E^{2(a + b x)})/(1 + c^2 - d^2 + 2\sqrt{-d^2})] - 4b^3 x^3 \operatorname{Log}[1 + ((1 + (c + d)^2)E^{2(a + b x)})/(-1 - c^2 + d^2 + 2\sqrt{-d^2})] + 6b^2 x^2 \operatorname{PolyLog}[2, ((1 + c^2 + 2cd + d^2)E^{2(a + b x)})/(1 + c^2 - d^2 + 2\sqrt{-d^2})] - 6b^2 x^2 \operatorname{PolyLog}[2, -(((1 + (c + d)^2)E^{2(a + b x)})/(-1 - c^2 + d^2 + 2\sqrt{-d^2}))] - 6b^2 x^2 \operatorname{PolyLog}[3, ((1 + c^2 + 2cd + d^2)E^{2(a + b x)})/(1 + c^2 - d^2 + 2\sqrt{-d^2})] + 6b^2 x^2 \operatorname{PolyLog}[3, -(((1 + (c + d)^2)E^{2(a + b x)})/(-1 - c^2 + d^2 + 2\sqrt{-d^2}))] - 3 \operatorname{PolyLog}[4, ((1 + c^2 + 2cd + d^2)E^{2(a + b x)})/(1 + c^2 - d^2 - 2\sqrt{-d^2})] + 3 \operatorname{PolyLog}[4, ((1 + c^2 + 2cd + d^2)E^{2(a + b x)})/(1 + c^2 - d^2 + 2\sqrt{-d^2})])]/(24b^3 \sqrt{-d^2})) \end{aligned}$$

Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 437, normalized size of antiderivative = 1.25, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5724, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(d \coth(a + bx) + c) dx \\
 & \quad \downarrow \text{5724} \\
 & \frac{1}{3}b(1 + i(c + d)) \int \frac{e^{2a+2bx} x^3}{-c - (-c - d + i)e^{2a+2bx} + d + i} dx - \frac{1}{3}b(1 - i(c + d)) \int \frac{e^{2a+2bx} x^3}{c - (c + d + i)e^{2a+2bx} - d + i} dx + \frac{1}{3}x^3 \arctan(d \coth(a + bx) + c) \\
 & \quad \downarrow \text{2620} \\
 & \frac{1}{3}b(1 + i(c + d)) \left(\frac{3 \int x^2 \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) dx}{2b(-c-d+i)} - \frac{x^3 \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) - \\
 & \frac{1}{3}b(1 - i(c + d)) \left(\frac{3 \int x^2 \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) dx}{2b(c+d+i)} - \frac{x^3 \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right) + \\
 & \quad \frac{1}{3}x^3 \arctan(d \coth(a + bx) + c) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$d)) \left(\frac{\frac{1}{3}b(1+i(c + \frac{x \operatorname{PolyLog}\left(3, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b})}{2b(-c-d+i)} - \frac{x^3 \log\left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} \right)$$

$$d)) \left(\frac{\frac{1}{3}b(1-i(c + \frac{x \operatorname{PolyLog}\left(3, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}\left(3, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right) de^{2a+2bx}}{b} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b})}{2b(c+d+i)} - \frac{x^3 \log\left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} \right)$$

$$\frac{1}{3}x^3 \arctan(d \coth(a + bx) + c)$$

↓ 7143

$$d)) \left(\frac{\frac{1}{3}x^3 \arctan(d \coth(a + bx) + c) + \frac{1}{3}b(1+i(c + \frac{x \operatorname{PolyLog}\left(3, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{4b^2} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b})}{2b(-c-d+i)} - \frac{x^3 \log\left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)} \right)$$

$$d)) \left(\frac{\frac{1}{3}b(1-i(c + \frac{x \operatorname{PolyLog}\left(3, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b} - \frac{\operatorname{PolyLog}\left(4, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{4b^2} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b})}{2b(c+d+i)} - \frac{x^3 \log\left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)} \right)$$

input

`Int[x^2*ArcTan[c + d*Coth[a + b*x]],x]`

output

```
(x^3*ArcTan[c + d*Coth[a + b*x]])/3 + (b*(1 + I*(c + d))*(-1/2*(x^3*Log[1
- ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)])/(b*(I - c - d)) + (3*(-1/2*(
x^2*PolyLog[2, ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)])/b + ((x*PolyLog
[3, ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)]/(2*b) - PolyLog[4, ((I - c
- d)*E^(2*a + 2*b*x))/(I - c + d)]/(4*b^2))/b))/(2*b*(I - c - d)))/3 - (
b*(1 - I*(c + d))*(-1/2*(x^3*Log[1 - ((I + c + d)*E^(2*a + 2*b*x))/(I + c
- d)])/(b*(I + c + d)) + (3*(-1/2*(x^2*PolyLog[2, ((I + c + d)*E^(2*a + 2*
b*x))/(I + c - d)])/b + ((x*PolyLog[3, ((I + c + d)*E^(2*a + 2*b*x))/(I +
c - d)]/(2*b) - PolyLog[4, ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)]/(4*
b^2))/b))/(2*b*(I + c + d)))/3
```

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```


rule 5724

```
Int[ArcTan[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Coth[a + b*x]]/(f*(m + 1))), x] + (-Simp[I*b*((I - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(I - c + d - (I - c - d)*E^(2*a + 2*b*x))], x], x] + Simp[I*b*((I + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x))/(I + c - d - (I + c + d)*E^(2*a + 2*b*x))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && NeQ[(c - d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 16.23 (sec) , antiderivative size = 6845, normalized size of antiderivative = 19.50

method	result	size
risch	Expression too large to display	6845

input

```
int(x^2*arctan(c+d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
result too large to display
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1269 vs. $2(259) = 518$.

Time = 0.19 (sec) , antiderivative size = 1269, normalized size of antiderivative = 3.62

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \text{Too large to display}$$

input `integrate(x^2*arctan(c+d*coth(b*x+a)),x, algorithm="fricas")`

output

```
1/6*(2*b^3*x^3*arctan((d*cosh(b*x + a) + c*sinh(b*x + a))/sinh(b*x + a)) +
3*I*b^2*x^2*dilog(sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(
cosh(b*x + a) + sinh(b*x + a))) + 3*I*b^2*x^2*dilog(-sqrt((c^2 - d^2 + 2*I
*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 3*I*b^
2*x^2*dilog(sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*
x + a) + sinh(b*x + a))) - 3*I*b^2*x^2*dilog(-sqrt((c^2 - d^2 - 2*I*d + 1)
/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - I*a^3*log(2*(
c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x
+ a) + 2*(c^2 - d^2 - 2*I*d + 1)*sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d
+ d^2 + 1))) - I*a^3*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2
+ 2*c*d + d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 - 2*I*d + 1)*sqrt((c^2 -
d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) + I*a^3*log(2*(c^2 + 2*c*d + d^
2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(c^2 -
d^2 + 2*I*d + 1)*sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) +
I*a^3*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 +
1)*sinh(b*x + a) - 2*(c^2 - d^2 + 2*I*d + 1)*sqrt((c^2 - d^2 - 2*I*d + 1)
/(c^2 - 2*c*d + d^2 + 1))) - 6*I*b*x*polylog(3, sqrt((c^2 - d^2 + 2*I*d +
1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 6*I*b*x*pol
ylog(3, -sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x +
a) + sinh(b*x + a))) + 6*I*b*x*polylog(3, sqrt((c^2 - d^2 - 2*I*d + 1)...
```

Sympy [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \text{Timed out}$$

input `integrate(x**2*atan(c+d*coth(b*x+a)),x)`

output `Timed out`

Maxima [F]

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \int x^2 \arctan(d \coth(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*coth(b*x+a)),x, algorithm="maxima")`

output `1/3*x^3*arctan2((c*e^(2*a) + d*e^(2*a))*e^(2*b*x) - c + d, e^(2*b*x + 2*a) - 1) + 4*b*d*integrate(1/3*x^3*e^(2*b*x + 2*a)/(c^2 - 2*c*d + d^2 + (c^2*e^(4*a) + 2*c*d*e^(4*a) + d^2*e^(4*a) + e^(4*a))*e^(4*b*x) - 2*(c^2*e^(2*a) - d^2*e^(2*a) + e^(2*a))*e^(2*b*x) + 1), x)`

Giac [F]

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \int x^2 \arctan(d \coth(bx + a) + c) dx$$

input `integrate(x^2*arctan(c+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan(d*coth(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \int x^2 \operatorname{atan}(c + d \coth(a + bx)) dx$$

input `int(x^2*atan(c + d*coth(a + b*x)),x)`output `int(x^2*atan(c + d*coth(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \arctan(c + d \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) d + c) x^2 dx$$

input `int(x^2*atan(c+d*coth(b*x+a)),x)`output `int(atan(coth(a + b*x)*d + c)*x**2,x)`

3.99 $\int x \arctan(c + d \coth(a + bx)) dx$

Optimal result	772
Mathematica [A] (warning: unable to verify)	773
Rubi [A] (verified)	773
Maple [C] (warning: unable to verify)	777
Fricas [B] (verification not implemented)	777
Sympy [F(-1)]	778
Maxima [F]	779
Giac [F]	779
Mupad [F(-1)]	779
Reduce [F]	780

Optimal result

Integrand size = 13, antiderivative size = 265

$$\begin{aligned}
 \int x \arctan(c + d \coth(a + bx)) dx &= \frac{1}{2}x^2 \arctan(c + d \coth(a + bx)) \\
 &+ \frac{1}{4}ix^2 \log\left(1 - \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right) \\
 &- \frac{1}{4}ix^2 \log\left(1 - \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right) \\
 &+ \frac{ix \operatorname{PolyLog}\left(2, \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right)}{4b} \\
 &- \frac{ix \operatorname{PolyLog}\left(2, \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right)}{4b} \\
 &- \frac{i \operatorname{PolyLog}\left(3, \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right)}{8b^2} \\
 &+ \frac{i \operatorname{PolyLog}\left(3, \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right)}{8b^2}
 \end{aligned}$$

output

$$\begin{aligned} & 1/2*x^2*\arctan(c+d*\coth(b*x+a))+1/4*I*x^2*\ln(1-(I-c-d)*\exp(2*b*x+2*a)/(I-c+d))-1/4*I*x^2*\ln(1-(I+c+d)*\exp(2*b*x+2*a)/(I+c-d))+1/4*I*x*\text{polylog}(2,(I-c-d)*\exp(2*b*x+2*a)/(I-c+d))/b-1/4*I*x*\text{polylog}(2,(I+c+d)*\exp(2*b*x+2*a)/(I+c-d))/b-1/8*I*\text{polylog}(3,(I-c-d)*\exp(2*b*x+2*a)/(I-c+d))/b^2+1/8*I*\text{polylog}(3,(I+c+d)*\exp(2*b*x+2*a)/(I+c-d))/b^2 \end{aligned}$$
Mathematica [A] (warning: unable to verify)

Time = 0.63 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.26

$$\int x \arctan(c + d \coth(a + bx)) dx = \frac{1}{2}x^2 \arctan(c + d \coth(a + bx)) + \frac{d \left(2b^2x^2 \log \left(1 - \frac{(1+(c+d)^2)e^{2(a+bx)}}{1+c^2-d^2+2\sqrt{-d^2}} \right) - 2b^2x^2 \log \left(1 + \frac{(1+(c+d)^2)e^{2(a+bx)}}{-1-c^2+d^2+2\sqrt{-d^2}} \right) + 2bx \text{PolyLog} \left(2, \frac{(1+c^2+2cd+d^2)}{1+c^2-d^2+2\sqrt{-d^2}} \right) \right)}{8b^2}$$

input

Integrate[x*ArcTan[c + d*Coth[a + b*x]],x]

output

$$\begin{aligned} & (x^2*\text{ArcTan}[c + d*\text{Coth}[a + b*x]])/2 + (d*(2*b^2*x^2*\text{Log}[1 - ((1 + (c + d)^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*\text{Sqrt}[-d^2])] - 2*b^2*x^2*\text{Log}[1 + ((1 + (c + d)^2)*E^(2*(a + b*x)))/(-1 - c^2 + d^2 + 2*\text{Sqrt}[-d^2])] + 2*b*x*\text{PolyLog}[2, ((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*\text{Sqrt}[-d^2])] - 2*b*x*\text{PolyLog}[2, -(((1 + (c + d)^2)*E^(2*(a + b*x)))/(-1 - c^2 + d^2 + 2*\text{Sqrt}[-d^2]))] + \text{PolyLog}[3, ((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 - 2*\text{Sqrt}[-d^2])] - \text{PolyLog}[3, ((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*\text{Sqrt}[-d^2])])/(8*b^2*\text{Sqrt}[-d^2]) \end{aligned}$$
Rubi [A] (verified)Time = 1.17 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.28, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {5724, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int x \arctan(d \coth(a + bx) + c) dx \\
& \quad \downarrow \text{5724} \\
& \frac{1}{2}b(1 + i(c + d)) \int \frac{e^{2a+2bx} x^2}{-c - (-c - d + i)e^{2a+2bx} + d + i} dx - \frac{1}{2}b(1 - i(c + \\
& d)) \int \frac{e^{2a+2bx} x^2}{c - (c + d + i)e^{2a+2bx} - d + i} dx + \frac{1}{2}x^2 \arctan(d \coth(a + bx) + c) \\
& \quad \downarrow \text{2620} \\
& \frac{1}{2}b(1 + i(c + d)) \left(\frac{\int x \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) dx}{b(-c-d+i)} - \frac{x^2 \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) - \frac{1}{2}b(1 - \\
& i(c + d)) \left(\frac{\int x \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) dx}{b(c+d+i)} - \frac{x^2 \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right) + \\
& \quad \frac{1}{2}x^2 \arctan(d \coth(a + bx) + c) \\
& \quad \downarrow \text{3011} \\
& \frac{1}{2}b(1 + i(c + \\
& d)) \left(\frac{\int \text{PolyLog} \left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) dx}{2b} - \frac{x \text{PolyLog} \left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b} - \frac{x^2 \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) - \\
& \frac{1}{2}b(1 - i(c + \\
& d)) \left(\frac{\int \text{PolyLog} \left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) dx}{2b} - \frac{x \text{PolyLog} \left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b} - \frac{x^2 \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right) + \\
& \quad \frac{1}{2}x^2 \arctan(d \coth(a + bx) + c) \\
& \quad \downarrow \text{2720}
\end{aligned}$$

$$d)) \left(\frac{\frac{1}{2}b(1 + i(c + \frac{\int e^{-2a-2bx} \text{PolyLog}\left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right) de^{2a+2bx}}{4b^2} - \frac{x \text{PolyLog}\left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b}}{b(-c-d+i)} - \frac{x^2 \log\left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)}}{\right) - \left(\frac{\frac{1}{2}b(1 - i(c + \frac{\int e^{-2a-2bx} \text{PolyLog}\left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right) de^{2a+2bx}}{4b^2} - \frac{x \text{PolyLog}\left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b}}{b(c+d+i)} - \frac{x^2 \log\left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)}}{\right) + \frac{1}{2}x^2 \arctan(d \coth(a + bx) + c)$$

↓ 7143

$$d)) \left(\frac{\frac{1}{2}x^2 \arctan(d \coth(a + bx) + c) + \frac{1}{2}b(1 + i(c + \frac{\text{PolyLog}\left(3, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{4b^2} - \frac{x \text{PolyLog}\left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b}}{b(-c-d+i)} - \frac{x^2 \log\left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i}\right)}{2b(-c-d+i)}}{\right) - \left(\frac{\frac{1}{2}b(1 - i(c + \frac{\text{PolyLog}\left(3, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{4b^2} - \frac{x \text{PolyLog}\left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b}}{b(c+d+i)} - \frac{x^2 \log\left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i}\right)}{2b(c+d+i)}}{\right)$$

input `Int[x*ArcTan[c + d*Coth[a + b*x]],x]`

output `(x^2*ArcTan[c + d*Coth[a + b*x]])/2 + (b*(1 + I*(c + d))*(-1/2*(x^2*Log[1 - ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)])/(b*(I - c - d)) + (-1/2*(x*PolyLog[2, ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)]/b + PolyLog[3, ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)]/(4*b^2)))/(b*(I - c - d)))/2 - (b*(1 - I*(c + d))*(-1/2*(x^2*Log[1 - ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)])/(b*(I + c + d)) + (-1/2*(x*PolyLog[2, ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)]/b + PolyLog[3, ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)]/(4*b^2)))/(b*(I + c + d)))/2`

Definitions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5724

```
Int[ArcTan[(c_) + Coth[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_))^(m_
), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Coth[a + b*x]]/(f*(m
+ 1))), x] + (-Simp[I*b*((I - c - d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(
E^(2*a + 2*b*x)/(I - c + d - (I - c - d)*E^(2*a + 2*b*x))), x], x] + Simp[I
*b*((I + c + d)/(f*(m + 1))) Int[(e + f*x)^(m + 1)*(E^(2*a + 2*b*x)/(I +
c - d - (I + c + d)*E^(2*a + 2*b*x))), x], x]) /; FreeQ[{a, b, c, d, e, f},
x] && IGtQ[m, 0] && NeQ[(c - d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.72 (sec) , antiderivative size = 6495, normalized size of antiderivative = 24.51

method	result	size
risch	Expression too large to display	6495

input `int(x*arctan(c+d*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output `result too large to display`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1051 vs. $2(195) = 390$.

Time = 0.25 (sec) , antiderivative size = 1051, normalized size of antiderivative = 3.97

$$\int x \arctan(c + d \coth(a + bx)) dx = \text{Too large to display}$$

input `integrate(x*arctan(c+d*coth(b*x+a)),x, algorithm="fricas")`

output

```

1/4*(2*b^2*x^2*arctan((d*cosh(b*x + a) + c*sinh(b*x + a))/sinh(b*x + a)) +
2*I*b*x*dilog(sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh
(b*x + a) + sinh(b*x + a))) + 2*I*b*x*dilog(-sqrt((c^2 - d^2 + 2*I*d + 1)/
(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) - 2*I*b*x*dilog(
sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sin
h(b*x + a))) - 2*I*b*x*dilog(-sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d +
d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) + I*a^2*log(2*(c^2 + 2*c*d + d^
2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(c^2 -
d^2 - 2*I*d + 1)*sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) +
I*a^2*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 +
1)*sinh(b*x + a) - 2*(c^2 - d^2 - 2*I*d + 1)*sqrt((c^2 - d^2 + 2*I*d + 1)
/(c^2 - 2*c*d + d^2 + 1))) - I*a^2*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x
+ a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(c^2 - d^2 + 2*I*d + 1)
*sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) - I*a^2*log(2*(c^2
+ 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a
) - 2*(c^2 - d^2 + 2*I*d + 1)*sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d +
d^2 + 1))) + (I*b^2*x^2 - I*a^2)*log(sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2
*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b^2*x^2 - I*a^2
)*log(-sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a
) + sinh(b*x + a)) + 1) + (-I*b^2*x^2 + I*a^2)*log(sqrt((c^2 - d^2 - 2*I*d +
1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1)

```

Sympy [F(-1)]

Timed out.

$$\int x \arctan(c + d \coth(a + bx)) dx = \text{Timed out}$$

input

```
integrate(x*atan(c+d*coth(b*x+a)), x)
```

output

Timed out

Maxima [F]

$$\int x \arctan(c + d \coth(a + bx)) dx = \int x \arctan(d \coth(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*coth(b*x+a)),x, algorithm="maxima")`

output `1/2*x^2*arctan2((c*e^(2*a) + d*e^(2*a))*e^(2*b*x) - c + d, e^(2*b*x + 2*a) - 1) + 2*b*d*integrate(x^2*e^(2*b*x + 2*a)/(c^2 - 2*c*d + d^2 + (c^2*e^(4*a) + 2*c*d*e^(4*a) + d^2*e^(4*a) + e^(4*a))*e^(4*b*x) - 2*(c^2*e^(2*a) - d^2*e^(2*a) + e^(2*a))*e^(2*b*x) + 1), x)`

Giac [F]

$$\int x \arctan(c + d \coth(a + bx)) dx = \int x \arctan(d \coth(bx + a) + c) dx$$

input `integrate(x*arctan(c+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan(d*coth(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + d \coth(a + bx)) dx = \int x \operatorname{atan}(c + d \coth(a + bx)) dx$$

input `int(x*atan(c + d*coth(a + b*x)),x)`

output `int(x*atan(c + d*coth(a + b*x)), x)`

Reduce [F]

$$\int x \arctan(c + d \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) d + c) x dx$$

input `int(x*atan(c+d*coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)*d + c)*x,x)`

3.100 $\int \arctan(c + d \coth(a + bx)) dx$

Optimal result	781
Mathematica [A] (verified)	782
Rubi [A] (verified)	782
Maple [B] (verified)	784
Fricas [B] (verification not implemented)	785
Sympy [F]	786
Maxima [F]	787
Giac [F]	787
Mupad [F(-1)]	787
Reduce [F]	788

Optimal result

Integrand size = 11, antiderivative size = 174

$$\int \arctan(c + d \coth(a + bx)) dx = x \arctan(c + d \coth(a + bx)) + \frac{1}{2}ix \log\left(1 - \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right) - \frac{1}{2}ix \log\left(1 - \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right) + \frac{i \operatorname{PolyLog}\left(2, \frac{(i - c - d)e^{2a+2bx}}{i - c + d}\right)}{4b} - \frac{i \operatorname{PolyLog}\left(2, \frac{(i + c + d)e^{2a+2bx}}{i + c - d}\right)}{4b}$$

output

```
x*arctan(c+d*coth(b*x+a))+1/2*I*x*ln(1-(I-c-d)*exp(2*b*x+2*a)/(I-c+d))-1/2
*I*x*ln(1-(I+c+d)*exp(2*b*x+2*a)/(I+c-d))+1/4*I*polylog(2,(I-c-d)*exp(2*b*
x+2*a)/(I-c+d))/b-1/4*I*polylog(2,(I+c+d)*exp(2*b*x+2*a)/(I+c-d))/b
```

Mathematica [A] (verified)

Time = 1.92 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.65

$$\int \arctan(c + d \coth(a + bx)) dx = x \arctan(c + d \coth(a + bx)) + \frac{4a\sqrt{-d^2} \arctan\left(\frac{1+c^2-d^2-(1+c^2+2cd+d^2)e^{2(a+bx)}}{2d}\right) + 2d(a+bx) \log\left(1 - \frac{(1+(c+d)^2)e^{2(a+bx)}}{1+c^2-d^2+2\sqrt{-d^2}}\right) - 2d(a+bx) \log\left(1 + \frac{(1+(c+d)^2)e^{2(a+bx)}}{1+c^2-d^2+2\sqrt{-d^2}}\right)}{4b\sqrt{-d^2}}$$

input

```
Integrate[ArcTan[c + d*Coth[a + b*x]], x]
```

output

```
x*ArcTan[c + d*Coth[a + b*x]] + (4*a*Sqrt[-d^2]*ArcTan[(1 + c^2 - d^2 - (1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(2*d)] + 2*d*(a + b*x)*Log[1 - ((1 + (c + d)^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2])] - 2*d*(a + b*x)*Log[1 + ((1 + (c + d)^2)*E^(2*(a + b*x)))/(-1 - c^2 + d^2 + 2*Sqrt[-d^2])] + d*PolyLog[2, ((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(1 + c^2 - d^2 + 2*Sqrt[-d^2])] - d*PolyLog[2, -(((1 + c^2 + 2*c*d + d^2)*E^(2*(a + b*x)))/(-1 - c^2 + d^2 + 2*Sqrt[-d^2]))]/(4*b*Sqrt[-d^2])
```

Rubi [A] (verified)Time = 0.69 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.36, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {5716, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \arctan(d \coth(a + bx) + c) dx$$

$$\downarrow 5716$$

$$b(1 + i(c + d)) \int \frac{e^{2a+2bx} x}{-c - (-c - d + i)e^{2a+2bx} + d + i} dx - b(1 - i(c + d)) \int \frac{e^{2a+2bx} x}{c - (c + d + i)e^{2a+2bx} - d + i} dx + x \arctan(d \coth(a + bx) + c)$$

$$\downarrow 2620$$

$$b(1+i(c+d)) \left(\frac{\int \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) dx}{2b(-c-d+i)} - \frac{x \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) - b(1-i(c+d)) \left(\frac{\int \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) dx}{2b(c+d+i)} - \frac{x \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right) + x \arctan(d \coth(a+bx) + c)$$

↓ 2715

$$b(1+i(c+d)) \left(\frac{\int e^{-2a-2bx} \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right) de^{2a+2bx}}{4b^2(-c-d+i)} - \frac{x \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) - b(1-i(c+d)) \left(\frac{\int e^{-2a-2bx} \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right) de^{2a+2bx}}{4b^2(c+d+i)} - \frac{x \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right) + x \arctan(d \coth(a+bx) + c)$$

↓ 2838

$$d) \left(-\frac{\text{PolyLog} \left(2, \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{4b^2(-c-d+i)} - \frac{x \log \left(1 - \frac{(-c-d+i)e^{2a+2bx}}{-c+d+i} \right)}{2b(-c-d+i)} \right) - b(1-i(c+d)) \left(-\frac{\text{PolyLog} \left(2, \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{4b^2(c+d+i)} - \frac{x \log \left(1 - \frac{(c+d+i)e^{2a+2bx}}{c-d+i} \right)}{2b(c+d+i)} \right)$$

input `Int[ArcTan[c + d*Coth[a + b*x]],x]`

output `x*ArcTan[c + d*Coth[a + b*x]] + b*(1 + I*(c + d))*(-1/2*(x*Log[1 - ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d]])/(b*(I - c - d)) - PolyLog[2, ((I - c - d)*E^(2*a + 2*b*x))/(I - c + d)]/(4*b^2*(I - c - d))) - b*(1 - I*(c + d))*(-1/2*(x*Log[1 - ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d]])/(b*(I + c + d)) - PolyLog[2, ((I + c + d)*E^(2*a + 2*b*x))/(I + c - d)]/(4*b^2*(I + c + d)))`

Defintions of rubi rules used

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2715

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2838

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 5716

```
Int[ArcTan[(c_) + Coth[(a_) + (b_)*(x_)]*(d_)], x_Symbol] := Simp[x*Arc
Tan[c + d*Coth[a + b*x]], x] + (-Simp[I*b*(I - c - d) Int[x*(E^(2*a + 2*b
*x))/(I - c + d - (I - c - d)*E^(2*a + 2*b*x))], x], x] + Simp[I*b*(I + c +
d) Int[x*(E^(2*a + 2*b*x))/(I + c - d - (I + c + d)*E^(2*a + 2*b*x))], x],
x] /; FreeQ[{a, b, c, d}, x] && NeQ[(c - d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 351 vs. $2(150) = 300$.

Time = 1.57 (sec) , antiderivative size = 352, normalized size of antiderivative = 2.02

method	result
derivativedivides	$\frac{\arctan(c+d \coth(bx+a))d \ln(-d \coth(bx+a)-d)}{2} - \frac{\arctan(c+d \coth(bx+a))d \ln(-d \coth(bx+a)+d)}{2} + \frac{d^2 \left(\frac{i \ln(-d \coth(bx+a)+d) \ln(-d \coth(bx+a)-d)}{2} \right)}{2}$
default	$\frac{\arctan(c+d \coth(bx+a))d \ln(-d \coth(bx+a)-d)}{2} - \frac{\arctan(c+d \coth(bx+a))d \ln(-d \coth(bx+a)+d)}{2} + \frac{d^2 \left(\frac{i \ln(-d \coth(bx+a)+d) \ln(-d \coth(bx+a)-d)}{2} \right)}{2}$
risch	Expression too large to display

```
input int(arctan(c+d*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

```
output 1/b/d*(1/2*arctan(c+d*coth(b*x+a))*d*ln(-d*coth(b*x+a)-d)-1/2*arctan(c+d*coth(b*x+a))*d*ln(-d*coth(b*x+a)+d)+1/2*d^2*(1/d*(1/2*I*ln(-d*coth(b*x+a)+d))*ln((I+d*coth(b*x+a)+c)/(I+c+d))-1/2*I*ln(-d*coth(b*x+a)+d)*ln((I-d*coth(b*x+a)-c)/(I-c-d))+1/2*I*dilog((I+d*coth(b*x+a)+c)/(I+c+d))-1/2*I*dilog((I-d*coth(b*x+a)-c)/(I-c-d)))-1/d*(1/2*I*ln(-d*coth(b*x+a)-d)*ln((I+d*coth(b*x+a)+c)/(I+c-d))-1/2*I*ln(-d*coth(b*x+a)-d)*ln((I-d*coth(b*x+a)-c)/(I-c+d)))+1/2*I*dilog((I+d*coth(b*x+a)+c)/(I+c-d))-1/2*I*dilog((I-d*coth(b*x+a)-c)/(I-c+d))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 813 vs. 2(128) = 256.

Time = 0.22 (sec) , antiderivative size = 813, normalized size of antiderivative = 4.67

$$\int \arctan(c + d \coth(a + bx)) dx = \text{Too large to display}$$

```
input integrate(arctan(c+d*coth(b*x+a)),x, algorithm="fricas")
```

output

```

1/2*(2*b*x*arctan((d*cosh(b*x + a) + c*sinh(b*x + a))/sinh(b*x + a)) - I*a
*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*s
inh(b*x + a) + 2*(c^2 - d^2 - 2*I*d + 1)*sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2
- 2*c*d + d^2 + 1))) - I*a*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) +
2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) - 2*(c^2 - d^2 - 2*I*d + 1)*sqrt((
c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))) + I*a*log(2*(c^2 + 2*c*d
+ d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2 + 1)*sinh(b*x + a) + 2*(c^
2 - d^2 + 2*I*d + 1)*sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))
) + I*a*log(2*(c^2 + 2*c*d + d^2 + 1)*cosh(b*x + a) + 2*(c^2 + 2*c*d + d^2
+ 1)*sinh(b*x + a) - 2*(c^2 - d^2 + 2*I*d + 1)*sqrt((c^2 - d^2 - 2*I*d +
1)/(c^2 - 2*c*d + d^2 + 1))) + (I*b*x + I*a)*log(sqrt((c^2 - d^2 + 2*I*d +
1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (I*b*x
+ I*a)*log(-sqrt((c^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b
*x + a) + sinh(b*x + a)) + 1) + (-I*b*x - I*a)*log(sqrt((c^2 - d^2 - 2*I*d
+ 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a)) + 1) + (-I*
b*x - I*a)*log(-sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cos
h(b*x + a) + sinh(b*x + a)) + 1) + I*dilog(sqrt((c^2 - d^2 + 2*I*d + 1)/(c
^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x + a))) + I*dilog(-sqrt((c
^2 - d^2 + 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(cosh(b*x + a) + sinh(b*x +
a))) - I*dilog(sqrt((c^2 - d^2 - 2*I*d + 1)/(c^2 - 2*c*d + d^2 + 1))*(...

```

Sympy [F]

$$\int \arctan(c + d \coth(a + bx)) dx = \int \operatorname{atan}(c + d \coth(a + bx)) dx$$

input

```
integrate(atan(c+d*coth(b*x+a)),x)
```

output

```
Integral(atan(c + d*coth(a + b*x)), x)
```

Maxima [F]

$$\int \arctan(c + d \coth(a + bx)) dx = \int \arctan(d \coth(bx + a) + c) dx$$

input `integrate(arctan(c+d*coth(b*x+a)),x, algorithm="maxima")`

output `4*b*d*integrate(x*e^(2*b*x + 2*a)/(c^2 - 2*c*d + d^2 + (c^2*e^(4*a) + 2*c*d*e^(4*a) + d^2*e^(4*a) + e^(4*a))*e^(4*b*x) - 2*(c^2*e^(2*a) - d^2*e^(2*a) + e^(2*a))*e^(2*b*x) + 1), x) + x*arctan2((c*e^(2*a) + d*e^(2*a))*e^(2*b*x) - c + d, e^(2*b*x + 2*a) - 1)`

Giac [F]

$$\int \arctan(c + d \coth(a + bx)) dx = \int \arctan(d \coth(bx + a) + c) dx$$

input `integrate(arctan(c+d*coth(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(d*coth(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(c + d \coth(a + bx)) dx = \int \operatorname{atan}(c + d \coth(a + bx)) dx$$

input `int(atan(c + d*coth(a + b*x)),x)`

output `int(atan(c + d*coth(a + b*x)), x)`

Reduce [F]

$$\int \arctan(c + d \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) d + c) dx$$

input `int(atan(c+d*coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)*d + c),x)`

3.101 $\int \frac{\arctan(c+d \coth(a+bx))}{x} dx$

Optimal result	789
Mathematica [N/A]	789
Rubi [N/A]	790
Maple [N/A]	790
Fricas [N/A]	791
Sympy [F(-1)]	791
Maxima [N/A]	791
Giac [N/A]	792
Mupad [N/A]	792
Reduce [N/A]	792

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + d \coth(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+d*coth(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 6.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \int \frac{\arctan(c + d \coth(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + d*Coth[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + d*Coth[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(d \coth(a + bx) + c)}{x} dx$$

↓ 7299

$$\int \frac{\arctan(d \coth(a + bx) + c)}{x} dx$$

input `Int[ArcTan[c + d*Coth[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + d \coth(bx + a))}{x} dx$$

input `int(arctan(c+d*coth(b*x+a))/x,x)`

output `int(arctan(c+d*coth(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \int \frac{\arctan(d \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*coth(b*x+a))/x,x, algorithm="fricas")`

output `integral(arctan(d*coth(b*x + a) + c)/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+d*coth(b*x+a))/x,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.99 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \int \frac{\arctan(d \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*coth(b*x+a))/x,x, algorithm="maxima")`

output `integrate(arctan(d*coth(b*x + a) + c)/x, x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \int \frac{\arctan(d \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+d*coth(b*x+a))/x,x, algorithm="giac")`

output `integrate(arctan(d*coth(b*x + a) + c)/x, x)`

Mupad [N/A]

Not integrable

Time = 0.64 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + d \coth(a + bx))}{x} dx$$

input `int(atan(c + d*coth(a + b*x))/x,x)`

output `int(atan(c + d*coth(a + b*x))/x, x)`

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\arctan(c + d \coth(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\coth(bx + a) d + c)}{x} dx$$

input `int(atan(c+d*coth(b*x+a))/x,x)`

output `int(atan(coth(a + b*x)*d + c)/x,x)`

3.102 $\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx$

Optimal result	794
Mathematica [A] (verified)	794
Rubi [A] (verified)	795
Maple [C] (warning: unable to verify)	798
Fricas [B] (verification not implemented)	799
Sympy [F(-2)]	800
Maxima [A] (verification not implemented)	800
Giac [F]	801
Mupad [F(-1)]	801
Reduce [F]	801

Optimal result

Integrand size = 19, antiderivative size = 142

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= -\frac{1}{12}ibx^4 + \frac{1}{3}x^3 \arctan(c + (i + c) \coth(a + bx)) + \frac{1}{6}ix^3 \log(1 - ice^{2a+2bx})$$

$$+ \frac{ix^2 \operatorname{PolyLog}(2, ice^{2a+2bx})}{4b} - \frac{ix \operatorname{PolyLog}(3, ice^{2a+2bx})}{4b^2} + \frac{i \operatorname{PolyLog}(4, ice^{2a+2bx})}{8b^3}$$

output

```
-1/12*I*b*x^4+1/3*x^3*arctan(c+(I+c)*coth(b*x+a))+1/6*I*x^3*ln(1-I*c*exp(2
*b*x+2*a))+1/4*I*x^2*polylog(2,I*c*exp(2*b*x+2*a))/b-1/4*I*x*polylog(3,I*c
*exp(2*b*x+2*a))/b^2+1/8*I*polylog(4,I*c*exp(2*b*x+2*a))/b^3
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.94

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= \frac{8b^3x^3 \arctan(c + (i + c) \coth(a + bx)) + 4ib^3x^3 \log\left(1 + \frac{ie^{-2(a+bx)}}{c}\right) - 6ib^2x^2 \operatorname{PolyLog}\left(2, -\frac{ie^{-2(a+bx)}}{c}\right) - \dots}{24b^3}$$

input `Integrate[x^2*ArcTan[c + (I + c)*Coth[a + b*x]],x]`

output $(8*b^3*x^3*ArcTan[c + (I + c)*Coth[a + b*x]] + (4*I)*b^3*x^3*Log[1 + I/(c*E^{(2*(a + b*x))})] - (6*I)*b^2*x^2*PolyLog[2, (-I)/(c*E^{(2*(a + b*x))})] - (6*I)*b*x*PolyLog[3, (-I)/(c*E^{(2*(a + b*x))})] - (3*I)*PolyLog[4, (-I)/(c*E^{(2*(a + b*x))})])/(24*b^3)$

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {5720, 25, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(c + (c + i) \coth(a + bx)) dx \\
 & \quad \downarrow \text{5720} \\
 & \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx)) - \frac{1}{3}b \int -\frac{x^3}{e^{2a+2bx}c+i} dx \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3}b \int \frac{x^3}{e^{2a+2bx}c+i} dx + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx)) \\
 & \quad \downarrow \text{2615} \\
 & \frac{1}{3}b \left(ic \int \frac{e^{2a+2bx}x^3}{e^{2a+2bx}c+i} dx - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx)) \\
 & \quad \downarrow \text{2620} \\
 & \frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \int x^2 \log(1 - ice^{2a+2bx}) dx}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx)) \\
 & \quad \downarrow \text{3011}
 \end{aligned}$$

$$\frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{\int x \text{PolyLog}(2, ice^{2a+2bx}) dx}{b} - \frac{x^2 \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx))$$

↓ 7163

$$\frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{\frac{x \text{PolyLog}(3, ice^{2a+2bx})}{2b} - \frac{\int \text{PolyLog}(3, ice^{2a+2bx}) dx}{b}}{2b} - \frac{x^2 \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx))$$

↓ 2720

$$\frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{\frac{x \text{PolyLog}(3, ice^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \text{PolyLog}(3, ice^{2a+2bx}) de^{2a+2bx}}{b}}{4b^2} - \frac{x^2 \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx))$$

↓ 7143

$$\frac{1}{3}b \left(ic \left(\frac{x^3 \log(1 - ice^{2a+2bx})}{2bc} - \frac{3 \left(\frac{\frac{x \text{PolyLog}(3, ice^{2a+2bx})}{2b} - \frac{\text{PolyLog}(4, ice^{2a+2bx})}{4b^2}}{b} - \frac{x^2 \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right)}{2bc} \right) - \frac{ix^4}{4} \right) + \frac{1}{3}x^3 \arctan(c + (c + i) \coth(a + bx)) +$$

input

```
Int[x^2*ArcTan[c + (I + c)*Coth[a + b*x]],x]
```

output

```
(x^3*ArcTan[c + (I + c)*Coth[a + b*x]])/3 + (b*((-1/4*I)*x^4 + I*c*((x^3*Log[1 - I*c*E^(2*a + 2*b*x)])/(2*b*c) - (3*(-1/2*(x^2*PolyLog[2, I*c*E^(2*a + 2*b*x)]))/b + ((x*PolyLog[3, I*c*E^(2*a + 2*b*x)])/(2*b) - PolyLog[4, I*c*E^(2*a + 2*b*x)]/(4*b^2))/b)/(2*b*c)))/3
```

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2615 $\text{Int}[(((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.}))))^{\text{n}_.}], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m} + 1} / (\text{a} * \text{d} * (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b} / \text{a} \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m}} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}} / (\text{a} + \text{b} * (\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}})), \text{x}], \text{x}] /; \text{FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2620 $\text{Int}[(((\text{F}_.)^{\text{g}_.}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.}))), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m}} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}} / \text{a})], \text{x}] - \text{Simp}[\text{d} * (\text{m} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m} - 1} * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}} / \text{a})], \text{x}], \text{x}] /; \text{FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2720 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{With}\{\text{v} = \text{FunctionOfExponential}[\text{u}, \text{x}]\}, \text{Simp}[\text{v} / \text{D}[\text{v}, \text{x}] \quad \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[\text{u}, \text{x}] / \text{x}, \text{x}], \text{x}, \text{v}], \text{x}]] /; \text{FunctionOfExponentialQ}[\text{u}, \text{x}] \&\& \text{!MatchQ}[\text{u}, (\text{w}_.) * ((\text{a}_.) * (\text{v}_.)^{\text{n}_.})^{\text{m}_.}] /; \text{FreeQ}\{\text{a}, \text{m}, \text{n}\}, \text{x}\} \&\& \text{IntegerQ}[\text{m} * \text{n}] \&\& \text{!MatchQ}[\text{u}, \text{E}^{\text{c}_.} * ((\text{a}_.) + (\text{b}_.) * \text{x})) * (\text{F}_.)[\text{v}_.] /; \text{FreeQ}\{\text{a}, \text{b}, \text{c}\}, \text{x}\} \&\& \text{InverseFunctionQ}[\text{F}[\text{x}]]]$
- rule 3011 $\text{Int}[\text{Log}[1 + (\text{e}_.) * ((\text{F}_.)^{\text{c}_.} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.}] * ((\text{f}_.) + (\text{g}_.) * (\text{x}_.)^{\text{m}_.}), \text{x_Symbol}] \rightarrow \text{Simp}[-(\text{f} + \text{g} * \text{x})^{\text{m}} * (\text{PolyLog}[2, (-\text{e}) * (\text{F}^{\text{c}}(\text{a} + \text{b} * \text{x}))^{\text{n}}] / (\text{b} * \text{c} * \text{n} * \text{Log}[\text{F}]))], \text{x}] + \text{Simp}[\text{g} * (\text{m} / (\text{b} * \text{c} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{f} + \text{g} * \text{x})^{\text{m} - 1} * \text{PolyLog}[2, (-\text{e}) * (\text{F}^{\text{c}}(\text{a} + \text{b} * \text{x}))^{\text{n}}], \text{x}], \text{x}] /; \text{FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{GtQ}[\text{m}, 0]$
- rule 5720 $\text{Int}[\text{ArcTan}[(\text{c}_.) + \text{Coth}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)] * (\text{d}_.)] * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{m}_.})], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{e} + \text{f} * \text{x})^{\text{m} + 1} * (\text{ArcTan}[\text{c} + \text{d} * \text{Coth}[\text{a} + \text{b} * \text{x}]] / (\text{f} * (\text{m} + 1))), \text{x}] - \text{Simp}[\text{b} / (\text{f} * (\text{m} + 1)) \quad \text{Int}[(\text{e} + \text{f} * \text{x})^{\text{m} + 1} / (\text{c} - \text{d} - \text{c} * \text{E}^{2 * \text{a} + 2 * \text{b} * \text{x}})], \text{x}], \text{x}] /; \text{FreeQ}\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{EqQ}[(\text{c} - \text{d})^2, -1]$

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol]
:> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x]
&& GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.12 (sec) , antiderivative size = 1405, normalized size of antiderivative = 9.89

method	result	size
risch	Expression too large to display	1405

input

```
int(x^2*arctan(c+(1+c)*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/2*I/b^3*a^2*dilog(1+I*exp(b*x+a)*(-I*c)^(1/2))+1/12*Pi*(csgn(I*(2*exp(2*
b*x+2*a)*c+2*I))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^2-csg
n(I*(2*exp(2*b*x+2*a)*c+2*I))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2
*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+
2*a)*c))*csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1)
)^2+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(2*I*exp(2*b*x+
2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))-cs
gn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^3+csgn(I*(2*exp(2*b*x+2*
a)*c+2*I)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(2*exp(2
*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))*csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(2*
b*x+2*a)-1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))*csgn((2
*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))+csgn(I*(2*I*exp(2*b*x+2*a)+2*ex
p(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b
*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*I*ex
p(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((2*I*exp(2*b*x+2
*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(2*I*exp(2*b*x+2*a)+2
*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*
x+2*a)*c)/(exp(2*b*x+2*a)-1))+csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)
/(exp(2*b*x+2*a)-1))^3-csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2
*b*x+2*a)-1))^2+csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^3-csg...

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs. $2(105) = 210$.

Time = 0.10 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.06

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= -i b^4 x^4 + 2i b^3 x^3 \log\left(-\frac{(c+i)e^{(2bx+2a)}}{ce^{(2bx+2a)}+i}\right) + 6i b^2 x^2 \operatorname{Li}_2\left(\frac{1}{2} \sqrt{4i} ce^{(bx+a)}\right) + 6i b^2 x^2 \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{4i} ce^{(bx+a)}\right) + i a^4$$

input

```
integrate(x^2*arctan(c+(I+c)*coth(b*x+a)),x, algorithm="fricas")
```


output

```
1/12*(-I*b^4*x^4 + 2*I*b^3*x^3*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) + I)) + 6*I*b^2*x^2*dilog(1/2*sqrt(4*I*c)*e^(b*x + a)) + 6*I*b^2*x^2*dilog(-1/2*sqrt(4*I*c)*e^(b*x + a)) + I*a^4 - 2*I*a^3*log(1/2*(2*c*e^(b*x + a) + I*sqrt(4*I*c))/c) - 2*I*a^3*log(1/2*(2*c*e^(b*x + a) - I*sqrt(4*I*c))/c) - 12*I*b*x*polylog(3, 1/2*sqrt(4*I*c)*e^(b*x + a)) - 12*I*b*x*polylog(3, -1/2*sqrt(4*I*c)*e^(b*x + a)) - 2*(-I*b^3*x^3 - I*a^3)*log(1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - 2*(-I*b^3*x^3 - I*a^3)*log(-1/2*sqrt(4*I*c)*e^(b*x + a) + 1) + 12*I*polylog(4, 1/2*sqrt(4*I*c)*e^(b*x + a)) + 12*I*polylog(4, -1/2*sqrt(4*I*c)*e^(b*x + a)))/b^3
```

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input

```
integrate(x**2*atan(c+(I+c)*coth(b*x+a)),x)
```

output

```
Exception raised: CoercionFailed >> Cannot convert _t0**2 - exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]
```

Maxima [A] (verification not implemented)

Time = 1.72 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.91

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx = \frac{1}{3} x^3 \arctan((c + i) \coth(bx + a) + c) + \frac{4}{9} \left(\frac{3x^4}{4ic - 4} - \frac{4b^3x^3 \log(-ice^{(2bx+2a)}) + 1}{-2b^4(-ic + 1)} + 6b^2x^2 \text{Li}_2(ice^{(2bx+2a)}) - 6bx \text{Li}_3(ice^{(2bx+2a)}) + 3 \text{Li}_4(ice^{(2bx+2a)}) \right)$$

input

```
integrate(x^2*arctan(c+(I+c)*coth(b*x+a)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctan((c + I)*coth(b*x + a) + c) + 4/9*(3*x^4/(4*I*c - 4) - (4*b^
3*x^3*log(-I*c*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog(I*c*e^(2*b*x + 2*a))
- 6*b*x*polylog(3, I*c*e^(2*b*x + 2*a)) + 3*polylog(4, I*c*e^(2*b*x + 2*a
)))/(b^4*(2*I*c - 2))*b*(c + I)
```

Giac [F]

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx = \int x^2 \arctan((c + i) \coth(bx + a) + c) dx$$

input

```
integrate(x^2*arctan(c+(I+c)*coth(b*x+a)),x, algorithm="giac")
```

output

```
integrate(x^2*arctan((c + I)*coth(b*x + a) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx = \int x^2 \operatorname{atan}(c + \coth(a + bx) (c + 1i)) dx$$

input

```
int(x^2*atan(c + coth(a + b*x)*(c + 1i)),x)
```

output

```
int(x^2*atan(c + coth(a + b*x)*(c + 1i)), x)
```

Reduce [F]

$$\int x^2 \arctan(c + (i + c) \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) c + \coth(bx + a) i + c) x^2 dx$$

input

```
int(x^2*atan(c+(I+c)*coth(b*x+a)),x)
```

output `int(atan(coth(a + b*x)*c + coth(a + b*x)*i + c)*x**2,x)`

3.103 $\int x \arctan(c + (i + c) \coth(a + bx)) dx$

Optimal result	803
Mathematica [A] (verified)	804
Rubi [A] (verified)	804
Maple [C] (warning: unable to verify)	807
Fricas [B] (verification not implemented)	808
Sympy [F(-2)]	808
Maxima [A] (verification not implemented)	809
Giac [F]	809
Mupad [F(-1)]	809
Reduce [F]	810

Optimal result

Integrand size = 17, antiderivative size = 113

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx = -\frac{1}{6}ibx^3 + \frac{1}{2}x^2 \arctan(c + (i + c) \coth(a + bx)) + \frac{1}{4}ix^2 \log(1 - ice^{2a+2bx}) + \frac{ix \operatorname{PolyLog}(2, ice^{2a+2bx})}{4b} - \frac{i \operatorname{PolyLog}(3, ice^{2a+2bx})}{8b^2}$$

output

```
-1/6*I*b*x^3+1/2*x^2*arctan(c+(I+c)*coth(b*x+a))+1/4*I*x^2*ln(1-I*c*exp(2*b*x+2*a))+1/4*I*x*polylog(2,I*c*exp(2*b*x+2*a))/b-1/8*I*polylog(3,I*c*exp(2*b*x+2*a))/b^2
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.91

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= \frac{2b^2 x^2 \left(2 \arctan(c + (i + c) \coth(a + bx)) + i \log \left(1 + \frac{ie^{-2(a+bx)}}{c} \right) \right) - 2ibx \operatorname{PolyLog} \left(2, -\frac{ie^{-2(a+bx)}}{c} \right) - i \operatorname{PolyLog} \left(3, -\frac{ie^{-2(a+bx)}}{c} \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c + (I + c)*Coth[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTan[c + (I + c)*Coth[a + b*x]] + I*Log[1 + I/(c*E^(2*(a + b*x))])) - (2*I)*b*x*PolyLog[2, (-I)/(c*E^(2*(a + b*x)))] - I*PolyLog[3, (-I)/(c*E^(2*(a + b*x)))]/(8*b^2)
```

Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.18, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {5720, 25, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(c + (c + i) \coth(a + bx)) dx$$

$$\downarrow \text{5720}$$

$$\frac{1}{2} x^2 \arctan(c + (c + i) \coth(a + bx)) - \frac{1}{2} b \int -\frac{x^2}{e^{2a+2bx} c + i} dx$$

$$\downarrow \text{25}$$

$$\frac{1}{2} b \int \frac{x^2}{e^{2a+2bx} c + i} dx + \frac{1}{2} x^2 \arctan(c + (c + i) \coth(a + bx))$$

$$\downarrow \text{2615}$$

$$\frac{1}{2} b \left(ic \int \frac{e^{2a+2bx} x^2}{e^{2a+2bx} c + i} dx - \frac{ix^3}{3} \right) + \frac{1}{2} x^2 \arctan(c + (c + i) \coth(a + bx))$$

$$\begin{array}{c} \downarrow 2620 \\ \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int x \log(1 - ice^{2a+2bx}) dx}{bc} \right) - \frac{ix^3}{3} \right) + \frac{1}{2}x^2 \arctan(c + (c + i) \coth(a + bx)) \end{array}$$

$$\begin{array}{c} \downarrow 3011 \\ \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int \text{PolyLog}(2, ice^{2a+2bx}) dx}{2b} - \frac{x \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right) - \frac{ix^3}{3} \right) + \\ \frac{1}{2}x^2 \arctan(c + (c + i) \coth(a + bx)) \end{array}$$

$$\begin{array}{c} \downarrow 2720 \\ \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int e^{-2a-2bx} \text{PolyLog}(2, ice^{2a+2bx}) de^{2a+2bx}}{4b^2} - \frac{x \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right) - \frac{ix^3}{3} \right) + \\ \frac{1}{2}x^2 \arctan(c + (c + i) \coth(a + bx)) \end{array}$$

$$\begin{array}{c} \downarrow 7143 \\ \frac{1}{2}x^2 \arctan(c + (c + i) \coth(a + bx)) + \\ \frac{1}{2}b \left(ic \left(\frac{x^2 \log(1 - ice^{2a+2bx})}{2bc} - \frac{\text{PolyLog}(3, ice^{2a+2bx})}{4b^2} - \frac{x \text{PolyLog}(2, ice^{2a+2bx})}{2b} \right) - \frac{ix^3}{3} \right) \end{array}$$

input `Int[x*ArcTan[c + (I + c)*Coth[a + b*x]],x]`

output `(x^2*ArcTan[c + (I + c)*Coth[a + b*x]])/2 + (b*((-1/3*I)*x^3 + I*c*((x^2*Log[1 - I*c*E^(2*a + 2*b*x)])/(2*b*c) - (-1/2*(x*PolyLog[2, I*c*E^(2*a + 2*b*x)])/b + PolyLog[3, I*c*E^(2*a + 2*b*x)]/(4*b^2))/(b*c)))/2`

Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 2615 $\text{Int}[(((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.}))))^{\text{n}_.}], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m} + 1} / (\text{a} * \text{d} * (\text{m} + 1)), \text{x}] - \text{Simp}[\text{b} / \text{a} \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m}} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}} / (\text{a} + \text{b} * (\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}})), \text{x}], \text{x}] /; \text{FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2620 $\text{Int}[(((\text{F}_.)^{\text{g}_.}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.} * ((\text{c}_.) + (\text{d}_.) * (\text{x}_.)^{\text{m}_.}) / ((\text{a}_.) + (\text{b}_.) * ((\text{F}_.)^{\text{g}_.}) * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{n}_.}))), \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c} + \text{d} * \text{x})^{\text{m}} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}} / \text{a})], \text{x}] - \text{Simp}[\text{d} * (\text{m} / (\text{b} * \text{f} * \text{g} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{c} + \text{d} * \text{x})^{\text{m} - 1} * \text{Log}[1 + \text{b} * ((\text{F}^{\text{g}}(\text{e} + \text{f} * \text{x}))^{\text{n}} / \text{a})], \text{x}], \text{x}] /; \text{FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0]$
- rule 2720 $\text{Int}[\text{u}_, \text{x_Symbol}] \rightarrow \text{With}\{\text{v} = \text{FunctionOfExponential}[\text{u}, \text{x}]\}, \text{Simp}[\text{v} / \text{D}[\text{v}, \text{x}] \quad \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[\text{u}, \text{x}] / \text{x}, \text{x}], \text{x}, \text{v}], \text{x}] /; \text{FunctionOfExponentialQ}[\text{u}, \text{x}] \&\& \text{!MatchQ}[\text{u}, (\text{w}_.) * ((\text{a}_.) * (\text{v}_.)^{\text{n}_.})^{\text{m}_.}] /; \text{FreeQ}\{\text{a}, \text{m}, \text{n}\}, \text{x}\} \&\& \text{IntegerQ}[\text{m} * \text{n}] \&\& \text{!MatchQ}[\text{u}, \text{E}^{\text{c}_.} * ((\text{a}_.) + (\text{b}_.) * \text{x})) * (\text{F}_.)[\text{v}_.] /; \text{FreeQ}\{\text{a}, \text{b}, \text{c}\}, \text{x}\} \&\& \text{InverseFunctionQ}[\text{F}[\text{x}]]]$
- rule 3011 $\text{Int}[\text{Log}[1 + (\text{e}_.) * ((\text{F}_.)^{\text{c}_.} * ((\text{a}_.) + (\text{b}_.) * (\text{x}_.)^{\text{n}_.})))^{\text{n}_.}] * ((\text{f}_.) + (\text{g}_.) * (\text{x}_.)^{\text{m}_.}), \text{x_Symbol}] \rightarrow \text{Simp}[-(\text{f} + \text{g} * \text{x})^{\text{m}} * (\text{PolyLog}[2, (-\text{e}) * (\text{F}^{\text{c}}(\text{a} + \text{b} * \text{x}))^{\text{n}}] / (\text{b} * \text{c} * \text{n} * \text{Log}[\text{F}]))], \text{x}] + \text{Simp}[\text{g} * (\text{m} / (\text{b} * \text{c} * \text{n} * \text{Log}[\text{F}])) \quad \text{Int}[(\text{f} + \text{g} * \text{x})^{\text{m} - 1} * \text{PolyLog}[2, (-\text{e}) * (\text{F}^{\text{c}}(\text{a} + \text{b} * \text{x}))^{\text{n}}], \text{x}], \text{x}] /; \text{FreeQ}\{\text{F}, \text{a}, \text{b}, \text{c}, \text{e}, \text{f}, \text{g}, \text{n}\}, \text{x}\} \&\& \text{GtQ}[\text{m}, 0]$
- rule 5720 $\text{Int}[\text{ArcTan}[(\text{c}_.) + \text{Coth}[(\text{a}_.) + (\text{b}_.) * (\text{x}_.)] * (\text{d}_.)] * ((\text{e}_.) + (\text{f}_.) * (\text{x}_.)^{\text{m}_.})], \text{x_Symbol}] \rightarrow \text{Simp}[(\text{e} + \text{f} * \text{x})^{\text{m} + 1} * (\text{ArcTan}[\text{c} + \text{d} * \text{Coth}[\text{a} + \text{b} * \text{x}]] / (\text{f} * (\text{m} + 1))), \text{x}] - \text{Simp}[\text{b} / (\text{f} * (\text{m} + 1)) \quad \text{Int}[(\text{e} + \text{f} * \text{x})^{\text{m} + 1} / (\text{c} - \text{d} - \text{c} * \text{E}^{2 * \text{a} + 2 * \text{b} * \text{x}})], \text{x}], \text{x}] /; \text{FreeQ}\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}, \text{f}\}, \text{x}\} \&\& \text{IGtQ}[\text{m}, 0] \&\& \text{EqQ}[(\text{c} - \text{d})^2, -1]$

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.58 (sec) , antiderivative size = 1369, normalized size of antiderivative = 12.12

method	result	size
risch	Expression too large to display	1369

input

```
int(x*arctan(c+(I+c)*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/4*I/b^2*polylog(2,I*c*exp(2*b*x+2*a))*a+1/8*Pi*(csgn(I*(2*exp(2*b*x+2*a)*c+2*I))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I))*csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^3+csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))*csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^2-csgn(I*(2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))*csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))+csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn((2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn((2*exp(2*b*x+2*a)*c+2*I)/(exp(2*b*x+2*a)-1))^3-csgn((2*exp...
```


Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(83) = 166$.

Time = 0.10 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.19

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= \frac{-2i b^3 x^3 + 3i b^2 x^2 \log\left(-\frac{(c+i)e^{(2bx+2a)}}{ce^{(2bx+2a)}+i}\right) - 2i a^3 + 6i bx \operatorname{Li}_2\left(\frac{1}{2} \sqrt{4i} ce^{(bx+a)}\right) + 6i bx \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{4i} ce^{(bx+a)}\right)}{b^2}$$

input `integrate(x*arctan(c+(I+c)*coth(b*x+a)),x, algorithm="fricas")`

output `1/12*(-2*I*b^3*x^3 + 3*I*b^2*x^2*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) + I)) - 2*I*a^3 + 6*I*b*x*dilog(1/2*sqrt(4*I*c)*e^(b*x + a)) + 6*I*b*x*dilog(-1/2*sqrt(4*I*c)*e^(b*x + a)) + 3*I*a^2*log(1/2*(2*c*e^(b*x + a) + I*sqrt(4*I*c))/c) + 3*I*a^2*log(1/2*(2*c*e^(b*x + a) - I*sqrt(4*I*c))/c) - 3*(-I*b^2*x^2 + I*a^2)*log(1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - 3*(-I*b^2*x^2 + I*a^2)*log(-1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - 6*I*polylog(3, 1/2*sqrt(4*I*c)*e^(b*x + a)) - 6*I*polylog(3, -1/2*sqrt(4*I*c)*e^(b*x + a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x*atan(c+(I+c)*coth(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 - exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.37 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.94

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= \left(\frac{2x^3}{3ic - 3} - \frac{2b^2x^2 \log(-ice^{(2bx+2a)} + 1) + 2bx \operatorname{Li}_2(ice^{(2bx+2a)}) - \operatorname{Li}_3(ice^{(2bx+2a)})}{-2b^3(-ic + 1)} \right) b(c + i)$$

$$+ \frac{1}{2} x^2 \arctan((c + i) \coth(bx + a) + c)$$

input `integrate(x*arctan(c+(I+c)*coth(b*x+a)),x, algorithm="maxima")`

output `(2*x^3/(3*I*c - 3) - (2*b^2*x^2*log(-I*c*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog(I*c*e^(2*b*x + 2*a)) - polylog(3, I*c*e^(2*b*x + 2*a)))/(b^3*(2*I*c - 2)))*b*(c + I) + 1/2*x^2*arctan((c + I)*coth(b*x + a) + c)`

Giac [F]

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx = \int x \arctan((c + i) \coth(bx + a) + c) dx$$

input `integrate(x*arctan(c+(I+c)*coth(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan((c + I)*coth(b*x + a) + c), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx = \int x \operatorname{atan}(c + \coth(a + bx) (c + 1i)) dx$$

input `int(x*atan(c + coth(a + b*x)*(c + 1i)),x)`

output `int(x*atan(c + coth(a + b*x)*(c + 1i)), x)`

Reduce [F]

$$\int x \arctan(c + (i + c) \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) c + \coth(bx + a) i + c) x dx$$

input `int(x*atan(c+(I+c)*coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)*c + coth(a + b*x)*i + c)*x,x)`

3.104 $\int \arctan(c + (i + c) \coth(a + bx)) dx$

Optimal result	811
Mathematica [A] (verified)	811
Rubi [A] (verified)	812
Maple [B] (verified)	814
Fricas [B] (verification not implemented)	815
Sympy [F(-2)]	815
Maxima [A] (verification not implemented)	816
Giac [F]	816
Mupad [F(-1)]	816
Reduce [F]	817

Optimal result

Integrand size = 15, antiderivative size = 79

$$\int \arctan(c + (i + c) \coth(a + bx)) dx = -\frac{1}{2}ibx^2 + x \arctan(c + (i + c) \coth(a + bx)) + \frac{1}{2}ix \log(1 - ice^{2a+2bx}) + \frac{i \operatorname{PolyLog}(2, ice^{2a+2bx})}{4b}$$

output

$$-1/2*I*b*x^2+x*\arctan(c+(I+c)*\coth(b*x+a))+1/2*I*x*\ln(1-I*c*\exp(2*b*x+2*a))+1/4*I*\operatorname{polylog}(2,I*c*\exp(2*b*x+2*a))/b$$

Mathematica [A] (verified)

Time = 1.29 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.90

$$\int \arctan(c + (i + c) \coth(a + bx)) dx = x \arctan(c + (i + c) \coth(a + bx)) + \frac{i \left(2bx \log \left(1 + \frac{ie^{-2(a+bx)}}{c} \right) - \operatorname{PolyLog} \left(2, -\frac{ie^{-2(a+bx)}}{c} \right) \right)}{4b}$$

input `Integrate[ArcTan[c + (I + c)*Coth[a + b*x]], x]`

output `x*ArcTan[c + (I + c)*Coth[a + b*x]] + ((I/4)*(2*b*x*Log[1 + I/(c*E^(2*(a + b*x))]) - PolyLog[2, (-I)/(c*E^(2*(a + b*x))]]))/b`

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5712, 25, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(c + (c + i) \coth(a + bx)) dx \\
 & \quad \downarrow \text{5712} \\
 & x \arctan(c + (c + i) \coth(a + bx)) - b \int -\frac{x}{e^{2a+2bx}c+i} dx \\
 & \quad \downarrow \text{25} \\
 & b \int \frac{x}{e^{2a+2bx}c+i} dx + x \arctan(c + (c + i) \coth(a + bx)) \\
 & \quad \downarrow \text{2615} \\
 & b \left(ic \int \frac{e^{2a+2bx}x}{e^{2a+2bx}c+i} dx - \frac{ix^2}{2} \right) + x \arctan(c + (c + i) \coth(a + bx)) \\
 & \quad \downarrow \text{2620} \\
 & b \left(ic \left(\frac{x \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int \log(1 - ice^{2a+2bx}) dx}{2bc} \right) - \frac{ix^2}{2} \right) + x \arctan(c + (c + i) \coth(a + bx)) \\
 & \quad \downarrow \text{2715} \\
 & b \left(ic \left(\frac{x \log(1 - ice^{2a+2bx})}{2bc} - \frac{\int e^{-2a-2bx} \log(1 - ice^{2a+2bx}) de^{2a+2bx}}{4b^2c} \right) - \frac{ix^2}{2} \right) + x \arctan(c + (c + i) \coth(a + bx))
 \end{aligned}$$

$$\begin{array}{c} \downarrow 2838 \\ b \left(ic \left(\frac{x \arctan(c + (c + i) \coth(a + bx)) + \text{PolyLog}(2, ice^{2a+2bx})}{4b^2c} + \frac{x \log(1 - ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right) \end{array}$$

input `Int[ArcTan[c + (I + c)*Coth[a + b*x]],x]`

output `x*ArcTan[c + (I + c)*Coth[a + b*x]] + b*((-1/2*I)*x^2 + I*c*((x*Log[1 - I*c*E^(2*a + 2*b*x)])/(2*b*c) + PolyLog[2, I*c*E^(2*a + 2*b*x)]/(4*b^2*c)))`

Definitions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2615 `Int[(((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5712

```
Int[ArcTan[(c_.) + Coth[(a_.) + (b_.)*(x_.)]*(d_.)], x_Symbol] := Simp[x*Arc
Tan[c + d*Coth[a + b*x]], x] - Simp[b Int[x/(c - d - c*E^(2*a + 2*b*x)),
x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, -1]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 544 vs. $2(65) = 130$.

Time = 0.62 (sec) , antiderivative size = 545, normalized size of antiderivative = 6.90

method	result
derivativedivides	$-\frac{\arctan(c+(i+c)\coth(bx+a))\ln(i+c+(i+c)\coth(bx+a))}{2i+2c} + \frac{2i\arctan(c+(i+c)\coth(bx+a))\ln(i+c+(i+c)\coth(bx+a))c}{2i+2c} + \frac{\arctan(c+(i+c)\coth(bx+a))\ln(i+c+(i+c)\coth(bx+a))}{2i+2c}$
default	$-\frac{\arctan(c+(i+c)\coth(bx+a))\ln(i+c+(i+c)\coth(bx+a))}{2i+2c} + \frac{2i\arctan(c+(i+c)\coth(bx+a))\ln(i+c+(i+c)\coth(bx+a))c}{2i+2c} + \frac{\arctan(c+(i+c)\coth(bx+a))\ln(i+c+(i+c)\coth(bx+a))}{2i+2c}$
risch	Expression too large to display

input

```
int(arctan(c+(I+c)*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```
1/b/(I+c)*(-arctan(c+(I+c)*coth(b*x+a))/(2*I+2*c)*ln(I+c+(I+c)*coth(b*x+a)
)+2*I*arctan(c+(I+c)*coth(b*x+a))/(2*I+2*c)*ln(I+c+(I+c)*coth(b*x+a))*c+ar
ctan(c+(I+c)*coth(b*x+a))/(2*I+2*c)*ln(I+c+(I+c)*coth(b*x+a))*c^2+arctan(c
+(I+c)*coth(b*x+a))/(2*I+2*c)*ln(c-(I+c)*coth(b*x+a)+I)-2*I*arctan(c+(I+c)
*coth(b*x+a))/(2*I+2*c)*ln(c-(I+c)*coth(b*x+a)+I)*c-arctan(c+(I+c)*coth(b*
x+a))/(2*I+2*c)*ln(c-(I+c)*coth(b*x+a)+I)*c^2-(I+c)^(1/2)/(I+c)*(-1/2*I*(
(ln(I+c+(I+c)*coth(b*x+a))-ln(-1/2*I*(I+c+(I+c)*coth(b*x+a))))*ln(-1/2*I*(
I-c-(I+c)*coth(b*x+a)))-dilog(-1/2*I*(I+c+(I+c)*coth(b*x+a))))+1/4*I*ln(I+
c+(I+c)*coth(b*x+a))^2-1/2/(I+c)*(-1/2*I*(dilog(-1/2*(I-c-(I+c)*coth(b*x+
a))/c)+ln(c-(I+c)*coth(b*x+a)+I)*ln(-1/2*(I-c-(I+c)*coth(b*x+a))/c))+1/2*I
*(dilog((-I-c-(I+c)*coth(b*x+a))/(-2*I-2*c))+ln(c-(I+c)*coth(b*x+a)+I)*ln(
(-I-c-(I+c)*coth(b*x+a))/(-2*I-2*c))))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(58) = 116$.

Time = 0.12 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.37

$$\int \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= \frac{-i b^2 x^2 + i b x \log\left(-\frac{(c+i)e^{(2bx+2a)}}{ce^{(2bx+2a)+i}}\right) + i a^2 + (i b x + i a) \log\left(\frac{1}{2} \sqrt{4i c} e^{(bx+a)} + 1\right) + (i b x + i a) \log\left(-\frac{1}{2} \sqrt{4i c} e^{(bx+a)} + 1\right)}{b}$$

input `integrate(arctan(c+(I+c)*coth(b*x+a)),x, algorithm="fricas")`

output `1/2*(-I*b^2*x^2 + I*b*x*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) + I)) + I*a^2 + (I*b*x + I*a)*log(1/2*sqrt(4*I*c)*e^(b*x + a) + 1) + (I*b*x + I*a)*log(-1/2*sqrt(4*I*c)*e^(b*x + a) + 1) - I*a*log(1/2*(2*c*e^(b*x + a) + I*sqrt(4*I*c))/c) - I*a*log(1/2*(2*c*e^(b*x + a) - I*sqrt(4*I*c))/c) + I*dilog(1/2*sqrt(4*I*c)*e^(b*x + a)) + I*dilog(-1/2*sqrt(4*I*c)*e^(b*x + a)))/b`

Sympy [F(-2)]

Exception generated.

$$\int \arctan(c + (i + c) \coth(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(atan(c+(I+c)*coth(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 - exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[b, _t0, exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.34 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01

$$\int \arctan(c + (i + c) \coth(a + bx)) dx$$

$$= 2b(c + i) \left(\frac{2x^2}{2ic - 2} - \frac{2bx \log(-ice^{(2bx+2a)} + 1) + \text{Li}_2(ice^{(2bx+2a)})}{-2b^2(-ic + 1)} \right)$$

$$+ x \arctan((c + i) \coth(bx + a) + c)$$

input `integrate(arctan(c+(I+c)*coth(b*x+a)),x, algorithm="maxima")`output `2*b*(c + I)*(2*x^2/(2*I*c - 2) - (2*b*x*log(-I*c*e^(2*b*x + 2*a) + 1) + di
log(I*c*e^(2*b*x + 2*a)))/(b^2*(2*I*c - 2))) + x*arctan((c + I)*coth(b*x +
a) + c)`**Giac [F]**

$$\int \arctan(c + (i + c) \coth(a + bx)) dx = \int \arctan((c + i) \coth(bx + a) + c) dx$$

input `integrate(arctan(c+(I+c)*coth(b*x+a)),x, algorithm="giac")`output `integrate(arctan((c + I)*coth(b*x + a) + c), x)`**Mupad [F(-1)]**

Timed out.

$$\int \arctan(c + (i + c) \coth(a + bx)) dx = \int \text{atan}(c + \coth(a + bx) (c + li)) dx$$

input `int(atan(c + coth(a + b*x)*(c + li)),x)`

output `int(atan(c + coth(a + b*x)*(c + 1i)), x)`

Reduce [F]

$$\int \arctan(c + (i + c) \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) c + \coth(bx + a) i + c) dx$$

input `int(atan(c+(I+c)*coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)*c + coth(a + b*x)*i + c),x)`

3.105 $\int \frac{\arctan(c+(i+c) \coth(a+bx))}{x} dx$

Optimal result	818
Mathematica [N/A]	818
Rubi [N/A]	819
Maple [N/A]	819
Fricas [N/A]	820
Sympy [F(-1)]	820
Maxima [N/A]	820
Giac [N/A]	821
Mupad [N/A]	821
Reduce [N/A]	822

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c + (i + c) \coth(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c+(I+c)*coth(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 3.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx$$

input `Integrate[ArcTan[c + (I + c)*Coth[a + b*x]]/x,x]`

output `Integrate[ArcTan[c + (I + c)*Coth[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c + (c + i) \coth(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c + (c + i) \coth(a + bx))}{x} dx$$

input `Int[ArcTan[c + (I + c)*Coth[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\arctan(c + (i + c) \coth(bx + a))}{x} dx$$

input `int(arctan(c+(I+c)*coth(b*x+a))/x,x)`

output `int(arctan(c+(I+c)*coth(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.95

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \int \frac{\arctan((c + i) \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+(I+c)*coth(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c + I)*e^(2*b*x + 2*a)/(c*e^(2*b*x + 2*a) + I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c+(I+c)*coth(b*x+a))/x,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 77, normalized size of antiderivative = 4.05

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \int \frac{\arctan((c + i) \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c+(I+c)*coth(b*x+a))/x,x, algorithm="maxima")`

output

```
I*b*x + 1/2*pi*log(x) - 1/4*(4*pi - 4*I*a - 2*arctan(c) - I*log(c^2 + 1))*
log(x) - 1/2*integrate(arctan(c*e^(2*b*x + 2*a))/x, x) - 1/4*I*integrate(1
og(c^2*e^(4*b*x + 4*a) + 1)/x, x)
```

Giac [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \int \frac{\arctan((c + i) \coth(bx + a) + c)}{x} dx$$

input

```
integrate(arctan(c+(I+c)*coth(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctan((c + I)*coth(b*x + a) + c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.97 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + \coth(a + bx) (c + 1i))}{x} dx$$

input

```
int(atan(c + coth(a + b*x)*(c + 1i))/x,x)
```

output

```
int(atan(c + coth(a + b*x)*(c + 1i))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.32

$$\int \frac{\arctan(c + (i + c) \coth(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\coth(bx + a) c + \coth(bx + a) i + c)}{x} dx$$

input `int(atan(c+(I+c)*coth(b*x+a))/x,x)`output `int(atan(coth(a + b*x)*c + coth(a + b*x)*i + c)/x,x)`

3.106 $\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx$

Optimal result	823
Mathematica [A] (verified)	824
Rubi [A] (verified)	824
Maple [C] (warning: unable to verify)	827
Fricas [B] (verification not implemented)	828
Sympy [F(-2)]	829
Maxima [A] (verification not implemented)	829
Giac [F]	830
Mupad [F(-1)]	830
Reduce [F]	830

Optimal result

Integrand size = 22, antiderivative size = 145

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx = \frac{1}{12} ibx^4 + \frac{1}{3} x^3 \arctan(c - (i - c) \coth(a + bx)) - \frac{1}{6} ix^3 \log(1 + ice^{2a+2bx}) - \frac{ix^2 \operatorname{PolyLog}(2, -ice^{2a+2bx})}{4b} + \frac{ix \operatorname{PolyLog}(3, -ice^{2a+2bx})}{4b^2} - \frac{i \operatorname{PolyLog}(4, -ice^{2a+2bx})}{8b^3}$$

output

```
1/12*I*b*x^4+1/3*x^3*arctan(c-(I-c)*coth(b*x+a))-1/6*I*x^3*ln(1+I*c*exp(2*b*x+2*a))-1/4*I*x^2*polylog(2,-I*c*exp(2*b*x+2*a))/b+1/4*I*x*polylog(3,-I*c*exp(2*b*x+2*a))/b^2-1/8*I*polylog(4,-I*c*exp(2*b*x+2*a))/b^3
```


Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.92

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx$$

$$= \frac{8b^3 x^3 \arctan(c + (-i + c) \coth(a + bx)) - 4ib^3 x^3 \log\left(1 - \frac{ie^{-2(a+bx)}}{c}\right) + 6ib^2 x^2 \text{PolyLog}\left(2, \frac{ie^{-2(a+bx)}}{c}\right) + \dots}{24b^3}$$

input

```
Integrate[x^2*ArcTan[c - (I - c)*Coth[a + b*x]],x]
```

output

```
(8*b^3*x^3*ArcTan[c + (-I + c)*Coth[a + b*x]] - (4*I)*b^3*x^3*Log[1 - I/(c*E^(2*(a + b*x)))] + (6*I)*b^2*x^2*PolyLog[2, I/(c*E^(2*(a + b*x)))] + (6*I)*b*x*PolyLog[3, I/(c*E^(2*(a + b*x)))] + (3*I)*PolyLog[4, I/(c*E^(2*(a + b*x)))])/(24*b^3)
```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.17, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$, Rules used = {5720, 2615, 2620, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(c - (-c + i) \coth(a + bx)) dx$$

$$\downarrow \text{5720}$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \int \frac{x^3}{i - ce^{2a+2bx}} dx$$

$$\downarrow \text{2615}$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \left(-ic \int \frac{e^{2a+2bx} x^3}{i - ce^{2a+2bx}} dx - \frac{ix^4}{4} \right)$$

$$\downarrow \text{2620}$$

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \left(-ic \left(\frac{3 \int x^2 \log(ie^{2a+2bx}c + 1) dx}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^4}{4} \right)$$

↓ 3011

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\int x \operatorname{PolyLog}(2, -ice^{2a+2bx}) dx}{b} - \frac{x^2 \operatorname{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^4}{4} \right)$$

↓ 7163

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\frac{x \operatorname{PolyLog}(3, -ice^{2a+2bx})}{2b} - \frac{\int \operatorname{PolyLog}(3, -ice^{2a+2bx}) dx}{b}}{2b} - \frac{x^2 \operatorname{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) \right)$$

↓ 2720

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\frac{x \operatorname{PolyLog}(3, -ice^{2a+2bx})}{2b} - \frac{\int e^{-2a-2bx} \operatorname{PolyLog}(3, -ice^{2a+2bx}) de^{2a+2bx}}{b}}{4b^2} - \frac{x^2 \operatorname{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) \right)$$

↓ 7143

$$\frac{1}{3}x^3 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{3}b \left(-ic \left(\frac{3 \left(\frac{\frac{x \operatorname{PolyLog}(3, -ice^{2a+2bx})}{2b} - \frac{\operatorname{PolyLog}(4, -ice^{2a+2bx})}{4b^2}}{b} - \frac{x^2 \operatorname{PolyLog}(2, -ice^{2a+2bx})}{2b} \right)}{2bc} - \frac{x^3 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^4}{4} \right)$$

input

```
Int[x^2*ArcTan[c - (I - c)*Coth[a + b*x]], x]
```

output

```
(x^3*ArcTan[c - (I - c)*Coth[a + b*x]]/3 - (b*((-1/4*I)*x^4 - I*c*(-1/2*(x^3*Log[1 + I*c*E^(2*a + 2*b*x)]))/(b*c) + (3*(-1/2*(x^2*PolyLog[2, (-I)*c*E^(2*a + 2*b*x)])/b + ((x*PolyLog[3, (-I)*c*E^(2*a + 2*b*x)])/(2*b) - PolyLog[4, (-I)*c*E^(2*a + 2*b*x)]/(4*b^2))/b)/(2*b*c)))/3
```

Defintions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

rule 5720

```
Int[ArcTan[(c_.) + Coth[(a_.) + (b_.)*(x_)]*(d_.)]*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Coth[a + b*x]]/(f*(m + 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c - d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.28 (sec) , antiderivative size = 1418, normalized size of antiderivative = 9.78

method	result	size
risch	Expression too large to display	1418

input

```
int(x^2*arctan(c-(1-c)*coth(b*x+a)),x,method=_RETURNVERBOSE)
```

output

```

1/6*I*x^3*ln(-2*exp(2*b*x+2*a)*c+2*I)-1/12*Pi*(-csgn(I*(2*exp(2*b*x+2*a)*c-2*I))*csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^2-csgn(I*(2*exp(2*b*x+2*a)*c-2*I))*csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c))*csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))*csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^2-csgn(I*(2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))*csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))-csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))-csgn((-2*I*exp(2*b*x+2*a)+2*exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn((2*exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^3-c...

```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs. $2(105) = 210$.

Time = 0.12 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.02

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx$$

$$= \frac{ib^4x^4 + 2ib^3x^3 \log\left(-\frac{ce^{(2bx+2a)} - i}{c-i}e^{(-2bx-2a)}\right) - 6ib^2x^2 \operatorname{Li}_2\left(\frac{1}{2}\sqrt{-4i}ce^{(bx+a)}\right) - 6ib^2x^2 \operatorname{Li}_2\left(-\frac{1}{2}\sqrt{-4i}ce^{(bx-a)}\right)}{b^4}$$

input

```
integrate(x^2*arctan(c-(I-c)*coth(b*x+a)),x, algorithm="fricas")
```

output

```
1/12*(I*b^4*x^4 + 2*I*b^3*x^3*log(-(c*e^(2*b*x + 2*a) - I)*e^(-2*b*x - 2*a)
)/(c - I)) - 6*I*b^2*x^2*dilog(1/2*sqrt(-4*I*c)*e^(b*x + a)) - 6*I*b^2*x^2
*dilog(-1/2*sqrt(-4*I*c)*e^(b*x + a)) - I*a^4 + 2*I*a^3*log(1/2*(2*c*e^(b*
x + a) + I*sqrt(-4*I*c))/c) + 2*I*a^3*log(1/2*(2*c*e^(b*x + a) - I*sqrt(-4
*I*c))/c) + 12*I*b*x*polylog(3, 1/2*sqrt(-4*I*c)*e^(b*x + a)) + 12*I*b*x*p
olylog(3, -1/2*sqrt(-4*I*c)*e^(b*x + a)) - 2*(I*b^3*x^3 + I*a^3)*log(1/2*s
qrt(-4*I*c)*e^(b*x + a) + 1) - 2*(I*b^3*x^3 + I*a^3)*log(-1/2*sqrt(-4*I*c)
*e^(b*x + a) + 1) - 12*I*polylog(4, 1/2*sqrt(-4*I*c)*e^(b*x + a)) - 12*I*p
olylog(4, -1/2*sqrt(-4*I*c)*e^(b*x + a))/b^3
```

Sympy [F(-2)]

Exception generated.

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input

```
integrate(x**2*atan(c-(I-c)*coth(b*x+a)),x)
```

output

```
Exception raised: CoercionFailed >> Cannot convert _t0**2 - exp(2*a) of ty
pe <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]
```

Maxima [A] (verification not implemented)

Time = 1.51 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.89

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx = \frac{1}{3} x^3 \arctan((c - i) \coth(bx + a) + c) - \frac{4}{9} \left(\frac{3x^4}{4ic + 4} - \frac{4b^3x^3 \log(i ce^{(2bx+2a)} + 1) + 6b^2x^2 \text{Li}_2(-i ce^{(2bx+2a)}) - 6bx \text{Li}_3(-i ce^{(2bx+2a)}) + 3 \text{Li}_4(-i ce^{(2bx+2a)})}{-2b^4(-ic - 1)} \right)$$

input

```
integrate(x^2*arctan(c-(I-c)*coth(b*x+a)),x, algorithm="maxima")
```

output

```
1/3*x^3*arctan((c - I)*coth(b*x + a) + c) - 4/9*(3*x^4/(4*I*c + 4) - (4*b^
3*x^3*log(I*c*e^(2*b*x + 2*a) + 1) + 6*b^2*x^2*dilog(-I*c*e^(2*b*x + 2*a))
- 6*b*x*polylog(3, -I*c*e^(2*b*x + 2*a)) + 3*polylog(4, -I*c*e^(2*b*x + 2
*a)))/(b^4*(2*I*c + 2))*b*(c - I)
```

Giac [F]

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx = \int x^2 \arctan((c - i) \coth(bx + a) + c) dx$$

input

```
integrate(x^2*arctan(c-(I-c)*coth(b*x+a)),x, algorithm="giac")
```

output

```
integrate(x^2*arctan((c - I)*coth(b*x + a) + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx = \int x^2 \operatorname{atan}(c + \coth(a + bx) (c - i)) dx$$

input

```
int(x^2*atan(c + coth(a + b*x)*(c - 1i)),x)
```

output

```
int(x^2*atan(c + coth(a + b*x)*(c - 1i)), x)
```

Reduce [F]

$$\int x^2 \arctan(c - (i - c) \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) c - \coth(bx + a) i + c) x^2 dx$$

input

```
int(x^2*atan(c-(I-c)*coth(b*x+a)),x)
```

output `int(atan(coth(a + b*x)*c - coth(a + b*x)*i + c)*x**2,x)`

3.107 $\int x \arctan(c - (i - c) \coth(a + bx)) dx$

Optimal result	832
Mathematica [A] (verified)	833
Rubi [A] (verified)	833
Maple [C] (warning: unable to verify)	836
Fricas [B] (verification not implemented)	837
Sympy [F(-2)]	837
Maxima [A] (verification not implemented)	838
Giac [F]	838
Mupad [F(-1)]	838
Reduce [F]	839

Optimal result

Integrand size = 20, antiderivative size = 116

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx = \frac{1}{6}ibx^3 + \frac{1}{2}x^2 \arctan(c - (i - c) \coth(a + bx)) - \frac{1}{4}ix^2 \log(1 + ice^{2a+2bx}) - \frac{ix \operatorname{PolyLog}(2, -ice^{2a+2bx})}{4b} + \frac{i \operatorname{PolyLog}(3, -ice^{2a+2bx})}{8b^2}$$

output

```
1/6*I*b*x^3+1/2*x^2*arctan(c-(I-c)*coth(b*x+a))-1/4*I*x^2*ln(1+I*c*exp(2*b*x+2*a))-1/4*I*x*polylog(2,-I*c*exp(2*b*x+2*a))/b+1/8*I*polylog(3,-I*c*exp(2*b*x+2*a))/b^2
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.89

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx$$

$$= \frac{2b^2x^2 \left(2 \arctan(c + (-i + c) \coth(a + bx)) - i \log \left(1 - \frac{ie^{-2(a+bx)}}{c} \right) \right) + 2ibx \operatorname{PolyLog} \left(2, \frac{ie^{-2(a+bx)}}{c} \right) + i \operatorname{PolyLog} \left(3, \frac{ie^{-2(a+bx)}}{c} \right)}{8b^2}$$

input

```
Integrate[x*ArcTan[c - (I - c)*Coth[a + b*x]],x]
```

output

```
(2*b^2*x^2*(2*ArcTan[c + (-I + c)*Coth[a + b*x]] - I*Log[1 - I/(c*E^(2*(a + b*x))])) + (2*I)*b*x*PolyLog[2, I/(c*E^(2*(a + b*x)))] + I*PolyLog[3, I/(c*E^(2*(a + b*x)))]/(8*b^2)
```

Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.16, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5720, 2615, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(c - (-c + i) \coth(a + bx)) dx$$

$$\downarrow \text{5720}$$

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{2}b \int \frac{x^2}{i - ce^{2a+2bx}} dx$$

$$\downarrow \text{2615}$$

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{2}b \left(-ic \int \frac{e^{2a+2bx} x^2}{i - ce^{2a+2bx}} dx - \frac{ix^3}{3} \right)$$

$$\downarrow \text{2620}$$

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{2}b \left(-ic \left(\frac{\int x \log(ie^{2a+2bx}c + 1) dx}{bc} - \frac{x^2 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

↓ 3011

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{2}b \left(-ic \left(\frac{\int \text{PolyLog}(2, -ice^{2a+2bx}) dx}{2b} - \frac{x \text{PolyLog}(2, -ice^{2a+2bx})}{2b} - \frac{x^2 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

↓ 2720

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{2}b \left(-ic \left(\frac{\int e^{-2a-2bx} \text{PolyLog}(2, -ice^{2a+2bx}) de^{2a+2bx}}{4b^2} - \frac{x \text{PolyLog}(2, -ice^{2a+2bx})}{2b} - \frac{x^2 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

↓ 7143

$$\frac{1}{2}x^2 \arctan(c - (-c + i) \coth(a + bx)) - \frac{1}{2}b \left(-ic \left(\frac{\text{PolyLog}(3, -ice^{2a+2bx})}{4b^2} - \frac{x \text{PolyLog}(2, -ice^{2a+2bx})}{2b} - \frac{x^2 \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^3}{3} \right)$$

input `Int[x*ArcTan[c - (I - c)*Coth[a + b*x]],x]`

output `(x^2*ArcTan[c - (I - c)*Coth[a + b*x]])/2 - (b*((-1/3*I)*x^3 - I*c*(-1/2*(x^2*Log[1 + I*c*E^(2*a + 2*b*x)]))/(b*c) + (-1/2*(x*PolyLog[2, (-I)*c*E^(2*a + 2*b*x)])/b + PolyLog[3, (-I)*c*E^(2*a + 2*b*x)]/(4*b^2))/(b*c)))/2`

Defintions of rubi rules used

rule 2615

```
Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2620

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 3011

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 5720

```
Int[ArcTan[(c_) + Coth[(a_) + (b_)*(x_)]*(d_)]*((e_) + (f_)*(x_))^(m_
), x_Symbol] := Simp[(e + f*x)^(m + 1)*(ArcTan[c + d*Coth[a + b*x]]/(f*(m
+ 1))), x] - Simp[b/(f*(m + 1)) Int[(e + f*x)^(m + 1)/(c - d - c*E^(2*a +
2*b*x)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && EqQ[(c -
d)^2, -1]
```

rule 7143

```
Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.80 (sec) , antiderivative size = 1382, normalized size of antiderivative = 11.91

method	result	size
risch	Expression too large to display	1382

input `int(x*arctan(c-(I-c)*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/4*I*x^2*\ln(1+I*c*\exp(2*b*x+2*a))-1/8*Pi*(-csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)) \\
 & *csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^2-csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)) \\
 & *csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c))*csgn \\
 & (I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c))*csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^3+csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))-csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))*csgn((2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^2-csgn(I*(2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))*csgn((2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))-csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2*csgn(I/(exp(2*b*x+2*a)-1))+csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2+csgn(I*(-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))*csgn((-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))-csgn((-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^3-csgn((-2*I*\exp(2*b*x+2*a)+2*\exp(2*b*x+2*a)*c)/(exp(2*b*x+2*a)-1))^2-csgn((2*\exp(2*b*x+2*a)*c-2*I)/(exp(2*b*x+2*a)-1))^3-csgn...
 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 247 vs. $2(83) = 166$.

Time = 0.13 (sec) , antiderivative size = 247, normalized size of antiderivative = 2.13

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx$$

$$= \frac{2i b^3 x^3 + 3i b^2 x^2 \log\left(-\frac{ce^{(2bx+2a)} - i}{c-i} e^{(-2bx-2a)}\right) + 2i a^3 - 6i bx \operatorname{Li}_2\left(\frac{1}{2} \sqrt{-4i} ce^{(bx+a)}\right) - 6i bx \operatorname{Li}_2\left(-\frac{1}{2} \sqrt{-4i} ce^{(bx+a)}\right)}{b^2}$$

input `integrate(x*arctan(c-(I-c)*coth(b*x+a)),x, algorithm="fricas")`

output `1/12*(2*I*b^3*x^3 + 3*I*b^2*x^2*log(-(c*e^(2*b*x + 2*a) - I)*e^(-2*b*x - 2*a)/(c - I)) + 2*I*a^3 - 6*I*b*x*dilog(1/2*sqrt(-4*I*c)*e^(b*x + a)) - 6*I*b*x*dilog(-1/2*sqrt(-4*I*c)*e^(b*x + a)) - 3*I*a^2*log(1/2*(2*c*e^(b*x + a) + I*sqrt(-4*I*c))/c) - 3*I*a^2*log(1/2*(2*c*e^(b*x + a) - I*sqrt(-4*I*c))/c) - 3*(I*b^2*x^2 - I*a^2)*log(1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) - 3*(I*b^2*x^2 - I*a^2)*log(-1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) + 6*I*polylog(3, 1/2*sqrt(-4*I*c)*e^(b*x + a)) + 6*I*polylog(3, -1/2*sqrt(-4*I*c)*e^(b*x + a)))/b^2`

Sympy [F(-2)]

Exception generated.

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(x*atan(c-(I-c)*coth(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 - exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[x,b,_t0,exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.63 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.92

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx =$$

$$-\left(\frac{2x^3}{3ic + 3} - \frac{2b^2x^2 \log(ice^{(2bx+2a)} + 1) + 2bx \operatorname{Li}_2(-ice^{(2bx+2a)}) - \operatorname{Li}_3(-ice^{(2bx+2a)})}{-2b^3(-ic - 1)} \right) b(c - i)$$

$$+ \frac{1}{2} x^2 \arctan((c - i) \coth(bx + a) + c)$$

input `integrate(x*arctan(c-(I-c)*coth(b*x+a)),x, algorithm="maxima")`output `-(2*x^3/(3*I*c + 3) - (2*b^2*x^2*log(I*c*e^(2*b*x + 2*a) + 1) + 2*b*x*dilog(-I*c*e^(2*b*x + 2*a)) - polylog(3, -I*c*e^(2*b*x + 2*a)))/(b^3*(2*I*c + 2)))*b*(c - I) + 1/2*x^2*arctan((c - I)*coth(b*x + a) + c)`**Giac [F]**

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx = \int x \arctan((c - i) \coth(bx + a) + c) dx$$

input `integrate(x*arctan(c-(I-c)*coth(b*x+a)),x, algorithm="giac")`output `integrate(x*arctan((c - I)*coth(b*x + a) + c), x)`**Mupad [F(-1)]**

Timed out.

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx = \int x \operatorname{atan}(c + \coth(a + bx) (c - i)) dx$$

input `int(x*atan(c + coth(a + b*x)*(c - 1i)),x)`

output `int(x*atan(c + coth(a + b*x)*(c - 1i)), x)`

Reduce [F]

$$\int x \arctan(c - (i - c) \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a) c - \coth(bx + a) i + c) x dx$$

input `int(x*atan(c-(I-c)*coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)*c - coth(a + b*x)*i + c)*x,x)`

3.108 $\int \arctan(c - (i - c) \coth(a + bx)) dx$

Optimal result	840
Mathematica [A] (verified)	840
Rubi [A] (verified)	841
Maple [B] (verified)	843
Fricas [B] (verification not implemented)	844
Sympy [F(-2)]	844
Maxima [A] (verification not implemented)	845
Giac [F]	845
Mupad [F(-1)]	845
Reduce [F]	846

Optimal result

Integrand size = 18, antiderivative size = 82

$$\int \arctan(c - (i - c) \coth(a + bx)) dx = \frac{1}{2} ibx^2 + x \arctan(c - (i - c) \coth(a + bx)) - \frac{1}{2} ix \log(1 + ice^{2a+2bx}) - \frac{i \operatorname{PolyLog}(2, -ice^{2a+2bx})}{4b}$$

output

```
1/2*I*b*x^2+x*arctan(c-(I-c)*coth(b*x+a))-1/2*I*x*ln(1+I*c*exp(2*b*x+2*a))
-1/4*I*polylog(2,-I*c*exp(2*b*x+2*a))/b
```

Mathematica [A] (verified)

Time = 1.27 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.87

$$\int \arctan(c - (i - c) \coth(a + bx)) dx = x \arctan(c + (-i + c) \coth(a + bx)) - \frac{i \left(2bx \log \left(1 - \frac{ie^{-2(a+bx)}}{c} \right) - \operatorname{PolyLog} \left(2, \frac{ie^{-2(a+bx)}}{c} \right) \right)}{4b}$$

input `Integrate[ArcTan[c - (I - c)*Coth[a + b*x]], x]`

output `x*ArcTan[c + (-I + c)*Coth[a + b*x]] - ((I/4)*(2*b*x*Log[1 - I/(c*E^(2*(a + b*x)))] - PolyLog[2, I/(c*E^(2*(a + b*x))]))/b`

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.17, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {5712, 2615, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(c - (-c + i) \coth(a + bx)) dx \\
 & \quad \downarrow \text{5712} \\
 & x \arctan(c - (-c + i) \coth(a + bx)) - b \int \frac{x}{i - ce^{2a+2bx}} dx \\
 & \quad \downarrow \text{2615} \\
 & x \arctan(c - (-c + i) \coth(a + bx)) - b \left(-ic \int \frac{e^{2a+2bx} x}{i - ce^{2a+2bx}} dx - \frac{ix^2}{2} \right) \\
 & \quad \downarrow \text{2620} \\
 & b \left(-ic \left(\frac{\int \log(i e^{2a+2bx} c + 1) dx}{2bc} - \frac{x \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right) \\
 & \quad \downarrow \text{2715} \\
 & b \left(-ic \left(\frac{\int e^{-2a-2bx} \log(i e^{2a+2bx} c + 1) de^{2a+2bx}}{4b^2c} - \frac{x \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right) \\
 & \quad \downarrow \text{2838}
 \end{aligned}$$

$$b \left(-ic \left(-\frac{x \arctan(c - (-c + i) \coth(a + bx)) - \text{PolyLog}(2, -ice^{2a+2bx})}{4b^2c} - \frac{x \log(1 + ice^{2a+2bx})}{2bc} \right) - \frac{ix^2}{2} \right)$$

input `Int[ArcTan[c - (I - c)*Coth[a + b*x]],x]`

output `x*ArcTan[c - (I - c)*Coth[a + b*x]] - b*((-1/2*I)*x^2 - I*c*(-1/2*(x*Log[1 + I*c*E^(2*a + 2*b*x)])/(b*c) - PolyLog[2, (-I)*c*E^(2*a + 2*b*x)]/(4*b^2*c)))`

Defintions of rubi rules used

rule 2615 `Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Simp[b/a Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2620 `Int((((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_.) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_.))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_.) + (e_.)*(x_.)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5712 `Int[ArcTan[(c_.) + Coth[(a_.) + (b_.)*(x_.)]*(d_.)], x_Symbol] := Simp[x*ArcTan[c + d*Coth[a + b*x]], x] - Simp[b Int[x/(c - d - c*E^(2*a + 2*b*x)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[(c - d)^2, -1]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 515 vs. $2(68) = 136$.

Time = 0.72 (sec) , antiderivative size = 516, normalized size of antiderivative = 6.29

method	result
derivativedivides	$\frac{\arctan(c+\coth(bx+a)(-i+c))\ln(-i+\coth(bx+a)(-i+c)+c)}{2i-2c} + \frac{2i\arctan(c+\coth(bx+a)(-i+c))\ln(-i+\coth(bx+a)(-i+c)+c)}{2i-2c} - \arctan(c+\coth(bx+a)(-i+c))\ln(-i+\coth(bx+a)(-i+c)+c)$
default	$\frac{\arctan(c+\coth(bx+a)(-i+c))\ln(-i+\coth(bx+a)(-i+c)+c)}{2i-2c} + \frac{2i\arctan(c+\coth(bx+a)(-i+c))\ln(-i+\coth(bx+a)(-i+c)+c)}{2i-2c} - \arctan(c+\coth(bx+a)(-i+c))\ln(-i+\coth(bx+a)(-i+c)+c)$
risch	Expression too large to display

input `int(arctan(c-(I-c)*coth(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
1/b/(-I+c)*(arctan(c+coth(b*x+a)*(-I+c))/(2*I-2*c)*ln(-I+coth(b*x+a)*(-I+c)+c)+2*I*arctan(c+coth(b*x+a)*(-I+c))/(2*I-2*c)*ln(-I+coth(b*x+a)*(-I+c)+c)*c-arctan(c+coth(b*x+a)*(-I+c))/(2*I-2*c)*ln(-I+coth(b*x+a)*(-I+c)+c)*c^2-arctan(c+coth(b*x+a)*(-I+c))/(2*I-2*c)*ln(coth(b*x+a)*(-I+c)-c+I)-2*I*arctan(c+coth(b*x+a)*(-I+c))/(2*I-2*c)*ln(coth(b*x+a)*(-I+c)-c+I)*c+arctan(c+coth(b*x+a)*(-I+c))/(2*I-2*c)*ln(coth(b*x+a)*(-I+c)-c+I)*c^2+(I-c)^2*(1/2/(I-c)*(-1/4*I*ln(-I+coth(b*x+a)*(-I+c)+c)^2+1/2*I*(dilog(-1/2*I*(coth(b*x+a)*(-I+c)+c+I))+ln(-I+coth(b*x+a)*(-I+c)+c)*ln(-1/2*I*(coth(b*x+a)*(-I+c)+c+I))))-1/2/(I-c)*(-1/2*I*(dilog((-I+coth(b*x+a)*(-I+c)+c)/(-2*I+2*c))+ln(coth(b*x+a)*(-I+c)-c+I)*ln((-I+coth(b*x+a)*(-I+c)+c)/(-2*I+2*c)))+1/2*I*(dilog(1/2*(coth(b*x+a)*(-I+c)+c+I)/c)+ln(coth(b*x+a)*(-I+c)-c+I)*ln(1/2*(coth(b*x+a)*(-I+c)+c+I)/c))))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(58) = 116$.

Time = 0.11 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.28

$$\int \arctan(c - (i - c) \coth(a + bx)) dx$$

$$= \frac{i b^2 x^2 + i b x \log\left(-\frac{(c e^{2bx+2a} - i) e^{-2bx-2a}}{c-i}\right) - i a^2 + (-i b x - i a) \log\left(\frac{1}{2} \sqrt{-4i c} e^{(bx+a)} + 1\right) + (-i b x - i a) \log\left(\frac{1}{2} \sqrt{-4i c} e^{(bx+a)} - 1\right)}{b}$$

input `integrate(arctan(c-(I-c)*coth(b*x+a)),x, algorithm="fricas")`

output `1/2*(I*b^2*x^2 + I*b*x*log(-(c*e^(2*b*x + 2*a) - I)*e^(-2*b*x - 2*a)/(c - I)) - I*a^2 + (-I*b*x - I*a)*log(1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) + (-I*b*x - I*a)*log(-1/2*sqrt(-4*I*c)*e^(b*x + a) + 1) + I*a*log(1/2*(2*c*e^(b*x + a) + I*sqrt(-4*I*c))/c) + I*a*log(1/2*(2*c*e^(b*x + a) - I*sqrt(-4*I*c))/c) - I*dilog(1/2*sqrt(-4*I*c)*e^(b*x + a)) - I*dilog(-1/2*sqrt(-4*I*c)*e^(b*x + a)))/b`

Sympy [F(-2)]

Exception generated.

$$\int \arctan(c - (i - c) \coth(a + bx)) dx = \text{Exception raised: CoercionFailed}$$

input `integrate(atan(c-(I-c)*coth(b*x+a)),x)`

output `Exception raised: CoercionFailed >> Cannot convert _t0**2 - exp(2*a) of type <class 'sympy.core.add.Add'> to QQ_I[b, _t0, exp(a)]`

Maxima [A] (verification not implemented)

Time = 1.70 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.98

$$\int \arctan(c - (i - c) \coth(a + bx)) dx$$

$$= -2b(c - i) \left(\frac{2x^2}{2ic + 2} - \frac{2bx \log(ice^{(2bx+2a)} + 1) + \text{Li}_2(-ice^{(2bx+2a)})}{-2b^2(-ic - 1)} \right)$$

$$+ x \arctan((c - i) \coth(bx + a) + c)$$

input `integrate(arctan(c-(I-c)*coth(b*x+a)),x, algorithm="maxima")`output `-2*b*(c - I)*(2*x^2/(2*I*c + 2) - (2*b*x*log(I*c*e^(2*b*x + 2*a) + 1) + di
log(-I*c*e^(2*b*x + 2*a)))/(b^2*(2*I*c + 2))) + x*arctan((c - I)*coth(b*x
+ a) + c)`**Giac [F]**

$$\int \arctan(c - (i - c) \coth(a + bx)) dx = \int \arctan((c - i) \coth(bx + a) + c) dx$$

input `integrate(arctan(c-(I-c)*coth(b*x+a)),x, algorithm="giac")`output `integrate(arctan((c - I)*coth(b*x + a) + c), x)`**Mupad [F(-1)]**

Timed out.

$$\int \arctan(c - (i - c) \coth(a + bx)) dx = \int \text{atan}(c + \coth(a + bx) (c - i)) dx$$

input `int(atan(c + coth(a + b*x)*(c - 1i)),x)`

output `int(atan(c + coth(a + b*x)*(c - 1i)), x)`

Reduce [F]

$$\int \arctan(c - (i - c) \coth(a + bx)) dx = \int \operatorname{atan}(\coth(bx + a)c - \coth(bx + a)i + c) dx$$

input `int(atan(c-(I-c)*coth(b*x+a)),x)`

output `int(atan(coth(a + b*x)*c - coth(a + b*x)*i + c),x)`

3.109 $\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx$

Optimal result	847
Mathematica [N/A]	847
Rubi [N/A]	848
Maple [N/A]	848
Fricas [N/A]	849
Sympy [F(-1)]	849
Maxima [N/A]	849
Giac [N/A]	850
Mupad [N/A]	850
Reduce [N/A]	851

Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \text{Int}\left(\frac{\arctan(c - (i - c) \coth(a + bx))}{x}, x\right)$$

output `Defer(Int)(arctan(c-(I-c)*coth(b*x+a))/x,x)`

Mathematica [N/A]

Not integrable

Time = 3.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx$$

input `Integrate[ArcTan[c - (I - c)*Coth[a + b*x]]/x,x]`

output `Integrate[ArcTan[c - (I - c)*Coth[a + b*x]]/x, x]`

Rubi [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c - (-c + i) \coth(a + bx))}{x} dx$$

↓ 7299

$$\int \frac{\arctan(c - (-c + i) \coth(a + bx))}{x} dx$$

input `Int[ArcTan[c - (I - c)*Coth[a + b*x]]/x,x]`

output `$Aborted`

Maple [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\arctan(c - (i - c) \coth(bx + a))}{x} dx$$

input `int(arctan(c-(I-c)*coth(b*x+a))/x,x)`

output `int(arctan(c-(I-c)*coth(b*x+a))/x,x)`

Fricas [N/A]

Not integrable

Time = 0.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.68

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \int \frac{\arctan((c - i) \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c-(I-c)*coth(b*x+a))/x,x, algorithm="fricas")`

output `integral(1/2*I*log(-(c*e^(2*b*x + 2*a) - I)*e^(-2*b*x - 2*a)/(c - I))/x, x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \text{Timed out}$$

input `integrate(atan(c-(I-c)*coth(b*x+a))/x,x)`

output `Timed out`

Maxima [N/A]

Not integrable

Time = 0.54 (sec) , antiderivative size = 72, normalized size of antiderivative = 3.27

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \int \frac{\arctan((c - i) \coth(bx + a) + c)}{x} dx$$

input `integrate(arctan(c-(I-c)*coth(b*x+a))/x,x, algorithm="maxima")`

output

```
-I*b*x - 1/4*(2*pi + 4*I*a - 2*arctan(c) + I*log(c^2 + 1))*log(x) - 1/2*integrate(arctan(c*e^(2*b*x + 2*a))/x, x) + 1/4*I*integrate(log(c^2*e^(4*b*x + 4*a) + 1)/x, x)
```

Giac [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \int \frac{\arctan((c - i) \coth(bx + a) + c)}{x} dx$$

input

```
integrate(arctan(c-(I-c)*coth(b*x+a))/x,x, algorithm="giac")
```

output

```
integrate(arctan((c - I)*coth(b*x + a) + c)/x, x)
```

Mupad [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \int \frac{\operatorname{atan}(c + \coth(a + bx) (c - i))}{x} dx$$

input

```
int(atan(c + coth(a + b*x)*(c - 1i))/x,x)
```

output

```
int(atan(c + coth(a + b*x)*(c - 1i))/x, x)
```

Reduce [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{\arctan(c - (i - c) \coth(a + bx))}{x} dx = \int \frac{\operatorname{atan}(\coth(bx + a) c - \coth(bx + a) i + c)}{x} dx$$

input `int(atan(c-(I-c)*coth(b*x+a))/x,x)`output `int(atan(coth(a + b*x)*c - coth(a + b*x)*i + c)/x,x)`

3.110 $\int \arctan(e^x) dx$

Optimal result	852
Mathematica [A] (verified)	852
Rubi [A] (verified)	853
Maple [B] (verified)	854
Fricas [B] (verification not implemented)	854
Sympy [F]	855
Maxima [B] (verification not implemented)	855
Giac [F]	856
Mupad [B] (verification not implemented)	856
Reduce [F]	856

Optimal result

Integrand size = 4, antiderivative size = 31

$$\int \arctan(e^x) dx = \frac{1}{2}i \operatorname{PolyLog}(2, -ie^x) - \frac{1}{2}i \operatorname{PolyLog}(2, ie^x)$$

output `1/2*I*polylog(2,-I*exp(x))-1/2*I*polylog(2,I*exp(x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.90

$$\int \arctan(e^x) dx = x \arctan(e^x) - \frac{1}{2}i(x(\log(1 - ie^x) - \log(1 + ie^x)) - \operatorname{PolyLog}(2, -ie^x) + \operatorname{PolyLog}(2, ie^x))$$

input `Integrate[ArcTan[E^x],x]`

output `x*ArcTan[E^x] - (I/2)*(x*(Log[1 - I*E^x] - Log[1 + I*E^x]) - PolyLog[2, (-I)*E^x] + PolyLog[2, I*E^x])`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {2720, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \arctan(e^x) dx \\ & \quad \downarrow \text{2720} \\ & \int e^{-x} \arctan(e^x) de^x \\ & \quad \downarrow \text{5355} \\ & \frac{1}{2}i \int e^{-x} \log(1 - ie^x) de^x - \frac{1}{2}i \int e^{-x} \log(1 + ie^x) de^x \\ & \quad \downarrow \text{2838} \\ & \frac{1}{2}i \operatorname{PolyLog}(2, -ie^x) - \frac{1}{2}i \operatorname{PolyLog}(2, ie^x) \end{aligned}$$

input `Int[ArcTan[E^x], x]`

output `(I/2)*PolyLog[2, (-I)*E^x] - (I/2)*PolyLog[2, I*E^x]`

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5355 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.))/(x_), x_Symbol] :> Simp[a*Log[x], x] + (Simp[I*(b/2) Int[Log[1 - I*c*x]/x, x], x] - Simp[I*(b/2) Int[Log[1 + I*c*x]/x, x], x]) /; FreeQ[{a, b, c}, x]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(21) = 42$.

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.71

method	result	size
parts	$x \arctan(e^x) + \frac{ix \ln(1+ie^x)}{2} - \frac{ix \ln(1-ie^x)}{2} + \frac{i \operatorname{dilog}(1+ie^x)}{2} - \frac{i \operatorname{dilog}(1-ie^x)}{2}$	53
derivativedivides	$\ln(e^x) \arctan(e^x) + \frac{i \ln(e^x) \ln(1+ie^x)}{2} - \frac{i \ln(e^x) \ln(1-ie^x)}{2} + \frac{i \operatorname{dilog}(1+ie^x)}{2} - \frac{i \operatorname{dilog}(1-ie^x)}{2}$	59
default	$\ln(e^x) \arctan(e^x) + \frac{i \ln(e^x) \ln(1+ie^x)}{2} - \frac{i \ln(e^x) \ln(1-ie^x)}{2} + \frac{i \operatorname{dilog}(1+ie^x)}{2} - \frac{i \operatorname{dilog}(1-ie^x)}{2}$	59
risch	$-\frac{ix \ln(1+ie^x)}{2} - \frac{i \ln(-i(-e^x+i)) \ln(-ie^x)}{2} + \frac{i \ln(-i(-e^x+i))x}{2} - \frac{i \operatorname{dilog}(-ie^x)}{2} - \frac{i \operatorname{dilog}(1-ie^x)}{2}$	69

input `int(arctan(exp(x)), x, method=_RETURNVERBOSE)`

output `x*arctan(exp(x))+1/2*I*x*ln(1+I*exp(x))-1/2*I*x*ln(1-I*exp(x))+1/2*I*dilog(1+I*exp(x))-1/2*I*dilog(1-I*exp(x))`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 40 vs. $2(15) = 30$.

Time = 0.11 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int \arctan(e^x) dx = x \arctan(e^x) + \frac{1}{2} i x \log(i e^x + 1) - \frac{1}{2} i x \log(-i e^x + 1) - \frac{1}{2} i \operatorname{Li}_2(i e^x) + \frac{1}{2} i \operatorname{Li}_2(-i e^x)$$

input `integrate(arctan(exp(x)),x, algorithm="fricas")`

output `x*arctan(e^x) + 1/2*I*x*log(I*e^x + 1) - 1/2*I*x*log(-I*e^x + 1) - 1/2*I*dilog(I*e^x) + 1/2*I*dilog(-I*e^x)`

Sympy [F]

$$\int \arctan(e^x) dx = \int \operatorname{atan}(e^x) dx$$

input `integrate(atan(exp(x)),x)`

output `Integral(atan(exp(x)), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(15) = 30$.

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.10

$$\int \arctan(e^x) dx = x \arctan(e^x) - \frac{1}{4} \pi \log(e^{2x} + 1) - \frac{1}{2} i \operatorname{Li}_2(i e^x + 1) + \frac{1}{2} i \operatorname{Li}_2(-i e^x + 1)$$

input `integrate(arctan(exp(x)),x, algorithm="maxima")`

output `x*arctan(e^x) - 1/4*pi*log(e^(2*x) + 1) - 1/2*I*dilog(I*e^x + 1) + 1/2*I*dilog(-I*e^x + 1)`

Giac [F]

$$\int \arctan(e^x) dx = \int \arctan(e^x) dx$$

input `integrate(arctan(exp(x)),x, algorithm="giac")`

output `integrate(arctan(e^x), x)`

Mupad [B] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.68

$$\int \arctan(e^x) dx = \frac{\text{polylog}(2, -e^x \text{li}) \text{li}}{2} - \frac{\text{polylog}(2, e^x \text{li}) \text{li}}{2}$$

input `int(atan(exp(x)),x)`

output `(polylog(2, -exp(x)*1i)*1i)/2 - (polylog(2, exp(x)*1i)*1i)/2`

Reduce [F]

$$\int \arctan(e^x) dx = \int \text{atan}(e^x) dx$$

input `int(atan(exp(x)),x)`

output `int(atan(e**x),x)`

3.111 $\int x \arctan(e^x) dx$

Optimal result	857
Mathematica [A] (verified)	857
Rubi [A] (verified)	858
Maple [A] (verified)	859
Fricas [A] (verification not implemented)	860
Sympy [F]	860
Maxima [F]	860
Giac [F]	861
Mupad [F(-1)]	861
Reduce [F]	861

Optimal result

Integrand size = 6, antiderivative size = 63

$$\int x \arctan(e^x) dx = \frac{1}{2}ix \operatorname{PolyLog}(2, -ie^x) - \frac{1}{2}ix \operatorname{PolyLog}(2, ie^x) - \frac{1}{2}i \operatorname{PolyLog}(3, -ie^x) + \frac{1}{2}i \operatorname{PolyLog}(3, ie^x)$$

output

```
1/2*I*x*polylog(2,-I*exp(x))-1/2*I*x*polylog(2,I*exp(x))-1/2*I*polylog(3,-I*exp(x))+1/2*I*polylog(3,I*exp(x))
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int x \arctan(e^x) dx = \frac{1}{2}i(x \operatorname{PolyLog}(2, -ie^x) - x \operatorname{PolyLog}(2, ie^x) - \operatorname{PolyLog}(3, -ie^x) + \operatorname{PolyLog}(3, ie^x))$$

input

```
Integrate[x*ArcTan[E^x],x]
```

output

```
(I/2)*(x*PolyLog[2, (-I)*E^x] - x*PolyLog[2, I*E^x] - PolyLog[3, (-I)*E^x] + PolyLog[3, I*E^x])
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.90, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5666, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x \arctan(e^x) dx \\
 & \quad \downarrow \text{5666} \\
 & \frac{1}{2}i \int x \log(1 - ie^x) dx - \frac{1}{2}i \int x \log(1 + ie^x) dx \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{2}i \left(\int \text{PolyLog}(2, ie^x) dx - x \text{PolyLog}(2, ie^x) \right) - \\
 & \frac{1}{2}i \left(\int \text{PolyLog}(2, -ie^x) dx - x \text{PolyLog}(2, -ie^x) \right) \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2}i \left(\int e^{-x} \text{PolyLog}(2, ie^x) de^x - x \text{PolyLog}(2, ie^x) \right) - \\
 & \frac{1}{2}i \left(\int e^{-x} \text{PolyLog}(2, -ie^x) de^x - x \text{PolyLog}(2, -ie^x) \right) \\
 & \quad \downarrow \text{7143} \\
 & \frac{1}{2}i(\text{PolyLog}(3, ie^x) - x \text{PolyLog}(2, ie^x)) - \frac{1}{2}i(\text{PolyLog}(3, -ie^x) - x \text{PolyLog}(2, -ie^x))
 \end{aligned}$$

input

```
Int [x*ArcTan[E^x] , x]
```

output

```
(-1/2*I)*(-(x*PolyLog[2, (-I)*E^x]) + PolyLog[3, (-I)*E^x]) + (I/2)*(-(x*PolyLog[2, I*E^x]) + PolyLog[3, I*E^x])
```

Definitions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5666 `Int[ArcTan[(a_) + (b_)*(f_)^((c_) + (d_)*(x_))]*(x_)^(m_), x_Symbol] := Simp[I/2 Int[x^m*Log[1 - I*a - I*b*f^(c + d*x)], x], x] - Simp[I/2 Int[x^m*Log[1 + I*a + I*b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IntegerQ[m] && m > 0`

rule 7143 `Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.70

method	result	size
risch	$\frac{ix \operatorname{polylog}(2, -ie^x)}{2} - \frac{ix \operatorname{polylog}(2, ie^x)}{2} - \frac{i \operatorname{polylog}(3, -ie^x)}{2} + \frac{i \operatorname{polylog}(3, ie^x)}{2}$	44

input `int(x*arctan(exp(x)), x, method=_RETURNVERBOSE)`

output `1/2*I*x*polylog(2, -I*exp(x))-1/2*I*x*polylog(2, I*exp(x))-1/2*I*polylog(3, -I*exp(x))+1/2*I*polylog(3, I*exp(x))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\int x \arctan(e^x) dx = \frac{1}{2} x^2 \arctan(e^x) + \frac{1}{4} i x^2 \log(i e^x + 1) - \frac{1}{4} i x^2 \log(-i e^x + 1) - \frac{1}{2} i x \operatorname{Li}_2(i e^x) + \frac{1}{2} i x \operatorname{Li}_2(-i e^x) + \frac{1}{2} i \operatorname{polylog}(3, i e^x) - \frac{1}{2} i \operatorname{polylog}(3, -i e^x)$$

input `integrate(x*arctan(exp(x)),x, algorithm="fricas")`output `1/2*x^2*arctan(e^x) + 1/4*I*x^2*log(I*e^x + 1) - 1/4*I*x^2*log(-I*e^x + 1) - 1/2*I*x*dilog(I*e^x) + 1/2*I*x*dilog(-I*e^x) + 1/2*I*polylog(3, I*e^x) - 1/2*I*polylog(3, -I*e^x)`**Sympy [F]**

$$\int x \arctan(e^x) dx = \int x \operatorname{atan}(e^x) dx$$

input `integrate(x*atan(exp(x)),x)`output `Integral(x*atan(exp(x)), x)`**Maxima [F]**

$$\int x \arctan(e^x) dx = \int x \arctan(e^x) dx$$

input `integrate(x*arctan(exp(x)),x, algorithm="maxima")`output `1/2*x^2*arctan(e^x) - integrate(1/2*x^2*e^x/(e^(2*x) + 1), x)`

Giac [F]

$$\int x \arctan(e^x) dx = \int x \arctan(e^x) dx$$

input `integrate(x*arctan(exp(x)),x, algorithm="giac")`

output `integrate(x*arctan(e^x), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(e^x) dx = \int x \operatorname{atan}(e^x) dx$$

input `int(x*atan(exp(x)),x)`

output `int(x*atan(exp(x)), x)`

Reduce [F]

$$\int x \arctan(e^x) dx = \int \operatorname{atan}(e^x) x dx$$

input `int(x*atan(exp(x)),x)`

output `int(atan(e**x)*x,x)`

3.112 $\int x^2 \arctan(e^x) dx$

Optimal result	862
Mathematica [A] (verified)	862
Rubi [A] (verified)	863
Maple [A] (verified)	865
Fricas [A] (verification not implemented)	865
Sympy [F]	866
Maxima [F]	866
Giac [F]	866
Mupad [F(-1)]	867
Reduce [F]	867

Optimal result

Integrand size = 8, antiderivative size = 91

$$\int x^2 \arctan(e^x) dx = \frac{1}{2}ix^2 \text{PolyLog}(2, -ie^x) - \frac{1}{2}ix^2 \text{PolyLog}(2, ie^x) - ix \text{PolyLog}(3, -ie^x) + ix \text{PolyLog}(3, ie^x) + i \text{PolyLog}(4, -ie^x) - i \text{PolyLog}(4, ie^x)$$

output

```
1/2*I*x^2*polylog(2,-I*exp(x))-1/2*I*x^2*polylog(2,I*exp(x))-I*x*polylog(3,-I*exp(x))+I*x*polylog(3,I*exp(x))+I*polylog(4,-I*exp(x))-I*polylog(4,I*exp(x))
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.88

$$\int x^2 \arctan(e^x) dx = \frac{1}{2}i(x^2 \text{PolyLog}(2, -ie^x) - x^2 \text{PolyLog}(2, ie^x) + 2(-x \text{PolyLog}(3, -ie^x) + x \text{PolyLog}(3, ie^x) + \text{PolyLog}(4, -ie^x) - \text{PolyLog}(4, ie^x)))$$

input

```
Integrate[x^2*ArcTan[E^x],x]
```

output

```
(I/2)*(x^2*PolyLog[2, (-I)*E^x] - x^2*PolyLog[2, I*E^x] + 2*(-(x*PolyLog[3, (-I)*E^x]) + x*PolyLog[3, I*E^x] + PolyLog[4, (-I)*E^x] - PolyLog[4, I*E^x]))
```

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {5666, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(e^x) dx \\
 & \quad \downarrow \text{5666} \\
 & \frac{1}{2}i \int x^2 \log(1 - ie^x) dx - \frac{1}{2}i \int x^2 \log(1 + ie^x) dx \\
 & \quad \downarrow \text{3011} \\
 & \frac{1}{2}i \left(2 \int x \text{PolyLog}(2, ie^x) dx - x^2 \text{PolyLog}(2, ie^x) \right) - \\
 & \frac{1}{2}i \left(2 \int x \text{PolyLog}(2, -ie^x) dx - x^2 \text{PolyLog}(2, -ie^x) \right) \\
 & \quad \downarrow \text{7163} \\
 & \frac{1}{2}i \left(2 \left(x \text{PolyLog}(3, ie^x) - \int \text{PolyLog}(3, ie^x) dx \right) - x^2 \text{PolyLog}(2, ie^x) \right) - \\
 & \frac{1}{2}i \left(2 \left(x \text{PolyLog}(3, -ie^x) - \int \text{PolyLog}(3, -ie^x) dx \right) - x^2 \text{PolyLog}(2, -ie^x) \right) \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2}i \left(2 \left(x \text{PolyLog}(3, ie^x) - \int e^{-x} \text{PolyLog}(3, ie^x) de^x \right) - x^2 \text{PolyLog}(2, ie^x) \right) - \\
 & \frac{1}{2}i \left(2 \left(x \text{PolyLog}(3, -ie^x) - \int e^{-x} \text{PolyLog}(3, -ie^x) de^x \right) - x^2 \text{PolyLog}(2, -ie^x) \right) \\
 & \quad \downarrow \text{7143}
 \end{aligned}$$

$$\frac{1}{2}i(2(x \operatorname{PolyLog}(3, ie^x) - \operatorname{PolyLog}(4, ie^x)) - x^2 \operatorname{PolyLog}(2, ie^x)) - \frac{1}{2}i(2(x \operatorname{PolyLog}(3, -ie^x) - \operatorname{PolyLog}(4, -ie^x)) - x^2 \operatorname{PolyLog}(2, -ie^x))$$

input `Int[x^2*ArcTan[E^x], x]`

output `(-1/2*I)*(-(x^2*PolyLog[2, (-I)*E^x]) + 2*(x*PolyLog[3, (-I)*E^x] - PolyLog[4, (-I)*E^x])) + (I/2)*(-(x^2*PolyLog[2, I*E^x]) + 2*(x*PolyLog[3, I*E^x] - PolyLog[4, I*E^x]))`

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5666 `Int[ArcTan[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol] := Simp[I/2 Int[x^m*Log[1 - I*a - I*b*f^(c + d*x)], x], x] - Simp[I/2 Int[x^m*Log[1 + I*a + I*b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IntegerQ[m] && m > 0`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.))]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163

```
Int[((e._) + (f._)*(x._))^(m._)*PolyLog[n_, (d._)*((F_)^((c._)*((a._) + (b._)
)*(x._)))^(p._)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.77

method	result
risch	$\frac{ix^2 \operatorname{polylog}(2, -ie^x)}{2} - \frac{ix^2 \operatorname{polylog}(2, ie^x)}{2} - ix \operatorname{polylog}(3, -ie^x) + ix \operatorname{polylog}(3, ie^x) + i \operatorname{polylog}(4, -ie^x)$

input

```
int(x^2*arctan(exp(x)),x,method=_RETURNVERBOSE)
```

output

```
1/2*I*x^2*polylog(2,-I*exp(x))-1/2*I*x^2*polylog(2,I*exp(x))-I*x*polylog(3
,-I*exp(x))+I*x*polylog(3,I*exp(x))+I*polylog(4,-I*exp(x))-I*polylog(4,I*
exp(x))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.96

$$\int x^2 \arctan(e^x) dx = \frac{1}{3} x^3 \arctan(e^x) + \frac{1}{6} i x^3 \log(i e^x + 1) - \frac{1}{6} i x^3 \log(-i e^x + 1) \\ - \frac{1}{2} i x^2 \operatorname{Li}_2(i e^x) + \frac{1}{2} i x^2 \operatorname{Li}_2(-i e^x) + i x \operatorname{polylog}(3, i e^x) \\ - i x \operatorname{polylog}(3, -i e^x) - i \operatorname{polylog}(4, i e^x) + i \operatorname{polylog}(4, -i e^x)$$

input

```
integrate(x^2*arctan(exp(x)),x, algorithm="fricas")
```

output

```
1/3*x^3*arctan(e^x) + 1/6*I*x^3*log(I*e^x + 1) - 1/6*I*x^3*log(-I*e^x + 1)
- 1/2*I*x^2*dilog(I*e^x) + 1/2*I*x^2*dilog(-I*e^x) + I*x*polylog(3, I*e^x
) - I*x*polylog(3, -I*e^x) - I*polylog(4, I*e^x) + I*polylog(4, -I*e^x)
```

Sympy [F]

$$\int x^2 \arctan(e^x) dx = \int x^2 \operatorname{atan}(e^x) dx$$

input `integrate(x**2*atan(exp(x)),x)`

output `Integral(x**2*atan(exp(x)), x)`

Maxima [F]

$$\int x^2 \arctan(e^x) dx = \int x^2 \arctan(e^x) dx$$

input `integrate(x^2*arctan(exp(x)),x, algorithm="maxima")`

output `1/3*x^3*arctan(e^x) - integrate(1/3*x^3*e^x/(e^(2*x) + 1), x)`

Giac [F]

$$\int x^2 \arctan(e^x) dx = \int x^2 \arctan(e^x) dx$$

input `integrate(x^2*arctan(exp(x)),x, algorithm="giac")`

output `integrate(x^2*arctan(e^x), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(e^x) dx = \int x^2 \operatorname{atan}(e^x) dx$$

input `int(x^2*atan(exp(x)), x)`output `int(x^2*atan(exp(x)), x)`**Reduce [F]**

$$\int x^2 \arctan(e^x) dx = \int \operatorname{atan}(e^x) x^2 dx$$

input `int(x^2*atan(exp(x)), x)`output `int(atan(e**x)*x**2, x)`

3.113 $\int \arctan(e^{a+bx}) dx$

Optimal result	868
Mathematica [A] (verified)	868
Rubi [A] (verified)	869
Maple [B] (verified)	870
Fricas [B] (verification not implemented)	871
Sympy [F]	871
Maxima [B] (verification not implemented)	872
Giac [F]	872
Mupad [B] (verification not implemented)	873
Reduce [F]	873

Optimal result

Integrand size = 8, antiderivative size = 45

$$\int \arctan(e^{a+bx}) dx = \frac{i \operatorname{PolyLog}(2, -ie^{a+bx})}{2b} - \frac{i \operatorname{PolyLog}(2, ie^{a+bx})}{2b}$$

output

```
1/2*I*polylog(2,-I*exp(b*x+a))/b-1/2*I*polylog(2,I*exp(b*x+a))/b
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.84

$$\int \arctan(e^{a+bx}) dx = x \arctan(e^{a+bx}) - \frac{i(bx(\log(1 - ie^{a+bx}) - \log(1 + ie^{a+bx})) - \operatorname{PolyLog}(2, -ie^{a+bx}) + \operatorname{PolyLog}(2, ie^{a+bx}))}{2b}$$

input

```
Integrate[ArcTan[E^(a + b*x)], x]
```

output

```
x*ArcTan[E^(a + b*x)] - ((I/2)*(b*x*(Log[1 - I*E^(a + b*x)] - Log[1 + I*E^(a + b*x)]) - PolyLog[2, (-I)*E^(a + b*x)] + PolyLog[2, I*E^(a + b*x)]))/b
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2720, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \arctan(e^{a+bx}) dx \\
 \downarrow 2720 \\
 \frac{\int e^{-a-bx} \arctan(e^{a+bx}) de^{a+bx}}{b} \\
 \downarrow 5355 \\
 \frac{\frac{1}{2}i \int e^{-a-bx} \log(1 - ie^{a+bx}) de^{a+bx} - \frac{1}{2}i \int e^{-a-bx} \log(1 + ie^{a+bx}) de^{a+bx}}{b} \\
 \downarrow 2838 \\
 \frac{\frac{1}{2}i \text{PolyLog}(2, -ie^{a+bx}) - \frac{1}{2}i \text{PolyLog}(2, ie^{a+bx})}{b}
 \end{array}$$

input `Int[ArcTan[E^(a + b*x)],x]`

output `((I/2)*PolyLog[2, (-I)*E^(a + b*x)] - (I/2)*PolyLog[2, I*E^(a + b*x)])/b`

Defintions of rubi rules used

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5355 `Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.)/(x_), x_Symbol] := Simp[a*Log[x], x] + (Simp[I*(b/2) Int[Log[1 - I*c*x]/x, x], x] - Simp[I*(b/2) Int[Log[1 + I*c*x]/x, x], x]) /; FreeQ[{a, b, c}, x]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 94 vs. 2(35) = 70.

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.11

method	result
derivativedivides	$\frac{\ln(e^{bx+a}) \arctan(e^{bx+a}) + \frac{i \ln(e^{bx+a}) \ln(1+ie^{bx+a})}{2} - \frac{i \ln(e^{bx+a}) \ln(1-ie^{bx+a})}{2} + \frac{i \operatorname{dilog}(1+ie^{bx+a})}{2} - \frac{i \operatorname{dilog}(1-ie^{bx+a})}{2}}{b}$
default	$\frac{\ln(e^{bx+a}) \arctan(e^{bx+a}) + \frac{i \ln(e^{bx+a}) \ln(1+ie^{bx+a})}{2} - \frac{i \ln(e^{bx+a}) \ln(1-ie^{bx+a})}{2} + \frac{i \operatorname{dilog}(1+ie^{bx+a})}{2} - \frac{i \operatorname{dilog}(1-ie^{bx+a})}{2}}{b}$
parts	$x \arctan(e^{bx+a}) - \frac{i(bx+a) \ln(1+ie^{bx+a})}{2} + \frac{i(bx+a) \ln(1-ie^{bx+a})}{2} - \frac{i \operatorname{dilog}(1+ie^{bx+a})}{2} + \frac{i \operatorname{dilog}(1-ie^{bx+a})}{2} - a \arctan(e^{bx+a})$
risch	$-\frac{ix \ln(1+ie^{bx+a})}{2} + \frac{i \ln(-i(-e^{bx+a}+i))x}{2} - \frac{ia \ln(1+ie^{bx+a})}{2b} - \frac{i \ln(-i(-e^{bx+a}+i)) \ln(-ie^{bx+a})}{2b} + \frac{i \ln(-i(-e^{bx+a}+i)) \ln(-ie^{bx+a})}{2b}$

input `int(arctan(exp(b*x+a)), x, method=_RETURNVERBOSE)`

output `1/b*(ln(exp(b*x+a))*arctan(exp(b*x+a))+1/2*I*ln(exp(b*x+a))*ln(1+I*exp(b*x+a))-1/2*I*ln(exp(b*x+a))*ln(1-I*exp(b*x+a))+1/2*I*dilog(1+I*exp(b*x+a))-1/2*I*dilog(1-I*exp(b*x+a)))`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(29) = 58$.

Time = 0.10 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.29

$$\int \arctan(e^{a+bx}) dx$$

$$= \frac{2bx \arctan(e^{(bx+a)}) + ia \log(e^{(bx+a)} + i) - ia \log(e^{(bx+a)} - i) + (ibx + ia) \log(ie^{(bx+a)} + 1) + (-ibx + ia) \log(-ie^{(bx+a)} + 1)}{2b}$$

input `integrate(arctan(exp(b*x+a)),x, algorithm="fricas")`

output `1/2*(2*b*x*arctan(e^(b*x + a)) + I*a*log(e^(b*x + a) + I) - I*a*log(e^(b*x + a) - I) + (I*b*x + I*a)*log(I*e^(b*x + a) + 1) + (-I*b*x - I*a)*log(-I*e^(b*x + a) + 1) - I*dilog(I*e^(b*x + a)) + I*dilog(-I*e^(b*x + a)))/b`

Sympy [F]

$$\int \arctan(e^{a+bx}) dx = \int \operatorname{atan}(e^{a+bx}) dx$$

input `integrate(atan(exp(a + b*x)), x)`

output `Integral(atan(exp(a + b*x)), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 63 vs. $2(29) = 58$.

Time = 0.13 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.40

$$\begin{aligned} & \int \arctan(e^{a+bx}) dx \\ &= \frac{(bx+a) \arctan(e^{(bx+a)})}{b} \\ & \quad - \frac{\pi \log(e^{(2bx+2a)} + 1) + 2i \operatorname{Li}_2(i e^{(bx+a)} + 1) - 2i \operatorname{Li}_2(-i e^{(bx+a)} + 1)}{4b} \end{aligned}$$

input `integrate(arctan(exp(b*x+a)),x, algorithm="maxima")`

output `(b*x + a)*arctan(e^(b*x + a))/b - 1/4*(pi*log(e^(2*b*x + 2*a) + 1) + 2*I*dilog(I*e^(b*x + a) + 1) - 2*I*dilog(-I*e^(b*x + a) + 1))/b`

Giac [F]

$$\int \arctan(e^{a+bx}) dx = \int \arctan(e^{(bx+a)}) dx$$

input `integrate(arctan(exp(b*x+a)),x, algorithm="giac")`

output `integrate(arctan(e^(b*x + a)), x)`

Mupad [B] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82

$$\int \arctan(e^{a+bx}) dx = -\frac{\operatorname{Li}_2(1 - e^{bx} e^a) \operatorname{li}}{2b} + \frac{\operatorname{Li}_2(1 + e^{bx} e^a) \operatorname{li}}{2b}$$

input `int(atan(exp(a + b*x)),x)`output `(dilog(exp(b*x)*exp(a)*1i + 1)*1i)/(2*b) - (dilog(1 - exp(b*x)*exp(a)*1i)*1i)/(2*b)`**Reduce [F]**

$$\int \arctan(e^{a+bx}) dx = \int \operatorname{atan}(e^{bx+a}) dx$$

input `int(atan(exp(b*x+a)),x)`output `int(atan(e**(a + b*x)),x)`

3.114 $\int x \arctan (e^{a+bx}) dx$

Optimal result	874
Mathematica [A] (verified)	874
Rubi [A] (verified)	875
Maple [B] (verified)	877
Fricas [B] (verification not implemented)	877
Sympy [F]	878
Maxima [F]	878
Giac [F]	879
Mupad [F(-1)]	879
Reduce [F]	879

Optimal result

Integrand size = 10, antiderivative size = 91

$$\int x \arctan (e^{a+bx}) dx = \frac{ix \operatorname{PolyLog} (2, -ie^{a+bx})}{2b} - \frac{ix \operatorname{PolyLog} (2, ie^{a+bx})}{2b} - \frac{i \operatorname{PolyLog} (3, -ie^{a+bx})}{2b^2} + \frac{i \operatorname{PolyLog} (3, ie^{a+bx})}{2b^2}$$

output

```
1/2*I*x*polylog(2,-I*exp(b*x+a))/b-1/2*I*x*polylog(2,I*exp(b*x+a))/b-1/2*I
*polylog(3,-I*exp(b*x+a))/b^2+1/2*I*polylog(3,I*exp(b*x+a))/b^2
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.78

$$\int x \arctan (e^{a+bx}) dx = \frac{i(bx \operatorname{PolyLog} (2, -ie^{a+bx}) - bx \operatorname{PolyLog} (2, ie^{a+bx}) - \operatorname{PolyLog} (3, -ie^{a+bx}) + \operatorname{PolyLog} (3, ie^{a+bx}))}{2b^2}$$

input

```
Integrate[x*ArcTan[E^(a + b*x)],x]
```

output

$$\left(\frac{i}{2} (b*x*\text{PolyLog}[2, (-I)*E^{(a + b*x)}] - b*x*\text{PolyLog}[2, I*E^{(a + b*x)}] - \text{PolyLog}[3, (-I)*E^{(a + b*x)}] + \text{PolyLog}[3, I*E^{(a + b*x)}]) \right) / b^2$$
Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5666, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(e^{a+bx}) dx$$

$$\downarrow 5666$$

$$\frac{1}{2}i \int x \log(1 - ie^{a+bx}) dx - \frac{1}{2}i \int x \log(1 + ie^{a+bx}) dx$$

$$\downarrow 3011$$

$$\frac{1}{2}i \left(\frac{\int \text{PolyLog}(2, ie^{a+bx}) dx}{b} - \frac{x \text{PolyLog}(2, ie^{a+bx})}{b} \right) -$$

$$\frac{1}{2}i \left(\frac{\int \text{PolyLog}(2, -ie^{a+bx}) dx}{b} - \frac{x \text{PolyLog}(2, -ie^{a+bx})}{b} \right)$$

$$\downarrow 2720$$

$$\frac{1}{2}i \left(\frac{\int e^{-a-bx} \text{PolyLog}(2, ie^{a+bx}) de^{a+bx}}{b^2} - \frac{x \text{PolyLog}(2, ie^{a+bx})}{b} \right) -$$

$$\frac{1}{2}i \left(\frac{\int e^{-a-bx} \text{PolyLog}(2, -ie^{a+bx}) de^{a+bx}}{b^2} - \frac{x \text{PolyLog}(2, -ie^{a+bx})}{b} \right)$$

$$\downarrow 7143$$

$$\frac{1}{2}i \left(\frac{\text{PolyLog}(3, ie^{a+bx})}{b^2} - \frac{x \text{PolyLog}(2, ie^{a+bx})}{b} \right) -$$

$$\frac{1}{2}i \left(\frac{\text{PolyLog}(3, -ie^{a+bx})}{b^2} - \frac{x \text{PolyLog}(2, -ie^{a+bx})}{b} \right)$$

input `Int[x*ArcTan[E^(a + b*x)],x]`

output $(-1/2*I)*(-((x*PolyLog[2, (-I)*E^(a + b*x)])/b) + PolyLog[3, (-I)*E^(a + b*x)]/b^2) + (I/2)*(-((x*PolyLog[2, I*E^(a + b*x)])/b) + PolyLog[3, I*E^(a + b*x)]/b^2)$

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n, x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5666 `Int[ArcTan[(a_.) + (b_.)*(f_)^(c_.) + (d_.)*(x_)]*(x_)^(m_.), x_Symbol] := Simp[I/2 Int[x^m*Log[1 - I*a - I*b*f^(c + d*x)], x], x] - Simp[I/2 Int[x^m*Log[1 + I*a + I*b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IntegerQ[m] && m > 0`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 348 vs. $2(71) = 142$.

Time = 0.13 (sec) , antiderivative size = 349, normalized size of antiderivative = 3.84

method	result
risch	$\frac{ia^2 \ln(1+ie^{bx+a})}{2b^2} - \frac{i \operatorname{polylog}(3, -ie^{bx+a})}{2b^2} - \frac{i \ln(1-ie^{bx+a})ax}{2b} + \frac{i \operatorname{dilog}(-ie^{bx+a})a}{2b^2} - \frac{ia^2 \ln(1-ie^{bx+a})}{2b^2} + \frac{i \operatorname{polylog}(2, -ie^{bx+a})}{2b^2}$

input `int(x*arctan(exp(b*x+a)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/2*I/b^2*a^2*\ln(1+I*\exp(b*x+a))-1/2*I*\operatorname{polylog}(3,-I*\exp(b*x+a))/b^2-1/2*I/b* \\ & \ln(1-I*\exp(b*x+a))*a*x+1/2*I/b^2*\operatorname{dilog}(-I*\exp(b*x+a))*a-1/2*I/b^2*a^2*\ln \\ & (1-I*\exp(b*x+a))+1/2*I/b^2*\operatorname{polylog}(2,-I*\exp(b*x+a))*a+1/2*I/b*\ln(-I*(\exp(b \\ & *x+a)+I))*a*x+1/2*I*x*\operatorname{polylog}(2,-I*\exp(b*x+a))/b+1/2*I*\operatorname{polylog}(3,I*\exp(b*x \\ & +a))/b^2-1/2*I/b^2*\operatorname{polylog}(2,I*\exp(b*x+a))*a+1/2*I/b^2*\ln(-I*\exp(b*x+a))* \\ & \ln(-I*(-\exp(b*x+a)+I))*a+1/2*I/b^2*\operatorname{dilog}(-I*(\exp(b*x+a)+I))*a-1/2*I/b^2*\ln \\ & (-I*(-\exp(b*x+a)+I))*a^2-1/2*I*x*\operatorname{polylog}(2,I*\exp(b*x+a))/b+1/2*I/b^2*\ln(-I* \\ & (\exp(b*x+a)+I))*a^2-1/2*I/b*\ln(-I*(-\exp(b*x+a)+I))*a*x+1/2*I/b*\ln(1+I*\exp \\ & (b*x+a))*a*x \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(61) = 122$.

Time = 0.10 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.66

$$\int x \arctan(e^{a+bx}) dx = \frac{2b^2x^2 \arctan(e^{(bx+a)}) - 2ibx \operatorname{Li}_2(ie^{(bx+a)}) + 2ibx \operatorname{Li}_2(-ie^{(bx+a)}) - ia^2 \log(e^{(bx+a)} + i) + ia^2 \log(e^{(bx+a)} - i)}{2b^2}$$

input `integrate(x*arctan(exp(b*x+a)),x, algorithm="fricas")`

output

```
1/4*(2*b^2*x^2*arctan(e^(b*x + a)) - 2*I*b*x*dilog(I*e^(b*x + a)) + 2*I*b*x*dilog(-I*e^(b*x + a)) - I*a^2*log(e^(b*x + a) + I) + I*a^2*log(e^(b*x + a) - I) + (I*b^2*x^2 - I*a^2)*log(I*e^(b*x + a) + 1) + (-I*b^2*x^2 + I*a^2)*log(-I*e^(b*x + a) + 1) + 2*I*polylog(3, I*e^(b*x + a)) - 2*I*polylog(3, -I*e^(b*x + a)))/b^2
```

Sympy [F]

$$\int x \arctan(e^{a+bx}) dx = \int x \operatorname{atan}(e^a e^{bx}) dx$$

input

```
integrate(x*atan(exp(b*x+a)),x)
```

output

```
Integral(x*atan(exp(a)*exp(b*x)), x)
```

Maxima [F]

$$\int x \arctan(e^{a+bx}) dx = \int x \arctan(e^{(bx+a)}) dx$$

input

```
integrate(x*arctan(exp(b*x+a)),x, algorithm="maxima")
```

output

```
1/2*x^2*arctan(e^(b*x + a)) - b*integrate(1/2*x^2*e^(b*x + a)/(e^(2*b*x + 2*a) + 1), x)
```

Giac [F]

$$\int x \arctan(e^{a+bx}) dx = \int x \arctan(e^{(bx+a)}) dx$$

input `integrate(x*arctan(exp(b*x+a)),x, algorithm="giac")`

output `integrate(x*arctan(e^(b*x + a)), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(e^{a+bx}) dx = \int x \operatorname{atan}(e^{a+bx}) dx$$

input `int(x*atan(exp(a + b*x)),x)`

output `int(x*atan(exp(a + b*x)), x)`

Reduce [F]

$$\int x \arctan(e^{a+bx}) dx = \int \operatorname{atan}(e^{bx+a}) x dx$$

input `int(x*atan(exp(b*x+a)),x)`

output `int(atan(e**(a + b*x))*x,x)`

3.115 $\int x^2 \arctan(e^{a+bx}) dx$

Optimal result	880
Mathematica [A] (verified)	880
Rubi [A] (verified)	881
Maple [B] (verified)	883
Fricas [A] (verification not implemented)	884
Sympy [F]	885
Maxima [F]	885
Giac [F]	885
Mupad [F(-1)]	886
Reduce [F]	886

Optimal result

Integrand size = 12, antiderivative size = 133

$$\int x^2 \arctan(e^{a+bx}) dx = \frac{ix^2 \operatorname{PolyLog}(2, -ie^{a+bx})}{2b} - \frac{ix^2 \operatorname{PolyLog}(2, ie^{a+bx})}{2b} - \frac{ix \operatorname{PolyLog}(3, -ie^{a+bx})}{b^2} + \frac{ix \operatorname{PolyLog}(3, ie^{a+bx})}{b^2} + \frac{i \operatorname{PolyLog}(4, -ie^{a+bx})}{b^3} - \frac{i \operatorname{PolyLog}(4, ie^{a+bx})}{b^3}$$

output

```
1/2*I*x^2*polylog(2,-I*exp(b*x+a))/b-1/2*I*x^2*polylog(2,I*exp(b*x+a))/b-I*x*polylog(3,-I*exp(b*x+a))/b^2+I*x*polylog(3,I*exp(b*x+a))/b^2+I*polylog(4,-I*exp(b*x+a))/b^3-I*polylog(4,I*exp(b*x+a))/b^3
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.86

$$\int x^2 \arctan(e^{a+bx}) dx = \frac{i(b^2 x^2 \operatorname{PolyLog}(2, -ie^{a+bx}) - b^2 x^2 \operatorname{PolyLog}(2, ie^{a+bx}) + 2(-bx \operatorname{PolyLog}(3, -ie^{a+bx}) + bx \operatorname{PolyLog}(3, ie^{a+bx})))}{2b^3}$$

input `Integrate[x^2*ArcTan[E^(a + b*x)],x]`

output $((I/2)*(b^2*x^2*PolyLog[2, (-I)*E^(a + b*x)] - b^2*x^2*PolyLog[2, I*E^(a + b*x)] + 2*(-b*x*PolyLog[3, (-I)*E^(a + b*x)]) + b*x*PolyLog[3, I*E^(a + b*x)] + PolyLog[4, (-I)*E^(a + b*x)] - PolyLog[4, I*E^(a + b*x)]))/b^3$

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5666, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 \arctan(e^{a+bx}) dx \\
 & \quad \downarrow 5666 \\
 & \frac{1}{2}i \int x^2 \log(1 - ie^{a+bx}) dx - \frac{1}{2}i \int x^2 \log(1 + ie^{a+bx}) dx \\
 & \quad \downarrow 3011 \\
 & \frac{1}{2}i \left(\frac{2 \int x \text{PolyLog}(2, ie^{a+bx}) dx}{b} - \frac{x^2 \text{PolyLog}(2, ie^{a+bx})}{b} \right) - \\
 & \frac{1}{2}i \left(\frac{2 \int x \text{PolyLog}(2, -ie^{a+bx}) dx}{b} - \frac{x^2 \text{PolyLog}(2, -ie^{a+bx})}{b} \right) \\
 & \quad \downarrow 7163 \\
 & \frac{1}{2}i \left(\frac{2 \left(\frac{x \text{PolyLog}(3, ie^{a+bx})}{b} - \frac{\int \text{PolyLog}(3, ie^{a+bx}) dx}{b} \right) - \frac{x^2 \text{PolyLog}(2, ie^{a+bx})}{b}}{b} \right) - \\
 & \frac{1}{2}i \left(\frac{2 \left(\frac{x \text{PolyLog}(3, -ie^{a+bx})}{b} - \frac{\int \text{PolyLog}(3, -ie^{a+bx}) dx}{b} \right) - \frac{x^2 \text{PolyLog}(2, -ie^{a+bx})}{b}}{b} \right) \\
 & \quad \downarrow 2720
 \end{aligned}$$

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \operatorname{PolyLog}(3, ie^{a+bx})}{b} - \frac{\int e^{-a-bx} \operatorname{PolyLog}(3, ie^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, ie^{a+bx})}{b} \right) -$$

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \operatorname{PolyLog}(3, -ie^{a+bx})}{b} - \frac{\int e^{-a-bx} \operatorname{PolyLog}(3, -ie^{a+bx}) de^{a+bx}}{b^2} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, -ie^{a+bx})}{b} \right)$$

↓ 7143

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \operatorname{PolyLog}(3, ie^{a+bx})}{b} - \frac{\operatorname{PolyLog}(4, ie^{a+bx})}{b^2} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, ie^{a+bx})}{b} \right) -$$

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \operatorname{PolyLog}(3, -ie^{a+bx})}{b} - \frac{\operatorname{PolyLog}(4, -ie^{a+bx})}{b^2} \right)}{b} - \frac{x^2 \operatorname{PolyLog}(2, -ie^{a+bx})}{b} \right)$$

input `Int[x^2*ArcTan[E^(a + b*x)],x]`

output `(-1/2*I)*(-(x^2*PolyLog[2, (-I)*E^(a + b*x)]/b) + (2*((x*PolyLog[3, (-I)*E^(a + b*x)]/b - PolyLog[4, (-I)*E^(a + b*x)]/b^2))/b) + (I/2)*(-(x^2*PolyLog[2, I*E^(a + b*x)]/b) + (2*((x*PolyLog[3, I*E^(a + b*x)]/b - PolyLog[4, I*E^(a + b*x)]/b^2))/b)`

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.))]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x))))^n]/(b*c*n*Log[F]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 5666 `Int[ArcTan[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol] :> Simp[I/2 Int[x^m*Log[1 - I*a - I*b*f^(c + d*x)], x], x] - Simp[I/2 Int[x^m*Log[1 + I*a + I*b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IntegerQ[m] && m > 0`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_)^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 406 vs. $2(111) = 222$.

Time = 0.18 (sec) , antiderivative size = 407, normalized size of antiderivative = 3.06

method	result
risch	$-\frac{i \operatorname{dilog}(-ie^{bx+a})a^2}{2b^3} - \frac{i \operatorname{polylog}(4, ie^{bx+a})}{b^3} + \frac{i \ln(-i(-e^{bx+a}+i))a^2 x}{2b^2} - \frac{ix^2 \operatorname{polylog}(2, ie^{bx+a})}{2b} - \frac{i \ln(1+ie^{bx+a})a^2 x}{2b^2} - \dots$

input `int(x^2*arctan(exp(b*x+a)),x,method=_RETURNVERBOSE)`

output

```
-1/2*I/b^3*dilog(-I*exp(b*x+a))*a^2-I*polylog(4,I*exp(b*x+a))/b^3+1/2*I/b^
2*ln(-I*(-exp(b*x+a)+I))*a^2*x-1/2*I*x^2*polylog(2,I*exp(b*x+a))/b-1/2*I/b
^2*ln(1+I*exp(b*x+a))*a^2*x-1/2*I/b^3*ln(-I*exp(b*x+a))*ln(-I*(-exp(b*x+a)
+I))*a^2+1/2*I/b^3*ln(-I*(-exp(b*x+a)+I))*a^3+1/2*I*x^2*polylog(2,-I*exp(b
*x+a))/b+I*polylog(4,-I*exp(b*x+a))/b^3-1/2*I/b^3*a^3*ln(1+I*exp(b*x+a))+I
*x*polylog(3,I*exp(b*x+a))/b^2+1/2*I/b^3*polylog(2,I*exp(b*x+a))*a^2+1/2*I
/b^3*ln(1-I*exp(b*x+a))*a^3-I*x*polylog(3,-I*exp(b*x+a))/b^2-1/2*I/b^3*dil
og(-I*(exp(b*x+a)+I))*a^2-1/2*I/b^3*polylog(2,-I*exp(b*x+a))*a^2-1/2*I/b^2
*ln(-I*(exp(b*x+a)+I))*x*a^2+1/2*I/b^2*ln(1-I*exp(b*x+a))*x*a^2-1/2*I/b^3*
ln(-I*(exp(b*x+a)+I))*a^3
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.41

$$\int x^2 \arctan(e^{a+bx}) dx$$

$$= \frac{2b^3x^3 \arctan(e^{(bx+a)}) - 3ib^2x^2 \text{Li}_2(ie^{(bx+a)}) + 3ib^2x^2 \text{Li}_2(-ie^{(bx+a)}) + ia^3 \log(e^{(bx+a)} + i) - ia^3 \log(e^{(bx+a)} - i)}{b^3}$$

input

```
integrate(x^2*arctan(exp(b*x+a)),x, algorithm="fricas")
```

output

```
1/6*(2*b^3*x^3*arctan(e^(b*x + a)) - 3*I*b^2*x^2*dilog(I*e^(b*x + a)) + 3*
I*b^2*x^2*dilog(-I*e^(b*x + a)) + I*a^3*log(e^(b*x + a) + I) - I*a^3*log(e
^(b*x + a) - I) + 6*I*b*x*polylog(3, I*e^(b*x + a)) - 6*I*b*x*polylog(3, -
I*e^(b*x + a)) + (I*b^3*x^3 + I*a^3)*log(I*e^(b*x + a) + 1) + (-I*b^3*x^3
- I*a^3)*log(-I*e^(b*x + a) + 1) - 6*I*polylog(4, I*e^(b*x + a)) + 6*I*pol
ylog(4, -I*e^(b*x + a)))/b^3
```

Sympy [F]

$$\int x^2 \arctan(e^{a+bx}) dx = \int x^2 \operatorname{atan}(e^a e^{bx}) dx$$

input `integrate(x**2*atan(exp(b*x+a)),x)`

output `Integral(x**2*atan(exp(a)*exp(b*x)), x)`

Maxima [F]

$$\int x^2 \arctan(e^{a+bx}) dx = \int x^2 \arctan(e^{(bx+a)}) dx$$

input `integrate(x^2*arctan(exp(b*x+a)),x, algorithm="maxima")`

output `1/3*x^3*arctan(e^(b*x + a)) - b*integrate(1/3*x^3*e^(b*x + a)/(e^(2*b*x + 2*a) + 1), x)`

Giac [F]

$$\int x^2 \arctan(e^{a+bx}) dx = \int x^2 \arctan(e^{(bx+a)}) dx$$

input `integrate(x^2*arctan(exp(b*x+a)),x, algorithm="giac")`

output `integrate(x^2*arctan(e^(b*x + a)), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(e^{a+bx}) dx = \int x^2 \operatorname{atan}(e^{a+bx}) dx$$

input `int(x^2*atan(exp(a + b*x)),x)`output `int(x^2*atan(exp(a + b*x)), x)`**Reduce [F]**

$$\int x^2 \arctan(e^{a+bx}) dx = \int \operatorname{atan}(e^{bx+a}) x^2 dx$$

input `int(x^2*atan(exp(b*x+a)),x)`output `int(atan(e**(a + b*x))*x**2,x)`

3.116 $\int \arctan(a + bf^{c+dx}) dx$

Optimal result	887
Mathematica [A] (verified)	888
Rubi [A] (verified)	888
Maple [A] (verified)	891
Fricas [A] (verification not implemented)	891
Sympy [F]	892
Maxima [A] (verification not implemented)	892
Giac [F]	893
Mupad [F(-1)]	893
Reduce [F]	893

Optimal result

Integrand size = 12, antiderivative size = 196

$$\int \arctan(a + bf^{c+dx}) dx = -\frac{\arctan(a + bf^{c+dx}) \log\left(\frac{2}{1-i(a+bf^{c+dx})}\right)}{d \log(f)} + \frac{\arctan(a + bf^{c+dx}) \log\left(\frac{2bf^{c+dx}}{(i-a)(1-i(a+bf^{c+dx}))}\right)}{d \log(f)} + \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bf^{c+dx})}\right)}{2d \log(f)} - \frac{i \operatorname{PolyLog}\left(2, 1 - \frac{2bf^{c+dx}}{(i-a)(1-i(a+bf^{c+dx}))}\right)}{2d \log(f)}$$

output `-arctan(a+b*f^(d*x+c))*ln(2/(1-I*(a+b*f^(d*x+c))))/d/ln(f)+arctan(a+b*f^(d*x+c))*ln(2*b*f^(d*x+c)/(I-a)/(1-I*(a+b*f^(d*x+c))))/d/ln(f)+1/2*I*polylog(2,1-2/(1-I*(a+b*f^(d*x+c))))/d/ln(f)-1/2*I*polylog(2,1-2*b*f^(d*x+c)/(I-a)/(1-I*(a+b*f^(d*x+c))))/d/ln(f)`

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.85

$$\int \arctan(a + bf^{c+dx}) dx = x \arctan(a + bf^{c+dx}) - \frac{b \left(dx \log(f) \left(\log \left(1 + \frac{b^2 f^{c+dx}}{ab - \sqrt{-b^2}} \right) - \log \left(1 + \frac{b^2 f^{c+dx}}{ab + \sqrt{-b^2}} \right) \right) + \text{PolyLog} \left(2, -\frac{b^2 f^{c+dx}}{ab - \sqrt{-b^2}} \right) - \text{PolyLog} \left(2, -\frac{b^2 f^{c+dx}}{ab + \sqrt{-b^2}} \right) \right)}{2\sqrt{-b^2}d \log(f)}$$

input `Integrate[ArcTan[a + b*f^(c + d*x)],x]`

output `x*ArcTan[a + b*f^(c + d*x)] - (b*(d*x*Log[f]*(Log[1 + (b^2*f^(c + d*x))/(a*b - Sqrt[-b^2]]) - Log[1 + (b^2*f^(c + d*x))/(a*b + Sqrt[-b^2]]) + PolyLog[2, -((b^2*f^(c + d*x))/(a*b - Sqrt[-b^2]))] - PolyLog[2, -((b^2*f^(c + d*x))/(a*b + Sqrt[-b^2]))])))/(2*Sqrt[-b^2]*d*Log[f])`

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.90, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {2720, 5570, 25, 27, 5381, 2849, 2752, 2897}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \arctan(a + bf^{c+dx}) dx \\ & \quad \downarrow 2720 \\ & \frac{\int f^{-c-dx} \arctan(bf^{c+dx} + a) df^{c+dx}}{d \log(f)} \\ & \quad \downarrow 5570 \\ & \frac{\int f^{-c-dx} \arctan(bf^{c+dx} + a) d(bf^{c+dx} + a)}{bd \log(f)} \\ & \quad \downarrow 25 \end{aligned}$$

$$\begin{aligned}
& \frac{\int -f^{-c-dx} \arctan(bf^{c+dx} + a) d(bf^{c+dx} + a)}{bd \log(f)} \\
& \quad \downarrow 27 \\
& \frac{\int -\frac{f^{-c-dx} \arctan(bf^{c+dx} + a)}{b} d(bf^{c+dx} + a)}{d \log(f)} \\
& \quad \downarrow 5381 \\
& \frac{-\int \frac{\log\left(\frac{2}{1-i(bf^{c+dx} + a)}\right)}{f^{2c+2dx+1}} d(bf^{c+dx} + a) + \int \frac{\log\left(\frac{2bf^{c+dx}}{(i-a)(1-i(bf^{c+dx} + a))}\right)}{f^{2c+2dx+1}} d(bf^{c+dx} + a) + \arctan(a + bf^{c+dx}) \log\left(\frac{2}{1-i(bf^{c+dx} + a)}\right)}{d \log(f)} \\
& \quad \downarrow 2849 \\
& \frac{-i \int \frac{\log\left(\frac{2}{1-i(bf^{c+dx} + a)}\right)}{1-\frac{2}{1-i(bf^{c+dx} + a)}} d\frac{1}{1-i(bf^{c+dx} + a)} + \int \frac{\log\left(\frac{2bf^{c+dx}}{(i-a)(1-i(bf^{c+dx} + a))}\right)}{f^{2c+2dx+1}} d(bf^{c+dx} + a) + \arctan(a + bf^{c+dx}) \log\left(\frac{2}{1-i(bf^{c+dx} + a)}\right)}{d \log(f)} \\
& \quad \downarrow 2752 \\
& \frac{\int \frac{\log\left(\frac{2bf^{c+dx}}{(i-a)(1-i(bf^{c+dx} + a))}\right)}{f^{2c+2dx+1}} d(bf^{c+dx} + a) + \arctan(a + bf^{c+dx}) \log\left(\frac{2}{1-i(a+bf^{c+dx})}\right) - \arctan(a + bf^{c+dx}) \log\left(\frac{2}{1-i(bf^{c+dx} + a)}\right)}{d \log(f)} \\
& \quad \downarrow 2897 \\
& \frac{\arctan(a + bf^{c+dx}) \log\left(\frac{2}{1-i(a+bf^{c+dx})}\right) - \arctan(a + bf^{c+dx}) \log\left(\frac{2bf^{c+dx}}{(-a+i)(1-i(a+bf^{c+dx}))}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, 1 - \frac{2}{1-i(a+bf^{c+dx})}\right)}{d \log(f)}
\end{aligned}$$

input `Int[ArcTan[a + b*f^(c + d*x)],x]`

output `-((ArcTan[a + b*f^(c + d*x)]*Log[2/(1 - I*(a + b*f^(c + d*x))]) - ArcTan[a + b*f^(c + d*x)]*Log[(2*b*f^(c + d*x))/((I - a)*(1 - I*(a + b*f^(c + d*x)))] - (I/2)*PolyLog[2, 1 - 2/(1 - I*(a + b*f^(c + d*x)))] + (I/2)*PolyLog[2, 1 - (2*b*f^(c + d*x))/((I - a)*(1 - I*(a + b*f^(c + d*x)))])/(d*Log[f]))`

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]`
- rule 2752 `Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]`
- rule 2849 `Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Simp[-e/g Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]`
- rule 2897 `Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]`
- rule 5381 `Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-a + b*ArcTan[c*x])*(Log[2/(1 - I*c*x)]/e), x] + (Simp[(a + b*ArcTan[c*x])*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x] + Simp[b*(c/e) Int[Log[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - Simp[b*(c/e) Int[Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(1 + c^2*x^2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]`

rule 5570

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[1/d Subst[Int[((d*e - c*f)/d + f*(x/d))^m*(a + b*ArcTan[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && I GtQ[p, 0]
```

Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\ln(-b f^{dx+c}) \arctan(a+b f^{dx+c}) - \frac{i \ln(-b f^{dx+c}) \ln\left(\frac{i+b f^{dx+c}+a}{a+i}\right)}{2} + \frac{i \ln(-b f^{dx+c}) \ln\left(\frac{i-b f^{dx+c}-a}{i-a}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{i+b f^{dx+c}+a}{a+i}\right)}{2}}{d \ln(f)}$
default	$\frac{\ln(-b f^{dx+c}) \arctan(a+b f^{dx+c}) - \frac{i \ln(-b f^{dx+c}) \ln\left(\frac{i+b f^{dx+c}+a}{a+i}\right)}{2} + \frac{i \ln(-b f^{dx+c}) \ln\left(\frac{i-b f^{dx+c}-a}{i-a}\right)}{2} - \frac{i \operatorname{dilog}\left(\frac{i+b f^{dx+c}+a}{a+i}\right)}{2}}{d \ln(f)}$
risch	$-\frac{i x \ln(1+i(a+b f^{dx+c}))}{2} + \frac{i \operatorname{dilog}\left(\frac{b f^{dx} f^c+a-i}{a-i}\right)}{2 \ln(f) d} + \frac{i \ln\left(\frac{b f^{dx} f^c+a-i}{a-i}\right) x}{2} + \frac{i \ln\left(\frac{b f^{dx} f^c+a-i}{a-i}\right) c}{2 d} - \frac{i c \ln(i f^{dx+c})}{2 d}$

input

```
int(arctan(a+b*f^(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/d/ln(f)*(ln(-b*f^(d*x+c))*arctan(a+b*f^(d*x+c))-1/2*I*ln(-b*f^(d*x+c))*ln((I+b*f^(d*x+c)+a)/(a+I))+1/2*I*ln(-b*f^(d*x+c))*ln((I-b*f^(d*x+c)-a)/(I-a))-1/2*I*dilog((I+b*f^(d*x+c)+a)/(a+I))+1/2*I*dilog((I-b*f^(d*x+c)-a)/(I-a)))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.08

$$\int \arctan(a + b f^{c+dx}) dx$$

$$= \frac{2 dx \arctan(b f^{dx+c} + a) \log(f) + i c \log(b f^{dx+c} + a + i) \log(f) - i c \log(b f^{dx+c} + a - i) \log(f) + (i dx \arctan(a + b f^{c+dx}) - \frac{1}{2} \ln(-b f^{dx+c}) \ln\left(\frac{i+b f^{dx+c}+a}{a+i}\right) + \frac{1}{2} \ln(-b f^{dx+c}) \ln\left(\frac{i-b f^{dx+c}-a}{i-a}\right) - \frac{1}{2} \operatorname{dilog}\left(\frac{i+b f^{dx+c}+a}{a+i}\right) + \frac{1}{2} \operatorname{dilog}\left(\frac{i-b f^{dx+c}-a}{i-a}\right))}{d \ln(f)}$$

input

```
integrate(arctan(a+b*f^(d*x+c)),x, algorithm="fricas")
```

output

```
1/2*(2*d*x*arctan(b*f^(d*x + c) + a)*log(f) + I*c*log(b*f^(d*x + c) + a +
I)*log(f) - I*c*log(b*f^(d*x + c) + a - I)*log(f) + (I*d*x + I*c)*log(f)*l
og((a^2 + (a*b + I*b)*f^(d*x + c) + 1)/(a^2 + 1)) + (-I*d*x - I*c)*log(f)*
log((a^2 + (a*b - I*b)*f^(d*x + c) + 1)/(a^2 + 1)) + I*dilog(-(a^2 + (a*b
+ I*b)*f^(d*x + c) + 1)/(a^2 + 1) + 1) - I*dilog(-(a^2 + (a*b - I*b)*f^(d*
x + c) + 1)/(a^2 + 1) + 1))/(d*log(f))
```

Sympy [F]

$$\int \arctan(a + bf^{c+dx}) dx = \int \operatorname{atan}(a + bf^{c+dx}) dx$$

input

```
integrate(atan(a+b*f**(d*x+c)),x)
```

output

```
Integral(atan(a + b*f**(c + d*x)), x)
```

Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.96

$$\int \arctan(a + bf^{c+dx}) dx = \frac{(dx + c) \arctan(bf^{dx+c} + a)}{d} - \frac{2(dx + c) \arctan\left(\frac{b^2 f^{dx+c} + ab}{b}\right) \log(f) + \left(\pi - \arctan\left(\frac{1}{a}\right)\right) \log(b^2 f^{2dx+2c} + 2abf^{dx+c} + a^2 + 1) - \arctan\left(\frac{b^2 f^{dx+c} + ab}{b}\right)}{2d \log(f)}$$

input

```
integrate(arctan(a+b*f^(d*x+c)),x, algorithm="maxima")
```

output

```
(d*x + c)*arctan(b*f^(d*x + c) + a)/d - 1/2*(2*(d*x + c)*arctan((b^2*f^(d*
x + c) + a*b)/b)*log(f) + (pi - arctan(1/a))*log(b^2*f^(2*d*x + 2*c) + 2*a
*b*f^(d*x + c) + a^2 + 1) - arctan(b*f^(d*x + c) + a)*log(b^2*f^(2*d*x + 2
*c)/(a^2 + 1)) + I*dilog((I*b*f^(d*x + c) + I*a + 1)/(I*a + 1)) - I*dilog(
(I*b*f^(d*x + c) + I*a - 1)/(I*a - 1)))/(d*log(f))
```

Giac [F]

$$\int \arctan(a + bf^{c+dx}) dx = \int \arctan(bf^{dx+c} + a) dx$$

input `integrate(arctan(a+b*f^(d*x+c)),x, algorithm="giac")`

output `integrate(arctan(b*f^(d*x + c) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \arctan(a + bf^{c+dx}) dx = \int \operatorname{atan}(a + bf^{c+dx}) dx$$

input `int(atan(a + b*f^(c + d*x)),x)`

output `int(atan(a + b*f^(c + d*x)), x)`

Reduce [F]

$$\int \arctan(a + bf^{c+dx}) dx = \int \operatorname{atan}(f^{dx+c}b + a) dx$$

input `int(atan(a+b*f^(d*x+c)),x)`

output `int(atan(f**(c + d*x)*b + a),x)`

3.117 $\int x \arctan(a + bf^{c+dx}) dx$

Optimal result	894
Mathematica [A] (verified)	895
Rubi [A] (verified)	895
Maple [B] (verified)	898
Fricas [A] (verification not implemented)	898
Sympy [F]	899
Maxima [F]	899
Giac [F]	900
Mupad [F(-1)]	900
Reduce [F]	900

Optimal result

Integrand size = 14, antiderivative size = 232

$$\int x \arctan(a + bf^{c+dx}) dx = \frac{1}{2}x^2 \arctan(a + bf^{c+dx}) - \frac{1}{4}ix^2 \log\left(1 - \frac{ibf^{c+dx}}{1 - ia}\right) + \frac{1}{4}ix^2 \log\left(1 + \frac{ibf^{c+dx}}{1 + ia}\right) - \frac{ix \operatorname{PolyLog}\left(2, \frac{ibf^{c+dx}}{1 - ia}\right)}{2d \log(f)} + \frac{ix \operatorname{PolyLog}\left(2, -\frac{ibf^{c+dx}}{1 + ia}\right)}{2d \log(f)} + \frac{i \operatorname{PolyLog}\left(3, \frac{ibf^{c+dx}}{1 - ia}\right)}{2d^2 \log^2(f)} - \frac{i \operatorname{PolyLog}\left(3, -\frac{ibf^{c+dx}}{1 + ia}\right)}{2d^2 \log^2(f)}$$

output

```
1/2*x^2*arctan(a+b*f^(d*x+c))-1/4*I*x^2*ln(1-I*b*f^(d*x+c)/(1-I*a))+1/4*I*x^2*ln(1+I*b*f^(d*x+c)/(1+I*a))-1/2*I*x*polylog(2,I*b*f^(d*x+c)/(1-I*a))/d/ln(f)+1/2*I*x*polylog(2,-I*b*f^(d*x+c)/(1+I*a))/d/ln(f)+1/2*I*polylog(3,I*b*f^(d*x+c)/(1-I*a))/d^2/ln(f)^2-1/2*I*polylog(3,-I*b*f^(d*x+c)/(1+I*a))/d^2/ln(f)^2
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.02

$$\int x \arctan(a + bf^{c+dx}) dx$$

$$= \frac{i(d^2x^2 \log^2(f) \log(1 - ia - ibf^{c+dx}) - d^2x^2 \log^2(f) \log(1 + ia + ibf^{c+dx}) - d^2x^2 \log^2(f) \log\left(\frac{i+a+bf^{c+dx}}{i+a}\right))}{d^2}$$

input

```
Integrate[x*ArcTan[a + b*f^(c + d*x)],x]
```

output

```
((I/4)*(d^2*x^2*Log[f]^2*Log[1 - I*a - I*b*f^(c + d*x)] - d^2*x^2*Log[f]^2*Log[1 + I*a + I*b*f^(c + d*x)] - d^2*x^2*Log[f]^2*Log[(I + a + b*f^(c + d*x))/(I + a)] + d^2*x^2*Log[f]^2*Log[1 + (b*f^(c + d*x))/(-I + a)] + 2*d*x*Log[f]*PolyLog[2, (b*f^(c + d*x))/(I - a)] - 2*d*x*Log[f]*PolyLog[2, -(b*f^(c + d*x))/(I + a)]) - 2*PolyLog[3, (b*f^(c + d*x))/(I - a)] + 2*PolyLog[3, -(b*f^(c + d*x))/(I + a)))/(d^2*Log[f]^2)
```

Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5666, 3012, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \arctan(a + bf^{c+dx}) dx$$

$$\downarrow \text{5666}$$

$$\frac{1}{2}i \int x \log(-ibf^{c+dx} - ia + 1) dx - \frac{1}{2}i \int x \log(ibf^{c+dx} + ia + 1) dx$$

$$\downarrow \text{3012}$$

$$\frac{1}{2}i \left(\int x \log \left(1 - \frac{ibf^{c+dx}}{1-ia} \right) dx + \frac{1}{2}x^2 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\int x \log \left(\frac{ibf^{c+dx}}{ia+1} + 1 \right) dx + \frac{1}{2}x^2 \log(ia + ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 3011

$$\frac{1}{2}i \left(\frac{\int \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right) dx}{d \log(f)} - \frac{x \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} + \frac{1}{2}x^2 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\frac{\int \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right) dx}{d \log(f)} - \frac{x \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} + \frac{1}{2}x^2 \log(ia + ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 2720

$$\frac{1}{2}i \left(\frac{\int f^{-c-dx} \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right) df^{c+dx}}{d^2 \log^2(f)} - \frac{x \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} + \frac{1}{2}x^2 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\frac{\int f^{-c-dx} \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right) df^{c+dx}}{d^2 \log^2(f)} - \frac{x \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} + \frac{1}{2}x^2 \log(ia + ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 7143

$$\frac{1}{2}i \left(\frac{\text{PolyLog} \left(3, -\frac{bf^{c+dx}}{a+i} \right)}{d^2 \log^2(f)} - \frac{x \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} + \frac{1}{2}x^2 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\frac{\text{PolyLog} \left(3, \frac{bf^{c+dx}}{i-a} \right)}{d^2 \log^2(f)} - \frac{x \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} + \frac{1}{2}x^2 \log(ia + ibf^{c+dx} + 1) - \frac{1}{2}x^2 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

input `Int[x*ArcTan[a + b*f^(c + d*x)],x]`

output `(-1/2*I)*((x^2*Log[1 + I*a + I*b*f^(c + d*x)]/2 - (x^2*Log[1 - (b*f^(c + d*x))/(I - a)]/2 - (x*PolyLog[2, (b*f^(c + d*x))/(I - a)]/(d*Log[f]) + PolyLog[3, (b*f^(c + d*x))/(I - a)]/(d^2*Log[f]^2)) + (I/2)*((x^2*Log[1 - I*a - I*b*f^(c + d*x)]/2 - (x^2*Log[1 + (b*f^(c + d*x))/(I + a)]/2 - (x*PolyLog[2, -(b*f^(c + d*x))/(I + a)]/(d*Log[f]) + PolyLog[3, -(b*f^(c + d*x))/(I + a)]/(d^2*Log[f]^2))`

Definitions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3012 `Int[Log[(d_) + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(f + g*x)^(m + 1)*(Log[d + e*(F^(c*(a + b*x)))^n]/(g*(m + 1))), x] + (Int[(f + g*x)^m*Log[1 + (e/d)*(F^(c*(a + b*x)))^n], x] - Simp[(f + g*x)^(m + 1)*(Log[1 + (e/d)*(F^(c*(a + b*x)))^n]/(g*(m + 1))), x]) /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && NeQ[d, 1]`

rule 5666 `Int[ArcTan[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol] := Simp[I/2 Int[x^m*Log[1 - I*a - I*b*f^(c + d*x)], x], x] - Simp[I/2 Int[x^m*Log[1 + I*a + I*b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] && IntegerQ[m] && m > 0`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 671 vs. 2(200) = 400.

Time = 0.68 (sec) , antiderivative size = 672, normalized size of antiderivative = 2.90

method	result
risch	$-\frac{ic^2 \ln(1-ia-if^{dx}f^cb)}{4d^2} + \frac{i \ln\left(1-\frac{ibf^{dx}f^c}{-ia-1}\right)cx}{2d} + \frac{ic^2 \ln\left(\frac{bf^{dx}f^c+a+i}{a+i}\right)}{2d^2} + \frac{i \operatorname{polylog}\left(2, \frac{ibf^{dx}f^c}{-ia-1}\right)c}{2 \ln(f)d^2} - \frac{ix^2 \ln(1+i(a+bf^{dx}f^c))}{4}$

```
input int(x*arctan(a+b*f^(d*x+c)), x, method=_RETURNVERBOSE)
```

```
output -1/4*I/d^2*c^2*ln(1-I*a-I*f^(d*x)*f^c*b)+1/2*I/d*ln(1-I*b/(-I*a-1)*f^(d*x)*f^c)*c*x+1/2*I/d^2*c^2*ln((b*f^(d*x)*f^c+a+I)/(a+I))+1/2*I/ln(f)/d^2*polylog(2,I*b/(-I*a-1)*f^(d*x)*f^c)*c-1/4*I*x^2*ln(1+I*(a+b*f^(d*x+c)))-1/2*I/ln(f)^2/d^2*polylog(3,I*b/(-I*a-1)*f^(d*x)*f^c)+1/2*I/ln(f)^2/d^2*polylog(3,I*b/(1-I*a)*f^(d*x)*f^c)+1/4*I/d^2*ln(1-I*b/(-I*a-1)*f^(d*x)*f^c)*c^2-1/4*I*I*ln(1-I*b/(1-I*a)*f^(d*x)*f^c)*x^2+1/4*I/d^2*c^2*ln(I*f^(d*x)*f^c*b+I*a+1)+1/2*I/ln(f)/d*polylog(2,I*b/(-I*a-1)*f^(d*x)*f^c)*x-1/4*I/d^2*ln(1-I*b/(1-I*a)*f^(d*x)*f^c)*c^2-1/2*I/d^2*c^2*ln((b*f^(d*x)*f^c+a-I)/(a-I))-1/2*I/d*c*ln((b*f^(d*x)*f^c+a-I)/(a-I))*x-1/2*I/ln(f)/d*polylog(2,I*b/(1-I*a)*f^(d*x)*f^c)*x+1/4*I*x^2*ln(1-I*(a+b*f^(d*x+c)))-1/2*I/d*ln(1-I*b/(1-I*a)*f^(d*x)*f^c)*c*x+1/2*I/d*c*ln((b*f^(d*x)*f^c+a+I)/(a+I))*x+1/2*I/ln(f)/d^2*c*dilog((b*f^(d*x)*f^c+a+I)/(a+I))+1/4*I*ln(1-I*b/(-I*a-1)*f^(d*x)*f^c)*x^2-1/2*I/ln(f)/d^2*c*dilog((b*f^(d*x)*f^c+a-I)/(a-I))-1/2*I/ln(f)/d^2*polylog(2,I*b/(1-I*a)*f^(d*x)*f^c)*c
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.31

$$\int x \arctan(a + bf^{c+dx}) dx$$

$$= \frac{2d^2x^2 \arctan(bf^{dx+c} + a) \log(f)^2 - ic^2 \log(bf^{dx+c} + a + i) \log(f)^2 + ic^2 \log(bf^{dx+c} + a - i) \log(f)^2}{4}$$

input `integrate(x*arctan(a+b*f^(d*x+c)),x, algorithm="fricas")`

output `1/4*(2*d^2*x^2*arctan(b*f^(d*x + c) + a)*log(f)^2 - I*c^2*log(b*f^(d*x + c) + a + I)*log(f)^2 + I*c^2*log(b*f^(d*x + c) + a - I)*log(f)^2 + 2*I*d*x*dilog(-(a^2 + (a*b + I*b)*f^(d*x + c) + 1)/(a^2 + 1) + 1)*log(f) - 2*I*d*x*dilog(-(a^2 + (a*b - I*b)*f^(d*x + c) + 1)/(a^2 + 1) + 1)*log(f) + (I*d^2*x^2 - I*c^2)*log(f)^2*log((a^2 + (a*b + I*b)*f^(d*x + c) + 1)/(a^2 + 1)) + (-I*d^2*x^2 + I*c^2)*log(f)^2*log((a^2 + (a*b - I*b)*f^(d*x + c) + 1)/(a^2 + 1)) - 2*I*polylog(3, -(a*b + I*b)*f^(d*x + c)/(a^2 + 1)) + 2*I*polylog(3, -(a*b - I*b)*f^(d*x + c)/(a^2 + 1)))/(d^2*log(f)^2)`

Sympy [F]

$$\int x \arctan(a + bf^{c+dx}) dx = \int x \operatorname{atan}(a + bf^{c+dx}) dx$$

input `integrate(x*atan(a+b*f**(d*x+c)),x)`

output `Integral(x*atan(a + b*f**(c + d*x)), x)`

Maxima [F]

$$\int x \arctan(a + bf^{c+dx}) dx = \int x \arctan(bf^{dx+c} + a) dx$$

input `integrate(x*arctan(a+b*f^(d*x+c)),x, algorithm="maxima")`

output `-b*d*f^c*integrate(1/2*f^(d*x)*x^2/(b^2*f^(2*d*x)*f^(2*c) + 2*a*b*f^(d*x)*f^c + a^2 + 1), x)*log(f) + 1/2*x^2*arctan(b*f^(d*x)*f^c + a)`

Giac [F]

$$\int x \arctan(a + bf^{c+dx}) dx = \int x \arctan(bf^{dx+c} + a) dx$$

input `integrate(x*arctan(a+b*f^(d*x+c)),x, algorithm="giac")`

output `integrate(x*arctan(b*f^(d*x + c) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x \arctan(a + bf^{c+dx}) dx = \int x \operatorname{atan}(a + bf^{c+dx}) dx$$

input `int(x*atan(a + b*f^(c + d*x)),x)`

output `int(x*atan(a + b*f^(c + d*x)), x)`

Reduce [F]

$$\int x \arctan(a + bf^{c+dx}) dx = \int \operatorname{atan}(f^{dx+c}b + a) x dx$$

input `int(x*atan(a+b*f^(d*x+c)),x)`

output `int(atan(f**(c + d*x)*b + a)*x,x)`

3.118 $\int x^2 \arctan(a + bf^{c+dx}) dx$

Optimal result	901
Mathematica [A] (verified)	902
Rubi [A] (verified)	902
Maple [B] (verified)	906
Fricas [A] (verification not implemented)	906
Sympy [F]	907
Maxima [F]	907
Giac [F]	908
Mupad [F(-1)]	908
Reduce [F]	908

Optimal result

Integrand size = 16, antiderivative size = 302

$$\begin{aligned} \int x^2 \arctan(a + bf^{c+dx}) dx &= \frac{1}{3}x^3 \arctan(a + bf^{c+dx}) - \frac{1}{6}ix^3 \log\left(1 - \frac{ibf^{c+dx}}{1-ia}\right) \\ &+ \frac{1}{6}ix^3 \log\left(1 + \frac{ibf^{c+dx}}{1+ia}\right) - \frac{ix^2 \operatorname{PolyLog}\left(2, \frac{ibf^{c+dx}}{1-ia}\right)}{2d \log(f)} \\ &+ \frac{ix^2 \operatorname{PolyLog}\left(2, -\frac{ibf^{c+dx}}{1+ia}\right)}{2d \log(f)} \\ &+ \frac{ix \operatorname{PolyLog}\left(3, \frac{ibf^{c+dx}}{1-ia}\right)}{d^2 \log^2(f)} - \frac{ix \operatorname{PolyLog}\left(3, -\frac{ibf^{c+dx}}{1+ia}\right)}{d^2 \log^2(f)} \\ &- \frac{i \operatorname{PolyLog}\left(4, \frac{ibf^{c+dx}}{1-ia}\right)}{d^3 \log^3(f)} + \frac{i \operatorname{PolyLog}\left(4, -\frac{ibf^{c+dx}}{1+ia}\right)}{d^3 \log^3(f)} \end{aligned}$$

output

```
1/3*x^3*arctan(a+b*f^(d*x+c))-1/6*I*x^3*ln(1-I*b*f^(d*x+c)/(1-I*a))+1/6*I*x^3*ln(1+I*b*f^(d*x+c)/(1+I*a))-1/2*I*x^2*polylog(2,I*b*f^(d*x+c)/(1-I*a))/d/ln(f)+1/2*I*x^2*polylog(2,-I*b*f^(d*x+c)/(1+I*a))/d/ln(f)+I*x*polylog(3,I*b*f^(d*x+c)/(1-I*a))/d^2/ln(f)^2-I*x*polylog(3,-I*b*f^(d*x+c)/(1+I*a))/d^2/ln(f)^2-I*polylog(4,I*b*f^(d*x+c)/(1-I*a))/d^3/ln(f)^3+I*polylog(4,-I*b*f^(d*x+c)/(1+I*a))/d^3/ln(f)^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 299, normalized size of antiderivative = 0.99

$$\int x^2 \arctan(a + bf^{c+dx}) dx$$

$$= \frac{i(d^3 x^3 \log^3(f) \log(1 - ia - ibf^{c+dx}) - d^3 x^3 \log^3(f) \log(1 + ia + ibf^{c+dx}) - d^3 x^3 \log^3(f) \log\left(\frac{i+a+bf^{c+dx}}{i+a}\right))}{1}$$

input `Integrate[x^2*ArcTan[a + b*f^(c + d*x)],x]`

output

```
((I/6)*(d^3*x^3*Log[f]^3*Log[1 - I*a - I*b*f^(c + d*x)] - d^3*x^3*Log[f]^3*Log[1 + I*a + I*b*f^(c + d*x)] - d^3*x^3*Log[f]^3*Log[(I + a + b*f^(c + d*x))/(I + a)] + d^3*x^3*Log[f]^3*Log[1 + (b*f^(c + d*x))/(-I + a)] + 3*d^2*x^2*Log[f]^2*PolyLog[2, (b*f^(c + d*x))/(I - a)] - 3*d^2*x^2*Log[f]^2*PolyLog[2, -((b*f^(c + d*x))/(I + a))] - 6*d*x*Log[f]*PolyLog[3, (b*f^(c + d*x))/(I - a)] + 6*d*x*Log[f]*PolyLog[3, -((b*f^(c + d*x))/(I + a))] + 6*PolyLog[4, (b*f^(c + d*x))/(I - a)] - 6*PolyLog[4, -((b*f^(c + d*x))/(I + a))])/(d^3*Log[f]^3)
```

Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5666, 3012, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \arctan(a + bf^{c+dx}) dx$$

$$\downarrow \text{5666}$$

$$\frac{1}{2}i \int x^2 \log(-ibf^{c+dx} - ia + 1) dx - \frac{1}{2}i \int x^2 \log(ibf^{c+dx} + ia + 1) dx$$

$$\downarrow \text{3012}$$

$$\frac{1}{2}i \left(\int x^2 \log \left(1 - \frac{ibf^{c+dx}}{1-ia} \right) dx + \frac{1}{3}x^3 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\int x^2 \log \left(\frac{ibf^{c+dx}}{ia+1} + 1 \right) dx + \frac{1}{3}x^3 \log(ia + ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 3011

$$\frac{1}{2}i \left(\frac{2 \int x \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right) dx}{d \log(f)} - \frac{x^2 \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} + \frac{1}{3}x^3 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\frac{2 \int x \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right) dx}{d \log(f)} - \frac{x^2 \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} + \frac{1}{3}x^3 \log(ia + ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 7163

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \text{PolyLog} \left(3, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} - \frac{\int \text{PolyLog} \left(3, -\frac{bf^{c+dx}}{a+i} \right) dx}{d \log(f)} \right)}{d \log(f)} - \frac{x^2 \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} + \frac{1}{3}x^3 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\frac{2 \left(\frac{x \text{PolyLog} \left(3, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} - \frac{\int \text{PolyLog} \left(3, \frac{bf^{c+dx}}{i-a} \right) dx}{d \log(f)} \right)}{d \log(f)} - \frac{x^2 \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} + \frac{1}{3}x^3 \log(ia + ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 2720

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \text{PolyLog} \left(3, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} - \frac{\int f^{-c-dx} \text{PolyLog} \left(3, -\frac{bf^{c+dx}}{a+i} \right) df^{c+dx}}{d^2 \log^2(f)} \right)}{d \log(f)} - \frac{x^2 \text{PolyLog} \left(2, -\frac{bf^{c+dx}}{a+i} \right)}{d \log(f)} + \frac{1}{3}x^3 \log(-ia - ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 + \frac{bf^{c+dx}}{a+i} \right) \right) - \frac{1}{2}i \left(\frac{2 \left(\frac{x \text{PolyLog} \left(3, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} - \frac{\int f^{-c-dx} \text{PolyLog} \left(3, \frac{bf^{c+dx}}{i-a} \right) df^{c+dx}}{d^2 \log^2(f)} \right)}{d \log(f)} - \frac{x^2 \text{PolyLog} \left(2, \frac{bf^{c+dx}}{i-a} \right)}{d \log(f)} + \frac{1}{3}x^3 \log(ia + ibf^{c+dx} + 1) - \frac{1}{3}x^3 \log \left(1 - \frac{bf^{c+dx}}{-a+i} \right) \right)$$

↓ 7143

$$\frac{1}{2}i \left(\frac{2 \left(\frac{x \operatorname{PolyLog}\left(3, -\frac{bf^{c+dx}}{a+i}\right)}{d \log(f)} - \frac{\operatorname{PolyLog}\left(4, -\frac{bf^{c+dx}}{a+i}\right)}{d^2 \log^2(f)} \right)}{d \log(f)} - \frac{x^2 \operatorname{PolyLog}\left(2, -\frac{bf^{c+dx}}{a+i}\right)}{d \log(f)} + \frac{1}{3}x^3 \log\left(-ia - ibf^{c+dx} + 1\right) \right) \\ \frac{1}{2}i \left(\frac{2 \left(\frac{x \operatorname{PolyLog}\left(3, \frac{bf^{c+dx}}{i-a}\right)}{d \log(f)} - \frac{\operatorname{PolyLog}\left(4, \frac{bf^{c+dx}}{i-a}\right)}{d^2 \log^2(f)} \right)}{d \log(f)} - \frac{x^2 \operatorname{PolyLog}\left(2, \frac{bf^{c+dx}}{i-a}\right)}{d \log(f)} + \frac{1}{3}x^3 \log\left(ia + ibf^{c+dx} + 1\right) - \frac{1}{3}x^3 \right)$$

input `Int[x^2*ArcTan[a + b*f^(c + d*x)],x]`

output `(-1/2*I)*((x^3*Log[1 + I*a + I*b*f^(c + d*x)]/3 - (x^3*Log[1 - (b*f^(c + d*x))/(I - a)]/3 - (x^2*PolyLog[2, (b*f^(c + d*x))/(I - a)]/(d*Log[f]) + (2*((x*PolyLog[3, (b*f^(c + d*x))/(I - a)]/(d*Log[f]) - PolyLog[4, (b*f^(c + d*x))/(I - a)]/(d^2*Log[f]^2)))/(d*Log[f])) + (I/2)*((x^3*Log[1 - I*a - I*b*f^(c + d*x)]/3 - (x^3*Log[1 + (b*f^(c + d*x))/(I + a)]/3 - (x^2*PolyLog[2, -((b*f^(c + d*x))/(I + a))]/(d*Log[f]) + (2*((x*PolyLog[3, -((b*f^(c + d*x))/(I + a))]/(d*Log[f]) - PolyLog[4, -((b*f^(c + d*x))/(I + a))]/(d^2*Log[f]^2)))/(d*Log[f]))`

Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

rule 3012

```
Int[Log[(d_) + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g
_.)*(x_))^(m_.), x_Symbol] := Simp[(f + g*x)^(m + 1)*(Log[d + e*(F^(c*(a +
b*x)))^n]/(g*(m + 1))), x] + (Int[(f + g*x)^m*Log[1 + (e/d)*(F^(c*(a + b*x)
))^n], x] - Simp[(f + g*x)^(m + 1)*(Log[1 + (e/d)*(F^(c*(a + b*x)))^n]/(g*(
m + 1))), x]) /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && NeQ[
d, 1]
```

rule 5666

```
Int[ArcTan[(a_.) + (b_.)*(f_)^((c_.) + (d_.)*(x_))]*(x_)^(m_.), x_Symbol] :
> Simp[I/2 Int[x^m*Log[1 - I*a - I*b*f^(c + d*x)], x], x] - Simp[I/2 In
t[x^m*Log[1 + I*a + I*b*f^(c + d*x)], x], x] /; FreeQ[{a, b, c, d, f}, x] &
& IntegerQ[m] && m > 0
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

rule 7163

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 757 vs. $2(268) = 536$.

Time = 1.11 (sec) , antiderivative size = 758, normalized size of antiderivative = 2.51

method	result
risch	$\frac{i \operatorname{polylog}\left(2, \frac{ib f^{dx} f^c}{-ia+1}\right) c^2}{2d^3 \ln(f)} + \frac{i \operatorname{polylog}\left(2, \frac{ib f^{dx} f^c}{-ia-1}\right) x^2}{2d \ln(f)} + \frac{i \operatorname{polylog}\left(4, \frac{ib f^{dx} f^c}{-ia-1}\right)}{d^3 \ln(f)^3} - \frac{ic^2 \ln\left(\frac{b f^{dx} f^c + a + i}{a + i}\right) x}{2d^2} - \frac{i \ln\left(1 - \frac{ib f^{dx} f^c}{-ia+1}\right)}{6}$

input

```
int(x^2*arctan(a+b*f^(d*x+c)),x,method=_RETURNVERBOSE)
```

output

```
1/2*I/d^3/ln(f)*polylog(2,I*b/(1-I*a)*f^(d*x)*f^c)*c^2+1/2*I/d/ln(f)*polylog(2,I*b/(-I*a-1)*f^(d*x)*f^c)*x^2+I/d^3/ln(f)^3*polylog(4,I*b/(-I*a-1)*f^(d*x)*f^c)-1/2*I/d^2*c^2*ln((b*f^(d*x)*f^c+a+I)/(a+I))*x-1/6*I*ln(1-I*b/(1-I*a)*f^(d*x)*f^c)*x^3-1/2*I/d^2*ln(1-I*b/(-I*a-1)*f^(d*x)*f^c)*x*c^2+1/2*I/d^2*ln(1-I*b/(1-I*a)*f^(d*x)*f^c)*x*c^2+I/d^2/ln(f)^2*polylog(3,I*b/(1-I*a)*f^(d*x)*f^c)*x-1/2*I/d^3/ln(f)*polylog(2,I*b/(-I*a-1)*f^(d*x)*f^c)*c^2-1/2*I/d/ln(f)*polylog(2,I*b/(1-I*a)*f^(d*x)*f^c)*x^2+1/6*I*ln(1-I*b/(-I*a-1)*f^(d*x)*f^c)*x^3-I/d^2/ln(f)^2*polylog(3,I*b/(-I*a-1)*f^(d*x)*f^c)*x-1/2*I/d^3/ln(f)*c^2*dilog((b*f^(d*x)*f^c+a+I)/(a+I))-I/d^3/ln(f)^3*polylog(4,I*b/(1-I*a)*f^(d*x)*f^c)+1/2*I/d^3/ln(f)*c^2*dilog((b*f^(d*x)*f^c+a-I)/(a-I))-1/2*I/d^3*c^3*ln((b*f^(d*x)*f^c+a+I)/(a+I))+1/6*I*x^3*ln(1-I*(a+b*f^(d*x+c)))-1/6*I/d^3*c^3*ln(I*f^(d*x)*f^c*b+I*a+1)+1/2*I/d^2*c^2*ln((b*f^(d*x)*f^c+a-I)/(a-I))*x-1/3*I/d^3*ln(1-I*b/(-I*a-1)*f^(d*x)*f^c)*c^3-1/6*I*x^3*ln(1+I*(a+b*f^(d*x+c)))+1/3*I/d^3*ln(1-I*b/(1-I*a)*f^(d*x)*f^c)*c^3+1/2*I/d^3*c^3*ln((b*f^(d*x)*f^c+a-I)/(a-I))+1/6*I/d^3*c^3*ln(1-I*a-I*f^(d*x)*f^c*b)
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 378, normalized size of antiderivative = 1.25

$$\int x^2 \arctan(a + b f^{c+dx}) dx$$

$$= \frac{2 d^3 x^3 \arctan(b f^{dx+c} + a) \log(f)^3 + 3i d^2 x^2 \operatorname{Li}_2\left(-\frac{a^2+(ab+ib)f^{dx+c}+1}{a^2+1} + 1\right) \log(f)^2 - 3i d^2 x^2 \operatorname{Li}_2\left(-\frac{a^2+(ab-ib)f^{dx+c}+1}{a^2+1} + 1\right) \log(f)^2}{6}$$

input `integrate(x^2*arctan(a+b*f^(d*x+c)),x, algorithm="fricas")`

output `1/6*(2*d^3*x^3*arctan(b*f^(d*x + c) + a)*log(f)^3 + 3*I*d^2*x^2*dilog(-(a^2 + (a*b + I*b)*f^(d*x + c) + 1)/(a^2 + 1) + 1)*log(f)^2 - 3*I*d^2*x^2*dilog(-(a^2 + (a*b - I*b)*f^(d*x + c) + 1)/(a^2 + 1) + 1)*log(f)^2 + I*c^3*log(b*f^(d*x + c) + a + I)*log(f)^3 - I*c^3*log(b*f^(d*x + c) + a - I)*log(f)^3 + (I*d^3*x^3 + I*c^3)*log(f)^3*log((a^2 + (a*b + I*b)*f^(d*x + c) + 1)/(a^2 + 1)) + (-I*d^3*x^3 - I*c^3)*log(f)^3*log((a^2 + (a*b - I*b)*f^(d*x + c) + 1)/(a^2 + 1)) - 6*I*d*x*log(f)*polylog(3, -(a*b + I*b)*f^(d*x + c)/(a^2 + 1)) + 6*I*d*x*log(f)*polylog(3, -(a*b - I*b)*f^(d*x + c)/(a^2 + 1)) + 6*I*polylog(4, -(a*b + I*b)*f^(d*x + c)/(a^2 + 1)) - 6*I*polylog(4, -(a*b - I*b)*f^(d*x + c)/(a^2 + 1)))/(d^3*log(f)^3)`

Sympy [F]

$$\int x^2 \arctan(a + bf^{c+dx}) dx = \int x^2 \operatorname{atan}(a + bf^{c+dx}) dx$$

input `integrate(x**2*atan(a+b*f**(d*x+c)),x)`

output `Integral(x**2*atan(a + b*f**(c + d*x)), x)`

Maxima [F]

$$\int x^2 \arctan(a + bf^{c+dx}) dx = \int x^2 \arctan(bf^{dx+c} + a) dx$$

input `integrate(x^2*arctan(a+b*f^(d*x+c)),x, algorithm="maxima")`

output `-b*d*f^c*integrate(1/3*f^(d*x)*x^3/(b^2*f^(2*d*x)*f^(2*c) + 2*a*b*f^(d*x)*f^c + a^2 + 1), x)*log(f) + 1/3*x^3*arctan(b*f^(d*x)*f^c + a)`

Giac [F]

$$\int x^2 \arctan(a + bf^{c+dx}) dx = \int x^2 \arctan(bf^{dx+c} + a) dx$$

input `integrate(x^2*arctan(a+b*f^(d*x+c)),x, algorithm="giac")`

output `integrate(x^2*arctan(b*f^(d*x + c) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \arctan(a + bf^{c+dx}) dx = \int x^2 \operatorname{atan}(a + bf^{c+dx}) dx$$

input `int(x^2*atan(a + b*f^(c + d*x)),x)`

output `int(x^2*atan(a + b*f^(c + d*x)), x)`

Reduce [F]

$$\int x^2 \arctan(a + bf^{c+dx}) dx = \int \operatorname{atan}(f^{dx+c}b + a) x^2 dx$$

input `int(x^2*atan(a+b*f^(d*x+c)),x)`

output `int(atan(f**(c + d*x)*b + a)*x**2,x)`

3.119 $\int e^{-x} \arctan(e^x) dx$

Optimal result	909
Mathematica [A] (verified)	909
Rubi [A] (verified)	910
Maple [A] (verified)	912
Fricas [A] (verification not implemented)	912
Sympy [A] (verification not implemented)	912
Maxima [A] (verification not implemented)	913
Giac [A] (verification not implemented)	913
Mupad [B] (verification not implemented)	913
Reduce [B] (verification not implemented)	914

Optimal result

Integrand size = 10, antiderivative size = 25

$$\int e^{-x} \arctan(e^x) dx = x - e^{-x} \arctan(e^x) - \frac{1}{2} \log(1 + e^{2x})$$

output `x-arctan(exp(x))/exp(x)-1/2*ln(1+exp(2*x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int e^{-x} \arctan(e^x) dx = x - e^{-x} \arctan(e^x) - \frac{1}{2} \log(1 + e^{2x})$$

input `Integrate[ArcTan[E^x]/E^x,x]`

output `x - ArcTan[E^x]/E^x - Log[1 + E^(2*x)]/2`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.32, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5730, 25, 2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{-x} \arctan(e^x) dx \\
 & \quad \downarrow \text{5730} \\
 & - \int -\frac{1}{1+e^{2x}} dx - e^{-x} \arctan(e^x) \\
 & \quad \downarrow \text{25} \\
 & \int \frac{1}{1+e^{2x}} dx - e^{-x} \arctan(e^x) \\
 & \quad \downarrow \text{2720} \\
 & \frac{1}{2} \int \frac{e^{-2x}}{1+e^{2x}} de^{2x} - e^{-x} \arctan(e^x) \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\int e^{-2x} de^{2x} - \int \frac{1}{1+e^{2x}} de^{2x} \right) - e^{-x} \arctan(e^x) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(\log(e^{2x}) - \int \frac{1}{1+e^{2x}} de^{2x} \right) - e^{-x} \arctan(e^x) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(e^{2x}) - \log(e^{2x} + 1)) - e^{-x} \arctan(e^x)
 \end{aligned}$$

input

Int[ArcTan[E^x]/E^x, x]

output

-(ArcTan[E^x]/E^x) + (Log[E^(2*x)] - Log[1 + E^(2*x)])/2

Definitions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 5730 `Int[((a_.) + ArcTan[u_]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

method	result	size
derivativedivides	$-\arctan(e^x)e^{-x} + \ln(e^x) - \frac{\ln(1+e^{2x})}{2}$	23
default	$-\arctan(e^x)e^{-x} + \ln(e^x) - \frac{\ln(1+e^{2x})}{2}$	23
parallelrisch	$\frac{(-\ln(1+e^{2x})e^x + 2xe^x - 2\arctan(e^x))e^{-x}}{2}$	29
risch	$\frac{ie^{-x}\ln(1+ie^x)}{2} - \frac{\ln(1+e^{2x})}{2} + x - \frac{ie^{-x}\ln(1-ie^x)}{2}$	42

input `int(arctan(exp(x))/exp(x),x,method=_RETURNVERBOSE)`output `-arctan(exp(x))/exp(x)+ln(exp(x))-1/2*ln(exp(x)^2+1)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.12

$$\int e^{-x} \arctan(e^x) dx = \frac{1}{2} (2xe^x - e^x \log(e^{2x} + 1) - 2 \arctan(e^x))e^{(-x)}$$

input `integrate(arctan(exp(x))/exp(x),x, algorithm="fricas")`output `1/2*(2*x*e^x - e^x*log(e^(2*x) + 1) - 2*arctan(e^x))*e^(-x)`**Sympy [A] (verification not implemented)**

Time = 1.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int e^{-x} \arctan(e^x) dx = x - \frac{\log(e^{2x} + 1)}{2} - e^{-x} \operatorname{atan}(e^x)$$

input `integrate(atan(exp(x))/exp(x),x)`

output `x - log(exp(2*x) + 1)/2 - exp(-x)*atan(exp(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int e^{-x} \arctan(e^x) dx = -\arctan(e^x) e^{(-x)} - \frac{1}{2} \log(e^{(-2x)} + 1)$$

input `integrate(arctan(exp(x))/exp(x),x, algorithm="maxima")`

output `-arctan(e^x)*e^(-x) - 1/2*log(e^(-2*x) + 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int e^{-x} \arctan(e^x) dx = -\arctan(e^x) e^{(-x)} + x - \frac{1}{2} \log(e^{(2x)} + 1)$$

input `integrate(arctan(exp(x))/exp(x),x, algorithm="giac")`

output `-arctan(e^x)*e^(-x) + x - 1/2*log(e^(2*x) + 1)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int e^{-x} \arctan(e^x) dx = x - \frac{\ln(e^{2x} + 1)}{2} - \operatorname{atan}(e^x) e^{-x}$$

input `int(atan(exp(x))*exp(-x),x)`

output `x - log(exp(2*x) + 1)/2 - atan(exp(x))*exp(-x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.32

$$\int e^{-x} \arctan(e^x) dx = \frac{-2\operatorname{atan}(e^x) - e^x \log(e^{2x} + 1) + 2e^x x}{2e^x}$$

input `int(atan(exp(x))/exp(x),x)`

output `(- 2*atan(e**x) - e**x*log(e**(2*x) + 1) + 2*e**x*x)/(2*e**x)`

3.120 $\int \frac{\arctan(x)}{(-1+x)^3} dx$

Optimal result	915
Mathematica [A] (verified)	915
Rubi [A] (verified)	916
Maple [A] (verified)	917
Fricas [A] (verification not implemented)	918
Sympy [B] (verification not implemented)	918
Maxima [A] (verification not implemented)	919
Giac [A] (verification not implemented)	919
Mupad [B] (verification not implemented)	919
Reduce [B] (verification not implemented)	920

Optimal result

Integrand size = 8, antiderivative size = 45

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = \frac{1}{4(1-x)} - \frac{\arctan(x)}{2(1-x)^2} - \frac{1}{4} \log(1-x) + \frac{1}{8} \log(1+x^2)$$

output

```
1/(4-4*x)-1/2*arctan(x)/(1-x)^2-1/4*ln(1-x)+1/8*ln(x^2+1)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = \frac{1}{8} \left(-\frac{2}{-1+x} - \frac{4 \arctan(x)}{(-1+x)^2} - 2 \log(1-x) + \log(1+x^2) \right)$$

input

```
Integrate[ArcTan[x]/(-1+x)^3,x]
```

output

```
(-2/(-1+x) - (4*ArcTan[x])/(-1+x)^2 - 2*Log[1-x] + Log[1+x^2])/8
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.18, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5387, 480, 657, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\arctan(x)}{(x-1)^3} dx \\
 & \quad \downarrow \text{5387} \\
 & \frac{1}{2} \int \frac{1}{(1-x)^2(x^2+1)} dx - \frac{\arctan(x)}{2(1-x)^2} \\
 & \quad \downarrow \text{480} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \frac{x+1}{(1-x)(x^2+1)} dx + \frac{1}{2(1-x)} \right) - \frac{\arctan(x)}{2(1-x)^2} \\
 & \quad \downarrow \text{657} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \left(\frac{x}{x^2+1} + \frac{1}{1-x} \right) dx + \frac{1}{2(1-x)} \right) - \frac{\arctan(x)}{2(1-x)^2} \\
 & \quad \downarrow \text{2009} \\
 & \frac{1}{2} \left(\frac{1}{2} \left(\frac{1}{2} \log(x^2+1) - \log(1-x) \right) + \frac{1}{2(1-x)} \right) - \frac{\arctan(x)}{2(1-x)^2}
 \end{aligned}$$

input `Int[ArcTan[x]/(-1 + x)^3,x]`

output `-1/2*ArcTan[x]/(1 - x)^2 + (1/(2*(1 - x))) + (-Log[1 - x] + Log[1 + x^2])/2`
`/2)/2`

Definitions of rubi rules used

rule 480 $\text{Int}[\frac{(c_+) + (d_+)(x_+)^{n_+}}{(a_+) + (b_+)(x_+)^2}, x_Symbol] \rightarrow \text{Simp}[d*((c + d*x)^{n+1}/((n+1)*(b*c^2 + a*d^2))), x] + \text{Simp}[b/(b*c^2 + a*d^2) \text{Int}[(c + d*x)^{n+1}*((c - d*x)/(a + b*x^2)), x], x] /;$ $\text{FreeQ}\{a, b, c, d, x\}$ && $\text{ILtQ}[n, -1]$

rule 657 $\text{Int}[\frac{((d_+) + (e_+)(x_+)^{m_+})*((f_+) + (g_+)(x_+)^{n_+})}{(a_+) + (c_+)(x_+)^2}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d + e*x)^m*((f + g*x)^n/(a + c*x^2)), x], x] /;$ $\text{FreeQ}\{a, c, d, e, f, g, m\}, x\}$ && $\text{IntegersQ}[n]$

rule 2009 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /;$ $\text{SumQ}[u]$

rule 5387 $\text{Int}[\frac{(a_+) + \text{ArcTan}[c_+](x_+)]*(b_+)*((d_+) + (e_+)(x_+)^{q_+}), x_Symbol] \rightarrow \text{Simp}[(d + e*x)^{q+1}*((a + b*\text{ArcTan}[c*x])/(e*(q+1))), x] - \text{Simp}[b*(c/(e*(q+1))) \text{Int}[(d + e*x)^{q+1}/(1 + c^2*x^2), x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, q\}, x\}$ && $\text{NeQ}[q, -1]$

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.71

method	result
default	$-\frac{\arctan(x)}{2(-1+x)^2} - \frac{1}{4(-1+x)} - \frac{\ln(-1+x)}{4} + \frac{\ln(x^2+1)}{8}$
parts	$-\frac{\arctan(x)}{2(-1+x)^2} - \frac{1}{4(-1+x)} - \frac{\ln(-1+x)}{4} + \frac{\ln(x^2+1)}{8}$
paralelrisch	$-\frac{2 \ln(-1+x)x^2 - \ln(x^2+1)x^2 - 2 - 4 \ln(-1+x)x + 2 \ln(x^2+1)x + 2 \ln(-1+x) - \ln(x^2+1) + 2x + 4 \arctan(x)}{8(-1+x)^2}$
risch	$\frac{i \ln(ix+1)}{4(-1+x)^2} - \frac{i(-2i \ln(-1+x)x^2 + i \ln(x^2+1)x^2 + 4i \ln(-1+x)x - 2i \ln(x^2+1)x - 2i \ln(-1+x) + i \ln(x^2+1) - 2ix + 2i + 2 \ln(-1+x))}{8(-1+x)^2}$

input `int(arctan(x)/(-1+x)^3,x,method=_RETURNVERBOSE)`

output $-1/2/(-1+x)^2*\arctan(x)-1/4/(-1+x)-1/4*\ln(-1+x)+1/8*\ln(x^2+1)$

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = \frac{(x^2 - 2x + 1) \log(x^2 + 1) - 2(x^2 - 2x + 1) \log(x - 1) - 2x - 4 \arctan(x) + 2}{8(x^2 - 2x + 1)}$$

input `integrate(arctan(x)/(-1+x)^3,x, algorithm="fricas")`

output `1/8*((x^2 - 2*x + 1)*log(x^2 + 1) - 2*(x^2 - 2*x + 1)*log(x - 1) - 2*x - 4*arctan(x) + 2)/(x^2 - 2*x + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(31) = 62$.

Time = 0.30 (sec) , antiderivative size = 153, normalized size of antiderivative = 3.40

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = -\frac{2x^2 \log(x-1)}{8x^2 - 16x + 8} + \frac{x^2 \log(x^2+1)}{8x^2 - 16x + 8} + \frac{4x \log(x-1)}{8x^2 - 16x + 8} - \frac{2x \log(x^2+1)}{8x^2 - 16x + 8} - \frac{2x}{8x^2 - 16x + 8} - \frac{2 \log(x-1)}{8x^2 - 16x + 8} + \frac{\log(x^2+1)}{8x^2 - 16x + 8} - \frac{4 \operatorname{atan}(x)}{8x^2 - 16x + 8} + \frac{2}{8x^2 - 16x + 8}$$

input `integrate(atan(x)/(-1+x)**3,x)`

output `-2*x**2*log(x - 1)/(8*x**2 - 16*x + 8) + x**2*log(x**2 + 1)/(8*x**2 - 16*x + 8) + 4*x*log(x - 1)/(8*x**2 - 16*x + 8) - 2*x*log(x**2 + 1)/(8*x**2 - 16*x + 8) - 2*x/(8*x**2 - 16*x + 8) - 2*log(x - 1)/(8*x**2 - 16*x + 8) + log(x**2 + 1)/(8*x**2 - 16*x + 8) - 4*atan(x)/(8*x**2 - 16*x + 8) + 2/(8*x**2 - 16*x + 8)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.69

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = -\frac{1}{4(x-1)} - \frac{\arctan(x)}{2(x-1)^2} + \frac{1}{8} \log(x^2+1) - \frac{1}{4} \log(x-1)$$

input `integrate(arctan(x)/(-1+x)^3,x, algorithm="maxima")`output `-1/4/(x - 1) - 1/2*arctan(x)/(x - 1)^2 + 1/8*log(x^2 + 1) - 1/4*log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.71

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = -\frac{1}{4(x-1)} - \frac{\arctan(x)}{2(x-1)^2} + \frac{1}{8} \log(x^2+1) - \frac{1}{4} \log(|x-1|)$$

input `integrate(arctan(x)/(-1+x)^3,x, algorithm="giac")`output `-1/4/(x - 1) - 1/2*arctan(x)/(x - 1)^2 + 1/8*log(x^2 + 1) - 1/4*log(abs(x - 1))`**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.69

$$\int \frac{\arctan(x)}{(-1+x)^3} dx = \frac{\ln(x^2+1)}{8} - \frac{\ln(x-1)}{4} - \frac{\frac{x}{4} + \frac{\operatorname{atan}(x)}{2} - \frac{1}{4}}{(x-1)^2}$$

input `int(atan(x)/(x - 1)^3,x)`output `log(x^2 + 1)/8 - log(x - 1)/4 - (x/4 + atan(x)/2 - 1/4)/(x - 1)^2`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.58

$$\int \frac{\arctan(x)}{(-1+x)^3} dx$$

$$= \frac{-4\operatorname{atan}(x) + \log(x^2 + 1)x^2 - 2\log(x^2 + 1)x + \log(x^2 + 1) - 2\log(x - 1)x^2 + 4\log(x - 1)x - 2\log(x - 1)}{8x^2 - 16x + 8}$$

input

```
int(atan(x)/(-1+x)^3,x)
```

output

```
( - 4*atan(x) + log(x**2 + 1)*x**2 - 2*log(x**2 + 1)*x + log(x**2 + 1) - 2
*log(x - 1)*x**2 + 4*log(x - 1)*x - 2*log(x - 1) - x**2 + 1)/(8*(x**2 - 2*
x + 1))
```

3.121 $\int \frac{\arctan(1+2x)}{(4+3x)^3} dx$

Optimal result	921
Mathematica [C] (verified)	921
Rubi [A] (verified)	922
Maple [A] (verified)	924
Fricas [A] (verification not implemented)	924
Sympy [B] (verification not implemented)	925
Maxima [A] (verification not implemented)	926
Giac [B] (verification not implemented)	926
Mupad [B] (verification not implemented)	927
Reduce [B] (verification not implemented)	928

Optimal result

Integrand size = 14, antiderivative size = 64

$$\int \frac{\arctan(1 + 2x)}{(4 + 3x)^3} dx = -\frac{1}{34(4 + 3x)} + \frac{8}{867} \arctan(1 + 2x) - \frac{\arctan(1 + 2x)}{6(4 + 3x)^2} + \frac{5}{289} \log(4 + 3x) - \frac{5}{578} \log(1 + 2x + 2x^2)$$

output

```
-1/34/(4+3*x)+8/867*arctan(1+2*x)-1/6*arctan(1+2*x)/(4+3*x)^2+5/289*ln(4+3*x)-5/578*ln(2*x^2+2*x+1)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.03 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.27

$$\int \frac{\arctan(1 + 2x)}{(4 + 3x)^3} dx = \frac{-289 \arctan(1 + 2x) + (4 + 3x)(-51 - (15 - 8i)(4 + 3x) \log(i + (1 + i)x) - (15 + 8i)(4 + 3x) \log(1 + 2x))}{1734(4 + 3x)^2}$$

input

```
Integrate[ArcTan[1 + 2*x]/(4 + 3*x)^3,x]
```

output

```
(-289*ArcTan[1 + 2*x] + (4 + 3*x)*(-51 - (15 - 8*I)*(4 + 3*x)*Log[I + (1 + I)*x] - (15 + 8*I)*(4 + 3*x)*Log[1 + (1 + I)*x] + 120*Log[4 + 3*x] + 90*x*Log[4 + 3*x]))/(1734*(4 + 3*x)^2)
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.16, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5568, 2081, 1145, 27, 1200, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(2x+1)}{(3x+4)^3} dx$$

$$\downarrow 5568$$

$$\frac{1}{3} \int \frac{1}{(3x+4)^2 ((2x+1)^2+1)} dx - \frac{\arctan(2x+1)}{6(3x+4)^2}$$

$$\downarrow 2081$$

$$\frac{1}{3} \int \frac{1}{(3x+4)^2 (4x^2+4x+2)} dx - \frac{\arctan(2x+1)}{6(3x+4)^2}$$

$$\downarrow 1145$$

$$\frac{1}{3} \left(\frac{1}{34} \int \frac{2(1-3x)}{(3x+4)(2x^2+2x+1)} dx - \frac{3}{34(3x+4)} \right) - \frac{\arctan(2x+1)}{6(3x+4)^2}$$

$$\downarrow 27$$

$$\frac{1}{3} \left(\frac{1}{17} \int \frac{1-3x}{(3x+4)(2x^2+2x+1)} dx - \frac{3}{34(3x+4)} \right) - \frac{\arctan(2x+1)}{6(3x+4)^2}$$

$$\downarrow 1200$$

$$\frac{1}{3} \left(\frac{1}{17} \int \left(\frac{-30x-7}{17(2x^2+2x+1)} + \frac{45}{17(3x+4)} \right) dx - \frac{3}{34(3x+4)} \right) - \frac{\arctan(2x+1)}{6(3x+4)^2}$$

$$\downarrow 2009$$

$$\frac{1}{3} \left(\frac{1}{17} \left(\frac{8}{17} \arctan(2x+1) - \frac{15}{34} \log(2x^2+2x+1) + \frac{15}{17} \log(3x+4) \right) - \frac{3}{34(3x+4)} \right) - \frac{\arctan(2x+1)}{6(3x+4)^2}$$

input `Int[ArcTan[1 + 2*x]/(4 + 3*x)^3,x]`

output `-1/6*ArcTan[1 + 2*x]/(4 + 3*x)^2 + (-3/(34*(4 + 3*x)) + ((8*ArcTan[1 + 2*x])/17 + (15*Log[4 + 3*x])/17 - (15*Log[1 + 2*x + 2*x^2])/34)/17)/3`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1145 `Int[((d_.) + (e_.)*(x_))^(m_)/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[e*((d + e*x)^(m + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))], x] + Simp[1/(c*d^2 - b*d*e + a*e^2) Int[(d + e*x)^(m + 1)*(Simp[c*d - b*e - c*e*x, x]/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[m, -1]`

rule 1200 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)^n/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && IntegerQ[n]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2081 `Int[(u_)^(m_.)*(v_)^(p_.), x_Symbol] := Int[ExpandToSum[u, x]^m*ExpandToSum[v, x]^p, x] /; FreeQ[{m, p}, x] && LinearQ[u, x] && QuadraticQ[v, x] && ! (LinearMatchQ[u, x] && QuadraticMatchQ[v, x])`

rule 5568

```
Int[((a_.) + ArcTan[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_), x_Symbol] := Simp[(e + f*x)^(m + 1)*((a + b*ArcTan[c + d*x])^p/(f*(m +
1))), x] - Simp[b*d*(p/(f*(m + 1))) Int[(e + f*x)^(m + 1)*((a + b*ArcTan[
c + d*x])^(p - 1)/(1 + (c + d*x)^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x
] && IGtQ[p, 0] && ILtQ[m, -1]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.84

method	result
derivativdivides	$-\frac{2 \arctan(1+2x)}{3(8+6x)^2} - \frac{1}{17(8+6x)} + \frac{5 \ln(8+6x)}{289} - \frac{5 \ln((1+2x)^2+1)}{578} + \frac{8 \arctan(1+2x)}{867}$
default	$-\frac{2 \arctan(1+2x)}{3(8+6x)^2} - \frac{1}{17(8+6x)} + \frac{5 \ln(8+6x)}{289} - \frac{5 \ln((1+2x)^2+1)}{578} + \frac{8 \arctan(1+2x)}{867}$
parts	$-\frac{1}{34(4+3x)} + \frac{8 \arctan(1+2x)}{867} - \frac{\arctan(1+2x)}{6(4+3x)^2} + \frac{5 \ln(4+3x)}{289} - \frac{5 \ln(2x^2+2x+1)}{578}$
parallelrisc	$\frac{810 \ln(\frac{4}{3}+x)x^2 - 405 \ln(x^2+x+\frac{1}{2})x^2 + 432 \arctan(1+2x)x^2 - 612 + 2160 \ln(\frac{4}{3}+x)x - 1080 \ln(x^2+x+\frac{1}{2})x + 1152 \arctan(1+2x)x - 5202(4+3x)^2}{5202(4+3x)^2}$
risc	$\frac{i \ln(1+i(1+2x))}{12(4+3x)^2} - \frac{i(960i \ln(4+3x) - 270i \ln(2x+1-i)x^2 - 720i \ln(2x+1-i)x + 1440i \ln(4+3x)x - 144 \ln(2x+1+i)x^2 - 144 \ln(2x+1+i)x)}{12(4+3x)^2}$

input

```
int(arctan(1+2*x)/(4+3*x)^3,x,method=_RETURNVERBOSE)
```

output

```
-2/3/(8+6*x)^2*arctan(1+2*x)-1/17/(8+6*x)+5/289*ln(8+6*x)-5/578*ln((1+2*x)
^2+1)+8/867*arctan(1+2*x)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.20

$$\int \frac{\arctan(1 + 2x)}{(4 + 3x)^3} dx$$

$$= \frac{(48x^2 + 128x - 11) \arctan(2x + 1) - 5(9x^2 + 24x + 16) \log(2x^2 + 2x + 1) + 10(9x^2 + 24x + 16) \ln(4 + 3x)}{578(9x^2 + 24x + 16)}$$

input

```
integrate(arctan(1+2*x)/(4+3*x)^3,x, algorithm="fricas")
```

output

```
1/578*((48*x^2 + 128*x - 11)*arctan(2*x + 1) - 5*(9*x^2 + 24*x + 16)*log(2
*x^2 + 2*x + 1) + 10*(9*x^2 + 24*x + 16)*log(3*x + 4) - 51*x - 68)/(9*x^2
+ 24*x + 16)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 223 vs. $2(56) = 112$.

Time = 0.32 (sec) , antiderivative size = 223, normalized size of antiderivative = 3.48

$$\int \frac{\arctan(1+2x)}{(4+3x)^3} dx = \frac{90x^2 \log(3x+4)}{5202x^2 + 13872x + 9248} - \frac{45x^2 \log(2x^2 + 2x + 1)}{5202x^2 + 13872x + 9248}$$

$$+ \frac{48x^2 \operatorname{atan}(2x+1)}{5202x^2 + 13872x + 9248} + \frac{240x \log(3x+4)}{5202x^2 + 13872x + 9248}$$

$$- \frac{120x \log(2x^2 + 2x + 1)}{5202x^2 + 13872x + 9248} + \frac{128x \operatorname{atan}(2x+1)}{5202x^2 + 13872x + 9248}$$

$$- \frac{51x}{5202x^2 + 13872x + 9248} + \frac{160 \log(3x+4)}{5202x^2 + 13872x + 9248}$$

$$- \frac{80 \log(2x^2 + 2x + 1)}{5202x^2 + 13872x + 9248} - \frac{11 \operatorname{atan}(2x+1)}{5202x^2 + 13872x + 9248}$$

$$- \frac{68}{5202x^2 + 13872x + 9248}$$

input

```
integrate(atan(1+2*x)/(4+3*x)**3,x)
```

output

```
90*x**2*log(3*x + 4)/(5202*x**2 + 13872*x + 9248) - 45*x**2*log(2*x**2 + 2
*x + 1)/(5202*x**2 + 13872*x + 9248) + 48*x**2*atan(2*x + 1)/(5202*x**2 +
13872*x + 9248) + 240*x*log(3*x + 4)/(5202*x**2 + 13872*x + 9248) - 120*x*
log(2*x**2 + 2*x + 1)/(5202*x**2 + 13872*x + 9248) + 128*x*atan(2*x + 1)/(
5202*x**2 + 13872*x + 9248) - 51*x/(5202*x**2 + 13872*x + 9248) + 160*log(
3*x + 4)/(5202*x**2 + 13872*x + 9248) - 80*log(2*x**2 + 2*x + 1)/(5202*x**
2 + 13872*x + 9248) - 11*atan(2*x + 1)/(5202*x**2 + 13872*x + 9248) - 68/(
5202*x**2 + 13872*x + 9248)
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.84

$$\int \frac{\arctan(1+2x)}{(4+3x)^3} dx = -\frac{1}{34(3x+4)} - \frac{\arctan(2x+1)}{6(3x+4)^2} + \frac{8}{867} \arctan(2x+1) - \frac{5}{578} \log(2x^2+2x+1) + \frac{5}{289} \log(3x+4)$$

input `integrate(arctan(1+2*x)/(4+3*x)^3,x, algorithm="maxima")`

output `-1/34/(3*x + 4) - 1/6*arctan(2*x + 1)/(3*x + 4)^2 + 8/867*arctan(2*x + 1) - 5/578*log(2*x^2 + 2*x + 1) + 5/289*log(3*x + 4)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 657 vs. $2(54) = 108$.

Time = 0.24 (sec) , antiderivative size = 657, normalized size of antiderivative = 10.27

$$\int \frac{\arctan(1+2x)}{(4+3x)^3} dx = \text{Too large to display}$$

input `integrate(arctan(1+2*x)/(4+3*x)^3,x, algorithm="giac")`

output

```
-1/578*(252*arctan(2*x + 1)*tan(1/2*arctan(2*x + 1))^4 - 125*log(2*(25*tan
(1/2*arctan(2*x + 1))^4 - 60*tan(1/2*arctan(2*x + 1))^3 - 14*tan(1/2*arcta
n(2*x + 1))^2 + 60*tan(1/2*arctan(2*x + 1)) + 25)/(tan(1/2*arctan(2*x + 1)
))^4 + 2*tan(1/2*arctan(2*x + 1))^2 + 1))*tan(1/2*arctan(2*x + 1))^4 + 320*
arctan(2*x + 1)*tan(1/2*arctan(2*x + 1))^3 + 300*log(2*(25*tan(1/2*arctan(
2*x + 1))^4 - 60*tan(1/2*arctan(2*x + 1))^3 - 14*tan(1/2*arctan(2*x + 1))^
2 + 60*tan(1/2*arctan(2*x + 1)) + 25)/(tan(1/2*arctan(2*x + 1))^4 + 2*tan(
1/2*arctan(2*x + 1))^2 + 1))*tan(1/2*arctan(2*x + 1))^3 + 45*tan(1/2*arcta
n(2*x + 1))^4 - 696*arctan(2*x + 1)*tan(1/2*arctan(2*x + 1))^2 + 70*log(2*
(25*tan(1/2*arctan(2*x + 1))^4 - 60*tan(1/2*arctan(2*x + 1))^3 - 14*tan(1/
2*arctan(2*x + 1))^2 + 60*tan(1/2*arctan(2*x + 1)) + 25)/(tan(1/2*arctan(2
*x + 1))^4 + 2*tan(1/2*arctan(2*x + 1))^2 + 1))*tan(1/2*arctan(2*x + 1))^2
+ 96*tan(1/2*arctan(2*x + 1))^3 - 320*arctan(2*x + 1)*tan(1/2*arctan(2*x
+ 1)) - 300*log(2*(25*tan(1/2*arctan(2*x + 1))^4 - 60*tan(1/2*arctan(2*x +
1))^3 - 14*tan(1/2*arctan(2*x + 1))^2 + 60*tan(1/2*arctan(2*x + 1)) + 25)
/(tan(1/2*arctan(2*x + 1))^4 + 2*tan(1/2*arctan(2*x + 1))^2 + 1))*tan(1/2*
arctan(2*x + 1)) - 270*tan(1/2*arctan(2*x + 1))^2 + 252*arctan(2*x + 1) -
125*log(2*(25*tan(1/2*arctan(2*x + 1))^4 - 60*tan(1/2*arctan(2*x + 1))^3 -
14*tan(1/2*arctan(2*x + 1))^2 + 60*tan(1/2*arctan(2*x + 1)) + 25)/(tan(1/
2*arctan(2*x + 1))^4 + 2*tan(1/2*arctan(2*x + 1))^2 + 1)) - 96*tan(1/2*...
```

Mupad [B] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.72

$$\int \frac{\arctan(1 + 2x)}{(4 + 3x)^3} dx = \frac{5 \ln(x + \frac{4}{3})}{289} - \frac{5 \ln(x^2 + x + \frac{1}{2})}{578} + \frac{8 \operatorname{atan}(2x + 1)}{867} - \frac{\frac{3x}{34} + \frac{\operatorname{atan}(2x+1)}{6} + \frac{2}{17}}{(3x + 4)^2}$$

input

```
int(atan(2*x + 1)/(3*x + 4)^3,x)
```

output

```
(5*log(x + 4/3))/289 - (5*log(x + x^2 + 1/2))/578 + (8*atan(2*x + 1))/867
- ((3*x)/34 + atan(2*x + 1)/6 + 2/17)/(3*x + 4)^2
```


Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.86

$$\int \frac{\arctan(1+2x)}{(4+3x)^3} dx$$

$$= \frac{384\operatorname{atan}(2x+1)x^2 + 1024\operatorname{atan}(2x+1)x - 88\operatorname{atan}(2x+1) + 720\log(3x+4)x^2 + 1920\log(3x+4)x - 1280\log(3x+4) - 360\log(2x^2+2x+1)x^2 - 960\log(2x^2+2x+1)x - 640\log(2x^2+2x+1) + 153x^2 - 272}{41616x^2 + 12480x + 1280}$$

input

```
int(atan(1+2*x)/(4+3*x)^3,x)
```

output

```
(384*atan(2*x + 1)*x**2 + 1024*atan(2*x + 1)*x - 88*atan(2*x + 1) + 720*log(3*x + 4)*x**2 + 1920*log(3*x + 4)*x + 1280*log(3*x + 4) - 360*log(2*x**2 + 2*x + 1)*x**2 - 960*log(2*x**2 + 2*x + 1)*x - 640*log(2*x**2 + 2*x + 1) + 153*x**2 - 272)/(4624*(9*x**2 + 24*x + 16))
```

3.122 $\int \arctan(\sqrt{1+x}) dx$

Optimal result	929
Mathematica [A] (verified)	929
Rubi [A] (verified)	930
Maple [A] (verified)	931
Fricas [A] (verification not implemented)	932
Sympy [A] (verification not implemented)	932
Maxima [A] (verification not implemented)	933
Giac [A] (verification not implemented)	933
Mupad [B] (verification not implemented)	933
Reduce [B] (verification not implemented)	934

Optimal result

Integrand size = 8, antiderivative size = 30

$$\int \arctan(\sqrt{1+x}) dx = -\sqrt{1+x} + 2 \arctan(\sqrt{1+x}) + x \arctan(\sqrt{1+x})$$

output

```
-(1+x)^(1/2)+2*arctan((1+x)^(1/2))+x*arctan((1+x)^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.73

$$\int \arctan(\sqrt{1+x}) dx = -\sqrt{1+x} + (2+x) \arctan(\sqrt{1+x})$$

input

```
Integrate[ArcTan[Sqrt[1 + x]],x]
```

output

```
-Sqrt[1 + x] + (2 + x)*ArcTan[Sqrt[1 + x]]
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {5726, 90, 27, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \arctan(\sqrt{x+1}) dx \\
 & \quad \downarrow 5726 \\
 & x \arctan(\sqrt{x+1}) - \int \frac{x}{\sqrt{x+1}(2x+4)} dx \\
 & \quad \downarrow 90 \\
 & 2 \int \frac{1}{2\sqrt{x+1}(x+2)} dx + x \arctan(\sqrt{x+1}) - \sqrt{x+1} \\
 & \quad \downarrow 27 \\
 & \int \frac{1}{\sqrt{x+1}(x+2)} dx + x \arctan(\sqrt{x+1}) - \sqrt{x+1} \\
 & \quad \downarrow 73 \\
 & 2 \int \frac{1}{x+2} d\sqrt{x+1} + x \arctan(\sqrt{x+1}) - \sqrt{x+1} \\
 & \quad \downarrow 216 \\
 & x \arctan(\sqrt{x+1}) + 2 \arctan(\sqrt{x+1}) - \sqrt{x+1}
 \end{aligned}$$

input

```
Int[ArcTan[Sqrt[1 + x]], x]
```

output

```
-Sqrt[1 + x] + 2*ArcTan[Sqrt[1 + x]] + x*ArcTan[Sqrt[1 + x]]
```

Definitions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntegerQ[a, b, c, d, m, n, x]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 5726 `Int[ArcTan[u_], x_Symbol] := Simp[x*ArcTan[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/(1 + u^2)), x], x] /; InverseFunctionFreeQ[u, x]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$(1 + x) \arctan(\sqrt{1 + x}) - \sqrt{1 + x} + \arctan(\sqrt{1 + x})$	25
default	$(1 + x) \arctan(\sqrt{1 + x}) - \sqrt{1 + x} + \arctan(\sqrt{1 + x})$	25
parts	$-\sqrt{1 + x} + 2 \arctan(\sqrt{1 + x}) + x \arctan(\sqrt{1 + x})$	25

input `int(arctan((1+x)^(1/2)),x,method=_RETURNVERBOSE)`

output `(1+x)*arctan((1+x)^(1/2))-(1+x)^(1/2)+arctan((1+x)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.60

$$\int \arctan(\sqrt{1+x}) dx = (x+2) \arctan(\sqrt{x+1}) - \sqrt{x+1}$$

input `integrate(arctan((1+x)^(1/2)),x, algorithm="fricas")`

output `(x + 2)*arctan(sqrt(x + 1)) - sqrt(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

$$\int \arctan(\sqrt{1+x}) dx = x \operatorname{atan}(\sqrt{x+1}) - \sqrt{x+1} + 2 \operatorname{atan}(\sqrt{x+1})$$

input `integrate(atan((1+x)**(1/2)),x)`

output `x*atan(sqrt(x + 1)) - sqrt(x + 1) + 2*atan(sqrt(x + 1))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \arctan(\sqrt{1+x}) dx = (x+1) \arctan(\sqrt{x+1}) - \sqrt{x+1} + \arctan(\sqrt{x+1})$$

input `integrate(arctan((1+x)^(1/2)),x, algorithm="maxima")`output `(x + 1)*arctan(sqrt(x + 1)) - sqrt(x + 1) + arctan(sqrt(x + 1))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \arctan(\sqrt{1+x}) dx = (x+1) \arctan(\sqrt{x+1}) - \sqrt{x+1} + \arctan(\sqrt{x+1})$$

input `integrate(arctan((1+x)^(1/2)),x, algorithm="giac")`output `(x + 1)*arctan(sqrt(x + 1)) - sqrt(x + 1) + arctan(sqrt(x + 1))`**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \arctan(\sqrt{1+x}) dx = \operatorname{atan}(\sqrt{x+1}) - \sqrt{x+1} + \operatorname{atan}(\sqrt{x+1})(x+1)$$

input `int(atan((x + 1)^(1/2)),x)`output `atan((x + 1)^(1/2)) - (x + 1)^(1/2) + atan((x + 1)^(1/2))*(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

$$\int \arctan(\sqrt{1+x}) dx = \operatorname{atan}(\sqrt{x+1})x + 2\operatorname{atan}(\sqrt{x+1}) - \sqrt{x+1}$$

input `int(atan((1+x)^(1/2)),x)`

output `atan(sqrt(x + 1))*x + 2*atan(sqrt(x + 1)) - sqrt(x + 1)`

$$3.123 \quad \int \frac{1}{(1+x^2)(2+\arctan(x))} dx$$

Optimal result	935
Mathematica [A] (verified)	935
Rubi [A] (verified)	936
Maple [A] (verified)	936
Fricas [A] (verification not implemented)	937
Sympy [A] (verification not implemented)	937
Maxima [A] (verification not implemented)	938
Giac [A] (verification not implemented)	938
Mupad [B] (verification not implemented)	938
Reduce [B] (verification not implemented)	939

Optimal result

Integrand size = 14, antiderivative size = 5

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(2 + \arctan(x))$$

output `ln(2+arctan(x))`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(2 + \arctan(x))$$

input `Integrate[1/((1 + x^2)*(2 + ArcTan[x])),x]`

output `Log[2 + ArcTan[x]]`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$, Rules used = {5417}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 1)(\arctan(x) + 2)} dx$$

↓ 5417

$$\log(\arctan(x) + 2)$$

input `Int[1/((1 + x^2)*(2 + ArcTan[x])),x]`

output `Log[2 + ArcTan[x]]`

Defintions of rubi rules used

rule 5417 `Int[1/(((a_.) + ArcTan[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)), x_Symbol]
:> Simp[Log[RemoveContent[a + b*ArcTan[c*x], x]]/(b*c*d), x] /; FreeQ[{a,
b, c, d, e}, x] && EqQ[e, c^2*d]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$\ln(2 + \arctan(x))$	6
default	$\ln(2 + \arctan(x))$	6
parallelrisch	$\ln(2 + \arctan(x))$	6
risch	$\ln(-\ln(-ix + 1) + \ln(ix + 1) + 4i)$	21

input `int(1/(x^2+1)/(2+arctan(x)),x,method=_RETURNVERBOSE)`

output `ln(2+arctan(x))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(\arctan(x) + 2)$$

input `integrate(1/(x^2+1)/(2+arctan(x)),x, algorithm="fricas")`

output `log(arctan(x) + 2)`

Sympy [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(\operatorname{atan}(x) + 2)$$

input `integrate(1/(x**2+1)/(2+atan(x)),x)`

output `log(atan(x) + 2)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(\arctan(x) + 2)$$

input `integrate(1/(x^2+1)/(2+arctan(x)),x, algorithm="maxima")`output `log(arctan(x) + 2)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(\arctan(x) + 2)$$

input `integrate(1/(x^2+1)/(2+arctan(x)),x, algorithm="giac")`output `log(arctan(x) + 2)`**Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \ln(\operatorname{atan}(x) + 2)$$

input `int(1/((x^2 + 1)*(atan(x) + 2)),x)`output `log(atan(x) + 2)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1+x^2)(2+\arctan(x))} dx = \log(\operatorname{atan}(x) + 2)$$

input `int(1/(x^2+1)/(2+atan(x)),x)`

output `log(atan(x) + 2)`

3.124 $\int \frac{1}{(a+ax^2)(b-2b \arctan(x))} dx$

Optimal result	940
Mathematica [A] (verified)	940
Rubi [A] (verified)	941
Maple [A] (verified)	941
Fricas [A] (verification not implemented)	942
Sympy [A] (verification not implemented)	942
Maxima [A] (verification not implemented)	943
Giac [A] (verification not implemented)	943
Mupad [B] (verification not implemented)	943
Reduce [B] (verification not implemented)	944

Optimal result

Integrand size = 19, antiderivative size = 17

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(1 - 2 \arctan(x))}{2ab}$$

output

```
-1/2*ln(1-2*arctan(x))/a/b
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(-1 + 2 \arctan(x))}{2ab}$$

input

```
Integrate[1/((a + a*x^2)*(b - 2*b*ArcTan[x])),x]
```

output

```
-1/2*Log[-1 + 2*ArcTan[x]]/(a*b)
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {5417}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(ax^2 + a)(b - 2b \arctan(x))} dx$$

↓ 5417

$$-\frac{\log(1 - 2 \arctan(x))}{2ab}$$

input `Int[1/((a + a*x^2)*(b - 2*b*ArcTan[x])),x]`

output `-1/2*Log[1 - 2*ArcTan[x]]/(a*b)`

Defintions of rubi rules used

rule 5417 `Int[1/(((a_.) + ArcTan[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)), x_Symbol]
:> Simp[Log[RemoveContent[a + b*ArcTan[c*x], x]]/(b*c*d), x] /; FreeQ[{a,
b, c, d, e}, x] && EqQ[e, c^2*d]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
parallelrisch	$-\frac{\ln(\arctan(x) - \frac{1}{2})}{2ba}$	14
default	$-\frac{\ln(2b \arctan(x) - b)}{2ab}$	19
risch	$-\frac{\ln(-i - \ln(-ix+1) + \ln(ix+1))}{2ba}$	29

input `int(1/(a*x^2+a)/(b-2*b*arctan(x)),x,method=_RETURNVERBOSE)`

output `-1/2*ln(arctan(x)-1/2)/b/a`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(2 \arctan(x) - 1)}{2ab}$$

input `integrate(1/(a*x^2+a)/(b-2*b*arctan(x)),x, algorithm="fricas")`

output `-1/2*log(2*arctan(x) - 1)/(a*b)`

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(\arctan(x) - \frac{1}{2})}{2ab}$$

input `integrate(1/(a*x**2+a)/(b-2*b*atan(x)),x)`

output `-log(atan(x) - 1/2)/(2*a*b)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(|2 \arctan(x) - 1|)}{2ab}$$

input `integrate(1/(a*x^2+a)/(b-2*b*arctan(x)),x, algorithm="maxima")`output `-1/2*log(abs(2*arctan(x) - 1))/(a*b)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(|2 \arctan(x) - 1|)}{2ab}$$

input `integrate(1/(a*x^2+a)/(b-2*b*arctan(x)),x, algorithm="giac")`output `-1/2*log(abs(2*arctan(x) - 1))/(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\ln(2 \operatorname{atan}(x) - 1)}{2ab}$$

input `int(1/((a + a*x^2)*(b - 2*b*atan(x))),x)`output `-log(2*atan(x) - 1)/(2*a*b)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + ax^2)(b - 2b \arctan(x))} dx = -\frac{\log(2\operatorname{atan}(x) - 1)}{2ab}$$

input `int(1/(a*x^2+a)/(b-2*b*atan(x)),x)`

output `(- log(2*atan(x) - 1))/(2*a*b)`

$$3.125 \quad \int \frac{x+x^3+(1+x)^2 \arctan(x)}{(1+x)^2(1+x^2)} dx$$

Optimal result	945
Mathematica [A] (verified)	945
Rubi [A] (verified)	946
Maple [A] (verified)	947
Fricas [A] (verification not implemented)	947
Sympy [B] (verification not implemented)	947
Maxima [A] (verification not implemented)	948
Giac [B] (verification not implemented)	948
Mupad [B] (verification not implemented)	949
Reduce [B] (verification not implemented)	949

Optimal result

Integrand size = 26, antiderivative size = 18

$$\int \frac{x+x^3+(1+x)^2 \arctan(x)}{(1+x)^2(1+x^2)} dx = \frac{1}{1+x} + \frac{\arctan(x)^2}{2} + \log(1+x)$$

output `1/(1+x)+1/2*arctan(x)^2+ln(1+x)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x+x^3+(1+x)^2 \arctan(x)}{(1+x)^2(1+x^2)} dx = \frac{1}{1+x} + \frac{\arctan(x)^2}{2} + \log(1+x)$$

input `Integrate[(x + x^3 + (1 + x)^2*ArcTan[x])/((1 + x)^2*(1 + x^2)),x]`

output `(1 + x)^(-1) + ArcTan[x]^2/2 + Log[1 + x]`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x+1)^2 \arctan(x) + x^3 + x}{(x+1)^2 (x^2+1)} dx$$

$$\downarrow \text{7276}$$

$$\int \left(\frac{\arctan(x)}{x^2+1} + \frac{x}{(x+1)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\arctan(x)^2}{2} + \frac{1}{x+1} + \log(x+1)$$

input `Int[(x + x^3 + (1 + x)^2*ArcTan[x])/((1 + x)^2*(1 + x^2)),x]`

output `(1 + x)^(-1) + ArcTan[x]^2/2 + Log[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7276 `Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpanD[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
default	$\frac{1}{1+x} + \frac{\arctan(x)^2}{2} + \ln(1+x)$	17
parts	$\frac{1}{1+x} + \frac{\arctan(x)^2}{2} + \ln(1+x)$	17
parallelrisch	$\frac{\arctan(x)^2 x + 2 \ln(1+x)x + 2 + \arctan(x)^2 + 2 \ln(1+x)}{2+2x}$	33
risch	$-\frac{\ln(ix+1)^2}{8} + \frac{\ln(-ix+1)\ln(ix+1)}{4} + \frac{-\ln(-ix+1)^2 x + 8 \ln(1+x)x - \ln(-ix+1)^2 + 8 \ln(1+x) + 8}{8+8x}$	74

input `int((x+x^3+(1+x)^2*arctan(x))/(1+x)^2/(x^2+1),x,method=_RETURNVERBOSE)`

output `1/(1+x)+1/2*arctan(x)^2+ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.44

$$\int \frac{x + x^3 + (1+x)^2 \arctan(x)}{(1+x)^2 (1+x^2)} dx = \frac{(x+1) \arctan(x)^2 + 2(x+1) \log(x+1) + 2}{2(x+1)}$$

input `integrate((x+x^3+(1+x)^2*arctan(x))/(1+x)^2/(x^2+1),x, algorithm="fricas")`

output `1/2*((x + 1)*arctan(x)^2 + 2*(x + 1)*log(x + 1) + 2)/(x + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(15) = 30.

Time = 0.31 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.94

$$\int \frac{x + x^3 + (1+x)^2 \arctan(x)}{(1+x)^2 (1+x^2)} dx = \frac{2x \log(x+1)}{2x+2} + \frac{x \operatorname{atan}^2(x)}{2x+2} + \frac{2 \log(x+1)}{2x+2} + \frac{\operatorname{atan}^2(x)}{2x+2} + \frac{2}{2x+2}$$

input `integrate((x+x**3+(1+x)**2*atan(x))/(1+x)**2/(x**2+1),x)`

output `2*x*log(x + 1)/(2*x + 2) + x*atan(x)**2/(2*x + 2) + 2*log(x + 1)/(2*x + 2) + atan(x)**2/(2*x + 2) + 2/(2*x + 2)`

Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{x + x^3 + (1 + x)^2 \arctan(x)}{(1 + x)^2 (1 + x^2)} dx = \frac{1}{2} \arctan(x)^2 + \frac{1}{x + 1} + \log(x + 1)$$

input `integrate((x+x^3+(1+x)^2*arctan(x))/(1+x)^2/(x^2+1),x, algorithm="maxima")`

output `1/2*arctan(x)^2 + 1/(x + 1) + log(x + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs. 2(16) = 32.

Time = 0.13 (sec) , antiderivative size = 104, normalized size of antiderivative = 5.78

$$\int \frac{x + x^3 + (1 + x)^2 \arctan(x)}{(1 + x)^2 (1 + x^2)} dx$$

$$= \frac{(x + 1)\left(\frac{1}{x+1} - 1\right) \arctan\left((x + 1)\left(\frac{1}{x+1} - 1\right)\right)^2 + 2(x + 1)\left(\frac{1}{x+1} - 1\right) \log\left(- (x + 1)\left(\frac{1}{x+1} - 1\right) + 1\right) - \arctan\left((x + 1)\left(\frac{1}{x+1} - 1\right)\right)}{2\left((x + 1)\left(\frac{1}{x+1} - 1\right) - 1\right)}$$

input `integrate((x+x^3+(1+x)^2*arctan(x))/(1+x)^2/(x^2+1),x, algorithm="giac")`

output `1/2*((x + 1)*(1/(x + 1) - 1)*arctan((x + 1)*(1/(x + 1) - 1)))^2 + 2*(x + 1)*(1/(x + 1) - 1)*log(-(x + 1)*(1/(x + 1) - 1) + 1) - arctan((x + 1)*(1/(x + 1) - 1)))^2 - 2*log(-(x + 1)*(1/(x + 1) - 1) + 1) - 2)/((x + 1)*(1/(x + 1) - 1) - 1)`

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{x + x^3 + (1 + x)^2 \arctan(x)}{(1 + x)^2 (1 + x^2)} dx = \ln(x + 1) + \frac{1}{x + 1} + \frac{\arctan(x)^2}{2}$$

input `int((x + atan(x)*(x + 1)^2 + x^3)/((x^2 + 1)*(x + 1)^2),x)`output `log(x + 1) + 1/(x + 1) + atan(x)^2/2`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.94

$$\int \frac{x + x^3 + (1 + x)^2 \arctan(x)}{(1 + x)^2 (1 + x^2)} dx$$

$$= \frac{\arctan(x)^2 x + \arctan(x)^2 + 2 \log(x + 1) x + 2 \log(x + 1) - 2x}{2x + 2}$$

input `int((x+x^3+(1+x)^2*atan(x))/(1+x)^2/(x^2+1),x)`output `(atan(x)**2*x + atan(x)**2 + 2*log(x + 1)*x + 2*log(x + 1) - 2*x)/(2*(x + 1))`

3.126 $\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx$

Optimal result	950
Mathematica [A] (verified)	950
Rubi [A] (verified)	951
Maple [A] (verified)	953
Fricas [A] (verification not implemented)	954
Sympy [F(-1)]	954
Maxima [A] (verification not implemented)	955
Giac [A] (verification not implemented)	955
Mupad [B] (verification not implemented)	956
Reduce [B] (verification not implemented)	956

Optimal result

Integrand size = 21, antiderivative size = 68

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = -\frac{\sqrt{x}}{8} + \frac{x^{3/2}}{24} - \frac{x^{5/2}}{40} + \frac{x^{7/2}}{56} + \frac{\pi x^4}{16} + \frac{\arctan(\sqrt{x})}{8} - \frac{1}{8}x^4 \arctan(\sqrt{x})$$

output

```
-1/8*x^(1/2)+1/24*x^(3/2)-1/40*x^(5/2)+1/56*x^(7/2)+1/16*Pi*x^4+1/8*arctan(x^(1/2))-1/8*x^4*arctan(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.85

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\arctan(\sqrt{x})}{8} - \frac{1}{840}\sqrt{x}(105 - 35x + 21x^2 - 15x^3 + 210x^{7/2} \arctan(\sqrt{x} - \sqrt{1+x}))$$

input

```
Integrate[-(x^3*ArcTan[Sqrt[x] - Sqrt[1 + x]]),x]
```

output

```
ArcTan[Sqrt[x]]/8 - (Sqrt[x]*(105 - 35*x + 21*x^2 - 15*x^3 + 210*x^(7/2)*ArcTan[Sqrt[x] - Sqrt[1 + x]]))/840
```

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.09, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.476$, Rules used = {25, 5682, 15, 5361, 60, 60, 60, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -x^3 \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 25 \\
 & - \int x^3 \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 5682 \\
 & \frac{\pi \int x^3 dx}{4} - \frac{1}{2} \int x^3 \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 15 \\
 & \frac{\pi x^4}{16} - \frac{1}{2} \int x^3 \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 5361 \\
 & \frac{1}{2} \left(\frac{1}{8} \int \frac{x^{7/2}}{x+1} dx - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{8} \left(\frac{2x^{7/2}}{7} - \int \frac{x^{5/2}}{x+1} dx \right) - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{8} \left(\int \frac{x^{3/2}}{x+1} dx + \frac{2x^{7/2}}{7} - \frac{2x^{5/2}}{5} \right) - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 60 \\
& \frac{1}{2} \left(\frac{1}{8} \left(- \int \frac{\sqrt{x}}{x+1} dx + \frac{2x^{7/2}}{7} - \frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{3} \right) - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16} \\
& \downarrow 60 \\
& \frac{1}{2} \left(\frac{1}{8} \left(\int \frac{1}{\sqrt{x}(x+1)} dx + \frac{2x^{7/2}}{7} - \frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{3} - 2\sqrt{x} \right) - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16} \\
& \downarrow 73 \\
& \frac{1}{2} \left(\frac{1}{8} \left(2 \int \frac{1}{x+1} d\sqrt{x} + \frac{2x^{7/2}}{7} - \frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{3} - 2\sqrt{x} \right) - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16} \\
& \downarrow 216 \\
& \frac{1}{2} \left(\frac{1}{8} \left(2 \arctan(\sqrt{x}) + \frac{2x^{7/2}}{7} - \frac{2x^{5/2}}{5} + \frac{2x^{3/2}}{3} - 2\sqrt{x} \right) - \frac{1}{4} x^4 \arctan(\sqrt{x}) \right) + \frac{\pi x^4}{16}
\end{aligned}$$

input `Int[-(x^3*ArcTan[Sqrt[x] - Sqrt[1 + x]]), x]`

output `(Pi*x^4)/16 + (-1/4*(x^4*ArcTan[Sqrt[x]]) + (-2*Sqrt[x] + (2*x^(3/2))/3 - (2*x^(5/2))/5 + (2*x^(7/2))/7 + 2*ArcTan[Sqrt[x]])/8)/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 60 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m + 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]`

rule 5682 `Int[ArcTan[(v_) + (s_.)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2 + 1]`

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.66

method	result	size
default	$-\frac{x^4 \arctan(\sqrt{x} - \sqrt{1+x})}{4} + \frac{x^{\frac{7}{2}}}{56} - \frac{x^{\frac{5}{2}}}{40} + \frac{x^{\frac{3}{2}}}{24} - \frac{\sqrt{x}}{8} + \frac{\arctan(\sqrt{x})}{8}$	45
parts	$-\frac{x^4 \arctan(\sqrt{x} - \sqrt{1+x})}{4} + \frac{x^{\frac{7}{2}}}{56} - \frac{x^{\frac{5}{2}}}{40} + \frac{x^{\frac{3}{2}}}{24} - \frac{\sqrt{x}}{8} + \frac{\arctan(\sqrt{x})}{8}$	45

input `int(-x^3*arctan(x^(1/2)-(1+x)^(1/2)),x,method=_RETURNVERBOSE)`

output `-1/4*x^4*arctan(x^(1/2)-(1+x)^(1/2))+1/56*x^(7/2)-1/40*x^(5/2)+1/24*x^(3/2)-1/8*x^(1/2)+1/8*arctan(x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.59

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{4} (x^4 - 1) \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{840} (15x^3 - 21x^2 + 35x - 105)\sqrt{x}$$

input `integrate(-x^3*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="fricas")`

output `1/4*(x^4 - 1)*arctan(sqrt(x + 1) - sqrt(x)) + 1/840*(15*x^3 - 21*x^2 + 35*x - 105)*sqrt(x)`

Sympy [F(-1)]

Timed out.

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \text{Timed out}$$

input `integrate(-x**3*atan(x**(1/2)-(1+x)**(1/2)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.65

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{4} x^4 \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{56} x^{\frac{7}{2}} - \frac{1}{40} x^{\frac{5}{2}} + \frac{1}{24} x^{\frac{3}{2}} - \frac{1}{8} \sqrt{x} + \frac{1}{8} \arctan(\sqrt{x})$$

input `integrate(-x^3*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="maxima")`output `1/4*x^4*arctan(sqrt(x + 1) - sqrt(x)) + 1/56*x^(7/2) - 1/40*x^(5/2) + 1/24*x^(3/2) - 1/8*sqrt(x) + 1/8*arctan(sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.65

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = -\frac{1}{4} x^4 \arctan(-\sqrt{x+1} + \sqrt{x}) + \frac{1}{56} x^{\frac{7}{2}} - \frac{1}{40} x^{\frac{5}{2}} + \frac{1}{24} x^{\frac{3}{2}} - \frac{1}{8} \sqrt{x} + \frac{1}{8} \arctan(\sqrt{x})$$

input `integrate(-x^3*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="giac")`output `-1/4*x^4*arctan(-sqrt(x + 1)+ sqrt(x)) + 1/56*x^(7/2) - 1/40*x^(5/2) + 1/24*x^(3/2) - 1/8*sqrt(x) + 1/8*arctan(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 1.72 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.06

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{x^{3/2}}{24} - \frac{\sqrt{x}}{8} - \frac{x^{5/2}}{40} + \frac{x^{7/2}}{56} + \frac{\operatorname{atan}(\sqrt{x+1} - \sqrt{x}) \left(\frac{x^5}{2} + \frac{x^4}{2}\right)}{2x+2} + \frac{\ln\left(\frac{(-1+\sqrt{x}i)^2}{x+1}\right) i}{16}$$

input `int(x^3*atan((x + 1)^(1/2) - x^(1/2)),x)`output `(log((x^(1/2)*i - 1)^2/(x + 1))*i)/16 - x^(1/2)/8 + x^(3/2)/24 - x^(5/2)/40 + x^(7/2)/56 + (atan((x + 1)^(1/2) - x^(1/2))*(x^4/2 + x^5/2))/(2*x + 2)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.75

$$\int -x^3 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right) x^4}{4} - \frac{\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right)}{4} + \frac{\sqrt{x} x^3}{56} - \frac{\sqrt{x} x^2}{40} + \frac{\sqrt{x} x}{24} - \frac{\sqrt{x}}{8}$$

input `int(-x^3*atan(x^(1/2)-(1+x)^(1/2)),x)`output `(210*atan(1/(sqrt(x + 1) + sqrt(x)))*x**4 - 210*atan(1/(sqrt(x + 1) + sqrt(x))) + 15*sqrt(x)*x**3 - 21*sqrt(x)*x**2 + 35*sqrt(x)*x - 105*sqrt(x))/84`
0

3.127 $\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx$

Optimal result	957
Mathematica [A] (verified)	957
Rubi [A] (verified)	958
Maple [A] (verified)	960
Fricas [A] (verification not implemented)	961
Sympy [A] (verification not implemented)	961
Maxima [A] (verification not implemented)	962
Giac [A] (verification not implemented)	962
Mupad [B] (verification not implemented)	963
Reduce [B] (verification not implemented)	963

Optimal result

Integrand size = 21, antiderivative size = 59

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\sqrt{x}}{6} - \frac{x^{3/2}}{18} + \frac{x^{5/2}}{30} + \frac{\pi x^3}{12} - \frac{\arctan(\sqrt{x})}{6} - \frac{1}{6}x^3 \arctan(\sqrt{x})$$

output

```
1/6*x^(1/2)-1/18*x^(3/2)+1/30*x^(5/2)+1/12*Pi*x^3-1/6*arctan(x^(1/2))-1/6*x^3*arctan(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{90} \left(-15 \arctan(\sqrt{x}) - \sqrt{x}(-15 + 5x - 3x^2 + 30x^{5/2} \arctan(\sqrt{x} - \sqrt{1+x})) \right)$$

input

```
Integrate[-(x^2*ArcTan[Sqrt[x] - Sqrt[1 + x]]),x]
```

output

```
(-15*ArcTan[Sqrt[x]] - Sqrt[x]*(-15 + 5*x - 3*x^2 + 30*x^(5/2)*ArcTan[Sqrt[x] - Sqrt[1 + x]]))/90
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {25, 5682, 15, 5361, 60, 60, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -x^2 \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 25 \\
 & - \int x^2 \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 5682 \\
 & \frac{\pi \int x^2 dx}{4} - \frac{1}{2} \int x^2 \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 15 \\
 & \frac{\pi x^3}{12} - \frac{1}{2} \int x^2 \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 5361 \\
 & \frac{1}{2} \left(\frac{1}{6} \int \frac{x^{5/2}}{x+1} dx - \frac{1}{3} x^3 \arctan(\sqrt{x}) \right) + \frac{\pi x^3}{12} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{6} \left(\frac{2x^{5/2}}{5} - \int \frac{x^{3/2}}{x+1} dx \right) - \frac{1}{3} x^3 \arctan(\sqrt{x}) \right) + \frac{\pi x^3}{12} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{6} \left(\int \frac{\sqrt{x}}{x+1} dx + \frac{2x^{5/2}}{5} - \frac{2x^{3/2}}{3} \right) - \frac{1}{3} x^3 \arctan(\sqrt{x}) \right) + \frac{\pi x^3}{12}
 \end{aligned}$$

$$\begin{aligned} & \downarrow 60 \\ & \frac{1}{2} \left(\frac{1}{6} \left(- \int \frac{1}{\sqrt{x}(x+1)} dx + \frac{2x^{5/2}}{5} - \frac{2x^{3/2}}{3} + 2\sqrt{x} \right) - \frac{1}{3} x^3 \arctan(\sqrt{x}) \right) + \frac{\pi x^3}{12} \\ & \downarrow 73 \\ & \frac{1}{2} \left(\frac{1}{6} \left(-2 \int \frac{1}{x+1} d\sqrt{x} + \frac{2x^{5/2}}{5} - \frac{2x^{3/2}}{3} + 2\sqrt{x} \right) - \frac{1}{3} x^3 \arctan(\sqrt{x}) \right) + \frac{\pi x^3}{12} \\ & \downarrow 216 \\ & \frac{1}{2} \left(\frac{1}{6} \left(-2 \arctan(\sqrt{x}) + \frac{2x^{5/2}}{5} - \frac{2x^{3/2}}{3} + 2\sqrt{x} \right) - \frac{1}{3} x^3 \arctan(\sqrt{x}) \right) + \frac{\pi x^3}{12} \end{aligned}$$

input `Int[-(x^2*ArcTan[Sqrt[x] - Sqrt[1 + x]]),x]`

output `(Pi*x^3)/12 + ((2*Sqrt[x] - (2*x^(3/2)))/3 + (2*x^(5/2))/5 - 2*ArcTan[Sqrt[x]])/6 - (x^3*ArcTan[Sqrt[x]]/3)/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
 rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
 , 0] || GtQ[b, 0])`

rule 5361 `Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
 Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
 1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
 x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
 & IntegerQ[m])) && NeQ[m, -1]`

rule 5682 `Int[ArcTan[(v_) + (s_.)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u
 , x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2
 + 1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.68

method	result	size
default	$-\frac{x^3 \arctan(\sqrt{x}-\sqrt{1+x})}{3} + \frac{x^{\frac{5}{2}}}{30} - \frac{x^{\frac{3}{2}}}{18} + \frac{\sqrt{x}}{6} - \frac{\arctan(\sqrt{x})}{6}$	40
parts	$-\frac{x^3 \arctan(\sqrt{x}-\sqrt{1+x})}{3} + \frac{x^{\frac{5}{2}}}{30} - \frac{x^{\frac{3}{2}}}{18} + \frac{\sqrt{x}}{6} - \frac{\arctan(\sqrt{x})}{6}$	40

input `int(-x^2*arctan(x^(1/2)-(1+x)^(1/2)),x,method=_RETURNVERBOSE)`

output `-1/3*x^3*arctan(x^(1/2)-(1+x)^(1/2))+1/30*x^(5/2)-1/18*x^(3/2)+1/6*x^(1/2)
 -1/6*arctan(x^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.59

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{3}(x^3 + 1) \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{90}(3x^2 - 5x + 15)\sqrt{x}$$

input `integrate(-x^2*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="fricas")`

output `1/3*(x^3 + 1)*arctan(sqrt(x + 1) - sqrt(x)) + 1/90*(3*x^2 - 5*x + 15)*sqrt(x)`

Sympy [A] (verification not implemented)

Time = 85.40 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.78

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{x^{5/2}}{30} - \frac{x^{3/2}}{18} + \frac{\sqrt{x}}{6} - \frac{x^3 \operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{3} - \frac{\operatorname{atan}(\sqrt{x})}{6}$$

input `integrate(-x**2*atan(x**(1/2)-(1+x)**(1/2)),x)`

output `x**(5/2)/30 - x**(3/2)/18 + sqrt(x)/6 - x**3*atan(sqrt(x) - sqrt(x + 1))/3 - atan(sqrt(x))/6`

Maxima [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.66

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{3} x^3 \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{30} x^{\frac{5}{2}} - \frac{1}{18} x^{\frac{3}{2}} + \frac{1}{6} \sqrt{x} - \frac{1}{6} \arctan(\sqrt{x})$$

input `integrate(-x^2*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="maxima")`

output `1/3*x^3*arctan(sqrt(x + 1) - sqrt(x)) + 1/30*x^(5/2) - 1/18*x^(3/2) + 1/6*sqrt(x) - 1/6*arctan(sqrt(x))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.66

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = -\frac{1}{3} x^3 \arctan(-\sqrt{x+1} + \sqrt{x}) + \frac{1}{30} x^{\frac{5}{2}} - \frac{1}{18} x^{\frac{3}{2}} + \frac{1}{6} \sqrt{x} - \frac{1}{6} \arctan(\sqrt{x})$$

input `integrate(-x^2*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="giac")`

output `-1/3*x^3*arctan(-sqrt(x + 1)+ sqrt(x)) + 1/30*x^(5/2) - 1/18*x^(3/2) + 1/6*sqrt(x) - 1/6*arctan(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.10

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\sqrt{x}}{6} - \frac{x^{3/2}}{18} + \frac{x^{5/2}}{30} + \frac{\operatorname{atan}(\sqrt{x+1} - \sqrt{x}) \left(\frac{2x^4}{3} + \frac{2x^3}{3}\right)}{2x+2} + \frac{\ln\left(\frac{(\sqrt{x}-i)^2}{x+1}\right) i}{12}$$

input `int(x^2*atan((x + 1)^(1/2) - x^(1/2)),x)`output `(log((x^(1/2) - 1i)^2/(x + 1))*1i)/12 + x^(1/2)/6 - x^(3/2)/18 + x^(5/2)/30 + (atan((x + 1)^(1/2) - x^(1/2))*((2*x^3)/3 + (2*x^4)/3))/(2*x + 2)`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.75

$$\int -x^2 \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right) x^3}{3} + \frac{\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right)}{3} + \frac{\sqrt{x} x^2}{30} - \frac{\sqrt{x} x}{18} + \frac{\sqrt{x}}{6}$$

input `int(-x^2*atan(x^(1/2)-(1+x)^(1/2)),x)`output `(30*atan(1/(sqrt(x + 1) + sqrt(x)))*x**3 + 30*atan(1/(sqrt(x + 1) + sqrt(x))) + 3*sqrt(x)*x**2 - 5*sqrt(x)*x + 15*sqrt(x))/90`

3.128 $\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx$

Optimal result	964
Mathematica [A] (verified)	964
Rubi [A] (verified)	965
Maple [A] (verified)	967
Fricas [A] (verification not implemented)	967
Sympy [A] (verification not implemented)	968
Maxima [A] (verification not implemented)	968
Giac [A] (verification not implemented)	968
Mupad [B] (verification not implemented)	969
Reduce [B] (verification not implemented)	969

Optimal result

Integrand size = 19, antiderivative size = 50

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = -\frac{\sqrt{x}}{4} + \frac{x^{3/2}}{12} + \frac{\pi x^2}{8} + \frac{\arctan(\sqrt{x})}{4} - \frac{1}{4}x^2 \arctan(\sqrt{x})$$

output

```
-1/4*x^(1/2)+1/12*x^(3/2)+1/8*Pi*x^2+1/4*arctan(x^(1/2))-1/4*x^2*arctan(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{12} \left(3 \arctan(\sqrt{x}) - \sqrt{x} \left(3 - x + 6x^{3/2} \arctan(\sqrt{x} - \sqrt{1+x}) \right) \right)$$

input

```
Integrate[-(x*ArcTan[Sqrt[x] - Sqrt[1 + x]]),x]
```

output

```
(3*ArcTan[Sqrt[x]] - Sqrt[x]*(3 - x + 6*x^(3/2)*ArcTan[Sqrt[x] - Sqrt[1 + x]]))/12
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {25, 5682, 15, 5361, 60, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -x \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 25 \\
 & - \int x \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 5682 \\
 & \frac{\pi}{4} \int x dx - \frac{1}{2} \int x \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 15 \\
 & \frac{\pi x^2}{8} - \frac{1}{2} \int x \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 5361 \\
 & \frac{1}{2} \left(\frac{1}{4} \int \frac{x^{3/2}}{x+1} dx - \frac{1}{2} x^2 \arctan(\sqrt{x}) \right) + \frac{\pi x^2}{8} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{4} \left(\frac{2x^{3/2}}{3} - \int \frac{\sqrt{x}}{x+1} dx \right) - \frac{1}{2} x^2 \arctan(\sqrt{x}) \right) + \frac{\pi x^2}{8} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{4} \left(\int \frac{1}{\sqrt{x}(x+1)} dx + \frac{2x^{3/2}}{3} - 2\sqrt{x} \right) - \frac{1}{2} x^2 \arctan(\sqrt{x}) \right) + \frac{\pi x^2}{8} \\
 & \quad \downarrow 73 \\
 & \frac{1}{2} \left(\frac{1}{4} \left(2 \int \frac{1}{x+1} d\sqrt{x} + \frac{2x^{3/2}}{3} - 2\sqrt{x} \right) - \frac{1}{2} x^2 \arctan(\sqrt{x}) \right) + \frac{\pi x^2}{8} \\
 & \quad \downarrow 216
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(2 \arctan(\sqrt{x}) + \frac{2x^{3/2}}{3} - 2\sqrt{x} \right) - \frac{1}{2} x^2 \arctan(\sqrt{x}) \right) + \frac{\pi x^2}{8}$$

input `Int[-(x*ArcTan[Sqrt[x] - Sqrt[1 + x]]),x]`

output `(Pi*x^2)/8 + (-1/2*(x^2*ArcTan[Sqrt[x]]) + (-2*Sqrt[x] + (2*x^(3/2))/3 + 2*ArcTan[Sqrt[x]])/4)/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 60 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 5361

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

rule 5682

```
Int[ArcTan[(v_) + (s_)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u
, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2
+ 1]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.70

method	result	size
default	$-\frac{x^2 \arctan(\sqrt{x}-\sqrt{1+x})}{2} + \frac{x^{\frac{3}{2}}}{12} - \frac{\sqrt{x}}{4} + \frac{\arctan(\sqrt{x})}{4}$	35
parts	$-\frac{x^2 \arctan(\sqrt{x}-\sqrt{1+x})}{2} + \frac{x^{\frac{3}{2}}}{12} - \frac{\sqrt{x}}{4} + \frac{\arctan(\sqrt{x})}{4}$	35

input

```
int(-x*arctan(x^(1/2)-(1+x)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
-1/2*x^2*arctan(x^(1/2)-(1+x)^(1/2))+1/12*x^(3/2)-1/4*x^(1/2)+1/4*arctan(x
^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.56

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{2} (x^2 - 1) \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{12} (x - 3)\sqrt{x}$$

input

```
integrate(-x*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="fricas")
```

output

```
1/2*(x^2 - 1)*arctan(sqrt(x + 1) - sqrt(x)) + 1/12*(x - 3)*sqrt(x)
```


Sympy [A] (verification not implemented)

Time = 21.45 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.78

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{x^{\frac{3}{2}}}{12} - \frac{\sqrt{x}}{4} - \frac{x^2 \operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{2} + \frac{\operatorname{atan}(\sqrt{x})}{4}$$

input `integrate(-x*atan(x**(1/2)-(1+x)**(1/2)),x)`output `x**(3/2)/12 - sqrt(x)/4 - x**2*atan(sqrt(x) - sqrt(x + 1))/2 + atan(sqrt(x))/4`**Maxima [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.68

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{2} x^2 \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{12} x^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} + \frac{1}{4} \arctan(\sqrt{x})$$

input `integrate(-x*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="maxima")`output `1/2*x^2*arctan(sqrt(x + 1) - sqrt(x)) + 1/12*x^(3/2) - 1/4*sqrt(x) + 1/4*arctan(sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.68

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = -\frac{1}{2} x^2 \arctan(-\sqrt{x+1} + \sqrt{x}) + \frac{1}{12} x^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} + \frac{1}{4} \arctan(\sqrt{x})$$

input `integrate(-x*arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="giac")`

output $-1/2*x^2*\arctan(-\sqrt{x + 1} + \sqrt{x}) + 1/12*x^{(3/2)} - 1/4*\sqrt{x} + 1/4*\arctan(\sqrt{x})$

Mupad [B] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.16

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{x^{3/2}}{12} - \frac{\sqrt{x}}{4} + \frac{\operatorname{atan}(\sqrt{x+1} - \sqrt{x})(x^3 + x^2)}{2x + 2} + \frac{\ln\left(\frac{(-1+\sqrt{x}i)^2}{x+1}\right) i}{8}$$

input $\text{int}(x*\operatorname{atan}((x + 1)^{(1/2)} - x^{(1/2)}), x)$

output $(\log((x^{(1/2)}*i - 1)^2/(x + 1)*i)/8 - x^{(1/2)}/4 + x^{(3/2)}/12 + (\operatorname{atan}((x + 1)^{(1/2)} - x^{(1/2)})*(x^2 + x^3))/(2*x + 2)$

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.74

$$\int -x \arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right) x^2}{2} - \frac{\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right)}{2} + \frac{\sqrt{x} x}{12} - \frac{\sqrt{x}}{4}$$

input $\text{int}(-x*\operatorname{atan}(x^{(1/2)}-(1+x)^{(1/2)}), x)$

output $(6*\operatorname{atan}(1/(\sqrt{x + 1} + \sqrt{x}))*x**2 - 6*\operatorname{atan}(1/(\sqrt{x + 1} + \sqrt{x})) + \sqrt{x}*x - 3*\sqrt{x})/12$

3.129 $\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx$

Optimal result	970
Mathematica [A] (verified)	970
Rubi [A] (verified)	971
Maple [A] (verified)	973
Fricas [A] (verification not implemented)	973
Sympy [A] (verification not implemented)	974
Maxima [A] (verification not implemented)	974
Giac [A] (verification not implemented)	974
Mupad [B] (verification not implemented)	975
Reduce [B] (verification not implemented)	975

Optimal result

Integrand size = 18, antiderivative size = 37

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\sqrt{x}}{2} + \frac{\pi x}{4} - \frac{\arctan(\sqrt{x})}{2} - \frac{1}{2}x \arctan(\sqrt{x})$$

output

```
1/2*x^(1/2)+1/4*Pi*x-1/2*arctan(x^(1/2))-1/2*x*arctan(x^(1/2))
```

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.84

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\sqrt{x}}{2} - (1+x) \arctan(\sqrt{x} - \sqrt{1+x})$$

input

```
Integrate[-ArcTan[Sqrt[x] - Sqrt[1 + x]],x]
```

output

```
Sqrt[x]/2 - (1 + x)*ArcTan[Sqrt[x] - Sqrt[1 + x]]
```

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {25, 5682, 24, 5345, 60, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -\arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 25 \\
 & - \int \arctan(\sqrt{x} - \sqrt{x+1}) dx \\
 & \quad \downarrow 5682 \\
 & \frac{\pi}{4} \int 1 dx - \frac{1}{2} \int \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 24 \\
 & \frac{\pi x}{4} - \frac{1}{2} \int \arctan(\sqrt{x}) dx \\
 & \quad \downarrow 5345 \\
 & \frac{1}{2} \left(\frac{1}{2} \int \frac{\sqrt{x}}{x+1} dx - x \arctan(\sqrt{x}) \right) + \frac{\pi x}{4} \\
 & \quad \downarrow 60 \\
 & \frac{1}{2} \left(\frac{1}{2} \left(2\sqrt{x} - \int \frac{1}{\sqrt{x}(x+1)} dx \right) - x \arctan(\sqrt{x}) \right) + \frac{\pi x}{4} \\
 & \quad \downarrow 73 \\
 & \frac{1}{2} \left(\frac{1}{2} \left(2\sqrt{x} - 2 \int \frac{1}{x+1} d\sqrt{x} \right) - x \arctan(\sqrt{x}) \right) + \frac{\pi x}{4} \\
 & \quad \downarrow 216 \\
 & \frac{1}{2} \left(\frac{1}{2} (2\sqrt{x} - 2 \arctan(\sqrt{x})) - x \arctan(\sqrt{x}) \right) + \frac{\pi x}{4}
 \end{aligned}$$

input

```
Int[-ArcTan[Sqrt[x] - Sqrt[1 + x]], x]
```

output $(\text{Pi} \cdot x)/4 + ((2 \cdot \text{Sqrt}[x] - 2 \cdot \text{ArcTan}[\text{Sqrt}[x]])/2 - x \cdot \text{ArcTan}[\text{Sqrt}[x]])/2$

Defintions of rubi rules used

rule 24 $\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a \cdot x, x] /; \text{FreeQ}[a, x]$

rule 25 $\text{Int}[-(F x_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F x, x], x]$

rule 60 $\text{Int}[(a_. + (b_.)(x_))^{(m_.)}((c_.) + (d_.)(x_))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b \cdot x)^{(m + 1)}((c + d \cdot x)^n / (b \cdot (m + n + 1))), x] + \text{Simp}[n \cdot ((b \cdot c - a \cdot d) / (b \cdot (m + n + 1))) \text{Int}[(a + b \cdot x)^m (c + d \cdot x)^{(n - 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (! \text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& ! \text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 73 $\text{Int}[(a_. + (b_.)(x_))^{(m_.)}((c_.) + (d_.)(x_))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{Subst}[\text{Int}[x^{(p \cdot (m + 1) - 1)}(c - a \cdot (d/b) + d \cdot (x^p/b))^{(n)}, x], x, (a + b \cdot x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 216 $\text{Int}[(a_. + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1 / (\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x / \text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{GtQ}[a, 0] || \text{GtQ}[b, 0])$

rule 5345 $\text{Int}[(a_. + \text{ArcTan}[(c_.)(x_)^{(n_.)}] \cdot (b_.))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot \text{ArcTan}[c \cdot x^n])^p, x] - \text{Simp}[b \cdot c \cdot n \cdot p \text{Int}[x^n \cdot ((a + b \cdot \text{ArcTan}[c \cdot x^n])^{(p - 1)}) / (1 + c^2 \cdot x^{(2 \cdot n)}), x], x] /; \text{FreeQ}[\{a, b, c, n\}, x] \&\& \text{IGtQ}[p, 0] \&\& (\text{EqQ}[n, 1] || \text{EqQ}[p, 1])$

rule 5682

```
Int[ArcTan[(v_) + (s_)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u
, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2
+ 1]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.76

method	result	size
default	$-x \arctan(\sqrt{x} - \sqrt{1+x}) + \frac{\sqrt{x}}{2} - \frac{\arctan(\sqrt{x})}{2}$	28
parts	$-x \arctan(\sqrt{x} - \sqrt{1+x}) + \frac{\sqrt{x}}{2} - \frac{\arctan(\sqrt{x})}{2}$	28

input

```
int(-arctan(x^(1/2)-(1+x)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
-x*arctan(x^(1/2)-(1+x)^(1/2))+1/2*x^(1/2)-1/2*arctan(x^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.59

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = (x+1) \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{2} \sqrt{x}$$

input

```
integrate(-arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="fricas")
```

output

```
(x + 1)*arctan(sqrt(x + 1) - sqrt(x)) + 1/2*sqrt(x)
```

Sympy [A] (verification not implemented)

Time = 7.75 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int -\arctan\left(\sqrt{x} - \sqrt{1+x}\right) dx = \frac{\sqrt{x}}{2} - x \operatorname{atan}\left(\sqrt{x} - \sqrt{x+1}\right) - \frac{\operatorname{atan}\left(\sqrt{x}\right)}{2}$$

input `integrate(-atan(x**(1/2)-(1+x)**(1/2)),x)`output `sqrt(x)/2 - x*atan(sqrt(x) - sqrt(x + 1)) - atan(sqrt(x))/2`**Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.70

$$\int -\arctan\left(\sqrt{x} - \sqrt{1+x}\right) dx = x \arctan\left(\sqrt{x+1} - \sqrt{x}\right) + \frac{1}{2}\sqrt{x} - \frac{1}{2}\arctan\left(\sqrt{x}\right)$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="maxima")`output `x*arctan(sqrt(x + 1) - sqrt(x)) + 1/2*sqrt(x) - 1/2*arctan(sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int -\arctan\left(\sqrt{x} - \sqrt{1+x}\right) dx = -x \arctan\left(-\sqrt{x+1} + \sqrt{x}\right) + \frac{1}{2}\sqrt{x} - \frac{1}{2}\arctan\left(\sqrt{x}\right)$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="giac")`output `-x*arctan(-sqrt(x + 1) + sqrt(x)) + 1/2*sqrt(x) - 1/2*arctan(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.08

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = x \operatorname{atan}(\sqrt{x+1} - \sqrt{x}) + \frac{\sqrt{x}}{2} - \frac{\ln\left(\frac{(-1+\sqrt{x}i)^2}{x+1}\right) i}{4}$$

input `int(atan((x + 1)^(1/2) - x^(1/2)),x)`output `x*atan((x + 1)^(1/2) - x^(1/2)) - (log((x^(1/2)*1i - 1)^2/(x + 1))*1i)/4 + x^(1/2)/2`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.73

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = \operatorname{atan}\left(\frac{1}{\sqrt{x+1} + \sqrt{x}}\right) x + \operatorname{atan}\left(\frac{1}{\sqrt{x+1} + \sqrt{x}}\right) + \frac{\sqrt{x}}{2}$$

input `int(-atan(x^(1/2)-(1+x)^(1/2)),x)`output `(2*atan(1/(sqrt(x + 1) + sqrt(x)))*x + 2*atan(1/(sqrt(x + 1) + sqrt(x))) + sqrt(x))/2`

3.130 $\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} dx$

Optimal result	976
Mathematica [A] (verified)	976
Rubi [A] (verified)	977
Maple [B] (verified)	979
Fricas [F]	979
Sympy [F]	980
Maxima [A] (verification not implemented)	980
Giac [F]	980
Mupad [F(-1)]	981
Reduce [F]	981

Optimal result

Integrand size = 21, antiderivative size = 42

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} dx = \frac{1}{4}\pi \log(x) - \frac{1}{2}i \operatorname{PolyLog}(2, -i\sqrt{x}) + \frac{1}{2}i \operatorname{PolyLog}(2, i\sqrt{x})$$

output `1/4*Pi*ln(x)-1/2*I*polylog(2,-I*x^(1/2))+1/2*I*polylog(2,I*x^(1/2))`

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.00

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} dx = -\arctan(\sqrt{x}-\sqrt{1+x}) \log(x) + \frac{1}{4}i((\log(1-i\sqrt{x})-\log(1+i\sqrt{x})) \log(x) - 2 \operatorname{PolyLog}(2, -i\sqrt{x}) + 2 \operatorname{PolyLog}(2, i\sqrt{x}))$$

input `Integrate[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x), x]`

output

```
-(ArcTan[Sqrt[x] - Sqrt[1 + x]]*Log[x]) + (I/4)*((Log[1 - I*Sqrt[x]] - Log
[1 + I*Sqrt[x]])*Log[x] - 2*PolyLog[2, (-I)*Sqrt[x]] + 2*PolyLog[2, I*Sqrt
[x]])
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {25, 5682, 14, 5359, 5355, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -\frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x} dx \\
 & \quad \downarrow \text{5682} \\
 & \frac{1}{4}\pi \int \frac{1}{x} dx - \frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x} dx \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{4}\pi \log(x) - \frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x} dx \\
 & \quad \downarrow \text{5359} \\
 & \frac{1}{4}\pi \log(x) - \int \frac{\arctan(\sqrt{x})}{\sqrt{x}} d\sqrt{x} \\
 & \quad \downarrow \text{5355} \\
 & -\frac{1}{2}i \int \frac{\log(1 - i\sqrt{x})}{\sqrt{x}} d\sqrt{x} + \frac{1}{2}i \int \frac{\log(i\sqrt{x} + 1)}{\sqrt{x}} d\sqrt{x} + \frac{1}{4}\pi \log(x) \\
 & \quad \downarrow \text{2838} \\
 & -\frac{1}{2}i \text{PolyLog}(2, -i\sqrt{x}) + \frac{1}{2}i \text{PolyLog}(2, i\sqrt{x}) + \frac{1}{4}\pi \log(x)
 \end{aligned}$$

input `Int[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x),x]`

output `(Pi*Log[x])/4 - (I/2)*PolyLog[2, (-I)*Sqrt[x]] + (I/2)*PolyLog[2, I*Sqrt[x]]`

Defintions of rubi rules used

rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 5355 `Int[((a_) + ArcTan[(c_)*(x_)])*(b_)/(x_), x_Symbol] := Simp[a*Log[x], x] + (Simp[I*(b/2) Int[Log[1 - I*c*x]/x, x], x] - Simp[I*(b/2) Int[Log[1 + I*c*x]/x, x], x]) /; FreeQ[{a, b, c}, x]`

rule 5359 `Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_)/(x_), x_Symbol] := Simp[1/n Subst[Int[(a + b*ArcTan[c*x])^p/x, x], x, x^n], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0]`

rule 5682 `Int[ArcTan[(v_) + (s_)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2 + 1]`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 373 vs. $2(28) = 56$.

Time = 0.26 (sec) , antiderivative size = 374, normalized size of antiderivative = 8.90

method	result
default	$2 \arctan(\sqrt{x} - \sqrt{1+x}) \ln\left(1 - \frac{1+i(\sqrt{x}-\sqrt{1+x})}{\sqrt{(\sqrt{x}-\sqrt{1+x})^2+1}}\right) - 2i \operatorname{polylog}\left(2, \frac{1+i(\sqrt{x}-\sqrt{1+x})}{\sqrt{(\sqrt{x}-\sqrt{1+x})^2+1}}\right) + 2 \arctan$

input `int(-arctan(x^(1/2)-(1+x)^(1/2))/x,x,method=_RETURNVERBOSE)`

output `2*arctan(x^(1/2)-(1+x)^(1/2))*ln(1-(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))-2*I*polylog(2,(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))+2*arctan(x^(1/2)-(1+x)^(1/2))*ln(1+(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))-I*polylog(2,-(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))+2*arctan(x^(1/2)-(1+x)^(1/2))*ln(1+(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))-2*I*polylog(2,-(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))-2*arctan(x^(1/2)-(1+x)^(1/2))*ln(1+(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))+1/2*I*polylog(2,-(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))-1/2*I*polylog(2,(1+I*(x^(1/2)-(1+x)^(1/2)))/((x^(1/2)-(1+x)^(1/2))^2+1)^(1/2))`

Fricas [F]

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x} dx = \int -\frac{\arctan(-\sqrt{x+1} + \sqrt{x})}{x} dx$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x,x, algorithm="fricas")`

output `integral(arctan(sqrt(x + 1) - sqrt(x))/x, x)`

Sympy [F]

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x} dx = -\int \frac{\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{x} dx$$

input `integrate(-atan(x**(1/2)-(1+x)**(1/2))/x,x)`

output `-Integral(atan(sqrt(x) - sqrt(x + 1))/x, x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.02

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x} dx = \frac{1}{4} \pi \log(x+1) + \arctan(\sqrt{x+1} - \sqrt{x}) \log(x) + \frac{1}{2} i \operatorname{Li}_2(i\sqrt{x}+1) - \frac{1}{2} i \operatorname{Li}_2(-i\sqrt{x}+1)$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x,x, algorithm="maxima")`

output `1/4*pi*log(x + 1) + arctan(sqrt(x + 1) - sqrt(x))*log(x) + 1/2*I*dilog(I*sqrt(x) + 1) - 1/2*I*dilog(-I*sqrt(x) + 1)`

Giac [F]

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x} dx = \int -\frac{\arctan(-\sqrt{x+1} + \sqrt{x})}{x} dx$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x,x, algorithm="giac")`

output `integrate(-arctan(-sqrt(x + 1) + sqrt(x))/x, x)`

Mupad [F(-1)]

Timed out.

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x} dx = \int \frac{\operatorname{atan}(\sqrt{x+1} - \sqrt{x})}{x} dx$$

input `int(atan((x + 1)^(1/2) - x^(1/2))/x,x)`output `int(atan((x + 1)^(1/2) - x^(1/2))/x, x)`**Reduce [F]**

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x} dx = \int \frac{\operatorname{atan}(\sqrt{x+1} - \sqrt{x})}{x} dx$$

input `int(-atan(x^(1/2)-(1+x)^(1/2))/x,x)`output `int(atan(sqrt(x + 1) - sqrt(x))/x,x)`

3.131 $\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^2} dx$

Optimal result	982
Mathematica [A] (verified)	982
Rubi [A] (verified)	983
Maple [B] (verified)	985
Fricas [A] (verification not implemented)	985
Sympy [B] (verification not implemented)	986
Maxima [A] (verification not implemented)	987
Giac [A] (verification not implemented)	987
Mupad [B] (verification not implemented)	988
Reduce [B] (verification not implemented)	988

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^2} dx = -\frac{\pi}{4x} + \frac{1}{2\sqrt{x}} + \frac{\arctan(\sqrt{x})}{2} + \frac{\arctan(\sqrt{x})}{2x}$$

output -1/4*Pi/x+1/2/x^(1/2)+1/2*arctan(x^(1/2))+1/2*arctan(x^(1/2))/x

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.98

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^2} dx = \frac{1}{2\sqrt{x}} + \frac{\arctan(\sqrt{x})}{2} + \frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x}$$

input Integrate[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x^2), x]

output 1/(2*Sqrt[x]) + ArcTan[Sqrt[x]]/2 + ArcTan[Sqrt[x] - Sqrt[1 + x]]/x

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {25, 5682, 15, 5361, 61, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -\frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x^2} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x^2} dx \\
 & \quad \downarrow \text{5682} \\
 & \frac{1}{4}\pi \int \frac{1}{x^2} dx - \frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x^2} dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x^2} dx - \frac{\pi}{4x} \\
 & \quad \downarrow \text{5361} \\
 & \frac{1}{2} \left(\frac{\arctan(\sqrt{x})}{x} - \frac{1}{2} \int \frac{1}{x^{3/2}(x+1)} dx \right) - \frac{\pi}{4x} \\
 & \quad \downarrow \text{61} \\
 & \frac{1}{2} \left(\frac{1}{2} \left(\int \frac{1}{\sqrt{x}(x+1)} dx + \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{x} \right) - \frac{\pi}{4x} \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(\frac{1}{2} \left(2 \int \frac{1}{x+1} d\sqrt{x} + \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{x} \right) - \frac{\pi}{4x} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \left(\frac{\arctan(\sqrt{x})}{x} + \frac{1}{2} \left(2 \arctan(\sqrt{x}) + \frac{2}{\sqrt{x}} \right) \right) - \frac{\pi}{4x}
 \end{aligned}$$

input `Int[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x^2),x]`

output `-1/4*Pi/x + (ArcTan[Sqrt[x]]/x + (2/Sqrt[x] + 2*ArcTan[Sqrt[x]])/2)/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 5361

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :>
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

rule 5682

```
Int[ArcTan[(v_) + (s_)*Sqrt[w_]]*(u_), x_Symbol] :> Simp[Pi*(s/4) Int[u
, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2
+ 1]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(27) = 54$.

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.39

method	result	size
default	$\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} + \frac{1}{2\sqrt{x}} + \frac{\operatorname{arctanh}(\sqrt{1+x})}{2} + \frac{\arctan(\sqrt{x})}{2} - \frac{\ln(1+\sqrt{1+x})}{4} + \frac{\ln(\sqrt{1+x}-1)}{4}$	57
parts	$\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x} + \frac{1}{2\sqrt{x}} + \frac{\operatorname{arctanh}(\sqrt{1+x})}{2} + \frac{\arctan(\sqrt{x})}{2} - \frac{\ln(1+\sqrt{1+x})}{4} + \frac{\ln(\sqrt{1+x}-1)}{4}$	57

input

```
int(-arctan(x^(1/2)-(1+x)^(1/2))/x^2,x,method=_RETURNVERBOSE)
```

output

```
arctan(x^(1/2)-(1+x)^(1/2))/x+1/2/x^(1/2)+1/2*arctanh((1+x)^(1/2))+1/2*arc
tan(x^(1/2))-1/4*ln(1+(1+x)^(1/2))+1/4*ln((1+x)^(1/2)-1)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^2} dx = -\frac{2(x+1)\arctan(\sqrt{x+1}-\sqrt{x})-\sqrt{x}}{2x}$$

input

```
integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^2,x, algorithm="fricas")
```

output

```
-1/2*(2*(x + 1)*arctan(sqrt(x + 1) - sqrt(x)) - sqrt(x))/x
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 537 vs. $2(31) = 62$.

Time = 43.71 (sec) , antiderivative size = 537, normalized size of antiderivative = 13.10

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^2} dx = -\frac{2x^{\frac{5}{2}}\sqrt{x+1}\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} + \frac{x^{\frac{5}{2}}}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} - \frac{4x^{\frac{3}{2}}\sqrt{x+1}\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} + \frac{x^{\frac{3}{2}}}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} - \frac{2\sqrt{x}\sqrt{x+1}\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} + \frac{2x^3\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} - \frac{x^2\sqrt{x+1}}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} + \frac{4x^2\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} - \frac{x\sqrt{x+1}}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2} + \frac{2x\operatorname{atan}(\sqrt{x} - \sqrt{x+1})}{-2x^{\frac{5}{2}}\sqrt{x+1} - 2x^{\frac{3}{2}}\sqrt{x+1} + 2x^3 + 2x^2}$$

input

```
integrate(-atan(x**(1/2)-(1+x)**(1/2))/x**2,x)
```

output

```
-2*x**(5/2)*sqrt(x + 1)*atan(sqrt(x) - sqrt(x + 1))/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) + x**(5/2)/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) - 4*x**(3/2)*sqrt(x + 1)*atan(sqrt(x) - sqrt(x + 1))/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) + x**(3/2)/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) - 2*sqrt(x)*sqrt(x + 1)*atan(sqrt(x) - sqrt(x + 1))/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) + 2*x**3*atan(sqrt(x) - sqrt(x + 1))/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) - x**2*sqrt(x + 1)/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) + 4*x**2*atan(sqrt(x) - sqrt(x + 1))/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) - x*sqrt(x + 1)/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2) + 2*x*atan(sqrt(x) - sqrt(x + 1))/(-2*x**(5/2)*sqrt(x + 1) - 2*x**(3/2)*sqrt(x + 1) + 2*x**3 + 2*x**2)
```

Maxima [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.71

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^2} dx = -\frac{\arctan(\sqrt{x+1} - \sqrt{x})}{x} + \frac{1}{2\sqrt{x}} + \frac{1}{2} \arctan(\sqrt{x})$$

input

```
integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^2,x, algorithm="maxima")
```

output

```
-arctan(sqrt(x + 1) - sqrt(x))/x + 1/2/sqrt(x) + 1/2*arctan(sqrt(x))
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.68

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^2} dx = \frac{\arctan(-\sqrt{x+1} + \sqrt{x})}{x} + \frac{1}{2\sqrt{x}} + \frac{1}{2} \arctan(\sqrt{x})$$

input

```
integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^2,x, algorithm="giac")
```

output

```
arctan(-sqrt(x + 1) + sqrt(x))/x + 1/2/sqrt(x) + 1/2*arctan(sqrt(x))
```

Mupad [B] (verification not implemented)

Time = 1.40 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.07

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^2} dx = -\frac{\operatorname{atan}(\sqrt{x+1} - \sqrt{x}) - \frac{\sqrt{x}}{2}}{x} + \frac{\ln\left(\frac{(-1+\sqrt{x}i)^2}{x+1}\right) i}{4}$$

input `int(atan((x + 1)^(1/2) - x^(1/2))/x^2,x)`output `(log((x^(1/2)*i - 1)^2/(x + 1))*i)/4 - (atan((x + 1)^(1/2) - x^(1/2)) - x^(1/2)/2)/x`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^2} dx = \frac{-2\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right) x - 2\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right) + \sqrt{x}}{2x}$$

input `int(-atan(x^(1/2)-(1+x)^(1/2))/x^2,x)`output `(- 2*atan(1/(sqrt(x + 1) + sqrt(x)))*x - 2*atan(1/(sqrt(x + 1) + sqrt(x))) + sqrt(x))/(2*x)`

3.132 $\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^3} dx$

Optimal result	989
Mathematica [A] (verified)	989
Rubi [A] (verified)	990
Maple [A] (verified)	992
Fricas [A] (verification not implemented)	992
Sympy [F(-1)]	993
Maxima [A] (verification not implemented)	993
Giac [A] (verification not implemented)	993
Mupad [B] (verification not implemented)	994
Reduce [B] (verification not implemented)	994

Optimal result

Integrand size = 21, antiderivative size = 50

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^3} dx = -\frac{\pi}{8x^2} + \frac{1}{12x^{3/2}} - \frac{1}{4\sqrt{x}} - \frac{\arctan(\sqrt{x})}{4} + \frac{\arctan(\sqrt{x})}{4x^2}$$

output

$$-1/8*\text{Pi}/x^2+1/12/x^{(3/2)}-1/4/x^{(1/2)}-1/4*\arctan(x^{(1/2)})+1/4*\arctan(x^{(1/2)})/x^2$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^3} dx = -\frac{\sqrt{x}(-1+3x)+3x^2\arctan(\sqrt{x})-6\arctan(\sqrt{x}-\sqrt{1+x})}{12x^2}$$

input

$$\text{Integrate}[-(\text{ArcTan}[\text{Sqrt}[x] - \text{Sqrt}[1 + x]]/x^3), x]$$

output

$$-1/12*(\text{Sqrt}[x]*(-1+3*x)+3*x^2*\text{ArcTan}[\text{Sqrt}[x]]-6*\text{ArcTan}[\text{Sqrt}[x]-\text{Sqrt}[1+x]])/x^2$$

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.12, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {25, 5682, 15, 5361, 61, 61, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int -\frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x^3} dx \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x^3} dx \\
 & \quad \downarrow \text{5682} \\
 & \frac{1}{4}\pi \int \frac{1}{x^3} dx - \frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x^3} dx \\
 & \quad \downarrow \text{15} \\
 & -\frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x^3} dx - \frac{\pi}{8x^2} \\
 & \quad \downarrow \text{5361} \\
 & \frac{1}{2} \left(\frac{\arctan(\sqrt{x})}{2x^2} - \frac{1}{4} \int \frac{1}{x^{5/2}(x+1)} dx \right) - \frac{\pi}{8x^2} \\
 & \quad \downarrow \text{61} \\
 & \frac{1}{2} \left(\frac{1}{4} \left(\int \frac{1}{x^{3/2}(x+1)} dx + \frac{2}{3x^{3/2}} \right) + \frac{\arctan(\sqrt{x})}{2x^2} \right) - \frac{\pi}{8x^2} \\
 & \quad \downarrow \text{61} \\
 & \frac{1}{2} \left(\frac{1}{4} \left(-\int \frac{1}{\sqrt{x}(x+1)} dx + \frac{2}{3x^{3/2}} - \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{2x^2} \right) - \frac{\pi}{8x^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{2} \left(\frac{1}{4} \left(-2 \int \frac{1}{x+1} d\sqrt{x} + \frac{2}{3x^{3/2}} - \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{2x^2} \right) - \frac{\pi}{8x^2}
 \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(-2 \arctan(\sqrt{x}) + \frac{2}{3x^{3/2}} - \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{2x^2} \right) - \frac{\pi}{8x^2}$$

input `Int[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x^3), x]`

output `-1/8*Pi/x^2 + ((2/(3*x^(3/2)) - 2/Sqrt[x] - 2*ArcTan[Sqrt[x]])/4 + ArcTan[Sqrt[x]]/(2*x^2))/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 61 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 5361

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

rule 5682

```
Int[ArcTan[(v_) + (s_)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u
, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2
+ 1]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{2x^2} + \frac{1}{12x^{\frac{3}{2}}} - \frac{1}{4\sqrt{x}} - \frac{\arctan(\sqrt{x})}{4}$	35
parts	$\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{2x^2} + \frac{1}{12x^{\frac{3}{2}}} - \frac{1}{4\sqrt{x}} - \frac{\arctan(\sqrt{x})}{4}$	35

input

```
int(-arctan(x^(1/2)-(1+x)^(1/2))/x^3,x,method=_RETURNVERBOSE)
```

output

```
1/2*arctan(x^(1/2)-(1+x)^(1/2))/x^2+1/12/x^(3/2)-1/4/x^(1/2)-1/4*arctan(x^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.70

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^3} dx = \frac{6(x^2-1)\arctan(\sqrt{x+1}-\sqrt{x})-(3x-1)\sqrt{x}}{12x^2}$$

input

```
integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^3,x, algorithm="fricas")
```

output

```
1/12*(6*(x^2 - 1)*arctan(sqrt(x + 1) - sqrt(x)) - (3*x - 1)*sqrt(x))/x^2
```

Sympy [F(-1)]

Timed out.

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^3} dx = \text{Timed out}$$

input `integrate(-atan(x**(1/2)-(1+x)**(1/2))/x**3,x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.68

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^3} dx = -\frac{1}{4\sqrt{x}} - \frac{\arctan(\sqrt{x+1} - \sqrt{x})}{2x^2} + \frac{1}{12x^{\frac{3}{2}}} - \frac{1}{4} \arctan(\sqrt{x})$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^3,x, algorithm="maxima")`output `-1/4/sqrt(x) - 1/2*arctan(sqrt(x + 1) - sqrt(x))/x^2 + 1/12/x^(3/2) - 1/4*arctan(sqrt(x))`**Giac [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.68

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^3} dx = -\frac{3x-1}{12x^{\frac{3}{2}}} + \frac{\arctan(-\sqrt{x+1} + \sqrt{x})}{2x^2} - \frac{1}{4} \arctan(\sqrt{x})$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^3,x, algorithm="giac")`output `-1/12*(3*x - 1)/x^(3/2) + 1/2*arctan(-sqrt(x + 1) + sqrt(x))/x^2 - 1/4*arctan(sqrt(x))`

Mupad [B] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.98

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^3} dx = -\frac{\operatorname{atan}(\sqrt{x+1}-\sqrt{x})}{2x^2} - \frac{\sqrt{x}}{12} + \frac{x^{3/2}}{4} + \frac{\ln\left(\frac{(\sqrt{x}-i)^2}{x+1}\right) 1i}{8}$$

input `int(atan((x + 1)^(1/2) - x^(1/2))/x^3,x)`output `(log((x^(1/2) - 1i)^2/(x + 1))*1i)/8 - (atan((x + 1)^(1/2) - x^(1/2))/2 - x^(1/2)/12 + x^(3/2)/4)/x^2`**Reduce [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.76

$$\begin{aligned} & \int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^3} dx \\ &= \frac{-6\operatorname{atan}(\sqrt{x+1} + \sqrt{x})x^2 - 6\operatorname{atan}\left(\frac{1}{\sqrt{x+1} + \sqrt{x}}\right) - 3\sqrt{x}x + \sqrt{x}}{12x^2} \end{aligned}$$

input `int(-atan(x^(1/2)-(1+x)^(1/2))/x^3,x)`output `(- 6*atan(sqrt(x + 1) + sqrt(x))*x**2 - 6*atan(1/(sqrt(x + 1) + sqrt(x))) - 3*sqrt(x)*x + sqrt(x))/(12*x**2)`

3.133 $\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^4} dx$

Optimal result	995
Mathematica [A] (verified)	995
Rubi [A] (verified)	996
Maple [A] (verified)	998
Fricas [A] (verification not implemented)	999
Sympy [F(-1)]	999
Maxima [A] (verification not implemented)	999
Giac [A] (verification not implemented)	1000
Mupad [B] (verification not implemented)	1000
Reduce [B] (verification not implemented)	1001

Optimal result

Integrand size = 21, antiderivative size = 59

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^4} dx = -\frac{\pi}{12x^3} + \frac{1}{30x^{5/2}} - \frac{1}{18x^{3/2}} + \frac{1}{6\sqrt{x}} + \frac{\arctan(\sqrt{x})}{6} + \frac{\arctan(\sqrt{x})}{6x^3}$$

output

```
-1/12*Pi/x^3+1/30/x^(5/2)-1/18/x^(3/2)+1/6/x^(1/2)+1/6*arctan(x^(1/2))+1/6*arctan(x^(1/2))/x^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.86

$$\int -\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{x^4} dx = \frac{1}{90} \left(-\frac{-3+5x-15x^2}{x^{5/2}} + 15 \arctan(\sqrt{x}) + \frac{30 \arctan(\sqrt{x}-\sqrt{1+x})}{x^3} \right)$$

input

```
Integrate[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x^4), x]
```

output

$$\frac{(-((-3 + 5x - 15x^2)/x^{5/2}) + 15\text{ArcTan}[\text{Sqrt}[x]] + (30\text{ArcTan}[\text{Sqrt}[x] - \text{Sqrt}[1 + x]])/x^3)/90}$$
Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {25, 5682, 15, 5361, 61, 61, 61, 73, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int -\frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x^4} dx \\ & \quad \downarrow \text{25} \\ & -\int \frac{\arctan(\sqrt{x} - \sqrt{x+1})}{x^4} dx \\ & \quad \downarrow \text{5682} \\ & \frac{1}{4}\pi \int \frac{1}{x^4} dx - \frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x^4} dx \\ & \quad \downarrow \text{15} \\ & -\frac{1}{2} \int \frac{\arctan(\sqrt{x})}{x^4} dx - \frac{\pi}{12x^3} \\ & \quad \downarrow \text{5361} \\ & \frac{1}{2} \left(\frac{\arctan(\sqrt{x})}{3x^3} - \frac{1}{6} \int \frac{1}{x^{7/2}(x+1)} dx \right) - \frac{\pi}{12x^3} \\ & \quad \downarrow \text{61} \\ & \frac{1}{2} \left(\frac{1}{6} \left(\int \frac{1}{x^{5/2}(x+1)} dx + \frac{2}{5x^{5/2}} \right) + \frac{\arctan(\sqrt{x})}{3x^3} \right) - \frac{\pi}{12x^3} \\ & \quad \downarrow \text{61} \\ & \frac{1}{2} \left(\frac{1}{6} \left(-\int \frac{1}{x^{3/2}(x+1)} dx - \frac{2}{3x^{3/2}} + \frac{2}{5x^{5/2}} \right) + \frac{\arctan(\sqrt{x})}{3x^3} \right) - \frac{\pi}{12x^3} \end{aligned}$$

$$\begin{aligned} & \downarrow 61 \\ & \frac{1}{2} \left(\frac{1}{6} \left(\int \frac{1}{\sqrt{x}(x+1)} dx - \frac{2}{3x^{3/2}} + \frac{2}{5x^{5/2}} + \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{3x^3} \right) - \frac{\pi}{12x^3} \\ & \downarrow 73 \\ & \frac{1}{2} \left(\frac{1}{6} \left(2 \int \frac{1}{x+1} d\sqrt{x} - \frac{2}{3x^{3/2}} + \frac{2}{5x^{5/2}} + \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{3x^3} \right) - \frac{\pi}{12x^3} \\ & \downarrow 216 \\ & \frac{1}{2} \left(\frac{1}{6} \left(2 \arctan(\sqrt{x}) - \frac{2}{3x^{3/2}} + \frac{2}{5x^{5/2}} + \frac{2}{\sqrt{x}} \right) + \frac{\arctan(\sqrt{x})}{3x^3} \right) - \frac{\pi}{12x^3} \end{aligned}$$

input `Int[-(ArcTan[Sqrt[x] - Sqrt[1 + x]]/x^4), x]`

output `-1/12*Pi/x^3 + (ArcTan[Sqrt[x]]/(3*x^3) + (2/(5*x^(5/2)) - 2/(3*x^(3/2)) + 2/Sqrt[x] + 2*ArcTan[Sqrt[x]])/6)/2`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 5361

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*(x_)^(m_.), x_Symbol] :=
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Simp[b*c*n*(p/(m +
1)) Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x],
x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &
& IntegerQ[m])) && NeQ[m, -1]
```

rule 5682

```
Int[ArcTan[(v_) + (s_.)*Sqrt[w_]]*(u_), x_Symbol] := Simp[Pi*(s/4) Int[u
, x], x] + Simp[1/2 Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2
+ 1]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.68

method	result	size
default	$\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{3x^3} + \frac{1}{30x^{\frac{5}{2}}} - \frac{1}{18x^{\frac{3}{2}}} + \frac{1}{6\sqrt{x}} + \frac{\arctan(\sqrt{x})}{6}$	40
parts	$\frac{\arctan(\sqrt{x}-\sqrt{1+x})}{3x^3} + \frac{1}{30x^{\frac{5}{2}}} - \frac{1}{18x^{\frac{3}{2}}} + \frac{1}{6\sqrt{x}} + \frac{\arctan(\sqrt{x})}{6}$	40

input

```
int(-arctan(x^(1/2)-(1+x)^(1/2))/x^4,x,method=_RETURNVERBOSE)
```

output

```
1/3*arctan(x^(1/2)-(1+x)^(1/2))/x^3+1/30/x^(5/2)-1/18/x^(3/2)+1/6/x^(1/2)+
1/6*arctan(x^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.68

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^4} dx$$

$$= -\frac{30(x^3 + 1)\arctan(\sqrt{x+1} - \sqrt{x}) - (15x^2 - 5x + 3)\sqrt{x}}{90x^3}$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^4,x, algorithm="fricas")`output `-1/90*(30*(x^3 + 1)*arctan(sqrt(x + 1) - sqrt(x)) - (15*x^2 - 5*x + 3)*sqrt(x))/x^3`**Sympy [F(-1)]**

Timed out.

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^4} dx = \text{Timed out}$$

input `integrate(-atan(x**(1/2)-(1+x)**(1/2))/x**4,x)`output `Timed out`**Maxima [A] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.66

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^4} dx = \frac{1}{6\sqrt{x}} - \frac{1}{18x^{\frac{3}{2}}} - \frac{\arctan(\sqrt{x+1} - \sqrt{x})}{3x^3}$$

$$+ \frac{1}{30x^{\frac{5}{2}}} + \frac{1}{6}\arctan(\sqrt{x})$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^4,x, algorithm="maxima")`

output $\frac{1}{6}\sqrt{x} - \frac{1}{18}x^{3/2} - \frac{1}{3}\arctan(\sqrt{x+1} - \sqrt{x})/x^3 + \frac{1}{30}x^{5/2} + \frac{1}{6}\arctan(\sqrt{x})$

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.66

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^4} dx = \frac{15x^2 - 5x + 3}{90x^{5/2}} + \frac{\arctan(-\sqrt{x+1} + \sqrt{x})}{3x^3} + \frac{1}{6}\arctan(\sqrt{x})$$

input `integrate(-arctan(x^(1/2)-(1+x)^(1/2))/x^4,x, algorithm="giac")`

output $\frac{1}{90}(15x^2 - 5x + 3)/x^{5/2} + \frac{1}{3}\arctan(-\sqrt{x+1} + \sqrt{x})/x^3 + \frac{1}{6}\arctan(\sqrt{x})$

Mupad [B] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.95

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^4} dx = -\frac{\frac{\operatorname{atan}(\sqrt{x+1}-\sqrt{x})}{3} - \frac{\sqrt{x}}{30} + \frac{x^{3/2}}{18} - \frac{x^{5/2}}{6}}{x^3} + \frac{\ln\left(\frac{(-1+\sqrt{x}i)^2}{x+1}\right) \operatorname{li}}{12}$$

input `int(atan((x + 1)^(1/2) - x^(1/2))/x^4,x)`

output $\frac{\log((x^{1/2}i - 1)^2/(x+1)i)}{12} - \frac{\operatorname{atan}((x+1)^{1/2} - x^{1/2})}{3} - \frac{x^{1/2}}{30} + \frac{x^{3/2}}{18} - \frac{x^{5/2}}{6}/x^3$

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.83

$$\int -\frac{\arctan(\sqrt{x} - \sqrt{1+x})}{x^4} dx$$

$$= \frac{-30\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right)x^3 - 30\operatorname{atan}\left(\frac{1}{\sqrt{x+1}+\sqrt{x}}\right) + 15\sqrt{x}x^2 - 5\sqrt{x}x + 3\sqrt{x}}{90x^3}$$

input `int(-atan(x^(1/2)-(1+x)^(1/2))/x^4,x)`output `(- 30*atan(1/(sqrt(x + 1) + sqrt(x)))*x**3 - 30*atan(1/(sqrt(x + 1) + sqrt(x))) + 15*sqrt(x)*x**2 - 5*sqrt(x)*x + 3*sqrt(x))/(90*x**3)`

3.134
$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

Optimal result	1002
Mathematica [A] (verified)	1002
Rubi [A] (verified)	1003
Maple [A] (verified)	1004
Fricas [B] (verification not implemented)	1005
Sympy [F]	1005
Maxima [F(-2)]	1006
Giac [F]	1006
Mupad [B] (verification not implemented)	1006
Reduce [B] (verification not implemented)	1007

Optimal result

Integrand size = 39, antiderivative size = 63

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^{1+m}}{c(1+m)\sqrt{d-\frac{c^2dx^2}{a}}}$$

output

```
(-c^2*x^2+a)^(1/2)*arctan(c*x/(-c^2*x^2+a)^(1/2))^(1+m)/c/(1+m)/(d-c^2*d*x^2/a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^{1+m}}{c(1+m)\sqrt{d-\frac{c^2dx^2}{a}}}$$

input

```
Integrate[ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^m/Sqrt[d - (c^2*d*x^2)/a],x]
```

output

```
(Sqrt[a - c^2*x^2]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^(1 + m))/(c*(1 + m)*Sqrt[d - (c^2*d*x^2)/a])
```

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

↓ 5680

$$\frac{\sqrt{a-c^2x^2} \int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{a-c^2x^2}} dx}{\sqrt{d-\frac{c^2dx^2}{a}}}$$

↓ 5678

$$\frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^{m+1}}{c(m+1)\sqrt{d-\frac{c^2dx^2}{a}}}$$

input

```
Int[ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^m/Sqrt[d - (c^2*d*x^2)/a], x]
```

output

```
(Sqrt[a - c^2*x^2]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^(1 + m))/(c*(1 + m)*Sqrt[d - (c^2*d*x^2)/a])
```

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] :> Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] :> Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m}
, x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.94

method	result	size
default	$\frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^{1+m} \sqrt{-c^2x^2+a}}{c(1+m)\sqrt{\frac{d(-c^2x^2+a)}{a}}}$	59

input

```
int(arctan(c*x/(-c^2*x^2+a)^(1/2))^(1+m)/(d-c^2*d*x^2/a)^(1/2),x,method=_RETUR
NVERBOSE)
```

output

```
arctan(c*x/(-c^2*x^2+a)^(1/2))^(1+m)/c/(1+m)*(-c^2*x^2+a)^(1/2)/(d*(-c^2*x
^2+a)/a)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 126 vs. $2(57) = 114$.

Time = 0.13 (sec) , antiderivative size = 126, normalized size of antiderivative = 2.00

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

$$= -\frac{\sqrt{-c^2x^2+a} \arctan\left(\frac{\sqrt{-c^2x^2+acx}}{c^2x^2-a}\right)^m \sqrt{-\frac{c^2dx^2-ad}{a}} \arctan\left(\frac{\sqrt{-c^2x^2+acx}}{c^2x^2-a}\right)}{acdm + acd - (c^3dm + c^3d)x^2}$$

input

```
integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))^m/(d-c^2*d*x^2/a)^(1/2),x, algorithm="fricas")
```

output

```
-sqrt(-c^2*x^2 + a)*a*(-arctan(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a)))^m*sqrt(-(c^2*d*x^2 - a*d)/a)*arctan(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a))/(a*c*d*m + a*c*d - (c^3*d*m + c^3*d)*x^2)
```

Sympy [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\operatorname{atan}^m\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{-d\left(-1+\frac{c^2x^2}{a}\right)}} dx$$

input

```
integrate(atan(c*x/sqrt(a - c**2*x**2))**m/sqrt(-d*(-1 + c**2*x**2/a)), x)
```

output

```
Integral(atan(c*x/sqrt(a - c**2*x**2))**m/sqrt(-d*(-1 + c**2*x**2/a)), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))^m/(d-c^2*d*x^2/a)^(1/2), x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^m}{\sqrt{-\frac{c^2dx^2}{a}+d}} dx$$

input `integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))^m/(d-c^2*d*x^2/a)^(1/2), x, algorithm="giac")`

output `integrate(arctan(c*x/sqrt(-c^2*x^2+a))^m/sqrt(-c^2*d*x^2/a+d), x)`

Mupad [B] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.90

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\operatorname{atan}\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^{m+1} \sqrt{a-c^2x^2}}{c(m+1) \sqrt{d-\frac{c^2dx^2}{a}}}$$

input `int(atan((c*x)/(a-c^2*x^2)^(1/2))^m/(d-(c^2*d*x^2)/a)^(1/2), x)`

output $(\operatorname{atan}((c*x)/(a - c^2*x^2))^{(1/2)})^{(m + 1)}*(a - c^2*x^2)^{(1/2)}/(c*(m + 1)*(d - (c^2*d*x^2)/a)^{(1/2)})$

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^m}{\sqrt{d - \frac{c^2dx^2}{a}}} dx = \frac{\sqrt{d} \sqrt{a} \operatorname{atan}\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^m \operatorname{atan}\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)}{cd(m+1)}$$

input $\operatorname{int}(\operatorname{atan}(c*x/(-c^2*x^2+a))^{(1/2)})^m/(d-c^2*d*x^2/a)^{(1/2)}, x)$

output $(\operatorname{sqrt}(d)*\operatorname{sqrt}(a)*\operatorname{atan}((c*x)/\operatorname{sqrt}(a - c**2*x**2))**m*\operatorname{atan}((c*x)/\operatorname{sqrt}(a - c**2*x**2)))/(c*d*(m + 1))$

$$3.135 \quad \int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

Optimal result	1008
Mathematica [A] (verified)	1008
Rubi [A] (verified)	1009
Maple [A] (verified)	1010
Fricas [F]	1010
Sympy [F]	1011
Maxima [F]	1011
Giac [F]	1012
Mupad [F(-1)]	1012
Reduce [B] (verification not implemented)	1012

Optimal result

Integrand size = 39, antiderivative size = 59

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^3}{3c\sqrt{d-\frac{c^2dx^2}{a}}}$$

output

```
1/3*(-c^2*x^2+a)^(1/2)*arctan(c*x/(-c^2*x^2+a)^(1/2))^3/c/(d-c^2*d*x^2/a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^3}{3c\sqrt{d-\frac{c^2dx^2}{a}}}$$

input

```
Integrate[ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^2/Sqrt[d - (c^2*d*x^2)/a],x]
```

output $(\text{Sqrt}[a - c^2*x^2]*\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]^3)/(3*c*\text{Sqrt}[d - (c^2*d*x^2)/a])$

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d - \frac{c^2dx^2}{a}}} dx$$

↓ 5680

$$\frac{\sqrt{a - c^2x^2} \int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{a-c^2x^2}} dx}{\sqrt{d - \frac{c^2dx^2}{a}}}$$

↓ 5678

$$\frac{\sqrt{a - c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^3}{3c\sqrt{d - \frac{c^2dx^2}{a}}}$$

input $\text{Int}[\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]^2/\text{Sqrt}[d - (c^2*d*x^2)/a], x]$

output $(\text{Sqrt}[a - c^2*x^2]*\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]^3)/(3*c*\text{Sqrt}[d - (c^2*d*x^2)/a])$

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.97

method	result	size
default	$\frac{\sqrt{\frac{d(-c^2x^2+a)}{a}} \arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^3 a}{3\sqrt{-c^2x^2+a} dc}$	57

input

```
int(arctan(c*x/(-c^2*x^2+a)^(1/2))^2/(d-c^2*d*x^2/a)^(1/2),x,method=_RETUR
NVERBOSE)
```

output

```
1/3/(-c^2*x^2+a)^(1/2)*(d*(-c^2*x^2+a)/a)^(1/2)/d/c*arctan(c*x/(-c^2*x^2+a)
^(1/2))^3*a
```

Fricas [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^2}{\sqrt{-\frac{c^2dx^2}{a}+d}} dx$$

input

```
integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))^2/(d-c^2*d*x^2/a)^(1/2),x, algori
thm="fricas")
```

output `integral(-a*sqrt(-(c^2*d*x^2 - a*d)/a)*arctan(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a))^2/(c^2*d*x^2 - a*d), x)`

Sympy [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d - \frac{c^2dx^2}{a}}} dx = \int \frac{\operatorname{atan}^2\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{-d\left(-1 + \frac{c^2x^2}{a}\right)}} dx$$

input `integrate(atan(c*x/(-c**2*x**2+a)**(1/2))**2/(d-c**2*d*x**2/a)**(1/2), x)`

output `Integral(atan(c*x/sqrt(a - c**2*x**2))**2/sqrt(-d*(-1 + c**2*x**2/a)), x)`

Maxima [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d - \frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^2}{\sqrt{-\frac{c^2dx^2}{a} + d}} dx$$

input `integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))^2/(d-c^2*d*x^2/a)^(1/2), x, algorithm="maxima")`

output `integrate(arctan(c*x/sqrt(-c^2*x^2 + a))^2/sqrt(-c^2*d*x^2/a + d), x)`

Giac [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^2}{\sqrt{-\frac{c^2dx^2}{a}+d}} dx$$

input `integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))^2/(d-c^2*d*x^2/a)^(1/2), x, algorithm="giac")`

output `integrate(arctan(c*x/sqrt(-c^2*x^2 + a))^2/sqrt(-c^2*d*x^2/a + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\operatorname{atan}\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

input `int(atan((c*x)/(a - c^2*x^2)^(1/2))^2/(d - (c^2*d*x^2)/a)^(1/2), x)`

output `int(atan((c*x)/(a - c^2*x^2)^(1/2))^2/(d - (c^2*d*x^2)/a)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.53

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{d}\sqrt{a}\operatorname{atan}\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^3}{3cd}$$

input `int(atan(c*x/(-c^2*x^2+a)^(1/2))^2/(d-c^2*d*x^2/a)^(1/2), x)`

output `(sqrt(d)*sqrt(a)*atan((c*x)/sqrt(a - c**2*x**2))**3)/(3*c*d)`

3.136
$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

Optimal result	1013
Mathematica [A] (verified)	1013
Rubi [A] (verified)	1014
Maple [A] (verified)	1015
Fricas [F]	1015
Sympy [F]	1016
Maxima [F]	1016
Giac [F]	1017
Mupad [F(-1)]	1017
Reduce [B] (verification not implemented)	1017

Optimal result

Integrand size = 37, antiderivative size = 59

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{2c\sqrt{d-\frac{c^2dx^2}{a}}}$$

output `1/2*(-c^2*x^2+a)^(1/2)*arctan(c*x/(-c^2*x^2+a)^(1/2))^2/c/(d-c^2*d*x^2/a)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{2c\sqrt{d-\frac{c^2dx^2}{a}}}$$

input `Integrate[ArcTan[(c*x)/Sqrt[a - c^2*x^2]]/Sqrt[d - (c^2*d*x^2)/a],x]`

output

```
(Sqrt[a - c^2*x^2]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^2)/(2*c*Sqrt[d - (c^2*d*x^2)/a])
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.054$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

↓ 5680

$$\frac{\sqrt{a-c^2x^2} \int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{a-c^2x^2}} dx}{\sqrt{d-\frac{c^2dx^2}{a}}}$$

↓ 5678

$$\frac{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2}{2c\sqrt{d-\frac{c^2dx^2}{a}}}$$

input

```
Int[ArcTan[(c*x)/Sqrt[a - c^2*x^2]]/Sqrt[d - (c^2*d*x^2)/a], x]
```

output

```
(Sqrt[a - c^2*x^2]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^2)/(2*c*Sqrt[d - (c^2*d*x^2)/a])
```

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.97

method	result	size
default	$\frac{\sqrt{\frac{d(-c^2x^2+a)}{a}} \arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^2}{2\sqrt{-c^2x^2+a} dc} a$	57

input

```
int(arctan(c*x/(-c^2*x^2+a)^(1/2))/(d-c^2*d*x^2/a)^(1/2),x,method=_RETURNV
ERBOSE)
```

output

```
1/2/(-c^2*x^2+a)^(1/2)*(d*(-c^2*x^2+a)/a)^(1/2)/d*c*arctan(c*x/(-c^2*x^2+a
)^(1/2))^2*a
```

Fricas [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)}{\sqrt{-\frac{c^2dx^2}{a}+d}} dx$$

input

```
integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))/(d-c^2*d*x^2/a)^(1/2),x, algorith
m="fricas")
```


output

```
integral(a*sqrt(-(c^2*d*x^2 - a*d)/a)*arctan(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a))/(c^2*d*x^2 - a*d), x)
```

Sympy [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d - \frac{c^2dx^2}{a}}} dx = \int \frac{\operatorname{atan}\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{-d\left(-1 + \frac{c^2x^2}{a}\right)}} dx$$

input

```
integrate(atan(c*x/(-c**2*x**2+a)**(1/2))/(d-c**2*d*x**2/a)**(1/2), x)
```

output

```
Integral(atan(c*x/sqrt(a - c**2*x**2))/sqrt(-d*(-1 + c**2*x**2/a)), x)
```

Maxima [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d - \frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)}{\sqrt{-\frac{c^2dx^2}{a} + d}} dx$$

input

```
integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))/(d-c^2*d*x^2/a)^(1/2), x, algorithm m="maxima")
```

output

```
integrate(arctan(c*x/sqrt(-c^2*x^2 + a))/sqrt(-c^2*d*x^2/a + d), x)
```

Giac [F]

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)}{\sqrt{-\frac{c^2dx^2}{a}+d}} dx$$

input `integrate(arctan(c*x/(-c^2*x^2+a)^(1/2))/(d-c^2*d*x^2/a)^(1/2),x, algorithm m="giac")`

output `integrate(arctan(c*x/sqrt(-c^2*x^2 + a))/sqrt(-c^2*d*x^2/a + d), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \int \frac{\operatorname{atan}\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx$$

input `int(atan((c*x)/(a - c^2*x^2)^(1/2))/(d - (c^2*d*x^2)/a)^(1/2),x)`

output `int(atan((c*x)/(a - c^2*x^2)^(1/2))/(d - (c^2*d*x^2)/a)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.53

$$\int \frac{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)}{\sqrt{d-\frac{c^2dx^2}{a}}} dx = \frac{\sqrt{d}\sqrt{a}\operatorname{atan}\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^2}{2cd}$$

input `int(atan(c*x/(-c^2*x^2+a)^(1/2))/(d-c^2*d*x^2/a)^(1/2),x)`

output `(sqrt(d)*sqrt(a)*atan((c*x)/sqrt(a - c**2*x**2))**2)/(2*c*d)`

3.137
$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx$$

Optimal result	1018
Mathematica [A] (verified)	1018
Rubi [A] (verified)	1019
Maple [A] (verified)	1020
Fricas [A] (verification not implemented)	1020
Sympy [F]	1021
Maxima [F]	1021
Giac [F]	1022
Mupad [B] (verification not implemented)	1022
Reduce [B] (verification not implemented)	1022

Optimal result

Integrand size = 39, antiderivative size = 55

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \frac{\sqrt{a - c^2 x^2} \log\left(\arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)\right)}{c \sqrt{d - \frac{c^2 dx^2}{a}}}$$

output $(-c^2*x^2+a)^{(1/2)}*\ln(\arctan(c*x/(-c^2*x^2+a)^{(1/2)}))/c/(d-c^2*d*x^2/a)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \frac{\sqrt{a - c^2 x^2} \log\left(\arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)\right)}{c \sqrt{d - \frac{c^2 dx^2}{a}}}$$

input `Integrate[1/(Sqrt[d - (c^2*d*x^2)/a]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]),x]`

output $(\text{Sqrt}[a - c^2*x^2]*\text{Log}[\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]])/(c*\text{Sqrt}[d - (c^2*d*x^2)/a])$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {5680, 5676}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right) \sqrt{d - \frac{c^2dx^2}{a}}} dx$$

↓ 5680

$$\frac{\sqrt{a - c^2x^2} \int \frac{1}{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)} dx}{\sqrt{d - \frac{c^2dx^2}{a}}}$$

↓ 5676

$$\frac{\sqrt{a - c^2x^2} \log\left(\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)\right)}{c\sqrt{d - \frac{c^2dx^2}{a}}}$$

input $\text{Int}[1/(\text{Sqrt}[d - (c^2*d*x^2)/a]*\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]),x]$

output $(\text{Sqrt}[a - c^2*x^2]*\text{Log}[\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]])/(c*\text{Sqrt}[d - (c^2*d*x^2)/a])$

Defintions of rubi rules used

rule 5676

```
Int[1/(ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*Sqrt[(a_.) + (b_.)*(x_)^2]), x_Symbol] :> Simp[(1/c)*Log[ArcTan[c*(x/Sqrt[a + b*x^2])]], x] /; FreeQ[{a, b, c}, x] && EqQ[b + c^2, 0]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] :> Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{\sqrt{\frac{d(-c^2x^2+a)}{a}} \ln\left(\arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)\right) a}{\sqrt{-c^2x^2+a} dc}$	55

input

```
int(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2)),x,method=_RETURNVERBOSE)
```

output

```
1/(-c^2*x^2+a)^(1/2)*(d*(-c^2*x^2+a)/a)^(1/2)/d/c*ln(arctan(c*x/(-c^2*x^2+a)^(1/2)))*a
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.51

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx$$

$$= -\frac{\sqrt{-c^2 x^2 + a} a \sqrt{-\frac{c^2 dx^2 - ad}{a}} \log\left(2 \arctan\left(\frac{\sqrt{-c^2 x^2 + acx}}{c^2 x^2 - a}\right)\right)}{c^3 dx^2 - acd}$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2)),x, algorithm="fricas")`

output `-sqrt(-c^2*x^2 + a)*a*sqrt(-(c^2*d*x^2 - a*d)/a)*log(2*arctan(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a)))/(c^3*d*x^2 - a*c*d)`

Sympy [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \int \frac{1}{\sqrt{-d\left(-1 + \frac{c^2 x^2}{a}\right)} \operatorname{atan}\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx$$

input `integrate(1/(d-c**2*d*x**2/a)**(1/2)/atan(c*x/(-c**2*x**2+a)**(1/2)),x)`

output `Integral(1/(sqrt(-d*(-1 + c**2*x**2/a))*atan(c*x/sqrt(a - c**2*x**2))), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \int \frac{1}{\sqrt{-\frac{c^2 dx^2}{a} + d} \arctan\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right)} dx$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2)),x, algorithm="maxima")`

output `integrate(1/(sqrt(-c^2*d*x^2/a + d)*arctan(c*x/sqrt(-c^2*x^2 + a))), x)`

Giac [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \int \frac{1}{\sqrt{-\frac{c^2 dx^2}{a} + d} \arctan\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right)} dx$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2)),x, algorithm="giac")`

output `integrate(1/(sqrt(-c^2*d*x^2/a + d)*arctan(c*x/sqrt(-c^2*x^2 + a))), x)`

Mupad [B] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \frac{\ln\left(\operatorname{atan}\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)\right) \sqrt{a - c^2 x^2}}{c \sqrt{d - \frac{c^2 dx^2}{a}}}$$

input `int(1/(atan((c*x)/(a - c^2*x^2)^(1/2))*(d - (c^2*d*x^2)/a)^(1/2)),x)`

output `(log(atan((c*x)/(a - c^2*x^2)^(1/2)))*(a - c^2*x^2)^(1/2))/(c*(d - (c^2*d*x^2)/a)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.53

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx = \frac{\sqrt{d} \sqrt{a} \log\left(\operatorname{atan}\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right)\right)}{cd}$$

input `int(1/(d-c^2*d*x^2/a)^(1/2)/atan(c*x/(-c^2*x^2+a)^(1/2)),x)`

output $(\sqrt{d}\sqrt{a}\log(\operatorname{atan}((c*x)/\sqrt{a - c**2*x**2}))) / (c*d)$

3.138
$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx$$

Optimal result	1024
Mathematica [A] (verified)	1024
Rubi [A] (verified)	1025
Maple [A] (verified)	1026
Fricas [A] (verification not implemented)	1026
Sympy [F]	1027
Maxima [A] (verification not implemented)	1027
Giac [F]	1027
Mupad [B] (verification not implemented)	1028
Reduce [B] (verification not implemented)	1028

Optimal result

Integrand size = 39, antiderivative size = 57

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = -\frac{\sqrt{a - c^2 x^2}}{c \sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)}$$

output `-(-c^2*x^2+a)^(1/2)/c/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = -\frac{\sqrt{a - c^2 x^2}}{c \sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)}$$

input `Integrate[1/(Sqrt[d - (c^2*d*x^2)/a]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^2),x]`

output `-(Sqrt[a - c^2*x^2]/(c*Sqrt[d - (c^2*d*x^2)/a]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]))`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2 \sqrt{d-\frac{c^2dx^2}{a}}} dx$$

↓ 5680

$$\frac{\sqrt{a-c^2x^2} \int \frac{1}{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2} dx}{\sqrt{d-\frac{c^2dx^2}{a}}}$$

↓ 5678

$$-\frac{\sqrt{a-c^2x^2}}{c \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right) \sqrt{d-\frac{c^2dx^2}{a}}}$$

input `Int[1/(Sqrt[d - (c^2*d*x^2)/a]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^2),x]`

output `-(Sqrt[a - c^2*x^2]/(c*Sqrt[d - (c^2*d*x^2)/a]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]))`

Defintions of rubi rules used

rule 5678 `Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]`

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m}
, x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{\sqrt{\frac{d(-c^2x^2+a)}{a}} a}{\sqrt{-c^2x^2+a} dc \arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)}$	57

input

```
int(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^2,x,method=_RET
URNVERBOSE)
```

output

```
-1/(-c^2*x^2+a)^(1/2)*(d*(-c^2*x^2+a)/a)^(1/2)/d/c*a/arctan(c*x/(-c^2*x^2+
a)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.44

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = -\frac{\sqrt{-c^2 x^2 + a} a \sqrt{-\frac{c^2 dx^2 - ad}{a}}}{(c^3 dx^2 - acd) \arctan\left(\frac{\sqrt{-c^2 x^2 + acx}}{c^2 x^2 - a}\right)}$$

input

```
integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^2,x, algo
rithm="fricas")
```

output

```
-sqrt(-c^2*x^2 + a)*a*sqrt(-c^2*d*x^2 - a*d)/a/((c^3*d*x^2 - a*c*d)*arct
an(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a))
```

Sympy [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = \int \frac{1}{\sqrt{-d\left(-1 + \frac{c^2 x^2}{a}\right)} \operatorname{atan}^2\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx$$

input `integrate(1/(d-c**2*d*x**2/a)**(1/2)/atan(c*x/(-c**2*x**2+a)**(1/2))**2,x)`

output `Integral(1/(sqrt(-d*(-1 + c**2*x**2/a))*atan(c*x/sqrt(a - c**2*x**2))**2), x)`

Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.51

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = -\frac{\sqrt{a}}{c\sqrt{d} \arctan\left(cx, \sqrt{-c^2 x^2 + a}\right)}$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^2,x, algorithm="maxima")`

output `-sqrt(a)/(c*sqrt(d)*arctan2(c*x, sqrt(-c^2*x^2 + a)))`

Giac [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = \int \frac{1}{\sqrt{-\frac{c^2 dx^2}{a} + d} \arctan\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right)^2} dx$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^2,x, algorithm="giac")`

output `integrate(1/(sqrt(-c^2*d*x^2/a + d)*arctan(c*x/sqrt(-c^2*x^2 + a))^2), x)`

Mupad [B] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = -\frac{\sqrt{a - c^2 x^2}}{c \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right) \sqrt{d - \frac{c^2 dx^2}{a}}}$$

input `int(1/(atan((c*x)/(a - c^2*x^2)^(1/2)))^2*(d - (c^2*d*x^2)/a)^(1/2),x)`

output `-(a - c^2*x^2)^(1/2)/(c*atan((c*x)/(a - c^2*x^2)^(1/2))*(d - (c^2*d*x^2)/a)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.54

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2} dx = -\frac{\sqrt{d} \sqrt{a}}{\arctan\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right) cd}$$

input `int(1/(d-c^2*d*x^2/a)^(1/2)/atan(c*x/(-c^2*x^2+a)^(1/2))^2,x)`

output `(- sqrt(d)*sqrt(a))/(atan((c*x)/sqrt(a - c**2*x**2))*c*d)`

3.139
$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx$$

Optimal result	1029
Mathematica [A] (verified)	1029
Rubi [A] (verified)	1030
Maple [A] (verified)	1031
Fricas [A] (verification not implemented)	1031
Sympy [F]	1032
Maxima [A] (verification not implemented)	1032
Giac [F]	1033
Mupad [B] (verification not implemented)	1033
Reduce [B] (verification not implemented)	1033

Optimal result

Integrand size = 39, antiderivative size = 59

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = -\frac{\sqrt{a - c^2 x^2}}{2c \sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2}$$

output `-1/2*(-c^2*x^2+a)^(1/2)/c/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^2`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = -\frac{\sqrt{a - c^2 x^2}}{2c \sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2}$$

input `Integrate[1/(Sqrt[d - (c^2*d*x^2)/a]*ArcTan[(c*x)/Sqrt[a - c^2*x^2]]^3),x]`

output $-1/2*\text{Sqrt}[a - c^2*x^2]/(c*\text{Sqrt}[d - (c^2*d*x^2)/a]*\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]^2)$

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^3 \sqrt{d - \frac{c^2dx^2}{a}}} dx$$

↓ 5680

$$\frac{\sqrt{a-c^2x^2} \int \frac{1}{\sqrt{a-c^2x^2} \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^3} dx}{\sqrt{d - \frac{c^2dx^2}{a}}}$$

↓ 5678

$$-\frac{\sqrt{a-c^2x^2}}{2c \arctan\left(\frac{cx}{\sqrt{a-c^2x^2}}\right)^2 \sqrt{d - \frac{c^2dx^2}{a}}}$$

input $\text{Int}[1/(\text{Sqrt}[d - (c^2*d*x^2)/a]*\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]^3), x]$

output $-1/2*\text{Sqrt}[a - c^2*x^2]/(c*\text{Sqrt}[d - (c^2*d*x^2)/a]*\text{ArcTan}[(c*x)/\text{Sqrt}[a - c^2*x^2]]^2)$

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.97

method	result	size
default	$-\frac{\sqrt{\frac{d(-c^2x^2+a)}{a}}}{2\sqrt{-c^2x^2+a} d \arctan\left(\frac{cx}{\sqrt{-c^2x^2+a}}\right)^2}$	57

input

```
int(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^3,x,method=_RET
URNVERBOSE)
```

output

```
-1/2/(-c^2*x^2+a)^(1/2)*(d*(-c^2*x^2+a)/a)^(1/2)/d/c*a/arctan(c*x/(-c^2*x^
2+a)^(1/2))^2
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.39

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = \frac{\sqrt{-c^2 x^2 + a} a \sqrt{-\frac{c^2 dx^2 - ad}{a}}}{2(c^3 dx^2 - acd) \arctan\left(\frac{\sqrt{-c^2 x^2 + acx}}{c^2 x^2 - a}\right)^2}$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^3,x, algo
rithm="fricas")`

output `1/2*sqrt(-c^2*x^2 + a)*a*sqrt(-(c^2*d*x^2 - a*d)/a)/((c^3*d*x^2 - a*c*d)*a
rctan(sqrt(-c^2*x^2 + a)*c*x/(c^2*x^2 - a))^2)`

Sympy [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = \int \frac{1}{\sqrt{-d\left(-1 + \frac{c^2 x^2}{a}\right)} \operatorname{atan}^3\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)} dx$$

input `integrate(1/(d-c**2*d*x**2/a)**(1/2)/atan(c*x/(-c**2*x**2+a)**(1/2))**3,x)`

output `Integral(1/(sqrt(-d*(-1 + c**2*x**2/a))*atan(c*x/sqrt(a - c**2*x**2))**3),
x)`

Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.49

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = -\frac{\sqrt{a}}{2c\sqrt{d} \arctan\left(cx, \sqrt{-c^2 x^2 + a}\right)^2}$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^3,x, algo
rithm="maxima")`

output `-1/2*sqrt(a)/(c*sqrt(d)*arctan2(c*x, sqrt(-c^2*x^2 + a))^2)`

Giac [F]

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = \int \frac{1}{\sqrt{-\frac{c^2 dx^2}{a} + d} \arctan\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right)^3} dx$$

input `integrate(1/(d-c^2*d*x^2/a)^(1/2)/arctan(c*x/(-c^2*x^2+a)^(1/2))^3,x, algo
rithm="giac")`

output `integrate(1/(sqrt(-c^2*d*x^2/a + d)*arctan(c*x/sqrt(-c^2*x^2 + a))^3), x)`

Mupad [B] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = -\frac{\sqrt{a - c^2 x^2}}{2c \operatorname{atan}\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^2 \sqrt{d - \frac{c^2 dx^2}{a}}}$$

input `int(1/(atan((c*x)/(a - c^2*x^2)^(1/2))^3*(d - (c^2*d*x^2)/a)^(1/2)),x)`

output `-(a - c^2*x^2)^(1/2)/(2*c*atan((c*x)/(a - c^2*x^2)^(1/2))^2*(d - (c^2*d*x^2)/a)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.53

$$\int \frac{1}{\sqrt{d - \frac{c^2 dx^2}{a}} \arctan\left(\frac{cx}{\sqrt{a - c^2 x^2}}\right)^3} dx = -\frac{\sqrt{d} \sqrt{a}}{2 \operatorname{atan}\left(\frac{cx}{\sqrt{-c^2 x^2 + a}}\right)^2 cd}$$

input `int(1/(d-c^2*d*x^2/a)^(1/2)/atan(c*x/(-c^2*x^2+a)^(1/2))^3,x)`

output `(- sqrt(d)*sqrt(a))/(2*atan((c*x)/sqrt(a - c**2*x**2))*2*c*d)`

3.140
$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx$$

Optimal result	1034
Mathematica [A] (verified)	1034
Rubi [A] (verified)	1035
Maple [F]	1036
Fricas [A] (verification not implemented)	1036
Sympy [F]	1037
Maxima [F(-2)]	1037
Giac [F]	1038
Mupad [F(-1)]	1038
Reduce [B] (verification not implemented)	1038

Optimal result

Integrand size = 40, antiderivative size = 72

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx = \frac{\sqrt{-\frac{ae^2}{b}-e^2x^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^{1+m}}{e(1+m)\sqrt{a+bx^2}}$$

output

```
(-a*e^2/b-e^2*x^2)^(1/2)*arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^(1+m)/e/(1+m)
)/(b*x^2+a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.92

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx = \frac{\sqrt{-\frac{e^2(a+bx^2)}{b}} \arctan\left(\frac{ex}{\sqrt{-\frac{e^2(a+bx^2)}{b}}}\right)^{1+m}}{e(1+m)\sqrt{a+bx^2}}$$

input `Integrate[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^m/Sqrt[a + b*x^2],x]`

output `(Sqrt[-((e^2*(a + b*x^2))/b)]*ArcTan[(e*x)/Sqrt[-((e^2*(a + b*x^2))/b)]]^(1 + m))/(e*(1 + m)*Sqrt[a + b*x^2])`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^m}{\sqrt{a + bx^2}} dx$$

↓ 5680

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \int \frac{\arctan\left(\frac{ex}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}}\right)^m}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}} dx}{\sqrt{a + bx^2}}$$

↓ 5678

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^{m+1}}{e(m+1)\sqrt{a + bx^2}}$$

input `Int[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^m/Sqrt[a + b*x^2],x]`

output `(Sqrt[-((a*e^2)/b) - e^2*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^(1 + m))/(e*(1 + m)*Sqrt[a + b*x^2])`

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] :> Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] :> Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m}
, x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{bx^2+a}} dx$$

input

```
int(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^m/(b*x^2+a)^(1/2),x)
```

output

```
int(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^m/(b*x^2+a)^(1/2),x)
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.83

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx$$

$$= -\frac{\sqrt{bx^2+a}\left(-\arctan\left(\frac{bx\sqrt{-\frac{be^2x^2+ae^2}{b}}}{be^2x^2+ae}\right)\right)^m \sqrt{-\frac{be^2x^2+ae^2}{b}} \arctan\left(\frac{bx\sqrt{-\frac{be^2x^2+ae^2}{b}}}{be^2x^2+ae}\right)}{aem + (bem + be)x^2 + ae}$$

input

```
integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^m/(b*x^2+a)^(1/2),x, algori
thm="fricas")
```

output

```
-sqrt(b*x^2 + a)*(-arctan(b*x*sqrt(-(b*e^2*x^2 + a*e^2)/b)/(b*e*x^2 + a*e)
))^m*sqrt(-(b*e^2*x^2 + a*e^2)/b)*arctan(b*x*sqrt(-(b*e^2*x^2 + a*e^2)/b)/
(b*e*x^2 + a*e))/(a*e*m + (b*e*m + b*e)*x^2 + a*e)
```

Sympy [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx = \int \frac{\operatorname{atan}^m\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx$$

input

```
integrate(atan(e*x/(-a*e**2/b-e**2*x**2)**(1/2))**m/(b*x**2+a)**(1/2),x)
```

output

```
Integral(atan(e*x/sqrt(-a*e**2/b - e**2*x**2))**m/sqrt(a + b*x**2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx = \text{Exception raised: RuntimeError}$$

input

```
integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^m/(b*x^2+a)^(1/2),x, algori
thm="maxima")
```

output

```
Exception raised: RuntimeError >> ECL says: sign: argument cannot be imagi
nary; found sqrt((-SAGE_VAR_b*SAGE_VAR_x^2)-SAGE_VAR_a)
```

Giac [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx = \int \frac{\arctan\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)^m}{\sqrt{bx^2+a}} dx$$

input `integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^m/(b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate(arctan(e*x/sqrt(-e^2*x^2 - a*e^2/b))^m/sqrt(b*x^2 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx = \int \frac{\operatorname{atan}\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)^m}{\sqrt{bx^2+a}} dx$$

input `int(atan((e*x)/(-e^2*x^2 - (a*e^2)/b)^(1/2))^m/(a + b*x^2)^(1/2),x)`

output `int(atan((e*x)/(-e^2*x^2 - (a*e^2)/b)^(1/2))^m/(a + b*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.44

$$\begin{aligned} & \int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^m}{\sqrt{a+bx^2}} dx \\ &= -\frac{\sqrt{b}(-1)^m \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+a}ix+bi x^2}{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2}\right)^m \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+a}ix+bi x^2}{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2}\right)^i}{b(m+1)} \end{aligned}$$

input `int(atan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^m/(b*x^2+a)^(1/2),x)`

output `(- sqrt(b)*(- 1)**m*atan((sqrt(b)*sqrt(a + b*x**2)*i*x + b*i*x**2)/(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2))**m*atan((sqrt(b)*sqrt(a + b*x**2)*i*x + b*i*x**2)/(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2))*i)/(b*(m + 1))`

3.141
$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx$$

Optimal result	1040
Mathematica [A] (verified)	1040
Rubi [A] (verified)	1041
Maple [F]	1042
Fricas [F]	1042
Sympy [F]	1043
Maxima [F(-2)]	1043
Giac [F]	1043
Mupad [F(-1)]	1044
Reduce [B] (verification not implemented)	1044

Optimal result

Integrand size = 40, antiderivative size = 68

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \frac{\sqrt{-\frac{ae^2}{b}-e^2x^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^3}{3e\sqrt{a+bx^2}}$$

output `1/3*(-a*e^2/b-e^2*x^2)^(1/2)*arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^3/e/(b*x^2+a)^(1/2)`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.91

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \frac{\sqrt{-\frac{e^2(a+bx^2)}{b}} \arctan\left(\frac{ex}{\sqrt{-\frac{e^2(a+bx^2)}{b}}}\right)^3}{3e\sqrt{a+bx^2}}$$

input `Integrate[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^2/Sqrt[a + b*x^2],x]`

output `(Sqrt[-((e^2*(a + b*x^2))/b)]*ArcTan[(e*x)/Sqrt[-((e^2*(a + b*x^2))/b)]]^3)/(3*e*Sqrt[a + b*x^2])`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^2}{\sqrt{a + bx^2}} dx$$

↓ 5680

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \int \frac{\arctan\left(\frac{ex}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}}\right)^2}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}} dx}{\sqrt{a + bx^2}}$$

↓ 5678

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^3}{3e\sqrt{a + bx^2}}$$

input `Int[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^2/Sqrt[a + b*x^2],x]`

output `(Sqrt[-((a*e^2)/b) - e^2*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^3)/(3*e*Sqrt[a + b*x^2])`

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{bx^2+a}} dx$$

input

```
int(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/(b*x^2+a)^(1/2), x)
```

output

```
int(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/(b*x^2+a)^(1/2), x)
```

Fricas [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \int \frac{\arctan\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)^2}{\sqrt{bx^2+a}} dx$$

input

```
integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/(b*x^2+a)^(1/2), x, algori
thm="fricas")
```

output

```
integral(arctan(b*x*sqrt(-(b*e^2*x^2 + a*e^2)/b)/(b*e*x^2 + a*e))^2/sqrt(b
*x^2 + a), x)
```

Sympy [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \int \frac{\operatorname{atan}^2\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx$$

input `integrate(atan(e*x/(-a*e**2/b-e**2*x**2)**(1/2))**2/(b*x**2+a)**(1/2),x)`

output `Integral(atan(e*x/sqrt(-a*e**2/b - e**2*x**2))**2/sqrt(a + b*x**2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/(b*x^2+a)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: sign: argument cannot be imaginary; found sqrt((-_SAGE_VAR_b*_SAGE_VAR_x^2)-_SAGE_VAR_a)`

Giac [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \int \frac{\arctan\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)^2}{\sqrt{bx^2+a}} dx$$

input `integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/(b*x^2+a)^(1/2),x, algorithm="giac")`

output `integrate(arctan(e*x/sqrt(-e^2*x^2 - a*e^2/b))^2/sqrt(b*x^2 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = \int \frac{\operatorname{atan}\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)^2}{\sqrt{bx^2+a}} dx$$

input `int(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))^2/(a + b*x^2)^(1/2), x)`

output `int(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))^2/(a + b*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.78

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{\sqrt{a+bx^2}} dx = -\frac{\sqrt{b} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+a}ix+bi x^2}{\sqrt{b}\sqrt{bx^2+ax+a+bx^2}}\right)^3}{3b} i$$

input `int(atan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/(b*x^2+a)^(1/2), x)`

output `(- sqrt(b)*atan((sqrt(b)*sqrt(a + b*x**2)*i*x + b*i*x**2)/(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2))**3*i)/(3*b)`

3.142
$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx$$

Optimal result	1045
Mathematica [A] (verified)	1045
Rubi [A] (verified)	1046
Maple [F]	1047
Fricas [F]	1047
Sympy [F]	1048
Maxima [F(-2)]	1048
Giac [F]	1048
Mupad [F(-1)]	1049
Reduce [B] (verification not implemented)	1049

Optimal result

Integrand size = 38, antiderivative size = 68

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \frac{\sqrt{-\frac{ae^2}{b}-e^2x^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}{2e\sqrt{a+bx^2}}$$

output

$1/2*(-a*e^2/b-e^2*x^2)^(1/2)*\arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2/e/(b*x^2+a)^(1/2)$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.91

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \frac{\sqrt{-\frac{e^2(a+bx^2)}{b}} \arctan\left(\frac{ex}{\sqrt{-\frac{e^2(a+bx^2)}{b}}}\right)^2}{2e\sqrt{a+bx^2}}$$

input

`Integrate[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]/Sqrt[a + b*x^2],x]`

output

```
(Sqrt[-((e^2*(a + b*x^2))/b)]*ArcTan[(e*x)/Sqrt[-((e^2*(a + b*x^2))/b)]]^2)/(2*e*Sqrt[a + b*x^2])
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)}{\sqrt{a + bx^2}} dx$$

↓ 5680

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \int \frac{\arctan\left(\frac{ex}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}}\right)}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}} dx}{\sqrt{a + bx^2}}$$

↓ 5678

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^2}{2e\sqrt{a + bx^2}}$$

input

```
Int[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]/Sqrt[a + b*x^2], x]
```

output

```
(Sqrt[-((a*e^2)/b) - e^2*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^2)/(2*e*Sqrt[a + b*x^2])
```

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{bx^2+a}} dx$$

input

```
int(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))/(b*x^2+a)^(1/2),x)
```

output

```
int(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))/(b*x^2+a)^(1/2),x)
```

Fricas [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \int \frac{\arctan\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)}{\sqrt{bx^2+a}} dx$$

input

```
integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))/(b*x^2+a)^(1/2),x, algorithm
m="fricas")
```

output

```
integral(-arctan(b*x*sqrt(-(b*e^2*x^2 + a*e^2)/b)/(b*e*x^2 + a*e))/sqrt(b*
x^2 + a), x)
```


Sympy [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \int \frac{\operatorname{atan}\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx$$

input `integrate(atan(e*x/(-a*e**2/b-e**2*x**2)**(1/2))/(b*x**2+a)**(1/2),x)`

output `Integral(atan(e*x/sqrt(-a*e**2/b - e**2*x**2))/sqrt(a + b*x**2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))/(b*x^2+a)^(1/2),x, algorithm m="maxima")`

output `Exception raised: RuntimeError >> ECL says: sign: argument cannot be imaginary; found sqrt((-_SAGE_VAR_b*_SAGE_VAR_x^2)-_SAGE_VAR_a)`

Giac [F]

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \int \frac{\arctan\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)}{\sqrt{bx^2+a}} dx$$

input `integrate(arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))/(b*x^2+a)^(1/2),x, algorithm m="giac")`

output `integrate(arctan(e*x/sqrt(-e^2*x^2 - a*e^2/b))/sqrt(b*x^2 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \int \frac{\operatorname{atan}\left(\frac{ex}{\sqrt{-e^2x^2-\frac{ae^2}{b}}}\right)}{\sqrt{bx^2+a}} dx$$

input `int(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))/(a + b*x^2)^(1/2), x)`

output `int(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))/(a + b*x^2)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.78

$$\int \frac{\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}{\sqrt{a+bx^2}} dx = \frac{\sqrt{b} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+a}ix+bi x^2}{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2}\right)^2 i}{2b}$$

input `int(atan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))/(b*x^2+a)^(1/2), x)`

output `(sqrt(b)*atan((sqrt(b)*sqrt(a + b*x**2)*i*x + b*i*x**2)/(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2))**2*i)/(2*b)`

3.143
$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)} dx$$

Optimal result	1050
Mathematica [A] (verified)	1050
Rubi [A] (verified)	1051
Maple [F]	1052
Fricas [A] (verification not implemented)	1052
Sympy [F]	1053
Maxima [F(-2)]	1053
Giac [F]	1054
Mupad [F(-1)]	1054
Reduce [B] (verification not implemented)	1054

Optimal result

Integrand size = 40, antiderivative size = 64

$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)} dx = \frac{\sqrt{-\frac{ae^2}{b}-e^2x^2} \log\left(\arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)\right)}{e\sqrt{a+bx^2}}$$

output

$(-a*e^2/b-e^2*x^2)^{(1/2)}*\ln(\arctan(e*x/(-a*e^2/b-e^2*x^2)^{(1/2)}))/e/(b*x^2+a)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)} dx = \frac{\sqrt{-\frac{e^2(a+bx^2)}{b}} \log\left(\arctan\left(\frac{ex}{\sqrt{-\frac{e^2(a+bx^2)}{b}}}\right)\right)}{e\sqrt{a+bx^2}}$$

input `Integrate[1/(Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]),x]`

output `(Sqrt[-((e^2*(a + b*x^2))/b)]*Log[ArcTan[(e*x)/Sqrt[-((e^2*(a + b*x^2))/b)]]]/(e*Sqrt[a + b*x^2]))`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {5680, 5676}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)} dx$$

↓ 5680

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \int \frac{1}{\sqrt{-x^2e^2 - \frac{ae^2}{b}} \arctan\left(\frac{ex}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}}\right)} dx}{\sqrt{a + bx^2}}$$

↓ 5676

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \log\left(\arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)\right)}{e\sqrt{a + bx^2}}$$

input `Int[1/(Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]),x]`

output `(Sqrt[-((a*e^2)/b) - e^2*x^2]*Log[ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]]/(e*Sqrt[a + b*x^2]))`

Definitions of rubi rules used

rule 5676

```
Int[1/(ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]*Sqrt[(a_.) + (b_.)*(x_)^2]), x_Symbol] := Simp[(1/c)*Log[ArcTan[c*(x/Sqrt[a + b*x^2])]], x] /; FreeQ[{a, b, c}, x] && EqQ[b + c^2, 0]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [F]

$$\int \frac{1}{\sqrt{bx^2 + a} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx$$

input

```
int(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2)),x)
```

output

```
int(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2)),x)
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.30

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx$$

$$= \frac{\sqrt{bx^2 + a} \sqrt{-\frac{be^2x^2 + ae^2}{b}} \log\left(2 \arctan\left(\frac{bx\sqrt{-\frac{be^2x^2 + ae^2}{b}}}{be^2x^2 + ae}\right)\right)}{be^2x^2 + ae}$$

input

```
integrate(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2)),x, algorithm="fricas")
```

output `sqrt(b*x^2 + a)*sqrt(-(b*e^2*x^2 + a*e^2)/b)*log(2*arctan(b*x*sqrt(-(b*e^2*x^2 + a*e^2)/b)/(b*e*x^2 + a*e)))/(b*e*x^2 + a*e)`

Sympy [F]

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx = \int \frac{1}{\sqrt{a + bx^2} \operatorname{atan}\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx$$

input `integrate(1/(b*x**2+a)**(1/2)/atan(e*x/(-a*e**2/b-e**2*x**2)**(1/2)),x)`

output `Integral(1/(sqrt(a + b*x**2)*atan(e*x/sqrt(-a*e**2/b - e**2*x**2))), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: sign: argument cannot be imaginary; found sqrt((-_SAGE_VAR_b*_SAGE_VAR_x^2)-_SAGE_VAR_a)`

Giac [F]

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx = \int \frac{1}{\sqrt{bx^2 + a} \arctan\left(\frac{ex}{\sqrt{-e^2x^2 - \frac{ae^2}{b}}}\right)} dx$$

input `integrate(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2)),x, algorithm="giac")`

output `integrate(1/(sqrt(b*x^2 + a)*arctan(e*x/sqrt(-e^2*x^2 - a*e^2/b))), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx = \int \frac{1}{\operatorname{atan}\left(\frac{ex}{\sqrt{-e^2x^2 - \frac{ae^2}{b}}}\right) \sqrt{bx^2 + a}} dx$$

input `int(1/(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))*(a + b*x^2)^(1/2)),x)`

output `int(1/(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))*(a + b*x^2)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx = \frac{\sqrt{b} \log\left(-\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+ax+bx^2}}{\sqrt{b}\sqrt{bx^2+ax+bx^2}}\right)\right) i}{b}$$

input `int(1/(b*x^2+a)^(1/2)/atan(e*x/(-a*e^2/b-e^2*x^2)^(1/2)),x)`

output $(\sqrt{b} \cdot \log(-\operatorname{atan}(\sqrt{b} \sqrt{a + b x^2}) i x + b i x^2) / (\sqrt{b} \sqrt{a + b x^2}) x + a + b x^2) i) / b$

3.144
$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2} dx$$

Optimal result	1056
Mathematica [A] (verified)	1056
Rubi [A] (verified)	1057
Maple [F]	1058
Fricas [A] (verification not implemented)	1058
Sympy [F]	1059
Maxima [F(-2)]	1059
Giac [F]	1059
Mupad [F(-1)]	1060
Reduce [B] (verification not implemented)	1060

Optimal result

Integrand size = 40, antiderivative size = 66

$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2} dx = -\frac{\sqrt{-\frac{ae^2}{b}-e^2x^2}}{e\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)}$$

output

$$-(-a*e^2/b-e^2*x^2)^(1/2)/e/(b*x^2+a)^(1/2)/\arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2} dx = \frac{e\sqrt{a+bx^2}}{b\sqrt{-\frac{e^2(a+bx^2)}{b}} \arctan\left(\frac{ex}{\sqrt{-\frac{e^2(a+bx^2)}{b}}}\right)}$$

input `Integrate[1/(Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^2),x]`

output `(e*Sqrt[a + b*x^2])/(b*Sqrt[-((e^2*(a + b*x^2))/b)]*ArcTan[(e*x)/Sqrt[-((e^2*(a + b*x^2))/b)]])`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^2} dx$$

↓ 5680

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \int \frac{1}{\sqrt{-x^2e^2 - \frac{ae^2}{b}} \arctan\left(\frac{ex}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}}\right)^2} dx}{\sqrt{a + bx^2}}$$

↓ 5678

$$-\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}{e\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)}$$

input `Int[1/(Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^2),x]`

output `-(Sqrt[-((a*e^2)/b) - e^2*x^2]/(e*Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]))`

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [F]

$$\int \frac{1}{\sqrt{bx^2 + a} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx$$

input

```
int(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2,x)
```

output

```
int(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2,x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.24

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx = \frac{\sqrt{bx^2 + a} \sqrt{-\frac{be^2x^2 + ae^2}{b}}}{(be^2x^2 + ae) \arctan\left(\frac{bx \sqrt{-\frac{be^2x^2 + ae^2}{b}}}{be^2x^2 + ae}\right)}$$

input

```
integrate(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2,x, algo
rithm="fricas")
```

output

```
sqrt(b*x^2 + a)*sqrt(-(b*e^2*x^2 + a*e^2)/b)/((b*e*x^2 + a*e)*arctan(b*x*s
qrt(-(b*e^2*x^2 + a*e^2)/b)/(b*e*x^2 + a*e))
```

Sympy [F]

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx = \int \frac{1}{\sqrt{a + bx^2} \operatorname{atan}^2\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx$$

input `integrate(1/(b*x**2+a)**(1/2)/atan(ex/(-a*e**2/b-e**2*x**2)**(1/2))**2,x)`

output `Integral(1/(sqrt(a + b*x**2)*atan(ex/sqrt(-a*e**2/b - e**2*x**2))**2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(1/(b*x^2+a)^(1/2)/arctan(ex/(-a*e^2/b-e^2*x^2)^(1/2))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: sign: argument cannot be imaginary; found sqrt((-SAGE_VAR_b*SAGE_VAR_x^2)-SAGE_VAR_a)`

Giac [F]

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx = \int \frac{1}{\sqrt{bx^2 + a} \arctan\left(\frac{ex}{\sqrt{-e^2x^2 - \frac{ae^2}{b}}}\right)^2} dx$$

input `integrate(1/(b*x^2+a)^(1/2)/arctan(ex/(-a*e^2/b-e^2*x^2)^(1/2))^2,x, algorithm="giac")`

output `integrate(1/(sqrt(b*x^2 + a)*arctan(e*x/sqrt(-e^2*x^2 - a*e^2/b))^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx = \int \frac{1}{\operatorname{atan}\left(\frac{ex}{\sqrt{-e^2x^2 - \frac{ae^2}{b}}}\right)^2 \sqrt{bx^2 + a}} dx$$

input `int(1/(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))^2*(a + b*x^2)^(1/2)),x)`

output `int(1/(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))^2*(a + b*x^2)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^2} dx = \frac{\sqrt{b}i}{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+a}ix+bi x^2}{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2}\right) b}$$

input `int(1/(b*x^2+a)^(1/2)/atan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2,x)`

output `(sqrt(b)*i)/(atan((sqrt(b)*sqrt(a + b*x**2)*i*x + b*i*x**2)/(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2))*b)`

3.145
$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^3} dx$$

Optimal result	1061
Mathematica [A] (verified)	1061
Rubi [A] (verified)	1062
Maple [F]	1063
Fricas [A] (verification not implemented)	1063
Sympy [F]	1064
Maxima [F(-2)]	1064
Giac [F]	1064
Mupad [F(-1)]	1065
Reduce [B] (verification not implemented)	1065

Optimal result

Integrand size = 40, antiderivative size = 68

$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^3} dx = -\frac{\sqrt{-\frac{ae^2}{b}-e^2x^2}}{2e\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^2}$$

output

```
-1/2*(-a*e^2/b-e^2*x^2)^(1/2)/e/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^2
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b}-e^2x^2}}\right)^3} dx = -\frac{\sqrt{-\frac{e^2(a+bx^2)}{b}}}{2e\sqrt{a+bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{e^2(a+bx^2)}{b}}}\right)^2}$$

input `Integrate[1/(Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^3),x]`

output `-1/2*Sqrt[-((e^2*(a + b*x^2))/b)]/(e*Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((e^2*(a + b*x^2))/b)]^2)`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {5680, 5678}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^3} dx$$

↓ 5680

$$\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}} \int \frac{1}{\sqrt{-x^2e^2 - \frac{ae^2}{b}} \arctan\left(\frac{ex}{\sqrt{-x^2e^2 - \frac{ae^2}{b}}}\right)^3} dx}{\sqrt{a + bx^2}}$$

↓ 5678

$$-\frac{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}{2e\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{e^2(-x^2) - \frac{ae^2}{b}}}\right)^2}$$

input `Int[1/(Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^3),x]`

output `-1/2*Sqrt[-((a*e^2)/b) - e^2*x^2]/(e*Sqrt[a + b*x^2]*ArcTan[(e*x)/Sqrt[-((a*e^2)/b) - e^2*x^2]]^2)`

Definitions of rubi rules used

rule 5678

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(a_.) + (b_.
)*(x_)^2], x_Symbol] := Simp[ArcTan[c*(x/Sqrt[a + b*x^2])]^(m + 1)/(c*(m +
1)), x] /; FreeQ[{a, b, c, m}, x] && EqQ[b + c^2, 0] && NeQ[m, -1]
```

rule 5680

```
Int[ArcTan[((c_.)*(x_))/Sqrt[(a_.) + (b_.)*(x_)^2]]^(m_.)/Sqrt[(d_.) + (e_.
)*(x_)^2], x_Symbol] := Simp[Sqrt[a + b*x^2]/Sqrt[d + e*x^2] Int[ArcTan[c
*(x/Sqrt[a + b*x^2])]^m/Sqrt[a + b*x^2], x], x] /; FreeQ[{a, b, c, d, e, m},
x] && EqQ[b + c^2, 0] && EqQ[b*d - a*e, 0]
```

Maple [F]

$$\int \frac{1}{\sqrt{bx^2 + a} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx$$

input

```
int(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^3,x)
```

output

```
int(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^3,x)
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.22

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx = -\frac{\sqrt{bx^2 + a} \sqrt{-\frac{be^2x^2 + ae^2}{b}}}{2 (bex^2 + ae) \arctan\left(\frac{bx \sqrt{-\frac{be^2x^2 + ae^2}{b}}}{bex^2 + ae}\right)^2}$$

input

```
integrate(1/(b*x^2+a)^(1/2)/arctan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^3,x, algo
rithm="fricas")
```

output

```
-1/2*sqrt(b*x^2 + a)*sqrt(-(b*e^2*x^2 + a*e^2)/b)/((b*e*x^2 + a*e)*arctan(
b*x*sqrt(-(b*e^2*x^2 + a*e^2)/b)/(b*e*x^2 + a*e))^2)
```


Sympy [F]

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx = \int \frac{1}{\sqrt{a + bx^2} \operatorname{atan}^3\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)} dx$$

input `integrate(1/(b*x**2+a)**(1/2)/atan(ex/(-a*e**2/b-e**2*x**2)**(1/2))**3,x)`

output `Integral(1/(sqrt(a + b*x**2)*atan(ex/sqrt(-a*e**2/b - e**2*x**2))**3), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(1/(b*x^2+a)^(1/2)/arctan(ex/(-a*e^2/b-e^2*x^2)^(1/2))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: sign: argument cannot be imaginary; found sqrt((-SAGE_VAR_b*SAGE_VAR_x^2)-SAGE_VAR_a)`

Giac [F]

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx = \int \frac{1}{\sqrt{bx^2 + a} \arctan\left(\frac{ex}{\sqrt{-e^2x^2 - \frac{ae^2}{b}}}\right)^3} dx$$

input `integrate(1/(b*x^2+a)^(1/2)/arctan(ex/(-a*e^2/b-e^2*x^2)^(1/2))^3,x, algorithm="giac")`

output `integrate(1/(sqrt(b*x^2 + a)*arctan(e*x/sqrt(-e^2*x^2 - a*e^2/b))^3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx = \int \frac{1}{\operatorname{atan}\left(\frac{ex}{\sqrt{-e^2x^2 - \frac{ae^2}{b}}}\right)^3 \sqrt{bx^2 + a}} dx$$

input `int(1/(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))^3*(a + b*x^2)^(1/2)),x)`

output `int(1/(atan((e*x)/(- e^2*x^2 - (a*e^2)/b)^(1/2))^3*(a + b*x^2)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.78

$$\int \frac{1}{\sqrt{a + bx^2} \arctan\left(\frac{ex}{\sqrt{-\frac{ae^2}{b} - e^2x^2}}\right)^3} dx = -\frac{\sqrt{b}i}{2\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{bx^2+a}ix+bi x^2}{\sqrt{b}\sqrt{bx^2+a}x+a+bx^2}\right)^2 b}$$

input `int(1/(b*x^2+a)^(1/2)/atan(e*x/(-a*e^2/b-e^2*x^2)^(1/2))^3,x)`

output `(- sqrt(b)*i)/(2*atan((sqrt(b)*sqrt(a + b*x**2)*i*x + b*i*x**2)/(sqrt(b)*sqrt(a + b*x**2)*x + a + b*x**2))**2*b)`

3.146 $\int \frac{\arctan(c(a+bx)) \log(d(a+bx))}{a+bx} dx$

Optimal result	1066
Mathematica [A] (verified)	1067
Rubi [A] (verified)	1067
Maple [C] (warning: unable to verify)	1069
Fricas [F]	1070
Sympy [F]	1071
Maxima [F(-2)]	1071
Giac [F]	1071
Mupad [F(-1)]	1072
Reduce [F]	1072

Optimal result

Integrand size = 24, antiderivative size = 101

$$\int \frac{\arctan(c(a+bx)) \log(d(a+bx))}{a+bx} dx = \frac{i \log(d(a+bx)) \operatorname{PolyLog}(2, -ic(a+bx))}{2b} - \frac{i \log(d(a+bx)) \operatorname{PolyLog}(2, ic(a+bx))}{2b} - \frac{i \operatorname{PolyLog}(3, -ic(a+bx))}{2b} + \frac{i \operatorname{PolyLog}(3, ic(a+bx))}{2b}$$

output

```
1/2*I*ln(d*(b*x+a))*polylog(2,-I*c*(b*x+a))/b-1/2*I*ln(d*(b*x+a))*polylog(
2,I*c*(b*x+a))/b-1/2*I*polylog(3,-I*c*(b*x+a))/b+1/2*I*polylog(3,I*c*(b*x+
a))/b
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.78

$$\int \frac{\arctan(c(a+bx)) \log(d(a+bx))}{a+bx} dx$$

$$= \frac{i(\log(d(a+bx)) \operatorname{PolyLog}(2, -ic(a+bx)) - \log(d(a+bx)) \operatorname{PolyLog}(2, ic(a+bx)) - \operatorname{PolyLog}(3, -ic(a+bx)) + \operatorname{PolyLog}(3, ic(a+bx)))}{2b}$$

input `Integrate[(ArcTan[c*(a + b*x)]*Log[d*(a + b*x)]/(a + b*x),x]`

output `((I/2)*(Log[d*(a + b*x)]*PolyLog[2, (-I)*c*(a + b*x)] - Log[d*(a + b*x)]*PolyLog[2, I*c*(a + b*x)] - PolyLog[3, (-I)*c*(a + b*x)] + PolyLog[3, I*c*(a + b*x)]))/b`

Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {5732, 2894, 2881, 2821, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\arctan(c(a+bx)) \log(d(a+bx))}{a+bx} dx$$

$$\downarrow \text{5732}$$

$$\frac{1}{2}i \int \frac{\log(d(a+bx)) \log(1-ic(a+bx))}{a+bx} dx - \frac{1}{2}i \int \frac{\log(d(a+bx)) \log(ic(a+bx)+1)}{a+bx} dx$$

$$\downarrow \text{2894}$$

$$\frac{1}{2}i \int \frac{\log(d(a+bx)) \log(-iac-ibxc+1)}{a+bx} dx - \frac{1}{2}i \int \frac{\log(d(a+bx)) \log(iac+ibxc+1)}{a+bx} dx$$

$$\downarrow \text{2881}$$

$$\frac{i \int \frac{\log(d(a+bx)) \log(1-ic(a+bx))}{a+bx} d(a+bx)}{2b} - \frac{i \int \frac{\log(d(a+bx)) \log(ic(a+bx)+1)}{a+bx} d(a+bx)}{2b}$$

$$\begin{aligned}
 & \downarrow 2821 \\
 & \frac{i\left(\int \frac{\text{PolyLog}(2, ic(a+bx))}{a+bx} d(a+bx) - \text{PolyLog}(2, ic(a+bx)) \log(d(a+bx))\right)}{2b} - \\
 & \frac{i\left(\int \frac{\text{PolyLog}(2, -ic(a+bx))}{a+bx} d(a+bx) - \text{PolyLog}(2, -ic(a+bx)) \log(d(a+bx))\right)}{2b} \\
 & \downarrow 7143 \\
 & \frac{i(\text{PolyLog}(3, ic(a+bx)) - \text{PolyLog}(2, ic(a+bx)) \log(d(a+bx)))}{2b} - \\
 & \frac{i(\text{PolyLog}(3, -ic(a+bx)) - \text{PolyLog}(2, -ic(a+bx)) \log(d(a+bx)))}{2b}
 \end{aligned}$$

input `Int[(ArcTan[c*(a + b*x)]*Log[d*(a + b*x)])/(a + b*x),x]`

output `((-1/2*I)*(-Log[d*(a + b*x)]*PolyLog[2, (-I)*c*(a + b*x)]) + PolyLog[3, (-I)*c*(a + b*x)])/b + ((I/2)*(-Log[d*(a + b*x)]*PolyLog[2, I*c*(a + b*x)]) + PolyLog[3, I*c*(a + b*x)])/b`

Defintions of rubi rules used

rule 2821 `Int[(Log[(d_.)*((e_) + (f_.)*(x_)^(m_.))]*((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.))/(x_), x_Symbol] := Simp[(-PolyLog[2, (-d)*f*x^m]*((a + b*Log[c*x^n])^p/m), x] + Simp[b*n*(p/m) Int[PolyLog[2, (-d)*f*x^m]*((a + b*Log[c*x^n])^(p - 1)/x), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && IGtQ[p, 0] && EqQ[d*e, 1]`

rule 2881 `Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.) + Log[(h_.)*((i_.) + (j_.)*(x_)^(m_.))]*(g_.))*((k_.) + (l_.)*(x_)^(r_.), x_Symbol] := Simp[1/e Subst[Int[(k*(x/d)^r*(a + b*Log[c*x^n])^p*(f + g*Log[h*((e*i - d*j)/e + j*(x/e))^m]), x], x, d + e*x], x] /; FreeQ[{a, b, c, d, e, f, g, h, i, j, k, l, n, p, r}, x] && EqQ[e*k - d*1, 0]`

rule 2894

```
Int[((a_.) + Log[(c_.)*(v_)^(n_.)]*(b_.))^(p_.)*(u_.), x_Symbol] := Int[u*(
a + b*Log[c*ExpandToSum[v, x]^n])^p, x] /; FreeQ[{a, b, c, n, p}, x] && Lin
earQ[v, x] && !LinearMatchQ[v, x] && !(EqQ[n, 1] && MatchQ[c*v, (e_.)*(f
_) + (g_.)*x] /; FreeQ[{e, f, g}, x])]
```

rule 5732

```
Int[(ArcTan[v_]*Log[w_] / ((a_.) + (b_.)*(x_)), x_Symbol] := Simp[I/2 Int[
Log[1 - I*v]*(Log[w]/(a + b*x)), x], x] - Simp[I/2 Int[Log[1 + I*v]*(Log[
w]/(a + b*x)), x], x] /; FreeQ[{a, b}, x] && LinearQ[v, x] && LinearQ[w, x]
&& EqQ[Simplify[D[v/(a + b*x), x]], 0] && EqQ[Simplify[D[w/(a + b*x), x]],
0]
```

rule 7143

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)] / ((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p] / (e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 6.99 (sec) , antiderivative size = 1087, normalized size of antiderivative = 10.76

method	result	size
risch	Expression too large to display	1087

input

```
int(arctan(c*(b*x+a))*ln(d*(b*x+a))/(b*x+a), x, method=_RETURNVERBOSE)
```

output

```

1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(b*x+a)*Pi*csgn(I*d*(b*x+a))^3-1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(-I*c*(b*x+a))*Pi*csgn(I*d*(b*x+a))^3+1/4/b*dilog(-I*c*(b*x+a))*Pi*csgn(I*d)*csgn(I*d*(b*x+a))^2+1/4/b*dilog(-I*c*(b*x+a))*Pi*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))^2+1/2*I/b*ln(-I*(-c*(b*x+a)+I))*ln(b*x+a)*ln(d)-1/2*I/b*ln(-I*(-c*(b*x+a)+I))*ln(-I*c*(b*x+a))*ln(d)-1/2*I*polylog(3,-I*c*(b*x+a))/b+1/2*I*polylog(3,I*c*(b*x+a))/b-1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(-I*c*(b*x+a))*Pi*csgn(I*d)*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))-1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(b*x+a)*Pi*csgn(I*d)*csgn(I*d*(b*x+a))^2-1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(b*x+a)*Pi*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))^2+1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(-I*c*(b*x+a))*Pi*csgn(I*d)*csgn(I*d*(b*x+a))^2+1/4/b*ln(-I*(-c*(b*x+a)+I))*ln(-I*c*(b*x+a))*Pi*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))^2+1/2*I/b*ln(b*x+a)*polylog(2,-I*c*(b*x+a))-1/2*I/b*dilog(-I*c*(b*x+a))*ln(d)-1/4/b*dilog(-I*c*(b*x+a))*Pi*csgn(I*d*(b*x+a))^3+1/4*I/b*ln(b*x+a)^2*ln(1+I*c*(b*x+a))-1/4/b*dilog(-I*c*(b*x+a))*Pi*csgn(I*d)*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))+1/4*I/b*ln(-I*(I+c*(b*x+a)))*ln(b*x+a)^2-1/4*I/b*ln(b*x+a)^2*ln(1-I*c*(b*x+a))-1/2*I/b*ln(d)*dilog(-I*(I+c*(b*x+a)))-1/2*I/b*ln(b*x+a)*polylog(2,I*c*(b*x+a))-1/4*I*(-I*Pi*ln(b*x+a)*csgn(I*d)*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))+I*Pi*ln(b*x+a)*csgn(I*d)*csgn(I*d*(b*x+a))^2+I*Pi*ln(b*x+a)*csgn(I*(b*x+a))*csgn(I*d*(b*x+a))^2-I*Pi*ln(b*x+a)*csgn(I*d*(b*x+a))^3+2*ln(d)*ln(b*x+a)+ln(b*x+a)^2)/b*ln(1+I*c*(b*x+a))-1/4/b*Pi*csgn(I*d...

```

Fricas [F]

$$\int \frac{\arctan(c(a+bx)) \log(d(a+bx))}{a+bx} dx = \int \frac{\arctan((bx+a)c) \log((bx+a)d)}{bx+a} dx$$

input

```
integrate(arctan(c*(b*x+a))*log(d*(b*x+a))/(b*x+a),x, algorithm="fricas")
```

output

```
integral(arctan(b*c*x + a*c)*log(b*d*x + a*d)/(b*x + a), x)
```

Sympy [F]

$$\int \frac{\arctan(c(a + bx)) \log(d(a + bx))}{a + bx} dx = \int \frac{\log(ad + bdx) \operatorname{atan}(ac + bcx)}{a + bx} dx$$

input `integrate(atan(c*(b*x+a))*ln(d*(b*x+a))/(b*x+a),x)`

output `Integral(log(a*d + b*d*x)*atan(a*c + b*c*x)/(a + b*x), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\arctan(c(a + bx)) \log(d(a + bx))}{a + bx} dx = \text{Exception raised: RuntimeError}$$

input `integrate(arctan(c*(b*x+a))*log(d*(b*x+a))/(b*x+a),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [F]

$$\int \frac{\arctan(c(a + bx)) \log(d(a + bx))}{a + bx} dx = \int \frac{\arctan((bx + a)c) \log((bx + a)d)}{bx + a} dx$$

input `integrate(arctan(c*(b*x+a))*log(d*(b*x+a))/(b*x+a),x, algorithm="giac")`

output `integrate(arctan((b*x + a)*c)*log((b*x + a)*d)/(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(c(a + bx)) \log(d(a + bx))}{a + bx} dx = \int \frac{\operatorname{atan}(c(a + bx)) \ln(d(a + bx))}{a + bx} dx$$

input `int((atan(c*(a + b*x))*log(d*(a + b*x)))/(a + b*x),x)`

output `int((atan(c*(a + b*x))*log(d*(a + b*x)))/(a + b*x), x)`

Reduce [F]

$$\int \frac{\arctan(c(a + bx)) \log(d(a + bx))}{a + bx} dx = \int \frac{\operatorname{atan}(bcx + ac) \log(bdx + ad)}{bx + a} dx$$

input `int(atan(c*(b*x+a))*log(d*(b*x+a))/(b*x+a),x)`

output `int((atan(a*c + b*c*x)*log(a*d + b*d*x))/(a + b*x),x)`

3.147 $\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx$

Optimal result	1073
Mathematica [A] (verified)	1073
Rubi [A] (verified)	1074
Maple [C] (warning: unable to verify)	1075
Fricas [A] (verification not implemented)	1076
Sympy [F(-1)]	1077
Maxima [A] (verification not implemented)	1077
Giac [A] (verification not implemented)	1077
Mupad [B] (verification not implemented)	1078
Reduce [B] (verification not implemented)	1078

Optimal result

Integrand size = 20, antiderivative size = 48

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx$$

$$= \frac{e^{ac+bcx} \arctan(\sinh(c(a + bx)))}{bc} - \frac{\log(1 + e^{2c(a+bx)})}{bc}$$

output `exp(b*c*x+a*c)*arctan(sinh(c*(b*x+a)))/b/c-ln(1+exp(2*c*(b*x+a)))/b/c`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.27

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx$$

$$= -\frac{e^{c(a+bx)} \arctan\left(\frac{1}{2}e^{-c(a+bx)} - \frac{1}{2}e^{c(a+bx)}\right) + \log(1 + e^{2c(a+bx)})}{bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTan[Sinh[a*c + b*c*x]],x]`

output `-((E^(c*(a + b*x))*ArcTan[1/(2*E^(c*(a + b*x))) - E^(c*(a + b*x))]/2] + Log[1 + E^(2*c*(a + b*x))])/(b*c)`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {7281, 5730, 2720, 27, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx \\
 & \quad \downarrow \text{7281} \\
 & \frac{\int e^{ac+bcx} \arctan(\sinh(ac + bcx)) d(ac + bcx)}{bc} \\
 & \quad \downarrow \text{5730} \\
 & \frac{e^{ac+bcx} \arctan(\sinh(ac + bcx)) - \int e^{ac+bcx} \operatorname{sech}(ac + bcx) d(ac + bcx)}{bc} \\
 & \quad \downarrow \text{2720} \\
 & \frac{e^{ac+bcx} \arctan(\sinh(ac + bcx)) - \int \frac{2e^{ac+bcx}}{1+e^{2ac+2bcx}} de^{ac+bcx}}{bc} \\
 & \quad \downarrow \text{27} \\
 & \frac{e^{ac+bcx} \arctan(\sinh(ac + bcx)) - 2 \int \frac{e^{ac+bcx}}{1+e^{2ac+2bcx}} de^{ac+bcx}}{bc} \\
 & \quad \downarrow \text{240} \\
 & \frac{e^{ac+bcx} \arctan(\sinh(ac + bcx)) - \log(e^{2ac+2bcx} + 1)}{bc}
 \end{aligned}$$

input `Int[E^(c*(a + b*x))*ArcTan[Sinh[a*c + b*c*x]],x]`

output `(E^(a*c + b*c*x)*ArcTan[Sinh[a*c + b*c*x]] - Log[1 + E^(2*a*c + 2*b*c*x)]) / (b*c)`

Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 5730 `Int[((a_.) + ArcTan[u_]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]]`

rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.89 (sec) , antiderivative size = 1299, normalized size of antiderivative = 27.06

method	result	size
risch	Expression too large to display	1299

input `int(exp(c*(b*x+a))*arctan(sinh(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

```

2*a/b+1/2/b/c*exp(c*(b*x+a))*Pi-ln(1+exp(2*c*(b*x+a)))/b/c-1/4/b/c*Pi*csgn
(exp(-c*(b*x+a))*(exp(c*(b*x+a))+I)^2)^2*exp(c*(b*x+a))-I/b/c*exp(c*(b*x+a)
))*ln(exp(c*(b*x+a))-I)+1/4/b/c*Pi*csgn(exp(-c*(b*x+a))*(exp(c*(b*x+a))-I)
^2)^3*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(exp(-c*(b*x+a))*(exp(c*(b*x+a))+I)^2)
^3*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a))*(exp(c*(b*x+a))-I)^2)^
3*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*(exp(c*(b*x+a))-I)^2)^3*exp(c*(b*x+a))+
1/4/b/c*Pi*csgn(I*(exp(c*(b*x+a))+I)^2)^3*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(I
*exp(-c*(b*x+a))*(exp(c*(b*x+a))+I)^2)^3*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(ex
p(-c*(b*x+a))*(exp(c*(b*x+a))-I)^2)^2*exp(c*(b*x+a))+I/b/c*exp(c*(b*x+a))*
ln(exp(c*(b*x+a))+I)-1/4/b/c*Pi*csgn(I*(exp(c*(b*x+a))-I)^2)*csgn(I*exp(-c
*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(exp(c*(b*x+a))-I)^2)*exp(c*(b*x+a))+1/4
/b/c*Pi*csgn(I*(exp(c*(b*x+a))+I)^2)*csgn(I*exp(-c*(b*x+a)))*csgn(I*exp(-c
*(b*x+a))*(exp(c*(b*x+a))+I)^2)*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(I*(exp(c*(b
*x+a))+I))^2*csgn(I*(exp(c*(b*x+a))+I)^2)*exp(c*(b*x+a))-1/2/b/c*Pi*csgn(I
*(exp(c*(b*x+a))+I))*csgn(I*(exp(c*(b*x+a))+I)^2)^2*exp(c*(b*x+a))-1/4/b/c
*Pi*csgn(I*(exp(c*(b*x+a))+I)^2)*csgn(I*exp(-c*(b*x+a))*(exp(c*(b*x+a))+I)
^2)^2*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a))*(exp(c*(b*x+a))-I)^
2)*csgn(exp(-c*(b*x+a))*(exp(c*(b*x+a))-I)^2)*exp(c*(b*x+a))+1/4/b/c*Pi*cs
gn(I*exp(-c*(b*x+a))*(exp(c*(b*x+a))+I)^2)*csgn(exp(-c*(b*x+a))*(exp(c*(b
*x+a))+I)^2)*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a))*(exp(c*(b*...

```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.56

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx$$

$$= \frac{(\cosh(bcx + ac) + \sinh(bcx + ac)) \arctan(\sinh(bcx + ac)) - \log\left(\frac{2 \cosh(bcx+ac)}{\cosh(bcx+ac) - \sinh(bcx+ac)}\right)}{bc}$$

input

```
integrate(exp(c*(b*x+a))*arctan(sinh(b*c*x+a*c)),x, algorithm="fricas")
```

output

```
((cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*arctan(sinh(b*c*x + a*c)) - log(2
*cosh(b*c*x + a*c)/(cosh(b*c*x + a*c) - sinh(b*c*x + a*c)))/(b*c)
```

Sympy [F(-1)]

Timed out.

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx = \text{Timed out}$$

input `integrate(exp(c*(b*x+a))*atan(sinh(b*c*x+a*c)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx = \frac{\arctan(\sinh(bc x + ac)) e^{((bx+a)c)}}{bc} - \frac{\log(e^{(2bcx+2ac)} + 1)}{bc}$$

input `integrate(exp(c*(b*x+a))*arctan(sinh(b*c*x+a*c)),x, algorithm="maxima")`

output `arctan(sinh(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) - log(e^(2*b*c*x + 2*a*c) + 1)/(b*c)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.38

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx = \frac{(\arctan(\frac{1}{2}(e^{(2bcx+2ac)} - 1))e^{(-bcx-ac)}) e^{(bcx)} - e^{(-ac)} \log(e^{(2bcx+2ac)} + 1)) e^{(ac)}}{bc}$$

input `integrate(exp(c*(b*x+a))*arctan(sinh(b*c*x+a*c)),x, algorithm="giac")`

output

$$(\arctan(1/2*(e^{(2*b*c*x + 2*a*c)} - 1))*e^{(-b*c*x - a*c)}*e^{(b*c*x)} - e^{(-a*c)}*\log(e^{(2*b*c*x + 2*a*c)} + 1))*e^{(a*c)}/(b*c)$$
Mupad [B] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.38

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx = \frac{e^{bcx} e^{ac} \operatorname{atan}\left(\frac{e^{bcx} e^{ac}}{2} - \frac{e^{-bcx} e^{-ac}}{2}\right)}{bc} - \frac{\ln(e^{2bcx} e^{2ac} + 1)}{bc}$$

input

$$\text{int}(\exp(c*(a + b*x))*\operatorname{atan}(\sinh(a*c + b*c*x)),x)$$

output

$$(\exp(b*c*x)*\exp(a*c)*\operatorname{atan}((\exp(b*c*x)*\exp(a*c))/2 - (\exp(-b*c*x)*\exp(-a*c))/2))/(b*c) - \log(\exp(2*b*c*x)*\exp(2*a*c) + 1)/(b*c)$$
Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.35

$$\int e^{c(a+bx)} \arctan(\sinh(ac + bcx)) dx = \frac{e^{bcx+ac} \operatorname{atan}\left(\frac{e^{2bcx+2ac}-1}{2e^{bcx+ac}}\right) - \log(e^{2bcx+2ac} + 1)}{bc}$$

input

$$\text{int}(\exp(c*(b*x+a))*\operatorname{atan}(\sinh(b*c*x+a*c)),x)$$

output

$$(e^{**}(a*c + b*c*x)*\operatorname{atan}((e^{**}(2*a*c + 2*b*c*x)} - 1)/(2*e^{**}(a*c + b*c*x)))) - \log(e^{**}(2*a*c + 2*b*c*x)} + 1))/(b*c)$$

3.148 $\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx$

Optimal result	1079
Mathematica [C] (verified)	1079
Rubi [A] (verified)	1080
Maple [C] (warning: unable to verify)	1082
Fricas [B] (verification not implemented)	1083
Sympy [F]	1084
Maxima [A] (verification not implemented)	1084
Giac [A] (verification not implemented)	1085
Mupad [B] (verification not implemented)	1085
Reduce [B] (verification not implemented)	1086

Optimal result

Integrand size = 20, antiderivative size = 103

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx = \frac{e^{ac+bcx} \arctan(\cosh(c(a + bx)))}{bc} - \frac{(1 - \sqrt{2}) \log(3 - 2\sqrt{2} + e^{2c(a+bx)})}{2bc} - \frac{(1 + \sqrt{2}) \log(3 + 2\sqrt{2} + e^{2c(a+bx)})}{2bc}$$

output

```
exp(b*c*x+a*c)*arctan(cosh(c*(b*x+a)))/b/c-1/2*(1-2^(1/2))*ln(3-2*2^(1/2)+exp(2*c*(b*x+a)))/b/c-1/2*(1+2^(1/2))*ln(3+2*2^(1/2)+exp(2*c*(b*x+a)))/b/c
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.08 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.42

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx = \frac{-4c(a + bx) + 2e^{c(a+bx)} \arctan\left(\frac{1}{2}e^{-c(a+bx)}(1 + e^{2c(a+bx)})\right) + \text{RootSum}\left[1 + 6\#1^2 + \#1^4 \&, \frac{ac+bcx-\log\left(e^{c(a+bx)}\right)}{2bc}\right]}{2bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTan[Cosh[a*c + b*c*x]],x]`

output $(-4*c*(a + b*x) + 2*E^(c*(a + b*x))*ArcTan[(1 + E^(2*c*(a + b*x)))/(2*E^(c*(a + b*x))]) + RootSum[1 + 6*#1^2 + #1^4 \& , (a*c + b*c*x - Log[E^(c*(a + b*x)) - #1] + 7*a*c*#1^2 + 7*b*c*x*#1^2 - 7*Log[E^(c*(a + b*x)) - #1]*#1^2)/(1 + 3*#1^2) \&])/(2*b*c)$

Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {7281, 5730, 2720, 27, 1576, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \arctan(\cosh(ac + bxc)) dx \\
 & \quad \downarrow \text{7281} \\
 & \frac{\int e^{ac+bx} \arctan(\cosh(ac + bxc)) d(ac + bxc)}{bc} \\
 & \quad \downarrow \text{5730} \\
 & \frac{e^{ac+bx} \arctan(\cosh(ac + bxc)) - \int \frac{e^{ac+bx} \sinh(ac+bx)}{\cosh^2(ac+bx)+1} d(ac + bxc)}{bc} \\
 & \quad \downarrow \text{2720} \\
 & \frac{e^{ac+bx} \arctan(\cosh(ac + bxc)) - \int -\frac{2e^{ac+bx}(1-e^{2ac+2bxc})}{1+6e^{2ac+2bxc}+e^{4ac+4bxc}} de^{ac+bx}}{bc} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 \int \frac{e^{ac+bx}(1-e^{2ac+2bxc})}{1+6e^{2ac+2bxc}+e^{4ac+4bxc}} de^{ac+bx} + e^{ac+bx} \arctan(\cosh(ac + bxc))}{bc} \\
 & \quad \downarrow \text{1576} \\
 & \frac{\int \frac{-ac-bxc+1}{1+7e^{2ac+2bxc}} de^{2ac+2bxc} + e^{ac+bx} \arctan(\cosh(ac + bxc))}{bc}
 \end{aligned}$$

$$\int \left(-\frac{2-\sqrt{2}}{2(4-3\sqrt{2}-\sqrt{2}e^{2ac+2bcx})} - \frac{1+\sqrt{2}}{2(3+2\sqrt{2}+e^{2ac+2bcx})} \right) de^{2ac+2bcx} + e^{ac+bcx} \arctan(\cosh(ac+bcx))$$

1141

bc

2009

$$\frac{e^{ac+bcx} \arctan(\cosh(ac+bcx)) - \frac{1}{2}(1+\sqrt{2}) \log(e^{2ac+2bcx} + 3 + 2\sqrt{2}) - \frac{1}{2}(1-\sqrt{2}) \log(-\sqrt{2}e^{2ac+2bcx} + 4 - 3\sqrt{2})}{bc}$$

input `Int[E^(c*(a + b*x))*ArcTan[Cosh[a*c + b*c*x]],x]`

output `(E^(a*c + b*c*x)*ArcTan[Cosh[a*c + b*c*x]] - ((1 + Sqrt[2])*Log[3 + 2*Sqrt[2] + E^(2*a*c + 2*b*c*x)])/2 - ((1 - Sqrt[2])*Log[4 - 3*Sqrt[2] - Sqrt[2]*E^(2*a*c + 2*b*c*x)])/2)/(b*c)`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1141 `Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1576 `Int[(x_)*((d_.) + (e_.)*(x_)^2)^(q_)*((a_.) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

rule 5730

```
Int[((a_.) + ArcTan[u]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]},
  Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]
/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x]
&& InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.) /; Fre
eQ[{c, d, m}, x]] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]
```

rule 7281

```
Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]]
  Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 4.48 (sec) , antiderivative size = 1371, normalized size of antiderivative = 13.31

method	result	size
risch	Expression too large to display	1371

input

```
int(exp(c*(b*x+a))*arctan(cosh(b*c*x+a*c)),x,method=_RETURNVERBOSE)
```

output

```

1/2*I/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a)))+1/4/b/c
*Pi*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))^3*exp
(c*(b*x+a))+1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(-ex
p(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))^2*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(I*exp(2*c*(b*x+a))+I+2*exp(c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))*csgn(exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))^2*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a)))*csgn(I*exp(2*c*(b*x+a))+I+2*exp(c*(b*x+a)))*csgn(I*exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a))))^3*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a)))*csgn(I*exp(-c*(b*x+a)))*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a)))*csgn(-I*exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-I)*csgn(I*exp(-c*(b*x+a)))*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a))))*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a)))*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a))))^3*exp(c*(b*x+a))+1/4/b/c*Pi*csgn(-I*exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-I)*csgn(I*exp(-c*(b*x+a)))*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(-c*(b*x+a)))*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a))))*csgn(exp(-c*(b*x+a))*(-exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a))))^2*exp(c*(b*x+a))...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(86) = 172$.

Time = 0.11 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.15

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx$$

$$= \frac{2(\cosh(bcx + ac) + \sinh(bcx + ac)) \arctan(\cosh(bcx + ac)) + \sqrt{2} \log\left(-\frac{3(2\sqrt{2}-3) \cosh(bcx+ac)^2 - 4(3\sqrt{2}-4)}{\cos}\right)}{}$$

input

```
integrate(exp(c*(b*x+a))*arctan(cosh(b*c*x+a*c)),x, algorithm="fricas")
```

output

```
1/2*(2*(cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*arctan(cosh(b*c*x + a*c)) +
sqrt(2)*log(-(3*(2*sqrt(2) - 3)*cosh(b*c*x + a*c)^2 - 4*(3*sqrt(2) - 4)*c
osh(b*c*x + a*c)*sinh(b*c*x + a*c) + 3*(2*sqrt(2) - 3)*sinh(b*c*x + a*c)^2
+ 2*sqrt(2) - 3)/(cosh(b*c*x + a*c)^2 + sinh(b*c*x + a*c)^2 + 3)) - log(2
*(cosh(b*c*x + a*c)^2 + sinh(b*c*x + a*c)^2 + 3)/(cosh(b*c*x + a*c)^2 - 2*
cosh(b*c*x + a*c)*sinh(b*c*x + a*c) + sinh(b*c*x + a*c)^2)))/(b*c)
```

Sympy [F]

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atan}(\cosh(ac + bcx)) dx$$

input

```
integrate(exp(c*(b*x+a))*atan(cosh(b*c*x+a*c)),x)
```

output

```
exp(a*c)*Integral(exp(b*c*x)*atan(cosh(a*c + b*c*x)), x)
```

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.27

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx = \frac{\arctan(\cosh(bc x + ac)) e^{((bx+a)c)}}{bc} - \frac{\sqrt{2} \log\left(-\frac{2\sqrt{2}-e^{(-2bcx-2ac)}-3}{2\sqrt{2}+e^{(-2bcx-2ac)}+3}\right)}{2bc} - \frac{2(bc x + ac)}{bc} - \frac{\log(6e^{(-2bcx-2ac)} + e^{(-4bcx-4ac)} + 1)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctan(cosh(b*c*x+a*c)),x, algorithm="maxima")
```

output

```
arctan(cosh(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) - 1/2*sqrt(2)*log(-(2*sqrt(
2) - e^(-2*b*c*x - 2*a*c) - 3)/(2*sqrt(2) + e^(-2*b*c*x - 2*a*c) + 3))/(b
*c) - 2*(b*c*x + a*c)/(b*c) - 1/2*log(6*e^(-2*b*c*x - 2*a*c) + e^(-4*b*c*x
- 4*a*c) + 1)/(b*c)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.50

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx$$

$$= \frac{\left(\sqrt{2}e^{(-ac)} \log\left(-\frac{2\sqrt{2}e^{(2ac)} - e^{(2bcx+4ac)} - 3e^{(2ac)}}{2\sqrt{2}e^{(2ac)} + e^{(2bcx+4ac)} + 3e^{(2ac)}}\right) + 2 \arctan\left(\frac{1}{2}\left(e^{(2bcx+2ac)} + 1\right)e^{(-bcx-ac)}\right) e^{(bcx)} - e^{(-ac)} \log\right)}{2bc}$$

input `integrate(exp(c*(b*x+a))*arctan(cosh(b*c*x+a*c)),x, algorithm="giac")`

output `1/2*(sqrt(2)*e^(-a*c)*log(-(2*sqrt(2)*e^(2*a*c) - e^(2*b*c*x + 4*a*c) - 3*e^(2*a*c))/(2*sqrt(2)*e^(2*a*c) + e^(2*b*c*x + 4*a*c) + 3*e^(2*a*c))) + 2*arctan(1/2*(e^(2*b*c*x + 2*a*c) + 1)*e^(-b*c*x - a*c))*e^(b*c*x) - e^(-a*c)*log(e^(4*b*c*x + 4*a*c) + 6*e^(2*b*c*x + 2*a*c) + 1))*e^(a*c)/(b*c)`

Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.29

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx$$

$$= \frac{\ln(-8e^{2c(a+bx)} - 2\sqrt{2} - 6\sqrt{2}e^{2c(a+bx)}) (\sqrt{2} - 1)}{2bc}$$

$$- \frac{\ln(2\sqrt{2} - 8e^{2c(a+bx)} + 6\sqrt{2}e^{2c(a+bx)}) (\sqrt{2} + 1)}{2bc}$$

$$+ \frac{e^{ac+bcx} \operatorname{atan}\left(\frac{e^{bcx}e^{ac}}{2} + \frac{e^{-bcx}e^{-ac}}{2}\right)}{bc}$$

input `int(exp(c*(a + b*x))*atan(cosh(a*c + b*c*x)),x)`

output `(log(- 8*exp(2*c*(a + b*x)) - 2*2^(1/2) - 6*2^(1/2)*exp(2*c*(a + b*x)))*(2^(1/2) - 1))/(2*b*c) - (log(2*2^(1/2) - 8*exp(2*c*(a + b*x)) + 6*2^(1/2)*exp(2*c*(a + b*x)))*(2^(1/2) + 1))/(2*b*c) + (exp(a*c + b*c*x)*atan((exp(b*c*x)*exp(a*c))/2 + (exp(-b*c*x)*exp(-a*c))/2))/(b*c)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.73

$$\int e^{c(a+bx)} \arctan(\cosh(ac + bcx)) dx$$

$$= \frac{2e^{bcx+ac} \operatorname{atan}\left(\frac{e^{2bcx+2ac}+1}{2e^{bcx+ac}}\right) + \sqrt{2} \log(e^{bcx+ac} - \sqrt{2}i + i) + \sqrt{2} \log(e^{bcx+ac} + \sqrt{2}i - i) - \sqrt{2} \log(e^{2bcx+2ac})}{2bc}$$

input `int(exp(c*(b*x+a))*atan(cosh(b*c*x+a*c)),x)`output `(2*e**(a*c + b*c*x)*atan((e**(2*a*c + 2*b*c*x) + 1)/(2*e**(a*c + b*c*x))) + sqrt(2)*log(e**(a*c + b*c*x) - sqrt(2)*i + i) + sqrt(2)*log(e**(a*c + b*c*x) + sqrt(2)*i - i) - sqrt(2)*log(e**(2*a*c + 2*b*c*x) + 2*sqrt(2) + 3) - log(e**(a*c + b*c*x) - sqrt(2)*i + i) - log(e**(a*c + b*c*x) + sqrt(2)*i - i) - log(e**(2*a*c + 2*b*c*x) + 2*sqrt(2) + 3))/(2*b*c)`

3.149 $\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$

Optimal result	1087
Mathematica [C] (verified)	1088
Rubi [A] (verified)	1088
Maple [C] (warning: unable to verify)	1092
Fricas [A] (verification not implemented)	1093
Sympy [F]	1093
Maxima [A] (verification not implemented)	1094
Giac [A] (verification not implemented)	1094
Mupad [B] (verification not implemented)	1095
Reduce [B] (verification not implemented)	1095

Optimal result

Integrand size = 20, antiderivative size = 134

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx = \frac{\arctan(1 - \sqrt{2}e^{ac+bcx})}{\sqrt{2}bc} - \frac{\arctan(1 + \sqrt{2}e^{ac+bcx})}{\sqrt{2}bc} + \frac{e^{ac+bcx} \arctan(\tanh(c(a + bx)))}{bc} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}e^{ac+bcx}}{1+e^{2c(a+bx)}}\right)}{\sqrt{2}bc}$$

output

```
-1/2*arctan(-1+2^(1/2)*exp(b*c*x+a*c))*2^(1/2)/b/c-1/2*arctan(1+2^(1/2)*exp(b*c*x+a*c))*2^(1/2)/b/c+exp(b*c*x+a*c)*arctan(tanh(c*(b*x+a)))/b/c+1/2*arctanh(2^(1/2)*exp(b*c*x+a*c)/(1+exp(2*c*(b*x+a))))*2^(1/2)/b/c
```


Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.66

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$$

$$= \frac{2e^{c(a+bx)} \arctan\left(\frac{-1+e^{2c(a+bx)}}{1+e^{2c(a+bx)}}\right) + \text{RootSum}\left[1 + \#1^4 \&, \frac{ac+bcx-\log(e^{c(a+bx)}-\#1)}{\#1} \&\right]}{2bc}$$

input

```
Integrate[E^(c*(a + b*x))*ArcTan[Tanh[a*c + b*c*x]], x]
```

output

```
(2*E^(c*(a + b*x))*ArcTan[(-1 + E^(2*c*(a + b*x)))/(1 + E^(2*c*(a + b*x))]  
] + RootSum[1 + #1^4 & , (a*c + b*c*x - Log[E^(c*(a + b*x)) - #1]/#1 & ]  
/(2*b*c)
```

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.31, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {7281, 5730, 27, 2679, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$$

$$\downarrow \text{7281}$$

$$\frac{\int e^{ac+bx} \arctan(\tanh(ac + bxc)) d(ac + bxc)}{bc}$$

$$\downarrow \text{5730}$$

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - \int \frac{2e^{3(ac+bxc)}}{1+e^{4(ac+bxc)}} d(ac + bxc)}{bc}$$

$$\downarrow \text{27}$$

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \int \frac{e^{3(ac+bcx)}}{1+e^{4(ac+bcx)}} d(ac + bcx)}{bc}$$

↓ 2679

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \int \frac{e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx}}{bc}$$

↓ 826

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \left(\frac{1}{2} \int \frac{1+e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right)}{bc}$$

↓ 1476

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \left(\frac{1}{2} \int \frac{1}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx} + \frac{1}{2} \int \frac{1}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx} \right) - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx}}{bc}$$

↓ 1082

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \left(\frac{1}{2} \left(\int \frac{\frac{1}{-1-e^{2ac+2bcx}} d(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} - \int \frac{\frac{1}{-1-e^{2ac+2bcx}} d(1+\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right)}{bc}$$

↓ 217

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right)}{bc}$$

↓ 1479

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \left(\frac{1}{2} \left(\int \frac{-\frac{\sqrt{2}-2e^{ac+bcx}}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} + \int \frac{\sqrt{2}(1+\sqrt{2}e^{ac+bcx})}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} \right) + \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right)}{bc}$$

↓ 25

$$\frac{e^{ac+bcx} \arctan(\tanh(ac + bcx)) - 2 \left(\frac{1}{2} \left(-\int \frac{\frac{\sqrt{2}-2e^{ac+bcx}}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} - \int \frac{\sqrt{2}(1+\sqrt{2}e^{ac+bcx})}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} \right) + \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right)}{bc}$$

↓ 27

$$e^{ac+bcx} \arctan(\tanh(ac+bcx)) - 2 \left(\frac{1}{2} \left(- \int \frac{\sqrt{2}-2e^{ac+bcx}}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx} - \frac{1}{2} \int \frac{1+\sqrt{2}e^{ac+bcx}}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx} \right) + \frac{1}{2} \right)$$

$$bc$$

↓ 1103

$$e^{ac+bcx} \arctan(\tanh(ac+bcx)) - 2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}+1)}{2\sqrt{2}} + \frac{\log(\sqrt{2}e^{ac+bcx}+1)}{2\sqrt{2}} \right) \right)$$

$$bc$$

input `Int[E^(c*(a + b*x))*ArcTan[Tanh[a*c + b*c*x]],x]`

output `(E^(a*c + b*c*x)*ArcTan[Tanh[a*c + b*c*x]] - 2*((-(ArcTan[1 - Sqrt[2]*E^(a*c + b*c*x)]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*E^(a*c + b*c*x)]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*E^(a*c + b*c*x) + E^(2*a*c + 2*b*c*x)]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*E^(a*c + b*c*x) + E^(2*a*c + 2*b*c*x)]/(2*Sqrt[2]))/2)/(b*c)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2679 `Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_))))^(p_)*(G_)^((h_)*((f_) + (g_)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log[G]))]}, Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Denominator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`

rule 5730 `Int[((a_) + ArcTan[u]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_) /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]`

rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.91 (sec) , antiderivative size = 1355, normalized size of antiderivative = 10.11

method	result	size
risch	Expression too large to display	1355

input `int(exp(c*(b*x+a))*arctan(tanh(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/4/b/c*\exp(c*(b*x+a))*\text{Pi}+1/2*I/b/c*\exp(c*(b*x+a))*\ln(\exp(2*c*(b*x+a))+I)- \\ & 1/4/b/c*\text{Pi}*c\text{sgn}(I/(1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(2*c*(b*x+a))-I))*c\text{sgn}(\\ & I*(\exp(2*c*(b*x+a))-I)/(1+\exp(2*c*(b*x+a))))*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn} \\ & n(I/(1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(2*c*(b*x+a))+I))*c\text{sgn}(I*(\exp(2*c*(b* \\ & x+a))+I)/(1+\exp(2*c*(b*x+a))))*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(2*c*(\\ & b*x+a))-I)/(1+\exp(2*c*(b*x+a))))*c\text{sgn}((1-I)*(\exp(2*c*(b*x+a))-I)/(1+\exp(2* \\ & c*(b*x+a))))^2*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I/(1+\exp(2*c*(b*x+a))))*c\text{sgn} \\ & (I*(\exp(2*c*(b*x+a))-I)/(1+\exp(2*c*(b*x+a))))^2*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}* \\ & c\text{sgn}(I*(\exp(2*c*(b*x+a))-I))*c\text{sgn}(I*(\exp(2*c*(b*x+a))-I)/(1+\exp(2*c*(b*x+a) \\ &))))^2*\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I/(1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(\\ & 2*c*(b*x+a))+I)/(1+\exp(2*c*(b*x+a))))^2*\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I*(\\ & \exp(2*c*(b*x+a))+I))*c\text{sgn}(I*(\exp(2*c*(b*x+a))+I)/(1+\exp(2*c*(b*x+a))))^2*e \\ & xp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(2*c*(b*x+a))+I)/(1+\exp(2*c*(b*x+a)))) \\ & *c\text{sgn}((1+I)*(\exp(2*c*(b*x+a))+I)/(1+\exp(2*c*(b*x+a))))^2*\exp(c*(b*x+a))-1/ \\ & 4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(2*c*(b*x+a))-I)/(1+\exp(2*c*(b*x+a))))*c\text{sgn}((1-I)*(\exp \\ & (2*c*(b*x+a))-I)/(1+\exp(2*c*(b*x+a))))*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I*(e \\ & xp(2*c*(b*x+a))+I)/(1+\exp(2*c*(b*x+a))))*c\text{sgn}((1+I)*(\exp(2*c*(b*x+a))+I)/(\\ & 1+\exp(2*c*(b*x+a))))*\exp(c*(b*x+a))+1/4/b/c*\ln(\exp(c*(b*x+a))+1/2*2^(1/2)+ \\ & 1/2*I*2^(1/2))*2^(1/2)-1/4/b/c*\ln(\exp(c*(b*x+a))-1/2*I*2^(1/2)-1/2*2^(1/2) \\ &)*2^(1/2)-1/4/b/c*\ln(\exp(c*(b*x+a))+1/2*I*2^(1/2)-1/2*2^(1/2))*2^(1/2)+\dots \end{aligned}$$

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.57

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$$

$$= \frac{4(\cosh(bcx + ac) + \sinh(bcx + ac)) \arctan\left(\frac{\sinh(bcx+ac)}{\cosh(bcx+ac)}\right) - 2\sqrt{2} \arctan(\sqrt{2} \cosh(bcx + ac) + \sqrt{2} \sinh(bcx + ac))}{b}$$

input `integrate(exp(c*(b*x+a))*arctan(tanh(b*c*x+a*c)),x, algorithm="fricas")`

output `1/4*(4*(cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*arctan(sinh(b*c*x + a*c)/cosh(b*c*x + a*c)) - 2*sqrt(2)*arctan(sqrt(2)*cosh(b*c*x + a*c) + sqrt(2)*sinh(b*c*x + a*c) + 1) - 2*sqrt(2)*arctan(sqrt(2)*cosh(b*c*x + a*c) + sqrt(2)*sinh(b*c*x + a*c) - 1) + sqrt(2)*log((sqrt(2) + 2*cosh(b*c*x + a*c))/(cosh(b*c*x + a*c) - sinh(b*c*x + a*c))) - sqrt(2)*log(-(sqrt(2) - 2*cosh(b*c*x + a*c))/(cosh(b*c*x + a*c) - sinh(b*c*x + a*c))))/(b*c)`

Sympy [F]

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atan}(\tanh(ac + bcx)) dx$$

input `integrate(exp(c*(b*x+a))*atan(tanh(b*c*x+a*c)),x)`

output `exp(a*c)*Integral(exp(b*c*x)*atan(tanh(a*c + b*c*x)), x)`

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.25

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx = \frac{\arctan(\tanh(bc x + ac)) e^{((bx+a)c)}}{bc} - \frac{\sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2 e^{(bcx+ac)})\right)}{2bc} - \frac{\sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2 e^{(bcx+ac)})\right)}{2bc} + \frac{\sqrt{2} \log(\sqrt{2} e^{(bcx+ac)} + e^{(2bcx+2ac)} + 1)}{4bc} - \frac{\sqrt{2} \log(-\sqrt{2} e^{(bcx+ac)} + e^{(2bcx+2ac)} + 1)}{4bc}$$

input `integrate(exp(c*(b*x+a))*arctan(tanh(b*c*x+a*c)),x, algorithm="maxima")`

output `arctan(tanh(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^(b*c*x + a*c)))/(b*c) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^(b*c*x + a*c)))/(b*c) + 1/4*sqrt(2)*log(sqrt(2)*e^(b*c*x + a*c) + e^(2*b*c*x + 2*a*c) + 1)/(b*c) - 1/4*sqrt(2)*log(-sqrt(2)*e^(b*c*x + a*c) + e^(2*b*c*x + 2*a*c) + 1)/(b*c)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.76

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx = \frac{4 \pi e^{(bcx+ac)} \left[\frac{\pi+4 \arctan\left(\frac{e^{(2bcx+2ac)}}{4 \pi}\right)}{4 \pi} \right] + \pi e^{(bcx+ac)} - (4 \arctan(e^{(2bcx+2ac)})) e^{(bcx)} - (2 \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2 e^{(bcx+ac)})\right)) e^{(bcx+a)}}{4 \pi}$$

input `integrate(exp(c*(b*x+a))*arctan(tanh(b*c*x+a*c)),x, algorithm="giac")`

output

```
-1/4*(4*pi*e^(b*c*x + a*c)*floor(1/4*(pi + 4*arctan(e^(2*b*c*x + 2*a*c)))/
pi) + pi*e^(b*c*x + a*c) - (4*arctan(e^(2*b*c*x + 2*a*c))*e^(b*c*x) - (2*
sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-a*c) + 2*e^(b*c*x))*e^(a*c))*e^(-3*
a*c) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-a*c) - 2*e^(b*c*x))*e^(a
*c))*e^(-3*a*c) - sqrt(2)*e^(-3*a*c)*log(sqrt(2)*e^(b*c*x - a*c) + e^(2*b*
c*x) + e^(-2*a*c)) + sqrt(2)*e^(-3*a*c)*log(-sqrt(2)*e^(b*c*x - a*c) + e^(
2*b*c*x) + e^(-2*a*c)))*e^(2*a*c))*e^(a*c))/(b*c)
```

Mupad [B] (verification not implemented)

Time = 2.03 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.22

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$$

$$= \frac{4e^{ac+bcx} \operatorname{atan}\left(\frac{e^{2bcx}e^{2ac}-1}{e^{2bcx}e^{2ac}+1}\right) + \sqrt{2} \ln\left(\sqrt{2}(-4-4i) + e^{bcx}e^{ac}8i\right)(-1-i) + \sqrt{2} \ln\left(\sqrt{2}(-4+4i) - e^{bcx}e^{ac}8i\right)(-1+i)}{4}$$

input

```
int(exp(c*(a + b*x))*atan(tanh(a*c + b*c*x)),x)
```

output

```
(2^(1/2)*log(2^(1/2)*(4 - 4i) - exp(b*c*x)*exp(a*c)*8i)*(1 - 1i) - 2^(1/2)
*log(- 2^(1/2)*(4 - 4i) - exp(b*c*x)*exp(a*c)*8i)*(1 - 1i) - 2^(1/2)*log(e
xp(b*c*x)*exp(a*c)*8i - 2^(1/2)*(4 + 4i))*(1 + 1i) + 2^(1/2)*log(2^(1/2)*
(4 + 4i) + exp(b*c*x)*exp(a*c)*8i)*(1 + 1i) + 4*exp(a*c + b*c*x)*atan((exp(
2*b*c*x)*exp(2*a*c) - 1)/(exp(2*b*c*x)*exp(2*a*c) + 1)))/(4*b*c)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.26

$$\int e^{c(a+bx)} \arctan(\tanh(ac + bcx)) dx$$

$$= \frac{4e^{bcx+ac} \operatorname{atan}\left(\frac{e^{2bcx+2ac}-1}{e^{2bcx+2ac}+1}\right) - 2\sqrt{2} \operatorname{atan}\left(\frac{2e^{bcx+ac}-\sqrt{2}}{\sqrt{2}}\right) - 2\sqrt{2} \operatorname{atan}\left(\frac{2e^{bcx+ac}+\sqrt{2}}{\sqrt{2}}\right) - \sqrt{2} \log(e^{2bcx+2ac} - e^{bcx+ac})}{4bc}$$

input

```
int(exp(c*(b*x+a))*atan(tanh(b*c*x+a*c)),x)
```


output

```
(4*exp(a*c + b*c*x)*atan((exp(2*a*c + 2*b*c*x) - 1)/(exp(2*a*c + 2*b*c*x) + 1)) - 2*sqrt(2)*atan((2*exp(a*c + b*c*x) - sqrt(2))/sqrt(2)) - 2*sqrt(2)*atan((2*exp(a*c + b*c*x) + sqrt(2))/sqrt(2)) - sqrt(2)*log(exp(2*a*c + 2*b*c*x) - exp(a*c + b*c*x)*sqrt(2) + 1) + sqrt(2)*log(exp(2*a*c + 2*b*c*x) + exp(a*c + b*c*x)*sqrt(2) + 1))/(4*b*c)
```

3.150 $\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx$

Optimal result	1097
Mathematica [C] (verified)	1098
Rubi [A] (verified)	1098
Maple [C] (warning: unable to verify)	1102
Fricas [A] (verification not implemented)	1103
Sympy [F]	1103
Maxima [A] (verification not implemented)	1104
Giac [B] (verification not implemented)	1104
Mupad [B] (verification not implemented)	1105
Reduce [B] (verification not implemented)	1105

Optimal result

Integrand size = 20, antiderivative size = 135

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx = -\frac{\arctan(1 - \sqrt{2}e^{ac+bcx})}{\sqrt{2}bc} + \frac{\arctan(1 + \sqrt{2}e^{ac+bcx})}{\sqrt{2}bc} + \frac{e^{ac+bcx} \arctan(\coth(c(a + bx)))}{bc} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}e^{ac+bcx}}{1+e^{2c(a+bx)}}\right)}{\sqrt{2}bc}$$

output

```
1/2*arctan(-1+2^(1/2)*exp(b*c*x+a*c))*2^(1/2)/b/c+1/2*arctan(1+2^(1/2)*exp
(b*c*x+a*c))*2^(1/2)/b/c+exp(b*c*x+a*c)*arctan(coth(c*(b*x+a)))/b/c-1/2*ar
ctanh(2^(1/2)*exp(b*c*x+a*c)/(1+exp(2*c*(b*x+a))))*2^(1/2)/b/c
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.66

$$\int e^{c(a+bx)} \arctan(\coth(ac+bcx)) dx$$

$$= \frac{2e^{c(a+bx)} \arctan\left(\frac{1+e^{2c(a+bx)}}{-1+e^{2c(a+bx)}}\right) + \text{RootSum}\left[1 + \#1^4 \&, \frac{-ac-bcx+\log(e^{c(a+bx)}-\#1)}{\#1} \&\right]}{2bc}$$

input

```
Integrate[E^(c*(a + b*x))*ArcTan[Coth[a*c + b*c*x]],x]
```

output

```
(2*E^(c*(a + b*x))*ArcTan[(1 + E^(2*c*(a + b*x)))/(-1 + E^(2*c*(a + b*x))]) + RootSum[1 + #1^4 & , (-a*c) - b*c*x + Log[E^(c*(a + b*x)) - #1]/#1 & ])/(2*b*c)
```

Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.30, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {7281, 5730, 27, 2679, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^{c(a+bx)} \arctan(\coth(ac+bcx)) dx$$

$$\downarrow 7281$$

$$\frac{\int e^{ac+bx} \arctan(\coth(ac+bcx)) d(ac+bcx)}{bc}$$

$$\downarrow 5730$$

$$\frac{e^{ac+bcx} \arctan(\coth(ac+bcx)) - \int \frac{2e^{3(ac+bcx)}}{1+e^{4(ac+bcx)}} d(ac+bcx)}{bc}$$

$$\downarrow 27$$

$$\frac{2 \int \frac{e^{3(ac+bcx)}}{1+e^{4(ac+bcx)}} d(ac+bcx) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 2679

$$\frac{2 \int \frac{e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 826

$$\frac{2 \left(\frac{1}{2} \int \frac{1+e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 1476

$$\frac{2 \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx} + \frac{1}{2} \int \frac{1}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx} \right) - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 1082

$$\frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{1}{-1-e^{2ac+2bcx}} d(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} - \frac{\int \frac{1}{-1-e^{2ac+2bcx}} d(1+\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 217

$$\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-e^{2ac+2bcx}}{1+e^{4ac+4bcx}} de^{ac+bcx} \right) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 1479

$$\frac{2 \left(\frac{1}{2} \left(\frac{\int -\frac{\sqrt{2}-2e^{ac+bcx}}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} + \frac{\int -\frac{\sqrt{2}(1+\sqrt{2}e^{ac+bcx})}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) \right) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 25

$$\frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2e^{ac+bcx}}{1-\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(1+\sqrt{2}e^{ac+bcx})}{1+\sqrt{2}e^{ac+bcx}+e^{2ac+2bcx}} de^{ac+bcx}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) \right) + e^{ac+bcx} \arctan(\coth(ac+bcx))}{bc}$$

↓ 27

$$2 \left(\frac{1}{2} \left(- \int \frac{\sqrt{2} - 2e^{ac+bcx}}{1 - \sqrt{2}e^{ac+bcx} + e^{2ac+2bcx}} de^{ac+bcx} - \frac{1}{2} \int \frac{1 + \sqrt{2}e^{ac+bcx}}{1 + \sqrt{2}e^{ac+bcx} + e^{2ac+2bcx}} de^{ac+bcx} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) \right) \frac{1}{bc}$$

↓ 1103

$$2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}e^{ac+bcx} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}e^{ac+bcx})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(-\sqrt{2}e^{ac+bcx} + e^{2ac+2bcx} + 1)}{2\sqrt{2}} - \frac{\log(\sqrt{2}e^{ac+bcx} + e^{2ac+2bcx} + 1)}{2\sqrt{2}} \right) \right) \frac{1}{bc}$$

input `Int[E^(c*(a + b*x))*ArcTan[Coth[a*c + b*c*x]],x]`

output `(E^(a*c + b*c*x)*ArcTan[Coth[a*c + b*c*x]] + 2*((-(ArcTan[1 - Sqrt[2]*E^(a*c + b*c*x)]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*E^(a*c + b*c*x)]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*E^(a*c + b*c*x) + E^(2*a*c + 2*b*c*x)]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*E^(a*c + b*c*x) + E^(2*a*c + 2*b*c*x)]/(2*Sqrt[2]))/2)/(b*c)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2679 `Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_))))^(p_)*(G_)^((h_)*((f_) + (g_)*(x_))), x_Symbol] := With[{m = FullSimplify[d*e*(Log[F]/(g*h*Log[G]))]}, Simp[Denominator[m]/(g*h*Log[G]) Subst[Int[x^(Denominator[m] - 1)*(a + b*F^(c*e - d*e*(f/g))*x^Numerator[m])^p, x], x, G^(h*((f + g*x)/Denominator[m]))], x] /; LtQ[m, -1] || GtQ[m, 1] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x]`

rule 5730 `Int[((a_) + ArcTan[u_]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_) /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]`

rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.84 (sec) , antiderivative size = 1355, normalized size of antiderivative = 10.04

method	result	size
risch	Expression too large to display	1355

input `int(exp(c*(b*x+a))*arctan(coth(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

```
-1/2*I/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))+I)+1/4/b/c*exp(c*(b*x+a))*Pi
+1/4/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))-I)/(-1+exp(2*c*(b*x+a))))^3*exp(c*(b*
x+a))+1/4/b/c*Pi*csgn((1+I)*(exp(2*c*(b*x+a))-I)/(-1+exp(2*c*(b*x+a))))^3*
exp(c*(b*x+a))+1/4/b/c*Pi*csgn((1-I)*(exp(2*c*(b*x+a))+I)/(-1+exp(2*c*(b*x
+a))))^3*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))+I)/(-1+exp(2*c
*(b*x+a))))^3*exp(c*(b*x+a))-1/4/b/c*Pi*csgn((1+I)*(exp(2*c*(b*x+a))-I)/(-
1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*csgn((1-I)*(exp(2*c*(b*x+
a))+I)/(-1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))+1/4*I/b/c*ln(exp(c*(b*x+a))
+1/2*I*2^(1/2)-1/2*2^(1/2))*2^(1/2)-1/4*I/b/c*ln(exp(c*(b*x+a))+1/2*2^(1/2)
)-1/2*I*2^(1/2))*2^(1/2)+1/2*I/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))-I)+1
/4*I/b/c*ln(exp(c*(b*x+a))+1/2*2^(1/2)+1/2*I*2^(1/2))*2^(1/2)-1/4*I/b/c*ln
(exp(c*(b*x+a))-1/2*I*2^(1/2)-1/2*2^(1/2))*2^(1/2)+1/4/b/c*Pi*csgn(I*(exp(
2*c*(b*x+a))-I))*csgn(I/(-1+exp(2*c*(b*x+a))))*csgn(I*(exp(2*c*(b*x+a))-I)
/(-1+exp(2*c*(b*x+a))))*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*(exp(2*c*(b*x+a)
+I))*csgn(I/(-1+exp(2*c*(b*x+a))))*csgn(I*(exp(2*c*(b*x+a))+I)/(-1+exp(2*c
*(b*x+a))))*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))-I))*csgn(I*
(exp(2*c*(b*x+a))-I)/(-1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*cs
gn(I/(-1+exp(2*c*(b*x+a))))*csgn(I*(exp(2*c*(b*x+a))-I)/(-1+exp(2*c*(b*x+a)
))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))-I)/(-1+exp(2*c*(
b*x+a))))*csgn((1+I)*(exp(2*c*(b*x+a))-I)/(-1+exp(2*c*(b*x+a))))^2*exp(...
```

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.56

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx$$

$$= \frac{4(\cosh(bcx + ac) + \sinh(bcx + ac)) \arctan\left(\frac{\cosh(bcx+ac)}{\sinh(bcx+ac)}\right) + 2\sqrt{2} \arctan(\sqrt{2} \cosh(bcx + ac) + \sqrt{2} \sinh(bcx + ac))}{b}$$

input `integrate(exp(c*(b*x+a))*arctan(coth(b*c*x+a*c)),x, algorithm="fricas")`

output `1/4*(4*(cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*arctan(cosh(b*c*x + a*c)/sinh(b*c*x + a*c)) + 2*sqrt(2)*arctan(sqrt(2)*cosh(b*c*x + a*c) + sqrt(2)*sinh(b*c*x + a*c) + 1) + 2*sqrt(2)*arctan(sqrt(2)*cosh(b*c*x + a*c) + sqrt(2)*sinh(b*c*x + a*c) - 1) - sqrt(2)*log((sqrt(2) + 2*cosh(b*c*x + a*c))/(cosh(b*c*x + a*c) - sinh(b*c*x + a*c))) + sqrt(2)*log(-(sqrt(2) - 2*cosh(b*c*x + a*c))/(cosh(b*c*x + a*c) - sinh(b*c*x + a*c)))/(b*c)`

Sympy [F]

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atan}(\coth(ac + bcx)) dx$$

input `integrate(exp(c*(b*x+a))*atan(coth(b*c*x+a*c)),x)`

output `exp(a*c)*Integral(exp(b*c*x)*atan(coth(a*c + b*c*x)), x)`

Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.24

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx = \frac{\arctan(\coth(bc x + ac)) e^{((bx+a)c)}}{bc} + \frac{\sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2 e^{(bcx+ac)})\right)}{2bc} + \frac{\sqrt{2} \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2} - 2 e^{(bcx+ac)})\right)}{2bc} - \frac{\sqrt{2} \log(\sqrt{2} e^{(bcx+ac)} + e^{(2bcx+2ac)} + 1)}{4bc} + \frac{\sqrt{2} \log(-\sqrt{2} e^{(bcx+ac)} + e^{(2bcx+2ac)} + 1)}{4bc}$$

input `integrate(exp(c*(b*x+a))*arctan(coth(b*c*x+a*c)),x, algorithm="maxima")`

output `arctan(coth(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) + 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^(b*c*x + a*c)))/(b*c) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^(b*c*x + a*c)))/(b*c) - 1/4*sqrt(2)*log(sqrt(2)*e^(b*c*x + a*c) + e^(2*b*c*x + 2*a*c) + 1)/(b*c) + 1/4*sqrt(2)*log(-sqrt(2)*e^(b*c*x + a*c) + e^(2*b*c*x + 2*a*c) + 1)/(b*c)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(118) = 236.

Time = 0.13 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.77

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx = \frac{4 \pi e^{(bcx+ac)} \left[\frac{3 \pi + 4 \arctan(e^{(2bcx+2ac)})}{4 \pi} \right] - \pi e^{(bcx+ac)} - (4 \arctan(e^{(2bcx+2ac)}) e^{(bcx)} - (2 \sqrt{2} \arctan(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2 e^{(bcx+ac)}))) e^{(bcx+ac)}}}{4 \pi}$$

input `integrate(exp(c*(b*x+a))*arctan(coth(b*c*x+a*c)),x, algorithm="giac")`

output

```
1/4*(4*pi*e^(b*c*x + a*c)*floor(1/4*(3*pi + 4*arctan(e^(2*b*c*x + 2*a*c)))/pi) - pi*e^(b*c*x + a*c) - (4*arctan(e^(2*b*c*x + 2*a*c))*e^(b*c*x) - (2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-a*c) + 2*e^(b*c*x))*e^(a*c))*e^(-3*a*c) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-a*c) - 2*e^(b*c*x))*e^(a*c))*e^(-3*a*c) - sqrt(2)*e^(-3*a*c)*log(sqrt(2)*e^(b*c*x - a*c) + e^(2*b*c*x) + e^(-2*a*c)) + sqrt(2)*e^(-3*a*c)*log(-sqrt(2)*e^(b*c*x - a*c) + e^(2*b*c*x) + e^(-2*a*c)))*e^(2*a*c))*e^(a*c))/(b*c)
```

Mupad [B] (verification not implemented)

Time = 1.13 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.21

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx$$

$$= \frac{4e^{ac+bcx} \operatorname{atan}\left(\frac{e^{2bcx}e^{2ac}+1}{e^{2bcx}e^{2ac}-1}\right) + \sqrt{2} \ln\left(\sqrt{2}(-4-4i) - e^{bcx}e^{ac}8i\right)(-1-i) + \sqrt{2} \ln\left(\sqrt{2}(-4+4i) + e^{bcx}e^{ac}8i\right)(-1+i)}{4bc}$$

input

```
int(exp(c*(a + b*x))*atan(coth(a*c + b*c*x)),x)
```

output

```
(2^(1/2)*log(2^(1/2)*(4 - 4i) + exp(b*c*x)*exp(a*c)*8i)*(1 - 1i) - 2^(1/2)*log(exp(b*c*x)*exp(a*c)*8i - 2^(1/2)*(4 - 4i))*(1 - 1i) - 2^(1/2)*log(-2^(1/2)*(4 + 4i) - exp(b*c*x)*exp(a*c)*8i)*(1 + 1i) + 2^(1/2)*log(2^(1/2)*(4 + 4i) - exp(b*c*x)*exp(a*c)*8i)*(1 + 1i) + 4*exp(a*c + b*c*x)*atan((exp(2*b*c*x)*exp(2*a*c) + 1)/(exp(2*b*c*x)*exp(2*a*c) - 1)))/(4*b*c)
```

Reduce [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.25

$$\int e^{c(a+bx)} \arctan(\coth(ac + bcx)) dx$$

$$= \frac{4e^{bcx+ac} \operatorname{atan}\left(\frac{e^{2bcx+2ac}+1}{e^{2bcx+2ac}-1}\right) + 2\sqrt{2} \operatorname{atan}\left(\frac{2e^{bcx+ac}-\sqrt{2}}{\sqrt{2}}\right) + 2\sqrt{2} \operatorname{atan}\left(\frac{2e^{bcx+ac}+\sqrt{2}}{\sqrt{2}}\right) + \sqrt{2} \log(e^{2bcx+2ac} - e^{bcx+ac})}{4bc}$$

input

```
int(exp(c*(b*x+a))*atan(coth(b*c*x+a*c)),x)
```

output

```
(4*e**(a*c + b*c*x)*atan((e**(2*a*c + 2*b*c*x) + 1)/(e**(2*a*c + 2*b*c*x)
- 1)) + 2*sqrt(2)*atan((2*e**(a*c + b*c*x) - sqrt(2))/sqrt(2)) + 2*sqrt(2)
*atan((2*e**(a*c + b*c*x) + sqrt(2))/sqrt(2)) + sqrt(2)*log(e**(2*a*c + 2*
b*c*x) - e**(a*c + b*c*x)*sqrt(2) + 1) - sqrt(2)*log(e**(2*a*c + 2*b*c*x)
+ e**(a*c + b*c*x)*sqrt(2) + 1))/(4*b*c)
```

3.151 $\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx$

Optimal result	1107
Mathematica [C] (verified)	1107
Rubi [A] (verified)	1108
Maple [C] (warning: unable to verify)	1110
Fricas [B] (verification not implemented)	1111
Sympy [F(-1)]	1112
Maxima [A] (verification not implemented)	1112
Giac [A] (verification not implemented)	1113
Mupad [B] (verification not implemented)	1113
Reduce [B] (verification not implemented)	1114

Optimal result

Integrand size = 20, antiderivative size = 103

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx = \frac{e^{ac+bcx} \arctan(\operatorname{sech}(c(a + bx)))}{bc} + \frac{(1 - \sqrt{2}) \log(3 - 2\sqrt{2} + e^{2c(a+bx)})}{2bc} + \frac{(1 + \sqrt{2}) \log(3 + 2\sqrt{2} + e^{2c(a+bx)})}{2bc}$$

output

```
exp(b*c*x+a*c)*arctan(sech(c*(b*x+a)))/b/c+1/2*(1-2^(1/2))*ln(3-2*2^(1/2)+exp(2*c*(b*x+a)))/b/c+1/2*(1+2^(1/2))*ln(3+2*2^(1/2)+exp(2*c*(b*x+a)))/b/c
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.08 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.41

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx = \frac{4c(a + bx) + 2e^{c(a+bx)} \arctan\left(\frac{2e^{c(a+bx)}}{1+e^{2c(a+bx)}}\right) + \operatorname{RootSum}\left[1 + 6\#1^2 + \#1^4 \&, \frac{-ac-bcx+\log(e^{c(a+bx)}-\#1)-7ac\#1}{1+e^{2c(a+bx)}}\right]}{2bc}$$

input `Integrate[E^(c*(a + b*x))*ArcTan[Sech[a*c + b*c*x]],x]`

output $(4*c*(a + b*x) + 2*E^{c*(a + b*x)}*ArcTan[(2*E^{c*(a + b*x)})/(1 + E^{2*c*(a + b*x)})] + RootSum[1 + 6*#1^2 + #1^4 \& , (-a*c) - b*c*x + Log[E^{c*(a + b*x)} - #1] - 7*a*c*#1^2 - 7*b*c*x*#1^2 + 7*Log[E^{c*(a + b*x)} - #1]*#1^2)/(1 + 3*#1^2) \&])/(2*b*c)$

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {7281, 5730, 25, 2720, 27, 1576, 1141, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bxc)) dx \\
 & \quad \downarrow 7281 \\
 & \int e^{ac+bx} \arctan(\operatorname{sech}(ac + bxc)) d(ac + bxc) \\
 & \quad \downarrow 5730 \\
 & \frac{e^{ac+bx} \arctan(\operatorname{sech}(ac + bxc)) - \int -\frac{e^{ac+bx} \operatorname{sech}(ac+bx) \tanh(ac+bx)}{\operatorname{sech}^2(ac+bx)+1} d(ac + bxc)}{bc} \\
 & \quad \downarrow 25 \\
 & \int \frac{e^{ac+bx} \operatorname{sech}(ac+bx) \tanh(ac+bx)}{\operatorname{sech}^2(ac+bx)+1} d(ac + bxc) + e^{ac+bx} \arctan(\operatorname{sech}(ac + bxc)) \\
 & \quad \downarrow 2720 \\
 & \int -\frac{2e^{ac+bx}(1-e^{2ac+2bx})}{1+6e^{2ac+2bx}+e^{4ac+4bx}} de^{ac+bx} + e^{ac+bx} \arctan(\operatorname{sech}(ac + bxc)) \\
 & \quad \downarrow 27 \\
 & \frac{e^{ac+bx} \arctan(\operatorname{sech}(ac + bxc)) - 2 \int \frac{e^{ac+bx}(1-e^{2ac+2bx})}{1+6e^{2ac+2bx}+e^{4ac+4bx}} de^{ac+bx}}{bc}
 \end{aligned}$$

$$\begin{array}{c}
 \downarrow 1576 \\
 \frac{e^{ac+bcx} \arctan(\operatorname{sech}(ac+bcx)) - \int \frac{-ac-bxc+1}{1+7e^{2ac+2bcx}} de^{2ac+2bcx}}{bc} \\
 \downarrow 1141 \\
 \frac{e^{ac+bcx} \arctan(\operatorname{sech}(ac+bcx)) - \int \left(-\frac{2-\sqrt{2}}{2(4-3\sqrt{2}-\sqrt{2}e^{2ac+2bcx})} - \frac{1+\sqrt{2}}{2(3+2\sqrt{2}+e^{2ac+2bcx})} \right) de^{2ac+2bcx}}{bc} \\
 \downarrow 2009 \\
 \frac{e^{ac+bcx} \arctan(\operatorname{sech}(ac+bcx)) + \frac{1}{2}(1+\sqrt{2}) \log(e^{2ac+2bcx} + 3 + 2\sqrt{2}) + \frac{1}{2}(1-\sqrt{2}) \log(-\sqrt{2}e^{2ac+2bcx} + 4 - 3\sqrt{2})}{bc}
 \end{array}$$

input `Int[E^(c*(a + b*x))*ArcTan[Sech[a*c + b*c*x]], x]`

output `(E^(a*c + b*c*x)*ArcTan[Sech[a*c + b*c*x]] + ((1 + Sqrt[2])*Log[3 + 2*Sqrt[2] + E^(2*a*c + 2*b*c*x)])/2 + ((1 - Sqrt[2])*Log[4 - 3*Sqrt[2] - Sqrt[2]*E^(2*a*c + 2*b*c*x)])/2)/(b*c)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1141 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[1/c^p Int[ExpandIntegrand[(d + e*x)^m*(b/2 - q/2 + c*x)^p*(b/2 + q/2 + c*x)^p, x], x], x] /; EqQ[p, -1] || !FractionalPowerFactorQ[q] /; FreeQ[{a, b, c, d, e}, x] && ILtQ[p, 0] && IntegerQ[m] && NiceSqrtQ[b^2 - 4*a*c]`

rule 1576 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 5730 `Int[((a_) + ArcTan[u_]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]]`

rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 4.10 (sec) , antiderivative size = 838, normalized size of antiderivative = 8.14

method	result	size
risch	Expression too large to display	838

input `int(exp(c*(b*x+a))*arctan(sech(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output

```

-1/2*I/b/c*exp(c*(b*x+a))*ln(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a)))-1/4/b/
c*Pi*csgn(I*exp(2*c*(b*x+a))+I+2*exp(c*(b*x+a)))*csgn(I/(1+exp(2*c*(b*x+a)
)))*csgn(I*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a)))
)*exp(c*(b*x+a))-1/4/b/c*Pi*csgn(I*exp(2*c*(b*x+a))+I+2*exp(c*(b*x+a)))*csgn
(I*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a))))^2*exp(c*
(b*x+a))-1/4/b/c*Pi*csgn(I/(1+exp(2*c*(b*x+a))))*csgn(I*(-exp(2*c*(b*x+a))
-1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))+1/4/b/c*Pi*c
sgn(-I*exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-I)*csgn(I/(1+exp(2*c*(b*x+a))))*c
sgn(I*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a))))*exp(c*
(b*x+a))+1/4/b/c*Pi*csgn(I/(1+exp(2*c*(b*x+a))))*csgn(I*(exp(2*c*(b*x+a))+
1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a))))^2*exp(c*(b*x+a))-1/4/b/c*Pi*c
sgn(I*(-exp(2*c*(b*x+a))-1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a))))^3*exp(
c*(b*x+a))-1/4/b/c*Pi*csgn(-I*exp(2*c*(b*x+a))+2*exp(c*(b*x+a))-I)*csgn(I*
(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*(b*x+a))))^2*exp(c*(b*x
+a))-1/4/b/c*Pi*csgn(I*(exp(2*c*(b*x+a))+1+2*I*exp(c*(b*x+a)))/(1+exp(2*c*
(b*x+a))))^3*exp(c*(b*x+a))-1/2/b/c*2^(1/2)*ln(exp(2*c*(b*x+a))+(2^(1/2)-1
)^2)+1/2/b/c*2^(1/2)*ln(exp(2*c*(b*x+a))+(1+2^(1/2))^2)-2*a/b+1/2*I/b/c*ex
p(c*(b*x+a))*ln(exp(2*c*(b*x+a))+1-2*I*exp(c*(b*x+a)))+1/2/b/c*ln(exp(2*c*
(b*x+a))+(2^(1/2)-1)^2)+1/2/b/c*ln(exp(2*c*(b*x+a))+(1+2^(1/2))^2)

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 276 vs. $2(86) = 172$.

Time = 0.12 (sec) , antiderivative size = 276, normalized size of antiderivative = 2.68

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx$$

$$= \frac{2(\cosh(bc x + ac) + \sinh(bc x + ac)) \arctan\left(\frac{2(\cosh(bc x + ac) + \sinh(bc x + ac))}{\cosh(bc x + ac)^2 + 2 \cosh(bc x + ac) \sinh(bc x + ac) + \sinh(bc x + ac)^2 + 1}\right) + \sqrt{2} \log\left(\frac{2(\cosh(bc x + ac) + \sinh(bc x + ac))}{\cosh(bc x + ac)^2 + 2 \cosh(bc x + ac) \sinh(bc x + ac) + \sinh(bc x + ac)^2 + 1}\right)}{2}$$

input

```
integrate(exp(c*(b*x+a))*arctan(sech(b*c*x+a*c)),x, algorithm="fricas")
```


output

```
1/2*(2*(cosh(b*c*x + a*c) + sinh(b*c*x + a*c))*arctan(2*(cosh(b*c*x + a*c)
+ sinh(b*c*x + a*c))/(cosh(b*c*x + a*c)^2 + 2*cosh(b*c*x + a*c)*sinh(b*c*
x + a*c) + sinh(b*c*x + a*c)^2 + 1)) + sqrt(2)*log((3*(2*sqrt(2) + 3)*cosh
(b*c*x + a*c)^2 - 4*(3*sqrt(2) + 4)*cosh(b*c*x + a*c)*sinh(b*c*x + a*c) +
3*(2*sqrt(2) + 3)*sinh(b*c*x + a*c)^2 + 2*sqrt(2) + 3)/(cosh(b*c*x + a*c)^
2 + sinh(b*c*x + a*c)^2 + 3)) + log(2*(cosh(b*c*x + a*c)^2 + sinh(b*c*x +
a*c)^2 + 3)/(cosh(b*c*x + a*c)^2 - 2*cosh(b*c*x + a*c)*sinh(b*c*x + a*c) +
sinh(b*c*x + a*c)^2)))/(b*c)
```

Sympy [F(-1)]

Timed out.

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx = \text{Timed out}$$

input

```
integrate(exp(c*(b*x+a))*atan(sech(b*c*x+a*c)),x)
```

output

Timed out

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.64

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx = \frac{\arctan(\operatorname{sech}(bcx + ac)) e^{((bx+a)c)}}{bc} - \frac{3\sqrt{2} \log\left(-\frac{2\sqrt{2}-e^{(2bcx+2ac)}-3}{2\sqrt{2}+e^{(2bcx+2ac)}+3}\right)}{8bc} + \frac{\sqrt{2} \log\left(-\frac{2\sqrt{2}-e^{(-2bcx-2ac)}-3}{2\sqrt{2}+e^{(-2bcx-2ac)}+3}\right)}{8bc} + \frac{\log(e^{(4bcx+4ac)} + 6e^{(2bcx+2ac)} + 1)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctan(sech(b*c*x+a*c)),x, algorithm="maxima")
```

output

```
arctan(sech(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) - 3/8*sqrt(2)*log(-(2*sqrt(2) - e^(2*b*c*x + 2*a*c) - 3)/(2*sqrt(2) + e^(2*b*c*x + 2*a*c) + 3))/(b*c) + 1/8*sqrt(2)*log(-(2*sqrt(2) - e^(-2*b*c*x - 2*a*c) - 3)/(2*sqrt(2) + e^(-2*b*c*x - 2*a*c) + 3))/(b*c) + 1/2*log(e^(4*b*c*x + 4*a*c) + 6*e^(2*b*c*x + 2*a*c) + 1)/(b*c)
```

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.50

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx = \frac{\left(\sqrt{2}e^{(-ac)} \log\left(-\frac{2\sqrt{2}e^{(2ac)} - e^{(2bcx+4ac)} - 3}{2\sqrt{2}e^{(2ac)} + e^{(2bcx+4ac)} + 3}e^{(2ac)}\right) - 2 \arctan\left(\frac{2e^{(bcx+ac)}}{e^{(2bcx+2ac)} + 1}\right) e^{(bcx)} - e^{(-ac)} \log\left(e^{(4bcx+4ac)} + 6e^{(2bcx+2ac)} + 1\right)\right)}{2bc}$$

input

```
integrate(exp(c*(b*x+a))*arctan(sech(b*c*x+a*c)),x, algorithm="giac")
```

output

```
-1/2*(sqrt(2)*e^(-a*c)*log(-(2*sqrt(2)*e^(2*a*c) - e^(2*b*c*x + 4*a*c) - 3*e^(2*a*c))/(2*sqrt(2)*e^(2*a*c) + e^(2*b*c*x + 4*a*c) + 3*e^(2*a*c))) - 2*arctan(2*e^(b*c*x + a*c)/(e^(2*b*c*x + 2*a*c) + 1))*e^(b*c*x) - e^(-a*c)*log(e^(4*b*c*x + 4*a*c) + 6*e^(2*b*c*x + 2*a*c) + 1))*e^(a*c)/(b*c)
```

Mupad [B] (verification not implemented)

Time = 0.81 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.31

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx = \frac{e^{ac+bcx} \operatorname{atan}\left(\frac{1}{\frac{e^{bcx}e^{ac}}{2} + \frac{e^{-bcx}e^{-ac}}{2}}\right)}{bc} + \frac{\ln(8e^{2c(a+bx)} - 2\sqrt{2} - 6\sqrt{2}e^{2c(a+bx)}) (\sqrt{2} + 1)}{2bc} - \frac{\ln(8e^{2c(a+bx)} + 2\sqrt{2} + 6\sqrt{2}e^{2c(a+bx)}) (\sqrt{2} - 1)}{2bc}$$

input `int(atan(1/cosh(a*c + b*c*x))*exp(c*(a + b*x)),x)`

output
$$\frac{(\exp(a*c + b*c*x)*\operatorname{atan}(1/((\exp(b*c*x)*\exp(a*c))/2 + (\exp(-b*c*x)*\exp(-a*c))/2)))/(b*c) + (\log(8*\exp(2*c*(a + b*x)) - 2*2^{(1/2)} - 6*2^{(1/2)*\exp(2*c*(a + b*x))})*(2^{(1/2)} + 1))/(2*b*c) - (\log(8*\exp(2*c*(a + b*x)) + 2*2^{(1/2)} + 6*2^{(1/2)*\exp(2*c*(a + b*x))})*(2^{(1/2)} - 1))/(2*b*c)}$$

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.68

$$\int e^{c(a+bx)} \arctan(\operatorname{sech}(ac + bcx)) dx$$

$$= \frac{2e^{bcx+ac} \operatorname{atan}\left(\frac{2e^{bcx+ac}}{e^{2bcx+2ac}+1}\right) - \sqrt{2} \log(e^{bcx+ac} - \sqrt{2}i + i) - \sqrt{2} \log(e^{bcx+ac} + \sqrt{2}i - i) + \sqrt{2} \log(e^{2bcx+2ac} - \sqrt{2}i + i) + \sqrt{2} \log(e^{2bcx+2ac} + \sqrt{2}i - i)}{2bc}$$

input `int(exp(c*(b*x+a))*atan(sech(b*c*x+a*c)),x)`

output
$$\frac{(2*e^{**}(a*c + b*c*x)*\operatorname{atan}((2*e^{**}(a*c + b*c*x))/(e^{**}(2*a*c + 2*b*c*x) + 1)) - \sqrt{2}*\log(e^{**}(a*c + b*c*x) - \sqrt{2}*i + i) - \sqrt{2}*\log(e^{**}(a*c + b*c*x) + \sqrt{2}*i - i) + \sqrt{2}*\log(e^{**}(2*a*c + 2*b*c*x) + 2*\sqrt{2} + 3) + \log(e^{**}(a*c + b*c*x) - \sqrt{2}*i + i) + \log(e^{**}(a*c + b*c*x) + \sqrt{2}*i - i) + \log(e^{**}(2*a*c + 2*b*c*x) + 2*\sqrt{2} + 3))/(2*b*c)}$$

3.152 $\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx$

Optimal result	1115
Mathematica [A] (verified)	1115
Rubi [A] (verified)	1116
Maple [C] (warning: unable to verify)	1117
Fricas [B] (verification not implemented)	1118
Sympy [F]	1119
Maxima [A] (verification not implemented)	1119
Giac [A] (verification not implemented)	1120
Mupad [B] (verification not implemented)	1120
Reduce [B] (verification not implemented)	1121

Optimal result

Integrand size = 20, antiderivative size = 47

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx = \frac{e^{ac+bcx} \arctan(\operatorname{csch}(c(a + bx)))}{bc} + \frac{\log(1 + e^{2c(a+bx)})}{bc}$$

output

```
exp(b*c*x+a*c)*arctan(csch(c*(b*x+a)))/b/c+ln(1+exp(2*c*(b*x+a)))/b/c
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.21

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx = \frac{e^{c(a+bx)} \arctan\left(\frac{2e^{c(a+bx)}}{-1+e^{2c(a+bx)}}\right) + \log(1 + e^{2c(a+bx)})}{bc}$$

input

```
Integrate[E^(c*(a + b*x))*ArcTan[Csch[a*c + b*c*x]], x]
```

output

```
(E^(c*(a + b*x))*ArcTan[(2*E^(c*(a + b*x)))/(-1 + E^(2*c*(a + b*x))]) + Log[1 + E^(2*c*(a + b*x))]/(b*c)
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {7281, 5730, 25, 2720, 27, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int e^{c(a+bx)} \arctan(\operatorname{csch}(ac+bcx)) dx \\
 & \quad \downarrow \text{7281} \\
 & \frac{\int e^{ac+bcx} \arctan(\operatorname{csch}(ac+bcx)) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{5730} \\
 & \frac{e^{ac+bcx} \arctan(\operatorname{csch}(ac+bcx)) - \int -e^{ac+bcx} \operatorname{sech}(ac+bcx) d(ac+bcx)}{bc} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int e^{ac+bcx} \operatorname{sech}(ac+bcx) d(ac+bcx) + e^{ac+bcx} \arctan(\operatorname{csch}(ac+bcx))}{bc} \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int \frac{2e^{ac+bcx}}{1+e^{2ac+2bcx}} de^{ac+bcx} + e^{ac+bcx} \arctan(\operatorname{csch}(ac+bcx))}{bc} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 \int \frac{e^{ac+bcx}}{1+e^{2ac+2bcx}} de^{ac+bcx} + e^{ac+bcx} \arctan(\operatorname{csch}(ac+bcx))}{bc} \\
 & \quad \downarrow \text{240} \\
 & \frac{e^{ac+bcx} \arctan(\operatorname{csch}(ac+bcx)) + \log(e^{2ac+2bcx} + 1)}{bc}
 \end{aligned}$$

input

```
Int[E^(c*(a + b*x))*ArcTan[Csch[a*c + b*c*x]], x]
```

output

```
(E^(a*c + b*c*x)*ArcTan[Csch[a*c + b*c*x]] + Log[1 + E^(2*a*c + 2*b*c*x)]) / (b*c)
```

Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]`
- rule 5730 `Int[((a_.) + ArcTan[u_]*(b_.))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Simp[(a + b*ArcTan[u]) w, x] - Simp[b Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_.) + (d_.)*x)^(m_.)] /; FreeQ[{c, d, m}, x] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]`
- rule 7281 `Int[u_, x_Symbol] := With[{lst = FunctionOfLinear[u, x]}, Simp[1/lst[[3]] Subst[Int[lst[[1]], x], x, lst[[2]] + lst[[3]]*x], x] /; !FalseQ[lst]]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.54 (sec) , antiderivative size = 885, normalized size of antiderivative = 18.83

method	result	size
risch	Expression too large to display	885

input `int(exp(c*(b*x+a))*arctan(csch(b*c*x+a*c)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & I/b/c*\exp(c*(b*x+a))*\ln(\exp(c*(b*x+a))-I)-1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a)) \\ &)+I)^2)^3*\exp(c*(b*x+a))+1/2/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))+I))*c\text{sgn}(I*(\exp \\ & (c*(b*x+a))+I)^2)^2*\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))+I)^2) \\ & *c\text{sgn}(I/(-1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(c*(b*x+a))+I)^2/(-1+\exp(2*c*(b* \\ & x+a))))*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))+I)^2)*c\text{sgn}(I*(\exp \\ & (c*(b*x+a))+I)^2/(-1+\exp(2*c*(b*x+a))))^2*\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I \\ & *(\exp(c*(b*x+a))+I)^2*c\text{sgn}(I*(\exp(c*(b*x+a))+I)^2)*\exp(c*(b*x+a))+1/4/b/c \\ & * \text{Pi}*c\text{sgn}(I/(-1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(c*(b*x+a))+I)^2/(-1+\exp(2*c* \\ & (b*x+a))))^2*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2)*c\text{sgn}(I \\ & /(-1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2/(-1+\exp(2*c*(b*x+a)))) \\ & *\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I/(-1+\exp(2*c*(b*x+a))))*c\text{sgn}(I*(\exp(c*(b* \\ & x+a))-I)^2/(-1+\exp(2*c*(b*x+a))))^2*\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(\\ & c*(b*x+a))+I)^2/(-1+\exp(2*c*(b*x+a))))^3*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I* \\ & (\exp(c*(b*x+a))-I))^2*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2)*\exp(c*(b*x+a))-1/2/b/c* \\ & \text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))-I))*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2)^2*\exp(c*(b*x+a) \\ &)+1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2)^3*\exp(c*(b*x+a))-1/4/b/c*\text{Pi}*c\text{sgn} \\ & (I*(\exp(c*(b*x+a))-I)^2)*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2/(-1+\exp(2*c*(b*x+a))) \\ &)^2*\exp(c*(b*x+a))+1/4/b/c*\text{Pi}*c\text{sgn}(I*(\exp(c*(b*x+a))-I)^2/(-1+\exp(2*c*(b*x \\ & +a))))^3*\exp(c*(b*x+a))-2*a/b-I/b/c*\exp(c*(b*x+a))*\ln(\exp(c*(b*x+a))+I)+\ln \\ & (1+\exp(2*c*(b*x+a)))/b/c \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. $2(45) = 90$.

Time = 0.10 (sec) , antiderivative size = 131, normalized size of antiderivative = 2.79

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx$$

$$= \frac{(\cosh(bc x + ac) + \sinh(bc x + ac)) \arctan\left(\frac{2(\cosh(bc x + ac) + \sinh(bc x + ac))}{\cosh(bc x + ac)^2 + 2 \cosh(bc x + ac) \sinh(bc x + ac) + \sinh(bc x + ac)^2 - 1}\right) + \log\left(\frac{\cosh(bc x + ac) + \sinh(bc x + ac)}{\cosh(bc x + ac) - \sinh(bc x + ac)}\right)}{bc}$$

input `integrate(exp(c*(b*x+a))*arctan(csch(b*c*x+a*c)),x, algorithm="fricas")`

output $((\cosh(b*c*x + a*c) + \sinh(b*c*x + a*c))*\arctan(2*(\cosh(b*c*x + a*c) + \sinh(b*c*x + a*c))/(\cosh(b*c*x + a*c)^2 + 2*\cosh(b*c*x + a*c)*\sinh(b*c*x + a*c) + \sinh(b*c*x + a*c)^2 - 1)) + \log(2*\cosh(b*c*x + a*c)/(\cosh(b*c*x + a*c) - \sinh(b*c*x + a*c))))/(b*c)$

Sympy [F]

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx = e^{ac} \int e^{bcx} \operatorname{atan}(\operatorname{csch}(ac + bcx)) dx$$

input `integrate(exp(c*(b*x+a))*atan(csch(b*c*x+a*c)),x)`

output `exp(a*c)*Integral(exp(b*c*x)*atan(csch(a*c + b*c*x)), x)`

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx = \frac{\arctan(\operatorname{csch}(bcx + ac)) e^{((bx+a)c)}}{bc} + \frac{\log(e^{(2bcx+2ac)} + 1)}{bc}$$

input `integrate(exp(c*(b*x+a))*arctan(csch(b*c*x+a*c)),x, algorithm="maxima")`

output `arctan(csch(b*c*x + a*c))*e^((b*x + a)*c)/(b*c) + log(e^(2*b*c*x + 2*a*c) + 1)/(b*c)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.38

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx$$

$$= \frac{\left(\arctan\left(\frac{2e^{(bcx+ac)}}{e^{(2bcx+2ac)}-1}\right) e^{(bcx)} + e^{(-ac)} \log(e^{(2bcx+2ac)} + 1)\right) e^{(ac)}}{bc}$$

input `integrate(exp(c*(b*x+a))*arctan(csch(b*c*x+a*c)),x, algorithm="giac")`output `(arctan(2*e^(b*c*x + a*c)/(e^(2*b*c*x + 2*a*c) - 1))*e^(b*c*x) + e^(-a*c)*log(e^(2*b*c*x + 2*a*c) + 1))*e^(a*c)/(b*c)`**Mupad [B] (verification not implemented)**

Time = 0.68 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.43

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx = \frac{\ln(e^{2bcx} e^{2ac} + 1)}{bc}$$

$$+ \frac{e^{bcx} e^{ac} \operatorname{atan}\left(\frac{1}{\frac{e^{bcx} e^{ac}}{2} - \frac{e^{-bcx} e^{-ac}}{2}}\right)}{bc}$$

input `int(atan(1/sinh(a*c + b*c*x))*exp(c*(a + b*x)),x)`output `log(exp(2*b*c*x)*exp(2*a*c) + 1)/(b*c) + (exp(b*c*x)*exp(a*c)*atan(1/((exp(b*c*x)*exp(a*c))/2 - (exp(-b*c*x)*exp(-a*c))/2)))/(b*c)`

Reduce [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.34

$$\int e^{c(a+bx)} \arctan(\operatorname{csch}(ac + bcx)) dx = \frac{e^{bcx+ac} \operatorname{atan}\left(\frac{2e^{bcx+ac}}{e^{2bcx+2ac}-1}\right) + \log(e^{2bcx+2ac} + 1)}{bc}$$

input `int(exp(c*(b*x+a))*atan(csch(b*c*x+a*c)),x)`

output `(e**(a*c + b*c*x)*atan((2*e**(a*c + b*c*x))/(e**(2*a*c + 2*b*c*x) - 1)) + log(e**(2*a*c + 2*b*c*x) + 1))/(b*c)`

3.153 $\int \frac{(a+b \arctan(cx^n))(d+e \log(fx^m))}{x} dx$

Optimal result	1122
Mathematica [C] (verified)	1123
Rubi [A] (verified)	1123
Maple [C] (warning: unable to verify)	1124
Fricas [B] (verification not implemented)	1125
Sympy [F(-1)]	1126
Maxima [F]	1126
Giac [F]	1126
Mupad [F(-1)]	1127
Reduce [F]	1127

Optimal result

Integrand size = 24, antiderivative size = 163

$$\int \frac{(a+b \arctan(cx^n))(d+e \log(fx^m))}{x} dx$$

$$= ad \log(x) + \frac{ae \log^2(fx^m)}{2m} + \frac{ibd \operatorname{PolyLog}(2, -icx^n)}{2n}$$

$$+ \frac{ibe \log(fx^m) \operatorname{PolyLog}(2, -icx^n)}{2n}$$

$$- \frac{ibd \operatorname{PolyLog}(2, icx^n)}{2n} - \frac{ibe \log(fx^m) \operatorname{PolyLog}(2, icx^n)}{2n}$$

$$- \frac{ibem \operatorname{PolyLog}(3, -icx^n)}{2n^2} + \frac{ibem \operatorname{PolyLog}(3, icx^n)}{2n^2}$$

output

```
a*d*ln(x)+1/2*a*e*ln(f*x^m)^2/m+1/2*I*b*d*polylog(2,-I*c*x^n)/n+1/2*I*b*e*
ln(f*x^m)*polylog(2,-I*c*x^n)/n-1/2*I*b*d*polylog(2,I*c*x^n)/n-1/2*I*b*e*1
n(f*x^m)*polylog(2,I*c*x^n)/n-1/2*I*b*e*m*polylog(3,-I*c*x^n)/n^2+1/2*I*b*
e*m*polylog(3,I*c*x^n)/n^2
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.18 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.71

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx$$

$$= -\frac{bcemx^n {}_4F_3\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}, \frac{3}{2}; -c^2x^{2n}\right)}{n^2}$$

$$+ \frac{bcx^n {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}; -c^2x^{2n}\right)(d + e \log(fx^m))}{n}$$

$$+ \frac{1}{2}a \log(x) (2d - em \log(x) + 2e \log(fx^m))$$

input

```
Integrate[((a + b*ArcTan[c*x^n])*(d + e*Log[f*x^m]))/x,x]
```

output

```
-((b*c*e*m*x^n*HypergeometricPFQ[{1/2, 1/2, 1/2, 1}, {3/2, 3/2, 3/2}, -(c^2*x^(2*n))])/n^2) + (b*c*x^n*HypergeometricPFQ[{1/2, 1/2, 1}, {3/2, 3/2}, -(c^2*x^(2*n))]*(d + e*Log[f*x^m]))/n + (a*Log[x]*(2*d - e*m*Log[x] + 2*e*Log[f*x^m]))/2
```

Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx$$

$$\downarrow \text{7293}$$

$$\int \left(\frac{d(a + b \arctan(cx^n))}{x} + \frac{e \log(fx^m)(a + b \arctan(cx^n))}{x} \right) dx$$

$$\downarrow \text{2009}$$

$$ad \log(x) + \frac{ae \log^2(fx^m)}{2m} + \frac{ibd \operatorname{PolyLog}(2, -icx^n)}{2n} - \frac{ibd \operatorname{PolyLog}(2, icx^n)}{2n} + \frac{ibe \operatorname{PolyLog}(2, -icx^n) \log(fx^m)}{2n} - \frac{ibe \operatorname{PolyLog}(2, icx^n) \log(fx^m)}{2n} - \frac{ibem \operatorname{PolyLog}(3, -icx^n)}{2n^2} + \frac{ibem \operatorname{PolyLog}(3, icx^n)}{2n^2}$$

input `Int[((a + b*ArcTan[c*x^n])*(d + e*Log[f*x^m]))/x,x]`

output `a*d*Log[x] + (a*e*Log[f*x^m]^2)/(2*m) + ((I/2)*b*d*PolyLog[2, (-I)*c*x^n])/n + ((I/2)*b*e*Log[f*x^m]*PolyLog[2, (-I)*c*x^n])/n - ((I/2)*b*d*PolyLog[2, I*c*x^n])/n - ((I/2)*b*e*Log[f*x^m]*PolyLog[2, I*c*x^n])/n - ((I/2)*b*e*m*PolyLog[3, (-I)*c*x^n])/n^2 + ((I/2)*b*e*m*PolyLog[3, I*c*x^n])/n^2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 159.48 (sec) , antiderivative size = 547, normalized size of antiderivative = 3.36

method	result
risch	$\frac{\left(-\frac{i\pi \operatorname{csgn}(if) \operatorname{csgn}(ix^m) \operatorname{csgn}(ifx^m)}{4} + \frac{i\pi \operatorname{csgn}(if) \operatorname{csgn}(ifx^m)^2}{4} + \frac{i\pi \operatorname{csgn}(ix^m) \operatorname{csgn}(ifx^m)^2}{4} - \frac{i\pi \operatorname{csgn}(ifx^m)^3}{4} + \frac{e \ln(f)}{2} + \frac{d}{2}\right)}{n} (-ib \operatorname{di} \dots)$

input `int((a+b*arctan(c*x^n))*(d+e*ln(f*x^m))/x,x,method=_RETURNVERBOSE)`

output

```
(-1/4*I*e*Pi*csgn(I*f)*csgn(I*x^m)*csgn(I*f*x^m)+1/4*I*e*Pi*csgn(I*f)*csgn
(I*f*x^m)^2+1/4*I*e*Pi*csgn(I*x^m)*csgn(I*f*x^m)^2-1/4*I*e*Pi*csgn(I*f*x^m
)^3+1/2*e*ln(f)+1/2*d)/n*(-I*b*dilog(1-I*c*x^n)+2*ln(x^n)*a+I*b*dilog(1+I*
c*x^n))-1/2*I*e*b/n*ln(-I*(-c*x^n+I))*ln(-I*c*x^n)*ln(x^m)-1/2*I*b*e*m*pol
ylog(3,-I*c*x^n)/n^2-1/2*I*e*b*ln(x)*ln(-I*(c*x^n+I))*ln(x^m)+1/2*I*e*b*ln
(x)*ln(1-I*c*x^n)*ln(x^m)+1/2*I*e*b*ln(x)^2*ln(-I*(c*x^n+I))*m-1/2*I*e*b*l
n(x)^2*ln(1-I*c*x^n)*m+1/2*I*e*b/n*dilog(-I*(c*x^n+I))*m*ln(x)+1/2*I*b*e*m
*polylog(3,I*c*x^n)/n^2+1/2*e*a/m*ln(x^m)^2+1/2*I*e*b*ln(x)*ln(-I*(-c*x^n+
I))*ln(x^m)+1/2*I*e*b*m/n*ln(x)*polylog(2,-I*c*x^n)+1/2*I*e*b*ln(x)^2*ln(1
+I*c*x^n)*m-1/2*I*e*b*m/n*ln(x)*polylog(2,I*c*x^n)+1/2*I*e*b/n*ln(-I*(-c*x
^n+I))*ln(-I*c*x^n)*m*ln(x)+1/2*I*e*b/n*dilog(-I*c*x^n)*m*ln(x)-1/2*I*e*b*
ln(x)*ln(1+I*c*x^n)*ln(x^m)-1/2*I*e*b/n*dilog(-I*c*x^n)*ln(x^m)-1/2*I*e*b*
ln(x)^2*ln(-I*(-c*x^n+I))*m-1/2*I*e*b/n*dilog(-I*(c*x^n+I))*ln(x^m)
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 250 vs. $2(121) = 242$.

Time = 0.13 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.53

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx$$

$$= \frac{2aemn^2 \log(x)^2 + 2i bempolylog(3, icx^n) - 2i bempolylog(3, -icx^n) + 2(bemn^2 \log(x)^2 + 2(ben^2 \log$$

input

```
integrate((a+b*arctan(c*x^n))*(d+e*log(f*x^m))/x,x, algorithm="fricas")
```

output

```
1/4*(2*a*e*m*n^2*log(x)^2 + 2*I*b*e*m*polylog(3, I*c*x^n) - 2*I*b*e*m*poly
log(3, -I*c*x^n) + 2*(b*e*m*n^2*log(x)^2 + 2*(b*e*n^2*log(f) + b*d*n^2)*lo
g(x))*arctan(c*x^n) - 2*(I*b*e*m*n*log(x) + I*b*e*n*log(f) + I*b*d*n)*dilo
g(I*c*x^n) - 2*(-I*b*e*m*n*log(x) - I*b*e*n*log(f) - I*b*d*n)*dilog(-I*c*x
^n) + (I*b*e*m*n^2*log(x)^2 - 2*(-I*b*e*n^2*log(f) - I*b*d*n^2)*log(x))*lo
g(I*c*x^n + 1) + (-I*b*e*m*n^2*log(x)^2 - 2*(I*b*e*n^2*log(f) + I*b*d*n^2)
*log(x))*log(-I*c*x^n + 1) + 4*(a*e*n^2*log(f) + a*d*n^2)*log(x))/n^2
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx = \text{Timed out}$$

input `integrate((a+b*atan(c*x**n))*(d+e*ln(f*x**m))/x,x)`

output `Timed out`

Maxima [F]

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx = \int \frac{(b \arctan(cx^n) + a)(e \log(fx^m) + d)}{x} dx$$

input `integrate((a+b*arctan(c*x^n))*(d+e*log(f*x^m))/x,x, algorithm="maxima")`

output `1/2*a*e*log(f*x^m)^2/m + a*d*log(x) - 1/2*(b*e*m*log(x)^2 - 2*b*e*log(x)*log(x^m) - 2*(b*e*log(f) + b*d)*log(x))*arctan(c*x^n) - integrate(-1/2*(b*c*e*m*n*x^n*log(x)^2 - 2*b*c*e*n*x^n*log(x)*log(x^m) - 2*(b*c*e*log(f) + b*c*d)*n*x^n*log(x))/(c^2*x*x^(2*n) + x), x)`

Giac [F]

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx = \int \frac{(b \arctan(cx^n) + a)(e \log(fx^m) + d)}{x} dx$$

input `integrate((a+b*arctan(c*x^n))*(d+e*log(f*x^m))/x,x, algorithm="giac")`

output `integrate((b*arctan(c*x^n) + a)*(e*log(f*x^m) + d)/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx = \int \frac{(a + b \operatorname{atan}(cx^n))(d + e \ln(fx^m))}{x} dx$$

input `int(((a + b*atan(c*x^n))*(d + e*log(f*x^m)))/x,x)`

output `int(((a + b*atan(c*x^n))*(d + e*log(f*x^m)))/x, x)`

Reduce [F]

$$\int \frac{(a + b \arctan(cx^n))(d + e \log(fx^m))}{x} dx$$

$$= \frac{2 \left(\int \frac{\operatorname{atan}(x^n c)}{x} dx \right) b d m + 2 \left(\int \frac{\operatorname{atan}(x^n c) \log(x^m f)}{x} dx \right) b e m + \log(x^m f)^2 a e + 2 \log(x) a d m}{2m}$$

input `int((a+b*atan(c*x^n))*(d+e*log(f*x^m))/x,x)`

output `(2*int(atan(x**n*c)/x,x)*b*d*m + 2*int((atan(x**n*c)*log(x**m*f))/x,x)*b*e*m + log(x**m*f)**2*a*e + 2*log(x)*a*d*m)/(2*m)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 1128
4.2 Links to plain text integration problems used in this report for each CAS . 1146

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
          ]
        ,(*ELSE*)
          finalresult={"C","Result contains complex when optimal does not."}
        ]
      ,(*ELSE*)(*result does not contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
          ]
        ]
      ,(*ELSE*)(*expnResult>expnOptimal*)
        If[FreeQ[result,Integrate] && FreeQ[result,Int],
          finalresult={"C","Result contains higher order function than in optimal. Order "
          ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
If[AppellFunctionQ[Head[expn]],
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
If[Head[expn]===RootSum,
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
If[Head[expn]===Integrate || Head[expn]===Int,
Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
9]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
MemberQ[{
Exp, Log,
Sin, Cos, Tan, Cot, Sec, Csc,
ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]
```

```
SpecialFunctionQ[func_] :=
MemberQ[{
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
}, func]
```

```
HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]
```

```
AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]
```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9

```



```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```



```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file